A 17-variable degree-twenty equation

Nonnegative integer unknowns; 104 binary-operation abbreviations; exact degree 20.

The 17 unknowns are the following distinct names, each a nonnegative integer:
`argJ, argFuel, argAlpha, argPellF, argIndex, argTime, argX`
`argY, argOrdinate, argEta, argZeta, argAuxOrd, argGamma, argRank`
`argFirstQ, argMainQ, argVroot`
Each monospaced G-name below is an abbreviation, not an additional unknown. The fixed integers β, K, μ, ω are defined after the equation.

$\texttt{G001} = \texttt{argFuel} + 1$
$\texttt{G002} = \texttt{argAlpha} + 1$
$\texttt{G003} = \texttt{argIndex} + 1$
$\texttt{G004} = \texttt{argX} + 1$
$\texttt{G005} = \texttt{argY} + 1$
$\texttt{G006} = \texttt{argOrdinate} + 1$
$\texttt{G007} = \texttt{argEta} + 1$
$\texttt{G008} = \texttt{argZeta} + 1$
$\texttt{G009} = \texttt{argGamma} + 1$
$\texttt{G010} = \texttt{argRank} + 1$
$\texttt{G011} = \texttt{argJ} \cdot \beta$
$\texttt{G012} = 1 + \texttt{G011}$
$\texttt{G013} = \texttt{G012} \cdot \texttt{G012}$
$\texttt{G014} = \texttt{G012} \cdot \texttt{G013}$
$\texttt{G015} = \texttt{G005} \cdot \texttt{G013}$
$\texttt{G016} = \texttt{G012} - \texttt{G001}$
$\texttt{G017} = \texttt{G016} - \texttt{G002}$
$\texttt{G018} = \texttt{G004} \cdot \texttt{G005}$
$\texttt{G019} = \texttt{G007} + \texttt{G008}$
$\texttt{G020} = \texttt{G005} \cdot \texttt{G019}$
$\texttt{G021} = \texttt{G007} + \texttt{G020}$
$\texttt{G022} = \texttt{G021} \cdot \texttt{G021}$
$\texttt{G023} = \texttt{G018} + \texttt{G005}$
$\texttt{G024} = 4 \cdot \texttt{G023}$
$\texttt{G025} = 3 + \texttt{G024}$
$\texttt{G026} = \texttt{G023} \cdot \texttt{G023}$
$\texttt{G027} = \texttt{G025} + \texttt{G026}$
$\texttt{G028} = \texttt{G023} \cdot \texttt{G021}$
$\texttt{G029} = \texttt{G004} + \texttt{G028}$
$\texttt{G030} = \texttt{G018} \cdot \texttt{G020}$
$\texttt{G031} = K \cdot \texttt{G012}$
$\texttt{G032} = \texttt{G004} + \texttt{G031}$
$\texttt{G033} = \texttt{G017} \cdot \texttt{G032}$
$\texttt{G034} = \texttt{G001} \cdot \texttt{G012}$
$\texttt{G035} = \texttt{G033} - \texttt{G034}$
$\texttt{G036} = \texttt{G030} + \texttt{G019}$
$\texttt{G037} = \texttt{G036} + \texttt{G006}$
$\texttt{G038} = \texttt{G019} \cdot \texttt{G035}$
$\texttt{G039} = \texttt{G037} - \texttt{G038}$
$\texttt{G040} = \texttt{G005} \cdot \texttt{G011}$
$\texttt{G041} = \texttt{argFirstQ} \cdot \texttt{argFirstQ}$
$\texttt{G042} = \texttt{G030} \cdot \texttt{G036}$
$\texttt{G043} = 1 + \texttt{G042}$
$\texttt{G044} = \texttt{G041} - \texttt{G043}$
$\texttt{G045} = \texttt{G016} \cdot \texttt{G011}$
$\texttt{G046} = \texttt{G002} + \texttt{G045}$
$\texttt{G047} = \texttt{G013} - 1$
$\texttt{G048} = \texttt{G046} \cdot \texttt{G047}$
$\texttt{G049} = \omega \cdot \texttt{G012}$
$\texttt{G050} = \mu + \texttt{G049}$
$\texttt{G051} = \texttt{argJ} \cdot \texttt{G050}$
$\texttt{G052} = \texttt{G048} + \texttt{G051}$
$\texttt{G053} = \texttt{G019} - \texttt{G010}$
$\texttt{G054} = \texttt{G053} - 1$
$\texttt{G055} = \texttt{G015} \cdot \texttt{G009}$
$\texttt{G056} = \texttt{G018} \cdot \texttt{G012}$
$\texttt{G057} = \texttt{G055} + \texttt{G056}$
$\texttt{G058} = \texttt{G057} - \texttt{G054}$
$\texttt{G059} = \texttt{G025} \cdot \texttt{G058}$
$\texttt{G060} = \texttt{argMainQ} \cdot \texttt{argMainQ}$
$\texttt{G061} = \texttt{G027} \cdot \texttt{G022}$
$\texttt{G062} = 1 + \texttt{G061}$
$\texttt{G063} = \texttt{G060} - \texttt{G062}$
$\texttt{G064} = \texttt{argPellF} \cdot \texttt{argPellF}$
$\texttt{G065} = \texttt{G064} - 1$
$\texttt{G066} = \texttt{G027} \cdot \texttt{G065}$
$\texttt{G067} = \texttt{G022} \cdot \texttt{G003}$
$\texttt{G068} = \texttt{G067} \cdot \texttt{G067}$
$\texttt{G069} = \texttt{G068} - \texttt{G066}$
$\texttt{G070} = \texttt{argTime} \cdot \texttt{G021}$
$\texttt{G071} = \texttt{G070} - \texttt{G010}$
$\texttt{G072} = \texttt{G064} \cdot \texttt{G071}$
$\texttt{G073} = \texttt{G072} - \texttt{G021}$
$\texttt{G074} = \texttt{argAuxOrd} \cdot \texttt{argAuxOrd}$
$\texttt{G075} = \texttt{argVroot} \cdot \texttt{argVroot}$
$\texttt{G076} = \texttt{G074} - \texttt{G075}$
$\texttt{G077} = \texttt{G066} \cdot \texttt{G076}$
$\texttt{G078} = \texttt{G074} - \texttt{G077}$
$\texttt{G079} = \texttt{G078} - 1$
$\texttt{G080} = \texttt{G044} \cdot \texttt{G044}$
$\texttt{G081} = \texttt{G063} \cdot \texttt{G063}$
$\texttt{G082} = \texttt{G079} \cdot \texttt{G079}$
$\texttt{G083} = \texttt{G069} \cdot \texttt{G069}$
$\texttt{G084} = \texttt{G080} + \texttt{G081}$
$\texttt{G085} = \texttt{G084} + \texttt{G082}$
$\texttt{G086} = \texttt{G085} + \texttt{G083}$
$\texttt{G087} = \texttt{G010} - \texttt{G052}$
$\texttt{G088} = \texttt{G087} \cdot \texttt{G087}$
$\texttt{G089} = \texttt{G086} + \texttt{G088}$
$\texttt{G090} = \texttt{argFirstQ} - \texttt{G039}$
$\texttt{G091} = \texttt{G011} \cdot \texttt{G090}$
$\texttt{G092} = \texttt{G091} + \texttt{G035}$
$\texttt{G093} = \texttt{G014} \cdot \texttt{G092}$
$\texttt{G094} = \texttt{G093} - \texttt{G040}$
$\texttt{G095} = \texttt{G094} \cdot \texttt{G094}$
$\texttt{G096} = \texttt{argMainQ} - \texttt{G029}$
$\texttt{G097} = \texttt{G015} \cdot \texttt{G096}$
$\texttt{G098} = \texttt{G097} - \texttt{G059}$
$\texttt{G099} = \texttt{G098} \cdot \texttt{G098}$
$\texttt{G100} = \texttt{argVroot} - \texttt{G073}$
$\texttt{G101} = \texttt{G100} \cdot \texttt{G100}$
$\texttt{G102} = \texttt{G089} + \texttt{G095}$
$\texttt{G103} = \texttt{G102} + \texttt{G099}$
$\texttt{G104} = \texttt{G103} + \texttt{G101}$

The single equation is $\boxed{\texttt{G104} = 0}$.

This is an external source-mapping candidate. Its zero set has been related to an earlier 20-variable equation; a Lean proof of the new 104-operation endpoint has not been completed.

The four fixed integers. Every index, set, sum, product and floor below defines a constant, not another unknown. Put

$$\rho = 2^{47}, \quad B = \rho^{2011}, \quad \beta = B - 1,$$
$$G_m(x) = \sum_{j=0}^{m-1} x^j, \quad \Lambda = \rho^{1005}.$$

$$\begin{aligned}\mu &= \beta - G_{951}(\rho) \\ &\quad - \rho^{960} G_{26}(\rho) - \rho^{1004}, \\ \omega &= \beta + 8G_{917}(\rho) - 4 + (2^{45} - 4)\Lambda \\ &\quad + \rho^{960} G_9(\rho) + \rho^{987} G_8(\rho) + \rho^{1004}, \\ K &= \rho^{18} + \rho^{1004} + \rho^{2020} \\ &\quad + 2\Lambda\Phi(1/\rho),\end{aligned}$$

where

$$\begin{aligned}\chi_i &= 1 + 2^{10}(1 + w_i + 2^8 n_i \\ &\quad + 2^{17} p_i + 2^{25} s_i) \quad (0 \le i \le 941),\end{aligned}$$
$$\pi(0) = 0, \quad \pi(i) = 1 + \sum_{j=1}^{941} \left\lfloor \frac{2^{44} - 1 + \chi_j - \chi_i}{2^{44}} \right\rfloor \quad (1 \le i \le 941).$$

$$\begin{aligned}\Phi(x) = &\sum_{i=0}^{941} \chi_i x^{\pi(i)} \\ &+ 2^{10} x^{969}(x^9 + 2^{17}) G_8(2x) \\ &+ 2^{18} x^{942}(x^{18} + 2^{17}) G_9(2x).\end{aligned}$$

The rational intermediate in K cancels exactly; all four results are positive integers.

The 942 indexed fixed tiles. Use the actions u_j, v_j, d_j decoded below. For $d = 0, 1$, let

$$h(q,b) = 2q + b + 2, \quad \mathcal{S}_d = \{u_j : d_j = d,\ 0 \le j < 430\},$$
$$r_{d,q} = 205 - 199d + |\mathcal{S}_d \cap \{0, \dots, q-1\}|.$$

An interval ending at -1 is empty. The table defines t_i for $0 \le i \le 941$; all index ranges are fixed:

index i	tile $t_i = (w_i, n_i, p_i, s_i)$
0	$(2, 435, 3, 1)$
$j+1$ $(j = 2q + b < 430)$	$((1-d_j) r_{0,u_j}, h(q,b), d_j r_{1,u_j}, v_j)$
$433 - 2a + b$ $(a, b = 0, 1)$	$(0, 432a + b, 0, 432a + b)$
$\substack{423+2r_{d,q}+b \\ q \in \mathcal{S}_d,\ d,b=0,1}$	$(dr_{d,q}, b, (1-d) r_{d,q}, h(q,b))$
$923 + 2a + b$ $(a, b = 0, 1)$	$(0, 432a + b, 0, 438)$
927	$(2, 434, 2, 0)$
928	$(3, 436, 4, 4)$
929	$(4, 436, 4, 0)$
$496 + t$ $(t = 434, 435, 436)$	$(1, 438, 1, t)$
$933 + 3a + b$ $(a, b = 0, 1, 2)$	$(a^2, 437, b, 437)$

A closed decoder for the 430 actions. Put $D_b(A, k) = \lfloor A / b^k \rfloor \bmod b$, where $a \bmod b = a - b\lfloor a/b \rfloor$. For $0 \le q < 215$, $b = 0, 1$, $j = 2q + b$:

$$\mathcal{N} = \mathcal{A} \bmod 2^{636}, \quad \mathcal{T} = \lfloor \mathcal{A} / 2^{636} \rfloor,$$
$$\nu_j = D_2(\mathcal{N}, j), \quad \kappa_j = (\sigma_j + 2)^{1-\nu_j} \quad (0 \le j < 430),$$
$$\sigma_j = \sum_{k=0}^{j-1} \nu_k, \quad \Pi_j = \prod_{k=0}^{j-1} \kappa_k \quad (0 \le j \le 430),$$
$$m_q = D_2(\mathcal{N}, 430 + q), \quad \tau_q = \sum_{k=0}^{q-1} m_k,$$
$$F_q = 14 + m_q (D_{16}(\lfloor \mathcal{T} / \Pi_{430} \rfloor, \tau_q) - 14),$$
$$u_j = (\sigma_{j+1} - (\lfloor \mathcal{T} / \Pi_j \rfloor \bmod \kappa_j)) \bmod 216,$$
$$v_j = D_2(F_q, 2b), \quad d_j = D_2(F_q, 2b + 1).$$

Empty sums/products are 0/1. Here $(v_0, d_0, u_0) = (1, 1, 1)$; the constructor tile uses this actual first target, including its state renaming.

The integer $\mathcal{A}$ in base 10. Its complete decimal expansion is the concatenation of these lines:

```
12889260410359561746736540743515845134244260640012702033212621
95398074180784058571518488282119861023093355389406939414042892
97455773992193784795705001618777025966369137954859764532832722
75411843410809038713502633204617333246733616940964578272360901
56186216508504953649192752154519268024509952082950238865468307
22999632482769253980274091217460707829023970817244790340704449
09085581271192071656170732004937337082171141082486384269072884
72778087419663417942892520175630649792329265834490253035577533
96449689808375788457488971017043689875433162972390177936123032
89025657757489526342497438772045714099661532481123050235898973
27443609903800541199351405985754278307227777584454508422717998
33411356440933534607489268032843 7
```