

Iterated Euler diagonals (OEIS A290353 and A290354)

Kotěšovec’s asymptotic form, the first logarithmic correction,
and the logarithmic height crossover

Three research reports of one external research session
(bundle Reports 225, 226 and 228), merged into one ProveIt report

All three Parts: 5 October 2026 merged: 6 October 2026

Abstract

Let $F_0(z) = z$ and $1 + F_{m+1}(z) = \prod_{d \geq 1} (1 - z^d)^{-[z^d]F_m(z)}$, so that $A(n, m) = [z^n]F_m(z)$ counts unlabeled rooted trees with n leaves, all at level m (OEIS A290353), and $a_n = A(n, n)$ is OEIS A290354. This report prints three manuscripts as Parts I–III, in the order in which each depends on the one before. Part I proves $a_n \sim c(2n/\pi)^n n^{-4/3}$ with $c = T^{1/3}e^{-B}h(1) > 0$, $T = \pi/2$, $B = \frac{1}{2} - \frac{\pi}{6} - \frac{\pi^2}{16} - \frac{\log 2}{6}$, where h is the smooth, strictly positive density of a positive measure whose Laplace transform is the entire parametrization P of $w \mapsto e^w - 1$ with normalized Fatou coordinate $-2/w + \frac{1}{3} \log w + O(w)$; this proves the form of the asymptotic that Václav Kotěšovec stated as a conjecture on A290354 (August 14, 2017), while the digits of $c \approx 4.4923$ remain uncertified. Part I also proves proportional-height asymptotics for $A(n, m)$ with n/m in a compact subset of $(0, \infty)$, fixed and square-root height shifts with the floor phase, and a Lambert- W inverse at exact sequence values. Part II proves the first logarithmic correction $1 + (-\frac{1}{18} \log^2 n + A \log n + C)/n + O(n^{-7/5})$, with A and C exact in $h(1), h'(1), h''(1)$ and a new clock constant κ in closed form in $\pi, \log 2$ and Catalan’s constant, the corresponding uniform proportional-height correction and a sharpened inverse. Part III proves $A(n, m)/(2T^{-n}m^{n-1}) \rightarrow e^{-1/(3\lambda)}$ when $m/(n \log n) \rightarrow \lambda \in (0, \infty)$, with every fixed inverse-logarithm order, through an exact Hankel representation of h near zero, $h(0) = 2$, a smooth germ $t^{t/3}h(t)$ and rational-plus-zeta coefficients; it also gives Kaneiwa’s normalization and smooth height prescriptions. Validated digits of c , higher corrections, effective constants, the regime $n/m \rightarrow \infty$ and a conjectured identity linking h with the amplitude of the labelled iterated Bell numbers (a neighbouring report) remain open. The report is unrefereed, and nothing in it is formalized.

Contents

Guide to this report	4
What is proved, and where	5
The OEIS statements, and the inverses	6
Notation across the three Parts	7
Provenance and merge decisions	8
What was checked, and what was not	10
Relation to neighbouring reports and formal projects	10

I	Diagonal and proportional height asymptotics for iterated Euler transforms	12
1	Definition and statement	12
2	Subsidiary terms and the nonautonomous clock	14
3	Matching and the positive real orbit	15
4	The repelling coordinate with uniform sectors	17
5	The local limit and angular tails	19
6	The positive measure and its density	21
7	Coefficient extraction and the amplitude	22
8	Proportional height and fixed shifts	23
9	An inverse consequence	24
10	Exact computation and numerical diagnostics	25
11	Further questions and quantitative extensions	27
II	First logarithmic correction for iterated Euler diagonals	29
12	Definition and main results	29
13	Analytic foundation from Report 225	31
14	Subsidiary forcing through fifth scaled order	33
15	Two further nonautonomous clock terms	34
16	Weighted defect and accumulated clock error	36
17	Finite Fatou expansion and forced matching	37
18	Uniform local limit on a growing strip	38
19	Derivative extraction and classical Fourier inversion	40
20	Uniform proportional height correction	42
21	A sharpened inverse formula	42
22	Exact computation and numerical diagnostics	43
23	Further questions and the boundary of the theorem	45
III	Logarithmic height crossover for iterated Euler transforms	47
24	The array and the crossover	47
25	Imported analytic interfaces and the scope of the result	49

26	Finite Fatou expansions with sectorial remainders	52
27	The endpoint value by direct Fourier inversion	52
28	An exact Hankel representation	53
29	A smooth endpoint germ and the logarithmic transseries	54
30	The rational and zeta coefficient structure	56
31	Transfer to shrinking degree to height ratios	57
32	Exact logarithmic cancellation and the crossover proof	58
33	Kaneiwa normalization and its first correction	59
34	Smooth suppression ratio inverses	59
35	Exact finite comparisons and reproducibility	62
36	Further questions and limits	63
37	Further questions and research (all three Parts)	65

Guide to this report

The three Parts are three manuscripts of the session bundle of Reports 1–243 (arrival commit 60f54ea06), placed in this directory by commit 47fc7a069 (batch 106). They study one object, the iterated Euler transforms F_m of $1 + x$ and their coefficient array $A(n, m)$:

Part	Bundle report	Delivered title	Labels
I	225 (base)	Diagonal and proportional height asymptotics for iterated Euler transforms	<code>ied:lead:</code>
II	226	First logarithmic correction for iterated Euler diagonals	<code>ied:corr:</code>
III	228	Logarithmic height crossover for iterated Euler transforms	<code>ied:log:</code>

Arrangement. The base of the merge is Report 225, whose `report225.tex` was staged as this `article.tex`. It proves the leading theorems and constructs the analytic foundation (the nonautonomous clock, the sectorial Fatou coordinate, the positive Laplace density h), so it is printed first, as Part I, with its section, statement and equation numbers unchanged. Report 226 uses Part I’s analytic interface and refines its clock and its coefficient extraction to the first correction; it is Part II. Report 228 uses both, Part I’s coordinate and density and Part II’s absolute derivative estimates, and pushes the transfer to degree-to-height ratios shrinking like a power of $1/\log m$; it is Part III. The three Parts thus appear in dependency order, which is also the order of writing.

Embedded copies. Report 226’s archive carries Report 225’s source and PDF under `supporting/`, and Report 228’s carries those of Reports 225 and 226; all six files are byte-identical to the standalone deliveries (SHA-256 at placement, `cmp` again at the write). They are printed once each, as Parts I and II, and are not shipped. Report 226’s archive also carries byte copies of eight code and data files of Report 225 (its ledger says “copied unchanged”); they are shipped once, under Report 225’s prefix.

Restatements. Part II’s Section 13 and Part III’s Section 25 state, in their own words, the results they import (from Part I, and for Part III also from Part II). They are printed in full, with notes naming the Part I or Part II statement behind each item, because their displays are cited by the later proofs, because Part II states as interface items several estimates that Part I proves only inside its proofs (the inverse-orbit bound and the disk and neighbourhood forms of the tail and outer bounds), and because Part III’s contains material that is not a restatement (its L^1 consequence (III.16) and its comparison with the iterated Bell report). (Wording corrected after the independent check of 6 October 2026; the first wording read “because they contain material that is not a restatement (Part II’s bound on inverse orbits, Part III’s L^1 consequence ...)”). Part II’s bound on inverse orbits is the estimate $|G^j(u)| \geq c(|u| + j)$ of Part I’s proof of Lemma 4.1.)

Numbering. Sections are numbered continuously and statements within sections, so a statement number moves with its section only. Part I keeps Report 225’s Sections 1–11. Report 226’s Section k is Section $k + 11$ here (Sections 12–23): its Theorems 1.1 and 1.2 are Theorems 12.1 and 12.2, its Corollary 10.1 is Corollary 21.1. Report 228’s Section k is Section $k + 23$ here (Sections 24–36): its Theorem 1.1 is Theorem 24.1, its Lemma 3.1, Proposition 4.1, Theorems 5.1, 6.1, 8.1 are 26.1, 27.1, 28.1, 29.1, 31.1, and its Proposition 11.1 and Theorems 11.2–11.3 are 34.1, 34.2 and 34.3. The manuscripts number equations consecutively through each document; each Part keeps its manuscript’s equation numbers, prefixed in Parts II and III: Report 226’s equation (k) is (II. k) here and Report 228’s is (III. k). Section 37 was added in the write; no delivered number moved.

Notes added when the manuscripts were merged are set in small type and tagged [**write**], with their date (6 October 2026). They give cross-references between the Parts, mark statements that a later Part extends or answers, mark results proved twice, and record what the write checked. Superseded statements and answered questions are printed as delivered, with a dated note. No delivered statement was changed, and no delivered symbol was renamed. Each Part has a section of open questions near its end: Part I its Section 11, Part II its Section 23, Part III its Section 36; Section 37 gathers them for the whole report under Vladimir's standing rule of 4 October 2026, with their sources, sketches and what is missing, and adds the question shared with the iterated Bell report.

What is proved, and where

Here $T = \pi/2$, $B = \frac{1}{2} - \frac{\pi}{6} - \frac{\pi^2}{16} - \frac{\log 2}{6}$, $f(w) = e^w - 1$, P is the entire solution of $P(s+1) = e^{P(s)} - 1$ whose inverse is the normalized Fatou coordinate $\Phi(w) = -2/w + \frac{1}{3} \log w + O(w)$, h is the density with $P(s) = \int_0^\infty e^{st} h(t) dt$, and $\beta = n/m$.

Statement	Where
P entire, the Laplace transform of a finite positive measure with every exponential moment, no atom at zero, support $[0, \infty)$; its density h is C^∞ and strictly positive on $(0, \infty)$	Part I, Theorem 1.1, Lemma 4.1, Proposition 6.1, Corollary 6.2
$a_n \sim c(n/T)^n n^{-4/3}$, $c = T^{1/3} e^{-B} h(1) > 0$, with an absolutely convergent Fourier formula for $h(1)$ (Kotěšovec's conjectured form on A290354)	Part I, Theorem 1.1
$A(n, m) \sim h(\beta) m^{-1} (m/T)^n m^{-\beta/3} T^{\beta/3} e^{-\beta B}$ uniformly for β in a compact subset of $(0, \infty)$; ratio e^k for fixed shifts; floor phase $e^{-\{\lambda n\}/\lambda}$; Gaussian-shaped profile for $ k \leq C\sqrt{n}$	Part I, Theorem 8.1, Corollaries 8.2 and 8.3
Lambert- W inverse	Part I, Corollary 9.1; sharpened in Part II, Corollary 21.1
$n = X + (\frac{4}{3} \log X - \log c)/q + o(1/\log X)$, nearest-integer recovery at exact sequence values (no onset)	
Subsidiary forcing through fifth scaled order; two further clock terms M, N ; the clock constant	Part II, Lemma 14.1, Section 15, (II.3)
$\kappa = \frac{\pi^3}{96} + \frac{\pi^2}{12} - \frac{13\pi}{48} - \frac{11}{36} - \frac{G}{6} - \frac{\pi \log 2}{24}$ (G Catalan)	
$a_n = c(n/T)^n n^{-4/3} \{1 + (-\frac{1}{18} \log^2 n + A \log n + C)/n + O(n^{-7/5})\}$, A, C exact in $h(1), h'(1), h''(1)$, κ	Part II, Theorem 12.1
The uniform first correction $\mathcal{Q}_\beta(d_m)/m$ for β in a compact subset of $(0, \infty)$	Part II, Theorem 12.2
Finite Fatou expansions to every order with rational a_j and differentiated sectorial remainders	Part II, Lemma 17.1 (to w^2); Part III, Lemma 26.1 (every order)
$h(0) = 2$; exact Hankel representation of h near zero; $t^{t/3} h(t)$ has a one-sided C^∞ germ; power-log expansion of h at zero (h is not C^1 at 0); $\log(h/2)$ coefficients	Part III, Proposition 27.1, Theorems 28.1 and 29.1, Corollary 29.2, Proposition 30.1
$c_j = \rho_j - \zeta(j)/(j^{3j})$, $\rho_j \in \mathbb{Q}$	
Transfer $m A(n, m) r_m^n = h(\beta) + O(\log m/(m\beta^{30}))$ for all n ; relative equivalent on $(\log m)^{-D} \leq \beta \leq B_0$	Part III, Theorem 31.1
$A(n, m)/(2T^{-n} m^{n-1}) \rightarrow e^{-1/(3\lambda)}$ for $m/(n \log n) \rightarrow \lambda$, every fixed order in $1/(\lambda \log n)$, $c_2 = (9 - \pi^2)/108$; Kaneiwa's normalization shifts K to $K+1$	Part III, Theorem 24.1, Section 33

Statement	Where
Smooth suppression-ratio inverses, rounded height prescriptions, observed-ratio inverses	Part III, Proposition 34.1, Theorems 34.2, 34.3

Open in all three Parts

Validated digits of c , $h(1)$, $h'(1)$, $h''(1)$; corrections beyond the first and an all-orders theorem; effective constants, onsets and certified thresholds; the regime $n/m \rightarrow \infty$ and the large-argument behaviour of h ; real analyticity of h on $(0, \infty)$ and analyticity of the germ at zero; polynomially small ratios and the transition to Kaneiwa's fixed-degree regime; other arrays and initial functions; the conjectured identity $I(\beta) = \beta 2^{\beta/3} h(\beta)$ with the iterated Bell amplitude. See Section 37.

The finite computations of the three companions (exact integer recurrences, exact rational formal coordinates, symbolic clock identities) check identities, normalizations and coefficient algebra; no asymptotic statement rests on them, and every decimal value of c , $h(1)$, $h'(1)$, $h''(1)$ in the report is an uncertified floating diagnostic.

The OEIS statements, and the inverses

The entries as read by the write on 5 October 2026 (A290354 revision #26 of June 21, 2018; A290353 revision #28 of September 3, 2026):

- **A290354** (Alois P. Heinz, July 28, 2017): “a(n) is the n-th term of the n-th Euler transform of the sequence with g.f. $1+x$ ”, also “the number of unlabeled rooted trees with exactly n leaves, all in level n”; formula “a(n) = A290353(n,n)” and the formula line, verbatim,

Conjecture: $a(n) \sim c * 2^n * n^{(n-4/3)} / \pi^n$, where $c = 4.4923\dots$ - _Vaclav Kotesovec_, Aug 14 2017

Since $2^n n^{n-4/3} / \pi^n = (n/T)^n n^{-4/3}$, Theorem 1.1 proves this form, with $c = T^{1/3} e^{-B} h(1) > 0$ identified analytically. It does not certify the digits: the value $c \approx 4.4923299$ (Part I, Section 10; Part II's diagnostics give the same) is a floating-point evaluation without error control, and agrees with the OEIS's 4.4923... only numerically, as Part I says. The entry's b-file covers $n = 0..414$; the manuscripts compare only the 22 displayed terms ($n = 0..21$) and an independently generated fixture through 414, as they state. The entry cross-references A139383, the diagonal of the labelled iterated Bell numbers (see “Relation to neighbouring reports”).

- **A290353** (Alois P. Heinz, July 28, 2017): “Square array A(n,k), $n \geq 0$, $k \geq 0$, read by antidiagonals, where column k is the k-th Euler transform of the sequence with g.f. $1+x$ ”; “A(n,k) is the number of unlabeled rooted trees with exactly n leaves, all in level k”; generating function of column $k > 0$: $\prod_{j>0} 1/(1-x^j)^{A(j,k-1)}$. Its k is the manuscripts' height m (Part II writes $b_{n,m}$). The entry's cross-references name rows 2, 3, 4 as A001477, A000217, A000330, that is $A(2,m) = m$, $A(3,m) = m(m+1)/2$, $A(4,m) = m(m+1)(2m+1)/6$; these are the coefficients $[q^2]F_h$, $[q^3]F_h$, $[q^4]F_h$ computed in Part II's Lemma 14.1 (checked by the write against the exact recurrence for $m \leq 12$). The entry carries no asymptotic statement.

The inverses. Part I's Corollary 9.1 and Part II's Corollary 21.1 start from $X \log(X/T) = Y$. This is the factorial core **p0:prop:factorial-core** of the transseries volume [11] with $\kappa = 1$, its d equal to $-\log T$ and $L = Y$: its solution $\exp(W_0(Y/T) + \log T) = T e^{W_0(Y/T)}$ equals $Y/W_0(Y/T) = X$; after taking logarithms it is the Lambert core **p0:thm:lambert-core** with $a = b = 1$ and $L = \log(Y/T)$ in the variable $\log X - \log T$ (L named after the independent check of 6 October 2026). The

corrections $X + \delta_0$ (Part I) and $X + \delta_0 + \delta_1/X$ (Part II) are proved directly, by Taylor expansion of an explicit smooth model and the mean-value theorem at the true integer; they parallel the first terms of `p0:thm:perturbed-inversion` but are not instances of it, because the perturbation $-\frac{4}{3}\log x + \log c + Q(\log x)/x$ is not a small analytic perturbation in that theorem's sense. The nearest-integer recovery at exact values $y = a_n$ resembles part (3) of `p0:thm:staircase`, but is not an instance either: its smooth root $x(y)$ solves a model equation and is not an admissible interpolation of (a_n) , and the manuscripts state no threshold for a general y between two terms. Part III's inverses (Section 34) involve no Lambert W : the model $\tau = \chi_n\beta - C(\beta)$ has the linear core $\chi_n\beta = \tau$ (the case $b = 0$ of `p0:thm:lambert-core`), perturbed by a germ C known to be C^∞ but not known to be analytic, so its finite reversion (Theorem 34.2) is proved by Taylor's theorem and the mean-value theorem, again an analogue rather than an instance of `p0:thm:perturbed-inversion`; its rounded height prescriptions are not instances of `p0:thm:staircase` (no monotonicity of $R(n, m)$ in m is available). None of the manuscripts claims novelty for the inversion mechanics.

Notation across the three Parts

Common to all Parts: F_h (the height index is h in Parts I–II and m in Part III's recurrence), $f(w) = e^w - 1$, $T = \pi/2$, B , Φ , P , the density h and its measure ν , $\beta = n/m$, the radius r , and the contour parameter s . Part II's $b_{n,m}$ is Parts I and III's $A(n, m)$. Part II's $d_\star = \frac{1}{3}\log T - B$, so its $D_n = n - \frac{1}{3}\log n + d_\star$ is Part I's D_n , and Part III's d_m is Part II's d_m . The symbols below change meaning between the Parts or inside one; each Part's text fixes its meaning there, and each Part opens with a short table of its own readings. No symbol was renamed.

Symbol	Meanings
h	The density of ν (all Parts) and the iteration height in F_h , R_h , E_h (Parts I, II; Part II's $t = hz$). <i>Tempting false reading</i> : in $F_h(z)$ and $[q^2]F_h = h$ the letter is the height, not the density.
K	Parts I, II: an integer entrance index ($M_0 \leq K \leq n$ in Lemma 3.3; $K = \lfloor \sqrt{n} \rfloor$ in Proposition 5.1 and (II.38); $K = \lceil At \rceil$ in Proposition 5.2); Part II also K_0 . Part III: the constant $K = \frac{1-\gamma+\log(T/2)}{3} - B$ (III.2), shifted to $K + 1$ by Kaneiwa's normalization; R_K is Kaneiwa's ratio. <i>False reading</i> : Part III's K is not an index.
κ	Parts II, III: the clock constant (II.3), $\kappa \approx -0.2543$. Part I: a small fixed $\kappa > 0$ bounding $K \leq \kappa n$ (before (23)).
ρ_j	Part II: $\rho_1 = h'(1)/h(1)$, $\rho_2 = h''(1)/h(1)$ and $\rho_j(\beta) = h^{(j)}(\beta)/h(\beta)$ (II.6). Part III: the rational numbers ρ_j , $j \geq 2$, of (III.6). <i>False reading</i> : Part III's $\rho_2 = 1/12$ is not Part II's $\rho_2 \approx -0.153$.
$c, C, \mathcal{C}$	$c = T^{1/3}e^{-B}h(1)$ the amplitude (Parts I, II). Part I: $C_z(w)$ the clock (10), C_λ (32), C_0 a fixed bound (Lemma 2.1). Part II: C the constant term of the correction (II.8), $C_z(w)$ the extended clock (II.32), C_0 a real-width bound, C_{30} . Part III: $C_0 = 1 - \gamma - \log 2$, $\mathcal{C}$ Catalan's constant (Part II's G), $C(\beta)$ the smooth germ (III.43); c_j the log-density coefficients (III.6). All Parts: c_k the coefficients in the exact recurrence; C , c generic constants. The iterated Bell report's $C \approx 2.8654$ is a different amplitude.
$G, \mathcal{G}, g$	Part I: $G(u) = 2/\log(1 + 2/u)$ (Lemma 4.1), Part III's $\mathcal{G}(u)$; $G_n(s) = F_n(T/(D_n - s))$ (24); $g(w) = \log(1 + w)$ (also Part II). Part II: G Catalan's constant; $G_n(s) = F_n(re^{s/n})$ (exponential, not Part I's rational parametrization); $G_3(y)$. Part III: $G_m(s) = F_m(r_me^{s/m})$; $g(t) = t^{t/3}h(t)$ the germ.
L	Parts I, II: the clock function $L(y)$ (9), and a fixed integer L in Propositions 5.1 and 18.1; Part I also $L(x) = x \log(x/T)$ in the proof of Corollary 9.1. Part III: $L = \log n$.
W	The positive Lambert function (Parts I, II). Part III: $W(x) = P(-x)$ (proof of Theorem 28.1); Part III uses no Lambert function.

Symbol	Meanings
λ	Part I: $m = \lfloor \lambda n \rfloor$ (Corollary 8.2), so $\lambda \approx m/n$. Part III: $m/(n \log n) \rightarrow \lambda$. <i>False reading:</i> Part III's λ is not a slope m/n . The iterated Bell report's λ is m/n .
α, β	$\beta = n/m$ in all Parts; Part I also uses $0 < \alpha < \beta < \pi$ for angles (proof of Lemma 4.1), β_0, β_1 interval ends, $\alpha_n = \{\lambda n\}$. Part II: $\alpha = 4/3$ (Section 21), $\alpha = n - H(z) - \kappa z$ (proof of Proposition 18.1). Part III: B_0 the upper end.
H	Parts I, II: the matching function $H(z) = T/z + \frac{1}{3} \log(1/z) + B$. Part III: $H = H_{\theta,a}$ a Hankel contour, $H_j(X)$ the log-polynomials of (III.27). The iterated Bell report's $H(n, m)$ is its array.
a, b	a_n the diagonal (all). Part II: $a(y) = (1 + y^2)/2$ (Part I's $A(y)$), $b(y, t)$, u, v step coefficients, $b_{n,m}$ the array. Parts I–III: $b_j = \sum_{d j} d[z^d]F_h$. Part III: a_j the rational Fatou coefficients (Part I's Section 10 lists the first three), b_j the germ's Taylor coefficients. Part I: b a small constant (Proposition 5.2).
A	$A(n, m)$ the array (Parts I, III). Part II: A the log n coefficient (II.8). Part I: $A(y)$, and A a large constant (Proposition 5.2). Part III: $\mathcal{A}(w)$ (III.20).
d, D	Part I: $d = n - h$ (Lemma 3.2), D_n . Part II: $d = -\frac{1}{3} \log n + d_*$, $d_m, d_y = 1 + y^2$, $D_n = n + d$. Part III: $d_m, D_N(u), D_t(v), D$ the band exponent (III.33).
R, r	Parts I, II: $R_h(z) = \sum_{j \geq 2} F_h(z^j)/j$; Part II also $R_n(s)$. Part III: the ratio $R(n, m)$ (III.3), R_2, R_K , the interval $\mathcal{R}$, and a radius R . r, r_m, r_n radii (all); Part III also a target ratio r .
Q	Part I: $Q(u) = u + \frac{1}{3} \log u$ (proof of Lemma 4.1), $Q_m(t)$ (proof of Theorem 8.1). Part II: $Q_2, Q_3, Q_\beta(d), Q_n(s), Q(u) = -u^2/18 + Au + C$. Part III: $Q_m(s)$, Part II's Q_n at height m .
M, N	Part II: the clock functions $M(y), N(y)$; also $M = n^{1/20}, M_n, M_0$. Part I: constants M, M_0 . Part III: $M = m^{1/20}$; N a truncation order.
q	Part I: the entrance constant $q, q = \log(X/T) + 1, q_m$. Part II: the derivative order q ($= 30$), q_X . Part III: a truncation degree q .
E	Parts I, II: E_h the clock error. Part III: $E(m, \beta)$ the transfer error (III.32).
δ, τ	Part I: δ the tail range, τ a first crossing time (Lemma 3.2). Part II: δ_0, δ_1 inverse coefficients. Part III: $\tau = -\log r$.
X, y	Parts I, II: $X = Y/W(Y/T)$; $y = w/z$ the clock ratio; $y = a_n$ in the inverses. Part III: $X = \log t$; $y = m/n, y_*, y_J$ height variables.
ℓ	Part II: $\ell = \log c$. Part III: $\ell_n = E_{n-1}/(n-1)!$ Kaneiwa's coefficient.
I, Ψ	Part II: I a compact interval of ratios. Part I: Ψ the reciprocal Abel coordinate; Part III also $\Psi_{\text{IBD}} = -\Phi + \frac{1}{3} \log 2$, the iterated Bell report's coordinate, and $I(\beta)$ that report's amplitude (Section 25).

Provenance and merge decisions

Three manuscripts of the session bundle of Reports 1–243. Their author lines read “Report 225”, “Report 226” and “Report 228” (PDF author field “Research report”); none names a person, a tool or an addressee, says it is AI-assisted, or carries “prepared for private review” wording. All three are dated 5 October 2026. Report 225 cites the iterated Bell report at commit 65969409c and Report 228 at commit 5fdc0d68a; both commits exist in the repository, and that report's `article.tex` is unchanged at HEAD (blob ab5014ec), so their section numbers refer to its current text. Report 226 pins no commit. They arrived in commit 60f54ea06 and were placed by 47fc7a069 (batch 106).

- **Part I**, bundle Report 225, archive `Report225.zip` (`src/report225.tex`, 639 lines, 16 pages); the base. Contributes the subsidiary-forcing estimate, the nonautonomous clock and its matching constant B , the noncircular real entrance, the sectorial Fatou coordinate and

entire parametrization, the local limit and angular tails, the positive measure and smooth strictly positive density, the diagonal equivalent, proportional height, fixed and square-root shifts, the floor phase, the Lambert inverse, and Kaneiwa’s and Bechtloff Weising’s antecedents.

- **Part II**, bundle Report 226, archive `Report226.zip` (`src/report226.tex`, 715 lines, 17 pages). Its `supporting/` holds Report 225’s source and PDF byte for byte, and eight of its code and data files are byte copies of Report 225’s. Contributes fifth-order forcing, the clock functions M and N , the closed form of κ , the weighted fourth-power defect, the finite Fatou expansion and forced matching, the growing-strip local limit, thirtieth-derivative extraction, the first correction on the diagonal and uniformly in compact ratios, and the sharpened inverse.
- **Part III**, bundle Report 228, archive `Report228.zip` (`src/report228.tex`, 729 lines, 17 pages). Its `supporting/` holds the sources and PDFs of Reports 225 and 226 byte for byte. Contributes the L^1 form of Part II’s estimates, finite Fatou expansions to every order, $h(0) = 2$, the Hankel representation, the endpoint germ and its rational-plus-zeta coefficients, the transfer to logarithmic bands, the crossover theorem, Kaneiwa’s normalization, the smooth inverses, and the local identification with the iterated Bell coordinate together with the conjectured amplitude identity.

Where the merge had to choose:

1. *Order and base*. Dependency order I–II–III, with the base first (see the Guide). Statements that a later Part extends stay as delivered, with dated notes.
2. *Restatements*. Part II’s Section 13 restates Part I’s Lemma 2.1, the clock and entrance (Sections 2–3, (23)), the coordinate (Lemma 4.1), the tail and outer bounds (Proposition 5.2, Lemma 5.3) and the density (Proposition 6.1), and re-derives Corollary 6.2. Part III’s Section 25 restates Part I’s coordinate and density and Part II’s absolute estimates, then derives a new L^1 bound. The intake dossier suggested replacing these sections by pointers, as the iterated Bell report did for its Part II; the write printed them in full, as the batch-105 model report did, because nearly every display in them is cited by the later proofs, because Part II’s states as items estimates that Part I proves only inside proofs, and because Part III’s contains material found nowhere else in the merge ((III.16) and the iterated Bell comparison). (Wording corrected after the independent check of 6 October 2026; the first wording read “and because each contains material found nowhere else in the merge”.) No passage of one manuscript repeats another verbatim.
3. *Front matter and bibliography*. The three title blocks and abstracts are printed at the heads of their Parts; one table of contents replaces three. The three bibliographies are merged into one, keeping every detail of every entry; where two manuscripts describe an entry differently both descriptions are kept. The citation keys were renamed (Report 228’s `oeis` is A290353, Reports 225 and 226’s `oeis` is A290354); Reports 226 and 228’s entries for Reports 225 and 226 now point to Parts I and II. The write added the entry for the transseries volume [11].
4. *Part III’s table*. Report 228’s `\input{crossover_table.tex}` is replaced by the table’s text, byte for byte apart from its two leading comment lines, which are kept as $\text{\LaTeX}$ comments; the generated file is shipped as `data/228-logheight-results-crossover_table.tex`.
5. *Build-specific lines*. The manuscripts’ `\pdfmapfile` lines (font-map additions for their TeX Live build) and Report 228’s table-of-contents format tweak are not carried into the merged preamble; they affect no text.
6. *Write additions*. This front matter, the reading-conventions tables of the Parts, Section 37 with Remarks 37.1 and 37.2, the dated notes, labels for unlabelled sections, and the label prefixes. Everything else is delivered text.

What was checked, and what was not

The report is unrefereed and not formalized. At intake (dossier of batch 106, 5 October 2026) the three manuscripts were read in full and no step was found false. Checked independently there: the first twelve terms of A290354 and the normalized ratios $a_n(T/n)^n n^{4/3} = 4.6013, 4.5529, 4.5178$ at $n = 40, 80, 200$ that Part I quotes, from an own implementation of the recurrence; the closed form of κ against a direct quadrature of the defining equation (II.23), -0.2543481698511 against -0.2543481698560 (difference from the truncated tail); that with Part II’s diagnostic constants the corrected prediction tracks the exact a_n far better than the leading term (residual 1.4×10^{-4} against 2.5×10^{-2} at $n = 200$); and $2^{1/3}h(1)$ against the iterated Bell amplitude at $\beta = 1$ (Remark 37.1). The delivered suites pass on copies (README). The write checked the rows $A(2, m)$, $A(3, m)$, $A(4, m)$ against Part II’s Lemma 14.1, the reversion coefficients (III.46)–(III.47), c_2 and c_3 from $S(t)$ and the Gamma expansion, the linear coefficient $C_0/3$ and K in (III.36), the reduction of Theorem 12.2 to Theorem 12.1 at $\beta = 1$, the floor amplitude C_λ of (32), the two orders of the inverse (II.53), and recomputed $c = 4.4923298968$ from the shipped diagnostic value of $h(1)$ (added after the independent check of 6 October 2026: an independent evaluation of $h(1)$ gives $c = 4.4923298971$; Remark 37.1).

What was *not* done: the analytic estimates (the clocks and their defects, the sector bootstraps, the tails, the Hankel deformations) were read and their displayed algebra spot-checked, but they are the manuscripts’ proofs, corroborated numerically, not an independent verification. The write did not read Kaneiwa (1980), Bechtloff Weising, Dudko–Sauzin, Prellberg–Mishna or Nagaev–Vakhtel; those citations and their page and theorem numbers are as the manuscripts and their source ledgers give them.

Relation to neighbouring reports and formal projects

The labelled analogue. SetTheory/Cardinals/docs/reports/generating-functions-and-asymptotics/oeis-sequence-asymptotics/a139383-iterated-bell-diagonals (labels ibd:) treats $H(n, m) = n![z^n](e^z - 1)^{om}$, OEIS A139383 and A261280: a fixed-shift diagonal theorem to every order (its Part I) and proportional depth $m = \lambda n + \delta$ for λ in compact subsets of $(0, \infty)$ (its Part II), by the same parabolic map $w \mapsto e^w - 1$ without subsidiary forcing. No theorem is shared: the arrays differ, and Part I here says the labelled theorem “cannot be directly transferred” (Section 1.1), because the forcing changes the leading scale ($(n/T)^n n^{-4/3}$ here, $n^{2n-5/6}/(2^{n-1}e^n)$ there) and the clock constant. The two reports share the normalized orbit: Part III records $\Psi_{\text{IBD}}(w) = -\Phi(w) + \frac{1}{3} \log 2$ and the unproved relation $I(\beta) = \beta 2^{\beta/3} h(\beta)$ between that report’s amplitude I and the density h here (Section 25). The values agree to eight significant digits at $\beta = 1$ (Remark 37.1), and the relation, if proved, would bear on that report’s open endpoint $\lambda \rightarrow \infty$ (Section 37, item 9, and Remark 37.2). That report answers no question asked here, and this one answers none asked there; a reciprocal note is proposed for it.

Other neighbours. No other report of the collection treats A290353 or A290354 (repository search, 6 October 2026; A290354 occurs elsewhere only in a cross-reference line of the iterated Bell report’s OEIS data). The batch-106 reports a252782-diagonal-euler-transforms (one Euler transform with weights j^n) and a005121-strict-partition-chains (the depth sum $\sum_m 2^{-m-1} H(n, m)$ of the labelled array) share only words with this one. The transseries volume supplies the factorial core of which the leading inverses are instances (“The OEIS statements, and the inverses”). Placement in the collection confers no formal status: no Lean or Rocq file in the

repository treats these sequences, and no statement of this report is formalized.

Part I

Diagonal and proportional height asymptotics for iterated Euler transforms

Bundle Report 225, archive `Report225.zip`, delivered as `src/report225.tex` (16 pages), the base of the merge. Delivered title block “Diagonal and proportional height asymptotics for iterated Euler transforms / Report 225 / 5 October 2026”. Printed as delivered, with `[write]` notes; section, statement and equation numbers are those of the delivery.

Reading conventions of Part I	Elsewhere in the report
$A(n, m) = [z^n]F_m$; $a_n = A(n, n)$	Part II: $b_{n,m}$ for $A(n, m)$
K an integer entrance index; κ a small constant in $K \leq \kappa n$	Part III: K a constant; Parts II, III: κ the clock constant
$L(y)$ the clock function; $C_z(w)$ the clock; E_h	Part II: $C_z(w)$; Part III: $L = \log n$, $E(m, \beta)$
$G(u)$, $g(w) = \log(1 + w)$,	Part II: G Catalan, $G_n(s) = F_n(re^{s/n})$; Part III:
$G_n(s) = F_n(T/(D_n - s))$	$\mathcal{G}(u)$, $g(t) = t^{t/3}h(t)$
$m = \lfloor \lambda n \rfloor$ (Corollary 8.2)	Part III: $m/(n \log n) \rightarrow \lambda$

Abstract of this Part (as delivered). We study the number a_n of unlabeled rooted trees with n leaves, all at level n , equivalently the diagonal of the iterated Euler transform in OEIS A290354. We prove

$$a_n \sim c \left(\frac{2n}{\pi} \right)^n n^{-4/3}, \quad c > 0,$$

and identify c by a normalized repelling parametrization of $w \mapsto e^w - 1$. The main estimate is a uniform intermediate-angle bound, obtained from an explicit nonautonomous clock, a positive-coefficient Cauchy entrance, and a forced sector bootstrap. A positive Laplace measure identifies the coefficient amplitude and proves its strict positivity without assuming a density expansion at zero. Uniform proportional-height asymptotics for the array A290353, fixed and square-root height shifts, and a Lambert W inverse corollary are included. Numerical diagnostics give $c \approx 4.4923$, agreeing with Kotesovec’s conjecture; the decimal computation is not a validated enclosure. This is an unrefereed mathematical research report, not a proof-assistant formalization.

1 Definition and statement

Let

$$F_0(z) = z, \quad F_{h+1}(z) = \exp \left(\sum_{j \geq 1} \frac{F_h(z^j)}{j} \right) - 1. \quad (1)$$

Each F_h is analytic in $|z| < 1$, has nonnegative coefficients, vanishes at zero, and has linear coefficient one. These facts follow by induction; the sum over j converges normally on compact subdisks because $F_h(z) = O(z)$ at zero. For $n \geq 1$ put $a_n = [z^n]F_n(z)$ and set $a_0 = 1$ separately. The constant-term exception matches the OEIS convention and has no asymptotic effect.

For clarity, write $A(n, m) = [z^n]F_m(z)$ for $n \geq 1$, and put $A(0, m) = 1$ by convention. This is the square array A290353 [1], with degree n and height m ; the sequence A290354 is its main diagonal. A tree whose leaves all have level $m + 1$ is a root carrying a nonempty unordered multiset

of trees whose leaves all have level m . If there are $[z^d]F_m(z)$ types of a subtree with d leaves, the ordinary multiset generating function is

$$1 + F_{m+1}(z) = \prod_{d \geq 1} (1 - z^d)^{-[z^d]F_m(z)}. \quad (2)$$

Taking its formal logarithm gives (1). This also proves integrality and explains why there is no factorial normalization. The first diagonal values are

$$1, 1, 2, 6, 30, 170, 1337, 12166, 133476, 1676364.$$

The separate constant row is a generating-function convention, rather than a claim about a nonempty tree with no leaves.

Set

$$T = \frac{\pi}{2}, \quad B = \frac{1}{2} - \frac{\pi}{6} - \frac{\pi^2}{16} - \frac{\log 2}{6}, \quad D_n = n - \frac{1}{3} \log n + \frac{1}{3} \log T - B. \quad (3)$$

The logarithms used near the positive axis have their real determination there.

Theorem 1.1. *There is an entire function P satisfying*

$$P(s+1) = e^{P(s)} - 1, \quad (4)$$

whose inverse coordinate near zero is normalized by

$$\Phi(w) = -\frac{2}{w} + \frac{1}{3} \log w + O(w), \quad \Phi(e^w - 1) = \Phi(w) + 1.$$

It is the Laplace transform of a finite positive measure ν on $[0, \infty)$ with every exponential moment. This measure has a smooth, strictly positive density h on $(0, \infty)$. Moreover,

$$a_n \sim c(n/T)^n n^{-4/3}, \quad c = T^{1/3} e^{-B} h(1) > 0. \quad (5)$$

For every real σ ,

$$h(1) = \frac{e^{-\sigma}}{2\pi} \int_{-\infty}^{\infty} P'(\sigma + it) e^{-it} dt, \quad (6)$$

and this integral is absolutely convergent.

The normalization of Φ is essential: translating a Fatou coordinate translates P and changes the density amplitude. Formula (3) is tied to the zero constant term in the displayed normalization.

1.1 Prior work and the scope of the result

The OEIS entry attributes (5), with $c = 4.4923\dots$, to Václav Kotěšovec on 14 August 2017 and labels it a conjecture [2]. Kaneiwa's 1980 paper [3], Propositions 1 and 2 on printed page 19, proves that the unlabeled r -fold partition number $p(r; n)$ is a polynomial of degree $n - 1$ in r and identifies its leading coefficient through $\tan x + \sec x$. His indexing gives

$$p(r; n) = [z^n]F_{r+1}(z), \quad a_n = p(n - 1; n) \quad (n \geq 1).$$

Thus polynomiality, the tangent leading profile, and its $\pi/2$ pole are prior results. Fixed- n polynomial information does not alone justify the growing- n diagonal limit. Kaneiwa credits Bell's

1938 polynomiality result for the separately defined *labeled* function $\tilde{p}(r; n)$ on page 18; that sentence should not be read as attributing his unlabeled Proposition 1 to Bell.

Bechtloff Weising's higher-order Bell symmetric functions [4], Definition 2.1 and Proposition 2.12, specialize in one variable to $1 + F_{m+1}$; in particular $a_n = [m_{(n)}]B_n^{(n-1)}$. This is an algebraic and species connection, not a diagonal asymptotic theorem in that source. The related iterated-Bell report [5] concerns $n![z^n](e^z - 1)^{om}$ and cannot be directly transferred: the subsidiary forcing in (1) changes both the leading scale and the clock constant. No general priority claim for parabolic-coordinate methods is made here.

[write] **The OEIS statement and the neighbouring report, 6 October 2026.** The front matter quotes A290354's formula line verbatim as read on 5 October 2026: "Conjecture: $a(n) \sim c * 2^n * n^{(n-4/3)} / \Pi^n$, where $c = 4.4923\dots$ - _Vaclav Kotesovec_, Aug 14 2017". Theorem 1.1 proves this form with c identified; the digits 4.4923 are matched only by an uncertified evaluation (Subsection 10.2). Kaneiwa's indexing $p(r; n) = [z^n]F_{r+1}$ is used again in Part III, Section 33, which compares his leading coefficient with the crossover. The iterated Bell report [5] is the labelled analogue; Part III, Section 25, records the common normalized orbit and a conjectured identity between its amplitude and the density h here (Section 37, item 9).

2 Subsidiary terms and the nonautonomous clock

Write

$$R_h(z) = \sum_{j \geq 2} \frac{F_h(z^j)}{j}, \quad f(w) = e^w - 1.$$

The subsidiary estimate below is independent of the size of the main value $F_h(z)$.

Lemma 2.1. *Uniformly for $h|q|$ sufficiently small and $|q|$ small,*

$$F_h(q) = q + hq^2 + O((h + h^2)|q|^3). \quad (7)$$

Consequently, for each fixed C_0 , uniformly for $h \leq C_0/|z|$ as $z \rightarrow 0$,

$$R_h(z) = \frac{z^2}{2} + \frac{z^3}{3} + \frac{hz^4}{2} + O(|z|^4). \quad (8)$$

Proof. For real $0 < q < 1$, coefficient positivity and the absence of a constant term give $F_h(q^j) \leq q^{j-1}F_h(q)$. Hence the exponent defining $F_{h+1}(q)$ is at most $F_h(q)[- \log(1 - q)]/q$. While $F_h(q) \leq 2q$, the one-step increment is at most $CF_h(q)^2$. The elementary reciprocal comparison, or induction against $q/(1 - C'hq)$ with C' fixed large, proves $F_h(q) \leq 2q$ for $hq \leq c$ and c sufficiently small. Positivity gives the corresponding complex bound.

Summing one-step $O(|q|^2)$ increments gives $F_h(q) = q + O(h|q|^2)$. Substituting this back into the recurrence gives the sharper increment $F_{h+1}(q) - F_h(q) = q^2 + O((h + 1)|q|^3)$, because the quadratic main term and the leading subsidiary term each contribute $q^2/2$. Summation proves (7). Apply it at $q = z^j$ for $j \geq 2$. The $j = 2$ error is $O(h^2|z|^6) = O(|z|^4)$; the baseline tail $j \geq 4$ and the remaining corrections are bounded by geometric series. This proves (8). $\square$

[write] **Extended in Part II, 6 October 2026.** Part II's Lemma 14.1 extends (8) to fifth scaled order, from the exact coefficients $[q^2]F_h = h$, $[q^3]F_h = h(h + 1)/2$, $[q^4]F_h = h(h + 1)(2h + 1)/6$, which are the rows $n = 2, 3, 4$ of A290353 (checked by the write against the exact recurrence for $h \leq 12$).

Define, in a narrow sector about the positive real axis,

$$J(y) = 2 \arctan y - \frac{\pi}{2}, \quad L(1) = 0, \quad L'(y) = \frac{y^3/3 - y - 4/3 - 2J(y)}{(1+y^2)^2}. \quad (9)$$

Put

$$C_z(w) = \frac{J(w/z)}{z} + L(w/z). \quad (10)$$

The clock domain will be $|\arg(w/z)| \leq \eta$, $|w/z| \geq 1/2$, with a slightly wider fixed sector available for Taylor segments. In particular $C_z(z) = 0$.

Lemma 2.2. *Suppose $h \leq C_0/|z|$, $w_h = F_h(z)$ remains sufficiently small, and w_h/z remains in the stated clock domain through time k . Then, with $E_h = C_z(w_h) - h$,*

$$E_{h+1} - E_h = \frac{z^2(L(w_h/z) - E_h)}{1 + (w_h/z)^2} + O(|w_h|^2 + |z|^2), \quad (11)$$

with uniform constants. Consequently

$$|E_k| \leq C \left(\sum_{h < k} |F_h(z)|^2 + k|z|^2 \right). \quad (12)$$

Proof. By Lemma 2.1,

$$w_{h+1} = w_h + \frac{w_h^2 + z^2}{2} + \frac{w_h^3}{6} + \frac{w_h z^2}{2} + \frac{z^3}{3} + \frac{h z^4}{2} + O((|w_h| + |z|)^4). \quad (13)$$

Write $y = w_h/z$, $A(y) = (1+y^2)/2$ and $B_1(y, hz) = y^3/6 + y/2 + 1/3 + hz/2$. The increment of y is $zA + z^2B_1 + O((|w_h| + |z|)^4/|z|)$. Since $J' = 1/A$, the coefficient of z in the clock increment is

$$\frac{B_1}{A} - \frac{y}{2} + AL'(y) = \frac{hz - J(y)}{1 + y^2}.$$

But $hz - J(y) = z(L(y) - E_h)$, proving the cancellation in (11).

For completeness the remainder is analytic and uniform. Derivatives of $J(w/z)/z$ of order $r \geq 1$ are $O((|w| + |z|)^{-r-1})$, and derivatives of $L(w/z)$ are $O((|w| + |z|)^{-r})$. The displacement in (13), divided by $|w|$, is $O(|w| + |z|)$ because $|w/z|$ is bounded below. Nested clock cones therefore keep the full Taylor segment away from $w = \pm iz$. The omitted recurrence terms, the mixed quadratic Taylor terms, and the cubic Taylor remainder all contribute $O((|w| + |z|)^2)$ to the clock increment.

Finally $L(y)/(1+y^2)$ is bounded on the clock cone and $|1+y^2|$ is bounded below. The multiplicative discrete Gronwall factor is at most $\exp(Ck|z|^2)$, uniformly bounded in the stated range. Since $E_0 = 0$, this gives (12). $\square$

3 Matching and the positive real orbit

Lemma 3.1. *Uniformly in a fixed narrow sector about the positive real axis,*

$$L(y) = \frac{1}{3} \log y + B + O(y^{-2}) \quad (|y| \rightarrow \infty). \quad (14)$$

Consequently, when $|w/z|$ is sufficiently large in that sector,

$$C_z(w) = H(z) - \frac{2}{w} + \frac{1}{3} \log w + O\left(\frac{|z|^2}{|w|^3} + \frac{|z|^2}{|w|^2}\right), \quad H(z) = \frac{T}{z} + \frac{1}{3} \log \frac{1}{z} + B. \quad (15)$$

For real $0 < x \leq w \leq \epsilon$, with ϵ small fixed,

$$H(x) - C_x(w) \asymp \frac{1}{w}, \quad C'_x(w) > 0. \quad (16)$$

Proof. The derivative in (9) is $1/(3y) + O(y^{-3})$; in particular there is no y^{-2} term. To compute its constant, integrate the part $(y^3/3 - y)/(1 + y^2)^2$ first. After subtraction of $(\log y)/3$ its contribution from 1 to infinity is $-(\log 2)/6 - 1/3$. In the rest set $u = \arctan y$. Its contribution is

$$\int_{\pi/4}^{\pi/2} (\pi - 4/3 - 4u) \cos^2 u \, du = \frac{5}{6} - \frac{\pi}{6} - \frac{\pi^2}{16}.$$

Their sum is B in (3). The arctangent expansion gives (15).

For real $w \geq x$, the arctangent identity gives

$$H(x) - C_x(w) = \frac{2}{x} \arctan \frac{x}{w} - \frac{1}{3} \log w + O((x/w)^2).$$

The first term is between fixed positive multiples of $1/w$; the remaining terms are smaller for small w , uniformly for $x/w \leq 1$. The expansion of L may be bounded on all $y \geq 1$ by enlarging its constant on a compact interval. Finally the main derivative $2/(x^2 + w^2)$ dominates the $O(1/(x + w))$ derivative of the correction for small ϵ . This proves (16). $\square$

Lemma 3.2. Fix $\sigma \in \mathbb{R}$ and let $r = T/(D_n - \sigma)$. There are constants $M_0, c_0, c, C > 0$ such that, for all sufficiently large n , if $d = n - h \geq M_0$ and $x > 0$ satisfies $|x - r| \leq c_0 d/n^2$, then every real iterate through h stays in the small-clock range and

$$\frac{c}{d} \leq F_h(x) \leq \frac{C}{d}. \quad (17)$$

Proof. We have $H(r) = n - \sigma + O(\log n/n)$ and $H'(x) = O(n^2)$ when $x \asymp n^{-1}$. Make c_0 small, then M_0 large compared with $|\sigma|$. Uniformly in the displayed range,

$$d/2 \leq H(x) - h \leq 2d. \quad (18)$$

It remains to justify that the small-clock hypotheses are not circular. Let τ be the first time $F_\tau(x) \geq \epsilon$, if it occurs by time h . Start with a slightly larger admissible clock radius, so that one-step overshoot gives $F_\tau(x) \leq 2\epsilon$. Positivity gives

$$F_{j+1}(x) - F_j(x) \geq \frac{1}{2} F_j(x)^2, \quad \sum_{j < \tau} F_j(x)^2 \leq 2(F_\tau(x) - x).$$

Lemma 2.2 therefore gives $E_\tau = O(\epsilon + \tau x^2)$; here $\tau \leq h \leq n$ and $x \asymp n^{-1}$. Lemma 3.1 and monotonicity of C_x imply $\tau \geq H(x) - C/\epsilon$. If M_0 is large enough, this contradicts (18). Thus no such first crossing occurs. Apply (12) again through h and use (16); the clock error is $O(F_h(x) + nx^2)$ and can be absorbed into the $1/F_h(x)$ comparison. Equation (18) now gives (17). $\square$

Lemma 3.3. *With r as in Lemma 3.2, there is $q > 0$ such that for every integer $M_0 \leq K \leq n$ and every $|z - r| \leq qK/n^2$, the orbit through $n - K$ lies in the clock cone, satisfies $|F_h(z)| \asymp (n - h)^{-1}$, and obeys*

$$C_z(F_{n-K}(z)) = n - K + O(K^{-1} + n^{-1}). \quad (19)$$

Proof. At height $h \leq n - K$, use the disk about r of radius $R_h = c_0(n - h)/n^2$. Positivity and Lemma 3.2 bound its supremum by $F_h(r + R_h) \leq C/(n - h)$. On its inner half, Cauchy's estimate gives $|F'_h| \leq Cn^2/(n - h)^2$. Hence

$$|F_h(z) - F_h(r)| \leq C|z - r| \frac{n^2}{(n - h)^2}.$$

Divide by the positive lower bound in (17). Taking $q \leq c_0/2$ and then decreasing it makes the relative error at most $CqK/(n - h) \leq Cq$. Also $|z - r|/r \leq CqK/n$. Thus $F_h(z)/z$ lies in the required fixed narrow cone, with modulus at least $1/2$, for every $h \leq n - K$. The two-sided comparison follows. Finally $\sum_{h < n-K} |F_h(z)|^2 \leq C/K$, so Lemma 2.2 gives (19). $\square$

4 The repelling coordinate with uniform sectors

The needed coordinate is constructed here to make the sector and branch choices explicit.

Lemma 4.1. *For each closed angular sector strictly contained in $|\arg w| < \pi$, there is a sufficiently small radius on which an analytic function Φ satisfies*

$$\Phi(w) = -2/w + \frac{1}{3} \log w + O(w), \quad \Phi'(w) = 2/w^2 + O(1/w), \quad \Phi(f(w)) = \Phi(w) + 1, \quad (20)$$

whenever both arguments lie in the chosen domain. These functions agree on common smaller sectors. There is a local inverse on every sufficiently distant closed subsector of $|\arg(-s)| < \pi$, with

$$P_{\text{loc}}(s) = -2/s + O(\log |s|/|s|^2), \quad P'_{\text{loc}}(s) = 2/s^2 + O(\log |s|/|s|^3). \quad (21)$$

For small w in a smaller sector, $P_{\text{loc}}(\Phi(w)) = w$. The local inverse extends to the entire P in Theorem 1.1 and agrees with it on its sector of definition.

Proof. Let $g(w) = \log(1 + w)$ and conjugate by $u = 2/w$. The resulting map satisfies

$$G(u) = \frac{2}{\log(1 + 2/u)} = u + 1 - \frac{1}{3u} + O(u^{-2}).$$

For $Q(u) = u + (\log u)/3$,

$$Q(G(u)) - Q(u) = 1 + O(u^{-2}). \quad (22)$$

We give the uniform geometric estimate behind summation of this defect. Fix $0 < \alpha < \beta < \pi$. If $|\arg u| \leq \alpha$, then

$$|u + j| \geq \cos(\alpha/2)(|u| + j), \quad j \geq 0,$$

and the straight ray has distance at least $c_{\alpha,\beta}(|u| + j)$ from the complement of $|\arg v| < \beta$. Distinguish an inner analyticity radius R_0 from a larger starting threshold R_1 . Bootstrap

$$|G^j(u) - (u + j)| \leq a(|u| + j),$$

where a is less than half the preceding radial and angular margins. This gives $|G^j(u)| \geq c(|u|+j) > R_0$ if R_1 is large enough. Summing the one-step $O(1/|G^j(u)|)$ error improves the bound to

$$|G^j(u) - (u + j)| \leq C \left(\frac{1}{|u|} + \log(1 + j/|u|) \right).$$

Divided by $|u| + j$, this is uniformly $O(R_1^{-1})$ because $\log(1+x)/(1+x)$ is bounded. Increase R_1 to obtain a strict improvement and $cR_1 > 2R_0$. This closes the induction, including times when the straight ray decreases in modulus. Only the angular sector is invariant; the starting-radius boundary need not be.

The defects in (22) are now absolutely summable with total $O(1/|u|)$. Therefore

$$\Psi(u) = \lim_{m \rightarrow \infty} (Q(G^m(u)) - m) = Q(u) + O(1/|u|)$$

exists locally uniformly, is analytic, and satisfies $\Psi(G(u)) = \Psi(u) + 1$ wherever both sides are defined. Nested sectors and thresholds justify shifting the limit. Cauchy estimates on smaller sectors give derivative bounds. Define

$$\Phi(w) = -\Psi(2/w) + \frac{1}{3} \log 2.$$

This proves (20); $g \circ f$ is the identity near zero.

For the inverse, solve $\Psi(u) = -s + (\log 2)/3$. A disk of radius $C \log |s|$ centered at the right side stays inside a slightly larger allowed sector for large $|s|$. On its boundary the logarithmic correction to the identity map is smaller than the radius for C large. Rouché's theorem gives exactly one root, and $\Psi'(u) = 1 + O(1/|u|)$ is nonzero. The roots depend analytically on s , agree on overlaps by uniqueness, and give (21). If a small w already satisfies $\Phi(w) = s$ in a smaller sector, then $u = 2/w = -s + O(\log |s|)$ lies in the same uniqueness disk. Thus $P_{\text{loc}}(\Phi(w)) = w$; global injectivity of Φ is neither assumed nor needed.

First restrict P_{loc} to a strict left half-plane $\text{Re } s < -R_2$ so that $f(P_{\text{loc}}(s)) = P_{\text{loc}}(s + 1)$ holds when both parameters lie there. On $\text{Re } s < m - R_2$, define $P(s) = f^{\circ m}(P_{\text{loc}}(s - m))$. Successive definitions agree on overlaps, so they produce an entire function obeying (4). Agreement with P_{loc} on every connected larger inverse sector follows from the identity theorem, since there is agreement on a nonempty open part of the left half-plane. Positivity first far to the left and then forward iteration show $P(s) > 0$ for real s . $\square$

[write] Finite expansions of Φ elsewhere in the report, 6 October 2026. Part II's Lemma 17.1 proves $\Phi(w) = -2/w + \frac{1}{3} \log w - w/36 + w^2/540 + O(w^3)$ with differentiated sectorial remainders, and Part III's Lemma 26.1 the expansion to every order with rational coefficients a_j , both by summing the Abel defect along the inverse orbits constructed in this proof. The coefficients displayed in Subsection 10.2 are those a_1, a_2, a_3 .

Combining Lemmas 3.1, 3.3, and 4.1 gives the crucial entrance phase. Restrict $M_0 \leq K \leq \kappa n$ with $\kappa > 0$ small fixed, so that $|F_{n-K}(z)/z| \asymp n/K$ lies in the large-ratio matching range. Uniformly on the entrance disk,

$$\Phi(F_{n-K}(z)) = n - K - H(z) + O(K^{-1} + n^{-1} + K^3/n^2). \quad (23)$$

Both matching scales below satisfy this restriction.

5 The local limit and angular tails

Proposition 5.1. *Locally uniformly for $s \in \mathbb{C}$,*

$$G_n(s) := F_n\left(\frac{T}{D_n - s}\right) \longrightarrow P(s). \quad (24)$$

Here G_n is defined on each fixed compact set for all sufficiently large n .

Proof. Fix a compact set S and take $K = \lfloor \sqrt{n} \rfloor$. The point $z = T/(D_n - s)$ differs from the real reference T/D_n by $O_S(n^{-2})$, so Lemma 3.3 applies. Exactly,

$$H\left(\frac{T}{D_n - s}\right) = n - s + \frac{1}{3} \log \frac{D_n - s}{n}.$$

Thus (23) gives entrance phase $s - K + o(1)$ uniformly on S .

Choose a fixed integer L larger than the inverse-sector threshold and $2 \max_{s \in S}(|s| + 1)$. For $0 \leq j \leq K - L$, the prototype $s - K + j$ lies in a fixed interior left sector, has modulus comparable to $K - j$, and has real part bounded away from zero. Bootstrap phase error at most one. By (8), the actual w -step differs from $f(w)$ by $O(n^{-2})$. The disk of this radius about the unforced image lies in the coordinate domain: the image has distance at least $c/(K - j)$ from its angular boundary and zero. On that disk $|\Phi'| \leq C(K - j)^2$. Lemma 4.1 identifies the perturbed point with the same inverse branch. Thus the phase error at step j is at most $C(K - j)^2/n^2$, and the total is $O(K^3/n^2) = o(1)$. Together with the initial uniform error this strictly improves the bootstrap for large n .

At time $n - L$ the value tends uniformly to $P(s - L)$. In the remaining fixed L steps the subsidiary forcing is still $O(n^{-2})$. Finite compositions of the entire function f preserve convergence on these compact images, proving (24). $\square$

Proposition 5.2. *Fix $\sigma \in \mathbb{R}$ and put $r = T/(D_n - \sigma)$. There are $M, \delta, C > 0$ such that, for all large n and $M \leq t \leq \delta n$,*

$$|F_n(re^{it/n})| \leq C/t, \quad \left| \frac{re^{it/n}}{n} F'_n(re^{it/n}) \right| \leq C/t^2. \quad (25)$$

The same holds for negative t with $|t|$ in place of t .

Proof. Choose A large and $K = \lceil At \rceil$. Let $z_0 = re^{it/n}$. Since $|z_0 - r| \leq Ct/n^2$, choosing A after the entrance constant q and then choosing $b > 0$ small places

$$|z - z_0| \leq bt/n^2 \quad \text{inside} \quad |z - r| \leq qK/n^2.$$

Choose δ small enough that $K/n \leq \kappa$ and $K \leq n/2$. Expansion of H , using $H'(z) = O(n^2)$ on this disk and $H(r) = n - \sigma + O(\log n/n)$, gives

$$s_0 := \Phi(F_{n-K}(z)) = -K + it + O(1 + bt + t^2/n + t \log n/n + K^3/n^2). \quad (26)$$

The bounded term includes fixed σ . First choose b and δ to control proportional errors, then M to control fixed constants and the local inverse radius, and finally n to control $\log n/n$. The error in (26) is at most $t/10$.

Use the fixed prototype $p_j = -K + j + it$, $0 \leq j \leq K$, and bootstrap $|s_j - p_j| \leq t/4$. Then

$$\operatorname{Im} s_j \geq 3t/4, \quad \operatorname{Re} s_j \leq t/4, \quad |\arg(-s_j)| \leq \pi/2 + \arctan(1/3) < 2\pi/3, \quad ct \leq |s_j| \leq CK.$$

After increasing M , the inverse estimate gives

$$|\arg P_{\text{loc}}(s_j)| \leq 2\pi/3, \quad c/K \leq |P_{\text{loc}}(s_j)| \leq C/t \leq \epsilon/4.$$

Construct Φ beforehand on the wider sector $|\arg w| < 3\pi/4$, $|w| < \epsilon$, with further harmless room for nested-sector estimates. The same bounds hold for $s_j + 1$, with the next prototype index. The unforced image $f(P_{\text{loc}}(s_j)) = P_{\text{loc}}(s_j + 1)$ has distance at least c'/K from zero and the angular boundary, and at least $\epsilon/2$ from the outer circle.

The actual forcing is $O(n^{-2})$ by Lemma 2.1. Its full perturbation disk lies inside the wider coordinate domain for large n , since $K \leq n/2$. On this disk $|\Phi'| \leq CK^2$, so one step changes the ideal unit phase increment by at most CK^2/n^2 . The inverse-identification clause of Lemma 4.1, with a fixed slight angular enlargement, ensures that the actual perturbed value is $P_{\text{loc}}(s_{j+1})$. It does not switch branches.

The total additional phase error is at most $CK^3/n^2 \leq C'A^3\delta^2t$. Make this at most $t/20$. Combined with the initial $t/10$ error it is at most $3t/20$, strictly improving the $t/4$ bootstrap. Therefore $|F_n(z)| \leq C/t$ throughout the disk about z_0 . Cauchy's derivative estimate there yields $|F'_n(z_0)| \leq Cn^2/t^2$, proving (25). Conjugation proves the negative-angle case.

The noncircular order of constants is: fixed nested sectors and clock cones; small ϵ and inner analyticity radii; c_0, q ; large A ; small b ; small δ ; large M ; large n . All may depend on fixed σ , but none depends on n, t once chosen. $\square$

Lemma 5.3. *For each fixed $\delta > 0$, uniformly for $\delta \leq |\theta| \leq \pi$ on the same moving circle,*

$$F_n(re^{i\theta}) = O(n^{-1}), \quad F'_n(re^{i\theta}) = O(1). \quad (27)$$

Proof. Put $z = \zeta/n$ and $v_h = nF_h(z)$. On a bounded v -domain, the subsidiary estimate gives

$$v_{h+1} = v_h + \frac{v_h^2 + \zeta^2}{2n} + O(n^{-2}), \quad v_0 = \zeta.$$

The limiting ordinary differential equation has solution

$$v(u) = \zeta \tan(\zeta u/2 + \pi/4), \quad 0 \leq u \leq 1.$$

A pole at positive real $u \leq 1$ requires $\zeta = (\pi/2 + 2k\pi)/u$. On $|\zeta| = T$ only $\zeta = T$ causes a pole; the negative pole family has modulus at least $3T$. Thus the closed arc away from T has compact complex neighborhoods on which the full ODE trajectory is bounded.

Choose a bounded disk strictly containing those trajectories. Its vector field is uniformly Lipschitz, and the local discretization error is $O(n^{-2})$. Discrete Gronwall gives $O(n^{-1})$ error while the numerical orbit stays in the disk; this estimate prevents a first exit for large n . Hence $F_n = O(n^{-1})$ uniformly on the complex neighborhoods. Cauchy in the ζ variable gives $F'_n = O(1)$. The moving arc is contained in those neighborhoods because $nr \rightarrow T$. $\square$

6 The positive measure and its density

Proposition 6.1. *The entire function P is the Laplace transform of a finite positive measure ν on $[0, \infty)$ with every exponential moment, no atom at zero, and support $[0, \infty)$. Its restriction to $(0, \infty)$ has a smooth strictly positive density h .*

Proof. Put $r_n = T/D_n$ and define the finite positive measures

$$\nu_n = \sum_{m \geq 1} [z^m] F_n(z) r_n^m \delta_{m/n}.$$

Their actual Laplace transforms are $F_n(r_n e^{s/n})$. The rational parameter in Proposition 5.1 is

$$D_n - \frac{T}{r_n e^{s/n}} = D_n(1 - e^{-s/n}) = s + O_S(\log n/n)$$

on every compact S . Hence these transforms converge locally uniformly to $P(s)$.

The masses tend to $P(0) > 0$. Lévy's continuity theorem applied to the normalized characteristic functions gives a finite weak limit ν , supported on $[0, \infty)$. For every real u , the transforms at $u + 1$ are uniformly bounded. On $t > R$, the integral of e^{ut} is bounded by e^{-R} times the integral of $e^{(u+1)t}$. This proves exponential uniform integrability, so $P(u) = \int e^{ut} \nu(dt)$. Analyticity and the exponential moments extend this identity to every complex u .

By Lemma 4.1, $P(-x) \sim 2/x$. Thus $\nu(\{0\}) = 0$. Its support has positive points arbitrarily close to zero: a positive gap would imply exponential decay of $P(-x)$, contradicting $2/x$.

For fixed real σ , Lemma 4.1 also gives $P'(\sigma + it) = O(t^{-2})$ as $|t| \rightarrow \infty$, since a vertical line eventually lies in an interior inverse sector. This is the Fourier transform of the finite measure $ue^{\sigma u} \nu(du)$ and is integrable. Fourier inversion gives this weighted measure a continuous density. Dividing by $ue^{\sigma u}$ shows that $\nu(du) = h(u) du$ on $(0, \infty)$, with h continuous and nonnegative.

The functional equation (4) and uniqueness of finite measures give

$$(e^u - 1)\nu = \sum_{k \geq 2} \frac{\nu^{*k}}{k!}. \quad (28)$$

Both sides are finite: the left side has mass $P(1) - P(0)$, and the positive series on the right has finite total mass. In particular $\nu * \nu \leq 2(e^u - 1)\nu$, so its closed support S is closed under addition. Positive support points tending to zero have integer multiples approaching any prescribed positive number. Therefore $S = [0, \infty)$.

Fix $t > 0$. Full support and continuity yield an interval $I \subset (0, t)$ on which $h > 0$. Full support also yields a point $b \in t - I$ with $h(b) > 0$. Then $h(t - b) > 0$, and continuity implies $(h * h)(t) > 0$. The convolution is finite and continuous on $(0, \infty)$: near any fixed t , split the defining integral into two small endpoint pieces and a compact interior. On an endpoint piece one factor is bounded away from zero's neighborhood while the other has arbitrarily small L^1 mass near zero; the interior follows by dominated convergence. Thus the density inequality from (28), initially almost everywhere, holds everywhere by continuity:

$$(e^t - 1)h(t) \geq \frac{1}{2}(h * h)(t) > 0.$$

This proves strict positivity, without a pointwise density expansion at zero. $\square$

[write] **The density at zero, 6 October 2026.** Part III supplies the expansion this proof does without: $h(0) = 2$ (Proposition 27.1), an exact Hankel representation near zero (Theorem 28.1), a C^∞ germ $t^{t/3}h(t)$ (Theorem 29.1) and the power-log expansion (III.28); in particular h is continuous at 0 but not C^1 there.

Corollary 6.2. *The density h is C^∞ on $(0, \infty)$.*

Proof. Fix real σ . Cauchy's estimates in the inverse sectors of Lemma 4.1, applied on disks of radius proportional to $|s|$, give

$$P^{(q)}(\sigma + it) = O_q(|t|^{-q-1}) \quad (|t| \rightarrow \infty)$$

for every integer $q \geq 1$. This is the Fourier transform of the finite measure $u^q e^{\sigma u} \nu(du)$. Given $k \geq 0$, choose $q > k$. Then $|t|^k P^{(q)}(\sigma + it)$ is integrable. Differentiation under Fourier inversion gives a C^k density for this weighted measure. On $(0, \infty)$ it agrees with $u^q e^{\sigma u} h(u)$ by uniqueness and continuity. Divide by the positive smooth weight on every compact subinterval of $(0, \infty)$. Since k was arbitrary, h is smooth there. $\square$

[write] **Proved twice, 6 October 2026.** Part II restates this corollary with the same argument after its interface list (Section 13, around (II.16)).

7 Coefficient extraction and the amplitude

Proof of Theorem 1.1. All structural claims about P and ν were proved above. Fix real σ and put $r = T/(D_n - \sigma)$. Differentiating the Cauchy coefficient formula, or integrating it by parts over the full circle, gives exactly

$$na_n r^n = \frac{1}{2\pi} \int_{-\pi n}^{\pi n} \left[\frac{z}{n} F'_n(z) \right] e^{-it} dt, \quad z = r e^{it/n}. \quad (29)$$

For the derivative normalization, at $z = T/(D_n - s)$ one has

$$\frac{z}{n} F'_n(z) = \frac{D_n - s}{n} G'_n(s).$$

Locally uniform holomorphic convergence in Proposition 5.1 gives $G'_n \rightarrow P'$. On the circle, the actual parameter is $s_n = D_n - (D_n - \sigma)e^{-it/n} \rightarrow \sigma + it$, uniformly for bounded t . The central integrand therefore tends to $P'(\sigma + it)e^{-it}$.

Proposition 5.2 bounds the intermediate tails by $C \int_M^\infty t^{-2} dt = O(M^{-1})$, uniformly in n . Lemma 5.3 bounds $z F'_n(z)/n$ by $O(n^{-2})$ on outer arcs, whose t -length is $O(n)$, so their contribution is $O(n^{-1})$. First let $n \rightarrow \infty$ and then $M \rightarrow \infty$ in (29). Fourier inversion of the density of $u e^{\sigma u} \nu(du)$ at $u = 1$ gives

$$na_n r^n \longrightarrow \frac{1}{2\pi} \int_{\mathbb{R}} P'(\sigma + it) e^{-it} dt = e^\sigma h(1) > 0. \quad (30)$$

This also proves (6).

Finally, expansion of (3) yields

$$r^{-n} = (n/T)^n n^{-1/3} T^{1/3} e^{-B-\sigma} (1 + o(1)).$$

The factor $e^{-\sigma}$ from the contour radius cancels the factor e^σ in (30). Division by n proves (5) with the stated positive constant. $\square$

[write] **Refined in Part II, 6 October 2026.** Part II's Theorem 12.1 adds the first relative correction, with remainder $O(n^{-7/5})$, by extracting coefficients from the thirtieth derivative (Section 19) instead of the first.

8 Proportional height and fixed shifts

Theorem 8.1. *Let $0 < \beta_0 < \beta_1 < \infty$ be fixed. Uniformly over positive integer pairs (n, m) with $m \rightarrow \infty$ and $\beta = n/m \in [\beta_0, \beta_1]$,*

$$A(n, m) \sim \frac{h(\beta)}{m} \left(\frac{m}{T}\right)^n m^{-\beta/3} T^{\beta/3} e^{-\beta B}. \quad (31)$$

The exact integer height m and exact ratio $\beta = n/m$ occur in this formula.

Proof. Fix real σ , let $r_m = T/(D_m - \sigma)$, and extend

$$Q_m(t) = \frac{z}{m} F'_m(z), \quad z = r_m e^{it/m}, \quad |t| \leq \pi m,$$

by zero elsewhere. Proposition 5.1 gives convergence to $Q(t) = P'(\sigma + it)$ uniformly on each compact interval. Proposition 5.2 gives $|Q_m(t)| \leq C/t^2$ in the intermediate range; Lemma 5.3 gives $O(m^{-2})$ in the outer range. The limiting derivative also has an integrable $O(t^{-2})$ tail. Splitting the integral at a fixed M , then letting $m \rightarrow \infty$ and $M \rightarrow \infty$, proves

$$\|Q_m - Q\|_{L^1(\mathbb{R})} \rightarrow 0.$$

Their Fourier transforms consequently converge uniformly for all real frequencies, by the L^1 norm bound. At the exact frequency $\beta = n/m$, the coefficient identity is

$$nA(n, m)r_m^n = \frac{1}{2\pi} \int_{\mathbb{R}} Q_m(t) e^{-i\beta t} dt = \beta e^{\beta\sigma} h(\beta) + o(1).$$

Fourier inversion applies to the continuous density of $ue^{\sigma u} \nu(du)$. On the chosen compact interval, β is bounded away from zero and h has a strictly positive minimum. The absolute error therefore gives the uniform relative estimate

$$A(n, m) = \frac{e^{\beta\sigma} h(\beta)}{m} r_m^{-n} (1 + o(1)).$$

Write $q_m = -(\log m)/3 + (\log T)/3 - B - \sigma$. Since $n = \beta m$ exactly,

$$\begin{aligned} r_m^{-n} &= (m/T)^n \exp(\beta m \log(1 + q_m/m)) \\ &= (m/T)^n m^{-\beta/3} T^{\beta/3} e^{-\beta B - \beta\sigma} \left(1 + O((\log m)^2/m)\right), \end{aligned}$$

uniformly for these β . Cancellation of the σ factors proves the theorem. $\square$

Corollary 8.2. *For every fixed integer k , as $n \rightarrow \infty$,*

$$\frac{[z^n] F_{n+k}(z)}{a_n} \rightarrow e^k.$$

For fixed real $\lambda > 0$, put $\alpha_n = \{\lambda n\}$ and $m = \lfloor \lambda n \rfloor$. Then

$$A(n, m) \sim C_\lambda e^{-\alpha_n/\lambda} (\lambda n/T)^n n^{-1-1/(3\lambda)}, \quad C_\lambda = h(1/\lambda) \lambda^{-1-1/(3\lambda)} T^{1/(3\lambda)} e^{-B/\lambda}. \quad (32)$$

Proof. For $m = n + k$, continuity gives $h(n/m) \rightarrow h(1)$, while $(m/n)^n \rightarrow e^k$. All remaining ratios of prefactors in (31) and (5) tend to one. For the second assertion, $m = \lambda n - \alpha_n$ gives

$$(m/(\lambda n))^n = e^{-\alpha_n/\lambda}(1 + O(n^{-1})), \quad n/m = 1/\lambda + O(n^{-1}).$$

Substitution into (31), using continuity of h and $O(\log n/n)$ for changes in the logarithmic powers, proves (32). $\square$

The oscillating floor factor need not tend to one. Dropping it changes the leading equivalent. Neither theorem asserts uniformity as $\beta \downarrow 0$ or $\beta \rightarrow \infty$.

[write] **Extended in Parts II and III, 6 October 2026.** Part II's Theorem 12.2 adds the first correction on the same compact ranges. Part III's Theorem 31.1 extends the leading equivalent of Theorem 8.1 (the same formula, since $e^{\beta d_m} = m^{-\beta/3} T^{\beta/3} e^{-\beta B}$) toward $\beta \downarrow 0$, uniformly on every band $(\log m)^{-D} \leq \beta \leq B_0$, with an explicit relative error; this yields the crossover of Theorem 24.1. Uniformity as $\beta \rightarrow \infty$ remains open (Section 37, item 5).

Corollary 8.3. *For each fixed $C > 0$, uniformly over integers $|k| \leq C\sqrt{n}$,*

$$e^{-k} \frac{[z^n] F_{n+k}(z)}{a_n} = \exp\left(-\frac{k^2}{2n}\right) (1 + o(1)).$$

Proof. Put $m = n + k$ in Theorem 8.1. Uniformly in this range, $\beta = n/m = 1 + O(n^{-1/2})$, so $h(\beta)/h(1) \rightarrow 1$. The other prefactor ratios tend uniformly to one: their logarithms, aside from the continuous density ratio, are $O((1 + |k|) \log n/n)$. The exponential ratio is determined by

$$n \log(1 + k/n) = k - \frac{k^2}{2n} + O_C(n^{-1/2}).$$

Substitution and division by the diagonal equivalent prove the claim. $\square$

This is a Gaussian-shaped asymptotic profile for height shifts. It is not a probability central limit theorem; no probability distribution on heights has been introduced.

9 An inverse consequence

Corollary 9.1. *Let $y = a_n$ with $n \rightarrow \infty$, put $Y = \log y$, and let W denote the positive real Lambert function. Put*

$$X = \frac{Y}{W(Y/T)}, \quad q = \log(X/T) + 1.$$

Then

$$n = X + \frac{\frac{4}{3} \log X - \log c}{q} + o(1/\log X). \quad (33)$$

In particular $n - X \rightarrow 4/3$. The explicit corrected expression in (33), and the large real root $x(y)$ of

$$\log y = x \log(x/T) - \frac{4}{3} \log x + \log c,$$

recover n by nearest-integer rounding for all sufficiently large exact sequence values $y = a_n$.

Proof. Let $L(x) = x \log(x/T)$. Then $L(X) = Y$ and $L'(X) = q$. Theorem 1.1 gives $L(n) - L(X) = \frac{4}{3} \log n - \log c + o(1)$. First comparison gives $n/X \rightarrow 1$, and the mean-value theorem then gives $n - X = O(1)$. Thus

$$L(n) - L(X) = q(n - X) + O(X^{-1}), \quad \log n = \log X + O(X^{-1}).$$

Division by q proves (33). Applying the same mean-value argument to the full leading model gives $x(a_n) = n + o(1/\log n)$; the rounding assertions follow. $\square$

No effective onset is asserted. This is recovery from exact sequence values, not a universal threshold-rounding rule for arbitrary y between them. The leading equivalent alone does not provide a higher-order error term.

[write] **Sharpened in Part II; the transseries volume, 6 October 2026.** Part II's Corollary 21.1 sharpens (33) to an error $O(X^{-7/5}/\log X)$; its $\delta_0 = (\frac{4}{3} \log X - \log c)/q_X$ is the correction here. $X = Y/W(Y/T)$ solves $X \log(X/T) = Y$, an instance of the factorial core `p0:prop:factorial-core` of [11] ($\kappa = 1$, its $d = -\log T$). The correction and the rounding at exact values are proved directly by the mean-value theorem; they parallel `p0:thm:perturbed-inversion` and part (3) of `p0:thm:staircase` but are not instances of them (front matter, “The OEIS statements, and the inverses”).

10 Exact computation and numerical diagnostics

10.1 The coefficient algorithm

Logarithmically differentiating (2) gives

$$\frac{zF'_{h+1}(z)}{1 + F_{h+1}(z)} = \sum_{j \geq 1} b_j z^j, \quad b_j = \sum_{d|j} d[z^d]F_h(z).$$

Therefore, writing $1 + F_{h+1}(z) = \sum_{k \geq 0} c_k z^k$, one obtains

$$c_0 = 1, \quad c_k = \frac{1}{k} \sum_{j=1}^k b_j c_{k-j} \quad (k \geq 1). \quad (34)$$

Truncation at degree N is exact for every coefficient retained. The public standard-library program checks divisibility before each integer division. It stores F_h with constant term zero and restores the convention $A(0, h) = 1$ only when reporting the array. Degree and height are each capped at 640; this is an implementation and testing limit, not a limitation of the theorems. The diagonal calculation uses $O(N^3)$ integer arithmetic operations and $O(N)$ stored integers; operation costs grow with the bit lengths.

An independent product implementation expands each factor using

$$(1 - z^d)^{-a} = \sum_{k \geq 0} \binom{a + k - 1}{k} z^{dk} \quad (a \geq 1)$$

and multiplies the finite truncations. It checks every entry through degree and height 12. A separate generated exact fixture contains diagonal values through 414. The public tests also compare the 22 displayed OEIS values, $n = 0, \dots, 21$. The full OEIS b-file was not successfully retrieved in the

source review; the 415-entry fixture is therefore explicitly described as generated mathematical data, not as a 415-term external comparison.

All public pass/fail checks use explicit exceptions, so Python optimization does not remove them. Local chunked decimal conversion works with Python’s minimum configurable integer-string digit limit of 640 without modifying that process-wide limit. This matters because the largest exact integers exceed that many digits.

10.2 The amplitude computation is not certified

The normalized ratios $a_n(T/n)^n n^{4/3}$ decrease slowly over the tested range: approximately 4.60133 at $n = 40$, 4.55287 at $n = 80$, 4.52400 at $n = 160$, 4.51777 at $n = 200$, and 4.50437 at $n = 414$. These finite values alone neither prove a limit nor certify its decimal digits.

A separate optional numerical script evaluates the repelling parametrization. The first terms of its normalized formal coordinate are

$$\Phi(w) = -2/w + \frac{1}{3} \log w - \frac{w}{36} + \frac{w^2}{540} + \frac{w^3}{7776} + \cdots.$$

The script uses nine correction coefficients, solves the truncated inverse equation near $s - M$ by Newton iteration, applies M forward iterates of f , and propagates the derivative by multiplication by e^w . Subtracting $2/(1-s)^2$ from the derivative before quadrature improves the observed tail: its known contribution to $h(1)$ is $2/e$ for $\sigma < 1$. Tests varying the forward depth, vertical contour, cutoff, and mesh give values near 4.4923299, consistent with the conjectured 4.4923... Floating-point agreement is not an interval enclosure. There is no certified digit count in this report, and the analytic theorem does not depend on these calculations.

[write] **Checks, 6 October 2026.** The shipped diagnostic `data/225-leading-data-amplitude_diagnostic.json` records $h(1) = 1.8146095596784502$ and $c = 4.492329896751906$; the write recomputed $T^{1/3}e^{-B}h(1) = 4.4923298968$ from that $h(1)$. Part II’s derivative diagnostics give $h(1) \approx 1.81460956$ as well (Subsection 22.2). The coefficients above agree with Part III’s $a_1 = -1/36$, $a_2 = 1/540$, $a_3 = 1/7776$, and the write’s rerun of `coordinate_series.py--order9` also gives $a_4 = -71/435456$. All values remain uncertified.

10.3 Reproducibility and limits

The accompanying archive contains the complete editable TeX, exact programs and tests, numerical diagnostic script, exact fixture, source ledger, and a deterministic builder. The builder verifies an input manifest, runs explicit tests, creates a private pdfLaTeX format, compiles three times without shell escape, rejects layout and reference warnings, and packages sorted members with fixed timestamps. It preserves the source tree and writes outputs only to a new separate directory. Ordinary and optimized cap-640 runs are compared by full ZIP bytes and by each member. Byte identity is a claim about the same toolchain; arbitrary TeX, font, or numerical-library versions need not reproduce identical bytes.

[write] **The companion here, 6 October 2026.** Report 225’s companion is shipped as `code/225-leading-*.py`, `data/225-leading-*` and `225-leading-sources-SOURCE_LEDGER.md`; its delivered README was replaced by the report README, and its PDF and `MANIFEST.sha256` are not shipped. The builder and tests run only in the delivered layout; the report README gives the rerun route from the arrival commit.

11 Further questions and quantitative extensions

11.1 Validated evaluation of the positive constant

The exact expression (6) is a useful starting point for a certificate, but a validated computation requires more than increased precision. One must bound the omitted Fatou-coordinate terms on a stated sector, enclose the inverse branch, control the forward propagation of both value and derivative, validate the finite-interval quadrature, and give an explicit integrable bound for the remaining vertical tail. The sector proof above supplies qualitative constants; it does not extract numerical values for all of them. A practical goal is a reproducible interval for c narrow enough to validate the conjectured initial digits, with each of these error sources accounted for separately.

The rational subtraction used in the diagnostic script could be extended to several terms derived from the large- s inverse expansion. This can reduce cancellation and the cutoff length, but every subtracted term and its Fourier contribution must be explicit. Mesh or precision stability cannot replace bounds for truncation of the asymptotic coordinate and the infinite integral.

[write] **Still open, 6 October 2026.** Not settled by Parts II or III; Section 37, item 1. Part II’s closed form of κ removes one constant from the first correction, which still needs $h(1)$, $h'(1)$, $h''(1)$.

11.2 Higher corrections to the diagonal

The logarithmic center D_n is determined only to the precision needed for a leading equivalent. Higher corrections require retaining more subsidiary terms in $R_h(z)$, including their dependence on h , and constructing a corresponding hierarchy of nonautonomous clocks. Their matching constants interact with the inverse-coordinate expansion. It is natural to investigate corrections involving powers of $1/n$ and polynomials in $\log n$, but no complete expansion of this form is asserted here.

A formal expansion at bounded s is insufficient for a coefficient theorem. The error must be controlled in an integrable norm on the growing contour, with the intermediate-angle and outer-angle arguments strengthened to the same order. The present choice $K = \lfloor \sqrt{n} \rfloor$ balances errors only for convergence; a higher-order theorem would need a new quantified matching analysis. This identifies a concrete obstruction to promoting formal series directly into an all-orders claim.

[write] **Partly answered by Part II, 6 October 2026.** Part II proves the first correction, $(-\frac{1}{18} \log^2 n + A \log n + C)/n$ with remainder $O(n^{-7/5})$ (Theorem 12.1), by two further clock terms and a quantified growing-strip matching with $K = \lfloor n^{1/2} \rfloor$; it confirms the expected shape, a polynomial in $\log n$ of degree two over n . Corrections beyond the first and an all-orders theorem remain open (Section 37, item 2).

11.3 Effective bounds and inverse thresholds

Making the constants in the clock, stopping, and sector estimates explicit would give finite- n bounds and a computable onset for the leading equivalent. Such bounds would in turn permit certified recovery of integer thresholds for arbitrary input sizes. Corollary 9.1 currently supplies eventual rounding at exact sequence values; it does not say how large “eventually” is, nor does it control the location of a general input between adjacent terms. A rigorous threshold algorithm should combine exact small-index computation with explicit upper and lower bounds, rather than treat a floating-point evaluation of c as exact.

One can also seek inverse statements for $A(n, \lfloor \lambda n \rfloor)$. Formula (32) shows why the bounded floor phase must enter such statements. A single smooth approximation that discards this phase can

have a nonvanishing multiplicative error.

[write] **Still open, 6 October 2026.** Part II’s Corollary 21.1 sharpens the inverse at exact sequence values, still without onset; Part III’s inverses concern ratios in the window $m \asymp n \log n$, not $A(n, \lfloor \lambda n \rfloor)$. Section 37, items 3 and 4.

11.4 Other height scales and the density

Theorem 8.1 treats n/m in a fixed compact subset of $(0, \infty)$. Extending it toward zero or infinity requires information about $h(t)$ at the corresponding endpoint and renewed uniformity estimates. In particular, the strict-positivity proof does not give an expansion of $h(t)$ at zero. The measure identity

$$(e^t - 1)\nu = \sum_{k \geq 2} \nu^{*k}/k!$$

is a possible starting point for endpoint analysis and stronger regularity, such as real analyticity. Smoothness away zero is already established in Corollary 6.2, but differentiation of this convolution identity or exchange of limits with its series needs separate justification.

Kaneiwa’s fixed-leaf polynomial regime, proportional height, and fixed-height partition regimes are genuinely different limiting questions. A uniform transition theory between them would explain how the polynomial and parabolic descriptions fit together. It may also clarify which parts of this proof extend to other multiset recurrences or other initial generating functions. Such a generalization must recompute the subsidiary forcing and the clock constant; the unforced map alone does not determine the answer.

[write] **Partly answered by Part III, 6 October 2026.** At the endpoint zero Part III proves the expansion asked for here: $h(0) = 2$, the Hankel representation and the power-log series (Proposition 27.1, Theorem 28.1, Corollary 29.2), and it extends the height theorem toward $n/m \rightarrow 0$ on logarithmic bands (Theorem 31.1), with Kaneiwa’s normalization compared in Section 33. Theorem 28.1 also gives real analyticity of h on some interval $(0, \epsilon)$. Open: $n/m \rightarrow \infty$, real analyticity on all of $(0, \infty)$, the transition to the fixed-degree regime, other recurrences (Section 37, items 5–8).

Part II

First logarithmic correction for iterated Euler diagonals

Bundle Report 226, archive `Report226.zip`, delivered as `src/report226.tex` (17 pages). Delivered title block “First logarithmic correction for iterated Euler diagonals / Report 226 / 5 October 2026”. Printed as delivered, with `[write]` notes. Delivered Section k is Section $k + 11$ here, its statement $k.j$ is $(k + 11).j$, and its equation (k) is (II. k).

Reading conventions of Part II	Elsewhere in the report
“Report 225”, [6]	Part I
$b_{n,m}$ the array; d_* , d , d_m	Parts I, III: $A(n, m)$; Part I: D_n , $d = n - h$
$\rho_1, \rho_2 = h'(1)/h(1), h''(1)/h(1)$; $\rho_j(\beta)$	Part III: ρ_j rational
A, C the correction constants; G Catalan’s constant	Parts I, III: $A(n, m)$; Part III: $\mathcal{C}$ Catalan, C_0 , $C(\beta)$
$M(y)$, $N(y)$ clock functions; $M = n^{1/20}$; $\mathcal{C}_z(w)$	Part III: $M = m^{1/20}$, N an order
$q = 30$ the derivative order; $\alpha = 4/3$, $\ell = \log c$	Part I: q entrance constant; Part III: ℓ_n Kaneiwa

Abstract of this Part (as delivered). For the iterated Euler diagonal a_n in OEIS A290354, we prove

$$a_n = c(2n/\pi)^n n^{-4/3} \left\{ 1 + \frac{-\log^2 n/18 + A \log n + C}{n} + O(n^{-7/5}) \right\}.$$

The constants A and C are exact expressions in derivatives of the positive density associated with the normalized repelling parametrization of $w \mapsto e^w - 1$. A new nonautonomous clock constant is evaluated in terms of π , $\log 2$, and Catalan’s constant. The proof constructs two further clock terms, obtains a weighted fourth-power defect, and matches a forced Fatou correction on a growing vertical strip. Thirtieth-derivative coefficient extraction turns these estimates into the stated relative remainder. A uniform correction for compact positive degree-to-height ratios and a sharpened Lambert W inverse are included. This companion uses the analytic foundation of the preceding unrefereed Report 225, whose unchanged PDF and TeX accompany the archive. It is an unrefereed research proof, not a proof-assistant formalization. Floating-point diagnostics are not certified decimal enclosures; no all-orders theorem is claimed.

12 Definition and main results

Define

$$F_0(z) = z, \quad F_{h+1}(z) = \exp \left(\sum_{j \geq 1} \frac{F_h(z^j)}{j} \right) - 1. \quad (\text{II.1})$$

For $n \geq 1$ and $m \geq 0$, put $b_{n,m} = [z^n]F_m(z)$ and $a_n = b_{n,n}$; set $a_0 = 1$ and $b_{0,m} = 1$ separately when reporting the integer arrays. Thus a_n is A290354 and $b_{n,m}$ is A290353 [2, 1]. The analytic generating functions themselves have constant term zero. They count unlabeled rooted trees whose leaves all have the stated height: a new root carries a nonempty unordered multiset of trees at the previous height. Equivalently,

$$1 + F_{h+1}(z) = \prod_{d \geq 1} (1 - z^d)^{-[z^d]F_h(z)}.$$

There is no factorial normalization. The diagonal begins

$$1, 1, 2, 6, 30, 170, 1337, 12166, 133476, 1676364.$$

The exact constants and functions used throughout are

$$T = \frac{\pi}{2}, \quad B = \frac{1}{2} - \frac{\pi}{6} - \frac{\pi^2}{16} - \frac{\log 2}{6}, \quad d_\star = \frac{1}{3} \log T - B, \quad (\text{II.2})$$

$$\kappa = \frac{\pi^3}{96} + \frac{\pi^2}{12} - \frac{13\pi}{48} - \frac{11}{36} - \frac{G}{6} - \frac{\pi \log 2}{24}, \quad (\text{II.3})$$

where $G = \sum_{j \geq 0} (-1)^j / (2j+1)^2$ is Catalan's constant. Let P denote the entire repelling parametrization normalized by

$$P(s+1) = e^{P(s)} - 1, \quad \Phi(w) = -\frac{2}{w} + \frac{1}{3} \log w + O(w), \quad \Phi(P(s)) = s \quad (\text{II.4})$$

in the inverse sector near negative real s . Report 225 [6] constructs this normalization and proves

$$P(s) = \int_0^\infty e^{sx} h(x) dx, \quad h \in C^\infty(0, \infty), \quad h(x) > 0 \quad (x > 0). \quad (\text{II.5})$$

The density has every exponential moment. Set

$$c = T^{1/3} e^{-B} h(1), \quad \rho_1 = \frac{h'(1)}{h(1)}, \quad \rho_2 = \frac{h''(1)}{h(1)}. \quad (\text{II.6})$$

The zero additive constant in (II.4) is essential: a translation of the Fatou coordinate would change P and h .

Theorem 12.1. *As $n \rightarrow \infty$,*

$$a_n = c(n/T)^n n^{-4/3} \left(1 + \frac{-\log^2 n / 18 + A \log n + C}{n} + O(n^{-7/5}) \right), \quad (\text{II.7})$$

where $c > 0$ and

$$A = \frac{d_\star}{3} + \frac{4}{9} + \frac{\rho_1}{3}, \quad (\text{II.8})$$

$$C = -\frac{d_\star^2}{2} - d_\star \left(\frac{4}{3} + \rho_1 \right) - T\kappa - \frac{1}{3} - \frac{4\rho_1}{3} - \frac{\rho_2}{2}.$$

The error is relative to the displayed leading expression. Its exponent is a convenient nonoptimal choice.

With $d = -\frac{1}{3} \log n + d_\star$, the numerator of the correction can equivalently be written

$$-\frac{d^2}{2} - d \left(\frac{4}{3} + \rho_1 \right) - T\kappa - \frac{1}{3} - \frac{4\rho_1}{3} - \frac{\rho_2}{2}. \quad (\text{II.9})$$

In particular the squared-logarithm coefficient is exactly $-1/18$. Retaining only a more accurate contour radius would miss some terms in A and C .

Theorem 12.2. *Let $I = [\beta_0, \beta_1] \subset (0, \infty)$ be fixed. Uniformly for integers $m \rightarrow \infty$ and $n \geq 1$ with $\beta = n/m \in I$, put $d_m = -\frac{1}{3} \log m + d_\star$ and $\rho_j(\beta) = h^{(j)}(\beta)/h(\beta)$ for $j = 1, 2$. Then*

$$b_{n,m} = h(\beta) T^{\beta/3} e^{-\beta B} (m/T)^n m^{-1-\beta/3} \left(1 + \frac{\mathcal{Q}_\beta(d_m)}{m} + O_I(m^{-7/5}) \right), \quad (\text{II.10})$$

$$\begin{aligned} \mathcal{Q}_\beta(d) = & -\frac{\beta d^2}{2} - d \left(1 + \frac{\beta}{3} + \beta \rho_1(\beta) \right) - \beta \kappa T \\ & - \frac{1}{3} - \left(1 + \frac{\beta}{3} \right) \rho_1(\beta) - \frac{\beta}{2} \rho_2(\beta). \end{aligned}$$

The exact height and exact ratio occur in the formula; no additional factor of β belongs in its leading amplitude.

No uniform conclusion is asserted as β approaches zero or infinity. In particular replacing an integer height by an unfloored real height without retaining its phase is not justified.

[write] **Toward $\beta \rightarrow 0$ in Part III, 6 October 2026.** Part III extends the *leading* term of (II.10) to the bands $(\log m)^{-D} \leq \beta \leq B_0$ (Theorem 31.1); no Part extends the correction $\mathcal{Q}_\beta$ there, and $\beta \rightarrow \infty$ is open (Section 37, item 5).

12.1 Dependence on the preceding report and prior work

Report 225 proves the leading equivalent and its analytic foundation. The present report extends its clock and coefficient analysis; it does not independently reprove all entrance and angular-tail estimates. Section 13 records the full interface used, and the accompanying archive includes the unchanged complete proof of Report 225. Neither report has undergone journal refereeing.

The OEIS attributes the leading equivalent with numerical amplitude 4.4923... to Václav Kotěšovec, dated 14 August 2017 [2]. The fixed-degree polynomiality and tangent leading profile have earlier antecedents explained and sourced in Report 225. The correction proved here is not inferred merely from that fixed-degree information. We make no worldwide priority claim for the result or for parabolic-coordinate methods.

[write] **Reading Part II against Part I, 6 October 2026.** “Report 225” and [6] are Part I of this report; its Sections 2–7 are Part I’s Sections 2–7, with the same numbers. The included copy is byte-identical to Part I and is not shipped. At leading order Theorem 12.1 is Part I’s Theorem 1.1 and Theorem 12.2 is Part I’s Theorem 8.1. Here $b_{n,m}$ is Part I’s $A(n, m)$, and $d_\star = \frac{1}{3} \log T - B$, so D_n is Part I’s D_n . Part II’s ρ_j are not Part III’s rational ρ_j .

13 Analytic foundation from Report 225

All branches agree with real values on the positive axis. Constants in local analytic estimates may depend on fixed sectors or fixed real-width bounds, but not on the growing indices. The following are the precise results imported from Report 225, Sections 2–7 [6].

- (i) Each F_h is analytic in $|z| < 1$, with nonnegative coefficients and linear coefficient one. If $(h+1)|q|$ is sufficiently small, then $|F_h(q)| \leq C|q|$ uniformly. This follows by positive-coefficient comparison on the real radius and a reciprocal recurrence bound; it does not assume that the main value $F_h(z)$ is small.

(ii) Define

$$J(y) = 2 \arctan y - T, \quad L(1) = 0, \quad L'(y) = \frac{y^3/3 - y - 4/3 - 2J(y)}{(1+y^2)^2},$$

$$H(z) = \frac{T}{z} + \frac{1}{3} \log \frac{1}{z} + B. \quad (\text{II.11})$$

The old clock $J(w/z)/z + L(w/z)$ gives a noncircular positive-real entrance bound and its Cauchy extension. More precisely, let $D_n = n - \frac{1}{3} \log n + d_\star$ and $r = T/D_n$. For suitable fixed $q > 0$, K_0 , and $\epsilon > 0$, if $K_0 \leq K \leq \epsilon n$, then on $|z - r| \leq qK/n^2$ the orbit through $n - K$ is small and

$$|F_h(z)| \asymp (n - h)^{-1} \quad (0 \leq h \leq n - K), \quad F_h(z)/z \text{ lies in a fixed narrow positive cone.} \quad (\text{II.12})$$

The cone has a slightly larger fixed cone available for Taylor segments. At entrance the old phase estimate is

$$\Phi(F_{n-K}(z)) = n - K - H(z) + O(K^{-1} + n^{-1} + K^3/n^2). \quad (\text{II.13})$$

(iii) The sector coordinate Φ in (II.4) satisfies the Abel equation $\Phi(f(w)) = \Phi(w) + 1$, $f(w) = e^w - 1$. Its inverse is uniquely selected in fixed interior sectors about the closed left half-plane for s of large modulus, and

$$P(s) = -\frac{2}{s} + O\left(\frac{\log |s|}{|s|^2}\right). \quad (\text{II.14})$$

A small point w in a smaller coordinate sector satisfying $\Phi(w) = s$ is the same inverse value $P(s)$; global injectivity is unnecessary. The inverse iterates $g^{\circ j}(w)$, $g(w) = \log(1 + w)$, obey $|g^{\circ j}(w)| \leq C/(|w|^{-1} + j)$ in nested sectors. Extension by finitely many iterates of f gives the entire P .

(iv) There are fixed $b, \delta, M_0, C > 0$ such that for $M_0 \leq |t| \leq \delta n$,

$$|z - re^{it/n}| \leq b|t|/n^2 \implies |F_n(z)| \leq C/|t|. \quad (\text{II.15})$$

For the outer angles $\delta \leq |\theta| \leq \pi$, the compact arc $\zeta \approx Te^{i\theta}$ has fixed complex neighborhoods on which $F_n(\zeta/n) = O(n^{-1})$. These neighborhoods follow from the bounded ODE profile $\zeta \tan(\zeta u/2 + \pi/4)$ for $0 \leq u \leq 1$ away from its positive pole. Both are bounds on two-dimensional analytic neighborhoods, not merely on one-dimensional arcs.

(v) The positive-measure construction yields (II.5), with no atom at zero, support $[0, \infty)$, and strict positivity of h on $(0, \infty)$. Its leading coefficient consequence is $a_n \sim c(n/T)^n n^{-4/3}$.

[write] Where Part I proves each item, 6 October 2026. (i) Lemma 2.1 and the first step of its proof. (ii) (9)–(10), Lemmas 2.2–3.3 and (23) (Part I writes $K \leq \kappa n$ for the $K \leq \epsilon n$ here; this κ is not the clock constant). (iii) Lemma 4.1, (20)–(21); the bound on inverse iterates is the estimate $|G^j(u)| \geq c(|u| + j)$ of its proof, read in $w = 2/u$. (iv) Proposition 5.2, whose proof (not its statement) bounds F_n on the whole disk about z_0 , and Lemma 5.3, whose proof (not its statement, which concerns the moving circle) gives the bound on complex neighbourhoods of the arc. (Extended after the independent check of 6 October 2026; the first wording read “Proposition 5.2, whose proof bounds F_n on the whole disk about z_0 , and Lemma 5.3”.) (v) Proposition 6.1 and Theorem 1.1. The paragraph below repeats Part I’s Corollary 6.2 with its proof. The section is printed in full because its displays are cited below.

For clarity, the derivative regularity needed below also follows directly from this interface. Sector Cauchy estimates in (II.14) give, for each fixed $q \geq 1$,

$$P^{(q)}(\sigma + it) = O_q(|t|^{-q-1}) \quad (\text{II.16})$$

uniformly for σ in a bounded interval. This is the Fourier transform of $x^q e^{\sigma x} h(x)$. If $q > k$, its product with t^k is integrable, so Fourier differentiation gives a C^k weighted density. Dividing by the nonzero smooth weight on compact positive intervals proves $h \in C^\infty(0, \infty)$. No boundary behavior of h or its derivatives at zero is used.

14 Subsidiary forcing through fifth scaled order

Write

$$R_h(z) = \sum_{j \geq 2} F_h(z^j)/j, \quad w = F_h(z), \quad y = w/z, \quad t = hz.$$

The map is $w_{h+1} = f(w_h + R_h(z))$.

Lemma 14.1. *Uniformly for $h \leq C_0/|z|$, as $z \rightarrow 0$,*

$$\begin{aligned} R_h(z) = & \frac{z^2}{2} + z^3 \left(\frac{1}{3} + \frac{t}{2} \right) + z^4 \left(\frac{1}{4} + \frac{t^2}{4} \right) \\ & + z^5 \left(\frac{1}{5} + \frac{7t}{12} + \frac{t^3}{6} \right) + O(|z|^6). \end{aligned} \quad (\text{II.17})$$

Proof. Coefficient extraction in one Euler step gives

$$[q^2]F_h = h, \quad [q^3]F_h = \frac{h(h+1)}{2}, \quad [q^4]F_h = \frac{h(h+1)(2h+1)}{6}.$$

Indeed, if the current coefficients are a, b, c , their increments are respectively 1, $a+1$, and $b+a^2/2+3a/2+1$; the latter becomes $(h+1)^2$. All three start at zero.

Take $\varrho = c_0/(h+1)$ with c_0 small. The small-subsidiary bound gives $\sup_{|q| \leq \varrho} |F_h(q)| \leq C\varrho$. Cauchy and a geometric sum therefore give, for $|q| \leq \varrho/2$,

$$F_h(q) = q + hq^2 + \frac{h(h+1)}{2}q^3 + \frac{h(h+1)(2h+1)}{6}q^4 + O((h+1)^4|q|^5). \quad (\text{II.18})$$

For small z , every $q = z^j$, $j \geq 2$, lies in this disk. The $j = 2$ remainder is $O((h+1)^4|z|^{10}) = O(|z|^6)$, and $j \geq 6$ contributes $O(|z|^6)$. At order z^5 , the baseline is $1/5$; the linear- h part of $[q^3]F_h$ at $j = 2$ contributes $t/4$, $[q^2]F_h$ at $j = 3$ contributes $t/3$, and the cubic- h part of $[q^4]F_h$ at $j = 2$ contributes $t^3/6$. This proves (II.17), including the otherwise easily lost $7t/12$ term. $\square$

[write] **Extends Part I, 6 October 2026.** This extends Part I's (8), whose $hz^4/2$ is the term $z^3t/2$ here. The three coefficients $[q^2]F_h$, $[q^3]F_h$, $[q^4]F_h$ are the rows $n = 2, 3, 4$ of A290353 (its cross-references name A001477, A000217, A000330); the write checked them against the exact recurrence for $h \leq 12$.

For the clock algebra write $d_y = 1 + y^2$ and $a(y) = d_y/2$. Expansion of the exponential gives

$$\Delta y = za + z^2b(y, t) + z^3v(y, t) + z^4u(y, t) + O(|z|^5(1 + |y|)^6), \quad (\text{II.19})$$

where

$$\begin{aligned} b(y, t) &= y^3/6 + y/2 + 1/3 + t/2, \\ v(y, t) &= y^4/24 + y^2/4 + y/3 + yt/2 + 3/8 + t^2/4, \\ u(y, t) &= y^5/120 + y^3/12 + y^2/6 + y^2t/4 + 3y/8 + yt^2/4 \\ &\quad + 11/30 + 5t/6 + t^3/6. \end{aligned} \quad (\text{II.20})$$

The estimate is uniform for bounded t when $|w| + |z|$ is small. These are finite Taylor identities with analytic remainders, not extrapolations from numerical values.

15 Two further nonautonomous clock terms

Define $M(1) = N(1) = 0$ on the same simply connected cone as J, L . First set

$$Q_2(y, t) = J'v + J''ab + \frac{J'''}{6}a^3 + L'b + \frac{L''}{2}a^2. \quad (\text{II.21})$$

The required equation for M is

$$aM' = -Q_2(y, J) - \frac{L}{d_y}. \quad (\text{II.22})$$

Its expanded form is

$$M' = -\frac{1}{36d_y^3} \{ 36J^2(y^2 - 1) + 12J(5y^3 - 3y - 8) + 72Ld_y \\ + y^6 + 9y^4 + 40y^3 - 21y^2 - 24y - 29 \}. \quad (\text{II.23})$$

The feedback term L/d_y is necessary. Setting $hz = J$ before retaining the self-consistency terms would give a wrong next clock.

Next define

$$Q_3(y, t) = J'u + J''(av + b^2/2) + \frac{J'''}{2}a^2b + \frac{J''''}{24}a^4 \\ + L'v + L''ab + \frac{L'''}{6}a^3 + M'b + \frac{M''}{2}a^2, \quad (\text{II.24})$$

$$G_3(y) = Q_3(y, J) + L \partial_t Q_2(y, J) + M/d_y, \quad N' = -G_3/a. \quad (\text{II.25})$$

Here $\partial_t Q_2(y, J) = J/d_y + L'/2$. Equations (II.22) and (II.25), with the zero values at one, are exact recursive definitions involving only differentiation, integration, and rational operations. They specify the functions without a fitted constant.

15.1 Large ratio expansions

Uniformly on each closed smaller cone,

$$J = T - 2/y + 2/(3y^3) + O(y^{-5}), \\ L = \frac{1}{3} \log y + B + \frac{5}{6y^2} + \frac{4/9 + 2T/3}{y^3} - \frac{7}{4y^4} + O(y^{-5}). \quad (\text{II.26})$$

Substitution in (II.23) gives

$$M' = -\frac{1}{36} - \frac{1}{6y^2} - \frac{5T/3 + 10/9}{y^3} \\ + \frac{-(2/3) \log y - 2B - T^2 + 9/2}{y^4} + O(y^{-5}), \quad (\text{II.27})$$

and integration yields

$$M = -\frac{y}{36} + \kappa + \frac{1}{6y} + \frac{5T/6 + 5/9}{y^2} + \frac{(2/9)\log y + 2B/3 + T^2/3 - 77/54}{y^3} + O(y^{-4}). \quad (\text{II.28})$$

Thus M has neither a divergent $\log y$ nor a divergent $\log^2 y$ term. The constant is the convergent integral

$$\kappa = \frac{1}{36} + \int_1^\infty \left(M'(y) + \frac{1}{36} \right) dy. \quad (\text{II.29})$$

For the next clock, direct expansion in (II.25) gives

$$\begin{aligned} G_3 &= -\frac{y^3}{540} - \frac{y}{360} + \frac{T}{12} + \frac{1}{18} \\ &\quad + \frac{(5/18)\log y + 5B/6 + 5T^2/12 - 3/20}{y} + O(\log y/y^2), \\ N' &= \frac{y}{270} + \frac{1}{540y} - \frac{T/6 + 1/9}{y^2} \\ &\quad + \frac{-(5/9)\log y - 5B/3 - 5T^2/6 + 161/540}{y^3} + O(\log y/y^4), \\ N &= \frac{y^2}{540} + \frac{\log y}{540} + n_0 + \frac{T/6 + 1/9}{y} \\ &\quad + \frac{(5/18)\log y + 5B/6 + 5T^2/12 - 11/1080}{y^2} + O(\log y/y^3). \end{aligned} \quad (\text{II.30})$$

The finite n_0 is fixed by $N(1) = 0$ and is not needed at the present order. Unlike M , N contains a divergent logarithm. Its clock contribution is $z^2 \log y$, below the first correction but relevant to future orders.

15.2 Exact evaluation of the new clock constant

Put $u = \arctan y$. An exact primitive, checked by differentiation and its value at $u = T/2$, is

$$\begin{aligned} L(\tan u) &= -\frac{1}{3} \log \cos u + \frac{1}{3} - \frac{1}{6} \cos 2u + (T - 2/3)u - u^2 \\ &\quad + (T/2 - 1/3 - u) \sin 2u - \frac{1}{6} \log 2 - \frac{1}{4} T^2 + T/3. \end{aligned} \quad (\text{II.31})$$

Substituting it and $J = 2u - T$ in (II.23) gives

$$(M' + 1/36) dy = \left(\frac{2}{3} \cos^2 u \log \cos u + \mathcal{P}(u) \right) du,$$

where the finite trigonometric polynomial is

$$\mathcal{P} = p_0 + p_2 \cos 2u + p_4 \cos 4u + q_2 \sin 2u + q_4 \sin 4u$$

and

$$\begin{aligned} p_0 &= 2u^2 - 2Tu + T^2/2 + 8u/3 - 4T/3 + 1/12 + \log 2/6, \\ p_2 &= 3u^2 - 3Tu + 3T^2/4 + 10u/3 - 5T/3 + 1/3 + \log 2/6, \\ p_4 &= u^2 - Tu + T^2/4 + 2u/3 - T/3 + 1/12, \\ q_2 &= 2u/3 - T/3 + 2/9, \quad q_4 = 7u/6 - 7T/12 + 7/18. \end{aligned}$$

Elementary termwise integration yields

$$\int_{T/2}^T \mathcal{P}(u) du = -\frac{5\pi}{16} - \frac{5}{12} - \frac{\log 2}{12} + \frac{\pi \log 2}{24} + \frac{\pi^3}{96} + \frac{\pi^2}{12}.$$

The two logarithmic integrals are

$$\begin{aligned} \int_{\pi/4}^{\pi/2} \log \cos u du &= -\frac{\pi}{4} \log 2 - \frac{G}{2}, \\ \int_{\pi/4}^{\pi/2} \cos 2u \log \cos u du &= \frac{\log 2}{4} + \frac{\pi}{8} + \frac{1}{4}. \end{aligned}$$

The first follows by combining the half-period log-cosine integral with $G = -\int_0^{\pi/4} \log \tan u du$; the latter identity also follows from the Fourier series of the logarithm. For the second, integration by parts leaves the boundary contribution $\log 2/4$ and $\int_{\pi/4}^{\pi/2} \sin^2 u du = \pi/8 + 1/4$. The other endpoint term tends to zero. Using $\cos^2 u = (1 + \cos 2u)/2$ in (II.29), including its $1/36$ normalization, proves exactly (II.3).

[write] **Checks, 6 October 2026.** At intake the closed form was compared with a direct quadrature of (II.29), using the primitive (II.31) (itself checked against quadrature of (9)): -0.2543481698511 against -0.2543481698560 , the difference coming from the truncated tail. The delivered symbolic suite (`code/226-firstcorr-code-clock_checks.py`) reproduced `data/226-firstcorr-data-symbolic_checks.json` on a copy at the write.

16 Weighted defect and accumulated clock error

Define

$$\mathcal{C}_z(w) = J(w/z)/z + L(w/z) + zM(w/z) + z^2N(w/z), \quad E_h = \mathcal{C}_z(F_h(z)) - h. \quad (\text{II.32})$$

All four functions vanish at ratio one, so $E_0 = 0$.

Lemma 16.1. *Suppose $h \leq C_0/|z|$, the orbit is in the small-clock range, and $y_h = F_h(z)/z$ remains in the fixed cone. Then*

$$|E_{h+1} - E_h| \leq \frac{C|z|^2|E_h|}{(1 + |y_h|)^2} + C(|F_h(z)|^4 + |z|^4). \quad (\text{II.33})$$

Consequently, on the entrance disks of (II.12),

$$\mathcal{C}_z(F_{n-K}(z)) = n - K + O(K^{-3} + n^{-3}). \quad (\text{II.34})$$

Proof. Put $Y = 1 + |y|$ on a slightly wider cone. There $|d_y| \asymp Y^2$ and $|w| + |z| \asymp |z|Y$. The definitions and sector expansions give, for the fixed derivative orders required below,

$$\begin{aligned} J^{(r)} &= O(Y^{-r-1}), & L^{(r)} &= O(Y^{-r}), \\ M^{(r)} &= O(Y^{1-r}), & N^{(r)} &= O(Y^{2-r}) \quad (r \geq 1), \end{aligned}$$

and $J = O(1)$, $L = O(1 + \log Y)$, $M = O(Y)$, $N = O(Y^2)$. Since $\Delta y = O(|z|Y^2)$, its relative size is $O(|w| + |z|)$; all Taylor segments stay in the larger cone for a sufficiently small clock radius.

Temporarily replace R_h by its polynomial through degree five, keeping a bounded independent t . The discarded $O(z^6)$ forcing changes the clock by $O(z^6/(|w| + |z|)^2) = O(z^4)$. Expand J/z to fourth order in the displacement, L to third, zM to second, and z^2N to first. Their omitted terms are bounded respectively by

$$|z|^{-1}Y^{-6}(|z|Y^2)^5, \quad Y^{-4}(|z|Y^2)^4, \quad |z|Y^{-2}(|z|Y^2)^3, \quad |z|^2(|z|Y^2)^2.$$

Each is $O(|z|^4Y^4)$. The step remainder $O(|z|^5Y^6)$ in (II.19), multiplied by the leading ratio derivative $O(|z|^{-1}Y^{-2})$, has the same bound.

The retained clock increment is exactly

$$1 + z \frac{t - J}{d_y} + z^2\{Q_2(y, t) + aM'\} + z^3\{Q_3(y, t) + aN'\} + O(|z|^4Y^4).$$

Set $t_0 = z\mathcal{C}_z(w) = J + zL + z^2M + z^3N$. It is bounded, and $t_0 - J = O(|z|(1 + \log Y))$ because $|z|Y$ is small. Expansion in t about J gives the terms

$$z^2\{L/d_y + Q_2(y, J) + aM'\} + z^3\{M/d_y + L\partial_t Q_2(y, J) + Q_3(y, J) + aN'\}.$$

They vanish by (II.22) and (II.25). All discarded t -Taylor terms are $O(|z|^4Y^4)$: representative largest bounds are $|z|^4Y^3$ and $|z|^4(1 + \log Y)^2Y^2$, both admissible.

The actual variable satisfies $t = hz = t_0 - zE_h$ exactly. Both t and t_0 are bounded before any bound on E_h is used, so their connecting segment is bounded. On it $\partial_t R_{\text{polynomial}} = O(z^3)$ and $\partial_w \mathcal{C}_z = O((|w| + |z|)^{-2})$. The derivative of the increment with respect to t is therefore $O(|z|/Y^2)$. Multiplication by $-zE_h$ gives the feedback term in (II.33); $|z|^4Y^4 \asymp |w|^4 + |z|^4$ gives the remainder.

Discrete Gronwall has multiplier at most $\exp(Cn|z|^2) = O(1)$. The already established entrance orbit gives

$$\sum_{h < n-K} |F_h(z)|^4 = O(K^{-3}), \quad n|z|^4 = O(n^{-3}).$$

Together with $E_0 = 0$ this proves (II.34). The orbit bounds come from the old clock, so the improved defect is not being used to bootstrap its own hypotheses. $\square$

17 Finite Fatou expansion and forced matching

Lemma 17.1. *In fixed smaller coordinate sectors,*

$$\Phi(w) = -\frac{2}{w} + \frac{1}{3} \log w - \frac{w}{36} + \frac{w^2}{540} + O(w^3). \quad (\text{II.35})$$

For

$$U_0(w) = \frac{2}{3w^3} + \frac{5}{6w^2} + \frac{1}{6w}, \quad (\text{II.36})$$

one has $U_0(f(w)) - U_0(w) = -\Phi'(w)/2 + O(w)$.

Proof. Let Φ_2 be the four displayed terms in (II.35). Direct series division gives

$$\Phi_2(f(w)) - \Phi_2(w) - 1 = -w^4/5184 + O(w^5).$$

Sum this defect along the inverse orbit from Section 13. Its absolute total is bounded by $C \sum_{j \geq 0} (|w|^{-1} + j)^{-4} = O(w^3)$. This constructs a normalized Abel coordinate with (II.35). Its

difference from the original normalized coordinate is inverse-map invariant and tends to zero along inverse iterates, hence vanishes. Sector Cauchy estimates justify differentiated remainders. A second finite calculation gives

$$U_0(f(w)) - U_0(w) + \Phi'_2(w)/2 = -w/1080 + O(w^2).$$

Replacing Φ'_2 by Φ' changes this by $O(w^2)$. Both lower pole terms in U_0 are needed, including for cancellation of the constant term. $\square$

[write] **Generalized in Part III, 6 October 2026.** Part III's Lemma 26.1 proves the expansion to every order, with rational a_j and differentiated remainders, by the same summation of the Abel defect along inverse orbits; its $a_1 = -1/36$, $a_2 = 1/540$ are the coefficients here.

Combining the expansions of J, L, M, N gives, when $|w/z|$ is large in the clock cone,

$$\begin{aligned} \mathcal{C}_z(w) &= H(z) + \kappa z + \Phi(w) + z^2 U_0(w) \\ &\quad + O\left(|w|^3 + |z|^2(1 + \log |w/z|) + \frac{|z|^3}{|w|^3} + \frac{|z|^4}{|w|^5}\right). \end{aligned} \quad (\text{II.37})$$

For example, the z^2/w^3 , z^2/w^2 , and z^2/w terms come respectively from J , L , and zM and assemble exactly into $z^2 U_0$. Lower M, N tails are absorbed by the displayed remainder in the small-clock range.

Take

$$K = \lfloor n^{1/2} \rfloor, \quad z \asymp n^{-1}, \quad w = F_{n-K}(z) \asymp K^{-1}. \quad (\text{II.38})$$

The four matching errors in (II.37) have orders at most K^{-3} , $n^{-2} \log n$, $n^{-3} K^3$, and $n^{-4} K^5$; each is $O(n^{-3/2})$. The clock error in (II.34) has the same order. Thus, for $\Theta_z = \Phi + z^2 U_0$,

$$\Theta_z(F_{n-K}(z)) = n - K - H(z) - \kappa z + O(n^{-3/2}). \quad (\text{II.39})$$

The sign of the new phase constant is negative because $+\kappa z$ is moved to the other side of the clock identity.

For an actual late step, use the Abel equation before Taylor expansion: $\Phi(w_{h+1}) = \Phi(w_h + R_h) + 1$. Since $R_h = z^2/2 + O(z^3)$, Lemma 17.1 yields

$$\Theta_z(w_{h+1}) - \Theta_z(w_h) - 1 = O\left(|z|^2 |w_h| + \frac{|z|^3}{|w_h|^2} + \frac{|z|^4}{|w_h|^4}\right). \quad (\text{II.40})$$

Indeed the $z^2 \Phi'/2$ term cancels the leading U_0 difference. The second-order Taylor error for Φ is $O(z^4/w_h^3)$, absorbed by the last term, and perturbing the argument in U_0 gives $O(z^4/w_h^4)$.

18 Uniform local limit on a growing strip

Proposition 18.1. *For every fixed real-width bound C_0 , uniformly at*

$$z = \frac{T}{D_n - s}, \quad |\operatorname{Re} s| \leq C_0, \quad |\operatorname{Im} s| \leq n^{1/20} + 2,$$

one has

$$F_n(z) = P(n - H(z) - \kappa z) + O(n^{-3/2}). \quad (\text{II.41})$$

Proof. Write $M_n = n^{1/20} + 2$ and use K in (II.38). The distance to $r = T/D_n$ is $O(M_n/n^2)$, so the whole parameter set lies in the old entrance disk of radius qK/n^2 . All entrance and new matching estimates are uniform there.

Let $\alpha = n - H(z) - \kappa z$. The exact identity

$$H\left(\frac{T}{D_n - s}\right) = n - s + \frac{1}{3} \log \frac{D_n - s}{n}$$

gives $\alpha = s + O((\log n + M_n)/n)$; its real part stays bounded and its imaginary part differs from $\text{Im } s$ by $o(1)$.

Choose one fixed integer L large relative to C_0 and the inverse-sector threshold. It does not depend on M_n . For $0 \leq j \leq K - L$, set $m = K - j$ and use the prototype phase $\alpha - m$. Its real part is at most $C_0 + 1 - m < 0$ and

$$|\alpha - m| \asymp m + |\text{Im } s|.$$

It lies in a fixed interior inverse sector about the closed left half-plane. A bootstrap phase error at most one preserves fixed angular and radial margins. The corresponding inverse values satisfy

$$|w_j| \asymp (m + |\text{Im } s|)^{-1}. \quad (\text{II.42})$$

The actual $O(n^{-2})$ forcing changes the uncorrected Φ phase by at most $Cn^{-2}(m + M_n)^2$. Its total is

$$O\left(n^{-2} \sum_{m=L}^K (m + M_n)^2\right) = O(K^3/n^2) = o(1),$$

because $M_n = o(K)$. The old entry error in (II.13) is also $o(1)$. These strictly improve the bootstrap for large n . The inverse-identification clause ensures every perturbed point uses the same branch. There is no new injectivity assumption.

The bounds just obtained imply the sums needed for the corrected phase:

$$\sum |w_j| = O(\log K), \quad \sum |w_j|^{-2} = O(K^3), \quad \sum |w_j|^{-4} = O(K^5).$$

Summing (II.40) to time $n - L$ gives

$$O(n^{-2} \log n + n^{-3} K^3 + n^{-4} K^5) = O(n^{-3/2}),$$

of the same order as the error in (II.39). Removing $z^2 U_0$ at time $n - L$ costs $O(n^{-2}(1 + M_n)^3) = O(n^{-37/20})$. Hence

$$\Phi(F_{n-L}(z)) = \alpha - L + O(n^{-3/2}).$$

The inverse derivative is uniformly bounded on the shifted strip, by analyticity on its bounded part and the sector estimate on its unbounded part. We obtain $F_{n-L}(z) = P(\alpha - L) + O(n^{-3/2})$.

The values $P(\alpha - L)$ form a bounded set and tend to zero at large imaginary height. Its closure is compact. The fixed entire composition $f^{\circ L}$ and its derivatives are bounded on a compact enlargement, so the remaining L iterates preserve the error; their subsidiary forcing contributes only $O(n^{-2})$. This proves (II.41). Large imaginary parts remain in the inverse sector, which is why one fixed L suffices. $\square$

18.1 Exponential parameter and explicit remainders

Set $d = -\frac{1}{3} \log n + d_*$, $D_n = n + d$, $r = T/D_n$, and

$$G_n(s) = F_n(re^{s/n}), \quad Q_n(s) = (d + 1/3)s - s^2/2 - d/3 - \kappa T.$$

The exact phase identity is

$$\begin{aligned} n - H(re^{s/n}) - \kappa re^{s/n} &= D_n(1 - e^{-s/n}) - \frac{1}{3} \log(1 + d/n) \\ &\quad + s/(3n) - \kappa T e^{s/n}/(n + d). \end{aligned} \quad (\text{II.43})$$

It implies

$$n - H(re^{s/n}) - \kappa re^{s/n} = s + Q_n(s)/n + R_n(s), \quad |R_n(s)| \leq Cn^{-2}(1 + |d| + |s|)^3. \quad (\text{II.44})$$

The rational parameter for $re^{s/n}$ is $D_n(1 - e^{-s/n})$; this preserves a slightly enlarged fixed-width strip and increases its height by $o(1)$. Both P' and P'' are uniformly bounded on such strips. Taylor expansion in (II.44) has additional error bounded by $Cn^{-2}(1 + |d| + |s|)^4$. On $|\operatorname{Im} s| \leq n^{1/20} + 2$ it is smaller than $n^{-3/2}$. Proposition 18.1 therefore gives

$$G_n(s) = P(s) + \frac{Q_n(s)P'(s)}{n} + O(n^{-3/2}). \quad (\text{II.45})$$

Cauchy's estimate on fixed-radius disks inside the enlarged strip gives the same error for every fixed derivative, in particular for derivative thirty. These degree-three and degree-four bounds avoid an unspecified polynomial whose degree could compromise the growing-strip argument.

19 Derivative extraction and classical Fourier inversion

For every nonnegative integer q , termwise coefficient extraction gives exactly

$$na_n r^n = \frac{1}{2\pi} \int_{-\pi n}^{\pi n} G_n^{(q)}(it) e^{-it} dt. \quad (\text{II.46})$$

Indeed a term of degree k acquires $(k/n)^q$, and orthogonality over this full period selects $k = n$, whose factor is one. Analyticity in a neighborhood of the circle justifies the termwise step. No integration by parts or endpoint assumption is needed.

19.1 Transport of analytic disks

The estimates imported from Report 225 are disk estimates, which permit arbitrary fixed derivative order. Choose a sufficiently small fixed $a > 0$. For $|t| \leq \delta n$,

$$|s - it| \leq a|t| \implies |re^{s/n} - re^{it/n}| \leq re^{a\delta} a|t|/n \leq b|t|/n^2.$$

Thus (II.15) and Cauchy give

$$|G_n^{(30)}(it)| \leq C_{30}|t|^{-31} \quad (M_0 \leq |t| \leq \delta n). \quad (\text{II.47})$$

For the outer angles, the fixed neighborhoods in $\zeta = nz$ admit s disks of radius $a_0 n$ for one fixed small $a_0 > 0$. The bound $G_n = O(n^{-1})$ there gives $G_n^{(30)} = O(n^{-31})$. Disks crossing the endpoints $\pm \pi n$ cause no problem because the exponential parametrization is periodic.

Put $M = n^{1/20}$. The finite-height intermediate and outer integrals outside $[-M, M]$ are

$$O(M^{-30} + n^{-30}) = O(n^{-3/2}).$$

The central integral of the differentiated error in (II.45) is

$$O(Mn^{-3/2}) = O(n^{-29/20}).$$

For the limiting transforms, (II.16) and Leibniz give

$$(Q_n P')^{(30)}(it) = O(|t|^{-30} + \log n |t|^{-31})$$

for large $|t|$. Its omitted tail, divided by n , is smaller still, and that of $P^{(30)}$ is $O(M^{-30})$. Both central integrals can consequently be replaced by full-line integrals with error $O(n^{-7/5})$. The margin between $29/20$ and $7/5$ is deliberate.

19.2 Signs and derivative weights in inversion

Write $\mathcal{I}[V](x) = (2\pi)^{-1} \int_{\mathbb{R}} V(it) e^{-itx} dt$ whenever the integral is absolutely convergent. We only apply it to the differentiated transforms with $q = 30$. At positive x , Fourier inversion and its justified ordinary derivatives give

$$\begin{aligned} \mathcal{I}[P^{(q)}](x) &= x^q h(x), \\ \mathcal{I}[(P')^{(q)}](x) &= x^{q+1} h(x), \\ \mathcal{I}[(sP')^{(q)}](x) &= -(x^{q+1} h)'(x) + qx^q h(x), \\ \mathcal{I}[(s^2 P')^{(q)}](x) &= (x^{q+1} h)''(x) - 2q(x^q h)'(x) + q(q-1)x^{q-1} h(x). \end{aligned} \tag{II.48}$$

For example, $(sP')^{(q)} = sP^{(q+1)} + qP^{(q)}$ and $(s^2 P')^{(q)} = s^2 P^{(q+1)} + 2qsP^{(q)} + q(q-1)P^{(q-1)}$. The t - and t^2 -weighted integrals are absolutely convergent by (II.16); multiplication by s corresponds to minus ordinary differentiation in x for this Laplace sign convention. These identities invoke no endpoint distributions and no regularity at zero.

At $x = 1$, the four inversions in (II.48) are respectively

$$h(1), \quad h(1), \quad -h(1) - h'(1), \quad 2h'(1) + h''(1).$$

In particular the $h(1)$ coefficient in the last expression cancels. Substituting into (II.46) and (II.45) gives

$$\begin{aligned} na_n r^n &= h(1) + \frac{1}{n} \{(-d/3 - \kappa T)h(1) \\ &\quad - (d + 1/3)(h(1) + h'(1)) - h'(1) - h''(1)/2\} + O(n^{-7/5}). \end{aligned} \tag{II.49}$$

Finally,

$$\begin{aligned} r^{-n} &= (n/T)^n e^d \left(1 - \frac{d^2}{2n} + O(\log^4 n/n^2)\right), \\ e^d &= n^{-1/3} T^{1/3} e^{-B}. \end{aligned}$$

Multiplication, division by the positive $h(1)$, and substitution of d give (II.9), hence (II.7) and (II.8). Cross-products and the radius remainder are $O(\log^4 n/n^2)$ and are smaller than the stated error. This completes the proof of Theorem 12.1.

20 Uniform proportional height correction

Proof of Theorem 12.2. Use the height-scaled variables $r = T/(m + d_m)$ and $G_m(s) = F_m(re^{s/m})$. The exact period identity at $\beta = n/m$ is

$$m\beta^q b_{n,m} r^n = \frac{1}{2\pi} \int_{-\pi m}^{\pi m} G_m^{(q)}(it) e^{-i\beta t} dt, \quad q = 30. \quad (\text{II.50})$$

A degree- k term acquires $(k/m)^q$, and the full period selects $k = n$, yielding exactly β^q . All growing-strip and tail estimates depend on height m , not on the target degree. Changing the oscillatory factor has modulus one for real β ; division by β^q is uniformly harmless on I .

Evaluate (II.48) at $x = \beta$ and divide by β^q . The first result is $h(\beta)$; the correction terms become

$$\beta h(\beta), \quad -h(\beta) - \beta h'(\beta), \quad 2h'(\beta) + \beta h''(\beta).$$

For example the third identity simplifies to $-h - xh'$ after division by x^q , and in the fourth the coefficient of h cancels identically, leaving $2h' + xh''$. Hence

$$\begin{aligned} mb_{n,m} r^n &= h(\beta) + \frac{J_\beta(d_m)}{m} + O_I(m^{-7/5}), \\ J_\beta(d) &= (-d/3 - \kappa T)\beta h(\beta) - (d + 1/3)(h(\beta) + \beta h'(\beta)) \\ &\quad - h'(\beta) - \beta h''(\beta)/2. \end{aligned}$$

Since $n = \beta m$ exactly,

$$r^{-n} = (m/T)^n e^{\beta d_m} \left(1 - \frac{\beta d_m^2}{2m} + O_I(\log^4 m/m^2) \right), \quad e^{\beta d_m} = m^{-\beta/3} T^{\beta/3} e^{-\beta B}.$$

Multiplication gives $\mathcal{Q}_\beta = J_\beta/h - \beta d_m^2/2$ in (II.10). Continuity and positivity imply $\min_I h > 0$; the derivatives are bounded on I . This converts the uniform absolute coefficient error into the asserted relative error. At $\beta = 1$ the expression reduces exactly to Theorem 12.1. $\square$

[write] **Checked, 6 October 2026.** The write checked the reduction: $\mathcal{Q}_1(d)$ is (II.9). The theorem refines Part I's Theorem 8.1, whose leading term it contains.

The exact-height formula can be used directly for $m = n + k$, for square-root height windows, or for $m = \lfloor \lambda n \rfloor$ with fixed $\lambda > 0$. In the last case the leading floor phase remains $\exp(-\{\lambda n\}/\lambda)$, as in Report 225. Formula (II.10) retains all such dependence by keeping the exact m and n/m ; replacing them prematurely by their limits can discard contributions as large as the correction itself.

21 A sharpened inverse formula

Set $\alpha = 4/3$, $\ell = \log c$, and

$$Q(u) = -u^2/18 + Au + C.$$

Theorem 12.1 implies

$$\log a_n = n \log(n/T) - \alpha \log n + \ell + \frac{Q(\log n)}{n} + O(n^{-7/5}), \quad (\text{II.51})$$

because the square of the first relative correction is $O(\log^4 n/n^2)$.

Corollary 21.1. *For an exact sequence value $y = a_n$, let $Y = \log y$ and, for large Y , define*

$$X = \frac{Y}{W(Y/T)}, \quad U = \log X, \quad q_X = \log(X/T) + 1, \quad (\text{II.52})$$

where W is the positive real Lambert branch. Put

$$\delta_0 = \frac{\alpha U - \ell}{q_X}, \quad \delta_1 = -\frac{\delta_0^2/2 - \alpha\delta_0 + Q(U)}{q_X}. \quad (\text{II.53})$$

Then

$$n = X + \delta_0 + \frac{\delta_1}{X} + O\left(\frac{X^{-7/5}}{\log X}\right). \quad (\text{II.54})$$

Proof. The definition of W gives $X \log(X/T) = Y$. The leading equivalent first gives $n = X + O(1)$. Define the explicit smooth model

$$F(x) = x \log(x/T) - \alpha \log x + \ell + Q(\log x)/x.$$

For bounded δ , Taylor expansion about X gives uniformly

$$\begin{aligned} F(X + \delta) - Y &= q_X \delta - \alpha U + \ell \\ &+ \frac{\delta^2/2 - \alpha\delta + Q(U)}{X} + O((1 + U^2)/X^2). \end{aligned}$$

Now $\delta_0 = O(1)$ and $\delta_1 = O(U)$, so substituting $\delta = \delta_0 + \delta_1/X$ cancels both retained orders. The residual is $O((1 + U^2)/X^2) = O(X^{-7/5})$. At the true integer n , (II.51) gives $F(n) - Y = O(n^{-7/5})$. On the bounded comparison interval $F'(x) \sim \log X$. The mean value theorem proves (II.54). Only F is differentiated; the asymptotic remainder is used solely at the true integer. $\square$

The constants δ_0, δ_1 in this inverse depend on X and are distinct from the fixed clock constant $d_\star$. The large real solution of $F(x) = \log y$ has the same error at $y = a_n$. With exact constants this yields eventual nearest-integer recovery, but neither an effective onset nor a certificate for arbitrary thresholds between consecutive sequence values. A fixed numerical error in $\log c$ creates an index error of order $1/\log X$, much larger than the fine rate in (II.54); the diagnostic decimal must not be substituted as if exact.

[write] **Relation to Part I and to the transseries volume, 6 October 2026.** δ_0 is the correction of Part I's (33); the write rechecked the two orders (II.53) by expanding $F(X + \delta_0 + \delta_1/X) - Y$. As in Part I, X is an instance of `p0:prop:factorial-core` of [11]; the two Newton-type corrections are proved directly and are not instances of `p0:thm:perturbed-inversion` (front matter).

22 Exact computation and numerical diagnostics

22.1 Exact recurrence and finite checks

Writing $1 + F_{h+1}(z) = \sum_{k \geq 0} c_k z^k$ and logarithmically differentiating the Euler product gives

$$b_j = \sum_{d|j} d[z^d]F_h(z), \quad c_0 = 1, \quad c_k = \frac{1}{k} \sum_{j=1}^k b_j c_{k-j}. \quad (\text{II.55})$$

Degree truncation is exact. The public code checks divisibility before every integer division. An independent finite-product implementation uses the binomial expansion of $(1 - z^d)^{-a}$ and agrees through degree and height twelve. A separately generated exact fixture contains all diagonal values through $n = 414$; only the first twenty-two displayed OEIS terms are an external sequence comparison. The fixture is not presented as a retrieved OEIS b-file.

The code also verifies finite formal-coordinate identities and the new clock algebra exactly over rational symbolic expressions. Such checks protect against algebraic transcription errors; they do not prove analytic remainder estimates, sector entrance, or the all-orders convergence of any formal series. All pass/fail checks use explicit exceptions and remain active under Python optimization. Local chunked decimal serialization handles long exact integers under `PYTHONINTMAXSTRDIGITS=640` without changing the global limit. The implementation caps and tested ranges are stated in the archive README.

22.2 Uncertified amplitude and derivative experiments

The accompanying optional floating-point programs solve a truncated Fatou inverse, take finitely many forward iterates, propagate derivatives, and numerically invert the transform. For example, P''' is the transform of $x^3h(x)$. Its absolutely convergent inversion and its first two ordinary x derivatives give $h(1)$, $h'(1)$, and $h''(1)$. Subtracting $12/(1-s)^4$ improves the observed tail; its inverse is $2x^3e^{-x}$, with values and derivatives at one equal to $2/e$, $4/e$, and $2/e$. After subtraction and quadrature, the derivative weights are undone by

$$h' = (x^3h)'|_1 - 3h(1), \quad h'' = (x^3h)''|_1 - 6h'(1) - 6h(1).$$

These are diagnostic computations with no interval error control for the truncated coordinate, inverse Newton step, forward propagation, finite contour, or quadrature.

Representative numerical values, not certified digit claims, are

$$\begin{aligned} h(1) &\approx 1.81460956, & \rho_1 &\approx -0.43688195, & \rho_2 &\approx -0.15328008, \\ \kappa &\approx -0.25434817, & A &\approx 0.60098418, & C &\approx -0.49816124. \end{aligned}$$

The associated leading amplitude is approximately 4.4923299. At $n = 414$, floating-point normalization of the exact integer gives

$$a_n(T/n)^n n^{4/3} \approx 4.50436880933,$$

while the unfitted first-correction prediction gives approximately 4.50433122939, a difference about 3.76×10^{-5} . No coefficient was fitted to these finite sequence values. Agreement and parameter sensitivity are useful diagnostics, not proofs of a remainder or of displayed digits.

[write] Checks and a comparison, 6 October 2026. At intake, with these constants the corrected prediction matched the exact a_n with residual 1.4×10^{-4} at $n = 200$, against 2.5×10^{-2} for the leading term. With Part I's diagnostic $h(1)$, $2^{1/3}h(1) = 2.2862647816$ agrees to eight significant digits with the iterated Bell report's exploratory amplitude $I(1)$ (Remark 37.1); both values are uncertified.

22.3 Reproducible archive

The archive contains this complete editable TeX, exact programs and tests, optional diagnostic programs and recorded examples, the exact fixture, source provenance, and a deterministic source-preserving builder. It also includes the unchanged Report 225 PDF and TeX as a clearly labeled supporting reference. The new report does not modify that earlier deliverable.

The builder validates its source manifest, runs the exact checks, creates a private pdfL^AT_EX format, compiles without shell escape, rejects overfull boxes, missing glyphs, and unresolved references, checks that source bytes were preserved, and writes a new PDF and sorted fixed-timestamp ZIP to a separate new directory. Ordinary and optimized runs under the restricted digit limit are compared as actual complete ZIPs and member bytes. Identical-toolchain byte reproducibility is distinct from mathematical correctness; different TeX or font versions may change PDF bytes. Every rendered page is also inspected before release.

[write] The companion here, 6 October 2026. Report 226’s own files are shipped as `code/226-firstcorr-*.py`, `data/226-firstcorr-*` and `226-firstcorr-sources-SOURCE_LEDGER.md`. Its eight byte copies of Report 225’s files and its `supporting/` copies of Report 225 are shipped once, as Report 225’s files and as Part I. Its builder and guard tests expect those files beside them; the report README gives the rerun route from the arrival commit.

23 Further questions and the boundary of the theorem

23.1 A genuine all orders expansion

The natural formal expectation is

$$\frac{a_n}{c(n/T)^n n^{-4/3}} \sim 1 + \sum_{r \geq 1} \frac{P_r(\log n)}{n^r}, \quad \deg P_r \leq 2r.$$

Only $P_1(X) = -X^2/18 + AX + C$ is proved here. If an all-orders construction establishes the stated degree bound and proves that no further contribution reaches degree $2r$, the leading coefficient would be

$$[X^{2r}]P_r(X) = \frac{(-1/18)^r}{r!}.$$

In particular the formal second-order highest term would be $\log^4 n / (648n^2)$. This conditional inference is not an all-orders theorem and does not follow from the first correction alone.

A proof needs an induction controlling arbitrary fixed scaled subsidiary order, large-ratio clock expansions and resonant logarithms, forced Fatou cohomological corrections, and growing-strip errors with extraction derivative order chosen for each target precision. The $\log y/540$ in N already shows why clocks cannot be treated as a plain power series in z : a $z^2 \log(1/z)$ contribution can enter the global phase. Exponentiating the radius alone misses this structure.

[write] Still open, 6 October 2026. Not settled by Part III, which works at one polynomial rate in m (Subsection 36.4). The conditional coefficient $(-1/18)^r/r!$ is recorded as unproved in Section 37, item 2.

23.2 Effective constants and certified inversion

Validated values of $h(1)$ and its derivatives require explicit sector remainder bounds, an enclosed inverse branch, controlled forward derivative propagation, interval quadrature, and an integrable certified tail. The present proof supplies qualitative constants and no finite onset. The exact formula for κ is elementary apart from Catalan’s constant, but certifying the full correction still requires the density derivative calculations.

An effective relative-error bound could turn the scalar inverse model into a certified threshold algorithm by combining exact small-index counts with rigorous upper and lower bounds. Theo-

rem 12.1 alone does not decide where a general input lies between a_n and a_{n+1} . One should separate this computational threshold problem from eventual rounding at exact sequence values.

[write] **Still open, 6 October 2026.** Section 37, items 1 and 3.

23.3 Endpoint height regimes and density structure

The compact-positive ratio theorem leaves open $n/m \rightarrow 0$ and $n/m \rightarrow \infty$. These require endpoint information about h and renewed uniform contour estimates. Smooth strict positivity away from zero supplies neither an expansion at zero nor a large-argument asymptotic. The positive-measure identity from Report 225,

$$(e^x - 1)\nu = \sum_{k \geq 2} \nu^{*k}/k!, \quad \nu(dx) = h(x) dx,$$

is a possible starting point, but exchanging endpoint limits or many derivatives with this convolution series requires separate arguments.

A transition theory connecting fixed-degree polynomiality, proportional height, and fixed-height partition asymptotics would also be useful. The forcing calculations here demonstrate that another initial generating function or multiset recurrence cannot simply inherit the same clock constants from the autonomous map. Its subsidiary terms and matching normalization must be recomputed.

[write] **Partly answered by Part III, 6 October 2026.** Part III treats $n/m \rightarrow 0$ at leading order on logarithmic bands (Theorem 31.1) and proves an expansion of h at zero (Proposition 27.1, Theorem 29.1, Corollary 29.2), so “an expansion at zero” is no longer missing. Open: $n/m \rightarrow \infty$ and a large-argument asymptotic of h , the correction on shrinking bands, the transition to fixed degree, other recurrences (Section 37, items 5, 7, 8).

Part III

Logarithmic height crossover for iterated Euler transforms

Bundle Report 228, archive `Report228.zip`, delivered as `src/report228.tex` (17 pages). Delivered title block “Logarithmic height crossover for iterated Euler transforms / Report 228 / 5 October 2026”. Printed as delivered, with `[write]` notes. Delivered Section k is Section $k + 23$ here, its statement $k.j$ is $(k + 23).j$, and its equation (k) is (III. k). Section 37, which closes the report, is added in the write.

Reading conventions of Part III	Elsewhere in the report
“Reports 225 and 226”, [6], [7]	Parts I and II
$m/(n \log n) \rightarrow \lambda$; $L = \log n$; $R(n, m)$ the ratio	Part I: $m = \lfloor \lambda n \rfloor$; $L(y)$ the clock; $R_h(z)$
K a constant; $C_0 = 1 - \gamma - \log 2$; $\mathcal{C}$ Catalan; $C(\beta)$	Parts I, II: K an index; C_0 bounds; Part II: G
the germ	Catalan
$\rho_j \in \mathbb{Q}$, c_j log-density coefficients; a_j Fatou	Part II: ρ_j ratios of derivatives of h ; all: c_k, b_j in
coefficients; b_j germ coefficients	the exact recurrence
$g(t) = t^{t/3}h(t)$; $W(x) = P(-x)$; $\mathcal{G}(u)$	Parts I, II: $g(w) = \log(1 + w)$; W Lambert; Part I:
	$G(u)$
$\tau = -\log r$; χ_n ; $y = m/n$	Part I: τ a time; Parts I, II: $y = w/z$

Abstract of this Part (as delivered). For the iterated Euler-transform array $A(n, m)$ in OEIS A290353, we prove

$$\frac{A(n, m)}{2(2/\pi)^n m^{n-1}} \rightarrow \exp\left(-\frac{1}{3\lambda}\right) \quad \text{if} \quad \frac{m}{n \log n} \rightarrow \lambda \in (0, \infty).$$

The result extends the positive fixed-slope analysis of Reports 225 and 226 to degree-to-height ratios decreasing as any fixed inverse power of the logarithm. We prove an exact small-argument Hankel representation for their normalized positive density h , show that $t^{t/3}h(t)$ has a one-sided smooth germ, and obtain every fixed logarithmic endpoint order with differentiated remainders. A rational coefficient generator isolates a single zeta value at each order of the logarithmic density. Exact cancellation of two logarithms yields every fixed inverse-logarithm order in the crossover window, without $\log \log n$ terms. We also invert the smooth suppression model and quantify rounded integer-height prescriptions and observed-ratio inverses. The exact Kaneiwa leading-monomial normalization changes the first correction constant by +1.

This unrefereed research proof uses explicitly stated analytic interfaces from the two preceding reports, included unchanged in the archive. It does not assert convergence of an infinite endpoint series, an all-orders expansion in $1/m$, an effective onset, discrete monotonicity, or exact integer-height recovery. Numerical comparisons are unvalidated floating diagnostics built from exact finite coefficients.

24 The array and the crossover

Define

$$F_0(z) = z, \quad F_{m+1}(z) = \exp\left(\sum_{j \geq 1} \frac{F_m(z^j)}{j}\right) - 1, \quad A(n, m) = [z^n]F_m(z) \quad (n \geq 1). \quad (\text{III.1})$$

This is the height-indexed array in OEIS A290353 [1]. It counts unlabeled rooted trees whose leaves all have the prescribed height. Equivalently, the new root carries a nonempty unordered multiset of trees from the preceding height, and

$$1 + F_{m+1}(z) = \prod_{d \geq 1} (1 - z^d)^{-A(d,m)}.$$

The generating functions have constant term zero; the tabular convention $A(0, m) = 1$ is imposed separately. There is no factorial normalization.

Put

$$\begin{aligned} T &= \frac{\pi}{2}, & B &= \frac{1}{2} - \frac{\pi}{6} - \frac{\pi^2}{16} - \frac{\log 2}{6}, \\ C_0 &= 1 - \gamma - \log 2, & K &= \frac{1 - \gamma + \log(T/2)}{3} - B, \end{aligned} \quad (\text{III.2})$$

where γ is Euler's constant. Our primary normalization is

$$R(n, m) = \frac{A(n, m)}{2T^{-n}m^{n-1}}. \quad (\text{III.3})$$

All asymptotics with two indices below take $n, m \rightarrow \infty$ in their specified ranges.

Theorem 24.1. *If $m/(n \log n) \rightarrow \lambda \in (0, \infty)$, then*

$$R(n, m) \longrightarrow e^{-1/(3\lambda)}. \quad (\text{III.4})$$

More precisely, let $\Lambda \subset (0, \infty)$ be compact, put $L = \log n$, and set $m = \lfloor \lambda n L \rfloor$. For every fixed $N \geq 1$, uniformly for $\lambda \in \Lambda$,

$$\log R(n, m) = -\frac{1}{3\lambda} + \frac{K}{\lambda L} + \sum_{j=2}^N \frac{c_j}{(\lambda L)^j} + O_{\Lambda, N}(L^{-N-1}), \quad (\text{III.5})$$

where the sum is empty for $N = 1$, and

$$c_j = \rho_j - \frac{\zeta(j)}{j3^j}, \quad \rho_j \in \mathbb{Q} \quad (j \geq 2). \quad (\text{III.6})$$

A finite coefficient algorithm determines every ρ_j . In particular,

$$c_2 = \frac{9 - \pi^2}{108}, \quad c_3 = \frac{17}{1080} - \frac{\zeta(3)}{81}.$$

For example, exponentiation gives

$$R(n, m) = e^{-1/(3\lambda)} \left\{ 1 + \frac{K}{\lambda L} + \frac{K^2/2 + (9 - \pi^2)/108}{\lambda^2 L^2} + O_{\Lambda}(L^{-3}) \right\}. \quad (\text{III.7})$$

Every order here means a fixed finite truncation followed by its asymptotic limit. It says neither that the infinite series converges nor that there is an all-orders expansion in powers of m^{-1} . One polynomial rate in the coefficient-transfer error will suffice for all fixed inverse-logarithm orders.

The smooth suppression-ratio inverses are proved in Section 34. With $\tau = -\log r$ and $\chi_n = (\log n)/3 - K$, their first height prescription is

$$\frac{m}{n} \approx \frac{\chi_n}{\tau} - \frac{c_2}{\chi_n} - \frac{c_3\tau}{\chi_n^2} - \frac{c_2^2\tau + c_4\tau^2}{\chi_n^3}. \quad (\text{III.8})$$

The inverse concerns a smooth asymptotic ratio; no unique discrete solution, least-height threshold, or exact recovery by rounding is claimed.

[write] Reading Part III against Parts I and II, 6 October 2026. “Report 225”, [6] and “Report 226”, [7] are Parts I and II; their included copies are byte-identical and not shipped. Here λ measures $m/(n \log n)$, not Part I’s slope m/n of Corollary 8.2; K is a constant, not Parts I and II’s entrance index; ρ_j are rational numbers, not Part II’s $h^{(j)}(1)/h(1)$; $L = \log n$, not the clock function.

25 Imported analytic interfaces and the scope of the result

The analytic foundation comes from Reports 225 and 226 [6, 7]. Their unchanged complete TeX and PDF files accompany this report, with hashes in the source ledger. We state exactly what is used; the present report supplies the endpoint, shrinking-ratio, and inverse arguments in full.

25.1 The normalized coordinate and density

Report 225, Sections 4 and 6, constructs an entire function P with

$$P(s+1) = e^{P(s)} - 1, \quad \Phi(w) = -\frac{2}{w} + \frac{1}{3} \text{Log } w + O(w). \quad (\text{III.9})$$

The coordinate Φ is analytic on every sufficiently small closed proper subsector of $|\arg w| < \pi$, with compatible branches on overlaps. It has zero additive constant in (III.9). The local inverse agrees with P on every sufficiently distant proper subsector of $|\arg(-s)| < \pi$; in particular

$$P(\Phi(w)) = w, \quad P'(s) = \frac{2}{s^2} + O\left(\frac{\log |s|}{|s|^3}\right) \quad (\text{III.10})$$

there. For the reciprocal inverse iteration

$$\mathcal{G}(u) = \frac{2}{\text{Log}(1 + 2/u)} = u + 1 - \frac{1}{3u} + O(u^{-2}), \quad (\text{III.11})$$

the normalized coordinate satisfies

$$\Psi(\mathcal{G}(u)) = \Psi(u) + 1, \quad \Psi(u) = u + \frac{1}{3} \text{Log } u + O(u^{-1}), \quad \Phi(w) = -\Psi(2/w) + \frac{1}{3} \log 2.$$

On nested proper sectors its inverse orbits satisfy $|\mathcal{G}^{\circ j}(u)| \geq c(|u| + j)$, and the normalized orbit limit is available on these sectors. These sector estimates, rather than a solely real-axis expansion, are used below.

The same report proves the positive Laplace representation

$$P(s) = \int_0^\infty e^{st} h(t) dt, \quad h \in C^\infty(0, \infty), \quad h(t) > 0 \ (t > 0). \quad (\text{III.12})$$

The density has every exponential moment. Fourier inversion applies to P' on vertical lines and to the fixed higher derivatives used below. We do not import endpoint smoothness or an endpoint expansion for h .

[write] **Where Part I proves this, 6 October 2026.** “Report 225, Sections 4 and 6” are Part I’s Sections 4 and 6: Lemma 4.1 (whose proof constructs $G = \mathcal{G}$, Ψ and $\Phi = -\Psi(2/w) + \frac{1}{3}\log 2$, with the orbit bound $|G^j(u)| \geq c(|u| + j)$), Proposition 6.1 and Corollary 6.2; Fourier inversion of P' is (30).

25.2 The absolute derivative estimates before division

For large m , set

$$\begin{aligned} d_m &= -\frac{1}{3}\log m + \frac{1}{3}\log T - B, & r_m &= \frac{T}{m + d_m}, \\ G_m(s) &= F_m(r_m e^{s/m}), & Q_m(s) &= (d_m + \frac{1}{3})s - \frac{1}{2}s^2 - \frac{1}{3}d_m - \kappa T. \end{aligned} \quad (\text{III.13})$$

Here κ is the fixed real constant of Report 226. Its value is immaterial for the transfer in this report; for a fully specified interface it is

$$\kappa = \frac{\pi^3}{96} + \frac{\pi^2}{12} - \frac{13\pi}{48} - \frac{11}{36} - \frac{\mathcal{C}}{6} - \frac{\pi \log 2}{24}, \quad \mathcal{C} = \sum_{j \geq 0} \frac{(-1)^j}{(2j+1)^2}.$$

Report 226, Sections “Derivative extraction and classical Fourier inversion” and “Uniform proportional height correction,” proves the following absolute estimates, all depending on height m , not on the target degree. With $M = m^{1/20}$, the thirtieth derivative of

$$G_m(s) - P(s) - m^{-1}Q_m(s)P'(s)$$

is $O(m^{-3/2})$ on $s = iu$, $|u| \leq M$, by analytic disk estimates on an enlarged strip. The finite-height intermediate derivative is $O(|u|^{-31})$, and the outer derivative is $O(m^{-31})$ on an interval of length $O(m)$. Their absolute tails are

$$O(M^{-30} + m^{-30}) = O(m^{-3/2}). \quad (\text{III.14})$$

The limiting derivative has $P^{(30)}(iu) = O(|u|^{-31})$, and

$$(Q_m P')^{(30)}(iu) = O(|u|^{-30} + \log m |u|^{-31}) \quad (\text{III.15})$$

for large $|u|$. No degree-to-height division has yet occurred.

Write $\tilde{G}_{m,30}(u)$ for $G_m^{(30)}(iu)$ on $[-\pi m, \pi m]$ and zero elsewhere. This notation means extension of the derivative’s values, not differentiation of a zero extension. Taking absolute values before integrating the preceding bounds proves

$$\left\| \tilde{G}_{m,30} - P^{(30)}(i \cdot) - \frac{(Q_m P')^{(30)}(i \cdot)}{m} \right\|_{L^1(\mathbb{R})} = O(m^{-7/5}). \quad (\text{III.16})$$

Indeed the central error is $O(Mm^{-3/2}) = O(m^{-29/20})$; (III.14) and the limiting P tail are $O(m^{-3/2})$; the correction tail after division by m is

$$O(m^{-1}(M^{-29} + \log m M^{-30})),$$

which is smaller still. These estimates even give the exponent 29/20 here; the weaker 7/5 is convenient and already sufficient. Equation (III.16) is an absolute L^1 consequence of the proved estimates, not an extrapolation of Report 226’s compact-slope relative theorem.

[write] **Where Part II proves this, 6 October 2026.** Report 226’s sections “Derivative extraction and classical Fourier inversion” and “Uniform proportional height correction” are Part II’s Sections 19 and 20: (II.45), (II.47), the outer-angle bound and the tail estimates following it, there at height n and here at height m (Part II’s proof of Theorem 12.2 makes the same change). d_m and Q_m are Part II’s d_m and Q_n with n replaced by m . The L^1 bound (III.16) itself is derived here.

25.3 Classical background and attribution

The endpoint calculation belongs to the universal unforced parabolic map $w \mapsto e^w - 1$. Finite sectorial Fatou expansions, their rational formal recursion, and Hankel inversion are classical methods. Dudko and Sauzin [8] give general sectorial asymptotics, including nonzero resiter; the present reciprocal germ has resiter 1/3. We give a direct finite-defect proof adapted to the precise normalization above.

The public iterated Bell report [5] already contains all-finite-order inverse-coordinate machinery and a related contour amplitude. Its inspected proportional-depth transfer fixes n/m in a positive compact interval and explicitly does not assert endpoint uniformity. In its notation, renamed here to avoid collisions, the small-variable coordinate is

$$\Psi_{\text{IBD}}(w) = -\Phi(w) + \frac{1}{3} \log 2.$$

This local identity follows from the common normalized reciprocal orbit limit. A formal contour match suggests an amplitude relation $I(\beta) = \beta 2^{\beta/3} h(\beta)$, but that identification requires its own analytic contour correspondence and is not used here. We neither attribute an endpoint theorem to that source nor claim a new array-specific origin for the universal coefficient calculus.

Prellberg’s 2002 seminar, as summarized by Mishna [9], already presents the parabolic logarithmic mechanism and its 1/3 specialization. Nagaev and Vakhel [10] give an important unforced antecedent: their critical branching large-deviation correction has the form $\exp[-\gamma u \log u/m]$, with $\gamma = 1/3$ in the Poisson case, in an appropriate moderate-deviation range. The heuristic high-moment saddle $u \asymp n$ places the logarithmic crossover in that range when $m/n \rightarrow \infty$. Passing from those probability estimates to a full moment equivalent needs tail domination, which is not supplied by this comparison; they do not furnish the forced Euler coefficient transfer proved here. The new scope relative to Report 225 is the quantitative transfer as n/m decreases to logarithmic scales, using the forced-array contour estimates and its constants T, B . No worldwide priority claim is made for the parabolic mechanism or universal endpoint calculus.

[write] **The iterated Bell report, 6 October 2026.** [5] is `a139383-iterated-bell-diagonals` in this repository; its article is unchanged since the pin, so the sections named are its current Sections 2, 4 (Part I there) and 12, 13, 17 (Part II there). Its Ψ is Ψ_{IBD} here and its $I(\beta)$ is the amplitude of its Theorem `ibd:pd:thm:main`. The relation $I(\beta) = \beta 2^{\beta/3} h(\beta)$ is unproved; at $\beta = 1$ the two sides agree to eight significant digits (Remark 37.1), and Section 37, item 9, records it as a question shared with that report, with a conditional consequence for its open endpoint (Remark 37.2). That report’s own literature section places the Nagaev–Vakhel range outside its compact-slope regime $m \asymp n$; here $m/n \rightarrow \infty$, which is the range named above.

26 Finite Fatou expansions with sectorial remainders

Lemma 26.1. *For every integer $N \geq 0$ there are unique rational numbers $a_1, \dots, a_N$ such that*

$$\Phi(w) = -\frac{2}{w} + \frac{1}{3} \operatorname{Log} w + \sum_{j=1}^N a_j w^j + O(w^{N+1}) \quad (\text{III.17})$$

uniformly on each sufficiently small closed proper sector. For every fixed derivative order k , the k th derivative of the remainder is $O(|w|^{N+1-k})$ on smaller closed sectors. The first values are

$$a_1 = -\frac{1}{36}, \quad a_2 = \frac{1}{540}, \quad a_3 = \frac{1}{7776}, \quad a_4 = -\frac{71}{435456}.$$

Proof. Choose a finite reciprocal model

$$V_N(u) = u + \frac{1}{3} \operatorname{Log} u + \sum_{j=1}^N v_j u^{-j}.$$

The expansion of (III.11) has rational coefficients. In $V_N(\mathcal{G}(u)) - V_N(u) - 1$, the new coefficient v_j first contributes $-j v_j u^{-j-1}$. Cancellation is therefore triangular, unique and rational, and gives

$$D_N(u) := V_N(\mathcal{G}(u)) - V_N(u) - 1 = O(u^{-N-2}).$$

Along an inverse orbit, $V_N(u) - u - (\operatorname{Log} u)/3 \rightarrow 0$, so the normalized orbit limit is the same Ψ as in Section 25. Telescoping gives the sign-sensitive identity

$$\Psi(u) - V_N(u) = \sum_{j \geq 0} D_N(\mathcal{G}^{\circ j}(u)). \quad (\text{III.18})$$

The sign is positive: the finite sum equals $V_N(\mathcal{G}^{\circ J}(u)) - J - V_N(u)$ before taking $J \rightarrow \infty$. The orbit bound gives absolute convergence and

$$\sum_{j \geq 0} |D_N(\mathcal{G}^{\circ j}(u))| \leq C \sum_{j \geq 0} (|u| + j)^{-N-2} = O(|u|^{-N-1}).$$

Transforming by $\Phi(w) = -\Psi(2/w) + (\log 2)/3$ proves (III.17); thus $a_j = -v_j/2^j$. Cauchy's derivative estimates on disks of radius proportional to $|w|$ inside nested sectors give every fixed differentiated remainder. Direct triangular substitution gives the stated coefficients. No convergence of the infinite formal coordinate series is required. $\square$

27 The endpoint value by direct Fourier inversion

The leading endpoint value follows already from the first-order interface in Report 225.

Proposition 27.1. *The density extends continuously to zero with*

$$h(t) = 2 + O(t \log(1/t)), \quad h(0) = 2. \quad (\text{III.19})$$

Proof. Fourier inversion of P' on the line $\operatorname{Re} s = -1/t$ gives

$$th(t) = \frac{e}{2\pi} \int_{\mathbb{R}} P'(-1/t + iu) e^{-iut} du.$$

The entire line is in the fixed inverse sector $|\arg(-s)| \leq \pi/2$. The leading term $2/s^2$ in (III.10) inverts to $2t$. The absolute error is bounded by a constant times

$$\int_{\mathbb{R}} \frac{\log \sqrt{t^{-2} + u^2}}{(t^{-2} + u^2)^{3/2}} du = O(t^2 \log(1/t)),$$

by $u = v/t$. Division by t proves the assertion. $\square$

[write] **Answers Parts I and II, 6 October 2026.** This answers, at the endpoint zero, the remark closing the proof of Part I's Proposition 6.1 and the questions of Subsections 11.4 and 23.3 that ask for an expansion of h at zero.

28 An exact Hankel representation

Set

$$\mathcal{A}(w) = \Phi(w) + 2/w - \frac{1}{3} \operatorname{Log} w = O(w). \quad (\text{III.20})$$

Fix $\pi/2 < \theta < \pi$ and $a > 0$. The contour $H = H_{\theta,a}$ starts at infinity on argument $-\theta$, travels inward to $ae^{-i\theta}$, follows the circle $|v| = a$ counterclockwise through the positive axis to $ae^{i\theta}$, and leaves along argument θ . All logarithms in the contour formula use the principal branch.

Theorem 28.1. *For sufficiently small positive t ,*

$$h(t) = \frac{2(2t)^{-t/3}}{2\pi i} \int_H e^v v^{t/3-2} \exp[-t\mathcal{A}(2t/v)] dv. \quad (\text{III.21})$$

The integral is absolutely convergent and is an exact identity. It supplies analytic continuation to a sufficiently small punctured sector about positive t .

Proof. Let $W(x) = P(-x)$. With an upward vertical contour, the absolutely convergent inverse of P' is

$$th(t) = -\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} W'(x) e^{tx} dx, \quad c > 0. \quad (\text{III.22})$$

To check the sign, replacing s by $-x$ reverses the vertical orientation and $P'(-x) = -W'(x)$. Equivalently, change u to $-u$ in the absolutely convergent Fourier formula. We take $c = 1/t$.

Choose $\theta < \theta_1 < \pi$. For small t , the line and scaled contour $x = v/t$ lie beyond the inverse threshold in the sector $|\arg x| \leq \theta_1$, apart from no problematic compact portion since $|v| \geq a$. Truncate both at common large x -radius R . The circular connectors from the vertical endpoints to the rays lie to the left of the vertical line, so $|e^{tx}| \leq e$ there. The estimate $W'(x) = O(R^{-2})$ and connector length $O(R)$ give integral $O(R^{-1})$. Thus these connectors vanish as $R \rightarrow \infty$. The vertical integral converges absolutely by $W' = O(|x|^{-2})$, and the ray integrals by $\cos \theta < 0$. Cauchy's theorem deforms the upward line to the stated lower-inward, upper-outward Hankel orientation.

For the second deformation put

$$\begin{aligned} D_t(v) &= \frac{1}{3}(\text{Log } v - \log(2t)) - \mathcal{A}(2t/v), \\ x_\ell(v) &= v/t + \ell D_t(v), \quad 0 \leq \ell \leq 1. \end{aligned}$$

Uniformly for $|v| \geq a$ on H ,

$$|D_t(v)| \leq C(1 + |\log |v|| + |\log t|), \quad \sup_{v \in H} |tD_t(v)/v| \longrightarrow 0.$$

Consequently this homotopy stays in the larger proper inverse sector and satisfies $|x_\ell(v)| \geq |v|/(2t)$ for small t . Its endpoint at $\ell = 1$ is exactly

$$x(v) = -\Phi(2t/v) = v/t + \frac{1}{3}(\text{Log } v - \log(2t)) - \mathcal{A}(2t/v). \quad (\text{III.23})$$

The compatible principal branches are valid since $\arg(2t/v) = -\arg v$ stays strictly inside $(-\pi, \pi)$. In particular no negative-real branch cut is crossed.

At ray endpoints $v = Re^{\pm i\theta}$ the homotopy connectors have length $O(\log R + |\log t|)$ and

$$\text{Re}(tx_\ell(v)) = R \cos \theta + O(t \log R + t |\log t|).$$

Their integrals tend to zero exponentially as $R \rightarrow \infty$, for fixed small t . On each finite truncated homotopy the integrand $W'(x)e^{tx}$ is entire, so Cauchy's theorem applies. This is a deformation of paths; it does not require global univalence of the map $v \mapsto x(v)$.

By the imported inverse identification,

$$W(x(v)) = P(\Phi(2t/v)) = 2t/v, \quad W'(x(v)) dx = -2tv^{-2} dv.$$

Substitution into (III.22), retaining its minus sign, gives (III.21), including the factor 2. As an orientation check, $(2\pi i)^{-1} \int_H e^v v^{-2} dv = 1$, which gives the positive endpoint value.

Finally choose $\eta > 0$ with $\eta + \theta < \pi$. For complex t with $0 < |t| < \epsilon$ and $|\arg t| < \eta$, every $2t/v$ stays in a fixed coordinate sector. Use the corresponding branch of $\text{Log } t$ in (III.21). Uniform convergence on compact subsets follows because $e^{\text{Re } v}$ on the rays dominates all fixed polynomial and logarithmic factors. The resulting holomorphic function agrees with the real density for positive t by the identity already proved. $\square$

[write] **A consequence, 6 October 2026.** Since the continuation is holomorphic on a punctured sector containing $(0, \epsilon)$ and agrees with h there, h is real analytic on $(0, \epsilon)$. Real analyticity on all of $(0, \infty)$, asked in Part I's Subsection 11.4, is not shown (Section 37, item 6).

29 A smooth endpoint germ and the logarithmic transseries

Here “endpoint transseries” refers only to finite asymptotic expansions in powers $t^j(\log t)^r$ and their differentiated remainders. No summability, exponentially small completion, or convergence assertion is attached to the term.

Theorem 29.1. *Let $g(t) = t^{t/3}h(t)$ for $t > 0$. There are real coefficients b_j , with $b_0 = 2$, such that for every fixed $N, k \geq 0$,*

$$\frac{d^k}{dt^k} \left(g(t) - \sum_{j=0}^N b_j t^j \right) = O(t^{N+1-k}) \quad (t \downarrow 0). \quad (\text{III.24})$$

Consequently g has a one-sided C^∞ extension to $[0, \epsilon)$ and $b_j = g^{(j)}(0)/j!$. This does not assert analyticity at zero.

Proof. Cancel the explicit $t^{-t/3}$ in (III.21) to obtain

$$g(t) = \frac{2 \cdot 2^{-t/3}}{2\pi i} \int_H e^v v^{-2} \exp \left[\frac{t}{3} \operatorname{Log} v - t\mathcal{A}(2t/v) \right] dv. \quad (\text{III.25})$$

Use Lemma 26.1 to any required finite order. On H , $|v| \geq a$, and the remainder after retaining $\mathcal{A}$ through degree q is uniformly $O(|t|^{q+1}|v|^{-q-1})$. Multiplication by t improves its order once more. The remaining factors $2^{-t/3}$ and $v^{t/3}$ are entire in t . Their finite Taylor remainders are bounded by fixed powers of $1 + |\operatorname{Log} v|$ times a small fixed power of $|v|$. Along the rays these are integrably dominated by $e^{\operatorname{Re} v}$; the inner arc is compact and avoids zero. Finite expansion and integration thus yield, uniformly in a smaller punctured t -sector,

$$g(t) = \sum_{j=0}^N b_j t^j + O(|t|^{N+1}).$$

Apply Cauchy's estimate to the actual holomorphic remainder on disks of radius δt centered at positive t . This proves (III.24), rather than formally differentiating an unsupported pointwise expansion. For a prescribed derivative order choose $N \geq k$; its limit is $k!b_k$. Choosing one additional order establishes compatibility of these endpoint derivatives. The constant follows from Proposition 27.1, or directly from the orientation check in Theorem 28.1. $\square$

The classical reciprocal-Gamma Hankel identity, on the orientation specified above, is

$$\frac{1}{2\pi i} \int_H e^v v^{-z} dv = \frac{1}{\Gamma(z)}.$$

For a finite nonnegative multi-index $\mathbf{r} = (r_1, r_2, \dots)$, let $R_{\mathbf{r}} = \sum r_j$ and $J_{\mathbf{r}} = \sum j r_j$. Expanding the last exponential in (III.25) gives the finite-order generator

$$g(t) \sim 2 \cdot 2^{-t/3} \sum_{\mathbf{r}} \frac{(-1)^{R_{\mathbf{r}}} 2^{J_{\mathbf{r}}}}{\Gamma(2 + J_{\mathbf{r}} - t/3)} \left(\prod_j \frac{a_j^{r_j}}{r_j!} \right) t^{R_{\mathbf{r}} + J_{\mathbf{r}}}. \quad (\text{III.26})$$

At any specified degree only finitely many multi-indices occur. Equation (III.26) is interpreted by truncation; Theorem 29.1 proves its remainder, and no convergence of the full multi-index sum is used.

Corollary 29.2. *Writing $X = \log t$, for every fixed $N \geq 0$,*

$$h(t) = \sum_{j=0}^N t^j H_j(X) + O(t^{N+1}(1 + |X|)^{N+1}), \quad H_j(X) = \sum_{r=0}^j b_{j-r} \frac{(-X/3)^r}{r!}. \quad (\text{III.27})$$

The k th derivative of its remainder is $O(t^{N+1-k}(1 + |\log t|)^{N+1})$ for every fixed k . In particular H_j has degree exactly j , with leading coefficient $2(-1/3)^j/j!$.

Proof. Use $h(t) = \exp[-t \operatorname{Log} t/3]g(t)$ in the punctured sector and multiply finite expansions. The exponential remainder obeys the stated power-log bound. Cauchy estimates on the same proportional disks give the differentiated bound; equivalently one may differentiate the finite product with its controlled remainder. Since $b_0 = 2$, its highest logarithmic coefficient is the stated nonzero number. $\square$

In particular,

$$h(t) = 2 + \frac{2}{3}t(C_0 - \log t) + \frac{t^2}{9} \left((\log t - C_0)^2 + \frac{3}{2} - \frac{\pi^2}{6} \right) + O(t^3(1 + |\log t|)^3). \quad (\text{III.28})$$

The differentiated estimates imply

$$h'(t) = -\frac{2}{3}(\log t + \gamma + \log 2) + O(t \log^2 t), \quad h''(t) = -\frac{2}{3t} + O(\log^2 t).$$

Thus the original density h is not C^1 at zero, although the explicitly renormalized germ g is smooth.

30 The rational and zeta coefficient structure

Because $g(t) = 2 + O(t)$ on a sector, a branch of $\log(g(t)/2)$ exists near the endpoint, approaching zero. Theorem 29.1 gives ordinary finite Taylor remainders for this logarithm and each fixed derivative. Therefore

$$\log \frac{h(t)}{2} = -\frac{t}{3} \log t + \frac{C_0}{3}t + \sum_{j=2}^N c_j t^j + O(t^{N+1}). \quad (\text{III.29})$$

After removing the single explicit $t \log t$ term, every k th derivative of the remainder is $O(t^{N+1-k})$.

Proposition 30.1. *The coefficients in (III.29) have the form (III.6). Their first rational parts are*

$$\begin{aligned} \rho_2 &= \frac{1}{12}, & \rho_3 &= \frac{17}{1080}, & \rho_4 &= \frac{13}{3888}, \\ \rho_5 &= \frac{2071}{2449440}, & \rho_6 &= \frac{5131261}{22044960000}, & \rho_7 &= \frac{261791507}{3968092800000}. \end{aligned}$$

Proof. Factor $1/\Gamma(2 - t/3)$ out of (III.26). The remaining formal ordinary series $S(t)$ has rational coefficients and constant term one, because each Gamma ratio is exactly

$$\frac{\Gamma(2 - t/3)}{\Gamma(2 + J - t/3)} = \prod_{k=2}^{J+1} (k - t/3)^{-1},$$

with empty product one when $J = 0$. Thus

$$\frac{g(t)}{2} \sim \frac{2^{-t/3}}{\Gamma(2 - t/3)} S(t), \quad \log S(t) \in t^2 \mathbb{Q}[[t]]. \quad (\text{III.30})$$

At each finite order this is justified by the already proved expansion. The familiar local Gamma expansion yields

$$-\log \Gamma(2 - t/3) = \frac{1 - \gamma}{3}t + \sum_{j \geq 2} \frac{1 - \zeta(j)}{j 3^j} t^j.$$

Combining it with (III.30) proves the rational/zeta form and the linear coefficient $C_0/3$. For example,

$$S(t) = 1 + \frac{t^2}{36} + \frac{11t^3}{3240} + O(t^4),$$

which immediately gives c_2 and c_3 . Finite multi-index enumeration with the rational a_j gives the further displayed ρ_j . The archive supplies exact-arithmetic reproduction of these identities, including a separate finite reversion check. The analytic remainders follow from the preceding proofs, not from symbolic checking. $\square$

31 Transfer to shrinking degree to height ratios

Theorem 31.1. *Uniformly over positive integer degrees n as $m \rightarrow \infty$, writing $\beta = n/m$, one has the absolute estimate*

$$mA(n, m)r_m^n = h(\beta) + O\left(\frac{\log m}{m\beta^{30}}\right). \quad (\text{III.31})$$

For every fixed $B_0 > 0$, uniformly for $0 < \beta \leq B_0$,

$$\begin{aligned} A(n, m) &= \frac{h(\beta)}{m} (m/T)^n e^{\beta d_m} \{1 + O(E(m, \beta))\}, \\ E(m, \beta) &= \frac{\log m}{m\beta^{30}} + \frac{\beta \log^2 m}{m}. \end{aligned} \quad (\text{III.32})$$

The relative expression is an asymptotic equivalent whenever its stated error tends to zero. In particular this holds uniformly on every fixed logarithmic band

$$(\log m)^{-D} \leq \beta \leq B_0, \quad D > 0 \text{ fixed}. \quad (\text{III.33})$$

Proof. Cauchy's estimates for the inverse-sector expansion of P give $P^{(q)}(iu) = O(|u|^{-q-1})$ at infinity for every fixed $q \geq 1$. Leibniz' rule, the quadratic form of Q_m , and $d_m = O(\log m)$ imply

$$\|(Q_m P')^{(30)}(i \cdot)\|_1 = O(\log m).$$

The bounded interval uses analyticity; on the tails (III.15) is integrable. Hence the pre-division estimate (III.16) implies

$$\|\tilde{G}_{m,30} - P^{(30)}(i \cdot)\|_1 = O(\log m/m). \quad (\text{III.34})$$

For every positive integer n , termwise full-period extraction is exact:

$$m\beta^{30} A(n, m)r_m^n = \frac{1}{2\pi} \int_{-\pi m}^{\pi m} G_m^{(30)}(iu) e^{-i\beta u} du. \quad (\text{III.35})$$

Indeed a monomial of degree k acquires $(k/m)^{30}$; orthogonality over a period selects $k = n$, leaving β^{30} . The contour is inside the disk of analyticity for large m . No integration by parts or endpoint matching assumption is needed.

Fourier inversion of $P^{(30)}$ gives $\beta^{30}h(\beta)$ for each $\beta > 0$. The oscillatory factor has absolute value one, so (III.34) and division by β^{30} give (III.31). The constant before that division is independent of the target degree. The displayed β^{-30} loss is therefore genuine and has not been discarded. Only this absolute assertion is globally uniform in β .

By Proposition 27.1, strict positivity, and continuity, $\min_{[0, B_0]} h > 0$. Thus division by $h(\beta)$ is uniform on this interval. Furthermore

$$\begin{aligned} r_m^{-n} &= (m/T)^n \exp\{n \log(1 + d_m/m)\} \\ &= (m/T)^n e^{\beta d_m} \left\{1 + O\left(\frac{\beta \log^2 m}{m}\right)\right\} \end{aligned}$$

uniformly for $\beta \leq B_0$. This proves (III.32). On (III.33),

$$E(m, \beta) = O\left(\frac{(\log m)^{30D+1}}{m} + \frac{\log^2 m}{m}\right) = o(1).$$

The constants may depend on B_0, D , and the fixed derivative order. There is no uniform relative theorem for arbitrary $\beta \downarrow 0$ here. $\square$

[write] **Relation to Parts I and II, 6 October 2026.** On a fixed compact β -range (III.32) is Part I's (31) with the rate $O(\log m/m + \log^2 m/m)$ in place of $o(1)$; Part II's Theorem 12.2 is sharper there. The new content is the uniformity down to $(\log m)^{-D}$.

32 Exact logarithmic cancellation and the crossover proof

Combining (III.3) and (III.32) gives

$$R(n, m) = \frac{h(\beta)}{2} e^{\beta d_m} \{1 + O(E(m, \beta))\}.$$

In a logarithmic band with sufficiently small upper endpoint, its logarithm is legitimate because $E = o(1)$ and $h > 0$. Equation (III.29) then gives, for every fixed $N \geq 1$,

$$\log R(n, m) = -\frac{\beta}{3} \log n + K\beta + \sum_{j=2}^N c_j \beta^j + O(\beta^{N+1} + E(m, \beta)). \quad (\text{III.36})$$

The identity behind this formula is exact:

$$-\frac{\beta}{3} \log \beta - \frac{\beta}{3} \log m = -\frac{\beta}{3} \log n, \quad \log \beta + \log m = \log n.$$

It removes potential $\log \log n$ terms before any asymptotic substitution.

Proof of Theorem 24.1. If $m/(n \log n) \rightarrow \lambda \in (0, \infty)$, then $\beta \rightarrow 0$, $\beta \log n \rightarrow 1/\lambda$, and $\log m \sim \log n$. The ratio belongs, for example, to a fixed logarithmic band with $D = 2$ for all sufficiently large n , and $E \rightarrow 0$. Equation (III.36) gives (III.4).

For the uniform fixed-order assertion let $m = \lfloor \lambda n L \rfloor$, $L = \log n$, and $\lambda \in \Lambda$. Then

$$\beta = \frac{1}{\lambda L} + O_\Lambda \left(\frac{1}{n L^2} \right), \quad E(m, \beta) = O_\Lambda(L^{30}/n).$$

Integer rounding changes βL by $O_\Lambda(1/(nL))$. These errors are smaller than every fixed inverse power of L . Substitute the finite expansion (III.36); its endpoint remainder becomes $O_{\Lambda, N}(L^{-N-1})$. This proves (III.5) and, by ordinary finite exponentiation, (III.7). No estimate of arbitrarily high order in m^{-1} was used. $\square$

More generally, whenever $\beta \rightarrow 0$ inside a fixed logarithmic band,

$$R(n, m) = \exp \left[-\frac{n}{3m} \log n \right] \{1 + O(\beta + E(m, \beta))\}. \quad (\text{III.37})$$

Thus $R \rightarrow 1$ if $m/(n \log n) \rightarrow \infty$ and m/n grows at most as a fixed power of $\log n$. Also $R \rightarrow 0$ if $m/n \rightarrow \infty$ while $m/(n \log n) \rightarrow 0$; this range lies in a logarithmic band because $m/n = o(\log n)$. Fixed degree followed by height tending to infinity is a different limit and is not covered by this transfer.

33 Kaneiwa normalization and its first correction

Kaneiwa [3] proves that his $p(r; n)$ is a degree- $(n - 1)$ polynomial in r with leading coefficient

$$\ell_n = \frac{E_{n-1}}{(n-1)!} = [x^{n-1}](\tan x + \sec x). \quad (\text{III.38})$$

His indexing is $p(r; n) = A(n, r + 1)$, so $r = m - 1$. His fixed-degree polynomial theorem alone provides no uniform error as n and r grow together.

The nearest pole of $\tan x + \sec x$ is $T = \pi/2$, with principal part $2/(T - x)$. The apparent singularity at $-T$ is removable; the next pole is $-3T$. Subtract the principal part and use Cauchy's coefficient estimate on any fixed circle of radius strictly between T and $3T$. For every fixed $1 < c < 3$ this gives

$$\frac{\ell_n}{2T^{-n}} = 1 + O(c^{-n}). \quad (\text{III.39})$$

If $n \rightarrow \infty$ and $\beta = n/m \rightarrow 0$, then

$$(n - 1) \log(1 - 1/m) = -\beta + O(1/m + n/m^2) \rightarrow 0.$$

Consequently the crossover limit is unchanged when (III.3) is replaced by

$$R_K(n, m) = \frac{A(n, m)}{\ell_n(m - 1)^{n-1}}. \quad (\text{III.40})$$

For higher corrections the height shift must be retained. Directly,

$$\log R_K(n, m) = \log R(n, m) + \beta + O(1/m + n/m^2 + c^{-n}). \quad (\text{III.41})$$

In the crossover window these residual errors are smaller than every fixed inverse power of $\log n$. Thus Theorem 24.1 and its exponential version hold for R_K with K replaced everywhere by $K + 1$, while every c_j for $j \geq 2$ is unchanged. In particular the second-order exponential coefficient becomes $((K + 1)^2/2 + c_2)/(\lambda^2 L^2)$. If one instead normalizes by $\ell_n m^{n-1}$, K stays unchanged.

[write] **Kaneiwa in Part I, 6 October 2026.** The indexing $p(r; n) = A(n, r + 1)$ is Part I's $p(r; n) = [z^n]F_{r+1}$ (Subsection 1.1), where the same Propositions 1 and 2 of Kaneiwa are credited.

34 Smooth suppression ratio inverses

Fix a compact interval $\mathcal{R} = [r_0, r_1] \subset (0, 1)$ and write

$$\tau = -\log r \in [\tau_0, \tau_1] = [-\log r_1, -\log r_0], \quad \chi_n = \frac{\log n}{3} - K. \quad (\text{III.42})$$

The suppression logarithm τ is distinct from the density argument. Define the actual smooth germ

$$C(\beta) = \log \frac{g(\beta)}{2} - \frac{C_0}{3}\beta. \quad (\text{III.43})$$

It has $C(0) = C'(0) = 0$ and finite Taylor expansions $C(\beta) = \sum_{j=2}^N c_j \beta^j + O(\beta^{N+1})$, with all fixed differentiated remainders. This definition uses the actual density, not an assumed convergent formal sum.

34.1 The unique small model root

Proposition 34.1. *There is $\beta_0 > 0$ such that for every sufficiently large n and every $r \in \mathcal{R}$ the equation*

$$\tau = \chi_n \beta_* - C(\beta_*) \quad (\text{III.44})$$

has exactly one solution $\beta_(n, r) \in (0, \beta_0)$. Uniformly in r , $\beta_* \sim \tau/\chi_n$.*

Proof. Choose β_0 so that $g > 0$ and C is smooth on $[0, \beta_0]$. For large n , the derivative of the right side is $\chi_n - C'(\beta) \geq \chi_n/2$ throughout that interval. Its values at the endpoints are zero and a quantity tending to infinity. The intermediate value theorem and strict increase prove existence and uniqueness in this endpoint interval. With $C(\beta) = O(\beta^2)$ and τ in a fixed positive compact interval, first $\beta_* = O(\chi_n^{-1})$, and then $\beta_* = \tau/\chi_n + O(\chi_n^{-3})$. No uniqueness outside the stated interval is asserted. $\square$

34.2 Every fixed reversion order

Put $x = 1/\chi_n$ and $u = \beta_*/x$. Equation (III.44) becomes

$$u = \tau + C(xu). \quad (\text{III.45})$$

Finite substitution determines every coefficient recursively. The first terms are

$$\beta_* = \frac{\tau}{\chi_n} + \frac{c_2 \tau^2}{\chi_n^3} + \frac{c_3 \tau^3}{\chi_n^4} + \frac{2c_2^2 \tau^3 + c_4 \tau^4}{\chi_n^5} + O_{\mathcal{R}}(\chi_n^{-6}), \quad (\text{III.46})$$

$$y_* := \frac{1}{\beta_*} = \frac{\chi_n}{\tau} - \frac{c_2}{\chi_n} - \frac{c_3 \tau}{\chi_n^2} - \frac{c_2^2 \tau + c_4 \tau^2}{\chi_n^3} + O_{\mathcal{R}}(\chi_n^{-4}). \quad (\text{III.47})$$

The absent χ_n^{-2} term in (III.46) and absent constant term in (III.47) are consequences of $C'(0) = 0$. Since $c_2 < 0$, the first displayed correction to the height model is positive.

Theorem 34.2. *For every fixed $J \geq 0$ there are explicitly generated polynomials $d_j(\tau)$ such that*

$$y_J(n, r) = \frac{\chi_n}{\tau} + \sum_{j=1}^J d_j(\tau) \chi_n^{-j}, \quad (\text{III.48})$$

$$y_*(n, r) - y_J(n, r) = O_{\mathcal{R}, J}(\chi_n^{-J-1}).$$

Here $d_1 = -c_2$, $d_2 = -c_3 \tau$, and $d_3 = -c_2^2 \tau - c_4 \tau^2$. The coefficient d_J requires only $c_2, \dots, c_{J+1}$.

Proof. Take a finite formal truncation $U_M(\tau, x)$ of (III.45) through degree M . Taylor's theorem for the smooth C gives a residual

$$U_M - \tau - C(xU_M) = O_{\mathcal{R}, M}(x^{M+1}).$$

Both the true u and this truncation stay in a common positive compact interval, uniformly in τ . There

$$\frac{\partial}{\partial u} \{u - \tau - C(xu)\} = 1 - xC'(xu) \geq \frac{1}{2}$$

for small x . The real mean-value theorem bounds $u - U_M$ by twice the residual. Since u stays bounded away from zero, finite reciprocal expansion then gives every claimed order for $y_* = 1/(xu)$.

Write $\beta_* = \tau x V$. The equation becomes $V = 1 + \sum_{j \geq 2} c_j \tau^{j-1} x^j V^j$ at each finite order. Every nonconstant coefficient of V is therefore a polynomial divisible by τ . The same holds for $1/V - 1$, so division by τx introduces no negative powers of τ beyond the leading $1/(\tau x)$. Formal recursion shows that the coefficient of x^j in y_* first involves c_{j+1} , proving polynomiality and the stated dependency. In particular direct substitution gives (III.46) and (III.47) with the displayed signs.

Equivalently one can solve the truncated polynomial model $\tau = \chi_n \beta - \sum_{j=2}^{J+1} c_j \beta^j$ in its unique small-root interval. Its omitted C remainder is $O(\beta^{J+2})$; division by the model derivative costs χ_n^{-1} and reciprocation costs χ_n^2 . Thus its height differs from y_* by $O(\chi_n^{-J-1})$, and it has the same y_J . Other polynomial roots are irrelevant. No convergence of an infinite reversion series is invoked. $\square$

34.3 Rounded height prescriptions and observed ratios

Theorem 34.3. *Choose any integer $m_J(n, r)$ with $|m_J - ny_J(n, r)| \leq 1$. Then, uniformly for $r \in \mathcal{R}$,*

$$\log \frac{R(n, m_J)}{r} = O_{\mathcal{R}, J} \left(\chi_n^{-J-2} + \frac{\chi_n^{30}}{n} \right). \quad (\text{III.49})$$

Rounding the exact smooth model ny_ instead gives error $O_{\mathcal{R}}(\chi_n^{30}/n)$ in this logarithmic ratio.*

Conversely, suppose integer heights have $m/n \asymp \log n$ and the actual observed ratio $r_n = R(n, m)$ lies in $\mathcal{R}$. Then

$$\frac{m}{n} = y_J(n, r_n) + O_{\mathcal{R}, J} \left(\chi_n^{-J-1} + \frac{\chi_n^{31}}{n} \right), \quad (\text{III.50})$$

with constants also depending on the fixed comparison bounds in $m/n \asymp \log n$.

Proof. Using the exact smooth germ in (III.32), rather than a Taylor truncation, gives on this height window

$$\log R(n, m) = -\chi_n \beta + C(\beta) + O(\chi_n^{30}/n). \quad (\text{III.51})$$

Indeed $m \asymp n\chi_n$, $\beta \asymp \chi_n^{-1}$, and $E(m, \beta) = O(\chi_n^{30}/n)$. As a function of $y = m/n$, the smooth part is

$$F_n(y) = -\chi_n/y + C(1/y), \quad F'_n(y) = \frac{\chi_n - C'(1/y)}{y^2} \asymp \chi_n^{-1}$$

uniformly on any common crossover-height window. The error $y_J - y_* = O(\chi_n^{-J-1})$ therefore costs $O(\chi_n^{-J-2})$ in the log ratio. Integer rounding costs $O(1/(n\chi_n))$, which is absorbed by χ_n^{30}/n . Since $F_n(y_*) = -\tau = \log r$, this proves (III.49) and the exact-model statement.

For the converse, take $\tau_n = -\log r_n$ at the actual observed ratio. Equation (III.51) and the lower derivative bound show that $m/n - y_*(n, r_n) = O(\chi_n^{31}/n)$. Add Theorem 34.2 to obtain (III.50). Inverting the ratio error costs precisely one extra factor χ_n . $\square$

These formulas use a specified target r , or the actual r_n at each n . Replacing r_n by its mere limiting value generally loses higher-order information unless an appropriate convergence rate is known separately. The available coefficient theorem gives no discrete monotonicity, exact ratio solvability, least integer height, or effective onset. Thus the estimates are smooth prescriptions and asymptotic observed-ratio inverses, not threshold or exact-integer-recovery theorems.

For the Kaneiwa normalization (III.40), replace K by $K + 1$ throughout this section, hence replace χ_n by $(\log n)/3 - K - 1$. The residual shift in (III.41) is smaller than χ_n^{30}/n on the window. The coefficients c_j , reversion signs, and stated orders remain unchanged.

[write] **No Lambert function here, 6 October 2026.** Unlike Parts I and II, these inverses involve no Lambert W : the core of (III.44) is linear (the case $b = 0$ of `p0:thm:lambert-core` in [11]), and the reversion with the smooth germ C is proved by Taylor’s theorem; it is an analogue, not an instance, of `p0:thm:perturbed-inversion`, whose hypotheses ask for analyticity, and the rounded prescriptions are not instances of `p0:thm:staircase`. The write rechecked (III.46)–(III.47) by substitution.

35 Exact finite comparisons and reproducibility

35.1 An exact finite coefficient algorithm

Logarithmic differentiation of the Euler product gives, with the preceding-height coefficients $f_d = A(d, m)$,

$$b_j = \sum_{d|j} df_d, \quad c_0 = 1, \quad c_k = \frac{1}{k} \sum_{j=1}^k b_j c_{k-j}, \quad A(k, m+1) = c_k \quad (k \geq 1). \quad (\text{III.52})$$

Truncation at degree N is exact for all coefficients retained. The public implementation uses integers and checks divisibility at each division. A separate finite-product implementation checks small tables using $(1 - z^d)^{-a} = \sum_{k \geq 0} \binom{a+k-1}{k} z^{dk}$. The working limits, degree and height at most 640, are explicit implementation caps, not theorem restrictions. Computing height M through degree N uses $O(MN^2)$ integer operations and $O(N)$ live coefficients apart from saved outputs; bit costs grow with coefficient size.

The code uses explicit exceptions for all pass/fail conditions, so checks remain active under Python’s `-O`. Chunked integer decimal conversion works with the minimum configurable integer-string digit limit 640 and does not change any process-wide limit. A guard test exercises a value exceeding that digit count. The exact coefficient generation is independent of any decimal normalization or asymptotic theorem.

35.2 Finite crossover diagnostics

Table 1 uses exact recurrence values at $n = 20, 40, 80$ and $m = \lfloor \lambda n \log n \rfloor$ for $\lambda = 1/2, 1, 3/2$. All heights are below 640. For finite-height comparison define

$$R_2(n, \lambda) = e^{-1/(3\lambda)} \left(1 + \frac{K}{\lambda \log n} + \frac{K^2/2 + c_2}{(\lambda \log n)^2} \right). \quad (\text{III.53})$$

This is the explicit second-order ratio expansion, rather than the exponential of a truncated logarithm. The exact coefficient uses the displayed integer height. The nominal floor choices agree at 80 and 160 decimal digits; this is a computational check, not an interval certificate for the transcendental floor. Values of R and R_2 are rounded floating evaluations, not interval enclosures; only $A(n, m)$ is computed exactly. The column $e^{-1/(3\lambda)}$ is the limiting ratio. These finite examples illustrate the size of the correction and do not establish an effective error bound or a certified asymptotic onset. The accompanying JSON records exact decimal coefficients, the chosen integer heights, normalization formulas, and diagnostic precision. The coefficient generator separately verifies the rational values through ρ_7 and the finite inverse identities. Symbolic and exact-integer checks are reproducible algebra; floating agreement is not an analytic remainder certificate.

[write] **The labelled counterpart, 6 October 2026.** Remark 37.2 gives the same comparison for the labelled array $H(n, m)$ at the same heights, against a conditional prediction.

n	λ	m	$R(n, m)$	$e^{-1/(3\lambda)}$	$R_2(n, \lambda)$
20	0.5	29	0.89828117	0.51341712	0.86765860
20	1	59	0.94696428	0.71653131	0.93775982
20	1.5	89	0.96417635	0.80073740	0.95910776
40	0.5	73	0.80413195	0.51341712	0.78974074
40	1	147	0.89724904	0.71653131	0.89222870
40	1.5	221	0.93039388	0.80073740	0.92738192
80	0.5	175	0.74764498	0.51341712	0.73949701
80	1	350	0.86485251	0.71653131	0.86215689
80	1.5	525	0.90778645	0.80073740	0.90621689

Table 1: Unvalidated floating comparisons built from exact finite coefficients. The two displayed approximations have different purposes: the limit records the crossover theorem, and R_2 retains the first two crossover ratio corrections.

35.3 Reproducing the archive

The archive includes editable TeX, exact and symbolic programs, generated comparisons, the source ledger, and unchanged Reports 225 and 226. Its builder verifies a source manifest, reruns the public checks, copies inputs to a new external output directory, creates a private pdfL^AT_EX format, compiles without shell escape, and rejects unresolved references, missing characters, or overfull boxes. Sources are preserved. Sorted ZIP members have fixed timestamps and permissions. Normal and optimized runs, including actual extraction and rebuilding of the ZIP, are compared by complete archive bytes and member hashes. Byte identity is a same-toolchain claim; different TeX, font, Python, or numerical-library versions can change output bytes.

[write] **The companion here, 6 October 2026.** Report 228’s companion is shipped as `code/228-logheight-*.py`, `data/228-logheight-*`, `228-logheight-code-README.md` and `228-logheight-sources-SOURCE_LEDGER.md`; its `supporting/` copies are Parts I and II. The delivered `code/reproduce.py--check` regenerated both result files byte for byte on a copy at the write; the report README gives the rerun route.

36 Further questions and limits

36.1 Polynomially small ratios

The explicit β^{-30} loss identifies an obstacle to much smaller ratios. The present theorem is organized around fixed logarithmic bands and the height window $m \asymp n \log n$. A broad transfer theory for polynomially small β , especially one approaching the fixed-degree polynomial regime, is not proved here. A narrow range may be tested against the displayed absolute error, but that observation alone does not supply a sharp uniform transition theorem, optimal range, or a high-height error matching Kaneiwa’s polynomial expansion. A useful next step is to exploit lower derivative orders or the retained first correction in the absolute Fourier norm, with explicitly balanced endpoint errors. Such a result would need its own statement and proof rather than a claim of arbitrary-small- β uniformity.

[write] **Still open, 6 October 2026.** Section 37, item 7.

36.2 Effective bounds and validated evaluation

Our remainder constants and onset are qualitative. A numerical certificate would require explicit bounds for the sectorial coordinate remainder, its inverse identification, contour tails, the finite-height derivative errors, and floating evaluation of the chosen model. Higher numerical precision or repeated mesh agreement cannot replace these bounds. An effective counterpart of (III.49) could produce rigorously enclosed height intervals for target ratios. Determining a least height would additionally require suitable discrete information about $R(n, m)$; no monotonicity result is available in this report.

[write] **Still open, 6 October 2026.** Section 37, items 1 and 3.

36.3 Stronger endpoint structure

The smooth germ g has an all-fixed-order Taylor asymptotic expansion on a sector, but analyticity at zero and convergence of its Taylor series have not been shown. Classical resurgence theory suggests studying sharper coefficient growth and summability questions, with attention to the nonzero resiter. These would concern the universal map, not a new Euler-specific density. Likewise a complete analytic comparison with the public IBD contour amplitude would clarify the global contour normalization beyond the local coordinate identity recorded above.

[write] **Still open; shared with the iterated Bell report, 6 October 2026.** Section 37, items 6 and 9.

36.4 Other arrays and other height windows

The constants T and B arise from the subsidiary Euler forcing. For another multiset recurrence, or another initial function, the universal endpoint calculation may persist while the clock and transfer change. Extending this method requires proving the new forced entrance and angular-tail estimates. At the opposite endpoint $\beta \rightarrow \infty$, a different density regime is needed. Finally, every fixed inverse-logarithm order here coexists with only one polynomial-in-height transfer rate; an all-orders $1/m$ theorem would require a further hierarchy of quantitative forced corrections and is not a consequence of the present proof.

[write] **Still open, 6 October 2026.** Section 37, items 2, 5 and 8.

37 Further questions and research (all three Parts)

[write] **Added in the write, 6 October 2026.** This section gathers the open questions of the three Parts under Vladimir’s standing rule of 4 October 2026: every claim that a manuscript states without proof, every limitation it records as a question, and the question shared with the iterated Bell report. Each item names its sources, the sketch they give, and what is missing. Questions answered inside the merge are marked at the questions themselves. No claim of the three manuscripts was found to be wrong, and none is refuted here.

1. *Validated constants.* Sources: Part I, Subsection 11.1; Part II, Subsection 23.2; Part III, Subsection 36.2. The amplitude c has the absolutely convergent formula (6); the first correction needs $h(1)$, $h'(1)$, $h''(1)$, since κ is now in closed form (II.3). Sketch: interval enclosures of the truncated Fatou coordinate on a stated sector (Lemma 26.1 gives the rational coefficients to every order), of the inverse branch, of the forward propagation of values and derivatives, of the finite quadrature and of the vertical tail. Missing: explicit numerical values for every constant of the sector proofs. Current values ($c \approx 4.4923299$, $h(1) \approx 1.81460956$) are floating diagnostics only.
2. *Corrections beyond the first, and an all-orders theorem.* Sources: Part I, Subsection 11.2; Part II, Subsection 23.1; Part III, Subsection 36.4 (an all-orders theorem in $1/m$). The expected form is $1 + \sum_r P_r(\log n)/n^r$ with $\deg P_r \leq 2r$; Part II proves P_1 . Part II’s conditional inference, that $[X^{2r}]P_r = (-1/18)^r/r!$ (so $\log^4 n/(648n^2)$ at second order) *if* an all-orders construction gives the degree bound and no other contribution reaches degree $2r$, is unproved. Missing: an induction over subsidiary orders, clock hierarchies with resonant logarithms (Part II’s N already has $\frac{1}{540} \log y$), forced Fatou cohomological corrections, and growing-strip errors with an extraction order chosen per precision.
3. *Effective constants, onsets and certified thresholds.* Sources: Part I, Subsection 11.3; Part II, Subsection 23.2; Part III, Subsection 36.2. All three Parts give qualitative constants only; their inverses recover n at exact sequence values eventually, without onset, and Part III’s prescriptions are smooth. Missing: explicit clock, stopping and sector constants; for least heights in Part III, discrete information (monotonicity) about $R(n, m)$, which no Part supplies.
4. *Inverse statements for $A(n, \lfloor \lambda n \rfloor)$.* Source: Part I, Subsection 11.3, second paragraph. The floor phase $e^{-\{\lambda n\}/\lambda}$ of (32) must enter; a smooth approximation that drops it has a non-vanishing multiplicative error. Not treated by any Part (Part III’s inverses concern $m \asymp n \log n$).
5. *The regime $n/m \rightarrow \infty$, and corrections near $n/m \rightarrow 0$.* Sources: Part I, Subsection 11.4; Part II, Subsection 23.3; Part III, Subsection 36.4. The side $n/m \rightarrow 0$ is answered at leading order for logarithmic bands by Part III (Theorem 31.1); the large-argument behaviour of h and any transfer as $\beta \rightarrow \infty$ are open, and no Part proves a first correction of Part II’s kind on a shrinking band.
6. *Analytic structure of h .* Sources: Part I, Subsection 11.4 (“stronger regularity, such as real analyticity”); Part III, Subsection 36.3 (analyticity of the germ g at zero, convergence of its Taylor series, coefficient growth and summability, with the nonzero resiter $1/3$). Known: $h \in C^\infty(0, \infty)$ (Corollary 6.2); h continues holomorphically to a small punctured sector about $(0, \epsilon)$, so h is real analytic on $(0, \epsilon)$ (Theorem 28.1); g has a one-sided C^∞ germ (Theorem 29.1). Missing: real analyticity on all of $(0, \infty)$ (the measure identity (28) is the suggested route; item 9 would give it), and any statement on the Taylor series of g .
7. *Polynomially small ratios and Kaneiwa’s regime.* Sources: Part III, Subsection 36.1; Part I, Subsection 11.4, second paragraph; Part II, Subsection 23.3, second paragraph. Part III’s transfer loses β^{-30} and stops at logarithmic bands; a uniform theory linking proportional height, the

crossover and Kaneiwa's fixed-degree polynomials (Section 33) is open. Part III's suggested step: lower derivative orders or the retained first correction in the absolute Fourier norm, with balanced endpoint errors.

8. *Other arrays and initial functions.* Sources: Part I, Subsection 11.4; Part II, Subsection 23.3; Part III, Subsection 36.4. The forcing and clock constants (T, B, κ) must be recomputed for another multiset recurrence or initial function; the endpoint calculus of the unforced map may persist. Missing: the forced entrance and tail estimates in each new case.
9. *The amplitude identity with the iterated Bell report* (a question shared with `a139383-iterated-bell-diagonals`). Source: Part III, Section 25 ("a formal contour match suggests an amplitude relation $I(\beta) = \beta 2^{\beta/3} h(\beta)$, but that identification requires its own analytic contour correspondence and is not used here") and Subsection 36.3 ("a complete analytic comparison with the public IBD contour amplitude"). Here I is the iterated Bell report's Part II amplitude, $H(n, m) \sim (n-1)! 2^{-n} (\lambda n)^n (\lambda n)^{-1/(3\lambda)} I(1/\lambda)$ for $m = \lambda n$ (its `ibd:pd:eq:factorial`), defined there by a fixed-radius contour functional. Sketch: both reports use the same normalized reciprocal orbit, and $\Psi_{\text{IBD}} = -\Phi + \frac{1}{3} \log 2$ holds locally; the iterated Bell report's own literature section derives $I(\beta) = \beta J_P(\beta)$ from Prellberg's contour integral under the same kind of unproved contour correspondence. Evidence: agreement to eight significant digits at $\beta = 1$ (Remark 37.1). Missing: an analytic identification of that report's contour functional with the Laplace inversion of P (orientation, normalization, vanishing endpoint terms). Consequences if proved: (a) h is real analytic on $(0, \infty)$, because that report proves I real analytic there (item 6); (b) Part III's endpoint results ($h(0) = 2$, Corollary 29.2, (III.29)) give the behaviour of $I(\beta)$ as $\beta \rightarrow 0$, that is, the amplitude half of the endpoint $\lambda \rightarrow \infty$ that the iterated Bell report leaves open; a labelled analogue of Theorem 31.1 would still be needed for a coefficient theorem (Remark 37.2).

Remark 37.1 ([**write**] The identity at $\beta = 1$, numerically, 6 October 2026). The iterated Bell report's shipped exploratory runs (`data/53-iterated-bell-exploratory-amplitude.json` there) give $I(1) = 2.2862647982, 2.2862647847, 2.2862647863$ (its $C = \sqrt{2\pi} I(1)/2$ between 2.865407976 and 2.865407993). Part I's shipped diagnostic $h(1) = 1.8146095596784502$ (`data/225-leading-data-amplitude_diagnostic.json`) gives $2^{1/3} h(1) = 2.2862647816$. The two sides agree to eight significant digits (2.2862648); their differences, 3×10^{-9} to 1.7×10^{-8} , are of the size of the spread of the three iterated Bell runs (1.4×10^{-8}). (Added after the independent check of 6 October 2026: an independent evaluation of (6) by the verification (four values of σ , Gauss–Legendre quadrature to $|u| = 4 \cdot 10^4$ with an integration-by-parts tail; uncertified) gives $h(1) = 1.814609559835$ and $2^{1/3} h(1) = 2.2862647818$, so the shipped diagnostic is low by 1.6×10^{-10} , from its truncation of the vertical tail at $|u| = 3000$; the comparison is unchanged, with differences 2.9×10^{-9} to 1.65×10^{-8} and agreement to eight significant digits.) Both sides are floating values without error control; this is evidence for item 9, not a proof of it, and it says nothing about $\beta \neq 1$.

Remark 37.2 ([**write**] A conditional consequence for the labelled array, 6 October 2026). Let $H(n, m) = n! [z^n] (e^z - 1)^{\circ m}$ and $R_{\text{lab}}(n, m) = H(n, m) / (n! m^{n-1} 2^{1-n})$. Suppose, for some $\beta_0 > 0$ and $D > 1$, (a) $I(\beta) = \beta 2^{\beta/3} h(\beta)$ for $0 < \beta \leq \beta_0$, and (b) $H(n, m) = (n-1)! 2^{-n} m^n m^{-\beta/3} I(\beta) \{1 + \epsilon(n, m)\}$ uniformly for $(\log m)^{-D} \leq \beta = n/m \leq \beta_0$, with $\epsilon \rightarrow 0$ (the iterated Bell report's leading form `ibd:pd:eq:factorial` with $\lambda n = m$, extended to logarithmic bands). If $m/(n \log n) \rightarrow \lambda \in (0, \infty)$, then $R_{\text{lab}}(n, m) \rightarrow e^{-1/(3\lambda)}$, the same limit as Part III's Theorem 24.1; and if $\epsilon = O(L^{-N-1})$ with $L = \log n$ and $m = \lfloor \lambda n L \rfloor$, then

$$\log R_{\text{lab}}(n, m) = -\frac{1}{3\lambda} + \frac{K_{\text{lab}}}{\lambda L} + \sum_{j=2}^N \frac{c_j}{(\lambda L)^j} + O(L^{-N-1}), \quad K_{\text{lab}} = \frac{1-\gamma}{3},$$

with the same c_j as (III.6).

Proof. By (a) and (b), since $(n-1)!m^n\beta = n!m^{n-1}$,

$$R_{\text{lab}}(n, m) = 2^{\beta/3} m^{-\beta/3} \frac{h(\beta)}{2} \{1 + \epsilon(n, m)\}.$$

By (III.29), $\log(h(\beta)/2) = -\frac{\beta}{3} \log \beta + \frac{C_0}{3} \beta + \sum_{j=2}^N c_j \beta^j + O(\beta^{N+1})$ with $C_0 = 1 - \gamma - \log 2$. Hence, using $\log \beta + \log m = \log n$ exactly as in (III.36),

$$\log R_{\text{lab}} = -\frac{\beta}{3} \log n + \frac{\log 2 + C_0}{3} \beta + \sum_{j=2}^N c_j \beta^j + O(\beta^{N+1}) + \log(1 + \epsilon),$$

and $\log 2 + C_0 = 1 - \gamma$. If $m/(n \log n) \rightarrow \lambda$, then $\beta \rightarrow 0$, $\beta \log n \rightarrow 1/\lambda$, and β eventually lies in the band of (b): $\beta \leq \beta_0$, and $\beta \geq 1/(2\lambda \log n) \geq (\log m)^{-D}$ because $D > 1$ and $\log m \sim \log n$. So the right side tends to $-1/(3\lambda)$. For $m = \lfloor \lambda n L \rfloor$, $\beta = 1/(\lambda L) + O(1/(nL^2))$, whose effect is smaller than every power of L , as in the proof of Theorem 24.1; substituting gives the display. $\square$

Hypothesis (b) is not proved anywhere: the iterated Bell report's Theorem (its `ibd:pd:thm:main`) is uniform only for $\lambda = m/n$ in compact subsets of $(0, \infty)$, and it records the endpoints $\lambda \rightarrow 0, \infty$ as open. A consistency check, not a proof: with $h(0) = 2$ the right side of (b) is $n!m^{n-1}2^{1-n}(1 + o(1))$ when $\beta \log m \rightarrow 0$, the leading term of the fixed- n asymptotic $H(n, r+1) \sim n!r^{n-1}/2^{n-1}$ of Skau and Kristensen that the iterated Bell report cites. Exact values of $H(n, m)$, computed by the write from $H(n, m+1) = \sum_j S(n, j)H(j, m)$ with Stirling numbers $S(n, j)$ and evaluated in 30-digit floating arithmetic (uncertified), behave as the remark predicts, with the heights $m = \lfloor \lambda n \log n \rfloor$ of Part III's Table 1:

n	λ	m	$R_{\text{lab}}(n, m)$	$e^{-1/(3\lambda)}$	$e^{-1/(3\lambda)}(1 + K_{\text{lab}}/(\lambda \log n))$
20	0.5	29	0.56666331	0.51341712	0.56172244
20	1	59	0.75860855	0.71653131	0.75023906
20	1.5	89	0.83316752	0.80073740	0.82585011
40	0.5	73	0.55659049	0.51341712	0.55264578
40	1	147	0.74875697	0.71653131	0.74390532
40	1.5	221	0.82522543	0.80073740	0.82113138
80	0.5	175	0.54955621	0.51341712	0.54644061
80	1	350	0.74202367	0.71653131	0.73957531
80	1.5	525	0.81979276	0.80073740	0.81790547

At $\lambda = 1$ the distance to the limit is 0.042, 0.032, 0.025 at $n = 20, 40, 80$, and after the K_{lab} term 0.0084, 0.0049, 0.0024. The Euler constant $K \approx 0.816$ of Part III and $K_{\text{lab}} \approx 0.141$ differ, as the subsidiary forcing leads one to expect. These finite values neither prove the limit nor certify any digit.

[write] **Independent check of the write, 6 October 2026.** An adversarial check made by the intake after the write (commit `84fc1aead`) re-pulled A290354 (live revision #26 of 21 June 2018) and A290353 (#28) and found every quoted line character for character, including the formula line in the front matter, in the note at the start of Part I and in the bibliography. The entry's history places the conjecture in Kotěšovec's revision #23 of 14 August 2017, so the date "Aug 14 2017" that the write gives is the date of the conjecture, not of the revision read, and it is correct. The check recomputed a_n as the n -th Euler transform of $1 + x$ in integer arithmetic (equal to the b-file for $n \leq 40$; $a_{40}(T/40)^{40}40^{4/3} = 4.6013347$), confirmed that Theorem 1.1 is exactly the conjectured form, and the rows $A(2, m)$, $A(3, m)$, $A(4, m)$ for $m \leq 12$. It

re-derived Remark 37.2 from (a), (b) and (III.29), found it labelled conditional wherever it appears (title, hypotheses, the paragraph after the proof, the README) and not listed as proved in the status table, and reproduced its table to every printed digit from an own exact computation of $H(n, m)$; extended to $n = 160$ it gives $m = 812$, $R_{\text{lab}} = 0.73782182$ at $\lambda = 1$ (distance 0.0213 to the limit, 0.0014 after the K_{lab} term) and $m = 406$, $R_{\text{lab}} = 0.54374893$ at $\lambda = 0.5$ (0.0303 and 0.0018). It evaluated the closed form (II.3) to 35 digits, $\kappa = -0.25434816985107480385506489650373339$, and matched it against an independent quadrature of the defining equation (II.23) to 5×10^{-28} . It recomputed $h(1)$ independently (Remark 37.1, dated note) and confirmed the classification of the inverses in the front matter: the factorial core and the Lambert core with $a = b = 1$ are instances; the corrections, the rounding at exact values and Part III's reversion and prescriptions are not. Two wordings are corrected, each with a dated note keeping the first wording: the front matter's justification of the printed restatements (twice) overstated Part II's novelty, since its bound on inverse orbits is in Part I's proof of Lemma 4.1; and the note after Part II's Section 13 now flags the neighbourhood form of the outer bound, as it already flagged the disk form of the tail bound, as proved only inside Part I's proofs. Added: the name $L = \log(Y/T)$ in the front matter, and the independent values of $h(1)$ and $c = 4.4923298971$. No mathematical error was found.

References

- [1] The OEIS Foundation Inc., *A290353*, square array of iterated Euler transforms, entry by Alois P. Heinz, 28 July 2017. Retrieved 5 October 2026. <https://oeis.org/A290353>.
[As cited in Report 228: *A290353*, iterated Euler-transform array, entry by Alois P. Heinz, 28 July 2017. Source checked 5 October 2026. Report 226: Source checked for Report 225 on 5 October 2026.]
- [2] The OEIS Foundation Inc., *A290354*, entry by Alois P. Heinz, 28 July 2017; asymptotic conjecture by Václav Kotěšovec, 14 August 2017. Retrieved 5 October 2026. <https://oeis.org/A290354>.
[As cited in Report 226: leading asymptotic conjecture by Václav Kotěšovec, 14 August 2017. Source checked for Report 225 on 5 October 2026.]
- [3] R. Kaneiwa, *On r -fold Partitions and a Certain Form of Infinite Products*, Otaru University of Commerce, Humanities Research **61** (1980), 11–26. <https://barrel.repo.nii.ac.jp/records/2331>; persistent identifier <https://hdl.handle.net/10252/2056>.
[Report 228 adds: Propositions 1 and 2, printed p. 19.]
- [4] M. Bechtloff Weising, *Higher Order Bell Symmetric Functions*, arXiv:2505.18504v2. <https://arxiv.org/abs/2505.18504v2>. Definitions and statements checked in the versioned HTML on 5 October 2026.
- [5] *Asymptotics of Iterated Bell Diagonals (OEIS A139383, A261280)*, unrefered research report in VladimirReshetnikov/ProveIt, combined fixed-shift and proportional-depth manuscripts, 2 October 2026. Catalogue checked 5 October 2026; indexed commit 65969409cfc007047182e9303e1d2d0106324dea. [Repository report](#).
[As cited in Report 228: unrefered report in VladimirReshetnikov/ProveIt, 2 October 2026. Snapshot at main commit 5fdc0d68a062c815100ac27bd173a8c4b901c08d, checked 5 October 2026. [Pinned public source](#). Sections 4, 12, 13, and 17.] **[[write]]** In this repository: `SetTheory/Cardinals/docs/reports/generating-functions-and-asymptotics/oeis-sequence-asymptotics/a139383-iterated-bell-diagonals`; its `article.tex` is the same blob at both pins and at HEAD, and its Sections 4, 12, 13 and 17 are the sections Report 228 names.]
- [6] *Diagonal and proportional height asymptotics for iterated Euler transforms*, Report 225, 5 October 2026. **[[write]]** Printed in this report as Part I.] As cited in Report 226: preceding unrefered research report; unchanged PDF and editable TeX included in the archive's **supporting** directory; exact provenance in `sources/SOURCE_LEDGER.md`. As cited in Report 228: unrefered research report, 5 October 2026; unchanged complete TeX and PDF in the accompanying archive; provenance in `sources/SOURCE_LEDGER.md`. **[[write]]** The included copies are byte-identical to Part I and

are not shipped; the ledgers are shipped as `226-firstcorr-sources-SOURCE_LEDGER.md` and `228-logheight-sources-SOURCE_LEDGER.md`.]

- [7] *First logarithmic correction for iterated Euler diagonals*, Report 226, unrefereed research report, 5 October 2026. Unchanged complete TeX and PDF in the accompanying archive; provenance in `sources/SOURCE_LEDGER.md`. `[[write]]` Printed in this report as Part II; the included copy is byte-identical to it and is not shipped.]
- [8] A. Dudko and D. Sauzin, *The resurgent character of the Fatou coordinates of a simple parabolic germ*, C. R. Acad. Sci. Paris I **352** (2014), 255–261. <https://doi.org/10.1016/j.crma.2013.12.005>.
- [9] T. Prellberg, *On the Asymptotic Analysis of a Class of Linear Recurrences*, September 2002 seminar, summary by M. Mishna, Algorithms Seminar 2002–2004 (2005), 47–50. Sections 2.2 and 3.1. <https://algo.inria.fr/seminars/summary/Prellberg2002a.pdf>.
- [10] S. V. Nagaev and V. I. Vakhtel, *Limit theorems for probabilities of large deviations of a Galton–Watson process*, Discrete Math. Appl. **13**(1) (2003), 1–26. Theorems 1–2 and their Poisson specialization; the source-comparison limitations are recorded in the ledger. <https://doi.org/10.1515/156939203321669537>; <https://www.mathnet.ru/eng/dm183>.
- [11] `[[write]]` *Transseries and Inversion*, the transseries volume of this repository, `Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/transseries_and_inversion.tex`: Proposition `p0:prop:factorial-core`, Theorems `p0:thm:lambert-core`, `p0:thm:perturbed-inversion` and `p0:thm:staircase`.