

Merged research report for the Surreal project

Omnific-Preserving Automorphisms

Convex-scale stabilizers, definable constants,
nondefinable monomials, and algebraic-parameter rigidity

Built from eleven manuscripts dated September 23, 2026

Abstract

Which symmetries of a Hahn field preserve its omnific-type integer part, and what does that integer part remember? Let k have characteristic zero, let $\Gamma \neq 0$ be a divisible ordered abelian group, write C_δ for the principal convex subgroup generated by $\delta > 0$, and let $\mathfrak{o} \subseteq k$ be a unital subring. A strong coefficient-fixing 1-automorphism σ of $k((t^\Gamma))$ preserves $\mathfrak{o} \oplus \Pi$, where Π is the purely infinite ideal, exactly when, for every $\delta > 0$, the shift- δ coefficient of its logarithmic character $\log(t^{-g}\sigma(t^g))$ vanishes on C_δ . The proof uses no exponential–logarithm correspondence and applies verbatim to the surreal field and $\mathbf{Oz}$. Where that correspondence is available, the same condition for the operator logarithm $\text{Log } \sigma$ is equivalent; the two coefficient functionals are different.

Consequences: the stabilizer is trivial exactly in Archimedean rank one; its common fixed field is the Hahn field on the bottom convex subgroup, and for $(\mathbf{No}, \mathbf{Oz})$ it is $\mathbb{R}$; in finite rank r its derived length is at most $r - 1$, with equality for $\mathbb{Q}^r$; explicit successive-scale flows generate a faithful $\mathbb{Z} \wr \mathbb{Z}$; for every set of surreal parameters the pointwise stabilizer is nonsolvable and carries a faithful action of $\mathbb{R}^\kappa$ for every set cardinal κ . Consequently no set of parameters makes the Conway monomials, the omega-map, any valuation-representative rule, the simplicity relation or the Gonshor exponential definable in $(\mathbf{No}, +, \cdot, <, \mathbf{Oz})$, although the pair defines the three zero-cut projections and the floor map; a monomial can even be sent to a two-term omnific integer. An equivalent form of the criterion bounds the support of the principal unit $t^{-g}\sigma(t^g)$ itself, and every element of the stabilizer keeps each monomial in the signed Archimedean block of its exponent. A second part proves algebraic-parameter rigidity of $\mathbf{Oz}$: embeddings into polynomial, Laurent and finite-type domains land in the coefficients, strong cancellation holds, and $\text{ML}(\mathbf{Oz}[T_1, \dots, T_n]) = \mathbf{Oz}$, while explicit commuting derivations integrate to formal flows that no polynomial family realizes. A third part proves that every ring automorphism of $\mathbf{Oz}$ extends to a strong automorphism of $\mathbf{No}$, as constant-term tests detect Hahn summability, and classifies the strong monomial-preserving embeddings that preserve the integer parts; proper self-embeddings of $\mathbf{No}$ preserving and reflecting $\mathbf{Oz}$ can move a real number. A fourth part adds a third proof of the constant-term detection, by induced matchings, detection inside subfields containing enough all-ones series, and equivariance of formal power-series evaluation. A fifth part shows that the Hahn summation cannot be topologized: for a densely ordered exponent group every ring topology realizing all Hahn sums as limits is pulled back from the constants, so on $\mathbf{Oz}$ it sees only $\mathbb{Z}$ and on $\mathbf{No}$ it is indiscrete, and a set-sized full Hahn field has a Hausdorff one exactly when its exponent group is 0 or $\mathbb{Z}$. A sixth part recovers the coefficients after adjoining external monomials of any rank: $\exists y (y^2 = 1 + x^4)$ defines the coefficient field of $\text{Frac}(F[G])$ for $F = \mathbf{No}, \mathbf{No}(i)$, a sparse power bound gives coefficient rigidity and cancellation at every finite rank, cancellation fails at infinite rank, and every field automorphism is monomial exactly when $\text{Hom}(G, \mathbb{Z}) = 0$ (with a published defect theorem).

This report merges eleven independently written manuscripts, prints shared results once with every source named, and keeps every source’s limitations. It is an AI-assisted draft: it has not been refereed, and nothing in it has been formalized in Lean.

Sources: manuscripts 09 (the base text), 04 and 02, pinned to repository commit [fb5c4b5](#), 08, pinned to [befe739](#), the later addition 10, pinned to [58cd8e1](#), the batch-30 additions 11, 12 and 13, pinned to [cf56b89](#), [a6c68ac](#) and [0865f04](#), the batch-31 addition 14, pinned to [3d40856](#), and the batch-32 additions 15, which names blobs of this report’s source at [b3fa9e2](#) and [20c4c9f](#), and 16, pinned to [1ab41af](#); their titles are in [sections 1.2, 20.1, 33.1, 38.1 and 43.1](#). Results already printed in the omnific Diophantine report [docs/surreal/omnific-diophantine-geometry](#) are cited, not reprinted. The provenance of every section is recorded in [section A](#); non-claims are collected in [section B](#). Finite checks accompany the report and check finite identities only ([section C](#)).

Contents

1	Scope, results, and conventions	5
1.1	Principal results	5
1.2	Provenance of this report	7
1.3	Imported results	11
1.4	Class size and foundations	12
1.5	Notation and conventions	12
I	Automorphisms preserving the omnific integers	13
2	Hahn support, the boundary ring, and strong maps	14
2.1	Summability	14
2.2	The rings and the zero cut	14
3	The convex-support criterion	15
3.1	Two coefficient functionals	15
3.2	The criterion	16
3.3	The support-cut form and a second route [10]	18
4	Shears, flows, and the rank dichotomy	20
4.1	Homogeneous shears	20
4.2	Families of shifts and a proved correspondence	21
4.3	Single-shift flows without divisibility	22
4.4	Why translations fail	23
4.5	The rank dichotomy and the bottom subgroup	23
4.6	Phase shears	24
4.7	Twists with a general profile [10]	24
5	The strong stabilizer and all pair automorphisms	26
6	Convex rank, solvability, and a wreath product	27
6.1	The bracket law	27
6.2	A filtration by convex scales	28
6.3	Sharpness, including the abstract group	28
6.4	Metabelian and not nilpotent: one group, printed once	29
7	Fixed fields and arbitrarily large stabilizers	31
7.1	Exponent coefficient functionals	31
7.2	Relative fixed fields and the Hahn hull	31
7.3	Product actions and nonsolvable stabilizers	32
7.4	A definable-closure bound in set-sized Hahn fields [10]	32
8	What the pair (No, Oz) remembers	33
9	What the pair forgets: monomials and representatives	34
9.1	No definable representative rule	35
9.2	Two-term witnesses, simplicity and the exponential [10]	36
10	Surcomplex and Gaussian consequences	39
11	Scope, answered questions, and further questions	41
11.1	What is settled, and status in other reports	41
11.2	Further questions	42
II	Algebraic-parameter rigidity of the omnific integers	43

12 Conventions and the root-covered property	44
12.1 Conventions of this part	44
13 The monomial-divisibility criterion	44
14 Finite-type domains carry no varying embedding	45
15 Polynomial and Laurent coefficient rigidity	46
16 Locally nilpotent derivations, coactions and gradings	48
17 Euler derivations and formal flows	48
17.1 The monomial orbit field	49
17.2 An omnific family of commuting directions	50
18 Set-sized workspaces and a fraction-field warning	51
19 Relation to Part I and the questions of this part	51
 III Automatic strongness and proper embeddings of the omnific pair	 52
20 Sources, results and conventions of this part	52
20.1 Three sources and where the merge chose	52
20.2 What this part settles	54
20.3 Conventions and notation of this part	54
21 Constant-term tests detect Hahn summability	56
22 Constant-term-covariant isomorphisms are strong	58
23 Automatic strongness of the omnific stabilizer	59
23.1 What the integer part reconstructs	59
23.2 The main theorem	59
24 The whole stabilizer	61
25 Constant-term adjoints and a proper-class obstruction	62
26 The bottom-gap classification of exact-monomial embeddings	64
26.1 Taylor sections and the rigidity boundary	66
27 Explicit proper embeddings of (No, Oz)	67
27.1 Two levels of normal-form substitution	67
27.2 One map, three instances	68
28 Closed images and the two ends of the value group	69
29 Constants, elementarity, conjugacy and parameters	70
29.1 Elementary reduct, nonelementary expansion	70
29.2 Continuum many copies separated by the omnific integers	71
29.3 Proper copies fixing prescribed parameters	72
30 Surcomplex and Gaussian versions	73
31 Boundaries of the method	75
32 Status of the questions, and questions of this part	76
32.1 Answered and re-scoped questions	76
32.2 Questions	76

IV A fourth derivation: induced matchings, detector subfields and formal evaluation	81
33 Source 14, what it re-derives, and its notation	81
33.1 The manuscript and its pin	81
33.2 Printed once	81
33.3 Conventions and notation of this part	83
34 Induced matchings: a third route to scalar detection	84
35 Detection in subfields with enough detectors	85
36 Further consequences	86
37 Questions of source 14, and stale statements	88
37.1 Status of the source's questions	88
37.2 Stale and checked statements	89
V Topological collapse of Hahn summation	89
38 Source 15, what it re-derives, and its notation	90
38.1 The manuscript and what it read	90
38.2 Printed once	90
38.3 Conventions and notation of this part	92
39 Hahn-compatible ring topologies	93
40 The algebraic half: a disjoint-row variant and four additions	97
41 A pairing topology that realizes exactly the Hahn sums	99
42 Questions of source 15, and stale statements	100
42.1 Status of the source's questions	100
42.2 Stale and checked statements	101
VI Coefficient recovery in monomial extensions of arbitrary rank	102
43 Source 16, its relation to Part II, and its notation	102
43.1 The manuscript and its pin	102
43.2 What it generalizes, and what is printed once	103
43.3 Conventions and notation of this part	103
44 Finite-support monomial extensions	105
45 A quartic equation recovers the coefficient field	106
46 Sparse powers under arbitrary monomial extensions	108
47 Root-covered rings at arbitrary rank	109
48 Automorphisms of group and monoid algebras	110
49 Rational group fields: an exact automorphism criterion	111
50 Finite-rank cancellation and the exact infinite-rank failure	112
51 Boundaries of the method	113

52 Questions of source 16, relations, and stale statements	114
52.1 Status of the source's questions	114
52.2 Relations and stale statements	116
A Provenance ledger	116
A.1 Section by section	116
A.2 Printed once, and where the merge chose	119
A.3 Corrections and stale statements	121
A.4 Symbols renamed from the sources	123
B Non-claims, by source	124
C Finite checks	133
References	136

1 Scope, results, and conventions

A Conway normal form carries three kinds of information: coefficients, exponents, and the monomials that represent the exponents. The omnific integers keep the purely infinite part of a normal form and an ordinary integer constant coefficient. This report asks how much of the normal-form presentation is determined by the surreal field together with this distinguished ring, and which automorphisms preserve it.

The answer is asymmetric. The coefficient field, the cut between negative and nonnegative Hahn exponents, the constant coefficient and the floor map are recoverable from the pair $(\mathbf{No}, \mathbf{Oz})$; much of this was already proved in the omnific Diophantine report (section 8). The monomial cross-section is not recoverable. Even an arbitrary *set* of named parameters leaves enough automorphisms to move a monomial to an infinite series and to defeat every definable choice of valuation representatives. The mechanism is a family of derivations that change a higher Archimedean scale by a lower one without ever moving a purely infinite term across the constant-term boundary. Part I classifies these automorphisms and derives the consequences. Part II, from a separate manuscript, shows the complementary rigidity: $\mathbf{Oz}$ admits no nonconstant *algebraic* family of embeddings, although it admits formal flows of all orders. Part III, from three later manuscripts, removes the strongness hypothesis from the classification for $(\mathbf{No}, \mathbf{Oz})$ and shows that the rigidity of automorphisms does not pass to proper embeddings. Part IV, from a ninth manuscript, derives the same automatic strongness a third time and adds a third detection proof and detection in subfields. Part V, from a tenth manuscript, contrasts this algebraic preservation of all Hahn sums with their topological realization: every jointly continuous ring topology that realizes them forgets Π . Part VI, from an eleventh, carries the coefficient rigidity of part II to monomial parameters of arbitrary rank.

1.1 Principal results

Bracketed numbers name the source manuscripts (section 1.2); [merge] marks statements assembled or generalized by the merge, each with a complete proof. The Hahn conventions are fixed in section 1.5: $K = k((t^\Gamma))$ with well-ordered supports, $t^\gamma = \omega^{-\gamma}$ on the surreals, Π the series supported in $\Gamma^{<0}$, $\mathcal{A}_k = k \oplus \Pi$ and $\mathcal{R}_\mathfrak{o} = \mathfrak{o} \oplus \Pi$, so that $\mathcal{R}_\mathbb{Z} = \mathbf{Oz}$ when $k = \mathbb{R}$ and $\Gamma = \mathbf{No}$.

- (1) **The convex-support criterion** [04, 09][merge]. For every characteristic-zero k , every unital $\mathfrak{o} \subseteq k$, every nonzero divisible Γ (set-sized, or $\Gamma = \mathbf{No}$), and every strong k -linear 1-automorphism σ : $\sigma(\mathcal{R}_\mathfrak{o}) = \mathcal{R}_\mathfrak{o}$ if and only if the logarithmic-character coefficients $a_{\sigma, \delta}$ vanish on C_δ for every $\delta > 0$, if and only if σ commutes with the three zero-cut projections (theorem 3.3). The proof is 04's and imports no exponential-logarithm correspondence. For contracting strong derivations the analogous criterion holds with 09's coefficients c_δ (theorem 3.4); where the correspondence is available the operator form follows (corollary 3.5). The functionals $a_{\sigma, \delta}$ and c_δ differ; their vanishing conditions are equivalent (example 3.2). For single fixed-shift flows 08 proves the criterion without divisibility (theorem 4.10). Source 10 proves it again in an equivalent form, on the support of the principal unit $t^{-g}\sigma(t^g)$ rather than its logarithm, with a least-forbidden-shift proof kept as a second route; the inclusion $\sigma(\mathcal{R}_\mathfrak{o}) \subseteq \mathcal{R}_\mathfrak{o}$ already suffices, every signed Archimedean block of exponents is preserved, and the necessity half needs no strongness (theorem 3.8, proposition 3.10, and remark 3.9).
- (2) **Rank and fixed fields** [04, 08, 09, 10]. The stabilizer $\mathcal{U}_{\mathcal{R}_\mathfrak{o}}(\Gamma)$ is trivial exactly when Γ is Archimedean (theorem 4.13); for set-sized Γ its common fixed field is the Hahn field on the bottom convex subgroup (theorem 4.15); and $\text{Fix}(\mathcal{U}_{\mathbf{Oz}}) = \mathbb{R}$, also for all field

- automorphisms of **No** preserving **Oz** (theorem 7.2). For every set A of parameters the elementary shears fixing A have common fixed field $\mathbb{R}((t^{V_A}))$, V_A the rational span of the supports (theorem 7.4). Binomial twists $t^g \mapsto t^g F(t^\delta)^{\theta(g)}$ with an arbitrary profile F have fixed field $k((t^{\ker \theta}))$, no nontrivial finite orbits and a relatively algebraically closed fixed field (theorems 4.18 and 4.20 and corollary 4.21), and in set-sized fields definable elements obey a support bound (proposition 7.8) [10].
- (3) **Group structure** [04, 08, 09]. In finite ordered rank r the stabilizer and its Lie algebra have derived length at most $r - 1$, with equality for $\mathbb{Q}^r$ (theorems 6.3 and 6.6). Two successive-scale flows generate a faithful $\mathbb{Z} \wr \mathbb{Z}$ with exact commutator leading terms (theorems 6.8 and 6.10). For every set P of surreal parameters, the pointwise stabilizer of P in $\mathcal{U}_{\mathbf{Oz}}$ carries a faithful action of the full product $\mathbb{R}^\kappa$ for every set cardinal κ and is nonsolvable (theorems 7.6 and 7.7).
 - (4) **What the pair remembers** [08, 09]. Beyond the reconstruction package of the Diophantine report, the three zero-cut projections P_-, P_0, P_+ on all of **No**, the floor map and truncation at a named monomial are parameter-free definable in $(\mathbf{No}, \mathbf{Oz})$, and every field automorphism preserving **Oz** commutes with them, without strongness (theorem 8.2 and corollary 5.2).
 - (5) **What it forgets** [04, 08, 09, 10]. For every set P of parameters, some strong 1-automorphism preserving **Oz** fixes P pointwise and moves a Conway monomial to an infinite series; so neither the monomial class nor the omega-map is P -definable (theorem 9.1), and no P -definable rule selects a representative of each natural-valuation class (theorem 9.4). Source 10 moves ω^{ω^β} to the two-term omnific integer $\omega^{\omega^\beta} + \omega^{\omega^\beta - \delta}$ and shows that neither the simplicity relation nor the Gonshor exponential is P -definable, even with all reals and the zero-cut projections named (theorems 9.8 and 9.10).
 - (6) **Surcomplex consequences** [04, 08, 09, 10]. The criterion holds for $\mathbf{Oz}[i]$; real shears commute with conjugation, and 10's two-term witnesses also preserve the order of the real axis and the projections (proposition 10.5); an imaginary-time shear fixing any set moves the real axis inside the leading-term kernel (theorem 10.2); a wild coefficient lift moves an ordinary real (proposition 10.3); and the common fixed field of all automorphisms preserving $\mathbf{Oz}[i]$ is $\mathbb{Q}$ (corollary 10.4).
 - (7) **Algebraic-parameter rigidity** [02]. Every nonzero element of **Oz** or $\mathbf{Oz}[i]$ divides a monomial whose roots of all orders lie in the ring (corollary 13.3). Consequently injective homomorphisms into polynomial and Laurent rings over a domain land in the coefficients (theorems 15.1 and 15.3), polynomial cylinders over **Oz** recover **Oz** under every isomorphism (theorem 15.4), $\text{ML}(\mathbf{Oz}[\mathbf{T}]) = \mathbf{Oz}$ (theorem 16.2), and embeddings of set-sized root-covered rings into finite-type domains land in a finite field of algebraic constants (theorem 14.4). Commuting derivations of **Oz** integrate to formal flows of all orders that no polynomial family realizes, with an exact transcendence degree for their monomial orbit fields (theorems 17.3 and 17.5).
 - (8) **Automatic strongness and proper embeddings** [11, 12, 13; 14]. A set-indexed family in a full Hahn field is summable exactly when, for every y , only finitely many $\text{ct}(x_j y)$ are nonzero, and countable $\{0, 1\}$ -valued tests suffice; so a field isomorphism of full Hahn fields commuting with ct up to a coefficient isomorphism is strong (theorems 21.4 and 22.1). Every ring automorphism of **Oz** extends uniquely to a strongly $\mathbb{R}$ -linear automorphism of **No** (theorem 23.2), which answers question 11.1; every automorphism of $(\mathbf{No}, \mathbf{Oz})$, or of a full real Hahn pair with divisible exponent group, is $M_{\chi, \tau} \circ u$ with u as in theorem 3.3 (theorem 24.1). Strong exact-monomial embeddings preserving the integer parts correspond

to coefficient sections into the Hahn field on the bottom gap, and move reals exactly when the positive value image is not cointial ([theorem 26.3](#) and [corollary 26.7](#)); explicit proper embeddings preserve and reflect $\mathbf{Oz}$ and send a real b to $b + \omega^{-1}$ ([theorem 27.3](#)). Further: constant-term adjoints at set size and a proper-class obstruction, closed images and a continuity dichotomy, nonelementarity, continuum many nonconjugate copies, parameter-fixed copies and surcomplex versions ([sections 25](#) and [28 to 30](#)).

- (9) **A fourth derivation** [14]. Source 14 proves the automatic strongness of the real pair again, in less generality, and is credited beside 12 and 13. It adds a third proof of detection: an induced matching yields a detector with all coefficients 1, so monomials and all-ones series on increasing sequences suffice ([lemma 34.1](#) and [corollary 34.2](#)). Detection and transport of sums hold in every subfield containing those series ([proposition 35.2](#)), but not in the Puiseux field ([example 35.3](#), [merge]). Pair automorphisms commute with formal power-series evaluation at infinitesimals ([theorem 36.1](#)), the adjoint is a sign-reversed transpose ([proposition 36.2](#)), and an extension of a ring automorphism of $\mathcal{R}_\Gamma$ is unique ([proposition 36.5](#), [merge]).
- (10) **Topological collapse** [15]. For a densely ordered Γ , the ring topologies on $\mathfrak{o} \oplus \Pi$ with jointly continuous multiplication in which every Hahn sum is the limit of its finite partial sums are exactly the pullbacks along ct of the ring topologies on $\mathfrak{o}$ ([theorem 39.4](#)); so on $\mathbf{Oz}$ and $\mathbf{Oz}[i]$ they see only $\mathbb{Z}$ and $\mathbb{Z}[i]$, countable monomial sums already force this, and on $\mathbf{No}$ and $\mathbf{No}(i)$ only the indiscrete topology qualifies ([theorem 39.7](#), [proposition 39.8](#), and [corollary 39.9](#)). A set-sized full Hahn field has a Hausdorff such topology exactly when $\Gamma = 0$ or $\Gamma \cong \mathbb{Z}$ ([theorem 39.12](#)). The congruence topologies already in the collection are instances ([remark 39.6](#), [merge]). Source 15 also re-derives automatic strongness a fourth time, credited in [part III](#); describes the strong $\mathbb{R}$ -linear dual of $\mathbf{No}$ by class coefficient functions ([proposition 40.5](#)); and gives a pairing topology realizing exactly the Hahn sums, with separately but not jointly continuous multiplication ([theorems 41.1](#) and [41.3](#)).
- (11) **Coefficient recovery at arbitrary rank** [16]. For a torsion-free abelian group G and F real closed or algebraically closed of characteristic zero, $\exists y (y^2 = 1 + x^4)$ defines F in $F(U^G) = \text{Frac}(F[U^G])$, and a denominator-cleared form defines $\mathbf{Oz}$ and $\mathbf{Oz}[i]$ in their monoid algebras ([theorem 45.4](#)). A nonmonomial P/Q that is an n th power in any monomial extension has $n \leq \max(\text{wt } P, \text{wt } Q) - 1$, sharply ([theorem 46.2](#)). Consequently [theorems 15.1](#) and [15.3 to 15.5](#) extend to monoid algebras of arbitrary rank, cancellation of finite rank ([theorems 47.1](#), [48.1](#) and [50.1](#)), with exact failure for $\mathbb{Q}^{(\kappa)}$, κ infinite ([theorem 50.4](#)); and, granted a defect theorem of Lin and Wang, every field automorphism of $F(U^G)$ is monomial exactly when $\text{Hom}(G, \mathbb{Z}) = 0$ ([theorem 49.2](#)). These are finite-support extensions, not Hahn fields ([example 51.1](#)).

These statements are proved here, but proving a statement here is not a claim that nobody proved it earlier. The general theory of strong Hahn-field automorphisms and of the exponential-derivation correspondence is prior work ([section 1.3](#)); each source names its most distinctive contributions, and no exhaustive priority claim is made ([section B](#)).

1.2 Provenance of this report

The report is built from eleven manuscripts written independently: five for [parts I](#) and [II](#), described here, three for [part III](#), described in [section 20.1](#), one for [part IV](#), described in [section 33.1](#), and one each for [parts V](#) and [VI](#), described in [sections 38.1](#) and [43.1](#). Four were placed together in commit `f4c9504` (batch 26, items 02, 04, 08 and 09; they keep these numbers here, as their shipped file prefixes do). Manuscripts 02, 04 and 09 pin repository

commit `fb5c4b53` (12 commits before the placement); manuscript 08 pins `befe739b` (10 commits before). Both pins contain the batch-24 placement `be06fc8` but precede the batch-24 and batch-25 writes, so no source saw the reconstruction package now in the omnific Diophantine report (its section `odg:def:sec:reconstruction`, placed after both pins). The fifth, manuscript 10, is item 05 of batch 28, placed as an addition in commit `c6359e4` under the file prefix `10-support-cut-`. It pins `58cd8e1`, a commit on a line that does not contain `f4c9504`: its author never saw this report, and its repository comparison predates it. Its material carries the label sub-prefix `opa:sc:`.

Source	Manuscript	Principal contribution to this report
09	<i>Omnific-Preserving Automorphisms: convex-scale stabilizers, definable constants, and nondefinable monomials</i>	Base text and structure; the criterion for any characteristic-zero k and unital $\mathfrak{o}$; the derivation form with c_δ ; the Archimedean dichotomy; the strong factorization; the finite-rank derived-length bound and its sharpness for the abstract group; set-parameter nondefinability and the absence of any definable representative rule; $\mathbb{R}^\kappa$ actions and nonsolvable stabilizers; projections and truncation.
04	<i>Automorphisms Preserving the Omnific Integers: a convex-support criterion, fixed fields, and nondefinability of Conway monomials</i>	The import-free proof of the criterion through the logarithmic character (the proof of record here); exact displacement and fixed fields; the bottom subgroup and the relative Hahn hull; the common-positive-shift exponential-logarithm correspondence; the explicit commutator formula; the one-term class and the RV sort; the real axis inside the leading-term kernel.
08	<i>What the Omnific Integer Part Remembers: coefficient reconstruction, convex-scale automorphisms, nondefinable monomials, and an explicit wreath product</i>	The divisibility-free threshold for single fixed-shift flows; phase shears; the faithful $\mathbb{Z} \wr \mathbb{Z}$ and exact commutator leading terms; the floor map and the order formula on $\mathbf{Oz}$; the wild coefficient lift and the Gaussian fixed field $\mathbb{Q}$.
02	<i>Algebraic-Parameter Rigidity of the Omnific Integers: strong cancellation, finite-type embeddings, and nonalgebraizable formal flows</i>	All of part II .

Source	Manuscript	Principal contribution to this report
10	<i>Omnific Integers Do Not Determine Surreal Monomials: exact support-cut stabilizers, fixed fields, and parameterwise nondefinability</i>	A fifth proof of the criterion, in the support-cut form, with the least-forbidden-shift argument as a second route; necessity without strongness; Archimedean blocks; twists with a general profile, their fixed fields, finite orbits and relative algebraic closedness; the set-sized definable-closure bound; the two-term witness; nondefinability of simplicity and of the Gonshor exponential; the question on other integer parts.
11	<i>Coefficient Gaps and Proper Self-Embeddings of the Surreal Field: omnific preservation, topological dichotomies, and nonconjugate copies</i>	Base of the embedding sections of part III : the bottom-gap classification, Taylor admissibility, closed images and the continuity dichotomy, nonelementarity, nonconjugate copies, parameter-fixed copies, conjugation-compatible surcomplex embeddings.
12	<i>Automatic Strongness and Proper Embeddings of the Omnific Integers: constant-term duality, full stabilizers, and the surreal–surcomplex boundary</i>	Base of the automatic-strongness sections of part III : scalar detection, the constant-term isomorphism theorem, automatic strongness of $\text{Aut}(\mathbf{Oz})$, the factorization at set size and the Archimedean corollary, adjoints and retractions, the proper-class functional, regular cardinals.
13	<i>Automatic Hahn Linearity of Omnific Isomorphisms: residue duality, full stabilizers, and coefficient-drifting self-embeddings</i>	The same main theorems as 12, printed once; its countable binary witness, isomorphisms between different groups, the coefficient-matrix formula, the example that constant-term preservation does not preserve $\mathbf{Oz}$, the small-integer-part example and the failure of pair homogeneity.
14	<i>Automatic Summability from Omnific Arithmetic: constant-term duality, the full omnific stabilizer, and a sharp set–class distinction</i>	A third derivation of 12’s and 13’s main theorems, printed once with them; all of part IV : the induced-matching route to detection, detection in subfields with the countable detector property, formal evaluation, the general adjoint matrix, the case $\Gamma = \mathbb{R}$ and one new question.
15	<i>Algebraic Rigidity and Topological Collapse of Surreal Arithmetic: Hahn summation, omnific integer topologies, and the discrete-exponent boundary</i>	All of part V : the classification of Hahn-compatible ring topologies on $\mathfrak{o} \oplus \Pi$ for dense Γ , the omnific and field specializations, the cyclic boundary for full Hahn fields, a disjoint-row detector with interpolation, the class description of the strong dual of $\mathbf{No}$, the pairing topology and six new questions; a fourth derivation of 12’s and 13’s main theorems, printed once with them.

Source	Manuscript	Principal contribution to this report
16	<i>Recovering Surreal Coefficients from Monomial Extensions: quartic definitions, arbitrary-rank rigidity, and sharp cancellation</i>	All of part VI : the quartic coefficient definitions, the sparse power bound, arbitrary-rank coefficient rigidity and automorphism formulas, the $\text{Hom}(G, \mathbb{Z}) = 0$ criterion, the finite/infinite cancellation threshold and ten new questions; it generalizes theorems 15.3 to 15.5 .

Manuscripts 04, 08 and 09 prove one classification at three generalities ([section 3](#)) and the same consequences: rank triviality, fixed field $\mathbb{R}$, set-parameter nondefinability and a nonnilpotent stabilizer. They are printed once. 09 is the base because its hypotheses on $(k, \mathfrak{o})$ are the weakest and its group-theoretic results the widest; 04's proof is used for the criterion itself because it needs no imported correspondence, and 08's single-flow version is kept as a marked case because it needs no divisibility. Manuscript 10, added later, is a fifth derivation of the same classification under the same hypotheses ([section 3.3](#)). Their reconstruction and definability sections re-derive results now in the Diophantine report; those are cited, and only the remaining extras are printed ([section 8](#)). Manuscript 02 has a different subject, algebraic families rather than automorphisms preserving a support condition, and forms [part II](#) with the sub-prefix `opa:par:`; its relation to [part I](#) is recorded in [section 19](#). None of the four contradicts another or any report in the collection. [Section A](#) records, section by section, what each source proved, where the merge had to choose, and every symbol renamed from a source.

Manuscript 10 re-derives the criterion, the rank dichotomy, the strong factorization, the fixed field of a shear, set-parameter nondefinability of the monomials and the Gaussian witnesses. It has the hypotheses of [theorem 3.3](#) exactly: characteristic-zero k , an arbitrary unital $\mathfrak{o}$, divisible Γ , a set or **No**. None of this is a strengthening. These results are printed once, with 10 added to their sources. Its genuinely new material is appended at the end of the sections it belongs to, so that no existing number changes: [section 3.3](#) (support-cut form, second proof, blocks), [section 4.7](#) (general twists and their fixed fields), [example 6.13](#), [section 7.4](#), [section 9.2](#) (two-term witnesses, simplicity and the exponential), [proposition 10.5](#) and [question 11.7](#). 10 contradicts neither this report nor any other; its novelty statement, written before this report existed, is corrected in [section 11.1](#).

Manuscripts 11, 12 and 13, items 03, 04 and 05 of batch 30, were placed as an addition in commit 21375f8. They form [part III](#), placed after [part II](#) and before the appendices, so that no existing theorem, section or equation number changes; their labels carry the sub-prefix `opa:as:`. 12 and 13 prove the same main theorems and are printed once, with 12 as the base; 11 is the base of the embedding sections, which contain the embeddings of 12 and 13 as instances. Where the merge chose is recorded in [section 20.1](#). None of the three contradicts this report or another; they answer [question 11.1](#), and the passages of [part I](#) that recorded it as open now carry status notes.

Manuscript 14, item 07 of batch 31, was placed as an addition in commit 9d28e28. It pins 3d40856, which contains the batch-26 text of this report but none of the material of sources 10–13, and it proves the main theorems of 12 and 13 again, for $\Gamma = \Delta$ divisible. Those results are printed once in [part III](#), with 14 added to their sources; its genuinely new material forms [part IV](#), placed after [part III](#) and before the appendices, so that no existing number changes, with the sub-prefix `opa:cm:`. It contradicts neither this report nor any other; its statement

that it removes the strongness hypothesis was accurate at its pin and is stale at the merge (section 37.2).

Manuscripts 15 and 16, items 03 and 08 of batch 32, were placed as additions in commit 7d04483. They form parts V and VI, placed after part IV and before the appendices, so that no existing number changes, with the sub-prefixes `opa:tc:` and `opa:cr:`. 15 names no commit; it read this report's source as written in b3fa9e2 and then in 20c4c9f, so it saw part III but not part IV. Its topological results are new; its algebraic half re-derives part III and is printed there once, with 15 added to the sources (section 38.2). 16 pins 1ab41af, which contains part III but not part IV; its prior manuscript is source 02, and it generalizes part II to arbitrary rank (section 43.2). Neither contradicts this report or another report.

1.3 Imported results

Conway normal forms, real closedness of **No** and the omnific integers are classical [6, 10]; factorization theory of **Oz** is due to L'Innocente and Mantova [14]. The monomial/character/kernel decomposition of valued Hahn automorphisms is due to Kuhlmann and Serra [13], and its surreal form, with the class-size conventions used here, to Kaplan, Krapp and Serra [12]; their rigidity of order together with simplicity [12, Proposition 5.1] is used in theorem 9.10. Ehrlich and Kaplan's initial subdomains and canonical integer part [7] are a different notion from a truncation-closed subring or a subring stabilized by an automorphism, and none is substituted for another here [10]. The collection's surcomplex automorphism report (`docs/surcomplex/surcomplex-field-automorphisms`) already constructs the fixed-shift derivations and flows used below, in greater generality (its `saut:thm:shiftflow`), and the four-layer decomposition of valued automorphisms (`saut:thm:decomp`).

The correspondence between contracting strong derivations and strong 1-automorphisms is due to Bagayoko, Krapp, Kuhlmann, Panazzolo and Serra.

Cited theorem 1.1 ([2, Theorem 3.13]; [12, Fact 4.2]). *Let k have characteristic zero and Γ be an ordered abelian group. If ∂ is a contracting strongly k -linear derivation of $k((t^\Gamma))$, then*

$$\text{Exp}(\partial)(f) = \sum_{n \geq 0} \frac{\partial^n(f)}{n!}$$

is a strongly k -linear 1-automorphism, and $\partial \mapsto \text{Exp}(\partial)$ is a bijection between contracting strongly k -linear derivations and strongly k -linear 1-automorphisms, with inverse $\text{Log}(\sigma) = \sum_{n \geq 1} (-1)^{n+1} (\sigma - \text{id})^n / n$. Composition corresponds to the Baker–Campbell–Hausdorff operation.

Remark 1.2 (What is proved where). [2] proves the correspondence for set-sized Γ . Source 09 attributes the surreal case to [12, Fact 4.2]; the placement dossier, reading an HTML rendering, doubted that this Fact states the bijection. The merge re-read the PDF of [12] (version 3): Fact 4.2 does state the bijection, cites [2, Theorem 3.13], and adds that the correspondence “was established in the case that G is a set, but it also holds for G a proper class”, without a separate proof. So 09's citation is accurate, and for $\Gamma = \mathbf{No}$ the operator-logarithm statements rest on that assertion. The criterion itself does not: its proof (theorem 3.3) and 04's correspondence on the common-positive-shift class (theorem 4.8) are complete and elementwise, and the explicit constructions on **No** use one set-sized well-ordered set of shifts (proposition 4.4). Every statement below that uses cited theorem 1.1 says so.

1.4 Class size and foundations

Unless $\Gamma = \mathbf{No}$ is explicitly under discussion, exponent groups are sets, so fields and automorphism groups are ordinary sets. For $\Gamma = \mathbf{No}$ we work in NBG with global choice [04, 08, 09]; every individual Hahn support and every summation family is a set. An automorphism of $\mathbf{No}$ is an individual class map; no set or NBG class whose members are all such proper-class graphs is formed. A statement about $\mathcal{U}_{\mathbf{Oz}}$ is a statement about individual maps, their compositions, or a set-indexed action encoded by one class relation; this is the convention of [12]. A common-fixed-field statement is pointwise; nonsolvability means that no ordinary finite derived length annihilates all iterated commutators; valuation representatives are treated through an equivalence relation on nonzero elements, not through proper-class cosets [09]. A quotient notation G/H in a full-class application abbreviates an additive map on G vanishing on H [08]. Bases are chosen only for set-sized vector spaces (lemma 7.3).

1.5 Notation and conventions

Part I uses the small- t Hahn convention of sources 04 and 09. Part II keeps its source's large-monomial convention, which is the convention of the other omnific reports; see section 12.1. Part III uses the t -convention; its further symbols are fixed in section 20.3, and those of parts IV to VI in sections 33.3, 38.3 and 43.3; the monomials U^g of part VI are external indeterminates, not Hahn monomials. For actual surreals

$$t^\gamma = \omega^{-\gamma}. \quad (1.1)$$

Thus negative t -exponents are purely infinite terms and positive ones are infinitesimal. The tempting false reading is the opposite: in this Part, II is supported at *negative* t -exponents, and a “positive shift” $\delta > 0$ moves a term towards the infinitesimal side. Source 08 wrote in the ω -convention with decreasing supports; its statements have been translated, and its derivation $D_{\varphi, \delta}$ with $D_{\varphi, \delta}(\omega^g) = \varphi(g)\omega^{g-\delta}$ is $D_{\delta, \theta}$ below with $\theta(\gamma) = \varphi(-\gamma) = -\varphi(\gamma)$.

Notation	Meaning in part I, and the sources' symbols
$K = k((t^\Gamma))$	Hahn field, k of characteristic zero; $v(f) = \min \text{supp } f$, $v(0) = \infty$; $\text{lc}(f)$ the coefficient at $v(f)$.
Π	Series supported in $\Gamma^{<0}$, zero included (the purely infinite ideal). Sources: $\mathcal{J}$ (08, 09), $\mathcal{I}$ (04).
$\mathcal{A}_k = k \oplus \Pi$	Series supported in $\Gamma^{\leq 0}$, the boundary ring. Sources: B (09), $\mathcal{A}$ (04), $\mathcal{B}_k$ (08); the Diophantine report's $\mathcal{A}_{\mathbb{k}}(\Gamma)$.
$\mathcal{R}_{\mathbf{o}} = \mathbf{o} \oplus \Pi$	The integer-part ring for a unital subring $\mathbf{o} \subseteq k$; $\mathcal{R}_{\mathbb{Z}} = \mathbf{Oz}$ for $(\mathbb{R}, \mathbf{No})$ and $\mathcal{R}_{\mathbb{Z}[i]} = \mathbf{Oz}[i]$ for $(\mathbb{C}, \mathbf{No})$. Sources: A_D (09), O_D (04), A_k (08); the Diophantine report's $\mathcal{R}_{\mathbf{o}}(\mathbb{k}, \Gamma)$. The letter D of the sources is freed for derivations.
$\mathcal{O}, \mathfrak{m}$	Series supported in $\Gamma^{\geq 0}$ and $\Gamma^{>0}$: the valuation ring and its maximal ideal.
P_-, P_0, P_+	Projections onto Π , k and $\mathfrak{m}$ in $K = \Pi \oplus k \oplus \mathfrak{m}$; P_0 restricted to $\mathcal{A}_k$ is the retraction ct .
C_δ	Principal convex subgroup $\{\gamma : \gamma \leq n\delta \text{ for some } n \geq 1\}$ of $\delta > 0$. Sources: C_δ (04, 09), H_δ (08), $H(a)$ (02); the transcendence report's $C(a)$.

Notation	Meaning in part I , and the sources' symbols
$\delta \ll \epsilon$	$n\delta < \epsilon$ for every ordinary $n \geq 1$.
H_g	$\{h : n h < g \text{ for all } n \geq 1\}$ for $g \neq 0$ [04]; the same convex subgroup as the quotient report's H_a with $a = g $.
$a_{\sigma,\delta}$	Logarithmic-character coefficient $[t^\delta] \log(t^{-g}\sigma(t^g))$ of a 1-automorphism [04].
c_δ	Operator coefficient $[t^{\gamma+\delta}] \partial(t^\gamma)$ of a derivation ∂ [09]; for $\partial = \text{Log } \sigma$ it differs from $a_{\sigma,\delta}$ (example 3.2).
$D_{\delta,\theta}$	The derivation $t^\gamma \mapsto \theta(\gamma)t^{\gamma+\delta}$ for additive $\theta : \Gamma \rightarrow k$ and $\delta > 0$. Sources: $D_{\delta,\theta}$ (09), $D_{\delta,a}$ (04), $D_{\varphi,\delta}$ (08, sign as above).
$\mathcal{U}_{\mathcal{R}_\circ}(\Gamma), \mathcal{U}_{\mathbf{Oz}}$	Strong k -linear 1-automorphisms preserving $\mathcal{R}_\circ$ (independent of $\mathfrak{o}$ by theorem 3.3). Sources: $\mathcal{U}_{A_D}$ (09), $\mathcal{U}_D(G)$ (04).
$q_\beta(\gamma)$	The coefficient of ω^β in the Conway normal form of the <i>exponent</i> $\gamma \in \mathbf{No}$. Sources: q_η (09), $[\omega^\beta]g$ (04), c_a (08), χ_b (02), ℓ_η (quotient report).
$\exp(u), \text{Exp}, \text{Log}$	The Hahn series $\sum u^n/n!$ for $v(u) > 0$ (E in 08); operator exponential and logarithm.

Two further collisions are resolved in the new material. Source 09's parameter monomial $H = \omega^\eta$ and its family H_j are written $\eta = \omega^\beta$ and η_j here, leaving H_g to 04's convex subgroup; and 08's automorphisms A_1, B_1 are written in 04's normalization A_s, B_u , with $A_1^{(08)} = A_{-1}$ and $B_1^{(08)} = B_{-1}$ ([section 6](#)). A 1-automorphism is a field automorphism that preserves the valuation and the leading coefficient of every nonzero series. *Strong* means that every Hahn-summable family is sent to a summable family with the same sum; a strong automorphism is one whose inverse is also strong (04's convention); *k-linear* includes fixing k pointwise.

Source 10's symbols. Source 10 writes in the large-monomial convention $X^g = \omega^g$ with reverse-well-ordered supports, that is $X^g = t^{-g}$; its statements are translated into the t -convention, and a t -exponent $\gamma < 0$ is its exponent $g = -\gamma > 0$. Two of its letters collide with this report's, and the tempting false readings are wrong:

- 10's H_h , the principal convex subgroup generated by h , is C_δ here, while 10's C_g , the elements infinitesimal relative to g , is H_g here. The letters C and H are *swapped*: 10's C_g is not a principal convex subgroup, and H_δ here is not 10's H_h .
- 10's projection pr_+ onto the positive ω -exponents, the purely infinite part, is P_- here, and its pr_- is P_+ ; its ct is P_0 . The subscripts are reversed by the sign convention.

Further renamings: 10's positive-exponent ideal $\mathcal{I}$ is Π ; its $\mathfrak{m}$ is $\mathfrak{m}$, the same set; its $R_D = D + \mathcal{I}$ is $\mathcal{R}_\circ$ with $D = \mathfrak{o}$; its multiplier ring B is $\mathcal{A}_k$; its stabilizer $\mathcal{U}_D$ is $\mathcal{U}_{\mathcal{R}_\circ}(\Gamma)$; its shift h is δ , and its condition " $h \ll g$ " is $\delta \ll |\gamma|$; its principal unit $u_g = 1 + \varepsilon_g$ is $1 + \varepsilon_\sigma(-g)$ ([section 3.3](#)); its exponent functional $\lambda_b = [\omega^b] \cdot$ is q_b ; its additive profile P and its formal variable T are Θ and ξ , since P and T are taken here; its twist $\sigma_{\lambda,F,h}$ is $\sigma_{h,-\lambda}^F$ of (4.4), the sign coming from $g = -\gamma$; its class twist $\Sigma_{b,h}$ is $\Sigma_{b,\delta}$ of (9.2); its expanded structure $\mathcal{N}^\dagger$ is $\mathfrak{S}^\dagger$ of [theorem 9.10](#).

Part I

Automorphisms preserving the omnific integers

2 Hahn support, the boundary ring, and strong maps

2.1 Summability

An element of $K = k((t^\Gamma))$ is a formal sum $f = \sum_{\gamma \in S} a_\gamma t^\gamma$ with $S \subseteq \Gamma$ well ordered. A set-indexed family $(f_i)_{i \in I}$ is *Hahn summable* if the union of its supports is well ordered and each exponent lies in only finitely many of the supports; its sum is formed coefficientwise with finite sums in k . No convergence in the fine topology of the surreals is involved; a strong sum is a support-controlled algebraic construction [04, 08, 09].

We use the Neumann support principle in its word form: if S is well ordered and $T \subseteq \Gamma^{>0}$ is well ordered, then $S + \langle T \rangle_{\mathbb{N}}$ is well ordered, and each exponent receives only finitely many contributions from a choice of an element of S and a finite word in T , counted across all word lengths [2]. The one-shift case is proved directly.

Lemma 2.1 (One-shift summation [04, 08, 10]). *If $T \subseteq \Gamma$ is well ordered and $\delta > 0$, then $T + \mathbb{N}\delta$ is well ordered, and each $\gamma \in \Gamma$ has only finitely many representations $\gamma = g + n\delta$ with $g \in T$ and $n \in \mathbb{N}$.*

Proof. An infinite strictly decreasing sequence $g_j + n_j\delta$ has an infinite subsequence on which the n_j are nondecreasing; on it the g_j strictly decrease, contradicting well-ordering of T . Infinitely many representations of one γ would have unbounded, hence eventually increasing, values of n , and the corresponding $g = \gamma - n\delta$ would strictly decrease in T . $\square$

For $h \in \mathfrak{m}$ the series $\exp(h) = \sum_{n \geq 0} h^n/n!$ and $\log(1+h) = \sum_{n \geq 1} (-1)^{n+1} h^n/n$ are Hahn summable, and they are inverse group isomorphisms between $(\mathfrak{m}, +)$ and $(1 + \mathfrak{m}, \cdot)$. This local exponential is the only exponential used in the criterion; neither $\exp$ nor Exp asserts anything about the Gonshor exponential of $\mathbf{No}$.

Lemma 2.2 (Coefficient polynomial [04]). *For $h \in \mathfrak{m}$ and $\delta > 0$ the expression $P_{h,\delta}(X) = \sum_{n \geq 1} \frac{X^n}{n!} [t^\delta] h^n$ is a polynomial over k . For every $q \in \mathbb{Q}$ one has $P_{h,\delta}(q) = [t^\delta] \exp(qh)$, and its coefficient of X is $[t^\delta] h$.*

Proof. The support principle gives only finitely many contributions to the exponent δ across all powers h^n , not merely within each power. So only finitely many displayed terms are nonzero. The rest is coefficientwise. $\square$

2.2 The rings and the zero cut

The additive decompositions

$$K = \Pi \oplus k \oplus \mathfrak{m}, \quad \mathcal{A}_k = k \oplus \Pi, \quad \mathcal{R}_\circ = \circ \oplus \Pi, \quad \mathcal{O} = k \oplus \mathfrak{m} \quad (2.1)$$

define the projections P_-, P_0, P_+ [04, 09]. Membership in Π means that *all* exponents are negative, which is stronger than $v(f) < 0$: the series $t^{-1} + t$ has negative valuation and is not in Π [04]. Π is a nonunital ideal of $\mathcal{A}_k$ and of $\mathcal{R}_\circ$; $\mathcal{A}_k$ is not a valuation ring; P_0 is a ring retraction $\mathcal{A}_k \rightarrow k$ restricting to $\mathcal{R}_\circ \rightarrow \circ$, but not a ring map on K : $P_0(t^{-1})P_0(t) = 0 \neq 1 = P_0(1)$. For

$k = \mathbb{R}$, $\mathfrak{o} = \mathbb{Z}$ the ring $\mathcal{R}_{\mathbb{Z}}$ is an integer part of K ; for $\Gamma = \mathbf{No}$ it is $\mathbf{Oz}$ and its floor function is the Diophantine report's `odg:thm:floor`.

The units $\mathcal{A}_k^\times = k^\times$, and more generally the fact that a product of two elements of $\mathcal{A}_k$ lying in $k^\times$ has constant factors, are the Diophantine report's constant-product lemma (`odg:lem:constantproduct`, `odg:def:lem:units`); sources 04, 08 and 09 reprove it, and it is not reprinted.

Lemma 2.3 (The divisible part [04, 08]). *If $\bigcap_{n \geq 1} n\mathfrak{o} = 0$ (for example $\mathfrak{o} = \mathbb{Z}$ or $\mathbb{Z}[i]$), then $\Pi = \bigcap_{n \geq 1} n\mathcal{R}_{\mathfrak{o}}$, the largest divisible subgroup of $\mathcal{R}_{\mathfrak{o}}$; it is preserved by every ring automorphism of $\mathcal{R}_{\mathfrak{o}}$. This holds for every ordered abelian Γ , divisible or not.*

Proof. A series in Π is n times a series with the same support. Conversely the constant coefficient of an element of the intersection lies in $\bigcap_n n\mathfrak{o} = 0$. Every divisible subgroup lies in the intersection. $\square$

For $\mathbf{Oz}$ this is the identity `odg:eq:divkernel` of the Diophantine report; it is printed because the threshold theorem uses it for non-divisible Γ .

Definition 2.4 (Principal convex scale). For $\delta > 0$ let $C_\delta = \{\gamma \in \Gamma : |\gamma| \leq n\delta \text{ for some } n \geq 1\}$, and write $\delta \ll \epsilon$ if $n\delta < \epsilon$ for all $n \geq 1$. Then $\gamma \notin C_\delta$ if and only if $\delta \ll |\gamma|$; $C_\delta \subsetneq C_\epsilon$ if and only if $\delta \ll \epsilon$; and C_δ is divisible when Γ is. For $g \neq 0$ the set $H_g = \{h : n|h| < |g| \text{ for all } n \geq 1\}$ is a convex subgroup, and a finite sum of positive elements of H_g is still smaller than $|g|$ [04].

3 The convex-support criterion

3.1 Two coefficient functionals

Let σ be a strong k -linear 1-automorphism. Since $t^{-g}\sigma(t^g)$ has leading term 1, its local logarithm is defined [04]:

$$h_\sigma(g) = \log(t^{-g}\sigma(t^g)) \in \mathfrak{m}, \quad a_{\sigma,\delta}(g) = [t^\delta]h_\sigma(g) \quad (\delta > 0), \quad (3.1)$$

so that $\sigma(t^g) = t^g \exp(h_\sigma(g))$. For a contracting strong k -derivation ∂ (that is, $v(\partial f) > v(f)$ for $f \neq 0$) put [09]

$$c_\delta(\gamma) = [t^{\gamma+\delta}]\partial(t^\gamma) \quad (\delta > 0), \quad \text{so} \quad \partial(t^\gamma) = t^\gamma \sum_{\delta > 0} c_\delta(\gamma)t^\delta. \quad (3.2)$$

For fixed γ the nonzero terms have well-ordered set support; no assumption is made that all shifts together form one well-ordered set.

Lemma 3.1 (Additivity [04, 09]). *The maps $h_\sigma : \Gamma \rightarrow (\mathfrak{m}, +)$, $a_{\sigma,\delta}$ and c_δ are additive, hence $\mathbb{Q}$ -linear when Γ is divisible. If Γ is divisible and an additive $\theta : \Gamma \rightarrow k$ vanishes on $(-\delta, 0)$, it vanishes on C_δ .*

Proof. The quotient at $g+g'$ is the product of the quotients at g and g' , and $\log$ takes products to sums. For c_δ the product rule gives $t^{-\gamma-\eta}\partial(t^{\gamma+\eta}) = t^{-\gamma}\partial(t^\gamma) + t^{-\eta}\partial(t^\eta)$. For the last claim, given $0 \neq h \in C_\delta$ choose N with $|h|/N < \delta$; one of $\pm h/N$ lies in $(-\delta, 0)$, and additivity gives $\theta(h) = 0$. $\square$

Example 3.2 (The functionals differ [merge]). Let $\Gamma = \mathbf{No}$, $k = \mathbb{R}$, and $\partial = D_{1,q_1} + D_{\omega,q_2}$ (notation of section 1.5; the t -exponent γ has $q_1(\gamma)$ and $q_2(\gamma)$ as its coefficients at ω and ω^2). Take $\sigma = \text{Exp}(\partial)$, an element of $\mathcal{U}_{\mathbf{Oz}}$ by proposition 4.4, and $g = \omega^2$. Then $\partial(t^g) = t^{g+\omega}$ and

$\partial^2(t^g) = t^{g+\omega+1} + t^{g+2\omega}$, and the only word in the shifts $\{1, \omega\}$ reaching the exponent $\omega + 1$ applies the shift ω first; $\log(1 + u) = u - u^2/2 + \dots$ adds nothing at $\omega + 1$, since u^2 starts at 2ω . Hence

$$t^{-g}\sigma(t^g) = 1 + t^\omega + \frac{1}{2}t^{\omega+1} + (\text{terms at exponents } > \omega + 1), \quad a_{\sigma, \omega+1}(\omega^2) = \frac{1}{2},$$

whereas $c_{\omega+1}(\omega^2) = 0$: the operator coefficients of ∂ at g are $c_\omega(g) = 1$ and zero at every other shift. So $a_{\sigma, \delta} \neq c_\delta$ in general; 04's condition concerns the logarithm of the monomial image, 09's the logarithm of the operator. Both vanishing conditions hold here, and by [theorem 3.3](#) and [corollary 3.5](#) each is equivalent to preservation of $\mathbf{Oz}$, hence to the other. The placement dossier checked the example by exact computation.

3.2 The criterion

The following statement combines 09's generality with 04's proof. 04 proved it for $k = \mathbb{R}, \mathbb{C}$ and $\mathfrak{o} = \mathbb{Z}, \mathbb{Z}[i]$, passing through [lemma 2.3](#); 09 proved it for every characteristic-zero k and unital $\mathfrak{o}$, but through the operator logarithm ([cited theorem 1.1](#)). The first step below is 09's and removes the need for $\bigcap n\mathfrak{o} = 0$; the rest is 04's.

Theorem 3.3 (Convex-support criterion [04, 09, 10][merge]). *Let k have characteristic zero, let $\mathfrak{o} \subseteq k$ be a unital subring, let $\Gamma \neq 0$ be divisible (a set, or $\Gamma = \mathbf{No}$), and let σ be a strong k -linear 1-automorphism of $K = k((t^\Gamma))$. The following are equivalent:*

- (i) $\sigma(\mathcal{R}_\mathfrak{o}) = \mathcal{R}_\mathfrak{o}$;
- (ii) $\sigma(\mathcal{A}_k) \subseteq \mathcal{A}_k$;
- (iii) $\sigma(\Pi) \subseteq \Pi$;
- (iv) $a_{\sigma, \delta}(C_\delta) = 0$ for every $\delta > 0$;
- (v) for every $g \neq 0$, every exponent $\delta \in \text{supp } h_\sigma(g)$ satisfies $\delta \ll |g|$;
- (vi) σ commutes with P_- , P_0 and P_+ .

When they hold, $\sigma(\Pi) = \Pi$, $\sigma(\mathcal{A}_k) = \mathcal{A}_k$, $\sigma(\mathfrak{m}) = \mathfrak{m}$, the constant coefficient of every series is preserved, and every purely infinite monomial stays purely infinite, though not necessarily a monomial. In particular the group $\mathcal{U}_{\mathcal{R}_\mathfrak{o}}(\Gamma)$ does not depend on $\mathfrak{o}$.

Proof. (i) $\Rightarrow$ (ii) [09]. For $\gamma < 0$, $t^\gamma \in \mathcal{R}_\mathfrak{o}$, so $\sigma(t^\gamma) \in \mathcal{R}_\mathfrak{o} \subseteq \mathcal{A}_k$. An element of $\mathcal{A}_k$ is $c + \sum_{\gamma < 0} a_\gamma t^\gamma$ with $c \in k$; strong k -linearity writes its image as $c + \sum a_\gamma \sigma(t^\gamma)$, a summable family of series supported in $\Gamma^{\leq 0}$, whose sum is supported there.

(ii) $\Rightarrow$ (iv) [04]. Fix $\delta > 0$ and $0 \neq g \in C_\delta$; put $x = -|g| < 0$, so $h_\sigma(x) = \pm h_\sigma(g)$, and choose $N \geq 1$ with $|g| \leq N\delta$. By [lemma 2.2](#), $P(X) = [t^\delta] \exp(Xh_\sigma(x))$ is a polynomial. For rational $0 < q < 1/N$, divisibility gives $qx \in \Gamma^{< 0}$ and additivity gives $\sigma(t^{qx}) = t^{qx} \exp(qh_\sigma(x))$, whose coefficient at $qx + \delta$ is $P(q)$. Since $q|g| < \delta$, this exponent is strictly positive, while $\sigma(t^{qx}) \in \mathcal{A}_k$ has no positive exponents. So P has infinitely many zeros and vanishes; its linear coefficient is $a_{\sigma, \delta}(x)$, so $a_{\sigma, \delta}(g) = 0$. This tests every coefficient of the logarithm; testing only the first nonzero one would not suffice.

(iv) $\Leftrightarrow$ (v) [04]. $g \notin C_\delta$ means $\delta \ll |g|$; consider each nonzero coefficient.

(v) $\Rightarrow$ (iii) [04]. For $g < 0$, $\text{supp } h_\sigma(g) \subseteq H_g^{> 0}$, so every exponent of $\exp(h_\sigma(g))$ is 0 or a finite sum of elements of $H_g^{> 0}$, hence smaller than $|g|$; thus $\sigma(t^g) = t^g \exp(h_\sigma(g)) \in \Pi$. For $f \in \Pi$ write $\sigma(f) = \sum c_g \sigma(t^g)$ by strongness; all supports are negative.

(iii) $\Rightarrow$ (vi) [04]. A 1-automorphism maps $\mathfrak{m}$ into $\mathfrak{m}$, and σ fixes k ; so each summand of (2.1) is mapped into itself and σ commutes with the projections.

(vi) $\Rightarrow$ (i) [04]. If $\sigma(y) \in \Pi$ with $y = y_- + c + y_+$, then $0 = P_0\sigma(y) = c$ and $0 = P_+\sigma(y) = \sigma(y_+)$, so $y \in \Pi$. With (iii) this gives $\sigma(\Pi) = \Pi$, and since σ fixes $\mathfrak{o} \subseteq k$, $\sigma(\mathcal{R}_{\mathfrak{o}}) = \mathcal{R}_{\mathfrak{o}}$. The same argument gives $\sigma(\mathcal{A}_k) = \mathcal{A}_k$ and $\sigma(\mathfrak{m}) = \mathfrak{m}$, and (vi) preserves constant coefficients. Condition (ii) does not mention $\mathfrak{o}$, which proves the last assertion. $\square$

Only (ii) $\Rightarrow$ (iv) uses divisibility; it divides an *exponent*, while characteristic zero separately permits rational scalars [09]. The argument is elementwise with set-sized supports, so it applies verbatim to $\Gamma = \mathbf{No}$, $k = \mathbb{R}$, $\mathfrak{o} = \mathbb{Z}$, that is to $\mathbf{Oz}$, and to $\mathbf{Oz}[i]$ with $(k, \mathfrak{o}) = (\mathbb{C}, \mathbb{Z}[i])$ [04]. The theorem classifies *existing* strong maps: an arbitrary additive $h : \Gamma \rightarrow \mathfrak{m}$ need not define a strong surjective field map by $t^g \mapsto t^g \exp h(g)$ [04] (question 11.2).

Theorem 3.4 (Boundary-preserving derivations [09]). *Let $\Gamma \neq 0$ be divisible and let ∂ be a contracting strong k -derivation of K . Then $\partial(\mathcal{A}_k) \subseteq \mathcal{A}_k$ if and only if $c_\delta(C_\delta) = 0$ for every $\delta > 0$. In that case $\partial(\mathcal{A}_k) \subseteq \Pi$, $\partial(\Pi) \subseteq \Pi$, and ∂ commutes with P_- , P_0 , P_+ .*

Proof. If $\partial(\mathcal{A}_k) \subseteq \mathcal{A}_k$ and $-\delta < \gamma < 0$, then $t^\gamma \in \mathcal{A}_k$ and $\gamma + \delta > 0$, so $c_\delta(\gamma) = 0$; lemma 3.1 extends this to C_δ . Conversely, if $\gamma < 0$ and $c_\delta(\gamma) \neq 0$, then $\gamma \notin C_\delta$, so $-\gamma > n\delta$ for all n and in particular $\gamma + \delta < 0$. Every term of $\partial(t^\gamma)$ is negative; constants are killed; strongness sums over any negative support. Positive exponents stay positive under positive shifts, which gives the projection identities. $\square$

The condition excludes a boundary term as well as a crossing term: testing $-\delta/2$ gives $c_\delta(-\delta) = 0$. This matters when constant coefficients are restricted to $\mathfrak{o}$ rather than k [09]. Source 04 proves the same criterion for derivations with common positive shifts (theorem 4.7), testing t^{-u} for $0 < u \leq \delta$.

Corollary 3.5 (Operator form [09]). *Let σ be a strong k -linear 1-automorphism of the form $\sigma = \text{Exp}(\partial)$ for a contracting strong k -derivation ∂ with $\text{Log}(\sigma) = \partial$. Then the conditions of theorem 3.3 are equivalent to $\partial(\mathcal{A}_k) \subseteq \mathcal{A}_k$ and to $c_\delta(C_\delta) = 0$ for all $\delta > 0$. For set-sized Γ every strong k -linear 1-automorphism has this form, so*

$$\mathcal{U}_{\mathcal{R}_{\mathfrak{o}}}(\Gamma) = \text{Exp}(\mathfrak{L}_A), \quad \mathfrak{L}_A = \{\partial \text{ contracting strong } k\text{-derivation} : c_\delta(C_\delta) = 0 \text{ for all } \delta > 0\}.$$

For $\Gamma = \mathbf{No}$ the same equality holds if the correspondence of cited theorem 1.1 is granted for proper-class Γ , as [12, Fact 4.2] asserts (remark 1.2); on the common-positive-shift class it is proved in theorem 4.8.

Proof. If the conditions hold, $\sigma(\mathcal{A}_k) = \mathcal{A}_k$; then each $(\sigma - \text{id})^n$ maps $\mathcal{A}_k$ into $\mathcal{A}_k$, and $\text{Log}(\sigma)$ is a strongly summable family of series supported in $\Gamma^{\leq 0}$, so $\partial(\mathcal{A}_k) \subseteq \mathcal{A}_k$. This step is taken in the k -algebra $\mathcal{A}_k$, not in $\mathcal{R}_{\mathfrak{o}}$, so no division by integers inside $\mathfrak{o}$ is needed. Theorem 3.4 converts this into the coefficient condition. Conversely, under the coefficient condition every iterate of ∂ maps $\mathcal{A}_k$ into Π and kills k , so $\text{Exp}(\pm\partial)$ preserve $\mathcal{R}_{\mathfrak{o}}$. The last assertions are cited theorem 1.1. $\square$

The two routes are genuinely different. 04's functional is read off the monomial images and needs no correspondence; 09's is read off the generator and gives the Lie-theoretic structure of section 6. Neither condition is stated in terms of the other: each is equivalent to preservation of $\mathcal{R}_{\mathfrak{o}}$ (example 3.2).

Remark 3.6 (What has and has not been classified [04, 09]). The classification is inside the *strong, coefficient-fixing, leading-term-preserving* automorphisms. It does not assert that every abstract field automorphism preserving $\mathcal{R}_{\mathfrak{o}}$ is strong; for the real pair, fixing of $\mathbb{R}$ is automatic (corollary 5.2), strongness is not known to be. The coefficient functionals come from an existing strong map; arbitrary functions satisfying the vanishing condition cannot be assembled into an operator without checking summability and invertibility.

Status (batch 30). For the real pair, strongness is now known to be automatic as well: every field automorphism of **No** preserving **Oz**, and every automorphism of a full real Hahn pair with divisible exponent group, is strong ([theorem 23.2](#) and [corollary 23.4](#)), so the classification covers them all ([theorem 24.1](#)). The remark stands as written for a general $(k, \mathfrak{o})$ ([question 32.13](#)), and its last sentence is unaffected ([remark 24.2](#)).

3.3 The support-cut form and a second route [10]

Source 10 proves [theorem 3.3](#) a fifth time, without knowledge of this report and under the same hypotheses. Its condition bounds the support of the principal unit itself rather than of its logarithm, and its necessity proof isolates the least forbidden shift instead of testing a polynomial in a rational parameter. The two forms are equivalent, so the statement is printed once, as further equivalent conditions, and 10's proof is kept as a marked second route. For a strong k -linear 1-automorphism σ put

$$\varepsilon_\sigma(\gamma) = t^{-\gamma}\sigma(t^\gamma) - 1 \in \mathfrak{m}, \quad 1 + \varepsilon_\sigma(\gamma) = \exp(h_\sigma(\gamma)). \quad (3.3)$$

Multiplicativity makes $\gamma \mapsto 1 + \varepsilon_\sigma(\gamma)$ a character $\Gamma \rightarrow (1 + \mathfrak{m}, \cdot)$ [13]. For $u \in 1 + \mathfrak{m}$ and $q \in \mathbb{Q}$ let $u^q = \sum_{r \geq 0} \binom{q}{r} (u - 1)^r = \exp(q \log u)$; the sum is Hahn summable by the support principle, and $(u^q)^n = u^{qn}$ for integers n is a formal identity of binomial series [10].

Lemma 3.7 (Leading-one roots [10]). *If $u, w \in 1 + \mathfrak{m}$ and $u^n = w^n$ for some integer $n \geq 1$, then $u = w$. Hence, for a character $c : \Gamma \rightarrow 1 + \mathfrak{m}$ on a divisible Γ , $c(q\gamma) = c(\gamma)^q$ for all $q \in \mathbb{Q}$ and $\gamma \in \Gamma$.*

Proof. Otherwise $u/w = 1 + \eta$ with $0 \neq \eta \in \mathfrak{m}$, and the term of least exponent of $(1 + \eta)^n - 1$ is n times that of η , which is nonzero in characteristic zero. For $q = a/b$ with $b \geq 1$, both $c(q\gamma)$ and $c(\gamma)^q$ have b th power $c(a\gamma) = c(\gamma)^a$. $\square$

Theorem 3.8 (Support-cut form [10]). *Under the hypotheses of [theorem 3.3](#), each of the following conditions is equivalent to the conditions (i)–(vi) there:*

- (vii) $\sigma(\mathcal{R}_\mathfrak{o}) \subseteq \mathcal{R}_\mathfrak{o}$;
- (viii) *for every $\gamma < 0$, every $\delta \in \text{supp } \varepsilon_\sigma(\gamma)$ satisfies $\delta \ll |\gamma|$, that is $\text{supp } \varepsilon_\sigma(\gamma) \subseteq H_\gamma$;*
- (ix) $\text{supp } \varepsilon_\sigma(\gamma) \subseteq H_\gamma$ for every $\gamma \neq 0$.

In source 10's notation, (viii) reads: every shift $h > 0$ occurring in ε_g , for $g > 0$, satisfies $nh < g$ for every $n \geq 1$.

First route, from the proof of [theorem 3.3](#). (i) $\Rightarrow$ (vii) is trivial, and (vii) $\Rightarrow$ (ii) is the proof of (i) $\Rightarrow$ (ii) there, which uses only $\sigma(t^\gamma) \in \mathcal{R}_\mathfrak{o}$ for $\gamma < 0$. For fixed $\gamma \neq 0$ the series $\varepsilon_\sigma(\gamma) = \exp(h_\sigma(\gamma)) - 1$ has its support among the finite sums of elements of $\text{supp } h_\sigma(\gamma)$, and $h_\sigma(\gamma) = \log(1 + \varepsilon_\sigma(\gamma))$ has its support among the finite sums of elements of $\text{supp } \varepsilon_\sigma(\gamma)$. Since H_γ is a subgroup, (ix) is equivalent to condition (v) of [theorem 3.3](#); this is 10's logarithmic corollary. Finally (ix) $\Rightarrow$ (viii) is trivial, and (viii) $\Rightarrow$ (ix) because, for $\gamma > 0$, $1 + \varepsilon_\sigma(\gamma) = (1 + \varepsilon_\sigma(-\gamma))^{-1} = \sum_{n \geq 0} (-\varepsilon_\sigma(-\gamma))^n$ has its support among the finite sums of elements of $\text{supp } \varepsilon_\sigma(-\gamma) \subseteq H_{-\gamma} = H_\gamma$. $\square$

Second route: least forbidden shift [10]. We prove (vii) $\Rightarrow$ (viii) directly. Fix $\gamma < 0$ and write $\varepsilon_\sigma(\gamma) = \sum_{\delta \in \mathcal{S}} b_\delta t^\delta$ with $\mathcal{S} \subseteq \Gamma^{>0}$ well ordered and all $b_\delta \neq 0$. Suppose some $\delta \in \mathcal{S}$ lies outside H_γ , and let δ be the *least* such element; the elements of $\mathcal{S}$ below δ lie in H_γ . As $\delta \notin H_\gamma$ there is $N \geq 1$ with $|\gamma| \leq N\delta$. Choose a rational $0 < q < 1/N$; divisibility gives $q\gamma \in \Gamma$ with

$0 < q|\gamma| < \delta$, and [lemma 3.7](#) gives $t^{-q\gamma}\sigma(t^{q\gamma}) = (1 + \varepsilon_\sigma(\gamma))^q$. In this binomial series the linear term contributes qb_δ at the exponent δ . A term of degree $r \geq 2$ contributes there only through $\delta = \delta_1 + \dots + \delta_r$ with $\delta_j \in \mathcal{S}$; each δ_j is then smaller than δ , hence in H_γ , and so is their sum, which is impossible. So the coefficient of $\sigma(t^{q\gamma})$ at the exponent $q\gamma + \delta > 0$ is exactly $qb_\delta \neq 0$, whereas $t^{q\gamma} \in \Pi \subseteq \mathcal{R}_\mathfrak{o}$ and no element of $\mathcal{R}_\mathfrak{o}$ has a positive exponent. This contradicts (vii).

Conversely, under (viii) every exponent of $\sigma(t^\gamma)$, $\gamma < 0$, is γ or $\gamma + \delta$ with $\delta \ll |\gamma|$, hence negative; strongness gives $\sigma(\Pi) \subseteq \Pi$, which is (iii) of [theorem 3.3](#), and 10 concludes by the same injectivity and surjectivity argument as (iii) $\Rightarrow$ (vi) $\Rightarrow$ (i) there. $\square$

The two necessity proofs are genuinely different. That of [theorem 3.3](#) (from 04) shows, for each shift, that the coefficient of $\exp(qh_\sigma(x))$ is a polynomial in q with infinitely many zeros, and so tests every coefficient of the logarithm. 10's proof needs one rational q : at the least shift outside H_γ no nonlinear term of the binomial series can contribute. Both use divisibility of Γ only to form $q\gamma$, and neither uses $\mathfrak{o}$ beyond $\mathfrak{o} \subseteq k$. The inclusion form (vii) is implicit in the proof of [theorem 3.3](#) but not in its statement; 10 states it.

Remark 3.9 (Necessity without strongness [10]). The second route applies σ only to monomials, and uses only that σ is a field automorphism with $\sigma(t^\gamma) \in t^\gamma(1 + \mathfrak{m})$ for every γ : multiplicativity on monomials, [lemma 3.7](#), and $\sigma(t^{q\gamma}) \in \mathcal{R}_\mathfrak{o}$. So every k -fixing 1-automorphism σ of K with $\sigma(\mathcal{R}_\mathfrak{o}) \subseteq \mathcal{R}_\mathfrak{o}$, strong or not, satisfies (viii), and (ix) by inversion of units. What fails without strongness is the passage from monomials to arbitrary series, which the converse and all consequences for nonmonomial elements use. For $(\mathbf{No}, \mathbf{Oz})$ fixing of $\mathbb{R}$ is automatic ([corollary 5.2](#)), so every field automorphism of $\mathbf{No}$ that preserves $\mathbf{Oz}$ and the leading terms of monomials satisfies the monomial condition [10]. The merge notes that the necessity half of 04's route is equally monomial: it evaluates σ only at the monomials t^{qx} . This is partial information on [question 11.1](#), not an answer.

Status (batch 30). The remark is subsumed: [question 11.1](#) is now answered ([theorem 23.2](#)), every such automorphism of $(\mathbf{No}, \mathbf{Oz})$ is strong, and the passage from monomials to arbitrary series is automatic ([corollary 24.4](#)).

Proposition 3.10 (Signed Archimedean blocks [10]). *Call nonzero $\gamma, \gamma' \in \Gamma$ block-equivalent if they have the same sign and $|\gamma| \leq n|\gamma'|$, $|\gamma'| \leq n|\gamma|$ for some $n \geq 1$; the exponent 0 forms a block of its own. If σ satisfies the conditions of [theorem 3.3](#), every exponent of $\sigma(t^\gamma)$ lies in the block of γ . Consequently σ commutes with the projection onto the terms whose exponents lie in any prescribed union of blocks, and $\sigma(k((t^\Delta))) = k((t^\Delta))$ for every convex subgroup $\Delta \subseteq \Gamma$.*

Proof. By (ix) of [theorem 3.8](#), the exponents of $\sigma(t^\gamma)$ are $\gamma + \delta$ with $\delta = 0$ or $\delta \in H_\gamma^{>0}$, and then $2\delta < |\gamma|$. For $\gamma < 0$ this gives $|\gamma|/2 < |\gamma + \delta| \leq |\gamma|$, and for $\gamma > 0$ it gives $\gamma \leq \gamma + \delta < 3\gamma/2$; either way $\gamma + \delta$ is in the block of γ . If Λ is a union of blocks and $f = f_\Lambda + f'$ splits the terms of f according to Λ , strongness puts the support of $\sigma(f_\Lambda)$ inside Λ and that of $\sigma(f')$ outside it, so the projection of $\sigma(f)$ is $\sigma(f_\Lambda)$. A convex subgroup is a union of blocks, so σ maps $k((t^\Delta))$ into itself; σ^{-1} satisfies the same conditions, which gives equality. $\square$

The zero cut is a union of blocks, so this recovers commutation with P_-, P_0, P_+ ; [theorem 4.10](#) fixes $k((t^{C_\delta}))$ pointwise for single flows, whereas here the whole stabilizer preserves every convex Hahn subfield setwise. This is not preservation of every truncation: a twist may send t^γ to $t^\gamma + t^{\gamma+\delta}$ ([theorem 9.8](#)), and a cut strictly between the two exponents separates the two terms. The preserved blocks are determined by Archimedean scale, not by individual exponents [10].

4 Shears, flows, and the rank dichotomy

4.1 Homogeneous shears

Let $\theta : \Gamma \rightarrow (k, +)$ be additive and $\delta > 0$. The formula

$$D_{\delta, \theta} \left(\sum_{\gamma} a_{\gamma} t^{\gamma} \right) = \sum_{\gamma} a_{\gamma} \theta(\gamma) t^{\gamma + \delta} \quad (4.1)$$

defines a contracting strong derivation: translation keeps supports well ordered and families summable, and additivity of θ gives the product rule on monomials, hence on all series. This is the fixed-shift derivation of the surcomplex automorphism report, `saut:thm:shiftflow`, which constructs its exponential for every additive θ and every $\delta > 0$; its n th iterate has coefficient $\prod_{j < n} \theta(\gamma + j\delta)$. Sources 04, 08 and 09 re-derive it; only 04 credited it. The integer-part condition is the additional restriction $\theta(C_{\delta}) = 0$, not merely $\theta(\delta) = 0$.

Theorem 4.1 (Homogeneous shears [04, 09]; after `saut:thm:shiftflow`). *Suppose $\theta(C_{\delta}) = 0$. Then $D_{\delta, \theta}^n(t^{\gamma}) = \theta(\gamma)^n t^{\gamma + n\delta}$ and*

$$\sigma_{\delta, \theta}(t^{\gamma}) := \text{Exp}(D_{\delta, \theta})(t^{\gamma}) = t^{\gamma} \exp(\theta(\gamma) t^{\delta}). \quad (4.2)$$

The strong extension $\sigma_{\delta, \theta}$ lies in $\mathcal{U}_{\mathcal{R}_o}(\Gamma)$ for every $\mathfrak{o}$, with inverse $\sigma_{\delta, -\theta}$, and $\sigma_{\delta, \theta} \sigma_{\delta, \psi} = \sigma_{\delta, \theta + \psi}$ when θ, ψ both vanish on C_{δ} . With $\Phi_s = \sigma_{\delta, s\theta}$ ($s \in k$), $s \mapsto \Phi_s$ is a homomorphism, injective if $\theta \neq 0$. Conversely $\text{Exp}(D_{\delta, \theta})$ preserves $\mathcal{R}_o$ only if $\theta(C_{\delta}) = 0$.

Proof. $\theta(\delta) = 0$ gives $\theta(\gamma + n\delta) = \theta(\gamma)$ and the iterate formula. For a series supported on S , all terms of the image lie in $S + \mathbb{N}\delta$; lemma 2.1 gives well-ordering and finite contributions across all n , for single inputs and for summable families. On monomials the prescription is multiplicative by additivity of θ and the exponential law; strongness extends this to all series. The map fixes t^{δ} and every coefficient of $\exp(\cdot)$, so applying it termwise gives $\sigma_{\delta, \theta} \sigma_{\delta, \psi}(t^{\gamma}) = t^{\gamma} \exp((\theta + \psi)(\gamma) t^{\delta})$, which gives the group law and the inverse. Every correction has larger exponent, so leading terms are fixed. Here $h_{\sigma}(\gamma) = \theta(\gamma) t^{\delta}$, whose only coefficient $a_{\sigma, \delta} = \theta$ annihilates C_{δ} ; apply theorem 3.3. The coefficient at $\gamma + \delta$ of $\Phi_s(t^{\gamma})$ is $s\theta(\gamma)$, which gives injectivity. The converse is the same criterion, since $a_{\sigma, \delta} = \theta$ is the only functional. $\square$

The summability above is proved directly; no appeal to cited theorem 1.1 is made. For $\gamma < 0$ with $\theta(\gamma) \neq 0$, every $\gamma + n\delta$ stays negative, because $-\gamma > n\delta$ for all ordinary n : the whole exponential, not only a Taylor polynomial, stays on the correct side of the boundary [09].

Theorem 4.2 (Exact displacement and fixed field [04, 08, 09, 10]). *Let $\theta(C_{\delta}) = 0$, $s \neq 0$, and $f = \sum c_{\gamma} t^{\gamma}$. If $T_{\theta}(f) = \{\gamma \in \text{supp } f : \theta(\gamma) \neq 0\}$ is empty, $\Phi_s(f) = f$. Otherwise let $\gamma_0 = \min T_{\theta}(f)$; then*

$$v(\Phi_s(f) - f) = \gamma_0 + \delta, \quad [t^{\gamma_0 + \delta}](\Phi_s(f) - f) = s \theta(\gamma_0) c_{\gamma_0}.$$

In particular $\text{Fix}(\Phi_s) = k((t^{\ker \theta}))$.

Proof. Only exponents with $\theta(\gamma) \neq 0$ and powers $n \geq 1$ contribute to the difference, at $\gamma + n\delta \geq \gamma_0 + \delta$, with equality only for $\gamma = \gamma_0$, $n = 1$. $\ker \theta$ is a divisible subgroup, so the Hahn field on it is a subfield. $\square$

This first-displacement computation appears three times in the sources: as 04's exact displacement, as 08's leading term of $T(x) - x$, and inside 09's fixed-field proof. It is printed once.

Example 4.3 (A nonmonomial image [04, 09]). Let $\Gamma = \mathbf{No}$, $\delta = 1$, $\theta = q_1$, the coefficient of ω in an exponent. Every member of C_1 is finite, so $q_1(C_1) = 0$. The flow fixes $\mathbb{R}$ and $\omega = t^{-1}$, and

$$\Phi_s(\omega^\omega) = \omega^\omega \exp(-s\omega^{-1}) = \sum_{n \geq 0} \frac{(-s)^n}{n!} \omega^{\omega-n}.$$

For $s \neq 0$ this is not a monomial; every exponent $\omega - n$ is positive, so it is an omnific integer. The sum is a Hahn sum with strictly decreasing exponents, not a limit in the fine topology.

4.2 Families of shifts and a proved correspondence

Proposition 4.4 (Uniformly supported shifts [09]). *Let $S \subseteq \Gamma^{>0}$ be a set-sized well-ordered set and $(\theta_s)_{s \in S}$ additive maps $\Gamma \rightarrow k$. Then $D(f) = \sum_\gamma \sum_{s \in S} a_\gamma \theta_s(\gamma) t^{\gamma+s}$ is a contracting strong derivation, the terms of $\text{Exp}(D)(f)$ are supported in $\text{supp } f + \langle S \rangle_{\mathbb{N}}$ and form a summable family, and if $\theta_s(C_s) = 0$ for all s then $\text{Exp}(D) \in \mathcal{U}_{\mathcal{R}_o}(\Gamma)$.*

Proof. The support principle applies to $\text{supp } f + S$ and to all finite words in S ; every coefficient of D , its iterates and products is a finite sum. Additivity of each θ_s gives the product rule; replacing one input support by the union of the supports of a summable input family gives strongness; each output exponent exceeds its input exponent. Preservation of $\mathcal{R}_o$ follows from the no-crossing argument of [theorem 3.4](#) applied at each step of a word in S , which shows that $\text{Exp}(\pm D)$ preserve Π and fix k . $\square$

All explicit constructions on $\mathbf{No}$ below use one set-sized well-ordered set of shifts, so they do not depend on the proper-class case of [cited theorem 1.1](#) [09]. Source 04 proves the correspondence itself on a class of this kind.

Definition 4.5 (Common positive shifts [04]). A strong k -linear operator L has *common positive shifts* if there is a well-ordered set $S \subseteq \Gamma^{>0}$ with $\text{supp } L(t^g) \subseteq g + S$ for every g . Write S^* for the additive monoid generated by S .

Lemma 4.6 (Iterated operator summability [04]). *If L has common positive shifts, then $(L^n f)_{n \geq 0}$ is Hahn summable for every f , formal power series in L define strong operators, and coefficientwise formal identities between such series hold under composition.*

Proof. $\text{supp } L^n f \subseteq \text{supp } f + nS$. Translate $\text{supp } f$ by its minimum and apply the support principle to the positive well-ordered set $S \cup ((\text{supp } f - \min \text{supp } f) \setminus \{0\})$; a base exponent is used once and later shifts come from S , so every exponent receives finitely many contributions across all paths and all n . For a summable input family use the union of supports; regrouping is justified because every coefficient has finitely many contributions. $\square$

Theorem 4.7 (Derivations with common shifts [04]). *Let $S \subseteq \Gamma^{>0}$ be well ordered and $a_\delta : \Gamma \rightarrow k$ additive for $\delta \in S$. Then $D = \sum_{\delta \in S} D_{\delta, a_\delta}$ is a strong k -linear derivation with common positive shifts, and every such derivation has a unique expression of this form. When $\bigcap_n n\mathbf{no} = 0$, D maps $\mathcal{R}_o$ into itself if and only if $a_\delta(C_\delta) = 0$ for all $\delta \in S$.*

Proof. Summability and the Leibniz rule are termwise. Conversely put $a_\delta(g) = [t^{g+\delta}]D(t^g)$; the Leibniz rule for t^{g+h} gives additivity, and strong linearity the expression. If $D(\mathcal{R}_o) \subseteq \mathcal{R}_o$ and $x \in \Pi$, then $D(x) = nD(x/n) \in n\mathcal{R}_o$ for all n , so $D(x) \in \Pi$ by [lemma 2.3](#). For $0 < u \leq \delta$ the coefficient of $D(t^{-u})$ at the nonnegative exponent $\delta - u$ is $a_\delta(-u)$, with no cancellation between distinct shifts, so it vanishes; additivity and divisibility extend this to C_δ . The converse is the no-crossing argument. $\square$

In this theorem D is a derivation; the sources' coefficient ring D is $\mathfrak{o}$ here, which removes the collision 04 flagged.

Theorem 4.8 (Exponential and logarithm on common shifts [04]). *On operators with common positive shifts, $D \mapsto \text{Exp}(D)$ and $\sigma \mapsto \text{Log}(\sigma) = \sum_{n \geq 1} (-1)^{n+1} (\sigma - \text{id})^n / n$ are inverse bijections between strong k -linear derivations and strong k -linear field automorphisms whose difference from the identity has common positive shifts; the shift set may enlarge to $S^* \setminus \{0\}$. They restrict to a bijection between derivations preserving $\mathcal{R}_{\mathfrak{o}}$ and automorphisms preserving $\mathcal{R}_{\mathfrak{o}}$ (for $\bigcap_n n\mathfrak{o} = 0$).*

Proof. All series are justified by lemma 4.6. For a derivation D , the iterated Leibniz rule gives multiplicativity of $\text{Exp } D$ after regrouping, and $\text{Exp } D \text{Exp}(-D) = \text{id}$ gives bijectivity; leading terms are fixed since all other shifts are positive. Conversely let $L = \sigma - \text{id}$ and $\sigma^{[X]} = \sum_{n \geq 0} \binom{X}{n} L^n$. For each ordinary $m \geq 0$ this is the binomial expansion of σ^m , so for every exponent γ the t^γ coefficient of $\sigma^{[X]}(fg) - \sigma^{[X]}(f)\sigma^{[X]}(g)$, a polynomial in X by lemma 4.6, vanishes at all $m \geq 0$ and is zero. Differentiating at $X = 0$, with $\frac{d}{dX} \binom{X}{n} \big|_{X=0} = (-1)^{n-1}/n$, shows that $\text{Log } \sigma$ is a derivation; one-variable formal identities give the inverse correspondence. If D preserves $\mathcal{R}_{\mathfrak{o}}$ it maps Π into Π , so $\text{Exp}(\pm D)$ do; if $\sigma(\Pi) = \Pi$, then $(\sigma - \text{id})(\Pi) \subseteq \Pi$ and the strong logarithmic sum has negative support. $\square$

4.3 Single-shift flows without divisibility

Source 08 treats one flow at a time but allows any ordered abelian group and does not require $\theta(\delta) = 0$. Its statements are translated from the ω -convention (section 1.5). For an ordered abelian group G , $(k, \mathfrak{o}) = (\mathbb{R}, \mathbb{Z})$ or $(\mathbb{C}, \mathbb{Z}[i])$, let Π_G be the series supported in $G^{<0}$ and $\mathcal{R}_G = \mathfrak{o} \oplus \Pi_G$.

Proposition 4.9 (Falling-factorial iterates [08]). *For any additive $\theta : G \rightarrow k$ and $\delta > 0$, $D_{\delta, \theta}^n(t^\gamma) = \prod_{j=0}^{n-1} (\theta(\gamma) + j\theta(\delta)) t^{\gamma+n\delta}$, and $T^s = \sum_{n \geq 0} s^n D_{\delta, \theta}^n / n!$ ($s \in k$) is a strong k -linear 1-automorphism with $T^s T^u = T^{s+u}$. If $\theta(\delta) = 0$, $T^s(t^\gamma) = t^\gamma \exp(s\theta(\gamma)t^\delta)$; if $c = \theta(\delta) \neq 0$, $T^s(t^\gamma) = t^\gamma (1 + sct^\delta)^{\theta(\gamma)/c}$ as a formal binomial series. In either case $T^s(t^{-\delta}) = t^{-\delta} - s\theta(\delta)$.*

Proof. The iterate formula is induction. Summability, strongness and the group law follow from lemma 2.1 as in theorem 4.1, and the pairs (n, m) with $n + m = r$ are finite in number; multiplicativity follows from the iterated Leibniz rule. The two monomial formulas are the substitutions of the iterate. For $\gamma = -\delta$ every product of length at least two has the factor $\theta(-\delta) + \theta(\delta) = 0$. $\square$

This is `saut:thm:shiftflow` once more, over an arbitrary ordered abelian G ; 08 reproves the summability for that reason.

Theorem 4.10 (Threshold for single flows [08]). *Let G be any ordered abelian group, $\delta > 0$, $\theta : G \rightarrow k$ additive and $s \in k \setminus \{0\}$. The following are equivalent: (a) $T^s(\mathcal{R}_G) \subseteq \mathcal{R}_G$; (b) $T^s(\mathcal{R}_G) = \mathcal{R}_G$; (c) $\theta(C_\delta) = 0$. When they hold, T^s fixes every series supported in C_δ and preserves the constant coefficient of every series.*

Proof. Assume (a). By lemma 2.3, $\Pi_G = \bigcap_n n\mathcal{R}_G$, so $T^s(\Pi_G) \subseteq \Pi_G$. The constant term of $T^s(t^{-\delta})$ is $-s\theta(\delta)$, so $\theta(\delta) = 0$. If $0 > \gamma \in C_\delta$ and $\theta(\gamma) \neq 0$, choose n with $\gamma + n\delta > 0$; the coefficient of $T^s(t^\gamma)$ there is $(s\theta(\gamma))^n / n! \neq 0$, and the exponents $\gamma + m\delta$ are distinct, contradicting $T^s(t^\gamma) \in \Pi_G$. So θ vanishes on the negative part of C_δ , hence on C_δ . Conversely, under (c), monomials with exponents in C_δ are fixed; if $\gamma < 0$ lies outside C_δ , then $\gamma < -n\delta$ for all n , so all $\gamma + n\delta$ stay negative; positive exponents outside C_δ stay positive. So no term enters or leaves exponent 0, $T^{\pm s}(\mathcal{R}_G) \subseteq \mathcal{R}_G$, and (b) follows. (b) $\Rightarrow$ (a) is trivial. $\square$

Only divisibility of *coefficients* by ordinary integers is used, so G need not be divisible. The theorem covers the elementary family only: even at rank one, exponent rescalings and characters preserve the integer part, and it does not assert that every leading-term-preserving automorphism is the exponential of one fixed-shift derivation [08].

Remark 4.11 (Divisibility and the ring $\mathcal{A}_k$ [09]; corrected). Source 09 remarks that divisibility of Γ is essential, because with $\Gamma = \mathbb{Z}$ the translation below preserves $\mathcal{A}_k(\mathbb{Z}) = k[t^{-1}]$. That is right for the boundary ring $\mathcal{A}_k$ and wrong if read for the integer-part ring: the translation $t^{-1} \mapsto t^{-1} - 1$ sends $\frac{1}{2}t^{-1} \in \mathcal{R}_{\mathbb{Z}}(\mathbb{Z}) = \mathbb{Z} + t^{-1}\mathbb{R}[t^{-1}]$ to $\frac{1}{2}t^{-1} - \frac{1}{2} \notin \mathcal{R}_{\mathbb{Z}}(\mathbb{Z})$, as [theorem 4.10](#) predicts. So for $\Gamma = \mathbb{Z}$ the stabilizers of $\mathcal{A}_k$ and of $\mathcal{R}_{\mathbb{Z}}$ differ, and the independence of $\mathfrak{o}$ in [theorem 3.3](#) genuinely needs divisibility, whereas the $\mathcal{R}_{\mathbb{Z}}$ -form of the criterion holds without divisibility for single flows. Whether the general criterion for $\mathcal{R}_{\mathbb{Z}}$ holds for every non-divisible Γ is left open ([question 11.6](#)).

4.4 Why translations fail

Example 4.12 (Translations [04, 08, 09]; the saut example). Let $\Gamma = \mathbb{Q}$, $\delta = 1$, $\theta(\gamma) = \gamma$. Then $D^n(t^\gamma) = \gamma(\gamma+1) \cdots (\gamma+n-1)t^{\gamma+n}$ and $\text{Exp}(D)(t^\gamma) = t^\gamma(1-t)^{-\gamma}$; on t this is $t \mapsto t/(1-t)$, and on t^{-1} it is $t^{-1} \mapsto t^{-1} - 1$, which preserves $k[t^{-1}]$. But

$$\text{Exp}(D)(t^{-1/2}) = t^{-1/2}(1-t)^{1/2} = t^{-1/2} - \frac{1}{2}t^{1/2} - \frac{1}{8}t^{3/2} - \dots$$

has positive support, so neither $\mathcal{A}_k$ nor $\mathcal{R}_{\mathfrak{o}}$ is preserved: $\theta(C_1) \neq 0$. The inverse $t \mapsto t/(1+t)$ is 04's example and $X \mapsto X + s$, $X = t^{-1}$, is 08's. This is exactly the transparent example $T_1(t) = t/(1-t)$ of the surcomplex automorphism report (its $\lambda = \ell$, the constant-coefficient functional, with $\delta = 1$, after `saut:thm:shiftflow`); on the surreals $\ell(1) = 1 \neq 0$, so T_1 does *not* preserve **Oz**: it sends $\omega^{1/2}$ to $\omega^{1/2} - \frac{1}{2}\omega^{-1/2} - \dots \notin \mathbf{Oz}$. Preservation of a few familiar infinite integers is not enough; all rational monomial powers must be tested.

The Taylor coefficient flows of that report (`saut:thm:coeffflow`) move real constants and never preserve **Oz**: this follows from [corollary 5.2](#), and directly, for $d(b) = 1$ they send $bt^{-\delta/2}$ to $bt^{-\delta/2} + t^{\delta/2}$. Composed with a proper exponent embedding whose positive values dominate every multiple of δ , they do preserve and reflect **Oz**, as embeddings ([theorems 26.6](#) and [27.3](#)).

4.5 The rank dichotomy and the bottom subgroup

Theorem 4.13 (Rank dichotomy [04, 09, 10]; [08] for single flows). *For a nonzero divisible ordered abelian group Γ , any characteristic-zero k and unital $\mathfrak{o} \subseteq k$: $\mathcal{U}_{\mathcal{R}_{\mathfrak{o}}}(\Gamma) = \{\text{id}\}$ if and only if Γ is Archimedean. For every ordered abelian G , the single flows of [theorem 4.10](#) preserving $\mathcal{R}_G$ are trivial if G is Archimedean, and some are nontrivial if G is a set with a proper C_δ .*

Proof. If Γ is Archimedean, $C_\delta = \Gamma$ for all δ , so by [theorem 3.3](#) every $h_\sigma(g)$ vanishes; σ fixes all monomials, hence everything. If some C_δ is proper, choose (by a basis of the set-sized $\mathbb{Q}$ -space Γ/C_δ , or of the rationalization of the torsion-free G/C_δ for 08's case) a nonzero $\mathbb{Q}$ -linear θ into $\mathbb{Q} \subseteq k$ vanishing on C_δ ; [theorem 4.1](#) or [theorem 4.10](#) gives a nonidentity element. For $\Gamma = \mathbf{No}$ explicit functionals q_β replace the choice ([section 7](#)). $\square$

For $(k, \mathfrak{o}) = (\mathbb{R}, \mathbb{Z})$ and a set-sized Archimedean Γ , every field automorphism of $\mathbb{R}((t^\Gamma))$ preserving $\mathcal{R}_{\mathbb{Z}}$, strong or not, is therefore a monomial map $M_{\chi, \tau}$ ([corollary 24.3](#)).

Definition 4.14 (Bottom convex subgroup [04]). $B(\Gamma) = \bigcap_{\delta > 0} C_\delta$, possibly zero.

Theorem 4.15 (Common fixed field at arbitrary set-sized rank [04]). *For a set-sized nonzero divisible Γ , $\text{Fix}(\mathcal{U}_{\mathcal{R}_{\mathfrak{o}}}(\Gamma)) = k((t^{B(\Gamma)}))$.*

Proof. If $g \in B(\Gamma)$, every $a_{\sigma,\delta}(g)$ vanishes by [theorem 3.3](#), so t^g is fixed, and strong linearity fixes the Hahn field on $B(\Gamma)$. If $g \in \text{supp } f$ lies outside $B(\Gamma)$, choose δ with $g \notin C_\delta$ and a $\mathbb{Q}$ -linear θ with $\theta(C_\delta) = 0$ and $\theta(g) = 1$; [theorem 4.2](#) shows Φ_1 moves f . $\square$

Example 4.16 ([04, 09]). Let $\Gamma = \mathbb{Q}^r$, ordered so that the *last* nonzero coordinate decides the sign (09's reverse-lexicographic order; 04 orders by the first coordinate, which is the same group with coordinates reversed). Then the coordinate vectors satisfy $0 < e_1 \ll e_2 \ll \dots \ll e_r$, $C_{e_j} = \mathbb{Q}e_1 \oplus \dots \oplus \mathbb{Q}e_j$, and $B(\Gamma) = \mathbb{Q}e_1$. For $r = 1$ the stabilizer is trivial; for $r \geq 2$ it is nontrivial and fixes exactly $k((t^{\mathbb{Q}e_1}))$. The presence of a smallest Archimedean layer, not the number of layers, controls the fixed field. For $\Gamma = \mathbf{No}$, $B(\mathbf{No}) = 0$: for $g \neq 0$ take $\delta = |g|/\omega$.

4.6 Phase shears

Theorem 4.17 (Finite-dimensional phase shears [08]). *Let H be a convex subgroup of an ordered abelian group G , let $\mathfrak{m}_H$ be the series supported in $H^{>0}$, and let L be an additive map on G vanishing on H , with values in $\mathfrak{m}_H$ and image in a finite-dimensional k -subspace. Then $T_L(t^g) = t^g \exp(L(g))$ extends strongly to a k -linear 1-automorphism of the Hahn field that fixes $k((t^H))$ pointwise, preserves $\mathcal{R}_G$ and constant coefficients, and satisfies $T_L T_M = T_{L+M}$, $T_L^{-1} = T_{-L}$, and $T_L = \text{id} \Leftrightarrow L = 0$.*

Proof. Let $u_1, \dots, u_d$ span the image of L and $U = \bigcup_j \text{supp } u_j$, a well-ordered subset of $H^{>0}$. All supports of all powers in all $\exp(L(g))$ lie in the monoid generated by U , which is well ordered with finite representations; with the input support this gives one summable bound, hence the strong extension. Additivity of L gives multiplicativity. Exponents in H have zero phase, so $k((t^H))$ is fixed; in particular each $M(g)$ and its exponential are fixed, which gives $T_L T_M = T_{L+M}$ on monomials and then strongly. Injectivity follows from injectivity of $\exp$ on $\mathfrak{m}$. If $g \notin H$, every exponent of $t^g \exp(L(g))$ lies in $g + H$, which has constant strict sign by convexity; so positive and negative supports are preserved and no term reaches exponent 0. $\square$

The finite-dimensionality is a sufficient uniform support condition and must not be dropped silently: independent phases for infinitely many exponents need not form a summable family [08]. Taking $H = C_\delta$ and $L(g) = s\theta(g)t^\delta$ recovers the flows of [theorem 4.10](#).

4.7 Twists with a general profile [10]

Source 10 builds its explicit automorphisms directly, without derivations. Fix $\delta > 0$, a formal variable ξ , and an additive map $\Theta : \Gamma \rightarrow \xi k[[\xi]]$ into the power series without constant term. Define

$$\sigma_{\delta,\Theta} \left(\sum_{\gamma} a_{\gamma} t^{\gamma} \right) = \sum_{\gamma} a_{\gamma} t^{\gamma} \exp(\Theta(\gamma)(t^{\delta})), \quad (4.3)$$

where $\Theta(\gamma)(t^{\delta})$ is $\Theta(\gamma)$ evaluated at $\xi = t^{\delta}$. For $\Theta(\gamma) = \theta(\gamma)\xi$ this is the shear $\sigma_{\delta,\theta}$ of (4.2). Writing $\Theta(\gamma) = \sum_{n \geq 1} \theta_n(\gamma)\xi^n$ with additive θ_n , the map is Exp of $\sum_n D_{n\delta,\theta_n}$, a derivation with the one well-ordered shift set $\mathbb{N}_{>0}\delta$ as in [proposition 4.4](#) (an identification by the merge); 10's proof below is direct.

Theorem 4.18 (Twists [10]). *If $\Theta(\delta) = 0$, then $\sigma_{\delta,\Theta}$ is a strong k -linear 1-automorphism of K with strong inverse $\sigma_{\delta,-\Theta}$, and $\sigma_{\delta,\Theta}\sigma_{\delta,\Psi} = \sigma_{\delta,\Theta+\Psi}$ for every second such Ψ ; the map $\Theta \mapsto \sigma_{\delta,\Theta}$ is injective. If $\Theta(C_\delta) = 0$, then $\sigma_{\delta,\Theta} \in \mathcal{U}_{\mathcal{R}_\delta}(\Gamma)$ for every unital $\mathfrak{o} \subseteq k$; conversely [merge] $\sigma_{\delta,\Theta} \in \mathcal{U}_{\mathcal{R}_\delta}(\Gamma)$ forces $\Theta(C_\delta) = 0$. For set-sized Γ , the additive group $\text{Hom}_{\mathbb{Q}}(\Gamma/C_\delta, \xi k[[\xi]])$ thus embeds in $\mathcal{U}_{\mathcal{R}_\delta}(\Gamma)$.*

Proof. The support of $t^\gamma \exp(\Theta(\gamma)(t^\delta))$ lies in $\gamma + \mathbb{N}\delta$. For an input supported in S , [lemma 2.1](#) makes $S + \mathbb{N}\delta$ well ordered with finitely many pairs (γ, n) at each exponent, which defines (4.3); for a summable family of inputs use the union of their supports. So the map is strong. Additivity of Θ gives multiplicativity on monomials, and strongness extends it to products of series, which are summable sums of monomial products; k is fixed. Since $\Theta(\delta) = 0$, the map fixes t^δ and hence every series in t^δ with coefficients in k , so $\sigma_{\delta, \Theta\sigma_\delta, \Psi}(t^\gamma) = t^\gamma \exp(\Theta(\gamma)(t^\delta)) \exp(\Psi(\gamma)(t^\delta)) = \sigma_{\delta, \Theta+\Psi}(t^\gamma)$, and strongness extends the identity; $\Psi = -\Theta$ gives the inverse. Every correction lies at a larger exponent, and no term arising from a larger input exponent reaches the least one, so leading terms are fixed. Evaluation at $\xi = t^\delta$ and $\exp$ are injective, which gives injectivity. If $\Theta(C_\delta) = 0$ and $\Theta(\gamma) \neq 0$, then $\gamma \notin C_\delta$, so every shift $n\delta$ satisfies $n\delta \ll |\gamma|$; thus $\text{supp } h_\sigma(\gamma) \subseteq H_\gamma$ for every $\gamma \neq 0$, and [theorem 3.3](#) applies. Conversely, if $0 \neq \gamma \in C_\delta$, say $|\gamma| \leq N\delta$, no shift $n\delta$ lies in H_γ , so condition (v) there forces $h_\sigma(\gamma) = \Theta(\gamma)(t^\delta) = 0$. $\square$

The field-automorphism part needs only $\Theta(\delta) = 0$; the stronger $\Theta(C_\delta) = 0$ is exactly the integer-part condition [10]. (For a single linear profile, [proposition 4.9](#) even allows $\theta(\delta) \neq 0$.) For additive $\theta : \Gamma \rightarrow k$ with $\theta(\delta) = 0$ and a profile $F \in 1 + \xi k[[\xi]]$, the choice $\Theta = \theta \log F$ gives the *binomial twist*

$$\sigma_{\delta, \theta}^F \left(\sum_{\gamma} a_{\gamma} t^{\gamma} \right) = \sum_{\gamma} a_{\gamma} t^{\gamma} F(t^{\delta})^{\theta(\gamma)}, \quad F^a = \exp(a \log F) = \sum_{n \geq 0} \binom{a}{n} (F - 1)^n. \quad (4.4)$$

Its inverse is $\sigma_{\delta, -\theta}^F = \sigma_{\delta, \theta}^{1/F}$, composition adds θ for fixed F, δ , and $\sigma_{\delta, \theta}^{\exp \xi} = \sigma_{\delta, \theta}$. The most economical profile is $F = 1 + \xi$:

$$\sigma_{\delta, \theta}^{1+\xi}(t^{\gamma}) = \sum_{n \geq 0} \binom{\theta(\gamma)}{n} t^{\gamma + n\delta}, \quad (4.5)$$

whose coefficients are polynomials over $\mathbb{Q}$ in $\theta(\gamma)$, with no choice of branch even for $k = \mathbb{C}$ [10]. Source 10's $\sigma_{\lambda, F, h}$ is $\sigma_{h, -\lambda}^F$ ([section 1.5](#)).

Example 4.19 (Two scales [10]). Let $\Gamma = \mathbb{Q}^2$ as in [example 4.16](#), so $e_1 \ll e_2$, and let θ be minus the e_2 -coordinate, which vanishes on $C_{e_1} = \mathbb{Q}e_1$. Then $\sigma = \sigma_{e_1, \theta}^{1+\xi}$ sends $t^{xe_1 + ye_2}$ to $t^{xe_1 + ye_2} (1 + t^{e_1})^{-y}$; for instance $\sigma(t^{-e_2}) = t^{-e_2} + t^{-e_2 + e_1}$ and

$$\sigma(t^{-e_2/2}) = t^{-e_2/2} + \frac{1}{2}t^{-e_2/2 + e_1} - \frac{1}{8}t^{-e_2/2 + 2e_1} + \frac{1}{16}t^{-e_2/2 + 3e_1} - \dots$$

Every exponent $-e_2/2 + ne_1$ is negative, however large n is. With $\Gamma = \mathbb{Q}$ the prescription $t^{-1} \mapsto t^{-1}(1 + t)$ would force a positive exponent at $t^{-1/2}$ (compare [example 4.12](#)).

Theorem 4.20 (Fixed field and first displacement [10]). *Let $\theta(C_\delta) = 0$, let $F = 1 + c_m \xi^m + c_{m+1} \xi^{m+1} + \dots$ with $m \geq 1$ and $c_m \neq 0$, and put $\sigma = \sigma_{\delta, \theta}^F$. Then $\text{Fix}(\sigma) = k((t^{\ker \theta}))$. More precisely, if $f = \sum a_{\gamma} t^{\gamma}$ is not supported in $\ker \theta$ and $\gamma_* = \min\{\gamma \in \text{supp } f : \theta(\gamma) \neq 0\}$, then*

$$v(\sigma(f) - f) = \gamma_* + m\delta, \quad [t^{\gamma_* + m\delta}](\sigma(f) - f) = a_{\gamma_*} \theta(\gamma_*) c_m.$$

For every integer $n \neq 0$, $\text{Fix}(\sigma^n) = \text{Fix}(\sigma)$.

Proof. Monomials with $\theta(\gamma) = 0$ are fixed, hence, by strongness, all of $k((t^{\ker \theta}))$. For $a \neq 0$ the first nonconstant term of F^a is $a c_m \xi^m$, so for $\theta(\gamma) \neq 0$ the difference $\sigma(t^{\gamma}) - t^{\gamma}$ has least term $\theta(\gamma) c_m t^{\gamma + m\delta}$ and all other terms at exponents at least $\gamma + (m+1)\delta$. Exponents $\gamma < \gamma_*$ of f contribute no difference, and exponents $\gamma > \gamma_*$ contribute only beyond $\gamma + m\delta > \gamma_* + m\delta$. So nothing cancels the term at $\gamma_* + m\delta$. The composition law gives $\sigma^n = \sigma_{\delta, n\theta}^F$, and $\ker(n\theta) = \ker \theta$ in characteristic zero. $\square$

For $F = \exp \xi$ this is [theorem 4.2](#); the case of a general profile and the statement for powers are 10's.

Corollary 4.21 (No finite motion [10]). *Every finite orbit of σ in [theorem 4.20](#) is a fixed point, and $\text{Fix}(\sigma)$ is relatively algebraically closed in K .*

Proof. A finite orbit gives $\sigma^n(f) = f$ for some $n \geq 1$, hence $\sigma(f) = f$. If f is algebraic over $\text{Fix}(\sigma)$, all $\sigma^n(f)$ are roots of its minimal polynomial, whose coefficients are fixed; so the orbit is finite. $\square$

This is relative algebraic closedness only: with real coefficients the fixed field need not contain a square root of -1 [10].

Proposition 4.22 (Independent exponent cosets [10]). *Let $V = \ker \theta$ and let $(\gamma_j)_{j \in J}$ have $\mathbb{Q}$ -linearly independent images in Γ/V . Then the monomials t^{γ_j} are algebraically independent over $k((t^V))$.*

Proof. A relation involves finitely many indices and has the form $\sum_{\nu \in A} b_\nu t^{\nu_1 \gamma_1 + \dots + \nu_r \gamma_r} = 0$ with $b_\nu \in k((t^V))$, distinct ν and some $b_\nu \neq 0$. For $\nu \neq \mu$, $\sum_j (\nu_j - \mu_j) \gamma_j \notin V$, so the nonzero summands have supports in distinct cosets of V and cannot cancel. $\square$

Remark 4.23 (Rank, divisibility and what is classified [10]). Source 10 proves [theorem 4.13](#) with a binomial witness: if $\delta \ll |\gamma_0|$ with $\gamma_0 < 0$ and θ is $\mathbb{Q}$ -linear with $\theta(C_\delta) = 0$ and $\theta(\gamma_0) = 1$, then $\sigma_{\delta, \theta}^{1+\xi}(t^{\gamma_0}) = t^{\gamma_0} + t^{\gamma_0+\delta}$, and the map fixes $k((t^{C_\delta}))$ pointwise. Its divisibility example is stated for $\mathfrak{o} = k$: $t^{-1} \mapsto t^{-1} + 1$ preserves $\mathcal{A}_k(\mathbb{Z}) = k[t^{-1}]$ for $\Gamma = \mathbb{Z}$, and 10 says explicitly that it does not preserve $\mathbb{Z} + t^{-1}\mathbb{R}[t^{-1}]$. This agrees with [remark 4.11](#) as corrected there; since 10 assumes divisibility throughout, it does not bear on [question 11.6](#). At Archimedean rank the monomial maps $t^\gamma \mapsto t^{2\gamma}$ and the character rescalings still preserve $\mathcal{R}_\mathfrak{o}$; they lie outside the leading-term-fixing factor and are no counterexample to [theorem 4.13](#).

5 The strong stabilizer and all pair automorphisms

Call σ *valued* if $v(\sigma f) = \tau(v(f))$ for an order automorphism τ of Γ . For a character $\chi : (\Gamma, +) \rightarrow (k^\times, \cdot)$ put

$$M_{\chi, \tau} \left(\sum a_\gamma t^\gamma \right) = \sum a_\gamma \chi(\gamma) t^{\tau(\gamma)}. \quad (5.1)$$

It is a strong field automorphism preserving $\mathcal{R}_\mathfrak{o}$, since τ preserves signs of exponents and $\chi(0) = 1$; $M_{\chi, \tau} M_{\chi', \tau'} = M_{\chi' \cdot (\chi \circ \tau'), \tau \tau'}$ [04, 09].

Theorem 5.1 (Strong valued factorization [04, 09, 10]). *Every strong valued k -fixing automorphism σ with $\sigma(\mathcal{R}_\mathfrak{o}) = \mathcal{R}_\mathfrak{o}$ factors uniquely as $\sigma = M_{\chi, \tau} \circ u$ with $u \in \mathcal{U}_{\mathcal{R}_\mathfrak{o}}(\Gamma)$, where τ is the induced exponent automorphism and $\chi(\gamma) = \text{lc}(\sigma(t^\gamma))$. The monomial maps form a subgroup normalizing $\mathcal{U}_{\mathcal{R}_\mathfrak{o}}(\Gamma)$, and*

$$M_{\chi, \tau} D_{\delta, \theta} M_{\chi, \tau}^{-1} = D_{\tau(\delta), \chi(\delta) \theta \circ \tau^{-1}}, \quad (5.2)$$

so the convex condition is invariant, since $\tau(C_\delta) = C_{\tau(\delta)}$.

Proof. Multiplicativity of leading coefficients gives that χ is a character. $u = M_{\chi, \tau}^{-1} \sigma$ fixes the leading term of each monomial; since the images of distinct monomial terms have strictly ordered valuations, strongness shows that u fixes the leading term of every series, and u preserves $\mathcal{R}_\mathfrak{o}$. The data τ, χ are forced by σ , hence so is u . Conjugating a 1-automorphism by a monomial map gives a 1-automorphism preserving $\mathcal{R}_\mathfrak{o}$. Formula (5.2) is checked on monomials. $\square$

The ambient decomposition is due to Kuhlmann–Serra [13], and its surcomplex form is the four-layer decomposition `saut:thm:decomp` $U \rtimes \text{Hom}(\Gamma, \mathbb{C}^\times) \rtimes (\text{Aut } \mathbb{C} \times \text{Aut } \Gamma)$ of the surcomplex automorphism report; [theorem 5.1](#) is its restriction to the $\mathcal{R}_o$ -stabilizer with the residue layer trivial. Sources 04, 09 and 10 credit [13]; the new information is the description of u by [theorem 3.3](#). Source 10 states the factorization as $\text{Stab}(\mathcal{R}_o) = \mathcal{U}_{\mathcal{R}_o}(\Gamma) \rtimes \{M_{\chi, \tau}\}$, restricting to positive characters when k is ordered and only order-preserving maps are considered; it is not a new decomposition theorem for $\text{Aut}(\mathbf{No})$, and coefficient-moving or nonstrong automorphisms lie outside it [10]. For the real pair neither kind exists: coefficients are fixed by [corollary 5.2](#) and strongness is automatic by [theorem 23.2](#), so the splitting covers every automorphism ([theorem 24.1](#)). For $k = \mathbb{R}$ and divisible Γ every character is positive, $\chi(\gamma) = \chi(\gamma/2)^2$; for $\Gamma = \mathbb{Q}$ the factor u is absent and every such real automorphism is $t^q \mapsto c^q t^{\lambda q}$ with $c > 0$, $\lambda \in \mathbb{Q}_{>0}$, so the higher-rank shears are a new phenomenon, not complicated one-variable substitutions [09]. The monomial maps include the dilations S_a of the Diophantine report (`odg:def:ex:dilation`) and the character twists ([remark 9.5](#)).

Corollary 5.2 (All pair automorphisms [08, 09, 10]). *Every field automorphism of $\mathbf{No}$ preserving $\mathbf{Oz}$ setwise fixes $\mathbb{R}$ pointwise, preserves order and the natural valuation ring, and commutes with P_- , P_0 , P_+ and with the floor map of $\mathbf{Oz}$; this needs no strongness. Consequently, among strong automorphisms, the $\mathbf{Oz}$ -stabilizer is exactly $\{M_{\chi, \tau}\} \rtimes \mathcal{U}_{\mathbf{Oz}}$, with χ positive; by [theorem 23.2](#) every automorphism preserving $\mathbf{Oz}$ is strong, so this is the whole stabilizer ([theorem 24.1](#)).*

Proof. By `odg:def:thm:autreal` of the Diophantine report, every automorphism of $\mathbf{Oz}$ extends uniquely to $\mathbf{No}$, preserves order, $\mathbb{R}$ and the finite ring, and fixes $\mathbb{R}$ pointwise; restriction and extension are inverse correspondences [08, 09]. The sets and graphs of [theorem 8.2](#) are parameter-free definable, so they are preserved. Fixing $\mathbb{R}$ pointwise makes a strong such map k -fixing, and preserving the valuation ring makes it valued, so [theorem 5.1](#) applies. The last clause is [theorem 23.2](#), proved in [part III](#) from the first sentence of this corollary and [theorem 22.1](#), with no circularity. $\square$

6 Convex rank, solvability, and a wreath product

6.1 The bracket law

Lemma 6.1 (Homogeneous brackets [04, 08, 09]). *For additive $\theta, \psi : \Gamma \rightarrow k$ and $\delta, \epsilon > 0$,*

$$[D_{\delta, \theta}, D_{\epsilon, \psi}] = D_{\delta + \epsilon, \theta(\epsilon)\psi - \psi(\delta)\theta}. \quad (6.1)$$

If $\theta(C_\delta) = 0$ and $\psi(C_\epsilon) = 0$, the bracket is zero when $C_\delta = C_\epsilon$; if $C_\delta \subsetneq C_\epsilon$ it is a multiple of $D_{\delta + \epsilon, \psi}$, whose shift lies in the C_ϵ scale and whose functional vanishes on $C_{\delta + \epsilon} = C_\epsilon$.

Proof. On t^γ the two compositions differ at $\gamma + \delta + \epsilon$ by $\psi(\gamma)\theta(\gamma + \epsilon) - \theta(\gamma)\psi(\gamma + \delta) = \theta(\epsilon)\psi(\gamma) - \psi(\delta)\theta(\gamma)$; strongness extends the identity. Equal scales: each functional kills the other's shift. If $C_\delta \subsetneq C_\epsilon$, then $\psi(\delta) = 0$. $\square$

Source 08 records the same identity in its own normalization (its bracket $[D_{\varphi, \delta}, D_{\psi, \epsilon}] = D_{\psi(\delta)\varphi - \varphi(\epsilon)\psi, \delta + \epsilon}$, which is (6.1) after the sign change of [section 1.5](#)). Elementary derivations with shifts in the same Archimedean class commute; noncommutativity is an interaction between layers [04]. For general strong derivations the computation applies after expansion into shift coefficients: the two compositions are strongly summable and every output coefficient is a finite sum [09].

6.2 A filtration by convex scales

Say that Γ has *finite ordered rank* r if its distinct nonzero principal convex subgroups are exactly $0 < C_1 < \cdots < C_r = \Gamma$. The layer of a shift δ is the j with $C_\delta = C_j$. A shift in layer r contributes nothing to $\mathfrak{L}_A$ (corollary 3.5), since its functional vanishes on Γ . Let $\mathfrak{F}_j \subseteq \mathfrak{L}_A$ consist of the derivations whose nonzero shifts have layers at least j , so $\mathfrak{L}_A = \mathfrak{F}_1 \supseteq \cdots \supseteq \mathfrak{F}_r = 0$. Selecting the terms of prescribed layers from the images of monomials keeps a summable subfamily, and the new coefficient functions stay additive, so the $\mathfrak{F}_j$ are strongly closed Lie ideals [09].

Lemma 6.2 ([09]). $[\mathfrak{F}_j, \mathfrak{F}_j] \subseteq \mathfrak{F}_{j+1}$ and $[\mathfrak{L}_A, \mathfrak{F}_j] \subseteq \mathfrak{F}_j$.

Proof. Apply lemma 6.1 to pairs of shifts: equal layers cancel and unequal layers give a term in the larger layer. $\square$

Theorem 6.3 (Finite-rank bound [09]). *If the set-sized group Γ has finite ordered rank r , then $\mathfrak{L}_A$ and $\mathcal{U}_{\mathcal{R}_o}(\Gamma)$ have derived length at most $r - 1$.*

Proof. Inductively $\mathfrak{L}_A^{(n)} \subseteq \mathfrak{F}_{n+1}$. For the group, put $G_j = \text{Exp}(\mathfrak{F}_j)$; by the Baker–Campbell–Hausdorff part of cited theorem 1.1 each G_j is a subgroup, and the logarithm of a commutator of two elements of G_j is a strongly summable Lie series each of whose terms contains a bracket of two elements of $\mathfrak{F}_j$, hence lies in $\mathfrak{F}_{j+1}$. So $[G_j, G_j] \subseteq G_{j+1}$, and $G_r = \{\text{id}\}$. $\square$

The Lie bound is elementary; the group bound uses the imported correspondence, which [2] proves for set-sized Γ .

6.3 Sharpness, including the abstract group

In $\mathbb{Q}^r$ ordered as in example 4.16, let θ_i extract the i th coordinate, $X_{j,i} = D_{e_j, \theta_i}$ for $j < i$, and

$$a_0 = 0, \quad a_k = \sum_{h=1}^k 2^{k-h} e_h, \quad Y_{k;j,i} = D_{a_k + e_j, \theta_i} \quad (k < j < i \leq r).$$

These satisfy the criterion, since $C_{e_j} = \mathbb{Q}e_1 \oplus \cdots \oplus \mathbb{Q}e_j$.

Lemma 6.4 (Balanced brackets [09]). $Y_{k;j,i} \in \mathfrak{L}_A^{(k)}$, and $[Y_{k-1;k,j}, Y_{k-1;j,i}] = Y_{k;j,i}$ for $k \geq 1$.

Proof. At $k = 0$ this is the definition. In the bracket, $\theta_j(a_{k-1} + e_j) = 1$ and $\theta_i(a_{k-1} + e_k) = 0$, and the shift is $2a_{k-1} + e_k + e_j = a_k + e_j$; apply (6.1) and induct. $\square$

Lemma 6.5 (Mixed-parameter detection [09]). *Take finitely many homogeneous positive-shift derivations and an iterated Lie bracket W of them. Replace each leaf occurrence D_ℓ by $\text{Exp}(s_\ell D_\ell)$ with independent scalar parameters and each bracket by the group word $ghg^{-1}h^{-1}$, giving $\mathcal{W}(\mathbf{s})$. In its action on a monomial, the coefficient of the product of all leaf parameters is the action of W . If $W \neq 0$, some specialization $\mathbf{s} \in k^m$ gives a nonidentity group word.*

Proof. Formally, $(1+sD+\cdots)(1+uE+\cdots)(1-sD+\cdots)(1-uE+\cdots) = 1+su[D, E] + \text{terms of other multidegrees}$; induct on the bracket tree, using that the subtrees have disjoint parameter sets and that a group word is the identity when a required leaf parameter is zero. Choose a monomial and an output exponent on which W acts nontrivially. Only a finite set of shifts occurs, so by the support principle the coefficient at that output is a polynomial in the parameters, nonzero by the first part. An infinite field has a point where it does not vanish. Repeated leaves are handled by their separate parameters. $\square$

Theorem 6.6 (Sharp derived length [09]). *For $\Gamma = \mathbb{Q}^r$ as above, every characteristic-zero k and every unital $\mathfrak{o} \subseteq k$, the group $\mathcal{U}_{\mathfrak{o}}(\Gamma)$ and its Lie algebra $\mathfrak{L}_A$ have derived length exactly $r - 1$.*

Proof. For $r = 1$ use [theorem 4.13](#). For $r \geq 2$, $0 \neq Y_{r-2;r-1,r} \in \mathfrak{L}_A^{(r-2)}$ by [lemma 6.4](#), and [lemma 6.5](#) turns its balanced bracket into a nonidentity element of the $(r - 2)$ nd derived subgroup. The upper bound is [theorem 6.3](#). $\square$

The lower bound does not identify the abstract derived subgroup with the exponential of the derived Lie algebra; it uses independent parameters and a nonzero mixed coefficient. The upper bound uses BCH and strongly closed ideals [09].

6.4 Metabelian and not nilpotent: one group, printed once

On $\Gamma = \mathbf{No}$ put $D = D_{1,q_1}$, $E = D_{\omega,q_2}$ and $E_n = D_{\omega+n,q_2}$. Both D and E satisfy the criterion: q_1 kills the finite exponents and q_2 kills every exponent bounded by an ordinary multiple of ω .

Proposition 6.7 ([04, 09]). *$[D, E_n] = E_{n+1}$ and $[E_m, E_n] = 0$; the E_n are nonzero and linearly independent over $\mathbb{R}$. So the Lie algebra spanned by $D, E_0, E_1, \dots$ is metabelian and not nilpotent. The same holds in rank three: in $\mathbb{Q}^3$ with $X = D_{e_1,\theta_2}$, $Y = D_{e_2,\theta_3}$ one has $(\text{ad } X)^n Y = D_{e_2+n e_1,\theta_3} \neq 0$.*

Proof. $q_1(\omega+n) = 1$, $q_2(1) = q_2(\omega+n) = 0$; apply (6.1). $E_n(t^{\omega^2}) = t^{\omega^2+\omega+n}$ gives independence. The rank-three case is the same computation [09]. $\square$

Put $A_s = \text{Exp}(sD)$ and $B_u = \text{Exp}(uE)$ for $s, u \in \mathbb{R}$, and $[F, G] = FGF^{-1}G^{-1}$. Here e^{st} denotes the Hahn series $\exp(st)$ in the monomial $t = \omega^{-1}$.

Theorem 6.8 (Exact commutators [04, 08]). *$A_s B_u A_s^{-1} = \text{Exp}(u e^{st} E)$ and $[A_s, B_u] = \text{Exp}(u(e^{st} - 1)E)$. With $C_0 = B_u$ and $C_{n+1} = [A_s, C_n]$,*

$$C_n = \text{Exp}(u(e^{st} - 1)^n E), \quad C_n(t^{-\omega^2}) = t^{-\omega^2} \exp(-u(e^{st} - 1)^n t^\omega),$$

so for $s, u \neq 0$, $C_n(t^{-\omega^2}) - t^{-\omega^2}$ has leading term $-us^n t^{-\omega^2+\omega+n}$. Hence $\langle A_s, B_u \rangle$ is metabelian and not nilpotent. At $s = u = -1$, the leading term is $(-1)^n \omega^{\omega^2-\omega-n}$.

Proof. D fixes t (as $q_1(1) = 0$) and $A_s(t^\omega) = t^\omega e^{st}$. On monomials, $A_s E A_s^{-1}(t^\gamma) = q_2(\gamma) t^{\gamma+\omega} e^{st}$, because E kills every series in t and $q_1(\gamma + \omega) - q_1(\gamma) = 1$; so $A_s E A_s^{-1} = e^{st} E$, and exponentiating gives the first formula. E kills $\mathbb{R}[[t]]$, so the derivations $h(t)E$, $h \in \mathbb{R}[[t]]$, commute and their exponentials add coefficients, which gives the second formula and, by induction, C_n . All shifts lie in the monoid generated by the well-ordered set $\{1\} \cup (\omega + \mathbb{N})$, so [theorem 4.8](#) justifies all sums. The maps $\text{Exp}(h(t)E)$ form an abelian group normalized by A_s , which acts by $h \mapsto e^{st}h$; so the generated group is abelian-by-abelian. Since $q_2(-\omega^2) = -1$, $C_n(t^{-\omega^2})$ is as displayed, and $(e^{st} - 1)^n$ has leading term $s^n t^n$; the leading term of the difference is therefore nonzero at every depth, so the lower central series never terminates. $\square$

Source 08 works with $A_1^{(08)} = A_{-1}$ and $B_1^{(08)} = B_{-1}$ and proves faithfulness, which identifies the group. Write $v = t = \omega^{-1}$, $u_0 = t^\omega = \omega^{-\omega}$, $P = t^{-\omega} = \omega^\omega$, $M = t^{-\omega^2} = \omega^{\omega^2}$, and for h supported in $C_\omega^{>0}$ let $N_h(t^\gamma) = t^\gamma \exp(-q_2(\gamma)h)$, a phase shear ([theorem 4.17](#)) with $N_h N_{h'} = N_{h+h'}$, fixing v, u_0, P and with $N_h(M) = M \exp(h)$. Then $B_{-1} = N_{u_0}$, $A_{-1}(u_0) = u_0 e^{-v}$, $A_{-1}(P) = P e^v$, and $A_{-1} N_h A_{-1}^{-1} = N_{A_{-1}(h)}$.

Lemma 6.9 (Finite exponential independence [08]). *For a Laurent polynomial $p \in \mathbb{Z}[X, X^{-1}]$ put $h_p = u_0 p(e^{-v})$ and $\mathcal{N}_p = N_{h_p}$. Then $p \mapsto p(e^{-v})$ is injective, $p \mapsto \mathcal{N}_p$ is an injective homomorphism, and $A_{-1}\mathcal{N}_p A_{-1}^{-1} = \mathcal{N}_{Xp}$.*

Proof. If $\sum_{n \in F} k_n e^{-nv} = 0$ with $|F| = r$ distinct integers, comparing the coefficients of v^j , $0 \leq j < r$, gives a Vandermonde system with distinct nodes $-n$, so all $k_n = 0$. Injectivity of $\mathcal{N}$ follows from $\mathcal{N}_p(M) = M \exp(h_p)$, and $A_{-1}(h_p) = u_0 e^{-v} p(e^{-v}) = h_{Xp}$. $\square$

Theorem 6.10 (A faithful $\mathbb{Z} \wr \mathbb{Z}$ [08]). *The map $\mathbb{Z}[X, X^{-1}] \rtimes \mathbb{Z} \rightarrow \mathcal{U}_{\mathbf{Oz}}$, $(p, m) \mapsto \mathcal{N}_p A_{-1}^m$, with $(p, m)(q, n) = (p + X^m q, m + n)$, is an injective homomorphism onto $\langle A_{-1}, B_{-1} \rangle$. So this group is the restricted wreath product $\mathbb{Z} \wr \mathbb{Z}$, of derived length exactly two. It extends $\mathbb{C}$ -linearly to $\mathbf{No}(i)$ preserving $\mathbf{Oz}[i]$, and it already lives on the set-sized exponent group $\mathbb{Q}\omega^2 + \mathbb{Q}\omega + \mathbb{Q}$.*

Proof. Lemma 6.9 gives the multiplication law, and $\mathcal{N}_{X^n} = A_{-1}^n B_{-1} A_{-1}^{-n}$ shows that the image is generated by A_{-1}, B_{-1} . If $\mathcal{N}_p A_{-1}^m = \text{id}$, applying it to P gives $P e^{mv} = P$, so $m = 0$, and then to M gives $h_p = 0$, so $p = 0$. Finite Laurent polynomials under multiplication by X are $\bigoplus_{\mathbb{Z}} \mathbb{Z}$ with the shift. The group is nonabelian and its normal subgroup $\mathcal{N}$ is abelian with cyclic quotient. $\square$

Remark 6.11 (All nonzero times [merge]). The same proof gives $\langle A_s, B_u \rangle \cong \mathbb{Z} \wr \mathbb{Z}$ for all real $s, u \neq 0$: here $B_u = N_{-ut\omega}$, $A_s N_h A_s^{-1} = N_{A_s(h)}$ and $A_s(t^\omega) = t^\omega e^{st}$, so with $h_p = -ut^\omega p(e^{st})$ the map $(p, m) \mapsto N_{h_p} A_s^m$ obeys the same law; the Vandermonde nodes ns are distinct, and $A_s^m(P) = P e^{-mst}$ forces $m = 0$.

Corollary 6.12 (Every set stabilizer is not nilpotent [04, 08]). *For every set $P \subseteq \mathbf{No}$, the pointwise stabilizer of P in $\mathcal{U}_{\mathbf{Oz}}$ contains a subgroup isomorphic to $\mathbb{Z} \wr \mathbb{Z}$, metabelian and not nilpotent.*

Proof. Choose ordinals $0 < \beta < \beta'$ above every inner exponent of every outer exponent of every member of P (lemma 7.1), and replace ω, ω^2 by $\epsilon = \omega^\beta, \omega^{\beta'}$, using the functionals $q_\beta, q_{\beta'}$ and the shifts $1, \epsilon$. Both functionals vanish on the exponents of P and satisfy the criterion, and $q_\beta(\epsilon) = 1, q_\beta(1) = q_{\beta'}(\epsilon) = q_{\beta'}(1) = 0$; the proofs are unchanged. $\square$

These examples live in rank three, consistent with theorem 6.3 (derived length at most two there) [08, 09].

Example 6.13 (Binomial twists at three scales [10]). Source 10 shows noncommutativity with binomial twists at the scales of $x = \omega, y = \omega^\omega$ and $z = \omega^{\omega^2}$. Let $\sigma = \sigma_{1, -q_1}^{1+\xi}$ and $\tau = \sigma_{\omega, -q_2}^{1+\xi}$, that is $\sigma(\omega^g) = \omega^g(1 + \omega^{-1})^{q_1(g)}$ and $\tau(\omega^g) = \omega^g(1 + \omega^{-\omega})^{q_2(g)}$; both lie in $\mathcal{U}_{\mathbf{Oz}}$ by theorem 4.18, since $q_1(C_1) = 0$ and $q_2(C_\omega) = 0$. With $u = 1 + x^{-1}$,

$$\sigma(x) = x, \sigma(y) = yu, \sigma(z) = z, \quad \tau(x) = x, \tau(y) = y, \tau(z) = z(1 + y^{-1}),$$

so $\sigma\tau(z) - \tau\sigma(z) = -z/(xy(1 + x^{-1})) \neq 0$ and $[\sigma, \tau](z) = z(1 + 1/(yu))/(1 + 1/y) \neq z$, with $[\sigma, \tau] = \sigma\tau\sigma^{-1}\tau^{-1}$. The rational expressions denote the corresponding Hahn series. The example proves only that the integer-part stabilizer is not abelian and gives no presentation of it; the exponential shears at the same scales have the exact structure of theorems 6.8 and 6.10.

7 Fixed fields and arbitrarily large stabilizers

7.1 Exponent coefficient functionals

For $\beta \in \mathbf{No}$ let $q_\beta(\gamma) = [\omega^\beta]\gamma$ be the coefficient of ω^β in the Conway normal form of the exponent γ (the *inner* normal form; the outer support of $f = \sum c_\gamma t^\gamma$ consists of the exponents γ). Normal-form addition makes $q_\beta : \mathbf{No} \rightarrow \mathbb{R}$ additive; it need not commute with infinite sums inside the exponent group, and only finite additivity is used [04]. If $0 < \delta \ll \omega^\beta$ then $q_\beta(C_\delta) = 0$: a nonzero γ with $|\gamma| \leq n\delta$ has leading Conway exponent at most that of δ , which is below β [04, 08, 09].

Lemma 7.1 (Parameter avoidance [04, 08, 09, 10]). *Let $P \subseteq \mathbf{No}(i)$ be a set, E the union of the outer supports of the real and imaginary parts of its members, and T the union of the inner supports of the members of E . Then E and T are sets, and for every ordinal $\beta > 0$ above all members of T , q_β vanishes on E .*

Proof. Replacement and union; every set of surreals has a strict upper bound, for example an ordinal above all their birthdays. This bounds a set of supports; it does not choose a bound for all exponents at once. $\square$

Theorem 7.2 (The common fixed field [04, 08, 09]). *$\text{Fix}(\mathcal{U}_{\mathbf{Oz}}) = \mathbb{R}$, and already the elementary shears σ_{δ, q_β} have common fixed field $\mathbb{R}$. The same holds for all field automorphisms of $\mathbf{No}$ preserving $\mathbf{Oz}$. The common fixed elements in $\mathbf{Oz}$ are $\mathbb{Z}$, and the common fixed subring of all ring automorphisms of $\mathbf{Oz}$ is $\mathbb{Z}$.*

Proof. All these maps fix $\mathbb{R}$ (corollary 5.2). If $f \notin \mathbb{R}$, some nonzero $\gamma \in \text{supp } f$ has $q_\beta(\gamma) \neq 0$ for some β , for instance its leading inner exponent [08]. Put $\theta = q_\beta$ and $\delta = \omega^{\beta-1}$; then $\theta(C_\delta) = 0$ and $\theta(\delta) = 0$, and theorem 4.2 shows that $\sigma_{\delta, \theta}$ moves f : its first displacement is $a_\lambda \theta(\lambda) t^{\lambda+\delta}$, λ the least exponent of f with $\theta(\lambda) \neq 0$. Ring automorphisms of $\mathbf{Oz}$ are restrictions of field automorphisms preserving $\mathbf{Oz}$ (odg: def: thm: autreal). $\square$

Source 04 reaches the same conclusion through lemma 7.3 with $V = 0$; the explicit choice above is 08's and 09's.

7.2 Relative fixed fields and the Hahn hull

Lemma 7.3 (Set-sized coefficient separation [04]). *Let $V \subseteq \mathbf{No}$ be a set-sized $\mathbb{Q}$ -subspace and $g \in \mathbf{No} \setminus V$. There are $\delta > 0$ and an additive class map $a : \mathbf{No} \rightarrow \mathbb{R}$ with $a(V) = 0$, $a(C_\delta) = 0$ and $a(g) = 1$.*

Proof. Let $T = \bigcup_{x \in V \cup \{g\}} \text{supp}_\omega(x)$, a nonempty set of inner exponents, and $p_T(x) = ([\omega^\beta]x)_{\beta \in T} \in \mathbb{R}^T$. It is injective on the set-sized space $W = V + \mathbb{Q}g$, so $\ell_0(p_T(v + qg)) = q$ is a well-defined $\mathbb{Q}$ -linear functional on $p_T(W)$; extend it through a basis of the set $\mathbb{R}^T$ to ℓ and put $a = \ell \circ p_T$. Choose β_0 below every member of T and $\delta = \omega^{\beta_0}$; then $p_T(C_\delta) = 0$. $\square$

For a set $A \subseteq k((t^{\mathbf{No}}))$ ($k = \mathbb{R}$ or $\mathbb{C}$) let S_A be the union of the outer supports, $V_A = \text{span}_{\mathbb{Q}}(S_A)$ and $\mathcal{H}(A) = k((t^{V_A}))$; let $\mathcal{E}_A$ be the group generated by the elementary shears Φ_s (with real s) that fix A pointwise, in the sense of section 1.4.

Theorem 7.4 (Exact elementary hull [04]). *For every set $A \subseteq k((t^{\mathbf{No}}))$, $\text{Fix}(\mathcal{E}_A) = \mathcal{H}(A) = k((t^{V_A}))$.*

Proof. By [theorem 4.2](#) a nonidentity shear with functional a fixes all of A iff a vanishes on S_A , iff $a(V_A) = 0$; such shears fix every series supported in V_A . If $f \notin \mathcal{H}(A)$, take $g \in \text{supp } f \setminus V_A$ and apply [lemma 7.3](#). $\square$

Corollary 7.5 (Monomial parameters [04]). *For a set $S \subseteq \mathbf{No}$ with $V = \text{span}_{\mathbb{Q}}(S)$, the pointwise stabilizer in $\mathcal{U}_{\mathbf{Oz}}$ of $\{t^g : g \in S\}$ has common fixed field exactly $\mathbb{R}((t^V))$.*

Proof. The elementary subgroup already has this fixed field. Conversely an ordered field automorphism fixing t^g fixes its positive rational powers, hence all t^v , $v \in V$, and strong linearity fixes every series supported on V . $\square$

Naming $\omega = t^{-1}$ gives the fixed field $\mathbb{R}((t^{\mathbb{Q}}))$, much larger than $\mathbb{R}(\omega)$; naming all monomials with real exponents gives $\mathbb{R}((t^{\mathbb{R}}))$ [04]. On sets, $A \mapsto \mathcal{H}(A)$ is extensive, monotone and idempotent, and $\mathcal{H}(A)$ is the smallest Hahn subfield with divisible exponent group containing every monomial of the normal forms of A [04]. For nonmonomial parameters [theorem 7.4](#) computes the fixed field of $\mathcal{E}_A$ only: a general strong automorphism fixing a series may do so by cancellation, and $\mathcal{H}(A)$ is not asserted to be the model-theoretic definable closure [04] ([question 11.3](#)).

7.3 Product actions and nonsolvable stabilizers

Theorem 7.6 (Faithful $\mathbb{R}^\kappa$ actions [09]). *For every set $P \subseteq \mathbf{No}$ and every set cardinal κ there is a faithful action of the full product $\mathbb{R}^\kappa$ by strong 1-automorphisms preserving $\mathbf{Oz}$ and fixing P pointwise.*

Proof. Let E, T be as in [lemma 7.1](#), and choose an ordinal β above κ and all of T . For $j < \kappa$ put $\delta_j = \omega^j$, $\eta_j = \omega^{\beta+j}$ (ordinal addition in the index) and $\theta_j = q_{\beta+j}$. Then $\theta_j|_E = 0$, $\theta_j(C_{\delta_j}) = 0$, $\theta_j(\delta_\ell) = 0$ and $\theta_j(\eta_\ell)$ is the Kronecker delta. For $\mathbf{a} \in \mathbb{R}^\kappa$ let $D_{\mathbf{a}}(t^\gamma) = t^\gamma \sum_{j < \kappa} a_j \theta_j(\gamma) t^{\delta_j}$; the shifts form one well-ordered set, so [proposition 4.4](#) applies with no finiteness condition on $\mathbf{a}$. $D_{\mathbf{a}}$ kills the exponents of P . The $D_{\mathbf{a}}$ commute by (6.1) and add, so $\mathbf{a} \mapsto \text{Exp}(D_{\mathbf{a}})$ is a homomorphism, and $\text{Exp}(D_{\mathbf{a}})(t^{\eta_j}) = t^{\eta_j} \exp(a_j t^{\delta_j})$ detects every a_j . The result is one class relation encoding a set-indexed action. $\square$

Theorem 7.7 (Nonsolvable stabilizers [09]). *For every set $P \subseteq \mathbf{No}$ the pointwise stabilizer of P in $\mathcal{U}_{\mathbf{Oz}}$ is nonsolvable; for every $r \geq 2$ it contains a subgroup whose derived length is at least $r - 1$, and exactly $r - 1$ granted [cited theorem 1.1](#) for $\Gamma = \mathbf{No}$.*

Proof. Choose ordinals $\beta \leq \eta_1 < \dots < \eta_r$ above T , and put $e_j = \omega^{\eta_j}$, $\theta_i = q_{\eta_i}$. Then $e_1 \ll \dots \ll e_r$, $\theta_i(e_j)$ is the Kronecker delta, and each D_{e_j, θ_i} ($j < i$) satisfies the criterion and kills P . The balanced brackets of [lemma 6.4](#) are unchanged, so [lemma 6.5](#) shows that the group generated by all $\text{Exp}(s D_{e_j, \theta_i})$, $s \in \mathbb{R}$, has derived length at least $r - 1$; this already gives nonsolvability. The upper bound applies the filtration by the $r - 1$ scales $C_{e_1}, \dots, C_{e_{r-1}}$ as in [theorem 6.3](#); its group step uses BCH. $\square$

7.4 A definable-closure bound in set-sized Hahn fields [10]

For a set-sized Γ consider the structure $K^\dagger = (K; 0, 1, +, -, \cdot, \mathcal{R}_o, (c)_{c \in k}, P_-, P_0, P_+)$, with the Hahn order added when k is ordered. For a set $A \subseteq K$ of parameters let S_A be the union of the supports of its members, as in [theorem 7.4](#), and put $V_\delta(A) = \text{span}_{\mathbb{Q}}(S_A) + C_\delta$ for $\delta > 0$ (10's S , A_S and $V_h(S)$).

Proposition 7.8 (Definable-closure support bound [10]). *Every element of K definable over A in $K^\dagger$ is supported in $V_\delta(A)$ for every $\delta > 0$; that is, $\text{dcl}(A) \subseteq \bigcap_{\delta > 0} k((t^{V_\delta(A)}))$.*

Proof. Let $g \in \text{supp } f$ lie outside the $\mathbb{Q}$ -subspace $V_\delta(A)$. A basis of the set-sized $\mathbb{Q}$ -space $\Gamma/V_\delta(A)$ gives a $\mathbb{Q}$ -linear $\theta : \Gamma \rightarrow \mathbb{Q} \subseteq k$ with $\theta(V_\delta(A)) = 0$ and $\theta(g) = 1$. The twist $\sigma_{\delta, \theta}^{1+\xi}$ lies in $\mathcal{U}_{\mathcal{R}_o}(\Gamma)$ (theorem 4.18); it fixes k , preserves $\mathcal{R}_o$, commutes with P_-, P_0, P_+ (theorem 3.3), preserves lc and hence the order, and fixes every monomial with exponent in S_A , hence every member of A . By theorem 4.20 it moves f . An element definable over A is fixed by every automorphism of $K^\dagger$ fixing A . $\square$

The bound may be strict. It is not a quantifier-elimination result and does not describe the whole definable closure; it says only that support excluded by a separating exponent functional cannot be introduced by a definition [10]. For $\Gamma = \mathbf{No}$ see corollary 9.11.

8 What the pair $(\mathbf{No}, \mathbf{Oz})$ remembers

Sources 04, 08 and 09 each reconstruct the coefficient field from $\mathbf{Oz}$; 08 and 09 add first-order definitions of the ordinary integers and of Π . All of this is now in the omnific Diophantine report, placed after the three pins, and is cited rather than reprinted:

- Π is defined in $\mathbf{Oz}$ and $\mathbf{Oz}[i]$ by $\exists y(x^2 = 2y^2)$ (`odg:def:thm:ideal`); the first-order definitions of Π and of the graph of ct through ordinary integers are `odg:cor:definablect`, and a positive-existential definition of $\mathbb{Z}$ is `odg:thm:guard`; Pell rigidity is `odg:cor:pell`;
- the multiplier identity $(\Pi : \Pi) = \mathcal{A}_k$, $k = \{0\} \cup \mathcal{A}_k^\times$ and its internal first-order form (`odg:def:thm:multiplier`, `odg:def:cor:internal`); $\text{Frac } \mathbf{Oz} = \mathbf{No}$ (`odg:thm:fractions`);
- the pure ring $\mathbf{Oz}$ interprets $\mathbf{No}$ with $\mathbb{R}$, the order, the finite ring, the infinitesimals and the standard part (`odg:def:thm:realrecovery`); the value group (`odg:def:cor:valuegroup`); every automorphism of $\mathbf{Oz}$ extends uniquely to $\mathbf{No}$ and fixes $\mathbb{R}$ pointwise (`odg:def:thm:autreal`);
- for $\mathbf{Oz}[i]$: $\mathbb{C}$ is definable (`odg:def:cor:internal`), every field automorphism of $\mathbb{C}$ lifts coefficientwise and restriction to $\mathbb{C}$ splits (`odg:def:prop:coeffaut`); a phase twist shows that neither $\mathbf{No}$ nor conjugation is definable from $\mathbf{Oz}[i]$ (`odg:def:lem:twist`, `odg:def:thm:norealaxis`).

In the sources: 04's intrinsic coefficient invariance, 08's characteristic reconstruction, its restriction correspondence and its split restriction to $\text{Aut } \mathbb{C}$, and 09's corollary on all pair automorphisms re-derive `odg:def:thm:multiplier`, `odg:def:thm:autreal` and `odg:def:prop:coeffaut`; source 10's coefficient-rigidity proposition re-derives the first two, with credit (section 9.2). 08's formula $J(x) :\Leftrightarrow \forall n (\text{Int}(n) \wedge n \neq 0 \Rightarrow \exists y x = ny)$ is the definition `odg:cor:definablect` in structure (present at 08's pin); 09's order-free formula $\Phi_J(a) :\Leftrightarrow a \in \mathbf{Oz} \wedge \forall u, v \in \mathbf{Oz} [u^2 - 2v^2 = 1 \Rightarrow \exists b \in \mathbf{Oz} a = ub]$ is a variant using all Pell first coordinates; both are dominated by the one-witness existential formula `odg:def:thm:ideal`, and 08's existential definition of $\mathbb{Z}$ (six witnesses, two inequations) by `odg:thm:guard`. 09 observes that the order symbol can be omitted, positivity being definable by nonzero squares. The following extras are not in the Diophantine report.

Lemma 8.1 (Order on $\mathbf{Oz}$ through fractions [08]). *For $z \in \mathbf{Oz}$, $z \geq 0$ if and only if $\exists a, b \in \mathbf{Oz} (b \neq 0 \wedge a^2 = zb^2)$.*

Proof. If $a^2 = zb^2$ with $b \neq 0$ then $z = (a/b)^2 \geq 0$. Conversely a square root of $z \geq 0$ exists in the real closed field $\mathbf{No}$ and is a quotient a/b of omnific integers (`odg:thm:fractions`). $\square$

The formula is existential with one inequation, consistent with the Diophantine report's result that the nonnegative cone is not positive-existentially definable (`odg:prop:notdiophantine`).

Theorem 8.2 (Zero-cut projections and floor [08, 09]). *In $(\mathbf{No}, +, \cdot, <, \mathbf{Oz})$, equivalently in the fraction field interpreted in the pure ring $\mathbf{Oz}$, the graphs of the projections P_-, P_0, P_+ on all of $\mathbf{No}$ (so $P_0 = \text{ct}$ extended to $\mathbf{No}$) and of the floor map $x \mapsto \lfloor x \rfloor_{\mathbf{Oz}}$ are parameter-free definable. For every monomial $m = t^\alpha$, truncation below α , $T_{<\alpha}(f) = m P_-(f/m)$, is definable with the single parameter m . The same definitions work in $(\mathbb{R}((t^\Gamma)), \mathcal{R}_\mathbb{Z})$ for every nonzero divisible ordered abelian Γ .*

Proof. $\Pi, \mathbb{R}, \mathcal{O}$ and $\mathfrak{m}$ are definable by the results cited above (in the t -convention $\mathcal{O}$ is the finite ring and $\mathfrak{m}$ the infinitesimals). For $x \in \mathbf{No}$, $P_-(x)$ is the unique $j \in \Pi$ with $x - j \in \mathcal{O}$: existence by splitting the support at 0, uniqueness from $\Pi \cap \mathcal{O} = 0$. Then $P_0(x)$ is the unique $c \in \mathbb{R}$ with $x - j - c \in \mathfrak{m}$, since $\mathbb{R} \cap \mathfrak{m} = 0$, and $P_+ = \text{id} - P_- - P_0$ [09]. The floor graph is $y \in \mathbf{Oz} \wedge y \leq x < y + 1$, unique because distinct omnific integers differ by at least one [08]; its value is `odg:thm:floor`. Dividing by m translates exponents by $-\alpha$, P_- keeps those below 0, and multiplying by m restores them [09]. For the set-sized field the proof uses only real coefficients, the Hahn order (the field is real closed since Γ is divisible), and the definitions of Π and $\mathbb{R}$: the formula of `odg:def:thm:ideal` works for every Γ , and the multiplier identity `odg:def:thm:multiplier` holds inside the whole Hahn field by the same one-monomial proof [09]. $\square$

Every pair automorphism therefore preserves $\Pi \oplus \mathbb{R} \oplus \mathfrak{m}$ summandwise, fixes the real summand, and satisfies $\text{ct}(\sigma x) = \text{ct}(x)$ and $\sigma(\lfloor x \rfloor) = \lfloor \sigma x \rfloor$ ([corollary 5.2](#)).

Remark 8.3 (Zero truncation is not the normal form [08, 09]). P_0 is definable on all of $\mathbf{No}$ but is not multiplicative there. Definability of P_- does not define the set of monomials, and truncation at a named monomial is not a uniform definable choice of such a boundary at every valuation; the distinction matters in comparison with work on truncation [4]. Recovering every coefficient by $[\omega^g]x = \text{ct}(\omega^{-g}x)$ would need the monomial map $g \mapsto \omega^g$, and [section 9](#) shows that this is a genuine obstruction.

Source 08's Question 2, whether the reconstruction of $\mathbb{C}$ from the pure ring $\mathbf{Oz}[i]$ is first-order, is answered positively by `odg:def:cor:internal`, whose formula $\exists y (x^2 = 2y^2)$ defines $\Pi_\mathbb{C}$ in $\mathbf{Oz}[i]$ and whose multiplier formula then defines $\mathbb{C}$; 09 gives an independent route, its Pell-divisibility formula with the Gaussian Pell solutions (such as (i, i)) handled, followed by the idealizer. The question is not repeated here.

9 What the pair forgets: monomials and representatives

Theorem 9.1 (Set-parameter invisibility [04, 08, 09, 10]). *For every set $P \subseteq \mathbf{No}$ there is a nonidentity strong real-linear 1-automorphism σ of $\mathbf{No}$ with $\sigma(\mathbf{Oz}) = \mathbf{Oz}$ that fixes $P \cup \mathbb{R} \cup \{\omega\}$ pointwise and sends a positive Conway monomial in $\mathbf{Oz}$ to a nonmonomial in $\mathbf{Oz}$. Explicitly, with β as in [lemma 7.1](#) and $\eta = \omega^\beta$, one may take $\sigma = \sigma_{1,q_\beta}$, and then*

$$\sigma(t^{-\eta}) = t^{-\eta} \exp(-t) = \sum_{n \geq 0} \frac{(-1)^n}{n!} t^{-\eta+n}, \quad \text{that is} \quad \sigma(\omega^{\omega^\beta}) = \sum_{n \geq 0} \frac{(-1)^n}{n!} \omega^{\omega^\beta-n}. \quad (9.1)$$

Consequently, in $\mathfrak{S} = (\mathbf{No}, 0, 1, +, \cdot, <, \mathbf{Oz})$ neither the monomial class $\mathfrak{M} = \{\omega^x : x \in \mathbf{No}\}$ nor the graph of $x \mapsto \omega^x$ is definable with parameters from any set, even after naming $\mathbb{R}$ as a predicate, every ordinary real, ω , and any further set of surreal constants.

Proof. q_β kills C_1 since $\beta > 0$, so $\sigma \in \mathcal{U}_{\mathbf{Oz}}$ by [theorem 4.1](#). It fixes each monomial with exponent in E , hence every member of P by strongness, and it fixes $\mathbb{R}$ and $t^{-1} = \omega$ because $q_\beta(-1) = 0$. Since $q_\beta(-\eta) = -1$, [\(9.1\)](#) follows; all exponents $-\eta + n$ are negative because η is infinite, and the terms at $-\eta$ and $-\eta + 1$ show that the image is not a monomial. A class definable from parameters fixed pointwise by an automorphism of $\mathfrak{S}$ is invariant under it; σ preserves every symbol and moves an element of $\mathfrak{M}$ out of $\mathfrak{M}$. A definable graph of the omega-map would make its range definable by an existential quantifier. $\square$

The three sources give the same construction: 04 with $a = [\omega^\beta]$ and explicit fixing of $\mathbb{R} \cup \{\omega\}$, 08 with the witness $\omega^{\omega^\alpha} \mapsto \sum_n \omega^{\omega^\alpha - n}/n!$ (its sign convention), 09 with q_η ; 10 uses a binomial twist instead of the exponential shear, and its image has exactly two terms ([theorem 9.8](#)). An individual formula uses finitely many parameters; the theorem is the stronger statement that one automorphism fixes an entire prescribed set. Nothing is claimed after naming a proper class of parameters: naming every surreal would remove the symmetry [04, 08]. Expanding the structure by the already definable order, floor or zero-truncation maps cannot remove the obstruction [08].

Corollary 9.2 ([04, 09, 10]). *The monomial class is not determined, as an automorphism-invariant subclass, by $\mathbf{Oz}$, its ordered fraction field and any set of parameters; the standard normal-form section of the natural valuation is not first-order recoverable from these data. The Conway monomials lying in $\mathbf{Oz}$ are not definable in the pure ring $\mathbf{Oz}$ with parameters from any set.*

Proof. The range of the section is $\mathfrak{M}$. For the pure ring, restrict the witness of [theorem 9.1](#) to $\mathbf{Oz}$, with parameters in $\mathbf{Oz}$; automorphisms of $\mathbf{Oz}$ and field automorphisms of $\mathbf{No}$ preserving $\mathbf{Oz}$ correspond (`odg: def: thm: autreal`). $\square$

9.1 No definable representative rule

Write $x \sim_v y$ if $v(x) = v(y)$ on $\mathbf{No}^\times$. A *valuation representative rule* is a function $s : \mathbf{No}^\times \rightarrow \mathbf{No}^\times$ with $v(s(x)) = v(x)$ and $s(x) = s(y)$ whenever $x \sim_v y$. For a set-sized field this is a section of the quotient by $\mathcal{O}^\times$; the formulation also makes sense on the class $\mathbf{No}$ without treating proper-class equivalence classes as elements. No multiplicativity, positivity or monomial-image assumption is imposed [09].

Lemma 9.3 (A whole fiber moves [09]). *For σ and η as in [theorem 9.1](#), every f with $v(f) = -\eta$ satisfies $v(\sigma f - f) = -\eta + 1$ and $\text{lc}(\sigma f - f) = -\text{lc}(f)$.*

Proof. This is [theorem 4.2](#) with $\gamma_0 = -\eta$ and $\theta(-\eta) = -1$: later input terms contribute beyond $-\eta + 1$, higher iterates at $-\eta + n$, $n \geq 2$. $\square$

Theorem 9.4 (No definable section [09]). *For every set $P \subseteq \mathbf{No}$, no P -definable function on $\mathbf{No}^\times$ is a valuation representative rule.*

Proof. Let $x = t^{-\eta}$. Since σ fixes valuations, $\sigma x \sim_v x$ and $s(\sigma x) = s(x)$, while P -definability gives $s(\sigma x) = \sigma(s(x))$. So $s(x)$ would be a fixed point of σ with valuation $-\eta$, contradicting [lemma 9.3](#). $\square$

This is the strongest nondefinability statement of the sources; it covers any representative rule, including one-term ones, and does not follow from character twists. The nondefinability is relative to the field-and-integer-part language; Conway's normal forms remain set-theoretically defined in the ambient foundation [09].

Remark 9.5 (Character twists, one-term series and the RV sort [04, 02]). Bare nondefinability of the coefficient-one monomials already follows from character twists: for a nonzero additive $a : \mathbf{No} \rightarrow \mathbb{R}$ with $a(V_A) = 0$ ([lemma 7.3](#)), $\sum c_g t^g \mapsto \sum c_g e^{a(g)} t^g$ preserves $\mathbf{Oz}$, fixes A and sends some monomial to a non-unit multiple of itself [04]. These twists are the $M_{\chi, \text{id}}$ of [theorem 5.1](#); they are 02's $\tau_{\chi, t}$ ([remark 17.10](#)), the real form of the Diophantine report's phase twist (`odg: def: lem: twist`), and the collection's character twists (`saut: thm: decomp`); they are established and not claimed new. What the shears add is motion inside the leading-term kernel with more than one nonzero term. In particular the class of nonzero one-term series $\{c\omega^g : c \in \mathbb{R}^\times, g \in \mathbf{No}\}$ is not definable from the ordered field, $\mathbf{Oz}$ and set parameters: the shear sends a one-term series to a multiterm one, which rescaling cannot do [04]. Nondefinability also survives adding the sort $\text{RV}(K) = K^\times / (1 + \mathfrak{m})$ with its multiplication, the quotient map and every element of that sort named, since a 1-automorphism acts trivially on it (for the class $\mathbf{No}$ this is a metatheoretic expansion, read formula by formula). This must be distinguished from naming a representative map $\text{RV}(K) \rightarrow K$, which the Conway monomials partly constitute and which would destroy the symmetry by design: $\Phi_s(t^{-\omega})$ has leading term $t^{-\omega}$, yet Φ_s moves $t^{-\omega}$ [04].

Remark 9.6 (Contrasts in the collection). The single dilation $S_2(\sum a_g t^g) = \sum a_g t^{2g}$ defines the canonical monomials, every coefficient and truncation in the language of fields with one automorphism (`dsup: thm: definability` in `docs/surcomplex/single-dilation-hahn-support`); $(\mathbf{No}, \mathbf{Oz})$ does not, even with set parameters. The two expansions differ exactly on the monomial section, and there is no conflict: the shears do not commute with S_2 , in line with that report's centralizer theorem (`dsup: thm: centralizer`). The shears are not exponential: by `cor: question54` of the rigidity report `docs/surreal/exponential-automorphism-rigidity` (from its `thm: no`), every exponential 1-automorphism of $\mathbf{No}$ is the identity, which is stronger than [12, Proposition 5.2] (strongly linear exponential 1-automorphisms are trivial), the result cited by 09. The shears are compatible with that report's displacement lemma: their displacement $-t^{-\eta+1}$ is infinite. Nor do they preserve the omega-map, whose compatibility questions [12, Questions 5.6, 5.7] are untouched here.

9.2 Two-term witnesses, simplicity and the exponential [10]

For a surreal $b > 0$ and a surreal δ with $0 < \delta < 1$, the functional q_b vanishes on C_δ : every member of C_δ is finite, and a finite surreal has no positive exponent in its normal form. Source 10's class twist is the binomial twist $\Sigma_{b, \delta} = \sigma_{\delta, -q_b}^{1+\xi}$; in normal forms,

$$\Sigma_{b, \delta} \left(\sum_g a_g \omega^g \right) = \sum_g a_g \omega^g (1 + \omega^{-\delta})^{q_b(g)}, \quad (9.2)$$

with the real binomial series. Each image has a set-sized support inside $\{g - n\delta\}$, and the proof of [theorem 4.18](#) is elementwise, so $\Sigma_{b, \delta}$ is a strong real-linear 1-automorphism of $\mathbf{No}$ with inverse given by $-q_b$; it fixes $\mathbb{R}$, preserves the order and $\mathbf{Oz}$, and commutes with P_-, P_0, P_+ [10].

Remark 9.7 (An exponent index is not a fixed element [10]). $\Sigma_{b, \delta}$ fixes the monomial $\omega^{-\delta}$ but need not fix the surreal δ : the equality $q_b(-\delta) = 0$ controls the monomial with exponent $-\delta$, not the exponents inside the normal form of δ . (For $\delta = \omega^{-\omega^b}$ one has $\Sigma_{b, \delta}(\delta) \neq \delta$.) This is the inner/outer distinction of [section 7](#), and it is why an automorphism can fix all valuations without commuting with the omega-map.

Theorem 9.8 (Two-term witnesses [10]). *For every set $S \subseteq \mathbf{No}$ there is an ordinal $\beta > 0$ such that, for every surreal δ with $0 < \delta < 1$, $\Sigma_{\beta, \delta}$ fixes S pointwise and*

$$\Sigma_{\beta, \delta}(\omega^{\omega^\beta}) = \omega^{\omega^\beta} + \omega^{\omega^\beta - \delta}, \quad (9.3)$$

a two-term omnific integer and not a Conway monomial. These images are pairwise distinct as δ ranges over $(0, 1)$, and they form a proper class.

Proof. Take β as in [lemma 7.1](#), above the inner supports of the outer supports of the members of S . Then q_β vanishes on every exponent of every member of S , so (9.2) fixes their monomials and, by strongness, the members themselves. Since $q_\beta(\omega^\beta) = 1$, the binomial exponent at ω^{ω^β} is one, which gives (9.3) with no further terms. Both ω^β and $\omega^\beta - \delta$ are positive, since ω^β is infinite, and they differ, so by uniqueness of normal forms the image is an omnific integer and not a monomial. Distinct δ give distinct second exponents. The class $(0, 1)$ is proper, since $\alpha \mapsto \omega^{-(\alpha+1)}$ injects the ordinals into it, and δ is recovered from the image, so the images form a proper class. $\square$

The witness of [theorem 9.1](#) is an infinite series; here two terms suffice. The theorem gives, for each set of parameters, one uniform formula for individual automorphisms indexed by the set parameters β, δ and a proper class of values at one input; it forms no proper class of automorphisms [10].

Corollary 9.9 (Bounded birthdays [10]). *For every ordinal α there is a nonidentity strong ordered-field automorphism of $\mathbf{No}$ that preserves $\mathbf{Oz}$ and fixes every surreal of birthday less than α .*

Proof. The surreals of birthday less than α form a set; apply [theorem 9.8](#). $\square$

The automorphism is not asserted to preserve the birthdays of other elements: the statement is pointwise fixing of a stage, not preservation of the simplicity hierarchy [10], which by [theorem 9.10](#) no such automorphism preserves.

The proof of [theorem 4.20](#) uses only the set-sized support of one input, so the fixed class of a class twist is explicit [10]:

$$\text{Fix}(\Sigma_{b,\delta}) = \left\{ \sum_g a_g \omega^g : q_b(g) = 0 \text{ for every } g \in \text{supp} \right\}, \quad (9.4)$$

and for a nonfixed f , with g_* its greatest exponent having $q_b(g_*) \neq 0$, the leading term of $\Sigma_{b,\delta}(f) - f$ is $a_{g_*} q_b(g_*) \omega^{g_* - \delta}$. The fixed class field is relatively algebraically closed in $\mathbf{No}$, in the finite-polynomial sense, by the proof of [corollary 4.21](#). If $(r_j)_{j \in J}$ is a $\mathbb{Q}$ -linearly independent family of reals with $|J| = 2^{\aleph_0}$, the monomials $\omega^{r_j \omega^b}$ are algebraically independent over (9.4) by [proposition 4.22](#). No exact transcendence degree of the class extension is claimed [10].

Let $\mathfrak{S}^\dagger = (\mathbf{No}; 0, 1, +, -, \cdot, <, \mathbf{Oz}, (r)_{r \in \mathbb{R}}, P_-, P_0, P_+)$, the structure $\mathfrak{S}$ of [theorem 9.1](#) with every real and the three zero-cut projections named; naming all reals uses a set-sized language. Definability has the usual finite-formula meaning, formula by formula in the class structure; no truth predicate for classes is assumed. Simplicity is the relation of being a proper initial segment of a sign sequence, not comparison of birthdays [10].

Theorem 9.10 (What remains undefinable [10]). *For every set $S \subseteq \mathbf{No}$ none of the following is first-order definable over S in $\mathfrak{S}^\dagger$: (i) the monomial class $\mathfrak{M}$; (ii) the graph of $x \mapsto \omega^x$; (iii) the simplicity relation; (iv) the graph of the Gonshor exponential. The class $\mathfrak{M} \cap \mathbf{Oz}$ is not definable over any set of parameters in the ordered ring $(\mathbf{Oz}; 0, 1, +, -, \cdot, <)$.*

Proof. Take $\Sigma = \Sigma_{\beta,\delta}$ from [theorem 9.8](#). It fixes S and every named real and preserves every symbol of $\mathfrak{S}^\dagger$. Items (i) and (ii) follow as in [theorem 9.1](#), since Σ moves the monomial $Z = \omega^{\omega^\beta}$ out of $\mathfrak{M}$.

(iii) A bijection of **No** preserving order and simplicity is the identity [12, Proposition 5.1]: the simplicity tree has one root, each node has two immediate successors distinguished by the order, and each limit sign sequence has a unique continuation, so induction on length fixes every node. A definition of simplicity would make Σ such a bijection, but $\Sigma \neq \text{id}$.

(iv) Put $d = \Sigma(Z) - Z = \omega^{\omega^\beta - \delta}$, which is positive and infinite. The Gonshor exponential is strictly increasing, extends the real exponential and satisfies $\exp(x + y) = \exp(x)\exp(y)$, so $\exp(d)$ is positive infinite and $v(\exp d) \neq 0$. As Σ fixes valuations, $v(\Sigma(\exp Z)) = v(\exp Z)$, whereas $v(\exp(\Sigma Z)) = v(\exp Z) + v(\exp d) \neq v(\exp Z)$. So Σ does not commute with the exponential, while the graph of a definable function is invariant.

For the ordered ring take $S \subseteq \mathbf{Oz}$: the restriction of Σ is an automorphism of $(\mathbf{Oz}; 0, 1, +, -, \cdot, <)$ fixing S , and Z and $\Sigma(Z)$ both lie in $\mathbf{Oz}$, but only Z lies in $\mathfrak{M}$. $\square$

Items (i), (ii) and the ordered-ring clause are [theorem 9.1](#) and [corollary 9.2](#), the order of **Oz** being definable in its pure ring ([lemma 8.1](#)), and the projections being definable in the pair ([theorem 8.2](#)); 10 re-derives them. Items (iii) and (iv) are new here. They need only some nonidentity 1-automorphism of $\mathfrak{S}^\dagger$ fixing S : the merge notes that for (iv) the rigidity report's [cor:question54](#) (every exponential 1-automorphism of **No** is the identity, from [thm:no](#), whose proof is the same valuation comparison) would equally serve with the witness of [theorem 9.1](#). The formal exp of principal units is not the Gonshor exponential, and the class twists are not claimed to preserve the latter; they provably do not. The displacement mechanism is related to the exponential rigidity arguments of [12] and of that report; what is added is that the witness preserves **Oz** and fixes an arbitrary set of parameters [10].

Corollary 9.11 (Support bounds for definable elements [10]; outer bound [merge]). *Let $A \subseteq \mathbf{No}$ be a set and f definable over A in $\mathfrak{S}^\dagger$. Then $f \in \mathbb{R}((t^{V_A}))$, with $V_A = \text{span}_{\mathbb{Q}}(S_A)$ as in [theorem 7.4](#). In particular $B_+(\{f\}) \subseteq B_+(A)$, where $B_+(A) = \bigcup_{s \in A} \bigcup_{g \in \text{supp } s} (\text{supp } g \cap \mathbf{No}^{>0})$ is the positive second-level support: definability creates no new positive inner support position.*

Proof. Every elementary shear fixing A is an automorphism of $\mathfrak{S}^\dagger$ fixing A , so $f \in \text{Fix}(\mathcal{E}_A) = \mathbb{R}((t^{V_A}))$ by [theorem 7.4](#). Each exponent of f is then a finite rational combination of exponents of members of A , and the normal-form support of a finite sum lies in the union of the supports. Source 10's direct proof of the inner bound: if $b > 0$ is in the inner support of an exponent of f but not in $B_+(A)$, then $\Sigma_{b,\delta}$ fixes A and moves f by (9.4); b need not be an ordinal. $\square$

Both are upper bounds only. They do not say that every element obeying them is definable, nor that every coefficient is recoverable from the named projections [10]; compare [question 11.3](#).

Floor, fractions and coefficient rigidity [10]. Source 10 also proves that every ordered-field automorphism stabilizing **Oz** commutes with the floor map, that $\text{Frac } \mathbf{Oz} = \mathbf{No}$ makes any automorphism fixing **Oz** pointwise the identity, and that every ring automorphism of **Oz** extends uniquely to **No** and fixes $\mathbb{R}$ pointwise, through $\Pi = \bigcap_n n\mathbf{Oz}$, the multiplier ring $\mathcal{A}_k$ of Π and its units $\mathbb{R}^\times$, with no strongness assumption. These are [corollary 5.2](#), [odg:thm:fractions](#), [odg:def:thm:multiplier](#) and [odg:def:thm:autreal](#), which 10 credits as prior repository results; they are not reprinted. There is no conflict between $\text{Frac } \mathbf{Oz} = \mathbf{No}$ and the moving of monomials: a ring determines its fraction field up to isomorphism, not a distinguished monomial section inside it; the automorphisms preserve **Oz** setwise and move some of its elements [10].

10 Surcomplex and Gaussian consequences

Identify $\mathbf{No}(i)$ with $\mathbb{C}((t^{\mathbf{No}}))$ by taking coefficientwise real and imaginary parts; the union of the two supports is well ordered. Then $\mathbf{Oz}[i] = \mathbb{Z}[i] \oplus \Pi_{\mathbb{C}} = \mathbf{Oz} + i\mathbf{Oz}$, which is not an ordered ring; conjugation is coefficientwise [04, 08, 09].

Theorem 10.1 (Gaussian stabilizer and conjugation-compatible witnesses [04, 09, 10]). *Theorem 3.3 applies to $(k, \mathfrak{o}) = (\mathbb{C}, \mathbb{Z}[i])$ with the same condition. The real-coefficient shears and actions of sections 7 and 9 extend $\mathbb{C}$ -linearly, commute with conjugation, preserve $\mathbf{No}$ and $\mathbf{Oz}[i]$; for every set of surcomplex parameters they fix the parameters and give nondefinability of the monomials and of any valuation representative rule, even with the real axis, conjugation and every ordinary complex constant named. Their common fixed field is $\mathbb{C}$, and for every set $A \subseteq \mathbf{No}(i)$ the real elementary shears fixing A have common fixed field $\mathbb{C}((t^{V_A}))$.*

Proof. The criterion was proved for every characteristic-zero k and unital $\mathfrak{o}$, and its proof uses no order on coefficients. The functionals q_β are real, so the derivations and their exponentials commute with conjugation and restrict to the real automorphisms on $\mathbf{No}$. Use the union of the real and imaginary supports in lemma 7.1; the moved monomial is real, and lemma 9.3 holds with any nonzero complex leading coefficient. For the fixed field, the first displacement of theorem 4.2 is a product of a nonzero real functional value and a nonzero complex coefficient. The hull statement is theorem 7.4 with $k = \mathbb{C}$ [04]. $\square$

For these real-axis-preserving maps the surreal modulus is equivariant, $|\sigma(z)| = \sigma(|z|)$, since $|z|$ is the nonnegative square root of $z\bar{z}$; this is not a statement that moduli are fixed [09].

Theorem 10.2 (The real axis inside the leading-term kernel [04]). *For every set $A \subseteq \mathbf{No}(i)$ there is a strong $\mathbb{C}$ -linear 1-automorphism of $\mathbf{No}(i)$ that preserves $\mathbf{Oz}[i]$, fixes A pointwise and does not preserve $\mathbf{No}$. Hence $\mathbf{No}$ is not definable in $(\mathbf{No}(i), 0, 1, +, \cdot, \mathbf{Oz}[i])$ with parameters from any set, even with all ordinary complex constants named and the Hahn valuation ring as a predicate.*

Proof. Take β above the inner exponents of the outer supports of A , and the flow Φ_s of theorem 4.1 with $\theta = q_\beta$, $\delta = 1$ and the imaginary time $s = i$. It fixes A and is strong, $\mathbb{C}$ -linear, leading-term preserving and $\mathbf{Oz}[i]$ -preserving. With $\eta = \omega^\beta$, $\Phi_i(t^{-\eta}) = t^{-\eta} \exp(-it) = t^{-\eta} - it^{-\eta+1} - \frac{1}{2}t^{-\eta+2} + \dots$ has a nonreal coefficient, although $t^{-\eta}$ is real; all its exponents are negative. A 1-automorphism preserves every valuation, hence the valuation ring. $\square$

Phase twists preserving $\mathbf{Oz}[i]$ and moving the real axis are already in the collection (`saut:thm:phase`; `odg:def:lem:twist`, `odg:def:thm:norealaxis`); the new points are arbitrary set-parameter fixing *inside the complete leading-term kernel* and multiterm images. The real shears commute with conjugation and the imaginary-time shear does not; the two conclusions must not be combined into a claim that one automorphism commutes with conjugation and moves its fixed field [04].

Proposition 10.3 (A wild coefficient lift [08]). *There is a strong automorphism of $\mathbf{No}(i)$ that preserves $\mathbf{Oz}[i]$, fixes $\mathbb{Z}[i]$ and every Conway monomial, and sends an ordinary real number outside $\mathbb{R}$. So neither $\mathbb{R}$ nor $\mathbf{No}$ is invariant under all field automorphisms preserving $\mathbf{Oz}[i]$.*

Proof. Choose a real r transcendental over $\mathbb{Q}$ and a transcendence basis of $\mathbb{C}$ over $\overline{\mathbb{Q}}$ containing it. On the rational function field it generates, fix $\overline{\mathbb{Q}}$ and the other basis elements and send $r \mapsto r + i$ (inverse $r \mapsto r - i$), and extend to an automorphism ρ of $\mathbb{C}$ by the set-sized algebraic-closure extension theorem; the extension is onto, since its image is algebraically

closed and $\mathbb{C}$ is algebraic over the base. The coefficientwise lift $\hat{\rho}$ (`odg:def:prop:coeffaut`) preserves $\mathbf{Oz}[i]$ and monomials and sends r to $r + i$. $\square$

Corollary 10.4 (Gaussian fixed field [08, 09]). *The common fixed elements in $\mathbf{No}(i)$ of all field automorphisms preserving $\mathbf{Oz}[i]$ are exactly $\mathbb{Q}$; in $\mathbf{Oz}[i]$ they are $\mathbb{Z}$.*

Proof. The shears of [theorem 10.1](#) reduce a common fixed point to $\mathbb{C}$. Every field automorphism of $\mathbb{C}$ lifts coefficientwise and preserves $\mathbf{Oz}[i]$ (`odg:def:prop:coeffaut`). A nonrational algebraic number is moved by a conjugate embedding extended to $\mathbb{C}$; a transcendental one by a translation of a transcendence basis element; rationals are fixed. Finally $\mathbb{Q} \cap \mathbf{Oz}[i] = \mathbb{Z}$. $\square$

This agrees with `saut:prop:fixedfields` of the surcomplex automorphism report, $\text{Fix}(\text{Aut } \mathbf{No}(i)) = \mathbb{Q}$, for the larger collection of all field automorphisms. The real and complex answers differ because a real automorphism preserving $\mathbf{Oz}$ fixes every ordinary real, $\mathbb{R}$ being parameter-free definable in the pair ([corollary 5.2](#), `odg:def:thm:autreal`), whereas a complex one preserving $\mathbf{Oz}[i]$ can move reals unless real structure is required; the explicit witnesses of [theorem 10.1](#) fix all of $\mathbb{C}$ [09]. *Correction* (batch 30): this sentence formerly gave as the reason that a real automorphism “preserves order and cannot move an ordinary real”. Read literally that is false: the Taylor automorphism of `saut:thm:coeffflow` is a strong order-preserving automorphism of $\mathbf{No}$ sending a real b to $b + t^\delta$. It does not preserve $\mathbf{Oz}$ ([example 4.12](#)); preservation of $\mathbf{Oz}$, not of the order, is what fixes $\mathbb{R}$. The comparison and the fixed fields above are unchanged, so this corrects the stated reason and retracts no result. Three mechanisms move the real axis while preserving $\mathbf{Oz}[i]$: the phase twist of the Diophantine report (fixes $\mathbb{C}$, preserves supports), the imaginary-time shear of [theorem 10.2](#) (fixes $\mathbb{C}$ and all leading terms), and the wild lift of [proposition 10.3](#) (fixes monomials, moves $\mathbb{R}$). The real shears of [theorem 10.1](#) preserve the real axis.

Proposition 10.5 (Two-term Gaussian witnesses [10]). *For every set $S \subseteq \mathbf{No}(i)$ there is an ordinal $\beta > 0$ such that for every surreal $0 < \delta < 1$ the $\mathbb{C}$ -linear extension $\tilde{\Sigma}_{\beta,\delta}(x + iy) = \Sigma_{\beta,\delta}(x) + i \Sigma_{\beta,\delta}(y)$ is a strong $\mathbb{C}$ -linear 1-automorphism of $\mathbf{No}(i)$ which (i) fixes S and $\mathbb{C}$ pointwise; (ii) preserves $\mathbf{No}$ and $\mathbf{Oz}[i]$ and commutes with conjugation; (iii) preserves the order of $\mathbf{No}$ and commutes with P_-, P_0, P_+ ; (iv) sends ω^{ω^β} to the two-term element (9.3). Consequently, with parameters from any set, neither the monomial class nor, on the real axis, the graph of the omega-map, the simplicity relation or the graph of the Gonshor exponential is definable in $\mathbf{No}(i)$ expanded by conjugation, the predicate $\mathbf{No}$, the order of $\mathbf{No}$, the predicate $\mathbf{Oz}[i]$, every complex constant and P_-, P_0, P_+ .*

Proof. Apply [theorem 9.8](#) to the set of real and imaginary parts of the members of S . The binomial coefficients in (9.2) are real, so $\tilde{\Sigma}_{\beta,\delta}$ is the same formula on complex-coefficient normal forms; it commutes with conjugation because it acts by one real map on both components and fixes i . [Theorem 3.3](#) with $(k, \mathfrak{o}) = (\mathbb{C}, \mathbb{Z}[i])$ gives (ii) and (iii) for $\mathbf{Oz}[i]$ and the projections. The witness is real. A definition of the monomial class would be preserved, contradicting (iv); for the real-axis statements restrict a putative definition to $\mathbf{No}$ and argue as in [theorem 9.10](#). No global surcomplex exponential is introduced. $\square$

Its fixed class field is the complex-coefficient analogue of (9.4), with the same first displacement, no nontrivial finite orbits and relative algebraic closedness [10]. Beyond [theorem 10.1](#), the proposition adds the two-term witness, the order of the real axis, the projections, and simplicity and the exponential. It differs from the phase twists of the Diophantine report, which move the real axis (`odg:def:lem:twist`): here the real axis and conjugation are preserved, and the obstruction lies inside the fixed real axis [10].

11 Scope, answered questions, and further questions

11.1 What is settled, and status in other reports

For a strong infinitesimal automorphism, preservation of the integer part is exactly a convex-scale annihilation condition; the size of the residual symmetry is measured by a sharp solvability bound in finite rank and by arbitrarily large parameter-fixing actions on **No**; and the omnific predicate recovers the coefficient fields and zero-cut truncation but no definable valuation cross-section [09]. An attempt to reconstruct Conway monomials by a first-order formula in the pair cannot succeed with any set of constants; a classification of strong **Oz**-automorphisms by coefficient and exponent actions alone misses the higher-rank factor $\mathcal{U}_{\mathbf{Oz}}$; and an alleged rank-one nontrivial 1-automorphism preserving a divisible-exponent integer part must violate a support hypothesis, as [example 4.12](#) shows [09].

The results bear on four questions of other reports. They are recorded here; the other reports are not edited by this merge.

- `osq:q:invisible` of `docs/surreal/set-sized-quotients-of-omnific-integers` asks, in its second clause, how much an automorphism of **Oz** can do inside Π while fixing `ct`. *Partly answered*: every automorphism of **Oz** fixes $\mathbb{R}$ and `ct` (`odg:def:thm:autreal`, [corollary 5.2](#)); the *strong* ones are exactly $\{M_{\chi,\tau}\} \ltimes \mathcal{U}_{\mathbf{Oz}}$, with $\mathcal{U}_{\mathbf{Oz}}$ classified by [theorem 3.3](#); they move monomials to infinite series, fix any set of parameters, carry $\mathbb{R}^\kappa$ and have nonsolvable stabilizers ([theorems 7.6, 7.7 and 9.1](#)). Nonstrong automorphisms, and the first clause of that question, remain open. *Status* (batch 30): there are no nonstrong automorphisms of **Oz** ([theorem 23.2](#)), so the second clause is *answered* ([corollary 24.4](#) and [section 32.1](#)); the first clause remains open.
- `odg:def:q:realform` of the Diophantine report asks which natural structure on $\mathbf{Oz}[i]$ rules out the coefficient-fixing twists and determines the real omnific subring. *Negative information, not an answer*: naming the valuation ring, every complex constant, the RV sort, or requiring leading-term fixing does not suffice ([theorem 10.2](#)); and in $(\mathbf{No}(i), \mathbf{Oz}[i])$ even $\mathbb{R} \subseteq \mathbb{C}$ is not invariant ([proposition 10.3](#)).
- The surcomplex automorphism report’s research direction “effective descriptions inside the leading-term kernel” (its section `saut:sec:questions`) is *not* answered. Its **Oz**-stabilizing part is now described by coefficient functionals ([theorem 3.3](#) and [corollary 3.5](#)), with finite support certificates on the common-positive-shift class ([theorem 4.8](#)); no exhaustion or generation theorem is proved ([question 11.4](#)). Its omega-map compatibility questions are untouched.
- `dsn:q:languages` of `docs/foundations-and-computation/definable-surreals-and-omnific-integers` asks about languages between the ordered field with **Oz** and set theory. *Further information, not an answer* [10]: in $(\mathbf{No}, +, \cdot, <, \mathbf{Oz})$, even with every real, the zero-cut projections and any set of parameters named, neither the simplicity relation nor the graph of the Gonshor exponential is definable ([theorem 9.10](#)), whereas order and simplicity together already define ω (`dsn:prop:simplicity-omega`); definable elements obey the support bound of [corollary 9.11](#).

The novelty statements of the batch-26 sources were checked against the tree at their merge: no report contains a classification of **Oz**-preserving automorphisms, the rank dichotomy or set-parameter nondefinability of monomials, and `osq:q:invisible` was recorded as open. Their reconstruction and definability claims were stale ([section 8](#)). Source 10’s statement that none of the report guides it inspected states the support-cut criterion, and its proposal of the criterion, the rank boundary, the fixed fields and the parameter witnesses as its contribution,

were accurate at its pin 58cd8e1, which does not contain this report. They are stale at the merge: this report was placed in f4c9504 and written in 9b80a30, and those results are [theorems 3.3, 4.2, 4.13](#) and [9.1](#). 10 is recorded as an independent re-derivation of them, and its new material is the list in [section 1.2](#). Its other repository statements (the fraction field and the multiplier and coefficient results of the Diophantine report, the phase twists of its reconstruction audit) are accurate.

11.2 Further questions

These questions arise from the proofs; they are not presented as named historical problems [04, 08, 09, 10].

Question 11.1 (Strongness [04, 09, 10]). Which ring automorphisms of $\mathbf{Oz}$ extend to strong field automorphisms of $\mathbf{No}$? Does preservation of $\mathbf{Oz}$ force strongness, or is there a nonstrong $\mathbf{Oz}$ -preserving example? Equivalently for 09: characterize the nonstrong automorphisms of $(\mathbf{No}, \mathbf{Oz})$ that commute with all set-indexed normal-form sums.

[Corollary 5.2](#) removes coefficient motion from this question but does not settle preservation of infinite sums; nothing is inferred from the existence of nonstrong automorphisms of the ambient field [12]. Source 10 asks what characterizes the nonstrong $\mathbf{Oz}$ -preserving 1-automorphisms, and whether every one is strong. *Partial information* [10]: the necessity half of the criterion holds without strongness on every monomial ([remark 3.9](#)); the passage to arbitrary series is what remains. The question stays open. *Status* (batch 30): *answered* by sources 12 and 13 (and independently by source 14, batch 31, [part IV](#)). Every ring automorphism of $\mathbf{Oz}$ extends uniquely to a strongly $\mathbb{R}$ -linear automorphism of $\mathbf{No}$ with strong inverse ([theorem 23.2](#)): all of them extend to strong automorphisms, preservation of $\mathbf{Oz}$ forces strongness, there is no nonstrong example, 09’s characterization is vacuous, and 10’s question has the answer yes. The analogous question for $\mathbf{Oz}[i]$ has a positive answer under conjugation or valuation compatibility and is open otherwise ([theorems 30.1](#) and [30.2](#) and [question 32.1](#)); for general $(k, \mathfrak{o})$ see [question 32.13](#).

Question 11.2 (Admissible logarithmic characters [04]). Give testable necessary and sufficient conditions on an additive $h : \Gamma \rightarrow \mathfrak{m}$ for $t^g \mapsto t^g \exp h(g)$ to define a strong bijective field map. Within that class the integer-part condition is exactly $a_\delta(C_\delta) = 0$.

Common positive support is one sufficient regime; the convex condition alone is not asserted to solve the global problem. This is the problem the surcomplex report describes under “what monomial prescriptions do not automatically give”. Sources 12 and 13 ask it again; it stays open ([section 32.1](#)). Source 14 asks it again as a parameter space for the whole leading-term-fixing factor ([section 37.1](#)).

Question 11.3 (Full relative fixed fields [04]). For a set A of parameters, determine $\text{Fix}((\mathcal{U}_{\mathbf{Oz}})_A)$. It is contained in $\mathbb{R}((t^{V_A}))$, with equality for monomial parameters ([corollary 7.5](#)); which conditions on A force equality?

Source 10 asks the definable-closure analogue: when is the bound of [proposition 7.8](#) sharp, and which support and coefficient conditions characterize definable closure in $\mathfrak{S}^\dagger$? Its upper bounds ([proposition 7.8](#) and [corollary 9.11](#)) are obstructions, not converse criteria; a full answer would have to separate algebraic support generation from the information contributed by the order and by $\mathbf{Oz}$ [10].

Question 11.4 (Generation and exhaustion [04, 08, 10, 15]). How much of $\mathcal{U}_{\mathbf{Oz}}$ is generated by elementary shears and their support-controlled limits, and can an intrinsic summation

convention make this precise without one common shift set for all inputs? Can all strong leading-term-fixing **Oz**-preserving automorphisms be described by a necessary and sufficient uniform support condition on their monomial phases?

No density or generation theorem is claimed; fine-topology convergence is not a substitute for a strong summation rule; the finite-dimensional phase condition of [theorem 4.17](#) is sufficient only. Source 10 asks the same for set-sized divisible Γ with its twists as generators, and for compositions indexed by a well-ordered set: [example 6.13](#) shows that one additive family is not enough, and a transfinite composition is not justified merely because each factor is strong; the accumulated changes must be controlled and a genuine limit map with an inverse proved [10]. Source 15 asks it again for every leading-term-fixing omnific automorphism, now that all are strong, asking for an admissibility condition that gives both global summability and an inverse (its Question 10); a topology realizing all Hahn sums offers no separated topological-group shortcut, since on **No** every one is indiscrete ([corollary 39.9](#)) [15]. *Status* (batch 32): open.

Question 11.5 (Weaker normal-form data [08]). Which normal-form data weaker than the omega-map remove the monomial nondefinability? Set-sized collections of constants do not ([theorem 9.1](#)).

Nor do every real constant, the zero-cut projections and the floor map, and simplicity and the Gonshor exponential are not definable either ([theorem 9.10](#)) [10].

Question 11.6 (Non-divisible exponent groups [merge]). For non-divisible Γ and $\mathfrak{o} = \mathbb{Z}$, is a strong k -linear 1-automorphism preserving $\mathcal{R}_{\mathbb{Z}}$ characterized by a convex-annihilator condition, as single flows are ([theorem 4.10](#))?

This records the scope correction of [remark 4.11](#). Source 10 assumes divisibility throughout, and its example with $\Gamma = \mathbb{Z}$ is stated for $\mathfrak{o} = k$ ([remark 4.23](#)); the question stays open.

Question 11.7 (Other integer parts [10]). Which parts of the support-cut criterion survive for noncanonical integer parts of a real closed Hahn field, or for subrings whose positive-support condition is replaced by a more general support family?

The proof uses closure under every positive rational power of a monomial and the exact boundary at exponent zero. An arbitrary integer part need not present that boundary termwise, and the criterion should not be transferred to it without a replacement for the least-forbidden-shift argument [10]. Source 10 also records that its construction does not answer the question of [12] on nontrivial omega-map-preserving automorphisms, since its witnesses do not preserve the omega-map, nor factorization problems for omnific integers, nor the order-unit question of the holonomic-rigidity report ([docs/surcomplex/holonomic-rigidity-for-entire-hahn-functions](#)). Source 12 asks the analogous question for automatic strongness ([question 32.13](#)).

Part II

Algebraic-parameter rigidity of the omnific integers

This part is manuscript 02. Its question is different from that of [part I](#): not which automorphisms preserve a support condition, but whether **Oz** admits nonconstant *algebraic* families of embeddings. It does not: the decisive property is that every nonzero element divides an element with roots of every order. It does admit formal families of all orders. The relation to [part I](#) is recorded in [section 19](#).

12 Conventions and the root-covered property

12.1 Conventions of this part

Ring homomorphisms are unital, domains are commutative integral domains, and $\mathbb{N} = \{0, 1, 2, \dots\}$; degrees of polynomials are ordinary integers. This part keeps its source's *large-monomial* convention, which is that of the other omnific reports: for a characteristic-zero field k , a unital subring $\mathfrak{o} \subseteq k$ and a nonzero divisible ordered abelian group Γ ,

$$k((X^\Gamma)) = \left\{ \sum_{g \in S} a_g X^g : S \subseteq \Gamma \text{ reverse well ordered} \right\}, \quad X^g = t^{-g},$$

so the leading exponent $\ell(x) = \max \text{supp } x$ is the largest, and $\ell(xy) = \ell(x) + \ell(y)$. The ideal $\Pi_k(\Gamma)$ consists of the series supported in $\Gamma^{>0}$ (the same set as Π in [part I](#)), and $\mathcal{R}_{\mathfrak{o}}(k, \Gamma) = \mathfrak{o} + \Pi_k(\Gamma)$ is the ring $\mathcal{R}_{\mathfrak{o}}$ of [part I](#), the Diophantine report's $\mathcal{R}_{\mathfrak{o}}(\mathbb{k}, \Gamma)$. For surreals $X = \omega$, so $\mathbf{Oz} = \mathcal{R}_{\mathbb{Z}}(\mathbb{R}, \mathbf{No})$ and $\mathbf{Oz}[i] = \mathcal{R}_{\mathbb{Z}[i]}(\mathbb{C}, \mathbf{No})$. Renamings from the source: its $R_\Gamma(D, k)$ and $\mathcal{P}_\Gamma(k)$ are $\mathcal{R}_{\mathfrak{o}}(k, \Gamma)$ and $\Pi_k(\Gamma)$; its principal convex subgroup $H(a)$ is $C(a)$, the transcendence report's notation (C_δ in [part I](#); the quotient report's H_a means the opposite, the elements infinitesimal relative to a); its covering element c is z ; its bounded-scale field K_b is K_{bd} and its stages H_j are $\Gamma_{\leq j}$. Elsewhere in this part the letters R, B, A, K, F, L denote rings and fields local to each statement.

The general theorems are proved for sets. Their conclusions for $\mathbf{Oz}$, $\mathbf{Oz}[i]$, $\mathbf{No}$ and $\mathbf{No}(i)$ use class functions and set-supported normal forms. A statement about one polynomial contains finitely many coefficients and variables, so the degree arguments apply to a class-sized coefficient domain unchanged. No proper-class scheme $\text{Spec}(\mathbf{Oz})$ and no class compactness or Noetherian theorem is used; the finite-type theorem is a theorem about sets, applied to the set-sized models of [section 18](#). Expressions involving $\text{Aut}(\mathbf{Oz})$ or ML of a class ring abbreviate statements about each automorphism or derivation; $(b, x) \mapsto D_b(x)$ is one definable class operation. For cancellation against a class-sized B , the transcendence-degree arguments involve only finite tuples of indeterminates inside one rational-function field [02].

Definition 12.1 ([02, 16]). A domain R is *root-covered* if for every $0 \neq a \in R$ there are $b \in R$ and $z = ab \neq 0$ such that for every $n \geq 1$ some $z_n \in R$ has $z_n^n = z$. (No compatibility between the z_n is required; the terminology is local to this report.)

Fields are root-covered. The interesting examples have few units: $\mathbf{Oz}^\times = \{\pm 1\}$ and $\mathbf{Oz}[i]^\times = \{\pm 1, \pm i\}$, and in general $\mathcal{R}_{\mathfrak{o}}(k, \Gamma)^\times = \mathfrak{o}^\times$ by additivity of leading exponents (the Diophantine report's `odg:prop:units`, `odg:def:lem:units`).

Lemma 12.2 (Localization [02]). *If R is root-covered and $S \subseteq R \setminus \{0\}$ is multiplicative, then $S^{-1}R$ is root-covered.*

Proof. $(a/s)(sb) = z$, and the roots z_n remain. □

Root-covered is sufficient for the conclusions below and is not asserted necessary; failure of the monomial criterion is not used to claim failure of cancellation or stable rigidity [02].

13 The monomial-divisibility criterion

For $a > 0$ in Γ let $C(a) = \{g : |g| \leq na \text{ for some } n \geq 1\}$; a is an *order unit* if $C(a) = \Gamma$.

Theorem 13.1 (Monomial divisibility [02, 16]; after `bst:thm:fieldcriterion`). *For $\Gamma \neq 0$ divisible and $R = \mathcal{R}_\alpha(k, \Gamma)$ the following are equivalent: (i) Γ has no order unit; (ii) for every $0 \neq x \in R$ some monomial X^c with $c > 0$ is divisible by x in R . Then the quotient can be chosen in $\Pi_k(\Gamma)$, and R is root-covered.*

Proof. Assume (i). The support of x lies in $[0, a]$ for some $a > 0$. The inverse of x in the Hahn field has support in the subgroup generated by $\text{supp } x$, hence in $C(a)$ (write $x = \alpha X^h(1 + \varepsilon)$ and expand the geometric series). Choose $c > na$ for all n ; then $X^c x^{-1}$ has strictly positive support, lies in $\Pi_k(\Gamma)$, and $x \cdot X^c x^{-1} = X^c$. Divisibility of Γ gives $(X^{c/n})^n = X^c$. Conversely, if a is an order unit and $x = 1 + X^a$, the unique Hahn quotient $X^c/(1 + X^a) = \sum_{n \geq 1} (-1)^{n-1} X^{c-na}$ has a nonzero coefficient at a negative exponent, so it is not in R . $\square$

This is the transcendence report’s field criterion `bst:thm:fieldcriterion` (`docs/surreal/transcendence-over-bounded-support`: the bounded-support ring is a field if and only if Γ has no order unit, by the same two arguments, the inverse support in a proper convex subgroup and the geometric series of $1 - t^u$), restricted to positive-support elements; its “order unit” is defined identically. Source 02’s audit read only the openings of the omnific reports and did not see it. The positive-support phrasing is what the rigidity theorems use.

Example 13.2 ([02]; [14]). In **No**, $\omega > n$ for all ordinary n , so $\omega^\omega/(1 + \omega) = \omega^{\omega-1} - \omega^{\omega-2} + \omega^{\omega-3} - \dots$ is an omnific integer: $1 + \omega$ divides ω^ω . With exponent group $\mathbb{Q}$ no exponent can absorb the tail. The example is L’Innocente and Mantova’s; the distinction is a larger Archimedean scale, not convergence.

Corollary 13.3 (Omnific root coverage [02, 16]). **Oz** and **Oz**[i] are root-covered: every nonzero element divides a positive real monomial ω^c , with quotient of strictly positive support.

Proof. This is immediate from the global monomial clearing `odg:thm:fractions` of the Diophantine report, applied to x^{-1} : some ω^c makes $\omega^c x^{-1}$ purely infinite. Source 02 gives the same argument through [theorem 13.1](#), using a strict upper bound of the set $\{na : n \geq 1\}$; the coefficients’ order is not used. $\square$

14 Finite-type domains carry no varying embedding

The results of this section concern set-sized fields and domains.

Lemma 14.1 (Unbounded roots [02]). *Let F/K be finitely generated and $u \in F^\times$. If u has n th roots in F for an unbounded set of n , then u is algebraic over K .*

Proof. Otherwise extend u to a transcendence basis $u, t_2, \dots, t_r$, with $[F : E] = d < \infty$ for $E = K(u, t_2, \dots, t_r)$. $Z^n - u$ is Eisenstein at the prime u of $K(t_2, \dots, t_r)[u]$, hence irreducible over E by Gauss’s lemma, so an n th root has degree $n \leq d$. $\square$

Lemma 14.2 (Algebraic constants [02]). *If F/K is finitely generated, the relative algebraic closure K' of K in F is finite over K ; if A is an integral K -algebra with $\text{Frac } A = F$, then $C_A = A \cap K'$ is a field finite over K .*

Proof. With a transcendence basis $\mathbf{t}$ and $[F : K(\mathbf{t})] = d$, every finitely generated $L \subseteq K'$ has $[L : K] = [L(\mathbf{t}) : K(\mathbf{t})] \leq d$ (adjoining algebraically independent indeterminates preserves linear independence; $\mathbf{t}$ stays independent over L). An L of maximal degree contains every element of K' . For $0 \neq a \in C_A$ the minimal polynomial has nonzero constant term, which expresses a^{-1} as a polynomial in a over K . $\square$

Lemma 14.3 (Finitely many unit translates [02]). *Let A be a finitely generated integral K -algebra and $f \in A$ transcendental over K . Only finitely many $\lambda \in K$ make $f - \lambda$ a unit of A .*

Proof. Extend f to a transcendence basis $f, t_2, \dots, t_r$ of $\text{Frac } A$, and put $B_0 = K[f, t_2, \dots, t_r]$. Generators of A are algebraic over $\text{Frac } B_0$; clearing denominators gives $0 \neq d \in B_0$ with A integral over $B = B_0[1/d]$ in the common overring. If $(f - \lambda)^{-1} \in A$, it lies in $\text{Frac } B_0$ and is integral over the integrally closed B , so $f - \lambda$ divides a power of d in the UFD B_0 , hence divides d ; the primes $f - \lambda$ are pairwise nonassociate. The argument does not assume $B \subseteq A$. $\square$

Theorem 14.4 (Finite-type embedding rigidity [02]). *Let R be a characteristic-zero root-covered domain, K a characteristic-zero field, and A a finitely generated integral K -algebra. Every injective homomorphism $\phi : R \rightarrow A$ has image in the field $C_A = \{x \in A : x \text{ algebraic over } K\}$, finite over K , and extends uniquely to $\text{Frac } R \rightarrow C_A$. If K is relatively algebraically closed in $\text{Frac } A$, then $C_A = K$. Neither K -linearity of ϕ nor a map $K \rightarrow R$ is assumed.*

Proof. For $a \neq 0$ take $ab = z$ and $z_n^n = z$. Then $\phi(z) \neq 0$ has roots of all orders, so it is algebraic (lemma 14.1) and a unit of A (lemma 14.2); hence $\phi(a)$ is a unit. If $f = \phi(a)$ were transcendental, every $f - m = \phi(a - m)$, $m \in \mathbb{Z}$, would be a unit (as $a - m \neq 0$), contradicting lemma 14.3. So $\phi(R) \subseteq C_A$, and nonzero images are invertible there. $\square$

The theorem is nontrivial only for set-sized R , such as the Hahn models of section 18; it is not a disguised statement that a proper class cannot inject into a set. One must not replace C_A by the whole relative algebraic closure: $A = \mathbb{Q}[x, \sqrt{2}x]$ has $\text{Frac } A = \mathbb{Q}(\sqrt{2})(x)$ but $C_A = \mathbb{Q}$, since elements of A have rational constant term; no normality of A is assumed [02]. If K is relatively algebraically closed in $\text{Frac } A$, every embedding $R \rightarrow A$ has image in K , and none exists if $\text{Frac } R$ does not embed in K .

Theorem 14.5 (Pointed parameter spaces [02]). *Let R be a set-sized characteristic-zero root-covered domain, $F = \text{Frac } R$, A a finite-type integral F -algebra with an F -algebra map $\varepsilon : A \rightarrow F$. If a ring homomorphism $\psi : R \rightarrow A$ has $\varepsilon\psi$ equal to the inclusion, then $\psi(x) = x$ for all x .*

Proof. ψ is injective, so $\psi(R) \subseteq C_A$. The restriction of ε to the field C_A is injective and fixes F ; for $a \in C_A$, $a - \varepsilon(a) \in C_A$ has image 0, so $C_A = F$. $\square$

Examples are $A = F[s_1, \dots, s_r, 1/h]$ with an F -point where $h \neq 0$; relations, singularities and localizations are allowed, and relative algebraic closedness is not required. The hypothesis is one common finite-type algebra containing the images of all of R , not separate formulas in unrelated algebras. For a finite extension L/F , $L[s]$ contains the root-covered L as constants and has no F -valued augmentation when $L \neq F$ [02].

15 Polynomial and Laurent coefficient rigidity

These proofs work over arbitrary coefficient domains, need no finite generation and assume no preservation of infinite sums [02].

Theorem 15.1 (Polynomial coefficient rigidity [02]). *If R is root-covered and B a domain, every injective $\phi : R \rightarrow B[T_1, \dots, T_n]$ has image in B .*

Proof. $\deg \phi(z) = n \deg \phi(z_n)$ for all n forces $\deg \phi(z) = 0$, and $\deg \phi(a) + \deg \phi(b) = 0$. $\square$

Lemma 15.2 (Laurent units [02]). *For a domain B , the units of $B[U_1^{\pm 1}, \dots, U_q^{\pm 1}]$ are the $b\mathbf{U}^{\mathbf{e}}$ with $b \in B^\times$ and $\mathbf{e} \in \mathbb{Z}^q$.*

Proof. The widths $M_j - m_j$ of the U_j -exponents are nonnegative and additive, so a unit has a single exponent vector. $\square$

Theorem 15.3 (Laurent coefficient rigidity [02]). *If R is root-covered and B a domain, every injective $\phi : R \rightarrow B[\mathbf{U}^{\pm 1}]$ has image in B ; hence so does every injective $\phi : R \rightarrow B[\mathbf{T}, \mathbf{U}^{\pm 1}]$.*

Proof. The largest and smallest U_j -exponents of $\phi(z)$ are divisible by every n , so $\phi(z)$ is a nonzero constant, a unit over $\text{Frac } B$, and $\phi(a)$ is a Laurent monomial by lemma 15.2. If $a \neq -1$ the same holds for $\phi(a) + 1$, which forces $\phi(a)$ to be constant. For mixed targets apply theorem 15.1 over $B[\mathbf{U}^{\pm 1}]$ first. $\square$

Injectivity and a domain target are needed: $x \mapsto x + sD(x)$ into $R[s]/(s^2)$ can be nonconstant with identity specialization (section 17). No noninjective analogue for all targets is claimed [02].

Theorem 15.4 (Strong coefficient invariance [02]). *Let R be root-covered, B any commutative unital ring, and $\Phi : R[X_1, \dots, X_p, U_1^{\pm 1}, \dots, U_q^{\pm 1}] \rightarrow B[Y_1, \dots, Y_{p'}, V_1^{\pm 1}, \dots, V_{q'}^{\pm 1}]$ an isomorphism with $p + q = p' + q' < \infty$. Then $\Phi(R) = B$, $q = q'$ and $p = p'$. In particular, for $R = \mathbf{Oz}$ or $\mathbf{Oz}[i]$, every isomorphism $R[\mathbf{X}_n] \cong B[\mathbf{Y}_n]$ carries R onto B : R is strongly invariant and cancels against an arbitrary coefficient ring with the same number of variables. If B is also root-covered, the equality of total variable counts can be dropped.*

Proof. The target is a domain, so B is, and theorem 15.3 gives $\Phi(R) \subseteq B$. With $F = \text{Frac } \Phi(R) \subseteq L = \text{Frac } B \subseteq E$, $\text{trdeg}_F E = p + q = p' + q' = \text{trdeg}_L E$, so L/F is algebraic. E is a rational function field over F , which has no new algebraic constants (a nonconstant rational function is transcendental, one variable at a time), so $B \subseteq F$, and comparing coefficients gives $B \subseteq \Phi(R)$. Polynomial variables add no units, and the quotient unit groups give $\mathbb{Z}^q \cong \mathbb{Z}^{q'}$. If both rings are root-covered, apply the first step to Φ and Φ^{-1} . $\square$

Equality of total variable counts cannot be dropped for arbitrary B : $R[X_1, X_2] = B[X_2]$ with $B = R[X_1]$. The general statements do not solve the unrestricted cancellation problem for finitely generated algebras; the strong-invariance terminology and history are older [11] [02].

Theorem 15.5 (Automorphisms of polynomial and torus rings [02]). *For a root-covered R , every automorphism of $R[\mathbf{T}]$ or $R[\mathbf{U}^{\pm 1}]$ preserves R ; for polynomial extensions, restriction to R is split surjective onto $\text{Aut } R$, with kernel the R -algebra automorphisms. Every automorphism of $R[\mathbf{T}]$ is $T \mapsto uT + b$ with $u \in R^\times$, $b \in R$, over a unique $\sigma \in \text{Aut } R$; every automorphism of $R[\mathbf{U}^{\pm 1}]$ is $U_i \mapsto u_i \prod_j U_j^{m_{ji}}$ over σ , with $u_i \in R^\times$ and $(m_{ji}) \in \text{GL}_q(\mathbb{Z})$. For $\mathbf{Oz}$, $u \in \{\pm 1\}$; for $\mathbf{Oz}[i]$, $u \in \{\pm 1, \pm i\}$.*

Proof. Apply theorem 15.3 to an automorphism and its inverse; coefficientwise extension is the section. In one variable, $\deg g \deg f = 1$; units give the torus formula via lemma 15.2. $\square$

The affine and torus formulas are standard once coefficients are preserved; the new omnific conclusion is that every ring automorphism preserves $\mathbf{Oz}$. No classification of multivariable polynomial automorphisms over R and no claim about the Jacobian conjecture follows [02].

Batch 32. Source 16 carries this section to monomial parameters of arbitrary rank (part VI): theorems 15.1 and 15.3 hold for every cancellative torsion-free monoid algebra $S[U^{\mathbf{M}}]$ (theorem 47.1), the torus part of theorem 15.5 for every torsion-free exponent group (theorem 48.1),

and [theorem 15.4](#) at every finite rational rank, with exact failure for $\mathbb{Q}^{(\kappa)}$, κ infinite ([theorems 50.1](#) and [50.4](#)).

16 Locally nilpotent derivations, coactions and gradings

A derivation D of a domain is *locally nilpotent* if every element is killed by some iterate, and $\text{ML}(A) = \bigcap_{D \in \text{LND}(A)} \ker D$. Derivations are ring derivations, hence $\mathbb{Z}$ -linear; A need not contain $\mathbb{Q}$.

Lemma 16.1 ([02]). *For a locally nilpotent D of a characteristic-zero domain, $\deg_D(a) = \max\{j : D^j a \neq 0\}$ satisfies $\deg_D(ab) = \deg_D a + \deg_D b$.*

Proof. $D^{r+s}(ab) = \binom{r+s}{r} D^r a D^s b \neq 0$, and higher iterates vanish. $\square$

Theorem 16.2 (Stable locally nilpotent rigidity [02]). *Let R be a characteristic-zero root-covered domain. Every locally nilpotent derivation of any domain containing R kills R , and $\text{ML}(R[T_1, \dots, T_n]) = R$ for every finite n . In particular $\text{ML}(\mathbf{Oz}[\mathbf{T}]) = \mathbf{Oz}$, $\text{ML}(\mathbf{Oz}[i][\mathbf{T}]) = \mathbf{Oz}[i]$, and $\mathbf{Oz}$ has no nonzero locally nilpotent derivation.*

Proof. $\deg_D(z) = m \deg_D(z_m)$ for all m forces $\deg_D z = 0$, hence $\deg_D a = 0$. The partial derivatives $\partial/\partial T_j$ are locally nilpotent with common kernel R . $\square$

Corollary 16.3 (No coactions or gradings [02]). *If R is root-covered, a homomorphism $\psi : R \rightarrow R[s_1, \dots, s_r]$ with identity specialization at $\mathbf{s} = 0$, or $\theta : R \rightarrow R[u_1^{\pm 1}, \dots, u_r^{\pm 1}]$ with identity specialization at $\mathbf{u} = 1$, is the constant inclusion; so there are no nontrivial additive-group or split-torus coactions of finite rank, and every $\mathbb{Z}^r$ -grading of R is concentrated in degree zero.*

Proof. Identity specialization makes the map injective; apply [theorems 15.1](#) and [15.3](#). A grading gives $\sum x_{\mathbf{m}} \mapsto \sum x_{\mathbf{m}} \mathbf{u}^{\mathbf{m}}$. $\square$

A Hahn expansion $\sum a_g X^g$ is not a grading: it has infinitely many terms, and the no-grading result concerns finite homogeneous decompositions. [Theorem 16.2](#) does not say that $R[\mathbf{T}]$ has no locally nilpotent derivation, and it is not a solution of a general stable-rigidity conjecture [5] [02].

17 Euler derivations and formal flows

The parameter s in this section is a new formal indeterminate, not a surreal and not a Hahn monomial.

Proposition 17.1 (Support-preserving Euler derivations [02]). *For additive $\chi : \Gamma \rightarrow (k, +)$, $D_\chi(\sum a_g X^g) = \sum \chi(g) a_g X^g$ is a k -linear derivation of $k((X^\Gamma))$ that preserves $\mathcal{R}_0(k, \Gamma)$, kills $\mathfrak{o}$ and maps $\mathcal{R}_0(k, \Gamma)$ into $\Pi_k(\Gamma)$; and $D_\chi D_\psi = D_\psi D_\chi$ for additive χ, ψ .*

Proof. The coefficient of X^h in xy is a finite sum of $a_g b_{h-g}$, and $\chi(h) = \chi(g) + \chi(h-g)$ splits it; $\chi(0) = 0$. $\square$

In the notation of [part I](#), D_χ is the shift-zero derivation $t^\gamma \mapsto \chi(-\gamma)t^\gamma$. Division by $n!$ is allowed on $\Pi_k(\Gamma)$, and $D_\chi^n(x) \in \Pi_k(\Gamma)$ for $n \geq 1$, so

$$\mathcal{E}_{\chi,s}(x) = \sum_{n \geq 0} \frac{D_\chi^n(x)}{n!} s^n \quad (17.1)$$

has coefficients in $\mathcal{R}_o(k, \Gamma)$; no division of an integer constant occurs.

Theorem 17.2 (Compatible formal flows [02]). *$\mathcal{E}_{\chi,s}$ is a unital ring homomorphism $\mathcal{R}_o(k, \Gamma) \rightarrow \mathcal{R}_o(k, \Gamma)[[s]]$ with identity specialization, satisfying $\mathcal{E}_{\chi,s}\mathcal{E}_{\chi,t} = \mathcal{E}_{\chi,s+t}$; its coefficientwise extension fixing s is an automorphism of $\mathcal{R}_o(k, \Gamma)[[s]]$ with inverse from $-s$, and it induces compatible automorphisms of $\mathcal{R}_o(k, \Gamma)[s]/(s^{N+1})$ for every N .*

Proof. $D_\chi^n(xy)/n! = \sum_r D_\chi^r(x)/r! \cdot D_\chi^{n-r}(y)/(n-r)!$, first in the Hahn field; the coefficient of $s^r t^m$ in both sides of the flow law is $D_\chi^{r+m}(x)/(r!m!)$; each coefficient uses finitely many inputs. Substituting $t = -s$ is valid since both parameters have positive order. $\square$

Theorem 17.3 (Nonalgebraization at the first jet [02]). *If Γ has no order unit and $\chi \neq 0$, the reduction of $\mathcal{E}_{\chi,s}$ modulo s^2 is nonconstant and is induced by no homomorphism $\psi : \mathcal{R}_o(k, \Gamma) \rightarrow \mathcal{R}_o(k, \Gamma)[s]$ with $\text{ev}_{s=0}\psi = \text{id}$, and D_χ is not locally nilpotent.*

Proof. Some $g > 0$ has $\chi(g) \neq 0$; then $D_\chi^n(X^g) = \chi(g)^n X^g \neq 0$. By [theorem 13.1](#) the ring is root-covered, and by [corollary 16.3](#) every such ψ is constant. $\square$

Formal automorphisms exist to *all* orders; the obstruction appears only when a single polynomial family must realize even the first-order part. A formal power series ring is not of finite type, and $R[s]/(s^2)$ is not a domain, so there is no conflict with [theorem 14.4](#) [02]. The flows involve no analytic convergence, no fine-topology limit and no identification of s with a surreal infinitesimal, and they are not an exponentiation of a derivation *inside* the Hahn field [02].

17.1 The monomial orbit field

Let $R = \mathcal{R}_o(k, \Gamma)$ be set sized, $F = \text{Frac } R$ (which contains k : $a = aX^g/X^g$) and $W = \chi(\Gamma)$, a $\mathbb{Q}$ -subspace of k . In $F((s))$ put

$$L_\chi = F\left(\frac{\mathcal{E}_{\chi,s}(X^g)}{X^g} : g > 0\right) = F(e^{ws} : w \in W), \quad e^{ws} = \sum_n w^n s^n / n!. \quad (17.2)$$

It uses only transformed monomials; it is not claimed to contain the image of every Hahn series [02].

Lemma 17.4 (Formal exponential independence [02]). *For distinct $\lambda_j \in F$ the $e^{\lambda_j s}$ are F -linearly independent; for $\mathbb{Q}$ -independent $w_1, \dots, w_r$ the $e^{w_j s}$ are algebraically independent over F .*

Proof. Comparing coefficients of s^n , $n < m$, gives a Vandermonde system. A polynomial relation is a linear relation among $e^{(\alpha \cdot \mathbf{w})s}$ with distinct exponents. $\square$

Theorem 17.5 (Monomial orbit field [02]). $\text{trdeg}_F L_\chi = \dim_{\mathbb{Q}} W$. *If $\chi \neq 0$, L_χ lies in no finitely generated extension of F inside $F((s))$, so $\mathcal{E}_{\chi,s} : R \rightarrow F[[s]]$ factors through no finite-type F -algebra with an F -algebra second map.*

Proof. A $\mathbb{Q}$ -basis (w_j) of W gives algebraically independent $e^{w_j s}$, and each e^{ws} has a power equal to a monomial in finitely many of them, so L_χ is algebraic over $F(e^{w_j s})$. For $0 \neq w \in W$, e^{ws} is transcendental and has roots $e^{(w/n)s}$ in L_χ , so [lemma 14.1](#) excludes a finitely generated overfield, even when $\dim_{\mathbb{Q}} W = 1$. A factorization would put the image of R in a finite-type domain $B \subseteq F[[s]]$ with $\text{Frac } B \supseteq L_\chi$. $\square$

17.2 An omnific family of commuting directions

Theorem 17.6 (Commuting omnific derivations [02]; after `osq:prop:classder`). *For $b \in \mathbf{No}$, $D_b(\sum a_g \omega^g) = \sum q_b(g) a_g \omega^g$ is an $\mathbb{R}$ -linear derivation of $\mathbf{No}$ and, with complex coefficients, a $\mathbb{C}$ -linear derivation of $\mathbf{No}(i)$ commuting with conjugation; it preserves $\mathbf{Oz}$ and $\mathbf{Oz}[i]$. The D_b commute, and every finite family with distinct indices is linearly independent over $\mathbf{Oz}$, respectively $\mathbf{Oz}[i]$.*

Proof. [Proposition 17.1](#) with $\chi = q_b$. For $m_c = \omega^{\omega^c} \in \mathbf{Oz}$, $D_b(m_c) = m_c$ if $b = c$ and 0 otherwise; applying $\sum_j a_j D_{b_j} = 0$ to m_{b_ℓ} gives $a_\ell m_{b_\ell} = 0$ in a domain. $\square$

These derivations, with the test elements m_c , are the quotient report's class-valued derivations `osq:prop:classder` (`docs/surreal/set-sized-quotients-of-omnific-integers`), which combine that report's sources 06, 08 and 10 and state independence over $\mathbb{R}$; the same evaluation gives independence over the omnific ring, as 02 states, and 02 adds their commutation and flows. In the t -convention of [part I](#), $D_b(t^\gamma) = -q_b(\gamma)t^\gamma$.

Corollary 17.7 (Multi-parameter flows [02]). *For distinct $b_1, \dots, b_r$ and $R = \mathbf{Oz}$ or $\mathbf{Oz}[i]$, the map*

$$x \mapsto \sum_{\alpha \in \mathbb{N}^r} \frac{D_{b_1}^{\alpha_1} \cdots D_{b_r}^{\alpha_r}(x)}{\alpha_1! \cdots \alpha_r!} s_1^{\alpha_1} \cdots s_r^{\alpha_r} \quad \text{into } R[[s_1, \dots, s_r]]$$

obeys the additive formal group law; its coordinate directions are independent, and no nonzero coordinate direction is induced by a polynomial family on R .

Proof. Compose one-parameter flows, which commute; independence is [theorem 17.6](#), and the last claim is [theorem 17.3](#) in one parameter. $\square$

Example 17.8 ([02]). For $b = 0$, $q_0(g)$ is the real constant coefficient of the exponent g : $D_0(\omega^r) = r\omega^r$ for real r , $D_0(\omega^\omega) = 0$, and $\mathcal{E}_{0,s}(\omega) = \omega + \omega s + \frac{\omega}{2}s^2 + \frac{\omega}{6}s^3 + \cdots$, every coefficient in $\mathbf{Oz}$ although $\frac{1}{2}, \frac{1}{6} \notin \mathbf{Oz}$; “ ωe^s ” abbreviates this series and is not the surreal exponential. Also $D_0(\omega^\omega/(1+\omega)) = -\omega^{\omega+1}/(1+\omega)^2$, again omnific, supported at $\omega - n$ for positive ordinary n (checked coefficientwise to 29 terms by the placement dossier).

Corollary 17.9 (A continuum of independent orbit elements [02]). *For $R = \mathbf{Oz}$, $F = \mathbf{No}$ (or $R = \mathbf{Oz}[i]$, $F = \mathbf{No}(i)$), the monomial orbit field $F(e^{rs} : r \in \mathbb{R})$ of D_0 has transcendence degree $2^{\aleph_0}$ over F , in the sense of a set-sized transcendence basis with finite algebraic relations.*

Proof. $q_0(r) = r$, and $e^{rs} = \mathcal{E}_{0,s}(\omega^r)/\omega^r$ for $r > 0$. A $\mathbb{Q}$ -basis of $\mathbb{R}$, made positive, has $2^{\aleph_0}$ elements; [lemma 17.4](#) holds over the class field F since each relation uses finitely many coefficients. $\square$

Only a lower bound is asserted for the field generated by the formal images of all Hahn series [02].

Remark 17.10 (Actual automorphisms remain abundant [02]). For additive $\chi : \mathbf{No} \rightarrow \mathbb{R}$ and ordinary $t \in \mathbb{R}$, $\tau_{\chi,t}(\sum a_g \omega^g) = \sum e^{t\chi(g)} a_g \omega^g$ is a support-preserving automorphism of $\mathbf{No}$ and $\mathbf{No}(i)$ preserving $\mathbf{Oz}$ and $\mathbf{Oz}[i]$, with $\tau_{\chi,t} \tau_{\chi,u} = \tau_{\chi,t+u}$ and positive multipliers. These character twists are established [12] and not claimed new (remark 9.5); they prevent reading the rigidity theorems as $\text{Aut}(\mathbf{Oz}) = 1$. What fails is realization of a family by a homomorphism into a polynomial or finite-type parameter algebra: with $\chi = q_0$ the twists send ω^r to $e^{tr} \omega^r$ for all real r at once, which no single polynomial parameter algebra can carry.

18 Set-sized workspaces and a fraction-field warning

Let $\Gamma = \bigoplus_{j \geq 0} \mathbb{Q}e_j$, ordered by the sign of the coefficient at the largest index, so $e_{j+1} > ne_j$ for all n : Γ is divisible with no order unit. For $(\mathfrak{o}, k) = (\mathbb{Z}, \mathbb{R})$ or $(\mathbb{Z}[i], \mathbb{C})$, $R = \mathcal{R}_{\mathfrak{o}}(k, \Gamma)$ is a root-covered set (theorem 13.1), so all the results of this part apply to it in ordinary set theory. The order embedding $e_j \mapsto \omega^j$ of exponent groups induces $X^{e_j} \mapsto \omega^{\omega^j}$ and embeds R in $\mathbf{Oz}$, respectively $\mathbf{Oz}[i]$; the coordinate projections give commuting derivations already here, and replacing $\mathbb{Q}e_j$ by $\mathbb{R}e_j$ gives a set-sized realization of corollary 17.9 [02].

Let $\Gamma_{\leq j} = \bigoplus_{\ell \leq j} \mathbb{Q}e_{\ell}$ (convex) and $K_{\text{bd}} = \bigcup_j k((X^{\Gamma_{\leq j}}))$.

Proposition 18.1 (Bounded-scale fraction field [02]; after `bst:prop:localdensity`). *For this R , $\text{Frac } R = K_{\text{bd}} \subsetneq k((X^{\Gamma}))$.*

Proof. An element of R with positive leading exponent in $\Gamma_{\leq j}$ has all its exponents in $\Gamma_{\leq j}$ by convexity, so quotients lie in K_{bd} . Conversely $y \in k((X^{\Gamma_{\leq j}}))$ gives $X^{e_{j+1}}y \in R$. Finally $\sum_{j \geq 0} X^{-e_j}$ is a Hahn series in no $k((X^{\Gamma_{\leq j}}))$. $\square$

This is the transcendence report's localization proposition `bst:prop:localdensity` together with its field criterion (condition (2) of `bst:thm:fieldcriterion`) for this group. For the full class ring, $\text{Frac } \mathbf{Oz} = \mathbf{No}$ and $\text{Frac } \mathbf{Oz}[i] = \mathbf{No}(i)$ (`odg:thm:fractions`); that fact does not justify replacing K_{bd} by the full Hahn field, since a fixed set-sized group may lack an upper bound for a set of its scales even without an order unit [02].

19 Relation to Part I and the questions of this part

The two parts are complementary. [Part I](#) concerns derivations with *positive* shifts and their exponentials inside the Hahn field; this part concerns shift-zero (support-preserving) derivations and needs an external formal parameter.

- 02's question on formal directions below is partly answered by [part I](#): the positive-shift derivations preserving $\mathbf{Oz}$ are classified exactly, by $c_{\delta}(C_{\delta}) = 0$ (theorem 3.4) and, on common positive shifts, by $a_{\delta}(C_{\delta}) = 0$ (theorem 4.7). Those with a common well-ordered set of shifts integrate *inside* $\mathbf{No}$ to automorphisms of $\mathbf{Oz}$ (theorem 4.8), so they are integrable by an honest automorphism rather than only by a formal flow; for an arbitrary contracting strong one this holds granted [cited theorem 1.1](#) for $\Gamma = \mathbf{No}$ (remark 1.2). The shift-zero derivations, the Hasse–Schmidt extensions of other derivations of $\mathbf{Oz}$, and derivations with no common well-ordered shift set are not covered.
- There is no conflict with theorem 16.2: the shift derivations $D_{\delta,\theta}$ restrict to derivations of $\mathbf{Oz}$ but are not locally nilpotent, since $D_{\delta,\theta}^n(t^{\gamma}) = \theta(\gamma)^n t^{\gamma+n\delta} \neq 0$ when $\theta(\gamma) \neq 0$; and the flows Φ_s are not polynomial in s , since $\Phi_s(t^{-n})$ involves every power of s .
- The character twists of [remark 17.10](#) are the factor $M_{\chi,\text{id}}$ of [theorem 5.1](#).

Theorem 15.5 concerns $\text{Aut}(\mathbf{Oz}[T])$, not $\text{Aut}(\mathbf{Oz})$, and does not bear on `osq:q:invisible`. The rational self-maps of $\mathbf{Oz}$ studied in the quotient report are maps of the ring given by rational functions, not ring automorphisms of $\mathbf{Oz}[T]$; the objects differ. Derivations into set-sized modules vanish (`osq:thm:derivations`), which does not conflict with [theorem 17.6](#), whose values lie in the class module Π .

Question 19.1 (Beyond the scale criterion [02]). For a nonzero divisible Γ *with* an order unit, which rings $\mathcal{R}_o(k, \Gamma)$ remain strongly invariant or stably rigid?

Theorem 13.1 shows that their elements need not divide monomials; it does not show that cancellation or locally nilpotent rigidity fails.

Question 19.2 (Formal directions [02]; re-scoped). Which derivations of $\mathbf{Oz}$ preserve its purely infinite ideal, and which admit an iterative integral Hasse–Schmidt extension $\mathbf{Oz} \rightarrow \mathbf{Oz}[[s]]$? How much larger is this class than the Euler family of [theorem 17.6](#)?

Status. Partly answered for positive shifts by [part I](#), as recorded above; open otherwise. The factorial argument of [theorem 17.2](#) is a sufficient condition for the displayed family and is not asserted to classify all integrable derivations or all formal automorphisms tangent to the identity.

Question 19.3 (Reduced parameter algebras [02, 16]). How far does coefficient rigidity extend to reduced parameter rings with several irreducible components when a prescribed specialization stays injective on the source?

The present proofs apply to domain quotients on which the map stays injective; a component can kill root-cover elements, and the dual-number example shows that nilpotent parameter rings need a different statement. Source 16 asks it again (its Question 12.8): its arbitrary-rank proofs ([part VI](#)) are for domains and do not address components on which the source map has a kernel; it suggests finding conditions under which enough root covers survive in every relevant domain quotient and then comparing the coefficient values across components [16]. *Status* (batch 32): open.

Part III

Automatic strongness and proper embeddings of the omnific pair

This part is built from three manuscripts of batch 30, numbered 11, 12 and 13 here. It has two themes. The first is automatic strongness: every ring automorphism of $\mathbf{Oz}$ extends to a strong automorphism of $\mathbf{No}$, which answers [question 11.1](#) and extends [theorem 5.1](#) and [corollary 5.2](#) to all automorphisms. The second is embeddings: proper self-embeddings of $\mathbf{No}$ can preserve and reflect $\mathbf{Oz}$ and still move a real number, and the strong exact-monomial ones are classified by a convex subgroup of the target value group. Its labels carry the sub-prefix `opa:as:`.

20 Sources, results and conventions of this part

20.1 Three sources and where the merge chose

The three manuscripts were placed together in commit 21375f8 as items 03, 04 and 05 of batch 30. They are numbered 11, 12 and 13 here, after their file prefixes:

- **11** *Coefficient Gaps and Proper Self-Embeddings of the Surreal Field: omnific preservation, topological dichotomies, and nonconjugate copies*, pinned to [cf56b89](#); files prefixed `11-coefficient-gaps-`.
- **12** *Automatic Strongness and Proper Embeddings of the Omnific Integers: constant-term duality, full stabilizers, and the surreal-surcomplex boundary*, pinned to [a6c68ac](#); files prefixed `12-automatic-strongness-`.
- **13** *Automatic Hahn Linearity of Omnific Isomorphisms: residue duality, full stabilizers, and coefficient-drifting self-embeddings*, pinned to [0865f04](#); files prefixed `13-omnific-isomorphisms-`.

The three pins are commits of one line that forked at [c6359e4](#), the placement of source 10. None contains [b3fa9e2](#), the batch-28 write of source 10 into this report. So no author saw [section 3.3](#), [remark 3.9](#) or the paragraph of partial information after [question 11.1](#); their comparisons were with the report of batch 26, and they cite its strong factorization ([theorem 5.1](#)) and its strongness-free [corollary 5.2](#) accurately.

Sources 12 and 13 prove the same main theorems, in nearly the same order: scalar detection of summability, automatic strongness of constant-term-covariant isomorphisms, automatic strongness of $\text{Aut}(\mathbf{Oz})$, the factorization of the whole stabilizer, scalar adjoints, a proper-class obstruction, a proper embedding moving a real, and surcomplex versions. They are printed once, with both sources named. The merge made these choices:

- *Base of the automatic-strongness sections: 12.* Its abstract theorem assumes only a coefficient isomorphism α and derives $F|_k = \alpha$, where 13 assumes $\sigma(k) = \ell$ ([theorem 22.1](#)); it also proves the factorization for set-sized groups, the Archimedean corollary, the regular-cardinal case and retractions for arbitrary strong embeddings. 13's genuinely different material is added with its proofs: the countable binary witness and its finite-row lemma, kept as a second route to detection ([lemma 21.3](#) and [theorem 21.4](#)); isomorphisms between different, possibly non-divisible groups ([proposition 23.1](#) and [corollary 23.4](#)); the coefficient-matrix formula ([proposition 22.3](#)); the example that constant-term preservation does not give preservation of $\mathbf{Oz}$ ([example 23.6](#)); the small-integer-part fraction example ([example 31.2](#)); and the failure of homogeneity ([corollary 27.5](#)).
- *Base of the embedding sections: 11.* Its bottom-gap classification ([theorem 26.3](#)) contains the embeddings of 12 and 13 as instances ([theorem 27.3](#)). The three explicit maps are the same construction with different order isomorphisms $\mathbf{No} \rightarrow \mathbf{No}_{>0}$ or $\mathbf{No} \rightarrow (1, \infty)$; they are printed once. 12's image intersection and annihilator and 13's non-homogeneity are kept.
- *Printed once, not reprinted.* The reconstruction ingredients that all three reprove (the ideal $\Pi = \bigcap_n n\mathbf{Oz}$, the existential test, the multiplier ring, its units, $\text{Frac } \mathbf{Oz} = \mathbf{No}$, fixing of $\mathbb{R}$) are cited from the Diophantine report and [section 8](#); the convex-support criterion, which 12 and 13 reprove, is [theorems 3.3](#) and [3.8](#); the Taylor coefficient automorphism, which 11 reproves, is `saut:thm:coeffflow` of the surcomplex automorphism report.
- *Credit added.* 12 and 13 do not credit `saut:thm:coeffflow`, which already moves a real b to $b + t^\delta$ by a strong automorphism of $\mathbf{No}$; 11 does. The Taylor motion in every embedding below is that theorem's ([section 26.1](#)).

No mathematical error was found in the three manuscripts. They agree with one another and with the rest of the report: automorphisms preserving $\mathbf{Oz}$ fix $\mathbb{R}$ ([corollary 5.2](#)), embeddings need not. Two statements in their delivered audits are wrong and are recorded in [section A.3](#); neither affects the mathematics.

Batch 31. Source 14, written independently of 11–13, re-derives the automatic-strongness results of this part; where it does, 14 is added to the source tag, and its own material and its notation are in [part IV](#).

20.2 What this part settles

- [Question 11.1](#) is *answered* in all its clauses ([theorem 23.2](#)): every ring automorphism of $\mathbf{Oz}$ extends uniquely to a strongly $\mathbb{R}$ -linear automorphism of $\mathbf{No}$ whose inverse is strong, so preservation of $\mathbf{Oz}$ forces strongness. There is no nonstrong $\mathbf{Oz}$ -preserving automorphism, so the clause of 09 (characterize the nonstrong automorphisms of $(\mathbf{No}, \mathbf{Oz})$ that commute with normal-form sums) is vacuous, and 10's variant (is every $\mathbf{Oz}$ -preserving 1-automorphism strong?) has the answer yes.
- [Theorem 5.1](#) and [corollary 5.2](#) hold for all automorphisms of $(\mathbf{No}, \mathbf{Oz})$ and of the full real Hahn pairs with nonzero divisible set-sized exponent group ([theorem 24.1](#)). For Archimedean groups every stabilizer element is a monomial map ([corollary 24.3](#)), which upgrades [theorem 4.13](#) from 1-automorphisms to the whole stabilizer.
- The second clause of `osq:q:invisible` (how much can an automorphism of $\mathbf{Oz}$ do inside Π while fixing `ct`?) is answered ([corollary 24.4](#)); its first clause is untouched.
- Scope. The answers concern $(k, \mathfrak{o}) = (\mathbb{R}, \mathbb{Z})$: $\mathbf{Oz}$ and the full real Hahn pairs. For $(\mathbb{C}, \mathbb{Z}[i])$ they hold when conjugation or the natural valuation is preserved ([section 30](#)); whether every automorphism of $\mathbf{Oz}[i]$ is strong stays open ([question 32.1](#)). For a general characteristic-zero k and unital $\mathfrak{o} \subseteq k$, as in [theorem 3.3](#), nothing here removes the strongness hypothesis ([question 32.13](#)).
- Embeddings: a strong field embedding of the full real Hahn pairs sending monomials to monomials and preserving the integer parts is classified by a coefficient section into the Hahn field on the *bottom gap*; real numbers can move exactly when the positive part of the value-group image is not coinital ([theorem 26.3](#) and [corollary 26.7](#)). Whether every embedding of $(\mathbf{No}, \mathbf{Oz})$ reflecting $\mathbf{Oz}$ is strong has a negative answer if there is a measurable cardinal ([question 32.2](#)).

20.3 Conventions and notation of this part

All three sources use the small- t convention (1.1) of [part I](#), $t^\gamma = \omega^{-\gamma}$, so the purely infinite series have *negative* t -exponents. For a field k and an ordered abelian group Γ put $K_\Gamma = k((t^\Gamma))$, Π_Γ for the series supported in $\Gamma^{<0}$ and $\mathfrak{m}_\Gamma$ for those supported in $\Gamma^{>0}$. For $k = \mathbb{R}$ put

$$\mathcal{R}_\Gamma = \mathbb{Z} \oplus \Pi_\Gamma, \quad \mathcal{A}_\Gamma = \mathbb{R} \oplus \Pi_\Gamma,$$

as in [theorem 4.10](#); for $\Gamma = \mathbf{No}$ these are $\mathbf{Oz}$ and $\mathcal{A}_k$. For $k = \mathbb{C}$ write $\Pi_\Gamma^\mathbb{C}, \mathcal{R}_\Gamma^\mathbb{C} = \mathbb{Z}[i] \oplus \Pi_\Gamma^\mathbb{C}$ and $K_\Gamma^\mathbb{C} = \mathbb{C}((t^\Gamma))$; for $\Gamma = \mathbf{No}$ these are $\Pi_\mathbb{C}, \mathbf{Oz}[i]$ and $\mathbf{No}(i)$. The constant term $\text{ct}(x) = [t^0]x$ is the projection P_0 on all of K_Γ ; it is k -linear and not multiplicative, since $\text{ct}(t^\gamma) \text{ct}(t^{-\gamma}) = 0 \neq \text{ct}(1)$ for $\gamma \neq 0$. The *constant-term pairing* is

$$\langle x, y \rangle_0 = \text{ct}(xy), \quad [t^\gamma]x = \langle x, t^{-\gamma} \rangle_0. \quad (20.1)$$

It is symmetric, k -bilinear and nondegenerate, and not positive definite: it is not an inner product, a trace or a residue integral [12, 13].

Strong maps. An additive map is *strongly additive* if it sends every Hahn summable family to a summable family and commutes with its sum; this is one-directional. A strongly additive

map need not be $\mathbb{R}$ -linear: the Taylor maps below move real constants. As in [section 1.5](#), a *strong automorphism* is one whose inverse is also strongly additive; every theorem below states both directions where it asserts them. For scalar families in k “summable” means “only finitely many nonzero terms”; this is not convergence in $\mathbb{R}$ [13].

Exponent maps. Here τ is an *ordered additive embedding* of exponent groups; when it is onto it is the order automorphism τ of (5.1). The monomial map $M_{\chi,\tau}$ of (5.1) is defined by the same formula for an embedding τ and then is a strong field embedding. A field embedding is *exact-monomial* if it sends every t^γ to a monomial $t^{\tau(\gamma)}$ with coefficient one (11 says “monomial-preserving”). This does *not* mean commuting with the omega-map $x \mapsto \omega^x$, and it excludes characters and principal-unit corrections [11].

Notation	Meaning in part III , and the sources’ symbols
$\mathcal{R}_\Gamma, \mathcal{A}_\Gamma$	$\mathbb{Z} \oplus \Pi_\Gamma$ and $\mathbb{R} \oplus \Pi_\Gamma$. Sources: I_Γ, A_Γ (11); $A_G, B_\mathbb{R}$ (12); $\mathcal{J}_\Gamma, \mathcal{A}$ (13).
$\langle x, y \rangle_0$	$\text{ct}(xy)$. Sources: $\langle x, y \rangle_0$ (12, 13); 11 writes $[x]_0$ for $\text{ct } x$.
$h_\sigma(g), a_{\sigma,\delta}$	The logarithmic character (3.1). Sources: ℓ_g, a_δ (12); L_γ, a_δ (13).
G_τ	The <i>bottom gap</i> of $\tau : \Gamma \hookrightarrow \Delta$ (definition 26.1). Source: C_τ (11).
ρ	An ordered field embedding $\mathbb{R} \rightarrow \mathbb{R}((t^{G_\tau}))$, a <i>coefficient section</i> (11). 13’s $\rho = \sigma _k$ is α here, 12’s letter.
$J_{\tau,\rho}$	The classified embedding (26.1). Source: $\Phi_{\tau,\rho}$ (11).
d	A $\mathbb{Q}$ -derivation of $\mathbb{R}$. Sources: d (11 and the surcomplex report), D (12, 13); D stays reserved for derivations of Hahn fields.
$\rho_{d,\delta}, \mathcal{T}_{d,\delta}$	The Taylor section (26.3) and the ambient Taylor automorphism (26.4), which is Φ_d of <code>saut:thm:coeffflow</code> . Sources: $\rho_{d,\delta}, E_{d,\delta}$ (11); ρ_D (12); E_D (13).
ι_μ	The inner lift (27.1) of an order embedding $\mu : \mathbf{No} \hookrightarrow \mathbf{No}$. Sources: T_h (11); 12’s θ is $\iota_{\mu_{12}}$ and 13’s second τ is $\iota_{\mu_{13}}$.
$\mu_{11}, \mu_{12}, \mu_{13}$	The order isomorphisms (27.2). Sources: p (11), ψ (12), p (13).
$J_{\mu,d}$	The explicit embedding (27.3). Sources: Φ (11), J_D (12), F_D (13).
$T^\dagger, e_\phi, \text{ret}_\phi$	Constant-term adjoint, $\phi^\dagger(1)$ and the linear retraction of a strong embedding ϕ (section 25). Sources: $T^\dagger, d = j^\dagger(1), E_j$ (12); T^*, A (13).
$\lambda_{\mathbf{On}}$	The class functional (25.2). Sources: Λ (12), λ (13).
$\mathcal{V}_\lambda$	The valuation ball $\{x : v(x) > \lambda\}$. Source: $\mathcal{B}_\lambda$ (11).

The collisions are resolved in the new material only, and the tempting false readings are wrong:

- 11’s C_τ is *not* a principal convex subgroup C_δ . It is the largest convex subgroup of the target group meeting $\tau(\Gamma)$ only in 0; in the notation of [definition 2.4](#), $G_\tau = \bigcap_{h \in \tau(\Gamma) > 0} H_h$. For the explicit maps below it happens to be a principal convex subgroup, C_1 or C_ω

(lemma 27.2). 11's $H = \tau(\Gamma)$ is written $\tau(\Gamma)$, leaving H_g to definition 2.4; 11's class $\mathcal{F}$ of finite surreals is C_1 .

- 12's θ is an *exponent embedding*, not an additive profile as in $D_{\delta,\theta}$; it is written $\iota_{\mu_{12}}$. 13 uses τ both for the order automorphism of $M_{\chi,\tau}$ and for its exponent embedding; the second is $\iota_{\mu_{13}}$.
- 11's and 13's order isomorphisms are both called p , and 11's ρ and 13's ρ are different maps; the renamings above separate them.
- 13 writes the factorization as $\sigma = U \circ M$, 12 and theorem 5.1 as $\sigma = M \circ u$; the two agree with $U = MuM^{-1}$, since the monomial maps normalize $\mathcal{U}_{\mathbf{Oz}}$.
- 11's target group Λ is Δ here, 13's letter, so that λ and Λ stay free; 11's Gaussian ring J_Γ and field L_Γ are $\mathcal{R}_\Gamma^\mathbb{C}$ and $K_\Gamma^\mathbb{C}$.
- The large-cardinal report's sign-sequence embedding, called J there and in its dossier, is written $\hat{j}$ here, to keep J for the strong embeddings of this part.

21 Constant-term tests detect Hahn summability

This section is 12's Section 3 and 13's Section 3, printed once. Let k be any field and $K = k((t^\Gamma))$ a full Hahn field; Γ may be a set or $\mathbf{No}$, with set-sized supports. The point is that one scalar functional, ct, together with multiplication, recognizes Hahn summability.

Lemma 21.1 (Finite intersection [12, 13, 14, 15]). *If $S, T \subseteq \Gamma$ are well ordered, then $S \cap (-T)$ is finite.*

Proof. An infinite subset of the well-ordered S contains a strictly increasing sequence $s_0 < s_1 < \dots$, and then $-s_0 > -s_1 > \dots$ would be a descending sequence in T . $\square$

Lemma 21.2 (Triangular detector [12]). *Let $(x_j)_{j \in I}$ be a set-indexed family in K in which each exponent occurs in only finitely many supports. If $\bigcup_j \text{supp } x_j$ is not well ordered, there is $y \in K$ with countable support and all coefficients in $\{0, 1\}$ such that $\langle x_j, y \rangle_0 \neq 0$ for infinitely many j .*

Proof. Choose a strictly decreasing sequence in the union of the supports. Each support meets it in finitely many places, since an infinite subsequence would descend inside a well order. Select recursively a decreasing subsequence $g_0 > g_1 > \dots$ and distinct indices $j_0, j_1, \dots$ with

$$[t^{g_n}]x_{j_n} \neq 0, \quad [t^{g_m}]x_{j_n} = 0 \quad (m > n) : \quad (21.1)$$

at each stage discard the finitely many places lying in the previously selected supports, and pick a later place and an index whose support contains it. Choose $c_n \in \{0, 1\}$ recursively with $\sum_{m \leq n} c_m [t^{g_m}]x_{j_m} \neq 0$; the new variable c_n has the nonzero coefficient $[t^{g_n}]x_{j_n}$, so at most one of the two choices is forbidden, over every field. Put $y = \sum_n c_n t^{-g_n}$, a Hahn series because the exponents $-g_n$ increase. By (21.1), $\langle x_{j_n}, y \rangle_0$ is exactly that finite sum, nonzero for every n . $\square$

Lemma 21.3 (Binary finite-row lemma [13]). *Let $(L_n)_{n \in \mathbb{N}}$ be nonzero linear forms on k^J , each using only finitely many coordinates. There is $a \in \{0, 1\}^J$ with $L_n(a) \neq 0$ for infinitely many n .*

Proof. Let $E_n \neq \emptyset$ be the finite support of L_n . If $E = \bigcup_n E_n$ is finite, some coordinate j has a nonzero coefficient in infinitely many rows, and $a = e_j$ works. If E is infinite, select rows $n_0 < n_1 < \dots$ inductively. At stage r finitely many coordinates have been assigned, and the rows after n_{r-1} still use infinitely many coordinates; choose n_r whose support contains an unassigned coordinate j_r , set the other new coordinates of that row to 0, and choose $a_{j_r} \in \{0, 1\}$ so that $L_{n_r}(a) \neq 0$ (the two choices change $L_{n_r}(a)$ by a nonzero field element). Mark all of E_{n_r} as assigned; later choices do not change this row. Finally set the remaining coordinates to 0. $\square$

The lemma holds in every characteristic, including over $\mathbb{F}_2$, and uses no uncountability, genericity or analytic argument. Asking for *every* row to be nonzero would be false: over $\mathbb{F}_2$ the forms $x, y, x + y$ cannot all be nonzero [13].

Theorem 21.4 (Scalar detection of Hahn summability [12, 13, 14, 15]). *For a set-indexed family $(x_j)_{j \in I}$ in the full Hahn field K the following are equivalent:*

- (i) (x_j) is Hahn summable;
- (ii) for every $y \in K$, only finitely many j have $\langle x_j, y \rangle_0 \neq 0$;
- (iii) the same holds for every y with countable support and coefficients in $\{0, 1\}$.

If they hold, then for every y

$$\left\langle \sum_j x_j, y \right\rangle_0 = \sum_j \langle x_j, y \rangle_0, \quad (21.2)$$

a finite scalar sum, and $\sum_j x_j$ is the unique element with this property.

Proof. (i) $\Rightarrow$ (ii): with $S = \bigcup_j \text{supp } x_j$ and $T = \text{supp } y$, only exponents in the finite set $S \cap (-T)$ contribute to a constant term (lemma 21.1), and each occurs in finitely many x_j ; all rearrangements in (21.2) are finite. (ii) $\Rightarrow$ (iii) is trivial. (iii) $\Rightarrow$ (i): the tests $y = t^{-g}$ give pointwise finiteness, and if the union of the supports were not well ordered, lemma 21.2 would give a countable binary y violating (iii). Uniqueness follows from the tests t^{-g} and nondegeneracy. $\square$

Second route to (iii) $\Rightarrow$ (i) [13]. Assume (iii); monomial tests give pointwise finiteness. If the union S is not well ordered, choose $\gamma_0 > \gamma_1 > \dots$ in S . Each support meets $\{\gamma_n\}$ finitely, so infinitely many distinct indices $j_0, j_1, \dots$ meet it. The forms $L_r(a) = \sum_{n: \gamma_n \in \text{supp } x_{j_r}} a_n [t^{\gamma_n}] x_{j_r}$ on $k^{\mathbb{N}}$ are nonzero and use finitely many coordinates, so lemma 21.3 gives binary a_n with infinitely many $L_r(a) \neq 0$. For $y = \sum_n a_n t^{-\gamma_n}$, a Hahn series, one has $\text{ct}(y x_{j_r}) = L_r(a)$, contradicting (iii). $\square$

The two routes differ in how cancellation is prevented: 12 first makes the family triangular and then chooses coefficients; 13 keeps the family and chooses coefficients row by row. 12's witness is also countable and binary, although 12 does not state it; 13 does. Every witness has countable set support, so the theorem holds for $\Gamma = \mathbf{No}$ [12, 13]. A third route, 14's induced matching, whose witness has all coefficients 1 and involves no choice of coefficients, is printed in section 34, and 15's disjoint-row variant of it in section 40.

Corollary 21.5 (Countable certificates [13]). *Every nonsummable set-indexed family has a countable subfamily and a countably supported binary y for which infinitely many products $x_j y$ have nonzero constant term.*

Proof. An exponent occurring in infinitely many supports gives a monomial witness; otherwise use lemma 21.2. $\square$

This does not say that summable families are countable: a Hahn sum may have an arbitrarily large set-sized support. It says that a failure of summability has a countable certificate.

Example 21.6 (Monomial tests are not enough [13, 15]). The family $x_n = t^{-n}$, $n \geq 1$, has each exponent in one member, but its supports descend. Each monomial test detects at most one member, whereas the Hahn series $y = \sum_{n \geq 1} t^n$ has $\text{ct}(yx_n) = 1$ for every n .

Remark 21.7 (Finiteness spaces [12, 13, 14]). [Lemma 21.1](#) is finite-intersection orthogonality, and its linearization is the finiteness-space duality of Ehrhard, applied to generalized power series by Blute, Cockett, Jacqmin and Scott [3]. Neither support duality nor the notion of a strong linear map is new; what the sources prove directly is the scalar criterion with its cancellation argument and set supports over the class **No**. It is an external characterization of summation, not first-order definability of summation, and not convergence in the valuation topology: a Hahn-summable family need not converge in that topology [13].

22 Constant-term-covariant isomorphisms are strong

Theorem 22.1 (Automatic strongness from constant-term covariance [12, 13, 14, 15]). *Let $K = k((t^\Gamma))$ and $L = l((t^\Delta))$ be full Hahn fields, the exponent groups sets or **No**, with set supports. Let $F : K \rightarrow L$ be a field isomorphism and $\alpha : k \rightarrow l$ a field isomorphism with*

$$\text{ct}(Fx) = \alpha(\text{ct } x) \quad (x \in K). \quad (22.1)$$

Then $F(r) = \alpha(r)$ for all $r \in k$, F is α -semilinear, and F and F^{-1} are strongly additive.

Proof. For $r \in k$ and $x \in K$, multiplicativity and (22.1) give $\text{ct}((F(r) - \alpha(r))F(x)) = \alpha(\text{ct}(rx)) - \alpha(r)\alpha(\text{ct } x) = 0$. As F is onto, $F(r) - \alpha(r)$ pairs to zero with all of L , so $F(r) = \alpha(r)$; semilinearity follows. Let (x_j) be summable with sum s . For $y \in L$ write $y = F(z)$; then $\langle F(x_j), y \rangle_0 = \alpha(\langle x_j, z \rangle_0)$ is nonzero for finitely many j ([theorem 21.4](#) in K), so $(F(x_j))$ is summable ([theorem 21.4](#) in L). With $w = \sum_j F(x_j)$, $\langle F(s), y \rangle_0 = \alpha(\sum_j \langle x_j, z \rangle_0) = \sum_j \langle F(x_j), y \rangle_0 = \langle w, y \rangle_0$ for all y , so $F(s) = w$. The inverse satisfies the hypotheses with α^{-1} . $\square$

13, 14 and 15 prove this with the extra hypothesis $F(k) = l$ and $\alpha = F|_k$; 12 derives it. The proof detects summability of the image family before any target sum is used, and it does not represent arbitrary strong functionals, which would fail for **No** ([theorem 25.6](#)). Surjectivity enters twice: it supplies a preimage of every target test y , and it identifies the image of k . An embedding supplies only the tests in its image; nondegeneracy of the pairing on the image does not detect all target support obstructions. This is a real boundary, not a removable word: the embeddings of [section 27](#) satisfy (22.1) with $\alpha = \text{id}$ and move real coefficients; they are strongly additive, so they contradict coefficient rigidity without surjectivity, not strongness [12, 13].

Corollary 22.2 (The summation operation is recovered [12, 14]). *An isomorphism of full Hahn fields that carries the coefficient field onto the coefficient field and commutes with the constant term up to that coefficient isomorphism respects the whole set-indexed Hahn summation. It need not preserve the monomial sections.*

Proposition 22.3 (Coefficient matrix [13]). *If in [theorem 22.1](#) $k = l$ and $\alpha = \text{id}$, then for all $f \in K$ and $\delta \in \Delta$*

$$[t^\delta]F(f) = \text{ct}(F^{-1}(t^{-\delta})f) = \sum_{\gamma \in \text{supp } f} [t^\gamma]f [t^{-\gamma}]F^{-1}(t^{-\delta}), \quad [t^\delta]F(t^\gamma) = [t^{-\gamma}]F^{-1}(t^{-\delta}), \quad (22.2)$$

with finite sums. So the coefficient matrix of F is the inverse transpose of that of F^{-1} , with exponent reversal from the pairing.

Proof. $[t^\delta]F(f) = \text{ct}(F(f)t^{-\delta}) = \text{ct}(F(fF^{-1}(t^{-\delta}))) = \text{ct}(fF^{-1}(t^{-\delta}))$ by (22.1); expand the last constant term, which has finitely many contributions by lemma 21.1. $\square$

The formula presupposes the global well-ordering of the image, which theorem 22.1 proves; a term-by-term formal argument alone would leave both the coefficient and the well-ordering gaps open [13].

23 Automatic strongness of the omnific stabilizer

23.1 What the integer part reconstructs

All three sources reprove the reconstruction of the coefficients from the integer part: $\Pi = \bigcap_n n\mathbf{Oz}$ (lemma 2.3), the existential test $\exists y (x^2 = 2y^2)$ (odg: def: thm: ideal), the multiplier ring $(\Pi : \Pi) = \mathcal{A}_k$ with units $\mathbb{R}^\times$ (odg: def: thm: multiplier), $\text{Frac } \mathbf{Oz} = \mathbf{No}$ (odg: thm: fractions; also [14, Proposition 2.4.5], which records it as [6, Theorem 32]), and the fixing of $\mathbb{R}$ by every automorphism (odg: def: thm: autreal, corollary 5.2). They credit these as prior work, and they are not reprinted. The one form needed below that is not already stated in the report is 13's version for isomorphisms between different, possibly non-divisible groups.

Proposition 23.1 (Pair isomorphisms preserve the constant term [12, 13, 14, 15]). *Let Γ, Δ be nonzero ordered abelian groups, sets or $\mathbf{No}$, and let $F : K_\Gamma \rightarrow K_\Delta$ be an ordered field isomorphism of the real Hahn fields with $F(\mathcal{R}_\Gamma) = \mathcal{R}_\Delta$. Then F fixes $\mathbb{R}$ pointwise, maps $\Pi_\Gamma, \mathbb{R}$ and $\mathfrak{m}_\Gamma$ onto $\Pi_\Delta, \mathbb{R}$ and $\mathfrak{m}_\Delta$, and $\text{ct}(Fx) = \text{ct}(x)$ for all x . If Γ and Δ are divisible, “ordered” may be omitted.*

Proof. By lemma 2.3, $\Pi_\Gamma = \bigcap_n n\mathcal{R}_\Gamma$ for every ordered abelian group, so $F(\Pi_\Gamma) = \Pi_\Delta$. The multiplier description $\{x : x\Pi_\Gamma \subseteq \Pi_\Gamma\} = \mathcal{A}_\Gamma$ holds for every nonzero Γ : a nonzero coefficient of x at $\delta > 0$ gives $\text{ct}(xt^{-\delta}) \neq 0$. As F is onto, $F(\mathcal{A}_\Gamma) = \mathcal{A}_\Delta$, and the units $\mathbb{R}^\times$ correspond, so F maps $\mathbb{R}$ onto $\mathbb{R}$; a field isomorphism $\mathbb{R} \rightarrow \mathbb{R}$ is the identity. The order is preserved by hypothesis, or because divisible groups give real closed Hahn fields whose order is defined by squares. The infinitesimals $\{x : |x| < 1/n \text{ for all } n\}$ are $\mathfrak{m}_\Gamma$, so $F(\mathfrak{m}_\Gamma) = \mathfrak{m}_\Delta$, and the decomposition $K = \Pi \oplus \mathbb{R} \oplus \mathfrak{m}$ gives the rest. $\square$

For $\Gamma = \Delta$ divisible this is also a consequence of the parameter-free definitions of theorem 8.2; for $\Gamma = \Delta = \mathbf{No}$ it is part of corollary 5.2. The coefficient field is reconstructed as a set and then fixed pointwise; an ordered embedding $\mathbb{R} \hookrightarrow \mathbf{No}$ need not have image $\mathbb{R}$, and section 27 gives embeddings that move reals by infinitesimals [13].

23.2 The main theorem

Theorem 23.2 (Automatic strongness of omnific automorphisms [12, 13, 14, 15]). *Every unital ring automorphism σ of $\mathbf{Oz}$ extends uniquely to a field automorphism $\tilde{\sigma}$ of $\mathbf{No}$. The extension preserves the order and fixes $\mathbb{R}$ pointwise, and $\tilde{\sigma}$ and $\tilde{\sigma}^{-1}$ are strongly $\mathbb{R}$ -linear: for every Hahn-summable family (f_j) the family $(\tilde{\sigma}(f_j))$ is summable and $\tilde{\sigma}(\sum_j f_j) = \sum_j \tilde{\sigma}(f_j)$. Consequently*

$$\text{Aut}(\mathbf{Oz}) \cong \text{Aut}(\mathbf{No}; \mathbf{Oz}) = \text{Aut}_{\mathbb{R}}^{\text{str}}(\mathbf{No}; \mathbf{Oz}), \quad (23.1)$$

by restriction, with extension to fractions as inverse; here $\text{Aut}(\mathbf{No}; \mathbf{Oz})$ denotes the field automorphisms of $\mathbf{No}$ preserving $\mathbf{Oz}$ setwise and the right-hand side those among them that are strongly $\mathbb{R}$ -linear with strong inverse.

Proof. Extension, uniqueness, order and fixing of $\mathbb{R}$ are `odg:def:thm:autreal`. By [corollary 5.2](#) (or [proposition 23.1](#)) the extension commutes with $P_0 = \text{ct}$, without any strongness assumption. [Theorem 22.1](#) with $k = l = \mathbb{R}$ and $\alpha = \text{id}$ gives strongness of $\tilde{\sigma}$ and its inverse. $\square$

The dependencies are not circular [13, 14, 15]:

$$\mathbf{Oz} \longrightarrow \Pi \longrightarrow \mathcal{A}_k \longrightarrow \mathbb{R} \longrightarrow (\Pi, \mathbb{R}, \mathfrak{m}) \longrightarrow \text{ct} \longrightarrow \text{Hahn summation}.$$

The first five steps use only algebra, the order and finite decompositions; the residue criterion is used only after the constant term has been recovered. The group notation in [\(23.1\)](#) is schematic, in the sense of [section 1.4](#): it concerns individual class maps and their compositions, and forms no class of all of them.

Remark 23.3 (What is and is not asserted [12, 13, 14, 15]). The theorem does not say that every automorphism of the field $\mathbf{No}$ is strong: Kaplan, Krapp and Serra construct automorphisms of $\mathbf{No}$ that are not strongly additive [[12](#), Example 4.4], and the theorem shows that none of them preserves $\mathbf{Oz}$. Nor does it say that the automorphisms of $(\mathbf{No}, \mathbf{Oz})$ are few: [part I](#) constructs large families of them, and they are strong by construction. The theorem is a regularity statement. It identifies the summation operation as preserved by these maps; invariance under automorphisms is not first-order definability of summation. It also gives an intrinsic certificate for a transported sum: a proposed value g of $\tilde{\sigma}(\sum_j f_j)$ is correct exactly when $\text{ct}(yg) = \sum_j \text{ct}(y\tilde{\sigma}(f_j))$ for every $y \in \mathbf{No}$, each a finite equation [13]. With [theorems 3.3](#) and [5.1](#) the theorem answers [question 11.1](#); see [section 32.1](#).

Corollary 23.4 (Full real Hahn pairs [12, 13, 14, 15]). *Every isomorphism F in [proposition 23.1](#) is strongly $\mathbb{R}$ -linear with strongly $\mathbb{R}$ -linear inverse. In particular, for a nonzero divisible set-sized Γ , every field automorphism of K_Γ stabilizing $\mathcal{R}_\Gamma$ is strongly $\mathbb{R}$ -linear.*

Proof. [Proposition 23.1](#) and [theorem 22.1](#) with $\alpha = \text{id}$. $\square$

The corollary assumes an isomorphism of the *ambient* Hahn fields. It does not assert that an abstract isomorphism $\mathcal{R}_\Gamma \cong \mathcal{R}_\Delta$, or an automorphism of $\mathcal{R}_\Gamma$, extends to the Hahn fields when $\text{Frac } \mathcal{R}_\Gamma$ is smaller than K_Γ , which can happen ([example 31.2](#), [[14](#), Proposition 2.4.5]); [12](#) states the automorphism case, [13](#) the isomorphism case [12, 13].

Corollary 23.5 (Determination by monomials [12, 13, 14, 15]). *Two automorphisms of $\mathbf{Oz}$ whose extensions agree on every Conway monomial are equal. An $\mathbf{Oz}$ -preserving field automorphism of $\mathbf{No}$ fixing every monomial is the identity.*

Proof. Both extensions fix $\mathbb{R}$ and commute with the normal-form sum of every element. $\square$

Example 23.6 (Constant-term preservation does not give preservation of $\mathbf{Oz}$ [13]). Choose a $\mathbb{Q}$ -linear $q : \mathbb{R} \rightarrow \mathbb{R}$ with $q(1) = 0$ and $q(\sqrt{2}) = 1$, by extending $\{1, \sqrt{2}\}$ to a Hamel basis, and put $\theta(\gamma) = q(\text{ct } \gamma)$, the constant term of the exponent's own normal form. Then θ is additive, $\theta(1) = 0$ and $\theta(\sqrt{2}) = 1$, and $\Psi(t^\gamma) = t^\gamma \exp(\theta(\gamma)t)$ defines a strong $\mathbb{R}$ -linear 1-automorphism of $\mathbf{No}$ with inverse given by $-\theta$: it is the flow T^1 of [proposition 4.9](#) with $\delta = 1$, which needs only $\theta(1) = 0$, not $\theta(C_1) = 0$. A term at exponent 0 in $\Psi(t^\gamma)$ requires $\gamma = -n$ with $n \in \mathbb{N}$, and $\theta(-n) = 0$; so $\text{ct} \circ \Psi = \text{ct}$ on monomials and, by strongness, everywhere. But

$$\Psi(t^{-\sqrt{2}}) = t^{-\sqrt{2}} \exp(-t)$$

has coefficient $\frac{1}{2}$ at the positive exponent $2 - \sqrt{2}$, so $\Psi(\mathbf{Oz}) \not\subseteq \mathbf{Oz}$. Constant-term preservation suffices for automatic strongness of isomorphisms, but not for preservation of $\mathbf{Oz}$. The merge notes the agreement with [theorem 4.1](#): here $\theta(C_1) \neq 0$, since $\sqrt{2} \in C_1$.

24 The whole stabilizer

Theorem 24.1 (Factorization of every pair automorphism [12, 13, 14, 15]). *Let $\Gamma = \mathbf{No}$, or a nonzero divisible set-sized ordered abelian group. Every field automorphism σ of $K_\Gamma = \mathbb{R}((t^\Gamma))$ with $\sigma(\mathcal{R}_\Gamma) = \mathcal{R}_\Gamma$ factors uniquely as*

$$\sigma = M_{\chi, \tau} \circ u, \quad u \in \mathcal{U}_{\mathcal{R}_\mathbb{Z}}(\Gamma),$$

where $\tau(\gamma) = v(\sigma(t^\gamma))$ is an order automorphism of Γ and $\chi(\gamma) = \text{lc}(\sigma(t^\gamma))$ is a positive character; u satisfies the equivalent conditions of [theorem 3.3](#), in particular $a_{u, \delta}(C_\delta) = 0$ for every $\delta > 0$. Conversely every such composition preserves $\mathcal{R}_\Gamma$. So the stabilizer is $\{M_{\chi, \tau}\} \ltimes \mathcal{U}_{\mathcal{R}_\mathbb{Z}}(\Gamma)$, and $\text{Aut}(\mathbf{Oz}) \cong \mathcal{U}_{\mathbf{Oz}} \rtimes \{M_{\chi, \tau}\}$.

Proof. By [theorem 23.2](#) or [corollary 23.4](#), σ is strong and fixes $\mathbb{R}$. It preserves the order, hence Archimedean equivalence, which is expressed with ordinary integers; so it maps the valuation ring onto itself and is valued in the sense of [section 5](#). Every character into $\mathbb{R}^\times$ of a divisible group is positive. [Theorem 5.1](#) now applies, with $k = \mathbb{R}$ and $\mathfrak{o} = \mathbb{Z}$. $\square$

This is [theorem 5.1](#) and the second sentence of [corollary 5.2](#) without the hypothesis of strongness. 12 proves the factorization for $\mathbf{No}$ and set-sized groups, 13 for $\mathbf{No}$, in the order $\sigma = U \circ M_{\chi, \tau}$ ([section 20.3](#)); 14 proves it for both, in the order $M \circ u$, reading the leading data from order and the fixed reals, so the existence of the factorization needs no strongness, and the composition law it states is the one after (5.1). All three, and 15, reprove the convex-support criterion: 12 by the least forbidden shift, which is source 10's route ([theorem 3.8](#)), and 13, 14 and 15 by a polynomial in the rational parameter q with infinitely many zeros, which is 04's route ([theorem 3.3](#)); those proofs are printed there. 13's nontrivial example, the shear $t^\gamma \mapsto t^\gamma \exp(q_1(\gamma)t)$ with $\omega^\omega \mapsto \omega^\omega \exp(-\omega^{-1})$, is [example 4.3](#).

Remark 24.2 (Admissibility is retained [12, 13, 14, 15]). [Theorem 24.1](#) classifies *existing* automorphisms. An arbitrary array of additive a_δ with $a_\delta(C_\delta) = 0$ is not thereby shown to define one: the monomial substitution must be summable on every well-ordered support, multiplicative and invertible with the same admissibility. Local convex vanishing is not proved sufficient, and no class-wide exponential–logarithm correspondence is used or established ([question 11.2](#)). The theorem removes a hypothesis on actual automorphisms; it licenses no unchecked infinite substitution.

Corollary 24.3 (Archimedean groups [12, 14, 15]). *For a nonzero divisible Archimedean set-sized Γ , every field automorphism of $\mathbb{R}((t^\Gamma))$ stabilizing $\mathcal{R}_\Gamma$ is a monomial map $M_{\chi, \tau}$.*

Proof. In [theorem 24.1](#), $u \in \mathcal{U}_{\mathcal{R}_\mathbb{Z}}(\Gamma) = \{\text{id}\}$ by [theorem 4.13](#). $\square$

[Theorem 4.13](#) gave this among strong maps; automatic strongness removes the qualification. No claim is made that omnific-preserving automorphisms are trivial: the monomial factor, and in higher rank the factor $\mathcal{U}$, can be large [12]. For $\Gamma = \mathbb{Q}$ this is the form $t^q \mapsto c^q t^{\lambda q}$ after [theorem 5.1](#), and for $\Gamma = \mathbb{R}$ the character may be discontinuous ([remark 36.3](#)) [14].

Corollary 24.4 (What an automorphism of $\mathbf{Oz}$ does inside Π [merge]). *Every ring automorphism σ of $\mathbf{Oz}$ satisfies $\text{ct} \circ \sigma = \text{ct}$; its extension fixes $\mathbb{R}$ and is $M_{\chi, \tau} \circ u$, where τ is an*

order automorphism of $\mathbf{No}$, χ a positive character and $u \in \mathcal{U}_{\mathbf{Oz}}$. Thus on Π it acts first by u , then by the exponent automorphism and positive monomial rescaling $M_{\chi, \tau}$. Here u is a strong leading-term-fixing automorphism whose logarithmic character has, for every $g \neq 0$, support in H_g , the elements infinitesimal relative to g ([theorem 3.3](#), condition (v)); every such composition occurs, and every monomial image stays in the signed Archimedean block of its exponent up to the action of τ ([proposition 3.10](#)).

Proof. [Theorem 24.1](#) with $\Gamma = \mathbf{No}$, and [theorem 3.3](#) and [proposition 3.10](#) for u . $\square$

With [theorems 7.6](#), [7.7](#) and [9.1](#) this describes the automorphisms of $\mathbf{Oz}$ inside Π as fully as [part I](#) describes the strong ones; the admissibility of logarithmic data remains [question 11.2](#). Source 10's [remark 3.9](#) is subsumed: its monomial condition now holds for every automorphism because every automorphism is strong.

25 Constant-term adjoints and a proper-class obstruction

In this section Γ, Δ are sets, $K = k((t^\Gamma))$ and $L = k((t^\Delta))$ share the coefficient field k , and linearity is over k , not over K . This is the scalar k -linear dual of a full Hahn field; the three-duals report ([docs/surcomplex/three-duals-of-hahn-vector-spaces](#), [duals:](#)) studies K -linear maps on vector-valued Hahn spaces and topological duals, a different convention, and no identification with its functionals is asserted [12, 13]. A k -linear functional $\lambda : K \rightarrow k$ is *strong* if it sends every summable family to a finitely supported scalar family and commutes with the finite sum. T denotes a k -linear map $K \rightarrow L$ in this section only.

Proposition 25.1 (Scalar representability at set size [12, 13, 14, 15]). *Every strong k -linear functional λ on K has a unique representation $\lambda(x) = \text{ct}(ax)$ with $a \in K$; conversely every such functional is strong.*

Proof. Put $a_g = \lambda(t^{-g})$, a function on the set Γ . If its support contained $g_0 > g_1 > \dots$, the monomials t^{-g_n} , with increasing exponents, would form a summable family with all images nonzero; so $a = \sum a_g t^g \in K$. Strongness on the normal-form family of x gives $\lambda(x) = \sum_g [t^g]x a_{-g} = \text{ct}(ax)$, a finite sum; uniqueness is nondegeneracy, and the converse is (21.2). 13 argues on the set $\{\gamma : \lambda(t^\gamma) \neq 0\}$, which meets every well-ordered set finitely and is therefore reverse well ordered, by dependent choice. $\square$

Theorem 25.2 (Adjoint characterization [12, 13, 14, 15]). *A k -linear map $T : K \rightarrow L$ is strongly k -linear if and only if it has a constant-term adjoint $T^\dagger : L \rightarrow K$,*

$$\langle Tx, y \rangle_0 = \langle x, T^\dagger y \rangle_0 \quad (x \in K, y \in L). \quad (25.1)$$

The adjoint is unique, k -linear and strong; $T^{\dagger\dagger} = T$, $(ST)^\dagger = T^\dagger S^\dagger$ and $(T + T')^\dagger = T^\dagger + T'^\dagger$. Multiplication by a fixed series is self-adjoint, and a coefficient-fixing, constant-term-preserving field isomorphism has adjoint its inverse.

Proof. If T is strong, $x \mapsto \text{ct}(T(x)y)$ is a strong functional, represented by a unique $T^\dagger y$ ([proposition 25.1](#)); uniqueness gives linearity. Conversely, given $T^\dagger$ and a summable (x_j) , the scalars $\langle Tx_j, y \rangle_0 = \langle x_j, T^\dagger y \rangle_0$ vanish for almost all j , so (Tx_j) is summable by [theorem 21.4](#), and (21.2) on both sides of (25.1) gives $T(\sum x_j) = \sum Tx_j$. Symmetry of the pairing makes T the adjoint of $T^\dagger$, so $T^\dagger$ is strong. The identities are substitutions; for an isomorphism F with $\text{ct} \circ F = \text{ct}$, $\text{ct}(F(x)y) = \text{ct}(xF^{-1}(y))$. $\square$

The direction “adjoint implies strong” uses only [theorem 21.4](#) and holds for **No** as well; the converse genuinely needs set-sized groups ([theorem 25.6](#)) [13]. The theorem can be read as a concrete transpose statement for the linearized finiteness space; transpose duality in finiteness spaces is not claimed new [12].

Theorem 25.3 (A canonical linear retraction [12]). *Let $\phi : K \hookrightarrow L$ be a strong k -linear field embedding and $e_\phi = \phi^\dagger(1)$. Then $e_\phi \neq 0$ and $\phi^\dagger(\phi(a)y) = a\phi^\dagger(y)$ for $a \in K$, $y \in L$. The map*

$$\text{ret}_\phi(y) = e_\phi^{-1}\phi^\dagger(y)$$

is a strong K -linear retraction (L a K -space through ϕ), with $\text{ret}_\phi \circ \phi = \text{id}_K$. If $\text{ct} \circ \phi = \text{ct}$ then $e_\phi = 1$ and $\text{ret}_\phi = \phi^\dagger$. If ϕ is not onto, ret_ϕ is not multiplicative.

Proof. For $b \in K$, $\text{ct}(b\phi^\dagger(\phi(a)y)) = \text{ct}(\phi(ba)y) = \text{ct}(ba\phi^\dagger(y))$, which gives the projection formula by nondegeneracy; $\text{ct}(e_\phi) = \text{ct}(\phi(1)) = 1$. With $y = 1$, $\phi^\dagger(\phi(a)) = ae_\phi$. Multiplication by a fixed series is strong, so ret_ϕ is strong and K -linear. If $\text{ct} \circ \phi = \text{ct}$, then $\text{ct}(xe_\phi) = \text{ct}(x)$ for all x and $e_\phi = 1$. A multiplicative unital retraction would be an injective field homomorphism with $\text{ret}_\phi(y - \phi(\text{ret}_\phi y)) = 0$, forcing $y = \phi(\text{ret}_\phi y)$ for all y . $\square$

13 proves the case $\text{ct} \circ \phi = \text{ct}$ with ϕ coefficient-fixing (its residue splitting); it adds that $\ker \phi^\dagger$ is the annihilator of $\phi(K)$ for the pairing, and that $L = \phi(K) \oplus \ker \phi^\dagger$ as $\phi(K)$ -spaces. More generally a proper field extension admits no unital ring retraction fixing the subfield [13].

Corollary 25.4 (Tower law and complement [12]). *For composable constant-term-preserving strong k -linear field embeddings $K \xrightarrow{\phi} L \xrightarrow{\psi} P$, $\text{ret}_{\psi\phi} = \text{ret}_\phi \circ \text{ret}_\psi$. For one such embedding $L = \phi(K) \oplus \ker \text{ret}_\phi$ as K -spaces, and $\phi \circ \text{ret}_\phi$ is a strong projection onto $\phi(K)$.*

Proof. Adjoints reverse composition and all normalizing elements are 1; the complement follows from $\text{ret}_\phi \circ \phi = \text{id}$. $\square$

ret_ϕ is not a field retraction and not a positive conditional expectation; neither multiplicativity nor positivity is part of its conclusion, and it is canonical only relative to the two pairings and to ϕ [12].

Example 25.5 (Monomial embeddings [12, 13]). For an ordered group embedding $\tau : \Gamma \hookrightarrow \Delta$ and a character $\chi : \Gamma \rightarrow k^\times$, the strong embedding $M_{\chi,\tau}$ preserves constant terms and

$$M_{\chi,\tau}^\dagger\left(\sum_{\eta} y_{\eta} t^{\eta}\right) = \sum_{g \in \Gamma} \chi(g)^{-1} y_{\tau(g)} t^g,$$

supported in the preimage under an order embedding of a well-ordered set; compare constant coefficients, using $\chi(-g) = \chi(g)^{-1}$. The same formula works for class groups with set supports, by replacement on the target support [13]. If $\eta \notin \tau(\Gamma)$, then $M_{\chi,\tau}^\dagger(t^\eta) = M_{\chi,\tau}^\dagger(t^{-\eta}) = 0$ while $M_{\chi,\tau}^\dagger(1) = 1$, so the retraction is not multiplicative on two monomials. For instance $\Gamma = \Delta = \mathbb{Z}$, $\tau(g) = 2g$, $\chi(g) = 2^g$ over $k = \mathbb{Q}$ give a weighted even-coefficient extraction [12].

For a constant-term-compatible embedding ϕ the source tests show only that $\text{ct}(\phi(h)\phi(x_j))$ is finitely supported in j for each $h \in K$; detection in L needs all $y \in L$. An adjoint supplies a source representative of every target test, so the question is whether all target residue functionals can be pulled back; this locates the remaining embedding questions without answering them [13] ([questions 32.2](#) and [32.3](#)).

Theorem 25.6 (A strong functional with no representing surreal [12, 13, 14, 15]). *For $x \in \mathbf{No}$ put*

$$\lambda_{\mathbf{On}}(x) = \sum_{\alpha \in \mathbf{On}} [t^{-\alpha}]x. \quad (25.2)$$

For each x only finitely many terms are nonzero. $\lambda_{\mathbf{On}}$ is a strongly $\mathbb{R}$ -linear functional, but no $a \in \mathbf{No}$ satisfies $\lambda_{\mathbf{On}}(x) = \text{ct}(ax)$ for all x . Hence the strong $\mathbb{R}$ -linear map $x \mapsto \lambda_{\mathbf{On}}(x) \cdot 1$ of $\mathbf{No}$ has no constant-term adjoint.

Proof. A well-ordered set of exponents meets $-\mathbf{On}$ finitely, since an infinite intersection would give a strictly decreasing sequence of ordinals. The same applies to the total support of a summable family, and only finitely many members contribute at each of those exponents; this gives strongness. As $\lambda_{\mathbf{On}}(t^{-\alpha}) = 1$ for every ordinal α , a representing a would have $[t^\alpha]a = 1$ for all α , a proper-class support. An adjoint evaluated at 1 would be such an a . $\square$

The definition is not a proper-class Hahn sum: for each input one first finds the finite subset of its support lying in $-\mathbf{On}$ and adds finitely many reals; the class only selects coefficients [12]. There is no conflict with [theorem 22.1](#): there the inverse of an isomorphism supplies each needed test element, and no arbitrary strong functional is represented. So the isomorphism theorem extends to $\mathbf{No}$ while the self-duality of [proposition 25.1](#) and [theorem 25.2](#) does not [12, 13].

26 The bottom-gap classification of exact-monomial embeddings

This section and the next three are source 11's, with the embeddings of 12 and 13 as instances. Automorphisms preserving $\mathbf{Oz}$ fix $\mathbb{R}$ because $\mathbb{R}$ is parameter-free definable in the pair (`odg:def:cor:internal`, [theorem 8.2](#)); the definition uses a universally quantified multiplier condition, which a proper embedding need not preserve [11, 12], as the Diophantine report already cautions after `odg:def:thm:autreal`. The existential part survives: by `odg:def:thm:ideal` and `odg:def:thm:homct`, every unital ring homomorphism $\mathcal{R}_\Gamma \rightarrow \mathcal{R}_\Delta$ maps Π_Γ into Π_Δ and preserves the constant term [11]. Throughout, Γ and Δ are nonzero divisible ordered abelian groups (sets, or $\mathbf{No}$ with set supports), so K_Γ and K_Δ are real closed, and $\tau : \Gamma \hookrightarrow \Delta$ is an ordered additive embedding.

Definition 26.1 (Bottom gap [11]). The *bottom gap* of τ is

$$\mathbf{G}_\tau = \{c \in \Delta : n|c| < h \text{ for every } h \in \tau(\Gamma)^{>0} \text{ and every } n \geq 1\}.$$

Since $\tau(\Gamma)$ is divisible, this is $\{c : |c| < h \text{ for every } h \in \tau(\Gamma)^{>0}\}$.

The word “gap” refers to a convex subgroup of the value group, not to an omitted cut of the field [11].

Lemma 26.2 (Intrinsic description [11]). *$\mathbf{G}_\tau$ is the largest convex subgroup of Δ meeting $\tau(\Gamma)$ only in 0; it is divisible, and $\mathbf{G}_\tau = 0$ if and only if $\tau(\Gamma)^{>0}$ is coinitial in $\Delta^{>0}$. The sum $\tau(\Gamma) + \mathbf{G}_\tau$ is direct and ordered lexicographically with $\tau(\Gamma)$ outer: $h < h'$ in $\tau(\Gamma)$ and $c, c' \in \mathbf{G}_\tau$ give $h + c < h' + c'$. Hence a well-ordered family of well-ordered blocks lying in distinct cosets $h + \mathbf{G}_\tau$ has well-ordered union.*

Proof. For $c, c' \in \mathbf{G}_\tau$ and $0 < h \in \tau(\Gamma)$, $|c + c'| < h/2 + h/2$; symmetry and convexity are clear, and $0 < c \in \tau(\Gamma)$ fails the condition at $h = c$. If D is convex with $D \cap \tau(\Gamma) = 0$, then $h \leq n|c|$

with $c \in D$ would put $h \in D$, so $D \subseteq G_\tau$. Convex subgroups of divisible groups are divisible. If $\tau(\Gamma)^{>0}$ is cointial no $c > 0$ lies in G_τ ; otherwise some $c > 0$ lies below every $h \in \tau(\Gamma)^{>0}$, hence below every h/n . For the blocks, $h' - h > |c - c'|$; a nonempty subset of a union of blocks has a least outer coordinate and a least element in that block. $\square$

Theorem 26.3 (Bottom-gap classification [11]). *There is a bijection between*

- (i) *strongly additive field embeddings $J : K_\Gamma \hookrightarrow K_\Delta$ with $J(t^\gamma) = t^{\tau(\gamma)}$ for all γ and $J(\mathcal{R}_\Gamma) \subseteq \mathcal{R}_\Delta$; and*
- (ii) *ordered field embeddings $\rho : \mathbb{R} \hookrightarrow \mathbb{R}((t^{G_\tau}))$,*

given by $\rho = J|_{\mathbb{R}}$ and

$$J_{\tau,\rho}\left(\sum_{\gamma} r_{\gamma} t^{\gamma}\right) = \sum_{\gamma} \rho(r_{\gamma}) t^{\tau(\gamma)}. \quad (26.1)$$

Every $J = J_{\tau,\rho}$ satisfies

$$\begin{aligned} v(Jx) &= \tau(v(x)), & \text{lc}(Jx) &= \text{lc}(x), & \text{ct}(Jx) &= \text{ct}(x), \\ J^{-1}(\mathcal{R}_\Delta) &= \mathcal{R}_\Gamma, & J^{-1}(\Pi_\Delta) &= \Pi_\Gamma. \end{aligned} \quad (26.2)$$

Without strongness, every field embedding with $J(t^\gamma) = t^{\tau(\gamma)}$ and $J(\mathcal{R}_\Gamma) \subseteq \mathcal{R}_\Delta$ still has $J(\mathbb{R}) \subseteq \mathbb{R}((t^{G_\tau}))$.

Here $\text{ct}(Jx) = \text{ct}(x)$ is an equality of coefficient extractions, not multiplicativity of ct .

Proof. Step 1: residues. An ordered embedding of $\mathbb{R}$ into a real Hahn field fixes $\mathbb{Q}$; a rational bound on $|r|$ bounds $|J(r)|$, so $J(r)$ has nonnegative support, and its constant term lies between all rational bounds of r . So $J(r) = r + (\text{a series with positive support})$, with first exponent 0 and first coefficient r when $r \neq 0$. (An embedding of a real closed field into an ordered field preserves positivity, positive elements being squares; so every field embedding here is ordered.)

Step 2: the gap is necessary. For $\gamma > 0$ and $r \in \mathbb{R}$, $rt^{-\gamma} \in \Pi_\Gamma$, so $J(r)t^{-\tau(\gamma)} \in \Pi_\Delta$ by the existential test. Hence every $c \in \text{supp } J(r)$ has $c < \tau(\gamma)$ for all $\gamma > 0$; with $c \geq 0$ and divisibility, $c \in G_\tau$. No strongness is used. If J is strong, expanding x into its summable family of terms gives (26.1), hence uniqueness.

Step 3: every ρ is admissible. For a support S , the blocks $\tau(\gamma) + \text{supp } \rho(r_\gamma)$, $\gamma \in S$, lie in distinct cosets of G_τ in the order of S (lemma 26.2), so (26.1) is a Hahn series. For a summable family (x_j) , the outer exponents form a well-ordered set, finitely many j contribute at each outer exponent, and each block is a finite union of well-ordered supports; so the images are summable, and additivity of ρ gives strong additivity.

Step 4: field operations. $J(1) = 1$. Products of terms map to $\rho(r)t^{\tau(\gamma)}\rho(s)t^{\tau(\eta)} = \rho(rs)t^{\tau(\gamma+\eta)}$, and the product family is summable (the support principle of section 2); regrouping by Step 3 gives multiplicativity. The first block of $J(x)$ starts at $\tau(v(x))$ with coefficient $\text{lc}(x)$, which gives injectivity, the order and the valuation formula.

Step 5: preservation and reflection. For $\gamma < 0$ the block at $\tau(\gamma)$ is negative, for $\gamma > 0$ positive; the zero block is $\rho(r_0)$ with constant term r_0 ; blocks cannot cancel. So $\text{ct}(Jx) = \text{ct}(x)$. If $x \in \mathcal{R}_\Gamma$ its zero block is the integer r_0 and all other blocks are negative. Conversely, if $J(x) \in \mathcal{R}_\Delta$, every positive block is absent and $r_0 \in \mathbb{Z}$, so $x \in \mathcal{R}_\Gamma$; for Π also $r_0 = 0$. $\square$

So these embeddings automatically *reflect* the integer part and Π . The classification is inside the exact-monomial category. It does not cover maps $t^\gamma \mapsto c_\gamma t^{\tau(\gamma)}(1 + u_\gamma)$ with characters or unit corrections, which already occur among the automorphisms of part I, and it does not assert that omnific-preserving embeddings are strong; beyond the category it gives only the necessary coefficient restriction of Step 2 [11] (questions 32.2 and 32.5).

Corollary 26.4 (Surjectivity [11]). *$J_{\tau,\rho}$ is onto if and only if τ is onto. Then $G_\tau = 0$, $\rho = \text{id}_\mathbb{R}$ and $J = M_{1,\tau}$.*

Proof. The valuation formula forces τ onto. Then $G_\tau = 0$ by lemma 26.2, and an ordered embedding $\mathbb{R} \rightarrow \mathbb{R}$ is the identity; reindexing by τ^{-1} inverts $M_{1,\tau}$. $\square$

This agrees with theorem 24.1: an automorphism of the pair that is exact-monomial is $M_{1,\tau}$.

26.1 Taylor sections and the rigidity boundary

A $\mathbb{Q}$ -derivation $d : \mathbb{R} \rightarrow \mathbb{R}$ is additive with $d(rs) = r d(s) + s d(r)$; it need not be continuous. Given a transcendence basis B of $\mathbb{R}/\mathbb{Q}$, arbitrary values $d(b)$, $b \in B$, extend uniquely to $\mathbb{Q}(B)$ and then through the algebraic extension $\mathbb{R}/\mathbb{Q}(B)$: if $p(X) = \sum_j c_j X^j$ is the minimal polynomial of a , then $d(a) = -\sum_j d(c_j) a^j / p'(a)$, with $p'(a) \neq 0$ in characteristic zero, and the extensions agree on finite subextensions. So for a transcendental b there is d with $d(b) = 1$, and every such d vanishes on the real algebraic numbers; the choice uses a transcendence basis of the set $\mathbb{R}/\mathbb{Q}$ and is not computable [11, 12, 13]. This is also the derivation used in `saut:thm:coeffflow`.

Lemma 26.5 (Taylor section [11, 12, 13]). *For $\delta > 0$ in an ordered group and a $\mathbb{Q}$ -derivation d of $\mathbb{R}$,*

$$\rho_{d,\delta}(r) = \sum_{n \geq 0} \frac{d^n(r)}{n!} t^{n\delta} \quad (26.3)$$

is an ordered field embedding of $\mathbb{R}$ into $\mathbb{R}((t^{\mathbb{Z}\delta}))$, with constant term r ; if $d(b) = 1$ then $\rho_{d,\delta}(b) = b + t^\delta$.

Proof. The iterated Leibniz rule $d^n(rs) = \sum_j \binom{n}{j} d^j(r) d^{n-j}(s)$ gives multiplicativity coefficient-wise; the constant term r gives injectivity and the sign; $d^2(b) = d(1) = 0$. $\square$

The ambient version is `saut:thm:coeffflow` of the surcomplex automorphism report, restricted to real coefficients (a derivation of $\mathbb{R}$ extends to $\mathbb{C}$ with $d(i) = 0$, and then the map preserves **No**):

$$\mathcal{T}_{d,\delta}\left(\sum_\lambda r_\lambda t^\lambda\right) = \sum_{\lambda, n \geq 0} \frac{d^n(r_\lambda)}{n!} t^{\lambda+n\delta} \quad (26.4)$$

is a strong automorphism of K_Δ fixing every monomial, preserving valuations and leading coefficients, with inverse $\mathcal{T}_{-d,\delta}$ and $v(\mathcal{T}_{d,\delta}(x) - x) \geq v(x) + \delta$. 11 reproves it and credits the surcomplex report; 12 and 13 use its Taylor ingredient without that credit. On its own it never preserves **Oz** (the remark after example 4.12): with $\delta = 1$, $\Delta = \mathbf{No}$ and $d(b) = 1$ it sends $\frac{b}{2}\omega = \frac{b}{2}t^{-1}$ to $\frac{b}{2}\omega + \frac{1}{2} \notin \mathbf{Oz}$ [11]. What changes below is its interaction with the image of a proper exponent embedding.

Theorem 26.6 (Exact Taylor admissibility [11]). *Let $\delta > 0$ and let $d \neq 0$ be a derivation of $\mathbb{R}$. The strong exact-monomial embedding $\mathcal{T}_{d,\delta} \circ M_{1,\tau}$ preserves $\mathcal{R}_\Gamma$ if and only if $\delta \in G_\tau$, that is $n\delta < \tau(\gamma)$ for all $\gamma > 0$ and $n \geq 1$. It is then $J_{\tau,\rho_{d,\delta}}$ and also reflects $\mathcal{R}_\Gamma$.*

Proof. $\mathcal{T}_{d,\delta} \circ M_{1,\tau}$ is $\sum_\gamma r_\gamma t^\gamma \mapsto \sum_\gamma \rho_{d,\delta}(r_\gamma) t^{\tau(\gamma)}$. If $\delta \in G_\tau$, then $\rho_{d,\delta}$ takes values in $\mathbb{R}((t^{G_\tau}))$ and theorem 26.3 applies. Conversely, for r with $d(r) \neq 0$ the coefficient of $\rho_{d,\delta}(r)$ at δ is nonzero, and Step 2 of that theorem forces $\delta \in G_\tau$. $\square$

Corollary 26.7 (Coinitiality is exactly coefficient rigidity [11]). *For fixed τ , every embedding in [theorem 26.3](#) fixes $\mathbb{R}$ pointwise if and only if $\tau(\Gamma)^{>0}$ is coinitial in $\Delta^{>0}$. Otherwise some such embedding sends a chosen transcendental real b to $b + t^\delta$ with $\delta > 0$.*

Proof. Coinitial means $G_\tau = 0$, and an ordered embedding $\mathbb{R} \rightarrow \mathbb{R}$ is the identity. Otherwise take $0 < \delta \in G_\tau$ and $d(b) = 1$ in [theorem 26.6](#). $\square$

Example 26.8 (A set-sized model [11]). Let $\Gamma = \mathbb{Q}$, $\Delta = \mathbb{Q} \oplus \mathbb{Q}$ ordered lexicographically with the first coordinate dominant, and $\tau(q) = (q, 0)$. Then $G_\tau = \{0\} \oplus \mathbb{Q}$ and $\delta = (0, 1)$ is admissible; for $q > 0$, $\rho_{d,\delta}(r)t^{(-q,0)} = \sum_n \frac{d^n(r)}{n!} t^{(-q,n)}$ has only negative exponents. No proper class is involved.

27 Explicit proper embeddings of $(\mathbf{No}, \mathbf{Oz})$

27.1 Two levels of normal-form substitution

At the *inner* level write an exponent $g \in \mathbf{No}$ in its large- ω Conway normal form $g = \sum_{a \in S} r_a \omega^a$, S reverse well ordered. For an order embedding $\mu : \mathbf{No} \hookrightarrow \mathbf{No}$ put

$$\iota_\mu(g) = \sum_{a \in S} r_a \omega^{\mu(a)}. \quad (27.1)$$

Lemma 27.1 (Inner lift [11]). *ι_μ is an $\mathbb{R}$ -linear ordered embedding of $(\mathbf{No}, +)$ into itself, with the same leading coefficient; its image is the class of normal forms supported in $\mu(\mathbf{No})$, and $\iota_\mu \iota_{\mu'} = \iota_{\mu\mu'}$.*

Proof. An order embedding preserves reverse well-ordering and identifies no exponents; addition is coefficientwise. Reindex a normal form supported in $\mu(\mathbf{No})$ by μ^{-1} for the image. $\square$

At the *outer* level ι_μ is an exponent embedding, and M_{1,ι_μ} is a strong exact-monomial field embedding fixing $\mathbb{R}$ ([theorem 26.3](#) with $\rho = \text{id}$). ι_μ is only additive: it is not a field map, and μ need not be additive. The three sources use three order isomorphisms:

$$\mu_{11}(a) = \begin{cases} (1-a)^{-1}, & a < 0, \\ a+1, & a \geq 0, \end{cases} \quad \mu_{13}(a) = a + \sqrt{a^2 + 1}, \quad \mu_{12}(a) = 1 + a + \sqrt{a^2 + 1}. \quad (27.2)$$

μ_{11} maps $\mathbf{No}$ onto $\mathbf{No}^{>0}$, its branches covering $(0, 1)$ and $[1, \infty)$. μ_{13} maps $\mathbf{No}$ onto $\mathbf{No}^{>0}$ with inverse $y \mapsto (y - y^{-1})/2$, since $(\sqrt{a^2 + 1} + a)(\sqrt{a^2 + 1} - a) = 1$ and $y \mapsto (y - y^{-1})/2$ is strictly increasing. So $\mu_{12} = 1 + \mu_{13}$ maps $\mathbf{No}$ onto $(1, \infty)$. They use only the ordered-field operations of $\mathbf{No}$ [11, 12, 13].

Lemma 27.2 (The three value images and their gaps). *The images $\iota_{\mu_{11}}(\mathbf{No}) = \iota_{\mu_{13}}(\mathbf{No})$ are the additive group of purely infinite surreals, the normal forms with all exponents positive; $\iota_{\mu_{12}}(\mathbf{No})$ consists of the normal forms with all exponents greater than 1. The gaps are*

$$G_{\iota_{\mu_{11}}} = G_{\iota_{\mu_{13}}} = C_1 \text{ [11, 13]}, \quad G_{\iota_{\mu_{12}}} = C_\omega \text{ [merge]},$$

the finite surreals and the surreals bounded by an ordinary multiple of ω . All three images are cofinal in $\mathbf{No}$ and none of their positive parts is coinitial.

Proof. The images follow from [lemma 27.1](#). A positive purely infinite surreal exceeds every real, hence every ordinary multiple of a finite surreal; so $C_1 \subseteq G$. An infinite c has leading exponent $a > 0$, and $\omega^{a/2}$ is a positive image element below $|c|$; so $c \notin G$. For μ_{12} : a positive

image element has leading term $r\omega^a$ with $r > 0$, $a > 1$, so it exceeds $n\omega$ for every n , and $C_\omega \subseteq \mathbf{G}$. If $c \notin C_\omega$, its leading exponent a exceeds 1, and $\omega^{(1+a)/2}$ is an image element below $|c|$. Arbitrarily large exponents give cofinality; no positive image element is at most 1. $\square$

The purely infinite surreals here form a subgroup of the *value group*; it is the same set as the ideal Π of $\mathbf{No}$, but it is used additively and is not a convex subgroup [11, 13].

27.2 One map, three instances

Theorem 27.3 (Proper embeddings moving a real [11, 12, 13]). *Let b be a transcendental real, d a derivation of $\mathbb{R}$ with $d(b) = 1$, and $\mu \in \{\mu_{11}, \mu_{12}, \mu_{13}\}$. Then*

$$J_{\mu,d}\left(\sum_{\gamma} r_{\gamma} t^{\gamma}\right) = \sum_{\gamma} \sum_{n \geq 0} \frac{d^n(r_{\gamma})}{n!} t^{\iota_{\mu}(\gamma)+n} \quad (27.3)$$

is $J_{\mu,\rho_{d,1}} = \mathcal{T}_{d,1} \circ M_{1,\iota_{\mu}}$. It is a proper, strongly additive, exact-monomial ordered field embedding $\mathbf{No} \hookrightarrow \mathbf{No}$ with

$$v(J_{\mu,d}x) = \iota_{\mu}(v(x)), \quad \text{ct} \circ J_{\mu,d} = \text{ct}, \quad J_{\mu,d}^{-1}(\mathbf{Oz}) = \mathbf{Oz}, \quad J_{\mu,d}(b) = b + \omega^{-1},$$

and it fixes every real algebraic number. Moreover $J_{\mu,d}(\omega) = \omega^{\omega}$ for $\mu = \mu_{11}, \mu_{13}$ and $J_{\mu_{12},d}(\omega) = \omega^{\omega^2}$. Its restriction is a proper ring embedding $\mathbf{Oz} \hookrightarrow \mathbf{Oz}$; it is not $\mathbb{R}$ -linear, and no automorphism of $(\mathbf{No}, \mathbf{Oz})$ agrees with it on b .

Proof. $1 \in \mathbf{G}_{\iota_{\mu}}$ by lemma 27.2, so theorems 26.3 and 26.6 give the embedding, strongness, valuation, constant term and reflection. $\rho_{d,1}(b) = b + t$ and $t = \omega^{-1}$; d kills the real algebraic numbers. $\iota_{\mu}(1) = \omega^{\mu(0)}$ with $\mu_{11}(0) = \mu_{13}(0) = 1$ and $\mu_{12}(0) = 2$, and $J(\omega) = J(t^{-1}) = t^{-\iota_{\mu}(1)}$. The value $1 = v(t)$ is not in $\iota_{\mu}(\mathbf{No})$, so $t \notin J(\mathbf{No})$; if the restriction to $\mathbf{Oz}$ were onto, extension to fractions (odg:thm:fractions) would make J onto. Every $\mathbf{Oz}$ -preserving automorphism fixes b (corollary 5.2), and the restriction to $\mathbf{Oz}$ determines the fraction-field map. $\square$

So the rigidity of automorphisms does not pass to embeddings: $(\mathbf{No}, +, \cdot, \mathbf{Oz})$ and $\mathbf{Oz}$ have proper self-embeddings, and an ordered-field embedding of $\mathbf{No}$ preserving and reflecting $\mathbf{Oz}$ need not preserve $\mathbb{R}$ setwise, even when it is strongly additive and preserves constant terms exactly [12]. 11 proves the theorem for μ_{11} through its classification; 12 (for μ_{12}) and 13 (for μ_{13}) give the same block argument directly. The three maps differ only in the order isomorphism: 11's and 13's have the same value image and send $\omega \mapsto \omega^{\omega}$. For μ_{12} , $\mathbf{G} = C_{\omega}$, so even $\delta = \omega$ would be admissible (lemma 27.2). For example

$$J_{\mu,d}(bt^g) = bt^{\iota_{\mu}(g)} + t^{\iota_{\mu}(g)+1} \quad (g < 0), \quad J_{\mu_{11},d}(b\omega) = b\omega^{\omega} + \omega^{\omega-1} \in \mathbf{Oz},$$

both terms purely infinite although $J(b)$ is not real; and $J(b^2) = b^2 + 2bt + t^2$, $J(b^{-1}) = \sum_n (-1)^n b^{-n-1} t^n$, $J(t^g) = t^{\iota_{\mu}(g)}$ [11, 12]. The ordinary Taylor section alone is not a map on arbitrary normal forms, since supports could meet its integer shifts; the proper exponent embedding separates the scales [12, 13].

Corollary 27.4 (Image, reals and a missing test [12]; for all three [merge]). *For each $J = J_{\mu,d}$: $J(\mathbf{No}) \cap \mathbb{R} = \ker d$, and $\text{ct}(tJ(x)) = 0$ for every $x \in \mathbf{No}$, although $t \neq 0$. So the image does not supply all constant-term tests.*

Proof. If $J(x)$ is constant, every block at $g \neq 0$ vanishes, so $x = r$ is real, and $\rho_{d,1}(r)$ is constant exactly when $d(r) = 0$ (see also proposition 29.2). An exponent $\iota_{\mu}(g) + n = -1$ would make $\iota_{\mu}(g)$ a negative integer; but $\iota_{\mu}(g)$ is 0 or of absolute value above every integer (lemma 27.2). So no image has a term at -1 . $\square$

12 proves the corollary for μ_{12} ; the merge notes that its proof uses only that nonzero values of ι_μ are infinite, true for all three. The annihilator is the missing detector in the isomorphism proof: tests from the image cannot distinguish t from 0, whereas the ambient test t^{-1} does [12].

Corollary 27.5 (Failure of pair homogeneity [13]). *The ordered embedding $\rho_{d,1} : \mathbb{R} \rightarrow \mathbf{No}$ is an embedding of the induced pairs $(\mathbb{R}, \mathbb{Z}) \rightarrow (\mathbf{No}, \mathbf{Oz})$ that extends to the proper pair embedding $J_{\mu,d}$ but to no automorphism of $(\mathbf{No}, \mathbf{Oz})$. So $(\mathbf{No}, \mathbf{Oz})$ is not strongly homogeneous for embeddings of set-sized induced substructures.*

Proof. Restrict the reflection property of [theorem 27.3](#) to $\mathbb{R}$; every $\mathbf{Oz}$ -preserving automorphism fixes b , whereas $\rho_{d,1}(b) = b + t$. $\square$

“Strongly homogeneous” means extension of isomorphisms between set-sized induced substructures, not extension of elementary partial maps in a saturated expansion. The pure ordered field $\mathbf{No}$ has this extension property for set-sized ordered subfields [12, Remark 2.10], so $\rho_{d,1}$ does extend to an ordered-field automorphism of $\mathbf{No}$, and every such extension fails to preserve $\mathbf{Oz}$; the obstruction lies in the integer part [13]. The map also satisfies $\text{ct} \circ J = \text{ct}$ but not $J \circ \text{ct} = \text{ct}$ nor $J \circ \text{ct} = \text{ct} \circ J$: for b these compare b with $b + t$. It violates no hypothesis of [theorem 22.1](#), since it is not onto [13]. It does not commute with the omega-map: $J(\omega^1) = \omega^\omega$ for μ_{11} whereas $\omega^{J(1)} = \omega$; no question of [12] on the omega-map is addressed [11].

28 Closed images and the two ends of the value group

Use the valuation topology with basic neighbourhoods $\mathcal{V}_\lambda = \{x : v(x) > \lambda\}$. On a real Hahn field with nonzero value group it is the order topology; for $\mathbf{No}$ it is the fine topology. No countable basis is assumed, and for proper classes the topological statements are local conditions on balls; no object containing all open subclasses is formed [11].

Theorem 28.1 (Closed images [11]). *Every image $F = J_{\tau,\rho}(K_\Gamma)$ of [theorem 26.3](#) is closed in K_Δ ; if J is proper, F is nowhere dense.*

Proof. Put $G = G_\tau$ and $D = \tau(\Gamma) + G$; image series are supported in D . If x has a nonzero coefficient at $\lambda \notin D$, then $v(x - y) \leq \lambda$ for all $y \in F$. If $\text{supp } x \subseteq D$, split $x = \sum_{h \in S} t^h x_h$ with $x_h \in \mathbb{R}((t^G))$ by the unique decomposition; the projection $D \rightarrow \tau(\Gamma)$ is nondecreasing, so S is well ordered. Put $r_h = \text{ct}(x_h)$. If $x_h = \rho(r_h)$ for all h , then $x = J_{\tau,\rho}(\sum_h r_h t^{\tau^{-1}(h)}) \in F$. Otherwise pick h and $c \in G$ where $x_h - \rho(r_h)$ has a nonzero coefficient, and put $L = \max\{h, h + c\}$. The block of any $y \in F$ at h is $\rho(s)$ for one $s \in \mathbb{R}$ (possibly 0); if $s \neq r_h$ then $x - y$ has a nonzero coefficient at h , and if $s = r_h$ at $h + c$. So $x + \mathcal{V}_L$ misses F , and F is closed. A subfield containing a ball $\mathcal{V}_L$ contains every $z \neq 0$, as $z = (zt^\gamma)/t^\gamma$ with both in $\mathcal{V}_L$ for large γ ; so a proper subfield has empty interior, and a closed one is nowhere dense. $\square$

Lemma 28.2 (Induced value embedding [11]). *Every field embedding $f : \mathbf{No} \hookrightarrow \mathbf{No}$ is ordered and induces an ordered additive embedding τ_f of the natural value group with $v(f(x)) = \tau_f(v(x))$. No monomial or strongness hypothesis is needed.*

Proof. Positivity is being a nonzero square. Archimedean equivalence and strict domination are expressed by inequalities with ordinary integers, which f preserves and reflects; multiplicativity gives additivity. $\square$

Theorem 28.3 (Continuity or external discreteness [11]). *Let $f : \mathbf{No} \hookrightarrow \mathbf{No}$ be any field embedding. Exactly one of the following holds.*

- (i) $\tau_f(\mathbf{No})$ is cofinal in $\mathbf{No}$. Then f is continuous, a homeomorphism onto its image, and the image is nondiscrete.
- (ii) $\tau_f(\mathbf{No})$ is not cofinal. Then there is L with $v(y - z) < L$ for all distinct $y, z \in f(\mathbf{No})$; the image is closed and uniformly discrete, and f is nowhere continuous.

Proof. (i) Given L , choose γ with $\tau_f(\gamma) > L$; then $f(\mathcal{V}_\gamma) = f(\mathbf{No}) \cap \mathcal{V}_{\tau_f(\gamma)} \subseteq \mathcal{V}_L$. (ii) Take L above $\tau_f(\mathbf{No})$: nonzero differences in the image have value below L , so $f^{-1}(\mathcal{V}_L) = \{0\}$, not open, and additivity gives discontinuity everywhere. A ball of radius L contains at most one image point, which gives closedness and uniform discreteness. $\square$

There is no contradiction in an order-isomorphic copy of the nondiscrete $\mathbf{No}$ being discrete: the intrinsic order topology of a non-cofinal ordered subfield can be coarser than its subspace topology; the inverse from the image stays continuous [11].

Theorem 28.4 (The two ends are independent [11]). *For the order embeddings*

$$\mu^{++}(a) = \begin{cases} a, & a \leq 0, \\ a + 1, & a > 0, \end{cases} \quad \mu^{+-} = \mu_{11},$$

$$\mu^{-+}(a) = -\mu_{11}(-a), \quad \mu^{--}(a) = \frac{\mu_{11}(a)}{1 + \mu_{11}(a)},$$

with ranges $(-\infty, 0] \cup (1, \infty)$, $(0, \infty)$, $(-\infty, 0)$ and $(0, 1)$, the value group $\iota_\mu(\mathbf{No})$ is respectively cofinal and coinital, cofinal and not coinital, coinital and not cofinal, neither; here “coinital” refers to the positive part. In each case there is a proper strong exact-monomial self-embedding preserving and reflecting $\mathbf{Oz}$: M_{1, ι_μ} in the coinital cases, where coefficient motion is impossible, and $\mathcal{T}_{d,1} \circ M_{1, \iota_\mu}$, moving b to $b + \omega^{-1}$, in the others. The embeddings are continuous with closed nondiscrete image in the cofinal cases and nowhere continuous with discrete image in the others; all four images are nowhere dense.

Proof. By lemma 27.1 the value image is the class of normal forms supported in the range. The range of μ^{++} has arbitrarily high and low levels and misses $(0, 1]$, so the image is proper, cofinal and coinital. μ^{+-} is lemma 27.2. For μ^{-+} all values are infinitesimal, so the group is bounded by 1, but monomials at arbitrarily negative levels make it coinital. For μ^{--} all nonzero values are purely infinite and below ω in absolute value, so the group is neither; its gap is C_1 , since a finite value is dominated by every positive image value and an infinite value with leading level $a > 0$ dominates ω^c for some $0 < c < \min\{a, 1\}$. Apply corollary 26.7 and theorems 28.1 and 28.3. $\square$

So cofinality must not replace coinitality in the rigidity criterion: $J_{\mu_{11}, d}$ is continuous with cofinal values and still moves $\mathbb{R}$ [11].

29 Constants, elementarity, conjugacy and parameters

29.1 Elementary reduct, nonelementary expansion

$\mathbb{R}$ is parameter-free definable in $(K_\Gamma, \mathcal{R}_\Gamma)$ by the existential test, the multiplier condition and its units (`odg: def: thm: ideal`, `odg: def: thm: multiplier`; the internal form is `odg: def: cor: internal`); 11 reproves this with explicit formulas D , M , Const , and derives the rigidity of automorphisms (corollary 5.2).

Theorem 29.1 (Logical separation [11]; [12] for its example). *Every field embedding $\mathbf{No} \hookrightarrow \mathbf{No}$ is elementary as an embedding of ordered fields. An embedding of [theorem 26.3](#) (with $\Gamma = \Delta = \mathbf{No}$) that moves some real is not elementary as an embedding of $(\mathbf{No}, \mathbf{Oz})$, although it preserves and reflects $\mathbf{Oz}$, and its restriction to $\mathbf{Oz}$ is not elementary as an embedding of rings.*

Proof. The first assertion is quantifier elimination for real closed ordered fields [16], read formula by formula for the class structure. If $\Phi(b) \notin \mathbb{R}$, the formula $\text{Const}(x)$ defining $\mathbb{R}$ holds of b and fails of $\Phi(b)$. For $\Phi(b) = b + t$ the new target witness is visible: $\omega \in \Pi$ but $(b + t)\omega = b\omega + 1 \notin \Pi$; the universal multiplier condition quantifies over more elements in the target. For the ring: $\text{Frac } \mathbf{Oz} = \mathbf{No}$ (`odg:thm:fractions`), the fraction field is uniformly interpretable in the ring by pairs modulo $ad = bc$, with the ring as the class of fractions $(z, 1)$, so an elementary ring embedding would induce an elementary embedding of the interpreted pair; represent b as $(b\omega)/\omega$. $\square$

The first assertion is classical. The new point is that a strong, exact-monomial, predicate-reflecting embedding can lose elementarity in the omnific expansion; coefficient fixing is not proved sufficient for elementarity ([question 32.6](#)) [11]. 12 records the same failure: the multiplier definition of $\mathcal{A}_k$ quantifies over all of Π , which an embedding need not preserve. This sharpens the Diophantine report's caution after `odg:def:thm:autreal` that its argument does not apply unchanged to endomorphisms: there are proper ring embeddings $\mathbf{Oz} \hookrightarrow \mathbf{Oz}$ whose fraction-field extensions move a real.

Proposition 29.2 (Ordinary overlap [11]; [12] for Taylor sections). *For $F = J_{\tau, \rho}(K_\Gamma)$: $F \cap \mathbb{R}((t^{\mathbb{G}_\tau})) = \rho(\mathbb{R})$ and $F \cap \mathbb{R} = \{r : \rho(r) = r\}$, a real closed subfield of $\mathbb{R}$; for $\rho = \rho_{d, \delta}$ it is $\ker d$.*

Proof. A nonzero block at $h \neq 0$ lies in $h + \mathbb{G}_\tau$, disjoint from $\mathbb{G}_\tau$, so an image in $\mathbb{R}((t^{\mathbb{G}_\tau}))$ is its zero block $\rho(r)$; if that is a real constant, its standard part gives r . If r is algebraic over the fixed field k , then $\rho(r)$ is a root of the same polynomial, all real roots lie in $\mathbb{R}$, and the residue singles out r ; so k is relatively algebraically closed in the real closed $\mathbb{R}$, hence real closed. $\rho_{d, \delta}(r) = r$ exactly when $d(r) = 0$. $\square$

29.2 Continuum many copies separated by the omnific integers

Lemma 29.3 (One-sided infinitesimal neighbourhoods [11]). *Let $F = J_{\tau, \rho}(K_\Gamma)$ and $\rho(r) \neq r$. Every $y \in F$ with $\text{st}(y) = r$ has $\text{sgn}(y - r) = \text{sgn}(\rho(r) - r)$, and $\rho(r)$ is such a y . For $\rho = \rho_{d, \delta}$ with $d(r) \neq 0$ the sign is that of $d(r)$.*

Proof. $y = J(x)$ with $x = r + \varepsilon$, $v(\varepsilon) > 0$ (by the valuation and constant-term formulas). The nonzero $\rho(r) - r$ has value in $\mathbb{G}_\tau^{>0}$, while $v(J\varepsilon) \in \tau(\Gamma)^{>0}$ is larger; so $y - r = (\rho(r) - r) + J(\varepsilon)$ has the leading term of $\rho(r) - r$, which is $d(r)t^\delta$ for a Taylor section. $\square$

The sign is invariant under every field automorphism of $\mathbf{No}$ fixing $\mathbb{R}$ pointwise, in particular under every $\mathbf{Oz}$ -preserving one ([corollary 5.2](#)), since these preserve order and standard part.

Theorem 29.4 (One field-conjugacy class, continuum many pair classes [11]). *There are proper strong exact-monomial embeddings $J_a : \mathbf{No} \hookrightarrow \mathbf{No}$, $a \in \mathbb{R}$, with images F_a , such that*

- (i) $J_a^{-1}(\mathbf{Oz}) = \mathbf{Oz}$;
- (ii) *all F_a have the cofinal value image $\iota_{\mu_{11}}(\mathbf{No})$ and are closed, nondiscrete and nowhere dense;*

- (iii) all F_a are conjugate under strong valuation-fixing field automorphisms of $\mathbf{No}$;
- (iv) for $a \neq a'$ no field automorphism of $\mathbf{No}$ fixing $\mathbb{R}$ pointwise, in particular none preserving $\mathbf{Oz}$, maps F_a onto $F_{a'}$.

So the pairs $(F_a, F_a \cap \mathbf{Oz})$ are all isomorphic to $(\mathbf{No}, \mathbf{Oz})$ but lie in pairwise inequivalent positions.

Proof. Choose distinct b_1, b_2 in a transcendence basis of $\mathbb{R}/\mathbb{Q}$ and let d_a be the derivation with $d_a(b_1) = 1$, $d_a(b_2) = a$ and $d_a = 0$ on the other basis elements; no independence of named constants is assumed. Put $J_a = \mathcal{T}_{d_a,1} \circ M_{1,\iota_{\mu_{11}}}$. (i) and (ii) are [theorems 27.3, 28.1](#) and [28.3](#). For (iii), $\mathcal{T}_{d_{a'},1} \mathcal{T}_{d_a,1}^{-1}$ maps F_a onto $F_{a'}$; the derivations need not commute, and only the individual inverses are used. For (iv) let $a < q < a'$ with $q \in \mathbb{Q}$ and $r = b_2 - qb_1$: $d_a(r) = a - q < 0 < a' - q = d_{a'}(r)$, so by [lemma 29.3](#) the points of F_a with standard part r lie below r and those of $F_{a'}$ above, and both sets are nonempty. Finally $F_a \cap \mathbf{Oz} = J_a(\mathbf{Oz})$ by reflection. $\square$

This is a lower bound of continuum on the pair-conjugacy types inside one field-conjugacy class, not a classification or an upper bound; the images have identical value data, so no classification can use the value image and its cuts alone [11]. It partly answers 13's question on the conjugacy of its drift images ([question 32.8](#)).

29.3 Proper copies fixing prescribed parameters

Lemma 29.5 (Two-level support control [11]). *For a set $A \subseteq \mathbf{No}$ let B be the union of the inner supports of the outer exponents of the members of A , a set. Every order embedding μ fixing B pointwise has $M_{1,\iota_\mu}(a) = a$ for all $a \in A$.*

Proof. ι_μ fixes every outer exponent of every member of A ; this is the inner/outer distinction of [lemma 7.1](#). $\square$

Theorem 29.6 (Parameter-fixed discrete and nondiscrete copies [11]). *For every set $A \subseteq \mathbf{No}$ there are proper strong exact-monomial embeddings $f_A, g_A : \mathbf{No} \hookrightarrow \mathbf{No}$ fixing $A \cup \mathbb{R}$ pointwise with $f_A^{-1}(\mathbf{Oz}) = g_A^{-1}(\mathbf{Oz}) = \mathbf{Oz}$. The image of f_A is closed and uniformly discrete and f_A is nowhere continuous; the image of g_A is closed, nondiscrete and nowhere dense and g_A is a homeomorphism onto it; no field automorphism of $\mathbf{No}$ maps one image onto the other. There is an explicit $\Omega > 1$ with $\Omega^{-1} < |y| < \Omega$ for every $0 \neq y \in f_A(\mathbf{No})$.*

Proof. Take B as in [lemma 29.5](#) and u above B . The map $\mu^{\text{cmp}}(x) = x$ for $x \leq u$, $u + \frac{x-u}{1+x-u}$ for $x > u$, is an order isomorphism onto $(-\infty, u+1)$ fixing B , so every value h of $\iota_{\mu^{\text{cmp}}}$ has $|h| < \beta = \omega^{u+1}$; put $f_A = M_{1,\iota_{\mu^{\text{cmp}}}}$. The map $\mu^{\text{jmp}}(x) = x$ for $x \leq u$, $x+1$ for $x > u$, fixes B , omits $(u, u+1]$ and has cofinal range; put $g_A = M_{1,\iota_{\mu^{\text{jmp}}}}$. [Theorems 26.3, 28.1](#) and [28.3](#) and [lemma 29.5](#) give the properties. A field automorphism of $\mathbf{No}$ is a homeomorphism (its value map is onto; [lemma 28.2](#) and [theorem 28.3](#)) and cannot exchange a discrete and a nondiscrete subspace. For $\Omega = \omega^\beta$: a nonzero $y \in f_A(\mathbf{No})$ has natural value h with $|h| < \beta$, so its leading monomial has exponent below β and $|y| < \Omega$; apply this to $1/y$. $\square$

The annulus bounds are external: Ω is not in the image, which is internally isomorphic to $\mathbf{No}$ and not Archimedean [11].

Extension elsewhere (batch 31). The independent-copies report `docs/surreal/independent-surreal-copies` extends the fixing clause, with $\rho = \text{id}$, from two maps to an $\mathbf{On}$ -indexed family with a prescribed intersection (its `isc:cor:parameters`, from `isc:main:copies`).

Let H be the $\mathbb{Q}$ -span of the supports of the members of A . There are proper strong exact-monomial embeddings Φ_α ($\alpha \in \mathbf{On}$) preserving and reflecting $\mathbf{Oz}$ and fixing the full Hahn field $\mathbb{R}((t^H)) \supseteq A \cup \mathbb{R}$ pointwise; any two images meet exactly in $\mathbb{R}((t^H))$, and every finite subfamily is jointly linearly disjoint over it. That field is the full support closure of A , not the field generated by A or its real closure. In the terms of [theorem 26.3](#) these maps are $J_{h_\alpha, \text{id}}$ for exponent embeddings h_α fixing H with ranges independent over H (`isc:thm:group`), obtained by class bookkeeping. For $H = 0$ that report also gives explicit maps (`isc:thm:explicit`): $M_{1, \iota_{\sigma_\alpha}}$ with σ_α an order isomorphism of $\mathbf{No}$ onto the interval $(3\alpha, 3\alpha + 1)$. These intervals are bounded and pairwise disjoint, and the bottom gap is nonzero. That report makes no topological assertion. [merge]: the exponent map of each explicit copy has bounded range, so by [theorem 28.3\(ii\)](#) its image is closed and uniformly discrete, as for f_A ; for the general family the alternative is not decided.

Theorem 29.7 (Relative coefficient motion [11]). *Let $A \subseteq \mathbf{No}$ be a set and k_A the field generated over $\mathbb{Q}$ by the real coefficients of the members of A . If $b \in \mathbb{R}$ is transcendental over k_A , there is a proper strong exact-monomial embedding fixing A pointwise, preserving and reflecting $\mathbf{Oz}$, with cofinal value image and $b \mapsto b + t^\delta$ for some $\delta > 0$.*

Proof. Choose d zero on k_A with $d(b) = 1$, through a transcendence basis of $\mathbb{R}/k_A$ containing b . With B as in [lemma 29.5](#), choose v below B and $c < v$, and let $\mu^{\text{rel}}(x) = v - (v - c) \frac{v-x}{1+v-x}$ for $x < v$ and x for $x \geq v$, an order isomorphism onto (c, ∞) fixing B . Positive values of $\iota_{\mu^{\text{rel}}}$ have leading exponent above c , so $\delta = \omega^c$ lies in the gap. Put $\mathcal{T}_{d,\delta} \circ M_{1, \iota_{\mu^{\text{rel}}}}$; it fixes A since the exponent lift fixes each member and d kills its coefficients. Apply [theorem 26.6](#). $\square$

An ordered embedding fixing a real field fixes the reals algebraic over it, so one cannot fix every real and move another; the transcendence hypothesis is needed [11].

30 Surcomplex and Gaussian versions

Identify $\mathbf{No}(i) = \mathbb{C}((t^{\mathbf{No}}))$ coefficientwise with conjugation $\overline{x + iy} = x - iy$; the union of the real and imaginary supports is well ordered, and a family of surcomplex series is summable if and only if its real and imaginary families are. The general Gaussian problem is kept separate from two restricted ones: with conjugation named, and with the natural valuation assumed preserved. They must not be merged into an unconditional classification [12, 13].

Theorem 30.1 (Conjugation-compatible automorphisms [12, 13, 14, 15]). *Every field automorphism F of $\mathbf{No}(i)$ with $F(\mathbf{Oz}[i]) = \mathbf{Oz}[i]$ that commutes with conjugation has the form $F(x + iy) = \sigma(x) + \varepsilon i \sigma(y)$ with σ an $\mathbf{Oz}$ -preserving automorphism of $\mathbf{No}$ and $\varepsilon = \pm 1$, and F and F^{-1} are strongly additive. Conversely every such map has these properties, so this stabilizer is $\text{Aut}(\mathbf{Oz}) \times C_2$; with $F(i) = i$ imposed, only $\varepsilon = 1$ occurs and F is strongly $\mathbb{C}$ -linear.*

Proof. Commuting with conjugation preserves its fixed field $\mathbf{No}$, and $\mathbf{Oz}[i] \cap \mathbf{No} = \mathbf{Oz}$. $F(i)$ is a root of $X^2 + 1$. [Theorem 23.2](#) makes σ strong, and summability is coordinatewise. Real lifts commute with conjugation, which makes the product direct. $\square$

Theorem 30.2 (Valuation-compatible Gaussian automorphisms [12, 13, 14, 15]). *Let F be a field automorphism of $\mathbf{No}(i)$ with $F(\mathbf{Oz}[i]) = \mathbf{Oz}[i]$ that maps the Hahn valuation ring onto itself (equivalently $v(Fx) = \tau(v(x))$ for an ordered automorphism τ of $\mathbf{No}$; values need not be fixed). Then $F(\mathbb{C}) = \mathbb{C}$; with $\alpha = F|_{\mathbb{C}}$, F is α -semilinear, $\text{ct}(Fx) = \alpha(\text{ct } x)$, and F and F^{-1} are strongly additive. The same holds for the fraction-field extension of a ring*

automorphism of $\mathbf{Oz}[i]$ whenever that extension preserves the valuation, and for valuation-compatible isomorphisms between full complex Hahn fields with nonzero exponent groups preserving their Gaussian integer parts [13].

Proof. $\Pi_{\mathbb{C}} = \bigcap_n n\mathbf{Oz}[i]$ (lemma 2.3); its multiplier ring is $\mathbb{C} \oplus \Pi_{\mathbb{C}}$ with units $\mathbb{C}^{\times}$ (odg: def: thm: multiplier), so $F(\mathbb{C}) = \mathbb{C}$. Valuation compatibility preserves $\mathfrak{m}_{\mathbb{C}}$, and $\mathbf{No}(i) = \Pi_{\mathbb{C}} \oplus \mathbb{C} \oplus \mathfrak{m}_{\mathbb{C}}$ gives $\text{ct}(Fx) = \alpha(\text{ct } x)$. Apply theorem 22.1. The fraction-field statement uses $\text{Frac } \mathbf{Oz}[i] = \mathbf{No}(i)$ (odg: thm: fractions); the isomorphism statement uses only the ambient fields. $\square$

α may be a wild automorphism of $\mathbb{C}$: coefficientwise application of any automorphism of $\mathbb{C}$ is strong and preserves $\mathbf{Oz}[i]$, since it fixes $\mathbb{Q}$ and sends i to $\pm i$ [13]. This is consistent with the wild lift of proposition 10.3 and with odg: def: prop: coeffaut, which are strong. Neither theorem shows that every automorphism of $\mathbf{Oz}[i]$ has a valuation-compatible extension; conjugation is not assumed intrinsic, and maps moving the real axis are already in the collection (theorem 10.2, odg: def: lem: twist). Since $\text{Frac } \mathbf{Oz}[i] = \mathbf{No}(i)$, the open issue is compatibility, not existence of the extension [12, 13] (question 32.1).

Theorem 30.3 (Conjugation-compatible embeddings [11]). *The strong field embeddings $\Psi : K_{\Gamma}^{\mathbb{C}} \hookrightarrow K_{\Delta}^{\mathbb{C}}$ that commute with conjugation, satisfy $\Psi(t^{\gamma}) = t^{\tau(\gamma)}$ and preserve $\mathcal{R}_{\Gamma}^{\mathbb{C}}$ are exactly $\Psi(x + iy) = J_{\tau, \rho}(x) + \varepsilon i J_{\tau, \rho}(y)$, $\varepsilon = \pm 1$, with (τ, ρ) as in theorem 26.3; each satisfies $\Psi^{-1}(\mathcal{R}_{\Delta}^{\mathbb{C}}) = \mathcal{R}_{\Gamma}^{\mathbb{C}}$, and $\Psi(i) = i$ forces $\varepsilon = 1$.*

Proof. Commuting with conjugation maps K_{Γ} into K_{Δ} , and $\mathcal{R}_{\Gamma}^{\mathbb{C}} \cap K_{\Gamma} = \mathcal{R}_{\Gamma}$, so the restriction is classified; $\Psi(i) = \pm i$. The converse is coordinatewise, and $\mathcal{R}^{\mathbb{C}}$ -membership is membership of both coordinates. $\square$

12's and 13's surcomplex embeddings $x + iy \mapsto J(x) + iJ(y)$ are the case $\varepsilon = 1$ of theorem 27.3; they are proper and strong, commute with conjugation, reflect $\mathbf{Oz}[i]$, move b to $b + t$, and their image meets $\mathbb{C}$ in $\ker d + i \ker d$ [12, 13].

Proposition 30.4 (Complex constants and nonelementarity [11]). *$\mathbb{C}$ is parameter-free definable in $(K_{\Gamma}^{\mathbb{C}}, \mathcal{R}_{\Gamma}^{\mathbb{C}})$. An embedding of theorem 30.3 with nontrivial ρ is elementary as an embedding of algebraically closed fields, but not in the Gaussian expansion; for $\Gamma = \Delta = \mathbf{No}$ its restriction to $\mathbf{Oz}[i]$ is not elementary as a ring embedding.*

Proof. For $z \in \mathcal{R}_{\Gamma}^{\mathbb{C}}$, $z \in \Pi_{\Gamma}^{\mathbb{C}}$ if and only if $z^2 = 2w^2$ for some $w \in \mathcal{R}_{\Gamma}^{\mathbb{C}}$, since $\sqrt{2} \notin \mathbb{Q}(i)$; the multiplier ring and its units define $\mathbb{C}$ as in the real case (odg: def: cor: internal). A moved real b has $\rho(b) \notin \mathbb{C}$. Pure-field elementarity is quantifier elimination for algebraically closed fields [16]; for the ring use $\text{Frac } \mathbf{Oz}[i] = \mathbf{No}(i)$ and the interpreted fraction field as in theorem 29.1. $\square$

Corollary 30.5 (Transferred copies [11]). *The explicit coefficient-moving embedding, the four topological cases and the parameter-fixed copies extend to $\mathbf{No}(i)$, preserving and reflecting $\mathbf{Oz}[i]$; parameters in a set $A \subseteq \mathbf{No}(i)$ are handled through their real and imaginary parts. The family of theorem 29.4 gives field-conjugate surcomplex images that are pairwise nonconjugate under automorphisms preserving both conjugation and $\mathbf{Oz}[i]$.*

Proof. Use $\varepsilon = 1$; the valuation of $x + iy$ is the minimum of the coordinate valuations, so the topology is the product topology. An automorphism preserving conjugation and $\mathbf{Oz}[i]$ restricts to an $\mathbf{Oz}$ -preserving automorphism of $\mathbf{No}$, and each extended image meets $\mathbf{No}$ in the real image; apply theorem 29.4. $\square$

No nonconjugacy is claimed for automorphisms preserving $\mathbf{Oz}[i]$ but not conjugation, which need not fix the real axis pointwise, and non-conjugation-compatible surcomplex embeddings are not classified [11].

31 Boundaries of the method

Proposition 31.1 (Regular cardinal support bounds [12]). *Let κ be a regular uncountable cardinal, Γ a set, and $K_{<\kappa}$ the field of series in $k((t^\Gamma))$ with support of size $< \kappa$, where summability also requires total support of size $< \kappa$. A family in $K_{<\kappa}$ is summable there if and only if fewer than κ members are nonzero and for every $y \in K_{<\kappa}$ only finitely many $\text{ct}(x_j y)$ are nonzero. Consequently a constant-term-covariant field isomorphism between two such fields is strong for this restricted summation.*

Proof. A summable family has fewer than κ nonzero members, since each contributes at an exponent of a total support of size $< \kappa$ and each exponent receives finitely many; pairing finiteness is [theorem 21.4](#). Conversely, regularity bounds the union of fewer than κ supports of size $< \kappa$; monomials test pointwise finiteness, and the countable witness of [lemma 21.2](#) lies in $K_{<\kappa}$ because κ is uncountable. An isomorphism preserves the number of nonzero members, and the proof of [theorem 22.1](#) carries over. $\square$

The cardinal condition cannot be dropped: a strictly increasing $(g_\alpha)_{\alpha < \kappa}$ in Γ gives (t^{g_α}) with finite pairings against every $y \in K_{<\kappa}$ ([lemma 21.1](#)) but total support of size κ [12]. For singular κ the union argument fails, and no extension is claimed; the proposition identifies no birthday cutoff with a support-bounded Hahn field [12]. 13 records the regular case only inside its singular-cardinal question ([question 32.12](#)).

Example 31.2 (Small integer parts [13, 14]). The proof of $\text{Frac } \mathbf{Oz} = \mathbf{No}$ uses an upper bound in $\mathbf{No}$ for every set of exponents, and does not transfer to set-sized groups. For $\Gamma = \mathbb{Z}$ and $u = t^{-1}$, $\mathcal{R}_{\mathbb{Z}} = \mathbb{Z} \oplus u\mathbb{R}[u]$ and $\text{Frac } \mathcal{R}_{\mathbb{Z}} = \mathbb{R}(u) \neq \mathbb{R}((t))$: $ru/u = r$ gives $\mathbb{R}(u)$, and $f = \sum_{n \geq 1} t^{n!}$ is not rational, since a rational power series has coefficients obeying an eventual linear recurrence, and arbitrarily long zero blocks followed by a 1 contradict one.

This is the phenomenon of [proposition 18.1](#) and of [14, Proposition 2.4.5]; it is why [corollary 23.4](#) assumes an ambient isomorphism [11, 12, 13].

Remark 31.3 (Distinctions kept [12, 13]). The detector must lie in the field tested: automatic for full Hahn fields and for the countable witness in $K_{<\kappa}$, not for arbitrary subfields (a sufficient condition is 14's countable detector property, [proposition 35.2](#), and [example 35.3](#) is a subfield where detection fails). Strongness is not $\mathbb{R}$ -linearity: $J_{\mu,d}$ is strong but not $\mathbb{R}$ -linear, so the adjoint theorem, which is for k -linear maps, does not apply to it by treating its coefficient lift as scalar multiplication. The retraction ret_ϕ is linear, never a field retraction for a proper extension. The adjoint theorem needs a set of exponents; automatic strongness of isomorphisms does not. No result of this part concerns the Gonshor exponential, logarithm, the Berarducci–Mantova derivation, the omega-map, birthdays or simplicity; the square roots in (27.2) are ordered-field operations reindexing an additive group. No proper-class Zorn lemma and no basis of the class $\mathbf{No}$ is used; the only bases chosen are a transcendence basis and a Hamel basis of the set $\mathbb{R}$ over $\mathbb{Q}$ [13].

32 Status of the questions, and questions of this part

32.1 Answered and re-scoped questions

- [Question 11.1](#) is *answered*. Every ring automorphism of $\mathbf{Oz}$ extends to a strong automorphism of $\mathbf{No}$ ([theorem 23.2](#)), so preservation of $\mathbf{Oz}$ forces strongness and there is no nonstrong $\mathbf{Oz}$ -preserving example; 09’s characterization of the nonstrong pair automorphisms is vacuous, and 10’s question whether every $\mathbf{Oz}$ -preserving 1-automorphism is strong has the answer yes. The analogous question for $\mathbf{Oz}[i]$ has a positive answer under conjugation or valuation compatibility and is open otherwise ([question 32.1](#)).
- `osq:q:invisible` of `docs/surreal/set-sized-quotients-of-omnific-integers`: its second clause is *answered* by [corollary 24.4](#); the nonstrong case left open in [section 11.1](#) does not exist. Its first clause, on class quotient domains, is untouched. The quotient report is not edited by this merge; since batch 31 it records the question as partly answered, with this corollary.
- [Question 11.2](#): unchanged. 12 and 13 ask it again, in the forms “for which support-controlled families of logarithmic characters does $a_\delta(C_\delta) = 0$ imply two-sided substitution admissibility?” and “give necessary and sufficient support conditions on additive $\gamma \mapsto L_\gamma \in \mathfrak{m}$ for $t^\gamma \mapsto t^\gamma \exp L_\gamma$ to extend to a strong automorphism, assuming $a_\delta(C_\delta) = 0$ ”, suggesting common-shift and finite-rank triangular classes. Automatic strongness settles the recognition of existing automorphisms, not their construction.
- [Question 11.4](#) now concerns every automorphism of $(\mathbf{No}, \mathbf{Oz})$ with trivial monomial factor, not only the strong ones; it is otherwise unchanged.
- [Question 11.7](#) is related to 12’s question on more intrinsic scalar tests, merged into [question 32.13](#).
- Scope of the answers: $\mathbf{Oz}$, the full real Hahn pairs, and $\mathbf{Oz}[i]$ with conjugation or valuation data. For general characteristic-zero k and unital $\mathfrak{o}$ the non-claims of 04, 09 and 10 that strongness is not shown ([section B](#)) still apply ([question 32.13](#)).

The novelty statements of 11, 12 and 13 were checked against the tree at the merge. No report proves automatic strongness, the scalar detection of summability, the adjoint theorems, or coefficient-moving embeddings of $(\mathbf{No}, \mathbf{Oz})$; their statements about this report and the Diophantine report were accurate at their pins. What they reprove (reconstruction, the criterion, the strong factorization, the Taylor automorphism) is credited above. Priority remains provisional, as each source says. Source 14’s novelty statement (batch 31), accurate at its pin but stale at the merge, is recorded in [section 37.2](#).

32.2 Questions

Each question is a proposed research direction, not a published open problem [11, 12, 13]. Duplicates across the three sources are merged, with every source named.

Question 32.1 (The unrestricted Gaussian stabilizer [12, 13, 14, 15]). Does every ring automorphism of $\mathbf{Oz}[i]$ have a valuation-compatible extension to $\mathbf{No}(i)$, that is, is $\mathfrak{m}_{\mathbb{C}}$ invariant? Can the positive-support component be recovered from $\mathbf{Oz}[i]$ and its fraction field without naming conjugation or a valuation?

$\mathbb{C}$ and $\Pi_{\mathbb{C}}$ are already intrinsic, so the missing object is $\mathfrak{m}_{\mathbb{C}}$; a positive answer with [theorem 30.2](#) would give automatic strongness for all automorphisms of $\mathbf{Oz}[i]$. A negative

answer should exhibit an automorphism of the ring violating a valuation inequality, not merely moving the real axis or being an unrelated automorphism of the algebraically closed fraction field [12, 13]. 14 asks the same, adding whether equivariance of the full constant term at least follows, and proposes first targets: the full field $\mathbb{C}((t^{\mathbb{Q}}))$, then a rank-two lexicographic exponent group. A counterexample must do more than move the real axis, since the strong phase twists of [remark 36.6](#) do that [14]. *Status* (batch 31): open. 15 asks it again (its Question 4) and notes that its topology does not help: a compatible ring topology cannot reconstruct the missing valuation, since on $\mathbf{No}(i)$ every one is indiscrete ([corollary 39.9](#)); 16 recovers $\mathbf{No}(i)$ as a coefficient field but not its conjugation ([question 52.9](#)). *Status* (batch 32): open.

Question 32.2 (Strongness of omnific embeddings [11, 12, 13, 14]). Let $f : \mathbf{No} \hookrightarrow \mathbf{No}$ be a field embedding with $f^{-1}(\mathbf{Oz}) = \mathbf{Oz}$. Must f be strongly additive? What changes if also $\text{ct} \circ f = \text{ct}$? Without strongness, which coefficient and image restrictions survive; can such an f have a proper dense image, or one that is neither closed nor discrete?

The embeddings of [section 27](#) are strong and do not answer the first clause; a counterexample needs a summable family whose image is not summable or has the wrong sum, and the missing step in the isomorphism proof is access to all target tests ([corollary 27.4](#)) [12, 13]. *Status*. The first two clauses have a negative answer if there is a measurable cardinal, more generally an elementary embedding $j : V \rightarrow M$ into a transitive class with a critical point, available as a class. The large-cardinal report ([docs/foundations-and-computation/large-cardinal-embeddings-and-normal-forms](#)) shows that the map $\hat{j}$ obtained by applying j to sign sequences is an ordered field embedding of $\mathbf{No}$ that fixes $\mathbb{R}$, preserves and reflects $\mathbf{Oz}$ and preserves constant terms, while it sends a summable family to a summable family whose sum is not the image of the sum; so it is not strongly additive. It is not onto, so it does not conflict with [theorem 23.2](#). Consequently a positive answer is not provable in the base theory unless that theory with a measurable cardinal is inconsistent; whether a counterexample exists without large cardinals stays open. The necessary gap restriction of [theorem 26.3](#) and the dichotomy of [theorem 28.3](#) remain valid without strongness, whereas the closed-image proof uses the strong expansion formula. The questions on dense images and on images neither closed nor discrete are not settled here [11].

Question 32.3 (Coefficient-fixing embeddings and target tests [13, 14]). For $\mathbb{R}$ -linear embeddings preserving the omnific predicate, can every target test $\text{ct}(y \cdot)$ be represented, after pullback, by a source multiplier?

Such adjoints suffice for strongness and are necessary at set size ([theorem 25.2](#)); the functional $\lambda_{\mathbf{On}}$ shows that a proof for $\mathbf{No}$ must use the multiplicativity of the embedding, not a theorem about all strong linear operators [13]. 14 asks which additional hypotheses make a constant-term-compatible embedding of full Hahn fields strong, in particular for embeddings of $\mathbf{No}$ preserving the integer part. The isomorphism proof fails exactly because a target detector need not lie in the image; at set size an adjoint of the underlying semilinear map suffices, and 14 proposes seeking a criterion that the image supplies all detectors needed for its families, weaker than surjectivity. No embedding theorem is deduced [14].

Question 32.4 (Coefficient-moving embeddings [12]; re-scoped). Which subfields $C_0 \subseteq \mathbb{R}$ occur as $f(\mathbf{No}) \cap \mathbb{R}$ for strongly additive embeddings f with $f^{-1}(\mathbf{Oz}) = \mathbf{Oz}$? Which arise from $\ker d$, and are higher-derivation coefficient lifts needed?

Partly answered by 11: in the exact-monomial strong class the intersections are the fixed fields of the coefficient sections ρ , which are real closed, and Taylor sections give $\ker d$ ([proposition 29.2](#)). Which real closed subfields occur, and the classes with characters, unit

corrections or no monomial condition, stay open. A classification should separate the coefficient homomorphism, the value-group embedding and the admissibility of their combination; block separation is sufficient, not necessary [12].

Question 32.5 (Removing exact monomial preservation [11]). Classify the strong embeddings $f : \mathbf{No} \hookrightarrow \mathbf{No}$ preserving $\mathbf{Oz}$ with $f(t^\gamma) = c_\gamma t^{\tau(\gamma)}(1 + u_\gamma)$, $c_\gamma > 0$, $v(u_\gamma) > 0$. Which compatibility and summability conditions are necessary and sufficient, and can every such embedding be normalized by an $\mathbf{Oz}$ -preserving automorphism?

The criterion of [part I](#) is a starting point, but surjectivity is unavailable: corrections on the image need not extend to all target scales. A first step would keep the corrections on one common positive shift set [11].

Question 32.6 (Elementarity of coefficient-fixing pair embeddings [11]). For an ordered additive $\tau : \mathbf{No} \hookrightarrow \mathbf{No}$, when is $M_{1,\tau} : (\mathbf{No}, \mathbf{Oz}) \hookrightarrow (\mathbf{No}, \mathbf{Oz})$ elementary? Is fixing $\mathbb{R}$ sufficient in the exact-monomial strong category?

[Theorem 29.1](#) gives the necessary condition $\rho = \text{id}$; quantifier elimination for real closed fields cannot decide the expanded structure [11].

Question 32.7 (Coefficient sections [11]). Describe the ordered sections $\rho : \mathbb{R} \hookrightarrow \mathbb{R}((t^C))$ with residue the identity. Which come from one Taylor derivation, compositions of such, or summable families of higher derivations, and what distinguishes sections with the same fixed subfield?

The classification reduces the coefficient part of the embedding problem exactly to this question; the one-sided neighbourhood invariant is only a first invariant [11].

Question 32.8 (Conjugacy of proper copies [11, 13]). Classify the images $J_{\tau,\rho}(\mathbf{No})$ up to $\mathbf{Oz}$ -preserving automorphisms of $\mathbf{No}$. How much do the value image, its cuts, the fixed coefficient field and the one-sided neighbourhood data determine? When are two images $J_{\mu,d}(\mathbf{No})$ conjugate, and which scale embeddings admit Taylor deformations with several independently prescribed motions?

Partly answered by [theorem 29.4](#): the value image alone is insufficient, and continuum many types occur in one field-conjugacy class. 13 notes that since every omnific automorphism fixes $\mathbb{R}$, the positions of the deformed coefficient fields may give finer invariants; a restricted target is $\mathcal{T}_{d,\delta} \circ M_{1,\tau}$ for fixed τ and one shift [11, 13]. *Related* (batch 31): the independent-copies report asks the pair version at a fixed Hahn core: which invariants decide whether an $\mathbf{Oz}$ -preserving automorphism of $\mathbf{No}$ fixing $\mathbb{R}((t^H))$ carries one independent pair of copies with that common field to another (its second research question, “Relative classification of pairs of copies”, unlabelled). It proves no conjugacy result, and the present question stays open.

Question 32.9 (Homogeneity of the pair [12, 13]). Which isomorphisms between set-sized substructures of $(\mathbf{No}, \mathbf{Oz})$ extend to automorphisms of the pair?

Fixing the reals of the domain is necessary ([corollary 27.5](#)); no sufficiency theorem is proved. The question differs from extension of elementary partial maps in a first-order language. A first target is a back-and-forth theorem for truncation-closed Hahn subfields with prescribed real coefficients and compatible integer parts [12, 13].

Question 32.10 (Surcomplex embeddings without a real form [11]). Classify the embeddings of $(\mathbf{No}(i), \mathbf{Oz}[i])$ without assuming commutation with conjugation, and decide which of the nonconjugate copies of [corollary 30.5](#) stay nonconjugate under the larger group.

$\mathbf{Oz}[i]$ reconstructs $\mathbb{C}$ but gives no pointwise real rigidity for the sign invariant [11].

Question 32.11 (The proper-class strong dual [12, 14, 15]). Can the strong $\mathbb{R}$ -linear dual of $\mathbf{No}$ be described by class coefficient functionals meeting every well-ordered set finitely, and which maps have transposes that stay set-supported on every argument?

Such a description must avoid unlicensed collections of classes; the issue is foundational as well as algebraic [12]. 14 asks for a natural enlargement of surreal kernels representing every strong $\mathbb{R}$ -linear functional on $\mathbf{No}$: formal kernels supported on a proper class whose pairing with every set-sized normal form is finite, which recover the set-sized duality on restriction and explain which strong class operators have surreal-valued adjoints. Such kernels are not surreal numbers, and merely writing a proper-class sum is not a definition [14]. *Status* (batch 32): the first clause is *answered* by 15's [proposition 40.5](#). The strong $\mathbb{R}$ -linear functionals on $\mathbf{No}$ are exactly the λ_a of class functions a whose support contains no strictly increasing sequence, equivalently meets every well-ordered set finitely; these are 14's kernels, and they restrict to the set-sized duality. The second clause, which strong operators have transposes that stay set-supported, is 15's Question 9, which asks whether the obstruction is a uniform bound on the supports of the adjoint coefficient classes; it stays open.

Question 32.12 (Singular support bounds [12, 13, 15]). For singular uncountable κ , when does a coefficient- and constant-term-compatible automorphism of $K_{<\kappa}$ preserve summability with total support of size $< \kappa$?

Countable binary witnesses lie in $K_{<\kappa}$ and detect well-ordering and local finiteness, but not the cardinal bound on unions; a sharp theorem may need uniform bounds on supports of monomial images, and a counterexample must preserve the restricted fields themselves [12, 13]. *Partial information* (batch 31): by 14's detection in subfields ([proposition 35.2](#) and [remark 35.4](#)) such an automorphism sends every family summable in $K_{<\kappa}$ to a Hahn-summable family with sum the image of the sum; only the size of the union of the image supports remains. The question stays open. 15 asks it again for support-bounded fields (its Question 6), asking that any large-cardinal dependence be kept apart from its ZFC topological obstruction, which uses only countable sequences ([part V](#)). *Status* (batch 32): open.

Question 32.13 (Other rings and coefficient fields [12, 14]; [merge]). For which integer parts, truncation subrings or distinguished subrings of a real closed Hahn field can one reconstruct a scalar pairing that forces automatic strongness of their stabilizers? In particular, for which characteristic-zero k and unital $\mathfrak{o} \subseteq k$ does $\sigma(\mathcal{R}_{\mathfrak{o}}) = \mathcal{R}_{\mathfrak{o}}$ force a field automorphism of $k((t^{\Gamma}))$ to be strong?

The omnific proof uses a divisible ideal, its multiplier ring and units to recover coefficients, then the order to recover the infinitesimals; it would be useful to find the weakest hypotheses under which this works and examples of failure [12]. The merge adds the second sentence: for $\mathfrak{o} = k$ the divisible-ideal step fails, and for non-real k without valuation data the order step is missing; compare [question 11.7](#). 14 asks the same for integer parts A of a real closed Hahn field, and whether the full constant term is reconstructible from A by the same chain; it suggests the checkable sufficient condition that the divisibility core of A recover a negative-support complement whose multiplier units form the coefficient field, and asks when that condition is necessary and stable under conjugation [14].

Question 32.14 (Isomorphisms of small integer parts [13, 14]). For which ordered groups Γ, Δ does every ring isomorphism $\mathcal{R}_{\Gamma} \cong \mathcal{R}_{\Delta}$ extend to an isomorphism of the full real Hahn fields preserving order and integer parts?

Example 31.2 shows the fraction field is not the Hahn field in general. One route is an intrinsic summation completion compatible with the pairing; another is to find two Hahn fields with indistinguishable small integer parts [13]. 14 asks the automorphism case for divisible Γ , with uniqueness: first determine $\text{Frac } \mathcal{R}_\Gamma$ and its position in K_Γ , since extension through the remaining series is a separate problem [14]. *Status* (batch 31): an extension, if it exists, is unique ([proposition 36.5](#), [merge]); existence stays open.

Question 32.15 (Exponential, omega-map and differential structure [11, 12, 13]). Which proper embeddings preserving **Oz**, and which gap sections and value embeddings, are compatible with the surreal exponential, the omega-map, a surreal derivation or a simplicity relation? Intersect the full omnific stabilizer with the automorphisms preserving such a structure.

The Taylor block embeddings give flexibility only in the algebraic pair and fail omega-map compatibility at 1; the questions must be compared with those of [12], not inferred from the construction, and the exponential rigidity already in the collection is not to be re-solved [11, 12, 13].

Question 32.16 (Restricted workspaces [11]). For a support-restricted subfield of a full Hahn field, when does the bottom-gap extension stay inside it, and which restrictions make the canonical integer part have the whole workspace as fraction field?

The block construction preserves well-ordering but not bounds on support cardinality, order type or boundedness; this matters for birthday cutoffs, computable presentations and transseries subfields [11].

Question 32.17 (Compositions [11]). Describe composition and image inclusion of classified maps in terms of the data (τ, ρ) ; the composite of (τ_1, ρ_1) and (τ_2, ρ_2) has exponent map $\tau_2\tau_1$ and section $J_{\tau_2, \rho_2} \circ \rho_1$. Can image inclusion and ambient extendability be decided from smaller invariants?

Status (batch 31): image inclusion is settled in a special case. Take $\rho = \text{id}$ and the exponent ranges $G_i = h_i(\mathbf{No})$, $i \in I$, of the family in the independent-copies report, which fix a common set-sized divisible group H and are independent over it (`isc:thm:group`). Put $E_J = \mathbb{R}((t^{G_J}))$ with $G_J = H + \sum_{j \in J} G_j$, so that $E_{\{i\}} = J_{h_i, \text{id}}(\mathbf{No})$. That report proves $E_J \cap E_T = E_{J \cap T}$, and $E_J \subseteq E_T$ exactly when $J \subseteq T$ (`isc:thm:boolean`). This decides inclusion and intersection among these images and their full Hahn joins. Composition is not treated there, and the general question stays open.

Question 32.18 (Formalization [11, 12, 13, 14, 15]). Formalize the finite-intersection lemma, the triangular or binary detector, the detection theorem and the constant-term isomorphism theorem for arbitrary full Hahn fields; then the multiplier reconstruction and the omnific specialization; and, separately, the bottom-gap classification for set-sized Hahn fields, the closed-image proof and the two-level index lift, isolating the class-size assumptions for **No**.

The sources propose this order; finite checks remain a separate test layer. Nothing in this part is formalized in Lean [11, 12, 13]. 14 proposes the same order, beginning with an abstract matrix lemma with finite row and column supports ([lemma 34.1](#)), and asks that the full-surreal corollary go through a stated universe-relative or class-coded normal-form interface, not by treating **No** as a set, and that the failure of class duality ([theorem 25.6](#)) be formalized separately; its proposed module names are plans only [14]. 15 asks that the topological collapse and the cyclic boundary be formalized first over abstract ordered monomial systems (the dense-interval and cyclic-cover lemmas, the closure-of-zero facts and the partial-sum argument), then instantiated in set-sized Hahn rings and in the repository's surreal field, with no opaque

imported obstruction and a minimal interface for Hahn-summable families (its Question 12) [15]. 16’s formalization question is [question 52.10](#).

Part IV

A fourth derivation: induced matchings, detector subfields and formal evaluation

This part records manuscript 14, the addition of batch 31. It proves again the main theorems of [part III](#) for the real pair, in less generality; those results are printed once, in [part III](#), with 14 added to their sources, and [section 33.2](#) lists them. What is printed here is what 14 adds: a third proof of scalar detection, by induced matchings, whose witness has all coefficients equal to 1 and needs no choice of coefficients ([section 34](#)); detection inside subfields that contain enough such witnesses ([section 35](#)); and equivariance of formal power-series evaluation, the general adjoint-matrix formula, the rank-one group $\Gamma = \mathbb{R}$ and the source’s shear and truncation examples ([section 36](#)). Its labels carry the sub-prefix `opa:cm:.`

33 Source 14, what it re-derives, and its notation

33.1 The manuscript and its pin

Manuscript 14 is *Automatic Summability from Omnific Arithmetic: constant-term duality, the full omnific stabilizer, and a sharp set-class distinction*, dated 23 September 2026, 24 pages. It is item 07 of batch 31, delivered as the archive `omnific_automatic_summability` in commit `39fe674` and placed as an addition in commit `9d28e28`; it is numbered 14 here after its file prefix, continuing this report’s local sequence. Its finite checks, their recorded run and its source statement are shipped as `code/14-automatic-summability-verify.py`, `data/14-automatic-summability-verification.json` and `14-automatic-summability-SOURCES.md`; its `article.tex`, PDF, README and checksum list are not shipped.

It pins repository commit `3d40856`. That commit contains this report’s `article.tex` exactly as written in batch 26 (`9b80a30`), that is [parts I](#) and [II](#) without the material of source 10, together with the placement `c6359e4` of source 10’s files. It contains neither `b3fa9e2`, the batch-28 write of source 10, nor `21375f8` and `20c4c9f`, the placement and the write of sources 11–13. So 14’s author saw [question 11.1](#) open and saw none of [part III](#); the three batch-30 authors and 14 worked independently of one another. 14 describes the batch-26 report accurately: its criterion concerns strong coefficient-fixing 1-automorphisms, and 14 credits its reconstruction and its convex-scale criterion as antecedents.

33.2 Printed once

The table maps 14’s statements to the results where they are printed. “Same” means the same statement with at most notational differences; otherwise the difference is named. 14’s theorem numbers are those of its own PDF.

14	Printed as	Comparison
Thm. 1.1, Thm. 5.3, Cor. 5.5	theorem 23.2 and corollary 23.4	Less general: 14 treats $\Gamma = \Delta$ divisible, while proposition 23.1 allows ordered isomorphisms between different, possibly non-divisible groups (13).
Lem. 2.2; the Neumann lemma	lemma 2.1 and the support principle of section 2 ; lemma 21.1 is its case of exponent 0	Same.
Lem. 3.1, Thm. 3.3, Cor. 3.4	theorem 21.4	Same statement; 14's proof is the third route of section 34 .
Thm. 4.1, Cor. 4.2	theorem 22.1 and corollary 22.2	14 assumes $\sigma(k) = l$ with $\alpha = \sigma _k$, as 13 does; 12 derives it.
Props. 5.1, 5.2	The reconstruction cited before proposition 23.1 , and that proposition	Re-derivation.
Lem. 5.4 and the remark after Cor. 5.5	<code>odg:thm:fractions</code> ; example 31.2	Same, including the example $\Gamma = \mathbb{Z}$.
Cor. 5.6	corollary 23.5	Same.
The route (5.4)	The dependency chain after theorem 23.2	Same chain.
Thms. 6.1, 6.2, Cor. 6.3	proposition 25.1 and theorem 25.2	Same; the general matrix formula (6.2) is printed as proposition 36.2 .
Thm. 7.1 and the operator after it	theorem 25.6	Same.
Thm. 8.1 and the composition law	theorem 24.1 ; the law after (5.1)	Same.
Thm. 8.2	theorem 3.3 , by 04's polynomial route (lemma 2.2)	Re-proof for $(\mathbb{R}, \mathbb{Z})$.
Cor. 8.3	corollary 24.3 ; the form for $\Gamma = \mathbb{Q}$ after theorem 5.1	Same; the case $\Gamma = \mathbb{R}$ is remark 36.3 .
Prop. 9.1 and the truncation example	example 4.19 , (4.5)	The inverse of that example with the two letters swapped; remark 36.4 .
Prop. 10.1, Thm. 10.2 Thm. 10.3	theorem 30.2 and its proof theorem 30.1	Same. Equivalent hypothesis (remark 36.6).
Ex. 10.4	The phase twists recorded after theorem 10.2	An instance.
Questions 12.1–12.8 and the question of its Section 10	section 37	One new question; the others are merged.

So 14 is a third independent derivation of the automatic-strongness theorem, after 12 and 13, and adds nothing to its generality. Its route $\mathcal{R}_\Gamma \rightarrow \Pi \rightarrow \mathcal{A}_\Gamma \rightarrow \mathbb{R} \rightarrow \mathfrak{m} \rightarrow \text{ct} \rightarrow \text{summation}$ is the chain printed after [theorem 23.2](#). Where 14's proof of a shared result differs only in presentation it is not reprinted. Its genuinely different proof, the induced matching, is kept as a marked third route.

33.3 Conventions and notation of this part

This part uses the conventions of [section 20.3](#): the small- t convention $t^\gamma = \omega^{-\gamma}$, $K_\Gamma = k((t^\Gamma))$, Π_Γ , $\mathcal{R}_\Gamma = \mathbb{Z} \oplus \Pi_\Gamma$, $\mathcal{A}_\Gamma = \mathbb{R} \oplus \Pi_\Gamma$ and the pairing $\langle x, y \rangle_0 = \text{ct}(xy)$. Manuscript 14 also writes in the t -convention. Its symbols are renamed as follows.

14's symbol	Here
$A_\Gamma = \mathbb{Z} \oplus \Pi$, and A	$\mathcal{R}_\Gamma$; for $\Gamma = \mathbf{No}, \mathbf{Oz}$.
$B = k \oplus \Pi$	$\mathcal{A}_k$; for $k = \mathbb{R}, \mathcal{A}_\Gamma$.
$\langle x, y \rangle$	$\langle x, y \rangle_0$.
$\tau : k \rightarrow l$, the coefficient isomorphism	α , as in theorem 22.1 .
ϕ , the exponent automorphism; $M_{\chi, \phi}$	τ ; $M_{\chi, \tau}$ of (5.1) .
ρ , a ring automorphism of $\mathbf{Oz}$; $\tilde{\rho}$	σ and $\tilde{\sigma}$, as in theorem 23.2 .
ν , a normalized 1-automorphism; $\mathcal{U}, \mathcal{G}, \mathcal{M}$	u ; $\mathcal{U}_{\mathcal{R}_\mathbb{Z}}(\Gamma)$ (and $\mathcal{U}_{\mathbf{Oz}}$); the stabilizer of $\mathcal{R}_\Gamma$; the monomial maps.
$L_\nu(g)$, $a_{\nu, \delta}$	$h_u(g)$, $a_{u, \delta}$ of (3.1) .
T^*	$T^\dagger$; the matrix $T_{h, g} = [t^h]T(t^g)$ keeps its name.
Λ	$\lambda_{\mathbf{On}}$ of (25.2) ; 14's $[\omega^\alpha]x$ is $[t^{-\alpha}]x$.
F , a power series	$\mathbf{F}$.
F , a subfield	K' .
$P_{<\gamma}$, truncation	$T_{<\gamma}$, as in theorem 8.2 .
e_1, e_2 of its shear (e_1 the larger scale); $s = t^{e_2}$	e_2, e_1 of example 4.16 ($e_1 \ll e_2$); s is written t^{e_1} .
h , an additive map	θ .
$u_1, \dots, u_m$, u , infinitesimals	$z_1, \dots, z_m$, z ; u is the leading-term-fixing factor.
$d > 0$ in its cancellation example	$\delta > 0$; d is a derivation of $\mathbb{R}$ in part III .
$\mathbf{No}[i]$, $\mathbf{Oz}[i]$; $A_\mathbb{C}$, $K_\mathbb{C}$, $\Pi_\mathbb{C}$, $B_\mathbb{C}$	$\mathbf{No}(i)$, $\mathbf{Oz}[i]$; $\mathcal{R}_\Gamma^\mathbb{C}$, $K_\Gamma^\mathbb{C}$, $\Pi_\Gamma^\mathbb{C}$, $\mathbb{C} \oplus \Pi_\Gamma^\mathbb{C}$.
$L = l((u^\Delta))$	$L = l((t^\Delta))$, with t for the monomials of both fields.
$a \ll b$	$a \ll b$, the same relation.

The collisions are resolved here only, and the tempting false readings are wrong:

- 14's A_Γ is *not* $\mathcal{A}_\Gamma$: 14's $A_\Gamma = \mathbb{Z} \oplus \Pi$ is $\mathcal{R}_\Gamma$, and $\mathcal{A}_\Gamma = \mathbb{R} \oplus \Pi$ is 14's B .
- 14's τ is a coefficient isomorphism; τ here is the exponent automorphism, 14's ϕ . 14's ρ is a ring automorphism of $\mathbf{Oz}$; ρ in [part III](#) is a coefficient section.
- 14's F has two meanings, neither of which is the isomorphism F of [theorem 22.1](#) or the profile F of [\(4.4\)](#).
- In 14's shear e_1 is the *larger* scale; in [example 4.16](#) it is the smaller. The letters are swapped here.
- 14's truncation $P_{<\gamma}$ is not one of the zero-cut projections P_-, P_0, P_+ .
- 14's additive h is not the logarithmic character h_σ .

14 calls an additive map *strong* when it sends summable families to summable families with the same sum; this is “strongly additive” here, and 14 states both directions where it asserts them. Its “normalized 1-automorphism” is a 1-automorphism in the sense of [section 1.5](#); 14 does not assume it strong, and by [corollary 23.4](#) every one preserving $\mathcal{R}_\Gamma$ is.

34 Induced matchings: a third route to scalar detection

Lemma 34.1 (An infinite induced matching [14]). *Let $(a_{n,j})_{n \in \mathbb{N}, j \in I}$ be a matrix over a field, I a set. Suppose that every row is nonzero and has finite support, and that every column has finite support. Then there are row indices $n_0 < n_1 < \dots$ and distinct column indices $j_0, j_1, \dots$ with*

$$a_{n_r, j_r} \neq 0, \quad a_{n_r, j_s} = 0 \quad (r \neq s).$$

Proof. Choose a nonzero entry in row $n_0 = 0$, in column j_0 . After r choices let E be the union of the supports of the chosen rows, a finite set of columns. Only finitely many rows meet E , since each column is finite. Choose a row n_r beyond all earlier chosen rows that does not meet E , and a nonzero entry in it, in column j_r . Then $j_r \notin E$, so column j_r vanishes on the earlier chosen rows, and row n_r vanishes on the earlier chosen columns, which lie in E . Induction gives the matching. The selection uses only the pattern of nonzero entries, not the characteristic or the values of the coefficients. $\square$

Third route to (iii) $\Rightarrow$ (i) in [theorem 21.4](#) [14]. Assume (iii). The tests $y = t^{-g}$ give pointwise finiteness. Suppose the union S of the supports is not well ordered, and choose $g_0 > g_1 > \dots$ in S . Put $a_{n,j} = [t^{g_n}]x_j$. Row n is nonzero because $g_n \in S$, and finite by pointwise finiteness. Column j is finite, since infinitely many g_n in $\text{supp } x_j$ would descend in a well-ordered set. [Lemma 34.1](#) gives n_r and j_r ; put

$$y = \sum_{r \geq 0} t^{-g_{n_r}}, \tag{34.1}$$

a Hahn series, since its exponents increase, with countable support and every coefficient equal to 1. Then $\langle x_{j_r}, y \rangle_0 = \sum_{s \geq 0} [t^{g_{n_s}}]x_{j_r} = a_{n_r, j_r} \neq 0$ for every r : the sum is finite, and only its term $s = r$ is nonzero. The distinct j_r contradict (iii). $\square$

The three routes differ in how cancellation is prevented. 12's triangular detector ([lemma 21.2](#)) makes the family triangular and then chooses coefficients in $\{0, 1\}$ one at a time; 13's binary lemma ([lemma 21.3](#)) keeps the family and chooses coefficients row by row; 14 passes to an induced matching, a diagonal submatrix, so each selected pairing is one matrix entry and

nothing is chosen. 14's witness is therefore a special case of the tests in (iii): countable support and all coefficients 1. Every construction uses a countable subset of one set-sized support, so the route works for $\Gamma = \mathbf{No}$ and for $\mathbf{No}(i)$ [14].

Corollary 34.2 (Monomial and all-ones tests suffice [14]). *A set-indexed family (x_j) in a full Hahn field $K = k((t^\Gamma))$, Γ a set or $\mathbf{No}$, is Hahn summable if and only if every exponent lies in only finitely many supports and, for every strictly increasing sequence $(h_n)_{n \in \mathbb{N}}$ in Γ , only finitely many j have $\langle x_j, \sum_n t^{h_n} \rangle_0 \neq 0$.*

Proof. Necessity is (i) $\Rightarrow$ (ii) of [theorem 21.4](#). For sufficiency the third route uses only monomial tests and the series (34.1), whose exponents $-g_{n_r}$ increase strictly. $\square$

14 states this through the proof of its detection theorem and its Proposition 11.3 ([proposition 35.2](#)).

Example 34.3 (A nonexistent sum disguised as zero [14]). Let $\delta > 0$ in Γ and, for $n \geq 1$, $x_{n,+} = t^{-n\delta}$ and $x_{n,-} = -t^{-n\delta}$. Each exponent lies in exactly two members, and grouping the members in pairs would cancel every coefficient. The family is not Hahn summable, since $\{-n\delta : n \geq 1\}$ descends. The all-ones series $y = \sum_{n \geq 1} t^{n\delta}$ gives $\langle x_{n,+}, y \rangle_0 = 1$ and $\langle x_{n,-}, y \rangle_0 = -1$ for every n : the test fails before any cancellation can hide the infinitely many contributions.

Here each exponent occurs twice with cancelling coefficients, whereas in [example 21.6](#) each exponent occurs once. The detection theorem recognizes admissibility of the family, not a value assigned by some other summation convention; neither conditional convergence nor an ordering of the index set is used [14].

Remark 34.4 (Summability is not convergence [14]). The family $(t^{n/(n+1)})_{n \geq 1}$ is Hahn summable over $\mathbb{Q}$, since its exponents increase, but its terms do not tend to zero in the valuation topology. A valuation-preserving automorphism is continuous, but continuity speaks only about families that converge in that topology, and the detector also handles bounded increasing supports; for $\mathbf{No}$, set-indexed topological convergence meets further size obstructions. The detection theorem reduces summability to scalar product tests, without completeness or net arguments. This agrees with [remark 21.7](#). 14 identifies the dual supports, the sets meeting every well-ordered set finitely, with the reverse-well-ordered sets, as the total-order case of [3, Theorem 3 and Lemmas 4.1–4.2], and claims no origin of that duality. *Batch 32*: for ring topologies with jointly continuous multiplication the obstruction on $\mathbf{No}$ is complete and is one of density, not of size: only the indiscrete topology realizes every Hahn sum as a limit ([corollary 39.9](#)).

35 Detection in subfields with enough detectors

Definition 35.1 (Countable detector property [14]). Let K' be a subfield of $K = k((t^\Gamma))$, Γ a set or $\mathbf{No}$, containing k and every monomial. K' has the *countable detector property* if $\sum_{n \geq 0} t^{h_n} \in K'$ for every strictly increasing sequence $(h_n)_{n \in \mathbb{N}}$ in Γ .

Proposition 35.2 (Detection in a subfield [14]). *Let K' have the countable detector property.*

- (a) *A set-indexed family in K' is Hahn summable in K if and only if, for every $y \in K'$, only finitely many $\langle x_j, y \rangle_0$ are nonzero.*
- (b) *Let σ be a field automorphism of K' with $\sigma(k) = k$ and $\text{ct}(\sigma x) = \alpha(\text{ct } x)$ for $x \in K'$, where $\alpha = \sigma|_k$. If (x_j) in K' is Hahn summable in K , so is $(\sigma(x_j))$. If moreover $s = \sum_j x_j$ lies in K' , then $\sum_j \sigma(x_j) = \sigma(s)$; in particular this sum lies in K' .*

Proof. (a) Necessity holds because $K' \subseteq K$ (theorem 21.4). The tests of corollary 34.2 are monomials and series (34.1), which lie in K' ; this gives sufficiency. (b) For $y \in K'$ write $y = \sigma(z)$ with $z \in K'$. Then $\langle \sigma x_j, y \rangle_0 = \text{ct}(\sigma(x_j z)) = \alpha(\langle x_j, z \rangle_0)$, nonzero for finitely many j by (a), so $(\sigma(x_j))$ is summable in K by (a). Let s' be its sum in K . By (21.2) in K , for every $y \in K'$,

$$\langle \sigma(s), y \rangle_0 = \alpha(\langle s, z \rangle_0) = \sum_j \alpha(\langle x_j, z \rangle_0) = \sum_j \langle \sigma x_j, y \rangle_0 = \langle s', y \rangle_0,$$

with finite sums. So $\sigma(s) - s'$ pairs to zero with every $y \in K'$, in particular with every monomial, and $\sigma(s) = s'$. $\square$

The inverse satisfies the same hypotheses with α^{-1} . As in the proof of theorem 22.1, the hypothesis $\sigma(k) = k$ may be replaced by covariance $\text{ct}(\sigma x) = \alpha(\text{ct} x)$ for some field automorphism α of k : since K' contains every monomial the pairing separates K' , and $\sigma(r) = \alpha(r)$ follows for $r \in k$ [merge]. The proposition says nothing about sums that leave K' , and it concerns the summation of the ambient full Hahn field, not a summation intrinsic to K' .

Example 35.3 (Puisseux series have too few detectors [merge]). Let $K = \mathbb{R}((t^{\mathbb{Q}}))$ and let $K' = \bigcup_{N \geq 1} \mathbb{R}((t^{\frac{1}{N}\mathbb{Z}}))$ be the field of Puiseux series; it contains $\mathbb{R}$ and every monomial. The family $x_n = t^{-n/(n+1)}$, $n \geq 1$, is not Hahn summable in K , since its exponents descend. For $y \in K'$ with $\text{supp } y \subseteq \frac{1}{N}\mathbb{Z}$, $\langle x_n, y \rangle_0 = [t^{n/(n+1)}]y$ is nonzero only if $Nn/(n+1) \in \mathbb{Z}$, that is only if $n+1$ divides N , since n and $n+1$ are coprime. So every test in K' gives finitely many nonzero pairings, and (a) of proposition 35.2 fails for K' . Accordingly $\sum_n t^{n/(n+1)}$, the sum of the summable family of remark 34.4, is not a Puiseux series.

Remark 35.4 (Relation to support-bounded fields [merge]). For every uncountable cardinal κ , regular or singular, the field $K_{<\kappa}$ of proposition 31.1 has the countable detector property. So proposition 35.2 gives for all such κ what proposition 31.1 gives for regular κ as far as ambient summability goes: a coefficient- and constant-term-compatible automorphism of $K_{<\kappa}$ sends every family that is summable in the sense of $K_{<\kappa}$ to a family that is Hahn summable in K , with fewer than κ nonzero members and with sum the image of the sum. For singular κ what question 32.12 still asks is only whether the union of the image supports has size $< \kappa$. The warning of remark 31.3, that the detector must lie in the field tested and that this fails for arbitrary subfields, stands: example 35.3 is such a failure, and the countable detector property is a sufficient condition, not a characterization (question 37.1). 14 does not claim the property for transseries, Puiseux or left-finite subfields; it notes that a series on an increasing sequence of exponents bounded above can fail a left-finiteness restriction [14]. Question 32.16 (11) asks the analogous question for the bottom-gap embeddings.

36 Further consequences

Theorem 36.1 (Formal evaluation is transported [14]). *Let Γ be **No** or a nonzero divisible set-sized ordered abelian group, and let σ be a field automorphism of $K_\Gamma = \mathbb{R}((t^\Gamma))$ with $\sigma(\mathcal{R}_\Gamma) = \mathcal{R}_\Gamma$. For $F \in \mathbb{R}[[X_1, \dots, X_m]]$ and $z_1, \dots, z_m \in \mathfrak{m}_\Gamma$,*

$$\sigma(F(z_1, \dots, z_m)) = F(\sigma(z_1), \dots, \sigma(z_m)). \quad (36.1)$$

In particular $\sigma(\exp z) = \exp(\sigma z)$ and $\sigma(\log(1 + z)) = \log(1 + \sigma z)$ for $z \in \mathfrak{m}_\Gamma$.

Proof. Write $F = \sum_\nu c_\nu X^\nu$ over $\nu \in \mathbb{N}^m$. The union of the supports of the z_j is a well-ordered subset of $\Gamma^{>0}$, so the Neumann support principle (section 2) makes $(c_\nu z^\nu)_\nu$ Hahn summable; its sum is the formal evaluation. By proposition 23.1 and corollary 23.4 (and theorem 23.2

for $\mathbf{No}$), σ fixes $\mathbb{R}$, maps $\mathfrak{m}_\Gamma$ onto itself and is strongly $\mathbb{R}$ -linear, so $\sigma(\sum_\nu c_\nu z^\nu) = \sum_\nu c_\nu \sigma(z)^\nu$, and the $\sigma(z_j)$ are again infinitesimal. $\square$

No radius of convergence enters: a formal series with arbitrarily fast-growing real coefficients is evaluated at Hahn infinitesimals through support finiteness. For a valuation-compatible Gaussian automorphism as in [theorem 30.2](#), with $\alpha = \sigma|_{\mathbb{C}}$, the same proof gives $\sigma(F(z)) = (\alpha F)(\sigma(z))$, where α acts on the coefficients; for the conjugation-compatible automorphisms of [theorem 30.1](#), α is the identity or complex conjugation [14]. The exponential here is the local one at infinitesimals ([section 2](#)); nothing is claimed about the Gonshor exponential of infinite arguments, whose compatibility is a further condition studied in [12].

Proposition 36.2 (The adjoint as a sign-reversed transpose [14]). *In the setting of [section 25](#), let $T : K \rightarrow L$ be strongly k -linear and put $T_{\eta,\gamma} = [t^\eta]T(t^\gamma)$ for $\gamma \in \Gamma$, $\eta \in \Delta$. Then for every $y = \sum_\eta y_\eta t^\eta \in L$ and $\gamma \in \Gamma$*

$$[t^\gamma]T^\dagger(y) = \sum_{\eta \in \Delta} T_{-\eta, -\gamma} y_\eta, \quad \text{so} \quad (T^\dagger)_{\gamma, \eta} = T_{-\eta, -\gamma}, \quad (36.2)$$

with finitely many nonzero terms.

Proof. $[t^\gamma]T^\dagger(y) = \langle t^{-\gamma}, T^\dagger y \rangle_0 = \langle T(t^{-\gamma}), y \rangle_0$ by (25.1), and the last constant term is $\sum_\eta [t^{-\eta}]T(t^{-\gamma}) y_\eta$, finite by [lemma 21.1](#). $\square$

For a coefficient-fixing isomorphism F with $\text{ct} \circ F = \text{ct}$ one has $F^\dagger = F^{-1}$ ([theorem 25.2](#)), and (36.2) becomes the coefficient-matrix formula (22.2) of 13. The general formula is 14's; it presupposes the strongness of T , which [theorem 25.2](#) supplies, and 14's suite checks it only on finite matrices ([section C](#)).

Remark 36.3 (Rank one with $\Gamma = \mathbb{R}$ [14, 15]). By [corollary 24.3](#) every field automorphism of $\mathbb{R}((t^\mathbb{R}))$ preserving $\mathcal{R}_\mathbb{R}$ is a monomial map $M_{\chi, \tau}$. Here $\tau(\gamma) = \lambda\gamma$ with $\lambda > 0$, since an order-preserving additive bijection of $\mathbb{R}$ is monotone and additive, hence $\mathbb{R}$ -linear; and $\chi(\gamma) = \exp(\theta(\gamma))$ with $\theta : \mathbb{R} \rightarrow \mathbb{R}$ additive, since χ is positive. θ need not be continuous: any $\mathbb{Q}$ -linear map defined on a Hamel basis will do, and every such pair (λ, θ) gives an automorphism by (5.1). So rank one removes the factor u , not all automorphisms and not all nonclassical coefficient scalings. For $\Gamma = \mathbb{Q}$ the character is $q \mapsto c^q$ (the paragraph after [theorem 5.1](#)).

Remark 36.4 (The shear of source 14 and truncation [14, 15]). Let $\Gamma = \mathbb{Q}e_1 \oplus \mathbb{Q}e_2$ with $e_1 \ll e_2$ as in [example 4.16](#), and let θ be the e_2 -coordinate. 14's automorphism, in these letters,

$$t^{ae_2 + be_1} \mapsto t^{ae_2 + be_1} (1 + t^{e_1})^a \quad (a, b \in \mathbb{Q}),$$

is the binomial twist $\sigma_{e_1, \theta}^{1+\xi}$ of (4.5), so it is a strong $\mathbb{R}$ -linear 1-automorphism preserving $\mathcal{R}_\Gamma$ ([theorem 4.18](#), since $\theta(C_{e_1}) = 0$); it is the inverse of [example 4.19](#). It sends t^{-e_2} to $\sum_{n \geq 0} (-1)^n t^{-e_2 + ne_1}$, a monomial to an infinite series. With $\gamma = -e_2 + \frac{1}{2}e_1$ and $T_{<\gamma}$ the truncation below γ ,

$$T_{<\gamma}(\sigma(t^{-e_2})) = t^{-e_2}, \quad \sigma(T_{<\gamma}(t^{-e_2})) = \sigma(t^{-e_2}) \neq t^{-e_2}.$$

So automatic strongness does not give commutation with truncation at a monomial, even when the exponent automorphism is the identity. This does not contradict [theorem 8.2](#): the definable truncation $T_{<\gamma}(f) = mP_-(f/m)$ uses the parameter $m = t^\gamma$, and σ carries it to $f \mapsto \sigma(m)P_-(f/\sigma(m))$, truncation relative to $\sigma(m) = t^\gamma(1 + t^{e_1})^{-1}$, which is not a monomial. The tempting reading “every $\mathbf{Oz}$ -preserving automorphism commutes with truncation at every monomial” is false; the true statements are commutation with P_-, P_0, P_+ ([corollary 5.2](#))

and with the projection onto any union of signed Archimedean blocks ([proposition 3.10](#)), in agreement with the paragraph after that proposition. 14 relates naming truncations to the complexity studied by Camacho [4].

Proposition 36.5 (Extensions from $\mathcal{R}_\Gamma$ are unique [merge]). *Let Γ be a nonzero divisible set-sized ordered abelian group. A ring automorphism σ of $\mathcal{R}_\Gamma$ has at most one extension to a field automorphism of K_Γ ; an extension, if one exists, fixes $\mathbb{R}$ and is strongly $\mathbb{R}$ -linear with strongly $\mathbb{R}$ -linear inverse.*

Proof. An extension $\tilde{\sigma}$ satisfies $\tilde{\sigma}(\mathcal{R}_\Gamma) = \sigma(\mathcal{R}_\Gamma) = \mathcal{R}_\Gamma$, so by [corollary 23.4](#) it fixes $\mathbb{R}$ and is strongly $\mathbb{R}$ -linear with strong inverse. Its values on monomials are forced: $\tilde{\sigma}(t^\gamma) = \sigma(t^\gamma)$ for $\gamma < 0$, since $t^\gamma \in \Pi_\Gamma$, and $\tilde{\sigma}(t^\gamma) = \sigma(t^{-\gamma})^{-1}$ for $\gamma > 0$. Applied to the normal-form family $(a_\gamma t^\gamma)$ of any f , strongness gives $\tilde{\sigma}(f) = \sum_\gamma a_\gamma \tilde{\sigma}(t^\gamma)$, which depends only on σ . $\square$

This answers the uniqueness clause of 14's Question 12.7 without assuming $\text{Frac } \mathcal{R}_\Gamma = K_\Gamma$, which fails in general ([example 31.2](#)); 14 notes only that uniqueness on the fraction field is automatic. Existence is [question 32.14](#).

Remark 36.6 (Gaussian statements of source 14 [14]). 14's real-structure theorem assumes $\sigma(\mathbf{No}) = \mathbf{No}$, where [theorem 30.1](#) assumes commutation with conjugation. The hypotheses are equivalent: conjugation fixes exactly $\mathbf{No}$, and conversely $\sigma(i) = \pm i$ gives $\sigma(x + iy) = \sigma(x) \pm i\sigma(y)$, which commutes with conjugation when $\sigma(\mathbf{No}) = \mathbf{No}$. 14's example over $\Gamma = \mathbb{Q}$, the strong map $\sum a_q t^q \mapsto \sum a_q e^{i\pi q/2} t^q$, preserves $\mathcal{R}_\mathbb{Q}^\mathbb{C}$, fixes $\mathbb{C}$ and sends t to it ; on $\mathbf{No}(i)$ it uses $\chi(\gamma) = \exp(i\pi\theta(\gamma)/2)$ with $\theta(\gamma) = \text{ct}(\gamma)$. It is a phase twist of the kind recorded after [theorem 10.2](#) (odg: def: lem: twist, saut: thm: phase). Since it is strong, it does not bear on [question 32.1](#); it only shows that preservation of $\mathbf{Oz}[i]$ alone does not give the hypothesis of [theorem 30.1](#).

37 Questions of source 14, and stale statements

37.1 Status of the source's questions

14 poses eight questions (its 12.1–12.8), proposed as research problems, not as published open problems, and repeats its Gaussian question at the end of its Section 10. Seven are merged into questions already in the report, with 14 added to their sources:

- 12.1 and the Section-10 question (must every automorphism of $\mathbf{No}(i)$ preserving $\mathbf{Oz}[i]$ be strong; does $\mathbf{Oz}[i]$ determine the valuation ring, or at least ct -equivariance?): [question 32.1](#), open; 14's suggested first targets are recorded there.
- 12.2 (which hypotheses make a constant-term-compatible embedding strong; embeddings of $\mathbf{No}$ preserving $\mathbf{Oz}$): [questions 32.2](#) and [32.3](#). The first clauses of the former have a negative answer given a measurable cardinal, as recorded there.
- 12.4 (class-supported kernels for strong functionals on $\mathbf{No}$): [question 32.11](#).
- 12.5 (other integer parts): [question 32.13](#), and [question 11.7](#) for the criterion.
- 12.6 (a parameter space for the normalized factor): [question 11.2](#), with [question 11.4](#); open. 14 adds that the convex criterion recognizes existing automorphisms and does not construct them, and that a parameterization must distinguish pointwise positive support from uniform control on summable families.
- 12.7 (does every automorphism of $\mathcal{R}_\Gamma$ extend to K_Γ , uniquely?): the automorphism case of [question 32.14](#). Uniqueness is answered by [proposition 36.5](#); existence stays open.

- 12.8 (a kernel-checked automatic-strongness theorem): [question 32.18](#).

One question is new:

Question 37.1 (Minimal closure for algebraic detection [14, 15]). Can the countable detector property be weakened to a necessary and sufficient closure condition on a subfield K' for the product detection of [proposition 35.2\(a\)](#)? Which natural computable or transseries subfields satisfy that condition?

The induced-matching proof needs one all-ones detector for a subsequence of each bad configuration, not every countable series with arbitrary coefficients, so a closure principle could quantify over bad support configurations rather than over all increasing sequences. The question has algebraic and computability aspects: the existence of a detector need not give an algorithm that recognizes or constructs it from a chosen representation [14]. [Example 35.3](#) shows that some closure condition is needed [merge]. 15 asks the same (its Question 5): which proper subfields admit the detector, whether closure under the countable sparse detectors is equivalent to a natural support condition, and with what effect on constant-term-preserving embeddings and isomorphisms; it asks for necessary and sufficient support conditions rather than sufficient completeness hypotheses [15]. *Status* (batch 32): partly addressed by the sufficient condition of [definition 35.1](#) and [proposition 35.2](#); open.

37.2 Stale and checked statements

- *Stale novelty*. 14’s abstract says that “the strong-additivity hypothesis can be removed” from this report’s real stabilizer classification, its introduction that it “supplies a missing implication”, and its questions section that its main result “closes the automatic-strongness question for the real omnific pair”. At its pin 3d40856 this was accurate: [question 11.1](#) was open. At the merge it is stale: sources 12 and 13 answered [question 11.1](#) in 20c4c9f ([theorem 23.2](#)). 14 is recorded as an independent third derivation, and its priority statement is not printed as current; its own caveat that global priority is not certified is kept.
- *Checked true*. 14’s description of the pinned report (the criterion concerns strong coefficient-fixing 1-automorphisms; reconstruction and criterion are credited), the example $\text{Frac}(\mathbb{Z} \oplus t^{-1}\mathbb{R}[t^{-1}]) = \mathbb{R}(t)$ for $\Gamma = \mathbb{Z}$, and the finite checks of its suite, rerun for the merge ([section C](#)). No mathematical error was found in 14.
- *Not rechecked*. 14’s pinpoint citations of [13] (an example of a non-strongly additive automorphism) and of [3] (Theorem 3, Lemmas 4.1–4.2) are reported as 14 gives them.

14’s limitations are listed with the other sources’ in [section B](#).

Part V

Topological collapse of Hahn summation

This part records manuscript 15, an addition of batch 32. Its new subject is topological: which ring topologies, with jointly continuous multiplication, realize every Hahn sum as the limit of its net of finite partial sums? For a densely ordered exponent group every such topology on $\mathfrak{o} \oplus \Pi$ is pulled back from the constants along ct , so none is Hausdorff; on $\mathbf{No}$ and $\mathbf{No}(i)$ only the indiscrete topology qualifies; and a set-sized full Hahn field has a Hausdorff one exactly

when its exponent group is 0 or order-isomorphic to $\mathbb{Z}$ (section 39). The algebraic half of 15, a fourth derivation of the automatic strongness of part III, is printed once there with 15 added to the sources; what it adds to that half is in section 40, and its pairing topology, which realizes exactly the Hahn sums but has only separately continuous multiplication, in section 41. Its labels carry the sub-prefix `opa:tc:`.

38 Source 15, what it re-derives, and its notation

38.1 The manuscript and what it read

Manuscript 15 is *Algebraic Rigidity and Topological Collapse of Surreal Arithmetic: Hahn summation, omnific integer topologies, and the discrete-exponent boundary*, dated 23 September 2026, 29 pages. It is item 03 of batch 32, delivered as the archive `Surreal_Arithmetic_Topological_Collapse` in commit `aa268a4` and placed as an addition in commit `7d04483`; it is numbered 15 here, continuing this report's local sequence. Its finite checks, their recorded run, its build record, its `Makefile` and its source audit are shipped as `code/15-topological-collapse-verify.py`, `data/15-topological-collapse-verification.json`, `data/15-topological-collapse-BUILD_AUDIT.json`, `code/15-topological-collapse-Makefile` and `15-topological-collapse-SOURCE_AUDIT.md`; its `.tex`, PDF and README are not shipped.

15 names no repository commit. Its pins are two blob identifiers of this report's `article.tex`, read through a connector: `059939b5`, the file as written in `b3fa9e2` (batch 28: parts I and II with source 10's material), in which question 11.1 was still open, and `60f3b172`, the file as written in `20c4c9f` (batch 30, with part III), which it found in a later directory read together with the audits of 11, 12 and 13 (blobs `050b52f`, `0e4e3a1`, `c4d3939`, as placed in `21375f8`). The merge checked all five identifiers against the history. So 15 saw part III but neither part IV (`d4d72d7`) nor the batch-31 reciprocal notes (`ab18444`). Its account of this report is accurate for what it read: it credits automatic strongness, scalar detection, constant-term covariance, adjoints, the class-size failure of duality and the qualified Gaussian versions to 12 and 13, and the criterion, the reconstruction, the strong factorization and the shear examples to this report; it claims as new only its Section 3, the topology. Its reading was incomplete, as its audit says: it does not mention 14's induced-matching route or detection in subfields, and it does not mention the congruence topologies of the quotient and Diophantine reports, which are instances of its classification (remark 39.6).

38.2 Printed once

The table maps 15's statements to the results where they are printed; 15's numbers are those of its own PDF. "Same" means the same statement up to notation.

15	Printed as	Comparison
Lem. 2.2	lemma 2.1, the support principle of section 2; lemma 21.1	Same.

15	Printed as	Comparison
Thm. 4.2	theorem 21.4	Same, except that 15's (iii) allows all countably supported tests, without the restriction to coefficients in $\{0, 1\}$; its disjoint-row proof is a variant of the third route (lemma 40.1).
Rem. 4.3	example 40.3	New example, beside example 34.3 .
Cor. 4.4	corollary 40.2	New.
Prop. 4.5	example 21.6	The same family; 15 states it for all finite-support tests.
Thm. 5.1, Rem. 5.2	theorem 22.1 and the paragraph after it	15 assumes $\sigma(k) = \ell$ and $\alpha = \sigma _k$, as 13 and 14 do.
Prop. 5.3	"Adjoint implies strong" in theorem 25.2	Same, for full Hahn fields of any size, as 13 notes.
Lems. 6.1, 6.2, Rem. 6.3	lemma 2.3 , <code>odg: def: thm: multiplier</code> , <code>odg: def: thm: autreal</code> ; proposition 23.1	Re-derivation.
Thm. 6.4	proposition 23.1 and corollary 23.4	Same, for one group.
Lem. 7.1	<code>odg: thm: fractions</code>	Same.
Thm. 1.2, its proof in Section 7, Cors. 7.2, 7.4	theorem 23.2 , (23.1)	Same; 15 credits 12 and 13.
Cor. 7.3; Rem. 7.5	corollary 23.5 ; remark 23.3	Same.
Thm. 8.1	theorem 24.1	Same.
Prop. 8.2	theorem 3.3 , conditions (i), (iv), (v)	Re-proof for $(\mathbb{R}, \mathbb{Z})$ by 04's polynomial route; 15 credits this report.
Rem. 8.3	remark 24.2	Same.
Cor. 8.4, the forms for $\mathbb{Q}$ and $\mathbb{R}$	corollary 24.3 , the paragraph after theorem 5.1 , remark 36.3	Same.
Ex. 8.5	remark 36.4	The same map, the inverse of example 4.19 .
Lem. 9.1	lemma 2.3 , <code>odg: def: thm: multiplier</code>	Re-derivation.
Thms. 9.2, 9.3	theorems 30.1 and 30.2	Same.
Cor. 9.4	corollary 40.4	New, the ring form.
Thm. 10.2, Cor. 10.3	proposition 25.1 and theorem 25.2	Same.
Prop. 10.4, Rem. 10.6	theorem 25.6 and the paragraph after it	Same.
Prop. 10.5	proposition 40.5	New.
Section 3; Section 11	section 39 ; section 41	New.

15	Printed as	Comparison
Questions 1–12	section 42	Six new, six merged.

So 15 is a fourth independent derivation of the automatic strongness of the real pair, after 12, 13 and 14, and it says so: it keeps its proof “for the comparison with the topological collapse theorem, not as a new resolution”. Where its proofs of shared results differ only in presentation they are not reprinted.

38.3 Conventions and notation of this part

This part uses the small- t convention (1.1) and the notation of [section 20.3](#): $K_\Gamma = k((t^\Gamma))$, Π_Γ , $\mathfrak{m}_\Gamma$, and for $k = \mathbb{R}$ $\mathcal{R}_\Gamma = \mathbb{Z} \oplus \Pi_\Gamma$ and $\mathcal{A}_\Gamma = \mathbb{R} \oplus \Pi_\Gamma$, with the pairing $\langle x, y \rangle_0 = \text{ct}(xy)$. For a field k and a unital subring $\mathfrak{o} \subseteq k$, $\mathcal{R}_\mathfrak{o} = \mathfrak{o} \oplus \Pi$ is the ring of [section 1.5](#), but in [section 39](#) the exponent group is only assumed *densely ordered* (nonzero, with no least positive element), not divisible; divisible groups are densely ordered, and $\mathcal{R}_\mathfrak{o}$ is then 15’s “support ring”. $\text{ct} : \mathcal{R}_\mathfrak{o} \rightarrow \mathfrak{o}$ is a ring homomorphism with kernel Π . 15’s symbols are renamed as follows.

15’s symbol	Here
$\mathcal{T}, \mathcal{T}_0, \mathcal{T}_\beta$, topologies	$\mathfrak{T}, \mathfrak{T}_0, \mathfrak{T}_\beta$.
$\mathcal{R}_{\mathfrak{o}, \Gamma} = \mathfrak{o} \oplus \Pi_\Gamma$	$\mathcal{R}_\mathfrak{o} = \mathfrak{o} \oplus \Pi$, with Γ densely ordered.
$N = \overline{\{0\}}$	$\overline{\{0\}}$, the closure of zero; $\mathbb{N}$ stays the natural numbers.
D, E (sets of exponents); D_U, D_V	$A, B; A_U, A_V$.
$R_\Gamma, A, \mathfrak{m}; \Pi_\mathbb{C}, A_\mathbb{C}$	$\mathcal{R}_\Gamma, \mathcal{A}_\Gamma, \mathfrak{m}_\Gamma; \Pi_\mathbb{C}, \mathbb{C} \oplus \Pi_\mathbb{C}$.
$\rho = \sigma _k$	α , as in theorem 22.1 .
$\mathcal{U}_{R_\Gamma}; h(g), a_\delta$	$\mathcal{U}_{\mathcal{R}_\mathbb{Z}}(\Gamma); h_u(g), a_{u, \delta}$ of (3.1).
$\langle f, h \rangle$; summation functional; K^Σ	$\langle f, h \rangle_0$; strong k -linear functional (section 25); the space of these.
$\Lambda; D = \{g : \lambda(t^g) \neq 0\};$ D_a	$\lambda_{\mathbf{On}}$ of (25.2); $S_\lambda; S_a$.
$\sigma_{c, r}$	$M_{\chi, \tau}$ with $\chi(q) = c^q, \tau(q) = rq$.
$\mathbf{No}[i], \mathbf{Oz}[i]; c(z)$	$\mathbf{No}(i), \mathbf{Oz}[i]; \bar{z}$.
$\text{Fin}(J)$	$\text{Fin}(J)$, the finite subsets of J directed by inclusion.

The collisions are resolved here only, and the tempting false readings are wrong:

- 15’s $\mathcal{T}$ is a topology. In [part III](#) $\mathcal{T}_{d, \delta}$ is the Taylor automorphism, T a linear map with adjoint $T^\dagger$, $T_{< \gamma}$ a truncation and T^1 a flow; none of them is a topology, and the topologies of this part are written $\mathfrak{T}$.
- 15’s N is the closure of zero, not the natural numbers.

- 15's sets D and E are not derivations and not the flow $\mathcal{E}$; the letters A, B used instead are sets of exponents, not the rings $\mathcal{A}_k, \mathcal{A}_\Gamma$.
- The neighbourhoods U, V, W of zero are local to this part: V is not the span V_A of [theorem 7.4](#) or a valuation ball $\mathcal{V}_\lambda$, and U is neither $\mathcal{U}$ nor a monomial U^g of [part VI](#).
- 15's "Automorphisms invisible to every compatible omnific topology" is [corollary 39.10](#). It is *not* [corollary 24.4](#), which describes algebraically what an automorphism of $\mathbf{Oz}$ does inside Π ; the topological statement says that no compatible topology can see that action at all.
- 15's $\mathbf{No}[i]$ is the field $\mathbf{No}(i) = \mathbf{No}(i)$ of this report, the same field. Its Hahn-field macro prints $k((t^\Gamma))$, as here; it has no symbol for the relative Hahn hull $\mathcal{H}(A)$ of [theorem 7.4](#).

For proper-class rings the topological statements are read with class-coded neighbourhood systems: a class of indices and one class relation describing membership in each basic neighbourhood. No class whose members are arbitrary proper-class open sets is formed; proofs use individual neighbourhoods, countable sequences and set-supported normal forms [15], in the sense of [section 1.4](#).

39 Hahn-compatible ring topologies

Definition 39.1 (Hahn-compatible ring topology [15]). A *Hahn-compatible ring topology* on a Hahn field, or on one of its subrings A , is a topology $\mathfrak{T}$ making addition, additive inversion and multiplication continuous, such that for every Hahn-summable family $(f_j)_{j \in J}$ of elements of A whose sum lies in A ,

$$\sum_{j \in F} f_j \longrightarrow \sum_{j \in J} f_j \quad (F \in \text{Fin}(J)).$$

Hausdorffness is not assumed; multiplication is required to be *jointly* continuous; the coefficients need not carry their usual topology, and no continuity of ct is assumed.

A much weaker consequence drives the negative results. If $g_0 < g_1 < \dots$ and the monomials at^{g_n} and their Hahn sum lie in A , then

$$at^{g_n} \longrightarrow 0. \tag{39.1}$$

Indeed the partial sums over $\{0, \dots, n\}$ form a cofinal subnet and converge to the sum; their consecutive differences are the terms [15]. So the obstructions below are already forced by countable monomial series.

Lemma 39.2 (Two opposite well orders cannot cover a dense interval [15]). *Let I be a nonempty open interval of a densely ordered set. There are no A, B with $I = A \cup B$ such that A contains no strictly increasing sequence and B contains no strictly decreasing sequence.*

Proof. If B met every nonempty open subinterval of I , density would give recursively a strictly decreasing sequence in B : after $b_n \in B \cap I$, the interval between the left end of I and b_n is nonempty and open. So some nonempty open subinterval $I' \subseteq I$ misses B , hence lies in A ; but I' contains a strictly increasing sequence. $\square$

Lemma 39.3 (Monomial collapse [15]). *Let k be any field, $\mathfrak{o} \subseteq k$ a unital subring and Γ a densely ordered abelian group. If a ring topology on $\mathcal{R}_{\mathfrak{o}}$ satisfies (39.1) for every $a \in k$ and every strictly increasing sequence of negative exponents, then every at^γ with $\gamma < 0$ lies in the closure of zero.*

Proof. Let $a \neq 0$ and $\gamma < 0$, and suppose a neighbourhood W of 0 misses at^γ . Joint continuity gives neighbourhoods U, V of 0 with $UV \subseteq W$. On the interval $I = (\gamma, 0)$ put

$$A = \{x \in I : at^x \notin U\}, \quad B = \{x \in I : t^{\gamma-x} \notin V\}.$$

By (39.1) A contains no strictly increasing sequence, and B none strictly decreasing, since $x \mapsto \gamma - x$ reverses order and maps I into the negative exponents. For $x \in I$ both monomials lie in $\mathcal{R}_\mathfrak{o}$ and $(at^x)t^{\gamma-x} = at^\gamma \notin W$, so $x \in A \cup B$, contradicting lemma 39.2. Hence every neighbourhood of 0 contains at^γ , which in a topological group is membership in $\overline{\{0\}}$. $\square$

Only countable monomial null sequences are used: no metric, first countability or description of the neighbourhoods [15].

Theorem 39.4 (Topological collapse onto the constants [15]). *Let k be a field, $\mathfrak{o} \subseteq k$ a unital subring and Γ a densely ordered abelian group. The Hahn-compatible ring topologies on $\mathcal{R}_\mathfrak{o} = \mathfrak{o} \oplus \Pi$ are exactly the pullbacks*

$$\mathfrak{T} = \text{ct}^{-1}(\mathfrak{T}_0) = \{\text{ct}^{-1}(O) : O \in \mathfrak{T}_0\} \quad (39.2)$$

of the ring topologies $\mathfrak{T}_0$ on $\mathfrak{o}$, and $\mathfrak{T}_0$ is determined by $\mathfrak{T}$. In particular:

- (i) every such topology has $\Pi \subseteq \overline{\{0\}}$;
- (ii) none is Hausdorff when $\Pi \neq 0$;
- (iii) the finest one is the pullback of the discrete topology of $\mathfrak{o}$;
- (iv) $\overline{\{0\}}^\mathfrak{T} = \text{ct}^{-1}(\overline{\{0\}}^{\mathfrak{T}_0})$, and the separated quotient of $(\mathcal{R}_\mathfrak{o}, \mathfrak{T})$ is that of $(\mathfrak{o}, \mathfrak{T}_0)$.

Proof. Let $\mathfrak{T}$ be compatible. The closure $\overline{\{0\}}$ is a closed ideal: a closed subgroup, and multiplication by a fixed element is continuous and fixes 0. By lemma 39.3 every negative monomial lies in $\overline{\{0\}}$. Every $p \in \Pi$ is the Hahn sum of its terms; the finite partial sums lie in $\overline{\{0\}}$ and converge to p , so $p \in \overline{\{0\}}$. Thus $\Pi \subseteq \overline{\{0\}}$.

In a topological group every open set O is invariant under translation by $\overline{\{0\}}$: for $x \in O$ choose a neighbourhood V of 0 with $x + V \subseteq O$; as $\overline{\{0\}} \subseteq V$, $x + \overline{\{0\}} \subseteq O$. So every open set is a union of Π -cosets, that is $O = \text{ct}^{-1}(O_0)$ for a unique $O_0 \subseteq \mathfrak{o}$. The sets O_0 form the quotient ring topology on $\mathcal{R}_\mathfrak{o}/\Pi \cong \mathfrak{o}$, which is also the subspace topology on the constants $\mathfrak{o} \subseteq \mathcal{R}_\mathfrak{o}$ (since $O \cap \mathfrak{o} = O_0$); this gives joint continuity of the induced operations, uniqueness and (39.2).

Conversely, the pullback of a ring topology along the ring homomorphism ct is a ring topology. In a Hahn-summable family only finitely many members have a nonzero constant coefficient, so the constant terms of the finite partial sums are eventually equal to the constant term of the sum; hence the partial sums converge in every pullback topology. (i)–(iv) follow from (39.2). $\square$

The theorem already applies to the set-sized ring $\mathbb{Z} \oplus \Pi$ with $\Gamma = \mathbb{Q}$. So its force is not a proper-class cardinality obstruction of the kind that governs set-sized quotients (`osq:thm:universal` of the quotient report); it comes from a dense exponent interval, formal summation and jointly continuous multiplication together [15].

Corollary 39.5 (A universal separated recipient [15]). *Let B be a Hausdorff topological ring and $\varphi : \mathcal{R}_\mathfrak{o} \rightarrow B$ a ring homomorphism taking every Hahn sum to the limit of the corresponding finite partial sums. Then φ factors uniquely through $\text{ct} : \mathcal{R}_\mathfrak{o} \rightarrow \mathfrak{o}$.*

Proof. The initial topology of φ is a Hahn-compatible ring topology, and its closure of zero is $\ker \varphi$ because B is Hausdorff. By [theorem 39.4](#), $\Pi \subseteq \ker \varphi$. $\square$

The hypothesis is on the treatment of sums. Arbitrary ring maps from a set-sized support ring need not factor through ct : the identity does not [15].

Remark 39.6 (The congruence topologies are instances [merge]). The collection already has compatible topologies on $\mathbf{Oz}$. The finite-congruence topology and, for an ordinary prime p , the p -adic topology are pulled back along ct from the profinite and the p -adic topologies of $\mathbb{Z}$ (`odg:eq:profinite` of the Diophantine report and the paragraph after `osq:thm: completions` of the quotient report, both recording that they are not Hausdorff and that $\Pi = \{0\}$). The Lean modules `Surreal/Foundations/OmnificCongruenceTopology.lean` and `Surreal/Foundations/OmnificSeparationTopology.lean` prove that Π is the closure of zero, that neither topology is T_2 , and that ct induces each topology from the corresponding topology of $\mathbb{Z}$ (`isInducing_omnificConstantCoeff_congruence` and its p -adic analogue; see the rows of `docs/FORMALIZATION.md` following `odg:eq:profinite`). By the converse half of [theorem 39.4](#) these topologies are Hahn-compatible, and by [theorem 39.7](#) below their failure to be Hausdorff is forced: every Hahn-compatible ring topology on $\mathbf{Oz}$ is such a pullback. The Lean results concern those two topologies; nothing of this part is formalized.

Theorem 39.7 (Omnific topologies [15]). *Every Hahn-compatible ring topology on $\mathbf{Oz}$ is the pullback along $\text{ct} : \mathbf{Oz} \rightarrow \mathbb{Z}$ of a unique ring topology on $\mathbb{Z}$, and every Hahn-compatible ring topology on $\mathbf{Oz}[i]$ is the pullback of a unique ring topology on $\mathbb{Z}[i]$. No such topology on either ring is Hausdorff.*

Proof. $\mathbf{No}$ is divisible, hence densely ordered. The proof of [theorem 39.4](#) is elementwise, with $(k, \mathfrak{o}) = (\mathbb{R}, \mathbb{Z})$ or $(\mathbb{C}, \mathbb{Z}[i])$; the descended topology is a topology on the set $\mathbb{Z}$ or $\mathbb{Z}[i]$, and no proper-class sum is formed. $\square$

Proposition 39.8 (Countable monomial tests suffice for the full omnific rings [15]). *In [theorem 39.7](#) it suffices to require (39.1) for countable strictly increasing sequences of negative exponents and fixed coefficients: every ring topology on $\mathbf{Oz}$ or $\mathbf{Oz}[i]$ with this property is a pullback along ct , hence Hahn-compatible.*

Proof. [Lemma 39.3](#) still puts every negative monomial into the closed ideal $\overline{\{0\}}$. For $p \in \Pi$, the support of p is a set of negative surreals; choose a positive surreal δ below every $-g$, $g \in \text{supp } p$, by the cut $\{0\} \mid \{-g : g \in \text{supp } p\}$. Then $t^\delta p \in \Pi$ and $p = t^{-\delta}(t^\delta p) \in \overline{\{0\}}$, an ideal. This replaces the convergence of a possibly uncountable normal form; the second paragraph of the proof of [theorem 39.4](#) finishes. $\square$

The positive buffer δ is a property of the class $\mathbf{No}$. For $\Gamma = \mathbb{Q}$ the support $\{-1/n : n \geq 1\}$ has none, so the proof does not transfer unchanged to set-sized groups [15]; whether the conclusion does is part of [question 42.2](#).

Corollary 39.9 (The surreal and surcomplex fields collapse [15]). *The only Hahn-compatible ring topology on $\mathbf{No}$ or on $\mathbf{No}(i)$ is the indiscrete topology; already the countable condition (39.1) forces it. In particular no Hausdorff topological-field structure on either field realizes every Hahn sum as a topological limit.*

Proof. [Lemma 39.3](#) applies inside either field and puts $t^{-1} \neq 0$ into the ideal $\overline{\{0\}}$, which is therefore the whole field; a topological group in which 0 is dense is indiscrete. The indiscrete topology is trivially compatible. $\square$

Corollary 39.10 (Automorphisms are topologically invisible [15]). *For every ring automorphism σ of $\mathbf{Oz}$ and every Hahn-compatible ring topology on $\mathbf{Oz}$, the elements x and $\sigma(x)$ have the same neighbourhoods. So σ is a homeomorphism and acts trivially on the separated quotient.*

Proof. σ fixes $\mathbb{Z}$ and preserves $\Pi = \bigcap_n n\mathbf{Oz}$ (lemma 2.3), so $\text{ct} \circ \sigma = \text{ct}$ on $\mathbf{Oz}$; by theorem 39.7 every open set is $\text{ct}^{-1}(O_0)$. $\square$

Algebraically the same automorphisms are large and act on every Archimedean scale (corollary 24.4); every compatible topology is blind to all of it.

Lemma 39.11 (Covers of an ordered group by opposite well orders [15]). *Let $\Gamma \neq 0$ be a set-sized ordered abelian group with $\Gamma = A \cup B$, where A is reverse well ordered and B is well ordered. Then $\Gamma \cong \mathbb{Z}$ as an ordered group.*

Proof. $A \neq \emptyset$, since Γ is not well ordered ($-n\delta$, $n \geq 1$, descends for $\delta > 0$); so A has a greatest element d , and $(d, \infty) \subseteq B$ is well ordered. Translating $\Gamma^{\geq 0}$ into (d, ∞) by an element above d shows that $\Gamma^{\geq 0}$ is well ordered. Let ϵ be the least positive element. For $g > 0$ some n has $g < n\epsilon$, since otherwise $g - n\epsilon$ would descend in $\Gamma^{\geq 0}$; for the least such n , $0 \leq g - (n-1)\epsilon < \epsilon$ forces $g = (n-1)\epsilon$. So $\Gamma = \mathbb{Z}\epsilon$. $\square$

Theorem 39.12 (Exact boundary for full Hahn fields [15]). *Let Γ be a set-sized ordered abelian group and k a field. The full Hahn field $k((t^\Gamma))$ admits a Hausdorff Hahn-compatible ring topology if and only if $\Gamma = 0$ or $\Gamma \cong \mathbb{Z}$ as an ordered group. More strongly, if Γ is neither, every ring topology on $k((t^\Gamma))$ in which every strictly increasing sequence of monomials t^{g_n} tends to 0 is indiscrete.*

Proof. Suppose such a topology is not indiscrete. Its closure of zero is a proper ideal of a field, hence 0. Choose a neighbourhood W of 0 with $1 \notin W$ and neighbourhoods U, V of 0 with $UV \subseteq W$, and put $A_U = \{g : t^g \notin U\}$, $A_V = \{g : t^g \notin V\}$. Neither contains a strictly increasing sequence, so both are reverse well ordered. Since $t^g t^{-g} = 1 \notin W$, $\Gamma = A_U \cup (-A_V)$, and $-A_V$ is well ordered; lemma 39.11 gives $\Gamma \cong \mathbb{Z}$ if $\Gamma \neq 0$.

Conversely, for $\Gamma = 0$ take the discrete topology of k . For $\Gamma = \mathbb{Z}\epsilon$ put $s = t^\epsilon$ and take the s -adic valuation topology of $k((s))$, a Hausdorff field topology. A well-ordered subset of $\mathbb{Z}$ is bounded below and has finitely many elements below any integer; with pointwise finiteness, for each N only finitely many members of a summable family have a term below N , so the net of finite partial sums converges to the Hahn sum. $\square$

Remark 39.13 (Scope [15]). The cyclic half is an existence statement: the compatible topologies of $k((s))$ are not classified. The theorem separates cyclic from merely discrete groups: lexicographic $\mathbb{Z}^2$ has a least positive element and is still excluded. The field theorem needs neither density nor divisibility but uses both t^g and t^{-g} , so it does not settle the negative-support ring for non-dense groups (question 42.1); for $\Gamma = \mathbb{Z}$ that ring is $\mathfrak{o} + t^{-1}k[t^{-1}]$, whose summable families are finite up to zero members, so its discrete topology is compatible and Hausdorff.

Nothing here conflicts with the order and valuation topologies, with formal Hahn calculus or with strong automorphisms: familiar topologies do not realize every Hahn sum. In $\mathbb{R}((t^\mathbb{Q}))$ the monomials $t^{-1/n}$ have a Hahn sum but do not tend to 0 in the valuation topology; the foundations report (docs/foundations-and-computation/foundations, found:sub:tsum) records such failures for particular topologies, including the rank-one found:ex:boundedrankone over $\mathbb{C}((t^\mathbb{Q}))$, and so do remarks 21.7 and 34.4 here. Theorem 39.12 is the general form [merge]: outside $\Gamma \in \{0, \mathbb{Z}\}$ no Hausdorff ring topology at all does better. Nor is there a conflict with finiteness spaces [3]: a monoid object for a summation-sensitive tensor product

does not have jointly continuous multiplication for a Cartesian product topology. And the theorems do not say that these rings lack Hausdorff ring topologies: the discrete topology always exists; it fails to realize the nontrivial Hahn sums [15].

As a consequence, a unital homomorphism φ from **No** (or **No**(i)) into a nonzero Hausdorff topological ring cannot take every Hahn sum to the corresponding limit: the initial topology of φ would be compatible and would separate 1 from 0, against [corollary 39.9](#). For **Oz** and **Oz**[i] every ring topology on $\mathbb{Z}$ or $\mathbb{Z}[i]$ gives a compatible topology by pullback, which accounts at once for the existence of non-Hausdorff compatible topologies and for their loss of all infinite-scale information. Topology remains usable if one keeps a valuation topology with a smaller class of convergent sums, keeps all sums with only separately continuous multiplication ([section 41](#)), or uses a summation-aware tensor or bornological framework; these choices preserve different structures and should be stated [15].

Remark 39.14 (Two proof chains [15]). The topological theorems do not use the automorphism classifications of this report: convergence of monomial sums gives excluded exponent sets with opposite well-order properties, then the dense-interval or cyclic-group obstruction, then the collapse. For the omnific rings one uses only the support description of **Oz** and its constant-term quotient; the countable strengthening uses the positive buffer of **No**. The algebraic chain (well-ordered supports, then scalar detection, then covariance implies strongness, with the reconstruction of Π , $\mathcal{A}_k$, $\mathbb{R}$ and $\mathfrak{m}$) is that of [part III](#). Together: the omnific ring algebraically forces preservation of all Hahn sums ([theorem 23.2](#)), while every jointly continuous ring topology realizing those sums forgets Π .

40 The algebraic half: a disjoint-row variant and four additions

15's detection theorem is [theorem 21.4](#). Its proof selects disjoint rows of an incidence system; this is a variant of 14's induced matching ([lemma 34.1](#)) that selects family members rather than exponents, and it yields prescribed pairings.

Lemma 40.1 (Disjoint rows [15]). *Let $(F_i)_{i \in I}$ be infinitely many nonempty finite subsets of $\mathbb{N}$ such that each $n \in \mathbb{N}$ lies in only finitely many F_i . Then there are distinct $i_0, i_1, \dots$ with the F_{i_r} pairwise disjoint.*

Proof. Choose one row. Its finitely many elements meet only finitely many other rows; discard those and repeat. At each stage finitely many rows have been discarded and infinitely many remain. $\square$

Variant of the third route in [theorem 21.4](#) [15]. We show that countably supported tests suffice. Assume that every y with countable support has $\langle x_j, y \rangle_0 \neq 0$ for only finitely many j . Monomial tests give pointwise finiteness. If the union S of the supports is not well ordered, choose $\gamma_0 > \gamma_1 > \dots$ in S and put $F_j = \{n : [t^{\gamma_n}]x_j \neq 0\}$. Each F_j is finite (a well-ordered support meets a descending sequence finitely), each n lies in finitely many F_j , and infinitely many F_j are nonempty. [Lemma 40.1](#) gives pairwise disjoint nonempty F_{j_r} ; pick $n_r \in F_{j_r}$ and put

$$y = \sum_{r \geq 0} ([t^{\gamma_{n_r}}]x_{j_r})^{-1} t^{-\gamma_{n_r}}, \quad (40.1)$$

a Hahn series with countable support. By disjointness, $[t^{\gamma_{n_s}}]x_{j_r} = 0$ for $s \neq r$, so $\langle x_{j_r}, y \rangle_0 = 1$ for every r , contradicting the assumption for this y . $\square$

With $a_{n,j} = [t^{\gamma_n}]x_j$, disjoint rows give $a_{n_r, j_r} \neq 0$ and $a_{n_s, j_r} = 0$ for $s \neq r$: an induced matching in the sense of [lemma 34.1](#), found by selecting the members j , with the stronger

property that their supports along the sequence are disjoint, rather than the exponents n . With coefficients 1 instead of the inverses in (40.1) each pairing would be the single entry $a_{n_r, j_r} \neq 0$, 14's all-ones witness. 15's normalization to 1 is what gives interpolation. The detector (40.1) has countable support but not coefficients in $\{0, 1\}$, so on its own the variant shows that countably supported tests with arbitrary coefficients suffice; the binary form of (iii) is 12's, 13's and 14's. The route works over every field, including finite fields and positive characteristic [15].

Corollary 40.2 (Sparse interpolation inside a bad family [15]). *Let (x_j) be pointwise finite with a union of supports that is not well ordered. There are distinct j_r and a countable strictly increasing $\Lambda \subseteq \Gamma$ such that for every sequence $(c_r) \in k^{\mathbb{N}}$ some y supported in Λ has $\langle x_{j_r}, y \rangle_0 = c_r$ for all r .*

Proof. Use the rows and exponents of (40.1), with numerators c_r in place of 1; the set $\Lambda = \{-\gamma_{n_r}\}$ does not depend on (c_r) . $\square$

Example 40.3 (The naive detector fails [15]). Putting coefficient 1 at *every* opposite exponent of a descending sequence does not detect non-summability. For $x_n = t^{-n} - t^{-(n+1)}$, $n \geq 1$, the family is not summable (its supports descend), but $y = \sum_{m \geq 1} t^m$ gives $\langle x_n, y \rangle_0 = 1 - 1 = 0$ for every n . A selection is needed: the members x_n with n odd have pairwise disjoint supports, and selecting them, as in lemma 40.1 or 14's matching, gives $y = \sum_{r \geq 0} t^{2r+1}$ with $\langle x_{2r+1}, y \rangle_0 = 1$.

In example 34.3 the cancellation is between members sharing an exponent, and the all-ones test on all exponents already works; here it is between adjacent exponents of one member, and the all-ones test on all exponents fails. 15's Proposition 4.5 is the family of example 21.6: $t^{-n\delta}$ is not summable, a finite-support test pairs nonzero with finitely many members, and $\sum_{n \geq 1} t^{n\delta}$ pairs to 1 with all. So finite-support tests do not suffice and countable ones do. This concerns the support of a test, not whether countable families determine every uncountable sum [15].

Proposition 23.1 transports the *full* constant term. On bounded elements ct is the standard part, but on $p + r + \varepsilon$ with $p \in \Pi$ the standard part is undefined while $\text{ct} = r$, so the invariant Π is essential; invariance of the residue on bounded elements would not suffice [15].

Corollary 40.4 (The Gaussian ring with its involution [15]). *Every unital ring automorphism of $\mathbf{Oz}[i]$ that commutes with conjugation extends uniquely to a field automorphism of $\mathbf{No}(i)$ commuting with conjugation, and the extension is strongly additive with strongly additive inverse.*

Proof. For $x + iy \in \mathbf{No}(i)$, `odg:thm:fractions` gives one $d \in \mathbf{Oz} \setminus \{0\}$ with $dx, dy \in \mathbf{Oz}$, so $\text{Frac } \mathbf{Oz}[i] = \mathbf{No}(i)$ and the automorphism extends uniquely to fractions; commutation with conjugation passes from numerators and denominators, and the extension preserves $\mathbf{Oz}[i]$. Theorem 30.1 gives the form and strongness. $\square$

For the bare pair $(\mathbf{No}(i), \mathbf{Oz}[i])$ strongness is not concluded. The proved alternatives are preservation of the valuation ring, commutation with conjugation, or covariance of the full constant term; no inference from “the constants are invariant” to “ ct is invariant” is made (question 32.1) [15].

Proposition 40.5 (Class coefficient description of the strong dual [15]). *An $\mathbb{R}$ -linear strong functional λ on $\mathbf{No}$ is determined by the class function $a(g) = \lambda(t^g)$, whose support $S_a = \{g : a(g) \neq 0\}$ contains no strictly increasing sequence, and*

$$\lambda\left(\sum_g c_g t^g\right) = \sum_g c_g a(g), \quad (40.2)$$

a finite sum. Conversely every class function $a : \mathbf{No} \rightarrow \mathbb{R}$ whose support contains no strictly increasing sequence defines by (40.2) a strong $\mathbb{R}$ -linear functional λ_a . It is $\text{ct}(h \cdot)$ for a surreal h exactly when S_a is a set, and then $h = \sum_g a(g)t^{-g}$.

Proof. A strictly increasing (g_n) in S_a would make (t^{g_n}) a summable family with infinitely many nonzero values of λ ; strongness on the normal-form family of an element gives (40.2). Conversely, a set with no strictly increasing sequence meets every well-ordered set finitely, since an infinite subset of a well-ordered set contains one; so the sum in (40.2) is finite, and for a summable family the union of the supports meets S_a in finitely many exponents, each in finitely many members, which gives the finiteness and the interchange of the two sums. If S_a is a set it is reverse well ordered, h is a surreal, and $\text{ct}(hf) = \lambda_a(f)$; a representing h has support exactly $-S_a$, which must be a set. $\square$

$\lambda_{\mathbf{On}}$ of theorem 25.6 is λ_a for the indicator a of $-\mathbf{On}$. A support “with no strictly increasing sequence” is the same as one meeting every well-ordered set finitely, so the proposition describes the strong $\mathbb{R}$ -linear dual of $\mathbf{No}$ exactly by class coefficient functionals meeting every well-ordered set finitely, each an individual class function in the sense of section 1.4; no class of such functionals is formed. This answers the first clause of question 32.11; the second, which strong operators have transposes that stay set-supported, is 15’s Question 9 and stays open (section 42.1). The proof of theorem 21.4 uses only countably supported tests and never represents a class functional, so this enlarged dual places no restriction on theorem 23.2; it only blocks the overgeneralization of proposition 25.1 and theorem 25.2 to $\mathbf{No}$ [15].

41 A pairing topology that realizes exactly the Hahn sums

In this section Γ is a set, $K = k((t^\Gamma))$, and k carries the discrete topology. Let $\mathfrak{T}_\beta$ be the weak topology generated by the functionals $f \mapsto \langle f, h \rangle_0$, $h \in K$; a basis of neighbourhoods of 0 is

$$U(h_1, \dots, h_m) = \{f : \langle f, h_1 \rangle_0 = \dots = \langle f, h_m \rangle_0 = 0\}. \quad (41.1)$$

Nondegeneracy of the pairing makes $(K, \mathfrak{T}_\beta)$ a Hausdorff topological vector space over the discrete field k . Over $\mathbb{R}$ or $\mathbb{C}$ the usual topology of the coefficients is not used, and $\mathfrak{T}_\beta$ is not identified with the Hahn-field topologies of Flynn and Shamseddine [8] [15].

Theorem 41.1 (Unordered sums in the pairing topology [15]). *For a family $(f_j)_{j \in J}$ in K , the net $s_F = \sum_{j \in F} f_j$, $F \in \text{Fin}(J)$, converges to f in $\mathfrak{T}_\beta$ if and only if (f_j) is Hahn summable with sum f .*

Proof. In a discrete field a net of finite partial sums of scalars converges if and only if all but finitely many summands vanish: beyond some F_0 the net is constant, and comparing F_0 with $F_0 \cup \{j\}$ kills every summand outside F_0 . Applied to $\langle f_j, h \rangle_0$ for each h , convergence of s_F gives the finiteness condition of theorem 21.4, hence summability, and the limit identity with nondegeneracy identifies the sum. Conversely a Hahn sum gives eventually constant scalar nets with the right limits (21.2), which is convergence in the initial topology. $\square$

Corollary 41.2 (Continuous linear maps [15]). *For set-sized full Hahn fields K, L over k , a k -linear map $K \rightarrow L$ is continuous for the pairing topologies if and only if it is strongly k -linear.*

Proof. A continuous map preserves convergent nets of finite partial sums (theorem 41.1). A strong map has a constant-term adjoint (theorem 25.2), so each target test functional composed with it is a source test functional. $\square$

In particular multiplication by a fixed a is continuous, as $\langle af, h \rangle_0 = \langle f, ah \rangle_0$, and the isomorphisms of [theorem 22.1](#) are homeomorphisms, the coefficient isomorphism being one of discrete fields [15].

Theorem 41.3 (Multiplication is not jointly continuous [15]). *If $\Gamma \neq 0$, multiplication $K \times K \rightarrow K$ is not jointly continuous for $\mathfrak{T}_\beta$; so $\mathfrak{T}_\beta$ is not a ring topology, although it realizes exactly the Hahn sums and multiplication is separately continuous.*

Proof. $\text{ct} = \langle \cdot, 1 \rangle_0$ is continuous and $\text{ct}^{-1}(0)$ is open. Joint continuity at $(0, 0)$ would give basic neighbourhoods U, V of the form (41.1) with $\text{ct}(uv) = 0$ for $u \in U, v \in V$, so $U \subseteq V^\perp = \{x : \langle x, v \rangle_0 = 0 \text{ for all } v \in V\}$. By nondegeneracy $x \mapsto \langle x, \cdot \rangle_0$ injects $V^\perp$ into $(K/V)^*$, which is finite dimensional since V has finite codimension. But U has finite codimension in K , which is infinite dimensional ($t^{n\delta}, n \geq 0$, are independent for $\delta > 0$), so U is infinite dimensional, a contradiction. $\square$

This particular obstruction is weaker than [theorem 39.12](#), which excludes every Hausdorff compatible ring topology outside the cyclic case; it shows that joint continuity cannot be weakened to separate continuity in that theorem. A summation-detecting topology does not turn Hahn constructions into topological-field constructions [15].

42 Questions of source 15, and stale statements

42.1 Status of the source's questions

15 poses twelve questions, proposed as research directions, not as established historical conjectures. Six are merged into questions of this report, with 15 added to their sources:

- Question 4 (Gaussian automatic strongness without extra structure): [question 32.1](#), open. 15 adds that the topological collapse does not settle it: a compatible ring topology cannot reconstruct the missing valuation, since on $\mathbf{No}(i)$ every one is indiscrete ([corollary 39.9](#)).
- Question 5 (proper Hahn subfields and detector closure): [question 37.1](#). It is *partly addressed* by 14's countable detector property ([definition 35.1](#) and [proposition 35.2](#)), a sufficient condition, and [example 35.3](#) shows that some condition is needed; the necessary-and-sufficient condition and its effect on constant-term-compatible embeddings stay open.
- Question 6 (countable versus unrestricted preservation in support-bounded fields, singular bounds): [question 32.12](#), open.
- Question 9 (class summation duals and adjoints): [question 32.11](#). Its first clause is answered by [proposition 40.5](#); 15's question is the second clause, open.
- Question 10 (generation of the full stabilizer): [question 11.4](#), open.
- Question 12 (a verified core and transfer layer): [question 32.18](#).

Six are new.

Question 42.1 (Nondense truncated support rings [15]). For which nonzero ordered abelian groups Γ does $\mathfrak{o} \oplus \Pi$ admit a Hausdorff Hahn-compatible ring topology? Dense groups are excluded ([theorem 39.4](#)) and $\Gamma = \mathbb{Z}$ admits the discrete topology ([remark 39.13](#)); the first test is lexicographic $\mathbb{Z}^2$.

The full-field obstruction does not settle it, because inverse monomials leave the negative-support ring. An answer should identify an intrinsic order property of the negative cone; the proof of monomial collapse uses bounded intervals, not the whole group [15].

Question 42.2 (Minimal forbidden summation families [15]). For a fixed dense exponent group, what is the smallest specified collection of countable monomial sums whose convergence already forces the constant-term collapse? Does an explicitly listed countable collection suffice for $\mathbb{Z} \oplus \Pi$ with $\Gamma = \mathbb{Q}$, or is more needed when nothing countable is assumed of the topology?

The present proof chooses its sequences after examining the neighbourhoods; removing that dependence would make the obstruction effective and show how much summation data must be named [15]. [Proposition 39.8](#) answers the question for the full omnific rings, where countable monomial sums suffice; its buffer argument fails for $\Gamma = \mathbb{Q}$ [merge].

Question 42.3 (Maximal topologically realizable summation laws [15]). Fix a Hausdorff valuation topology on a Hahn field. Classify the largest subfamilies of the Hahn sums that converge in it and are closed under reindexing, subfamilies, finite products and multiplication by fixed elements. How do they change when the valuation is coarsened?

The criterion that only finitely many summands contribute below each valuation threshold is a starting point, not a classification of alternative topologies and compatible summation structures [15].

Question 42.4 (A non-Cartesian multiplication framework [15]). Can the algebra of surreal Hahn sums be organized in a summation-sensitive tensor category in which multiplication is a continuous morphism, scalar duality has the correct class-size qualification, and the strong automorphisms are exactly the invertible morphisms? Which completion or boundedness conditions are needed?

Finiteness spaces and formal summation algebras are precedents [3, 2]; the problem is a precise full-surreal implementation that does not mistake tensor continuity for joint continuity in a product topology [15].

Question 42.5 (Separating a chosen family of scales [15]). Given a set $E \subseteq \Pi$, which parts of the Hahn summation law must be discarded to obtain a Hausdorff ring topology separating every nonzero member of E from 0? Is the trade-off governed by a support invariant of E ?

The unrestricted law cannot separate even one negative monomial; a sharp restricted result would serve asymptotic models that keep selected scales [15].

Question 42.6 (Definability beyond automorphism invariance [15]). In a language with sorts for admissible set-indexed families, how much of Hahn summation can be recovered from the field and the omnific predicate? Is there a useful definability theorem strictly stronger than automorphism invariance ([theorem 23.2](#)) and weaker than naming the Conway monomial section or full set theory?

The topological quotient and the nondefinability of monomials ([theorem 9.1](#)) exclude several naive proposals; an answer must fix the language, the size of its sorts and the interpretation of family parameters [15]. Compare [question 11.5](#) and `dsn:q:languages` [merge].

42.2 Stale and checked statements

- *Novelty, checked.* 15 first answered [question 11.1](#) itself, found 12 and 13 in a later read, and changed its claim: automatic strongness is credited to them and not claimed.

Its remaining claim, the topological classification, is new at the merge: no report of the collection classifies Hahn-compatible ring topologies; the collection had particular topologies only ([remarks 39.6](#) and [39.13](#), [section 28](#)). Its statement that the inspected audits do not contain the classification is true, and the merge found no other report containing it.

- *Checked true.* The two article blobs and the three audit blobs it cites ([section 38.1](#)); its description of [question 11.1](#) as open in the first blob and answered in the second; its credits to the criterion, [theorem 5.1](#) and the shear examples; the finite checks of its suite, rerun for the merge ([section C](#)). No mathematical error was found in 15.
- *Not rechecked.* 15’s descriptions of [8] (whose topologies it does not identify with $\mathfrak{T}_\beta$) and of [3, 2] are reported as 15 gives them.

15’s limitations are listed with the other sources’ in [section B](#).

Part VI

Coefficient recovery in monomial extensions of arbitrary rank

This part records manuscript 16, the second addition of batch 32. Its question is that of [part II](#), asked at arbitrary rank: after adjoining external monomial parameters, of any rank and not necessarily finitely generated, can the ring or field still recover its surreal or omnific coefficients? It can. One existential quartic equation defines the coefficient field in the fraction field of a group algebra ([section 45](#)); a sparse power bound survives every monomial refinement ([section 46](#)); coefficient rigidity, automorphism formulas and finite-rank cancellation follow at arbitrary rank, cancellation failing exactly at infinite rank ([sections 47, 48](#) and [50](#)); and, with a published defect theorem, every field automorphism is monomial exactly when $\text{Hom}(G, \mathbb{Z}) = 0$ ([section 49](#)). The parameters are finite-support external monomials, not Hahn sums. Its labels carry the sub-prefix `opa:cr:`.

43 Source 16, its relation to Part II, and its notation

43.1 The manuscript and its pin

Manuscript 16 is *Recovering Surreal Coefficients from Monomial Extensions: quartic definitions, arbitrary-rank rigidity, and sharp cancellation*, dated 23 September 2026, 24 pages. It is item 08 of batch 32, delivered as the archive `surreal_coefficient_rigidity` in commit `aa268a4` and placed as an addition in commit `7d04483`; it is numbered 16 here. Its checks, their recorded run, its build script, its hash-bearing build record and its source audit are shipped as `code/16-coefficient-recovery-verify.py`, `data/16-coefficient-recovery-verification.json`, `code/16-coefficient-recovery-build.py`, `data/16-coefficient-recovery-build_report.json` and `16-coefficient-recovery-SOURCE_AUDIT.md`; its `article.tex`, PDF, README and checksum list are not shipped (the checksum list and the hashes of the build record were verified at placement).

It pins `1ab41af`, the last commit it observed, which contains [parts I](#) to [III](#) of this report (`20c4c9f`) but not the placement `9d28e28` or the write `d4d72d7` of source 14. It read the catalogue, the audits of sources 11 and 12 and a few guides, not this article. Its “prior

manuscript” is source 02 itself, the base of [part II](#), retrieved separately with 02’s pin `fb5c4b5`. 16 describes 02 accurately and credits it for the root-covered framework, the monomial-divisibility criterion, polynomial and Laurent coefficient rigidity, strong cancellation, the finite-type obstructions and the formal flows; it claims none of these.

43.2 What it generalizes, and what is printed once

16’s numbers are those of its own PDF; its main theorems are lettered A–D.

16	Printed as	Relation to part II
Main Thm. A(iii), Thm. 6.4	theorems 45.4 and 47.1	Generalize theorems 15.1 and 15.3 from $B[\mathbf{T}, \mathbf{U}^{\pm 1}]$ with finitely many variables to $S[U^M]$ for every cancellative torsion-free monoid, of any rank.
Thm. 7.1	theorem 48.1	Generalizes the torus part of theorem 15.5 , with $\mathrm{GL}_q(\mathbb{Z}) = \mathrm{Aut}(\mathbb{Z}^q)$, to $\mathrm{Aut}(G)$ for every torsion-free G .
Thm. 9.1	theorem 50.1	Generalizes theorem 15.4 from $\mathbb{N}^p \times \mathbb{Z}^q$ to monoids of finite rational rank, and to coefficient rings with closed fraction field.
Thm. 9.3, Main Thm. D	theorems 50.3 and 50.4	Exact failure at infinite rank, parallel to the example $R[X_1, X_2] = B[X_2]$ after theorem 15.4 .
Prop. 6.2	theorem 13.1 and corollary 13.3	Reproved and credited; printed once, 16 added.
Def. 6.1; Ex. 6.3	definition 12.1 ; the workspace of section 18	Same, with the same fraction-field warning.
Prop. 4.1	<code>odg:thm:fractions</code> , <code>odg:prop:units</code> , <code>odg:def:lem:units</code>	Reproved (“classical”); cited.
Question 12.8	question 19.3	Merged.
All else	This part	New.

The proofs of [part II](#) use degrees in finitely many variables. That argument does not reach arbitrary rank: an element and its roots can lie in different finitely generated exponent lattices, and no finitely generated target need contain all the witnesses. 16’s substitute is the sparse bound of [theorem 46.2](#), which is uniform under changes of lattice [16].

43.3 Conventions and notation of this part

Rings are commutative and unital, homomorphisms unital, embeddings injective, and domains have characteristic zero unless a counterexample says otherwise. G is a set-sized torsion-free

abelian group and $\mathbf{M}$, $\mathbf{N}$ set-sized cancellative torsion-free commutative monoids, written additively, with group completion $\mathrm{gp}(\mathbf{M})$ and rational rank $\mathrm{rk}_{\mathbb{Q}} \mathbf{M} = \dim_{\mathbb{Q}}(\mathbb{Q} \otimes \mathrm{gp} \mathbf{M})$. The *external monomials* are written U^g :

$$k[U^G] = \left\{ \sum_{g \in S} a_g U^g : S \subseteq G \text{ finite} \right\}, \quad U^g U^h = U^{g+h}, \quad k(U^G) = \mathrm{Frac} k[U^G],$$

and $R[U^{\mathbf{M}}] \subseteq R[U^{\mathrm{gp} \mathbf{M}}]$ is the monoid algebra. A *monomial* is cU^g with $c \neq 0$; $\mathrm{wt} f$ is the number of nonzero coefficients of $f \in k[U^G]$. So $B[\mathbf{U}^{\pm 1}] = B[U^{\mathbb{Z}^q}]$ and $B[\mathbf{T}] = B[U^{\mathbb{N}^p}]$ in the notation of [theorems 15.1](#) and [15.3](#). An element of $k(U^G)$ is a quotient of two finite sums, not a Hahn series [16].

16's symbol	Here
$X^g, k[G],$ $K_k(G) = \mathrm{Frac} k[G]$	$U^g, k[U^G], k(U^G).$
M, N (monoids); H (a second group)	$\mathbf{M}, \mathbf{N}; G'.$
$v(f), d(f),$ $w(f) = d(f) - v(f)$	$\min \mathrm{supp} f, \max \mathrm{supp} f, \mathrm{wd}(f),$ for an auxiliary order on G .
ℓ , the specializing homomorphism	$\lambda.$
$D = T d/dT$	$\vartheta.$
$\mathcal{U}_{\infty}(B); D(G) = \bigcap_n nG$	$\mathrm{Rt}_{\infty}(B); G_{\mathrm{div}}.$
$\mathcal{D}(B), \mathcal{C}(B)$	$\mathfrak{D}(B), \mathfrak{C}(B).$
$\Delta_k(G) =$ $K_k(G)^{\times} / (k^{\times} X^G)$	$\mathrm{Def}_k(G) = k(U^G)^{\times} / (k^{\times} U^G).$
$\sigma \in \mathrm{Aut}(F), \tau \in \mathrm{Aut}(G),$ $\chi \in \mathrm{Hom}(G, F^{\times})$	$\alpha, \tau, \chi.$
$T = \phi^{-1}(S)$ in the cancellation proof	$R_1.$
$R_{\Gamma}(D, k), \mathcal{P}_{\Gamma}(k)$	$\mathcal{R}_{\mathfrak{o}}(k, \Gamma), \Pi_k(\Gamma),$ as in section 12.1 .
$\mathbf{Oz}[i]; \mathbf{No}(i)$	$\mathbf{Oz}[i]; \mathbf{No}(i),$ the same.

The collisions are resolved here only, and the tempting false readings are wrong:

- 16's X^g is an *external indeterminate*. In [part II](#) $X^g = t^{-g} = \omega^g$ is a Hahn monomial of $k((X^{\Gamma}))$. So U^g here is not a surreal monomial: in $\mathbf{No}(U^G)$ with $g \neq 0$, U^g is transcendental over $\mathbf{No}$, while $\omega^g \in \mathbf{No}$ is a coefficient ([example 51.1](#)).
- 16's $K_k(G)$ is the rational group field $k(U^G)$, not the Hahn field $K_{\Gamma} = k((t^{\Gamma}))$ of [part III](#); no Hahn summation is involved.
- 16's $\Delta_k(G)$ is a defect group, not the target group Δ of [part III](#); 16's $D(G)$ is not a derivation; 16's $\mathcal{D}(B)$ and $\mathcal{C}(B)$ are sets of ring elements, not the convex subgroups C_{δ} and $C(a)$ or the operator coefficients c_{δ} ; 16's $\mathcal{U}_{\infty}$ is not the stabilizer $\mathcal{U}$.
- 16's monoid M is not a monomial map $M_{\chi, \tau}$. Its $\tau \in \mathrm{Aut}(G)$ is a group automorphism, with no order, and its χ need not be positive; the automorphisms ([48.1](#)) are the finite-support analogue of $M_{\chi, \tau}$ composed with a coefficient automorphism α , 16's σ .

- 16's v is the minimum of a finite support for an auxiliary order, not the Hahn valuation, and its d is not a derivation of $\mathbb{R}$; its ℓ is not the leading exponent $\ell(x)$ of [part II](#).

Coefficient rings may be the classes **No**, **No**(i), **Oz**, **Oz**[i]; the exponent groups and monoids are sets. An element of a monoid algebra is a finite list of exponents and coefficients; fractions are coded by the sets of minimum-rank representatives (Scott's trick), not by proper-class equivalence classes. No proper-class spectrum, Hahn completion or set of all class automorphisms is formed, and a displayed automorphism group is schematic: each class automorphism is specified by the displayed data, and each admissible choice gives one. The algebraic arguments involve finitely many supports and coefficients at a time; the one step that uses a published set-theoretic theorem for all $g \in G$ at once descends to a set-sized coefficient subfield ([section 49](#)) [16].

44 Finite-support monomial extensions

A torsion-free abelian group admits a compatible total order (embed it in $\mathbb{Q} \otimes G$ and order a basis lexicographically). Fix one; for $0 \neq f \in S[U^G]$ the least and greatest support exponents and the width $\text{wd}(f) = \max \text{supp } f - \min \text{supp } f$ are additive under products in a domain S , since the extreme coefficients of a product are products of extreme coefficients [16].

Lemma 44.1 (Monomial factors and units [16]). *Let S be a domain. If fg is a nonzero monomial in $S[U^G]$, then f and g are monomials. So $S[U^G]^\times = S^\times U^G$, and $S[U^M]^\times = S^\times U^{M^\times}$ with $M^\times = M \cap (-M)$ in $\text{gp } M$.*

Proof. A monomial has width 0, widths are nonnegative and additive, and a finite support of width 0 is a point. If $fg = 1$ the coefficients are units and the exponents cancel. $\square$

Lemma 44.2 (No new algebraic constants [16]). *For every field k , k is relatively algebraically closed in $k(U^G)$. If $k \subseteq L$ and $G \subseteq G'$, then $k(U^G) \cap L^\times U^{G'} = k^\times U^G$ inside $L(U^{G'})$.*

Proof. $v_0(P/Q) = \min \text{supp } P - \min \text{supp } Q$ is a valuation of $k(U^G)$ with residue field k (16's v ; it is not the Hahn valuation). Let $z \neq 0$ be algebraic over k , with minimal polynomial of nonzero constant term. If $v_0(z) > 0$ the constant term is the unique term of least valuation, and if $v_0(z) < 0$ the term of highest degree is; either contradicts the equation, so $v_0(z) = 0$. Let $c \in k$ be the residue of z ; if $z \neq c$, then $z - c$ is algebraic with $v_0(z - c) > 0$, which is impossible. For the second claim, $P/Q = cU^h$ gives $\text{supp } P = h + \text{supp } Q$, so $h \in G$, and a coefficient comparison gives $c \in k$. No closedness of k is used. $\square$

The valuation is auxiliary; no automorphism below is assumed to preserve it.

Lemma 44.3 (One-variable specialization [16]). *Given finitely many elements of $k[U^G]$ and a finite set of exponent combinations, there is a homomorphism λ from the subgroup generated by their supports to $\mathbb{Z}$ that is injective on the prescribed set, so the substitution $U^g \mapsto T^{\lambda(g)}$ keeps prescribed denominators nonzero. Moreover, for $P, Q \in k[U^G] \setminus \{0\}$:*

- if $P/Q \notin k$, λ can be chosen so that the specialized fraction is not constant;*
- if P/Q is not a monomial, λ can be chosen so that the specialized fraction is not a monomial and $\text{wt } P, \text{wt } Q$ are unchanged.*

Proof. The subgroup is finitely generated and torsion-free, so $\cong \mathbb{Z}^d$, and an integer functional outside finitely many hyperplanes separates a finite set: try $(1, m, \dots, m^{d-1})$, each coincidence

excluding finitely many integers m . Include the supports of all denominators. For (i) require injectivity on $\text{supp } P \cup \text{supp } Q$: proportional specializations then force $P = cQ$. For (ii) require injectivity on $\text{supp } P - \text{supp } Q$, which also separates $\text{supp } P$ and $\text{supp } Q$; a specialized quotient cT^j would give $\lambda(\text{supp } P) = j + \lambda(\text{supp } Q)$, every difference $a - b$ in the resulting bijection would map to j and so be one fixed g , and the coefficients would give $P = cU^gQ$. $\square$

The specialization is applied to one identity with its denominators cleared, and chosen to keep them nonzero; it is never a field homomorphism on all of $k(U^G)$ [16].

45 A quartic equation recovers the coefficient field

Lemma 45.1 (Mason–Stothers in characteristic zero [16]). *If nonzero pairwise coprime $A, B, C \in k[T]$, not all constant, satisfy $A + B = C$, then $\max(\deg A, \deg B, \deg C) \leq \deg \text{rad}(ABC) - 1$, where rad is the product of the distinct monic irreducible factors.*

Proof. $W = A'B - AB'$ is nonzero: otherwise $(A/B)' = 0$, so $A/B \in k$ in characteristic zero and coprimality makes all three constant. Since $C = A + B$, also $W = A'C - AC' = C'B - CB'$. A factor of multiplicity m of A , B or C divides W to multiplicity at least $m - 1$, so by coprimality $ABC/\text{rad}(ABC)$ divides W , and $\deg W \leq \deg A + \deg B - 1$ gives the bound for $\deg C$; permute for the others. This is the standard Wronskian proof [1]. $\square$

Lemma 45.2 (The quartic has only constant rational points [16]). *If $a, b \in k(T)$ satisfy $b^2 = 1 + a^4$, then $a, b \in k$.*

Proof. Write $a = P/Q$ in lowest terms. Then $(bQ^2)^2 = P^4 + Q^4$, and a rational function whose square is a polynomial is a polynomial, so $V_0 = bQ^2 \in k[T]$ with $V_0^2 = P^4 + Q^4$. If $P = 0$ we are done; if $V_0 = 0$, coprimality makes P, Q constant. Otherwise P, Q, V_0 are pairwise coprime; if $d = \max(\deg P, \deg Q) > 0$, then $\deg V_0 \leq 2d$ and lemma 45.1 for $P^4 + Q^4 = V_0^2$ gives $4d \leq \deg P + \deg Q + \deg V_0 - 1 \leq 4d - 1$. So $a \in k$, and $b \in k$ by lemma 44.2. $\square$

Theorem 45.3 (Quartic rigidity in every rational group field [16]). *For every characteristic-zero field k and torsion-free abelian group G , every solution of $b^2 = 1 + a^4$ in $k(U^G)$ has $a, b \in k$.*

Proof. If $a \notin k$, choose λ by lemma 44.3(i) keeping a nonconstant and the denominators of a, b nonzero; the identity with cleared denominators specializes to a nonconstant solution in $k(T)$, against lemma 45.2. Then $b \in k$ by lemma 44.2. $\square$

Theorem 45.4 (Uniform coefficient definitions [16]; 16's Main Theorem A). *Let G be a set-sized torsion-free abelian group and $\mathbf{M}$ a set-sized cancellative torsion-free monoid.*

(i) *If F is real closed, or algebraically closed of characteristic zero, then in $F(U^G)$*

$$x \in F \iff \exists y (y^2 = 1 + x^4). \quad (45.1)$$

(ii) *If R is a domain whose fraction field has one of these properties, then in $R[U^{\mathbf{M}}]$*

$$x \in R \iff \exists u \exists v (v \neq 0 \wedge u^2 = (1 + x^4)v^2). \quad (45.2)$$

(iii) *(i) applies to $F = \mathbf{No}, \mathbf{No}(i)$ and (ii) to $R = \mathbf{Oz}, \mathbf{Oz}[i]$. Every embedding of $\mathbf{No}$ or $\mathbf{No}(i)$ into a field $k(U^G)$ lands in k , and every embedding of $\mathbf{Oz}$ or $\mathbf{Oz}[i]$ into $S[U^{\mathbf{M}}]$, S a domain, lands in S .*

Proof of (i). In a real closed F , $1 + x^4 > 0$ has a square root; in an algebraically closed one every element does. The converse is [theorem 45.3](#). More generally, if $\phi : F \hookrightarrow k(U^G)$ is any field embedding, choose $b \in F$ with $b^2 = 1 + a^4$ for each $a \in F$; the images satisfy the same equation, so $\phi(a) \in k$. No order, valuation, linearity or summability of ϕ is assumed. $\square$

Corollary 45.5 (Coefficient recovery under isomorphism [16]). *If F, E are real closed or algebraically closed of characteristic zero, every field isomorphism $F(U^G) \cong E(U^{G'})$ restricts to an isomorphism $F \cong E$. For set-sized structures, an elementary embedding of the ambient fields restricts to an elementary embedding of the coefficient fields.*

Proof. The same parameter-free formula defines both coefficient fields; for elementarity relativize every quantifier to [\(45.1\)](#). $\square$

Remark 45.6 (Why a quartic [16]). The conic $Y^2 = 1 + X^2$ fails: $X = (T - T^{-1})/2$, $Y = (T + T^{-1})/2$ is a nonconstant point in $k(T)$. The quartic has positive genus and $1 + x^4$ has a square root at *every* real closed coefficient, not only on a bounded interval; this gives one formula for the real and the complex cases.

The omnific rings enter through $\text{Frac } \mathbf{Oz} = \mathbf{No}$ and $\text{Frac } \mathbf{Oz}[i] = \mathbf{No}(i)$ with units $\{\pm 1\}$ and $\{\pm 1, \pm i\}$ (`odg:thm:fractions`, `odg:prop:units`), which 16 reproves by the usual monomial clearing (its Proposition 4.1, credited to Conway through [14, Proposition 2.4.5]).

Theorem 45.7 (Existential coefficient recovery for rings [16]). *Let R be a domain whose fraction field F is real closed or algebraically closed of characteristic zero. Then [\(45.2\)](#) defines R in $R[U^M]$, and every embedding $R \hookrightarrow S[U^M]$, S a domain, has image in S .*

Proof. $\text{Frac}(R[U^M]) = F(U^G)$ with $G = \text{gp } M$, since fractions give F and the differences of exponents, and $F \cap R[U^M] = R$ by comparing coefficients. For $x \in R$ choose $b \in F$ with $b^2 = 1 + x^4$ and write $b = u/v$ with $v \neq 0$. Conversely, witnesses give $(u/v)^2 = 1 + x^4$ in $F(U^G)$, so $x \in F$ by [theorem 45.3](#) and $x \in R$. An embedding $R \hookrightarrow S[U^M]$ extends to $F \hookrightarrow \text{Frac}(S)(U^G)$, whose image lies in $\text{Frac } S$ by (i), and $\text{Frac}(S) \cap S[U^M] = S$. $\square$

This completes the proof of [theorem 45.4](#): the same two-witness formula works for every $\mathbf{Oz}[U^M]$ and $\mathbf{Oz}[i][U^M]$, whatever the rank, presentation or divisibility of M .

Corollary 45.8 (Uniform interpretation [16]). *Among monoid algebras whose coefficient rings have real closed or algebraically closed fraction fields of characteristic zero, the coefficient ring is uniformly parameter-free existentially definable. Isomorphisms of such algebras preserve the coefficient rings, even when the monoids have different or infinite ranks; for set-sized models, elementary embeddings restrict to elementary embeddings of the coefficient rings.*

Proof. Use the same formula on both sides, and relativize quantifiers as in [corollary 45.5](#). $\square$

The ring formula contains an inequation, so it is existential but not one polynomial equation; deleting $v \neq 0$ admits the witnesses $u = v = 0$ for every x . The field formula is one equation with one witness. Neither formula defines $\mathbf{Oz}$ inside the pure field $\mathbf{No}$, or $\mathbb{R}$ inside $\mathbf{No}$: the ambient structures are the monoid algebra and its fraction field, and with trivial parameters the ring formula only restates $x \in R$. No algorithm recognizing surreal coefficients is inferred [16].

46 Sparse powers under arbitrary monomial extensions

Lemma 46.1 (Sparse multiplicity bound [16]). *A nonzero Laurent polynomial $f = \sum_{j=1}^s c_j T^{m_j}$ over a characteristic-zero field, with distinct m_j and nonzero c_j , has no nonzero root of multiplicity $\geq s$.*

Proof. With $\vartheta = T d/dT$, a root $\alpha \neq 0$ of multiplicity $\geq s$ gives $\vartheta^i f(\alpha) = \sum_j m_j^i c_j \alpha^{m_j} = 0$ for $0 \leq i < s$, since ϑ^i is a combination of derivatives of order $\leq i$. The Vandermonde determinant $\prod_{j < l} (m_l - m_j)$ is nonzero in characteristic zero, so every $c_j \alpha^{m_j} = 0$. This classical bound is discussed by Mattarei [17]. $\square$

Theorem 46.2 (Uniform sparse power obstruction [16]; 16's Main Theorem B). *Let $k \subseteq L$ be fields of characteristic zero, $G \hookrightarrow G'$ an embedding of torsion-free abelian groups, and $0 \neq r = P/Q \in k(U^G)$ with $P, Q \in k[U^G] \setminus \{0\}$. If r is not a monomial and $r = h^n$ for some $h \in L(U^{G'})$ and $n \geq 1$, then*

$$n \leq \max\{\text{wt } P, \text{wt } Q\} - 1. \quad (46.1)$$

P and Q need not be coprime. The bound is sharp.

Proof. Write $h = A/B$. Only finitely many exponents occur in P, Q, A, B ; by lemma 44.3(ii) specialize them to one variable keeping P, Q, A, B nonzero, the weights of P, Q , and r non-monomial. Then $r_0 = h_0^n$ in $L(T)$. A nonmonomial rational function of one variable has a zero or pole at some $\alpha \neq 0$ in an algebraic closure of L (otherwise numerator and denominator are, after cancellation, powers of T up to scalars). Its order $e \neq 0$ there is a multiple of n , and by lemma 46.1 $|e| = |\text{ord}_\alpha P_0 - \text{ord}_\alpha Q_0| \leq \max\{\text{wt } P, \text{wt } Q\} - 1$; multiplying by powers of T changes no order at $\alpha \neq 0$, and common factors cancel in the difference. So $n \leq |e|$. Sharpness: $r = (1 + T)^n$, or $((1 + T)/(1 - T))^n$, whose numerator and denominator have $n + 1$ terms each. $\square$

The root h may use new exponents, with arbitrarily fine denominators, and new coefficients; the bound does not depend on the rank of G' [16].

Corollary 46.3 (Unbounded roots force a monomial [16]). *If $0 \neq r \in k(U^G)$ has roots of unbounded orders in $L(U^{G'})$, then r is a monomial; in particular if the roots lie in $k(U^G)$ or $k[U^G]$.*

Proof. Fix one presentation P/Q and take a root order above (46.1). $\square$

Lemma 46.4 (Roots of monomials [16]). *If $h^n = cU^g$ in $k(U^G)$ with $n \geq 1$, $c \neq 0$, then $h = dU^a$ with $d \in k^\times$, $a \in G$, $d^n = c$ and $na = g$.*

Proof. In the Laurent polynomial ring of a finite free subgroup containing g and the supports of a numerator and denominator of h , a factorial domain, write $h = P/Q$ coprime; $P^n = cU^g Q^n$ forces P and Q to be units, and lemma 44.1 applies. $\square$

For a domain B put $\text{Rt}_\infty(B) = \{b \neq 0 : \text{for every } n \geq 1 \text{ some } c \in B \text{ has } c^n = b\}$, the *all-order root elements* (the roots need not be compatible), and for a torsion-free group $G_{\text{div}} = \bigcap_{n \geq 1} nG$.

Proposition 46.5 (The all-order root locus [16]). *For a characteristic-zero field k , $\text{Rt}_\infty(k(U^G)) = \{cU^g : c \in \text{Rt}_\infty(k), g \in G_{\text{div}}\}$, and G_{div} is the largest divisible subgroup of G . For a domain S the same holds in $S[U^G]$ with $\text{Rt}_\infty(S)$.*

Proof. [Corollary 46.3](#) makes such an element a monomial and [lemma 46.4](#) identifies its roots; in $S[U^G]$ the roots are monomials with coefficients in S ([lemma 44.1](#)). The converse takes roots of the coefficient and solutions of $na = g$. If $g = nh \in G_{\text{div}}$ and $g = nmu$, torsion-freeness gives $h = mu$; so G_{div} is divisible, and it contains every divisible subgroup. $\square$

Remark 46.6 (Two detectors [16]). For k real closed or algebraically closed, k is also the set of x such that x^2 and $1 + x^2$ both lie in Rt_∞ or vanish: they are then monomials differing by 1, so $x^2 \in k$, and [lemma 44.2](#) applies. This description is infinitary; the quartic replaces the countable family of root requirements by one equation.

47 Root-covered rings at arbitrary rank

A domain R is root-covered ([definition 12.1](#)) exactly when every nonzero element divides an element of $\text{Rt}_\infty(R)$; 16 uses this form. That $\mathbf{Oz}$ and $\mathbf{Oz}[i]$ are root-covered, and that $\mathcal{R}_o(k, \Gamma)$ is so exactly when the divisible $\Gamma \neq 0$ has no order unit, are [corollary 13.3](#) and [theorem 13.1](#), which 16 reproves with the same geometric-series argument and credits to source 02 (its Proposition 6.2), together with the workspace of [section 18](#) and its warning that the fraction field is not the full Hahn field (its Example 6.3).

Theorem 47.1 (Arbitrary-monoid coefficient rigidity [16]). *Let R be a root-covered domain of characteristic zero and S a domain. Every embedding $\phi : R \hookrightarrow S[U^M]$, for a cancellative torsion-free monoid M of any rank, has image in S .*

Proof. For $a \neq 0$ choose $ab = w \in \text{Rt}_\infty(R)$. Then $\phi(w) \neq 0$ has roots of all orders in $S[U^M] \subseteq S[U^{\text{gp}M}]$, so it is a monomial ([proposition 46.5](#)), and so is $\phi(a)$ ([lemma 44.1](#)). For $a \neq 0, -1$ the same holds for $\phi(a) + 1 = \phi(a + 1)$; two monomials differing by 1 both have exponent 0, since otherwise the finite supports disagree at the constant term. So $\phi(a) \in S$; and $\phi(0) = 0$, $\phi(-1) = -1$. $\square$

With $M = \mathbb{N}^p \times \mathbb{Z}^q$ this is [theorems 15.1](#) and [15.3](#); the new cases are the monoids that are not finitely generated, such as $\mathbb{Q}^d$, $\mathbb{Q}_{\geq 0}^d$ or $\mathbb{Q}^{(\kappa)}$, where the degree argument of [part II](#) does not apply ([section 43.2](#)) [16].

Proposition 47.2 (An intrinsic root-divisor description [16]). *For a domain B let $\mathfrak{D}(B)$ be the set of nonzero a dividing some element of $\text{Rt}_\infty(B)$, and*

$$\mathfrak{C}(B) = \{0, -1\} \cup \{a \neq 0, -1 : a \in \mathfrak{D}(B), a + 1 \in \mathfrak{D}(B)\}.$$

If R is root-covered of characteristic zero, then $\mathfrak{C}(R[U^M]) = R$ for every cancellative torsion-free M . So isomorphisms between such monoid algebras preserve the coefficient rings.

Proof. As in [theorem 47.1](#), $a \in \mathfrak{D}(R[U^M])$ is a monomial, and a and $a + 1$ both monomials force $a \in R$. Root-coveredness gives the reverse inclusion; the definition is intrinsic to the ring. $\square$

$\mathfrak{C}$ is not claimed first order, and a finite truncation is vacuous: for orders $n_1, \dots, n_s$ and $N = \text{lcm}(n_j)$, every nonzero a divides a^N , which has the roots a^{N/n_j} . For $\mathbf{Oz}$ and $\mathbf{Oz}[i]$ the finite alternative is ([45.2](#)) [16].

48 Automorphisms of group and monoid algebras

Theorem 48.1 (All group-algebra automorphisms [16]). *Let R be a characteristic-zero domain that is root-covered or has a real closed or algebraically closed fraction field, and G torsion-free abelian. Every automorphism of $R[U^G]$ is uniquely*

$$\phi\left(\sum_g a_g U^g\right) = \sum_g \alpha(a_g) \chi(g) U^{\tau(g)}, \quad \alpha \in \text{Aut } R, \tau \in \text{Aut } G, \chi \in \text{Hom}(G, R^\times), \quad (48.1)$$

and every such triple gives an automorphism.

Proof. Coefficient rigidity (theorem 47.1 or theorem 45.7) for $\phi|_R$ and $\phi^{-1}|_R$ gives $\phi(R) = R$. By lemma 44.1, $R[U^G]^\times/R^\times \cong G$, so ϕ induces $\tau \in \text{Aut } G$ and $\phi(U^g) = \chi(g) U^{\tau(g)}$ with $\chi(g) \in R^\times$; multiplicativity makes χ a character, additivity gives (48.1), and the monomial basis gives uniqueness. Conversely the formula is multiplicative, with inverse from α^{-1} , τ^{-1} and the inverse unit factors. $\square$

Corollary 48.2 (Reconstruction of the exponent group [16]). *If R and S each satisfy a hypothesis of theorem 48.1, an isomorphism $R[U^G] \cong S[U^{G'}]$ forces $R \cong S$ and $G \cong G'$, and these suffice.*

Proof. Apply coefficient rigidity in both directions, then compare unit groups modulo the coefficient units. $\square$

For the omnific rings the characters take values in $\{\pm 1\}$ ($R = \mathbf{Oz}$) or $\{\pm 1, \pm i\}$ ($R = \mathbf{Oz}[i]$), which are trivial on divisible G . So, schematically in the sense of section 43.3,

$$\text{Aut}(R[U^G]) \cong \text{Aut}(R) \times \text{Aut}(G) \quad (R \in \{\mathbf{Oz}, \mathbf{Oz}[i]\}, G \text{ divisible}), \quad (48.2)$$

with $\text{Aut}(\mathbb{Q}^d) = \text{GL}_d(\mathbb{Q})$. The coefficient automorphisms $\text{Aut}(R)$ are not classified here; for $\mathbf{Oz}$ they are the subject of parts I and III [16].

Theorem 48.3 (Divisible monoids [16]). *Let R, S be as in corollary 48.2 and $\mathbf{M}, \mathbf{N}$ divisible cancellative torsion-free monoids. Every isomorphism $\phi : R[U^{\mathbf{M}}] \rightarrow S[U^{\mathbf{N}}]$ is uniquely $\phi(\sum_m a_m U^m) = \sum_m \alpha(a_m) \chi(m) U^{f(m)}$ with a ring isomorphism $\alpha : R \rightarrow S$, a monoid isomorphism $f : \mathbf{M} \rightarrow \mathbf{N}$ and a character $\chi : \mathbf{M} \rightarrow S^\times$; conversely such data give an isomorphism. For the omnific and Gaussian omnific rings every such character is trivial.*

Proof. Coefficient rigidity gives α . As $\mathbf{M}$ is divisible, U^m has roots $U^{m/n}$ of every order, so its image is a monomial $c_m U^{f(m)}$ with $c_m \in S$ (corollary 46.3); the same holds for ϕ^{-1} on each U^n , and composing shows that the exponent maps are inverse and the c_m are units. Products make f additive and c_m a character. A character of a divisible monoid into a finite group of exponent e is trivial: $\chi(m) = \chi(m/e)^e = 1$. $\square$

Example 48.4 (Rational orthants [16]). For $\mathbf{M} = \mathbb{Q}_{\geq 0}^d$, a monoid automorphism permutes the extreme rays (a ray is extreme when every decomposition of one of its elements into two elements of $\mathbf{M}$ stays on it) and scales each by a positive rational; it extends to a $\mathbb{Q}$ -linear bijection of $\mathbb{Q}^d$. So $\text{Aut}(\mathbf{M}) \cong (\mathbb{Q}_{>0})^d \rtimes \text{Sym}(d)$ and, for $R = \mathbf{Oz}$ or $\mathbf{Oz}[i]$, $\text{Aut}(R[U^{\mathbf{M}}]) \cong \text{Aut}(R) \times ((\mathbb{Q}_{>0})^d \rtimes \text{Sym}(d))$, in contrast with the large nonlinear automorphism groups of polynomial rings in several variables. (16 writes $\mathfrak{S}_d$ for the symmetric group; $\mathfrak{S}^\dagger$ is source 10's structure here.)

49 Rational group fields: an exact automorphism criterion

In $F[U^{\pm 1}]$ the units force monomial images, but $F(U)$ has $U \mapsto U + 1$. Divisibility of the exponent group can prevent such a change from extending to all monomial roots [16]. The general criterion uses one external theorem.

Cited theorem 49.1 ([15, Theorem 4.5]). *For a set-sized field k of characteristic zero and a torsion-free abelian group G , the defect group $\text{Def}_k(G) = k(U^G)^\times / (k^\times U^G)$ is free abelian.*

[Cited theorem 49.1](#) is from an unrefereed preprint of August 2026. 16 uses it as an identified input and does not reprove it; neither does this report, and no peer-review status is inferred. The results of [sections 44 to 48](#) do not depend on it. For a class field k and a homomorphism $g \mapsto r_g \in k(U^{G'})^\times$ from a set group G , choose numerator and denominator codes for the set of values; their coefficients generate a set subfield k_0 , all r_g lie in $k_0(U^{G'})$, and by [lemma 44.2](#) being a monomial over k is being one over k_0 . So the kernel of $G \rightarrow \text{Def}_{k_0}(G')$ is the monomial kernel, and no proper-class version of [cited theorem 49.1](#) is used [16].

Theorem 49.2 (Exact monomial-automorphism criterion [16]; 16's Main Theorem C). *Let F be real closed or algebraically closed of characteristic zero and G a set-sized torsion-free abelian group. The following are equivalent:*

- (i) $\text{Hom}(G, \mathbb{Z}) = 0$;
- (ii) *every field automorphism of $F(U^G)$ is uniquely of the form (48.1), extended to quotients, with $\alpha \in \text{Aut } F$, $\tau \in \text{Aut } G$ and $\chi \in \text{Hom}(G, F^\times)$.*

The implication (i) $\Rightarrow$ (ii) uses [cited theorem 49.1](#); for groups generated by elements with unbounded root orders it has the independent proof of [proposition 49.3](#).

Proof. An automorphism ϕ preserves F ([corollary 45.5](#)). Assume (i). By [cited theorem 49.1](#), with the descent above if F is a class, $g \mapsto [\phi(U^g)]$ is a homomorphism from G into a free abelian group, hence zero, since each coordinate projection to $\mathbb{Z}$ vanishes. So every $\phi(U^g)$ is a monomial; the same for ϕ^{-1} gives inverse exponent maps, and coefficient comparison gives (48.1) as in [theorem 48.1](#), extended uniquely to fractions. Conversely, if $\text{Hom}(G, \mathbb{Z}) \neq 0$, a nonzero map has image $d\mathbb{Z} \cong \mathbb{Z}$ and splits, so $G \cong G_0 \oplus \mathbb{Z}$ and $F(U^G) \cong F(U^{G_0})(T)$; the automorphism fixing $F(U^{G_0})$ with $T \mapsto T + 1$ is not monomial. $\square$

The condition says that G has no direct summand $\cong \mathbb{Z}$; it does not say that G is divisible [16].

Proposition 49.3 (An elementary range [16]). *If G is generated by elements g for which $\{n \geq 1 : g \in nG\}$ is unbounded, the description (ii) of [theorem 49.2](#) holds for F real closed or algebraically closed of characteristic zero, without [cited theorem 49.1](#).*

Proof. For such a generator U^g has roots of unbounded orders, and so has its image, which is a monomial by [corollary 46.3](#). Products give every g , and ϕ^{-1} gives the inverse exponent map; F is preserved by the quartic formula. $\square$

Examples are the divisible groups and the $\mathbb{Z}[1/p]$ -modules, and direct sums of groups of this kind, even though sums of generators may have only bounded root orders; so divisibility of every element is not needed [16]. For divisible G and $F = \mathbf{No}$, every character in (48.1) is positive, as $\chi(g) = \chi(g/2)^2$, and for $G = \mathbb{Q}^d$ it is determined by d positive surreal values. Over $\mathbf{No}(i)$ compatible roots matter: $U^q \mapsto e^{2\pi i q} U^q$, $q \in \mathbb{Q}$, is a nontrivial automorphism fixing U^1 ;

its coefficients are ordinary roots of unity, and no surcomplex exponential is used. Triples compose as

$$(\alpha, \tau, \chi) \circ (\alpha', \tau', \chi') = (\alpha\alpha', \tau\tau', g \mapsto \alpha(\chi'(g))\chi(\tau'(g))).$$

Lemma 49.4 (Relative rank is relative transcendence degree [16]). *If $A \leq G$ are torsion-free abelian groups, $\text{trdeg}_{k(U^A)} k(U^G) = \dim_{\mathbb{Q}}((G/A) \otimes \mathbb{Q})$; in particular $\text{trdeg}_k \text{Frac}(k[U^M]) = \text{rk}_{\mathbb{Q}} M$.*

Proof. If the images of g_j form a basis of $(G/A) \otimes \mathbb{Q}$, the U^{g_j} are algebraically independent over $k(U^A)$ (clear denominators and compare the distinct cosets of exponents), and every U^g is algebraic over them and $k(U^A)$, since $ng = a + \sum_j m_j g_j$ for some $n > 0$ and $a \in A$. The monoid case uses the group completion. This computation is also used in [15]. $\square$

Theorem 49.5 (Coefficient field and exponent group together [16]). *Let F, E be real closed or algebraically closed of characteristic zero and G, G' set-sized torsion-free abelian groups. Then $F(U^G) \cong E(U^{G'})$ if and only if $F \cong E$ and $G \cong G'$. The forward implication uses cited theorem 49.1 and the two-frame argument of Lin and Wang; it does not say that every isomorphism is monomial.*

Proof. An isomorphism ϕ maps F onto E . Put $A = \{g : \phi(U^g) \in E^{\times} U^{G'}\}$ and $A' = \{h : \phi^{-1}(U^h) \in F^{\times} U^G\}$. Monomial comparison in both directions gives $A \cong A'$ and $\phi(F(U^A)) = E(U^{A'})$. The defect maps embed G/A and G'/A' into free abelian groups (cited theorem 49.1), so both are free and the sequences split; by lemma 49.4 their ranks agree. Hence $G \cong A \oplus \mathbb{Z}^{(I)} \cong A' \oplus \mathbb{Z}^{(I)} \cong G'$. Class coefficients use the descent above; the converse is the coefficientwise monomial construction. $\square$

So rational monomial extensions of **No** or **No**(i) determine their parameter groups up to isomorphism. This extends a credited reconstruction argument and is not a new proof of Lin and Wang's theorem; it leaves $\text{Aut}(\mathbf{No})$ and $\text{Aut}(\mathbf{No}(i))$ as coefficient data [16].

50 Finite-rank cancellation and the exact infinite-rank failure

Theorem 50.1 (Finite-rank coefficient invariance [16]). *Let R be a characteristic-zero domain that is root-covered or has a real closed or algebraically closed fraction field, let M, N be cancellative torsion-free monoids of the same finite rational rank d , and S any domain. Every isomorphism $\phi : R[U^M] \rightarrow S[U^N]$ has $\phi(R) = S$; in particular $R \cong S$.*

Proof. Coefficient rigidity gives $\phi(R) \subseteq S$. Put $R_1 = \phi^{-1}(S)$, so $R \subseteq R_1 \subseteq R[U^M]$, and let $F = \text{Frac } R$, $E = \text{Frac } R_1$. Pulling the N -presentation back along ϕ gives $\text{trdeg}_E \text{Frac}(R[U^M]) = d$, and $\text{trdeg}_F \text{Frac}(R[U^M]) = d$ by lemma 49.4. Additivity gives $d = \text{trdeg}_F E + d$, so E/F is algebraic as d is finite, and $E = F$ by lemma 44.2. So $R_1 \subseteq F \cap R[U^M] = R$. $\square$

Corollary 50.2 (Finite-rank field coefficient invariance [16]). *Let F be real closed or algebraically closed of characteristic zero, G, G' of the same finite rational rank and E any field. Every isomorphism $F(U^G) \cong E(U^{G'})$ maps F onto E .*

Proof. The field clause of theorem 45.4 puts the image of F inside E ; pull E back to an intermediate field and argue as above. $\square$

These are coefficient-invariance statements, stronger than an isomorphism of coefficient rings. Finite generation is not needed, only finite rank; this is the first difference from the polynomial-cylinder statements of theorem 15.4, which for finitely many polynomial and

Laurent variables also give the equality of the numbers of variables of each kind. Laurent cancellation has a large literature [9] [16].

Theorem 50.3 (Infinite-rank counterexamples [16]). *Let κ be an infinite cardinal and $G_\kappa = \mathbb{Q}^{(\kappa)}$.*

- (i) *If R is a root-covered domain of characteristic zero and $S = R[U^\mathbb{Q}]$, then $S \not\cong R$ but $R[U^{G_\kappa}] \cong S[U^{G_\kappa}]$.*
- (ii) *If F is real closed or algebraically closed of characteristic zero and $E = F(U^\mathbb{Q})$, then $E \not\cong F$ but $F(U^{G_\kappa}) \cong E(U^{G_\kappa})$.*

Proof. $\mathbb{Q} \oplus G_\kappa \cong G_\kappa$ gives both isomorphisms. In (i), $1 + U$ is not a monomial, so by [proposition 46.5](#) and [lemma 44.1](#) it divides no element of $\text{Rt}_\infty(S)$; so S is not root-covered, while R is, and root-coveredness is an isomorphism invariant. In (ii), F satisfies $\forall x \exists y (y^2 = 1 + x^4)$ and E does not, at $x = U^1$ ([theorem 45.3](#)). $\square$

Theorem 50.4 (A sharp rank threshold [16]; 16's Main Theorem D). *For $R = \mathbf{Oz}$ or $\mathbf{Oz}[i]$, the property “for every domain S , every isomorphism $R[U^{G_\kappa}] \cong S[U^{G_\kappa}]$ maps R onto S ” holds exactly when κ is finite. For $F = \mathbf{No}$ or $\mathbf{No}(i)$ and arbitrary comparison fields E , the analogous property of $F(U^{G_\kappa})$ has the same threshold. For infinite κ the witnesses are $S = R[U^\mathbb{Q}]$ and $E = F(U^\mathbb{Q})$.*

Proof. [Theorem 50.1](#) and [corollary 50.2](#) for finite κ ($\text{rk}_\mathbb{Q} \mathbb{Q}^\kappa = \kappa$); [theorem 50.3](#) for infinite κ , where no isomorphism can map R onto $S \not\cong R$. $\square$

In the counterexamples the enlarged comparison ring is not root-covered and the enlarged comparison field is neither real closed nor algebraically closed, so [corollary 48.2](#) and [theorem 49.5](#) do not apply to that side; within the original family reconstruction holds at every rank. Cardinal arithmetic, $\kappa + 1 = \kappa$, not a defect of the finite-rank proof, allows the absorption [16].

51 Boundaries of the method

Example 51.1 (Hahn completion destroys the detector [16]). Let k be real closed or algebraically closed of characteristic zero. In the Hahn field $k((t^\mathbb{Q}))$,

$$b = \sum_{n \geq 0} \binom{1/2}{n} t^{4n} \quad \text{satisfies} \quad b^2 = 1 + t^4,$$

coefficientwise, so the nonconstant t satisfies the quartic detector there, while no nonconstant solution exists in $k(U^\mathbb{Q})$. The results of this part must not be transferred to Hahn fields. There is no conflict with $\sqrt{1 + \omega^4} \in \mathbf{No}$: in $\mathbf{No}(U^G)$ the element ω is a coefficient, and an external U^g with $g \neq 0$ is not a surreal magnitude. Adjoining every rational power of a parameter can increase rigidity, whereas allowing arbitrary Hahn sums destroys the detector [16].

One missing divisibility hypothesis allows translations [16]. For $G = \mathbb{Z}$, $F(U^G) = F(T)$ has $T \mapsto T + 1$. For $G = \mathbb{Q}$ [theorem 49.2](#) excludes it; elementarily, T has every root $T^{1/n}$, while $T + 1$ has two terms and is no n th power in $F(U^\mathbb{Q})$ for $n > 1$ by [\(46.1\)](#). At the ring level even $R[T^{\pm 1}]$ has no translation, since $T + 1$ is not a unit; field and ring hypotheses must not be interchanged.

Example 51.2 (Positive characteristic [16]). In characteristic $p > 0$, $(1 + T)^{p^r} = 1 + T^{p^r}$ has two terms for every r , so (46.1) fails. For $G = \mathbb{Z}[1/p]$, $\text{Hom}(G, \mathbb{Z}) = 0$, but $T^{1/p^n} \mapsto T^{1/p^n} + 1$ ($n \geq 0$) are compatible under p th powers and define a nonmonomial automorphism of $k(U^G)$, with inverse from -1 ; so characteristic zero is substantive in theorem 49.2. In characteristic 2, $1 + x^4 = (1 + x^2)^2$ for every x , and the quartic formula fails completely.

Torsion, zero divisors and injectivity [16]. Torsion-free cancellative monoids give domains and separate finite supports. A monoid with a nontrivial idempotent e has $U^e(1 - U^e) = 0$; the two-element idempotent monoid gives $R[Y]/(Y^2 - Y) \cong R \times R$, whose independent factor automorphisms need not preserve the diagonal copy of R . Such examples are outside the theorems. Injectivity keeps the root element nonzero in the target; no classification of noninjective maps into reduced parameter rings is given, and question 19.3 stays open.

What the results say about surreal automorphisms [16]. They classify automorphisms of the constructed extensions relative to the remaining coefficient automorphism. They do not compute $\text{Aut}(\mathbf{No})$ or $\text{Aut}(\mathbf{No}(i))$, and they do not decide whether an automorphism of $\mathbf{No}(i)$ preserving $\mathbf{Oz}[i]$ preserves conjugation or the natural valuation (question 32.1). Hahn summation is not part of $F[U^G]$ or $F(U^G)$; every displayed sum is finite.

Remark 51.3 (Dependencies and proof obligations [16]). The quartic path uses field algebra and lemma 45.1; the sparse path uses finite specialization and lemma 46.1; both are proved here. Arbitrary-monoid rigidity rests on 02's root-cover framework and the sparse bound; the group-algebra and divisible-monoid formulas on coefficient recovery, units and roots; the general field criterion and theorem 49.5 on cited theorem 49.1; the divisible and prime-local criterion on the sparse bound alone; the cancellation threshold on embedding rigidity, relative transcendence degree and group absorption. 16's checklist of proof obligations is honoured above: monoids are cancellative and torsion-free before completion; specialization acts on a finite identity with nonzero denominators, injective on $\text{supp } P - \text{supp } Q$ where needed; a root order is compared with a fixed presentation, without a single lattice containing all roots; zero cases precede Mason–Stothers; ring witnesses satisfy $v \neq 0$ and $\text{Frac}(R[U^M]) = \text{Frac}(R)(U^{\text{gp } M})$ is used explicitly; set-sized Hahn root-covered rings are not identified with $\mathbf{No}$ or with their Hahn fraction field; the field criterion recovers coefficients first and then uses the set-subfield form of the defect theorem; the infinite-rank counterexamples leave the recovered family; characters of the omnific *rings* lie in a finite unit group and vanish on divisible groups, those of the surcomplex *field* need not; all-order root membership is not treated as first order.

52 Questions of source 16, relations, and stale statements

52.1 Status of the source's questions

16 poses eleven questions (its 12.1–12.11), proposed as research problems, not as published open problems. Question 12.8 (reduced parameter rings and kernels) is question 19.3, with 16 added to its sources; 16's domain proofs do not reach it. The other ten are new.

Question 52.1 (Beyond rational monomial fields [16]). Which intermediate fields $k(U^G) \subseteq L \subseteq k((t^G))$ (for an ordered G , with $U^g = t^g$) still satisfy quartic rigidity, so that $\exists y (y^2 = 1 + x^4)$ detects only k ?

A first task is to find the smallest algebraic or support-closed extensions with a nonconstant point of the quartic; adjoining the square root of example 51.1 already breaks the detector, so inclusion in a Hahn field is no useful sufficient condition [16].

Question 52.2 (Single-equation ring definitions [16]). For $R = \mathbf{Oz}$ or $\mathbf{Oz}[i]$, can the inequation in (45.2) be removed using further polynomial witnesses, uniformly in M and without naming a nonunit parameter?

A purely equational definition would be a different Diophantine result; deleting $v \neq 0$ is invalid [16].

Question 52.3 (Optimal root certificates [16]). Can (46.1) be refined through the Newton polytopes or coefficient relations of P, Q , keeping invariance under arbitrary torsion-free monomial extensions?

The sharp worst case uses weights only; a refinement would rule out a given root order without multivariate factoring, with finite separating functionals as witnesses, over effective coefficient fields and exponent presentations, not as an algorithm for all surreals [16].

Question 52.4 (Positive characteristic [16]). What replaces $\text{Hom}(G, \mathbb{Z}) = 0$ for monomial automorphism rigidity in characteristic p , and which prime-to- p root tests remain valid?

[Example 51.2](#) rules out the characteristic-zero criterion; Frobenius-divisible monomial and nonmonomial directions should be separated first [16].

Question 52.5 (Cancellation for nonabsorbing infinite groups [16]). For which infinite-rank groups G does coefficient invariance against arbitrary comparison rings still hold for $R[U^G]$?

$\mathbb{Q}^{(\kappa)}$ fails because it absorbs a $\mathbb{Q}$ summand; the finite-rank proof cannot cancel infinite cardinals, but that does not prove failure for every infinite-rank group [16].

Question 52.6 (Monoids without divisibility [16]). How much of a cancellative torsion-free monoid M can be recovered from $R[U^M]$, once the coefficient ring is recovered, without assuming M divisible?

[Theorem 48.3](#) needs all-order roots; nonlinear polynomial automorphisms obstruct the same formula otherwise, and the monoid-ring isomorphism literature should be compared [16].

Question 52.7 (Root-covered coefficient definability [16]). Which root-covered domains admit a uniform first-order definition of the coefficient ring in every $R[U^M]$, beyond the closed-fraction-field case?

$\mathfrak{C}$ is infinitary and its finite truncations are vacuous; the quartic settles the omnific examples, not every root-covered domain [16].

Question 52.8 (Intrinsic monomial sets [16]). Under $\text{Hom}(G, \mathbb{Z}) = 0$, is the monomial subgroup $F^\times U^G$ of $F(U^G)$ first-order definable for interesting classes of groups, rather than only invariant under all automorphisms?

For divisible G its all-order-root description is explicit but not first order; invariance never establishes definability [16].

Question 52.9 (Surcomplex real-form recovery [16]). Does additional Gaussian omnific structure in larger extensions recover a preferred conjugation or real form of $\mathbf{No}(i)$, and under exactly which valuation or summation hypotheses?

The present results recover $\mathbf{No}(i)$ as a coefficient field, not its copy of $\mathbf{No}$ or its conjugation, so they do not answer [question 32.1](#) [16]. The question is a relative of `odg:def:q:realform` of the Diophantine report and of [question 32.1](#) [merge].

Question 52.10 (Formalization in stages [16]). Can the specialization lemma, the sparse power bound, the quartic detector and monoid reconstruction be formalized over set-sized coefficient fields and monoids, and then connected to the repository’s surreal construction?

16 proposes the order: sparse support separation, the Mason–Stothers quartic corollary, fraction and monoid compatibility, coefficient interpretation, finite-rank cancellation, and only then class descent, keeping the general criterion a separate dependency until defect freeness is formally available. The Lean formalization of Mason–Stothers by Baek and Lee [1] is a possible starting point, not a certificate for 16 [16].

52.2 Relations and stale statements

- *Not answered.* [Question 19.1](#) (groups with an order unit) is not addressed. 16 does not answer `odg: def: q: coeffpe` of the Diophantine report: that question asks for a lower-complexity recovery of $\mathbb{R}$ or $\mathbb{C}$ from the integer-part ring alone, whereas (45.2) recovers R in $R[U^M]$; with trivial M it is vacuous, and it defines neither **Oz** nor $\mathbb{R}$ in **No** [merge]. [Question 19.3](#) and [question 19.2](#) stay as they were.
- *Checked true.* 16’s description of source 02 and of its pin; the pin `1ab41af` and its tree `fba47f2`, which 16 correctly distinguishes; the SHA-256 hashes in its build record, which match the delivered files; its statement that the reducible-parameter question remains open; its suite, rerun for the merge ([section C](#)). No mathematical error was found in 16.
- *Not rechecked.* [cited theorem 49.1](#) and the two-frame method are Lin and Wang’s; 16 read the preprint and states Theorem 4.5 as displayed, and the merge did not recheck the preprint. The pinpoint descriptions of [17], [1] and [9] are reported as 16 gives them.

16’s limitations are listed with the other sources’ in [section B](#).

A Provenance ledger

A.1 Section by section

Section	Sources and choices
1	Structure from 09; imported results from all three automorphism sources; the Fact 4.2 recheck is the merge’s (remark 1.2). 10’s provenance and symbols were added with it (batch 28); item (8) of section 1.1 and the rows and paragraph for 11, 12 and 13 in section 1.2 were added in batch 30; item (9), the row and paragraph for 14, and the counts on the title page and in the abstract were updated in batch 31; items (10) and (11), the rows and paragraph for 15 and 16, the abstract and the counts in batch 32.
2	04’s one-shift lemma and coefficient polynomial; 09’s and 04’s ring decompositions; 04’s and 08’s divisible part. Units and constant products are cited from the Diophantine report, not reprinted. 10’s one-shift lemma is lemma 2.1 .

Section	Sources and choices
3	The criterion is printed once. <i>Choice:</i> 09's generality (k of characteristic zero, unital $\mathfrak{o}$) with 04's import-free proof; the first implication is 09's, which removes 04's use of $\bigcap n\mathfrak{o} = 0$. 09's derivation form is printed as theorem 3.4 ; its operator form, which uses cited theorem 1.1 , as corollary 3.5 . Example 3.2 is from the placement dossier's check. Section 3.3 is 10's: its support-cut form printed as further equivalent conditions, its least-forbidden-shift proof as a marked second route, necessity without strongness, and Archimedean blocks.
4	Shears from 04 and 09 (after <code>saut:thm:shiftflow</code>); exact displacement from 04 (08 and 09 have the same computation); 09's uniform shifts; 04's common-shift correspondence as a second, proved route; 08's divisibility-free threshold as a marked case; the translation example from 04, 08, 09 and the surcomplex report; rank dichotomy from 04 and 09 (08's for single flows); bottom subgroup from 04; phase shears from 08. Section 4.7 is 10's: twists with a general profile, the binomial family, their fixed fields, finite orbits and algebraic independence; 10's rank witness and divisibility example are recorded in remark 4.23 .
5	Factorization from 04 and 09 (credited to [13] and <code>saut:thm:decomp</code>); all pair automorphisms from 08 and 09 through <code>odg:def:thm:autreal</code> . 10 re-derives both.
6	Bracket law from 04, 08, 09; filtration, rank bound, balanced brackets, mixed-parameter lemma and sharpness from 09; 04's commutator formula and 08's faithful $\mathbb{Z} \wr \mathbb{Z}$ printed as one group in 04's normalization; the all-times remark is the merge's. Example 6.13 is 10's.
7	Fixed field $\mathbb{R}$ from 04, 08, 09 (explicit choice of 08, 09); separation, hull, monomial parameters and closure from 04; $\mathbb{R}^\kappa$ and nonsolvability from 09. Section 7.4 is 10's.
8	Reconstruction and definitions of $\mathbb{Z}$ and Π from 04, 08, 09 are cited to the Diophantine report; printed extras: 08's order formula, the projections (09), floor (08), truncation and the set-sized version (09).
9	Set-parameter nondefinability from 04, 08, 09 printed once with 04's strengthening; representative rules from 09; twists, one-term class and RV from 04; contrasts added by the merge. Section 9.2 is 10's; the outer bound in corollary 9.11 and the remark on <code>cor:question54</code> are the merge's.
10	Complex criterion and conjugation-compatible witnesses from 09 and 04; real axis inside the kernel from 04; wild lift and Gaussian fixed field from 08 and 09 (09's definition of $\mathbb{C}$ is cited to the Diophantine report); proposition 10.5 from 10.

Section	Sources and choices
11	Status records and questions: 04's four, 08's first and third, 09's nonstrong question (merged with 04's first), and the merge's non-divisibility question. 10's four questions: nonstrong maps and generation are merged into questions 11.1 and 11.4 , definable closure is recorded under question 11.3 , and other integer parts is question 11.7 ; the item on <code>dsn:q:languages</code> and the correction of 10's novelty statement are the merge's.
part II	Manuscript 02 throughout, with credits to <code>bst:thm:fieldcriterion</code> , <code>bst:prop:localdensity</code> , <code>osq:prop:classder</code> and <code>odg:thm:fractions</code> ; section 19 is the merge's.
Part III	Manuscripts 11, 12 and 13 (batch 30). 20 : the merge's. 21–24 : 12's spine with 13's binary witness (second route), set-pair isomorphisms, coefficient matrix and example 23.6 ; the reconstruction is cited, not reprinted; the convex criterion reproved by 12 and 13 is theorems 3.3 and 3.8 ; corollary 24.4 is the merge's. 25 : 12's and 13's adjoint theory, 12's general retraction and tower law, 13's kernel statement. 26–29 : 11's, with the embeddings of 12 and 13 as the instances of theorem 27.3 , 12's image corollary (extended to all three by the merge), 13's non-homogeneity; the Taylor automorphism is <code>saut:thm:coeffflow</code> (credit added for 12 and 13); the gap C_ω in lemma 27.2 is the merge's. 30 : 12 and 13 for automorphisms, 11 for embeddings. 31 : 12's regular cardinals, 13's small integer parts, both sources' distinctions. 32 : the thirty questions of the three sources (ten each) merged into eighteen, with 12's and 13's admissibility question recorded under question 11.2 ; the status records and the note on the large-cardinal report are the merge's. Batch 31: 14 added to the sources of the results it re-derives, pointers to part IV , and 14's comments and status notes after questions 32.1 , 32.3 , 32.11 to 32.14 and 32.18 .
part IV	Manuscript 14 (batch 31). 33 : the merge's, from 14's comparison with the report. 34 : 14's matching lemma and detection proof (third route), its cancellation example and topology remark. 35 : 14's definition and proposition; the Puiseux example, the weakened hypothesis after the proposition and remark 35.4 are the merge's. 36 : 14's formal evaluation, matrix formula, $\Gamma = \mathbb{R}$ remark, shear and truncation example and Gaussian statements, translated; proposition 36.5 is the merge's. 37 : 14's eight questions, one new, seven merged, and the stale-novelty record.

Section	Sources and choices
part V	Manuscript 15 (batch 32). 38: the merge's, from 15's comparison and the blob check. 39: 15's Section 3, with its scope and audit remarks; remark 39.6 and the comparison with <code>found:sub:tsum</code> in remark 39.13 are the merge's. 40: 15's disjoint-row lemma (a variant of the third route), interpolation, cancellation example, Gaussian ring corollary and class description; the comparison with 14's matching and the reading of proposition 40.5 as an answer to the first clause of question 32.11 are the merge's. 41: 15's Section 11. 42: 15's twelve questions, six new, six merged. Its algebraic half is printed in part III, with 15 added to the sources.
part VI	Manuscript 16 (batch 32). 43: the merge's, from 16's comparison with source 02. 44–51: 16's Sections 2–10, translated to external monomials U^g ; its Proposition 4.1, Definition 6.1, Proposition 6.2 and Example 6.3 are printed in part II and the Diophantine report. 52: 16's eleven questions, ten new, one merged; the relation to <code>odg:def:q:coeffpe</code> is the merge's. The status notes and pointers in part II are the merge's.

A.2 Printed once, and where the merge chose

- *The classification* (04 thm. criterion, 08 thm. threshold, 09 thms. derivation and stabilizer): theorem 3.3 with 04's proof, theorem 3.4, corollary 3.5, and 08's theorem 4.10 as the divisibility-free single-flow case. 04's $a_{\sigma,\delta}$ and 09's c_δ are kept as different functionals with equivalent conditions (example 3.2).
- *Rank triviality* (04, 09; 08 for single flows): theorem 4.13. *Common fixed field* $\mathbb{R}$ (04, 08, 09): theorem 7.2. *Set-parameter nondefinability* (04, 08, 09): theorem 9.1.
- *The metabelian non-nilpotent group*: 04's $\langle A_s, B_u \rangle$ and 08's $\langle A_1, B_1 \rangle$ are the same group at $s = u = -1$; printed once as theorems 6.8 and 6.10.
- *Strong factorization* (04, 09): theorem 5.1.
- *Not reprinted* (cited to the Diophantine report): units and constant products, Pell rigidity, $\text{Frac } \mathbf{Oz} = \mathbf{No}$, the multiplier ring, the reconstruction of $\mathbb{R}$ and of the order, automorphisms fix $\mathbb{R}$, the split restriction to $\text{Aut } \mathbb{C}$, the first-order and existential definitions of $\mathbb{Z}$, Π and $\mathbb{C}$.
- *Kept as second routes*: 04's common-shift correspondence (theorem 4.8) beside the imported one; 04's separation lemma beside the explicit choice in theorem 7.2; 09's Pell-divisibility definition of $\Pi_{\mathbb{C}}$ is mentioned beside the Diophantine report's.
- *Added by the merge* (tagged [merge], complete proofs): the combined statement of theorem 3.3; example 3.2; remark 6.11; question 11.6; with source 10, the converse in theorem 4.18 and the outer bound in corollary 9.11.
- *Source 10, printed once* (batch 28): its criterion (10 thm. criterion, cor. logcut) as theorem 3.3 with the further equivalent conditions of theorem 3.8; its rank theorem as theorem 4.13; its stabilizer splitting as theorem 5.1; its fixed field for $F = \exp \xi$ as theorem 4.2; its parameter theorem as theorem 9.1 (with the two-term witness added as

theorem 9.8); monomials and the omega-map in its nondefinability theorem as theorem 9.1 and corollary 9.2; its inner-support bound as a consequence of theorem 7.4; its Gaussian theorem as theorem 10.1 with the extras of proposition 10.5; its nonabelian example beside theorem 6.8; its one-shift lemma as lemma 2.1; its floor, fraction-field and coefficient-rigidity statements as corollary 5.2, `odg:thm:fractions`, `odg:def:thm:multiplier` and `odg:def:thm:autreal`, not reprinted.

- *Source 10, kept as a second route:* its least-forbidden-shift proof of necessity (theorem 3.8), beside 04's polynomial argument; its direct proof of the inner-support bound (corollary 9.11).
- *Sources 12 and 13, printed once* (batch 30): scalar detection (12 thm. 3.3, 13 thm. 3.3) as theorem 21.4; the constant-term isomorphism theorem (12 thm. 4.1, 13 thm. 4.1) as theorem 22.1, with 12's weaker hypothesis; automatic strongness of $\text{Aut}(\mathbf{Oz})$ (12 thm. 5.5, 13 thm. 6.1) as theorem 23.2; the set pairs (12 cor. 5.6, 13 prop. 5.5 and cor. 6.2) as proposition 23.1 and corollary 23.4; monomial determination (12 cor. 5.7, 13 cor. 6.3); the factorization (12 thms. 6.1–6.2, 13 thms. 7.1–7.2) as theorem 24.1; scalar representability and adjoints (12 prop. 7.2, thm. 7.3; 13 thms. 8.1, 8.3); the retraction (12 thm. 7.4, 13 prop. 8.4) as theorem 25.3; monomial adjoints (12 ex. 7.5, 13 ex. 8.5); the class functional (12 thm. 8.1, 13 ex. 11.2); the surcomplex automorphism theorems (12 thms. 10.1–10.2, 13 thms. 10.1–10.2). Their reprinted reconstruction (12 lems. 5.1–5.2 and lem. 5.4, prop. 5.3; 13 lems. 5.1–5.2 and prop. 5.3) is cited from the Diophantine report and corollary 5.2, and their reproofs of the criterion are the two routes of theorems 3.3 and 3.8.
- *Sources 11, 12 and 13, printed once:* the embedding moving a real (11 thm. 6.3, 12 thm. 9.5, 13 thm. 9.5) as theorem 27.3, an instance of 11's classification; $J(\mathbf{No}) \cap \mathbb{R} = \ker d$ (12, 11 prop. 8.5) as corollary 27.4 and proposition 29.2; nonelementarity (12 rem. 9.8, 11 thm. 8.3) as theorem 29.1; the surcomplex embeddings (12 prop. 10.3, 13 cor. 10.4) as the case $\varepsilon = 1$ of theorem 30.3. 11's reproof of the Taylor automorphism is `saut:thm:coeffflow`; its constants and automorphism rigidity (11 prop. 8.1, cor. 8.2) are `odg:def:thm:multiplier`, `odg:def:cor:internal` and corollary 5.2.
- *Kept as a second route* (batch 30): 13's binary finite-row proof of detection beside 12's triangular detector (theorem 21.4).
- *Added by the merge* (batch 30): corollary 24.4, the gap C_ω for 12's exponent embedding (lemma 27.2), the extension of corollary 27.4 to all three maps, and the second sentence of question 32.13.
- *Source 14, printed once* (batch 31; its numbering): Thm. 1.1, Thm. 5.3 and Cor. 5.5 as theorem 23.2 and corollary 23.4; Lem. 2.2 as lemma 2.1 and lemma 21.1; Thm. 3.3 and Cor. 3.4 as theorem 21.4; Thm. 4.1 and Cor. 4.2 as theorem 22.1 and corollary 22.2; Props. 5.1–5.2 as the cited reconstruction and proposition 23.1; Lem. 5.4 as `odg:thm:fractions`; Cor. 5.6 as corollary 23.5; Thms. 6.1–6.2 and Cor. 6.3 as proposition 25.1 and theorem 25.2; Thm. 7.1 as theorem 25.6; Thm. 8.1 as theorem 24.1; Thm. 8.2 as theorem 3.3; Cor. 8.3 as corollary 24.3; Prop. 9.1 as the inverse of example 4.19; Prop. 10.1 and Thms. 10.2–10.3 as theorems 30.1 and 30.2; Ex. 10.4 as the phase twists of `odg:def:lem:twist` (table in section 33.2).
- *Source 14, kept as a third route:* its induced-matching proof of detection (lemma 34.1), beside 12's triangular and 13's binary detectors.
- *Added by the merge* (batch 31): example 35.3, remark 35.4, and proposition 36.5, the weakened hypothesis after proposition 35.2, and the status notes after questions 32.12

and 32.14.

- *Source 15, printed once* (batch 32; its numbering; table in [section 38.2](#)): Lem. 2.2 as [lemma 2.1](#) and [lemma 21.1](#); Thm. 4.2 as [theorem 21.4](#); Prop. 4.5 as [example 21.6](#); Thm. 5.1 as [theorem 22.1](#); Prop. 5.3 as a direction of [theorem 25.2](#); Lems. 6.1–6.2 as the cited reconstruction and [proposition 23.1](#); Thm. 6.4 as [proposition 23.1](#) and [corollary 23.4](#); Lem. 7.1 as `odg:thm:fractions`; Thm. 1.2, Cors. 7.2 and 7.4 as [theorem 23.2](#); Cor. 7.3 as [corollary 23.5](#); Thm. 8.1 as [theorem 24.1](#); Prop. 8.2 as [theorem 3.3](#); Rem. 8.3 as [remark 24.2](#); Cor. 8.4 as [corollary 24.3](#); Ex. 8.5 as [remark 36.4](#); Thms. 9.2–9.3 as [theorems 30.1](#) and [30.2](#); Thm. 10.2, Cor. 10.3 and Prop. 10.4 as [proposition 25.1](#) and [theorems 25.2](#) and [25.6](#).
- *Source 15, kept as a variant of the third route*: its disjoint-row proof of detection ([lemma 40.1](#)), beside 14’s induced matching.
- *Source 16, printed once* (batch 32): Def. 6.1, Prop. 6.2 and Ex. 6.3 as [definition 12.1](#), [theorem 13.1](#) and [corollary 13.3](#) and the workspace of [section 18](#); Prop. 4.1 as `odg:thm:fractions`, `odg:prop:units` and `odg:def:lem:units`; Question 12.8 as [question 19.3](#). Its generalizations of [theorems 15.1](#) and [15.3](#) to [15.5](#) are printed as new results in [part VI](#), since they are stronger.
- *Added by the merge* (batch 32): [remark 39.6](#); the comparison with the foundations report in [remark 39.13](#); the reading of [proposition 40.5](#) as an answer to the first clause of [question 32.11](#); the remark after [question 42.2](#); the relation after [question 42.6](#); the relation after [question 52.9](#) and to `odg:def:q:coeffpe` ([section 52.2](#)); the status notes after [questions 11.4](#), [19.3](#), [32.1](#), [32.11](#), [32.12](#), [32.18](#) and [37.1](#); the pointers after [theorem 15.5](#) and in [remark 34.4](#).

A.3 Corrections and stale statements

1. *Divisibility* (09): its remark that divisibility is essential is correct for $\mathcal{A}_k$ and not for $\mathcal{R}_{\mathbb{Z}}$: the translation does not preserve $\mathbb{Z} + t^{-1}\mathbb{R}[t^{-1}]$ ([remark 4.11](#)).
2. *Fact 4.2 of [12]* (09): the placement dossier doubted that it states the bijection; the merge re-read the PDF and found that it does, with only an assertion for proper classes ([remark 1.2](#)). 09’s claims $\mathcal{U}_{\mathbf{Oz}} = \text{Exp}(\mathcal{L}_A)$ and the upper half of “exactly $r - 1$ ” in its nonsolvability theorem rest on that assertion and are stated so ([corollary 3.5](#) and [theorem 7.7](#)).
3. *Stale novelty* (04, 08, 09): their reconstruction, first-order recovery of $\mathbb{R}$ and definitions of Π were written at pins `fb5c4b53` and `befe739b`, before the Diophantine report’s reconstruction section was placed; they are cited, not claimed ([section 8](#)). 04’s remark that its reconstruction is not first order is superseded by `odg:def:cor:internal`. 08’s definition of Π has the structure of `odg:cor:definablect`, which was in the tree at 08’s pin.
4. *08’s Question 2* is answered ([section 8](#)) and dropped.
5. *Credits added*: `saut:thm:shiftflow` (08 and 09 did not credit it), `saut:thm:decomp` (04 and 09 credited only [13]), the rigidity report’s `thm:no` and `cor:question54` (09 cited only [12, Proposition 5.2]), `bst:thm:fieldcriterion`, `bst:prop:localdensity`, `osq:prop:classder` and `odg:thm:fractions` (for 02); and the surcomplex report’s $T_1(t) = t/(1 - t)$ is noted not to preserve $\mathbf{Oz}$ ([example 4.12](#)).
6. *Repository statements checked true*: at both pins the catalogue listed 51 reports (04, 08); the fixed-shift flow is `saut:thm:shiftflow` (04); phase twists preserve $\mathbf{Oz}[z]$ and move

- the real axis (04); the exponential-rigidity question is answered in its report (04); Pell rigidity and a definition of $\mathbb{Z}$ were in the Diophantine report (08); $\text{Frac } \mathbf{Oz} = \mathbf{No}$ was there (09, 02); no report contained an $\mathbf{Oz}$ -automorphism classification (04, 08, 09). 04 reports two date fields for [12] v3 (23 April and 24 August 2026); it is cited as version 3 of 23 April 2026, as elsewhere in the collection.
7. *Stale novelty* (10): at its pin 58cd8e1, which does not contain this report, 10 stated that no inspected report guide contains the support-cut criterion and proposed the criterion, the rank boundary, the specialized fixed fields and the parameter witnesses as new. At the merge these are [theorems 3.3, 4.2, 4.13](#) and [9.1](#); 10 is credited as an independent fifth source, and only the material listed in [section 1.2](#) is printed as its contribution ([section 11.1](#)).
 8. *Checked true* (10): $\text{Frac } \mathbf{Oz} = \mathbf{No}$ is in the Diophantine report (`odg:thm:fractions`); the multiplier identity, real-coefficient rigidity and the Gaussian phase twists recorded in that report’s reconstruction audit are now `odg:def:thm:multiplier`, `odg:def:thm:autreal` and `odg:def:lem:twist`; 10 credits them and its own proofs of them are not claimed as new. No mathematical error was found in 10.
 9. *Status notes* (batch 30): [question 11.1](#) is answered by [theorem 23.2](#). The statements that recorded it as open or partly answered now carry status notes: [remark 3.6](#), [remark 3.9](#), the paragraph after [theorem 5.1](#), [corollary 5.2](#), the item on `osq:q:invisible` in [section 11.1](#), and the paragraph after [question 11.1](#). No label, statement number or earlier text was removed; the non-claims of 04, 09 and 10 are kept as source records with a status note ([section B](#)).
 10. *A false reason, corrected* (09; merge text of batch 26): after [corollary 10.4](#) the report said that a real automorphism “preserves order and cannot move an ordinary real”. Read literally this is false: the Taylor automorphism `saut:thm:coeffflow` is strong and order preserving and moves a real b to $b + t^\delta$. The correct reason is that an automorphism preserving $\mathbf{Oz}$ fixes $\mathbb{R}$, which is definable in the pair. The sentence is corrected in place; no result depended on it, so this is a correction, not a retraction.
 11. *Credits added* (batch 30): `saut:thm:coeffflow` for the Taylor motion used by 12 and 13, which do not credit it (13’s audit calls it an established mechanism); 11 credits it.
 12. *Repository statements checked* (batch 30): 12’s statement that this report’s Theorem 5.1 is the strong factorization and that its Corollary 5.2 fixes $\mathbb{R}$ and commutes with the projections and floor without strongness, and that 10’s audit disclaims a classification of nonstrong automorphisms, is accurate at its pin `a6c68ac`. 13’s statements on [theorems 3.3](#) and [5.1](#) and on the Diophantine reconstruction, and 11’s on the surcomplex Taylor motions and exponent lifts, the Diophantine ideal test, multiplier and fraction field and the automorphism rigidity of this report, are accurate at their pins. Their statements that no report proves automatic strongness, scalar detection or coefficient-moving embeddings are still true at the merge.
 13. *Wrong statements in delivered files* (batch 30), disclosed and not edited, since delivered files are kept byte-identical: 11’s audit (`11-coefficient-gaps-SOURCE_AUDIT.md`, Section 2) calls `a6c68ac3826ac337762b71a2ef901997d019260c` “a tree SHA, not a commit SHA”; it is a commit (it is 12’s pin), whose tree is `12bfef3`. 11’s pin `cf56b89`, its tree `21d04b0` and its four blob hashes are correct. 13’s audit (`13-omnific-isomorphisms-SOURCE_AUDIT.md`) gives `71c1710` as the blob of the top-level `README.md` at its pin `0865f04`; that blob is the file at the earlier commit `b895e86`, and at `0865f04` it is `8c636c7`. Neither affects the mathematics.

14. *Stale novelty* (14, batch 31): at its pin 3d40856, which contains the batch-26 text of this report but not 20c4c9f, 14 stated that the strongness hypothesis “can be removed” from this report’s real stabilizer classification and that it supplies the missing implication. That was accurate at the pin and is stale at the merge: sources 12 and 13 answered [question 11.1](#) in 20c4c9f. 14 is credited as an independent third derivation ([section 37.2](#)); no mathematical error was found in it.
15. *Delivered-file behaviour* (14, batch 31): 14’s script, shipped as `code/14-automatic-summability-verify.py`, always writes `data/verification.json` under the parent of its own directory and has no output option; its delivered README says it writes `data/verification.json`. Run in place it would create an unprefixed `data/verification.json` in this report; the file is not edited, and the collection README gives rerun instructions on a copy.
16. *Repository statements checked* (15, batch 32): 15 pins no commit but two blobs of this report’s source; 059939b5 is the file written in b3fa9e2 and 60f3b172 the file written in 20c4c9f, and its three audit blobs are those of 21375f8. Its statements that [question 11.1](#) was open in the first and answered by 12 and 13 in the second, and that the inspected audits do not contain its topological classification, are accurate. Its reading omitted [part IV](#) and the congruence topologies of the neighbouring reports ([section 38.1](#)). No mathematical error was found in 15 ([section 42.2](#)).
17. *Repository statements checked* (16, batch 32): its pin 1ab41af and tree fba47f2 are as stated, its description of source 02 is accurate, and the SHA-256 hashes of its build record match its delivered files. No mathematical error was found in 16 ([section 52.2](#)); its general automorphism criterion and pair reconstruction rest on an unrefereed preprint ([cited theorem 49.1](#)).
18. *Delivered-file behaviour* (15, 16, batch 32): `code/15-topological-collapse-verify.py` writes `verification.json` beside itself unless `--output` is given, so run in place it would create `code/verification.json`; `code/15-topological-collapse-Makefile` names the delivered `Surreal_Arithmetic_Topological_Collapse.tex` and `verify.py`. `code/16-coefficient-recovery-verify.py` likewise writes `verification.json` beside itself unless `--output` is given. `code/16-coefficient-recovery-build.py` creates `build_logs/` beside itself, then runs `verify.py` and `pdflatexarticle.tex` in its own directory, rewriting `article.pdf` and `verification.json` there; in this report those delivered names are absent, so it creates `code/build_logs/` and stops. The files are not edited; the collection README gives rerun instructions on a copy.

A.4 Symbols renamed from the sources

$\mathcal{J}$ (08, 09) and $\mathcal{I}$ (04) are Π ; B (09), $\mathcal{A}$ (04), $\mathcal{B}_k$ (08) are $\mathcal{A}_k$; A_D (09), O_D (04), A_k (08), $R_\Gamma(D, k)$ (02) are $\mathcal{R}_\circ$ or $\mathcal{R}_\circ(k, \Gamma)$, with the coefficient ring D renamed $\circ$ because D names derivations; $\mathcal{P}_\Gamma(k)$ (02) is $\Pi_k(\Gamma)$; H_δ (08) and $H(a)$ (02) are C_δ and, in [part II](#), $C(a)$; $\mathcal{U}_{A_D}$ (09) and $\mathcal{U}_D(G)$ (04) are $\mathcal{U}_{\mathcal{R}_\circ}(\Gamma)$; $D_{\varphi, \delta}$ (08) is $D_{\delta, -\varphi}$; q_η (09), $[\omega^\beta]$ (04), c_a (08) and χ_b (02) are q_β ; 09’s $H = \omega^\eta$ and H_j are $\eta = \omega^\beta$ and η_j ; 08’s A_1, B_1, u, E are $A_{-1}, B_{-1}, u_0, \exp$; 02’s covering element c is z , its K_b and H_j are K_{bd} and $\Gamma_{\leq j}$. 08’s ω -convention statements are translated to the t -convention ([section 1.5](#)); [part II](#) keeps 02’s large-monomial convention ([section 12.1](#)). Source 10 (large-monomial convention, $X^g = t^{-g}$, translated): H_h is C_δ and C_g is H_g (letters swapped); $\text{pr}_+, \text{ct}, \text{pr}_-$ are P_-, P_0, P_+ (subscripts reversed); $\mathcal{I}$ is Π ; R_D is $\mathcal{R}_\circ$; B is $\mathcal{A}_k$; $\mathcal{U}_D$ is $\mathcal{U}_{\mathcal{R}_\circ}(\Gamma)$; the shift h is δ ; u_g is $1 + \varepsilon_\sigma(-g)$; λ_b is q_b ; P and T are Θ and ξ ; $\sigma_{\lambda, F, h}$ is $\sigma_{h, -\lambda}^F$; $\mathcal{N}^\dagger$ is $\mathfrak{S}^\dagger$ ([section 1.5](#)). Sources 11, 12 and 13 (all in the t -convention; the table of [section 20.3](#)): $I_\Gamma, A_G, \mathcal{J}_\Gamma$ are $\mathcal{R}_\Gamma$; $A_\Gamma, B_\mathbb{R}, \mathcal{A}$ are $\mathcal{A}_\Gamma$; 12’s ℓ_g and 13’s L_γ are $h_\sigma(g)$;

11's C_τ is $\mathbf{G}_\tau$ and its H is $\tau(\Gamma)$, its $\mathcal{F}$ is C_1 , its Λ is Δ , its $\Phi_{\tau,\rho}$ is $J_{\tau,\rho}$, its $E_{d,\delta}$ is $\mathcal{T}_{d,\delta}$, its S_τ is $M_{1,\tau}$, its T_h is ι_μ , its p is μ_{11} , its h_{jk} are $\mu^{\pm\pm}$, its h_c, h_j, h in the parameter theorems are $\mu^{\text{cmp}}, \mu^{\text{jmp}}, \mu^{\text{rel}}$, its $\mathcal{B}_\lambda$ is $\mathcal{V}_\lambda$, its J_Γ, L_Γ are $\mathcal{R}_\Gamma^\mathbb{C}, K_\Gamma^\mathbb{C}$; 12's θ is $\iota_{\mu_{12}}$, its ψ is μ_{12} , its D, ρ_D and J_D are $d, \rho_{d,1}$ and $J_{\mu_{12},d}$, its $j^\dagger(1)$ and E_j are e_ϕ and ret_ϕ , its Λ is $\lambda_{\mathbf{On}}$, its $\mathbf{Sc}$ is $\mathbf{No}(i)$; 13's second τ is $\iota_{\mu_{13}}$, its p is μ_{13} , its $\rho = \sigma|_k$ is α , its D, E_D and F_D are $d, \rho_{d,1}$ and $J_{\mu_{13},d}$, its T^* is $T^\dagger$, its λ is $\lambda_{\mathbf{On}}$, its $\mathcal{P}$ is Π , and its factorization $U \circ M$ is $M \circ u$ with $U = MuM^{-1}$. The large-cardinal report's J is $\hat{j}$. No normalization changes. Source 14 (the table of [section 33.3](#)): its $A_\Gamma = \mathbb{Z} \oplus \Pi$ and A are $\mathcal{R}_\Gamma$; its B is $\mathcal{A}_k$ ($\mathcal{A}_\Gamma$ for $k = \mathbb{R}$); its $\langle x, y \rangle$ is $\langle x, y \rangle_0$; its coefficient isomorphism τ is α and its exponent automorphism ϕ is τ ; its $\rho, \tilde{\rho}$ are $\sigma, \tilde{\sigma}$; its $\nu, \mathcal{U}, \mathcal{G}, \mathcal{M}$ are $u, \mathcal{U}_{\mathcal{R}_\mathbb{Z}}(\Gamma)$, the stabilizer and the monomial maps; its $L_\nu(g), a_{\nu,\delta}$ are $h_u(g), a_{u,\delta}$; its T^* is $T^\dagger$; its Λ is $\lambda_{\mathbf{On}}$; its power series F is $\mathbf{F}$ and its subfield F is K' ; its $P_{<\gamma}$ is $T_{<\gamma}$; its shear letters e_1, e_2 are e_2, e_1 and its s is t^{e_1} ; its additive h is θ ; its infinitesimals u_j are z_j ; its $d > 0$ is δ ; its $\mathbf{No}[i], \mathbf{Oz}[i], A_\mathbb{C}, K_\mathbb{C}, \Pi_\mathbb{C}, B_\mathbb{C}$ are $\mathbf{No}(i), \mathbf{Oz}[i], \mathcal{R}_\Gamma^\mathbb{C}, K_\Gamma^\mathbb{C}, \Pi_\Gamma^\mathbb{C}, \mathbb{C} \oplus \Pi_\Gamma^\mathbb{C}$; its target monomial u is t . No normalization changes. Source 15 (the table of [section 38.3](#); t -convention): its topologies $\mathcal{T}, \mathcal{T}_0, \mathcal{T}_\beta$ are $\mathfrak{T}, \mathfrak{T}_0, \mathfrak{T}_\beta$; its $R_{\mathfrak{o},\Gamma} = \mathfrak{o} \oplus \Pi_k$ is $\mathcal{R}_{\mathfrak{o}}$ with Γ densely ordered; its closure of zero N is $\{0\}$; its exponent sets D, E, D_U, D_V are $\mathbf{A}, \mathbf{B}, \mathbf{A}_U, \mathbf{A}_V$; its $R_\Gamma, A, \mathfrak{m}$ are $\mathcal{R}_\Gamma, \mathcal{A}_\Gamma, \mathfrak{m}_\Gamma$; its $\rho = \sigma|_k$ is α ; its $\mathcal{U}_{R_\Gamma}, h(g), a_\delta$ are $\mathcal{U}_{\mathcal{R}_\mathbb{Z}}(\Gamma), h_u(g), a_{u,\delta}$; its $\langle f, h \rangle$ is $\langle f, h \rangle_0$; its summation functionals are strong functionals; its Λ is $\lambda_{\mathbf{On}}$; its D and D_a (supports of functionals) are S_λ and S_a ; its $\sigma_{c,r}$ is $M_{\chi,\tau}$; its $\mathbf{No}[i], \mathbf{Oz}[i]$ and $c(z)$ are $\mathbf{No}(i), \mathbf{Oz}[i]$ and $\bar{z}$; its invisibility corollary is [corollary 39.10](#), not [corollary 24.4](#). No normalization changes. Source 16 (the table of [section 43.3](#)): its external $X^g, k[G], K_k(G)$ are $U^g, k[U^G], k(U^G)$; its monoids M, N are $\mathbf{M}, \mathbf{N}$ and its second group H is G' ; its v, d, w are $\text{min supp}, \text{max supp}, \text{wd}$; its ℓ is λ and its $T d/dT$ is ϑ ; its $\mathcal{U}_\infty(B)$ and $D(G)$ are $\text{Rt}_\infty(B)$ and G_{div} ; its $\mathcal{D}(B), \mathcal{C}(B)$ are $\mathfrak{D}(B), \mathfrak{C}(B)$; its $\Delta_k(G)$ is $\text{Def}_k(G)$; its coefficient automorphism σ is α ; its $T = \phi^{-1}(S)$ is R_1 ; its $\mathfrak{S}_d$ is $\text{Sym}(d)$; its $R_\Gamma(D, k), \mathcal{P}_\Gamma(k)$ are $\mathcal{R}_{\mathfrak{o}}(k, \Gamma), \Pi_k(\Gamma)$; its $\mathbf{Oz}[i]$ is $\mathbf{Oz}[i]$. No normalization changes.

B Non-claims, by source

No source claims to solve a named conjecture, and none claims Lean verification, peer review or historical priority; the finite suites check identities only. The report keeps every limitation stated by each source.

Source 09.

1. The classification is only within strong, coefficient-fixing, leading-term-preserving automorphisms; it does not assert that an abstract automorphism preserving $\mathcal{R}_{\mathfrak{o}}$ is strong, and nonstrong pair automorphisms are not classified (the definability recovery applies to them; the logarithmic classification does not).
2. Coefficient functionals come from an existing strong derivation; arbitrary functions satisfying the vanishing condition are not asserted to assemble into an operator.
3. The shears are plain-field automorphisms and are not claimed to preserve the Gonshor exponential.
4. No set or NBG class of class-function graphs is formed; actions are encoded by single class relations; supports are sets.
5. The exponential–logarithm correspondence and the factorization pattern are imported, not reproved or claimed; the explicit many-shift constructions avoid the class version of the import.

6. No new factorization theorem for omnific integers is claimed; [14] addresses a different problem.
7. **Oz** is not identified with a model of full ordinary arithmetic.
8. Truncation at a named monomial is not a uniform definable choice of boundary.
9. Nondefinability is relative to the field-and-integer-part language; Conway normal forms remain set-theoretically defined.
10. Hahn-sum formulas are not claims of convergence in the fine topology.
11. Modulus equivariance is not a claim that moduli are fixed.
12. The repository comparison was bounded and targeted; search coverage is not a proof of novelty.
13. The exact-arithmetic program checks finite identities only; it does not decide well-ordering, prove the correspondence, verify class-theoretic statements or certify novelty; the formalization order is a proposal, not implemented modules.

Source 04.

1. It does not classify abstract (nonstrong) automorphisms of **Oz**; the fraction-field extension of an abstract automorphism is not known to be strong.
2. It imposes no exponential, omega-map, birthday or simplicity structure, and does not revisit the collection's exponential-rigidity question as if unanswered.
3. The criterion classifies existing maps; it is not a construction for arbitrary logarithmic characters, and the convex condition alone is not asserted to solve global admissibility.
4. The relative fixed field is computed for the elementary subgroup only; the full stabilizer only for monomial parameters; the Hahn hull is not asserted to be definable closure.
5. Class maps are treated individually; no class of all automorphisms is formed.
6. Nothing is claimed after naming a proper class of parameters.
7. Bare nondefinability of coefficient-one monomials already follows from character twists; the nondefinability statement alone is not advertised as a new mechanism.
8. $\text{Frac } \mathbf{Oz} = \mathbf{No}$ is a full-class statement; a set-sized Hahn field need not be the fraction field of its integer-part ring.
9. Its intrinsic coefficient invariance was not claimed to be first order (now superseded by the Diophantine report).
10. Real and imaginary shear times preserve different structures; its Gaussian result is not the first surcomplex automorphism moving the real axis.
11. No generation or density theorem; fine-topology convergence is not strong summation.
12. The fixed-shift exponential, the general monomial/kernel decomposition and the exponential-derivation correspondence are prior work.
13. Its finite checks are not proofs and do not implement arbitrary normal forms; no Lean file, no repository checkout or build; historical priority unverified.

Source 08.

1. The threshold covers the elementary family only; rank-one rescalings and characters preserve the integer part; not every leading-term-preserving automorphism is claimed to be the exponential of one fixed-shift derivation.
2. Finite-dimensional phase images are a sufficient condition only.
3. No first-order Gaussian reconstruction was claimed (one now exists, [section 8](#)).
4. The automorphisms are not asserted to commute with the surreal exponential, the Berarducci–Mantova derivation or the omega-map; the wreath product does not contradict rigidity for those structures.
5. Parameter avoidance fails for set-sized fields with the whole field named; a universe-relative collection is not identified with the proper class.
6. No classification of the full stabilizer, and no solution of additional-structure problems.
7. The reconstruction is not a homomorphism $\mathbf{Oz} \rightarrow \mathbb{R}$.
8. The nondefinability is not a statement about adding an arbitrary class predicate, nor about automorphisms required to commute with the omega-map.
9. Strong summation is not topological convergence.
10. Pell rigidity, the existence of nontrivial Hahn automorphisms, the derivation–exponential construction and the monomial-character viewpoint are not claimed new; some elementary lemmas may be known; priority is provisional.
11. Its questions are not named historical problems.
12. No Lean; not refereed; the checks are regression tests, not proofs of well-ordering, parameter avoidance or definability.

Source 02.

1. Root-covered is sufficient, not claimed necessary; failure of the monomial criterion is not used to claim failure of cancellation or rigidity.
2. The finite-type theorems are theorems about sets; no $\text{Spec}(\mathbf{Oz})$ and no class compactness; “ $\text{Aut}(\mathbf{Oz})$ ” and ML of class rings are shorthand; the finite-type theorem is not a statement that a class cannot inject into a set.
3. It does not classify $\text{Aut}(\mathbf{Oz})$, all derivations, or multivariable polynomial automorphisms, and makes no Jacobian-conjecture claim.
4. Not every homomorphism into every target is constant: injectivity and domain targets are required; no noninjective analogue is given.
5. It does not claim that $R[\mathbf{T}]$ has no locally nilpotent derivation; a Hahn expansion is not a grading.
6. The orbit-field formula concerns monomials; only a lower bound holds for the full formal image.
7. Formal flows involve no analytic convergence and no fine-topology limits, s is not a surreal, and they are not exponentiation inside a Hahn field.
8. The cancellation theorem does not solve the unrestricted cancellation problem; $\text{ML}(R[\mathbf{T}]) = R$ is not a solution of a general stable-rigidity conjecture.

9. Not new: character twists [12], strong-invariance terminology [11], the Makar–Limanov framework [5], Euler derivations, Vandermonde arguments, and the example of [14].
10. No named factorization conjecture is claimed solved; the literature search is not a priority certificate.
11. The questions are not presented as published open problems; no Lean, no peer review; the checks are finite identities.

Source 10.

1. The classification is within strong, coefficient-fixing, leading-term-fixing automorphisms; nonstrong and coefficient-moving automorphisms are not classified, and strongness is not a harmless omitted hypothesis.
2. The criterion classifies existing strong maps; an abstract character obeying the cut condition is not asserted to give an automorphism, since summability and an inverse must be proved.
3. The rank theorem concerns the leading-term-fixing factor only; it does not say that all rank-one automorphisms are trivial, and monomial maps such as $t^\gamma \mapsto t^{2\gamma}$ and character rescalings are not counterexamples. Archimedean rank is not the order-unit condition of the holonomic report.
4. The strong valued decomposition, principal-unit characters and their crossed law [13], simplicity rigidity [12], the existence of nontrivial strongly linear 1-automorphisms, Hahn normal forms and the support description of **Oz** [14] are prior work; the splitting is not a new decomposition theorem for $\text{Aut}(\mathbf{No})$.
5. Not every truncation is preserved, only signed Archimedean blocks.
6. The fixed field is relatively, not absolutely, algebraically closed; no exact transcendence degree of a class extension is claimed.
7. The nonabelian example gives no presentation or exhaustive description of the stabilizer; no transfinite composition or closure theorem is claimed.
8. No proper-class-indexed sum is formed and no class of automorphisms is formed; class maps are individual, in NBG with global choice.
9. The bounded-birthday automorphism does not preserve birthdays or the simplicity hierarchy globally.
10. The definable-closure statements are upper bounds, not descriptions of definable closure and not quantifier elimination; they do not recover every coefficient from the projections.
11. The formal exponential is not the Gonshor exponential.
12. $\text{Frac } \mathbf{Oz} = \mathbf{No}$, the multiplier identity and coefficient rigidity are prior repository results, credited and not claimed; automatic fixing of coefficients is asserted only for the full surreal field, not for arbitrary $\mathcal{R}_o \subseteq K$.
13. $\mathcal{R}_o$ is called integer-part-shaped, not an integer part in every generality; no ordering of the complex field is asserted and no global surcomplex exponential is introduced.
14. The divisibility example is for $o = k$ and does not assert that translation preserves $\mathbb{Z} + t^{-1}\mathbb{R}[t^{-1}]$; the criterion is not transferred to other integer parts.
15. Initial subdomains, truncation-closed subrings and subrings stabilized by an automorphism are different notions, and none is substituted for another.

16. It does not answer the omega-map question of [12], factorization problems for omnific integers, or the holonomic order-unit question; no named published conjecture is solved, and its questions are research questions.
17. Its finite checks do not prove summability, well-ordering of arbitrary supports, the existence of separating functionals, class-size statements or nondefinability.
18. No Lean, no repository clone or build, no peer review; novelty is provisional, and the repository and literature comparison was targeted, not exhaustive.

Source 11.

1. It is an unrefereed research draft with no Lean verification; priority is provisional and the novelty assessment limited; the existence of proper Hahn-field embeddings is not claimed as new.
2. The classification carries the exact-monomial hypothesis: it is not a classification of all surreal embeddings, does not assert that omnific-preserving embeddings are strong, and does not cover characters or unit corrections.
3. Taylor automorphisms, exponent lifts, the reconstruction of constants and the classical inputs (Hahn fields, normal forms, real and algebraic closedness, quantifier elimination, extension of derivations) are prior work.
4. The derivations of $\mathbb{R}$ are choice-dependent and noncomputable.
5. “Monomial-preserving” does not mean commuting with the omega-map, and “strong” does not include $\mathbb{R}$ -linearity; no omega-map or exponential question of [12] is claimed, and the explicit map fails omega-map compatibility at 1.
6. Elementarity of pure-field embeddings is classical; fixing the coefficients is not proved sufficient for elementarity of the pair.
7. The continuum of conjugacy types is a lower bound, not a classification.
8. No nonconjugacy is claimed for automorphisms preserving $\mathbf{Oz}[i]$ but not conjugation, and non-conjugation-compatible surcomplex embeddings are not classified.
9. $\text{Frac } \mathcal{R}_\Gamma = K_\Gamma$ is not assumed for set-sized Γ .
10. No collection of class automorphisms or of open subclasses is formed; the topological claims are local.
11. The annulus bounds are external.
12. The finite tests prove no infinite, class, conjugacy or model-theoretic statement.
13. No repository was modified or built.

Source 12.

1. It is unrefereed, has no Lean verification, solves no named published conjecture and certifies no priority; the repository comparison was limited to four files and five public sources, the full articles and the ledger were not read, and a local clone failed.
2. Support duality and finiteness spaces are not new; $\text{Frac } \mathbf{Oz} = \mathbf{No}$ is classical; the reconstruction and the convex criterion are prior repository results.
3. Admissibility is retained: arbitrary coefficient arrays are not claimed to define automorphisms.

4. The proper embeddings are not a counterexample to strongness of all embeddings, and are not claimed initial, exponential or omega-compatible; the derivation is not computable.
5. The adjoint theorem is for set-sized groups only, fails for **No**, and does not apply to non- $\mathbb{R}$ -linear maps.
6. No unrestricted Gaussian valuation-preservation theorem is proved, and preservation of the real axis is not assumed.
7. The support-bound result is for regular uncountable κ only; no singular case, and no identification with a birthday cutoff.
8. ret_ϕ is not a field retraction or a positive conditional expectation.
9. No preservation of the exponential, logarithm, omega-map, birthdays or simplicity is claimed.
10. Invariance under automorphisms is not first-order definability.
11. The set-pair corollary does not assert that automorphisms of $\mathcal{R}_\Gamma$ extend when $\text{Frac } \mathcal{R}_\Gamma$ is smaller.
12. Omnific-preserving automorphisms are not claimed to be trivial.
13. The distinction from the three-duals report is not an exhaustive non-overlap claim.
14. Finite checks are not proofs.

Source 13.

1. It is AI-assisted and unrefereed, not proof-assistant checked, not a resolution of a named conjecture, and makes no priority claim; a failed search is not novelty, and it does not claim to settle every question posed.
2. Earlier repository results are credited: fractions, reconstruction, the strong stabilizer and the shears.
3. The scalar strong-dual representation is set-size only; the automorphism theorem is not.
4. The residue criterion is not valuation-topological convergence and not first-order definability of summation.
5. Not all automorphisms of **No** are strong (the examples of [12] stand), and omnific automorphisms are not trivial.
6. Local convex vanishing is not proved sufficient, and no class-wide operator exponential-logarithm correspondence is used.
7. It does not prove that every automorphism of **Oz**[i] is valuation-compatible, and conjugation is not intrinsic; its two Gaussian theorems must not be merged into an unconditional classification.
8. The full-pair theorem assumes an ambient isomorphism; isomorphisms of the small integer parts are not claimed to extend.
9. No Gonshor exponential, Berarducci–Mantova derivation or omega-map is used; its exponent map is not a field map; no proper-class Zorn lemma or basis is used.
10. The drift embedding comes with no uniqueness claim; the failure of homogeneity concerns set-sized induced substructures, not elementary maps.
11. Finite checks do not verify infinite summability, transcendence bases or surjectivity.
12. No repository build was run, and the formalization ledger was not read.

Source 14.

1. It is an AI-assisted research draft, not refereed and not Lean-verified; “theorem” means that a proof is supplied. Global priority is not certified: the literature search is not a certificate of absence, and the priority of the omnific application, of the detector’s formulation and of the class counterexample needs independent review.
2. Finiteness-space duality [3] and the general decompositions of Hahn-field automorphisms [13, 12] are prior work; no priority is claimed for those frameworks, and the underlying support duality is not claimed to originate there. The reconstruction and the convex-scale criterion are credited to this report and re-proved.
3. The unrestricted Gaussian problem is not solved: neither automatic strongness of every automorphism preserving **Oz**[*i*] nor a nonstrong counterexample is claimed, and the phase example is strong, so it disproves nothing.
4. No embedding theorem: the isomorphism proof uses surjectivity essentially, and no statement about constant-term-compatible embeddings is inferred.
5. No compatibility with the global Gonshor exponential is claimed; the formal exponential and logarithm are evaluated at infinitesimals only. Continuity, the Conway monomial cross-section and arbitrary truncations are neither assumed nor concluded, and commutation with every truncation is false.
6. The representation of strong functionals and the adjoint theory hold for set-sized exponent groups only and fail for **No**.
7. It does not assert that every ring automorphism of every $\mathcal{R}_\Gamma$ extends to its full Hahn field; the fraction-field argument is for **Oz** only.
8. The convex criterion classifies existing normalized automorphisms; it does not construct automorphisms from additive data satisfying the vanishing condition, since joint support conditions and invertibility are needed.
9. Summability, the monomials and a cross-section are not claimed first-order definable; the intersection formula for Π is not a single first-order formula; the conclusion is invariance of an externally quantified operation, not definability.
10. The class-group notation $\text{Aut}(\mathbf{No}, \mathbf{Oz})$ and its factorization are elementwise; no NBG class of proper-class graphs is formed; every summed family is set-indexed and every series has set support, in NBG with global choice.
11. The countable detector property is a sufficient closure condition only; it is not claimed for transseries, Puiseux or left-finite subfields.
12. The adjoints are for a bilinear algebraic pairing, not a Hilbert-space inner product; $\sigma^\dagger = \sigma^{-1}$ is not analytic unitarity, and the pairing is not a Hermitian norm.
13. Formal summability is not convergence of partial sums or in the valuation topology; no ordering of the index set is used; the automorphism–derivation correspondence of [2] is not used.
14. Its finite checks validate finite identities only: they do not establish well-ordering of arbitrary supports, quantify over all series, verify class replacement or prove the main theorem. No Lean compilation was performed, and the proposed Lean modules and names are plans only.
15. Conway’s irreducibility conjectures are not addressed and no famous open conjecture is claimed settled; its questions are proposed research problems, not published open problems.

16. The repository was read, not modified; its comparison was targeted, not an audit of every manuscript; the Wikipedia article it lists was an orientation source only.

Source 15.

1. It is an AI-assisted manuscript, not independently refereed and not checked in a proof assistant; “theorem” means that a written proof is supplied. Historical priority is not certified; the source comparison was targeted, not an exhaustive audit of the repository or the literature, and its reading of this report was incomplete (the updated article was not read in its entirety). No repository build, Lean run or modification was performed; a local clone failed.
2. The topological results are about compatibility with *every* Hahn sum. Ordinary ring topologies without this summation requirement are not excluded, and the rings are not claimed to lack Hausdorff ring topologies: the discrete topology always exists and fails only to realize the nontrivial sums.
3. Joint continuity of multiplication is essential: the pairing topology of [section 41](#) is Hausdorff, realizes exactly the Hahn sums, and has only separately continuous multiplication.
4. The support-ring classification assumes a densely ordered exponent group. For non-dense groups such as lexicographic $\mathbb{Z}^2$ the truncated ring $\mathfrak{o} \oplus \Pi$ is open; the full-field theorem does not settle it.
5. The cyclic case is an existence statement, not a classification of all compatible topologies.
6. The countable-only strengthening uses the positive buffer of the class **No**; it is not asserted for set-sized groups, and its proof fails for $\Gamma = \mathbb{Q}$.
7. Automatic strongness, scalar detection, covariance, adjoints, the class-size failure of duality and the qualified Gaussian results are credited to sources 12 and 13; the disjoint-row detector is an alternative proof, not a priority claim; the convex criterion, the reconstruction, the factorization and the shear examples are credited to this report; its reconstruction proofs are not claimed new.
8. Unrestricted Gaussian automatic strongness is not proved: valuation or conjugation compatibility is assumed explicitly, the bare pair $(\mathbf{No}(i), \mathbf{Oz}[i])$ is not claimed strong, and the collapse theorem does not settle it.
9. Global admissibility of arbitrary monomial prescriptions is not solved; the convex criterion is not an existence theorem.
10. Invariance is not definability: no first-order definition of families or sums and no definable monomial section is claimed.
11. Class statements concern individual class maps and class-coded neighbourhood systems; no class of class objects and no proper-class sum is formed; families are set-indexed and supports are sets.
12. The self-duality and the equivalence of strongness with adjointability need set-sized groups and fail for **No**.
13. The detector needs a full Hahn field or a subfield containing its countable detectors; the isomorphism argument uses surjectivity and is not stated for embeddings.
14. The pairing topology uses the discrete topology of the coefficients, not the usual topology of $\mathbb{R}$ or $\mathbb{C}$, and is not identified with the topologies of [\[8\]](#).

15. Finiteness spaces and formal summation algebras [3, 2] are prior work; being a monoid object for a summation-sensitive tensor product is not joint continuity, and no class operator-exponential theorem is imported.
16. Its finite checks do not test arbitrary topologies, infinite supports, proper classes or the infinite covering lemmas: a finite rational mesh can always be covered by two finite sets.
17. No named historical conjecture is claimed resolved; its questions are proposed directions, not established conjectures.

Source 16.

1. It is a proof-bearing research draft, not peer reviewed and with no Lean verification of its new results; historical priority is not certified, some proposed claims may already follow from published results, and the repository comparison was targeted (catalogue, guides and audits, earlier reads from the then-current `main`), not an exhaustive review; a local clone failed and no repository build or Wolfram run took place.
2. No named Conway factorization conjecture and no unrestricted automorphism problem is claimed solved: $\text{Aut}(\mathbf{No})$ and $\text{Aut}(\mathbf{No}(i))$ are left as coefficient data.
3. The general criterion [theorem 49.2](#) and [theorem 49.5](#) depend on [cited theorem 49.1](#), from an unrefereed preprint; no peer-review status is inferred, the theorem is not reproved, and the two-frame method is credited to Lin and Wang.
4. The field formula does not define $\mathbf{Oz}$ (or $\mathbb{R}$) inside the pure field $\mathbf{No}$; the ring formula is interpreted in $R[U^M]$, uses an inequation, and is not a single Diophantine equation.
5. The root-divisor set $\mathfrak{C}$ is infinitary and not first order; its finite truncations are vacuous; all-order root membership is not treated as a first-order formula; no algorithm recognizing surreal coefficients is inferred.
6. Nothing is claimed about Gaussian stabilizers (whether automorphisms of $\mathbf{No}(i)$ preserving $\mathbf{Oz}[i]$ preserve conjugation or the valuation), Hahn summation or conjugation: the extensions have finite supports and are not Hahn completions, and the results must not be transferred to Hahn fields.
7. Everything is in characteristic zero, for domains and cancellative torsion-free monoids; positive characteristic, torsion and idempotents are outside the theorems.
8. Injectivity is essential; noninjective maps are not classified, and reduced parameter rings are not addressed ([question 19.3](#) stays open).
9. The infinite-rank failure is proved only for $\mathbb{Q}^{(\kappa)}$; other infinite-rank groups are open.
10. The quartic obstruction, the sparse multiplicity bound, the monomial unit theorem, normal forms, the transcendence-rank computation and defect freeness are antecedents, not inventions; the root-cover framework, the scale criterion and finite-polynomial cancellation are source 02's.
11. Class-sized coefficient cases are elementwise, with Scott-trick codes and set-subfield descent; no proper-class spectrum, Hahn completion or set of all class automorphisms is formed, and automorphism-group formulas are schematic.
12. An external monomial is not identified with a surreal magnitude.
13. Its finite checks are finite tests, not proofs: they cannot prove the arbitrary-rank results, quantify over all roots, validate class foundations or establish novelty. The Lean formalization of Mason–Stothers [1] and the repository's Lean code are not verification of 16.

14. Its questions are proposals, not published open problems.

Status note (batch 30). Non-claim 1 of 09, non-claim 1 of 04 and non-claim 1 of 10 say that nonstrong automorphisms are not classified and that strongness is not shown. They are kept as the sources’ records. For **(No, Oz)** and the full real Hahn pairs with divisible exponent group, [theorem 23.2](#) and [corollary 23.4](#) now show that every automorphism is strong, so [theorem 24.1](#) classifies them all; for **Oz**[*i*] without conjugation or valuation data, and for a general $(k, \mathfrak{o})$, those non-claims still describe the state of knowledge ([questions 32.1](#) and [32.13](#)).

Counts: 13 non-claims from 09, 13 from 04, 12 from 08, 11 from 02, 18 from 10, 13 from 11, 14 from 12, 12 from 13, 16 from 14, 17 from 15, 14 from 16.

C Finite checks

Eleven suites ship with the report, byte-identical to their delivery, under the prefixes of their manuscripts. They check finite identities in exact arithmetic; they do not prove Hahn summability, class-sized statements, definability or novelty. They were rerun on copies for the merges (10’s for the batch-28 addition, 11’s, 12’s and 13’s for the batch-30 addition, 14’s for the batch-31 addition, 15’s and 16’s for the batch-32 additions) with Python 3.14.4 and SymPy 1.14.0, with the recorded results reproduced: 15’s rewritten record is identical to the shipped one, and 16’s differs only in its `python` and `elapsed_seconds` fields.

Script and record	Content and behaviour
code/09-omnific-preservin g-verify_finite_identities.py; data/09-omnific-preservin g-verification_output.txt	Standard library. Product rule and bracket on finite Laurent polynomials, balanced brackets for ranks 2 to 8, mixed group-commutator coefficients with square-zero parameters, shear iterates, exponential laws, the rank-one escape, Pell growth. Twelve PASS lines and “ALL CHECKS PASSED”. Prints only.
code/04-preserving-automorphisms-verification.py; data/04-preserving-automorphisms-verification_report.json	SymPy 1.14.0 (<code>data/04-preserving-automorphisms-requirements.txt</code>). 7,062 assertions in 17 groups: exponential group law, bracket formula, coefficient polynomial, first corrections, commutator coefficients in lexicographic $\mathbb{Q}^3$. <i>Writes</i> its report to <code>--output</code> , by default <code>verification_report.json</code> in the working directory.
code/08-integer-part-symmetries-verify.py; data/08-integer-part-symmetries-verification.json	Standard library. 9,469 checks in 15 categories: derivation rule, falling-factorial iterates, flow composition, bracket, Vandermonde determinants, the semidirect-product law, commutator phases. <i>Always writes</i> <code>verification.json</code> beside itself.
code/02-parameter-rigidity-verify.py; data/02-parameter-rigidity-verification.json	Standard library. 7,041 assertions in 13 families: Euler product rule, factorial Leibniz, truncated flow composition, commutation, geometric remainders, Vandermonde determinants, binomial identity. <i>Writes</i> <code>verification.json</code> beside itself unless <code>--output</code> is given.

Script and record	Content and behaviour
<code>code/10-support-cut-verify.py; data/10-support-cut-verification.json</code>	Standard library, exact rational arithmetic. 28,668 assertions in 19 categories: binomial coefficients, formal additive composition and inverses by two independent power algorithms, rank-two composition, inverse and multiplicativity, signed support projections, exact first displacement, the least-bad-shift coefficient and the rational root crossing zero, the three-scale commutator. Writes <code>verification.json</code> beside itself unless <code>--output</code> is given.
<code>code/11-coefficient-gap-s-verify_finite_models.py; data/11-coefficient-gap-s-verification.json</code>	Standard library, exact fractions, seed 20260923. 3,880 assertions in 15 categories: truncated Taylor identities over $\mathbb{Q}[b, b^{-1}]$, multiplication of finite outer Hahn polynomials with lexicographic shifts, Taylor inversion, integer-part preservation and reflection, signs of negative blocks, rational samples of the index maps, parameter fixing. Prints to standard output; writes a file only with <code>--output</code> .
<code>code/12-automatic-strongness-verify.py; data/12-automatic-strongness-verification.json</code>	Standard library, exact fractions, seed 20260923. 17,774 assertions: 10,800 triangular-detector checks, 4,502 monomial adjoint and transfer checks, 2,472 rank-two Taylor-block and binomial checks. Writes <code>verification.json</code> beside itself unless <code>--output</code> is given.
<code>code/13-omnific-isomorphisms-verify.py; data/13-omnific-isomorphisms-verification.json</code>	Standard library, exact fractions, seed 20260923. 13,885 assertions in 16 categories: binary finite-row choices and their persistence, descending-support detection, coefficient reconstruction, monomial adjoints and retractions, scale-separated Taylor models. Writes <code>../data/verification.json</code> relative to its own directory unless <code>--output</code> is given.
<code>code/14-automatic-summability-verify.py; data/14-automatic-summability-verification.json</code>	Standard library, exact fractions, seed 20260923 for the adjoint trials. Generalized binomial identities to degree 20 for 31 rational parameters (961 multiplication and 31 inverse cases); the induced matching on a banded 256-row prefix (rows meeting columns $n, n+1, n+3$), with a matching of size 64 and its all-ones detector; the sign-reversed transpose (36.2) in 100 random trials with 13 source and 9 target exponents; the finite prefixes of example 34.3 for 1, 2, 5, 16 and 64 pairs. Prints its report and <i>always writes</i> <code>../data/verification.json</code> relative to its own directory; it has no output option.

Script and record	Content and behaviour
<code>code/15-topological-colla</code> <code>pse-verify.py;</code> <code>data/15-topological-colla</code> <code>pse-verification.json</code>	Standard library, exact fractions, seed 20260923. 27,139 assertions in 14 categories: constant-term pairings and coefficient extraction, disjoint rows and the sparse detector with interpolation, the cancellation boundary of example 40.3 , monomial transport, rank-two Vandermonde and inverse identities for the shear, order reflection $x \mapsto \gamma - x$ and fixed products on negative intervals, bounded increasing supports, valuation non-escape and cyclic remainders. It does not test arbitrary topologies or the infinite covering lemmas. <i>Writes verification.json</i> beside itself unless <code>--output</code> is given.
<code>code/16-coefficient-recov</code> <code>ery-verify.py;</code> <code>data/16-coefficient-recov</code> <code>ery-verification.json</code>	Standard library, exact integers and fractions, seed 20260923. 1,314 checks in 12 categories: polynomial <i>abc</i> instances, quartic finite tests, sparse polynomial and rational powers and sharpness, Vandermonde determinants, support-difference separation, specialization multiplicativity, automorphism composition and inverses, truncated Hahn roots of $1 + t^4$, and the boundary examples. <i>Writes verification.json</i> beside itself unless <code>--output</code> is given. 16's <code>code/16-coefficient-recovery-build.py</code> is its delivered build script (section A.3).

The placement dossiers also recorded independent checks (281 for the automorphism cluster, including [example 3.2](#), the commutator formula for symbolic s, u , the leading terms of [theorem 6.8](#) for $n \leq 4$, the conjugation lemma and the failure example of [remark 4.11](#); for 02 the quotient-rule identity of [example 17.8](#) to 29 terms; and 8,724 assertions for sources 11–13, including the Taylor-block embedding in a lexicographic $\mathbb{Z}^2$ model, the failure of the ambient Taylor automorphism to preserve **Oz** ([section 26.1](#)), the binary finite-row algorithm over $\mathbb{F}_2$ and $\mathbb{Q}$, the triangular detector, the least-forbidden-shift coefficient and [example 23.6](#)); these scripts are not shipped.

References

- [1] J. Baek and S. Lee, *Formalizing Mason–Stothers theorem and its corollaries in Lean 4*, [arXiv:2408.15180v1](#), 2024.
- [2] V. Bagayoko, L. S. Krapp, S. Kuhlmann, D. Panazzolo and M. Serra, *Automorphisms and derivations on algebras endowed with formal infinite sums*, [arXiv:2403.05827v2](#), September 22, 2025; especially Theorems 3.11 and 3.13.
- [3] R. Blute, R. Cockett, P.-A. Jacqmin and P. Scott, *Finiteness spaces and generalized power series*, [arXiv:1805.09836](#), 2018.
- [4] S. Camacho, *Truncation in Hahn fields is undecidable and wild*, [arXiv:1706.03722](#), 2017.
- [5] M. Chitayat and D. Daigle, *On the rigidity of certain Pham–Brieskorn rings*, *J. Algebra* **550** (2020), 290–308; [arXiv:1907.13259](#).
- [6] J. H. Conway, *On Numbers and Games*, Academic Press, 1976; second edition, A K Peters, 2001.
- [7] P. Ehrlich and E. Kaplan, *Number systems with simplicity hierarchies: a generalization of Conway’s theory of surreal numbers II*, [arXiv:1512.04001](#), 2015.
- [8] D. Flynn and K. Shamseddine, *On the topological structure of the Hahn field and convergence of power series*, [arXiv:1901.09137v1](#), 2019.
- [9] G. Freudenburg, *Laurent cancellation for rings of transcendence degree one*, [arXiv:1309.4737v2](#), 2013.
- [10] H. Gonshor, *An Introduction to the Theory of Surreal Numbers*, London Mathematical Society Lecture Note Series 110, Cambridge University Press, 1986.
- [11] Y. Hirano, *On the uniqueness of rings of coefficients in skew polynomial rings*, *Publ. Math. Debrecen* **54** (1999), 489–495.
- [12] E. Kaplan, L. S. Krapp and M. Serra, *Decomposing the automorphism group of the surreal numbers*, [arXiv:2509.22374v3](#), April 23, 2026; especially Remark 2.10, Fact 4.2, Example 4.4, Propositions 5.1 and 5.2, and Questions 5.4, 5.6, 5.7.
- [13] S. Kuhlmann and M. Serra, *The automorphism group of a valued field of generalised formal power series*, *J. Algebra* **605** (2022), 339–376; [arXiv:2107.03362](#).
- [14] S. L’Innocente and V. Mantova, *A factorisation theory for generalised power series and omnific integers*, *Adv. Math.* **442** (2024), 109513; [arXiv:1710.07304v5](#).
- [15] J. Lin and X. Wang, *Reconstruction of torsion-free abelian groups from rational group fields*, [arXiv:2608.10381v1](#), August 11, 2026 (preprint; cited for Theorem 4.5, not rechecked here).
- [16] D. Marker, *Model Theory: An Introduction*, Graduate Texts in Mathematics 217, Springer, 2002.
- [17] S. Mattarei, *Root multiplicities and number of nonzero coefficients of a polynomial*, *J. Algebra Appl.* **6** (2007), 469–475; [arXiv:math/0512239](#).

Reports of the collection are cited by directory and label: the omnific Diophantine report `docs/surreal/omnific-diophantine-geometry (odg:)`, the quotient report `docs/surreal/set-sized-quotients-of-omnific-integers (osq:)`, the surcomplex automorphism report `docs/surcomplex/surcomplex-field-automorphisms (saut:)`, the rigidity report `docs/surreal/exponential-automorphism-rigidity (bare labels)`, the transcendence report `docs/surreal/transcendence-over-bounded-support (bst:)`, the single-dilation report `docs/surcomplex/single-dilation-hahn-support (dsup:)` and the definable-surreals report `docs/foundations-and-computation/definable-surreals-and-omnific-integer`

s (dsn:), the three-duals report docs/surcomplex/three-duals-of-hahn-vector-spaces (duals:), the independent-copies report docs/surreal/independent-surreal-copies (isc:), and the large-cardinal report docs/foundations-and-computation/large-cardinal-embeddings-and-normal-forms, cited by directory only. Part V also cites the foundations report docs/foundations-and-computation/foundations (found:).