

Unitary divisor sum colored partitions and their asymptotics

Research article and reproducibility companion

2 October 2026

Part II: 4 October 2026 (UTC)

Abstract

The products defining OEIS A301981 and A301982 have color multiplicities equal to the sum of the unitary divisors. Their recorded pure stretched-exponential equivalents are false: neither coefficient-to-model ratio is eventually bounded above and below by positive constants. A least-height-zero argument supplies the necessary noncancellation, and an Abelian implication transfers coefficient assumptions to Mellin holomorphy. For each sequence separately, a logarithmic error of order $O_\varepsilon(n^{1/4+\varepsilon})$ for every $\varepsilon > 0$ is equivalent to the Riemann hypothesis. Unconditionally the error is $o(n^{1/3})$ and cannot be $O(n^{1/4-\varepsilon})$ for any $0 < \varepsilon < 1/4$. We give a corrected multiplicative equivalent that evaluates the exact product at the explicit point $(2A/n)^{1/3}$, an all-fixed-order expansion using the exact saddle and cumulants, and shrinking inverse-threshold brackets. An off-center Hermite version gives the same fixed-order and inverse accuracy at an explicit evaluation point, without an inner saddle solve. The arguments apply established Mellin and saddle methods to these two specific products. They neither prove nor disprove the Riemann hypothesis, and they require no simplicity of zeta zeros or infinite zero-residue expansion.

Part II (added 3 October 2026) answers the third further question of Part I. For each sign, $\liminf a_n^\pm/m_n^\pm = 0$ and $\limsup a_n^\pm/m_n^\pm = \infty$: the logarithm of the ratio exceeds a positive multiple of $n^{\sigma_*}/3(\log n)^{m-1}$, and falls below a negative one, infinitely often, where $s_* = \sigma_* + i\tau_*$ with $\sigma_* \geq 3/4$ is the surviving Mellin pole of Part I and m its order. Hence both centered logarithms are $\Omega_\pm(n^{1/4})$, and the error of inverting either pure model is $\Omega_\pm((\log y)^{7/8})$. The lower constants are explicit in the leading Laurent coefficient at that pole. The method is Landau's positivity principle; no Riemann hypothesis, simplicity of zeros or sum over zeros is used.

Contents

I	Corrected equivalents, exact saddles, and an RH criterion	2
1	The two products and the conclusions	2
1.1	Prior work and the scope of the contribution	4
2	Arithmetic and Mellin preliminaries	6
2.1	Convergence and elementary estimates	6
2.2	The arithmetic Dirichlet series and the kernels	6
3	Noncancellation and the false pure equivalents	7

4	Unconditional contour estimates	8
5	The exact saddle and all fixed orders	9
5.1	A uniform angular estimate	9
5.2	Finite cumulant polynomials	10
6	An equivalent at the explicit leading saddle	12
7	The logarithmic obstruction and the RH equivalence	12
7.1	A conditional reciprocal zeta bound	13
8	Model inverses and integer thresholds	14
9	All fixed orders without an inner saddle solve	16
9.1	The explicit-point model inverse	17
10	Computation and reproducibility	18
10.1	Exact coefficient recurrence and independent expansion	18
10.2	Cumulants and finite product tails	19
10.3	Interpretation of the numerical checks	19
11	Further questions and precise limits	20
II	Two-sided fluctuations of unitary-divisor partitions	22
12	Two-sided arithmetic oscillations in A301981 and A301982	24
12.1	The question and the advance over the repository	24
12.2	The prior coefficient bridge, with its analytic justification	25
12.3	A positivity lemma with the exact pole constant	26
12.4	Noncancellation and the coefficient theorem	27
12.5	A consequence for inversion	29
12.6	Scope and further questions	30
13	Verification, reproducibility, and further directions	30
13.1	Make the partition oscillations effective	31
13.2	Formalization and an independent submission path	31

Part I

Corrected equivalents, exact saddles, and an RH criterion

1 The two products and the conclusions

A positive divisor d of m is unitary if $\gcd(d, m/d) = 1$. Write

$$b_m = \sigma^*(m) = \sum_{d|m, \gcd(d, m/d)=1} d = \prod_{p^a \parallel m} (1 + p^a), \quad b_1 = 1. \quad (1.1)$$

There are b_m colors available for a part of numerical size m . The two generating functions are

$$F_-(z) = \prod_{m \geq 1} (1 - z^m)^{-b_m} = \sum_{n \geq 0} a_n^- z^n, \quad \text{A301981,} \quad (1.2)$$

$$F_+(z) = \prod_{m \geq 1} (1 + z^m)^{b_m} = \sum_{n \geq 0} a_n^+ z^n, \quad \text{A301982.} \quad (1.3)$$

The minus product allows repeated parts of each color. In the plus product, each individual color may be used at most once; different colors of the same numerical size may occur together. All coefficients are positive, and $a_0^\pm = 1$.

For either sign set $f_\pm(t) = \log F_\pm(e^{-t})$ for real $t > 0$, and put

$$A_- = \frac{\pi^2}{6}, \quad A_+ = \frac{\pi^2}{8}, \quad c_\pm = 3(A_\pm/4)^{1/3}, \quad t_0^\pm(n) = (2A_\pm/n)^{1/3}. \quad (1.4)$$

Unless signs matter, a_n, f, A, c, t_0 refer to one fixed choice. The conjectures in Kotěšovec's 2018 OEIS entries, as inspected on 2 October 2026, assert $a_n^\pm \sim m_n^\pm$, where

$$m_n^+ = \frac{\exp(c_+ n^{2/3})}{2^{4/3} \sqrt{3} \pi^{1/6} n^{2/3}}, \quad (1.5)$$

$$m_n^- = \frac{\mathcal{G}^6 \exp(c_- n^{2/3} - 1/2)}{2 \cdot 3^{5/6} \pi^{1/3} n^{5/6}}, \quad (1.6)$$

and $\mathcal{G}$ is the Glaisher–Kinkelin constant [1, 2]. Define

$$\begin{aligned} B_+ &= -\frac{1}{2} \log 2, & d_+ &= 0, \\ B_- &= 6 \log \mathcal{G} - \frac{1}{2} - \frac{1}{2} \log(2\pi), & d_- &= \frac{1}{2}, \end{aligned} \quad (1.7)$$

$$R_\pm(t) = f_\pm(t) - A_\pm t^{-2} - d_\pm \log t - B_\pm. \quad (1.8)$$

These are exact definitions of $R_\pm$, not bounded-error assertions.

Theorem 1.1 (Summary of the asymptotic conclusions). *For each sequence separately the following statements hold.*

- (i) *The conjecture $a_n^\pm \sim m_n^\pm$ is false. More strongly, no constants $0 < C_1 < C_2 < \infty$ bound $a_n^\pm/m_n^\pm$ between C_1 and C_2 for every sufficiently large n .*
- (ii) *Unconditionally,*

$$\log a_n^\pm = c_\pm n^{2/3} + o(n^{1/3}),$$

whereas, for every $0 < \varepsilon < 1/4$, the error is not $O(n^{1/4-\varepsilon})$.

- (iii) *The Riemann hypothesis is equivalent to*

$$\text{for every } \varepsilon > 0, \quad \log a_n^\pm = c_\pm n^{2/3} + O_\varepsilon(n^{1/4+\varepsilon}) \quad (n \rightarrow \infty).$$

The constants in the O_ε notation may depend on the sign and on ε .

- (iv) *There is an unconditional corrected equivalent at an explicit point:*

$$a_n^\pm \sim \frac{(t_0^\pm)^2}{\sqrt{12\pi A_\pm}} \exp(f_\pm(t_0^\pm) + nt_0^\pm) = m_n^\pm \exp(R_\pm(t_0^\pm)). \quad (1.9)$$

The equality between the two right-hand expressions in (1.9) is exact. The equivalence has relative error $o(1)$, without a rate stated here. Its product value retains the arithmetic fluctuations. A more precise all-fixed-order expansion at the *exact* saddle, and its inverse consequences, appear in Sections 5 and 8. Section 9 also gives all fixed orders at the explicit point, retaining the exact mean mismatch and all odd correction grades.

1.1 Prior work and the scope of the contribution

The Dirichlet series of σ^* is classical; Tóth records it unnumbered in Section 4.9, printed page 28 [4]. His equation (50), on page 29, is an alternating series and is not the identity used below. Arithmetic partition products, nonreal zeta-zero corrections, saddle expansions and Mellin-based RH characterizations have substantial prior history.

Bodini, Duchon, Jacquot and Mutařchiev studied the totient-weight product $\prod(1 - q^m)^{-\varphi(m)}$, whose Mellin kernel has denominator $\zeta(s)$ [6]. Bureaux and Enriquez proved a zeta-zero-dependent equivalent and an RH equivalence for lattice convex chains, with main scale $n^{2/3}$ and logarithmic error exponent $1/6 + \varepsilon$ [5]. Their Section 3.2 is a direct antecedent for the Mellin-holomorphy converse used here; Section 4.1 explains why false pure equivalents can look convincing over very long numerical ranges. Their Section 4.2 also corrects a prefactor in the earlier totient treatment. We do not import an unchecked zero-residue sum or constants from that treatment.

Dong, Robles, Zaharescu and Zeindler study squarefree-part partitions with Dirichlet series $\zeta(s)/\zeta(2s)$ [7]. Their Theorem 1.11 relates logarithmic coefficient errors to zero locations; Lemma 6.8 and Section 6.6 distinguish retaining the exact saddle from absorbing zero effects into an explicit logarithmic error. General arbitrary-order infinite-product expansions are developed by Bridges, Brindle, Bringmann and Franke [8]; the reality-of-poles hypotheses (P1)–(P2) in their published theorem cannot simply be applied after discarding the nonreal poles present here. Ahmadi, Gómez-Aíza and Ward treat both signs and higher-order real poles [9]; Berndt, Robles, Zaharescu and Zeindler treat divisor-weight multiplicities [10]. Basak, Robles and Zaharescu give further denominator-zero formulas for Möbius-convolution products [11].

The contribution here is the sequence-specific correction of (1.5)–(1.6), with cancellation addressed, the precise $1/4 + \varepsilon$ RH criterion, and the accompanying exact-saddle and inverse formulas. A targeted literature search did not find an earlier treatment of these exact products and conclusions. That is a limited search observation, not an exhaustive priority claim or a claim of a new general Mellin/RH method.

[write] Added 2 October 2026, batch 77. This report is a single manuscript, printed in full: manuscript 70 of batch 77 of ProveIt’s incoming-reports intake, delivered as the archive `unitary-partition-report.zip` in commit 096ee7b87 (main file `unitary-partitions.tex`, now `article.tex`) and placed in commit d0e6008d9 in the research-report collection, among the OEIS-sequence asymptotics reports. The manuscript names no ProveIt commit and continues no repository path, so it has no pin; its author line (above) names neither an author nor a tool. Nothing in this report is formalized in Lean or Rocq, and placement in the collection confers no formal status on any statement here. The writing step prefixed every label with `udp:`, restored the diacritic in “Kotěšovec” (printed “Kotesovec” in the delivery), and added these **[write]** notes and one reference [14]; no statement, proof or number of the manuscript was altered.

Claims about OEIS text. Theorem 3.3 and Theorem 1.1(i) are this manuscript’s theorems about the conjectured equivalents printed in the OEIS entries A301981 and A301982, as the manuscript inspected them on 2 October 2026. They refute no statement of ProveIt: at intake an untruncated search of the research-report collection, the transseries volumes, `0eis/` and `docs/reports/` found no file naming either sequence or stating either formula, so nothing was retracted. The delivery does not say that a correction was submitted to the OEIS, and nothing here implies one. The refutation is analytic (the least-height noncancellation of Lemma 3.1); as Section 10 says, finite numerical agreement or disagreement with $m_n^\pm$ could not decide it.

The RH criterion is an equivalence. Theorem 7.4 (and Theorem 1.1(iii)) states that RH holds if and only if the logarithmic error is $O_\varepsilon(n^{1/4+\varepsilon})$ for every $\varepsilon > 0$; it neither proves nor disproves RH. RH is assumed only in Lemma 7.3 and in the direction “RH $\Rightarrow$ bound” of Theorem 7.4. Every other result of the report, including the refutation, the obstruction of Theorem 7.2, the explicit-point equivalent and all expansions and inverse brackets, is unconditional.

Neighbouring reports. The collection’s other partition-product reports, `a033552-catalan-partitions`, `a022629-distinct-partition-norms`, `a097356-sqrt-restricted-partitions` and `a238016-restricted-`

partitions-cubic-boundary, and the batch-77 reports `a174065-radix-layer-partitions` (where a pure OEIS equivalent likewise omits an oscillating factor, there a log-periodic one from Mellin poles on a vertical line) and `a372395-acyclic-orientation-partitions`, treat other products. None of them treats unitary-divisor weights, a zeta-zero denominator or an RH criterion, and no result is shared.

Notation. Several letters carry more than one meaning in the manuscript. Each is printed as delivered; the table fixes the meanings, with the tempting false readings. No symbol was renamed.

symbol	meaning here (and what it is <i>not</i>)
$c, c_{\pm}, c_*$	$c_{\pm} = 3(A_{\pm}/4)^{1/3}$, the growth constant of (1.4); $c_* > 2$ is a contour abscissa (Section 2); $c_1, \dots, c_4$ are positive constants in the angular estimates. None of them is another.
$C_j(t)$	the cumulant correction terms (5.6), functions of t . Not the constants $0 < C_1 < C_2$ of Theorem 1.1(i), the C_r of (2.3), or the $C_M, \widehat{C}_M$ of the error and bracket bounds.
$A, A_{\pm}$	$A_- = \pi^2/6, A_+ = \pi^2/8$; in Lemmas 3.2 and 7.1 $A > 0$ is arbitrary (applied with $A = A_{\pm}$).
$K, K_r(s)$	$K > 0$ is the amplitude in Lemma 3.2; K_r the Mellin kernels (2.7); $K = 2M + 2$ in the proofs of Theorems 5.2 and 9.1; $K > 4\pi$ in the proof of Lemma 5.1.
$\mathcal{G}, G_M$	the Glaisher–Kinkelin constant; the exact-saddle model (8.1). Unrelated (the manuscript says so).
$\varepsilon, \epsilon_M, \varepsilon$	ε an arbitrarily small exponent; $\epsilon_M(y)$ the bracket half-width of Theorem 8.2; $\varepsilon = \pm 1$ the sign in (10.4), where $\varepsilon = 1$ is the <i>minus</i> product. It is not the subscript sign $\pm$.
δ	the mean mismatch $\delta(x)$ of (9.1) (Section 9); the error exponent in Lemma 7.1; the strip width in Lemma 7.3.
H, H_d	a holomorphic function in (3.2); the probabilists’ Hermite polynomials (Section 9).
E, E_1, E_r	the auxiliary exponential $E(u)$ in the proof of Theorem 5.2; the boundary errors $E_1, E_r = \kappa_r - A(r+1)!t^{-r-2}$.
$R_{\pm}, \mathcal{R}$	$R_{\pm}(t)$ is the exact remainder (1.8), which is <i>not</i> bounded (that is the content of Theorem 3.3); $\mathcal{R}$ a Taylor remainder; R_*, r_* disc radii.
$m_n^{\pm}, m_r$	the refuted OEIS models (1.5)–(1.6); m_r multiplicities in (5.6), (9.2).
$d_{\pm}, d_r, D$	$d_{\pm}$ the logarithmic coefficient in (1.7); d_r the tail constants in (10.3); $D(s)$ the Dirichlet series (2.4) and $D = \sum r m_r$ a degree.
σ, σ^*	$\sigma(m)$ the ordinary divisor sum in (2.1), $\sigma^*(m) = b_m$ the unitary one; $\sigma \in (3/4, 1)$ a contour abscissa in the proof of Theorem 7.4.
t_x, t_n, t_0	t_x the exact saddle (5.1), t_n its value at $x = n$; $t_0 = t_0(n) = (2A/n)^{1/3}$ the explicit point. t_0 is <i>not</i> t_x at $x = 0$.
$N(y), N$	the integer threshold of Theorem 8.2; $N = \lfloor 1/t \rfloor$ in the proof of Lemma 5.1.
L, L_M	the product cutoff length (Section 10); the explicit-point model (9.6).
P_r, P_M	the cumulant polynomials (10.4); a polynomial bound in (5.11).
a_n, a_w, b_m	$a_n = a_n^{\pm}$ the coefficients; $a_w(t)$ standardized cumulants in the proof of Theorem 5.2; $b_m = \sigma^*(m)$ the colour count. The interval ends $a < 2A < b$ in the proof of Lemma 7.1 are neither.
ϕ_t, ϕ	the characteristic function (5.3); $\phi(v) = cv^{2/3} - v$ in the proof of Lemma 3.2; φ Euler’s totient (Section 1.1).

[write] **Added 3 October 2026, batch 86.** Since 3 October 2026 this report has two Parts. Part I (Sections 1–11) is the batch-77 manuscript described in the preceding note, printed as before: its sections, equations, theorems and labels keep their numbers, so “Section 11, question 3”, “Theorem 3.3”, “Lemma 4.1”, “Theorem 6.1” and “Theorem 8.2” still mean what the manuscript of Part II cites. The preceding note’s “single manuscript, printed in full” and “no pin” now describe Part I; its statement that no neighbouring report shares a result remains true. Part II (Sections 12 and 13) is the partition subject of manuscript 01 of batch 86, prepared with ChatGPT, unrefereed and not formalized, pinned at commit `eaf08931c`. It answers question 3 of Section 11: for each sign $\liminf a_n^{\pm}/m_n^{\pm} = 0$ and $\limsup a_n^{\pm}/m_n^{\pm} = \infty$, with two-sided logarithmic oscillations of order at least $n^{1/4}$ (Theorem 12.5). Part II’s symbols $m, W, s_*, \sigma_*, C_*, C_{\pm}, K_{\pm}, L_{\pm}, h, H$ and $T_{\pm}$ do not mean what the table above might suggest; in particular its $W(n)$ is not the Lambert W function. A separate table at the start of Part II fixes them.

2 Arithmetic and Mellin preliminaries

2.1 Convergence and elementary estimates

The elementary bounds

$$m \leq b_m \leq \sigma(m), \quad \sum_{m \leq X} b_m \leq \sum_{d \leq X} d \lfloor X/d \rfloor \leq X^2 \quad (X \geq 1) \quad (2.1)$$

will be useful throughout. They prove absolute convergence of the logarithmic product series for $\operatorname{Re} t > 0$, locally uniformly, and hence analyticity there. For every fixed nonnegative integer r , partial summation gives

$$\sum_{m \geq 1} b_m m^r e^{-mu} = O_r(u^{-r-2}) \quad (u > 0). \quad (2.2)$$

For $u \geq 1$ the exponential decay can be absorbed into the same bound. The logarithmic expansions are

$$f_-(t) = \sum_{m, \ell \geq 1} \frac{b_m}{\ell} e^{-m\ell t}, \quad f_+(t) = \sum_{m, \ell \geq 1} \frac{(-1)^{\ell+1} b_m}{\ell} e^{-m\ell t}.$$

In particular $f_+(t) = f_-(t) - f_-(2t)$. After r derivatives, (2.2) leaves the convergent sum $\sum_{\ell \geq 1} \ell^{-3}$. Thus for every fixed $r \geq 0$,

$$|f^{(r)}(w)| \leq C_r t^{-r-2} \quad (0 < t \leq 1, \operatorname{Re} w \geq t/2). \quad (2.3)$$

No sign condition on higher cumulants is needed for this estimate.

2.2 The arithmetic Dirichlet series and the kernels

For $\operatorname{Re} s > 2$, the prime-power Euler factor is

$$1 + \sum_{a \geq 1} (1 + p^a) p^{-as} = \frac{1 - p^{1-2s}}{(1 - p^{-s})(1 - p^{1-s})}.$$

Consequently the classical identity [4, Section 4.9, p. 28] is

$$D(s) = \sum_{m \geq 1} \frac{b_m}{m^s} = \frac{\zeta(s)\zeta(s-1)}{\zeta(2s-1)}. \quad (2.4)$$

Let $\chi_-(s) = 1$ and $\chi_+(s) = 1 - 2^{-s}$. Absolute convergence permits termwise Mellin integration and gives

$$M_{\pm}(s) := \int_0^\infty f_{\pm}(t) t^{s-1} dt = \Gamma(s) \chi_{\pm}(s) \frac{\zeta(s+1)\zeta(s)\zeta(s-1)}{\zeta(2s-1)} \quad (\operatorname{Re} s > 2). \quad (2.5)$$

Mellin inversion on any line $\operatorname{Re} s = c_* > 2$ follows either from the standard inverse formula for e^{-t} and absolute summation, or directly from the absolutely convergent kernels. To avoid confusion, c_* here is a contour location, whereas c in (1.4) is the coefficient growth constant.

Define the exact cumulants, including $r = 0$, by

$$\kappa_r(t) = (-1)^r f^{(r)}(t), \quad V(t) = \kappa_2(t). \quad (2.6)$$

Differentiation of Mellin inversion gives the useful kernels

$$K_r(s) = \Gamma(s+r)\chi(s) \frac{\zeta(s+1)\zeta(s)\zeta(s-1)}{\zeta(2s-1)}, \quad (2.7)$$

$$\kappa_r(t) = \frac{1}{2\pi i} \int_{c_*-i\infty}^{c_*+i\infty} K_r(s) t^{-s-r} ds. \quad (2.8)$$

The pole at $s = 2$ has residue $A(r+1)!$. At $s = 1$ the two zeta poles cancel:

$$\frac{\zeta(s)}{\zeta(2s-1)} \longrightarrow 2, \quad K_r(1) = -r!\chi(1)\zeta(2). \quad (2.9)$$

For orientation, the principal residue at zero produces the constants in (1.7). Indeed,

$$D(0) = -\frac{1}{2}, \quad D'(0) = -\frac{1}{2} \log(2\pi) - 6\zeta'(-1), \quad \zeta'(-1) = \frac{1}{12} - \log \mathcal{G}.$$

The s^{-1} coefficient in $\Gamma(s)\zeta(s+1)$ vanishes. Thus the double pole of M_- contributes $\frac{1}{2} \log t + B_-$, and subtraction at $2t$ contributes B_+ for the plus case. These are only principal residue terms. Nonreal poles obstruct a bounded remainder, as we now prove.

3 Noncancellation and the false pure equivalents

We use the classical facts that nontrivial zeta zeros exist, are isolated, have $0 < \operatorname{Re} \rho < 1$, have nonzero imaginary part, and are symmetric about the critical line and the real axis [13]. No numerical zero location is needed.

Lemma 3.1 (A surviving denominator pole). *Both kernels M_+ and M_- have a genuine pole at some s_0 with $3/4 \leq \operatorname{Re} s_0 < 1$ and $\operatorname{Im} s_0 > 0$.*

Proof. Choose a nontrivial zero $\rho = \beta + i\gamma$ of least positive imaginary part, reflecting in the critical line if necessary so that $\beta \geq 1/2$. Such a least positive height exists by isolation, compactness of bounded-height portions of the critical strip, and absence of real nontrivial zeros. Set $s_0 = (1 + \rho)/2$. The denominator in (2.5) vanishes at s_0 to the multiplicity of ρ .

Every numerator factor is analytic and nonzero at this point. The gamma function has no zeros and no pole there. The argument $s_0 + 1$ has real part greater than one. The factor $\zeta(s_0)$ cannot vanish, because its positive imaginary part is $\gamma/2$, below the least positive zero height. The number $s_0 - 1$ has real part strictly between $-1/2$ and 0 and nonzero imaginary part, so is neither a trivial nor a nontrivial zero. Finally $|2^{-s_0}| < 1$. The pole therefore survives with the full multiplicity of ρ . $\square$

Lemma 3.2 (Abelian equivalent). *Let $A, K > 0$, $p \in \mathbb{R}$ and $c = 3(A/4)^{1/3}$. As $t \downarrow 0$,*

$$\sum_{n \geq 1} K n^{-p} e^{cn^{2/3} - tn} \sim K \sqrt{12\pi A} (2A)^{-p} t^{3p-2} e^{A/t^2}. \quad (3.1)$$

An eventual relative $o(1)$ perturbation of the summands' coefficients preserves the equivalent.

Proof. Write $v = t^3 n$ and $\phi(v) = cv^{2/3} - v$. Its unique maximum is at $v_0 = 2A$, where $\phi(v_0) = A$ and $\phi''(v_0) = -1/(3v_0)$. With $v = v_0 + tu$ the exponent is $A/t^2 - u^2/(6v_0) + O(t|u|^3)$ on bounded u intervals. The mesh in u is t^2 , so the Gaussian Riemann sum yields

$$K t^{3p-2} v_0^{-p} e^{A/t^2} \int_{\mathbb{R}} e^{-u^2/(6v_0)} du,$$

which is (3.1). A fixed neighborhood of v_0 supplies Gaussian domination; outside it ϕ loses a fixed amount. Near $v = 0$, where v^{-p} may be singular, first remove finitely many n , then bound the remaining polynomial factors by powers of t^{-1} and n . At infinity, the negative linear term in ϕ controls every such factor. These ranges are exponentially smaller than the saddle contribution. This also proves the required Riemann-sum tail estimates.

For a relative coefficient perturbation, choose a large fixed index beyond which it lies between $1 - \eta$ and $1 + \eta$. Positivity sandwiches the tail sums; the finite prefix is negligible. Let $\eta \downarrow 0$. $\square$

Theorem 3.3 (Refutation of the recorded equivalents). *Neither $a_n^+ \sim m_n^+$ nor $a_n^- \sim m_n^-$ holds. For either sign, the ratio cannot eventually remain in a compact subinterval of $(0, \infty)$.*

Proof. Applying Lemma 3.2 to the constants in (1.5)–(1.6) shows that a proposed coefficient equivalent would imply

$$f_{\pm}(t) = A_{\pm}t^{-2} + d_{\pm} \log t + B_{\pm} + o(1).$$

Specifically the Abelian prefactor is $2^{-1/2}$ for the plus case and $\mathcal{G}^6 e^{-1/2}/\sqrt{2\pi}$ for the minus case. Eventual bounds $C_1 \leq a_n^{\pm}/m_n^{\pm} \leq C_2$ likewise imply the same formula with $O(1)$ instead of $o(1)$, by positivity; finite initial terms are negligible.

If $R_{\pm}$ in (1.8) were bounded near zero, split its Mellin integral at one. The tail $\int_1^{\infty} f(t)t^{s-1} dt$ is entire, since f decays exponentially. Uniform integrability on compact subsets gives a holomorphic function H on $\operatorname{Re} s > 0$ such that

$$M(s) = \frac{A}{s-2} - \frac{d}{s^2} + \frac{B}{s} + H(s). \quad (3.2)$$

This agrees with (2.5) on $\operatorname{Re} s > 2$ and hence by uniqueness of meromorphic continuation wherever both sides are defined. It forbids any nonreal pole in $\operatorname{Re} s > 0$, contradicting Lemma 3.1. $\square$

Remark 3.4. The theorem does not determine which one-sided divergence occurs. In particular, it does not assert both $\liminf a_n/m_n = 0$ and $\limsup a_n/m_n = \infty$. No two-sided oscillation theorem is being used.

[write] **Added 3 October 2026, batch 86.** Part II settles the alternative that this remark leaves open: both occur. For each sign, $\liminf a_n^{\pm}/m_n^{\pm} = 0$ and $\limsup a_n^{\pm}/m_n^{\pm} = \infty$ (12.3); more precisely $\log(a_n^{\pm}/m_n^{\pm})$ lies above $(K_{\pm} - \varepsilon)W(n)$ and below $-(K_{\pm} - \varepsilon)W(n)$ infinitely often, with $W(n) = n^{\sigma_*/3}(\log n)^{m-1} \geq n^{1/4}$ (Theorem 12.5). The proof uses the pole of Lemma 3.1 together with its order m and its leading Laurent coefficient (Lemma 12.4), and Landau's positivity principle (Lemmas 12.2 and 12.3); it assumes neither RH nor simple zeros. The remark remains true of Theorem 3.3 by itself. Part II is unrefereed, prepared with AI assistance and not formalized.

4 Unconditional contour estimates

An exact shift to the line $\operatorname{Re} s = 1$ sharpens what a shift merely to $\operatorname{Re} s > 1$ supplies.

Lemma 4.1 (Derivative estimates on the boundary line). *For each fixed integer $r \geq 0$, unconditionally,*

$$\kappa_r(t) = A(r+1)!t^{-r-2} + o(t^{-r-1}) \quad (t \downarrow 0). \quad (4.1)$$

Proof. We need a reciprocal-zeta estimate uniform on a closed strip. Trudgian's unconditional bound [12] states

$$\left| \frac{1}{\zeta(\sigma + iT)} \right| \leq 6.9 \times 10^6 \log T \quad (T \geq 45, \sigma \geq 1 - (8 \log T)^{-1}). \quad (4.2)$$

It covers $\sigma \geq 1$ in particular. Conjugation handles negative heights. At bounded heights on a fixed closed strip, the classical zero-free line and continuity give boundedness, interpreting $1/\zeta$ at its analytic zero at one. Any uniform polynomial bound on this half-plane would suffice for the proof.

On $1 \leq \operatorname{Re} s \leq c_*$, the denominator argument $2s - 1$ has real part at least one. There are no denominator-zero poles. The point $s = 1$ is removable by (2.9), and the only pole crossed in (2.8) is at $s = 2$. Standard polynomial vertical bounds for the other zeta factors, together with (4.2) and Stirling's exponential factor $e^{-\pi|\operatorname{Im} s|/2}$, make the horizontal sides of the contour vanish. The shifted kernel is absolutely integrable, including at zero height because the cancellation at one was treated as a quotient. Thus

$$\kappa_r(t) - A(r+1)!t^{-r-2} = \frac{t^{-r-1}}{2\pi} \int_{\mathbb{R}} K_r(1+iu)e^{iu \log(1/t)} du. \quad (4.3)$$

Here $K_r(1+iu) \in L^1(\mathbb{R})$. Apply the Riemann–Lebesgue lemma separately for each fixed r to obtain (4.1). No contour below one or sum over zeros has been used. $\square$

5 The exact saddle and all fixed orders

For each $x > 0$, define t_x by the exact equation

$$\kappa_1(t_x) = x. \quad (5.1)$$

To see existence and uniqueness, use independent geometric random variables for each color in the minus case, with parameter e^{-mt} and weight m . In the plus case use independent Bernoulli variables with success probability $e^{-mt}/(1+e^{-mt})$ and weight m . The resulting nonnegative integer random variable S_t has finite expectation and variance; finiteness follows from (2.1). Its expectation is $\kappa_1(t)$ and its variance is $V(t) > 0$. Hence $\kappa'_1(t) = -V(t) < 0$. Its limits are infinity and zero at the two endpoints, by Lemma 4.1 and the product. In particular,

$$t_x \sim (2A/x)^{1/3}, \quad V(t) \sim 6At^{-4}, \quad \frac{dt_x}{dx} = -\frac{1}{V(t_x)}. \quad (5.2)$$

The centered characteristic function is

$$\phi_t(\theta) = \exp(f(t - i\theta) - f(t) - i\kappa_1(t)\theta). \quad (5.3)$$

At $t = t_n$, Fourier inversion gives

$$a_n = \frac{e^{f(t_n)+nt_n}}{2\pi} \int_{-\pi}^{\pi} \phi_{t_n}(\theta) d\theta. \quad (5.4)$$

5.1 A uniform angular estimate

Lemma 5.1. *For small $t > 0$, uniformly in $|\theta| \leq \pi$,*

$$|\phi_t(\theta)| \leq \exp\{-c_1 \min(t^{-4}\theta^2, t^{-2})\} \quad (5.5)$$

with a positive constant c_1 depending only on the sign.

Proof. Let $N = \lfloor 1/t \rfloor$ and keep only factors with $N \leq m \leq 2N$. There $q = e^{-mt}$ stays in a fixed compact subset of $(0, 1)$. A Bernoulli factor has squared modulus

$$\left| \frac{1 + qe^{im\theta}}{1 + q} \right|^2 = 1 - \frac{4q}{(1 + q)^2} \sin^2(m\theta/2),$$

and a geometric factor has squared modulus

$$\left| \frac{1 - q}{1 - qe^{im\theta}} \right|^2 = \left(1 + \frac{4q}{(1 - q)^2} \sin^2(m\theta/2) \right)^{-1}.$$

The corresponding logarithms are at most $-c_2 \sin^2(m\theta/2)$ uniformly. Raising them to $b_m \geq m$ proves

$$\log |\phi_t(\theta)| \leq -c_3 \sum_{m=N}^{2N} m \sin^2(m\theta/2).$$

For $|\theta| \leq 1/N$, the sine arguments have modulus at most one, so this sum is at least a constant times $N^4 \theta^2$. For $1/N \leq |\theta| \leq K/N$, divide the sum by N^2 and write $u = N|\theta|$. The Riemann sums converge uniformly for $u \in [1, K]$ to

$$\int_1^K v \sin^2(vu/2) dv.$$

This continuous function is strictly positive on that compact interval: zero would require a sine to vanish on a full interval. The sum is therefore at least a positive constant times N^2 .

Finally take $K > 4\pi$ and $K/N \leq |\theta| \leq \pi$. The geometric-sum estimate gives

$$\left| \sum_{m=N}^{2N} e^{im\theta} \right| \leq \frac{1}{|\sin(\theta/2)|} \leq \frac{\pi}{|\theta|} \leq \frac{\pi N}{K}.$$

Thus $\sum_{m=N}^{2N} \sin^2(m\theta/2) \geq N/4$ for sufficiently large N . Multiplying by $m \geq N$ finishes the last range. Since $N \asymp 1/t$, all three estimates combine into (5.5). $\square$

5.2 Finite cumulant polynomials

Set $C_0(t) = 1$. For $j \geq 1$, define

$$C_j(t) = \sum_{\substack{m_3, \dots, m_{2j+2} \geq 0 \\ \sum_{r=3}^{2j+2} (r-2)m_r = 2j}} (-1)^{D/2} (D-1)!! \prod_{r=3}^{2j+2} \frac{1}{m_r!} \left(\frac{\kappa_r(t)}{r! V(t)^{r/2}} \right)^{m_r}, \quad D = \sum_{r=3}^{2j+2} r m_r. \quad (5.6)$$

The constraint makes D even. The sum is finite, and (2.3) and (5.2) imply $C_j(t) = O_j(t^{2j})$. The first two are

$$C_1 = \frac{\kappa_4}{8V^2} - \frac{5\kappa_3^2}{24V^3}, \quad (5.7)$$

$$C_2 = -\frac{\kappa_6}{48V^3} + \frac{7\kappa_3\kappa_5}{48V^4} + \frac{35\kappa_4^2}{384V^4} - \frac{35\kappa_3^2\kappa_4}{64V^5} + \frac{385\kappa_3^4}{1152V^6}. \quad (5.8)$$

All cumulants in these formulas are exact product derivatives.

Theorem 5.2 (All-fixed-order exact-saddle expansion). *For every fixed integer $M \geq 0$ and either sign,*

$$a_n = \frac{\exp(f(t_n) + nt_n)}{\sqrt{2\pi V(t_n)}} \left\{ \sum_{j=0}^M C_j(t_n) + O_M(t_n^{2M+2}) \right\}. \quad (5.9)$$

The expansion is valid with $n \rightarrow \infty$ after M is fixed; uniformity as $M \rightarrow \infty$ is not asserted.

Proof. Put $K = 2M + 2$, $z = \theta\sqrt{V}$, and choose a fixed $0 < \eta < 1$. We first work on $|z| \leq t^{-\eta}$. Taylor expansion of $f(t - i\theta)$ through degree $K + 1$, using (2.3), gives

$$\log \phi_t(z/\sqrt{V}) = -z^2/2 + \sum_{w=1}^{K-1} t^w a_w(t) (iz)^{w+2} + \mathcal{R}, \quad |\mathcal{R}| \leq C_M t^K |z|^{K+2}, \quad (5.10)$$

where

$$a_w(t) = \frac{\kappa_{w+2}(t)}{(w+2)!V(t)^{(w+2)/2}t^w}$$

are uniformly bounded for the finitely many indices involved. The remainder follows because $V \asymp t^{-4}$, so the $(K+2)$ nd derivative term is $O(t^{-K-4}\theta^{K+2}) = O(t^K|z|^{K+2})$.

For a precise expansion by weighted degree, introduce a real auxiliary variable u and the finite exponential

$$E(u) = \exp \left\{ \sum_{w=1}^{K-1} u^w a_w(t) (iz)^{w+2} \right\}.$$

For $0 \leq u \leq t$ and $|z| \leq t^{-\eta}$ the exponent has modulus at most

$$C_M z^2 \sum_{w=1}^{K-1} (t|z|)^w = o(1)z^2.$$

Every derivative in u of fixed order is a polynomial in derivatives of that exponent times $E(u)$. Because each exponent derivative is a finite sum of nonnegative powers of u times polynomials in z , it is bounded by a fixed polynomial in $|z|$ for $0 \leq u \leq t \leq 1$. For small enough t , Taylor's integral remainder for E through degree $K-1$, evaluated at $u = t$, is therefore bounded by

$$C_M t^K P_M(|z|) e^{z^2/8} \quad (5.11)$$

for a fixed polynomial P_M . Also $|\mathcal{R}| \leq C_M z^2 (t|z|)^K = o(1)z^2$, and $|e^{\mathcal{R}} - 1| \leq |\mathcal{R}|e^{|\mathcal{R}|}$. Incorporating this factor in (5.11) replaces $e^{z^2/8}$ by at most $e^{z^2/4}$ and possibly enlarges the polynomial. Multiplication by $e^{-z^2/2}$ is integrable uniformly. Hence the integrated remainder on the local interval is $O_M(t^K)$.

The coefficients of u^k in E are exactly the terms with $\sum (r-2)m_r = k$. Since r and $r-2$ have the same parity, odd k gives an odd imaginary polynomial in real z . It integrates to zero on the symmetric interval. Extending each retained polynomial integral to all of $\mathbb{R}$ adds an exponentially small Gaussian tail. The even moments

$$\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} z^D e^{-z^2/2} dz = (D-1)!!$$

then produce precisely (5.6), including its factor $i^D = (-1)^{D/2}$.

It remains to justify discarding the other Fourier angles. By (5.5) and $V \asymp t^{-4}$, throughout $|z| > t^{-\eta}$ the characteristic function is at most $\exp(-c_4 t^{-2\eta})$ in modulus. The entire z interval has length $O(t^{-2})$, so its integral is smaller than any fixed power of t . Substituting the local integral into (5.4) proves (5.9). $\square$

6 An equivalent at the explicit leading saddle

Theorem 6.1. For $t_0 = (2A/n)^{1/3}$,

$$t_n - t_0 = o(t_0^2), \quad a_n \sim \frac{t_0^2}{\sqrt{12\pi A}} \exp(f(t_0) + nt_0).$$

Consequently (1.9) holds and $\log a_n = cn^{2/3} + o(n^{1/3})$ unconditionally.

Proof. By Lemma 4.1,

$$\kappa_1(t) = 2At^{-3} + E_1(t), \quad E_1(t) = o(t^{-2}).$$

Since $\kappa_1(t_n) = n = 2At_0^{-3}$ and $t_n \sim t_0$, the mean value theorem gives

$$6A\xi^{-4}|t_n - t_0| = |E_1(t_n)| = o(t_n^{-2})$$

for some ξ between t_n and t_0 . Therefore $t_n - t_0 = o(t_0^2)$.

Let $\Phi_n(t) = f(t) + nt$. The exact saddle is stationary: $\Phi'_n(t_n) = 0$. Its second derivative is $V(t) > 0$. Uniformly between t_n and t_0 , $V(t) = O(t_0^{-4})$, so Taylor's theorem gives

$$0 \leq \Phi_n(t_0) - \Phi_n(t_n) \leq \frac{1}{2} \sup_{u \text{ between } t_n, t_0} V(u) (t_n - t_0)^2 = o(1).$$

Also $V(t_n) \sim 6At_0^{-4}$. The $M = 0$ case of Theorem 5.2 now proves the equivalent. Replacing the two large exponent terms separately would not justify this step; stationarity of their sum is essential.

Finally $At_0^{-2} + nt_0 = cn^{2/3}$. The $r = 0$ case of (4.1) implies $f(t_0) = At_0^{-2} + o(t_0^{-1})$, and the logarithm of the prefactor is $O(\log n)$. This yields the logarithmic refinement. Substituting the definitions (1.7)–(1.8) into the displayed equivalent verifies its exact algebraic reformulation in (1.9). $\square$

Remark 6.2. The Riemann–Lebesgue step gives no stated rate of convergence. Thus this theorem has only a relative $o(1)$ remainder. It does not replace the quantified higher-order and inverse results that use t_n . The correction $R(t_0)$ is computable by a convergent positive-real product; it is not defined by a zero-residue series.

7 The logarithmic obstruction and the RH equivalence

Lemma 7.1 (Abelian transfer of logarithmic errors). *Let positive coefficients satisfy*

$$\log a_n = cn^{2/3} + O(n^\delta), \quad 0 < \delta < 2/3, \quad c = 3(A/4)^{1/3}.$$

Then

$$\log \sum_{n \geq 0} a_n e^{-tn} = At^{-2} + O(t^{-3\delta}). \quad (7.1)$$

Proof. Fix $0 < a < 2A < b$ such that outside $a \leq v \leq b$ the function $cv^{2/3} - v$ is at most $A - \eta_0$ for some $\eta_0 > 0$. On $at^{-3} \leq n \leq bt^{-3}$ the perturbation in the logarithm is $O(t^{-3\delta})$. The unperturbed sum on this interval is $\exp(At^{-2} + O(\log(1/t)))$ by the same Gaussian localization as Lemma 3.2; this supplies both lower and upper bounds on its logarithm.

Outside this interval, the lower-order term Cn^δ cannot undo the fixed saddle loss as $t \downarrow 0$. More explicitly, on the low range it is at most $Ca^\delta t^{-3\delta} = o(t^{-2})$, and on the high range $n^\delta = o(n^{2/3})$ uniformly once $n \geq bt^{-3}$. Increase b if necessary so the negative $-tn$ term controls that tail and its sum. The intervening compact v ranges retain a fixed loss. Polynomial counting factors contribute $O(\log(1/t))$, absorbed by $t^{-3\delta}$ because $\delta > 0$. Finite initial coefficients are negligible. Combining the bounds proves (7.1). $\square$

Theorem 7.2 (Unconditional lower obstruction). *For either sign and every $0 < \varepsilon < 1/4$,*

$$\log a_n^\pm - c_\pm n^{2/3} \neq O(n^{1/4-\varepsilon}).$$

Proof. If this bound held, Lemma 7.1 with $\delta = 1/4 - \varepsilon$ would give $f(t) - At^{-2} = O(t^{-3/4+3\varepsilon})$. Splitting the Mellin integral at one then makes $M(s) - A/(s-2)$ holomorphic on $\operatorname{Re} s > 3/4 - 3\varepsilon$. This contradicts the pole of Lemma 3.1, whose real part is at least $3/4$. $\square$

[write] **Added 3 October 2026, batch 86.** Part II makes this obstruction two-sided: by Theorem 12.5, $\log a_n^\pm - c_\pm n^{2/3}$ is $\Omega_\pm(n^{1/4})$, that is, it exceeds a positive multiple of $n^{1/4}$ and falls below a negative multiple infinitely often, which contains the present theorem. With Theorem 7.4 below, under RH the error is both $\Omega_\pm(n^{1/4})$ and $O_\varepsilon(n^{1/4+\varepsilon})$ (the note after Theorem 12.5).

7.1 A conditional reciprocal zeta bound

We provide the conditional growth estimate used in the contour argument rather than presupposing a sharp RH bound.

Lemma 7.3. *Assume RH. For each $0 < \delta < 1/2$, the reciprocal $1/\zeta(w)$ has at most polynomial vertical growth on each fixed closed vertical strip with $\operatorname{Re} w \geq 1/2 + \delta$, interpreting its value at $w = 1$ by continuation.*

Proof. For large $|T|$, center a disc at $w_c = 2 + iT$, with radii

$$R_* = 3/2 - \delta/2, \quad r_* = 3/2 - \delta.$$

Under RH the outer disc has no zero, since its real parts exceed $1/2 + \delta/2$; it also has no pole at these heights. Choose an analytic logarithm of ζ on the disc, agreeing at the center with the bounded Euler-product logarithm. The elementary continuation formula

$$\zeta(w) = \frac{w}{w-1} - w \int_1^\infty \{u\} u^{-w-1} du \quad (\operatorname{Re} w > 0)$$

shows $|\zeta(w)| = O_\delta(1 + |T|)$ uniformly on the outer disc. Therefore $\operatorname{Re} \log \zeta(w) \leq C_\delta \log(2 + |T|)$ there. Borel–Carathéodory bounds the modulus of this analytic logarithm throughout the inner disc by $O_\delta(\log(2 + |T|))$. At its leftmost point $1/2 + \delta + iT$, exponentiation gives

$$|1/\zeta(1/2 + \delta + iT)| \leq (2 + |T|)^{C'_\delta}.$$

The same inner disc covers all real parts between this value and 2 at height T . For real parts at least two the Euler product supplies a uniform bound. At bounded heights use compactness and zero-freeness, treating the reciprocal's zero at one analytically. No global branch of $\log \zeta$ through its pole is required. $\square$

Theorem 7.4 (A sequence-specific RH criterion). *For either choice of sign, considered separately, RH holds if and only if*

$$\forall \varepsilon > 0, \quad \log a_n^\pm = c_\pm n^{2/3} + O_\varepsilon(n^{1/4+\varepsilon}). \quad (7.2)$$

Proof. First assume (7.2). Take arbitrarily small positive ε with $1/4 + \varepsilon < 2/3$. Lemma 7.1 and the Mellin splitting argument continue $M(s) - A/(s-2)$ holomorphically to $\operatorname{Re} s > 3/4 + 3\varepsilon$. These continuations agree on their overlaps, giving holomorphy throughout $\operatorname{Re} s > 3/4$.

Suppose RH fails. By symmetry there is a zero with real part greater than $1/2$ and positive imaginary part. Choose such a zero ρ at the least positive height among this subset. At $s_0 = (1+\rho)/2$,

we have $\operatorname{Re} s_0 > 3/4$. As in Lemma 3.1, every numerator factor except possibly $\zeta(s_0)$ is nonzero. If $\zeta(s_0) = 0$, it is itself an RH-violating zero at half the selected height, a contradiction. Thus M has a genuine pole strictly right of $3/4$, contradicting holomorphy. This proves RH, using symmetry. The argument deliberately does not claim that every denominator zero is uncanceled.

Conversely assume RH and fix $3/4 < \sigma < 1$. Lemma 7.3, with $\delta = 2\sigma - 3/2$, supplies polynomial growth for $1/\zeta(2s - 1)$ on the relevant closed contour strip. Stirling decay and polynomial bounds for the other zeta factors justify shifting (2.8) to $\operatorname{Re} s = \sigma$, including vanishing horizontal segments and differentiation to every fixed order. The point one is removable and the only pole crossed is at two. Hence

$$\kappa_r(t) = A(r+1)!t^{-r-2} + O_{\sigma,r}(t^{-r-\sigma}). \quad (7.3)$$

The saddle equation gives

$$t_n = t_0\{1 + O(t_0^{2-\sigma})\}.$$

The function $nt + At^{-2}$ is stationary at t_0 . Its change from t_0 to t_n is $O(t_0^{2-2\sigma})$, while the remainder in $f(t_n)$ is $O(t_0^{-\sigma})$. The $M = 0$ formula of Theorem 5.2 contributes only $O(\log n)$ from its prefactor and $O(t_n^2)$ from its relative error. Therefore

$$\log a_n = cn^{2/3} + O_\sigma(n^{\sigma/3}).$$

Given any $\varepsilon > 0$, choose $\sigma \in (3/4, 1)$ with $\sigma/3 < 1/4 + \varepsilon$. This proves (7.2). $\square$

Theorem 7.4 is an equivalence to RH, not a resolution of it. Neither direction assumes simple zeros, a numerical zero certification, or absolute convergence of a zero-residue sum. Together with Theorems 3.3, 6.1 and 7.2, it proves Theorem 1.1.

8 Model inverses and integer thresholds

Define $S_M(t) = \sum_{j=0}^M C_j(t)$, which is positive for sufficiently small t because $S_M(t) = 1 + O(t^2)$. For large real x put

$$G_M(x) = f(t_x) + xt_x - \frac{1}{2} \log(2\pi V(t_x)) + \log S_M(t_x). \quad (8.1)$$

The notation G_M is unrelated to the Glaisher constant $\mathcal{G}$.

Lemma 8.1. *For fixed M ,*

$$G'_M(x) = t_x - \frac{\kappa_3(t_x)}{2V(t_x)^2} + O_M(t_x^5) = t_x + O_M(t_x^3) > 0 \quad (8.2)$$

for large x . For $M = 0$ the first equality has no $O(t_x^5)$ term.

Proof. The exact identities $\kappa'_r = -\kappa_{r+1}$ and $V' = -\kappa_3$ show that differentiation of a standardized cumulant costs at most one power of t . Termwise differentiation in the finite sum (5.6) gives $C'_j(t) = O(t^{2j-1})$. Thus $S'_M(t)/S_M(t) = O_M(t)$, absent for $M = 0$.

When differentiating $f(t_x) + xt_x$, the terms involving t'_x cancel by the saddle equation and the result is t_x . Using $t'_x = -1/V$ for the other terms gives exactly

$$G'_M(x) = t_x - \frac{\kappa_3}{2V^2} - \frac{S'_M}{VS_M}.$$

The stated bounds follow from $V \asymp t^{-4}$ and $\kappa_3 = O(t^{-5})$. $\square$

Theorem 8.2 (Eventual increase and inverse localization). *For each sign the sequence a_n is eventually strictly increasing. For sufficiently large y , define the integer threshold*

$$N(y) = \min\{n \geq 0 : a_n \geq y\}.$$

For fixed $M \geq 0$, there is a unique sufficiently large solution $x_M(y)$ of

$$G_M(x_M(y)) = \log y. \quad (8.3)$$

There are constants $C_M > 0$ and Y_M such that for $y \geq Y_M$, on writing

$$\epsilon_M(y) = C_M t_{x_M(y)}^{2M+1} = O(x_M(y)^{-(2M+1)/3}),$$

one has

$$\lceil x_M(y) - \epsilon_M(y) \rceil \leq N(y) \leq \lceil x_M(y) + \epsilon_M(y) \rceil. \quad (8.4)$$

In particular $N(y) \sim (\log y/c)^{3/2}$.

Proof. Theorem 5.2 gives, for integer $n \rightarrow \infty$,

$$\log a_n - G_M(n) = O_M(t_n^{2M+2}). \quad (8.5)$$

At $M = 0$, subtract the formulas at $n + 1$ and n . By Lemma 8.1 and $t'_x = -1/V = O(t_x^4)$,

$$G_0(n+1) - G_0(n) = t_n + O(t_n^3),$$

whereas the difference of the two error terms is $O(t_n^2)$. Consequently

$$\log a_{n+1} - \log a_n = t_n + O(t_n^2) > 0$$

for large n . This establishes eventual increase without assuming it in advance.

Lemma 8.1 shows G_M is eventually strictly increasing, and its main term is $cx^{2/3}$, so it tends to infinity. Thus (8.3) has a unique solution in that tail for large y . In any bounded neighborhood of $x = x_M(y)$, the quantities t_u are uniformly comparable to t_x , and $G'_M(u) \sim t_x$.

Let $e = C_M t_x^{2M+1}$ with C_M larger than a suitable multiple of the uniform constant in (8.5). The mean value theorem then shows that an integer at or above $x + e$ within distance one has $\log a_n > \log y$, while an integer below $x - e$ within distance one has $\log a_n < \log y$. The strict inequalities follow by increasing C_M once; they also cover exact integer coincidence of a bracket endpoint. Choose the upper integer $\lceil x + e \rceil$ and the lower excluded integer $\lceil x - e \rceil - 1$. Eventual monotonicity extends the conclusions through the entire increasing tail. For large y , the finite exceptional prefix has all its coefficients below y . This proves the global threshold bracket (8.4).

Finally (8.5) and the leading logarithmic equivalent imply $G_M(x) \sim cx^{2/3}$; inversion gives the stated leading scale. $\square$

Remark 8.3 (What the rounding statement does and does not say). The constants C_M, Y_M are existence constants here, not numerically certified global cutoffs. The bracket shrinks, and its two ceilings agree except near integers. It is not an unconditional exact-ceiling identity $N(y) = \lceil x_M(y) \rceil$. Numerical solutions of (8.3) test the model inverse; by themselves they do not certify the asymptotic remainder constant or its range of validity.

[write] Added 2 October 2026, batch 77: relation to the repository's inversion apparatus.

The inverse of this section is taken of the exact-saddle model G_M , which keeps the arithmetic fluctuations of f ; it is not a linear-logarithmic phase, so the Lambert-core theorems `p0:thm:lambert-core` and `p0:thm:lambert-centered` of ProveIt's transseries volume [14] do not apply to it as stated. They apply to the refuted pure models: with $X = x^{2/3}$, $\log m_x^\pm = c_\pm X + b_\pm \log X + \text{const}$ with $b_+ = -1$, $b_- = -5/4$, which is `p0:thm:lambert-core` with $a = c_\pm$, $b = b_\pm$ and the branch W_{-1} . That inverse cannot locate $N(y)$ to within a bounded distance: the slope of $\log m_x$ tends to zero, while by Theorem 3.3 the logarithmic error does not. What is an instance of repository results is the integer step: the width ϵ_M (local error divided by a slope comparable to t_x) is the residual-to-root conversion `p0:thm:backward-error`, and the bracket (8.4), with the remark that its ceilings agree except near integers, is `p0:thm:staircase`, parts (1)–(2), with $x_* = x_M(y)$ and $\eta = \epsilon_M(y)$; the manuscript proves it directly from the mean value theorem and eventual monotonicity, without an interpolation. The same holds for Theorem 9.2 with L_M and $\hat{x}_M$. No novelty is claimed for that step; what is specific to these sequences is the model G_M (or L_M) and its error $O_M(t^{2M+2})$.

9 All fixed orders without an inner saddle solve

The explicit point t_0 can also support a quantified expansion if its mean mismatch is retained. This gives an alternative to solving (5.1); it is not a constant-coefficient expansion in powers of $n^{-1/3}$. For real $x > 0$, set

$$t = t_0(x) = (2A/x)^{1/3}, \quad \delta(x) = \frac{x - \kappa_1(t)}{\sqrt{V(t)}}, \quad \lambda_r(t) = \frac{\kappa_r(t)}{V(t)^{r/2}} \quad (r \geq 3). \quad (9.1)$$

Lemma 4.1 gives $\delta(x) = o(1)$ unconditionally, while $\lambda_r(t) = O(t^{r-2})$. In particular $|\delta(x)| \leq 1$ eventually.

Let H_d denote the probabilists' Hermite polynomial,

$$H_d(v) = (-1)^d e^{v^2/2} \frac{d^d}{dv^d} e^{-v^2/2}, \quad H_0 = 1, \quad H_1 = v, \quad H_{d+1} = vH_d - dH_{d-1}.$$

Set $Q_0 = 1$ and, for $j \geq 1$, define the finite correction grade

$$Q_j(t, \delta) = \sum_{\substack{m_3, \dots, m_{j+2} \geq 0 \\ \sum_{r=3}^{j+2} (r-2)m_r = j}} H_D(\delta) \prod_{r=3}^{j+2} \frac{1}{m_r!} \left(\frac{\lambda_r(t)}{r!} \right)^{m_r}, \quad D = \sum_{r=3}^{j+2} r m_r. \quad (9.2)$$

Uniformly for $|\delta| \leq 1$, $Q_j = O_j(t^j)$. The first four grades, with every Hermite argument equal to δ , are

$$\begin{aligned} Q_1 &= \lambda_3 H_3 / 6, \\ Q_2 &= \lambda_4 H_4 / 24 + \lambda_3^2 H_6 / 72, \\ Q_3 &= \lambda_5 H_5 / 120 + \lambda_3 \lambda_4 H_7 / 144 + \lambda_3^3 H_9 / 1296, \\ Q_4 &= \lambda_6 H_6 / 720 + (\lambda_3 \lambda_5 / 720 + \lambda_4^2 / 1152) H_8 \\ &\quad + \lambda_3^2 \lambda_4 H_{10} / 1728 + \lambda_3^4 H_{12} / 31104. \end{aligned} \quad (9.3)$$

At zero mismatch, $Q_{2j}(t, 0) = C_j(t)$ and the odd grades vanish. At t_0 the mismatch is generally nonzero, and odd grades must be retained: for example $Q_1 = o(t)$ need not be $O(t^2)$.

Theorem 9.1 (Explicit-point expansion). *For fixed $M \geq 0$ put*

$$U_M(t, \delta) = \sum_{j=0}^{2M+1} Q_j(t, \delta).$$

For either sequence, with $t = t_0(n)$ and $\delta = \delta(n)$,

$$a_n = \frac{e^{f(t)+nt-\delta^2/2}}{\sqrt{2\pi V(t)}} \left\{ U_M(t, \delta) + O_M(t^{2M+2}) \right\}. \quad (9.4)$$

More generally the formula holds uniformly for small positive t and integer n with $|n - \kappa_1(t)| \leq \sqrt{V(t)}$, using the corresponding mismatch. The sum $U_M = 1 + O_M(t)$ is eventually positive.

Proof. At a general t , Fourier inversion differs from (5.4) by the factor $e^{-i(n-\kappa_1(t))\theta}$. With $z = \theta\sqrt{V}$, this is $e^{-i\delta z}$, of modulus one. The uniform derivative and angular bounds from Section 5 are therefore unchanged.

Put $K = 2M + 2$. Use exactly the auxiliary-variable Taylor expansion in the proof of Theorem 5.2, retaining every weighted degree through $K - 1$. Its remainder is bounded, after multiplication by the Gaussian, by

$$C_M t^K P_M(|z|) e^{-z^2/4}.$$

The extra factor has modulus one, so this integrated absolute error remains $O_M(t^K)$ uniformly in real δ . The tails and extension to the real line remain exponentially small. Unlike at the centered saddle, odd terms no longer integrate to zero. Instead the Gaussian Fourier identity is

$$\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-z^2/2 - i\delta z} (iz)^D dz = H_D(\delta) e^{-\delta^2/2}. \quad (9.5)$$

Indeed multiplication by $(iz)^D$ corresponds to $(-d/d\delta)^D$; the sign from the Gaussian derivative cancels this sign. Thus there is no additional $(-1)^D$ in (9.2). Integrating each retained monomial gives Q_j . Dividing the absolute scaled-integral remainder by $e^{-\delta^2/2} \geq e^{-1/2}$ preserves its order on $|\delta| \leq 1$, proving (9.4). The fact that $\delta(n) = o(1)$ verifies this condition for the explicit point. No rate for that convergence is needed. $\square$

9.1 The explicit-point model inverse

For large real x , with $t = t_0(x)$, define

$$L_M(x) = f(t) + xt - \frac{1}{2} \log(2\pi V(t)) - \frac{1}{2} \delta(x)^2 + \log U_M(t, \delta(x)). \quad (9.6)$$

It involves an explicit evaluation point and convergent cumulant sums, without an inner saddle equation.

Theorem 9.2. *For each fixed M , $L'_M(x) = t_0(x) + o(t_0(x)^2) > 0$ eventually. For sufficiently large y let $\hat{x}_M(y)$ be the unique large solution of $L_M(x) = \log y$. There are positive constants $\hat{C}_M, \hat{Y}_M$ such that, for $y \geq \hat{Y}_M$,*

$$\left[\hat{x}_M(y) - \hat{C}_M t_0(\hat{x}_M(y))^{2M+1} \right] \leq N(y) \leq \left[\hat{x}_M(y) + \hat{C}_M t_0(\hat{x}_M(y))^{2M+1} \right]. \quad (9.7)$$

This has the same inverse-location scale as (8.4) and the same qualification about rounding near integers.

Proof. Write $E_r(t) = \kappa_r(t) - A(r+1)!t^{-r-2}$, so $E_r = o(t^{-r-1})$ for each fixed r . We do not differentiate a bare little- o estimate. Instead let $W(t) = 2At^{-3} - \kappa_1(t)$ and use exact identities:

$$W'(t) = V(t) - 6At^{-4} = E_2(t), \quad V' = -\kappa_3.$$

Consequently

$$\frac{d\delta}{dt} = \frac{E_2(t)}{\sqrt{V}} + \frac{\delta\kappa_3}{2V} = o(t^{-1}), \quad t'_0(x) = -\frac{t_0^4}{6A}, \quad \delta'(x) = o(t_0^3). \quad (9.8)$$

Similarly exact differentiation gives

$$\frac{d\lambda_r}{dt} = -\kappa_{r+1}V^{-r/2} + \frac{r}{2}\kappa_r\kappa_3V^{-r/2-1} = O_r(t^{r-3}).$$

Thus $\partial_t Q_j = O_j(t^{j-1})$ and $\partial_\delta Q_j = O_j(t^j)$ for bounded mismatch. Along x , each derivative is $O_j(t_0^{j+3})$, and $d \log U_M / dx = O_M(t_0^4)$. Direct differentiation of (9.6) now yields

$$\begin{aligned} L'_M(x) &= t_0 + (x - \kappa_1(t_0))t'_0 + \frac{\kappa_3(t_0)}{2V(t_0)}t'_0 - \delta\delta' + \frac{d}{dx} \log U_M \\ &= t_0 + o(t_0^2) + O(t_0^3) + o(t_0^3) + O_M(t_0^4). \end{aligned}$$

Here the second term uses $(x - \kappa_1)t'_0 = \delta\sqrt{V}O(t_0^4) = o(t_0^2)$. This proves the derivative assertion. The leading term of $L_M(x)$ is $cx^{2/3}$, so the model increases to infinity and has the stated eventual inverse.

Theorem 9.1 gives $\log a_n - L_M(n) = O_M(t_0(n)^{2M+2})$. Dividing the local error by a derivative comparable to t_0 gives width $O(t_0^{2M+1})$. Eventual coefficient increase is already established in Theorem 8.2. Apply precisely its local comparison to the two integers immediately outside this real interval, and use monotonicity for more distant integers and a sufficiently large y for the finite prefix. This proves (9.7), including the endpoint convention. $\square$

Arithmetic fluctuations remain in f , the exact cumulants and δ . Deleting the odd grades on the sole ground that $\delta \rightarrow 0$ would generally invalidate the asserted remainder.

10 Computation and reproducibility

The companion archive contains original executable checks and recorded data, the article source, a PDF rebuild script, pinned dependency information and checksum manifests. It does not contain full third-party papers or private working reports. The calculations are ancillary checks; the arguments above are analytical proofs.

10.1 Exact coefficient recurrence and independent expansion

Let

$$h_k^- = \sum_{m|k} mb_m, \quad h_k^+ = \sum_{m|k} mb_m(-1)^{k/m+1}.$$

Logarithmic differentiation of the products yields the exact recurrence

$$a_0^\pm = 1, \quad na_n^\pm = \sum_{k=1}^n h_k^\pm a_{n-k}^\pm. \quad (10.1)$$

Only factors with $m \leq n$ contribute to a coefficient of degree n . Thus a product truncated at length $L \geq n$ gives that coefficient exactly, not approximately. The code checks divisibility by n at every step. A separate method multiplies the finite binomial factors: $\binom{b_m}{j}$ for the plus case and $\binom{b_m+j-1}{j}$ for the minus case. It also independently computes unitary divisor sums by trial division. The displayed OEIS prefixes contain 36 terms for A301981 and 37 for A301982, including a_0 , and match these independent expansions.

10.2 Cumulants and finite product tails

To evaluate exact-product formulas numerically, a finite positive-real product is used with a rigorous deterministic tail bound. For $r \geq 0$, the absolute differentiated logarithmic-series tail beyond L is bounded by

$$\sum_{m>L} b_m m^r \sum_{\ell \geq 1} \ell^{r-1} e^{-m\ell t}. \quad (10.2)$$

For $r = 0$, use $\ell^{-1} \leq 1$. For $r \geq 1$, the elementary inequality

$$\sum_{\ell \geq 1} \ell^{r-1} q^\ell \leq (r-1)! \frac{q}{(1-q)^r} \quad (0 < q < 1)$$

follows from $\ell^{r-1} \leq (r-1)! \binom{\ell+r-2}{r-1}$ (with the evident $r = 1$ case). Since $b_m \leq m^2$, one convenient common bound, for $r \geq 0$, is

$$T_{r,L}(t) \leq \frac{d_r}{(1 - e^{-(L+1)t})^{\max(1,r)}} \sum_{m>L} m^{r+2} e^{-mt}, \quad d_0 = 1, \quad d_r = (r-1)! \ (r \geq 1). \quad (10.3)$$

The remaining sum can be evaluated by the finite polynomial recursion for geometric power sums or bounded by an incomplete gamma integral once L exceeds the function's turning point. These bounds cover both signs; signs in the plus logarithmic series are discarded only for bounding the tail.

Cumulant evaluation itself uses a rational polynomial recursion. Let $q = e^{-mt}$, and write $\varepsilon = 1$ for the minus product and $\varepsilon = -1$ for the plus product. Starting with $P_1(q) = q$, set

$$P_{r+1}(q) = q(1 - \varepsilon q)P'_r(q) + r\varepsilon q P_r(q). \quad (10.4)$$

Then the r th cumulant contribution of size m is $b_m m^r P_r(q)/(1 - \varepsilon q)^r$ for $r \geq 1$. This supplies exact symbolic polynomials followed by floating-point evaluation, without finite-differencing the product.

10.3 Interpretation of the numerical checks

The replay compares exact coefficients with the $M = 0, 1, 2$ saddle formulas, the explicit- t_0 equivalent and numerical model inverses. Separate off-center checks compute all Hermite grades, compare independent binomial products and integer recurrences through degree 500, and verify the Gaussian Fourier signs directly. They also test that deleting odd grades damages the claimed accuracy. Repeating at higher precision and larger product cutoffs tests stability; deterministic bounds such as (10.3) bound discarded infinite tails. These are distinct from interval certification of floating-point arithmetic. Unless explicitly marked as an exact integer assertion, recorded numerical checks are diagnostics, not certified enclosures.

For illustration, at $n = 1000$ for A301982 the relative error $a_n/\hat{a}_n - 1$ is approximately -2.05977×10^{-3} at $M = 0$, -2.99391×10^{-6} at $M = 1$ and -7.89389×10^{-9} at $M = 2$. These values are consistent with the proved t_n^2, t_n^4, t_n^6 orders. The products and coefficients are computed independently

of fitting any asymptotic parameters. Finite numerical agreement with the old models cannot establish a pure equivalent and cannot decide RH. No numerical zero search is needed in the replay or in the proof.

The archive’s README gives exact commands. Invoke shell scripts with `bash`, so a clean extraction need not preserve executable mode bits. A supplied safe Python extraction utility rejects absolute paths, parent traversal, duplicate names and symlinks before writing a new destination. Checksums pin the supplied sources and data, and a clean replay recomputes the diagnostics and rebuilds the PDF. TeX engine versions can change PDF bytes, so rebuild verification also checks mathematical text and a successful warning-free typeset result rather than presuming cross-engine binary identity.

[write] **Added 2 October 2026, batch 77: the shipped files.** In ProveIt the “companion archive” is this report’s directory, with a different layout: the programs are in `code/` (with the delivered `reproduce.sh` and `build.sh`), the recorded data, the requirements pin and the four delivered receipts in `data/` (the receipts as `data/receipts-*`). The two checksum manifests the text mentions (`MANIFEST.sha256`, `SOURCE-MANIFEST.sha256`) and the delivered PDF are not shipped; they were verified at intake and survive in the archive of commit `096ee7b87`. “The archive’s README” is the delivery README, which the report README replaces; the report README gives rerun commands that work with the shipped names. At intake (2 October 2026) the replay was run on a copy: twelve recomputed JSON files equalled the recorded ones; `inverse_diagnostics.json` and `offcenter_inverse_diagnostics.json` were not regenerated (each run exceeded the intake’s three-minute limit), and the PDF rebuild was not run.

11 Further questions and precise limits

Several extensions would require additional work beyond the results proved here.

1. Quantify the Fourier decay in (4.3) to give a useful explicit rate in the fixed- t_0 equivalent, or analyze Newton-corrected explicit evaluation points. The present proof only supplies $o(1)$ relative error there.
2. Make the remainder constants and starting ranges in Theorem 5.2 effective enough to certify integer thresholds for prescribed large y . The current inverse theorem gives asymptotic existence of shrinking brackets.
3. Determine one-sided or two-sided oscillation properties of $\log a_n - cn^{2/3}$ and a_n/m_n . The non- O and noncompactness conclusions here do not settle the stronger alternatives.
4. Investigate carefully formulated contour-based zero corrections with multiplicities and cancellation handled. An explicit full zero-residue transseries would require separate convergence and remainder arguments and is not supplied here.
5. Compare other arithmetic color weights with denominator zeros and possible numerator cancellation, keeping exact-family claims separate from the already established general methods.

The results are compatible with every currently undecided aspect of RH. They correct two particular multiplicative conjectures while preserving the valid leading logarithmic scale and replacing the missing arithmetic information by exact product or saddle data.

[write] **Added 2 October 2026, batch 77.** As of this write no report of the collection and no transseries volume of ProveIt answers any of these questions; in particular the repository supplies no zero-residue transseries for these products (item 4).

[write] **Added 3 October 2026, batch 86.** Question 3 is answered by Part II (unrefereed, prepared with AI assistance, not formalized). Both quantities oscillate two-sidedly: for each sign $\liminf a_n^\pm/m_n^\pm = 0$, $\limsup a_n^\pm/m_n^\pm = \infty$, and $\log a_n^\pm - c_\pm n^{2/3}$ and $\log(a_n^\pm/m_n^\pm)$ are $\Omega_\pm(n^{1/4})$, with explicit lower constants $K_\pm$ from one uncanceled pole (Theorem 12.5); the error of inverting either pure model oscillates likewise on the scale $(\log y)^{7/8}$ (Corollary 12.6). Finer forms of the question stay open and are Part II’s own questions (Section 12.6): how often each sign occurs, effective indices and certified constants, the sharp scale and

amplitude, and the joint behaviour of the two signs. Question 4 is unchanged and is restated as Part II's question 4. Questions 1, 2 and 5 are unchanged. The preceding note, that no report answers any of these questions, was true on 2 October 2026 and remains true of questions 1, 2, 4 and 5.

Part II

Two-sided fluctuations of unitary-divisor partitions

Summary. We strengthen the existing refutation of the pure asymptotic models for A301981 and A301982: each coefficient-to-model ratio has lower limit zero and upper limit infinity, with two-sided logarithmic oscillations of at least order $n^{1/4}$ and corresponding inverse errors.

From the manuscript’s introduction. This Part strengthens an existing refutation of two partition asymptotics to quantitative oscillation in both directions. The proof strategy has a common feature with the manuscript’s other advances: an exact structural identity is retained long enough to control information that a leading estimate loses. The repository snapshot used throughout is

eaf08931cbfd0132dd36ab82a7743ccb02506dc7, 4 October 2026, 00:59:12 UTC.

The related OEIS entries and the cited primary literature were inspected on 4 October 2026 UTC. The manuscript supplies written proofs, independent internal mathematical reviews, exact finite checks, and an interval certificate. These are distinct forms of evidence: a finite computation does not establish an infinite asymptotic statement. External peer review and proof-assistant formalization remain future work.

Let a_n^- and a_n^+ be the coefficients of

$$\prod_{j \geq 1} (1 - z^j)^{-\sigma^*(j)}, \quad \prod_{j \geq 1} (1 + z^j)^{\sigma^*(j)},$$

respectively. Their pure OEIS models $m_n^\pm$ are defined precisely in Section 12. The prior report (Part I) already proves that these models are not asymptotic equivalents. The new result is

$$\liminf \frac{a_n^\pm}{m_n^\pm} = 0, \quad \limsup \frac{a_n^\pm}{m_n^\pm} = \infty, \quad \log(a_n^\pm/m_n^\pm) = \Omega_\pm(n^{1/4}).$$

The notation $\Omega_\pm$ means that positive and negative multiples of the indicated scale occur along separate infinite subsequences. The proof gives sharper lower constants from one noncancelled Mellin pole. It requires neither the Riemann hypothesis nor simplicity of zeta zeros. The error of inverting either pure model likewise has both signs on a scale at least $(\log y)^{7/8}$.

The two-sided partition theorem answers its predecessor’s Section 11, question 3; it does not claim the earlier refutation as new. The general methods of analytic inversion, Lagrange inversion, and Landau positivity are also established methods. The search did not identify earlier proofs of these particular new conclusions. It does not establish exhaustive historical priority. Each Part states its dependencies explicitly so that its correctness can be assessed independently of any claim about importance or novelty.

[write] **Added 3 October 2026, batch 86: provenance of Part II.** Part II is one subject of a three-subject manuscript: manuscript 01 of batch 86 of ProveIt’s incoming-reports intake, *Three advances on OEIS conjectures and asymptotics. Bridgeless toroidal maps, golden-ratio moment laws, and unitary-divisor partitions*, dated 4 October 2026 (UTC), whose author line reads “Prepared with ChatGPT” and “Building on Vladimir Reshetnikov’s ProveIt research repository” (a 2,894-line source and a 40-page PDF). It arrived as the archive `oeis_advances.zip` (wrapper directory `oeis_advances/`, main file `oeis_advances.tex`) in commit `ae9baa422` and was split by subject in placement commit `0f084afa9`. Its Part III, the single Section 7, continues this report and is printed here in full as Section 12. The manuscript pins the repository at commit

eaf08931cbfd0132dd36ab82a7743ccb02506dc7 and cites this report’s `article.tex` at that commit, which is the text printed as Part I: the report did not change between the pin and the placement. The manuscript’s other subjects are printed elsewhere in the collection: its Part I (Section 2, the bridgeless toroidal maps of A343093) is the new report `enumerative-combinatorics/a343093-bridgeless-toroidal-maps`, and its Part II (Sections 3–6, A088714 and A088713) is Part IV of `oeis-sequence-asymptotics/a088714-bell-scale-growth`. Of its shared material this Part prints the passages on the partition subject: from its abstract and Section 1 (above), from Sections 8 and 9 (Section 13), and the bibliography entries that Section 7 cites. Its program `verify_partitions.py` and that program’s recorded output are shipped as `code/02-two-sided-verify_partitions.py` and `data/02-two-sided-partition_verification.json`. The manuscript’s L^AT_EX source and its README were placed as the starting `article.tex` and `README.md` of the A343093 report; they, the delivered PDF and every other delivered file survive in the archive of commit `ae9baa422`. The manuscript is unrefereed and was prepared with AI assistance. The internal reviews it records (Section 13) are not refereeing. Nothing in it is formalized in Lean or Rocq, and its place in this report confers no formal status.

What it answers. Part II answers question 3 of Section 11: both $\liminf a_n^\pm/m_n^\pm = 0$ and $\limsup a_n^\pm/m_n^\pm = \infty$ hold, and both centered logarithms are $\Omega_\pm(n^{1/4})$ (Theorem 12.5). It thereby settles the alternative that the remark after Theorem 3.3 leaves open, and makes the one-sided obstruction of Theorem 7.2 two-sided. Dated notes at those three places of Part I say so.

Overlap with Part I, printed and credited. Proposition 12.1 restates Lemma 4.1 (its (12.5) is (4.1)) and Theorem 6.1 (its (12.6) is (1.9) in logarithmic form), as the manuscript says, and its proof re-derives both in outline. Within that proof (12.7) is (2.4)–(2.5), (12.8) is Lemma 5.1, and (12.9) is the case $M = 0$ of Theorem 5.2. The definitions (12.1), (12.2) and (12.4) are (1.4), (1.5)–(1.6) and (1.7)–(1.8). The zero selected in (12.12) is the zero chosen in the proof of Lemma 3.1, and s_* is that lemma’s s_0 : Lemma 12.4 re-proves Lemma 3.1 and adds what is new, the order m of the pole and its leading Laurent coefficient $C_\pm$. Corollary 12.6 uses the eventual increase of Theorem 8.2. New to the repository are Lemma 12.3 (a written-out form of Landau’s classical positivity method, for which no novelty is claimed), Theorem 12.5, Corollary 12.6 and the constant (12.13).

Editorial changes. The manuscript’s Section 7 is Section 12 here; its theorem and equation numbers 7. k are printed as 12. k . Every label carries the prefix `udp:tw:` in place of the manuscript’s `part:`, and the closing subsection received the label `udp:tw:sec:scope`. Its citations of the pinned report (its key `unitaryprior`, whose pin and path this note records) now point to Part I, and its cited section, theorem and lemma numbers (Section 11, Theorem 3.3, Lemma 4.1, Theorem 6.1, Theorem 8.2) are cross-references, which print the same numbers. Its OEIS, Tóth, Trudgian and DLMF citations use Part I’s bibliography entries (the manuscript inspected the OEIS entries again on 4 October 2026 UTC); Landau 1905 and Diamond–Pintz 2009 are added. In the passages taken from the abstract and Section 1 above, “Finally, we strengthen” became “We strengthen”, “The third strengthens” became “This Part strengthens”, “The subjects differ, but the proof strategy has a common feature” became “The proof strategy has a common feature with the manuscript’s other advances”, “Part III” became the section reference, the citation of the prior report became Part I, a clause about older map-enumeration transformations was omitted, and the display (1.3) is unnumbered. The interval certificate named there belongs to the A088714 subject (Part IV of `a088714-bell-scale-growth`); this Part uses none. The `[write]` notes are new. No statement, proof or number of the manuscript’s Section 7 was altered.

[write] Added 3 October 2026, batch 86: notation of Part II. Part II keeps the manuscript’s symbols. Most agree with Part I’s: $A_\pm$, $c_\pm$, $d_\pm$, $B_\pm$, $R_\pm$, $f_\pm$, $m_n^\pm$, $M_\pm$, κ_r , V , t_n , ϕ_t , $K_r(s)$, the angular constants c_1, c_2, c_3 , and $\rho = \beta + i\gamma$ mean the same in both Parts; $t_{0,\pm}(n)$ is Part I’s $t_0^\pm(n)$, $\chi_\pm$ its $\chi_\pm$, $\mathcal{G}$ its $\mathcal{G}$, $N_\pm(y)$ its $N(y)$, and $\Re$ its $\Re$. The following do not, and each is printed as delivered; the right column gives the tempting false reading.

symbol	meaning in Part II	not to be confused with
j, b_j	part size, $b_j = \sigma^*(j)$	Part I writes m, b_m
m	order of the pole at s_* (multiplicity of ρ)	a part size; the multiplicities m_r of (5.6)
$W(n)$	$n^{\sigma^*/3}(\log n)^{m-1}$, the oscillation scale of Theorem 12.5	<i>not</i> the Lambert W function of the neighbouring report a088714-bell-scale-growth ; not $W(t) = 2At^{-3} - \kappa_1(t)$ in the proof of Theorem 9.2
$s_* = \sigma_* + i\tau_*$	$(1 + \rho)/2$, the pole of Lemma 12.4 (Part I's s_0); a general nonreal pole in Lemma 12.3	σ_* (subscript) is not σ^* (superscript, the unitary divisor sum), $\sigma(m)$ or the contour abscissa σ
$C_*, C_\pm$	leading Laurent coefficient of H at s_* (Lemma 12.3); that of $M_\pm$ ((12.13))	the contour abscissa c_* ; $C_j(t)$, C_1, C_2, C_r , C_M
$K_\pm, L_\pm$	lower constants of Theorem 12.5 and Corollary 12.6	K (four meanings), L, L_M
$h, H; h_\pm, H_\pm$	a function on $[1, \infty)$ and its Mellin transform (Lemma 12.3); $h_\pm(x) = R_\pm(1/x)$ and (12.15)	H in (3.2); Hermite H_d
q, Q, α, δ, r	the nonnegative function, its transform, its abscissa of convergence, a shift and a disc radius in the proof of Lemma 12.2	$q = e^{-mt}$; Q_j ; the three δ 's; the derivative order r
$T_\pm(s), E_X(s)$	$\int_1^\infty f_\pm(t)t^{s-1} dt$; an entire correction	the tail bound $T_{r,L}(t)$; $E(u)$, E_1, E_r
E_n, g, ξ_n	$\log(a_n^\pm/m_n^\pm)$, $\log m^\pm(x)$, a mean-value point (proof of Corollary 12.6)	E_r ; the Glaisher constant
$x_n, X_\pm(y)$	$(n/(2A_\pm))^{1/3} = 1/t_{0,\pm}(n)$; the inverse of the <i>pure</i> model $m^\pm(x)$	$x_M(y)$, $\hat{x}_M(y)$, inverses of the exact models G_M, L_M

12 Two-sided arithmetic oscillations in A301981 and A301982

12.1 The question and the advance over the repository

Let

$$b_j = \sigma^*(j) = \sum_{\substack{d|j \\ (d,j/d)=1}} d, \quad F_-(z) = \prod_{j \geq 1} (1 - z^j)^{-b_j}, \quad F_+(z) = \prod_{j \geq 1} (1 + z^j)^{b_j},$$

and write $a_n^\pm = [z^n]F_\pm(z)$. These are A301981 and A301982, respectively; the weights form A034448 [1, 2, 3]. For the minus sign each color can be repeated; for the plus sign each individual color can be used once. All coefficients are positive.

Put $\mathcal{G}$ for the Glaisher–Kinkelin constant and define

$$A_- = \frac{\pi^2}{6}, \quad A_+ = \frac{\pi^2}{8}, \quad c_\pm = 3(A_\pm/4)^{1/3}, \quad t_{0,\pm}(n) = (2A_\pm/n)^{1/3}, \quad (12.1)$$

$$m_n^- = \frac{\mathcal{G}^6 e^{c_- n^{2/3} - 1/2}}{2^3 5^{5/6} \pi^{1/3} n^{5/6}}, \quad m_n^+ = \frac{e^{c_+ n^{2/3}}}{2^{4/3} \sqrt{3} \pi^{1/6} n^{2/3}}. \quad (12.2)$$

The OEIS entries label $a_n^\pm \sim m_n^\pm$ as conjectural. The existing ProveIt report (Part I) already refutes both equivalents: neither ratio eventually lies in a compact subinterval of $(0, \infty)$. Its Section 11, question 3, explicitly leaves the direction of divergence and two-sided oscillation open. Its remark after Theorem 3.3 stresses that it does not prove both a zero lower limit and an infinite upper limit.

The result here answers that question. For each sign,

$$\liminf_{n \rightarrow \infty} \frac{a_n^\pm}{m_n^\pm} = 0, \quad \limsup_{n \rightarrow \infty} \frac{a_n^\pm}{m_n^\pm} = +\infty. \quad (12.3)$$

More quantitatively, there are positive constants such that $\log(a_n^\pm/m_n^\pm)$ exceeds a positive multiple of $n^{1/4}$ infinitely often and is below a negative multiple of $n^{1/4}$ infinitely often. We give sharper constants depending on an uncanceled zeta zero and its multiplicity. No Riemann hypothesis, simple-zero assumption, or expansion over all zeros is used.

The method is the classical positivity argument associated with Landau [15]; a related Mellin application is in Diamond–Pintz [16]. The contribution is its rigorous application to this explicit repository question, including the coefficient sampling step and the resulting inverse error. We make no claim to a new general oscillation principle or an exhaustive historical priority determination.

12.2 The prior coefficient bridge, with its analytic justification

In this subsection fix one sign, and omit it from A, c, f, R, t_0 when convenient. Set

$$\begin{aligned} f_\pm(t) &= \log F_\pm(e^{-t}), \quad t > 0, \\ d_- &= \frac{1}{2}, \quad B_- = 6 \log \mathcal{G} - \frac{1}{2} - \frac{1}{2} \log(2\pi), \quad d_+ = 0, \quad B_+ = -\frac{1}{2} \log 2, \\ R_\pm(t) &= f_\pm(t) - A_\pm t^{-2} - d_\pm \log t - B_\pm. \end{aligned} \tag{12.4}$$

Thus $R_\pm$ is an exactly defined function, not an error symbol with an assumed sign or bound.

Proposition 12.1 (Prior coefficient bridge). *For either sign, as $t \downarrow 0$ and $n \rightarrow \infty$,*

$$(-1)^r f_\pm^{(r)}(t) = A_\pm(r+1)! t^{-r-2} + o(t^{-r-1}) \quad \text{for every fixed } r \geq 0, \tag{12.5}$$

$$\log \frac{a_n^\pm}{m_n^\pm} = R_\pm(t_{0,\pm}(n)) + o(1). \tag{12.6}$$

In particular $h_\pm(x) := R_\pm(1/x)$ satisfies $h_\pm(x) = o(x)$ and $h'_\pm(x) = o(1)$.

These are Lemma 4.1 and Theorem 6.1 of Part I. The following argument records why the bridge is available without an assumption about zeta zeros inside the critical strip.

Proof. The elementary bounds

$$j \leq b_j \leq \sigma(j), \quad \sum_{j \leq X} b_j \leq \sum_{d \leq X} d \lfloor X/d \rfloor \leq X^2$$

give, by partial summation, $\sum_j b_j j^r e^{-ju} = O_r(u^{-r-2})$ for $u > 0$. Consequently the product logarithms and their derivatives converge locally uniformly on $\Re t > 0$, and $|f^{(r)}(w)| = O_r(t^{-r-2})$ when $\Re w \geq t/2$. The classical Dirichlet series [4, Section 4.9] gives

$$\begin{aligned} D(s) &= \sum_{j \geq 1} b_j j^{-s} = \frac{\zeta(s)\zeta(s-1)}{\zeta(2s-1)}, \quad \Re s > 2, \\ M_\pm(s) &= \int_0^\infty f_\pm(t) t^{s-1} dt = \Gamma(s) \chi_\pm(s) \frac{\zeta(s+1)\zeta(s)\zeta(s-1)}{\zeta(2s-1)}, \\ \chi_-(s) &= 1, \quad \chi_+(s) = 1 - 2^{-s}. \end{aligned} \tag{12.7}$$

For completeness, at p^a the weight is $1 + p^a$, so its Euler factor is $(1 - p^{1-2s})/((1 - p^{-s})(1 - p^{1-s}))$. Termwise Mellin integration of $f_-(t) = \sum_{j,\ell \geq 1} b_j e^{-j\ell t}/\ell$ proves (12.7); $f_+(t) = f_-(t) - f_-(2t)$ proves the plus case.

The kernel for $(-1)^r f^{(r)}(t)$ is obtained by replacing $\Gamma(s)$ by $\Gamma(s+r)$ and t^{-s} by t^{-s-r} . It has residue $A(r+1)!$ at $s = 2$. At $s = 1$ the quotient $\zeta(s)/\zeta(2s-1)$ tends to 2, so that point is removable.

The line $\Re(2s - 1) = 1$ is zero-free. The bound $1/\zeta(\sigma + iT) = O(\log(2 + |T|))$ for $\sigma \geq 1$ and large $|T|$ follows, for example, from Trudgian [12]. Together with the gamma factor and the usual polynomial bounds for zeta on fixed strips, it permits shifting the Mellin contour to $\Re s = 1$. The remaining integral is

$$t^{-r-1} \frac{1}{2\pi} \int_{\mathbb{R}} K_r(1 + iu) e^{iu \log(1/t)} du, \quad K_r(1 + iu) \in L^1(\mathbb{R}).$$

The Riemann–Lebesgue lemma proves (12.5).

Here are the coefficient details needed below. Let $\kappa_r(t) = (-1)^r f^{(r)}(t)$, $V(t) = \kappa_2(t)$, and let t_n be the unique solution of $\kappa_1(t_n) = n$. Existence and uniqueness follow by realizing each factor as a geometric, or Bernoulli, probability generating function: κ_1 is the total mean, $V > 0$ is its variance, and $\kappa_1' = -V$. Equation (12.5) gives $t_n \sim t_0$ and $V(t) \sim 6At^{-4}$.

The centered characteristic function of this independent sum obeys

$$|\phi_t(\theta)| \leq \exp\{-c_1 \min(t^{-4}\theta^2, t^{-2})\}, \quad |\theta| \leq \pi, \quad (12.8)$$

with $c_1 > 0$. Indeed keep the factors with $N \leq j \leq 2N$, $N = \lfloor 1/t \rfloor$. Their individual logarithmic moduli are at most $-c_2 \sin^2(j\theta/2)$, and $b_j \geq j$. For $|\theta| \leq 1/N$, the weighted sine-square sum is $\gg N^4 \theta^2$. For $1/N \leq |\theta| \leq K/N$, its quotient by N^2 tends uniformly to the strictly positive integral $\int_1^2 v \sin^2(vN|\theta|/2) dv$, for any fixed $K > 1$. For $K/N \leq |\theta| \leq \pi$, choosing $K > 4\pi$ and using $|\sum_{j=N}^{2N} e^{ij\theta}| \leq \pi/|\theta|$ gives a lower bound $\gg N^2$. This proves (12.8).

On the local scale $z = \theta\sqrt{V}$, Taylor's theorem gives $\log \phi_t(z/\sqrt{V}) = -z^2/2 + O(t|z|^3)$ on every fixed z interval. The bound (12.8) supplies an integrable Gaussian majorant until $|z| \asymp t^{-1}$; beyond it the total integral is $O(t^{-2}e^{-c_3 t^{-2}})$. Fourier inversion and dominated convergence therefore yield

$$a_n = \frac{\exp(f(t_n) + nt_n)}{\sqrt{2\pi V(t_n)}} (1 + o(1)). \quad (12.9)$$

Writing $\kappa_1(t) = 2At^{-3} + o(t^{-2})$ and comparing the saddle equations shows $t_n - t_0 = o(t_0^2)$. The phase $f(t) + nt$ is stationary at t_n and its second derivative is $O(t_0^{-4})$ between t_n and t_0 . Its change on that interval is consequently $o(1)$. It follows from (12.9) that

$$a_n = \frac{t_0^2}{\sqrt{12\pi A}} e^{f(t_0) + nt_0} (1 + o(1)) = m_n e^{R(t_0)} (1 + o(1)).$$

The last equality is exact before the factor $1 + o(1)$, by (12.2) and (12.4). Taking logarithms proves (12.6). Finally $R(t) = o(t^{-1})$ and $R'(t) = o(t^{-2})$ imply the assertions about h by the chain rule. $\square$

12.3 A positivity lemma with the exact pole constant

The next two lemmas isolate the classical mechanism so that eventual one-sided bounds are not confused with absolute bounds.

Lemma 12.2 (Landau's positivity principle). *Let $q : [1, \infty) \rightarrow [0, \infty)$ be locally integrable and of at most polynomial growth. If the real abscissa of convergence α of $Q(s) = \int_1^\infty q(x)x^{-s-1} dx$ is finite, then Q has a singularity at the real point $s = \alpha$.*

Proof. Suppose Q were holomorphic near α . Set $x = e^u$. For $s_1 > \alpha$,

$$(-1)^j Q^{(j)}(s_1) = \int_0^\infty u^j q(e^u) e^{-s_1 u} du \geq 0.$$

Holomorphy in the right half-plane together with a disc about α allows s_1 sufficiently close to α and $\delta > s_1 - \alpha$ such that the Taylor series at s_1 converges at $s_1 - \delta$. For example, if the disc about α has radius r , take $s_1 = \alpha + r/4$ and $r/4 < \delta < r/2$. Tonelli's theorem then identifies the finite value of that Taylor series with

$$\sum_{j \geq 0} \frac{\delta^j}{j!} \int_0^\infty u^j q(e^u) e^{-s_1 u} du = \int_0^\infty q(e^u) e^{-(s_1 - \delta)u} du.$$

This contradicts the definition of α . □

Lemma 12.3 (A quantitative two-sided consequence). *Let $h : [1, \infty) \rightarrow \mathbb{R}$ be locally integrable and of at most polynomial growth. Suppose $H(s) = \int_1^\infty h(x) x^{-s-1} dx$, initially defined far to the right, continues meromorphically to $\Re s > 0$ and is holomorphic near every positive real point. Suppose $s_* = \sigma_* + i\tau_*$, where $\sigma_* > 0$ and $\tau_* \neq 0$, is a pole of order $m \geq 1$, with*

$$C_* = \lim_{s \rightarrow s_*} (s - s_*)^m H(s) \neq 0.$$

Then

$$\begin{aligned} \limsup_{x \rightarrow \infty} \frac{h(x)}{x^{\sigma_*} (\log x)^{m-1}} &\geq \frac{|C_*|}{(m-1)!}, \\ \liminf_{x \rightarrow \infty} \frac{h(x)}{x^{\sigma_*} (\log x)^{m-1}} &\leq -\frac{|C_*|}{(m-1)!}. \end{aligned} \tag{12.10}$$

Proof. If the first inequality failed, choose $0 < c < |C_*|/(m-1)!$ exceeding the alleged upper limit. Then $q(x) = cx^{\sigma_*} (\log x)^{m-1} - h(x) \geq 0$ for $x \geq X$, with $X > 1$. Set $q = 0$ on $[1, X)$. Its Mellin transform has the meromorphic continuation

$$Q(s) = \frac{c(m-1)!}{(s - \sigma_*)^m} - H(s) + E_X(s), \tag{12.11}$$

where E_X is entire, being the integral over a finite interval. This expression is holomorphic near every real $s > \sigma_*$. Lemma 12.2 shows that the abscissa of convergence of the nonnegative integral is at most σ_* ; if that abscissa is $-\infty$, the conclusion is immediate. Hence for real $\sigma > \sigma_*$, $|Q(\sigma + i\tau_*)| \leq Q(\sigma)$. The identity (12.11) holds throughout this half-plane by uniqueness of continuation. Multiply that inequality by $(\sigma - \sigma_*)^m$ and let $\sigma \downarrow \sigma_*$. The limits of its two sides are, respectively, $|C_*|$ and $c(m-1)!$: the artificial pole in (12.11) is regular at s_* , whereas H is regular at the real point σ_* . The resulting inequality contradicts the choice of c . Apply the same argument to $-h$ to obtain the lower limit. □

No rightmost-pole hypothesis appears here. If a further pole lies to the right, the assumed one-sided envelope already contradicts the holomorphy of its nonnegative integral in that larger half-plane. The proof needs only a single nonreal pole whose leading Laurent coefficient is nonzero.

12.4 Noncancellation and the coefficient theorem

Choose a nontrivial zeta zero $\rho = \beta + i\gamma$ of least positive imaginary part, choosing its reflection about the critical line so that $\beta \geq 1/2$. Such a least height exists because zeta zeros are isolated, bounded-height portions of the closed critical strip are compact away from the pole at 1, and there are no real nontrivial zeros. Let its multiplicity be m and put

$$s_* = \frac{1 + \rho}{2} = \sigma_* + i\tau_*, \quad \frac{3}{4} \leq \sigma_* < 1. \tag{12.12}$$

Only the usual location, symmetry, and existence facts about zeta zeros are needed [13].

Lemma 12.4 (The surviving pole and its constant). *For each sign, $M_{\pm}$ in (12.7) has a pole of order m at s_* , and its leading Laurent coefficient is*

$$C_{\pm} = \frac{m! \Gamma(s_*) \chi_{\pm}(s_*) \zeta(s_* + 1) \zeta(s_*) \zeta(s_* - 1)}{2^m \zeta^{(m)}(\rho)} \neq 0. \quad (12.13)$$

Proof. The denominator has leading term $2^m \zeta^{(m)}(\rho)(s - s_*)^m / m!$. Gamma has neither a zero nor a pole at s_* , and $\zeta(s_* + 1) \neq 0$ because $\Re(s_* + 1) > 1$. If $\zeta(s_*) = 0$, its positive imaginary part $\gamma/2$ would contradict the selection of γ . The point $s_* - 1$ has real part in $[-1/4, 0)$ and nonzero imaginary part, so it is neither a trivial nor a nontrivial zero of zeta. Finally $|2^{-s_*}| < 1$, so $\chi_+(s_*) \neq 0$. Division of the leading terms proves the formula. $\square$

Theorem 12.5 (Quantified two-sided oscillation). *With (12.12)–(12.13), define*

$$K_{\pm} = \frac{|C_{\pm}|}{(m-1)!} (2A_{\pm})^{-\sigma_*/3} 3^{1-m} > 0, \quad W(n) = n^{\sigma_*/3} (\log n)^{m-1}.$$

For each sign separately,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{\log(a_n^{\pm}/m_n^{\pm})}{W(n)} &\geq K_{\pm}, \\ \liminf_{n \rightarrow \infty} \frac{\log(a_n^{\pm}/m_n^{\pm})}{W(n)} &\leq -K_{\pm}. \end{aligned} \quad (12.14)$$

The same inequalities hold after replacing $\log(a_n^{\pm}/m_n^{\pm})$ by $\log a_n^{\pm} - c_{\pm} n^{2/3}$. Consequently (12.3) holds, and both centered logarithms are $\Omega_{\pm}(n^{1/4})$.

Proof. For $h_{\pm}(x) = R_{\pm}(1/x)$ and $\Re s > 2$, splitting the Mellin integral at 1 gives

$$H_{\pm}(s) = M_{\pm}(s) - \frac{A_{\pm}}{s-2} + \frac{d_{\pm}}{s^2} - \frac{B_{\pm}}{s} - T_{\pm}(s), \quad T_{\pm}(s) = \int_1^{\infty} f_{\pm}(t) t^{s-1} dt. \quad (12.15)$$

The function $T_{\pm}$ is entire because $f_{\pm}(t)$ decays exponentially at infinity. Formula (12.15) is therefore a meromorphic continuation to $\Re s > 0$ and has the same pole and Laurent coefficient as $M_{\pm}$ at s_* .

It is holomorphic near every positive real point. The pole at 2 has been removed, the point 1 is removable as already checked, and the rational terms have poles only at 0. At every other positive real s , $\zeta(2s-1)$ has no zero: its argument is in $(-1, \infty)$, an interval without real zeta zeros. The pole of that denominator at $s = 1$ is already covered. There are no other positive real singularities of the numerator. Proposition 12.1 supplies the polynomial growth of $h_{\pm}$. Thus Lemma 12.3 applies.

It remains to preserve both signs when sampling. Let $x_n = (n/(2A_{\pm}))^{1/3}$. Consecutive values differ by $O(x_n^{-2})$. For any large x , choose n with $x_n \leq x < x_{n+1}$. The derivative estimate $h'_{\pm} = o(1)$ gives $h_{\pm}(x) - h_{\pm}(x_n) = o(x_n^{-2})$. Also $x/x_n \rightarrow 1$ and $\log x / \log x_n \rightarrow 1$. The continuous upper and lower bounds therefore hold along these discrete samples. Since

$$x_n^{\sigma_*} (\log x_n)^{m-1} = (2A_{\pm})^{-\sigma_*/3} 3^{1-m} n^{\sigma_*/3} (\log n)^{m-1} (1 + o(1)),$$

(12.6) proves (12.14). The difference between the two centered logarithms is $O(\log n)$, which is negligible on the displayed scale. Finally $\sigma_*/3 \geq 1/4$ and $m \geq 1$; each of the positive and negative subsequences is unbounded in magnitude. Exponentiation proves (12.3). $\square$

The constants in (12.14) are rigorous lower bounds on oscillation amplitudes, not exact amplitudes or estimates of the first index where a sign occurs. In particular the proof does not assert that a single zero dominates the full remainder.

[write] **Added 3 October 2026, batch 86: Theorem 12.5 beside Part I.** Theorem 12.5 and Theorems 7.2 and 7.4 bound the same centered logarithm $\log a_n^\pm - c_\pm n^{2/3}$ from the two sides. Unconditionally it is $o(n^{1/3})$ (Theorem 6.1) and, by Theorem 12.5, exceeds $(K_\pm - \varepsilon)W(n)$ and falls below $-(K_\pm - \varepsilon)W(n)$ infinitely often, with $W(n) \geq n^{1/4}$; this contains Theorem 7.2. Under RH every zero has $\beta = 1/2$, so $\sigma_* = 3/4$ and $W(n) = n^{1/4}(\log n)^{m-1}$, while Theorem 7.4 gives $O_\varepsilon(n^{1/4+\varepsilon})$; under RH the exponent $1/4$ is therefore exact. These combinations are immediate from the stated theorems; the manuscript itself uses no numerical information about zeta zeros, and neither does this note.

12.5 A consequence for inversion

Extend either expression $m_n^\pm$ in (12.2) to a positive function $m^\pm(x)$ for real $x > 0$, and let $X_\pm(y)$ be its inverse on its eventually increasing tail. Let

$$N_\pm(y) = \min\{n \geq 0 : a_n^\pm \geq y\}.$$

The earlier report proves that $a_n^\pm$ is eventually strictly increasing (Part I, Theorem 8.2). This is the only additional prior input in the following corollary.

Corollary 12.6 (Oscillating error of the pure inverse). *Put*

$$L_\pm = \frac{2^{1-m}|C_\pm|}{(m-1)!(3A_\pm)^{(\sigma_*+1)/2}} > 0.$$

Then

$$\begin{aligned} \limsup_{y \rightarrow \infty} \frac{N_\pm(y) - X_\pm(y)}{(\log y)^{(\sigma_*+1)/2} (\log \log y)^{m-1}} &\geq L_\pm, \\ \liminf_{y \rightarrow \infty} \frac{N_\pm(y) - X_\pm(y)}{(\log y)^{(\sigma_*+1)/2} (\log \log y)^{m-1}} &\leq -L_\pm. \end{aligned} \tag{12.16}$$

In particular this inverse error is $\Omega_\pm((\log y)^{7/8})$.

Proof. It suffices to use $y_n = a_n^\pm$, for which $N_\pm(y_n) = n$ once the increasing tail exceeds the finite initial prefix. Write $g(x) = \log m^\pm(x)$ and $E_n = \log(a_n^\pm/m_n^\pm)$. The mean value theorem gives

$$X_\pm(y_n) - n = \frac{E_n}{g'(\xi_n)}.$$

Here $\log a_n^\pm = c_\pm n^{2/3} + o(n^{1/3})$ by Proposition 12.1, so $X_\pm(y_n)/n \rightarrow 1$, $\xi_n/n \rightarrow 1$, and $g'(\xi_n) = (2A_\pm)^{1/3} n^{-1/3} (1 + o(1))$. Apply Theorem 12.5, with the sign reversed. Since $n \sim (\log y_n/c_\pm)^{3/2}$, conversion of the scale yields the stated constant:

$$K_\pm (2A_\pm)^{-1/3} c_\pm^{-(\sigma_*+1)/2} (3/2)^{m-1} = L_\pm.$$

□

Thus a leading logarithmic equivalent can support a correct relative inverse scale while its pure-model inverse has large errors of both signs. The exact-product inverse already constructed in Part I retains the arithmetic remainder and is unaffected by this obstruction.

12.6 Scope and further questions

The proven advance is the resolution of the repository’s two-sided oscillation question, with the stronger $n^{1/4}$ lower scale and an inverse consequence. The existing refutation, Mellin kernel, least-height noncancellation argument, and coefficient bridge are credited prior inputs; the bridge and noncancellation have been rederived above to expose every analytic hypothesis. The accompanying `code/verify_partitions.py` compares direct unitary-divisor enumeration with prime-power factorization, then compares binomial product expansion with the logarithmic-derivative recurrence. All comparisons pass through $n = 160$, including the first 21 displayed OEIS terms of each sequence. The recorded exact output is `data/partition_verification.json`. These computations check definitions and finite data; they cannot certify either infinitely occurring sign.

Several substantive questions remain.

1. Determine how often each sign occurs. The proof produces arbitrarily large indices of both signs; it supplies no natural or logarithmic density and no bound on gaps between sign changes.
2. Obtain effective indices and certified positive lower bounds for $K_{\pm}$. This requires certified information on a selected zero and quantitative control of the positivity argument, beyond simply evaluating (12.13) in floating arithmetic.
3. Determine the sharp oscillation scale and amplitude. The bound uses one surviving pole, which need not be rightmost. Even under RH, a limiting distribution would require control of the combined zero contributions and possible relations between their imaginary parts.
4. Construct a rigorously truncated zero-residue formula with an explicit remainder. Pole locations alone do not justify summing all residues, exchanging limits, or claiming a convergent oscillatory expansion.
5. Study joint behavior of the two signs. Their exact relation $f_{+}(t) = f_{-}(t) - f_{-}(2t)$ links the remainders at different scales, but the separate $\liminf$ / $\limsup$ assertions above imply no simultaneous sign pattern.

13 Verification, reproducibility, and further directions

The manuscript’s accompanying archive contains the complete L^AT_EX source, the PDF, four verification programs, their exact or interval output, and a short reproduction guide. The programs do not access the network. The stored OEIS prefixes and the repository pin make their finite reference data explicit. Standard-library integer and rational arithmetic suffice for the partition checks. The partition program, `verify_partitions.py`, computes unitary-divisor weights by divisor enumeration and by prime-power factorization; compares direct product expansion with the logarithmic-derivative recurrence through $n = 160$, including all 21 stored initial terms of each sequence.

Separate mathematical reviews checked the noncancellation, positivity argument, saddle bridge, and inverse scaling in this Part. These reviews are internal checks of a new manuscript. They are not external refereeing or machine-checked formal proofs.

There is deliberately no plot presented as evidence of the partition sign theorem. The argument is an infinite-subsequence result. A short coefficient prefix need not exhibit its full oscillation scale, and an uncertified finite sum over zeta zeros would add no proof of the claimed lower and upper limits.

[write] **Added 3 October 2026, batch 86: the shipped files of Part II.** In ProveIt the program is `code/02-two-sided-verify_partitions.py` and its recorded output `data/02-two-sided-partition_verification.json`, both byte-identical to the delivery; the manuscript’s other three programs and their outputs belong to the other subjects and are shipped with the A343093 report and with Part IV of the

A088714 report. The program prints its report and writes a file only when given `-output`; the README gives a rerun on a copy. At placement (3 October 2026) a run on a copy reproduced the recorded output up to line endings, and at this write a second run on a copy did the same. Its 21 stored terms of each sequence agree with the first 21 of the 36 and 37 terms of Part I's `data/oeis-prefixes.json`, and an independent recomputation at this write by the recurrence (10.1) gives the same $a_{160}^{\pm}$. These are checks of definitions and finite data only. The delivery has no checksum manifest.

The finite ratios recorded in Part I (`data/unitary_diagnostics.json`) do not show the oscillation: for $20 \leq n \leq 1000$, a_n^-/m_n^- rises from about 0.690 to 0.838 and a_n^+/m_n^+ falls from about 1.100 to 1.067. This does not contradict Theorem 12.5, which is a statement about infinite subsequences and gives no index at which either sign must appear (question 2 of Section 12.6); it is the situation that the manuscript's paragraph on the absence of a plot describes.

13.1 Make the partition oscillations effective

The two-sided theorem provides explicit symbolic lower constants from a genuine Mellin pole, but no first index at which either bound must occur. A useful next result would combine a certified zero enclosure, a residue enclosure, and a quantitative version of the positivity argument to produce a finite search interval containing an index of each sign.

A different question concerns statistics rather than existence: do the normalized logarithmic errors have limiting distributions, and what are the densities or gaps of sign changes? Any answer must control the combined zero contributions and the remainder of a truncated explicit formula. The exact identity $f_+(t) = f_-(t) - f_-(2t)$ also makes joint sign patterns a distinct problem. Separate oscillation of the two sequences gives no theorem about their simultaneous behavior.

13.2 Formalization and an independent submission path

The density and Mellin arguments require a broader analytic foundation, making them later targets for formalization. For external mathematical review, the three Parts of the manuscript can be separated into articles with their dependencies preserved. Updates for the partition entries should distinguish the earlier refutation from the new two-sided strengthening. No update or submission has been made as part of this work. Independent refereeing remains appropriate before asserting historical priority.

[write] **Added 3 October 2026, batch 86.** Section 13 prints the partition passages of the manuscript's Sections 8 and 9: the opening of Section 8 (its sentence on the arithmetic of the checks restricted to the partition program), the partition row of its table of checks as a sentence, the partition item of its list of internal reviews ("in Part III" became "in this Part"), its paragraph on the absence of a plot, its Section 9.5 in full, and the sentences of its Section 9.7 that concern the partition subject, lightly recast where they listed several subjects ("targets" became "targets for formalization", and the clause about the bibliographic comparison with the map literature was omitted). Its Section 9's other directions concern the other subjects. The questions of Section 12.6 and the two directions above are open; the manuscript supplies no zero-residue formula, so question 4 of Section 11 also stays open. The delivery says that nothing was submitted to the OEIS, and nothing here implies a submission.

References

- [1] The OEIS Foundation Inc., *A301981, Euler transform of A034448*, entry by V. Kotěšovec, 30 March 2018, inspected 2 October 2026. <https://oeis.org/A301981>.
- [2] The OEIS Foundation Inc., *OEIS entry A301982*, entry by V. Kotěšovec, 30 March 2018, inspected 2 October 2026. <https://oeis.org/A301982>.

- [3] The OEIS Foundation Inc., *A034448, sum of unitary divisors*. <https://oeis.org/A034448>.
- [4] L. Tóth, Alternating sums concerning multiplicative arithmetic functions, *Journal of Integer Sequences* **20** (2017), Article 17.2.1. Ordinary Dirichlet series: Section 4.9, p. 28, unnumbered. <https://cs.uwaterloo.ca/journals/JIS/VOL20/Toth/toth25.pdf>.
- [5] J. Bureaux and N. Enriquez, On the number of lattice convex chains, *Discrete Analysis* 2016:19, 15 pp. <https://doi.org/10.19086/da.1013>; <https://arxiv.org/abs/1603.09587>.
- [6] O. Bodini, P. Duchon, A. Jacquot and L. Mutafchiev, Asymptotic analysis and random sampling of digitally convex polyominoes, in *Discrete Geometry for Computer Imagery*, LNCS 7749, Springer, 2013, pp. 95–106. Extended preprint: <https://arxiv.org/abs/1306.2108>. See also the correction in [5, Section 4.2].
- [7] A. Dong, N. Robles, A. Zaharescu and D. Zeindler, *Exponential sums twisted by general arithmetic functions*, arXiv:2412.20101 (2024), version 1. <https://arxiv.org/abs/2412.20101>.
- [8] W. Bridges, B. Brindle, K. Bringmann and J. Franke, Asymptotic expansions for partitions generated by infinite products, *Mathematische Annalen* **390** (2024), 2593–2632. <https://doi.org/10.1007/s00208-024-02807-x>.
- [9] L. Ahmadi, R. Gómez-Aíza and M. D. Ward, A unified treatment of families of partition functions, *La Matematica* **3** (2024), 1257–1296. <https://doi.org/10.1007/s44007-024-00134-w>.
- [10] B. C. Berndt, N. Robles, A. Zaharescu and D. Zeindler, Partitions with multiplicities associated with divisor functions, *Journal of Mathematical Analysis and Applications* **533**(1) (2024), Article 127987. <https://doi.org/10.1016/j.jmaa.2023.127987>.
- [11] D. Basak, N. Robles and A. Zaharescu, Exponential sums over Möbius convolutions with applications to partitions, *Canadian Journal of Mathematics*. <https://doi.org/10.4153/S0008414X24000701>; preprint <https://arxiv.org/abs/2312.17435>.
- [12] T. S. Trudgian, Explicit bounds on the logarithmic derivative and the reciprocal of the Riemann zeta-function, *Functiones et Approximatio Commentarii Mathematici* **52**(2) (2015), 253–261. <https://doi.org/10.7169/facm/2015.52.2.5>. Exact bound and range also recorded at <https://researchportalplus.anu.edu.au/en/publications/explicit-bounds-on-the-logarithmic-derivative-and-the-reciprocal-/>.
- [13] NIST Digital Library of Mathematical Functions, Section 25.10, *Zeros of the Riemann zeta function*. <https://dlmf.nist.gov/25.10>.
- [14] [write] ProveIt, *Transseries: the polynomial-logarithmic calculus, series reversal at infinity, and inversion*, Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/transseries_and_inversion.tex; Theorems p0:thm:lambert-core, p0:thm:lambert-centered, p0:thm:backward-error, p0:thm:staircase.
- [15] E. Landau, Über einen Satz von Tschebyschef, *Mathematische Annalen* **61** (1905), 527–550. https://geodesic.mathdoc.fr/item/MAN_1905__61_158244/.
- [16] H. G. Diamond and J. Pintz, Oscillation of Mertens’ product formula, *Journal de théorie des nombres de Bordeaux* **21** (2009), no. 3, 523–533. <https://doi.org/10.5802/jtnb.687>.