

Partitions on maximal Schreier supports

Full coefficient asymptotics for OEIS A239950

Report 195 Prepared for Vladimir

4 October 2026

Abstract

Let p_n count partitions of n whose least part is the number of distinct part sizes. Equivalently, arbitrary positive multiplicities are placed on a maximal Schreier support and the objects are counted by total size. We prove

$$p_n = \frac{S^{1/4}}{2\sqrt{6\pi(1-a)}} n^{-3/4} e^{2\sqrt{S}n} \left(1 + \frac{c}{\sqrt{n}} + O(n^{-1}) \right),$$

where $a = \log(3/2)$, $S = \text{Li}_2(2/3) - \text{Li}_2(1/3) + a \log 2$, and

$$c = -\frac{26a^2 - 82a + 29}{144(a-1)^2} \sqrt{S} - \frac{3}{16\sqrt{S}} < 0.$$

The second correction is also evaluated explicitly, with relative remainder $O(n^{-3/2})$. The expansion continues to every fixed half-power order, with relative coefficients in $\mathbb{Q}(a, \sqrt{S})$. The proof includes global radial localization, a full-torus joint lattice local limit theorem, and an absolutely convergent three-variable saddle expansion with a shifted-lattice remainder estimate. We also obtain a Gaussian least-part law, eventual strict increase, a Lambert W_{-1} inverse sandwich with integer rounding, and an exact rational certificate ruling out finite eta quotients for $F(q)/q$. Classical support terminology, occupancy products, and saddle methods are explicitly credited. The accompanying code separates exact standard-library verification from optional symbolic algebra.

[write] **7 October 2026.** This is bundle Report 195 as placed in ProveIt (batch 111), with every delivered label prefixed by `mss:`. Its provenance, what the write read and checked, the collected non-claims and a table of reading conventions are in Section 1.2; the additions marked [write] are Remarks 1.3 and 10.2, the note at the end of Section 8.2, and a note on the questions of Section 12.1. No delivered statement, section or equation number has changed.

Contents

1	The sequence and the results	3
2	Exact enumeration and support weights	7
3	The continuous saddle and radial localization	9
4	A joint occupancy and size model	11
5	Full-torus Fourier control and the lattice limit	12
6	The coefficient equivalent and its normalization	14
7	A three-variable saddle to every fixed order	15
8	The first two explicit corrections	19
9	The least part in the constrained ensemble	23
10	Eventual monotonicity and a rounded inverse	23
11	A rational interior zero and the product obstruction	26
12	Reproducibility and further questions	27
A	Finite-jet instructions for higher coefficients	29
B	A normalization check by Gaussian integration	29

1 The sequence and the results

A nonempty integer partition is a finite multiset of positive integers. Its *support* is the set of part sizes that occur; its size is the sum of all parts, with multiplicity. Throughout,

$$p_n = \#\{\lambda \vdash n : \min \lambda = |\text{supp } \lambda|\}, \quad p_0 = 0, \quad F(q) = \sum_{n \geq 0} p_n q^n. \quad (1.1)$$

This is OEIS A239950 [1]. The empty partition is excluded. The distinction between support cardinality and total number of parts is essential.

Define the positive constants

$$\begin{aligned} a &= \log(3/2), & S &= \text{Li}_2(2/3) - \text{Li}_2(1/3) + a \log 2, \\ D &= \frac{1}{\sqrt{6(1-a)}}, & K &= \frac{DS^{1/4}}{2\sqrt{\pi}}, \end{aligned} \quad (1.2)$$

where $\text{Li}_2(z) = \sum_{k \geq 1} z^k/k^2$ for $|z| \leq 1$. Also put

$$B_* = -\frac{26a^2 - 82a + 29}{144(a-1)^2}, \quad c = B_*\sqrt{S} - \frac{3}{16\sqrt{S}}. \quad (1.3)$$

Here B_* is a scalar correction constant; it will be kept distinct from the occupancy integral $B(x, s)$ below. Positivity of S follows directly from its displayed definition, and $0 < a < 1$.

Theorem 1.1 (Coefficient expansion). *There are effectively computable real constants c_j , with $c_0 = 1$, $c_1 = c$, and c_2 given by (8.13), such that for every fixed integer $R \geq 0$,*

$$p_n = Kn^{-3/4}e^{2\sqrt{S}n} \left(\sum_{j=0}^R c_j n^{-j/2} + O_R(n^{-(R+1)/2}) \right). \quad (1.4)$$

In particular,

$$p_n = Kn^{-3/4}e^{2\sqrt{S}n} \left(1 + \frac{c}{\sqrt{n}} + O(n^{-1}) \right), \quad c < 0. \quad (1.5)$$

In particular, the first two relative corrections are explicit, with the three-term form and $O(n^{-3/2})$ remainder in (8.14). Every relative coefficient c_j belongs to the evaluated field $\mathbb{Q}(a, \sqrt{S})$; no algebraic independence of these constants is presumed. The quantifier “for every fixed R ” is part of the assertion. No convergence of the infinite series, uniformity as R grows, or effective numerical onset is claimed. The proof of coefficient extraction is separate from the real-radial expansion

$$F(e^{-t}) = De^{S/t} \left(1 + B_*t + O(t^2) \right). \quad (1.6)$$

A radial equivalent alone would not justify (1.4).

For the distributional result, introduce

$$V = 2a - \frac{1}{2}, \quad \kappa = \frac{3(1-a)}{V}, \quad W = 2S - a^2\kappa, \quad \kappa_{\text{eff}} = \frac{2S\kappa}{W}. \quad (1.7)$$

All four constants are positive; this will follow from positive-definite covariance matrices, rather than numerical evaluation.

Theorem 1.2 (Least part and inversion). *Let M_n be the least part of a partition drawn uniformly from the p_n objects in (1.1), and let $t = \sqrt{S/n}$. Then*

$$\sqrt{t\kappa_{\text{eff}}} \left(M_n - \frac{a}{t} \right) \implies N(0, 1). \quad (1.8)$$

The corresponding local mass estimate is uniform on every bounded interval of the standardized variable, as stated in Theorem 9.1.

The sequence p_n is eventually strictly increasing. For large Y , let

$$r(Y) = \left[-\frac{3}{4\sqrt{S}} W_{-1} \left(-\frac{4\sqrt{S}}{3} (K/Y)^{2/3} \right) \right]^2, \quad (1.9)$$

and $N(Y) = \min\{n \geq 0 : p_n \geq Y\}$. For some $C > 0$ and all sufficiently large Y ,

$$\left[r(Y) - \frac{c}{\sqrt{S}} - \frac{C}{\sqrt{r(Y)}} \right] \leq N(Y) \leq \left[r(Y) - \frac{c}{\sqrt{S}} + \frac{C}{\sqrt{r(Y)}} \right]. \quad (1.10)$$

The two ceilings are necessary in this level of conclusion. An arbitrarily small real error can still change a ceiling near an integer. The statements do not give an explicit least index of monotonicity.

1.1 Relation to classical work and the scope of the article

The Schreier family consists of finite sets $E \subset \mathbb{N}$ with $|E| \leq \min E$, together with the empty set. A nonempty member is inclusion-maximal exactly when $|E| = \min E$. The terminology *maximal Schreier set* is classical; see Chu and Vasseur [2]. Our support family is precisely this family of maximal sets. We put the geometric weight $q^j/(1-q^j)$ on each selected size j , and extract coefficients by the sum of all parts. This weighting and indexing must not be confused with enumeration by the greatest support element.

Schreier-type sets, partitions, and compositions are studied by Beanland and Chu [3]; bounded-multiplicity Schreier multisets are studied by Chu et al. [4]. Those models provide important context. The present constraint uses the number of distinct sizes and permits unbounded positive multiplicities. Counting ordered compositions by total length is also a different problem.

The occupancy product used here is standard in the study of distinct summands; Hwang and Yeh [7] state this product and study general measures of distinctness. Goh and Schmutz [5] prove a central limit theorem for the number of distinct sizes in ordinary random partitions. Hwang [6] treats central and local limit theorems and large deviations through two-dimensional saddle methods in a Meinardus setting. The full theorem of that paper was not available in the bounded source review for this article, so its exact applicability to the moving lower cutoff used here remains unresolved. We do not claim that general partition methods fail to cover some of the analysis.

Grabner, Knopfmacher, and Wagner [8] provide a general asymptotic scheme for partition statistics. Their quoted $P(q)H(q)$ transfer hypotheses should not be assumed for this sequence merely by writing $H = F/P$: its exponential radial scale differs from that of the ordinary partition product. Archibald et al. [9] give related generating functions involving part sizes and largest-part multiplicities. Conjugating a partition preserves its number of distinct sizes and transforms its least part into the multiplicity of the greatest part, yielding the conjugate interpretation in A239950.

The contribution here is a self-contained, sequence-specific derivation with explicit constants and reproducible finite checks. Neither the bounded source review nor the absence of an asymptotic

on an inspected OEIS page establishes worldwide novelty. No novelty of the Schreier family, the occupancy product, or the general saddle and local-limit methodology is asserted.

Remark 1.3 ([write] The OEIS entry, quoted, 7 October 2026). OEIS A239950, read on 7 October 2026 in the internal format, is at revision #18 (17 November 2015), author Clark Kimberling (30 March 2014), offset 0. Its name is “Number of partitions of n such that (number of distinct parts) = least part.”; a comment of Joerg Arndt (28 April 2014) gives the conjugate reading “(number of distinct parts) = multiplicity of the greatest part” for $n > 0$; its formula line is “A239948(n) + a(n) + A239951(n) = A000041(n) for $n \geq 0$.”; it carries Alois P. Heinz’s Maple program and b-file ($0 \leq n \leq 1000$) and Jean-François Alcover’s Mathematica programs. It states no asymptotic formula and no conjecture. Its 58 data terms are exactly those of the shipped `data/oeis_prefix.json`. The write fetched the b-file (1001 terms, SHA-256 72e8fed9e454...1cf55f), which the package did not retrieve, and recomputed $p_0, \dots, p_{1500}$ by a route of its own, the signed Euler expansion

$$C_m(q) = \frac{q^m(q; q)_m}{(1 - q^m)(q; q)_\infty} \sum_{k \geq m-1} (-1)^{k+m-1} \binom{k}{m-1} \frac{q^{k(k+1)/2+km}}{(q; q)_k}$$

of (2.1) and (2.4), which shares nothing with the positive recurrence (2.5): all 1001 b-file terms and all 1501 terms of the shipped `data/generated-exact_terms.txt` agree, as do brute-force counts for $n \leq 30$. The name quoted in the bibliography entry [1], “Number of partitions of n such that the number of different parts is equal to the smallest part.”, is a paraphrase, not the entry’s name at any revision the write saw; the delivered text is kept. Nothing was submitted to the OEIS.

1.2 Provenance, sources and reading conventions

[write] **Added 7 October 2026, batch 111.** *Provenance.* This report is Report 195 of the session bundle that ProveIt’s docs/incoming intake processed as batch 111: the archive `Report195_TeX_and_Reproducible_Code.zip` (687,972 bytes, SHA-256 561bd3b3a4ce...a968e68412c0122, 25 files, no wrapper directory), arrival commit 60f54ea06, placed in d451ef3d8. The text is the delivered `Report195.tex` (1084 lines, dated 4 October 2026), shipped as `article.tex`. Its title page and PDF author field read “Prepared for Vladimir”; the package carries no “prepared for private review” line, no e-mail address and no other personal data, and records no ProveIt commit. The delivered programs are shipped under `code/`, the observed OEIS prefix and the recorded outputs under `data/` (the latter as `data/generated-*`), and the delivered `DATA_SOURCES.md`, `README_REPRODUCIBILITY.md`, `code/PROVENANCE.md`, `code/README.md` and `data/README.md` as `DATA_SOURCES.md`, `README_REPRODUCIBILITY.md`, `code-PROVENANCE.md`, `code-README.md` and `data-README.md`; the report README maps every name. Not shipped, and retrievable from the arrival commit: the delivered 27-page PDF, the checksum manifest `SHA256SUMS.json` (24 entries, verified at the write), and the delivered README, replaced by the report README. It is a single-source report, so no merge choice was made. The write prefixed the 145 delivered labels with `mss:` and updated the 109 references to them (90 `\eqref`, 19 `\ref`); it labelled Section 1 and Subsections 1.1 and 12.1, and added this subsection, the notes marked [write] and Remarks 1.3 and 10.2. The added remarks are the last statements of their sections, this subsection follows the last delivered text of Section 1, and the added displays are unnumbered, so every theorem, section and equation of the source keeps its delivered number (the label tables of a build of the delivered text and of this one were compared).

The sources, as the write read them (7 October 2026).

- OEIS A239950 with its b-file, in the internal format (Remark 1.3).
- The transseries volume `Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/transseries_and_inversion.tex`: `p0:def:model`, `p0:def:core`, `p0:thm:lambert-core`, `p0:thm:lambert-centered`, `p0:thm:flattening`, `p0:def:three-inverses` and `p0:thm:staircase`, read for Remark 10.2.

- Not read by the write: the cited literature [2, 3, 4, 5, 6, 7, 8, 9]. The source’s account of it, and its delivered `DATA_SOURCES.md` (which says what the source inspected of each item, for instance only the publisher abstract of Goh and Schmutz and the abstract of Hwang), are reported, not confirmed.

What was checked at intake. The batch-111 dossier (7 October 2026) read the manuscript, found no error in it, confirmed $c_1 < 0$ and the value of c_2 against the exact terms to $n = 1500$, and reran the delivered suite on copies: its outputs reproduce, the only failures being Windows artifacts (line-ending bytes in hashed outputs, a recorded Python version string, a console code page), none mathematical. *What the write checked.* The proofs line by line, with no error found; in particular $B(a, s(a)) = a$ from (3.9), $V = 2a - \frac{1}{2}$ as B_s , $-G''(a) = \kappa = 3(1 - a)/V$, $A_0 = 2^{-1/2}$, the normalization (6.7), $K = DS^{1/4}/(2\sqrt{\pi})$, $-h_v(a, s_0)/12 = \frac{1}{8}$, the cap (2.6), and the reduction of $r(Y)$ to W_{-1} . With SymPy 1.14.0 it verified $\det H = -2SV\kappa$, $C = -H^{-1}$ in (8.2), that the quadratic form (7.7) is $\frac{1}{2}w^T H w$ on the plane (7.6), $W = L_0 - J_0^2/V$, both forms of (6.5), $VW\kappa_{\text{eff}} = 2SV\kappa$, that the five pieces (8.7) sum to (8.8), and that the radial pieces (8.9) sum to B_* . The six-dimensional contraction (8.17) itself was not redone symbolically; c_2 was tested numerically (note at the end of Section 8.2). In exact arithmetic it reproduced the sign certificate (8.10)–(8.11) and both Horner certificates (11.5)–(11.6) (left sides $0.2799\dots$ and $-0.5046\dots$; tail bounds $3.4 \cdot 10^{-25}$ and $1.0 \cdot 10^{-13}$), using its own counts. With mpmath 1.3.0 at 60 digits it evaluated the constants, the residuals and the inverse coefficients (the note at the end of Section 8.2 and Remark 10.2). Finite and floating computations are observations, not certificates.

Relation to the repository. Before batch 111 no file of the repository named A239950. No statement of this report is formalized in Lean or Rocq, and its place in the collection confers no formal status. The nearest reports are in the same directory `SetTheory/Cardinals/docs/reports/generating-functions-and-asymptotics/oeis-sequence-asymptotics/: a239964-sizes-equal-max-multiplicity` (batch 111: the number of distinct sizes equal to the *maximum* multiplicity, a different statistic on the same occupancy product) and `a098131-minimum-length-compositions` (ordered compositions whose parts are at least their number, the Schreier-type condition on length, which the source’s `DATA_SOURCES.md` distinguishes from this support-cardinality model). Neither shares a result with this report. The inverses of Section 10 are compared with the transseries volume in Remark 10.2.

[write] **What is not claimed, collected from the source.** No worldwide novelty: neither the bounded source review nor the absence of an asymptotic on the OEIS page establishes it, and no novelty of the Schreier family, the occupancy product, or the saddle and local-limit methodology is asserted; the applicability of Hwang’s theorems [6] to the moving cutoff is unresolved, and the source does not claim that general partition methods fail to cover some of the analysis. The expansion (1.4) holds for every fixed order only: no convergence, no uniformity as R grows, no effective numerical onset, no explicit least index of monotonicity, and no algebraic independence of a and S . The radial equivalent alone would not justify the coefficient expansion. Theorem 9.1 gives no higher-order centering and no relative local law beyond bounded windows; the full-target uniformity of Theorem 5.3 is absolute, not relative. The inverse brackets are smooth-envelope statements, not an identification with piecewise-linear interpolation; no convergent inverse series and no unconditional rounded equality are asserted. Proposition 11.1 excludes finite eta quotients (and any representation holomorphic and nonvanishing in the disk), not sums of products, numerators with zeros or q -hypergeometric identities. The optional symbolic programs compute the first two corrections only; they do not prove localization, tails or transfer, and finite checks do not prove the asymptotics. Determinism is a same-stack statement. The OEIS b-file was not retrieved by the source. The write adds: its own checks are floating or finite computations, except the exact certificates it reproduced; Remark 10.2 claims no novelty for any inversion.

Reading conventions (letters the source reuses, with the tempting false readings; no symbol of the source was renamed; symbols of the transseries volume carry the subscript “vol” in Remark 10.2).

Symbol	Meaning here (where)	Not to be read as
a, b	$\log(3/2)$; the slope $3/(2V)$ of the linear radius (7.2); in (11.4), b is the integer with $R = (b - 1)/b$	each other; the volume's $a_{\text{vol}}, b_{\text{vol}}$
c, c_j, c_0, c_1, c_2	$c = c_1$ the first relative correction; the coefficients of (1.4); but c_0 of Lemma 5.1, $c_1 < c_2$ in the proof of Lemma 5.2 and c_1, c_2 in (7.9) are unrelated positive constants	each other
C	the contraction matrix (8.2); elsewhere generic constants	a covariance of real variables
D, D_c	the constant $1/\sqrt{6(1-a)}$; the speed coordinate $X + aZ$ (Section 8.2); a fixed displacement in the proof of Theorem 10.1	each other
$p_n, p_{n,m}, p(v, s), p_0, p(n)$	the counts; with least part m ; the occupancy density; its value at $v = x$; the partition numbers	each other
$P, P_j, P_k, P(R)$	the phase (7.4); the expansion polynomials (7.16); the homogeneous phase terms (Section 8); $\prod(1 - R^j)^{-1}$	each other
B, B_*, B_{**}, B_{2r}	the occupancy integral; the radial correction; the second-order constant (8.12); Bernoulli numbers	each other
$V, V(x, s)$	$2a - \frac{1}{2}$; the variance integral	each other
W, W_{-1}	$2S - a^2\kappa$; Lambert's lower branch	each other
K, K_m	the leading constant; the occupancy count	each other
$N, N(Y)$	the total size variable; the threshold index; a truncation order in (7.16) and (11.4)	each other
Q, Q_ℓ	e^{-x} in (3.2); the quadratic form (7.7); the polynomial of (8.11); the geometric sum of Lemma 5.1	each other
T, T_N, T_J	the integral (4.7); the Horner truncation; the inverse model (10.7)	each other
J, J_0, L, L_0	the integrals (4.7) and their saddle values; J is also a compact interval (Theorem 5.3) and an inverse order	each other; the volume's L_{vol}
R, R_j	a fixed order (also a window bound and a radius); the multiplicities (4.2)	each other
ℓ_j, ℓ_1, ℓ_2	the logarithmic coefficients (10.6) in Section 10; the log-amplitude terms (8.1) in Section 8	each other
$E, \mathbb{E}, E_j, \mathbb{E}, \mathcal{E}_H$	a block of integers; expectation; the pieces (8.7); the Euler–Maclaurin series; the Wick contraction	each other
$S, \mathcal{G}, G$	the exponent constant; $F(q)/q$; the radial phase $\Psi(x, s(x))$	each other

2 Exact enumeration and support weights

For $m \geq 1$, let $p_{n,m}$ count admissible partitions with least part m , and let $C_m(q) = \sum_n p_{n,m} q^n$. The size m must occur positively; precisely $m - 1$ larger sizes must occur. Thus

$$C_m(q) = [u^m] \frac{uq^m}{1 - q^m} \prod_{j>m} \left(1 + \frac{uq^j}{1 - q^j}\right), \quad F(q) = \sum_{m \geq 1} C_m(q). \quad (2.1)$$

Each factor $q^j/(1 - q^j)$ selects a positive multiplicity for size j . Equivalently,

$$F(q) = \sum_{\substack{E \subset \mathbb{N} \text{ finite, } E \neq \emptyset \\ |E| = \min E}} \prod_{j \in E} \frac{q^j}{1 - q^j}. \quad (2.2)$$

To verify the maximality description: if $|E| < \min E$, a sufficiently large new element can be adjoined without leaving the Schreier family. If $|E| = \min E$, adjoining a larger element increases cardinality past the old minimum; adjoining a smaller element reduces the minimum as well. No proper superset is in the family.

The smallest possible total for fixed minimum m is

$$m + (m + 1) + \cdots + (2m - 1) = \frac{m(3m - 1)}{2}. \quad (2.3)$$

Therefore every coefficient in (2.1) involves finitely many m and j . These are formal power series identities. Analytically, $0 \leq p_n \leq p(n)$, where $p(n)$ is the ordinary partition number. Thus F and the relevant sums converge normally on every closed disk $|q| \leq r < 1$.

In q -Pochhammer notation $(z; q)_\infty = \prod_{j \geq 0} (1 - zq^j)$, the exact product is

$$\prod_{j > m} \left(1 + \frac{uq^j}{1 - q^j} \right) = \frac{((1 - u)q^{m+1}; q)_\infty}{(q^{m+1}; q)_\infty}. \quad (2.4)$$

This standard factorwise identity is not a simplification of the subsequent diagonal sum to a single product. Section 11 gives a precise obstruction to one tempting type of product formula.

2.1 An exact positive recurrence

The public program computes $p_0, \dots, p_N$ simultaneously. Process possible sizes j in descending order. Just before size j , let $d_{k,s}$ count partitions of s with k distinct sizes, all greater than j . The partitions that acquire positive multiplicity of j are

$$e_{k,s} = \sum_{r \geq 1} d_{k-1, s-rj} = e_{k, s-j} + d_{k-1, s-j}, \quad d_{k,s}^{\text{new}} = d_{k,s} + e_{k,s}. \quad (2.5)$$

Negative subscripts contribute zero. Update k in descending order so that the old $k - 1$ row remains available. When $k = j$, add $e_{j,s}$ to the desired answer. Each admissible partition is added once, at its minimum.

Rows above

$$k_{\max} = \left\lfloor \frac{1 + \sqrt{1 + 24N}}{6} \right\rfloor \quad (2.6)$$

are unnecessary, by (2.3). In a row k while processing j , every entry with $s < kj + k(k - 1)/2$ is zero. Integer square roots and explicit integer inequalities implement these cutoffs without floating arithmetic. The recurrence is positive and uses arbitrary-precision integers.

For orientation, the first 31 values are

n	0	1	2	3	4	5	6	7	8	9
p_n	0	1	1	1	1	2	2	3	4	4
n	10	11	12	13	14	15	16	17	18	19
p_n	6	6	9	8	14	11	19	18	25	24
n	20	21	22	23	24	25	26	27	28	29
p_n	37	31	50	46	61	64	86	79	112	115
n	30									
p_n	136									

These early decreases illustrate why eventual strict increase requires proof. The package regenerates all coefficients through 1500, checks the observed OEIS prefix through 57, and independently enumerates all ordinary partitions through 35, directly testing the defining condition. The inspected OEIS b-file was not retrieved; agreement with its entire contents is not claimed.

3 The continuous saddle and radial localization

For $t > 0$, $u = e^s > 0$, and $x = tm$, define

$$\begin{aligned} h(v, s) &= \log \left(1 + \frac{e^s}{e^v - 1} \right), & I(x, s) &= \int_x^\infty h(v, s) dv, \\ p(v, s) &= h_s(v, s) = \frac{e^s}{e^v + e^s - 1}, & B(x, s) &= \int_x^\infty p(v, s) dv, \\ V(x, s) &= \int_x^\infty p(v, s)(1 - p(v, s)) dv, & \Psi(x, s) &= I(x, s) - xs. \end{aligned} \quad (3.1)$$

Differentiation under the integral is legitimate for $x > 0$ on compact parameter sets. The derivatives have integrable exponential tails. With $Q = e^{-x}$,

$$I(x, s) = \text{Li}_2(Q) - \text{Li}_2((1 - u)Q). \quad (3.2)$$

For $u > 1$, the second dilogarithm is understood on the negative real axis by its real analytic continuation, or simply by the defining integral in (3.1).

At every $x > 0$, $B(x, s)$ is strictly increasing in s , since $B_s = V > 0$. Its limits as $s \rightarrow -\infty$ and $s \rightarrow +\infty$ are 0 and $+\infty$. For the first limit use $p(v, s) \leq u/(e^v - 1)$; for the second, p is bounded below on an interval whose length tends to infinity with $\log u$. Hence there is a unique smooth solution $s = s(x)$ of

$$B(x, s(x)) = x. \quad (3.3)$$

This is the minimizer of $\Psi(x, \cdot)$. Write

$$G(x) = \Psi(x, s(x)), \quad A(x) = \frac{u(x)e^{-x}}{1 - e^{-x}} \exp\{-h(x, s(x))/2\}. \quad (3.4)$$

Lemma 3.1 (Global strict concavity). *The function G is strictly concave on $(0, \infty)$ and has its unique global maximum at $a = \log(3/2)$. At that point $u(a) = 1/2$, $G(a) = S$, and*

$$\begin{aligned} p(a, s(a)) &= \frac{1}{2}, & h(a, s(a)) &= \log 2, & V(a, s(a)) &= V = 2a - \frac{1}{2}, \\ -G''(a) &= \kappa, & A(a) &= A_0 = 2^{-1/2}. \end{aligned} \quad (3.5)$$

Proof. Put $p_0 = p(x, s(x))$. Direct differentiation gives

$$G' = -h - s, \quad s'(x) = \frac{1 + p_0}{V(x, s(x))}, \quad G'' = \frac{p_0}{1 - e^{-x}} - \frac{(1 + p_0)^2}{V(x, s(x))}. \quad (3.6)$$

The identity

$$-p_v(v, s) = \frac{p(v, s)(1 - p(v, s))}{1 - e^{-v}} \quad (3.7)$$

implies $V(x, s(x)) \leq p_0 \leq 1$, and (3.3) implies $V(x, s(x)) \leq x$. Therefore

$$\frac{p_0 V(x, s(x))}{1 - e^{-x}} \leq \frac{p_0 \min(x, 1)}{1 - e^{-x}} \leq \frac{p_0}{1 - e^{-1}} < 4p_0 \leq (1 + p_0)^2. \quad (3.8)$$

For the middle inequality, $x/(1 - e^{-x})$ increases up to $x = 1$, and $1/(1 - e^{-x})$ decreases thereafter. The last inequality is $(1 - p_0)^2 \geq 0$. This proves $G'' < 0$.

For $u \neq 1$, integration yields

$$B(x, s) = -\frac{u}{1-u} \log(1 - (1-u)e^{-x}). \quad (3.9)$$

At $(x, u) = (a, 1/2)$ it equals a . Also $p_0 = 1/2$ and $h = \log 2 = -s$, so $G'(a) = 0$. Strict concavity proves global uniqueness. Substituting in (3.2) gives $G(a) = S$. Differentiating (3.9) in s gives $V = 2a - 1/2$; the curvature and amplitude then follow from (3.6) and (3.4). In particular $V > 0$ because it is an integral of a strictly positive function. $\square$

Lemma 3.2 (Global tails). *Fix a neighborhood J_0 of a whose closure lies in $(0, \infty)$. There is $\eta > 0$ such that, as $t \downarrow 0$,*

$$\sum_{m: tm \notin J_0} C_m(e^{-t}) = O\left(t^{-1} \log(1/t) e^{(S-\eta)/t}\right). \quad (3.10)$$

Consequently the corresponding coefficient contribution is bounded by e^{tn} times this quantity for every n .

Proof. Positivity of the product and monotonicity of h give, for every chosen $u > 0$,

$$C_m(e^{-t}) \leq \frac{ue^{-x}}{1-e^{-x}} \exp\{[I(x, \log u) - x \log u]/t\}. \quad (3.11)$$

For small x , first choose fixed $u \in (0, 1)$ so small that $I(0, \log u) < S/4$. This is possible by dominated convergence in the integral, since $h(v, \log u) \leq -\log(1 - e^{-v})$, an integrable function. Then choose $\delta > 0$ with $\delta |\log u| < S/4$. On $x \leq \delta$ the exponent is at most $S/(2t)$, and the prefactor is at most $u/(tm)$. Summation costs $O(t^{-1} \log(1/t))$.

For $x \geq M$, use $u = 1$. Here $I(x, 0) = \text{Li}_2(e^{-x}) < S/2$ when M is sufficiently large, while the prefactors $1/(e^{tm} - 1)$ sum to $O(t^{-1})$. On $[\delta, M] \setminus J_0$, choose $u = u(x)$. The prefactor is bounded, and the unique maximum of G supplies a uniform positive gap below S . There are $O(t^{-1})$ terms. Combining the three ranges proves the assertion. Finally $p_{n,m} e^{-tn} \leq C_m(e^{-t})$. $\square$

3.1 A radial local estimate

On a compact x -interval in $(0, \infty)$, Euler–Maclaurin gives uniformly

$$\sum_{j>m} h(tj, s(x)) = \frac{I(x, s(x))}{t} - \frac{1}{2} h(x, s(x)) + O(t). \quad (3.12)$$

The endpoint sign comes from starting the sum at $m+1$. Let Z_j , $j > m$, be independent Bernoulli variables with success probabilities $p(tj, s(x))$. Their sum Z is almost surely finite. Normalizing the occupancy product gives the exact identity

$$C_m(e^{-t}) = \frac{ue^{-x}}{1-e^{-x}} \exp\left\{\sum_{j>m} h(tj, s) - ms\right\} \mathbb{P}(Z = m-1). \quad (3.13)$$

Its mean and variance satisfy

$$\mathbb{E}Z = m - \frac{p_0}{2} + O(t), \quad \text{Var } Z = \frac{V(x, s(x))}{t} + O(1). \quad (3.14)$$

A direct Bernoulli Fourier bound is

$$\left| \mathbb{E} e^{i\phi Z} \right| \leq \exp \left\{ -2 \sin^2(\phi/2) \sum_{j>m} p_j(1-p_j) \right\}. \quad (3.15)$$

On the scale $\phi = \sqrt{t}\beta$, the centered logarithm converges to $-V(x, s(x))\beta^2/2$. Third moments give the uniform remainder $O(\sqrt{t}|\beta|^3)$ on bounded β ; (3.15) provides integrable Gaussian domination and suppresses the rest of the torus. Since $m-1-\mathbb{E}Z = O(1)$, Fourier inversion proves

$$C_m(e^{-t}) = A(x) \sqrt{\frac{t}{2\pi V(x, s(x))}} e^{G(x)/t} (1 + o(1)), \quad (3.16)$$

uniformly on compact intervals.

Summing near a , whose x -mesh is t and Gaussian width is $\sqrt{t}$, and using Lemma 3.2, yields

$$F(e^{-t}) \sim \frac{A_0}{\sqrt{V_K}} e^{S/t} = D e^{S/t}. \quad (3.17)$$

For rigor in the intermediate tails, the atom bound from (3.15) is $O(\sqrt{t})$ uniformly on compact x -sets; combined with $G(x) \leq S - c(x-a)^2$ it supplies a Gaussian Riemann-sum majorant. This avoids applying a local equivalent outside its stated uniform range.

4 A joint occupancy and size model

To extract coefficients, total size must be controlled together with occupancy. Define

$$H_m(t, u) = \frac{ue^{-tm}}{1-e^{-tm}} \prod_{j>m} \left(1 + \frac{ue^{-tj}}{1-e^{-tj}} \right). \quad (4.1)$$

For $j > m$, independently choose nonnegative multiplicities R_j with

$$\mathbb{P}(R_j = 0) = \frac{1}{1 + u/(e^{tj} - 1)}, \quad \mathbb{P}(R_j = r) = \frac{ue^{-tjr}}{1 + u/(e^{tj} - 1)} \quad (r \geq 1). \quad (4.2)$$

Independently choose $R_m \geq 1$ with

$$\mathbb{P}(R_m = r) = (1 - e^{-x})e^{-x(r-1)}, \quad r \geq 1. \quad (4.3)$$

Set

$$N = mR_m + \sum_{j>m} jR_j, \quad K_m = 1 + \sum_{j>m} \mathbf{1}_{\{R_j>0\}}. \quad (4.4)$$

We write K_m for the occupancy random variable to distinguish it from the leading constant K . Since $\sum_{j>m} \mathbb{P}(R_j > 0) < \infty$, only finitely many sizes are occupied almost surely.

Every configuration of total size n and occupancy k has probability $e^{-tn}u^k/H_m(t, u)$. Consequently the exact tilted identity is

$$p_{n,m} = e^{tn}u^{-m}H_m(t, u)\mathbb{P}_{t,u,m}(N = n, K_m = m). \quad (4.5)$$

The mandatory minimum factor, including its power of u , has not been dropped. From (3.12), when $u = u(x)$,

$$u^{-m}H_m(t, u) = A(x)e^{G(x)/t}(1 + O(t)) \quad (4.6)$$

uniformly on compact x -intervals.

4.1 Moments and a positive covariance matrix

In the following integrals $s = s(x)$, and derivatives in v hold s fixed. Define

$$\begin{aligned} T(x) &= I(x, s) + xh(x, s), \\ J(x) &= \int_x^\infty v \frac{p(v, s)(1 - p(v, s))}{1 - e^{-v}} dv, \quad L(x) = \int_x^\infty v^2 h_{vv}(v, s) dv. \end{aligned} \quad (4.7)$$

At an optional site with $v = tj$, logarithmic derivatives identify

$$-h_v = \mathbb{E}R_j, \quad h_{vv} = \text{Var } R_j, \quad -p_v = \text{Cov}(R_j, \mathbf{1}_{\{R_j > 0\}}). \quad (4.8)$$

Integration by parts, with the exponentially vanishing upper endpoints, gives

$$J = xp_0 + B = x(1 + p_0), \quad L = \frac{x^2 p_0}{1 - e^{-x}} + 2T. \quad (4.9)$$

Uniform Euler–Maclaurin, including the mandatory site, now yields

$$\begin{aligned} \mathbb{E}N &= t^{-2}T(x) + O(t^{-1}), & \mathbb{E}K_m &= t^{-1}x + 1 - p_0/2 + O(t), \\ \text{Var } N &= t^{-3}L(x) + O(t^{-2}), & \text{Cov}(N, K_m) &= t^{-2}J(x) + O(t^{-1}), \\ \text{Var } K_m &= t^{-1}V(x, s(x)) + O(1). \end{aligned} \quad (4.10)$$

The mandatory site contributes $O(t^{-1})$ to $\mathbb{E}N$ and $O(t^{-2})$ to $\text{Var } N$, while its occupancy is identically one.

Let

$$\mathcal{M}(x) = \begin{pmatrix} L(x) & J(x) \\ J(x) & V(x, s(x)) \end{pmatrix}. \quad (4.11)$$

This matrix is positive definite. Indeed it is the integral of the covariance matrix of $(vR, \mathbf{1}_{\{R > 0\}})$ under the optional-site law. The atoms $R = 0, 1, 2$ have positive mass and yield the noncollinear points $(0, 0), (v, 1), (2v, 1)$. Every nonzero linear form has positive variance at every $v > 0$. Continuity then gives a uniformly positive least eigenvalue on each compact x -interval.

At the saddle,

$$T(a) = S, \quad J_0 = \tfrac{3}{2}a, \quad L_0 = 2S + \tfrac{3}{2}a^2, \quad W = L_0 - J_0^2/V = 2S - a^2\kappa > 0. \quad (4.12)$$

Also the envelope identity $G' = -h - s$ gives

$$T = G - xG', \quad T'(a) = -aG''(a) = a\kappa. \quad (4.13)$$

These formulas account for the nonzero correlation between size and occupancy. Ignoring it would give an incorrect least-part variance and prefactor.

5 Full-torus Fourier control and the lattice limit

For a real number z , write $\|z\|_{2\pi}$ for its distance to $2\pi\mathbb{Z}$.

Lemma 5.1 (Consecutive integers). *There is an absolute $c_0 > 0$ such that, for every sufficiently large integer ℓ , every interval E of ℓ consecutive integers, and every $\theta \in [-\pi, \pi]$,*

$$\sum_{j \in E} \|j\theta\|_{2\pi}^2 \geq c_0 \ell \min(1, \ell^2 \theta^2). \quad (5.1)$$

Proof. Let $Q_\ell = \sum_{j \in E} e^{ij\theta}$. Its modulus is independent of the starting point of E , and

$$\sum_{j \in E} \sin^2(j\theta/2) \geq \frac{1}{2}(\ell - |Q_\ell|).$$

If $|\theta| \geq 2\pi/\ell$, the geometric sum and $|\sin(\theta/2)| \geq |\theta|/\pi$ imply $|Q_\ell| \leq \pi/|\theta| \leq \ell/2$. If $|\theta| \leq 2\pi/\ell$, use

$$\ell^2 - |Q_\ell|^2 = 2 \sum_{d=1}^{\ell-1} (\ell - d)(1 - \cos(d\theta)).$$

Retain only $1 \leq d \leq \lfloor \ell/(4\pi) \rfloor$. On these terms $|d\theta| \leq 1/2$, so the sum is at least $c\theta^2\ell^4$. Dividing by $\ell + |Q_\ell| \leq 2\ell$ gives $\ell - |Q_\ell| \geq c\theta^2\ell^3$. Finally $\|z\|_{2\pi}^2 \geq 4\sin^2(z/2)$ proves the claim in both ranges, including $\theta = 0$. $\square$

Lemma 5.2 (Joint minor arcs). *Let $x = tm$ and u range over fixed compact subintervals of $(0, \infty)$. There is $c > 0$, uniform in those parameters, such that*

$$\left| \mathbb{E} e^{i\theta N + i\phi K_m} \right| \leq \exp \left\{ -c \left[t^{-1} \min \left(1, \frac{\theta^2}{t^2} \right) + \frac{\phi^2}{t} \right] \right\}, \quad \theta, \phi \in [-\pi, \pi]. \quad (5.2)$$

In particular,

$$\sup_{r, k \in \mathbb{Z}} \mathbb{P}(N = r, K_m = k) = O(t^2). \quad (5.3)$$

Proof. Choose constants $c_1 < c_2$ with c_1 greater than every allowed x , and use the consecutive block $E_t = \{j : c_1/t \leq j \leq c_2/t\}$. Its length is comparable to t^{-1} . For all these optional sites, the probabilities of $R_j = 0, 1, 2$ have a common positive lower bound.

For a single-site characteristic function χ_j , the pairs $(0, 1)$ and $(1, 2)$ in its squared modulus give

$$|\chi_j(\theta, \phi)| \leq \exp\{-c[\|j\theta + \phi\|_{2\pi}^2 + \|j\theta\|_{2\pi}^2]\}. \quad (5.4)$$

This follows from $1 - \cos z \geq 2\|z\|_{2\pi}^2/\pi^2$ and $\sqrt{1-v} \leq e^{-v/2}$, with a sufficiently small uniform c . The torus triangle inequality gives

$$\|\phi\|_{2\pi}^2 \leq 2(\|j\theta + \phi\|_{2\pi}^2 + \|j\theta\|_{2\pi}^2).$$

Allocate a fixed fraction of the exponent to this bound, and another fraction to Lemma 5.1. Summing over E_t proves (5.2); all other factors have modulus at most one. Fourier inversion and the elementary Gaussian integrals give $O(t^{3/2})$ from θ and $O(t^{1/2})$ from ϕ , hence (5.3). $\square$

This estimate covers rational resonances as well as generic angles. The difference between multiplicities 1 and 2 removes the occupancy phase; consecutive sizes then remove every secondary size peak. No unverified aperiodicity assumption is needed.

Theorem 5.3 (Uniform bivariate lattice limit). *Let $J \subset (0, \infty)$ be compact, let $x = tm \in J$, and choose $u = u(x)$. For integers r, k , put*

$$z_{t,m}(r, k) = \begin{pmatrix} t^{3/2}(r - \mathbb{E}N) \\ t^{1/2}(k - \mathbb{E}K_m) \end{pmatrix}. \quad (5.5)$$

Uniformly in all these m, r, k ,

$$t^{-2} \mathbb{P}(N = r, K_m = k) = \frac{\exp\{-\frac{1}{2} z_{t,m}^T \mathcal{M}(x)^{-1} z_{t,m}\}}{2\pi \sqrt{\det \mathcal{M}(x)}} + o_J(1). \quad (5.6)$$

Proof. Center the variables and put $\theta = t^{3/2}\alpha$, $\phi = t^{1/2}\beta$. For bounded (α, β) , (4.10) gives the limiting quadratic logarithm

$$-\frac{1}{2}(\alpha, \beta)\mathcal{M}(x)(\alpha, \beta)^T.$$

The third-order remainder is uniform. Indeed $\mathbb{E}R_j^r \leq C_r e^{-tj}$ at every optional site, for each fixed $r \geq 1$: $tj \geq \min J > 0$ and u is bounded above and below. Summing third absolute moments of the scaled centered linear forms gives

$$O_J(\sqrt{t})(|\alpha| + |\beta|)^3.$$

The mandatory site's contribution is smaller. These estimates also justify taking logarithms of the individual characteristic factors and passing to their convergent sum.

After scaling, (5.2) bounds the modulus by $e^{-c(\alpha^2 + \beta^2)}$ on $|\alpha| \leq t^{-1/2}$. In the remaining strip the α -length is $O(t^{-3/2})$ and its modulus has a factor $e^{-c/t}$; integration in β remains bounded. Extend the characteristic functions by zero outside the scaled torus. They converge in $L^1(\mathbb{R}^2)$ to the corresponding Gaussian characteristic functions, uniformly in $x \in J$. Uniformity follows from the uniform local expansion, the common Gaussian bound, and compactness of J .

Fourier inversion now proves (5.6). The factor t^2 is precisely the Jacobian $d\theta d\phi = t^2 d\alpha d\beta$. The L^1 error controls the inverse transform uniformly over every lattice target, not just targets with bounded z . $\square$

6 The coefficient equivalent and its normalization

Choose $t = \sqrt{S/n}$, so $n = S/t^2$ exactly. Near the saddle write

$$y = \frac{tm - a}{\sqrt{t}}. \quad (6.1)$$

For bounded y , (4.10) and (4.13) give

$$z_{t,m}(n, m) = (-a\kappa y, 0)^T + o(1). \quad (6.2)$$

Since the first diagonal entry of $\mathcal{M}(a)^{-1}$ is $1/W$, Theorem 5.3 yields

$$\mathbb{P}(N = n, K_m = m) = \frac{t^2}{2\pi\sqrt{VW}} \exp\{-a^2\kappa^2 y^2/(2W)\}(1 + o(1)), \quad (6.3)$$

uniformly on bounded y -intervals. The density is bounded below there, so the absolute local-limit error can be written relatively in this window.

Combining (4.5), (4.6), and

$$G(tm) = S - \frac{1}{2}\kappa t y^2 + O(t^{3/2}|y|^3),$$

gives

$$p_{n,m} = e^{2S/t} \frac{A_0 t^2}{2\pi\sqrt{VW}} \exp\{-\kappa_{\text{eff}} y^2/2\}(1 + o(1)), \quad (6.4)$$

where

$$\kappa_{\text{eff}} = \kappa + \frac{a^2\kappa^2}{W} = \frac{2S\kappa}{W}. \quad (6.5)$$

The y -mesh is $\sqrt{t}$. Therefore for fixed R ,

$$e^{-2S/t} t^{-3/2} \sum_{|y| \leq R} p_{n,m} \longrightarrow \frac{A_0}{2\pi\sqrt{VW}} \int_{-R}^R e^{-\kappa_{\text{eff}} y^2/2} dy.$$

To pass to the whole sum, take a small fixed neighborhood of a on which $G(x) \leq S - c(x - a)^2$. Equations (4.6) and (5.3) bound each term there by

$$p_{n,m} \leq C e^{2S/t} t^2 e^{-cy^2}. \quad (6.6)$$

The resulting normalized Gaussian Riemann sums are uniformly integrable. Outside the neighborhood, Lemma 3.2 is exponentially smaller than $e^{2S/t} t^{3/2}$. Hence $R \rightarrow \infty$ is legitimate. The prefactor is

$$\frac{A_0}{\sqrt{2\pi VW \kappa_{\text{eff}}}} = \frac{A_0}{2\sqrt{\pi S V \kappa}} = \frac{D}{2\sqrt{\pi S}}. \quad (6.7)$$

We have proved

$$p_n \sim \frac{D}{2\sqrt{\pi S}} t^{3/2} e^{2S/t} = K n^{-3/4} e^{2\sqrt{S}n}. \quad (6.8)$$

This is coefficient extraction from a genuinely joint lattice estimate, including the entire Fourier torus.

7 A three-variable saddle to every fixed order

The previous proof establishes the leading constant with transparent probabilistic normalization. For higher orders, it is more efficient to expand the exact double Fourier integral and the discrete minimum simultaneously. A carefully chosen *linear* positive radius makes the resulting three-variable Gaussian absolutely integrable.

7.1 An exact contour with a linear occupancy radius

For complex z with $\Re z > 0$, use $H_m(z, e^s)$ for the analytic version of (4.1). For any positive real occupancy radius e^s , coefficient inversion gives

$$p_{n,m} = \frac{1}{(2\pi)^2} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} e^{nz - m(s+i\phi)} H_m(z, e^{s+i\phi}) d\theta d\phi, \quad z = t - i\theta. \quad (7.1)$$

The radius can be chosen separately for every integer m . Near a/t , set

$$x = tm, \quad s_0 = -\log 2, \quad b = \frac{3}{2V}, \quad s_{\text{lin}}(x) = s_0 + b(x - a). \quad (7.2)$$

This is an exact choice of contour; it does not replace an exact coefficient by an approximation.

Put $X = x - a$, $U = s_{\text{lin}}(x) - s_0 + i\phi$, and $Z = (z - t)/t = -i\theta/t$. The moving endpoint is

$$v = zm = (a + X)(1 + Z). \quad (7.3)$$

With $n = S/t^2$, the large phase in (7.1) is P/t , where

$$P(X, U, Z) = \frac{I((a + X)(1 + Z), s_0 + U)}{1 + Z} - (a + X)(s_0 + U) + S(1 + Z). \quad (7.4)$$

At the origin, $P = 2S$, its gradient vanishes, and its Hessian is

$$H = \begin{pmatrix} 3/2 & -3/2 & 3a/2 \\ -3/2 & V & -3a/2 \\ 3a/2 & -3a/2 & 2S + 3a^2/2 \end{pmatrix}, \quad \det H = -2SV\kappa \neq 0. \quad (7.5)$$

Writing $\lambda = \sqrt{t}$, parameterize the integration plane by

$$X = \lambda y, \quad U = \lambda(by + i\beta), \quad Z = -i\lambda\alpha. \quad (7.6)$$

The quadratic term in the exponent becomes

$$Q(y, \beta, \alpha) = -\frac{\kappa}{2}y^2 - \frac{1}{2}(L_0\alpha^2 + 2J_0\alpha\beta + V\beta^2) + ia\kappa y\alpha. \quad (7.7)$$

Its real part is strictly negative definite because $\kappa > 0$ and $\mathcal{M}(a)$ is positive definite. Thus all polynomial Gaussian moments used below are absolutely convergent on this plane.

The real radial phase on the linear radius is

$$G_{\text{lin}}(x) = I(x, s_{\text{lin}}(x)) - xs_{\text{lin}}(x). \quad (7.8)$$

Its value, first derivative, and second derivative at a are $S, 0, -\kappa$. The second derivative agrees with the optimized curvature because $b = s'(a)$. Hence $G_{\text{lin}}(x) \leq S - c_0(x - a)^2$ on a sufficiently small fixed neighborhood.

7.2 Localization before expansion

Lemma 5.2 requires only compact positive ranges for x and u , not the equation $B = x$. It therefore applies unchanged to the linear radius. Its characteristic-function bound, multiplied by the real normalization in (7.1), yields an absolute Gaussian bound

$$Ce^{-c_1y^2 - c_2(\alpha^2 + \beta^2)} \quad (7.9)$$

after removing $e^{2S/t}$, on $|\theta| \leq t$. On $|\theta| > t$, the factor $e^{-c/t}$ dominates the polynomial length of the scaled α -interval. The β -integral retains a Gaussian bound. For tm outside this fixed neighborhood, revert to the positive bounds of Lemma 3.2; choosing different radii in different m ranges is legitimate in the exact formula.

Fix $0 < \varepsilon < 1/10$. The sum and integral can consequently be restricted to

$$|y|, |\alpha|, |\beta| \leq t^{-\varepsilon} \quad (7.10)$$

with normalized error $O(t^M)$ for every fixed M . This follows by integrating Gaussian tails and summing their shifted-grid analogues. It is an absolute estimate made before Taylor expansion, so it does not presume cancellation on minor arcs.

7.3 Complex Euler–Maclaurin with a moving endpoint

The identity

$$h(v, s) = \log(1 + (e^s - 1)e^{-v}) - \log(1 - e^{-v}) \quad (7.11)$$

defines the required branches by the power-series logarithm. In the local box, $v = zm$ stays near a , e^s stays near $1/2$, and along the ray $z[m, \infty)$ both logarithmic arguments remain in their zero-free domains. More explicitly, $\Re(zr) = tr \geq tm$ is bounded below and $|e^s - 1|$ stays close to $1/2$, so both $|e^{-zr}| < 1$ and $|(e^s - 1)e^{-zr}| < 1$ uniformly. Derivatives decay exponentially in $\Re v$.

Applying Euler–Maclaurin to the real j variable gives, for every fixed integer $p \geq 1$,

$$\sum_{j>m} h(zj, s) = \frac{I(zm, s)}{z} - \frac{1}{2}h(zm, s) - \sum_{r=1}^{p-1} \frac{B_{2r}}{(2r)!} z^{2r-1} h_{v, 2r-1}(zm, s) + O_p(t^{2p-1}), \quad (7.12)$$

where B_{2r} denotes the Bernoulli number. The remainder is uniform in (7.10): the periodic-Bernoulli remainder is bounded by a constant times

$$|z|^{2p} \int_m^\infty |h_{v, 2p}(zr, s)| dr = O(t^{2p-1}), \quad (7.13)$$

since $|z| \asymp t$ there. The ray integral equals $I(zm, s)/z$ by contour deformation in the zero-free right half-plane; the exponentially decaying tail permits closing the deformation at infinity. Alternatively, integrating the uniformly convergent exponential series in (7.11) proves the same identity term by term.

The leading amplitude is now the analytic function

$$\mathcal{A}(v, s) = \frac{e^s e^{-v}}{1 - e^{-v}} e^{-h(v, s)/2}. \quad (7.14)$$

Its value at the saddle is $A_0 = 2^{-1/2}$. The first Euler–Maclaurin amplitude correction is $-zh_v(v, s)/12$; at the saddle its coefficient of t is

$$-\frac{h_v(a, s_0)}{12} = \frac{1}{8}. \quad (7.15)$$

7.4 Uniform Taylor remainders and the shifted lattice

After factoring out $A_0 e^{2S/t}$, Taylor expansion of (7.4), (7.14), and (7.12) on (7.6) gives, for each fixed N ,

$$e^{Q(w)} \left(\sum_{j=0}^N \lambda^j P_j(w) + \mathcal{R}_N(t, w) \right), \quad w = (y, \beta, \alpha), \quad (7.16)$$

where $P_0 = 1$ and every P_j is an explicitly calculable polynomial. The parity is

$$P_j(-w) = (-1)^j P_j(w). \quad (7.17)$$

Indeed a phase monomial of degree k contributes λ^{k-2} times a homogeneous polynomial of degree k , and an amplitude monomial of degree k contributes λ^k . Each base Euler–Maclaurin correction has even λ -order $4r - 2$. Products preserve (7.17).

Here the expansion is asymptotic, not merely formal. To see this, analytically continue the phase and amplitude to a fixed neighborhood of the origin. Their order- k Taylor terms are bounded by constants times $|w|^k$ after substitution. Beyond the quadratic, the phase perturbation is bounded by $C\lambda|w|^3$ on the local box. Because $\lambda|w| \rightarrow 0$ uniformly there, its exponential can be absorbed in

a fixed fraction of the negative Gaussian $\Re Q$. Taylor's integral remainder in λ , after the removable singularity of $(P - 2S)/\lambda^2$ has been removed, is bounded by λ^{N+1} times a fixed polynomial in $|w|$, multiplied by this slightly weakened Gaussian. Taking the Euler–Maclaurin order large enough and treating its uniform residual separately yields constants $C_N, c_N > 0$ and an integer d_N such that

$$|\mathcal{R}_N(t, w)| e^{\Re Q(w)} \leq C_N \lambda^{N+1} (1 + |w|^{d_N}) e^{-c_N |w|^2}. \quad (7.18)$$

No derivatives of the Euler–Maclaurin residual are needed: its uniform $O(t^{2p-1})$ bound can be chosen smaller than the requested order before expansion.

The y values lie on a shifted lattice of mesh λ . After integrating α, β , each coefficient $e^Q P_j$ is a Schwartz function of y . For any such function f and every fixed integer $M \geq 1$,

$$\lambda \sum_{k \in \mathbb{Z}} f(\lambda(k + \delta)) = \int_{\mathbb{R}} f(y) dy + O_M(\lambda^M), \quad 0 \leq \delta < 1, \quad (7.19)$$

uniformly in the shift. One proof applies Euler–Maclaurin on the whole real line with arbitrarily many derivatives; all endpoint terms vanish and the remainder is bounded by a constant times $\lambda^M \|f^{(M)}\|_1$. Truncating or extending the local grid costs less than every prescribed power by Gaussian tails.

Thus the shifted grid does not need to be symmetric. First replace its coefficient sums by integrals using (7.19); then every odd coefficient vanishes by the simultaneous sign reversal $w \mapsto -w$. Applying (7.18) with $N = 2R + 1$ leaves a normalized error $O(\lambda^{2R+2}) = O(t^{R+1})$.

The exact measure is $d\theta d\phi = t^2 d\alpha d\beta$, together with a y -mesh $\sqrt{t}$. The overall scale is therefore $t^{3/2}$. The Gaussian normalization in Section 6, or direct integration of (7.7), gives

$$p_n = \frac{D}{2\sqrt{\pi S}} t^{3/2} e^{2S/t} \left(1 + b_1 t + \cdots + b_R t^R + O_R(t^{R+1}) \right). \quad (7.20)$$

As $t = \sqrt{S/n}$, setting $c_j = b_j S^{j/2}$ proves the existence and remainder in Theorem 1.1. The construction is effective: take finite jets of the analytic functions, multiply finite polynomials to the desired order, and integrate their Gaussian moments. The same argument with the size-Fourier variable absent gives a real-radial expansion to every fixed order. The coefficient proof has not inferred this expansion from an unproved complex continuation of the radial answer.

7.5 An explicit coefficient prescription and its field

For a compact formal prescription, use independent formal variables ξ, v, ζ and write $w = (\xi, v, \zeta)$. Define

$$v_\lambda = (a + \lambda\xi)(1 + \lambda\zeta), \quad s_\lambda = s_0 + \lambda v. \quad (7.21)$$

All expressions below are formal series in λ with polynomial coefficients in w . Set

$$\begin{aligned} \mathcal{R}(\lambda, w) = \frac{\mathcal{A}(v_\lambda, s_\lambda)}{A_0} \exp \left\{ \frac{P(\lambda\xi, \lambda v, \lambda\zeta) - 2S}{\lambda^2} - \frac{1}{2} w^T H w \right. \\ \left. - \sum_{r \geq 1} \frac{B_{2r}}{(2r)!} \lambda^{4r-2} (1 + \lambda\zeta)^{2r-1} h_{v^{2r-1}}(v_\lambda, s_\lambda) \right\}. \end{aligned} \quad (7.22)$$

The apparent division by λ^2 is removable after subtracting the constant and using the vanishing gradient. The quadratic term is explicitly subtracted, so the exponent inside the first line starts at

positive λ -order. For a specified coefficient, only finitely many r in the second line matter. Let $\mathcal{E}_H$ be the normalized Wick contraction with covariance $-H^{-1}$, defined explicitly in Section 8. Then the relative coefficients in (7.20) are

$$b_j = \mathcal{E}_H([\lambda^{2j}]\mathcal{R}(\lambda, w)), \quad c_j = S^{j/2}b_j. \quad (7.23)$$

This is a finite prescription for each fixed j . The integration plane in (7.6) explains the normalized Gaussian contraction, but need not be substituted again in the invariant formula (7.23).

Corollary 7.1 (Field of the relative coefficients). *For every $j \geq 0$, $b_j \in \mathbb{Q}(a, S)$ and $c_j \in \mathbb{Q}(a, \sqrt{S})$.*

Proof. At $q = e^{-v} = 2/3$, $u = e^s = 1/2$, every positive-order derivative of h is rational, because differentiating its two logarithms gives rational functions of q, u . Derivatives of I involving v and of total degree at least two are therefore rational. Pure s derivatives of I come from

$$I_s = -\frac{u}{1-u} \log(1 - (1-u)q);$$

the only undifferentiated logarithm has saddle value $\log(2/3) = -a$. These derivatives belong to $\mathbb{Q}(a)$.

The exceptional low-degree values are $I_0 = S - a \log 2$, $I_v = -\log 2$, and $I_s = a$. With $\Delta v = (a + X)(1 + Z) - a$, the $\log 2$ terms in the full I jet are $-(a + \Delta v) \log 2$. Dividing by $1 + Z$ makes them $-(a + X) \log 2$, which cancels exactly with the s_0 part of $-(a + X)(s_0 + U)$ in P . Thus every coefficient of the complete phase is in $\mathbb{Q}(a, S)$.

The positive-order derivatives of $\log \mathcal{A}$ are rational at the saddle. After A_0 has been factored out, its amplitude jet and the Euler–Maclaurin jets have coefficients in $\mathbb{Q}(a)$ after the moving-endpoint substitution. Bernoulli numbers are rational. Finally the entries of $-H^{-1}$ lie in $\mathbb{Q}(a, S)$. Finite products, formal exponentiation to a fixed order, and Wick contractions preserve this field. The Gaussian determinant and A_0 are already absorbed into the leading factor, so they create no additional roots in these relative coefficients. All denominators are well-defined at the stated constants because the analytic functions and H^{-1} exist there. Algebraic independence of a, S is neither asserted nor needed. $\square$

8 The first two explicit corrections

We give the algebra in a form that can be checked by finite polynomial operations. This also specifies how higher coefficients are obtained.

Let P_k be the homogeneous degree- k term in the Taylor expansion of P in (7.4), with coordinates (X, U, Z) . Let ℓ_1, ℓ_2 be the homogeneous degree-one and degree-two terms in

$$\log \mathcal{A}((a + X)(1 + Z), s_0 + U) - \log A_0. \quad (8.1)$$

Let $\mathcal{E}_H$ denote normalized Gaussian contraction on the plane (7.6), with the quadratic form (7.7). Its covariance in the complex coordinates is $C = -H^{-1}$:

$$C = \begin{pmatrix} \frac{-4Sa + S - 3a^3 + 3a^2}{6S(a-1)} & -\frac{1}{2(a-1)} & \frac{a}{2S} \\ -\frac{1}{2(a-1)} & -\frac{1}{2(a-1)} & 0 \\ \frac{a}{2S} & 0 & -\frac{1}{2S} \end{pmatrix}. \quad (8.2)$$

This is a contraction notation, not a positive real probability covariance. Absolute Gaussian convergence of the plane was proved in (7.7). Integration by parts, or completing the square, proves the usual Wick recurrence:

$$\mathcal{E}_H[v_i v_1^{e_1} \cdots v_d^{e_d}] = \sum_{j=1}^d e_j C_{ij} \mathcal{E}_H[v_1^{e_1} \cdots v_j^{e_j-1} \cdots v_d^{e_d}], \quad (8.3)$$

with the evident interpretation that one chosen factor v_i was removed before applying the formula. Equivalently, even monomial moments are sums over all pairings of products of covariance entries; odd moments vanish.

The coefficient of t in (7.20) is exactly

$$b_1 = \frac{1}{8} + \mathcal{E}_H \left[\ell_2 + \frac{1}{2} \ell_1^2 + \ell_1 P_3 + P_4 + \frac{1}{2} P_3^2 \right]. \quad (8.4)$$

The first term is (7.15). There is no omitted varying Jacobian: the plane transformation is linear. The minimum-size factor is already part of $\mathcal{A}$, and the full size term nz is already part of P .

For a completely specified derivative recipe, write $q = e^{-v}$ and $u = e^s$. Then

$$\begin{aligned} h &= \log(1 - (1 - u)q) - \log(1 - q), \\ I_v &= -h, & I_s &= -\frac{u}{1 - u} \log(1 - (1 - u)q), \\ \partial_v &= -q \partial_q, & \partial_s &= u \partial_u, \\ \log \mathcal{A} &= \log u + \log q - \frac{1}{2} \log(1 - q) - \frac{1}{2} \log(1 - (1 - u)q). \end{aligned} \quad (8.5)$$

Apply these operators at $(q, u) = (2/3, 1/2)$, taking $I(a, s_0) = S - a \log 2$. Taylor-expand I to degree four and $\log \mathcal{A}$ to degree two, compose the endpoint shift

$$v - a = X + aZ + XZ, \quad (8.6)$$

and insert the geometric truncation $1/(1 + Z) = 1 - Z + Z^2 - Z^3 + Z^4 + O(Z^5)$. These instructions determine every term in (8.4) without a fitted numerical constant.

To display the simplification compactly, put $d = a - 1$. The five respective contributions in (8.4) are

$$\begin{aligned} E_0 &= \frac{1}{8}, \\ E_1 &= -\frac{45Sa - 20S + 18a^2 - 18a}{16Sd}, \\ E_2 &= \frac{3(7Sa - 3S + 3a^2 - 3a)}{8Sd}, \\ E_3 &= \frac{S^2(76a^2 - 11a - 8) + S(42a^3 - 12a^2 - 69a + 39) + 27a^2d^2}{48S^2d^2}, \\ E_4 &= -\frac{S^2(245a^2 - 106a + 5) + S(126a^3 - 9a^2 - 261a + 144) + 81a^2d^2}{144S^2d^2}. \end{aligned} \quad (8.7)$$

Here E_1 is the contraction of $\ell_2 + \ell_1^2/2$, E_2 of $\ell_1 P_3$, E_3 of P_4 , and E_4 of $P_3^2/2$. Summing gives

$$\begin{aligned} b_1 &= -\frac{26Sa^2 - 82Sa + 29S + 27a^2 - 54a + 27}{144S(a - 1)^2} \\ &= B_* - \frac{3}{16S}. \end{aligned} \quad (8.8)$$

Thus $c = b_1\sqrt{S}$, as asserted.

For comparison, performing the same contraction in the two-dimensional radial phase $I(a + X, s_0 + U) - (a + X)(s_0 + U)$ gives the five pieces

$$\frac{1}{8}, \quad -\frac{5(9a-4)}{16d}, \quad \frac{3(7a-3)}{8d}, \quad \frac{76a^2-11a-8}{48d^2}, \quad -\frac{245a^2-106a+5}{144d^2}. \quad (8.9)$$

Their sum is B_* . This independently evaluates the radial coefficient in (1.6). It does not replace the three-variable extraction that produced the $-3/(16S)$ contribution in (8.8).

8.1 An exact sign proof

The positive atanh series gives

$$a = 2 \sum_{k \geq 0} \frac{5^{-(2k+1)}}{2k+1} < \frac{2}{5} + \frac{2}{375} + \frac{1}{7500} = \frac{3041}{7500} < \frac{811}{2000} < 1. \quad (8.10)$$

For the tail $k \geq 2$, replace denominators by 5 and sum the geometric series. The bound is strict because at least one original denominator exceeds 5. The polynomial $Q(z) = 26z^2 - 82z + 29$ is strictly decreasing on $[0, 1]$, and exact arithmetic gives

$$Q(811/2000) = \frac{48373}{2000000} > 0. \quad (8.11)$$

Therefore $Q(a) > 0$, so $B_* < 0$ and $c < 0$. In particular the leading expression $Kn^{-3/4}e^{2\sqrt{Sn}}$ eventually overestimates p_n . This sign conclusion does not depend on approximate values of the saddle constants.

8.2 The second coefficient by a finite contraction

The same invariant prescription gives an explicit second coefficient. Put

$$B_{**} = \frac{676a^4 + 3416a^3 - 10200a^2 + 22172a - 7559}{41472(a-1)^4}. \quad (8.12)$$

Then

$$c_2 = SB_{**} - \frac{15}{16}B_* - \frac{15}{512S}, \quad (8.13)$$

and the stronger explicit expansion is

$$p_n = Kn^{-3/4}e^{2\sqrt{Sn}} \left(1 + \frac{c}{\sqrt{n}} + \frac{c_2}{n} + O(n^{-3/2}) \right). \quad (8.14)$$

The symbol B_{**} in this decomposition is an explicit rational function of a ; the calculation below extracts coefficients directly from the three-variable phase.

Here are finite algebraic details sufficient to reconstruct this calculation. Retain the homogeneous phase terms P_k through $k = 6$ and the log-amplitude terms ℓ_k through $k = 4$, extending the definitions in (8.1). Let E_j be the degree- j part of

$$E(X, U, Z) = -\frac{1+Z}{12} h_v((a+X)(1+Z), s_0+U).$$

The required Euler–Maclaurin terms are

$$\begin{aligned} E_0 &= \frac{1}{8}, \\ E_1 &= \frac{U}{16} - \frac{7X}{16} + \left(\frac{1}{8} - \frac{7a}{16}\right) Z, \\ E_2 &= -\frac{UX}{8} + \left(\frac{1}{16} - \frac{a}{8}\right) UZ + \frac{19X^2}{16} \\ &\quad + \left(\frac{19a}{8} - \frac{7}{8}\right) XZ + \left(\frac{19a^2}{16} - \frac{7a}{16}\right) Z^2. \end{aligned} \tag{8.15}$$

Define, in the same abstract complex-coordinate basis as (8.2),

$$\begin{aligned} r_1 &= P_3 + \ell_1, & r_2 &= P_4 + \ell_2 + E_0, \\ r_3 &= P_5 + \ell_3 + E_1, & r_4 &= P_6 + \ell_4 + E_2. \end{aligned} \tag{8.16}$$

Under scaling by λ , the logarithm of the normalized factor is $\lambda r_1 + \lambda^2 r_2 + \lambda^3 r_3 + \lambda^4 r_4 + \dots$. The coefficient of $t^2 = \lambda^4$ is therefore exactly

$$b_2 = \mathcal{E}_H \left[r_4 + r_1 r_3 + \frac{1}{2} r_2^2 + \frac{1}{2} r_1^2 r_2 + \frac{1}{24} r_1^4 \right]. \tag{8.17}$$

This includes $E_0^2/2$ and all mixed phase, amplitude, and endpoint terms. The next Bernoulli term starts at t^3 and cannot contribute. The polynomial in brackets has degree at most twelve; Wick pairing or six applications of the finite Gaussian heat operator evaluate it exactly. Using (8.5) for the phase and amplitude derivatives, this calculation gives

$$b_2 = B_{**} - \frac{15B_*}{16S} - \frac{15}{512S^2}. \tag{8.18}$$

Multiplication by S gives (8.13).

The optional second-coefficient program computes the derivative jets and the contraction in (8.17). It uses the linear coordinate $D_c = X + aZ$, whose covariance with Z vanishes, to keep the polynomial operations small; it explicitly checks this transformed covariance against the transformed Hessian. The resulting rational identity is checked against (8.12) and (8.18). This is exact symbolic algebra, not a fit to finite sequence values.

For the analytic error in (8.14), invoke the already proved $R = 2$ case of Section 7. Computing phase degree six and log-amplitude degree four identifies the λ^4 coefficient; it does not alone establish an unintegrated $O(\lambda^6)$ error. In the remainder proof take $N = 5$: the next λ^5 polynomial exists, is odd, and has zero full Gaussian integral, while its shifted-lattice error is smaller than every fixed power. The remaining normalized error is $O(\lambda^6) = O(t^3)$. All denominators in the exact answer are nonzero since $S > 0$ and $0 < a < 1$.

[write] Decimals and a numerical check, 7 October 2026. The source prints no decimal value. For orientation, the write’s values (mpmath at 60 digits from the closed forms, every digit a truncation) are $a = 0.4054651081\dots$, $S = 0.7481056530\dots$, $D = 0.5294631036\dots$, $K = 0.1389061592\dots$, $B_* = -0.0005169346\dots$, $c = c_1 = -0.2172274082\dots$, $B_{**} = 0.0000099188\dots$, $c_2 = -0.0386693672\dots$, $V = 0.3109302162\dots$, $\kappa = 5.7363504177\dots$, $W = 0.5531440891\dots$ and $\kappa_{\text{eff}} = 15.5163772317\dots$. With $\varrho_n = p_n/(Kn^{-3/4}e^{2\sqrt{S}n})$ and the exact counts, the quantity $(\varrho_n - 1 - c/\sqrt{n} - c_2/n)n^{3/2}$ is $-0.0155\dots$, $-0.0202\dots$, $-0.0204\dots$, $-0.0203\dots$ at $n = 400, 800, 1200, 1500$: it stays bounded, as (8.14) requires, while $(\varrho_n - 1 - c/\sqrt{n})n$ is $-0.0394\dots$, $-0.0393\dots$, $-0.0391\dots$ at $n = 400, 800, 1500$, close to c_2 . This is a finite numerical observation; it neither proves the expansion nor determines c_3 .

9 The least part in the constrained ensemble

Theorem 9.1 (Gaussian law and bounded-window local masses). *Choose an admissible partition uniformly at size n , let M_n be its least part, and set $t = \sqrt{S/n}$. With*

$$z_m = \sqrt{t\kappa_{\text{eff}}} \left(m - \frac{a}{t} \right), \quad \varphi(z) = \frac{e^{-z^2/2}}{\sqrt{2\pi}}, \quad (9.1)$$

for every fixed $R < \infty$,

$$\mathbb{P}(M_n = m) = \sqrt{t\kappa_{\text{eff}}} \varphi(z_m)(1 + o(1)) \quad (9.2)$$

uniformly for integer m with $|z_m| \leq R$. Moreover $z_{M_n} \Rightarrow N(0, 1)$.

Proof. Divide (6.4) by (6.8). Its constants simplify using (6.7), giving (9.2). The standardized mesh is $\sqrt{t\kappa_{\text{eff}}}$.

To prove tightness, (6.6) and the total equivalent imply, throughout a fixed neighborhood of $tm = a$,

$$\mathbb{P}(M_n = m) \leq C\sqrt{t}e^{-cy^2}, \quad y = (tm - a)/\sqrt{t}.$$

Outside that neighborhood, the total probability is exponentially small, up to harmless polynomial factors, by Lemma 3.2. The Gaussian Riemann sums therefore have uniformly vanishing tails as their standardized window grows. On a bounded window, (9.2) turns sums against bounded continuous test functions into the corresponding Gaussian integral. Taking the window to infinity proves weak convergence. $\square$

Thus the least part is centered to first order at $a\sqrt{n/S}$ and fluctuates on the $n^{1/4}$ scale. The theorem supplies neither a higher-order centering constant nor a relative local approximation uniform in unbounded standardized windows. The full-target uniformity in Theorem 5.3 is an *absolute* joint local-limit estimate; it must not be misread as such a stronger relative law for M_n .

10 Eventual monotonicity and a rounded inverse

Taking logarithms of (1.5) gives

$$\log p_n = 2\sqrt{Sn} - \frac{3}{4}\log n + \log K + \frac{c}{\sqrt{n}} + O(n^{-1}). \quad (10.1)$$

Subtracting adjacent instances,

$$\log p_{n+1} - \log p_n = \frac{\sqrt{S}}{\sqrt{n}} + O(n^{-1}) > 0 \quad (10.2)$$

for sufficiently large n . No derivative estimate for an unknown remainder sequence has been used: the difference of two $O(n^{-1})$ quantities is still $O(n^{-1})$, which is already smaller than the main increment.

Let

$$f_0(x) = 2\sqrt{Sx} - \frac{3}{4}\log x + \log K. \quad (10.3)$$

For sufficiently large Y , its large inverse root $r = r(Y)$ satisfies $f_0(r) = \log Y$. Setting $v = \sqrt{r}$, the equation is $Kv^{-3/2}e^{2\sqrt{S}v} = Y$. Raising to power $-2/3$ and multiplying by $-4\sqrt{S}/3$ gives

$$\left(-\frac{4\sqrt{S}}{3}v\right)e^{-4\sqrt{S}v/3} = -\frac{4\sqrt{S}}{3}(K/Y)^{2/3}.$$

The lower real branch W_{-1} selects the large solution and proves (1.9).

Choose $C_0 > 0$ so that, for all sufficiently large integers n ,

$$f_-(n) \leq \log p_n \leq f_+(n), \quad f_{\pm}(x) = f_0(x) + \frac{c}{\sqrt{x}} \pm \frac{C_0}{x}. \quad (10.4)$$

Both smooth envelopes are increasing eventually. Their roots at height $\log Y$ obey

$$x_{\pm}(Y) = r(Y) - \frac{c}{\sqrt{S}} + O(r(Y)^{-1/2}). \quad (10.5)$$

Indeed $f'_0(r) = \sqrt{S}/\sqrt{r} + O(r^{-1})$, and a bounded displacement d changes f_0 by $d\sqrt{S}/\sqrt{r} + O(r^{-1})$. The choice $d = -c/\sqrt{S}$ cancels the correction $c/\sqrt{r}$. The residual is $O(r^{-1})$, which corresponds to an index displacement $O(r^{-1/2})$ by the mean value theorem.

The upper envelope reaches the target no later than the true sequence can; the lower envelope reaches it no earlier than necessary for the upper bound. More explicitly, any integer $n < x_+(Y)$ has $\log p_n < \log Y$, while every sufficiently large integer $n \geq x_-(Y)$ has $\log p_n \geq \log Y$. Hence

$$\lceil x_+(Y) \rceil \leq N(Y) \leq \lceil x_-(Y) \rceil.$$

Once Y exceeds the finitely many early sequence values, those early terms cannot affect $N(Y)$. Enlarging a fixed C in (10.5) proves (1.10). The correction $-c/\sqrt{S}$ is positive, consistent with the eventual overestimate by the leading model.

10.1 Inverse envelopes to every fixed order

The preceding constant-shift bracket extends to arbitrary fixed order, with an exact indexing convention. Define the logarithmic coefficients by the formal identity

$$\log \left(1 + \sum_{j \geq 1} c_j z^j \right) = \sum_{j \geq 1} \ell_j z^j. \quad (10.6)$$

Thus $\ell_1 = c_1 = c$ and $\ell_2 = c_2 - c_1^2/2$. These are finite rational polynomials in the c_j at each order.

Theorem 10.1 (All fixed inverse orders). *There are consistently defined constants $\delta_0, \delta_1, \dots$ in $\mathbb{Q}(a, \sqrt{S})$ such that, for every fixed integer $J \geq 0$, putting*

$$T_J(r) = r + \sum_{k=0}^{J-1} \delta_k r^{-k/2}, \quad (10.7)$$

with an empty sum for $J = 0$, there is $A_J > 0$ for which

$$\left\lceil T_J(r(Y)) - A_J r(Y)^{-J/2} \right\rceil \leq N(Y) \leq \left\lceil T_J(r(Y)) + A_J r(Y)^{-J/2} \right\rceil \quad (10.8)$$

for all sufficiently large Y . The first two formal coefficients are

$$\delta_0 = -\frac{c_1}{\sqrt{S}}, \quad \delta_1 = \frac{3\delta_0/4 - \ell_2}{\sqrt{S}}. \quad (10.9)$$

Here $\ell_2 = c_2 - c_1^2/2$ uses the explicit coefficient (8.13), so both displayed inverse corrections are given by exact expressions in a, S .

Proof. The forward expansion gives, for each fixed J ,

$$\log p_n = f_0(n) + \sum_{j=1}^J \ell_j n^{-j/2} + O_J(n^{-(J+1)/2}).$$

Add and subtract $C_J x^{-(J+1)/2}$ to the smooth expression to obtain envelopes $g_J^\pm(x)$. Their derivatives are asymptotic to $\sqrt{S}x^{-1/2}$, so both are increasing eventually. Their large roots $\rho_J^\pm$ at level $\log Y$ bracket the integer threshold after ceilings, exactly as in (10.4). Comparing at $r \pm D$ for a sufficiently large fixed D shows $\rho_J^\pm = r + O_J(1)$; their derivatives there are bounded below by a positive constant times $r^{-1/2}$.

For the explicit recursion put $q = r^{-1/2}$ and $x = r + d$. Subtracting $f_0(r)$, the envelope-free residual is

$$\begin{aligned} & 2\sqrt{S}q^{-1}((1+q^2d)^{1/2} - 1) - \frac{3}{4}\log(1+q^2d) \\ & + \sum_{j=1}^J \ell_j q^j (1+q^2d)^{-j/2}. \end{aligned} \quad (10.10)$$

For bounded d this is analytic in q at zero after removal of the displayed removable singularity. For $J \geq 1$, divide the residual by q . Its leading term is $\sqrt{S}d + \ell_1$. Substitute

$$d = \delta_0 + \delta_1 q + \cdots + \delta_{J-1} q^{J-1}$$

and cancel successive coefficients. At each step the coefficient of the new δ_k is $\sqrt{S} \neq 0$; only $\ell_1, \dots, \ell_{k+1}$ are needed. This proves consistency as J increases, and gives (10.9) from the first two equations.

The finite substitution leaves the unscaled residual $O_J(q^{J+1})$. The envelope term has that same order. Dividing by the lower bound of order q for the derivative, the mean value theorem gives

$$\rho_J^\pm = T_J(r) + O_J(q^J).$$

For $J = 0$, use $d = 0$: the envelope residual is $O(q)$, so the root error is $O(1)$ without referring to undefined coefficients. Enlarging A_J and applying the monotonicity of ceilings proves (10.8). Corollary 7.1, (10.6), and the rational recursion show $\delta_k \in \mathbb{Q}(a, \sqrt{S})$. $\square$

The $J = 0$ bracket has bounded real width; shrinking begins at $J = 1$. Keeping δ_0 through δ_R requires $J = R + 1$ and gives real error $O(r^{-(R+1)/2})$. These are smooth-envelope inverse expansions, not an identification with ordinary piecewise-linear interpolation of the integer sequence. Neither convergence of an infinite inverse series nor one unconditional rounded equality is asserted.

Remark 10.2 ([write] The inverses and the transseries volume, 7 October 2026). Statement by statement, against the volume named in Section 1.2 (its symbols carry the subscript “vol”).

- (a) *Instance.* Let n_1 be an index from which p_n is strictly increasing (it exists by Section 10). For $Y > \max_{n < n_1} p_n$, $N(Y)$ is the integer staircase $N_*(Y)$ of `p0:def:three-inverses` with $A_n = p_n$, so `p0:thm:staircase(1)` applies: $N(Y) = \lceil \nu(Y) \rceil$ for every admissible interpolation. In the exact counts $p_{n+1} > p_n$ for $33 \leq n < 1500$, and the last decrease is $p_{33} = 184 < p_{32} = 190$; that is a finite observation, not a proof that $n_1 = 33$.
- (b) *Instance.* The root $r(Y)$ of (1.9) is `p0:thm:lambert-core(3)`, case $b_{\text{vol}} < 0$, after the substitution $X_{\text{vol}} = \sqrt{r}$: with $a_{\text{vol}} = 2\sqrt{S}$, $b_{\text{vol}} = -\frac{3}{2}$, $L_{\text{vol}} = \log(Y/K)$ the core equation $a_{\text{vol}}X + b_{\text{vol}} \log X = L_{\text{vol}}$ is $f_0(X^2) = \log Y$, and the large solution $(|b_{\text{vol}}|/a_{\text{vol}})(-W_{-1}(z))$ with $z = -(a_{\text{vol}}/|b_{\text{vol}}|)e^{-L_{\text{vol}}/|b_{\text{vol}}|} = -\frac{4}{3}\sqrt{S}(K/Y)^{2/3}$ is exactly the bracket in (1.9); it exists once L_{vol} exceeds the fold value L_c of `p0:eq:core-threshold`.
- (c) *Formal instance.* In the variable $X = \sqrt{x}$ the forward expansion is the exponential–power model of `p0:def:model` with data $(K, 2\sqrt{S}, 1, -\frac{3}{2}, 1, \sum_j \ell_j t^j)$ (equivalently $(K, 2\sqrt{S}, \frac{1}{2}, -\frac{3}{4}, \frac{1}{2}, \sum_j \ell_j t^j)$ in x). The coefficients δ_k of Theorem 10.1 are those of the Lambert-centred series of `p0:thm:lambert-centered(4)` (formal exactness), squared: if $\sqrt{x} = X + \sum_{n \geq 1} \gamma_n X^{-n}$ is that series about $X = \sqrt{r(Y)}$, then $\delta_k = 2\gamma_{k+1} + \sum_{i+j=k, i, j \geq 1} \gamma_i \gamma_j$. Both are the unique formal solutions of the same residual equation, so they agree; the write checked $\delta_0 = 2\gamma_1$ and $\delta_1 = 2\gamma_2$ symbolically from the volume’s Bell recurrence ($\gamma_1 = -\ell_1/(2\sqrt{S})$, $\gamma_2 = -(-\frac{3}{2}\gamma_1 + \ell_2)/(2\sqrt{S})$) and numerically, $\delta_0 = 0.2511499825\dots$, $\delta_1 = 0.2897638360\dots$. That coefficient identity is formal. *Instance:* the smooth envelopes $g_j^\pm$ in the proof of Theorem 10.1 are exactly of exponential–power type in X , with data $(K, 2\sqrt{S}, 1, -\frac{3}{2}, 1, \sum_{j \leq J} \ell_j t^j \pm C_J t^{J+1})$, so the root estimate $\rho_j^\pm = T_J(r) + O_J(r^{-J/2})$ there is `p0:thm:lambert-centered(5)` with $N = J$, followed by squaring (the term $\pm C_J t^{J+1}$ changes no γ_n with $n \leq J$). The counts themselves are not of that type, and (5) says nothing about $N(Y)$.
- (d) *Analogues.* The brackets (1.10) and (10.8) are, as stated and proved, analogues of `p0:thm:staircase(2)`: two-ceiling brackets proved directly at the integers from smooth envelopes, not deduced from an admissible interpolation of the counts. (Diagnostic: at record values $Y = p_n$, sampled for $200 \leq n < 1500$, $|T_2(r(Y)) - n| < 0.0019$.)

No novelty is claimed for any of these inversions.

11 A rational interior zero and the product obstruction

Set

$$\mathcal{G}(q) = F(q)/q = \sum_{n \geq 0} p_{n+1} q^n. \quad (11.1)$$

This function is holomorphic on $|q| < 1$, with $\mathcal{G}(0) = 1$.

Proposition 11.1 (Exact sign certificate). *The inequalities*

$$\mathcal{G}(-3/4) > 1/4, \quad \mathcal{G}(-4/5) < -1/2 \quad (11.2)$$

hold. Consequently $\mathcal{G}$ has a real zero in $(-4/5, -3/4)$ and is not a finite eta quotient after removal of a monomial factor.

Proof. The two finite inequalities below are verified by exact rational arithmetic in the accompanying standard-library certificate. We first prove the uniform error bound that makes them statements about the infinite series.

Let $P(R) = \prod_{j \geq 1} (1 - R^j)^{-1}$ for $0 < R < 1$. Expanding the logarithm gives

$$\log P(R) = \sum_{k \geq 1} \frac{R^k}{k(1 - R^k)} \leq -\frac{\log(1 - R)}{1 - R}. \quad (11.3)$$

For $R = (b - 1)/b$ this yields $P(R) \leq b^b$. Let $T_N(q) = \sum_{n=0}^N p_{n+1}q^n$ and $r = |q| < R$. Since $p_{n+1} \leq p(n + 1)$,

$$\begin{aligned} |\mathcal{G}(q) - T_N(q)| &\leq (r/R)^{N+1} \sum_{n > N} p(n + 1)R^n \\ &\leq (r/R)^{N+1} \frac{P(R)}{R} \leq (r/R)^{N+1} \frac{b^b}{R}. \end{aligned} \quad (11.4)$$

For $N = 1000$, use $(r, b) = (3/4, 5)$ and $(4/5, 6)$. The exact rational quantity on the right is less than 10^{-12} in both cases. Horner evaluation with rational numbers, using coefficients regenerated from (2.5), verifies

$$T_{1000}(-3/4) - \left(\frac{3/4}{4/5}\right)^{1001} \frac{5^5}{4/5} > \frac{1}{4}, \quad (11.5)$$

and

$$T_{1000}(-4/5) + \left(\frac{4/5}{5/6}\right)^{1001} \frac{6^6}{5/6} < -\frac{1}{2}. \quad (11.6)$$

Equations (11.4)–(11.6) prove (11.2). Continuity gives an interior negative real zero.

A finite eta quotient, after removing its power of q , is a finite product of integral powers of $(q^d; q^d)_\infty$, $d \geq 1$, multiplied by a nonzero constant. Each factor is holomorphic and nonvanishing in $|q| < 1$: its logarithmic product converges normally on compact subdisks. Negative integral powers remain holomorphic and nonvanishing there. The product therefore has no interior zero, contradicting (11.2). $\square$

This is a finite-computation-assisted exact certificate, not a numerical root estimate. It excludes finite eta quotients and, more generally, any proposed representation known to be holomorphic and nonvanishing throughout the disk. It does not exclude sums or differences of eta quotients, a numerator with zeros, or other basic-hypergeometric identities. Failure to recognize an alternative identity is not a theorem of nonexistence.

12 Reproducibility and further questions

The companion archive contains the editable L^AT_EX manuscript, authored verification programs, observed finite reference data, and build instructions. Its mandatory verification path uses only the Python standard library for exact arithmetic and replay; producing the PDF additionally requires a `pdflatex` installation and the listed TeX packages. The full exact coefficient list through 1500 is regenerated rather than accepted as evidence merely because a saved file exists.

The mandatory checks are:

1. regenerate the positive dynamic program and compare the observed 58-term OEIS prefix;
2. independently enumerate ordinary partitions through 35 and test the defining condition;
3. evaluate the rational sign proof (8.10)–(8.11);

4. recalculate the two rational Horner certificates and their proven tails;
5. replay the checks in normal and optimized Python, requiring identical exact outputs;
6. verify the explicit archive inventory and deterministic same-stack rebuilds.

All public guards use explicit exceptions so that they remain active under Python's `-O` option. The optional SymPy verification differentiates the elementary formulas in (8.5) and performs exact Wick contractions. One program checks the first correction in two and three dimensions; another checks the explicit second coefficient. SymPy is an additional dependency and is used only when requested. These are fixed-order verifiers, not an unrestricted all-orders engine. Saved receipts are not substituted for running the computations. The analytic proofs of the Fourier tails and asymptotic remainders are in the article, not delegated to finite coefficient agreement.

Determinism is a same-stack statement: fixed source files, Python, TeX engine, packages, and fonts give identical rebuilt bytes under the documented build. It is not a claim that unrelated TeX installations produce identical PDFs. The build uses a fresh absolute output path and refuses unsafe paths or existing destinations; the README gives the exact commands. The archive excludes downloaded literature and nonpublic review material.

12.1 Questions suggested by the analysis

The following are extensions rather than results proved here.

1. Can one derive explicit c_j for $j \geq 3$ in a compact recurrence, and give effective remainder constants or a computable monotonicity onset? The finite-jet construction proves existence at every fixed order but does not provide those quantitative error bounds.
2. What are the next centering and variance corrections for the least part? A marked three-variable expansion should contain them, but Theorem 9.1 only gives the stated first-order centering and bounded-window local law.
3. Can one obtain relative moderate-deviation estimates for the least part, beyond fixed standardized windows, with explicit ranges? This requires additional uniformity beyond the present Gaussian-tail domination.
4. Does the diagonal generating function admit a useful nonproduct basic-hypergeometric identity? Proposition 11.1 rules out finite eta quotients but leaves many identities possible.
5. How do the constants and phases change for constraints such as $\min \lambda = r |\text{supp } \lambda| + s$, or for bounded multiplicities? Each variant needs its own exact model and global saddle check.
6. How much of the moving-cutoff analysis can be placed directly under general partition local-limit theorems, in particular the full hypotheses of Hwang [6]? Resolving that applicability question would clarify the relation with existing frameworks.

[write] 7 October 2026. Under Vladimir's standing rule of 4 October 2026 (unproved claims are moved to questions, wrong ones are refuted on the record), the write found no claim of the source that is unproved as stated and none that is wrong, so nothing was moved here or refuted. Two finite observations bear on the questions without settling them: for Question 1, the exact counts increase strictly for $33 \leq n < 1500$ (Remark 10.2(a)), which proves no onset; for Question 4, the shipped exact Euler exponents of F through degree 200 show no pure period up to 50, which excludes no identity.

A Finite-jet instructions for higher coefficients

Here is a concrete algebraic implementation of the effectiveness assertion in Theorem 1.1. To determine terms through t^R :

1. Expand $I(v, s)$ about (a, s_0) through total degree $2R + 2$ using $I_v = -h$ and the elementary I_s in (8.5). Expand $\log \mathcal{A}$ through total degree $2R$.
2. Compose $v - a = X + aZ + XZ$, $s - s_0 = U$, and expand $(1 + Z)^{-1}$ to the required degree in (7.4). Keep the Hessian term separate.
3. Include all Euler–Maclaurin terms $-B_{2r} z^{2r-1} h_{v, 2r-1} / (2r)!$ whose substitution can contribute through λ^{2R} . Terms with base exponent $4r - 2 > 2R$ can be discarded.
4. Substitute the formal scaled coordinates $(X, U, Z) = (\lambda\xi, \lambda v, \lambda\zeta)$ in (7.22), keeping (ξ, v, ζ) abstract. Expand through λ^{2R} and divide the amplitude by A_0 . This gives finite coefficient polynomials in the complex-coordinate basis.
5. Contract the coefficient of λ^{2j} with the complex-coordinate covariance (8.2). Equivalently, substitute the plane first and integrate against the normalized Gaussian (7.7); do not apply C directly to a polynomial already expressed in the real plane coordinates. The result is b_j , and $c_j = S^{j/2} b_j$.

To prove a remainder through this order one expands one further odd λ -order, as in Section 7; that odd integral vanishes but allows the $O(t^{R+1})$ bound. The computational jet truncation for the coefficients themselves does not replace that analytic remainder argument.

The entries of H and all required derivatives are exact expressions built from rational operations and the constants in (1.2). At the working point the exponential variables are rational, $e^{-a} = 2/3$ and $e^{s_0} = 1/2$, which keeps finite symbolic calculations small. The package’s optional verifiers implement the $R = 1$ and $R = 2$ calculations directly from these instructions. Coefficients c_j for $j \geq 3$ are prescribed but not explicitly evaluated here. No numerical fitting to the coefficient list is used.

B A normalization check by Gaussian integration

The integral of the leading three-dimensional Gaussian can be checked without determinants of complex matrices. Integrate β in (7.7) first:

$$\int_{\mathbb{R}} e^{-V\beta^2/2 - J_0\alpha\beta} d\beta = \sqrt{\frac{2\pi}{V}} e^{J_0^2\alpha^2/(2V)}.$$

The remaining α exponent is $-W\alpha^2/2 + iak_y\alpha$, so

$$\int_{\mathbb{R}} e^{-W\alpha^2/2 + iak_y\alpha} d\alpha = \sqrt{\frac{2\pi}{W}} e^{-a^2\kappa^2 y^2/(2W)}.$$

Then the y integral has curvature κ_{eff} . Thus

$$\int_{\mathbb{R}^3} e^{Q(w)} dw = \frac{(2\pi)^{3/2}}{\sqrt{VW\kappa_{\text{eff}}}} = \frac{(2\pi)^{3/2}}{\sqrt{2SV\kappa}}. \quad (\text{B.1})$$

Multiplication by $A_0/(2\pi)^2$, the Fourier Jacobian t^2 , and the reciprocal minimum-lattice mesh $t^{-1/2}$ gives exactly

$$\frac{D}{2\sqrt{\pi S}} t^{3/2} e^{2S/t}.$$

This checks the mandatory amplitude, both Fourier normalizations, the lattice spacing, and the conditional covariance correction in one calculation.

References

- [1] OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, A239950, “Number of partitions of n such that the number of different parts is equal to the smallest part.” Entry inspected 4 October 2026. <https://oeis.org/A239950>.
- [2] H. V. Chu and Z. L. Vasseur, *Weighted Schreier-type sets and the Fibonacci sequence*, *Fibonacci Quarterly* **62**(4) (2024), 305–315. <https://www.fq.math.ca/Papers/62-4/chu08302024rev.pdf>.
- [3] K. Beanland and H. V. Chu, *On Schreier-type sets, partitions, and compositions*, *Journal of Integer Sequences* **27** (2024). Primary preprint: <https://arxiv.org/abs/2311.01926>.
- [4] H. V. Chu, Y. Geng, J. King, S. J. Miller, G. Tresch, and Z. L. Vasseur, *Linear recurrences from counting Schreier-type multisets*, *Integers* **26** (2026), A53. <https://math.colgate.edu/~integers/aa53/aa53.pdf>.
- [5] W. M. Y. Goh and E. Schmutz, *The number of distinct part sizes in a random integer partition*, *Journal of Combinatorial Theory, Series A* **69** (1995), 149–158. Publisher abstract: <https://www.sciencedirect.com/science/article/pii/0097316595901116>.
- [6] H.-K. Hwang, *Limit theorems for the number of summands in integer partitions*, *Journal of Combinatorial Theory, Series A* **96** (2001), 89–126. <https://doi.org/10.1006/jcta.2000.3170>.
- [7] H.-K. Hwang and Y.-N. Yeh, *Measures of distinctness for random partitions and compositions of an integer*. Primary author copy: https://mathweb.ucsd.edu/~ebender/comp.papers/102-distinct_parts.pdf.
- [8] P. J. Grabner, A. Knopfmacher, and S. Wagner, *A general asymptotic scheme for the analysis of partition statistics*, *Combinatorics, Probability and Computing* **23** (2014), 1057–1086. <https://doi.org/10.1017/S0963548314000418>. Primary author copy: <https://www.math.tugraz.at/~grabner/Publications/PartitionStatistics.pdf>.
- [9] M. Archibald, A. Blecher, C. Brennan, A. Knopfmacher, and T. Mansour, *Partitions according to multiplicities and part sizes*, *Australasian Journal of Combinatorics* **66**(1) (2016), 104–119. https://ajc.maths.uq.edu.au/pdf/66/ajc_v66_p104.pdf.