Concise axiomatizations of intuitionistic primitive recursive arithmetic

November 2026

Abstract

Intuitionistic first-order primitive recursive arithmetic is bi-interpretable with a calculus that has variables, zero, successor and one iterator as its terms, equations and two binders as its formulas, and eight inference rules. Both numerical domains are whole, equality is actual equality, and both composite maps have graph $x = y$. This article states that calculus and the bridge once, and then collects every refinement of it that the research packages of the `ConciseIPRA` directory of the `ProveIt` repository contain: changes of the rule basis, guarded formula languages, rules that merge computation with equality, normal forms, clause calculi, rules that discharge a witness, introductions that take no witness, other recursors, obstruction results, and proof-certificate formats with their checkers. Each definition, rule inventory, theorem and proof is given once, in one notation. Statements that the packages leave unproved are collected at the end of each chapter.

Contents

Introduction **10**

1 The base theories and their bridge **12**
1.1 Terms and substitution . . . 12
1.2 Native formulas and their ordinary reading . . . 13
1.3 The native scalar core . . . 14
1.4 Structural properties and derived rules . . . 15
1.4.1 Weakening, substitution and cut . . . 15
1.4.2 Closure by cases and numerical cases . . . 16
1.4.3 Equality . . . 17
1.4.4 Congruence of the binders and the Leibniz rule . . . 18
1.4.5 Gates and selectors of formulas . . . 18
1.4.6 The connectives as native formulas . . . 20
1.5 The ordinary theory T_{eq} . . . 21
1.6 Intuitionistic primitive recursive arithmetic . . . 22
1.7 Arithmetic inside T_{eq} . . . 22
1.7.1 The equational fragment . . . 23
1.7.2 Addition . . . 23
1.7.3 Zero test, interpolation and parity . . . 24
1.7.4 A three-cycle and a ramp . . . 25
1.7.5 The factor map and the comparator . . . 26
1.7.6 The saturating counter and the predecessor . . . 27
1.7.7 Successor laws, subtraction and the reflection of equality . . . 28
1.7.8 Induction for quantifier-free formulas . . . 29
1.7.9 Order, powers of two and pairing . . . 30
1.7.10 Primitive recursion . . . 31
1.8 The bridge . . . 32
1.8.1 The native core and T_{eq} . . . 32
1.8.2 Term interpretations . . . 34
1.8.3 From T_{eq} to iPRA . . . 35
1.8.4 From iPRA to T_{eq} . . . 36
1.8.5 The two round trips on terms . . . 37
1.8.6 The bi-interpretation theorem . . . 37
1.9 Clause contexts . . . 39
1.10 The measurement protocol . . . 39
1.11 Further questions . . . 40

2 Refinements of the eight rules **41**
2.1 Conventions and the comparison method . . . 41
2.2 The introduction rules . . . 43
2.2.1 Progress through a witness and progress at the closure . . . 43
2.2.2 One closure introduction with a mode read off the matrix . . . 44
2.2.3 What cannot be removed from the introductions . . . 44
2.2.4 One introduction with a mode read off the matrix and the witness . . . 46
2.3 The transport rule . . . 47
2.3.1 One template, two templates, a formula template . . . 47
2.3.2 Sources other than the literal zero . . . 48
2.3.3 Reversal of the root equation . . . 49
2.3.4 The same restriction for the uncrossed rule . . . 50
2.3.5 Replacement with no template . . . 53
2.3.6 Replacement of one occurrence . . . 54
2.3.7 A restricted inventory of computation instances . . . 55
2.4 Induction and uniqueness of iteration . . . 56
2.4.1 The rule and the two theorems . . . 56
2.4.2 First consequences of uniqueness . . . 57
2.4.3 Retractions and induction for zero equations . . . 58
2.4.4 A predecessor from uniqueness . . . 59
2.4.5 Truncated subtraction and diagonal descent . . . 61
2.4.6 An accumulator and the commutativity of the maximum . . . 62
2.4.7 Reflection of equality and the proofs of the theorems . . . 62
2.4.8 Other forms of the uniqueness rule . . . 63
2.5 The progressive reading of the closure . . . 66
2.5.1 The calculus with orbit elimination . . . 67
2.5.2 Derived rules of the orbit calculus . . . 68
2.5.3 Soundness for the progressive reading . . . 69
2.5.4 The embedding of the core . . . 69
2.5.5 The reading inside the core and the two round trips . . . 70
2.5.6 Transport in a formula template . . . 72
2.5.7 Elimination at a target . . . 73
2.5.8 The calculi of the chapter . . . 74
2.6 Two results on other recursors . . . 75
2.6.1 Counter congruence at a zero seed . . . 75
2.6.2 Recursors seeded at zero . . . 77
2.7 Checker sizes . . . 78
2.8 Further questions . . . 82

3 Guards and public languages **86**
3.1 Guards as literal shapes . . . 87
3.1.1 The root-determined introduction . . . 87
3.1.2 Structural operations, the kernel, and equality at admissible sources . . . 88
3.1.3 Deleting the choice is not conservative . . . 89
3.1.4 The shapes . . . 90
3.1.5 Public languages . . . 92
3.2 The guarded bootstrap and the translation into a public language . . . 92
3.2.1 Guard schemes . . . 93
3.2.2 The translation and its two composites . . . 95
3.2.3 The guarded core . . . 97
3.3 Which transport sources are needed . . . 98

3.4 The failure of composition and its repairs . . . 100
3.4.1 The obstruction . . . 101
3.4.2 First repair: auxiliary formulas . . . 102
3.4.3 Second repair: the administrative orbit . . . 103
3.4.4 Third repair: the mirror guard . . . 104
3.4.5 Fourth repair: protected atoms . . . 105
3.5 Guards on the full native language . . . 107
3.5.1 Selectors and the criterion . . . 107
3.5.2 The diagonal guard . . . 109
3.5.3 The mirror selector and six rules . . . 110
3.6 Comparison of the guards . . . 111
3.7 Checkers and their sizes . . . 112
3.8 Further questions . . . 114

4 Computation, equality and normal forms 117
4.1 Conventions and shared lemmas . . . 117
4.1.1 Notation . . . 117
4.1.2 Equality from reflexivity and crossed transport . . . 118
4.1.3 The single introduction . . . 119
4.1.4 Congruence below binders and provability of conversion . . . 120
4.2 The rule Eval on raw terms . . . 120
4.2.1 The operation Φ and the calculi . . . 121
4.2.2 The identity frame . . . 121
4.2.3 Computation, equality and transport recovered . . . 122
4.2.4 Equivalence with the core . . . 123
4.2.5 Substitution . . . 124
4.2.6 Irredundancy . . . 124
4.3 Normalizers and the transfer of derivations . . . 125
4.3.1 Admissible normalizers and normalizing substitution . . . 125
4.3.2 The kernel on normal forms . . . 126
4.4 Zero-normal terms . . . 128
4.4.1 Zero normalization . . . 128
4.4.2 The kernel without the zero computation rule . . . 129
4.4.3 Six rules with the single introduction . . . 130
4.5 The introduction selected by a reflexive base . . . 131
4.5.1 The calculus . . . 131
4.5.2 The structural rules, without circularity . . . 131
4.5.3 Progress recovered by two decoders . . . 132
4.5.4 Equivalence . . . 133
4.5.5 The same guard on raw terms, and two variants . . . 134
4.6 Equality as replacement of an initializer . . . 135
4.6.1 The rule Seed and the calculus . . . 135
4.6.2 Equality recovered . . . 136
4.6.3 Equivalence . . . 137
4.7 One successor step . . . 137
4.7.1 The rule Step . . . 137
4.7.2 Five patterns with clause contexts . . . 139
4.8 Seed evaluation and suspended application . . . 140
4.8.1 The rule SeedEval . . . 140
4.8.2 Suspended application . . . 141
4.9 Full normalization . . . 145

4.9.1 Termination and unique normal forms . . . 145
4.9.2 The kernel without computation rules . . . 147
4.9.3 Six rules on full normal forms . . . 147
4.9.4 What normalization costs . . . 149
4.10 A smaller checker for the unchanged core . . . 150
4.11 Checker sizes . . . 150
4.12 Further questions . . . 154
4.12.1 What is not achieved . . . 154
4.12.2 Claims stated without proof . . . 155
4.12.3 Open questions . . . 155

5 Clause calculi **157**
5.1 From Q formulas to clauses . . . 157
5.1.1 The Q-free form of a formula . . . 157
5.1.2 Clause occurrences and their origins . . . 158
5.2 The five-rule calculus . . . 158
5.2.1 Rules . . . 159
5.2.2 Derived rules . . . 159
5.2.3 Equivalence with the native scalar core . . . 161
5.3 Variants of the arithmetic interface . . . 162
5.3.1 Two evaluation cases . . . 162
5.3.2 Zero-normal terms and one evaluation scheme . . . 163
5.3.3 Reflexivity as one fixed clause . . . 166
5.3.4 Four schemes on zero-normal terms . . . 168
5.4 Variants of the introduction . . . 168
5.4.1 The diagonal guard . . . 168
5.4.2 Guards with a derivable base . . . 170
5.4.3 The fixed zero base on zero-normal terms . . . 171
5.5 Variants of the context . . . 172
5.5.1 The prefix representation . . . 172
5.5.2 Rank from variable support . . . 174
5.5.3 One body per context entry . . . 175
5.5.4 Homogeneous contexts with an asymmetric gate . . . 177
5.6 Fusion: evaluation at the discharge . . . 178
5.6.1 The raw calculus . . . 179
5.6.2 The zero-normal companion . . . 180
5.7 The ordinary realization . . . 181
5.8 Successor rigidity . . . 181
5.9 Checker sizes . . . 182
5.10 Further questions . . . 185

6 Discharge of a witness and restricted equality rules **187**
6.1 Normal forms, clause contexts, and the reference calculus . . . 187
6.1.1 Conventions on full normal forms . . . 187
6.1.2 Formulas, clauses, and the two conventions . . . 189
6.1.3 The shared rules and the reference calculus . . . 189
6.1.4 Structural operations . . . 190
6.1.5 Closures, numerical cases, and the two introductions . . . 191
6.1.6 Equality from one term transport . . . 192
6.1.7 Two-row packets . . . 192
6.1.8 Interchangeability of the two conventions . . . 193

6.2 Naming packets: equality at the discharge of a witness . . . 193
6.2.1 The rule NAME . . . 193
6.2.2 Transport from one discharge . . . 194
6.2.3 Equivalence . . . 194
6.3 The inferred witness . . . 195
6.3.1 The selector and the two forms of the rule . . . 195
6.3.2 The export form . . . 197
6.3.3 The open form . . . 197
6.3.4 Summary and comparison of the four packages . . . 199
6.4 Substitutive discharge on the canonical recursor syntax . . . 200
6.4.1 Syntax . . . 200
6.4.2 Four rule families . . . 201
6.4.3 The bootstrap of equality . . . 202
6.4.4 The canonical basis and the bridge . . . 202
6.5 Restricted substitution . . . 204
6.5.1 Substitution for a variable . . . 204
6.5.2 The newest variable . . . 205
6.5.3 Substitution for a parameter on raw terms . . . 207
6.5.4 Unary congruence does not suffice . . . 210
6.6 Specialization without a term . . . 212
6.6.1 The calculus . . . 212
6.6.2 Naming a term . . . 213
6.6.3 Persistent proofs . . . 214
6.6.4 Equivalence . . . 215
6.7 A constancy rule in place of induction . . . 216
6.7.1 The theory . . . 216
6.7.2 Recovery of quantifier-free induction . . . 217
6.7.3 Both interpretations . . . 218
6.8 Checkers and program equivalences . . . 219
6.9 Checker sizes . . . 221
6.10 Further questions . . . 222

7 Introduction without a supplied witness 225
7.1 Restricted witnesses . . . 225
7.1.1 Scopes and the restricted rules . . . 225
7.1.2 What a restricted witness loses . . . 226
7.2 Obstructions . . . 227
7.2.1 Each shape alone is incomplete . . . 227
7.2.2 The witness and the target cannot both be restricted . . . 228
7.3 Names . . . 230
7.3.1 Naming matrices . . . 230
7.3.2 Derived rules with variable witnesses . . . 230
7.3.3 The introduction at a name . . . 231
7.4 The coverage packet . . . 232
7.4.1 The predecessor . . . 232
7.4.2 The packet and the names . . . 232
7.4.3 A mode selected by the witness . . . 234
7.5 The newest parameter . . . 234
7.5.1 Stable derivations and hereditary families . . . 235
7.5.2 Closure by cases, the Leibniz rule and equation cases . . . 236
7.5.3 The packet in every scope . . . 237

7.5.4 The equivalence theorem . . . 237
7.5.5 The seven-rule diagonal variant . . . 239
7.5.6 One bit . . . 240
7.5.7 The five-family clause calculus . . . 241
7.6 The raw guarded calculus . . . 242
7.6.1 The calculus and its three introductions . . . 243
7.6.2 The naming exception . . . 243
7.6.3 Uniform introduction: the stable fragment . . . 245
7.6.4 Uniform introduction: names and completeness . . . 247
7.7 A recursor whose predecessor computes . . . 248
7.7.1 Terms in normal form . . . 248
7.7.2 Three rule families . . . 249
7.7.3 Constructions that precede equality . . . 250
7.7.4 Completeness . . . 251
7.7.5 The ordinary theory and the bridge . . . 252
7.7.6 Checking a template through its two instances . . . 253
7.8 Conditional copy . . . 254
7.8.1 Use, Close and Export . . . 254
7.8.2 Preparations without equality . . . 255
7.8.3 The conditional-copy theorem . . . 255
7.8.4 Removal of the witness . . . 256
7.8.5 The liberal calculus presents the ordinary theory . . . 257
7.8.6 Quantifier-free induction and the bridge . . . 258
7.8.7 Discharge by a supplied name . . . 259
7.9 Checkers and sizes . . . 260
7.10 Further questions . . . 263

8 Recursors with a built-in seed 265
8.1 The successor-seeded recursor . . . 266
8.1.1 Terms, equations and the two theories . . . 266
8.1.2 Equality, cases and induction before the predecessor . . . 267
8.1.3 Boolean terms and contextual case analysis . . . 268
8.1.4 The predecessor . . . 269
8.1.5 Recovery of the iterator . . . 273
8.1.6 Comparison of the two term languages . . . 273
8.2 Tied-seed recursors . . . 275
8.2.1 The recursor and its elementary terms . . . 275
8.2.2 The one-coordinate bound . . . 276
8.2.3 The theory with a Boolean equality test . . . 277
8.2.4 A compiler with caps . . . 280
8.2.5 The functions denoted by terms . . . 283
8.2.6 Truncated subtraction in place of the equality test . . . 284
8.2.7 The native calculus with truncated subtraction . . . 285
8.3 The self-seeded recursor . . . 287
8.3.1 The theories . . . 287
8.3.2 Without injectivity of the successor . . . 288
8.3.3 With injectivity: predecessor and order . . . 291
8.3.4 Bounded iteration of finite-case functions . . . 292
8.3.5 Arithmetic by finite-case graphs . . . 293
8.4 Three observations on the native core . . . 296
8.4.1 One native binder over a literal equation . . . 296

8.4.2 The case rule at a literal equation cannot be deleted . . . 297
8.4.3 Transport from the literal zero alone . . . 297
8.5 Checker sizes . . . 298
8.6 Further questions . . . 300

9 Canonical and counter-aware recursors 302
9.1 Term languages and normal forms . . . 302
9.1.1 Three recursors without a successor constructor . . . 302
9.1.2 Zero-body canonical terms . . . 303
9.1.3 Full normal forms . . . 304
9.1.4 The status of the successor and of each computation . . . 306
9.2 What the counter gives . . . 306
9.2.1 A decoder that computes . . . 306
9.2.2 Primitive recursion without pairing . . . 307
9.2.3 The register iterator . . . 308
9.2.4 A register with one bound variable . . . 310
9.2.5 Without a counter . . . 311
9.3 Translations to iterator terms and the transfer of the bridge . . . 311
9.3.1 The ordinary theories . . . 311
9.3.2 The translations . . . 312
9.4 The seeded recursor: unrestricted calculi and six rules . . . 315
9.4.1 The raw and the canonical unrestricted calculus . . . 315
9.4.2 The six rules . . . 317
9.4.3 The bridge for the seeded calculi . . . 319
9.5 Five rules with clause contexts . . . 320
9.5.1 The calculus . . . 320
9.5.2 Closure as an abbreviation . . . 321
9.5.3 Comparison with the six rules . . . 322
9.6 Counter evaluation . . . 323
9.6.1 The clause syntax and a liberal calculus . . . 323
9.6.2 The five patterns of counter evaluation . . . 324
9.6.3 Ordinary logic with the decoder . . . 326
9.6.4 Adequacy and the main theorem . . . 327
9.7 Three families on normal forms . . . 328
9.7.1 Use, Close and Name . . . 329
9.7.2 Use, Introduce and Open with an inferred witness . . . 330
9.7.3 The scalar normal forms embed exactly . . . 332
9.7.4 The main theorem for the two calculi . . . 333
9.8 Uniform binding and one traversal . . . 333
9.8.1 The concrete syntax . . . 334
9.8.2 One traversal . . . 335
9.8.3 The checkers . . . 336
9.8.4 Three refinements of checkers for the kernel on iterator terms . . . 337
9.9 Boundary results . . . 337
9.9.1 A recursor with neither counter nor initializer . . . 338
9.9.2 The two computation equations of the iterator are independent . . . 339
9.9.3 Induction on zero equations does not give induction on equations . . . 340
9.9.4 The last computation rule . . . 342
9.10 Checker sizes . . . 342
9.11 Further questions . . . 344

10 Obstructions **347**
10.1 Conventions and the presentations under test . . . 348
10.1.1 Three kinds of exclusion . . . 348
10.1.2 The paired-introduction form of the native core . . . 349
10.1.3 Reduced term languages . . . 349
10.1.4 The progressive calculus . . . 351
10.2 Coordinatewise powers and truth transfer . . . 351
10.2.1 The powers . . . 351
10.2.2 Equation induction does not replace numerical cases . . . 354
10.2.3 Boolean cases are not admissible in the progressive calculus . . . 355
10.3 Symmetry . . . 355
10.4 Majorants: almost unary term languages . . . 357
10.4.1 Zero-seeded recursion is monotone . . . 358
10.4.2 Bounding coordinates . . . 358
10.4.3 Single-body constructors . . . 358
10.4.4 Adaptive query circuits . . . 359
10.5 Finite-dimensional term interpretations . . . 361
10.6 Finite profiles and relational bi-interpretations . . . 363
10.6.1 Finite control . . . 363
10.6.2 An existential-free presentation . . . 365
10.6.3 No compression of scalars . . . 366
10.6.4 No finite-dimensional bi-interpretation . . . 367
10.6.5 The selector boundary . . . 368
10.7 Zero profiles of positive iteration . . . 368
10.8 Rule-level obstructions . . . 369
10.8.1 Atomic case introduction cannot be made automatic . . . 369
10.8.2 Elimination majors must be proved, not only retained . . . 370
10.8.3 Root replacement does not give congruence . . . 370
10.8.4 The witness order of the paired binder . . . 371
10.8.5 Two side conditions that cannot be dropped . . . 371
10.9 Independence of the progress rule . . . 372
10.9.1 Coherent iterator structures . . . 372
10.9.2 Native terms in the tower . . . 374
10.9.3 The theorem . . . 375
10.9.4 No successor cycle without progress . . . 376
10.10 The progressive binder presentation . . . 378
10.11 The counter-congruence form of the calculus . . . 381
10.12 Constructed sources and certificate interfaces . . . 383
10.12.1 Successor sources constructed from predecessors . . . 383
10.12.2 Statements about the checkers of the packages . . . 385
10.13 Two partial routes . . . 386
10.13.1 From uniqueness of iteration toward induction . . . 387
10.13.2 What fixed countdown uniqueness determines . . . 388
10.14 Checker sizes . . . 388
10.15 Further questions . . . 390

11 Certificates and checkers **393**
11.1 The calculi that are checked . . . 394
11.2 The sorted arity encoding . . . 395
11.3 The instruction format and the certified zero . . . 398
11.3.1 The base protocol . . . 398

11.3.2 Restricted computation and the certified zero . . . 399
11.3.3 Correspondence with the calculus . . . 400
11.4 Uniform zero and constructed majors . . . 403
11.4.1 The recoding . . . 403
11.4.2 Constructed majors and local floors . . . 404
11.4.3 Sizes of certificates . . . 405
11.5 Neutral major premises . . . 406
11.5.1 The protocol and its hazard . . . 406
11.5.2 The invariant of successful runs . . . 406
11.5.3 Correspondence . . . 407
11.5.4 Composition with local floors . . . 408
11.6 Fixing an elimination before its major is proved . . . 409
11.6.1 The global-floor version . . . 409
11.6.2 The local-floor version . . . 410
11.7 Scalar neutral premises and loops . . . 411
11.7.1 The guarded-source protocol . . . 411
11.7.2 Loops over immutable contexts . . . 413
11.7.3 The root test is a logical boundary . . . 413
11.7.4 Local rewritings of the program text . . . 414
11.8 Validated presentations . . . 414
11.8.1 Input contracts and the public grammar . . . 414
11.8.2 Scope is checked by the first opening . . . 415
11.8.3 Typed checkers . . . 416
11.8.4 Goal-validated checkers . . . 417
11.8.5 A rejected saving . . . 419
11.9 Inferred majors, eager selection and the phase of validation . . . 419
11.9.1 The controls . . . 419
11.9.2 The inferred major . . . 419
11.9.3 Eager selection of the goal . . . 421
11.9.4 The remainder selector . . . 422
11.9.5 Truth-valued results and the tail of a packet . . . 422
11.9.6 The phase of validation . . . 423
11.9.7 A typed native checker and typed guarded checkers . . . 424
11.9.8 The audit of sources . . . 425
11.10 Results on designs that were examined . . . 425
11.10.1 Projection boxes lose reflexivity . . . 426
11.10.2 Eager initialization has a two-element model . . . 426
11.10.3 Positive iteration in the standard model . . . 427
11.10.4 Conditional and progressive closure of an equation . . . 429
11.10.5 Further rejected designs . . . 430
11.11 Checker sizes . . . 430
11.11.1 The coordinatewise envelope . . . 434
11.12 Further questions . . . 435

A The packages **438**

Introduction

The question

The question behind all of this work is how small a formal system can be while it still is, in the strongest reasonable sense, intuitionistic primitive recursive arithmetic iPRA. "Is" means a bi-interpretation in which the numerical domain of each interpretation is the whole domain, equality is interpreted by equality, and each composite of the two numerical maps is the identity, provably. "Small" has two readings. The mathematical one counts term constructors, formula constructors and inference rules, and asks which of them can be removed, restricted or fused. The empirical one fixes a measurement protocol for complete proof checkers and compares their sizes (Section 1.10). The two readings do not always agree, and the text keeps them apart: a smaller rule basis can need a larger checker, and several of the smallest checkers implement an unchanged rule basis.

How the article is organized

Section 1.1 to Section 1.10 of the first chapter state the common core: the terms, the native formulas and their ordinary reading, the eight rules of the native scalar core N, its structural properties and derived connectives, the ordinary theory T_{eq}, the arithmetic inside T_{eq}, and the bridge to iPRA. No other chapter restates this material.

The next six chapters stay on iterator terms. The second treats refinements of the eight rules on the unchanged language: the introductions, the transport rule, uniqueness of iteration in place of induction, and the progressive reading of the closure binder. The third treats guards: formula languages in which the shape of a matrix determines the mode of its introduction. The fourth treats rules that merge computation with equality and calculi on normal forms. The fifth treats clause calculi, in which contexts hold reusable clauses and the closure binder is not a formula. The sixth treats rules that fuse equality reasoning with the discharge of an opened witness, and restricted substitution rules. The seventh treats introductions that take no witness.

Two chapters change the term language: recursors with a built-in seed, and canonical and counter-aware recursors in which the successor is defined. The tenth chapter collects the obstruction and independence results, with the models and invariants that prove them. The last chapter treats proof-certificate formats and checkers, and records the measured sizes.

Each chapter ends with a table of the checker sizes that its packages report and with a section of further questions: claims stated without proof, conjectures, and what was tried and not achieved.

Conventions

The notation of the first chapter is used throughout. Names of calculi other than N, T_{eq} and iPRA are local to the chapter that introduces them; the same letter can denote different calculi

in different chapters. Every chapter compares its calculi with the eight-rule core of the first chapter. A research package is cited by its directory name, as in `witness_rule`; Chapter A lists every package with the chapters that draw on it. Where a result is formalized in Lean in the repository, the name of the Lean declaration is given in typewriter type; the status of the formalization is recorded in the README of the directory and is not repeated here in full. Editorial repairs of the source reports are stated where they occur.

Chapter 1

The base theories and their bridge

Three formal systems recur in every chapter of the article: the native scalar core N, a calculus with eight rules and no connectives; the ordinary theory T_{eq}, intuitionistic logic with equality over the same terms; and iPRA, intuitionistic first-order primitive recursive arithmetic with quantifier-free induction. This chapter states the three systems, develops the derived rules of N and the arithmetic of T_{eq}, and proves that the three systems are bi-interpretable with whole domains, actual equality and identity composites (Theorem 1.78). It also fixes the notion of a clause context and the protocol by which proof checkers are measured. The later chapters use these statements without repeating them.

Notation that is fixed here and used throughout: $1 := S0$, $2 := SS0$, $3 := SSS0$; a term or formula written $h(v)$ has a distinguished variable v, and $h(s)$ abbreviates $h[s/v]$; $A \dashv\vdash B$ means that each of A, B is derivable from the other in the system under discussion.

1.1 Terms and substitution

This section draws on the reference statement of the inherited basis filed with `certified_zero` and on the syntax section of `raw_clausal`.

Definition 1.1 (Terms)**.** There is one numerical sort. The terms are

$$t ::= x \mid 0 \mid St \mid I(n, a;\, u.\, h),$$

where x ranges over a countable set of variables. In the *iterator* $I(n, a;\, u.\, h)$ the term n is the *counter*, a is the *seed* and h is the *body*. The iterator binds the variable u in h and nowhere else; u is not bound in n or in a. The body may contain other free variables, which act as fixed numerical parameters. Terms that differ only in the names of bound variables are identified. $\mathrm{FV}(t)$ is the set of free variables of t.

A *substitution* θ is a map from variables to terms that is the identity outside a finite set. The term $t[\theta]$ is defined by

$$x[\theta] = \theta(x), \qquad 0[\theta] = 0, \qquad (St)[\theta] = S(t[\theta]), \qquad I(n, a;\, u.\, h)[\theta] = I(n[\theta], a[\theta];\, u.\, h[\theta]),$$

where in the last clause the bound variable u is first renamed so that $\theta(u) = u$ and $u \notin \mathrm{FV}(\theta(x))$ for every $x \neq u$. The substitution that sends x to s and fixes all other variables is written $[s/x]$. Substitution is capture-free by construction, and the following standard facts hold by induction on t.

Lemma 1.2 (Substitution)**.** (i) *If* $x \notin \mathrm{FV}(t)$ *then* $t[s/x] = t$.

(ii) $t[\theta][\theta'] = t[\theta'']$, *where* $\theta''(x) = \theta(x)[\theta']$.

(iii) *If* $x \neq z$ *and* $x \notin \mathrm{FV}(r)$*, then* $t[s/x][r/z] = t[r/z][s[r/z]/x]$.

Definition 1.3 (Selector, zero test, parity)**.** For terms c, a, b put

$$\mathrm{sel}(c,a,b) := I(c,a;\, u.\, b) \quad (u \notin \mathrm{FV}(b)), \qquad Z(c) := \mathrm{sel}(c,1,0), \qquad N(n) := I(n,0;\, u.\, Z(u)).$$

The italic letter N names the parity term; the calculus is written with the sans-serif letter N.

The two *computation equations* of the iterator are

$$I(0,a;\, u.\, h) = a, \tag{1.1}$$
$$I(Sn,a;\, u.\, h) = h[I(n,a;\, u.\, h)/u]. \tag{1.2}$$

They are equations between distinct terms, to be derived or assumed in a calculus; no term is identified with its value. For the selector they read $\mathrm{sel}(0,a,b) = a$ and $\mathrm{sel}(Sc,a,b) = b$, because the bound variable does not occur in the body.

Remark 1.4 (What the language does not contain)*.* Every term is finite syntax. No numerical variable ranges over bodies or over descriptions of functions, and there is no term that evaluates descriptions uniformly. The body of an iterator receives the previous value only; it does not receive the counter. The reference statements present the same language with function symbols instead of a binder: at stage zero the symbols are 0 and S, and for every earlier-stage term $h(u,\vec{p})$ a symbol $i_h(n,a,\vec{p})$ is added with the equations $i_h(0,a,\vec{p}) = a$ and $i_h(Sn,a,\vec{p}) = h(i_h(n,a,\vec{p}),\vec{p})$. The formal parameters of a symbol are distinct and its actual arguments are arbitrary. The two presentations have the same terms up to this renaming, and the article uses the binder.

1.2 Native formulas and their ordinary reading

This section draws on the reference statements filed with `certified_zero` (the inherited basis and the inherited ordinary bridge).

Definition 1.5 (Native formulas)**.** The native formulas are

$$A ::= (s = t) \mid Q_y A \mid D_{x,y} A.$$

Q_y binds y; $D_{x,y}$ binds the two distinct variables x and y. Formulas that differ only in the names of bound variables are identified, and $A[\theta]$ is capture-free substitution. A *literal equation* is a formula of the form $s = t$; an equation below a binder is not a literal equation. The designated contradiction is the equation $0 = S0$.

There are no connectives, no falsity constant and no ordinary quantifier in the native language.

Definition 1.6 (Ordinary formulas and the ordinary reading)**.** The *ordinary formulas* are the first-order formulas over the terms of Theorem 1.1, built from equations by $\bot$, $\wedge$, $\vee$, $\to$, $\forall$ and $\exists$. The *ordinary reading* A° of a native formula A is the ordinary formula defined by

$$\begin{aligned} (s = t)^\circ &:= (s = t), \\ (Q_y B)^\circ &:= B^\circ(0) \to \forall y\, B^\circ(y), \\ (D_{x,y} B)^\circ &:= \exists x \left(B^\circ(x,0) \to \forall y\, B^\circ(x,y)\right). \end{aligned}$$

For a list Γ of native formulas, Γ° is the list of the readings of its members.

Thus Q_yB says that B holds everywhere provided it holds at zero, and $D_{x,y}B$ says that some x has this property. The reading uses the whole numerical domain and reads the equality sign as equality. By induction on A,

$$(A[\theta])^\circ = A^\circ[\theta]. \tag{1.3}$$

In the Lean formalization the native formulas are `NFm` and the reading is `NFm.toFm`.

1.3 The native scalar core

This section draws on the reference statement of the inherited basis filed with `certified_zero`; the two rules Tr and QI are taken in the forms of `one_context` and `direct_closure`, as in the Lean kernel.

A *sequent* is $\Gamma \vdash A$, where the *context* Γ is a finite list of native formulas and A is a native formula.

Definition 1.7 (The native scalar core N)**.** The derivable sequents of N are generated by the following eight rules. In Tr, $h(v)$ is a term with a distinguished variable v.

$$\frac{A \in \Gamma}{\Gamma \vdash A}\ \textsc{Hyp} \qquad \frac{}{\Gamma \vdash I(0,a;\,u.\,h) = a}\ \mathrm{I0}$$

$$\frac{}{\Gamma \vdash I(Sn,a;\,u.\,h) = h[I(n,a;\,u.\,h)/u]}\ \mathrm{IS}$$

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash h(s) = r}{\Gamma \vdash r = h(t)}\ \textsc{Tr} \qquad \frac{\Gamma \vdash Q_yB(y) \qquad \Gamma \vdash B(0)}{\Gamma \vdash B(t)}\ \mathrm{QE}$$

$$\frac{\Gamma \vdash D_{x,y}B(x,y) \qquad \Gamma,\ Q_yB(b,y) \vdash A}{\Gamma \vdash A}\ \mathrm{DE}$$

$$\frac{\Gamma,\ B(t,0) \vdash B(t,Sz)}{\Gamma \vdash D_{x,y}B(x,y)}\ \mathrm{DI} \qquad \frac{\Gamma,\ E(z) \vdash E(Sz)}{\Gamma \vdash Q_yE(y)}\ \mathrm{QI}$$

Side conditions. In DE the variable b is not free in Γ, in A or in $D_{x,y}B$. In DI the variable z is not free in Γ, in t or in $D_{x,y}B$; the term t is the *witness*. In QI the variable z is not free in Γ or in Q_yE, and E is a literal equation. The displayed assumptions $Q_yB(b,y)$, $B(t,0)$ and $E(z)$ are discharged.

The rules are named as follows: hypothesis; the two computation equations; transport in one term; elimination of Q; elimination of D; introduction of D *by cases*; introduction of Q *by progress*. The matrix of DI is an arbitrary native formula, and its premise has one assumption, the instance at zero. The matrix of QI is a literal equation, and its premise has one assumption, the instance at the eigenvariable.

Remark 1.8 (The complete inventory). The eight rules are all there is. There is no reflexivity axiom and no other equality rule; no axiom states that a number is zero or a successor, that the successor is injective or that $Sx \neq 0$; there is no induction rule other than QI, whose matrix is one equation; there is no logical rule for a connective, since there is no connective; and there are no structural rules. Weakening, substitution and cut are admissible (Theorem 1.10). Every one of the omitted principles is derived in Section 1.4 or, through the bridge, in Section 1.7. In Lean the calculus is the inductive predicate `NPrf`, with de Bruijn indices: a fresh variable is the new index 0 over a context whose formulas are shifted.

Remark 1.9 (The calculus as the reports state it). The research packages state the calculus with two rules in a stronger form: transport in an arbitrary native formula template,

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash F(s)}{\Gamma \vdash F(t)},$$

and a witnessed introduction by progress, which infers $D_{x,y}E(x,y)$ from $\Gamma, E(t,z) \vdash E(t,Sz)$ for a literal equation E. Both are derived rules of N (Theorems 1.16 and 1.23). Conversely, in the calculus of the reports reflexivity follows by transporting $I(0,t;\, u.\, 0) = t$ along itself in the template $v = t$, symmetry and transitivity follow by transport in the templates $v = s$ and $s = v$, the rule Tr follows from the transport of $r = h(s)$ in the template $r = h(v)$, and QI follows by introducing $D_{x,y}E(y)$ with x not free in E and opening it by DE, which yields the hypothesis $Q_yE(y)$ itself. The two calculi therefore derive the same sequents (Lean: `iprf_iff_nprf`). The reports also write the calculus with clause contexts instead of the binder Q; that form is described in Section 1.9. The comparison of rule bases is the subject of later chapters and is not pursued here.

1.4 Structural properties and derived rules

This section draws on the reference statement of the inherited basis filed with `certified_zero` for the logical interface, and follows the Lean modules `NativeStruct`, `NativeEq`, `NativeGate` and `NativeLogic` in the organization of the proofs. Throughout the section $\vdash$ is derivability in N.

1.4.1 Weakening, substitution and cut

Lemma 1.10 (Weakening and substitution). *If $\Gamma \vdash A$, then $\Delta \vdash A[\theta]$ for every substitution θ and every context Δ that contains $B[\theta]$ for each $B \in \Gamma$. In particular $\Gamma \vdash A$ implies $\Gamma[\theta] \vdash A[\theta]$, and $\Gamma \vdash A$ with $\Gamma \subseteq \Delta$ implies $\Delta \vdash A$.*

Proof. By induction on the derivation of $\Gamma \vdash A$, for all θ and Δ at once.

Hyp: $A \in \Gamma$ gives $A[\theta] \in \Delta$.

I0, IS: an instance of a computation equation under θ is again an instance, for (1.2) by Theorem 1.2: $(h[I(n,a;\, u.\, h)/u])[\theta] = h[\theta][I(n[\theta], a[\theta];\, u.\, h[\theta])/u]$.

Tr: choose the template variable v fixed by θ and not free in its values. Then $h(s)[\theta] = h[\theta](s[\theta])$ and $h(t)[\theta] = h[\theta](t[\theta])$, so the two premises given by the induction hypothesis are premises of Tr with the template $h[\theta]$.

QE: $B(t)[\theta] = B[\theta](t[\theta])$ and $B(0)[\theta] = B[\theta](0)$, and the induction hypothesis gives $\Delta \vdash Q_yB[\theta]$ and $\Delta \vdash B[\theta](0)$.

DE: choose b' not free in Δ, $A[\theta]$, $D_{x,y}B[\theta]$ and the values of θ, and let θ' be θ changed to send b to b'. Because b is not free in Γ, A and $D_{x,y}B$, the substitutions θ and θ' agree on them, and $(Q_yB(b,y))[\theta'] = Q_yB[\theta](b',y)$. The induction hypothesis for the second premise, with θ' and the context $\Delta, Q_yB[\theta](b',y)$, gives $\Delta, Q_yB[\theta](b',y) \vdash A[\theta]$; for the first premise it gives $\Delta \vdash D_{x,y}B[\theta]$. DE applies.

DI: choose z' fresh and let θ' send z to z' and agree with θ elsewhere. The induction hypothesis with θ' and the context $\Delta, B[\theta](t[\theta],0)$ gives $\Delta, B[\theta](t[\theta],0) \vdash B[\theta](t[\theta],Sz')$, and DI with the witness $t[\theta]$ applies.

QI: as for DI, with the premise $\Delta, E[\theta](z') \vdash E[\theta](Sz')$; the formula $E[\theta]$ is a literal equation because substitution preserves the root of a formula. □

Lemma 1.11 (Cut). *If $\Gamma \vdash A$ and $\Delta \vdash B$ for every $B \in \Gamma$, then $\Delta \vdash A$.*

Proof. The following more general statement is proved by induction on the derivation of $\Gamma \vdash A$: if $\Delta \vdash B[\theta]$ for every $B \in \Gamma$, then $\Delta \vdash A[\theta]$. Hyp is the assumption. I0 and IS are instances. Tr and QE follow from the induction hypotheses for their premises as in Theorem 1.10. For DE choose b' and θ' as in that proof. Every $C \in \Gamma$ satisfies $C[\theta'] = C[\theta]$, hence $\Delta, Q_y B[\theta](b', y) \vdash C[\theta']$ by weakening, and the new hypothesis is derivable from itself. The induction hypothesis for the second premise gives $\Delta, Q_y B[\theta](b', y) \vdash A[\theta]$, and DE applies. DI and QI are treated in the same way. □

In Lean the three properties are `NPrf.weaken`, `NPrf.subst` and `NPrf.cut`. They are used below without mention. For a context Γ, formulas A and B are *interderivable over* Γ if $\Gamma, A \vdash B$ and $\Gamma, B \vdash A$; for the empty Γ this is $A \dashv\vdash B$. By cut, interderivable formulas may replace each other as hypotheses and as conclusions.

1.4.2 Closure by cases and numerical cases

Lemma 1.12 (A witness that is not used)**.** *If $x \notin \mathrm{FV}(B)$, then $D_{x,y}B \dashv\vdash Q_y B$.*

Proof. From $D_{x,y}B$, DE with a fresh b adds the hypothesis $Q_y B[b/x]$, which is $Q_y B$, and Hyp derives the conclusion $Q_y B$. From $Q_y B$, DI with the witness 0 asks for $Q_y B,\ B(0) \vdash B(Sz)$, which is QE at Sz. □

Lemma 1.13 (Introduction of Q by cases)**.** *If $\Gamma, B(0) \vdash B(Sz)$ with z not free in Γ and $Q_y B$, then $\Gamma \vdash Q_y B(y)$. Here B is an arbitrary native formula.*

Proof. Choose x not free in B. DI with the witness 0 and the matrix B derives $\Gamma \vdash D_{x,y}B$ from the given premise, and Theorem 1.12 gives $\Gamma \vdash Q_y B$. □

Corollary 1.14 (Numerical cases)**.** *If $\Gamma \vdash B(0)$ and $\Gamma, B(0) \vdash B(Sz)$ with z fresh, then $\Gamma \vdash B(t)$ for every term t. In particular, if $\Gamma \vdash B(0)$ and $\Gamma \vdash B(Sz)$ with z fresh, then $\Gamma \vdash B(t)$.*

Proof. Theorem 1.13 and QE. □

Lemma 1.15 (Introduction of D from a closure)**.** *If $\Gamma \vdash Q_y B(t, y)$, then $\Gamma \vdash D_{x,y}B(x, y)$.*

Proof. DI with the witness t asks for $\Gamma, B(t, 0) \vdash B(t, Sz)$, which is QE at Sz applied to the given closure and the assumption. □

Corollary 1.16 (The witnessed and the combined introduction)**.** *Let $E(x, y)$ be a literal equation and z a fresh variable.*

(i) *If $\Gamma, E(t, z) \vdash E(t, Sz)$, then $\Gamma \vdash D_{x,y}E(x, y)$.*

(ii) *If $\Gamma, E(t, 0), E(t, z) \vdash E(t, Sz)$, then $\Gamma \vdash D_{x,y}E(x, y)$.*

Proof. (i) QI gives $\Gamma \vdash Q_y E(t, y)$, and Theorem 1.15 applies. (ii) Over the context $\Gamma, E(t, 0)$, QI gives $Q_y E(t, y)$, and QE at Sz with the hypothesis $E(t, 0)$ gives $E(t, Sz)$. This is the premise of DI with the witness t. □

In Lean the rules of this subsection are `NPrf.qI`, `NPrf.cases` and `NPrf.dIProg`.

Remark 1.17 (The older paired introduction). Part (ii) is the introduction rule of the oldest reports, which retains the instance at zero and, for an equation, the instance at the eigenvariable. The inherited basis replaces it by a choice between the two assumptions, one per inference, and proves that the consequence relation is unchanged. In one direction each choice is the old rule after a weakening. In the other direction only an old introduction at an equation

needs a construction, and it is the one in the proof of (ii). The witness t is kept as a parameter of the matrix $E(t, y)$, and the closure is introduced without a witness; an introduction and elimination of $D_{x,y}E(x, y)$ itself would return a fresh witness, not t. The reference counts inference nodes for its own presentation, in which the inner step is a witnessed introduction with an unused coordinate followed by DE: an old derivation with n nodes, r of them introductions at an equation, becomes a derivation with exactly $n + 5r \leq 6n$ nodes, the six nodes being the outer introduction by cases, the inner introduction by progress, DE, QE and two uses of Hyp. In N the inner introduction and its elimination are the single inference QI, so four nodes replace one. The bound counts nodes of the inference tree; it is not a bound on the size of formulas or of serialized certificates.

1.4.3 Equality

Lemma 1.18 (Equality rules)**.** *The following rules are derivable, each by the stated number of applications of* Tr*.*

(i) *Reflexivity:* $\Gamma \vdash t = t$ *(one application, on two instances of I0).*

(ii) *Symmetry:* $\Gamma \vdash s = t$ *implies* $\Gamma \vdash t = s$ *(one more).*

(iii) *Transitivity:* $\Gamma \vdash s = t$ *and* $\Gamma \vdash t = r$ *imply* $\Gamma \vdash s = r$ *(one more).*

(iv) *Congruence:* $\Gamma \vdash s = t$ *implies* $\Gamma \vdash h(s) = h(t)$ *for every term template* $h(v)$ *(one more).*

(v) *Transport in an equation template:* $\Gamma \vdash s = t$ *and* $\Gamma \vdash a(s) = b(s)$ *imply* $\Gamma \vdash a(t) = b(t)$ *(two).*

Proof. Write Tr_h for the rule with the template $h(v)$. (i) Apply Tr_v to the premises $I(0, t; u.\, u) = t$ and $I(0, t; u.\, u) = t$, both instances of I0; the conclusion is $t = t$. (ii) Apply Tr_v to $s = s$ and $s = t$; the conclusion is $t = s$. (iii) Apply Tr_v to $t = r$ and $t = s$, the latter by (ii); the conclusion is $s = r$. (iv) Apply Tr_h to $s = t$ and $h(s) = h(s)$; the conclusion is $h(s) = h(t)$. (v) Tr_a applied to $s = t$ and $a(s) = b(s)$ gives $b(s) = a(t)$, and Tr_b applied to $s = t$ and $b(s) = a(t)$ gives $a(t) = b(t)$. □

By (ii)–(iv), provable equality over a fixed context is a congruence for every term template, and chains of equations are used below in the usual way. The selector equations

$$\Gamma \vdash \mathrm{sel}(0, a, b) = a, \qquad \Gamma \vdash \mathrm{sel}(Sc, a, b) = b \tag{1.4}$$

are instances of I0 and IS.

Lemma 1.19 (Simultaneous congruence)**.** *If* $\Gamma \vdash s_i = t_i$ *for* $i = 1, \ldots, k$*, then* $\Gamma \vdash a[\vec{s}/\vec{x}] = a[\vec{t}/\vec{x}]$ *for every term* a*.*

Proof. Let a_j be a with $t_1, \ldots, t_j, s_{j+1}, \ldots, s_k$ substituted for $\vec{x}$. With a fresh variable v and the template $h_j(v)$ obtained by substituting $t_1, \ldots, t_{j-1}, v, s_{j+1}, \ldots, s_k$, one has $a_{j-1} = h_j(s_j)$ and $a_j = h_j(t_j)$ as terms, so Theorem 1.18(iv) gives $\Gamma \vdash a_{j-1} = a_j$. Transitivity joins $a_0 = a[\vec{s}/\vec{x}]$ to $a_k = a[\vec{t}/\vec{x}]$. □

Lemma 1.20 (Equation induction)**.** *Let* $E(y)$ *be a literal equation and* z *a variable not free in* Γ *and* $Q_y E$*. If* $\Gamma \vdash E(0)$ *and* $\Gamma, E(z) \vdash E(Sz)$*, then* $\Gamma \vdash E(t)$ *for every term* t*.*

Proof. QI and QE. □

In Lean this rule is `NPrf.eqInd`.

1.4.4 Congruence of the binders and the Leibniz rule

Lemma 1.21 (Congruence of Q and D)**.** *Let B and B' be interderivable over Γ.*

(i) *If y is not free in Γ, then Q_yB and Q_yB' are interderivable over Γ.*

(ii) *If x and y are not free in Γ, then $D_{x,y}B$ and $D_{x,y}B'$ are interderivable over Γ.*

Proof. Both parts rest on one derivation. Let $C(y)$ and $C'(y)$ be interderivable over Γ, with y not free in Γ, and let z be fresh. Then

$$\Gamma,\ Q_yC(y),\ C'(0) \vdash C'(Sz). \tag{1.5}$$

Indeed, substituting 0 for y in $\Gamma, C' \vdash C$ gives $\Gamma, C'(0) \vdash C(0)$, because y is not free in Γ; QE at Sz gives $C(Sz)$; and substituting Sz for y in $\Gamma, C \vdash C'$ gives $\Gamma, C(Sz) \vdash C'(Sz)$.

(i) Theorem 1.13 applied to (1.5) for $C = B$ and $C' = B'$ gives $\Gamma, Q_yB \vdash Q_yB'$. The converse is symmetric.

(ii) Over $\Gamma, D_{x,y}B$ apply DE with a fresh b; the new hypothesis is $Q_yB(b,y)$. Substituting b for x shows that $B(b,y)$ and $B'(b,y)$ are interderivable over Γ. DI with the witness b and the matrix B' asks for $\Gamma, Q_yB(b,y), B'(b,0) \vdash B'(b,Sz)$, which is (1.5) for $C = B(b,y)$ and $C' = B'(b,y)$. Hence $\Gamma, D_{x,y}B \vdash D_{x,y}B'$, and the converse is symmetric. □

The binder Q is contravariant in its instance at zero, which is why both directions of the hypothesis are used in each direction of the conclusion.

Theorem 1.22 (Leibniz rule)**.** *Let A be a native formula and let $\Gamma \vdash s_i = t_i$ for $i = 1, \dots, k$. Then $A[\vec{s}/\vec{x}]$ and $A[\vec{t}/\vec{x}]$ are interderivable over Γ. In particular $\Gamma \vdash A[\vec{s}/\vec{x}]$ implies $\Gamma \vdash A[\vec{t}/\vec{x}]$.*

Proof. By induction on A, for all Γ, $\vec{s}$ and $\vec{t}$. If A is $a = b$, then Theorem 1.19 gives $\Gamma \vdash a[\vec{s}/\vec{x}] = a[\vec{t}/\vec{x}]$ and the same for b, and from the hypothesis $a[\vec{s}/\vec{x}] = b[\vec{s}/\vec{x}]$ symmetry and transitivity give $a[\vec{t}/\vec{x}] = b[\vec{t}/\vec{x}]$; the converse is the same argument with $\vec{s}$ and $\vec{t}$ exchanged. If A is Q_yB, rename y so that it is not free in Γ, $\vec{s}$, $\vec{t}$ and differs from $\vec{x}$. The induction hypothesis makes $B[\vec{s}/\vec{x}]$ and $B[\vec{t}/\vec{x}]$ interderivable over Γ, and Theorem 1.21(i) applies. The case $D_{x,y}B$ is the same with Theorem 1.21(ii). □

Corollary 1.23 (Transport in a formula template)**.** *If $\Gamma \vdash s = t$ and $\Gamma \vdash F(s)$, then $\Gamma \vdash F(t)$, for every native formula template $F(v)$.*

In Lean these are `NPrf.leibniz` and `NPrf.tr`; the two-step rule of Theorem 1.18(v) is `NPrf.trEq`. Transport never has to be primitive below a binder: the template is taken apart by QE, DE and DI.

1.4.5 Gates and selectors of formulas

A case distinction on a term cannot be expressed by a connective. It is pushed into a formula, down to its equations.

Definition 1.24 (Selector and gate of formulas)**.** For a term v and native formulas A, B the native formula $\mathrm{sel}(v, A, B)$ is defined by recursion, first on A and, for an equation A, on B. Bound variables are chosen fresh for v and for the other formula.

$$\begin{aligned}
\mathrm{sel}(v,\ s = t,\ r = w) &:= \big(\mathrm{sel}(v,s,r) = \mathrm{sel}(v,t,w)\big),\\
\mathrm{sel}(v,\ s = t,\ Q_yB) &:= Q_y\ \mathrm{sel}\big(v,\ \mathrm{sel}(y,0,s) = \mathrm{sel}(y,0,t),\ B\big),\\
\mathrm{sel}(v,\ s = t,\ D_{x,y}B) &:= D_{x,y}\ \mathrm{sel}\big(v,\ \mathrm{sel}(y,0,s) = \mathrm{sel}(y,0,t),\ B\big),\\
\mathrm{gate}(v,B) &:= \mathrm{sel}(v,\ 0 = 0,\ B),\\
\mathrm{sel}(v,\ Q_yA,\ B) &:= Q_y\ \mathrm{sel}\big(v,\ A,\ \mathrm{gate}(y,B)\big),\\
\mathrm{sel}(v,\ D_{x,y}A,\ B) &:= D_{x,y}\ \mathrm{sel}\big(v,\ A,\ \mathrm{gate}(y,B)\big).
\end{aligned}$$

The intention is that $\mathrm{sel}(v, A, B)$ is A when v is zero and B when v is a successor, and that $\mathrm{gate}(v, B)$ is trivially derivable when v is zero and is B when v is a successor. When a binder of one formula has no counterpart in the other, the other formula is guarded by the bound variable, so that the instance of the binder at zero is trivial. By induction on the definition, substitution commutes with both operations:

$$\mathrm{sel}(v, A, B)[\theta] = \mathrm{sel}(v[\theta], A[\theta], B[\theta]), \qquad \mathrm{gate}(v, B)[\theta] = \mathrm{gate}(v[\theta], B[\theta]). \tag{1.6}$$

Lemma 1.25 (A guarded equation under a binder)**.** *If x and y are not free in s and t, then*

$$Q_y\big(\mathrm{sel}(y, 0, s) = \mathrm{sel}(y, 0, t)\big) \dashv\vdash (s = t) \dashv\vdash D_{x,y}\big(\mathrm{sel}(y, 0, s) = \mathrm{sel}(y, 0, t)\big).$$

Proof. From the closure: its instance at 0 is $\mathrm{sel}(0, 0, s) = \mathrm{sel}(0, 0, t)$, which is derivable because both sides equal 0 by (1.4). QE at 1 gives $\mathrm{sel}(1, 0, s) = \mathrm{sel}(1, 0, t)$, and (1.4) turns it into $s = t$. From $s = t$: (1.4) gives $\mathrm{sel}(Sz, 0, s) = s = t = \mathrm{sel}(Sz, 0, t)$, which is the premise of Theorem 1.13. The second equivalence follows from the first by Theorem 1.12. □

Lemma 1.26 (Laws of the gate and the selector)**.** *For all terms v and native formulas A, B:*

(i) $\vdash \mathrm{gate}(0, B)$*;*

(ii) $\mathrm{gate}(Sv, B) \dashv\vdash B$*;*

(iii) $Q_y\,\mathrm{gate}(y, B) \dashv\vdash B \dashv\vdash D_{x,y}\,\mathrm{gate}(y, B)$ *for $x, y \notin \mathrm{FV}(B)$;*

(iv) $\mathrm{sel}(0, A, B) \dashv\vdash A$*;*

(v) $\mathrm{sel}(Sv, A, B) \dashv\vdash B$*.*

Proof. First let A be an equation $s = t$. The two statements

$$\mathrm{sel}(0,\ s = t,\ B) \dashv\vdash (s = t), \qquad \mathrm{sel}(Sv,\ s = t,\ B) \dashv\vdash B$$

are proved by induction on B, for all s, t and v. If B is $r = w$, they hold by (1.4) and the equality rules. If B is $Q_y B'$, the induction hypothesis for B' and the equation $\mathrm{sel}(y, 0, s) = \mathrm{sel}(y, 0, t)$ gives

$$\mathrm{sel}\big(0,\ \mathrm{sel}(y, 0, s) = \mathrm{sel}(y, 0, t),\ B'\big) \dashv\vdash \big(\mathrm{sel}(y, 0, s) = \mathrm{sel}(y, 0, t)\big),$$
$$\mathrm{sel}\big(Sv,\ \mathrm{sel}(y, 0, s) = \mathrm{sel}(y, 0, t),\ B'\big) \dashv\vdash B'.$$

By Theorem 1.21(i) the first formula of the statement is interderivable with $Q_y(\mathrm{sel}(y, 0, s) = \mathrm{sel}(y, 0, t))$, hence with $s = t$ by Theorem 1.25, and the second with $Q_y B'$. The case $D_{x,y}B'$ is the same with Theorem 1.21(ii) and the second equivalence of Theorem 1.25.

(i) and (ii) are the two statements for the equation $0 = 0$, which is derivable by reflexivity.

(iii) From $Q_y\,\mathrm{gate}(y, B)$: the instance at 0 is derivable by (i), QE at 1 gives $\mathrm{gate}(1, B)$, and (ii) gives B. From B: (ii) gives $\mathrm{gate}(Sz, B)$, the premise of Theorem 1.13. The second equivalence follows by Theorem 1.12.

(iv) and (v) are proved by induction on A, for all v and B. For an equation they are the two statements above. If A is $Q_y A'$, then $\mathrm{sel}(0, A, B) = Q_y\,\mathrm{sel}(0, A', \mathrm{gate}(y, B))$ is interderivable with $Q_y A'$ by the induction hypothesis and Theorem 1.21(i); and $\mathrm{sel}(Sv, A, B) = Q_y\,\mathrm{sel}(Sv, A', \mathrm{gate}(y, B))$ is interderivable with $Q_y\,\mathrm{gate}(y, B)$ by the induction hypothesis and Theorem 1.21(i), hence with B by (iii). The case $D_{x,y}A'$ is the same with Theorem 1.21(ii). □

Lemma 1.27 (Selector at an arbitrary index)**.** (i) *If $\Gamma \vdash A$ and $\Gamma \vdash B$, then $\Gamma \vdash \mathrm{sel}(w, A, B)$ for every term w.*

(ii) *If $\Gamma \vdash v = w$ and $\Gamma \vdash \mathrm{sel}(v, A, B)$, then $\Gamma \vdash \mathrm{sel}(w, A, B)$.*

Proof. (i) Apply Theorem 1.14 to the formula $\mathrm{sel}(y, A, B)$ with y fresh: its instance at 0 follows from A by Theorem 1.26(iv), and its instance at Sz follows from B by Theorem 1.26(v). The instances are computed by (1.6). (ii) Theorem 1.23 with the template $\mathrm{sel}(y, A, B)$. □

1.4.6 The connectives as native formulas

Definition 1.28 (Native connectives)**.** With fresh bound variables and the parity term N of Theorem 1.3,

$$\begin{aligned} \bot_* &:= (0 = S0), \\ A \Rightarrow_* B &:= Q_y\ \mathrm{sel}(y, A, B), \\ A \wedge_* B &:= Q_y\ \mathrm{gate}\big(y,\ \mathrm{sel}(N(y), A, B)\big), \\ \exists_* x\, A &:= D_{x,y}\ \mathrm{gate}(y, A), \\ A \vee_* B &:= \exists_* x\ \mathrm{sel}(x, A, B), \\ \forall_* x\, A &:= A(0) \wedge_* Q_x A. \end{aligned}$$

By (1.6) each of these operations commutes with substitution. The parity term satisfies

$$\vdash N(0) = 0, \qquad \vdash N(1) = 1, \qquad \vdash N(2) = 0, \tag{1.7}$$

by I0, by $N(S0) = Z(N(0)) = Z(0) = \mathrm{sel}(0, 1, 0) = 1$, and by $N(S1) = Z(N(1)) = Z(1) = \mathrm{sel}(S0, 1, 0) = 0$. The conjunction reads its two operands off the instances at 1 and at 2.

Theorem 1.29 (Natural deduction for the native connectives)**.** *The following rules are derivable in* N*, for all native formulas.*

(i) $\bot_* E$*:* $\Gamma \vdash \bot_*$ *implies* $\Gamma \vdash A$*.*

(ii) $\Rightarrow_* I$*:* $\Gamma, A \vdash B$ *implies* $\Gamma \vdash A \Rightarrow_* B$*.* $\Rightarrow_* E$*:* $\Gamma \vdash A \Rightarrow_* B$ *and* $\Gamma \vdash A$ *imply* $\Gamma \vdash B$*.*

(iii) $\wedge_* I$*:* $\Gamma \vdash A$ *and* $\Gamma \vdash B$ *imply* $\Gamma \vdash A \wedge_* B$*.* $\wedge_* E$*:* $\Gamma \vdash A \wedge_* B$ *implies* $\Gamma \vdash A$ *and* $\Gamma \vdash B$*.*

(iv) $\exists_* I$*:* $\Gamma \vdash A(t)$ *implies* $\Gamma \vdash \exists_* x\, A(x)$*.* $\exists_* E$*:* $\Gamma \vdash \exists_* x\, A(x)$ *and* $\Gamma, A(b) \vdash C$*, with* b *not free in* Γ*,* C *and* $\exists_* x\, A$*, imply* $\Gamma \vdash C$*.*

(v) $\vee_* I$*: each of* $\Gamma \vdash A$*,* $\Gamma \vdash B$ *implies* $\Gamma \vdash A \vee_* B$*.* $\vee_* E$*:* $\Gamma \vdash A \vee_* B$*,* $\Gamma, A \vdash C$ *and* $\Gamma, B \vdash C$ *imply* $\Gamma \vdash C$*.*

(vi) $\forall_* I$*:* $\Gamma \vdash A(x)$ *with* x *not free in* Γ *implies* $\Gamma \vdash \forall_* x\, A(x)$*.* $\forall_* E$*:* $\Gamma \vdash \forall_* x\, A(x)$ *implies* $\Gamma \vdash A(t)$*.*

Proof. (i) From $0 = S0$ every equation $s = t$ follows: $s = \mathrm{sel}(0, s, t) = \mathrm{sel}(S0, s, t) = t$, by (1.4) and congruence in the first argument of the selector. Every formula follows by induction on the number of its binders, a measure that substitution does not change. For $Q_y B$ use Theorem 1.13, whose premise $B(Sz)$ holds by the induction hypothesis; for $D_{x,y} B$ use DI with the witness 0, whose premise $B(0, Sz)$ holds by the induction hypothesis.

(ii) Introduction. By Theorem 1.13 it suffices to derive $\Gamma, \mathrm{sel}(0, A, B) \vdash \mathrm{sel}(Sz, A, B)$. Theorem 1.26(iv) gives A, the premise gives B, and Theorem 1.26(v) gives $\mathrm{sel}(Sz, A, B)$. Elimination. Theorem 1.26(iv) turns A into $\mathrm{sel}(0, A, B)$, QE at 1 gives $\mathrm{sel}(1, A, B)$, and Theorem 1.26(v) gives B.

(iii) Introduction. By Theorem 1.13 it suffices to derive $\mathrm{gate}(Sz, \mathrm{sel}(N(Sz), A, B))$ from Γ, and by Theorem 1.26(ii) it suffices to derive $\mathrm{sel}(N(Sz), A, B)$, which is Theorem 1.27(i). Elimination. The instance of the matrix at 0 is derivable by Theorem 1.26(i). QE at Sc and Theorem 1.26(ii) give $\Gamma \vdash \mathrm{sel}(N(Sc), A, B)$ for every term c. For $c = 1$, (1.7) and Theorem 1.27(ii) give $\mathrm{sel}(0, A, B)$, hence A. For $c = 0$ they give $\mathrm{sel}(1, A, B)$, hence B.

(iv) Introduction. DI with the witness t asks for $\mathrm{gate}(Sz, A(t))$, which follows from $A(t)$ by Theorem 1.26(ii). Elimination. DE with the fresh variable b adds the hypothesis $Q_y\, \mathrm{gate}(y, A(b))$, which gives $A(b)$ by Theorem 1.26(iii); cut with the second premise gives C.

(v) Introduction is (iv) with the witnesses 0 and 1 and Theorem 1.26(iv), (v). Elimination. By (iv) it suffices to derive C from $\Gamma, \mathrm{sel}(b, A, B)$ with b fresh. Apply Theorem 1.14 to the formula $\mathrm{sel}(y, A, B) \Rightarrow_* C$ with y fresh. Its instance at 0 follows by (ii) from $\Gamma, \mathrm{sel}(0, A, B) \vdash C$, which holds by Theorem 1.26(iv) and the premise $\Gamma, A \vdash C$; its instance at Sz follows in the same way from Theorem 1.26(v) and $\Gamma, B \vdash C$. The instance at b and (ii) give C.

(vi) Introduction. Substituting 0 for x in the premise gives $\Gamma \vdash A(0)$, and substituting Sz gives $\Gamma \vdash A(Sz)$, hence $\Gamma \vdash Q_x A$ by Theorem 1.13; (iii) combines the two. Elimination. By (iii) the premise gives $A(0)$ and $Q_x A$, and QE at t gives $A(t)$. □

Remark 1.30 (Which rules the logical interface uses). No derivation in this section applies QI, except the derivations of Theorem 1.20 and Theorem 1.16. The equality rules, the Leibniz rule, numerical cases and the whole of Theorem 1.29 are derived from the seven other rules. Cases supply the logic and progress supplies induction for equations; the two do not mix. If QI were allowed at a compound matrix, the proof of Theorem 1.20 would give induction for every native formula. The universal quantifier needs no numerical case distinction: its elimination is QE. The reference statement of the inherited basis, which has no binder Q, defines implication and the universal quantifier through D with an unused coordinate and eliminates the universal quantifier through the derived coverage $x = 0 \vee_* \exists_* v\,(x = Sv)$. Its abbreviations differ in detail from the ones above (the selector aligns the binders of its two formulas, and the matrix of the universal quantifier is gated), but they satisfy the same introduction and elimination rules and are therefore interderivable with them. In Lean the connectives and their rules are the module `NativeLogic`.

1.5 The ordinary theory T_{eq}

This section draws on the reference statement of the inherited ordinary bridge filed with `certified_zero` and on the section on the ordinary realization of `raw_clausal`.

Definition 1.31 (The logic). *Intuitionistic logic with equality* over a language of terms is the natural deduction calculus for sequents $\Gamma \vdash \varphi$ of ordinary formulas with the rules: hypothesis; $\bot$E; introduction and elimination for $\wedge$, $\vee$, $\to$, $\forall$ and $\exists$, with the usual eigenvariable conditions; reflexivity $\Gamma \vdash t = t$; and the Leibniz rule, from $\Gamma \vdash s_i = t_i$ for all i and $\Gamma \vdash \varphi[\vec{s}/\vec{x}]$ infer $\Gamma \vdash \varphi[\vec{t}/\vec{x}]$. A *theory* is a set of formulas, its axioms; every substitution instance of an axiom is derivable in every context. Derivability is closed under weakening, substitution and cut.

Definition 1.32 (T_{eq}). T_{eq} is the theory in intuitionistic logic with equality over the terms of Theorem 1.1 with the following axioms.

(T1) The computation equations (1.1) and (1.2), for all terms.

(T2) Coverage: $x = 0 \ \vee \ \exists y\,(x = Sy)$.

(T3) Separation: $0 = S0 \to \bot$.

(T4) Induction for literal equations: for every equation $E(z)$,

$$E(0) \to \big(\forall z\,(E(z) \to E(Sz)) \to \forall z\,E(z)\big).$$

Injectivity of the successor, $Sx \neq 0$ and induction for compound formulas are not axioms. In Lean the axioms are `TeqAx`.

Lemma 1.33 (Cases and contextual induction in T_{eq}). (i) *For every ordinary formula $\varphi(y)$: if $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} \varphi(0)$ and $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} \varphi(Sz)$ with z not free in Γ and φ, then $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} \forall y\,\varphi(y)$.*

(ii) *For every equation* $E(z)$*: if* $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} E(0)$ *and* $\Gamma, E(z) \vdash_{\mathsf{T}_{\mathrm{eq}}} E(Sz)$ *with* z *not free in* Γ*, then* $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} E(t)$ *for every term* t*.*

Proof. (i) Let y be fresh and eliminate the disjunction (T2) at $x = y$. In the branch $y = 0$ the Leibniz rule turns $\varphi(0)$ into $\varphi(y)$. In the other branch eliminate the existential quantifier at the fresh variable z; from $y = Sz$ the Leibniz rule turns $\varphi(Sz)$ into $\varphi(y)$. Generalize over y. (ii) Discharge $E(z)$, generalize over z, apply (T4) and instantiate at t. The context Γ is arbitrary, so the rule holds with quantified side hypotheses. □

Remark 1.34 (Separation is an axiom). The computation equations, coverage and equation induction hold in the structure with one element, in which every equation is true. Separation fails there. Hence separation does not follow from the other axioms, and the native contradiction $0 = S0$ has to be linked to $\bot$ explicitly: the reading of the native calculus proves every formula from $0 = S0$, and the translation of $\bot$ is that equation, but neither fact removes (T3) from T_{eq}.

1.6 Intuitionistic primitive recursive arithmetic

This section fixes the usual presentation, as in the reference statement on equality reflection filed with `certified_zero` and in the Lean module `PR`.

Definition 1.35 (iPRA)**.** The *function symbols* are generated as follows, each with an arity: zero of arity 0; succ of arity 1; the projections π^k_i of arity k, for $1 \le i \le k$; the composition $\mathrm{comp}(f; g_1, \dots, g_m)$ of arity k, for f of arity m and $g_1, \dots, g_m$ of arity k; and the recursion $\mathrm{prec}(f, g)$ of arity $k+1$, for f of arity k and g of arity $k+2$. The terms are the variables and the applications $f(t_1, \dots, t_k)$; 0 is zero() and St is succ(t). iPRA is the theory in intuitionistic logic with equality over these terms with the axioms

$$\begin{aligned} \pi^k_i(\vec{x}) &= x_i, & \mathrm{comp}(f; \vec{g})(\vec{x}) &= f(g_1(\vec{x}), \dots, g_m(\vec{x})), \\ \mathrm{prec}(f, g)(0, \vec{x}) &= f(\vec{x}), & \mathrm{prec}(f, g)(Sy, \vec{x}) &= g\big(y, \mathrm{prec}(f, g)(y, \vec{x}), \vec{x}\big), \\ Sx = 0 &\to \bot, & Sx = Sy &\to x = y, \end{aligned}$$

and induction for quantifier-free formulas: for every quantifier-free $\varphi(z)$,

$$\varphi(0) \to \big(\forall z\, (\varphi(z) \to \varphi(Sz)) \to \forall z\, \varphi(z)\big).$$

The recursion argument of $\mathrm{prec}(f, g)$ comes first. Distinct descriptions are distinct symbols, even when they compute the same function on the natural numbers. The quantifiers of the logic are unrestricted; only the induction formula is quantifier free. In Lean the axioms are `PRAAx`.

1.7 Arithmetic inside T_{eq}

This section draws on the reference statements filed with `certified_zero` (scalar predecessor, equality reflection, primitive recursion) and, for the predecessor, on the equational derivations of `uniform_raw`, `newest_parameter`, `parameter_witness` and `no_witness`. The language has no predecessor and no pairing, and the body of an iterator does not see the counter. The arithmetic is developed in this order: a predecessor term; truncated subtraction and a term that reflects equality; induction for quantifier-free formulas; pairing; primitive recursion. No later item is used for an earlier one.

1.7.1 The equational fragment

Definition 1.36 (The fragment $\mathcal{E}$)**.** A sequent of $\mathcal{E}$ is $\Gamma \vdash s = t$ with Γ a finite list of equations. The rules of $\mathcal{E}$ are Hyp, I0, IS, Tr and *equation induction*:

$$\frac{\Gamma \vdash E(0) \qquad \Gamma, E(z) \vdash E(Sz)}{\Gamma \vdash E(t)}\ \text{EI} \qquad (E \text{ an equation, } z \text{ not free in } \Gamma \text{ and in } E(0)).$$

Lemma 1.37 (Properties of $\mathcal{E}$)**.** (i) *$\mathcal{E}$ is closed under weakening, substitution and cut.*

(ii) *Reflexivity, symmetry, transitivity and congruence (Theorems 1.18 and 1.19) are derived rules of $\mathcal{E}$.*

(iii) Equational cases*: if $\Gamma \vdash E(0)$ and $\Gamma \vdash E(Sz)$ with z fresh, then $\Gamma \vdash E(t)$.*

(iv) *If $\Gamma \vdash_{\mathcal{E}} s = t$, then $\Gamma \vdash_{\mathsf{N}} s = t$ and $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} s = t$.*

Proof. (i) The proofs of Theorems 1.10 and 1.11 apply; EI is treated like QI, with the eigenvariable renamed. (ii) The derivations use I0 and Tr only. (iii) Weaken the second premise by $E(z)$ and apply EI. (iv) EI is the derived rule of Theorem 1.20 in N and of Theorem 1.33(ii) in T_{eq}; Tr follows in T_{eq} from the Leibniz rule; the other rules are shared. □

By (i), an equation derived in $\mathcal{E}$ with variables that are not free in the context may be instantiated at arbitrary terms. Every case distinction below proves an equation with an independent case variable and is an instance of (iii); no hypothesis is ever specialized to a case. The fragment has no coverage axiom, no disjunction, no separation and no rule for D. Until Theorem 1.48 all derivations are in $\mathcal{E}$, in the empty context unless a context is shown.

Lemma 1.38 (Uniqueness of iteration)**.** *Let $f(z)$ be a term template and a, $h(u)$ terms in which z is not free, and let z be not free in Γ. If $\Gamma \vdash f(0) = a$ and $\Gamma \vdash f(Sz) = h(f(z))$, then $\Gamma \vdash f(c) = I(c, a;\, u.\, h)$ for every term c.*

Proof. EI on the equation $f(z) = I(z, a;\, u.\, h)$. The base is $f(0) = a = I(0, a;\, u.\, h)$. In the step, $f(Sz) = h(f(z)) = h(I(z, a;\, u.\, h)) = I(Sz, a;\, u.\, h)$, by the hypothesis, congruence on the induction equation, and IS. □

In Lean the rule is `NPrf.uniq`.

Corollary 1.39 (Rotation and congruence of the body)**.** (i) $\vdash I(Sy, a;\, u.\, h) = I(y, h(a);\, u.\, h)$.

(ii) *If $\Gamma \vdash h(u) = h'(u)$ with u not free in Γ, then $\Gamma \vdash I(c, a;\, u.\, h) = I(c, a;\, u.\, h')$.*

Proof. (i) Take $f(z) := I(Sz, a;\, u.\, h)$. Then $f(0) = h(I(0, a;\, u.\, h)) = h(a)$ and $f(Sz) = h(f(z))$ by IS, I0 and congruence, and Theorem 1.38 gives the claim. (ii) Take $f(z) := I(z, a;\, u.\, h)$. Then $f(0) = a$ and $f(Sz) = h(f(z)) = h'(f(z))$, the last step by instantiating the hypothesis at $u = f(z)$. Theorem 1.38 with the body h' gives the claim. □

Thus no extensionality rule for bodies is assumed; it is an equation induction.

1.7.2 Addition

Put $a + b := I(b, a;\, u.\, Su)$, $2a := a + a$ and $3a := (a + a) + a$; the same terms are also written $2 \cdot a$ and $3 \cdot a$.

Lemma 1.40 (Addition)**.** $\mathcal{E}$ *derives*

$$a+0=a, \qquad a+Sb=S(a+b), \qquad 0+b=b, \qquad Sa+b=S(a+b),$$
$$a+b=b+a, \qquad (a+b)+c=a+(b+c).$$

Consequently $Sa=a+1$, $2(Sa)=SS(2a)$, $3(Sa)=SSS(3a)$, $3(a+b)=3a+3b$, $2\cdot 0=0$ *and* $3\cdot 0=0$.

Proof. The first two equations are I0 and IS. The others are equation inductions on the last variable shown. $0+b=b$: the base is $0+0=0$, and $0+Sb=S(0+b)=Sb$. $Sa+b=S(a+b)$: the base is $Sa+0=Sa=S(a+0)$, and $Sa+Sb=S(Sa+b)=SS(a+b)=S(a+Sb)$. Commutativity: the base is $a+0=a=0+a$, and $a+Sb=S(a+b)=S(b+a)=Sb+a$. Associativity: the base is $(a+b)+0=a+b=a+(b+0)$, and $(a+b)+Sc=S((a+b)+c)=S(a+(b+c))=a+S(b+c)=a+(b+Sc)$. The consequences follow by these equations. □

Sums are rearranged below by Theorem 1.40 without comment. No cancellation law is used, and there is no multiplication.

1.7.3 Zero test, interpolation and parity

Lemma 1.41 (Zero test and interpolation)**.** $\mathcal{E}$ *derives the following, for all terms and every term template* $F(v)$.

(i) $Z(0)=1$ *and* $Z(Sc)=0$.

(ii) $\mathrm{sel}(c,a,a)=a$ *and* $\mathrm{sel}(c,Sa,Sb)=S\,\mathrm{sel}(c,a,b)$.

(iii) $Z(Z(Z(c)))=Z(c)$.

(iv) Equational interpolation*:* $F(Z(c))=\mathrm{sel}\big(c,F(1),F(0)\big)$.

(v) Two-point rule*: if* $\Gamma\vdash F(0)=F'(0)$, $\Gamma\vdash F(1)=F'(1)$ *and* $\Gamma\vdash Z(Z(b))=b$, *then* $\Gamma\vdash F(b)=F'(b)$.

Proof. (i) is (1.4). (ii)–(iv) are equational cases on a fresh variable in the place of c. For (ii): at 0 both sides of each equation are a, respectively Sa; at Sz they are a, respectively Sb. For (iii): $Z(Z(Z(0)))=Z(Z(1))=Z(0)$, and $Z(Z(Z(Sz)))=Z(Z(0))=Z(1)=0=Z(Sz)$. For (iv): at 0, $F(Z(0))=F(1)=\mathrm{sel}(0,F(1),F(0))$; at Sz, $F(Z(Sz))=F(0)=\mathrm{sel}(Sz,F(1),F(0))$. (v) Instantiate (iv) for F and for F' at $c=Z(b)$:

$$F(Z(Z(b)))=\mathrm{sel}\big(Z(b),F(1),F(0)\big)=\mathrm{sel}\big(Z(b),F'(1),F'(0)\big)=F'(Z(Z(b))),$$

the middle step by congruence from the two hypotheses. Congruence along $Z(Z(b))=b$ on both ends gives $F(b)=F'(b)$. □

The two-point rule is the replacement for a case distinction on a Boolean value. It is an equality rule, not an elimination of a disjunction: the term b and the context Γ are arbitrary, and Γ may contain equations in which the variables of b occur.

Lemma 1.42 (Parity)**.** $\mathcal{E}$ *derives*

$$N(0)=0, \qquad N(Sn)=Z(N(n)), \qquad N(1)=1, \qquad Z(Z(N(n)))=N(n),$$
$$N(n)+Z(N(n))=1, \qquad N(SSn)=N(n),$$
$$N(2n)=0, \qquad N(S(2n))=1, \qquad N(3n)=N(n).$$

Proof. The first two equations are computations and give $N(1) = Z(0) = 1$. $Z(Z(N(n))) = N(n)$ is an equation induction on n: the base is $Z(Z(0)) = Z(1) = 0 = N(0)$, and $Z(Z(N(Sn))) = Z(Z(Z(N(n)))) = Z(N(n)) = N(Sn)$ by Theorem 1.41(iii); the induction equation is not used. The equation $N(n) + Z(N(n)) = 1$ is the two-point rule for $F(v) := v + Z(v)$, $F'(v) := 1$ and $b := N(n)$: $F(0) = 0 + 1 = 1$ and $F(1) = 1 + 0 = 1$. Two computations give $N(SSn) = Z(Z(N(n))) = N(n)$. $N(2n) = 0$ is an induction: $N(2 \cdot 0) = N(0) = 0$, and $N(2(Sn)) = N(SS(2n)) = N(2n) = 0$. Then $N(S(2n)) = Z(N(2n)) = Z(0) = 1$. $N(3n) = N(n)$ is an induction: $N(3 \cdot 0) = N(0)$, and $N(3(Sn)) = N(SSS(3n)) = Z(N(SS(3n))) = Z(N(3n)) = Z(N(n)) = N(Sn)$. □

No statement that a number is even or odd is involved.

1.7.4 A three-cycle and a ramp

Put

$$j(u) := Z(u) + 2N(u), \qquad M(n) := I(n, 0;\, u.\, j(u)), \qquad G(n) := I(n, 0;\, u.\, S(u + M(u))).$$

Lemma 1.43 (Three-cycle)**.** $\mathcal{E}$ *derives* $j(0) = 1$, $j(1) = 2$, $j(2) = 0$, $M(0) = 0$ *and* $M(Sn) = j(M(n))$.

Proof. $j(0) = Z(0) + 2N(0) = 1 + (0 + 0) = 1$; $j(1) = Z(1) + 2N(1) = 0 + (1 + 1) = 2$; $j(2) = Z(S1) + 2N(SS0) = 0 + 2N(0) = 0$. The last two equations are I0 and IS. □

Only these three values of j are used; no theorem about remainders modulo three is needed.

Lemma 1.44 (Ramp)**.** $\mathcal{E}$ *derives*

(i) $M(G(n)) = N(n)$;

(ii) $G(0) = 0$, $G(Sn) = S(G(n) + N(n))$ *and* $G(SSn) = SSSG(n)$;

(iii) $G(2n) = 3n$;

(iv) $G(3n) = 3G(n) + N(n)$.

Proof. (i) Two identities are derived first, for variables g and b, in the context that consists of the one equation $Z(Z(b)) = b$:

$$M(S(g + b)) = \mathrm{sel}\big(b,\ j(M(g)),\ j(j(M(g)))\big), \tag{1.8}$$

$$\mathrm{sel}\big(b,\ j(b),\ j(j(b))\big) = Z(b). \tag{1.9}$$

Both are instances of the two-point rule. For (1.8) the templates are $F(v) := M(S(g + v))$ and $F'(v) := \mathrm{sel}(v, j(M(g)), j(j(M(g))))$: $F(0) = M(Sg) = j(M(g)) = F'(0)$ and $F(1) = M(SSg) = j(j(M(g))) = F'(1)$. For (1.9) the templates are $\mathrm{sel}(v, j(v), j(j(v)))$ and $Z(v)$: at 0 both sides are $j(0) = 1 = Z(0)$, and at 1 they are $j(j(1)) = j(2) = 0 = Z(1)$, by Theorem 1.43. By Theorem 1.42 the hypothesis is derivable for $b = N(n)$, so by substitution and cut both identities hold at $g = G(n)$ and $b = N(n)$ in every context.

Now EI on n. The base is $M(G(0)) = M(0) = 0 = N(0)$. In the step the induction equation is $M(G(n)) = N(n)$, and

$$\begin{aligned} M(G(Sn)) &= M\big(S(G(n) + M(G(n)))\big) = M\big(S(G(n) + N(n))\big) \\ &= \mathrm{sel}\big(N(n),\ j(M(G(n))),\ j(j(M(G(n))))\big) \\ &= \mathrm{sel}\big(N(n),\ j(N(n)),\ j(j(N(n)))\big) = Z(N(n)) = N(Sn). \end{aligned}$$

The second and fourth steps use the induction equation by congruence, the third is (1.8) and the fifth is (1.9). The value $N(n)$ is not split under the induction hypothesis.

(ii) The first equation is I0; the second is IS and (i). Applying it twice, $G(SSn) = S(S(G(n)+N(n))+Z(N(n))) = SS\big(G(n)+(N(n)+Z(N(n)))\big) = SS(G(n)+1) = SSSG(n)$, by Theorems 1.40 and 1.42.

(iii) EI: $G(2\cdot 0) = G(0) = 0 = 3\cdot 0$, and $G(2(Sn)) = G(SS(2n)) = SSSG(2n) = SSS(3n) = 3(Sn)$.

(iv) EI. The base is $G(3\cdot 0) = 0 = 3G(0) + N(0)$. In the step, by (ii), the induction equation and $N(3n) = N(n)$,

$$G(3(Sn)) = G(SSS(3n)) = SSS\,S\big(G(3n) + N(3n)\big) = SSSS\big(3G(n) + N(n) + N(n)\big),$$

and, by (ii) and $N(n) + Z(N(n)) = 1$,

$$\begin{aligned} 3G(Sn) + N(Sn) &= 3\,S\big(G(n) + N(n)\big) + Z(N(n)) = SSS\big(3G(n) + 3N(n) + Z(N(n))\big) \\ &= SSS\big(3G(n) + N(n) + N(n) + 1\big). \end{aligned}$$

The two right-hand sides are equal by Theorem 1.40. □

1.7.5 The factor map and the comparator

Put

$$H(u) := \mathrm{sel}\big(N(u), G(u), u\big), \qquad 2^x := I(x, 1;\, u.\, u + u), \qquad L(x,y) := N\big(I(y, 2^x;\, u.\, H(u))\big).$$

The term H replaces a factor two by a factor three and fixes the numbers of parity one; 2^x is a term, not an operation of the language.

Lemma 1.45 (Factor map)**.** $\mathcal{E}$ *derives* $H(2u) = 3u$, $H(1) = 1$ *and* $H(3u) = 3H(u)$.

Proof. $H(2u) = \mathrm{sel}(N(2u), G(2u), 2u) = \mathrm{sel}(0, 3u, 2u) = 3u$ by Theorems 1.42 and 1.44. $H(1) = \mathrm{sel}(N(1), G(1), 1) = \mathrm{sel}(1, G(1), 1) = 1$. For the third equation the two-point rule gives, for variables a, c and b in the context $Z(Z(b)) = b$,

$$\mathrm{sel}(b,\ 3a + b,\ 3c) = 3\ \mathrm{sel}(b, a, c) :$$

at $b = 0$ both sides are $3a$, and at $b = 1$ both sides are $3c$. Instantiating at $b = N(u)$, $a = G(u)$ and $c = u$,

$$H(3u) = \mathrm{sel}\big(N(3u), G(3u), 3u\big) = \mathrm{sel}\big(N(u),\ 3G(u)+N(u),\ 3u\big) = 3\ \mathrm{sel}\big(N(u), G(u), u\big) = 3H(u).$$

The identity is proved with an independent variable b and then instantiated; the dependence of $N(u)$ on u causes no difficulty. □

Lemma 1.46 (Orbits of the factor map)**.** $\mathcal{E}$ *derives* $I(y, 3a;\, u.\, H(u)) = 3\, I(y, a;\, u.\, H(u))$ *and* $I(y, 1;\, u.\, H(u)) = 1$.

Proof. Both are equation inductions on y with a fixed. The bases are I0. The steps are $I(Sy, 3a;\, u.\, H(u)) = H(I(y, 3a;\, u.\, H(u))) = H(3\, I(y, a;\, u.\, H(u))) = 3H(I(y, a;\, u.\, H(u))) = 3\, I(Sy, a;\, u.\, H(u))$ and $I(Sy, 1;\, u.\, H(u)) = H(I(y, 1;\, u.\, H(u))) = H(1) = 1$, by Theorem 1.45. □

Lemma 1.47 (Comparator)**.** $\mathcal{E}$ *derives*

$$L(0, y) = 1, \qquad L(Sx, 0) = 0, \qquad L(Sx, Sy) = L(x, y), \qquad L(Sn, n) = 0.$$

Proof. $2^0 = 1$ and $2^{Sx} = 2 \cdot 2^x$ are computations. Hence $L(0,y) = N(I(y,1;\, u.\, H(u))) = N(1) = 1$ by Theorem 1.46, and $L(Sx,0) = N(2 \cdot 2^x) = 0$ by I0 and Theorem 1.42. For the third equation,

$$\begin{aligned} L(Sx,Sy) &= N\big(I(Sy, 2\cdot 2^x;\, u.\, H(u))\big) = N\big(I(y, H(2\cdot 2^x);\, u.\, H(u))\big) \\ &= N\big(I(y, 3\cdot 2^x;\, u.\, H(u))\big) = N\big(3\, I(y, 2^x;\, u.\, H(u))\big) = N\big(I(y, 2^x;\, u.\, H(u))\big) = L(x,y), \end{aligned}$$

by rotation (Theorem 1.39), $H(2u) = 3u$, Theorem 1.46 and $N(3n) = N(n)$. The fourth is an equation induction: $L(S0,0) = 0$ by the second equation, and $L(SSn,Sn) = L(Sn,n)$ by the third. □

On the natural numbers $L(x,y)$ is the indicator of $x \le y$. That reading is not used: only the four equations are.

1.7.6 The saturating counter and the predecessor

Put

$$\beta(k,u) := \mathrm{sel}\big(L(SSu,k),\, u,\, Su\big), \qquad W(t,k) := I(t,0;\, u.\, \beta(k,u)), \qquad P(m) := W(m,m).$$

The counter $W(t,k)$ counts up to t and stops below k. Written out,

$$P(m) = I(m,0;\, u.\, \mathrm{sel}\big(L(SSu,m),\, u,\, Su\big)),$$

a term in the one variable m, which occurs both as the counter and in the body.

Theorem 1.48 (Predecessor)**.** *The equational fragment derives, in the empty context,*

$$\vdash_{\mathcal{E}} P(0) = 0, \qquad \vdash_{\mathcal{E}} P(Sx) = x.$$

Consequently $Sx = Sy \vdash_{\mathcal{E}} x = y$, and the same sequents are derivable in N *and in* T_{eq}.

Proof. By I0 and IS,

$$W(0,k) = 0, \qquad W(St,k) = \beta(k, W(t,k)). \tag{1.10}$$

By Theorems 1.41 and 1.47(ii),

$$\beta(Sk,Su) = \mathrm{sel}\big(L(SSSu,Sk), Su, SSu\big) = \mathrm{sel}\big(L(SSu,k), Su, SSu\big) = S\beta(k,u), \tag{1.11}$$

$$\beta(SSk,0) = \mathrm{sel}\big(L(SS0,SSk), 0, 1\big) = \mathrm{sel}\big(L(0,k), 0, 1\big) = \mathrm{sel}(1,0,1) = 1. \tag{1.12}$$

Conjugacy. $W(St,SSk) = SW(t,Sk)$, by EI on t with k fixed. The base is $W(S0,SSk) = \beta(SSk, W(0,SSk)) = \beta(SSk,0) = 1 = SW(0,Sk)$, by (1.10) and (1.12). The step is

$$\begin{aligned} W(SSt,SSk) &= \beta(SSk,\ W(St,SSk)) = \beta(SSk,\ SW(t,Sk)) \\ &= S\beta(Sk,\ W(t,Sk)) = SW(St,Sk), \end{aligned}$$

where the second step uses the induction equation and the third is (1.11).

Below its cap the counter does not saturate. $W(n,Sn) = n$, by EI on n. The base is $W(0,S0) = 0$. In the step, $W(Sn,SSn) = SW(n,Sn) = Sn$, by the conjugacy instantiated at $t = k = n$ and the induction equation. The instantiation is made in an identity that is already derived with independent variables; no parameter of an active induction hypothesis is changed.

The two equations. $P(0) = W(0,0) = 0$. Substituting Sx for both variables in (1.10),

$$\begin{aligned} P(Sx) = W(Sx,Sx) &= \beta(Sx,\ W(x,Sx)) = \beta(Sx,x) \\ &= \mathrm{sel}\big(L(SSx,Sx),\, x,\, Sx\big) = \mathrm{sel}\big(L(Sx,x),\, x,\, Sx\big) = \mathrm{sel}(0,x,Sx) = x, \end{aligned}$$

by Theorem 1.47. Finally, from $Sx = Sy$ congruence gives $P(Sx) = P(Sy)$, hence $x = y$. The transfer to N and T_{eq} is Theorem 1.37(iv). □

Remark 1.49 (Sources and the use of coverage in the inherited proof). The construction (parity, three-cycle, ramp, factor map, comparator, saturating counter) is the scalar predecessor of the reference statement filed with `certified_zero`. That proof works in the theory with numerical cases. It uses them to show that $Z(n)$ and $N(n)$ take the values 0 and 1 only, and then splits the value $N(n)$ under the induction hypothesis in the ramp invariant Theorem 1.44(i), again in Theorem 1.44(iv), and in $H(3u) = 3H(u)$; it also proves $\mathrm{sel}(c, Sa, Sb) = S\,\mathrm{sel}(c, a, b)$ by cases, which is an equational case. The splits are replaced here by the equation $Z(Z(N(n))) = N(n)$ and the two-point rule, applied to identities that are derived in advance with independent variables, which is the contribution of `uniform_raw`, `newest_parameter`, `parameter_witness` and `no_witness`. The four notes differ in presentation. `uniform_raw` states the interpolation as $F(Z(c)) = \mathrm{sel}(c, F(1), F(0))$, as above; `no_witness` states it as $F(Z(Z(y))) = \mathrm{sel}(y, F(0), F(1))$; `parameter_witness` uses a finite-value transfer lemma for $Z(t)$ and $N(t)$ proved by equational cases; `newest_parameter` instantiates two identities in $Z(v)$ at $v = Z(N(n))$ and replaces the split in $H(3u)$ by $\mathrm{sel}(c, a + b, a' + b') = \mathrm{sel}(c, a, a') + \mathrm{sel}(c, b, b')$ and $\mathrm{sel}(c, c, 0) = 0$. The reference defines the counter with the body $\mathrm{sel}(L(Su, m), u, Su)$ and then identifies the body of $P(Sm)$ with it by a separate induction; the form with $L(SSu, k)$ and the conjugacy between the caps SSk and Sk, taken from `no_witness`, avoids that induction. The stronger statement matters where a calculus is closed under substitution only on its equational part, or restricts the witnesses of DI: the fragment $\mathcal{E}$ has no rule for D, so the predecessor is available before any naming of terms.

Both forms are formalized in Lean. The fragment $\mathcal{E}$ is the calculus `EqPrf` of the modules `EqKernel`, `EqArith`, `EqParity`, `EqCompare` and `EqPred`; it has exactly the five rules of Theorem 1.36: hypothesis, the two computations, transport in one term template, and induction for a literal equation with parameters. Theorem 1.48 is `EqPrf.pred_zero`, `EqPrf.pred_succ` (in the form `EqPrf Γ (Tm.pred (succ t)) t`, for every term and every context) and `EqPrf.succ_inj`. The calculus embeds in the kernel under any restriction of the witness of DI (`EqPrf.toDW`), because its derivations contain no introduction of D. The older modules `TeqParity`, `TeqCompare` and `TeqPred` prove the two equations in T_{eq} and split the Boolean value by coverage.

1.7.7 Successor laws, subtraction and the reflection of equality

From here on the derivations are in T_{eq} and use coverage and separation. "By cases on r" means the elimination of the disjunction (T2) at r followed by the Leibniz rule; unlike equational cases it may be applied under hypotheses that mention r.

Lemma 1.50 (Successor laws). T_{eq} *proves* $Sx = Sy \to x = y$, $Sx = 0 \to \bot$ *and* $x = 0 \vee x = SP(x)$.

Proof. The first is Theorem 1.48. From $Sx = 0$ congruence gives $Z(Sx) = Z(0)$, that is $0 = S0$, and (T3) gives $\bot$. For the third, eliminate (T2): if $x = 0$ the first disjunct holds, and if $x = Sy$ then $P(x) = y$ and $x = SP(x)$. □

Put $x \dot{-} y := I(y, x;\, u.\, P(u))$ and $K(u) := \mathrm{sel}(u, 1, P(u))$, and define the *equality test*

$$\varepsilon(a, b) := K(Sa \dot{-} b).$$

Lemma 1.51 (Truncated subtraction). T_{eq} *proves*

(i) $x \dot{-} 0 = x$, $x \dot{-} Sy = P(x \dot{-} y)$ *and* $Sx \dot{-} Sy = x \dot{-} y$;

(ii) $Sa \dot{-} a = 1$;

(iii) $\mathrm{sel}(x \dot{-} y,\ x,\ y + (x \dot{-} y)) = x$;

(iv) $Sa \dot{-} b = 1 \leftrightarrow a = b$.

Proof. (i) The first two are computations. The equation $P(Sx \dot{-} y) = x \dot{-} y$ holds by induction on y: the base is $P(Sx) = x$, and $P(Sx \dot{-} Sy) = P(P(Sx \dot{-} y)) = P(x \dot{-} y) = x \dot{-} Sy$. Its left side is $Sx \dot{-} Sy$.

(ii) Induction on a: $S0 \dot{-} 0 = 1$, and $SSa \dot{-} Sa = Sa \dot{-} a$ by (i).

(iii) Induction on y with x fixed. The base is $\mathrm{sel}(x, x, 0 + x) = \mathrm{sel}(x, x, x) = x$, by Theorem 1.41(ii). For the step write $r := x \dot{-} y$; the induction equation is $\mathrm{sel}(r, x, y + r) = x$ and the goal is $\mathrm{sel}(P(r), x, Sy + P(r)) = x$. Argue by cases on r. If $r = 0$, then $P(r) = 0$ and the goal is $\mathrm{sel}(0, x, \dots) = x$. If $r = Sz$, the induction equation gives $y + Sz = x$, and $P(r) = z$, so $Sy + P(r) = Sy + z = y + Sz = x$ and the goal is $\mathrm{sel}(z, x, x) = x$.

(iv) If $Sa \dot{-} b = 1$, then (iii) at $x = Sa$, $y = b$ gives $\mathrm{sel}(1, Sa, b + 1) = Sa$, that is $Sb = Sa$, and Theorem 1.50 gives $a = b$. The converse is (ii). □

The equation (iii) says, in one literal equation, that a nonzero difference is an actual difference.

Theorem 1.52 (Reflection of equality)**.** T_{eq} *proves, for all terms a and b,*

$$\varepsilon(a, b) = 0 \;\leftrightarrow\; a = b.$$

Consequently $\mathsf{T}_{\mathrm{eq}} \vdash a = b \vee (a = b \to \bot)$.

Proof. First, $K(u) = 0 \leftrightarrow u = 1$. If $u = S0$ then $K(u) = P(S0) = 0$. Conversely assume $K(u) = 0$ and argue by cases on u. If $u = 0$ then $K(u) = 1$, so $S0 = 0$, which is contradictory by Theorem 1.50. If $u = Sz$ then $K(u) = P(Sz) = z$, so $z = 0$ and $u = S0$. Hence $\varepsilon(a, b) = 0$ is equivalent to $Sa \dot{-} b = 1$, and to $a = b$ by Theorem 1.51(iv). For the consequence argue by cases on $\varepsilon(a, b)$: if it is 0 then $a = b$; if it is Sz, then $a = b$ would give $Sz = 0$. □

The theorem concerns the equality of the logic on the whole domain. It is not restricted to numerals, and no faithful comparison symbol is assumed.

Remark 1.53 (Variants). The reference statement uses the sign $\mathrm{sg}(u) := \mathrm{sel}(u, 0, 1)$, the invariant $\mathrm{sel}(\mathrm{sg}(x \dot{-} y), x, y + (x \dot{-} y)) = x$ and the test $\mathrm{sg}(K(Sa \dot{-} b))$, which takes the values 0 and 1 only, and it carries the disjunction "$b_\varphi = 0 \vee b_\varphi = 1$" through the construction below. The form above, which is the one of the Lean modules `TeqEq` and `TeqReflect`, omits the sign: the selector tests its first argument for zero, so a two-valued characteristic is not needed.

1.7.8 Induction for quantifier-free formulas

Definition 1.54 (Characteristic term)**.** For a quantifier-free ordinary formula φ the term χ_φ is

$$\chi_{s=t} := \varepsilon(s, t), \qquad \chi_\bot := 1, \qquad \chi_{\varphi\wedge\psi} := \mathrm{sel}(\chi_\varphi, \chi_\psi, 1),$$
$$\chi_{\varphi\vee\psi} := \mathrm{sel}(\chi_\varphi, 0, \chi_\psi), \qquad \chi_{\varphi\to\psi} := \mathrm{sel}(\chi_\varphi, \chi_\psi, 0).$$

The definition is a recursion on finite syntax, and $\chi_{\varphi[\theta]} = \chi_\varphi[\theta]$.

Proposition 1.55 (Reflection of quantifier-free formulas)**.** *For every quantifier-free φ,* $\mathsf{T}_{\mathrm{eq}} \vdash \chi_\varphi = 0 \leftrightarrow \varphi$.

Proof. By induction on φ. An equation is Theorem 1.52. For $\bot$: $S0 = 0$ implies $\bot$ by Theorem 1.50, and $\bot$ implies everything.

Conjunction. Assume $\mathrm{sel}(\chi_\varphi, \chi_\psi, 1) = 0$ and argue by cases on χ_φ. If $\chi_\varphi = 0$ the hypothesis is $\chi_\psi = 0$, and the induction hypotheses give φ and ψ. If χ_φ is a successor the hypothesis is

$1 = 0$, a contradiction. Conversely, φ and ψ give $\chi_\varphi = 0$ and $\chi_\psi = 0$, hence $\mathrm{sel}(0, \chi_\psi, 1) = \chi_\psi = 0$.

Disjunction. Assume $\mathrm{sel}(\chi_\varphi, 0, \chi_\psi) = 0$ and argue by cases on χ_φ. If it is 0, then φ. If it is a successor, the hypothesis is $\chi_\psi = 0$, hence ψ. Conversely, φ gives $\mathrm{sel}(0, 0, \chi_\psi) = 0$, and ψ gives $\mathrm{sel}(\chi_\varphi, 0, 0) = 0$ by Theorem 1.41(ii).

Implication. Assume $\mathrm{sel}(\chi_\varphi, \chi_\psi, 0) = 0$ and φ. Then $\chi_\varphi = 0$, the hypothesis is $\chi_\psi = 0$, and ψ follows. Conversely assume $\varphi \to \psi$ and argue by cases on χ_φ. If it is 0, then φ, hence ψ, hence $\chi_\psi = 0$ and $\mathrm{sel}(0, \chi_\psi, 0) = 0$. If it is a successor, the selector is 0. □

No excluded middle for arbitrary formulas is used: the case distinctions are on terms.

Theorem 1.56 (Quantifier-free induction)**.** *For every quantifier-free $\varphi(z)$,*

$$\mathsf{T}_{\mathrm{eq}} \vdash \varphi(0) \to (\forall z\,(\varphi(z) \to \varphi(Sz)) \to \forall z\,\varphi(z)).$$

Equivalently, if $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} \varphi(0)$ and $\Gamma, \varphi(z) \vdash_{\mathsf{T}_{\mathrm{eq}}} \varphi(Sz)$ with z not free in Γ, then $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} \varphi(t)$ for every t. The context Γ may contain quantified formulas.

Proof. Assume the two antecedents. By Theorem 1.55 the first gives $\chi_\varphi(0) = 0$. Under the further assumption $\chi_\varphi(z) = 0$ the proposition gives $\varphi(z)$, the second antecedent gives $\varphi(Sz)$, and the proposition gives $\chi_\varphi(Sz) = 0$. Theorem 1.33(ii) for the equation $\chi_\varphi(z) = 0$ gives $\chi_\varphi(z) = 0$ for all z, and the proposition gives $\varphi(z)$. □

In Lean this is `Teq.qfInd`.

1.7.9 Order, powers of two and pairing

Write $x \le y$ for the equation $x \mathbin{\dot-} y = 0$.

Lemma 1.57 (Order)**.** T_{eq} *proves*

(i) $x \mathbin{\dot-} x = 0$, $(a + b) \mathbin{\dot-} a = b$, $a \mathbin{\dot-} (a + b) = 0$ *and* $S(x + d) \mathbin{\dot-} x = Sd$;

(ii) $\mathrm{sel}(x \mathbin{\dot-} y,\ x + (y \mathbin{\dot-} x),\ y) = y$;

(iii) $x \le y \to x + (y \mathbin{\dot-} x) = y$;

(iv) $x \le y \to x \le y + z$; $\quad Sx \le y \to x \le y$; $\quad x \le y \wedge y \le z \to x \le z$.

Proof. (i) By induction: $0 \mathbin{\dot-} 0 = 0$ and $Sx \mathbin{\dot-} Sx = x \mathbin{\dot-} x$; $(0 + b) \mathbin{\dot-} 0 = b$ and $(Sa + b) \mathbin{\dot-} Sa = S(a + b) \mathbin{\dot-} Sa = (a + b) \mathbin{\dot-} a$, by induction on a; $a \mathbin{\dot-} (a + 0) = a \mathbin{\dot-} a = 0$ and $a \mathbin{\dot-} (a + Sb) = P(a \mathbin{\dot-} (a + b)) = P(0) = 0$, by induction on b. The fourth equation is the second for $x + Sd$.

(ii) Induction on y with x fixed. The base $\mathrm{sel}(x, x + (0 \mathbin{\dot-} x), 0) = 0$ holds by cases on x: at 0 the left side is $0 + (0 \mathbin{\dot-} 0) = 0$, and at a successor it is 0. For the step write $r := x \mathbin{\dot-} y$; the induction equation is $\mathrm{sel}(r, x + (y \mathbin{\dot-} x), y) = y$ and the goal is

$$\mathrm{sel}(P(r),\ x + (Sy \mathbin{\dot-} x),\ Sy) = Sy.$$

Argue by cases on r. If $r = 0$, the induction equation gives $x + (y \mathbin{\dot-} x) = y$. By (i), $Sy \mathbin{\dot-} x = S(x + (y \mathbin{\dot-} x)) \mathbin{\dot-} x = S(y \mathbin{\dot-} x)$, so $x + (Sy \mathbin{\dot-} x) = S(x + (y \mathbin{\dot-} x)) = Sy$, which is the goal since $P(r) = 0$. If $r = Sz$, then $P(r) = z$ and Theorem 1.51(iii) gives $y + Sz = x$. Argue by cases on z. If $z = 0$, then $x = Sy$ and the goal is $Sy + (Sy \mathbin{\dot-} Sy) = Sy$, which holds by (i). If z is a successor, the selector returns Sy.

(iii) is (ii) with $x \mathbin{\dot-} y = 0$.

(iv) By (iii), $y + z = x + ((y \mathbin{\dot-} x) + z)$, and (i) applies. If $Sx \le y$, then $y = Sx + d = x + Sd$ with $d := y \mathbin{\dot-} Sx$, and (i) applies. If $x \le y$ and $y \le z$, then $z = y + (z \mathbin{\dot-} y)$ and the first part applies. □

Lemma 1.58 (Powers of two)**.** T_{eq} *proves* $2^0 = 1$, $2^{Sx} = 2^x + 2^x$, $2^x \le 2^{x+d}$, $x \le y \to 2^x \le 2^y$ *and* $Sx \le 2^x$.

Proof. The first two are computations. $2^x \le 2^{x+d}$ by induction on d: the base is Theorem 1.57(i), and $2^{x+Sd} = 2^{x+d} + 2^{x+d}$, so the step is Theorem 1.57(iv). Monotonicity follows by $y = x + (y \dot{-} x)$. $Sx \le 2^x$ by induction on x: the base is $1 \dot{-} 1 = 0$. In the step let $p := 2^x$ and $d := p \dot{-} Sx$, so that $p = Sx + d$ by the induction equation and Theorem 1.57(iii). Then $p + p = SSx + (x + d + d)$ by Theorem 1.40, and $SSx \le p + p$ by Theorem 1.57(i). □

Put

$$\ell(k, v) := \mathrm{sel}\big(2^{Sv} \dot{-} k,\ Sv,\ v\big), \qquad \Lambda(t, k) := I(t, 0;\, v.\, \ell(k, v)), \qquad \lg(w) := \Lambda(w, w),$$
$$\langle a, b\rangle := 2^{a+b} + a, \qquad \mathrm{fst}(w) := w \dot{-} 2^{\lg(w)}, \qquad \mathrm{snd}(w) := \lg(w) \dot{-} \mathrm{fst}(w).$$

The step $\ell(k, v)$ increments v while $2^{v+1} \le k$, so $\Lambda(t, k)$ is a saturating counter for the binary logarithm of k.

Theorem 1.59 (Pairing)**.** T_{eq} *proves* $\lg\langle a, b\rangle = a + b$, $\mathrm{fst}\langle a, b\rangle = a$ *and* $\mathrm{snd}\langle a, b\rangle = b$.

Proof. Fix a, b and put $c := a + b$ and $k := 2^c + a$.

Below the exponent the counter follows its index: $\mathrm{sel}(i \dot{-} c,\ \Lambda(i, k),\ i) = i$, by induction on i. The base is $\mathrm{sel}(0 \dot{-} c, 0, 0) = 0$. In the step argue by cases on $Si \dot{-} c$. If it is a successor, the goal $\mathrm{sel}(Si \dot{-} c, \Lambda(Si, k), Si) = Si$ is a computation. If it is 0, then $Si \le c$, hence $i \le c$ by Theorem 1.57(iv), and the induction equation gives $\Lambda(i, k) = i$. Moreover $2^{Si} \le 2^c \le k$ by Theorems 1.57 and 1.58, so $\Lambda(Si, k) = \ell(k, i) = \mathrm{sel}(0, Si, i) = Si$, which is the goal. At $i = c$ this gives

$$\Lambda(c, k) = c. \tag{1.13}$$

At the exponent the counter stops: $\ell(k, c) = c$. By Theorem 1.58, $2^c = Sc + d$ with $d := 2^c \dot{-} Sc$, hence $2^{Sc} = 2^c + 2^c = (2^c + a) + S(b + d)$ and $2^{Sc} \dot{-} k = S(b + d)$ by Theorem 1.57(i). The selector in $\ell(k, c)$ therefore returns c.

Beyond the exponent: $\Lambda(c + j, k) = c$, by induction on j. The base is (1.13), and $\Lambda(S(c + j), k) = \ell(k, \Lambda(c + j, k)) = \ell(k, c) = c$.

Conclusion. $Sc \le 2^c$ gives $c \le 2^c \le k$, so $k = c + (k \dot{-} c)$ and $\lg(k) = \Lambda(k, k) = c$. Then $\mathrm{fst}(k) = (2^c + a) \dot{-} 2^c = a$ and $\mathrm{snd}(k) = (a + b) \dot{-} a = b$ by Theorem 1.57(i). □

Surjectivity of the pairing is not needed and not claimed. The numerical domain is never replaced by the range of the coding; a pair only stores the state of one recursion.

Remark 1.60 (The pairing of the reference statement)*.* The reference statement filed with `certified_zero` uses the pairing $(a + b)^2 + a$, with multiplication $a \cdot b := I(b, 0;\, u.\, u + a)$ and a bounded search for the integer square root, $q(n) := I(n, 0;\, v.\, \mathrm{sel}((Sv)^2 \dot{-} n, Sv, v))$, whose invariants are proved by quantifier-free induction; it appeals to "the usual linear-order and semiring identities" without deriving them. The pairing above is the one of the Lean modules `TeqOrder` and `TeqPair`. It needs no multiplication, and all its inductions are on literal equations, so Theorem 1.56 is not used for it. Either pairing serves the next theorem.

1.7.10 Primitive recursion

Definition 1.61 (The derived recursor)**.** Let a be a term and $g(y, v)$ a term with two distinguished variables. With a fresh variable w put

$$s_g(w) := \big\langle S\,\mathrm{fst}(w),\ g(\mathrm{fst}(w), \mathrm{snd}(w))\big\rangle, \qquad R(c, a;\, y\, v.\, g) := \mathrm{snd}\big(I(c, \langle 0, a\rangle;\, w.\, s_g(w))\big).$$

The recursor binds y and v in g.

The iterator carries the pair of the counter and the value. For a function symbol f of iPRA of arity k the term $\ulcorner f \urcorner$, with free variables among the fixed variables $x_1, \dots, x_k$, is defined by recursion on f; $\ulcorner f \urcorner(t_1, \dots, t_k)$ abbreviates $\ulcorner f \urcorner[\vec{t}/\vec{x}]$.

$$\ulcorner \mathrm{zero} \urcorner := 0, \qquad \ulcorner \mathrm{succ} \urcorner := Sx_1, \qquad \ulcorner \pi_i^k \urcorner := x_i, \qquad \ulcorner \mathrm{comp}(f; \vec{g}) \urcorner := \ulcorner f \urcorner(\ulcorner g_1 \urcorner, \dots, \ulcorner g_m \urcorner),$$
$$\ulcorner \mathrm{prec}(f, g) \urcorner := R\big(x_1,\ \ulcorner f \urcorner(x_2, \dots, x_{k+1});\ y\, v. \ulcorner g \urcorner(y, v, x_2, \dots, x_{k+1})\big).$$

The translation σ of the terms of iPRA into the terms of T_{eq} is

$$\sigma(x) := x, \qquad \sigma\big(f(t_1, \dots, t_k)\big) := \ulcorner f \urcorner\big(\sigma(t_1), \dots, \sigma(t_k)\big).$$

It commutes with substitution literally, $\sigma(t[\theta]) = \sigma(t)[\sigma \circ \theta]$, and $\sigma(0) = 0$, $\sigma(St) = S\sigma(t)$.

Theorem 1.62 (Primitive recursion)**.** (i) T_{eq} *proves* $R(0, a;\, y\, v.\, g) = a$ *and* $R(Sc, a;\, y\, v.\, g) = g\big(c,\ R(c, a;\, y\, v.\, g)\big)$.

(ii) *For every defining equation* $s = t$ *of a function symbol of* iPRA *(Theorem 1.35),* $\mathsf{T}_{\mathrm{eq}} \vdash \sigma(s) = \sigma(t)$.

Proof. (i) Write $A(c) := I(c, \langle 0, a\rangle;\, w.\, s_g(w))$. By induction on c, $\mathrm{fst}(A(c)) = c$: the base is $\mathrm{fst}\langle 0, a\rangle = 0$, and $\mathrm{fst}(A(Sc)) = \mathrm{fst}(s_g(A(c))) = S\,\mathrm{fst}(A(c)) = Sc$, by Theorem 1.59. Then $R(0, a;\, y\, v.\, g) = \mathrm{snd}\langle 0, a\rangle = a$, and

$$R(Sc, a;\, y\, v.\, g) = \mathrm{snd}\big(s_g(A(c))\big) = g\big(\mathrm{fst}(A(c)), \mathrm{snd}(A(c))\big) = g\big(c,\ R(c, a;\, y\, v.\, g)\big).$$

(ii) For a projection, $\sigma(\pi_i^k(\vec{x}))$ is the term x_i. For a composition, $\sigma(\mathrm{comp}(f; \vec{g})(\vec{x}))$ and $\sigma(f(g_1(\vec{x}), \dots, g_m(\vec{x})))$ are the same term. For a recursion the two equations are (i) for the seed $\ulcorner f \urcorner(\vec{x})$ and the step $\ulcorner g \urcorner(y, v, \vec{x})$. □

The construction is an external recursion on each finite description. It is not an internal quantification over functions, and no universal evaluator is introduced. In Lean the recursor is the content of the module `TeqRec`.

1.8 The bridge

This section draws on the reference statements filed with `certified_zero` (the inherited basis, the inherited ordinary bridge and the section on primitive recursion) and follows the Lean modules `NativeTranslate`, `NativeComposite`, `NativeRoundTrip`, `ScalarToPRA`, `PRAToScalar`, `RoundTripScalar` and `RoundTripPRA` in the organization of the proofs. The bridge has two spans: between N and T_{eq} the terms are the same and the formulas are translated; between T_{eq} and iPRA the logic is the same and the terms are translated.

1.8.1 The native core and T_{eq}

Definition 1.63 (Native reading of an ordinary formula)**.** For an ordinary formula φ over the terms of Theorem 1.1, the native formula φ^* replaces every connective by its native abbreviation (Theorem 1.28):

$$(s = t)^* := (s = t), \qquad \bot^* := \bot_*, \qquad (\varphi \wedge \psi)^* := \varphi^* \wedge_* \psi^*, \qquad (\varphi \vee \psi)^* := \varphi^* \vee_* \psi^*,$$
$$(\varphi \to \psi)^* := \varphi^* \Rightarrow_* \psi^*, \qquad (\forall x\, \varphi)^* := \forall_* x\, \varphi^*, \qquad (\exists x\, \varphi)^* := \exists_* x\, \varphi^*.$$

By (1.6), $(\varphi[\theta])^* = \varphi^*[\theta]$.

Proposition 1.64 (Soundness of the ordinary reading)**.** *If* $\Gamma \vdash_{\mathsf{N}} A$*, then* $\Gamma^\circ \vdash_{\mathsf{T}_{\mathrm{eq}}} A^\circ$.

Proof. By induction on the derivation, using (1.3) for the instances. Hyp is hypothesis; I0 and IS are (T1). Tr: from $s = t$ and $h(s) = r$ the equality rules give $r = h(t)$. QE: the premises read $B^\circ(0) \to \forall y\, B^\circ$ and $B^\circ(0)$, which give $B^\circ(t)$. DE: the first premise reads $\exists x\, (B^\circ(x,0) \to \forall y\, B^\circ(x,y))$ and the hypothesis $Q_y B(b,y)$ of the second reads $B^\circ(b,0) \to \forall y\, B^\circ(b,y)$, so the rule is existential elimination. DI: the premise reads $\Gamma^\circ, B^\circ(t,0) \vdash B^\circ(t,Sz)$. Over $\Gamma^\circ, B^\circ(t,0)$, Theorem 1.33(i) for the formula $B^\circ(t,y)$ gives $\forall y\, B^\circ(t,y)$; discharge $B^\circ(t,0)$ and introduce the existential quantifier at t. QI: the premise reads $\Gamma^\circ, E(z) \vdash E(Sz)$; over $\Gamma^\circ, E(0)$, Theorem 1.33(ii) gives $E(y)$ for a fresh y; generalize and discharge $E(0)$. □

Coverage is used for DI only and equation induction for QI only.

Proposition 1.65 (Completeness of the native reading)**.** *If* $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} \varphi$*, then* $\Gamma^* \vdash_{\mathsf{N}} \varphi^*$.

Proof. By induction on the derivation. Every logical rule of Theorem 1.31 for a connective translates into the rule of the same name in Theorem 1.29; the translation commutes with the substitutions in the quantifier rules. Reflexivity is Theorem 1.18(i), and the Leibniz rule is Theorem 1.22. It remains to derive the native reading of each axiom in the empty context; by Theorem 1.10 all substitution instances then follow in every context.

(T1) The readings are the conclusions of I0 and IS.

(T2) Apply Theorem 1.14 to the formula $C(y) := (y = 0) \vee_* \exists_* v\, (y = Sv)$. $C(0)$ follows from $0 = 0$ by $\vee_*$I. $C(Sz)$ follows from $Sz = Sz$ by $\exists_*$I with the witness z and $\vee_*$I. Hence $\vdash C(x)$.

(T3) The reading is $(0 = S0) \Rightarrow_* (0 = S0)$, which is $\Rightarrow_*$I applied to Hyp.

(T4) Let $U := \forall_* z\, (E(z) \Rightarrow_* E(Sz))$. By $\Rightarrow_*$I twice it suffices to derive $E(0), U \vdash E(0) \wedge_* Q_z E(z)$, the conclusion being $\forall_* z\, E(z)$ by definition. The first component is a hypothesis. For the second apply QI: over $E(0), U, E(z)$ with z fresh, $\forall_*$E gives $E(z) \Rightarrow_* E(Sz)$ and $\Rightarrow_*$E gives $E(Sz)$. □

Proposition 1.66 (The two formula round trips)**.** (i) *For every native formula* A*:* $(A^\circ)^* \dashv\vdash A$ *in* N.

(ii) *For every ordinary formula* φ*:* $\mathsf{T}_{\mathrm{eq}} \vdash \varphi \leftrightarrow (\varphi^*)^\circ$.

Proof. (i) By induction on A. An equation is its own round trip.

Let A be $Q_y B$ and put $C := (B^\circ)^*$, so that $C \dashv\vdash B$ by the induction hypothesis and $(A^\circ)^* = C(0) \Rightarrow_* \forall_* y\, C$. From this formula derive $Q_y B$ by Theorem 1.13: under the assumption $B(0)$ one has $C(0)$, hence $\forall_* y\, C$, hence $C(Sz)$ by $\forall_*$E, hence $B(Sz)$. Conversely, from $Q_y B$ and the assumption $C(0)$ one has $B(0)$, hence $B(y)$ for a fresh y by QE, hence $C(y)$; $\forall_*$I and $\Rightarrow_*$I give $(A^\circ)^*$.

Let A be $D_{x,y} B$. Then $(A^\circ)^* = \exists_* x\, Y$, where $Y := ((Q_y B)^\circ)^*$ and $Y \dashv\vdash Q_y B$ by the previous case, with x as a parameter. From $\exists_* x\, Y$: $\exists_*$E at a fresh b gives $Y(b)$, hence $Q_y B(b,y)$, and Theorem 1.15 gives $D_{x,y} B$. From $D_{x,y} B$: DE at a fresh b gives $Q_y B(b,y)$, hence $Y(b)$, and $\exists_*$I gives $\exists_* x\, Y$.

(ii) A derivation in N can be used inside T_{eq}: if $\Delta \vdash_{\mathsf{N}} A$ and $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} B^\circ$ for every $B \in \Delta$, then $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} A^\circ$, by Theorem 1.64 and cut. This is called *transfer* below. The following equivalences are provable in T_{eq} for all native formulas X, Y.

(a) $(\bot_*)^\circ \leftrightarrow \bot$: by (T3) and $\bot$E.

(b) $(X \wedge_* Y)^\circ \leftrightarrow X^\circ \wedge Y^\circ$: by transfer of $\wedge_*$E and of $\wedge_*$I.

(c) $(\exists_* x\, X)^\circ \leftrightarrow \exists x\, X^\circ$: the left side is $\exists x\, (Q_y \,\mathrm{gate}(y, X))^\circ$, and $(Q_y \,\mathrm{gate}(y, X))^\circ \leftrightarrow X^\circ$ by transfer of Theorem 1.26(iii).

(d) $(X \Rightarrow_* Y)^\circ \leftrightarrow (X^\circ \to Y^\circ)$: from left to right by transfer of $\Rightarrow_*$E. From right to left, the goal $(X \Rightarrow_* Y)^\circ$ is $\mathrm{sel}(0, X, Y)^\circ \to \forall y\, \mathrm{sel}(y, X, Y)^\circ$. Assume $\mathrm{sel}(0, X, Y)^\circ$. Transfer of Theorem 1.26(iv) gives X°, hence Y°. Transfer of $Y \vdash X \Rightarrow_* Y$ gives $(X \Rightarrow_* Y)^\circ$, and its antecedent is the assumption.

(e) $(X \vee_* Y)^\circ \leftrightarrow X^\circ \vee Y^\circ$: by (c) the left side is equivalent to $\exists x\, \mathrm{sel}(x, X, Y)^\circ$. Given $\mathrm{sel}(x, X, Y)^\circ$, eliminate (T2) at x. If $x = 0$, transfer of $x = 0,\ \mathrm{sel}(x, X, Y) \vdash X$ gives X°; if $x = Sz$, transfer of $x = Sz,\ \mathrm{sel}(x, X, Y) \vdash Y$ gives Y°. Both native sequents hold by Theorem 1.27(ii) and Theorem 1.26(iv), (v). The converse is transfer of $\vee_*$I.

(f) $(\forall_* x\, X)^\circ \leftrightarrow \forall x\, X^\circ$: transfer of $\wedge_*$E gives $X^\circ(0)$ and $(Q_x X)^\circ = X^\circ(0) \to \forall x\, X^\circ$ from the left side, hence the right side. Conversely $\forall x\, X^\circ$ gives $X^\circ(0)$ and $(Q_x X)^\circ$, and transfer of $\wedge_*$I gives the left side.

The claim follows by induction on φ, because provable equivalence is a congruence for the connectives and quantifiers of the logic. □

Corollary 1.67 (N and T_{eq} prove the same)**.** $\Gamma \vdash_{\mathsf{N}} A$ *if and only if* $\Gamma^\circ \vdash_{\mathsf{T}_{\mathrm{eq}}} A^\circ$*, and* $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} \varphi$ *if and only if* $\Gamma^* \vdash_{\mathsf{N}} \varphi^*$.

Proof. If $\Gamma^\circ \vdash_{\mathsf{T}_{\mathrm{eq}}} A^\circ$, then $(\Gamma^\circ)^* \vdash_{\mathsf{N}} (A^\circ)^*$ by Theorem 1.65, and Theorem 1.66(i) with cut gives $\Gamma \vdash_{\mathsf{N}} A$. If $\Gamma^* \vdash_{\mathsf{N}} \varphi^*$, then $(\Gamma^*)^\circ \vdash_{\mathsf{T}_{\mathrm{eq}}} (\varphi^*)^\circ$ by Theorem 1.64, and Theorem 1.66(ii) gives $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} \varphi$. The other two implications are the two propositions. □

In Lean these results are `NPrf.sound`, `NPrf.ofPrf`, `NEquiv.ofFm_toFm`, `Teq.iff_toFm_ofFm`, `NPrf.iff_teq` and `Teq.iff_native`.

1.8.2 Term interpretations

Definition 1.68 (Term interpretation)**.** Let T and T' be theories in intuitionistic logic with equality over term languages $\mathcal{L}$ and $\mathcal{L}'$. A *term interpretation* of T in T' is a map F from the terms of $\mathcal{L}$ to the terms of $\mathcal{L}'$ such that

(i) $F(x) = x$ for every variable x, and $\mathrm{FV}(F(t)) \subseteq \mathrm{FV}(t)$;

(ii) $T' \vdash F(t[\theta]) = F(t)[F \circ \theta]$ for every term t and substitution θ;

(iii) $T' \vdash \varphi^F$ for every axiom φ of T,

where φ^F is φ with each equation $s = t$ replaced by $F(s) = F(t)$.

A term interpretation does not relativize quantifiers and reads equality as equality: its domain is the whole domain.

Lemma 1.69 (Term interpretations preserve derivability)**.** *Let F be a term interpretation of T in T'.*

(i) $T' \vdash (\varphi[\theta])^F \leftrightarrow \varphi^F[F \circ \theta]$.

(ii) *If* $\Gamma \vdash_T \varphi$*, then* $\Gamma^F \vdash_{T'} \varphi^F$.

Proof. (i) By induction on φ. For an equation the two sides are $F(s[\theta]) = F(t[\theta])$ and $F(s)[F \circ \theta] = F(t)[F \circ \theta]$, which are equivalent by condition (ii) of the definition. The connectives and quantifiers are handled by the congruence of provable equivalence, the bound variables being fixed by θ and by F. (ii) By induction on the derivation. The propositional rules, $\forall$I and $\exists$E translate literally. For $\forall$E the premise gives $\forall x\, \varphi^F$, hence $\varphi^F[F(t)/x]$, which is equivalent to $(\varphi[t/x])^F$ by (i); $\exists$I is similar. Reflexivity is reflexivity. For the Leibniz rule

the premises give $F(s_i) = F(t_i)$ and, by (i), $\varphi^F[F(\vec{s})/\vec{x}]$; the Leibniz rule of T' and (i) give $(\varphi[\vec{t}/\vec{x}])^F$. An instance $\varphi[\theta]$ of an axiom translates to a formula equivalent to $\varphi^F[F \circ \theta]$, an instance of a theorem of T'. □

Lemma 1.70 (Round trips give faithfulness)**.** *Let F be a term interpretation of T in T' and G one of T' in T, with $T \vdash G(F(t)) = t$ and $T' \vdash F(G(t')) = t'$ for all terms. Then*

(i) *$T \vdash \varphi \leftrightarrow (\varphi^F)^G$ and $T' \vdash \psi \leftrightarrow (\psi^G)^F$;*

(ii) *$\Gamma \vdash_T \varphi$ if and only if $\Gamma^F \vdash_{T'} \varphi^F$, and $\Gamma \vdash_{T'} \psi$ if and only if $\Gamma^G \vdash_T \psi^G$.*

Proof. (i) By induction on the formula; for an equation $s = t$ the round trip is $G(F(s)) = G(F(t))$, equivalent to $s = t$ by the hypothesis. (ii) One direction is Theorem 1.69. For the other, if $\Gamma^F \vdash_{T'} \varphi^F$ then $(\Gamma^F)^G \vdash_T (\varphi^F)^G$, and (i) gives $\Gamma \vdash_T \varphi$. □

1.8.3 From T_{eq} to iPRA

A term of iPRA has no binder. An iterator term is therefore compiled into a function symbol of its free variables, and the iterator becomes a primitive recursion whose step ignores the counter. For a symbol A of arity k and a symbol H of arity $k+1$ put

$$\iota(A, H) := \mathrm{prec}\big(A,\ \mathrm{comp}(H; \pi_2^{k+2}, \pi_3^{k+2}, \ldots, \pi_{k+2}^{k+2})\big),$$

a symbol of arity $k+1$. By the axioms for recursion, composition and projection,

$$\mathsf{iPRA} \vdash \iota(A,H)(0,\vec{x}) = A(\vec{x}), \qquad \mathsf{iPRA} \vdash \iota(A,H)(Sy,\vec{x}) = H\big(\iota(A,H)(y,\vec{x}),\vec{x}\big). \tag{1.14}$$

Definition 1.71 (The translation τ)**.** Fix an enumeration of the variables. For a list $\vec{x} = x_1, \ldots, x_k$ of distinct variables and a term t with $\mathrm{FV}(t) \subseteq \vec{x}$, the symbol $[t]_{\vec{x}}$ of arity k is

$$[x_i]_{\vec{x}} := \pi_i^k, \qquad [0]_{\vec{x}} := \mathrm{comp}(\mathrm{zero};\ ), \qquad [St]_{\vec{x}} := \mathrm{comp}(\mathrm{succ}; [t]_{\vec{x}}),$$
$$[I(c,a;\ u.\,h)]_{\vec{x}} := \mathrm{comp}\big(\iota([a]_{\vec{x}}, [h]_{u\vec{x}});\ [c]_{\vec{x}}, \pi_1^k, \ldots, \pi_k^k\big) \qquad (u \notin \vec{x}).$$

The translation of terms is

$$\tau(x) := x, \qquad \tau(0) := 0, \qquad \tau(St) := S\tau(t), \qquad \tau(I(c,a;\ u.\,h)) := \iota\big([a]_{\vec{y}}, [h]_{u\vec{y}}\big)\big(\tau(c), \vec{y}\big),$$

where $\vec{y}$ lists the variables of $\mathrm{FV}(a) \cup (\mathrm{FV}(h) \setminus \{u\})$ in the order of the enumeration.

The translation does not commute with substitution literally: a substitution into the body of an iterator changes the symbol. It commutes provably.

Lemma 1.72 (Extensionality of the simulated iterator)**.** *Let $\vec{z}$ be a list of distinct variables, $\vec{s}$ a list of terms, and u a variable that is not in $\vec{z}$ and not free in $\vec{s}$. If $\mathsf{iPRA} \vdash A'(\vec{z}) = A(\vec{s})$ and $\mathsf{iPRA} \vdash H'(u,\vec{z}) = H(u,\vec{s})$, then $\mathsf{iPRA} \vdash \iota(A',H')(t,\vec{z}) = \iota(A,H)(t,\vec{s})$ for every term t.*

Proof. By quantifier-free induction on a fresh variable n in the equation $\iota(A',H')(n,\vec{z}) = \iota(A,H)(n,\vec{s})$. The base is $A'(\vec{z}) = A(\vec{s})$ by (1.14). In the step,

$$\begin{aligned}\iota(A',H')(Sn,\vec{z}) &= H'\big(\iota(A',H')(n,\vec{z}),\vec{z}\big) = H\big(\iota(A',H')(n,\vec{z}),\vec{s}\big)\\ &= H\big(\iota(A,H)(n,\vec{s}),\vec{s}\big) = \iota(A,H)(Sn,\vec{s}),\end{aligned}$$

by (1.14), the second hypothesis instantiated at $u = \iota(A',H')(n,\vec{z})$, and the induction equation. Instantiate n at t. □

Lemma 1.73 (Evaluation and substitution)**.** (i) *If $\mathrm{FV}(t) \subseteq \vec{x}$, then $\mathsf{iPRA} \vdash [t]_{\vec{x}}(\vec{r}) = \tau(t)[\vec{r}/\vec{x}]$ for all terms $\vec{r}$. In particular $\mathsf{iPRA} \vdash [t]_{\vec{x}}(\vec{x}) = \tau(t)$.*

(ii) $\mathsf{iPRA} \vdash \tau(t[\theta]) = \tau(t)[\tau \circ \theta]$ *for every substitution* θ.

Proof. (i) By induction on t, for all lists $\vec{x}$. For a variable, 0 and a successor the claim follows from the axioms for projection and composition and the induction hypothesis. Let t be $I(c, a;\, u.\, h)$, and let $\vec{y}$ be the list of Theorem 1.71, a sublist of $\vec{x}$. By the induction hypothesis for a and h with variables as arguments, $[a]_{\vec{x}}(\vec{x}) = \tau(a) = [a]_{\vec{y}}(\vec{y})$ and $[h]_{u\vec{x}}(u, \vec{x}) = \tau(h) = [h]_{u\vec{y}}(u, \vec{y})$. By Theorem 1.72,

$$\iota\big([a]_{\vec{x}}, [h]_{u\vec{x}}\big)(n, \vec{x}) = \iota\big([a]_{\vec{y}}, [h]_{u\vec{y}}\big)(n, \vec{y})$$

for a fresh variable n. By the axioms for composition and projection and the induction hypothesis for c, $[t]_{\vec{x}}(\vec{r}) = \iota([a]_{\vec{x}}, [h]_{u\vec{x}})(\tau(c)[\vec{r}/\vec{x}], \vec{r})$. Substituting $\vec{r}$ for $\vec{x}$ and $\tau(c)[\vec{r}/\vec{x}]$ for n in the displayed equation identifies this term with $\tau(t)[\vec{r}/\vec{x}]$.

(ii) By induction on t. The cases other than the iterator are immediate. Let t be $I(c, a;\, u.\, h)$ with u fixed by θ and not free in its values, so that $t[\theta] = I(c[\theta], a[\theta];\, u.\, h[\theta])$. Let $\vec{y}$ be the list for a, h and $\vec{z}$ the list for $a[\theta]$, $h[\theta]$. By (i) and the induction hypothesis,

$$[a[\theta]]_{\vec{z}}(\vec{z}) = \tau(a[\theta]) = \tau(a)[\tau \circ \theta] = [a]_{\vec{y}}(\tau(\theta(\vec{y}))),$$
$$[h[\theta]]_{u\vec{z}}(u, \vec{z}) = \tau(h[\theta]) = \tau(h)[\tau \circ \theta] = [h]_{u\vec{y}}(u, \tau(\theta(\vec{y}))).$$

Theorem 1.72 with $\vec{s} := \tau(\theta(\vec{y}))$ and the induction hypothesis for c give

$$\begin{aligned}\tau(t[\theta]) &= \iota\big([a[\theta]]_{\vec{z}}, [h[\theta]]_{u\vec{z}}\big)\big(\tau(c[\theta]), \vec{z}\big)\\ &= \iota\big([a]_{\vec{y}}, [h]_{u\vec{y}}\big)\big(\tau(c)[\tau \circ \theta], \tau(\theta(\vec{y}))\big) = \tau(t)[\tau \circ \theta].\end{aligned}$$

□

Proposition 1.74 (T_{eq} in iPRA)**.** τ *is a term interpretation of* T_{eq} *in* iPRA.

Proof. Conditions (i) and (ii) of Theorem 1.68 are the definition and Theorem 1.73(ii). The axioms:

(T1) With the notation of Theorem 1.71, $\tau(I(0, a;\, u.\, h)) = \iota(\dots)(0, \vec{y}) = [a]_{\vec{y}}(\vec{y}) = \tau(a)$ by (1.14) and Theorem 1.73(i). Writing $q := \tau(I(n, a;\, u.\, h))$,

$$\tau(I(Sn, a;\, u.\, h)) = \iota(\dots)(S\tau(n), \vec{y}) = [h]_{u\vec{y}}(q, \vec{y}) = \tau(h)[q/u] = \tau\big(h[I(n, a;\, u.\, h)/u]\big),$$

by (1.14) and both parts of Theorem 1.73.

(T2) Let $\mathrm{pd} := \mathrm{prec}(\mathrm{zero}, \pi_1^2)$, so that $\mathrm{pd}(0) = 0$ and $\mathrm{pd}(Sy) = y$. Quantifier-free induction on x proves $x = 0 \vee x = S\mathrm{pd}(x)$: the base is the left disjunct, and the step is the right disjunct, since $Sx = S\mathrm{pd}(Sx)$. Existential introduction gives the image of (T2).

(T3) is the axiom $Sx = 0 \to \bot$ at $x = 0$, with symmetry.

(T4) For an equation $E(z)$, written $s = t$, the image of the axiom is

$$\tau(s[0/z]) = \tau(t[0/z]) \to \Big(\forall z\,(\tau(s) = \tau(t) \to \tau(s[Sz/z]) = \tau(t[Sz/z])) \to \forall z\,\tau(s) = \tau(t)\Big),$$

which by Theorem 1.73(ii) is equivalent to the induction axiom of iPRA for the equation $\tau(s) = \tau(t)$. □

1.8.4 From iPRA to T_{eq}

Proposition 1.75 (iPRA in T_{eq})**.** *The translation* σ *of Section 1.7 is a term interpretation of* iPRA *in* T_{eq}.

Proof. σ fixes variables and commutes with substitution literally. The defining equations hold by Theorem 1.62(ii). The images of $Sx = 0 \to \bot$ and $Sx = Sy \to x = y$ are those formulas themselves, which hold by Theorem 1.50. For a quantifier-free φ the formula φ^σ is quantifier free and $(\varphi[t/z])^\sigma = \varphi^\sigma[\sigma(t)/z]$, so the image of the induction axiom for φ is the instance of Theorem 1.56 for φ^σ. □

1.8.5 The two round trips on terms

Lemma 1.76 (A recursion that ignores its counter is the iterator)**.** *If y is not free in $h(v)$, then $\mathsf{T}_{\mathrm{eq}} \vdash R(c,a;\, y\, v.\, h(v)) = I(c,a;\, u.\, h(u))$.*

Proof. Theorem 1.38 for $f(z) := R(z,a;\, y\, v.\, h(v))$: by Theorem 1.62(i), $f(0) = a$ and $f(Sz) = h(f(z))$. □

Proposition 1.77 (Round trips on terms)**.** (i) *$\mathsf{T}_{\mathrm{eq}} \vdash \sigma(\tau(t)) = t$ for every term t of T_{eq}.*

(ii) *$\mathsf{iPRA} \vdash \tau(\sigma(t)) = t$ for every term t of iPRA.*

Proof. (i) By the definition of $\ulcorner\cdot\urcorner$ the term $\ulcorner \mathrm{comp}(H; \pi_2^{k+2}, \ldots, \pi_{k+2}^{k+2})\urcorner(y, v, \vec{x})$ is $\ulcorner H\urcorner(v, \vec{x})$, so by Theorem 1.76

$$\mathsf{T}_{\mathrm{eq}} \vdash \ulcorner \iota(A,H)\urcorner(c,\vec{x}) = I(c, \ulcorner A\urcorner(\vec{x});\, u.\, \ulcorner H\urcorner(u,\vec{x})). \tag{1.15}$$

First, $\mathsf{T}_{\mathrm{eq}} \vdash \ulcorner [t]_{\vec{x}}\urcorner(\vec{x}) = t$, by induction on t. For a variable, 0 and a successor both sides are the same term up to the induction hypothesis. For an iterator,

$$\begin{aligned}\ulcorner [I(c,a;\, u.\, h)]_{\vec{x}}\urcorner(\vec{x}) &= \ulcorner \iota([a]_{\vec{x}}, [h]_{u\vec{x}})\urcorner(\ulcorner [c]_{\vec{x}}\urcorner(\vec{x}), \vec{x}) \\ &= I(\ulcorner [c]_{\vec{x}}\urcorner(\vec{x}), \ulcorner [a]_{\vec{x}}\urcorner(\vec{x});\, u.\, \ulcorner [h]_{u\vec{x}}\urcorner(u,\vec{x})),\end{aligned}$$

by (1.15), and the last term equals $I(c,a;\, u.\, h)$ by the induction hypothesis, congruence in the counter and the seed, and Theorem 1.39(ii) for the body. Now the claim follows by induction on t; for an iterator, with the notation of Theorem 1.71,

$$\begin{aligned}\sigma(\tau(I(c,a;\, u.\, h))) &= \ulcorner \iota([a]_{\vec{y}}, [h]_{u\vec{y}})\urcorner(\sigma(\tau(c)), \vec{y}) \\ &= I(\sigma(\tau(c)), \ulcorner [a]_{\vec{y}}\urcorner(\vec{y});\, u.\, \ulcorner [h]_{u\vec{y}}\urcorner(u,\vec{y})) = I(c,a;\, u.\, h).\end{aligned}$$

(ii) First, $\mathsf{iPRA} \vdash \tau(\ulcorner f\urcorner) = f(x_1, \ldots, x_k)$ for every symbol f of arity k, by induction on f. For zero and succ the two sides are the same term, and for a projection the claim is its axiom. For a composition, by Theorem 1.73(ii) and the induction hypotheses,

$$\tau(\ulcorner f\urcorner(\ulcorner g_1\urcorner, \ldots, \ulcorner g_m\urcorner)) = \tau(\ulcorner f\urcorner)[\tau(\ulcorner \vec{g}\urcorner)/\vec{x}] = f(g_1(\vec{x}), \ldots, g_m(\vec{x})) = \mathrm{comp}(f;\vec{g})(\vec{x}).$$

For a recursion $p := \mathrm{prec}(f,g)$ write $q := \tau(\ulcorner p\urcorner)$ and $\vec{x}' := x_2, \ldots, x_{k+1}$. By Theorem 1.62(i), T_{eq} proves $\ulcorner p\urcorner[0/x_1] = \ulcorner f\urcorner(\vec{x}')$ and $\ulcorner p\urcorner[Sx_1/x_1] = \ulcorner g\urcorner(x_1, \ulcorner p\urcorner, \vec{x}')$. By Theorem 1.74, Theorem 1.69 and Theorem 1.73(ii), iPRA proves the images of these equations, and with the induction hypotheses for f and g they read

$$q[0/x_1] = f(\vec{x}'), \qquad q[Sx_1/x_1] = g(x_1, q, \vec{x}').$$

Quantifier-free induction on x_1 in the equation $q = p(x_1, \vec{x}')$ now succeeds: the base is $q[0/x_1] = f(\vec{x}') = p(0,\vec{x}')$, and under the induction equation $q[Sx_1/x_1] = g(x_1, q, \vec{x}') = g(x_1, p(x_1,\vec{x}'), \vec{x}') = p(Sx_1, \vec{x}')$.

The claim follows by induction on t: $\tau(\sigma(f(\vec{t}))) = \tau(\ulcorner f\urcorner)[\tau(\sigma(\vec{t}))/\vec{x}] = f(\tau(\sigma(\vec{t}))) = f(\vec{t})$, by Theorem 1.73(ii), the first step and the induction hypothesis. □

1.8.6 The bi-interpretation theorem

Theorem 1.78 (Bi-interpretation)**.** *The systems N, T_{eq} and iPRA are pairwise bi-interpretable. Precisely:*

(i) *N and T_{eq}. $\Gamma \vdash_{\mathsf{N}} A$ if and only if $\Gamma^\circ \vdash_{\mathsf{T}_{\mathrm{eq}}} A^\circ$; $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} \varphi$ if and only if $\Gamma^* \vdash_{\mathsf{N}} \varphi^*$; $(A^\circ)^* \dashv\vdash A$ in N; and $\mathsf{T}_{\mathrm{eq}} \vdash \varphi \leftrightarrow (\varphi^*)^\circ$.*

(ii) T_{eq} and iPRA. *τ and σ are term interpretations in opposite directions;* $\mathsf{T}_{\mathrm{eq}} \vdash \sigma(\tau(t)) = t$ *and* $\mathsf{iPRA} \vdash \tau(\sigma(t)) = t$*;* $\Gamma \vdash_{\mathsf{T}_{\mathrm{eq}}} \varphi$ *if and only if* $\Gamma^{\tau} \vdash_{\mathsf{iPRA}} \varphi^{\tau}$*;* $\Gamma \vdash_{\mathsf{iPRA}} \psi$ *if and only if* $\Gamma^{\sigma} \vdash_{\mathsf{T}_{\mathrm{eq}}} \psi^{\sigma}$*;* $\mathsf{T}_{\mathrm{eq}} \vdash \varphi \leftrightarrow (\varphi^{\tau})^{\sigma}$*; and* $\mathsf{iPRA} \vdash \psi \leftrightarrow (\psi^{\sigma})^{\tau}$.

(iii) N and iPRA. *With* $A^{\flat} := (A^{\circ})^{\tau}$ *and* $\psi^{\sharp} := (\psi^{\sigma})^{*}$*:* $\Gamma \vdash_{\mathsf{N}} A$ *if and only if* $\Gamma^{\flat} \vdash_{\mathsf{iPRA}} A^{\flat}$*;* $\Gamma \vdash_{\mathsf{iPRA}} \psi$ *if and only if* $\Gamma^{\sharp} \vdash_{\mathsf{N}} \psi^{\sharp}$*;* $(A^{\flat})^{\sharp} \dashv\vdash A$ *in* N*; and* $\mathsf{iPRA} \vdash \psi \leftrightarrow (\psi^{\sharp})^{\flat}$.

(iv) Shape of the interpretations. *Every translation fixes the variables, leaves the range of every quantifier and binder the whole numerical domain, and reads equality as equality. On terms the composites are provably the identity, so the definable map between the domain and its double interpretation is the identity, with graph $x = y$; it is total, bijective and commutes with every term former and function symbol.*

(v) Equiconsistency. $\vdash_{\mathsf{N}} 0 = S0$ *if and only if* $\vdash_{\mathsf{T}_{\mathrm{eq}}} \bot$ *if and only if* $\vdash_{\mathsf{iPRA}} \bot$.

Proof. (i) is Theorems 1.66 and 1.67. (ii) is Theorems 1.74, 1.75 and 1.77 and Theorem 1.70. (iii) The two equivalences are the compositions of those in (i) and (ii). For the first round trip, $(A^{\flat})^{\sharp} = (((A^{\circ})^{\tau})^{\sigma})^{*}$. By (ii), T_{eq} proves $A^{\circ} \leftrightarrow ((A^{\circ})^{\tau})^{\sigma}$, so by Theorem 1.65 the native readings of the two sides are interderivable in N, and $(A^{\circ})^{*} \dashv\vdash A$ by (i). For the second, $(\psi^{\sharp})^{\flat} = (((\psi^{\sigma})^{*})^{\circ})^{\tau}$. By (i), T_{eq} proves $\psi^{\sigma} \leftrightarrow ((\psi^{\sigma})^{*})^{\circ}$; applying τ, iPRA proves $(\psi^{\sigma})^{\tau} \leftrightarrow (\psi^{\sharp})^{\flat}$, and $\mathsf{iPRA} \vdash \psi \leftrightarrow (\psi^{\sigma})^{\tau}$ by (ii). (iv) The first sentence holds by the definitions of A°, φ^{*}, τ and σ; the second is Theorem 1.77, where for the span between N and T_{eq} the terms are not translated at all. (v) If $\vdash_{\mathsf{N}} 0 = S0$ then $\vdash_{\mathsf{T}_{\mathrm{eq}}} 0 = S0$ by Theorem 1.64, and (T3) gives $\bot$. If $\vdash_{\mathsf{T}_{\mathrm{eq}}} \bot$ then $\vdash_{\mathsf{N}} \bot^{*}$, which is $0 = S0$. Since $\bot^{\tau} = \bot = \bot^{\sigma}$, (ii) gives the second equivalence. □

The translations act on finite derivations effectively. No claim is made that any of the systems proves its own consistency. The interpretations use no restriction to standard numerals, no retract or quotient of the domain, and no assumed symbol for equality: the reflection of equality is Theorem 1.52, a theorem.

In Lean, part (ii) is `biInterp` with `teq_iff_pra`, `pra_iff_teq`, `teq_roundTrip`, `pra_roundTrip` and `teq_consistent_iff`; part (iii) is `native_iff_pra`, `pra_iff_native`, `native_roundTrip` and `pra_native_roundTrip`; and part (v) is `native_inconsistent_iff`. The final theorems depend on no axiom beyond `propext`, `Classical.choice` and `Quot.sound`, and part (ii) on `propext` and `Quot.sound` alone.

Remark 1.79 (Consistency). Read over the natural numbers, with the iterator as iteration, every rule of N preserves truth, and $0 = S0$ is false. Hence N does not derive $0 = S0$, and by Theorem 1.78(v) neither T_{eq} nor iPRA derives $\bot$. This argument is carried out in the metatheory and is formalized in Lean as `NPrf.sound_std`, `native_consistent`, `teq_consistent` and `pra_consistent`.

Remark 1.80 (How the packages state the bridge). Every package repeats this theorem for its own calculus, by composing an equivalence of its calculus with the inherited one and the inherited bridge. The reference statement composes two steps: the native calculus against an ordinary theory T_I, which is T_{eq} with the iterators presented as staged function symbols (Theorem 1.4), and T_I against iPRA. Its proof of the second step identifies the double translation of an operation with the operation by uniqueness of iteration and of primitive recursion, as in Theorem 1.77; the explicit treatment of substitution (Theorems 1.72 and 1.73) is spelled out in the Lean formalization and is given here in full.

1.9 Clause contexts

This section draws on the definition in the main report of `raw_clausal` and on the reference statement of the inherited basis filed with `certified_zero`. Only the notion and its reading are fixed here; the clause calculi are the subject of other chapters, beginning with Section 5.2.

Definition 1.81 (Clauses and clause contexts)**.** A *clause* is an expression

$$[A;\, y.\, B(y)], \qquad y \notin \mathrm{FV}(A),$$

where A and B are native formulas. The clause binds y in its consequent B only. It is a member of a context, not a formula: it cannot occur inside a formula or as the conclusion of a sequent. A *clause context* is a finite list of clause occurrences. Its *ordinary reading* is

$$[A;\, y.\, B(y)]^{\circ} := A^{\circ} \to \forall y\, B^{\circ}(y).$$

An assumption of a formula A is the *constant clause*

$$[\,0 = 0;\, v.\, A\,], \qquad v \notin \mathrm{FV}(A),$$

and Γ, A abbreviates $\Gamma, [\,0 = 0;\, v.\, A\,]$.

A clause is used by the inference that, for an occurrence $[A;\, y.\, B(y)]$ in Γ, passes from $\Gamma \vdash A$ to $\Gamma \vdash B(t)$ for a term t. Three observations connect clauses to the formulas of this chapter.

(i) The reading of the constant clause is $0 = 0 \to \forall v\, A^{\circ}$, which is equivalent to A° in logic with equality. The use of a constant clause is the use of the hypothesis A, preceded by a derivation of $0 = 0$.

(ii) The clause $[B(b, 0);\, y.\, B(b, y)]$ has the reading of $Q_y B(b, y)$. It is the clause that the elimination of $D_{x,y}B$ adds for a fresh b, and its use is QE. A calculus with clause contexts therefore needs no binder Q: its formulas are generated from equations by D alone.

(iii) A native formula has a form without Q: by Theorem 1.12, $Q_y B$ is interderivable with $D_{x,y}B$ for x not free in B.

The reference statement of the inherited basis uses the antecedent $I(0, 0;\, u.\, 0) = 0$, an instance of I0, in the constant clause; `raw_clausal` and the later packages use $0 = 0$. A context retains clause occurrences, not only their values: an inference that uses a clause names an occurrence.

1.10 The measurement protocol

This section states the convention common to all packages; the wording follows `certified_zero` and `evaluation_discharge`.

Definition 1.82 (Measured object and coordinates)**.** The measured object is a *complete standalone proof checker*: one Python module that reads a certificate in JSON and accepts exactly the derivable sequents of the calculus it presents. The whole module is counted: imports, input handling, syntax and scope checks, all rules, and the acceptance driver. The source is first *normalized*: docstrings and comments are removed by reconstruction from the syntax tree, the tree is printed by `ast.unparse` of CPython 3.12.14, and one final line feed is appended. The three coordinates of the normalized source are

(i) the number of significant *tokens*, excluding whitespace, comments, indentation markers and end markers;

(ii) the number of *AST nodes* of the whole module;

(iii) the number of *normalized bytes* in UTF-8.

The size of the normalized source under `gzip` at level 9, with the modification time set to zero and no file name, is reported for information and is not a coordinate. A presentation qualifies only if its ordinary reading is bi-interpretable with iPRA in the sense of Theorem 1.78: whole numerical domains, actual equality, identity composites with graph $x = y$, and both formula round trips. Each package compares its checker with a *matched control*, an implementation of the calculus it improves on, written with the same input contract, variable encoding and context representation, so that a difference in size is attributable to the change of rules. Where different programs attain the minima of different coordinates, they are reported as separate programs and no row combines measurements of different files.

Three limits apply to every size in the article. A measured size is an achieved upper bound on an unknown optimum; it is not a lower bound for all calculi or all implementations. A different Python version can change the normalization. The finite tests that accompany a checker supplement the written proofs and are not proofs.

No checker belongs to this chapter: the calculus N and its bridge are the common reference, and the sizes of the checkers that implement N or its variants are listed in the chapters on those variants.

1.11 Further questions

Question 1.83 (Independence of the progress rule)**.** Is QI derivable from the seven other rules of N? The repository records that each of the other seven rules is independent, and that a reading over the natural numbers cannot separate QI: there the soundness conditions of DI and DE imply the condition of QI. A countermodel needs nonstandard elements.

Question 1.84 (Progress at compound matrices)**.** If QI is allowed for an arbitrary matrix, the proof of Theorem 1.20 gives induction for every native formula, and so, by Theorem 1.29, for the native reading of every ordinary formula. The reference statement of the inherited basis asserts that this changes the theory. It gives no proof that the resulting theory is not bi-interpretable with iPRA in the sense of Theorem 1.78.

Question 1.85 (Other term languages)**.** The reference statement of the inherited basis asserts that its argument applies unchanged when the iterator terms are replaced by the terms of iPRA, with the defining equations of projection, composition and recursion in place of I0 and IS. The assertion is stated without proof. The derivations of Section 1.4 use the iterator only through the selector and the parity term.

Question 1.86 (A shorter bridge)**.** Three constructions account for most of the length of the proof of Theorem 1.78: the predecessor from iteration alone (Theorem 1.48), the conjunction of two formulas through the parity term (Theorem 1.28), and the selector of two formulas by a double recursion on their binders (Theorem 1.24). No shorter replacement for any of them is known. On the side of iPRA, the extensionality of the simulated iterator (Theorem 1.72) and the round trip of a recursion (Theorem 1.77(ii)) are two separate inductions; in T_{eq} their counterparts are both instances of Theorem 1.38. Is there one lemma of iPRA that covers both?

Question 1.87 (The square-root pairing)**.** The reference statement on primitive recursion obtains pairing from a bounded square-root search and refers to the linear-order and semiring identities of multiplication as usual without deriving them in T_{eq}. The article replaces that pairing by the one of Theorem 1.59. A derivation of the square-root pairing in T_{eq} from Theorem 1.56 is not written out in any package.

Chapter 2

Refinements of the eight rules

The native scalar core N has eight rules (Section 1.3). This chapter collects the changes of that rule basis which keep the language of equations, Q and $D_,$ as it is: there are no clause contexts, no restricted class of public formulas and, with the one exception of the calculus $\mathsf{U_c}$ in Section 2.4.8, no normal forms of terms. The changes fall into four groups. Section 2.2 treats the two introduction rules: progress may introduce a closure directly or through a witnessed $D_,$, and the two introductions may be merged into one rule whose mode is read off the matrix and the witness. Section 2.3 treats the transport rule: the number of templates, the crossing of the conclusion, the exclusion of the literal 0 as a source, the reversal of the root equation, replacement with no template, and replacement of a single occurrence. Section 2.4 replaces induction for equations by uniqueness of iteration. Section 2.5 changes the reading of Q at a literal equation to progressivity and presents the seven-rule calculi that result. Section 2.6 records two results on recursors other than the iterator, which two of the packages prove on the side.

Every calculus of Sections 2.2 to 2.4 derives exactly the sequents of N, on the same formulas. The calculi of Section 2.5 do not; they are equivalent to N through a pair of translations with both formula round trips. Table 2.1 at the end of Section 2.5 lists all calculi of the chapter.

2.1 Conventions and the comparison method

This section fixes what all thirteen packages of the chapter restate before their own results; it draws on no package in particular.

The rules of N are named as in Section 1.3: Hyp, I0, IS, Tr, QE, DE, DI (introduction of $D_,$ by cases) and QI (introduction of Q by progress at a literal equation). A *term template* $h(v)$ is a term with a designated variable v, which may occur several times, once, or not at all; in the last case the template is *constant*. An *equation template* is $a(v) = b(v)$ for two term templates, and a *formula template* $F(v)$ is a native formula with a designated variable. The sign $\equiv$ is identity of syntax (up to the names of bound variables), so that $s \not\equiv 0$ says that s is not the term 0. This is a test on syntax and not on values: $I(0, 0; u.\, 0) \not\equiv 0$, although the term equals 0 by I0. A formula is *compound* if it is not a literal equation, that is, if its root is Q or $D_,$. Substitution preserves the root of a formula and may change whether a term is the literal 0 or a literal successor.

All calculi of the chapter have the terms of Section 1.1, the formulas of Section 1.2, finite lists of formulas as contexts, and the rules Hyp, I0, IS and DE in their forms of Section 1.3. A rule inventory below therefore names these four rules and displays the others. The one exception is the calculus $\mathsf{U_c}$ of Theorem 2.83, which is stated on normal terms and has reflexivity in place of the two computation equations. Derivability in a calculus X is written

$\vdash_{\mathsf{X}}$, and $\vdash$ without a subscript is derivability in N.

Definition 2.1 (Equivalence on the same formulas)**.** Two calculi X and Y on the native language are *equivalent* if $\Gamma \vdash_{\mathsf{X}} A$ holds exactly when $\Gamma \vdash_{\mathsf{Y}} A$ holds, for every context Γ and every native formula A.

The equivalence is about sequents with arbitrary contexts and arbitrary formulas, not about closed equations. It is always proved by two translations of derivations that copy the common rules and replace each instance of a changed rule by a fixed finite derivation; the translations are effective and leave terms, formulas, contexts and conclusions unchanged.

Proposition 2.2 (Transfer of the bridge)**.** *Let* X *be equivalent to* N*. Then every statement of Theorem 1.78 holds for* X *with the same translations:* X*,* T_{eq} *and* iPRA *are bi-interpretable with whole domains, actual equality and identity numerical composites, both formula round trips hold, and* $\vdash_{\mathsf{X}} 0 = S0$ *holds exactly when* iPRA *is inconsistent. Weakening, substitution of terms, cut and the Leibniz rule are admissible in* X*.*

Proof. The statements of Theorem 1.78 and Section 1.4 about N are statements about its derivable sequents, and X has the same derivable sequents. □

Each package of the chapter restates this transfer for its own calculus; it is not repeated below. The last sentence of the proposition concerns the consequence relation of X. Whether an operation maps each rule instance to an instance of the same rule is a different question.

Lemma 2.3 (Structural operations by direct induction)**.** *In every calculus of this chapter, weakening of the context, injective renaming of free variables, and cut are admissible by a direct induction on derivations. Substitution of terms for variables maps rule instances to rule instances of the same rule in a calculus none of whose side conditions tests a term for being the literal* 0*, a literal successor or a syntactic match of another term.*

Proof. The side conditions of the rules below are of four kinds: freshness of an eigenvariable; a test on the root of the matrix; a test whether a term of the instance is the literal 0 or a literal successor; and, for one rule, a syntactic match of subterms. An injective renaming preserves all four. For weakening, rename the eigenvariables of the derivation away from the added formulas and keep the added formulas in every context; each inference is still an instance of its rule with the same terms. For cut, let $\Gamma \vdash_{\mathsf{X}} A$ and $\Gamma, A \vdash_{\mathsf{X}} C$. Rename the eigenvariables of the second derivation away from the first, and replace every use of Hyp for the assumption A by the first derivation, weakened by the local assumptions. No term of an inference changes, so every side condition is preserved. For substitution the first two kinds of side condition are preserved after a renaming of eigenvariables, as in the proof for N in Section 1.4; the last two kinds need not be. □

Several derived rules of Section 1.4 use only part of N, and they are used in that form.

Lemma 2.4 (Derived rules and the rules they use)**.** *Let* X *be a calculus of this chapter.*

(i) *If* X *contains QE and DI, then* X *derives:* $D_{x,y}B \dashv\vdash Q_yB$ *for* $x \notin \mathrm{FV}(B)$*; the introduction of* Q_yB *by cases, from* $\Gamma, B(0) \vdash B(Sz)$ *with* z *fresh, for an arbitrary matrix* B*; numerical cases, from* $\Gamma \vdash B(0)$ *and* $\Gamma \vdash B(Sz)$ *with* z *fresh to* $\Gamma \vdash B(t)$*; and* $D_{x,y}B(x,y)$ *from* $Q_yB(t,y)$*.*

(ii) *If* X *contains QE and QI, then* X *derives equation induction: from* $\Gamma \vdash E(0)$ *and* $\Gamma, E(z) \vdash E(Sz)$*, with* E *a literal equation and* z *fresh, to* $\Gamma \vdash E(t)$*.*

(iii) *If* X *contains* Tr*, then* X *derives reflexivity, symmetry, transitivity, congruence in every term template, and transport in an equation template, from* $s = t$ *and* $a(s) = b(s)$ *to* $a(t) = b(t)$.

Proof. The derivations of these rules in Section 1.4 use Hyp, DE, weakening and the rules named in each item; for (iii) they use I0 and Tr alone. □

In (iii) each of the four equality rules is one application of Tr, and transport in an equation template is two. Symmetry needs a constant template: Tr with the template $h(v) := s$, any derivable equation as its first premise and $s = t$ as its second premise concludes $t = s$.

2.2 The introduction rules

This section draws on `direct_closure`, `direct_closure_qprogress` and `witness_rule`, and on the obstruction section of `reversal_core`.

2.2.1 Progress through a witness and progress at the closure

The reports state induction for equations as an introduction of D, with a witness:

$$\frac{\Gamma,\ E(t,z) \vdash E(t,Sz)}{\Gamma \vdash D_{x,y}E(x,y)}\ \mathrm{DP} \qquad (E \text{ a literal equation}),$$

where z is not free in Γ, in t or in $D_{x,y}E$.

Definition 2.5. N_D is the calculus with the rules Hyp, I0, IS, Tr, QE, DE, DI and DP. It is N with QI replaced by DP.

The rule DP has a matrix in two variables and a witness term; the rule QI has a matrix in one variable and no witness.

Theorem 2.6 (Direct closure)**.** N_D *and* N *are equivalent. The proof uses only the rules Hyp, QE, DE, DI and weakening; it holds for any common set of further rules for which weakening and renaming are admissible.*

Proof. All rules but one are common. *QI in* N_D. Let $\Gamma, E(z) \vdash E(Sz)$ be the translated premise. Choose a variable x not free in E and Γ. The rule DP with the matrix $E(y)$, in which the coordinate x is not used, and the witness 0 has exactly this premise and concludes $\Gamma \vdash D_{x,y}E(y)$. Apply DE with a fresh b: the opened hypothesis is $Q_yE(y)$, because x does not occur in E, and the second premise $\Gamma, Q_yE(y) \vdash Q_yE(y)$ is Hyp. The conclusion is $\Gamma \vdash Q_yE(y)$.

DP in N. Let $\Gamma, E(t,z) \vdash E(t,Sz)$ be the translated premise. The formula $E(t,y)$ is a literal equation in y with t as a parameter, and z is not free in t, so QI gives $\Gamma \vdash Q_yE(t,y)$. Let w be fresh. Over $\Gamma, E(t,0)$, weakening and QE at Sw give $E(t,Sw)$. This is the premise of DI with the witness t, which concludes $\Gamma \vdash D_{x,y}E(x,y)$. No equality rule, no computation equation and no substitution into the given premise is used. □

Proposition 2.7 (Node counts)**.** *Count every instance of a rule as one node. A derivation of* N *with* n *nodes,* r *of them instances of QI, translates to a derivation of* N_D *with* $n + 2r \le 3n$ *nodes. A derivation of* N_D *with* n *nodes,* r *of them instances of DP, translates to a derivation of* N *with* $n + 3r \le 4n$ *nodes.*

Proof. An instance of QI becomes DP, DE and Hyp over the same premise. An instance of DP becomes QI, QE, Hyp and DI over the same premise. Each premise derivation is used once, and weakening and renaming do not change the number of nodes. □

The bounds count nodes of the inference tree; they do not bound the size of formulas, of serialized certificates or of a checker.

Remark 2.8 (Sources). `direct_closure` and `direct_closure_qprogress` are two independent accounts of Theorems 2.6 and 2.7, with the same two constructions and the same counts; the second contains the proof twice, once as a proof note and once as an audit. Both state the calculi with transport in a formula template in place of Tr, which the theorem permits. Both stress that the rule QI was a derived rule of the calculus before it was taken as primitive, so that the change is one of the choice of primitives. Their example is $F(n) := I(n, 0; u.\, Su)$ with the equation $F(y) = y$: with transport in a formula template the derivation of $F(n) = n$ by QI and QE has six nodes, its translation to N_D has eight, and the translation of that back has eleven. In Lean the rule QI is the constructor `qIInd` of the kernel, adopted from these packages, and DP is the derived rule `NPrf.dIProg`; the equivalence of the calculus of the reports, with DP and transport in a formula template, is `iprf_iff_nprf`.

2.2.2 One closure introduction with a mode read off the matrix

For a matrix $B(y)$ and a variable z put $\eta_B(z) := z$ if B is a literal equation and $\eta_B(z) := 0$ if B is compound.

Definition 2.9. N_det is N with QI replaced by

$$\frac{\Gamma,\ B(\eta_B(z)) \vdash B(Sz)}{\Gamma \vdash Q_y B(y)}\ \mathrm{QI}_\eta,$$

where B is an arbitrary native formula and z is not free in Γ and $Q_y B$. Its rules are Hyp, I0, IS, Tr, QE, DE, DI and QI_η.

A certificate for QI_η carries no choice between cases and progress: the root of the matrix decides.

Corollary 2.10. N_det, N *and* N_D *are equivalent, and the bounds of Theorem 2.7 hold between* N_det *and* N_D.

Proof. At a literal equation QI_η is QI. At a compound matrix it is the introduction of Q by cases, which is derivable in N by Theorem 2.4(i): DI with an unused coordinate and the witness 0, then DE and Hyp. This is the same three-node construction as in Theorem 2.6, with DI in place of DP. Conversely N_det contains QI. □

The introduction of $Q_y E$ by cases for a literal equation E is not an instance of QI_η, and it remains derivable through DI and DE. For example $Q_y(y = 1)$ is derivable in all three calculi from the premise $0 = 1 \vdash Sz = 1$ of the introduction by cases, while the premise $z = 1 \vdash Sz = 1$ of progress is not derivable.

2.2.3 What cannot be removed from the introductions

Proposition 2.11 (The guard of QI)**.** *If QI is allowed at every matrix, the resulting calculus derives induction for every native formula: from* $\Gamma \vdash B(0)$ *and* $\Gamma, B(z) \vdash B(Sz)$ *with* z *fresh to* $\Gamma \vdash B(t)$.

Proof. The unrestricted rule gives $\Gamma \vdash Q_y B(y)$, and QE gives $B(t)$. □

Through the native connectives of Section 1.4 this is induction for every first-order formula, and the bridge of Theorem 1.78, whose target has induction for quantifier-free formulas only, no longer applies. The rule QI_η avoids this by using the instance at zero at a compound matrix.

Proposition 2.12 (DI cannot be dropped)**.** *Let* N^- *be* N *without DI, or* $\mathsf{N}_{\mathrm{det}}$ *without DI. If no formula of* Γ *has the root* $D_{,}$*, then* N^- *derives from* Γ *no formula with the root* $D_{,}$*. In particular* N^- *does not derive* $D_{x,y}(0 = 0)$*, which* N *derives.*

Proof. Suppose a derivation of $\Gamma \vdash_{\mathsf{N}^-} A$ with A of root $D_{,}$ and Γ as stated, and take one with the least number of nodes. Its last rule is not Hyp, by the assumption on Γ; not a computation equation or Tr, whose conclusions are equations; and not an introduction of Q, whose conclusion has the root Q. If it is QE with the conclusion $B(t)$, then its second premise $\Gamma \vdash B(0)$ has the same root $D_{,}$ and the same context, and a smaller derivation. If it is DE, then its first premise $\Gamma \vdash D_{x,y}B$ has the root $D_{,}$ and the same context, and a smaller derivation. Both contradict minimality. In N the formula $D_{x,y}(0 = 0)$ follows by DI with the witness 0 from the premise $0 = 0 \vdash 0 = 0$. □

The assumptions inside a subderivation of an introduction of Q do not affect the argument, because such an introduction is never the last rule of a derivation of a formula with the root $D_{,}$.

Proposition 2.13 (DI cannot be restricted to compound matrices)**.** *Restrict DI to compound matrices* B*.*

(i) *The calculus with Hyp, I0, IS, Tr, QE, DE, QI and the restricted DI does not derive* $D_{x,y}(0 = 0)$*.*

(ii) *The calculus with Hyp, I0, IS, Tr, QE, DE, DP and the restricted DI does not derive* $D_{x,y}(y = 1)$*.*

Both formulas are derivable in N*. The statements remain true with transport in a formula template in place of Tr.*

Proof. Both parts interpret terms in the standard natural numbers, equations as equality and Q_yB by its ordinary reading, and give a formula $D_{x,y}B$ with a compound matrix its ordinary reading. The interpretation of a formula $D_{x,y}E$ with a literal equation E is changed. It commutes with substitution, because substitution preserves the root of a matrix, and truth of a sequent means that every assignment satisfying the context satisfies the conclusion.

(i) Let $D_{x,y}E$ be false for every literal equation E. The rules Hyp, I0, IS, transport and QE preserve truth as they do for the ordinary reading. QI preserves truth by induction on the natural numbers. DE at an atomic matrix has a false first premise, and at a compound matrix it is the elimination of an existential quantifier. The restricted DI preserves truth because every natural number is zero or a successor. The formula $D_{x,y}(0 = 0)$ is false.

(ii) Let $D_{x,y}E$ mean $\exists x \forall y\, (E(x, y) \to E(x, Sy))$ for a literal equation E. DP preserves truth: its premise states the progress of $E(t, y)$, and t is a witness. DE at an atomic matrix preserves truth: if $\forall y\, (E(b, y) \to E(b, Sy))$ holds, then $E(b, 0) \to \forall y\, E(b, y)$ holds by induction on the natural numbers, and this is the reading of the opened hypothesis $Q_yE(b, y)$. The other rules preserve truth as in (i). The formula $D_{x,y}(y = 1)$ is false, because at $y = 1$ the equation $1 = 1$ holds and $2 = 1$ does not.

In N, $D_{x,y}(0 = 0)$ is derived in Theorem 2.12, and $D_{x,y}(y = 1)$ follows by DI with the witness 0 from $0 = 1 \vdash Sz = 1$, which holds because $0 = 1$ derives every equation (Section 1.4). □

Remark 2.14 (Sources). Theorems 2.11 and 2.12 are from `direct_closure_qprogress`; `direct_closure` states the first. Theorem 2.13(i) is the obstruction section of `reversal_core`. Part (ii) is stated in `direct_closure_qprogress` with the interpretation and a reference to an earlier package for the verification of the rules. `direct_closure_qprogress` also recalls why the ordinary theory T_{eq} needs its coverage axiom: the product of two copies of the standard

model, with all operations coordinatewise, satisfies the computation equations, separation and induction for equations, because the solution set of an equation is a product $U_0 \times U_1$, the base puts 0 into both factors, and the step applied at $(r, 0)$ and at $(0, r)$ closes each factor under successor; and the element $(0, 1)$ is neither zero nor a successor.

2.2.4 One introduction with a mode read off the matrix and the witness

The two introductions DI and QI can be replaced by one introduction of $D_{,}$. For a matrix $B(x, y)$, a witness t and a variable z put

$$\eta_{B,t}(z) := \begin{cases} z, & B \text{ is a literal equation and } t \equiv 0, \\ 0, & \text{otherwise.} \end{cases}$$

Definition 2.15 (The calculus W)**.** W has the seven rules Hyp, I0, IS, Tr, QE, DE and

$$\frac{\Gamma,\ B(t, \eta_{B,t}(z)) \vdash B(t, Sz)}{\Gamma \vdash D_{x,y}B(x, y)}\ \mathrm{DI_w},$$

where z is not free in Γ, in t or in $D_{x,y}B$.

The rule selects progress exactly when the matrix is a literal equation and the witness is the literal 0, and cases otherwise. A witness such as $I(0, 0;\ u.\, 0)$, which equals 0, selects cases. The calculus has no rule that introduces Q.

Lemma 2.16 (A second name for zero)**.** *Let $c := I(0, 0;\ u.\, 0)$. Every calculus containing I0 and Tr derives $c = 0$ and $0 = c$, and $c \not\equiv 0$.*

Proof. $c = 0$ is an instance of I0, and $0 = c$ follows by symmetry (Theorem 2.4(iii)). □

Theorem 2.17 (Witness-selected introduction)**.** W *and* N *are equivalent.*

Proof. From W *to* N. An instance of $\mathrm{DI_w}$ in the mode of cases is an instance of DI. An instance in the mode of progress has a literal equation $E(x, y)$ as its matrix, the witness 0 and the premise $\Gamma, E(0, z) \vdash E(0, Sz)$; it is an instance of DP with the witness 0, which N derives by Theorem 2.6.

From N *to* W. The rule QI with the premise $\Gamma, E(z) \vdash E(Sz)$: choose x not free in E and Γ. The rule $\mathrm{DI_w}$ with the matrix $E(y)$ and the witness 0 is in the mode of progress, has this premise and concludes $D_{x,y}E(y)$; DE and Hyp give $Q_y E(y)$ as in Theorem 2.6.

The rule DI with a compound matrix, or with a witness $t \not\equiv 0$, is an instance of $\mathrm{DI_w}$. There remains DI with a literal equation $E(x, y)$ as its matrix and the witness 0. Let

$$\pi\colon \quad \Gamma,\ E(0, 0) \vdash_{\mathsf{W}} E(0, Sz)$$

be the translated premise, with z fresh. Work over $\Gamma, E(c, 0)$ with c as in Theorem 2.16. Transport in the equation template $E(v, 0)$ along $c = 0$ gives $E(0, 0)$ (Theorem 2.4(iii)). Cut with π, which is admissible in W by Theorem 2.3, gives $E(0, Sz)$. Transport in the equation template $E(v, Sz)$ along $0 = c$ gives $E(c, Sz)$. Hence $\Gamma, E(c, 0) \vdash_{\mathsf{W}} E(c, Sz)$. Since $c \not\equiv 0$, the rule $\mathrm{DI_w}$ with the witness c is in the mode of cases and has exactly this premise. It concludes $D_{x,y}E(x, y)$; the conclusion does not record the witness. □

The construction changes an assumption by a derived transport and a cut. It does not substitute a term into the derivation π, so it cannot change the mode of an introduction inside π.

Remark 2.18 (Substitution in W). A substitution can turn a witness that is a variable into the literal 0, and then an instance of DI_{w} in the mode of cases is mapped to a sequent that is not the premise of an instance in the mode of progress. Substitution is admissible in W by Theorem 2.2, through N, and is not obtained by a direct induction on the derivations of W. For this reason the Lean kernel keeps the two introductions of N.

Remark 2.19 (Sources and variants). Theorem 2.17 is the theorem of `witness_rule`; in Lean it is `wprf_iff_nprf`, for the calculus `WPrf`. The package compares W with the calculus that has DI and DP, so its proof has four cases: DI repeated; DI at an equation with the witness 0, as above; DP with the witness 0 repeated; and DP with a witness $t \not\equiv 0$, which is the second construction of Theorem 2.6 with the inner closure introduced through an unused coordinate. The package works with a restricted inventory of computation instances and therefore uses $I(2,1;\, u.\,0)$ as its second name for zero; that inventory is treated in Theorem 2.53.

The argument does not use any property of the number zero in the witness coordinate. If p is a closed term and $q \not\equiv p$ is a closed term with $p = q$ and $q = p$ derivable from the common rules, then progress may be selected exactly at the witness p: an introduction by cases at p is repaired at the witness q, and every auxiliary introduction with an unused coordinate takes the witness p. The coordinate y of the induction still starts at 0. The literal 0 is chosen because it is the cheapest witness to recognize.

`one_context` states the same calculus with the two-template transport of Section 2.3.1 replaced by Tr and calls it $\mathcal{O}_7$; since N already has Tr, its calculus is W as defined here.

2.3 The transport rule

This section draws on `one_context`, `one_sided_transport`, `reversal_core`, `source_dispatch`, `template_free` and the note on single occurrences in `kernel_refinement`.

Throughout the section $J(a) := I(0, a;\, u.\,0)$ and $c := J(0)$. The equation $J(a) = a$ is an instance of I0, and $J(a) \not\equiv 0$ for every term a; in particular $c = 0$ is an instance of I0 and $c \not\equiv 0$.

2.3.1 One template, two templates, a formula template

The rule Tr of N has one term template and a crossed conclusion: from $s = t$ and $h(s) = r$ it infers $r = h(t)$. The reports use two stronger forms,

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash a(s) = b(s)}{\Gamma \vdash a(t) = b(t)}\ \mathrm{Tr}_2 \qquad\qquad \frac{\Gamma \vdash s = t \qquad \Gamma \vdash F(s)}{\Gamma \vdash F(t)}\ \mathrm{Tr}_F,$$

with two term templates $a(v)$, $b(v)$ and with a native formula template $F(v)$.

Definition 2.20. N_2 is N with Tr replaced by Tr_2, and N_F is N with Tr replaced by Tr_F. Each has the rules Hyp, I0, IS, its transport rule, QE, DE, DI and QI.

Theorem 2.21 (One term template suffices)**.** N, N_2 *and* N_F *are equivalent. An instance of* Tr_2 *is simulated in* N *by two instances of* Tr *with the same first premise. An instance of* Tr *is simulated in* N_2 *by one instance of* Tr_2 *with a constant second template, followed by symmetry.*

Proof. Tr_2 *in* N. Tr with the template a, applied to $s = t$ and $a(s) = b(s)$, gives $b(s) = a(t)$. Tr with the template b, applied to $s = t$ and $b(s) = a(t)$, gives $a(t) = b(t)$; the term $a(t)$ is the fixed side of the second step.

Tr *in* N_2. First, N_2 derives reflexivity and symmetry: Tr_2 with the templates v and a, applied to $J(a) = a$ and $J(a) = a$, gives $a = a$; and Tr_2 with the templates v and s, applied

to $s = t$ and $s = s$, gives $t = s$. Now let $s = t$ and $h(s) = r$ be given. Tr_2 with the templates $h(v)$ and the constant r gives $h(t) = r$, and symmetry gives $r = h(t)$.

N_F. An instance of Tr_2 is the instance of Tr_F at the template $a(v) = b(v)$, so N_F derives every sequent of N_2. Conversely Tr_F is a derived rule of N, by the Leibniz rule of Section 1.4.

In each direction the seven other rules are copied, so the induction on derivations applies to open sequents and to premises under both binders. □

The crossing of the conclusion is what makes two successive replacements in one template reproduce a simultaneous replacement in both sides of an equation. The rule Tr has no side condition, so substitution maps its instances to instances (Theorem 2.3).

Remark 2.22 (Size of the simulation). The simulation of Tr_2 uses the derivation of $s = t$ twice. A proof represented as a directed acyclic graph shares it; a proof represented as a tree copies it, and repeated translation can enlarge a tree substantially. No linear bound on the size of translated trees is claimed in `one_context` or `one_sided_transport`.

Remark 2.23 (Sources). The comparison of N with N_2 is the theorem of `one_context`, whose reference calculus is N_2. In Lean the rule Tr is the constructor `NPrf.trOne` of the kernel, adopted from that package, and Tr_2 is the derived rule `NPrf.trEq`. The comparison with N_F is the remark on the calculus of the reports in Section 1.3.

2.3.2 Sources other than the literal zero

Let Tr^- be the rule Tr with the side condition $s \not\equiv 0$ on the left side of its first premise, the *source*:

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash h(s) = r}{\Gamma \vdash r = h(t)}\ \mathrm{Tr}^- \qquad (s \not\equiv 0).$$

Every variable and every term with the root S or I is a permitted source, including terms whose value is zero.

Definition 2.24 (The calculi O and M)**.** O is N with Tr replaced by Tr^-; its rules are Hyp, I0, IS, Tr^-, QE, DE, DI and QI. M is W with Tr replaced by Tr^-; its seven rules are Hyp, I0, IS, Tr^-, QE, DE and DI_w.

Lemma 2.25 (Equality from I0 and Tr^-)**.** *The following are derivable from I0 and* Tr^- *alone, over an arbitrary context.*

(i) *Reflexivity:* $a = a$.

(ii) *Symmetry: from* $a = b$ *to* $b = a$, *also when* a *or* b *is the literal* 0.

(iii) *Restricted transport in an equation template: if* $s \not\equiv 0$, *from* $s = t$ *and* $a(s) = b(s)$ *to* $a(t) = b(t)$.

(iv) *Joining through zero: if* $p \not\equiv 0$ *and* $q \not\equiv 0$, *from* $p = 0$ *and* $q = 0$ *to* $p = q$.

(v) *Transitivity: from* $r = s$ *and* $s = t$ *to* $r = t$.

(vi) *Congruence: from* $s = t$ *to* $f(s) = f(t)$, *for every term template* f.

(vii) *The rule* Tr *without the side condition.*

Proof. (i) Tr^- with the template v, applied to $J(a) = a$ and $J(a) = a$, gives $a = a$; the source is $J(a) \not\equiv 0$.

(ii) Tr^- with the constant template a, applied to $c = 0$ and $a = b$, gives $b = a$; the source is $c \not\equiv 0$.

(iii) As in Theorem 2.21: the two steps have the same first premise $s = t$.

(iv) $\textsc{Tr}^-$ with the template $J(v)$, applied to $p = 0$ and $J(p) = p$, gives $p = J(0)$, that is $p = c$. In the same way $q = c$. By (ii), $c = p$ and $c = q$. $\textsc{Tr}^-$ with the template v, applied to $c = q$ and $c = p$, gives $p = q$. The sources are p, q and c.

(v) If $s \not\equiv 0$, reverse $r = s$ by (ii) and apply $\textsc{Tr}^-$ with the template v to $s = t$ and $s = r$. Let $s \equiv 0$. If $r \equiv 0$, the second premise is the conclusion; if $t \equiv 0$, the first premise is. Otherwise $r \not\equiv 0$ and $t \not\equiv 0$, the premises are $r = 0$ and, after (ii), $t = 0$, and (iv) applies.

(vi) If $s \not\equiv 0$, apply $\textsc{Tr}^-$ with the template f to $s = t$ and $f(s) = f(s)$. If $s \equiv 0$ and $t \not\equiv 0$, reverse the premise, obtain $f(t) = f(0)$ and reverse again. If both sides are the literal 0, use (i).

(vii) From $s = t$ and $h(s) = r$: (vi) gives $h(s) = h(t)$, (ii) gives $r = h(s)$, and (v) gives $r = h(t)$. □

The case distinctions in this proof are on the syntax of the given terms and belong to the construction of the derivation. The proof uses no induction, no numerical case distinction, no successor computation and no substitution into a derivation. It uses the constant template in (ii) and the crossing of the conclusion in (ii) and (iv).

Proposition 2.26 (Retraction criterion)**.** *Let a term language have a literal* 0 *and a term context* $J(v)$ *such that* $J(a) \not\equiv 0$ *for every term* a*, and let* $J(a) = a$ *be available as an axiom scheme in every context. Then the crossed rule restricted to sources other than the literal* 0 *derives the crossed rule at all sources, and with it reflexivity, symmetry, transitivity, congruence and transport in an equation template.*

Proof. The proof of Theorem 2.25 uses nothing else. □

Theorem 2.27 (Transport at sources other than the literal zero)**.** O*,* M *and* N *are equivalent.*

Proof. Every instance of $\textsc{Tr}^-$ is an instance of Tr, and Tr is derivable in O and in M by Theorem 2.25(vii). This proves the equivalence of O and N. For M, the rule $\mathrm{DI_w}$ is derivable in N by Theorem 2.17. Conversely the proof of Theorem 2.17 derives QI and DI in M: it uses $\mathrm{DI_w}$, DE, Hyp, cut, which is admissible in M by Theorem 2.3, the equations $c = 0$ and $0 = c$, and transport in an equation template along them, which M derives from Tr by Theorem 2.4(iii). □

Remark 2.28 (Substitution and the choice of the kernel)*.* Substituting 0 for a variable can turn a permitted source into the literal 0, so substitution does not map instances of $\textsc{Tr}^-$ to instances. Substitution is admissible in O and M as an operation on derivable sequents (Theorem 2.2); what fails is the preservation of each rule instance. `reversal_core` makes the same correction to an earlier wording about the Lean calculus `ZPrf`. The Lean kernel keeps unrestricted sources so that its substitution theorem is a direct induction.

Remark 2.29 (Sources)*.* Theorems 2.25 and 2.27 are from `one_sided_transport`, which calls M the calculus $\mathcal{C}$ and reserves the letter $\mathcal{M}$ for the same seven rules with $\textsc{Tr}_2$ at sources other than the literal 0. The package proves the equivalence twice: by simulating that two-template rule with Theorem 2.25(iii) and quoting an earlier equivalence, and directly as above. In Lean the statement that I0 and $\textsc{Tr}^-$ recover Tr is `OKernel.trOneF`, for the class `OKernel`; the equivalences are `oprf_iff_nprf` for O and `mprf_iff_nprf`, `mprf_iff_iprf` for M. Theorem 2.26 is stated in `reversal_core`.

2.3.3 Reversal of the root equation

Define an operation on the root of a native formula:

$$(s = t)^\dagger := (t = s), \qquad (Q_y B)^\dagger := Q_y B, \qquad (D_{x,y} B)^\dagger := D_{x,y} B.$$

It does not descend into a matrix, it satisfies $A^{\dagger\dagger} = A$, and it commutes with substitution.

Definition 2.30 (The calculus X)**.** X has the eight rules Hyp, I0, IS, DE, DI, QI and

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash F(s)}{\Gamma \vdash F(t)^{\dagger}}\ \textsc{TrX} \quad (s \not\equiv 0),$$

$$\frac{\Gamma \vdash Q_y B(y) \qquad \Gamma \vdash B(0)}{\Gamma \vdash B(t)^{\dagger}}\ \mathrm{QX},$$

where $F(v)$ is an arbitrary native formula template, constant templates included.

Lemma 2.31 (Reversal)**.** *If* $\Gamma \vdash_{\mathsf{X}} A$*, then* $\Gamma \vdash_{\mathsf{X}} A^{\dagger}$*.*

Proof. Apply TrX with the first premise $c = 0$ and the constant template $F(v) := A$. □

Theorem 2.32 (Root-reversing eliminations)**.** X *and* N *are equivalent. The derivation of the rules of* N *in* X *uses neither IS nor an introduction rule nor DE.*

Proof. From X *to* N. For TrX, transport in the formula template F (Section 1.4) gives $F(t)$; if $F(t)$ is an equation, symmetry gives $F(t)^{\dagger}$, and otherwise $F(t)^{\dagger}$ is $F(t)$. For QX, use QE and the same step.

From N *to* X. The rule $\textsc{Tr}^{-}$ is the instance of TrX at the template $F(v) := \big(h(v) = r\big)$, since $F(t)^{\dagger}$ is $r = h(t)$. Hence Tr is derivable in X by Theorem 2.25(vii). The rule QE is QX followed by Theorem 2.31; the two reversals cancel. The six other rules are common. The constructions hold over arbitrary contexts, so they apply inside the premises of DE and of the two introductions. □

Only the equational instances of transport have to be recovered in X; transport in a compound template is used in the other direction, where it is available.

Remark 2.33 (Sources). Theorem 2.32 is the theorem of `reversal_core`, which proves the items of Theorem 2.25 directly for TrX, with the templates $v = a$, $J(v) = p$, $r = v$ and $f(s) = f(v)$, and derives $\textsc{Tr}_2$ from them. `one_sided_transport` observes that $\textsc{Tr}^{-}$ is the restriction of TrX to equations with one constant side, needs no reversal of arbitrary formulas and leaves QE unchanged. In Lean the calculus X is the module `NativeReversal`, adopted from the package, with `rprf_iff_nprf`, `reversal_iff_pra` and `pra_iff_reversal`. The package reports one comparison of proof sizes: a certificate of the same closed goal, symmetry of an equation with the source 0, has 114 instruction packets and 5435 bytes when produced by the method of Section 2.3.4 and 37 packets and 1259 bytes when produced with reversal.

2.3.4 The same restriction for the uncrossed rule

For transport in a formula template with an uncrossed conclusion the restriction of the source is also harmless, but the proof needs an induction. Let

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash F(s)}{\Gamma \vdash F(t)}\ \textsc{Tr}_F^{-} \qquad (s \not\equiv 0).$$

Definition 2.34 (The calculus Z)**.** Z is N_F with $\textsc{Tr}_F$ replaced by $\textsc{Tr}_F^{-}$; its rules are Hyp, I0, IS, $\textsc{Tr}_F^{-}$, QE, DE, DI and QI.

The sans-serif letter Z names the calculus; the italic letters Z and N name the zero test $Z(c) = \mathrm{sel}(c, 1, 0)$ and the parity term $N(n) = I(n, 0;\, u.\, Z(u))$ of Section 1.1.

In Z the two eliminations with a template can be merged into one rule whose first premise is selected by the source. For a template $F(v)$ put $M(s,t;F) := Q_v F(v)$ if $s \equiv 0$ and $M(s,t;F) := (s = t)$ otherwise; then

$$\frac{\Gamma \vdash M(s,t;F) \qquad \Gamma \vdash F(s)}{\Gamma \vdash F(t)}\ \mathrm{SE}$$

is QE when s is the literal 0 and TR_F^- otherwise. This is the form in which `source_dispatch` states the calculus, and the reason for its name.

Theorem 2.35 (Literal-zero sources of uncrossed transport)**.** Z *and* N *are equivalent.*

Every instance of TR_F^- is derivable in N by the Leibniz rule. For the converse it suffices to derive TR in Z; the seven other rules are common. The proof does not assume that substitution is admissible in Z: every auxiliary equation is derived at an arbitrary term by a derived rule applied at that term. By Theorem 2.4(i) and (ii), numerical cases and equation induction are derivable in Z, with no use of transport.

Lemma 2.36 (Equality at permitted sources)**.** Z *derives: $r = r$; from $p = q$ with $p \not\equiv 0$, the equations $q = p$ and $f(p) = f(q)$ for every term template f; from $p = q$ with $p \not\equiv 0$ and $p = r$, the equation $q = r$; from $r = s$ and $s = t$ with $s \not\equiv 0$, the equation $r = t$.*

Proof. Transport $J(r) = r$ along itself in the template $v = r$; the source is $J(r)$. For the other statements transport, along $p = q$ or along $s = t$, the equations $p = p$ in the template $v = p$, $f(p) = f(p)$ in the template $f(p) = f(v)$, $p = r$ in the template $v = r$, and $r = s$ in the template $r = v$. □

Lemma 2.37 (Joining through zero)**.** *If $p \not\equiv 0$ and $q \not\equiv 0$, then* Z *derives $p = q$ from $p = 0$ and $q = 0$.*

Proof. Transport $J(p) = J(p)$ along $p = 0$ in the template $J(v) = J(p)$; the result is $c = J(p)$. Symmetry at the source c gives $J(p) = c$, and transport along $J(p) = p$ in the template $v = c$ gives $p = c$. In the same way $q = c$. Symmetry at the source q gives $c = q$, and transport of $p = c$ along $c = q$ in the template $p = v$ gives $p = q$. The sources are p, q, c, $J(p)$ and $J(q)$. □

Define, with the selector of Section 1.1,

$$K(v) := \mathrm{sel}(v,0,1), \qquad H(v) := \mathrm{sel}(v,1,2), \qquad R(n) := I(n,0;\, u.\, H(u)), \qquad L(n) := N(R(n)).$$

In the standard model R takes the values $0,1,2,2,\ldots$ and L is 1 at 1 and 0 elsewhere; the proof derives what it needs and does not use this description.

Lemma 2.38 (A test for the argument one)**.** *For every term v,* Z *derives*

$$H(v) = SK(v), \quad R(1) = 1, \quad R(SSv) = 2, \quad N(1) = 1, \quad N(2) = 0, \quad L(1) = 1, \quad L(SSv) = 0, \quad L(Sv) = Z$$

Proof. All chains of equations below pass through middle terms with the root I or S, so each step is one of the operations of Theorem 2.36.

$H(v) = SK(v)$: by numerical cases on the matrix $H(y) = SK(y)$ at the term v. At 0: $H(0) = 1$ and $K(0) = 0$ by I0, hence $SK(0) = 1$ by congruence at the source $K(0)$, and $H(0) = SK(0)$ by symmetry at the source $SK(0)$ and transitivity through 1. At Sz: $H(Sz) = 2$ and $K(Sz) = 1$ by IS, hence $SK(Sz) = 2$, and the two equations are joined through 2.

$N(1) = Z(N(0))$ by IS, $N(0) = 0$ by I0, hence $Z(N(0)) = Z(0)$ by congruence at the source $N(0)$, and $Z(0) = 1$ by I0; so $N(1) = 1$. Likewise $N(2) = Z(N(1)) = Z(1) = 0$, the last by IS. In the same way $R(1) = H(R(0)) = H(0) = 1$.

IS and the first equation at the term $R(v)$ give $R(Sv) = H(R(v)) = SK(R(v))$. Then $R(SSv) = H(R(Sv)) = H(SK(R(v))) = 2$, by IS, congruence at the source $R(Sv)$, and IS for the selector.

$L(1) = N(R(1)) = N(1) = 1$ and $L(SSv) = N(R(SSv)) = N(2) = 0$, by congruence at the sources $R(1)$ and $R(SSv)$.

$L(Sv) = Z(v)$: by numerical cases on the matrix $L(Sy) = Z(y)$ at the term v. At 0, $L(1) = 1$ and $Z(0) = 1$ are joined through 1. At Sz, $L(SSz) = 0$ and $Z(Sz) = 0$, the latter by IS, are joined by Theorem 2.37; both left sides have the root I. □

Lemma 2.39 (Filter)**.** *For every term t, Z derives $t = \mathrm{sel}(Z(t), t, 0)$ in every context.*

Proof. By numerical cases on the matrix $y = \mathrm{sel}(Z(y), y, 0)$ at the term t. At 0: $Z(0) = 1$, so $\mathrm{sel}(Z(0), 0, 0) = \mathrm{sel}(1, 0, 0)$ by congruence at the source $Z(0)$, and $\mathrm{sel}(1, 0, 0) = 0$ by IS; symmetry at the left side, which has the root I, gives $0 = \mathrm{sel}(Z(0), 0, 0)$. At Sz: $Z(Sz) = 0$, so $\mathrm{sel}(Z(Sz), Sz, 0) = \mathrm{sel}(0, Sz, 0) = Sz$ by congruence at the source $Z(Sz)$ and I0, and symmetry at the left side gives the instance. □

Lemma 2.40 (Symmetry at the source zero)**.** *For every term t and every context Γ containing $0 = t$, Z derives $\Gamma \vdash t = 0$.*

Proof. If $t \equiv 0$, use reflexivity. Otherwise t is a permitted source. Put $A(n) := I(n, t; u.\, Su)$ and prove $n = A(n)$ by equation induction on n, with t as a parameter and $0 = t$ in the context.

Base. $A(0) = t$ by I0, and $t = A(0)$ by symmetry at the source $A(0)$. Transport of $0 = t$ along $t = A(0)$ in the template $0 = v$ gives $0 = A(0)$; the source is t.

Step. Assume $z = A(z)$ with z fresh. The source z is a variable, so congruence gives $Sz = SA(z)$. IS gives $A(Sz) = SA(z)$, and symmetry at the source $A(Sz)$ gives $SA(z) = A(Sz)$. Transport of $Sz = SA(z)$ along this equation in the template $Sz = v$ gives $Sz = A(Sz)$.

At the term 1 the induction gives $1 = A(1)$. IS gives $A(1) = SA(0)$, and congruence at the source $A(0)$ gives $SA(0) = St$; so $A(1) = St$, and transport at the source $A(1)$ gives $1 = St$. Congruence at the source 1 gives $L(1) = L(St)$. With $L(1) = 1$ and $L(St) = Z(t)$ from Theorem 2.38, the third operation of Theorem 2.36 at the source $L(1)$ gives $1 = L(St)$, and transport at the source $L(St)$ gives $1 = Z(t)$. Symmetry at the source 1 gives $Z(t) = 1$.

Transport of the equation $t = \mathrm{sel}(Z(t), t, 0)$ of Theorem 2.39 along $Z(t) = 1$ in the template $t = \mathrm{sel}(v, t, 0)$ gives $t = \mathrm{sel}(1, t, 0)$, and transport along the instance $\mathrm{sel}(1, t, 0) = 0$ of IS in the template $t = v$ gives $t = 0$. □

The chain of the proof is $0 = t \;\Rightarrow\; 1 = St \;\Rightarrow\; Z(t) = 1 \;\Rightarrow\; t = 0$. Numerical cases are used only for equations that hold in every context, and the one induction that keeps the assumption $0 = t$ has the literal equation $n = A(n)$ as its matrix.

Proof of Theorem 2.35. Symmetry holds in Z without restriction: at a source other than the literal 0 by Theorem 2.36, and at the source 0 by Theorem 2.40 and cut. Congruence from $s = t$: if $s \not\equiv 0$ use Theorem 2.36; if $s \equiv t \equiv 0$ use reflexivity; if $s \equiv 0$ and $t \not\equiv 0$, derive $t = 0$, then $f(t) = f(0)$ at the source t, and reverse. Transitivity from $r = s$ and $s = t$: if $s \not\equiv 0$ use Theorem 2.36; if $s \equiv 0$ and one of r, t is the literal 0, a premise is the conclusion; otherwise reverse $0 = t$ and apply Theorem 2.37 to $r = 0$ and $t = 0$. These constructions are not circular, since symmetry at the source 0 was proved before unrestricted congruence and transitivity. Finally, from $s = t$ and $h(s) = r$, congruence gives $h(s) = h(t)$, symmetry gives $r = h(s)$ and transitivity gives $r = h(t)$, which is Tr. □

Remark 2.41 (Sources and comparison). Theorem 2.35 and its lemmas are from `source_dispatch`, whose calculus has DP in place of QI (Theorem 2.6). The package also derives unrestricted transport in a formula template inside its calculus, by induction on the template; here that step is the Leibniz rule of N, since the equivalence needs Tr only. In Lean the theorem is `zprf_iff_nprf`, and the recovery of unrestricted transport is `ZKernel.trF` for a class `ZKernel` that has equation induction and no logical rule. The contrast with Theorem 2.25 is the point of the two later packages: for the crossed rule, and for the reversing rule, a constant template reverses an equation in one step, and neither the test terms nor an induction is needed.

2.3.5 Replacement with no template

A transport rule can be given no template at all. For terms s, t, u let $\mathrm{Rep}_s^t(u)$ be the result of one traversal of u from the root: a subterm that is identical with s is replaced by t and is not traversed further; at any other subterm the constructor is kept and its immediate subterms are traversed; a variable or a 0 that is not replaced is unchanged. Under the binder of an iterator the bound variable is renamed away from s and t, so that only free occurrences of s are matched and no variable of t is captured. The test at a subterm precedes the descent, inserted copies of t are not traversed, and the operation is total even when t contains s. It is one pass and not rewriting to a normal form. In particular $\mathrm{Rep}_s^t(s) = t$.

Definition 2.42 (The calculi G and G_7)**.** The rule of global replacement is

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash u = r}{\Gamma \vdash r = \mathrm{Rep}_s^t(u)}\ \textsc{Rep}.$$

G is N with Tr replaced by Rep; its rules are Hyp, I0, IS, Rep, QE, DE, DI and QI. G_7 is W with Tr replaced by Rep; its seven rules are Hyp, I0, IS, Rep, QE, DE and DI_{w}.

The rule cannot choose which occurrences of s to replace. A chosen template is recovered by planting a marker that cannot occur by accident.

Lemma 2.43 (Equality from I0 and Rep)**.** *From I0 and Rep alone the following are derivable: from $s = t$ and $s = u$ the equation $u = t$; reflexivity; symmetry; transitivity.*

Proof. Rep applied to $s = t$ and $s = u$ concludes $u = \mathrm{Rep}_s^t(s)$, which is $u = t$. This joining, applied to $J(a) = a$ and $J(a) = a$, gives $a = a$; applied to $a = a$ and $a = b$ it gives $b = a$. For transitivity let $a = b$ and $b = e$ be given: symmetry gives $b = a$, and the joining applied to $b = e$ and $b = a$ gives $a = e$. □

Only matches at the root are used in this proof. Let $|T|$ be the number of nodes of a term T, the body of every iterator included; renaming of variables does not change it.

Lemma 2.44 (Exact marking)**.** *Let $h(v)$ be a term template, s and t terms, and*

$$n := 1 + |h| + |s| + |t|, \qquad K := I(0, s;\, w.\, S^n 0).$$

Then the occurrences of K in $h(K)$ are exactly the copies inserted for v, and

$$\mathrm{Rep}_K^s(h(K)) = h(s), \qquad \mathrm{Rep}_K^t(h(K)) = h(t).$$

Proof. K has more nodes than h. Two subterm occurrences of a term are disjoint or nested. An occurrence in $h(K)$ of a subterm identical with K that is disjoint from all inserted copies lies in a part of h and has at most $|h|$ nodes. One that lies properly inside an inserted copy has fewer nodes than K. One that properly contains an inserted copy has more nodes than K. None of these is identical with K, so the occurrences are the inserted copies. Below a binder of h the inserted copy and the term compared with it are subject to the same renaming, which preserves the number of nodes. The traversal replaces exactly these copies and stops at each, which gives the two identities. □

The identities are facts about syntax; no rule of the calculus is applied in them. The term K keeps the free variables of s, and $K = s$ is an instance of I0.

Lemma 2.45 (Chosen templates from global replacement)**.** *From I0 and* Rep *alone: from $s = t$ the equation $h(s) = h(t)$, for every term template h; and the rule* Tr*.*

Proof. Let K be as in Theorem 2.44. From $K = s$ and $s = t$, transitivity gives $K = t$. Reflexivity gives $h(K) = h(K)$. Rep applied to $K = s$ and $h(K) = h(K)$ concludes $h(K) = h(s)$, and applied to $K = t$ and $h(K) = h(K)$ it concludes $h(K) = h(t)$, by Theorem 2.44. The joining of Theorem 2.43, applied to $h(K) = h(t)$ and $h(K) = h(s)$, gives $h(s) = h(t)$. For Tr, from $s = t$ and $h(s) = r$: symmetry gives $r = h(s)$ and transitivity gives $r = h(t)$. □

For example let $s = t$ be given and let $h(v) := I(v, s;\, w.\, s)$, in which the two displayed copies of s are to stay. Replacement of s in $h(s)$ would change all three copies. In $h(K) = I(K, s;\, w.\, s)$ the marker is too large to occur in the two fixed copies, and the two replacements produce $I(s, s;\, w.\, s)$ and $I(t, s;\, w.\, s)$.

Lemma 2.46 (Rep in N)**.** *Every instance of* Rep *is an instance of* Tr*.*

Proof. The occurrences of s that the traversal of u replaces are disjoint. Replace them by a fresh variable v to obtain a term template $h(v)$; then $u \equiv h(s)$ and $\mathrm{Rep}_s^t(u) \equiv h(t)$. If no occurrence is replaced, h is constant. Tr with this template, applied to $s = t$ and $h(s) = r$, concludes $r = h(t)$. □

Thus Rep is Tr with the template computed from the two premises.

Theorem 2.47 (Replacement with no template)**.** G, G_7 *and* N *are equivalent.*

Proof. Rep is derivable in N by Theorem 2.46, and Tr is derivable in G and in G_7 by Theorem 2.45. This settles G. For G_7 the introductions are treated as in Theorem 2.17: an instance of DI_w is derivable in N, and QI and DI are derivable in G_7 by the proof given there, with the alias c, transport in an equation template derived from Tr, and cut, which is admissible in G_7 by Theorem 2.3. □

Remark 2.48 (Substitution and certificates)*.* A substitution can create new occurrences of the source in the second premise of Rep, so it does not map instances to instances; injective renaming does. Substitution is admissible through Theorem 2.2. An instance of Rep in a certificate consists of its two premise derivations; the price is paid in the derivations, which grow by the marker.

Remark 2.49 (Sources)*.* The section is `template_free`, whose calculus is G_7. In Lean the calculus G is `TPrf`, with `tprf_iff_nprf`; the combination G_7 with the single introduction is not formalized. The compact implementation of the package works with the restricted inventory of Theorem 2.53 and uses the aliases $C_s := I(2, 1;\, u.\, s)$ and $K := I(0, C_s;\, u.\, S^n 0)$, so that $C_s = s$ is a permitted instance of IS and $K = C_s$ a permitted instance of I0, also when s is the literal 0.

2.3.6 Replacement of one occurrence

A *term context with one hole* $C[\,]$ is a term with exactly one occurrence of a hole; the hole may lie in the body of an iterator, and $C[s]$ plugs s into the hole after the bound variables of C have been renamed away from s. The context consisting of the hole alone is allowed.

Definition 2.50 (The calculi L and N_u)**.** L is N with Tr replaced by

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash C[s] = r}{\Gamma \vdash C[t] = r}\ \mathrm{LOR},$$

for a term context C with one hole. N_u is N with Tr replaced by the same rule for an arbitrary term template $h(v)$ in place of C: from $s = t$ and $h(s) = r$ infer $h(t) = r$. Each has the rules Hyp, I0, IS, its replacement rule, QE, DE, DI and QI.

Theorem 2.51 (One occurrence suffices)**.** L, N_u *and* N *are equivalent.*

Proof. An instance of LOR is an instance of the rule of N_u, with the template obtained by putting a fresh variable into the hole. The rule of N_u is derivable in N: Tr gives $r = h(t)$ and symmetry gives $h(t) = r$.

It remains to derive Tr in L. LOR with the context consisting of the hole, applied to $J(a) = a$ and $J(a) = a$, gives $a = a$; applied to $s = t$ and $s = s$ it gives $t = s$. Now let $s = t$ and $h(s) = r$ be given. Rename the bound variables of h away from s and t and number the occurrences of v in h from 1 to k. Let h_j be h with the first j occurrences replaced by t and the others by s, and let C_j be h with the first $j-1$ occurrences replaced by t, the j-th by the hole and the others by s. Then $h_{j-1} \equiv C_j[s]$ and $h_j \equiv C_j[t]$, and C_j has one hole. Hence k applications of LOR lead from $h(s) = r$, which is $h_0 = r$, to $h_k = r$, which is $h(t) = r$, and symmetry gives $r = h(t)$. The occurrences are those of the template variable; a subterm of h that happens to be identical with s is part of every C_j and is not replaced. □

Remark 2.52 (Sources)*.* The theorem for L is the auxiliary note of `kernel_refinement`, which states it against N_2: the occurrences in the left template are replaced one at a time, the equation is reversed, the occurrences in the right template are replaced, and the equation is reversed again. It answers the question of `one_context` whether a restriction to one occurrence is adequate under the binder of the iterator. In Lean, L is `LPrf` with `lprf_iff_nprf`; contexts with one hole are `Tm.Hole`, and the chain $h_0, \ldots, h_k$ is `Reach.subst`, where below a binder the hole and the bound variable are exchanged (`Reach.iterH`). The calculus N_u is `UPrf` with `uprf_iff_nprf`. The package measures no checker for L: a traversal by position has a cost of its own, and the compact checkers share one opening routine between transport and QE.

2.3.7 A restricted inventory of computation instances

Several compact checkers of the packages accept a computation equation only when its seed, and for IS also the predecessor of its counter, is not the literal 0. Call an instance of I0 *permitted* if $a \not\equiv 0$, and an instance of IS permitted if $n \not\equiv 0$ and $a \not\equiv 0$. Let $c_* := I(2, 1;\, u.\, 0)$.

Lemma 2.53 (Repair of computation instances)**.** *Let a calculus have the permitted instances of I0 and IS and a transport rule from which transport in an equation template at sources other than the literal* 0 *is derivable without a computation equation. Then it derives every instance of I0 and of IS.*

Proof. $c_* = 0$ is a permitted instance of IS: the predecessor of the counter and the seed are 1, and the body is the constant 0. Write an instance that is not permitted as $P(0)$, where the equation template $P(v)$ has v at the offending positions:

$$I(0, v;\, u.\, h) = v, \qquad I(Sn, v;\, u.\, h) = h[I(n, v;\, u.\, h)/u],$$
$$I(Sv, a;\, u.\, h) = h[I(v, a;\, u.\, h)/u], \qquad I(Sv, v;\, u.\, h) = h[I(v, v;\, u.\, h)/u].$$

The variable v is fresh, and other occurrences of 0 are left alone. In each case $P(c_*)$ is a permitted instance. Transport in the template P along $c_* = 0$, at the source $c_* \not\equiv 0$, gives $P(0)$. □

The hypothesis on the transport rule holds for TR, TR^{-} (Theorem 2.25(iii)), TRX, TR_2 and TR^{-}_{F}. Hence every result of this chapter that uses $c = I(0, 0;\, u.\, 0)$ or another computation instance with a literal 0 holds for the restricted inventory as well: the instance is first derived by the lemma. The repair is one transport on two computation instances, and its only source is c_*.

Remark 2.54 (Sources). The lemma is proved in `witness_rule`, where c_* serves also as the second name for zero in Theorem 2.17, and in `source_dispatch`, which checks that the repair does not use transport at the source 0 before Theorem 2.35 is available. `reversal_core` and `template_free` use it in their certificate producers. The restricted inventory itself belongs to an earlier package, `certified_zero`.

2.4 Induction and uniqueness of iteration

This section draws on `scalar_uniqueness`, `orbit_closure` and `crossed_uniqueness`.

Induction for literal equations can be replaced by a rule that has no induction hypothesis in its premise: a term that satisfies the recurrence of an iterator is that iterator. The three packages state the rule in three forms. `scalar_uniqueness` concludes the equation at a supplied target; this is the rule UR below, and Sections 2.4.1 to 2.4.7 prove that it replaces equation induction. `orbit_closure` concludes the closure of the equation, with no target, and isolates what the restriction of the rule to the body Su already gives. `crossed_uniqueness` adds a premise for the seed and crosses the conclusion; on fully normalized terms that rule also replaces transport and the two computation equations. The second and third forms are treated in Section 2.4.8, which reduces each to UR; the calculus of `crossed_uniqueness` is the one calculus of the chapter that is stated on normal forms.

2.4.1 The rule and the two theorems

Let $f(z)$ and $h(u)$ be term templates, and write $h(f(z))$ for $h[f(z)/u]$. The rule of *uniqueness of iteration* is

$$\frac{\Gamma \vdash f(Sz) = h(f(z))}{\Gamma \vdash f(t) = I(t, f(0);\, u.\, h(u))}\ \mathrm{UR},$$

where z is not free in Γ, in t or in h. The body h may have other free variables, which are parameters. The premise is derived in Γ itself, with no added assumption; the seed $f(0)$ is read off the template, so there is no premise for the base; and the body receives one state and no counter. The rule is a scheme over terms: z is a numerical variable and nothing ranges over functions.

Definition 2.55 (The calculus U and the theory T_{UR})**.** U is N with QI replaced by UR; its eight rules are Hyp, I0, IS, Tr, QE, DE, DI and UR. The ordinary theory T_{UR} is T_{eq} (Section 1.5) with its rule of induction for literal equations replaced by UR, read for contexts of ordinary formulas: intuitionistic logic with equality, the two computation equations, coverage $x = 0 \vee \exists y\, x = Sy$, separation $0 = S0 \to \bot$, and UR.

In U the introduction DI treats atomic and compound matrices alike, and no rule looks at a witness; substitution maps instances of UR to instances after a renaming of z (Theorem 2.3).

Definition 2.56 (Successor uniqueness)**.** SU is the rule UR restricted to the body $h(u) := Su$: from $\Gamma \vdash f(Sz) = Sf(z)$ infer $\Gamma \vdash f(t) = I(t, f(0);\, u.\, Su)$. The theory T_{SU} is T_{UR} with UR replaced by SU.

An instance of SU supplies no body. Every derivation of T_{SU} is a derivation of T_{UR}, and the lemmas below say which of them use SU only.

Proposition 2.57 (Uniqueness from induction)**.** *UR is a derived rule of* N *and of* T_{eq}.

Proof. Let $g(y) := I(y, f(0);\, u.\, h(u))$ and apply equation induction to $f(y) = g(y)$. Base: $g(0) = f(0)$ by I0, and symmetry. Step: assume $f(z) = g(z)$. The premise gives $f(Sz) = h(f(z))$, congruence gives $h(f(z)) = h(g(z))$, and IS gives $g(Sz) = h(g(z))$; hence $f(Sz) = g(Sz)$. □

In Lean this derived rule is `NPrf.uniq`.

Theorem 2.58 (Uniqueness of iteration in place of equation induction)**.** T_{UR} *and* T_{eq} *derive the same sequents of ordinary formulas. Consequently* T_{UR} *is bi-interpretable with* iPRA *with whole domains, actual equality and identity numerical composites.*

Theorem 2.59 (The native form)**.** U *and* N *are equivalent.*

One half of each theorem is Theorem 2.57. The other half, the derivation of induction for literal equations from UR, occupies Sections 2.4.2 to 2.4.7. Its order avoids a circle: uniqueness gives induction for equations of the form $q(n) = 0$ and, separately, a predecessor; truncated subtraction and an explicit accumulator give a term that reflects equality; and only then is induction for arbitrary equations recovered. Pairing and the compiler for primitive recursion (Theorem 1.62) are not used; they become available afterwards. All derivations in Sections 2.4.2 to 2.4.7 are in T_{UR}, over an arbitrary fixed context Γ of ordinary formulas, and variables that are generalized or used as eigenvariables are not free in Γ.

2.4.2 First consequences of uniqueness

Lemma 2.60 (Common update, constancy, identity)**.** (i) *If* $f(Sz) = h(f(z))$, $g(Sz) = h(g(z))$ *and* $f(0) = g(0)$, *then* $f(t) = g(t)$.

(ii) *If* $f(Sz) = f(z)$, *then* $f(t) = f(0)$.

(iii) $I(n, 0;\, u.\, Su) = n$.

(iv) *If* $h(c) = c$, *then* $I(n, c;\, u.\, h(u)) = c$.

Proof. (i) UR for f and for g gives two iterators that differ only in the equal seeds. (ii) is (i) for f and the constant template $f(0)$, with $h(u) := u$. (iii) is UR for $f(z) := z$ and $h(u) := Su$, whose premise is reflexivity. (iv) is UR for the constant template c. □

Part (iii), and part (i) for the body Su, are derivable in T_{SU}. Part (ii) is uniqueness for the body u, and it is not an instance of SU.

Put $a + b := I(b, a;\, u.\, Su)$, $2a := a + a$ and $3a := a + a + a$. The computation equations give $a + 0 = a$ and $a + Sb = S(a + b)$, so $a + 1 = Sa$.

Lemma 2.61 (Addition and iteration)**.** *Addition is associative and commutative with the neutral element* 0, *and* $Sa + n = S(a + n)$. *For every body* h,

$$I(m + n, a;\, u.\, h) = I(n, I(m, a;\, u.\, h);\, u.\, h), \qquad I(Sn, a;\, u.\, h) = I(n, h(a);\, u.\, h).$$

Proof. Each statement is Theorem 2.60(i) or UR for functions of n. $0 + n = n$ is Theorem 2.60(iii). $S(a + n)$ has the seed Sa and the update S, hence equals $Sa + n$. The term $n + a$ has the seed $0 + a = a$ and satisfies $Sn + a = S(n + a)$, so it has the update S and equals $a + n$. The term $a + (b + n)$ has the seed $a + b$ and the update S, hence equals $(a + b) + n$. The term $I(m + n, a;\, u.\, h)$ has the seed $I(m, a;\, u.\, h)$ and the update h. The term $I(Sn, a;\, u.\, h)$ has the seed $I(1, a;\, u.\, h) = h(a)$ and the update h. □

The laws of addition in this lemma compare terms with the update S only, so they are derivable in T_{SU}; the two equations for an arbitrary body h use UR for h.

2.4.3 Retractions and induction for zero equations

Lemma 2.62 (Rigidity of retractions)**.** *Let $e(x)$ be a term template such that*

$$e(0) = 0, \qquad e(e(x)) = e(x), \qquad e(Se(x)) = Se(x)$$

are derivable for a variable x not free in Γ. Then $e(t) = t$ for every term t. The derivation is in T_{SU} and does not use separation.

Proof. Put $h(u) := e(Se(u))$ and $r(n) := I(n, 0; u.\, h(u))$. First, $e(r(n)) = r(n)$, by coverage on n. At 0 it is $e(0) = 0$, since $r(0) = 0$. At Sm, the term $r(Sm) = h(r(m)) = e(Se(r(m)))$ is a value of e, and the second equation gives

$$e(r(Sm)) = e\big(e(Se(r(m)))\big) = e(Se(r(m))) = r(Sm).$$

The third equation now gives $r(Sn) = e(Se(r(n))) = Se(r(n)) = Sr(n)$, and $r(0) = 0$. SU and Theorem 2.60(iii) give $r(n) = n$, and then $e(n) = e(r(n)) = r(n) = n$. □

Lemma 2.63 (Induction from a numerical split)**.** *Let $A(n)$ be an ordinary formula with $\Gamma \vdash A(0)$ and $\Gamma, A(z) \vdash A(Sz)$. Suppose that $\Gamma \vdash A(n) \vee \neg A(n)$ and that a term template $e(n)$ satisfies $\Gamma, A(n) \vdash e(n) = n$ and $\Gamma, \neg A(n) \vdash e(n) = 0$, with n not free in Γ. Then $\Gamma \vdash A(t)$ for every term t.*

Proof. By cases, $A(e(n))$: if $A(n)$, then $e(n) = n$; otherwise $e(n) = 0$ and $A(0)$ holds. The first property of e at the argument $e(n)$ gives $e(e(n)) = e(n)$. The step at $e(n)$ gives $A(Se(n))$, hence $e(Se(n)) = Se(n)$. From $A(0)$, $e(0) = 0$. Theorem 2.62 gives $e(n) = n$, and $A(e(n))$ becomes $A(n)$. □

The lemma is a statement about a split that has been justified; it gives no induction for formulas that are not decided by a term.

Lemma 2.64 (Induction for zero equations)**.** *Let $q(n)$ be a term template. If $\Gamma \vdash q(0) = 0$ and $\Gamma, q(z) = 0 \vdash q(Sz) = 0$ with z fresh, then $\Gamma \vdash q(t) = 0$.*

Proof. Separation gives $Sv \neq 0$: from $Sv = 0$, the template $\mathrm{sel}(w, 0, 1)$ gives $1 = 0$. With coverage, $q(n) = 0$ is decided. The template $e(n) := \mathrm{sel}(q(n), n, 0)$ equals n if $q(n) = 0$ and equals 0 if $q(n)$ is a successor. Theorem 2.63 applies. □

Theorems 2.63 and 2.64 use uniqueness only through Theorem 2.62. Hence:

Theorem 2.65 (Zero equations from successor uniqueness)**.** *T_{SU} derives induction for zero equations, in the form of Theorem 2.64, over arbitrary contexts of ordinary formulas.*

Remark 2.66 (Sources). Theorems 2.62 to 2.64 are reproduced in `scalar_uniqueness` from an earlier package, `scope_kernel`, which leaves the predecessor and the reflection of equality as an open bootstrap. The next three subsections close it. In those two packages the first step of the proof of Theorem 2.62 is an instance of UR for the body $e(Se(u))$: the template $e(r(n))$ has the seed 0 and satisfies $e(r(Sn)) = h(r(n)) = h(e(r(n)))$. The proof given above, with coverage in place of that instance, is from `orbit_closure`, and Theorem 2.65 is its theorem. The decision of $q(n) = 0$ in Theorem 2.64 needs separation; no other step of the three lemmas does.

2.4.4 A predecessor from uniqueness

The construction follows the predecessor of Theorem 1.48 (parity, three-cycle, ramp, comparator, saturating counter), with a ramp modified so that every induction of the original becomes an instance of Theorem 2.60. It uses coverage and does not use separation.

Coverage and the selector equations give, by cases on e,

$$\mathrm{sel}(e, Sa, Sb) = S\ \mathrm{sel}(e, a, b). \tag{2.1}$$

Recall $Z(c) = \mathrm{sel}(c, 1, 0)$ and $N(n) = I(n, 0;\, u.\, Z(u))$, and define

$$j(u) := Z(u)+2N(u), \qquad M(n) := I(n, 0;\, u.\, j(u)), \qquad g(u) := u+N(M(u))+1, \qquad G(n) := I(n, 0;\, u.\, g(u)).$$

Lemma 2.67 (Two clocks)**.** (i) $N(n) = 0 \vee N(n) = 1$*;* $Z(Z(N(n))) = N(n)$*;* $N(n) + Z(N(n)) = 1$*;* $N(N(n)) = N(n)$*;* $N(n+2) = N(n)$*.*

(ii) $N(2n) = 0$*,* $N(2n+1) = 1$ *and* $N(3n) = N(n)$*.*

(iii) $M(n) = 0 \vee M(n) = 1 \vee M(n) = 2$*;* $j(0) = 1$*,* $j(1) = 2$*,* $j(2) = 0$*;* $M(n+3) = M(n)$*.*

(iv) $M(3n) = 0$ *and* $M(3n+1) = 1$*.*

Proof. (i) By coverage on n: $N(0) = 0$, and $N(Sm) = Z(N(m))$ is a value of Z, which is 1 at 0 and 0 at a successor. The next three equations hold in the two cases $N(n) = 0$ and $N(n) = 1$, by $Z(0) = 1$, $Z(1) = 0$, $N(0) = 0$ and $N(1) = Z(N(0)) = 1$. Two computations give $N(n+2) = Z(Z(N(n)))$.

(ii) The template $N(2n)$ satisfies $N(2(Sn)) = N(2n+2) = N(2n)$, so Theorem 2.60(ii) gives $N(2n) = N(0) = 0$; then $N(2n+1) = Z(N(2n)) = 1$. The template $N(3n)$ has the seed 0 and satisfies $N(3(Sn)) = N(3n+3) = Z(N(3n+2)) = Z(N(3n))$, so UR gives $N(3n) = I(n, 0;\, u.\, Z(u)) = N(n)$.

(iii) $j(0) = Z(0) + 2N(0) = 1$. At a successor u, $Z(u) = 0$ and $2N(u)$ is 0 or 2. So j takes values in $\{0, 1, 2\}$, and so does M, by coverage on n. $j(1) = 0 + 2N(1) = 2$ and $j(2) = 0 + 2N(2) = 0$, since $N(2) = Z(N(1)) = 0$. Three computations give $M(n+3) = j(j(j(M(n))))$, and j permutes $0, 1, 2$ cyclically.

(iv) $M(3(Sn)) = M(3n+3) = M(3n)$, so Theorem 2.60(ii) gives $M(3n) = M(0) = 0$, and $M(3n+1) = j(M(3n)) = 1$. □

Lemma 2.68 (The ramp)**.** (i) $M(g(u)) = Z(M(u))$*.*

(ii) $M(G(n)) = N(n)$*.*

(iii) $G(Sn) = G(n) + N(n) + 1$*,* $G(n+2) = G(n) + 3$ *and* $G(n+3) = G(n) + N(n) + 4$*.*

(iv) $G(2n) = 3n$ *and* $G(3n) = 3G(n) + N(n)$*.*

Proof. (i) By the three cases for $M(u)$. If $M(u) = 0$, then $g(u) = Su$ and $M(Su) = j(0) = 1 = Z(0)$. If $M(u) = 1$, then $g(u) = SSu$ and $M(SSu) = j(j(1)) = 0 = Z(1)$. If $M(u) = 2$, then $N(2) = 0$, $g(u) = Su$ and $M(Su) = j(2) = 0 = Z(2)$.

(ii) The template $M(G(n))$ has the seed $M(0) = 0$ and, by (i), satisfies $M(G(Sn)) = M(g(G(n))) = Z(M(G(n)))$. UR gives $M(G(n)) = I(n, 0;\, u.\, Z(u)) = N(n)$.

(iii) $G(Sn) = g(G(n)) = G(n) + N(M(G(n))) + 1 = G(n) + N(N(n)) + 1$, and $N(N(n)) = N(n)$. Two steps give $G(n+2) = G(n) + N(n) + 1 + Z(N(n)) + 1 = G(n) + 3$, by $N(n) + Z(N(n)) = 1$. A third step gives $G(n+3) = G(n) + 3 + N(n+2) + 1 = G(n) + N(n) + 4$.

(iv) $G(2n)$ and $3n$ have the seed 0 and the update $v \mapsto v + 3$, by (iii); Theorem 2.60(i) gives $G(2n) = 3n$. For the second equation put

$$f(n) := G(3n), \qquad k(n) := 3G(n) + N(n), \qquad \kappa(v) := v + M(v) + 4.$$

By (ii) and Theorem 2.67(ii), $M(f(n)) = N(3n) = N(n)$. By Theorem 2.67(iv) and the two cases for $N(n)$, $M(k(n)) = N(n)$. By (iii), $f(Sn) = G(3n+3) = G(3n)+N(3n)+4 = \kappa(f(n))$. By (iii) and Theorem 2.67(i),

$$k(Sn) = 3G(n) + 3N(n) + 3 + Z(N(n)) = 3G(n) + 2N(n) + 4 = \kappa(k(n)).$$

Both templates have the seed 0, and Theorem 2.60(i) gives $f(n) = k(n)$. □

The update g differs from the update $u + M(u) + 1$ of the ramp in Theorem 1.48 away from the orbit of 0; the difference is what makes (i) hold at every argument, so that (ii) is an instance of UR with no induction hypothesis.

Now define the comparison step, the comparator, the saturating step and the predecessor:

$$\chi(u) := \mathrm{sel}(N(u), G(u), u), \qquad \varepsilon(x) := I(x, 1;\, u.\, 2u), \qquad \lambda(x, y) := N\big(I(y, \varepsilon(x);\, u.\, \chi(u))\big),$$
$$\beta_m(u) := \mathrm{sel}(\lambda(Su, m), u, Su), \qquad \sigma(t, m) := I(t, 0;\, u.\, \beta_m(u)),$$
$$P(m) := I(m, 0;\, u.\, \mathrm{sel}(\lambda(SSu, m), u, Su)).$$

Lemma 2.69 (The comparator)**.** (i) $\chi(2u) = 3u$, $\chi(1) = 1$ *and* $\chi(3u) = 3\chi(u)$.

(ii) $I(y, 3a;\, u.\, \chi(u)) = 3\, I(y, a;\, u.\, \chi(u))$ *and* $I(y, 1;\, u.\, \chi(u)) = 1$.

(iii) $\lambda(0, y) = 1$, $\lambda(Sx, 0) = 0$ *and* $\lambda(Sx, Sy) = \lambda(x, y)$.

Proof. (i) $N(2u) = 0$ and $G(2u) = 3u$ give $\chi(2u) = 3u$; $N(1) = 1$ gives $\chi(1) = 1$. For the third equation, $N(3u) = N(u)$. If $N(u) = 0$, then $\chi(3u) = G(3u) = 3G(u)+N(u) = 3G(u) = 3\chi(u)$. If $N(u) = 1$, then $\chi(3u) = 3u = 3\chi(u)$.

(ii) The template $3\, I(y, a;\, u.\, \chi(u))$ has the seed $3a$ and, by (i), the update χ; UR gives the first equation. The second is Theorem 2.60(iv).

(iii) $\varepsilon(0) = 1$, so $\lambda(0, y) = N(1) = 1$ by (ii). $\lambda(Sx, 0) = N(\varepsilon(Sx)) = N(2\varepsilon(x)) = 0$. By the second equation of Theorem 2.61, by (i) and by (ii),

$$I(Sy, 2\varepsilon(x);\, u.\, \chi(u)) = I(y, \chi(2\varepsilon(x));\, u.\, \chi(u)) = I(y, 3\varepsilon(x);\, u.\, \chi(u)) = 3\, I(y, \varepsilon(x);\, u.\, \chi(u)),$$

and $N(3v) = N(v)$ gives $\lambda(Sx, Sy) = \lambda(x, y)$. □

Theorem 2.70 (Predecessor from uniqueness)**.** *The computation equations, coverage and UR derive* $P(0) = 0$ *and* $P(Sm) = m$*, and hence that* $Sx = Sy$ *implies* $x = y$.

Proof. By (2.1) and Theorem 2.69(iii),

$$\beta_{Sm}(Su) = \mathrm{sel}(\lambda(Su, m), Su, SSu) = S\beta_m(u), \qquad \beta_{Sm}(0) = \mathrm{sel}(\lambda(0, m), 0, 1) = 1.$$

As functions of t, the terms $\sigma(St, Sm)$ and $S\sigma(t, m)$ have the seed 1 and the update β_{Sm}: for the first by IS, with $\sigma(1, Sm) = \beta_{Sm}(0) = 1$, and for the second by $S\beta_m(\sigma(t, m)) = \beta_{Sm}(S\sigma(t, m))$. Theorem 2.60(i) gives

$$\sigma(St, Sm) = S\sigma(t, m). \tag{2.2}$$

The template $\lambda(Sn, n)$ satisfies $\lambda(SSn, Sn) = \lambda(Sn, n)$ and has the value $\lambda(1, 0) = 0$ at 0, so $\lambda(Sn, n) = 0$ by Theorem 2.60(ii). The template $\sigma(n, n)$ has the seed 0 and, by (2.2), the update S, so $\sigma(n, n) = n$ by UR and Theorem 2.60(iii). Hence

$$\sigma(Sn, n) = \beta_n(\sigma(n, n)) = \mathrm{sel}(\lambda(Sn, n), n, Sn) = n. \tag{2.3}$$

$P(0) = 0$ is an instance of I0. For a variable m put $\sigma'(t, m) := I(t, 0;\, u.\, \mathrm{sel}(\lambda(SSu, Sm), u, Su))$, so that $P(Sm)$ is $\sigma'(Sm, m)$. By Theorem 2.69(iii) the body of σ' equals $\beta_m(u)$ at every u, so $\sigma'(t, m)$ and $\sigma(t, m)$ have the same seed and the same update, and Theorem 2.60(i) gives $\sigma'(t, m) = \sigma(t, m)$. With (2.3), $P(Sm) = \sigma(Sm, m) = m$. Applying P to $Sx = Sy$ gives $x = y$. □

Every appeal to induction in the construction of Theorem 1.48 has been replaced by an instance of Theorem 2.60, and every case distinction uses coverage and a finite range.

2.4.5 Truncated subtraction and diagonal descent

Put $d(a,b) := I(b,a;\, u.\, P(u))$ and $\mathrm{sg}(a) := \mathrm{sel}(a,0,1)$.

Lemma 2.71 (Truncated subtraction)**.** *$d(a,0) = a$, $d(a,Sb) = P(d(a,b))$, $d(Sa,Sb) = d(a,b)$, $d(a,a) = 0$, $d(0,a) = 0$, $P(d(Sn,a)) = d(n,a)$ and $d(a+n,n) = a$. Addition is cancellative: $a+n = b+n$ implies $a = b$.*

Proof. The first two are the computation equations. As functions of b, the terms $d(Sa,Sb)$ and $d(a,b)$ have the seed a, by $d(Sa,1) = P(Sa) = a$, and the update P. The template $d(a,a)$ satisfies $d(Sa,Sa) = d(a,a)$ and is 0 at 0; Theorem 2.60(ii) applies. The template $d(0,a)$ and the constant 0 have the seed 0 and the update P. For the sixth equation use coverage on a: at 0 it is $P(Sn) = n$, and at Sa' it is $P(d(n,a')) = d(n,Sa')$. The template $d(a+n,n)$ satisfies $d(a+Sn,Sn) = d(a+n,n)$ and is a at 0. Cancellation follows by applying $d(\,\cdot\,,n)$. □

Lemma 2.72 (Diagonal descent)**.** *Let $q(x_1,\dots,x_r)$ be a term template, $r \geq 1$, such that*

$$q(Sx_1,\dots,Sx_r) = q(x_1,\dots,x_r)$$

and such that $q(x_1,\dots,x_r) = 0$ whenever some x_i is 0. Then $q(x_1,\dots,x_r) = 0$.

Proof. Coverage in each argument gives $q(P(x_1),\dots,P(x_r)) = q(x_1,\dots,x_r)$: if some x_i is 0, both sides are 0, and otherwise the first hypothesis applies. The template $f(k) := q(d(x_1,k),\dots,d(x_r,k))$ therefore satisfies $f(Sk) = f(k)$, and Theorem 2.60(ii) gives $f(k) = f(0)$. At $k = x_1$ the first argument is $d(x_1,x_1) = 0$, so $f(x_1) = 0$; and $f(0)$ is $q(x_1,\dots,x_r)$. □

The lemma uses uniqueness for one scalar state; it does not quantify over tuples and uses no rule for simultaneous recursion.

A term is *Boolean* if it is derivably 0 or 1. Put $\mathrm{or}(p,q) := \mathrm{sel}(p,q,1)$ and $\mathrm{xor}(p,q) := \mathrm{sel}(p,q,Z(q))$. For Boolean p and q, four cases show that $\mathrm{xor}(p,q) = 0$ implies $p = q$.

Lemma 2.73 (Three Boolean identities)**.** (i) $Z(d(Sn,a)) = \mathrm{sg}(d(a,n))$.

(ii) $Z\big(d(d(x,a),d(b,a))\big) = \mathrm{or}\big(Z(d(x,a)),Z(d(x,b))\big)$.

(iii) $\mathrm{or}(\mathrm{sg}(u),\mathrm{sg}(v)) = \mathrm{sg}(u+v)$.

Proof. (i) Let $q(n,a)$ be the xor of the two sides, which are Boolean. By Theorem 2.71, $q(Sn,Sa) = q(n,a)$. If $a = 0$, the sides are $Z(Sn) = 0$ and $\mathrm{sg}(0) = 0$. If $n = 0$ and $a = Sb$, the sides are $Z(d(1,Sb)) = Z(d(0,b)) = 1$ and $\mathrm{sg}(Sb) = 1$. Theorem 2.72 gives $q(n,a) = 0$.

(ii) Let $q(x,a,b)$ be the xor of the two sides. Simultaneous successors cancel in every subtraction, so $q(Sx,Sa,Sb) = q(x,a,b)$. If $x = 0$, both sides are 1. If $a = 0$, the sides are $Z(d(x,b))$ and $\mathrm{or}(Z(x),Z(d(x,b)))$; they agree by coverage on x, since at $x = 0$ both are 1 and at a successor $Z(x) = 0$. If $b = 0$, the sides are $Z(d(x,a))$ and $\mathrm{or}(Z(d(x,a)),Z(x))$; they agree by coverage on x and the two values of the Boolean term $Z(d(x,a))$. Theorem 2.72 gives $q(x,a,b) = 0$.

(iii) By coverage on u: at 0 both sides are $\mathrm{sg}(v)$, and at a successor both are 1, by $Su' + v = S(u'+v)$. □

2.4.6 An accumulator and the commutativity of the maximum

Define

$$\mu(n,a) := d(n,d(n,a)), \qquad \varphi(n,a,b) := \mu(n,a)+\mu\big(d(n,a),d(b,a)\big), \qquad m(a,b) := a+d(b,a).$$

These terms are compositions of d and addition; none is introduced by a recursion whose body sees the counter.

Lemma 2.74 (The accumulator)**.** (i) $\mu(Sn,a) = \mu(n,a) + Z(d(Sn,a))$.

(ii) $\varphi(0,a,b) = 0$ *and* $\varphi(Sn,a,b) = \varphi(n,a,b) + \mathrm{sg}\big(d(a,n) + d(b,n)\big)$.

(iii) $m(a,b) = m(P(a),P(b)) + \mathrm{sg}(a+b)$.

Proof. (i) Split $d(Sn,a)$ by coverage. If it is 0, then $d(n,a) = P(0) = 0$ by Theorem 2.71, so $\mu(Sn,a) = Sn$ and $\mu(n,a) = n$, and $Z(0) = 1$. If it is Sr, then $d(n,a) = r$ and $\mu(Sn,a) = d(Sn,Sr) = d(n,r) = \mu(n,a)$, and $Z(Sr) = 0$.

(ii) $\varphi(0,a,b) = \mu(0,a) + \mu(0,d(b,a)) = 0$, because $d(0,x) = 0$. Put $w := d(b,a)$ and split $d(Sn,a)$ again. If it is 0, then $d(n,a) = 0$, $\varphi(Sn,a,b) = Sn$, $\varphi(n,a,b) = n$ and $Z(d(0,w)) = 1$. If it is Sr, then $d(n,a) = r$, the first summand is unchanged by (i), and (i) for the second summand gives $\mu(Sr,w) = \mu(r,w) + Z(d(Sr,w))$. In both cases

$$\varphi(Sn,a,b) = \varphi(n,a,b) + Z\big(d(d(Sn,a),d(b,a))\big).$$

Theorem 2.73(ii) at $x = Sn$, then (i) of that lemma for each of the two Boolean terms, then (iii), turn the last summand into $\mathrm{sg}(d(a,n) + d(b,n))$.

(iii) By coverage on a and b. At $(0,0)$ both sides are 0. At $(0,Sb)$ and at $(Sa,0)$ both sides are the successor of $m(0,b) = b$ and of $m(a,0) = a$. At (Sa,Sb), $m(Sa,Sb) = Sa + d(b,a) = Sm(a,b)$ and $\mathrm{sg}(Sa+Sb) = 1$. □

Lemma 2.75 (Commutativity of the maximum)**.** $a + d(b,a) = b + d(a,b)$.

Proof. For fixed a, b put

$$F(n) := m(d(a,n),d(b,n)) + \varphi(n,a,b), \qquad F'(n) := m(d(b,n),d(a,n)) + \varphi(n,a,b).$$

By Theorem 2.74(iii) at the arguments $d(a,n)$, $d(b,n)$, by Theorem 2.74(ii) and by the laws of addition, $F(Sn) = F(n)$, and likewise $F'(Sn) = F'(n)$. Theorem 2.60(ii) gives $F(n) = F(0) = m(a,b)$ and $F'(n) = F'(0) = m(b,a)$. At $n = b$, $d(b,b) = 0$ and $m(x,0) = x = m(0,x)$, so $F(b)$ and $F'(b)$ both equal $d(a,b) + \varphi(b,a,b)$. Hence $m(a,b) = m(b,a)$. □

The familiar proof of this identity is an induction whose hypothesis is used at the predecessor. Here the two sides are joined by a term φ whose recurrence depends on the counter, although its definition is a composition; the recurrence is Theorem 2.74(ii).

2.4.7 Reflection of equality and the proofs of the theorems

Lemma 2.76 (Reflection of equality)**.** *Let* $\Delta(a,b) := \mathrm{sg}(d(a,b) + d(b,a))$. *Then* $\Delta(a,b) = 0 \vee \Delta(a,b) = 1$*, and* $\Delta(a,b) = 0$ *holds if and only if* $a = b$.

Proof. By coverage and separation, $\mathrm{sg}(u) = 0$ if and only if $u = 0$, and $u + v = 0$ if and only if $u = 0$ and $v = 0$, since a sum with a successor summand is a successor. If $\Delta(a,b) = 0$, then $d(a,b) = 0 = d(b,a)$, and Theorem 2.75 gives $a = a + d(b,a) = b + d(a,b) = b$. If $a = b$, both subtractions are $d(a,a) = 0$. □

The equality reflected is the equality of the logic on the whole numerical domain.

Proof of Theorem 2.58. By Theorem 2.57, every derivation of T_{UR} is a derivation of T_{eq} after each instance of UR is replaced by its derivation. Conversely let $\Gamma \vdash s(0) = r(0)$ and $\Gamma, s(z) = r(z) \vdash s(Sz) = r(Sz)$ in T_{UR}. Put $q(n) := \Delta(s(n), r(n))$. By Theorem 2.76 the two premises give $\Gamma \vdash q(0) = 0$ and $\Gamma, q(z) = 0 \vdash q(Sz) = 0$; Theorem 2.64 gives $q(t) = 0$; and Theorem 2.76 gives $s(t) = r(t)$. The context Γ is arbitrary, so this is the contextual induction rule of T_{eq}. The two theories have the same language and the same derivable sequents, and Theorem 1.78 applies to T_{UR}. In particular induction for quantifier-free formulas (Theorem 1.56) holds in T_{UR}. □

Proof of Theorem 2.59. UR is derivable in N by Theorem 2.57. Conversely it suffices to derive equation induction in U: given $\Gamma, E(z) \vdash_{\mathsf{U}} E(Sz)$, equation induction over the context $\Gamma, E(0)$ gives $E(Sz')$ for a fresh z', and the introduction of Q by cases (Theorem 2.4(i)) gives $\Gamma \vdash_{\mathsf{U}} Q_y E(y)$.

The native connectives of Section 1.4 obey their natural deduction rules in U, and the equality rules, the Leibniz rule and numerical cases hold in U: by Theorem 1.30, their derivations use the seven rules of N other than QI. With these, U derives the reading of every axiom and rule of T_{UR}: the computation equations, coverage by numerical cases, separation because the native falsity is the equation $0 = S0$, and UR itself. Hence the derivation of the induction rule in the proof of Theorem 2.58, written with the native connectives, is a derivation in U. □

Remark 2.77 (Sources and status). Sections 2.4.1 to 2.4.7 are `scalar_uniqueness`, except for the proof of Theorem 2.62 and for Theorem 2.65, which are from `orbit_closure` (Theorem 2.66). `scalar_uniqueness` attributes the idea of replacing induction by uniqueness of recursion to the literature (Okada and Scott study uniqueness of iteration in a typed setting) and claims the completion for the scalar language only: one state, no counter in the body, no pairing. Theorem 2.71 also yields the Mal'cev term $d(x + z, y)$, which the package notes. In Lean, Theorem 2.57 is proved; the converse direction of Theorems 2.58 and 2.59 is not formalized.

2.4.8 Other forms of the uniqueness rule

Three variations of UR occur in `orbit_closure` and `crossed_uniqueness`. They are related to UR as follows. The rule ORB concludes the closure of the equation of UR and takes no target; it is interderivable with UR by the rules for the closure alone (Theorem 2.79). The rule SU of Theorem 2.56 is the instance of UR at one body; it gives induction for zero equations (Theorem 2.65) and the laws of addition, and whether it gives induction for all equations is open. The rule UX has a premise for the seed in place of the seed $f(0)$ and a crossed conclusion; UR is its instance at the seed $f(0)$, and with transport it follows from UR. Taken as the only arithmetic rule on fully normalized terms, UX derives transport as well, so that it replaces I0, IS, TR and QI together (Theorem 2.86).

The closure of the orbit equation

For term templates f and h as in UR, and a variable y not free in h, the ordinary and the native form of the rule are

$$\frac{\Gamma \vdash f(Sz) = h(f(z))}{\Gamma \vdash \forall y\,(f(y) = I(y, f(0);\, u.\, h(u)))} \qquad \frac{\Gamma \vdash f(Sz) = h(f(z))}{\Gamma \vdash Q_y(f(y) = I(y, f(0);\, u.\, h(u)))}\ \textsc{Orb},$$

where z is not free in Γ or in h.

Definition 2.78 (The calculus U_o and the theory T_{Orb})**.** U_o is N with QI replaced by the native form of ORB; its eight rules are HYP, I0, IS, TR, QE, DE, DI and ORB. T_{Orb} is T_{UR} with UR replaced by the ordinary form of ORB.

Proposition 2.79 (Orbit uniqueness)**.** *T_{Orb} and T_{UR} derive the same sequents of ordinary formulas, and U_o and U are equivalent. Hence T_{Orb} derives the sequents of T_{eq}, and U_o is equivalent to N.*

Proof. Let $E(y)$ be the equation $f(y) = I(y, f(0); u.\, h(u))$. In the ordinary theories, an instance of UR is ORB followed by the elimination of $\forall$ at t; and UR at a fresh variable w as its target, followed by the introduction of $\forall$, is ORB.

In the native calculi, ORB is derived in U thus: UR at the target Sw, with w fresh, gives $E(Sw)$, and the introduction of Q by cases (Theorem 2.4(i)), whose assumption $E(0)$ is not needed, gives $Q_y E(y)$. Conversely UR is derived in U_o by QE at t: its second premise $E(0)$ is the instance $I(0, f(0); u.\, h(u)) = f(0)$ of I0, reversed. Both constructions hold over arbitrary contexts. The last sentence of the proposition is Theorems 2.58 and 2.59. □

The base $E(0)$ of the orbit equation is a theorem, so $Q_y E$ and the unconditional $\forall y\, E$ say the same; in a calculus with clause contexts (Section 1.9) ORB adds the clause of E with a trivial antecedent, and the term at which it is used is chosen when the clause is used. The rule removes a parameter of the certificate and not the principle.

Remark 2.80 (The body must not see the counter)*.* In ORB the variable y of the conclusion is not free in h. A checker that reads f and h off a proposed closure has to test this, also below nested binders. Without the test, $f(y) := y$ and the body Sy have the premise $Sz = Sz$ and the conclusion $Q_y\big(y = I(y, 0;\, u.\, Sy)\big)$, whose instance at 1 is $1 = 2$.

The successor body

Proposition 2.81 (Addition from successor uniqueness)**.** *T_{SU} derives that addition is associative and commutative with the neutral element 0. If $\Gamma \vdash k(Sz) = k(z)$ for a term template k and a fresh z, then T_{SU} derives $\Gamma \vdash n + k(n) = n + k(0)$.*

Proof. The laws of addition are those of Theorem 2.61, whose proofs compare terms with the update S. For the second statement let $f(n) := n + k(n)$. Then $f(0) = 0 + k(0) = k(0)$ and

$$f(Sn) = Sn + k(Sn) = Sn + k(n) = S(n + k(n)) = Sf(n),$$

so SU gives $f(n) = I(n, k(0);\, u.\, Su) = k(0) + n = n + k(0)$. □

If addition were cancellative in T_{SU}, the proposition would give $k(n) = k(0)$, which is Theorem 2.60(ii), uniqueness for the body u. Cancellation is proved in Theorem 2.71 from the predecessor, whose construction uses other bodies.

Remark 2.82 (What successor uniqueness is not known to give)*.* To pass from Theorem 2.65 to induction for all equations by the route of Section 2.4.7, a term Δ with $\Delta(a, b) = 0 \leftrightarrow a = b$ is needed. The construction of Sections 2.4.4 to 2.4.6 uses UR for the bodies u, $Z(u)$, $j(u)$, $g(u)$, $\chi(u)$, $\beta_m(u)$ and $P(u)$, among others, and cannot be repeated in T_{SU} as it stands. `orbit_closure` does not claim that T_{SU} is equivalent to T_{eq}. It also records that a model used in another package, `canonical_recursors`, to separate induction for zero equations from induction for all equations does not decide the question: in that model, for a nonstandard counter p, the term $e(a) := I(p, a;\, u.\, u)$ satisfies $e(0) = 0$ and $e(Sa) = Se(a)$ and is not the identity, whereas SU with Theorem 2.60(iii) gives $e(a) = a$. The model therefore refutes SU, and it is not a model of T_{SU} in which equation induction fails. The model is not reproduced here.

A seed premise and a crossed conclusion, on normal forms

Let $\mathrm{nf}(t)$ be the full normal form of a term t for the two computation equations, read from left to right, and let $\mathrm{nf}(A)$ and $\mathrm{nf}(\Gamma)$ be obtained by normalizing every term. The following facts are taken from the packages `full_normalization` and `computation_normal`, which belong to another chapter: nf is total; $\mathrm{nf}(\mathrm{nf}(t)[\theta]) = \mathrm{nf}(t[\theta])$ for every substitution θ; and N derives $t = \mathrm{nf}(t)$. A term, a formula or a context is *normal* if nf fixes it. For normal terms the selector satisfies $\mathrm{nf}(\mathrm{sel}(0, a, b)) = a$ and $\mathrm{nf}(\mathrm{sel}(Sc, a, b)) = b$, and $\mathrm{nf}(I(1, a;\, u.\, h)) = \mathrm{nf}(h(a))$.

Definition 2.83 (The calculus U_c)**.** The sequents of U_c consist of normal formulas, and every instance $B(t)$, $f(n)$ or $h(f(z))$ in a rule denotes the normal form of the substitution. Its six rules are Hyp, QE, DE, DI in their forms of Section 1.3, reflexivity,

$$\frac{}{\Gamma \vdash t = t}\ \textsc{Ref},$$

and *seeded crossed uniqueness*,

$$\frac{\Gamma \vdash f(0) = a \qquad \Gamma \vdash f(Sz) = h(f(z))}{\Gamma \vdash I(n, a;\, u.\, h(u)) = f(n)}\ \mathrm{UX},$$

where $f(z)$ and $h(u)$ are term templates and z is not free in Γ, a, n or h.

There is no computation rule, no transport rule and no rule for progress: computation is in the syntax, and normalization is part of the specification of the calculus and of any checker for it. The introduction DI is by cases at every matrix. The body h receives the state and no counter, and the second premise has no induction hypothesis. This is the one calculus of the chapter without the rules I0 and IS.

Lemma 2.84 (Structural rules on normal forms)**.** *Weakening and cut are admissible in* U_c*, and if* $\Gamma \vdash_{\mathsf{U}_\mathrm{c}} A$ *then* $\mathrm{nf}(\Gamma[\theta]) \vdash_{\mathsf{U}_\mathrm{c}} \mathrm{nf}(A[\theta])$.

Proof. By induction on derivations, with eigenvariables renamed. No rule has a mode or a test on a term, and $\mathrm{nf}(\mathrm{nf}(t)[\theta]) = \mathrm{nf}(t[\theta])$ identifies the substituted premises and conclusions of each rule with an instance of the same rule. □

Lemma 2.85 (Equality from seeded crossed uniqueness)**.** U_c *derives, over an arbitrary context:*

(i) *numerical cases and the introduction of* Q *by cases;*

(ii) *symmetry: from* $s = t$ *to* $t = s$*;*

(iii) *congruence: from* $s = t$ *to* $H(s) = H(t)$*, for every term template* H*;*

(iv) *joining: from* $s = t$ *and* $s = r$ *to* $t = r$*;*

(v) *transitivity, and the rule* Tr*: from* $s = t$ *and* $H(s) = r$ *to* $r = H(t)$*;*

(vi) *the rule UR.*

None of these derivations uses an induction, a predecessor or a test of equality.

Proof. (i) Theorem 2.4(i); its derivations use Hyp, QE, DE and DI.

(ii) Apply UX with the constant template $f(z) := s$, the seed $a := t$, the body $h(u) := u$ and the counter $n := 0$. The first premise is $s = t$, the second is $s = s$, and the conclusion $I(0, t;\, u.\, u) = s$ is, in normal form, $t = s$.

(iii) Apply UX with $f(z) := I(z, s; u.\, H(u))$, $a := t$, $h := H$ and $n := 1$. The first premise is $s = t$, because $f(0)$ has the normal form s. The two sides of the second premise, $I(Sz, s; u.\, H(u))$ and $H(I(z, s; u.\, H(u)))$, have the same normal form, so it is an instance of Ref. The conclusion is $H(t) = H(s)$ in normal form, and (ii) reverses it. The template may use its variable several times or not at all, also below binders.

(iv) Apply UX with $f(z) := \mathrm{sel}(z, s, r)$, $a := t$, $h(u) := u$ and $n := 1$. The first premise is $s = t$. The second premise is $r = \mathrm{sel}(z, s, r)$, since $f(Sz)$ has the normal form r; it follows by numerical cases on z, from $r = s$ at 0, which is (ii) applied to $s = r$, and from $r = r$ at a successor. The conclusion is $t = r$ in normal form.

(v) From $s = t$ and $t = r$: (ii) gives $t = s$, and (iv) gives $s = r$. From $s = t$ and $H(s) = r$: (iii) gives $H(s) = H(t)$, and (iv) gives $r = H(t)$.

(vi) UX with the seed $a := f(0)$ and the first premise by Ref, followed by (ii). □

The counter 0 makes UX a replacement at the root, the counter 1 makes it an application of a template, and the selector joins two equations; each conclusion is read after normalization, which is why the calculus is stated on normal forms.

Theorem 2.86 (Seeded crossed uniqueness)**.** *For every context* Γ *and every native formula* A*,*

$$\Gamma \vdash A \text{ in } \mathsf{N} \quad \Longleftrightarrow \quad \mathrm{nf}(\Gamma) \vdash_{\mathsf{U_c}} \mathrm{nf}(A).$$

Hence the bridge of Theorem 1.78 holds for $\mathsf{U_c}$*, with the translations composed with* nf*.*

Proof. From right to left. Every rule of $\mathsf{U_c}$ is derivable in N on normal formulas. Ref is reflexivity. For UX apply equation induction to $I(y, a; u.\, h(u)) = f(y)$: the base follows from I0 and the first premise, and in the step IS, congruence under h applied to the induction hypothesis, and the second premise give $I(Sz, a; u.\, h(u)) = h(I(z, a; u.\, h(u))) = h(f(z)) = f(Sz)$. The normalization of an instance is justified by $t = \mathrm{nf}(t)$ and the Leibniz rule. The same two facts make A and $\mathrm{nf}(A)$ interderivable in N, so $\mathrm{nf}(\Gamma) \vdash \mathrm{nf}(A)$ gives $\Gamma \vdash A$.

From left to right. First, if $\Gamma \vdash_{\mathsf{U}} A$ in the calculus U of Theorem 2.55, then $\mathrm{nf}(\Gamma) \vdash_{\mathsf{U_c}} \mathrm{nf}(A)$, by induction on the derivation. Hyp is clear. The two sides of an instance of I0 or IS have the same normal form, so the instance becomes Ref. Tr becomes Theorem 2.85(v) with the template $\mathrm{nf}(h)$, because $\mathrm{nf}(h(s)) = \mathrm{nf}(\mathrm{nf}(h)(\mathrm{nf}(s)))$. QE, DE and DI become the same rules, by the same identity. UR becomes Theorem 2.85(vi). Second, $\Gamma \vdash A$ in N implies $\Gamma \vdash_{\mathsf{U}} A$ by Theorem 2.59. □

Remark 2.87 (Sources, overlaps and status). Theorems 2.79 to 2.82 are from `orbit_closure`, which states Orb for the ordinary theory and describes the native form with its checker. Its translator maps a derivation with UR to one with Orb by the second construction in the proof of Theorem 2.79. Theorems 2.83, 2.85 and 2.86 are from `crossed_uniqueness`. That package also derives transport in a formula template inside $\mathsf{U_c}$ by induction on the template, and restates the native connectives through $D_,$ with an unused coordinate; here both follow from Theorem 2.86 and Section 1.4. Both packages take the passage from UR to equation induction from `scalar_uniqueness`, as the proofs above do, and neither claims it as new. Neither package is formalized in Lean.

2.5 The progressive reading of the closure

This section draws on `uniform_progress`, `target_closure` and the note on mixed closure in `scope_and_closure`.

The single introduction of Section 2.2.4 tests the witness. A single introduction whose mode depends on the root of the matrix alone is possible if the meaning of $Q_y E$ at a literal

equation E is changed from $E(0) \to \forall y\, E(y)$ to progressivity, $\forall y\, (E(y) \to E(Sy))$. The calculi of this section have seven rules, keep all native formulas, and are stable under substitution or under renaming; they are equivalent to N through two translations and not on the same formulas.

2.5.1 The calculus with orbit elimination

Extend the selector of modes of Section 2.2.2 to terms: $\eta_B(a) := a$ if the matrix B is a literal equation and $\eta_B(a) := 0$ if B is compound. The *orbit term* is

$$\omega(a, n) := I(a, n;\, u.\, Su),$$

with the counter a and the seed n.

Definition 2.88 (The calculus R)**.** R has the seven rules Hyp, I0, IS, Tr_2, DE and

$$\frac{\Gamma \vdash Q_y B(y) \qquad \Gamma \vdash B(\eta_B(a))}{\Gamma \vdash B(\omega(\eta_B(a), n))}\ \mathrm{QE}_\omega$$

$$\frac{\Gamma,\ B(t, \eta_B(z)) \vdash B(t, Sz)}{\Gamma \vdash D_{x,y} B(x, y)}\ \mathrm{DI}_\eta,$$

where a and n are arbitrary terms, and z is not free in Γ, in t or in $D_{x,y}B$. The designated contradiction is $0 = S0$.

At a compound matrix QE_ω passes from $B(0)$ to $B(\omega(0, n))$, and $\omega(0, n) = n$ is an instance of I0. At a literal equation E it passes from $E(a)$ to $E(\omega(a, n))$, the instance n steps further. Transport in a compound formula is not a rule, and there is no rule of numerical cases.

Definition 2.89 (The progressive reading)**.** The ordinary formula F^{p} is defined by

$$\begin{aligned}
(s = t)^{\mathrm{p}} &:= (s = t), \\
(Q_y E)^{\mathrm{p}} &:= \forall y\, (E(y) \to E(Sy)) && (E \text{ a literal equation}), \\
(Q_y B)^{\mathrm{p}} &:= B^{\mathrm{p}}(0) \to \forall y\, B^{\mathrm{p}}(y) && (B \text{ compound}), \\
(D_{x,y} B)^{\mathrm{p}} &:= \exists x\, (Q_y B)^{\mathrm{p}}.
\end{aligned}$$

The reading commutes with substitution, because substitution preserves the root of a matrix. The elimination and the reading have to change together:

Proposition 2.90 (The two readings differ)**.** *Let $E(y)$ be the equation $N(y) = 1$ for the parity term N of Section 1.1.*

(i) *In the standard model $E(0) \to \forall y\, E(y)$ is true and $\forall y\, (E(y) \to E(Sy))$ is false.*

(ii) *The rule QE_ω does not preserve truth for the ordinary reading of Section 1.2.*

(iii) N *derives $Q_y E(y)$, and* R *does not.*

Proof. $N(0) = 0$, $N(1) = 1$ and $N(2) = 0$. (i) The premise $E(0)$ is false, and progress fails at $y = 1$. (ii) With $a := 1$ and $n := 1$ the two premises $Q_y E$ and $E(1)$ are true in the ordinary reading and the conclusion $E(\omega(1, 1))$, which is $E(2)$, is false. (iii) In N the closure follows by the introduction by cases from $N(0) = 1 \vdash N(Sz) = 1$, which holds because $N(0) = 0$ and $0 = 1$ derives every equation. In R it is not derivable by Theorem 2.95 below and (i). □

2.5.2 Derived rules of the orbit calculus

Lemma 2.91 (Structural rules)**.** *Weakening, cut and substitution of terms are admissible in* R*, and substitution maps every rule instance to an instance of the same rule.*

Proof. Theorem 2.3: no rule of R tests a term, and $\eta_B(a)[\theta] = \eta_{B[\theta]}(a[\theta])$. □

Lemma 2.92 (Equality, closure introduction, equation induction)**.** R *derives:*

(i) *reflexivity, symmetry, transitivity, congruence in a term template, and* $\textsc{Tr}$*;*

(ii) *the introduction* QI_η *of Theorem 2.9: from* $\Gamma, B(\eta_B(z)) \vdash B(Sz)$ *with* z *fresh to* $\Gamma \vdash Q_y B(y)$*;*

(iii) *equation induction: from* $\Gamma \vdash E(0)$ *and* $\Gamma, E(z) \vdash E(Sz)$*, for a literal equation* E *and* z *fresh, to* $\Gamma \vdash E(t)$*.*

The derivations of (i) and (ii) use $\textsc{Hyp}$*, I0,* $\textsc{Tr}_2$*, DE and* DI_η *only.*

Proof. (i) Reflexivity and symmetry are derived from I0 and $\textsc{Tr}_2$ in the proof of Theorem 2.21. Transitivity and congruence are $\textsc{Tr}_2$ applied to $r = s$ in the template $r = v$ and to $f(s) = f(s)$ in the template $f(s) = f(v)$. $\textsc{Tr}$ follows as in that proof.

(ii) Apply DI_η with the matrix $B(y)$, a coordinate x not free in B and the witness 0; then DE, whose opened hypothesis is $Q_y B(y)$, and $\textsc{Hyp}$.

(iii) By (ii) the step gives $Q_y E(y)$. QE_ω with $a := 0$ and $n := t$ gives $E(\omega(0,t))$ from the base, and $\omega(0,t) = t$ is an instance of I0; transport in the equation template E gives $E(t)$. □

The order of the arguments of ω is what makes (iii) work: $\omega(0,t) = t$ is a computation, whereas $\omega(t,0) = t$ is a theorem that needs the induction under construction.

Lemma 2.93 (Orbit identities)**.** *In every calculus or theory with the equality rules, the computation equations and equation induction, in particular in* R*,* N *and* T_{eq}*,*

$$\omega(0,n) = n, \qquad \omega(Sa,n) = S\,\omega(a,n), \qquad \omega(a,0) = a,$$
$$\omega(a,Sn) = S\,\omega(a,n), \qquad \omega(a,n) = \omega(n,a), \qquad \omega(a,1) = Sa.$$

Proof. The first two are I0 and IS. The third is by induction on a: $\omega(0,0) = 0$, and $\omega(Sa,0) = S\,\omega(a,0) = Sa$. The fourth is by induction on a with the parameter n: both sides are Sn at 0, and $\omega(Sa,Sn) = S\,\omega(a,Sn) = SS\,\omega(a,n) = S\,\omega(Sa,n)$. The fifth is by induction on a: the base is $n = \omega(n,0)$, and $\omega(Sa,n) = S\,\omega(a,n) = S\,\omega(n,a) = \omega(n,Sa)$. The last is $\omega(a,1) = S\,\omega(a,0) = Sa$. □

Lemma 2.94 (The two interfaces of a closure, and transport)**.** *In* R*, over an arbitrary context:*

(i) *one step: from* $Q_y E(y)$ *and* $E(a)$ *to* $E(Sa)$*, for a literal equation* E*;*

(ii) *transport: from* $s = t$ *and* $F(s)$ *to* $F(t)$*, for every native formula template* F*;*

(iii) *instance from the base: from* $Q_y B(y)$ *and* $B(0)$ *to* $B(t)$*, for every matrix* B*;*

(iv) *homogeneous congruence: if* B *and* C *are interderivable over* Γ *and are both literal equations or both compound, then* $Q_y B$ *and* $Q_y C$ *are interderivable over* Γ*, and so are* $D_{x,y}B$ *and* $D_{x,y}C$*, where the bound variables are not free in* Γ*.*

Proof. (i) QE_ω with $n := 1$ gives $E(\omega(a,1))$, and $\omega(a,1) = Sa$ by Theorem 2.93.

(ii) and (iii) are proved together by induction on the number of binders of the formula, a measure that substitution does not change; (iii) for B uses (ii) for B. For an equation, (ii) is $\textsc{Tr}_2$. For (iii): at a literal equation, QE_ω with $a := 0$ and $n := t$ gives $B(\omega(0,t))$; at a compound matrix it gives the same from $B(0)$; and (ii) for B along $\omega(0,t) = t$ gives $B(t)$.

(ii) for $Q_y B(v,y)$. Derive the target $Q_y B(t,y)$ by QI_η: under the assumption $B(t,\eta_B(z))$, transport in B along $t = s$ gives $B(s,\eta_B(z))$; then (i), if B is a literal equation, or (iii) at Sz, if B is compound, gives $B(s,Sz)$ from the given closure $Q_y B(s,y)$; and transport in B along $s = t$ gives $B(t,Sz)$.

(ii) for $D_{x,y}B(v,x,y)$. Apply DE to the given formula with a fresh b, and derive the target by DI_η with the witness b: the premise $B(t,b,\eta_B(z)) \vdash B(t,b,Sz)$ follows from the opened closure $Q_y B(s,b,y)$ exactly as in the previous case. The conclusion does not contain b.

Every use of transport or of (iii) inside these two cases is for the matrix B, which has fewer binders.

(iv) Repeat the two cases of (ii) with the given derivations between B and C, instantiated by substitution (Theorem 2.91), in place of transport. Because B and C have the same kind of root, $\eta_B = \eta_C$, so the assumptions of the two introductions match. □

The restriction in (iv) is necessary. An equation E and a compound formula C can be interderivable while $Q_y E$, which is progressive, and $Q_y C$, which is conditional, are not; the padding of Theorem 2.96 gives such pairs. Transport is unaffected, because it preserves the root of the matrix.

2.5.3 Soundness for the progressive reading

Theorem 2.95 (Soundness)**.** *If* $\Gamma \vdash_{\mathsf{R}} F$*, then* $\Gamma^{\mathrm{p}} \vdash_{\mathsf{T}_{\mathrm{eq}}} F^{\mathrm{p}}$*.*

Proof. By induction on the derivation. Hyp, the computation equations, $\textsc{Tr}_2$ and DE have their ordinary meanings; for DE the opened hypothesis reads $(Q_y B(b,y))^{\mathrm{p}}$, the matrix of the existential quantifier.

DI_η at a literal equation: the premise gives $\forall z\,(E(t,z) \to E(t,Sz))$, and t is a witness. At a compound matrix: the premise gives $B^{\mathrm{p}}(t,Sz)$ under $B^{\mathrm{p}}(t,0)$, so coverage gives $B^{\mathrm{p}}(t,0) \to \forall y\, B^{\mathrm{p}}(t,y)$.

QE_ω at a compound matrix: the first premise reads $B^{\mathrm{p}}(0) \to \forall y\, B^{\mathrm{p}}(y)$, the second gives $B^{\mathrm{p}}(0)$, and $\omega(0,n) = n$. At a literal equation: assume $\forall y\,(E(y) \to E(Sy))$ and $E(a)$. Induction on k for the literal equation $E(\omega(k,a))$, with these two assumptions in the context, gives $E(\omega(k,a))$ for every k: the base is $E(a)$ by $\omega(0,a) = a$, and the step is the assumed progress at $\omega(k,a)$ with $\omega(Sk,a) = S\,\omega(k,a)$. At $k := n$, the equation $\omega(n,a) = \omega(a,n)$ of Theorem 2.93 gives $E(\omega(a,n))$. □

The use of the commutativity of ω in this proof is harmless: it is a theorem of T_{eq}, whose induction is given. It is separate from the recovery of induction inside R, which uses $\omega(0,t) = t$ only.

2.5.4 The embedding of the core

For a literal equation E and fresh variables r, v let

$$\mathrm{pad}(E) := D_{r,v}\,\mathrm{gate}(v,E), \qquad \text{that is,} \qquad D_{r,v}(\mathrm{sel}(v,0,s) = \mathrm{sel}(v,0,t)) \quad \text{for } E \equiv (s = t),$$

and for a native formula F let $F^\sharp := \mathrm{pad}(F)$ if F is a literal equation and $F^\sharp := F$ if F is compound. The formula $F^\sharp$ is always compound. The translation $A^\uparrow$ of a native formula is

$$(s = t)^\uparrow := (s = t), \qquad (Q_y B)^\uparrow := Q_y (B^\uparrow)^\sharp, \qquad (D_{x,y}B)^\uparrow := D_{x,y}(B^\uparrow)^\sharp.$$

It fixes terms and equations, preserves the kind of the root, commutes with substitution, and puts a compound matrix below every binder.

Lemma 2.96 (Padding)**.** *For a literal equation E, the formulas* $\mathrm{pad}(E)$ *and E are interderivable in* R *over every context. Hence $F^\sharp$ and F are interderivable for every F.*

Proof. Let E be $s = t$. The computation equations give $\mathrm{gate}(0, E)$, both sides being 0, and make $\mathrm{gate}(Sz, E)$ and E interderivable. From E: the matrix $\mathrm{gate}(v, E)$ is a literal equation, so DI_η asks for $\mathrm{gate}(Sz, E)$ under the assumption $\mathrm{gate}(z, E)$, which is not needed. From $\mathrm{pad}(E)$: DE opens $Q_v\,\mathrm{gate}(v, E)$, and QE_ω with $a := 0$, $n := 1$ and the second premise $\mathrm{gate}(0, E)$ gives $\mathrm{gate}(\omega(0, 1), E)$. I0 and $\textsc{Tr}_2$ give $\mathrm{gate}(1, E)$, hence E. □

Theorem 2.97 (Embedding)**.** *If $\Gamma \vdash A$ in* N*, then $\Gamma^\uparrow \vdash_{\mathsf{R}} A^\uparrow$.*

Proof. By induction on the derivation. Hyp, I0, IS translate to themselves, and Tr is derivable in R (Theorem 2.92(i)).

QE. The premises translate to $Q_y(B^\uparrow)^\sharp$ and $B^\uparrow(0)$. Theorem 2.96 gives $(B^\uparrow)^\sharp(0)$, Theorem 2.94(iii) gives $(B^\uparrow)^\sharp(t)$, and Theorem 2.96 gives $B^\uparrow(t)$, which is $B(t)^\uparrow$.

DE. The first premise translates to $D_{x,y}(B^\uparrow)^\sharp$, and the hypothesis opened by DE in R is $Q_y(B^\uparrow)^\sharp(b, y)$, the translation of $Q_y B(b, y)$.

DI. The translated premise is $\Gamma^\uparrow, B^\uparrow(t, 0) \vdash B^\uparrow(t, Sz)$. The target has the compound matrix $(B^\uparrow)^\sharp$, so DI_η asks for $(B^\uparrow)^\sharp(t, Sz)$ under $(B^\uparrow)^\sharp(t, 0)$; remove the padding of the assumption, cut with the premise, and pad the conclusion.

QI. The translated premise is $\Gamma^\uparrow, E(z) \vdash E(Sz)$, and the target is $Q_y\,\mathrm{pad}(E(y))$, with a compound matrix. By Theorem 2.92(ii) it suffices to derive $\mathrm{pad}(E(Sz'))$ from $\mathrm{pad}(E(0))$. The assumption gives $E(0)$; equation induction in R (Theorem 2.92(iii)) with the translated premise as its step gives $E(Sz')$; and Theorem 2.96 pads it. □

2.5.5 The reading inside the core and the two round trips

The translation $F^\downarrow$ expresses the progressive reading with the native connectives $\Rightarrow_*$, $\forall_*$, $\exists_*$ of Section 1.4. It fixes equations and is defined by

$$(Q_y E)^\downarrow := \forall_* y\,(E(y) \Rightarrow_* E(Sy)), \qquad (Q_y B)^\downarrow := Q_y B^\downarrow,$$
$$(D_{x,y} E)^\downarrow := \exists_* x\,\forall_* y\,(E(x, y) \Rightarrow_* E(x, Sy)), \qquad (D_{x,y} B)^\downarrow := D_{x,y} B^\downarrow,$$

where E is a literal equation and B is compound. It maps compound formulas to compound formulas and commutes with substitution.

Theorem 2.98 (The reading is derivation-preserving)**.** *If $\Gamma \vdash_{\mathsf{R}} F$, then $\Gamma^\downarrow \vdash F^\downarrow$ in* N*.*

Proof. By Theorem 1.78 the ordinary reading of a native connective is equivalent in T_{eq} to the corresponding connective of the readings. By induction on F, therefore, $(F^\downarrow)^\circ$ is equivalent in T_{eq} to F^{p}. Theorem 2.95 gives $\Gamma^{\mathrm{p}} \vdash_{\mathsf{T}_{\mathrm{eq}}} F^{\mathrm{p}}$, hence $(\Gamma^\downarrow)^\circ \vdash_{\mathsf{T}_{\mathrm{eq}}} (F^\downarrow)^\circ$, and the faithfulness of the ordinary reading (Theorem 1.78) gives $\Gamma^\downarrow \vdash F^\downarrow$. □

A formula of the form $A^\uparrow$ is an *image*. The rules of the native connectives are available in R for images, over arbitrary contexts of R.

Lemma 2.99 (Logic on images)**.** *Let Δ be a context of* R*, let A, A_i, B, C be native formulas, and let F be a formula of* R*.*

(i) *If $A_1, \ldots, A_k \vdash C$ in* N *and $\Delta \vdash_{\mathsf{R}} A_i^\uparrow$ for every i, then $\Delta \vdash_{\mathsf{R}} C^\uparrow$.*

(ii) *If $\Delta, C(0)^\uparrow \vdash_{\mathsf{R}} C(Sz)^\uparrow$ with z fresh, then $\Delta \vdash_{\mathsf{R}} (Q_y C(y))^\uparrow$.*

(iii) *If* $\Delta, A^{\uparrow} \vdash_{\mathsf{R}} B^{\uparrow}$*, then* $\Delta \vdash_{\mathsf{R}} (A \Rightarrow_* B)^{\uparrow}$.

(iv) *If* $\Delta \vdash_{\mathsf{R}} C(z)^{\uparrow}$ *with* z *not free in* Δ*, then* $\Delta \vdash_{\mathsf{R}} (\forall_* y\, C(y))^{\uparrow}$.

(v) *If* $\Delta \vdash_{\mathsf{R}} (\exists_* x\, C(x))^{\uparrow}$ *and* $\Delta, C(b)^{\uparrow} \vdash_{\mathsf{R}} F$ *with* b *not free in* Δ*,* F *and* C*, then* $\Delta \vdash_{\mathsf{R}} F$.

Proof. (i) Theorem 2.97, weakening and cut. Every rule of the native connectives without a discharged assumption is of this form.

(ii) The target is $Q_y(C^{\uparrow})^{\sharp}(y)$, with a compound matrix. By Theorem 2.92(ii) it suffices to derive $(C^{\uparrow})^{\sharp}(Sz)$ from $(C^{\uparrow})^{\sharp}(0)$, which follows from the hypothesis and Theorem 2.96.

(iii) $A \Rightarrow_* B$ is $Q_y \operatorname{sel}(y, A, B)$. By (ii) it suffices to derive $\operatorname{sel}(Sz, A, B)^{\uparrow}$ from $\operatorname{sel}(0, A, B)^{\uparrow}$. By (i) and $\operatorname{sel}(0, A, B) \vdash A$, the assumption gives $A^{\uparrow}$; the hypothesis gives $B^{\uparrow}$; and (i) with $B \vdash \operatorname{sel}(Sz, A, B)$ gives the conclusion.

(iv) $\forall_* y\, C$ is $C(0) \wedge_* Q_y C$. Substitution in R gives $C(0)^{\uparrow}$ and $C(Sz)^{\uparrow}$, (ii) gives $(Q_y C)^{\uparrow}$, and (i) with the introduction of $\wedge_*$ gives the conclusion.

(v) $\exists_* x\, C$ is $D_{x,y} \operatorname{gate}(y, C)$, and its image is $D_{x,y}(\operatorname{gate}(y, C)^{\uparrow})^{\sharp}$. DE in R with the fresh variable b opens the image of $Q_y \operatorname{gate}(y, C(b))$. By (i) and $Q_y \operatorname{gate}(y, C(b)) \vdash C(b)$, this gives $C(b)^{\uparrow}$, and cut with the second hypothesis gives F. The formula F is carried through DE and is not analyzed. □

The lemma does not say that the selector and the connectives of N can be applied to arbitrary formulas of R; that would ignore the dependence of Q on the root of its matrix.

Lemma 2.100 (Normalization of atomic closures)**.** *For a literal equation* E*, over every context of* R*,*

$$Q_y E(y) \dashv\vdash_{\mathsf{R}} (Q_y E(y))^{\downarrow\uparrow}, \qquad D_{x,y} E(x, y) \dashv\vdash_{\mathsf{R}} (D_{x,y} E(x, y))^{\downarrow\uparrow}.$$

Proof. Equations are their own images. From $Q_y E(y)$ and $E(z)$, one step (Theorem 2.94(i)) gives $E(Sz)$; Theorem 2.99(iii) and (iv) give the image of $\forall_* y\, (E(y) \Rightarrow_* E(Sy))$. Conversely, from that image and $E(z)$, Theorem 2.99(i) with the eliminations of $\forall_*$ and $\Rightarrow_*$ gives $E(Sz)$, and QI_η at the literal equation E gives $Q_y E(y)$.

For $D_,$: apply DE at a fresh b, convert the opened $Q_y E(b, y)$ by the first part, and apply Theorem 2.99(i) with the introduction of $\exists_*$. Conversely apply Theorem 2.99(v) with $F := D_{x,y} E(x, y)$: under the image of $\forall_* y\, (E(b, y) \Rightarrow_* E(b, Sy))$ and the assumption $E(b, z)$ the eliminations give $E(b, Sz)$, which is the premise of DI_η with the witness b. □

Theorem 2.101 (Round trips)**.** *For every native formula* A *and every formula* F *of* R*,*

$$A \dashv\vdash A^{\uparrow\downarrow} \text{ in } \mathsf{N}, \qquad F \dashv\vdash_{\mathsf{R}} F^{\downarrow\uparrow},$$

over arbitrary contexts.

Proof. In N. First, $\operatorname{pad}(E)^{\downarrow}$ is interderivable with E: it is $\exists_* r\, \forall_* v\, (\operatorname{gate}(v, E) \Rightarrow_* \operatorname{gate}(Sv, E))$ with r unused. From E every $\operatorname{gate}(Sv, E)$ follows. Conversely $\operatorname{gate}(0, E)$ is derivable, and progress at 0 gives $\operatorname{gate}(1, E)$, hence E. Now induct on A. Equations are fixed. The matrix $(B^{\uparrow})^{\sharp}$ is compound, so $(Q_y B)^{\uparrow\downarrow}$ is $Q_y((B^{\uparrow})^{\sharp})^{\downarrow}$. If B is an equation, the matrix is $\operatorname{pad}(B)^{\downarrow}$, interderivable with B. If B is compound, the matrix is $B^{\uparrow\downarrow}$, interderivable with B by the induction hypothesis. The congruence of Q in N (Section 1.4) gives the claim, and $D_,$ is the same.

In R. Induct on F. Equations are fixed. If the matrix of F is a literal equation, the claim is Theorem 2.100. Let the matrix B be compound. The induction hypothesis makes B and $B^{\downarrow\uparrow}$ interderivable, and both are compound. Theorem 2.94(iv) makes $Q_y B$ and $Q_y B^{\downarrow\uparrow}$ interderivable, and since $B^{\downarrow\uparrow}$ is compound, $Q_y B^{\downarrow\uparrow}$ is $(Q_y B)^{\downarrow\uparrow}$ with no padding inserted. $D_,$ is the same. No congruence across different kinds of root is used, and the normalization is proved for every formula before the connectives are extended to all of R. □

Theorem 2.102 (The orbit calculus and the core)**.** (i) $\Gamma \vdash A$ *in* N *if and only if* $\Gamma^{\uparrow} \vdash_{\mathsf{R}} A^{\uparrow}$.

(ii) $\Gamma \vdash_{\mathsf{R}} F$ *if and only if* $\Gamma^{\downarrow} \vdash F^{\downarrow}$ *in* N.

(iii) R *derives* $0 = S0$ *if and only if* N *does.*

(iv) *With the progressive reading,* R *is bi-interpretable with* T_{eq} *and with* iPRA*, with whole domains, actual equality, identity numerical composites and both formula round trips.*

Proof. (i) One direction is Theorem 2.97. For the other, apply Theorem 2.98 to $\Gamma^{\uparrow} \vdash_{\mathsf{R}} A^{\uparrow}$ and remove $\uparrow\downarrow$ by Theorem 2.101 and cut. (ii) is symmetric, with Theorem 2.98, Theorem 2.97 and the second round trip. (iii) is (i) or (ii) for the equation $0 = S0$, which both translations fix. (iv) The translations $\uparrow$ and $\downarrow$ fix every term, so composing them with the interpretations of Theorem 1.78 adds no change of domain; the formula round trips compose. □

Corollary 2.103 (Connectives for all formulas of the orbit calculus)**.** *For formulas* F*,* G *of* R *put* $F \Rightarrow_{\mathsf{R}} G := (F^{\downarrow} \Rightarrow_{*} G^{\downarrow})^{\uparrow}$*, and likewise for* $\wedge$*,* $\vee$*,* $\exists$ *and* $\forall$*; falsity is* $0 = S0$*. These operations obey the rules of intuitionistic natural deduction in* R*, and* R *derives coverage and equation induction.*

Proof. By Theorem 2.101, $(F \Rightarrow_{\mathsf{R}} G)^{\downarrow}$ is interderivable in N with $F^{\downarrow} \Rightarrow_{*} G^{\downarrow}$. By Theorem 2.102(ii) each rule for $\Rightarrow_{\mathsf{R}}$ is therefore the rule for $\Rightarrow_{*}$ in N, applied to the readings. The other connectives are the same. □

Remark 2.104 (Sources and Lean). The calculus and Theorem 2.102 are from `uniform_progress`. The package compares R with an earlier progressive presentation that reads every closure progressively, has a recursor whose body sees the counter, two eliminations and a rule of Boolean cases; here the iterator is unchanged, compound closures keep the conditional reading, and one elimination serves both. In Lean (modules `NativeProgressive` and following) the calculus is `RPrf`, the translation $\downarrow$ is `NFm.rd` and $\uparrow$ is `NFm.emb`. Theorem 2.98 is `RPrf.toNPrf`, with `rprf_sound_pra` and `rprf_consistent`; Theorem 2.91 is `RPrf.subst`; Theorem 2.92(iii) is `RPrf.eqInd`; Theorem 2.94(ii) is `RPrf.leibniz`; Theorem 2.96 is `RPrf.padI` and `RPrf.padE`; Theorem 2.97 is `NPrf.toR`; Theorem 2.102(i) and (ii) are `rprf_emb_iff` and `rprf_iff_rd`; and the two round trips are `NEquiv.rd_emb` and `REquivIn.emb_rd`. In the Lean proof of Theorem 2.98 the only rule that needs an argument is QE_{ω} at an equation, which is an equation induction in the kernel (`NPrf.progE`).

2.5.6 Transport in a formula template

Definition 2.105 (The calculus R_F)**.** R_F is R with $\textsc{Tr}_2$ replaced by $\textsc{Tr}_F$; its seven rules are $\textsc{Hyp}$, I0, IS, $\textsc{Tr}_F$, QE_{ω}, DE and DI_{η}.

Proposition 2.106 (Mixed closure)**.** R_F *and* R *derive the same sequents. Hence Theorems 2.95 and 2.102 hold for* R_F*.*

Proof. $\textsc{Tr}_2$ is the instance of $\textsc{Tr}_F$ at an equation template, and $\textsc{Tr}_F$ is derivable in R by Theorem 2.94(ii). □

Remark 2.107 (The account of `scope_and_closure`). R_F is the "mixed closure" calculus of `scope_and_closure`, written there with the orbit term $I(n, t; u.\, Su)$ and the selector d_B for η_B. The package proves the bi-interpretation with T_{eq} independently of `uniform_progress`, and its proof differs in three places. The gate is extended through binders, $G_v(Q_y B) := Q_y G_v B$ and $G_v(D_{x,y} B) := D_{x,y} G_v B$, with $\vdash G_0 B$ and $G_{Sv} B \dashv\vdash B$ proved by induction on B from homogeneous congruence. Every matrix is padded, $P(B) := D_{r,v} G_v B$, compound or not. And the second formula round trip is proved through T_{eq}: with ρ the progressive reading and

$K(\varphi)$ the padded native translation of an ordinary formula φ, the package shows $\rho(K\varphi) \leftrightarrow \varphi$ in T_{eq} and $B \dashv\vdash K(B^{\rho})$ in its calculus, the latter by the cases of Theorems 2.100 and 2.101. For QE_{ω} it needs only $\omega(a,1) = Sa$ inside the calculus, as here. In Lean the calculus is `MCPrf`, with `mcprf_iff_rprf`, `mcprf_iff_rd`, `mcprf_emb_iff` and `mcprf_consistent`. The package measures no checker for it and states that the single introduction moves a cost into the elimination, which builds a term and takes a second numerical parameter.

2.5.7 Elimination at a target

The orbit elimination takes two terms, a source and a seed. It can be replaced by an elimination that takes the target alone. For a matrix B and a term t put

$$\delta_B(t) := \begin{cases} r, & B \text{ is a literal equation and } t \equiv Sr, \\ 0, & \text{otherwise.} \end{cases}$$

The test is on the syntax of t; terms are not normalized before it, and $J(Sr) = I(0, Sr;\, u.\,0)$ is not a literal successor.

Definition 2.108 (The calculus T)**.** T has the seven rules Hyp, I0, IS, Tr_2, DE, DI_{η} and

$$\frac{\Gamma \vdash Q_y B(y) \qquad \Gamma \vdash B(\delta_B(t))}{\Gamma \vdash B(t)}\ \mathrm{QE}_{\delta}.$$

A target that is a literal successor asks for the instance at its predecessor; every other target of an atomic closure, and every target of a compound closure, asks for the instance at zero.

Lemma 2.109 (Interfaces and induction in T)**.** T *derives the equality rules and* QI_{η}*, and for a literal equation* E*, a compound matrix* B *and every term* t*:*

(i) *one step: from* $Q_y E(y)$ *and* $E(t)$ *to* $E(St)$*;*

(ii) *instance from the base: from* $Q_y E(y)$ *and* $E(0)$ *to* $E(t)$*, and from* $Q_y B(y)$ *and* $B(0)$ *to* $B(t)$*;*

(iii) *equation induction.*

Proof. The equality rules and QI_{η} are derived as in Theorem 2.92. (i) QE_{δ} at the target St. (ii) For B compound the rule is QE_{δ} itself. For E, take the target $J(t)$, which is not a literal successor: QE_{δ} gives $E(J(t))$ from $E(0)$, and $J(t) = t$ with Tr_2 gives $E(t)$. (iii) QI_{η} and (ii). □

This bootstrap uses no predecessor, no numerical case distinction and no identity for the orbit term.

Lemma 2.110 (Substitution in T)**.** *If* $\Gamma \vdash_{\mathsf{T}} A$*, then* $\Gamma[\theta] \vdash_{\mathsf{T}} A[\theta]$*. Injective renaming maps rule instances to rule instances; substitution in general does not.*

Proof. By induction on the derivation, with eigenvariables renamed. All rules but QE_{δ} commute with substitution. Consider QE_{δ} with the matrix B and the target t. If B is compound, or $t \equiv Sr$, or neither t nor $t[\theta]$ is a literal successor, the substituted inference is an instance. In the remaining case B is a literal equation, the second premise is the instance at 0, and $t[\theta]$ is a literal successor; then Theorem 2.109(ii) gives $B[\theta](t[\theta])$ from the substituted premises. That lemma does not depend on the present one. □

Lemma 2.111 (Transport in T)**.** *The statements (ii) and (iv) of Theorem 2.94 hold in* T*.*

Proof. The proof of Theorem 2.94 uses TR_2, QI_η, DE, DI_η, one step at a literal equation, the instance from the base at a compound matrix, and substitution. In T these are Theorems 2.109 and 2.110, and the instance from the base at a compound matrix is primitive, so no redex has to be removed. □

Theorem 2.112 (Target elimination and orbit elimination)**.** T *and* R *derive the same sequents. Hence Theorems 2.95 and 2.102 hold for* T*.*

Proof. The two calculi differ in the elimination of Q only.

QE_δ in R. At a literal equation and a target Sr it is one step (Theorem 2.94(i)). In the other cases its second premise is the instance at 0, and it is Theorem 2.94(iii); for a compound matrix this uses transport in a compound formula of R, which is why that lemma precedes the comparison.

QE_ω in T. At a compound matrix, QE_δ with the target $\omega(0, n)$ gives the conclusion from $B(0)$ at once. At a literal equation, let $Q_y E(y)$ and $E(a)$ be given. Equation induction on k in T gives $E(\omega(k, a))$: the base is $E(a)$ with $\omega(0, a) = a$, and the step is one step at $\omega(k, a)$ with $\omega(Sk, a) = S\,\omega(k, a)$. At $k := n$, the equation $\omega(n, a) = \omega(a, n)$, which holds in T by Theorem 2.93, gives $E(\omega(a, n))$. □

The orbit identities are needed to simulate the orbit rule in T. They are not needed by the rule QE_δ, by its soundness or by the induction of Theorem 2.109. Directly, QE_δ is sound for the progressive reading as follows: at a literal successor it uses one instance of progress, at another target of an atomic closure it uses equation induction from the instance at zero, and at a compound matrix it is the elimination of the conditional reading.

Remark 2.113 (Sources and Lean). The subsection is the first part of `target_closure`. In Lean the calculus is `TQPrf`, with `tqprf_iff_rprf`, the orbit rule derived as `TQPrf.orbit`, and the consequences `tqprf_emb_iff`, `tqprf_iff_rd` and `tqprf_consistent`. The second part of the package concerns the typing of certificates for the calculus with the conditional reading and is summarized in Section 2.7.

2.5.8 The calculi of the chapter

Table 2.1 lists the calculi of Sections 2.2 to 2.5. Each has HYP, I0, IS and DE; the second column gives its other rules. The calculus U_c is the exception: it is stated on normal terms and has neither I0 nor IS. The column "subst." says whether substitution maps rule instances to rule instances (Theorem 2.3); for U_c substitution is followed by normalization (Theorem 2.84).

Table 2.1: The calculi of the chapter.

Calculus	Rules besides HYP, I0, IS, DE	Rules	Relation to N	Subst.
N	TR, QE, DI, QI	8		yes
N_D	TR, QE, DI, DP	8	equivalent, Theorem 2.6	yes
N_det	TR, QE, DI, QI_η	8	equivalent, Theorem 2.10	yes
W	TR, QE, DI_w	7	equivalent, Theorem 2.17	no
N_2	TR_2, QE, DI, QI	8	equivalent, Theorem 2.21	yes
N_F	TR_F, QE, DI, QI	8	equivalent, Theorem 2.21	yes

Calculus	Rules besides Hyp, I0, IS, DE	Rules	Relation to N	Subst.
O	Tr$^-$, QE, DI, QI	8	equivalent, Theorem 2.27	no
M	Tr$^-$, QE, DI$_\mathrm{w}$	7	equivalent, Theorem 2.27	no
X	TrX, QX, DI, QI	8	equivalent, Theorem 2.32	no
Z	Tr^-_F, QE, DI, QI	8	equivalent, Theorem 2.35	no
G	Rep, QE, DI, QI	8	equivalent, Theorem 2.47	no
G$_7$	Rep, QE, DI$_\mathrm{w}$	7	equivalent, Theorem 2.47	no
L	LOR, QE, DI, QI	8	equivalent, Theorem 2.51	yes
N$_\mathrm{u}$	uncrossed replacement in a term template, QE, DI, QI	8	equivalent, Theorem 2.51	yes
U	Tr, QE, DI, UR	8	equivalent, Theorem 2.59	yes
U$_\mathrm{o}$	Tr, QE, DI, Orb	8	equivalent, Theorem 2.79	yes
U$_\mathrm{c}$	Ref, QE, DI, UX, on normal terms; no I0, IS	6	through nf, Theorem 2.86	yes
R	Tr$_2$, QE$_\omega$, DI$_\eta$	7	through $\uparrow$ and $\downarrow$, Theorem 2.102	yes
R$_F$	Tr$_F$, QE$_\omega$, DI$_\eta$	7	same sequents as R, Theorem 2.106	yes
T	Tr$_2$, QE$_\delta$, DI$_\eta$	7	same sequents as R, Theorem 2.112	no

2.6 Two results on other recursors

This section draws on the appendix of `witness_rule` on counter congruence and on the obstruction note of `scope_and_closure`. Both concern term languages other than that of Section 1.1; the formulas and the logical rules are the native ones.

2.6.1 Counter congruence at a zero seed

The appendix of `witness_rule` refines an earlier package, `counter_congruence`, whose term language has a recursor that sees the counter. The terms are variables, 0, St and $\mathrm{Rec}(n, a;\, u, v.\, h)$, which binds u, the previous counter, and v, the previous state, in h. The computation equations are

$$\mathrm{Rec}(0, a;\, u, v.\, h) = Sa, \qquad \mathrm{Rec}(Sn, a;\, u, v.\, h) = h[n/u][\mathrm{Rec}(n, a;\, u, v.\, h)/v].$$

The formulas are equations and $D_{x,y}A$; a closure $Q_y B$ occurs in a context only, as a resource in the sense of Section 1.9.

Definition 2.114 (The calculus $\mathcal{C}_0$)**.** The rules of $\mathcal{C}_0$ are: Hyp for a formula of the context; lookup of a closure of the context; the two computation equations; QE, with its first premise

a lookup; DE; DI; DP; root replacement on the left,

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash s = r}{\Gamma \vdash t = r}\ \text{RL};$$

and counter congruence at a zero seed,

$$\frac{\Gamma \vdash s = t}{\Gamma \vdash \mathrm{Rec}(s, 0;\, u, v.\, h) = \mathrm{Rec}(t, 0;\, u, v.\, h)}\ \text{ZCC} \qquad (v \text{ not free in } h).$$

The seed of ZCC is the literal 0, and its body does not depend on the state, also not below nested binders. There is no reflexivity, no general congruence and no transport. The reference calculus of `counter_congruence` has, in place of ZCC, the rule CC with an arbitrary seed a and an arbitrary body. Count one node for every rule instance, lookups included.

Lemma 2.115 (Root equality and cases)**.** *In $\mathcal{C}_0$, without ZCC:*

(i) *reflexivity is derivable with three nodes, and symmetry and transitivity are derivable;*

(ii) *from $l = r$, $l = f$ and $r = g$, two instances of RL give $f = g$;*

(iii) *numerical cases, from $A(0)$ and $A(Sz)$ to $A(t)$, are derivable with four nodes besides the two premises, with no equality rule and no DP.*

Proof. (i) $\mathrm{Rec}(Ss, 0;\, u, v.\, u) = s$ is a computation equation, and RL applied to two copies of it gives $s = s$. RL applied to $s = t$ and $s = s$ gives $t = s$, and applied to $t = s$ and $t = r$ it gives $s = r$. (ii) RL on $l = r$ and $l = f$ gives $r = f$, and on $r = f$ and $r = g$ it gives $f = g$. (iii) DI with an unused coordinate and the witness 0, DE, a lookup of the opened closure, and QE. □

Theorem 2.116 (Congruence from ZCC)**.** *For every term template $F(x)$, $\mathcal{C}_0$ derives $F(s) = F(t)$ from $s = t$, with exactly 20 nodes besides the derivation of $s = t$ and with no instance of DP. Consequently $\mathcal{C}_0$ derives transport in every formula template, and it derives the same sequents as the calculus with CC and as the calculus with unrestricted transport.*

Proof. Put $U(n) := \mathrm{Rec}(n, 0;\, u, v.\, SSu)$. Its computation equations are $U(0) = S0$ and $U(Sz) = SSz$, so numerical cases give $U(n) = Sn$ for every term n, with six nodes. ZCC applies to U and gives $U(s) = U(t)$; with $U(s) = Ss$ and $U(t) = St$, Theorem 2.115(ii) gives $Ss = St$. This stage has $1 + 6 + 6 + 2 = 15$ nodes.

Put $V_F(n) := \mathrm{Rec}(Sn, 0;\, u, v.\, F(u))$. One computation equation gives $V_F(n) = F(n)$. ZCC, applied to $Ss = St$ with the body $F(u)$, gives $V_F(s) = V_F(t)$, and Theorem 2.115(ii) with $V_F(s) = F(s)$ and $V_F(t) = F(t)$ gives $F(s) = F(t)$. This stage has $1 + 1 + 1 + 2 = 5$ nodes.

Transport in a formula template $A(w)$ follows by induction on the number of binders of A. For an equation $p(w) = q(w)$, join $p(s) = p(t)$ and $q(s) = q(t)$ with $p(s) = q(s)$ by Theorem 2.115. For $D_{x,y}B(w, x, y)$, apply DE at a fresh b and DI at the witness b: the assumption $B(t, b, 0)$ is transported back to $B(s, b, 0)$, QE with the opened closure gives $B(s, b, Sz)$, and transport forward gives $B(t, b, Sz)$.

Every instance of ZCC is an instance of CC, and CC is the congruence just proved for the template $\mathrm{Rec}(x, a;\, u, v.\, h)$. In the calculus with transport, RL is transport at the left root and ZCC is transport of one occurrence in an instance of reflexivity. □

The construction of `counter_congruence` has 56 nodes for the same congruence; replacing its final assembly by Theorem 2.115(ii) gives 44, and the representation through U and V_F gives 20. These are sizes of the constructions described, not lower bounds, and the terms in the nodes grow with F.

Proposition 2.117 (A wrapped rule)**.** *Let ZCC$^+$ infer* $\mathrm{Rec}(Ss, 0;\, u, v.\, h) = \mathrm{Rec}(St, 0;\, u, v.\, h)$ *from* $s = t$*, with* v *not free in* h*. Replacing ZCC by ZCC$^+$ gives the same derivable sequents, and congruence in a term template then has* 5 *nodes besides the derivation of* $s = t$*.*

Proof. ZCC$^+$ with the body $F(u)$ gives $V_F(s) = V_F(t)$ directly; two computation equations and two instances of RL finish. Conversely Theorem 2.116 derives ZCC$^+$, as congruence for the template $\mathrm{Rec}(Sx, 0;\, u, v.\, h)$. □

ZCC$^+$ wraps arbitrary premises in successors. It is not ZCC with a side condition on its premise, and the difference matters:

Proposition 2.118 (Successor premises do not suffice)**.** *Restrict ZCC to premises of the form* $Sp = Sq$*, and keep the other rules of* $\mathcal{C}_0$*. The resulting calculus does not derive* $0 = S0 \vdash S0 = SS0$*, which* $\mathcal{C}_0$ *derives.*

Proof. Interpret terms in the standard natural numbers, with the recursor read as primitive recursion with the seed $a + 1$. Interpret an equation $s = t$ as $e(s, t)$, where $e(m, n)$ holds if $m = n$ or both m and n are in $\{0, 1\}$; interpret closures and $D_{,}$ by their ordinary readings over this interpretation of equations. The relation e is an equivalence, so RL preserves truth. Computation equations are true because their two sides have equal values. For the restricted rule the two counters are positive, and on positive numbers e is equality, so a true premise gives equal counters and equal values of the recursor. Lookup, QE and DE preserve truth by the readings, DI because every number is zero or a successor, and DP by induction on the natural numbers for the interpreted predicate. The assumption $0 = S0$ is true, since $e(0, 1)$ holds, and $S0 = SS0$ is false, since $e(1, 2)$ fails. $\mathcal{C}_0$ derives the sequent by congruence for S. □

The interpretation is an interpretation of the rules; it does not read the equation sign as equality and is not a model of the ordinary theory. The package assigns no checker size to $\mathcal{C}_0$ or to the wrapped rule.

2.6.2 Recursors seeded at zero

The iterator has three arguments: a counter, a seed and a body. The note of `scope_and_closure` examines what is lost if the seed is fixed to zero. Consider two term languages over variables, 0 and S. The *scalar* language has $\mathrm{It}_0(n;\, u.\, h)$ with

$$\mathrm{It}_0(0;\, u.\, h) = 0, \qquad \mathrm{It}_0(Sn;\, u.\, h) = h[\mathrm{It}_0(n;\, u.\, h)/u].$$

The *counter-aware* language has $\mathrm{Rec}_0(n;\, k, u.\, h)$ with

$$\mathrm{Rec}_0(0;\, k, u.\, h) = 0, \qquad \mathrm{Rec}_0(Sn;\, k, u.\, h) = h[n/k][\mathrm{Rec}_0(n;\, k, u.\, h)/u].$$

All statements below are about the denotations of terms in the standard natural numbers.

Theorem 2.119 (Monotonicity)**.** *Every term of either language denotes a function that is nondecreasing in each of its free variables.*

Proof. By induction on the term. Variables, 0 and S are clear. For the recursor, let the counter n and the body h denote nondecreasing functions, fix an assignment ρ of the parameters, and let a^ρ_j be the state after j steps: $a^\rho_0 = 0$ and $a^\rho_{j+1} = h(j, a^\rho_j, \rho)$. The sequence is nondecreasing in j: $a^\rho_0 = 0 \le a^\rho_1$, and $a^\rho_j \le a^\rho_{j+1}$ gives $a^\rho_{j+1} = h(j, a^\rho_j, \rho) \le h(j + 1, a^\rho_{j+1}, \rho) = a^\rho_{j+2}$. For $\rho \le \sigma$, induction on j gives $a^\rho_j \le a^\sigma_j$. Since $n(\rho) \le n(\sigma)$,

$$a^\rho_{n(\rho)} \le a^\sigma_{n(\rho)} \le a^\sigma_{n(\sigma)}.$$

The scalar case is the same without the argument j of h. The argument covers a variable that occurs both in the counter and in the body. □

Theorem 2.120 (Zero sets in the scalar language)**.** *For a term $t(x_1, \dots, x_r)$ of the scalar language let $\mathcal{Z}(t) := \{\vec{x} : t(\vec{x}) = 0\}$. Then $\mathcal{Z}(t)$ is the whole of $\mathbb{N}^r$ or a union $\bigcup_{j \in J}\{\vec{x} : x_j = 0\}$ of coordinate hyperplanes, for some $J \subseteq \{1, \dots, r\}$, possibly empty. Every set of this form is the zero set of a term.*

Proof. By induction on t. The zero set of 0 is everything, that of a variable is its hyperplane, and that of a successor is empty. For $\mathrm{It}_0(n;\, u.\, h)$ fix the parameters and let b be the value of h at $u = 0$. If $b = 0$, every iterate from 0 is 0. If $b > 0$, monotonicity of h gives $h(v) \geq b > 0$ for every v, so every iterate after the first step is positive. Hence

$$\mathcal{Z}\big(\mathrm{It}_0(n;\, u.\, h)\big) = \mathcal{Z}(n) \cup \mathcal{Z}\big(h[0/u]\big).$$

The zero set of $h[0/u]$ is everything if the zero set of h is everything or contains the hyperplane $u = 0$, and otherwise it is the union of the same hyperplanes of the parameters. A union of two sets of the stated form is of that form. Conversely 0 and $S0$ realize the whole set and the empty set, and $\mathrm{It}_0(c;\, u.\, a)$ with u not in a has the zero set $\mathcal{Z}(c) \cup \mathcal{Z}(a)$, so nested terms of this kind realize every finite union of hyperplanes. □

Corollary 2.121 (What a zero seed loses)**.** (i) *No term of the scalar language denotes the predecessor, whose zero set is $\{0, 1\}$, or addition, whose zero set is $\{(0, 0)\}$.*

(ii) *No term of either language denotes the function that is 1 at 0 and 0 elsewhere. Hence no term denotes the selector $\mathrm{sel}(c, a, b)$, and no translation of terms recovers the iterator of Section 1.1 from either language on the same domain with the standard 0 and S.*

(iii) *In the counter-aware language $\mathrm{Rec}_0(n;\, k, u.\, k)$ denotes the predecessor, so Theorem 2.120 does not extend to it.*

Proof. (i) Theorem 2.120. (ii) The function decreases from 0 to 1, against Theorem 2.119; it is $\mathrm{sel}(c, 1, 0)$. (iii) The computation equations give 0 at 0 and n at Sn. □

These are statements about definability by terms on the same domain in the standard model. They do not exclude graphs of functions defined by formulas, definable subdomains, domains of tuples or quotient equalities; an interpretation of that kind would need its own proofs of totality and functionality, of the transfer of induction and of both composites. Whether addition is denoted by a term of the counter-aware language is not decided in the package. In Lean, Theorem 2.119 and part (ii) of the corollary are `ZTm.mono` and `ZTm.no_zero_test` in the module `ZeroSeedObstruction`, for a zero-seeded recursor whose body sees the counter, with `Tm.val_zf` for the selector of the kernel's language.

2.7 Checker sizes

This section collects the measurements reported by `direct_closure`, `direct_closure_qprogress`, `witness_rule`, `source_dispatch`, `reversal_core`, `one_context`, `one_sided_transport`, `template_free`, `scalar_uniqueness`, `orbit_closure`, `crossed_uniqueness`, `uniform_progress` and `target_closure`. All numbers are for complete standalone Python programs under the protocol of Section 1.10, as the packages state them. Table 2.2 gives, for each program, its significant tokens, AST nodes and normalized bytes, and the same three numbers for the control it is matched against. Programs in different blocks have different input contracts and are not comparable across blocks. The compact programs of `kernel_refinement` and of `scope_and_closure` implement calculi of other chapters and are reported there; neither package measures a checker for the calculi L and R_F of this chapter.

Table 2.2: Checker sizes reported by the packages of the chapter: tokens, AST nodes, normalized bytes; the control is given as tokens / AST nodes / bytes.

Package	Program	Tok.	AST	Bytes	Matched control
`direct_closure`	strict tagged checker with QI	1169	1337	5116	with DP: 1187 / 1358 / 5199
	compact checker with counter vectors (experiment)	271	314	533	byte profile of `neutral_scope`: 266 / 310 / 529
`direct_closure_qprogress`	compact checker, QI at literal equations only	285	331	568	`neutral_scope`: 266 / 310 / 529
	compact checker for $\mathsf{N}_{\mathrm{det}}$	278	322	552	`neutral_scope`: 266 / 310 / 529
`witness_rule`	compact checker, two introductions, Boolean dispatch	264	310	550	byte profile of `certified_zero`: 264 / 310 / 558
	compact checker for W	266	312	555	the same; joint profile of `certified_zero`: 262 / 296 / 598
	the same with an explicit type test	266	312	559	the same
	the same with an explicit list test	266	310	563	the same
`source_dispatch`	byte profile	266	305	526	266 / 310 / 529
	AST profile	263	283	607	263 / 289 / 613
	token profile	259	289	572	261 / 292 / 576
`reversal_core`	typed checker for X	456	520	1055	old typed control: 532 / 618 / 1365; same refactoring without reversal: 450 / 514 / 1049
	X with a validator for goals of equations and $D_{,}$	379	432	830	old: 450 / 521 / 1109; without reversal: 373 / 426 / 824
	X with a validator for all goals	405	462	879	old: 481 / 560 / 1189; without reversal: 399 / 456 / 873
	compact byte profile with a scalar neutral premise	265	304	525	266 / 305 / 526
	compact AST profile with a scalar neutral premise	262	282	606	263 / 283 / 607
	compact checker for X	271	310	531	none stated
`one_context`	tagged checker for N	1089	1150	3358	for N_2: 1101 / 1160 / 3382
	strict validator, equations and $D_{,}$	359	414	803	373 / 426 / 824
	strict validator, all formulas	391	444	863	399 / 456 / 873
	full typing of the certificate	442	502	1039	450 / 514 / 1049
	strict validator with reversal, equations and $D_{,}$	365	420	809	379 / 432 / 830
	strict validator with reversal, all formulas	397	450	869	405 / 462 / 879

Package	Program	Tok.	AST	Bytes	Matched control
	full typing with reversal	448	508	1045	456 / 520 / 1055
`one_sided_ transport`	strict checker for M	1171	1339	3246	$\textsc{Tr}_2$ at the same sources: 1196 / 1364 / 3302
`template_free`	strict tagged checker for G_7	1027	1072	3147	W with three traversals: 1029 / 1089 / 3181; 1059 / 1124 / 3215; 1044 / 1096 / 3206
	compact forward checker with $\textsc{Rep}$	397	464	865	crossed rule, another packet layout: 384 / 448 / 830
`scalar_ uniqueness`	tagged checker for U	1124	1175	3393	for N: 1084 / 1141 / 3322
	validated codec with scope reuse, calculus unchanged	381	439	866	383 / 442 / 870
`orbit_closure`	tagged checker for U_{o}, templates supplied	1132	1178	3400	for U: 1124 / 1175 / 3393
	the same, templates read off the closure	1145	1194	3455	the same
`crossed_ uniqueness`	strict checker for U_{c}, normalizer included	1909	2087	5618	none stated
	compact checker of the guarded calculus, not of U_{c}	269	301	502	byte profile of `deterministic_ introduction`: 267 / 303 / 506
`uniform_progress`	typed checker for R	1206	1366	4553	for N: 1235 / 1407 / 4692; for W: 1173 / 1334 / 4456
	compact token profile, guarded-source calculus	256	283	593	257 / 285 / 568
	compact AST profile, the same calculus	260	278	602	261 / 279 / 603
	compact byte profile, the same calculus	263	300	521	264 / 301 / 522
`target_closure`	typed checker for T	1195	1358	4532	for R: 1206 / 1366 / 4553
	typed checker of the conditional calculus with a lexical gate	399	453	895	recursive typing: 442 / 502 / 1039
	the same with a descriptive error	404	455	963	the same

The following paragraphs summarize what the programs do beyond the rules.

`direct_closure`. The two strict checkers are built from one body and two fragments for the progress rule; they differ in that branch only. The saving is the witness, its scope check and one substitution; the normalized source compressed by gzip is 19 bytes larger. The experiment with counter vectors passes the counter of a computation as an empty or nonempty list, so that $I(1, a;\, u.\, h)$ is certified without the repair of Theorem 2.53; it costs four bytes. A variant of 528 bytes writes the empty list where the protocol uses the scalar zero, a canonical certificate fails on it, and it is not counted as a complete checker.

`direct_closure_qprogress`. The compact programs keep the arity encoding of the con-

trol. A failed test on the matrix must reject and not return a false value, which in this encoding would enter the context as the scalar zero; the first program uses a sentinel that is not JSON for the failure, and the second gives a compound matrix the mode of cases.

witness_rule. Every selected first premise is truthy and every generated computation equation is a nonempty list, so one Boolean expression selects the branch; null markers become empty lists. This saves eight bytes with the calculus unchanged, and the test of the witness adds five. The package proves that every accepting run is simulated, after a fixed change of markers, by an accepting run of the control.

source_dispatch. In canonical certificates the source is falsy exactly when it is the literal 0, so the comparison of the first premise against two patterns becomes one comparison selected by the source. The byte and AST profiles also move the pure computation of the replaced goal before the check of the first premise, and the byte profile uses the checker function itself as the invalid image of a substitution.

reversal_core. The typed checker shares one validator between terms and formulas and writes the neutral premise as a scalar; these two changes give the reduction of 310 bytes against the old control, and reversal itself adds six bytes. The compact programs of 525 bytes and 282 AST nodes check certificates without reversal, for the equivalent calculus.

one_context. In the tagged checker only the branch of transport changes. Independently, the strict validator of the goal omits the upper bound on variable indices: for a tree of the right shape that bound holds exactly when the first traversal of the unchanged kernel succeeds, which happens before any instruction is read, so the sets of accepted certificates are equal. The sizes after gzip increase in all six pairs.

one_sided_transport. The fixed side r of $\textsc{Tr}^{-}$ is read off the conclusion, so the instruction carries three terms where the control carries four.

template_free. One traversal serves lifting, substitution and replacement; an instruction for Rep consists of its two premise derivations. Against the control with the same traversal the saving is 59 bytes. The compact forward program synthesizes its conclusion and is 35 bytes larger than its control, which has another packet layout.

scalar_uniqueness. The instruction for UR carries the two templates, the target and the premise, checked with no added assumption; the direct replacement of QI by UR costs 71 bytes. The second pair keeps the calculus of `N`: a fully typed codec whose goal validator relaxes the test `2 < a < d` on a variable index to `a > 2` and leaves the upper bound to the first scope traversal of the kernel, the factorization described for `one_context`; the change is four bytes.

orbit_closure. The instruction for Orb carries a proposed closure and one premise. The selected checker reads f and h off the closure, compares the seed with the instance of f at zero, and rejects a body that mentions the counter (Theorem 2.80); it is 62 bytes larger than its control, the checker for `U` of `scalar_uniqueness`. The translator of the package is not trusted; its output is checked by the target checker. The package also audits every Python source of the corpus, 5,825 paths with 1,380 distinct contents, by parsing and measuring without executing them, and finds the least values of the three coordinates, 256 tokens, 276 AST nodes and 506 bytes, in three different programs of other packages; it records which smaller files are not complete standalone checkers.

crossed_uniqueness. The checker for U_{c} counts the normalizer, substitution and all scope checks, introduces the eigenvariable of UX itself, and checks scopes before normalizing, so that a subterm erased by a computation is still checked. Separately, the package composes three known refinements (delayed elimination, eager selection from a pair, a shorter selector of the mode) into a compact checker of the guarded calculus of another chapter with 502 bytes against 506, and a validating companion with 1172 bytes against 1183; neither implements U_{c}, and the least token and AST counts of the corpus are not improved.

uniform_progress. The typed checker for `R` is 139 bytes smaller than the matched checker

for N and 97 bytes larger than the matched checker for W. Its three compact programs are not checkers for R; they turn the continuation into a loop over an immutable context and write the neutral premise as a scalar.

target_closure. The checker for T saves 21 bytes against that for R by dropping the source and the seed of the orbit, and remains 76 bytes larger than that for W. The lexical gate replaces the recursive test that every value of a certificate is an integer or a list by a test on the input text: a text that the JSON parser accepts decodes to such a value exactly when all its characters are among the brackets, the comma, the minus sign, the digits and the four whitespace characters of JSON.

2.8 Further questions

The packages of the chapter state the following questions, limits and unproved claims. None of them claims a smallest calculus or a lower bound on the size of a checker.

Question 2.122 (Checkers for the direct closure)**.** The compact checkers of `direct_closure_qprogress` with QI are larger than their control, because the two openings at the root must be distinguished. Is there an encoding in which the direct closure rule gives a smaller complete checker, with malformed, aliased and scope-violating inputs counted?

Question 2.123 (The variant with counter vectors)**.** The 528-byte variant in `direct_closure` fails on a canonical certificate because the base counter is written as an empty list. Is there a conversion between the two spellings of zero that makes it complete? No false theorem is known to be accepted by it.

Question 2.124 (Deleting an inference case)**.** The direct closure rule removes a witness and keeps eight rules. Is there a basis on the same formulas that deletes a case of the checker and preserves arbitrary contexts and both formula round trips? (`direct_closure_qprogress`.)

Question 2.125 (Focused derivations)**.** Can a normalization theorem that permutes introductions and eliminations justify a simpler protocol for first premises? Equation induction makes the theorem nontrivial, and it has to precede any removal of checking. (`direct_closure_qprogress`.)

Question 2.126 (One-template transport in a compact protocol)**.** Can a certificate format expose TR or TR^{-} and keep the opening routine shared with QE, so that the complete checker is smaller than the compact programs under the same contract for the goal? (`one_context`, `one_sided_transport`.)

Question 2.127 (Sharing in certificates)**.** What is the least overhead in the size of certificates when TR_2 is replaced by TR? A certificate with sharing avoids the copy of the first premise (Theorem 2.22); can a checker for such certificates be smaller? (`one_context`, `one_sided_transport`.)

Question 2.128 (Constant templates)**.** Symmetry is derived from TR with a constant template. Is there an equality rule simpler than TR^{-} that keeps this step? A restriction to templates in which the variable occurs must replace it. (`one_sided_transport`.)

Question 2.129 (Removing both syntactic tests)**.** The test $s \not\equiv 0$ of TR^{-} and the test of the witness in DI_{w} serve the dispatch of compact checkers; the unrestricted rules are admissible. Can both tests be removed while a compact complete checker is kept? (`one_sided_transport`.)

Question 2.130 (The forward computation equation)**.** If the successor computation is replaced by $I(Sn, a;\, u.\, h) = I(n, h(a);\, u.\, h)$, is the equation IS derivable? The induction that suggests itself needs its hypothesis at the changed seed $h(a)$, while equation induction keeps parameters fixed. `one_sided_transport` leaves this open and claims no independence.

Question 2.131 (A fixed input contract)**.** Which contract should later comparisons fix: goals of equations and $D_{,}$, goals of all native formulas, or exact typing of the whole certificate? The minima reported by the packages are attained by different programs under different contracts. (`one_context`, `scalar_uniqueness`.)

Question 2.132 (Replacement without a template in a compact checker)**.** Does the rule Rep give a smaller complete checker under a fixed compact contract? The compact forward program of `template_free` is larger than its control.

Question 2.133 (Replacement of one occurrence in a checker)**.** Does LOR help a readable or a formally verified checker, even though a traversal by position costs more than it saves in the compact programs? (`kernel_refinement`.)

Question 2.134 (Uniqueness with fewer bodies)**.** Can UR be restricted to a family of bodies and still give the predecessor of Theorem 2.70? The reflection of equality uses mostly the case of a constant recurrence (Theorem 2.60(ii)), while the predecessor uses other bodies. For the retractions the question is answered: Theorem 2.62 needs the body Su only. (`scalar_uniqueness`, `orbit_closure`.)

Question 2.135 (The strength of successor uniqueness)**.** Does T_{SU} derive induction for all literal equations? A positive answer has to supply a predecessor and a term that reflects equality from the body Su alone; a negative answer has to give a model of the computation equations for every body and of every instance of SU over arbitrary contexts in which equation induction fails. The model that separates induction for zero equations from induction for all equations is not such a model (Theorem 2.82). (`orbit_closure`.)

Question 2.136 (Successor and identity bodies)**.** Does SU together with uniqueness for the body u (Theorem 2.60(ii)) give induction for all equations? Does T_{SU} prove that addition is cancellative, which by Theorem 2.81 would give uniqueness for the body u? (`orbit_closure`.)

Question 2.137 (Compact encodings for the other forms of uniqueness)**.** Can Orb be combined with the compact encodings of goals without adding more recognition of syntax than it removes? Can UX receive a compact encoding that keeps the freshness of its eigenvariable and the independence of its body from the counter, and can the normalizer share the traversal by arity while respecting sorted positions, protected slots and reindexing? The checkers for U_{o} and U_{c} are larger than their comparators. (`orbit_closure`, `crossed_uniqueness`.)

Question 2.138 (Crossed uniqueness with clauses)**.** Can a calculus with clause contexts fuse UX with the discharge of a witness without complicating the derivation of equality in Theorem 2.85? (`crossed_uniqueness`.)

Question 2.139 (Uses of the accumulator)**.** The term φ of Theorem 2.74 is a composition of subtraction and addition with a recurrence that depends on the counter. Does it give a more direct compiler for primitive recursion, or a shorter predecessor, without coding tuples? (`scalar_uniqueness`.)

Question 2.140 (A checker for uniqueness)**.** The checker for U is 71 bytes larger than its control. Is there an architecture that shares the check of a computation equation with the check of UR and reverses the comparison? (`scalar_uniqueness`.)

Question 2.141 (Cost of the root-sensitive closure)**.** Can R, R_F or T be checked with less total work than the test of the witness in W? The matched checkers for R and T are larger than that for W, and the comparison has to include the instances at the source and at the target, the construction of the orbit term and the test of the root. (`uniform_progress`, `target_closure`, `scope_and_closure`.)

Question 2.142 (A zero seed through graphs)**.** Theorem 2.121 excludes a translation of terms. Can an interpretation by graphs of functions recover an arbitrary seed from a language with the seed fixed to zero, with proofs of totality and functionality, the transfer of induction and both composites? Is addition denoted by a term of the counter-aware language of Section 2.6.2? (`scope_and_closure`.)

Question 2.143 (Lower bounds)**.** Is there a lower bound on the size of presentations in a fixed class, with a fixed encoding and cost model? A census of existing programs is not such a bound. (`one_context`, `scope_and_closure`.)

The following statements of the packages are recorded without proof or as results that were not achieved.

- `direct_closure_qprogress` states that replacing the two computation equations by a rule of conversion in a context would still need both rewriting patterns and an account of reflexivity, and that a conversion engine left out of the count would not be a smaller checker. No calculus of this kind is given.
- `witness_rule`: no program of the package is below all three earlier coordinate minima; the restriction of the introduction is not credited with a saving of bytes.
- `one_context` states that removing a primitive equality test from the self-seeded proposals of other packages is a missing theorem and does not follow from coverage of a protected orbit.
- `scalar_uniqueness`: the converse of Theorem 2.57 is a written proof; it is not formalized in Lean, where only Theorem 2.57 is proved.
- `orbit_closure`: the equivalence of T_{SU} with T_{eq} is not established, and the package does not claim it. Its statement about the model of `canonical_recursors` (Theorem 2.82) rests on the construction of that package. The rule Orb removes a field of the certificate and makes the matched checker larger, by 7 bytes with supplied templates and by 62 bytes with templates read off the closure.
- `crossed_uniqueness`: six rule names do not by themselves make a smaller presentation than the three-family clause calculi of other packages; the checker for U_{c} is the largest of Table 2.2, and its normalizer is counted. The 502-byte program lowers one coordinate for a calculus of another chapter. The package asks whether a relational bi-interpretation on a changed domain could reduce a complete checker; the obstructions to definability by terms do not exclude it, and both composites would need proofs.
- `orbit_closure`, `crossed_uniqueness`: neither is formalized in Lean, and both rest on the written proof of Theorem 2.58.
- `template_free`: the calculus G_7, with Rep and the single introduction together, is not formalized in Lean.
- `uniform_progress`, `target_closure`: neither calculus gives a new minimum in any coordinate of the matched comparison; the compact minima of other packages are not improved by a checker for R or T.
- `target_closure` reports two routes that gave no result: making the guard of an atomic matrix implicit simplifies the datatype of goals and adds a cost of elaboration that loses against a validator factored in the same way; and a stopped-clock construction in a self-seeded arithmetic gives a diagonal equation and does not prove that the successor is injective. No independence result is inferred from either.

- `scope_and_closure`: the obstruction of Section 2.6.2 is about terms in the standard model and is not an impossibility theorem for simpler axiomatizations.
- The proofs of this chapter are written proofs; the Lean names in the remarks mark the statements that are formalized. The Python checkers are not formally verified, and the tests reported by the packages concern particular certificates.

Chapter 3

Guards and public languages

The native scalar core has two introduction rules: DI by cases, at every matrix, and QI by progress, at literal equations (Section 1.3). This chapter treats calculi with a single introduction of $D_{,}$. Its premise retains one local assumption, the instance of the matrix at zero (*cases*) or the instance at the eigenvariable (*progress*), and the choice between the two is made by the shape of the matrix alone: no certificate bit, no test of the witness. At a literal equation the two modes prove different formulas. They prove the same formula when the instance at zero is provable, and a *guard* is a literal shape of an equation whose instance at zero is provable by computation and into which every equation can be lifted. The guard is used in two ways. Either the formula language is restricted to a *public language* in which every atomic matrix below $D_{,}$ has the guarded shape, and an arbitrary native formula is translated into it; or the full language is kept and the progress mode is used exactly at the guarded shape.

The chapter is organized by idea. Section 3.1 defines the guarded calculi, the shapes, and what substitution does to them. Section 3.2 recovers numerical cases, implication between equations and equation induction from a guard, once for all guards, and translates the native language into a public language with both composites. Section 3.3 determines which transport sources are needed: the literal 0 is not. Section 3.4 shows that the two restrictions together are incomplete in a strictly guarded language, and gives four repairs: auxiliary formulas, the administrative orbit, the mirror guard, and protected atoms. Section 3.5 treats the guards that work on the full native language, the diagonal guard and the mirror guard with six rules. Section 3.6 compares the guards, and Section 3.7 records the checkers and their sizes.

Throughout the chapter 1 abbreviates $S0$ and 2 abbreviates $SS0$; $s \equiv t$ is literal identity of terms; a term is *iterator-headed* if it has the form $I(n, a;\, u.\, h)$ and *successor-headed* if it has the form St; and

$$W(r) := I(0, r;\, u.\, 0)$$

is the *wrapper* of r, an iterator-headed term with $W(r) = r$ by the first computation equation. For a calculus K and formulas X, Y of its language, $X \dashv\vdash_{\mathsf{K}} Y$ means that $\Gamma \vdash_{\mathsf{K}} X$ if and only if $\Gamma \vdash_{\mathsf{K}} Y$, for every context Γ of K; the subscript is omitted when the calculus is clear.

In Sections 3.1 to 3.4 transport is the uncrossed rule in a formula template,

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash F(s)}{\Gamma \vdash F(t)},$$

as the packages of those sections state it. For N it is a derived rule (Theorem 1.23). It is the rule written $\textsc{Tr}_F$ in Section 2.3.4, and its restriction to the sources $s \not\equiv 0$, written $\textsc{Tr}^{-}$ below, is the rule $\textsc{Tr}^{-}_F$ of that section, not the crossed rule of Theorem 2.24. The orientation of the conclusion matters as soon as the sources s are restricted (Theorem 3.34). In Section 3.5 the equality rule is the rule $\textsc{Tr}$ of the kernel.

3.1 Guards as literal shapes

This section collects the definitions common to `guarded_core`, `guarded_source`, `guarded_orbit`, `deterministic_auxiliary`, `mirror_guard`, `mirror_core`, `protected_core` and `diagonal_guard`.

3.1.1 The root-determined introduction

Definition 3.1 (Root-determined introduction)**.** For a matrix $B(x,y)$ put

$$\eta_B(z) := \begin{cases} z, & B \text{ is a literal equation,} \\ 0, & B \text{ is compound (its root is } Q \text{ or } D_{,}). \end{cases}$$

The *root-determined introduction* is the rule

$$\frac{\Gamma,\ B(t,\eta_B(z)) \vdash B(t,Sz)}{\Gamma \vdash D_{x,y}B(x,y)}\ \mathrm{DI}_{\mathrm{root}},$$

where z is not free in Γ, in the witness t or in $D_{x,y}B$, and the displayed assumption is discharged. The rule is in *progress mode* at a literal equation and in *case mode* at a compound matrix.

The mode is read off the matrix before the witness is substituted, and it is the same for every substitution instance of the matrix, because substitution of terms for variables preserves and reflects the root of a formula.

Definition 3.2 (Guarded calculus)**.** Let $\mathcal{F}$ be a set of native formulas, the *proof language*, and $\mathcal{T} \subseteq \mathcal{F}$ a set of *templates*. Both contain every equation and are closed under substitution of terms for free variables; $\mathcal{F}$ contains $B(t,r)$ and $Q_yB(t,y)$ whenever it contains $D_{x,y}B(x,y)$, and $B(t)$ whenever it contains $Q_yB(y)$. Let $\mathcal{S}$ be a set of terms closed under injective renaming of variables, the *admissible sources*. The *guarded calculus* $\mathsf{K}(\mathcal{F},\mathcal{S})$ derives sequents $\Gamma \vdash A$, with Γ a finite list of formulas of $\mathcal{F}$ and $A \in \mathcal{F}$, by the following seven rules.

$$\frac{A \in \Gamma}{\Gamma \vdash A}\ \textsc{Hyp} \qquad \frac{}{\Gamma \vdash I(0,a;\,u.\,h) = a}\ \mathrm{I0} \qquad \frac{}{\Gamma \vdash I(Sn,a;\,u.\,h) = h[I(n,a;\,u.\,h)/u]}\ \mathrm{IS}$$

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash F(s)}{\Gamma \vdash F(t)}\ \textsc{Tr}_{\mathcal{S}} \quad (s \in \mathcal{S},\ F \in \mathcal{T}) \qquad \frac{\Gamma \vdash Q_yB(y) \qquad \Gamma \vdash B(0)}{\Gamma \vdash B(t)}\ \mathrm{QE}$$

$$\frac{\Gamma \vdash D_{x,y}B(x,y) \qquad \Gamma,\ Q_yB(b,y) \vdash A}{\Gamma \vdash A}\ \mathrm{DE} \qquad \frac{\Gamma,\ B(t,\eta_B(z)) \vdash B(t,Sz)}{\Gamma \vdash D_{x,y}B(x,y)}\ \mathrm{DI}_{\mathrm{root}}$$

In DE the variable b is not free in Γ, A or $D_{x,y}B$; in $\mathrm{DI}_{\mathrm{root}}$ the conclusion belongs to $\mathcal{F}$ and z is not free in Γ, t or $D_{x,y}B$. There is no reflexivity axiom, no successor axiom, no rule of induction, no rule that introduces Q, and no structural rule.

Three classes of sources occur: all terms; the terms $s \not\equiv 0$; and the iterator-headed terms. The condition $s \not\equiv 0$ is a test of syntax: a variable, or an iterator whose value is zero, is an admissible source. The rule with the sources $s \not\equiv 0$ is written $\textsc{Tr}^{-}$ and the rule with iterator-headed sources $\textsc{Tr}_I$. Two kinds of proof language occur: the full native language $\mathcal{L}$, with $\mathcal{T} = \mathcal{L}$; and the proof language $\mathcal{P}^Q$ of a public language $\mathcal{P}$ (Theorem 3.16), in which Q occurs only at the root of a hypothesis and the templates are the equations and the formulas of $\mathcal{P}$.

Remark 3.3 (Source-selected elimination). With the sources $s \not\equiv 0$ the two rules $\textsc{Tr}^{-}$ and QE are the two branches of the rule SE of Section 2.3.4, whose first premise is $Q_vF(v)$ at the source 0 and $s = t$ at every other source. In this form, which is the one of `guarded_source`,

`guarded_orbit` and `deterministic_auxiliary`, a guarded calculus has six rule patterns: the source selects the elimination and the root of the matrix selects the introduction. The calculus Z of Theorem 2.34 has SE with the two introductions of N; the calculi of this chapter have it with $\mathrm{DI}_{\mathrm{root}}$ alone.

Remark 3.4 (Clause contexts). `guarded_core` and `mirror_guard` state their calculi with the clause contexts of Section 1.9: DE adds the clause $[B(b,0);\, y.\, B(b,y)]$, an assumption is a constant clause, and one rule Use infers $B(t)$ from a derivation of A and a clause $[A;\, y.\, B(y)]$ of the context. A hypothesis $Q_y B(y)$ used by Hyp and QE is the clause $[B(0);\, y.\, B(y)]$ used by Use, so the calculi below are the clause calculi of those packages for contexts of assumptions and opened clauses. The statements and proofs of this chapter hold verbatim for contexts with arbitrary clauses $[A;\, y.\, B(y)]$ of public formulas, with a translation of clauses componentwise.

3.1.2 Structural operations, the kernel, and equality at admissible sources

Lemma 3.5 (Structural operations)**.** *In every guarded calculus $\mathsf{K}(\mathcal{F},\mathcal{S})$ weakening, injective renaming of free variables and cut are admissible: if $\Gamma \vdash A$ and $\Delta \vdash C$ for every $C \in \Gamma$, then $\Delta \vdash A$. If $\mathcal{S}$ is closed under substitution, as are the class of all terms and the class of iterator-headed terms, then substitution is admissible: $\Gamma \vdash A$ implies $\Gamma[\theta] \vdash A[\theta]$.*

Proof. Each statement is proved by induction on the derivation, with the eigenvariables of DE and $\mathrm{DI}_{\mathrm{root}}$ renamed away from the data that are inserted, as for N (Section 1.4). Weakening and cut do not change the matrix of an introduction or the source of a transport, so every inference keeps its rule and its mode. A renaming, and a substitution θ, preserve the root of every formula, so an introduction in progress mode remains one in progress mode and likewise for case mode; the languages $\mathcal{F}$ and $\mathcal{T}$ are closed under substitution; and the source $s[\theta]$ of a transported equation is admissible when $\mathcal{S}$ is closed under θ, which holds for every injective renaming and, in the two named cases, for every substitution. □

The class of terms $s \not\equiv 0$ is not closed under substitution: a variable is an admissible source and its instance 0 is not. A derivation with Tr^{-} therefore cannot be substituted into rule by rule. The arguments of Sections 3.3 and 3.4 are arranged so that no such substitution is used before the unrestricted rule has been recovered.

Lemma 3.6 (Inclusion in the kernel)**.** *Every derivation of a guarded calculus $\mathsf{K}(\mathcal{F},\mathcal{S})$ is transformed, rule by rule, into a derivation of the same sequent in N.*

Proof. Hyp, I0, IS, QE and DE are rules of N. $\mathrm{Tr}_{\mathcal{S}}$ is an instance of transport in a formula template (Theorem 1.23). $\mathrm{DI}_{\mathrm{root}}$ at a compound matrix is DI, and at a literal equation it is the witnessed introduction by progress, which N derives (Theorem 1.16(i)). □

Consequently every guarded calculus is sound for the ordinary reading (Section 1.2): a derivable sequent of $\mathsf{K}(\mathcal{F},\mathcal{S})$ is derivable in N, hence its reading is provable in T_{eq} and in iPRA (Theorem 1.78).

Lemma 3.7 (Equality at admissible sources)**.** *Let $\mathsf{K} = \mathsf{K}(\mathcal{F},\mathcal{S})$ and let $\mathcal{S}$ contain every iterator-headed term. The following hold over every context Γ, by derivations that use only I0, IS and $\mathrm{Tr}_{\mathcal{S}}$ in equation templates.*

(i) *$\Gamma \vdash r = r$ for every term r.*

(ii) *If $\Gamma \vdash p = q$ and $p \in \mathcal{S}$, then $\Gamma \vdash q = p$; $\Gamma \vdash f(p) = f(q)$ for every term template $f(v)$; $\Gamma \vdash r = p$ implies $\Gamma \vdash r = q$; and $\Gamma \vdash E(p)$ implies $\Gamma \vdash E(q)$ for every equation template $E(v)$.*

(iii) *Every instance of I0 and IS can be reversed.*

(iv) Join at zero. *If* $p, q \in \mathcal{S}$, $\Gamma \vdash p = 0$ *and* $\Gamma \vdash q = 0$, *then* $\Gamma \vdash p = q$.

(v) Lifting. *Let* $p, a, b \in \mathcal{S}$, *let* $q \in \mathcal{S}$ *or* $q \equiv 0$, *and let* $\Gamma \vdash a = p$ *and* $\Gamma \vdash b = q$. *Then* $\Gamma \vdash p = q$ *if and only if* $\Gamma \vdash a = b$. *The implication from* $a = b$ *to* $p = q$ *needs no condition on* q.

(vi) Expansion on the right. *Let* $\mathcal{S}$ *contain every term other than the literal* 0, *let* $q \not\equiv 0$ *and* $\Gamma \vdash q = r$. *Then, for every term* s, $\Gamma \vdash s = q$ *if and only if* $\Gamma \vdash s = r$.

Proof. (i)–(iv) For the sources $s \not\equiv 0$ these are Theorems 2.36 and 2.37; the wrapper $W(r)$ is the term $J(r)$ used there. The derivations given there transport only at the sources p and q named in each item and at $W(0)$, $W(p)$, $W(q)$ and $W(r)$, and the left side of a computation equation is iterator-headed. Hence they are derivations of K for every class $\mathcal{S}$ that contains the iterator-headed terms, under the stated conditions $p, q \in \mathcal{S}$. No equation with the source 0 is used, and in particular transitivity through the literal 0 is avoided.

(v) Let $\Gamma \vdash p = q$. Transitivity through p gives $a = q$. If $q \in \mathcal{S}$, reverse $b = q$ at the source b and pass through q: $a = b$. If $q \equiv 0$, then $a = 0$ and $b = 0$, and (iv) gives $a = b$. Conversely let $\Gamma \vdash a = b$. Transport along $a = p$ in the template $v = b$ gives $p = b$, and transport along $b = q$ in the template $p = v$ gives $p = q$; the sources are a and b.

(vi) From $s = q$, transport along $q = r$ in the template $s = v$ gives $s = r$. Conversely let $\Gamma \vdash s = r$. If $r \not\equiv 0$, reverse $q = r$ at the source q and transport at the source r in the template $s = v$. If $r \equiv 0$ and $s \not\equiv 0$, then $s = 0$ and $q = 0$, and (iv) gives $s = q$. If $r \equiv 0$ and $s \equiv 0$, the required equation $0 = q$ is the reverse of $q = 0$ at the source q. The case distinction is on the syntax of the terms r and s; it is not a case distinction inside the calculus. □

With all terms as sources, (i) and (ii) are the ordinary equality rules, and the calculus has reflexivity, symmetry, transitivity, congruence and transport in every equation template.

3.1.3 Deleting the choice is not conservative

The administrative calculus is $\mathsf{U} := \mathsf{K}(\mathcal{L}, \text{all terms})$: the full native language with the root-determined introduction. It is the calculus that the compact checkers of `guarded_core` implement, and by Theorem 3.6 it is a subcalculus of N. It is a proper one.

Definition 3.8 (Mixed reading)**.** The *mixed reading* $A^\dagger$ of a native formula is

$$
\begin{aligned}
(s = t)^\dagger &:= (s = t),\\
(Q_y B)^\dagger &:= B^\dagger(0) \to \forall y\, B^\dagger(y),\\
(D_{x,y} B)^\dagger &:= \begin{cases} \exists x\, \forall y\, (B(x,y) \to B(x,Sy)), & B \text{ a literal equation},\\ \exists x\, (B^\dagger(x,0) \to \forall y\, B^\dagger(x,y)), & B \text{ compound}. \end{cases}
\end{aligned}
$$

The mixed reading depends on the syntax of the matrix, not on its meaning: it is not invariant under the replacement of a matrix by an interderivable one. It commutes with substitution, because substitution preserves the root of a matrix.

Lemma 3.9 (Soundness for the mixed reading)**.** *If* $\Gamma \vdash_{\mathsf{U}} A$, *then* T_{eq} *proves* $A^\dagger$ *from* $\Gamma^\dagger$.

Proof. By induction on the derivation. Hyp, the computation equations and QE are immediate, and transport is sound because the reading commutes with substitution. At an atomic $\mathrm{DI}_{\mathrm{root}}$ the premise proves $B(t,z) \to B(t,Sz)$ for a fresh z, and generalization on z followed by existential introduction gives the progressive reading. At a compound $\mathrm{DI}_{\mathrm{root}}$

the premise proves $B^\dagger(t,0) \to B^\dagger(t,Sz)$, and coverage $y = 0 \vee \exists z\, y = Sz$ extends it to $B^\dagger(t,0) \to \forall y\, B^\dagger(t,y)$. For DE at a literal equation E, fix x with $\forall y\,(E(x,y) \to E(x,Sy))$; induction for the literal equation $E(x,y)$ in T_{eq} gives $E(x,0) \to \forall y\, E(x,y)$, which is the reading of the hypothesis $Q_y E(b,y)$ that the rule supplies. DE at a compound matrix is existential elimination. □

Proposition 3.10 (The root-determined introduction alone is weaker)**.** *With x not free in the matrix,* N *derives $D_{x,y}(y=1)$ and* U *does not.*

Proof. The rule $\mathrm{DI}_{\mathrm{root}}$ is the introduction by cases restricted to compound matrices together with the witnessed introduction by progress at literal equations. Hence U is the calculus of Theorem 2.13(ii) with transport in a formula template, and the statement is that proposition. Its interpretation is the truth of the mixed reading in the natural numbers, which Theorem 3.9 strengthens to provability in T_{eq}: the mixed reading $\exists x \forall y\,(y = 1 \to Sy = 1)$ of the formula fails at $y = 1$. □

Remark 3.11. `guarded_core` gives the example with the matrix $p(y) = 1$, where $p(y) := I(y,0;\, u.\,\mathrm{sel}(u,1,0))$ has the values $0,1,0$ at $0,1,2$: the introduction by cases has the base $0 = 1$, and the progressive reading fails between 1 and 2. The formula $D_{x,y}(y=1)$ is the one of its implementation section. The formula of the proposition shows two readings of one raw binder that disagree in the natural numbers, so no formula composite can identify the atomic binder of U with its ordinary reading; no impossibility for other interpretations of U is claimed.

Hence the single introduction is complete only together with a restriction: of the language, so that the two readings agree at every atomic matrix; or of the progress mode, to matrices at which they agree.

3.1.4 The shapes

Definition 3.12 (Guarded shapes)**.** Let $B(x,y)$ be a literal equation with the distinguished variables x and y. It is

(a) *zero-seeded in y* if it is literally $I(y,0;\, u.\,h) = I(y,0;\, u'.\,k)$;

(b) *mirrored in y* if it is literally $a = I(y,a;\, u.\,h)$, the left side and the seed being the same raw term a;

(c) *diagonal in y* if it is literally $I(y,y;\, u.\,h) = I(y,y;\, u'.\,k)$;

(d) an *orbit matrix in y* if it is literally $y = I(y,a;\, u.\,Su)$ with y not free in a.

The bodies h and k are arbitrary: they may contain x, y, their own state variable and further variables; in (b) the term a may contain x and y. The two state binders of (a) and (c) are independent. An equation $p = q$ is *protected* if p and q are iterator-headed. For an equation $E \equiv (s = t)$ and a term v, the *gate* and the *mirror gate* of E at v are

$$G_v E := \big(\mathrm{sel}(v,0,s) = \mathrm{sel}(v,0,t)\big), \qquad H_v E := \big(s = \mathrm{sel}(v,s,t)\big),$$

where $G_v E$ is $\mathrm{gate}(v,E)$ of Theorem 1.24 at an equation; the *wrapped* equation is $W(s) = W(t)$; and for an equation $E(y) \equiv (s(y) = t(y))$ the *diagonal lift* is

$$\Delta_y E := \big(I(y,y;\, u.\,s(y)) = I(y,y;\, u.\,t(y))\big), \qquad u \notin \mathrm{FV}(s,t).$$

For a variable y, the gate G_yE is zero-seeded in y, the mirror gate H_yE is mirrored in y and Δ_yE is diagonal in y, whether or not y occurs in E; in each the update does not use its state. Each shape is a condition on the syntax. None asks for an arithmetical equality or for the provability of a base case.

Lemma 3.13 (Instances at zero)**.** *Let $\mathsf{K}(\mathcal{F},\mathcal{S})$ be a guarded calculus whose sources include the iterator-headed terms, and let $B(x,y)$ be zero-seeded, mirrored or diagonal in y. Then $\Gamma \vdash B(t,0)$ for every context Γ and every term t, by I0 and transport in equation templates at iterator-headed sources.*

Proof. If B is zero-seeded or diagonal, both sides of $B(t,0)$ have the form $I(0,0;\,u.\,h')$; each is equal to 0 by I0, and Theorem 3.7(iv) joins them. If B is mirrored, $B(t,0)$ is $a' = I(0,a';\,u.\,h')$, the reverse of an instance of I0 (Theorem 3.7(iii)). □

An orbit matrix has no such property: its instance at zero is $0 = I(0,a;\,u.\,Su)$, which is interderivable with $0 = a$.

Lemma 3.14 (Cases at a matrix with a provable base)**.** *Let $B(x,y)$ be a literal equation such that $D_{x,y}B$ belongs to the proof language of a guarded calculus and $\Gamma \vdash B(t,0)$ for every Γ and t. If $\Gamma, B(t,0) \vdash B(t,Sz)$ with z fresh, then $\Gamma \vdash D_{x,y}B$.*

Proof. Cut the derivation of $B(t,0)$ into the premise: $\Gamma \vdash B(t,Sz)$. Weaken by $B(t,z)$ and apply $\mathrm{DI}_{\mathrm{root}}$ in progress mode. The variable z stays fresh because the base that was removed does not contain it. □

Proposition 3.15 (Substitution and the shapes)**.** *Let σ be a capture-free substitution of terms for variables other than the distinguished x and y, with x and y not free in its values.*

(i) *The root of a formula is preserved and reflected by σ: B is a literal equation if and only if $B\sigma$ is.*

(ii) *The zero-seeded and the mirrored shape are preserved by σ. Neither is reflected: for variables p and q the matrix $I(y,p;\,u.\,0) = I(y,0;\,u.\,0)$ is not zero-seeded and becomes zero-seeded under $p \mapsto 0$, and the matrix $p = I(y,q;\,u.\,0)$ is not mirrored and becomes mirrored under $q \mapsto p$.*

(iii) *The diagonal shape is preserved and reflected: $B\sigma$ is diagonal in y if and only if B is. The same holds for the substitution of a witness for x, if y is not free in the witness.*

(iv) *The classes of protected equations and of iterator-headed terms are closed under σ; the class of terms other than the literal 0 is not.*

Proof. (i), (ii) and (iv) are read off the definition of substitution. In (ii) the shapes are preserved because the literal 0, the variable y and the two equal occurrences of a are mapped to 0, to y and to two equal occurrences of $a\sigma$.

(iii) Rename the binders away from the values of σ. A constructor-headed term remains headed by the same constructor. A term $r\sigma$ is the distinguished variable y if and only if r is y: another variable that is kept differs from y; the value of a substituted variable does not contain y; and a constructor-headed term does not become a variable. Hence an iterator $I(n,a;\,u.\,h)\sigma$ has the pair (y,y) as counter and seed if and only if $I(n,a;\,u.\,h)$ has. A side of the equation that is a substituted variable may become iterator-headed, but its value has no free y, so it does not acquire the shape. A side that is an iterator has the shape after σ exactly when it has it before. The bodies are unrestricted and impose no condition. The substitution of a witness is the same argument with x among the substituted variables. □

The proposition is the reason for the two treatments of a guard. In a public language the progress mode is selected by the root, which is stable, and the shape is a requirement on the language, for which preservation suffices. On the full language the progress mode is selected by the shape itself, and a shape that substitution can create changes the mode of an inference under substitution (Section 3.5).

3.1.5 Public languages

Definition 3.16 (Public language and its proof language)**.** Let $\mathfrak{A}$ be a class of equations and $\mathfrak{M}$ a class of atomic matrices, both closed under substitution, with every instance $B(t, r)$ of a matrix $B \in \mathfrak{M}$ in $\mathfrak{A}$. The *public language* $\mathcal{P}(\mathfrak{A}, \mathfrak{M})$ is the least set of formulas such that

(1) every equation of $\mathfrak{A}$ is public;

(2) $D_{x,y}B$ is public if B is a compound public formula;

(3) $D_{x,y}B$ is public if $B(x, y)$ is a matrix of $\mathfrak{M}$ in y.

A *matrix of $\mathcal{P}$ in y* is a compound public formula or a matrix of $\mathfrak{M}$ in y. The *proof language* $\mathcal{P}^Q$ consists of the public formulas, all equations, and the *resources* $Q_y B(y)$ for the matrices B of $\mathcal{P}$ in y.

A public language has no Q. Clause (3) is the restriction: an atomic matrix immediately below $D_,$ has a prescribed literal shape, and no decision procedure and no proof is involved in recognizing it. Compound matrices are unrestricted within the grammar. By Theorem 3.15 the languages below are closed under substitution. The public languages of the chapter are

$$\begin{aligned}
\mathcal{P}_0 &:= \mathcal{P}(\text{all equations, zero-seeded}),\\
\mathcal{P}_0^+ &:= \mathcal{P}(\text{all equations, zero-seeded or orbit}),\\
\mathcal{P}_m &:= \mathcal{P}(\text{all equations, mirrored}),\\
\mathcal{P}_W &:= \mathcal{P}(\text{protected equations, zero-seeded}),
\end{aligned}$$

and the calculi are listed in Table 3.1. The designated contradiction of $\mathcal{P}_0$, $\mathcal{P}_0^+$ and $\mathcal{P}_m$ is $0 = 1$; that of $\mathcal{P}_W$ is $W(0) = W(1)$.

Calculus	Proof language	Sources	Package
G	$\mathcal{P}_0^Q$	all terms	`guarded_core`
U	$\mathcal{L}$	all terms	`guarded_core`
U^-	$\mathcal{L}$	$s \not\equiv 0$	`guarded_source`, `deterministic_auxiliary`
G^-	$\mathcal{P}_0^Q$	$s \not\equiv 0$	incomplete (Theorem 3.37)
$\mathsf{G}^-_{\mathrm{orb}}$	$(\mathcal{P}_0^+)^Q$	$s \not\equiv 0$	`guarded_orbit`
M	$\mathcal{P}_m^Q$	$s \not\equiv 0$	`mirror_guard`
H	$\mathcal{P}_W^Q$	iterator-headed	`protected_core`

Table 3.1: The guarded calculi $\mathsf{K}(\mathcal{F}, \mathcal{S})$ of Sections 3.1 to 3.4. Each has the seven rules of Theorem 3.2.

3.2 The guarded bootstrap and the translation into a public language

This section draws on `guarded_core` for the gate, the padding, the implication between equations and the translation; on `mirror_guard` for the form of the induction argument that

uses no substitution; on `guarded_orbit` and `protected_core` for the translation of Q; and on `deterministic_auxiliary` for the same translation. The argument is given once, for an arbitrary guard, and is instantiated in Sections 3.2.3 and 3.4.

3.2.1 Guard schemes

Definition 3.17 (Guard scheme)**.** Let $\mathsf{K} = \mathsf{K}(\mathcal{F}, \mathcal{S})$ be a guarded calculus and $\mathcal{P} = \mathcal{P}(\mathfrak{A}, \mathfrak{M})$ a public language with $\mathcal{P}^Q \subseteq \mathcal{F}$. A *guard scheme* for K and $\mathcal{P}$ on a class $\mathfrak{E} \subseteq \mathfrak{A}$ of equations, closed under substitution, consists of

- an equation $\gamma_v U \in \mathfrak{E}$ for every $U \in \mathfrak{E}$ and every term v, and
- a relation of *compatible pairs* on $\mathfrak{E}$, closed under substitution, with an equation $\mu_w(U, V) \in \mathfrak{E}$ for every compatible pair (U, V) and every term w,

both commuting with substitution, such that for all U, V, all terms w and all variables y:

(G1) $\gamma_y U$ is a matrix of $\mathfrak{M}$ in y, whether or not y occurs in U;

(G2) $\Gamma \vdash \gamma_0 U$ for every context Γ, and $\gamma_{Sw} U \dashv\vdash U$;

(G3) $\mu_0(U, V) \dashv\vdash U$ and $\mu_{Sw}(U, V) \dashv\vdash V$;

(G4) $(\gamma_y U,\, U)$ is a compatible pair.

Two schemes are used. The *zero-seeded scheme* has

$$\gamma_v(s = t) := G_v(s = t) = \big(\mathrm{sel}(v, 0, s) = \mathrm{sel}(v, 0, t)\big), \qquad \mu_w(s = t,\ r = q) := \big(\mathrm{sel}(w, s, r) = \mathrm{sel}(w, t, q)\big),$$

with every pair compatible. The *mirror scheme* has

$$\gamma_v(s = t) := H_v(s = t) = \big(s = \mathrm{sel}(v, s, t)\big), \qquad \mu_w(s = a,\ s = b) := \big(s = \mathrm{sel}(w, a, b)\big),$$

with (U, V) compatible when U and V have the same left side. In both, (G1) and (G4) hold by inspection: $G_y U$ is zero-seeded and $H_y U$ is mirrored in y, and $H_y(s = t)$ has the left side s. Conditions (G2) and (G3) are statements about a calculus and are verified where the scheme is used.

Fix a calculus, a public language and a guard scheme on $\mathfrak{E}$. For $U \in \mathfrak{E}$ and a compatible pair (U, V) put

$$L(U) := D_{a,v} \gamma_v U, \qquad (U \Rightarrow V) := D_{a,w} L\big(\mu_w(U, V)\big),$$

with fresh bound variables, a not occurring in the matrix. Both are compound public formulas, by (G1). For a formula A that is an equation of $\mathfrak{E}$ or a compound public formula, the *padding* is

$$P(A) := \begin{cases} L(A), & A \in \mathfrak{E}, \\ A, & A \text{ compound.} \end{cases}$$

$P(A)$ is always compound.

Lemma 3.18 (Padding)**.** *$L(U) \dashv\vdash U$ for every $U \in \mathfrak{E}$, and hence $P(A) \dashv\vdash A$.*

Proof. Let $\Gamma \vdash U$. The matrix $\gamma_v U$ is a literal equation, so the introduction of $L(U)$ is in progress mode; with the witness 0 its premise is $\Gamma, \gamma_z U \vdash \gamma_{Sz} U$, which follows from U by (G2) without the local assumption. Conversely let $\Gamma \vdash L(U)$. DE adds the resource $Q_v \gamma_v U$; its base $\gamma_0 U$ is derivable by (G2), QE at 1 gives $\gamma_1 U$, and (G2) gives U. The eigenvariable of DE does not occur in U. □

Lemma 3.19 (Numerical cases)**.** *Let $A(y)$ be a compound public formula or an equation of $\mathfrak{E}$, and let z be not free in Γ and A. If $\Gamma \vdash A(0)$ and $\Gamma \vdash A(Sz)$, then $\Gamma \vdash A(t)$ for every term t.*

Proof. Let A be compound. Introduce $D_{a,y}A(y)$, with a not free in A, in case mode with the witness 0: the premise $\Gamma, A(0) \vdash A(Sz)$ is the second hypothesis, weakened. DE adds $Q_y A(y)$, and QE with the first hypothesis gives $A(t)$. If A is an equation of $\mathfrak{E}$, pad the two hypotheses by Theorem 3.18, apply the compound case to $L(A(y))$, and remove the padding of $L(A(t))$. □

The context of the two hypotheses is the same Γ and does not contain the case variable. The lemma is not a rule for a case distinction on a variable that occurs in an assumption.

Lemma 3.20 (Implication between equations)**.** *For a compatible pair (U, V) the following rules are admissible:*

$$\frac{\Gamma,\ U \vdash V}{\Gamma \vdash U \Rightarrow V}, \qquad \frac{\Gamma \vdash U \Rightarrow V \qquad \Gamma \vdash U}{\Gamma \vdash V}.$$

Proof. Write μ_w for $\mu_w(U, V)$. The matrix $L(\mu_w)$ of $U \Rightarrow V$ is compound, so its introduction is in case mode, with the premise $\Gamma, L(\mu_0) \vdash L(\mu_{Sz})$. Over $\Gamma, L(\mu_0)$: Theorem 3.18 gives μ_0, (G3) gives U, the given derivation and cut give V, (G3) gives μ_{Sz}, and Theorem 3.18 gives $L(\mu_{Sz})$. For the elimination, DE adds $Q_w L(\mu_w)$; from U, (G3) and padding give $L(\mu_0)$; QE at 1 gives $L(\mu_1)$; and padding and (G3) give V. The conversions use transport outside the fresh binders only; no congruence of a binder is assumed. □

Theorem 3.21 (Equation induction from a guard)**.** *Let $E(y)$ be an equation of $\mathfrak{E}$ and z a variable not free in Γ and E. If $\Gamma \vdash E(0)$ and $\Gamma, E(z) \vdash E(Sz)$, then $\Gamma \vdash E(t)$ for every term t. The derivation uses the step premise at its own eigenvariable only: no substitution into the given derivations is made.*

Proof. Put

$$J(y) := \gamma_y E(y), \qquad \Phi(y) := \big(J(y) \Rightarrow E(y)\big),$$

which is defined by (G4) and is a compound public formula.

The implication at every term. $\Gamma \vdash \Phi(0)$ by the introduction rule of Theorem 3.20: the base $E(0)$ is derivable over Γ, and the assumption $J(0)$ is not used. For a fresh w, $\Gamma \vdash \Phi(Sw)$ by the same rule: the assumption $J(Sw)$ is $\gamma_{Sw}E(Sw)$, which gives $E(Sw)$ by (G2). Both derivations are over the context Γ, which does not contain the case variable, so Theorem 3.19 applies to the compound formula Φ and gives $\Gamma \vdash \Phi(v)$ for every term v.

Progress of the guard. Over $\Gamma, J(z)$ the elimination rule and $\Phi(z)$ give $E(z)$; the step premise and cut give $E(Sz)$; and (G2) gives $\gamma_{Sz}E(Sz)$, which is $J(Sz)$. Thus $\Gamma, J(z) \vdash J(Sz)$.

Conclusion. By (G1) the matrix $J(y)$ is atomic and admissible, so $\mathrm{DI}_{\mathrm{root}}$ in progress mode, with an unused witness coordinate a and the witness 0, gives $\Gamma \vdash D_{a,y}J(y)$. DE adds $Q_y J(y)$. The base $J(0) = \gamma_0 E(0)$ is derivable by (G2), and QE at t gives $J(t)$. With $\Phi(t)$ the elimination rule gives $E(t)$. □

Remark 3.22 (Two forms of the argument)*.* `guarded_core`, `guarded_orbit` and `protected_core` use the implication $J(y) \Rightarrow E(Sy)$ in place of Φ. Its two branches then need the step premise at 0 and at Sw, that is, a substitution into the given derivation, and the conclusion needs a second application of numerical cases. That form is correct wherever substitution is admissible: in G and H by Theorem 3.5, and in $\mathsf{G}^{-}_{\mathrm{orb}}$ after the unrestricted transport has been recovered, where `guarded_orbit` performs the substitution in the calculus with the derived rule as an inference and then expands that rule. The form of the proof above is the one of

`mirror_guard`, where it is needed because substitution is not available under the source restriction; it applies to every scheme. In each form the case distinction is made on a compound implication over a context that does not contain the case variable; a case distinction made while $J(z)$ is retained would violate the eigenvariable condition.

3.2.2 The translation and its two composites

Definition 3.23 (Translation into a public language)**.** Let a guard scheme on $\mathfrak{E}$ be given, and let ε map every equation to an equation of $\mathfrak{E}$, commuting with substitution. The translation T of native formulas is

$$\begin{aligned} T(s=t) &:= \varepsilon(s=t), \\ T(Q_y B) &:= D_{a,y} P(TB), \qquad a \text{ fresh}, \\ T(D_{x,y} B) &:= D_{x,y} P(TB). \end{aligned}$$

T leaves every term unchanged outside ε, commutes with substitution up to the names of bound variables, and maps every native formula to a public formula: every translated matrix is padded, hence compound, and the atomic matrices inside a padding are admissible by (G1). The embedding ε is the identity except for protected atoms, where it is the wrapping $\varepsilon(s=t) = (W(s) = W(t))$.

Theorem 3.24 (The public-language theorem)**.** *Let $\mathsf{K} = \mathsf{K}(\mathcal{F}, \mathcal{S})$, $\mathcal{P} = \mathcal{P}(\mathfrak{A}, \mathfrak{M})$ with $\mathcal{P}^Q \subseteq \mathcal{F}$, a guard scheme on $\mathfrak{E}$ and an embedding ε be given, and assume:*

(T1) *$\varepsilon(E) \dashv\vdash_{\mathsf{N}} E$ for every equation E;*

(T2) *K derives $\varepsilon(E)$ for every instance E of I0 and IS, and $\Gamma \vdash \varepsilon(s=t)$ and $\Gamma \vdash \varepsilon(h(s)=r)$ imply $\Gamma \vdash \varepsilon(r = h(t))$;*

(T3) *$U \dashv\vdash_{\mathsf{K}} \varepsilon(U)$ for every equation $U \in \mathfrak{A}$;*

(T4) *$\Gamma \vdash_{\mathsf{K}} B(t,0)$ for every matrix $B \in \mathfrak{M}$, every Γ and every t.*

Then:

(a) *$\Gamma \vdash_{\mathsf{N}} A$ implies $T\Gamma \vdash_{\mathsf{K}} TA$, for all native Γ and A;*

(b) *$A \dashv\vdash_{\mathsf{N}} TA$ for every native formula A;*

(c) *$B \dashv\vdash_{\mathsf{K}} TB$ for every public formula B;*

(d) *for public Γ and B: $\Gamma \vdash_{\mathsf{K}} B$ if and only if $\Gamma \vdash_{\mathsf{N}} B$.*

The transformations of derivations are effective.

Proof. For a native matrix B, the translated closure is $\mathrm{Cl}_y B := D_{a,y} P(TB)$, so $T(Q_y B) = \mathrm{Cl}_y B$.

(a). By induction on the derivation in N.

Hyp: $TA \in T\Gamma$.

I0, IS, Tr: these are (T2); the rule Tr of N infers $r = h(t)$ from $s = t$ and $h(s) = r$.

QE: the induction hypotheses are $T\Gamma \vdash \mathrm{Cl}_y B$ and $T\Gamma \vdash TB(0)$. DE on the first adds the resource $Q_y P(TB)(y)$. Padding gives $P(TB)(0)$, QE at t gives $P(TB)(t)$, and removal of the padding gives $TB(t) = T(B(t))$.

DE: the induction hypotheses are $T\Gamma \vdash D_{x,y} P(TB)$ and $T\Gamma, \mathrm{Cl}_y B(b,y) \vdash TA$. DE in K on the first adds $Q_y P(TB)(b,y)$. Over this context derive $\mathrm{Cl}_y B(b,y)$: its matrix is compound, the introduction is in case mode with the witness 0, and its premise $P(TB)(b,0) \vdash P(TB)(b, Sz)$

is QE on the resource and the local base. Cut $\mathrm{Cl}_y B(b,y)$ into the second hypothesis, which gives TA, and close DE; b is not free in $T\Gamma$ and TA. This treats every use of the hypothesis $Q_y B(b,y)$ of the native derivation, not only its uses as the major premise of QE.

DI: the induction hypothesis is $T\Gamma, TB(t,0) \vdash TB(t,Sz)$. The matrix $P(TB)$ is compound, so the introduction of $D_{x,y}P(TB)$ is in case mode, with the premise $T\Gamma, P(TB)(t,0) \vdash P(TB)(t,Sz)$: remove the padding of the base, cut it into the induction hypothesis, and pad the result.

QI: the induction hypothesis is $T\Gamma, \varepsilon E(z) \vdash \varepsilon E(Sz)$, and the goal is $D_{a,y}L(\varepsilon E(y))$. Its introduction is in case mode, with the premise $T\Gamma, L(\varepsilon E(0)) \vdash L(\varepsilon E(Sz'))$ for a fresh z'. Over $T\Gamma, L(\varepsilon E(0))$ the padding gives the base $\varepsilon E(0)$; Theorem 3.21 for the equation $\varepsilon E(y) \in \mathfrak{E}$, with the step premise weakened, gives $\varepsilon E(Sz')$; and padding gives the goal.

(b). By induction on A, using three facts about N: $Q_y B \dashv\vdash D_{a,y}B$ for a not free in B (Theorem 1.12); interderivable matrices give interderivable formulas under Q and under D, (Theorem 1.21); and $L(U) \dashv\vdash_{\mathsf{N}} U$. The third holds because (G2) holds in N by Theorem 3.6 and cut, and the proof of Theorem 3.18 then applies in N, where the introduction by cases is available at every matrix. The case of an equation is (T1). For $Q_y B$: $Q_y B \dashv\vdash D_{a,y}B \dashv\vdash D_{a,y}TB \dashv\vdash D_{a,y}P(TB)$. The case of $D_{x,y}B$ is the same without the first step.

(c). By induction on the number of binders of B, for all instances of B at once. For an equation of $\mathfrak{A}$ this is (T3).

Let $B = D_{x,y}C$ with C compound. Then TC is compound and $TB = D_{x,y}TC$. From $\Gamma \vdash D_{x,y}C$, DE adds $Q_y C(b,y)$; the introduction of $D_{x,y}TC$ at the witness b is in case mode, and its premise $TC(b,0) \vdash TC(b,Sz)$ is obtained by converting the base to $C(b,0)$ with the induction hypothesis, applying QE at Sz, and converting $C(b,Sz)$ to $TC(b,Sz)$. The converse is symmetric.

Let $B = D_{x,y}C$ with $C \in \mathfrak{M}$. Then $TB = D_{x,y}L(\varepsilon C)$, and every instance of C is an equation of $\mathfrak{A}$. From $\Gamma \vdash D_{x,y}C$, DE adds $Q_y C(b,y)$. The introduction of $D_{x,y}L(\varepsilon C)$ at the witness b is in case mode; its local base is not used: $C(b,0)$ is derivable by (T4), QE gives $C(b,Sz)$, (T3) gives $\varepsilon C(b,Sz)$, and padding gives $L(\varepsilon C(b,Sz))$. Conversely, from $\Gamma \vdash D_{x,y}L(\varepsilon C)$, DE adds $Q_y L(\varepsilon C(b,y))$. The introduction of $D_{x,y}C$ at the witness b is in progress mode; its local assumption $C(b,z)$ is not used: $C(b,0)$ is derivable by (T4), (T3) and padding give $L(\varepsilon C(b,0))$, QE gives $L(\varepsilon C(b,Sz))$, and removal of the padding and (T3) give $C(b,Sz)$. This direction uses the provable base; it fails for an arbitrary atomic matrix, by Theorem 3.10.

(d). The implication from left to right is Theorem 3.6. Conversely let $\Gamma \vdash_{\mathsf{N}} B$. By (a), $T\Gamma \vdash_{\mathsf{K}} TB$. By (c), $\Gamma \vdash_{\mathsf{K}} TC$ for every $C \in \Gamma$, so cut gives $\Gamma \vdash_{\mathsf{K}} TB$, and (c) gives $\Gamma \vdash_{\mathsf{K}} B$. □

Corollary 3.25 (Bi-interpretation)**.** *Under the hypotheses of Theorem 3.24, the theory of the public sequents of* K*, with the ordinary reading* $(D_{x,y}B)^\circ = \exists x\,(B^\circ(x,0) \to \forall y\, B^\circ(x,y))$*, is bi-interpretable with* iPRA*. Both numerical domains are whole, equality is equality, both numerical composites have the graph* $x = y$*, and both formula composites are provably equivalent to the identity. In particular* K *derives its contradiction* $\varepsilon(0=1)$ *if and only if* iPRA *proves* $\bot$*.*

Proof. The inclusion of the public language into the native language and the translation T are the identity on the numerical sort and on equality; if ε is the wrapping, the identity graph is expressed by $W(x) = W(y)$, which is $x = y$ in N by (T1). They preserve derivability by Theorem 3.6 and by (a), and their two composites are (b) and (c). Composition with the bi-interpretation of N, T_{eq} and iPRA (Theorem 1.78) gives the statement. For the contradiction: $\varepsilon(0=1)$ is public, so by (d) it is derivable in K if and only if it is derivable in N, that is, by (T1), if and only if N derives $0 = 1$. □

The arithmetic of the bridge, that is, the predecessor, the reflection of equality, quantifier-free induction, pairing and the compiler for primitive recursion (Theorems 1.48, 1.52 and 1.62), is not redone in any public language: it is transferred by (a) and (d).

3.2.3 The guarded core

This subsection draws on `guarded_core`. The guarded core is $\mathsf{G} = \mathsf{K}(\mathcal{P}_0^Q, \text{all terms})$: public formulas are equations and paired formulas whose atomic matrices are zero-seeded, transport is unrestricted, and the one introduction is determined by the root. Its seven rules are those of Theorem 3.2.

Theorem 3.26 (The guarded core)**.** (i) G *admits weakening, substitution and cut.*

(ii) *The zero-seeded scheme on all equations is a guard scheme for* G *and* $\mathcal{P}_0$*. Hence* G *admits numerical cases for every public formula, implication between equations, and contextual induction for every literal equation.*

(iii) *With* ε *the identity,* T *translates derivations of* N *into derivations of* G*;* $A \dashv\vdash_{\mathsf{N}} TA$ *for every native* A*;* $B \dashv\vdash_{\mathsf{G}} TB$ *for every* $B \in \mathcal{P}_0$*; and for* Γ, B *in* $\mathcal{P}_0$*,* $\Gamma \vdash_{\mathsf{G}} B$ *if and only if* $\Gamma \vdash_{\mathsf{N}} B$*.*

(iv) G *on* $\mathcal{P}_0$ *is bi-interpretable and equiconsistent with* iPRA*, with whole domains, equality as equality and identity numerical composites.*

Proof. (i) is Theorem 3.5. For (ii), all terms are sources, so Theorem 3.7(i),(ii) give the ordinary equality rules. The two sides of G_0U are equal to 0 and the two sides of $G_{Sw}(s = t)$ are equal to s and to t, by the selector equations; this is (G2). The two sides of $\mu_0(s = t, r = q)$ are equal to s and t, and those of μ_{Sw} to r and q; this is (G3). The consequences are Theorems 3.19 to 3.21. For (iii) and (iv) the hypotheses of Theorem 3.24 hold: (T1) and (T3) are trivial; in (T2) the computation equations are axioms and $r = h(t)$ follows from $s = t$ and $h(s) = r$ by transport in the template $h(v) = r$ and symmetry; (T4) is Theorem 3.13. Then apply Theorems 3.24 and 3.25. □

Corollary 3.27 (Administrative proofs are conservative)**.** *For* Γ *and* B *in* $\mathcal{P}_0$*,*

$$\Gamma \vdash_{\mathsf{U}} B \iff \Gamma \vdash_{\mathsf{G}} B \iff \Gamma \vdash_{\mathsf{N}} B.$$

On $\mathcal{P}_0$ *the mixed and the ordinary reading agree:* T_{eq} *proves* $A^\dagger \leftrightarrow A^\circ$ *for every* $A \in \mathcal{P}_0$*.*

Proof. A derivation of G is a derivation of U, a derivation of U is transformed into one of N by Theorem 3.6, and Theorem 3.26(iii) closes the circle. For the readings, at a zero-seeded matrix $E(x, y)$ the instance $E(x, 0)$ is provable, so with x fixed $\forall y\,(E(x, y) \to E(x, Sy))$ and $E(x, 0) \to \forall y\, E(x, y)$ are equivalent: from the first, by induction for the literal equation $E(x, y)$; from the second, because the base gives every instance. The equivalence passes under the existential quantifier, and at a compound matrix the two readings have the same form, so induction on A applies. □

A derivation may therefore use unguarded atomic matrices and Q formulas as intermediate formulas without changing the public theory. The converse statement fails for U as a whole (Theorem 3.10): the full language of U is not claimed to be bi-interpretable with iPRA under either reading.

Remark 3.28 (Versions)*.* `guarded_core` states the comparison with the calculus of the reports on the language of equations and D, with clause contexts (Theorem 3.4); the clause $T(Q_y B) = D_{a,y} P(TB)$ is added in `guarded_orbit`, `deterministic_auxiliary` and `protected_core`,

and `guarded_source` first replaces every Q_yB by $D_{a,y}B$. `guarded_core` derives conservativity of U from Theorem 3.9 and the agreement of the readings, through iPRA; the direct inclusion of Theorem 3.6 is the argument of `guarded_orbit` and `protected_core`. In Lean the aim of the package, one root-determined introduction, is reached on the full language by the witness-selected introduction (Theorem 2.17, `wprf_iff_nprf`); its own route is formalized in the common form described in Theorem 3.47.

3.3 Which transport sources are needed

This section draws on `guarded_source` for the dependency theorem; on `deterministic_auxiliary`, `guarded_orbit` and `mirror_guard`, each of which reproduces the same numerical argument inside its own calculus; and on `guarded_orbit` for the observation that the argument needs induction at one equation only. The singleton test, the filter and the join at zero are taken by all four from `source_dispatch`; they are proved in Section 2.3.4 for the calculus Z, which has both introductions of N, and are cited from there. This section proves what the guarded calculi need beyond that: the argument goes through with two closure conditions in place of the introductions.

Transport with the sources $s \not\equiv 0$ is as strong as unrestricted transport, on equations, as soon as a small amount of induction is available. The amount is made explicit, because in Section 3.4 it is exactly what a strictly guarded language fails to supply.

Definition 3.29 (Restricted equational kernel)**.** A *restricted equational kernel* is a relation $\Gamma \vdash A$ between finite contexts and formulas of a language that contains all equations, closed under weakening and under the rules Hyp for equations, I0, IS and $\textsc{Tr}^{-}$ in equation templates: from $\Gamma \vdash s = t$ with $s \not\equiv 0$ and $\Gamma \vdash E(s)$ infer $\Gamma \vdash E(t)$, for a literal equation template $E(v)$.

The derivability relation of every guarded calculus whose sources include the terms $s \not\equiv 0$ is a restricted equational kernel, and Theorem 3.7 holds for every restricted equational kernel, since its proof uses these rules only. Consider the following two closure conditions on a restricted equational kernel.

(C) *Cases at stable left sides.* If $E(v)$ is an equation template whose left side is iterator-headed, z is a variable not in E, and $\vdash E(0)$ and $\vdash E(Sz)$ in the empty context, then $\vdash E(r)$ for every term r.

(O) *One step of an orbit.* For every term $t \not\equiv 0$ and every context Γ, if $\Gamma \vdash 0 = I(0, t;\, u.\, Su)$, then $\Gamma \vdash 1 = I(1, t;\, u.\, Su)$.

Define the terms

$$\begin{aligned} \zeta(v) &:= \mathrm{sel}(v, 1, 0), & \kappa(v) &:= \mathrm{sel}(v, 0, 1), & \chi(v) &:= \mathrm{sel}(v, 1, 2), \\ N(n) &:= I(n, 0;\, u.\, \zeta(u)), & R(n) &:= I(n, 0;\, u.\, \chi(u)), & \lambda(n) &:= N(R(n)). \end{aligned}$$

These are the terms Z, K, H, N, R and L of Section 2.3.4; three are renamed because H_v and $L(\cdot)$ denote the mirror gate and the padding in this chapter. The term ζ is the zero test, and $\lambda(Sv)$ tests whether v is 0.

Lemma 3.30 (Tests and filter)**.** *In a restricted equational kernel with* (C)*, the following equations are derivable over every context, for every term v:*

$$\chi(v) = S\kappa(v), \qquad R(1) = 1, \qquad R(SSv) = 2, \qquad N(1) = 1, \qquad N(2) = 0,$$
$$\lambda(1) = 1, \qquad \lambda(SSv) = 0, \qquad \lambda(Sv) = \zeta(v), \qquad \mathrm{sel}(\zeta(v), v, 0) = v.$$

Proof. The first eight equations are Theorem 2.38. The derivations given there are made over the empty context and use I0, IS, the operations of Theorem 3.7(i)–(iv) at sources that are iterator-headed, successor-headed or the numerals 1 and 2, and numerical cases at the two matrices $\chi(y) = S\kappa(y)$ and $\lambda(Sy) = \zeta(y)$. Both matrices have an iterator-headed left side, so the two case distinctions are instances of (C), and the derivations are valid in every restricted equational kernel with (C).

The filter of Theorem 2.39 is the equation $v = \mathrm{sel}(\zeta(v), v, 0)$, proved by cases at a matrix whose left side is a variable; (C) does not cover it. In the orientation stated here the left side is iterator-headed and (C) applies. At 0, replace $\zeta(0)$ by 1; then $\mathrm{sel}(1, 0, 0) = 0$ by IS. At Sz, replace $\zeta(Sz)$ by 0; then $\mathrm{sel}(0, Sz, 0) = Sz$ by I0. Both replacements are congruences at an iterator-headed source. □

Theorem 3.31 (Equational source elimination)**.** *Let $\vdash$ be a restricted equational kernel with* (C) *and* (O)*. Then $\vdash$ has unrestricted equality: over every context, symmetry, transitivity and congruence hold for all equations, and $\Gamma \vdash s = t$ and $\Gamma \vdash E(s)$ imply $\Gamma \vdash E(t)$ for every equation template $E(v)$ and every source s. Both* (C) *and* (O) *hold if $\vdash$ is closed under contextual induction for literal equations: from $\Gamma \vdash E(0)$ and $\Gamma, E(z) \vdash E(Sz)$, with z fresh, infer $\Gamma \vdash E(r)$.*

Proof. From induction to (C) *and* (O). For (C), weaken the derivation of $E(Sz)$ by the assumption $E(z)$ and apply induction; the induction hypothesis is not used. For (O), write $A(n) := I(n, t;\, u.\, Su)$ and apply induction to the equation $n = A(n)$ at the target 1: the base is the hypothesis of (O), and the step $\Gamma, z = A(z) \vdash Sz = A(Sz)$ is the step of the proof of Theorem 2.40, whose transports have the sources z, $A(Sz)$ and $SA(z)$.

Symmetry at the source 0. Let $\Gamma \vdash 0 = t$ with $t \not\equiv 0$; the claim is $\Gamma \vdash t = 0$. Read the proof of Theorem 2.40 with the given derivation of $0 = t$ in place of the hypothesis. It derives $0 = A(0)$ from $0 = t$, uses its induction for the single conclusion $1 = A(1)$, which is (O), and derives $\zeta(t) = 1$ from $1 = A(1)$ by IS and Theorem 3.30. With the filter in the orientation of Theorem 3.30 the last step reads: put $q := \mathrm{sel}(\zeta(t), t, 0)$; the filter gives $q = t$; the replacement of $\zeta(t)$ by 1 and IS give $q = \mathrm{sel}(1, t, 0) = 0$; reverse $q = t$ at the source q and pass through q to obtain $t = 0$.

Unrestricted equality. Symmetry, congruence and transitivity at every source follow from symmetry at the source 0 and Theorem 3.7(ii),(iv) as in the proof of Theorem 2.35, which uses nothing else; the cut made there is not needed, because symmetry at the source 0 has been obtained as a derived rule. For an equation template $a(v) = b(v)$, congruence gives $a(s) = a(t)$ and $b(s) = b(t)$, and symmetry and transitivity turn $a(s) = b(s)$ into $a(t) = b(t)$. □

The proof uses induction in two places only: through (C), where the induction hypothesis is ignored, and through (O), at the single equation $n = I(n, t;\, u.\, Su)$ with t and the context fixed. The variable source z of the step of (O) is never replaced by 0: the target 1 is supplied by the induction rule, not by a substitution into the step. No logical rule occurs: no binder, no introduction by cases. This removes the apparent circularity between the two reductions, in which the recovery of transport at the source 0 would need logical cases and the recovery of logical cases at an equation would need unrestricted transport.

Corollary 3.32 (The equational fragment)**.** *Let $\mathcal{E}^-$ be the calculus whose formulas are the equations and whose rules are* Hyp*, I0, IS, Tr$^-$ in equation templates and contextual induction for literal equations. Unrestricted transport in equation templates is admissible in $\mathcal{E}^-$. The restriction $s \not\equiv 0$ changes neither its theorems nor its consequences from hypotheses.*

Theorem 2.35 is the case of Theorem 3.31 for the calculus Z, which has QI and hence induction; what Theorem 3.31 adds is that the two closure conditions suffice, which is what

the calculi of Section 3.4 need. The Lean statements for the fragment, `ZKernel.trF` and `zprf_iff_nprf`, are recorded in Theorem 2.41.

Proposition 3.33 (Transport in public templates)**.** *Let* K *be a guarded calculus whose proof language contains* $\mathcal{P}^Q$ *for a public language* $\mathcal{P} = \mathcal{P}(\mathfrak{A}, \mathfrak{M})$*. Assume that* K *has unrestricted equality in the sense of Theorem 3.31, and that for every* $B \in \mathfrak{M}$ *either* $\Gamma \vdash B(t, 0)$ *for all* Γ *and* t*, or* $\Gamma \vdash D_{x,y}B'$ *for all* Γ *and all substitution instances* B' *of* B*. Then for every template* $F(v) \in \mathcal{P}$*,*

$$\Gamma \vdash s = t \quad \textit{and} \quad \Gamma \vdash F(s) \quad \textit{imply} \quad \Gamma \vdash F(t).$$

Proof. By induction on the number of binders of F, for all contexts and all s, t. For an equation this is the hypothesis. Let $F(v) = D_{x,y}B(v, x, y)$. If every instance of $D_{x,y}B$ is derivable, nothing is to be done. Otherwise apply DE to $F(s)$, which adds $Q_yB(s, b, y)$, and introduce $F(t)$ at the witness b.

If B is compound the introduction is in case mode. From its base $B(t, b, 0)$ the induction hypothesis, with the reversed equation $t = s$, gives $B(s, b, 0)$; QE gives $B(s, b, Sz)$; and the induction hypothesis gives $B(t, b, Sz)$.

If B is atomic the introduction is in progress mode and its local assumption is not used: $B(s, b, 0)$ is derivable, QE gives $B(s, b, Sz)$, and transport in the equation template gives $B(t, b, Sz)$. □

Hence, under the hypotheses of the proposition, a calculus $\mathsf{K}(\mathcal{P}^Q,\ s \not\equiv 0)$ derives the same sequents as $\mathsf{K}(\mathcal{P}^Q$, all terms): every unrestricted transport is replaced by its derivation and the other inferences are kept. The proposition is about public templates. No transport at the source 0 is asserted in a template that contains an unguarded atomic matrix below a binder.

Remark 3.34 (The crossed rule)*.* The induction of Theorem 3.31 is the price of the uncrossed conclusion. For the rule Tr of the kernel the restriction $s \not\equiv 0$ costs nothing: Theorem 2.25(ii) reverses an equation with the source 0 in one inference, by a constant template, and needs neither the tests nor an induction; the same holds for the reversing rule of Theorem 2.32. The calculi of Sections 3.1 to 3.4 have the uncrossed rule, and for it the obstruction of Theorem 3.36 is real.

Remark 3.35 (Substitution under a source restriction)*.* As for the calculi of Theorem 2.28, substitution is admissible in a source-restricted guarded calculus once its equivalence with the unrestricted calculus is known, and what fails is the rule-by-rule substitution of Theorem 3.5; `guarded_source`, `guarded_orbit` and `protected_core` make the same point. The equivalence must be proved first, and none of the proofs of this chapter substitutes into a derivation with Tr^- before that point.

3.4 The failure of composition and its repairs

This section draws on `mirror_guard`, `guarded_orbit` and `deterministic_auxiliary` for the obstruction; on `guarded_source` and `deterministic_auxiliary` for the repair by auxiliary formulas; on `guarded_orbit` for the repair by the orbit; on `mirror_guard` for the mirror guard; and on `protected_core` for protected atoms.

The guarded core removes the choice of the introduction mode (Theorem 3.26), and the source restriction removes the transports at the literal 0 (Theorem 3.31). The first result is stated for unrestricted transport and the second for a calculus with induction at arbitrary equations. In the strictly guarded calculus $\mathsf{G}^- = \mathsf{K}(\mathcal{P}_0^Q,\ s \not\equiv 0)$ neither hypothesis is available, and the two results do not compose.

3.4.1 The obstruction

An equation is *zero-left* if its left side is the literal 0. A formula of a proof language $\mathcal{P}^Q$ is *safe* if it is not a zero-left equation: it is a public compound formula, a resource, or an equation whose left side is not the literal 0.

Proposition 3.36 (An inert zero-left hypothesis)**.** *Let $\mathcal{P} = \mathcal{P}(\text{all equations}, \mathfrak{M})$, where the left side of every matrix of $\mathfrak{M}$ is iterator-headed or successor-headed, and let $\mathsf{K} = \mathsf{K}(\mathcal{P}^Q,\ s \not\equiv 0)$. If Γ consists of safe formulas, F is safe and $\Gamma,\ 0 = t \vdash_{\mathsf{K}} F$, then $\Gamma \vdash_{\mathsf{K}} F$.*

Proof. By induction on the derivation, for all safe Γ and F. Every instance of a matrix of $\mathcal{P}$ is safe: a compound formula stays compound, and the left side of an instance of a matrix of $\mathfrak{M}$ keeps its head.

Hyp: the conclusion is safe, so it is not the hypothesis $0 = t$ and belongs to Γ. I0, IS: no hypothesis is used.

Tr$^-$, with the premises $p = q$ and $F(p)$ and the safe conclusion $F(q)$: the major premise is safe because $p \not\equiv 0$. The minor premise is safe: if F is compound this is clear; if $F(v)$ is $a(v) = b(v)$ and $a(p) \equiv 0$, then $a(v)$ is the constant 0, because a constructor-headed term and the variable v do not become the literal 0 when v is replaced by $p \not\equiv 0$; then $a(q) \equiv 0$, against the safety of $F(q)$. The induction hypothesis applies to both premises.

QE, with the premises $Q_y B(y)$ and $B(0)$: both are safe, the first as a resource and the second as an instance of a matrix.

DE: the major premise is compound; the minor premise has the same conclusion and one more hypothesis, the resource $Q_y B(b, y)$, which is safe.

$\mathrm{DI}_{\mathrm{root}}$: the local assumption $B(t', \eta_B(z))$ and the conclusion $B(t', Sz)$ of the premise are instances of a matrix, hence safe. □

Corollary 3.37 (The strictly guarded composition is incomplete)**.** *In G^- the sequents $0 = 1 \vdash 1 = 0$ and $0 = x \vdash x = 0$, for a variable x, are not derivable. Both are derivable in G. The same holds for every calculus of Theorem 3.36, for instance with the atomic matrices $\mathrm{sel}(y, t, s) = t$.*

Proof. The conclusions $1 = 0$ and $x = 0$ are safe. By Theorem 3.36 a derivation would give $\vdash 1 = 0$, respectively $\vdash x = 0$, in the empty context, hence in N by Theorem 3.6; both fail in the natural numbers. In G symmetry is Theorem 3.7(ii). □

The proposition says where the composition breaks. Every equation that a strictly guarded derivation can obtain from a binder has a stable left side: the instances of a zero-seeded matrix, the local assumptions and the premises of the introductions. A transport with a source $s \not\equiv 0$ cannot turn a zero-left equation into a safe one. A zero-left hypothesis therefore has no path to a safe conclusion. The proposition does not apply to a context with a clause whose antecedent is zero-left and whose consequent is safe, and each of the repairs below supplies such a clause or removes the zero-left hypotheses from the public language.

Remark 3.38 (Versions of the obstruction). Theorem 3.36 is stated in `mirror_guard`, for every guard with a constructor-headed left side, and in `guarded_orbit` for the hypothesis $0 = x$. `deterministic_auxiliary` proves the first sequent of Theorem 3.37 underivable by a counterinterpretation over the natural numbers: every zero-left equation is read as true, every other equation as equality, and the compound formulas by their ordinary reading. Every equation below a binder of a formula of $\mathcal{P}_0^Q$ has an iterator-headed left side, so it is read as equality and the reading commutes with substitution; the computation equations, both modes of the introduction and QE and DE are sound; and Tr$^-$ is sound because its major premise $p = q$ is not zero-left and an equation template whose instance at p is zero-left has the constant left side 0, so that its instance at q is true by convention. The hypothesis $0 = 1$

is true and the conclusion $1 = 0$ is false. In Lean the calculus `PGPrf` with the root-determined introduction and transport at iterator-headed sources does not derive $x = z$ from $x = y$ and $y = z$ between variables (`pgprf_not_trans`), by a reading in which a value is a number with a colour and equality is not transitive; hence it is strictly weaker than the kernel on arbitrary sequents (`pgprf_strictly_weaker`).

3.4.2 First repair: auxiliary formulas

This subsection draws on `guarded_source` and `deterministic_auxiliary`. The public language remains $\mathcal{P}_0$, and derivations are made in $\mathsf{U}^- = \mathsf{K}(\mathcal{L},\, s \not\equiv 0)$: the proof language is the full native language, in which an atomic matrix below $D_,$ need not be guarded. A public theorem is a public formula with a derivation in U^-; its intermediate formulas need not be public. This convention is part of the calculus. By Theorem 3.3 its rules are Hyp, the two computation equations, SE, DE and $\mathrm{DI}_{\mathrm{root}}$.

Theorem 3.39 (Source selection and root-determined introduction, with auxiliary formulas). (i) U^- *admits contextual induction for every literal equation, by a derivation without transport, and numerical cases for every native formula.*

(ii) U^- *has unrestricted equality and admits transport, at every source, in every template of* $\mathcal{P}_0$.

(iii) *With the zero-seeded scheme and* ε *the identity,* T *translates derivations of* N *into derivations of* U^-, *and* $B \dashv\vdash_{\mathsf{U}^-} TB$ *for every* $B \in \mathcal{P}_0$.

(iv) *For* Γ *and* B *in* $\mathcal{P}_0$,

$$\Gamma \vdash_{\mathsf{U}^-} B \iff \Gamma \vdash_{\mathsf{G}} B \iff \Gamma \vdash_{\mathsf{N}} B,$$

and U^- *on* $\mathcal{P}_0$ *is bi-interpretable and equiconsistent with* iPRA, *with whole domains, equality as equality and identity numerical composites.*

Proof. (i) Let $\Gamma \vdash E(0)$ and $\Gamma, E(z) \vdash E(Sz)$. The formula $D_{a,y}E(y)$, with a not free in E, belongs to $\mathcal{L}$, and its introduction is in progress mode: with the witness 0 its premise is the given step. DE adds $Q_y E(y)$, in which the eigenvariable does not occur, and QE with the base gives $E(t)$. For numerical cases at an equation, use this induction and ignore the induction hypothesis; at a compound formula $A(y)$, introduce $D_{a,y}A(y)$ in case mode and proceed in the same way.

(ii) By (i) and Theorem 3.31, U^- has unrestricted equality; additional hypotheses of the context are passive in that proof. Theorem 3.33 with Theorem 3.13 gives the transport in public templates.

(iii) and (iv). With unrestricted equality the zero-seeded scheme satisfies (G2) and (G3) in U^- exactly as in Theorem 3.26, and (T1) to (T4) hold for the same reasons. Apply Theorems 3.24 and 3.25; the middle member of the chain is Theorem 3.26(iii). □

The induction of (i) is made at an arbitrary literal equation, and the matrix E is in general not guarded. The proof of Theorem 3.31 uses it at the equation $n = I(n, t;\, u.\, Su)$, whose left side is a variable. By Theorem 3.37 this use of an unguarded matrix cannot be replaced by guarded ones: the auxiliary formulas are a necessary part of the calculus, and they are why the proof language is stated explicitly. The theorem asserts the equivalence and the composites for the public language. It asserts neither a formula composite nor transport at the source 0 for all auxiliary formulas; by Theorem 3.10 the former fails.

Remark 3.40 (Versions). `guarded_source` proves Theorem 3.31 for the fragment $\mathcal{E}^-$ and instantiates it in U^-, with the filter in the orientation $v = \mathrm{sel}(\zeta(v), v, 0)$; `deterministic_auxiliary`

runs the numerical argument directly in U^-, with the filter as in Theorem 3.30. Both prove the implication from U^- to G in (iv) through iPRA, by Theorem 3.9, the agreement of the readings on $\mathcal{P}_0$ (Theorem 3.27) and the faithfulness of the bridge; Theorem 3.6 gives it directly. In Lean the combination of source selection with a single introduction is proved on the full native language, with the witness-selected introduction and the crossed rule (the seven-rule calculus of Theorem 2.24, `mprf_iff_nprf`); the auxiliary formulas of the two packages are not formalized.

3.4.3 Second repair: the administrative orbit

This subsection draws on `guarded_orbit`. The auxiliary formulas of the first repair are reduced to one family. The proof language is $(\mathcal{P}_0^+)^Q$: the guarded language together with the paired formulas over orbit matrices $y = I(y, a;\, u.\, Su)$, y not free in a. The calculus is $\mathsf{G}^-_{\mathrm{orb}} = \mathsf{K}((\mathcal{P}_0^+)^Q,\, s \not\equiv 0)$, with the rules of Theorem 3.2; no rule is added for the orbit matrices. The public language remains $\mathcal{P}_0$.

An equation is *stable-left* if its left side is iterator-headed or successor-headed. The class of stable-left equations is closed under substitution, and none of its members is zero-left.

Theorem 3.41 (The orbit bootstrap). (i) *In G^-, hence in $\mathsf{G}^-_{\mathrm{orb}}$, the zero-seeded scheme on the stable-left equations is a guard scheme for $\mathcal{P}_0$. Hence numerical cases hold for every compound public formula and every stable-left equation, and condition* (C) *of Section 3.3 holds.*

(ii) *For every term t with y not free in it and every context, $\mathsf{G}^-_{\mathrm{orb}}$ derives the* orbit closure

$$D_{a,y}(y = I(y, t;\, u.\, Su)),$$

and condition (O) *holds.*

(iii) *$\mathsf{G}^-_{\mathrm{orb}}$ has unrestricted equality, admits transport at every source in every template of $\mathcal{P}_0^+$, and derives the same sequents as $\mathsf{K}((\mathcal{P}_0^+)^Q$, all terms).*

(iv) *With the zero-seeded scheme on all equations and ε the identity, T translates derivations of N into derivations of $\mathsf{G}^-_{\mathrm{orb}}$; $B \dashv\vdash TB$ for every $B \in \mathcal{P}_0$; and for Γ and B in $\mathcal{P}_0$,*

$$\Gamma \vdash_{\mathsf{G}^-_{\mathrm{orb}}} B \iff \Gamma \vdash_{\mathsf{G}} B \iff \Gamma \vdash_{\mathsf{N}} B.$$

The public theory is bi-interpretable and equiconsistent with iPRA, with whole domains, equality as equality and identity numerical composites.

Proof. (i) (G1) and (G4) hold by inspection, and G_vU and $\mu_w(U, V)$ are stable-left because their left sides are iterator-headed. For (G2): the two sides of G_0U are iterator-headed and equal to 0 by I0, so Theorem 3.7(iv) proves G_0U. For $U \equiv (s = t)$ put $a := \mathrm{sel}(Sw, 0, s)$ and $b := \mathrm{sel}(Sw, 0, t)$; then $a = s$ and $b = t$ by IS, the term s is not the literal 0 because U is stable-left, and Theorem 3.7(v) gives $U \dashv\vdash G_{Sw}U$. (G3) is the same lemma: for $\mu_0(s = t, r = q)$ with $a := \mathrm{sel}(0, s, r)$, $b := \mathrm{sel}(0, t, q)$, and for μ_{Sw} with $a := \mathrm{sel}(Sw, s, r)$, $b := \mathrm{sel}(Sw, t, q)$, where $r \not\equiv 0$ because the second equation is stable-left. The consequences are Theorem 3.19, and (C) is its case of an equation with an iterator-headed left side over the empty context.

(ii) Write $A(n) := I(n, t;\, u.\, Su)$. The matrix $y = A(y)$ is an orbit matrix, so the formula belongs to $\mathcal{P}_0^+$ and its introduction is in progress mode. With the witness 0 the premise is $\Gamma, z = A(z) \vdash Sz = A(Sz)$, which is the step derived in the proof of Theorem 3.31: the source z is a variable. No base is needed. For (O), let $\Gamma \vdash 0 = A(0)$. DE on the orbit closure adds $Q_y(y = A(y))$, and QE with the base at the target 1 gives $1 = A(1)$.

(iii) By (i), (ii) and Theorem 3.31 the calculus has unrestricted equality. Theorem 3.33 applies: a zero-seeded matrix has provable instances at zero (Theorem 3.13), and every paired formula over an orbit matrix is derivable by (ii).

(iv) With unrestricted equality the zero-seeded scheme on all equations satisfies (G2) and (G3) as in Theorem 3.26, and (T1) to (T4) hold. Apply Theorems 3.24 and 3.25 to the calculus $\mathsf{G}^-_{\mathrm{orb}}$ and the public language $\mathcal{P}_0$. □

In the derivation of $0 = t \vdash t = 0$ the orbit closure is eliminated with the zero-left base $0 = A(0)$ and yields the stable-left equation $1 = A(1)$. This is the clause that Theorem 3.36 excludes from a strictly guarded context, and it is the only use of an unguarded matrix: by (i) the cases needed by Theorem 3.30 are available in G^- itself. The filter must be taken in the orientation of Theorem 3.30, with the iterator on the left; the equation $v = \mathrm{sel}(\zeta(v), v, 0)$ is not stable-left.

Remark 3.42 (Scope and versions). The theorem also holds with $\mathcal{P}_0^+$ as the public language: for an orbit matrix O both $D_{x,y}O$ and its translation $D_{x,y}L(O)$ are derivable, the second in case mode from its base by the elimination of the orbit closure, so the two are interderivable. The mixed reading of Theorem 3.8 agrees with the ordinary reading on $\mathcal{P}_0^+$ as well: at an orbit matrix both readings are provable, the progressive one by the step of (ii) and the conditional one from it by induction, so their agreement does not depend on a provable base. `guarded_orbit` states the calculus with the full native language as proof language, that is U^-, and observes that the derivations constructed lie in $\mathsf{G}^-_{\mathrm{orb}}$; it proves the induction of Theorem 3.21 in the form discussed in Theorem 3.22. The named derivation of $0 = t \vdash t = 0$ recorded by the package has 326 distinct nodes, of which 171 are transports, 122 computation equations and 10 introductions, exactly one of them at an orbit matrix.

3.4.4 Third repair: the mirror guard

This subsection draws on `mirror_guard`. The guard is turned around: an atomic matrix below $D_,$ has the mirrored shape $a = I(y, a;\, u.\, h)$, with one iterator, the seed repeated on the left. The calculus is $\mathsf{M} = \mathsf{K}(\mathcal{P}_m^Q,\, s \not\equiv 0)$ and is strictly guarded: every formula of a derivation is public or a resource. The left side of a mirrored matrix is arbitrary, so the hypothesis of Theorem 3.36 fails, and a zero-left equation $0 = t$ has the mirror gate $0 = \mathrm{sel}(v, 0, t)$, which is again zero-left: it can be padded without a transport at the source 0.

Theorem 3.43 (The mirror guard)**.** (i) *The mirror scheme on all equations is a guard scheme for* M *and* $\mathcal{P}_m$*. Hence* M *admits numerical cases for every public formula, implication between equations with a common left side, and contextual induction for every literal equation, by derivations that substitute into no given derivation.*

(ii) M *has unrestricted equality, admits transport at every source in every template of* $\mathcal{P}_m$*, and derives the same sequents as* $\mathsf{K}(\mathcal{P}_m^Q, \textit{all terms})$*.*

(iii) *With the mirror scheme and* ε *the identity,* T *translates derivations of* N *into derivations of* M*;* $A \dashv\vdash_{\mathsf{N}} TA$ *for every native* A*;* $B \dashv\vdash_{\mathsf{M}} TB$ *for every* $B \in \mathcal{P}_m$*; and for* Γ, B *in* $\mathcal{P}_m$*,* $\Gamma \vdash_{\mathsf{M}} B$ *if and only if* $\Gamma \vdash_{\mathsf{N}} B$*.*

(iv) M *on* $\mathcal{P}_m$ *is bi-interpretable and equiconsistent with* iPRA*, with whole domains, equality as equality and identity numerical composites.*

Proof. (i) (G1) and (G4) hold by inspection. (G2): $H_0(s = t)$ is $s = \mathrm{sel}(0, s, t)$, the reverse of an instance of I0 (Theorem 3.7(iii)). For $H_{Sw}(s = t)$ put $q := \mathrm{sel}(Sw, s, t)$; then $q \not\equiv 0$ and $q = t$ by IS, and Theorem 3.7(vi) gives $(s = q) \dashv\vdash (s = t)$. (G3): for $\mu_0(s = a, s = b)$ take $q := \mathrm{sel}(0, a, b)$ with $q = a$, and for μ_{Sw} take $q := \mathrm{sel}(Sw, a, b)$ with $q = b$; Theorem 3.7(vi)

gives $(s = q) \dashv\vdash (s = a)$ and $(s = q) \dashv\vdash (s = b)$. Only the right side of an equation is expanded or contracted; the left side s is never touched, and it may be the literal 0 or a variable. The consequences are Theorems 3.19 to 3.21.

(ii) By (i), M is closed under induction for literal equations, so Theorem 3.31 gives unrestricted equality. A mirrored matrix has provable instances at zero (Theorem 3.13), so Theorem 3.33 gives the transport in public templates and the equality of the two consequence relations.

(iii) and (iv). (G2) and (G3) hold by (i); (T1) and (T3) are trivial; (T2) holds by unrestricted equality; (T4) is Theorem 3.13. Apply Theorems 3.24 and 3.25. □

In the induction of Theorem 3.31 for the equation $n = I(n, t;\, u.\, Su)$, the guard $J(n)$ of Theorem 3.21 is $n = \mathrm{sel}(n, n, I(n, t;\, u.\, Su))$: a mirrored matrix whose left side is the induction variable. The zero-seeded guard has no such matrix, which is the content of Theorem 3.36; the orbit matrix of the second repair supplies one by hand. The order of the recovery is: equality at admissible sources; expansion on the right; padding and cases; implication with a common left side; induction; equality at the source 0; transport in public templates; the translation. The induction is recovered before the source restriction is lifted, and it uses the implication $J(y) \Rightarrow E(y)$, not $J(y) \Rightarrow E(Sy)$ (Theorem 3.22).

Remark 3.44 (Scope). The mirrored shape shows one iterator where the zero-seeded shape shows two, and it repeats the seed; no statement about the length of expanded terms is made. `mirror_guard` states the calculus with clause contexts and proves its theorem against the calculus of the reports on the language of equations and $D_{,}$ (Theorems 3.4 and 3.28).

3.4.5 Fourth repair: protected atoms

This subsection draws on `protected_core`. The zero-left equations are removed from the public language. A public atom is a protected equation $p = q$, both sides iterator-headed, and ordinary equality is expressed by the wrapped equation:

$$s \simeq t \;:=\; \big(W(s) = W(t)\big), \qquad \bot_W \;:=\; \big(W(0) = W(1)\big).$$

The public language is $\mathcal{P}_W$, with zero-seeded atomic matrices, and the calculus is $\mathsf{H} = \mathsf{K}(\mathcal{P}_W^{Q}, \text{iterator-headed terms})$: its seven rules are Hyp, I0, IS, Tr_I, QE, DE and $\mathrm{DI}_{\mathrm{root}}$. Its proof language contains all equations as auxiliary formulas, which are needed for the computation equation $W(t) = t$ when t is not iterator-headed; they are not public atoms. The source condition is stronger than $s \not\equiv 0$: a variable is not a source. In return the class of sources is closed under substitution.

Lemma 3.45 (Lifting a wrapped equation)**.** *For all terms s, t and every term template $f(v)$, in H,*

$$\Gamma \vdash W(s) = W(t) \quad \textit{implies} \quad \Gamma \vdash W(f(s)) = W(f(t)).$$

Proof. Congruence along $W(s) = W(t)$ in the template $W(f(v))$ gives $W(f(W(s))) = W(f(W(t)))$. Congruence along the computation equations $W(s) = s$ and $W(t) = t$ in the same template gives $W(f(W(s))) = W(f(s))$ and $W(f(W(t))) = W(f(t))$. All sources and all six sides are iterator-headed, so Theorem 3.7(ii) reverses the first of the last two equations and composes the three. The argument does not depend on whether s is the literal 0. □

Theorem 3.46 (The protected core)**.** (i) H *admits weakening, substitution and cut.*

(ii) *The zero-seeded scheme on the protected equations is a guard scheme for H and $\mathcal{P}_W$. Hence H admits numerical cases for every public formula, implication between protected equations, and contextual induction for every protected equation.*

(iii) *With* $\varepsilon(s = t) := (W(s) = W(t))$, T *translates derivations of* N *into derivations of* H*, all of whose transports have iterator-headed sources;* $A \dashv\vdash_{\mathsf{N}} TA$ *for every native* A*;* $B \dashv\vdash_{\mathsf{H}} TB$ *for every* $B \in \mathcal{P}_W$*; and for* Γ, B *in* $\mathcal{P}_W$*,* $\Gamma \vdash_{\mathsf{H}} B$ *if and only if* $\Gamma \vdash_{\mathsf{N}} B$*.*

(iv) H *on* $\mathcal{P}_W$ *is bi-interpretable and equiconsistent with* iPRA*. Both numerical domains are whole; equality is equality, expressed in the public language by* $\simeq$*; both numerical composites are the identity, with the graph* $x \simeq y$*; and*

$$\vdash_{\mathsf{H}} \bot_W \iff \vdash_{\mathsf{N}} 0 = 1 \iff \mathsf{iPRA} \vdash \bot.$$

Proof. (i) is Theorem 3.5: an iterator-headed source keeps its head under substitution, and the zero-seeded shape is preserved. The argument would fail for the sources $s \not\equiv 0$.

(ii) (G1) and (G4) hold by inspection, and the gates and selectors of protected equations are protected. (G2): the two sides of G_0U are iterator-headed and equal to 0, so Theorem 3.7(iv) proves G_0U; and for $U \equiv (p = q)$, Theorem 3.7(v) with $a := \mathrm{sel}(Sw, 0, p)$ and $b := \mathrm{sel}(Sw, 0, q)$ gives $U \dashv\vdash G_{Sw}U$, since p and q are sources. (G3) is the same lemma applied to $\mathrm{sel}(0, p, r)$, $\mathrm{sel}(0, q, s)$ and to $\mathrm{sel}(Sw, p, r)$, $\mathrm{sel}(Sw, q, s)$.

(iii) and (iv). (T1): $W(s) = s$ and $W(t) = t$ in N. (T2): for an instance $c = d$ of I0 or IS the source c is iterator-headed and congruence in the template $W(v)$ gives $W(c) = W(d)$; and from $W(s) = W(t)$ and $W(h(s)) = W(r)$, Theorem 3.45 gives $W(h(s)) = W(h(t))$, and symmetry and transitivity at the source $W(h(s))$ give $W(r) = W(h(t))$. (T3): for a protected $p = q$, Theorem 3.7(v) with $a := W(p)$ and $b := W(q)$ gives $(p = q) \dashv\vdash (W(p) = W(q))$. (T4) is Theorem 3.13. Apply Theorems 3.24 and 3.25; the contradiction $\varepsilon(0 = 1)$ is $\bot_W$. □

The compatibility of the two reductions is obtained here without recovering symmetry at the source 0: every equation that must later act as a major premise is represented at stable iterator roots, the zero computations are joined through $W(0)$, and the equalities of the native derivation are moved by Theorem 3.45. Theorem 3.36 does not apply because no public equation is zero-left. No statement is made about an auxiliary hypothesis $0 = t$ in H.

Remark 3.47 (The executable extension and the form in Lean). The compact checkers of `protected_core`, and those of `guarded_source`, `guarded_orbit`, `deterministic_auxiliary` and `mirror_guard`, accept derivations of U^-. Since $\mathsf{H} \subseteq \mathsf{U}^- \subseteq \mathsf{N}$ as sets of derivable sequents, by inclusion and Theorem 3.6, the public consequences of U^- on $\mathcal{P}_W$ are those of H (Theorem 3.46(iii)), and likewise on $\mathcal{P}_m$ by Theorem 3.43(iii). `protected_core` and `mirror_guard` note that for this inclusion the ordinary reading suffices and the mixed reading is not needed.

In Lean the public languages of this section are formalized in a common form (`NativePublic`). The calculus `PGPrf` is the kernel with the root-determined introduction and with transport at iterator-headed sources only. A gating translation `pub` replaces every equation $s = t$ by the closure of its gate, $Q_w G_w(s = t)$, so that every atomic matrix is guarded (`NFm.pub_guarded`) and the two sides of an opened gate are iterators, and `pgprf_pub_iff`: `PGPrf` $(\Gamma.\mathrm{map}\ pub)\ (pub\ A) \leftrightarrow$ `NPrf` $\Gamma\ A$. The theorem is proved for any gating of equations (`Gating`); the mirror gate $s = I(w, s;\, u.\, t)$ is a second instance (`Gating.mirror`, `NFm.pub_mirrored`), with iterator-headed sources, which is stronger than the exclusion of the literal 0 in Theorem 3.43. The variant `NFm.pubD` uses the paired binder with an unused witness for every closure and reaches the grammar of the packages (`pgprf_pubD_iff`, `NFm.pubD_noQ`, `NFm.pubD_guarded`). The proofs use neither `Classical.choice` nor the bi-interpretation. The differences from the packages are: an equation at the root is gated, not left bare or wrapped as $W(s) = W(t)$; clause contexts and the auxiliary formulas of Section 3.4.2 and Section 3.4.3 are not formalized; and the certificate formats are not modelled.

3.5 Guards on the full native language

This section draws on `diagonal_guard` for the criterion, the diagonal guard and the lemmas on cases and implication, and on `mirror_core` for the mirror selector, its equational decoder, the repair of substitution and the six-rule form. `mirror_core` takes the repair of substitution from `reflexive_base` and the mirrored shape from `mirror_guard`.

In this section the language is the full native language, the contexts are lists of native formulas, and the readings of Q and $D_,$ are the ordinary ones. The progress mode of the single introduction is used at the guarded shape and nowhere else, so the atomic matrices that are not guarded keep their introduction by cases, and Theorem 3.10 does not arise. The equality rule is the rule Tr of the kernel.

3.5.1 Selectors and the criterion

Definition 3.48 (Selector)**.** A *selector* is a property P of matrices $B(x, y)$ that holds only of literal equations, is decided from the syntax of B, and is invariant under renaming of bound variables and under injective renaming of the variables other than x and y. Put $\eta_B^P(z) := z$ if $P(B)$ and $\eta_B^P(z) := 0$ otherwise. The calculus N_P has the rules Hyp, I0, IS, Tr, QE and DE of N (Section 1.3) and the one introduction

$$\frac{\Gamma,\ B(t, \eta_B^P(z)) \vdash B(t, Sz)}{\Gamma \vdash D_{x,y}B(x, y)}\ \mathrm{DI}_P,$$

with z not free in Γ, t and $D_{x,y}B$. It has seven rules. The test P is applied to the matrix before the witness is substituted.

Lemma 3.49 (Structure and equality in N_P)**.** *N_P admits weakening, injective renaming of free variables and cut. Reflexivity, symmetry, transitivity, congruence and transport in an equation template are derivable.*

Proof. The structural statements are proved by induction on derivations as in Theorem 3.5: an injective renaming preserves P and its negation, and weakening and cut do not change a matrix. The equality rules are Theorem 2.4(iii): their derivations use I0 and Tr only. □

Consider two conditions on a selector.

(S1) If $P(B)$, then $\Gamma \vdash B(t, 0)$ for every Γ and t, by I0, IS and Tr.

(S2) Every equation template $E(y)$ has a *lift* $\Lambda_E(y)$: an equation with $P(\Lambda_E)$, as a matrix in y with an unused witness coordinate, such that $\Lambda_E(Sv) \dashv\vdash E(Sv)$ for every term v, by I0, IS and Tr.

Lemma 3.50 (Cases from (S1))**.** *Let P satisfy* (S1)*. In N_P, for every native matrix B and every native formula $F(y)$, with z fresh:*

(i) *if $\Gamma, B(t, 0) \vdash B(t, Sz)$, then $\Gamma \vdash D_{x,y}B$;*

(ii) *if $\Gamma, F(0) \vdash F(Sz)$, then $\Gamma \vdash Q_y F(y)$;*

(iii) *if $\Gamma \vdash F(0)$ and $\Gamma, F(0) \vdash F(Sz)$, then $\Gamma \vdash F(t)$ for every term t.*

Proof. (i) If $P(B)$ fails, this is DI_P. If $P(B)$ holds, cut the derivation of $B(t, 0)$ given by (S1) into the premise, weaken by $B(t, z)$, and apply DI_P in progress mode. (ii) and (iii) By (i) the rule DI is admissible in N_P, and the introduction of Q by cases and numerical cases follow from DI, DE, Hyp and QE as in N (Theorem 2.4(i)). □

No induction for an equation is used in the lemma. For equations $U \equiv (a = b)$ and $V \equiv (c = d)$ and a fresh w put

$$(U \Rightarrow_= V) := Q_w\big(\mathrm{sel}(w, a, c) = \mathrm{sel}(w, b, d)\big).$$

This is the native implication $U \Rightarrow_* V$ of Theorem 1.28 at two equations. Its rules are proved for N in Section 1.4; the next lemma derives them in N_P, where QI is not yet available.

Lemma 3.51 (Implication between equations, and cases on a term of the context)**.** *Let P satisfy* (S1)*. In N_P:*

(i) *if $\Gamma, U \vdash V$, then $\Gamma \vdash U \Rightarrow_= V$; and $\Gamma \vdash U \Rightarrow_= V$ and $\Gamma \vdash U$ imply $\Gamma \vdash V$;*

(ii) *for an equation E and a term r, with v not free in Γ, r and E: if $\Gamma, r = 0 \vdash E$ and $\Gamma, r = Sv \vdash E$, then $\Gamma \vdash E$. The context Γ may contain r and its variables.*

Proof. (i) Let $J(w)$ be the matrix of $U \Rightarrow_= V$. By the selector equations $J(0) \dashv\vdash U$ and $J(Sz) \dashv\vdash V$. For the introduction, over $\Gamma, J(0)$ convert $J(0)$ to U, cut into the given derivation, convert V to $J(Sz)$, and apply Theorem 3.50(ii). For the elimination, convert U to $J(0)$, apply QE at 1 and convert $J(1)$ to V.

(ii) For a new variable y put $F(y) := ((r = y) \Rightarrow_= E)$. By (i) the two hypotheses give $\Gamma \vdash F(0)$ and $\Gamma \vdash F(Sv)$. Theorem 3.50(iii) at the target r gives $(r = r) \Rightarrow_= E$, and reflexivity and (i) give E. The variable of the case distinction is the new y, not a variable of Γ. □

Theorem 3.52 (The selector criterion)**.** *Let P be a selector with* (S1) *and* (S2)*. Then the rule QI of N is admissible in N_P, and for every context Γ and every native formula A,*

$$\Gamma \vdash_{\mathsf{N}_P} A \iff \Gamma \vdash_{\mathsf{N}} A.$$

Both transformations are effective and change no term, formula or assumption. Hence N_P is bi-interpretable and equiconsistent with iPRA *by the interpretations of N (Theorem 1.78).*

Proof. From N_P to N: six rules are shared. An inference DI_P at a matrix without P is DI. At a matrix with P the matrix is a literal equation, and the inference is derived as in Theorem 3.6: QI, QE and DI.

From N to N_P: DI is Theorem 3.50(i), and it remains to derive QI. Let $\Gamma, E(z) \vdash E(Sz)$ with z fresh, let Λ be the lift of E, and put $\Delta := \Gamma, E(0)$.

Decoding the lift. $\Delta, \Lambda(z) \vdash E(z)$, by Theorem 3.51(ii) for the term z. In the branch $z = 0$, transport of the hypothesis $E(0)$ along $0 = z$ gives $E(z)$. In the branch $z = Sw$, transport of $\Lambda(z)$ gives $\Lambda(Sw)$, (S2) gives $E(Sw)$, and transport along $Sw = z$ gives $E(z)$.

Progress of the lift. Cut $E(z)$ into the step premise: $\Delta, \Lambda(z) \vdash E(Sz)$, and by (S2) $\Delta, \Lambda(z) \vdash \Lambda(Sz)$. The variable z is not free in Δ.

Conclusion. Since $P(\Lambda)$, the rule DI_P in progress mode with the witness 0 gives $\Delta \vdash D_{x,y}\Lambda(y)$, and DE and Hyp give $\Delta \vdash Q_y\Lambda(y)$. With $\Lambda(0)$ from (S1), QE at Sz' gives $\Lambda(Sz')$, and (S2) gives $E(Sz')$. Thus $\Gamma, E(0) \vdash E(Sz')$, and Theorem 3.50(ii) gives $\Gamma \vdash Q_y E(y)$.

The base $E(0)$ is retained and discharged at the end: the conditional closure is never identified with unconditional progress, and the argument does not need $E(0)$ to be derivable. □

Proposition 3.53 (Substitution in N_P)**.** (i) *If P is preserved and reflected by substitution, then substitution is admissible in N_P by an induction that keeps the rule and the mode of every inference.*

(ii) *If P is preserved by substitution and satisfies* (S1)*, then substitution is admissible in N_P.*

Proof. Induct on the derivation, renaming each eigenvariable away from the values of the substitution. All rules other than DI_P commute with substitution as in N. For DI_P in progress mode, preservation gives the same mode after substitution. For DI_P in case mode at B: in (i) the substituted matrix does not have P, by reflection, and the same inference applies; in (ii) the induction hypothesis gives the substituted premise by cases, and Theorem 3.50(i) derives the conclusion whether or not the substituted matrix has P. □

Remark 3.54 (Versions of the proof). The criterion is the *guard criterion* of `diagonal_guard`. That package derives QI with the implication $\Lambda(y) \Rightarrow_{=} E(Sy)$, proved by cases on a new variable, and needs the step premise at 0 and at Sv; it substitutes into the derivation of the premise, which is justified for the diagonal selector by Theorem 3.53(i) and, for a selector without the stability property, by translating native derivations by induction on their height and substituting into the native premise first. `mirror_core` decodes the lift by two unconditional equations (Theorem 3.57) and uses the step premise at its own eigenvariable. The proof above takes the order of `mirror_core` and decodes with Theorem 3.51(ii), which is a lemma of `diagonal_guard`; it needs neither a substitution nor a special property of the lift.

3.5.2 The diagonal guard

Theorem 3.55 (The diagonal guard)**.** *Let P hold of the matrices that are diagonal in y (Theorem 3.12), and write $\mathsf{N}_{\mathrm{diag}}$ for N_P.*

(i) *P is a selector that satisfies* (S1) *and* (S2)*, the latter with the diagonal lift $\Delta_y E$.*

(ii) *$\Gamma \vdash_{\mathsf{N}_{\mathrm{diag}}} A$ if and only if $\Gamma \vdash_{\mathsf{N}} A$, for all native Γ and A.*

(iii) *$\mathsf{N}_{\mathrm{diag}}$ admits weakening, substitution and cut, and the substitution keeps the rule and the mode of every inference.*

Proof. (i) (S1): at $y = 0$ both sides of a diagonal matrix have the form $I(0, 0;\, u.\, h')$ and are equal to 0 by I0. (S2): the lift $\Delta_y E$ is diagonal in y, and at $y = Sv$ its two sides are $I(Sv, Sv;\, u.\, s(Sv))$ and $I(Sv, Sv;\, u.\, t(Sv))$, which are equal to $s(Sv)$ and $t(Sv)$ by IS, the updates being constant. No identity between arbitrary iterator bodies is asserted. (ii) is Theorem 3.52. (iii) By Theorem 3.15(iii) the diagonal shape is preserved and reflected, so Theorem 3.53(i) and Theorem 3.49 apply. □

In Lean the calculus is `GPrf` (`NativeDiagonal`), with `gprf_iff_nprf`, `gprf_iff_iprf` and substitution by direct induction (`GPrf.subst`); the equivalence uses neither `Classical.choice` nor the bi-interpretation. The formalization recovers progress by an equational decoder in place of the implication of the package: with $E(y) \equiv (s(y) = t(y))$, numerical cases prove the unconditional equation

$$\mathrm{sel}(y,\ s(0),\ I(y, y;\, u.\, s(y))) = s(y)$$

and the same for t, and from $E(0)$ and $\Delta_v E$ they give $s(v) = \mathrm{sel}(v, s(0), I(v, v;\, u.\, s(v))) = \mathrm{sel}(v, t(0), I(v, v;\, u.\, t(v))) = t(v)$.

The diagonal guard keeps seven rules, the full language and the readings of both binders. The witness-selected calculus W (Theorems 2.15 and 2.17) also has seven rules on the full language, and its mode can change when a substitution turns a witness into the literal 0 (Theorem 2.18); the calculus with orbit elimination (Section 2.5.1) changes the reading of Q at equations and its elimination; and the guarded core restricts the public language. The diagonal guard does none of these.

3.5.3 The mirror selector and six rules

Theorem 3.56 (The mirror selector)**.** *Let* Mir *hold of the matrices that are mirrored in y, and write* $\mathsf{N}_{\mathrm{mir}}$ *for* $\mathsf{N}_{\mathrm{Mir}}$*.*

(i) Mir *is a selector that satisfies* (S1) *and* (S2)*, the latter with the lift* $H_y E(y) = \big(s(y) = \mathrm{sel}(y, s(y), r(y))\big)$ *of* $E(y) \equiv (s(y) = r(y))$*.*

(ii) $\Gamma \vdash_{\mathsf{N}_{\mathrm{mir}}} A$ *if and only if* $\Gamma \vdash_{\mathsf{N}} A$*.*

(iii) *Substitution is admissible in* $\mathsf{N}_{\mathrm{mir}}$*. It does not keep the mode of every inference.*

Proof. (i) (S1): the instance at zero is $a' = I(0, a';\, u.\, h')$, the reverse of an instance of I0. (S2): $\mathrm{sel}(Sv, s, r) = r$ by IS. (ii) is Theorem 3.52. (iii) The shape is preserved (Theorem 3.15(ii)), so Theorem 3.53(ii) applies. It is not reflected: the matrix $p = I(y, q;\, u.\, u)$ is not mirrored for distinct variables p, q and is mirrored after both are replaced by one variable, so an introduction by cases at this matrix becomes, under that substitution, an instance of Theorem 3.50(i), which cuts the provable base and weakens to progress. □

The selector is tested on the matrix before the witness is substituted: the matrix $x = I(y, 0;\, u.\, 0)$ is not mirrored, although its instance at the witness 0 is $0 = I(y, 0;\, u.\, 0)$.

Lemma 3.57 (The equational decoder of the mirror lift)**.** *Let* $E(y) \equiv (s(y) = r(y))$ *and* $\Lambda(y) \equiv \big(s(y) = \mathrm{sel}(y, s(y), r(y))\big)$*. In* $\mathsf{N}_{\mathrm{mir}}$*, for every term* v*,* $\Gamma \vdash E(0)$ *and* $\Gamma \vdash \Lambda(v)$ *imply* $\Gamma \vdash E(v)$*, by numerical cases for equations and equality alone.*

Proof. The equations

$$\mathrm{sel}\big(v,\ s(0),\ s(v)\big) = s(v),$$
$$\mathrm{sel}\big(v,\ r(0),\ \mathrm{sel}(v, s(v), r(v))\big) = r(v)$$

are derivable over the empty context by Theorem 3.50(iii): at 0 both left sides are equal to their seed by I0, and at a successor the first uses one application of IS and the second two. They are proved before any assumption about v is retained, and then weakened. From $\Lambda(v)$, congruence in the template $\mathrm{sel}(v, s(0), -)$ gives $\mathrm{sel}(v, s(0), s(v)) = \mathrm{sel}(v, s(0), \mathrm{sel}(v, s(v), r(v)))$, and the base $s(0) = r(0)$ changes the seed on the right. Hence

$$s(v) = \mathrm{sel}\big(v, s(0), s(v)\big) = \mathrm{sel}\big(v, s(0), \mathrm{sel}(v, s(v), r(v))\big) = \mathrm{sel}\big(v, r(0), \mathrm{sel}(v, s(v), r(v))\big) = r(v).$$

□

The lemma replaces the first step of the proof of Theorem 3.52 for the mirror selector, without an implication between equations.

Definition 3.58 (The six-rule mirror calculus)**.** The calculus M_6 is the calculus E_2 of Theorem 4.9 with its introduction WI replaced by the mirror-selected one: its rules are Hyp, Refl, the two cases $\textsc{Eval}_0$ and $\textsc{Eval}_S$ of Eval, QE, DE and

$$\frac{\Gamma,\ B(t, \eta_B^{\mathrm{Mir}}(z)) \vdash B(t, Sz)}{\Gamma \vdash D_{x,y} B(x, y)}\ \mathrm{DI}_{\mathrm{Mir}}.$$

It has six rule families and seven inference constructors, counted as in Theorem 4.10. There is no computation axiom, no transport rule, no structural rule and no rule of induction.

Theorem 3.59 (Six rules with the mirror selector)**.** *For every context* Γ *and every native formula* A*,*

$$\Gamma \vdash_{\mathsf{M}_6} A \iff \Gamma \vdash_{\mathsf{N}_{\mathrm{mir}}} A \iff \Gamma \vdash_{\mathsf{N}} A.$$

M_6 *admits weakening, injective renaming, cut and substitution, and is bi-interpretable and equiconsistent with* iPRA *by the interpretations of* N*.*

Proof. The derivations of symmetry and transitivity, of I0 and IS, and of Tr in E_2 (Theorems 4.13, 4.14 and 4.16) use Refl and the two cases of Eval only, so they are derivations of M_6; the other rules of $\mathsf{N}_{\mathrm{mir}}$ are rules of M_6. Conversely Refl and both cases of Eval are derived from I0, IS and Tr as in the proof of Theorem 4.18, hence in $\mathsf{N}_{\mathrm{mir}}$. The second equivalence is Theorem 3.56. Weakening, injective renaming and cut are proved for M_6 as for E_3 in the proof of Theorem 4.18, with the observation that an injective renaming preserves and reflects Mir. Substitution is proved as in Theorem 3.53(ii): Eval commutes with substitution (Theorem 4.21), and the derived introduction by cases of Theorem 3.50(i) is available in M_6 by the first equivalence. □

Remark 3.60 (The derivations of `mirror_core`)*.* `mirror_core` derives the equality rules of M_6 directly. It joins $s = t$ and $s = r$ to $r = t$ through $I(0, s;\, u.\, u)$ by two instances of $\textsc{Eval}_0$, which is the identity frame of Theorem 4.11, and it derives Tr from the two iterators $I(0, s;\, v.\, h(v))$ and $I(1, s;\, v.\, h(v))$ by the composition described in the remark after Theorem 4.16.

In Lean the seven-rule calculus $\mathsf{N}_{\mathrm{mir}}$ is `MiPrf` (`miprf_iff_nprf`) and the six-rule calculus is `M6Prf` (`m6prf_iff_nprf`), in the module `NativeMirror`. The structural rules proved there are weakening, injective renaming (`MiPrf.ren`) and cut, which suffice for weakening below a binder and for the equivalence; the progress of an arbitrary equation uses the lift $s = \mathrm{sel}(y, s, t)$. Relative to the six-rule calculus with the diagonal selector (`SPrf`), the mirror selector shows one iterator where the diagonal shows two, and it loses the reflection of the shape.

3.6 Comparison of the guards

This section compares the guards of `guarded_core`, `protected_core`, `mirror_guard`, `mirror_core` and `diagonal_guard`; Table 3.2 restates results of the preceding sections.

Guard	Atomic matrix	Lift of $s = t$	Under substitution	Used in
zero-seeded	$I(y, 0;\, u.\, h) = I(y, 0;\, u'.\, k)$	$\mathrm{sel}(y, 0, s) = \mathrm{sel}(y, 0, t)$	preserved, not reflected	$\mathcal{P}_0$: G, U^-, $\mathsf{G}^-_{\mathrm{orb}}$
wrapped	zero-seeded; root atoms protected	$W(s) = W(t)$ at the root	preserved; sources closed	$\mathcal{P}_W$: H
mirror	$a = I(y, a;\, u.\, h)$	$s = \mathrm{sel}(y, s, t)$	preserved, not reflected	$\mathcal{P}_m$: M; full language: $\mathsf{N}_{\mathrm{mir}}$, M_6
diagonal	$I(y, y;\, u.\, h) = I(y, y;\, u'.\, k)$	$I(y, y;\, u.\, s) = I(y, y;\, u.\, t)$	preserved and reflected	full language: $\mathsf{N}_{\mathrm{diag}}$

Table 3.2: The four guards.

The base. Every guard has an instance at zero that is derivable by I0 and equality at iterator-headed sources (Theorem 3.13). For the mirror guard it is the reverse of one computation equation. For the zero-seeded and the diagonal guard two computations are joined; under a source restriction the join passes through $W(0)$ (Theorem 3.7(iv)).

The left side. The zero-seeded, wrapped and diagonal shapes have an iterator on the left. With the sources $s \not\equiv 0$ and the uncrossed transport this makes a zero-left hypothesis inert (Theorem 3.36): the zero-seeded guard then needs an unguarded matrix in the proof language

(Theorems 3.39 and 3.41), or public atoms that are never zero-left (Theorem 3.46). The mirror guard leaves the left side free and is complete in a strictly guarded language (Theorem 3.43).

Sources. The guarded core uses all sources. The mirror guard, and the zero-seeded guard with auxiliary or orbit formulas, use the sources $s \not\equiv 0$, a class that is not closed under substitution; their proofs avoid substitution until equality is unrestricted. The protected core uses iterator-headed sources, a class closed under substitution, and pays with a wrapper on every public atom and with a larger validator of the public grammar (Section 3.7).

Substitution of the shape. In a public language the mode is selected by the root and the shape is a condition on the language, so preservation is enough. As a selector on the full language the shape must not be created by substitution if the mode of an inference is to be stable: the diagonal shape is the only one of the four with this property (Theorem 3.15), and for it substitution is a rule-preserving induction (Theorem 3.55). The mirror selector is closed under substitution with a repair (Theorem 3.56). The zero-seeded shape also satisfies (S1) and (S2), with the gate $G_y E$ as the lift, so by Theorems 3.52 and 3.53 it is a selector on the full language with the same repair; `diagonal_guard` records the matrix $I(y, p;\, u.\,0) = I(y, 0;\, u.\,0)$ as the witness that its mode is not stable.

Cost of recognition. The zero-seeded and the diagonal test inspect fixed positions of the two sides. The mirror test compares two term trees, which is linear in their size; `mirror_core` notes that its smaller checker source does not imply faster checking or shorter proofs.

3.7 Checkers and their sizes

This section records the programs of `guarded_core`, `guarded_source`, `guarded_orbit`, `deterministic_auxil` `mirror_guard`, `protected_core`, `mirror_core` and `diagonal_guard`. Sizes are measured by the protocol of Section 1.10.

The compact checkers. `guarded_core`, `guarded_source`, `guarded_orbit`, `deterministic_auxiliary`, `mirror_guard` and `protected_core` modify the arity-coded checkers of the earlier packages; the codec is inherited and is not changed. A formula array of length one is $D_,$, of length two an equation, of length three an administrative Q; a term array of length one is a successor and of length three an iterator; the other lengths denote zero or truth. The root-determined introduction replaces the comparison of the parity of the matrix with a field of the certificate, `len(a) % 2 == u`, by the parity alone, `not len(a) % 2`: among the meaningful roots only the equation has even length, so an equation retains its instance at the eigenvariable and a compound matrix its instance at zero. `guarded_source` writes the test as `len(a) == 2`, and `deterministic_auxiliary` has an optional spelling `~len(a) % 2` that saves three bytes. The byte profile of `guarded_core` also moves the check of the major premise into the expression that supplies the hypothesis, which preserves every successful run. Source selection is the test `m == ([s, u] if s else [t] * 3)`: the canonical literal 0 is the integer 0, which is false, and selects the Q major, and every other source selects the equation. Computation packets that cannot have a literal 0 as initializer or as predecessor are obtained from the *certified zero* $c_0 := I(2, 1;\, u.\,0)$, with $c_0 = 0$, by a transport at the iterator-headed source c_0, so that no transport at the literal 0 is reintroduced. The byte and AST profiles use lexical scopes and the token profiles a global supply of fresh names; their certificates are not interchangeable.

The contract of the compact checkers. A compact checker accepts derivations of U (`guarded_core`) or of U^- (the other five) in the administrative syntax. Each package states that, for a closed public formula A in canonical encoding and in the ideal unbounded execution, an accepting certificate exists if and only if A is derivable in its calculus. The soundness half rests on the correctness of the inherited codec (opening, freshness and the readings of the

surplus arities), which these packages cite and do not reprove, together with Theorems 3.27, 3.39, 3.41, 3.43 and 3.46 and Theorem 3.47; the completeness half encodes a derivation with fresh integer names. A compact checker does not test that the goal is public, and it accepts some goals that are not. Each package therefore supplies a *strict* checker, which validates the closed public goal and then runs the same kernel, and counts it in full. The strict checker of `mirror_guard` compares the two occurrences of the seed after raising the left side across the binder of the iterator; a comparison of the raw arrays would reject matrices with bound variables. One inherited negative control that changes only the retired mode field is accepted by the new checkers; an old certificate that introduces an unguarded equation by cases is rejected by them, which is Theorem 3.10 and the reason why the raw certificate languages of the old and the new checkers are not claimed to coincide.

The typed checkers. `mirror_core` and `diagonal_guard` measure checkers with tagged terms, formulas and proof trees in de Bruijn form, which validate every term, formula, scope and proof node of the full native language. The diagonal checker tests, after opening the witness, whether both sides begin with an iterator whose counter and seed are the bound index 0; by Theorem 3.15(iii) the test after opening is equivalent to the test on the matrix. The mirror checker tests the matrix before the witness is substituted. The two packages differ from their controls in the guard of the introduction only. The translator of `diagonal_guard` implements both directions of Theorem 3.55(ii); three translated certificates grow from 485, 885 and 1,965 bytes to 11,506, 14,218 and 20,283 bytes. `mirror_core` supplies the translation from M_6 to the kernel, the derived introduction by cases and the repaired substitution as programs; the translation of QI into M_6 is given by the written proof only.

Recorded runs. The packages record the following executions of their programs on positive and negative certificates, in ordinary and optimized Python, all with the expected outcome: 800 (`guarded_core`), 150 (`guarded_source`), 224 (`guarded_orbit`), 242 (`deterministic_auxiliary`), 480 (`mirror_guard`), 608 (`protected_core`); for `mirror_core`, 92 command-line checks of 23 fixtures, 36 replays in the native checker and 72 replays of numerical substitutions; for `diagonal_guard`, 192 processes on 32 fixtures, 93 validations of translated certificates and 240 finite checks of the preservation and reflection of the guard. The runs test the programs and the certificates; the theorems of this chapter are the written proofs.

Table 3.3: Programs and sizes as the packages state them. The last column is the matched control with its tokens, AST nodes and normalized bytes. The smallest values of one package are attained by different programs.

Package	Program	Tokens	AST	Bytes	Matched control
`guarded_core`	direct	265	308	528	byte profile, 266/310/529
	byte profile	266	307	527	byte profile, 266/310/529
	token profile	260	290	575	token profile, 261/292/576
	AST profile	263	286	607	AST profile, 263/289/613
	readable	292	336	1275	none
	strict	491	568	1208	old core with goal parser, 4
`guarded_source`	token profile	257	285	568	`source_dispatch`, 259/289
	AST profile	261	279	603	`source_dispatch`, 263/283
	byte profile	264	301	522	`source_dispatch`, 266/305
	strict	490	561	1202	`guarded_core` strict, 491/5

Package	Program	Tokens	AST	Bytes	Matched control
`guarded_orbit`	token profile	258	287	571	`source_dispatch`, 259/289
	AST profile	262	281	606	`source_dispatch`, 263/283
	byte profile	265	303	525	`source_dispatch`, 266/305
	strict	491	563	1205	guarded 491/568/1208; sca
`deterministic_auxiliary`	direct	265	303	525	`source_dispatch`, 266/305
	byte profile	265	303	522	`source_dispatch`, 266/305
	AST profile	262	281	606	`source_dispatch`, 263/283
	token profile	258	287	571	`source_dispatch`, 259/289
	strict	491	563	1205	guarded 491/568/1208; nat
`mirror_guard`	token profile	258	287	571	`source_dispatch`, 259/289
	AST profile	262	281	606	`source_dispatch`, 263/283
	byte profile	265	303	525	`source_dispatch`, 266/305
	readable	279	319	1272	none
	strict	578	661	1503	strict derivative of the sour
`protected_core`	token profile	258	287	571	`source_dispatch`, 259/289
	AST profile	262	281	606	`source_dispatch`, 263/283
	byte profile	265	303	525	`source_dispatch`, 266/305
	readable	284	326	1301	none
	strict	505	582	1251	`guarded_core` strict, 491/5
`mirror_core`	mirror selector, six rules	1012	1082	3109	diagonal selector, six rules,
`diagonal_guard`	diagonal, seven rules	1069	1124	3244	native, eight rules, 1101/11 witness-selected, seven rule

The gains recorded in Table 3.3 are small. The compact profiles of `guarded_orbit`, `deterministic_auxiliary`, `mirror_guard` and `protected_core` have the same three sizes, since they share the two dispatch tests and differ in the calculus that justifies them and in the strict validator. The strict checkers do not improve on the unrestricted control of 1109 bytes; those of `mirror_guard` and `protected_core` are larger than their matched controls. The gzip size of the byte profile of `guarded_orbit`, `deterministic_auxiliary` (direct), `mirror_guard` and `protected_core` is 331 bytes against 330 for the control. The diagonal checker is 39 bytes larger than the witness-selected control; its advantage is the stability of the mode under substitution. The typed checkers are not smaller than the compact ones, which have a different input contract.

3.8 Further questions

This section collects the open questions and the unproved or negative statements of `guarded_core`, `guarded_source`, `guarded_orbit`, `deterministic_auxiliary`, `mirror_guard`, `protected_core`, `mirror_core` and `diagonal_guard`.

Question 3.61 (Minimal proof language)**.** For the zero-seeded public language $\mathcal{P}_0$ with the sources $s \not\equiv 0$, Theorem 3.37 shows that some unguarded matrix is needed and Theorem 3.41 shows that the orbit matrices suffice. Which substitution-stable families of auxiliary equation invariants suffice for the repair of equality and for the arithmetic of the bridge? Is there a comparably short checker that enforces the smaller intermediate grammar, with a simpler soundness proof than that of the full arity codec? (`guarded_source`, `guarded_orbit`, `deterministic_auxiliary`.)

Question 3.62 (The smallest stable guard)**.** Is there a syntactic class of equations that is cheaper to recognize than the diagonal shape and has provable instances at zero, a lift of every equation, and preservation and reflection under substitution? What is the smallest family of base recognizers that supports an equational decoder for every equation? The mirror and the diagonal shape are two instances; neither gives a lower bound. (`diagonal_guard`, `mirror_core`.)

Question 3.63 (A public grammar with a small validator)**.** Is there a substitution-stable public grammar that makes the root-determined introduction complete, keeps both composites, and has a strict checker no larger than the unrestricted control of 1109 bytes under the same complete-input contract? (`guarded_orbit`, `deterministic_auxiliary`; `guarded_source` asks whether the validator can share more of the kernel.)

Question 3.64 (Zero-source symmetry)**.** Can the derivation of $t = 0$ from $0 = t$ with the uncrossed rule (Theorem 3.31) be made shorter than the one through the singleton test and the filter? A shorter uniform derivation would shorten certificates without changing a checker. (`guarded_source`, `deterministic_auxiliary`.)

Question 3.65 (Sizes of translated derivations)**.** Give explicit bounds for the expansion caused by the join at zero, the padding, the implication between equations and the lift of an equation, and compare proof trees with a checked format with sharing. No polynomial bound on the size of a translated certificate is claimed. (`guarded_orbit`, `diagonal_guard`.)

The following items are stated by the packages without proof, or record what was not achieved.

- *No minimality.* No package claims a least number of rules, a least checker, or a lower bound for a class of checkers; the observed sizes are upper bounds, and the three smallest coordinates of a package belong to different programs. Whether one checker improves all three coordinates of its controls at once is open (`guarded_source`), as is a lower bound for a precisely specified class of complete checkers (`mirror_core`).

- *The codec.* The theorems on the compact checkers are stated relative to the inherited arity codec, whose soundness is cited and not reproved in these packages, and neither the codec nor a certificate format is formalized in a proof assistant.

- *Administrative formulas.* Completeness is not claimed for goals outside the public language, and the full language of U or U^- is not claimed to be bi-interpretable with iPRA; by Theorem 3.10 the two sound readings of an unguarded atomic binder differ. No impossibility is claimed for other interpretations of the administrative language (`guarded_core`). Whether the independent choice between cases and progress can be removed on every administrative formula with its ordinary reading and without a validator is asked by `guarded_orbit`; the selectors of Section 3.5 remove it on the full native language at the cost of a test of the shape.

- *Formalization.* In Lean the dependency theorem (`ZKernel.trF`), the diagonal guard (`GPrf`), the mirror selector (`MiPrf`, `M6Prf`) and the public languages in the gated form (`PGPrf`) are proved. Not formalized: the calculi with clause contexts; the auxiliary formulas of Section 3.4.2 and the orbit bootstrap of Section 3.4.3, including numerical cases at stable-left equations; the mirror calculus M with the sources $s \not\equiv 0$ in the form of Theorem 3.43; and the wrapped root atoms of H. `deterministic_auxiliary` asks whether the auxiliary progress of an equation can be a separate proof judgment with a smaller derivation datatype, the target being the equality of the public consequence relations and not a round trip for auxiliary formulas; `mirror_core` proposes its two

unconditional selector equations (Theorem 3.57) as the route with the fewest dependencies. `mirror_guard` observes that a formalization of M cannot start by restricting the transport rule of the kernel and using its substitution theorem (Theorem 3.35).

- *Programs that are not complete translators.* The proof generators of `mirror_guard` and `protected_core` are collections of worked constructions; the normalizer of `protected_core` rejects an unsupported equation leaf with the source 0 instead of translating it; `mirror_core` has no program for the translation of QI.

- *Selectors in the compact interface.* Whether the mirror or the diagonal selector reduces the size of a compact checker with a complete validator of the goal is not answered (`mirror_core`, `diagonal_guard`); nor is the combination of the diagonal introduction with the crossed one-term transport in a matched typed checker (`diagonal_guard`), or a constant-time mirror test by sharing of syntax without a larger trusted reader (`mirror_core`).

- *Other bases.* `guarded_orbit` asks whether a different numerical basis can remove a constructor or a binding operation without a richer arithmetic axiom or a more complex parser, and notes that the earlier register, tied-seed and zero-start results constrain the answer.

Chapter 4

Computation, equality and normal forms

Three of the eight rules of the native scalar core N (Section 1.3) concern numerical terms: the zero computation equation, the successor computation equation and the transport rule. This chapter collects the calculi in which these three rules are merged, and the calculi in which one or both computation equations leave the rule inventory and become part of the syntax of terms. Two ideas organize the material.

The first idea keeps the terms, the formulas and ordinary substitution unchanged. Reflexivity and one rule Eval, which composes an equation with one computation step, replace the three rules (Section 4.2).

The second idea restricts the terms to the normal forms of a computation and lets every substitution of a rule be followed by normalization. A computation that is normalized needs no rule. Section 4.3 proves once, for an arbitrary normalizer that satisfies five conditions, that derivations transfer between N and the kernel on normal forms. The zero computation alone is normalized in Section 4.4; on the resulting zero-normal terms the remaining computation and the equality reasoning are presented by the rules of Sections 4.5 to 4.8. Both computations are normalized in Section 4.9.

Every calculus K of the chapter is given with its complete rule inventory and with a theorem of one of two forms: K derives exactly the sequents of N, or K derives the normal form of a sequent exactly when N derives the sequent. In both cases the bi-interpretation of Theorem 1.78 transfers (Theorem 4.31). Table 4.1 lists the calculi; the rule names are explained in the sections cited.

4.1 Conventions and shared lemmas

This section fixes the notation common to the nine packages of the chapter (`computational_equality`, `equality_from_iteration`, `uniform_step`, `seed_evaluation`, `zero_normal`, `reflexive_base`, `initializer_equality`, `full_normalization`, `computation_normal`) and proves once the arguments that several of them repeat.

4.1.1 Notation

Raw terms are the terms of Section 1.1, with ordinary capture-free substitution; no computation is performed by substitution, so $I(0, a;\, u.\, h)$ is a raw term distinct from a. The relation $s \equiv t$ is literal identity of syntax, up to the renaming of bound variables; in a de Bruijn representation it is equality of finite trees. Every test in a rule of this chapter is a test of literal identity and never a test of provable equality. A formula is a *literal equation* if its outermost

Calculus	Terms	Rules	Equivalence
E_3	raw	Hyp, Refl, Eval (three cases), QE, DE, WI	Theorem 4.18
E_2	raw	the same, Eval with two cases	Theorem 4.12
$\mathsf{R}^{\mathrm{raw}}$	raw	Hyp, I0, IS, Tr, QE, DE, RI	Theorem 4.53
N^0	zero-normal	Hyp, IS, Tr, QE, DE, DI, QI	Theorem 4.38
Z_6	zero-normal	Hyp, IS, Tr, QE, DE, WI	Theorem 4.42
R	zero-normal	Hyp, IS, Tr, QE, DE, RI	Theorem 4.51
S	zero-normal	Hyp, IS, Seed, QE, DE, WI	Theorem 4.64
U	zero-normal	Hyp, Refl, Step, QE, DE, WI	Theorem 4.70
$\mathsf{C}_{\mathrm{Step}}$	zero-normal, clauses	Use, Refl, Step, DE, WI	Theorem 4.73
$\mathsf{C}_{\mathrm{Seed}}$	zero-normal, clauses	Use, Refl, SeedEval, DE, WI	Theorem 4.77
$\mathsf{C}_{\mathrm{App}}$	unit-normal, clauses	Use, Refl, App, DE, WI	Theorem 4.85
N^{f}	full normal	Hyp, Refl, Tr, QE, DE, DI, QI	Theorem 4.94
F	full normal	Hyp, Refl, Tr, QE, DE, WI	Theorem 4.98

Table 4.1: The calculi of the chapter.

constructor is equality; an equation below Q_y or $D_{x,y}$ is not a literal equation. The numeral 1 is $S0$.

A *template* is a term h with a distinguished variable v; it may contain v several times, not at all, or below iterator binders, and it may contain other variables. Its instance at s is written $h(s)$. A *matrix* is a formula B with one or two distinguished variables; $B(r)$ and $B(t, r)$ are its instances. Instances are formed with the substitution of the calculus at hand: ordinary substitution on raw terms and the normalizing substitution of Theorem 4.25 on normal forms.

The rules of N are those of Section 1.3, with their names: Hyp, I0, IS, Tr (from $s = t$ and $h(s) = r$ infer $r = h(t)$), QE, DE, DI (by cases) and QI (by progress, for literal equations). The first premise of Tr is its *major* premise and the second its *minor* premise. Weakening and substitution of terms in derivations (Theorem 1.10), cut (Theorem 1.11), the equality rules (Theorem 1.18) and the Leibniz rule (Theorem 1.22) are admissible in N.

Definition 4.1 (Witness-selected introduction)**.** WI is the rule DI_{w} of Theorem 2.15, with the function $\eta_{B,t}$ defined there, read with the terms and the substitution of the calculus at hand; the test $t \equiv 0$ is applied to the witness as it is supplied in that syntax. An instance with $\eta_{B,t}(z) = z$ is in *progress mode*; every other instance is in *case mode* and is an instance of DI.

On raw terms WI replaces DI and QI by Theorem 2.17, whose proof repairs the one missing instance of DI at the term $I(0, 0; u.\, 0)$ of Theorem 2.16. That term is not available on normal forms. Theorem 4.3 states the argument for an arbitrary syntax and an arbitrary term that is provably 0 and not literally 0; the sections below supply the term.

4.1.2 Equality from reflexivity and crossed transport

Lemma 4.2. *Let K be a calculus that has Tr for every template, with a substitution for which the instance of the identity template at s is s.*

(a) If $\Gamma \vdash_{\mathsf{K}} \ell = r$ for some term ℓ, then $\Gamma \vdash_{\mathsf{K}} r = r$.

(b) If K derives $r = r$ for every term r, then the following rules are admissible in K, by derivations that use only reflexivity and Tr: symmetry; transitivity; congruence, from

$s = t$ infer $h(s) = h(t)$; and transport in a literal equation, from $s = t$ and $a(s) = b(s)$ infer $a(t) = b(t)$.

Proof. (a) is the derivation of Theorem 1.18(i) with the equation $\ell = r$ in place of the instance of I0: Tr with the identity template and $\ell = r$ as both premises concludes $r = r$. (b) The derivations of Theorem 1.18(ii)–(v) use reflexivity and Tr only, one application for each of symmetry, transitivity and congruence and two for transport in an equation; they apply to K with the instances of the templates formed by the substitution of K. □

For a calculus on raw terms that contains I0 the lemma is Theorem 2.4(iii). It is stated here for the calculi in which I0 is not a rule and reflexivity has another source.

4.1.3 The single introduction

Lemma 4.3 (The single introduction)**.** *Let K be a calculus on native formulas over a term syntax with a substitution for which instances compose (ordinary substitution on raw terms, or a normalizing substitution, by Theorem 4.26). Assume that K has Hyp, QE and DE; that weakening, injective renaming of free variables and cut on an assumption are admissible in K; and that symmetry of equality and transport in literal equations are admissible in K.*

(i) If K has WI, then QI is admissible in K, and so is the introduction of a closure by cases, for every matrix B:

$$\frac{\Gamma,\, B(0) \vdash B(Sz)}{\Gamma \vdash Q_y B(y)} \qquad (z \text{ fresh}).$$

Every instance of DI is an instance of WI, except when B is a literal equation and $t \equiv 0$.

(ii) If K has WI, $E(x, y)$ is a literal equation and Z is a term with $Z \not\equiv 0$ and $\Gamma \vdash_{\mathsf{K}} Z = 0$, then

$$\Gamma,\, E(0,0) \vdash_{\mathsf{K}} E(0, Sz) \quad (z \text{ fresh}) \qquad \text{implies} \qquad \Gamma \vdash_{\mathsf{K}} D_{x,y} E(x, y).$$

(iii) If K has DI and QI, then WI is admissible in K.

Proof. The three parts are the three constructions in the proof of Theorem 2.17, which use of the calculus only the rules and the admissible operations assumed here.

(i) For QI, the construction given there applies: WI for the matrix $E(y)$ with an unused witness coordinate, at the witness 0, is in progress mode and has the given premise, and DE with Hyp returns $Q_y E(y)$. For the closure by cases use the same construction with $B(y)$ in place of $E(y)$ and the witness 1: since $1 \not\equiv 0$, WI is in case mode for every B, and its premise is $\Gamma, B(0) \vdash B(Sz)$. In N this rule is Theorem 1.13, with the witness 0; with WI the witness must not be the literal 0. The last assertion is the definition of the two modes.

(ii) This is the repair in the proof of Theorem 2.17, with Z in place of the term c of Theorem 2.16, after z is renamed so that it is fresh for Z: in the context $\Gamma, E(Z, 0)$, transport along $Z = 0$, cut with the given derivation, and transport along $0 = Z$ give $E(Z, Sz)$, and WI at the witness Z is in case mode. Only transport in literal equations is used, and no term is substituted into the given derivation.

(iii) An instance of WI in case mode is an instance of DI. An instance in progress mode, with the premise $\Gamma, E(0, z) \vdash E(0, Sz)$, is derived as in Theorem 1.16(i), by QI, QE under the assumption $E(0, 0)$, and DI at the witness 0. □

Remark 4.4. The term Z of part (ii) is $I(0, 0;\, u.\, 0)$ on raw terms (Theorem 2.16), the terms $I(1, 0;\, u.\, 0)$ or $I(1, 0;\, u.\, u)$ on zero-normal terms (Theorem 4.40), and a variable b opened by DE on full normal forms (Theorem 4.97). As on raw terms (Theorem 2.18), no calculus with

WI is shown closed under the substitution of terms by a direct induction on derivations; on normal forms a substitution can in addition create a redex whose contraction turns a witness into the literal 0. Substitution is transferred from an equivalent calculus with DI and QI. Weakening, injective renaming and cut leave the witness and the root of the matrix unchanged and are proved directly, as in Theorem 2.3.

4.1.4 Congruence below binders and provability of conversion

Transport in N replaces a term in a template by capture-free substitution. It does not replace the body of an iterator by a pointwise equal body, and it does not replace a matrix below Q_y or $D_{x,y}$. Both replacements are derived rules of N: the congruence of the body of an iterator is Theorem 1.39(ii), which rests on an equation induction, and the congruence of the binders is Theorem 1.21. Every comparison of N with a calculus on normal forms depends on them, because a normalizer rewrites below binders.

Definition 4.5 (Computation reduction)**.** The relation $\to$ on raw terms is the closure, under every term context including the bodies of iterators, of the two contractions

$$I(0, a;\, u.\, h) \to a, \qquad I(Sn, a;\, u.\, h) \to h[I(n, a;\, u.\, h)/u].$$

The first is the *zero contraction* and the second the *successor contraction*; $\to^*$ is the reflexive and transitive closure.

Lemma 4.6 (Conversion is provable)**.** *If $t \to^* t'$, then $\vdash_{\mathsf{N}} t = t'$, with all free variables of t as parameters. If the formula A' is obtained from the formula A by replacing each term by a reduct of it, then $\Gamma, A \vdash_{\mathsf{N}} A'$ and $\Gamma, A' \vdash_{\mathsf{N}} A$ for every Γ.*

Proof. For one step $t \to t'$ argue by induction on the position of the contracted redex. At the root the equation is an instance of I0 or IS. Below a successor, in the counter or in the initializer of an iterator, it follows from the induction hypothesis by congruence in the template with one hole at that position. In the body of $I(n, a;\, u.\, h)$, with $h \to h'$, the induction hypothesis is $\vdash_{\mathsf{N}} h = h'$ with u free, in the empty context, and Theorem 1.39(ii) gives $I(n, a;\, u.\, h) = I(n, a;\, u.\, h')$. Transitivity extends the statement to $\to^*$.

For formulas argue by induction on A. At a literal equation use the two term equations and the equality rules. At $Q_y B$ and $D_{x,y} B$ the induction hypothesis holds for the matrix with the bound variables free, in both directions, and Theorem 1.21 applies. □

Remark 4.7. The congruence of the body is restated as a lemma in `zero_normal`, `initializer_equality`, `full_normalization` and `computation_normal`, each time by a direct equation induction on $I(z, a;\, u.\, h) = I(z, a;\, u.\, h')$ in place of the uniqueness lemma used for Theorem 1.39; its version in T_{eq} is formalized as `Q.iter_body_congr` and `Q.iter_congr`. The pointwise equation of the bodies is derived in a context without u; it is not a hypothesis that may vary with the state of the iteration. The equality reasoning of the calculi below is recovered without induction, whereas the comparison of each calculus on normal forms with N uses the equation induction inside Theorem 1.39(ii). The packages `uniform_step` and `seed_evaluation` state this dependency explicitly.

4.2 The rule Eval on raw terms

This section draws on `computational_equality`, which introduces the rule Eval with three cases, and on `equality_from_iteration`, which shows that the first case is redundant and that neither of the other two is. The terms, the formulas and substitution are those of N; nothing is normalized.

4.2.1 The operation Φ and the calculi

Definition 4.8. For raw terms s, t, b the partial operation $\Phi(s, t, b)$ is given by the rows

shape of b	required source s	$\Phi(s, t, b)$
$b \equiv s$	s	t
$b \equiv I(0, a;\, u.\, h)$	$s \equiv a$	t
$b \equiv I(Sc, a;\, u.\, h)$	$s \equiv I(c, a;\, u.\, h)$	$h[t/u]$

and is undefined if no row applies.

The rows are disjoint. In the second row s is a proper subterm of b, and in the third row b has exactly one node more than s, so neither overlaps the first; the second and third rows have different constructors at the root of the counter. Hence Φ is a deterministic partial operation on syntax. The third row requires an equation for the whole preceding iterate $I(c, a;\, u.\, h)$; it does not compare initializers, and it applies unchanged when $c \equiv 0$.

Definition 4.9 (The calculi E_3 and E_2)**.** The calculus E_3 has raw terms, native formulas, ordinary substitution, finite lists of reusable assumptions as contexts, and exactly the six rules Hyp, QE, DE, WI (Theorem 4.1) and

$$\frac{}{\Gamma \vdash s = s}\ (\textsc{Refl}), \qquad \frac{\Gamma \vdash s = t \qquad \Gamma \vdash b = r}{\Gamma \vdash r = \Phi(s, t, b)}\ (\textsc{Eval}),$$

the latter whenever $\Phi(s, t, b)$ is defined. The three rows of Φ give the three cases of Eval:

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash s = r}{\Gamma \vdash r = t}\ (\textsc{Root})$$

$$\frac{\Gamma \vdash a = t \qquad \Gamma \vdash I(0, a;\, u.\, h) = r}{\Gamma \vdash r = t}\ (\textsc{Eval}_0)$$

$$\frac{\Gamma \vdash I(c, a;\, u.\, h) = t \qquad \Gamma \vdash I(Sc, a;\, u.\, h) = r}{\Gamma \vdash r = h[t/u]}\ (\textsc{Eval}_S)$$

The calculus E_2 is E_3 without the case Root. In DE the eigenvariable is fresh for the context, the conclusion and the parameters of the matrix. No other logical, equality, successor, computation, induction or structural rule is primitive.

The conclusion of Eval is crossed: its left side is the right side of the second premise. The rule takes no template, no occurrence path, no counter and no body as a parameter; these are read off the two premises, which determine the conclusion or the failure of the rule.

Remark 4.10 (Counting). E_3 has six rules if Eval counts once and eight if its cases count separately; E_2 has six and seven. The Lean formalization `EPrf` has the two cases of E_2 as constructors (`evalZero`, `evalSucc`) and Root as the derived rule `EPrf.evalId`. The calculus U of Section 4.7 has one case and six rules under either convention. The number of arithmetic and equality schemas, including reflexivity, is four for E_3, three for E_2 and two for U.

4.2.2 The identity frame

Lemma 4.11 (Identity frame)**.** *Suppose that a calculus has reflexivity and, for every term s, a term $K(s)$ such that the inference*

$$\frac{s = t \qquad K(s) = r}{r = t}$$

is a rule. Then Root is derivable, by two instances of that rule and one reflexivity. The equation $K(s) = s$ is not needed as an axiom.

Proof. Let p derive $s = t$ and q derive $s = r$. The rule applied to q and to the reflexivity $K(s) = K(s)$ gives $K(s) = r$. The rule applied to p and to $K(s) = r$ gives $r = t$. Each of p and q is used once. □

Theorem 4.12 (ROOT is redundant)**.** *For every context* Γ *and native formula* A*,*

$$\Gamma \vdash_{\mathsf{E}_2} A \quad \Longleftrightarrow \quad \Gamma \vdash_{\mathsf{E}_3} A.$$

The translations leave every term and formula of the sequent unchanged. A derivation of E_3 *with* N *inference nodes,* k *of which are instances of* ROOT*, is translated to a derivation of* E_2 *with exactly* $N + 2k$ *nodes.*

Proof. From right to left, apply Theorem 4.11 with $K(s) := I(0, s;\, u.\, 0)$: its rule is EVAL_0 with the body 0. Explicitly, an instance of ROOT with premises $p\colon s = t$ and $q\colon s = r$ is replaced by

$$\mathrm{RC}(p, q) := \mathrm{Ev}_0(p,\ \mathrm{Ev}_0(q,\ \mathrm{R}_{K(s)})),$$

where R_w is the reflexivity leaf $w = w$ and Ev_0 is EVAL_0 with its premises in the displayed order. The replacement is carried out recursively; every other inference is copied. Each replaced node becomes two instances of EVAL_0 and one reflexivity, and its premises occur once, which gives the count. From left to right the translation is the inclusion. □

The count is of inference nodes. The added reflexivity leaf contains the term $K(s)$, whose size depends on s, so the theorem bounds neither the size of a serialized derivation nor the time of checking it.

4.2.3 Computation, equality and transport recovered

The derivations of this subsection are valid in E_2 over an arbitrary context and use only REFL, EVAL_0 and EVAL_S, apart from the given premises. No binder rule, no numerical case distinction and no induction is used. Write $\mathrm{Ev}(p, q)$ for EVAL with first premise p and second premise q.

Lemma 4.13 (Equality rules)**.** *Symmetry and transitivity are derivable in* E_2*: for* $p\colon s = t$ *put* $\mathrm{Sym}(p) := \mathrm{RC}(\mathrm{R}_s, p)$*, which derives* $t = s$*; for* $p\colon s = t$ *and* $q\colon t = v$ *put* $\mathrm{Trans}(p, q) := \mathrm{RC}(q, \mathrm{Sym}(p))$*, which derives* $s = v$*.*

Proof. RC derives $r = t$ from $s = t$ and $s = r$ (Theorem 4.12). With $s = s$ and $s = t$ it derives $t = s$. With $t = v$ and $t = s$ it derives $s = v$. □

Lemma 4.14 (Both computation equations)**.** *For all raw* c, a, h *the equations*

$$I(0, a;\, u.\, h) = a, \qquad I(Sc, a;\, u.\, h) = h[I(c, a;\, u.\, h)/u]$$

have derivations in E_2 *with three nodes.*

Proof. Put $A := I(0, a;\, u.\, h)$, $C := I(c, a;\, u.\, h)$ and $B := I(Sc, a;\, u.\, h)$. Then $\mathrm{Ev}(\mathrm{R}_a, \mathrm{R}_A)$ is an instance of EVAL_0 and concludes $A = a$, and $\mathrm{Ev}(\mathrm{R}_C, \mathrm{R}_B)$ is an instance of EVAL_S and concludes $B = h[C/u]$. The second derivation is the same when $c \equiv 0$. □

Theorem 4.15 (Congruence in every template)**.** *For every template* $h(u)$ *the rule*

$$\frac{\Gamma \vdash s = t}{\Gamma \vdash h[s/u] = h[t/u]}$$

is admissible in E_2*. The derivation adds eight inference nodes to the derivation of* $s = t$*, which occurs once.*

Proof. Let p derive $s = t$ and put $A_s := I(0, s;\, u.\, h)$ and $B_s := I(1, s;\, u.\, h)$. Four instances of Eval give

$$\begin{aligned} z_s &:= \mathrm{Ev}(\mathrm{R}_s, \mathrm{R}_{A_s}) && : A_s = s, \\ c_s &:= \mathrm{Ev}(z_s, \mathrm{R}_{B_s}) && : B_s = h[s/u], \\ z_{s,t} &:= \mathrm{Ev}(p, \mathrm{R}_{A_s}) && : A_s = t, \\ \mathrm{Cong}_h(p) &:= \mathrm{Ev}(z_{s,t}, c_s) && : h[s/u] = h[t/u]. \end{aligned}$$

The first and third lines are instances of $\textsc{Eval}_0$. In the second and fourth lines the second premise has the left side B_s, whose preceding iterate is literally A_s, the left side of the first premise; these are instances of $\textsc{Eval}_S$. The substitutions into h are the ordinary substitutions performed by Eval itself, so no congruence is assumed inside the derivation. The tree has four reflexivity leaves, of which R_{A_s} occurs twice, and four instances of Eval. □

Corollary 4.16 (Transport)**.** *The rule Tr, from $s = t$ and $h[s/u] = r$ infer $r = h[t/u]$, is admissible in E_2 for all templates and all terms s, including the literal 0. In E_3 the derivation adds nine inference nodes to the derivations of the premises; in E_2 it adds eleven. Each premise occurs once.*

Proof. Apply Root to $\mathrm{Cong}_h(p)$ and the second premise. In E_2 the instance of Root is the three nodes of RC. □

Remark 4.17. `equality_from_iteration` derives transport in E_2 by a different composition: from $C_0 := I(0, s;\, u.\, h)$ and $C_1 := I(1, s;\, u.\, h)$ it forms $C_0 = t$ and $C_1 = r$ by Theorems 4.13 and 4.14 and concludes with one instance of $\textsc{Eval}_S$. The template h occurs in these derivations as the body of an iterator; it is not a parameter of a primitive rule. The node counts say nothing about the size of the term labels, which contain copies of h, s and t.

4.2.4 Equivalence with the core

Theorem 4.18 (Equivalence)**.** *For every finite context Γ of native formulas and every native formula A,*

$$\Gamma \vdash_{\mathsf{E}_3} A \quad\Longleftrightarrow\quad \Gamma \vdash_{\mathsf{E}_2} A \quad\Longleftrightarrow\quad \Gamma \vdash_{\mathsf{N}} A.$$

The translations are effective on finite derivations and leave every term, formula and assumption of the sequent unchanged.

Proof. The first equivalence is Theorem 4.12. For the second it suffices, by Theorem 2.17, to compare E_2 and E_3 with the calculus W of Theorem 2.15, which has the rules Hyp, QE, DE and WI in common with them.

From W to E_2: I0 and IS are derived by Theorem 4.14 and Tr by Theorem 4.16; every other inference is copied, and each instance of WI keeps its witness and its mode.

From E_3 to W: reflexivity and the equality rules are derivable in W (Theorem 2.4(iii)). For an instance of Eval with premises $s = t$ and $b = r$ distinguish the rows of Φ. If $b \equiv s$, Tr with the identity template gives $r = t$. If $b \equiv I(0, s;\, u.\, h)$, then I0 gives $b = s$, and the equality rules give $r = t$ from $b = s$, $s = t$ and $b = r$. If $b \equiv I(Sc, a;\, u.\, h)$ and $s \equiv I(c, a;\, u.\, h)$, then IS gives $b = h[s/u]$; with $b = r$ this gives $h[s/u] = r$, and Tr with the template h and the major premise $s = t$ gives $r = h[t/u]$. Neither translation uses a structural rule or a substitution of terms in derivations. □

Remark 4.19. This is the proof of `computational_equality`, which states the theorem for E_3 and cites `wprf_iff_nprf` for the step from W to N; `equality_from_iteration` adds E_2. The Lean statements are `eprf_iff_nprf` and `eprf_iff_iprf`. Weakening, injective renaming and

cut are admissible in E_3 and E_2 also by the direct induction of Theorem 2.3: the operation Φ is a match of literal shapes, which an injective renaming preserves and a cut does not touch.

Corollary 4.20 (Linear bound on inference nodes)**.** *A derivation with N inference nodes in W is translated to a derivation of E_3 with at most $9N$ nodes.*

Proof. Each leaf I0 or IS becomes three nodes (Theorem 4.14). Each instance of Tr becomes the translations of its premises, each once, and nine further nodes (Theorem 4.16). Every other node is copied. Induction on the derivation gives the bound. □

The bound holds for this translation. It is not a bound on the size of formulas or of serialized derivations, and it is not a statement of optimality.

4.2.5 Substitution

Proposition 4.21 (Eval is stable under substitution)**.** *If $\Phi(s, t, b)$ is defined, then for every capture-free substitution σ the term $\Phi(s[\sigma], t[\sigma], b[\sigma])$ is defined by the same row and*

$$\Phi(s[\sigma], t[\sigma], b[\sigma]) \equiv \Phi(s, t, b)[\sigma].$$

Proof. The first row is immediate. In the second row substitution keeps the outer iterator with the literal counter 0, and its initializer is the substituted source. In the third row $b[\sigma] \equiv I(S(c[\sigma]), a[\sigma];\, u.\, h[\sigma])$ and $s[\sigma] \equiv I(c[\sigma], a[\sigma];\, u.\, h[\sigma])$, where σ acts on the parameters of h and leaves u bound; the composition law of ordinary substitution identifies $(h[t/u])[\sigma]$ with $h[\sigma][t[\sigma]/u]$. No computation takes place: if $c[\sigma]$ is the literal 0, the preceding iterate is still the raw term $I(0, a[\sigma];\, u.\, h[\sigma])$. The size comparison after Theorem 4.8 excludes a collapse into the first row. □

The proposition concerns Refl and Eval. The calculus E_3 keeps WI, so Theorem 4.4 applies: substitution of terms in derivations of E_3, the Leibniz rule, numerical cases and equation induction are admissible by Theorem 4.18 and Theorem 2.2, and Eval adds no repair of its own.

4.2.6 Irredundancy

Proposition 4.22 (Neither remaining case can be deleted)**.** *The calculus E_2 without Eval_0 does not derive $I(0, 0;\, u.\, 0) = 0$. The calculus E_2 without Eval_S does not derive $I(1, 0;\, u.\, Su) = S(I(0, 0;\, u.\, Su))$.*

Proof. Interpret the formulas in the standard natural numbers, with the standard 0 and S, equality as equality and the readings of Section 1.2 for Q_y and $D_{x,y}$. Under any interpretation of the iterator as a function of its counter, its initializer and its parameters, Hyp, Refl, QE and DE are sound; WI is sound in progress mode by induction on the standard numbers and in case mode by the distinction between 0 and successors.

Without Eval_0: interpret $I(n, a;\, u.\, h)$ as the n-fold iterate of h applied to $a + 1$. Then $I(Sc, a;\, u.\, h)$ is h applied to $I(c, a;\, u.\, h)$, so Eval_S is sound, while $I(0, 0;\, u.\, 0)$ has the value 1.

Without Eval_S: interpret $I(n, a;\, u.\, h)$ as a. Eval_0 is sound, and the displayed successor equation reads $0 = 1$. □

Proposition 4.23 (Reflexivity cannot be deleted)**.** *In E_2 without Refl, no literal equation is derivable from a context that contains no literal equation. In particular no equation is derivable from the empty context.*

Proof. By induction on a derivation whose context contains no literal equation and whose conclusion is one. Hyp cannot be the last rule. Either case of Eval has a literal equation as a premise in the same context. An instance of QE whose conclusion $B(v)$ is a literal equation has a matrix that is a literal equation, so its second premise $B(0)$ is a literal equation in the same context. DE adds an assumption with root Q_y, so the context of its second premise still contains no literal equation and its conclusion is the same equation. WI concludes a formula with root $D_{x,y}$. Each case contradicts the induction hypothesis. □

The two propositions concern the deletion of these rules from this inventory. They exclude neither another term language nor another rule basis.

4.3 Normalizers and the transfer of derivations

This section states once the comparison argument that `zero_normal`, `initializer_equality`, `full_normalization` and `computation_normal` each carry out for their own normalizer, and on which `equality_from_iteration`, `uniform_step`, `seed_evaluation` and `reflexive_base` rely by citation. The argument uses five properties of the normalizer and nothing else.

4.3.1 Admissible normalizers and normalizing substitution

Definition 4.24 (Admissible normalizer)**.** An *admissible normalizer* is a map ν from raw terms to raw terms such that for all raw terms t, all renamings ρ of variables and all capture-free simultaneous substitutions σ:

(ν1) $\nu(x) \equiv x$, $\nu(0) \equiv 0$ and $\nu(St) \equiv S\nu(t)$;

(ν2) $\nu(t[\rho]) \equiv \nu(t)[\rho]$ and $\mathrm{FV}(\nu(t)) \subseteq \mathrm{FV}(t)$;

(ν3) $\nu(\nu(t)) \equiv \nu(t)$;

(ν4) $\nu(t[\sigma]) \equiv \nu\big(\nu(t)[\nu \circ \sigma]\big)$;

(ν5) $\vdash_{\mathsf{N}} t = \nu(t)$, with the free variables of t as parameters.

A term t is *ν-normal* if $\nu(t) \equiv t$. The map ν is extended to formulas by $\nu(s = t) := (\nu(s) = \nu(t))$, $\nu(Q_y B) := Q_y \nu(B)$ and $\nu(D_{x,y} B) := D_{x,y} \nu(B)$, and to contexts entry by entry. A formula is ν-normal if $\nu(A) \equiv A$.

Condition (ν4) is the *coherence of normalization with substitution*: the normal form of an instance depends only on the normal forms of the term and of the substituted terms. Condition (ν5) says that ν is the identity inside N; it is a statement about each fixed term and does not say that N or iPRA proves a uniform theorem about ν. By (ν2) the extension to formulas does not depend on the names of bound variables, and ν never changes the root of a formula; in particular $\nu(A)$ is a literal equation exactly when A is.

Definition 4.25 (Normalizing substitution)**.** For a raw term or formula T and a substitution σ,

$$T\langle\sigma\rangle := \nu(T[\sigma]).$$

For a template h and a matrix B of ν-normal syntax, the instances $h(s)$, $B(r)$ and $B(t, r)$ are formed with this substitution.

Lemma 4.26 (Laws of normalizing substitution)**.** *Let ν be an admissible normalizer, T a raw term or formula, and σ, τ substitutions.*

(a) $\nu(\nu(T)[\sigma]) \equiv \nu(T[\sigma])$.

(b) $(T\langle\sigma\rangle)\langle\tau\rangle \equiv T\langle x \mapsto \sigma(x)\langle\tau\rangle\rangle \equiv \nu(T[\sigma][\tau])$.

(c) If T is ν-normal, then $T\langle\rho\rangle \equiv T[\rho]$ for every renaming ρ; in particular the identity substitution leaves T unchanged.

In particular instances compose: for a matrix $B(x, y)$ of ν-normal syntax, the instance at t of the template $B(v, r)$ is $B(t, r)$.

Proof. For terms, (a) follows from (ν4) applied to $\nu(T)$, from (ν3) and from (ν4) applied to T: $\nu(\nu(T)[\sigma]) \equiv \nu(\nu(\nu(T))[\nu \circ \sigma]) \equiv \nu(\nu(T)[\nu \circ \sigma]) \equiv \nu(T[\sigma])$. For (b), by (a), $(T\langle\sigma\rangle)\langle\tau\rangle \equiv \nu(\nu(T[\sigma])[\tau]) \equiv \nu(T[\sigma][\tau])$. Let $\theta(x) := \sigma(x)[\tau]$, so that $T[\sigma][\tau] \equiv T[\theta]$ by the composition law of ordinary substitution. By (ν4), (ν3) and (ν4) again, $\nu(T[\nu \circ \theta]) \equiv \nu(\nu(T)[\nu \circ \nu \circ \theta]) \equiv \nu(\nu(T)[\nu \circ \theta]) \equiv \nu(T[\theta])$, and the left side is $T\langle x \mapsto \sigma(x)\langle\tau\rangle\rangle$. (c) is ($\nu$2) with ($\nu$3). For formulas use induction on the formula: below a binder the substitution is lifted, and the lifting of $\nu \circ \sigma$ is ν composed with the lifting of σ by (ν1) and (ν2). □

Lemma 4.27 (A normalizer is the identity in the core)**.** *Let ν be an admissible normalizer. For every raw formula A and every context Γ, both $\Gamma, A \vdash_{\mathsf{N}} \nu(A)$ and $\Gamma, \nu(A) \vdash_{\mathsf{N}} A$. For every raw term t and substitution σ, $\vdash_{\mathsf{N}} t[\sigma] = t\langle\sigma\rangle$.*

Proof. The term statement is (ν5) for $t[\sigma]$. For formulas argue by induction on A. A literal equation uses (ν5) at its two sides and the equality rules. For $Q_y B$ and $D_{x,y} B$ the induction hypothesis holds for B with its bound variables free, in both directions, and Theorem 1.21 gives both directions for the formula. □

The inclusion of ν-normal syntax into raw syntax therefore commutes with substitution up to provable equality: the term $t\langle\sigma\rangle$ is by definition the normal form of $t[\sigma]$, the two terms need not be identical, and their equation is a theorem of N.

4.3.2 The kernel on normal forms

Definition 4.28 (The kernel N^ν)**.** Let ν be an admissible normalizer. The calculus N^ν has the ν-normal native formulas as formulas, finite lists of them as contexts, the substitution of Theorem 4.25, and the following rules, in which every term is ν-normal:

- Hyp, Tr, QE, DE, DI and QI, read with normalizing substitution;
- Comp$^\nu$: $\Gamma \vdash \nu(e)$ for every instance e of I0 or of IS on raw terms.

Lemma 4.29. *Weakening, cut on an assumption and normalizing substitution are admissible in N^ν: if $\Gamma \vdash_{\mathsf{N}^\nu} A$, then $\Gamma\langle\sigma\rangle \vdash_{\mathsf{N}^\nu} A\langle\sigma\rangle$ for every σ.*

Proof. By induction on derivations, with eigenvariables renamed away from σ. No rule of N^ν tests a term or a formula for a literal shape other than the root of the matrix in QI, which substitution preserves. By Theorem 4.26(b) the instance under σ of a premise or conclusion of Tr, QE, DE, DI or QI is the corresponding premise or conclusion of the same rule at the substituted parameters. For Comp$^\nu$, $\nu(e)\langle\sigma\rangle \equiv \nu(e[\sigma])$ by Theorem 4.26(a), and $e[\sigma]$ is an instance of the same computation equation. Weakening and cut are standard. □

Theorem 4.30 (Transfer)**.** *Let ν be an admissible normalizer. For every finite context Γ of raw native formulas and every raw native formula A,*

$$\Gamma \vdash_{\mathsf{N}} A \quad \Longleftrightarrow \quad \nu(\Gamma) \vdash_{\mathsf{N}^\nu} \nu(A).$$

Both directions are effective on finite derivations.

Proof. From left to right, by induction on the derivation in N, normalizing every formula. An assumption of Γ becomes an assumption of $\nu(\Gamma)$. A leaf I0 or IS with conclusion e becomes $\textsc{Comp}^\nu$ with conclusion $\nu(e)$. For $\textsc{Tr}$ with premises $s = t$ and $h[s/v] = r$, the induction hypotheses are $\nu(s) = \nu(t)$ and $\nu(h[s/v]) = \nu(r)$. By $(\nu 4)$, $\nu(h[s/v]) \equiv \nu(h)\langle\nu(s)/v\rangle$, so $\textsc{Tr}$ of N^ν with the template $\nu(h)$ gives $\nu(r) = \nu(h)\langle\nu(t)/v\rangle$, which is $\nu(r) = \nu(h[t/v])$ by $(\nu 4)$. In the same way $(\nu 4)$, with $(\nu 1)$ for the terms 0 and Sz, turns the premises and the conclusion of QE, DI and QI into those of the same rule at the matrix $\nu(B)$ and the parameters $\nu(t)$; the matrix of QI remains a literal equation. For DE the local assumption $Q_y B[b/x]$ becomes $Q_y \nu(B)[b/x]$ by $(\nu 2)$. Freshness conditions are preserved, because ν introduces no free variable by $(\nu 2)$.

From right to left, read a derivation of $\Delta \vdash B$ in N^ν as raw syntax and show $\Delta \vdash_{\mathsf{N}} B$ by induction. A leaf $\textsc{Comp}^\nu$ with conclusion $\nu(e)$: the equation e is an axiom of N, and Theorem 4.27 gives $\nu(e)$. For $\textsc{Tr}$ with premises $s = t$ and $h\langle s/v\rangle = r$: Theorem 4.27 gives $h[s/v] = h\langle s/v\rangle$, hence $h[s/v] = r$; $\textsc{Tr}$ of N gives $r = h[t/v]$, and Theorem 4.27 gives $r = h\langle t/v\rangle$. For QE convert the premise $B\langle 0/y\rangle$ to $B[0/y]$ by Theorem 4.27, apply QE of N, and convert $B[t/y]$ to $B\langle t/y\rangle$. For DE the local assumption is unchanged, since $B\langle b/x\rangle \equiv B[b/x]$ by Theorem 4.26(c). For DI work in the context $\Delta, B[t/x, 0/y]$: convert the assumption to $B\langle t/x, 0/y\rangle$, cut with the translated premise, convert the result to $B[t/x, Sz/y]$, and apply DI of N. QI is treated in the same way. The conversions are uniform in the parameters, so no eigenvariable enters a context or a conclusion. This gives $\nu(\Gamma) \vdash_{\mathsf{N}} \nu(A)$ from $\nu(\Gamma) \vdash_{\mathsf{N}^\nu} \nu(A)$; Theorem 4.27 and cut, applied to each assumption and to the conclusion, give $\Gamma \vdash_{\mathsf{N}} A$. □

A derivation with the single introduction must not be normalized rule by rule: a witness that is not literally 0 can have the normal form 0, and the mode of WI then changes. The theorem is therefore stated for the kernel with DI and QI, and each calculus with WI on normal forms is compared with N^ν afterwards.

Corollary 4.31 (Calculi equivalent to the kernel on normal forms)**.** *Let ν be an admissible normalizer and let K be a calculus on ν-normal native formulas that derives exactly the sequents of N^ν. Let J and H be the translations of formulas from N to iPRA and from iPRA to N of Theorem 1.78, and put $H_\nu := \nu \circ H$.*

(a) $\Gamma \vdash_{\mathsf{N}} A$ if and only if $\nu(\Gamma) \vdash_{\mathsf{K}} \nu(A)$, for raw Γ and A. For ν-normal Δ and B, $\Delta \vdash_{\mathsf{K}} B$ if and only if $\Delta \vdash_{\mathsf{N}} B$.

(b) Weakening, cut, normalizing substitution and the Leibniz rule, from $s = t$ and $F(s)$ infer $F(t)$ for every matrix F, are admissible in K.

(c) For ν-normal Δ, B and formulas Θ, φ of iPRA,

$$\begin{aligned} \Delta \vdash_{\mathsf{K}} B \quad &\iff \quad J\Delta \vdash_{\mathsf{iPRA}} JB, \\ \Theta \vdash_{\mathsf{iPRA}} \varphi \quad &\iff \quad H_\nu\Theta \vdash_{\mathsf{K}} H_\nu\varphi, \end{aligned}$$

the formulas B and $H_\nu JB$ are interderivable in K, and $\mathsf{iPRA} \vdash \varphi \leftrightarrow JH_\nu\varphi$. The numerical domains are whole, equality is interpreted by equality, and the numerical composites have the graph $x = y$. In particular $\vdash_{\mathsf{K}} 0 = S0$ if and only if $\mathsf{iPRA} \vdash \bot$.

For a calculus on raw terms that derives exactly the sequents of N, the same statements, with ν the identity, are Theorem 2.2.

Proof. (a) The first statement is Theorem 4.30. For the second, apply it to Δ and B, which are their own normal forms. (b) Weakening, cut and substitution are Theorem 4.29. For the Leibniz rule, N derives $F[t/v]$ from $s = t$ and $F[s/v]$ (Theorem 1.22), and normalizing this

sequent gives the rule on normal forms by (ν4). (c) The first equivalence is (a) composed with Theorem 1.78. For the second, $\Theta \vdash_{\mathsf{iPRA}} \varphi$ if and only if $H\Theta \vdash_{\mathsf{N}} H\varphi$, if and only if $\nu(H\Theta) \vdash_{\mathsf{K}} \nu(H\varphi)$ by (a). The formulas B and HJB are interderivable in N; normalizing the two sequents gives the interderivability of B and $H_\nu JB$ in K. By Theorem 4.27, $H\varphi$ and $\nu(H\varphi)$ are interderivable in N, hence $\mathsf{iPRA} \vdash JH\varphi \leftrightarrow JH_\nu\varphi$, and $\mathsf{iPRA} \vdash \varphi \leftrightarrow JH\varphi$. The map ν fixes the variables and changes neither a domain formula nor the interpretation of equality: normal forms are representatives of expressions, not a restriction or a quotient of the domain of numbers. The formula $0 = S0$ is ν-normal by (ν1). □

Every calculus on normal forms in the following sections is compared with N^ν on the same ν-normal sequents, and Theorem 4.31 is then not repeated. The normalizer and the normalizing substitution are part of the specification of such a calculus. A checker for it contains them, and the measurements of Section 4.11 count them: the deletion of a rule does not delete the cost of the computation that the rule expressed.

4.4 Zero-normal terms

This section draws on `zero_normal`, which deletes the zero computation rule by normalizing it in the syntax, and on the restatements of the same normalization layer in `initializer_equality`, `equality_from_iteration`, `uniform_step`, `seed_evaluation` and `reflexive_base`.

4.4.1 Zero normalization

Definition 4.32 (Zero normalization)**.** For terms c, a, h the *smart constructor* is

$$\overline{I}(c, a;\, u.\, h) := \begin{cases} a, & c \equiv 0, \\ I(c, a;\, u.\, h), & c \not\equiv 0. \end{cases}$$

The map nf_0 on raw terms is defined by structural recursion:

$$\mathrm{nf}_0(x) := x, \qquad \mathrm{nf}_0(0) := 0, \qquad \mathrm{nf}_0(St) := S\,\mathrm{nf}_0(t),$$
$$\mathrm{nf}_0(I(c, a;\, u.\, h)) := \overline{I}(\mathrm{nf}_0(c), \mathrm{nf}_0(a);\, u.\, \mathrm{nf}_0(h)).$$

A term is *zero-normal* if no iterator in it, at any depth, has the literal 0 as its counter.

The test in the last clause is on the *normalized* counter. For example $\mathrm{nf}_0(I(I(0, 0;\, v.\, 0), a;\, u.\, h)) \equiv \mathrm{nf}_0(a)$, although the outer counter is not literally 0; the equivalence of "$\mathrm{nf}_0(t) \equiv 0$" with "$t \equiv 0$" fails. Only the zero computation is normalized: $I(1, a;\, u.\, h)$ is zero-normal and is not identified with $h[a/u]$. In $I(c, a;\, u.\, h)$ the binder u covers h only, so the initializer returned by a contraction keeps its scope and no variable index is adjusted; for instance $\mathrm{nf}_0(I(n, a;\, u.\, I(0, u;\, v.\, 0))) \equiv I(n, a;\, u.\, u)$ for a variable n.

Lemma 4.33. *(a)* $\mathrm{nf}_0(t)$ *is zero-normal;* $\mathrm{nf}_0(t) \equiv t$ *if and only if* t *is zero-normal;* nf_0 *is idempotent.*

(b) nf_0 *commutes with renaming, and* $\mathrm{FV}(\mathrm{nf}_0(t)) \subseteq \mathrm{FV}(t)$.

(c) $\mathrm{nf}_0(t)$ *has at most as many nodes as* t*, and* nf_0 *is a primitive recursive operation on codes of finite terms.*

(d) $t \to^* \mathrm{nf}_0(t)$ *by zero contractions. Zero contraction, in any position and in any order, terminates, and* $\mathrm{nf}_0(t)$ *is its unique normal form.*

Proof. (a) By structural induction: the children of a rebuilt iterator are zero-normal, and the smart constructor returns either the zero-normal initializer or an iterator whose counter is not 0. On a zero-normal term every clause returns its argument. (b) A renaming sends no variable, successor or iterator to the literal 0, so it commutes with the test of the smart constructor; a contraction deletes subterms and creates no variable. (c) A contraction deletes an iterator node with its counter and body and keeps the normalized initializer; otherwise the normalized children are kept below the same constructor. A structural recursion with these size bounds is primitive recursive. It never performs a successor contraction: $I(Sn, a;\, u.\, h)$ is kept even for a closed numeral n, so no evaluator for arbitrary primitive recursive programs is involved. (d) The bottom-up computation of nf_0 is a sequence of zero contractions. A zero contraction decreases the number of nodes, hence terminates. If $t \to t'$ by a zero contraction, then $\mathrm{nf}_0(t) \equiv \mathrm{nf}_0(t')$, by induction on the position: at the root both sides are the normalized initializer; in a child, the normal form of the child is unchanged by the induction hypothesis, and the result is unchanged even if the child is erased. A term without a zero redex is zero-normal and fixed by nf_0, so every normal form of t is $\mathrm{nf}_0(t)$. □

Lemma 4.34 (Coherence with substitution)**.** *For every raw term t and every capture-free simultaneous substitution σ,*

$$\mathrm{nf}_0(t[\sigma]) \equiv \mathrm{nf}_0\big(\mathrm{nf}_0(t)[\mathrm{nf}_0 \circ \sigma]\big).$$

Proof. By induction on t. For a variable both sides are $\mathrm{nf}_0(\sigma(x))$, by idempotence. The cases 0 and St are immediate. Let $t \equiv I(c, a;\, u.\, h)$ and let σ' be the lifting of σ below u; by Theorem 4.33(b) the lifting of $\mathrm{nf}_0 \circ \sigma$ is $\mathrm{nf}_0 \circ \sigma'$. The left side is $\overline{I}(\mathrm{nf}_0(c[\sigma]), \mathrm{nf}_0(a[\sigma]); u.\, \mathrm{nf}_0(h[\sigma']))$. If $\mathrm{nf}_0(c) \equiv 0$, then $\mathrm{nf}_0(t) \equiv \mathrm{nf}_0(a)$ and the right side is $\mathrm{nf}_0(\mathrm{nf}_0(a)[\mathrm{nf}_0 \circ \sigma]) \equiv \mathrm{nf}_0(a[\sigma])$ by the induction hypothesis for a; the induction hypothesis for c gives $\mathrm{nf}_0(c[\sigma]) \equiv \mathrm{nf}_0(0[\mathrm{nf}_0 \circ \sigma]) \equiv 0$, so the left side is $\mathrm{nf}_0(a[\sigma])$ too. If $\mathrm{nf}_0(c) \not\equiv 0$, then $\mathrm{nf}_0(t) \equiv I(\mathrm{nf}_0(c), \mathrm{nf}_0(a);\, u.\, \mathrm{nf}_0(h))$ and the right side is the smart constructor applied to $\mathrm{nf}_0(\mathrm{nf}_0(c)[\mathrm{nf}_0 \circ \sigma])$, $\mathrm{nf}_0(\mathrm{nf}_0(a)[\mathrm{nf}_0 \circ \sigma])$ and $\mathrm{nf}_0(\mathrm{nf}_0(h)[\mathrm{nf}_0 \circ \sigma'])$, which by the induction hypotheses are the three arguments of the smart constructor on the left side. The substituted counter may or may not be 0; both sides apply the same test to the same term. □

Proposition 4.35. nf_0 *is an admissible normalizer, and the* nf_0*-normal terms are the zero-normal terms.*

Proof. (ν1) holds by definition, (ν2) and (ν3) by Theorem 4.33, (ν4) by Theorem 4.34, and (ν5) by Theorem 4.33(d) and Theorem 4.6. □

In the rest of this section and in Sections 4.5 to 4.8, terms and formulas are zero-normal and $t\langle\sigma\rangle$ is $\mathrm{nf}_0(t[\sigma])$. The substitution is computed by rebuilding the term bottom-up with the smart constructor: in an iterator the substitution acts on the counter and the initializer in the current scope and, lifted, on the body. A substitution into a zero-normal term can create a counter 0, which is then contracted. Neither the normalizer nor any rule tests whether a term *denotes* 0.

4.4.2 The kernel without the zero computation rule

Definition 4.36 (The kernel N^0)**.** The calculus N^0 has zero-normal native formulas, finite lists of them as contexts, normalizing substitution, and the seven rules Hyp, Tr, QE, DE, DI, QI and

$$\frac{}{\Gamma \vdash I(Sc, a;\, u.\, h) = h\langle \overline{I}(c, a;\, u.\, h)/u\rangle} \qquad \text{(IS)}$$

for all zero-normal c, a, h. The left side is zero-normal because Sc is not the literal 0; at $c \equiv 0$ the right side is $h\langle a/u\rangle$.

Lemma 4.37. N^0 *derives* $r = r$ *for every zero-normal* r*, and* N^0 *derives exactly the sequents of* $\mathsf{N}^{\mathrm{nf}_0}$*.*

Proof. Put $J_r := I(1, r;\, u.\, u)$. IS gives $J_r = r$, because the right side is $u\langle \overline{I}(0, r; u.u)/u\rangle \equiv r$, and Theorem 4.2(a) gives $r = r$. For the second statement compare IS with $\textsc{Comp}^{\mathrm{nf}_0}$. If e is the instance $I(0, a;\, u.\, h) = a$ of I0, then $\mathrm{nf}_0(e)$ is $\mathrm{nf}_0(a) = \mathrm{nf}_0(a)$, which N^0 derives. If e is an instance of IS at raw c, a, h, then by Theorem 4.34 $\mathrm{nf}_0(e)$ is the instance of IS of N^0 at $\mathrm{nf}_0(c), \mathrm{nf}_0(a), \mathrm{nf}_0(h)$. Conversely an instance of IS of N^0 at zero-normal c, a, h is nf_0 of the raw instance at the same terms. The other rules are common. □

Theorem 4.38 (The zero computation rule is deleted)**.** *For every finite context* Γ *of raw native formulas and every raw native formula* A*,*

$$\Gamma \vdash_{\mathsf{N}} A \quad\Longleftrightarrow\quad \mathrm{nf}_0(\Gamma) \vdash_{\mathsf{N}^0} \mathrm{nf}_0(A).$$

N^0 *is closed under weakening, cut and normalizing substitution, and Theorem 4.31 applies to it.*

Proof. Theorem 4.35, Theorem 4.30 and Theorems 4.29 and 4.37. □

Remark 4.39. The theorem is the faithfulness corollary of `zero_normal` for its seven-rule calculus and the raw/canonical equivalence of `initializer_equality`, whose comparison calculus lists reflexivity as an eighth rule; `equality_from_iteration`, `uniform_step`, `seed_evaluation` and `reflexive_base` cite it. The normalization layer of `zero_normal` is the Lean module `ZeroNormalization`, with `nf_subst`, `hsub_comp`, `nf_correct` and `inclusion_subst_coherent`; the theorem is `nzprf_iff_nprf` for the calculus `NZPrf`, with `NZPrf.normal` and `nzprf_iff_pra`. The Lean calculus admits raw rule parameters and proves as a theorem that it derives only zero-normal formulas from zero-normal assumptions; here this holds by the definition of the syntax. Definitional zero computation is proposed earlier in the package `prevalidated_elimination`, where it does not reduce the checker because the arity coding places the initializer below a dummy binder and a contraction must lower indices, and the package `uniform_zero` asks whether another layout of binders makes it cheaper; `zero_normal` carries it out for the present layout, in which the initializer lies outside the binder.

4.4.3 Six rules with the single introduction

Lemma 4.40 (Zero-normal aliases of zero)**.** *The closed terms* $I(1, 0;\, u.\, 0)$ *and* $I(1, 0;\, u.\, u)$ *are zero-normal and are not literally* 0*, and IS of* N^0 *derives* $I(1, 0;\, u.\, 0) = 0$ *and* $I(1, 0;\, u.\, u) = 0$*.*

Proof. The counter $S0$ is not 0, so neither term is contracted. The right sides of the two instances of IS are $0\langle \overline{I}(0, 0; u.0)/u\rangle \equiv 0$ and $u\langle \overline{I}(0, 0; u.u)/u\rangle \equiv 0$. □

The raw alias $I(0, 0;\, u.\, 0)$ of Theorem 4.18 is not available: it normalizes to 0 and selects the progress mode. The packages use $I(1, 0;\, u.\, 0)$ (`zero_normal`, `initializer_equality`, `equality_from_iteration`) or $I(1, 0;\, u.\, u)$ (`uniform_step`, `seed_evaluation`); the only property used is that the term is provably 0 and not literally 0.

Definition 4.41 (The calculus Z_6)**.** Z_6 has the syntax and the substitution of N^0 and the six rules Hyp, IS, Tr, QE, DE and WI; the test $t \equiv 0$ of WI is applied to the zero-normal witness as supplied. There is no primitive reflexivity, symmetry, substitution, cut, induction, coverage or separation rule.

Theorem 4.42. Z_6 *and* N^0 *derive the same sequents. Hence* $\Gamma \vdash_{\mathsf{N}} A$ *if and only if* $\mathrm{nf}_0(\Gamma) \vdash_{\mathsf{Z}_6} \mathrm{nf}_0(A)$*.*

Proof. Weakening, injective renaming and cut are admissible in Z_6 by direct induction on derivations: they preserve the root of a matrix and the literal-zero status of a witness, and cut substitutes no term. Reflexivity is derived as in Theorem 4.37, and Theorem 4.2(b) gives symmetry and transport in literal equations; the instances match by Theorem 4.26. By Theorem 4.3(i), QI is admissible in Z_6; by Theorem 4.3(ii) with the alias $Z := I(1, 0; u.\, 0)$ of Theorem 4.40, so is the one instance of DI that is not an instance of WI. Conversely WI is admissible in N^0 by Theorem 4.3(iii). The last statement is Theorem 4.38. □

Corollary 4.43 (Equation induction)**.** *For a literal equation E the rule that infers $\Gamma \vdash E(t)$ from $\Gamma \vdash E(0)$ and $\Gamma, E(z) \vdash E(Sz)$, with z fresh, is admissible in Z_6.*

Proof. QI, admissible by Theorem 4.42, and QE. □

Remark 4.44. `zero_normal` states Theorem 4.42 with a seven-rule bridge, the calculus N^0, and warns that a derivation of the calculus W of Theorem 2.15 on raw terms must not be normalized with its modes kept. The six-rule calculus is not formalized in Lean in this form: the formalized six-rule calculus on zero-normal terms is `NRPrf` of Section 4.5, with a different mode test, and the calculus U of Section 4.7, with the present test.

4.5 The introduction selected by a reflexive base

This section draws on `reflexive_base`. The single introduction is kept, and its mode is selected by the formula that the case mode would assume, in place of the pair of tests on the root of the matrix and on the witness. No alias of zero is needed.

4.5.1 The calculus

Definition 4.45 (The calculus R)**.** For a zero-normal matrix $B(x, y)$ and a zero-normal term t, let $\mathsf{r}(B, t)$ hold if the instance $B(t, 0)$ is literally $r = r$ for some term r, and put

$$\eta^{\mathsf{r}}_{B,t}(z) := \begin{cases} z, & \mathsf{r}(B,t), \\ 0, & \text{otherwise.} \end{cases}$$

The rule RI is

$$\frac{\Gamma,\ B(t, \eta^{\mathsf{r}}_{B,t}(z)) \vdash B(t, Sz)}{\Gamma \vdash D_{x,y}B(x, y)} \qquad (z \text{ fresh for } \Gamma,\ t \text{ and the parameters of } B).$$

The calculus R has the syntax and the substitution of N^0 and the six rules Hyp, IS, Tr, QE, DE and RI.

The predicate r compares two finite trees. Since the instance $B(t, 0)$ has the root of B, the condition $\mathsf{r}(B, t)$ implies that B is a literal equation. The instance $B(t, 0)$ is normalized, so the predicate depends on the zero normalization of the calculus.

4.5.2 The structural rules, without circularity

The lemmas are proved in the following order: laws of normalizing substitution; weakening and cut; equality; unrestricted introduction by cases; substitution. None of them presupposes the equivalence theorem.

Lemma 4.46. *Weakening, injective renaming of free variables and cut on an assumption are admissible in R. Reflexivity, symmetry, transitivity, congruence and transport in literal equations are derivable in R from IS and Tr.*

Proof. The first three operations leave the matrix and the witness of each instance of RI unchanged or apply an injective renaming to them. A reflexive equation stays reflexive, and an equation that is not reflexive does not become reflexive under an injective renaming, so no mode changes. The equality rules are those of Theorems 4.2 and 4.37; their derivations use no introduction rule. □

In particular every base selected by r is derivable in every context.

Lemma 4.47 (Every introduction by cases is admissible)**.** *For every zero-normal matrix B and witness t, if $\Gamma, B(t,0) \vdash_{\mathsf{R}} B(t,Sz)$ with z fresh, then $\Gamma \vdash_{\mathsf{R}} D_{x,y}B$.*

Proof. If $\mathsf{r}(B,t)$ fails, this is RI. If it holds, the assumption $B(t,0)$ is literally $r = r$; derive it by reflexivity and remove it by cut. Weaken the resulting derivation of $B(t,Sz)$ by the assumption $B(t,z)$ and apply RI in progress mode. □

Lemma 4.48 (Substitution)**.** *Normalizing substitution is admissible in* R.

Proof. If $B(t,0)$ is literally $r = r$, then for every σ its instance is $r\langle\sigma\rangle = r\langle\sigma\rangle$, and by Theorem 4.26 this is the zero instance of the substituted matrix at the substituted witness. Hence a true guard remains true. Now argue by induction on a derivation. IS, Tr, QE and DE commute with normalizing substitution by Theorem 4.26, with fresh eigenvariables. For an instance of RI in progress mode the guard remains true and the induction hypothesis is the premise of RI in progress mode at the substituted data. For an instance in case mode the induction hypothesis is the substituted case premise, and Theorem 4.47 applies whether or not the guard has become true. □

The guard is preserved by substitution and not reflected: if the zero instance of an equation is $p = q$ for distinct variables p, q, the guard fails, and it holds after the substitution of 0 for both. A second way in which a false guard becomes true is the contraction of a counter that the substitution turns into 0. The proof repairs exactly this change of mode.

4.5.3 Progress recovered by two decoders

By Theorem 4.47, DE and the construction of Theorem 4.3(i), the closure by cases is admissible in R:

$$\frac{\Lambda,\ F(0) \vdash F(Sz)}{\Lambda \vdash Q_y F(y)} \qquad (z \text{ fresh for } \Lambda \text{ and the parameters of } F). \tag{4.1}$$

Recall the selector $\mathrm{sel}(v,a,b) = I(v,a;\,u.\,b)$, with u not in b. In zero-normal syntax the instance $\mathrm{sel}(0,a,b)$ is a, and IS gives $\mathrm{sel}(Sv,a,b) = b$.

Lemma 4.49 (Case decoder)**.** *Let $r(y)$ be a zero-normal template and a a term without y, and let $\mathrm{Dec}_{r,a}(y)$ be the equation*

$$\mathrm{sel}(y,\ a,\ \mathrm{sel}(y,0,r(y))) = r(y).$$

Then $\vdash_{\mathsf{R}} Q_y\mathrm{Dec}_{r,a}(y)$. Consequently $\Lambda \vdash_{\mathsf{R}} a = r(0)$ implies $\Lambda \vdash_{\mathsf{R}} \mathrm{Dec}_{r,a}(v)$ for every term v.

Proof. The zero instance of $\mathrm{Dec}_{r,a}$ is literally $a = r(0)$. At $y = Sz$ two instances of IS and transitivity give

$$\mathrm{sel}(Sz,a,\mathrm{sel}(Sz,0,r(Sz))) = \mathrm{sel}(Sz,0,r(Sz)) = r(Sz),$$

a derivation that does not use the assumption of the base. Rule (4.1) gives $Q_y\mathrm{Dec}_{r,a}(y)$ in the empty context. Weaken into Λ and apply QE with the given base and the target v. □

The lemma is a numerical case distinction in a fixed context: Λ does not contain the generalized variable, and the conclusion can be instantiated at a variable that occurs in Λ, because QE imposes no freshness condition on its target.

Theorem 4.50 (Equation progress)**.** *Let $E(y)$ be the zero-normal literal equation $s(y) = t(y)$. If $\Gamma, E(z) \vdash_{\mathsf{R}} E(Sz)$ with z fresh for Γ and the parameters of E, then $\Gamma \vdash_{\mathsf{R}} Q_y E(y)$.*

Proof. Let π be the given derivation, write $s_0 := s(0)$ and $t_0 := t(0)$, and keep the base as an assumption: $\Delta := \Gamma, E(0)$, where $E(0)$ is $s_0 = t_0$. The *progressive lift* of E is

$$H(y) := \big(\mathrm{sel}(y, 0, s(y)) = \mathrm{sel}(y, 0, t(y))\big).$$

Its zero instance is literally $0 = 0$. Consider the two decoders

$$\mathrm{Dec}_{s,s_0}(y)\colon\ \ \mathrm{sel}\big(y, s_0, \mathrm{sel}(y, 0, s(y))\big) = s(y), \qquad \mathrm{Dec}_{t,s_0}(y)\colon\ \ \mathrm{sel}\big(y, s_0, \mathrm{sel}(y, 0, t(y))\big) = t(y).$$

The zero instance of the first is $s_0 = s_0$, derivable by reflexivity, and that of the second is $s_0 = t_0$, the assumption $E(0)$. By Theorem 4.49, $\Delta \vdash \mathrm{Dec}_{s,s_0}(z)$ and $\Delta \vdash \mathrm{Dec}_{t,s_0}(z)$ at the fresh variable z.

Add the assumption $H(z)$. Congruence in the template $w \mapsto \mathrm{sel}(z, s_0, w)$ turns it into

$$\mathrm{sel}\big(z, s_0, \mathrm{sel}(z, 0, s(z))\big) = \mathrm{sel}\big(z, s_0, \mathrm{sel}(z, 0, t(z))\big).$$

The symmetric form of $\mathrm{Dec}_{s,s_0}(z)$, this equation and $\mathrm{Dec}_{t,s_0}(z)$ give $s(z) = t(z)$, which is $E(z)$. Weaken π into the context $\Delta, H(z)$ and cut $E(z)$ into its assumption; the result is $\Delta, H(z) \vdash E(Sz)$. Nothing is substituted into π: its variable is still z. IS on each side of $H(Sz)$ converts $E(Sz)$ to $H(Sz)$, so

$$\Delta,\, H(z) \vdash H(Sz).$$

Regard $H(y)$ as a matrix with an unused witness coordinate. Its zero instance at any witness is $0 = 0$, so RI is in progress mode and gives $\Delta \vdash D_{x,y} H(y)$; DE returns $\Delta \vdash Q_y H(y)$, since the witness is unused. Reflexivity derives $H(0)$, QE at Sz gives $H(Sz)$, and IS gives $E(Sz)$. This is a derivation of $\Gamma, E(0) \vdash E(Sz)$, and rule (4.1) gives $\Gamma \vdash Q_y E(y)$. □

The proof uses equality, the closure by cases, weakening, renaming and cut. It uses no implication between formulas, no substitution of terms into the given derivation and not Theorem 4.48.

4.5.4 Equivalence

Theorem 4.51. *For every zero-normal context Γ and formula A,*

$$\Gamma \vdash_{\mathsf{R}} A \quad \Longleftrightarrow \quad \Gamma \vdash_{\mathsf{N}^0} A.$$

Both translations are effective and are the identity on the sequent. Hence $\Gamma \vdash_{\mathsf{N}} A$ if and only if $\mathrm{nf}_0(\Gamma) \vdash_{\mathsf{R}} \mathrm{nf}_0(A)$ for raw Γ, A.

Proof. From R to N^0: five rules are common. An instance of RI in case mode is an instance of DI. In progress mode the matrix is a literal equation; put $E(y) := B(t, y)$. QI gives $Q_y E(y)$; in the context $\Gamma, E(0)$, QE gives $E(Sz)$; DI at the witness t concludes $D_{x,y} B$. From N^0 to R: DI is admissible by Theorem 4.47 and QI by Theorem 4.50. The last statement is Theorem 4.38. □

Remark 4.52. The Lean formalization is `NRPrf`, with `nrprf_iff_nprf`, `NRPrf.normal` and `nrprf_iff_pra`; its completeness proof translates the calculus of the package `diagonal_guard`, with the invariant that a literally reflexive assumption may be dropped. The method of a guarded introduction is that of the package `diagonal_guard`; the guard on a computed zero instance, the proof of Theorem 4.48 for a guard that can change from false to true, and Theorem 4.53 are specific to `reflexive_base`. The rule that keeps both $B(t,0)$ and, at a literal equation, $B(t,z)$ as assumptions of one introduction is a different, older rule.

4.5.5 The same guard on raw terms, and two variants

Proposition 4.53 (Seven rules without normalization)**.** *Let* $\mathsf{R}^{\mathrm{raw}}$ *be the calculus on raw terms with ordinary substitution and the rules* Hyp*, I0, IS,* Tr*, QE, DE and RI, where* $\mathsf{r}(B,t)$ *tests the literal syntax of the raw instance* $B(t,0)$*. Then* $\mathsf{R}^{\mathrm{raw}}$ *derives exactly the sequents of* N*.*

Proof. Reflexivity follows from I0 and Theorem 4.2(a) applied to $I(0,r;\,u.\,u)=r$. Weakening, injective renaming and cut are direct inductions; Theorem 4.47 and rule (4.1) are obtained as before, and a true guard is preserved by ordinary substitution, so Theorem 4.48 holds with the same proof. The translation from $\mathsf{R}^{\mathrm{raw}}$ to N is that of Theorem 4.51.

Only the lift in the proof of progress changes, because the raw zero instance of H is not reflexive. For $E(y) \equiv (s(y)=t(y))$ use the asymmetric equation

$$K(y) := \big(\mathrm{sel}(y,t(y),s(y)) = \mathrm{sel}(0,t(y),s(y))\big).$$

Its raw zero instance is literally $\mathrm{sel}(0,t(0),s(0)) = \mathrm{sel}(0,t(0),s(0))$, so it satisfies the guard before any computation. For every v, IS computes the left side of $K(Sv)$ to $s(Sv)$ and I0 computes its right side to $t(Sv)$, so $K(Sv)$ and $E(Sv)$ are interderivable. In the context $\Gamma, E(0)$ the two decoder equations are

$$\mathrm{sel}\big(y,s(0),\mathrm{sel}(y,t(y),s(y))\big)=s(y), \qquad \mathrm{sel}\big(y,s(0),\mathrm{sel}(0,t(y),s(y))\big)=t(y).$$

Their zero instances follow from I0 and, respectively, reflexivity and the assumption $s(0)=t(0)$; their successor instances follow from IS for the outer selector and IS, respectively I0, for the inner one. The closure by cases and QE give both decoders at every term. Congruence of $K(z)$ in the template $w \mapsto \mathrm{sel}(z,s(0),w)$ and the decoders give $E(z)$. Cut it into the given derivation of $E(Sz)$, convert $E(Sz)$ to $K(Sz)$, and conclude with RI in progress mode, DE, QE and the closure by cases as in Theorem 4.50. □

The asymmetric lift also serves on zero-normal terms, where its base contracts to $t(0)=t(0)$. The guard is thus independent of the normalization of the zero computation. The Lean formalization of $\mathsf{R}^{\mathrm{raw}}$ is `RRPrf`, with `rrprf_iff_nprf`; its structural rules are weakening, injective renaming and cut.

Proposition 4.54 (Two variants of the guard)**.** *Theorem 4.51 remains true if* $\mathsf{r}(B,t)$ *is replaced by either of the following predicates:*

(a) $B(t,0)$ *is literally* $0=0$*;*

(b) $B(x,0)$*, with* x *still a variable, is literally* $r=r$*.*

Proof. The proofs of this section use three properties of the guard: a selected base is derivable; a true guard is preserved by substitution; and the lift H of Theorem 4.50 satisfies the guard. For (a) the base is $0=0$, which is preserved by substitution, and the zero instance of H is $0=0$. For (b), if $B(x,0)$ is $r=r$ then every $B(t,0)$ is reflexive, the guard is preserved by the substitution of the parameters, and the zero instance of H with an unused coordinate x is $0=0$. □

Variant (a) compares with one fixed formula and recognizes fewer matrices. Variant (b) removes the witness from the test and costs a checker one more normalizing substitution, since the instance $B(t, 0)$ is needed for the assumption of the case mode in any event. The proof of Theorem 4.53 does not establish variant (a) on raw terms, where the base of the asymmetric lift is reflexive and in general not $0 = 0$.

4.6 Equality as replacement of an initializer

This section draws on `initializer_equality`. Successor computation is kept as a rule, and transport in an arbitrary template is replaced by a rule that changes only the initializer of an iterator. Because an iterator with the counter 0 is its initializer, the same rule replaces at the root.

4.6.1 The rule Seed and the calculus

Definition 4.55 (Seed replacement)**.** For zero-normal terms s, t, b the partial operation $\rho_{s,t}(b)$ is

$$\rho_{s,t}(b) := \begin{cases} t, & b \equiv s, \\ I(n, t;\, u.\, h), & b \equiv I(n, s;\, u.\, h), \\ \text{undefined}, & \text{otherwise.} \end{cases}$$

The two branches are disjoint, since a finite term is not its own proper subterm. The second branch inspects one constructor and its initializer and leaves the counter and the body unchanged; its result is zero-normal because the counter is the same term, which is not 0. No bound variable is substituted or renamed.

Definition 4.56 (The calculus S)**.** S has the syntax and the substitution of N^0 and the six rules Hyp, IS (as in Theorem 4.36), QE, DE, WI and

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash b = r}{\Gamma \vdash \rho_{s,t}(b) = r} \quad (\rho_{s,t}(b) \text{ defined}) \qquad (\textsc{Seed}).$$

The conclusion of Seed is not crossed. The rule has no parameter: the two premises determine the conclusion or the failure of the rule. Substitution is not removed from the calculus, since IS and the binder rules use it.

Lemma 4.57 (Schematic form)**.** *Seed is exactly the schema*

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash \overline{I}(n, s;\, u.\, h) = r}{\Gamma \vdash \overline{I}(n, t;\, u.\, h) = r}$$

for arbitrary zero-normal n and h. Its instance at $n \equiv 0$ is the rule of root replacement*: from $s = t$ and $s = r$ infer $t = r$.*

Proof. At $n \equiv 0$ the schema is the first branch of ρ; the erased body is irrelevant. At any other n the second premise displays the iterator from which the counter and the body are read, and the schema is the second branch. Conversely the first branch is the schema at $n \equiv 0$, $h \equiv 0$, and the second branch is the schema at the counter and body of b. □

Lemma 4.58 (Seed is stable under substitution)**.** *If $\rho_{s,t}(b)$ is defined, then for every substitution σ the term $\rho_{s\langle\sigma\rangle,t\langle\sigma\rangle}(b\langle\sigma\rangle)$ is defined and equals $\rho_{s,t}(b)\langle\sigma\rangle$.*

Proof. In the root branch this is immediate. Let $b \equiv I(n, s;\, u.\, h)$. Then $b\langle\sigma\rangle$ is the smart constructor applied to the substituted counter, initializer and body. If the substituted counter is 0, both $b\langle\sigma\rangle$ and $\rho_{s,t}(b)\langle\sigma\rangle$ lose their iterator node and become $s\langle\sigma\rangle$ and $t\langle\sigma\rangle$, and the rule applies in its root branch. Otherwise both keep the node, with the same counter and body, and the rule applies in its second branch. The rule is stable although normalization can change the branch. □

4.6.2 Equality recovered

The derivations of this subsection hold in S over an arbitrary context and use only IS and Seed, apart from the given premises.

Lemma 4.59 (Root equality)**.** *Reflexivity, symmetry and transitivity are derivable in* S.

Proof. IS gives $I(1, a;\, u.\, u) = a$. Root replacement with this equation as both premises gives $a = a$. For $s = t$, root replacement with the second premise $s = s$ gives $t = s$. For $s = t$ and $t = r$, symmetry gives $t = s$, and root replacement with the premises $t = s$ and $t = r$ gives $s = r$. □

Lemma 4.60 (One-step application)**.** *For every zero-normal template $f(u)$ and term a, IS derives $I(1, a;\, u.\, f) = f\langle a/u\rangle$.*

Proof. The right side of IS at the counter 0 is $f\langle\overline{I}(0, a; u.f)/u\rangle \equiv f\langle a/u\rangle$. This is the point at which the definitional zero computation is used. □

Theorem 4.61 (Congruence in every template)**.** *For every zero-normal template $f(u)$ the rule that infers $f\langle s/u\rangle = f\langle t/u\rangle$ from $s = t$ is admissible in* S. *If the derivation of $s = t$ has P inference nodes, the derivation of the conclusion has $P + 4$ nodes, independently of the depth and the number of occurrences of u in f.*

Proof. Write $f(a)$ for $f\langle a/u\rangle$ and put $A_a := I(1, a;\, u.\, f)$. The derivation is

$$\dfrac{\dfrac{s = t \qquad A_s = f(s)}{A_t = f(s)} \qquad A_t = f(t)}{f(s) = f(t)}$$

The equations $A_s = f(s)$ and $A_t = f(t)$ are instances of IS (Theorem 4.60). The upper inference is Seed in its second branch, which replaces the initializer s of A_s by t. The lower inference is root replacement. The derivation adds two instances of IS and two of Seed; it uses neither reflexivity nor symmetry nor transitivity. The occurrences of u in f, including those below iterator binders, are handled by the substitution of IS. □

Corollary 4.62 (Transport)**.** *The following rules are admissible in* S*, for all terms and contexts: from $s = t$ and $f(s) = r$ infer $f(t) = r$; the rule* Tr*; and transport in a literal equation, from $s = t$ and $a(s) = b(s)$ infer $a(t) = b(t)$.*

Proof. Theorem 4.61 gives $f(s) = f(t)$, and root replacement with the second premise $f(s) = r$ gives $f(t) = r$. Symmetry gives Tr. Two uses of Tr, as in Theorem 4.2(b), give transport in an equation. □

Remark 4.63. The node count is for the displayed tree; its labels contain copies of f, s and t, so it bounds neither the size of a serialized derivation nor the time of checking. The package `witness_rule` has a congruence rule restricted to the counter position of a different recursor, with a separate root rule and a construction with $P+5$ nodes; here the lifting and the first root replacement are one inference, and the instance of the same rule at the counter 0 is the root replacement. The two term languages differ, so the comparison is not a uniform shortening of derivations.

4.6.3 Equivalence

Theorem 4.64. *For every zero-normal context* Γ *and formula* A*,*

$$\Gamma \vdash_{\mathsf{S}} A \quad \Longleftrightarrow \quad \Gamma \vdash_{\mathsf{N}^0} A.$$

Hence $\Gamma \vdash_{\mathsf{N}} A$ *if and only if* $\mathrm{nf}_0(\Gamma) \vdash_{\mathsf{S}} \mathrm{nf}_0(A)$ *for raw* Γ, A*.*

Proof. Weakening, injective renaming and cut are admissible in S by direct induction: they leave the terms of each rule instance unchanged up to an injective renaming, so neither the mode of WI nor the branch of Seed changes.

From N^0 to S: Tr is admissible by Theorem 4.62, which with Theorem 4.59 also supplies the symmetry and the transport in literal equations required by Theorem 4.3. By Theorem 4.3(i), QI is admissible, and by Theorem 4.3(ii) with the alias $Z := I(1, 0;\, u.\, 0)$, for which IS derives $Z = 0$, so is the instance of DI that WI lacks. The other rules are common.

From S to N^0: by Theorem 4.57 an instance of Seed has the premises $s = t$ and $\overline{I}(n, s; u.h) = r$. Tr of N^0 in the template $\overline{I}(n, v; u.h)$, which is v if $n \equiv 0$ and $I(n, v;\, u.\, h)$ otherwise, gives $r = \overline{I}(n, t; u.h)$, and symmetry gives the conclusion. WI is admissible in N^0 by Theorem 4.3(iii). The last statement is Theorem 4.38. □

By Theorem 4.31(b), substitution and the Leibniz rule are admissible in S. `initializer_equality` also proves the Leibniz rule directly, by induction on the matrix: at $Q_y B$ it introduces the closure by cases, transports the base backwards, applies the given closure and transports the result forwards; at $D_{x,y} B$ it opens the given formula and introduces the target by cases at the same witness. Only the literal-equation case uses Theorem 4.62.

Remark 4.65. The Lean formalization of the rule is `NIPrf`, with `niprf_iff_nprf`; it keeps DI and QI and therefore has seven rules. A successor-seeded canonical recursor, as in the package `canonical_recursors`, erases the body of a recursor at the counter 0 and keeps the node; it does not make the zero step the identity of the initializer, and the root replacement is then not an instance of replacement of an initializer.

4.7 One successor step

This section draws on `uniform_step`, which presents the rule Step with clause contexts, and on the companion calculus of `equality_from_iteration`, where the same rule is called UE and the contexts are lists of native formulas. On zero-normal terms the case $\textsc{Eval}_0$ of Theorem 4.9 is absorbed by the syntax, and reflexivity with the one remaining case gives all computation and all equality reasoning.

4.7.1 The rule Step

Definition 4.66 (The rule Step and the calculus U)**.** On zero-normal terms the rule Step is

$$\frac{\Gamma \vdash \overline{I}(c, a;\, u.\, h) = t \qquad \Gamma \vdash I(Sc, a;\, u.\, h) = r}{\Gamma \vdash r = h\langle t/u\rangle}.$$

The calculus U has the syntax and the substitution of N^0 and the six rules Hyp, Refl, Step, QE, DE and WI.

The second premise shows c, a and h: its left side is an iterator whose counter is a successor. The left side of the first premise is compared with the predecessor iterate built by the smart constructor; at $c \equiv 0$ it is a. The rule has one case, takes no template and compares no term with an arbitrary context. On raw terms the same rule is $\textsc{Eval}_S$.

Lemma 4.67 (Computation and equality from Refl and Step)**.** *The following are derivable from Refl and Step over an arbitrary context.*

(a) Successor computation: $I(Sc, a;\, u.\, h) = h\langle \overline{I}(c, a; u.h)/u\rangle$*, by one Step on two reflexivities.*

(b) The pivot: from $s = t$ *and* $s = r$ *infer* $r = t$*, by two instances of Step and one reflexivity.*

(c) Symmetry and transitivity.

(d) Congruence: from $s = t$ *infer* $h\langle s/v\rangle = h\langle t/v\rangle$*, by two instances of Step and two reflexivities.*

(e) Tr: from $s = t$ *and* $h\langle s/v\rangle = r$ *infer* $r = h\langle t/v\rangle$*, by four instances of Step and three reflexivities; and transport in a literal equation.*

The derivations (b)–(e) use Step at the counter $c \equiv 0$ *only.*

Proof. (a) Apply Step to the reflexivities of $\overline{I}(c, a; u.h)$ and of $I(Sc, a;\, u.\, h)$.

(b) Put $J_s := I(1, s;\, v.\, v)$; its predecessor iterate is $\overline{I}(0, s; v.v) \equiv s$. Step with the premises $s = r$ and $J_s = J_s$ gives $J_s = r$. Step with the premises $s = t$ and $J_s = r$ gives $r = t$:

$$\dfrac{s = t \qquad \dfrac{s = r \qquad J_s = J_s}{J_s = r}}{r = t}$$

This is Theorem 4.11 with the frame $K(s) := J_s$, which is not normalized away. The derivation is valid for every s, including 0.

(c) The pivot applied to $s = s$ and $s = t$ gives $t = s$. From $s = t$ and $t = r$, symmetry gives $t = s$, and the pivot applied to $t = r$ and $t = s$ gives $s = r$.

(d) Put $H_s := I(1, s;\, v.\, h)$. Step with the premises $s = s$ and $H_s = H_s$ gives $H_s = h\langle s/v\rangle$. Step with the premises $s = t$ and $H_s = h\langle s/v\rangle$ gives $h\langle s/v\rangle = h\langle t/v\rangle$:

$$\dfrac{s = t \qquad \dfrac{s = s \qquad H_s = H_s}{H_s = h\langle s/v\rangle}}{h\langle s/v\rangle = h\langle t/v\rangle}$$

The first result is reused as the second premise; it is not necessary to derive both $H_s = h\langle s/v\rangle$ and $H_s = h\langle t/v\rangle$ and to pivot them.

(e) The pivot applied to the conclusion of (d) and to $h\langle s/v\rangle = r$ gives $r = h\langle t/v\rangle$. Transport in a literal equation follows by Theorem 4.2(b). □

Remark 4.68. The short congruence (d) is that of `seed_evaluation` and of the package `clausal_single_evaluation`. `uniform_step` and `equality_from_iteration` derive both $H_s = h\langle s/v\rangle$ and $H_s = h\langle t/v\rangle$ and pivot them; their transport has six instances of Step and five reflexivities. None of the derivations uses equation induction, a numerical case distinction, the injectivity of the successor, a binder rule or the substitution of terms in derivations.

Proposition 4.69 (Step is stable under substitution)**.** *The instance under a normalizing substitution* σ *of the premises and the conclusion of an instance of Step are the premises and the conclusion of the instance of Step at the substituted counter, initializer and body.*

Proof. The counter of the second premise stays a successor. For zero-normal c, a, h the term $\overline{I}(c, a; u.h)$ is $\mathrm{nf}_0(I(c, a;\, u.\, h))$, so by Theorem 4.26(a) its instance under σ is the smart constructor applied to the substituted counter, initializer and body, including when the substituted counter is 0. By Theorem 4.26(b) the instance of $h\langle t/u\rangle$ is the instance of the substituted body at $t\langle\sigma\rangle$. □

Theorem 4.70. *For every zero-normal context* Γ *and formula* A*,*

$$\Gamma \vdash_{\mathsf{U}} A \quad \Longleftrightarrow \quad \Gamma \vdash_{\mathsf{N}^0} A.$$

Hence $\Gamma \vdash_{\mathsf{N}} A$ *if and only if* $\mathrm{nf}_0(\Gamma) \vdash_{\mathsf{U}} \mathrm{nf}_0(A)$ *for raw* Γ, A*. The equivalence holds also for the calculus with the seven rules* Hyp*,* Refl*,* Step*, QE, DE, DI and QI.*

Proof. Weakening, injective renaming and cut are admissible in U by direct induction, as in Theorem 4.42. From N^0 to U: IS and Tr are derived by Theorem 4.67. The term $Z := I(1, 0; u.\, 0)$ is not literally 0, and Step on the reflexivities of 0 and of Z derives $Z = 0$; Theorem 4.3(i) and (ii) give QI and DI. From U to N^0: reflexivity is Theorem 4.37. For an instance of Step, IS gives $I(Sc, a; u.\, h) = h\langle \overline{I}(c, a; u.h)/u\rangle$; congruence along the first premise gives $h\langle \overline{I}(c, a; u.h)/u\rangle = h\langle t/u\rangle$, and with the second premise the equality rules give $r = h\langle t/u\rangle$. WI is admissible by Theorem 4.3(iii). For the seven-rule calculus the arithmetic rules are translated in the same way and the introductions are common. □

Remark 4.71. U is the companion calculus of `equality_from_iteration`, formalized as `ZEPrf` with `zeprf_iff_nprf`; the missing introduction by cases is derived there with the alias $I(1, 0; u.\, 0)$, with no weakening and no cut. The seven-rule calculus is the formalization `NSPrf` of the arithmetic interface of `uniform_step`, with `nsprf_iff_nprf`, `NSPrf.normal` and `nsprf_iff_pra`. The theorem is an equivalence of consequence relations under arbitrary assumptions; it is more than the agreement of the two interfaces on true equations between numerals.

4.7.2 Five patterns with clause contexts

`uniform_step` combines Step with the clause contexts of Section 1.9: the formulas are the equations and the formulas $D_{x,y}B$, a context is a finite list of clauses, and an assumption of a formula A is the constant clause $\mathrm{ord}(A)$ of Section 5.1.2.

Definition 4.72 (The calculus $\mathsf{C}_{\mathrm{Step}}$)**.** $\mathsf{C}_{\mathrm{Step}}$ is the calculus C^0 of Theorem 5.20: zero-normal terms, normalizing substitution, clause contexts, and the five rules Use, Refl, the rule Step of Theorem 4.66 (called Eval there), the elimination DE that retains the clause $[B(b, 0);\, y.B(b, y)]$ (called Elim there), and WI with its assumption stored as a constant clause (called Intro there).

Theorem 4.73 (Faithfulness of the five patterns)**.** *For every finite context* Γ *of raw native formulas and every raw native formula* A*,*

$$\Gamma \vdash_{\mathsf{N}} A \quad \Longleftrightarrow \quad \mathrm{ord}(\mathrm{nf}_0(\tau\Gamma)) \vdash_{\mathsf{C}_{\mathrm{Step}}} \mathrm{nf}_0(\tau A),$$

where τ *is the translation of Theorem 5.1, which replaces each* $Q_y B$ *by* $D_{w,y}B$ *with an unused* w*. The ordinary realization of* $\mathsf{C}_{\mathrm{Step}}$ *is bi-interpretable and equiconsistent with* iPRA*, with whole numerical domains, equality interpreted by equality and numerical composites with the graph* $x = y$*.*

Proof. Theorem 5.22(iv), with Theorems 5.12 and 5.70 for the composites and the bridge. □

The derived rules of the calculus are those of Theorem 5.21; its items (ii)–(v) are the derivations of Theorem 4.67.

Remark 4.74. The calculus of `uniform_step` is C^0; the package differs from Section 5.3.2 in its proof of Theorem 4.73, in three respects. First, from N it translates the derivations of N^0 (Theorem 4.38) rule by rule, where Theorem 5.22(i) normalizes derivations of the raw clause calculus: Hyp is Use applied to the reflexivity of 0, IS and Tr are Theorem 4.67, QE

and QI are the closure interfaces of Theorem 5.9, and for DE the retained clause derives the translated closure (Theorem 5.9(iii)) and a cut supplies the branch. Second, its alias of zero is $q := I(1,0;\, u.\, u)$, and it recovers the introduction by cases of a literal equation $E(x,y)$ at the witness 0 without a cut: the given premise is closed at the unused witness q into $D_{w,y}E(0,y)$, the elimination retains the clause $[E(0,0);\, y.E(0,y)]$, and under the assumption $E(q,0)$ a transport along $q = 0$, a Use of that clause and a transport along $0 = q$ give $E(q,Sz)$, the premise of WI at q in case mode. Third, toward N it argues by the soundness of the five rules in T_{eq} under the ordinary reading, recorded after Theorem 5.70, and by Theorem 1.78, in place of the translation of annotated contexts in Theorem 5.22(ii). The five patterns keep the choice between cases and progress inside WI; the distinction between coverage and equation induction is not removed. The Lean formalization of the package is the arithmetic interface with native contexts (`NSPrf`, Theorem 4.70).

4.8 Seed evaluation and suspended application

This section draws on `seed_evaluation`: its rule SeedEval, which fuses the replacement of an initializer (Section 4.6) with the successor step (Section 4.7), and its second construction, in which every iteration at a successor counter is factored into a suspended one-step application.

4.8.1 The rule SeedEval

Definition 4.75. On zero-normal terms the rule SeedEval is

$$\frac{\Gamma \vdash a = b \qquad \Gamma \vdash I(Sc,a;\, u.\, h) = r}{\Gamma \vdash r = h\langle \overline{I}(c,b;\, u.\, h)/u\rangle}.$$

The calculus $\mathsf{C}_{\mathrm{Seed}}$ is $\mathsf{C}_{\mathrm{Step}}$ (Theorem 4.72) with SeedEval in place of Step.

The counter, the initializer and the body are read off the second premise, and the left side of the first premise is compared with the initializer a alone. In comparison with Step, the first premise no longer mentions the preceding iterate, and the conclusion changes at the same time: it iterates from the equal seed b before the last application of the body. At $c \equiv 0$ the two rules are the same inference,

$$\frac{a = b \qquad I(1,a;\, u.\, h) = r}{r = h\langle b/u\rangle},$$

and by Theorem 4.67 the pivot, symmetry, transitivity, congruence and Tr are derived from reflexivity and this shared instance. They are therefore available with either rule.

Lemma 4.76 (The two rules are interderivable)**.** *Over an arbitrary context and with reflexivity:*

(a) SeedEval derives successor computation, by one instance on two reflexivities;

(b) every instance of SeedEval is derivable from Step, by two instances at the counter 0 *and one further instance;*

(c) every instance of Step is derivable from SeedEval, by seven instances.

At $c \equiv 0$ no expansion is needed in (b) and (c). None of the derivations uses a binder rule, a case distinction or induction.

Proof. Put $P := \overline{I}(c, a; u.h)$ and $B := I(Sc, a;\, u.\, h)$.

(a) SeedEval applied to $a = a$ and $B = B$ concludes $B = h\langle P/u\rangle$. Its first premise does not contain P.

(b) Let $a = b$ and $B = r$ be given. Congruence (Theorem 4.67(d)) in the template $v \mapsto \overline{I}(c, v; u.h)$, with v fresh for c and h, gives $\overline{I}(c, a; u.h) = \overline{I}(c, b; u.h)$ by two instances of Step at the counter 0. Step with this equation and $B = r$ gives $r = h\langle\overline{I}(c, b; u.h)/u\rangle$. The body h is fixed throughout; no agreement of two different bodies is proved.

(c) Let $P = t$ and $B = r$ be given. By (a), $B = h\langle P/u\rangle$. The pivot applied to $B = r$ and $B = h\langle P/u\rangle$ gives $h\langle P/u\rangle = r$. Tr along $P = t$ in the template h gives $r = h\langle t/u\rangle$. The derivation has one instance for the computation, two for the pivot and four for Tr. □

Theorem 4.77. *For every finite context* Γ *of clauses and every formula* A *of the clause calculi,*

$$\Gamma \vdash_{\mathsf{C}_{\mathrm{Step}}} A \quad \Longleftrightarrow \quad \Gamma \vdash_{\mathsf{C}_{\mathrm{Seed}}} A.$$

The sequent is literally unchanged. The same equivalence holds for the calculi with native contexts: the calculus with the rules Hyp, Refl, SeedEval, QE, DE, DI and QI, and the calculus with WI in place of the last two, derive the sequents of N^0*. Consequently Theorem 4.73 holds for* $\mathsf{C}_{\mathrm{Seed}}$*.*

Proof. Replace each arithmetic inference of a derivation by its expansion in Theorem 4.76 and keep every other inference. The two calculi have the same terms, formulas, clauses, contexts, substitution and freshness conditions, and an expansion has the same premises and conclusion as the inference that it replaces. No witness is changed, so the mode of each WI is unchanged. The statement for native contexts follows in the same way from Theorem 4.70. □

The seven-rule calculus with native contexts is formalized as `NVPrf`, with `nvprf_iff_nprf`, `NVPrf.normal` and `nvprf_iff_pra`. The fusion differs from the rule of `initializer_equality`, which keeps successor computation as a separate rule, and from the package `counter_evaluation`, which fuses evaluation with an equation between counters for a recursor whose body binds the counter and the state.

Proposition 4.78 (Size of the computation certificate)**.** *Let* $c \not\equiv 0$*. The derivation of* $B = h\langle P/u\rangle$ *by Step is* $\mathrm{Step}(\mathrm{Refl}(P), \mathrm{Refl}(B))$*, and by SeedEval it is* $\mathrm{SeedEval}(\mathrm{Refl}(a), \mathrm{Refl}(B))$*. The two have the same number of inference nodes, and the second contains* $|P| - |a| = 1 + |c| + |h|$ *fewer nodes of term syntax, where* $|\cdot|$ *counts one node for each variable, constant and constructor. In a serialization that writes* $I(c, a;\, u.\, h)$ *as the JSON list* `["i",c,a,h]` *without optional whitespace, the second derivation is shorter by* $8 + \ell(c) + \ell(h)$ *bytes, where* ℓ *is the length of the serialized term. At* $c \equiv 0$ *the two derivations coincide.*

Proof. $|P| = 1 + |c| + |a| + |h|$, and the only difference between the two derivations is the term of the first reflexivity. The brackets, the tag and the separators of the list contribute eight bytes, and the counter and the body are no longer repeated. The conclusion is the same in both derivations. □

The saving grows without bound with the body. It is established for these derivations of a single successor computation; the translation of Theorem 4.76(c) expands other derivations, and a format that shares subterms changes the comparison.

4.8.2 Suspended application

The second construction of `seed_evaluation` removes the counter from the arithmetic rule altogether. The price is a normalization between the zero normalization and the full normalization of Section 4.9: an iterator at a successor counter is factored into a one-step application whose body is never executed by the normalizer.

Definition 4.79 (Unit normalization)**.** Extend the raw terms by the constructor $U(a;\, u.\, h)$, which binds u in h and not in a. Its *expansion* is $U(a; u.h)^\circ := I(1, a^\circ;\, u.\, h^\circ)$, and $(\cdot)^\circ$ commutes with the other constructors. On terms n, a, h define

$$K(n, a;\, u.\, h) := \begin{cases} a, & n \equiv 0, \\ U(K(m, a;\, u.\, h);\, u.\, h), & n \equiv Sm, \\ I(n, a;\, u.\, h), & \text{otherwise}, \end{cases}$$

by recursion on n, and define nf_U by structural recursion: $\mathrm{nf}_U(x) := x$, $\mathrm{nf}_U(0) := 0$, $\mathrm{nf}_U(St) := S\,\mathrm{nf}_U(t)$,

$$\mathrm{nf}_U(U(a; u.h)) := U(\mathrm{nf}_U(a); u.\,\mathrm{nf}_U(h)), \qquad \mathrm{nf}_U(I(n, a;\, u.\, h)) := K(\mathrm{nf}_U(n), \mathrm{nf}_U(a);\, u.\, \mathrm{nf}_U(h)).$$

A term is *unit-normal* if it is fixed by nf_U: its subterms are unit-normal and the counter of each of its iterators has neither 0 nor S at its root.

For example, for a variable x,

$$\mathrm{nf}_U\big(I(SSx, a;\, u.\, h)\big) \equiv U\big(U(I(x, \mathrm{nf}_U(a);\, u.\, \mathrm{nf}_U(h)); u.\,\mathrm{nf}_U(h)); u.\,\mathrm{nf}_U(h)\big).$$

Each displayed u binds its own copy of the body. The result is obtained by inspecting the leading successors of the counter; neither $h[a/u]$ nor any other execution of h occurs. A suspended application is always kept, also when its body is constant or the identity. In T_{eq} the expansion of $U(a; u.h)$ equals $h^\circ[a^\circ/u]$, so the constructor adds no numerical freedom and no strength.

Lemma 4.80 (Termination and uniqueness)**.** *Consider the two contractions*

$$I(0, a;\, u.\, h) \to a, \qquad I(Sn, a;\, u.\, h) \to U(I(n, a;\, u.\, h);\, u.\, h),$$

closed under every context, with no contraction for U. Every reduction sequence is finite, every term reduces to $\mathrm{nf}_U(t)$, and $\mathrm{nf}_U(t)$ is the unique normal form of t. In particular the reduction is confluent, and nf_U is idempotent and primitive recursive on codes of terms.

Proof. Assign positive integer weights by $w(x) = w(0) = 1$, $w(St) = w(t) + 1$,

$$w(U(a; u.h)) = w(a) + w(h) + 1, \qquad w(I(n, a;\, u.\, h)) = (w(n) + 1)(w(a) + w(h) + 1).$$

Write $A = w(a)$, $H = w(h)$ and $q = w(n)$. The zero contraction decreases the weight from $2(A + H + 1)$ to A. The second contraction changes it from $(q + 2)(A + H + 1)$ to $(q + 1)(A + H + 1) + H + 1$, a decrease by exactly $A > 0$; the duplication of the body is accounted for. Every constructor is strictly increasing in the weight of each child, so the decrease persists in every context, including bodies. Hence every reduction decreases a positive integer.

nf_U is invariant under both contractions at the root: for the first by its definition, and for the second because

$$\mathrm{nf}_U(I(Sn, a;\, u.\, h)) \equiv U\big(K(\mathrm{nf}_U(n), \mathrm{nf}_U(a); u.\,\mathrm{nf}_U(h)); u.\,\mathrm{nf}_U(h)\big) \equiv \mathrm{nf}_U\big(U(I(n, a;\, u.\, h); u.h)\big).$$

A structural induction extends the invariance to contractions in context. The irreducible terms are exactly the unit-normal terms. By termination every term has an irreducible reduct, which by invariance is $\mathrm{nf}_U(t)$. Any two reducts of t reduce to this term.

With a primitive recursive coding of terms, the weight and a deterministic choice of one contraction are primitive recursive, and fewer than $w(t)$ contractions occur; the bounded iteration of the step computes nf_U. The normalizer is not an evaluator of primitive recursive programs: it never substitutes an argument into the body of U. □

Lemma 4.81 (Coherence with substitution)**.** $\mathrm{nf}_U(t[\sigma]) \equiv \mathrm{nf}_U(\mathrm{nf}_U(t)[\mathrm{nf}_U \circ \sigma])$ *for all terms* t *and substitutions* σ*. Hence the normalizing substitution* $t\langle\sigma\rangle_U := \mathrm{nf}_U(t[\sigma])$ *satisfies the laws of Theorem 4.26;* nf_U *commutes with renaming and introduces no free variable. On terms without* U*,* $\mathrm{nf}_U \circ \mathrm{nf}_0 = \mathrm{nf}_U$*.*

Proof. Both contractions are preserved by every capture-free substitution, and contracting inside the terms substituted into a fixed term is a reduction of the instance. Hence $t[\sigma]$ reduces to $\mathrm{nf}_U(t)[\mathrm{nf}_U \circ \sigma]$, and both have the normal form of the left side, by Theorem 4.80. Below U and the body of an iterator the substitution is lifted. The laws follow as in Theorem 4.26, whose proof uses only $(\nu 1)$–$(\nu 4)$. The last statement holds because $t \to^* \mathrm{nf}_0(t)$ by zero contractions, which are among the two contractions. □

Definition 4.82 (The rule App and the calculus $\mathsf{C}_{\mathrm{App}}$)**.** On unit-normal terms the rule App is

$$\frac{\Gamma \vdash a = b \qquad \Gamma \vdash U(a;\, u.\, h) = r}{\Gamma \vdash r = h\langle b/u\rangle_U}.$$

The calculus $\mathsf{C}_{\mathrm{App}}$ has unit-normal terms, the formulas and clauses of Section 4.7.2 over them, the substitution $\langle\cdot\rangle_U$, and the five rules Use, Refl, App, DE and WI, the last three logical rules as in Theorem 4.72.

App inspects no counter and builds no predecessor iterate; that work is done by the constructor K.

Lemma 4.83. *From Refl and App, over an arbitrary context:*

(a) the pivot, symmetry, transitivity, congruence in every unit-normal template and Tr are derivable, without induction or case distinction;

(b) for unit-normal c, a, h*,* $K(Sc, a; u.h) = h\langle K(c, a; u.h)/u\rangle_U$ *is derivable by one App on two reflexivities;*

(c) the image under nf_U *of an instance of Step on zero-normal terms is an instance of App.*

Proof. (a) As in Theorem 4.67, with the frame $J_s := U(s; v.v)$ for the pivot and $H_s := U(s; v.h)$ for congruence: App on $s = r$ and $J_s = J_s$ gives $J_s = r$, and App on $s = t$ and $J_s = r$ gives $r = t$; App on $s = s$ and $H_s = H_s$ gives $H_s = h\langle s/v\rangle_U$, and App on $s = t$ and this equation gives $h\langle s/v\rangle_U = h\langle t/v\rangle_U$.

(b) By definition $K(Sc, a; u.h) \equiv U(K(c, a; u.h); u.h)$. Apply App to the reflexivities of $K(c, a; u.h)$ and of this term.

(c) Let the instance have the premises $\overline{I}(c, a; u.h) = t$ and $I(Sc, a;\, u.\, h) = r$ and the conclusion $r = \mathrm{nf}_0(h[t/u])$. Write c', a', h' for the images of c, a, h under nf_U. Then $\mathrm{nf}_U(\overline{I}(c, a; u.h)) \equiv K(c', a'; u.h')$, in both cases of the smart constructor, and $\mathrm{nf}_U(I(Sc, a;\, u.\, h)) \equiv U(K(c', a'; u.h'); u.h')$. App concludes $\mathrm{nf}_U(r) = h'\langle \mathrm{nf}_U(t)/u\rangle_U$, and by Theorem 4.81 this term is $\mathrm{nf}_U(h[t/u]) \equiv \mathrm{nf}_U(\mathrm{nf}_0(h[t/u]))$. □

Lemma 4.84. *For every term* t *without* U*,* $\mathrm{nf}_U(t) \equiv 0$ *if and only if* $\mathrm{nf}_0(t) \equiv 0$*. In particular, for a zero-normal* t*,* $\mathrm{nf}_U(t) \equiv 0$ *if and only if* $t \equiv 0$*.*

Proof. By structural induction. A variable and a successor do not normalize to 0 under either map. For an iterator, nf_U yields 0 exactly when the normalized counter and the normalized initializer are both 0: otherwise the result is the initializer, which is not 0, or has U or an iterator at its root. The same holds for nf_0, and the induction hypotheses for the counter and the initializer apply. □

Theorem 4.85. *Let Γ be a context of clauses and A a formula of $\mathsf{C}_{\mathrm{Step}}$.*

(a) If $\Gamma \vdash_{\mathsf{C}_{\mathrm{Step}}} A$, then $\mathrm{nf}_U(\Gamma) \vdash_{\mathsf{C}_{\mathrm{App}}} \mathrm{nf}_U(A)$.

(b) $\mathsf{C}_{\mathrm{App}}$ is sound for T_{eq} under the expansion $(\cdot)^\circ$ and the readings of clauses and formulas, and T_{eq} proves $t^\circ = \mathrm{nf}_U(t)^\circ$ for every term t.

(c) If Γ consists of assumption clauses and $\mathrm{nf}_U(\Gamma) \vdash_{\mathsf{C}_{\mathrm{App}}} \mathrm{nf}_U(A)$, then $\Gamma \vdash_{\mathsf{C}_{\mathrm{Step}}} A$.

(d) For a unit-normal t, $\mathrm{nf}_U(t^\circ) \equiv t$.

Consequently the ordinary realization of $\mathsf{C}_{\mathrm{App}}$ is bi-interpretable and equiconsistent with iPRA*, with whole domains, equality interpreted by equality and numerical composites with the graph $x = y$.*

Proof. (a) By induction on the derivation, translating terms by nf_U, formulas termwise and contexts clause by clause. Refl is immediate, and each Step becomes one App by Theorem 4.83(c). Use and DE are translated by Theorem 4.81. For WI the root of the matrix is preserved, and by Theorem 4.84 the witness is literally 0 after the translation exactly when it was before, so the mode is the same; freshness is preserved because nf_U introduces no variable.

(b) In T_{eq} the zero contraction is the zero computation equation. For the second contraction both sides equal $h[I(n, a;\, u.\, h)/u]$: the left by successor computation, the expansion of the right by the successor and the zero computation at the counter 1. Contractions in the counter or the initializer are lifted by congruence, and contractions in a body by body congruence (Theorem 1.39(ii), transferred to T_{eq}), which uses the induction for literal equations. Hence $\mathsf{T}_{\mathrm{eq}} \vdash t^\circ = \mathrm{nf}_U(t)^\circ$. App is sound by the computation at the counter 1 and equality, and the logical rules are sound under the ordinary reading as recorded after Theorem 5.70: Use and DE are ordinary eliminations, WI in case mode uses coverage and in progress mode the induction for a literal equation under the assumed base, and a normalized instance and the ordinary instance are provably equal.

(c) By (b), T_{eq} proves the reading of A from the readings of Γ. If Γ is $\mathrm{ord}(\Delta)$, then Δ and A are zero-normal native formulas without Q_y, so $\Delta \vdash_{\mathsf{N}} A$ by Theorem 1.78, and Theorem 5.22(iii) gives $\Gamma \vdash_{\mathsf{C}_{\mathrm{Step}}} A$.

(d) By structural induction: the expansion $I(1, a^\circ;\, u.\, h^\circ)$ of an application normalizes to $U(a; u.h)$, and an iterator of a unit-normal term has a counter with neither 0 nor S at its root, whose normalized expansion is the same counter.

The last statement follows from (a)–(d) and Theorem 4.73: the constructor U is an abbreviation of a fixed term, and normalization changes representatives of expressions only. □

Proposition 4.86 (The variant without a new constructor)**.** *Let $U(a; u.h)$ abbreviate $I(1, a;\, u.\, h)$ in the unchanged term grammar, and define K and nf_U by the clauses of Theorem 4.79, without the clause for U. The fixed points of nf_U are the terms in which every iterator counter is the literal 1 or has a root other than 0 and S. The normal forms are those of the contractions*

$$I(0, a;\, u.\, h) \to a, \qquad I(Sn, a;\, u.\, h) \to I(1, I(n, a;\, u.\, h);\, u.\, h) \quad (n \not\equiv 0),$$

which terminate; Theorems 4.81 and 4.84 hold; the rule App is the instance of Step at the counter 0, followed by normalization; and Theorem 4.85 holds for the resulting calculus.

Proof. The side condition on the second contraction is necessary: without it $I(1, a;\, u.\, h)$ would be contracted to $I(1, I(0, a;\, u.\, h);\, u.\, h)$, which the zero contraction contracts back. For termination use the weights $w(x) = w(0) = 1$, $w(St) = w(t) + 1$ and, with $A = w(a)$ and $H = w(h)$,

$$w(I(n, a;\, u.\, h)) = \begin{cases} A + H + 1, & n \equiv 1, \\ (w(n) + 1)(A + H + 1), & \text{otherwise.} \end{cases}$$

The zero contraction decreases the weight from $2(A+H+1)$ to A. In the second contraction the left counter Sn is not 1, so the left weight is $(w(n)+2)(A+H+1)$. If $n \not\equiv 1$ the right weight is $(w(n)+1)(A+H+1)+H+1$, smaller by A. If $n \equiv 1$ the right weight is $A+2H+2$ and the left weight is $4(A+H+1)$. The weight is strictly increasing in the initializer and the body. In the counter position the one exceptional case is a contraction that makes the counter the literal 1, which replaces the product by the smaller sum; a counter that is the irreducible term 1 is not contracted. Hence every contraction in context decreases the weight.

Both contractions preserve nf_U; for the second this is the identity $K(Sn, a; u.h) \equiv U(K(n, a; u.h); u.h)$, which holds also at $n \equiv 0$. For coherence with substitution the conditional contraction is not closed under substitution, since a substitution can turn n into a term with the normal form 0. Argue in two steps. First, replacing each substituted term by its normal form does not change the normal form of the instance, by the invariance in context. Second, if $t \to t'$ by one contraction, then $\mathrm{nf}_U(t[\sigma]) \equiv \mathrm{nf}_U(t'[\sigma])$: for the zero contraction literally, and for the second by the displayed identity for K, which is valid at 0. Following a reduction of t to $\mathrm{nf}_U(t)$ gives the statement. The remaining proofs are those of Theorems 4.83 to 4.85; the unit iterator is never removed and its body is never exposed. □

4.9 Full normalization

This section draws on `full_normalization` and `computation_normal`. The two packages arrive at the same six-rule calculus on full normal forms; they differ in the proof of termination, in the repair of the single introduction and in their measurements. Both computations are moved into the syntax, and no computation rule remains.

4.9.1 Termination and unique normal forms

The reduction $\to$ is that of Theorem 4.5. It is stable under substitution: if $t \to t'$, then $t[\sigma] \to t'[\sigma]$; and if $\sigma(x) \to^* \sigma'(x)$ for every x, then $t[\sigma] \to^* t[\sigma']$. A term is *strongly normalizing* if it has no infinite reduction sequence. A term has finitely many one-step reducts, so for a strongly normalizing term t the reduction sequences from t have a maximal length $d(t)$. The number of nodes of t is $|t|$.

Lemma 4.87 (Iterator closure)**.** *Let n and a be strongly normalizing, and let $h[v/u]$ be strongly normalizing for every strongly normalizing term v. Then $I(n, a; u.\, h)$ is strongly normalizing.*

Proof. With $v \equiv u$ the body h is strongly normalizing. Argue by induction on the triple $(d(n), |n|, d(a)+d(h))$ in the lexicographic order; it suffices to show that every one-step reduct of $I(n, a; u.\, h)$ is strongly normalizing. A reduction in n decreases the first component. A reduction in a or in h leaves the first two components and decreases the third; if $h \to h'$, then $h'[v/u]$ is a reduct of $h[v/u]$ for every v, so the hypothesis on the body persists. If $n \equiv 0$, the root reduct is a. If $n \equiv Sp$, then $d(p) = d(n)$ and $|p| < |n|$, so $I(p, a; u.\, h)$ is strongly normalizing by the induction hypothesis, and the root reduct $h[I(p, a; u.\, h)/u]$ is strongly normalizing by the hypothesis on the body. □

Theorem 4.88 (Strong normalization)**.** *If σ assigns strongly normalizing terms to the variables, then $t[\sigma]$ is strongly normalizing for every raw term t. In particular every raw term is strongly normalizing.*

Proof. By induction on t. Variables, 0 and successors are immediate. For $I(n, a; u.\, h)$, with u renamed away from σ, the instances $n[\sigma]$ and $a[\sigma]$ are strongly normalizing by the induction hypotheses. The induction hypothesis for h applies to every extension of σ by an assignment

of a strongly normalizing v to u; this is the hypothesis on the body in Theorem 4.87 for the term $I(n[\sigma], a[\sigma]; u.\, h[\sigma])$. The last statement is the case of the identity substitution. □

Theorem 4.89 (Confluence)**.** *The reduction $\to$ is confluent. Hence every raw term t has a unique normal form* $\mathrm{nf}(t)$*, up to the renaming of bound variables.*

Proof. Define the *parallel reduction* $\Rightarrow$ inductively: $x \Rightarrow x$; $0 \Rightarrow 0$; if $t \Rightarrow t'$ then $St \Rightarrow St'$; and if $n \Rightarrow n'$, $a \Rightarrow a'$ and $h \Rightarrow h'$, then

$$I(n, a; u.\, h) \Rightarrow I(n', a'; u.\, h'), \qquad I(0, a; u.\, h) \Rightarrow a', \qquad I(Sn, a; u.\, h) \Rightarrow h'[I(n', a'; u.\, h')/u].$$

Then $t \Rightarrow t$, and $\to \subseteq \Rightarrow \subseteq \to^*$. By induction on $h \Rightarrow h'$, if also $r \Rightarrow r'$ then $h[r/u] \Rightarrow h'[r'/u]$; the case of a contracted successor redex uses the composition law of substitution.

Define the *complete development* $t^\bullet$ by $x^\bullet := x$, $0^\bullet := 0$, $(St)^\bullet := St^\bullet$, $(I(0, a; u.\, h))^\bullet := a^\bullet$, $(I(Sn, a; u.\, h))^\bullet := h^\bullet[I(n^\bullet, a^\bullet; u.\, h^\bullet)/u]$, and $(I(n, a; u.\, h))^\bullet := I(n^\bullet, a^\bullet; u.\, h^\bullet)$ for every other n. The claim is that $t \Rightarrow t'$ implies $t' \Rightarrow t^\bullet$; it is proved by induction on $t \Rightarrow t'$. Consider $I(n, a; u.\, h) \Rightarrow I(n', a'; u.\, h')$, with $n' \Rightarrow n^\bullet$, $a' \Rightarrow a^\bullet$ and $h' \Rightarrow h^\bullet$ by the induction hypotheses. If $n \equiv 0$, then $n' \equiv 0$ and $I(0, a'; u.\, h') \Rightarrow a^\bullet$. If $n \equiv Sm$, then $n' \equiv Sm'$ with $m' \Rightarrow m^\bullet$, and $I(Sm', a'; u.\, h') \Rightarrow h^\bullet[I(m^\bullet, a^\bullet; u.\, h^\bullet)/u]$. Otherwise the congruence rule applies. For $I(0, a; u.\, h) \Rightarrow a'$ the induction hypothesis is $a' \Rightarrow a^\bullet$. For $I(Sn, a; u.\, h) \Rightarrow h'[I(n', a'; u.\, h')/u]$ the substitution property, applied to $h' \Rightarrow h^\bullet$ and $I(n', a'; u.\, h') \Rightarrow I(n^\bullet, a^\bullet; u.\, h^\bullet)$, gives the reduct $(I(Sn, a; u.\, h))^\bullet$. The remaining cases are immediate.

By the claim, two parallel reducts of t both reduce in parallel to $t^\bullet$, so $\Rightarrow$ has the diamond property, and its transitive closure $\to^*$ is confluent. With Theorem 4.88, every term has exactly one normal form. □

Remark 4.90. The proof of Theorem 4.88 is that of `full_normalization`, which obtains confluence from local confluence and termination. `computation_normal` obtains termination by an embedding into the simply typed calculus with a recursor at the type of numbers (Gödel's system T): the iterator is translated as a recursor whose step ignores its counter, a zero contraction becomes one recursor contraction, a successor contraction becomes one recursor contraction followed by two β-contractions, and the translation respects contexts and substitution, so an infinite reduction would contradict the strong normalization of T, for which the package cites the mechanized proof of S. Urciuoli (EPTCS 376, 2023); it obtains confluence by parallel reduction, as above. The Lean formalization defines the normalizer by evaluation: the term is evaluated in an environment of terms by structural recursion (`Tm.nbe`), the iterator by a recursion on the value of its counter that unfolds a numeral prefix and leaves a neutral counter in place (`Tm.recV`), and `Tm.full` is the value in the identity environment. Totality and uniqueness of normal forms then need no proof. The system T and the evaluation are used in the metatheory; neither is added to the arithmetic.

Lemma 4.91 (Full normal forms)**.** *The normal terms are generated by*

$$ne ::= x \mid I(ne, nt; u.\, nt), \qquad nt ::= ne \mid 0 \mid S\, nt.$$

The terms ne are the neutral *terms. The counter of every iterator of a normal term is neutral; it has neither 0 nor S at its root. Every closed normal term is a numeral.*

Proof. A term is normal exactly when its subterms are normal and no contraction applies at its root, which gives the grammar. A neutral term has a free variable: following the counters from its root leads, outside every binder, to a variable. Hence a closed normal term is built from 0 and S. □

For example $I(x, 0; u.\, 0)$ is normal, although its value is 0 for every x, while $I(Sx, 0; u.\, 0)$ has the normal form 0. Normalization does not decide equality.

4.9.2 The kernel without computation rules

Proposition 4.92. nf *is an admissible normalizer.*

Proof. (ν1): variables and 0 are normal, and the reducts of St are the terms St' with t' a reduct of t. (ν2): reduction commutes with renaming, the renaming of a normal term is normal by Theorem 4.91, and a contraction creates no free variable. (ν3) is immediate. (ν4): by the two stability properties, $t[\sigma] \to^* \mathrm{nf}(t)[\sigma] \to^* \mathrm{nf}(t)[\mathrm{nf} \circ \sigma]$, so both sides of ($\nu$4) are the normal form of $t[\sigma]$. (ν5) is Theorem 4.6. □

In the rest of this section terms and formulas are in full normal form and $t\langle\sigma\rangle$ is $\mathrm{nf}(t[\sigma])$. An expression $I(n, a;\, u.\, h)$ in a rule denotes the normal form of the displayed term, also when a substitution exposes a redex. Formula normalization changes terms and never removes Q_y or $D_{x,y}$.

Definition 4.93 (The kernel N^{f})**.** N^{f} has native formulas over full normal forms, normalizing substitution, and the seven rules Hyp, Refl (for every normal term), Tr, QE, DE, DI and QI.

Theorem 4.94 (Both computation rules are deleted)**.** *For every finite context* Γ *of raw native formulas and every raw native formula* A*,*

$$\Gamma \vdash_{\mathsf{N}} A \quad \Longleftrightarrow \quad \mathrm{nf}(\Gamma) \vdash_{\mathsf{N}^{\mathrm{f}}} \mathrm{nf}(A).$$

N^{f} *is closed under weakening, cut and normalizing substitution, and Theorem 4.31 applies to it.*

Proof. By Theorem 4.92 and Theorem 4.30 it suffices that N^{f} and N^{nf} have the same rules. The left side of an instance e of I0 or IS reduces to its right side, so $\mathrm{nf}(e)$ is literally $r = r$ for a normal r, an instance of Refl. Conversely, for a normal r the reflexivity $r = r$ is $\mathrm{nf}(e)$ for the instance $I(0, r;\, u.\, 0) = r$ of I0. □

Remark 4.95. The theorem is formalized as `fnprf_iff_nprf` for the calculus `FNPrf`; the laws of the normalizer are `Tm.full_subst_congr` and `NFm.full_subst_congr`, and (ν5) is `NPrf.eqFull`. Both packages state the theorem with the auxiliary seven-rule calculus as a bridge to their six rules, and both stress that the comparison uses properties of the raw calculus that are established before it: the equality rules, the Leibniz rule, cut and equation induction of N.

4.9.3 Six rules on full normal forms

Definition 4.96 (The calculus F)**.** F has the syntax and the substitution of N^{f} and the six rules Hyp, Refl, Tr, QE, DE and WI. The template of Tr is any normal term with one distinguished variable, used any number of times; there is no restriction on the terms of its first premise. The test $t \equiv 0$ of WI is applied to the normal witness.

There is no computation axiom: both equations are enforced by the normalizing substitution. Weakening, injective renaming and cut are admissible in F by direct induction on derivations, since they preserve normality, the literal 0, the root of a matrix and freshness, and no guard depends on the context. The equality rules and transport in literal equations are derivable by Theorem 4.2(b), the instances matching by Theorem 4.26.

On full normal forms Theorem 4.3(ii) has no closed term Z to use: by Theorem 4.91 a closed normal term is a numeral, and the aliases of Theorem 4.40 have the normal form 0. The packages give two repairs.

Lemma 4.97 (A fresh name for zero)**.** *Let $Z(x,y)$ be the equation $I(y,0;\,u.\,x)=0$. Then $\vdash_{\mathsf{F}} D_{x,y}Z(x,y)$, and $Q_yZ(b,y)\vdash_{\mathsf{F}} b=0$ for a variable b.*

Proof. Introduce $D_{x,y}Z$ by WI at the witness 0. The matrix is a literal equation, so the mode is progress. The conclusion of the premise is $Z(0,Sz)$, the normal form of $I(Sz,0;\,u.\,0)=0$, which is $0=0$; reflexivity derives it without the assumption. For the second statement the base $Z(b,0)$ is the normal form of $I(0,0;\,u.\,b)=0$, which is $0=0$. QE with this base, derived by reflexivity, at the target 1 gives $Z(b,1)$, the normal form of $I(1,0;\,u.\,b)=0$, which is $b=0$. □

The witness opened by DE is a variable, hence not literally 0, and it is provably 0. It is discharged before the conclusion; no constant and no element of the domain is added.

Theorem 4.98. *For every context Γ and formula A in full normal form,*

$$\Gamma\vdash_{\mathsf{F}} A \quad\Longleftrightarrow\quad \Gamma\vdash_{\mathsf{N}^{\mathrm{f}}} A.$$

Hence $\Gamma\vdash_{\mathsf{N}} A$ if and only if $\mathrm{nf}(\Gamma)\vdash_{\mathsf{F}}\mathrm{nf}(A)$ for raw Γ,A, and substitution and the Leibniz rule are admissible in F.

Proof. From N^{f} to F: QI is admissible by Theorem 4.3(i). An instance of DI is an instance of WI unless its matrix is a literal equation $E(x,y)$ and its witness is 0. In that case let $\Gamma,E(0,0)\vdash E(0,Sz)$ be given. Choose b fresh and work in the context $\Gamma':=\Gamma,Q_yZ(b,y)$, where $\Gamma'\vdash b=0$ by Theorem 4.97. By Theorem 4.3(ii) with the term b, applied to the weakened premise, $\Gamma'\vdash D_{x,y}E$. DE with the first premise $\vdash D_{x,y}Z$ closes the scope of b, which occurs neither in Γ nor in $D_{x,y}E$. From F to N^{f}, WI is admissible by Theorem 4.3(iii). The remaining statements are Theorems 4.31 and 4.94. □

Only transport in literal equations, weakening, injective renaming and cut are used in the repair; the Leibniz rule and the substitution of terms in derivations of F are consequences of the theorem and not premises of it.

Lemma 4.99 (The compound envelope)**.** *For a literal equation $E\equiv(s=t)$ in full normal form put*

$$G(v,E):=\big(I(v,0;\,u.\,s)=I(v,0;\,u.\,t)\big),\qquad L(E):=Q_vG(v,E),$$

with u,v not free in E. Then $G(0,E)\equiv(0=0)$ and $G(Sw,E)\equiv E$, and for every context Γ both $\Gamma,E\vdash_{\mathsf{F}} L(E)$ and $\Gamma,L(E)\vdash_{\mathsf{F}} E$.

Proof. The two identities are computations of normal forms; in the second, u is not free in s and t. For the first derivation apply the closure by cases of Theorem 4.3(i) to the matrix $G(v,E)$: its premise asks for $G(Sz,E)$, which is the assumption E. For the second apply QE to $L(E)$ with the base $0=0$ at the target 1. Both are uniform in the parameters of E and use no induction. □

The envelope yields a second proof of the missing case in Theorem 4.98, without Theorem 4.97. Let $\Gamma,E(0,0)\vdash E(0,Sz)$ and put $F(x,y):=L(E(x,y))$, a matrix with the root Q_v. In the context $\Gamma,F(0,0)$ the envelope gives $E(0,0)$, cut gives $E(0,Sz)$, and the envelope gives $F(0,Sz)$. WI at the witness 0 is in case mode, because F is not a literal equation, and derives $D_{x,y}F$. Open it by DE at a fresh b, retaining $Q_yF(b,y)$. To introduce $D_{x,y}E$ at the witness b, in case mode since a variable is not 0, assume $E(b,0)$; the envelope gives $F(b,0)$, QE gives $F(b,Sz)$, and the envelope gives $E(b,Sz)$. DE closes the derivation.

Remark 4.100. The fresh-zero lemma is the repair of `computation_normal`, and the compound envelope that of `full_normalization`. The Lean formalization of the six rules, `F6Prf` with `f6prf_iff_nprf`, uses a third repair, `F6Prf.dIZero`: with $T(x) := \mathrm{sel}(x, 0, 0)$, the formula $D_{x,y}B(T(x), y)$ is introduced at the witness 1, in case mode, from the given premise, because $T(1)$ has the normal form 0; it is opened at a fresh b; and $D_{x,y}B$ is introduced at the neutral witness $T(b)$ from the opened closure by QE.

Example 4.101. The formula $D_{x,y}(y = 1)$ is derivable: under its reading the antecedent is $0 = 1$. On raw terms it is introduced in case mode at the witness $I(0, 0; u.\, 0)$, from the premise $0 = 1 \vdash Sz = 1$. Full normalization turns this witness into 0, and the same inference would be in progress mode, with the premise $z = 1 \vdash Sz = 1$, which is not derivable. A derivation normalized rule by rule is therefore rejected, and the derivation produced by either repair is accepted. The package `full_normalization` ships both certificates.

4.9.4 What normalization costs

Proposition 4.102 (No primitive recursive uniform normalizer)**.** *The map $t \mapsto \mathrm{nf}(t)$ on codes of raw terms is total recursive and not primitive recursive.*

Proof. Totality is Theorems 4.88 and 4.89. Suppose that the map is primitive recursive. Enumerate the raw terms with one free variable, replacing a code that is not well formed by the code of the term 0. The check of well-formedness, the formation of the closed instance of a term at a numeral and the reading of the value of a numeral are primitive recursive. By Theorem 4.91 and (ν5) the normal form of a closed instance is the numeral of its value. Hence the function $V(e, n)$, the value at n of the term with the code e, is primitive recursive. By Theorem 1.62 every primitive recursive unary function is the value function of such a term. In particular $n \mapsto V(n, n) + 1$ has a code e, and $V(e, e) = V(e, e) + 1$, a contradiction. □

Each fixed term denotes a primitive recursive function; the obstruction is to one evaluator for all descriptions. A checker for F or N^{f} is a total recursive program that contains a normalizer, and its running time and the size of its intermediate terms are not bounded by a primitive recursive function of the input: by Theorem 4.91 every closed term is expanded to a numeral, so a short closed expression can have a normal form of enormous size. No primitive recursive bound and no proof in iPRA of a uniform normalization theorem is claimed. The statements of this section are external statements about each finite derivation; both interpretations of Theorem 4.31 act on fixed finite expressions.

Proposition 4.103 (Fuel)**.** *Let the checker of F be modified so that a certificate carries a number f, each successor contraction spends one unit, and exhaustion rejects. The relation "the certificate with fuel f is accepted" is primitive recursive, and a formula has an accepted certificate for some fuel exactly when it has an accepted certificate of the unbounded checker.*

Proof. Let m be the size of the certificate. An instance of a term or formula of size at most M at terms of size at most M has size at most M^2. A successor contraction replaces a term of size at most M by a term of size at most M^2, since the body and the recursive iterator are subterms of the redex and the iterator is copied once for each occurrence of the bound variable. A zero contraction and the traversals between contractions do not increase the size, and at most M zero contractions occur in a row. The checker performs at most m inferences, each with a number of substitutions bounded by a constant k, and at most f successor contractions. Hence every term that occurs has size at most $m^{2^{km+f}}$, and the length of the computation is bounded by a primitive recursive function of m and f; a computation with such bounds on its length and on the size of its configurations is primitive recursive. An accepting run of the unbounded checker is finite and performs finitely many successor contractions, so it is accepted with that much fuel; a run accepted with fuel is a run of the unbounded checker. □

`computation_normal` ships the fuelled checker and measures it (Section 4.11); `full_normalization` observes that a relation of this kind can also be obtained from explicit reduction traces checked step by step, and does not build that checker.

4.10 A smaller checker for the unchanged core

This section draws on the second result of `computation_normal`, which concerns a program and no calculus. The rules, the certificates and the kernel function of the checker are those of the package `validated_presentation`; only the validator of the public goal changes, by a composition of that validator with the reuse of the scope traversal of the package `scope_kernel`.

The input is a JSON array. Its last element is the closed public goal, in the arity coding of the native formulas, and the earlier elements are proof packets consumed from the end; an unused prefix is allowed. A lexical gate accepts the whole input only if every value in it is an integer or a list. The public validator of the baseline carries a depth d, even at formula positions and odd at numerical positions; a node with r children changes the depth by $5 - r$, a formula node has one, two or three children, of which the node with two children is the equation and has numerical children, and a numerical node has one or three children. A numerical leaf a is accepted if d is odd and $a = 0$ or a is odd with $2 < a < d$. The kernel, before it reads any packet, opens the goal by a scope traversal whose substitution argument is, in this initial call, the kernel function itself and not a term: at a leaf within the scope the traversal returns the leaf, and at a leaf outside the scope it attempts to use the substitution argument as a term and fails.

The new validator keeps only the Boolean "numerical position", which becomes true below a node with two children and is otherwise inherited, and accepts a numerical leaf a if $a = 0$ or a is odd with $a > 2$.

Theorem 4.104 (Same accepted inputs)**.** *In the execution model without resource limits, the new checker and the baseline accept exactly the same inputs.*

Proof. The lexical gate is the same. On inputs of integers and lists, induction on the tree shows that the Boolean of the new validator is the parity of the depth of the old one wherever the shape is valid, by the table of arities above. All old tests remain except the upper bound $a < d$ at numerical leaves, so every goal accepted by the old validator is accepted by the new one. Let a goal be accepted by the new validator and violate an upper bound. The kernel is unchanged and computes the initial opening of the goal before it tests membership in the context or reads a packet. Every numerical position has an odd depth at least three, and the leaf is 0 or an odd integer greater than two; at the violating leaf the opening uses its substitution argument, which is a function, as a term, and the operation raises an exception. An exception is a rejection in both programs, and no packet can prevent it. A formula-position integer is rejected by both validators. On goals accepted by the old validator the initial opening is the identity, and all later steps of the two programs are the same. □

Hence every soundness, completeness and interpretation result for the certificate system of the baseline holds for the new checker. The result is an implementation change for an unchanged calculus; its sizes are in Section 4.11.

4.11 Checker sizes

This section collects the measurements of all nine packages, by the protocol of Section 1.10. Every row of Table 4.2 is a complete standalone program, with its input handling, its validation of syntax and scope, its substitution and normalization, all rules and its output. The

gzip column, at level 9 with the timestamp zero, is reported by the packages for information and is not an objective. The reference interpreter is CPython 3.12.14. `uniform_step`, `seed_evaluation` and `initializer_equality` remove docstrings before normalization; the other packages do not. A dash marks a number that the package does not state.

Table 4.2: Complete checkers and their matched controls, as stated by the packages.

Complete program	Tokens	AST nodes	Bytes	Gzip-9
`computational_equality` (tagged, native contexts)				
Archived seven-rule control	1024	1080	3145	964
Matched seven-rule control	1008	1064	3121	957
E_3	959	1026	3039	966
`equality_from_iteration` (same format)				
E_3, upstream	959	1026	3039	966
E_2	954	1019	2996	955
Zero-normal control with ROOT kept	987	1059	3064	982
U	979	1047	3011	965
`uniform_step` (clause contexts)				
Matched zero-normal control, computation and transport	860	949	3048	1018
Matched raw control, three-case EVAL	853	969	3094	1026
Incoming canonical clausal checker	946	1072	3305	1073
$\mathsf{C}_{\mathrm{Step}}$	829	902	2950	1005
Native contexts: matched zero-normal control	1010	1080	3118	–
Native contexts: REFL and STEP	981	1051	3029	–
`seed_evaluation` (clause contexts)				
Computation and transport control	860	949	3048	–
STEP, helper style	829	902	2950	–
SEEDEVAL, helper style	829	902	2950	–
STEP, shared forward traversal	758	815	2804	–
SEEDEVAL, shared forward traversal	756	815	2802	–
STEP, goal-directed	741	826	2283	–
SEEDEVAL, goal-directed	741	826	2283	–
STEP, shared goal-directed traversal	709	787	2265	–
SEEDEVAL, shared goal-directed traversal	709	787	2265	–
APP with the constructor U, shared forward traversal	803	864	2902	–
`zero_normal` (tagged, native contexts)				
Raw, two introductions (8 rules)	1084	1141	3322	984
N^0 (7 rules)	1073	1136	3272	996
Raw, single introduction (7 rules)	1024	1080	3145	964
Z_6 (6 rules)	1013	1075	3095	977
`reflexive_base` (same format)				
N^0 (7 rules)	1073	1136	3272	996
Z_6, the witness-selected control (6 rules)	1013	1075	3095	977
R (6 rules)	1013	1073	3084	981
`initializer_equality` (closed goals)				

Complete program	Tokens	AST nodes	Bytes	Gzip-9
One-template control	1202	1360	4029	1173
S, selected	1192	1347	3979	1175
S, explicit shape tests	1236	1404	4156	1226
Canonical recursor checker, inherited	1338	1485	4611	1205
Canonical recursor checker, shared traversal	1224	1378	4079	1188
`full_normalization` (goal-directed)				
Raw computations, matched control	895	930	2569	941
Zero normalization, matched control	890	930	2532	945
F, block form	904	962	2546	960
F, expression form	908	965	2529	963
`computation_normal`				
Compact arity envelope (reference)	265	306	516	332
Public goal, baseline	383	442	870	507
Public goal, scope reuse (Section 4.10)	377	433	860	508
Z_6, original	1013	1075	3095	977
Z_6, term as computation certificate	991	1049	3029	971
F, initial	993	1046	3048	965
F, neutral-counter test	992	1046	3044	964
F, explicit fuel	1024	1086	3204	1003
Z_6, goal-directed	972	1004	2982	965
F, goal-directed	982	1008	3063	961
F, goal-directed, fuel	1010	1044	3189	1001

The matched differences. The effect of each change of rules, against the control that shares everything else, is as follows.

- `computational_equality`: E_3 against the matched seven-rule control saves 82 bytes, 49 tokens and 38 AST nodes; gzip grows by 9 bytes. The matched control differs from the archived one only by the same use of the invariant on proved formulas.

- `equality_from_iteration`: E_2 against E_3 saves 5 tokens, 7 AST nodes, 43 bytes and 11 gzip bytes. U against its control with ROOT saves 8 tokens, 12 AST nodes and 53 bytes. U against E_3 has 28 fewer bytes, 20 more tokens and 21 more AST nodes.

- `uniform_step`: C_{Step} against the matched zero-normal control saves 98 bytes, 31 tokens and 47 AST nodes. The delivered file with documentation has 3507 bytes and 1274 gzip bytes. With native contexts the checker with REFL and STEP is smaller than its matched control and larger in tokens and AST nodes than the checker of E_3 (959, 1026, 3039).

- `seed_evaluation`: the change from STEP to SEEDEVAL ties in the helper style and in both goal-directed pairs, and saves 2 bytes and 2 tokens with the shared forward traversal; of the 148 bytes between the 2950-byte and the 2802-byte checker, 146 come from the shared traversal, which benefits both rules. The package reports gzip sizes of 818 and 822 bytes for the goal-directed pair and of 872 and 877 bytes for the shared-traversal pair, STEP first. The checker with the constructor U is larger than its STEP control in all three coordinates.

- `zero_normal`: the deletion of the zero computation rule saves 50 bytes, 11 tokens and 5

AST nodes in each of the two pairs; gzip grows by 12 and 13 bytes. The frozen upstream checkers have 3358 and 3382 bytes.

- `reflexive_base`: R against Z_6 saves 11 bytes and 2 AST nodes with the same number of tokens; gzip grows by 4 bytes.
- `initializer_equality`: S against the one-template control saves 50 bytes, 10 tokens and 13 AST nodes; gzip grows by 2 bytes. The two rows for the canonical recursor checker measure a refactoring of an inherited program and no change of rules.
- `full_normalization`: the expression form of F against the control with zero normalization has 3 fewer bytes, 18 more tokens, 35 more AST nodes and 18 more gzip bytes, and 40 fewer bytes than the raw control. Its auxiliary forward checker has 3100 bytes against 3095 for the forward checker of Z_6.
- `computation_normal`: the scope reuse saves 10 bytes, 6 tokens and 9 AST nodes; gzip grows by 1 byte. For the calculi, the apparent gain of 47 bytes of F over the original checker of Z_6 disappears once the latter also takes a term as its computation certificate: F with the neutral-counter test is then 15 bytes and 1 token larger, 3 AST nodes and 7 gzip bytes smaller. The goal-directed checker of F is larger than that of Z_6, and fuel adds to both.

The checkers. All checkers are single Python modules that import only `json` and `sys`, read one JSON value, use explicit tests in place of `assert`, and accept by printing `ACCEPT` with exit status zero. Proof translators and test runners are not imported and are not trusted. The checkers of `computational_equality`, `equality_from_iteration`, `zero_normal` and `reflexive_base` take an object with the fields `variables`, `context`, `goal` and `proof`, with tagged terms in de Bruijn notation and six proof tags; they check open sequents. Their Eval instruction takes two premise proofs and nothing else. It relies on the invariant that the inference function returns a well-scoped native formula, so that a proved formula of length three is an equation and a term of length four is an iterator; the matched controls use the same invariant. The checkers on zero-normal terms reject every supplied iterator with the literal counter 0, so that a malformed subterm cannot be erased before it is validated, and rebuild terms with the smart constructor during substitution; in `zero_normal` this is one conditional after the rebuilding of a node. The test of `reflexive_base` compares the slices of a validated formula after its tag, which is exact because an equation has three fields and a binder formula two.

The clause checkers of `uniform_step` and `seed_evaluation` take the fields `goal` and `proof` and check closed goals. Bound variables are positive indices and eigenvariables are negative levels; every substituted term has only negative levels free, so the opening of a binder needs no shift. In the helper style the change from Step to SeedEval is two lines: the comparison of the first source with the initializer in place of the predecessor iterate, and the predecessor iterate from the new seed inside the conclusion. The checker of `initializer_equality` implements Seed by one comparison of the two left sides and, failing that, one comparison with the initializer of an iterator; no substitution is performed by that instruction.

The selected checker of `full_normalization` is goal-directed: one recursive traversal checks arities and sorts, renames variables and contracts exposed redexes, and validation is the requirement that the identity action returns its input, with all children checked before a redex is erased. A goal that is a retained assumption or a reflexive equation closes without reading its proof field, in the checker and in its controls. In `computation_normal` a normal term is its own reflexivity certificate, and the fuelled variant adds the field `fuel`.

Executed checks. The packages record the following runs, none with an unexpected outcome; they support the implementations and are not proofs of the theorems. `computational_equality`: 38 inherited fixtures, 14 translation round trips within the bound $9N$, 29 further round trips, 51 adversarial cases and 206 process executions. `equality_from_iteration`: 460 process checks, 62 translation checks, 1280 normalization checks and 320 substitution checks. `uniform_step`: 376 process checks of the clause checker, 100 of the native checkers with 38 translation round trips, and 2104 comparisons of the opening operation. `seed_evaluation`: 398 process checks, 166 translation round trips and 92 rejections, among others, and 83 certificates for the calculus with U. `zero_normal`: 500 process checks, the translation of 64 inherited certificates and 4832 checks of the coherence with substitution. `reflexive_base`: 13 translation cases, 16 malformed inputs and 86 process checks, including two substitutions that turn the guard from false to true. `initializer_equality`: 600 process checks and 31 translation round trips. `full_normalization`: 680 process checks of the goal-directed family and 132 of the forward checker, with the two certificates of Theorem 4.101. `computation_normal`: 272 runs and 9000 generated goals for Theorem 4.104, and a recovered case introduction at the witness 0 with the context $a = b$ and the matrix $I(y, a;\, u.\, b) = I(y, x;\, u.\, 0)$.

4.12 Further questions

This section collects, from all nine packages, the questions that they leave open, the claims that they state without proof and the results that they report as not achieved.

4.12.1 What is not achieved

- No package proves a lower bound or a minimality statement, for rule counts or for checker sizes. Every package states that its checker does not improve on the compact arity-coded checker of the repository (516 bytes; 521 bytes at the snapshot used by `zero_normal`) or on the fully validated profiles of 866, 870 and 788 bytes that have different certificate formats and validation contracts. A port of Eval, of the identity frame, of Seed or of zero normalization to those encodings is not established.

- Moving a computation into the syntax does not by itself reduce the size of a complete checker. The measurements of Section 4.11 for U against E_2, for the constructor U, and for F against Z_6 are mixed or negative, and gzip sizes grow in most matched pairs.

- `seed_evaluation` reports that SeedEval gives no convincing reduction of checker size: two of its three matched pairs tie.

- `zero_normal` examines the rule of common-source congruence, from $s = t$ and $s = u$ infer $h(t) = h(u)$, in place of Tr. With reflexivity it is interderivable with Tr: the instance whose first premise is a reflexivity is congruence, and the instance with the identity template composes two equations with a common left side. An apparent saving of 36 bytes disappears after the same factoring is applied to the control; both programs measure 3322 bytes, 1084 tokens and 1141 AST nodes.

- `computation_normal` recommends that F not replace the zero-normal kernel on the ground of its shorter rule table: the matched checker size does not improve uniformly, normal forms can be enormous, and the uniform normalizer is not primitive recursive (Theorem 4.102).

- The Lean formalization covers the calculi as recorded in the remarks of this chapter. It does not cover, in the forms stated here: the clause calculi $\mathsf{C}_{\mathrm{Step}}$ and $\mathsf{C}_{\mathrm{Seed}}$ with their

clause contexts, the calculus $\mathsf{C}_{\mathrm{App}}$ and the unit normalization, Z_6 and S with the single introduction, the fresh-zero lemma and the compound envelope, the fuelled relation of Theorem 4.103, and any of the Python checkers.

4.12.2 Claims stated without proof

- `seed_evaluation` states that the interpretation in which every literal equation is false obstructs the deletion of reflexivity from the clause calculi, and that moving reflexivity into an initial universal clause still leaves a reflexivity axiom to count. Theorem 4.23 proves the corresponding statement for E_2; the statement for the clause calculi is not proved in the package.

- Theorem 4.85(c) is asserted in the notes of `seed_evaluation` for arbitrary clause contexts, by reference to the completeness of the clause calculus for T_{eq}; the argument given covers contexts of assumption clauses. The general statement requires the completeness of $\mathsf{C}_{\mathrm{Step}}$ for arbitrary clauses.

- `full_normalization` argues that its executable traversal, which fuses substitution with normalization and passes the unevaluated recursive iterator into the substitution of the body, computes $\mathrm{nf}(t[\sigma])$: every step contracts a redex of the materialized instance, termination is Theorem 4.88, and the result is normal. The argument is given in prose and is not carried out as a proof about the program.

4.12.3 Open questions

Question 4.105 (Lower bounds)**.** For a fixed term grammar, certificate language, validation contract and set of primitive operations, what is the least size of a checker of a calculus bi-interpretable with iPRA? No package fixes such a class. `computational_equality` asks for an order of simplicity that separates rule counts, constructors of the syntax, tokens, bytes and certificate sizes; `initializer_equality` and `full_normalization` ask for lower bounds once a class of representations is declared.

Question 4.106 (A smaller basis for arithmetic and equality)**.** Can the interface of reflexivity and one evaluation rule be reduced further while a small normalization layer is counted in full? Theorems 4.22 and 4.23 show that no rule of E_2 can be deleted from that inventory; they are not lower bounds across term languages. `equality_from_iteration` notes that a different basis has to account for the freedom of the initializer, for the reflection of equality (Theorem 1.52) and for both round trips; `reflexive_base` asks whether another arithmetic or logical rule can be removed with a net reduction of the complete checking algorithm.

Question 4.107 (An introduction that is stable under substitution)**.** Eval, Step and Seed are stable under substitution (Theorems 4.21, 4.58 and 4.69), and WI is not (Theorem 4.4). Is there a calculus of six rules in which substitution in derivations is proved by a direct induction, with a single introduction that needs no test on the witness, without enlarging the complete checker and without induction beyond literal equations? The guard of Section 4.5 is preserved and not reflected. `uniform_step` asks whether the logical rules can be merged further while the distinction between coverage and equation induction and the eigenvariable conditions are kept.

Question 4.108 (Sizes of derivations)**.** The congruence of Theorem 4.15 adds eight nodes, that of Theorem 4.61 four, that of Theorem 4.67 four, and the elimination of Root adds two per instance (Theorem 4.12); in each case the labels contain copies of the template and of the endpoints. What are the least numbers of nodes for these fragments, for trees and for

derivations with shared subproofs, and what are the sizes of serialized derivations and the checking times on an arithmetic library? Can sharing reduce the cost without a decoder that is larger than the rule it removes? Can an interface based on the initializer shorten broader families of derivations than the single computations of Theorem 4.78, without the sevenfold expansion of Theorem 4.76(c)?

Question 4.109 (Normalization between zero and full)**.** Is there a normalization between the zero normalization and the full normalization that reduces both the arithmetic interface and the complete trusted implementation? The unit normalization of Section 4.8.2 has a primitive recursive normalizer and a rule without a counter, and its measured checker is larger than its control.

Question 4.110 (Classes of provable bases)**.** Which decidable classes of bases that are derivable in every context give the best exchange between the size of a checker and the length of derivations for the guard of Theorem 4.45? The classes “$r = r$” and “$0 = 0$” of Theorem 4.54 already differ. Does $\mathsf{R}^{\mathrm{raw}}$, which needs no normalization, lower the size of a complete compact checker, and does the guard “$0 = 0$” suffice on raw terms?

Question 4.111 (Other positions of a recursor)**.** Which arithmetic primitives support both the identity and the application of an arbitrary template through one distinguished argument, as the initializer of the scalar iterator does at the counters 0 and 1 (Theorems 4.57 and 4.60)? `initializer_equality` requires equality interpreted by equality and both formula round trips in every such comparison.

Question 4.112 (Narrower induction)**.** `seed_evaluation` asks whether the approach by rigidity of the successor, which derives induction for equations with the right side 0, can be completed to the reflection of equality without importing general equation induction at an intermediate step, and notes that a recursor whose body sees the counter gives a simpler predecessor (Theorem 1.48) and does not by itself supply the reflection argument.

Question 4.113 (Formalization and a common benchmark)**.** Formalize the statements listed above as not covered, in particular the conversion relation with finite witnesses and the fuelled relation, the fresh-zero lemma and the compound envelope next to `F6Prf.dIZero`, and the clause calculi with Step, SeedEval and App. Separately, measure the compact, the goal-directed and the forward checkers under common contracts for the validity of syntax, open sequents, sharing in certificates and the behavior on errors, so that a bound for one encoding is not read as a bound for all presentations.

Chapter 5

Clause calculi

This chapter treats the presentations whose formulas are equations and the paired binder only. The closure binder Q_y is not a formula constructor in them. Its work is done by the context: a context entry is a reusable clause $[A;\, y.\, B(y)]$ (Section 1.9), and the elimination of $D_{x,y}B$ adds a clause where the kernel adds the formula $Q_yB(b,y)$. Section 5.1 relates sequents of the native scalar core N to clause sequents. Section 5.2 gives the five-rule calculus and its equivalence with N. The later sections vary one component at a time: the arithmetic interface (Section 5.3), the introduction rule (Section 5.4), the form of a context entry (Section 5.5), and the number of rule families (Section 5.6). Section 5.8 records a separate arithmetic observation of one of the packages, and Section 5.9 lists the checker sizes.

Every calculus of the chapter has the terms of Section 1.1, the designated contradiction $0 = S0$ and the ordinary reading of Section 1.2. Throughout, $\equiv$ is literal identity of syntax up to the names of bound variables. A matrix $B(x,y)$ displays the two variables that an enclosing $D_{x,y}$ binds, and $B(t,r)$ is the result of substituting t for x and r for y. Substitution avoids capture. A *literal equation* is a formula whose root is an equation; an equation below a binder D is not one. The numeral $S0$ is also written 1. The rules of N are cited by their names in Section 1.3: Hyp, the two computation equations (I0, IS), Tr, QE, DE, the introduction by cases (DI) and the introduction of Q by progress (QI).

5.1 From Q formulas to clauses

This section draws on `raw_clausal`, `clausal_two_case`, `reduced_clause_calculus` and `clause_guards`.

5.1.1 The Q-free form of a formula

A native formula is *Q-free* if it is built from equations by $D_{x,y}$ alone. The Q-free formulas are the formulas of every calculus of this chapter.

Definition 5.1 (Q-free form)**.** The translation τ from native formulas to Q-free formulas is

$$\tau(s=t) := (s=t), \qquad \tau(D_{x,y}B) := D_{x,y}\tau(B), \qquad \tau(Q_yB) := D_{w,y}\tau(B),$$

where each inserted w is fresh and does not occur in B. Terms are unchanged. The inclusion of Q-free formulas into native formulas is written j and is usually left implicit. For a matrix $B(y)$ and a fresh w the *closure abbreviation* is

$$q_yB(y) := D_{w,y}B(y), \qquad w \notin \mathrm{FV}(B).$$

The translation preserves free variables, commutes with substitution up to the names of bound variables, sends equations to equations and all other formulas to formulas with root D, and satisfies $\tau j F \equiv F$ for every Q-free F. The ordinary reading of $q_y B$ is $\exists w\,(B(0) \to \forall y\, B(y))$, which is equivalent to $B(0) \to \forall y\, B(y)$ because the domain is inhabited by 0. In the Lean development τ is `unQ`.

Lemma 5.2 (The round trip in `N`)**.** *The following hold in* `N`*, in every context.*

(i) $Q_y B \dashv\vdash D_{w,y} B$ *for* $w \notin \mathrm{FV}(B)$.

(ii) If $\Gamma, B \vdash B'$ *and* $\Gamma, B' \vdash B$*, where the variables* x, y *are not free in* Γ*, then* $\Gamma, D_{x,y} B \vdash D_{x,y} B'$ *and* $\Gamma, Q_y B \vdash Q_y B'$*, and conversely.*

(iii) $A \dashv\vdash j\tau A$ *for every native formula* A.

Proof. (i) From $Q_y B$: under the assumption $B(0)$ the rule QE gives $B(Sz)$, so the introduction by cases at an arbitrary witness gives $D_{w,y} B$. From $D_{w,y} B$: the rule DE opens $Q_y B(b, y)$, which is $Q_y B$ because w does not occur in B, and Hyp closes the branch.

(ii) Assume $D_{x,y} B$. Apply DE at a fresh b; the branch has the assumption $Q_y B(b, y)$. Introduce $D_{x,y} B'$ by cases at the witness b. Its branch assumes $B'(b, 0)$. Substitution in the derivation of $\Gamma, B' \vdash B$ (Section 1.4) and cut give $B(b, 0)$, QE gives $B(b, Sz)$, and the substituted derivation of $\Gamma, B \vdash B'$ gives $B'(b, Sz)$. For Q, pass from $Q_y B$ to $D_{w,y} B$ by (i), apply the case just proved with the unused variable w in the place of x, and return by (i). The converse statements are symmetric.

(iii) Induction on A. Equations are unchanged. For $A \equiv D_{x,y} B$ use (ii) and the induction hypothesis for B. For $A \equiv Q_y B$ use (ii) for Q and then (i). □

5.1.2 Clause occurrences and their origins

A context of a clause calculus is a finite list of clause *occurrences* $[A;\, y.\, B(y)]$ with $y \notin \mathrm{FV}(A)$ (Section 1.9). The clause is a context object and not a formula. Its reading is $A \to \forall y\, B(y)$. A rule that uses a clause selects an occurrence, so two occurrences with the same text are distinct. An ordinary assumption F is the constant clause

$$\mathrm{ord}(F) := [\, 0 = 0;\, v.\, F\,], \qquad v \notin \mathrm{FV}(F),$$

and Γ, F abbreviates $\Gamma, \mathrm{ord}(F)$. For a list Δ of formulas, $\mathrm{ord}(\Delta)$ applies ord to each member.

In a derivation that starts from ordinary assumptions a clause occurrence arises in two ways only.

Definition 5.3 (Origins)**.** An *assumption occurrence* is an occurrence of the form $\mathrm{ord}(G)$. An *opened occurrence* is an occurrence of the form $[B(b, 0);\, y.\, B(b, y)]$ for a matrix $B(x, y)$ and a variable b. An *annotated* context is a context in which each occurrence is marked as one of the two and has the corresponding form. The *native image* $\Gamma^{\natural}$ of an annotated context replaces each assumption occurrence $\mathrm{ord}(G)$ by the formula G and each opened occurrence $[B(b, 0);\, y.\, B(b, y)]$ by the formula $Q_y B(b, y)$.

The mark belongs to the occurrence. A clause such as $[0 = 0;\, y.\, 0 = 0]$ has both forms, and either mark may be given to it. The rules of the calculi below add an opened occurrence at each elimination and an assumption occurrence at each introduction, so every context reachable from $\mathrm{ord}(\Delta)$ has a canonical annotation.

5.2 The five-rule calculus

This section draws on `raw_clausal`; the same lemmas are restated in `rootfree_clausal`, `reduced_clause_calculus`, `clausal_two_case` and `uniform_clauses`.

5.2.1 Rules

The evaluation operation $\Phi(s,t,b)$ on raw terms is that of Theorem 4.8. Its three rows are called the *root row* ($b \equiv s$, value t), the *zero row* ($b \equiv I(0,a;\,u.\,h)$, $s \equiv a$, value t) and the *successor row* ($b \equiv I(Sn,a;\,u.\,h)$, $s \equiv I(n,a;\,u.\,h)$, value $h[t/u]$); they are disjoint, and the three cases of the rule Eval that they give are Root, Eval_0 and Eval_S of Theorem 4.9. The operation Φ_2 is Φ without the root row. Chapter 4 studies the rule with contexts of native formulas; here it is combined with clause contexts.

Definition 5.4 (The calculus C_3)**.** Formulas are Q-free, contexts are lists of clause occurrences, and the rules are the following five.

$$\frac{}{\Gamma \vdash s = s}\ \textsc{Refl} \qquad \frac{\Gamma \vdash s = t \quad \Gamma \vdash b = r}{\Gamma \vdash r = \Phi(s,t,b)}\ \textsc{Eval}$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash B(t)}\ \textsc{Use} \qquad \frac{\Gamma \vdash D_{x,y}B(x,y) \quad \Gamma, [B(b,0);\, y.\,B(b,y)] \vdash F}{\Gamma \vdash F}\ \textsc{Elim}$$

$$\frac{\Gamma, B(t,\eta) \vdash B(t,Sz)}{\Gamma \vdash D_{x,y}B(x,y)}\ \textsc{Intro}$$

Eval requires $\Phi(s,t,b)$ to be defined. Use requires an occurrence $[A;\,y.\,B(y)]$ in Γ and allows every term t. In Elim the variable b is fresh for Γ, F and B. In Intro the variable z is fresh for Γ, t and B, the witness t is an arbitrary term, and η is the term $\eta_{B,t}(z)$ of the witness-selected introduction (Theorem 2.15): it is z if B is a literal equation and $t \equiv 0$ (*progress mode*) and 0 otherwise (*case mode*). The calculus C is C_3 with Φ_2 in the place of Φ.

There is no hypothesis rule, no computation axiom, no transport rule, no induction rule and no structural rule. The three rows of Φ and the mode test of Intro are part of the specification. The calculus C is `CPrf` in the Lean development.

5.2.2 Derived rules

The lemmas of this subsection hold in C_3 and in C under arbitrary clause contexts.

Lemma 5.5 (Structural operations)**.** *An ordinary assumption can be used: $\Gamma, F \vdash F$. Weakening by clause occurrences and injective renaming of free variables are admissible. Cut for ordinary assumptions is admissible: if $\Gamma \vdash F$ and $\Gamma, \mathrm{ord}(F), \Gamma' \vdash A$ then $\Gamma, \Gamma' \vdash A$.*

Proof. Refl proves $0 = 0$, and Use on $[0 = 0;\, v.\,F]$ at any term proves F. Weakening and injective renaming are inductions on the derivation after the eigenvariables are made fresh; an injective renaming preserves the root of a matrix and the test $t \equiv 0$, so it preserves the mode of Intro, and it preserves the literal matches of Φ.

For cut, induct on the derivation that has the distinguished occurrence $\mathrm{ord}(F)$. At a Use that selects this occurrence, replace the node by the given derivation of F, weakened to the local context; the derivation of its premise $0 = 0$ is discarded. At every other node transform the premises and reapply the rule. The eigenvariables are made fresh first. No substitution of a term in a derivation is performed. □

Lemma 5.6 (Equality and computation)**.** *Refl and the zero and successor rows of Eval derive, over every clause context, symmetry and transitivity of equality, the inference Root, the two computation equations*

$$I(0,a;\,u.\,h) = a, \qquad I(Sn,a;\,u.\,h) = h[I(n,a;\,u.\,h)/u],$$

congruence in every term template, and transport in a term and in a literal equation:

$$\frac{s = t}{h[s/u] = h[t/u]} \qquad \frac{s = t \quad h[s/u] = r}{r = h[t/u]} \qquad \frac{s = t \quad a[s/u] = b[s/u]}{a[t/u] = b[t/u]}$$

with no restriction on s.

Proof. The derivations of Section 4.2.3 (Theorems 4.11 and 4.13 to 4.16) consist of REFL, EVAL_0, EVAL_S and the given premises. They contain no rule that reads the context, so they are derivations of C and of C_3 over every clause context. Transport in an equation is the one-term transport with the template a, which gives $b[s/u] = a[t/u]$, followed by the one-term transport with the template b. □

Lemma 5.7 (All introductions by cases)**.** *For every matrix B and every witness t the rule*

$$\frac{\Gamma, B(t,0) \vdash B(t,Sz)}{\Gamma \vdash D_{x,y}B(x,y)} \qquad (z \text{ fresh})$$

is admissible.

Proof. INTRO has this form unless B is a literal equation and $t \equiv 0$. That case is repaired as in the proof of Theorem 2.17, at the second name $c \equiv I(0,0;\, u.\, 0)$ of zero (Theorem 2.16): in the context $\Gamma, B(c,0)$ transport along $c = 0$ (Theorem 5.6) gives $B(0,0)$, the cut of Theorem 5.5 with the given derivation gives $B(0,Sz)$, transport along $0 = c$ gives $B(c,Sz)$, and INTRO at the witness c is in case mode. The cut is the cut for a constant clause; only transport in an equation is used. □

Lemma 5.8 (Transport in every formula)**.** *For every Q-free formula F with a distinguished variable v, from $\Gamma \vdash s = t$ and $\Gamma \vdash F[s/v]$ one derives $\Gamma \vdash F[t/v]$.*

Proof. Induction on F. Equations are Theorem 5.6. Let $F \equiv D_{x,y}B(v,x,y)$. Apply ELIM to $F[s/v]$ at a fresh b, which adds $[B(s,b,0);\, y.\, B(s,b,y)]$. Introduce $F[t/v]$ by Theorem 5.7 at the witness b. Its branch assumes $B(t,b,0)$. The induction hypothesis along $t = s$ gives $B(s,b,0)$, USE at Sz gives $B(s,b,Sz)$, and the induction hypothesis along $s = t$ gives $B(t,b,Sz)$. Each use of the induction hypothesis concerns the matrix B. □

Lemma 5.9 (The three closure interfaces)**.** *With $q_yB \equiv D_{w,y}B$, w fresh, the following are derivable.*

(i) From $\Gamma, B(0) \vdash B(Sz)$, z fresh, derive $\Gamma \vdash q_yB$.

(ii) From $\Gamma \vdash q_yB$ and $\Gamma \vdash B(0)$ derive $\Gamma \vdash B(t)$ for every term t.

(iii) If Γ contains an occurrence $[B(0);\, y.\, B(y)]$ then $\Gamma \vdash q_yB$.

(iv) For a literal equation $E(y)$: from $\Gamma, E(z) \vdash E(Sz)$, z fresh, derive $\Gamma \vdash q_yE$.

Proof. (i) is Theorem 5.7 with the unused variable w in the place of x. For (ii) apply ELIM to q_yB; the fresh witness does not occur in B, so the added clause is $[B(0);\, y.\, B(y)]$, and USE with the given derivation of $B(0)$ proves $B(t)$. For (iii), under the assumption $B(0)$ use the retained clause at Sz and apply (i). For (iv) apply INTRO to the matrix $E(y)$, in which w does not occur, at the witness 0; the matrix is a literal equation and the witness is literally 0, so the rule is in progress mode. □

Corollary 5.10 (Equation induction and numerical cases)**.** *For a literal equation $E(y)$, from $\Gamma \vdash E(0)$ and $\Gamma, E(z) \vdash E(Sz)$ one derives $\Gamma \vdash E(t)$. For an arbitrary formula $F(y)$, from $\Gamma \vdash F(0)$ and $\Gamma \vdash F(Sz)$, z fresh, one derives $\Gamma \vdash F(t)$. For a literal equation $E(x,y)$ and a term t, from $\Gamma, E(t,z) \vdash E(t,Sz)$ one derives $\Gamma \vdash D_{x,y}E$.*

Proof. The first two statements are (iv) and (ii), respectively (i) and (ii), of Theorem 5.9. For the third, (iv) gives $q_y E(t, y)$, and Theorem 5.7 at the witness t gives $D_{x,y}E$, since under the assumption $E(t, 0)$ interface (ii) gives $E(t, Sz)$. □

Progress is available for literal equations only. Extending the progress mode to matrices with root D would give induction for arbitrary formulas by the same argument and would be a different theory.

5.2.3 Equivalence with the native scalar core

Theorem 5.11 (The five-rule calculus and N)**.** *Let* X *be* C_3 *or* C*.*

(i) If Γ *is an annotated clause context and* $\Gamma \vdash_{\mathsf{X}} F$*, then* $\Gamma^{\natural} \vdash_{\mathsf{N}} F$*.*

(ii) If Γ *is a native context and* $\Gamma \vdash_{\mathsf{N}} A$*, then* $\mathrm{ord}(\tau\Gamma) \vdash_{\mathsf{X}} \tau A$*.*

(iii) For Q-free Δ *and* F*:* $\mathrm{ord}(\Delta) \vdash_{\mathsf{X}} F$ *if and only if* $\Delta \vdash_{\mathsf{N}} F$*.*

(iv) For native Γ *and* A*:* $\mathrm{ord}(\tau\Gamma) \vdash_{\mathsf{X}} \tau A$ *if and only if* $\Gamma \vdash_{\mathsf{N}} A$*.*

All translations are effective on finite derivations and are the identity on terms.

Proof. (i) Induction on the derivation. N derives reflexivity, symmetry, transitivity and Leibniz replacement (Section 1.4). Refl is reflexivity. The three rows of Eval are derived in N from I0, IS and Tr as in the proof of Theorem 4.18. A Use of an assumption occurrence $\mathrm{ord}(G)$ is Hyp. A Use of an opened occurrence $[B(b, 0); y.\, B(b, y)]$ is QE applied to the assumption $Q_y B(b, y)$ and the translated derivation of $B(b, 0)$. Elim is DE, which adds $Q_y B(b, y)$, the image of the opened occurrence. Intro in case mode is the introduction by cases, with the new occurrence marked as an assumption. In progress mode the matrix is a literal equation $E(x, y)$ and $t \equiv 0$; the translated premise is $\Gamma^{\natural}, E(0, z) \vdash E(0, Sz)$, the introduction of Q by progress gives $Q_y E(0, y)$, and the introduction by cases at the witness 0 gives $D_{x,y}E$, because under the assumption $E(0, 0)$ the rule QE gives $E(0, Sz')$.

(ii) Induction on the derivation in N. Hyp becomes Refl at 0 followed by Use on the constant clause. The computation equations and Tr are Theorem 5.6. QE is interface (ii) of Theorem 5.9, since $\tau(Q_y B) \equiv q_y \tau(B)$ and τ commutes with substitution. The introduction by cases is Theorem 5.7. The introduction of $Q_y E$ by progress is interface (iv), whose conclusion is $q_y E \equiv \tau(Q_y E)$. For DE with major premise $D_{x,y}B$ and branch $\Gamma, Q_y B(b, y) \vdash A$, translate the major premise and apply Elim, which adds $[\tau B(b, 0); y.\, \tau B(b, y)]$. In the extended context interface (iii) derives $q_y \tau B(b, y)$, which is $\tau(Q_y B(b, y))$. Weaken the translated branch to the extended context and cut this derivation in for its ordinary assumption (Theorem 5.5). Elim then concludes τA.

(iii) From left to right use (i) with the canonical annotation, for which $\mathrm{ord}(\Delta)^{\natural} = \Delta$. From right to left use (ii); τ is the identity on Q-free formulas.

(iv) From right to left is (ii). From left to right, (i) gives $\tau\Gamma \vdash_{\mathsf{N}} \tau A$, and Theorem 5.2(iii) with cut in N replaces $\tau\Gamma$ by Γ and τA by A. □

Corollary 5.12 (Both formula composites)**.** *The composite* τj *is the identity on Q-free formulas, and* $A \dashv\vdash_{\mathsf{N}} j\tau A$ *for every native formula* A*. Both term translations are identities. In particular* $\vdash_{\mathsf{C}} 0 = S0$ *if and only if* $\vdash_{\mathsf{N}} 0 = S0$*.*

Remark 5.13. In the Lean development (iv) is `cprf_iff_nprf` and (iii) is `cprf_iff_nprf_of_qfree`, for the calculus C (`NativeClausal`). There the clause calculus is defined over all native formulas, and `CPrf.qfree` states that it derives only Q-free formulas from Q-free clauses. `raw_clausal` states the theorem for C_3 against the six-rule calculus with Refl and Eval of

`computational_equality`; `clausal_two_case` and `uniform_clauses` state it for C against the same calculus with two evaluation rows, and `clause_guards` against the six-rule calculus with the diagonal introduction. The proof above compares with N directly; the cases are the same. The Lean proof of completeness translates the six-rule calculus rule by rule.

Proposition 5.14 (Substitution)**.** *If $\Gamma \vdash_{\mathsf{X}} F$ and σ is a substitution of terms for free variables, then $\Gamma\sigma \vdash_{\mathsf{X}} F\sigma$, for X either of C_3, C and every clause context Γ. The rule at a node of the derivation is not always preserved.*

Proof. Induction on the derivation, with eigenvariables made fresh for σ. Refl is immediate, and $\Phi(s\sigma, t\sigma, b\sigma) \equiv \Phi(s,t,b)\sigma$ by the same row (Theorem 4.21). Use and Elim commute with σ. An Intro in progress mode keeps its equation root and its literal witness 0. An Intro in case mode whose matrix has root D, or whose witness t has $t\sigma \not\equiv 0$, remains in case mode. In the remaining case the matrix is a literal equation, $t \not\equiv 0$ and $t\sigma \equiv 0$; the substituted premise is the premise of an introduction by cases at the witness 0, and Theorem 5.7 applies. □

A substitution can thus change the mode of an introduction, and the derivation after substitution may contain the nodes of Theorem 5.7 in the place of one Intro. Theorem 5.37 gives a calculus in which the tree is preserved.

Remark 5.15 (Clauses that are not of the two origins)*.* Theorem 5.11(i) concerns annotated contexts. An arbitrary clause $[A;\, y.\, B]$ supplied from outside is represented in N by the derived implication and universal quantifier of the native logical interface (Theorem 1.78), as the formula that reads $A \to \forall y\, B$, and Use becomes the elimination of the implication followed by the elimination of the universal quantifier. The equivalences (iii) and (iv) do not need such clauses. They cover every sequent with ordinary assumptions and not only closed theorems.

5.3 Variants of the arithmetic interface

This section draws on `rootfree_clausal`, `reduced_clause_calculus`, `clausal_two_case`, `clausal_single_evaluation`, `raw_clausal`, `uniform_clauses` and `clause_contexts`; each subsection names its sources. The variants are: an evaluation rule with two rows, an evaluation rule with one scheme on zero-normal terms, and reflexivity stated as a clause, on raw and on zero-normal terms.

5.3.1 Two evaluation cases

This subsection draws on `rootfree_clausal`, `reduced_clause_calculus` and `clausal_two_case`, which present the same calculus C. Its two evaluation rows are the rules $\textsc{Eval}_0$ and $\textsc{Eval}_S$ of Theorem 4.9. The root row is derived by the identity frame of Theorem 4.11: with $K(s) \equiv I(0, s;\, u.\, 0)$, the derivation

$$\mathrm{RC}(p,q) := \mathrm{Ev}_0(p,\ \mathrm{Ev}_0(q,\ \mathrm{R}_{K(s)}))$$

of Theorem 4.12 turns $p : s = t$ and $q : s = r$ into $r = t$ by two instances of $\textsc{Eval}_0$ and one Refl leaf. It contains no rule that reads the context, so it is a derivation of C over every clause context. The body of $K(s)$ is never evaluated; any body serves.

Theorem 5.16 (Deletion of the root row)**.** *For every clause context Γ and every formula F,*

$$\Gamma \vdash_{\mathsf{C}_3} F \quad\Longleftrightarrow\quad \Gamma \vdash_{\mathsf{C}} F.$$

The translation from C_3 to C leaves the end sequent unchanged and turns a derivation with N nodes, k of them uses of the root row, into a derivation with exactly $N + 2k \le 3N$ nodes. The converse translation is the inclusion.

Proof. Every rule of C is a rule of C_3. Conversely replace each use of the root row by RC and copy every other node, as in Theorem 4.12; the count is the count of that theorem. What the clause setting adds is that the replacement is made under arbitrary retained clauses, also those opened inside ELIM and INTRO: no variable is bound or freed, and the matrices and witnesses of INTRO are unchanged, so its modes and all freshness conditions are unchanged. □

The count is of inference nodes and bounds no serialized size (Theorem 4.12).

Proposition 5.17 (Three new nodes are necessary)**.** *Let x, y, z be distinct variables, and consider the fragment with the rules* REFL, EVAL_0 *and* EVAL_S *and with two given leaves $p : x = y$ and $q : x = z$. Every derivation of $z = y$ in this fragment contains at least two* EVAL *nodes and one* REFL *leaf.* $\mathrm{RC}(p, q)$ *attains the bound.*

Proof. The conclusion $z = y$ is not p, not q and not reflexive, so the last inference is an EVAL. In both rows the second premise has the form $b = r$ with b an iterator, and the conclusion has r as its left side. The second premise of the last inference is therefore $b = z$ with b an iterator. It is not p or q, whose left side is the variable x, and it is not a REFL leaf, since an iterator is not the variable z. So it ends in another EVAL. Follow second premises upwards from there. Each has an iterator as its left side, so the path does not end at p or q; it ends at a REFL leaf. □

The bound concerns this fragment and this cost convention. It does not bound derivations that use the logical rules or previously derived lemmas, and it is not a bound on the size of a checker.

Proposition 5.18 (Neither remaining row is redundant)**.** *In* C *without* EVAL_0 *the equation $I(0, 0; u.\, 0) = 0$ is not derivable. In* C *without* EVAL_S *the equation $I(S0, 0; u.\, Su) = S0$ is not derivable.*

Proof. The two interpretations of Theorem 4.22 apply, with a clause read as $A \to \forall y\, B(y)$: under any compositional interpretation of the iterator in the standard natural numbers REFL, USE, ELIM and both modes of INTRO are sound, each interpretation makes the remaining row sound, and the displayed equations are false, the second because its left side has the value 0. □

Remark 5.19. The three packages state Theorem 5.16 with the frame $I(0, s; u.\, 0)$; `uniform_clauses`, `clause_guards`, `one_body` and `prefix_clauses` use $I(0, s; u.\, u)$. The frame argument is that of `equality_from_iteration` (Theorem 4.11), applied inside the clause calculus. Theorem 5.17 is in `reduced_clause_calculus` and, in outline, in `clausal_two_case`; Theorem 5.18 is in `clausal_two_case`. `rootfree_clausal` also gives the direct proof of Theorem 5.14.

From here on Φ_2 has two rows, and the lemmas of Section 5.2 are used for C.

5.3.2 Zero-normal terms and one evaluation scheme

This subsection draws on `clausal_single_evaluation`, on the companion calculi of `reduced_clause_calcul` and `rootfree_clausal`, and on `clause_contexts`. The zero computation $I(0, a; u.\, h) = a$ is moved from the rules into the syntax.

Zero-normal terms, the smart constructor $\overline{I}$ and the normalization nf_0 are those of Theorem 4.32, and normal substitution is the normalizing substitution $t\langle s/x\rangle := \mathrm{nf}_0(t[s/x])$ of Theorem 4.25. The map nf_0 acts on formulas, clauses and contexts through their terms, and $B\langle t, r\rangle$ is the corresponding instance of a matrix. Two facts are used below. Normalization is coherent with substitution,

$$\mathrm{nf}_0(t\sigma) \equiv \mathrm{nf}_0\big(\mathrm{nf}_0(t)\,(\mathrm{nf}_0 \circ \sigma)\big)$$

(Theorem 4.34), so normal substitutions compose (Theorem 4.26). And N derives $t = \mathrm{nf}_0(t)$ for every raw term t (Theorem 4.6), so that $\Gamma, F \vdash_{\mathsf{N}} \mathrm{nf}_0(F)$ and $\Gamma, \mathrm{nf}_0(F) \vdash_{\mathsf{N}} F$ for every formula F and context Γ (Theorems 4.27 and 4.35).

Definition 5.20 (The calculus C^0)**.** Terms, formulas and clauses are zero-normal, contexts and the notion of origin are as before, and every substitution performed by a rule is normal substitution. The rules are

$$\frac{}{\Gamma \vdash s = s}\ \textsc{Refl} \qquad \frac{\Gamma \vdash A}{\Gamma \vdash B\langle t/y\rangle}\ \textsc{Use}$$

$$\frac{\Gamma \vdash \overline{I}(n, a;\, u.\, h) = t \quad \Gamma \vdash I(Sn, a;\, u.\, h) = r}{\Gamma \vdash r = h\langle t/u\rangle}\ \textsc{Eval}$$

$$\frac{\Gamma \vdash D_{x,y}B \quad \Gamma, [B\langle b, 0\rangle;\, y.\, B\langle b, y\rangle] \vdash F}{\Gamma \vdash F}\ \textsc{Elim}$$

$$\frac{\Gamma, B\langle t, \eta\rangle \vdash B\langle t, Sz\rangle}{\Gamma \vdash D_{x,y}B}\ \textsc{Intro}$$

with the side conditions of Theorem 5.4; in particular η is z if B is a literal equation and $t \equiv 0$, and 0 otherwise.

There is one evaluation scheme, the rule Step of Theorem 4.66; it has no root row and no zero row. The calculus is $\mathsf{C5}$ of `clausal_single_evaluation`, it is `ZCPrf` in the Lean development, and it coincides with the calculus $\mathsf{C}_{\mathrm{Step}}$ that Theorem 4.72 takes from `uniform_step`; Section 4.7.2 derives the native rules in it for that package and proves statement (iv) of Theorem 5.22 below through T_{eq}.

Lemma 5.21 (Derived rules of C^0)**.** *The following hold in C^0 under arbitrary clause contexts.*

(i) The structural operations of Theorem 5.5.

(ii) From $p : s = t$ and $q : s = r$ one derives $r = t$ by two Eval nodes and one Refl leaf. Hence symmetry and transitivity.

(iii) The successor computation $I(Sn, a;\, u.\, h) = h\langle \overline{I}(n, a;\, u.\, h)/u\rangle$.

(iv) From $p : s = t$ one derives $h\langle s/u\rangle = h\langle t/u\rangle$ by two Eval nodes and two Refl leaves, with p used once.

(v) From $p : s = t$ and $q : h\langle s/u\rangle = r$ one derives $r = h\langle t/u\rangle$ by four Eval nodes and three Refl leaves, with p and q used once each. Two such steps give transport in an equation.

(vi) The introduction by cases of Theorem 5.7 for every matrix and every witness.

(vii) Transport in every formula (Theorem 5.8), the closure interfaces (Theorem 5.9) and Theorem 5.10.

Proof. (i) The proofs of Theorem 5.5 apply; injective renaming preserves normal terms.

(ii) to (v) The rule Eval of C^0 is the rule Step of Theorem 4.66, and the derivations of Theorem 4.67 consist of Refl, Step and the given premises, so they are derivations of C^0 over every clause context: (ii) is the pivot with the frame $I(S0, s;\, u.\, u)$, with symmetry and transitivity; (iii) is the successor computation; (iv) is the congruence; (v) is the transport. Replacing one node of a one-term transport rule by the derivation of (v) adds six nodes and duplicates neither premise.

(vi) Only a literal equation at the witness 0 needs an argument. Let $Z \equiv I(S0, 0;\, u.\, 0)$. It is zero-normal and is not literally 0 (Theorem 4.40), and Eval on the reflexivities of 0 and

of Z proves $Z = 0$. The proof of Theorem 5.7 applies with Z in the place of c. The raw term $I(0, 0;\, u.\, 0)$ cannot be used: it normalizes to 0.

(vii) The proofs of Theorems 5.8 and 5.9 use only the items above, USE, ELIM and INTRO. □

Theorem 5.22 (The zero-normal calculus and N)**.** *(i) For every raw clause context* Γ *and raw formula* F*: if* $\Gamma \vdash_{\mathsf{C}} F$ *then* $\mathrm{nf}_0(\Gamma) \vdash_{\mathsf{C}^0} \mathrm{nf}_0(F)$*.*

(ii) If Γ *is an annotated zero-normal context and* $\Gamma \vdash_{\mathsf{C}^0} F$*, then* $\Gamma^\natural \vdash_{\mathsf{N}} F$*. Here an opened occurrence is* $[B\langle b, 0\rangle;\, y.\, B\langle b, y\rangle]$ *and its native image is* $Q_y B(b, y)$*.*

(iii) For raw Q-free Δ *and* F*:* $\mathrm{ord}(\mathrm{nf}_0\, \Delta) \vdash_{\mathsf{C}^0} \mathrm{nf}_0(F)$ *if and only if* $\Delta \vdash_{\mathsf{N}} F$*. In particular, for zero-normal* Δ *and* F*,* $\mathrm{ord}(\Delta) \vdash_{\mathsf{C}^0} F$ *if and only if* $\Delta \vdash_{\mathsf{N}} F$*.*

(iv) For native Γ *and* A*:* $\mathrm{ord}(\mathrm{nf}_0\, \tau\Gamma) \vdash_{\mathsf{C}^0} \mathrm{nf}_0(\tau A)$ *if and only if* $\Gamma \vdash_{\mathsf{N}} A$*.*

Proof. (i) Induction on the derivation in C, normalizing every sequent. REFL is REFL. For $\textsc{Eval}_0$ the normalized premises are $\mathrm{nf}_0(a) = \mathrm{nf}_0(t)$ and $\mathrm{nf}_0(a) = \mathrm{nf}_0(r)$, since $\mathrm{nf}_0(I(0, a;\, u.\, h)) \equiv \mathrm{nf}_0(a)$, and Theorem 5.21(ii) gives $\mathrm{nf}_0(r) = \mathrm{nf}_0(t)$. For $\textsc{Eval}_S$ the normalized premises are

$$\overline{I}(\mathrm{nf}_0\, n, \mathrm{nf}_0\, a;\, u.\, \mathrm{nf}_0\, h) = \mathrm{nf}_0(t), \qquad I(S\, \mathrm{nf}_0\, n, \mathrm{nf}_0\, a;\, u.\, \mathrm{nf}_0\, h) = \mathrm{nf}_0(r),$$

because the counter $S\, \mathrm{nf}_0(n)$ is not 0. EVAL of C^0 gives $\mathrm{nf}_0(r) = \mathrm{nf}_0(h)\langle \mathrm{nf}_0(t)/u\rangle$, which is $\mathrm{nf}_0(r) = \mathrm{nf}_0(h[t/u])$ by Theorem 4.34. A USE of $[A;\, y.\, B]$ at t becomes a USE of $[\mathrm{nf}_0\, A;\, y.\, \mathrm{nf}_0\, B]$ at $\mathrm{nf}_0(t)$, with conclusion $\mathrm{nf}_0(B)\langle \mathrm{nf}_0(t)/y\rangle \equiv \mathrm{nf}_0(B[t/y])$. For ELIM the normal form of the added clause $[B(b, 0);\, y.\, B(b, y)]$ is $[\mathrm{nf}_0(B)\langle b, 0\rangle;\, y.\, \mathrm{nf}_0(B)\langle b, y\rangle]$, the clause that ELIM of C^0 adds for $D_{x,y}\, \mathrm{nf}_0(B)$. For INTRO with matrix B and witness t the normalized premise is $\mathrm{nf}_0(\Gamma), \mathrm{nf}_0(B)\langle \mathrm{nf}_0\, t, \eta\rangle \vdash \mathrm{nf}_0(B)\langle \mathrm{nf}_0\, t, Sz\rangle$. The matrix $\mathrm{nf}_0(B)$ is a literal equation exactly when B is one. If the mode of INTRO for $(\mathrm{nf}_0\, B, \mathrm{nf}_0\, t)$ is the mode for (B, t), apply INTRO of C^0. The modes differ only when B is a literal equation, $t \not\equiv 0$ and $\mathrm{nf}_0(t) \equiv 0$. Then the source is in case mode, the normalized premise is the premise of an introduction by cases at the witness 0, and Theorem 5.21(vi) applies.

(ii) Induction on the derivation, as in Theorem 5.11(i). REFL is reflexivity. For EVAL, N derives $I(n, a;\, u.\, h) = \overline{I}(n, a;\, u.\, h)$ by the zero computation or by reflexivity, hence $I(n, a;\, u.\, h) = t$ from the first premise; the successor computation, Leibniz replacement and Theorem 4.27 give $I(Sn, a;\, u.\, h) = h[t/u] = \mathrm{nf}_0(h[t/u])$, and the second premise gives $r = h\langle t/u\rangle$. A USE of an assumption occurrence is HYP. For a USE of an opened occurrence the translated premise is $B\langle b, 0\rangle$; Theorem 4.27 gives $B(b, 0)$, QE on $Q_y B(b, y)$ gives $B(b, t)$, and Theorem 4.27 gives $B\langle b, t\rangle$. ELIM is DE. For INTRO the translated premise is $\Gamma^\natural, B\langle t, \eta\rangle \vdash B\langle t, Sz\rangle$; Theorem 4.27 and cut in N give $\Gamma^\natural, B(t, \eta) \vdash B(t, Sz)$, and the two modes are treated as in Theorem 5.11(i).

(iii) If $\Delta \vdash_{\mathsf{N}} F$ then $\mathrm{ord}(\Delta) \vdash_{\mathsf{C}} F$ by Theorem 5.11(iii), and (i) gives $\mathrm{ord}(\mathrm{nf}_0\, \Delta) \vdash_{\mathsf{C}^0} \mathrm{nf}_0(F)$, because $\mathrm{nf}_0(\mathrm{ord}(G)) \equiv \mathrm{ord}(\mathrm{nf}_0\, G)$. Conversely (ii) gives $\mathrm{nf}_0(\Delta) \vdash_{\mathsf{N}} \mathrm{nf}_0(F)$, and Theorem 4.27 with cut gives $\Delta \vdash_{\mathsf{N}} F$.

(iv) Combine (iii) for $\tau\Gamma$ and τA with Theorem 5.2(iii). □

Corollary 5.23 (Substitution)**.** *Normal substitution of terms for free variables is admissible in* C^0 *for sequents with ordinary assumptions.*

Proof. By Theorem 5.22(iii), substitution in N (Section 1.4) and $\mathrm{nf}_0(F\sigma) \equiv \mathrm{nf}_0(F)\langle \mathrm{nf}_0 \circ \sigma\rangle$ from Theorem 4.34. A witness may become the literal 0 under the substitution, so the rule at a node is not always preserved. □

Remark 5.24. In the Lean development (iv) is `zcprf_iff_nprf`, (iii) is `zcprf_iff_nprf_of_qfree` and the translation of (i) is `CPrf.toZC` (`NativeZeroClausal`). `clausal_single_evaluation`

proves (iii) for zero-normal sequents by comparison with the kernel N^0 on zero-normal terms (Theorem 4.38); the cases are those of Theorem 5.11 with Theorem 5.21 in the place of the lemmas of Section 5.2. The proof above normalizes derivations of C, which is the route of the companion of `reduced_clause_calculus` and of the Lean proof. The change of mode in (i) and its repair at the witness $I(S0, 0; u.\, 0)$ are stated in `reduced_clause_calculus` and `clausal_single_evaluation`. The count in Theorem 5.21(iv) and (v) is from `clausal_single_evaluation`; the congruence of Theorem 5.6 has four evaluations because it joins two separately evaluated endpoints. `rootfree_clausal` states Theorem 5.21(ii) to (iv) for its companion, whose introduction is that of Section 5.4.3.

5.3.3 Reflexivity as one fixed clause

This subsection draws on `raw_clausal` for the obstruction and on `uniform_clauses` for the calculus.

Theorem 5.25 (Closed seeds do not replace reflexivity)**.** *Let S be a finite set of closed equations that are true in the standard natural numbers, and let $M \subseteq \mathbb{N}$ be the finite set of values of their sides. Let C_3^S be C_3 with* Refl *deleted and the members of S added as axioms in every context. If $k \notin M$, then C_3^S does not derive $S^k0 = S^k0$ from the empty context. In particular the calculus with the single axiom $0 = 0$ in the place of* Refl *does not derive $S0 = S0$, and the calculus with* Refl *deleted derives no equation from the empty context.*

Proof. Evaluate terms in the standard natural numbers, the iterator by ordinary iteration. Give the equation atom the truth condition

$$[\![s = t]\!]_\nu \quad :\Leftrightarrow \quad [\![s]\!]_\nu = [\![t]\!]_\nu \in M.$$

Interpret $D_{x,y}B$ as $\exists x\,([\![B(x,0)]\!] \to \forall y\,[\![B(x,y)]\!])$ and a clause $[A;\, y.\, B]$ as $[\![A]\!] \to \forall y\,[\![B(y)]\!]$, with two-valued truth. A sequent is valid if its formula is true under every assignment that makes its clauses true. Every rule preserves validity.

Each seed is true. For Eval let the premises $s = t$ and $b = r$ be true, so that $[\![s]\!] = [\![t]\!] \in M$ and $[\![b]\!] = [\![r]\!] \in M$. In each row $\Phi(s, t, b)$ has the value of b: in the root row $[\![t]\!] = [\![s]\!] = [\![b]\!]$; in the zero row $[\![t]\!] = [\![a]\!] = [\![I(0, a;\, u.\, h)]\!]$; in the successor row $[\![h[t/u]]\!] = [\![h[s/u]]\!] = [\![I(Sn, a;\, u.\, h)]\!]$, because $s \equiv I(n, a;\, u.\, h)$ and t have the same value. Hence the conclusion $r = \Phi(s, t, b)$ is true. Use is the elimination of an implication and of a universal quantifier. Elim is existential elimination; the fresh variable takes the value of the witness, and neither the context nor the conclusion depends on it. For Intro fix an assignment that satisfies Γ. The clause $\mathrm{ord}(F)$ is true whenever F is. In case mode, if $B(t, 0)$ is true then the premise makes $B(t, Sn)$ true for every standard n; every number is 0 or a successor, so every instance is true and the conditional closure holds at the value of t. If $B(t, 0)$ is false the closure holds trivially. In progress mode the premise states that $B(0, n)$ implies $B(0, Sn)$ for every standard n; induction on the standard numbers gives every instance from $B(0, 0)$. The argument does not use that the atoms mean identity.

So every theorem of C_3^S is true in the interpretation, and $S^k0 = S^k0$ is false for $k \notin M$. For the two special cases take $S = \{0 = 0\}$, $M = \{0\}$, and $S = \emptyset$. □

The interpretation is a counter-interpretation of an inference system. It is not a structure in which the equality symbol is identity, and it is not the arithmetic interpretation of the calculus. For contexts of native formulas the case $S = \emptyset$ is proved syntactically in Theorem 4.23; the theorem adds the closed seeds. The clause

$$[\, 0 = 0;\, y.\, y = y \,]$$

is false in it. Retained in every context, together with a derivation of $0 = 0$, it restores Refl: Use at t gives $t = t$. This relocates universal reflexivity into the initial context; the axiom and the cost of validating it remain part of the specification. The calculus that follows makes this choice and also removes the neutral antecedent.

Definition 5.26 (The calculus V)**.** A *uniform clause* is $[\Pi;\, y.\, B(y)]$ where Π is empty or one formula and y does not occur in Π; it reads $\forall y\, B(y)$, respectively $A \to \forall y\, B(y)$. The *reflexivity clause* and the clause of an assumption are

$$\mathcal{R} := [\,;\, y.\, y = y\,], \qquad \langle A\rangle := [\,;\, v.\, A\,] \quad (v \notin \mathrm{FV}(A)).$$

The rules are

$$\frac{\Gamma \vdash A_1 \;\cdots\; \Gamma \vdash A_k}{\Gamma \vdash B(t)}\ \textsc{Use} \qquad \frac{\Gamma \vdash s = t \quad \Gamma \vdash b = r}{\Gamma \vdash r = \Phi_2(s,t,b)}\ \textsc{Eval}$$

$$\frac{\Gamma \vdash D_{x,y}B \quad \Gamma, [B(b,0);\, y.\, B(b,y)] \vdash F}{\Gamma \vdash F}\ \textsc{Elim}$$

$$\frac{\Gamma, \langle B(t,\eta)\rangle \vdash B(t,Sz)}{\Gamma \vdash D_{x,y}B}\ \textsc{Intro}$$

where Use selects an occurrence $[A_1, \ldots, A_k;\, y.\, B]$ of Γ with $k \in \{0,1\}$, and the side conditions and η are those of Theorem 5.4. For a context Γ of uniform clauses, Γ° replaces each occurrence $[\,;\, y.\, B]$ by $[0 = 0;\, y.\, B]$.

There are four rule families and one fixed axiom clause. Use of $\mathcal{R}$ at t gives $t = t$, and Use of $\langle A\rangle$ gives A; both have no premise. Reflexivity is not eliminated as a principle; it is the clause $\mathcal{R}$. The calculus is `VPrf` in the Lean development.

Theorem 5.27 (Uniform clauses)**.** *For every context Γ of uniform clauses and every formula A,*

$$\mathcal{R}, \Gamma \vdash_{\mathsf{V}} A \quad\Longleftrightarrow\quad \Gamma^\circ \vdash_{\mathsf{C}} A.$$

Hence, for Q-free Δ and F, $\mathcal{R}, \langle\Delta\rangle \vdash_{\mathsf{V}} F$ if and only if $\Delta \vdash_{\mathsf{N}} F$. The translations leave terms and formulas unchanged. The translation from C to V does not increase the number of nodes; the converse adds at most one Refl node for each use of a clause without antecedent.

Proof. The occurrences of Γ and of Γ° correspond by position, and $\mathcal{R}$ is the first occurrence on the left. Both directions are inductions on derivations, over all Γ.

From right to left: Refl at t becomes Use of $\mathcal{R}$ at t. A Use of an occurrence $[0 = 0;\, y.\, B]$ that corresponds to $[\,;\, y.\, B]$ becomes the Use without premise, and the derivation of $0 = 0$ is discarded. A Use of an occurrence that corresponds to a clause with an antecedent keeps its translated premise. Eval is copied. Elim adds the same clause on both sides, and Intro adds $\mathrm{ord}(B(t,\eta))$, which is $\langle B(t,\eta)\rangle^\circ$. Witnesses, modes and eigenvariables are unchanged.

From left to right: a Use of $\mathcal{R}$ at t becomes Refl at t. A Use of $[\,;\, y.\, B]$ becomes a Use of $[0 = 0;\, y.\, B]$ with premise Refl at 0. The other rules are copied.

The second statement follows from Theorem 5.11(iii), since $\langle\Delta\rangle^\circ = \mathrm{ord}(\Delta)$. □

Remark 5.28. In the Lean development the first statement is `vprf_iff_cprf` and the second is `vprf_iff_nprf_of_qfree` (`NativeUniformClauses`). In the native reading of a context of V that is reachable from assumptions, $\mathcal{R}$ supplies reflexivity, $\langle A\rangle$ is the assumption A and an opened clause is $Q_y B(b,y)$, so Use is reflexivity, Hyp or QE.

5.3.4 Four schemes on zero-normal terms

This subsection draws on `clause_contexts`.

Definition 5.29 (The calculus V^0)**.** V^0 has zero-normal terms, normal substitution, uniform clauses and the clause $\mathcal{R}$, with the four schemes Use (as in V), Step (the rule Eval of C^0), Open (the rule Elim of C^0) and Close (the rule Intro of C^0 with the assumption stored as $\langle B\langle t, \eta\rangle\rangle$).

Theorem 5.30 (Clause factorization)**.** *For every zero-normal context Γ of uniform clauses and every zero-normal formula A, $\mathcal{R}, \Gamma \vdash_{\mathsf{V}^0} A$ if and only if $\Gamma^\circ \vdash_{\mathsf{C}^0} A$. Hence, for raw Q-free Δ and F, $\mathcal{R}, \langle \mathrm{nf}_0\, \Delta\rangle \vdash_{\mathsf{V}^0} \mathrm{nf}_0(F)$ if and only if $\Delta \vdash_{\mathsf{N}} F$.*

Proof. The proof of Theorem 5.27 applies; it does not inspect the evaluation rule or the terms. The second statement is Theorem 5.22(iii). □

In the Lean development V^0 is `ZVPrf` and the two statements are `zvprf_iff_zcprf` and `zvprf_iff_nprf`.

Remark 5.31 (The atomic case without cut). `clause_contexts` recovers the introduction by cases of a literal equation $E(x, y)$ at the witness 0 without the cut of Theorem 5.5; Section 4.7.2 gives the same derivation for `uniform_step`. Let $q \equiv I(S0, 0;\, u.\, u)$, which is not literally 0 and satisfies $q = 0$ by Theorem 5.21(ii). Given $\Gamma, E(0,0) \vdash E(0, Sz)$, apply Close to the matrix $E(0, y)$ with an unused first variable at the witness q; this is case mode, its premise is the given derivation, and it gives $D_{v,y}E(0, y)$. Open adds the clause $[E(0,0);\, y.\, E(0, y)]$. Now apply Close to $E(x, y)$ at the witness q: its assumption $E(q, 0)$ transports to $E(0, 0)$, Use gives $E(0, Sz)$, and transport gives $E(q, Sz)$.

5.4 Variants of the introduction

This section draws on `clause_guards` and `rootfree_clausal`. The introduction of C selects its mode by the root of the matrix and by the literal witness. A substitution can change the second test (Theorem 5.14). This section gives introductions whose mode is selected by a guard on the matrix alone, and one whose mode is selected by the instantiated zero base.

5.4.1 The diagonal guard

This subsection draws on `clause_guards`.

Definition 5.32 (The calculus K)**.** A matrix $B(x, y)$ is *diagonal*, written $\mathrm{diag}_y(B)$, if it has the literal shape

$$I(y, y;\, u.\, h) = I(y, y;\, v.\, k),$$

where the four displayed occurrences of y are the inner variable of the enclosing $D_{x,y}$ and h, k are arbitrary. The test is made on the matrix before the witness is substituted. K is C with the introduction

$$\frac{\Gamma, B(t, \eta_B(z)) \vdash B(t, Sz)}{\Gamma \vdash D_{x,y}B}\ \textsc{Intro}, \qquad \eta_B(z) := \begin{cases} z, & \mathrm{diag}_y(B), \\ 0, & \text{otherwise}, \end{cases}$$

with z fresh for Γ, t and B.

No rule of K inspects a witness. Theorem 5.5, Theorem 5.6 and Theorem 5.6 hold in K with the same proofs: they use Refl, Eval and Use only, and injective renaming preserves the diagonal test.

Lemma 5.33 (Derived rules of K). *(i) If* $\mathrm{diag}_y(B)$ *then* $\Gamma \vdash B(t,0)$ *for every* t *and* Γ.

(ii) The introduction by cases of Theorem 5.7 is admissible for every matrix and every witness.

(iii) The closure interfaces (i) to (iii) of Theorem 5.9, numerical cases for every formula, and transport in every formula (Theorem 5.8) hold.

Proof. (i) The instance is $I(0,0;\, u.\, h') = I(0,0;\, v.\, k')$. Both sides equal 0 by the zero computation, and symmetry and transitivity join them. Neither body is evaluated.

(ii) For a matrix that is not diagonal this is Intro. For a diagonal matrix cut the theorem $B(t,0)$ of (i) into the given branch $\Gamma, B(t,0) \vdash B(t,Sz)$. The result $\Gamma \vdash B(t,Sz)$ is weakened by the assumption $B(t,z)$, and Intro in progress mode applies.

(iii) The proofs of Theorem 5.9(i) to (iii) and of Theorem 5.8 use the introduction by cases, Use and Elim. Numerical cases follow from interfaces (i) and (ii). □

Equation induction is not immediate in K, since progress is restricted to diagonal matrices. For a literal equation $E(y) \equiv (s(y) = t(y))$ define the *diagonal lift*

$$H_E(y) := \big(L_s(y) = L_t(y)\big), \qquad L_s(y) := I(y,y;\, u.\, s(y)), \quad L_t(y) := I(y,y;\, u.\, t(y)),$$

with u not in s, t. The matrix H_E is diagonal and $H_E(0)$ is a theorem by Theorem 5.33(i). The successor computation gives $L_s(Sz) = s(Sz)$ and $L_t(Sz) = t(Sz)$, hence

$$H_E(Sz) \dashv\vdash E(Sz). \tag{5.1}$$

Numerical cases prove the two equations

$$\mathrm{sel}(y, s(0), L_s(y)) = s(y), \qquad \mathrm{sel}(y, t(0), L_t(y)) = t(y) \tag{5.2}$$

without induction: at 0 both sides of the first are $s(0)$ after the zero computation, and at Sz the selector and the iterator with constant body compute to $s(Sz)$; likewise for t. Under $E(0)$, congruence along $s(0) = t(0)$ and along $H_E(y)$ joins the left sides of (5.2), so

$$E(0),\ H_E(y) \vdash E(y). \tag{5.3}$$

Theorem 5.34 (Equation induction in K). *For every literal equation* $E(y)$*, every context* Γ *and every term* q*, derivations of* $\Gamma \vdash E(0)$ *and of* $\Gamma, E(z) \vdash E(Sz)$*,* z *fresh, yield a derivation of* $\Gamma \vdash E(q)$*. Consequently interface (iv) of Theorem 5.9 and Theorem 5.10 hold in* K.

Proof. Under the assumption $H_E(z)$, (5.3) and the given derivation of $E(0)$ give $E(z)$. Cut this into the given step derivation at its own variable z; the result is $E(Sz)$, and (5.1) gives $H_E(Sz)$. Intro on the diagonal matrix H_E with an unused first variable proves $q_y H_E$. Its zero instance is a theorem, so interface (ii) gives $H_E(q)$, and (5.3) gives $E(q)$. No term is substituted in the given step derivation, and the induction that is being derived is not used for (5.2).

For interface (iv), work in $\Gamma, E(0)$. The first part gives $E(Sw)$ for a fresh w, and interface (i) gives $\Gamma \vdash q_y E$. With a witness, Theorem 5.33(ii) gives the third statement of Theorem 5.10. □

Theorem 5.35 (The diagonal clause calculus and N). *The four statements of Theorem 5.11 hold for* K.

Proof. (i) is proved as there, except for INTRO. An introduction with a matrix that is not diagonal is the introduction by cases of N. A diagonal matrix is a literal equation $E(x, y)$; the translated premise is $\Gamma^\natural, E(t, z) \vdash E(t, Sz)$, the introduction of Q by progress gives $Q_y E(t, y)$, and the introduction by cases at t with QE gives $D_{x,y}E$. (ii) is proved as there, with Theorem 5.33(ii) for the introduction by cases and Theorem 5.34 for the introduction of Q by progress. (iii) and (iv) follow as there. □

Lemma 5.36 (The diagonal shape under substitution)**.** *Let σ substitute terms for free variables of a matrix $B(x, y)$, lifted under x and y. Then* $\mathrm{diag}_y(B\sigma)$ *if and only if* $\mathrm{diag}_y(B)$*.*

Proof. A substitution does not change an iterator constructor that is present and fixes the bound y. A term substituted for a free variable is free of the bound y, so it is neither the variable y nor an iterator whose counter and initializer are y. Hence neither of the two required iterator heads can be created or destroyed, and a matrix with root D keeps its root. □

Theorem 5.37 (Substitution preserves the derivation tree)**.** *If $\Gamma \vdash_{\mathsf{K}} A$ and σ is a substitution of terms for free variables, then $\Gamma\sigma \vdash_{\mathsf{K}} A\sigma$ by a derivation with the same tree of rules, the same selected clause occurrences and the same evaluation rows, after a consistent renaming of eigenvariables.*

Proof. Induction on the derivation. REFL is immediate. The literal matches of each row of Φ_2 are preserved and substitution commutes with the instantiation of the body. USE keeps its occurrence and its antecedent. ELIM uses a fresh variable in the substituted context. For INTRO, Theorem 5.36 fixes the mode, so the substituted branch is the branch that the same rule requires. □

Remark 5.38. In the Lean development K is `KPrf`, the equivalences are `kprf_iff_nprf` and `kprf_iff_nprf_of_qfree`, and Theorem 5.37 is `KPrf.subst` (`NativeClausalDiagonal`). `clause_guards` compares K with the six-rule native calculus that has the diagonal introduction (`SPrf`), for which the translation τ preserves and reflects the diagonal shape, so that introductions are copied. C has substitution as an admissible rule (Theorem 5.14); the statement for K is that the derivation is not repaired.

5.4.2 Guards with a derivable base

This subsection draws on `clause_guards`; the method of repairing an introduction after a one-way change of guard is that of `reflexive_base`.

Let $G_y(B)$ be a decidable predicate on matrices that is invariant under renaming of bound variables and under injective renaming of free variables. K_G is K with G in the place of diag. Consider the following conditions.

(a) Every matrix that satisfies G is a literal equation.

(b) For every matrix B that satisfies G and every t, the zero instance $B(t, 0)$ is derivable by REFL and EVAL.

(c) For every literal equation $E(y)$ there is a matrix $H_E(y)$ that satisfies G, with derivations of $H_E(Sz) \dashv\vdash E(Sz)$ and of $E(0), H_E(y) \vdash E(y)$ that use REFL, EVAL, USE, ELIM and introductions by cases only.

Theorem 5.39 (Guard criterion)**.** *Under (a), (b) and (c) the four statements of Theorem 5.11 hold for K_G. If moreover G is preserved by substitution of terms for free variables of the matrix, substitution is admissible in K_G by a direct transformation of derivations that inserts nodes only at introductions whose matrix acquires the guard. If G is also reflected by substitution, the transformation preserves the derivation tree.*

Proof. Condition (b) gives the introduction by cases for guarded matrices as in Theorem 5.33(ii), hence the closure interfaces, numerical cases and transport in formulas. Condition (c) and the proof of Theorem 5.34, with the given lift in the place of the diagonal lift, give equation induction and interface (iv). A guarded introduction is sound in N because by (a) it is an instance of progress for a literal equation; an unguarded one is an introduction by cases. The proof of Theorem 5.35 then applies.

For substitution induct on the derivation. A guarded matrix remains guarded, and an unguarded matrix that remains unguarded keeps its rule. If an unguarded matrix becomes guarded, the substituted branch assumes the substituted zero instance; (b) proves that instance, cut removes the assumption, the branch is weakened by the progress assumption, and the guarded Intro applies. If G is reflected this case does not occur. □

Proposition 5.40 (The mirror guard on the full language)**.** *Let* $\mathrm{mirror}_y(B)$ *hold if* $B \equiv (a = I(y, a;\, u.\, h))$*. The guard satisfies (a), (b) and (c) and is preserved by substitution. It is not reflected by substitution.*

Proof. (a) holds by definition. (b): the zero instance is $a' = I(0, a';\, u.\, h')$, the symmetric form of the zero computation. (c): for $E(y) \equiv (s(y) = t(y))$ let

$$M_E(y) := \big(s(y) = I(y, s(y);\, u.\, t(y))\big), \qquad u \notin \mathrm{FV}(t).$$

The successor instance computes to $E(Sz)$. Numerical cases prove

$$\mathrm{sel}(y, s(0), s(y)) = s(y), \qquad \mathrm{sel}(y, t(0), I(y, s(y);\, u.\, t(y))) = t(y),$$

since at 0 both sides are $s(0)$, respectively $t(0)$, and at Sz the selector returns its last argument, which computes to $t(Sz)$ in the second equation. Under $E(0)$ and $M_E(y)$ congruence joins the two left sides, hence $E(y)$. The shape $a = I(y, a;\, u.\, h)$ is preserved by substitution. For distinct free variables p, q the matrix $p = I(y, q;\, u.\, h)$ is not guarded, and the substitution of p for q makes it guarded. □

Every matrix that is not guarded takes the case branch, so the mirror calculus is defined on the whole Q-free language; `mirror_core` restricts the public language instead. By Theorem 5.39 the mirror calculus is equivalent to N and has admissible substitution with a local repair, and it lacks the exact statement of Theorem 5.37. A guard that tests one iterator side without its relation to the initializer does not satisfy (b).

5.4.3 The fixed zero base on zero-normal terms

This subsection draws on `rootfree_clausal`; the guard is that of `reflexive_base`.

Definition 5.41 (The calculus $\mathsf{C}^0_{\mathrm{b}}$)**.** $\mathsf{C}^0_{\mathrm{b}}$ is C^0 with the introduction

$$\frac{\Gamma, B\langle t, \eta\rangle \vdash B\langle t, Sz\rangle}{\Gamma \vdash D_{x,y}B}\ \textsc{Intro}$$

in which

$$\eta := \begin{cases} z, & B\langle t, 0\rangle \equiv (0 = 0), \\ 0, & \text{otherwise.} \end{cases}$$

The test is one comparison of the computed base with a fixed formula. Substitution and normalization preserve the root of a formula, so the test excludes matrices with root D; there is no separate test of the root or of the witness.

Theorem 5.42 (The fixed zero base)**.** *For every zero-normal clause context* Γ *and formula* F*,* $\Gamma \vdash_{\mathsf{C}^0_\mathrm{b}} F$ *if and only if* $\Gamma \vdash_{\mathsf{C}^0} F$*. Hence Theorem 5.22(iii) and (iv) hold for* C^0_b*.*

Proof. The two calculi differ in the introduction only. Theorem 5.21(i) to (v) hold in C^0_b with the same proofs.

Cases in C^0_b*.* Let $\Gamma, B\langle t, 0\rangle \vdash B\langle t, Sz\rangle$. If the base is not $0 = 0$ the rule applies. If it is $0 = 0$, cut Refl into the branch, weaken by the assumption $B\langle t, z\rangle$ and apply the rule in progress mode. No auxiliary witness is needed. The closure interfaces (i) to (iii) and transport in formulas follow.

Equation progress in C^0_b*.* Let $E(y) \equiv (s(y) = t(y))$ and let $\Gamma, E(z) \vdash E(Sz)$ be given. Write $s_0 \equiv s\langle 0\rangle$, and let the selector on zero-normal terms be $\mathrm{sel}(v, a, b) \equiv \overline{I}(v, a;\, u.\, b)$. Put

$$\begin{aligned} H(y) &:= \big(\mathrm{sel}(y, 0, s(y)) = \mathrm{sel}(y, 0, t(y))\big),\\ F_s(y) &:= \big(\mathrm{sel}(y, s_0, \mathrm{sel}(y, 0, s(y))) = s(y)\big),\\ F_t(y) &:= \big(\mathrm{sel}(y, s_0, \mathrm{sel}(y, 0, t(y))) = t(y)\big). \end{aligned}$$

In zero-normal syntax $H\langle 0\rangle$ is literally $0 = 0$, $F_s\langle 0\rangle$ is $s_0 = s_0$ and $F_t\langle 0\rangle$ is $E\langle 0\rangle$. Each successor instance of F_s and of F_t is derivable by two selector computations. Work in $\Gamma, E\langle 0\rangle$. The introduction by cases gives $q_y F_s$ and $q_y F_t$, and interface (ii) with the bases Refl and $E\langle 0\rangle$ gives $F_s\langle z\rangle$ and $F_t\langle z\rangle$. Under the further assumption $H\langle z\rangle$, congruence for the template $v \mapsto \mathrm{sel}(z, s_0, v)$ and the two decoder equations give $E\langle z\rangle$. Cut this into the given step derivation; the result is $E\langle Sz\rangle$, and one selector computation on each side gives $H\langle Sz\rangle$. The introduction of C^0_b on the matrix H with an unused first variable is in progress mode, because its base is $0 = 0$, and gives $q_y H$. Refl and interface (ii) give $H\langle Sw\rangle$, hence $E\langle Sw\rangle$, for a fresh w. The introduction by cases concludes $\Gamma \vdash q_y E$. No equation induction is used before the guarded rule is applied.

From C^0 *to* C^0_b*.* An Intro of C^0 in case mode is an introduction by cases. In progress mode the matrix is a literal equation and the witness is 0; equation progress gives $q_y B\langle 0, y\rangle$, and the introduction by cases at the witness 0 with interface (ii) gives $D_{x,y}B$.

From C^0_b *to* C^0*.* If the guard is false the rule is an introduction by cases (Theorem 5.21(vi)). If it is true the matrix is a literal equation, interface (iv) of C^0 applied to the matrix $B\langle t, y\rangle$ gives $q_y B\langle t, y\rangle$, its base is $0 = 0$, and interface (ii) gives $B\langle t, Sz\rangle$; the introduction by cases at the witness t concludes $D_{x,y}B$. □

If the base is literally $0 = 0$ it remains so under substitution, since $\mathrm{nf}_0(\mathrm{nf}_0(G)\sigma) \equiv \mathrm{nf}_0(G\sigma)$. A base that is not $0 = 0$ can become $0 = 0$, which changes an introduction by cases into a progressive one; the derived introduction by cases repairs that instance. So the guard is preserved and not reflected, as for the mirror guard.

5.5 Variants of the context

This section draws on `prefix_clauses`, `clause_contexts` and `one_body`. A context entry of C has an antecedent and a body. In V the antecedent is optional (Section 5.3.3). This section gives the representation of a formula by its prefix, with clauses that keep one antecedent, and two calculi whose context entries have no antecedent field at all.

5.5.1 The prefix representation

This subsection draws on `prefix_clauses` and on the ranked representation of `clause_contexts`.

Lemma 5.43 (Unique prefix)**.** *Every Q-free formula is, in exactly one way up to the names of bound variables,*

$$D_{x_1,y_1} \cdots D_{x_k,y_k}(s = t), \qquad k \geq 0.$$

The number k is unchanged by substitution of terms. A native formula has this form if and only if it is Q-free.

Proof. An equation has $k = 0$. A formula $D_{x,y}B$ has exactly one formula child; the grammar has no branching constructor, so removing binders reaches exactly one equation, and restoring them in order reconstructs the formula. Substitution acts on the two terms of the leaf. □

The formula is therefore the record $[k, s, t]$, written $D^k(s = t)$: a number and two terms, with $2k$ bound variables in scope at the leaf. Only term syntax is recursive. The record is a bijective encoding of the Q-free formulas; it is not a prenex normal form theorem for ordinary formulas, and each D keeps its conditional reading. With de Bruijn indices starting at 1, the two outer variables of the matrix of $D^{k+1}(s = t)$ are the indices $2k + 2$ (the witness) and $2k + 1$ at the leaf, and an iterator body adds one to the index sought.

Lemma 5.44 (Opening commutes with packing)**.** *The record of an instance $B(t, r)$ of the matrix of $D^{k+1}(s = t)$ is obtained from $[k, s, t]$ by substituting at the index $2k + 2$ and then at the index $2k + 1$, with larger indices decreased when a variable is removed and the index sought increased by one under each iterator body.*

Proof. Each removed pair of binders adds two to the depth at which the substitution acts. At the leaf induct on the two terms: a variable is replaced at the indicated index, S adds no binder and the body of an iterator adds one. These are the clauses of capture-avoiding substitution on the nested representation. □

In the Lean development the predicate `NFm.Pre k A` states that A is k binders D followed by one equation, with `NFm.qfree_iff_pre`, `NFm.Pre.unique` and `NFm.Pre.subst`; `VPrf.pre` states that V derives only such formulas from Q-free assumptions (`NativePrefix`).

Definition 5.45 (The calculus PF)**.** Formulas are records $D^k(s = t)$ on raw terms. A formula is *reflexive at its leaf* if it is $D^k(r = r)$ with literally identical terms. Clauses are $[A;\ y.\, B(y)]$ with one antecedent, an assumption A is $[0 = 0;\ v.\, A]$, and every context contains the clause

$$\rho := [\, 0 = 0;\ y.\, y = y\,].$$

The rules are Use, Eval with Φ_2, Elim and Intro of C, with one change: in Use of $[A;\ y.\, B]$ the premise is omitted when A is reflexive at its leaf. In Intro the test that the matrix is a literal equation is the test that the conclusion has $k = 1$.

There are four rule families and the fixed clause ρ. The test of reflexivity at the leaf compares two term trees and evaluates nothing.

Lemma 5.46 (Reflexive prefixes)**.** *Every formula $D^k(r = r)$ is derivable in every context in C, in V with $\mathcal{R}$, and in PF.*

Proof. Induction on k. For $k = 0$ use Refl in C, Use of $\mathcal{R}$ in V, and Use of ρ in PF, where the antecedent $0 = 0$ is reflexive and the premise is omitted. For $k > 0$ apply Intro at the witness $S0$, which is case mode whatever the matrix. The target at the fresh successor has $k - 1$ binders and identical terms at its leaf, since both receive the same substitutions; the induction hypothesis proves it, and the added clause is not used. □

Theorem 5.47 (The prefix calculus)**.** *For every clause context* Γ *and formula* A*, with formulas packed on the left,*

$$\rho, \Gamma \vdash_{\mathsf{PF}} A \quad \Longleftrightarrow \quad \Gamma \vdash_{\mathsf{C}} A.$$

Hence $\rho, \mathrm{ord}(\Delta) \vdash_{\mathsf{PF}} F$ *if and only if* $\Delta \vdash_{\mathsf{N}} F$*, for Q-free* Δ *and* F*. Both formula round trips between records and nested formulas are identities.*

Proof. From right to left: pack all formulas and add ρ. Refl at t becomes Use of ρ at t without premise. The other rules translate directly; by Theorem 5.44 substitutions agree, and the free variables are unchanged, so the eigenvariable conditions agree. From left to right: a Use of the distinguished ρ becomes Refl. Any other Use keeps its translated premise; if the premise was omitted, Theorem 5.46 supplies a derivation of the antecedent in C. Eval, Elim and Intro translate directly. No cut is needed. The second statement is Theorem 5.11(iii). □

By Theorem 5.27 the same holds against V, with the empty antecedent represented by $0 = 0$. `prefix_clauses` states the theorem against `CPrf` and `VPrf`.

5.5.2 Rank from variable support

This subsection draws on the optional notes of `clause_contexts`. It shows that the number k of a public record can be recovered from the variables of its leaf, at the price of identifying some theorems.

Work in V^0 with records and de Bruijn indices. For a term t let $\mu(t)$ be the greatest positive index that is free relative to the iterator binders of t, and 0 if there is none: $\mu(j) = \max(j, 0)$, $\mu(St) = \mu(t)$ and $\mu(I(n, a;\, u.\, h)) = \max(\mu(n), \mu(a), \max(\mu(h) - 1, 0))$. For a record $F = [k, s, t]$ let $r(F) := \lceil \max(\mu(s), \mu(t))/2 \rceil$, so that $r(F) \leq k$. Define

$$P[k, s, t] := \begin{cases} (s, t), & r(F) = k, \\ (0, 0), & r(F) < k, \end{cases} \qquad U(s, t) := \big[\lceil \max(\mu(s), \mu(t))/2 \rceil,\ s,\ t\big].$$

The pair $(0, 0)$ decodes to $0 = 0$. The quotient calculus has pairs as the formulas of its judgments and records in its clauses, which keep their rank: Use asks for derivations of $P(A_i)$ and returns $P(B\langle t/y \rangle)$; Step acts on pairs of rank zero; Open decodes a major premise of nonzero rank, adds its clause as a record and derives the same pair in the branch; Close decodes a goal of nonzero rank, adds its assumption as a record and asks for the packed successor instance. P is applied only to formulas all of whose clause variables are instantiated, and it is recomputed after every normal substitution.

Proposition 5.48 (Support quotient)**.** *For every record* F*,* F *and* $U(P(F))$ *are interderivable in* V^0 *in every context that contains* $\mathcal{R}$*, and* $P(U(a)) = a$ *for every pair* a*. The quotient calculus derives* $P(F)$ *from* $\mathcal{R}$ *if and only if* V^0 *derives* F *from* $\mathcal{R}$*.*

Proof. If $r(F) = k$ then $U(P(F))$ is F. If $r(F) < k$, the two variables of the outermost binder, with the indices $2k$ and $2k - 1$, do not occur in the leaf, so $F \equiv D_{x,y}B$ with x, y not in B. Close at the witness 0 has B both as its assumption and as its target in either mode, so F is a theorem; $0 = 0$ is a theorem; and each follows from the other by weakening. $P(U(a)) = a$ because U uses the rank that P tests.

Soundness of the quotient rules: by the first part a derivation of $U(P(A))$ converts to one of A and conversely, by cut and weakening. Use converts its packed premises, applies Use of V^0 and converts the result. Step involves rank zero only. Close converts the local base and the target and applies Close; the decoded goal has the rank derived from its support and packs to the same pair. Open decodes a record with root D and adds the clause of Open.

Completeness: transform a derivation of V^0, packing every conclusion. Use and Step transform directly. A Close whose goal keeps its rank is the same Close; if the goal collapses, Use of $\mathcal{R}$ at 0 proves $0 = 0$ and the premise is discarded. Open transforms directly when its major premise keeps its rank. If the major premise $D_{x,y}B$ collapses, the added clause is $[B;\, y.\, B]$, and each use of it returns the formula of its own premise; delete those uses and keep the premise derivations, removing the clause from the branch. The branch may still contain the eigenvariable b in intermediate terms. To discharge it, use the theorem $T \equiv D_{x,y}(x = x)$, which has rank 1 and support index 2 and does not collapse. Close at the witness 0 proves T, both instances being $0 = 0$. Open of T at the fresh b adds the clause $[b = b;\, y.\, b = b]$, the transformed branch is weakened by it, and the rule concludes the goal in the outer scope. □

The quotient contracts a binder only when both its variables are absent; theorems such as $D_{x,y}(x = x)$ are kept, and no completeness of the test for theorems is claimed. A clause body has one more bound variable, the clause variable, so the same support can belong to a body of rank zero with a clause variable or to a formula of rank one; clause bodies therefore keep an explicit rank.

5.5.3 One body per context entry

This subsection draws on `one_body`. An opened occurrence $[B(b, 0);\, y.\, B(b, y)]$ is determined by its body. An assumption can be stored in the same form.

Definition 5.49 (Gate and the calculus OB)**.** For $A \equiv D^d(s = t)$ and a variable v outside the $2d$ bound variables the *gate* is

$$\mathrm{gate}_v\, A := D^d\big(\mathrm{sel}(v, 0, s) = \mathrm{sel}(v, 0, t)\big).$$

A context of OB is a finite list of *bodies* $B(v)$, each a formula with one distinguished variable; a body reads $B(0) \to \forall v\, B(v)$, which is the reading of $Q_v B$. The rules are Refl for equations, $\textsc{Eval}_0$ and $\textsc{Eval}_S$, and

$$\frac{\Gamma \vdash B(0)}{\Gamma \vdash B(t)}\ \textsc{Use} \qquad \frac{\Gamma \vdash D_{x,y}B(x, y) \quad \Gamma, B(b, y) \vdash A}{\Gamma \vdash A}\ \textsc{Open}$$

$$\frac{\Gamma, \mathrm{gate}_v(B(w, \eta)) \vdash B(w, Sz)}{\Gamma \vdash D_{x,y}B(x, y)}\ \textsc{Intro}$$

where Use selects a body $B(v)$ of Γ, Open adds the body $y \mapsto B(b, y)$ with b fresh, and in Intro the variables z and v are fresh, w is the witness, and η is z if B is a literal equation and $w \equiv 0$, and 0 otherwise.

A context entry has no antecedent field, and there is no entry for an ordinary assumption. The gate is a construction on the existing grammar. Its variable v lies outside the binders of A, and the terms of the leaf are placed under the new iterator binder of the selector, which in a de Bruijn representation requires one lifting.

The subsystem with Refl and the two evaluation rows gives the equality reasoning of Theorem 5.6 and Theorem 5.6 in every context. Write $s \simeq t$ if this subsystem proves $s = t$. Its derivations have no mode test and are stable under substitution. In particular

$$\mathrm{sel}(0, 0, s) \simeq 0, \qquad \mathrm{sel}(Sz, 0, s) \simeq s. \tag{5.4}$$

For formulas A, A' with the same prefix, aligned, write $A \approx A'$ if their left terms are related by $\simeq$ and their right terms are related by $\simeq$, the bound variables of the prefix being treated as free. The definition does not presuppose transport in formulas.

Lemma 5.50 (Deep reflexivity)**.** *Every $D^d(t = t)$ is derivable in* OB *in every context. Weakening and injective renaming of free variables are admissible.*

Proof. Induction on d as in Theorem 5.46: at a binder apply Intro at the witness $S0$ and ignore the new gate body. Weakening and renaming are inductions on derivations. □

Lemma 5.51 (Conversion and access)**.** *For every d the following hold in* OB*.*

L_d *If $A \approx A'$ and both have d binders, a derivation of either in a context converts to a derivation of the other in that context.*

H_d *If A has d binders and $\mathrm{gate}_v A$ is a body of Γ, then $\Gamma \vdash A$.*

Proof. The statements are proved in the order $L_0, H_0, L_1, H_1, \ldots$.

L_0 is symmetry and transitivity. For L_d with $d > 0$ assume L_{d-1} and H_{d-1}, and let $A \equiv D_{x,y}B$, $A' \equiv D_{x,y}B'$. Apply Open to the given derivation of A at a fresh b, which adds the body $B(b, y)$. Apply Intro for A' at the witness b. A variable is not the literal 0, so this is case mode, and the branch has the body $\mathrm{gate}_v B'(b, 0)$. H_{d-1} gives $B'(b, 0)$. L_{d-1}, with the endpoint derivations instantiated at $x := b$, $y := 0$, gives $B(b, 0)$. Use of the body $B(b, y)$ at Sz gives $B(b, Sz)$, and L_{d-1} gives $B'(b, Sz)$. Intro concludes A' and Open discharges b. The other direction uses the symmetric endpoint equations.

For H_d, given L_d: let $T_d \equiv D^d(0 = 0)$ with the prefix of A. Theorem 5.50 proves T_d, and (5.4) gives $\mathrm{gate}_0 A \approx T_d$ and $\mathrm{gate}_1 A \approx A$. L_d gives $\mathrm{gate}_0 A$. Use of the body $\mathrm{gate}_v A$ at 1 gives $\mathrm{gate}_1 A$, and L_d gives A. □

Conversion at d binders uses access at $d-1$ binders only, so the induction is well founded. It is an induction on formula syntax; it adds no induction to the arithmetic and does not presuppose the access that it proves. The terms of the leaf may contain the bound variables of the prefix, so the endpoint equations live under the binders and cannot be replaced by one global substitution of equals.

Theorem 5.52 (From clauses to bodies)**.** *For Q-free $A_1, \ldots, A_n, F$: if $\mathrm{ord}(A_1), \ldots, \mathrm{ord}(A_n) \vdash_{\mathsf{C}} F$ then $\mathrm{gate}\, A_1, \ldots, \mathrm{gate}\, A_n \vdash_{\mathsf{OB}} F$.*

Proof. Induct on the derivation with its canonical annotation, translating an assumption occurrence $\mathrm{ord}(A)$ to the body $\mathrm{gate}_v A$ and an opened occurrence $[B(b, 0); y.\, B(b, y)]$ to the body $B(b, y)$. Refl and Eval are unchanged. A Use of an assumption occurrence is replaced by H_d of Theorem 5.51; its premise $0 = 0$ is discarded. A Use of an opened occurrence has the premise $B(b, 0)$ and is Use of OB. Elim adds an opened occurrence, as Open adds its body. Intro adds the assumption $B(w, \eta)$, whose translation is the gate body that Intro of OB adds; witness, mode and eigenvariables are unchanged. □

The translation uses no cut and no weakening of OB beyond Theorem 5.50, and no substitution in a derivation with Intro.

Lemma 5.53 (The gate in N)**.** *For every Q-free A and fresh v, in every context, $A \vdash_{\mathsf{N}} Q_v\, \mathrm{gate}_v A$ and $Q_v\, \mathrm{gate}_v A \vdash_{\mathsf{N}} A$.*

Proof. The equations (5.4) hold in N, so Theorem 5.2(ii), applied along the prefix, gives $\mathrm{gate}_{Sz} A \dashv\vdash A$ and $\mathrm{gate}_0 A \dashv\vdash D^d(0 = 0)$; the latter formula is derivable by d introductions by cases. From A one obtains $\mathrm{gate}_{Sz} A$, the introduction by cases with an unused first variable gives $D_{w,v}\, \mathrm{gate}_v A$, and Theorem 5.2(i) gives $Q_v\, \mathrm{gate}_v A$. Conversely QE with the derivable base $\mathrm{gate}_0 A$ gives $\mathrm{gate}_1 A$, hence A. □

Theorem 5.54 (One-body contexts and N)**.** *If Γ is a context of bodies and $\Gamma \vdash_{\mathsf{OB}} A$, then $\Gamma^Q \vdash_{\mathsf{N}} A$, where Γ^Q replaces each body $B(v)$ by $Q_v B$. For Q-free $A_1, \dots, A_n, F$ the following three statements are equivalent:*

$$\begin{aligned}&\operatorname{gate} A_1, \dots, \operatorname{gate} A_n \vdash_{\mathsf{OB}} F,\\ &A_1, \dots, A_n \vdash_{\mathsf{N}} F,\\ &\operatorname{ord}(A_1), \dots, \operatorname{ord}(A_n) \vdash_{\mathsf{C}} F.\end{aligned}$$

Proof. Soundness is by induction on the derivation. Refl and Eval are as in Theorem 5.11(i). Use is QE on the assumption $Q_v B$. Open is DE, which adds $Q_y B(b, y)$. For Intro the translated premise is $\Gamma^Q, Q_v \operatorname{gate}_v(B(w, \eta)) \vdash B(w, Sz)$; Theorem 5.53 and cut give $\Gamma^Q, B(w, \eta) \vdash B(w, Sz)$, and the two modes are treated as in Theorem 5.11(i). For the equivalences: soundness and Theorem 5.53 with cut lead from the first statement to the second; the second and the third are equivalent by Theorem 5.11(iii); and Theorem 5.52 leads from the third to the first. □

Remark 5.55. In the Lean development OB is `OBPrf`, the first equivalence is `obprf_iff_nprf`, its form for arbitrary native sequents through τ is `obprf_unQ_iff_nprf`, and the equivalence with C is `obprf_iff_cprf` (`NativeOneBody`); Theorem 5.51 is `OBPrf.conv` and `OBPrf.acc`, and Theorem 5.52 is `CPrf.toOB`. The Lean proof derives the gate at 0 directly (`OBPrf.gtZero`) and does not convert it from deep reflexivity. `one_body` proves soundness through T_{eq} and the bridge (Theorem 1.78); the proof above reads a body as a formula $Q_v B$ of N, as the Lean proof does.

Remark 5.56 (A padding trap). To align the numbers of binders of two formulas, a formula A is padded as $D_{x,y} \operatorname{gate}_y A$ with x unused, which is equivalent to A by Theorem 5.53. Padding by a binder both of whose variables are unused is incorrect: $D_{x,y} A$ then reads $\exists x\, (A \to A)$ and is a theorem (compare Theorem 5.48).

5.5.4 Homogeneous contexts with an asymmetric gate

This subsection draws on the appendix of `prefix_clauses`. It removes the antecedent field as OB does, keeps reflexivity as a fixed entry, and uses a gate whose zero instance is literally reflexive.

Definition 5.57 (The calculus HB)**.** For a fresh variable v let

$$\kappa_v(s = t) := \big(I(0, s;\, u.\, t) = I(v, s;\, u.\, t)\big), \qquad \kappa_v(D_{x,y} B) := D_{x,y} \kappa_v B,$$

with u not in t. A context is a list of bodies and always contains the body $\rho(y) \equiv (y = y)$. The rules are Eval with Φ_2, and

$$\frac{\Gamma \vdash B(0)}{\Gamma \vdash B(t)} \qquad \frac{B(0) \equiv D^j(r = r)}{\Gamma \vdash B(t)} \qquad \textsc{Use}$$

for a body $B(y)$ of Γ, the rule Open of OB, and Intro of OB with $\kappa_v\big(B(t, \eta)\big)$ as the added body.

There are four rule families, no rule Refl and no hypothesis rule. The second form of Use omits the premise exactly when the raw base $B(0)$ is reflexive at its leaf, for some $j \geq 0$; a theorem with root D of another shape does not qualify. The gate satisfies

$$\kappa_0(s = t) \text{ is literally reflexive}, \qquad \kappa_{Sz}(s = t) \dashv\vdash (s = t),$$

the second by the two computation equations, since u is not in t. The gate changes neither the number of binders nor the root of a formula.

Lemma 5.58 (Gate retrieval)**.** *In* HB*, in every context:*

(i) $t = t$ *is derivable, every* $D^j(r = r)$ *is derivable, and the equality rules of Theorem 5.6 and Theorem 5.6 hold;*

(ii) $\kappa_0 A$ *is derivable for every* A*;*

(iii) a derivation of $\kappa_1 A$ *converts to a derivation of* A*, and conversely;*

(iv) if $\kappa_v A$ *is a body of* Γ *then* $\Gamma \vdash A$*.*

Proof. (i) Use of ρ at t has the base $0 = 0$, so its premise is omitted and it concludes $t = t$. Reflexive prefixes are proved as in Theorem 5.46, with Intro at the witness $S0$ and the new gate body ignored. The equality rules use reflexivity and Eval only. (ii) $\kappa_0 A$ has the prefix of A and identical terms at its leaf, so it is a reflexive prefix.

(iii) and (iv) are proved together by induction on the number of binders of A. For an equation, (iii) is the two computations and transport. For (iv), Use of the body $\kappa_v A$ at 1 has a base that is reflexive at its leaf, so its premise is omitted and it concludes $\kappa_1 A$; (iii) at the same number of binders gives A. For (iii) at $A \equiv D_{x,y}B$, assume (iii) and (iv) for formulas with fewer binders. Given $\kappa_1 A \equiv D_{x,y}\kappa_1 B$, apply Open at a fresh b, which adds the body $\kappa_1 B(b, y)$, and Intro for A at the witness b, which is case mode and adds the body $\kappa_v(B(b, 0))$. (iv) gives $B(b, 0)$, (iii) gives $\kappa_1 B(b, 0)$, Use of the body $\kappa_1 B(b, y)$ at Sz gives $\kappa_1 B(b, Sz)$, and (iii) gives $B(b, Sz)$. The converse opens A, introduces $\kappa_1 A$ at the witness b, retrieves the local zero assumption by (iv), converts it by (iii), uses the body $B(b, y)$ at Sz and converts the result by (iii). □

Theorem 5.59 (Homogeneous contexts and N)**.** *For Q-free* $A_1, \ldots, A_n, F$*,*

$$\rho, \kappa A_1, \ldots, \kappa A_n \vdash_{\mathsf{HB}} F \quad \Longleftrightarrow \quad A_1, \ldots, A_n \vdash_{\mathsf{N}} F.$$

Proof. From right to left translate a derivation of C (Theorem 5.11(iii)) by the origins of its occurrences, as in Theorem 5.52: Refl becomes Use of ρ, a Use of an assumption occurrence becomes Theorem 5.58(iv), a Use of an opened occurrence becomes Use with its premise, and Elim and Intro become Open and Intro.

From left to right read a body $B(y)$ as $Q_y B$ and induct as in Theorem 5.54. The reading $Q_y(y = y)$ of ρ is derivable in N by progress. A Use with omitted premise is QE whose base $D^j(r = r)$ is derivable in N by j introductions by cases. For Intro use $A \vdash_{\mathsf{N}} Q_v \kappa_v A$, which is proved as Theorem 5.53 from $\kappa_{Sz} A \dashv\vdash A$. Finally $Q_v \kappa_v A_i$ follows from A_i, and cut gives the sequent on the right. □

Remark 5.60. `prefix_clauses` proves completeness from the seven-rule witness calculus of `witness_rule` by the same strengthened induction over contexts, and soundness through T_{eq}. The package states HB on records; the gate leaves k unchanged and acts on the leaf. This second theorem of the package is not formalized in Lean. OB and HB differ in three points: the gate (symmetric selector or asymmetric iterator), reflexivity (a rule for equations or the entry ρ with omitted premises), and the proof of access (conversion along $\approx$ from deep reflexivity or a literally reflexive zero gate).

5.6 Fusion: evaluation at the discharge

This section draws on `evaluation_discharge`. The two rows of Eval and the elimination of D become one rule family, which leaves three: Use, Intro and Discharge.

5.6.1 The raw calculus

Definition 5.61 (One evaluation step)**.** For formulas E and F the formula $\varrho_E(F)$ is defined by three disjoint cases, every match being literal:

$$\begin{aligned} \varrho_{\,s=t}\big(I(0,s;\,u.\,h)=r\big) &:= (r=t),\\ \varrho_{\,I(c,a;\,u.\,h)=t}\big(I(Sc,a;\,u.\,h)=r\big) &:= \big(r=h[t/u]\big),\\ \varrho_E(F) &:= F \qquad \text{in every other case.} \end{aligned}$$

The first case has a literal zero counter and the second a successor, so they do not overlap, and ϱ is a total function of (E,F). The replacement term, the body and the counter are read from the two formulas. An E with root D always selects the third case. In the Lean development the graph of ϱ is the relation `Rho`, with `Rho.functional` and `Rho.total`.

Definition 5.62 (The calculus ED)**.** Terms are raw, contexts are lists of uniform clauses, and the rules are Use and Intro of V (Theorem 5.26) and

$$\frac{\Gamma \vdash D_{x,y}B \quad \Gamma, \mathcal{B}_b \vdash E \quad \Gamma, \mathcal{B}_b \vdash F}{\Gamma \vdash \varrho_E(F)}\ \textsc{Discharge}$$

where $\mathcal{B}_b := [B(b,0);\, y.\,B(b,y)]$ and b is fresh for Γ and B and does not occur free in $\varrho_E(F)$. The formulas E and F may contain b.

The rule opens $D_{x,y}B$ at b, proves E and F in that scope, performs one evaluation step, and exports the result if the step has removed b or b was absent. The third case does not derive F without a premise: the rule requires a derivation of F. The two patterns of ϱ and its identity case are part of the specification.

Lemma 5.63. *Weakening and injective renaming of free variables are admissible in* ED *and in* V*. The formula* $\Theta \equiv D_{x,y}(0=0)$ *is derivable in both, in every context, also in one without* $\mathcal{R}$*.*

Proof. Weakening and renaming are inductions on derivations with eigenvariables made fresh; an injective renaming preserves and reflects the literal matches of ϱ, the constructors 0 and S, and the mode of Intro. For Θ apply Intro at the witness 0. The matrix is a literal equation, the assumption and the target are both $0=0$, and Use of the added clause $\langle 0=0\rangle$ closes the branch. □

Theorem 5.64 (Evaluation at the discharge)**.** *For every context* Γ *of uniform clauses and every formula* A*,*

$$\Gamma \vdash_{\mathsf{ED}} A \quad\Longleftrightarrow\quad \Gamma \vdash_{\mathsf{V}} A.$$

The translations leave the end sequent unchanged, and $\mathcal{R}$ *need not be in* Γ*. Hence, for* Q*-free* Δ *and* F*,* $\mathcal{R}, \langle\Delta\rangle \vdash_{\mathsf{ED}} F$ *if and only if* $\Delta \vdash_{\mathsf{N}} F$*.*

Proof. From left to right translate the three premises of a Discharge. If $\varrho_E(F)$ is given by the first or the second case, apply the corresponding row of Eval to the translated derivations of E and F in the context $\Gamma, \mathcal{B}_b$. If it is given by the third case, the translated derivation of F has the required conclusion. Then apply Elim; its eigenvariable condition is the condition that b is not free in $\varrho_E(F)$. Use and Intro are unchanged.

From right to left translate Elim with major premise $D_{x,y}B$ and branch conclusion A by Discharge with the same major premise, with $E := D_{x,y}B$, derived in the branch by weakening the translated major premise, and with $F := A$. Since E has root D, $\varrho_E(A) \equiv A$. Translate an Eval with premises E and F by Discharge with the major premise Θ of Theorem 5.63 and with the translated derivations of E and F weakened to the context $\Gamma, [0=0;\, y.\,0=0]$. The corresponding case of ϱ matches and gives the conclusion of the Eval, which contains no b. The second statement is Theorem 5.27. □

Under the ordinary reading DISCHARGE is sound directly: eliminate the existential at b; in the first case the zero computation and $s = t$ turn the second equation into $r = t$, in the second case the successor computation and the equation for the preceding iterate give $r = h[t/u]$, and the third case keeps the formula; the conclusion leaves the scope because b does not occur in it.

Remark 5.65. In the Lean development ED is `E3Prf`, with `e3prf_iff_vprf` on every clause context and `e3prf_iff_nprf_of_qfree` (`NativeEvalDischarge`). The Lean translation from V carries a renaming and an extension of the context through its induction, where the proof above uses Theorem 5.63. The translation of ELIM copies the derivation of its major premise into the scope of the witness.

Remark 5.66 (Checking without fuel). `evaluation_discharge` observes that the certificate predicate of ED is primitive recursive with no fuel field. On tree sizes raw substitution satisfies $|a[t/u]| \leq |a| \cdot \max(1, |t|)$, a step of ϱ performs at most one substitution in a body, each recursive call consumes a proper subcertificate, and the work at a node is bounded by a fixed polynomial in the sizes of its data, its context and the conclusions of its subderivations. The zero-normal companion adds structural deletions and at most one constructed iterator.

5.6.2 The zero-normal companion

Definition 5.67 (Seed evaluation and the calculus ED^0)**.** On zero-normal terms let

$$\sigma_E(F) := \begin{cases} r = h\langle \overline{I}(c, t;\, u.\, h)/u \rangle, & E \equiv (s = t),\ F \equiv (I(Sc, s;\, u.\, h) = r), \\ F, & \text{otherwise.} \end{cases}$$

ED^0 has zero-normal terms, normal substitution, uniform clauses, USE and CLOSE of V^0, and DISCHARGE with $\sigma_E(F)$ in the place of $\varrho_E(F)$. The calculus $\mathsf{V}^0_{\mathrm{s}}$ is V^0 with STEP replaced by

$$\frac{\Gamma \vdash s = t \quad \Gamma \vdash I(Sc, s;\, u.\, h) = r}{\Gamma \vdash r = h\langle \overline{I}(c, t;\, u.\, h)/u \rangle}\ \textsc{SeedEval}.$$

The first premise of SEEDEVAL equates the initializer, not the preceding iterate; the preceding iterate at the new initializer is constructed and is not a premise. The rule is that of `seed_evaluation`.

Lemma 5.68 (SEEDEVAL and STEP)**.** *On zero-normal terms, in every context that contains $\mathcal{R}$, SEEDEVAL is derivable from USE of $\mathcal{R}$ and STEP, and STEP is derivable from USE of $\mathcal{R}$ and SEEDEVAL. No logical rule is used.*

Proof. From STEP: let $s = t$ and $I(Sc, s;\, u.\, h) = r$. Theorem 5.21(iv) for the template $v \mapsto \overline{I}(c, v;\, u.\, h)$ gives $\overline{I}(c, s;\, u.\, h) = \overline{I}(c, t;\, u.\, h)$, and STEP on this equation and the second premise gives $r = h\langle \overline{I}(c, t;\, u.\, h)/u \rangle$.

From SEEDEVAL: at the counter 0 the rule reads $a = b,\ I(S0, a;\, u.\, h) = r \vdash r = h\langle b/u \rangle$. With $K \equiv I(S0, s;\, u.\, u)$, the rule on $s = r$ and $K = K$ gives $K = r$, and on $s = t$ and $K = r$ gives $r = t$; hence symmetry and transitivity. The rule on $a = a$ and the reflexivity of $I(Sc, a;\, u.\, h)$ gives the successor computation. With $J \equiv I(S0, s;\, u.\, h)$, the rule on $s = s$ and $J = J$ gives $J = h\langle s/u \rangle$, and on $s = t$ and this equation gives $h\langle s/u \rangle = h\langle t/u \rangle$. For STEP with premises $\overline{I}(c, a;\, u.\, h) = t$ and $I(Sc, a;\, u.\, h) = r$, the successor computation and congruence along the first premise give $I(Sc, a;\, u.\, h) = h\langle t/u \rangle$, and the second premise gives $r = h\langle t/u \rangle$. □

Theorem 5.69 (The zero-normal fusion)**.** *For every zero-normal context Γ of uniform clauses and every zero-normal formula A, $\Gamma \vdash_{\mathsf{ED}^0} A$ if and only if $\Gamma \vdash_{\mathsf{V}^0_{\mathrm{s}}} A$. If Γ contains $\mathcal{R}$, both are equivalent to $\Gamma \vdash_{\mathsf{V}^0} A$.*

Proof. The proof of Theorem 5.64 applies with SeedEval in the place of the two rows of Eval: a major premise with root D recovers Open, and the major premise Θ recovers SeedEval. No substitution in a derivation is used. The second statement is Theorem 5.68. □

The companion and its equivalence are not formalized in Lean. The arithmetic interface with SeedEval, with the introductions of the kernel and Q in the place of clause contexts, is `NVPrf` (`NativeZeroSeed`).

5.7 The ordinary realization

This section draws on all eleven packages of the chapter: `raw_clausal`, `rootfree_clausal`, `reduced_clause_calculus`, `clausal_two_case`, `clausal_single_evaluation`, `uniform_clauses`, `clause_guards`, `clause_contexts`, `prefix_clauses`, `one_body` and `evaluation_discharge`. Each restates the ordinary realization, the logical macros of the native interface (gate, selector and parity term, with q_y in the place of Q_y) and the bridge. These belong to the common core, and the statements reduce to it as follows.

Corollary 5.70 (Bi-interpretability)**.** *Let* X *be one of* C_3, C, C^0, C^0_b, V, V^0, K, *the mirror calculus,* PF, OB, HB, ED, ED^0, *with its fixed reflexivity clause or entry where it has one. Then* X *derives* $0 = S0$ *from no assumptions if and only if* N *does, if and only if* iPRA *derives falsity. The ordinary realization of* X *is bi-interpretable with* iPRA *with whole domains, actual equality and identity numerical composites.*

Proof. Each calculus has been shown equivalent to N on Q-free sequents by translations that are the identity on terms, or the map nf_0 on terms for the zero-normal calculi, and the formula composites are the identity and $A \mapsto j\tau A$, respectively $A \mapsto \mathrm{nf}_0(j\tau A)$, which are provably equivalent to A in N (Theorems 4.27 and 5.2). The claim follows from Theorem 1.78. □

Under the ordinary reading every rule is sound in T_eq directly: Refl, Eval, Step and SeedEval follow from the computation equations and equality, Use and the eliminations are ordinary eliminations, an introduction in case mode uses coverage, and an introduction in progress mode uses induction for a literal equation under the assumed base. The separation $0 = S0 \to \bot$ is an axiom of the ordinary realization and not a rule of a clause calculus, in which $0 = S0$ is the designated contradiction; positive equations alone admit a one-element structure.

5.8 Successor rigidity

This section draws on `raw_clausal`. It concerns an arithmetic principle and not a clause calculus. The background theory is intuitionistic first-order logic with equality over the scalar terms, with the two computation equations and coverage $\forall n\,(n = 0 \vee \exists m\, n = Sm)$. Neither the injectivity of S nor the separation of 0 from $S0$ is used. The rule is

$$\frac{\Gamma \vdash f(0) = 0 \quad \Gamma \vdash f(Sz) = S(f(z))}{\Gamma \vdash f(t) = t}\ \mathrm{SR}_0$$

where $f(z)$ is a term with a distinguished variable, not a function variable, and z is fresh for Γ, t and the parameters of f. The second premise has no local assumption. The rule is the uniqueness of iterators restricted to the initializer 0 and the update S. Induction for literal equations derives SR_0: induct on $f(n) = n$, where the step is $f(Sz) = S(f(z)) = Sz$ under $f(z) = z$. Unrestricted uniqueness of scalar iterators also derives it, applied to f and to the identity template with the common initializer 0.

Theorem 5.71 (Zero-equation induction from successor rigidity)**.** *Work in the background theory with* SR_0*. Let* $E(x)$ *be a formula with* $\Gamma \vdash E(0)$ *and* $\Gamma, E(z) \vdash E(Sz)$*,* z *fresh, and let* $e(x)$ *be a term such that, for a fresh* x*,*

$$\Gamma \vdash E(e(x)), \qquad \Gamma, E(x) \vdash e(x) = x.$$

Then $\Gamma \vdash E(t)$ *for every term* t*. In particular, for every term* $q(x)$ *the rule*

$$\frac{\Gamma \vdash q(0) = 0 \quad \Gamma, q(z) = 0 \vdash q(Sz) = 0}{\Gamma \vdash q(t) = 0}\ \mathrm{ZI}$$

is derivable.

Proof. Let $r(n) := I(n, 0; u.\, e(Su))$. First, $\Gamma \vdash E(r(n))$, by coverage on n. If $n = 0$ then $r(n) = 0$ and the base applies. If $n = Sm$ then $r(n) = e(S(r(m)))$, and the first condition on e at $S(r(m))$ applies. This step assumes nothing about $r(m)$ and uses no induction. Next instantiate the given step at $r(z)$; with $E(r(z))$ it gives $E(S(r(z)))$, and the second condition on e gives $e(S(r(z))) = S(r(z))$. The computation equations therefore prove $r(0) = 0$ and $r(Sz) = S(r(z))$ in the context Γ. SR_0 gives $r(t) = t$, and $E(r(t))$ transports to $E(t)$. The eigenvariable of SR_0 and the witness of the coverage elimination are fresh, so Γ may contain arbitrary formulas.

For ZI take $E(x) \equiv (q(x) = 0)$ and $e(x) := \mathrm{sel}(q(x), x, 0)$. If $q(x) = 0$ the selector computes to x, which is the second condition. For the first apply coverage to $q(x)$. If $q(x) = 0$ then $e(x) = x$ and $q(e(x)) = 0$. If $q(x) = Sv$ then $e(x) = 0$ and the base $q(0) = 0$ gives $q(e(x)) = 0$. These cases need no test of equality of two terms and no separation. □

The construction of e and r is effective on syntax, so the theorem is a transformation of derivations. The proof uses one trajectory and no idempotence lemma; `scope_kernel` obtains induction from unrestricted uniqueness by a retraction argument. The theorem does not give full scalar arithmetic from SR_0; see Theorem 5.72.

5.9 Checker sizes

This section collects the measurements of the eleven packages named in Section 5.7. All of them measure complete standalone Python checkers by the protocol of Section 1.10, under CPython 3.12.14: the module without its docstring, printed by `ast.unparse`, with one final line feed. Each count includes parsing, validation of syntax and scope, substitution, the rules and the acceptance interface. The checkers read one JSON value, a closed goal and a proof, and accept closed theorems; open sequents occur in the branches. The universal statements of this chapter are the written proofs; the packages' finite test suites concern the implementations.

Representation. With three exceptions the checkers share one codec: 0 is the integer 0, a free parameter is a negative level, a bound variable is a positive index, St is `[0,t]`, the iterator is `[n,a,h]`, an equation is `[s,t]` and $D_{x,y}B$ is `[B]`, the expected sort separating a successor from an equation. Every inserted term has no loose positive index, so opening a binder needs no shifting; an elimination or introduction assigns one new negative level and validates its conclusion in the outer scope, which enforces the eigenvariable condition. `clausal_two_case` uses tagged lists and infers the conclusion of an elimination; it proves, by induction on a successful computation, that a returned formula is well scoped and derivable in C from the retained context, and that every derivation of a closed theorem has a certificate. `clause_contexts` writes St as `[t]` and $D_{x,y}B$ as `["d",B]`. `prefix_clauses` stores a formula as `[k,s,t]` and St as `[t]`; the expected sort separates a record from an iterator.

Protocols. `clausal_single_evaluation` and `clause_contexts` check goal-directed certificates in which a validated reflexive goal closes without a proof node. The selected checker of `clause_contexts` reads a JSON array of two-field packets followed by the goal, from right to left; the shape of the first field selects Use, Close, Open or Step, an unconsumed prefix is rejected, and an unconditional local clause is stored with the antecedent $0 = 0$, which closes without input. The package proves that a successful check yields a derivation of V^0 and that every derivation of a closed theorem has an accepted certificate, by induction on the call tree and by the invariant that a call consumes exactly its own segment. In `prefix_clauses` and `evaluation_discharge` the premise field of a Use whose premise is omitted or absent is an inert field of the format. `evaluation_discharge` infers every intermediate conclusion from the proof tree and supplies no template to Discharge.

What the measurements show. The deletion of the root row saves 47 bytes, 5 tokens and 7 AST nodes in an unchanged format. Uniform clauses save 80 bytes against the same validator. Zero-normal syntax with one evaluation scheme is smaller than its raw control in `clausal_single_evaluation` and `rootfree_clausal` and larger in the companion of `reduced_clause_calculus`. The diagonal checker is 100 bytes larger than the witness-mode checker of `rootfree_clausal`. In `clause_contexts` the factorization of clauses alone saves 53 bytes, 2 tokens and 5 AST nodes against the strongest control, and the rest of its reduction comes from the packet protocol and shared validation; its ranked representation is larger than its recursive one. In `prefix_clauses` the prefix representation alone saves 3 tokens and 63 bytes and adds 10 AST nodes against its nested control. The one-body checker and the homogeneous checker are larger than their controls, by the cost of the gate and its lifting. The fusion at the discharge saves 103 bytes, 22 tokens and 22 AST nodes on raw terms and 21 bytes, 12 tokens and 10 AST nodes on zero-normal terms. `one_body` also reports a refinement of an existing checker for the native calculus on the equation and D language that changes no rule.

Table 5.1: Checker sizes as stated by the packages: tokens, AST nodes, normalized bytes. A control is given as tokens/AST nodes/bytes.

Package	Program	Tok.	AST	Bytes	Matched control
`raw_clausal`	five rules, three rows (C_3)	853	993	3046	canonical five-rule control 903/1035/3175; raw six-rule control 930/1072/3252; upstream canonical checker 946/1072/3305
`rootfree_clausal`	raw, two rows (C)	837	977	2956	matched raw control 842/984/3003; incoming raw clausal 853/993/3046
`rootfree_clausal`	zero-normal, fixed zero base ($\mathsf{C}^0_{\mathrm{b}}$)	852	987	2948	zero-normal control with root row 864/1006/2993
`rootfree_clausal`	zero-normal, witness test (C^0)	853	992	2948	the same control; tagged native normal comparator 979/1047/3011
`reduced_clause_calculus`	raw, two rows (C)	848	986	2999	exact raw clause control 853/993/3046
`reduced_clause_calculus`	companion, one scheme (C^0)	871	1011	3050	zero-normal with root row 879/1023/3107

Package	Program	Tok.	AST	Bytes	Matched control
`clausal_two_case`	two-case checker (C)	794	868	2787	matched three-case control 808/888/2861; pinned three-case checker 853/993/3046; the same with the root row deleted 848/986/2999
`clausal_single_evaluation`	single evaluation (C^0)	741	826	2283	matched raw control 744/842/2416; zero-normal control with root row 765/865/2426
`clausal_single_evaluation`	forward implementation	867	1008	3022	forward control 853/993/3046
`uniform_clauses`	uniform clauses (V)	807	931	2912	raw clause checker 853/993/3046; with shared validator 825/958/3039; two rows 848/986/2999; two rows, shared validator 820/951/2992
`clause_guards`	diagonal (K)	872	1013	3056	native diagonal control 975/1138/3395; witness-mode clause control 837/977/2956
`clause_guards`	mirror guard	874	1012	3059	the same
`clause_contexts`	four schemes, packets (V^0)	666	770	2282	pinned five-pattern tagged source 829/902/2950; five-pattern arity and scope control 753/828/2730
`clause_contexts`	packets, empty antecedent	665	770	2308	the same
`clause_contexts`	explicit tree checker	751	823	2677	the same
`clause_contexts`	ranked tree checker	781	853	2692	the same
`clause_contexts`	support-rank experiment, not selected	854	957	3069	none
`prefix_clauses`	prefix calculus (PF)	788	900	2700	matched nested control 791/890/2763; `uniform_clauses` 807/931/2912
`prefix_clauses`	homogeneous variant (HB)	840	965	2843	the same
`one_body`	one body (OB)	901	1041	3130	matched two-field control 848/986/2999
`one_body`	native checker, preferred refinement	357	412	778	existing native checker 358/413/788
`one_body`	native checker, byte-priority variant	357	415	777	the same

Package	Program	Tok.	AST	Bytes	Matched control
`evaluation_discharge`	raw fusion (ED)	643	695	1892	raw separate rules 665/717/1995
`evaluation_discharge`	zero-normal fusion (ED^0)	633	697	1858	zero-normal separate rules 645/707/1879; incoming controls 663/751/1983, 683/771/1990 and 636/708/2010

5.10 Further questions

Question 5.72 (Successor rigidity)**.** Do the computation equations, coverage, separation and SR_0 construct a term $\delta(a, b)$ with $\delta(a, b) = 0 \leftrightarrow a = b$ on the whole domain without using induction for arbitrary literal equations? Such a term would turn ZI into induction for every literal equation, and the bridge (Theorem 1.78) would apply. The known proofs of the predecessor (Theorem 1.48) and of the reflection of equality (Theorem 1.52) use inductions on general equations or uniqueness with updates other than S, so importing them is circular. `raw_clausal` claims no converse from ZI to SR_0, no equivalence of SR_0 with unrestricted uniqueness, and no checker for SR_0.

Question 5.73 (A stable introduction at no cost)**.** Is there an introduction whose mode is preserved and reflected by substitution, as in Theorem 5.37, with a complete checker no larger than the witness-mode checker? The diagonal checker of `clause_guards` is larger by 100 bytes. Is there a guard that satisfies the conditions of Theorem 5.39 and is cheaper to test than the diagonal shape?

Question 5.74 (One-body contexts without the cost of the gate)**.** Can the antecedent field be removed, as in Theorems 5.54 and 5.59, by a representation of assumptions that needs neither an origin tag nor a gate with lifting, so that a complete checker becomes smaller? The measured checkers of `one_body` and of the homogeneous variant of `prefix_clauses` are larger than their controls, and a variant of `one_body` with primitive deep reflexivity is larger still (3208 bytes).

Question 5.75 (The identity case of the discharge)**.** Can the identity case of ϱ be replaced by a single arithmetic pattern that is always active, without a supplied template and without enlarging the checker or the arithmetic assumptions?

Question 5.76 (Combinations not measured)**.** `prefix_clauses` and `one_body` ask for the diagonal introduction on the prefix representation and on one-body contexts, with a matched measurement. Neither combination is stated or measured in the packages.

Question 5.77 (Sizes of derivations)**.** Theorem 5.16 bounds inference nodes, not serialized size. `raw_clausal` reports that its translation from the six-rule raw calculus to C_3 produced at most nine times as many nodes on every executed example; no general bound is proved. No package bounds the serialized size of a translated derivation, the cost of closing and reopening a closure with an unused witness, or the duplication of the major premise in Theorem 5.64. `rootfree_clausal`, `reduced_clause_calculus` and `evaluation_discharge` ask whether certificates with sharing recover these costs once reference validation is charged.

The following statements of the packages are negative or unproved and are recorded as such.

- No package claims a smallest presentation or a lower bound over presentations, and none of the checkers of Table 5.1 is smaller than the compact codecs of other packages (516 and 508 normalized bytes under their own input contracts, 788, 780 and 778 bytes for validated public-goal checkers). The packages ask for a fixed common contract, comprising goal grammar, scope checking, normalization and measurement, before any such comparison.

- The formal statements not present in the Lean development are: the calculus $\mathsf{C}_{\mathrm{b}}^{0}$ on clauses (Theorem 5.42), the guard criterion and the mirror guard on clauses (Theorems 5.39 and 5.40), the support quotient (Theorem 5.48), the homogeneous calculus (Theorem 5.59), the zero-normal fusion (Theorem 5.69), the propositions Theorems 5.17 and 5.18, the obstruction Theorem 5.25, the theorem Theorem 5.71, and the correspondence of each Python checker with its calculus.

- `reduced_clause_calculus` implements the translation of certificates from raw to zero-normal syntax only; the converse compiler is not implemented, and the converse is the theorem Theorem 5.22.

- `clause_contexts` reports that neither the ranked representation nor the support quotient reduced its checker, and `prefix_clauses` that the prefix representation does not reduce every coordinate against its nested control.

- `reduced_clause_calculus` asks for a deletion that removes an arithmetic obligation, a term constructor or a logical operation while preserving both composites; the deletions of this chapter concern the proof interface and leave the arithmetic basis unchanged. Theorems 5.18 and 5.25 show that, with the other rules fixed, reflexivity cannot be replaced by finitely many closed seeds and neither evaluation row of C can be dropped.

Chapter 6

Discharge of a witness and restricted equality rules

A derivation in the native scalar core N (Section 1.3) supplies data at three places. The rule Tr takes an equation $s = t$ together with a term template. The rule DE opens a paired formula at a fresh witness and is separate from every equality rule. The rule QE, and the use of a clause that replaces it in a clause calculus (Section 1.9), takes an arbitrary term. The calculi of this chapter fuse or restrict these interfaces.

In Section 6.2 one rule, Name, discharges an opened witness b together with a proved equation $b = t$; it replaces both the equality rule and the elimination of $D_{x,y}$. In Section 6.3 neither the equation nor the term t is supplied: both are read off two rows of the matrix. In Section 6.4 the fusion is carried out on the canonical recursor syntax, where no normalization is available and reflexivity is derived. In Section 6.5 the equality rule substitutes for a variable only, for the newest variable only, or, on raw terms, for a free parameter; an obstruction shows that unary congruence is not enough. In Section 6.6 the use of a clause takes no term argument. Section 6.7 records an induction interface that one of the packages treats separately. Sections 6.8 and 6.9 summarize the checkers, and Section 6.10 lists what is left open.

Throughout, $\equiv$ is literal identity of syntax up to the names of bound variables, a *parameter* is a free numerical variable, and a *literal equation* is a formula whose root is an equation. The numerals $S0$ and $SS0$ are written 1 and 2, and H is the equation $0 = 0$.

6.1 Normal forms, clause contexts, and the reference calculus

This section fixes what is common to `witness_discharge`, `inferred_witness`, `deterministic_opening`, `first_instance`, `clause_discharge` and `variable_substitution`. Full normalization is the subject of Section 4.9 and the clause calculi on raw terms that of Section 5.2; they are cited, and only what is specific to the calculi of this chapter is stated here. The remark on the strict traversal is from `first_instance`.

6.1.1 Conventions on full normal forms

In Sections 6.1 to 6.3 and in Sections 6.5.1 and 6.5.2 every term and every formula is in full normal form: nf is the normal form for the reduction of Theorem 4.5, which exists and is unique by Theorems 4.88 and 4.89, and every displayed instance $B(t)$, $B(t, r)$ or $F[t/b]$ is formed with the normalizing substitution of Theorem 4.25. Substitution is simultaneous: an inserted term is not traversed again, so that $F[t/b]$ is defined also when t contains b. By Theorem 4.92 the map nf is an admissible normalizer; in particular it commutes with injective

renaming, it leaves the root of a formula unchanged, and it is coherent with substitution,

$$\mathrm{nf}(T\sigma) = \mathrm{nf}\big(\mathrm{nf}(T)[x \mapsto \mathrm{nf}(\sigma x)]\big), \tag{6.1}$$

so that instances compose (Theorem 4.26) and it is immaterial whether a compound expression is normalized before or after one of its variables is instantiated. The counter of an iterator in a normal form is neutral (Theorem 4.91).

Let P be the parity term of Theorem 1.3, written N there, $P(y) = I(y, 0;\, u.\, \mathrm{sel}(u, 1, 0))$. On normal forms the computations of Eq. (1.7) and of the selector are literal identities:

$$\mathrm{sel}(0, a, c) \equiv a, \qquad \mathrm{sel}(Sv, a, c) \equiv c, \qquad P(1) \equiv 1, \qquad P(2) \equiv 0. \tag{6.2}$$

No induction, and no statement about P at other arguments, is used anywhere in this chapter.

Remark 6.1 (Termination of a strict traversal). The checkers of this chapter normalize the counter, the seed and the body first and, at a successor counter, normalize the predecessor iterate $I(n, a;\, u.\, h)$ before inserting it into the body. If the body does not mention its state variable, that value is erased by the root contraction, so the strategy is not a reduction sequence of Theorem 4.5 as it stands. `first_instance` justifies it as follows. Add an administrative term $\mathsf{let}\ u = a\ \mathsf{in}\ h$ with the reduction to $h[a/u]$, and replace the successor contraction by the *strict expansion*

$$I(Sn, a;\, u.\, h) \longrightarrow \mathsf{let}\ v = I(n, a;\, u.\, h)\ \mathsf{in}\ h[v/u], \qquad v \text{ fresh},$$

reducing under all constructors and under both components of a let. No new normal forms arise: every let has a root contraction, and that contraction composes with a strict expansion to the original successor contraction. A let is strongly normalizing whenever its value is strongly normalizing and its body sends every strongly normalizing term substituted for u to a strongly normalizing term; this is proved by induction on the sum of the reduction heights of the two components, an internal contraction decreasing the sum and the root contraction being covered by the hypothesis on the body. The proof of Theorem 4.87 uses this fact at the successor root, after obtaining the predecessor iterate by the same lexicographic induction, and the structural induction of Theorem 4.88, extended by the new constructor, gives strong normalization of the auxiliary syntax. The strict traversal is an evaluation strategy of the auxiliary calculus: after a strict expansion it normalizes the value and then contracts the let. Hence it terminates, also when the value is erased, and since forgetting the administrative terms preserves conversion it returns the unique normal form $\mathrm{nf}(t)$.

Remark 6.2 (Bounded checkers). By Theorem 4.102 the normalizer is total recursive and not primitive recursive; it is part of the specification of every calculus on normal forms and is counted in every checker of such a calculus. `witness_discharge`, `inferred_witness`, `first_instance` and `clause_discharge` each supply a second checker with a fuel field, as in Theorem 4.103. The metering differs from the one there: one unit is consumed at every entry into the recursive traversal of syntax and into the recursive proof check, not at every successor contraction. The conclusions are the same, by the same argument. An accepting unbounded run with k metered entries is accepted with fuel k and rejected with fuel $k - 1$, and erasing the fuel turns an accepting bounded run into an accepting unbounded run, so that

$$\exists p\ \ \mathrm{Accept}(A, p) \quad \Longleftrightarrow \quad \exists p\, \exists k\ \ \mathrm{Accept}_{\mathrm{fuel}}(A, p, k);$$

and since every cycle of the call graph passes through a metered entry, the bounded acceptance relation is primitive recursive. Nothing is asserted about a primitive recursive bound on the fuel that a derivable formula needs.

6.1.2 Formulas, clauses, and the two conventions

The formulas of Sections 6.1 to 6.3 are the Q-free formulas of Theorem 5.1 over normal forms, with the closure abbreviation $q_y A(y) = D_{w,y}A(y)$, w unused, of that definition. Contexts are lists of clause occurrences (Sections 1.9 and 5.1.2). For a matrix $B(x, y)$ and a term r the *opened clause* is

$$\mathcal{C}_B(r) := [B(r, 0);\ y.\, B(r, y)].$$

Two conventions for assumptions and reflexivity occur in the packages.

- *The convention with* Refl. Every clause has exactly one antecedent, an assumption F is the clause $\mathrm{ord}(F) = [H;\ v.\, F]$ of Section 5.1.2, and reflexivity is a rule.
- *The convention with the reflexivity clause.* Clauses are the uniform clauses of Theorem 5.26, with at most one antecedent; an assumption F is the unconditional clause $[\ ;\ v.\, F]$; every context contains one distinguished occurrence of the clause $\mathcal{R} = [\ ;\ y.\, y = y]$ of that definition; and there is no reflexivity rule: $t = t$ is a use of $\mathcal{R}$ at t.

In both, Γ, F denotes Γ extended by the assumption clause of F. Theorem 6.13 shows that the two conventions are interchangeable. All proofs below are written in the convention with Refl.

6.1.3 The shared rules and the reference calculus

The rules Use, Elim and Intro are those of Theorem 5.4, read on full normal forms with normalizing substitution; Intro is the witness-selected introduction of Theorem 4.1 with a clause as its assumption. The equality rule of the reference calculus is Tr, as in N.

Definition 6.3 (Shared rules)**.**

$$\frac{}{\Gamma \vdash t = t}\ \textsc{Refl} \tag{6.3}$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash B(t)}\ \textsc{Use}, \qquad [A;\ y.\, B(y)] \text{ an occurrence in } \Gamma \tag{6.4}$$

$$\frac{\Gamma,\ B(t, \eta) \vdash B(t, Sz)}{\Gamma \vdash D_{x,y}B(x, y)}\ \textsc{Intro}, \qquad \eta = \begin{cases} z, & B \text{ is a literal equation and } t \equiv 0, \\ 0, & \text{otherwise} \end{cases} \tag{6.5}$$

$$\frac{\Gamma \vdash D_{x,y}B(x, y) \qquad \Gamma,\ \mathcal{C}_B(b) \vdash A}{\Gamma \vdash A}\ \textsc{Elim} \tag{6.6}$$

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash h(s) = r}{\Gamma \vdash r = h(t)}\ \textsc{Tr} \tag{6.7}$$

In Use the term t is arbitrary; for an unconditional clause the rule has no premise. In Intro the variable z is fresh for Γ, for the witness t and for $D_{x,y}B$; the instance with $\eta = 0$ is the *case mode* and the instance with $\eta = z$ the *progress mode*, and the mode is determined by the literal syntax of the normal forms, not by provable equality. In Elim the variable b is fresh for Γ, A and $D_{x,y}B$. In Tr, $h(w)$ is an arbitrary term template.

Definition 6.4 (The reference calculus)**.** $\mathcal{T}$ is the calculus with the five rules Refl, Use, Intro, Elim and Tr, in the convention with Refl.

$\mathcal{T}$ is the fully normalized clause calculus with which the packages of this section compare themselves. It is the clause form of the six-rule calculus F of Theorem 4.96.

Proposition 6.5 (The reference calculus and the kernel)**.** *For a native formula A let $A^\flat := \tau(\mathrm{nf}(A))$, with τ the Q-free form of Theorem 5.1. Then* N *derives $\Gamma \vdash A$ if and only if $\mathcal{T}$ derives $A^\flat$ from the assumption clauses of the formulas $G^\flat$, $G \in \Gamma$; and* N *derives A from $A^\flat$ and conversely. The translation is the identity on normalized Q-free formulas and fixes $0 = S0$.*

Proof. N derives A from $\mathrm{nf}(A)$ and conversely by Theorem 4.6, and $\mathrm{nf}(A)$ from $A^\flat$ and conversely by Theorem 5.2(iii). By Theorem 4.98 it remains to show, for Γ and A in full normal form, that $\Gamma \vdash_{\mathsf{F}} A$ if and only if $\mathrm{ord}(\tau\Gamma) \vdash_{\mathcal{T}} \tau A$. This is the argument of Theorem 5.11 with F in the place of N and $\mathcal{T}$ in the place of C_3; it is shorter, because F and $\mathcal{T}$ have the same equality rule and the same introduction.

From $\mathcal{T}$ to F, for an annotated context (Theorem 5.3) and its native image: Refl and Tr are rules of F; a Use of an assumption occurrence is Hyp, and a Use of an opened occurrence is QE; Elim is DE; and Intro is WI in the same mode, its new occurrence being an assumption.

From F to $\mathcal{T}$: Hyp is Refl at 0 followed by Use; Refl and Tr are rules; QE is the second rule of Theorem 6.7 below, since $\tau(Q_y B) \equiv q_y \tau B$; and WI is Intro in the same mode, because τ sends literal equations to literal equations and all other formulas to formulas with root D. For DE apply Elim to the translated major premise. In the extended context, under the assumption $\tau B(b, 0)$, the opened clause at Sz gives $\tau B(b, Sz)$, so the first rule of Theorem 6.7 gives $q_y \tau B(b, y)$, which is $\tau(Q_y B(b, y))$; weaken the translated branch and cut this formula into it (Theorem 6.6).

Finally, from $\mathrm{ord}(\tau\Gamma) \vdash_{\mathcal{T}} \tau A$ the first translation gives $\tau\Gamma \vdash_{\mathsf{F}} \tau A$, and Theorem 5.2(iii), which holds in F by Theorem 4.98, with cut replaces $\tau\Gamma$ and τA by Γ and A. □

6.1.4 Structural operations

Lemma 6.6 (Weakening, renaming, cut)**.** *Let $\mathcal{S}$ be a calculus whose rules are among those of Theorem 6.3 and the rules* Name (6.10)*,* Open (6.11)*,* Export (6.12) *and* VSub (6.17) *introduced below, and which contains* Refl *and* Use*. The following are admissible in $\mathcal{S}$, by direct induction on derivations.*

(a) *Weakening: if $\Gamma \vdash A$ and every occurrence of Γ is an occurrence of Γ', then $\Gamma' \vdash A$.*

(b) *Injective renaming: if $\Gamma \vdash A$ and ι is an injective map of parameters, then $\Gamma\iota \vdash A\iota$.*

(c) *Cut of an assumption: if $\Gamma \vdash F$ and $\Gamma, F \vdash A$, then $\Gamma \vdash A$.*

None of the three substitutes a term for a parameter in a derivation.

Proof. The inductions are those of Theorem 5.5; for Refl, Use, Elim, Intro and Tr nothing changes, since normalization commutes with injective renaming. Only the side conditions of the four new rules have to be checked under an injective renaming. Such a renaming preserves and reflects the property of being a variable, the literal identities tested by the selector of Theorem 6.23, and the absence of a parameter from a formula. Hence the selected witness of Open and Export, the source of VSub and the export conditions of Name and Export are preserved. For weakening and cut, no side condition of the new rules refers to the clauses of the context other than through freshness, and the selector depends on the matrix of the major premise alone. □

Substitution of a term for a parameter does not preserve rule instances in general. It can turn a witness into the literal 0 and so change the mode of Intro (Theorems 4.101 and 5.14), it can make the test of the selector succeed, and it can turn the variable source of VSub into a compound term. Admissibility of substitution is proved for the calculus $\mathcal{O}$ in Theorem 6.32 and follows for the others from the equivalences.

6.1.5 Closures, numerical cases, and the two introductions

The lemmas of this subsection and the next are the counterparts on full normal forms of Theorems 5.6, 5.7 and 5.9 and Theorem 5.8. They are restated because the calculi of Sections 6.2 and 6.3 do not have ELIM as a rule, and one of them has it at first for two kinds of matrices only; the hypotheses record what each argument uses.

Lemma 6.7 (Closure and numerical cases)**.** *Let $\mathcal{S}$ contain* USE *and* INTRO *and admit weakening, and suppose that* ELIM *is a primitive or derived rule of $\mathcal{S}$ for every matrix in which the witness coordinate does not occur. Then the following rules are derived, for every formula $A(y)$, every term t and z fresh:*

$$\frac{\Gamma,\ A(0) \vdash A(Sz)}{\Gamma \vdash q_y A(y)} \qquad \frac{\Gamma \vdash q_y A(y) \qquad \Gamma \vdash A(0)}{\Gamma \vdash A(t)} \qquad \frac{\Gamma \vdash A(0) \qquad \Gamma \vdash A(Sz)}{\Gamma \vdash A(t)}$$

No equality rule is used.

Proof. The second rule is Theorem 5.9(ii), whose proof eliminates a matrix with an unused witness coordinate, and the third follows from the first two as in Theorem 5.10. The first rule is obtained differently from Theorem 5.9(i), with no appeal to an unrestricted case introduction: apply INTRO to the matrix $A(y)$, with the unused witness coordinate, at the witness 1; since $1 \not\equiv 0$ the rule is in case mode at every root, and its premise is the given one. □

The third rule is a distinction of numerical cases, not an induction: its successor premise has no hypothesis.

For a literal equation E let $G(v, E)$ be the gate of Theorem 4.99, with $G(0, E) \equiv H$ and $G(Sw, E) \equiv E$, and let the *envelope* in the present formulas be $L(E) := q_v G(v, E)$.

Lemma 6.8 (The two unrestricted introductions)**.** *Let $\mathcal{S}$ contain* REFL*,* USE *and* INTRO*, admit weakening and cut, and have* ELIM *as a primitive or derived rule for every matrix in which the witness coordinate does not occur and for every matrix with root D. Then for every matrix B, every literal equation $E(x, y)$ and every witness t the rules*

$$\frac{\Gamma,\ B(t,0) \vdash B(t,Sz)}{\Gamma \vdash D_{x,y} B(x,y)} \qquad \frac{\Gamma,\ E(t,z) \vdash E(t,Sz)}{\Gamma \vdash D_{x,y} E(x,y)}$$

are derived, with z fresh as in INTRO*.*

Proof. Case introduction. INTRO is the first rule unless B is a literal equation $E(x, y)$ and $t \equiv 0$. For that case the derivation is the one given after Theorem 4.99 for the missing case of Theorem 4.98, read in the present formulas: Q_v becomes q_v; the two derivations $\Gamma, E \vdash L(E)$ and $\Gamma, L(E) \vdash E$ of Theorem 4.99 use the first and the second rule of Theorem 6.7, with REFL for the base H; the matrix $L(E(x, y))$ has root D, so it is introduced at the witness 0 in case mode and opened by ELIM at a fresh c; and the retained closure is the opened clause, used by USE at Sz. The derivation eliminates matrices with an unused witness coordinate and one matrix with root D, and no others.

Progress introduction. This is the third statement of Theorem 5.10, with the same proof: INTRO for the matrix $E(t, y)$ at the literal witness 0 of an unused coordinate is in progress mode and yields $q_y E(t, y)$; in the context $\Gamma, E(t, 0)$ the second rule of Theorem 6.7 gives $E(t, Sz')$; and the case introduction at the witness t concludes. □

6.1.6 Equality from one term transport

Uncrossed transport is the rule

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash h(s) = r}{\Gamma \vdash h(t) = r}\ \textsc{UTr} \tag{6.8}$$

for an arbitrary term template $h(w)$.

Lemma 6.9 (Equality algebra)**.** *Let $\mathcal{S}$ have Refl, as a rule or derived. If UTr is derivable in $\mathcal{S}$, or if Tr is, then so are both, and so are symmetry, transitivity, the congruence $s = t \vdash h(s) = h(t)$, and transport in every equation template: $s = t,\ p(s) = q(s) \vdash p(t) = q(t)$.*

Proof. From UTr: with $h(w) = w$ and $r \equiv s$ the premises are $s = t$ and the reflexive $s = s$, and the conclusion is $t = s$; this is symmetry, and UTr followed by symmetry is Tr. From Tr and reflexivity, Theorem 4.2(b) gives symmetry, transitivity, congruence and transport in an equation template, and Tr followed by symmetry is UTr. □

Lemma 6.10 (Transport in a formula)**.** *Let $\mathcal{S}$ contain Use and Intro, admit weakening, have Elim as a primitive or derived rule for all matrices, and derive symmetry and transport in every equation template. Then for every formula template $F(w)$*

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash F(s)}{\Gamma \vdash F(t)}$$

is derived in $\mathcal{S}$.

Proof. The proof of Theorem 5.8 applies, by induction on the syntax of the template. Its one appeal to Theorem 5.7 is an Intro at the fresh parameter opened by Elim; a parameter is not the literal 0, so that instance is in case mode whatever the root of the matrix is, and the case introduction at the witness 0 of Theorem 6.8 is not needed. □

6.1.7 Two-row packets

The device by which a calculus without a general equality rule recovers Tr is a paired formula whose matrix has prescribed rows at $y = 0, 1, 2$. It first occurs in `variable_substitution`.

Definition 6.11 (Packet)**.** Let $E_1(x) = (a_1 = c_1)$ and $E_2(x) = (a_2 = c_2)$ be equations in which x may occur and y does not. The *packet* with rows E_1, E_2 is the matrix

$$\Pi(x, y) := \Big(\mathrm{sel}(y, 0, \mathrm{sel}(P(y), a_2, a_1)) = \mathrm{sel}(y, 0, \mathrm{sel}(P(y), c_2, c_1))\Big).$$

By (6.2) its rows are literally

$$\Pi(x, 0) \equiv H, \qquad \Pi(x, 1) \equiv E_1(x), \qquad \Pi(x, 2) \equiv E_2(x). \tag{6.9}$$

Lemma 6.12 (Introduction of a packet)**.** *Let $\mathcal{S}$ contain Use and Intro, admit weakening, and have the numerical-case rule of Theorem 6.7. If $\Gamma \vdash E_1(s)$ and $\Gamma \vdash E_2(s)$, then $\Gamma \vdash D_{x,y}\Pi(x, y)$. In any context that contains the opened clause $\mathcal{C}_\Pi(b)$, Refl and Use derive $E_1(b)$ and $E_2(b)$.*

Proof. Apply Intro at the witness s. Its goal is

$$\Pi(s, Sz) \equiv (\mathrm{sel}(P(Sz), a_2(s), a_1(s)) = \mathrm{sel}(P(Sz), c_2(s), c_1(s))).$$

Let $A(v) := (\mathrm{sel}(v, a_2(s), a_1(s)) = \mathrm{sel}(v, c_2(s), c_1(s)))$. Then $A(0) \equiv E_2(s)$, $A(Sz') \equiv E_1(s)$ and, by (6.1), $A(P(Sz)) \equiv \Pi(s, Sz)$. The numerical-case rule at the target $P(Sz)$, with the two given derivations weakened to the local context and an eigenparameter z' chosen fresh independently of z, proves the goal. The assumption of Intro is not used, so the derivation is valid in either mode, in particular when $s \equiv 0$. The second statement is (6.9): the antecedent of $\mathcal{C}_\Pi(b)$ is H, proved by Refl, and Use at 1 and at 2 gives the two rows at b. □

6.1.8 Interchangeability of the two conventions

Lemma 6.13 (Reflexivity as a fixed clause)**.** *Let $\mathcal{S}$ be a calculus in the convention with* Refl *whose other rules are among* Use, Intro, Elim, Tr, Name, Open, Export *and* VSub*, and let $\mathcal{S}^{\mathcal{R}}$ have the same rules without* Refl*, in the convention with the reflexivity clause. For a context Δ of uniform clauses let $L(\Delta)$ be the context Δ° of Theorem 5.26, which replaces every unconditional occurrence $[\,;\, y.\, B]$ by $[H;\, y.\, B]$. Then for every formula F*

$$\mathcal{R}, \Delta \vdash_{\mathcal{S}^{\mathcal{R}}} F \quad \Longleftrightarrow \quad L(\Delta) \vdash_{\mathcal{S}} F.$$

The statement covers arbitrary contexts, not only those that arise in a derivation from $\mathcal{R}$ alone.

Proof. The two inductions of the proof of Theorem 5.27 apply. They translate only the uses of $\mathcal{R}$, the uses of unconditional clauses, Refl, and the assumption added by Intro, and they copy every other rule; occurrences correspond by position, also when two have the same value. The rules that are copied here are Tr, VSub, and the rules Elim, Name, Open and Export, each of which adds the clause $\mathcal{C}_B(r)$ with one antecedent on both sides and has the same premises and side conditions in the two conventions. □

6.2 Naming packets: equality at the discharge of a witness

This section draws on `witness_discharge`. The packet is that of `variable_substitution` (Section 6.5.1); its final substitution and the elimination of its witness become one inference here.

6.2.1 The rule Name

Equality substitution is performed only while the fresh witness of a paired formula is discharged:

$$\frac{\Gamma \vdash D_{x,y}B(x,y) \qquad \Gamma,\ \mathcal{C}_B(b) \vdash b = t \qquad \Gamma,\ \mathcal{C}_B(b) \vdash F}{\Gamma \vdash F[t/b]}\ \textsc{Name} \tag{6.10}$$

The variable b is fresh for Γ and $D_{x,y}B$, and the exported formula $F[t/b]$ must not contain b. The two branch premises have the same extended context. The term t and the formula F may contain b; in particular $t \equiv b$ is allowed. The side condition concerns the exported normal form and is not a prohibition of b in t.

Definition 6.14 (The calculus of witness discharge)**.** $\mathcal{W}$ is the calculus with the four rules Refl, Use, Intro and Name, in the convention with Refl. $\mathcal{W}^{\mathcal{R}}$ has the three rule families Use, Intro and Name and the fixed clause $\mathcal{R}$. There is no further logical, equality, computation, case, induction, weakening or cut rule.

Lemma 6.15 (Soundness of Name)**.** *Name is sound for the ordinary reading in T_{eq} (Section 1.5), for every formula F and also when t contains b.*

Proof. Eliminate the existential quantifier in the reading of the major premise at a fresh b; the local assumption is the reading of $\mathcal{C}_B(b)$. The two branch derivations give $b = t$ and F. Equality substitution in the template $F[w/b]$ gives the raw instance at t, and the computation equations identify it with its normal form $F[t/b]$. This step does not need $b \notin \mathrm{FV}(t)$: an equation $b = t(b)$ supports the simultaneous substitution of $t(b)$ for b in a fixed formula. What the elimination of the quantifier needs is the absence of b from the result, which is the side condition. □

USE has the reading of its clause, and INTRO is sound by coverage in case mode and by induction for literal equations in progress mode.

Lemma 6.16 (Ordinary elimination)**.** *ELIM is a derived rule of $\mathcal{W}$.*

Proof. Take $t \equiv b$ in NAME. The equality premise is REFL, the other premise is the given derivation of A, and $A[b/b] \equiv A$ does not contain b. □

By Theorems 6.6 and 6.16 the closure rules, the numerical cases, the two unrestricted introductions and the introduction of packets (Theorems 6.7, 6.8 and 6.12) are available in $\mathcal{W}$. None of them uses an instance of NAME other than with $t \equiv b$.

6.2.2 Transport from one discharge

Theorem 6.17 (Transport in $\mathcal{W}$)**.** *For every term template $h(w)$ the rule UTR (6.8) is derived in $\mathcal{W}$. Apart from ordinary eliminations, its derivation contains one instance of NAME.*

Proof. Let $\Gamma \vdash s = t$ and $\Gamma \vdash h(s) = r$, let x be fresh for h, r and t, and let Π be the packet with the rows

$$E_1(x) = (h(x) = r), \qquad E_2(x) = (x = t).$$

The two given equations are $E_1(s)$ and $E_2(s)$, so $\Gamma \vdash D_{x,y}\Pi$ by Theorem 6.12. Apply NAME with a fresh b. In the context $\Gamma, \mathcal{C}_\Pi(b)$ the same lemma gives the row $b = t$, which is the equality premise, and the row $h(b) = r$, which is taken as F. The exported formula is

$$(h(b) = r)[t/b] \equiv (h(t) = r)$$

by (6.1), because b occurs neither in h nor in r; it does not contain b. The identity is one of normal forms and not an application of an equality rule. □

Corollary 6.18 (Equality in $\mathcal{W}$)**.** *$\mathcal{W}$ derives symmetry, transitivity, congruence, TR, and transport in every formula template.*

Proof. Theorem 6.9 applies by Theorem 6.17; in particular symmetry is obtained from the packet with the identity template and a reflexive minor premise, and is not assumed by it. Theorem 6.10 applies by Theorem 6.16. □

6.2.3 Equivalence

Theorem 6.19 ($\mathcal{W}$ and the reference calculus)**.** *For every context Γ and formula F,*

$$\Gamma \vdash_{\mathcal{W}} F \quad \Longleftrightarrow \quad \Gamma \vdash_{\mathcal{T}} F.$$

Both directions are effective transformations of derivations that leave the end sequent unchanged.

Proof. From $\mathcal{T}$ to $\mathcal{W}$, translate a derivation recursively: ELIM is replaced by Theorem 6.16 and TR by Theorem 6.18; the other three rules are shared. The constructions that replace a TR node were derived without TR and without a translated node, so the recursion is on the given derivation alone.

From $\mathcal{W}$ to $\mathcal{T}$: $\mathcal{T}$ derives symmetry and UTR from TR by Theorem 6.9, and hence transport in every formula template by Theorem 6.10. At a NAME node the translated branch premises give $b = t$ and F in the context $\Gamma, \mathcal{C}_B(b)$. Transport in the template $F[w/b]$, w fresh, gives $F[t/b]$ in that context; the substitution is simultaneous, so this is correct also when t contains b. By the side condition the result does not contain b, and ELIM discharges the clause. The finitely many eigenparameters are renamed before each replacement. □

Corollary 6.20 (Three rule families)**.** *For every context Δ of clauses with at most one antecedent and every formula F, $\mathcal{R}, \Delta \vdash_{\mathcal{W}^{\mathcal{R}}} F$ if and only if $L(\Delta) \vdash_{\mathcal{T}} F$. Consequently* N *derives $\Gamma \vdash A$ if and only if $\mathcal{W}^{\mathcal{R}}$ derives $A^\flat$ from $\mathcal{R}$ and the unconditional assumption clauses of the formulas of $\Gamma^\flat$, and the ordinary reading of $\mathcal{W}^{\mathcal{R}}$ is bi-interpretable with* iPRA *in the sense of Theorem 1.78: whole domains, actual equality, both numerical composites with graph $x = y$, and both formula round trips.*

Proof. The first statement is Theorems 6.13 and 6.19; the second adds Theorem 6.5. The comparisons are the identity on normalized terms and formulas and fix $0 = S0$, so the interpretations and composites of Theorem 1.78 are unchanged. □

Remark 6.21 (What the count of three means; Lean). The number three counts inference patterns after reflexivity has been moved into one explicitly listed axiom clause; it does not say that reflexivity is proved from nothing. The reduction specific to `witness_discharge` is the localization of all equality elimination at the discharge of a witness; the clauses with at most one antecedent and the fixed clause are choices of presentation taken from `uniform_clauses`.

In Lean the kernel form is `NMPrf`: on full normal forms, with Name in place of Tr and DE and with the hypothesis $Q_y B(b, y)$ in the opened context, the kernel has the six rules Hyp, Refl, Name, QE, DI and QI, and `nmprf_iff_nprf` (`NativeWitnessDischarge`) proves it equivalent to N; the packet is `NMPrf.packet`. The clause form $\mathcal{W}^{\mathcal{R}}$ is `WDPrf`, with `wdprf_iff_nprf` (`NativeNameClauses`).

Remark 6.22 (Substitution in $\mathcal{W}$). The substitution inside Name acts on the proved formula and is justified by the proved equation; it is not a substitution into the branch derivation. Substitution of a term for a parameter in a derivation of $\mathcal{W}$ is admissible through Theorem 6.19 and the admissibility in $\mathcal{O}$ (Theorems 6.32 and 6.36), not by an induction that preserves the rule at each node.

6.3 The inferred witness

This section draws on `inferred_witness`, `deterministic_opening`, `first_instance` and `clause_discharge`. The four packages contain one idea in two forms; Theorem 6.37 records how they differ.

6.3.1 The selector and the two forms of the rule

Name has three premises, a supplied term t and a supplied formula F. The equation $b = t$ need not be proved: it can be read off the opened clause.

Definition 6.23 (Selector)**.** For a matrix $B(x, y)$ and a variable b that does not occur in $D_{x,y}B$, the *selected witness* is

$$\rho_B(b) := \begin{cases} r, & B(b,0) \equiv H \ \text{ and } \ B(b,1) \equiv (b = r), \\ b, & \text{otherwise.} \end{cases}$$

In the first case the test *succeeds*; in the second the selector *falls back*.

The test inspects two rows of the matrix, at $y = 0$ and at $y = 1$, and compares normal forms literally. The left side of the row at 1 must be the variable b itself; an equation $r = b$, an equation provably of that shape, and an equation whose left side merely contains b are not recognized. The right side r may contain b. There is no search, no decision of provable equality, and no choice left to the certificate.

Lemma 6.24 (Properties of the selector). (a) *For an injective renaming ι, $\rho_{B\iota}(\iota b) \equiv \rho_B(b)\iota$, and the test succeeds for $B\iota$ at ιb if and only if it succeeds for B at b.*

(b) *If x does not occur in B, or if B has root D, the selector falls back.*

(c) *If the test succeeds for B with $\rho_B(b) \equiv r$, and σ is a substitution with $\sigma b \equiv b$ under which b remains fresh for $D_{x,y}B\sigma$, then it succeeds for $B\sigma$ with $\rho_{B\sigma}(b) \equiv r\sigma$.*

(d) *A fallback is not stable under substitution.*

Proof. (a) Normalization commutes with ι, and ι preserves and reflects literal identity. (b) If x does not occur in B, the row $B(b,1)$ does not contain b and is not an equation with left side b. If B has root D, so has $B(b,0)$, because normalization does not remove a formula constructor, and it is not H. (c) By (6.1), $(B\sigma)(b,0) \equiv B(b,0)\sigma \equiv H\sigma \equiv H$ and $(B\sigma)(b,1) \equiv (b = r)\sigma \equiv (b = r\sigma)$. (d) For a parameter a and a term r let $B(x,y) := \big(\mathrm{sel}(y,a,x) = \mathrm{sel}(y,a,r)\big)$. Then $B(b,0) \equiv (a = a) \not\equiv H$ and the selector falls back, while under $a \mapsto 0$ the rows are H and $b = r$ and the test succeeds. □

Lemma 6.25 (The selected equation). *In every context that contains $\mathcal{C}_B(b)$, the equation $b = \rho_B(b)$ is derived by* Refl *and* Use.

Proof. If the selector falls back, the equation is $b = b$. If the test succeeds, the antecedent of $\mathcal{C}_B(b)$ is H, proved by Refl, and Use at 1 gives $B(b,1) \equiv (b = \rho_B(b))$. □

The witness can be used in two ways. Either the clause is opened at the selected witness and the goal is unchanged,

$$\frac{\Gamma \vdash D_{x,y}B(x,y) \qquad \Gamma,\ \mathcal{C}_B(\rho_B(b)) \vdash A}{\Gamma \vdash A}\ \textsc{Open} \tag{6.11}$$

with b fresh for Γ, A and $D_{x,y}B$; or the clause is opened at the fresh variable and the selected witness is substituted in the exported formula,

$$\frac{\Gamma \vdash D_{x,y}B(x,y) \qquad \Gamma,\ \mathcal{C}_B(b) \vdash F}{\Gamma \vdash F[\rho_B(b)/b]}\ \textsc{Export} \tag{6.12}$$

with b fresh for Γ and $D_{x,y}B$ and b not in the exported formula. In Open the parameter b stays available in the branch whether or not the selector returns it, and the clause $\mathcal{C}_B(\rho_B(b)) = [B(\rho_B(b),0);\ y.\, B(\rho_B(b),y)]$ may contain it. Both rules have two premises and no term argument. Against Name, the rule Export drops the equality premise and the term t; the rule Open drops in addition the formula F.

Definition 6.26 (The two calculi of the inferred witness). $\mathcal{O}$ has the rules Refl, Use, Intro and Open; $\mathcal{X}$ has the rules Refl, Use, Intro and Export. Their forms $\mathcal{O}^{\mathcal{R}}$ and $\mathcal{X}^{\mathcal{R}}$ with the reflexivity clause have three rule families each.

Proposition 6.27 (Soundness). *Open and Export are sound for the ordinary reading in T_{eq}, also when $\rho_B(b)$ contains b. Moreover every instance of Export is derivable in $\mathcal{W}$ with the same end sequent.*

Proof. Eliminate the existential quantifier of the major premise at b; the local assumption is the reading of $\mathcal{C}_B(b)$, and Theorem 6.25 gives $b = \rho_B(b)$. For Export, equality substitution gives $F[\rho_B(b)/b]$, which does not contain b. For Open, equality substitution in the template $\mathcal{C}_B(v)^\circ$ gives the reading of $\mathcal{C}_B(\rho_B(b))$, the branch gives A, and A does not contain b. No occurs check on the selected witness is needed: the rules do not assert that the equation $b = r(b)$ has a solution independently of the major premise. The last statement: keep the two premises of Export, insert the derivation of Theorem 6.25, and apply Name with $t \equiv \rho_B(b)$. □

6.3.2 The export form

Lemma 6.28 (Ordinary elimination in $\mathcal{X}$)**.** *ELIM is a derived rule of $\mathcal{X}$, by one EXPORT.*

Proof. Apply EXPORT with $F \equiv A$. Since A does not contain b, $A[\rho_B(b)/b] \equiv A$ whichever term the selector returns. The argument does not need the selector to fall back. □

Hence Theorems 6.7, 6.8 and 6.12 hold in $\mathcal{X}$.

Theorem 6.29 (Transport in $\mathcal{X}$)**.** *UTR is derived in $\mathcal{X}$ for every term template $h(w)$.*

Proof. Let $\Gamma \vdash s = t$ and $\Gamma \vdash h(s) = r$, and let Π be the packet with the rows, exchanged against Theorem 6.17,

$$E_1(x) = (x = t), \qquad E_2(x) = (h(x) = r).$$

Theorem 6.12 gives $\Gamma \vdash D_{x,y}\Pi$; the equation $s = t$ is used here, in the proof of the major premise, and not as a premise of the discharge. Apply EXPORT with a fresh b. By (6.9), $\Pi(b, 0) \equiv H$ and $\Pi(b, 1) \equiv (b = t)$, so the test succeeds and $\rho_\Pi(b) \equiv t$. In the branch, USE of $\mathcal{C}_\Pi(b)$ at 2 with the premise H gives $F \equiv (h(b) = r)$. The exported formula is $(h(b) = r)[t/b] \equiv (h(t) = r)$ by (6.1), and it does not contain b. □

The row that the rule must read is at the fixed position 1; the row that the branch supplies is read by USE at 2.

Theorem 6.30 ($\mathcal{X}$ and $\mathcal{W}$)**.** $\Gamma \vdash_{\mathcal{X}} F$ *if and only if* $\Gamma \vdash_{\mathcal{W}} F$*, for every context* Γ *and formula* F*, by effective transformations that leave the end sequent unchanged.*

Proof. From $\mathcal{X}$ to $\mathcal{W}$ use the last statement of Theorem 6.27 at every EXPORT node. From $\mathcal{W}$ to $\mathcal{X}$: by Theorems 6.9, 6.10, 6.28 and 6.29, $\mathcal{X}$ derives transport in every formula template. At a NAME node the translated branches give $b = t$ and F under the opened clause; transport in the template $F[w/b]$ gives $F[t/b]$; by the side condition of NAME it does not contain b, and Theorem 6.28 discharges the witness. This covers a term t that contains b. The order of the constructions is: ordinary elimination, numerical cases, the packet, transport in a term, transport in a formula; none uses a later one, and no substitution into a derivation is used. □

6.3.3 The open form

In $\mathcal{O}$ ordinary elimination is not immediate: the branch of OPEN assumes $\mathcal{C}_B(\rho_B(b))$ where ELIM assumes $\mathcal{C}_B(b)$, and it does not help that the goal is free of b. The assumption is changed by substituting in the branch derivation. The order of the argument avoids a circle: the instances of OPEN on which the selector necessarily falls back give numerical cases and the unrestricted case introduction; these give substitution; substitution gives ordinary elimination.

Lemma 6.31 (Fallback openings)**.** *If the witness coordinate does not occur in B, or if B has root D, then OPEN for B is literally ELIM for B. Consequently Theorems 6.7, 6.8 and 6.12 hold in $\mathcal{O}$.*

Proof. Theorem 6.24(b). The three lemmas require ELIM only for these two kinds of matrices, and weakening and cut hold by Theorem 6.6. □

Theorem 6.32 (Substitution in $\mathcal{O}$)**.** *If $\Gamma \vdash_{\mathcal{O}} A$ and σ is a simultaneous substitution of terms for parameters, then $\Gamma\sigma \vdash_{\mathcal{O}} A\sigma$. The transformation of derivations is effective. It does not preserve the rule instance at each node.*

Proof. By induction on the given derivation, with the statement quantified over all substitutions. The induction hypothesis is applied to proper subderivations of the given derivation only, never to a derivation that the construction has produced. A substitution may send a parameter to a term that contains that parameter; images are inserted once.

REFL is immediate. For USE of an occurrence $[A'; y.\, B]$ at t, the context $\Gamma\sigma$ contains $[A'\sigma; y.\, B\sigma]$, the induction hypothesis gives $A'\sigma$, and USE at $t\sigma$ gives $(B\sigma)(t\sigma) \equiv B(t)\sigma$ by (6.1).

For INTRO at the witness t with the matrix B, rename z away from the images of σ and let $B' \equiv B\sigma$, $t' \equiv t\sigma$. The induction hypothesis gives $\Gamma\sigma, B'(t', \eta) \vdash B'(t', Sz)$ with the old η. A progress instance remains one: the root is still an equation and the literal 0 is still 0. A case instance remains one unless B is a literal equation and $t \not\equiv 0 \equiv t'$. In that event the substituted premise is the premise of the unrestricted case introduction of Theorem 6.8, which holds in $\mathcal{O}$ by Theorem 6.31 and was proved before the present theorem.

For OPEN with the major premise $D_{x,y}B$ and the parameter b, rename b away from the images of σ, let $B' \equiv B\sigma$ and $\rho' \equiv \rho_{B'}(b)$, and translate the major premise by the induction hypothesis. There are two cases, according to the test for the *old* matrix.

If the old test succeeds with $\rho_B(b) \equiv r$, then by Theorem 6.24(c) the new test succeeds and $\rho' \equiv r\sigma$. Apply the induction hypothesis to the branch with σ, which fixes b. Its clause becomes

$$\mathcal{C}_B(r)\sigma \equiv [B'(r\sigma, 0);\ y.\, B'(r\sigma, y)] \equiv \mathcal{C}_{B'}(\rho')$$

by (6.1); this does not need r to be free of b.

If the old selector falls back, the old branch has the clause $\mathcal{C}_B(b)$. The new test may fail or succeed. In both events apply the induction hypothesis to the old branch with the substitution that agrees with σ on the other parameters and sends b to ρ'. The clause becomes $\mathcal{C}_{B'}(\rho')$, and the context and the goal become $\Gamma\sigma$ and $A\sigma$, because Γ and A do not contain b and no image of σ does.

In either case OPEN applies to the translated premises and proves $A\sigma$. □

The second case is necessary: by Theorem 6.24(d) the selector is not an invariant test, and an argument that treats it as one leaves a gap. The checkers of `inferred_witness` and `deterministic_opening` are tested on a derivation whose opening falls back and for which a substitution of a passive parameter makes the test succeed: the old branch is then rejected, and the old branch instantiated at the newly selected witness is accepted.

Corollary 6.33 (Ordinary elimination in $\mathcal{O}$)**.** *ELIM is admissible in $\mathcal{O}$ for every matrix.*

Proof. Given $\Gamma, \mathcal{C}_B(b) \vdash A$ with b fresh for Γ and A, apply Theorem 6.32 with the substitution $b \mapsto \rho_B(b)$ that fixes all other parameters. The result is $\Gamma, \mathcal{C}_B(\rho_B(b)) \vdash A$, the branch premise of OPEN. □

Theorem 6.34 (Transport in $\mathcal{O}$)**.** *UTR is derived in $\mathcal{O}$ for every term template $h(w)$.*

Proof. Take the packet of Theorem 6.29, with the rows $x = t$ and $h(x) = r$; by Theorems 6.12 and 6.31, $\Gamma \vdash D_{x,y}\Pi$. Apply OPEN with a fresh b. The test succeeds with $\rho_\Pi(b) \equiv t$, so the branch has the clause $\mathcal{C}_\Pi(t) = [H; y.\, \Pi(t, y)]$. USE at 2 with the premise H gives $\Pi(t, 2) \equiv (h(t) = r)$, the unchanged outer goal, which does not contain b. There is no equality premise and no substitution in an exported formula; the derivation does not use Theorem 6.32. □

Theorem 6.35 ($\mathcal{O}$ and the reference calculus)**.** *$\Gamma \vdash_{\mathcal{O}} A$ if and only if $\Gamma \vdash_{\mathcal{T}} A$, for every context Γ and formula A, by effective transformations that leave the end sequent unchanged.*

Proof. From $\mathcal{T}$ to $\mathcal{O}$: Elim is replaced by Theorem 6.33, and Tr by Theorem 6.34 followed by symmetry (Theorem 6.9).

From $\mathcal{O}$ to $\mathcal{T}$: $\mathcal{T}$ derives transport in every formula template. At an Open node translate the major premise and open it by Elim at b, keeping the clause $\mathcal{C}_B(b)$. Write $\rho \equiv \rho_B(b)$; Theorem 6.25 gives $b = \rho$. Derive $q_y B(\rho, y)$ by the first closure rule of Theorem 6.7: from its assumption $B(\rho, 0)$ transport along $\rho = b$ in the template $B(w, 0)$ to $B(b, 0)$, use $\mathcal{C}_B(b)$ at Sz, and transport along $b = \rho$ in the template $B(w, Sz)$ to $B(\rho, Sz)$. Elim for $q_y B(\rho, y)$, whose witness coordinate is unused, opens exactly the clause $[B(\rho, 0);\, y.\, B(\rho, y)] = \mathcal{C}_B(\rho)$. The translated branch, weakened by the clause $\mathcal{C}_B(b)$ that is still present, proves A. Both eliminations close, since A contains neither the unused witness nor b. The construction is valid when ρ contains b: throughout the inner part b is an available parameter, and only the outer elimination discharges it. □

6.3.4 Summary and comparison of the four packages

Theorem 6.36 (The inferred witness)**.** *The calculi $\mathcal{T}$, $\mathcal{W}$, $\mathcal{X}$ and $\mathcal{O}$ derive the same sequents. For every context Δ of clauses with at most one antecedent and every formula F the following are equivalent: $L(\Delta) \vdash_{\mathcal{T}} F$; $\mathcal{R}, \Delta \vdash_{\mathcal{W}^{\mathcal{R}}} F$; $\mathcal{R}, \Delta \vdash_{\mathcal{X}^{\mathcal{R}}} F$; $\mathcal{R}, \Delta \vdash_{\mathcal{O}^{\mathcal{R}}} F$. Substitution of terms for parameters is admissible in each of them.* N *derives $\Gamma \vdash A$ if and only if $\mathcal{O}^{\mathcal{R}}$, equivalently $\mathcal{X}^{\mathcal{R}}$, derives $A^\flat$ from $\mathcal{R}$ and the unconditional assumption clauses of $\Gamma^\flat$; and the ordinary readings of $\mathcal{O}^{\mathcal{R}}$ and $\mathcal{X}^{\mathcal{R}}$ are bi-interpretable with* iPRA *as in Theorem 6.20.*

Proof. Theorems 6.19, 6.30 and 6.35 give the first statement, Theorem 6.13 the second. Theorem 6.32 gives substitution in $\mathcal{O}$, and the other calculi inherit it through the equivalences, which leave end sequents unchanged. The rest is Theorem 6.5 and Theorem 1.78. □

In Lean the rule on the kernel, with the selected witness read off the normalized matrix in place of DE, is `FOPrf` with `foprf_iff_nprf` (`NativeInferredWitness`); its completeness proof translates a kernel derivation under an arbitrary substitution and at DE sends the fresh variable to the selected witness whatever it is, as in the second case of the proof of Theorem 6.32; no distinction of cases on the test is needed there. The clause form $\mathcal{O}^{\mathcal{R}}$ is `OCPrf` with `ocprf_iff_nprf` (`NativeOpenClauses`), and $\mathcal{X}^{\mathcal{R}}$ is `XCPrf` with `xcprf_iff_nprf` (`NativeExportClauses`).

Remark 6.37 (How the four packages differ). *Rows inspected.* There is no difference: all four inspect the rows $B(b, 0)$ and $B(b, 1)$ of the normalized matrix, require the first to be literally $0 = 0$ and the second to be literally an equation whose left side is the fresh variable, and take the right side of the second as the witness.

The reflexivity clause. There is no difference either: all four are stated in the convention with the fixed unconditional clause $[\,;\, y.\, y = y]$ and clauses with at most one antecedent, and have three rule families.

Use of the witness. `inferred_witness` and `deterministic_opening` state Open: the clause is opened at the selected witness and the branch proves the unchanged goal. Their proofs are those of Theorems 6.31 to 6.35. `first_instance` and `clause_discharge` state Export: the clause is opened at the fresh variable and the witness is substituted in the exported formula. Their proofs are those of Theorems 6.28 to 6.30, and they need no substitution theorem. `inferred_witness` keeps the export form as a second write-up and as a control checker, and `deterministic_opening` keeps it as an intermediate variant with its own checker.

Reference calculus. `inferred_witness` proves the equivalence with $\mathcal{T}$ in the convention with the reflexivity clause; the other three prove it with $\mathcal{W}^{\mathcal{R}}$. In the translation of Open back to the reference calculus, `inferred_witness` transports the base of the closure in both

directions, as in the proof of Theorem 6.35; `deterministic_opening` distinguishes the two cases of the test and, when it succeeds, uses that the base is H and ignores the assumption of the closure.

Additional material. `first_instance` compares its calculus with the raw clause calculus directly instead of through $\mathcal{W}$, by the argument cited in Theorem 6.5, and contributes Theorem 6.1. `clause_discharge` delivers, besides the checker of $\mathcal{X}^{\mathcal{R}}$, a smaller checker of the unchanged $\mathcal{W}^{\mathcal{R}}$. `deterministic_opening` delivers a reordering of an inherited compact checker that is independent of the new rule (Section 6.8).

Names. The rule is called O* in `inferred_witness`, O in `deterministic_opening`, Name in `first_instance` and Row in `clause_discharge`; the Lean development calls EXPORT Discharge.

Remark 6.38 (The conditions of the test). Three conditions cannot be removed. `deterministic_opening` exhibits, for each, a certificate that the correspondingly weakened checker accepts: without the requirement that the left side of the row at 1 is the fresh witness, and without the requirement that the row at 0 is $0 = 0$, the checker accepts $1 = 0$; without the requirement that the outer goal is in the outer scope, it accepts a formula in which the fresh parameter escapes. In `clause_discharge` the source of the replacement must be the witness opened by the same inference: an equation whose left side is an older parameter, a constant or a compound term triggers no replacement. With an older parameter as source, the argument of Theorem 6.28 fails, since the substitution then acts on a goal that mentions that parameter; the generalization is not part of any of the calculi (Section 6.10).

6.4 Substitutive discharge on the canonical recursor syntax

This section draws on `substitutive_discharge`. Its term syntax, and the passage from that syntax to iterator terms, are those of `canonical_recursors`.

The calculi of Sections 6.2 and 6.3 depend on full normalization: the rows of a packet must be literal before any equality rule is available. If the discharge rule is given a formula template, so that the discharged equation may have an arbitrary source, no packet is needed, the only normalization is a structural erasure, and reflexivity is derived from one computation schema.

6.4.1 Syntax

In this section the convention on normal forms is not in force. Raw terms are

$$t ::= x \mid 0 \mid R(n, a;\, u.\, h),$$

where u is bound in the body h only. Successor is defined,

$$S(a) := R(0, a;\, u.\, 0), \qquad 1 := S0,$$

and the intended recursor is seeded by a successor: $R(0, a; u.h) = Sa$ and $R(Sn, a; u.h) = h[R(n, a; u.h)/u]$. The first equation is made part of the syntax. The *canonicalization* cn canonicalizes all subterms and replaces the body of every R whose counter is the literal 0 by the literal 0; a term is *canonical* if every subterm $R(0, a; u.h)$ has $h \equiv 0$. The operation does not unfold a successor iteration, does not execute a body, and does not test whether a term denotes zero. It never changes the root constructor of a term, so

$$\mathrm{cn}(t) \equiv 0 \quad \Longleftrightarrow \quad t \equiv 0.$$

Terms and formulas of the calculus are canonical, and substitution means capture-avoiding raw substitution followed by cn.

Lemma 6.39 (Coherence of canonicalization)**.** *For every term or formula T and every capture-avoiding substitution σ,*

$$\mathrm{cn}(T\sigma) = \mathrm{cn}\big(\mathrm{cn}(T)[x \mapsto \mathrm{cn}(\sigma x)]\big).$$

Canonical substitution is associative, commutes with renaming, and introduces no free variables beyond those of its arguments.

Proof. By structural induction, after renaming bound variables apart. At a node R: if the counter is the literal 0, both sides erase the body; if a counter different from 0 becomes 0 under the substitution, both sides erase it again; otherwise both sides keep the constructor and the induction hypotheses identify the three subterms. Canonicalization does not turn a constructor other than a variable into the literal 0. Formula binders need only the usual lifted substitution. Associativity follows by applying the identity to two successive raw substitutions. □

Formulas are equations and $D_{x,y}B$. A context is a finite list of clauses $[A;\, y.\, B(y)]$ with one antecedent, and an assumption A is the clause $[0 = 0;\, v.\, A]$. The truth of $0 = 0$ is derived, not assumed, and no clause is fixed in advance. The contradiction is $0 = S0$.

6.4.2 Four rule families

Definition 6.40 (The calculus of substitutive discharge)**.** $\mathcal{K}$ has the following four rules. All displayed expressions are canonical.

$$\frac{}{\Gamma \vdash R(Sn, a;\, u.\, h) = h[R(n, a;\, u.\, h)/u]}\ \textsc{Calc} \tag{6.13}$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash B(t)}\ \textsc{Use}, \qquad [A;\, y.\, B(y)] \text{ an occurrence in } \Gamma$$

$$\frac{\Gamma,\ B(t, \eta) \vdash B(t, Sz)}{\Gamma \vdash D_{x,y}B(x, y)}\ \textsc{Intro}, \qquad \eta = \begin{cases} z, & B \text{ is a literal equation and } t \equiv 0, \\ 0, & \text{otherwise} \end{cases}$$

$$\frac{\Gamma \vdash D_{x,y}B \qquad \Gamma,\ \mathcal{C}_B(b) \vdash s = t \qquad \Gamma,\ \mathcal{C}_B(b) \vdash F(s)}{\Gamma \vdash F(t)}\ \textsc{SDis} \tag{6.14}$$

In Calc the predecessor iterate on the right is canonicalized; at $n \equiv 0$ it is Sa. In Intro, z is fresh for Γ, t and $D_{x,y}B$, and the test is literal. In SDis, b is fresh for Γ and $D_{x,y}B$, the exported formula $F(t)$ does not contain b, and $F(v)$ is an arbitrary formula template whose distinguished variable v is different from b; a constant template is allowed, and s, t and the template may contain b. The calculus $\mathcal{K}_{\mathrm{sep}}$ has Calc, Use, Intro, Elim and the transport rule

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash F(s)}{\Gamma \vdash F(t)}\ \textsc{TrF} \tag{6.15}$$

for every formula template F.

$\mathcal{K}$ has no hypothesis rule, no reflexivity axiom, no separate equality rule, no separate elimination of $D_{x,y}$, no coverage or separation axiom and no structural rule. The one computation schema is Calc; the progress mode of Intro carries the induction. Weakening, exchange, contraction, injective renaming and cut of an assumption are admissible by induction on derivations, as in Theorem 6.6; injective renaming preserves the literal test of Intro. Substitution of a term for a parameter can turn a witness into the literal 0; it is admissible through Theorem 6.46, not by an induction that preserves the rules.

6.4.3 The bootstrap of equality

Let, for a term t and variables u, x, y that do not occur in it,

$$W_t := R(1, 0;\, u.\, t), \qquad J := (W_0 = 0), \qquad \Theta := D_{x,y} J.$$

Calc proves $W_t = t$: the body is constant and ignores the predecessor iterate.

Lemma 6.41 (A closed paired theorem)**.** Θ *is derived by* Calc *and* Intro *alone.*

Proof. Apply Intro at the witness 1. The matrix does not depend on its bound variables, so the goal of the premise is J, which Calc proves without the assumption. Neither the use of an assumption nor reflexivity is involved. □

Lemma 6.42 (Transport in a formula)**.** *TrF is a derived rule of* $\mathcal{K}$*.*

Proof. Use Θ as the major premise of SDis, and weaken the two given derivations into the opened context; they do not mention the fresh witness, and $F(t)$ does not contain it. No law of equality is used. □

Lemma 6.43 (Reflexivity and assumptions)**.** $\Gamma \vdash t = t$ *for every term* t*, and an assumption can be used:* $\Gamma, A \vdash A$*.*

Proof. Calc gives $W_t = t$. Use it as both premises of TrF with the template $F(v) = (v = t)$: the instance at the source is $W_t = t$ and the instance at the target is $t = t$. For an assumption A, take $t \equiv 0$ and apply Use to $[0 = 0;\, v.\, A]$. □

Lemma 6.44 (Equality and ordinary elimination)**.** *Symmetry, transitivity, the congruence* $s = t \vdash h(s) = h(t)$ *and* Elim *are derived rules of* $\mathcal{K}$*.*

Proof. From $s = t$, transport the reflexive $s = s$ in the template $(v = s)$ to $t = s$. From $s = t$ and $t = r$, transport $t = r$ along $t = s$ in the template $(v = r)$ to $s = r$. From $s = t$, transport the reflexive $h(s) = h(s)$ in the template $(h(s) = h(v))$ to $h(s) = h(t)$. For Elim, let the opened branch prove a formula A without b. Apply SDis with the constant template $F(v) = A$ and with the equation J, proved by Calc, as the equality premise; the third premise is the given branch, and the exported formula is A. No packet and no normalization of a selector is needed. □

Theorem 6.45 ($\mathcal{K}$ and the calculus with separate rules)**.** *For every context* Γ *of canonical clauses and every formula* A*,* $\Gamma \vdash_{\mathcal{K}} A$ *if and only if* $\Gamma \vdash_{\mathcal{K}_{\mathrm{sep}}} A$*. The translations are effective and preserve the end sequent, including its parameters.*

Proof. An instance of SDis is TrF inside the opened context followed by Elim, whose side condition is that of SDis. Conversely TrF is Theorem 6.42 and Elim is Theorem 6.44. The other rules are shared. Induct on derivations, renaming eigenparameters when the fixed auxiliary derivations are inserted. □

6.4.4 The canonical basis and the bridge

Proposition 6.46 (Recovered rules)**.** *The following are derived in* $\mathcal{K}$*, with* $q_y B := D_{w,y} B$ *for an unused* w*.*

(a) *Case introduction for every matrix and witness: from* $\Gamma, B(t, 0) \vdash B(t, Sz)$ *infer* $\Gamma \vdash D_{x,y} B$*.*

(b) *From* $\Gamma, B(0) \vdash B(Sz)$ *infer* $\Gamma \vdash q_y B$*; from* $\Gamma \vdash q_y B$ *and* $\Gamma \vdash B(0)$ *infer* $\Gamma \vdash B(t)$*; and, for a literal equation* B*, from* $\Gamma, B(z) \vdash B(Sz)$ *infer* $\Gamma \vdash q_y B$*. Hence induction for literal equations under an arbitrary context and numerical cases for arbitrary formulas.*

(c) *A context that contains the clause* $[B(0);\, y.\, B(y)]$ *proves* $q_y B$*.*

Proof. (a) Only a literal equation at the witness 0 is missing from INTRO. The term $c := W_0$ is canonical, is not the literal 0, and satisfies $c = 0$ by CALC. Given the premise at the witness 0, start from the assumption $B(c, 0)$, transport it to $B(0, 0)$, insert the given derivation by cut, and transport its conclusion $B(0, Sz)$ back to $B(c, Sz)$. INTRO at c is in case mode and concludes $D_{x,y}B$. No induction is used.

(b) The first rule is (a) for the unused coordinate. For the second, ELIM opens $[B(0);\, y.\, B(y)]$, and USE with the proof of $B(0)$ gives $B(t)$. The third is INTRO at the literal witness 0 of the unused coordinate, in progress mode.

(c) Use (a) at the unused witness 1: under its assumption $B(0)$, use the clause at Sz. □

By (c) a derivation of the Eq/Q/D basis of `canonical_recursors` whose elimination of $D_{x,y}$ keeps a formula $Q_y B(b, y)$ is translated by deriving $q_y B(b, y)$ from the opened clause and cutting it into the translated branch; the two representations of a context are related by derivations and are not identified by convention. Reflexivity, symmetry and TRF give the equality rule of that basis, and its elimination of Q is the second rule of (b). Conversely that basis derives formula transport and the elimination of $D_{x,y}$, which validate SDIS, a clause $[A;\, y.\, B]$ being represented by its ordinary reading. So the two have the same consequences on formulas.

Theorem 6.47 (Bridge for $\mathcal{K}$)**.** *The ordinary reading of* $\mathcal{K}$ *and* iPRA *are bi-interpretable with whole domains, actual equality, both numerical composites with graph* $x = y$*, and both formula round trips. In particular they are equiconsistent.*

Proof. From $\mathcal{K}$ to iPRA: interpret each fixed term $R(n, a; u.h)$ by primitive recursion with the seed Sa and the displayed step. The erasure of a body at the counter 0 preserves the value. CALC is a defining equation, USE is implication and universal elimination, INTRO is coverage in case mode and induction for an equation under its assumed base in progress mode, and SDIS is equality substitution inside the scope of an existential elimination.

From iPRA to $\mathcal{K}$: `canonical_recursors` constructs, before any numerical selector, cancellation of successor or reflection of equality is available, a canonical term p with $p(0) = 0$ and $p(Sa) = a$. The iterator of Section 1.1 is then the term

$$\widehat{I}(n, a;\, u.\, h) := p\Big(R(n, a;\, v.\, S(h[p(v)/u]))\Big), \tag{6.16}$$

which satisfies the two computation equations of the iterator: at the counter 0 the value is $p(Sa) = a$, and at a successor CALC exposes an outer successor that p removes. By Theorems 6.43, 6.44 and 6.46 the rules needed by the logical macros and by the arithmetic of Theorems 1.48, 1.52 and 1.62 are available in $\mathcal{K}$, with $\mathrm{sel}(v, a, b)$ read as $\widehat{I}(v, a; u.b)$. This gives the defining equations and the quantifier-free induction of iPRA.

For the numerical composites, work through each fixed primitive recursive description in the order of its dependencies. Once the earlier descriptions are identified, a description and its return translation have the same seed and the same step, and induction for the equation between the two trajectories proves them equal at every number of the whole domain. For the body of an R, pointwise equality of bodies is lifted to the iteration by the same argument; it is not assumed as a congruence. The interpretation of successor returns Sa. Hence both composites are the identity. The logical round trips of Theorem 1.78 identify the reading of $D_{x,y}$ with its returned macro, and induction on formulas extends the numerical

identities to both formula composites. The contradiction $0 = S0$ corresponds to falsum in both directions. □

Remark 6.48 (Checking, the raw companion, Lean). The syntax recursion and the proof recursion of $\mathcal{K}$ are structural, and the only canonicalization erases a body at a literal counter 0. Under a standard coding of finite trees the certificate relation is therefore primitive recursive, with no fuel field, in contrast to Theorem 6.2.

The package contains a companion on raw iterator terms without any normalization: the two computation equations of Section 1.1, the use of a clause, Intro, and the same discharge rule. An assumption F is $[H_0; v.\, F]$ with $H_0 := (I(0,0;\, u.\, 0) = 0)$, one instance of the computation schema; $D_{x,y}H_0$ is introduced at the witness $S0$ without its assumption; transport is the discharge of this formula; reflexivity is the transport of $I(0,t;\, u.\, 0) = t$ along itself in the template $(v = t)$; and Elim is the discharge with a constant template and the equation H_0. The proofs are those of Theorems 6.41 to 6.44.

In Lean the discharge rule on the kernel, in place of Tr and DE, gives the seven-rule calculus `SDPrf`, with `sdprf_iff_nprf` (`NativeSubstitutiveDischarge`): the closed theorem is $D_{x,y}(I(0,0;\, u.\, u) = 0)$, proved by cases from a computation; a discharge of it that does not use the opened context is transport in an arbitrary formula (`SDPrf.tr`), and a constant template gives DE (`SDPrf.dE`). The four families on the canonical recursor syntax are not formalized.

6.5 Restricted substitution

This section draws on `variable_substitution` (Sections 6.5.1 and 6.5.2), `parameter_substitution` (Section 6.5.3) and `substitutive_discharge` (Section 6.5.4).

6.5.1 Substitution for a variable

The convention on normal forms of Section 6.1 is in force. The equality rule substitutes for a parameter and does nothing else:

$$\frac{\Gamma \vdash x = t \qquad \Gamma \vdash u = v}{\Gamma \vdash v = u[t/x]}\ \text{VSub}, \qquad x \text{ a parameter.} \tag{6.17}$$

The rule substitutes simultaneously for every occurrence of x in u. It does not remove x from the parameters and does not substitute in Γ or in v. The term t may contain x. No template, no selected occurrence, no compound source and no matching of a pattern is supplied. A variable bound by an iterator is not a source.

Definition 6.49. $\mathcal{V}$ is the calculus with the five rules Refl, VSub, Use, Elim and Intro, in the convention with Refl.

Theorems 6.6, 6.7 and 6.12 apply to $\mathcal{V}$, in which Elim is primitive. In particular the numerical cases are derived with no equality rule, and neither the case introduction at the witness 0 nor an envelope is needed for them.

Theorem 6.50 (Transport in $\mathcal{V}$)**.** *For every term template $h(w)$ the rule Tr is derived in $\mathcal{V}$. The derivation contains exactly one instance of VSub.*

Proof. Let $\Gamma \vdash s = t$ and $\Gamma \vdash h(s) = r$, and take the packet Π of Theorem 6.17, with the rows $h(x) = r$ and $x = t$. By Theorem 6.12, $\Gamma \vdash D_{x,y}\Pi$. Open it by Elim at a fresh b; in the branch the same lemma gives $b = t$ and $h(b) = r$. One VSub with the major premise $b = t$ and the minor premise $h(b) = r$ yields

$$r = (h(b))[t/b] \equiv h(t)$$

by (6.1). The conclusion does not contain b, and ELIM closes. □

Everything before the one VSUB uses the logical rules, reflexivity and literal computation only: no symmetry, transitivity or congruence, no substitution into a derivation, no reflection of equality and no induction. Occurrences of s in the passive part of h are not touched, because b is fresh for h; normalization may duplicate or erase the distinguished occurrences, and (6.1) accounts for this without counting them.

Theorem 6.51 ($\mathcal{V}$ and the reference calculus)**.** *$\Gamma \vdash_{\mathcal{V}} A$ if and only if $\Gamma \vdash_{\mathcal{T}} A$, for every context Γ and formula A, by effective transformations that are the identity on the end sequent.*

Proof. A VSUB node is one TR: with w fresh let $K(w) := u[w/x]$; then $K(x) \equiv u$ and $K(t) \equiv u[t/x]$, also when t contains x, and TR from $x = t$ and $K(x) = v$ concludes $v = K(t)$. Conversely a TR node is replaced by the derivation of Theorem 6.50 after its premises are translated. The fresh names of that derivation are chosen away from the context, the conclusion and the terms involved, so all conditions of ELIM and INTRO hold. □

Together with Theorem 6.5 and Theorem 1.78 this gives the comparison with N and the bridge to iPRA, as in Theorem 6.20.

Proposition 6.52 (The native form with six rules)**.** *On normal forms with the formulas $s = t$, $Q_y A$, $D_{x,y} A$ and contexts of formulas, the calculus with HYP, REFL, VSUB, QE, DE and the introduction INTRO of Theorem 6.3 derives the same sequents as the calculus with TR in place of VSUB, under the identity translation.*

Proof. VSUB is one TR as before. For the converse the numerical cases use no equality rule: introduce $D_{x,y} F(y)$, with x unused, at the witness 1, ignoring the assumption $F(0)$; the branch of DE has the hypothesis $Q_y F(y)$, and QE with the given proof of $F(0)$ gives $F(t)$. The packet is introduced and eliminated as in Theorem 6.50; the two uses of the clause $[H;\, y.\, \Pi(b, y)]$ become two applications of QE to the hypothesis $Q_y \Pi(b, y)$ with the reflexive premise H, and the final VSUB is unchanged. □

The calculus with TR in this proposition is the six-rule calculus of `full_normalization`. In Lean the kernel on full normal forms with VSUB in place of TR is `VSPrf`, with `vsprf_iff_nprf` (`NativeVariableSubstitution`) and the packet `VSPrf.packet`.

Remark 6.53 (What is reduced, and what the proof needs). The source of the equality rule is a variable, and the action of the rule is the substitution that the binders already require. The freedom of an ordinary equality rule is recovered by introducing, using and eliminating a fresh name. Full normalization is what makes the argument short: the rows of the packet at 0, 1 and 2 are literal before any congruence is available. The proof uses the logical rules; no completeness statement is made for the equality rules in isolation.

6.5.2 The newest variable

In this subsection the parameters of a sequent form a finite ordered list, and the eigenparameter of ELIM or INTRO is put at the front of the list of the premise it belongs to. $\mathcal{V}_0$ is $\mathcal{V}$ with the source of VSUB restricted to the front parameter of the list at its node; with de Bruijn indices the source is the variable of index 0.

Weakening is not admissible in $\mathcal{V}_0$ by an induction that preserves the rule at each node: a parameter inserted in front turns the former front parameter into an older one. Theorem 6.6 is not available, and the completeness proof uses an invariant instead.

Definition 6.54 (Internally sourced derivations)**.** A derivation in $\mathcal{V}_0$ with the initial parameter list Ξ is *internally sourced* if at every VSub node the source is the eigenparameter of an Elim node of the derivation, the VSub node lies in the branch of that node, and no other Elim or Intro node lies between the two.

So the source is never a member of Ξ, and it is the front parameter at its node.

Lemma 6.55 (Safe structural changes)**.** *Let an internally sourced derivation of $\Gamma \vdash A$ over Ξ be given, let ι be an injection of Ξ into a list Ξ', not necessarily monotone, and let Γ' be a context over Ξ' that contains every occurrence of $\Gamma\iota$. Then there is an internally sourced derivation of $\Gamma' \vdash A\iota$ over Ξ'.*

Proof. Rename the eigenparameters of the derivation away from Ξ' and Γ'. A node of the derivation has a sequent over a list $(e_k, \dots, e_1, \Xi)$, where $e_1, \dots, e_k$ are the eigenparameters of the Elim and Intro nodes on the path from the root, and a context Γ, Λ, where Λ consists of the clauses those nodes add. Replace it by the sequent over $(e_k, \dots, e_1, \Xi')$ with the context $\Gamma', \Lambda\iota$ and the formula renamed by ι, which is extended by the identity on the e_i. The initial list is not reset when the transformation descends into a premise. Every instance of Refl, Use, Elim and Intro is carried to an instance, as in Theorem 6.6: normalization and substitution commute with injective renaming, the mode of Intro is preserved, and the added clauses do not affect any side condition. At a VSub node the source is e_k with $k \geq 1$, and it is still the front parameter of the new list and still bound by the same Elim node; inserting, permuting or renaming members of the initial list changes neither. The result is internally sourced over Ξ'. □

The lemma does not give weakening for arbitrary derivations of $\mathcal{V}_0$.

Theorem 6.56 (The newest variable suffices)**.** *Every derivation in $\mathcal{T}$ over a parameter list Ξ has an internally sourced translation in $\mathcal{V}_0$ with the same end sequent and the same list. Consequently $\mathcal{V}_0$, $\mathcal{V}$ and $\mathcal{T}$ derive the same sequents, and weakening, substitution of terms for parameters and Tr are admissible in $\mathcal{V}_0$.*

Proof. By induction on the derivation in $\mathcal{T}$. Refl, Use, Elim and Intro are copied. The translated premises are internally sourced over their own lists; a source that is bound by an Elim node inside a premise is bound by a node inside the combined derivation, so the invariant is preserved.

At a Tr node apply the construction of Theorem 6.50 to the two translated premises. The construction places copies of them under the eigenparameters and the assumption clauses of the Intro of the packet and of the numerical-case rule; each copy is obtained by Theorem 6.55, which applies because the premises are internally sourced by the induction hypothesis. The remaining nodes of the construction are Refl, Use, Elim, Intro and the one VSub. That node stands at the end of the branch of the Elim which opens the packet at b; its premises $b = t$ and $h(b) = r$ are derived from the opened clause by Refl and Use, with no eigenparameter in between. So its source b is the front parameter and is bound by an Elim node of the constructed derivation, with no node of Elim or Intro between them.

Conversely a restricted VSub is an instance of VSub and hence one Tr (Theorem 6.51). So the three calculi agree. A derivation of $\mathcal{V}_0$ is weakened, or instantiated, or used as a premise of Tr, by translating it to $\mathcal{T}$, operating there, and translating back. The completeness proof itself uses none of these operations on arbitrary derivations. □

The same invariant and translation apply to the six-rule form of Theorem 6.52, with QE for Use and DE for Elim: the final VSub stands under its own newest eigenparameter of DE. In Lean the kernel form with the newest variable is `V0Prf`, with `v0prf_iff_nprf`

(`NativeNewestVariable`); the same proof applies, because the packet substitutes only for the witness it has just opened and no derivation is renamed. The clause form $\mathcal{V}_0$ is `VCPrf`, with `vcprf_iff_nprf` (`NativeVariableClauses`); there the premises needed under further variables are obtained from the induction hypothesis at a shifted renaming, which is the protected translation of Theorem 6.55.

6.5.3 Substitution for a parameter on raw terms

In this subsection terms are the raw terms of Section 1.1 and substitution is ordinary capture-avoiding substitution; $I(0, a;\, u.\, h)$ and a are different terms, and nothing is normalized. The rows of a packet are then not literal, and substitution for a variable has to be combined with the joining of two equations at the root. Formulas are equations and $D_{x,y}B$; contexts consist of clauses $[A;\, y.\, B(y)]$ with one antecedent; the assumption clause of F is $\mathrm{ord}(F) := [H;\, v.\, F]$.

For terms s, t, u define

$$\psi(s,t,u) := \begin{cases} u[t/s], & s \text{ is a parameter,} \\ t, & s \text{ is not a parameter and } u \equiv s, \\ u, & s \text{ is not a parameter and } u \not\equiv s. \end{cases}$$

The first case substitutes for a variable in one pass and is defined also when t contains s. For every s, $\psi(s,t,s) \equiv t$. For a source that is not a variable no proper subterm of u is inspected. Define also the partial one-step computation

$$\kappa(I(0,a;\, u.\, h)) := a, \qquad \kappa(I(Sn,a;\, u.\, h)) := h[I(n,a;\, u.\, h)/u],$$

undefined on all other terms; the iterator in the second result is left as it is.

Definition 6.57 (The calculus of parameter substitution)**.** $\mathcal{P}$ has the following five rule families.

$$\frac{}{\Gamma \vdash b = \kappa(b)}\ \textsc{Compute}, \qquad \kappa(b) \text{ defined} \tag{6.18}$$

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash u = r}{\Gamma \vdash r = \psi(s,t,u)}\ \textsc{Param} \tag{6.19}$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash B(t)}\ \textsc{Use}, \qquad [A;\, y.\, B(y)] \text{ an occurrence in } \Gamma$$

$$\frac{\Gamma \vdash D_{x,y}B(x,y) \qquad \Gamma,\ \mathcal{C}_B(b) \vdash F}{\Gamma \vdash F}\ \textsc{Open}$$

$$\frac{\Gamma,\ B(t,\eta) \vdash B(t,Sz)}{\Gamma \vdash D_{x,y}B(x,y)}\ \textsc{Close}, \qquad \eta = \begin{cases} z, & B \text{ is a literal equation and } t \equiv 0, \\ 0, & \text{otherwise} \end{cases}$$

Open is Elim and Close is Intro, on raw syntax and with the same freshness conditions; the tests of Close are tests on raw syntax. In this subsection Open denotes this rule and not the rule of Eq. (6.11).

There is no reflexivity axiom, no hypothesis rule, no template, no constructor Q, no case rule and no induction schema. Compute contains the two computation equations; counting it as one family is a grouping of the implementation and does not turn two equations into one.

Lemma 6.58 (Equality at the root)**.** *In $\mathcal{P}$ the following are derived under every context:*

$$s = t,\ s = r \vdash r = t; \qquad \vdash a = a; \qquad s = t \vdash t = s; \qquad s = t,\ t = r \vdash s = r.$$

Proof. The first, *joining at a common left side*, is PARAM with $u \equiv s$, since $\psi(s,t,s) \equiv t$. For reflexivity, COMPUTE proves $I(0,a;\,u.\,0) = a$, and joining this equation with itself gives $a = a$. For symmetry join the reflexive $s = s$, as major premise, with $s = t$. For transitivity take the symmetric $t = s$ and join $t = r$, as major premise, with $t = s$. Only COMPUTE and joining at the root are used. □

Lemma 6.59 (Elementary operations)**.** *The use of an assumption, weakening, injective renaming and cut of an assumption are admissible in $\mathcal{P}$.*

Proof. Reflexivity proves H, and USE of $[H;\,v.\,F]$ proves F. Weakening and injective renaming are inductions on derivations; an injective renaming preserves being a parameter, literal identity of terms, the root of a formula and the literal 0, hence the case of ψ and the mode of CLOSE. For cut, replace every USE of the occurrence $[H;\,v.\,F]$ by the given derivation of F, weakened to the local context, after renaming eigenparameters. No substitution of a term for a parameter in a derivation is used. □

Lemma 6.60 (Fresh aliases)**.** *For a term a and a variable b fresh for Γ, F and a,*

$$\frac{\Gamma,\ b = a \vdash F}{\Gamma \vdash F}$$

is admissible in $\mathcal{P}$. Its derivation uses the two computation equations, joining at the root, the logical rules and cut; it uses no substitution in a proper subterm.

Proof. Let $B(x,y) := (\mathrm{sel}(y,a,x) = a)$, where $\mathrm{sel}(y,a,x) = I(y,a;\,u.\,x)$. CLOSE at the witness a proves $D_{x,y}B$: its goal $\mathrm{sel}(Sz,a,a) = a$ is the successor case of COMPUTE for the constant body a. The assumption is not used, so the derivation is valid in either mode, also when $a \equiv 0$.

Open at a fresh b. The clause is $[\mathrm{sel}(0,a,b) = a;\ y.\ \mathrm{sel}(y,a,b) = a]$. Its antecedent is the zero case of COMPUTE, and USE at 1 gives $\mathrm{sel}(1,a,b) = a$. The successor case of COMPUTE gives $\mathrm{sel}(1,a,b) = b$. Joining, with the former as major premise, gives $b = a$. Cut this into the given derivation, weakened by the clause; OPEN discharges b. □

A second derivation uses the case mode of CLOSE only: take the witness $I(0,a;\,v.\,0)$, which is not the literal 0; the successor case of COMPUTE gives $\mathrm{sel}(Sz,a,I(0,a;\,v.\,0)) = I(0,a;\,v.\,0)$, the zero case gives $I(0,a;\,v.\,0) = a$, and transitivity joins them. So the lemma does not depend on the progress mode. The alias is the device of the optional argument in `template_free`; the point here is that it needs neither substitution in a proper subterm nor transport in a template nor induction, and therefore cannot hide the principle it is used to derive.

Theorem 6.61 (Substitution for a parameter gives every template)**.** *For every term template $h(v)$, if $\Gamma \vdash_{\mathcal{P}} s = t$ then $\Gamma \vdash_{\mathcal{P}} h(s) = h(t)$. Repeated occurrences of v, occurrences under the binder of an iterator and the absence of v are all permitted.*

Proof. Work under the assumption $b = s$, with b fresh for Γ, h, s and t. Transitivity gives $b = t$. Reflexivity gives $h(b) = h(b)$. PARAM with the major premise $b = s$ and this minor premise gives $h(b) = h(s)$: the source is the parameter b, and substitution for b marks exactly the distinguished occurrences of the template. In the same way $b = t$ gives $h(b) = h(t)$. Joining $h(b) = h(t)$, as major premise, with $h(b) = h(s)$ gives $h(s) = h(t)$, which does not contain b, and Theorem 6.60 discharges the alias. □

The argument is not a pass that replaces every occurrence of the possibly compound term s in $h(s)$; such a pass would also replace occurrences of s in the passive part of the template. Only the parameter case of ψ acts on a proper subterm.

Corollary 6.62 (The equality interface of $\mathcal{P}$)**.** *$\mathcal{P}$ derives* Tr*, $s = t,\ h(s) = r \vdash r = h(t)$, and transport in every equation template, $s = t,\ p(s) = q(s) \vdash p(t) = q(t)$. It derives the replacement rule of* `template_free`*: from $s = t$ and $u = v$ infer $v = u'$, where u' results from u by replacing, in one pass from the root, the occurrences of s by t.*

Proof. Join the congruence $h(s) = h(t)$, as major premise, with $h(s) = r$. Apply this to p and then to q, with symmetry, for the equation template. For the replacement rule, a pass from the root selects finitely many pairwise disjoint occurrences of s in u; replacing exactly these by a fresh variable gives a template h with $h(s) \equiv u$ and $h(t) \equiv u'$. The extraction of the template belongs to the producer of the proof and not to the checker. □

Lemma 6.63 (Every case introduction)**.** *For every matrix B and witness t, from $\Gamma, B(t, 0) \vdash B(t, Sz)$ one derives $\Gamma \vdash D_{x,y}B(x, y)$ in $\mathcal{P}$.*

Proof. Only a literal equation E at the witness 0 is not already the case mode of Close. Work under a fresh alias $b = 0$. From the assumption $E(b, 0)$, transport in an equation template gives $E(0, 0)$; cut in the given derivation to obtain $E(0, Sz)$; transport along the symmetric equation gives $E(b, Sz)$. Close at the variable b is in case mode and concludes $D_{x,y}E$, which does not contain b; Theorem 6.60 discharges the alias. Only transport in equations is used. □

Lemma 6.64 (Closures, induction, and transport in formulas)**.** *With $q_yB := D_{w,y}B$ for an unused w, the following are derived in $\mathcal{P}$.*

(a) *From $\Gamma, B(0) \vdash B(Sz)$ infer $\Gamma \vdash q_yB$.*

(b) *From $\Gamma \vdash q_yB$ and $\Gamma \vdash B(0)$ infer $\Gamma \vdash B(t)$.*

(c) *A context that contains $[B(0);\ y.\, B(y)]$ proves q_yB.*

(d) *For a literal equation E, from $\Gamma, E(z) \vdash E(Sz)$ infer $\Gamma \vdash q_yE$; with (b), induction for literal equations under an arbitrary context.*

(e) *Transport in every formula template.*

Proof. (a) Close at the unused witness 1 is in case mode. (b) Open for the unused witness adds $[B(0);\ y.\, B(y)]$, and Use gives $B(t)$. (c) In (a), under the assumption $B(0)$, use the clause at Sz. (d) Close at the unused literal witness 0 is in progress mode. (e) By induction on the template, as in Theorem 6.10: the equation case is Theorem 6.62; for $D_{x,y}K(v, x, y)$ open the source at b and introduce the target at the same b in case mode, converting the base backwards and the successor instance forwards. □

Theorem 6.65 ($\mathcal{P}$ and the kernel)**.** *Let τ be the translation of native formulas that leaves terms unchanged and satisfies $\tau(s = t) = (s = t)$, $\tau(D_{x,y}B) = D_{x,y}\tau B$ and $\tau(Q_yB) = D_{w,y}\tau B$ with w fresh, and let j be the inclusion of the formulas of $\mathcal{P}$ into native formulas. For every context Γ of native formulas and every native formula A,*

$$\Gamma \vdash_{\mathsf{N}} A \quad \Longleftrightarrow \quad \mathrm{ord}(\tau\Gamma) \vdash_{\mathcal{P}} \tau A.$$

Moreover $\tau j F \equiv F$ for every formula F of $\mathcal{P}$, and N *derives A from $j\tau A$ and conversely.*

Proof. τ commutes with substitution and preserves the property of being a literal equation.

From N *to $\mathcal{P}$*, by induction on the derivation. Hyp is the use of an assumption (Theorem 6.59). The two computation rules are the two cases of Compute. Tr is Theorem 6.62. QE is Theorem 6.64(b). DI is Theorem 6.63, and QI is Theorem 6.64(d). For DE, open the translated major premise at b; by Theorem 6.64(c) the opened clause proves $q_y\tau B(b, y)$, which is $\tau(Q_yB(b, y))$; weaken the translated branch and cut this formula into it; Open closes.

From $\mathcal{P}$ to N. Consider contexts each of whose occurrences is an assumption clause $\mathrm{ord}(G)$ or an opened clause $\mathcal{C}_B(b)$, with its origin recorded, and represent them by the native formulas jG and $Q_y jB(b,y)$ respectively. By induction on a derivation in $\mathcal{P}$ from such a context, N derives j of its conclusion from the representing formulas. A Use of an assumption clause is Hyp, and a Use of an opened clause is QE. Compute is a computation rule. Param is an instance of Tr: with the template $u[v/s]$, v fresh, if s is a parameter; with the identity template if s is not a parameter and $u \equiv s$; with the constant template u otherwise. Open is DE. A Close in case mode is DI. A Close in progress mode, at the literal witness 0 with the matrix E, is QI for $Q_y E(0,y)$, followed by QE at Sz under the assumption $E(0,0)$ and by DI.

Round trips. τj is the identity. For $j\tau$ the essential equivalence in N is $Q_y B(y) \dashv\vdash D_{w,y}B(y)$ for an unused w: from left to right use QE under the base and DI at any witness; from right to left DE opens exactly $Q_y B(y)$. The equivalence lifts through both binders by the congruence of binders in N (Section 1.4), and induction on A gives the interderivability of A and $j\tau A$. With cut in N this turns the second translation into the implication from right to left of the displayed equivalence. □

The proof of completeness uses the parameter case of ψ and joining at the root, never the third case, which only reverses the minor premise.

Corollary 6.66 (Bridge for $\mathcal{P}$)**.** *The ordinary reading of $\mathcal{P}$ is bi-interpretable and equiconsistent with* iPRA*, with whole domains, actual equality, and both numerical composites with graph $x = y$.*

Proof. Compose Theorem 6.65 with Theorem 1.78. Terms are unchanged, so the numerical maps are identities, and $0 = S0$ is fixed by τ and j. □

Remark 6.67 (Substitution, checking, Lean)*.* Substitution of a term for a parameter does not preserve the rules of $\mathcal{P}$: it can turn the source of Param from a variable into a compound term and can turn a witness of Close into the literal 0. Its admissibility follows from Theorem 6.65 and substitution in N (Section 1.4). The checker performs structural traversals, capture-avoiding substitutions, literal comparisons and one-step computations only, so its certificate relation is primitive recursive; proofs and substituted terms can still grow considerably, since the producer builds an alias around every transport in a template.

In Lean the kernel with Tr replaced by Param is `PPrf`, with `pprf_iff_nprf` (`NativeParameter`); the alias is discharged through $D_{x,y}(\mathrm{sel}(y,s,x) = s)$, proved by cases with the witness s and eliminated at $y = 1$ (`PPrf.aliasE`), and the calculus is closed under injective renaming (`PPrf.ren`). The five families of $\mathcal{P}$, which combine the rule with clause contexts and the introduction with a mode, are not formalized together.

6.5.4 Unary congruence does not suffice

VSub and Param are binary: they combine two equations. One may ask whether the equality rule can be the unary congruence. On raw and on zero-normal terms it cannot.

Proposition 6.68 (An obstruction)**.** *Let N^c be the calculus obtained from N by replacing Tr with reflexivity, symmetry and the rule*

$$\frac{\Gamma \vdash s = t}{\Gamma \vdash h(s) = h(t)}$$

for every term template h, all other rules being kept. Then N^c does not derive $x = y,\ y = z \vdash x = z$. The same holds on zero-normal terms, where $I(0,a;\,u.\,h)$ is identified with a.

Proof. Let $V = \{0, m, 2\}$, ordered $0 < m < 2$, and call two elements *near* if they are equal or adjacent; so 0 and 2 are not near, and m is near to every element. Interpret 0 as 0, successor as the constant function with value 2, and

$$I(n, a;\, u.\, h) \;\text{ as }\; \begin{cases} a, & n = 0, \\ m, & n \neq 0, \end{cases}$$

the body being ignored. An equation $s = t$ is *true* under an assignment of elements of V to the variables if the values of s and t are near. This is a truth assignment to judgments and not a structure with actual equality. $Q_y B$ and $D_{x,y} B$ are given the Boolean readings of Section 1.2 over V. A sequent is valid if every assignment that makes its context true makes its conclusion true. The value of a term commutes with substitution, since bodies are ignored, and the value of a term is unchanged by zero-normalization.

The rules other than the introductions. Hyp, reflexivity and symmetry are valid. The zero computation holds with equal values. In the successor computation the left side has the value m, which is near to every value. QE and DE are valid for the Boolean readings. For the congruence rule it suffices that the interpretation of the iterator preserves nearness in both arguments together: if both counters are 0 the outputs are the two seeds; if exactly one is 0 the outputs are a seed and m; otherwise both are m. Zero and successor are constants.

The shape of term functions. Every term denotes either the constant m, or a function that takes the value of a *payload*, which is 0, 2 or one variable, when all variables of a finite *gate* are 0, and the value m otherwise. This holds for variables, 0 and successors with an empty gate, and for $I(n, a;\, u.\, h)$ by cases on the counter: if n is constantly m or has the payload 2, the iterator is constantly m; if n has the payload 0 and the gate G, the iterator is a with G added to its gate; if n has a variable w as payload, the iterator is a with $G \cup \{w\}$ added to its gate. Consequently, as a function of one variable v with the others fixed, a term takes at $(0, m, 2)$ one of the triples

$$(0,0,0), \quad (m,m,m), \quad (2,2,2), \quad (0,m,2), \quad (0,m,m), \quad (2,m,m).$$

The key property. Every formula $B(v)$ satisfies, for every variable v and all values of the other variables,

$$B(0) \text{ and } B(2) \text{ imply } B(m). \tag{6.20}$$

For an equation: at $v = m$ each side has the value m or is constant in v. If one side has the value m there, the equation is true at m. If both are constant, its truth does not depend on v.

For $Q_y E$ with E an equation: the formula is false exactly if E is true at $y = 0$ and $y = m$ and false at $y = 2$. Falsity at 2 means that the two sides have the values 0 and 2 there, so neither side has y in its gate, all gate variables of both sides have the value 0, and each side is its payload. Inspecting the payloads, the formula is then false only if the payloads are y and 0, or y and a variable w whose value is 0. Hence $Q_y E$ is true, or false, or equivalent to the statement that some variable of a fixed finite set is different from 0. A formula of this last kind satisfies (6.20): if v belongs to the set it is true at $v = m$, and otherwise it does not depend on v. For $D_{x,y} E$, the reading is the existence of a value of x at which $Q_y E$ holds; if x is in the fixed finite set the formula is true, and otherwise it is the same statement as before. Finally, if B is true, false, or the statement that some variable of a finite set is different from 0, then $Q_y B$ is true: if y is not in the set the instances do not depend on y, and if it is, the instance at $y = 0$ implies all instances. So $D_{x,y} B$ is true as well, and so is every formula with more binders.

The introductions. For DI, take the value of the witness t for x. If $B(t, 0)$ is true, the premise gives $B(t, Sz)$, that is, B at $y = 2$, and (6.20) gives B at $y = m$. For QI, if $E(0)$ is true, the premise with z assigned 0 gives E at 2, and (6.20) gives E at m.

So every derivable sequent is valid. Under $x \mapsto 0$, $y \mapsto m$, $z \mapsto 2$ the equations $x = y$ and $y = z$ are true and $x = z$ is false. The contradiction $0 = S0$ is false as well. □

Full normalization of the successor computation is outside the obstruction: it identifies values that the interpretation only relates. The proposition excludes this one deletion in the neighborhood of the kernel; it does not exclude other presentations of arithmetic.

6.6 Specialization without a term

This section draws on `deterministic_specialization`. The calculus it restricts is the five-rule clause calculus of `uniform_step`.

6.6.1 The calculus

In this section terms are *zero-normal*: the normalization contracts $I(0, a;\, u.\, h)$ to a in all subterms and does not contract an iterator whose counter is a successor. Terms are the fixed points of this map, and every substitution is followed by it, including a substitution that makes a counter the literal 0; in particular $\mathrm{sel}(0, a, b) \equiv a$. Formulas are equations and $D_{x,y}F$, so that every formula is a chain of binders ending in one equation; its *rank* is the number of binders. A context is a finite list of clauses $[A;\, y.\, B(y)]$, written with the *newest* occurrence first, and the parameters of a sequent form a list whose last member is the newest. An assumption A is the clause $[0 = 0;\, v.\, A]$. The order is used only to choose the target of a specialization; clauses are reusable.

Definition 6.69 (Deterministic specialization)**.** The rules are

$$\frac{}{\Gamma \vdash t = t}\ \textsc{Refl} \qquad \frac{\Gamma \vdash I(c, a;\, u.\, h) = t \qquad \Gamma \vdash I(Sc, a;\, u.\, h) = r}{\Gamma \vdash r = h[t/u]}\ \textsc{Step} \tag{6.21}$$

$$\frac{\Gamma \vdash D_{x,y}B(x, y) \qquad [B(b, 0);\, y.\, B(b, y)],\ \Gamma \vdash F}{\Gamma \vdash F}\ \mathrm{DE}$$

$$\frac{[0 = 0;\, v.\, B(t, \eta)],\ \Gamma \vdash B(t, Sz)}{\Gamma \vdash D_{x,y}B(x, y)}\ \mathrm{DI}, \qquad \eta = \begin{cases} z, & B \text{ is a literal equation and } t \equiv 0, \\ 0, & \text{otherwise} \end{cases}$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash B(t)}\ \textsc{Use}, \qquad [A;\, y.\, B(y)] \text{ an occurrence in } \Gamma,\ t \text{ arbitrary}$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash B(v_i)}\ \textsc{Use}_*, \qquad [A;\, y.\, B(y)] = \Gamma_i, \quad v_i = \begin{cases} 1, & i = 0, \\ b_*, & i > 0. \end{cases} \tag{6.22}$$

In Step the first iterator is zero-normal, so it is a when $c \equiv 0$; repeated expressions are literally identical. In DE the variable b is fresh for Γ, F and $D_{x,y}B$, and in DI the variable z is fresh for Γ, t and $D_{x,y}B$; each becomes the newest parameter of its premise, and the displayed clause becomes the newest clause. In $\textsc{Use}_*$, Γ_0 is the newest clause and b_* is the newest parameter, or 0 if there is none. $\mathcal{U}$ is the calculus with Refl, Step, Use, DE and DI; $\mathcal{U}_*$ has $\textsc{Use}_*$ in place of Use.

So the newest clause is used at 1 and every older clause at the newest parameter. A certificate of $\textsc{Use}_*$ consists of the index of the clause and the proof of its antecedent; it contains no term and no mode. The witness of DI is still supplied, and the completeness proof uses that facility to name the desired term.

$\mathcal{U}$ is the calculus of `uniform_step` with clause contexts; its comparison with the kernel belongs to that package, and its arithmetic interface on the kernel is `NSPrf`, with `nsprf_iff_nprf`. The facts about it that are used here are the following.

Lemma 6.70 (Arithmetic from STEP)**.** *Using REFL and STEP only, and opening no scope, both $\mathcal{U}$ and $\mathcal{U}_*$ derive: the pivot $s = t,\ s = r \vdash r = t$; symmetry and transitivity; the congruence $s = t \vdash h(s) = h(t)$; transport in every equation template; and the successor computation $I(Sc, a;\ u.\, h) = h[I(c, a;\ u.\, h)/u]$, in particular* $\mathrm{sel}(Sz, a, b) = b$.

Proof. Let $J_s := I(1, s;\ u.\, u)$. Since $I(0, s;\ u.\, u) \equiv s$, STEP with $c \equiv 0$ sends $s = t$ and $J_s = r$ to $r = t$. Given $p\colon s = t$ and $q\colon s = r$, STEP applied to q and the reflexive $J_s = J_s$ gives $J_s = r$, and STEP applied to p and this gives $r = t$. Symmetry is the pivot with the first premise $s = s$, and transitivity follows. For a template h let $K := I(1, s;\ u.\, h)$; STEP applied to the reflexive $s = s$ and $K = K$ gives $K = h(s)$, and STEP applied to $s = t$ and $K = h(s)$ gives $h(s) = h(t)$. An equation template $p(v) = q(v)$ is transported through $p(t) = p(s)$, $p(s) = q(s)$ and $q(s) = q(t)$. The successor computation is STEP applied to the reflexivities of $I(c, a;\ u.\, h)$ and $I(Sc, a;\ u.\, h)$. □

Conversely the two computation equations and transport give STEP: in the computation equation for $I(Sc, a;\ u.\, h)$ replace $I(c, a;\ u.\, h)$ by t in the body and the successor iterator by r. So STEP adds no arithmetic strength.

Lemma 6.71 (The unrestricted calculus)**.** *In $\mathcal{U}$ cut of an assumption is admissible, and the following are derived: case introduction for every matrix and witness; for a literal equation E and every witness t, the progress introduction from $\Gamma, E(t, z) \vdash E(t, Sz)$ to $\Gamma \vdash D_{x,y}E$; and transport in every formula template.*

Proof. Cut is an induction on the derivation: a use of the assumption clause is replaced by the given proof. The term $c := I(1, 0;\ u.\, u)$ is zero-normal, is not the literal 0, and STEP proves $c = 0$. Given a premise $\Gamma, E(0, 0) \vdash E(0, Sz)$ for a literal equation E, start from $E(c, 0)$, transport to $E(0, 0)$, insert the given derivation, and transport back to $E(c, Sz)$; DI at c is in case mode. For progress at t, introduce $D_{w,y}E(t, y)$ with an unused w at the witness 0; DE adds $[E(t, 0);\ y.\, E(t, y)]$; under the base $E(t, 0)$ use it at Sz and apply the case introduction at t. Transport through a binder is as in Theorem 6.10. A progress introduction for a compound matrix is never used; it would change the induction of the theory. □

6.6.2 Naming a term

For a term t let

$$\mathrm{Nm}_t := D_{x,y}(\mathrm{sel}(y, 0, x) = \mathrm{sel}(y, 0, t)), \qquad \mathcal{M}_{b,t} := [0 = 0;\ y.\ \mathrm{sel}(y, 0, b) = \mathrm{sel}(y, 0, t)].$$

Lemma 6.72 (Naming)**.** *In every context, $\mathcal{U}_*$ derives* Nm_t. *Opening it by DE at a fresh b adds the clause $\mathcal{M}_{b,t}$, and in the context of that premise, where $\mathcal{M}_{b,t}$ is the newest clause, $\mathcal{U}_*$ derives $b = t$.*

Proof. Apply DI at the witness t. The goal $\mathrm{sel}(Sz, 0, t) = \mathrm{sel}(Sz, 0, t)$ has literally equal sides and is proved by REFL, in either mode. The antecedent of the opened clause is $\mathrm{sel}(0, 0, b) = \mathrm{sel}(0, 0, t)$, that is, $0 = 0$. Since the clause is newest, USE$_*$ applies it at 1, with REFL for its antecedent, and gives $\mathrm{sel}(1, 0, b) = \mathrm{sel}(1, 0, t)$; the computations $\mathrm{sel}(1, 0, b) = b$ and $\mathrm{sel}(1, 0, t) = t$ of Theorem 6.70 give $b = t$. □

This derivation of $b = t$ is valid immediately after the opening. It does not remain a derivation when the context is extended, because the clause then ceases to be newest. An extension of the scope changes b_*, and the insertion of a clause changes which clause is newest; so weakening is not an operation that preserves the rule at each node of a derivation of $\mathcal{U}_*$. The completeness proof replaces weakening by procedures that produce a proof again in every larger context.

6.6.3 Persistent proofs

A *situation* is a pair $(\Xi;\Gamma)$ of a parameter list and a context over it. An *extension* of $(\Xi;\Gamma)$ is a situation $(\Xi';\Gamma')$ together with an injective map ι of the members of Ξ to members of Ξ' and an injective map of the occurrences of Γ to occurrences of Γ' that sends an occurrence of a clause C to an occurrence of $C\iota$. Extensions compose, and neither map need preserve the order.

Definition 6.73 (Persistent proof)**.** A *persistent proof* of a formula A over $(\Xi;\Gamma)$ is a function that assigns to every extension of $(\Xi;\Gamma)$ a derivation in $\mathcal{U}_*$ of $\Gamma' \vdash A\iota$.

A persistent proof is a finite proof-producing procedure of the metatheory; it is not an additional rule.

Lemma 6.74 (Elementary persistence)**.** (a) *A persistent proof over a situation restricts to a persistent proof over each of its extensions.*

(b) *If* $[0 = 0;\, v.\, A]$ *occurs in* Γ*, then* A *has a persistent proof over* $(\Xi;\Gamma)$*.*

(c) *An equation* $t = t$ *has a persistent proof; if the two premises of a* Step *have persistent proofs, so has its conclusion. Hence the conclusions of the derivations of Theorem 6.70 have persistent proofs when their premises have.*

Proof. (a) Compose extensions. (b) In every extension apply $\textsc{Use}_*$ to the image of the occurrence, with Refl for the antecedent; the bound variable is unused, so either target gives $A\iota$. (c) Apply the rule in each extension; Refl and Step do not depend on the context or on the order of the parameters. □

Lemma 6.75 (Specialization of an equation clause)**.** *Let* $[A;\, y.\, E(y)]$ *occur in* Γ *with* E *a literal equation, let* A *have a persistent proof over* $(\Xi;\Gamma)$*, and let* t *be a term over* Ξ*. Then* $E(t)$ *has a persistent proof over* $(\Xi;\Gamma)$*. Consequently, if* $\mathcal{M}_{b,t}$ *occurs in* Γ*, at any position, then* $b = t$ *has a persistent proof over* $(\Xi;\Gamma)$*.*

Proof. Let an extension be given; write t, E, A for the renamed expressions. Derive Nm_t and open it by DE at a fresh b (Theorem 6.72). In the premise, b is the newest parameter and $\mathcal{M}_{b,t}$ the newest clause. The persistent proof of A, taken at this larger situation, gives A. The given clause is now an older clause, so $\textsc{Use}_*$ applies it at $b_* \equiv b$ and gives $E(b)$. Theorem 6.72 gives $b = t$, and transport in the equation template E (Theorem 6.70), which opens no scope, gives $E(t)$. This formula does not contain b, and DE closes. The construction is uniform in the extension, so it is a persistent proof.

For the second statement apply the first to the clause $\mathcal{M}_{b,t}$, the target 1 and the antecedent $0 = 0$, which has the persistent proof Refl; this gives $\mathrm{sel}(1, 0, b) = \mathrm{sel}(1, 0, t)$ persistently, and Theorem 6.74(c) with the selector computations gives $b = t$. The argument is not circular: the auxiliary naming clause of the first part is newest when it is used at 1, and the final transport there is between equations. □

Lemma 6.76 (Transport and specialization at every rank)**.** *For every* $r \geq 0$ *the following hold.*

(T_r) *Let* $F(w)$ *be a formula template of rank* r *and let* $s = t$ *have a persistent proof over* $(\Xi;\Gamma)$*. From a derivation in* $\mathcal{U}_*$ *of* $\Gamma \vdash F(s)$ *one obtains a derivation of* $\Gamma \vdash F(t)$*; and from a persistent proof of* $F(s)$ *over* $(\Xi;\Gamma)$ *a persistent proof of* $F(t)$*.*

(S_r) *Let* $[A;\, y.\, F(y)]$ *occur in* Γ *with* F *of rank* r*, let* A *have a persistent proof over* $(\Xi;\Gamma)$*, and let* t *be a term over* Ξ*. Then* $F(t)$ *has a persistent proof over* $(\Xi;\Gamma)$*.*

Proof. (T_0) is transport in an equation template (Theorem 6.70), with the persistent proof of $s = t$ taken at the identity extension; (S_0) is Theorem 6.75. In every (T_r) the second statement follows from the first, applied in each extension to the restricted persistent proofs.

(T_r) for $r > 0$, from (T_{r-1}) and (S_{r-1}). Let $F(w) = D_{x,y}G(w, x, y)$. Use the given derivation of $F(s)$ as the major premise of DE, with a fresh witness c; the context of the branch contains the clause $\mathcal{C} := [G(s, c, 0);\, y.\, G(s, c, y)]$. In the branch introduce $F(t)$ by DI at the witness c. As c is a parameter it is not the literal 0, so this is the case mode also when G is an equation, and the premise has the assumption clause of $G(t, c, 0)$ and the goal $G(t, c, Sz)$. In the situation of that premise:

- $G(t, c, 0)$ has a persistent proof by Theorem 6.74(b);
- $t = s$ has a persistent proof by Theorem 6.74(a),(c), so (T_{r-1}) for the template $G(w, c, 0)$ gives a persistent proof of $G(s, c, 0)$;
- (S_{r-1}) for the clause $\mathcal{C}$, this antecedent and the target Sz gives a persistent proof of $G(s, c, Sz)$;
- (T_{r-1}) for the template $G(w, c, Sz)$, applied to the derivation that this persistent proof yields at the identity extension, gives a derivation of $G(t, c, Sz)$.

DI and DE close; $F(t)$ does not contain c. The given derivation of $F(s)$ is used outside the new scopes and is not weakened.

(S_r) for $r > 0$, from (T_r) and Theorem 6.75. In a given extension derive Nm_t and open it at a fresh b. In the premise, the persistent proof of A gives A, and $\textsc{Use}_*$ applies the given clause, now older, at b, which gives a derivation of $F(b)$. By Theorem 6.75, $b = t$ has a persistent proof over the situation of the premise; it is needed in persistent form because (T_r) opens further scopes. (T_r) gives $F(t)$, and DE closes. The construction is uniform in the extension.

The recursion is well founded: (T_r) uses rank $r - 1$ only, and (S_r) uses (T_r) and rank 0. □

6.6.4 Equivalence

Theorem 6.77 (Deterministic specialization)**.** *For every parameter list, every context* Γ *and every formula* A*,*

$$\Gamma \vdash_{\mathcal{U}} A \quad \Longleftrightarrow \quad \Gamma \vdash_{\mathcal{U}_*} A.$$

The terms, the formulas and the ordinary reading are unchanged.

Proof. Every instance of $\textsc{Use}_*$ is an instance of Use. Conversely, by induction on a derivation in $\mathcal{U}$ of $\Gamma \vdash A$ over Ξ, the formula A has a persistent proof over $(\Xi; \Gamma)$.

Refl and Step: Theorem 6.74(c). A Use of $[A';\, y.\, B]$ at t: the induction hypothesis gives a persistent proof of A', and (S_r) of Theorem 6.76, with r the rank of B, gives a persistent proof of $B(t)$. DE with the major premise $D_{x,y}B$ and the branch over $(\Xi, b;\, \mathcal{C}_B(b), \Gamma)$: in a given extension choose a fresh b', take the derivation of the major premise that its persistent proof yields there, extend the extension by $b \mapsto b'$ and by sending the opened clause to the new opened clause, take the derivation that the persistent proof of the branch yields at this extension, and apply DE. DI is treated in the same way; an injective renaming preserves the root of the matrix and the literal test of the witness, hence the mode.

The compilation recurses on the height of the given derivation, and the auxiliary procedures recurse on the rank of formulas; there is no recursion on a derivation that the procedure has produced. So every derivation in $\mathcal{U}$ yields a finite derivation in $\mathcal{U}_*$: evaluate the persistent proof at the identity extension. □

Composed with the comparison of `uniform_step` between $\mathcal{U}$ and the kernel and with Theorem 1.78, the theorem gives the bi-interpretation of the ordinary reading of $\mathcal{U}_*$ with iPRA, with whole domains, actual equality and identity numerical composites; the equivalence itself changes no term, no formula and no reading.

Remark 6.78 (The cost). The argument of an arbitrary specialization is removed from the checker at the price of larger proofs: the translation wraps every use of a clause at a term in a naming formula, and at rank r it nests the constructions of ranks below r. In the test suite of the package the largest expansion is from 699 to 175,640 bytes of compact JSON. No polynomial bound on the size of the translation is claimed.

Remark 6.79 (The kernel counterpart in Lean). On the kernel the corresponding restriction concerns the term of QE. With QE allowed at a variable and at the numeral 1 only, the kernel derives the same sequents: this is `VQPrf` with `vqprf_iff_nprf` (`NativeVariableQE`). An arbitrary target t is named by a fresh alias $b = t$, whose discharge uses QE at 1 once; QE is applied at b, and $B(b)$ is moved to $B(t)$ by the Leibniz rule, which is proved together with QE at Sz by induction on the formula (`VQPrf.qELeib`, `VQPrf.leibniz`). The numeral cannot be dropped (`vqprf_var_incomplete`): with QE at variables only, the kernel does not derive $0 = 1$ from $Q_y(\mathrm{sel}(y, 0, 1) = 0)$, which N derives by QE at 1. The countermodel reads every variable as 0, every equation by its value in the natural numbers, and every formula with root Q or $D_,$ as true; each rule preserves truth, since $B(v)$ at a variable v has the truth value of $B(0)$. The form of the package, on zero-normal terms with clause contexts and the choice of the target by position, is not formalized; that choice is not stable under renaming, which is why the proof above regenerates the proofs of premises.

6.7 A constancy rule in place of induction

This section draws on an appendix of `substitutive_discharge`. It is independent of the discharge rule: it presents an arithmetic whose one arithmetical inference has no induction hypothesis and no step function. The price is a primitive Boolean equality.

6.7.1 The theory

Definition 6.80 (The constancy theory)**.** The logic is intuitionistic first-order logic with equality over one sort. Terms are

$$t ::= x \mid 0 \mid S(t) \mid \delta(t, t) \mid R(n, a;\, i, u.\, h),$$

where the recursor binds i and u in h only. With $1 := S0$ and $s \neq t := (s = t \to \bot)$, the axioms are the universal closures of

$$R(0, a;\, i, u.\, h) = a, \tag{6.23}$$

$$R(Sn, a;\, i, u.\, h) = h[n/i,\ R(n, a;\, i, u.\, h)/u], \tag{6.24}$$

$$0 = 1 \to \bot, \tag{6.25}$$

$$\big(s = t \wedge \delta(s, t) = 0\big) \vee \big(s \neq t \wedge \delta(s, t) = 1\big), \tag{6.26}$$

the substitution in (6.24) being simultaneous, and the one further rule is

$$\frac{\Gamma \vdash f(Sz) = f(z)}{\Gamma \vdash f(t) = f(0)}\ \textsc{Const} \tag{6.27}$$

for a term template $f(x)$, with z fresh for Γ, t and f. The context Γ consists of arbitrary first-order formulas. There is no coverage axiom, no injectivity axiom for successor, no induction schema and no assumed reflection of equality.

6.7.2 Recovery of quantifier-free induction

Define

$$\mathrm{sel}(v,a,b) := R(v,a;\, i,u.\, b), \qquad a+b := R(b,a;\, i,u.\, Su), \qquad pd(v) := R(v,0;\, i,u.\, i).$$

Lemma 6.81 (Elementary computations)**.** *Without* Const*:* $\mathrm{sel}(0,a,b) = a$ *and* $\mathrm{sel}(Sv,a,b) = b$*;* $a+0=a$*,* $a+Sb = S(a+b)$ *and* $a+1=Sa$*;* $pd(0)=0$ *and* $pd(Sv)=v$*; successor is injective;* $Sv \neq 0$*; and*

$$\delta(s,t) = 0 \vee \delta(s,t) = 1, \qquad \delta(s,t) = 0 \leftrightarrow s = t. \tag{6.28}$$

Proof. The equations are instances of (6.23) and (6.24), and injectivity follows from $pd(Sv) = v$. If $Sv = 0$, applying $w \mapsto \mathrm{sel}(w,0,1)$ gives $1 = 0$, against (6.25); no distinction of cases on a number is used. The disjunction in (6.28) is read off (6.26). If $s = t$, the second disjunct of (6.26) contradicts it, so $\delta(s,t) = 0$. If $\delta(s,t) = 0$, the second disjunct would give $0 = 1$, so $s = t$. □

For a quantifier-free formula A define a term q_A, with the value 0 meaning that A holds:

$$\begin{aligned} q_{s=t} &:= \delta(s,t), & q_\bot &:= 1,\\ q_{A\wedge B} &:= \mathrm{sel}(q_A, q_B, 1), & q_{A\vee B} &:= \mathrm{sel}(q_A, 0, q_B),\\ q_{A\to B} &:= \mathrm{sel}(q_A, q_B, 0). \end{aligned}$$

Lemma 6.82 (Boolean terms)**.** *For every quantifier-free* A*, without* Const*,*

$$q_A = 0 \vee q_A = 1, \qquad q_A = 0 \leftrightarrow A.$$

Proof. By induction on A in the metatheory, with the two computations of sel and (6.28). In the case of an implication distinguish the two values of q_A: if it is 0, then A holds and $A \to B$ is equivalent to B; if it is 1, then $\neg A$ follows and the implication holds. Only the decidability already proved for the quantifier-free antecedent is used. □

Theorem 6.83 (Quantifier-free induction from constancy)**.** *The theory of Theorem 6.80 derives*

$$\frac{\Gamma \vdash A(0) \qquad \Gamma,\ A(z) \vdash A(Sz)}{\Gamma \vdash A(t)}$$

for every quantifier-free A *and every context* Γ *of first-order formulas,* z *fresh. In particular it derives induction for literal equations under an arbitrary context.*

Proof. Write $q(n) := q_{A(n)}$ and define the terms

$$e(n) := \mathrm{sel}(q(n), 0, \mathrm{sel}(q(Sn), 1, 0)), \qquad a(n) := R(n,0;\, i,u.\, u{+}e(i)), \qquad F(n) := a(n){+}q(n).$$

These are terms of the language: e and the addition are expanded before the outer recursor is formed. The computations give $a(0) = 0$ and $a(Sn) = a(n) + e(n)$. By Theorem 6.82 the pair $(q(n), q(Sn))$ has four possible values. Three of them give $F(Sn) = F(n)$ with no induction hypothesis:

$q(n)$	$q(Sn)$	$e(n)$	$F(Sn) = F(n)$
0	0	0	$a(n) = a(n)$
1	1	0	$Sa(n) = Sa(n)$
1	0	1	$Sa(n) = Sa(n)$

The fourth, $(0,1)$, is impossible under Γ: $q(n) = 0$ gives $A(n)$, the step premise gives $A(Sn)$, and Theorem 6.82 gives $q(Sn) = 0$, against $0 \neq 1$. By elimination of the finitely many disjunctions, $\Gamma \vdash F(Sz) = F(z)$. Const gives $F(t) = F(0)$. The base $A(0)$ gives $q(0) = 0$, so $F(0) = 0$. If $q(t) = 1$, then $0 = F(t) = a(t) + 1 = Sa(t)$, against $Sv \neq 0$. Hence $q(t) = 0$, and $A(t)$ holds. □

The only identities of addition used are $a + 0 = a$ and $a + 1 = Sa$; commutativity and associativity are not used, so no induction is hidden in an algebraic rearrangement.

6.7.3 Both interpretations

Theorem 6.84 (The constancy theory and iPRA)**.** *The theory of Theorem 6.80 is bi-interpretable with* iPRA*, with whole domains, actual equality and both numerical composites provably the identity. In particular the two are equiconsistent.*

Proof. iPRA *in the constancy theory.* Theorem 6.83 applied to the quantifier-free formula $n = 0 \vee n = S(pd(n))$ gives coverage: the case 0 is immediate, and at a successor $pd(Sn) = n$ gives the second disjunct without the hypothesis. Injectivity of successor is in Theorem 6.81. Every primitive recursive definition compiles into the recursor with a counter: zero, successor, projections and composition are term operations, and $f(0, \vec{x}) = g(\vec{x})$, $f(Sn, \vec{x}) = h(n, f(n, \vec{x}), \vec{x})$ is represented by $R(n, g(\vec{x}); i, u.\, h(i, u, \vec{x}))$ after g and h are compiled; the defining equations are instances of (6.23) and (6.24). With separation, injectivity and Theorem 6.83 this interprets iPRA.

The constancy theory in iPRA. Each fixed recursor term denotes a primitive recursive function, and its two equations hold by construction. Const is admissible: under its premise, induction for the equation $f(n) = f(0)$ gives the conclusion. For δ an equality characteristic is constructed as follows. Let

$$d(a, b) := R(b, a;\, i, u.\, pd(u)).$$

The computation equations and induction for equations give

$$d(a, 0) = a, \quad d(a, Sb) = pd(d(a, b)), \quad d(Sa, Sb) = d(a, b), \quad d(Sn, n) = 1;$$

the third by induction on b at fixed a, with the base $pd(Sa) = a$ and the step obtained by applying pd to the preceding equation, the fourth by induction on n from the third. Inductions on the second argument of addition give $0 + b = b$ and $Sa + b = S(a + b)$. Quantifier-free induction on b, with a as a parameter, proves

$$d(a, b) \neq 0 \;\rightarrow\; a = b + d(a, b). \tag{6.29}$$

The base uses $0 + a = a$. For the step put $r := d(a, b)$ and suppose $pd(r) = d(a, Sb) \neq 0$. Coverage for r gives $r = S(pd(r))$, the case $r = 0$ contradicting $pd(r) \neq 0$. So $r \neq 0$, and the induction hypothesis gives $a = b + r = b + S(pd(r)) = Sb + pd(r)$. Now define

$$Z(a) := \mathrm{sel}(a, 1, 0), \qquad U(a) := \mathrm{sel}\big(a, 0, Z(pd(a))\big), \qquad E(a, b) := \mathrm{sel}\big(b, Z(a), U(d(a, pd(b)))\big).$$

Coverage and separation give that $Z(a)$ and $U(a)$ are 0 or 1, that $Z(a) = 1$ exactly if $a = 0$, and that $U(a) = 1$ exactly if $a = 1$. If $b = 0$, then $E(a, b) = 1$ is equivalent to $a = 0$. If $b = Sn$, then $E(a, b) = 1$ is equivalent to $d(a, n) = 1$, which implies $a = n + 1 = Sn = b$ by (6.29), and conversely $a = Sn$ gives $d(Sn, n) = 1$. So $E(a, b)$ is 0 or 1 and equals 1 exactly if $a = b$, and

$$\delta_*(a, b) := Z(E(a, b))$$

is 0 or 1 and equals 0 exactly if $a = b$. It satisfies (6.26). This derivation uses quantifier-free inductions of iPRA; it is not part of the bootstrap of Theorem 6.83.

Composites. In the constancy theory, induction for equations (Theorem 6.83) proves the uniqueness of every fixed primitive recursive definition under arbitrary parameters; by induction on terms the round trip of every recursor term has the value of the original. The same argument in iPRA treats the round trip of each primitive recursive function symbol. For

the primitive δ, both δ and the translated δ_* satisfy (6.26) in the constancy theory, and the distinction of the two cases of the first shows $\delta(a,b) = \delta_*(a,b)$: both are 0 if $a = b$ and both are 1 otherwise. Congruence and induction on formulas give both formula round trips. The numerical isomorphisms have graph $x = y$. □

Remark 6.85 (What the rule does and does not simplify). The arithmetical inference concerns one term and one unconditional recurrence with the identity as step; it has no discharged hypothesis and no step template, and the accumulator $a(n)$ is a construction of the proof and not of the rule. Against the kernel the presentation adds a binary term constructor, a primitive equality split in place of coverage, and a second bound variable of the recursor. The split (6.26) is essential to the proof above: replacing it by the assertion that an internally defined equality test is faithful, without proving that assertion in the induction-free base, would be circular. Likewise, for a tail-recursive iterator the equation $d(a, Sb) = d(pd(a), b)$ does not by itself give $d(a, Sb) = pd(d(a, b))$; the missing commutation is a fusion property. No checker is supplied for this theory, and nothing is claimed about its size.

6.8 Checkers and program equivalences

This section summarizes the programs of `witness_discharge`, `inferred_witness`, `deterministic_opening`, `first_instance`, `clause_discharge`, `substitutive_discharge`, `variable_substitution`, `parameter_substitution` and `deterministic_specialization`. Sizes are in Section 6.9.

Common contract. Every checker is a standalone Python module that reads one JSON value, verifies a closed theorem, prints `ACCEPT` and exits with status zero on success, and rejects on any exception; no decision depends on an `assert`. The checkers of `variable_substitution` and `deterministic_specialization` code syntax by tagged lists. The others use integers and bare lists: 0 is zero, a negative integer is a free parameter level, a positive integer is a de Bruijn index, `[0,t]` is a successor, `[n,a,h]` an iterator or recursor binding one variable in its last entry, `[s,t]` an equation and `[B]` a paired formula in whose matrix index 1 is y and index 2 is x; the expected sort distinguishes a successor from an equation. Subterms are validated before a contraction can erase them. Inserted terms have no loose positive index, so insertion below a binder needs no lifting. The theorems of this chapter concern open sequents; the public interfaces check closed goals.

`witness_discharge`. A goal-directed checker of $\mathcal{W}^{\mathcal{R}}$ with three packets, for Use (clause index, term, premise), Intro (witness, branch) and Name (matrix, local formula, target, major premise, equality premise, branch). The exported formula is compared with the already validated outer goal, which enforces the absence of the eigenparameter also below binders and after normalization has erased an occurrence.

`inferred_witness`. A forward checker of $\mathcal{O}^{\mathcal{R}}$ that infers every conclusion once. It stores the antecedent $B(b, 0)$ of the opened clause as computed and substitutes the selected witness in the body only: if the test succeeds the antecedent is H and is unchanged by the substitution, and if it fails the substitution is the identity, so the stored clause is $\mathcal{C}_B(\rho_B(b))$ in both cases.

`deterministic_opening`. The checker of `witness_discharge` with the packet of the third rule shortened to matrix, major premise and branch. On validated syntax the test of the row at 0 is that both entries are zero, and the test that the first entry of the row at 1 is the integer of b also establishes that the row is an equation, a paired formula having a list as its only entry.

`first_instance`. A checker of $\mathcal{X}^{\mathcal{R}}$ in which every proof node is a triple; the depth of the clause context equals the number of parameters in scope.

`clause_discharge`. Two goal-directed checkers on the input `[goal, proof]`, one of $\mathcal{W}^{\mathcal{R}}$ and one of $\mathcal{X}^{\mathcal{R}}$. The next fresh parameter is computed from the length of the context, and one traversal performs shifting and opening. The replacement acts on a parameter leaf exactly if

it is the left side of the row and is the fresh witness. A variant that infers conclusions was tried and was not smaller.

substitutive_discharge. A checker of $\mathcal{K}$ with four instructions and no fuel (Theorem 6.48), its matched control with separate rules, and the raw companion with its control.

variable_substitution. A goal-directed native checker of the six-rule form with the newest variable, obtained from the checker of `full_normalization` by changing the equality packet, and a forward checker of $\mathcal{V}_0$ whose equality packet consists of two premise proofs. In the native checker the leaves for hypothesis and reflexivity ignore an unused hint, as in its control.

parameter_substitution. A forward checker of $\mathcal{P}$ with five instructions. The certificate of $0 = 0$ joins two copies of the computation $I(0, 0; u.\, u) = 0$. Substitution in a proper subterm acts on negative integers only, so it cannot rewrite the zero that marks a successor.

deterministic_specialization. A goal-directed checker of $\mathcal{U}_*$ with one instruction for each of the five rules. The instruction for a clause contains the index and the proof of the antecedent; the target is computed from the index and the scope, and contexts are stored newest first. A translator in both directions is supplied and is not part of the checker.

Three statements about programs, independent of the rules, are proved in the packages.

Proposition 6.86 (Scope from the context)**.** *In a checker whose public entry starts from the empty context and scope* 0*, and in which the branch of every scope-extending rule adds exactly one clause and one parameter while no other call changes either, every reachable call satisfies* $scope = |context|$*. The scope argument can therefore be computed from the context.*

Proof. By induction on the tree of calls: the initial call satisfies the equation, the rules without a branch recur with both unchanged, and a branch increments both. □

`deterministic_specialization` applies this to the checker of $\mathcal{U}_*$ and to its control, and `clause_discharge` and `first_instance` build it into their checkers. The statement concerns the public entry; it is not claimed for a direct call of an internal function with a context and a scope that do not match.

Fusion of the rewrite with the continuation. `deterministic_opening` reorders, in three inherited compact checkers of an earlier guarded calculus, the rewriting of the goal by an elimination and the construction of the new hypothesis: the hypothesis is constructed first and the rewrite is an argument of the continuation call, evaluated before the scope is decremented by the left-to-right evaluation order of arguments. The two orders accept the same certificates. A rewrite changes the goal only if the major premise is a nonempty list, and then the construction of the hypothesis cannot perform the assignment of an introduction; conversely an introduction changes the goal only if the major premise is empty, and then the rewrite is the identity; so the two operations commute. The proof of the major premise runs in its own activation and does not affect the scope of the caller, and the rewrite consumes no proof packet. By induction on the number of packets each new program accepts exactly the certificates of its control and consumes the same suffix. These programs do not implement Open.

One traversal for grammar and scope. The supplement of `deterministic_specialization` modifies the validated compact checker of `scalar_uniqueness`, which checks the unchanged kernel. The control traverses the public goal twice, once for sorts and arities and once for scope; the new program does both in the first traversal, under a flag that is set for the public entry only and cleared before the first proof packet is read. By induction on the goal the fused traversal succeeds exactly if both old traversals succeed and then returns the goal; with the flag cleared every added guard is skipped and the remaining computation is that of the control.

6.9 Checker sizes

This section lists the sizes reported by the nine packages named at the beginning of Section 6.8. All figures follow the protocol of Section 1.10 and are those stated by the packages. A control is matched if it shares the term and formula coding, the normalizer, the context representation and the public contract with the program and differs in the rule under comparison; where a package compares with an unmatched program this is said.

Table 6.1: Complete checkers of the packages of this chapter.

Program	Tokens	AST	Bytes	Control (tokens / AST / bytes)
`witness_discharge`				
$\mathcal{W}^{\mathcal{R}}$	732	813	2,273	separate elimination and transport: 769 / 841 / 2,362
$\mathcal{W}^{\mathcal{R}}$ with fuel	777	868	2,486	the same with fuel: 814 / 896 / 2,575
`inferred_witness`				
$\mathcal{O}^{\mathcal{R}}$, forward	683	771	1,990	explicit equation at opening: 676 / 765 / 1,998; explicit equation at output: 663 / 751 / 1,983
$\mathcal{O}^{\mathcal{R}}$ with fuel	728	826	2,203	unmatched: `witness_discharge` with fuel, 777 / 868 / 2,486
`deterministic_opening`				
$\mathcal{O}^{\mathcal{R}}$, goal-directed	730	809	2,253	checker of `witness_discharge`: 732 / 813 / 2,273
intermediate export variant	745	828	2,296	the same control
compact profile, fused (inherited calculus)	267	302	508	265 / 306 / 516
AST profile, fused (inherited calculus)	263	276	580	260 / 278 / 602
validated native profile, fused (inherited calculus)	360	409	780	358 / 413 / 788
`first_instance`				
$\mathcal{X}^{\mathcal{R}}$, triples	704	804	2,157	$\mathcal{W}^{\mathcal{R}}$, triples: 701 / 813 / 2,228
$\mathcal{X}^{\mathcal{R}}$ with fuel	752	863	2,340	$\mathcal{W}^{\mathcal{R}}$ with fuel: 749 / 872 / 2,411
`clause_discharge`				
$\mathcal{W}^{\mathcal{R}}$ (explicit discharge)	636	708	2,010	checker of `witness_discharge`: 732 / 813 / 2,273
$\mathcal{X}^{\mathcal{R}}$ (deterministic discharge)	653	729	2,058	explicit discharge: 636 / 708 / 2,010
explicit discharge with fuel	673	755	2,139	
deterministic discharge with fuel	690	776	2,187	explicit discharge with fuel: 673 / 755 / 2,139
`substitutive_discharge`				
$\mathcal{K}$	713	798	2,267	$\mathcal{K}_{\mathrm{sep}}$: 726 / 809 / 2,318
raw companion	744	828	2,550	separate rules: 756 / 837 / 2,606
`variable_substitution`				
native six rules, newest variable	899	951	2,499	checker of `full_normalization` with Tr: 908 / 965 / 2,529

Program	Tokens	AST	Bytes	Control (tokens / AST / bytes)
$\mathcal{V}_0$	898	968	2,826	clausal Tr: 907 / 982 / 2,856
`parameter_substitution`				
$\mathcal{P}$	769	892	2,857	template rule, unused code removed: 765 / 881 / 2,907; template rule, shared opener: 777 / 895 / 2,937
`deterministic_specialization`				
$\mathcal{U}_*$, scope from context	819	896	2,914	$\mathcal{U}$, scope from context: 819 / 895 / 2,926
$\mathcal{U}_*$, explicit scope	829	903	2,938	$\mathcal{U}$, explicit scope (checker of `uniform_step`, newest first): 829 / 902 / 2,950
kernel, fused grammar and scope traversal	360	412	820	checker of `scalar_uniqueness`: 381 / 439 / 866

The comparisons are local. In its matched pair, Name against separate rules is smaller in all three coordinates; Open against an explicit equation at the opening saves 8 bytes and costs 7 tokens and 6 nodes, and is 7 bytes larger than the control with an explicit equation at the output; Export saves 71 bytes and 9 nodes and costs 3 tokens in `first_instance`, and costs 48 bytes, 17 tokens and 21 nodes in `clause_discharge`; SDis is smaller in all three; VSub for the newest variable saves 30 bytes, 9 tokens and 14 nodes in both profiles; Param saves 50 bytes and costs 4 tokens and 11 nodes; Use$_*$ saves 12 bytes, leaves the tokens unchanged and costs one node, and the computation of the scope from the context saves a further 24 bytes, 10 tokens and 7 nodes. `parameter_substitution` also supplies, as a comparison source and not as a new calculus, a refined checker of the two-case evaluation calculus with 741 tokens, 862 nodes and 2,731 bytes. Every package states that shorter checkers of the repository with other certificate contracts, in particular the compact checker of 516 bytes and, where it is named, the validated checker of 788 bytes, remain smaller than its programs, and none claims a minimum.

6.10 Further questions

This section collects the questions, limits and unproved claims stated by the nine packages named at the beginning of Section 6.8, and one conjecture recorded in the repository.

Question 6.87 (Substitution for a variable on raw terms)**.** On full normal forms substitution for a variable is the only equality rule needed (Theorem 6.51). On raw terms the rows of a packet are not literal, and Param, which also joins two equations at the root, suffices (Theorem 6.65). Is substitution for a variable alone, without joining at the root, strictly weaker on raw terms? The repository records the affirmative answer as a conjecture.

Question 6.88 (The least equality interface on raw terms)**.** The instances of Param that the completeness proof uses are joining at the root and substitution at a fresh variable; the third case of ψ is a reversal. Is there a smaller total operation that expresses exactly the needed instances without a larger test on syntax? An answer has to account for arbitrary equations between closed terms as assumptions, and not only for the computation equations.

Question 6.89 (Unary congruence on full normal forms)**.** Theorem 6.68 covers raw and zero-normal terms. Full normalization of the successor computation identifies values that its

interpretation only relates. Whether unary congruence suffices on full normal forms is not decided.

Question 6.90 (The source of the selected equation)**.** In OPEN and EXPORT the source of the selected equation is the witness opened by the same inference. A generalization in which the row at 1 may name an older parameter was considered in `clause_discharge` and discarded; the derivation of ordinary elimination by one EXPORT fails for it. Whether a sound and complete rule of that kind exists is open.

Question 6.91 (The normalization that naming needs)**.** The packets use the two computations of sel and the two closed values $P(1)$ and $P(2)$. Is there a smaller discipline of normalization, coherent with substitution and with nested binders, that makes exactly these computations literal and keeps the theorems of Sections 6.2, 6.3 and 6.5.1? Can a syntax expose the rows at 0 and 1 of a matrix without full normalization and keep both formula round trips?

Question 6.92 (A term template in the substitutive discharge)**.** Can the formula template of SDIS be restricted to a term template, with transport in compound formulas recovered, while the complete checker stays smaller than that of $\mathcal{K}_{\mathrm{sep}}$? The occurrences of the witness that the rule permits have to be tracked.

Question 6.93 (An introduction that is stable under substitution)**.** The mode of the single introduction depends on the literal spelling of the witness, and this is the reason why substitution does not preserve the rules of the calculi of this chapter. Is there a test that does not depend on the supplied witness, permits the recovery of the case introduction, and does not cost more in the checker than it saves?

Question 6.94 (The equality split)**.** Can the axiom (6.26) of Theorem 6.80 be derived from a small recursion base and CONST alone? The split must not be assumed during such a bootstrap.

Question 6.95 (Size of the translations)**.** Theorem 6.77 produces a finite derivation with no polynomial bound (Theorem 6.78). The translations of Theorems 6.50 and 6.61 add one packet or one alias per transport. What are the least expansions, for certificates as trees and with explicit sharing?

Question 6.96 (Lower bounds)**.** A lower bound on the size of a checker needs a fixed public grammar, a fixed contract for malformed input, a fixed source language and a fixed measure. No package of this chapter fixes these, and none proves a lower bound. Is there one complete program that improves the token, node and byte coordinates of the repository simultaneously under one contract? Can the rules of this chapter be represented in the smallest certificate languages based on arities without losing their savings to normalization and to the bookkeeping of scope?

The following are stated by the packages as not achieved or not claimed.

- *Formalization.* No package contains a Lean or Rocq proof of its theorems. The repository proves the kernel forms and the clause forms named in Theorems 6.21, 6.48, 6.67 and 6.79 and after Theorems 6.36 and 6.56. Not formalized are: the four families of $\mathcal{K}$ on the canonical recursor syntax, the five families of $\mathcal{P}$ together, the positional calculus $\mathcal{U}_*$, and the constancy theory of Section 6.7.

- *Effectiveness.* The proof translations of this chapter are effective; none is claimed to be primitive recursive. No primitive recursive bound on normalization, and no bound on the fuel needed by a derivable formula, is asserted (Theorem 6.2). `clause_discharge` does not claim an arithmetized equivalence of consistency statements in a specified weak metatheory.

- *Executable translations.* The tests of `deterministic_opening` supply, for each tested theorem, a matching proof in $\mathcal{W}^{\mathcal{R}}$; they do not contain a general executable compiler from $\mathcal{O}^{\mathcal{R}}$, nor certified compilers between the calculi on normal forms and the raw kernel with normalization traces and the repair of the introduction mode.
- *The equality fragment.* The proofs of Theorems 6.17 and 6.50 use the logical rules; no completeness statement is made for the equality rules alone.
- *Size.* No package claims a smallest presentation or a smallest checker, and the matched comparisons of Section 6.9 are not improvements in every coordinate.
- *Arithmetic.* The restrictions of this chapter change the equality interface and the use of clauses. They do not weaken the arithmetic: the construction of the predecessor and the reflection of equality (Theorems 1.48 and 1.52) remain what a proposal with weaker induction would have to redo, as `parameter_substitution` notes.

Chapter 7

Introduction without a supplied witness

The rule DI of the native scalar core takes a term: its witness. This chapter restricts that term. The witness is first required to be a variable or the literal 0, then to be the newest numerical parameter of the current scope, with 0 in an empty scope, so that the rule takes no term at all. Every restricted calculus of the chapter derives the same sequents as its unrestricted counterpart, on all open sequents and with the identity translation of terms and formulas. The restriction of the witness cannot be combined with a restriction of the targets of QE (Theorem 7.7).

The chapter is organized by idea. Section 7.1 states the restricted calculi and what they lose: a calculus with variable witnesses is closed under renaming and not under substitution, and a calculus with a scope-determined witness is not closed under the extension of its scope. Section 7.2 contains the negative results. Section 7.3 reduces the completeness of a restricted calculus to the derivability of a name for every term, and Section 7.4 derives the names from a formula proved by numerical cases. Section 7.5 treats the newest parameter, where derivations must be regenerated instead of weakened. Section 7.6 treats the raw guarded clause calculus, with and without a naming exception. Sections 7.7 and 7.8 treat two calculi on a recursor whose predecessor computes; the second combines the restricted introduction with an inferred discharge by means of conditional copy.

Throughout the chapter 1 abbreviates $S0$ and 2 abbreviates $SS0$.

7.1 Restricted witnesses

This section draws on `no_witness`, `newest_parameter`, `parameter_witness` and `deterministic_introducti`

7.1.1 Scopes and the restricted rules

In `N` as stated in Section 1.3 the variables available to a derivation are left implicit. A restriction that mentions the variables in scope needs them explicit.

Definition 7.1 (Scopes)**.** A *numerical scope* is a finite list $\Pi = (a_1, \ldots, a_m)$ of distinct variables, called *parameters*. A term or formula is *over* Π if its free variables are in Π. The *newest parameter* is

$$\nu(\Pi) := \begin{cases} a_m, & m > 0, \\ 0, & m = 0. \end{cases}$$

A judgment is $\Pi; \Gamma \vdash A$, where Γ and A are over Π. The scope and the context are separate lists: an assumption added to Γ does not change Π. The order of Π is proof syntax and has no numerical meaning.

With scopes the rules of N read as follows. Hyp, I0, IS, Tr and QE keep the scope, and the terms they mention are over Π. The three rules with an eigenvariable append it to the scope of their premise:

$$\frac{\Pi;\Gamma \vdash D_{x,y}B \qquad \Pi, b;\ \Gamma, Q_y B(b,y) \vdash A}{\Pi;\Gamma \vdash A}\ \mathrm{DE}$$

$$\frac{\Pi, z;\ \Gamma, B(t,0) \vdash B(t,Sz)}{\Pi;\Gamma \vdash D_{x,y}B}\ \mathrm{DI} \qquad \frac{\Pi, z;\ \Gamma, E(z) \vdash E(Sz)}{\Pi;\Gamma \vdash Q_y E(y)}\ \mathrm{QI}$$

The conclusions are over Π, the witness t of DI is over Π, and E is a literal equation. A derivation of N in the sense of Section 1.3 is a derivation in this sense over any scope that contains its free variables, and conversely.

Definition 7.2 (Restrictions of the witness)**.** Each of the following calculi has the terms, formulas and rules of N, except that the witness t of DI is restricted.

(i) N_0: t is the literal 0.

(ii) N_v: t is a parameter of Π.

(iii) $\mathsf{N}_{0\mathrm{v}}$: t is the literal 0 or a parameter of Π.

(iv) N_ν: t is $\nu(\Pi)$. The rule takes no term:

$$\frac{\Pi, z;\ \Gamma, B(\nu(\Pi),0) \vdash B(\nu(\Pi),Sz)}{\Pi;\Gamma \vdash D_{x,y}B}\ \mathrm{DI}_\nu$$

In DI_ν the witness is computed in the scope Π of the conclusion, before the eigenvariable z is appended. The eigenvariable is never the witness.

Every derivation of N_ν is a derivation of $\mathsf{N}_{0\mathrm{v}}$, and every derivation of N_0, N_v or $\mathsf{N}_{0\mathrm{v}}$ is a derivation of N. Hence on derivable judgments

$$\mathsf{N}_\nu \ \subseteq\ \mathsf{N}_{0\mathrm{v}} \ \subseteq\ \mathsf{N}. \tag{7.1}$$

The distinction between a variable and the constant 0 is syntactic: a term that is provably equal to 0 is not the literal 0.

The derivations of reflexivity, symmetry, transitivity and congruence of Section 1.4 use I0 and Tr only, and equation induction is QI followed by QE. These rules are therefore available, with the same derivations, in every calculus of Theorem 7.2; equations are transformed by them without further comment. In particular the selector equations $\mathrm{sel}(0,a,b) = a$ and $\mathrm{sel}(Sc,a,b) = b$ are available.

7.1.2 What a restricted witness loses

Lemma 7.3 (Structural rules for variable witnesses)**.** *In* $\mathsf{N}_{0\mathrm{v}}$ *and in* N_v *the following are admissible.*

(i) *Renaming and weakening: if* $\Pi;\Gamma \vdash A$*,* ρ *is an injective map from* Π *into a scope* Π'*, and* Δ *is a context over* Π' *that contains* $\rho\Gamma$*, then* $\Pi';\Delta \vdash \rho A$*.*

(ii) *Cut: if* $\Pi;\Gamma \vdash A$ *and* $\Pi;\Gamma, A \vdash B$*, then* $\Pi;\Gamma \vdash B$*.*

Proof. (i) By induction on the derivation, for all ρ and Δ. A renaming sends a variable witness to a variable witness and fixes the literal 0. At a rule with an eigenvariable, choose a fresh parameter of the receiving scope, extend ρ by it and extend Δ by the renamed assumption. (ii) By induction on the derivation of $\Pi; \Gamma, A \vdash B$. A use of Hyp for A is replaced by the derivation of A, renamed along the inclusion of Π into the scope of that leaf and weakened by the assumptions added on the way, by (i). All other rule data are unchanged. □

The proof of (i) does not extend to substitution. A derivation of $\mathsf{N}_{0\mathrm{v}}$ that introduces $D_{x,y}B$ at a parameter a is not turned into a derivation by substituting a compound term for a, because the image of the witness is not a variable. Closure under substitution holds for $\mathsf{N}_{0\mathrm{v}}$ (Theorem 7.21), but only as a consequence of its completeness. The arguments of the chapter therefore never substitute a term into a derivation of a restricted calculus; where a formula $B(t)$ is needed for a compound t, it is obtained by QE from a closure, or the derivation is constructed anew for that instance.

In N_ν less is available.

Lemma 7.4 (Weakening of the context at a fixed scope)**.** *If* $\Pi; \Gamma \vdash A$ *in* N_ν *and* $\Gamma \subseteq \Delta$ *with* Δ *over* Π*, then* $\Pi; \Delta \vdash A$ *in* N_ν*.*

Proof. By induction on the derivation. The scope of every node is unchanged, so every DI_ν selects the same witness; the eigenvariables are not in Π and hence not free in Δ. □

Example 7.5 (Extending the scope changes the witness)**.** In the empty scope N_ν derives $D_{x,y}(\mathrm{sel}(y, 0, x) = 0)$: the selected witness is 0, and the premise asks for $\mathrm{sel}(Sz, 0, 0) = 0$, an instance of IS. The same tree is not a derivation over the scope (a). There the rule selects a and its premise asks for $\mathrm{sel}(Sz, 0, a) = 0$ from $\mathrm{sel}(0, 0, a) = 0$, that is, for $a = 0$, which is not derivable because it is false at $a = 1$ in the standard reading. The judgment itself remains derivable over (a), by Theorem 7.35; its derivation there is a different tree.

Thus a derivation of N_ν cannot be weakened to a larger scope by replaying its tree, and the usual proof of cut fails as well: when the receiving derivation enters a branch with a fresh parameter, the derivation to be inserted would have to be weakened to the larger scope. Section 7.5 replaces the weakening of derivations by families of derivations that are constructed anew in every scope; in the raw guarded calculus with a naming exception, weakening is recovered by naming the old selection anew (Theorem 7.45).

7.2 Obstructions

This section draws on `newest_parameter` for the two shapes and on `constructed_computation` for the simultaneous restriction.

7.2.1 Each shape alone is incomplete

Proposition 7.6 (One shape is not enough)**.** (i) N_0 *does not derive* $D_{x,y}(\mathrm{sel}(y, 1, x) = 1)$ *in the empty scope and context.* N *derives it.*

(ii) N_v *derives no formula* $D_{x,y}B$ *in the empty scope and the empty context.* N *derives* $D_{x,y}(0 = 0)$ *there.*

Proof. (i) Read the terms in the natural numbers, read $Q_y B$ as $B(0) \to \forall y\, B(y)$, and read $D_{x,y}B$ at $x = 0$, as $B(0, 0) \to \forall y\, B(0, y)$. A judgment is valid if its conclusion is true under every assignment of its parameters that makes its context true. All rules of N_0 preserve validity. For Hyp, the computation equations, Tr and QE this is the usual verification, and QI is

induction in the natural numbers. For DE, the premise is valid at $b = 0$, where its assumption $Q_y B(0, y)$ is the reading of the major premise. For DI at the witness 0, the premise gives $B(0, Sz)$ for every z from $B(0, 0)$, and $B(0, 0)$ gives itself. The formula $D_{x,y}(\mathrm{sel}(y, 1, x) = 1)$ is false in this reading: its instance at $y = 0$ is $\mathrm{sel}(0, 1, 0) = 1$, which is true, and its instance at $y = 1$ is $\mathrm{sel}(1, 1, 0) = 1$, which says $0 = 1$. In N the rule DI with the witness 1 asks for $\mathrm{sel}(Sz, 1, 1) = 1$, an instance of IS.

(ii) Call a formula *simple* if it is an equation or a formula $Q_y E$ with E an equation. In N_v, every formula derived over the empty scope from a context of simple formulas is simple. This is proved by induction on the derivation, the claim being made only for the nodes whose scope is empty. Hyp yields a member of the context. I0, IS and Tr yield equations. The major premise $Q_y B$ of QE has the same scope and context and is simple by the induction hypothesis, so B is an equation and so is $B(t)$. The major premise of DE would be a formula $D_{x,y} B$ derived over the empty scope from the same context, against the induction hypothesis; hence DE does not occur at such a node. DI needs a parameter of the empty scope as its witness and does not occur. The conclusion $Q_y E$ of QI is simple by the form of the rule. In N the rule DI with the witness 0 derives $D_{x,y}(0 = 0)$, its premise being reflexivity. □

The Lean development contains both statements: the calculus with the witness restricted by shape is `DWPrf`, and the two parts are `dwprf_zero_incomplete` and `dwprf_var_incomplete` in the module `NativeWitnessShape`. Part (ii) explains why DI_ν uses 0 in the empty scope, and part (i) why the parameters matter. The proposition concerns the two shapes only; it is not a statement about all proof calculi.

7.2.2 The witness and the target cannot both be restricted

The target t of QE can be restricted to a variable or the numeral 1 without loss (Lean: `vqprf_iff_nprf`). The two restrictions do not combine.

Theorem 7.7 (Finite-range obstruction)**.** *Let* N_R *be* N *with the witness of DI restricted to* 0 *or a parameter in scope and the target of QE restricted to* 1 *or a parameter in scope. Then* N_R *does not derive*

$$D_{x,y}\big(\mathrm{sel}(y, 1, x) = 1\big),$$

which N *derives. The same holds when* N *is replaced by the calculus of the reports, with transport in formula templates and with the witnessed introduction by progress for literal equations, and for the introduction by progress with a compound matrix.*

Proof. Evaluate terms in the natural numbers. For a formula A and an assignment e of numbers to variables define a truth value $[A]_e$ by

$$\begin{aligned} [s = t]_e &:\Leftrightarrow s \text{ and } t \text{ have the same value under } e, \\ [Q_y B]_e &:\Leftrightarrow [B]_{e[y \mapsto 0]} \Rightarrow [B]_{e[y \mapsto 1]}, \\ [D_{x,y} B]_e &:\Leftrightarrow [B]_{e[x \mapsto 0,\, y \mapsto 0]} \Rightarrow [B]_{e[x \mapsto 0,\, y \mapsto 1]}. \end{aligned}$$

These clauses define an invariant of derivations; they are not an ordinary first-order interpretation. By induction on B, the value $[B(t)]_e$ equals $[B]_{e[y \mapsto n]}$, where n is the value of t under e; bound variables are renamed away from t before the substitution. Call a judgment *valid* if its conclusion is true under the assignment e_0 that sends every parameter to 0 whenever its context is true under e_0. Every rule of N_R preserves validity.

Hyp is immediate. The computation equations are true under every assignment, and Tr, as well as transport in a formula template, preserves truth by the substitution property.

DI: an allowed witness has the value 0 under e_0, and the eigenvariable z has the value 0, so Sz has the value 1. The premise therefore supplies exactly the implication that defines

$[D_{x,y}B]_{e_0}$. For the introduction by progress the premise has the assumption $B(t,z)$, which has the same value at $z=0$, and the same implication results; the matrix need not be an equation for this argument. QI is the same computation with the clause for Q. The context does not contain z and is unaffected.

DE: the fresh parameter b has the value 0. The assumption $Q_yB(b,y)$ then has the truth value of the major premise $D_{x,y}B$, so the validity of the second premise gives the conclusion.

QE: at a parameter, the conclusion $B(t)$ has the truth value of the premise $B(0)$. At the target 1 the rule is the elimination of the implication that defines $[Q_yB]_{e_0}$.

By induction on derivations every derivable judgment is valid. For the displayed formula the instance at $x=0$, $y=0$ is $\mathrm{sel}(0,1,0)=1$, which is true, and the instance at $x=0$, $y=1$ is $\mathrm{sel}(1,1,0)=1$, which is false. In N, DI with the witness 1 derives the formula from $\mathrm{sel}(Sz,1,1)=1$. □

Theorem 7.8 (Finitely many numeral targets do not help)**.** *Fix $N \geq 1$ and let $W = \{0,\dots,N\}$. Let $\mathsf{N}_{\mathrm{R},N}$ be N with the witness of DI restricted to 0 or a parameter and the target of QE restricted to a parameter or a numeral in W. Then $\mathsf{N}_{\mathrm{R},N}$ does not derive*

$$D_{x,y}\big(\mathrm{sel}(y,N+1,x)=N+1\big),$$

which N derives with the witness $N+1$.

Proof. Evaluate terms in the natural numbers. For an arbitrary assignment e of numbers to variables define $[s=t]_e$ as before and

$$[Q_yB]_e :\Leftrightarrow [B]_{e[y\mapsto 0]} \Rightarrow \bigwedge_{j\in W} [B]_{e[y\mapsto j]},$$

$$[D_{x,y}B]_e :\Leftrightarrow \bigvee_{i\in W} \Big([B]_{e[x\mapsto i,\, y\mapsto 0]} \Rightarrow \bigwedge_{j\in W} [B]_{e[x\mapsto i,\, y\mapsto j]}\Big).$$

The bound variables range over W; the values of terms, and hence the values at which a matrix is evaluated after a substitution, may lie outside W, and the substitution property holds as before. A judgment is *valid* if its conclusion is true under every assignment that sends its parameters into W and makes its context true. DI by cases preserves validity: the witness has a value $i \in W$, every positive $j \leq N$ is $S(j-1)$ with $j-1 \in W$, and the case $j=0$ is the assumption. The introductions by progress, and QI, preserve validity by the finite induction from 0 to N. DE uses a witness $i \in W$ and the validity of its second premise at the assignment $b \mapsto i$. QE has a target with a value in W. Transport and the computation equations are as before. The displayed formula is false: for every $i \in W$ the instance at $y=0$ is true, and the instance at $y=1$, which belongs to the conjunction because $1 \in W$, says $i = N+1$. □

Remark 7.9 (Scope of the obstruction)*.* The two theorems exclude the equality of the theorem sets in the identity language under the simultaneous restriction. They do not show that a differently interpreted theory with both restrictions fails to be bi-interpretable with iPRA. The invariant of Theorem 7.7 is not the case $N=1$ of Theorem 7.8: it fixes the witness at 0 and gives a smaller separating formula. Both invariants remain sound for the rule of `constructed_computation` that supplies a variable from a known instance at zero,

$$\frac{\Pi, w;\ \Gamma, H(w) \vdash A \qquad \Pi;\Gamma \vdash H(0)}{\Pi;\Gamma \vdash A} \qquad w \notin \mathrm{FV}(\Gamma, A).$$

In the first invariant $H(w)$ and $H(0)$ have the same truth value. In the second, extend the assignment by $w \mapsto 0$, which is allowed because $0 \in W$; the second premise makes $H(w)$ true there and the first supplies A. No rule that names an arbitrary term by a witness outside W is covered.

7.3 Names

This section draws on `deterministic_introduction` for the naming matrix, on `no_witness` and `parameter_witness` for the derived rules of the calculus with variable witnesses, and on the Lean module `NativeNamedWitness` for the organization.

A restricted introduction cannot introduce $D_{x,y}B$ at a compound term t. It can introduce it at a parameter b that is provably equal to t. A *name* of t is a formula whose opening at a fresh b yields the equation $b = t$.

7.3.1 Naming matrices

Definition 7.10 (Gate, naming matrix, alias matrix)**.** For an equation $E \equiv (s = t)$ and a term v the *gate* of E is

$$\gamma_v(E) := \big(\mathrm{sel}(v, 0, s) = \mathrm{sel}(v, 0, t)\big).$$

For a term r, and variables x, y not free in r, the *naming matrix* and the *alias matrix* of r are

$$\mathrm{Nm}_r(x, y) := \gamma_y(x = r) = \big(\mathrm{sel}(y, 0, x) = \mathrm{sel}(y, 0, r)\big), \qquad \mathrm{Al}_r(x, y) := \big(\mathrm{sel}(y, r, x) = r\big).$$

The formulas $D_{x,y}\mathrm{Nm}_r$ and $D_{x,y}\mathrm{Al}_r$ are the *names* of r.

By the selector equations, $\gamma_0(E)$ is derivable in every context, both sides being equal to 0, and $\gamma_{Sz}(E)$ is interderivable with E. The ordinary reading of either name is equivalent to $\exists x\,(x = r)$: the instance of the matrix at $y = 0$ is provable and its instance at $y = 1$ is equivalent to $x = r$. The gate cannot be omitted. The formula $D_{x,v}(x = r)$, whose matrix does not contain v, reads $\exists x\,(x = r \to x = r)$ and says nothing about r.

Lemma 7.11 (A name is derivable with the witness it names)**.** *For every term r over Π, N derives $\Pi; \Gamma \vdash D_{x,y}\mathrm{Nm}_r$ and $\Pi; \Gamma \vdash D_{x,y}\mathrm{Al}_r$ by DI with the witness r.*

Proof. The premise for the naming matrix asks for $\mathrm{sel}(Sz, 0, r) = \mathrm{sel}(Sz, 0, r)$, which is reflexivity. The premise for the alias matrix asks for $\mathrm{sel}(Sz, r, r) = r$, an instance of IS. Neither uses its assumption. □

Lemma 7.12 (Opening a name)**.** *Let Γ contain $Q_y\mathrm{Nm}_r(b, y)$ or $Q_y\mathrm{Al}_r(b, y)$. Then $\Gamma \vdash b = r$, by a derivation that uses Hyp, I0, IS, Tr and one QE with the target* 1.

Proof. For the naming matrix, the instance at 0 is $\mathrm{sel}(0, 0, b) = \mathrm{sel}(0, 0, r)$; both sides equal 0 by I0. QE at 1 gives $\mathrm{sel}(1, 0, b) = \mathrm{sel}(1, 0, r)$, and IS turns the two sides into b and r. For the alias matrix, the instance at 0 is $\mathrm{sel}(0, r, b) = r$, an instance of I0. QE at 1 gives $\mathrm{sel}(1, r, b) = r$, and IS gives $\mathrm{sel}(1, r, b) = b$. □

Opening the naming matrix uses reflexivity where opening the alias matrix uses one successor computation; otherwise the two are interchangeable. The naming matrix is the one that `deterministic_introduction` recognizes syntactically (Section 7.6); the alias matrix is the one of the Lean development.

7.3.2 Derived rules with variable witnesses

Let $\mathsf{N}_{\mathrm{v}}^{+}$ be N_{v} together with the axiom scheme

$$\Pi; \Gamma \vdash D_{x,y}\mathrm{Al}_a \qquad \text{for every term } a \text{ over } \Pi.$$

A renaming sends an instance of the scheme to an instance, so Theorem 7.3 holds for $\mathsf{N}_{\mathrm{v}}^{+}$ with the same proof. The next two lemmas are stated for $\mathsf{N}_{\mathrm{v}}^{+}$ and for $\mathsf{N}_{0\mathrm{v}}$ at once.

Lemma 7.13 (Closure by cases)**.** *If* $\Pi, z;\ \Gamma, B(0) \vdash B(Sz)$*, then* $\Pi; \Gamma \vdash Q_y B(y)$*. Here* B *is an arbitrary native formula.*

Proof. Choose r not free in B. In $\mathsf{N}_{0\mathrm{v}}$, DI with the witness 0 derives $D_{r,y}B(y)$ from the given premise. DE at a fresh b adds the assumption $Q_y B(y)$, which is the conclusion and does not contain b. In $\mathsf{N}_{\mathrm{v}}^{+}$ a variable is needed as the witness. Open the name $D_{x,y}\mathrm{Al}_0$ of 0 at a fresh parameter c and introduce $D_{r,y}B(y)$ at c; its premise is the given one, renamed along the inclusion of scopes and weakened by Theorem 7.3. Eliminate $D_{r,y}B(y)$ as before, and discharge c by DE: the conclusion $Q_y B(y)$ does not contain it. □

Lemma 7.14 (Leibniz rule)**.** *For every native formula* $F(v)$ *with a distinguished variable* v*: if* $\Pi; \Gamma \vdash s = t$ *and* $\Pi; \Gamma \vdash F(s)$*, then* $\Pi; \Gamma \vdash F(t)$*.*

Proof. By induction on the number of binders of F. The statement is taken for all scopes, contexts and terms at once, because the induction hypothesis is applied to new instances of the matrix; no term is substituted into a derivation.

If $F(v)$ is an equation $p(v) = q(v)$, congruence gives $p(s) = p(t)$ and $q(s) = q(t)$, and symmetry and transitivity give $p(t) = q(t)$.

Let $F(v) = Q_y B(v, y)$. By Theorem 7.13 it suffices to derive $B(t, Sz)$ over Π, z from $\Gamma, B(t, 0)$. The equation $t = s$ and the induction hypothesis for the matrix $B(v, 0)$ give $B(s, 0)$. QE at Sz, applied to the weakened premise $Q_y B(s, y)$, gives $B(s, Sz)$. The equation $s = t$ and the induction hypothesis for $B(v, Sz)$ give $B(t, Sz)$.

Let $F(v) = D_{x,y}B(v, x, y)$. Apply DE to the premise at a fresh parameter b, which adds $Q_y B(s, b, y)$. Introduce $D_{x,y}B(t, x, y)$ at the variable b. The premise of this introduction has the assumption $B(t, b, 0)$ and asks for $B(t, b, Sz)$: the induction hypothesis gives $B(s, b, 0)$, QE at Sz gives $B(s, b, Sz)$, and the induction hypothesis gives $B(t, b, Sz)$. The conclusion does not contain b, and DE discharges it. Where a premise is used below a new parameter it is renamed and weakened by Theorem 7.3. □

7.3.3 The introduction at a name

Theorem 7.15 (Variable witnesses and a name for every term)**.** *For every scope* Π*, context* Γ *and formula* A *over* Π*,*

$$\Pi; \Gamma \vdash_{\mathsf{N}_{\mathrm{v}}^{+}} A \quad \Longleftrightarrow \quad \Pi; \Gamma \vdash_{\mathsf{N}} A.$$

Proof. From left to right, an introduction at a variable is an instance of DI, and the axioms are derivable in N by Theorem 7.11.

From right to left it suffices to show that DI at an arbitrary term t over Π is admissible in $\mathsf{N}_{\mathrm{v}}^{+}$; the claim then follows by induction on the derivation in N. Let a derivation of

$$\Pi, z;\ \Gamma, B(t, 0) \vdash B(t, Sz)$$

be given. Take the axiom $D_{x,y}\mathrm{Al}_t$ and apply DE to it at a fresh parameter b. The branch has the scope Π, b and the context $\Gamma_b = \Gamma, Q_y \mathrm{Al}_t(b, y)$, and Theorem 7.12 gives $\Gamma_b \vdash b = t$. In this branch introduce $D_{x,y}B$ at the variable b. The premise of the introduction has the scope Π, b, z and the context $\Gamma_b, B(b, 0)$. The Leibniz rule gives $B(t, 0)$ from $B(b, 0)$ and $b = t$. Rename the given derivation along the inclusion of Π, z into Π, b, z, weaken it to the context $\Gamma_b, B(b, 0), B(t, 0)$, and cut $B(t, 0)$; the result is $B(t, Sz)$. The Leibniz rule with $t = b$ gives $B(b, Sz)$. The introduction at b yields $\Pi, b; \Gamma_b \vdash D_{x,y}B$, a formula over Π, and DE discharges b. □

Corollary 7.16 (Completeness is the derivability of the names)**.** $\mathsf{N}_{0\mathrm{v}}$ *derives the same judgments as* N *if and only if it derives* $\Pi; \varnothing \vdash D_{x,y}\mathrm{Al}_a$ *for every scope* Π *and every term* a *over* Π*.*

Proof. If the two calculi agree, the names are derivable by Theorem 7.11. Conversely, suppose the names are derivable in $\mathsf{N}_{0\mathrm{v}}$. A derivation of $\mathsf{N}_{\mathrm{v}}^{+}$ becomes a derivation of $\mathsf{N}_{0\mathrm{v}}$ when each axiom leaf is replaced by the derivation of that name, weakened to the context of the leaf by Theorem 7.3. With Theorem 7.15 and (7.1) this gives $\mathsf{N} \subseteq \mathsf{N}_{\mathrm{v}}^{+} \subseteq \mathsf{N}_{0\mathrm{v}} \subseteq \mathsf{N}$. □

Remark 7.17 (Lean status). Theorem 7.15 is `dnprf_iff_nprf` for the calculus `DNPrf`, and Theorem 7.16 is `dwprf_complete_iff_names`; both are in `NativeNamedWitness`. The steps of the Lean proof are the ones above: the name, its opening, the introduction at the name, and a Leibniz rule proved over one context without substitution into derivations; a dummy witness in a scope without variables is a name of 0, as in Theorem 7.13. The derivation of the names inside the restricted calculus, which is the subject of the next section, is formalized as well (Theorem 7.23).

7.4 The coverage packet

This section draws on `no_witness` (its restriction note), `parameter_witness` (its appendix on restricted witnesses) and `newest_parameter`.

By Theorem 7.16 it remains to derive the name of an arbitrary term t with variable witnesses. The name cannot be introduced at t. The route is indirect: a formula $X(y)$ that says "y is 0 or a successor" is proved for all y by numerical cases, where in the successor case the witness is the case variable; it is then instantiated at $y = St$ by QE, whose target is unrestricted; and the injectivity of the successor turns the opened instance into a name of t.

7.4.1 The predecessor

Let $\mathcal{E}$ be the equational fragment of Theorem 1.36: Hyp, the two computation equations, Tr and induction for an equation, on judgments whose assumptions and conclusions are equations. By Theorem 1.48, $\mathcal{E}$ derives, for the predecessor term written P there and pred in this chapter,

$$\mathrm{pred}(0) = 0, \qquad \mathrm{pred}(Sn) = n. \tag{7.2}$$

$\mathcal{E}$ is closed under substitution (Theorem 1.37), so for every term r it derives $\mathrm{pred}(Sr) = r$. In a calculus that has QI and QE, a derivation of $\mathcal{E}$ is a derivation without any introduction of $D_{,}$. Hence in N_0, $\mathsf{N}_{0\mathrm{v}}$ and N_ν, for all Π, Γ and terms r, s over Π,

$$\Pi; \Gamma \vdash \mathrm{pred}(Sr) = r, \qquad \Pi;\ \Gamma,\ Sr = Ss \vdash r = s, \tag{7.3}$$

the second by congruence with pred. The order of the argument matters. The predecessor is obtained before any term is named; obtaining it from the bridge of Theorem 1.78, which is proved for the unrestricted calculus, would be circular. That the derivation needs no case distinction under a hypothesis, hence no DI, is the contribution of the four packages recorded in Theorem 1.49. In Lean the fragment is `EqPrf`, with `EqPrf.pred_zero`, `EqPrf.pred_succ` and `EqPrf.succ_inj`, and `EqPrf.toDW` embeds it in the kernel under any restriction of the witness.

7.4.2 The packet and the names

Definition 7.18 (Coverage packet)**.** For a term y put

$$M_y(x, v) := \gamma_v\big(\mathrm{sel}(y, 0, Sx) = y\big) = \big(\mathrm{sel}(v, 0, \mathrm{sel}(y, 0, Sx)) = \mathrm{sel}(v, 0, y)\big), \qquad X(y) := D_{x,v} M_y(x, v).$$

The ordinary reading of $X(y)$ is equivalent to $\exists x\,(\mathrm{sel}(y, 0, Sx) = y)$, which holds at $y = 0$ for every x and says $y = Sx$ at a successor. The formula never asks for an introduction of $\exists x\,(x = t)$ at the witness t.

Lemma 7.19 (The packet with variable witnesses)**.** *In* $\mathsf{N}_{0\mathrm{v}}$*, for every term* t *over* Π*,* $\Pi;\Gamma \vdash X(t)$*.*

Proof. $X(0)$: introduce at the witness 0. The premise asks for $M_0(0, Sz)$, that is, for $\mathrm{sel}(Sz, 0, \mathrm{sel}(0, 0, 1)) = \mathrm{sel}(Sz, 0, 0)$. By IS and I0 the left side equals $\mathrm{sel}(0, 0, 1)$ and then 0, and the right side equals 0.

$X(Sz)$ over the scope Π, z: introduce at the variable z. The premise asks for $M_{Sz}(z, Sz')$, that is, for $\mathrm{sel}(Sz', 0, \mathrm{sel}(Sz, 0, Sz)) = \mathrm{sel}(Sz', 0, Sz)$; both sides equal Sz by IS. Neither premise uses its assumption.

By Theorem 7.13, with the assumption $X(0)$ unused, $\Pi;\Gamma \vdash Q_y X(y)$. QE with the base $X(0)$ and the target t gives $X(t)$. This is an application of QE, not a substitution into the derivation of $X(Sz)$. □

Lemma 7.20 (The alias)**.** *Let* t *be over* Π *and* $b \notin \Pi$*. If* Γ *contains* $Q_v M_{St}(b, v)$*, then* $\Gamma \vdash b = t$*, by a derivation in which no* $D_{,}$ *is introduced.*

Proof. The instance $M_{St}(b, 0)$ is a gate at 0 and is derivable. QE at 1 gives $\mathrm{sel}(1, 0, \mathrm{sel}(St, 0, Sb)) = \mathrm{sel}(1, 0, St)$, and IS, applied twice on the left and once on the right, gives $Sb = St$. By (7.3), $b = t$. □

Theorem 7.21 (The literal 0 and variables suffice)**.** *For every scope* Π*, context* Γ *and formula* A *over* Π*,*

$$\Pi;\Gamma \vdash_{\mathsf{N}_{0\mathrm{v}}} A \quad \Longleftrightarrow \quad \Pi;\Gamma \vdash_{\mathsf{N}} A.$$

Consequently $\mathsf{N}_{0\mathrm{v}}$ *is closed under substitution, and the bridge of Theorem 1.78 holds for it with the same translations.*

Proof. By Theorem 7.16 it suffices to derive $\Pi;\varnothing \vdash D_{x,y}\mathrm{Al}_t$ for t over Π. By Theorem 7.19, $X(St)$ is derivable. Apply DE to it at a fresh b; by Theorem 7.20 the branch derives $b = t$. Introduce $D_{x,y}\mathrm{Al}_t$ at the variable b: the premise asks for $\mathrm{sel}(Sz, t, b) = t$, and $\mathrm{sel}(Sz, t, b) = b$ by IS. The name does not contain b, and DE discharges it. The two calculi have the same terms and formulas and the comparison is the identity, so every property of the consequence relation of N transfers. □

Remark 7.22 (Versions of the packet and of the proof)*.* The formula $X(y)$ is $H(y)$ in `no_witness`, $X(y)$ in `newest_parameter` and $A(r)$ in `uniform_raw`. `parameter_witness` uses the ungated packet

$$T(v) := D_{x,y}\big(\mathrm{sel}(y, v, \mathrm{sel}(v, 0, Sx)) = v\big),$$

whose instance at $y = 0$ is $v = v$ and whose instances at successors are $\mathrm{sel}(v, 0, Sx) = v$; it serves in a calculus where the mode of the introduction depends on the witness (Theorem 7.24). The restriction note of `no_witness` and the appendix of `parameter_witness` prove Theorem 7.21 directly, with the naming matrix $D_{x,v}\gamma_v(x = t)$ and a derived rule

$$\frac{\Pi, b;\ \Gamma, b = t \vdash F}{\Pi;\Gamma \vdash F} \qquad b \notin \mathrm{FV}(\Gamma, t, F),$$

obtained by opening $X(St)$ and cutting the alias. The proof above factors the same argument through Theorem 7.15. The theorem answers the question whether DI at the literal 0 or at a variable is enough.

Remark 7.23 (Lean status)*.* Theorem 7.21 is formalized: `dwprf_iff_nprf`, in the module `NativeWitnessComplete`, states that the kernel with DI at the literal 0 or at a variable, and at no other term, derives exactly what the kernel derives. The names are derived there (`DWPrf.names`) from a packet in which the predecessor stands inside the matrix: the name

$D_{x,v}\mathrm{Al}_{\mathrm{pred}(y)}$ of $\mathrm{pred}(y)$ holds at $y = 0$ with the witness 0 and at $y = Sz$ with the witness z, by the closure by cases with the dummy witness 0 (`DWPrf.namePQ`); it is used at $y = Sa$, and opening that name gives a variable equal to $\mathrm{pred}(Sa) = a$ (`DWPrf.nameOf`). The packet $X(y)$ applies the predecessor after the opening instead; the two differ in this order only. The restriction to the newest parameter of Section 7.5 is not formalized.

7.4.3 A mode selected by the witness

`parameter_witness` states the restriction for a calculus with one introduction whose mode is read off the witness. Let R be the seven-rule calculus W of Theorem 2.15, with scopes, in which the witness t of its introduction is the literal 0 or a parameter of Π:

$$\frac{\Pi, z;\ \Gamma, B(t, \eta_{B,t}(z)) \vdash B(t, Sz)}{\Pi; \Gamma \vdash D_{x,y}B(x, y)}\ \mathrm{RDI},$$

where $\eta_{B,t}(z)$ is z if B is a literal equation and t is the literal 0, and 0 otherwise. There is no rule QI. Every introduction at a variable is by cases.

Theorem 7.24 (The calculus with a witness-selected mode)**.** R *and* N *derive the same judgments.*

Proof. Every derivation of R is a derivation of W, which is equivalent to N by Theorem 2.17. For the converse, the proof of Theorem 7.21 is repeated with the following changes.

Structural rules. Theorem 7.3 holds for R: a renaming sends variables to variables, fixes the literal 0 and preserves the root of a matrix, hence the mode.

Closure by progress. If $\Pi, z; \Gamma, E(z) \vdash E(Sz)$ for a literal equation E, then $\Pi; \Gamma \vdash Q_y E(y)$: RDI at the witness 0 on $D_{x,y}E(y)$ with x not free in E is in the mode by progress, and DE returns the opened assumption. With QE this is equation induction, so every derivation of $\mathcal{E}$ is a derivation of R rule by rule, and (7.3) holds in R.

Closure by cases. The introduction of $D_{r,y}B(y)$ at 0 is by progress when B is an equation, so a variable witness is used for every B. The formula $D_{r,w}(0 = 0)$ is derivable by RDI at 0, its premise being reflexivity. Open it at a fresh b and introduce $D_{r,y}B(y)$ at b; this is by cases. Eliminate the unused coordinate to obtain $Q_y B(y)$ and discharge b. This replaces Theorem 7.13.

Leibniz rule. Theorem 7.14 holds with the same proof; the reintroduction in the case of $D_,$ is at a variable and is by cases.

Packet. With T as in Theorem 7.22: $T(0)$ is introduced at 0 by progress, its premise asking for $\mathrm{sel}(Sz, 0, \mathrm{sel}(0, 0, 1)) = 0$, which holds by computation; $T(Sz)$ is introduced at z by cases, its premise asking for $\mathrm{sel}(Sz', Sz, \mathrm{sel}(Sz, 0, Sz)) = Sz$, which holds by computation. Closure by cases gives $Q_v T(v)$ and QE gives $T(t)$ for every t. Opening $T(St)$ at a fresh b gives a closure whose instance at 0 is $\mathrm{sel}(0, St, \cdot) = St$, an instance of I0; QE at 1 and two computations give $Sb = St$, hence $b = t$.

Recovery of the rules of N. QI is closure by progress. For DI at an arbitrary term t, with an arbitrary matrix, open $T(St)$ at b, introduce at b, which is by cases, and transport the assumption and the conclusion as in the proof of Theorem 7.15. This includes the introduction by cases of an equation at the literal 0, which RDI does not offer directly. □

7.5 The newest parameter

This section draws on `no_witness` and `newest_parameter` for the eight-rule calculus, on `no_witness` for the seven-rule diagonal variant, and on `parameter_witness` for the one-bit rule and its clause calculus.

In N_ν the introduction takes no term. The proof of Theorem 7.21 uses renaming, weakening to a larger scope and cut, none of which is available for the derivations of N_ν (Theorem 7.5). The remedy is to work with families of derivations that are constructed anew in every scope. A local derivation whose witness is a freshly chosen case variable z is used while z is the newest parameter; it is never moved to a scope in which z is older.

7.5.1 Stable derivations and hereditary families

Definition 7.25 (Stable derivations)**.** An application of DI_ν with the matrix $B(x, y)$ is *dummy* if x is not free in B. A derivation of N_ν is *stable* if all its introductions are dummy.

The premise of a dummy introduction does not depend on the selected witness.

Lemma 7.26 (Stability)**.** *Let $\Pi; \Gamma \vdash A$ have a stable derivation. Let θ assign to each parameter of Π a term over a scope Π', and let Δ be a context over Π' that contains $\Gamma[\theta]$. Then $\Pi'; \Delta \vdash A[\theta]$ has a stable derivation. If $\Pi; \Gamma \vdash A$ and $\Pi; \Gamma, A \vdash B$ have stable derivations, so has $\Pi; \Gamma \vdash B$.*

Proof. The first statement is proved as the substitution lemma of Section 1.4, by induction on the derivation for all θ and Δ. At a dummy introduction the premise is $\Gamma, B(0) \vdash B(Sz)$ whatever the witness is, and the rule applies in the scope Π' to the matrix $B[\theta]$, in which x is still not free. The second statement is the usual induction on the derivation of $\Pi; \Gamma, A \vdash B$; the inserted derivation of A is moved to the scope of each leaf by the first statement. □

Derivations without any introduction of $D_,$ are stable. In particular the judgments (7.3) and the derivation of Theorem 7.20 are stable in N_ν.

Definition 7.27 (Hereditary derivability)**.** Write $\Pi; \Gamma \Vdash A$ if for every scope Π', every injective map ρ from Π into Π' and every context Δ over Π' that contains $\rho\Gamma$, the calculus N_ν derives $\Pi'; \Delta \vdash \rho A$.

Hereditary derivability is a property of a family of derivations, one for each (Π', ρ, Δ); it is not a rule of the calculus. The member of the family at the identity is a derivation of $\Pi; \Gamma \vdash A$. The families below are given by constructions that take the receiving scope and context as input and choose their eigenvariables there.

Lemma 7.28 (Closure properties)**.** (i) *If $A \in \Gamma$, then $\Pi; \Gamma \Vdash A$.*

(ii) *If $\Pi; \Gamma \Vdash A$, ρ is an injective map from Π into Π' and $\Delta \supseteq \rho\Gamma$ is over Π', then $\Pi'; \Delta \Vdash \rho A$.*

(iii) *If $\Pi; \Gamma \vdash A$ has a stable derivation, then $\Pi; \Gamma \Vdash A$.*

(iv) *The rules I0, IS, T_R and QE hold with $\Vdash$ in place of $\vdash$.*

(v) *If $\Pi; \Gamma \Vdash D_{x,y}B$ and $\Pi, b;\ \Gamma, Q_y B(b, y) \Vdash A$ with A over Π, then $\Pi; \Gamma \Vdash A$.*

(vi) *If $\Pi, z;\ \Gamma, E(z) \Vdash E(Sz)$ for a literal equation E, then $\Pi; \Gamma \Vdash Q_y E(y)$.*

Proof. (i) is Hyp in every receiving context. (ii) holds because injective maps compose. (iii) is Theorem 7.26. For (iv), apply the rule to the members of the given families at (Π', ρ, Δ), with the terms renamed by ρ. For (v), let (Π', ρ, Δ) be given. Choose $b' \notin \Pi'$ and extend ρ by $b \mapsto b'$. The member of the second family at (Π', b') and the context $\Delta, Q_y \rho B(b', y)$ derives ρA, and DE applies to it and to the member of the first family. (vi) is proved in the same way with QI. □

Lemma 7.29 (Uniform cut)**.** *If* $\Pi;\Gamma \Vdash A$ *and* $\Pi;\Gamma, A \vdash B$ *in* N_ν*, then* $\Pi;\Gamma \vdash B$ *in* N_ν*. Consequently, if* $\Pi;\Gamma \Vdash A$ *and* $\Pi;\Gamma, A \Vdash B$*, then* $\Pi;\Gamma \Vdash B$*.*

Proof. By induction on the derivation of $\Pi;\Gamma, A \vdash B$, for all Π and Γ. At HYP for A, take the member of the family at the identity; any other HYP is kept. A rule whose premises have the same scope and context is reapplied to the results of the induction hypothesis. At a rule with an eigenvariable, say DI_ν with the premise $\Pi, z;\ \Gamma, A, B(w,0) \vdash B(w,Sz)$ and $w = \nu(\Pi)$, Theorem 7.28(ii) gives $\Pi, z;\ \Gamma, B(w,0) \Vdash A$, the induction hypothesis gives $\Pi, z;\ \Gamma, B(w,0) \vdash B(w,Sz)$, and the rule is reapplied in the scope Π, where it selects the same w. DE and QI are treated in the same way. Every node of the receiving derivation keeps its scope, and no derivation is moved to another scope: below a new parameter the family supplies a new member.

For the second statement let (Π', ρ, Δ) be given. The second family has a member deriving $\Pi';\Delta, \rho A \vdash \rho B$, and $\Pi';\Delta \Vdash \rho A$ by Theorem 7.28(ii). □

7.5.2 Closure by cases, the Leibniz rule and equation cases

Lemma 7.30 (Closure by cases)**.** *In* N_ν*, if* $\Pi, z;\ \Gamma, B(0) \vdash B(Sz)$*, then* $\Pi;\Gamma \vdash Q_y B(y)$*; if the given derivation is stable, so is the result. If* $\Pi, z;\ \Gamma, B(0) \Vdash B(Sz)$*, then* $\Pi;\Gamma \Vdash Q_y B(y)$*.*

Proof. Choose r not free in B and apply DI_ν to $D_{r,y}B(y)$. The introduction is dummy, and its premise is the given one whatever $\nu(\Pi)$ is. DE at a fresh b adds the assumption $Q_y B(y)$, which is the conclusion. For the hereditary statement, the member of the given family at (Π', z'), $\rho[z \mapsto z']$ and $\Delta, \rho B(0)$ is the premise needed in the scope Π'. □

Lemma 7.31 (Hereditary Leibniz rule)**.** *For every native formula* $F(v)$*: if* $\Pi;\Gamma \Vdash s = t$ *and* $\Pi;\Gamma \Vdash F(s)$*, then* $\Pi;\Gamma \Vdash F(t)$*.*

Proof. By induction on the number of binders of F, for all scopes, contexts and terms. For an equation the claim follows from Theorem 7.28(iv), as in Theorem 7.14.

Let $F(v) = Q_y B(v,y)$. By Theorem 7.30 it suffices to show $\Theta \Vdash B(t,Sz)$, where Θ stands for the scope Π, z with the context $\Gamma, B(t,0)$. By Theorem 7.28, $\Theta \Vdash t = s$, $\Theta \Vdash B(t,0)$ and $\Theta \Vdash Q_y B(s,y)$. The induction hypothesis gives $\Theta \Vdash B(s,0)$, QE at Sz gives $\Theta \Vdash B(s,Sz)$, and the induction hypothesis gives $\Theta \Vdash B(t,Sz)$.

Let $F(v) = D_{x,y}B(v,x,y)$, and let (Π', ρ, Δ) be given; write s', t', B' for the images under ρ. The first family gives $\Pi';\Delta \vdash D_{x,y}B'(s',x,y)$. Apply DE at a fresh b. In the scope Π', b the parameter b is the newest, so DI_ν for $D_{x,y}B'(t',x,y)$ selects b; its premise has the scope Π', b, z and the context $\Delta, Q_y B'(s',b,y), B'(t',b,0)$, denoted Θ. As in the previous case, $\Theta \Vdash B'(s',b,0)$ by the induction hypothesis, $\Theta \Vdash B'(s',b,Sz)$ by QE, and $\Theta \Vdash B'(t',b,Sz)$ by the induction hypothesis. The member at the identity is the premise of the introduction. The conclusion does not contain b, and DE discharges it. Whatever the receiving scope is, the parameter b is generated after all its parameters, so the same recipe always introduces at the newest parameter. □

The derivation of (7.2) needs only equation induction. `newest_parameter` observes in addition that case distinctions on a term of the context are available in N_ν for equational conclusions, without DI at a chosen witness. The statement is Theorem 3.51 for hereditary families, with the implication $U \Rightarrow_= V$ between equations defined there.

Lemma 7.32 (Contextual equation cases)**.** *In* N_ν*, for equations* U*,* V*,* F *and a term* e *over* Π*:*

(i) *if* $\Pi;\Gamma, U \Vdash V$*, then* $\Pi;\Gamma \Vdash U \Rightarrow_= V$*;*

(ii) *if* $\Pi;\Gamma \Vdash U \Rightarrow_= V$ *and* $\Pi;\Gamma \Vdash U$*, then* $\Pi;\Gamma \Vdash V$*;*

(iii) *if* $\Pi;\Gamma, e=0 \Vdash F$ *and* $\Pi, z;\ \Gamma, e=Sz \Vdash F$*, then* $\Pi;\Gamma \Vdash F$*.*

Proof. The proof of Theorem 3.51 applies, with Theorem 7.30 for the closure by cases and Theorem 7.29 for the cut; in (iii) the two hypotheses give $(e=y) \Rightarrow_= F$ at $y=0$ and at $y=Sz$ by (i), the closure by cases and QE give it at $y=e$, and (ii) with $e=e$ gives F. □

All introductions in the constructions of Theorems 7.30 and 7.32 are dummy; applied to stable derivations they yield stable derivations.

7.5.3 The packet in every scope

Lemma 7.33 (The packet with the newest parameter)**.** *In* N_ν*, for every scope* Π*, every context* Γ *and every term* t *over* Π*:* $\Pi;\Gamma \vdash X(St)$*. Hence* $\Pi;\Gamma \Vdash X(St)$*.*

Proof. $X(0)$: the rule selects $w=\nu(\Pi)$, and its premise asks for $M_0(w,Sz)$, that is, for $\mathrm{sel}(Sz,0,\mathrm{sel}(0,0,Sw))=\mathrm{sel}(Sz,0,0)$. Both sides equal 0 by computation, whatever w is. So $X(0)$ is derivable in the empty scope and also when w is an unrelated parameter.

$D_{r,y}X(y)$ with r unused: the premise of this introduction has the scope Π,z and asks for $X(Sz)$ from $X(0)$. In the scope Π,z the newest parameter is z, so the introduction of $X(Sz)$ selects z; its premise asks for $M_{Sz}(z,Sz')$, both sides of which equal Sz by computation. The assumptions of both premises are unused.

DE on $D_{r,y}X(y)$ gives $Q_yX(y)$, and QE with the base $X(0)$ and the target St gives $X(St)$. The construction works in every scope and every context, and $\rho X(St)=X(S\rho t)$; this is the hereditary statement. □

The construction is applied to the whole closure. It is not claimed that $\Pi,z;\Gamma \Vdash X(Sz)$: a derivation that introduces $X(Sz)$ at z is not a derivation in a scope in which z is no longer the newest parameter. In each receiving scope the case variable is chosen last.

Lemma 7.34 (The hereditary alias)**.** *Let* Θ *be a scope and context such that the context contains* $Q_vM_{St}(b,v)$*. Then* $\Theta \Vdash b=t$*.*

Proof. The derivation of Theorem 7.20 is stable. □

7.5.4 The equivalence theorem

Theorem 7.35 (No witness in the eight-rule calculus)**.** *For every scope* Π*, context* Γ *and formula* A *over* Π*,*

$$\Pi;\Gamma \vdash_{\mathsf{N}_\nu} A \quad\Longleftrightarrow\quad \Pi;\Gamma \vdash_{\mathsf{N}} A.$$

Both directions are effective transformations of derivations, and the terms and formulas of the judgment are unchanged.

Proof. From left to right every rule of N_ν is an instance of a rule of N. For the converse the following is proved by induction on a derivation of N:

$$\text{if } \Pi;\Gamma \vdash_{\mathsf{N}} A, \text{ then } \Pi;\Gamma \Vdash A.$$

Hyp, I0, IS, Tr and QE are Theorem 7.28(i) and (iv). DE and QI are Theorem 7.28(v) and (vi), applied to the families that the induction hypothesis gives for the premises.

Consider DI with the witness t and the matrix B. The induction hypothesis is

$$\Pi, z;\ \Gamma, B(t,0) \Vdash B(t,Sz). \tag{7.4}$$

Let (Π', ρ, Δ) be given and write $t' = \rho t$, $B' = \rho B$. By Theorem 7.33, $\Pi'; \Delta \vdash X(St')$. Apply DE to it at a fresh b; the branch has the scope Π', b and the context $\Delta_b = \Delta, Q_v M_{St'}(b, v)$. In this branch apply DI_ν to $D_{x,y}B'$. The newest parameter is b, so the rule selects b, and its eigenvariable z' is appended after this selection. Let Θ be the scope Π', b, z' with the context $\Delta_b, B'(b, 0)$; the premise to be derived is $\Theta \vdash B'(b, Sz')$.

(a) $\Theta \Vdash b = t'$ by Theorem 7.34, and $\Theta \Vdash B'(b, 0)$ by membership. By Theorem 7.31, $\Theta \Vdash B'(t', 0)$.

(b) The map $\rho[z \mapsto z']$ from Π, z into Π', b, z' is injective, and the context of Θ extended by $B'(t', 0)$ contains the image of $\Gamma, B(t, 0)$. By (7.4) and Theorem 7.28(ii), $\Theta, B'(t', 0) \Vdash B'(t', Sz')$.

(c) By (a), (b) and Theorem 7.29, $\Theta \Vdash B'(t', Sz')$.

(d) By Theorem 7.31 with the equation $t' = b$, $\Theta \Vdash B'(b, Sz')$.

The member of the last family at the identity is the premise. The introduction yields $\Pi', b; \Delta_b \vdash D_{x,y}B'$, a formula over Π', and DE discharges b. This was done for an arbitrary (Π', ρ, Δ), so $\Pi; \Gamma \Vdash D_{x,y}B$.

The construction is well founded. The induction hypothesis is used for the premises of the given derivation only. Theorem 7.31 is an induction on formulas. The packet, the alias and the predecessor are fixed constructions that precede the theorem. No derivation of N_ν is moved to another scope and no term is substituted into one. Finally the member at the identity of $\Pi; \Gamma \Vdash A$ is the required derivation. □

Corollary 7.36 (Structural rules and the three calculi)**.** N_ν, $\mathsf{N}_{0\mathrm{v}}$ *and* N *derive the same judgments. Substitution, weakening to a larger scope and cut are admissible in* N_ν *as transformations of derivable judgments. The ordinary realization of* N_ν *is bi-interpretable and equiconsistent with* iPRA *by the translations of Theorem 1.78, with whole domains, actual equality and the identity graph* $x = y$ *in both numerical composites.*

Proof. The first statement is Theorem 7.35 and (7.1). For the structural rules translate to N, apply its rule (Section 1.4) and translate back; they are consequences of the theorem, not steps of its proof, and they do not say that an unchanged derivation tree is a derivation after an extension of the scope. The comparison with N is the identity on terms, formulas and contexts, so the translations of the bridge, the derivations of both formula round trips and the preservation of $0 = S0$ transfer unchanged. □

Remark 7.37 (The three formulations of the proof families)*.* `no_witness` calls the families *portable*: a procedure that, for every extension of the scope by fresh parameters and of the context by assumptions, constructs a derivation of the unchanged formula. `newest_parameter` calls them *persistent builders*, with the same definition. Both state the induction on the source derivation for an arbitrary substitution σ of terms over a target scope for the source parameters, a target context Δ, and a family for $E[\sigma]$ for each source assumption E; the result is a family for $A[\sigma]$. The substitution takes the place of the renaming ρ, and stability under extension takes the place of stability under injective maps. `parameter_witness` and `uniform_raw` use hereditary derivability as in Theorem 7.27, and the proof above follows them. Theorem 7.29 is the persistent cut of `newest_parameter` and the uniform cut of `parameter_witness`. `no_witness` also derives the standalone name $D_{x,v}\gamma_v(x = t)$ in N_ν: while $b = t$ is available and b is the newest parameter, introduce the name, and then discharge the opening of $X(St)$. None of the packages claims a Lean proof of Theorem 7.35; a formalization has to encode the families, or an equivalent protected translation, and cannot assume the weakening of restricted derivations.

7.5.5 The seven-rule diagonal variant

The diagonal calculus is the seven-rule calculus $\mathsf{N}_{\mathrm{diag}}$ of Theorem 3.55: the calculus N_P of Theorem 3.48 for the selector P that holds of the matrices diagonal in y (Theorem 3.12). Its one introduction DI_P takes a witness t and reads its mode off the raw matrix, before the witness is inserted; write $\eta_B(z)$ for $\eta_B^P(z)$. It derives the same judgments as N (Theorem 3.55; Lean: `gprf_iff_nprf`). Let $\mathsf{N}_{\mathrm{diag},\nu}$ be $\mathsf{N}_{\mathrm{diag}}$ with scopes and with $t = \nu(\Pi)$:

$$\frac{\Pi, z;\ \Gamma, B(\nu(\Pi), \eta_B(z)) \vdash B(\nu(\Pi), Sz)}{\Pi; \Gamma \vdash D_{x,y}B(x,y)}.$$

The matrix selects the mode and the scope selects the witness. No guard, axiom or auxiliary syntax is added.

Theorem 7.38 (No witness in the seven-rule calculus)**.** $\mathsf{N}_{\mathrm{diag},\nu}$, $\mathsf{N}_{\mathrm{diag}}$ *and* N *derive the same judgments.*

Proof. Every rule of $\mathsf{N}_{\mathrm{diag},\nu}$ is a rule of $\mathsf{N}_{\mathrm{diag}}$. For the converse, Theorems 7.25 and 7.27 and Theorems 7.26, 7.28 and 7.29 are taken for $\mathsf{N}_{\mathrm{diag},\nu}$, with item (vi) of Theorem 7.28 omitted. Their proofs are unchanged, because the diagonal shape is preserved and reflected by the substitution of terms for parameters (Theorem 3.55(iii)). The following steps differ from the proof of Theorem 7.35; they are the steps of Theorem 3.52 carried out for families of derivations, since the structural rules of $\mathsf{N}_{\mathrm{diag}}$ are not available under the restriction.

Closure by cases. Let $\Pi, z; \Gamma, B(0) \vdash B(Sz)$ and r not free in B. If B is not diagonal, the introduction of $D_{r,y}B(y)$ is by cases and Theorem 7.30 applies. If B is diagonal, $B(0)$ is derivable in every scope and context by a stable derivation; this is condition (S1) of Theorem 3.55(i). Theorem 7.29 removes the assumption $B(0)$ from the given premise, and Theorem 7.4, which holds for $\mathsf{N}_{\mathrm{diag},\nu}$ with the same proof, adds the assumption $B(z)$. The result is the premise of the introduction by progress. In both cases DE yields $Q_yB(y)$.

Equation induction. There is no rule QI. For a term $s(y)$ let $L_s(y) := I(y, y;\, u.\, s(y))$ with u not free in s, so that $L_s(y) = L_t(y)$ is the diagonal lift $\Delta_y E$ of Theorem 3.12 for $E(y) \equiv (s(y) = t(y))$; then $L_s(0) = 0$ and $L_s(Sz) = s(Sz)$ by computation. The decoder equation

$$\mathrm{sel}\big(y, s(0), L_s(y)\big) = s(y) \tag{7.5}$$

holds at $y = 0$ by I0 and at $y = Sz$ by two successor computations; closure by cases and QE give it at every term, in every scope and context. Let $E(y) \equiv (s(y) = t(y))$ and suppose $\Pi, z; \Gamma, E(z) \Vdash E(Sz)$. Then $\Pi; \Gamma \Vdash Q_yE(y)$. Indeed, let (Π', ρ, Δ) be given, and drop ρ from the notation. By closure by cases it suffices to derive $E(Sz)$ in the scope Π', z from $\Delta_0 = \Delta, E(0)$. The matrix $L_s(y) = L_t(y)$ is diagonal. Introduce $D_{r,y}(L_s(y) = L_t(y))$ with r unused; this is by progress, and the premise has a further eigenvariable z_1 and the context $\Delta_0, L_s(z_1) = L_t(z_1)$, denoted Θ. By (7.5) for s and for t, the assumption and the base $E(0)$,

$$\Theta \Vdash s(z_1) = \mathrm{sel}(z_1, s(0), L_s(z_1)) = \mathrm{sel}(z_1, t(0), L_t(z_1)) = t(z_1).$$

The step family, at the map $z \mapsto z_1$, gives $\Theta, E(z_1) \Vdash E(Sz_1)$; uniform cut gives $\Theta \Vdash E(Sz_1)$, and computation gives $\Theta \Vdash L_s(Sz_1) = L_t(Sz_1)$. After the introduction, DE adds the closure $Q_y(L_s(y) = L_t(y))$. Its instance at 0 is derivable, QE at Sz gives $L_s(Sz) = L_t(Sz)$, and computation gives $E(Sz)$. With QE this is equation induction for hereditary families. Every rule of the fragment $\mathcal{E}$ therefore preserves $\Vdash$, and by induction on a derivation of $\mathcal{E}$ the judgments (7.3) hold with $\Vdash$. The construction uses closure by cases, computations and the equality rules; it uses no predecessor and no chosen witness.

Leibniz rule. In the case $F(v) = D_{x,y}B(v, x, y)$ of Theorem 7.31 the target is introduced at the opened parameter b. If its matrix is not diagonal, the proof is unchanged. If it

is diagonal, the premise has the assumption $B'(t', b, z)$ in place of $B'(t', b, 0)$; the latter is derivable outright, and the same three steps give $B'(t', b, Sz)$ without the assumption.

Packet. The matrices $M_y(x, v)$ have outer iterators with the seed 0, and the diagonal shape requires the seed to be the bound variable; the matrix of $D_{r,y}X(y)$ is not an equation. All introductions in the proof of Theorem 7.33 are therefore by cases, and the proof is unchanged.

Translation. The induction is on a derivation of $\mathsf{N}_{\mathrm{diag}}$. The mode of an introduction is the same in source and target, because the alias changes the selected witness and not the raw matrix. In the mode by progress, step (a) of the proof of Theorem 7.35 transports $B'(b, z')$ to $B'(t', z')$, step (b) uses the induction hypothesis with the assumption $B(t, z)$, and steps (c) and (d) are unchanged.

The equivalence with N is the cited one for $\mathsf{N}_{\mathrm{diag}}$. □

7.5.6 One bit

Let R_ν be the calculus R of Section 7.4.3 with the variable witnesses restricted to $\nu(\Pi)$ in a nonempty scope; the literal 0 remains available, and in the empty scope it is the only witness. For an implementation both choices are made total. The calculus R_β has, in place of RDI, the rule

$$\frac{\Pi, z;\ \Gamma, B(t_\beta, \eta) \vdash B(t_\beta, Sz)}{\Pi; \Gamma \vdash D_{x,y}B(x, y)}\ \textsc{Bit-DI}$$

$$t_\beta = \begin{cases} 0, & \beta = 0, \\ \nu(\Pi), & \beta = 1, \end{cases} \qquad \eta = \begin{cases} z, & \beta = 0 \text{ and } B \text{ is a literal equation}, \\ 0, & \text{otherwise}, \end{cases}$$

whose only datum is the bit $\beta \in \{0, 1\}$. The mode is selected by the bit and not by an inspection of the selected term: with $\beta = 1$ in the empty scope the rule is an introduction by cases at 0, also for an equation.

Theorem 7.39 (Zero or the newest parameter)**.** *R_ν, R_β and N derive the same judgments.*

Proof. Every rule of R_ν is an instance of Bit-DI: an introduction at the literal 0 is the bit 0, with the same mode, and an introduction at the newest parameter is the bit 1. Every instance of Bit-DI is an introduction by cases or an introduction by progress at the witness 0, and both are rules of N, the second by Theorem 1.16. It remains to show $\mathsf{N} \subseteq \mathsf{R}_\nu$. Hereditary derivability and Theorems 7.28 and 7.29 are taken for R_ν; a renaming preserves the mode of an introduction, and the proof of uniform cut is unchanged because each node keeps its scope. The following steps differ from the proof of Theorem 7.35.

Closure by progress and equation induction. If $\Pi, z; \Gamma, E(z) \Vdash E(Sz)$, then $\Pi; \Gamma \Vdash Q_y E(y)$, by the introduction of $D_{x,y}E(y)$ at the literal 0 in each receiving scope. This replaces Theorem 7.28(vi), and with QE it gives (7.3) with $\Vdash$ by induction on a derivation of $\mathcal{E}$.

Closure by cases. Let $\Pi, z; \Gamma, B(0) \Vdash B(Sz)$ and let (Π', ρ, Δ) be given. The formula $D_{r,w}(0 = 0)$ is derivable at the witness 0 by progress. Open it at a fresh b, which is then the newest parameter, and introduce $D_{r,y}\rho B(y)$ at b, by cases. Its premise is the member of the given family in the scope Π', b, z'; it is not obtained by renaming a derivation. Eliminate the two unused coordinates.

Leibniz rule. As Theorem 7.31. The reintroduction is at the parameter just opened, which is the newest, and it is by cases.

Packet. With T as in Theorem 7.22, $T(0)$ is derivable in every scope by progress at 0. For $Q_v T(v)$, expand the closure by cases: open $D_{r,w}(0 = 0)$ at a fresh b, introduce the dummy $D_{r,v}T(v)$ at b, and in its premise, where the eigenvariable z has just been appended and is the newest parameter although b was added before, introduce $T(Sz)$ at z by cases; the instance asked for holds by computation. Eliminate the two unused coordinates. QE gives $T(t)$ for

every t, in every scope and context. It is not claimed that $T(Sz)$ has a hereditary derivation. Opening $T(St)$ at a fresh b gives $b = t$ hereditarily, as in the proof of Theorem 7.24.

Translation. The induction is on a derivation of N. QI is closure by progress. For DI at t open $T(S\rho t)$ at a fresh b and introduce at b, by cases; steps (a) to (d) of the proof of Theorem 7.35 are unchanged.

Finally $\mathsf{N} \subseteq \mathsf{R}_\nu \subseteq \mathsf{R}_\beta \subseteq \mathsf{N}$ on derivable judgments. □

7.5.7 The five-family clause calculus

The checker of `parameter_witness` uses the raw clause presentation without the binder Q, and the restriction is proved directly for it: a translation of N_ν into clauses may insert parameters and need not preserve which parameter is the newest. Formulas are equations and $D_{x,y}B$. A clause $[A;\, y.B(y)]$ reads $A \to \forall y\, B(y)$ (Section 1.9); an assumption F is the clause $\mathsf{a}(F) := [0 = 0;\, v.F]$. Contexts are lists of clauses. The calculus C_β has five rule families:

$$\frac{}{\Pi;\Gamma \vdash t = t}\ \textsc{Refl} \qquad \frac{\Pi;\Gamma \vdash s = t \qquad \Pi;\Gamma \vdash b = r}{\Pi;\Gamma \vdash r = \Phi}\ \textsc{Eval} \qquad \frac{\Pi;\Gamma \vdash A}{\Pi;\Gamma \vdash B(t)}\ \textsc{Use}\ ([A;\, y.B(y)] \in \Gamma)$$

$$\frac{\Pi;\Gamma \vdash D_{x,y}B \qquad \Pi, b;\ \Gamma, [B(b,0);\, y.B(b,y)] \vdash A}{\Pi;\Gamma \vdash A}\ \textsc{Elim} \qquad \frac{\Pi, z;\ \Gamma, \mathsf{a}(B(t_\beta, \eta)) \vdash B(t_\beta, Sz)}{\Pi;\Gamma \vdash D_{x,y}B}\ \textsc{Bit-Intro}$$

Eval applies in exactly two cases, decided by literal identity of terms: b is $I(0, a;\, u.\, h)$, s is a and Φ is t; or b is $I(Sc, a;\, u.\, h)$, s is $I(c, a;\, u.\, h)$ and Φ is $h[t/u]$. In Bit-Intro, t_β and η are as in Bit-DI. There are no other axioms or rules; an assumption is used by Refl at 0 and Use. The comparison calculus C has the same rules, except that its introduction takes an arbitrary term t over Π and is by progress exactly when B is a literal equation and t is the literal 0.

The equality rules are derived from Refl and Eval alone, as in the packages on two-case evaluation. Eval on two reflexivities gives the two computation equations. With $J = I(0, s;\, u.\, u)$, Eval on $s = r$ and $J = J$ gives $J = r$, and Eval on $s = t$ and $J = r$ gives $r = t$; this joining rule yields symmetry and transitivity. For a term template $h(v)$ let $J = I(0, s;\, u.\, h(u))$ and $K = I(1, s;\, u.\, h(u))$: from $s = t$, Eval gives $J = t$ and then, with $K = K$, $K = h(t)$; the same with $s = s$ gives $K = h(s)$, and joining gives $h(s) = h(t)$. These derivations contain no Elim and no introduction and are valid in every scope.

Theorem 7.40 (The one-bit clause calculus)**.** *C_β and C derive the same judgments $\Pi;\Gamma \vdash A$, for all clause contexts Γ.*

Proof. From C_β to C. An introduction with the bit 0 is the introduction of C at the literal 0, with the same mode. With the bit 1 in a nonempty scope it is the introduction of C at a variable, by cases. With the bit 1 in the empty scope it is by cases at 0. For a compound matrix this is the introduction of C at 0. For an equation matrix, C introduces at the term $w = I(0, 0;\, u.\, 0)$, which is not the literal 0, so the mode is by cases; the assumption $B(w, 0)$ is transported along $w = 0$ to $B(0, 0)$, the given premise is applied, and its conclusion is transported back to $B(w, Sz)$. The calculus C is closed under injective renaming and weakening and admits the cut of an assumption, by the direct inductions, so this composition is a derivation.

From C to C_β. Write $\Pi;\Gamma \Vdash_c A$ if for every injective map ρ from Π into a scope Π' and every clause context Δ over Π' that contains $\rho\Gamma$, C_β derives $\Pi';\Delta \vdash \rho A$. The renaming acts on the antecedent and on the body of each clause, and the receiving context retains the clauses themselves.

Cut of an assumption. If $\Pi;\Gamma, \mathsf{a}(A) \vdash F$ and $\Pi;\Gamma \Vdash_c A$, then $\Pi;\Gamma \vdash F$; this is proved as Theorem 7.29. A Use of the clause $\mathsf{a}(A)$ is replaced by the member of the family in the scope

of that leaf, its premise $0 = 0$ being dropped; every other USE keeps its clause. ELIM and BIT-INTRO add their clauses in both contexts. No clause other than an assumption is cut.

Rules. REFL, EVAL and USE hold with $\Vdash_c$, the clause of a USE being a member of the renamed context; ELIM holds with $\Vdash_c$ by sending its eigenvariable to a fresh parameter of the receiving scope and retaining the new clause there.

Closures. Write $q_y A(y)$ for $D_{x,y}A(y)$ with x not free in A. Its introduction by cases is BIT-INTRO with the bit 1: the witness is unused, and the fallback in the empty scope is still by cases. For an equation E its introduction by progress is the bit 0. Its elimination is ELIM followed by USE of the clause $[A(0);\, y.A(y)]$. With hereditary premises these three operations hold for $\Vdash_c$, as in Theorem 7.30. In particular equation induction holds for $\Vdash_c$, and (7.3) holds with $\Vdash_c$ by induction on a derivation of $\mathcal{E}$.

Leibniz rule. If $\Pi;\Gamma \Vdash_c s = t$ and $\Pi;\Gamma \Vdash_c F(s)$, then $\Pi;\Gamma \Vdash_c F(t)$, by induction on F. For an equation this is the equality rules. For $D_{x,y}B(s,x,y)$, open the formula at a fresh b in the receiving scope, retaining the clause $[B(s,b,0);\, y.B(s,b,y)]$, and introduce the target with the bit 1, which selects b. From the assumption $B(t,b,0)$ the induction hypothesis gives $B(s,b,0)$, USE of the retained clause at Sz gives $B(s,b,Sz)$, and the induction hypothesis gives $B(t,b,Sz)$; the cut of an assumption composes these steps. ELIM discharges b.

Packet. With T as in Theorem 7.22, $T(0)$ is introduced with the bit 0 in every scope: the matrix is an equation, the introduction is by progress, and the instance asked for holds by computation. For $q_v T(v)$ use the bit 1 on the outer formula; in its premise the eigenvariable z is the newest parameter, and $T(Sz)$ is introduced with the bit 1, which selects z. The construction is repeated in every scope. ELIM and USE give $\Pi;\Gamma \Vdash_c T(t)$ for every t. Opening $T(St)$ at a fresh b and using its clause at 1 gives $Sb = St$, hence $b = t$, with $\Vdash_c$ from the opened clause context.

Translation. By induction on a derivation of C its conclusion is shown to be hereditarily derivable. REFL, EVAL, USE and ELIM are treated above. An introduction by progress at the literal 0 is the bit 0. For an introduction by cases at a term t, open $T(S\rho t)$ at a fresh b, retain $\mathsf{a}(b = \rho t)$ for the moment, and introduce the formula with the bit 1. In the premise the Leibniz rule turns the assumption $B(b,0)$ into $B(\rho t,0)$. The family given by the induction hypothesis is taken at the map that keeps the images of the source parameters, sends the source eigenvariable to the new one and skips b; the cut of an assumption supplies its assumption, and the Leibniz rule gives $B(b,Sz)$. After the introduction, the cut of an assumption replaces $\mathsf{a}(b = \rho t)$ by its hereditary derivation from the opened packet, and ELIM discharges b. □

Corollary 7.41. *The ordinary realizations of* R_ν, R_β, $\mathsf{N}_{\mathrm{diag},\nu}$ *and* C_β *are bi-interpretable and equiconsistent with* iPRA*, with whole numerical domains, actual equality and the identity graph in both numerical composites.*

Proof. Each calculus derives the same judgments, on the same terms and formulas, as a calculus for which the bridge of Theorem 1.78 is established: N for the first three, and the unrestricted clause calculus C for the last. The translations and the derivations of both formula round trips transfer unchanged. □

7.6 The raw guarded calculus

This section draws on `deterministic_introduction` for the naming exception, and on `uniform_raw` and the guarded companion of `newest_parameter` for the calculus without it.

7.6.1 The calculus and its three introductions

Terms are the raw terms of Section 1.1; no computation is built into syntactic identity or into substitution. A *canonical formula* is an equation or a formula $D_{x,y}B$ in which B is canonical and, if B is an equation, it has the literal shape

$$I(y, 0;\, u.\, h) = I(y, 0;\, v.\, k). \tag{7.6}$$

The bodies h and k may contain x, y and parameters. An equation at the root of a formula is unrestricted. Substitution of terms for parameters preserves canonical formulas. Contexts are lists of clauses $[A;\, y.B(y)]$ of canonical formulas, with y not free in A. Put $H := (I(0, 0;\, u.\, 0) = 0)$, an instance of I0; an assumption A is the clause $\mathsf{a}(A) := [H;\, v.A]$ and is used by Use with one computation.

The calculus G has the two computation equations as axioms in every context and four inference rules:

$$\frac{\Pi; \Gamma \vdash A}{\Pi; \Gamma \vdash B(t)}\ \textsc{Use}\ \big([A;\, y.B(y)] \in \Gamma\big) \qquad \frac{\Pi; \Gamma \vdash s = t \qquad \Pi; \Gamma \vdash F(s)}{\Pi; \Gamma \vdash F(t)}\ \textsc{Tr}$$

$$\frac{\Pi; \Gamma \vdash D_{x,y}B \qquad \Pi, b;\ \Gamma, [B(b, 0);\, y.B(b, y)] \vdash A}{\Pi; \Gamma \vdash A}\ \textsc{Open} \qquad \frac{\Pi, z;\ \Gamma, \mathsf{a}(B(t, \eta_B(z))) \vdash B(t, Sz)}{\Pi; \Gamma \vdash D_{x,y}B}\ \textsc{Intro}$$

with $\eta_B(z) = z$ if B is an equation and $\eta_B(z) = 0$ if B is compound. In Tr, $F(v)$ is a canonical formula template. The mode of Intro depends on the root of the matrix and not on the witness. There is no primitive hypothesis rule, reflexivity, weakening, cut, coverage or induction rule. The ordinary reading of $D_{x,y}B$ is that of Section 1.2; the comparison of G with N, by padding equation matrices, and hence the bridge to iPRA, is established for G in `guarded_core` and is inherited here.

Definition 7.42 (Three selections of the witness)**.** In G the witness t of Intro is an arbitrary term over Π. In G_ν it is $\nu(\Pi)$. In G_d it is

$$\omega(B, \Pi) := \begin{cases} r, & B \text{ is literally } \mathrm{Nm}_r \text{ with } r \text{ over } \Pi, \\ \nu(\Pi), & \text{otherwise.} \end{cases}$$

A match with a naming matrix has priority, and it is literal up to the names of bound variables: r contains neither x, y nor the bound variables of the two iterators. It is not a match up to provable equality.

Every naming matrix has the shape (7.6). The derivations of G_ν and of G_d are derivations of G once the selected witnesses are written in. In all three calculi reflexivity and symmetry are derived from I0 and Tr: with $q = I(0, r;\, u.\, 0)$, the equation $q = r$ used as both premises of Tr in the template $v = r$ gives $r = r$, and from $s = t$ the transport of $s = s$ in the template $v = s$ gives $t = s$.

7.6.2 The naming exception

In G_d the names are immediate, and the work is structural: weakening to a larger scope can change the selected witness, and the general cut needs the weakening of the inserted derivation. The order of the argument is a restricted cut, then weakening, then the general cut, then completeness.

Lemma 7.43 (Algebraic derivations and restricted cut)**.** *Call a derivation* algebraic *if it uses only the computation axioms,* Use *and* Tr*.*

(i) *An algebraic derivation of* $\Pi;\Gamma \vdash A$ *yields an algebraic derivation of* $\Pi';\Delta \vdash \rho A$ *for every injective map* ρ *from* Π *into* Π' *and every clause context* Δ *over* Π' *into which* $\rho\Gamma$ *is embedded.*

(ii) *If* $\Pi;\Gamma \vdash A$ *has an algebraic derivation and* $\Pi;\Gamma,\mathsf{a}(A) \vdash F$ *in* G_d*, then* $\Pi;\Gamma \vdash F$ *in* G_d*.*

Proof. (i) By induction on the derivation; its rules have no choice that depends on the scope and no eigenvariable. (ii) By induction on the derivation of F. A Use of the clause $\mathsf{a}(A)$ is replaced by the algebraic derivation of A, moved to the scope and context of that leaf by (i). At Intro and Open the induction continues in the extended branch. Deleting a clause does not change the scope, so every Intro selects the same witness before and after. No weakening of a derivation that contains an Intro is used. □

Lemma 7.44 (Every term has a name)**.** *For every term* r *over* Π*,* G_d *derives* $\Pi;\Gamma \vdash D_{x,y}\mathrm{Nm}_r$*. If a clause context contains* $[\mathrm{Nm}_r(b,0);\, y.\mathrm{Nm}_r(b,y)]$*, then* $b = r$ *and* $r = b$ *have algebraic derivations from it.*

Proof. The rule selects r by the naming alternative. The matrix is an equation, so the premise asks for $\mathrm{Nm}_r(r,Sz)$ from $\mathrm{Nm}_r(r,z)$; the conclusion is literally $\mathrm{sel}(Sz,0,r) = \mathrm{sel}(Sz,0,r)$ and is a reflexivity. For the second statement, both sides of $\mathrm{Nm}_r(b,0)$ equal 0 by I0, so the antecedent of the clause is derivable; Use at 1 gives $\mathrm{sel}(1,0,b) = \mathrm{sel}(1,0,r)$, and IS with Tr gives $b = r$; symmetry gives $r = b$. □

Theorem 7.45 (Weakening by naming the old selection anew)**.** *Let* $\Pi;\Gamma \vdash A$ *be derivable in* G_d*. For every injective map* ρ *from* Π *into a scope* Π' *and every clause context* Δ *over* Π' *into which* $\rho\Gamma$ *is embedded,* $\Pi';\Delta \vdash \rho A$ *is derivable in* G_d*, by an effective construction.*

Proof. By induction on the height of the given derivation, for all ρ and Δ. The computation axioms, Use and Tr are immediate. At Open, choose a fresh parameter of the receiving scope, extend ρ and the embedding of clauses, and apply the induction hypothesis to the two premises.

Consider an Intro with the matrix B and the selected witness w, and let $B' = \rho B$, $w' = \rho w$. A match with a naming matrix is preserved and reflected by ρ: a bound variable does not become a parameter or conversely, and a parameter does not become a numeral or a compound term. If $B = \mathrm{Nm}_r$, the witness selected in the receiving scope is $\rho r = w'$, and the induction hypothesis for the premise gives the rule directly. The same holds if B is not a naming matrix and $\nu(\Pi') = w'$.

Suppose B' is not a naming matrix and $\nu(\Pi') \neq w'$. Derive $D_{x,y}\mathrm{Nm}_{w'}$ by Theorem 7.44 and open it at a fresh parameter e, which is then the newest; its clause gives algebraic derivations of $e = w'$ and $w' = e$. The formula $D_{x,y}B'$ will be introduced in the scope Π', e, where the rule selects e. Let v be the eigenvariable of that introduction and put

$$A_\mathrm{old} = B'(w', \eta_{B'}(v)), \qquad A_\mathrm{new} = B'(e, \eta_{B'}(v)).$$

Apply the induction hypothesis to the premise of the given Intro, with ρ extended by sending its eigenvariable to v, and with the receiving context consisting of Δ, the clause of the name, $\mathsf{a}(A_\mathrm{old})$ and $\mathsf{a}(A_\mathrm{new})$. This is legitimate although e and v are new parameters of the receiving scope, because the induction statement quantifies over all injective maps of the smaller derivation. The result derives $B'(w', Sv)$. From A_new, Tr along $e = w'$ gives A_old; this derivation is algebraic. Theorem 7.43(ii) removes the clause $\mathsf{a}(A_\mathrm{old})$. Tr along $w' = e$ gives $B'(e, Sv)$. Apply Intro, which selects e, and discharge the name by Open; the conclusion does not contain e. If the old selection was 0 in an empty scope, the name is that of 0.

Every use of the induction hypothesis is for a proper subderivation of the given derivation. The naming and transport derivations built here are never fed back into the induction, and

the restricted cut was established before. The construction is therefore well founded and uses neither the general cut nor weakening. □

Corollary 7.46 (Cut of an assumption)**.** *If* $\Pi;\Gamma \vdash A$ *and* $\Pi;\Gamma,\mathsf{a}(A) \vdash F$ *in* G_d*, then* $\Pi;\Gamma \vdash F$ *in* G_d.

Proof. By induction on the derivation of F. A Use of $\mathsf{a}(A)$ is replaced by the derivation of A, moved to the scope and context of that leaf by Theorem 7.45. Deleting a clause leaves the scope unchanged, so the introductions of the receiving derivation keep their selections. □

Theorem 7.47 (The introduction with a naming exception is complete)**.** *For every scope* Π*, canonical clause context* Γ *and canonical formula* A*,*

$$\Pi;\Gamma \vdash_{\mathsf{G}_\mathrm{d}} A \quad\Longleftrightarrow\quad \Pi;\Gamma \vdash_{\mathsf{G}} A.$$

Proof. From left to right the selected witness is a witness of Intro in G. For the converse induct on the derivation in G. At an Intro whose matrix is a naming matrix Nm_r, discard the supplied witness and the premise and use Theorem 7.44. At any other Intro, with the witness t, translate the premise by the induction hypothesis, derive $D_{x,y}\mathrm{Nm}_t$ and open it at a fresh b. In the context with the opened name and the assumption $B(b,\eta_B(z))$, Tr along $b = t$ gives $B(t,\eta_B(z))$. Theorems 7.45 and 7.46 insert the translated premise and give $B(t,Sz)$, and Tr along $t = b$ gives $B(b,Sz)$. The newest parameter of the scope is b and B is not a naming matrix, so the rule selects b. Close the introduction and the opening of the name. □

Remark 7.48 (Why the qualifications matter)*.* The simulation changes the witness from t to b and must not change the assumption of the premise. In G the mode depends on the root of the matrix only. A rule that selects progress exactly when the witness is the literal 0 does not fit this argument as it stands: at $t = 0$ it has the assumption $B(0,z)$ and at b the assumption $B(b,0)$; the calculus R of Section 7.4.3 is treated by recovering the two modes separately. The diagonal selection of Section 7.5.5 is independent of the witness as well. The recognition of a naming matrix is a fixed syntactic match. Replacing it by the condition that some instance of the matrix names the witness would need additional data or a search, and a rule about a provable naming instance would reintroduce a premise. The proof uses transport in arbitrary canonical formulas: the donor of the restricted cut converts $B(e,\eta)$, which may be compound. Through the comparison, substitution is admissible in G_d as a transformation of derivable judgments; a certificate is not claimed to remain valid after a substitution.

7.6.3 Uniform introduction: the stable fragment

In G_ν the name $D_{x,y}\mathrm{Nm}_r$ is no longer immediate, and the names are generated as in Section 7.4. The guarded calculus has no rule for equation induction, so the predecessor has to be obtained first, in a part of the calculus that is insensitive to the selected witness.

Definition 7.49. An Intro of G_ν with the matrix $B(x,y)$ is *dummy* if x is not free in B; a derivation is *stable* if all its introductions are dummy.

Lemma 7.50 (Stability)**.** *Stable derivations of* G_ν *are closed under the substitution of terms over a scope* Π' *for the parameters, with any clause context over* Π' *that contains the substituted clauses, and under the cut of an assumption by a stable derivation.*

Proof. As Theorem 7.26. The premise of a dummy introduction does not depend on the witness; a substitution does not change whether the matrix is an equation, and it preserves the shape (7.6). □

For an equation E let the *padding* be $L(E) := D_{r,v}\gamma_v(E)$ with r, v not free in E; it is canonical. For equations $U \equiv (s = t)$ and $V \equiv (p = q)$ let

$$U \Rightarrow_g V := D_{r,w}L\big(\mathrm{sel}(w, s, p) = \mathrm{sel}(w, t, q)\big),$$

a canonical formula with a compound matrix and an unused coordinate r.

Lemma 7.51 (Padding, implication and cases without names)**.** *The following hold for stable derivations of* G_ν*, in every scope and context.*

(i) *E and $L(E)$ are interderivable.*

(ii) *If $\Pi; \Gamma, \mathsf{a}(U) \vdash V$, then $\Pi; \Gamma \vdash U \Rightarrow_g V$; and $U \Rightarrow_g V$ together with U gives V.*

(iii) *If $\Pi; \Gamma \vdash F(0)$ and $\Pi, z; \Gamma \vdash F(Sz)$, then $\Pi; \Gamma \vdash F(t)$ for every term t; here $F(y)$ is a compound formula or an equation.*

(iv) *If E is an equation and t a term over Π, and $\Pi; \Gamma, \mathsf{a}(t = 0) \vdash E$ and $\Pi, z;\ \Gamma, \mathsf{a}(t = Sz) \vdash E$, then $\Pi; \Gamma \vdash E$.*

Proof. (i) From E: introduce $L(E)$. The matrix $\gamma_v(E)$ is an equation, so the premise asks for $\gamma_{Sz}(E)$ from $\gamma_z(E)$; it follows from E by IS and Tr, and the assumption is unused. From $L(E)$: open it at a fresh b; the antecedent $\gamma_0(E)$ of the clause holds by I0, Use at 1 gives $\gamma_1(E)$, and IS gives E, which does not contain b.

(ii) Write row_w for $\mathrm{sel}(w, s, p) = \mathrm{sel}(w, t, q)$. The matrix is compound, so the introduction is by cases and asks for $L(\mathrm{row}_{Sz})$ from $L(\mathrm{row}_0)$. By (i) and I0 the assumption gives U; the cut of the given derivation gives V; IS gives row_{Sz}, and (i) pads it. For the elimination open the formula at a fresh b, derive $L(\mathrm{row}_0)$ from U by I0 and (i), use the clause at 1, and unpad and compute to obtain V.

(iii) For a compound F, introduce $D_{r,y}F(y)$ with r unused; it is by cases, and its premise is the given derivation of $F(Sz)$ weakened by the assumption $F(0)$. Open it at a fresh b and use the clause $[F(0); y.F(y)]$ at t, the antecedent being the given derivation moved to the scope Π, b by Theorem 7.50. For an equation $E(y)$ apply this to $L(E(y))$ and use (i).

(iv) Let $F(y)$ be $(t = y) \Rightarrow_g E$. By (ii) the hypotheses give $F(0)$ and $F(Sz)$, by (iii) $F(t)$, and by (ii) with the reflexivity $t = t$ the equation E. □

Lemma 7.52 (Equation induction in the stable fragment)**.** *Let $E(y) \equiv (s(y) = t(y))$. If $\Pi; \Gamma \vdash E(0)$ and $\Pi, z;\ \Gamma, \mathsf{a}(E(z)) \vdash E(Sz)$ have stable derivations in G_ν, then so has $\Pi; \Gamma \vdash E(a)$ for every term a over Π. Consequently every derivation of the equational fragment $\mathcal{E}$ is translated into a stable derivation of G_ν, and for all terms r, s*

$$\Pi; \Gamma \vdash \mathrm{pred}(Sr) = r, \qquad \Pi;\ \Gamma, \mathsf{a}(Sr = Ss) \vdash r = s$$

by stable derivations.

Proof. Let $G(y) := \gamma_y(E(y))$, an equation of the shape (7.6). First, $E(z)$ is derivable from $G(z)$: by Theorem 7.51(iv) with the term z, in the branch $z = 0$ transport the base $E(0)$ along $0 = z$, and in the branch $z = Sw$ transport $G(z)$ to $G(Sw)$, compute the two gates to obtain $E(Sw)$, and transport back. The step then gives $E(Sz)$, by Theorem 7.50, and computation gives $G(Sz)$. This is the premise of the introduction of $D_{r,y}G(y)$ with r unused, which is by progress. Open it at a fresh b; the antecedent $G(0)$ of its clause holds by computation, and Use at a gives $G(a)$. A second application of Theorem 7.51(iv), with the term a and with $E(0)$ in the branch $a = 0$, gives $E(a)$. All introductions are dummy.

For the translation of $\mathcal{E}$, an assumption E is the clause $\mathsf{a}(E)$, the equality rules are instances of Tr and reflexivity, and induction is the first statement; the instances of earlier lemmas that an equational derivation uses are available by Theorem 7.50. The last statement follows from Theorem 1.48. □

7.6.4 Uniform introduction: names and completeness

Hereditary derivability $\Pi;\Gamma \Vdash A$ is defined for G_ν as in Theorem 7.27, with clause contexts. The computation axioms, Use and Tr hold with $\Vdash$; Open holds with $\Vdash$ by allocating its parameter in each receiving scope; a stable derivation is a member of a hereditary family; and, as in Theorem 7.29,

$$\Pi;\Gamma \Vdash A \quad \text{and} \quad \Pi;\Gamma, \mathsf{a}(A) \vdash B \quad \text{imply} \quad \Pi;\Gamma \vdash B. \tag{7.7}$$

In the proof of (7.7) the donor is generated at each Use of the removed clause in the scope and context of that leaf, the other clause addresses are adjusted, and every other node keeps its scope and therefore its selected witness. The hypotheses of the donor must remain available after the deletion.

The packet $X(y)$ of Theorem 7.18 is canonical: both sides of its matrix are iterators with the counter v and the seed 0.

Lemma 7.53 (The packet in the guarded calculus)**.** *In G_ν, for every scope Π, clause context Γ and term t over Π: $\Pi;\Gamma \vdash D_{a,r}X(r)$ with a unused, and $\Pi;\Gamma \vdash X(St)$.*

Proof. $X(0)$ is introduced at whatever witness the scope selects. Its matrix is an equation, the introduction is by progress, and the instance asked for computes to $0 = 0$ whatever the witness and the assumption are. The matrix of $D_{a,r}X(r)$ is compound, so its introduction is by cases and asks for $X(Sz)$ with a new eigenvariable z. In that branch the scope ends in z, so the introduction of $X(Sz)$ selects z, and the instance asked for computes to $Sz = Sz$. The assumption $X(0)$ of the outer introduction is unused. Open $D_{a,r}X(r)$ at a fresh parameter; its clause is $[X(0);\, r.X(r)]$. Derive $X(0)$ again in the extended scope and use the clause at St. The result does not contain the opened parameter, and Open discharges it. □

The lemma is about the whole construction, which allocates z anew in each receiving scope. A separate derivation of $X(Sz)$ is not claimed to remain a derivation after z ceases to be the newest parameter.

Lemma 7.54 (The alias in the guarded calculus)**.** *If a clause context over a scope contains $[M_{St}(b,0);\, v.M_{St}(b,v)]$, then $b = t$ is hereditarily derivable from it.*

Proof. The antecedent holds by computation. Use at 1 and the successor computations give $Sb = St$, and Theorem 7.52 gives $b = t$. The derivation is stable. □

Theorem 7.55 (Uniform introduction on raw terms)**.** *For every scope Π, canonical clause context Γ and canonical formula A,*

$$\Pi;\Gamma \vdash_{\mathsf{G}_\nu} A \iff \Pi;\Gamma \vdash_{\mathsf{G}} A \iff \Pi;\Gamma \vdash_{\mathsf{G}_\mathrm{d}} A.$$

The naming exception of G_d is not needed.

Proof. The second equivalence is Theorem 7.47, and $\mathsf{G}_\nu \subseteq \mathsf{G}$ is immediate. For $\mathsf{G} \subseteq \mathsf{G}_\nu$ it is shown by induction on a derivation of G that $\Pi;\Gamma \Vdash A$. The computation axioms, Use and Tr are copied with their terms renamed. For Open, allocate a parameter in the receiving scope and apply the induction hypothesis to the branch with the extended map.

Consider an Intro at a term t with the matrix B. The induction hypothesis is $\Pi, z;\ \Gamma, \mathsf{a}(B(t, \eta_B(z))) \Vdash B(t, Sz)$. Let (Π', ρ, Δ) be given and write t', B' for the images. Derive $X(St')$ by Theorem 7.53 and open it at a fresh b. Introduce $D_{x,y}B'$ in the scope Π', b; the rule selects b and then allocates its eigenvariable z'. Let Θ be the scope Π', b, z' with the context consisting of Δ, the clause of the packet and $\mathsf{a}(B'(b, \eta_{B'}(z')))$. By Theorem 7.54, $\Theta \Vdash b = t'$, and Tr gives $\Theta \Vdash B'(t', \eta_{B'}(z'))$. The induction hypothesis at the map $\rho[z \mapsto z']$, with (7.7), gives

$\Theta \Vdash B'(t', Sz')$, and Tr along $t' = b$ gives $\Theta \Vdash B'(b, Sz')$. Its member at the identity is the premise of the introduction. The conclusion does not contain b, and Open discharges it. The two modes agree because η_B depends on the root of the matrix, which neither ρ nor the change of witness alters. All uses of the induction hypothesis are for proper subderivations, and the packet and the alias were constructed before. □

Corollary 7.56 (The literal naming theorem is derived)**.** *For every term t, the formula $D_{x,y}\mathrm{Nm}_t$ has a hereditary derivation in G_ν, without a selection by pattern.*

Proof. Derive $X(St)$, open it at a fresh b and extract $b = t$. Introduce $D_{x,y}\mathrm{Nm}_t$; the selected witness is b. In the premise the alias and the selector computations give $\mathrm{Nm}_t(b, Sz)$, the progress assumption being unused. Discharge the packet. □

By Theorem 7.55 and the inherited comparison of G with N, the ordinary realizations of G_ν and of G_d are bi-interpretable and equiconsistent with iPRA, with whole domains, actual equality and the identity graph in both numerical composites; weakening, renaming, cut and substitution are admissible in G_ν as consequences of the theorem.

Remark 7.57 (Versions). The stable fragment and the route to equation induction through Theorem 7.51(iv) are those of the guarded companion of `newest_parameter`, which states the translation with persistent builders and an arbitrary substitution, as in Theorem 7.37. `uniform_raw` proves instead, by induction on a derivation of $\mathcal{E}$ and for all substitutions σ at once, that $\Gamma\sigma \Vdash E\sigma$. At an induction with the invariant $E(y)$ it regenerates the step at $y = 0$ and at $y = Sw$ from the source derivation, proves $\Phi(y) := G(y) \Rightarrow_g E(Sy)$ at 0 and at Sw, obtains $\Phi(z)$ by cases, and concludes as above; no derivation of the target is substituted into. A technical note of `uniform_raw` also observes that, once the names are available, the proof of Theorem 7.45 applies to G_ν with no case for a naming matrix, which gives weakening and then cut directly. The checkers of `uniform_raw` and of `newest_parameter` for G_ν are obtained from the checker of G_d by deleting the function that selects the witness (Section 7.9).

7.7 A recursor whose predecessor computes

This section draws on `parameter_intro`; the normalization lemma and the extensionality lemma are also in `conditional_copy`.

On iterator terms no term p satisfies $p(Sz) = z$ by computation, and the names of Section 7.4 need the derived predecessor and an equality rule. With a recursor whose body receives the counter, and with terms taken in normal form, the predecessor computes, and the name of every term is obtained with clause specialization alone. This makes the restricted introduction available in a calculus of three rule families in which equality transport is itself part of the discharge rule and is not available beforehand.

7.7.1 Terms in normal form

Raw terms are variables, 0 and $R(n, a;\, i, u.\, h)$, where i and u are bound in h only. The successor is the abbreviation $Sa := R(0, a;\, i, u.\, 0)$. The computation equations are

$$R(0, a;\, i, u.\, h) = Sa, \qquad R(Sn, a;\, i, u.\, h) = h[n/i,\ R(n, a;\, i, u.\, h)/u], \tag{7.8}$$

with simultaneous substitution on the right. To define normal forms without the circular rewrite of $R(0, a;\, i, u.\, 0)$ to itself, take 0, S and R as distinct raw constructors, orient the two equations from left to right, allow reduction in every subterm including bodies, and print the normal constructor Sa as $R(0, a;\, i, u.\, 0)$. The *public terms* are the normal forms, and every substitution performed by a rule is followed by normalization, written nf. Computation is

thus part of the syntax, and the normalizer is part of a checker and of its measured size. The binding notation is a staged first-order language as in Section 1.1: variables range over numbers, and there is no term that evaluates descriptions.

Lemma 7.58 (Normalization)**.** *Every well-sorted term has a unique normal form. Reduction introduces no free variable. For every substitution σ,*

$$\mathrm{nf}(t[\sigma]) = \mathrm{nf}\big(\mathrm{nf}(t)[\mathrm{nf}\circ\sigma]\big),$$

and the same holds for formulas and for templates with one distinguished variable.

Proof. Embed the raw terms into the terms of ground type of the recursor over the natural numbers, with the starting value Sa and the step $(i,u)\mapsto h(i,u)$; no variable is applied as a function. The reducibility argument specializes to this fragment. A recursor with computable arguments and a computable body is computable, by induction on the computable counter and its reducts: at zero it reduces to the computable starting value; at a successor the recursive call is computable at the predecessor and the computable body preserves computability; at a neutral counter the recursor is neutral and its reducts are those of computable proper subterms. Induction on a term, with computable terms substituted for its variables, gives strong normalization; applying the argument under fresh variables covers reductions inside bodies. The two root rules are disjoint and left-linear, and parallel reduction commutes with substitution, so nested reductions and duplicated reducts join; this gives confluence, and with strong normalization the uniqueness of normal forms. A substitution instance of a conversion is a conversion, and normalizing its two sides gives the displayed law. The right sides of the rules contain only variables of their left sides. □

The lemma is a statement about finite terms made from outside. It gives no primitive recursive bound, uniform in the term, on the length of normalization.

Three abbreviations have useful normal forms. Put

$$\begin{aligned} P(n) &:= R(n,0;\, i,u.\, i), & P(0) &\equiv 1, & P(Sn) &\equiv n,\\ C(n,a,b) &:= P\big(R(n,a;\, i,u.\, Sb)\big), & C(0,a,b) &\equiv a, & C(Sn,a,b) &\equiv b, \end{aligned}$$

and, with fresh j,v and simultaneous substitution,

$$\begin{aligned} J(n,a;\, i,u.\, h) &:= P\big(R(n,a;\, j,v.\, S(h[j/i,P(v)/u]))\big),\\ J(0,a;\, i,u.\, h) &\equiv a, \qquad J(Sn,a;\, i,u.\, h) \equiv h[n/i,\ J(n,a;\, i,u.\, h)/u]. \end{aligned}$$

Here $\equiv$ is literal identity of normal forms. P is a left inverse of the successor whose value at 0 is 1; a predecessor in the usual sense is $p(n) := C(n,0,P(n))$, with $p(0)\equiv 0$ and $p(Sn)\equiv n$. The iterator of Section 1.1 is J with the counter coordinate unused. In this section and the next, C takes the place of sel in the gate and in the naming matrix of Theorem 7.10.

7.7.2 Three rule families

Formulas are equations and $D_{x,y}B$. A clause is $[\ ;\ y.B]$ or $[A;\ y.B]$ with y not free in A; the first reads $\forall y\, B$. Every derivation starts from the one clause

$$\mathsf{Rf} := [\ ;\ y.\, y = y],$$

and an assumption A is the clause $[\ ;\ v.A]$. For a matrix B and a term r write $B_r(y)$ for $\mathrm{nf}(B(r,y))$. The calculus P_ν has three rule families.

(i) Use: for a clause of the context and a term t over Π, derive the antecedent of the clause, if it has one, and conclude the normalized body at t.

(ii) INTRO: with $\ell = \nu(\Pi)$ and a fresh z,

$$\frac{\Pi, z;\ \Gamma, [\ ;\ v.B_\ell(\eta)] \vdash B_\ell(Sz)}{\Pi; \Gamma \vdash D_{x,y}B} \qquad \eta = \begin{cases} z, & B_\ell(0) \text{ is literally an equation } r = r, \\ 0, & \text{otherwise.} \end{cases}$$

The instruction supplies no witness, no mode and no proof of the instance at 0.

(iii) DISCHARGE: with a fresh b, the clause $\mathcal{C}_B(b) := [\mathrm{nf}(B(b,0));\ y.\,\mathrm{nf}(B(b,y))]$ and a template $F(v)$,

$$\frac{\Pi; \Gamma \vdash D_{x,y}B \qquad \Pi, b;\ \Gamma, \mathcal{C}_B(b) \vdash s = t \qquad \Pi, b;\ \Gamma, \mathcal{C}_B(b) \vdash \mathrm{nf}(F(s))}{\Pi; \Gamma \vdash \mathrm{nf}(F(t))}$$

provided $\mathrm{nf}(F(t))$ does not contain b. The terms s, t and the template may contain b.

There are no other logical, equality, arithmetic or structural rules; the computations (7.8) belong to the syntax. The designated contradiction is $0 = 1$.

Proposition 7.59 (Soundness)**.** *Every rule of* P_ν *is valid under the ordinary reading in intuitionistic logic with equality, the equations* (7.8)*, numerical coverage,* $0 = 1 \to \bot$ *and induction for literal equations.*

Proof. USE is the application of a clause. DISCHARGE is the elimination of the existential quantifier followed by the substitution of equals in the branch; the condition on $\mathrm{nf}(F(t))$ is the eigenvariable condition, and the rule is sound also when t contains b. For INTRO: if the instance at 0 is not reflexive, the mode is by cases, and coverage combines the retained instance at 0 with the derived instance at successors. If it is reflexive, the matrix is a literal equation, because normalization does not remove a binder of a formula; the instance at 0 is provable and induction for that equation gives all instances. In both cases ℓ is a witness of the existential quantifier. No induction on a quantified formula is used. □

A premise by cases can be used at any witness. If the rule selects progress instead, the discarded assumption at 0 is literally reflexive: derive it by USE of Rf and ignore the progress assumption. In the constructions below this is an operation on *builders*, that is, on constructions that can be carried out again after fresh parameters and clauses have been adjoined; it is not a weakening principle for derivation trees. A builder is called *persistent*.

7.7.3 Constructions that precede equality

Write $q_y A$ for $D_{w,y}A$ with w not free in A. The witness of its introduction is unused, so its instances do not depend on which parameter is the newest. The elimination of $D_{x,y}B$ without substitution is DISCHARGE with $s = t = 0$, the equation $0 = 0$ by USE of Rf, and the constant template whose value is the conclusion. Hence the introduction by cases and the elimination of q are available from the three rules, with the repair of the previous paragraph when the instance at 0 is reflexive; this gives numerical cases for all formulas, with the branches constructed in their own scopes. Neither equality transport nor equation induction is needed for these constructions.

For a formula A let $G_v A$ be A with each side s of its final equation replaced by $C(v, 0, s)$, the binders being kept. Then $G_{Sz}A \equiv A$, and $G_0 A$ is a tower of binders over $0 = 0$, which is derivable in every scope by INTRO and reflexivity, its witnesses being unused. For persistent premises the padding $L(A) := q_y G_y A$ is interderivable with A: one direction ignores its assumption at 0 and uses A, the other uses the closure at 1 with the provable instance at 0. Padding two formulas to the same number of binders permits their selector, sidewise by $C(v, \cdot, \cdot)$, and the implication $q_y\,\mathrm{Sel}(y, A, B)$ with its introduction and elimination. Every introduction in these constructions has an unused witness, except those of the towers over $0 = 0$, whose result does not depend on the witness.

Lemma 7.60 (A name without equality)**.** *For every term t over Π there is a persistent derivation in P_ν of*

$$\mathrm{Name}(t) := D_{x,u}\big(C(u,0,x) = C(u,0,t)\big),$$

and the clause obtained by opening $\mathrm{Name}(t)$ *at a fresh b gives $b = t$ by one* Use*.*

Proof. The opened clause is $[0 = 0;\ u.\, C(u,0,b) = C(u,0,t)]$; Use at 1, with the antecedent from Rf, gives $b = t$. For the derivation let

$$F(y) := D_{x,u}\Big(C\big(y,0,C(u,0,x)\big) = C\big(y,0,C(u,0,P(y))\big)\Big).$$

$F(0)$ normalizes to $D_{x,u}(0 = 0)$ and is introduced at whatever parameter is the newest. In the branch for $F(Sz)$ the parameter z is the newest, and $F(Sz)$ normalizes to $D_{x,u}(C(u,0,x) = C(u,0,P(Sz)))$, that is, to $\mathrm{Name}(z)$, because $P(Sz) \equiv z$. Its introduction selects the witness z; the instance at 0 is $0 = 0$, so the mode is by progress, and the instance asked for is $z = z$, by Use of Rf. Numerical cases give $q_y F(y)$, and its elimination at the term St gives $\mathrm{nf}(F(St))$, which is literally $\mathrm{Name}(t)$, since $C(St,0,b) \equiv b$ and $P(St) \equiv t$. The derivation is constructed anew in each scope. □

The proof uses no arbitrary witness, no equality substitution and no predecessor theorem. The shift by one is necessary: the value $P(0) = 1$ is erased by the outer selector.

7.7.4 Completeness

The source of the translation is the diagonal calculus $\mathsf{N}_{\mathrm{diag}}$ of Section 7.5.5, over the terms of Section 1.1 extended by the fixed descriptions R with the equations (7.8); it is stated with transport in formula templates, which is a derived rule of it. The iterator is translated to J with an unused counter coordinate, equations and binders are translated compositionally, $Q_y B$ is translated to $q_y B$, and all terms are normalized. A diagonal matrix has the normalized instance $0 = 0$ at $y = 0$, whatever its witness and its bodies are.

Theorem 7.61 (Completeness of the three families)**.** *Let $\Gamma \vdash A$ be derivable in $\mathsf{N}_{\mathrm{diag}}$ over the extended vocabulary. In every scope that contains the parameters, and from the clauses $[\ ;\ v.B^*]$ for $B \in \Gamma$, P_ν derives the normalized translation A^*. In particular closed theorems and the contradiction are preserved.*

Proof. By induction on the source derivation the following is shown: for every target scope and context, every assignment of target terms to the source parameters, and persistent builders of the translations of the source assumptions, there is a persistent builder of the translation of the conclusion. At the root the builder of an assumption is its use. Every replacement of a local source assumption made below uses only the use of a clause, reflexivity, and transport through the dummy major premise $D_{x,y}(0 = 0)$, whose introduction does not depend on the selected witness; the builders therefore remain persistent.

Computation axioms translate to reflexivity, their two sides having the same normal form. *Hypotheses* invoke their builders.

Elimination of $D_\cdot$. Translate the major premise and apply Discharge with a constant template and $0 = 0$. The source assumption $Q_y B(b,y)$ of the branch is represented by the clause $[B(b,0);\ y.B(b,y)]$. Turn the clause into the formula $q_y B(b,y)$ by an introduction with an unused witness: in its premise use the clause at Sz, the antecedent being the local assumption, or reflexivity if the rule selected progress. This builder is persistent and realizes the assumption that the induction hypothesis for the branch requires.

Elimination of Q. Translate the major premise to $q_y B(y)$ and open it first. Then apply the induction hypothesis to the derivation of $B(0)$ in the extended context, use the new clause

at the required term, and close the DISCHARGE with a constant template and $0 = 0$. The premise $B(0)$ is not translated outside the opening and then moved into the branch.

Transport. Open the dummy major premise $D_{x,y}(0 = 0)$, apply the induction hypothesis to the two source premises in the extended context, and finish the DISCHARGE with the template of the source inference.

Introduction at a witness t. Derive Name(t) by Theorem 7.60 and open it at a fresh b. The branch has the persistent equation $b = t$, and b is the newest parameter, so INTRO selects b. If the source introduction is by progress, its matrix is diagonal, its normalized instance at 0 is $0 = 0$ also at b, and the target rule selects progress: transport the assumption at b to the assumption at t, apply the induction hypothesis to the source premise, and transport the result back to b. If the source introduction is by cases and the target rule selects cases, do the same with the assumption at 0. If the target rule selects progress, the instance at 0 at the witness b is reflexive: derive it by USE of Rf, transport it to the instance at 0 at the witness t, apply the induction hypothesis, and transport back; the progress assumption is unused. For each transport back, the dummy major premise is opened before the induction hypothesis is applied to the source premise, so that the recursive call is made in the extended context; the builders of its assumptions are persistent there. The introduction yields $D_{x,y}B$, which does not contain b, and the opening of the name is closed with a constant template and $0 = 0$.

Every recursive call is for a proper subderivation of the source, and no completed derivation of the target is moved into a new scope. □

The translation covers formulas that contain R essentially and not only the image of the iterator terms: the logical comparison and the diagonal argument treat the two sides of an equation as arbitrary term templates, and the additional computation axioms translate to reflexivity.

7.7.5 The ordinary theory and the bridge

Let T_R be intuitionistic first-order logic with equality over the fixed descriptions R, with the equations (7.8), numerical coverage, $0 = 1 \to \bot$ and contextual induction for literal equations. The implication $0 = 1 \to \bot$ is stated explicitly: the equation $0 = 1$ is the native contradiction and is not by itself falsity in a structure with one element.

Lemma 7.62 (Extensionality for a fixed body)**.** *If $h(i, u) = k(i, u)$ is derivable in T_R uniformly in i and u, then $R(n, a;\, i, u.\, h) = R(n, a;\, i, u.\, k)$ is derivable for every n, with all other parameters fixed.*

Proof. By induction for the displayed equation. At 0 both sides are Sa. At a successor the computation exposes the two bodies; the induction equation replaces the previous value in one of them, and the given identity turns h into k. □

Consequently every finite reduction of a term t to $\mathrm{nf}(t)$, also inside a body, yields a derivation of $t = \mathrm{nf}(t)$ in T_R, and by induction on a formula its raw and its normalized readings are equivalent. No truth or evaluation predicate is used.

Theorem 7.63 (Bridge for the recursor calculus)**.** *P_ν is sound and complete for T_R, and T_R is bi-interpretable with iPRA: both domains are the whole numerical sort, equality is equality, both numerical comparison maps have the graph $x = y$, and every formula is provably equivalent to its double translation. In particular P_ν derives $0 = 1$ if and only if iPRA proves $\bot$.*

Proof. Soundness is Theorem 7.59 with Theorem 7.62. Completeness is Theorem 7.61 together with the equivalence of $\mathsf{N}_{\mathrm{diag}}$ with N and the presentation of the ordinary theory by N (Theorem 1.78), which are parametric in the additional fixed descriptions.

A description $R(n, a;\, i, u.\, h)$ is interpreted in iPRA by the primitive recursion with the base Sa and the step h, after its proper subdescriptions. Conversely

$$\begin{aligned} 0^* = 0, \qquad S^*(x) = R(0, x;\, i, u.\, 0), \qquad \pi^*_j(\vec{x}) = x_j, \\ (f \circ \vec{g})^*(\vec{x}) = f^*(g^*_1(\vec{x}), \dots, g^*_m(\vec{x})), \\ \mathrm{rec}(g, h)^*(n, \vec{x}) = J\big(n, g^*(\vec{x});\, i, u.\, h^*(i, u, \vec{x})\big). \end{aligned}$$

No pairing is needed, because the body receives the counter. The compiled equations hold by the normal forms of J. Applying P to $Sx = Sy$ gives the injectivity of the successor; applying $C(\cdot, 0, 1)$ to $Sx = 0$ gives $1 = 0$, hence the separation axiom. The iterator J carries the reflection of equality of Theorem 1.52, so every quantifier-free formula is equivalent to the vanishing of a term and its induction follows from induction for an equation (see also Theorem 7.74). For two descriptions with the same base and provably identical steps, induction for the equation of their values proves that they agree at every counter; by induction on a description from outside, its double translation has the same values as the original, for all arguments. The formula round trips follow by induction on formulas from the reading of $D_,$ and the logical macros (Section 7.8.5). Falsity corresponds to $0 = 1$, and derivations of a contradiction translate in both directions. □

7.7.6 Checking a template through its two instances

In DISCHARGE the checker receives a template $F(v)$, terms s, t and derivations of $s = t$ and $\mathrm{nf}(F(s))$, and its result $\mathrm{nf}(F(t))$ must equal the conclusion, which is checked. A separate check that F is well-scoped and normal is redundant.

Lemma 7.64 (Completion of an accepted template)**.** *Let F be a finite expression such that the substitution of the checked terms s and t for its distinguished variable, followed by normalization, succeeds, and such that the two results are well-scoped formulas in the local scope. Then there is a well-scoped template $F'(v)$ in normal form with*

$$\mathrm{nf}(F'(s)) = \mathrm{nf}(F(s)), \qquad \mathrm{nf}(F'(t)) = \mathrm{nf}(F(t))$$

literally.

Proof. Sorts. The traversal that substitutes and normalizes checks every proper child at its required sort, term or formula, before any reduction that erases it. At the root it may accept an unclassified expression. A root that is neither an equation, a binder nor a recursor does not reduce and yields no formula; a numerical root stays numerical under reduction. Since an instance is a formula, the root of F is an equation or a binder, and F is a sorted formula template that may contain additional numerical parameters and reducible subterms.

Additional parameters. Read every parameter of F outside the local scope, and every bound-variable index that exceeds the binders above it by more than the one position of the distinguished variable, as an additional free parameter; the reading is consistent under a binder because the excess is measured relative to the depth. Let ρ replace all additional parameters by 0 and fix v, the local parameters and the bound variables. Neither s nor t contains an additional parameter. For $r \in \{s, t\}$, by Theorem 7.58,

$$\mathrm{nf}\big((\rho F)[r/v]\big) = \mathrm{nf}\big(\rho(F[r/v])\big) = \mathrm{nf}\big(\rho\ \mathrm{nf}(F[r/v])\big) = \mathrm{nf}(F[r/v]),$$

the last step because the checked instance is normal and contains no additional parameter. A reduction that becomes possible after an additional parameter is closed does not affect the argument, which uses the law of Theorem 7.58 and not an identity of reduction sequences.

The template. Put $F' := \mathrm{nf}(\rho F)$. It is sorted, well-scoped and normal, and a second application of Theorem 7.58 gives the two identities. The distinguished variable may occur any number of times. □

Replacing F by F' leaves the two premises and the conclusion unchanged, so every accepted use of the relaxed interface is an instance of DISCHARGE, and every canonical template is still accepted. The equation $s = t$ must be checked before the template is instantiated, so that s and t are known to be terms, and the instance in the branch and the conclusion keep their checks; the lemma does not justify omitting either.

7.8 Conditional copy

This section draws on `conditional_copy`; the example of an unsound probe is from `deterministic_introduct`

The calculus P_ν takes no witness at its introduction and still takes an equation, with its derivation, and a template at its discharge. Other calculi infer the substitution of a discharge from two instances of the opened matrix and take an arbitrary witness at their introduction. The two restrictions are combined here. The composition is not immediate: with an inferred discharge, equality transport is recovered from packets that are introduced at arbitrary witnesses, and the names of Theorem 7.60 give an equation $b = t$, whose use would need the transport that is being recovered. The combination rests on a formula that copies an instance of a formula to a fresh parameter without any equation.

7.8.1 Use, Close and Export

Terms, normal forms, the abbreviations P, C, J, formulas, clauses and the initial clause Rf are those of Section 7.7. Write $\top_* := (0 = 0)$ and $\bot_* := (0 = 1)$. The calculus K has three rule families; all displayed instances are normalized.

(i) USE: as in P_ν.

(ii) CLOSE: with $\ell = \nu(\Pi)$ and a fresh z,

$$\frac{\Pi, z;\ \Gamma, [\ ;\ v.\,\mathrm{nf}(B(\ell,\eta))] \vdash \mathrm{nf}(B(\ell, Sz))}{\Pi; \Gamma \vdash D_{x,y}B} \qquad \eta = \begin{cases} z, & \mathrm{nf}(B(\ell,0)) \text{ is literally } \top_*, \\ 0, & \text{otherwise.} \end{cases}$$

The test for $\top_*$ is narrower than the test for a reflexive equation in P_ν and suffices.

(iii) EXPORT: with a fresh b, put

$$B_0 = \mathrm{nf}(B(b,0)), \qquad B_1 = \mathrm{nf}(B(b,1)), \qquad r_B = \begin{cases} t, & B_0 \equiv \top_* \text{ and } B_1 \equiv (b = t), \\ b, & \text{otherwise,} \end{cases}$$

where $\equiv$ is literal identity, including the orientation of the equation, and let $\mathcal{C}_B(b) = [B_0;\ y.\,\mathrm{nf}(B(b,y))]$. Then

$$\frac{\Pi; \Gamma \vdash D_{x,y}B \qquad \Pi, b;\ \Gamma, \mathcal{C}_B(b) \vdash F}{\Pi; \Gamma \vdash \mathrm{nf}(F[r_B/b])} \qquad b \notin \mathrm{FV}\big(\mathrm{nf}(F[r_B/b])\big).$$

F may contain b, and so may r_B; the substitution is one simultaneous pass. No equation, replacement term or template is supplied.

Only USE takes a term.

Lemma 7.65 (Soundness)**.** *Every rule of* K *is valid in* T_R *under the ordinary reading.*

Proof. USE and CLOSE are as in Theorem 7.59; if the instance at 0 is $\top_*$, the matrix is a literal equation, since normalization does not remove a binder. For EXPORT open the existential quantifier. In the selected case reflexivity proves B_0, the clause at 1 gives $b = r_B$, and the substitution of equals turns F into $F[r_B/b]$; in the default case the substitution is the identity. The elimination of the quantifier is legitimate because the normalized result does not contain b. Normalization is justified in T_R by Theorem 7.62. □

Example 7.66 (The instance at 0 must be tested)**.** An EXPORT that inspected only the instance at 1 would be unsound. Let f be a term with $f(0) = f(1) = 0$ and $f(2) = 1$, and let $B(x, y)$ be $x = f(y)$. The reading of $D_{x,y}B$, $\exists x\,(x = f(0) \to \forall y\, x = f(y))$, is provable with the witness 1, whose antecedent is false. The instance at 1 at a fresh b is $b = 0$. A rule that replaced b by 0 on this evidence would produce the clause $0 = 0 \to \forall y\, 0 = f(y)$, which gives $0 = 1$ at $y = 2$. The test $B_0 \equiv \top_*$ excludes this matrix. For the same reason the name $D_{x,y}\mathrm{Nm}_t$ of Section 7.6 is not silently combined with an inferred opening: opening a name directly at t removes the fresh alias $b = t$ on which the simulation of an arbitrary witness relies.

7.8.2 Preparations without equality

If F does not contain the opened parameter b, the substitution of EXPORT does nothing to it; so the ordinary elimination of $D_{x,y}B$ is available whatever r_B is. As in Section 7.7, $q_yA(y)$ is $D_{d,y}A(y)$ with d unused; opening it and using its clause gives $A(t)$ from q_yA and $A(0)$, and the witness selected by CLOSE for it is irrelevant. Given persistent constructions of $A(0)$ and of $A(Sz)$, CLOSE introduces $q_yA(y)$: if the instance at 0 is $\top_*$, supply reflexivity in place of the assumption at 0 and ignore the progress assumption; otherwise the rule is by cases. The same holds for a premise that assumes $A(0)$.

Every formula is a chain of binders ending in an equation. For a formula A with d binders let G_vA be as in Section 7.7; then

$$G_0A \equiv T_d, \qquad G_{Sz}A \equiv A, \qquad T_0 = \top_*, \quad T_{d+1} = D_{x,y}T_d.$$

Every tower T_d is derivable in every scope, by reflexivity at depth 0 and CLOSE above it; none of its witnesses occurs in its body. For formulas A and B with the same number of binders, align the binders and replace corresponding sides s, t of the final equations by $C(v, s, t)$. This defines $\mathrm{Sel}(v, A, B)$, with

$$\mathrm{Sel}(0, A, B) \equiv A, \qquad \mathrm{Sel}(Sz, A, B) \equiv B.$$

A substitution in terms does not change the number of binders, so $H(s)$ and $H(x)$ always have the same depth. For formulas of different depth the padding $L(A) = q_vG_vA$, interderivable with A, equalizes the depths. All of these are constructions carried out in the scope where they are needed.

7.8.3 The conditional-copy theorem

Theorem 7.67 (Conditional copy)**.** *For every formula $H(x)$ with a distinguished variable and every term t over Π, the formula*

$$\Delta_t(H) := D_{x,y}\,\mathrm{Sel}(y, H(t), H(x))$$

is derivable in K in every scope and context, by a construction that uses no equality transport and no equation other than reflexivity.

Proof. Construct the closure

$$q_vG_v\Delta_{P(v)}(H). \tag{7.9}$$

Its instance at $v = 0$ is a tower. In the branch for $v = Sz$, with z fresh, the conclusion normalizes to $\Delta_z(H)$, because $P(Sz) \equiv z$ and the gate disappears. The parameter z is the newest, so CLOSE for $\Delta_z(H)$ selects $x = z$. Its instance at 0 and its instance at a successor are both literally $H(z)$:

$$\mathrm{Sel}(0, H(z), H(z)) \equiv H(z), \qquad \mathrm{Sel}(Sw, H(z), H(z)) \equiv H(z).$$

If $H(z) \not\equiv \top_*$, the rule is by cases and its assumption is its conclusion. If $H(z) \equiv \top_*$, the rule is by progress and reflexivity derives the instance asked for. The outer introduction is by cases, since its instance at 0 has at least one binder; its assumption is unused. Open (7.9) by the ordinary elimination, supply the tower as the instance at 0, and use the clause at St. The gate disappears and $P(St) \equiv t$, so the result is literally $\Delta_t(H)$; it does not contain the opened parameter. Every step can be carried out in every scope. □

The ordinary reading of $\Delta_t(H)$ is $\exists x\,(H(t) \to H(x))$. The theorem neither proves $H(t)$ nor identifies x with t. Opening $\Delta_t(H)$ gives a fresh parameter b and the clause $[H(t);\, v.\,\mathrm{Sel}(v, H(t), H(b))]$, which yields $H(b)$ at $v = 1$ once $H(t)$ is supplied.

7.8.4 Removal of the witness

Let L, the *liberal* calculus, have the syntax, Use and Export of K and a Close that takes an arbitrary term t over Π in place of ℓ, with the same test: progress exactly when $\mathrm{nf}(B(t,0)) \equiv \top_*$.

Theorem 7.68 (Witness removal)**.** *For every scope* Π*, clause context* Γ *and formula* A*, the calculi* K *and* L *derive the same judgments* $\Pi; \Gamma \vdash A$*.*

Proof. A Close of K is a Close of L. For the converse a derivation of L is translated by induction on its tree. The induction statement admits an injective map θ from the source parameters into an arbitrary target scope and a map from the positions of the source clauses to positions of target clauses, such that the clause at each image position is literally the normalized θ-image of the source clause; further target parameters and clauses are allowed. This is an invariant of membership; it is not an assumption that completed target derivations can be weakened.

Use follows the map of positions and translates the derivation of the antecedent. Export translates its major premise, sends its eigenvariable to the new target parameter and extends the map of positions by the opened clause; its two tests are preserved and reflected by injective renaming and Theorem 7.58.

Consider a Close at a term t with the matrix B and the premise $\Gamma, [\ ;\, v.\,\mathrm{nf}(B(t,\eta))] \vdash \mathrm{nf}(B(t, Sz))$. Write t, B also for their images, and put $H(x) := q_y B(x,y)$.

(a) Derive $\Delta_t(H)$ by Theorem 7.67 and open it at a fresh b; its clause is $[H(t);\, v.\,\mathrm{Sel}(v, H(t), H(b))]$.

(b) Apply Close to $D_{x,y}B$. The newest parameter is b, so the rule selects b; let z' be its eigenvariable.

(c) In the premise derive $H(t)$ by a Close with an unused witness. Its test is the source test at t, and its premise, with a further eigenvariable, is the image of the source premise: the induction hypothesis applies with the source eigenvariable sent to the new one and the source assumption sent to the local clause of this Close.

(d) Use of the clause of (a) at 1 gives $H(b)$.

(e) Open $H(b)$; its clause is $[B(b,0);\, y.B(b,y)]$. If the Close of (b) is by cases, its assumption is $B(b,0)$. If it is by progress, $B(b,0)$ is literally $\top_*$ and reflexivity supplies it. Use at Sz' gives $B(b, Sz')$, and the unused witness of $H(b)$ is discharged.

The Close of (b) yields $D_{x,y}B$, which does not contain b, and the ordinary elimination discharges b. Every recursive call is for the premise of the source Close, a proper subderivation, although it may be made several times; the packet and the towers are explicit constructions. At each call the tests agree, because they are made on the same instantiated formula. □

Remark 7.69 (The modes need not agree). The test at t and the test at b may differ. The proof transfers the whole conditional closure H and not a progressive premise; at b a progressive mode has a reflexive instance at 0, so the closure still yields the instance at a successor. No equality reasoning repairs a failed test. The same proof applies with the rule NAME of Section 7.8.7 in place of EXPORT.

7.8.5 The liberal calculus presents the ordinary theory

Lemma 7.70 (Structural facts of the liberal calculus)**.** *In* L*, weakening by clauses, injective renaming of parameters and the cut of an assumption* $[\,;v.A]$ *are admissible.*

Proof. By induction on derivations. A supplied witness remains available after a renaming, an injective renaming preserves and reflects the literal tests of CLOSE and EXPORT, and normalization commutes with renaming. For the cut, a USE of the clause is replaced by the given derivation, weakened and renamed into the scope of the leaf. No equality transport and no substitution of terms into derivations is used. The lemma is about L and is not claimed for K at this stage. □

Lemma 7.71 (Transport from an inferred discharge)**.** *In* L*, from* $s = t$ *and* $h(s) = u$ *one derives* $h(t) = u$*. Consequently symmetry, transitivity, congruence and transport in arbitrary formulas are derived.*

Proof. Let

$$E(x,y) := \Big(C(y,0,C(P(y),x,h(x))) = C(y,0,C(P(y),t,u))\Big).$$

Its normalized instances are $E(x,0) \equiv \top_*$, $E(x,1) \equiv (x = t)$ and $E(x,SSv) \equiv (h(x) = u)$. Introduce $D_{x,y}E$ at the witness s; the mode is by progress, and the instance asked for, $C(z,s,h(s)) = C(z,t,u)$, is derived by numerical cases on z from the two given equations, the progress assumption being unused. Open the formula at b: the first two instances make EXPORT select t. USE at 2, with the reflexive antecedent, gives $h(b) = u$, and EXPORT concludes $h(t) = u$.

With reflexivity this rule gives symmetry, transitivity and congruence, and transport in an equation template is two such replacements. Transport in a formula $D_{x,y}B(v,x,y)$ follows by induction on the number of binders: open the given formula at its witness, introduce the target at that witness, transport the assumption at 0 backwards through the matrix, use the opened clause at the successor and transport the result forwards; if the new instance at 0 is $\top_*$, supply reflexivity instead. The result does not contain the opened witness. The premises are moved into the local scopes by Theorem 7.70. □

Lemma 7.72 (Cases and equation induction in the liberal calculus)**.** *In* L *the introduction by cases is available for every matrix and witness, and for a literal equation* $E(y) \equiv (s(y) = t(y))$*, from* $E(0)$ *and a derivation of* $E(Sz)$ *from* $E(z)$ *one derives* $E(a)$ *for every term* a*.*

Proof. If an introduction by cases selects progress because its instance at 0 is $\top_*$, replace the assumption at 0 by reflexivity and ignore the progress assumption. For the induction let

$$K(y) := (C(y,0,s(y)) = C(y,0,t(y))),$$

whose instance at 0 is literally $\top_*$. Numerical cases, without induction, give the two decoder equations

$$C(y,s(0),C(y,0,s(y))) = s(y), \qquad C(y,s(0),C(y,0,t(y))) = t(y);$$

the first has reflexive branches, and the second uses $E(0)$ at 0 and is reflexive at successors. From $K(z)$, congruence with $v \mapsto C(z,s(0),v)$ and the decoders give $E(z)$. The cut of this into

the step gives $E(Sz)$, which is literally $K(Sz)$. The introduction by progress gives $q_y K(y)$, Use with the reflexive antecedent gives $K(a)$, and the decoders give $E(a)$. No compound term is substituted into a derivation. A conditional closure $q_y E(y)$ is introduced by cases, with $E(0)$ as its assumption and the induction just derived for its successor instance. □

The ordinary logical vocabulary is given by the following macros, with the depths equalized by L before each binary selector and all bound variables fresh:

$$\begin{aligned} (A \to B)^* &= q_y \operatorname{Sel}(y, A^*, B^*), & (A \wedge B)^* &= q_y G_y \operatorname{Sel}(P(y), A^*, B^*), \\ (\exists x\, A)^* &= D_{x,y} G_y A^*, & (A \vee B)^* &= D_{v,y} G_y \operatorname{Sel}(v, A^*, B^*), \\ (\forall x\, A)^* &= q_y G_y A^*(P(y)), & \bot^* &= \bot_*. \end{aligned}$$

The instances of the conjunction are a tower at 0, A^* at 1 and B^* at every SSv; two numerical case splits introduce it and Use at 1 and 2 eliminates it. The implication has its antecedent at 0 and its consequent at successors. The existential gate has a tower at 0. Numerical cases give the elimination of the disjunction. The universal macro has a tower at 0 and $A^*(z)$ at Sz, because $P(Sz) \equiv z$; it is used at St to obtain $A^*(t)$, and its introduction needs a fresh renaming of a derivation of $A^*(z)$ only. Transport along $0 = 1$ turns the provable $G_0 F$ into $G_1 F \equiv F$, which is explosion for every F. These constructions give the introduction and elimination rules of intuitionistic logic under arbitrary assumptions, and by induction on a formula the ordinary reading of each macro is equivalent to the intended connective. `parameter_intro` uses the same macros with two differences: the conjunction selects by a parity term, $(A \wedge B)^* = q_y G_y \operatorname{Sel}(\operatorname{par}(y), A^*, B^*)$, and the universal quantifier is $(\forall x\, A)^* = A^*(0) \wedge q_x A^*(x)$.

Proposition 7.73 (The liberal calculus)**.** L *is sound and complete for* T_R*. Hence so is* K*.*

Proof. Soundness is Theorem 7.65 with a supplied witness. For completeness translate the diagonal calculus $\mathsf{N}_{\mathrm{diag}}$ over the vocabulary with the fixed descriptions R, as in Theorem 7.61: computation axioms become reflexivity after normalization; hypotheses are used; equality is Theorem 7.71; the eliminations of Q and $D_,$ are the derived ones; a diagonal matrix has the instance $0 = 0$ at 0, so its introduction by progress is the progress mode of Close; an introduction by cases is Theorem 7.72. The structural facts of Theorem 7.70 move derivations between scopes. This gives the full consequence relation and not only closed equations. The statement for K follows by Theorem 7.68, which regenerates the branches in the target scopes. □

7.8.6 Quantifier-free induction and the bridge

Lemma 7.74 (Reflection of equality in T_R)**.** *There is a term* $e(a, b)$ *such that* T_R *proves* $e(a, b) = 0 \vee e(a, b) = 1$ *and* $e(a, b) = 0 \leftrightarrow a = b$*. For every quantifier-free formula* A *there is a term* b_A *with the same parameters such that* T_R *proves* $b_A = 0 \vee b_A = 1$ *and* $A \leftrightarrow b_A = 0$*. Hence* T_R *proves every instance of contextual quantifier-free induction.*

Proof. Define addition and truncated subtraction by J, with

$$x + 0 = x, \quad x + Sy = S(x + y), \qquad d(x, 0) = x, \quad d(x, Sy) = p(d(x, y)),$$

and put $\operatorname{sg}(u) = C(u, 0, 1)$ and $K(u) = C(u, 1, p(u))$. Coverage and the separation of 0 and 1 give that $\operatorname{sg}(u)$ is 0 or 1, that $\operatorname{sg}(u) = 0 \leftrightarrow u = 0$ and that $K(u) = 0 \leftrightarrow u = 1$. Four inductions for literal equations give

$$0 + y = y, \qquad Sx + y = S(x + y), \qquad d(Sx, Sy) = d(x, y), \qquad d(Sa, a) = 1;$$

the step of the third applies p to the induction hypothesis, and the fourth follows from the third.

The invariant is the single equation

$$C\big(\mathrm{sg}(d(x,y)),\, x,\, y + d(x,y)\big) = x, \tag{7.10}$$

proved by induction on y with x fixed. At 0 both alternatives of the selector equal x, by $0 + x = x$. For the step put $r = d(x,y)$ and $r' = p(r) = d(x,Sy)$. If $r' = 0$, the selector returns x. Otherwise coverage gives $r = Sr'$, the alternative $r = 0$ contradicting $r' \neq 0$; the induction equation gives $y + r = x$, and $Sy + r' = y + Sr' = x$. All case distinctions are coverage, and the induction formula is (7.10).

Hence $d(Sa,b) = 1 \leftrightarrow a = b$: from left to right (7.10) gives $b + 1 = Sa$ and the injectivity of the successor gives $a = b$; from right to left this is the fourth identity. Put $e(a,b) = \mathrm{sg}(K(d(Sa,b)))$. For a quantifier-free formula define, by recursion on its syntax,

$$b_{s=t} = e(s,t), \quad b_{\bot} = 1, \quad b_{A\wedge B} = C(b_A, b_B, 1), \quad b_{A\vee B} = C(b_A, 0, b_B), \quad b_{A\to B} = C(b_A, b_B, 0).$$

By induction on the syntax b_A is 0 or 1 and $A \leftrightarrow b_A = 0$; for the implication split the two values of b_A. No excluded middle for formulas with quantifiers is used. Induction for the literal equation $b_{A(n)} = 0$ gives the induction for A, under arbitrary assumptions. □

Theorem 7.75 (Introduction and discharge without supplied terms)**.** K *is a sound and complete presentation of* T_R*, and* T_R *is bi-interpretable with* iPRA*, with the domains* $x = x$*, equality as equality, the graph* $x = y$ *in both numerical composites and a provable equivalence of every formula with its double translation. Hence* K *and* iPRA *are equiconsistent.*

Proof. The first statement is Theorem 7.73. The interpretations, the compilation of primitive recursion by J, the injectivity and separation of the successor and the uniqueness argument are those of the proof of Theorem 7.63; quantifier-free induction is Theorem 7.74; the formula round trips follow from the macros of Section 7.8.5 by induction on formulas. Falsity is translated to $0 = 1$, whose reading implies $\bot$ by the separation axiom of T_R. □

Here iPRA has quantifier-free induction, and T_R has induction for literal equations only; the words "first-order" refer to the logical language.

7.8.7 Discharge by a supplied name

Let $\mathsf{K_N}$ have the syntax, the clause Rf, Use and Close of K, and in place of Export the rule

$$\frac{\Pi;\Gamma \vdash D_{x,y}B \qquad \Pi, b;\ \Gamma, \mathcal{C}_B(b) \vdash b = t \qquad \Pi, b;\ \Gamma, \mathcal{C}_B(b) \vdash F}{\Pi;\Gamma \vdash \mathrm{nf}(F[t/b])}\ \textsc{Name} \qquad b \notin \mathrm{FV}\big(\mathrm{nf}(F[t/b])\big).$$

The term t is over Π, b and may contain b; it is inserted once. The left side of the supplied equation is fixed to the opened parameter, whereas Discharge of P_ν takes both sides of an equation and a template.

Proposition 7.76 (The name variant)**.** $\mathsf{K_N}$*,* K *and* T_R *have the same open consequences under their readings.*

Proof. Name is sound in T_R: equality gives the substituted formula and the condition on the scope permits the elimination of the quantifier. Every instance of Export is an instance of Name in the same scope: in the default case with the reflexivity $b = b$, in the selected case with the equation $b = t$ obtained from the opened clause by reflexivity and Use at 1. The replacement adds no parameter, so no Close changes its witness or its mode, and induction on the derivation gives $\mathsf{K} \subseteq \mathsf{K_N}$. The completeness of K and the soundness of $\mathsf{K_N}$ give the rest. □

A backward checker for K_N receives a template $T(v)$ and a term t, checks the conclusions $b = t$ and $F = \mathrm{nf}(T[b/v])$ in the branch, and requires the goal to be $A = \mathrm{nf}(T[t/v])$. It does not forbid occurrences of b in T.

Lemma 7.77 (Template representation)**.** *Let $T(v)$ be a well-scoped template and t a term of the local scope, and put $F = \mathrm{nf}(T[b/v])$ and $A = \mathrm{nf}(T[t/v])$. If A does not contain b, then $\mathrm{nf}(F[t/b]) = A$ literally.*

Proof. A renaming of parameters, injective or not, preserves normal forms: the reductions test only for the constructors 0 and S at a counter, and an identification of parameters creates neither; no reduction turns an equation with equal sides into $0 = 0$. Choose a parameter c not in T or t and let $U = \mathrm{nf}(T[v := t,\, b := c])$, with simultaneous substitution. Identifying c with b gives

$$U[c \mapsto b] = \mathrm{nf}(U[c \mapsto b]) = \mathrm{nf}(T[t/v]) = A.$$

Since A does not contain b, U contains neither b nor c, and $U = A$. Substituting t for c and Theorem 7.58 give $\mathrm{nf}(T[v := t,\, b := t]) = A$. Finally

$$\mathrm{nf}(F[t/b]) = \mathrm{nf}\big((T[b/v])[t/b]\big) = \mathrm{nf}(T[v := t,\, b := t]) = A.$$

No substitution is made inside an inserted copy of t. □

The checker validates a template through its two normalized instances, as in Theorem 7.64, and the same completion applies: the excess parameters of an accepted raw template are replaced by 0, which preserves both checked instances literally because neither they nor the checked arguments b, t contain an excess parameter; the normalized completed template is well-scoped, and Theorem 7.77 applies to it. The equation premise validates t, and each instance is validated in its own scope; these checks are needed. Conversely, abstracting all free occurrences of b in F into v represents every instance of NAME. The backward representation is therefore exact.

7.9 Checkers and sizes

This section draws on the checker reports of `no_witness`, `parameter_witness`, `deterministic_introduction` `newest_parameter`, `uniform_raw`, `parameter_intro` and `conditional_copy`. The checker of `constructed_computation` belongs to another chapter.

All checkers are standalone Python modules that read one JSON certificate, validate syntax, scopes and every inference, and accept or reject; none calls a prover. They share the encoding of syntax: the integer 0 is zero, negative integers are parameters counted from the outside, positive integers are de Bruijn indices, and lists encode the constructors. The sizes are measured by the protocol of Section 1.10 under CPython 3.12.14.

Table 7.1: Checkers of the chapter: significant tokens, AST nodes and normalized bytes, with the matched control as tokens / nodes / bytes.

Program	Tokens	Nodes	Bytes	Matched control
`no_witness`				
eight rules, N_ν	971	1078	2817	supplied witness: 979 / 1086 / 2834
seven rules, $\mathsf{N}_{\mathrm{diag},\nu}$	924	1021	2753	supplied witness: 932 / 1029 / 2770

Program	Tokens	Nodes	Bytes	Matched control
`parameter_witness`				
five families, C_{β}	833	971	2947	two-case raw clause checker: 837 / 977 / 2956
`deterministic_introduction`				
G_{d}	1159	1339	3688	G: 1078 / 1252 / 3442
inherited native calculus, shared first traversal	333	379	656	incoming native control: 360 / 409 / 780
inherited guarded compact module	267	303	506	before the change: 267 / 302 / 508
`newest_parameter`				
G_{ν}	1070	1243	3425	G: 1078 / 1252 / 3442; G_{d}: 1159 / 1339 / 3688
`uniform_raw`				
G_{ν}	1073	1248	3439	G_{d}: 1159 / 1339 / 3688
`parameter_intro`				
P_{ν}	630	701	1925	supplied witness: 636 / 707 / 1967
P_{ν} with a separate template check	646	717	1985	
P_{ν} with fuel	667	748	2056	supplied witness, with fuel: 673 / 754 / 2098
`conditional_copy`				
K	646	748	1844	L: 651 / 755 / 1863
K_{N}, backward	619	686	1907	P_{ν} of `parameter_intro`: 630 / 701 / 1925
explicit equation at NAME	634	742	1835	
K with fuel	685	796	1995	

The packages compare their programs with further programs of the corpus. `parameter_intro` states 636 / 708 / 2010 for the incoming checker with an explicit name and 683 / 771 / 1990 for the incoming checker with an inferred opening; its gzip sizes are 778 for its checker, 790 for the matched control, 792 with the separate template check, and 826 and 838 for the two variants with fuel. `conditional_copy` states 662 / 741 / 1950 for the checker of `uniform_recursion`. The compact programs of the corpus use other input contracts and are smaller: 267 tokens, 302 nodes and 508 bytes; 256, 283 and 594; 263, 276 and 580; and, with a validator of the public goal, 360, 409 and 780. The three minima 256, 276 and 508 belong to different programs. No package claims a global minimum or a lower bound.

`no_witness`. An introduction is the instruction `['di', matrix, premise]`; its witness is the integer $-n$ in a scope of n parameters, which is zero in the empty scope. The controls differ only by a supplied and validated witness term. Each matched pair differs by 8 tokens, 8 nodes and 17 bytes. In the tagged syntax of the pinned programs the same change saves 3 bytes and 1 node and adds 2 tokens. Eighteen accepted and 33 rejected certificates are run in two modes; the certificate for $X(2)$ checks the part of the naming that uses numerical cases and not the predecessor or the equivalence theorem.

`parameter_witness`. The introduction field is a bit, decoded by the truth value of the JSON value; it is never inserted as a term, and the witness is constructed from the scope. Certificates for the packet T at 0, for its closure and for its instances at 1 and 2 are checked, and a certificate that uses the eigenvariable of an introduction as its witness is rejected. A

separate evaluator checks 447 numerical equalities of the predecessor construction, among them the values of the predecessor at $0, \dots, 7$.

`deterministic_introduction`. The naming matrix is recognized as the literal list pattern of Nm_r with r valid in the parent scope. The checker of G_{d} is larger than its control by 81 tokens, 87 nodes and 246 bytes: the rule removes a datum from each introduction and costs more code to recognize the exception. The fixtures contain an introduction that is valid before an extension of the scope, the rejected replay of the same tree after the extension, and the repaired derivation of Theorem 7.45. The package also contains two changes to inherited programs that do not implement G_{d}. One checks the grammar of the public goal during the first scope traversal, using the parity of the traversal depth as the distinction between formulas and terms; an induction on the goal shows that the shared pass succeeds exactly when the two separate passes do and that the subsequent traces are equal. The other replaces $k - 1$ by the bitwise complement of $-k$ in the compact module, which saves two bytes and adds one node.

`newest_parameter` and `uniform_raw`. Both obtain the checker of G_{ν} from the checker of G_{d} by deleting the function that selects the witness and instantiating the matrix at $-n$. `newest_parameter` also contains an unmeasured checker of the eight rules of N_{ν}, with certificates for $X(St)$ at a parameter t in scopes of zero, one, two and five parameters, and a rejected replay of a closed derivation in a larger scope. `uniform_raw` checks the whole closure $D_{a,r}X(r)$ and its instance at the successor of a compound term.

`parameter_intro`. The recursor and the binder $D_{,}$ both bind two variables in their last child, so one traversal serves terms, formulas, scopes, substitution and normalization. The reduction of a successor counter substitutes the previous counter first, into the outer of the two bound positions, and the previous value second; the opposite order without a shift would capture variables. The witness of an introduction is the integer $1 - n$ for a context of n clauses. Omitting the witness field saves 42 bytes, 6 tokens and 6 nodes; omitting the separate template check, by Theorem 7.64, saves 60 bytes, 16 tokens and 16 nodes. The variant with fuel decrements a budget at every call of the traversal and of the inference checker; every accepted certificate has a sufficient budget, so the two interfaces have the same theorems, and the bounded relation is primitive recursive. The certificates that name 1 and 2 without a witness field need 1155 and 1345 units.

`conditional_copy`. The forward checker of K infers every conclusion and compares the goal with it. Omitting the witness saves 19 bytes, 5 tokens and 7 nodes against L. The checker of K has fewer bytes and more tokens and nodes than that of `parameter_intro`; the backward checker of K_{N} is smaller in all three coordinates, by 18 bytes, 11 tokens and 15 nodes. The package contains a translator from L to K that implements the proof of Theorem 7.68 on arbitrary derivations. Its outputs show the growth of certificates, in inference nodes and bytes of JSON: an introduction at the witness 1 grows from 2 nodes and 74 bytes to 15 and 529; a conditional copy of depth one from 7 and 368 to 46 and 2527; a derivation with a used progress assumption from 15 and 421 to 80 and 2868; the congruence of the successor under two universal quantifiers from 25 and 1335 to 142 and 8909; a derivation in an open scope from 4 and 76 to 17 and 501.

The restricted rules reduce the data of a certificate and, except for G_{d}, the measured checker by a small amount. They increase the length of certificates, because a witness term is replaced by a derivation of its name.

7.10 Further questions

The following items are stated by the packages without proof, are open, or mark what the results of the chapter do not establish.

Question 7.78 (Formalization of the names)**.** By Theorem 7.16, which is formalized, the completeness of $\mathsf{N}_{0\mathrm{v}}$ is the derivability of the names. The derivation of the names (Theorems 7.19 and 7.20) and the derivation of the predecessor in the fragment $\mathcal{E}$ are written proofs. Formalize them, with the predecessor as a theorem of the equational fragment separate from numerical coverage.

Question 7.79 (Formalization of the scope-determined witness)**.** None of Theorems 7.35, 7.38 to 7.40, 7.45, 7.47, 7.55, 7.61, 7.67 and 7.68 is formalized. A formalization of the first has to take derivations of the source with realized contexts, or hereditary families, as its induction interface. For G_{d} the selection with priority for a naming matrix, the clause contexts and the guarded language are not formalized. For the recursor calculi a syntax with a normalizer and its bi-interpretation with the iterator terms would be needed first; `conditional_copy` proposes a small interface for Theorem 7.67 consisting of a predecessor that computes on successors, gates, clause specialization, Close at the newest parameter and the ordinary elimination, so that the theorem can be reused. Theorem 7.64 and the translation of Theorem 7.61 are named by `parameter_intro` as the next formalization task.

Question 7.80 (Three families on raw iterator terms)**.** Can the introduction at the newest parameter be combined, on raw iterator terms without normalization, with a three-family calculus whose equality transport is recovered from packets? The proofs of Section 7.5 keep Tr as a primitive rule, and transferring them unchanged would be circular. Sections 7.7 and 7.8 answer the question on the recursor with normalization, where $P(Sz) \equiv z$ holds by computation; on iterator terms no term has this property. A technical note of `uniform_raw` sketches a raw calculus with a universal reflexivity clause, a deterministic introduction, Use and a discharge that evaluates, in which the opening and the evaluation rules are recovered through the dummy major premise $D_{x,y}(I(y,0;\,u.\,0) = I(y,0;\,u.\,0))$ and formula transport is proved before the names. The note is not part of the proved results of that package.

Question 7.81 (Transport in terms only)**.** In G_{d} and G_{ν} the rule Tr acts on arbitrary canonical formulas, and the restricted cut of Theorem 7.43 uses this. Can it be weakened to transport in terms before the bootstrap, with formula transport proved without unrestricted witnesses and without unrepaired weakening?

Question 7.82 (The size of certificates)**.** The translations are effective. No linear or polynomial bound on the size of the translated certificates is claimed; the translations regenerate a premise inside additional scopes and may duplicate it. Determine the growth, and whether sharing of regenerated subderivations in a checked directed acyclic graph reduces it without enlarging the trusted rules.

Question 7.83 (Modes without a guard)**.** Can the scope determine the witness while the mode of the introduction is also determined without a test on the matrix, with a direct proof of all native consequences? In N_{ν} the progress rule QI is kept; the result does not remove it, does not reduce the number of rules and does not settle the independence of progress.

Question 7.84 (The specialization term)**.** In all calculi of the chapter QE, or Use, still takes an arbitrary term; the proofs specialize a closure at St. By Theorems 7.7 and 7.8 the target cannot be restricted to a parameter or finitely many numerals together with the witness in the identity language. Whether a differently interpreted theory with both restrictions is bi-interpretable with iPRA is not decided, and replacing the specialization by a fixed one needs another way to deliver t.

Question 7.85 (Less normalization)**.** The proofs of Sections 7.7 and 7.8 use the literal identities $P(St) \equiv t$ and the two instances of the selector. Is there a canonical syntax that guarantees these reductions without full normalization, with its own coherence law in place of Theorem 7.58?

Further remarks of the packages on what is not shown:

- No package proves that its presentation is minimal, and none improves the compact programs of the corpus. Whether the restriction of the witness improves the compact records under a matched contract is open, and a comparison that fixes one validation contract for all competing calculi is proposed. The question of `deterministic_introduction` whether the selector by pattern can be recognized more cheaply is superseded by Theorem 7.55, which removes the selector.

- `deterministic_introduction` supplies a complete elaborator from certificates of G_{d} to certificates of G and illustrates the converse by one translated pair; `uniform_raw` supplies no translator; the translator of `conditional_copy` is not verified. The correctness of the Python programs is supported by finite tests only.

- In Theorems 7.63 and 7.73 the equivalence of the diagonal calculus with N and the bridge are used over the vocabulary extended by the fixed descriptions R. The packages argue that the proofs are parametric in additional terms; the Lean development covers the iterator terms.

- The predecessor is used in the form stated in Section 7.4: derivable in the equational fragment. The four audits recorded after Theorem 1.48 are written derivations, supported by numerical checks of the defining terms and, in `newest_parameter`, by replayed symbolic certificates of seven equation-case lemmas.

Chapter 8

Recursors with a built-in seed

The iterator $I(n, a;\, u.\, h)$ of Section 1.1 takes its initial value a as an independent argument and returns it unchanged at the counter 0. This chapter treats three arithmetic bases in which the value at 0 is tied to other data of the recursor.

- The *successor-seeded* recursor $R(n, a; u.h)$ starts at the successor of its initializer, $R(0, a; u.h) = Sa$. The successor is then the abbreviation $Sa := R(0, a; u.0)$, and the term language is $x \mid 0 \mid R(n, a; u.h)$.
- The *tied-seed* recursor $B(n; u, v.h)$ has no initializer. Its update sees the old counter u and the old state v, and its value at 0 is $S(h[0/u][0/v])$.
- The *self-seeded* recursor $F(n; v.h)$ has no initializer and its update sees only the old state; its value at 0 is $S(h[0/v])$.

The hypotheses differ from base to base, and Table 8.1 records which properties are axioms and which are derived. In every case the logical base is either ordinary intuitionistic first-order logic with equality or the native binders Q_y, $D_{x,y}$ of Section 1.3.

Theory	Logic	Axioms beyond the two recursor equations	Derived
N_R	native	none (the eight rule forms of N)	predecessor, iterator, hence everything derived in N
T_R	ordinary	coverage, $0 = S0 \to \bot$, induction for literal equations	predecessor, injectivity of S, iterator
$\mathsf{T}_{B\varepsilon}$	ordinary	Boolean equality test ε (range, reflection), $0 = S0 \to \bot$, induction for literal equations	predecessor, injectivity, quantifier-free induction, coverage
T_{Bd}	ordinary	two equations for d, coverage, $0 = S0 \to \bot$, induction for literal equations	predecessor, injectivity, reflection of equality, quantifier-free induction
N_{Bd}	native	two equations for d	the same, and the ordinary connectives
T_F	ordinary	injectivity of S, $0 = S0 \to \bot$, Boolean equality test, induction for literal equations	coverage, predecessor (as a finite-case graph)
T_F^-	ordinary	as T_F without injectivity	coverage, a right predecessor, $S^k a \neq a$ for each fixed $k \geq 1$

Table 8.1: Axioms and derived properties of the theories of this chapter.

All recursors are staged: the body h of a recursor term is a term built earlier, and each fixed body may be read as a function symbol whose arguments are the counter, the initializer

where there is one, and the free variables of the body. No theory of this chapter quantifies over bodies or has an evaluator for them. Substitution is capture-avoiding. The numerals are $1 := S0$, $2 := S1$, and so on; $S^k t$ denotes k written applications of S, and $\bar{k} := S^k 0$.

8.1 The successor-seeded recursor

This section draws on `scalar_successor`, `transient_seed` and `successor_seed_completion`, three write-ups of the same recursor.

8.1.1 Terms, equations and the two theories

Definition 8.1. The R-terms are

$$t ::= x \mid 0 \mid R(n, a; u.h),$$

where $R(n, a; u.h)$ binds u in h and nowhere else. The successor is the abbreviation $Sa := R(0, a; u.0)$. The two computation equations are

$$R(0, a; u.h) = Sa, \tag{8.1}$$
$$R(Sn, a; u.h) = h[R(n, a; u.h)/u]. \tag{8.2}$$

With the abbreviation expanded, (8.1) reads $R(0, a; u.h) = R(0, a; u.0)$ and the counter on the left of (8.2) is $R(0, n; u.0)$; no successor constructor is hidden.

(a) N_R is the native calculus with the eight rule forms of N (Section 1.3) over the R-terms: Hyp, (8.1) and (8.2) as the two computation equations, Tr, QE, DE, DI by cases and QI by progress for literal equations. The contradiction is $0 = S0$.

(b) T_R is ordinary intuitionistic first-order logic with equality over the R-terms, with (8.1), (8.2), coverage $\forall x\,(x = 0 \vee \exists y\, x = Sy)$, separation $0 = S0 \to \bot$, and induction for literal equations E with side assumptions: from $\Gamma \vdash E(0)$ and $\Gamma, E(z) \vdash E(Sz)$, with z fresh, infer $\Gamma \vdash E(t)$.

Neither theory has an axiom of successor injectivity, a predecessor, a selector or an initializer. The term language is that of the seeded recursor of Theorem 9.1, where it is studied on canonical terms with normalizing substitution; here the terms are raw and both computation equations are rules. Weakening, substitution and cut (Theorems 1.10 and 1.11) hold for N_R with the same proofs: the only cases that mention a term constructor are those of the two computation equations, and (8.1), (8.2) are closed under substitution like (1.1), (1.2). In the same way every result of Section 1.4 whose proof does not use a computation equation holds for N_R, and for the other native calculi of this chapter; these results are cited below for such calculi without further comment.

Throughout the section

$$\mathrm{sel}^+(n, a, b) := R(n, a; u.b) \quad (u \notin \mathrm{FV}(b)), \qquad Z(n) := \mathrm{sel}^+(n, 0, 0).$$

The computation equations give

$$\mathrm{sel}^+(0, a, b) = Sa, \qquad \mathrm{sel}^+(Sn, a, b) = b, \qquad Z(0) = 1, \qquad Z(Sn) = 0. \tag{8.3}$$

The zero branch of sel^+ adds one to the supplied value, so sel^+ is not the selector sel of Section 1.1.

8.1.2 Equality, cases and induction before the predecessor

Lemma 8.2. N_R *derives* $t = t$ *for every term* t*, symmetry and transitivity of equality, congruence* $s = t \vdash h(s) = h(t)$ *for every term template* h*, and transport of literal equations: from* $s = t$ *and* $l(s) = r(s)$ *infer* $l(t) = r(t)$*. It admits the rules*

$$\frac{\Gamma \vdash B(0) \qquad \Gamma \vdash B(Sz)}{\Gamma \vdash B(t)}\ (\textit{NC}), \qquad \frac{\Gamma \vdash E(0) \qquad \Gamma, E(z) \vdash E(Sz)}{\Gamma \vdash E(t)}\ (\textit{EI}),$$

where B *is any native formula,* E *is a literal equation and* z *is fresh for* Γ*,* t *and the parameters of the matrix. Under the hypothesis* $0 = S0$ *it derives every equation.*

Proof. Let $r := R(S0, 0; u.t)$ with $u \notin \mathrm{FV}(t)$. Equation (8.2) is $r = t$. Rule Tr (from $s = t$ and $h(s) = u$ infer $u = h(t)$) with the source equation $r = t$, the template $h(v) := v$ and the premise $r = t$ gives $t = t$. This replaces the two instances of (1.1) in Theorem 1.18(i). The proofs of Theorem 1.18(ii)–(v) use Tr and reflexivity only, and give symmetry, transitivity, congruence and transport of equations. Rule NC is Theorem 1.14 and rule EI is Theorem 1.20; their proofs use DI, DE, QI, QE and Hyp and no computation equation.

Under $0 = S0$, congruence gives $\mathrm{sel}^+(0, 0, s) = \mathrm{sel}^+(S0, 0, s)$, that is $S0 = s$ by (8.3), for every s. Hence $s = t$ for all s, t. □

In T_R reflexivity and replacement are rules of the logic, NC follows from coverage, EI is the induction rule, and every formula follows from $0 = S0$ by separation.

Lemma 8.3 (Uniqueness of iteration)**.** *Let* u *be a variable not free in* Γ*. If* $\Gamma \vdash h = k$*, then* $\Gamma \vdash R(n, a; u.h) = R(n, a; u.k)$ *for all terms* n*,* a*. More generally, if* $\Gamma \vdash Sa = Sb$ *and* $\Gamma \vdash h = k$*, then* $\Gamma \vdash R(n, a; u.h) = R(n, b; u.k)$*.*

Proof. By EI on a fresh counter z for the literal equation $R(z, a; u.h) = R(z, b; u.k)$. At 0 both sides are computed by (8.1) to Sa and Sb. At Sz the left side is $h[R(z, a; u.h)/u]$, which equals $h[R(z, b; u.k)/u]$ by the induction hypothesis and congruence, and this equals $k[R(z, b; u.k)/u]$ by the instance of $\Gamma \vdash h = k$ at $u := R(z, b; u.k)$, which exists because u is not free in Γ. Then substitute n for z. □

Two bodies with different syntax are thus identified by a proof, not by a primitive extensionality rule.

Lemma 8.4 (Replacement)**.** *Write* $A \approx A'$ *if* A *and* A' *are interderivable over the empty context; by weakening and substitution they are then interderivable over every context and under every substitution.*

(i) If $\vdash t = t'$ *is derivable in the empty context, and* A' *arises from* A *by replacing occurrences of* t *by* t'*, also in the scope of binders that bind variables of* t*, then* $A \approx A'$*, in* N_R *and in* N*.*

(ii) The Leibniz rule (Theorem 1.22) and transport in a formula template (Theorem 1.23) hold in N_R*.*

Proof. (ii) The proofs of Theorems 1.19, 1.21 and 1.22 use the equality rules, QE, DE, DI and the structural properties, and no computation equation.

(i) By induction on A. At a literal equation the bound variables of A that occur in t are free, and transport of equations along $t = t'$ applies. At a binder the matrices are interderivable over the empty context by the induction hypothesis, and Theorem 1.21 applies, since no variable is free in the empty context. Statement (i) is not an instance of Theorem 1.22, whose substitution is capture-avoiding. □

8.1.3 Boolean terms and contextual case analysis

The proof of the predecessor theorem splits on the value of a two-valued term inside an induction step, where the induction hypothesis is in the context and mentions that term. Rule NC does not allow this: its premises carry no equations $t = 0$, $t = Sz$, and its context may not mention the case variable. In T_R the split is a disjunction elimination. In N_R it is derived as follows, before any selector with unshifted branches exists.

Lemma 8.5 (Functional range)**.** *For every formula template $K(w)$ and every term r: if $\Gamma \vdash K(1)$ and $\Gamma \vdash K(0)$ then $\Gamma \vdash K(Z(r))$. The context may mention r.*

Proof. Apply NC to the template $w \mapsto K(Z(w))$ with w fresh. Its zero premise $K(Z(0))$ follows from $K(1)$ and $Z(0) = 1$, and its successor premise $K(Z(Sz))$ from $K(0)$ and $Z(Sz) = 0$, by Theorem 8.4(ii). Then instantiate w by r. □

Define the parity terms

$$U(n) := R(n, 0; u.Z(u)), \qquad N(n) := Z(U(n)).$$

Lemma 8.6 (Parity and the Boolean decoder)**.** N_R *derives*

$$Z(Z(Z(r))) = Z(r), \qquad N(0) = 0, \qquad N(Sn) = Z(N(n)), \qquad N(SSn) = N(n),$$
$$N(Z(r)) = Z(r), \qquad N(SSZ(r)) = Z(r).$$

Proof. The first equation is Theorem 8.5 for the template $Z(Z(w)) = w$: its instances are $Z(Z(1)) = Z(0) = 1$ and $Z(Z(0)) = Z(1) = 0$. By (8.1), $U(0) = 1$, so $N(0) = Z(1) = 0$. By (8.2), $U(Sn) = Z(U(n))$, so $N(Sn) = Z(Z(U(n))) = Z(N(n))$, and $N(SSn) = Z(Z(Z(U(n)))) = Z(U(n)) = N(n)$. In particular $N(1) = Z(0) = 1$. The equation $N(Z(r)) = Z(r)$ is Theorem 8.5 for the template $N(w) = w$, with instances $N(1) = 1$ and $N(0) = 0$. The last equation follows from the fourth and the fifth. □

Definition 8.7. A term b is *Boolean in* Γ if $\Gamma \vdash b = Z(r)$ for some term r. The numerals 0 and 1 are Boolean in every context, since $Z(1) = 0$ and $Z(0) = 1$; so is every term $N(n)$.

If b and c are Boolean in Γ, then $\Gamma, Sb = Sc \vdash b = c$: apply the template $w \mapsto N(Sw)$ to the hypothesis and use the last equation of Theorem 8.6 on both sides. This cancels a successor on Boolean terms only; it is not injectivity of S.

Lemma 8.8 (Restricted implication)**.** *Let b, c be Boolean in Γ, let $s = t$ be an equation, and put*

$$\mathcal{I}(b = c;\ s = t) := D_{x,y}\big(\mathrm{sel}^+(y, b, s) = \mathrm{sel}^+(y, c, t)\big)$$

with x, y fresh. Then

(i) if $\Gamma, b = c \vdash s = t$, then $\Gamma \vdash \mathcal{I}(b = c;\ s = t)$;

(ii) if $\Gamma \vdash \mathcal{I}(b = c;\ s = t)$ and $\Gamma \vdash b = c$, then $\Gamma \vdash s = t$.

Proof. (i) Use DI by cases with witness 0. The local assumption is the matrix at $y = 0$, which by (8.3) is $Sb = Sc$; the decoder gives $b = c$, and cutting in the given derivation gives $s = t$. By (8.3) this is the matrix at $y = Sz$.

(ii) Rule DE opens $Q_y\big(\mathrm{sel}^+(y, b, s) = \mathrm{sel}^+(y, c, t)\big)$; the opened parameter does not occur. Its zero instance is $Sb = Sc$, which follows from $b = c$ by congruence. Rule QE at the term 1 gives $\mathrm{sel}^+(1, b, s) = \mathrm{sel}^+(1, c, t)$, that is $s = t$. □

Lemma 8.9 (Contextual Boolean cases)**.** *Let b be Boolean in Γ and let E be a literal equation. If $\Gamma, b = 0 \vdash E$ and $\Gamma, b = 1 \vdash E$, then $\Gamma \vdash E$. The context and E may mention b.*

Proof. Let $\Gamma \vdash b = Z(r)$ and let w be a variable not in Γ, b, r, E. Apply NC to the native formula $B(w) := \mathcal{I}(b = Z(w); E)$. Its zero premise is Theorem 8.8(i) applied to $\Gamma, b = Z(0) \vdash E$, which is the second given branch because $Z(0) = 1$. Its successor premise is Theorem 8.8(i) applied to $\Gamma, b = Z(Sz) \vdash E$, which is the first given branch because $Z(Sz) = 0$. Rule NC gives $\Gamma \vdash B(r)$, and Theorem 8.8(ii) with $\Gamma \vdash b = Z(r)$ gives E. The context does not change during the argument; the variable that varies is the fresh w. □

A branch whose hypotheses yield $0 = S0$ is closed by the last statement of Theorem 8.2.

Remark 8.10. Theorems 8.6, 8.8 and 8.9 are the native bootstrap of `transient_seed`. `successor_seed_compl` proves the same case rule for terms with values among $0, 1, 2$: a term b has the *functional range rule* for a finite list V of numerals if $\Gamma \vdash K(r)$ for all $r \in V$ implies $\Gamma \vdash K(b)$; successor is cancelled on such terms by applying the three-cycle M of Theorem 8.12 and then j twice, using $M(w) = w$ and $j(j(j(w))) = w$ on $0, 1, 2$; and the guard formula is $D_{a,y}(\mathrm{sel}^+(y, x, u) = \mathrm{sel}^+(y, b, v))$ evaluated at $x = r$ for each $r \in V$. The predecessor proof below needs the two-valued case only. `scalar_successor` states its finite case analyses as abbreviations for repeated NC and substitution; inside an induction step they are instances of Theorem 8.9.

8.1.4 The predecessor

All statements of this subsection are derivable in N_R, by computation, the equality laws, NC, EI for the displayed literal equation with its other parameters fixed, and Theorem 8.9. They are therefore derivable in T_R. No step cancels a successor or uses a selector with unshifted branches.

Lemma 8.11 (Shifted sum)**.** *Let $a \oplus b := R(b, a; u.Su)$. Then $a \oplus 0 = Sa$, $a \oplus Sb = S(a \oplus b)$, and*

$$0 \oplus a = Sa, \qquad Sa \oplus b = S(a \oplus b), \qquad a \oplus b = b \oplus a, \qquad (a \oplus b) \oplus c = a \oplus (b \oplus c).$$

Proof. The first two equations are (8.1) and (8.2). For $0 \oplus a = Sa$ use EI on a: $0 \oplus 0 = 1$ and $0 \oplus Sa = S(0 \oplus a) = SSa$. For $Sa \oplus b = S(a \oplus b)$ use EI on b: $Sa \oplus 0 = SSa = S(a \oplus 0)$ and $Sa \oplus Sb = S(Sa \oplus b) = SS(a \oplus b) = S(a \oplus Sb)$. For commutativity use EI on b: $a \oplus 0 = Sa = 0 \oplus a$ and $a \oplus Sb = S(a \oplus b) = S(b \oplus a) = Sb \oplus a$. For associativity use EI on c: both sides are $S(a \oplus b)$ at 0, and $(a \oplus b) \oplus Sc = S((a \oplus b) \oplus c) = S(a \oplus (b \oplus c)) = a \oplus (b \oplus Sc)$. □

In the standard model $a \oplus b = a + b + 1$. Define the affine terms

$$A_2(a) := S^2(a \oplus a), \qquad A_3(a) := S^4((a \oplus a) \oplus a), \qquad A_2^\circ(a) := S^3(a \oplus a),$$

with standard values $2a + 3$, $3a + 6$ and $2a + 4$. Theorem 8.11 gives

$$A_2(0) = 3, \qquad A_2(Sa) = S^2 A_2(a), \qquad A_3(0) = 6, \qquad A_3(Sa) = S^3 A_3(a). \tag{8.4}$$

Lemma 8.12 (A three-cycle)**.** *Let*

$$j(u) := \mathrm{sel}^+(u, 0, \mathrm{sel}^+(Z(N(u)), 1, 0)), \qquad X(n) := R(n, 0; u.j(u)), \qquad M(n) := j(j(X(n))).$$

Then $j(0) = 1$, $j(1) = 2$, $j(2) = 0$, $M(0) = 0$ and $M(Sn) = j(M(n))$.

Proof. By (8.3), $j(0) = S0$ and $j(Sz) = \mathrm{sel}^+(Z(N(Sz)), 1, 0)$. Since $N(1) = 1$ and $N(2) = N(0) = 0$, this gives $j(1) = \mathrm{sel}^+(0, 1, 0) = 2$ and $j(2) = \mathrm{sel}^+(1, 1, 0) = 0$. Further $X(0) = 1$ and $X(Sn) = j(X(n))$, so $M(0) = j(j(1)) = 0$ and $M(Sn) = j(j(j(X(n)))) = j(M(n))$. □

Lemma 8.13 (The ramp)**.** *Let* $G(n) := R(n, 0; u.\, u \oplus M(S^2 u))$. *Then*

$$M(G(n)) = S(N(n)), \tag{8.5}$$

$$N(n) = 0 \vdash G(Sn) = SG(n), \qquad N(n) = 1 \vdash G(Sn) = S^2 G(n), \tag{8.6}$$

$$G(S^2 n) = S^3 G(n), \tag{8.7}$$

$$N(n) = 0 \vdash G(S^3 n) = S^4 G(n), \qquad N(n) = 1 \vdash G(S^3 n) = S^5 G(n). \tag{8.8}$$

Proof. By the computation equations, $G(0) = 1$ and $G(Sn) = G(n) \oplus M(S^2 G(n)) = G(n) \oplus j(j(M(G(n))))$.

Equation (8.5) is proved by EI on n. The base is $M(1) = j(M(0)) = j(0) = 1 = S(N(0))$. For the step keep the induction hypothesis $M(G(n)) = S(N(n))$ in the context and apply Theorem 8.9 to the Boolean term $N(n)$. If $N(n) = 0$, then $M(G(n)) = 1$, $j(j(1)) = 0$, so $G(Sn) = G(n) \oplus 0 = SG(n)$ and $M(G(Sn)) = j(M(G(n))) = j(1) = 2 = S(Z(0)) = S(N(Sn))$. If $N(n) = 1$, then $M(G(n)) = 2$, $j(j(2)) = 1$, so $G(Sn) = G(n) \oplus 1 = S^2 G(n)$ and $M(G(Sn)) = j(j(M(G(n)))) = j(j(2)) = 1 = S(Z(1)) = S(N(Sn))$.

The two computations of $G(Sn)$ just made, now with (8.5) as a theorem, are (8.6). For (8.7) apply Theorem 8.9 to $N(n)$: if $N(n) = 0$ then $N(Sn) = 1$ and $G(S^2 n) = S^2 G(Sn) = S^3 G(n)$; if $N(n) = 1$ then $N(Sn) = 0$ and $G(S^2 n) = SG(Sn) = S^3 G(n)$. Finally $G(S^3 n) = S^3 G(Sn)$ by (8.7), and (8.6) gives (8.8). □

From $G(0) = 1$, $N(0) = 0$ and (8.6), (8.7) one computes $G(1) = 2$, $G(2) = 4$, $G(3) = 5$, $G(4) = 7$, $G(6) = 10$. In the standard model $G(n) = \lfloor 3n/2 \rfloor + 1$.

Lemma 8.14 (Affine identities)**.**

$$N(A_2(a)) = 1, \qquad N(A_3(a)) = N(a), \qquad SG(A_2(a)) = A_3(a),$$
$$G(A_3(a)) = A_3(G(a)) \oplus N(a).$$

Proof. Each equation is proved by EI on a with (8.4).

$N(A_2(0)) = N(3) = N(1) = 1$, and $N(A_2(Sa)) = N(S^2 A_2(a)) = N(A_2(a))$.

$N(A_3(0)) = N(6) = N(0)$, and $N(A_3(Sa)) = N(S^3 A_3(a)) = Z(N(A_3(a))) = Z(N(a)) = N(Sa)$.

$SG(A_2(0)) = SG(3) = 6 = A_3(0)$, and $SG(A_2(Sa)) = SG(S^2 A_2(a)) = S^3 SG(A_2(a)) = S^3 A_3(a) = A_3(Sa)$ by (8.7).

For the last equation the base is $G(A_3(0)) = G(6) = 10$ and $A_3(G(0)) \oplus N(0) = A_3(1) \oplus 0 = S9 = 10$. For the step keep the induction hypothesis $G(A_3(a)) = A_3(G(a)) \oplus N(a)$ and apply Theorem 8.9 to $N(a)$; the left side of the goal is $G(S^3 A_3(a))$ and $N(A_3(a)) = N(a)$.

If $N(a) = 0$, then by (8.8) and the hypothesis

$$G(S^3 A_3(a)) = S^4 G(A_3(a)) = S^4(A_3(G(a)) \oplus 0) = S^5 A_3(G(a)),$$

while $G(Sa) = SG(a)$, $N(Sa) = 1$ and

$$A_3(G(Sa)) \oplus N(Sa) = S^3 A_3(G(a)) \oplus 1 = S^5 A_3(G(a)).$$

If $N(a) = 1$, then

$$G(S^3 A_3(a)) = S^5 G(A_3(a)) = S^5(A_3(G(a)) \oplus 1) = S^7 A_3(G(a)),$$

while $G(Sa) = S^2 G(a)$, $N(Sa) = 0$ and

$$A_3(G(Sa)) \oplus N(Sa) = S^6 A_3(G(a)) \oplus 0 = S^7 A_3(G(a)).$$ □

Lemma 8.15 (The factor map)**.** *Let* $H(u) := \mathrm{sel}^+(Z(N(u)), G(u), u)$*. Then*

$$N(u) = 0 \vdash H(u) = u, \qquad N(u) = 1 \vdash H(u) = SG(u),$$
$$H(2) = 2, \qquad H(A_2(a)) = A_3(a), \qquad H(A_3(a)) = A_3(H(a)).$$

Proof. The first line is (8.3) with $Z(0) = 1$ and $Z(1) = 0$. Since $N(2) = 0$, $H(2) = 2$. Since $N(A_2(a)) = 1$, $H(A_2(a)) = SG(A_2(a)) = A_3(a)$. For the last equation apply Theorem 8.9 to $N(a)$, using $N(A_3(a)) = N(a)$. If $N(a) = 0$, both sides are $A_3(a)$. If $N(a) = 1$, then by Theorem 8.14

$$H(A_3(a)) = SG(A_3(a)) = S(A_3(G(a)) \oplus 1) = S^3 A_3(G(a)) = A_3(SG(a)) = A_3(H(a)). \quad \square$$

On standard numbers H fixes even inputs and sends an odd u to $3(u+1)/2$. In the coordinate $u+3$ the terms A_2 and A_3 are doubling and tripling, and an active step of H multiplies by $3/2$. This explains the comparator below; the proof does not use it.

Lemma 8.16 (A comparator)**.** *Let*

$$\alpha(x) := R(x, 0; u.A_2^\circ(u)), \qquad e(x) := S\alpha(x), \qquad \Phi(x,y) := R(y, \alpha(x); u.H(u)), \qquad L(x,y) := Z(N(\Phi(x,y)))$$

Then $L(x,y)$ *is Boolean, and*

$$L(0,y) = 1, \qquad L(Sx, 0) = 0, \qquad L(Sx, Sy) = L(x,y), \qquad L(Sn, n) = 0.$$

Proof. By computation and Theorem 8.11, $e(0) = S\alpha(0) = 2$ and

$$e(Sx) = SA_2^\circ(\alpha(x)) = S^4(\alpha(x) \oplus \alpha(x)) = S^2(S\alpha(x) \oplus S\alpha(x)) = A_2(e(x)).$$

Also $\Phi(x,0) = S\alpha(x) = e(x)$ by (8.1), and $\Phi(x, Sy) = H(\Phi(x,y))$. The initializer of Φ is chosen so that the successor-seeded initial value is $e(x)$; no predecessor is needed to start the orbit there.

By EI on y, $\Phi(0,y) = 2$: the base is $e(0) = 2$ and the step is $H(2) = 2$. Hence $L(0,y) = Z(N(2)) = Z(0) = 1$. Next $\Phi(Sx, 0) = A_2(e(x))$ has N-value 1, so $L(Sx, 0) = Z(1) = 0$.

By EI on y with x fixed, $\Phi(Sx, Sy) = A_3(\Phi(x,y))$. At $y = 0$ the left side is $H(e(Sx)) = H(A_2(e(x))) = A_3(e(x)) = A_3(\Phi(x,0))$. In the step both orbits apply H, and $H(A_3(v)) = A_3(H(v))$ turns $H(A_3(\Phi(x,y)))$ into $A_3(\Phi(x,Sy))$. Applying $Z \circ N$ and $N(A_3(v)) = N(v)$ gives $L(Sx,Sy) = L(x,y)$.

Finally $L(Sn, n) = 0$ by EI on n from the second and third equations. $\square$

The standard value of $L(x,y)$ is 1 if $x \le y$ and 0 otherwise.

Lemma 8.17 (Saturation)**.** *Let*

$$K(c,u) := \mathrm{sel}^+(c,u,u), \qquad \beta_m(u) := K(Z(L(Su, m)), u), \qquad V(t,m) := R(t, 0; u.\beta_m(u)).$$

Then

$$V(St, S^2 k) = SV(t, Sk), \qquad V(n, Sn) = Sn, \qquad V(Sn, Sn) = V(S^2 n, Sn) = Sn.$$

Proof. $K(0,u) = Su$ and $K(Sv, u) = u$, so NC on c gives $K(c, Su) = SK(c,u)$. With the comparator equations,

$$\beta_{Sm}(Su) = K(Z(L(Su,m)), Su) = S\beta_m(u), \qquad \beta_{S^2k}(1) = K(Z(L(0,k)), 1) = K(0,1) = 2.$$

The first equation of the lemma is EI on t with k fixed. At $t = 0$, $V(1, S^2k) = \beta_{S^2k}(V(0, S^2k)) = \beta_{S^2k}(1) = 2 = SV(0, Sk)$. In the step, $V(S^2t, S^2k) = \beta_{S^2k}(SV(t,Sk)) = S\beta_{Sk}(V(t,Sk)) = SV(St, Sk)$.

The second equation is a separate EI on n: $V(0,1)=1$, and $V(Sn,S^2n)=SV(n,Sn)$ by the first equation at $t=k=n$, an instance of a completed lemma. For the third,

$$\beta_{Sn}(Sn)=K(Z(L(S^2n,Sn)),Sn)=K(Z(L(Sn,n)),Sn)=K(1,Sn)=Sn,$$

so one and then two further steps of the recursor from $V(n,Sn)=Sn$ leave the value unchanged. □

Lemma 8.18 (A capped predecessor)**.** *Let* $Q(m):=R(m,0;u.K(Z(L(S^2u,m)),u))$. *Then* $Q(0)=Q(1)=1$ *and* $Q(S^2k)=Sk$.

Proof. $Q(0)=1$ by (8.1), and $Q(1)=K(Z(L(3,1)),1)=K(Z(L(2,0)),1)=K(1,1)=1$. For $m=S^2k$ the comparator equations give $L(S^2u,S^2k)=L(u,k)=L(Su,Sk)$, so the body of Q at this parameter equals $\beta_{Sk}(u)$ for a generic u. By Theorem 8.3, at an independent counter t with k fixed,

$$R(t,0;u.K(Z(L(S^2u,S^2k)),u))=V(t,Sk).$$

Substituting $t=S^2k$ and Theorem 8.17 give $Q(S^2k)=V(S^2k,Sk)=Sk$. □

Theorem 8.19 (Predecessor)**.** *Let*

$$P(m):=\mathrm{sel}^{+}\big(Z(L(3,m)),\ Q(Q(m)),\ L(2,m)\big).$$

N_R *and* T_R *derive*

$$P(0)=0,\qquad P(Sn)=n.$$

Consequently both derive $Sx=Sy\vdash x=y$.

Proof. For $m\in\{0,1,2\}$ the comparator equations give $L(3,m)=0$, so $Z(L(3,m))=1$ and $P(m)=L(2,m)$ by (8.3); and $L(2,0)=0$, $L(2,1)=L(1,0)=0$, $L(2,2)=L(0,0)=1$. Hence $P(0)=0$, $P(1)=0$ and $P(2)=1$. For $m=S^3k$, $L(3,m)=L(0,k)=1$, so the zero branch of sel^{+} applies and, by Theorem 8.18,

$$P(S^3k)=SQ(Q(S^3k))=SQ(S^2k)=S^2k.$$

Rule NC on n, and in its successor case NC once more, give $P(Sn)=n$ from $P(1)=0$, $P(2)=1$ and $P(S^3k)=S^2k$. Injectivity follows by applying P to $Sx=Sy$. □

Remark 8.20 (The three constructions). The construction above is that of `successor_seed_completion`: the orbit G is started at the seed 1 that (8.1) provides, the affine terms absorb the shift, and the capped predecessor Q is corrected at three small arguments. The two other packages reach Theorem 8.19 by different terms.

`scalar_successor` first builds a term τ with $\tau(0)=1$, $\tau(1)=0$, $\tau(S^2n)=S^2n$ and $\tau(\tau(n))=n$, from the parity term U, a Boolean conjunction $Z(\mathrm{sel}^{+}(a,0,Z(b)))$ and two finite orbits. Conjugation gives an iterator that starts at zero,

$$K_0(n;u.h):=\tau(R(n,0;v.\tau(h[\tau(v)/u]))),\qquad K_0(0;u.h)=0,\quad K_0(Sn;u.h)=h[K_0(n;u.h)/u],$$

for every body h. With the shifted sum $a\oplus b$, the terms $a\oplus a$ and $(a\oplus a)\oplus a$, a ramp $F(n):=K_0(n;u.\,u\oplus M'(u))$ over a three-state orbit M' with the invariant $M'(F(n))=N(n)$, and a factor map, it obtains a strict comparator V with $V(x,0)=0$, $V(0,Sy)=1$, $V(Sx,Sy)=V(x,y)$. The predecessor is then the saturated zero-start orbit $P(m):=K_0\big(m;u.\,\mathrm{sel}^{+}(Z(V(Su,m)),u,u)\big)$, with no correction.

`transient_seed` works in T_R and observes that $f(R(n,a;u.f))$ starts at $f(Sa)$: a seed Sa outside an intended orbit is used once to enter it. Doubling and tripling orbits $d_2(Sn)=$

$S^2 d_2(n)$, $d_3(Sn) = S^3 d_3(n)$ that start at zero are obtained in this way, then a ramp and a factor map, and a comparator L with the equations of Theorem 8.16 and the persistence law $L(x,y) = 1 \to L(x,Sy) = 1$. The predecessor is a reset cycle: with $F_m(u) := \mathrm{sel}^+(Z(L(Su,m)),u,0)$, which increments below m and resets to 0 from m on, and

$$J_m(t) := F_m\big(R(t,m;u.F_m(u))\big), \qquad K_m(t) := F_m\big(R(t,m;u.F_m(F_m(u)))\big),$$

one has $J_m(0) = K_m(0) = 0$ because $F_m(Sm) = 0$. Induction on the literal equation $\mathrm{sel}^+(L(t,m),0,J_m(t)) = \mathrm{sel}^+(L(t,m),0,t)$ gives $L(t,m) = 1 \to J_m(t) = t$; two further inductions give the period $J_m(m \oplus t) = J_m(t)$ and $K_m(Sa) = J_m(a \oplus Sa)$; and $P(n) := K_n(n)$ satisfies $P(Sa) = J_{Sa}(a) = a$. The package then transfers the proof to N_R by Theorem 8.9.

The three proofs use the same resources: literal-equation induction with fixed parameters, numerical cases, and Boolean cases in a context.

8.1.5 Recovery of the iterator

Theorem 8.21 (Initialized iteration)**.** *For every body h and a fresh variable v put*

$$\widehat{I}(n,a;u.h) := P\big(R(n,a;\, v.\, S(h[P(v)/u]))\big).$$

N_R and T_R derive

$$\widehat{I}(0,a;u.h) = a, \qquad \widehat{I}(Sn,a;u.h) = h[\widehat{I}(n,a;u.h)/u].$$

In particular $\widehat{I}(c,a;u.b)$ with $u \notin \mathrm{FV}(b)$ is a selector with the unshifted values a and b.

Proof. Write $r(n) := R(n,a;v.S(h[P(v)/u]))$. By (8.1) and Theorem 8.19, $\widehat{I}(0,a;u.h) = P(Sa) = a$. By (8.2), $r(Sn) = S(h[P(r(n))/u])$, so $\widehat{I}(Sn,a;u.h) = P(S(h[P(r(n))/u])) = h[P(r(n))/u] = h[\widehat{I}(n,a;u.h)/u]$. The equation $P(Sx) = x$ is used at arbitrary terms; no invariant about the state $r(n)$ is needed. □

Remark 8.22. The recursor, the candidate equations and the compiler $\widehat{I}$ are stated in an earlier draft of the series, which leaves the term P open; the three packages supply it. In the opposite direction the predecessor is the only obstacle: a successor-seeded recursor whose update also receives the old counter has an immediate predecessor (see Theorem 8.28 for the analogous computation).

8.1.6 Comparison of the two term languages

Definition 8.23. The translation $t \mapsto t^\iota$ from R-terms to the terms of Section 1.1 and the translation $t \mapsto t^\rho$ in the opposite direction are defined by

$$x^\iota := x, \qquad 0^\iota := 0, \qquad R(n,a;u.h)^\iota := I(n^\iota, S(a^\iota);\, u.\, h^\iota),$$
$$x^\rho := x, \qquad 0^\rho := 0, \qquad (St)^\rho := S(t^\rho), \qquad I(n,a;\, u.\, h)^\rho := \widehat{I}(n^\rho, a^\rho; u.h^\rho),$$

where $S(t^\rho)$ on the right is the abbreviation $R(0,t^\rho;u.0)$. Formulas, native or ordinary, are translated by translating their terms. Both translations commute with substitution and preserve free variables and the property of being a literal equation.

Note that $(Sa)^\iota = I(0,S(a^\iota);\, u.\, 0)$, which is not literally $S(a^\iota)$ but equals it by the zero computation equation of the iterator.

Theorem 8.24 (Equivalence of the term languages)**.**

(a) If $\Gamma \vdash A$ is derivable in N_R, then $\Gamma^\iota \vdash A^\iota$ is derivable in N. If $\Gamma \vdash A$ is derivable in T_R, then $\Gamma^\iota \vdash A^\iota$ is derivable in T_{eq}.

(b) If $\Gamma \vdash A$ is derivable in N, then $\Gamma^\rho \vdash A^\rho$ is derivable in N_R. If $\Gamma \vdash A$ is derivable in T_{eq}, then $\Gamma^\rho \vdash A^\rho$ is derivable in T_R.

(c) N_R and T_R derive $t^{\iota\rho} = t$ for every R-term t; N and T_{eq} derive $t^{\rho\iota} = t$ for every iterator term t. Hence $A^{\iota\rho} \approx A$ in N_R and $A^{\rho\iota} \approx A$ in N for native formulas, and $A^{\iota\rho} \leftrightarrow A$, $A^{\rho\iota} \leftrightarrow A$ are provable in T_R, T_{eq} for ordinary formulas.

(d) Consequently $\Gamma \vdash A$ is derivable in N_R if and only if $\Gamma^\iota \vdash A^\iota$ is derivable in N, and likewise for T_R and T_{eq}, and for $(\cdot)^\rho$ in the other direction. The contradictions correspond: $(0 = S0)^\iota$ is interderivable with $0 = S0$, and $(0 = S0)^\rho$ is $0 = S0$.

Proof. (a) By induction on the derivation. Each rule instance other than a computation equation is translated into an instance of the same rule, because the translation commutes with substitution, keeps literal equations literal and keeps parameters fresh. One adjustment is needed where a rule displays a successor: the premise of DI by cases translates to $\Gamma^\iota, B^\iota(t^\iota, 0) \vdash B^\iota(t^\iota, I(0, Sz;\, u.\,0))$, and the premise of QI, or of the induction rule of T_R, to $\Gamma^\iota, E^\iota(z) \vdash E^\iota(I(0, Sz;\, u.\,0))$. Replacing $I(0, Sz;\, u.\,0)$ by Sz along the zero computation equation, by the Leibniz property of Section 1.4, gives the premise of the same rule in N or T_{eq}. The translation of (8.1) is $I(0, Sa^\iota;\, u.\,h^\iota) = I(0, Sa^\iota;\, u.\,0)$; both sides equal Sa^ι by the zero computation equation. The translation of (8.2) is

$$I(I(0, Sn^\iota;\, u.\,0), Sa^\iota;\, u.\,h^\iota) = h^\iota[I(n^\iota, Sa^\iota;\, u.\,h^\iota)/u],$$

which follows from $I(0, Sn^\iota;\, u.\,0) = Sn^\iota$, congruence and the successor computation equation. For T_R the logical rules are translated identically; coverage translates to $\forall x\,(x = 0 \vee \exists y\, x = I(0, Sy;\, u.\,0))$ and separation to $0 = I(0, S0;\, u.\,0) \to \bot$, both equivalent in T_{eq} to the corresponding axioms.

(b) By induction on the derivation. Here $(St)^\rho$ is literally $S(t^\rho)$, so every rule instance is translated into an instance of the same rule, and coverage, separation and induction are translated into themselves. The two computation equations of the iterator are translated into $\widehat{I}(0, a^\rho; u.h^\rho) = a^\rho$ and $\widehat{I}(Sn^\rho, a^\rho; u.h^\rho) = h^\rho[\widehat{I}(n^\rho, a^\rho; u.h^\rho)/u]$, which are instances of Theorem 8.21.

(c) First, $t^{\iota\rho} = t$ by induction on the R-term t. The cases of variables and 0 are trivial. For $t = R(n, a; u.h)$,

$$t^{\iota\rho} = P\Big(R(n^{\iota\rho}, S(a^{\iota\rho});\, v.\,S(h^{\iota\rho}[P(v)/u]))\Big).$$

By the induction hypothesis, congruence and Theorem 8.3 (for the body, whose induction hypothesis $h^{\iota\rho} = h$ is derivable in the empty context with u free), this equals $P\big(R(n, Sa; v.S(h[P(v)/u]))\big)$. The equation

$$P\big(R(z, Sa;\, v.\,S(h[P(v)/u]))\big) = R(z, a; u.h)$$

is proved by EI on z: at 0 the left side is $P(SSa) = Sa = R(0, a; u.h)$; at Sz it is $P(S(h[P(R(z, Sa; \ldots))/u])) = h[P(R(z, Sa; \ldots))/u]$, which by the induction hypothesis is $h[R(z, a; u.h)/u] = R(Sz, a; u.h)$.

Second, $t^{\rho\iota} = t$ by induction on the iterator term t. By (a) applied to Theorem 8.19, the term template $P^\iota(m) := (P(m))^\iota$ satisfies $P^\iota(0) = 0$ and $P^\iota(I(0, Sx;\, u.\,0)) = x$, hence $P^\iota(Sx) = x$, in N and T_{eq}. For $t = Ss$, $t^{\rho\iota} = I(0, S(s^{\rho\iota});\, u.\,0) = S(s^{\rho\iota}) = Ss$. For $t = I(n, a;\, u.\,h)$,

$$t^{\rho\iota} = P^\iota\Big(I(n^{\rho\iota}, S(a^{\rho\iota});\, v.\,I(0, S(h^{\rho\iota}[P^\iota(v)/u]);\, w.\,0))\Big),$$

which by the induction hypothesis, the zero computation equation and uniqueness of iteration for the iterator (the proof of Theorem 8.3 with the iterator equations) equals $P^\iota\big(I(n, Sa;\, v.\,S(h[P^\iota(v)/u]))\big)$. Induction for the literal equation $P^\iota\big(I(z, Sa;\, v.\,S(h[P^\iota(v)/u]))\big) = I(z, a;\, u.\,h)$ proves the rest: at 0 both sides are a, and at Sz the left side is $h[P^\iota(I(z, Sa;\, v.\,\ldots))/u]$, which by the induction hypothesis is $h[I(z, a;\, u.\,h)/u] = I(Sz, a;\, u.\,h)$.

The statements on formulas follow from Theorem 8.4(i), in N_R and in N, and from replacement of provably equal terms under quantifiers in ordinary logic.

(d) If $\Gamma^\iota \vdash A^\iota$ is derivable in N, then $\Gamma^{\iota\rho} \vdash A^{\iota\rho}$ is derivable in N_R by (b), and $\Gamma \vdash A$ follows by (c) and cut. The other cases are the same. For the contradiction, $(0 = S0)^\iota$ is $0 = I(0, S0; u.\, 0)$. □

Corollary 8.25. *N_R and T_R are bi-interpretable with N, T_{eq} and iPRA: the interpretations are the translations of Theorem 8.23 composed with those of Theorem 1.78; the domains are the whole domain, equality is actual equality, each term is provably equal to its double translation, so that the graph $x = y$ is the isomorphism of both numerical composites, and every formula is provably equivalent to its double translation. The theories are equiconsistent. In iPRA the recursor with body h is the function defined by primitive recursion with initial value Sa and a step that ignores the counter.*

Proof. Theorem 8.24 gives the statement for N and T_{eq} with the identity as numerical map. Composition with Theorem 1.78 gives the rest; the description in iPRA is the composite of $(\cdot)^\iota$ with the interpretation of the iterator. □

Consequently every result that Theorems 1.48, 1.52 and 1.62 state for T_{eq} holds in T_R for the translated terms: reflection of equality into a term, induction for quantifier-free formulas, pairing, and a term with the defining equations of each primitive recursive description. The predecessor of Theorem 1.48 is not needed: Theorem 8.19 provides one directly.

Remark 8.26 (Rule forms in the packages). The three packages state the native calculus with two rule forms that differ from Section 1.3: transport is formula transport, from $s = t$ and $F(s)$ infer $F(t)$ for every native formula template F; and progress introduces the paired binder, from $\Gamma, E(t, z) \vdash E(t, Sz)$ infer $\Gamma \vdash D_{x,y}E(x, y)$ for a literal equation E. Both calculi derive the same sequents. Formula transport is Theorem 8.4(ii). The progress rule on $D_{x,y}$ is derived from QI: the premise gives $Q_yE(t, y)$, and DI by cases with witness t follows because its premise $E(t, 0) \vdash E(t, Sz)$ is QE. Conversely QI is derived from the progress rule on $D_{v,y}E(y)$ with v unused, followed by DE and Hyp; and the transport rule of Section 1.3 is a case of formula transport combined with symmetry. `successor_seed_completion` and `scalar_successor` also describe a *computation-supply* presentation used by their checkers: instead of a computation leaf, a rule passes from $\Gamma, H \vdash A$ to $\Gamma \vdash A$ for a computation equation H. The two presentations derive the same sequents by cut in one direction and Hyp in the other.

8.2 Tied-seed recursors

This section draws on `bounded_initializer`, `monus_and_neutral_major` and the obstruction section of `transient_seed`.

8.2.1 The recursor and its elementary terms

Definition 8.27. The tied-seed recursor $B(n; u, v.h)$ binds the distinct variables u (old counter) and v (old state) in h. Its computation equations are

$$B(0; u, v.h) = S\big(h[0/u][0/v]\big), \tag{8.9}$$

$$B(Sn; u, v.h) = h[n/u][B(n; u, v.h)/v]. \tag{8.10}$$

The same body determines the value at 0 and every later transition. Put

$$C(n, a) := B(n; u, v.a), \quad W(n) := B(n; u, v.u), \quad P(n) := W(C(n, n)), \quad Z(n) := C(n, 0),$$

with $u, v \notin \mathrm{FV}(a)$.

Lemma 8.28. *From* (8.9), (8.10) *and the equality laws alone,*

$$C(0,a) = Sa, \quad C(Sn,a) = a, \quad W(0) = 1, \quad W(Sn) = n, \quad P(0) = 0, \quad P(Sn) = n, \quad Z(0) = 1, \quad Z(Sn) =$$

Hence $Sx = Sy \vdash x = y$, *and* $Sx = 0 \vdash 0 = S0$.

Proof. The first four equations are instances of the computation equations. $P(0) = W(C(0,0)) = W(1) = 0$ and $P(Sn) = W(C(Sn,Sn)) = W(Sn) = n$. Applying P to $Sx = Sy$ gives $x = y$. From $Sx = 0$, applying P gives $x = 0$, and substitution gives $S0 = 0$. □

The successor can be removed from the primitive signature: $\sigma(a) := C(0,a)$ satisfies $\sigma(a) = Sa$, and writing σ for S in (8.9), (8.10) and in every axiom gives an equivalent presentation on the terms $x \mid 0 \mid B(n;u,v.h)$ and the other primitive operations. The constant-body instance of (8.9) becomes tautological; (8.10) for it and all other instances keep their content. The section keeps S for readability.

Lemma 8.29 (Gate and zero-start recursion)**.** *Assume induction for literal equations (rule* EI *of Theorem 8.2). Let*

$$G(n,a) := P(P(B(a;u,v.C(Z(n),v)))), \qquad B_0(n;u,v.h) := B\big(Sn;\, r,s.\, G(r, h[P(r)/u][s/v])\big).$$

Then

$$G(0,a) = 0, \qquad G(Sn,a) = a, \qquad B_0(0;u,v.h) = 0, \qquad B_0(Sn;u,v.h) = h[n/u][B_0(n;u,v.h)/v].$$

Proof. For the control 0 the body of the inner recursor is $C(1,v) = v$, its value at 0 is $S(C(1,0)) = 1$, and EI on a shows that the value at a is 1; so $G(0,a) = P(P(1)) = 0$. For the control Sn the body is $C(0,v) = Sv$, the value at 0 is $S(S0) = 2$, and EI on a shows that the value at a is SSa; so $G(Sn,a) = P(P(SSa)) = a$. In both inductions n is a fixed parameter.

Let $k(r,s) := G(r, h[P(r)/u][s/v])$, so that $k(0,s) = 0$ and $k(Sr,s) = h[r/u][s/v]$. Then $B(0;r,s.k) = S(k(0,0)) = 1$, $B(1;r,s.k) = k(0,1) = 0$, and $B(SSn;r,s.k) = k(Sn, B(Sn;r,s.k)) = h[n/u][B(Sn;r,s.k)/v]$. These are the equations for B_0. □

The gate G is a one-sided selector: it returns 0 or its second argument. The term $P(P(B(a;u,v.C(n,v))))$ is the complementary mask, with values a at $n = 0$ and 0 at $n = Sm$.

8.2.2 The one-coordinate bound

The independent initializer cannot be recovered by a term.

Definition 8.30. A function $f\colon \mathbb{N}^r \to \mathbb{N}$ has the *bounding coordinate* i if there is a nondecreasing $F\colon \mathbb{N} \to \mathbb{N}$ with $f(x_1,\dots,x_r) \le F(x_i)$ for all arguments.

Theorem 8.31 (One-coordinate bound)**.** *Consider the terms built from variables,* 0, S, B, *and optionally a binary* ε *with values in* $\{0,1\}$ *and a binary* d *with* $d(x,y) \le x$, *in the standard model with* (8.9), (8.10). *The function denoted by a term is bounded by a constant or has a bounding coordinate among its free variables, with a nondecreasing primitive recursive bound that is obtained by recursion on the term.*

Proof. By induction on the term, treating the bound variables u, v of a body as further input coordinates while the body is processed. A variable has the bound $F(z) = z$ in its own coordinate; 0 and $\varepsilon(s,t)$ are bounded by constants; St and $d(s,t)$ inherit the coordinate of t and of s, with bounds $F+1$ and F.

Let $t = B(n;u,v.h)$ and write $b(m,\vec{p})$ for the value of the recursor with body h at the counter value m. By the induction hypothesis h, as a function of $u,v,\vec{p}$, is bounded by a constant or has a bounding coordinate with bound H.

1. If $h \le K$ for a constant K, then $b(0, \vec{p}) \le K + 1$ and $b(m + 1, \vec{p}) \le K$.
2. If a parameter p_j is the bounding coordinate, then $h(0, 0, \vec{p})$ and every $h(m, b(m, \vec{p}), \vec{p})$ are at most $H(p_j)$, so $b(m, \vec{p}) \le H(p_j) + 1$ for all m. This holds whatever the counter term is.
3. If u is the bounding coordinate, put $K(m) := 1 + \max_{q \le m} H(q)$. Then $b(0, \vec{p}) \le H(0) + 1 \le K(0)$ and $b(m + 1, \vec{p}) \le H(m) \le K(m + 1)$.
4. If v is the bounding coordinate, put $K(0) := H(0)+1$ and $K(m+1) := \max\{K(m), H(K(m))\}$. Then $b(0, \vec{p}) \le H(0) + 1$, because the state argument of $h(0, 0, \vec{p})$ is 0, and $b(m + 1, \vec{p}) \le H(b(m, \vec{p})) \le H(K(m)) \le K(m + 1)$ by induction on m and monotonicity of H.

In cases 3 and 4, K is nondecreasing and primitive recursive and $t \le K(n(\vec{p}))$. By the induction hypothesis for the counter term n, either $n \le N$ for a constant N, and $K(N)$ bounds t, or $n(\vec{p}) \le N(p_i)$ for a nondecreasing primitive recursive N, and $K(N(p_i))$ bounds t in the coordinate p_i. □

Corollary 8.32. *No term of these languages denotes, in the standard model, any of the following functions: the selector* $\mathrm{sel}(n, a, b)$ *with* $\mathrm{sel}(0, a, b) = a$ *and* $\mathrm{sel}(n + 1, a, b) = b$*; addition; an injection* $\mathbb{N}^2 \to \mathbb{N}$.

Proof. The selector is unbounded. The coordinate n does not bound it, since at $n = 0$ it is a; the coordinate a does not, since at $a = 0$, $n = 1$ it is b; the coordinate b does not, since at $b = 0$, $n = 0$ it is a. Addition is unbounded when either argument is fixed at 0. If a coordinate bounds a binary function, fixing that coordinate at 0 confines infinitely many pairs to finitely many values; a bounded function is not injective either. □

Remark 8.33. The theorem is in `transient_seed` for the terms $x \mid 0 \mid B$, and in `bounded_initializer` with S and ε; the case of d is immediate from $d(x, y) \le x$. The same four cases, with the value 1 at the counter 0 in place of $1 + h(0, 0, \vec{p})$, give the bound for the recursor $J(0; h) = 1$, $J(Sn; h) = h(n, J(n; h))$, which `scalar_successor` mentions as the reason for keeping the initializer in R; additional unary primitive recursive operations compose with the bound of their argument and do not change the argument. The bound concerns the functions denoted by terms. Sound logical rules cannot change it, so there is no compiler from the iterator language to these terms. It does not show that a theory on these terms fails to be bi-interpretable with iPRA: Theorem 8.44 interprets each primitive recursive function by a graph with finitely many value terms. The successor-seeded recursor $R(n, a; u.h)$ escapes the bound because $a \oplus b$ is unbounded in each of its two arguments. Related obstructions for tied seeds are in the packages `local_scopes` and `uniform_predecessor`.

8.2.3 The theory with a Boolean equality test

Definition 8.34. $\mathsf{T}_{B\varepsilon}$ is ordinary intuitionistic first-order logic with equality over the terms $t ::= x \mid 0 \mid St \mid \varepsilon(t, t) \mid B(n; u, v.h)$, with the axioms (8.9), (8.10),

$$\varepsilon(x, y) = 0 \vee \varepsilon(x, y) = 1, \tag{8.11}$$

$$\varepsilon(x, y) = 1 \leftrightarrow x = y, \tag{8.12}$$

$$0 = 1 \to \bot, \tag{8.13}$$

universally closed, and induction for literal equations with side assumptions. Coverage, injectivity of the successor and induction for compound formulas are not axioms.

The equality on the right of (8.12) is the equality of the logic on the whole domain. The axioms make numerical equality decidable and give $\varepsilon(s, t) = 0 \leftrightarrow s \neq t$; they do not give excluded middle for quantified formulas.

Lemma 8.35 (Quantifier-free induction and coverage)**.** *In* $\mathsf{T}_{B\varepsilon}$ *every quantifier-free formula* A *has a term* χ_A *with* $\chi_A = 0 \vee \chi_A = 1$ *and* $A \leftrightarrow \chi_A = 1$*. Induction with side assumptions holds for quantifier-free formulas, and* $x = 0 \vee x = S(P(x))$ *is provable.*

Proof. By recursion on A: $\chi_{s=t} := \varepsilon(s,t)$, $\chi_\bot := 0$, and on Boolean values Z is negation and G is conjunction by Theorems 8.28 and 8.29; disjunction is $Z(G(Z(b), Z(c)))$ and implication is $Z(G(b, Z(c)))$. The truth tables are checked on the two values. The clause for implication is valid intuitionistically because A is decidable by its characteristic term. Given a base and a step for $A(n)$, pass to the literal equation $\chi_{A(n)} = 1$, apply the induction rule and pass back. Apply the result to $x = 0 \vee x = S(P(x))$: the base is immediate, and the step holds by $P(Sx) = x$ without the induction hypothesis. □

For a Boolean b let $\mathrm{inc}(a,b) := C(Z(b), a)$, so that $\mathrm{inc}(a,0) = a$ and $\mathrm{inc}(a,1) = Sa$. Reflection and Theorem 8.28 give

$$\varepsilon(x,x) = 1, \qquad \varepsilon(x,y) = \varepsilon(y,x), \qquad \varepsilon(Sx,Sy) = \varepsilon(x,y), \qquad \varepsilon(0,Sx) = 0.$$

Define the *hit term* and the order relations

$$\mathrm{hit}(n,y) := B_0(n; r, v.\, \mathrm{inc}(v, \varepsilon(r,y))), \qquad x < y :\Leftrightarrow \mathrm{hit}(y,x) = 1, \qquad x \le y :\Leftrightarrow \mathrm{hit}(Sy,x) = 1.$$

Thus $\mathrm{hit}(0,y) = 0$ and $\mathrm{hit}(Sn,y) = \mathrm{inc}(\mathrm{hit}(n,y), \varepsilon(n,y))$.

Lemma 8.36 (Hit identities)**.** $\mathsf{T}_{B\varepsilon}$ *proves*

$$\begin{gathered}\mathrm{hit}(Sn,0) = 1, \quad \mathrm{hit}(Sn,Sy) = \mathrm{hit}(n,y), \quad \mathrm{hit}(n,n) = 0, \quad \mathrm{hit}(Sn,n) = 1,\\ \mathrm{hit}(n,y) = 0 \vee \mathrm{hit}(n,y) = 1, \qquad \mathrm{hit}(Sy,x) = Z(\mathrm{hit}(x,y)),\\ \mathrm{hit}(n,y) = 1 \to \mathrm{hit}(Sn,y) = 1, \qquad \mathrm{hit}(Sn,y) = 1 \to \mathrm{hit}(n,y) = 1 \vee n = y.\end{gathered}$$

Proof. $\mathrm{hit}(Sn,0) = 1$ by induction on n: the first step tests $\varepsilon(0,0) = 1$, and every later test $\varepsilon(Sn,0)$ is 0. $\mathrm{hit}(Sn,Sy) = \mathrm{hit}(n,y)$ by induction on n with y fixed: the base is $\mathrm{inc}(0, \varepsilon(0,Sy)) = 0$, and the step uses $\varepsilon(Sn,Sy) = \varepsilon(n,y)$. Then $\mathrm{hit}(n,n) = 0$ by induction, and $\mathrm{hit}(Sn,n) = \mathrm{inc}(0,1) = 1$.

For the range use quantifier-free induction on n: if $\varepsilon(n,y) = 0$ the state is unchanged; if $\varepsilon(n,y) = 1$ then $n = y$, the old state is $\mathrm{hit}(n,n) = 0$ and the new state is 1. If $\mathrm{hit}(n,y) = 1$, then $n \neq y$ by the diagonal value, so the next test is 0 and the hit persists. The same case split gives the last implication.

The duality $\mathrm{hit}(Sy,x) = Z(\mathrm{hit}(x,y))$ is proved by induction on x with y fixed. At $x = 0$ both sides are 1. In the step the left side is $\mathrm{hit}(Sy,Sx) = \mathrm{hit}(y,x)$ and the right side is $Z(\mathrm{inc}(\mathrm{hit}(x,y), \varepsilon(x,y)))$. If $\varepsilon(x,y) = 0$, the right side is $Z(\mathrm{hit}(x,y))$, and the induction hypothesis reads $\mathrm{inc}(\mathrm{hit}(y,x),0) = \mathrm{hit}(y,x) = Z(\mathrm{hit}(x,y))$. If $\varepsilon(x,y) = 1$, then $x = y$, the left side is 0 and the right side is $Z(\mathrm{hit}(Sx,x)) = Z(1) = 0$. □

Lemma 8.37 (Order)**.** *In* $\mathsf{T}_{B\varepsilon}$ *the relations* $<$ *and* $\le$ *form a discrete linear order with least element* 0*:* $x \le y \leftrightarrow (x = y \vee x < y)$*,* $x < Sy \leftrightarrow x \le y$*, exactly one of* $x < y$*,* $x = y$*,* $y < x$ *holds, both relations are transitive, the successor preserves and reflects them, and* $x \le Sy$ *with* $x \neq Sy$ *implies* $x \le y$*.*

Proof. The diagonal identities give $x \le x$, $x \not< x$ and $x < Sx$. The recurrence for $\mathrm{hit}(Sy,x)$, split on $\varepsilon(y,x)$, gives the first equivalence. If $x \neq y$, then $\mathrm{hit}(Sy,x) = \mathrm{hit}(y,x)$, and by duality $\mathrm{hit}(x,y)$ is its Boolean complement; this is trichotomy. For transitivity of $<$ keep $\mathrm{hit}(y,x) = 1$ and induct on z in the quantifier-free formula $\mathrm{hit}(z,y) = 1 \to \mathrm{hit}(z,x) = 1$. The base has a false antecedent. At Sz the last implication of Theorem 8.36 gives $\mathrm{hit}(z,y) = 1$, and then the induction hypothesis and persistence apply, or $z = y$, and then the retained $\mathrm{hit}(y,x) = 1$

and persistence apply. Transitivity of $\leq$ and the mixed laws follow by splitting equality from strict comparison, and antisymmetry from trichotomy. The identity $\mathrm{hit}(Sn, Sy) = \mathrm{hit}(n, y)$ gives preservation and reflection by the successor. $x < Sy \leftrightarrow x \leq y$ holds by definition, and the last claim follows from it and the first equivalence. Finally $\mathrm{hit}(Sx, 0) = 1$ is $0 \leq x$, and $x \leq 0$ gives $x = 0$ by trichotomy. □

Define the difference, the minimum, doubling and bounded addition by

$$\widehat{d}(x, y) := B_0(x; u, v.\, \mathrm{inc}(v, \mathrm{hit}(Su, y))), \qquad \mathrm{mn}(x, y) := \widehat{d}(x, \widehat{d}(x, y)), \qquad \mathrm{db}(n) := B_0(n; u, v.SSv),$$
$$K_b(a) := \widehat{d}(\mathrm{db}(b), \widehat{d}(b, a)), \qquad \mathrm{ad}_b(a, c) := \widehat{d}(K_b(a), \widehat{d}(b, c)).$$

Thus $\widehat{d}(0, y) = 0$ and $\widehat{d}(Sx, y) = \mathrm{inc}(\widehat{d}(x, y), \mathrm{hit}(Sx, y))$.

Lemma 8.38 (Difference)**.** $\mathsf{T}_{B\varepsilon}$ *proves*

$$\widehat{d}(x, 0) = x, \qquad \widehat{d}(Sx, Sy) = \widehat{d}(x, y), \qquad \widehat{d}(x, x) = 0, \qquad \widehat{d}(x, y) = 0 \leftrightarrow x \leq y,$$
$$\widehat{d}(x, Sy) = P(\widehat{d}(x, y)), \qquad \widehat{d}(x, y) \leq x,$$
$$y \leq x \to \widehat{d}(Sx, y) = S(\widehat{d}(x, y)), \qquad 0 < y \leq x \to \widehat{d}(x, P(y)) = S(\widehat{d}(x, y)),$$

and that $\widehat{d}$ *is monotone in its first argument and antitone in its second.*

Proof. $\widehat{d}(x, 0) = x$ by induction, since $\mathrm{hit}(Sx, 0) = 1$. For $\widehat{d}(Sx, Sy) = \widehat{d}(x, y)$ induct on x with y fixed: the base uses $\mathrm{hit}(1, Sy) = 0$, and the step uses $\mathrm{hit}(SSx, Sy) = \mathrm{hit}(Sx, y)$.

Quantifier-free induction on x with y fixed proves $\widehat{d}(x, y) = 0 \leftrightarrow \mathrm{hit}(x, y) = 0$. In the step, if $\mathrm{hit}(Sx, y) = 1$ the new difference is a successor; if $\mathrm{hit}(Sx, y) = 0$, then $\mathrm{hit}(x, y) = 0$ by persistence, the old difference is 0 and stays 0. By duality $\mathrm{hit}(x, y) = 0$ is $\mathrm{hit}(Sy, x) = 1$, which gives the characterization of $\leq$ and $\widehat{d}(x, x) = 0$.

For $\widehat{d}(x, Sy) = P(\widehat{d}(x, y))$ induct on x with y fixed. In the step the left side is $\widehat{d}(x, y)$ and the right side is $P(\mathrm{inc}(\widehat{d}(x, y), \mathrm{hit}(Sx, y)))$. If the Boolean is 1 the predecessor cancels the successor; if it is 0 the characterization gives $\widehat{d}(Sx, y) = 0$, hence $\widehat{d}(x, y) = 0$ and both sides are 0.

Quantifier-free induction on x proves $\widehat{d}(x, y) \leq x$: the new state is the old one or its successor. For monotonicity in the first argument induct on v in $u \leq v \to \widehat{d}(u, z) \leq \widehat{d}(v, z)$ with u, z fixed: at Sv either $u = Sv$, or $u \leq v$ by discreteness and the new state is at least the old one; the base uses $u \leq 0 \to u = 0$. For antitonicity keep $y \leq z$ and induct on x in $\widehat{d}(x, z) \leq \widehat{d}(x, y)$: if $\mathrm{hit}(Sx, z) = 0$ the left state stays and the right state does not decrease; if it is 1 then $z \leq x$, so $y \leq x$ and both states increase.

The first of the two final implications is the defining recurrence, since $y \leq x$ means $\mathrm{hit}(Sx, y) = 1$. For the second, $y = S(P(y))$ by coverage, $P(y) < x$, so $\widehat{d}(x, P(y)) \neq 0$, and coverage at this difference with $\widehat{d}(x, S(P(y))) = P(\widehat{d}(x, P(y)))$ gives the claim. □

Lemma 8.39 (Minimum, doubling, bounded addition)**.** $\mathsf{T}_{B\varepsilon}$ *proves:*

(a) $x \leq y \to \mathrm{mn}(x, y) = x$ *and* $y \leq x \to \mathrm{mn}(x, y) = y$*;*
(b) db *is monotone,* $n \leq \mathrm{db}(n)$ *and* $\widehat{d}(\mathrm{db}(n), n) = n$*;*
(c) $a \leq b \to \mathrm{ad}_b(a, 0) = a$*;*
(d) $a \leq b \wedge Sc \leq b \to \mathrm{ad}_b(a, Sc) = S(\mathrm{ad}_b(a, c))$*;*
(e) $c \leq b \to \mathrm{ad}_b(0, c) = c$*;*
(f) $\mathrm{ad}_b(b, b) = \mathrm{db}(b)$ *and* $\mathrm{ad}_b(a, c) \leq \mathrm{db}(b)$ *for all* a*,* c*;*
(g) ad_b *is monotone in its second argument;*
(h) if $a, c \leq b$ *and* $a, c \leq b'$ *then* $\mathrm{ad}_b(a, c) = \mathrm{ad}_{b'}(a, c)$*.*

Proof. (a) The first claim uses $\widehat{d}(x,y) = 0$ and $\widehat{d}(x,0) = x$. For the second induct on y in $y \le x \to \widehat{d}(x,\widehat{d}(x,y)) = y$ with x fixed. The base is $\widehat{d}(x,x) = 0$. If $Sy \le x$, then $y < x$ and $0 < \widehat{d}(x,y) \le x$, so by Theorem 8.38 $\widehat{d}(x,\widehat{d}(x,Sy)) = \widehat{d}(x,P(\widehat{d}(x,y))) = S(\widehat{d}(x,\widehat{d}(x,y))) = Sy$.

(b) $\mathrm{db}(0) = 0$ and $\mathrm{db}(Sn) = SS\,\mathrm{db}(n)$ give $n \le \mathrm{db}(n)$ by quantifier-free induction; monotonicity is an induction on the larger argument that splits equality at the successor from the discrete predecessor case. The identity is an induction on a literal equation: $\widehat{d}(\mathrm{db}(Sn), Sn) = \widehat{d}(S\,\mathrm{db}(n), n) = S(\widehat{d}(\mathrm{db}(n), n)) = Sn$, using $n \le \mathrm{db}(n)$.

(c)–(h) Since $\widehat{d}(b,a) \le b$, antitonicity and (b) give $b = \widehat{d}(\mathrm{db}(b), b) \le K_b(a) \le \mathrm{db}(b)$, and $K_b(0) = b$. If $Sa \le b$, put $q := \widehat{d}(b,a)$; then $0 < q \le b \le \mathrm{db}(b)$ and $\widehat{d}(b,Sa) = P(q)$, so $K_b(Sa) = \widehat{d}(\mathrm{db}(b), P(q)) = S(K_b(a))$. Quantifier-free induction on a proves (c) under $a \le b$: the base is $\widehat{d}(b,b) = 0$, and the step increments $\widehat{d}(K_b(a), b)$ by the last identity and $b \le K_b(a)$. For (d) put $q := \widehat{d}(b,c)$; then $0 < q \le b \le K_b(a)$ and $\mathrm{ad}_b(a, Sc) = \widehat{d}(K_b(a), P(q)) = S(\widehat{d}(K_b(a), q))$. Statement (e) is a quantifier-free induction on $c \le b$ from (c) and (d). For (f) substitute $a = c = b$ and use $\widehat{d}(b,b) = 0$ twice; the bound is $\widehat{d}(K_b(a), \widehat{d}(b,c)) \le K_b(a) \le \mathrm{db}(b)$. For (g), $c \le c'$ gives $\widehat{d}(b,c') \le \widehat{d}(b,c)$, and antitonicity of the outer difference gives the claim. For (h) keep $a \le b, b'$ and induct on c in $c \le \mathrm{mn}(b,b') \to \mathrm{ad}_b(a,c) = \mathrm{ad}_{b'}(a,c)$: the base gives a on both sides, and (d) applies to both sides whenever $Sc \le \mathrm{mn}(b,b')$. □

Every induction in Theorems 8.36 to 8.39 is on a quantifier-free formula with the other variables fixed. There is no total binary addition term; ad_b adds below the bound b.

Lemma 8.40 (Clipped initialized recursion)**.** *Let*

$$\mathrm{sel}_b(c,a,e) := \mathrm{ad}_b\big(G(Z(c),a), G(c,e)\big), \qquad \mathrm{Rec}_b(n,a;u,v.h) := B_0\Big(Sn;\, r,s.\ \mathrm{sel}_b\big(r, \mathrm{mn}(a,b), \mathrm{mn}(h[P(r)/u][s/$$

If $a, e \le b$, then $\mathrm{sel}_b(0,a,e) = a$ and $\mathrm{sel}_b(Sc,a,e) = e$. For all arguments,

$$\mathrm{Rec}_b(0,a;u,v.h) = \mathrm{mn}(a,b), \qquad \mathrm{Rec}_b(Sn,a;u,v.h) = \mathrm{mn}\big(h[n/u][\mathrm{Rec}_b(n,a;u,v.h)/v],\, b\big),$$

and $\mathrm{Rec}_b(n,a;u,v.h) \le b$.

Proof. By Theorem 8.29 one of the two summands of sel_b is 0 and the other is a at $c = 0$ and e at a successor; Theorem 8.39(c) and (e) give the values. The equations for Rec_b follow from those of B_0 and of sel_b, since both arguments $\mathrm{mn}(\cdot, b)$ are at most b; the bound is a quantifier-free induction. □

The cap b is a parameter that stays fixed during the recursion. Arbitrary initialization is thus available once a cap has been fixed, in agreement with Theorem 8.31: the value is bounded in the coordinate b.

8.2.4 A compiler with caps

Fix the primitive recursive descriptions built from zero, successor, projections, composition and primitive recursion. The following construction is a recursion on a description outside the theory; it produces terms and proofs of $\mathsf{T}_{B\varepsilon}$ and adds no evaluator for codes of descriptions.

Theorem 8.41 (Bounded realization)**.** *For every k-ary primitive recursive description f there are terms $\tau_f(b,x_1,\dots,x_k)$ and $M_f(b)$ such that $\mathsf{T}_{B\varepsilon}$ proves:*

(i) M_f is nondecreasing and $b \le M_f(b)$;

(ii) $\tau_f(b,\vec{x}) \le M_f(b)$ for all b, $\vec{x}$;

(iii) if $x_i \le b$ and $x_i \le c$ for all i, then $\tau_f(b,\vec{x}) = \tau_f(c,\vec{x})$.

Proof. By recursion on f.

Initial descriptions. $\tau_0 := 0$, $\tau_{\pi_i}(b, \vec{x}) := \mathrm{mn}(x_i, b)$, $\tau_S(b, x) := S\,\mathrm{mn}(x, b)$, with $M(b) := Sb$. The properties follow from Theorem 8.39(a).

Composition. Let $f(\vec{x}) = h(g_1(\vec{x}), \ldots, g_m(\vec{x}))$. Put $Q := M_{g_1} \circ \cdots \circ M_{g_m}$ (the identity if $m = 0$), $M_f := M_h \circ Q$ and

$$\tau_f(b, \vec{x}) := \tau_h(Q(b), \tau_{g_1}(b, \vec{x}), \ldots, \tau_{g_m}(b, \vec{x})).$$

Since the M_{g_i} are nondecreasing and $z \le M_{g_i}(z)$, $M_{g_i}(b) \le Q(b)$. (ii) is the bound for τ_h. For (iii) the inner values agree at the caps b and c by the hypothesis for the g_i, and they are bounded by $Q(b)$ and by $Q(c)$; (iii) for h gives the claim. (i) holds by composition.

Primitive recursion. Let $f(0, \vec{x}) = g(\vec{x})$ and $f(Sn, \vec{x}) = h(n, f(n, \vec{x}), \vec{x})$. Put

$$F := M_h \circ M_g \circ S, \qquad I_F(t) := B_0(t; u, v.F(v)), \qquad C_b := I_F(S\,\mathrm{db}(b)), \qquad M_f(b) := C_b.$$

F is nondecreasing with $Sz \le F(z)$ and $M_h(z) \le F(z)$. Then $I_F(0) = 0$, $I_F(St) = F(I_F(t))$, $t \le I_F(t)$ by induction, and I_F is nondecreasing by a quantifier-free induction that uses discreteness of the order; so $b \mapsto C_b$ is nondecreasing and $b \le C_b$. For $j \le b$ define the stage bound

$$\kappa_b(j) := I_F(S\,\mathrm{ad}_b(b, j)).$$

Theorem 8.39 gives

$$b \le \kappa_b(0), \quad M_g(b) \le \kappa_b(0), \qquad Sj \le b \to \kappa_b(Sj) = F(\kappa_b(j)), \qquad j \le b \to b \le \kappa_b(j) \le C_b. \tag{8.14}$$

Indeed $\kappa_b(0) = I_F(Sb) = F(I_F(b)) \ge F(b) \ge M_g(b)$; the recurrence is Theorem 8.39(d); the upper bound is Theorem 8.39(f) and the lower bound is monotonicity.

Write $n_b := \mathrm{mn}(n, b)$ and $\vec{x}_b := (\mathrm{mn}(x_i, b))_i$, and define

$$\tau_f(b, n, \vec{x}) := \mathrm{Rec}_{C_b}(n_b, \tau_g(b, \vec{x}_b);\, u, v.\, \tau_h(C_b, u, v, \vec{x}_b)).$$

(ii) is the bound of Theorem 8.40.

No clipping on bounded inputs. Fix $\vec{x} \le b$ and let $r_b(j) := \mathrm{Rec}_{C_b}(j, \tau_g(b, \vec{x}); u, v.\tau_h(C_b, u, v, \vec{x}))$. Quantifier-free induction on j proves

$$j \le b \to r_b(j) \le \kappa_b(j). \tag{8.15}$$

At 0, $\tau_g(b, \vec{x}) \le M_g(b) \le \kappa_b(0) \le C_b$, so the seed is not clipped. Let $Sj \le b$. The arguments j, $r_b(j)$, $\vec{x}$ of the step are bounded by $\kappa_b(j)$ and by C_b, so (iii) for h gives

$$\tau_h(C_b, j, r_b(j), \vec{x}) = \tau_h(\kappa_b(j), j, r_b(j), \vec{x}) \le M_h(\kappa_b(j)) \le F(\kappa_b(j)) = \kappa_b(Sj) \le C_b.$$

The value is not clipped, which proves (8.15) and

$$Sj \le b \to r_b(Sj) = \tau_h(C_b, j, r_b(j), \vec{x}). \tag{8.16}$$

Coherence. Fix b, c and $n, \vec{x}$ bounded by both. The minimum identities remove the clipping of the inputs. Quantifier-free induction on j proves $(j \le b \wedge j \le c) \to r_b(j) = r_c(j)$: the base is (iii) for g and the absence of seed clipping; if $Sj \le b, c$, the two states at j are equal and bounded by C_b and C_c, as are the other arguments, so (iii) for h equates the two step values and (8.16) removes their clipping. At $j = n$ this is (iii) for f. The parameters $b, c, n, \vec{x}$ are fixed and both induction formulas are quantifier-free.

Every body inserted into B_0 or Rec_b is the already compiled term τ_h of the original language. No recursion over a function introduced later by a graph is used. □

For a nonempty tuple $\vec{x} = (x_1, \ldots, x_k)$ let

$$C_i(\vec{x}) := \bigwedge_{j<i} x_j < x_i \wedge \bigwedge_{j>i} x_j \leq x_i$$

say that i is the first largest coordinate. By Theorem 8.37 these k quantifier-free cases are exhaustive and pairwise disjoint. Define the graph

$$\Phi_f(\vec{x}, z) := \bigvee_{i=1}^{k} (C_i(\vec{x}) \wedge z = \tau_f(x_i, \vec{x})), \tag{8.17}$$

and $z = \tau_f(0)$ for a nullary f.

Example 8.42. For addition, without the general compiler: $\Phi_+(x, y, z) := (y \leq x \wedge z = \mathrm{ad}_x(x, y)) \vee (x < y \wedge z = \mathrm{ad}_y(x, y))$. In each case the chosen cap bounds both arguments, and Theorem 8.39 gives the addition equations. Each of the two value terms is bounded by twice one input, whereas addition has no bounding coordinate.

Lemma 8.43 (Graphs)**.** $\mathsf{T}_{B\varepsilon}$ *proves that each* Φ_f *is total and single-valued. In the definitional extension by the functions* f^* *with graph* Φ_f*:*

(a) $f^*(\vec{x}) = \tau_f(b, \vec{x})$ *whenever* b *bounds all* x_i*;*
(b) the defining equations of the description f *hold;*
(c) induction holds for quantifier-free formulas that contain finitely many of the f^**.*

Proof. Exhaustiveness gives a value term in each case, and disjointness gives uniqueness. (a) On the case C_i both b and x_i bound all arguments, so Theorem 8.41(iii) applies.

(b) The initial equations follow from the initial terms. For composition take a cap b for the inputs; the inner values are bounded by $Q(b)$, and (a) at $Q(b)$ for the outer function gives the equation from the definition of τ_f. For recursion at the counter 0 take a cap b for $0, \vec{x}$: the seed is not clipped, so the value is $\tau_g(b, \vec{x}) = g^*(\vec{x})$. At Sn take a cap b for $Sn, \vec{x}$: then $n \leq b$, (8.16) applies, (a) identifies the previous state with $f^*(n, \vec{x})$, all arguments of the step are bounded by C_b, and (a) for h identifies the step value with $h^*(n, f^*(n, \vec{x}), \vec{x})$.

(c) Conservativity of the extension is not enough for this. By recursion on a term with occurrences of the f^* one obtains finitely many pairs (A_j, t_j) of a quantifier-free guard and a term, both in the original language, such that the guards cover all inputs and the term equals t_j on A_j: at a composition take the product of the cases of the arguments and then the cases of the outer graph. The guards need not be disjoint; the values agree on overlaps. For a quantifier-free formula take a common refinement for all its terms, replace the terms by their values on each guard, and keep the connectives. The disjunction of the guarded formulas is quantifier-free in the original language and provably equivalent to the given formula, also under implication and negation, because the guards cover all inputs. Theorem 8.35 applied to it gives the induction instance. A graph with an existential quantifier would not allow this step. □

Theorem 8.44 (Bi-interpretation)**.** $\mathsf{T}_{B\varepsilon}$ *and* iPRA *are bi-interpretable. Both interpretations use the whole domain and actual equality, the logic is translated identically, and the identity is the isomorphism of both composites. The theories are equiconsistent.*

Proof. In iPRA. Interpret 0, S as themselves, ε as the primitive recursive characteristic of equality, and each staged recursor with body h by primitive recursion with initial value $S(h(0, 0, \vec{p}))$ and step $h(n, v, \vec{p})$, after translating the body. The equations (8.9), (8.10) become defining equations, and (8.11)–(8.13) and induction for literal equations are provable.

In $\mathsf{T}_{B\varepsilon}$. Interpret each description f by its graph Φ_f, or work in the extension by the f^*. Theorem 8.43 gives the defining equations and every quantifier-free induction instance; Theorem 8.28 and (8.13) give injectivity of the successor and $0 \neq Sx$.

The composite on the side of iPRA. By recursion on a description, with quantifier-free induction on the counter at each recursion node, the double interpretation of a function equals the function: both satisfy the same initial and step equations.

The composite on the side of $\mathsf{T}_{B\varepsilon}$. By recursion on the stage of a term. 0 and S are preserved. The interpreted characteristic of equality and ε are both Boolean and both reflect equality, so they agree by cases. For a recursor with body h the hypothesis for the earlier stage identifies the interpreted body with h; the recursor and its double interpretation satisfy (8.9), (8.10) for the same body, and induction on the equation between them identifies them at every counter and parameter. Where the double interpretation is expressed with graph-defined functions, Theorem 8.43(c) provides the induction instance.

All operations and equality are preserved by the identity, so every formula is provably equivalent to its double translation by induction on the formula. Falsity is preserved in both directions. □

8.2.5 The functions denoted by terms

For $k \geq 1$ let $\mathcal{A}_k$ be the set of primitive recursive $f\colon \mathbb{N}^k \to \mathbb{N}$ for which there are a coordinate i and a unary primitive recursive F with $f(\vec{x}) \leq F(x_i)$ for all $\vec{x}$, and let $\mathcal{A}_0$ be the constants.

Theorem 8.45 (Exact class of term functions)**.** *The functions denoted in the standard model by the terms of* $\mathsf{T}_{B\varepsilon}$ *with k free variables are exactly the functions in $\mathcal{A}_k$.*

Proof. Every term denotes a primitive recursive function, and Theorem 8.31 gives the bound. Conversely let $f(\vec{x}) \leq F(x_i)$. The compiler provides the cases $C_j(\vec{x})$ and the value terms $v_j(\vec{x}) := \tau_f(x_j, \vec{x})$, with value $f(\vec{x})$ on C_j. Let $a_j(\vec{x}) := G(\chi_{C_j}(\vec{x}), v_j(\vec{x}))$: it is 0 off its case and $f(\vec{x})$ on it. Exactly one case holds, so each a_j is at most $F(x_i)$ and at most one is nonzero. Every unary primitive recursive function is a term: for one variable the only case is automatic and the value is $\tau_F(x, x)$. Put $b := \tau_F(x_i, x_i)$, $s_0 := 0$ and $s_j := \mathrm{ad}_b(s_{j-1}, a_j)$ for $1 \leq j \leq k$. Every partial sum is 0 or $f(\vec{x})$, so both arguments of each bounded addition are at most b, and s_k denotes f. For $k = 0$ use the numeral. □

The converse direction assumes that the majorant is valid in the standard model. To prove inside $\mathsf{T}_{B\varepsilon}$ that the single term equals the graph-defined function one needs a proof of the majorant in the theory; Theorem 8.44 needs no such hypothesis.

Proposition 8.46 (Number of value terms)**.** *The graph* (8.17) *of a k-ary function has k value terms. The k-ary sum is not, in the standard model, always equal to one of $r < k$ terms.*

Proof. Each of the r terms is bounded by a constant or by a function of one input. Choose an input that controls none of them, fix the others at 0 and let the chosen one grow: the terms stay bounded and the sum does not. □

This concerns a choice among finitely many value terms. It is not a bound on the number of clauses of a graph formula: the characteristic function of a primitive recursive graph is bounded, hence a term by Theorem 8.45, so every such graph has a definition by one equation. The value terms are what gives totality and the transfer of induction in Theorem 8.43.

Example 8.47 (The majorant must be primitive recursive)**.** $\mathcal{A}_2$ is strictly smaller than the set of primitive recursive functions that are bounded by an arbitrary function of one input. Let $u_0, u_1, \ldots$ enumerate the unary primitive recursive descriptions and let $g(x) := 1 + \max_{e \leq x} u_e(x)$. This g is total recursive and has no primitive recursive majorant, since $g(x) > u_e(x)$ for $x \geq e$. Fix a machine for g and let $s(x, t)$ be its output if it halts within t steps on x, and 0 otherwise; s is primitive recursive. Then $f(x, t) := x + s(x, t)$ is primitive recursive, and for fixed x its values lie in $\{x, x + g(x)\}$, so it is bounded by a function of x. A

primitive recursive bound in x would majorize g, and no bound in t exists because $f(x,t) \geq x$. Hence $f \notin \mathcal{A}_2$.

8.2.6 Truncated subtraction in place of the equality test

Definition 8.48. The terms are $t ::= x \mid 0 \mid St \mid d(t,t) \mid B(n;u,v.h)$. The computation equations are (8.9), (8.10) and

$$d(x,0) = x, \qquad d(x,Sy) = P(d(x,y)), \tag{8.18}$$

where P is the term of Theorem 8.27.

(a) T_{Bd} is ordinary intuitionistic first-order logic with equality over these terms, with the four computation equations, coverage $\forall x\,(x = 0 \vee \exists y\, x = Sy)$, separation $0 = S0 \to \bot$ and induction for literal equations with side assumptions.

(b) N_{Bd} is the native calculus over these terms with the four computation equations and the rule forms of Section 1.3; the contradiction is $0 = S0$.

In T_{Bd} coverage is an axiom, in contrast to $\mathsf{T}_{B\varepsilon}$. It is used in the proof of Theorem 8.50, and it does not follow from the computation equations and induction for literal equations: `monus_and_neutral_major` refers to the coordinatewise structure on $\mathbb{N}\times\mathbb{N}$, treated in `scope_kernel`, which satisfies those and refutes coverage. In N_{Bd} the case rule NC supplies this resource.

By Theorems 8.28 and 8.29 the terms P, Z, G and B_0 and their equations are available in both theories. In T_{Bd}, coverage and the values of Z give

$$Z(x) = 0 \vee Z(x) = 1, \qquad Z(x) = 1 \leftrightarrow x = 0. \tag{8.19}$$

Lemma 8.49. *With induction for literal equations only,*

$$d(0,y) = 0, \qquad P(d(Sx,y)) = d(x,y), \qquad d(Sx,Sy) = d(x,y), \qquad d(x,x) = 0, \qquad d(Sx,x) = 1.$$

Proof. $d(0,y) = 0$ by induction on y with $P(0) = 0$. For the second equation fix x and induct on y: the base is $P(Sx) = x$, and $P(d(Sx,Sy)) = P(P(d(Sx,y))) = P(d(x,y)) = d(x,Sy)$. The third equation follows from (8.18) and the second. The last two are inductions on x with bases $d(0,0) = 0$, $d(1,0) = 1$ and the third equation as step. □

Lemma 8.50 (Gated complement)**.** T_{Bd} *proves*

$$G\big(Z(d(y,x)),\, d(x,d(x,y))\big) = G\big(Z(d(y,x)),\, y\big), \tag{8.20}$$

and consequently $d(y,x) = 0 \to d(x,d(x,y)) = y$.

Proof. Fix x and apply induction for literal equations to (8.20) as an equation in y. At $y = 0$ the control is $Z(d(0,x)) = 1$, and both second arguments are 0, by $d(x,0) = x$ and $d(x,x) = 0$.

For the step keep the induction hypothesis and put $a := d(Sy,x)$. By coverage a is a successor or 0. If a is a successor, $Z(a) = 0$ and both sides of the goal are 0. Let $a = 0$. By Theorem 8.49, $d(y,x) = P(d(Sy,x)) = 0$, so the induction hypothesis, with both gates at the control 1, gives

$$d(x,t) = y, \qquad t := d(x,y).$$

If $t = 0$, then $d(x,t) = x$, so $x = y$ and $a = d(Sy,y) = 1$, contradicting $a = 0$. Hence $t = Sw$ for some w by coverage, and $w = P(t) = d(x,Sy)$ by (8.18). Put $v := d(x,w)$. Then $P(v) = d(x,Sw) = d(x,t) = y$. If $v = 0$, then $y = 0$, so $t = d(x,0) = x = Sw$ and

$v = d(Sw, w) = 1$, a contradiction. Hence $v = Sz$ by coverage, $z = P(v) = y$ and $v = Sy$. With $w = d(x, Sy)$ this reads

$$d(x, d(x, Sy)) = Sy,$$

which is the goal at the control $Z(a) = 1$. The contradictions use $0 \neq 1$ only. The only induction formula is the literal equation (8.20). If $d(y, x) = 0$, both gates have the control 1. □

Theorem 8.51 (Reflection of equality)**.** *Let* $\varepsilon(x, y) := G(Z(d(x, y)), Z(d(y, x)))$. T_{Bd} *proves*

$$d(x, y) = 0 \wedge d(y, x) = 0 \to x = y, \qquad \varepsilon(x, y) = 0 \vee \varepsilon(x, y) = 1, \qquad \varepsilon(x, y) = 1 \leftrightarrow x = y.$$

Proof. From $d(y, x) = 0$, Theorem 8.50 gives $d(x, d(x, y)) = y$; substituting $d(x, y) = 0$ and $d(x, 0) = x$ gives $x = y$. Each input of the gate is 0 or 1 by (8.19), and the gate equations give the range and $\varepsilon(x, y) = 1 \leftrightarrow d(x, y) = 0 \wedge d(y, x) = 0$: if the first input is 0 the output is 0, and otherwise it is the second input. The first statement gives $x = y$ from the right side; conversely $x = y$ makes both differences 0 by $d(x, x) = 0$. □

Corollary 8.52. T_{Bd} *is bi-interpretable with* iPRA*, with whole domains, actual equality, the logic translated identically and the identity as isomorphism of both composites.*

Proof. By Theorem 8.51 the term ε satisfies (8.11) and (8.12), so every derivation of $\mathsf{T}_{B\varepsilon}$ becomes a derivation of T_{Bd} when this term is substituted for the symbol; in particular Theorems 8.41 and 8.43 hold. In the other direction iPRA interprets the recursors as in Theorem 8.44 and d as truncated subtraction; it proves (8.18), coverage, separation and induction for literal equations.

For the composite on the side of T_{Bd}, first the recursors C and W, which do not contain d, are identified with their double interpretations by induction on the equation between them, and hence P is preserved. The double interpretation of d is the graph of the primitive recursive truncated subtraction; it satisfies the two equations (8.18) with the preserved P, and induction on the second argument identifies it with d. Likewise the term $\widehat{d}$ of Theorem 8.38, with ε expanded, satisfies $\widehat{d}(x, 0) = x$ and $\widehat{d}(x, Sy) = P(\widehat{d}(x, y))$, so $\widehat{d}(x, y) = d(x, y)$ by induction on y for this literal equation of the original language. The remaining recursors are identified as in Theorem 8.44, and the composite on the side of iPRA is unchanged. □

8.2.7 The native calculus with truncated subtraction

The proof of Theorem 8.50 splits on $t = 0$ or $t = Sw$ while an induction hypothesis that mentions t is in the context, and Theorem 8.32 excludes a selector term. N_{Bd} derives the split by a selector whose zero branch is Boolean.

Lemma 8.53. N_{Bd} *derives the equality laws,* NC *and* EI *of Theorem 8.2, the equations of Theorems 8.28, 8.29 and 8.49, every equation from* $0 = S0$*, and the functional range rule: if* $\Gamma \vdash K(0)$ *and* $\Gamma \vdash K(1)$ *then* $\Gamma \vdash K(Z(t))$*, for every formula template* K *and term* t*.*

Proof. Reflexivity follows from the computation $C(S0, p) = p$ as in Theorem 8.2, and the other equality laws, NC and EI are derived as there; Theorem 8.4 holds as well. The equations of the three lemmas use computation, equality and EI only. Under $0 = S0$, transport between $C(0, a) = Sa$ and $C(1, a) = a$ gives $Sa = a$; then EI on the equation $a = 0$ gives $a = 0$ for every a, hence every equation. The functional range rule is NC on the template $w \mapsto K(Z(w))$. □

By recursion on a native formula, with DI by cases at each $D_{x,y}$ and the representation of $Q_y B$ by $D_{v,y} B$ with v unused, every native formula follows from $0 = S0$.

A term b is called *two-valued* in Γ if the functional range rule holds for it: $\Gamma \vdash K(0)$ and $\Gamma \vdash K(1)$ imply $\Gamma \vdash K(b)$ for every template K. Terms $Z(t)$ are two-valued.

Lemma 8.54 (A selector with a Boolean zero branch)**.** *Let* $\mathrm{inc}(a,b) := C(Z(b),a)$ *and*

$$\eta(y,b,u) := \mathrm{inc}\big(G(y,u),\, G(Z(y),b)\big).$$

Then $\eta(Sy,b,u) = u$ *for all terms, and* $\eta(0,b,u) = b$ *for every two-valued* b*.*

Proof. $\eta(Sy,b,u) = \mathrm{inc}(u,G(0,b)) = \mathrm{inc}(u,0) = C(1,u) = u$. $\eta(0,b,u) = \mathrm{inc}(0,G(1,b)) = C(Z(b),0)$, which is $C(1,0) = 0$ at $b = 0$ and $C(0,0) = 1$ at $b = 1$; the functional range rule for the template $C(Z(w),0) = w$ gives $\eta(0,b,u) = b$. □

The zero branch of η is two-valued and the successor branch is arbitrary, so η does not contradict Theorem 8.31.

Lemma 8.55 (Contextual cases)**.** *In* N_{Bd}*, let* $s = t$ *be a literal equation.*
(a) If b *is two-valued in* Γ*,* $\Gamma, b = 0 \vdash s = t$ *and* $\Gamma, b = 1 \vdash s = t$*, then* $\Gamma \vdash s = t$*.*
(b) If z *is fresh,* $\Gamma, r = 0 \vdash s = t$ *and* $\Gamma, r = Sz \vdash s = t$*, then* $\Gamma \vdash s = t$*.*
The context and the equation may mention b *and* r*.*

Proof. (a) Let $K(x) := D_{a,y}(\eta(y,x,s) = \eta(y,b,t))$ with a unused. For $c \in \{0,1\}$ derive $K(c)$ by DI by cases: the local assumption $\eta(0,c,s) = \eta(0,b,t)$ is $c = b$ by Theorem 8.54, the given branch gives $s = t$, and this is the matrix at Sz. The range rule for b gives $K(b)$. Rule DE opens its closure, whose zero instance $\eta(0,b,s) = \eta(0,b,t)$ holds because both sides equal b, and QE at 1 gives $s = t$.

(b) Rule NC on fresh variables in literal equations gives

$$G(Z(r),r) = 0, \qquad G(Z(Z(r)),r) = G\big(Z(Z(r)), S(P(r))\big) :$$

for the first, the zero case is $G(1,0) = 0$ and a successor case has the control 0; for the second, the zero case has the control 0, and a successor case has the control 1 and $P(Sx) = x$. Apply (a) to $b := Z(r)$. Under $Z(r) = 1$ the first identity gives $r = 0$. Under $Z(r) = 0$ the second gives $r = S(P(r))$; substitute $P(r)$ for z in the second given branch. □

Theorem 8.56. N_{Bd} *derives* (8.20)*, the three statements of Theorem 8.51 in the forms*

$$d(x,y) = 0,\ d(y,x) = 0 \vdash x = y, \qquad s = t \vdash \varepsilon(s,t) = 1, \qquad \varepsilon(s,t) = 1 \vdash s = t,$$

with $\varepsilon(s,t)$ *two-valued, and the introduction and elimination rules of intuitionistic implication, conjunction, disjunction, both quantifiers and falsity for the formulas defined in the proof. It is bi-interpretable with* T_{Bd} *and with* iPRA*, with whole domains, actual equality, identity composites and both formula round trips.*

Proof. Every case split in the proof of Theorem 8.50 has an equation as its conclusion, so it is an instance of Theorem 8.55(b); the branches that end in $0 = 1$ are closed by Theorem 8.53. The induction is EI for the same literal equation. The term $\varepsilon(s,t)$ is two-valued by the range rules of its two inputs and the gate equations.

For a literal equation E, namely $s = t$, define the gate $\mathcal{G}_v(E) := \big(G(v,\varepsilon(s,t)) = G(v,1)\big)$; it is derivable at $v = 0$ and interderivable with E at $v = Sw$. For literal equations E $(s = t)$ and F $(r = w)$ define the formula selector $\mathcal{M}_v(E,F) := \big(\eta(v,\varepsilon(s,t),\varepsilon(r,w)) = 1\big)$; by Theorem 8.54, $\mathcal{M}_0(E,F)$ is interderivable with E and $\mathcal{M}_{Sz}(E,F)$ with F. Extend the gate through $D_{x,y}$ by $\mathcal{G}_v(D_{x,y}A) := D_{x,y}\mathcal{G}_v A$, replace each $Q_y A$ by $D_{a,y}A$ with a unused, align binder depths by the padding $A \approx D_{a,y}\mathcal{G}_y A$, and extend the selector through matched binders by recursion on the formulas; the endpoint properties are preserved by the binder

congruence of Theorem 8.4. With $p(z) := B_0(z; u, v.Z(v))$, so that $p(1) = 1$ and $p(2) = 0$, and with unused coordinates fresh, put

$$\begin{aligned} A \Rightarrow_* B &:= D_{a,z}\mathcal{M}_z(A, B), & A \wedge_* B &:= D_{a,z}\mathcal{G}_z\mathcal{M}_{p(z)}(A, B), \\ \exists_* x\, A(x) &:= D_{x,y}\mathcal{G}_y A(x), & A \vee_* B &:= \exists_* x\, \mathcal{M}_x(A, B), \\ \forall_* x\, A(x) &:= A(0) \wedge_* D_{a,z}\mathcal{G}_z A(z), & \bot_* &:= (0 = S0). \end{aligned}$$

Implication introduction is DI by cases, whose local assumption is the zero instance of the selector; implication elimination opens the closure and specializes at 1. The derivable zero instance of the gate gives the introductions of conjunction and existence, and specialization at 1 and 2 their eliminations. Once implication is available, NC on an implication template gives numerical cases in a context for arbitrary formulas, hence disjunction elimination and universal elimination. These are the constructions of Theorem 1.78 with the atomic gate and selector above in place of the numerical selector sel, which does not exist here.

Every rule of N_{Bd} is sound for T_{Bd} under the readings of Section 1.2: DI by cases uses coverage and QI uses induction for literal equations. Conversely N_{Bd} derives the translation of every theorem of T_{Bd} by the rules just established. The round trips are

$$Q_y B \approx B(0) \Rightarrow_* \forall_* y\, B(y), \qquad D_{x,y}B \approx \exists_* x\, \big(B(x, 0) \Rightarrow_* \forall_* y\, B(x, y)\big) :$$

forward, DE opens the witness, and under the zero instance QE gives every instance; backward, eliminate the existential, keep the zero instance of the matrix, and use the implication and the universal at Sz for DI by cases. Induction on formulas gives both formula composites, and the numerical composites are identities. The bi-interpretation with iPRA is Theorem 8.52. □

8.3 The self-seeded recursor

This section draws on `selfseed_scalar`, `scope_and_coverage` and `coverage_and_kernel`.

8.3.1 The theories

Definition 8.57. The terms are $t ::= x \mid 0 \mid St \mid \varepsilon(t, t) \mid F(n; v.h)$, where $F(n; v.h)$ binds v in h. The computation equations are

$$F(0; v.h) = S(h[0/v]), \qquad F(Sn; v.h) = h[F(n; v.h)/v]. \tag{8.21}$$

T_F^- is ordinary intuitionistic first-order logic with equality over these terms with (8.21), the axioms (8.11), (8.12), (8.13) for ε and separation, and induction for literal equations with side assumptions. T_F is T_F^- with the additional axiom $Sx = Sy \to x = y$. Coverage is not an axiom of either theory.

The update sees neither an initializer nor the old counter. Compared with $\mathsf{T}_{B\varepsilon}$, which derives injectivity from the counter argument (Theorem 8.28), T_F postulates it. As in Section 8.2, S can be removed from the signature by $Sa := F(0; v.a)$. Put

$$C(n, a) := F(n; v.a), \qquad Z(n) := C(n, 0),$$

so that $C(0, a) = Sa$, $C(Sn, a) = a$, $Z(0) = 1$, $Z(Sn) = 0$.

Lemma 8.58. *T_F^- proves $Sn \neq 0$. The terms*

$$\neg_B b := \varepsilon(b, 0), \qquad b \wedge_B c := \varepsilon\big(C(b, \varepsilon(c, 0)), 0\big), \qquad b \vee_B c := \neg_B(\neg_B b \wedge_B \neg_B c)$$

have the truth tables of negation, conjunction and disjunction on the values $0, 1$. Every quantifier-free formula A has a term χ_A with $\chi_A = 0 \vee \chi_A = 1$ and $\chi_A = 1 \leftrightarrow A$, and T_F^- admits induction with side assumptions for quantifier-free formulas.

Proof. If $Sn = 0$, then $0 = Z(Sn) = Z(0) = 1$. Reflection and the range axiom give $\varepsilon(0,0) = 1$ and $\varepsilon(1,0) = 0$. For the conjunction, $C(0, \varepsilon(c,0)) = S(\varepsilon(c,0)) \neq 0$, so the value at $b = 0$ is 0; and $C(1, \varepsilon(c,0)) = \varepsilon(c,0)$, so the value at $b = 1$ is $\neg_B \neg_B c = c$. The only non-Boolean intermediate value is a successor, and its distinctness from 0 does not cancel a successor. Define $\chi_{s=t} := \varepsilon(s,t)$, $\chi_\perp := 0$, and the connectives by the Boolean terms, implication as $\neg_B \chi_A \vee_B \chi_B$; the Boolean range comes from the range axiom and the truth tables. Induction for $A(n)$ is induction for the literal equation $\chi_{A(n)} = 1$. □

Definition 8.59. A *finite-case graph* is a formula $\bigvee_i (A_i(\vec{x}) \wedge y = t_i(\vec{x}))$ with quantifier-free guards A_i and terms t_i of the language, such that the guards are provably exhaustive and the values provably agree where guards overlap.

Lemma 8.60 (Finite-case elimination)**.** *A finite-case graph is provably total and single-valued. In the definitional extension by the functions of finitely many finite-case graphs, every quantifier-free formula is provably equivalent to a quantifier-free formula of the original language, and induction for quantifier-free formulas of the extension is admissible. Compositions of finite-case functions are finite-case.*

Proof. As in Theorem 8.43(c): expand each term into finitely many guarded terms of the original language by refining the cases of its arguments and then the cases of the outer graph, take a common refinement for a formula, and apply Theorem 8.58 to the resulting disjunction. □

The lemma concerns function terms in formulas. It does not allow a new body h of F that contains a function defined by a graph; bodies are always terms of the original language.

8.3.2 Without injectivity of the successor

All statements of this subsection are theorems of T^-_F.

Lemma 8.61 (No fixed point)**.** *$Sa \neq a$ for every a.*

Proof. The term $h(v) := \varepsilon(v,0)$ has no fixed point: $h(v) = v$ and the range of h give $v = 0$ or $v = 1$, contradicting $h(0) = 1$ and $h(1) = 0$. If $Sa = a$, then $F(a; v.h) = F(Sa; v.h) = h(F(a; v.h))$. □

Theorem 8.62 (No cycle of a fixed length)**.** *For each fixed integer $k \geq 1$, the equations* (8.21) *and the axioms* (8.11)*,* (8.12)*,* (8.13) *prove $\forall a\ S^k a \neq a$, without any instance of induction. Consequently distinct numerals are provably distinct, and $S^i a \neq S^j a$ for fixed $i < j$.*

Proof. The proofs are constructed by recursion on k outside the theory. Assume the statement for all lengths below k. Then $\bar{0}, \ldots, \bar{k}$ are pairwise distinct: $\bar{0}$ differs from a positive numeral by Theorem 8.58, and $\bar{i} = \bar{j}$ with $1 \leq i < j \leq k$ would be $S^{j-i}(\bar{i}) = \bar{i}$ with $1 \leq j - i < k$. For $k = 1$ only separation is used.

For a term b with $b = 0 \vee b = 1$ let $\mathrm{Inc}(b,a) := C(\varepsilon(b,0), a)$, so that $\mathrm{Inc}(0,a) = a$ and $\mathrm{Inc}(1,a) = Sa$. Define, as a written finite expansion,

$$T_0(v) := 0, \qquad T_{j+1}(v) := \mathrm{Inc}(\varepsilon(v, \bar{j}), \cdot)^{j+1}(T_j(v)) \quad (0 \leq j < k), \qquad c_k(v) := T_k(v),$$

where the exponent denotes $j+1$ written compositions. At $v = \bar{i}$ with $i < k$ the test with index i is 1 and the others are 0, so $i+1$ increments are performed; at $v = \bar{k}$ all tests are 0. Hence

$$c_k(\bar{i}) = \overline{i+1}\ (0 \leq i < k), \qquad c_k(\bar{k}) = 0, \qquad c_k(v) = \bar{0} \vee \cdots \vee c_k(v) = \bar{k},$$

the last by the finitely many cases of (8.11): if some test is 1, reflection identifies v with a numeral and distinctness makes the other tests 0.

If $c_k^k(w) = w$ for the k-fold composition, then w is one of $\bar{0}, \ldots, \bar{k}$ by the range, since $k \geq 1$. On these $k+1$ distinct values c_k is one cycle, and k applications of it end at a different numeral. So $c_k^k(w) \neq w$ for every w.

Now let $S^k a = a$ and $w := F(a; v.c_k)$. By k uses of the successor equation and congruence, $w = F(S^k a; v.c_k) = c_k^k(F(a; v.c_k)) = c_k^k(w)$, a contradiction. The last statement applies the theorem to $S^i a$ with the length $j - i$. □

The theorem excludes a cycle along one successor chain. It does not derive $a = b$ from $Sa = Sb$: two predecessor chains may merge without a cycle. The term c_k depends on the fixed k, and there is no term for cycles of a length given by a variable.

For the next results define, with a parameter a,

$$g_a(v) := \varepsilon(v, 0) \vee_B \varepsilon(Sv, a), \qquad \theta_a(v) := C(g_a(v), v), \qquad U(j, a) := F(j; v.\theta_a(v)), \qquad p_*(a) := U(a, a).$$

The update θ_a holds the state if it is 0 or if its successor is a, and increments it otherwise. Since $\theta_a(0) = 0$, the orbit starts at $U(0, a) = 1$, and $U(Sj, a) = \theta_a(U(j, a))$.

Lemma 8.63 (Protected orbit)**.** *$U(j, a) = Sj \vee S(U(j, a)) = a$, and $S(p_*(S^2 m)) = S^2 m$.*

Proof. The first statement is a quantifier-free induction on j with a fixed. The base is $U(0, a) = S0$. In the step put $v := U(j, a)$ and decide $Sv = a$. If it holds, θ_a fixes v and the second alternative persists. Otherwise the first alternative of the hypothesis gives $v = Sj \neq 0$, the guard is false and $U(Sj, a) = Sv = SSj$.

Specialize the proved statement to $j = m$, $a = S^2 m$. Both alternatives give $S(U(m, S^2 m)) = S^2 m$, so the state is fixed by θ_a and two further computation steps give $U(S^2 m, S^2 m) = U(m, S^2 m)$. No successor is cancelled. □

Theorem 8.64 (Coverage and a right predecessor)**.** *Let P be the function of the finite-case graph*

$$(n = 0 \wedge p = 0) \vee (n = 1 \wedge p = 0) \vee (n \neq 0 \wedge n \neq 1 \wedge p = p_*(n)).$$

T_F^- proves, by Theorem 8.60,

$$P(0) = P(1) = 0, \qquad n = 0 \vee n = S(P(n)), \qquad S(P(Sn)) = Sn,$$

and hence coverage $n = 0 \vee \exists m\, n = Sm$.

Proof. The graph is finite-case because equality is decidable and $0 \neq 1$. Apply quantifier-free induction to

$$n = 0 \ \vee \ n = 1 \ \vee \ S(p_*(n)) = n.$$

The base holds. In the step, $n = 0$ gives $Sn = 1$. If $n = 1$, then $Sn = S^2 0$ and Theorem 8.63 with $m = 0$ gives $S(p_*(Sn)) = Sn$. If $S(p_*(n)) = n$, then $Sn = S^2(p_*(n))$ and Theorem 8.63 with $m = p_*(n)$ gives $S(p_*(Sn)) = Sn$. This proves the displayed disjunction for every n. To pass to P, decide $n = 0$ and $n = 1$: if $n = 0$ the first alternative of $n = 0 \vee n = S(P(n))$ holds; if $n = 1$ then $n = S0 = S(P(1))$; and if $n \neq 0, 1$ the third alternative of the disjunction holds and $P(n) = p_*(n)$, so $n = S(P(n))$. Applying $n = 0 \vee n = S(P(n))$ to Sn and excluding its zero alternative gives $S(P(Sn)) = Sn$. □

The proof holds for all elements of a model, standard or not. The result is a right inverse: it chooses one predecessor of each nonzero element and does not prove that $P(Sa) = a$.

Proposition 8.65 (Standard numerals have unique predecessors)**.** *For every fixed integer $k \geq 1$, T_F^- proves*

$$Sx = \bar{k} \leftrightarrow x = \overline{k-1}. \tag{8.22}$$

Proof. By recursion on k outside the theory. The data carried are the pairwise distinctness of $\bar{0}, \ldots, \overline{k+1}$ and (8.22) for all smaller positive indices. Initially $0, 1, 2$ are distinct by Theorems 8.58 and 8.61.

Let h_k be a term with the values

$$h_k(v) = \begin{cases} \overline{i-1}, & v = \bar{i},\ 1 \leq i \leq k-1, \\ \overline{k-1}, & v = \overline{k+1}, \\ \bar{k}, & \text{otherwise.} \end{cases}$$

Such a term is obtained as follows. For $r = 1, \ldots, k$ let $b_r(v)$ be the Boolean negation of the disjunction of $\varepsilon(v, \bar{i})$ for $1 \leq i \leq \min(r, k-1)$, together with $\varepsilon(v, \overline{k+1})$ if $r = k$; it is the characteristic term of the condition that the displayed value is at least r. Put $t_0(v) := 0$, $t_r(v) := C(\neg_B b_r(v), t_{r-1}(v))$ and $h_k := t_k$: a true condition contributes one successor. For $k = 1$ the values of h_1 are those of $\neg_B \varepsilon(v, 2)$.

Let $A_k(n) := F(n; v.h_k)$ and let $q_0 := \overline{k+1}$, $q_j := \overline{k-j}$ for $1 \leq j \leq k$; these numerals are distinct. Since $h_k(0) = \bar{k}$, $A_k(0) = q_0$, and computation gives $A_k(\bar{j}) = q_j$ for $0 \leq j \leq k$. The term h_k never has the value q_0, and for $1 \leq j \leq k$,

$$h_k(v) = q_j \leftrightarrow v = q_{j-1}.$$

Quantifier-free induction gives $A_k(n) = q_0 \leftrightarrow n = 0$: at a successor both sides are false. For $1 \leq j < k$, recursion on j gives $A_k(n) = q_j \leftrightarrow n = \bar{j}$: by coverage (Theorem 8.64) $n = 0$ or $n = Sm$; at 0 both sides are false; at Sm the left side is equivalent to $A_k(m) = q_{j-1}$, hence to $m = \overline{j-1}$, and so is the right side by (8.22) for $j < k$.

Now let $Sx = \bar{k}$. Then $h_k(A_k(x)) = A_k(\bar{k}) = q_k$, so $A_k(x) = q_{k-1}$ and $x = \overline{k-1}$. The converse is immediate. For the distinctness of $\overline{k+2}$: an equality $\overline{k+2} = \bar{j}$ with $1 \leq j \leq k$ reduces by (8.22) for j to $\overline{k+1} = \overline{j-1}$; equality with 0 is excluded by Theorem 8.58 and equality with $\overline{k+1}$ by Theorem 8.61. □

This is a family of derivations, one for each k, and not one theorem that quantifies over k. It does not exclude a collision $Sx = Sy$, $x \neq y$, over a nonstandard target.

Lemma 8.66 (An equivalent form of injectivity)**.** *In the extension by P put $T(n) := P(Sn)$. Then*

$$S(T(n)) = Sn, \qquad T(T(n)) = T(n), \qquad T(0) = 0,$$

and over T_F^- injectivity of the successor is equivalent to $\forall n\,(T(n) = n \to T(Sn) = Sn)$.

Proof. The first equation is Theorem 8.64; applying P to it gives the second; the third is $P(1) = 0$. If the fixed points of T are closed under successor, quantifier-free induction on $T(n) = n$ gives $T(n) = n$ for all n, and from $Sa = Sb$ one gets $a = T(a) = T(b) = b$. Conversely injectivity applied to $S(T(n)) = Sn$ gives $T(n) = n$ for all n. □

Remark 8.67. `scope_and_coverage` and `coverage_and_kernel` contain the protected orbit with the same guard, under the names V, W and U, p_*; the first derives $n = 0 \vee n = S(P(n))$ by induction on that formula in the extension by P, the second by the induction of the proof above. Theorems 8.62 and 8.66 are from the first package and Theorems 8.61 and 8.65 from the second. Neither derives injectivity. The first package records why a comparison of the orbits with targets Sn and S^2n does not close the gap: for $v \neq 0$ the term $S(\theta_{Sn}(v))$ stops when $Sv = Sn$ and $\theta_{S^2n}(Sv)$ stops when $S^2v = S^2n$, so the intertwining of the two orbits needs $S^2v = S^2n \to Sv = Sn$ at the visited states, which is an instance of the missing cancellation.

8.3.3 With injectivity: predecessor and order

From here on the theory is T_F.

Theorem 8.68 (Predecessor)**.** T_F *proves* $p_*(S^2m) = Sm$*, and for the finite-case function* P *of Theorem 8.64*

$$P(0) = 0, \qquad P(Sn) = n, \qquad n = 0 \vee n = S(P(n)).$$

Proof. Injectivity applied to Theorem 8.63 gives the first equation. By coverage $n = 0$ or $n = Sm$. In the first case $P(Sn) = P(1) = 0$. In the second S^2m differs from 0 and, by injectivity and $Sm \neq 0$, from 1, so $P(S^2m) = p_*(S^2m) = Sm$. □

Write $w(a) := p_*(a)$. The constructions below use this term only through

$$w(S^3m) = S^2m, \tag{8.23}$$

that is, w removes one successor from every value that is at least 3, and $w(w(\cdot))$ removes two from every value that is at least 4.

Remark 8.69. This order of proof, coverage first and then the predecessor with corrections at 0 and 1, is that of `coverage_and_kernel`. `selfseed_scalar` uses the orbit without the guard at zero, $V(j,a) := F(j; v.C(\varepsilon(Sv,a),v))$ and $W(a) := V(a,a)$, with the invariant $a \neq 1 \to (V(j,a) = S^2j \vee S(V(j,a)) = a)$, which gives (8.23) for W and the values $W(0) = 2$, $W(1) = 2$, $W(2) = 4$. Its predecessor graph has the exceptional inputs $0, 1, 2$, and predecessor and coverage are proved together by quantifier-free induction on $(n = 0 \vee n = S(P(n))) \wedge P(Sn) = n$. Both proofs cancel a successor at the step from $S(V(m,a)) = a$ to the value of $V(m,a)$, which is where T_F uses its injectivity axiom.

For a parameter a define the capped counter and the comparison term

$$\lambda_a(v) := C(\varepsilon(v, S^2a), v), \qquad \mathrm{Cap}(n,a) := F(n; v.\lambda_a(v)), \qquad L(a,n) := \varepsilon(\mathrm{Cap}(n,a), S^2a),$$

and write $a \leq n$ for $L(a,n) = 1$. The update λ_a holds the value S^2a and increments every other value; since $S^2a \neq 0$, $\mathrm{Cap}(0,a) = 2$.

Lemma 8.70 (Capped counting and order)**.** T_F *proves*

$$\mathrm{Cap}(n,a) = S^2n \vee \mathrm{Cap}(n,a) = S^2a, \qquad \mathrm{Cap}(a,a) = S^2a, \qquad \mathrm{Cap}(Sn,Sa) = S(\mathrm{Cap}(n,a)),$$
$$L(0,n) = 1, \qquad L(Sa,0) = 0, \qquad L(Sa,Sn) = L(a,n),$$
$$a \leq Sb \leftrightarrow (a \leq b \vee a = Sb),$$

and that $\leq$ *is a discrete linear order with least element* 0 *which the successor preserves and reflects. The finite-case function* $\min(n,a)$*, with value* n *if* $n \leq a$ *and* a *otherwise, satisfies* $\mathrm{Cap}(n,a) = S^2\min(n,a)$.

Proof. The first statement is a quantifier-free induction on n with a fixed: the base is $\mathrm{Cap}(0,a) = S^20$; a state S^2a is held; and a state $S^2n \neq S^2a$ is incremented to S^2Sn. At $n = a$ both alternatives give the second statement. In particular $\mathrm{Cap}(n,0) = 2$.

Coverage gives $C(c,Sv) = S(C(c,v))$, and reflection with injectivity gives $\varepsilon(Sv,S^3a) = \varepsilon(v,S^2a)$; hence $\lambda_{Sa}(Sv) = S(\lambda_a(v))$. Also $\mathrm{Cap}(1,Sa) = \lambda_{Sa}(2) = 3$, because $S^3a \neq 2$. Induction on n for the literal equation, with a fixed, gives the third statement: the base is $3 = S2$ and the step applies λ_{Sa} and the last identity.

$L(0,n) = 1$ by $\mathrm{Cap}(n,0) = 2$; $L(Sa,0) = \varepsilon(2,S^3a) = 0$; and $L(Sa,Sn) = \varepsilon(S\,\mathrm{Cap}(n,a), SS^2a) = L(a,n)$.

For the equivalence, let $\mathrm{Cap}(Sb,a) = S^2a$. If $\mathrm{Cap}(b,a) = S^2a$, then $a \le b$. Otherwise $\mathrm{Cap}(b,a) = S^2b$ by the first statement, the next value is S^3b, and $S^3b = S^2a$ gives $a = Sb$. Conversely a reached cap persists, and $\mathrm{Cap}(Sb,Sb) = S^2Sb$.

Linearity. Fix a and induct on n in $a \le n \vee n \le a$. The base is $0 \le a$. In the step, $a \le n$ persists. If $n \le a$, split $a = 0$ or $a = Sb$ by coverage. If $a = 0$, then $\mathrm{Cap}(0,n) = 2 = S^2n$ gives $n = 0$ and $a \le Sn$. If $a = Sb$, the equivalence gives $n \le b$, and then $Sn \le Sb = a$, or $n = a$, and then $a \le Sn$.

Irreflexivity and descent. $Sn \not\le n$ by induction on the literal equation $L(Sn,n) = 0$. $Sn \le a \to n \le a$: by coverage $a = 0$ contradicts $L(Sn,0) = 0$, and $a = Sb$ gives $n \le b$ and then $n \le Sb$ by persistence.

Antisymmetry. Fix a and induct on n in $(a \le n \wedge n \le a) \to a = n$. At Sn the equivalence splits $a \le Sn$ into $a = Sn$, or $a \le n$; in the latter case $Sn \le a$ gives $n \le a$, the hypothesis gives $a = n$, and $Sn \le n$ is a contradiction.

Transitivity. Fix a, b and induct on c in $(a \le b \wedge b \le c) \to a \le c$. At Sc the equivalence splits $b \le Sc$ into $b \le c$, where the hypothesis and persistence apply, or $b = Sc$, where substitution applies.

For the last statement: if $n \le a$, the first statement gives $\mathrm{Cap}(n,a) = S^2n$ directly, or $\mathrm{Cap}(n,a) = S^2a$, which is $a \le n$ and gives $a = n$ by antisymmetry. If $n \not\le a$, then $a \le n$ by linearity, which is $\mathrm{Cap}(n,a) = S^2a$. □

Lemma 8.71 (Bounded selection)**.** *For Boolean c,p,q let $\mu(c,p,q) := (c \wedge_B p) \vee_B (\neg_B c \wedge_B q)$, and let*

$$\Theta(b,c,a,d) := F\big(b;\, v.\, C(\mu(c,\varepsilon(v,S^2a),\varepsilon(v,S^2d)),v)\big), \qquad \mathrm{Sel}(b,c,a,d) := w(w(\Theta(b,c,a,d))).$$

If $a = S^2a_0$ and $a \le b$, then $c = 1 \to \mathrm{Sel}(b,c,a,d) = a$. If $d = S^2d_0$ and $d \le b$, then $c = 0 \to \mathrm{Sel}(b,c,a,d) = d$.

Proof. If $c = 1$ the body equals $\lambda_a(v)$ for a generic v, so $\Theta(b,c,a,d) = \mathrm{Cap}(b,a)$ by induction on the equation between the two recursors at a counter variable; similarly $\Theta(b,c,a,d) = \mathrm{Cap}(b,d)$ if $c = 0$. Under the hypotheses the selected value is $S^2a = S^4a_0$ or $S^2d = S^4d_0$, and two applications of (8.23) remove two successors. □

8.3.4 Bounded iteration of finite-case functions

Write $e(x) := S^2x$. On an input $u = e(x)$ the finite cases

$$u = 2 :\ 0; \qquad u = 3 :\ 1; \qquad u \ne 2,3 :\ w(w(u)) \tag{8.24}$$

return x: in the last case $u = S^4x'$ by coverage. For $B = e(b)$ let $\mathrm{clip}_B(v) := w(w(\mathrm{Cap}(v,B)))$. For $v \ge 2$, Theorem 8.70 and (8.23) give $\mathrm{clip}_B(v) = \min(v,B)$ and $2 \le \mathrm{clip}_B(v) \le B$. The term is applied only to arguments $e(t)$ or to values in $[2,B]$.

Proposition 8.72 (Bounded iteration)**.** *Let $g(\vec{p})$ and $h(v,\vec{p})$ be finite-case functions. For every cap variable b there is a finite-case function $\mathrm{It}_b(n,\vec{p})$, obtained from terms of the original language by the cases* (8.24)*, such that T_F proves*

$$\mathrm{It}_b(0,\vec{p}) = \min(g(\vec{p}),b), \qquad \mathrm{It}_b(Sn,\vec{p}) = \min\big(h(\mathrm{It}_b(n,\vec{p}),\vec{p}),b\big), \qquad \mathrm{It}_b(n,\vec{p}) \le b.$$

Every occurrence of F in the construction has a body that is a term of the original language.

Proof. Put $B := e(b)$. Let g have the cases (A_i,t_i). Encode and clip each candidate as $\mathrm{Cap}(t_i,b) = e(\min(t_i,b))$, which lies in $[2,B]$ for all inputs, and fold the cases by nested

$\mathrm{Sel}(B, \chi_{A_i}, \cdot, \cdot)$ with the default value 2. By Theorem 8.71 this gives a term $a_B(\vec{p})$ in $[2, B]$ with $a_B(\vec{p}) = e(\min(g(\vec{p}), b))$.

For the update, compose the cases of h with the cases (8.24) for its state argument, refine the guards, and encode, clip and select in the same way. The result is a term $h_B(u, \vec{p})$ of the original language with $2 \le h_B(u, \vec{p}) \le B$ for every u, because every candidate is a value of $\mathrm{Cap}(\cdot, b)$, and

$$h_B(e(v), \vec{p}) = e(\min(h(v, \vec{p}), b)).$$

All guard tests are Boolean terms; nothing depends on an order of evaluation.

The seed of the recursor is tied to the body, so the initial value a_B is entered through two transient states. Let

$$H_B(u) := \mathrm{Sel}\Big(S^2B, \varepsilon(u, 0), B,\ \mathrm{Sel}(S^2B, \varepsilon(u, SB), S^2B,\ \mathrm{Sel}(S^2B, \varepsilon(u, S^2B), a_B, h_B(u)))\Big).$$

All candidates lie in $[2, S^2B]$. The sentinels SB and S^2B are distinct and exceed B, so

$$H_B(0) = B, \qquad H_B(SB) = S^2B, \qquad H_B(S^2B) = a_B, \qquad 2 \le u \le B \to H_B(u) = h_B(u),$$

and therefore $F(0; u.H_B) = SB$, $F(1; u.H_B) = S^2B$, $F(2; u.H_B) = a_B$. Put $Y_b(n, \vec{p}) := F(e(n); u.H_B(u))$. Quantifier-free induction gives $2 \le Y_b(n, \vec{p}) \le B$, $Y_b(0, \vec{p}) = a_B$ and $Y_b(Sn, \vec{p}) = h_B(Y_b(n, \vec{p}), \vec{p})$. Applying the cases (8.24) to Y_b gives It_b. □

The proposition gives initialized, capped iteration of finite-case functions whose update sees the state only. Each use below supplies a cap that is a term of the original language in one variable and a proof that clipping does not occur on the intended arguments.

8.3.5 Arithmetic by finite-case graphs

All functions introduced in this subsection are finite-case, every induction is quantifier-free after Theorem 8.60, and Theorem 8.72 is used whenever an update refers to one of them.

Lemma 8.73 (Difference and doubling)**.** *There is a finite-case function d with*

$$d(x, 0) = x, \qquad d(x, Sn) = P(d(x, n)), \qquad d(x, n) \le x, \qquad d(Sx, Sy) = d(x, y), \qquad d(n, n) = 0.$$

The term $Q(n) := F(n; v.S^2v)$ and the finite-case function $\mathrm{db}(n) := P(P(P(Q(n))))$ satisfy

$$Q(0) = 3, \quad Q(Sn) = S^2Q(n), \qquad \mathrm{db}(0) = 0, \quad \mathrm{db}(Sn) = S^2\,\mathrm{db}(n), \quad Q(n) = S^3\,\mathrm{db}(n),$$

and db *is monotone with $n \le \mathrm{db}(n)$.*

Proof. Let $d(x, n)$ be Theorem 8.72 with initial value x, update P and cap x. Since $P(v) \le v$, clipping is never active, which gives the first three statements. $d(Sx, Sy) = d(x, y)$ by induction on y with x fixed: the base is $P(Sx) = x$ and the step applies P. Then $d(n, n) = 0$ by induction. The equations for Q are (8.21). By induction $Q(n) \ge 3$, so the functional coverage $n = 0 \vee n = S(P(n))$ cancels three successors and gives the equations for db. $n \le \mathrm{db}(n)$ is a quantifier-free induction, and monotonicity is an induction on the larger argument in $a \le b \to \mathrm{db}(a) \le \mathrm{db}(b)$ with the equivalence of Theorem 8.70. □

Lemma 8.74 (Addition)**.** *There is a finite-case function $+$ with $0+a = a$, $Sn+a = S(n+a)$, commutative, associative and monotone, and with $n + a < Q(b)$ whenever $n, a \le b$.*

Proof. For a cap c let $A_c(n,a)$ be Theorem 8.72 with initial value a and update S, and write $s_c(v) := \min(Sv, c)$:

$$A_c(0,a) = \min(a,c), \qquad A_c(Sn,a) = s_c(A_c(n,a)).$$

For $a \le c$, induction on the equation gives the seed commutation $A_c(n, s_c(a)) = s_c(A_c(n,a))$: both sides have the same value at 0 and the same update. The minimum satisfies $\min(0,c) = 0$ and $\min(Sn,c) = s_c(\min(n,c))$ by the order laws, so $A_c(n,0) = \min(n,c)$ by uniqueness of iteration.

Fix $a \le c$ and induct on n in $n \le c \to A_c(n,a) = A_c(a,n)$. The base is the last identity. If $Sn \le c$, the step applies s_c on the left and the seed commutation on the right. Since the update increases and s_c preserves order, A_c is monotone in the counter (induction on the larger counter) and in the initial value within $[0,c]$ (induction on the common counter).

Fix an input bound b and take the cap $c := Q(b) = S^3\,\mathrm{db}(b)$, a term of the original language. Induct on n in $n \le b \to A_c(n,n) = \mathrm{db}(n)$: at a successor the seed commutation gives $A_c(Sn,Sn) = s_c(s_c(A_c(n,n)))$, and since $\mathrm{db}(Sn) \le \mathrm{db}(b) < c$ neither step is clipped, so the value is $S^2\,\mathrm{db}(n)$. By monotonicity

$$n, a \le b \to A_{Q(b)}(n,a) \le A_{Q(b)}(b,b) = \mathrm{db}(b) < Q(b),$$

so there is no clipping on bounded inputs. For the caps of two input bounds b, b', induction on n restricted by $n \le b, b'$ shows that the results agree for $a \le b, b'$: the values at 0 are a and both steps are unclipped successors.

Take b to be the first largest of the two arguments, by finitely many order cases; this defines the finite-case function $n + a$. Coherence across the caps proves $0 + a = a$ and $Sn + a = S(n + a)$, bounded commutativity proves commutativity, and associativity and monotonicity are quantifier-free inductions with fixed parameters. □

Lemma 8.75 (Multiplication and pairing)**.** *Let $E(n) := F(n; v.Q(v))$, so that $E(0) = 4$ and $E(Sn) = Q(E(n))$. There are finite-case functions $\cdot$, κ, π_0, π_1 with*

$$n \cdot a \le E(Q(b)) \text{ for } n, a \le b, \qquad \kappa(x,y) := (x+y)^2 + x \le Q(E(Q(Q(b)))) \text{ for } x, y \le b,$$
$$\pi_0(\kappa(x,y)) = x, \qquad \pi_1(\kappa(x,y)) = y, \qquad \pi_0(z) \le z, \qquad \pi_1(z) \le z,$$

and the semiring identities for $+$ and $\cdot$.

Proof. E is monotone, $E(n) \ge n$ and $E(Sn) = \mathrm{db}(E(n)) + 3$. Fix $n, a \le b$ and apply Theorem 8.72 with initial value 0, update $v \mapsto v + a$ and cap $E(Q(b))$. For the state v_j at a stage $j \le b$ the invariant $v_j \le E(b + j)$ holds: the base is immediate, and

$$v_j + a \le E(b+j) + b \le \mathrm{db}(E(b+j)) < E(S(b+j)),$$

using $E(b+j) \ge b$. Since $b + j \le \mathrm{db}(b) < Q(b)$, the stage bound is at most the cap, so clipping is inactive. Cap coherence as for addition gives a finite-case multiplication with the bound $E(Q(b))$; its equations at 0 and at a successor give the semiring identities by quantifier-free induction, in particular $w^2 + w < (Sw)^2$.

κ is finite-case with the stated bound, which is monotone and at least its argument, and both arguments of a code are at most the code. For the root, apply Theorem 8.72 with initial value 0, the update that sends u to Su if $(Su)^2 \le z$ and to u otherwise, the cap z and the counter z. The update preserves $u \le z$ and $u^2 \le z$, and the quantifier-free invariant "$u_j = j$ or the update fixes u_j" shows that the state at $j = z$ is fixed: if it equals z, then $(Sz)^2 > z$. So the result $r(z)$ satisfies $r(z)^2 \le z < (Sr(z))^2$, and it is unique with this property by monotonicity of squares. Put

$$\pi_0(z) := d(z, r(z)^2), \qquad \pi_1(z) := d(r(z), \pi_0(z)).$$

The identity $d(x+y,y) = x$ follows by induction on y from $d(Sx,Sy) = d(x,y)$ and the addition equations. For $z = \kappa(x,y)$ and $w := x+y$, the inequalities $w^2 \le z \le w^2+w < (Sw)^2$ give $r(z) = w$, hence the projection equations. The projections are total on all arguments and bounded by their argument, since a difference is bounded by its first argument. □

Theorem 8.76 (Primitive recursion)**.** *For every primitive recursive description f there are a finite-case function f^* in T_F, proofs of the defining equations of f for f^*, and a term M_f in one variable, monotone and with $b \le M_f(b)$, such that $\vec{x} \le b \to f^*(\vec{x}) \le M_f(b)$ is provable.*

Proof. By recursion on f. Zero, successor and projections have these data with the bound Q. For a composition $f = h(g_1,\dots,g_m)$ take a common refinement of the cases and the bound $M_h \circ M_{g_1} \circ \cdots \circ M_{g_m}$. Nullary descriptions have numeral values and constant shifts of b as bounds.

Let $f(0,\vec{p}) = g(\vec{p})$ and $f(Sn,\vec{p}) = h(n,f(n,\vec{p}),\vec{p})$. Write K_κ for the bound of κ in Theorem 8.75 and put

$$\Delta := K_\kappa \circ M_h \circ M_g \circ Q, \qquad \Upsilon(t) := F(t; v.\Delta(v)), \qquad \beta(b) := \Upsilon(SQ(b)).$$

Each factor is monotone and at least its argument, and $\Delta(v) \ge Sv$; so Υ is monotone with $\Upsilon(t) \ge St$, and β is monotone with $\beta(b) \ge b$, by quantifier-free inductions. The state update is the finite-case function

$$J(z,\vec{p}) := \kappa\big(S\pi_0(z),\, h^*(\pi_0(z),\pi_1(z),\vec{p})\big),$$

with the initial value $\kappa(0,g^*(\vec{p}))$. Apply Theorem 8.72 to them with the cap $\beta(b)$ and let z_j be the state at the stage j. For $n,\vec{p} \le b$ and $j \le b$,

$$z_j \le \Upsilon(Sb+j). \tag{8.25}$$

The seed satisfies $\kappa(0,g^*(\vec{p})) \le K_\kappa(M_g(b)) \le \Delta(b) \le \Delta(\Upsilon(b)) = \Upsilon(Sb)$. For the step write $s := \Upsilon(Sb+j)$. Then $s \ge b$, both projections of z_j are at most s, the next counter is at most $Ss \le Q(s)$ and the next value is at most $M_h(M_g(Q(s)))$, so $J(z_j,\vec{p}) \le \Delta(s) = \Upsilon(S(Sb+j))$. Since $j \le b$, the argument $Sb+j$ is at most $S\,\mathrm{db}(b) \le SQ(b)$, so every stage bound is at most $\beta(b)$ and no clipping occurs. The invariant (8.25) is quantifier-free in the extension.

By the projection equations and a quantifier-free induction guarded by $j \le b$, the first component of z_j is j, and the second component satisfies the base equation and, for $Sj \le b$, the step equation. Two caps give the same run up to their common bound on the counter: the seeds agree and the unclipped updates are identical. Take b to be the first largest among $n,\vec{p}$ and define $f^*(n,\vec{p}) := \pi_1(z_n)$. For any larger input bound, cap coherence identifies this value with the run at that cap, so $f^*(n,\vec{p}) \le \beta(b)$ for every valid b; put $M_f := \beta$. Cap coherence proves the defining equations also when a successor input changes the first largest coordinate.

Theorem 8.72 expands every use of g^*, h^*, κ, π_i inside the update into terms of the original language, encoded cases and bounded selections. There is no evaluator for codes of descriptions and no recursor over functions defined by graphs. □

Theorem 8.77 (Bi-interpretation)**.** T_F *and* iPRA *are bi-interpretable, with whole domains, actual equality, the logic translated identically and the identity as isomorphism of both composites. They are equiconsistent.*

Proof. In iPRA keep 0, S and equality, interpret ε as the primitive recursive characteristic of equality, and each recursor with body h by primitive recursion with initial value $S(h(0))$ and a step h that ignores the counter. All axioms of T_F and its induction rule hold. Conversely interpret each description by the finite-case function of Theorem 8.76; totality, single-valuedness, the defining equations and the quantifier-free induction instances are proved in T_F by Theorem 8.60, and injectivity and separation are axioms.

On the side of iPRA, recursion on a description reduces the identification of a function with its double interpretation to uniqueness from the base and step equations, which is a quantifier-free induction. On the side of T_F, 0 and S are unchanged; the two characteristics of equality are two-valued and reflect equality, so they agree by cases; and for a recursor with body h, the earlier stage identifies the body, both functions satisfy (8.21) for it, and induction on the equation between them identifies them, with Theorem 8.60 where graph-defined functions occur. Induction on formulas extends this to all formulas, and falsity is preserved. □

8.4 Three observations on the native core

The packages of this chapter contain three results about the calculus N on iterator terms that do not concern a seeded recursor. They are recorded here from `coverage_and_kernel`, `transient_seed` and `monus_and_neutral_major`.

8.4.1 One native binder over a literal equation

In this subsection formulas are read in iPRA through Section 1.2 and Theorem 1.78, and sel, the predecessor P of Theorem 1.48, the characteristic terms of Theorem 1.52 and the pairing of Theorem 1.62 are available. A formula is called Σ_2^{qf} if it has the form $\exists x\,\forall y\,\theta(x,y,\vec{p})$ with θ quantifier-free, and Π_1^{qf} if it has the form $\forall y\,\theta(y,\vec{p})$. These are classes of prenex formulas; no prenex normal form theorem for intuitionistic formulas is used.

Proposition 8.78 (One binder)**.** *Up to provable equivalence, the native formulas $D_{x,z}E(x,z,\vec{p})$ with E a literal equation are exactly the Σ_2^{qf} formulas, and the native formulas $Q_zE(z,\vec{p})$ with E a literal equation are exactly the Π_1^{qf} formulas.*

Proof. The reading of $D_{x,z}E$ is $\exists x\,(E(x,0)\to\forall z\,E(x,z))$, which is equivalent to $\exists x\,\forall z\,(E(x,0)\to E(x,z))$ because z does not occur in $E(x,0)$; the matrix is quantifier-free. Without the existential quantifier this is the statement for Q_z.

Conversely let $r(x,y,\vec{p})$ be a characteristic term with $r=0\leftrightarrow\theta$, and let $E(x,z,\vec{p})$ be the literal equation

$$\mathrm{sel}\big(z,0,r(x,P(z),\vec{p})\big)=0.$$

Its instance at $z=0$ is provable, and its instance at $z=Sy$ is equivalent to $r(x,y,\vec{p})=0$ by $P(Sy)=y$. By coverage, $\forall z\,E(x,z,\vec{p})\leftrightarrow\forall y\,\theta(x,y,\vec{p})$, and the provable antecedent $E(x,0,\vec{p})$ drops out of the reading of $D_{x,z}E$. □

Corollary 8.79. *If rule QI is extended to matrices that consist of one $D_{x,y}$ over a literal equation, the extended calculus derives every induction instance for Σ_2^{qf} formulas. If it is extended to matrices Q_yE with E a literal equation, it derives every induction instance for Π_1^{qf} formulas.*

Proof. Let $B(n,\vec{p})$ be the native formula of Theorem 8.78 for the given formula, and let $\Gamma\vdash B(0,\vec{p})$ and $\Gamma,B(z,\vec{p})\vdash B(Sz,\vec{p})$. The extended rule gives $Q_nB(n,\vec{p})$, and QE with the base gives $B(t,\vec{p})$ for every term t. The equivalence of the proposition transports the derivation to the given formula. □

A guard "at most one binder" is therefore not equivalent to the guard "literal equation" unless an elimination or conservativity theorem for the additional induction instances is proved. No statement on the strength of the extended calculus relative to iPRA is made.

For a native formula let $\delta(s=t):=0$ and $\delta(Q_yA):=\delta(D_{x,y}A):=1+\delta(A)$. For a fresh z let G_zA be A with every equation $s=t$ replaced by $\mathrm{sel}(z,0,s)=\mathrm{sel}(z,0,t)$. Then $\delta(G_zA)=\delta(A)$, G_0A is provable, and $G_{Su}A\leftrightarrow A$: for equations by computation, and for a binder by induction on the formula, with the witness 0 for $D_{x,y}$.

Theorem 8.80 (Alternating prefixes)**.** *For every $k \geq 0$, every formula $\exists x_1 \forall y_1 \cdots \exists x_k \forall y_k \, \theta(\vec{x}, \vec{y}, \vec{p})$ with θ quantifier-free is provably equivalent in* iPRA *to the reading of a native formula of depth $\delta \leq k$. Consequently an extension of QI to all matrices of depth at most k derives induction for this class.*

Proof. For $k = 0$ take the characteristic equation. If $A(x, y, \vec{p})$ translates the remaining prefix at depth at most $k-1$, put $\mathcal{B}(A) := D_{x,z} \, G_z A(x, P(z), \vec{p})$. Its matrix at $z = 0$ is provable and its matrix at $z = Sy$ is equivalent to $A(x, y, \vec{p})$, so by coverage $\mathcal{B}(A) \leftrightarrow \exists x \, \forall y \, A(x, y, \vec{p})$, and $\delta(\mathcal{B}(A)) \leq k$. The induction statement is the argument of Theorem 8.79. □

Blocks of several variables are packed by pairing, and dummy blocks can be added because the domain is inhabited; so the inclusion covers the prenex Σ_{2k} formulas and, for $k \geq 1$, the prenex Π_{2k-1} formulas. At depth greater than one only this inclusion is claimed: the reading of a nested native formula contains implications with quantified antecedents, and moving quantifiers through them is not valid intuitionistically.

8.4.2 The case rule at a literal equation cannot be deleted

Proposition 8.81. *Consider the calculus on iterator terms with paired introduction by cases and the progress rule on $D_{x,y}$ of Theorem 8.26, and restrict the introduction so that its local assumption is $E(t, z)$ whenever the matrix is a literal equation E, and $B(t, 0)$ otherwise. The restricted calculus does not derive $D_{x,y}(y = S0)$, which the unrestricted calculus derives.*

Proof. The unrestricted calculus derives the formula by DI by cases: under the local assumption $0 = S0$ every equation follows. For the restricted calculus read equations as equality on the standard numbers, $Q_y B$ as $B(0) \to \forall y \, B(y)$, and

$$(D_{x,y} B)^\dagger := \begin{cases} \exists x \, \forall y \, (B(x, y) \to B(x, Sy)), & B \text{ a literal equation}, \\ \exists x \, (B^\dagger(x, 0) \to \forall y \, B^\dagger(x, y)), & B \text{ compound}. \end{cases}$$

Every rule of the restricted calculus is sound for this reading: a progress witness for an equation gives its closure by induction on the standard numbers, so DE is sound; compound case introduction is sound because every positive number is a successor; and transport preserves the distinction between literal and compound matrices. The formula $D_{x,y}(y = S0)$ is false in this reading, because $y = 1$ holds at 1 and fails at 2. □

Under the standard reading the same formula is true, since its zero instance is false. Hence the two readings of a $D_{x,y}$ over an equation, closure and progress, cannot be exchanged by padding the equation into a compound formula without losing one direction of the formula round trip.

8.4.3 Transport from the literal zero alone

Proposition 8.82. *In the calculus on iterator terms with formula transport (Theorem 8.26) and no reflexivity axiom, replace transport by the rule: from $\Gamma \vdash 0 = t$ and $\Gamma \vdash F(0)$ infer $\Gamma \vdash F(t)$. The resulting calculus does not derive $0 = 0$.*

Proof. Interpret terms in $\{0, 1\}$ with $S0 = S1 = 1$ and every iterator term equal to 1, read an equation $s = t$ as "the value of s is 1", and read Q_y, $D_{x,y}$ by the clauses of Section 1.2. Every computation equation holds, because its left side is an iterator. No equation $0 = t$ holds, so the restricted rule is sound. Every element is 0 or a successor, and induction holds for every predicate on the domain: the base gives the value at 0 and the step at 0 gives the value at 1. So DI by cases and the progress rule are sound, as are Hyp, QE and DE. The equation $0 = 0$ does not hold. □

The relation used in the proof is not equality. With unrestricted transport $0 = 0$ follows from the computation $I(0, 0; v.\, 0) = 0$ used as both premises with the template $w = 0$. The proposition does not apply to a calculus that has $0 = 0$ as an axiom or as an initial context entry.

8.5 Checker sizes

All eight packages report sizes of complete standalone checkers under the protocol of Section 1.10. Table 8.2 lists them as stated, with the control each package measures against. Only the programs of the first three packages that are marked R check the successor-seeded calculus N_R. All other programs check the calculus N on iterator terms; no package reports a checker for $\mathsf{T}_{B\varepsilon}$, T_{Bd}, N_{Bd}, T_F or T_F^-, and the sizes below are not sizes of those theories.

Table 8.2: Checker sizes as stated by the packages (tokens, AST nodes, normalized bytes).

Package	Program	Tokens	AST
`scalar_successor`	`checker.py` (R)	279	319
	`checker_bytes.py` (R)	279	321
	`checker_ast.py` (R)	282	314
	control: generated assumptions, main	281	318
	control: generated assumptions, AST profile	283	315
	control: guarded elimination, token profile	281	325
`transient_seed`	`checker_sr_bytes.py` (R)	284	322
	`checker_bytes.py` (iterator)	282	320
	`checker_structure.py` (iterator)	280	311
	`checker_joint.py` (iterator, predecessor packets)	280	314
	control: generated assumptions, main	281	318
	control: generated assumptions, AST profile	283	315
`successor_seed_completion`	`checker_direct.py` (R)	283	320
	`checker.py` (R)	284	322
	`checker_ast.py` (R)	282	313
	`checker_ast_total.py` (R)	287	314
	control: prior generated-assumption checker	281	318
	control: prior AST-oriented checker	283	315
`bounded_initializer`	`checker.py`	266	310
	`checker_compact.py`	266	312
	`checker_ast.py`	267	298
	control: byte leader	269	311
	control: AST and token leader	268	300
`monus_and_neutral_major`	`checker.py`	263	307
	`checker_bytes.py`	263	309
	`checker_joint.py`	261	295
	control: certified zero, explicit list test	264	308
	control: certified zero, compact list test	264	310
	control: certified zero, generic traversal	262	296
`selfseed_scalar`	`local_argument.py`	267	309
	`local_argument_bytes.py`	267	311

Package	Program	Tokens	AST
	`local_loop_joint.py`	263	291
	`local_loop_joint_separate.py`	262	293
	control: `baseline_checker.py`	264	308
	control: `baseline_checker_bytes.py`	264	310
	control: `baseline_checker_joint.py`	262	296
`scope_and_coverage`	`checker_compact.py`	266	310
	`checker_explicit.py`	266	308
	control: neutral major, global floor (byte incumbent)	263	309
	control: guarded major, lexical floor	267	311
	control: neutral major, lexical floor, conditional dispatch	266	310
`coverage_and_kernel`	`checker.py`	266	308
	`checker_bytes.py`	266	310
	`checker_ast.py`	263	289
	control: byte leader	263	309
	control: token leader	261	295
	control: AST leader	263	291

In every package the minima over the three coordinates are attained by different programs, and each package states that no single one of its programs is below all three coordinates of its control, with one exception: `checker_joint.py` of `transient_seed`, at 280/314/684, is below 281/315/685, the coordinatewise minimum of its two controls. The packages state that these minima are sizes attained in a fixed corpus of earlier checkers and not lower bounds.

Successor-seeded checkers. The R programs derive from the checker with generated assumptions of an earlier draft: a certificate is a JSON array whose last element is the closed goal and whose earlier four-field packets are consumed from the end; a computation equation enters as a local assumption (Theorem 8.26). An arithmetic packet supplies the data $[n, a, h]$ and a flag, and the checker itself builds the counter 0 or Sn of the generated equation, so no shape test on a supplied counter is needed. The same change applied to the iterator gives the matched iterator programs of `transient_seed`, whose joint profile uses a changed packet format for predecessor computations. The AST profiles accept additional raw aliases and reject malformed nonempty strings only through the recursion limit of the interpreter; `checker_ast_total.py` rejects them directly. `transient_seed` ships kernel certificates for the Boolean decoder (45 commands) and for a restricted implication (149 commands); `successor_seed_completion` states that the full predecessor derivation has not been emitted as a kernel certificate.

Iterator checkers in the other packages. The programs of `bounded_initializer` compose an arity encoding of the raw syntax with a predecessor supply. Those of `monus_and_neutral_major` implement its neutral-major certificate protocol for N, in which every instruction proves a major premise and a derivable truth serves where none is needed; the protocol is not described in this chapter. Those of `selfseed_scalar`, `scope_and_coverage` and `coverage_and_kernel` compose lexical proof scopes, the arity encoding, the certified-zero dispatch and, in the last two, neutral majors; the loop variants keep contexts as immutable tuples. `selfseed_scalar` records two failed variants: a loop with in-place mutation of the context list accepts a certificate of $0 = S0$ by leaking an assumption of a subproof, and a variant with opaque objects as fresh parameters accepts a certificate of $0 = S0$ because a floating-point leaf 3.0 is treated

as a rigid parameter while comparing equal to the bound index 3; the stricter repair of the latter loses the canonical certificate of universal reflexivity.

8.6 Further questions

No result of this chapter has a Lean formalization recorded in the repository; the packages present written proofs with finite numerical checks of the defining terms on standard numbers.

Question 8.83 (Injectivity for the self-seeded recursor)**.** Does T_F^- prove $Sx = Sy \to x = y$? By Theorem 8.66 this is equivalent to the closure under successor of the fixed points of $n \mapsto P(Sn)$, and by Theorem 8.63 to the derivation of $p_*(S^2m) = Sm$ from $S(p_*(S^2m)) = S^2m$. `selfseed_scalar`, `scope_and_coverage` and `coverage_and_kernel` leave it open. A countermodel has to satisfy all staged recursor equations and all instances of induction for literal equations, together with a successor collision; a branching successor graph without cycles is not enough. Neither the bi-interpretability of T_F^- with iPRA nor its failure is established.

Question 8.84 (Uniform rigidity)**.** Theorems 8.62 and 8.65 are families of derivations indexed by a fixed integer. Is there a single derivation with a variable index, without a general compiler for primitive recursion in T_F^-?

Question 8.85 (Shorter predecessor terms)**.** Is there a substantially shorter predecessor term for the successor-seeded recursor than those of Theorems 8.19 and 8.20, for instance a reset cycle that needs no Boolean comparator? The three packages prove existence and correctness on the whole domain and make no claim on size.

Question 8.86 (Other seed maps)**.** For which maps σ in place of the successor does the recursor with $R(0, a; u.h) = \sigma(a)$ recover arbitrary initialization from its two equations and induction for literal equations? A criterion has to provide a proved inverse and not assume one. Related: is there a general class of finite reset mechanisms for which the transient-seed method works, and is the correction of a capped predecessor at finitely many arguments an instance of a general repair principle?

Question 8.87 (Cancellation on finite-valued terms)**.** In N_R and N_{Bd} a restricted implication and contextual case analysis are obtained from cancellation of the successor on two-valued terms, before a selector exists (Theorems 8.9 and 8.55). Which other weak term languages admit this route to the full logical interface?

Question 8.88 (Tied seeds without an additional operation)**.** Theorem 8.31 excludes terms for addition, selection and pairing. It does not exclude interpretations of iPRA by relations, quotients or several coordinates in a theory on the terms $x \mid 0 \mid B(n; u, v.h)$ alone; Theorem 8.44 needs ε and Theorem 8.52 needs d and the coverage axiom. An invariant that is stable under such interpretations is not known from the packages of this chapter.

Question 8.89 (Fewer axioms for the equality test)**.** T_{Bd} replaces the range and reflection axioms of $\mathsf{T}_{B\varepsilon}$ by two recursive equations, at the price of coverage as an axiom. Can range and reflection be replaced by recursive equations from which coverage is still derived? `monus_and_neutral_major` investigates a basis with three equations for the minimum, in which parity$(\min(2x, 2y + 1))$ has the standard meaning of a comparator; the derivation of the equations of d and of reflection on the whole domain from that basis, without induction over a shifted parameter, is left open there.

Question 8.90 (Internal majorants and sizes of the compilers)**.** The converse direction of Theorem 8.45 produces a single term for a function with a majorant that is valid in the

standard model; a proof in $\mathsf{T}_{B\varepsilon}$ that this term equals the graph-defined function needs a proof of the majorant in the theory. Bounds on the size of the terms and proofs produced by Theorems 8.41, 8.43, 8.72 and 8.76, on the branching of the case expansions for nested terms, and on the cost of translating the graph-based proofs into a native calculus are not given.

Question 8.91 (Extended progress)**.** Theorems 8.79 and 8.80 show that an extension of QI to matrices with binders derives induction for prenex classes. Whether such an extension is conservative over N, or strictly stronger, is not decided; at depth greater than one only the inclusion of Theorem 8.80 is proved.

Question 8.92 (Variants of the two obstructions)**.** Theorem 8.81 refutes the deletion of case introduction at literal equations for the two displayed readings; it does not exclude a calculus in which a $D_{x,y}$ over an equation is an auxiliary judgment and the object formulas are padded. Theorem 8.82 does not exclude transport from the literal zero together with a reflexivity axiom. Neither variant is decided in the packages of this chapter; calculi with a single introduction rule are the subject of `witness_rule` and `guarded_core`, and restrictions on the source of transport of `source_dispatch`.

Question 8.93 (Checkers for the new bases)**.** No package measures a complete checker for $\mathsf{T}_{B\varepsilon}$, T_{Bd}, N_{Bd} or T_{F}, in which the equality test or the subtraction equations, the ordinary logic where it is used, and the validation of the staged bodies are all charged. Whether any of these bases, or N_{R}, admits a single checker below the controls of Table 8.2 in all three coordinates is open, as are lower bounds for a fixed checker language and input grammar.

Question 8.94 (Formalization)**.** The packages name as first targets for a proof assistant: the contextual case lemmas (Theorems 8.9 and 8.55), the factor map identities and the predecessor (Theorems 8.15 and 8.19), the gated complement (Theorem 8.50), cap coherence and finite-case elimination (Theorems 8.41, 8.43 and 8.60), and the coverage and rigidity proofs of Theorems 8.64 and 8.65.

Chapter 9

Canonical and counter-aware recursors

The common core takes the successor S and the iterator $I(n, a;\, u.\, h)$ as term constructors and states both computation equations as rules (Sections 1.1 and 1.3). This chapter treats the term languages in which the successor is an instance of the recursor, or the body of the recursor sees its counter, or both, and in which part of the computation is moved into a canonical grammar of terms with a normalizing substitution. Section 9.1 fixes the term languages and their normal forms. Section 9.2 shows what a counter-aware body provides: a decoder that computes, a selector, and a compiler for primitive recursion that uses no pairing. Section 9.3 gives the translations to and from iterator terms and transfers the bi-interpretation of Theorem 1.78. Sections 9.4 to 9.7 treat the proof calculi stated on these syntaxes. Section 9.8 treats the concrete syntax in which the recursor and the paired binder bind the same number of variables. Section 9.9 collects the boundary results.

9.1 Term languages and normal forms

This section draws on `register_graph`, `canonical_recursors`, `clausal_canonical`, `counter_evaluation`, `uniform_binding`, `uniform_recursion` and `single_constructor`.

9.1.1 Three recursors without a successor constructor

Definition 9.1 (Term languages)**.** Each of the following grammars has one numerical sort, variables, the constant 0 and one binding constructor.

1. The *seeded recursor*. Terms are $t ::= x \mid 0 \mid R(n, a; u.h)$; the variable u is bound in h only. Put $Sa := R(0, a; u.0)$ and $1 := S0$. The computation equations are

$$R(0, a; u.h) = Sa, \tag{9.1}$$
$$R(Sn, a; u.h) = h[R(n, a; u.h)/u]. \tag{9.2}$$

2. The *counter-aware seeded recursor*. Terms are $t ::= x \mid 0 \mid R(n, a; i, u.h)$; the variables i, u are bound in h only. The first receives the old counter and the second the preceding state. Put $Sa := R(0, a; i, u.0)$ and $1 := S0$. The computation equations are

$$R(0, a; i, u.h) = Sa, \tag{9.3}$$
$$R(Sn, a; i, u.h) = h[n/i,\ R(n, a; i, u.h)/u], \tag{9.4}$$

   with simultaneous capture-avoiding substitution on the right.

3. The *register iterator*. Terms are $t ::= x \mid 0 \mid J(n; i, r.h)$; the variables i, r are bound in h only. Put $Sa := J(a; i, r.0)$ and $1 := S0$. The term $J(n; i, r.h)$ executes n instructions on a register that starts at 1. The body receives the old counter i and the register r and returns an instruction: the value 0 increments the register, and a value St loads t. The axioms are stated in Theorem 9.13.

The number of bound variables distinguishes the first two constructors, so both are written R. In every language the variables of a body other than the bound ones are passive numerical parameters, and substitution is capture avoiding.

In all three languages the successor is a defined operation, not a constructor. The initializer of the seeded recursors is an argument, but the iteration starts at its successor; this is what allows the successor to be an instance of the recursor. The register iterator has no initializer argument at all.

The binding notation is a notation for finite descriptions in a first-order language, as in Section 1.1. Each fixed body, built from earlier descriptions, determines one function symbol whose arguments are the counter, the initializer where there is one, and the passive parameters of the body. No variable ranges over bodies, and no evaluator for descriptions is postulated inside any theory of this chapter.

9.1.2 Zero-body canonical terms

The equations (9.1) and (9.3) say that a recursor with counter 0 does not depend on its body. A canonical grammar records this by one spelling.

Definition 9.2 (Zero-body normalization)**.** For raw terms of either seeded language define ν by $\nu(x) = x$, $\nu(0) = 0$ and

$$\nu\big(R(n, a; \vec{u}.h)\big) = \begin{cases} R(0, \nu a; \vec{u}.0), & n \equiv 0, \\ R(\nu n, \nu a; \vec{u}.\nu h), & n \not\equiv 0, \end{cases}$$

where $\vec{u}$ is u or i, u and $\equiv$ is literal identity of syntax. It is not a test for provable equality with 0. A term is *zero-body canonical* if it is a fixed point of ν, that is, if every subterm with literal counter 0 has literal body 0. The map ν extends to formulas, clauses and contexts by normalizing all terms. For zero-body canonical T and a substitution σ the *normalized substitution* is

$$T\langle\sigma\rangle := \nu\big(T[\sigma]\big).$$

The map ν is a terminating traversal of a finite tree. It preserves the root constructor of a term and of a formula, so

$$\nu(t) \equiv 0 \iff t \equiv 0, \qquad \nu(Sa) = S(\nu a). \tag{9.5}$$

It is idempotent, commutes with renaming of free and bound variables, and introduces no free variable. It performs no successor computation.

Lemma 9.3 (Substitution coherence)**.** *For every raw term or formula T and every capture-free simultaneous substitution σ,*

$$\nu\big(T[\sigma]\big) = \nu\big(\nu(T)[\nu \circ \sigma]\big).$$

Consequently $(T\langle\sigma\rangle)\langle\tau\rangle = T\langle\sigma\tau\rangle$ for zero-body canonical T, where $(\sigma\tau)(x) = \sigma(x)\langle\tau\rangle$.

Proof. Induct on T with bound names chosen fresh for σ. For a variable both sides are $\nu(\sigma(x))$ by idempotence, and the case 0 is trivial. Let $T = R(n, a; \vec{u}.h)$.

If $n \equiv 0$, then $n[\sigma] \equiv 0$ and both sides erase the body: the left side is $R(0, \nu(a[\sigma]); \vec{u}.0)$, the right side is $R(0, \nu(\nu(a)[\nu \circ \sigma]); \vec{u}.0)$, and the induction hypothesis for a applies.

If $n \not\equiv 0$ and $n[\sigma] \equiv 0$, then n is a variable x with $\sigma(x) \equiv 0$, because a substitution instance of a recursor node is a recursor node. The left side erases the body. On the right, $\nu(T) = R(x, \nu a; \vec{u}.\nu h)$, and after the substitution $\nu \circ \sigma$ the counter is $\nu(\sigma(x)) \equiv 0$ by (9.5), so the outer ν erases the body as well. The initializers agree by the induction hypothesis.

If $n[\sigma] \not\equiv 0$, then $n \not\equiv 0$, and by (9.5) neither $\nu(n[\sigma])$ nor $\nu(n)[\nu \circ \sigma]$ is literally 0. Both sides retain the node, and the induction hypotheses for the counter, the initializer and the body give the equality; for the body the substitution is extended by the identity on $\vec{u}$.

Formula constructors bind variables but introduce no further case. For the consequence, apply the identity twice: $\nu(\nu(T[\sigma])[\tau]) = \nu(T[\sigma][\tau])$ since τ is canonical, and $\nu(T[\sigma\tau]) = \nu(T[\sigma][\tau])$ since T is canonical and $\nu(\sigma(x)[\tau]) = (\sigma\tau)(x)$. □

A variable replaced by 0 may expose a zero counter. Normalizing only the original syntax, or testing the counter before substitution, is therefore not sufficient, and rejecting noncanonical input is not sufficient either: instances generated by a rule have to be normalized. In a display that belongs to a canonical calculus, an expression $R(n, a; \vec{u}.h)$ denotes its normalization; the convention matters only when $n \equiv 0$, where the written body is discarded.

Remark 9.4. Theorem 9.3 is the substitution-coherence lemma of `canonical_recursors` for the seeded recursor. `counter_evaluation` and `uniform_binding` state it for the counter-aware recursor in the form $\nu(t\sigma) = \nu(\nu(t)\,\nu(\sigma))$ with the same proof, and `clausal_canonical` states it for the opening operation of a de Bruijn representation (Theorem 9.81). The proof above covers both binding arities.

9.1.3 Full normal forms

The counter-aware language also admits a grammar in which both computation equations are built into the syntax.

Definition 9.5 (Normal terms)**.** In the counter-aware language, a *canonical successor* is a term $R(0, a; i, u.0)$ with a normal. A *neutral* term is a variable or a term $R(c, a; i, u.h)$ with c neutral and a, h normal. A term is *normal* if it is 0, a canonical successor or neutral. Thus a retained recursor node has a counter that is neither 0 nor a successor.

Every normal term has the form $S^k(c_0)$ with $c_0 \equiv 0$ or c_0 neutral; k is the length of its successor prefix. Every normal term is zero-body canonical.

Definition 9.6 (Normalization by evaluation)**.** An environment ρ maps variables to normal terms. Define $\mathrm{ev}(t, \rho)$ by recursion on the raw term t: $\mathrm{ev}(x, \rho) = \rho(x)$, $\mathrm{ev}(0, \rho) = 0$, and

$$\mathrm{ev}\big(R(n, a; i, u.h), \rho\big) = \mathrm{run}\big(\mathrm{ev}(n, \rho), \mathrm{ev}(a, \rho), H\big), \qquad H(d, w) = \mathrm{ev}\big(h, \rho[i \mapsto d, u \mapsto w]\big),$$

where, for normal c, v,

$$\mathrm{run}(c, v, H) = \begin{cases} Sv, & c \equiv 0, \\ H\big(d, \mathrm{run}(d, v, H)\big), & c \equiv Sd, \\ R(c, v; i, u.H(i, u)), & c \text{ neutral}, \end{cases} \tag{9.6}$$

with i, u fresh in the third clause. The notation $H(i, u)$ is evaluation of the body in the environment extended by the two fresh variables; it introduces no function object into the term language. Put $\mathrm{nf}(t) = \mathrm{ev}(t, \mathrm{id})$, and for formulas, clauses and substitutions normalize all terms.

Lemma 9.7 (Normalization and substitution)**.** *Evaluation is total on raw terms and normal environments, and its values are normal. Moreover:*

1. $\mathrm{nf}(t) = t$ *for every normal* t*; hence* nf *is idempotent.*
2. nf *respects* (9.3) *and* (9.4)*: both sides of every instance have the same normal form, and so do two terms that differ by an instance inside any context, including under a binder.*
3. *For every raw* t *and substitution* σ*,*

$$\mathrm{nf}(t[\sigma]) = \mathrm{nf}\big(\mathrm{nf}(t)[\mathrm{nf}\circ\sigma]\big). \tag{9.7}$$

The same statements hold for formulas and clauses; no formula binder is erased.

Proof. Totality. Induct on t. At a recursor node the values c, v are normal and the body evaluator H is total with normal values, by the induction hypothesis for the proper subterms. For fixed total H, induct on the successor prefix of c in (9.6): the first and third clauses return a normal term, and in the second clause the recursive call has a shorter prefix and H is total.

Item 1. Induct on a normal term. A canonical successor evaluates to $\mathrm{run}(0, a, H) = Sa$. A neutral $R(c, a; i, u.h)$ evaluates by the third clause to $R(c, a; i, u.H(i, u))$, and $H(i, u) = h$ by the induction hypothesis.

Two environment laws. For a substitution σ and environments ρ, τ write $(\rho \circ \sigma)(x) = \mathrm{ev}(\sigma(x), \rho)$ and $(\tau \bullet \rho)(x) = \mathrm{ev}(\rho(x), \tau)$. Then

$$\mathrm{ev}(t[\sigma], \rho) = \mathrm{ev}(t, \rho \circ \sigma), \tag{9.8}$$

$$\mathrm{ev}\big(\mathrm{ev}(t, \rho), \tau\big) = \mathrm{ev}(t, \tau \bullet \rho). \tag{9.9}$$

Both are proved by induction on t with bound names fresh. For (9.8) the recursor case holds because the body evaluator of $h[\sigma]$ in ρ is the body evaluator of h in $\rho \circ \sigma$, by the induction hypothesis. For (9.9) let H be the body evaluator of h in ρ and H^τ the one in $\tau \bullet \rho$; the induction hypothesis for h gives $\mathrm{ev}(H(d, w), \tau) = H^\tau(\mathrm{ev}(d, \tau), \mathrm{ev}(w, \tau))$. It suffices to show

$$\mathrm{ev}\big(\mathrm{run}(c, v, H), \tau\big) = \mathrm{run}\big(\mathrm{ev}(c, \tau), \mathrm{ev}(v, \tau), H^\tau\big)$$

by induction on the successor prefix of c. If $c \equiv 0$ both sides are $S(\mathrm{ev}(v, \tau))$. If $c \equiv Sd$ the left side is $H^\tau(\mathrm{ev}(d, \tau), \mathrm{ev}(\mathrm{run}(d, v, H), \tau))$, which by the hypothesis for d is the second clause of (9.6) at the counter $S(\mathrm{ev}(d, \tau)) = \mathrm{ev}(Sd, \tau)$. If c is neutral, the left side evaluates the residual node $R(c, v; i, u.H(i, u))$ in τ; its body evaluator is $(d, w) \mapsto \mathrm{ev}(H(i, u), \tau[i \mapsto d, u \mapsto w]) = H^\tau(d, w)$, so the left side is $\mathrm{run}(\mathrm{ev}(c, \tau), \mathrm{ev}(v, \tau), H^\tau)$ whichever clause applies to $\mathrm{ev}(c, \tau)$. This is the case in which a substitution exposes a new redex.

Item 2. In every environment, $\mathrm{ev}(R(0, a; i, u.h), \rho) = S(\mathrm{ev}(a, \rho))$ independently of h, and $\mathrm{ev}(R(Sn, a; i, u.h), \rho) = H(c, \mathrm{run}(c, v, H))$ with $c = \mathrm{ev}(n, \rho)$, which by (9.8) is the value of $h[n/i, R(n, a; i, u.h)/u]$ in ρ. Evaluation is compositional: the value of a term in all environments is determined by the values of its immediate subterms in all environments. Hence replacing a subterm by one with the same values does not change the value.

Item 3. By (9.8), $\mathrm{nf}(t[\sigma]) = \mathrm{ev}(t, \mathrm{nf}\circ\sigma)$. By (9.8), item 1 and (9.9), $\mathrm{nf}(\mathrm{nf}(t)[\mathrm{nf}\circ\sigma]) = \mathrm{ev}(\mathrm{ev}(t, \mathrm{id}), \mathrm{nf}\circ\sigma) = \mathrm{ev}(t, \mathrm{nf}\circ\sigma)$. □

Remark 9.8. `single_constructor` defines the normalizer by evaluation, as above. `uniform_recursion` obtains the same normal forms from rewriting: it reads $R(n, a; i, u.h)$ as the base-type recursor of Gödel's system T with initial value Sa and update $h(i, u)$, derives strong normalization from the reducibility argument for that recursor (a numerical expression is computable when it is strongly normalizing, an update is computable when it maps computable arguments to

computable expressions, and the recursor case is an induction on the successor spine of a normal counter), observes that the zero and successor rules have disjoint head patterns and commute with reductions in their arguments and with erasure of an unused body, and concludes unique normal forms from termination and local confluence. The two definitions agree, since by item 2 of Theorem 9.7 every reduction step preserves nf and by item 1 the normal terms are exactly the irreducible ones.

Remark 9.9 (No uniform primitive recursive normalizer). Each fixed term denotes a primitive recursive function of its parameters. It does not follow that normalization of arbitrary term descriptions is primitive recursive uniformly; a universal evaluator of that kind leads to the usual diagonal contradiction. The normalizer is an external total recursive operation on finite syntax, and no assertion that all descriptions normalize is an axiom of any theory here. `uniform_recursion` supplies in addition a checker with an explicit budget. Bounded simulation gives a primitive recursive certificate predicate with the same set of provable formulas, since every finite successful run has a sufficient budget.

9.1.4 The status of the successor and of each computation

The languages of this chapter differ from the iterator terms of Section 1.1, and from one another, in which of three items is a constructor or a rule and which is carried by the syntax. The following table summarizes the presentations treated below.

Terms	Successor	Zero computation	Successor computation
$0, S, I$ (Section 1.1)	constructor	rule	rule
raw seeded R, either arity	$R(0, a; \vec{u}.0)$	rule	rule
zero-body canonical R, one binder (Sections 9.4 and 9.5)	$R(0, a; u.0)$	identity of spelling	the one computation rule
zero-body canonical R, two binders (Section 9.6)	$R(0, a; i, u.0)$	identity of spelling	derived from reflexivity and counter evaluation
normal R, two binders (Section 9.7)	$R(0, a; i, u.0)$	identity of spelling	identity of spelling
register J (Section 9.2)	$J(a; i, r.0)$	axiom $J(0; i, r.h) = 1$	graph biconditional

A computation that is an identity of spelling has not left the trusted specification: the normalizer is applied at every substitution and comparison, and it is counted in every checker of Section 9.10. Its soundness in ordinary arithmetic is Theorem 9.23; it rests on an extensionality property of recursor bodies (Theorem 9.22) and not on syntactic idempotence alone.

9.2 What the counter gives

This section draws on `counter_evaluation`, `uniform_binding`, `uniform_recursion`, `single_constructor`, `register_graph` and, for the contrast with a body that does not see its counter, `canonical_recursors`.

9.2.1 A decoder that computes

In the counter-aware language define, with all auxiliary bound variables fresh,

$$\begin{aligned} P(n) &:= R(n, 0; i, u.i), \\ C(n, a, b) &:= P\big(R(n, a; i, u.Sb)\big), \\ \mathrm{Rec}(n, a; i, u.h) &:= P\big(R(n, a; i, v.S(h[P(v)/u]))\big), \\ p(n) &:= C(n, 0, P(n)), \qquad \mathrm{sg}(n) := C(n, 0, 1). \end{aligned} \tag{9.10}$$

Lemma 9.10 (Decoder, selector and recursion)**.** *The following equations are derivable from* (9.3)*,* (9.4)*, reflexivity and congruence, for arbitrary terms and passive parameters:*

$$\begin{aligned} P(0) &= 1, & P(Sn) &= n,\\ C(0,a,b) &= a, & C(Sn,a,b) &= b,\\ \mathrm{Rec}(0,a;i,u.h) &= a, & \mathrm{Rec}(Sn,a;i,u.h) &= h[n/i,\mathrm{Rec}(n,a;i,u.h)/u],\\ p(0) &= 0, & p(Sn) &= n,\\ \mathrm{sg}(0) &= 0, & \mathrm{sg}(Sn) &= 1. \end{aligned}$$

No induction, numerical case distinction or logical rule is used. On normal terms each equation is a literal identity of normal forms.

Proof. The equations for P are the two computations with the body that returns the old counter and ignores the state. In $C(0,a,b)$ the inner recursor is Sa and in $C(Sn,a,b)$ it is Sb; then apply $P(Sx)=x$. For recursion put $T(n)=R(n,a;i,v.S(h[P(v)/u]))$. Then $T(0)=Sa$ and $T(Sn)=S(h[n/i,P(T(n))/u])$, and applying P gives both equations. The equations for p and sg are instances of those for C and P. On normal terms each step is a computation, so both sides have the same normal form by Theorem 9.7. □

The decoder P is a left inverse of the successor obtained by one computation step, for an arbitrary open argument. Its value at 0 is 1, not 0; the predecessor of Theorem 1.48 is p. The state of Rec is stored as a successor and the outer P removes that wrapper after each observation, so the successor seed of R does not restrict the initial value.

Remark 9.11. `counter_evaluation` and `uniform_binding` use the names of (9.10). `uniform_recursion` writes W for the decoder, P for the predecessor p and J for Rec, and uses the term $K(y) := C(W(y),1,0)$, with $K(1)=1$ and $K(S1)=0$. `single_constructor` defines the predecessor and the selector from the recursion macro, as $p(t) := \mathrm{Rec}(t,0;i,v.i)$ and $C(t,a,b) := \mathrm{Rec}(t,a;i,v.b)$; they satisfy the same equations.

9.2.2 Primitive recursion without pairing

Theorem 9.12 (Direct compiler)**.** *Every fixed primitive recursive description compiles to a term of the counter-aware language whose defining equations are derivable from the computation equations and equality reasoning alone. Zero, successor and projections are terms, composition is substitution, and a recursion $f(0,\vec{p})=g(\vec{p})$, $f(Sn,\vec{p})=h(n,f(n,\vec{p}),\vec{p})$ compiles to*

$$f^*(n,\vec{p}) = \mathrm{Rec}\big(n,g^*(\vec{p});i,u.h^*(i,u,\vec{p})\big).$$

The compiler uses no pairing, no projection functions, no induction and no coverage. The iterator of Section 1.1 is the instance $I(n,a;\,u.\,h) := \mathrm{Rec}(n,a;i,u.h)$ with $i \notin \mathrm{FV}(h)$.

Proof. By external induction on the description. The recursion case is Theorem 9.10; the other cases are immediate. □

For example $x+y = \mathrm{Rec}(y,x;i,v.Sv)$, $x\cdot y = \mathrm{Rec}(y,0;i,v.v+x)$ and the truncated subtraction is $d(x,y) = \mathrm{Rec}(y,x;i,v.p(v))$. The compiler is an external recursion on each finite description. It is not a universal evaluator for description codes inside the theory. The compiler of Theorem 1.62 for iterator terms stores the counter beside the state by a pairing function and therefore comes after the construction of pairing; here the counter is an argument of the body.

9.2.3 The register iterator

The register iterator carries the same information in a different form: the body returns an instruction, and the axioms describe the transition graph.

Definition 9.13 (The theory T_J)**.** The language has the terms of Theorem 9.1(3) and the formulas

$$A ::= (s = t) \mid \bot \mid (A \wedge A) \mid (A \vee A) \mid (A \to A) \mid \forall x\, A \mid \exists x\, A.$$

The logical rules are those of intuitionistic natural deduction with equality: hypothesis, reflexivity, transport of an arbitrary formula along a proved equation, ex falso, and the introduction and elimination rules of $\wedge, \vee, \to, \forall, \exists$ with the usual eigenvariable conditions. There is no classical rule. For a fixed body $h(i, r, \vec{p})$ write $j_n = J(n; i, r.h)$ and $e_n = h(n, j_n, \vec{p})$. The arithmetic axioms are the universal closures of

$$J(0; i, r.h) = 1, \tag{J0}$$

$$J(Sn; i, r.h) = b \;\leftrightarrow\; \big((e_n = 0 \wedge b = Sj_n) \vee e_n = Sb\big), \tag{STEP}$$

$$0 = 1 \to \bot, \tag{Sep}$$

where b is a variable distinct from the passive parameters. The one arithmetic rule is induction for literal equations with an arbitrary context:

$$\frac{\Gamma \vdash s(0) = t(0) \qquad \Gamma,\ s(z) = t(z) \vdash s(Sz) = t(Sz)}{\Gamma \vdash s(a) = t(a)} \qquad (z \text{ fresh for } \Gamma, a \text{ and the parameters}). \tag{9.11}$$

An equation under a quantifier or a connective is not eligible as an invariant.

The right-to-left direction of (STEP) gives two conditional computation rules, whose premises are proved equations and not tests performed by an evaluator:

$$e_n = 0 \implies J(Sn; i, r.h) = Sj_n, \tag{9.12}$$

$$e_n = St \implies J(Sn; i, r.h) = t. \tag{9.13}$$

Lemma 9.14 (Functional coverage)**.** *Let $f(n, a) := J(n; i, r.a)$ with i, r fresh for a, and $W(a) := f(1, a)$. Without induction, $\mathsf{T}_J \vdash a = 0 \vee a = S(W(a))$. In particular $\forall a\,(a = 0 \vee \exists x\, a = Sx)$ is derivable.*

Proof. Take the constant body a, the counter $n = 0$ and $b = f(1, a)$ in (STEP). Its left side holds by reflexivity. Its right side is $(a = 0 \wedge b = S1) \vee a = Sb$; drop the second conjunct of the first alternative. □

Proposition 9.15 (The graph axiom)**.** *Over intuitionistic logic with equality, and without* (Sep)*, induction or injectivity of the successor:*

1. (J0) *and* (STEP) *are equivalent to* (J0)*,* (9.12)*,* (9.13) *and numerical coverage $\forall a\,(a = 0 \vee \exists x\, a = Sx)$;*
2. *both are equivalent to the single schema*

$$J(n; i, r.h) = b \;\leftrightarrow\; (n = 0 \wedge b = 1) \vee \exists m\,\big(n = Sm \wedge ((e_m = 0 \wedge b = Sj_m) \vee e_m = Sb)\big). \tag{9.14}$$

Proof. (1) From (STEP) the two conditional rules follow by its right-to-left direction and coverage is Theorem 9.14. Conversely the right-to-left direction of (STEP) is the two conditional rules. For the left-to-right direction assume $J(Sn; i, r.h) = b$ and split coverage for e_n. If $e_n = 0$, then (9.12) gives $b = Sj_n$. If $e_n = St$, then (9.13) gives $b = t$ and hence $e_n = Sb$.

(2) The right-to-left direction of (9.14) gives (J0) at $n = 0$, $b = 1$ and both conditional rules at $n = Sm$. Its left-to-right direction applied to the reflexive equation $J(n; i, r.0) = J(n; i, r.0)$ gives $n = 0 \vee \exists m\, n = Sm$. By (1) this proves (STEP). Conversely, assume (J0) and (STEP). The right-to-left direction of (9.14) is (J0) and the right-to-left direction of (STEP) after transport along $n = Sm$. For the left-to-right direction split coverage for n: at $n = 0$ use (J0), and at $n = Sm$ use (STEP). □

The biconditional is essential: the left-to-right direction of (STEP) alone does not assert that the computation is deterministic. A packaging such as (9.14) is also available for the iterator of Section 1.1, so a smaller number of named schemas is not by itself a simplification.

Lemma 9.16 (Bootstrap without induction)**.** *Using only equality,* (J0) *and the conditional rules,* T_J *proves*

$$f(0, a) = 1, \qquad f(n, 0) = Sn, \qquad f(Sn, Sa) = a, \tag{9.15}$$

and for

$$Z(x) := f(x, 1), \quad W(x) := f(1, x), \quad D(x) := f(x, Sx), \quad P(x) := W(D(x)),$$

$$\begin{aligned} Z(0) &= 1, & Z(Sx) &= 0, \\ W(0) &= S1, & W(Sx) &= x, \\ D(0) &= 1, & D(Sx) &= Sx, \\ P(0) &= 0, & P(Sx) &= x. \end{aligned}$$

Further, for

$$G(c, a) := P^2 f(Sa, Z(c)), \qquad C(c, a, b) := P^2 f\big(Sa,\, G(c, S^3 b)\big),$$

where P^2 *and* S^3 *are finite compositions,*

$$G(0, a) = 0, \quad G(Sn, a) = a, \qquad C(0, a, b) = a, \quad C(Sn, a, b) = b. \tag{9.16}$$

Consequently the successor is injective, and with (Sep) *no successor is* 0*.*

Proof. The first equation of (9.15) is (J0), the second is the definition of S, and the third is a load step whose premise is the reflexive equation $Sa = Sa$. The eight endpoint equations are instances: for example $Z(Sx) = f(Sx, S0) = 0$, $W(0) = f(1, 0) = S1$, $P(0) = W(1) = W(S0) = 0$ and $P(Sx) = W(Sx) = x$. For G: at $c = 0$ the inner term is $f(Sa, S0) = 0$ and $P^2(0) = 0$; at $c = Sn$ it is $f(Sa, 0) = S^2 a$ and P^2 returns a. For C: at $c = 0$ the instruction $G(c, S^3 b)$ is 0, so the inner term is $S^2 a$; at $c = Sn$ the instruction is $S^3 b$, and the counter Sa, which is visibly a successor, loads $S^2 b$. Applying P to $Sx = Sy$ gives $x = y$. An equation $Sx = 0$ gives $Z(Sx) = Z(0)$, that is $0 = 1$. □

The selector C returns arbitrary numerical outputs. It is not limited to Boolean results or to values bounded by a preselected initializer; the direct use of the old register in the increment branch is what makes this possible.

Theorem 9.17 (Register compiler)**.** *For terms* a *and* $h(u, v)$ *of the register language, with* i, r *fresh, put*

$$H(i, r) := S\big(C(i, a, h(P(i), r))\big), \qquad \mathrm{Rec}_J(n, a; u, v.h) := J(Sn; i, r.H(i, r)).$$

Without induction, coverage, pairing or reflection of equality,

$$\mathrm{Rec}_J(0, a; u, v.h) = a, \qquad \mathrm{Rec}_J(Sn, a; u, v.h) = h\big(n, \mathrm{Rec}_J(n, a; u, v.h)\big).$$

Hence every fixed primitive recursive description compiles to a term of the register language with derivable defining equations.

Proof. The instruction is always a successor, so every update is a load. The first update has counter $i = 0$ and old register 1; by (9.16) its instruction is Sa and (9.13) gives $J(1; i, r.H) = a$. For the second equation put $r_0 = J(Sn; i, r.H)$. The old counter is Sn, so (9.16) and $P(Sn) = n$ give $H(Sn, r_0) = S(h(n, r_0))$, and one load gives $J(S(Sn); i, r.H) = h(n, r_0)$. Initial functions and composition translate by substitution. □

The compact source grammar does not guarantee small expanded terms. The compiler shipped with `register_graph` uses the instruction $S(C(Z(i), h(P(i), r), a))$, which has the same endpoint values and a smaller expansion.

9.2.4 A register with one bound variable

Proposition 9.18 (One-variable register)**.** *Let $t ::= x \mid 0 \mid J_s(n; r.h)$ with $Sa := J_s(a; r.0)$, and let the axioms be* (J0)*,* (STEP) *with $e_n = h(j_n, \vec{p})$,* (Sep) *and* (9.11)*. Then Theorems 9.14 and 9.16 hold unchanged, and for*

$$H(v) := S^3\big(C(P(v), a, h(P^2 v))\big), \qquad K(n) := J_s(Sn; v.H(v)), \qquad \widehat{I}(n, a; r.h) := P^2 K(n)$$

the equations $\widehat{I}(0, a; r.h) = a$ and $\widehat{I}(Sn, a; r.h) = h(\widehat{I}(n, a; r.h))$ are derivable without induction or coverage.

Proof. The bootstrap uses constant bodies only. One load step gives, at every counter,

$$J_s(Sn; v.H) = S^2\big(C(P(J_s(n; v.H)), a, h(P^2 J_s(n; v.H)))\big),$$

so every state after the initial one is visibly a double successor. At the first step the previous register is 1 and $P(1) = 0$, so $K(0) = S^2 a$. At a later step $P(K(n))$ is a successor and the selector takes the update branch, so $K(Sn) = S^2 h(P^2 K(n))$. Applying P twice gives both equations. The shape of the state is a consequence of one computation, not an inferred invariant. □

The one-variable register keeps the binding arity of the iterator and still has neither a successor constructor nor an initializer. Its simulation of general primitive recursion passes through the iterator and therefore through the pairing of Theorem 1.62. The counter variable of the two-variable register is the cost of the direct compiler of Theorem 9.17.

Proposition 9.19 (Reflexivity from conditional computation)**.** *Consider* (J0)*,* (9.12) *and* (9.13) *as rules whose premises are derivations, together with transport of an equation along an equation. Then $t = t$ is derivable for every term t, without a primitive reflexivity rule.*

Proof. (i) (J0) for the zero body is literally $1 = 1$. (ii) A load step with the constant instruction $1 = S0$, whose premise is (i), gives $J(1; i, r.1) = 0$; transporting this equation along itself gives $0 = 0$. (iii) An increment step with the zero body at the counter t, whose premise is (ii), gives the literal equation $S(St) = S(St)$, since $J(t; i, r.0)$ is St by definition. (iv) A load of the constant instruction $S(St)$, whose premise is (iii), gives an equation with right side St, and transport along itself gives $St = St$. (v) A load of the constant instruction St gives an equation with right side t, and transport along itself gives $t = t$. □

By Theorem 9.19 the register terms are compatible with a native kernel in the style of Section 1.3, in which a conditional computation step takes a proved equation as premise. No checker for such a kernel is supplied (Section 9.11).

9.2.5 Without a counter

The seeded recursor with one bound variable has the partial selector $B(n,a,b) := R(n,a;u.b)$, $u \notin \mathrm{FV}(b)$, with $B(0,a,b) = Sa$ and $B(Sn,a,b) = b$, but its body has no access to the counter, and the decoder of Theorem 9.10 is not available. A predecessor exists nevertheless: by Theorem 8.19 there is a fixed term P of the seeded language with

$$P(0) = 0, \qquad P(Sa) = a \tag{9.17}$$

derivable in the native calculus and in the ordinary theory of the seeded recursor, by native numerical cases, equality reasoning and induction on literal equations. By Theorem 8.21 the term

$$\widehat{I}(n,a;u.h) := P\big(R(n,a;v.S(h[P(v)/u]))\big)$$

then satisfies the two equations of the iterator, and $C(n,a,b) := \widehat{I}(n,a;u.b)$ is a selector with $C(0,a,b) = a$ and $C(Sn,a,b) = b$.

The difference between the two arities is therefore the length of the bootstrap and the need for pairing, not expressive strength: with the counter the predecessor is one computation step and primitive recursion is Theorem 9.12; without it the predecessor is the construction of Section 8.1 and primitive recursion is compiled as in Theorem 1.62. If the independent initializer is removed as well and no equality test is added, no term denotes the predecessor (Theorem 9.85). The witness-free introductions that use the computing predecessor of the counter-aware recursor are treated in Sections 7.7 and 7.8, on the normal terms of Theorem 9.5.

9.3 Translations to iterator terms and the transfer of the bridge

This section draws on `register_graph`, `canonical_recursors`, `clausal_canonical`, `counter_evaluation`, `uniform_binding`, `uniform_recursion` and `single_constructor`: each of them proves, for its own syntax, a bi-interpretation with iPRA. Here the statement is proved once, relative to T_{eq}, and Theorem 1.78 is then applied.

9.3.1 The ordinary theories

Definition 9.20 (Ordinary theories)**.** The theories T_R^1 and T_R^2 have the raw terms of the seeded recursor with one and with two bound variables, the formulas and logical rules of Theorem 9.13, and the following arithmetic part: the two computation equations ((9.1), (9.2), respectively (9.3), (9.4)); coverage $\forall x\,(x = 0 \vee \exists y\, x = Sy)$; separation $0 = S0 \to \bot$; and the rule (9.11) of induction for literal equations with an arbitrary context. The theory T_J is that of Theorem 9.13; coverage is a theorem of it (Theorem 9.14). The theory T_R^1 is the theory T_R of Theorem 8.1.

These are the theory T_{eq} of Section 1.5 with the terms exchanged. The separation axiom is part of each theory: the positive formulas read in isolation admit a one-element structure. Since a fixed body is a function symbol of its parameters, ordinary equality logic gives congruence in the parameters of a body. Replacing a body by a different body that is provably equal to it pointwise is not an instance of congruence; it is the next lemma.

Lemma 9.21 (Recurrence uniqueness)**.** *Let* T *be* T_{eq}, T_R^1 *or* T_R^2, *and let* $F(z)$, $G(z)$, $k(z,w)$, $k'(z,w)$ *be terms with* z, w *fresh for* Γ. *If*

$$\Gamma \vdash F(0) = G(0), \quad \Gamma \vdash F(Sz) = k(z,F(z)), \quad \Gamma \vdash G(Sz) = k'(z,G(z)), \quad \Gamma \vdash k(z,w) = k'(z,w),$$

then $\Gamma \vdash F(t) = G(t)$ *for every term* t.

Proof. Apply (9.11) to the literal equation $F(z) = G(z)$, with all parameters fixed. The base is given. Under $F(z) = G(z)$,

$$F(Sz) = k(z, F(z)) = k(z, G(z)) = k'(z, G(z)) = G(Sz),$$

by congruence in the argument w of k and the instance $w := G(z)$ of the fourth hypothesis. □

Lemma 9.22 (Body extensionality)**.** *In* T^1_R *and* T^2_R*, if* $\Gamma \vdash a = b$ *and* $\Gamma \vdash h = k$ *with the bound variables* $\vec{u}$ *free in this premise and fresh for* Γ*, then* $\Gamma \vdash R(n, a; \vec{u}.h) = R(n, b; \vec{u}.k)$ *for every* n*.*

Proof. For one bound variable this is Theorem 8.3. In general take $F(z) = R(z, a; \vec{u}.h)$ and $G(z) = R(z, b; \vec{u}.k)$ in Theorem 9.21. At 0 the two sides are Sa and Sb, equal by congruence. The successor equations are the computation equations, and the updates agree by the second premise. □

Lemma 9.23 (Normalization is sound)**.** T^1_R *and* T^2_R *prove* $t = \nu(t)$ *for every raw term* t*, and* T^2_R *proves* $t = \mathrm{nf}(t)$*. Hence every formula is provably equivalent to its normalization, and a raw substitution instance is provably equivalent to the normalized one.*

Proof. For ν induct on t. The counter and the initializer change by congruence, the body by Theorem 9.22 and the induction hypothesis with the bound variables free. If the counter is literally 0, the zero computation, applied to the term and to its normalization, identifies both with Sa.

For nf prove $\mathsf{T}^2_R \vdash t[\rho] = \mathrm{ev}(t, \rho)$ for every normal environment ρ, by induction on t. At a recursor node the induction hypotheses give $n[\rho] = c$, $a[\rho] = v$ and, at the fresh variables, $h[\rho] = H(i, u)$; hence $R(n, a; i, u.h)[\rho] = R(c, v; i, u.H(i, u))$ by Theorem 9.22. It remains to prove $R(c, v; i, u.H(i, u)) = \mathrm{run}(c, v, H)$ by induction on the successor prefix of c. If $c \equiv 0$ this is the zero computation. If c is neutral the two sides are identical. If $c \equiv Sd$, the successor computation gives $H(i, u)[d/i, R(d, v; i, u.H(i, u))/u]$; replace the state by $\mathrm{run}(d, v, H) =: w$ using the hypothesis for d, and observe that $H(i, u)[d/i, w/u] = h[\rho][d/i, w/u] = H(d, w)$ by the induction hypothesis for h in the two environments. The statements on formulas follow by transport of formulas along equations. □

9.3.2 The translations

Definition 9.24 (Translations)**.** Let T be one of T^1_R, T^2_R, T_J. The translation $t \mapsto t^\iota$ from T to T_{eq} fixes variables and 0 and is defined on the binding constructor as follows.

1. For T^1_R: $R(n, a; u.h)^\iota = I(n^\iota, Sa^\iota;\, u.\, h^\iota)$, as in Theorem 8.23.
2. For T^2_R: $R(n, a; i, u.h)^\iota = F(n^\iota)$, where F is the term given by the compiler of Theorem 1.62 for the recursion $F(0) = Sa^\iota$, $F(Sz) = h^\iota[z/i, F(z)/u]$.
3. For T_J: $J(n; i, r.h)^\iota = F(n^\iota)$, where F is given by the same compiler for the recursion

$$F(0) = S0, \qquad F(Sz) = \mathrm{sel}(e,\ SF(z),\ \mathrm{pd}(e)), \qquad e = h^\iota[z/i, F(z)/r],$$

and pd is the predecessor term of Theorem 1.48, written P there.

The compiler of Theorem 1.62 provides, for terms a and $g(y, v)$ of T_{eq}, a defined term with the value a at 0 and the value $g(c, \cdot)$ of the preceding value at Sc; it is built with pairing. That derived recursor of T_{eq} is not the constructor R of this chapter, whose value at 0 is Sa.

The translation $s \mapsto s^\kappa$ from T_{eq} to T fixes variables and 0, sends Ss to the defined successor of s^κ, and sends the iterator to the initialized iteration of T:

$$I(n,a;\,u.\,h)^\kappa = \begin{cases} \widehat{I}(n^\kappa, a^\kappa; u.h^\kappa) & \text{in } \mathsf{T}^1_R \text{ (Theorem 8.21; the translation } \rho \text{ of Theorem 8.23)},\\ \mathrm{Rec}(n^\kappa, a^\kappa; i, u.h^\kappa),\ i \text{ fresh}, & \text{in } \mathsf{T}^2_R \text{ (Theorem 9.10)},\\ \mathrm{Rec}_J(n^\kappa, a^\kappa; i, u.h^\kappa),\ i \text{ fresh}, & \text{in } \mathsf{T}_J \text{ (Theorem 9.17)}. \end{cases}$$

Both translations commute with substitution and are extended to formulas so that they commute with every connective and quantifier.

Theorem 9.25 (Transfer of the bridge)**.** *Let* T *be* T^1_R, T^2_R *or* T_J.

1. *If* $\mathsf{T} \vdash A$ *then* $\mathsf{T}_{\mathrm{eq}} \vdash A^\iota$, *and if* $\mathsf{T}_{\mathrm{eq}} \vdash B$ *then* $\mathsf{T} \vdash B^\kappa$.
2. $\mathsf{T} \vdash t^{\iota\kappa} = t$ *for every term* t *of* T, *and* $\mathsf{T}_{\mathrm{eq}} \vdash s^{\kappa\iota} = s$ *for every term* s *of* T_{eq}.
3. *Consequently* $\mathsf{T} \vdash A^{\iota\kappa} \leftrightarrow A$ *and* $\mathsf{T}_{\mathrm{eq}} \vdash B^{\kappa\iota} \leftrightarrow B$; *both translations are faithful; and* T *is bi-interpretable with* T_{eq}, *hence by Theorem 1.78 with* N *and with* iPRA, *by interpretations whose domains are the whole numerical domain, whose equality is actual equality and whose numerical composites have the graph* $x = y$. *The four theories are equiconsistent.*

The same holds for the one-variable register of Theorem 9.18, with $J_s(n; r.h)^\iota = I(n^\iota, S0;\, v.\, \mathrm{sel}(h^\iota(v), Sv, \mathrm{pd}(h^\iota$ *and the iteration* $\widehat{I}$ *of that proposition for* κ.

Proof. For T^1_R the three statements are Theorems 8.24 and 8.25. The proof is written for T^2_R and T_J.

(1) *From* T *to* T_{eq}. The logical rules translate to themselves. The translation of the defined successor satisfies

$$\mathsf{T}_{\mathrm{eq}} \vdash (St)^\iota = S(t^\iota). \tag{9.18}$$

For the seeded recursor this is the base equation of the translated recursion. For the register, $(St)^\iota = F_0(t^\iota)$, where F_0 translates the zero body and satisfies $F_0(0) = S0$ and $F_0(Sz) = \mathrm{sel}(0, SF_0(z), \mathrm{pd}(0)) = SF_0(z)$; induction on the literal equation $F_0(z) = Sz$ gives (9.18). Thus the defined successor is interpreted as the successor of T_{eq} at every element of the domain, not only at numerals. With (9.18) and transport, the computation equations of the seeded recursor translate to the defining equations of the translated recursion, coverage and separation translate to the corresponding axioms of T_{eq}, and an instance of (9.11) translates to an instance of the induction rule of T_{eq} for a literal equation. For T_J, (J0) is the base equation. For (STEP) write $j = F(n)$ and e for the translated instruction, so that $F(Sn) = \mathrm{sel}(e, Sj, \mathrm{pd}(e))$, and split coverage for e. If $e = 0$ then $F(Sn) = Sj$, and both sides of the biconditional are equivalent to $b = Sj$, the second alternative being excluded by separation. If $e = Sy$ then $F(Sn) = y$, and both sides are equivalent to $b = y$, by injectivity of the successor. (Sep) is separation.

From T_{eq} *to* T. The two computation equations of the iterator translate to the equations of Rec and Rec_J, which are derivable by Theorems 9.10 and 9.17. Coverage and separation are axioms or theorems of T, and the induction rule for literal equations is (9.11).

(2) Induct externally on the term. Let $t = R(n, a; i, u.h)$ in T^2_R and let $F(z)$ be the κ-translation of the compiled recursion of Theorem 9.24(2). By part (1), T^2_R proves $F(0) = Sa^{\iota\kappa}$ and $F(Sz) = h^{\iota\kappa}[z/i, F(z)/u]$. By the induction hypotheses, $a^{\iota\kappa} = a$ and $h^{\iota\kappa} = h$ with i, u free. The term $G(z) = R(z, a; i, u.h)$ has the same base and the same update, so Theorem 9.21 gives $F(z) = G(z)$, and the instance at $n^{\iota\kappa} = n$ is the claim. In the other direction, for $s = I(n, a;\, u.\, h)$ the term $s^{\kappa\iota}$ is $F(n^{\kappa\iota})$ for a term F of T_{eq} that satisfies the two equations of

the iterator with initial value $a^{\kappa\iota}$ and update $h^{\kappa\iota}$, by part (1); Theorem 9.21 in T_{eq} and the induction hypotheses give $s^{\kappa\iota} = s$. For $s = Ss_0$ use (9.18).

For $t = J(n; i, r.h)$ in T_J the double translation is $F(n^{\iota\kappa})$ with $F(0) = 1$ and $F(Sz) = C'(e', SF(z), P'(e'))$, where $e' = h^{\iota\kappa}[z/i, F(z)/r]$ and C', P' are the translated selector and predecessor, which have their endpoint equations in T_J. Apply (9.11) to the equation $F(z) = J(z; i, r.h)$. The base is (J0). Under the induction equation and the induction hypothesis for h, the two instructions e' and e_z are equal. Split coverage for this common value. In the zero branch both sides increment equal registers, by (9.12); in the successor branch both load the same number, by (9.13). The invariant is a single equation, and the case split is supplied by coverage inside the step.

(3) Since the translations commute with the logical operations, part (2) and transport give the two formula equivalences by induction on formulas. Faithfulness follows: if $\mathsf{T}_{\mathrm{eq}} \vdash A^{\iota}$ then $\mathsf{T} \vdash A^{\iota\kappa}$ and hence $\mathsf{T} \vdash A$. Both translations fix the domain, equality and $\bot$, so the identity graph is a provably total bijection that preserves every operation in both composites. The one-variable register is treated in the same way: coverage validates its graph axiom for the displayed iterator, induction on a literal equation shows that its successor abbreviation agrees with the successor, and recurrence uniqueness gives both composites. □

Remark 9.26 (The direct interpretations of the packages). Each package states the theorem against iPRA directly, with the same two maps: a recursor or register description is interpreted by the primitive recursive function symbol with the same base and step, and a function symbol of iPRA is interpreted by the compiler of Theorem 9.12 or Theorem 9.17, without pairing. By Theorem 9.21 these maps agree provably with the composites of ι and κ with the interpretations of Theorem 1.78. `canonical_recursors` and `clausal_canonical` use the first clause of Theorem 9.24 and the iteration $\widehat{I}$, and compile primitive recursion with pairing as in Theorem 1.62. The mere agreement of two theories on standard numerals, or mutual interpretability, would not give Theorem 9.25; the composites are proved inside the theories by Theorem 9.21.

Proposition 9.27 (Quantifier-free induction). *Each of T^1_R, T^2_R and T_J proves the successor to be injective, proves equality decidable, and admits induction for all quantifier-free formulas with an arbitrary context.*

Proof. Let $A(z)$ be quantifier-free in the language of T. By Theorem 1.56, T_{eq} proves the induction principle for the quantifier-free formula A^{ι} as a sentence, with its parameters universally quantified. By Theorem 9.25(1) the theory T proves the κ-translation of this sentence, in which the successor is the defined successor, and by Theorem 9.25(3) the formula $A^{\iota\kappa}$ may be replaced by A. A sentence of this form yields the rule with an arbitrary context. Injectivity and decidability of equality transfer in the same way from Theorems 1.48 and 1.52. □

Remark 9.28 (The reflection argument in the packages). `register_graph`, `counter_evaluation`, `uniform_binding`, `uniform_recursion` and `single_constructor` do not transfer Theorems 1.52 and 1.56; they repeat the proofs with the terms of Section 9.2, where the predecessor and the selector are available before any induction. With addition and truncated subtraction compiled by Theorem 9.12, four inductions on literal equations give $0 + y = y$, $Sx + y = S(x + y)$, $d(Sx, Sy) = d(x, y)$ and $d(Sa, a) = 1$; a fifth, on the single equation

$$C\big(\mathrm{sg}(d(x, y)),\, x,\, y + d(x, y)\big) = x$$

with x fixed, in whose step every case split is supplied by coverage, gives $d(Sa, b) = 1 \leftrightarrow a = b$. The reflection term is $\epsilon(a, b) = \mathrm{sg}(K(d(Sa, b)))$ with $K(n) = C(n, 1, p(n))$, and the characteristic term of a quantifier-free formula is defined with 0 for truth by $b_{s=t} = \epsilon(s, t)$, $b_{\bot} = 1$, $b_{A\wedge B} = C(b_A, b_B, 1)$, $b_{A\vee B} = C(b_A, 0, b_B)$, $b_{A\to B} = C(b_A, b_B, 0)$. `register_graph`

uses the control $d(x, y)$ in place of its sign, the term $e(a, b) = Z(W(d(Sa, b)))$ and characteristic terms with 1 for truth. An independent audit shipped with `uniform_binding` confirms that the only induction instances in this argument are literal equations. The argument is that of Theorems 1.52 and 1.56 and is not repeated.

9.4 The seeded recursor: unrestricted calculi and six rules

This section draws on `canonical_recursors`.

The native formulas are those of Section 1.3, $A ::= (s = t) \mid Q_y A \mid D_{x,y} A$, over the terms of the seeded recursor with one bound variable, with the reading of Section 1.2. Contexts are finite reusable lists of formulas, and the designated contradiction is $0 = S0$. The six-rule calculus has two tests on literal syntax, which are not stable under substitution. The comparison with the raw calculus is therefore established first for two unrestricted calculi, and the six rules are recovered afterwards.

9.4.1 The raw and the canonical unrestricted calculus

Definition 9.29. The *raw calculus* $\mathcal{R}_{\mathrm{raw}}$ uses raw terms and ordinary substitution. Its rules are:

1. Hyp: $\Gamma \vdash A$ if $A \in \Gamma$;
2. the two computation equations (9.1) and (9.2), in every context;
3. transport in every native formula template: from $s = t$ and $F[s/v]$ infer $F[t/v]$;
4. QE: from $Q_y B(y)$ and $B(0)$ infer $B(t)$;
5. DE: from $D_{x,y} B(x, y)$ and a derivation of A from $\Gamma, Q_y B(b, y)$ infer A, where b is fresh for Γ, A and the parameters of B;
6. introduction by cases: from $\Gamma, B(t, 0) \vdash B(t, Sz)$ infer $D_{x,y} B(x, y)$;
7. introduction by progress: from $\Gamma, E(t, z) \vdash E(t, Sz)$ infer $D_{x,y} E(x, y)$, for a literal equation E.

In the two introductions z is fresh for Γ, the witness t and the parameters of the matrix. The *canonical calculus* $\mathcal{C}_\nu$ has the same rules on zero-body canonical formulas with normalized substitution, except that its only computation schema is

$$R(Sn, a; u.h) = h\langle R(n, a; u.h)/u\rangle. \tag{R_+}$$

The left counter of (R_+) is never literally 0, so all displayed terms are canonical. The raw zero computation has become an identity of spelling. The raw calculus is the kernel N on the seeded terms in an equivalent rule set: transport of formulas is the Leibniz rule of Section 1.4, and progress concludes a paired formula instead of a closure.

Lemma 9.30 (Structural rules)**.** *Weakening, renaming of eigenvariables, cut of an ordinary assumption and capture-free numerical substitution are admissible in $\mathcal{R}_{\mathrm{raw}}$ and $\mathcal{C}_\nu$.*

Proof. Induct on derivations after freshening the local eigenparameters. For $\mathcal{C}_\nu$, Theorem 9.3 shows that normalized substitution maps every rule instance to a rule instance, including the instances of (R_+). The distinction between the two introductions depends only on the root of the matrix and is stable. □

Lemma 9.31 (Elementary resources)**.** *Both unrestricted calculi derive reflexivity, symmetry, transitivity and congruence of equality, introduction of Q_yB by cases, numerical cases, and induction on literal equations with an arbitrary context.*

Proof. Let $\mathrm{wr}(a) := R(S0, 0; u.a)$ with $u \notin \mathrm{FV}(a)$. The successor computation gives $\mathrm{wr}(a) = a$, also in $\mathcal{C}_\nu$. Transport of this equation along itself in the template $v = a$ gives $a = a$. The templates $v = s$, $s = v$ and $f(s) = f(v)$ give symmetry, transitivity and congruence.

From $\Gamma, B(0) \vdash B(Sz)$, introduction by cases with an unused witness coordinate gives $D_{w,y}B(y)$, and DE followed by Hyp exports $Q_yB(y)$. With proofs of $B(0)$ and $B(Sz)$, QE gives $B(t)$; this is numerical cases. The same construction with introduction by progress gives $Q_yE(y)$ from $\Gamma, E(z) \vdash E(Sz)$, and with a proof of $E(0)$ it gives $E(t)$. □

Lemma 9.32 (Body extensionality, native form)**.** *In both unrestricted calculi, if $\Gamma \vdash h(v) = k(v)$ for a variable v fresh for Γ, then $\Gamma \vdash R(n, a; u.h(u)) = R(n, a; u.k(u))$.*

Proof. Apply induction on literal equations to the equation $R(z, a; u.h) = R(z, a; u.k)$ at a fresh counter z, with the initializer and the parameters fixed. At 0 both terms are Sa, by the zero computation in $\mathcal{R}_{\mathrm{raw}}$ and by identity of canonical spelling and reflexivity in $\mathcal{C}_\nu$. At a successor the computation exposes the two updates. Replace the old state in h by the induction equation, and use the premise at the second old state; this instance of the premise is justified by Theorem 9.30. □

The lemma is necessary. Changing a normalized body changes a finite description; it is not congruence for one fixed function symbol.

Lemma 9.33 (Normalization is provable in the raw calculus)**.** *For every raw term t, $\mathcal{R}_{\mathrm{raw}} \vdash t = \nu(t)$. For every raw native formula A, the raw calculus derives $A \vdash \nu(A)$ and $\nu(A) \vdash A$, uniformly in the free parameters.*

Proof. For terms induct on t. The counter and the initializer change by congruence. The body changes by Theorem 9.32 and the induction hypothesis with the bound variable free. If the counter is literally 0, the raw zero computation replaces the body by 0.

For formulas induct on the formula. An equation follows from the two endpoint identities. For $Q_yA(y)$, under this assumption introduce $Q_y\nu(A)(y)$ by cases: convert the assumed normalized base to the old base, apply QE at the fresh successor and convert back. The reverse direction is identical. For $D_{x,y}A$, eliminate the old formula at a fresh witness, introduce the new one at that witness by cases, convert the new base to the old base, use the exported closure, and convert the successor conclusion. These are derived congruence arguments for the two binders; no ordinary quantifier logic is assumed. □

Theorem 9.34 (Canonical and raw consequence)**.** *For zero-body canonical Γ, A,*

$$\Gamma \vdash_{\mathcal{C}_\nu} A \iff \Gamma \vdash_{\mathcal{R}_{\mathrm{raw}}} A.$$

More generally, ν sends every raw derivation to a canonical derivation of the normalized sequent.

Proof. From raw to canonical, normalize every sequent of the derivation. By Theorem 9.3 an instance of transport, QE, DE or an introduction becomes an instance of the same rule with normalized substitution. A raw zero computation becomes $Sa = Sa$, which is derivable by Theorem 9.31, and a successor computation becomes an instance of (R_+).

Conversely, regard canonical expressions as raw ones and recurse on the canonical derivation. A canonical rule instance differs from the raw instance only by normalization of substituted expressions. Theorem 9.33 converts premises, local assumptions and conclusions in

both directions, and cut handles the local assumptions. For example, canonical transport converts the minor premise $\nu(E[s/v])$ to $E[s/v]$, applies raw transport and converts $E[t/v]$ to its normalization. Freshness is preserved because ν introduces no free variable. No substitution theorem for a restricted rule enters either direction. □

9.4.2 The six rules

For a native formula define the root reversal by $(s = t)^\dagger = (t = s)$, $(Q_y B)^\dagger = Q_y B$ and $(D_{x,y} B)^\dagger = D_{x,y} B$. It commutes with normalized substitution.

Definition 9.35 (The calculus $\mathcal{K}_6$)**.** Contexts are finite reusable lists of zero-body canonical native formulas. The rules are:

1. Hyp: $\Gamma \vdash A$ if $A \in \Gamma$.
2. Successor computation: every instance of (R_+), in every context.
3. Crossed equational transport, for a template $E(v)$ that is a literal equation (constant templates are allowed):
$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash E\langle s/v\rangle}{\Gamma \vdash (E\langle t/v\rangle)^\dagger} \qquad (s \not\equiv 0). \tag{X}$$
4. Crossed closure elimination:
$$\frac{\Gamma \vdash Q_y B(y) \qquad \Gamma \vdash B\langle 0/y\rangle}{\Gamma \vdash (B\langle t/y\rangle)^\dagger}. \tag{QX}$$
5. Paired elimination, with b fresh for Γ, A and the parameters of B:
$$\frac{\Gamma \vdash D_{x,y} B(x,y) \qquad \Gamma,\ Q_y B\langle b/x\rangle \vdash A}{\Gamma \vdash A}. \tag{DE}$$
6. One introduction, with z fresh for Γ, t and the parameters of B:
$$\frac{\Gamma,\ B\langle t/x, \eta/y\rangle \vdash B\langle t/x, Sz/y\rangle}{\Gamma \vdash D_{x,y} B} \qquad \eta = \begin{cases} z, & B \text{ is a literal equation and } t \equiv 0, \\ 0, & \text{otherwise.} \end{cases} \tag{WI}$$

There is no primitive reflexivity, symmetry, numerical case rule, induction rule, zero-computation rule, logical connective or rule of numerical substitution. On iterator terms, one-term transport at sources other than the literal 0 together with the single witness-directed introduction is the seven-rule calculus `MPrf`, which is formalized and proved equivalent to the kernel (`mprf_iff_nprf`). $\mathcal{K}_6$ crosses its two eliminations, takes equation templates, and has no zero computation; what is specific to it is the canonical syntax. Weakening, renaming of eigenvariables and cut of an ordinary assumption are admissible in $\mathcal{K}_6$ by induction on derivations: cut replaces uses of an assumption by proofs and changes no witness, root or source term, and renaming preserves literal zero and roots. Numerical substitution is not claimed at this stage.

Lemma 9.36 (Equality in $\mathcal{K}_6$)**.** *$\mathcal{K}_6$ derives reflexivity, symmetry, transitivity, congruence, uncrossed transport in every literal-equation template without the restriction on the source, and the ordinary rule QE.*

Proof. Keep $\mathrm{wr}(a) = R(S0, 0; u.a)$ and put $c := \mathrm{wr}(0)$. Each $\mathrm{wr}(a)$ is not literally 0, and $\mathrm{wr}(a) = a$ is an instance of (R_+).

Symmetry. (X) with major premise $c = 0$ and a constant template reverses any proved equation. Since the reversal of a compound formula is the identity, (QX) followed by a reversal is ordinary QE, and (X) followed by a reversal is uncrossed transport at sources that are not literally 0.

Reflexivity. Use $\mathrm{wr}(a) = a$ as both premises of (X) with the template $v = a$.

Zero join. Let $p, q \not\equiv 0$ with $p = 0$ and $q = 0$. (X) with major $p = 0$, minor $\mathrm{wr}(p) = p$ and template $\mathrm{wr}(v) = p$ gives $p = c$; similarly $q = c$. Reverse the second and transport $p = c$ along $c = q$, whose source c is allowed, to get $p = q$.

Transitivity. Given $s = t$ and $t = r$: if $t \not\equiv 0$ transport $s = t$ along $t = r$. If $t \equiv 0$ and both endpoints are not literally 0, reverse the second equation and use the zero join. If an endpoint is literally 0, the conclusion is one of the premises.

Congruence. Along $s = t$ with $s \not\equiv 0$, transport the reflexive equation $f(s) = f(s)$ in the template $f(s) = f(v)$. For $0 = t$ with $t \not\equiv 0$, reverse to $t = 0$ and transport $f(t) = f(t)$ in the template $f(v) = f(t)$, which gives $f(0) = f(t)$. If both endpoints are literally 0 use reflexivity.

Transport. From $s = t$ and $a(s) = b(s)$, congruence and transitivity give $a(t) = a(s) = b(s) = b(t)$. Every step instantiates a schema at the terms at hand; none substitutes into a derivation. □

Lemma 9.37 (Introduction by cases)**.** *Every instance of introduction by cases is derivable in* $\mathcal{K}_6$*.*

Proof. (WI) is introduction by cases when the matrix is compound or the witness is not literally 0. Let E be an equation matrix with the premise $\Gamma, E(0,0) \vdash E(0, Sz)$. Use the witness c, which is not literally 0 and is provably 0. Under the assumption $E(c, 0)$, Theorem 9.36 gives $E(0, 0)$; cut with the premise gives $E(0, Sz)$, and transport back gives $E(c, Sz)$. (WI) at the witness c is in case mode and concludes the unchanged formula $D_{x,y}E(x, y)$. The witness c does not contain z. □

Lemma 9.38 (Introduction by progress)**.** *Every instance of introduction by progress for a literal equation is derivable in* $\mathcal{K}_6$*.*

Proof. Let the premise be $\Gamma, E(t, z) \vdash E(t, Sz)$ and let w be unused in $E(t, y)$. (WI) with witness 0 is in progress mode and proves $\Gamma \vdash D_{w,y}E(t, y)$. Under the additional assumption $E(t, 0)$ eliminate this formula. The exported closure is $Q_yE(t, y)$, independent of the fresh witness, and ordinary QE gives $E(t, Sz)$. Closing (DE) gives the premise of Theorem 9.37 at the witness t. □

Lemma 9.39 (Transport in compound templates)**.** $\mathcal{K}_6$ *derives transport in every native formula template.*

Proof. Induct on the template. Equations are Theorem 9.36. For Q_yB, introduce the new closure by cases with an unused witness as in Theorem 9.31: convert its base back along the equation by the induction hypothesis, apply the old closure at the fresh successor, and convert forward. For $D_{x,y}B$, (DE) opens the old formula at a fresh witness; introduce the new formula by cases at that witness, convert its base back, use the exported closure, and convert the successor forward. The other direction of the equation is obtained by symmetry. Only symmetry, cut and renaming are used. □

Theorem 9.40 (Six-rule equivalence)**.** *For zero-body canonical* Γ, A*,*

$$\Gamma \vdash_{\mathcal{K}_6} A \iff \Gamma \vdash_{\mathcal{C}_\nu} A.$$

Proof. Theorems 9.36 to 9.39 provide every rule of $\mathcal{C}_\nu$ inside $\mathcal{K}_6$. Conversely an instance of (WI) is one of the two unrestricted introductions, and (X) and (QX) are transport and QE followed by the derived symmetry when the conclusion is an equation; the other rules are common. No admissibility of numerical substitution in $\mathcal{K}_6$ is presupposed; it now transfers through the equivalence. □

Remark 9.41. Normalization preserves the literal-zero status of a whole term and the root of a formula, so it does not alter a test of (X) or (WI). An arbitrary numerical substitution can alter such a test: a variable witness can become 0. Keeping the unrestricted equivalence of Theorem 9.34 and the recovery of the restricted rules separate avoids an invalid rule-by-rule substitution argument.

9.4.3 The bridge for the seeded calculi

Theorem 9.42. *Extend ι and κ of Theorem 9.24(1) to native formulas, fixing $=$, Q_y and $D_{x,y}$.*

1. *If $\Gamma \vdash_{\mathcal{R}_{\mathrm{raw}}} A$ then $\Gamma^\iota \vdash_{\mathsf{N}} A^\iota$, and if $\Gamma \vdash_{\mathsf{N}} A$ then $\Gamma^\kappa \vdash_{\mathcal{R}_{\mathrm{raw}}} A^\kappa$.*
2. *$\mathcal{R}_{\mathrm{raw}} \vdash t^{\iota\kappa} = t$ and $\mathsf{N} \vdash s^{\kappa\iota} = s$; every native formula is interderivable with its double translation in both calculi.*
3. *For zero-body canonical Γ, A: $\Gamma \vdash_{\mathcal{K}_6} A$ if and only if $\Gamma^\iota \vdash_{\mathsf{N}} A^\iota$.*

Consequently $\mathcal{K}_6$, $\mathcal{C}_\nu$ and $\mathcal{R}_{\mathrm{raw}}$ are sound and faithful for T^1_R under the reading of Section 1.2, and each is equiconsistent and bi-interpretable with iPRA*, with whole domains, actual equality and numerical composites of graph $x = y$. On raw syntax the composite through canonical syntax is provably, not literally, the identity; on canonical syntax ν is literally idempotent.*

Proof. (1) From $\mathcal{R}_{\mathrm{raw}}$ to N: the computation equations translate to the computation equations of the iterator, using $\mathsf{N} \vdash (St)^\iota = S(t^\iota)$ and the Leibniz rule of Section 1.4; transport of formulas is that Leibniz rule; QE, DE and introduction by cases are rules of N; and introduction by progress is derived by forming $Q_y E(t, y)$ by progress, applying QE under the assumption $E(t, 0)$ and introducing by cases at t. From N to $\mathcal{R}_{\mathrm{raw}}$: the computation equations of the iterator are Theorem 8.21 with (9.17); the one-term transport of N is transport followed by symmetry; and the progress rule of N is introduction by progress with an unused witness followed by DE and Hyp, as in Theorem 9.31.

(2) By Theorems 9.30 and 9.31 the proof of Theorem 9.21 is available in $\mathcal{R}_{\mathrm{raw}}$ and in N: it uses equality reasoning, an instance of a derived equation and induction on a literal equation. The argument of Theorem 9.25(2) then gives the two term identities. A formula differs from its double translation by these provable identities at its terms, with the bound variables of the formula free; the binder-congruence argument of Theorem 9.33 gives interderivability.

(3) By Theorems 9.34 and 9.40, derivability in $\mathcal{K}_6$ is derivability in $\mathcal{R}_{\mathrm{raw}}$. One direction is (1). For the other, a derivation of $\Gamma^\iota \vdash_{\mathsf{N}} A^\iota$ gives $\Gamma^{\iota\kappa} \vdash A^{\iota\kappa}$ in $\mathcal{R}_{\mathrm{raw}}$, and (2) with cut gives $\Gamma \vdash A$. The consequences follow from Theorem 1.78 and Theorem 9.25. □

The mathematical change in this section is the canonical syntax and the proof that it preserves substitution, derivability and both composites. The inventory decreases from seven rules to six, and the successor constructor is gone, but the omitted zero computation is the law $R(0, a; u.h) \equiv R(0, a; u.0)$ of the syntax, which every substitution and comparison of a complete checker enforces.

9.5 Five rules with clause contexts

This section draws on `clausal_canonical`.

The calculus of this section is the clause presentation of Section 1.9 with the one-term transport of Section 1.3, stated on the zero-body canonical terms of the seeded recursor with one bound variable. The closure binder is no longer a formula constructor, and the hypothesis rule and the closure elimination become one rule of clause use. Only what the canonical syntax and the missing reflexivity rule require is proved.

9.5.1 The calculus

Definition 9.43 (The calculus $\mathcal{K}_5$)**.** Terms are zero-body canonical terms of the seeded recursor, and all substitutions are normalized. Formulas are $F ::= (s = t) \mid D_{x,y}F$. A context is a finite reusable list of clauses $[A;\, y.\, B(y)]$ with $y \notin \mathrm{FV}(A)$; clauses are context objects, not formulas or goals. Let

$$\mathrm{wr}(a) := R(S0, 0; u.a), \qquad c := \mathrm{wr}(0), \qquad H := (c = 0).$$

An ordinary assumption F is the constant clause $[H;\, v.\, F]$ with v unused; this applies also to the local assumption of rule (I). The rules are:

1. Computation, in every context:

$$R(Sn, a; u.h) = h\langle R(n, a; u.h)/u\rangle. \tag{R}$$

2. One-term crossed transport, for a term template $h(v)$ and an arbitrary source:

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash h(s) = r}{\Gamma \vdash r = h(t)}. \tag{T}$$

3. Clause use, for $[A;\, y.\, B(y)] \in \Gamma$:

$$\frac{\Gamma \vdash A}{\Gamma \vdash B(t)}. \tag{U}$$

4. Paired elimination, with b fresh for Γ, F and the parameters of B:

$$\frac{\Gamma \vdash D_{x,y}B(x, y) \qquad \Gamma,\ [B(b, 0);\, y.\, B(b, y)] \vdash F}{\Gamma \vdash F}. \tag{E}$$

5. Witness-directed introduction, with z fresh for Γ, t and the parameters of B:

$$\frac{\Gamma,\ B(t, \eta) \vdash B(t, Sz)}{\Gamma \vdash D_{x,y}B(x, y)} \qquad \eta = \begin{cases} z, & B \text{ is a literal equation and } t \equiv 0, \\ 0, & \text{otherwise.} \end{cases} \tag{I}$$

There is no separate hypothesis rule, zero-computation rule, closure constructor, closure elimination, reflexivity axiom, case rule or induction rule. The neutral antecedent is H and not $0 = 0$, because H is an instance of (R) and reflexivity is not yet available. The clause pair, its bound variable and its substitution are part of the specification and of the implementation; they are not a saving that goes uncounted.

Computation proves $\mathrm{wr}(a) = a$ in every context, in particular H. Use of an ordinary assumption is therefore computation of H followed by (U). Weakening, renaming of fresh parameters and cut of an ordinary assumption follow by induction on derivations; in a cut each use of the constant clause is replaced by the supplied proof of its conclusion. No substitution theorem for rule (I) is presupposed.

Lemma 9.44 (Equality from one term template)**.** *$\mathcal{K}_5$ derives reflexivity, symmetry, transitivity, congruence and transport in every literal-equation template.*

Proof. Use $\mathrm{wr}(a) = a$ as both premises of (T) with the identity template to get $a = a$. To reverse $s = t$, take the major premise H and the constant template $h(v) = s$; (T) returns $t = s$. Given $s = t$ and $t = r$, reverse the former and apply (T) with the identity template to the major $t = r$ and the minor $t = s$, which gives $s = r$. Congruence is (T) with the major $s = t$ and the reflexive minor $h(s) = h(s)$. For a template $a(v) = b(v)$ the two applications

$$s = t,\ a(s) = b(s) \implies b(s) = a(t), \qquad s = t,\ b(s) = a(t) \implies a(t) = b(t)$$

give transport. Each template is fixed finite syntax; no function variable occurs. □

Lemma 9.45 (All case introductions)**.** *For every formula B and term t the rule "from $\Gamma, B(t,0) \vdash B(t,Sz)$ infer $D_{x,y}B(x,y)$" is admissible in $\mathcal{K}_5$.*

Proof. The only instance that is not primitive has an equation matrix E and the witness 0. Use the witness c. Under $E(c,0)$, the theorem $c = 0$ and Theorem 9.44 give $E(0,0)$; cut with the supplied branch gives $E(0,Sz)$; transport back gives $E(c,Sz)$. Rule (I) at the witness c is in case mode and proves the original goal. □

Lemma 9.46 (Transport in every formula)**.** *For every template $F(v)$, from $s = t$ and $F(s)$ one derives $F(t)$ in $\mathcal{K}_5$, under arbitrary retained clauses.*

Proof. Induct on the template. Equations are Theorem 9.44. For $F(v) = D_{x,y}B(v,x,y)$, eliminate $F(s)$ at a fresh b, which retains the clause $[B(s,b,0);\ y.\, B(s,b,y)]$. Introduce the target by Theorem 9.45 at the witness b: convert the local $B(t,b,0)$ to $B(s,b,0)$ by symmetry and the induction hypothesis, apply (U) at Sz, and convert $B(s,b,Sz)$ to $B(t,b,Sz)$. The recursion uses only the smaller matrix. □

9.5.2 Closure as an abbreviation

For a fresh unused variable w write $q_y B(y) := D_{w,y}B(y)$.

Lemma 9.47. *In $\mathcal{K}_5$:*

1. *from $\Gamma, B(0) \vdash B(Sz)$ one derives $\Gamma \vdash q_y B$;*
2. *from $q_y B$ and $B(0)$ one derives $B(t)$;*
3. *in every context that contains $[B(0);\ y.\, B(y)]$ one derives $q_y B$;*
4. *for a literal equation E, from $\Gamma, E(z) \vdash E(Sz)$ one derives $\Gamma \vdash q_y E$; hence induction on literal equations with an arbitrary context, numerical cases for all formulas, and introduction by progress at every witness are derivable.*

Proof. (1) is Theorem 9.45 with an unused witness. (2) Eliminate the paired formula, use the opened clause at t, and close the elimination. (3) Under the assumption $B(0)$ of the case introduction use the clause at Sz. (4) Rule (I) at the witness 0 with an unused witness coordinate is in progress mode. With (2) this gives induction, and (1) with (2) gives numerical cases. For progress at a witness t, form $q_y E(t,y)$ by progress, and under the base $E(t,0)$ specialize it at Sz and introduce by cases at t. □

Items (1)–(3) are introduction, elimination and eta expansion of a closure resource. The third is what translates a branch of (DE) in $\mathcal{K}_6$, which assumes a closure formula.

9.5.3 Comparison with the six rules

Define T from the formulas of $\mathcal{K}_6$ to those of $\mathcal{K}_5$ by

$$T(s = t) = (s = t), \qquad T(D_{x,y}B) = D_{x,y}T(B), \qquad T(Q_yB) = q_yT(B),$$

with every inserted coordinate fresh and unused. T preserves free variables, the distinction between equation and compound roots, and normalized substitution. Let U be the inclusion of the formulas of $\mathcal{K}_5$ into the native formulas, and let $\mathrm{ord}(\Delta)$ replace each formula of a list Δ by its constant clause.

Theorem 9.48 (Preservation in both directions)**.** *For every native context* Γ *and formula* A*,*

$$\Gamma \vdash_{\mathcal{K}_6} A \implies \mathrm{ord}(T\Gamma) \vdash_{\mathcal{K}_5} T(A),$$

and for formulas Δ, F *of* $\mathcal{K}_5$*,*

$$\mathrm{ord}(\Delta) \vdash_{\mathcal{K}_5} F \implies \Delta \vdash_{\mathcal{K}_6} F.$$

Proof. Forward. Induct on the derivation in $\mathcal{K}_6$. Hyp uses H and the constant clause. Computation is unchanged. (X) is derived transport followed by symmetry. (QX) is Theorem 9.47(2), followed by symmetry when the conclusion is an equation. (WI) is (I): the translation preserves roots and the witness is unchanged. For (DE), the translated major premise proves $D_{x,y}T(B)$; rule (E) opens $[T(B)(b,0);\, y.\, T(B)(b,y)]$; Theorem 9.47(3) derives the translated assumption $q_yT(B)(b,y)$ of the old branch, and cut inserts this proof into the translated branch. All new variables are freshened first.

Backward. Track each context occurrence by the rule that introduced it. A constant clause is represented by its formula as an assumption of $\mathcal{K}_6$, and a clause opened by (E) by the assumption $Q_yB(b,y)$. (U) on a constant clause is Hyp, and (U) on an opened clause is ordinary QE, which $\mathcal{K}_6$ derives (Theorem 9.36). (R) is the same computation, (T) is derived one-term transport, (E) is (DE), and (I) is (WI). Induction on the derivation gives the claim. □

A clause $[A;\, y.\, B]$ supplied from outside, which is neither constant nor opened by (E), can be interpreted by the formula for $A \to \forall y\, B(y)$ built from the logical macros; (U) is then implication elimination followed by universal elimination. A checker for closed certificates needs no such clause.

Theorem 9.49 (Both composites)**.** $T(U(F)) = F$ *literally for every formula* F *of* $\mathcal{K}_5$*, and* $A \dashv\vdash_{\mathcal{K}_6} U(T(A))$ *for every native formula* A*. Both numerical translations are the identity. Consequently* $\mathcal{K}_5$ *is equiconsistent and bi-interpretable with* iPRA*, with whole domains, actual equality and numerical composites of graph* $x = y$*; the designated contradiction* $0 = S0$ *is unchanged.*

Proof. The first identity is immediate from the clauses of T. For the second use $Q_yB(y) \dashv\vdash D_{w,y}B(y)$ in $\mathcal{K}_6$ with w unused: forward, under $B(0)$ specialize the closure at Sz and introduce the paired formula by cases; backward, (DE) opens exactly Q_yB at an unused witness and Hyp closes the branch. The binder congruence of Theorem 9.33 lifts this through Q_y and $D_{x,y}$, by induction on the formula. No quotient, retract or reinterpretation of equality is introduced. The consequence follows from Theorems 9.42 and 9.48. □

Under the reading $[A;\, y.\, B]^\circ = A^\circ \to \forall y\, B^\circ(y)$ the five rules are directly sound in T^1_R: (R) is a defining equation, (T) is transport followed by symmetry, (U) is implication and universal elimination, (E) is existential elimination, and (I) is coverage in case mode and induction on an equation under the assumed zero instance in progress mode, followed by existential introduction. Certificates of $\mathcal{K}_5$ can be longer than those of $\mathcal{K}_6$ while the checker is shorter: an ordinary assumption needs the neutral computation premise, and the removal of the closure constructor adds elimination and eta-expansion steps.

9.6 Counter evaluation

This section draws on `counter_evaluation` and `uniform_binding`.

On the counter-aware recursor, the computation rule and the transport rule merge into one rule that places the supplied equation in the counter. The rule has no analogue on iterator terms, since it needs a body that sees the counter.

9.6.1 The clause syntax and a liberal calculus

Throughout this section and the next, terms are canonical terms of the counter-aware language, either zero-body canonical (with ν) or normal (with nf), and every displayed substitution is normalized accordingly. Formulas are $A ::= (s = t) \mid D_{x,y}B$ with the reading of Section 1.2. A context is a finite reusable list of clauses $[A;\, y.\, B(y)]$, $y \notin \mathrm{FV}(A)$, read as $A^\circ \to \forall y\, B^\circ(y)$ as in Section 1.9; a clause may also have an empty antecedent, $[\,;\, y.\, B(y)]$, read as $\forall y\, B^\circ(y)$. An ordinary assumption A is the constant clause $[0 = 0;\, v.\, A]$, or $[\,;\, v.\, A]$ where empty antecedents are used, with v unused. The designated contradiction is $0 = 1$.

The calculi below have rules with tests on literal syntax. The following calculus has none and serves as the common reference.

Definition 9.50 (The liberal clause calculus $\mathcal{L}$)**.** The rules of $\mathcal{L}$ are:

1. Refl: $\Gamma \vdash t = t$.
2. Comp: $\Gamma \vdash R(Sn, a; i, u.h) = h\langle n/i, R(n, a; i, u.h)/u\rangle$.
3. Tr: from $s = t$ and $h\langle s/v\rangle = r$ infer $r = h\langle t/v\rangle$, for a term template h.
4. Use: for $[A;\, y.\, B(y)] \in \Gamma$, from $\Gamma \vdash A$ infer $\Gamma \vdash B\langle t/y\rangle$; for a clause with empty antecedent there is no premise.
5. DE: from $D_{x,y}B(x, y)$ and a derivation of A from $\Gamma, [B(b, 0);\, y.\, B(b, y)]$, with b fresh for Γ, A and the parameters of B, infer A.
6. Case: from $\Gamma, B(t, 0) \vdash B(t, Sz)$ infer $D_{x,y}B(x, y)$, for every matrix and witness.
7. Prog: from $\Gamma, E(t, z) \vdash E(t, Sz)$ infer $D_{x,y}E(x, y)$, for a literal equation E.

In the last two rules z is fresh for Γ, t and the parameters of the matrix. On normal terms both sides of Comp have the same spelling, and the rule is an instance of Refl.

Lemma 9.51 (Structure of $\mathcal{L}$)**.** *The following are admissible in $\mathcal{L}$: weakening; injective renaming of parameters; numerical substitution; cut of an ordinary assumption; and cut of a clause, that is, from $\Gamma, [A;\, y.\, B(y)] \vdash F$ and $\Gamma, A \vdash B(y)$ with y fresh for Γ, A infer $\Gamma \vdash F$.*

Proof. Induct on derivations after freshening eigenparameters. Every rule is a schema without a test on the shape of a term, so Theorem 9.3, respectively Theorem 9.7, shows that a substitution instance of a rule instance is a rule instance. For the cut of an assumption replace each use of its constant clause by the supplied proof. For the cut of a clause replace each use of the clause at a term t, whose premise proves A, by the second derivation with t substituted for y, followed by a cut of the assumption A. □

Lemma 9.52 (Derived rules of $\mathcal{L}$)**.** *$\mathcal{L}$ derives:*

1. *symmetry, transitivity, congruence, and transport in every equation template;*
2. *numerical cases: from $\Gamma \vdash F(0)$ and $\Gamma \vdash F(Sz)$, z fresh, infer $\Gamma \vdash F(t)$;*

3. *equation induction: from* $\Gamma \vdash E(0)$ *and* $\Gamma, E(z) \vdash E(Sz)$*, for a literal equation* E *and* z *fresh, infer* $\Gamma \vdash E(t)$*;*

4. *transport in every formula template: from* $s = t$ *and* $F(s)$ *infer* $F(t)$*;*

5. *binder congruence: if* $B(x, y)$ *and* $B'(x, y)$ *are interderivable in every context uniformly in* x, y*, then so are* $D_{x,y}B$ *and* $D_{x,y}B'$*.*

Proof. (1) As in Theorem 9.44, with Refl in place of the two uses of wr: to reverse $s = t$ apply Tr to the major $0 = 0$ and the constant template.

(2), (3) For an unused coordinate w write $q_y A := D_{w,y}A(y)$. Case introduces $q_y F$ from $\Gamma \vdash F(Sz)$, ignoring the assumed base, and Prog introduces $q_y E$ from the step. DE opens the clause $[A(0);\, y.\, A(y)]$, the given base proves its antecedent, and Use at t finishes.

(4) Induct on the template; equations are (1). For $F(v) = D_{x,y}B(v, x, y)$, open $F(s)$ at a fresh b, which retains $[B(s, b, 0);\, y.\, B(s, b, y)]$. Introduce $F(t)$ by Case at the witness b: convert the local assumption $B(t, b, 0)$ backward along $s = t$ by the induction hypothesis, use the old clause at Sz, and convert the result forward to $B(t, b, Sz)$. Close Case and DE.

(5) The same construction with the two conversions given by the hypothesis. □

9.6.2 The five patterns of counter evaluation

Definition 9.53 (The calculus $\mathcal{K}_{\mathrm{CE}}$)**.** Terms are zero-body canonical. Write $\varrho(t, a, h)$ for the canonical term $R(t, a; i, u.h)$, which is Sa when $t \equiv 0$. Every clause has an antecedent, and an ordinary assumption A is $[0 = 0;\, v.\, A]$. The rules are Refl, Use and DE of Theorem 9.50, the rule of counter evaluation

$$\frac{\Gamma \vdash s = t \qquad \Gamma \vdash R(Ss, a; i, u.h) = r}{\Gamma \vdash r = h\langle t/i, \varrho(t, a, h)/u\rangle} \tag{CE}$$

and the witness-directed introduction, with z fresh for Γ, w and the parameters of B,

$$\frac{\Gamma,\ B(w, \eta) \vdash B(w, Sz)}{\Gamma \vdash D_{x,y}B(x, y)} \qquad \eta = \begin{cases} z, & B \text{ is a literal equation and } w \equiv 0, \\ 0, & \text{otherwise.} \end{cases} \tag{DI}$$

In (CE) the two substitutions in h are simultaneous. The seed, the body and the passive parameters are read from the second premise and are kept fixed: occurrences of s inside a or h are not replaced. The first endpoint s is an arbitrary term, including the literal 0. There is no further computation axiom, equality rule, induction rule, successor axiom or logical connective. The count of five patterns counts (DI) as one schema with two modes; it is not a claim under every convention of counting rules. `uniform_binding` calls DE and (DI) Open and Close.

Theorem 9.54 (Arithmetic reconstruction)**.** *Refl and* (CE) *alone derive the successor computation Comp, symmetry, transitivity, congruence in every term template, and the crossed transport Tr. The derivations use no logical rule, no numerical cases and no induction.*

Proof. Computation. Apply (CE) to the reflexive premises $n = n$ and $B = B$ with $B = R(Sn, a; i, u.h)$. The conclusion is $B = h\langle n/i, \varrho(n, a, h)/u\rangle$.

Root composition. Let p prove $s = t$ and q prove $s = r$, and put $K_s := R(Ss, 0; i, u.i)$. (CE) applied to q and $K_s = K_s$ gives $K_s = r$, and applied to p and this result it gives $r = t$. Write $\mathsf{J}(p, q)$ for this derivation of $r = t$ from $s = t$ and $s = r$.

Symmetry and transitivity. $\mathsf{J}(\mathrm{Refl}(s), p)$ proves $t = s$ from $p\colon s = t$. Given $p\colon s = t$ and $q\colon t = v$, the derivation $\mathsf{J}(q, \mathrm{Sym}(p))$ proves $s = v$.

Congruence. For a template $F(v)$ choose the binders fresh and put $B_F := R(Ss, 0; i, u.F(i))$. (CE) on two reflexive premises gives $B_F = F(s)$, and with first premise $p\colon s = t$ and this second premise it gives $F(s) = F(t)$. The template may have repeated occurrences and occurrences under the binders of nested recursors.

Crossed transport. Apply J to $F(s) = F(t)$ and the supplied $q\colon F(s) = r$; the result is $r = F(t)$. Every substitution in the argument is an instance of (CE); no congruence is presumed in order to establish congruence. □

Proposition 9.55 (Proof costs)**.** *Count every primitive inference, including a* Refl *leaf, as one node and count repeated subtrees repeatedly. Root composition adds three nodes to its premises, congruence adds four, and crossed transport adds seven. Each supplied premise occurs once.*

Proof. $\mathsf{J}(p, q)$ is two applications of (CE) and one leaf. Congruence is two applications and two leaves. Crossed transport is J applied to a congruence, $4+3 = 7$. The term labels contain copies of F and of the endpoints, so these are bounds on inference nodes, not on bytes or running time. □

Theorem 9.56 (Equality-interface equivalence)**.** *Let $\mathcal{K}_{\mathrm{CT}}$ be $\mathcal{K}_{\mathrm{CE}}$ with* Refl *and* (CE) *replaced by* Comp *and* Tr*. A sequent is derivable in $\mathcal{K}_{\mathrm{CE}}$ if and only if it is derivable in $\mathcal{K}_{\mathrm{CT}}$, and the two proof translations are effective and preserve end sequents literally.*

Proof. Theorem 9.54 replaces each Comp or Tr node. Conversely, in $\mathcal{K}_{\mathrm{CT}}$ the computation with the counter projection gives $K_s = s$, and Tr applied to two copies of this equation with the identity template gives $s = s$. Tr with a constant template gives symmetry, with the identity template and a reversed premise it gives transitivity, and with a reflexive minor premise it gives congruence. For (CE): from $s = t$, congruence in the counter gives $R(Ss, a; i, u.h) = R(St, a; i, u.h)$, and Comp gives $R(St, a; i, u.h) = h\langle t/i, \varrho(t, a, h)/u\rangle$; reverse the second premise $R(Ss, a; i, u.h) = r$ and compose. The three logical rules are common. The use of ordinary equality in the converse is legitimate, since it is derived in $\mathcal{K}_{\mathrm{CT}}$ before (CE) is simulated; the forward direction never assumes it. □

Lemma 9.57 (Both introductions)**.** *Case and Prog are derivable in $\mathcal{K}_{\mathrm{CE}}$ for every matrix and witness. Before this, weakening, renaming of fresh parameters and cut of an ordinary assumption are admissible in $\mathcal{K}_{\mathrm{CE}}$.*

Proof. The three structural operations do not change the literal-zero test of (DI); for cut, replace each Use of the constant clause $[0 = 0; v.\, A]$ by the supplied proof of A and discard the proof of its antecedent.

The only instance of Case that is not an instance of (DI) has an equation matrix E and the witness 0. Let $Z := R(S0, 0; i, u.0)$. It is not literally 0, and Theorem 9.54 proves $Z = 0$. Under $E(Z, 0)$, transport in the equation gives $E(0, 0)$; cut with the premise gives $E(0, Sz)$; transport back gives $E(Z, Sz)$. (DI) at the witness Z is in case mode and concludes $D_{x,y}E(x, y)$. Only transport in an equation is used, not transport through a binder.

For Prog at a witness w, introduce $D_{v,y}E(w, y)$ with v unused at the witness 0; this is (DI) in progress mode. DE opens $[E(w, 0); y.\, E(w, y)]$. Under $E(w, 0)$, Use gives $E(w, Sz)$, and Case at w gives $D_{x,y}E(x, y)$. Close DE. □

Theorem 9.58. *On zero-body canonical terms, $\mathcal{K}_{\mathrm{CE}}$, $\mathcal{K}_{\mathrm{CT}}$ and $\mathcal{L}$ derive the same sequents whose clauses have antecedents. In particular numerical substitution is admissible in $\mathcal{K}_{\mathrm{CE}}$.*

Proof. Every rule of $\mathcal{L}$ is derivable in $\mathcal{K}_{\mathrm{CE}}$ by Theorems 9.54 and 9.57. Conversely (CE) is derivable in $\mathcal{L}$ as in Theorem 9.56, and an instance of (DI) is an instance of Case or of Prog. Substitution is Theorem 9.51. □

The test of (DI) is not invariant under substitution: a variable witness can become 0. A progress introduction keeps its equation root and its literal witness 0, but a case introduction whose witness becomes 0 has to be replaced by the derived Case. Admissibility of substitution is obtained through $\mathcal{L}$, not by a rule-by-rule argument.

9.6.3 Ordinary logic with the decoder

The derivation of intuitionistic logic from the paired binder is that of the gates, selectors and connectives of Section 1.4; it needs a selector with the two endpoint equations. With the decoder P of Theorem 9.10 the conjunction and the universal quantifier take a shorter form: no parity iterator and no separate zero instance are needed. The construction is given in full for $\mathcal{L}$, on either kind of canonical terms.

Definition 9.59 (Gates, selection and the logical macros)**.** With binders renamed away from the control v, put

$$G_v(s=t) := \big(C(v,0,s) = C(v,0,t)\big), \qquad G_v(D_{x,y}B) := D_{x,y}G_vB,$$

and $L(F) := D_{w,y}G_yF$ with w, y fresh. Every formula is a finite chain of paired binders ending in an equation. For formulas A, B apply L to the shorter one until the depths agree, align the bound names, and at the equations put

$$M_v\big((s=t),(a=b)\big) := \big(C(v,s,a) = C(v,t,b)\big),$$

carrying the matched binders outside. With all auxiliary binders fresh and w unused define

$$\begin{aligned} \bot_* &:= (0=1), & A \to_* B &:= D_{w,y}M_y(A,B),\\ A \wedge_* B &:= D_{w,y}G_yM_{P(y)}(A,B), & \exists_* x\, A(x) &:= D_{x,y}G_yA(x),\\ A \vee_* B &:= \exists_* x\, M_x(A,B), & \forall_* x\, A(x) &:= D_{w,y}G_yA(P(y)). \end{aligned} \tag{9.19}$$

These are finite formulas of the same syntax, not further constructors.

Lemma 9.60 (Gates and selection)**.** *In $\mathcal{L}$, for all formulas and in every context: $\vdash G_0F$; $G_{Sz}F \dashv\vdash F$; $L(F) \dashv\vdash F$; $M_0(A,B) \dashv\vdash A$; and $M_{Sz}(A,B) \dashv\vdash B$.*

Proof. For an equation the first two statements are the endpoint equations of C, reflexivity and transport. For $D_{x,y}B$ the first is Case with the gate theorem of the matrix, the assumed base being unused, and the second is binder congruence (Theorem 9.52). For $L(F)$: forward, Case uses F to prove the successor gate; backward, DE, the theorem G_0F and Use at 1 give G_1F and hence F. For M: at the equations the endpoint equations of C give the two statements for the padded formulas, binder congruence carries them through the common binders, and $L(F) \dashv\vdash F$ removes the padding. □

Lemma 9.61 (Natural deduction)**.** *In $\mathcal{L}$ the macros* (9.19) *admit every introduction and elimination rule of intuitionistic first-order natural deduction, with the usual discharge and eigenvariable conditions, and ex falso for $\bot_*$. Moreover $\mathcal{L}$ proves coverage $n = 0 \vee_* \exists_* m\,(n = Sm)$.*

Proof. Implication. Given a derivation of B under the assumption A, Case introduces $A \to_* B$: the assumed zero instance gives A and the successor instance is equivalent to B. For elimination, DE retains the selector clause, the supplied A proves its antecedent, and Use at 1 gives B.

Conjunction. From A and B, numerical cases prove $M_v(A,B)$ for every term v, in particular for $v = P(Sz)$. The successor gate and Case give $A \wedge_* B$; the antecedent of its clause is

a gate theorem. For elimination, DE and Use at 1 and at $S1$ give the selector at $P(1) = 0$ and at $P(S1) = 1$, hence A and B by transport. No parity or range theorem is used.

Existential quantifier. From $A(t)$, Case at the witness t introduces the gated formula. Conversely DE at a fresh b, the zero gate and Use at 1 give $A(b)$; cut this into a branch whose conclusion and context do not contain b, and close DE.

Disjunction. The witnesses 0 and 1 give the two injections. For elimination, discharge the two branches as $A \to_* F$ and $B \to_* F$. Numerical cases, in the original context, prove $M_v(A, B) \to_* F$ for every v. Existential elimination supplies $M_b(A, B)$ at a fresh tag b, and implication elimination gives F. The tag is not substituted into a retained assumption, since the numerical cases are performed before that assumption is added.

Universal quantifier. A derivation of $A(z)$ with z fresh gives $G_{Sz}A(P(Sz))$ by $P(Sz) = z$ and the gate equivalence, and Case with an unused witness gives $\forall_* x\, A(x)$. For elimination, DE, the zero gate and Use at St give $G_{St}A(P(St))$ and hence $A(t)$. Every numerical target is covered directly; the value $P(0) = 1$ plays no role, because the outer gate makes the zero instance provable.

Falsity. Under $0 = 1$, transport in the control of $C(v, s, t)$ and the two endpoint equations give $s = t$ for arbitrary terms. Induction on the number of binders, with Case, gives every formula.

Coverage. Numerical cases on n: the left injection at 0, and the right injection with the witness z at Sz. □

Remark 9.62 (Variants of the macros). The form (9.19) is that of `uniform_binding` and `single_constructor`. `counter_evaluation` uses a term π with $\pi(1) = 1$ and $\pi(S1) = 0$ in the conjunction and defines $\forall_* x\, A(x) := A(0) \wedge_* D_{w,x}A(x)$. `uniform_recursion` uses $K(y) = C(P(y), 1, 0)$ in the conjunction and the form above for the universal quantifier. `clausal_canonical`, which has no decoder, uses $p(z) = \widehat{I}(z, 0; u.C(u, 1, 0))$ with $p(1) = 1$, $p(S1) = 0$ and the universal quantifier with a separate zero instance. `single_constructor` defines M_v by recursion on the pair of formulas, padding an equation E against a paired formula by $G_y E$ under the aligned binder; the result is interderivable with the definition above.

9.6.4 Adequacy and the main theorem

Theorem 9.63 (Adequacy of $\mathcal{L}$)**.** *Let $A \mapsto A^*$ translate the formulas of T^2_R to the clause syntax: an equation is normalized, $\bot$ becomes $\bot_*$, and the connectives and quantifiers become the macros* (9.19)*.*

1. *If $\Gamma \vdash_{\mathcal{L}} A$ then T^2_R proves A° from the readings of the clauses of Γ.*
2. *If T^2_R proves B from assumptions Δ, then $\Delta^* \vdash_{\mathcal{L}} B^*$, the formulas of Δ^* being ordinary assumptions.*
3. *$\mathsf{T}^2_R \vdash (B^*)^\circ \leftrightarrow B$ for every formula B of T^2_R, and $(A^\circ)^* \dashv\vdash_{\mathcal{L}} A$ for every formula A of the clause syntax, under arbitrary retained clauses.*

Hence $\mathcal{L}$ is faithful for T^2_R in both directions, and both numerical maps of this comparison are the identity.

Proof. (1) Induct on the derivation. By Theorem 9.23 a normalized instance is provably equivalent to the raw instance. Refl, Comp and Tr are equality reasoning and a computation equation. Use is implication and universal elimination, and DE is existential elimination. For Case, assume $B(t, 0)$; coverage and the premise give $B(t, y)$ for every y; then introduce the witness t. For Prog, assume $E(t, 0)$ and apply (9.11) with the premise as step.

(2) The logical rules are Theorem 9.61 and the equality rules are Theorem 9.52. The translation of the zero computation has the same canonical term on both sides, and that of the successor computation is Comp. Coverage is Theorem 9.61. Separation translates to $(0 = 1) \to_* (0 = 1)$. The rule (9.11) is Theorem 9.52(3), with the unchanged context.

(3) For the first statement induct on B: in T^2_R the gate and selector endpoints, coverage for an arbitrary control and $P(St) = t$ show that the reading of each macro is equivalent to the connective or quantifier it represents; for example $\forall y\,(M_y(A, B))^\circ$ is equivalent to $A^\circ \wedge B^\circ$ by coverage, so the reading of $A \to_* B$ is equivalent to $A^\circ \to B^\circ$. For the second induct on A. Equations are unchanged. For $D_{x,y}B$ apply the induction hypothesis to the matrix, uniformly, and prove

$$D_{x,y}B(x, y) \dashv\vdash \exists_* x(B(x, 0) \to_* \forall_* y\, B(x, y)).$$

Forward, open the left side at b; under $B(b, 0)$ its clause gives $B(b, z)$ for a fresh z; universal, implication and existential introduction build the right side; close DE. Backward, existential elimination gives a fresh b and the implication; introduce $D_{x,y}B$ by Case at b: the local base feeds the implication, and universal elimination at Sz gives the successor instance. Faithfulness follows: if T^2_R proves A° from the readings of Γ, then by (2) and (3) $\mathcal{L}$ derives A from Γ, each clause $[A_0;\, y.\, B_0]$ providing the translation of its reading by implication and universal introduction. □

Theorem 9.64 (Counter evaluation)**.** *The five-pattern calculus $\mathcal{K}_{\mathrm{CE}}$ is sound and faithful for T^2_R, and it is equiconsistent and bi-interpretable with* iPRA*: both numerical domains are the whole domain, equality is actual equality, the numerical composites have the graph $x = y$, and every formula is provably equivalent to its double translation. The translations preserve the full consequence relation, including open sequents and formulas with unbounded quantifiers.*

Proof. Combine Theorems 9.25, 9.58 and 9.63. The contradiction $0 = 1$ reads as an equation that T^2_R refutes by separation and that implies every formula, so inconsistency transfers in both directions. □

Remark 9.65. `counter_evaluation` and `uniform_binding` present the same calculus. `counter_evaluation` contributes Theorems 9.54 to 9.56 and a checker that synthesizes the conclusion of each proof node; its ordinary theory has the zero computation as an identity of canonical spelling. `uniform_binding` takes the ordinary theory over raw terms with both computation equations, reads canonical terms as raw ones, justifies the normalization by Theorem 9.23, and contributes the macros (9.19) and the goal-directed checker with one traversal (Section 9.8). The equivalence of the two certificate formats is Theorem 9.84.

9.7 Three families on normal forms

This section draws on `uniform_recursion` and `single_constructor`.

On normal terms (Theorem 9.5) both computation equations are identities of spelling, and the equations of Theorem 9.10 are literal identities of normal forms. A calculus then needs neither a computation rule nor, with a suitable elimination, a transport rule. Two such calculi are stated on the counter-aware recursor. Both take the clause syntax of Section 9.6 with empty antecedents, and both contain in every context the fixed clause

$$\mathcal{A} := [\,;\, y.\, y = y], \tag{9.20}$$

which is the initial context of a closed proof. Reflexivity is thus an axiom clause; it has not been deleted without replacement. An ordinary assumption F is $[\,;\, v.\, F]$. On iterator terms in

full normal form, the two calculi are the clause calculi of the packages `witness_discharge` and `inferred_witness`, which are formalized as `WDPrf` (`wdprf_iff_nprf`) and `OCPrf` (`ocprf_iff_nprf`), with the normalizer `Tm.full`; the counter-aware forms below are not formalized. What the new syntax requires is proved here: the packets that carry transport use the decoder, and the repair of the introduction at the witness 0 cannot use a closed term that is provably but not literally 0, because normal terms have none.

9.7.1 Use, Close and Name

Definition 9.66 (The calculus $\mathcal{K}_{\mathrm{Name}}$)**.** Terms are normal and all displayed instances are normalized by nf. The rules are:

1. Use: for $[A;\, y.\, B(y)] \in \Gamma$ and a term t, from $\Gamma \vdash A$ infer $\Gamma \vdash B(t)$; for an empty antecedent there is no premise. In particular $\mathcal{A}$ gives $t = t$.

2. Close: the rule (DI), its local assumption being an unconditional constant clause.

3. Name: for a fresh parameter b and the clause $\mathcal{B}_b := [B(b,0);\, y.\, B(b,y)]$,

$$\frac{\Gamma \vdash D_{x,y}B \qquad \Gamma, \mathcal{B}_b \vdash b = t \qquad \Gamma, \mathcal{B}_b \vdash F}{\Gamma \vdash F\langle t/b\rangle} \qquad b \notin \mathrm{FV}(\Gamma, F\langle t/b\rangle). \tag{9.21}$$

   The term t and the formula F may contain b. The substitution replaces the free occurrences of b once and does not revisit an inserted copy of t. The case $t \equiv b$ is permitted.

There is no transport rule, no hypothesis rule apart from Use, no computation inference, no successor axiom and no separate induction rule. Weakening, exchange, contraction, injective renaming of parameters and cut of an ordinary assumption are admissible by induction on derivations; these operations preserve the test of Close.

Lemma 9.67 (Elimination, closure and cases)**.** *In $\mathcal{K}_{\mathrm{Name}}$:*

1. *from $\Gamma \vdash D_{x,y}B$ and $\Gamma, \mathcal{B}_b \vdash A$ with $b \notin \mathrm{FV}(\Gamma, A)$ one infers $\Gamma \vdash A$;*

2. *writing $q_y A(y) := D_{x,y}A(y)$ for an unused x: from $\Gamma, A(0) \vdash A(Sz)$ one infers $\Gamma \vdash q_y A$, and from $q_y A$ and $A(0)$ one infers $A(t)$;*

3. *numerical cases: from $\Gamma \vdash A(0)$ and $\Gamma \vdash A(Sz)$, z fresh, one infers $\Gamma \vdash A(t)$.*

Proof. (1) In (9.21) take $t \equiv b$. The equality premise is $b = b$, from $\mathcal{A}$, and the export is literally A. (2) Introduce the unused witness at 1, which is not literally 0, so that Close is in case mode. For the elimination open the paired formula by (1); its clause is $[A(0);\, y.\, A(y)]$; use it with the supplied base at t. (3) Weaken the successor proof by the unused base assumption and compose the two steps of (2). □

Lemma 9.68 (Transport from one discharge)**.** *For every term template $h(w)$, from $\Gamma \vdash s = t$ and $\Gamma \vdash h(s) = r$ one infers $\Gamma \vdash h(t) = r$ in $\mathcal{K}_{\mathrm{Name}}$. Hence symmetry, transitivity, congruence and the crossed transport Tr are derivable.*

Proof. Let $K(y) := C(P(y), 1, 0)$, so that $K(1) \equiv 1$ and $K(S1) \equiv 0$ as normal forms. Choose x, y fresh and put

$$B(x,y) := \Big(C(y, 0, C(K(y), x, h(x))) = C(y, 0, C(K(y), t, r))\Big).$$

Its normalized rows are $B(x,0) \equiv (0 = 0)$, $B(x,1) \equiv (h(x) = r)$ and $B(x, S1) \equiv (x = t)$, and its row at Sz is $C(K(Sz), x, h(x)) = C(K(Sz), t, r)$. To introduce $D_{x,y}B$ at the witness

s, apply numerical cases to the equation $C(v,s,h(s)) = C(v,t,r)$ and instantiate it at $v = K(Sz)$: its zero case is $s = t$ and its successor case is $h(s) = r$. This derivation ignores the local assumption of Close and is valid in either mode, including $s \equiv 0$. Now apply (9.21) with a fresh b. In the opened context the antecedent $0 = 0$ holds by $\mathcal{A}$; Use at $S1$ gives the equality premise $b = t$, and Use at 1 gives the formula premise $h(b) = r$. The export is $h(t) = r$ by (9.7); the passive data do not contain b.

Symmetry is the case $h(w) = w$, $r \equiv s$ with the reflexive minor premise. Congruence uses a reflexive minor premise, and transitivity and the crossed form follow. □

The argument is not circular: ordinary elimination, closure and numerical cases (Theorem 9.67) are available before any equality rule, and the packet needs only the literal endpoint identities and no range theorem for K.

Lemma 9.69 (Formula transport)**.** *From $s = t$ and $F(s)$ one derives $F(t)$ in $\mathcal{K}_{\mathrm{Name}}$, for every formula template F.*

Proof. Induct on the number of paired binders. The atomic case is two applications of Theorem 9.68. For $F(v) = D_{x,y}B(v,x,y)$, eliminate $F(s)$ at a fresh c and introduce $F(t)$ at the witness c, which is not literally 0, so that Close is in case mode. Under $B(t,c,0)$ the induction hypothesis and symmetry give $B(s,c,0)$; use the old clause at Sz and transport the result to $B(t,c,Sz)$. □

Lemma 9.70 (The introduction at the witness 0)**.** *For an equation $E = (s = t)$ put $G_vE := (C(v,0,s) = C(v,0,t))$ and $L(E) := q_vG_vE$. Then $G_0E \equiv (0 = 0)$ and $G_{Sw}E \equiv E$ literally, and $E \dashv\vdash L(E)$ in every context. Consequently Case is derivable in $\mathcal{K}_{\mathrm{Name}}$ for every matrix and witness, and so are Prog and induction on literal equations.*

Proof. The interderivability is closure introduction and closure elimination at 1 (Theorem 9.67), without transport. Let $\Gamma, E(0,0) \vdash E(0,Sz)$ be a case premise for an equation matrix at the witness 0, the only instance of Case that Close does not provide. Consider the compound matrix $A(x,y) := L(E(x,y))$. Unwrapping the base, applying the premise and wrapping the result gives $\Gamma, A(0,0) \vdash A(0,Sz)$, and Close at the witness 0 is in case mode because A is compound; it proves $D_{x,y}A$. Open this formula at a fresh c (Theorem 9.67). Introduce $D_{x,y}E$ at the witness c, which is not literally 0: wrap the assumption $E(c,0)$, use the opened clause at Sz, and unwrap. The conclusion does not contain c and is exported.

For Prog at a witness t, Close in progress mode at the witness 0 with a matrix independent of the witness coordinate gives $q_yE(t,y)$; under $E(t,0)$ its clause gives $E(t,Sz)$, and Case at t concludes. Induction is the same closure used with a supplied base. □

Theorem 9.71. *On normal terms, with $\mathcal{A}$ in every context, $\mathcal{K}_{\mathrm{Name}}$ and $\mathcal{L}$ derive the same sequents. In particular numerical substitution is admissible in $\mathcal{K}_{\mathrm{Name}}$.*

Proof. Refl is Use of $\mathcal{A}$, Comp is an instance of it on normal terms, and Tr, DE, Case and Prog are Theorems 9.67, 9.68 and 9.70. Conversely Close is an instance of Case or Prog. For (9.21), apply DE; in the opened context derive F and $b = t$; transport in the template obtained from F by replacing the free occurrences of b by a new variable (Theorem 9.52) gives $F\langle t/b\rangle$, which does not contain b; close DE. □

9.7.2 Use, Introduce and Open with an inferred witness

Definition 9.72 (The calculus $\mathcal{K}_{\mathrm{Open}}$)**.** Terms are normal and all displayed instances are normalized. The rules are Use and the introduction (DI), as in Theorem 9.66, and the

following elimination. For a major premise $D_{x,y}B(x,y)$ and a parameter b fresh for Γ and the parameters of B, let $B_b(y)$ be the normalized instance $B(b,y)$ and

$$\tau_B(b) := \begin{cases} r, & B_b(0) \equiv (0=0) \text{ and } B_b(1) \equiv (b=r), \\ b, & \text{otherwise.} \end{cases} \tag{9.22}$$

The variable b has to be the entire left side of the row at 1; no search for a provable equality takes place. The rule is

$$\frac{\Gamma \vdash D_{x,y}B \qquad \Gamma,\ [B(\tau_B(b),0);\ y.\, B(\tau_B(b),y)] \vdash A}{\Gamma \vdash A} \qquad b \notin \mathrm{FV}(A). \tag{Open}$$

The branch is checked in the scope that contains b, also when the inferred term contains b. The replacement of b by $\tau_B(b)$ is simultaneous and occurs once.

The rule (Open) is sound for the ordinary reading. Open the existential statement at b, which gives $B(b,0) \to \forall y\, B(b,y)$. If the probe succeeds, reflexivity supplies $B(b,0)$ and the instance at 1 is $b = \tau_B(b)$; if it fails, this equation is $b = b$. Replacement of b by an equal term converts the opened clause to the clause of the rule, and the branch gives A, which does not contain b.

Weakening, injective renaming of parameters and cut of an ordinary assumption are admissible in $\mathcal{K}_{\mathrm{Open}}$ by induction on derivations; injective renaming preserves the literal tests.

Lemma 9.73 (Safe openings and the repair of cases)**.** *In $\mathcal{K}_{\mathrm{Open}}$:*

1. *if the witness coordinate is unused in B, or the root of B is compound, then $\tau_B(b) \equiv b$ and* (Open) *is the ordinary elimination DE;*
2. *the closure rules and numerical cases of Theorem 9.67(2),(3) hold;*
3. *Case is derivable for every matrix and witness.*

None of these uses a substitution theorem or an equality rule.

Proof. (1) If x is unused, the row at 1 does not contain the fresh b, so its left side is not b. If the root of B is compound, the row at 0 is compound and is not $0 = 0$. (2) Introduce $q_y A$ at the unused witness 1, in case mode; its opening is safe by (1). (3) The construction of Theorem 9.70 applies: the interderivability of E and $L(E)$ uses closures only, the formula $D_{x,y}L(E(x,y))$ has a compound matrix and is opened safely at a fresh c, and the final introduction at the witness c is in case mode. □

Lemma 9.74 (Substitution)**.** *If $\Gamma \vdash A$ in $\mathcal{K}_{\mathrm{Open}}$, then $\Gamma\langle\sigma\rangle \vdash A\langle\sigma\rangle$ for every simultaneous numerical substitution σ.*

Proof. Use induction on the original derivation, with the assertion quantified over all substitutions, after freshening the eigenparameters away from the images of σ. Use follows from (9.7). An introduction in progress mode remains in progress mode. An introduction in case mode remains a case introduction unless its witness becomes 0 under an equation matrix; then use Theorem 9.73(3).

At an instance of (Open) let B' be the normalized instance of B under σ and $\tau' = \tau_{B'}(b)$. If the old probe succeeded at r, its rows $0 = 0$ and $b = r$ remain rows of this form under σ with b fixed, so the new probe succeeds at $r\langle\sigma\rangle$, and the induction hypothesis for the branch with b fixed gives the new branch. If the old probe failed, the old branch has the clause at b, and the new probe may succeed. Apply the induction hypothesis to the original branch with the substitution that extends σ by $b \mapsto \tau'$. It produces exactly the new opened clause; the context and the conclusion did not contain b and are not affected by the extra component; and an inserted image is not substituted into again. The induction is on a proper subderivation of the original derivation, so no transformed and possibly larger proof is used circularly. □

Lemma 9.75 (Ordinary elimination, transport and induction)**.** *In $\mathcal{K}_{\mathrm{Open}}$:*

1. *DE is admissible;*
2. *for every term template $h(w)$, from $s = t$ and $h(s) = r$ one infers $h(t) = r$; hence symmetry, transitivity, congruence and transport in every formula template are derivable;*
3. *Prog and induction on literal equations are derivable.*

Proof. (1) Given a branch under the clause at b, apply Theorem 9.74 to it with $b \mapsto \tau_B(b)$ and then (Open); the context and the conclusion do not contain b. This is why the substitution lemma precedes unrestricted opening.

(2) Use the matrix

$$B(x, y) := \Big(C(y, 0, C(P(y), x, h(x))) = C(y, 0, C(P(y), t, r))\Big). \tag{9.23}$$

Its normalized rows are $B(x, 0) \equiv (0 = 0)$, $B(x, 1) \equiv (x = t)$ and $B(x, S1) \equiv (h(x) = r)$, and its row at Sz is literally $C(z, x, h(x)) = C(z, t, r)$, because $P(Sz)$ computes to z. At the witness s, numerical cases on z prove this row: the zero case is $s = t$ and the successor case is $h(s) = r$. The derivation ignores the local assumption of the introduction, so it is valid in either mode and proves $D_{x,y}B$. Now (Open) infers the witness t: the row at 0 is $0 = 0$ and the row at 1 is $b = t$. The opened clause has the base $0 = 0$, which $\mathcal{A}$ proves, and Use at $S1$ gives $h(t) = r$, which is in the outer scope. The remaining statements follow as in Theorems 9.68 and 9.69, with (1) for the eliminations.

(3) The introduction at the witness 0 with an unused witness coordinate gives $q_y E(y)$ from $\Gamma, E(z) \vdash E(Sz)$; its opening is safe, and Use with the base gives $E(t)$. Prog at a witness t follows with Case. □

The decoder makes the packet (9.23) simpler than the packet of Theorem 9.68 and than the parity packet of the calculus on iterator terms: one successor computation exposes the variable on which the numerical cases are taken.

Theorem 9.76. *On normal terms, with $\mathcal{A}$ in every context, $\mathcal{K}_{\mathrm{Open}}$ and $\mathcal{L}$ derive the same sequents.*

Proof. The rules of $\mathcal{L}$ are derivable by Theorems 9.73 and 9.75. Conversely the introduction is an instance of Case or Prog. For (Open) with a successful probe at r, apply DE at b; the opened clause has the antecedent $0 = 0$, and Use at 1 gives $b = r$. For a fresh y, Use gives $B(b, y)$ and transport along $b = r$ gives $B(r, y)$; hence the clause $[B(r, 0);\, y.\, B(r, y)]$ of the branch can be cut (Theorem 9.51), and the branch gives A. With a failed probe the rule is DE. □

9.7.3 The scalar normal forms embed exactly

The calculus of `inferred_witness` is $\mathcal{K}_{\mathrm{Open}}$ on iterator terms $0, S, I$ in full normal form: a normal iterator term $I(n, a;\, v.\, h)$ has a neutral counter, and the normalizer nf_I applies both computation equations of Section 1.1. Let T_{raw} translate variables and 0 identically, the successor by the defined successor, and $I(n, a;\, v.\, h)$ by $\mathrm{Rec}(n, a; i, v.h)$ with i fresh, and for a normal scalar term m put $T(m) := \mathrm{nf}(T_{\mathrm{raw}}(m))$.

Lemma 9.77 (Exact embedding of normal forms)**.** *For every scalar term t, $\mathrm{nf}(T_{\mathrm{raw}}(t)) = T(\mathrm{nf}_I(t))$. The raw translation commutes with substitution and T commutes with normalized substitution. On normal scalar terms T preserves and reflects the literal 0, each variable and the property of being a successor, and it preserves the set of free variables.*

Proof. The two scalar computation equations translate to the computations of Theorem 9.10, so $T_{\mathrm{raw}}(t)$ and $T_{\mathrm{raw}}(\mathrm{nf}_I(t))$ have the same normal form by Theorem 9.7. For a normal scalar term describe T by induction. Zero, variables and successors are immediate. A normal iterator has a neutral counter, and its image is

$$P(R(Tn, Ta; i, u.S((Th)[P(u)/v]))).$$

Here Tn is neutral, so the inner recursor is retained and neutral, and the outer P is retained. Substituting the neutral term $P(u)$ for the body variable does not turn a neutral counter into 0 or a successor, so the whole image is normal. The raw translation commutes with substitution by induction on terms, with the two new variables fresh at every translated iterator, and (9.7) gives the normalized statement. The four kinds of normal source term have distinct kinds of image: 0, the same variable, a canonical successor, a neutral recursor. No free variable of a normal source term is erased. □

Proposition 9.78 (Translation of the scalar calculus)**.** *Extended to formulas without change of roots, T maps every instance of* Use*, of the introduction and of* (Open) *of the normalized scalar calculus to an instance of the same rule of $\mathcal{K}_{\mathrm{Open}}$.*

Proof. For Use this is commutation with normalized substitution. For the introduction, reflection of the literal 0 preserves the mode, and the translated successor row is the required successor row. For (Open) the two tests on rows are preserved and reflected, so $\tau_{T(B)}(b) = T(\tau_B(b))$, and normalized substitution gives the translated branch clause. Preservation of free variables gives the scope conditions. □

The translation concerns derivations of the already normalized scalar calculus. If raw scalar proofs are normalized first, a witness can become 0; that separate step is handled by Theorem 9.73(3). The embedding preserves more than standard numerical values: a row becomes $0 = 0$, or has a given fresh variable as its entire left side, exactly when the source row did.

9.7.4 The main theorem for the two calculi

Theorem 9.79. *$\mathcal{K}_{\mathrm{Name}}$ and $\mathcal{K}_{\mathrm{Open}}$ are sound and faithful for T^2_R and are equiconsistent and bi-interpretable with* iPRA*, with whole domains, actual equality, numerical composites of graph $x = y$, and every formula provably equivalent to its double translation. Every fixed primitive recursive description compiles to a term whose defining equations hold by computation (Theorem 9.12).*

Proof. By Theorems 9.71 and 9.76 both calculi are $\mathcal{L}$ on normal terms, and Theorems 9.25 and 9.63 apply. In particular $\mathcal{K} \vdash 0 = 1$ if and only if $\mathsf{iPRA} \vdash \bot$; this is a metatheorem and not an assertion that either system proves its own consistency. □

On normal terms the coverage theorem of Theorem 9.61 strengthens to $t = 0 \vee_* t = S(p(t))$, since $p(Sz) \equiv z$; this is the form used by `uniform_recursion` and `single_constructor`.

9.8 Uniform binding and one traversal

This section draws on `clausal_canonical`, `counter_evaluation`, `uniform_binding`, `uniform_recursion` and `single_constructor`, and for the separate refinements on `canonical_recursors` and `register_graph`.

9.8.1 The concrete syntax

In the checkers of these packages the integer 0 is the numerical constant, a negative integer is a free parameter (a level), and a positive integer is a de Bruijn index of a bound variable. Without a successor constructor and without a closure constructor, the array arities distinguish the constructors:

Object	Array	Child depths, one binder in R	Child depths, two binders in R
$R(n, a; \vec{u}.h)$	$[n, a, h]$	$(d, d, d+1)$	$(d, d, d+2)$
$s = t$	$[s, t]$	(d, d)	(d, d)
$D_{x,y}B$	$[B]$	$(d+2)$	$(d+2)$

In a paired matrix y has index 1 and x index 2; in the body of the counter-aware recursor the state has index 1 and the counter index 2. With two binders in R, every constructor other than the equation binds exactly two variables in its last child, so that for child number j of a node of arity k the depth increment is

$$2\,[\,k \neq 2 \text{ and } j = k - 1\,]. \tag{9.24}$$

The sort is still checked: formula arities are 1 and 2, and the only arity of a compound term is 3. The arity convention does not identify a formula with a term. A successor is $[0, a, 0]$, and a compound canonical counter is a successor exactly when its first child is 0; the test that distinguishes a successor node from an iterator node on the terms of Section 1.1 is not needed.

A context clause is a pair (A, B) whose second component has one more bound variable. It is a separate data structure and is never passed to the opening operation as one object; clause use opens B alone.

Definition 9.80 (Opening)**.** For an expression e and a term v, the operation $O_d(e, v)$ removes the index $d + 1$ and preserves the d nearer indices: on an integer it replaces $d + 1$ by v, decrements an integer greater than $d + 1$ and leaves every other integer unchanged; on an array it processes the children at the depths of the table; and at each resulting recursor node it erases the body if the resulting counter is 0. O_d^{raw} is the same operation without the erasure.

Lemma 9.81 (Opening)**.** *Let e have free scope m and ambient depth $k > d$, and let v be a term of free scope m with no free positive index.*

1. *$O_d(e, v)$ has the sort of e, ambient depth $k-1$ and free scope m; it contains no parameter beyond those of e and v; and it is zero-body canonical if e and v are.*

2. *For canonical e, v, $O_d(e, v) = \nu(O_d^{\mathrm{raw}}(e, v))$, and for arbitrary raw e, v, $\nu(O_d^{\mathrm{raw}}(\nu e, \nu v)) = \nu(O_d^{\mathrm{raw}}(e, v))$.*

3. *For a matrix B and terms t, w without free positive indices, $O_0(O_1(B, t), w) = \nu(B[t/x, w/y])$, and the two openings may be performed in the opposite order.*

Proof. (1) Induct on e. The integer cases remove the selected index, lower an outer index, or keep a nearer index or a parameter. A replacement without free positive indices stays well scoped under any additional binders. The array cases use the binder increments of the table. Erasing a body removes occurrences only. (2) is Theorem 9.3 for the substitution that opening performs: a body discarded before the opening had a counter that was literally 0 and stays 0; a counter that becomes 0 only by the opening is erased on both sides by the outer ν; and $\nu(t)$ is 0 only if t is. (3) follows from the simultaneous-substitution lemma, since neither replacement contains a free positive index, and from (2). □

The hypothesis on v is essential. Because free parameters are negative levels and do not depend on the binder depth, inserting such a v under binders needs no lifting; inserting a term with a free positive index without lifting is incorrect. For the same reason an old context needs no shifting when a fresh parameter is allocated: the new negative integer is absent from everything validated before. A branch can use it, but its discharged result cannot contain it, because the outer goal and the witness are validated at the old scope.

Remark 9.82 (Unfolding under binders). With full normalization a successor contraction can occur under binders, and the inserted terms can then have free positive indices. The simultaneous substitution of (9.4) is implemented counter first: remove the counter slot at depth 1, under the retained state slot and with lifting of the inserted term, and then remove the state slot. Substituting the previous state first without lifting captures an ambient index under the counter binder that is still present. For example, under one ambient binder $R(1, v; j, w.w)$ normalizes to Sv, that is $[[0, 0, 0], 1, 1]$ normalizes to $[0, 1, 0]$, whereas the state-first order returns $S0$. This correction is recorded in the mathematical review shipped with `uniform_recursion`.

9.8.2 One traversal

Lemma 9.83 (The shared walker). *Let* walk *visit a tree with the depth increments* (9.24)*, check the sort and arity of every node, apply a given operation to the numerical leaves, and erase the body at every recursor node whose resulting counter is* 0*.*

1. *With the leaf operation that checks $-n \le i \le d$ for an exact integer i at scope n and depth d and returns i, the test that the result equals the input holds exactly for the members of the scoped zero-body canonical grammar.*

2. *With the leaf operation that maps $d + 1$ to v, an index $i > d + 1$ to $i - 1$ and every other leaf to itself, and with v canonical without free positive indices, the result is the canonical capture-avoiding opening O_d.*

Proof. Induct on the tree. The integer clause checks the sort and the scope interval. At an array the admissible arity determines whether the children are terms or formulas, and the last child of a recursor or a paired formula receives two more protected indices. Every child is visited before the erasure, so a malformed body is rejected before it can be discarded. In (1), equality with the returned tree excludes every zero-counter node with a nonzero body. In (2) the three index cases are opening of one slot, and the final erasure is that of O_d. □

Thus one traversal owns grammar validation, the canonical-form check and opening. For a certificate that starts from the empty context every branch of an elimination or introduction adds exactly one clause and one fresh parameter, and no other rule adds either; by induction on checker calls the scope equals the length of the context, so no separate scope counter is passed. This is an invariant of certificates built from the empty root, not a statement about abstract clause contexts. In the rule (CE) the checker opens the state slot with the canonical term $R(t, a; i, u.h)$ and then the former counter slot with t, which is the simultaneous substitution; the opening is computed before the two premises are checked, and this does not bypass validation, because proving the major premise validates both endpoints and proving the minor premise validates the seed and the original body, whose outer counter is a successor and cannot have been erased.

Proposition 9.84 (Checker correspondence). *Disregarding physical resource bounds:*

1. *A closed canonical formula has a certificate accepted by the goal-directed checker of* `uniform_binding` *if and only if it is a theorem of $\mathcal{K}_{\mathrm{CE}}$, and accepted certificates translate effectively to and from the certificates of the conclusion-synthesizing checker of* `counter_evaluation`*, with literally identical final formulas.*

2. *The checker of* `uniform_recursion` *accepts a certificate for a closed normal goal exactly when the certificate describes a derivation in* $\mathcal{K}_{\mathrm{Name}}$*, and every finite derivation has an accepted certificate; the same holds for* `single_constructor` *and* $\mathcal{K}_{\mathrm{Open}}$.

Proof. (1) Use the scope invariant and Theorem 9.83. An accepted reflexive goal is a Refl node, and every other accepted branch is the displayed instance of (CE), Use, DE or (DI) after translation of its premises; this is soundness, by induction on the accepted call tree. Conversely induct on a derivation: insert the derived endpoints and the seed and body of the minor premise as the hints of a (CE) node, copy the clause index and target of Use, the matrix of DE and the witness of the introduction. The opening invariant identifies every required premise goal, and each construction preserves the end formula.

(2) Induct on the proof packets. The formula synthesized at each node is checked for grammar, normal form and scope. In Use the term is checked before instantiation and a proof of the antecedent has to synthesize exactly the retained antecedent. In the introduction the matrix and the witness are checked, the generated assumption is the row selected by the rule, and the result of the branch has to equal the generated successor row. In Name the major premise has to be a paired formula, the equality has to have the freshly allocated parameter as its left endpoint, and the final scope check excludes that parameter from the normalized export. In (Open) the major premise is checked before the context is extended, the probe (9.22) is computed, and the conclusion of the branch is validated again in the outer scope; when the probe succeeds the stored base is $0 = 0$ and coincides with the substituted base of the rule. Theorem 9.7 and Theorem 9.82 identify the computed instances with the normalized substitutions. Conversely a derivation is encoded by its rule data and the certificates of its premises, with fresh negative levels in order of scope creation. □

9.8.3 The checkers

All checkers are standalone Python modules that read one JSON object with exactly the keys `goal` and `proof`, validate a closed goal against the full grammar, and accept by printing `ACCEPT` with exit status zero; none imports a proof producer or a theorem table. Exact integer tests exclude Booleans, and recursion and memory limits of the interpreter can reject a large valid certificate. Sizes are in Section 9.10.

`register_graph`. The checker implements the natural deduction of Theorem 9.13 with named variables, simultaneous substitution, alpha equivalence, the three arithmetic schemas and (9.11). It is a readable reference and is much larger than the compact checkers of the kernel. A separate ancillary module is a profile of the kernel `N` on iterator terms, not a checker for T_J: it replaces a recursive continuation of its parent by a loop and uses immutable tuple contexts, which lowers the node count of the inspected leader by one and raises its bytes. Its generic opener inherits a qualification on malformed input: in an interpreter without limits certain nonempty strings cause unbounded traversal, so soundness concerns finite successful executions.

`canonical_recursors`. Three modules share one proof-tree format, context handling, de Bruijn convention and driver: iterator terms with a successor and seven rules, raw seeded terms with seven rules, and $\mathcal{K}_6$. The canonical module applies the constructor $[n, a, 0 \text{ if } n = 0 \text{ else } h]$ at every newly built recursor node after its children have been normalized, validates canonical input and has no zero-computation rule. The scalar control uses crossed eliminations and the witness-directed introduction and is an equivalent presentation of the formalized seven-rule calculus named in Section 9.4, not a transcription of it.

`clausal_canonical`. The checker for $\mathcal{K}_5$ has the instruction tags `r`, `x`, `u`, `e`, `i` and one opening function for terms and formulas. Its matched control is $\mathcal{K}_6$ ported to the same variable convention and opening function, with the closure constructor, the restricted transport and

the root-reversing closure elimination retained, so that the comparison isolates the change of rules and formulas from the change of representation.

counter_evaluation. The checker for $\mathcal{K}_{\mathrm{CE}}$ has the tags `r` (reflexivity), `v` (counter evaluation), `u`, `e`, `i` and synthesizes the conclusion of every node. Its matched control is $\mathcal{K}_{\mathrm{CT}}$ on the same syntax. An untrusted translator implements Theorem 9.56: a computation node becomes (CE) on two reflexive leaves and a transport node becomes the construction of Theorem 9.55.

uniform_binding. The checker for $\mathcal{K}_{\mathrm{CE}}$ is goal directed: a validated reflexive goal closes at once and its payload is ignored, and the other instructions are `v`, `u`, `e`, `i` with hints. Validation and opening are the two uses of Theorem 9.83, and the scope is the length of the context. Its matched control applies the same organization to iterator terms.

uniform_recursion. The checker for $\mathcal{K}_{\mathrm{Name}}$ has the packets `u`, `i`, `e`; the fixed clause (9.20) is the initial context $([\,],[1,1])$, so that at context length n the parameters are $-1,\dots,-(n-1)$ and the next one is $-n$. The minimal proof is `{"goal":[0,0],"proof":["u",0,0,null]}`. A variant carries the budget of Theorem 9.9; its control is the scalar control with the same budget instrumentation.

single_constructor. The checker for $\mathcal{K}_{\mathrm{Open}}$ has the packets `u`, `i`, `e`, with the same three logical tests as its control on iterator terms; the changes are confined to the arithmetic syntax, the shared binder increment, the recursor computation and the successor abbreviation in the rules.

9.8.4 Three refinements of checkers for the kernel on iterator terms

Three packages of this chapter also contain refinements of existing checkers that do not concern the calculi of this chapter; no size of these programs is a size of a calculus on the new syntaxes.

`canonical_recursors` refines the compact guarded codec: it pairs the comparisons of the major premise and of the target that are already checked, uses a shorter discriminator between integers and lists, and composes the administrative truth convention of the scalar zero with the guarded checker. The byte and node improvements belong to different modules. The compact modules do not validate the public grammar themselves; a strict front end is counted separately. The eager paired comparison may inspect instruction fields that were unused before, so the completeness statement is made for the specified well-scoped language of integer and list certificates, and no equality of behavior on malformed JSON is claimed.

`uniform_recursion` fuses the grammar validation of the strict native checker of `deterministic_opening` into its first scope traversal. That walker increases its depth by $5-k$ at a node of arity k. At a formula position under p protected slots the depth is $2p$, and at a term position it is $3+2p$; hence even depth means formula and odd depth means term, and the admissible arities at depth d are exactly 1 and $2+(d \bmod 2)$. A numerical leaf is 0 or an odd integer at least 3 within the scope interval. The grammar predicate is enabled for the initial traversal without substitution image and disabled before any proof instruction is read. By induction on the tree the fused traversal succeeds exactly when the old grammar test and the old scope traversal both succeed, and it returns the same tree; every later transition of the checker is unchanged, so the translation of certificates in both directions is the identity. Thresholds of resource exhaustion and failure diagnostics are not equated. One variant states the predicate explicitly; a smaller one folds it into the existing cases.

`register_graph` supplies the ancillary profile described above.

9.9 Boundary results

This section draws on `canonical_recursors` and, for Theorem 9.93, on `clausal_canonical`.

The constructions above move a computation into the syntax or change the representation of the terms. The results of this section show that three nearby simplifications are not available: deleting the initializer together with the counter, deleting one of the two computation equations of the iterator, and restricting induction to equations with 0 on one side. A fourth shows that the single computation rule of $\mathcal{K}_5$ cannot be deleted without replacement.

9.9.1 A recursor with neither counter nor initializer

Consider the terms generated by variables, 0, S and a self-seeded constructor $F(n; v.h)$ with the standard interpretation

$$F(0; v.h) = S(h(0)), \qquad F(Sn; v.h) = h\big(F(n; v.h)\big),$$

with arbitrary nesting and passive parameters, and no further operation. The successor may be taken as the abbreviation $F(0; v.a)$, $v \notin \mathrm{FV}(a)$.

Theorem 9.85 (Unary terms of the self-seeded language)**.** *For every term $t(n)$ of this language with one free variable there is a natural number C such that*

$$\forall n \in \mathbb{N} \quad \big(t(n) \le C \ \vee \ t(n) \ge n\big).$$

Consequently no term defines the predecessor $\max(n-1, 0)$*, integer halving, or any function with unbounded values strictly below its argument.*

Let $\mathcal{U}$ be the class of functions $f\colon \mathbb{N} \to \mathbb{N}$ for which there is C with $f(n) \le C$ or $f(n) \ge n$ for every n.

Lemma 9.86 (Closure)**.** *Constants, the identity and the successor belong to $\mathcal{U}$. The class is closed under composition and under pointwise selection from a finite family of its members.*

Proof. For a pointwise selection take the maximum of the finitely many bounds. For $f \circ g$ with bounds C_f, C_g: if $g(n) \le C_g$ the value is at most $\max_{j \le C_g} f(j)$; otherwise $g(n) \ge n$, and the value is at most C_f or at least $g(n) \ge n$. □

Lemma 9.87 (Orbits)**.** *If $h \in \mathcal{U}$, then the sequence $r(0) = 1 + h(0)$, $r(n+1) = h(r(n))$ is a member of $\mathcal{U}$ as a function of n.*

Proof. If the orbit repeats a value it is eventually periodic and hence bounded. Otherwise it is injective. Let C be a bound for h. A step with $r(n+1) < r(n)$ ends in $\{0, \dots, C\}$, so an injective orbit has finitely many such steps. After the last one the orbit increases strictly, and it eventually exceeds every value of the earlier finite prefix. For all sufficiently large n the $n+1$ distinct values $r(0), \dots, r(n)$ lie in $[0, r(n)]$, so $r(n) \ge n$. Increase the constant to cover the finitely many earlier arguments. □

Proof of Theorem 9.85. Say that a function $g\colon \mathbb{N}^k \to \mathbb{N}$, $k \ge 1$, has *finite-control form* if there are a coordinate j, maps β_l from $\mathbb{N}$ into finite sets for $l = 1, \dots, k$, and functions $T_e \in \mathcal{U}$ indexed by the tuples e of feature values, such that

$$g(x_1, \dots, x_k) = T_{(\beta_1(x_1), \dots, \beta_k(x_k))}(x_j).$$

The coordinate j is called active. More generally the index may be any function of the feature values, and dummy coordinates may be added. A unary function of this form is a pointwise selection from finitely many members of $\mathcal{U}$ and belongs to $\mathcal{U}$ by Theorem 9.86. It therefore suffices to show that every term, as a function of any nonempty list of variables that contains its free variables, has finite-control form.

Composition. Let $g(m,\vec{x})$ and $n(\vec{x})$ have finite-control form, say $g(m,\vec{x}) = T_{(\beta(\vec{x}),\beta_m(m))}(y)$ with y the active coordinate of g, and $n(\vec{x}) = N_{\gamma(\vec{x})}(x_{j'})$. Then $g(n(\vec{x}),\vec{x})$ has finite-control form. Indeed the value $\beta_m(n(\vec{x})) = \beta_m(N_{\gamma(\vec{x})}(x_{j'}))$ is determined by $\gamma(\vec{x})$ and by the feature $x_{j'} \mapsto (\beta_m(N_c(x_{j'})))_c$, which has finitely many values because c ranges over a finite set. If y is a coordinate x_j, the value is $T_{(\beta(\vec{x}),\beta_m(n(\vec{x})))}(x_j)$ with an index determined by features of the individual coordinates. If y is m, the value is $(T_{(b,f)} \circ N_c)(x_{j'})$ with $b = \beta(\vec{x})$, $c = \gamma(\vec{x})$ and $f = \beta_m(n(\vec{x}))$; the compositions belong to $\mathcal{U}$ by Theorem 9.86, and the index is again determined by features of the coordinates.

Induction on terms. Variables, 0 and successors are immediate, the last by composition with the successor. For $F(n; v.h)$ first replace the counter by a fresh variable m. By the induction hypothesis $h(v,\vec{x}) = H_{(\beta_v(v),\beta(\vec{x}))}(y)$ with y active.

If y is the state v, fix the features $c = \beta(\vec{x})$. The update $h_c(v) = H_{(\beta_v(v),c)}(v)$ belongs to $\mathcal{U}$, and by Theorem 9.87 so does its orbit r_c. Then $F(m; v.h) = r_{\beta(\vec{x})}(m)$ has finite-control form with the counter active.

If y is a passive parameter p, form the finite catalogue of the functions $V_k(p)$ of p consisting of all update profiles $H_{(b,c)}(p)$ and all initial profiles $1 + H_{(b,c)}(p)$; every member belongs to $\mathcal{U}$. For fixed $\vec{x}$ every state of the orbit is a catalogue value at p: the initial state is $1 + H_{(\beta_v(0),c)}(p)$ and the state after a state w is $H_{(\beta_v(w),c)}(p)$, with $c = \beta(\vec{x})$. The finite-valued feature

$$\kappa(p) = \big(\beta_v(V_k(p))\big)_k$$

determines, together with c, every transition of the catalogue indices. For each pair (c,κ) let $\theta_{c,\kappa}(m)$ be the index reached by this finite-state process after m updates, and pack these finitely many processes into the finite-valued feature $\eta(m) = (\theta_{c,\kappa}(m))_{c,\kappa}$ of the counter. Then

$$F(m; v.h) = V_{\eta(m)_{\beta(\vec{x}),\kappa(p)}}(p),$$

which is finite-control form with p active. The argument uses only the displayed finite observations and no test of numerical equality.

Finally substitute the actual counter term n for m by the composition step. This completes the induction and proves the bound. For the predecessor choose $n > C + 1$: its value exceeds C and is less than n. Halving fails at every $n > 2C + 1$. □

Remark 9.88. The theorem concerns definitions by terms under the standard interpretation. It is not an obstruction to interpretations by graphs or on changed domains. The counter-aware tied-seed language of the package `local_scopes` has a finite-control obstruction of its own but defines the predecessor; the unary bound is specific to removing the access to the old counter together with the independent initializer.

9.9.2 The two computation equations of the iterator are independent

Proposition 9.89. *Neither computation equation of the iterator of Section 1.1 follows from the other, even over the standard natural numbers with coverage, separation and every first-order induction instance.*

Proof. Interpret every fixed body compositionally in $\mathbb{N}$. If $I(n,a;\, u.\, h)$ is interpreted as $h^n(0)$, the successor equation holds and $I(0,1;\, u.\, u) = 0 \neq 1$. If it is interpreted as a, the zero equation holds and the successor equation fails for $h(u) = Su$. □

Thus the canonical calculi succeed by changing the representation; they do not derive a deleted law from the remaining axioms.

9.9.3 Induction on zero equations does not give induction on equations

Let T_0 be the theory T_{eq} of Section 1.5 with its induction rule restricted to equations of the form $0 = t(n, \vec{p})$, or equivalently $t(n, \vec{p}) = 0$, with an arbitrary context. A calculus that uses the progress introduction only for matrix equations with the literal 0 on one side has this induction. The following theorem shows that such a restriction does not preserve the consequence relation.

Theorem 9.90 (A model separating the two inductions)**.** *There is a classical model of T_0, in which the successor is injective, and in which the sentence*

$$\forall n\, \forall a\ I(n, a;\, u.\, u) = a$$

is false, although its instance at $n = 0$ and its induction step are provable from the computation equations and equality logic. Consequently induction on arbitrary literal equations is not admissible in T_0. The same model satisfies induction for every equation $\bar{k} = t(n, \vec{p})$ with a fixed standard numeral $\bar{k}$ on one side.

A classical model is in particular a model of intuitionistic logic. Its construction uses compactness and choices of compatible paths in the metatheory; these are not principles granted to T_0.

Lemma 9.91 (Paths through finite fibers)**.** *Let $(B_j)_{j\in\mathbb{Z}}$ be finite nonempty sets and $f_j \colon B_j \to B_{j+1}$ total maps. There is a sequence $(b_j)_{j\in\mathbb{Z}}$ with $b_j \in B_j$ and $b_{j+1} = f_j(b_j)$ for every j.*

Proof. The product $\prod_j B_j$ of finite discrete spaces is compact, and each equation between adjacent coordinates is a closed condition. Every finite set of these conditions is satisfiable: choose a value at the least index involved, propagate it forward, and choose arbitrary values elsewhere. The finite intersection property gives a common solution. □

Proof of Theorem 9.90. The base. Fix a nonstandard model M of Peano arithmetic expanded by symbols for all primitive recursive descriptions; it exists by compactness. For $m \in M$ let $A_m = \{0\}$ if m is a standard numeral and $A_m = \{0, 1\}$ otherwise; the distinction is external. Put

$$D = \{(m, i) : m \in M,\ i \in A_m\}, \qquad \pi(m, i) = m, \qquad 0_D = (0_M, 0), \qquad S_D(m, i) = (S_M m, i).$$

The successor is well defined because it preserves standardness and nonstandardness. The map π is surjective and preserves zero and successor. Every standard numeral has a one-element fiber and every nonstandard element has two lifts. On D the successor is injective, zero is not a successor, and coverage holds: a nonstandard (m, i) has the predecessor $(m-1, i)$. The nonstandard part of M is a disjoint union of chains of the order type of $\mathbb{Z}$ under successor, and the label i is constant along each chain of D.

The iterator on actual functions. Let $\mathscr{H}$ be the set of functions $h \colon D \to D$ for which there is a unary function $f \colon M \to M$, primitive recursive with parameters, such that

$$\pi(h(x)) = f(\pi(x)) \qquad (x \in D). \tag{9.25}$$

By surjectivity of π the function f is unique. Write $\mathrm{It}_f(m, b)$ for its iteration in M with initial value b. This does not depend on the description chosen for f: two descriptions of the same function give iterations with the same base and pointwise equal updates, and an induction in M proves them equal at every counter. This induction is used in the base model, not in T_0. Define an operation $\mathcal{I} \colon D \times D \times \mathscr{H} \to D$ as follows. For a standard counter

k put $\mathcal{I}(k,a,h) = h^k(a)$, the finite iteration on D. For fixed a, h and a nonstandard chain $n_j = (m_j, i)$, $j \in \mathbb{Z}$, with $m_{j+1} = S_M m_j$, put

$$r_j = \mathrm{It}_f(m_j, \pi(a)), \qquad B_j = \pi^{-1}(r_j).$$

The sets B_j have one or two elements, $r_{j+1} = f(r_j)$ by the computation law in M, and (9.25) makes h a total map $B_j \to B_{j+1}$. Choose a path $b_j \in B_j$ by Theorem 9.91 and put $\mathcal{I}(n_j, a, h) = b_j$. Make these choices for all triples of a function, an initializer and a chain, with one specified family: for $h = \mathrm{id}_D$ choose on every nonstandard chain the constant path

$$\mathcal{I}(n, a, \mathrm{id}_D) = (\pi(a), 0). \tag{9.26}$$

This is legitimate, since the projected iteration is constantly $\pi(a)$, the point with label 0 always exists, and the identity fixes it. The choices depend on the actual function h, so every description of the identity body receives the same family. By construction

$$\mathcal{I}(0_D, a, h) = a, \qquad \mathcal{I}(S_D n, a, h) = h(\mathcal{I}(n, a, h)), \qquad \pi(\mathcal{I}(n, a, h)) = \mathrm{It}_f(\pi(n), \pi(a)),$$

the standard and the nonstandard counters never meeting under a finite number of successor steps.

Terms. Interpret $I(n, a;\, u.\, h)$ in D under a valuation ρ as $\mathcal{I}(\llbracket n \rrbracket, \llbracket a \rrbracket, v \mapsto \llbracket h \rrbracket_{\rho[u:=v]})$. By induction on the term, with a bound variable of a body treated as a further free variable,

$$\pi(\llbracket t \rrbracket_{D,\rho}) = \llbracket t \rrbracket_{M, \pi\circ\rho}, \tag{9.27}$$

where the right side uses the iterator of M; in particular every body function supplied to $\mathcal{I}$ satisfies (9.25) with the interpretation of the body in M and belongs to $\mathscr{H}$. The operation $\mathcal{I}$ is chosen for actual functions before this induction, so nested terms create no circular definition. Equal parameter values determine the same body function and the same iterator; hence D is a first-order structure for the staged language, the substitution lemma holds, and both computation schemas hold for every fixed body.

Zero equations. Since the fiber over 0_M has one element, $\llbracket t \rrbracket_D = 0_D$ if and only if $\llbracket t \rrbracket_M = 0_M$ at the projected valuation. Fix parameters $\vec{a}$ in D and suppose that in D

$$t(0, \vec{a}) = 0, \qquad \forall n\, (t(n, \vec{a}) = 0 \to t(Sn, \vec{a}) = 0).$$

By (9.27) and surjectivity of π the same two statements hold in M for the primitive recursive interpretation of t at $\pi\vec{a}$. Induction in M and reflection of zero along π give $\forall n \in D\; t(n, \vec{a}) = 0$. The rule with a context is valid as well: at a valuation that makes the context true the two premises supply the displayed base and universal step, and no projection of the context is needed. The same argument applies to an equation $\bar{k} = t(n, \vec{a})$, because every standard numeral has a one-element fiber.

Failure of the full rule. Choose a nonstandard $b \in M$ and $a = (b, 1)$, and let $E(n)$ be $I(n, a;\, u.\, u) = a$. Its base is a computation equation, and its step is provable from $I(Sn, a;\, u.\, u) = I(n, a;\, u.\, u)$ and transitivity. For every nonstandard counter n, however, (9.26) gives $I(n, a;\, u.\, u) = (b, 0) \neq (b, 1) = a$. The two elements have the same projection; tests against zero cannot distinguish the labels. □

Corollary 9.92 (No reflection of equality)**.** T_0 *proves for no term* $e(x, y)$ *the law* $e(x, y) = 0 \leftrightarrow x = y$.

Proof. In the model take $x = (b, 0)$ and $y = (b, 1)$ over a nonstandard b. By (9.27) the terms $e(x, y)$ and $e(x, x)$ have the same projection. If the law held, $e(x, x) = 0$, and the one-element fiber over zero would force $e(x, y) = 0$, contradicting $x \neq y$. □

The usual conversion of an equation into a zero equation therefore cannot justify the restriction: the reflection of equality of Theorem 1.52 is exactly the missing conclusion. The theorem refutes the recovery of the induction rule of T_{eq} in the presence of full intuitionistic logic with equality, all numerical cases, both computation schemas and injectivity of the successor. Under the reading of Section 1.2, the rules QE, DE, introduction by cases and progress for zero equations are valid in the same model, so the native rules do not evade the counterexample. The construction does not show that T_0 has no interpretation of arithmetic on a quotient or on a changed domain (Section 9.11).

9.9.4 The last computation rule

Proposition 9.93. *The calculus $\mathcal{K}_5$ without rule (R) derives no equation from the empty context, not even $0 = 0$.*

Proof. Assign the value false to every equation and the value true to every paired formula, and read a clause $[A;\, y.\, B]$ as the implication from the value of A to the value of B. This agrees with the reading of the paired binder as a conditional closure when all equations are false, and it does not impose the equality axioms. By induction on derivations, if all clauses of Γ are true, then the conclusion of $\Gamma \vdash A$ is true. Rule (T) has equations as premises, so it applies only when some clause is false. Rule (U) is modus ponens. In rule (E) the opened clause $[B(b, 0);\, y.\, B(b, y)]$ is true, because its two components have the same root. Rule (I) concludes a paired formula. With the empty context the conclusion is true and is therefore not an equation. □

Adding a reflexivity rule or another seed of equations repairs the loss, as in Sections 9.6 and 9.7, but that is a replacement and not a deletion. The proposition concerns this one deletion; it is not a theorem that a presentation of iPRA needs five rules.

9.10 Checker sizes

This section draws on the measurements stated by `register_graph`, `canonical_recursors`, `clausal_canonical`, `counter_evaluation`, `uniform_binding`, `uniform_recursion` and `single_constructo`

Each row is a complete standalone module in the sense of Section 1.10: imports, JSON decoding, validation of syntax and scope, normalization, substitution, every rule and the final comparison are counted. All measurements are stated by the packages for CPython 3.12.14, after removal of docstrings and `ast.unparse` with one final line feed. The role column names the new program of a package, its matched control, and the other programs that the package reports for orientation. The programs marked † implement the kernel on iterator terms and no calculus of this chapter.

Program	Tokens	AST	Bytes	Role
`register_graph`				
Reference checker for T_J	2814	3060	13633	new
Ancillary profile of the kernel†	261	288	604	new
Direct parent of the ancillary profile†	261	292	576	control
Earlier leader in nodes†	263	289	613	other
Earlier leader in bytes†	266	310	529	other
`canonical_recursors`, strict family				
$\mathcal{K}_6$ on canonical seeded terms, six rules	1338	1485	4611	new
Iterator terms with successor, seven rules	1433	1595	4923	control

Program	Tokens	AST	Bytes	Role
Raw seeded terms, seven rules	1378	1532	4753	control
`canonical_recursors`, compact guarded codec[†]				
Refined compact byte profile	265	306	516	new
Frozen compact byte control	265	303	522	control
Refined compact node profile	263	280	602	new
Frozen compact node control	262	281	606	control
Refined strict guarded checker	542	624	1376	new
Matched strict guarded control	541	620	1381	control
`clausal_canonical`				
$\mathcal{K}_5$, five rules	946	1072	3305	new
$\mathcal{K}_6$ in the same variable convention	1107	1266	3820	control
$\mathcal{K}_6$ as shipped by `canonical_recursors`	1338	1485	4611	other
Six rules on zero-normal iterator terms (`zero_normal`)	1013	1075	3095	other
`counter_evaluation`				
$\mathcal{K}_{\mathrm{CE}}$	854	987	3004	new
$\mathcal{K}_{\mathrm{CT}}$, same terms, computation and transport	929	1063	3272	control
Scalar zero-normal, evaluation with one case	884	1025	3061	other
Raw scalar clause checker with five rules, unchanged	853	993	3046	other
The same with the known deletion of the root case	848	986	2999	other
$\mathcal{K}_5$ of `clausal_canonical`, unchanged	946	1072	3305	other
`uniform_binding`				
$\mathcal{K}_{\mathrm{CE}}$, goal directed, one traversal	693	764	2238	new
Scalar checker with the same shared traversal	699	773	2284	control
Incoming scalar checker, exact source	741	826	2283	control
Conclusion-synthesizing checker of `counter_evaluation`	854	987	3004	other
`uniform_recursion`				
$\mathcal{K}_{\mathrm{Name}}$	662	741	1950	new
Scalar control with NAME	663	751	1983	control
$\mathcal{K}_{\mathrm{Name}}$ with a budget	710	800	2136	new
Scalar control with a budget	711	810	2169	control
Strict native checker, explicit fusion[†]	340	385	685	new
Strict native checker, integrated cases[†]	336	381	661	new
Strict native control[†]	360	409	780	control
Compact control with a different contract[†]	267	302	508	other
`single_constructor`				
$\mathcal{K}_{\mathrm{Open}}$	683	764	1965	new
Iterator terms with inferred opening	683	771	1990	control

The comparisons that isolate a change of calculus are those against a matched control. The checker for $\mathcal{K}_6$ is 312 bytes (6.34%) below the scalar control and 142 bytes below the raw control, with normalization counted. The checker for $\mathcal{K}_5$ is 515 bytes (13.48%), 161 tokens and 194 nodes below its matched control; the difference of 1306 bytes to the shipped checker for $\mathcal{K}_6$ combines the change of interface with the simpler variable representation and is not to be attributed to the deleted rule. The checker for $\mathcal{K}_{\mathrm{CE}}$ is 268 bytes (8.19%), 75 tokens and 76

nodes below $\mathcal{K}_{\mathrm{CT}}$, and the node overheads of the translation are those of Theorem 9.55. The goal-directed checker with one traversal is 46 bytes, 6 tokens and 9 nodes below the matched scalar control and 45 bytes, 48 tokens and 62 nodes below the exact incoming checker. The checker for $\mathcal{K}_{\mathrm{Name}}$ is 33 bytes, one token and 10 nodes below its control, and the checker for $\mathcal{K}_{\mathrm{Open}}$ is 25 bytes and 7 nodes below its control with the same token count. Compressed sizes are reported by the packages as supplementary data and are not an optimization target; they are omitted here.

9.11 Further questions

This section draws on `register_graph`, `canonical_recursors`, `clausal_canonical`, `counter_evaluation`, `uniform_binding`, `uniform_recursion` and `single_constructor`. It collects their open questions, the claims that they state without proof, and what they report as not achieved.

Status of formalization. None of the calculi and term languages of this chapter is formalized in Lean or Rocq. The theorems of Sections 9.1 to 9.7 and 9.9 are written proofs, and the executable tests of the packages are finite evidence for the implementations. The formalized results named in Sections 9.4 and 9.7 concern iterator terms.

Question 9.94 (Formalization)**.** Each term language of this chapter needs its own syntax, its normalizer where it has one, and the bi-interpretation with iterator terms of Theorem 9.25. The packages propose the following orders of work. For Theorems 9.34 and 9.40: body extensionality before the soundness of normalization, and the tests on literal sources and witnesses after the unrestricted equivalence. For $\mathcal{K}_5$: the invariant on context occurrences and the eta expansion of the closure, proved directly. For $\mathcal{K}_{\mathrm{CE}}$: a deep embedding with a checked equivalence to the kernel; the arithmetic reconstruction is small and the main work is the syntax with two binders and its canonicalization. For normal terms: extend normalization by evaluation from iterator terms to the counter-aware recursor, prove the environment laws (9.8)–(9.9), and lift the comparison of the inferred opening through Theorem 9.77. For the checkers: the invariant of the sorted walker (Theorem 9.83) together with the interpretation of the counter-aware recursor. For the register: the compiler and both round trips as translations of finite descriptions, keeping fixed syntactic bodies apart from arbitrary functions of the metatheory. Can these be carried out by extending the existing development?

Question 9.95 (A native register kernel)**.** Can a native checker with Q_y and $D_{x,y}$ share its verified equality premise with the increment and load steps (9.12), (9.13), so that the register grammar also reduces the complete trusted source, when all conditional premises, scope checks and syntax checks are counted? Theorem 9.19 shows compatibility of the basis with such a kernel; no checker for it is supplied.

Question 9.96 (The register selector and the cost of the counter)**.** Is there a substantially shorter numerical selector for the register basis than the term C of Theorem 9.16, which has a short proof and a large expansion? Source brevity, term size and certificate size are different objectives. Does the register with two bound variables, or the one-variable register of Theorem 9.18, yield the smaller complete checker after comparable implementations?

Question 9.97 (Weakening the graph axiom)**.** Which parts of (STEP) can be weakened while actual equality and the formula round trips on the whole domain are retained? `register_graph` states without proof that one-sided operational membership and agreement on the standard model are insufficient.

Question 9.98 (Zero-equation induction)**.** Which principle beyond induction on zero equations excludes the two-point fibers of Theorem 9.90? Reflection of actual equality suffices, at

a cost that has to be counted. Does a narrowly stated uniqueness or extensionality principle for the identity iterator already give full induction on equations, and what does it cost in a complete checker? Does T_0 interpret arithmetic on a quotient, or admit a bi-interpretation with iPRA on a changed domain? The countermodel decides neither.

Question 9.99 (Successor rigidity)**.** The package on raw clause calculi with ordinary substitution states a restricted successor-rigidity principle that yields induction on zero equations and leaves a faithful bootstrap of equality and predecessor unresolved. Does that narrower principle suffice? `counter_evaluation` records the question and does not use the unproved conclusion.

Question 9.100 (The self-seeded language)**.** Can Theorem 9.85 be strengthened to a normal form or an effective classification of the functions defined by self-seeded terms? The proof gives the existence of a bound and no algorithm that extracts an optimal bound from a term. Is there an obstruction for interpretations by graphs or on changed domains?

Question 9.101 (Tests on literal syntax)**.** The witness-directed introduction is concise but not stable under substitution: a substituted witness can become the literal 0. Is there a substitution-stable presentation with a comparable complete checker? In particular, guarded atomic matrices make the introduction depend on the root of the formula only and guarantee a reflexive zero row for the opening (Open); is the resulting complete grammar with its checker simpler than the unrestricted $\mathcal{K}_{\mathrm{Open}}$? Is there a uniform counter rule on raw syntax, with the zero computation stated explicitly instead of placed in the canonical substitution, whose checker is of comparable size?

Question 9.102 (Compact encodings)**.** Can an encoding of canonical seeded terms absorb the erasure of zero-counter bodies into a traversal that is needed anyway and improve the compact source minima, while validating the complete public language? Does the grammar with disjoint arities, without a successor constructor and without a closure constructor, yield a smaller fully validated compact program once normalization and every case of malformed input are counted? Can the combination of the counter-aware arithmetic with the three families be ported to the compact certificate architecture below the existing profiles? Can a uniform representation remove the partial opening of binders, given that the recursor and the paired binder bind two variables and a clause one, while reducing the whole checker? Can the two instantiations of the counter body in (CE) be given a simpler complete scope discipline? Can a pointfree representation of proofs reduce the substitution machinery without introducing pairing structure, when the typing and conversion costs of external tuple contexts are counted?

Question 9.103 (Proof size)**.** What are the optimal translations of proof graphs between $\mathcal{K}_6$ and $\mathcal{K}_5$, and how much sharing avoids repeated neutral premises? Can a comparison of proof complexity between counter evaluation and the evaluation rule on the state of an iterator quantify certificate lengths and label sizes in addition to the inference nodes of Theorem 9.55? Should checkers with an explicit budget be compared only with controls that carry the same budget?

Question 9.104 (Measures and lower bounds)**.** The coordinates used in Section 9.10 and the counts of constructors, rules and proof expansion disagree with one another. Is there a robust measure that compares different proof interfaces? Can a lower bound be proved for a restricted class of checkers, with the term grammar, the class of decidable operations on syntax, the permitted runtime primitives and the certificate interface fixed in advance? A count of named rules alone changes too easily, by merging cases or by moving work into the syntax.

Question 9.105 (A new arithmetic reduction)**.** Eliminating the initializer, changing the domain or replacing the recursor may still give a simpler basis. A candidate has to respect the self-seeded constructions and the obstruction of Theorem 9.85, and it has to supply both interpretation composites, not only expressiveness on the standard model.

Not achieved. The packages state the following limits.

- No program of Section 9.10 for a calculus of this chapter is below the compact codecs of the kernel (516 and 508 bytes), and no minimality of an axiomatization or of a checker is claimed. An observed minimum among programs is an upper bound on the optimum.
- The register theory does not improve the size of a complete checker: its graph axiom is richer, its body has two bound variables, and its reference checker has 13633 bytes. The ancillary profile lowers the node count by one and is larger in bytes.
- The refined compact byte profile does not improve the token count of its control, and the token frontier of 258 reported by `canonical_recursors` is not beaten.
- The checker for $\mathcal{K}_5$ has fewer rules and tokens but more bytes than the six-rule checker on zero-normal iterator terms of `zero_normal`; its positive claim in bytes is the matched comparison.
- The checker for $\mathcal{K}_{\mathrm{CE}}$ is smaller in all three coordinates than the scalar control with one evaluation case, but the raw scalar clause checker with the root case deleted is five bytes smaller.
- For the checker of `uniform_binding` the compressed size is larger than that of the exact incoming scalar checker.
- Two directions were examined in `uniform_binding` and discarded. A formula of the clause syntax is a chain of paired binders ending in an equation and can be flattened to a triple of a depth and two terms; the change is exact, but the implementation grew, because checking of ranks and opening of endpoints outweighed the saved traversal. A rule of parameter substitution used only in the reverse direction is sound, but the intended completeness argument by a fresh alias fails: once a goal no longer mentions the alias, the rule can require that same goal as its premise.
- The translator of test fixtures of `uniform_recursion` declines an inherited fixture whose witness becomes the literal 0 under normalization; a direct proof of the same formula is supplied, and no universal executable proof compiler is claimed.
- Concrete limits of the interpreter on recursion depth and memory can reject a valid large certificate for every checker of the chapter; completeness is a statement about the calculi and about the checkers without such limits.

Chapter 10

Obstructions

The chapter collects the negative results: what a smaller numerical basis or a smaller rule inventory cannot do, together with the model or invariant that proves it. Each result names the exact class of theories or presentations that it excludes and states what it does not exclude. The results are grouped by method. Section 10.2 uses coordinatewise powers of the standard model and exact truth transfer for conjunctive formulas. Section 10.3 uses the automorphisms of such powers. Sections 10.4 and 10.5 use growth invariants of term languages and a dimension argument for term interpretations. Section 10.6 counts definable values and reaches arbitrary relational bi-interpretations through witness extraction. Section 10.7 classifies zero sets. Section 10.8 collects small counter-interpretations of individual rule deletions. Section 10.9 builds a nonstandard model of all iterator computations, by a tower of ultrapowers, in which the progress rule fails. The packages that prove the obstructions also contain positive statements about the presentations that the obstructions are measured against; these are in Sections 10.10 to 10.13.

None of the results is a lower bound on the size of a proof checker, and none excludes equiconsistency. Table 10.1 lists the main statements.

Table 10.1: The obstructions of the chapter.

Result	Excluded	Not excluded
Theorem 10.16	case introduction with target z in place of Sz, with all other native rules	other readings of the weaker calculus
Theorem 10.17	deletion of Boolean cases from the progressive calculus	a weaker case principle
Theorem 10.22	ordinary bi-interpretation with iPRA of any theory true in an eventually constant power: equation induction, conjunctive induction, all true conjunctive sequents, many sorts	one-way interpretations, equiconsistency, disjunctive axioms
Theorem 10.28	a term compiler into the zero-seeded recursor	relational interpretations
Theorems 10.32 and 10.36	terms for addition, selection and pairing when the initializer is computed from the update	an independent input, relational interpretations

Result	Excluded	Not excluded
Theorem 10.41	finite-dimensional interpretations with a term-defined addition, any quotient equality	graph-defined operations
Theorem 10.48	terms for an equality test, minimum and truncated subtraction in the tied-seed language	formula-defined relations
Theorem 10.54	every finite-dimensional bi-interpretation of iPRA with an existential-free theory true in the standard tied-seed algebra	one-way interpretations, theories with positive existence axioms
Theorem 10.58	terms for predecessor, zero test, addition and initialized iteration from positive iteration	graph definitions
Theorems 10.61, 10.63 to 10.65 and 10.99	five individual rule deletions or replacements	redesigned calculi with their own interpretation proofs
Theorem 10.75	deletion of QI by progress from the core: the other seven rules do not derive $Q_y(I(y, 0;\, u.\, u) = 0)$	fusions of QI with another rule, weaker induction interfaces, equivalence of uniqueness of iteration with induction

10.1 Conventions and the presentations under test

This section fixes the vocabulary shared by `adaptive_seed`, `research`, `local_scopes`, `checker_and_barriers`, `compact_kernel`, `uniform_predecessor`, `scope_kernel`, `constructed_computation` and `counter_congruen`

10.1.1 Three kinds of exclusion

The statements of the chapter have three strengths.

Term definability. A reduced term language is given a standard interpretation: every fixed term denotes a primitive recursive function on $\mathbb{N}$, by structural recursion on the term followed by numerical primitive recursion. No uniform evaluator of all descriptions is used. A term-definability obstruction is a statement about these functions. It excludes a direct compiler of the primitive recursive signature into terms with their usual meanings. It does not exclude a definition of the missing function by a formula.

Derivability in a specified rule system. A rule system is shown to lose a theorem of another one by a reading under which every rule of the first is sound and the theorem is false. A classical structure is a one-node Kripke model, so soundness in a classical structure is a sound test of intuitionistic derivability. Such a result concerns the displayed rules with the identity translation of formulas. It does not exclude a different translation.

Ordinary bi-interpretability. This is the strongest kind.

Definition 10.1 (Ordinary interpretations)**.** An *ordinary interpretation* $\mathcal{I}$ of a theory U in a theory T has a dimension $d \geq 0$ and consists of formulas of T: a domain formula $\delta(\vec{x})$ in d variables for each sort of U, an equality formula $\varepsilon(\vec{x}, \vec{y})$, and a graph formula for each function and relation symbol of U. The formulas may contain finitely many parameters, subject to a parameter formula. T proves that ε is an equivalence on δ respected by the graphs, that the function graphs are total and single-valued modulo ε, and the translation of every axiom of U. The domain, the graphs and the quotient equality are arbitrary formulas.

A *bi-interpretation* of T and U is a pair of ordinary interpretations $\mathcal{I}$ of U in T and $\mathcal{J}$ of T in U such that T proves that a formula η defines an isomorphism of the identity interpretation of T onto $\mathcal{J}$ interpreted through $\mathcal{I}$, and U proves the corresponding statement for the other composite. The formula η of T is the *source-side return isomorphism.*

The bridge of Theorem 1.78 is a bi-interpretation in this sense, of dimension one, with identity domains, actual equality and return isomorphisms with graph $x = y$. The obstructions of Sections 10.3 and 10.6 exclude all bi-interpretations in the sense of Theorem 10.1, not only those of this special form. They use only the source-side return isomorphism.

10.1.2 The paired-introduction form of the native core

The packages of this chapter state the native calculus with a progress introduction that concludes a paired formula:

$$\frac{\Gamma,\ E(t,z) \vdash E(t,Sz)}{\Gamma \vdash D_{x,y}E(x,y)}\quad \text{DI-PROGRESS}, \qquad E \text{ a literal equation, } z \text{ fresh for } \Gamma, t.$$

The core N of Section 1.3 has QI by progress instead. The *paired-introduction form* N_{p} is N with QI replaced by DI-PROGRESS. Several packages also supply a computation instance as a discharged local assumption (rule COMP, called SUPPLY in some packages): from $\Gamma, H \vdash A$ with H an instance of a computation equation infer $\Gamma \vdash A$.

Lemma 10.2. N *and* N_{p} *derive the same sequents.* COMP *is admissible in both, and the computation equations are derivable from* COMP *and* HYP.

Proof. Let $\Gamma, E(t,z) \vdash E(t,Sz)$ with z fresh. In N, QI for the matrix $y \mapsto E(t,y)$ gives $\Gamma \vdash Q_y E(t,y)$. Weaken by $E(t,0)$. HYP gives $E(t,0)$, and QE at the target Sz' gives $\Gamma, E(t,0) \vdash E(t,Sz')$. DI by cases with witness t gives $\Gamma \vdash D_{x,y}E(x,y)$. Conversely, let $\Gamma, E(z) \vdash E(Sz)$. In N_{p}, DI-PROGRESS with a witness coordinate u that does not occur in E gives $\Gamma \vdash D_{u,y}E(y)$. DE supplies the clause $Q_y E(y)$, which does not contain the fresh witness, and HYP concludes $Q_y E(y)$. COMP is a cut with a computation equation (Section 1.4); conversely COMP followed by HYP derives the equation. □

In the paired-introduction form the formula $Q_y B$ is interderivable with $D_{u,y}B$ for an unused u: DE followed by HYP gives one direction, and DI by cases at the witness 0 with QE gives the other. The rule-level statements of Section 10.8 refer to N_{p} where the difference matters.

10.1.3 Reduced term languages

Definition 10.3 (Seedless recursors)**.** Each of the following languages has variables, 0 and one recursion constructor. Only the update body binds the displayed bound variables; its other variables are passive parameters, fixed through the recursion. Bodies are fixed finite earlier terms.

(a) *Zero-seeded recursion* has, besides 0 and S, the constructor $\mathsf{L}(n; u, v.h)$ with

$$\mathsf{L}(0; u, v.h) = 0, \qquad \mathsf{L}(Sn; u, v.h) = h[n/u][\mathsf{L}(n; u, v.h)/v].$$

(b) *Tied-seed recursion* has the constructor $\mathsf{B}(n; u, v.h)$ and the abbreviation $Sa := \mathsf{B}(0; u, v.a)$ for u, v not in a, with

$$\mathsf{B}(0; u, v.h) = S(h[0/u][0/v]), \tag{10.1}$$
$$\mathsf{B}(Sn; u, v.h) = h[n/u][\mathsf{B}(n; u, v.h)/v]. \tag{10.2}$$

The two substitutions are simultaneous. The initial state is tied to the update; there is no independent initializer. The standard interpretation is

$$\mathsf{B}_h(0,\vec{p}) = 1 + h(0,0,\vec{p}), \qquad \mathsf{B}_h(m+1,\vec{p}) = h(m,\mathsf{B}_h(m,\vec{p}),\vec{p}),$$

so that S denotes the successor. The *unary-update* variant $\mathsf{F}(n; v.h)$ is the case in which u does not occur in h.

(c) *Positive iteration* has, besides 0 and S, the constructor $\mathsf{R}^+(n, a; v.h)$, standing for $h^{n+1}(a)$:

$$\mathsf{R}^+(0,a;v.h) = h[a/v], \qquad \mathsf{R}^+(Sn,a;v.h) = h[\mathsf{R}^+(n,a;v.h)/v].$$

For reference, the *counter-aware initialized recursor* is $\mathsf{R}(n, a; u, v.h)$ with $\mathsf{R}(0, a; u, v.h) = a$ and $\mathsf{R}(Sn, a; u, v.h) = h[n/u][\mathsf{R}(n, a; u, v.h)/v]$. It is ordinary primitive recursion and compiles every primitive recursive description directly.

The tied-seed language is the most capable of the three. The next two lemmas record what it does define; they are the positive boundary of the obstructions of Sections 10.4 and 10.6.

Lemma 10.4 (Terms of the tied-seed language)**.** *Put*

$$W(n) := \mathsf{B}(n; u, v.u), \quad Z(n) := \mathsf{B}(n; u, v.0), \quad H(n,a) := \mathsf{B}(n; u, v.a), \quad P(n) := W(H(n,n)).$$

The computation equations (10.1)*,* (10.2) *and equality reasoning prove*

$$W(0) = 1,\ W(Sn) = n, \quad Z(0) = 1,\ Z(Sn) = 0, \quad H(0,a) = Sa,\ H(Sn,a) = a, \quad P(0) = 0,\ P(Sn) = n.$$

With induction for literal equations, the two gates

$$G(n,a) := P(P(\mathsf{B}(a; u, v.H(Z(n), v)))), \qquad G'(n,a) := P(P(\mathsf{B}(a; u, v.H(n, v))))$$

satisfy $G(0,a) = 0$*,* $G(Sq,a) = a$ *and* $G'(0,a) = a$*,* $G'(Sq,a) = 0$*.*

Proof. The equations for W, Z and H are instances of (10.1) and (10.2). Then $P(0) = W(H(0,0)) = W(S0) = 0$ and $P(Sn) = W(H(Sn,Sn)) = W(Sn) = n$. For G at the control 0, the update is $H(Z(0), v) = H(1, v) = v$. Induction on the literal equation $\mathsf{B}(a; u, v.H(Z(0), v)) = 1$ has the base $S(H(1,0)) = S0$ and a successor step that repeats the previous state. Two predecessors give $G(0,a) = 0$. At the control Sq the update is $H(0, v) = Sv$. Induction on $\mathsf{B}(a; u, v.H(Z(Sq), v)) = SSa$ has the base $S(H(0,0)) = SS0$ and a successor step that applies S. Two predecessors give $G(Sq,a) = a$. The argument for G' exchanges the two cases. The inductions keep the actual update bodies and the control parameter fixed; no extensionality principle for recursion descriptions and no induction on a compound formula is used. □

Lemma 10.5 (Zero-start counter recursion is a macro)**.** *For a body h with bound variables u, v and fresh r, s, put*

$$L(n; u, v.h) := \mathsf{B}(Sn;\ r, s.\, G(r,\ h[Pr/u][s/v])).$$

Then $L(0; u, v.h) = 0$ and $L(Sn; u, v.h) = h[n/u][L(n; u, v.h)/v]$ follow from Theorem 10.4 without further induction.

Proof. Let $K(r,s) = G(r, h[Pr/u][s/v])$. The gate equations and $P(Sr) = r$ give $K(0,s) = 0$ and $K(Sr,s) = h[r/u][s/v]$. Hence $\mathsf{B}_K(0) = S(K(0,0)) = 1$ and $\mathsf{B}_K(1) = K(0, \mathsf{B}_K(0)) = 0$, which is $L(0) = 0$. At a successor, $L(Sn) = \mathsf{B}_K(SSn) = K(Sn, \mathsf{B}_K(Sn)) = h[n/u][L(n)/v]$. □

Remark 10.6. Theorem 10.4 is in `counter_congruence` (with the gate G'), `research` (with $P(n) = W(H(Z(n), W(n)))$, which computes the same function) and `local_scopes` (with the gate G and the internal proof). Theorem 10.5 is in `local_scopes`. Predecessor, a zero test, a one-sided gate and zero-start recursion are therefore not what the tied-seed language lacks. The obstruction is to combining two unbounded inputs.

10.1.4 The progressive calculus

Definition 10.7 (Progressive binder calculus)**.** Numerical terms are variables, 0, St and the counter-aware recursor $\mathsf{R}(n, a; u, v.h)$ with its two computation equations. Object formulas are

$$A ::= (s = t) \mid \mathsf{H}_{x,y} A.$$

Internal clauses are $\mathsf{P}_y A(y)$ with A an object formula. Contexts are finite reusable lists of object formulas and clauses. Write $\Gamma \triangleright M$ when M is a member of Γ or a proved object formula; clauses are not goals. The ordinary readings are

$$(\mathsf{P}_y B)^\circ = \forall y\, (B^\circ(y) \to B^\circ(Sy)), \qquad (\mathsf{H}_{x,y} B)^\circ = \exists x\, \forall y\, (B^\circ(x, y) \to B^\circ(x, Sy)).$$

Besides the computation equations, the rules are

$$\frac{\Gamma \triangleright s = t \quad \Gamma \vdash A(s)}{\Gamma \vdash A(t)}\ \textsc{Tr} \qquad \frac{\Gamma,\ B(t, z) \vdash B(t, Sz)}{\Gamma \vdash \mathsf{H}_{x,y} B(x, y)}\ \text{HI}$$

$$\frac{\Gamma \triangleright \mathsf{H}_{x,y} B(x, y) \quad \Gamma,\ \mathsf{P}_y B(a, y) \vdash A}{\Gamma \vdash A}\ \text{HE} \qquad \frac{\Gamma \triangleright \mathsf{P}_y B(y) \quad \Gamma \vdash B(t)}{\Gamma \vdash B(St)}\ \textsc{Step}$$

$$\frac{\Gamma \triangleright \mathsf{P}_y E(y) \quad \Gamma \vdash E(0)}{\Gamma \vdash E(t)}\ \textsc{Iter} \qquad \frac{\Gamma,\ \mathrm{sg}(t) = 0 \vdash A \quad \Gamma,\ \mathrm{sg}(t) = 1 \vdash A}{\Gamma \vdash A}\ \textsc{BoolCases}$$

In Iter the matrix E is a literal equation. In HI the variable z is fresh for Γ, t and the passive parameters; in HE the variable a is fresh for Γ, A and the passive parameters. Here $1 = S0$ and $\mathrm{sg}(n) = \mathsf{R}(n, 0; u, v.1)$, so $\mathrm{sg}(0) = 0$ and $\mathrm{sg}(Sn) = 1$. There is no primitive hypothesis rule and no other induction rule. The *case-free progressive calculus* omits BoolCases.

HI is an unrestricted logical introduction; the only induction is Iter, at literal equations, in an elimination. The calculus with BoolCases is a complete presentation of iPRA (Theorem 10.89); Theorem 10.17 shows that BoolCases cannot be deleted.

10.2 Coordinatewise powers and truth transfer

This section draws on `research`, `compact_kernel`, `scope_kernel` and `checker_and_barriers`. One family of structures serves four results: equation induction does not imply numerical cases (Theorem 10.16), Boolean cases are not admissible in the progressive calculus (Theorem 10.17), and, in Section 10.3, positive axioms and conjunctive induction do not give a theory bi-interpretable with iPRA.

10.2.1 The powers

Definition 10.8 (Coordinatewise powers)**.** Let $\mathcal{N}$ be a finitary expansion of the standard structure $(\mathbb{N}, 0, S)$ by total functions and relations. The signature may be infinite; in particular it may contain a symbol for every primitive recursive description. $\mathcal{N}$ may be many-sorted, with nonempty sorts and with $(\mathbb{N}, 0, S)$ as a distinguished numerical sort. For a nonempty index set J, the structure $\mathcal{N}^{(J)}$ has, in each sort, the families $a = (a(i))_{i \in J}$ of elements of that

sort that are constant outside a finite subset of J. Functions are interpreted coordinatewise. A relation holds of families exactly when it holds at every coordinate. Equality is actual equality of families, and constants are constant families.

For $J = \{1, 2\}$ the structure is the full product $\mathcal{A} = \mathcal{N} \times \mathcal{N}$. For $J = \mathbb{N}$ it is the structure $\mathcal{E}$ of eventually constant sequences. For $i \in J$, the family e_i of the numerical sort has the value 1 at i and 0 elsewhere.

Finitely many families are simultaneously constant outside the union of their exceptional sets, so coordinatewise functions preserve the domain. Each element of $\mathcal{E}$ has a finite description, so $\mathcal{E}$ is countable when the sorts of $\mathcal{N}$ are.

Definition 10.9 (Conjunctive formulas)**.** The class $\mathcal{C}$ of *conjunctive formulas* is generated from atomic formulas and $\top$ by $\wedge$, $\exists$ and $\forall$, with unrestricted quantifier alternation. A conjunctive formula without $\forall$ is *primitive positive*. No disjunction, implication or negation occurs inside a conjunctive formula.

Theorem 10.10 (Conjunctive transfer)**.** *For every $\varphi(\vec{x}) \in \mathcal{C}$, every nonempty J and every tuple $\vec{a}$ of $\mathcal{N}^{(J)}$,*

$$\mathcal{N}^{(J)} \models \varphi(\vec{a}) \quad \Longleftrightarrow \quad \mathcal{N} \models \varphi(\vec{a}(i)) \ \text{ for every } i \in J.$$

Proof. Induction on the formula. Terms evaluate coordinatewise, so atomic formulas transfer by the definition of the relations and of equality. Conjunction is immediate.

Existential quantifier. If $\mathcal{N}^{(J)} \models \theta(b, \vec{a})$, the induction hypothesis gives $\mathcal{N} \models \theta(b(i), \vec{a}(i))$ at every i. Conversely, suppose $\mathcal{N} \models \exists y\, \theta(y, \vec{a}(i))$ at every i. There is a finite set $F \subseteq J$ such that $\vec{a}(i)$ is one fixed tuple $\vec{a}_\infty$ for all $i \notin F$. Choose a witness b_i for each of the finitely many $i \in F$ and one witness b_∞ for $\vec{a}_\infty$ if $J \neq F$. The family with the values b_i on F and b_∞ elsewhere belongs to $\mathcal{N}^{(J)}$, and the induction hypothesis gives $\mathcal{N}^{(J)} \models \theta(b, \vec{a})$. Only finitely many witnesses are chosen.

Universal quantifier. Suppose $\mathcal{N}^{(J)} \models \forall y\, \theta(y, \vec{a})$. Fix i and an element b of the relevant sort of $\mathcal{N}$. The constant family $\overline{b}$ is an element of $\mathcal{N}^{(J)}$; instantiate at $\overline{b}$ and apply the induction hypothesis at i to get $\mathcal{N} \models \theta(b, \vec{a}(i))$. Conversely, if every coordinate satisfies $\forall y\, \theta(y, \vec{a}(i))$, then for every family b every coordinate satisfies $\theta(b(i), \vec{a}(i))$, and the induction hypothesis gives $\mathcal{N}^{(J)} \models \theta(b, \vec{a})$. □

Corollary 10.11 (Regular axioms and conjunctive induction)**.** *Let J be nonempty.*

(i) $\mathcal{N}^{(J)}$ satisfies every sentence $\forall \vec{x}\, (\varphi \to \psi)$ and every sentence $\forall \vec{x}\, (\varphi \to \bot)$ with $\varphi, \psi \in \mathcal{C}$ that is true in $\mathcal{N}$.

(ii) For every $A(n, \vec{p}) \in \mathcal{C}$, with n of the numerical sort, $\mathcal{N}^{(J)}$ satisfies

$$\forall \vec{p} \Big[\big(A(0, \vec{p}) \wedge \forall n\, (A(n, \vec{p}) \to A(Sn, \vec{p})) \big) \to \forall n\, A(n, \vec{p}) \Big].$$

Consequently the contextual induction rule for matrices in $\mathcal{C}$ is sound in $\mathcal{N}^{(J)}$ under arbitrary retained assumptions.

Proof. (i) If $\varphi(\vec{a})$ holds in $\mathcal{N}^{(J)}$, then $\varphi(\vec{a}(i))$ holds in $\mathcal{N}$ at every i by Theorem 10.10; truth of the sentence in $\mathcal{N}$ gives $\psi(\vec{a}(i))$ at every i, and the converse direction of the theorem gives $\psi(\vec{a})$. For a negative constraint, a true antecedent in $\mathcal{N}^{(J)}$ gives a true antecedent in $\mathcal{N}$ at any one coordinate, which contradicts the constraint.

(ii) Fix $\vec{p}$ and suppose the base and the progress hypothesis hold in $\mathcal{N}^{(J)}$. Fix $i \in J$ and put $U_i = \{k : \mathcal{N} \models A(k, \vec{p}(i))\}$. Transfer of the base gives $0 \in U_j$ for every j. Let $k \in U_i$ and

let a be the family with the value k at i and 0 elsewhere. At i the formula $A(a(i), \vec{p}(i))$ holds by the choice of k, and at every other coordinate it is the transferred base; hence $\mathcal{N}^{(J)} \models A(a, \vec{p})$. The progress hypothesis gives $A(Sa, \vec{p})$, and transfer at i gives $k+1 \in U_i$. Ordinary induction in $\mathbb{N}$ gives $U_i = \mathbb{N}$ for every i, and the converse direction of the theorem gives $A(a, \vec{p})$ for every a.

For the contextual rule, fix an assignment that makes the retained assumptions true. The induction variable is fresh, so changing its value does not change their truth, and the argument applies under that assignment. The retained assumptions may be arbitrary formulas. Witnesses for A are not assumed to be terms. □

In the primitive recursive signature the corollary covers every defining equation, the injectivity of S, the separation $0 = Sx \to \bot$, every universal Horn sentence $\forall \vec{x}\,(E_1 \wedge \cdots \wedge E_r \to F)$ with equations E_k and an equation or $\bot$ as F that is true in $\mathbb{N}$, and induction for literal equations with parameters.

Definition 10.12. T_{alg} is intuitionistic first-order logic with equality in the signature with a symbol for every primitive recursive description, with all defining equations, the separation axioms and induction for every literal equation. Coverage is absent. For a structure $\mathcal{N}$ as in Theorem 10.8, $T_{\mathcal{C}}(\mathcal{N})$ is intuitionistic logic with equality together with all sentences of Theorem 10.11(i) that are true in $\mathcal{N}$ and all induction sentences of Theorem 10.11(ii); $T_{\mathrm{pp}}(\mathcal{N})$ is its restriction to primitive positive φ, ψ, A.

By Theorem 10.11, every $\mathcal{N}^{(J)}$ is a model of $T_{\mathcal{C}}(\mathcal{N})$, hence of $T_{\mathrm{pp}}(\mathcal{N})$, and, for the primitive recursive signature, of T_{alg} together with all universal Horn sentences true in $\mathbb{N}$.

Proposition 10.13 (What fails in a power)**.** *Let J have at least two elements and let the signature contain* sg*. In $\mathcal{N}^{(J)}$:*

(i) e_i is neither 0 nor a successor, so coverage $\forall x\,(x = 0 \vee \exists y\, x = Sy)$ fails;

(ii) the quantifier-free induction instance for $\varphi(n) := (\mathrm{sg}(n) = 0 \vee \mathrm{sg}(n) = 1)$ fails: $\varphi(0)$ and $\varphi(Sa)$ hold for every a, and $\varphi(e_i)$ fails;

(iii) no formula with finitely many parameters defines a linear order of the numerical sort when J is infinite (Theorem 10.19).

Proof. Every successor family is positive at every coordinate, and e_i is 0 at some coordinate and 1 at i. The family $\mathrm{sg}(Sa)$ is the constant 1, while $\mathrm{sg}(e_i) = e_i$ is neither the constant 0 nor the constant 1. For (iii), an order defined over a finite tuple of parameters is preserved by the involution of Theorem 10.19 that fixes the tuple, and an order-preserving involution of a linear order is the identity (Theorem 10.20). □

The same holds for the relation $x \le y$ of the primitive recursive signature: for $i \neq j$ neither $e_i \le e_j$ nor $e_j \le e_i$ holds, since each fails at one coordinate.

Remark 10.14. The missing feature is a disjunctive range assertion, not numerical computation. A primitive recursive characteristic of equality still reflects equality in a power: its value is the zero family exactly when its arguments agree at every coordinate. What fails is the assertion that the value is globally 0 or globally 1. In the same way, equation induction proves the algebraic identities $t(\mathrm{sel}(n, a, b)) = \mathrm{sel}(n, t(a), t(b))$ for each fixed term context t, and $\mathrm{sel}(b, 0, 1) = b$ for the Boolean value $b = \mathrm{sel}(n, 0, 1)$; these hold for the mixed values $(0, 1)$ and $(1, 0)$ of $\mathcal{A}$ and do not entail $b = 0 \vee b = 1$. Coordinatewise predecessor satisfies $P(0) = 0$ and $P(Sx) = x$ in every power, so adding valid predecessor equations does not restore coverage.

Remark 10.15. The product $\mathcal{A}$ with literal-equation induction is in `scope_kernel` and `checker_and_barriers`; `research` attributes the product model of non-coverage to an earlier draft. The structure $\mathcal{E}$ with induction for equations and for finite conjunctions of atoms, universal Horn sentences and arbitrary finitary expansions is in `research`. Primitive positive transfer, the universal quantifier and the many-sorted form are in `compact_kernel`. The statement for an arbitrary index set is the same proof.

10.2.2 Equation induction does not replace numerical cases

Consider the *direct-target* replacement of case introduction:

$$\frac{\Gamma,\ B(t,0) \vdash B(t,z)}{\Gamma \vdash D_{x,y}B(x,y)} \qquad z \text{ fresh for } \Gamma, t \text{ and the parameters of } B.$$

It is sound for the ordinary reading of $D_{x,y}$ in every structure, with no coverage axiom. Let N^{dt} be N (or N_{p}) with DI by cases replaced by this rule.

Theorem 10.16 (`scope_kernel`)**.** *Every theorem of* N^{dt} *is valid in every power* $\mathcal{N}^{(J)}$ *of the standard model of the iterator language, under the ordinary reading. Write* $1 = S0$ *and define*

$$E(n,y,p) := \big(\mathrm{sel}(p,\, \mathrm{sel}(n,y,1),\, \mathrm{sel}(n,1,0)) = 0\big), \qquad K(n,p) := Q_y E(n,y,p),$$
$$M(p) := Q_n K(n,p), \qquad F := Q_p M(p).$$

The closed formula F *is derivable in* N *without the progress rule and is false in* $\mathcal{A}$*. Hence* N^{dt} *does not derive* F*.*

Proof. Soundness. Interpret $Q_y B$ and $D_{x,y}B$ by their ordinary readings. Hyp, transport, QE and DE are sound in every structure with actual equality. The computation equations hold coordinatewise. The progress rule is sound by Theorem 10.11(ii). The direct-target rule is sound because its fresh variable already ranges over the whole domain.

Derivation of F *in* N*.* A closure $Q_y B(y)$ follows from a derivation of $B(Sz)$ with z fresh: DI by cases with an unused witness gives $D_{u,y}B(y)$, and DE with Hyp gives $Q_y B(y)$. At $n = Sz$ the computation equations give $\mathrm{sel}(Sz,y,1) = 1$ and $\mathrm{sel}(Sz,1,0) = 0$, so $E(Sz,y,p)$ is interderivable with $\mathrm{sel}(p,1,0) = 0$, independently of y. Hence $E(Sz,0,p) \vdash E(Sz,Sw,p)$ by transport, and $K(Sz,p)$ is derivable. This is the successor instance of $M(p)$, so $M(p)$ is derivable for a free p; its instance $M(Sw)$ is the successor instance of F. Only case introduction, DE, QE, the computations and transport are used.

F is false in $\mathcal{A}$. Let $p_* = (0,1)$. Coordinatewise computation gives

$$\mathcal{A} \models E(n,y,p_*) \quad\Longleftrightarrow\quad n_1 = 0,\ n_2 > 0,\ y_1 = 0.$$

Hence $K(0,p_*)$ holds, since its antecedent $E(0,0,p_*)$ is false, and $K((0,1),p_*)$ fails: its antecedent holds at $y = 0$ and its consequent fails at $y = (1,0)$. So $M(p_*)$ is false. At $p = 0$ the formula $E(0,y,0)$ is $y = 0$, so $K(0,0)$, which reads $0 = 0 \to \forall y\,(y = 0)$, is false and $M(0)$ is true. Therefore F, which reads $M(0) \to \forall p\, M(p)$, is false. □

The theorem excludes this replacement with the identity translation of formulas. It does not show that every interpretation of the weaker calculus in iPRA fails, and it does not show that no restricted form of numerical cases suffices.

10.2.3 Boolean cases are not admissible in the progressive calculus

Theorem 10.17 (`checker_and_barriers`)**.** *Let C and Z be the terms $C(n,a,b) = \mathsf{R}(n,a;u,v.b)$ and $Z(n) = \mathsf{R}(n,1;u,v.0)$, so that $C(0,a,b) = a$, $C(Sn,a,b) = b$, $Z(0) = 1$ and $Z(Sn) = 0$. For a numerical term t put*

$$F(t) := \mathsf{H}_{u,y}\,\mathsf{H}_{v,z}(C(t,Z(y),z) = 0),$$

with u, v unused. The case-free progressive calculus of Theorem 10.7 derives $F(0)$ and $F(1)$ and does not derive $F(\mathrm{sg}(t))$. Hence BoolCases*, with an arbitrary object formula as conclusion, is not admissible in the case-free calculus, in the presence of all primitive recursive symbols, their defining equations and contextual induction for literal equations.*

Proof. The endpoints. The derivations use the derived identity $\Gamma, A \vdash A$ of Theorem 10.83, which needs no cases. At $t = 0$ the inner matrix is interderivable with $Z(y) = 0$, independently of z. Its progress step is therefore an identity, and HI gives $\mathsf{H}_{v,z}(C(0,Z(y),z) = 0)$ for arbitrary y. The outer HI ignores its local assumption. At $t = 1$ the inner matrix is interderivable with $z = 0$, independently of y. For the outer HI, open the assumed inner formula at y by HE, introduce the inner formula at Sy by HI with the same witness, convert its local assumption to the matrix at y by the computation equations, apply Step, and convert back.

The model. Let $\mathcal{A}$ be the product of the standard model of the primitive recursive signature. Under the readings of Theorem 10.7, Tr, HI, HE and Step are logical rules, valid in every structure with actual equality, and Iter is induction for a literal equation, valid by Theorem 10.11(ii). The computation equations hold. So every theorem of the case-free calculus is valid in $\mathcal{A}$.

Let $e = (0,1)$, so $\mathrm{sg}(e) = e$. For $y = (y_1, y_2)$ and $z = (z_1, z_2)$ the inner equation $C(e,Z(y),z) = 0$ says $(Z(y_1), z_2) = (0,0)$. If $y_1 = 0$ it is false for every z, so the inner progress statement is vacuously true. If $y_1 > 0$ it says $z_2 = 0$; it holds at $z = (0,0)$ and fails at $Sz = (1,1)$, so the inner statement is false. Thus the inner formula is true at $y = (0,0)$ and false at $Sy = (1,1)$, which refutes the outer progress statement for every u. Hence $\mathcal{A} \not\models F(e)$, and $F(\mathrm{sg}(t))$ is not derivable.

Non-admissibility. From the assumption $\mathrm{sg}(t) = 0$, transport of $F(0)$ gives $F(\mathrm{sg}(t))$; from $\mathrm{sg}(t) = 1$, transport of $F(1)$ gives the same. An admissible BoolCases would derive $F(\mathrm{sg}(t))$. □

Proposition 10.18 (The binder depth is sharp for this separation)**.** *Let $G(t)$ be an equation or a formula $\mathsf{H}_{x,y}E(x,y,t)$ with E a literal equation. If $G(0)$ and $G(1)$ hold in the standard model, then $G((0,1))$ holds in $\mathcal{A}$.*

Proof. For an equation this is coordinatewise evaluation. Otherwise let a_0 and a_1 be standard witnesses for $t = 0$ and $t = 1$. If $E((a_0,a_1),(y_1,y_2),(0,1))$ holds in $\mathcal{A}$, then $E(a_0,y_1,0)$ and $E(a_1,y_2,1)$ hold in $\mathbb{N}$, so $E(a_0,y_1+1,0)$ and $E(a_1,y_2+1,1)$ hold, which is E at Sy in $\mathcal{A}$. □

The first failure arises when an implication tests a compound premise: one coordinate makes an inner implication vacuous while the other would fail it. Theorem 10.17 rejects the deletion of BoolCases. It is not a statement about other logical bases, other primitive operations or other interpretations; the stronger statement about bi-interpretations of the case-free calculus is Theorem 10.25.

10.3 Symmetry

This section draws on `research` and `compact_kernel`; `checker_and_barriers` applies the result to the case-free progressive calculus. `research` attributes the method, a finite-parameter

involution against a definable linear order, to earlier drafts on unary arithmetic. The obstruction of Theorem 10.13 concerns the original domain. An interpretation on a different definable domain is not touched by it; the symmetry argument handles that freedom.

Lemma 10.19 (Finite parameters leave an involution)**.** *Let J be infinite. For every finite tuple $\vec{c}$ of $\mathcal{N}^{(J)}$, the expansion $(\mathcal{N}^{(J)}, \vec{c})$ has an automorphism of order two that is not the identity on the numerical sort.*

Proof. Outside a finite set $F \subseteq J$ every component of $\vec{c}$ is constant. Choose distinct $i, j \notin F$ and exchange the coordinates i and j in every family of every sort. The exchange preserves the domain, commutes with every coordinatewise function and preserves every coordinatewise relation, whatever the size of the signature. It fixes $\vec{c}$, has order two, and moves e_i to e_j. □

Lemma 10.20 (Order rigidity)**.** *Every classical model of* iPRA *has a parameter-free definable strict linear order, for example $x < y :\Leftrightarrow Sx \dot{-} y = 0$ with truncated subtraction. An automorphism of finite order of a linearly ordered structure that preserves the order is the identity.*

Proof. The order laws are theorems of primitive recursive arithmetic with quantifier-free induction. If $a < \sigma(a)$, repeated application gives $a < \sigma(a) < \cdots < \sigma^r(a) = a$ for the order r of σ; the case $\sigma(a) < a$ is symmetric. Linearity gives $\sigma(a) = a$. □

Lemma 10.21 (Automorphisms under a bi-interpretation)**.** *Let a structure M interpret a structure N by an ordinary interpretation with parameters among $\vec{c}$. Every automorphism of M that fixes $\vec{c}$ acts coordinatewise on representing tuples and induces an automorphism of N; this is a group homomorphism $\mathrm{Aut}(M/\vec{c}) \to \mathrm{Aut}(N)$. If in addition N interprets a copy of M, and an isomorphism η of M onto that copy is definable in M, and $\vec{c}$ contains the parameters of η and representatives in M of the parameters of the second interpretation, then the homomorphism is injective.*

Proof. The domain, the equality and the graphs of the interpretation are definable over $\vec{c}$, so they are preserved by $\sigma \in \mathrm{Aut}(M/\vec{c})$; this gives the induced automorphism $\widehat{\sigma}$ and the homomorphism. The automorphism $\widehat{\sigma}$ fixes the parameters of the second interpretation, since σ fixes their representatives, and so induces an automorphism $\widehat{\widehat{\sigma}}$ of the copy of M. Since η is definable over parameters fixed by σ, $\eta(\sigma(a)) = \widehat{\widehat{\sigma}}(\eta(a))$. If $\widehat{\sigma}$ is the identity, so is $\widehat{\widehat{\sigma}}$, hence $\eta(\sigma(a)) = \eta(a)$, and injectivity of η gives $\sigma(a) = a$. Only this one composite is used. The interpretations may have any finite dimension, quotient equality and domains different from the original ones. □

Theorem 10.22 (Symmetry obstruction)**.** *Let J be infinite and let T be a theory, in the language of a structure $\mathcal{N}$ as in Theorem 10.8, that has $\mathcal{N}^{(J)}$ as a model. Then T and* iPRA *are not bi-interpretable in the sense of Theorem 10.1. Arbitrary formula-defined domains, graphs and quotient equality are allowed, in any finite dimension and over finite products of the sorts. Finite parameter tuples are allowed, including parameters of the return isomorphism, provided the interpretations and the isomorphism exist in $\mathcal{N}^{(J)}$; this covers a parameter condition that T proves to be inhabited and under which T proves all the required facts.*

Proof. Apply a hypothetical bi-interpretation in $\mathcal{N}^{(J)}$, which yields a classical model $\mathcal{M}$ of iPRA. Let $\vec{c}$ collect the parameters of the first interpretation, of the return isomorphism, and representatives in $\mathcal{N}^{(J)}$ of the parameters of the second interpretation. Theorem 10.19 gives a non-identity involution σ of $(\mathcal{N}^{(J)}, \vec{c})$. Its image in $\mathrm{Aut}(\mathcal{M})$ has order dividing two and preserves the definable order, so it is the identity by Theorem 10.20. By Theorem 10.21 the homomorphism is injective, so σ is the identity on every sort, which contradicts its choice on the numerical sort. Faithfulness of the action on the other sorts is not needed. □

Corollary 10.23. *None of the following theories is bi-interpretable with* iPRA *in the sense of Theorem 10.1:*

(i) T_{alg}*, also with every set of universal Horn sentences true in* $\mathbb{N}$ *adjoined;*

(ii) for every finitary expansion $\mathcal{N}$ *of* $(\mathbb{N}, 0, S)$*, the theory of all universal Horn sentences true in* $\mathcal{N}$ *with induction for atomic formulas or for finite conjunctions of atomic formulas;*

(iii) $T_{\mathrm{pp}}(\mathcal{N})$ *and* $T_{\mathcal{C}}(\mathcal{N})$*, and every subtheory of them, recursively axiomatized or not;*

(iv) the many-sorted theories $T_{\mathcal{C}}(\mathcal{A}')$ *for a many-sorted* $\mathcal{A}'$ *with numerical sort* $(\mathbb{N}, 0, S)$*, with induction on the numerical sort for conjunctive matrices whose parameters and witnesses may range over the other sorts.*

Proof. Each theory is true in $\mathcal{E}$ by Theorem 10.11. In the many-sorted case the family varied in the induction argument belongs to the numerical sort and the parameters of the other sorts stay fixed. □

Remark 10.24 (Function objects)*.* The many-sorted case applies when a sort consists of numerical functions and the signature has a total application operation. Extensionality, $(\forall n\ \mathrm{app}(f, n) = \mathrm{app}(g, n)) \to f = g$, has a conjunctive antecedent and is preserved. Adding function objects, application, extensionality and equational combinator laws does not by itself remove the symmetry.

Corollary 10.25 (The case-free progressive calculus)**.** *Every rule of the case-free progressive calculus is sound in* T_{alg} *under the readings of Theorem 10.7;* ITER *is induction for a literal equation. Hence* $\mathcal{E}$ *is a model of its ordinary first-order realization, and that realization is not bi-interpretable with* iPRA *in the sense of Theorem 10.1.*

Remark 10.26 (Scope)*.* Theorem 10.22 is about ordinary model-theoretic bi-interpretations, or about a native proposal whose claimed interpretations are proved to induce such interpretations under the displayed readings. Soundness of a native calculus in an ordinary theory does not show that the calculus derives the logical rules of that theory, and a proof translation that changes the meaning of the connectives need not give an ordinary interpretation; the theorem does not address such translations. It does not exclude equiconsistency, one-way interpretations, disjunctive axioms such as coverage or a Boolean range assertion, non-regular rules, or expansions that are not interpreted coordinatewise. A parameter condition that excludes $\mathcal{E}$ and is not proved inhabited there is outside the claim. Changing a term-defined operation to a positive existential graph does not escape the theorem, since existential witnesses and conjunctive induction of arbitrary alternation are allowed.

Remark 10.27*.* `research` proves Theorem 10.22 for T_{alg}, universal Horn sentences and conjunctions of atoms. `compact_kernel` proves it for T_{pp}, for $T_{\mathcal{C}}$ and for many sorts, with the order $x \le y :\Leftrightarrow \exists z\,(x + z = y)$ in place of the one in Theorem 10.20. `checker_and_barriers` restates the argument for Theorem 10.25.

10.4 Majorants: almost unary term languages

This section draws on `counter_congruence`, `research`, `local_scopes`, `adaptive_seed` and `uniform_predecessor`. All statements concern the standard interpretation of fixed finite terms. The bounding functions belong to the metatheory; they are not asserted to be terms of the reduced languages.

10.4.1 Zero-seeded recursion is monotone

Proposition 10.28 (`counter_congruence`)**.** *Every function denoted by a term of the zero-seeded language of Theorem 10.3(a) is nondecreasing in each coordinate. Hence no term denotes the zero test $z(0) = 1$, $z(Sn) = 0$, and the terms do not denote all primitive recursive functions.*

Proof. Induction on terms. Variables, 0 and S are monotone. For $\mathsf{L}(n; u, v.h)$ the update $h(u, v, \vec{p})$ is coordinatewise nondecreasing by the induction hypothesis. Fix $\vec{p}$ and write $r_0 = 0$, $r_{k+1} = h(k, r_k, \vec{p})$. Then $r_0 \le r_1$, and $r_k \le r_{k+1}$ gives $r_{k+1} = h(k, r_k, \vec{p}) \le h(k+1, r_{k+1}, \vec{p}) = r_{k+2}$. So the value is nondecreasing in the counter. Induction on k gives $r_k(\vec{p}) \le r_k(\vec{p}\,')$ for $\vec{p} \le \vec{p}\,'$ coordinatewise. Substitution of the counter term and of parameter terms preserves monotonicity by the induction hypothesis. The zero test is not monotone at $0 < 1$. □

The statement is formalized as `ZTm.mono` and `ZTm.no_zero_test` in the module `ZeroSeedObstruction`; `Tm.val_zf` records that $\mathrm{sel}(x, 1, 0)$ of the iterator language is the zero test. The proposition excludes the direct term compiler only. The tied-seed language of Theorem 10.3(b) escapes it, since Z of Theorem 10.4 is the zero test.

10.4.2 Bounding coordinates

Definition 10.29. A function $f\colon \mathbb{N}^r \to \mathbb{N}$ has the *bounding coordinate* i if there is a nondecreasing unary primitive recursive function F with $f(x_1, \dots, x_r) \le F(x_i)$ for all arguments. A function is *almost unary* if it is globally bounded or has a bounding coordinate.

The name is the one used in clone theory for functions bounded by a unary function of one coordinate, whose closure under composition is standard (Pinsker, *Clones containing all almost unary functions*, Algebra Universalis 51 (2004)). The primitive recursive majorants are the additional content here.

Lemma 10.30 (Envelopes and composition)**.** *(i) For a primitive recursive map $g\colon \mathbb{N}^r \to \mathbb{N}^s$ the function $\widehat{g}(z) = \max\{g(\vec{z})_k : z_1, \dots, z_r \le z,\ k \le s\}$ is a nondecreasing unary primitive recursive function that bounds every output coordinate of g whenever all inputs are at most z. In particular a unary bound F may be replaced by its nondecreasing envelope $\widehat{F}(z) = \max_{m \le z} F(m)$.*

(ii) Constants, projections, unary primitive recursive functions and globally bounded functions are almost unary.

(iii) If f and $g_1, \dots, g_r$ are almost unary, so is $f(g_1(\vec{x}), \dots, g_r(\vec{x}))$.

Proof. (i) and (ii) are immediate. For (iii), if f is globally bounded so is the composite. If $f \le F(x_j)$, only g_j matters: a constant bound c of g_j gives the constant bound $F(c)$, and $g_j \le N(x_i)$ gives $F(N(x_i))$, by monotonicity of F. □

10.4.3 Single-body constructors

Definition 10.31 (`uniform_predecessor`)**.** A *single-body constructor scheme* is given by integers $a, b, d, q \ge 1$ and fixed primitive recursive maps

$$A_i\colon \mathbb{N} \to \mathbb{N}^q\ (i \le a), \quad U_j\colon \mathbb{N}^{d+2} \to \mathbb{N}^q\ (j \le b), \quad F_0\colon \mathbb{N}^a \to \mathbb{N}^d, \quad F_1\colon \mathbb{N}^b \to \mathbb{N}^d, \quad G\colon \mathbb{N}^d \to \mathbb{N}.$$

Its supplied subdescriptions are a counter term t and one body $h(\vec{u}, \vec{p})$ with q local variables $\vec{u}$ and passive variables $\vec{p}$. With $n = t(\vec{p})$, the constructor denotes $G(\vec{r}_n)$, where

$$\vec{r}_0 = F_0\big(h(A_1(n), \vec{p}), \dots, h(A_a(n), \vec{p})\big), \tag{10.3}$$

$$\vec{r}_{m+1} = F_1\big(h(U_1(n, m, \vec{r}_m), \vec{p}), \dots, h(U_b(n, m, \vec{r}_m), \vec{p})\big). \tag{10.4}$$

The maps are fixed before the body is supplied. No passive parameter and no retained register enters F_0, F_1 or G directly: passive data and retained state reach the seed and update combiners only through evaluations of the one body h. The target counter n and the elapsed counter m may enter the probe locations. The *single-body term language* has variables, 0, any primitive operations that are almost unary, composition, and the constructors of all schemes, arbitrarily nested.

The tied-seed recursor is the scheme $d = 1$, $q = 2$, $a = b = 1$, $A_1(n) = (0, 0)$, $U_1(n, m, r) = (m, r)$, $F_0(z) = z + 1$, $F_1(z) = z$, $G(r) = r$.

Theorem 10.32 (Single-body information barrier)**.** *Every term of the single-body term language denotes an almost unary function.*

Proof. Induction on descriptions; the local and passive variables of a body are independent inputs for the induction hypothesis. The primitive cases hold by assumption and composition is Theorem 10.30(iii). Consider a constructor, and fix envelopes f_0, f_1, g of F_0, F_1, G by Theorem 10.30(i).

h is bounded by a constant c. Every probe value is at most c, so every register at every stage is at most $\max(f_0(c), f_1(c))$, and the result is bounded by g of that constant.

$h(\vec{u}, \vec{p}) \le H(p_i)$ for a passive coordinate. All seed and update probes obey this bound, whatever their local arguments. The result is at most $g\big(\max(f_0(H(p_i)), f_1(H(p_i)))\big)$, a nondecreasing primitive recursive function of p_i alone.

$h(\vec{u}, \vec{p}) \le H(u_\ell)$ for a local coordinate. Choose nondecreasing primitive recursive envelopes α and υ such that all coordinates of every $A_i(n)$ are at most $\alpha(n)$, and all coordinates of every $U_j(n, m, \vec{r})$ are at most $\upsilon(s)$ whenever $n, m, r_1, \dots, r_d \le s$. Define

$$w(0, n) = f_0(H(\alpha(n))), \qquad w(m + 1, n) = \max\Big(w(m, n),\ f_1\big(H(\upsilon(n + m + w(m, n)))\big)\Big).$$

Induction on m shows that every coordinate of $\vec{r}_m$ is at most $w(m, n)$: at the base the probe arguments are bounded by $\alpha(n)$; at the step the sum $n+m+w(m, n)$ bounds both counters and every register, so the probes are bounded by $H(\upsilon(n + m + w(m, n)))$ and the new registers by f_1 of that. Hence the result is at most $g(w(n, n))$, and $K(n) = \max_{z \le n} g(w(z, z))$ is a nondecreasing unary primitive recursive bound, independent of every passive parameter.

In the last case apply the induction hypothesis to the counter term: a constant bound c of t gives the bound $K(c)$, and $t(\vec{p}) \le N(p_i)$ gives $K(N(p_i))$. The cases exhaust the induction hypotheses for h. □

Remark 10.33 (Effectiveness)*.* Given bounding-coordinate or global-bound witnesses for the primitive operations of a description, the proof constructs a coordinate and a majorant effectively. A bound of an arbitrary globally bounded primitive recursive function cannot be computed from its bare description: for a machine e, the function $b_e(x, y)$ equal to the first halting time of e if it is at most $\max(x, y)$ and 0 otherwise is globally bounded, and a uniform extractor of bounds would decide halting. Unary primitives have computable envelopes and Boolean predicates have the bound 1, so these need no witness. The existence statement of the theorem needs no witnesses.

10.4.4 Adaptive query circuits

Definition 10.34 (`adaptive_seed`)**.** A *query circuit* P is a finite acyclic circuit with no input nodes. Its nodes are numerals, applications of fixed primitive recursive functions to earlier nodes, and queries $Q(a, b)$ whose two arguments are earlier nodes; one node is the output.

For an oracle $Q\colon \mathbb{N}^2 \to \mathbb{N}$ the output is P^Q. Later queries may depend on earlier answers. For an update term $h(u, v, \vec{p})$ put $h_{\vec{p}}(a, b) = h(a, b, \vec{p})$; the *tied-seed recursor with circuit* P is

$$\mathsf{R}_P(0; u, v.h) = P^{h_{\vec{p}}}, \qquad \mathsf{R}_P(n+1; u, v.h) = h(n, \mathsf{R}_P(n; u, v.h), \vec{p}).$$

The circuit reaches $\vec{p}$ only through queries to $h_{\vec{p}}$; it receives no separate initializer, no counter and no passive parameter. The *circuit term language* has constants, projections, any primitive operations that are almost unary, substitution, and the constructors R_P for all circuits P, arbitrarily nested. The combining operations inside a circuit are arbitrary primitive recursive functions.

The tied-seed recursor B is the circuit with one query $Q(0, 0)$ followed by successor. A circuit may, for example, compute $q = h(0, 0, \vec{p})$ and return $q + h(q, q+1, \vec{p})^2$.

Lemma 10.35. *Let P be a query circuit.*

(i) There is a unary primitive recursive C_P such that $P^Q \le C_P(M)$ whenever $Q(a, b) \le M$ for all a, b.

(ii) If $Q(a, b) \le F(a)$ for all a, b, with F a fixed unary primitive recursive function, then P^Q is bounded by a constant that depends on P and F only. The same holds with $F(b)$.

Proof. Traverse the circuit in topological order. For (i) give each numeral its value, each query the bound M, and each operation node the envelope $\widehat{g}$ of the maximum of the bounds of its inputs; every node bound is a unary primitive recursive function of M. For (ii) use constant bounds throughout: at a query, apply $\widehat{F}$ to the constant bound of the controlling argument, which is already available. □

Theorem 10.36 (Tied-seed circuit obstruction)**.** *Every term of the circuit term language denotes an almost unary function.*

Proof. Induction on terms, with Theorem 10.30 for the primitives and for substitution. Consider $\mathsf{R}_P(n; u, v.h)$ and write $r_j(\vec{p})$ for its state at the counter j.

If $h \le c$ or $h(u, v, \vec{p}) \le F(p_i)$ for a passive coordinate, write M for c or $F(p_i)$. Theorem 10.35(i) gives $r_0 \le C_P(M)$, and $r_{j+1} \le M$, so $\max(C_P(M), M)$ bounds the result independently of the counter.

If $h(u, v, \vec{p}) \le F(u)$, Theorem 10.35(ii) bounds r_0 by a constant b_0, and $r_{j+1} \le F(j)$. The nondecreasing function $M(n) = \max(b_0, \widehat{F}(n))$ bounds r_n for every n.

If $h(u, v, \vec{p}) \le F(v)$, Theorem 10.35(ii) again bounds r_0 by a constant b_0. Put $K(z) = \max(z, \widehat{F}(z))$. Then K is nondecreasing, $K \ge \mathrm{id}$, and induction gives $r_n \le K^n(b_0)$; the function $n \mapsto K^n(b_0)$ is primitive recursive and nondecreasing.

In the last two cases compose the bound in n with the bound of the counter term, as in Theorem 10.30(iii). □

Corollary 10.37 (The tied-seed language)**.** *Every term of the tied-seed language of Theorem 10.3(b), and of its unary-update variant, denotes an almost unary function. This remains true after adjoining all unary primitive recursive functions and any operations that are globally bounded or have a bounding coordinate, for example a primitive characteristic of equality with values in $\{0, 1\}$, predecessor, zero tests and minimum.*

Proof. The language is a sublanguage of both the single-body and the circuit term language. Directly: if $h \le c$ the values are at most $c + 1$; if $h \le F(p_i)$ they are at most $1 + F(p_i)$; if $h \le F(u)$ the value at the counter m is at most $1 + F(m)$; if $h \le F(v)$ it is at most $\kappa(m)$ with $\kappa(0) = 1 + F(0)$ and $\kappa(m+1) = \max(\kappa(m), F(\kappa(m)))$. □

Corollary 10.38 (What almost unary terms cannot denote)**.** *No term of the single-body term language or of the circuit term language denotes addition, maximum, the selector with* $\mathrm{sel}(0,a,b) = a$ *and* $\mathrm{sel}(Sn,a,b) = b$*, or an injection* $\mathbb{N}^2 \to \mathbb{N}$*. In particular the tied-seed language has no such term and cannot compile the initialized iterator, since* $I(b,a;\, v.\, Sv) = a+b$*.*

Proof. For addition or maximum, fix the supposed bounding coordinate and increase the other argument; a global bound fails likewise. For the selector, a bound in the control fails at the control 0 as a grows; a bound in a fails at $a = 0$ and control 1 as b grows; a bound in b fails at $b = 0$ and control 0 as a grows. Additional dummy coordinates can be fixed. For an injection, fixing the bounding coordinate at 0 leaves infinitely many pairs whose values lie in one finite interval. □

Remark 10.39 (The boundary of the hypotheses)*.* The data interface is what the theorems restrict. (i) If the initializer circuit receives one further passive input y, the update $h(u,v,x,y) = x$ and the initializer $y + h(0,0,x,y)$ give $x+y$ at the counter 0. (ii) A circuit with two independent update oracles queries the constant updates x and y and adds the answers. (iii) If a passive initializer enters a register directly, $r_0 = a$, $r_{m+1} = Sr_m$ computes $a+n$. (iv) Direct register retention escapes Theorem 10.32 even with a tied seed: $r_0 = h(0,\vec{p})$, $r_{m+1} = Sr_m$ with the constant body $h(u,a) = a$ computes $a+n$; here the old register reaches the update combiner without passing through the body. (v) Finiteness of the circuit matters: for the oracle that is 1 exactly at $(u,v) = (x,y)$, every answer is at most 1, and an unbounded search over all pairs finds (x,y) and returns $x+y$; such a search is not a fixed finite circuit and is not total for all oracles. These examples mark the boundary. None of them is claimed to give a complete basis or a smaller checker.

Remark 10.40 (Scope and sources)*.* The theorems are term-definability obstructions. They do not exclude quantified graph definitions, a relational interpretation, a different interpretation domain or additional axioms; an independent unbounded initializer is one escape and is not asserted to be the only one. The one-coordinate bound for B and the exclusion of addition and selection are in `counter_congruence`; `research` rederives them for F and B; `local_scopes` adds the bounded-valued primitives. Theorem 10.32 is from `uniform_predecessor` and Theorem 10.36 from `adaptive_seed`. The two classes are incomparable: the first has register vectors, several probes in the update and counter-dependent probe locations, but a non-adaptive seed; the second has an adaptive seed, but a scalar state and the fixed update interface.

10.5 Finite-dimensional term interpretations

This section draws on `adaptive_seed` and `local_scopes`, which prove the same theorem independently. It turns the bound of Section 10.4 into an obstruction to interpretations that use finite tuples and a quotient equality, as long as the interpreted addition is given by a tuple of terms. The theorem is about functions on $\mathbb{N}$; the domain and the equivalence need not be definable.

Theorem 10.41 (Finite quotient from coordinate bounds)**.** *Let* $d \geq 1$*,* $D \subseteq \mathbb{N}^d$*, let* $\sim$ *be an equivalence relation on* D *and* $e \in D$*. Let* $f\colon D \times D \to D$ *satisfy*

$$f(e,z) \sim z \sim f(z,e) \qquad (z \in D).$$

Suppose that for each output coordinate i *there are an input coordinate* $j_i \leq d$ *and a nondecreasing unary* g_i *such that, uniformly on* $D \times D$*, either* $f(x,y)_i \leq g_i(x_{j_i})$ *or* $f(x,y)_i \leq g_i(y_{j_i})$*, or that* $f(x,y)_i$ *has a constant bound. Then* $D/\!\sim$ *is finite. More precisely, with* $L(z) = f(e,z)$ *and* $R(z) = f(z,e)$*, a composition of at most* $\ell_d = \lceil \log_2(d+1) \rceil$ *maps chosen from* L, R *sends*

every element of D to a $\sim$-equivalent element of one finite box $\prod_{i\le d}\{0,\dots,C_i\}$, and $D/\sim$ has at most $\prod_i(C_i+1)$ elements.

Proof. A coordinate bounded through an x-input has the constant bound $g_i(e_{j_i})$ under L and the bound $g_i(z_{j_i})$ under R; for a coordinate bounded through a y-input the roles are exchanged; a constantly bounded coordinate is bounded under both. Regard $\{1,\dots,d\}$ as a set of states that track dependencies of bounds. A letter sends a state i to the state j_i if it leaves the dependence, and to an absorbing terminal state if it gives a constant bound. Each state is terminated by at least one of the two letters.

Choose the letters from the outside of the composition inward. Let S be the current set of live states, initially all d states. The number of states of S terminated by L plus the number terminated by R is at least $|S|$, so one letter terminates at least $\lceil |S|/2\rceil$ of them. Choose it and replace S by the set of images of the survivors; each survivor has one image, so the new set has at most $\lfloor |S|/2\rfloor$ elements. After at most ℓ_d choices S is empty, since $2^{\ell_d} > d$.

Let the chosen letters be $V_1,\dots,V_k$ and $W = V_1\circ\cdots\circ V_k$, so that V_k is applied first. By induction on the number of letters, each coordinate of the partial composition $V_1\circ\cdots\circ V_m$ is either uniformly bounded or bounded by a composition of the functions g_i along one live dependency path, applied to a live coordinate of the argument; appending the next letter on the inside replaces that coordinate by its bound, which is legitimate because the g_i are nondecreasing. When the path terminates the bound is a constant. The empty final set gives constants $C_1,\dots,C_d$ with $W(D)\subseteq\prod_i\{0,\dots,C_i\}$.

Both L and R map D into D and send every element to an equivalent one, so $W(z)\sim z$ by transitivity. Every class meets the finite set $D\cap\prod_i\{0,\dots,C_i\}$. □

Associativity, commutativity and compatibility of f with $\sim$ are not used. A unary bound may be replaced by its nondecreasing envelope, so monotonicity is no extra assumption for term majorants. For $d=0$ the domain has at most one element.

Proposition 10.42 (The word length is sharp)**.** *For every $d\ge 1$ there are $D=\mathbb{N}^d$, the universal equivalence, e the zero tuple and an f whose coordinates are projections or constants, such that the shortest composition of L,R with finite image has length exactly $\lceil\log_2(d+1)\rceil$.*

Proof. Let $k=\lfloor\log_2 d\rfloor$. If $k=0$ then $d=1$; take $f(x,y)=x$, so R is the identity and L is constant, and the minimum is 1. Let $k\ge 1$. Label 2^k of the coordinates by words $w\in\{L,R\}^k$ and let $\sigma(w)$ be w with its first letter removed and R appended. Put $f(x,y)_w = y_{\sigma(w)}$ if w starts with L and $f(x,y)_w = x_{\sigma(w)}$ if w starts with R; the remaining coordinates are 0. Under a letter, the state w survives exactly when the letter is its first letter, and then moves to $\sigma(w)$. For a word $\alpha\in\{L,R\}^k$, read from the outside inward, the state α survives all k letters, and the coordinate α of the composition is the projection to the coordinate R^k of the argument. So no word of length k has finite image, and neither has a shorter word, since extending a word with finite image on the inside keeps the image finite. The word L^{k+1} has finite image: a state with an R among its k letters is terminated when that letter is reached, and the state L^k has become R^k after k letters and is terminated by the last L. The minimum is $k+1=\lceil\log_2(d+1)\rceil$. □

Corollary 10.43 (No finite-dimensional term interpretation of iPRA)**.** *Let T be a theory in a numerical language whose standard model on $\mathbb{N}$ satisfies T and in which every term denotes an almost unary function; for example a theory true in the standard tied-seed algebra, in the single-body algebra or in the circuit algebra. There is no interpretation of* iPRA *in T such that the interpreted domain is a nonempty subset of a finite power of the source domain, the interpreted equality is an arbitrary equivalence relation on it, and the interpreted addition has one fixed tuple of terms as realizers. Fixed numerical parameters are allowed,*

and successor may be interpreted in any way consistent with the axioms. In particular there is no bi-interpretation with these properties.

Proof. Apply the interpretation in the standard model. Let e represent the interpreted zero. The two zero laws of addition are theorems of iPRA, so the addition tuple satisfies the unit hypothesis of Theorem 10.41; each of its coordinate terms is almost unary in the $2d$ input variables, and a majorant in a fixed parameter is a constant. So the interpreted quotient is finite. The interpreted successor is injective on the quotient and its image omits the class of zero, which is impossible on a finite nonempty set. □

Remark 10.44 (Limits). (i) Both unit laws are needed: on $\mathbb{N}$ with equality, $f(x, y) = x$ satisfies $f(z, e) = z$ and the coordinate bound, and the quotient is infinite. (ii) The realizers must be terms. With decidable equality to zero the quantifier-free graph $(x = 0 \wedge z = y) \vee (x \neq 0 \wedge z = x)$ is total and single-valued and defines an associative operation on $\mathbb{N}$ with the two-sided unit 0, which selects the first nonzero argument; it is not almost unary. Likewise $\max(x, y)$ has the unit 0 and is defined by cases that choose different controlling coordinates. Neither example is an interpretation of iPRA; they show that the relational route is not closed by Theorem 10.41. (iii) The qualifier about the standard model is needed: a theory whose axioms exclude the standard algebra is not covered. (iv) The logarithmic bound counts layers of unit multiplication in a metatheoretic construction; it is not a bound on term size, evaluation time or checker size.

Remark 10.45. `adaptive_seed` proves Theorem 10.41 by the halving argument given here and proves Theorem 10.42. `local_scopes` proves the theorem by counting: from each starting coordinate at most one binary word of length m keeps a dependence alive, so at most d of the 2^m words are admitted, and a word admitted from no coordinate exists when $2^m > d$.

10.6 Finite profiles and relational bi-interpretations

This section draws on `local_scopes`. The coordinate bound leaves two questions open: whether the tied-seed language computes a bounded comparison of two unbounded inputs, and whether an interpretation that defines its operations by formulas escapes. A stronger invariant of tied-seed terms answers the first, and, combined with witness extraction from proofs, answers the second for a specified class of theories.

10.6.1 Finite control

Definition 10.46. A function $f\colon \mathbb{N}^r \to \mathbb{N}$, $r \geq 1$, has *finite control* if there are an active coordinate i, finite nonempty sets A_j and unary maps $\alpha_j\colon \mathbb{N} \to A_j$ for $j \neq i$, and unary functions F_a indexed by $a \in \prod_{j \neq i} A_j$, such that

$$f(x_1, \ldots, x_r) = F_{(\alpha_j(x_j))_{j \neq i}}(x_i).$$

The value may depend without bound on one coordinate; every other coordinate supplies only a member of a fixed finite set. The functions F_a are the *profiles*. For primitive recursive f all displayed maps are taken primitive recursive, with finite encodings of the A_j; specified finite codomains replace exact ranges, and no decision procedure for attained ranges is assumed.

A constant function has finite control with any active coordinate, and the definition imposes no restriction at arity one.

Lemma 10.47. *(i) If f has finite control and ρ is unary with finite range, then $\rho \circ f$ factors through finite-valued unary maps of all input coordinates.*

(ii) The functions with finite control contain the constants and projections and are closed under composition.

Both preserve primitive recursiveness of the displayed maps.

Proof. (i) Attach to the active coordinate the finite feature $\beta_i(x) = (\rho(F_a(x)))_a$. Together with the $\alpha_j(x_j)$ it determines $\rho(f(\vec{x}))$.

(ii) Consider $f(g_1(\vec{x}), \dots, g_r(\vec{x}))$ where f has the active input i and g_i has the active input k. Every other g_j enters f through the finite-valued $\alpha_j \circ g_j$, which by (i) is determined by finite unary features of all coordinates of $\vec{x}$. Combine the features of the coordinates other than k with the finite controls of g_i. With these combined controls fixed, the expression is one unary function of x_k; the features that depend on x_k are evaluated inside it. There are finitely many combined controls. □

Theorem 10.48 (Finite control of tied-seed terms)**.** *Every function of positive arity denoted by a term of the tied-seed language of Theorem 10.3(b) has finite control, with primitive recursive feature maps and profiles obtained by recursion on the term. A closed term is a constant.*

Proof. Variables and 0 are covered by Theorem 10.47. It suffices to show that $\mathsf{B}_h(n, \vec{p})$ has finite control when the update $h(u, v, \vec{p})$ has; the actual counter term is then substituted by Theorem 10.47(ii). Write $\gamma(\vec{p})$ for a tuple of finite unary features, one for each passive coordinate.

The active input of h is the previous state: $h(u, v, \vec{p}) = F_{\alpha(u),\gamma(\vec{p})}(v)$. For each of the finitely many values c of γ define $R_c(0) = 1 + F_{\alpha(0),c}(0)$ and $R_c(m+1) = F_{\alpha(m),c}(R_c(m))$. Then $\mathsf{B}_h(n, \vec{p}) = R_{\gamma(\vec{p})}(n)$, with active coordinate n.

The active input of h is the old counter: $h(u, v, \vec{p}) = F_{\beta(v),\gamma(\vec{p})}(u)$. Define $R_c(0) = 1 + F_{\beta(0),c}(0)$ and $R_c(m+1) = F_{\beta(R_c(m)),c}(m)$. Again $\mathsf{B}_h(n, \vec{p}) = R_{\gamma(\vec{p})}(n)$.

The active input of h is a passive parameter p. Write $\vec{q}$ for the other passive parameters and $h(u, v, p, \vec{q}) = F_{\alpha(u),\beta(v),\gamma(\vec{q})}(p)$. Form a finite catalogue $(V_j)_{j \in I}$ of unary functions that contains every profile $F_{a,b,c}$ and every initial profile $V_{0,c}(p) = 1 + F_{\alpha(0),\beta(0),c}(p)$, with separate indices for separate occurrences. Every state of the recursion is the value at p of a catalogue function. The only observation of a state used by the update is its β-value; record all of them in the finite feature

$$\kappa(p) = \big(\beta(V_j(p))\big)_{j \in I}.$$

For each control value c and each possible vector k, define a finite-state process: its initial state is the index of $V_{0,c}$, and at the counter m the state j is followed by the index of $F_{\alpha(m),k_j,c}$. Let $\theta_{c,k}(n) \in I$ be its state after n transitions and pack the finitely many processes into the finite-valued feature $\eta(n) = (\theta_{c,k}(n))_{c,k}$. Induction on the counter gives

$$\mathsf{B}_h(n, p, \vec{q}) = V_{\eta(n)_{\gamma(\vec{q}),\kappa(p)}}(p) :$$

the initial case selects $V_{0,c}$, and the transition uses exactly the recorded observation $\beta(V_j(p)) = \kappa(p)_j$. With $H_{e,c}(p) = V_{e_{c,\kappa(p)}}(p)$ this reads $\mathsf{B}_h(n, p, \vec{q}) = H_{\eta(n),\gamma(\vec{q})}(p)$, with the active coordinate p and finite controls on n and on each remaining parameter. The processes are defined for all vectors k, including unrealized ones; only the realized ones are used. □

The bound of Theorem 10.37 follows from the theorem by taking the maximum of the finitely many profiles and its running maximum. The new content is that the other inputs supply only finitely many distinct profiles.

Corollary 10.49 (No equality test, minimum or truncated subtraction)**.** *No term of the tied-seed language denotes the characteristic function of equality or of inequality,* $\min(a, b)$*, or* $a \dot{-} b = \max(a - b, 0)$.

Proof. A binary function with finite control and active first coordinate has finitely many distinct sections $a \mapsto f(a,b)$; with active second coordinate it has finitely many sections $b \mapsto f(a,b)$. Each displayed function has infinitely many distinct sections in both directions: for equality the section at b is the characteristic function of $\{b\}$; the functions $x \mapsto \min(x,b)$ differ at an x above both parameters; for truncated subtraction, distinct second arguments are separated at a large first argument, and distinct first arguments at the second argument 0. □

Remark 10.50. The theorem does not decide whether every unary primitive recursive function is denoted by a tied-seed term, since finite control is no restriction at arity one. It does not concern formula-defined relations: the source has logical equality $a = b$, which is not a term that returns its characteristic value. Bounded-valued primitives preserve the coordinate bound (Theorem 10.37) and need not preserve finite control; an adjoined equality characteristic violates it.

10.6.2 An existential-free presentation

Let T_{B} be intuitionistic first-order logic with equality in the staged signature of the tied-seed language, with the computation equations (10.1), (10.2), coverage $\forall x\,(x = 0 \vee \exists y\, x = Sy)$, separation $0 = S0 \to \bot$, and the universal closures of

$$\big(E(0,\vec{p}) \wedge \forall n\,(E(n,\vec{p}) \to E(Sn,\vec{p}))\big) \to \forall n\, E(n,\vec{p})$$

for literal equations E. By Theorem 10.4 the term P satisfies $P(0) = 0$ and $P(Sy) = y$ by computation, so coverage is equivalent, without induction, to the existential-free sentence $\forall x\,(x = 0 \vee x = S(Px))$. Equality is treated as a binary predicate with universal axioms of reflexivity, symmetry, transitivity and congruence for every function symbol; transport through formulas is derived. Thus T_{B} has an existential-free axiomatization in its own language. An infinite axiom scheme causes no difficulty, since a proof uses finitely many axioms.

Lemma 10.51 (Finite witness extraction)**.** *Let the language contain the closed term* 0 *and let* Γ *be a finite set of existential-free formulas. If intuitionistic first-order logic proves* $\Gamma \vdash \exists z_1 \cdots \exists z_d\, R(\vec{x},\vec{z})$*, then there are finitely many tuples of terms* $\vec{t}_1(\vec{x}), \ldots, \vec{t}_M(\vec{x})$ *with*

$$\Gamma \vdash \bigvee_{\ell \le M} R(\vec{x}, \vec{t}_\ell(\vec{x})).$$

The formula R *is arbitrary; it may contain existential quantifiers. No uniqueness of witnesses is assumed.*

Proof. Take a cut-free intuitionistic sequent proof; cut elimination for intuitionistic first-order logic is standard (for instance Avigad, *Algebraic proofs of cut elimination*, J. Log. Algebr. Program. 49 (2001), Theorem 7.6). Equality is handled by the existential-free axioms, so no equality extension of the theorem is needed.

Starting at the end sequent, trace the branches whose succedent carries the displayed leading existential quantifiers, and keep the prefix of witness terms already selected. At a left implication rule keep the proof of the first premise unchanged and follow the second premise; at a left disjunction rule follow both premises; the other left and structural rules preserve the tracked succedent. Every antecedent formula on a tracked branch is a subformula instance of a formula of Γ and is existential-free, so no left existential rule occurs there. While a leading existential quantifier remains, an initial sequent cannot close the branch, because the succedent is not an antecedent formula; the only applicable right rule is existential introduction. Record its witness term and continue. When all d witnesses are introduced, stop and keep the

remaining proof of the matrix instance unchanged; the matrix is not analyzed. If a tracked branch closes by falsity elimination, fill the unselected witnesses with 0; the matrix instance follows from the same falsity. There are finitely many branches, so finitely many complete tuples $\vec{t}_\ell$ are recorded.

Replace each stopped branch by its kept proof of $R(\vec{x}, \vec{t}_\ell)$ followed by introductions into the common disjunction, and reassemble the tracked left and structural inferences. An existential introduction that supplied a recorded component is omitted, since the disjunction is already the succedent of its transformed premise. Left implication inferences are reconstructed from the unchanged first premise and the transformed second premise.

No numerical parameter is bound on a tracked branch: left existential rules are absent, and right universal introduction cannot have the tracked succedent. Extraneous free variables of the proof are replaced by 0 after renaming eigenvariables, so the witness terms contain only $\vec{x}$ and the retained parameters. □

The lemma does not give a single uniform witness term; a finite disjunction is what the next theorem needs.

10.6.3 No compression of scalars

Theorem 10.52 (No provable compression from k scalars to d coordinates)**.** *Let T be an intuitionistic theory in the tied-seed language with an existential-free axiomatization and with the standard tied-seed algebra as a model. Let $k > d \geq 1$ and let $R(\vec{x}, \vec{z})$ be any formula with k variables $\vec{x}$ and d variables $\vec{z}$. If $T \vdash \forall \vec{x}\, \exists \vec{z}\, R(\vec{x}, \vec{z})$, then in the standard model some output tuple $\vec{z}$ occurs with two different input tuples. In particular T does not also prove*

$$R(\vec{x}, \vec{z}) \wedge R(\vec{x}\,', \vec{z}) \to \bigwedge_{i \leq k} x_i = x_i'.$$

The relation may have many outputs for each input.

Proof. Instantiate at free $\vec{x}$ and collect the finitely many axioms used. Theorem 10.51 gives d-tuples of terms $\vec{t}_1, \ldots, \vec{t}_M$ such that for every input in the standard model some $\vec{t}_\ell(\vec{x})$ is an R-output. By Theorem 10.48 each coordinate term $t_{\ell j}$ has finite control; let $q_{\ell j}$ be the number of its profiles. On the grid $X_N = \{0, \ldots, N-1\}^k$ it takes at most $q_{\ell j} N$ values, since its active argument takes at most N values; constant terms are included by adding an unused active argument. So $\vec{t}_\ell$ takes at most $N^d \prod_j q_{\ell j}$ values on X_N, and the union of the images has at most CN^d elements with $C = \sum_\ell \prod_j q_{\ell j}$. If every output belonged to at most one input, the N^k inputs of X_N would need N^k distinct outputs in this union, which fails for $N > C$. Soundness in the standard model turns a proof of the displayed implication into exactly that condition. □

Corollary 10.53 (Scalar pairing is not provably total)**.** *In T_{B} no formula is provably a total coding of pairs of numbers by single numbers with unique decoding. This includes a pairing relation $\pi_1(c) = a \wedge \pi_2(c) = b$ with unary terms as projections, and arbitrary formulas as projection relations, provided each code determines at most one pair.*

Proof. A provably total coding relation $R(a, b, c)$ with provably unique decoding is the case $k = 2$, $d = 1$ of Theorem 10.52. □

Even a proof that every unary primitive recursive function is a tied-seed term would therefore not give pairing inside T_{B}.

10.6.4 No finite-dimensional bi-interpretation

Theorem 10.54. *Let T be an intuitionistic theory in the tied-seed language with an existential-free axiomatization and with the standard tied-seed algebra as a model; for example T_{B}, or T_{B} with further existential-free axioms true in the standard algebra. Then T and* iPRA *are not bi-interpretable in the sense of Theorem 10.1. The interpreted domains, the operation graphs and the quotient equality are arbitrary formulas. The statement is parameter-free, and it holds when fixed parameters are retained uniformly in the interpretation proofs without assumptions that are not existential-free.*

Proof. Let $\mathcal{I}$ interpret iPRA in T with dimension d, domain $D \subseteq \mathbb{N}^d$ and interpreted equality E, and let $\mathcal{J}$ interpret T in iPRA with dimension e. A zero-dimensional domain has finitely many tuples and cannot take part in the source-side isomorphism with the infinite source domain, so $d, e \geq 1$.

The composite represents source numbers by e-tuples of interpreted arithmetic elements, that is, by de source coordinates. The source-side return isomorphism provides a formula $\eta(a, \vec{u})$ of T, with $\vec{u} \in D^e$ and the domain conditions included, such that

$$T \vdash \forall a\, \exists \vec{u}\, \eta(a, \vec{u}), \qquad T \vdash \eta(a, \vec{u}) \wedge \eta(a', \vec{u}\,') \wedge E^{\mathcal{I}}_{\mathcal{J}}(\vec{u}, \vec{u}\,') \to a = a',$$

where $E^{\mathcal{I}}_{\mathcal{J}}$ is the translation under $\mathcal{I}$ of the equality of $\mathcal{J}$. Only this part of the bi-interpretation is used.

Fix $k \geq 1$. iPRA has a primitive recursive injective tupling of ke numbers with primitive recursive projections, and proves its decoding equations. Its translation under $\mathcal{I}$ is a provably total coding relation from $(D/E)^{ke}$ to D/E that is injective on input tuples. Define $R_k(a_1, \ldots, a_k, \vec{c})$ in T: there are η-representatives $\vec{u}_i \in D^e$ of all a_i, and the interpreted tupling codes the ke arithmetic components of $(\vec{u}_1, \ldots, \vec{u}_k)$ by $\vec{c} \in D$. From the totality of η and of the tupling, finite existential reasoning gives

$$T \vdash \forall a_1 \cdots \forall a_k\, \exists c_1 \cdots \exists c_d\, R_k(\vec{a}, \vec{c}).$$

The inputs a_i range over the whole source domain; no choice is used.

Suppose $R_k(\vec{a}, \vec{c})$ and $R_k(\vec{a}\,', \vec{c})$. The same tuple $\vec{c}$ represents the same arithmetic element, by reflexivity of E on D. Injectivity of the interpreted tupling gives componentwise E-equality of the arithmetic components of $\vec{u}_i$ and $\vec{u}\,'_i$, hence $E^{\mathcal{I}}_{\mathcal{J}}(\vec{u}_i, \vec{u}\,'_i)$ for each i, since interpretations respect equality. The second property of η gives $a_i = a'_i$. So T proves that the output fibers of R_k are disjoint. Surjectivity of η and uniqueness of representatives are not needed.

With $k = d + 1$ this contradicts Theorem 10.52. □

Corollary 10.55 (Provably inhabited parameter families)**.** *The exclusion holds for a bi-interpretation with a finite tuple $\vec{p}$ of r source parameters subject to a condition $\Theta(\vec{p})$, when $T \vdash \exists \vec{p}\, \Theta(\vec{p})$ and T proves the interpretation and return-isomorphism facts uniformly under $\Theta(\vec{p})$.*

Proof. The construction with the parameters retained proves totality and disjoint fibers of $R_{k,\vec{p}}$ under $\Theta(\vec{p})$. Put $\widehat{R}_k(\vec{a}, \vec{p}, \vec{c}) := \Theta(\vec{p}) \wedge R_{k,\vec{p}}(\vec{a}, \vec{c})$. Then $T \vdash \forall \vec{a}\, \exists \vec{p}\, \exists \vec{c}\, \widehat{R}_k$, and an identical output $(\vec{p}, \vec{c})$ uses identical parameters, so the fibers are disjoint. There are $d + r$ output coordinates; take $k = d + r + 1$. The formula Θ is inside the output matrix, so no restriction on its quantifiers is needed. □

Remark 10.56 (Scope). Positive existential domain formulas do not rescue a bi-interpretation: they occur inside the matrix R_k, which witness extraction does not analyze, and the return isomorphism removes every domain condition from the unrestricted source inputs. The theorem does not exclude equiconsistency or one-way interpretations on complicated domains;

the return isomorphism is essential. It does not cover a parameter condition that T does not prove inhabited, a theory with positive existence axioms that cannot be eliminated in the tied-seed language, or every axiomatization in that language. It does not extend automatically to additional bounded numerical primitives, which preserve the coordinate bound and need not preserve finite control.

10.6.5 The selector boundary

The obstruction has an exact positive counterpart: a bounded primitive does not help (Theorem 10.37), and one unbounded selector restores primitive recursion.

Proposition 10.57 (`local_scopes`)**.** *Adjoin to the tied-seed language an operation C with $C(0,a,b)=a$ and $C(Sn,a,b)=b$, allowed inside update bodies. For terms $g(\vec{p})$ and $h(n,z,\vec{p})$ and fresh u,v put*

$$K(u,v,\vec{p}) := C(u,\, g(\vec{p}),\, h(W(u),v,\vec{p})), \qquad J(n,\vec{p}) := \mathsf{B}(Sn; u, v.K(u,v,\vec{p})),$$

with W as in Theorem 10.4. Equality reasoning and the computation equations prove $J(0,\vec{p}) = g(\vec{p})$ and $J(Sn,\vec{p}) = h(n,J(n,\vec{p}),\vec{p})$, with no induction, predecessor search, equality characteristic, pairing or independent initializer.

Proof. Suppress $\vec{p}$. The first update discards the tied initial state: $J(0) = \mathsf{B}(S0;u,v.K) = K(0,\mathsf{B}(0;u,v.K)) = C(0,g,\dots) = g$. At a successor, $J(Sn) = \mathsf{B}(SSn;u,v.K) = K(Sn,\mathsf{B}(Sn;u,v.K)) = C(Sn,g,h(W(Sn),J(n))) = h(n,J(n))$. These hold for arbitrary parameters. □

Zero, successor and projections are terms and composition is substitution, so every primitive recursive description compiles into the enlarged language, and conversely each description of the enlarged language is primitive recursive. This is numerical completeness only; the logical and interpretation claims for such a basis need their own proofs and their own size accounting.

A formula definition of C does not supply this compiler. With coverage and $0 \neq S0$, the term Z decides $z=0$, and $(z=0 \wedge w=a) \vee (z \neq 0 \wedge w=b)$ is provably total and single-valued, so a symbol with this graph is a definitional extension. Two things do not follow. The tied-seed signature has recursion symbols for its original update terms only; a recursion whose update contains the new symbol is a new graph whose totality and uniqueness need proof. And literal-equation induction for terms with the new symbol translates into induction for formulas that are not literal equations of the old language.

10.7 Zero profiles of positive iteration

This section draws on `checker_and_barriers`. Positive iteration $\mathsf{R}^+(n,a;v.h) = h^{n+1}(a)$ of Theorem 10.3(c) is not almost unary: $\mathsf{R}^+(n,a;v.Sv) = n+a+1$ grows in both arguments. A different invariant applies.

Theorem 10.58 (Zero-profile theorem)**.** *For every term $t(x_1,\dots,x_r)$ built from variables, 0, S and R^+, the zero set of t in the standard interpretation is $\varnothing$, $\mathbb{N}^r$, or $\{\vec{x} : x_j = 0\}$ for a single j. The profile and the coordinate are obtained by recursion on the term.*

Proof. Induction on terms, with bound update variables fresh. 0 is always zero, a successor never, and x_j exactly when $x_j = 0$. Consider $\mathsf{R}^+(n,a;v.h)$ with update $h(v,\vec{x})$. By the induction hypothesis the zero set of h has one of four forms. If h is never zero, the iterator is never zero. If h is always zero, the iterator is always zero. If h is zero exactly when a passive variable x_j is zero, the iterator has the same profile. These three cases use that at least one update is applied. If h is zero exactly when its state v is zero, every update preserves the zero

status of the state, so the final state has the zero profile of the seed a, which is classified by the induction hypothesis. The zero profile of the counter is not used: the number of updates is positive. A variable that occurs both in the counter and in the seed or the update keeps its role in the respective profile. □

Corollary 10.59. *No term of this language denotes the predecessor (zero set $\{0,1\}$), the zero test that is 1 at 0 and 0 elsewhere (zero set $\mathbb{N}\setminus\{0\}$), addition (zero set $\{(0,0)\}$), or the initialized iterator, whose constant-update case $I(n,a;\,v.\,b)$ is the selector and gives the zero test at $a=1$, $b=0$.*

The definable shifted addition $n+a+1$ is never zero and does not contradict the corollary.

Remark 10.60 (Scope). This is a standard-model term-definability obstruction. In intuitionistic logic with coverage, injectivity of S and separation $0\neq Sm$, the formula

$$(n=0\wedge z=a)\ \vee\ \exists m\,(n=Sm\wedge z=\mathsf{R}^{+}(m,a;v.h))$$

defines a total single-valued graph of the initialized iteration of each update h of the old language. Separation is needed to exclude overlap of the two branches. The observation does not give a bi-interpretation with iPRA: induction for the graph-defined expanded language and closure under nesting of newly defined updates remain to be proved, and in the native calculus the connectives of the graph formula are themselves reconstructed from a selector, so assuming them at that point is circular. The invariant does not survive a zero-test primitive: $\mathsf{R}^{+}(n,0;v.Zv)$ alternates between 1 and 0, so zero status then depends on the iteration count.

The motivation is a raw encoding in which a two-field numerical array $[a,h]$ has the value a and a three-field array $[n,a,h]$ the value $\mathsf{R}^{+}(n,a;v.h)$, which would allow a uniform predecessor expansion in place of a zero/successor conditional. The theorem shows that this encoding cannot inherit the term compiler or the selector-based logical bootstrap.

10.8 Rule-level obstructions

This section draws on `research`, `adaptive_seed`, `local_scopes`, `counter_congruence` and `constructed_computation`. Each result rejects one tempting deletion or replacement of a rule by a reading that validates the remaining rules. Each concerns the stated rule system with the identity translation of formulas.

10.8.1 Atomic case introduction cannot be made automatic

In N_{p} a certificate chooses between the two introductions of $D_{x,y}$. The *forced-progress* variant removes the choice: it uses DI-PROGRESS when the matrix is a literal equation and DI by cases when the matrix is compound. Equivalently, it deletes DI by cases at literal-equation matrices.

Proposition 10.61. N_{p} *derives $D_{x,y}(y=1)$, with x unused. The forced-progress variant does not.*

Proof. In N_{p}, DI by cases needs $0=1\vdash Sz=1$. From $0=1$, transport through the selector gives $s=\mathrm{sel}(0,s,t)=\mathrm{sel}(1,s,t)=t$ for all terms s,t, in particular $Sz=1$. No induction on a compound formula is used.

For the variant, evaluate terms in $\mathbb{N}$, read equations as equality and Q_yB as $B(0)\to\forall y\,B(y)$, and read $D_{x,y}$ by

$$(D_{x,y}B)^{\dagger}=\begin{cases}\exists x\,\forall y\,(B^{\dagger}(x,y)\to B^{\dagger}(x,Sy)), & B \text{ a literal equation},\\ \exists x\,(B^{\dagger}(x,0)\to\forall y\,B^{\dagger}(x,y)), & B \text{ compound}.\end{cases}$$

The reading is defined on each fixed formula. Forced progress introduction is sound by its direct meaning. Case introduction at a compound matrix is sound by the zero-or-successor property of $\mathbb{N}$. Paired elimination at an atomic matrix is sound because induction in $\mathbb{N}$ turns the supplied progressiveness into the conditional closure $Q_y B(a, y)$; at a compound matrix it is sound directly. QE, Hyp and the computation equations are sound. Transport is sound because numerical substitution does not change whether a matrix is a literal equation. Under † the formula $D_{x,y}(y = 1)$ says $\exists x\, \forall y\, (y = 1 \to Sy = 1)$, which is false at $y = 1$. □

The padding equivalence between a formula and a gated paired closure does not repair the counterexample: padding moves a proof obligation into a compound matrix, and the elimination returns only the conditional closure, not the progressiveness needed to introduce a raw atomic paired formula. A conditional closure can hold because its zero antecedent is false, without its matrix being progressive. A public language in which every atomic matrix is gated evades the obstruction at the cost of new grammar and decoder obligations.

Remark 10.62. `research` rederives the proposition from an earlier draft on flat computation certificates; `compact_kernel` and `uniform_predecessor` restate the conclusion. For the root-determined introduction with unrestricted transport the corresponding strictness is formalized as `rdprf_not_cases` and `rdprf_strictly_weaker`, with the progressive reading over the natural numbers as the sound reading.

10.8.2 Elimination majors must be proved, not only retained

Proposition 10.63 (`adaptive_seed`)**.** N_p *derives* $Q_y(0 = 0)$. *Restrict* N_p *so that the major premise of QE and of DE and the equation of a transport must be members of the retained context, keep* Comp*, the two introductions and, optionally, direct induction at equation goals, and add no cut rule. The restricted calculus does not derive* $Q_y(0 = 0)$.

Proof. In N_p, DI by cases with the witness 0 derives $D_{x,y}(0 = 0)$ from $0 = 0 \vdash 0 = 0$; DE applied to this derived major supplies $Q_y(0 = 0)$, and Hyp concludes.

In the restricted calculus consider a goal that is a closure $Q_y B$ in a context of equations only, initially empty. Hyp does not close it. An introduction concludes a paired formula. Comp adds an equation and keeps the goal; induction applies to equation goals only. Transport along a retained equation leaves a minor that is again a closure, in the same context. QE and DE need a retained closure or paired formula, and there is none. Induction on the height of a derivation shows that no derivation exists. □

The proposition concerns this deletion with closures as public formulas and the paired-introduction form; in N the formula is immediate by QI. It does not show that the closure binder must remain an object connective, and an unrestricted cut repairs the displayed theorem. The package verifies the example by a mutation of a checker in which the recursive check of a major is replaced by a membership test.

10.8.3 Root replacement does not give congruence

A further merger encodes an equation $s = t$ as a clause that sends $r = s$ to $r = t$, so that equality transport and closure use become one clause elimination. This yields replacement at the root of an equation only.

Proposition 10.64 (`local_scopes`)**.** *The computation equations, zero/successor cases, induction, reflexivity and root replacement do not derive congruence of* S *for a primitive equation predicate.*

Proof. Take $\mathbb{N}$ with the standard functions and interpret the equation predicate by the equivalence relation $\approx$ whose only non-singleton class is $\{2, 3\}$. Every computation equation holds because both sides have the same value. Reflexivity and root replacement are sound for an equivalence relation. Cases and induction are sound on $\mathbb{N}$ for every interpreted formula. But $2 \approx 3$ and $S2 \not\approx S3$. □

The counter-model is not the equality of the complete theory, which is actual equality. Theorem 10.95 shows what must be added: congruence in one fixed position, the counter of a recursor, suffices. A Leibniz clause $P(s) \to P(t)$ schematic in a formula context also merges the two eliminations, and moves the work into context application and higher-order substitution; no smaller checker for that route is established.

10.8.4 The witness order of the paired binder

A reversed binder $\mathsf{E}_{x,y}B := \forall y\, \exists x\, (B(x, 0) \to B(x, y))$ would merge the two eliminations and remove the reusable clause. Consider the naive introduction that fixes a witness term $t(z)$ in the instruction and checks $B(t(z), 0) \vdash B(t(z), Sz)$, its progress variant with a witness independent of z at a literal equation, and the ordinary instance elimination.

Proposition 10.65 (`counter_congruence`)**.** *There is a literal equation matrix B such that the ordinary reading $\forall y\, \exists x\, (B(x, 0) \to B(x, y))$ is true in $\mathbb{N}$ and $\mathsf{E}_{x,y}B$ is false under a reading that validates the naive rules. Hence the naive rules do not derive the equivalence of every E formula with its ordinary reading.*

Proof. Read $\mathsf{E}_{x,y}B$, with free parameters fixed, as: there is a unary primitive recursive f with $\forall y\, (B(f(y), 0) \to B(f(y), y))$. The naive case rule is sound with $f(0) = 0$ and $f(Sz) = t(z)$; the progress rule is sound with a constant f; instance elimination and transport are sound. The gated macros for the connectives keep their two-valued meanings: an existential macro needs a constant selector, and the universal macro has an unused existential coordinate.

Let F be a total computable function that is not primitive recursive and let $T(n, c)$ be the primitive recursive predicate saying that c is a halting computation trace of a fixed program for F on the input n. Choose a literal equation B with $B(x, 0) \equiv \top$ and $B(x, Sn) \equiv T(n, x)$, by a primitive recursive characteristic and a case selector. The ordinary reading is true. The selector reading is false: a primitive recursive f would give the trace $f(Sn)$, and the primitive recursive output decoder would compute $F(n)$. □

The source witness obtained at a fresh universal parameter may have to be chosen inside the proof scope of that parameter, after an existential elimination; a witness fixed before the subproof does not allow this interleaving. The proposition excludes this rule package with this reading. A repair with delayed witness choice or an intermediate introduction judgment is not excluded.

10.8.5 Two side conditions that cannot be dropped

Example 10.66 (Freshness of the step variable for the witness)**.** The formula $D_{x,y}\big(I(y, 0;\, v.\, Sx) = y\big)$ is false in $\mathbb{N}$: its zero instance is true, and its reading requires every positive number to equal one fixed Sx. If an introduction could take the step variable z as its witness, the step $I(Sz, 0;\, v.\, Sz) = Sz$ would be a computation instance. The condition that z is fresh for the witness is therefore necessary (`local_scopes`).

Example 10.67 (Local assumptions do not escape a major)**.** Case introduction derives $D_{x,y}(0 = 1)$ while locally assuming $0 = 1$. A checker that lets this assumption leak from the subproof of a major into the context of the caller accepts $0 = 1$ (`research`).

`constructed_computation` proves, by a finite-range invariant, that restricting the witnesses of DI to 0 or variables and at the same time the targets of QE to 1, to finitely many numerals, or to variables loses the theorem $D_{x,y}(\mathrm{sel}(y, N+1, x) = N+1)$ for a suitable N; the statement and its proof are given with the specialization rules in another chapter.

10.9 Independence of the progress rule

This section draws on `progress_independence`. It shows that QI by progress is not admissible over the other seven rules of the core. Write N^- for N without QI.

The README of the repository records three facts that shape a countermodel. Each of the other seven rules is independent of the rest, by readings of the language over the natural numbers (`NativeModels`). Such a reading cannot work for QI: over the natural numbers the soundness conditions for DI and DE already imply the condition for QI, whatever an equation means (`Sem.Sound.qIInd_of`); so the numerical domain must be nonstandard. And N^- derives $0 = 1$ from every successor cycle of odd length (`NoInd.no_odd_cycle`), so the model has no such cycle; Theorem 10.81 extends this to every length. The structures used earlier in the chapter do not serve. Every power $\mathcal{N}^{(J)}$ satisfies induction for literal equations (Theorem 10.11), so QI is sound in it; what fails there is coverage, hence DI by cases (Theorem 10.13). An ultrapower of the standard model with its internal iterator satisfies induction as well, by the theorem of Łoś. The countermodel has to do the opposite of a power: keep coverage and both computation equations at every count, nonstandard counts included, and lose induction for one equation. Coverage and acyclicity force the nonstandard part of the successor graph to consist of chains infinite in both directions, so every step function that is definable with parameters needs an orbit infinite in both directions along each chain.

All arguments of this section are carried out in classical metamathematics with the axiom of choice. This adds no classical axiom to the object calculus: a classical structure is a sound test of intuitionistic derivability (Section 10.1.1).

10.9.1 Coherent iterator structures

Definition 10.68. An *iterator language* is a first-order language L with 0, S and iterator symbols $I_h(n, a, \vec{p})$, one for each member $h(u, \vec{p})$ of a set of *descriptors*: terms of L with a designated state variable u and a finite tuple of passive variables $\vec{p}$. The variable u need not occur in h. The language is built in stages, so that the body of an iterator symbol is a term of an earlier stage. An L-structure is *coherent* if it satisfies the following first-order sentences.

(a) *Successor conditions.* S is injective, 0 is not a successor, every element is 0 or a successor, and $S^k x \neq x$ for each $k \geq 1$.

(b) *Computation.* For every iterator symbol, $I_h(0, a, \vec{p}) = a$ and $I_h(Sn, a, \vec{p}) = h(I_h(n, a, \vec{p}), \vec{p})$.

(c) *Cross-extensionality.* For every two iterator symbols I_h, I_g of L, the case $h = g$ included,

$$\forall \vec{p} \forall \vec{q} \Big[\forall u \, (h(u, \vec{p}) = g(u, \vec{q})) \to \forall n \, \forall a \, (I_h(n, a, \vec{p}) = I_g(n, a, \vec{q})) \Big]. \tag{Ext}$$

By (Ext) an iterator depends on the unary function $u \mapsto h(u, \vec{p})$ and on its two numerical inputs, not on the presentation of the function; repetition, omission and reordering of parameters are included. An independent choice of an orbit for each written body would not have this property: substitution of terms for the passive parameters changes the written body without changing the function, and the substitution lemma for the native syntax would fail (Theorem 10.74).

Lemma 10.69 (Orbits in one ultrapower)**.** *Let A be a nonempty L-structure, $\mathcal{U}$ a nonprincipal ultrafilter on $\mathbb{N}$ and $B = A^{\mathbb{N}}/\mathcal{U}$. For every term $h(u, \vec{p})$ of L and every tuple $\vec{p}$ of B, the function $f(v) = h^B(v, \vec{p})$ has a two-sided orbit: a family $(x_j)_{j\in\mathbb{Z}}$ of elements of B with $f(x_j) = x_{j+1}$ for all j.*

Proof. Let $\vec{p} = [\vec{p}_i]$, put $f_i(v) = h^A(v, \vec{p}_i)$ and choose any $a_i \in A$. For $j \in \mathbb{Z}$ let $x_j = [f_i^{\max(0,i+j)}(a_i)]$, with finite external iteration in each coordinate. Term functions of B are computed coordinatewise, and $\max(0, i + j) + 1 = \max(0, i + j + 1)$ for all $i \geq -j$, a cofinite set of coordinates. Hence $f(x_j) = x_{j+1}$. □

The lemma concerns functions defined by terms with parameters. No induction axiom of A is used, and the orbit need not pass through a prescribed element. In the structure $\mathfrak{M}_0$ of Theorem 10.72 the function $u \mapsto S(I_{\mathrm{id}}(u, u))$ maps everything into the standard ray and has no two-sided orbit; this is why the ultrapower precedes every adjunction of iterators.

Lemma 10.70 (Chains)**.** *In a structure with the successor conditions, the elements $S^k 0$, $k \in \mathbb{N}$, are pairwise distinct and form one component of the successor graph, the* standard ray*; every other component is a copy of $(\mathbb{Z}, +1)$.*

Proof. Every nonzero element has exactly one predecessor and 0 has none. An element outside the standard ray does not reach 0 after finitely many predecessor steps, so its iterated predecessors and successors are all defined; they are pairwise distinct by injectivity and the absence of cycles. The same two properties make the numerals distinct. □

Lemma 10.71 (Extension of a coherent structure)**.** *Let A be a coherent L-structure, $B = A^{\mathbb{N}}/\mathcal{U}$ as in Theorem 10.69, and let $\mathcal{D}$ be a set of terms of L with designated state variables that have no iterator symbol in L. Then B is a coherent elementary extension of A, and B has an expansion to $L \cup \{I_h : h \in \mathcal{D}\}$ that is coherent for the enlarged language. The expansion changes no element and no function of B.*

Proof. The conditions of Theorem 10.68 are first-order, so they hold in B by the theorem of Łoś, also for the parameter tuples that are new in B, and the diagonal embedding is elementary.

By Theorem 10.70 the successor graph of B is the standard ray together with copies of $\mathbb{Z}$; choose an origin b in each copy. Call a function $f\colon B \to B$ *presented* if $f = h^B(-, \vec{p})$ for a term h of L and a tuple $\vec{p}$ of B, and *old* if h can be taken with an iterator symbol in L. Assign to every presented function f one map $It_f\colon B \times B \to B$ as follows.

If f is old, say $f = g^B(-, \vec{q})$ with I_g in L, put $It_f(n, a) = I_g^B(n, a, \vec{q})$. This does not depend on the choice of g and $\vec{q}$, by (Ext) in B.

If f is not old, put $It_f(S^k 0, a) = f^k(a)$ on the standard ray. For each copy of $\mathbb{Z}$ with origin b and each seed a choose a two-sided orbit $(x_j)_{j\in\mathbb{Z}}$ of f by Theorem 10.69 and put $It_f(S^j b, a) = x_j$, where negative powers are iterated predecessors. The axiom of choice provides the choices.

In both cases $It_f(0, a) = a$ and $It_f(Sn, a) = f(It_f(n, a))$: for old f by the computation axioms of I_g, on the standard ray by the definition, and on a copy of $\mathbb{Z}$ because $x_{j+1} = f(x_j)$. No condition links the values on a copy of $\mathbb{Z}$ to the seed, and none is required.

For $h \in \mathcal{D}$ define $I_h(n, a, \vec{p}) = It_f(n, a)$ with $f = h^B(-, \vec{p})$. The computation axioms of the new symbols hold. (Ext) holds for every pair of symbols of the enlarged language: for two old symbols it holds in B already, and in the remaining cases two instances with the same function f are both equal to It_f, an old instance by the first clause of the definition. □

Lemma 10.72 (The seed)**.** *Let $L_0 = \{0, S, I_{\mathrm{id}}\}$, where I_{id} is the iterator symbol of the descriptor u without parameters. Let $\mathfrak{M}_0 = \mathbb{N} \sqcup \mathbb{Z}$, with 0 and $1 = S0$ in the first summand, the ordinary successor on each summand, and*

$$I_{\mathrm{id}}(n, a) = \begin{cases} a, & n \in \mathbb{N}, \\ 1, & n \in \mathbb{Z}. \end{cases}$$

Then $\mathfrak{M}_0$ is coherent, and $I_{\mathrm{id}}(c, 0) = 1 \neq 0$ for every c in the second summand.

Proof. The successor conditions are evident. $I_{\mathrm{id}}(0, a) = a$, and $I_{\mathrm{id}}(Sn, a) = I_{\mathrm{id}}(n, a)$ because S preserves the summands. (Ext) for the one symbol without parameters is trivial. No predicate that distinguishes the summands is in the language. □

Proposition 10.73 (The tower)**.** *Let L_{i+1} be L_i together with an iterator symbol for every term of L_i with a designated state variable that has no symbol in L_i, and let $L_\omega = \bigcup_i L_i$. There is a coherent L_ω-structure $\mathfrak{M}_\omega$ that extends $\mathfrak{M}_0$, in which every term of L_ω with a designated state variable has an iterator symbol, and in which $I_{\mathrm{id}}(c, 0) = 1 \neq 0$ for the elements c of the second summand of $\mathfrak{M}_0$.*

Proof. Given the coherent L_i-structure $\mathfrak{M}_i$, let $\mathfrak{M}_{i+1}$ be the expansion to L_{i+1} of its ultrapower given by Theorem 10.71; the ultrapower is taken in the whole language L_i, and only descriptors over L_i receive symbols at this stage, so no body contains a symbol that is being introduced. Replace each ultrapower by an isomorphic copy that contains the preceding structure. Then for every j the reducts $(\mathfrak{M}_i | L_j)_{i \geq j}$ form an elementary chain, and by the elementary chain theorem the reduct to L_j of the union $\mathfrak{M}_\omega$ is an elementary extension of each of them. Every successor condition, every computation axiom and every instance of (Ext) mentions finitely many symbols, hence is a sentence of some L_j that holds in $\mathfrak{M}_j$, and so holds in $\mathfrak{M}_\omega$. The equation $I_{\mathrm{id}}(c, 0) = 1$ and the inequation $1 \neq 0$ hold in $\mathfrak{M}_0$ and are preserved. A term of L_ω is a term of some L_i and has a symbol in L_{i+1}. □

10.9.2 Native terms in the tower

Translate native terms into L_ω by $x^* = x$, $0^* = 0$, $(St)^* = St^*$ and

$$I(n, a;\, u.\, h)^* = I_{h^*}(n^*, a^*, \vec{p}),$$

where $\vec{p}$ lists the free variables of h other than u in a fixed order. Inner iterators are translated first, so h^* is a term of some L_i and its symbol belongs to L_{i+1}; the number of stages is the nesting depth of iterators. Parameters are numerical variables; there is no sort of functions and no internal evaluator. The value $[\![t]\!]_\rho$ of a native term under an assignment ρ in $\mathfrak{M}_\omega$ is the value of t^*. The body u is the descriptor of I_{id}, so $[\![I(y, 0;\, u.\, u)]\!]_\rho = I_{\mathrm{id}}(\rho(y), 0)$.

Lemma 10.74 (Substitution)**.** *For native terms t, s, capture-avoiding substitution and every assignment ρ in $\mathfrak{M}_\omega$,*

$$[\![t[s/x]]\!]_\rho = [\![t]\!]_{\rho[x \mapsto [\![s]\!]_\rho]},$$

and the value of a term does not change under renaming of bound variables. The same holds for native formulas under the ordinary readings of Q_y and $D_{x,y}$ with actual equality.

Proof. Induction on t. The cases of variables, 0 and S are immediate. Let $t = I(n, a;\, u.\, h)$ with u renamed away from x and s. The translation of $t[s/x]$ is $I_g(\ldots, \vec{q})$ for the descriptor $g = (h[s/x])^*$, whose passive variables $\vec{q}$ are those of h other than x together with those of s; in general g is a different descriptor from h^*. By the induction hypothesis the counter and the seed have the required values, and for every element v

$$[\![h[s/x]]\!]_{\rho[u \mapsto v]} = [\![h]\!]_{\rho[x \mapsto [\![s]\!]_\rho][u \mapsto v]},$$

so the unary function of g at the parameters $\rho(\vec{q})$ equals the unary function of h^* at the parameters given by $\rho[x \mapsto [\![s]\!]_\rho]$. (Ext) for the pair I_g, I_{h^*} gives equal iterator values. Renaming of the bound variable is the same argument with equal functions. The statement for formulas follows by induction on formulas. □

This is the point at which independent choices for written bodies would be insufficient: the two descriptors are syntactically unrelated, and only (Ext), imposed for all parameter tuples and preserved through all stages, identifies their iterators.

10.9.3 The theorem

Theorem 10.75 (Progress is not admissible)**.** *Let* $E(y) := (I(y, 0; u.\, u) = 0)$.

(i) N^- *derives* $E(0)$ *and the sequent* $E(z) \vdash E(Sz)$.

(ii) Every sequent derivable in N^- *is valid in* $\mathfrak{M}_\omega$ *under the ordinary readings, with actual equality.*

(iii) $Q_y E(y)$ *is false in* $\mathfrak{M}_\omega$*, and so is* $D_{x,y} E(y)$ *with* x *unused.*

Hence N^- *does not derive* $Q_y E(y)$*, which* N *derives by one application of QI: the rule QI is not admissible over the other seven rules, and* DI-PROGRESS *is not admissible over the other rules of the paired-introduction form. The countermodel is a classical structure with actual equality in which the successor is injective, coverage and separation hold, there is no successor cycle, and both computation equations hold for every finite iterator term at every count.*

Proof. (i) $E(0)$ is the zero computation. The successor computation gives $I(Sz, 0; u.\, u) = I(z, 0; u.\, u)$, and transitivity with the hypothesis gives $E(Sz)$. Reflexivity, symmetry and transitivity are derived from the zero computation and transport, without QI (Section 1.4).

(ii) A sequent is valid if every assignment that satisfies its context satisfies its conclusion. HYP is immediate. The zero computation is the computation axiom for I_{h^*}. For the successor computation, the left side has the value $h^*(I_{h^*}(n, a, \vec{p}), \vec{p})$ by the computation axiom, and the right side $h[I(n, a; u.\, h)/u]$ has the same value by Theorem 10.74. Transport, from $s = t$ and $h(s) = r$ to $r = h(t)$, is sound because equality is actual and $[\![h(s)]\!]_\rho = [\![h(t)]\!]_\rho$ by Theorem 10.74. QE is modus ponens and instantiation, with Theorem 10.74 for the instance $B(t)$. DE is existential elimination: the value of the fresh variable is the witness, and freshness keeps the context and the conclusion unchanged. For DI by cases let ρ satisfy Γ and take the value of t as witness. Suppose $B(t, 0)$ holds under ρ. Every element is 0 or Sd for some d; in the second case $\rho[z \mapsto d]$ still satisfies Γ and $B(t, 0)$ and gives t the same value, since z is fresh, so the premise gives $B(t, Sz)$ there. Hence $B(t, y)$ holds for every y. All seven arguments are first-order soundness arguments in one structure; coverage is used for DI only.

(iii) By the translation, $E(y)$ says $I_{\mathrm{id}}(y, 0) = 0$. It holds at $y = 0$ and fails at $y = c$ by Theorem 10.73. So $E(0) \to \forall y\, E(y)$ is false, and with an unused witness coordinate so is $\exists x\, (E(0) \to \forall y\, E(y))$. □

With the seven independence results recorded in the README (`NativeModels`), each of the eight rule families of N is independent of the other seven.

Remark 10.76 (What the construction needs)*.* Four requirements carry the proof, and dropping any of them leaves a gap. The iterator of a function that is not old is chosen once for the function, not for a written body. (Ext) is imposed for all pairs of symbols and all parameter tuples, so that it transfers to the parameters of later ultrapowers. Each ultrapower is taken in the whole current language, so that the earlier iterators and their coherence are carried along. And a stage adds symbols only for descriptors over the preceding language, so that the orbit

lemma applies to them. The construction makes no assertion about iteration of arbitrary functions; Theorem 10.69 is false for arbitrary maps.

Remark 10.77 (A countable model). The languages L_i and L_ω are countable. By the downward Löwenheim–Skolem theorem $\mathfrak{M}_\omega$ has a countable elementary substructure that contains c; it satisfies the same sentences, hence is coherent and refutes $E(c)$, and the soundness argument applies to it. This does not contradict the obstruction to readings over the natural numbers: the successor of the countable carrier has nonstandard chains and is not the map $n \mapsto n+1$ under any enumeration that would make it a reading in that sense. No computable presentation of the model is claimed.

Proposition 10.78 (A seed with cancellative addition). *Let $M_0 = \{(0,n) : n \in \mathbb{N}\} \cup \{(k,z) : k \geq 1,\ z \in \mathbb{Z}\}$ with componentwise addition, $0 = (0,0)$, $1 = (0,1)$ and $S(k,z) = (k, z+1)$. Let $I_{\mathrm{id}}(n,a)$ be a if the first coordinate of n is 0 and 1 otherwise, and let $K(n,a) = a+n$ be the iterator symbol of the descriptor Su. Then M_0 is coherent for the language $\{0, S, I_{\mathrm{id}}, K\}$, $K(n,0) = n$ holds, addition $a+n = K(n,a)$ is associative, commutative and cancellative with the unit 0, and $I_{\mathrm{id}}((1,0),0) = 1$. The tower over M_0 gives a countermodel as in Theorem 10.75 that satisfies these laws in addition.*

Proof. M_0 is a submonoid of the group $\mathbb{Z}^2$, so addition is associative, commutative and cancellative; it is conical, and $Sx = x+1$. The successor is injective, omits 0, has no cycle, and every nonzero element has a predecessor in M_0. The computation axioms of I_{id} hold because S preserves the first coordinate, and those of K are $a+0 = a$ and $a+Sn = S(a+n)$. (Ext) between the two symbols has a false antecedent, since the identity and the successor are different functions. The displayed laws are first-order sentences of the seed language and survive the tower. □

The stronger seed shows that the countermodel does not depend on a defect of addition. It does not separate uniqueness of iteration from induction: in M_0 the term $f(n) = n + I_{\mathrm{id}}(n,0)$ satisfies $f(0) = 0$ and $f(Sn) = Sf(n)$, and $f(c) = Sc \neq c = K(c,0)$ at $c = (1,0)$, so the instance of the rule UR of Section 10.13 for the update Su fails there. The instance of UR for the identity update and the constant template $f(z) = 0$ already excludes the witness of the theorem, since its conclusion is $0 = I(t, 0;\, u.\, u)$.

10.9.4 No successor cycle without progress

The countermodel has no successor cycle, and it cannot have one. Let T^- be intuitionistic logic with equality, the two computation equations and coverage; there is no induction. Write $Z(x) := \mathrm{sel}(x, 1, 0)$, so $Z(0) = 1$ and $Z(Sx) = 0$, write $\overline{j}$ for $S^j 0$, and $h^{[r]}$ for the r-fold composition of a unary term, which introduces no iterator. All inductions below are on the integer that specifies a term.

Lemma 10.79 (Finite clocks). *For every $m \geq 2$ there is a term $h_m(x)$, built from the leaves $\overline{0}, \ldots, \overline{m-1}$ by the selector with arbitrary counter terms, such that T^- proves*

$$\bigvee_{j<m} h_m(x) = \overline{j}, \qquad h_m(\overline{j}) = \overline{j+1}\ (j < m-1), \qquad h_m(\overline{m-1}) = 0.$$

Proof. Let $h_2 = Z$; coverage gives $Z(x) = 0 \vee Z(x) = 1$. Given h_m, put $T_m(x) := I(x, 0;\, u.\, h_m(u))$. By coverage on x and the two computations, $T_m(x)$ is 0 or a value of h_m, so it is one of $\overline{0}, \ldots, \overline{m-1}$; nothing is asserted about the preceding iterator value. Finite computation gives $T_m(\overline{j}) = \overline{j}$ for $j < m$ and $T_m(\overline{m}) = 0$. For $1 \leq r < m$ put

$$d_{m,r}(z) := Z\big(h_m^{[m-r]}(z)\big), \qquad V_{m,0}(z) := 0, \qquad V_{m,r}(z) := \mathrm{sel}\big(d_{m,r}(z),\, V_{m,r-1}(z),\, \overline{r+1}\big),$$

and $V_m := V_{m,m-1}$. Ground computation gives $d_{m,r}(\overline{s}) = 1$ for $s = r$ and $d_{m,r}(\overline{s}) = 0$ for $s < m$, $s \neq r$; hence $V_m(0) = 0$ and $V_m(\overline{r}) = \overline{r+1}$ for $1 \leq r < m$. At an arbitrary argument V_m takes one of the values $0, \overline{2}, \ldots, \overline{m}$, by coverage on the selector counters. Put

$$h_{m+1}(x) := \mathrm{sel}(x,\, 1,\, V_m(T_m(x))).$$

Its values are among $\overline{0}, \ldots, \overline{m}$. At 0 its value is 1; at $\overline{j}$ with $1 \leq j < m$ it is $V_m(\overline{j}) = \overline{j+1}$; at $\overline{m}$ it is $V_m(0) = 0$. □

For example $h_3(x) = \mathrm{sel}(x, 1, \mathrm{sel}(Z(Z(T_2(x))), 0, 2))$ with $T_2(x) = I(x, 0;\, u.\, Z(u))$.

Lemma 10.80 (Separation of numerals)**.** *For $m \geq 2$ and distinct $i, j < m$, T^- proves $\overline{i} = \overline{j} \to 0 = 1$, without injectivity of the successor.*

Proof. Apply $h_m^{[r]}$ with $r \equiv -i \pmod m$, $0 \leq r < m$. The result is $0 = \overline{s}$ with $1 \leq s < m$, and Z gives $1 = 0$. □

Theorem 10.81 (No finite successor cycle)**.** *For every $k \geq 1$ and every term c, T^- proves $S^k c = c \to 0 = 1$, and N^- derives the sequent $S^k c = c \vdash 0 = 1$.*

Proof. Let $m = k + 1$ and $v = T_m(c)$. The hypothesis and k successor computations give $v = T_m(S^k c) = h_m^{[k]}(v)$. By Theorem 10.79, $v = \overline{j}$ for some $j < m$, and the equation becomes $\overline{j} = \overline{(j+k) \bmod m}$. Since $k = m - 1$ the two residues differ, and Theorem 10.80 gives $0 = 1$.

For N^- the finite disjunctions are removed. The cases rule for equation templates, from $E(0)$ and $E(Sz)$ to $E(t)$, is derivable in N^-: DI by cases with an unused witness, DE and QE (`NoInd.cases`). First, if $E(\ell)$ is derivable for each leaf ℓ of a term built from finitely many leaves by the selector, then E of the term is derivable, by induction on the term with the cases rule at each selector counter. Second, since h_m is such a term over $\overline{0}, \ldots, \overline{m-1}$, the cases rule at the counter of T_m gives: if $E(\overline{j})$ is derivable for every $j < m$, then $E(T_m(c))$ is derivable; the zero branch is $E(0)$ and the successor branch is $E(h_m(T_m(z)))$. Third, put $b_r(a) := Z(h_m^{[(-r) \bmod m]}(a))$ for $r < m$, so that $b_r(\overline{j}) = 1$ exactly when $r = j$, and

$$\mathrm{xor}(p, q) := \mathrm{sel}(p, q, Z(q)), \qquad \mathrm{or}(p, q) := \mathrm{sel}(p, q, 1),$$

and let $D_m(a, b)$ be the left-nested or, starting at 0, of the terms $\mathrm{xor}(b_r(a), b_r(b))$, $r < m$. The cases rule for the template $\mathrm{xor}(Z(w), Z(w)) = 0$, instantiated at the arguments of Z in the terms $b_r(a)$, and the computation $\mathrm{or}(0, 0) = 0$ give $D_m(a, a) = 0$ for every a. Ground computation gives $D_m(\overline{j}, h_m^{[k]}(\overline{j})) = 1$ for $j < m$, so the second step gives $D_m(v, h_m^{[k]}(v)) = 1$ at $v = T_m(c)$. Under the hypothesis $v = h_m^{[k]}(v)$, and transport gives $D_m(v, v) = 1$, hence $0 = 1$. No disjunction, no characteristic of equality of arbitrary numbers and no QI is used. □

The README records the odd case in Lean (`NoInd.no_odd_cycle`, `NoInd.no_succ_fixed`) and leaves the even case open, because it needs a counter of another period whose behavior has to be established without induction; the clocks h_m are such counters. The theorem is a constraint on countermodels and does not make QI redundant.

Remark 10.82 (What is proved and what is not). Theorem 10.75 is an independence result for one rule of one fixed kernel, with the identity translation of formulas. It is proved by a written model construction that uses a nonprincipal ultrafilter and the axiom of choice. It is not formalized: the README lists the package as filed and not formalized, and the Lean readings of `NativeModels` fix the standard successor, so they cannot express the model. The result does not bound the size of a checker, does not concern presentations in which QI is fused with another rule or replaced by a weaker induction interface, and does not show that the eight rules are a minimum: equivalent presentations with fewer rule families exist. It does not decide whether the uniqueness rule for the successor update is equivalent to induction for literal equations, since the seed of Theorem 10.78 refutes that uniqueness instance.

10.10 The progressive binder presentation

This section draws on `research`. It shows that the calculus of Theorem 10.7, with BoolCases, is a complete presentation of iPRA, so that Theorem 10.17 isolates the one rule that cannot be removed. Structural weakening, contraction, exchange, substitution, renaming and assumption cut are admissible by induction on derivations, with eigenparameters renamed first.

Lemma 10.83 (Identity and equation induction, without cases)**.** *The case-free calculus derives $t = t$, symmetry, transitivity and numerical congruence; $\Gamma, A \vdash A$ for every object formula A; and contextual induction for literal equations: from $\Gamma \vdash E(0)$ and $\Gamma, E(z) \vdash E(Sz)$ with z fresh, $\Gamma \vdash E(t)$.*

Proof. The computation $\mathsf{R}(0, t; u, v.0) = t$, used as the major and as the minor of Tr with the template $w = t$, gives $t = t$; the templates $w = s$, $s = w$ and $f(s) = f(w)$ give the other equality rules. An available equation $s = t$ transports $s = s$ to $s = t$. For an available $\mathsf{H}_{x,y}B$, induct on the binder depth: HE supplies a fresh a and $\mathsf{P}_y B(a, y)$; reintroduce the formula by HI at the witness a, where the local assumption $B(a, z)$ yields a proof of itself by the induction hypothesis and Step gives $B(a, Sz)$. For induction, HI with an unused witness gives $\mathsf{H}_{x,y}E(y)$, HE supplies $\mathsf{P}_y E(y)$, and Iter gives $E(t)$. □

Lemma 10.84 (Boolean cases and constructor cases)**.** *Put $P(n) = \mathsf{R}(n, 0; u, v.u)$ and $C(n, a, b) = \mathsf{R}(n, a; u, v.b)$. The case-free calculus derives*

$$C(\mathrm{sg}(x), 0, x) = x, \qquad C(\mathrm{sg}(x), x, S(Px)) = x.$$

Over the case-free calculus, BoolCases is equivalent to the rule

$$\frac{\Gamma,\ t = 0 \vdash A \quad \Gamma,\ t = Sz \vdash A}{\Gamma \vdash A}\ \textsc{Cases}, \qquad z \text{ fresh for } \Gamma, t, A.$$

Proof. Both identities follow by literal-equation induction; each base is a zero computation and each successor step is a successor computation that does not use its induction hypothesis. Hence $\mathrm{sg}(t) = 0$ gives $t = 0$ and $\mathrm{sg}(t) = 1$ gives $t = S(Pt)$. For Cases from BoolCases, substitute $P(t)$ for z in the second premise. Conversely Cases gives BoolCases by the equations of sg. □

The missing principle is thus the two-value range of sg; equation induction supplies the numerical identities.

Lemma 10.85 (Binder congruence)**.** *If $A(x, y)$ and $B(x, y)$ are interderivable uniformly in x, y, with x, y not in the retained context, then $\mathsf{H}_{x,y}A$ and $\mathsf{H}_{x,y}B$ are interderivable. No cases and no compound induction are used.*

Proof. From $\mathsf{H}_{x,y}A$, HE gives $\mathsf{P}_y A(a, y)$. Introduce $\mathsf{H}_{x,y}B$ at the witness a: convert the local $B(a, z)$ to $A(a, z)$, apply Step, convert $A(a, Sz)$ to $B(a, Sz)$. □

Definition 10.86 (Gates, selectors and macros)**.** With bound variables renamed apart, define by recursion on formulas

$$\begin{aligned} G_v(s = t) &:= (C(v, 0, s) = C(v, 0, t)), & G_v(\mathsf{H}_{x,y}A) &:= \mathsf{H}_{x,y}G_v A, \\ M_v(s = t,\ r = w) &:= (C(v, s, r) = C(v, t, w)), & M_v(\mathsf{H}_{x,y}A, \mathsf{H}_{x,y}B) &:= \mathsf{H}_{x,y}M_v(A, B), \\ M_v(E, \mathsf{H}_{x,y}B) &:= \mathsf{H}_{x,y}M_v(G_y E, B), & M_v(\mathsf{H}_{x,y}A, E) &:= \mathsf{H}_{x,y}M_v(A, G_y E), \end{aligned}$$

for equations E; the maximal binder depth decreases in each recursive call. Let $\mathsf{T} = (0 = 0)$ and let p, q, d be the terms with

$$p(0) = 0,\ p(Sn) = C(p(n), 1, 0); \quad q(0) = 0,\ q(Sn) = C(p(n), q(n), S(q(n))); \quad d(0) = 0,\ d(Sn) = SS(d(n)),$$

which are parity, half and doubling in $\mathbb{N}$. The logical macros are

$$\begin{aligned} A \Rightarrow_{\mathsf{H}} B &:= \mathsf{H}_{u,z} M_z(A, B), & \exists_{\mathsf{H}} x\, A(x) &:= \mathsf{H}_{x,z} G_z A(x), \\ A \wedge_{\mathsf{H}} B &:= \mathsf{H}_{u,z} G_z M_{p(z)}(A, B), & A \vee_{\mathsf{H}} B &:= \exists_{\mathsf{H}} x\, M_x(A, B), \\ \forall_{\mathsf{H}} x\, A(x) &:= \mathsf{H}_{u,z} M_{p(z)}(\mathsf{T}, A(q(z))), & \bot_{\mathsf{H}} &:= (0 = S0), \end{aligned}$$

with all unused coordinates fresh. These are translations of fixed finite syntax, not truth predicates.

Lemma 10.87. *The calculus of Theorem 10.7 derives:*

(i) $G_0 A$, *and* $G_{Sv} A$ *is interderivable with* A;

(ii) A *is interderivable with* $\mathsf{H}_{u,y} G_y A$ *for fresh* u, y;

(iii) $M_0(A, B)$ *is interderivable with* A *and* $M_{Sv}(A, B)$ *with* B;

(iv) $p(d(n)) = 0$, $p(Sd(n)) = 1$, $q(d(n)) = n$ *and* $q(Sd(n)) = n$.

Items (i)–(iii) hold in the case-free calculus.

Proof. (i) By induction on the formula, with the computations of C at equations and Theorem 10.85 at binders; for G_0 at a binder, HI ignores its local assumption and uses the induction hypothesis for the successor matrix. (ii) Forward, HI ignores its assumption and uses $G_{Sz} A$ from A. Backward, HE supplies the progress clause, $G_0 A$ is a theorem, and one Step gives $G_1 A$, hence A. (iii) By computation, (ii) and congruence. (iv) Induct on the single equation $p(d(n)) = 0$: at a successor two parity updates send 0 to 1 and back. Then $p(Sd(n)) = 1$ by one computation. Induct on the single equation $q(d(n)) = n$: the two half updates at $d(n)$ and $Sd(n)$ first keep the value and then increment it, by the parity equations. Then $q(Sd(n)) = n$ by its defining update. Two single-equation inductions are used in sequence; no induction on a conjunction or a compound formula. □

Proposition 10.88 (All intuitionistic rules)**.** *The calculus of Theorem 10.7 derives the introduction and elimination rules of intuitionistic natural deduction for the macros of Theorem 10.86, with the usual eigenvariable conditions, explosion from* $\bot_{\mathsf{H}}$, *and coverage* $(n = 0) \vee_{\mathsf{H}} \exists_{\mathsf{H}} v\, (n = Sv)$. *No excluded middle for quantified formulas is used.*

Proof. By Theorem 10.84, Cases is available, and with Theorem 10.87(iii) it selects either endpoint of a selector at an arbitrary control, also when the context mentions the control; in particular proofs of A and of B give a proof of $M_t(A, B)$.

Implication. Given $\Gamma, A \vdash B$, apply HI; the local assumption is $M_z(A, B)$, and Cases on z gives A and then B at $z = 0$, and B at $z = Sw$; in both branches the successor matrix is B. For elimination, HE supplies the clause, A is its zero matrix, and Step at 0 gives B.

Existential quantifier. From $A(t)$, HI at the witness t ignores the local gate. For elimination, HE gives a fresh a and the clause; the zero gate is a theorem and Step at 0 gives $A(a)$.

Conjunction. If A and B are proved, Cases and the selector endpoints prove every successor matrix. Conversely the zero gate is a theorem, Step at 0 gives the matrix at 1, which is B since $p(1) = 1$, and a second Step gives the matrix at 2, which is A since $p(2) = 0$.

Disjunction. The introductions use the witnesses 0 and 1. For elimination, eliminate the existential at a fresh a and apply Cases to a.

Universal quantifier. Let $\Gamma \vdash A(x)$ with x fresh. Then T and $A(q(Sz))$ are proved, so Cases and the selector endpoints prove the successor matrix, and HI ignores its assumption. For specialization, HE gives the clause; the matrix at $d(t)$ is a theorem by Theorem 10.87(iv), and Step at $d(t)$ gives the matrix at $Sd(t)$, which is $A(t)$. No compound progress clause is iterated.

Falsity and coverage. Under $0 = 1$, computation and transport give $s = C(0, s, t) = C(1, s, t) = t$; induction on formulas then gives every object formula, since at a binder HI ignores its assumption. Cases proves coverage by its two introductions. □

Theorem 10.89 (`research`)**.** *Let T_{H} be intuitionistic first-order logic with equality in the staged signature of* R*, with the computation equations, induction for literal equations, coverage and $0 = 1 \to \bot$. The calculus of Theorem 10.7 is sound in T_{H} under its ordinary readings and derives the translations of all rules and axioms of T_{H}. For every matrix B,*

$$\mathsf{H}_{x,y} B(x, y) \;\dashv\vdash\; \exists_{\mathsf{H}} x \, \forall_{\mathsf{H}} y \, (B(x, y) \Rightarrow_{\mathsf{H}} B(x, Sy)).$$

T_{H} and iPRA *have the same theorems in the staged primitive recursive language. The progressive calculus is equiconsistent and bi-interpretable with* iPRA*, with whole domains, actual equality and return isomorphisms with graph $x = y$.*

Proof. Soundness. HI, HE and Step are logical rules under the readings; Iter is equation induction under the retained base; BoolCases is the range of sg, which follows from coverage.

Completeness. Theorem 10.88 gives the logical rules and coverage, Theorem 10.83 gives equation induction, and separation translates to the identity implication $(0 = 1) \Rightarrow_{\mathsf{H}} (0 = 1)$.

The composite. Forward, HE supplies a fresh a and the clause; at a fresh y assume $B(a, y)$ and use Step; introduce the implication, the universal and the existential macro and close HE. Backward, eliminate the existential at a fresh a, specialize the universal at the eigenparameter z of HI, and eliminate the implication under the assumption $B(a, z)$. The clause composite is the universal progress formula. In T_{H} the gate, selector and clock equations prove the intended meaning of every macro; for the universal macro, the even-to-odd progress implications give every instance $A(n)$, and all instances of A make every progress implication true, with no induction on A. Induction on formulas, with Theorem 10.85, gives both formula composites.

Arithmetic. T_{H} has the computation equations of ordinary counter-aware primitive recursion, coverage, separation and equation induction. The scalar iterator is the case of R whose update ignores its counter, so the construction of a characteristic of actual equality and the derivation of quantifier-free induction from equation induction (Theorem 1.52) apply to the terms of the larger language, and T_{H} proves every quantifier-free induction instance. Counter-aware recursion compiles every primitive recursive description directly, in place of the scalar compiler. Numerical descriptions are identified with themselves, and equation induction identifies two fixed recursions with the same equations, so both domain isomorphisms have the graph $x = y$. □

Remark 10.90. The presentation has one unrestricted introduction and pays for it with the explicit BoolCases and with the distinction between Step and Iter. The universal quantifier is recovered with one progress step through the clock p, q, d. No smaller complete checker for this presentation is established, and the presentation is not assigned the measurements of the checkers of its package.

10.11 The counter-congruence form of the calculus

This section draws on `counter_congruence`. General equality transport carries an arbitrary formula or term context as rule data. Here it is recovered from two specific operations: replacement of the complete left side of an equation, and congruence in the counter of a recursor. The result is a restriction of the primitive interface; it does not by itself give a shorter checker. Theorem 10.64 shows that root replacement alone is not enough, and Theorem 10.99 that counter congruence alone is not enough.

Definition 10.91 (The calculus $\mathcal{K}_{\mathrm{ctr}}$)**.** Terms are variables, 0 and the successor-seeded counter-aware recursor $\mathsf{R}_S(n, a; u, v.h)$, with $Sa := \mathsf{R}_S(0, a; u, v.0)$ and $1 := S0$. The computation schemas are

$$\mathsf{R}_S(0, a; u, v.h) = Sa, \tag{R0}$$

$$\mathsf{R}_S(Sn, a; u, v.h) = h[n/u][\mathsf{R}_S(n, a; u, v.h)/v]. \tag{RS}$$

Object formulas are $A ::= (s = t) \mid D_{x,y}A$. Contexts are finite lists of object formulas and internal clauses Q_yB; a clause is available by membership only, and an object formula by membership or by a proof, written $\Gamma \triangleright M$. The rules are Hyp for object formulas, R0, RS,

$$\frac{\Gamma \triangleright s = t \quad \Gamma \vdash s = r}{\Gamma \vdash t = r}\ \mathrm{RL} \qquad \frac{\Gamma \triangleright s = t}{\Gamma \vdash \mathsf{R}_S(s, a; u, v.h) = \mathsf{R}_S(t, a; u, v.h)}\ \mathrm{CC}$$

and QE, DE, DI by cases and DI-progress as in N_{p}, with majors available in the sense of $\triangleright$. In CC the initializer, the update and the passive parameters are the same on both sides. There is no primitive reflexivity, no general transport and no other congruence. The designated contradiction is $0 = 1$.

Weakening, exchange, contraction, renaming, substitution of a term for a parameter and cut of an object assumption are admissible by induction on derivations; substitution preserves the matching initializer and update of CC and the matching root of RL. These are operations on proof syntax and do not replace equal terms inside a formula.

Lemma 10.92 (Equality algebra from root replacement)**.** *Reflexivity, symmetry and transitivity are derivable from RS and RL, without CC.*

Proof. For a term s put $c_s := \mathsf{R}_S(Ss, 0; u, v.u)$; RS gives $c_s = s$. With one copy as the major and one as the minor, RL gives $s = s$. An available $s = t$ with the minor $s = s$ gives $t = s$. For $s = t$ and $t = r$, symmetry gives $t = s$, and RL with this major and the minor $t = r$ gives $s = r$. Paths of equations are composed at their roots; no proper subterm changes. □

Lemma 10.93 (Numerical cases)**.** *For every object template $A(y)$, from $\Gamma \vdash A(0)$ and $\Gamma \vdash A(Sz)$ with z fresh, $\Gamma \vdash A(t)$ is derivable without either equality rule and without DI-progress.*

Proof. Weaken the second premise by $A(0)$ and apply DI by cases with the witness 0 and an unused coordinate x, which gives $D_{x,y}A(y)$. DE supplies the clause $Q_yA(y)$, and QE with the first premise gives $A(t)$. □

Lemma 10.94 (Counter representation of a context)**.** *Let $W(n) := \mathsf{R}_S(n, 0; u, v.u)$, so that $W(0) = S0$ and $W(Sn) = n$ by R0 and RS. For a numerical template $F(x)$ put*

$$H_F(n) := \mathsf{R}_S(n, F(0); u, v.S(F(Su))), \qquad J_F(n) := W(H_F(n)),$$

with fresh bound variables. Then $\mathcal{K}_{\mathrm{ctr}}$ derives $J_F(n) = F(n)$ for arbitrary n in every context, by cases and without DI-progress.

Proof. The update of H_F does not contain v, so R0 and RS give literally $H_F(0) = S(F(0))$ and $H_F(Sz) = S(F(Sz))$. Let q be 0 or Sz. CC for the recursor W gives $W(H_F(q)) = W(S(F(q)))$, and RS gives $W(S(F(q))) = F(q)$. Composition by Theorem 10.92 gives $J_F(q) = F(q)$. Theorem 10.93 for the equation $J_F(y) = F(y)$ gives the claim. The only change of a proper subterm is the one CC instance through W; replacing $H_F(q)$ by $S(F(q))$ under W without it would assume the congruence that is being derived. □

Theorem 10.95 (All numerical congruence from counter congruence)**.** *For every numerical template $F(x)$, from $\Gamma \triangleright s = t$ the calculus $\mathcal{K}_{\mathrm{ctr}}$ derives $\Gamma \vdash F(s) = F(t)$. Hence for templates U, V, from $\Gamma \triangleright s = t$ and $\Gamma \vdash U(s) = V(s)$ it derives $\Gamma \vdash U(t) = V(t)$.*

Proof. CC with the initializer $F(0)$ and the update $S(F(Su))$ gives $H_F(s) = H_F(t)$; a second CC for W gives $J_F(s) = J_F(t)$. Theorem 10.94 and Theorem 10.92 give $F(s) = J_F(s) = J_F(t) = F(t)$. For the second claim, $U(t) = U(s) = V(s) = V(t)$. □

Proposition 10.96 (Constant overhead in inference nodes)**.** *Count one node for each primitive inference, hypothesis lookups included, and count repeated subproofs repeatedly. If the supplied proof of $s = t$ has p nodes, the derivation of $F(s) = F(t)$ has $p + 56$ nodes, independently of the syntactic depth of F.*

Proof. Reflexivity has three nodes; symmetry adds four nodes to its premise and transitivity five to the sum of its premises. Each of the two proofs of $J_F(q) = F(q)$ has eight nodes: a computation, a CC lifting, a second computation and the transitivity overhead. The cases construction adds DE, DI, QE and one clause lookup, so $J_F(t) = F(t)$ has twenty nodes. The two CC liftings of $s = t$ have $p + 2$ nodes, the reversed left representation $20 + 4$, the right one 20, and two transitivities give $(p + 2) + 24 + 20 + 5 + 5 = p + 56$. □

The count is not a bound on bytes or running time: term and formula labels contain copies of F. It is not a constant bound for the lifting to compound formulas below.

Theorem 10.97 (Transport in object formulas)**.** *For every object template $A(w)$, from $\Gamma \triangleright s = t$ and $\Gamma \vdash A(s)$ the calculus $\mathcal{K}_{\mathrm{ctr}}$ derives $\Gamma \vdash A(t)$, with no progress instance in the expansion. An available clause $Q_y B(s, y)$ together with an available $s = t$ has the force of $Q_y B(t, y)$: from a proof of $B(t, 0)$ one obtains $B(t, r)$ for every r.*

Proof. Induction on the number of binders of A, uniformly in contexts, terms and parameters. An equation is Theorem 10.95. Let $A(w) = D_{x,y}B(w, x, y)$ with the bound variables renamed apart. Eliminate $A(s)$ at a fresh a, which retains $Q_y B(s, a, y)$. Introduce $D_{x,y}B(t, x, y)$ by cases with the witness a and a fresh z: the local assumption $B(t, a, 0)$ is transported backward along $t = s$ by the induction hypothesis to $B(s, a, 0)$; the clause at Sz gives $B(s, a, Sz)$; the induction hypothesis transports it forward to $B(t, a, Sz)$. Close the introduction and the elimination. Both recursive uses concern the smaller matrix B, and the only introductions added are case introductions. For the clause, transport the base backward, specialize, and transport forward. □

Corollary 10.98 (Identity-language equivalence)**.** *Let $\mathcal{K}_{\mathrm{tr}}$ be $\mathcal{K}_{\mathrm{ctr}}$ with RL and CC replaced by unrestricted transport in object formulas. For every context Γ of object formulas and clauses and every object formula A, $\Gamma \vdash A$ in $\mathcal{K}_{\mathrm{ctr}}$ if and only if $\Gamma \vdash A$ in $\mathcal{K}_{\mathrm{tr}}$. The translations are effective and preserve end sequents literally.*

Proof. From $\mathcal{K}_{\mathrm{tr}}$, replace each transport by Theorem 10.97. Conversely RL is transport with the template $w = r$; reflexivity follows from a computation by the same instance; and CC is obtained by transporting the left occurrence of s in $\mathsf{R}_S(s, a; u, v.h) = \mathsf{R}_S(s, a; u, v.h)$ to t and applying symmetry. □

Since the language and the meaning of every primitive are unchanged, the bi-interpretation of $\mathcal{K}_{\mathrm{tr}}$ with iPRA, with whole domains, actual equality and identity return isomorphisms, transfers to $\mathcal{K}_{\mathrm{ctr}}$. In the successor-seeded basis the numerical bridge is short: with $C(n,a,b) := W(\mathsf{R}_S(n,a;u,v.Sb))$ and $P(n) := C(n,0,W(n))$ one has $C(0,a,b) = a$, $C(Sn,a,b) = b$, $P(0) = 0$, $P(Sn) = n$, and for terms g and h the term $W\big(\mathsf{R}_S(n,g(\vec{p});u,v.Sh(u,W(v),\vec{p}))\big)$ satisfies the equations of the primitive recursion with base g and step h, without predecessor search, pairing or induction. The package attributes this compiler, the logical macros and the lifting of Theorem 10.97 to earlier drafts; its own contribution is Theorem 10.94 and Theorem 10.95, with the computation schemas unchanged and the counter as the only congruence position. An earlier draft on fixed-position transport uses replacement in the initializer of a scalar iterator together with an additional first-step computation schema.

Theorem 10.99 (Root replacement is not derivable)**.** *Let $\mathcal{K}^-$ be $\mathcal{K}_{\mathrm{ctr}}$ without RL, with no reflexivity or other equality rule added. Then $\mathcal{K}^-$ does not derive $0 = 0$ from the empty context. In particular RL is not admissible in $\mathcal{K}^-$.*

Proof. Interpret the atom $s = t$ by a binary predicate, not assumed to be equality. The domain is $\{0,1\}$, 0 denotes 0, and every recursion term denotes 1 under every assignment, whatever its counter, initializer, update and parameters; this is compatible with substitution and renaming. So Sa is constantly 1. Let $s = t$ be true exactly when s has the value 1; the right side is immaterial. Read $Q_y B$ and $D_{x,y} B$ by their ordinary readings.

R0, RS and the conclusion of CC have a recursion term on the left and are true. Hyp, availability, QE and DE are sound as logical rules. For DI by cases let b be the value of the witness; a sound premise gives $B(b,0) \to B(b,1)$, because Sz denotes 1, and on the two-element domain this is $B(b,0) \to \forall y\, B(b,y)$. For DI-progress, with parameters and witness fixed, a term $U(y)$ has one of the value functions $y \mapsto 0$, $y \mapsto 1$, $y \mapsto y$, so the truth set of a literal equation $U(y) = V(y)$ is $\varnothing$, $\{0,1\}$ or $\{1\}$; in each case truth at 0 implies truth everywhere, and the closure holds for every witness. So every rule of $\mathcal{K}^-$ is sound, and $0 = 0$ is false. In $\mathcal{K}_{\mathrm{ctr}}$ two copies of $\mathsf{R}_S(S0,0;u,v.u) = 0$ as major and minor of RL prove $0 = 0$. □

The theorem concerns the specified inference system. The interpretation is not a model of an ordinary theory whose equality carries its usual axioms, and the theorem does not exclude a bi-interpretation with a different equality or domain.

10.12 Constructed sources and certificate interfaces

This section draws on `constructed_computation` and `uniform_predecessor` for the computation interface, and on `checker_and_barriers`, `compact_kernel`, `local_scopes` and `research` for statements about their checkers. The object calculus is N_{p} with Comp throughout; the results change the certificate format, not the theory.

10.12.1 Successor sources constructed from predecessors

Earlier certificate formats submit an iterator source $[c,a,h]$ and must check that the counter c has successor shape before using a successor computation; deleting that check admits a false equation. The constructed-source formats submit the predecessor.

Definition 10.100. A *constructed computation packet* supplies a predecessor p, an initializer a, an update h and a mode. In base mode the checker generates $I(0,a;\, v.\, h) = a$. In step mode it constructs the successor itself and generates

$$I(Sp,a;\, v.\, h) = h[I(p,a;\, v.\, h)/v].$$

In `constructed_computation` the mode is an explicit flag in a previously unused packet field, and the generated equation is opened at the fresh parameter of the continuation. In `uniform_predecessor` the mode is read off the raw representation of p (an empty representation selects the base equation), the packet is marked by a null major, and the generated equation is opened with the image zero. In both, the recursive image $I(p, a;\, v.\, h)$ is validated as a numerical term only when opening the update requests it.

Lemma 10.101 (Known-instance discharge)**.** *If $w \notin \mathrm{FV}(\Gamma, A)$, $\Gamma \vdash H(0)$ and $\Gamma, H(w) \vdash A$, then $\Gamma \vdash A$.*

Proof. Rename the eigenparameters of the second derivation, substitute 0 for w, which leaves Γ and A unchanged, and cut with the first derivation. □

Lemma 10.102 (Boxed zero)**.** *Put $B_0(p) := I(0, p;\, w.\, 0)$. The base computation and the successor computations at predecessors of the form $B_0(p)$ derive every instance of the successor computation.*

Proof. The base computation gives $B_0(p) = p$. Compute at $B_0(p)$ and transport along $B_0(p) = p$ with the template $I(Sw, a;\, u.\, h) = h[I(w, a;\, u.\, h)/u]$. The argument holds for every element of the domain. □

Theorem 10.103 (Constructed computation supply)**.** *(i) In the format of `constructed_computation`, every successfully generated equation $H(w)$ has a zero instance $H(0)$ derivable from the computation equations and transport. Hence its continuation is admissible by Theorem 10.101.*

(ii) In the format of `uniform_predecessor`, every successfully supplied equation is a theorem uniformly in every newly admitted parameter; no known-instance discharge is needed.

(iii) Both formats derive exactly the closed theorems of N_{p} with Comp*. In the first, the step at the predecessor 0 is the packet with $p = 0$ in step mode. In the second, a literal 0 selects the base equation, and the step at 0 is obtained by Theorem 10.102.*

Proof. Validation separates into two obligations: the source counter is a successor, and a recursive image used in the result is a valid term. Construction establishes the first, and demand-driven substitution the second.

(i) In base mode, after $w = 0$ the initializer and its copy on the right agree by structural zero completion, and the equation is the base computation. In step mode, suppose first that opening the update requests the recursive image. The image $[p, a, h]$ is then validated at numerical root; successful validation is idempotent, so validating on demand gives the same substituted value as validating eagerly. The additional dummy slot below the constructed successor introduces no dependency, since root validation of the image rejects a meaningful reference to it. The two occurrences of the predecessor agree and the zero instance is the successor computation. If the update never requests its image, it is independent of the state on every meaningful path, and after zero completion the equation is $I(Sp_0, a_0;\, v.\, h_0) = h_0$ with h_0 independent of v, which is a successor computation without the unused image. The old goal and context were checked before w was allocated and do not contain it.

(ii) With a null major the continuation opens the equation with the image zero, and the newly admitted parameters stay rigid. For a predecessor with empty representation the decoded counter is zero and the two occurrences of the initializer agree, so the equation is a base computation. Otherwise the source counter is the constructed successor, a successor whatever the value of the predecessor. If the update uses its state, the image is checked at the old parameter floor and the opening lemma of the inherited codec identifies the right side

with the update at the preceding iterator. If not, the update is independent of the state, and the successor computation with a state-independent update holds for every predecessor and initializer, including new parameters.

(iii) Soundness is (i), (ii) and the unchanged logical branches. For completeness, encode each instance of Comp by its packet and close its use by Hyp; the other instructions are unchanged. □

The raw codec is inherited. Input is a JSON array that ends with the closed goal; four-field packets are consumed from the end; membership in the context closes a goal without a packet. One list grammar is read in two sorts, with arities standing for truth or zero, closure or successor, equation, and paired binder or iterator. Negative coordinates name allocated rigid parameters and positive coordinates name protected binder slots. An opener substitutes a checked image at exposed numerical coordinates. These conventions are syntax, not arithmetic operations.

The uniform format shortens certificates: a step at a scalar parameter is one record where the earlier flat format needs three (boxing, computation, transport). For the same named derivations and the delivered encoders the record counts are as follows; they are not lower bounds on shortest proofs.

Named derivation	Earlier format	Uniform predecessor
Nonvacuous zero witness	4	2
Atomic progress closure	6	4
Arbitrary-term specialization	8	6
Contextual induction with seed congruence	48	40

10.12.2 Statements about the checkers of the packages

Composition of the arity codec with predecessor packets (`checker_and_barriers`, `compact_kernel`). The arity codec encodes $s = t$ as $[s, t]$, $D_{x,y}B$ as $[B]$ and Q_yB as $[B, B, B]$, with the depth increment $5 - \mathrm{len}$ for every array, numerical root depth three, and fixed truth and zero aliases at the other arities. The composed checker uses the packet $[\texttt{null}, p, a, h]$ of Theorem 10.100. For a truthy p it corresponds to the arity packet $[[\,], [p, a, h], 1, 0]$ and for a falsy p to $[[\,], [0, a, h], p, 0]$: the generated raw equation, the image checks at the old floor, the next floor and the opened goal and hypothesis are identical in the two executions, so the inherited supply proof applies. The literal 0 stays available as a witness of an introduction, and the step at the predecessor 0 is the three-node derivation of Theorem 10.102.

Theorem 10.104 (Composed verifier)**.** *Every successful execution of each of the supplied composed checkers has a decoded conclusion derivable in* N_p*, and every closed theorem of* N_p *has a finite accepted canonical certificate, given sufficient resources.*

Proof. Induction on the successful trace. Membership is Hyp, by raw equality and the opening invariant: at a meaningful sorted root a successful opening implements capture-free filling of the distinguished hole, by induction on the tree, and the interpretation of a constant alias is unchanged by traversal of its children. A proved equality or closure major with matching endpoints is transport or QE; a mismatch leaves the goal and adds truth. Triplication of a unary major is DE at a fresh parameter. The positional argument check makes an introduction a genuine instance, with the restriction of progress to literal equations. Arithmetic supply is Theorem 10.103. Parameters admitted inside a completed major and unused by its end sequent are replaced by 0. For completeness, encode a derivation rule by rule, with the boxed step at the predecessor 0. □

The invariant is conditional: opening is not idempotent on every raw tree below ignored annotations. The two packages obtain the same 612-byte program.

Local eigenparameter scopes (`local_scopes`). The inherited checker decreases one global integer at every allocation, so the proof of a major permanently advances the allocator of the continuation. The local-scope checker passes the current floor to each call by value.

Theorem 10.105 (Local scope preserves the calculus)**.** *With the same sorted decoding and rule correspondence, the local-scope checker is sound for* N_{p} *with* Comp*, and every finite canonical derivation has a local-scope certificate.*

Proof. At the entry of a call with floor k and context g the scope contains the labels $k, \dots, -1$, and $k = -|g|$, since each continuation appends one context entry and decreases the floor by one. Induct on a successful trace. The entry traversal checks the goal at the current scope before a name is allocated. A major is checked at the same scope, so its conclusion contains no name allocated inside its subproof, and context extension builds a new list, so its local assumptions do not escape. The caller then allocates $k - 1$; the discharged side proof imposes no freshness requirement on this independent binding. The openings, the transport, the closure use, the paired rules and the generated arithmetic assumption are justified as before; the generated assumption needs only a derivable zero instance, by Theorem 10.101. For completeness, rename the binders of a named derivation, label the active eigenparameters at a branch with d context entries by members of $-1, \dots, -d$, serialize a major with the same environment and discard its private extensions on return. □

The theorem states preservation of the calculus under scope-respecting renaming, not equality of accepted byte strings: two sibling subproofs may reuse a raw label, and an old packet may mention a name from a finished side proof.

Iterative continuations (`research`). The loop checker keeps recursion for majors and runs continuations in a loop over an immutable context.

Lemma 10.106 (Continuation simulation)**.** *On canonical certificates and with unbounded resources, the iterative checker has the same successful derivations as the recursive flat checker, with the same order of allocation of fresh names.*

Proof. The invariant is that the current goal, the contents of the context, the global freshness floor and the unconsumed instruction stack equal those of the corresponding recursive call. Hypothesis closure is the same membership test. The recursive call for a major uses the same preceding context, and its allocations persist in both programs. Every branch constructs the same next goal and added assumption; tuple concatenation has the contents of list concatenation and changes no earlier context. Opening the goal and opening the assumption have no effect on the floor, so their order is immaterial. Induction on the trace gives both directions. □

Persistence of the context is necessary by Theorem 10.67. The loop removes the dependence of recursion depth on the number of continuation steps, not on nested majors, syntax traversal or parsing.

10.13 Two partial routes

This section draws on the second research note of `scope_kernel` and on `local_scopes`. Both results are conditional: each isolates the statement that is missing for a complete replacement of a rule.

10.13.1 From uniqueness of iteration toward induction

Work in intuitionistic logic with equality over the scalar iterator, with the two computation equations. The *uniqueness rule* is

$$\frac{\Gamma \vdash f(Sz) = h(f(z))}{\Gamma \vdash f(t) = I(t, f(0);\, u.\, h(u))}\ \mathrm{UR},$$

where f and h are fixed term templates, z is fresh for Γ, t and the parameters, and $h(u)$ does not contain z. The premise has no discharged induction hypothesis and there is no separate base premise. Uniqueness of recursion as an arithmetic principle is classical for richer languages with counter-aware recursion and product types; the question here is the scalar update.

Lemma 10.107. *(i) Induction for literal equations derives UR, without coverage or an equality test.*

(ii) UR derives: from $f(0) = g(0)$, $f(Sz) = h(f(z))$ and $g(Sz) = h(g(z))$ with z fresh, $f(t) = g(t)$.

(iii) UR derives $I(t, 0;\, u.\, Su) = t$.

Proof. (i) Induct on $f(n) = I(n, f(0);\, u.\, h(u))$: the base is the zero computation, and $f(Sn) = h(f(n)) = h(I(n, f(0);\, u.\, h(u))) = I(Sn, f(0);\, u.\, h(u))$. (ii) Apply UR to f and to g and use congruence in the initializer. (iii) Take $f(z) = z$ and $h(u) = Su$; the premise is reflexivity. □

Part (i) is formalized for the kernel as `NPrf.uniq`.

Lemma 10.108 (Retraction lemma)**.** *Let $e(x)$ be a term template such that Γ proves, with x fresh,*

$$e(0) = 0, \qquad e(e(x)) = e(x), \qquad e(S(e(x))) = S(e(x)).$$

Then the computation equations, equality reasoning and UR prove $\Gamma \vdash e(t) = t$. No induction is used.

Proof. Put $h(u) = e(S(e(u)))$ and $r(n) = I(n, 0;\, u.\, h(u))$. Idempotence at $S(e(u))$ gives $e(h(u)) = h(u)$, and idempotence under S and e gives $h(e(u)) = h(u)$; the third hypothesis is $h(u) = S(e(u))$. For $f(n) = e(r(n))$: $f(0) = e(0) = 0$ and $f(Sn) = e(h(r(n))) = h(r(n)) = h(e(r(n))) = h(f(n))$. UR and replacement of the equal initializers give $e(r(n)) = r(n)$; so the whole orbit of h lies in the image of the retraction. Then $r(Sn) = h(r(n)) = S(e(r(n))) = S(r(n))$ and $r(0) = 0$, so UR for r with the update Su and Theorem 10.107(iii) give $r(n) = n$. Hence $e(n) = e(r(n)) = r(n) = n$. □

Proposition 10.109 (Induction from a term-defined split)**.** *Let $E(n)$ be a formula with $\Gamma \vdash E(0)$ and $\Gamma, E(z) \vdash E(Sz)$, z fresh. Suppose there is a term $e(n)$ such that, uniformly in a fresh n, $\Gamma \vdash E(n) \vee \neg E(n)$, $\Gamma, E(n) \vdash e(n) = n$ and $\Gamma, \neg E(n) \vdash e(n) = 0$. Then UR proves $\Gamma \vdash E(t)$.*

Proof. A case split proves $E(e(n))$: in the first branch by $e(n) = n$, in the second by $e(n) = 0$ and the base. The true-branch identity at $e(n)$ gives $e(e(n)) = e(n)$. The step at $e(n)$ gives $E(S(e(n)))$, and the true-branch identity gives $e(S(e(n))) = S(e(n))$. The base gives $e(0) = 0$. Theorem 10.108 gives $e(n) = n$, and transport of $E(e(n))$ gives $E(n)$. □

Corollary 10.110 (Induction for zero equations)**.** *With coverage and separation, UR derives: from $\Gamma \vdash q(0) = 0$ and $\Gamma, q(z) = 0 \vdash q(Sz) = 0$, $\Gamma \vdash q(t) = 0$. Only coverage, separation, the selector computations, equality reasoning and UR are used.*

Proof. Coverage gives the case distinction for every term, and $Sx = 0$ is contradictory: the template $v \mapsto \mathrm{sel}(v, 0, S0)$ gives $S0 = 0$. So $q(n) = 0$ is decidable. Put $e(n) = \mathrm{sel}(q(n), n, 0)$. If $q(n) = 0$ then $e(n) = n$; otherwise coverage for $q(n)$ leaves the successor alternative, where $e(n) = 0$. Apply Theorem 10.109. □

Coverage decides whether one number is zero; it does not give a term $\delta(a, b)$ with $\delta(a, b) = 0 \leftrightarrow a = b$. If the arithmetic with UR constructed such a term and proved the equivalence on its whole domain, Theorem 10.110 applied to $\delta(s(n), t(n)) = 0$ would recover induction for literal equations, and with Theorem 10.107(i) the two principles would be equivalent over the common background. The note does not supply this: the available proofs of predecessor and of reflection of equality (Theorems 1.48 and 1.52) use literal-equation inductions with non-constant right sides, and importing them would be circular. The note records one step of a translation. With $Z(u) = \mathrm{sel}(u, 1, 0)$, $N(n) = I(n, 0; u.\, Z(u))$, $a + b = I(b, a; u.\, Su)$, $j(u) = Z(u) + 2N(u)$, $M(n) = I(n, 0; u.\, j(u))$ and $G(n) = I(n, 0; u.\, u + M(u) + 1)$, the comparison $M(G(n)) = N(n)$ can be set up as a comparison of two recurrences with the common update $k(v) = I(Sv, v; u.\, j(u))$, and finite Boolean computations give $k(N(n)) = Z(N(n))$. The README of the repository records that the package `scalar_uniqueness` derives predecessor, reflection of equality and equation induction from UR and so closes this gap; that argument is not part of the note. A native kernel with UR in place of the progress rule would in addition have to derive its equality logic, case distinction and cut without the removed rule.

10.13.2 What fixed countdown uniqueness determines

A draft on scalar countdown uses a bouncing predecessor W and a countdown L with $L(0, a) = a$ and $L(Sn, a) = W(L(n, a))$, and the contextual rule

$$\frac{\Gamma \vdash g(Sz, \vec{p}) = W(g(z, \vec{p}))}{\Gamma \vdash g(t, \vec{p}) = L(t, g(0, \vec{p}))}\ (U_W),$$

with g a fixed template and z fresh. It completes its theory with an additional faithfulness sentence.

Lemma 10.111 (`local_scopes`)**.** *If $\Gamma \vdash f(0) = 0$ and $\Gamma \vdash f(Sz) = S(f(z))$ with z fresh, then $\Gamma \vdash L(f(t), a) = L(t, a)$ for every t and a, by U_W alone. Consequently, if the theory also proves*

$$\forall x\, \forall y\, \big((\forall a\, L(x, a) = L(y, a)) \to x = y\big),$$

it derives $f(t) = t$ for every such f.

Proof. Take z fresh for a and $g(n) = L(f(n), a)$. Then $g(Sz) = L(S(f(z)), a) = W(L(f(z), a)) = W(g(z))$ and $g(0) = L(0, a) = a$. Apply U_W and substitute $g(0) = a$. The second claim instantiates the displayed sentence at $x = f(t)$, $y = t$. □

For example $L(A(t, 0), a) = L(t, a)$ for an initialized successor iterator A. Equality of all countdown observations is a statement about an action and is not equality of the counters. No proof of the displayed faithfulness sentence from U_W is given, and no equivalence with the stronger single-observation sentence $\forall a\, \forall b\, (L(b, SSa) = 2 \to a = b)$ of the draft is claimed.

10.14 Checker sizes

Table 10.2 lists the complete standalone programs of the packages of this chapter with the sizes that the packages state, under the protocol of Section 1.10. The sizes of `scope_kernel`

and of `progress_independence` belong to their checker work and are listed with it in another chapter. The observed minima at the snapshots of the packages are: 277 tokens, 311 nodes and 672 bytes, attained by different programs, for `adaptive_seed`, `research`, `local_scopes` and `uniform_predecessor`; 268, 300 and 616 for `checker_and_barriers` and `compact_kernel`; 281, 315 and 685 for `constructed_computation` and `counter_congruence`. These minima are attained sizes, hence upper bounds on an unknown optimum; they are not lower bounds.

Table 10.2: Checker sizes as stated by the packages. The control column gives tokens / nodes / bytes of the matched control.

Program	Tokens	Nodes	Bytes	Matched control
`adaptive_seed`				
rejected local-scope prototype	279	322	685	277 / 317 / 684
`research`				
iterative checker	276	305	749	278 / 314 / 679 (flat computation)
byte profile	280	319	652	same
explanatory checker	303	339	1401	—
`local_scopes`				
local-scope checker	284	323	668	281 / 322 / 672 (zero-product control)
readable checker	320	363	1426	—
`checker_and_barriers`				
explicit list dispatch	266	310	612	265 / 312 / 616 (old arity, explicit lists, same driver)
compact list predicate	266	312	604	265 / 314 / 608 (old arity, compact lists, same driver)
Boolean-expression profile	267	315	593	274 / 318 / 601 (old affine, same driver)
guarded traversal, tuple context	269	300	690	see below
`compact_kernel`				
selected composition	266	310	612	269 / 306 / 628 (matched old-flat control)
shorter-byte composition	267	315	597	—
smaller-AST composition	267	298	639	—
immutable-context loop	266	296	689	—
`uniform_predecessor`				
selected checker	269	307	629	269 / 306 / 628 (flat computation, matched driver)
predecessor packet, two verdicts	278	315	680	278 / 314 / 679 (flat computation)
ancillary byte profile	269	311	616	see below
ancillary AST profile	268	300	649	see below
readable checker	312	359	1531	—
`constructed_computation`				
selected profile	279	314	682	283 / 315 / 683 (old generator, same initialization profile)
explicit assignments	277	317	684	281 / 318 / 685 (generated assumptions, principal)
AST profile	279	314	709	283 / 315 / 710 (generated assumptions, AST profile)
`counter_congruence`				
readable checker of $\mathcal{K}_{\mathrm{ctr}}$	1617	1822	6834	—
module import only	281	320	680	281 / 318 / 685 (frozen principal baseline)
module import, exact class	280	321	680	same
module import, exact type	281	322	676	same
module import, zero-product test	281	322	672	same

Program	Tokens	Nodes	Bytes	Matched control
inverted hypothesis guard	283	314	751	283 / 315 / 710 (frozen AST baseline)
compact, separate RL, CC, QE	343	395	881	296 / 333 / 704 (general transport)
compact, CC fused into local facts	341	388	892	same

For `checker_and_barriers` the further same-driver controls are the old arity token profile, 264 / 303 / 641, and the old research loop, 267 / 297 / 698; a screen of 105 old profiles under the same rejection driver gives the matched minima 264 tokens, 297 nodes and 601 bytes, of which only the byte minimum is improved, by the 593-byte profile. For `uniform_predecessor` the matched-driver controls are: token leader 268 / 309 / 633, AST leader 271 / 303 / 654, byte leader 272 / 314 / 621, flat computation 269 / 306 / 628, predecessor joint profile 271 / 306 / 637; the selected checker is one node and one byte larger than the matched flat checker, and its position below the unchanged minima comes from rejection by exit status in place of a printed verdict. In `constructed_computation` the arithmetic generator alone changes the explicit program from 281 / 318 / 685 to 277 / 317 / 684. The 612-byte programs of `checker_and_barriers` and `compact_kernel` are the same file.

The implementation devices are inherited source idioms and are not arithmetic principles. In short: the list test `a * 0 == []` on parsed JSON values; a short-circuit expression for guarded elimination; exception-based openers, which reject nonempty strings only through the recursion limit of the runtime and are therefore not compared with structurally terminating parsers; folding of initialization into a default argument; module import in place of a name import; and rejection by nonzero exit status. All programs import only the standard JSON library, and none imports a producer or a theorem library.

10.15 Further questions

The following statements of the packages are not results. First, what was not achieved.

- No package of this chapter obtains a complete axiomatization that is simpler than the common core under the measurement protocol. Where a checker improves a coordinate of the observed minima, the gain is an implementation change to an inherited theory. No delivered program is below all three coordinates in `research`, `local_scopes`, `checker_and_barriers`, `compact_kernel` or `counter_congruence`; `constructed_computation` is below its three minima by two tokens, one node and three bytes; `uniform_predecessor` is below the unchanged minima only through the rejection convention.

- No obstruction of the chapter is a lower bound on the size of a checker, and none excludes equiconsistency or all one-way interpretations. Apart from Theorem 10.28, Theorem 10.107(i), the strictness statement cited after Theorem 10.61 and the odd case of Theorem 10.81, the README of the repository lists none of them as formalized in Lean. In particular the countermodel of Theorem 10.75 and the finite clocks of Theorem 10.79 are written proofs only.

- `progress_independence`: the result is the independence of one rule of the fixed kernel. No smaller arithmetic basis follows from it, and the package does not decide whether uniqueness of iteration for the successor update is equivalent to induction for literal equations; its strengthened countermodel refutes that uniqueness instance and so gives no answer.

- No complete checker is given for the progressive calculus of Section 10.10. No implementation of $\mathcal{K}_{\mathrm{ctr}}$ is below the inherited programs; the compact attempts of the package have no universal audit of their raw aliases.
- `adaptive_seed`: the deletion of derived majors is rejected only in the form of Theorem 10.63. For a redesigned calculus without public closure goals, logical normalization does not justify deleting all derived majors, because equation induction can produce an equality that is then used for transport into a compound formula, and at an arbitrary parameter that induction is not a reducible successor case. This is an open proof obligation, not an incompleteness theorem.
- `compact_kernel`: two directions were examined without result. Replacing computation supply by numerical definitional conversion requires a reducer that respects dummy slots, the state binder, neutral counters and rigid parameters, and gave no smaller checker. Binders in the certificate grammar, with eigenparameters allocated by the opener, require a new proof-opening discipline, and no complete smaller construction was obtained.
- `uniform_predecessor`: binary-tree domains and arithmetic whose behavior depends on the syntax of a body were examined; no reduction of the complete checking mechanism and no valid smaller presentation was found.
- `scope_kernel`: the route from UR to induction is unfinished in the note (Section 10.13); the note claims no bi-interpretation and no smaller checker for a calculus with UR.

Question 10.112 (Relational escapes)**.** Theorem 10.41 and the theorems of Section 10.4 exclude term realizers only, and Theorem 10.54 requires an existential-free axiomatization and finite control. Is there a seedless basis, with positive existence axioms or with an additional primitive, that is bi-interpretable with iPRA through operations defined by graphs, and whose complete checker, with the defining equations and scope rules of the addition counted, is smaller? Under which witness-extraction property do graph interpretations reduce to the term case?

Question 10.113 (Leaving the almost unary class)**.** An independent initializer is one way to combine two unbounded inputs. Is there another operation, such as pairing or addition, that restores completeness of a seedless basis more cheaply, with both interpretation composites proved internally? Theorem 10.57 gives numerical completeness for an unbounded selector and no more.

Question 10.114 (Extent of finite control)**.** Does finite control, and with it Theorem 10.54, survive the adjunction of bounded numerical primitives other than an equality characteristic? Is every unary primitive recursive function denoted by a tied-seed term? Neither is decided by Theorem 10.48.

Question 10.115 (A common barrier)**.** Theorem 10.32 and Theorem 10.36 are incomparable. Does the conclusion hold for constructors in which both the seed and the update are computed by adaptive circuits of queries to one body, with register vectors and counter-dependent probe locations? No package states this.

Question 10.116 (Weaker case principles)**.** Is there a primitive rule weaker in syntax than BoolCases that supplies the mixed-parameter elimination refuted by the separator of Theorem 10.17, supports both formula round trips and admits a smaller checker? Is there a restriction of numerical cases that avoids the product model of Theorem 10.16 and still derives cases for all native formulas? A proposal must say where the global case distinction enters; predecessor identities do not supply it.

Question 10.117 (Positive iteration)**.** Is there a small operation, cheaper than the split into base and step, that destroys the zero-profile invariant of Theorem 10.58 and recovers a selector and the logical bootstrap? Can initialized iteration be introduced relationally from positive iteration without assuming the selector-based logic that it is meant to reconstruct, and how does induction transfer through the graph definitions when new updates are nested?

Question 10.118 (Majors and cuts)**.** Can atomic cuts be compiled away without induction on compound formulas, so that a focused calculus with fewer proved majors is complete? A normalization theorem would have to handle equation-induction proofs used for transport in arbitrary formulas, or an explicit generator of local induction clauses would have to be counted in the checker.

Question 10.119 (Uniqueness and countdown)**.** Does a complete checker for a calculus with UR improve the size coordinates? Removing an induction hypothesis does not decide this, since the conclusion of UR constructs several term instances and an iterator. Does U_W alone prove the faithfulness sentence of Theorem 10.111, so that the additional observation axiom of the countdown draft can be removed?

Question 10.120 (The countermodel to progress)**.** Can Theorem 10.71 and the tower of Theorem 10.73 be formalized in the Lean development, with arbitrary carriers and an interpreted successor in place of readings over the natural numbers? Which parts of the extension lemma survive when further laws of iteration, such as fusion or uniqueness, are imposed on the coherent structures? Computation and extensionality in the update are compatible with the countermodel; uniqueness for the identity update excludes its particular witness, and excluding one model does not give arithmetic strength. Is there a counterpart of the model that satisfies the uniqueness rule for the successor update and still refutes an induction for a literal equation, which would separate the two principles? Is there a presentation that combines computational equality with an induction interface weaker than QI without moving the cost into syntax validation?

Question 10.121 (Counter congruence and the reversed binder)**.** Can RL, CC and numerical computation be encoded through one checked representation that deletes a trusted operation, so that the restricted interface of Section 10.11 removes code? Can a staged introduction with delayed witness choice repair the reversed binder of Theorem 10.65, with the reverse formula composite proved for nested formulas?

Question 10.122 (Formalization)**.** The packages name as formalization targets: the bi-interpretation of the native calculus with its ordinary realization, with formula composites, freshness and arbitrary contexts; the scalar predecessor and the reflection of equality, without reintroducing quantifier-free induction before the characteristic is proved correct; the derivation transformer for $J_F = F$ of Theorem 10.94, without use of the ambient equality substitution; the sorted codec with its opener, freshness conditions and input contract; and a proved decoder that connects the arity checker with the Lean semantics of the native core. None is supplied.

Question 10.123 (Metric)**.** A coordinatewise envelope is not one program and not a lower bound. Would a fixed implementation grammar, a fixed semantic contract, a fixed treatment of the host runtime and a policy on aliases make strict-improvement claims more informative? Triplicated closures make explicitly serialized certificates grow as 3^r in closure depth, and the protocol charges checker source only; how should certificate size and checking time enter the comparison? The empirical target left by `checker_and_barriers` is one structurally terminating complete program below 593 bytes, 264 tokens and 297 nodes under matched drivers.

Chapter 11

Certificates and checkers

The packages of this chapter do not change a rule basis. Each of them fixes a calculus that is established elsewhere in the article, specifies a format of proof certificates for it, and gives a program that accepts exactly the certificates of derivable sequents. The mathematical content is the specification of the format, the invariants of successful runs, and two kinds of theorem: a checker is sound and complete for its calculus, or two checkers of one calculus accept the same certificates. The programs are small, and many of their steps are safe only because of an invariant of the whole run; these invariants are stated and proved here. Program text is quoted only where an invariant refers to it.

All checkers of this chapter share one encoding of syntax (Section 11.2) and one instruction format (Section 11.3). They differ in a small number of independent devices, each of which is introduced once:

Device	Section	Introduced by
initializer dispatch, certified zero	Section 11.3	`certified_zero`
uniform zero, constructed majors	Section 11.4	`uniform_zero`
neutral major premise (empty array)	Section 11.5	`uniform_premise`, `monus_and_neutral_major`
local floors, shared hypothesis expression	Section 11.5	`neutral_scope`
elimination fixed before its major	Section 11.6	`prevalidated_elimination`
scalar neutral premise, tuple loop	Section 11.7	`kernel_refinement`, `compact_update`
validators of goal and certificate	Section 11.8	`validated_core` and successors
inferred major, eager selection, remainder selector, phase by context	Section 11.9	`inferred_major`, `smaller_envelope`, `frontier_review`, `context_phase`

The packages `neutral_scope`, `scope_and_closure` (in its checker part), `compact_update`, `kernel_refinement`, `checked_refinement`, `validated_core`, `validated_presentation` and `scope_kernel` state that they contain no new rule and no new arithmetic; each composes devices of this list, and the composition is what is proved.

Throughout, a checker reads one finite array of JSON values. Its last element is the *goal*; the earlier elements are *packets*, consumed from the end. A run *accepts* if it terminates normally; a failed operation (a comparison of incomparable values, a missing packet, a wrong number of fields) rejects. An unused prefix of packets is permitted. Completeness statements refer to ideal executions without bounds on memory or recursion depth; a physical run can

reject a valid large certificate, and no statement below equates the resource limits of two programs.

11.1 The calculi that are checked

This section collects, from `certified_zero`, `kernel_refinement`, `compact_update` and `validated_presenta` the three rule bases for which the checkers of this chapter are written, and two admissible rules on which every soundness proof relies.

Definition 11.1 (The calculus $\mathcal{A}$)**.** Terms and native formulas are those of Sections 1.1 and 1.3; a context is a finite list of formulas. The calculus $\mathcal{A}$ has the rules

$$\frac{}{\Gamma \vdash A}\ (A \in \Gamma) \qquad \frac{}{\Gamma \vdash I(0,a;\,u.\,h) = a} \qquad \frac{}{\Gamma \vdash I(Sn,a;\,u.\,h) = h[I(n,a;\,u.\,h)/u]}$$

$$\frac{\Gamma \vdash s = t \quad \Gamma \vdash F(s)}{\Gamma \vdash F(t)} \qquad \frac{\Gamma \vdash Q_y B(y) \quad \Gamma \vdash B(0)}{\Gamma \vdash B(t)} \qquad \frac{\Gamma \vdash D_{x,y} B(x,y) \quad \Gamma, Q_y B(b,y) \vdash A}{\Gamma \vdash A}$$

$$\frac{\Gamma, B(t,0) \vdash B(t,Sz)}{\Gamma \vdash D_{x,y} B(x,y)} \qquad \frac{\Gamma, E(t,z) \vdash E(t,Sz)}{\Gamma \vdash D_{x,y} E(x,y)}$$

named Hyp, I0, IS, Tr, QE, DE, DI-case and DI-progress. In Tr, $F(v)$ is an arbitrary native formula template with one designated numerical variable, which may occur any number of times and below binders. In DE the parameter b is fresh for Γ, A and the parameters of B. In both introductions z is fresh for Γ, t and the parameters of the matrix, and in DI-progress the matrix E is a literal equation: its outermost constructor is equality.

The calculus $\mathcal{A}$ is the calculus `IPrf` of the Lean development, and `iprf_iff_nprf` states that it derives the same sequents as the native core N of Section 1.3. Consequently weakening, renaming of parameters, capture-free numerical substitution and cut of an assumption are admissible for $\mathcal{A}$ (Section 1.4), and $\mathcal{A}$ inherits the bridge of Theorem 1.78. The formula $0 = 0$ is derivable: with $r := I(0,0;\,u.\,0)$, I0 proves $r = 0$, and Tr with $F(v) \equiv (v = 0)$ applied to $r = 0$ and $F(r)$ gives $0 = 0$.

Lemma 11.2 (Zero-instance supply)**.** *Let $\vec{b}$ be parameters that do not occur in Γ or A. If $\vdash H(\vec{0})$ and $\Gamma, H(\vec{b}) \vdash A$, then $\Gamma \vdash A$.*

Proof. Rename the eigenparameters of the second derivation away from $\vec{b}$ and substitute 0 for every member of $\vec{b}$. Since $\vec{b}$ does not occur in Γ or A, the result is a derivation of $\Gamma, H(\vec{0}) \vdash A$. Cut the assumption $H(\vec{0})$ against its derivation. □

Lemma 11.3 (Neutral resource)**.** *If T is a closed theorem, then $\Gamma, T \vdash A$ if and only if $\Gamma \vdash A$.*

Proof. Weakening gives one direction and cut gives the other. Only renaming of eigenparameters is used, no numerical substitution. □

Theorem 11.2 is the reason why a checker may add a computation equation to its context without recognizing the equation as an axiom: it is enough that the instance at $\vec{0}$ of the added formula is an instance of I0 or IS. The open formula $H(\vec{b})$ need not be true. For example $H(b) \equiv (I(0,0;\,v.\,0) = b)$ has a computation instance at $b = 0$ and is false at $b = S0$.

Definition 11.4 (Source-selected elimination)**.** For a template $F(v)$ and terms s, t put

$$M(s,t;F) := \begin{cases} Q_v F(v), & s \text{ is literally } 0, \\ s = t, & \text{otherwise.} \end{cases}$$

The rule SE infers $F(t)$ from $M(s,t;F)$ and $F(s)$. The calculus $\mathcal{A}^-$ is $\mathcal{A}$ with Tr and QE replaced by SE.

Definition 11.5 (The guarded-source calculus $\mathcal{U}$)**.** A formula is *public* if it is an equation, or $D_{x,y}B$ with B public and compound, or $D_{x,y}E$ with E an equation of the shape

$$I(y, 0;\, u.\, h) = I(y, 0;\, u.\, k). \tag{11.1}$$

The calculus $\mathcal{U}$ has Hyp, I0, IS, SE, DE and one introduction

$$\frac{\Gamma, B(t, \eta_B(z)) \vdash B(t, Sz)}{\Gamma \vdash D_{x,y}B(x, y)}, \qquad \eta_B(z) := \begin{cases} z, & B \text{ a literal equation}, \\ 0, & B \text{ compound}, \end{cases}$$

with z fresh as before. Derivations may use all native formulas, including Q and unguarded atomic matrices, as auxiliary formulas; a *public sequent* has public assumptions and a public conclusion. The *auxiliary reading* A^* of a formula in T_{eq} is the ordinary reading of Section 1.2 except that $(D_{x,y}E)^* := \exists x\, \forall y\, (E(x, y) \to E(x, Sy))$ when E is a literal equation.

Remark 11.6 (Imported equivalences). The checkers of Sections 11.7 and 11.8 are written for $\mathcal{A}^-$ and $\mathcal{U}$, whose equivalence with N is not a result of this chapter. The package `source_dispatch` proves that $\mathcal{A}^-$ and $\mathcal{A}$ derive the same sequents (in Lean: `zprf_iff_nprf`); every omitted instance of Tr, with source the literal 0, is derivable. The packages `guarded_core` and `guarded_source` prove that every rule of $\mathcal{U}$ is valid in T_{eq} under the auxiliary reading, that A^* and the ordinary reading of A are equivalent in T_{eq} for public A (the zero instance of a guarded equation (11.1) is a theorem), and that $\mathcal{U}$ derives every public sequent whose ordinary reading is derivable in T_{eq}; the guarded public language is formalized in Lean in the form `pgprf_pub_iff`. Numerical substitution into a derivation is not admissible in $\mathcal{A}^-$ or $\mathcal{U}$ as a transformation of derivations, because it can turn a source into the literal 0. For this reason soundness of a checker of $\mathcal{U}$ is proved below through the reading in T_{eq}, where Theorem 11.2 is available, and not by a direct appeal to Theorem 11.2 in $\mathcal{U}$.

11.2 The sorted arity encoding

The encoding of this section is used by every compact checker of the chapter; it is restated, from an earlier checker, by `certified_zero`, `uniform_premise`, `prevalidated_elimination` and `monus_and_neutral_major`, whose statements are merged here.

Definition 11.7 (Raw trees)**.** A *raw tree* is a scalar or a finite array of raw trees. The *arity* of an array is its length. A scalar is *numeric* if it is an integer, a floating-point number or a Boolean; the Booleans are compared and added as 0 and 1. A raw value is *false-valued* if it is numerically zero, the empty array, or one of the remaining empty JSON values (`null`, the empty string, the empty object); otherwise it is *true-valued*.

A raw tree is read at a position of one of two sorts, term or formula, below some number $p \geq 0$ of *protected slots*. The *depth* of the position is $3 + 2p$ at a term position and $2p$ at a formula position, so that its parity records the sort. Slots are numbered from the innermost, starting with 0. A *floor* is an integer $k \leq 0$.

Definition 11.8 (Sorted reading)**.** Fix a floor k. At a term position below p slots a numeric scalar r reads as

$$\begin{cases} 0, & 0 \leq r < 3, \\ \text{slot } j, & 3 + 2j \leq r < 5 + 2j \text{ and } j < p, \\ \text{the rigid parameter named } r, & k \leq r < 0, \\ \text{the hole } w, & \text{otherwise}, \end{cases}$$

where w is one distinguished variable. Arrays read as follows.

Arity	At a term position	At a formula position
0	0	$0 = 0$
1	S of the child	$D_{x,y}$ of the child
2	0	equality of the two children
3	$I(n, a;\, u.\, h)$ of the children n, a, h	Q_y of the last child
≥ 4	0	$0 = 0$

The child of a unary term array is read below two additional *dummy* slots, which read as 0. The children n and a of a ternary term array are read below one additional dummy slot and the child h below one additional slot, the state u. The children of a binary formula array are read at term positions below the same slots. The child of a unary formula array is read below two additional slots, y innermost and x next; the last child of a ternary formula array is read below one additional slot y. A scalar at a formula position reads as $0 = 0$. Children that the table does not mention (those of a binary term array, of an array of arity at least 4, and the first two children of a ternary formula array) are *annotations* and have no reading.

With child depth $d+5-r$ for a child of an array of arity r at depth d, all the environments of Theorem 11.8 are obtained from one law: the increments are 4, 3 and 2 at arities 1, 2 and 3, that is, two slots, a change of sort, and one slot.

Definition 11.9 (Canonical encoding)**.** The *canonical encoding* of a term or formula whose free variables are rigid parameters with negative integer names writes 0 as the scalar 0, a parameter as its name, the slot j as $3 + 2j$, and

$$St \mapsto [t], \qquad I(n, a;\, u.\, h) \mapsto [n, a, h], \qquad s = t \mapsto [s, t], \qquad D_{x,y}B \mapsto [B], \qquad Q_yB \mapsto [B, B, B],$$

where the slot numbers of variables are counted through the dummy slots of Theorem 11.8. A canonical encoding contains no reference to a dummy slot, and its reading is the encoded expression.

The triplication of the matrix of Q serves the export step of the checker: repeating a unary array $[B]$ three times yields exactly the canonical closure $[B, B, B]$, and repeating any array of arity at least 2 yields an array of arity at least 6, which reads as $0 = 0$.

Definition 11.10 (Opener)**.** For a floor k, a raw tree a, a raw *image* t or the missing image $\perp$, and a depth d, the opening $O_k(a; t, d)$ is defined by

$$O_k(a; t, d) := \begin{cases} [\, O_k(b; t, d + 5 - r) : b \in a \,], & a \text{ an array of arity } r, \\ a, & a \text{ a scalar with } k \leq a < d, \\ O_k(t; \perp, 3), & a \text{ another scalar and } d \text{ odd}, \\ 0, & a \text{ another scalar and } d \text{ even}. \end{cases}$$

It is undefined if a comparison is applied to a value that is not numeric, or to $\perp$. The tree a is *validated at depth d and floor k* if $O_k(a; \perp, d)$ is defined, and *closed at depth d and floor k* if its reading at a position of depth d and floor k contains no hole. The depth is omitted when it is 0, the depth of the formula root.

The image is inserted after being validated at the root term depth 3 with a missing image, so an insertion never starts a further substitution; the opener terminates by structural recursion on finite trees. The checkers differ in the test that recognizes an array (an explicit type test, the equation `a*0 == []`, which separates arrays from the other JSON values on which it is defined, or an attempted traversal that falls back to the scalar case), and in the spelling of $\perp$ (`None`, the complex number `0j`, or a function object); each spelling makes every

comparison with $\perp$ fail. The profile that recognizes arrays by attempted traversal accepts a few further empty iterables as aliases of the empty array and has one qualification: on a nonempty string an ideal execution does not terminate, and an actual one rejects through its recursion limit. This qualification is attached to that profile wherever its size is quoted (Section 11.11).

Lemma 11.11 (Semantic opening)**.** *Let a be read at a position of depth d and floor k, and suppose that $a' = O_k(a; t, d)$ is defined.*

(a) The tree a' is closed at depth d and floor k.

(b) If $t = \perp$, then a is closed at depth d and floor k, and a' and a have the same reading.

(c) If $t \neq \perp$ and the hole occurs in the reading of a, then $t' = O_k(t; \perp, 3)$ is defined, the reading τ of t at the root term position is a term whose only free variables are parameters with names in $[k, 0)$, and the reading of a' is the reading of a with τ substituted, without capture, for the hole. If the hole does not occur, a' and a have the same reading.

Proof. Statement (b) is proved first, by induction on a. A scalar at a term position is retained exactly when it lies in $[k, d)$, that is, when it does not read as the hole; otherwise the comparison with $\perp$ fails. A scalar at a formula position is retained or replaced by 0, and both read as $0 = 0$. At an array of arity 1, 2 or 3 the children that have a reading are opened at exactly the depths of their positions, and the induction hypothesis applies; annotations are traversed, which can make the opening undefined but does not affect a reading. Arrays of other arities have a constant reading and keep their arity.

For (c), the same induction applies, with one new case: a scalar that reads as the hole is replaced by t'. By (b) for t at depth 3, the tree t' reads as τ at the root term position, and every scalar of t' at its root level lies in $[k, 3)$, so it reads as 0 or as a parameter in $[k, 0)$; a scalar of t' below q of its own binders lies in $[k, 3 + 2q)$ and refers to 0, to a parameter, or to one of these q binders. Hence t' has the reading τ at every term position, below any number of slots: copying it below further binders captures nothing and moves none of its indices. Statement (a) follows from (b) and (c), since every retained scalar at a term position lies in $[k, d)$. □

No claim is made that an opened tree is unchanged by a second opening: annotations below arrays of arity at least 4 are traversed at depths that may decrease, and a scalar at a formula position is not a fixed point of every traversal. Only readings at positions that have one are used. Since the reading is a function of the raw tree, equal raw trees have equal readings.

Lemma 11.12 (Exposure of the outermost slot)**.** *If a is closed at depth $d + 2$ and floor k, then its reading at depth d is its reading at depth $d + 2$ with the outermost slot replaced by the hole. Consequently $O_k(a; t, d)$, when defined, reads as the reading of a at depth $d + 2$ with τ substituted for the outermost slot.*

Proof. Let the position of depth $d + 2$ lie below $p + 1$ slots. A scalar r at a term position with a reading, below q further binders, lies in $[k, 3 + 2(p + 1 + q))$ because the reading has no hole. At depth d the same position has depth $3 + 2(p + q)$: a scalar in $[k, 3 + 2(p + q))$ keeps its reading, and a scalar in $[3 + 2(p + q), 5 + 2(p + q))$, which read as slot $p + q$, the outermost of the $p + 1$ slots, now reads as the hole. The second statement is Theorem 11.11(c). □

Theorem 11.12 covers the three openings of a checker. For a closed goal $[B]$, the matrix B is closed at depth 4 with slots y and x; $O_k(B; t, 2)$ instantiates the witness slot x and keeps y. For a matrix B' closed at depth 2, $O_k(B'; t, 0)$ instantiates its index slot y. For a closed

major $[B]$, the opening $O_k([B,B,B];t,0)$ opens each copy of B at depth 2: it instantiates x and leaves y bound by the closure.

Lemma 11.13 (Canonical opening)**.** *Let a be the canonical encoding of an expression below $p+1$ slots whose parameters have names in $[k,0)$, read at depth d below p slots, and let t be the canonical encoding of a term whose free variables are parameters with names in $[k,0)$. Then $O_k(a;t,d)$ is the canonical encoding of the expression with that term substituted for the outermost slot, and $O_k(a;\bot,d+2)=a$.*

Proof. A canonical encoding uses the scalar 0, parameter names in $[k,0)$ and slot scalars $3+2j$, all retained when in range; the exposed slot scalars are replaced by $O_k(t;\bot,3)=t$. The encoding of a term without free slots does not depend on the number of slots above it, so the result is canonical. □

Lemma 11.14 (Freshness)**.** *If a tree is closed at depth d and floor k, its reading contains only parameters with names in $[k,0)$, and it is closed at depth d with the same reading at every floor $k' \le k$. Hence the names in $[k',k)$ are fresh for every formula that is the reading of a tree closed at floor k.*

Proof. Lowering the floor changes the reading only of scalars below k, and a tree closed at floor k has none at a position with a reading, since such a scalar reads as the hole. □

For canonical data there is exactly one name in $[k-1,k)$, the integer $k-1$. An arbitrary certificate may contain fractional scalars; then finitely many names lie in $[k-1,k)$, they are distinct parameters, and the arguments below treat them as a finite family $\vec{b}$.

11.3 The instruction format and the certified zero

This section follows `certified_zero`: the base protocol of all compact checkers, the restriction of computation that lets the initializer field select the computation branch, and the correspondence between the checker and the calculus $\mathcal{A}$.

11.3.1 The base protocol

Definition 11.15 (Base protocol)**.** The state of a run consists of a *global floor* k, initially 0, and the stack of unread packets. The procedure Check(a,g), for a raw goal a and a list g of raw trees, performs the following steps.

(1) Compute $a^\circ = O_k(a;\bot,0)$. If a° is a member of g, return.

(2) Remove the last packet. It has exactly four fields $[m,s,u,t]$.

(3) If m is true-valued, run Check(m,g). Afterwards, if $m=[s,u]$, or if s is false-valued and $m=[t,t,t]$, and if moreover $O_k(t;u,0)=a$, replace a by $O_k(t;s,0)$.

(4) Form the hypothesis h.

  (i) If m is true-valued, h is the threefold repetition of m.

  (ii) If m is false-valued and u is true-valued, then $h=[[s,u,t],u]$ when s is false-valued, and $h=[[[s],u,t],\ O_k(t;[s,u,t],3)]$ when s is true-valued.

  (iii) If m and u are false-valued, a is an array $[B]$ of arity one; replace a by $O_k(B;t,2)$ and put $h=a$.

(5) Decrease k by 1.

(6) Let $\iota = k$ if m is true-valued, or if m is numerically zero and the arity of a is even; otherwise let ι be the scalar zero. Run Check$\big(O_k(a; [k], 0),\ g +\!\!+ [O_k(h; \iota, 0)]\big)$.

The checker reads the goal a_0 and runs Check$(a_0, [\,])$.

The three branches of step (4) are selected by data: a true-valued major selects an elimination, a true-valued initializer field u selects a computation, and otherwise the packet is an introduction. There is no operation code. A packet is used as follows on canonical encodings; fields without a role are written 0, and $[\,]$ stands for any false-valued marker that is not numerically zero.

Inference	Packet $[m, s, u, t]$	Effect
Tr with major $s = u$, template F	$[[s, u], s, u, F]$	goal $F(u)$ becomes $F(s)$
QE with major $Q_y B$	$[[B, B, B], 0, u, B]$	goal $B(u)$ becomes $B(0)$
DE with major $D_{x,y} B$	$[[B], 0, 0, 0]$	$Q_y B(b, y)$ is added
I0, initializer a, update h	$[[\,], 0, a, h]$	$I(0, a;\ v.\, h) = a$ is added
IS, predecessor n	$[[\,], n, a, h]$	the IS equation is added
DI-case, witness w	$[[\,], 0, 0, w]$	goal $D_{x,y} B$ becomes $B(w, Sz)$
DI-progress, witness w	$[0, 0, 0, w]$	the same, with $E(w, z)$ added

In a template F the designated variable is encoded as the outermost slot of a formula read below one slot, so that Theorem 11.12 applies; for QE the template is the matrix itself. A derivation is serialized with the packets of the major premise after the packet of the inference and the packets of the continuation after those, and the whole sequence is reversed, since packets are consumed from the end.

Two points of the definition depend on arguments given below. First, in the computation branch the field u must be true-valued, so the equations I0 and IS are available only for an initializer that is not the scalar 0; Theorem 11.17 shows that nothing is lost. Second, step (6) may open a generated computation equation at the fresh parameter instead of at zero (this happens when m is numerically zero and the goal has even arity); Theorem 11.18 and Theorem 11.2 show that this is sound.

11.3.2 Restricted computation and the certified zero

Definition 11.16. The calculus $\mathcal{A}_{\mathrm{r}}$ is $\mathcal{A}$ with its computation rules restricted by the syntax of their arguments: I0 is available only when the initializer a is not the literal term 0, and IS only when neither the predecessor n nor the initializer a is the literal term 0. A term that is not the literal 0 is permitted even if it denotes zero.

The restriction on the predecessor is present in earlier checkers of the same family (`uniform_predecessor`); the restriction on the initializer is the one introduced by `certified_zero`. The term

$$C := I(SS0, S0;\ v.\, 0)$$

is a permitted argument, and the instance of IS with predecessor $S0$, initializer $S0$ and update 0 is permitted and reads $C = 0$.

Theorem 11.17 (Certified zero)**.** *Every instance of I0 and IS is derivable in $\mathcal{A}_{\mathrm{r}}$, by one application of Tr to two permitted computation instances. Hence $\mathcal{A}_{\mathrm{r}}$ and $\mathcal{A}$ derive the same sequents, over the same terms and formulas.*

Proof. A permitted instance is immediate. For an excluded instance choose a fresh variable w, with the bound state variable renamed away from w, and an equation $E(w)$ as follows.

I0, initializer 0 :	$E(w) \equiv (I(0, w;\, v.\, h) = w);$
IS, initializer 0, predecessor $n \neq 0$:	$E(w) \equiv (I(Sn, w;\, v.\, h) = h[I(n, w;\, v.\, h)/v]);$
IS, predecessor 0, initializer $a \neq 0$:	$E(w) \equiv (I(Sw, a;\, v.\, h) = h[I(w, a;\, v.\, h)/v]);$
IS, both 0 :	$E(w) \equiv (I(Sw, w;\, v.\, h) = h[I(w, w;\, v.\, h)/v]).$

Only the displayed occurrences of the initializer and the predecessor are replaced by w; occurrences of 0 inside h stay. In each case $E(C)$ is a permitted computation instance, because C is not the literal 0, and $E(0)$ is the excluded instance. Tr with the template E, applied to the permitted instance $C = 0$ and to $E(C)$, gives $E(0)$. Replacing every excluded leaf of a derivation in this way gives the inclusion of $\mathcal{A}$ in $\mathcal{A}_{\mathrm{r}}$; the other inclusion is trivial. Contexts and end sequents are unchanged. □

The construction uses no induction, no earlier computation with initializer 0 and no decision about the value of a term. Its cost is explicit: if a tree derivation has N inference nodes of which r are excluded computation leaves, the repaired derivation has $N + 2r$ nodes; if the node proving $C = 0$ is shared in a directed acyclic representation and $r > 0$, it has $N + r + 1$ nodes. These are counts for this construction, not lower bounds.

11.3.3 Correspondence with the calculus

Lemma 11.18 (Zero-instance computation)**.** *In step (4ii) of Theorem 11.15, suppose that the opening $h' = O_{k-1}(h; \iota, 0)$ of step (6) is defined. Let $\vec{b}$ be the parameters with names in $[k-1, k)$ that occur in the reading $H(\vec{b})$ of h'. Then $H(\vec{0})$ is an instance of I0 or IS.*

Proof. The tree h is a binary array, so h' reads as an equation; its left side is a ternary term array and reads as an iterator with update η, the reading of t below the state slot.

Let s be false-valued, so $h = [[s, u, t], u]$. The counter s is traversed by the opening; a false-valued tree for which this succeeds is a numeric zero or the empty array and reads as 0. The initializer u is read on the left below the dummy slot of the iterator and on the right at the root. The two readings agree except that a scalar referring to the dummy slot reads as 0 on the left and as the hole, hence as the reading of ι, on the right. The reading of ι is 0 or the parameter named $k-1$. After the substitution of 0 for $\vec{b}$ both sides have the same initializer α, and $H(\vec{0})$ is $I(0, \alpha;\, v.\, \eta) = \alpha$.

Let s be true-valued, so $h = [[[s], u, t], O_k(t; [s, u, t], 3)]$. If the reading of t at depth 3 contains the hole, that is, if the update uses its state, then by Theorem 11.11(c) the image $[s, u, t]$ is validated at depth 3 and floor k. This validates s and u below the dummy slot and t below exactly one state slot, so no coordinate of t beyond its state is left unbound. The right side of h reads as $\eta[I(\sigma, \alpha;\, v.\, \eta)/v]$, where σ and α are the readings of s and u. On the left, s occurs below two further dummy slots of the constructed successor; all its scalars lie in $[k, 5)$, so its reading is again σ. Thus h is closed at floor k and reads as the instance of IS with predecessor σ. If the update does not use its state, the right side $O_k(t; [s, u, t], 3)$ is t validated at depth 3, and η does not contain v. The left side reads, after the substitution of 0 for $\vec{b}$ and for the hole, as $I(S\sigma^0, \alpha^0;\, v.\, \eta)$ for some terms σ^0, α^0, and $\eta = \eta[I(\sigma^0, \alpha^0;\, v.\, \eta)/v]$ because v does not occur in η. This is again an instance of IS. □

Theorem 11.19 (Soundness of the base protocol)**.** *Let every member of g be closed at floor k. If* Check(a, g)*, started at floor k, returns, then a is closed at floor k and the reading of a is derivable in $\mathcal{A}$ from the readings of the members of g. In particular, if the checker accepts a goal a_0, the reading of a_0 is a theorem of $\mathcal{A}$; for a canonical encoding this reading is the encoded formula.*

Proof. The proof is by induction on the number of calls in the run. Floors only decrease, and by Theorem 11.14 a tree that is closed at some floor keeps its reading at all lower floors; readings are written without mention of the floor. Step (1) validates a, so a is closed by Theorem 11.11(b). If the run returns in step (1), a° has the reading of a and is a member of g, and HYP applies.

Elimination. Let m be true-valued and let $k' \le k$ be the floor after step (3). By the induction hypothesis the reading of m is derivable from g, and m is closed at floor k. Suppose that the goal is replaced. If $m = [s, u]$, then s and u are closed at depth 3 with readings σ and υ, the major reads $\sigma = \upsilon$, and by Theorem 11.11(c) the old goal $O_{k'}(t; u, 0)$ and the new goal $O_{k'}(t; s, 0)$ read as $F(\upsilon)$ and $F(\sigma)$ for the template $F(w)$ that is the reading of t at floor k'. By TR it suffices to derive $F(\sigma)$. If s is false-valued and $m = [t, t, t]$, then t is closed at depth 2, the major reads $Q_y B(y)$ for the reading B of t below the slot y, and by Theorem 11.12 the old goal reads $B(\upsilon)$. A false-valued image that is used by an opening is a numeric zero or the empty array and reads 0, so the new goal reads $B(0)$, and by QE it suffices to derive $B(0)$. If the goal is not replaced there is nothing to show at this point. In each case the current goal a_1 is closed at floor k', and it remains to derive its reading from g.

The final call runs at floor $k' - 1$ with goal $O_{k'-1}(a_1; [k' - 1], 0)$, which has the reading of a_1 since a_1 contains no hole, and with the additional hypothesis $h' = O_{k'-1}(h; k' - 1, 0)$, where h is the threefold repetition of m. All members of the extended context are closed at floor $k' - 1$ by Theorem 11.11(a), so the induction hypothesis applies. If $m = [B]$ is unary, then $h = [B, B, B]$, and by Theorem 11.12 h' reads $Q_y B(b, y)$, where $D_{x,y}B$ is the reading of m and b is the parameter named $k' - 1$. By Theorem 11.14, b is fresh for g, for a_1 and for the parameters of B, and DE concludes. If m is an array of arity at least 2, then h has arity at least 6 and h' reads $0 = 0$; Theorem 11.3 concludes. If m is a true-valued scalar for which the run continues, it is numeric, h is a numeric scalar and h' reads $0 = 0$ as well; moreover no replacement of the goal has occurred, since a scalar is not equal to an array. (The program writes the second alternative of the test as a value that is the Boolean `False` when s is true-valued; a true-valued scalar is not equal to it either.)

Computation. Let m be false-valued and u true-valued. The goal is unchanged, and the final call derives its reading from g and the reading $H(\vec{b})$ of h', where $\vec{b}$ are the parameters with names in $[k - 1, k)$. They are fresh for g and a by Theorem 11.14. By Theorem 11.18 $H(\vec{0})$ is an axiom of $\mathcal{A}$, and Theorem 11.2 concludes.

Introduction. Let m and u be false-valued and $a = [B]$. Then B is closed at depth 4, and by Theorem 11.12 $a_1 = O_k(B; t, 2)$ reads, below the slot y, as $B(\tau, y)$, where $D_{x,y}B(x, y)$ is the reading of a and τ is the reading of t if the witness slot occurs, and 0 otherwise; the parameters of τ have names in $[k, 0)$. Let z be the parameter named $k - 1$. In the final call the goal $O_{k-1}(a_1; [k - 1], 0)$ reads $B(\tau, Sz)$ and the hypothesis $O_{k-1}(a_1; \iota, 0)$ reads $B(\tau, z)$ if $\iota = k - 1$ and $B(\tau, 0)$ if ι is zero. The parameter z is fresh for g, for τ and for the parameters of B. The opener preserves arities, so the arity of a_1 is the arity of B. If it is 2, B is a literal equation and DI-progress or DI-case applies. If it is 1 or 3, the arity is odd, ι is zero, and DI-case applies to the compound matrix. If it is 0 or at least 4, the conclusion reads $D_{x,y}(0 = 0)$, which is derivable by DI-case from the theorem $0 = 0$. If B is a scalar, step (6) is undefined and the run rejects. □

Remark 11.20. Three details of the programs lie outside Theorem 11.15 and do not affect Theorem 11.19. On the empty goal [ ] an introduction packet is processed with shifted arguments; the reading of that goal is the theorem $0 = 0$. In two of the three profiles the replacement of step (3) is written as a Boolean expression that keeps the old goal when the new one is false-valued; then the stronger, unchanged obligation is checked. Canonical formulas are nonempty arrays, so neither detail affects canonical certificates. Finally a major is always proved before it is used: a mismatch of the pattern of step (3) leaves the goal unchanged but

does not remove the obligation to prove m.

Theorem 11.21 (Completeness of the base protocol)**.** *Let A be a closed formula. If A is derivable in $\mathcal{A}$, there is a stack of packets on which the checker accepts the canonical encoding of A.*

Proof. By Theorem 11.17 there is a derivation in $\mathcal{A}_{\mathrm{r}}$. The following statement is proved by induction on the height of derivations: for every floor k, every list g of canonical encodings of formulas with parameters named in $[k, 0)$, and every derivation of $\Gamma \vdash A'$ in $\mathcal{A}_{\mathrm{r}}$ with Γ among the formulas encoded in g and parameters named in $[k, 0)$, there is a stack on which Check, called at floor k with the canonical encoding of A' and with g, returns. Weakening and renaming of parameters do not increase the height of a derivation, so the induction hypothesis may be applied to premises in a larger context and with renamed eigenparameters. By Theorem 11.13 the goal is unchanged by step (1). If it is a member of g, the empty stack suffices; this covers Hyp and also every sequent whose conclusion happens to be retained already. Otherwise take the last inference of the derivation and the packet of the table after Theorem 11.15.

For Tr, QE and DE the major is a nonempty array, and the stack for its derivation, which exists by the induction hypothesis, is consumed by step (3) and leaves a floor $k' \leq k$. By Theorem 11.13 the pattern test of step (3) succeeds for Tr and QE and replaces the goal by the canonical encoding of the minor premise; for DE it fails, because a unary array is equal neither to $[0, 0]$ nor to $[0, 0, 0]$. The final call runs at floor $k' - 1$ with the same goal and a context extended by a formula that reads $0 = 0$, or by the canonical encoding of $Q_y B(b, y)$ with the eigenparameter renamed to $k' - 1$. The induction hypothesis for the remaining premise, weakened and renamed, supplies the rest of the stack.

For a computation leaf, which in $\mathcal{A}_{\mathrm{r}}$ has an initializer that is not the literal 0 and, for IS, such a predecessor, the fields u and, for IS, s are true-valued: the canonical encoding of a term other than the literal 0 is a nonempty array or a negative integer. Step (4ii) forms $[[0, a, h], a]$ or $[[[n], a, h], O_k(h; [n, a, h], 3)]$, which by Theorem 11.13 is the canonical encoding of the computation equation; the canonical encoding of a term without free slots is the same below the dummy slots. This is the goal, so the final call returns in its step (1).

For an introduction with witness w the packet has false-valued m and u, step (4iii) opens the matrix at the canonical encoding of w, and step (6) forms the canonical encodings of $B(w, Sz)$ and of $B(w, z)$ or $B(w, 0)$ with z renamed to $k - 1$: the marker $m = 0$ selects $B(w, z)$ exactly when the matrix is an equation, as DI-progress requires, and the marker $[\,]$ selects $B(w, 0)$. The induction hypothesis supplies the stack for the premise. In all cases the stack consists of the stack of the continuation, followed by the stack of the major if there is one, followed by the packet of the last inference. The packet is then consumed first, the major call consumes exactly its own stack, and the final call consumes the rest; that each call consumes exactly the stack constructed for it, when this stack is the end of a longer one, is part of the statement proved by the induction. □

Remark 11.22 (Profiles). `certified_zero` delivers three complete programs of the base protocol, which differ only in the recognition of arrays described after Theorem 11.10: an explicit type test, the zero-product test, and attempted traversal. The first two terminate on every finite input; the third has the qualification on nonempty strings. All three are complete for canonical certificates. The package also isolates the effect of the initializer dispatch from other changes of the source: restoring the earlier test that the major is the null value, in the three final programs, costs exactly eight bytes, two tokens and three syntax-tree nodes in each.

11.4 Uniform zero and constructed majors

This section follows `uniform_zero`. The certified zero of Theorem 11.17 repairs a restriction that exists only because the scalar 0 is at once the canonical zero and a false-valued marker. A different canonical spelling of zero removes the restriction, and the same observation lets an elimination packet omit its major formula.

11.4.1 The recoding

By Theorem 11.8 the scalars 0, 1 and 2 all read as zero at a term position. Let R map a canonical encoding to the tree obtained by replacing each scalar 0 by the scalar 1; arrays, slot scalars and parameter names are unchanged. The trees $R(A)$ are the *canonical encodings with uniform zero*: the term 0 is the scalar 1, the term $S0$ is [1]. The recoding applies to expression fields of packets only; the markers 0 and [ ] in the major field keep their role.

Lemma 11.23 (Uniform truth value)**.** *Every canonical encoding with uniform zero of a term is true-valued, every such encoding of a formula is a nonempty array, and $R(A)$ has the same reading as A.*

Proof. The scalar 1 is true-valued, a parameter name is negative, a slot scalar is at least 3, and the encodings of S, of the iterator, of equations, of D and of Q are nonempty arrays. The scalars 0 and 1 have the same reading at every term position, and R changes nothing else. □

Truth value is a property of the syntax: a true-valued term may denote zero.

Let O_k^+ be the opener of Theorem 11.10 with one change: a scalar a with $k \le a < d$ is returned as a if it is true-valued and as the scalar 1 otherwise.

Lemma 11.24 (Opening commutes with the recoding)**.** *For canonical encodings A and t as in Theorem 11.13,*

$$O_k^+(R(A); R(t), d) = R(O_k(A; t, d)),$$

and likewise with the missing image when no image is used. Opening a canonical template with uniform zero at the raw image 0 *gives the same result as opening it at the raw image* 1*.*

Proof. First let no image be used; the proof is by induction on A. A scalar 0 of A occurs at a term position, where it is retained by O_k; in $R(A)$ it is the scalar 1, retained by O_k^+. Every other retained scalar is true-valued and unchanged on both sides. Arrays are traversed at the same depths on both sides. A canonical formula has no scalar at a formula position, so the formula-position clause does not arise. With an image, the only new case is an exposed scalar, replaced on the left by $O_k^+(R(t); \bot, 3)$ and on the right by $R(O_k(t; \bot, 3))$, which are equal by the first part. For the last statement, $O_k^+(0; \bot, 3) = 1 = O_k^+(1; \bot, 3)$ at every floor. □

Lemma 11.25 (Semantic opening for O^+)**.** *Theorems 11.11, 11.12 and 11.14 hold for O_k^+.*

Proof. The only changed result is a false-valued numeric scalar retained by the range test (the integer or floating-point zero, or the Boolean `False`), which is returned as 1. At a term position both read as 0; at a formula position both read as $0 = 0$. All other clauses are those of O_k. □

The identity of Theorem 11.24 is restricted to canonical encodings; for arbitrary raw trees only the statement about readings in Theorem 11.25 is available, and only it is used for soundness. In contrast with a definitional reduction of $I(0, a; v.\, h)$ to a, the recoding deletes no node and no slot, so no index is shifted.

11.4.2 Constructed majors and local floors

Definition 11.26 (Uniform-zero protocols)**.** The *uniform-zero protocol* is the base protocol with O^+ in place of O and with *local floors*: the floor is an argument $\mathrm{Check}(a, g, k)$ of the procedure, the call for the major in step (3) receives the caller's floor k and leaves it unchanged, and step (5) decreases the caller's own floor. The *constructed-major protocol* changes, in addition, step (3): if the field m is true-valued, the major is

$$M(m, s, u, t) := \begin{cases} [s, u], & s \text{ true-valued}, \\ [t, t, t], & s \text{ false-valued and } t \text{ true-valued}, \\ m, & s, t \text{ false-valued}; \end{cases}$$

the checker runs $\mathrm{Check}(M, g, k)$, and afterwards, if $O_k^+(t; u, 0) = a$ and $O_k^+(t; s, 0)$ is true-valued, replaces a by $O_k^+(t; s, 0)$. Steps (4)–(6) use M in place of m.

Local floors are taken over from the package `local_scopes`. They are justified by the same induction as the global floor.

Lemma 11.27 (Local floors)**.** *Theorems 11.19 and 11.21 hold for the base protocol with local floors.*

Proof. In the proof of Theorem 11.19 the induction hypothesis for the major call is the statement that the reading of m is derivable from g; it does not mention the names that the call allocates, which are eigenparameters of that derivation. With local floors $k' = k$ in the elimination case, the templates are opened at the caller's floor, and the final call at floor $k - 1$ has an eigenparameter that is fresh for g and for the goal, both closed at floor k. The rest of the proof is unchanged. For completeness the stack of the major derivation is produced at floor k and that of the continuation at floor $k - 1$; names used inside the major derivation are reused, which is a renaming of eigenparameters in disjoint subderivations. No call changes the context of its caller, since a context is extended by forming a new list. □

With local floors and an initial context of c entries, every call satisfies $k = c - |g|$: a major call keeps both, and a continuation appends one entry and decreases k. The floor is nevertheless not a function of the context that the program could recompute at entry for free; see Theorem 11.53.

Proposition 11.28 (Soundness of constructed majors)**.** *Theorem 11.19 holds for the constructed-major protocol.*

Proof. By Theorems 11.25 and 11.27 only step (3) needs an argument. If s is true-valued, the proved major is $[s, u]$ and reads $\sigma = \upsilon$; a successful comparison identifies the goal with $F(\upsilon)$ for the reading F of the template t, the new goal is $F(\sigma)$, and Tr applies. No separately supplied formula is claimed to have these endpoints. If s is false-valued and t true-valued, the proved major is $[t, t, t]$ and reads $Q_y F(y)$; a false-valued image that is used reads 0, so the new goal is $F(0)$ and QE applies. If s and t are false-valued, the supplied m is proved and exported as in the base protocol; the comparison $O_k^+(t; u, 0) = a$ can then succeed only for a template t that is a false-valued scalar or the empty array, in which case $O_k^+(t; s, 0)$ is false-valued and the goal is not replaced. □

Theorem 11.29 (Direct canonical compilation)**.** *Every derivation in $\mathcal{A}$, with unrestricted I0 and IS, of a closed formula has a certificate in the constructed-major protocol on canonical encodings with uniform zero. No computation leaf is replaced, and the compilation is the identity on readings.*

Proof. The induction of Theorem 11.21 is repeated with Theorem 11.24 in place of Theorem 11.13 and with the packets

Inference	Packet $[m, s, u, t]$
Tr through F, major $s = u$	$[1, R(s), R(u), R(F)]$
QE from $Q_y B$, target u	$[1, 0, R(u), R(B)]$
DE from $D_{x,y} B$	$[R(D_{x,y} B), 0, 0, 0]$
I0, initializer a, update h	$[0, 0, R(a), R(h)]$
IS, predecessor n	$[0, R(n), R(a), R(h)]$
DI-case, witness w	$[[\,], 0, 0, R(w)]$
DI-progress, witness w	$[0, 0, 0, R(w)]$

By Theorem 11.23 the source $R(s)$ of a transport is true-valued even when s is the term 0, and a matrix $R(B)$ is a nonempty array, so the constructed major is exactly the equation or the closure of the inference. The initializer $R(a)$ is true-valued for every term a, so every computation leaf selects the computation branch. In I0 the generated counter is the raw marker 0, which the opening of the hypothesis returns as the canonical 1; in IS the predecessor $R(n)$ is true-valued even for $n = 0$, and the generated counter [1] is the canonical successor of zero. The zero images of step (6) become the canonical 1 as well. The recoding does not change the root of a matrix, so the restriction of progress to literal equations is unchanged. The arrays of arity 6 or 9 that equality and closure majors add to the context are not used by the certificate. □

By Theorem 11.23 the consequence relation presented on canonical encodings is that of $\mathcal{A}$, read through the identity, so the bridge of Theorem 1.78 applies without change.

11.4.3 Sizes of certificates

Let $|X|$ be the length in bytes of the JSON text of X without whitespace. Then $|[X, Y]| = |X| + |Y| + 3$ and $|[B, B, B]| = 3|B| + 4$. In a transport packet the major field $[s, u]$ is replaced by the marker 1 while s and u remain in their own fields, which saves exactly $|s| + |u| + 2$ bytes; in a closure packet the field $[B, B, B]$ is replaced by 1, which saves exactly $3|B| + 3$ bytes. No inference and no premise is removed by these two changes, and for one derivation with one allocation of names the savings add over its packets, since the recoding of zero replaces one digit by another.

Separately, the direct serialization of computation leaves removes the repairs of Theorem 11.17: a derivation with N nodes and r excluded leaves has N packets instead of $N + 2r \le 3N$, when neither serialization closes an additional goal by a retained hypothesis. The constructed major copies fields that are present at most three times and C is a fixed term, so both translations are linear in the encoded proof; the triplication of matrices in the encoding of Q, which can be exponential relative to a shared representation, is not removed.

Example 11.30. The equation $I(0, 0; v.\, 0) = 0$ has, in the base protocol with the repair of Theorem 11.17, a certificate of three packets and 97 bytes. With uniform zero the certificate is

```
[[0,0,1,1],[[1,1,1],1]],
```

one packet and 23 bytes: the goal $[[1, 1, 1], 1]$ and the packet of I0 with initializer and update 1. The package reports, for sixteen selected examples, 120 packets and 5029 bytes in the base protocol, 90 packets and 3801 bytes with uniform zero, and 90 packets and 2955 bytes with constructed majors. These are measurements of particular certificates, not minimal sizes.

The gain is in the certificates. Under a comparison with a matched control that already uses local floors and the shared hypothesis expression of Section 11.5, uniform zero costs five bytes of checker source and constructed majors a further six (Section 11.11).

11.5 Neutral major premises

This section merges `uniform_premise` and the certificate protocol of `monus_and_neutral_major`, which are two write-ups of one device, and adds the composition with local floors of `neutral_scope`. In the base protocol a packet has a major premise only if its first field is true-valued. Here every packet has one: a computation or an introduction names a fixed proved truth, and the test for the presence of a major disappears. The safety of the resulting program rests on an invariant of successful runs.

11.5.1 The protocol and its hazard

Definition 11.31 (Neutral-major protocol)**.** The neutral-major protocol is the base protocol of Theorem 11.15 with the following changes.

(N1) The initial call is Check$(a_0, [[\,]])$: the context contains the empty array, which reads $0 = 0$.

(N2) In step (3), Check(m, g) is run for every packet, whether or not m is true-valued. The replacement test that follows is unchanged.

(N3) In step (4), branch (i) is taken if the threefold repetition of m is true-valued, branch (ii) if it is not and u is true-valued, and branch (iii) otherwise.

(N4) In step (6), $\iota = k$ if m is true-valued, or if the arity of a taken modulo 2 is numerically equal to u; otherwise ι is zero.

On canonical certificates a packet without a substantive major has $m = [\,]$, an introduction by cases has $u = [\,]$, and an introduction by progress has $u = 0$.

The call Check$([\,], g)$ returns in its step (1) without reading a packet and without changing the floor, because $[\,]$ is a member of every context of the run. The mode of an introduction moves from the first field to the initializer field, which is false-valued in an introduction.

The program evaluates the test of step (3) as membership of m in the pair consisting of $[s, u]$ and of the value that is $[t, t, t]$ if s is false-valued and the Boolean `False` otherwise. In the base protocol the test is reached only for a true-valued m. Now it is reached for every m, and the Boolean `False` is equal to the numeric scalar 0. If a numeric zero could be a proved major, a packet with $m = 0$ and a true-valued s would replace the goal $F(u)$ by $F(s)$ with no equation proved. The protocol is therefore sound only if a numeric scalar is never a major of a returning call, and this is what the invariant below establishes. It holds because the initial context is $[[\,]]$ and not $[0]$.

11.5.2 The invariant of successful runs

Lemma 11.32 (Root openings)**.** *For a numeric scalar r and any image t,*

$$O_k(r; t, 0) = \begin{cases} r, & k \le r < 0, \\ 0, & \textit{otherwise}, \end{cases}$$

which is numeric and does not depend on t. If $O_k(a; t, 0)$ is defined and a is an array, the result is an array; if the result is numeric, a is numeric.

Proof. The depth 0 is even, so a scalar outside $[k, 0)$ is replaced by 0 and no image is consulted. An array is mapped to an array by the first clause of Theorem 11.10, and a scalar that is not numeric fails the comparison. □

Lemma 11.33 (No numeric goal)**.** *In the neutral-major protocol there is no floor, no stack and no list g of arrays for which* Check(a, g) *returns with a numeric goal a.*

Proof. Suppose there is such a run, and choose one with the least number of calls, over all floors, stacks and contexts. By Theorem 11.32 the tree a° is numeric, and no numeric scalar is equal to an array, so the run does not return in step (1). It reads a packet, and Check(m, g) returns. If m were numeric, this call would be a smaller run of the excluded kind; so m is not numeric, and since its opening is defined, m is an array.

If the replacement test of step (3) succeeds, then $O_k(t; u, 0) = a$ is numeric, so t is numeric by Theorem 11.32, and $O_k(t; s, 0) = O_k(t; u, 0)$ by the same lemma. The goal is numeric after step (3) in every case.

If m is nonempty, h is its threefold repetition, an array. If m is empty and u is true-valued, h is a binary array. In the remaining case step (4iii) requires the goal to be an array of arity one, which a numeric scalar is not, and the run rejects. So h is an array, its opening in step (6) is an array, and the final call has a numeric goal $O_k(a; [k], 0)$ and a context of arrays. It is a smaller run of the excluded kind, a contradiction. □

Lemma 11.34 (Contexts of arrays)**.** *In a run of the neutral-major protocol from* Check$(a_0, [[\,]])$*, every call has a context all of whose members are arrays, and every major m whose call returns is an array.*

Proof. The initial context consists of the array $[\,]$. Suppose a call with a context of arrays reaches step (3). The major call receives the same context. If it returns, m is an array by Theorem 11.33. The hypothesis h is then the threefold repetition of a nonempty array, or a binary array, or, in branch (iii), the opened matrix $a = O_k(B; t, 2)$. In the last case m is the empty array, so step (6) evaluates the arity of a, which is undefined for a scalar; if the final call is reached, $h = a$ is an array. The opening of an array is an array, and the new context is formed by appending it to a copy of the old one, so no call changes the context of another. □

Corollary 11.35 (The unguarded test)**.** *At every point of a run that is reached after the return of a major call, the test of step (3) succeeds if and only if m is a nonempty array and either $m = [s, u]$, or s is false-valued and $m = [t, t, t]$.*

Proof. By Theorem 11.34 m is an array. An array is not equal to the Boolean `False`, and the empty array is equal neither to $[s, u]$ nor to $[t, t, t]$. □

The corollary states that, on the states that can occur, the test has the value of the same test guarded by the condition that m is true-valued, although the two differ on arbitrary values. Two mutations show that the hypotheses matter. If the initial context is [0], the numeric zero is a proved major, and a certificate of $S0 = 0$ is accepted. If the comparison in (N4) uses the field s in place of u, an introduction with $s = 1$ obtains the hypothesis $B(t, z)$ for a matrix of odd arity, that is, progress for a compound matrix; the field s is unrestricted in an introduction, whereas u is false-valued there and no false-valued value is numerically equal to 1. Both mutants are constructed and observed by the packages.

11.5.3 Correspondence

Theorem 11.36 (Neutral-major protocol)**.** *If the checker of the neutral-major protocol accepts a goal, the reading of the goal is a theorem of $\mathcal{A}$. If a closed formula is a theorem of $\mathcal{A}$, its canonical encoding is accepted on some stack.*

Proof. Soundness. The statement of Theorem 11.19 is proved for contexts whose members are arrays closed at the current floor, by the same induction. By Theorem 11.35 the elimination case applies verbatim when m is a nonempty array. If m is the empty array, its reading is the theorem $0 = 0$, the goal is not replaced, and the packet is a computation or an introduction. In an introduction u is false-valued, and by (N4) $\iota = k$ exactly when the arity of the opened matrix is even and u is numerically zero; so the hypothesis $B(\tau, z)$ is granted only to a matrix of even arity, and the case analysis on the arity in the proof of Theorem 11.19 applies. In a computation u is true-valued, and (N4) can give $\iota = k$ when u is numerically 1; Theorem 11.18 holds for either value of ι, so Theorem 11.2 applies as before. The result is a derivation of the reading of the goal from $0 = 0$, and Theorem 11.3 removes this assumption.

Completeness. By Theorem 11.17 take a derivation in $\mathcal{A}_{\mathrm{r}}$ and repeat the induction of Theorem 11.21 with contexts that contain, in addition, the empty array. The packets are those of the base protocol with [ ] in the major field of computations and introductions and with the mode in the field u. A canonical goal is a nonempty array and is never equal to [ ], so the additional member of the context closes no goal of the derivation; the major call of a packet without substantive major returns at once and consumes nothing. □

Remark 11.37 (A second sufficient argument). One may read every numeric scalar at a formula position as the contradiction $0 = S0$ instead of $0 = 0$. Theorem 11.11 remains true, since such a scalar is retained or replaced by the scalar 0. Under this reading a returning call with a numeric major would prove the contradiction, and the conclusion of its parent would follow by explosion whatever the test of step (3) does. This gives soundness without Theorem 11.33. The invariant is the stronger statement: numeric majors do not occur, so the reading of scalars need not be changed.

Remark 11.38 (Sources). `uniform_premise` proves Theorems 11.33 and 11.34 as one simultaneous induction on successful runs (every goal and every appended hypothesis is an array), for the profiles with an exact test for arrays, and states Theorem 11.37. `monus_and_neutral_major` proves them by the minimal counterexample used above and treats, in addition, the profile with attempted traversal: there the empty string and the empty object open to the empty array, an object whose only key is the empty string opens to [[ ]], a true-valued object cannot be a major because its threefold repetition is undefined, and a nonempty string has no finite successful opening. The statements above then hold with “array” read as “array or one of these empty aliases”. The two packages report different programs (Section 11.11); their protocols coincide.

11.5.4 Composition with local floors

Definition 11.39. The *local neutral-major protocol* is the neutral-major protocol with local floors (Theorem 11.26).

Theorem 11.40 (Composed checker)**.** *Theorems 11.33 to 11.35 and Theorem 11.36 hold for the local neutral-major protocol.*

Proof. The proofs of the three invariants do not refer to the floor except through Theorem 11.32, which holds at every floor; the minimal counterexample is chosen over all floors. The proof of Theorem 11.36 combines with Theorem 11.27. The context of a caller is not changed by a major call, because every extension forms a new list. □

The programs of `neutral_scope` write step (4), as modified by (N3), as one Boolean chain: the threefold repetition of m, or else the computation equation if u is true-valued, or else the introduction. By Theorem 11.34 the threefold repetition is true-valued exactly for a nonempty array m, and the computation equation is a binary array, hence true-valued; so the

chain takes the branch that (N3) prescribes, and the introduction is reached exactly when m and u are false-valued. The missing image is spelled `0j`, a value that the JSON reader cannot produce and that fails every comparison. A further profile of the package replaces the final call by a loop over an immutable context; this transformation is treated once, in Theorem 11.51. The package constructs two mutants: the one with initial context [0] described above, and one in which the loop extends a shared mutable list, which accepts a certificate of $0 = S0$.

11.6 Fixing an elimination before its major is proved

This section follows `prevalidated_elimination` and the checker part of `scope_and_closure`. The order of two steps of the base protocol is exchanged: the template of an elimination is opened, and the goal replaced, before the major premise is proved. The proof of the major can then be folded into the expression that forms the hypothesis.

11.6.1 The global-floor version

Definition 11.41 (Prevalidated protocol)**.** The prevalidated protocol is the base protocol of Theorem 11.15 with step (3) replaced by the following two steps.

(3′) If $m = [s, u]$, or if s is false-valued and $m = [t, t, t]$, and if moreover $O_k(t; u, 0) = a$ and $O_k(t; s, 0)$ is true-valued, replace a by $O_k(t; s, 0)$.

(3″) If m is true-valued, run Check(m, g).

The program evaluates the test of (3′) as membership of m in the pair consisting of $[s, u]$ and of the value that is $[t, t, t]$ if s is false-valued and the complex number `1j` otherwise, without a test that m is true-valued; and it forms the hypothesis of branch (4i) as `c(m,g) or m*3`. One of the three delivered profiles replaces the goal also when $O_k(t; s, 0)$ is false-valued; Theorem 11.43 covers both forms, and canonical minor premises are nonempty arrays.

Lemma 11.42 (Two properties of the program text)**.** *(a) No value produced by the JSON reader is equal to `1j`. Hence the membership test succeeds exactly under the condition of (3′), and then m is a nonempty array.*

(b) Every returning call of the checking procedure returns the null value. Hence `c(m,g) or m*3` proves the major and then has the value of the threefold repetition of m.

Proof. (a) The values of the reader are arrays, strings, objects, the null value, Booleans and real numbers, including its nonfinite numbers; none is equal to a complex number with nonzero imaginary part. If the test succeeds, m is $[s, u]$ or $[t, t, t]$, a nonempty array. (b) A call returns in step (1) with no value or after its final call with no value; this is an induction on the run. A failure inside the major call propagates and the hypothesis is not formed. □

Writing the Boolean `False` for the excluded alternative in (a) is unsound here: it is equal to the numeric 0, the marker of progress, so a packet with $m = 0$ and a true-valued s would pass the test and, having a false-valued major, would prove nothing. The package exhibits a false closed equation accepted after this replacement. Likewise, spelling the missing image as the integer 0 instead of a value that fails comparisons fills unbound holes and accepts a goal that contains an unallocated name.

Lemma 11.43 (Frozen elimination)**.** *In step (3′), let the goal a and the context be closed at floor k, and suppose that the goal is replaced by $a_1 = O_k(t; s, 0)$. If the major call of step (3″) returns, leaving a floor $k' \le k$, then a_1 is closed at floor k' with the same reading as at floor k, and the reading of a is derivable in $\mathcal{A}$ from the readings of m and a_1.*

Proof. By Theorem 11.11(a) a_1 is closed at floor k: every name in it was admitted at floor k, and so were the names of the inserted image. By Theorem 11.14 its reading is the same at floor k'. The major call returns, and its step (1) validates m at floor k, so m is closed at floor k. If $m = [s, u]$, the two openings of t at the one floor k read $F(v)$ and $F(\sigma)$ for one template F, and the reading of m is $\sigma = v$; Tr applies. If s is false-valued and $m = [t, t, t]$, the openings read $B(v)$ and $B(0)$ by Theorem 11.12, the reading of m is $Q_y B(y)$, and QE applies. □

Theorem 11.44 (Prevalidated protocol)**.** *The checker of the prevalidated protocol is sound and complete for $\mathcal{A}$ in the sense of Theorems 11.19 and 11.21.*

Proof. For soundness, the induction of Theorem 11.19 is repeated; its elimination case uses Theorem 11.43 in place of the argument about openings at the floor k', and the major is proved in step $(3'')$ by the induction hypothesis, by Theorem 11.42. A replacement in step $(3')$ is followed by step $(3'')$, since m is then true-valued; if the major call fails, the run rejects.

For completeness, the stacks of Theorem 11.21 are accepted without change. In a canonical packet the designated variable of a template is an exposed slot scalar, which is a hole at every floor, and all other names are admitted at floor k; so the openings of step $(3')$ at floor k are the trees that the base protocol computes at floor k'. Canonical minor premises are nonempty arrays, so the condition that $O_k(t; s, 0)$ is true-valued removes no canonical inference. □

Proposition 11.45 (Raw acceptance is not preserved)**.** *There is a stack that the prevalidated checker accepts and the base checker rejects.*

Proof. Let $o = [0]$, $r = [0, o, 0]$, $M = [r, o]$ and $c = [0, 0, o, 0]$, and consider the stack $[\,c,\ c,\ [M, r, o, [-1, o]],\ [o, o]\,]$ with goal $[o, o]$, which reads $S0 = S0$. The term r reads $I(0, S0;\, v.\, 0)$, and the packet c supplies the equation $r = S0$. In the prevalidated order the template $[-1, o]$ is opened at floor 0, where the scalar -1 is a hole: the opening at $u = o$ is the goal, and the goal becomes $[r, o]$. The major $M = [r, o]$ is then proved by one packet c, and the continuation proves the new goal $[r, o]$ by the other. In the base order the major is proved first, which lowers the floor to -1; the scalar -1 of the template is then an admitted name, the opening at u is $[-1, o]$, the test fails, the goal remains $[o, o]$, and the remaining packet does not prove it. □

The package states that the two sets of accepted stacks are incomparable and supplies a stack accepted only in the base order; that stack is not reproduced in its report (Section 11.12). The two checkers agree on canonical certificates and present the same consequence relation, by Theorem 11.44.

11.6.2 The local-floor version

With local floors the exchange is an identity of accepted inputs. This is the checker result of `scope_and_closure`, which applies it to the local neutral-major programs of Theorem 11.39 and to a loop profile of the same protocol.

Theorem 11.46 (Equality of accepted certificates)**.** *Consider a checker with local floors in which the major call is a separate statement before the replacement test, and the checker obtained from it by evaluating the replacement test first and placing the major call at the head of the Boolean chain that forms the hypothesis. The two checkers accept the same finite inputs.*

Proof. The replacement test reads the packet, the current goal and the opener at the caller's floor. It reads no packet from the stack and changes neither the context nor the floor; its only effect is the replacement of the caller's goal. The major call receives the same arguments m,

g and k in both programs, has its own floor and its own binding of the context, does not refer to the caller's goal, and extends contexts only by forming new lists or tuples. Its one shared effect is the consumption of packets, which the replacement test does not touch. Hence the two operations commute on every state. By Theorem 11.42(b) the major call contributes a false value to the chain, so the hypothesis is formed from the same remaining operands. An induction on the returning run, applied to the major calls and the continuations, shows that each program returns exactly when the other does, with the same goal, floor and remaining stack. A failure of either operation prevents acceptance in both orders; only the point of rejection can differ. □

Theorem 11.45 does not contradict Theorem 11.46: with a global floor the major call lowers the floor at which the test is evaluated, and with local floors it does not. The decrement of step (5) is not moved in either version; the witness of an introduction is opened before the fresh name is allocated.

11.7 Scalar neutral premises and loops

This section covers the checker parts of `kernel_refinement` and `compact_update`, the compact profiles of `validated_core`, and the compact changes of `checked_refinement` and `deterministic_introduction`. The programs are checkers of the guarded-source calculus $\mathcal{U}$ of Theorem 11.5. Their protocol is taken from `guarded_source`; the scalar spelling of the neutral premise is taken from `reversal_core`. `kernel_refinement` and `compact_update` apply that spelling to the same programs independently and report the same sizes; both describe the result as a composition.

11.7.1 The guarded-source protocol

Definition 11.47 (Guarded-source protocol)**.** The guarded-source protocol has local floors and the following steps for Check(a, g, k); steps (1), (2) and (5) are those of Theorem 11.15.

(3) Let M° be $[s, u]$ if s is true-valued and $[t, t, t]$ otherwise. If $m = M^\circ$, $O_k(t; u, 0) = a$ and $O_k(t; s, 0)$ is true-valued, replace a by $O_k(t; s, 0)$.

(4) Run Check(m, g, k). Form h as in (N3) of Theorem 11.31.

(6) Let $\iota = k$ if m is true-valued or the arity of a is 2, and zero otherwise. Run the final call as in Theorem 11.15.

The initial context is $[0]$, and a packet without substantive major has $m = 0$. The *native protocol* is the same with (N4) of Theorem 11.31 in place of step (6).

Step (3) implements SE: the source field selects the shape of the major, an equation for a true-valued source and a closure for a false-valued one. Step (6) implements the single introduction of $\mathcal{U}$: its mode is read off the root of the matrix, and a packet carries no mode. The native protocol keeps the free choice of mode and is a checker of $\mathcal{A}^-$; it is the proof kernel of the validated checkers of Section 11.8.

Proposition 11.48 (Scalar neutral premise)**.** *In the guarded-source protocol let every member of g be closed at floor k. If* Check(a, g, k) *returns, then a is closed at floor k and the auxiliary reading of a is derivable in T_{eq} from the auxiliary readings of the members of g. Conversely, every closed theorem of $\mathcal{U}$ has a certificate on which its canonical encoding is accepted. The same holds for the native protocol with the ordinary reading and $\mathcal{A}^-$.*

Proof. Soundness. A scalar at the formula root reads $0 = 0$ (Theorem 11.8), so the initial context is a theorem and a scalar goal is trivially derivable; no invariant on the roots of goals is needed. The induction is that of Theorem 11.19, carried out in T_{eq}. The tree M° is an array in every case. Hence the test of step (3) succeeds only for a nonempty array m that is $[s, u]$ with s true-valued, or $[t, t, t]$ with s false-valued; a scalar major never passes it. In the first case the major reads $\sigma = \upsilon$ and the replacement is a transport of equals, valid in T_{eq}. In the second the major reads $Q_y B(y)$, the goal $B(\upsilon)$ is replaced by $B(0)$, and the reading $B(0) \to \forall y\, B(y)$ of the closure justifies it. The openings are evaluated at the caller's floor, which the major call does not change.

By the induction hypothesis the reading of m is derivable. If $m = [B]$, the exported hypothesis reads $Q_y B(b, y)$ with b fresh. For a compound B the auxiliary reading of $D_{x,y}B$ is $\exists x$ of the reading of $Q_y B(x, y)$, and existential elimination applies. For a literal equation B it is $\exists x \forall y\, (B(x, y) \to B(x, Sy))$, from which $B(b, 0) \to \forall y\, B(b, y)$ follows by induction for the equation $B(b, y)$. If m is an array of arity at least 2, or a true-valued numeric scalar, the exported hypothesis reads $0 = 0$. If the threefold repetition of m is false-valued, the packet is a computation, treated by Theorems 11.2 and 11.18 in T_{eq}, or an introduction. In an introduction the hypothesis is $B(\tau, z)$ exactly when the opened matrix has arity 2. For a literal equation the premise gives $\forall y\, (B(\tau, y) \to B(\tau, Sy))$, the auxiliary reading with witness τ. For a matrix of arity 1 or 3 the premise gives $B(\tau, Sz)$ from $B(\tau, 0)$ for fresh z, hence $B(\tau, 0) \to \forall y\, B(\tau, y)$ by coverage. For other arities the matrix reads $0 = 0$ and the conclusion is valid outright. In the native protocol the ordinary reading is used throughout and the case analysis is that of Theorem 11.19.

Completeness. The induction of Theorem 11.21 is repeated with local floors (Theorem 11.27) and the packets

$$[[s, u], s, u, F], \qquad [[B, B, B], 0, u, B], \qquad [[B], 0, 0, 0], \qquad [0, s, u, t], \qquad [0, 0, 0, w]$$

for SE at a source other than the literal 0, SE at the literal 0, DE, a computation and the introduction; in the native protocol the introduction by cases has $u = [\,]$. The canonical encoding of a source other than the literal 0 is true-valued, so M° is the intended major in both cases of SE, and for DE the unary major differs from $M^\circ = [0, 0, 0]$. A neutral major 0 is closed by the initial member of the context without a packet. Computations with a literal zero initializer or predecessor are first replaced as in Theorem 11.17; the transport used there has the source C, which is not the literal 0, so the replacement is available in $\mathcal{U}$ and $\mathcal{A}^-$. A canonical goal is a nonempty array and is not closed by the scalar member of the context. □

By Theorem 11.6, for a canonical public formula A the checker of the guarded-source protocol accepts some stack with goal A if and only if A is a theorem of the public theory. The checker also accepts goals that are not canonical public formulas, for example scalars and auxiliary formulas; the compact programs assume that the goal offered to them is canonical, and this assumption is their input contract (Section 11.8).

Remark 11.49 (Comparison with the empty-array premise). The predecessor programs have the initial context $[[\,]]$ and the neutral major $[\,]$. The scalar spelling is shorter by one byte and one syntax-tree node. It does not need the invariant of Theorem 11.33: in the test of step (3) the candidate M° is always an array, whereas the test of Theorem 11.31 has a Boolean candidate that a numeric zero matches. The two spellings do not accept the same raw stacks, since a stack with the old neutral major can be rejected. The translation of certificates replaces the major field of a packet when that field is exactly the empty array and changes no other occurrence of an empty array; in particular the mode field of an introduction in the native protocol is kept.

Remark 11.50 (Substitution in the restricted calculus). In the soundness proof the fresh names of a generated computation equation are specialized to zero by Theorem 11.2 applied in

T_{eq}, after the continuation has been interpreted there. Applying a substitution theorem to derivations of $\mathcal{U}$ would be incorrect, as noted in Theorem 11.6.

11.7.2 Loops over immutable contexts

Theorem 11.51 (Tail call and loop)**.** *In any of the protocols of this chapter, replace the final call of step (6) by a return to step (1) with the new goal and the new context, and represent contexts by immutable tuples that are extended by forming a new tuple. The resulting checker has the same returning runs, with the same consumption of packets and the same floors, as the recursive checker.*

Proof. Relate a list and a tuple when they have the same members in the same order; membership tests agree on related contexts. A major call receives related contexts, the same stack and the same floor, and by induction on its run it returns after consuming the same packets and leaving the same floor. The recursive checker then calls itself on the opened goal and the extended context; the loop continues with exactly these objects at the same floor. The extension forms a new tuple and rebinds a local name, so the context held by the caller of a major call is unchanged. The two openings of step (6) are evaluated in the opposite order in the two programs, but they are pure at the fixed new floor. When step (1) closes the goal, the loop ends and the recursive checker returns through its pending frames, which do nothing further. Unrolling a finite run of the loop gives a run of the recursive checker, and conversely. □

The loop can check longer continuations than the recursion under a fixed recursion limit; the theorem compares ideal executions. The loop profile with the fewest tokens, in `kernel_refinement` and `validated_core`, is not a program of Theorem 11.47: it keeps a global floor, an empty initial context and the optional major of Theorem 11.41, with the test of step (3) and the mode of step (6) of Theorem 11.47. Its soundness is that of Theorem 11.44, read in T_{eq} as in Theorem 11.48. Certificates of the global and of the local protocols are not interchangeable, since they allocate names differently. Immutability is necessary. If the context is a list extended in place, a hypothesis that is local to the proof of a major premise remains in the context of the caller. `kernel_refinement` gives the stack

```
[[[],0,[],[0]],[[[0,[0]]],0,0,0],[0,[0]]]
```

with goal $0 = S0$: the major $D_{x,y}(0 = S0)$ is introduced by cases, the case assumption $0 = S0$ is appended during its proof, and with a shared list it closes the goal of the caller. The checker with tuples rejects this stack.

11.7.3 The root test is a logical boundary

Step (6) of Theorem 11.47 tests both the major and the arity of the matrix. A shorter test that inspects only the arity of the generated hypothesis grants the hypothesis $B(t, z)$ also to matrices of arity 3, that is, to matrices that are closures. `kernel_refinement` shows that this changes the theory.

Proposition 11.52 (Progress at closures gives full induction)**.** *Add to $\mathcal{A}$ the rule that infers $D_{x,y}B$ from $\Gamma, B(t, z) \vdash B(t, Sz)$, z fresh, for matrices B whose root is Q. Then induction is derivable for every native formula.*

Proof. Let $P(n)$ be a native formula. The logical reconstruction of Section 1.2 provides a gate $G_v A$ with $\vdash G_0 A$ and $G_{Sv} A \dashv\vdash A$. Put $G(n) := Q_z G_z P(n)$. Then $G(n) \dashv\vdash P(n)$: from $G(n)$ and the theorem $G_0 P(n)$, QE gives $G_{S0} P(n)$ and hence $P(n)$; conversely $P(n)$ gives $G_{Sz} P(n)$,

so DI-case with an unused witness gives $D_{u,z}G_zP(n)$, and DE exports the closure $Q_zG_zP(n)$, in which the fresh parameter does not occur. A derivation of $P(n) \vdash P(Sn)$ therefore yields $G(n) \vdash G(Sn)$. The added rule, with a matrix $B(x,y) \equiv G(y)$ that does not contain x, gives $D_{x,y}G(y)$; DE gives $Q_yG(y)$, and QE with the base $G(0)$ gives $G(t)$. Translating back, $P(0)$ and $P(n) \vdash P(Sn)$ give $P(t)$. □

The strengthening reaches public formulas through the translation of the guarded language, so a restriction of the goal grammar does not repair it. The same observation applies to an arbitrary quantified matrix in the progress rule (`checked_refinement`).

11.7.4 Local rewritings of the program text

Three reported changes are identities of the host language and involve no invariant of runs.

In the compact guarded checker, `checked_refinement` writes M° as `s and [s,u] or [t]*3`. If s is false-valued both spellings give $[t,t,t]$; if s is true-valued, $[s,u]$ is a nonempty array, hence true-valued, and both give $[s,u]$. The values agree for all JSON values. The change saves one byte and adds three syntax-tree nodes.

In a compact guarded checker with the decrement written inside the final call, `deterministic_introductio` writes $k-1$ as `~-k`. On integers the complement satisfies $\sim x = -x-1$, so $\sim(-k) = k-1$, and every floor of a run is an integer, being obtained from 0 by decrements. The change saves two bytes and adds one node.

Remark 11.53 (The floor is not a free function of the context). With local floors and initial context [0], every call satisfies $k = 1-|g|$. `checked_refinement` and `validated_presentation` record that this does not shorten the program. The opener depends on the current floor, and the floor is decreased in step (5) before the two openings of step (6), which must admit the new name. A variant that fixes the floor from the length of the context at entry and drops the decrement accepts simple computation certificates and rejects valid certificates with induction under a retained hypothesis or with transport below a binder. A correct variant that recomputes the floor at entry and keeps the decrement is 8 bytes longer than the program it was meant to shorten (529 bytes against 521; an invalid variant has 514).

11.8 Validated presentations

This section covers `validated_core`, `validated_presentation`, `scope_kernel` (its checker part), `checked_refinement`, and the separate implementation refinements reported in `deterministic_introd` `one_body` and `uniform_recursion`. The compact checkers accept goals that are not canonical encodings. A validated checker adds a test of its input, and the results of this section say that the test can share work with the proof kernel. Every result compares a program with a control that satisfies the same input contract.

11.8.1 Input contracts and the public grammar

Definition 11.54 (Public grammar)**.** For a raw tree a and a depth $e \geq 0$ the predicate $P(a,e)$ holds if

- a is an exact integer, e is odd, and $a = 0$ or a is odd with $3 \leq a < e$; or
- a is an array of arity r, where $r \in \{1,\ 2+(e \bmod 2)\}$ or $a = [a_0, a_0, a_0]$, and $P(b, e+5-r)$ holds for every child b.

The *shape predicate* $S(a,e)$ is defined in the same way with the condition $3 \leq a < e$ weakened to $3 \leq a$. The predicates P^{ed} and S^{ed} omit the alternative $a = [a_0, a_0, a_0]$. A raw tree is a

canonical closed formula of the language with equality, Q and D if $P(a, 0)$ holds, and of the language with equality and D if $P^{\mathrm{ed}}(a, 0)$ holds. It is a canonical closed *guarded* formula if $P^{\mathrm{ed}}(a, 0)$ holds and, in every unary array $[B]$ at a formula position whose child B is binary, each of the two children of B is a ternary array whose first two entries are 5 and 0.

At an even depth the admitted arities are 1 and 2, for D and equality, together with the diagonal triple for Q; at an odd depth they are 1 and 3, for S and the iterator, where the diagonal alternative adds nothing. The depth law of Section 11.2 makes one arity table serve both sorts, since the parity of the depth is the sort. A leaf is the scalar 0 or the odd scalar of a slot in scope. In the guarded condition the entries 5 and 0 are the encodings of the index y, below the dummy slot of the iterator, and of the initializer 0, so the condition is the guard (11.1). The language with equality and D expresses the language with Q: a closure $Q_y B$ is interderivable with $D_{x,y} B$ for x not in B, by DE in one direction and by DI-case and QE in the other; closures remain available as auxiliary formulas in both cases.

Definition 11.55 (Input contracts)**.** A checker satisfies

- the *raw contract* if it tests nothing about its goal beyond what the proof kernel computes; its claims concern goals that are canonical encodings;
- the *goal contract* for a grammar if it accepts only inputs whose goal is a canonical closed formula of that grammar;
- the *typed contract* for a grammar if it accepts only inputs that are nonempty arrays, all of whose nodes, including those of unused packets, are exact integers or arrays, and whose goal is a canonical closed formula of that grammar.

Sizes of programs under different contracts, or under one contract for different grammars, are not comparable; Section 11.11 lists the contract of every program. Let $J(a)$ state that every node of a is an exact integer or an array. Booleans are not exact integers.

11.8.2 Scope is checked by the first opening

Lemma 11.56 (The first opening supplies the upper bounds)**.** *For every raw tree a with $J(a)$ and every depth e,*

$$P(a, e) \iff S(a, e) \text{ and } O_0(a; \bot, e) \text{ is defined,}$$

and if this holds then $O_0(a; \bot, e) = a$. The same is true of P^{ed} and S^{ed}.

Proof. By induction on a. For an integer at an odd depth, S requires $a = 0$ or a odd with $a \geq 3$; the opening at floor 0 is defined exactly if $0 \leq a < e$, which under S is the missing bound $a < e$, and then the scalar is retained. At an even depth S and P fail for every integer. For an array, S and P impose the same condition on the arity, the children are opened at the depths $e + 5 - r$ at which the predicates are evaluated, and the induction hypothesis applies; the opening rebuilds the array from unchanged children. □

A kernel run starts with $O_0(a_0; \bot, 0)$ at floor 0, in step (1), before any packet is read and before the membership test. If a leaf violates its upper bound, the opener consults the missing image; in these programs the missing image is the checking function itself, to which the array test of the opener cannot be applied, and the run rejects. Theorem 11.56 therefore lets a validator omit every scope bound and keep only the shape.

11.8.3 Typed checkers

The typed checkers consist of the native protocol of Theorem 11.47 as proof kernel, preceded by a validation of the whole input. The four programs considered are the control of `validated_core`, an incoming checker with separate validators for raw types and for the goal, and three successive refinements.

Lemma 11.57 (Persistent type invariant)**.** *If the input satisfies J, every raw tree that the kernel handles afterwards satisfies J, with the exception of the missing image. On trees satisfying J the tests* `a*0 == []` *and* `a*0 != 0` *both hold exactly for arrays.*

Proof. The kernel constructs trees by forming arrays, repeating an array, decreasing the integer floor, and opening; each preserves J, by induction on the run. The product of an array with 0 is the empty array and that of an integer is 0. □

Theorem 11.58 (One validator with a raw mode)**.** *Let $V(a, d)$ be defined for integers a by: $d = 0$, or d is odd and ($a = 0$ or a is odd with $2 < a < d - 2$); and for other values by:* `a*0 == []`*, and ($d = 0$ or the arity is in $\{1, 2 + (d \bmod 2)\}$ or $a = [a_0, a_0, a_0]$), and $V(b, d')$ for all children b, where $d' = 0$ if $d = 0$ and $d' = d + 5 - r$ otherwise. Then, for every JSON value a,*

$$V(a, 0) \iff J(a), \qquad V(a, \delta + 2) \iff J(a) \text{ and } P(a, \delta) \quad (\delta \geq 0).$$

Consequently the program that removes the goal a_0 from the input q, requires $V(q, 0)$ and $V(a_0, 2)$, and then runs the kernel satisfies the typed contract and accepts the same inputs as its control.

Proof. Here V holds when the evaluation returns a true value; a failed operation counts as false. In raw mode, exact integers pass; among the other values of the JSON reader only arrays satisfy `a*0 == []`, the others giving a different product or a failure; the arity condition is vacuous and the children stay in raw mode. Induction gives $V(a, 0) \iff J(a)$. In scoped mode with $d = \delta + 2$, an integer passes exactly at odd δ with $a = 0$ or a odd and $2 < a < \delta$, which is the integer clause of $P(a, \delta)$; an array passes the arity condition of P, and its children are evaluated at $d + 5 - r = (\delta + 5 - r) + 2$, so the induction hypothesis applies. On an empty array the diagonal alternative fails by an undefined access, which is a rejection.

For the program: an input that is not an array has no last element to remove, and an empty array has none either. For a nonempty array the two requirements say that the prefix and the goal satisfy J and that $P(a_0, 0)$ holds. The control tests the same three conditions with two separate recursive functions and then runs the same kernel, up to the spelling of the array test, which agrees on J by Theorem 11.57. □

The control has 1049 bytes and the program of Theorem 11.58 has 897. Of the difference, 117 bytes are obtained by sharing the arity table between the sorts and by dropping a repeated type test, with two separate guards; the remaining 35 come from merging the two input tasks into the one predicate V and one diagnostic.

Theorem 11.59 (Typing by the printed form)**.** *Let Σ consist of the ten digits, the two square brackets, the comma, the space and the minus sign. For a value x of the JSON reader, $J(x)$ holds if and only if the printed representation of x in the host language contains only characters of Σ. Consequently the program that requires this of the whole input, removes the goal a_0, requires $v(a_0, 0)$, and runs the unchanged kernel accepts the same inputs as the program of Theorem 11.58. Here $v(a, d)$ requires, for a value a whose product with 0 is 0, that d is odd and that $a = 0$ or a is odd with $2 < a < d$; and for any other value a, of arity r, that $r \in \{1, 2 + (d \bmod 2)\}$ or $a = [a_0, a_0, a_0]$, and that $v(b, d + 5 - r)$ holds for all children b.*

Proof. An exact integer prints as decimal digits with an optional minus sign, and an array prints as the representations of its elements separated by a comma and a space, within brackets; so $J(x)$ implies the condition, by induction. For the converse, each other kind of node contributes a character outside Σ that enclosing arrays preserve: a string a quotation mark, an object a brace, the Booleans and the null value letters, a finite floating-point number a decimal point or an exponent marker, a nonfinite one letters. The program tests the condition by removing characters of Σ from both ends of the printed form and requiring that nothing remains, which holds exactly when no other character occurs anywhere. The test is applied to the whole input before the goal is removed, so an unused packet cannot hide a value of another type.

On values satisfying J the product with 0 is 0 exactly for integers, so v dispatches correctly, and $v(a, d)$ is $V(a, d{+}2)$ with the raw mode removed: the bound $a < (d{+}2){-}2$ is $a < d$, parities agree, and the children are evaluated at depths that differ by 2. Hence $v(a, 0) \iff P(a, 0)$ on such values. The goal, the stack and the kernel are then the same in both programs. □

The statement is about values of the standard reader, not about arbitrary objects of the host language. The two programs differ at resource limits: the recursive raw validator fails on an unused packet nested 600 levels deep that the printed-form test accepts. A variant that filters the input text before parsing, with the same characters and the JSON whitespace, has a direct lexical proof and is larger (885 bytes against 870).

Theorem 11.60 (Scope reuse and direct splitting)**.** *In the program of Theorem 11.58, replace the condition $2 < a < d - 2$ of V by $2 < a$, and replace the removal of the last element by a destructuring of the input into a prefix and a last element. The resulting program accepts the same inputs.*

Proof. The changed predicate tests J and S in place of J and P. A goal accepted by the old validator is accepted by the new one and leads to the same first state of the kernel. A goal accepted only by the new validator satisfies S and not P; by Theorem 11.56 its first opening is undefined, and the run rejects before a packet is read; no hypothesis can intervene, because the whole goal is traversed before the membership test. All later operations are unchanged.

The destructuring produces, for a nonempty array, the same prefix and last element. An empty array and a value that cannot be iterated are rejected. A nonempty string or object is destructured into pieces whose last element is a string, which fails the type test of the goal validator. So no input that was rejected is accepted. □

The three refinements have 897, 870 and 881 bytes; the second and the third are alternatives, each measured against the first.

11.8.4 Goal-validated checkers

Theorem 11.61 (Guarded goal validation)**.** *Let $W(a)$ state that every node of a is an exact integer or an exact array, that $S^{\mathrm{ed}}(a, 0)$ holds, and that for every unary array $[B]$ at a formula position with B binary, both children b of B satisfy* `b[:2] == [5,0]`*. A goal a is a canonical closed guarded formula if and only if $W(a)$ holds and $O_0(a; \bot, 0)$ is defined. Hence the checker that tests W and runs the compact guarded kernel accepts the same inputs as the checker with a complete separate validator of guarded formulas.*

Proof. On a child of a binary formula array that has passed the shape test, the prefix test has exactly the successes of the explicit guard test: such a child is an integer, for which the slice is undefined, or a unary array, whose prefix has length one, or a ternary array, for which equality of the prefix with $[5, 0]$ states the two coordinate conditions; its third child has been validated as a term. With Theorem 11.56 this gives the stated equivalence, and the opening

returns the goal unchanged. For the programs: the reader, the removal of the goal, the kernel and the verdict are the same. If the complete validator accepts, so does W, and the kernel starts in the same state. If W holds and the complete validator fails, the first opening is undefined and the run rejects before a packet is read. □

The complete validator carries the number of protected slots through the recursion and tests the iterator shape explicitly; the program with W has 921 bytes against 1201.

For the native calculus with public goals of equality and D, the goal contract is satisfied by a kernel preceded by a validator of S^{ed}, by Theorem 11.56. Two controls of this form occur, of 803 and of 780 bytes; they differ in the arrangement of the kernel (in the second the replacement test is evaluated inside the final call). The reported refinements are the following.

Theorem 11.62 (Grammar checked during the first opening)**.** *Give the opener a flag v. For $v = 0$ it is the opener of Theorem 11.10. For $v = 1$ an array is traversed only if its arity lies in $\{1, 2 + (d \bmod 2)\}$; a scalar is retained only if, in addition to $k \le a < d$, it is an exact integer, d is odd, and $a = 0$ or a is odd with $a > 1$; and in every other case the missing image is consulted, also at an even depth. Let the checker call the kernel on the goal with $v = 1$, and let the kernel set $v = 0$ after the membership test of step (1) of that call; every other call has $v = 0$. This checker accepts the same inputs as the control that validates S^{ed} by a separate function and then runs the kernel.*

Proof. By induction on the goal, the opening with $v = 1$ at floor 0 and depth e is defined exactly if $P^{\mathrm{ed}}(a, e)$ holds, and then returns a. For an integer the conditions are those of P^{ed}: at an odd depth, $a = 0$ or a odd with $1 < a < e$. An integer that fails them, or any scalar at an even depth, leads to the missing image, and the opening is undefined; the ordinary opener would return the scalar 0 at an even depth, which is why the flag forces the failure there. An array of a forbidden arity is not traversed and is handed to the scalar comparison, which is undefined for an array. An array of an admitted arity is traversed at the depths of P^{ed}. By Theorem 11.56 the control accepts the goal exactly under the same condition and starts its kernel in the same state. A goal with $P^{\mathrm{ed}}(a, 0)$ is a nonempty array; whether or not it is a member of the initial context, both programs agree on step (1). After step (1) of the first call the flag is 0 in every call, the opener is the ordinary one, and all packets, majors, contexts and floors coincide. An induction on the consumed stack gives the same verdict and, on acceptance, the same unused prefix. □

The parity of the depth takes the place of the sort argument of a separate validator, as in Theorem 11.54. `deterministic_introduction` realizes Theorem 11.62 against the control of 780 bytes with 656 bytes; `uniform_recursion` realizes it against the same control twice, with an explicit grammar guard (685 bytes) and with the guard merged into the cases of the opener (661 bytes), and notes that malformed goals are rejected at once and not through exhaustion of the recursion limit. Both packages state that the fusion of grammar and scope traversals is not new with them and claim only the measured refinement.

Two further changes are rewritings of a validator of S^{ed} that preserve its successes. `checked_refinement` replaces an early return on integers followed by a return on arrays by one conditional expression with the same two branches (803 to 788 bytes). `one_body` takes the result as control and changes, under the exact-integer branch, $a = 0$ to the negation of a and $a \ge 3$ to $a > 2$; replaces the Boolean sort flag by the numbers 0 and 1, of which only the truth value and the sum with 2 are used; and replaces the exact array test, after integers have been excluded, by `a*0 == []`, which holds exactly for arrays among the values of the reader and fails or is undefined otherwise (788 to 778 bytes). A variant writes M° as in Section 11.7 in addition (777 bytes, with two more syntax-tree nodes than the control). Failures may occur at a different operation than in the control; the accepted inputs are the same.

11.8.5 A rejected saving

The opener returns the scalar 0 for an out-of-range scalar at an even depth and consults the image only at an odd depth. `validated_core` records that removing this distinction of parities gives a shorter program that accepts the canonical guarded formula

$$D_{x,y}(I(y,0;\,u.\,0) = I(y,0;\,u.\,S0)),$$

which is false: its instance at $y = 0$ holds, and its instance at $y = S0$ is $0 = S0$. The accepted certificate uses a scalar template, which without the parity test is opened both as a formula and as a term. The parity test is the separation of the two sorts and cannot be removed.

11.9 Inferred majors, eager selection and the phase of validation

This section covers `inferred_major` and `smaller_envelope`, which overlap, `frontier_review`, `checked_presentation`, `context_phase`, the two checkers of `progress_independence`, the compact checker of `crossed_uniqueness`, and the audit of sources in `orbit_closure`. All programs are checkers of $\mathcal{U}$ or $\mathcal{A}^{-}$ in the protocols of Theorem 11.47; no rule is changed. Each device is compared with a control, and for each device the set of accepted stacks is unchanged, or shrinks, or changes in a way that is described exactly. All statements are about stacks whose nodes are integers and arrays.

11.9.1 The controls

The compact controls of this section are programs of the guarded-source protocol of Theorem 11.47 with two differences, both taken from packages outside this chapter. First, step (6) reads: $\iota = k$ if m is true-valued or the arity of a is even. This *parity selector* agrees with the test "the arity of a is 2" at the arities 1, 2 and 3; at the arities 0 and at least 4 the matrix reads $0 = 0$, and the proof of Theorem 11.48 applies without change. Second, there are two arrangements of step (3). In the control of 516 bytes the replacement test precedes step (4), as in Theorem 11.47. In the control of 506 bytes it follows step (4): it is evaluated at the floor k, with the goal as step (4) leaves it, immediately before the decrement of step (5). The control of 506 bytes is the compact program of `deterministic_introduction` in the table of Section 11.11.

Lemma 11.63 (Delayed replacement)**.** *In a checker with local floors, the two arrangements of the replacement test of step (3) have the same returning runs.*

Proof. Let m be true-valued. If the major call returns, the threefold repetition of m is true-valued, so step (4) takes branch (i) and does not change the goal. The test reads the packet, the goal and the opener at the caller's floor; the major call changes none of them, and the opener is pure. So the test has the same value and the same effect before and after step (4). Let m be false-valued. The candidate M° is a nonempty array and is not equal to m, so the test fails in both positions, although branch (iii) of step (4) may have replaced the goal in between; the opening $O_k(t; u, 0)$ that the test evaluates does not depend on the goal. An opening that is undefined in one arrangement is evaluated, and undefined, in the other. □

11.9.2 The inferred major

`inferred_major` and `smaller_envelope` deliver the same two programs for this device, a loop profile with a global floor and a loop profile with local floors and the scalar neutral premise; their controls are the two loop profiles of Section 11.7. In these controls step (3)

tests $m = M^\circ$ and $O_k(t; u, 0) = a$. The first field of an elimination packet is thus determined by its other fields.

Definition 11.64 (Inferred-major protocol)**.** The inferred-major protocol is obtained from its control by replacing step (3) with: if $O_k(t; u, 0) = a$, put $m := M^\circ$ and replace a by $O_k(t; s, 0)$. All later steps use the new value of m; in particular the major call of step (4) is $\mathrm{Check}(M^\circ, g, k)$. On a canonical certificate the first field of an elimination by SE is arbitrary, for instance 0; the first field of a DE packet is the major $[B]$ as before, and its template field is 0.

The major is reconstructed, not omitted: its proof is checked as before, in the unchanged context. The device differs from the constructed major of Theorem 11.26, which is triggered by a true-valued first field; here the trigger is the agreement of the target instance with the goal.

Theorem 11.65 (Repair)**.** *Let the inferred-major checker accept a stack. In each consumed packet at which the test $O_k(t; u, 0) = a$ succeeded, replace the first field by M°, and leave all other fields and packets unchanged. The control accepts the resulting stack, with the same goals, contexts, floors and consumption of packets. Hence the soundness statement of Theorem 11.48 holds for the inferred-major protocol with local floors, and the statement made after Theorem 11.51 for the profile with a global floor.*

Proof. At a packet where the test succeeded, the control finds $m = M^\circ$ and $O_k(t; u, 0) = a$ and replaces the goal by $O_k(t; s, 0)$, as the inferred run does. At a packet where the test failed, the conjunction tested by the control fails, whatever m is, and the first field is unchanged. In both cases the two runs continue with the same major, the same goal and the same floor; the reconstruction of M° is pure and precedes the major call, so with a global floor the floor seen by the major call is the same too. The major call consumes the repaired packets of its own subrun, by induction on the run, and the hypothesis, the decrement and the final call coincide. The repair is determined by the accepted run. □

Lemma 11.66 (Separation of terms and goals)**.** *Let t be the canonical encoding of a term below any number of slots. Then $O_k(t; u, 0)$ is defined for every u, without the image being consulted, and contains no binary array. Every canonical encoding of a formula contains a binary array. Hence $O_k(t; u, 0)$ is not a canonical formula.*

Proof. The arrays of t have arity 1 or 3, so from the depth 0 every node of t is traversed at an even depth, where a scalar is retained or replaced by 0 and the image is not consulted. The traversal preserves arities. A canonical formula ends in an equation, which is a binary array. □

Theorem 11.67 (Completeness with inferred majors)**.** *Every closed theorem of $\mathcal{U}$ has a certificate on which the inferred-major checker accepts its canonical encoding.*

Proof. Take the certificate of Theorem 11.48, with the template field of every DE packet equal to 0. At a packet of SE the test succeeds and M° is the major of the inference, so the run is that of the control. At a computation packet the field t is the encoding of the update, and at an introduction packet that of the witness; by Theorem 11.66 the test is defined and fails, since every goal of the run is a canonical formula. At a DE packet $O_k(0; u, 0)$ is the scalar 0, which is not a canonical formula. So no packet is taken for an elimination that is not one, and the run is that of the control. A neutral major 0 is closed by the initial context in the local profile and is not checked in the global profile. □

Proposition 11.68 (Change of the accepted stacks)**.** *Compare the inferred-major checker with its control on one stack.*

(a) *If at every consumed packet $O_k(t; u, 0)$ is defined, and is equal to the goal only when $m = M^\circ$, the two runs coincide. This holds for the certificates of Theorem 11.67.*

(b) *A stack accepted by the inferred-major checker and rejected by the control becomes a stack accepted by the control under the repair of Theorem 11.65. Such stacks exist: both packages replace the major field of eliminations by* 0 *in certificates of the control and obtain stacks that the inferred-major checker accepts and the control rejects.*

(c) *A stack accepted by the control can be rejected by the inferred-major checker only if, at some consumed packet with $m \neq M^\circ$, the opening $O_k(t; u, 0)$ is undefined or is equal to the goal.*

Proof. (a) Under the hypothesis the two tests have the same value at every packet, and where they succeed the assignment $m := M^\circ$ changes nothing; an induction on the run gives the claim. The control evaluates $O_k(t; u, 0)$ only when $m = M^\circ$ in some profiles; the hypothesis makes the additional evaluation harmless. Statement (c) is the contrapositive of (a), and (b) is Theorem 11.65 together with the cited certificates. □

In case (c) with equality, the control treats the packet as a DE, a computation or an introduction with the supplied m, while the inferred-major checker replaces the major and the goal; the packages exhibit no such stack and do not claim equality of the accepted sets. The set of accepted canonical goals is the same, by Theorems 11.65 and 11.67. The two programs improve the token and the node coordinate of their controls by three each and are five bytes longer.

11.9.3 Eager selection of the goal

`frontier_review` changes the form of the replacement in the control of 516 bytes; `checked_presentation`, `progress_independence` and `crossed_uniqueness` apply the same change to the control of 506 bytes. Write C for the condition $m = M^\circ$ and $O_k(t; u, 0) = a$, which these controls evaluate as one comparison of pairs, and R for $O_k(t; s, 0)$. The control takes the new goal to be R if C holds and R is true-valued, and a otherwise, and evaluates R only if C holds. The refinement evaluates R in every case and takes the new goal to be R if C holds and a otherwise; in the program this is the indexing of the pair (a, R) by the truth value of C.

Theorem 11.69 (Eager selection)**.** *(a) Every stack accepted by the refined checker is accepted by its control, with the same goals, contexts, floors and consumption of packets.*

(b) *The converse fails: there is a stack accepted by the control and rejected by the refined checker.*

(c) *Every closed theorem of $\mathcal{U}$ has a certificate accepted by the refined checker.*

Proof. (a) Whenever R is defined, the two forms select the same goal. This is clear if C fails, and if C holds and R is true-valued. Let C hold and R be false-valued. By Theorem 11.32 an opening of an array at the depth 0 is an array of the same arity, so a false-valued R arises only from the empty array t or from a numeric t; in both cases the opening does not depend on the image, so $R = O_k(t; u, 0) = a$. The opener is pure, so on an accepted run of the refined checker every operation of the control is available with the same value, and an induction on the run gives (a).

(b) Insert, before a certificate of a computation, the packet $[0, 3, [0], [5, 0]]$. It has $m = 0$, so C fails, and the control processes it as a computation packet that adds a hypothesis. The refined checker evaluates $R = O_k([5, 0]; 3, 0)$: the scalar 5 is exposed at the depth 3, the image 3 is validated at the depth 3, where it is out of range, and the run rejects.

(c) If $O_k(s; \bot, 3)$ is defined, then $O_k(t; s, d)$ is defined for every tree t of integers and arrays and every depth d: by induction on t, an array leads to finitely many calls on its children, and a scalar is retained, replaced by 0, or replaced by the validated image, which starts no further substitution. In the certificate of Theorem 11.48 choose the inert fields as follows: $s = u = t = 0$ in a DE packet and $s = 0$ in an introduction packet. Then in every packet s is 0 or the canonical encoding of a term, $O_k(s; \bot, 3)$ is defined, and so is R. The run of the refined checker is then the run of the control. □

Since C is evaluated as a comparison of pairs, $O_k(t; u, 0)$ is evaluated at every packet already in these controls; the refinement adds only the evaluation of R. The accepted canonical goals are unchanged by (a) and (c). The refinement saves three bytes in both controls. In the arrangement of Theorem 11.63 with the test after step (4), all openings of the test are evaluated at the floor k, before the decrement that the same expression performs.

11.9.4 The remainder selector

`smaller_envelope` replaces the parity selector of step (6) by the remainder of 3 modulo the arity of a: $\iota = k$ if m is true-valued or this remainder is not 0.

Proposition 11.70 (Remainder selector)**.** *The checker with the remainder selector satisfies Theorem 11.48: it is sound for the auxiliary reading and accepts a certificate of every closed theorem of $\mathcal{U}$. It rejects every stack on which step (6) is reached with a false-valued m and a goal of arity* 0*. Equality of the accepted stacks with the control is not claimed.*

Proof. At the arities 1, 2 and 3 both selectors have the values 0, 1 and 0, so every packet whose matrix or goal is a meaningful formula is treated as in the control; in particular a matrix that is a closure receives the zero instance, as Theorem 11.52 requires. At the arity 0 the remainder is undefined and the run rejects. At an arity of at least 4 the remainder is 3 and $\iota = k$; the parity selector gives $\iota = k$ at even and zero at odd arities. In an introduction such a matrix reads $0 = 0$, and the conclusion is valid for either hypothesis; in a computation Theorem 11.18 holds for either value of ι. So the soundness induction is unchanged. Canonical certificates contain only matrices and goals of arity 1, 2 or 3 at step (6), so the completeness part is unchanged. □

At an arity of at least 4 the two selectors can retain different raw hypotheses with the same reading $0 = 0$, and a later membership test can then differ; this is why only soundness and the equality of the accepted canonical goals are stated. The selector saves one byte. `checked_presentation` examines a further selector, which inspects the arity of the generated hypothesis, and rejects it for the reason given in Theorem 11.52; it also notes that in the native protocol the two false-valued modes 0 and [] of the field u must remain distinct, since identifying them removes the introduction by cases at equations.

The program of 502 bytes, which `checked_presentation`, `progress_independence` and `crossed_uniqueness` deliver with identical text, is the control of 506 bytes with eager selection and the remainder selector. By Theorems 11.69 and 11.70 it satisfies Theorem 11.48. Its accepted stacks are included in those of its control as long as no formula of arity 0 or at least 4 reaches step (6); without this restriction neither inclusion is claimed.

11.9.5 Truth-valued results and the tail of a packet

Two changes of `inferred_major` and `progress_independence` preserve every run.

Proposition 11.71 (Truth-valued result)**.** *Let the checking procedure of the control of* 506 *bytes return the truth value of the membership test of step (1) if it is true, and otherwise a*

one-element list whose element is the pair formed by the hypothesis of step (4) and the result of the final call; and let branch (i) of step (4) be written as the conjunction of the major call and the threefold repetition of m. The resulting checker has the same returning runs, with the same consumption of packets and the same floors, as the control.

Proof. By induction on the run, every returning call returns a true value: the value of a successful membership test, or a nonempty list. The control forms the hypothesis as the disjunction of the result of the major call, which is the null value, and the threefold repetition; the new form is the conjunction of a true value and the threefold repetition. Both have the value of the threefold repetition, and if that is false-valued both continue with the same later alternatives. The list has one element, obtained from one removal of a packet, and its pair is evaluated from left to right, so the hypothesis is formed before the goal is selected, the floor decreased and the final call made, in the order of the control. A failure in either component propagates. □

The new results retain data that the control discards, so the two programs need not behave alike under a fixed bound on memory. The program has 503 bytes with the complement spelling of the decrement (Section 11.7) and 505 with the ordinary one.

In the global loop profile with inferred majors, `progress_independence` destructures a packet into m and the list r of the remaining fields, then r into s, u and t, and uses r for the image $[s, u, t]$ of step (4ii) and its first two entries for the major $[s, u]$. The second destructuring succeeds exactly if r has three entries, the list r is not modified, and a packet of another arity is rejected by one of the two destructurings as it is by the control. All runs are preserved. The change saves two tokens and costs three nodes and six bytes.

11.9.6 The phase of validation

The control is the native goal-validated checker of Theorem 11.62 with 656 bytes, whose opener has the flag v.

Proposition 11.72 (Flag as an order)**.** *In the checker of Theorem 11.62 write each guard "$v = 0$ or p" as $v \le p$, and the disjunction of the parity of the depth with v as their bitwise disjunction. The accepted inputs and the unused prefixes are unchanged.*

Proof. The flag is 0 or 1, and so is the parity of the depth, which starts at 0 and changes by integers. Each guarded condition p has the value 0, `False` or `True`: the arity condition is a Boolean and is evaluated only for arrays, and the leaf condition is evaluated only after the range test has succeeded and begins with an exact-integer test. For such values $v \le p$ holds exactly if $v = 0$ or p is true, and the bitwise disjunction of two bits is their disjunction. The new form evaluates p also when $v = 0$; by the preceding remarks this evaluation is defined wherever the control reaches the guard. □

`context_phase` removes the flag. The context is an immutable tuple and is initially empty. The opener applies the grammar conditions of Theorem 11.62 exactly when the context is empty. After the membership test of step (1), every call that does not return appends the scalar 0 to its context, by forming a new tuple.

Theorem 11.73 (Phase by context)**.** *The checker of `context_phase` accepts the same inputs as the control of Theorem 11.62, with the same consumption of packets and the same unused prefix.*

Proof. In the first call the context is empty and the control has $v = 1$ and the context [0]. The guards of the two openers have the same truth values, so the first opening is defined in one program exactly if it is in the other, and returns the goal unchanged. A goal that passes

is a nonempty array; it is a member neither of the empty context nor of [0]. Both programs therefore continue; the control sets $v = 0$, and the new checker appends 0. The opener of the first call refers to the rebound context and from then on applies no grammar condition; every later call receives a nonempty context. So there is one validating traversal in each program.

For the remaining run, relate a call of the control and a call of the new checker if their floors and stacks agree, their goals agree up to the replacement of the integer 0 by the Boolean `False` at formula positions, and the tuple has the members of the list up to the same replacement, together with additional copies of 0. At an even depth the new opener can return `False` where the control returns 0; the two are equal, have the same truth value, and behave alike under every operation of the kernel, and the exact-integer test that separates them is applied only in the first opening. Related calls agree on the membership test, since membership does not depend on multiplicity. Otherwise both remove the same packet, their major calls are related, they select the same branch and form related hypotheses, and they decrease the floor at the same point. The additional 0 is a formula that the context already contains. The final calls are related, and an induction on the run in both directions gives the claim. □

The floor of this checker is not a function of the length of the context, because of the additional copies of 0; the observation after Theorem 11.27 does not apply to it, and the argument would fail for a variant that computed the floor from the context. The shortest variant tests exact integers by the identity of a leaf with its conversion to an integer, which is correct for the values of the standard reader of the measured runtime and is not a statement about other objects; a variant with the explicit type test is three bytes longer. A further variant extends a mutable list by the scalar 0 only. Since 0 is already a member after the first step, a major call that appends to the list of its caller adds no formula; actual hypotheses are still added by forming a new list. This variant is correct for one invocation of the program and is not offered as a reusable procedure, because its default context is no longer empty after the first call.

The checker of Theorem 11.72 and the checker of Theorem 11.73 have the same three sizes and are different programs.

11.9.7 A typed native checker and typed guarded checkers

`checked_presentation` composes devices of Section 11.8 and of the present section into a checker with the typed contract for the grammar with equality and D. The driver applies the printed-form test of Theorem 11.59 to the whole input and, if it fails, has no procedure with which to remove the goal. The kernel is that of Theorem 11.62, with the replacement test after step (4) and with eager selection. Because every node of the input is an exact integer or an array, the exact-integer test in the flagged opener is omitted and arrays are recognized by a product with 0 that differs from 0 (Theorem 11.57).

Theorem 11.74 (Typed native checker)**.** *This checker satisfies the typed contract for the grammar with equality and D. Every input that it accepts is accepted by the typed checker of Theorem 11.59 for that grammar, with the same run, and every closed theorem of $\mathcal{A}^{-}$ in that grammar has a certificate that it accepts.*

Proof. By Theorem 11.59 the driver reaches the kernel exactly for nonempty arrays whose nodes satisfy J; an input that is not an array has no last element to remove. By Theorems 11.56 and 11.62 the first opening is defined exactly if the goal satisfies P^{ed} at the depth 0, which is the condition that the control tests by a separate predicate; the omitted exact-integer test is implied by J for nodes of the input, and values generated by the kernel reach the opener only after the flag is 0. By Theorem 11.63 the position of the replacement test does not change the returning runs. By Theorem 11.69(a) every accepted run is a run of the

checker without eager selection, and by part (c) of that theorem, in the form for the native protocol, canonical certificates are accepted. The decrement is written with the complement, which is the same integer. □

The inclusion is proper in general: eager selection can reject a stack with an ill-scoped inert field. The program has 673 bytes against 849 for its control. The program of 656 bytes from which its kernel is taken satisfies the goal contract only and is not its control.

Four programs add the typed contract to a compact guarded kernel; they test J on the whole input, by a recursive predicate or by the printed form, test the guarded goal grammar as in Theorem 11.61, and run the kernel. Their correctness is that of the kernel together with Theorem 11.59 or the definition of J, and Theorem 11.61. `frontier_review` wraps its kernel of 513 bytes (1183 bytes against 1186 for the same wrapper around the control). `checked_presentation` and `crossed_uniqueness` wrap the kernel of 502 bytes with a recursive validator (1172 bytes; against 1176 for the wrapper around the control of 506 bytes in the first package and against the 1183 bytes of `frontier_review` in the second). `progress_independence` wraps the same kernel with the printed-form test (869 bytes, without a stated control). `smaller_envelope` supplies a readable program of this contract (2473 bytes) as an interface for audit.

11.9.8 The audit of sources

The audit in `orbit_closure` measures every Python source of the ninety package trees at one pinned revision: 5825 paths with 1380 distinct contents, read and normalized as data and not executed. Its normalization removes docstrings at every level of nesting; this correction of an earlier pass changes 173 measurements and none of the minima. Among the complete compact checkers of the guarded calculus it finds the minima 256 tokens, 276 nodes and 506 bytes, attained by three programs: the loop profile with a global floor of `kernel_refinement` (256/283/594), a loop profile of `deterministic_opening` (263/276/580), and the compact program of `deterministic_introduction` (267/303/506). All 106 programs that are smaller than one of these coordinates are classified one by one as builders, drivers, generators, measurement programs, converters or diagnostics, none of which is a complete checker. The audit thereby establishes two points that the package reports had left open: the triple 256/278/516 that `smaller_envelope` and `frontier_review` take as their baseline was not the minimum of the corpus at that revision, because the programs of 276 nodes and of 506 bytes, which lie in subdirectories of their packages, had been overlooked; and no smaller complete checker is hidden among the sources. `progress_independence` extends the audit to the 1416 Python files of three later archives with the same outcome below 506 bytes. The audit is a census of a finite corpus. It does not verify any checker, it is not a lower bound, and it keeps the compact contract apart from the validated ones.

11.10 Results on designs that were examined

The checker packages also record mathematical results obtained while other designs were examined: obstructions to two shortcuts in `certified_zero`, an analysis of eager iteration in the standard model in `prevalidated_elimination`, a criterion on closures in `compact_update`, and smaller observations in `prevalidated_elimination`. They are independent of the protocols above.

11.10.1 Projection boxes lose reflexivity

An experimental checker of `certified_zero` differs from the base protocol in its opener: the root depth is 1, the child depth of an array of arity r is $d+((r-2) \bmod 3)$, an exposed scalar is replaced by the binary array $[t', 0]$ formed from the validated image t', and S is represented by an array of arity 4. The binary array was intended to read as the identity at a term position and as the test $t = 0$ at a formula position, which removes a branch that depends on the sort. No projection equation is supplied by the computation rules.

Proposition 11.75. *Define V on raw trees by: $V([L, R])$ holds if R is numerically 0 or L is an array of arity 3; $V(a)$ holds for every a that is not a binary array. If the experimental checker accepts a goal a, then $V(a)$ holds. In particular it accepts no certificate for the equation $[\sigma, \sigma]$ with $\sigma = [0, 0, 0, 0]$, the encoding of $S0 = S0$, nor for $[p, p]$ with $p = [\tau, 0]$ a projection box that is not numerically 0.*

Proof. Every opening at a depth $d \geq 1$ preserves V. A retained scalar remains a scalar and an exposed one becomes a binary array $[t', 0]$, for which V holds. An array keeps its arity. For a binary array, the right child is numerically 0 after the opening exactly if it was before: the scalar 0 is retained because $k \leq 0 < d$, a retained nonzero scalar stays nonzero, and a replaced scalar becomes an array; the left child is a ternary array after the opening exactly if it was before, since a replaced scalar becomes a binary array. Child depths are at least 1 again.

Now let every member of the context satisfy V; then every goal of a returning call satisfies V, by induction on the run. A goal closed by a hypothesis is the opening of a tree with V. For a packet with a major, the major call is a smaller run; a replacement changes the goal from one opening of a template to another, and both satisfy V or neither does, by the first part; the exported hypothesis is a threefold repetition, of arity divisible by 3, or a scalar, and satisfies V. A supplied computation equation has a ternary array on the left and satisfies V. An introduction requires a unary goal, for which V holds whatever the premise is. The final openings preserve V. For the two equations of the statement V fails: the right side is not numerically 0, and the left side has arity 4, or 2. □

This is an obstruction to completeness, not a claim that the experiment accepts a false formula: the inserted boxes have no equations, so the intended reading as the identity is not available inside the calculus. It does not exclude a different interpretation into another fragment of raw trees, which would need its own compilation and round trips.

11.10.2 Eager initialization has a two-element model

A second shortcut applies the update already at the counter 0, so that both computation equations use the same opening:

$$F(0, a; v.h) = h[a/v], \qquad F(Sn, a; v.h) = h[F(n, a; v.h)/v]. \tag{11.2}$$

Let the terms be x, 0, St and $F(n, a; v.h)$, with v bound in h only, and let T_{e} be intuitionistic first-order logic with equality together with (11.2), coverage $\forall x\,(x = 0 \vee \exists y\, x = Sy)$, separation $0 \neq S0$, and induction for literal equations with parameters. Successor injectivity is not an axiom of T_{e}.

Theorem 11.76 (Finite model)**.** *Let $M = \{0, 1\}$ with $0 < 1$, interpret 0 by 0, S by the constant function 1, and the eager iterator by*

$$[\![F(n, a; v.h)]\!]_\rho := [\![h]\!]_{\rho[v \mapsto [\![a]\!]_\rho]}.$$

Then M is a model of T_{e}, of $\forall x\,(0 \neq Sx)$, and of induction for every first-order formula.

Proof. Every term is monotone in all its free variables. This is an induction on terms: variables are projections, 0 and S are constant, and for $F(n, a; v.h)$, if $A(\rho)$ and $H(b, \rho)$ denote the values of a and of h with $v \mapsto b$, then $\rho \leq \sigma$ gives $H(A(\rho), \rho) \leq H(A(\sigma), \sigma)$ by the induction hypotheses for a and h. The interpretation also satisfies the substitution lemma, by induction on terms with the bound variable renamed.

Every monotone map $H\colon M \to M$ is idempotent. The monotone maps are the constant 0, the identity and the constant 1.

Fix values of the parameters and let H be the resulting interpretation of $v \mapsto h$; it is monotone, hence idempotent. With α the value of the initializer, both sides of the first equation of (11.2) have the value $H(\alpha)$; the left side of the second has the value $H(\alpha)$, since the counter is ignored, and the right side $H(H(\alpha))$. The substitution lemma makes this a verification of the schemas and not only of closed instances. Coverage holds because the elements are 0 and $S0 = 1$, and $Sx = 1 \neq 0$. For induction, if $A(0)$ and $\forall x\,(A(x) \to A(Sx))$ hold, the step at $x = 0$ gives $A(1)$, and these are all elements. A classical structure is a one-world Kripke model, so all intuitionistic rules are valid in it. □

The native presentation with Q and D over the eager terms is covered as well: under the ordinary reading of Section 1.2 the rules Hyp, Tr, QE and DE are valid in M; for DI-case the retained base gives the instance $y = 0$ and the premise the instance $y = 1 = S0$; for progress the base and the step give all instances by the induction just verified.

Corollary 11.77. *There is no interpretation of* iPRA *in T_{e} whose domain is a definable subset of a finite power of the domain modulo a definable equivalence, with or without parameters. In particular T_{e} is not bi-interpretable with* iPRA.

Proof. Such an interpretation would turn M into a model of iPRA with at most 2^d elements for some d. But iPRA proves that the numerals are pairwise distinct, by successor injectivity and separation, and has no finite model. □

The result concerns the equations (11.2) with an independent initializer. It does not apply to an update that receives the counter, to the initialized iterator of Section 1.1, or to a successor-seeded iterator, and adding successor injectivity excludes the model M without providing an interpretation.

11.10.3 Positive iteration in the standard model

`prevalidated_elimination` determines the functions that the eager terms denote in the standard natural numbers, where $F(n, a; u.h)$ is the $(n+1)$-fold iterate of $b \mapsto h(b, \vec{x})$ applied to a. For $f\colon \mathbb{N}^r \to \mathbb{N}$ a coordinate j is *essential* if changing it can change the value with the other coordinates fixed, and f is *strictly increasing in j* if it is so for every choice of the other coordinates.

Theorem 11.78 (Constant, projection or growth)**.** *Every eager term $t(x_1, \ldots, x_r)$ denotes a function of exactly one of the following kinds:*

(i) a constant;

(ii) a projection x_j;

(iii) a function with a nonempty set U of essential coordinates that is strictly increasing in each coordinate of U and satisfies $t(\vec{x}) \geq x_j + 1$ for all $j \in U$.

In particular every term is nondecreasing in each coordinate. The kind and the set of essential coordinates are computed by recursion on the term, without evaluating it.

Proof. The three kinds are disjoint: a projection violates the bound of (iii) at its coordinate, and a constant has no essential coordinate. The classification is proved by induction on terms, together with monotonicity. The term 0 is constant and a variable is a projection. S maps a constant to a constant and a function of kind (ii) or (iii) to one of kind (iii) with the same essential coordinates, the bound $\geq x_j$ becoming $\geq x_j + 1$.

Consider $F(n, a; u.h)$ and apply the induction hypothesis to n, to a, and to h with its state u as a further coordinate. If h is constant, the result is that constant, since at least one update is performed. If h is the projection to u, the result is a. If h is a projection to a parameter, the result is that projection. If h is of kind (iii) and does not depend on u, the result is h.

There remains the case that h is of kind (iii) and depends on u, so that $h(b, \vec{x}) \geq b + 1$ and h is strictly increasing in b. Let U_n and U_a be the essential coordinates of n and a, let V be the essential parameter coordinates of h, and let $U = U_n \cup U_a \cup V$. The result does not depend on coordinates outside U; if U is empty it is constant. Let U be nonempty. The result is monotone: for $\rho \leq \sigma$ the initializer and the counter are ordered and the update under ρ is pointwise at most the update under σ, so the states after a common number of updates are ordered, and further updates under σ only increase the state. It is strictly increasing in each $j \in U$: an increase of an essential coordinate of a strictly increases the initial state, and the strict monotonicity of the update preserves a strict difference; an increase of an essential parameter coordinate of h strictly increases the state at the first update, which exists even for the counter 0; an increase of an essential coordinate of n adds updates, each of which raises the state by at least 1; and if a coordinate has several of these roles, its other effects are nondecreasing. Hence U is exactly the set of essential coordinates. Finally, with N and A the values of n and a, the result is at least $A + N + 1$; for $j \in U_a$ this is at least $x_j + 1$ since $A \geq x_j$, for $j \in U_n$ likewise since $N \geq x_j$, and for $j \in V$ the last application of h is at least $x_j + 1$.

The case distinction is a recursive procedure: it uses the kinds and essential sets of the parts and one test for emptiness of a union. □

Corollary 11.79. *A nonconstant eager term is unbounded in each essential coordinate. No eager term denotes a nonconstant function with finite range; in particular none denotes a zero test with values in* $\{0, 1\}$ *or the predecessor.*

Proof. A projection is unbounded, and in kind (iii) the bound $t \geq x_j + 1$ gives unboundedness. The predecessor is nonconstant and not strictly increasing from 0 to 1. □

Theorem 11.80 (No unit on a tuple domain with actual equality)**.** *Let* $\varnothing \neq D \subseteq \mathbb{N}^d$*, let* $e \in D$*, and let* $f \colon D \times D \to D$ *be given by a* d*-tuple of eager terms, possibly with numerical parameters, such that* $f(e, z) = z = f(z, e)$ *for all* $z \in D$*, with equality of tuples. Then* D *has exactly one element. No definability of* D *is assumed.*

Proof. Parameters are replaced by numerals, so Theorem 11.78 applies to each coordinate function $f_i(x_1, \ldots, x_d, y_1, \ldots, y_d)$. Let J be the set of coordinates that are not constant on D. For $i \in J$ the function f_i has an essential coordinate in each of the two argument tuples, since otherwise $f_i(e, z) = z_i$ or $f_i(z, e) = z_i$ would be constant on D; so f_i is of kind (iii). Moreover some essential coordinate j_i of the left argument lies in J: if all of them were constant on D, then $z \mapsto f_i(z, e)$ would be constant on D. The bound of (iii) and the right unit law give $z_i = f_i(z, e) \geq z_{j_i} + 1$ for all $z \in D$. If J were nonempty, the map $i \mapsto j_i$ on the finite set J would have a cycle, and adding the inequalities around a cycle of length $\ell \geq 1$ would give $z_i \geq z_i + \ell$. So J is empty, every coordinate is constant on D, and D is a singleton. □

Theorem 11.80 is not a statement that eager terms are almost unary. With

$$q(a, b) := F(a, b; u.Su) = a{+}b{+}1, \qquad d(a) := q(a, a) = 2a{+}1, \qquad \pi(a, b) := F(a, Sd(b); u.d(u)),$$

iteration of $u \mapsto 2u+1$ gives $\pi(a,b)+1 = 2^{a+1}(2b+3)$ with $2b+3$ odd, so π is an injective pairing in the standard model. Its two projections are not eager terms: a nonconstant unary eager term is strictly increasing, while $\pi(1,0) = 11 < 13 = \pi(0,2)$ and $\pi(0,1) = 9 < 11 = \pi(1,0)$ show that neither projection is monotone on the range of π.

Equality modulo a definable relation is a real exception to Theorem 11.80. With $m(a,b) := F(a,0;u.q(u,b)) = (a+1)(b+1)$ and

$$(a,b)\ E\ (c,d) :\Leftrightarrow q(a,d) = q(c,b),$$

the quotient $\mathbb{N}^2/E$ is identified with $\mathbb{Z}$ by $(a,b) \mapsto a-b$, and the tuples of eager terms

$$(a,b) \oplus (c,d) := \big(q(a,c),\, q(b,d)\big), \qquad (a,b) \otimes (c,d) := \big(q(m(a,c),m(b,d)),\, q(m(a,d),m(b,c))\big)$$

induce addition and multiplication of integers, because $(a+1)(c+1)+(b+1)(d+1)-(a+1)(d+1)-(b+1)(c+1) = (a-b)(c-d)$. These are identities of the standard model only. In the model M of Theorem 11.76 one has $q(a,b) = 1$ for all arguments, E identifies all pairs, and the quotient has one element; so the construction does not give an interpretation of iPRA in T_{e}, in accordance with Theorem 11.77.

11.10.4 Conditional and progressive closure of an equation

`compact_update` determines when the two readings of a closure that occur in this chapter agree. Let $E(y)$ be a literal equation with parameters, and write

$$B := E(0), \qquad C(E) := B \to \forall y\, E(y), \qquad P(E) := \forall y\, \big(E(y) \to E(Sy)\big).$$

The following statements are proved in T_{eq}; the only induction used is induction for the equation E.

Lemma 11.81 (Anchored progress)**.** *$P(E) \to C(E)$, and $C(E) \leftrightarrow (B \to P(E))$.*

Proof. From $P(E)$ and B, induction for E gives $\forall y\, E(y)$. From $C(E)$ and B one has $\forall y\, E(y)$, hence $P(E)$. From $B \to P(E)$ and B one has $P(E)$, and the first statement gives $\forall y\, E(y)$. □

Theorem 11.82 (Agreement criterion)**.** $\big(C(E) \leftrightarrow P(E)\big) \leftrightarrow \big(\neg B \to P(E)\big)$.

Proof. Let $C(E) \leftrightarrow P(E)$ and $\neg B$. Then $C(E)$ holds, since its antecedent is refuted, and so $P(E)$. Conversely let $\neg B \to P(E)$. By Theorem 11.81 it remains to derive $P(E)$ from $C(E)$. The equation B is decidable in T_{eq} (Theorem 1.52). If B holds, $C(E)$ gives $\forall y\, E(y)$ and hence $P(E)$; if $\neg B$ holds, the assumption gives $P(E)$. □

Decidability is used for the one equation B and not as excluded middle. In Lean the criterion is proved in T_{eq} for every quantifier-free formula in place of the equation, as `Teq.closure_iff_progress`, with `Teq.closure_of_progress` for the first statement of Theorem 11.81 and the decidability lemma `Teq.qfDec`.

A guarded equation (11.1) has a provable base and satisfies the criterion; this is the equivalence of the two readings used in Theorem 11.6. A provable base is not necessary: for $E(y) \equiv (0 = S0)$ both $C(E)$ and $P(E)$ are provable. The criterion fails for $E(y) \equiv (y = S0)$: the base is refutable, so $C(E)$ is provable, while $P(E)$ would give $SS0 = S0$ from $E(S0)$, so $\neg P(E)$ is provable. This is why an introduction by progress cannot replace the introduction by cases at unguarded equations.

Theorem 11.83 (Criterion below an existential quantifier)**.** *For an equation $E(x,y)$ let C_x and P_x be the two closures for fixed x. Then*

$$\big(\exists x\, C_x \leftrightarrow \exists x\, P_x\big) \leftrightarrow \big(\exists x\, \neg E(x,0) \to \exists x\, P_x\big).$$

Proof. By Theorem 11.81, $\exists x\, P_x \to \exists x\, C_x$ always. Let $\exists x\, C_x \to \exists x\, P_x$ and let $\neg E(a,0)$. Then C_a holds, so $\exists x\, C_x$ and hence $\exists x\, P_x$. Conversely let $\exists x\, \neg E(x,0) \to \exists x\, P_x$ and let C_a hold. Decide the equation $E(a,0)$. If it holds, C_a gives $\forall y\, E(a,y)$, hence P_a. If it fails, a witnesses the antecedent, and $\exists x\, P_x$ follows, possibly with another witness. □

The Lean form is `Teq.ex_closure_iff_progress`. Agreement for every x implies agreement of the existential statements, but not conversely: for $E(x,y) \equiv (\mathrm{sel}(x, S0, y) = S0)$ the equation holds for all y at $x = 0$, so both existential closures are provable with witness 0, while at $x = S0$ it is $y = S0$, where the two readings differ.

Proposition 11.84 (Base-retaining normal form)**.** *For $A(y) := B \wedge E(y)$ one has $C(A) \leftrightarrow P(A) \leftrightarrow C(E)$, without a decision of B. If $E(y)$ is $s(y) = t(y)$ and $\langle \cdot, \cdot \rangle$ is a provably injective pairing, $A(y)$ is equivalent to the literal equation $\langle s(0), s(y) \rangle = \langle t(0), t(y) \rangle$.*

Proof. $A(0)$ is equivalent to B, so $C(A)$ is equivalent to $B \to \forall y\, (B \wedge E(y))$, that is, to $C(E)$. From $C(E)$ and $A(y)$, the retained B gives $E(Sy)$ and hence $A(Sy)$; this is $P(A)$. From $P(A)$ and B, the formulas $A(y)$ and $E(y)$ are equivalent, so $P(A)$ is the progress step for E, and induction for E from the base B gives $\forall y\, E(y)$. The last statement is the injectivity of the pairing. □

The normal form is not a measured kernel: using it in a primitive presentation requires pairing and its injectivity before the completeness theorem that is being established, and the rule that retains both $E(0)$ and $E(z)$ is the older anchored introduction. Likewise the condition $\neg E(0) \to P(E)$ of Theorem 11.82 is a statement that needs a proof; a checker can test only a syntactic sufficient condition such as the guard.

11.10.5 Further rejected designs

`prevalidated_elimination` records three further attempts.

The remaining length of the certificate as the floor. Consuming a packet would then allocate the fresh name before the witness of an introduction is validated, and the witness could contain the name. The freshness of the witness is necessary: for $E(x,y) \equiv (\mathrm{sel}(y, 0, x) = y)$ the instance at $y = 0$ is $0 = 0$ and the instance at Sz is $x = Sz$, so $D_{x,y}E$ is false under the ordinary reading, because no x equals both $S0$ and $SS0$; yet an introduction by cases with the illegal witness $t = Sz$ has the premise $\mathrm{sel}(Sz, 0, Sz) = Sz$, which is a computation.

Zero computation by definition. Identifying $I(0, a;\, v.\, h)$ with a is legitimate, but in the encoding of Section 11.2 the initializer lies below a dummy slot. In $I(n, a;\, u.\, I(0, u;\, v.\, 0))$ the occurrence of u in the inner initializer has the scalar 5, and removing the inner iterator must lower it to 3; returning the raw initializer does not preserve scope. A normalizer that lowers indices, validates the result and keeps transport and hypothesis lookup correct modulo the conversion was larger than the dispatch it would remove.

Synthesis of the conclusion. A checker can compute its conclusion from the certificate instead of checking a goal at each call; hypotheses are then referred to by index and majors and endpoints are not repeated. Two experimental programs of this kind are larger than the compact checkers (Section 11.11): the inference needs a dispatch on hypotheses, a comparison of minor instances and of the inferred premise of an introduction with the required successor instance, and a test that an inferred conclusion contains no eigenparameter. Shorter certificates and a shorter checker are separate resources.

11.11 Checker sizes

The table lists every program whose size is reported by a package of this chapter, with the figures as the packages state them, under the measurement protocol of Section 1.10. Each

row is one complete program unless it is marked as an envelope, which combines coordinates of different programs and is not the size of any program. The second column gives the input contract of Theorem 11.55 and the calculus:

Code	Contract and calculus
R/$\mathcal{A}$	raw contract, calculus $\mathcal{A}$ of Theorem 11.1
R/$\mathcal{U}$	raw contract with a canonical guarded goal assumed, calculus $\mathcal{U}$ of Theorem 11.5
G/$\mathcal{U}$	goal contract for canonical closed guarded formulas, calculus $\mathcal{U}$
G/$\mathcal{A}^{-}$	goal contract for the grammar with equality and D, calculus $\mathcal{A}^{-}$
T/eqd	typed contract for the grammar with equality, Q and D, calculus $\mathcal{A}^{-}$
T/ed	typed contract for the grammar with equality and D, calculus $\mathcal{A}^{-}$
T/g	typed contract for canonical closed guarded formulas, calculus $\mathcal{U}$

A program is to be compared only with the control named in its own row, which has the same contract and the same calculus. A dagger marks a profile that recognizes arrays by attempted traversal and carries the qualification on nonempty strings stated after Theorem 11.10. The packages also report sizes after compression; these are informational in every package and are omitted. Controls are given as tokens / syntax-tree nodes / bytes.

Program	Code	Tokens	Nodes	Bytes	Matched control
`certified_zero`: base protocol, Section 11.3					
explicit array test	R/$\mathcal{A}$	264	308	566	envelopes below
zero-product array test	R/$\mathcal{A}$	264	310	558	envelopes below
attempted traversal†	R/$\mathcal{A}$	262	296	598	envelopes below
envelope of the supplied programs	R/$\mathcal{A}$	268	300	616	different programs
envelope of seven matched controls	R/$\mathcal{A}$	264	300	601	different programs
`uniform_zero`: local floors, Section 11.4					
control, zero-product test	R/$\mathcal{A}$	267	311	545	published 264/310/558
control, explicit test	R/$\mathcal{A}$	267	309	553	published 264/308/566
uniform zero, zero-product test	R/$\mathcal{A}$	269	314	550	267/311/545
uniform zero, explicit test	R/$\mathcal{A}$	269	312	558	267/309/553
constructed majors, zero-product test	R/$\mathcal{A}$	272	314	556	267/311/545
constructed majors, explicit test	R/$\mathcal{A}$	272	312	564	267/309/553
constructed majors, readable	R/$\mathcal{A}$	309	361	1600	none
`uniform_premise`: neutral major, global floor, Section 11.5					
explicit test, conditional replacement	R/$\mathcal{A}$	262	303	548	264/308/566
zero-product test, Boolean replacement	R/$\mathcal{A}$	263	308	537	264/310/558
`monus_and_neutral_major`: neutral major, global floor, Section 11.5					
explicit array test	R/$\mathcal{A}$	263	307	550	264/308/566
zero-product array test	R/$\mathcal{A}$	263	309	542	264/310/558
attempted traversal†	R/$\mathcal{A}$	261	295	578	262/296/598†
`neutral_scope`: neutral major, local floors, Section 11.5					
zero-product array test	R/$\mathcal{A}$	266	310	529	263/309/542
explicit array test	R/$\mathcal{A}$	266	308	537	none stated
loop, attempted traversal†	R/$\mathcal{A}$	263	290	610	263/291/642
`prevalidated_elimination`: global floor, Section 11.6					
explicit array test	R/$\mathcal{A}$	263	307	547	264/308/566

Program	Code	Tokens	Nodes	Bytes	Matched control
zero-product array test	R/$\mathcal{A}$	263	309	539	264/310/558
attempted traversal†	R/$\mathcal{A}$	261	292	576	262/296/598†
conclusion synthesis, experimental	other	330	390	951	none
synthesis, shared binder, experimental	other	334	392	777	none
`scope_and_closure`: neutral major, local floors, elimination first, Theorem 11.46					
zero-product array test	R/$\mathcal{A}$	267	309	528	266/310/529
loop, attempted traversal†	R/$\mathcal{A}$	264	288	608	263/289/613†
`kernel_refinement`, `compact_update`, `validated_core`: guarded-source protocol, Section 11.7					
loop, global floor† (not in `compact_update`)	R/$\mathcal{U}$	256	283	594	257/285/568†
loop, scalar neutral premise†	R/$\mathcal{U}$	260	278	602	261/279/603†
scalar neutral premise, zero-product test	R/$\mathcal{U}$	263	300	521	264/301/522
the same with a goal validator	G/$\mathcal{U}$	489	560	1201	490/561/1202
`checked_refinement`					
compact, M° rewritten	R/$\mathcal{U}$	263	303	520	263/300/521
guarded goal validation, Theorem 11.61	G/$\mathcal{U}$	407	459	921	489/560/1201
native goal validation, one expression	G/$\mathcal{A}^-$	358	413	788	359/414/803
`validated_core`: Theorem 11.58					
one validator with raw mode	T/eqd	401	468	897	450/514/1049
the same, smaller goal grammar	T/ed	390	454	878	424/484/1000
`validated_presentation`: Theorem 11.59					
typing by the printed form	T/eqd	383	442	870	401/468/897
the same, smaller goal grammar	T/ed	370	426	849	390/454/878
lexical filter before parsing	T/eqd	–	–	885	401/468/897
`scope_kernel`: Theorem 11.60					
scope reuse and direct splitting	T/eqd	393	460	881	401/468/897
the same, smaller goal grammar	T/ed	382	446	862	390/454/878
`deterministic_introduction`, separate refinement					
grammar in the first opening, Theorem 11.62	G/$\mathcal{A}^-$	333	379	656	360/409/780
compact, complement for the decrement	R/$\mathcal{U}$	267	303	506	267/302/508
`uniform_recursion`, separate refinement					
grammar in the first opening, explicit guard	G/$\mathcal{A}^-$	340	385	685	360/409/780
the same, merged into the cases	G/$\mathcal{A}^-$	336	381	661	360/409/780
`one_body`, separate refinement					
validator rewritten	G/$\mathcal{A}^-$	357	412	778	358/413/788
validator and M° rewritten	G/$\mathcal{A}^-$	357	415	777	358/413/788
`inferred_major` and `smaller_envelope`: inferred major, Theorem 11.64 (the same two programs in both packages)					
loop, global floor†	R/$\mathcal{U}$	253	280	599	256/283/594†
loop, local floors, scalar neutral premise†	R/$\mathcal{U}$	257	275	607	260/278/602†
`inferred_major`: Theorems 11.71 and 11.72					

Program	Code	Tokens	Nodes	Bytes	Matched control
truth-valued result, complement decrement	R/$\mathcal{U}$	278	310	503	267/303/506
truth-valued result, ordinary decrement	R/$\mathcal{U}$	278	309	505	267/303/506
flag as an order	G/$\mathcal{A}^{-}$	327	375	643	333/379/656
`smaller_envelope`: Theorem 11.70					
remainder selector	R/$\mathcal{U}$	264	304	515	265/306/516
readable typed program	T/g	580	666	2473	none
`frontier_review`: Theorem 11.69					
eager selection	R/$\mathcal{U}$	268	306	513	265/306/516
the same with typed wrapper	T/g	462	524	1183	459/524/1186
`checked_presentation`, `progress_independence`, `crossed_uniqueness`: compositions on the control of 506 bytes					
eager selection and remainder selector (all three packages)	R/$\mathcal{U}$	269	301	502	267/303/506
eager selection only (`progress_independence`, `crossed_uniqueness`)	R/$\mathcal{U}$	270	303	503	267/303/506
remainder selector only (`crossed_uniqueness`)	R/$\mathcal{U}$	266	301	505	267/303/506
typed wrapper, recursive validator (`checked_presentation`, `crossed_uniqueness`)	T/g	463	519	1172	461/521/1176 and 462/524/1183
typed wrapper, printed form (`progress_independence`)	T/g	421	471	869	none stated
`progress_independence`: inferred major with reuse of the packet tail					
loop, global floor†	R/$\mathcal{U}$	251	283	605	253/280/599†
`checked_presentation`: Theorem 11.74					
typed native checker	T/ed	344	393	673	370/426/849
`context_phase`: Theorem 11.73					
phase by context, immutable tuple	G/$\mathcal{A}^{-}$	327	375	643	333/379/656
the same, explicit type test	G/$\mathcal{A}^{-}$	327	375	646	333/379/656
mutable list, one invocation only	G/$\mathcal{A}^{-}$	325	375	641	333/379/656
Attested by the audit of `orbit_closure` and not listed above					
loop profile of `deterministic_opening`	R/$\mathcal{U}$	263	276	580	not in this chapter

Some remarks on the reading of the table.

The controls of `uniform_premise`, `monus_and_neutral_major` and `prevalidated_elimination` are the three programs of `certified_zero`; `uniform_premise` lists in addition an older program of 266 tokens, 296 nodes and 689 bytes that shares the minimum of nodes. The control of the byte profile of `neutral_scope` is the byte profile of `monus_and_neutral_major`, and its loop profile is compared with an older program outside this chapter, as is the loop profile of `scope_and_closure` (a program of `coverage_and_kernel`). The controls of the guarded-source rows are the programs of `guarded_source`. The control of 1000 bytes was constructed by `validated_core` by restricting the control of 1049 bytes to the smaller grammar. The control of 780 bytes is a program of `deterministic_opening`; the control of 803 bytes is the native goal-validated checker current at the time of `checked_refinement`, whose validator already omits the scope bounds in the manner that the report attributes to `one_context`. The two experimental synthesis programs have a certificate format of their own.

Under the raw contract no package claims a program that improves tokens, nodes and bytes at once against all programs of its contract, with one exception that is qualified: the traversal profile of `certified_zero` is below both envelopes of its package in all three coordinates. Wherever a package states a triple such as 256/278/521 or 257/278/521 for compact checkers, the three numbers belong to three programs. Under the goal contract and the typed contract single programs do improve all three coordinates of their controls; see the envelope below. In `uniform_zero` the comparison with a matched control shows that the new devices cost source: the control without them has 545 bytes, uniform zero adds 5 and constructed majors a further 6; the programs are smaller than the earlier published minimum of 558 only because the control already is. In `prevalidated_elimination` the saving of 19 bytes of the byte profile is decomposed by the package into 11 for the exchange of steps and the folding of the major call, 6 for the unguarded test, and 2 for the spelling of the missing image. The typed checkers and the compact guarded checker of 520 or 521 bytes satisfy different contracts, and `validated_presentation` and `scope_kernel` state explicitly that their figures are not to be set against it.

11.11.1 The coordinatewise envelope

The smallest value of each coordinate among the programs of the table, and of the audited corpus where the audit of Section 11.9 applies, is the following, separately for each contract. A triple in one line is the size of one program only where this is said.

Contract	Tokens	Nodes	Bytes	Attained by
R/$\mathcal{A}$	261	288	528	three programs: the traversal profiles of `monus_and_neutral_major` and `prevalidated_elimination` (261 tokens), the loop profile of `scope_and_closure` (288 nodes) and its zero-product profile (528 bytes)
R/$\mathcal{U}$	251	275	502	three programs: the global loop profile of `progress_independence` (251/283/605), the local loop profile with inferred major of `inferred_major` and `smaller_envelope` (257/275/607), and the program of `checked_presentation`, `progress_independence` and `crossed_uniqueness` (269/301/502)
G/$\mathcal{U}$	407	459	921	one program, of `checked_refinement`
T/g	421	471	869	one program, of `progress_independence`
G/$\mathcal{A}^{-}$	325	375	641	the one-invocation list variant of `context_phase`; among programs without that qualification 327/375/643, attained by one program in `inferred_major` and by another in `context_phase`
T/ed	344	393	673	one program, of `checked_presentation`
T/eqd	383	442	870	one program, of `validated_presentation`

The envelope under R/$\mathcal{U}$ has the following history. The audit of `orbit_closure` finds 256/276/506 in the corpus that precedes the packages of Section 11.9. The inferred major gives 253 tokens and 275 nodes; the truth-valued result gives 503 bytes; eager selection with the remainder selector gives 502 bytes; the reuse of the packet tail gives 251 tokens. The three attaining programs differ in more than size. The two loop profiles recognize arrays by attempted traversal and accept the changed set of stacks of Theorem 11.68; the program of 502 bytes uses the zero-product test and accepts stacks as described after Theorem 11.70. All three accept the same canonical goals, by Theorem 11.48. No program attains two of the three minima, and the packages state that no single program is below 253/275/506 in all coordinates. Smaller figures for T/eqd are reported by packages outside this chapter. Since T/g is a stronger contract than G/$\mathcal{U}$, the line for T/g also bounds the latter.

11.12 Further questions

The packages state the following questions, limits and unproved claims. Items that several packages raise are merged.

Question 11.85 (Verification of the checkers)**.** Formalize the sorted reading, the opening lemma (Theorem 11.11), the zero-instance supply (Theorems 11.2 and 11.18), the invariants of successful runs and the compilation of derivations, for the actual programs on raw input and not for a model restricted to canonical input, and connect them to the derivability relation of the Lean development. All soundness and completeness proofs of this chapter are written proofs; no package verifies a program in a proof assistant. (`certified_zero`, `uniform_zero`, `neutral_scope`, `kernel_refinement`, `compact_update`, `validated_presentation`, `checked_refinement`, `scope_kernel`.)

Question 11.86 (One program for all coordinates)**.** Is there, under a fixed contract, one complete program that is below the current minima in tokens, syntax-tree nodes and bytes simultaneously, and in particular one that recognizes arrays by an exact test? (`certified_zero`, `prevalidated_elimination`, `kernel_refinement`, `checked_refinement`.)

Question 11.87 (Lower bounds)**.** Is there a class of presentations, with a fixed implementation language, input contract, treatment of malformed input and charged dependencies, for which a lower bound on checker size can be proved? The recorded sizes are upper bounds attained by particular programs, and a census of existing programs is not a lower bound. A benchmark of this kind should report the size of certificates and the work of checking together with the size of the checker. (`uniform_zero`, `neutral_scope`, `validated_presentation`, `scope_and_closure`.)

Question 11.88 (Removing a rule and shrinking a validated checker)**.** Can a rule or a term constructor be removed, with every omitted instance recovered and both formula round trips of Theorem 1.78 preserved, in such a way that a fully validated checker under a fixed contract also becomes smaller? The packages of this chapter re-encode an unchanged basis. (`uniform_zero`, `neutral_scope`, `compact_update`, `kernel_refinement`, `validated_presentation`, `checked_refinement`.)

Question 11.89 (Constructed majors)**.** Can the construction of the major from the fields of the packet (Theorem 11.26) be obtained without the six additional bytes of source that it costs against a matched control, and without an unaccounted condition on witnesses or a changed induction rule? (`uniform_zero`.)

Question 11.90 (Definitional zero computation)**.** Is there a layout of binders in which the reduction of $I(0, a; v.\, h)$ to a needs no lowering of indices, so that the zero computation can be made definitional at a lower total cost than the dispatch it removes? (`uniform_zero`, `prevalidated_elimination`.)

Question 11.91 (Closures as a separate kind of resource)**.** Can closures be represented so that a certificate cannot forge them and a raw tree does not read as a closure in one use and as an object formula in another, at a lower total cost, so that one simpler dispatcher serves all eliminations and the introduction? Theorem 11.52 shows that the present identification forces the test on the root of the matrix. (`kernel_refinement`, `checked_refinement`.)

Question 11.92 (A checkable class for the closure criterion)**.** Is there a class of matrices that satisfies the criterion of Theorem 11.82, is cheaper to validate than the guard (11.1), and still gives full expressiveness of the public language? Can a root-determined introduction coexist with the full language with equality, Q and D under the ordinary readings, at a lower total cost than the witness-selected rule? The equation $y = S0$ excludes the direct replacement of cases by progress, not every redesign. (`compact_update`, `checked_refinement`.)

Question 11.93 (Scope carried by the certificate)**.** Would a tree-shaped certificate that allocates opaque eigenparameters when a binder is entered, so that raw input cannot name them, simplify the freshness argument at a lower total size? Can the remaining length of the certificate serve as the floor if the allocation is given a separate phase? (`compact_update`, `prevalidated_elimination`.)

Question 11.94 (Terms as atomic assertions)**.** Can an equation $s = t$ be replaced by an assertion $\delta(s, t) = 0$ for a fixed distance term δ at a lower total trusted cost, when the expansion of δ and the proofs of reflection and of the round trips are counted? (`validated_presentation`.)

Question 11.95 (Whole-certificate typing during the kernel run)**.** Theorems 11.56 and 11.62 let the first opening of the kernel check the scope and the grammar of the goal. Can the exact typing of the whole certificate, including unused packets, be shared with the kernel in the same way? (`kernel_refinement` asks for the sharing of public syntax validation with the opening traversal; for the goal this is answered by the results of Section 11.8.)

Question 11.96 (The envelope of Section 11.11.1)**.** Is there one complete program under the raw contract for $\mathcal{U}$ that is below 251 tokens, 275 nodes and 502 bytes at once, or at least one that attains two of these minima? The packages of Section 11.9 answer neither; each of their compact programs improves one coordinate of its control and worsens another. (`inferred_major`, `smaller_envelope`, `frontier_review`, `checked_presentation`, `progress_independence`.)

Question 11.97 (One verified decoder with the refinements as corollaries)**.** Can the sorted reading and the soundness induction for the compact checker be formalized once, so that the refinements of Section 11.9 (the inferred major, eager selection, the remainder selector, the phase by context) become verified corollaries, and can the acceptance relation of the programs, with its totalized decoder and its scope transitions, be linked to the Lean calculus by a verified compiler? The root-opening lemma (Theorem 11.32) is proposed as a starting point. (`smaller_envelope`, `frontier_review`, `checked_presentation`.)

Question 11.98 (Validation together with the kernel)**.** Can an invariant shared between parsing, binding and proof checking make the typing of the whole certificate and the goal grammar cheaper together with the kernel, with a soundness statement for the whole accepted input language? Theorem 11.74 still types the input by a separate test before the kernel runs. (`frontier_review`.)

Question 11.99 (The floor as part of the phase)**.** The length of the context determines the floor at the entry of a call (Theorem 11.53), but the continuation uses the next floor, and the checker of Theorem 11.73 gives up the relation altogether. Is there a design in which the floor is not a separate argument and the transition to the next floor and the immutability of contexts are still accounted for, at a smaller total size? (`frontier_review`, `context_phase`.)

Question 11.100 (Raw acceptance under inference)**.** Exhibit a stack that a control accepts and the corresponding inferred-major checker rejects (Theorem 11.68(c)), or show that none exists for one of the two profiles. For the program of 502 bytes, decide whether its accepted stacks are comparable with those of its control when formulas of arity 0 or at least 4 occur (Theorem 11.70). Neither point is settled by the packages, which claim only the equality of the accepted canonical goals.

The following items are limits and claims that the packages state without a proof in their reports, and results of these packages that belong to other chapters.

- `context_phase` contains a theorem that is not checker mathematics: the native kernel without the introduction of Q by progress does not derive $Q_y(I(y, 0;\, u.\, u) = 0)$, nor the public formula $D_{x,y}(I(y, 0;\, u.\, u) = 0)$, which the full kernel derives; the proof expands a countably saturated structure coherently with substitution in iterator bodies, after a method of Friedman. The package `progress_independence` treats the same independence question. The theorem belongs to the chapter on obstructions and is not proved here.

- `smaller_envelope` contains a finite-menu obstruction: restricting the witnesses of the introduction to 0 and parameters and the targets of closure elimination to compositions of a fixed finite collection of primitive recursive operations loses native consequences. It belongs to the chapter on obstructions as well, together with the questions that the package attaches to it (an infinite but structurally generated family of targets; a quotient or another numerical domain) and the related question of `frontier_review` whether introduction and specialization can both be automatic without an enumerator of terms.

- `checked_presentation` asks whether successor-only uniqueness of iteration gives a discriminator for actual equality, and whether the remaining question of independence of induction has a model of the binding iterator that respects substitution; `smaller_envelope` and `frontier_review` ask whether the arithmetic bridge (predecessor, formula selectors) or the term signature can be simplified together with the checker. These concern the rule basis and not the checkers.

- The correctness of the exact-integer test by identity with the conversion to an integer in `context_phase` is an argument about one runtime and its standard reader; the package says so and supplies the variant with the explicit type test.

- `prevalidated_elimination` states that the sets of stacks accepted in the base order and in the prevalidated order are incomparable. The report displays only the stack of Theorem 11.45; a stack accepted only in the base order is among its test data and is not exhibited.

- `validated_core` compares its programs with controls whose validators are not printed in the report; Theorem 11.58 assumes, as the report states, that the control tests J on the whole input and P on the goal.

- The statement that the checkers of the guarded-source protocol accept exactly the public theorems rests on the equivalence theorem of `guarded_source` (Theorem 11.6), which is a written proof outside this chapter; `kernel_refinement` and `compact_update` ask for its formalization together with the restriction on substitution.

- `neutral_scope` asks whether the self-seeded basis of another package can derive its additional equality and constructor principles, or whether a complete checker that charges them can still be smaller than the checkers of the initialized iterator; it reports no result.

- `kernel_refinement` asks whether its replacement of one occurrence in the left side of an equation helps a readable or verified checker; it does not help the compact programs.

- The experimental synthesis programs of `prevalidated_elimination` are not established as complete checkers.

- None of the packages claims a minimal program, a simplest axiomatization, or a result about the arithmetic basis. The statements of Section 11.10 about eager iteration concern the theory T_{e} and the standard model; they are not lower bounds for axiomatizations of iPRA.

Appendix A

The packages

The article merges the research reports that were delivered as separate packages. The table lists each package with the chapters that draw on it.

Package	Chapters
`adaptive_seed`	10 (Obstructions)
`bounded_initializer`	8 (Recursors with a built-in seed)
`canonical_recursors`	2 (Refinements of the eight rules), 4 (Computation, equality and normal forms), 6 (Discharge of a witness and restricted equality rules), 9 (Canonical and counter-aware recursors)
`certified_zero`	1 (The base theories and their bridge), 2 (Refinements of the eight rules), 11 (Certificates and checkers)
`checked_presentation`	11 (Certificates and checkers)
`checked_refinement`	11 (Certificates and checkers)
`checker_and_barriers`	10 (Obstructions)
`clausal_canonical`	9 (Canonical and counter-aware recursors)
`clausal_single_evaluation`	4 (Computation, equality and normal forms), 5 (Clause calculi)
`clausal_two_case`	5 (Clause calculi)
`clause_contexts`	5 (Clause calculi)
`clause_discharge`	6 (Discharge of a witness and restricted equality rules)
`clause_guards`	5 (Clause calculi)
`compact_kernel`	10 (Obstructions)
`compact_update`	11 (Certificates and checkers)
`computation_normal`	2 (Refinements of the eight rules), 4 (Computation, equality and normal forms)
`computational_equality`	4 (Computation, equality and normal forms), 5 (Clause calculi)
`conditional_copy`	7 (Introduction without a supplied witness)
`constructed_computation`	7 (Introduction without a supplied witness), 10 (Obstructions)
`context_phase`	11 (Certificates and checkers)
`counter_congruence`	2 (Refinements of the eight rules), 10 (Obstructions)
`counter_evaluation`	4 (Computation, equality and normal forms), 9 (Canonical and counter-aware recursors)
`coverage_and_kernel`	8 (Recursors with a built-in seed), 11 (Certificates and checkers)

Package	Chapters
`crossed_uniqueness`	2 (Refinements of the eight rules), 11 (Certificates and checkers)
`deterministic_auxiliary`	3 (Guards and public languages)
`deterministic_introduction`	7 (Introduction without a supplied witness), 11 (Certificates and checkers)
`deterministic_opening`	6 (Discharge of a witness and restricted equality rules), 9 (Canonical and counter-aware recursors), 11 (Certificates and checkers)
`deterministic_specialization`	6 (Discharge of a witness and restricted equality rules)
`diagonal_guard`	3 (Guards and public languages), 4 (Computation, equality and normal forms)
`direct_closure`	1 (The base theories and their bridge), 2 (Refinements of the eight rules)
`direct_closure_qprogress`	2 (Refinements of the eight rules)
`equality_from_iteration`	4 (Computation, equality and normal forms), 5 (Clause calculi)
`evaluation_discharge`	1 (The base theories and their bridge), 5 (Clause calculi)
`first_instance`	6 (Discharge of a witness and restricted equality rules)
`frontier_review`	11 (Certificates and checkers)
`full_normalization`	2 (Refinements of the eight rules), 4 (Computation, equality and normal forms), 6 (Discharge of a witness and restricted equality rules)
`guarded_core`	3 (Guards and public languages), 7 (Introduction without a supplied witness), 8 (Recursors with a built-in seed), 11 (Certificates and checkers)
`guarded_orbit`	3 (Guards and public languages)
`guarded_source`	3 (Guards and public languages), 11 (Certificates and checkers)
`inferred_major`	11 (Certificates and checkers)
`inferred_witness`	6 (Discharge of a witness and restricted equality rules), 9 (Canonical and counter-aware recursors)
`initializer_equality`	4 (Computation, equality and normal forms)
`kernel_refinement`	2 (Refinements of the eight rules), 11 (Certificates and checkers)
`local_scopes`	8 (Recursors with a built-in seed), 9 (Canonical and counter-aware recursors), 10 (Obstructions), 11 (Certificates and checkers)
`mirror_core`	3 (Guards and public languages), 5 (Clause calculi)
`mirror_guard`	3 (Guards and public languages)
`monus_and_neutral_major`	8 (Recursors with a built-in seed), 11 (Certificates and checkers)
`neutral_scope`	2 (Refinements of the eight rules), 11 (Certificates and checkers)
`newest_parameter`	1 (The base theories and their bridge), 7 (Introduction without a supplied witness)
`no_witness`	1 (The base theories and their bridge), 7 (Introduction without a supplied witness)
`one_body`	5 (Clause calculi), 11 (Certificates and checkers)

Package	Chapters
one_context	1 (The base theories and their bridge), 2 (Refinements of the eight rules), 11 (Certificates and checkers)
one_sided_transport	2 (Refinements of the eight rules)
orbit_closure	2 (Refinements of the eight rules), 11 (Certificates and checkers)
parameter_intro	7 (Introduction without a supplied witness)
parameter_substitution	6 (Discharge of a witness and restricted equality rules)
parameter_witness	1 (The base theories and their bridge), 7 (Introduction without a supplied witness)
prefix_clauses	5 (Clause calculi)
prevalidated_elimination	4 (Computation, equality and normal forms), 11 (Certificates and checkers)
progress_independence	10 (Obstructions), 11 (Certificates and checkers)
protected_core	3 (Guards and public languages)
raw_clausal	1 (The base theories and their bridge), 5 (Clause calculi)
reduced_clause_calculus	5 (Clause calculi)
reflexive_base	3 (Guards and public languages), 4 (Computation, equality and normal forms), 5 (Clause calculi)
register_graph	9 (Canonical and counter-aware recursors)
research	10 (Obstructions)
reversal_core	2 (Refinements of the eight rules), 11 (Certificates and checkers)
rootfree_clausal	5 (Clause calculi)
scalar_successor	8 (Recursors with a built-in seed)
scalar_uniqueness	2 (Refinements of the eight rules), 6 (Discharge of a witness and restricted equality rules), 10 (Obstructions)
scope_and_closure	2 (Refinements of the eight rules), 11 (Certificates and checkers)
scope_and_coverage	8 (Recursors with a built-in seed)
scope_kernel	2 (Refinements of the eight rules), 4 (Computation, equality and normal forms), 5 (Clause calculi), 8 (Recursors with a built-in seed), 10 (Obstructions), 11 (Certificates and checkers)
seed_evaluation	4 (Computation, equality and normal forms), 5 (Clause calculi)
selfseed_scalar	8 (Recursors with a built-in seed)
single_constructor	9 (Canonical and counter-aware recursors)
smaller_envelope	11 (Certificates and checkers)
source_dispatch	2 (Refinements of the eight rules), 3 (Guards and public languages), 8 (Recursors with a built-in seed), 11 (Certificates and checkers)
substitutive_discharge	6 (Discharge of a witness and restricted equality rules)
successor_seed_completion	8 (Recursors with a built-in seed)
target_closure	2 (Refinements of the eight rules)
template_free	2 (Refinements of the eight rules), 6 (Discharge of a witness and restricted equality rules)
transient_seed	8 (Recursors with a built-in seed)
uniform_binding	9 (Canonical and counter-aware recursors)

Package	Chapters
`uniform_clauses`	5 (Clause calculi), 6 (Discharge of a witness and restricted equality rules)
`uniform_predecessor`	8 (Recursors with a built-in seed), 10 (Obstructions), 11 (Certificates and checkers)
`uniform_premise`	11 (Certificates and checkers)
`uniform_progress`	2 (Refinements of the eight rules)
`uniform_raw`	1 (The base theories and their bridge), 7 (Introduction without a supplied witness)
`uniform_recursion`	7 (Introduction without a supplied witness), 9 (Canonical and counter-aware recursors), 11 (Certificates and checkers)
`uniform_step`	4 (Computation, equality and normal forms), 5 (Clause calculi), 6 (Discharge of a witness and restricted equality rules)
`uniform_zero`	4 (Computation, equality and normal forms), 11 (Certificates and checkers)
`validated_core`	11 (Certificates and checkers)
`validated_presentation`	4 (Computation, equality and normal forms), 11 (Certificates and checkers)
`variable_substitution`	6 (Discharge of a witness and restricted equality rules)
`witness_discharge`	6 (Discharge of a witness and restricted equality rules), 9 (Canonical and counter-aware recursors)
`witness_rule`	2 (Refinements of the eight rules), 4 (Computation, equality and normal forms), 5 (Clause calculi), 8 (Recursors with a built-in seed)
`zero_normal`	4 (Computation, equality and normal forms), 9 (Canonical and counter-aware recursors)