

Gonshor's Product-Birthday Conjecture

An exact formula for ordinal-support series

A partial solution, an equality classification,
and exact reproducible checks

Research draft prepared with ChatGPT
20 September 2026

Scope of the result. The conjectured inequality

$$\ell(xy) \leq \ell(x) \otimes \ell(y)$$

is proved here for all surreal normal forms with real coefficients and exponents that are negatives of ordinals, allowing arbitrary set-sized transfinite support. The proof also identifies every equality case in this ring. The conjecture for arbitrary surreal numbers is **not settled** by this article.

Abstract

Gonshor's product-birthday conjecture asks whether the canonical sign length of a surreal product is bounded by the natural product of the sign lengths of its factors. We investigate the ring $\mathcal{A} = \{\sum_{\alpha \in S} c_\alpha \omega^{-\alpha}\}$, where S is any set of ordinals and the nonzero coefficients are real. Specializing the classical reduced-sign-expansion theorem gives an exact birthday formula determined by the supremum of the support, the last coefficient when one exists, and one endpoint correction. This yields a sharp support bound for products and shows that multiplication does not increase the leading Cantor degree of the birthday inside $\mathcal{A}$. Consequently the conjecture holds throughout $\mathcal{A}$, strictly except for zero factors, factors equal to ± 1 , and pairs of ordinary integers. We also obtain a four-coefficient formula for finite-support product birthdays and describe natural set-sized local subrings. Exact rational computations check the endpoint formula on 64,401 series and the product results on 96,636 pairs. The transfinite statements are established by written proofs, not by these finite tests. Historical priority for these particular special-case results has not been established.

Contents

1	The problem, its status, and the outcome	2
2	Conventions and elementary ordinal facts	3
3	The ordinal-support ring	4
4	The classical sign rule in the all-minus case	5
5	An exact endpoint formula	6
6	Products: a support estimate and a degree gap	8
7	The product conjecture and all equality cases on the ring	9
8	A four-coefficient formula for finite supports	11
9	Examples, cancellations, and transfinite endpoints	11
10	Set-sized local subrings and sharp birthday ceilings	13
11	Computation and reproducibility	14
12	Assessment of the attempt and the remaining general problem	16
A	A compact proof audit	17
B	Source and artifact notes	17

1 The problem, its status, and the outcome

1.1 The conjecture actually being investigated

A surreal number has a canonical sign expansion of ordinal length. Write $\ell(x)$ for that length, with $\ell(0) = 0$. The question is

$$\boxed{\ell(xy) \leq \ell(x) \otimes \ell(y) \quad (x, y \in \mathbf{No}).} \quad (1)$$

Here $\otimes$ denotes the *natural*, or Hessenberg, product of ordinals. The corresponding additive bound uses natural sum: $\ell(x + y) \leq \ell(x) \oplus \ell(y)$.

The Open Problem Garden entry attributes (1) to Gonshor’s 1986 book [Gon86, Lán12]. More substantially, in an August 2024 account Philip Ehrlich identified the question as Gonshor’s conjecture and reported the monomial case proved with van den Dries: $x = r\omega^a$, $y = s\omega^b$, with arbitrary surreal a, b and real r, s . His account did not report a general solution [Ehr24]. No general resolution was located in the sources consulted for this draft on 20 September 2026. That is a statement about the literature search, not a certification that every relevant publication has been examined.

There is an elementary reason that the ordinal operations must be made explicit. With $x = \omega + 1$ and $y = 2$, their surreal product is the ordinal $\omega \cdot 2 + 2$. Its birthday is therefore $\omega \cdot 2 + 2$, whereas the *ordinary* ordinal product $(\omega + 1) \cdot 2 = \omega \cdot 2 + 1$ is smaller. The natural product has the correct value $(\omega + 1) \otimes 2 = \omega \cdot 2 + 2$.

1.2 Why recursive expression rank is not the answer

The rank of an unreduced recursive expression for a surreal number can exceed the canonical sign length of the number it represents. For example, Roughan’s calculation of the recursive product form for 2×3 gives generation 12 [Rou18]. The value of that form is the integer 6, whose canonical birthday is 6. Thus the form calculation does not contradict (1). In this article every occurrence of ℓ refers to the canonical number, never to an expression tree.

This distinction matters for both research and implementation. Expanding Conway’s recursive multiplication rule and measuring its depth does not by itself measure the birthday of its output value. Our computations instead evaluate a canonical normal-form sign-length rule.

1.3 The result and its logical dependencies

The main domain is

$$\mathcal{A} = \{0\} \cup \left\{ \sum_{\alpha \in S} c_\alpha \omega^{-\alpha} : S \subseteq \mathbf{On} \text{ is a nonempty set, } c_\alpha \in \mathbb{R} \setminus \{0\} \right\}. \quad (2)$$

The sum is a Conway normal form, ordered by increasing α . These exponents are *negative ordinals as surreal numbers*; they are not arbitrary negative surreal exponents. This ring contains $\mathbb{R}$, $\mathbb{R}[[\omega^{-1}]]$, and many series with uncountable or otherwise transfinite support. It does not contain ω , or even $\omega^{-1/2}$.

The route to the main theorem is

$$\begin{aligned} &\text{classical reduced sign expansions} \implies \text{exact endpoint formula} \\ &\implies \text{support and degree bounds} \implies (1) \text{ on } \mathcal{A}. \end{aligned}$$

The endpoint formula and the deductions from it are proved below. The substantial imported ingredient is the classical sign-expansion theorem. Accordingly, this is not a foundational reconstruction of all surreal arithmetic. It is a self-contained derivation of a special-case product theorem from an explicitly stated standard ingredient.

Research status. The general conjecture remains outside the conclusion of this draft. The results below are proposed as rigorous special-case theorems, with complete arguments relative to the classical normal-form sign theorem. They have not been peer reviewed or formally checked, and no claim of historical priority is made.

2 Conventions and elementary ordinal facts

2.1 Two different kinds of ordinal arithmetic

Ordinary ordinal addition and multiplication are written $\alpha + \beta$ and $\alpha \cdot \beta$. They describe ordered concatenation and its iteration. The operations $\oplus$ and $\otimes$ are natural sum and natural product. Surreal addition and multiplication use the usual symbols; when both operands are ordinals embedded in **No**, they agree with the natural operations [Ber20, Section 6].

In particular, throughout this paper $\omega \cdot \alpha$ in a *birthday formula* is an ordinary ordinal product. One must not replace it by $\omega \otimes \alpha$. Likewise, $1 + \rho$ in an exponent below is ordinary addition and can differ from $\rho + 1$.

If a common decreasing list of Cantor exponents is chosen, then

$$\alpha = \sum_{j=1}^m \omega^{\rho_j} a_j, \quad \beta = \sum_{j=1}^m \omega^{\rho_j} b_j, \quad \rho_1 > \cdots > \rho_m,$$

where zero coefficients are allowed for alignment, and

$$\alpha \oplus \beta = \sum_{j=1}^m \omega^{\rho_j} (a_j + b_j). \quad (3)$$

Natural product is polynomial multiplication, with exponent addition itself interpreted naturally:

$$\left(\sum_i \omega^{\xi_i} a_i \right) \otimes \left(\sum_j \omega^{\eta_j} b_j \right) = \bigoplus_{i,j} \omega^{\xi_i \oplus \eta_j} a_i b_j. \quad (4)$$

These formulas also fix the notation for arbitrarily large ordinals.

Definition 2.1. For a nonzero ordinal θ , write $\deg \theta$ for its leading Cantor exponent and $\text{lc } \theta$ for its leading positive integer coefficient. Thus

$$\theta = \omega^{\deg \theta} \cdot \text{lc } \theta + \theta', \quad \theta' < \omega^{\deg \theta}.$$

We leave $\deg 0$ undefined. In particular, a positive finite ordinal has degree zero.

Lemma 2.2. For nonzero ordinals θ, η ,

$$\deg(\theta \oplus \eta) = \max(\deg \theta, \deg \eta), \quad (5)$$

$$\deg(\theta \otimes \eta) = \deg \theta \oplus \deg \eta, \quad (6)$$

$$\deg(\omega \cdot \theta) = 1 + \deg \theta. \quad (7)$$

If $d, e > 0$, then

$$d \oplus e \geq \max(d, e) + 1. \quad (8)$$

Natural sum is strictly increasing in each argument and is therefore monotone in both arguments jointly.

Proof. Equation (5) follows by aligning Cantor normal forms. In (4), the largest exponent is the natural sum of the largest exponent from each factor, proving (6). Ordinary left multiplication by ω changes a leading term $\omega^d c$ to $\omega^{1+d} c$, proving (7).

For strict monotonicity, compare two ordinal Cantor normal forms at the greatest exponent where their coefficients differ. Adding the same coefficient list does not change that first comparison. Thus $a < a'$ implies $a \oplus b < a' \oplus b$. Taking $b > 0$ shows that $a \oplus b > a$; interchanging the arguments gives $a \oplus b > b$. For ordinals, being strictly greater than D means being at least $D + 1$, proving (8). $\square$

Lemma 2.3. *If $\delta \leq \alpha$, the interval $[\delta, \alpha)$, in its inherited order, has an order type η satisfying $\delta + \eta = \alpha$. For $\eta > 0$, deleting its first point leaves an order type η' satisfying $1 + \eta' = \eta$. Furthermore,*

$$\omega \cdot \delta + \omega \cdot \eta = \omega \cdot \alpha, \quad \sup_i \omega \cdot \alpha_i = \omega \cdot \sup_i \alpha_i$$

for an increasing nonempty family with no last index.

Proof. The ordinal α is the concatenation of its initial segment δ and the stated interval. The deletion statement is the same observation with initial segment 1. The identities follow respectively from distributivity of left multiplication over ordinary ordinal addition and from the limit clause in the definition of ordinal multiplication. $\square$

3 The ordinal-support ring

3.1 Normal forms and the meaning of the sums

We use Conway's normal-form representation: every surreal has a unique set-sized series $\sum_{i < \nu} r_i \omega^{a_i}$ with nonzero real coefficients and strictly decreasing surreal exponents. Addition and multiplication are the corresponding series operations, and $\omega^a \omega^b = \omega^{a+b}$ [Ber20, Sections 8–10]. The notation ω^a here means Conway's omega-map, not a substitution into a separately defined real exponential function.

For $x \in \mathcal{A}$, let $S(x)$ be its *ordinal-index support*:

$$S(x) = \{\alpha : c_\alpha \neq 0\}.$$

This is a set of nonnegative ordinals, not the usual support of actual exponents $\{-\alpha : \alpha \in S(x)\}$. A subset of the ordinals is automatically well ordered. Its increasing enumeration makes $-\alpha$ a decreasing sequence, so every sum in (2) is a legitimate normal form. No analytic convergence condition or boundedness condition on the real coefficients is imposed.

Proposition 3.1. *The class $\mathcal{A}$ is a commutative subring of $\mathbf{No}$. If $x = \sum c_\alpha \omega^{-\alpha}$ and $y = \sum d_\beta \omega^{-\beta}$, then*

$$xy = \sum_{\gamma} p_\gamma \omega^{-\gamma}, \quad p_\gamma = \sum_{\alpha \oplus \beta = \gamma} c_\alpha d_\beta, \quad (9)$$

with absent coefficients interpreted as zero. Each inner sum is finite.

Proof. Surreal addition on embedded ordinals is natural sum. Therefore

$$(-\alpha) + (-\beta) = -(\alpha \oplus \beta), \quad \omega^{-\alpha} \omega^{-\beta} = \omega^{-(\alpha \oplus \beta)}.$$

To check local finiteness directly, write $\gamma = \sum_{j=1}^m \omega^{\rho_j} n_j$ in Cantor normal form. A pair of ordinals with natural sum γ is obtained by splitting each integer coefficient n_j between its two components. No new Cantor exponent can appear, since the coefficients are nonnegative. Consequently there are exactly

$$\#\{(\alpha, \beta) : \alpha \oplus \beta = \gamma\} = \prod_{j=1}^m (n_j + 1) \quad (10)$$

ordered pairs. For $\gamma = 0$, the empty product is 1.

The candidate product indices form the set $\{\alpha \oplus \beta : \alpha \in S(x), \beta \in S(y)\}$. It is well ordered, and every coefficient is a finite sum. Thus the normal form product exists and belongs to $\mathcal{A}$. Addition and negation are coefficientwise and preserve membership. The real constants give the identity element. All ring laws are inherited from **No**. $\square$

In particular, $\mathcal{A}$ has no zero divisors. It is not a subfield: $\omega^{-1} \in \mathcal{A}$, whereas its inverse ω is not in $\mathcal{A}$. This failure is a genuine restriction of the result, not just a matter of terminology.

4 The classical sign rule in the all-minus case

4.1 Real coefficients

Put $\mathbb{D} = \mathbb{Z}[1/2]$. A surreal has finite birthday exactly when it is a dyadic real. A non-dyadic real has birthday ω [Ber20, Section 5]. For $0 \neq c = m/2^k \in \mathbb{D}$ in reduced form, $k \geq 0$, the finite birthday is

$$d(c) := \ell(c) = \lceil |c| \rceil + k. \quad (11)$$

For integer c , this says $d(c) = |c|$. The formula is also recorded as Lemma 1 of [Rou18]; it follows by the successive bisections in the dyadic sign tree.

For use in a sign block, define the coefficient-tail length

$$\tau(c) = \begin{cases} d(c) - 1, & c \in \mathbb{D} \setminus \{0\}, \\ \omega, & c \in \mathbb{R} \setminus \mathbb{D}. \end{cases} \quad (12)$$

Deleting the initial sign of a finite sign string shortens it by one. Deleting the initial sign of an ω -long real sign string leaves length ω . Reversing all signs, as needed for a negative coefficient, does not change any of these lengths.

4.2 Precisely the imported ingredient

We use Gonshor's reduced-sign-expansion theorem, in the formulation of Bournez and Guilmant, Definition 2.20 and Theorem 2.21 [BG22]; their theorem cites Gonshor's Theorems 5.11–5.12. Its needed specialization is as follows.

For successive normal-form exponents $-\alpha_i$, begin with their all-minus sign strings. Delete each minus position already shared with an earlier exponent. At a successor term, delete one additional minus after the preceding exponent if its coefficient is non-dyadic. Start each term with one initial plus, then append an ω -long block for each remaining minus. Append the sign string of the absolute coefficient without its initial plus. Reverse the whole term for a negative coefficient. Concatenate the term blocks in normal-form order.

Only the lengths of these blocks are needed. The specialization is particularly simple because $-\alpha$ has α minus signs and no plus signs. In the general sign theorem, plus signs in the exponent change the sizes of subsequent blocks; that complication is absent here.

4.3 A usable block-length formula

Write the support in increasing order as $S = \{\alpha_i : i < \nu\}$, with coefficient c_i , and put

$$\delta_i = \sup_{j < i} \alpha_j, \quad \sup \emptyset = 0.$$

The positions shared with earlier all-minus exponents are exactly $[0, \delta_i)$: a position ξ is shared precisely when $\xi < \alpha_j$ for some $j < i$. Therefore the remaining interval is $[\delta_i, \alpha_i)$. Let its order type be η_i .

When i is a successor and $c_{i-1} \notin \mathbb{D}$, delete the first point of this interval and call the resulting order type η'_i . The interval is nonempty in that case. In all other cases put $\eta'_i = \eta_i$. The specialized sign rule gives

$$B_i = 1 + \omega \cdot \eta'_i + \tau(c_i), \quad \ell(x) = \sum_{i < \nu} B_i, \quad (13)$$

where every operation on the right is ordinary ordinal arithmetic.

This equation is already a transfinite algorithm, but it appears to require the whole support. The next theorem compresses it to endpoint information. The point is not that earlier signs literally disappear. Rather, finite initial lengths are absorbed when later ω -blocks are appended, as in $7 + \omega = \omega$.

5 An exact endpoint formula

Theorem 5.1 (Exact birthday). *Let $0 \neq x = \sum_{\alpha \in S} c_\alpha \omega^{-\alpha} \in \mathcal{A}$ and put $\sigma = \sup S$.*

If S has no greatest element, then

$$\boxed{\ell(x) = \omega \cdot \sigma.} \quad (14)$$

If S has greatest element a and $c_a \notin \mathbb{D}$, then

$$\boxed{\ell(x) = \omega \cdot (a + 1).} \quad (15)$$

If S has greatest element a and $c_a \in \mathbb{D}$, put $n = d(c_a)$ and define $e(x) \in \{0, 1\}$ by

$$e(x) = 1 \iff \begin{cases} \sup(S \setminus \{a\}) = a, & \text{or} \\ a = b + 1 \text{ for some } b \in S \text{ with } c_b \notin \mathbb{D}. \end{cases} \quad (16)$$

Here $\sup \emptyset = 0$, so the first condition includes the constant case $a = 0$. Then

$$\boxed{\ell(x) = \omega \cdot a + (n - 1 + e(x)).} \quad (17)$$

There are two different reasons for the correction. The last exponent may be a limit of preceding support indices, leaving no new exponent signs; or it may be the immediate ordinal successor of a preceding index with a non-dyadic coefficient, causing the one new minus to be deleted. These mechanisms must not be conflated.

Proof. Enumerate S as in (13). We prove the statements for all initial normal forms by transfinite induction. For a segment with last index α_i , let P_i denote its sign length. Along the way the following bounds hold:

$$\omega \cdot \alpha_i \leq P_i \leq \omega \cdot (\alpha_i + 1). \quad (18)$$

They are immediate from the asserted formulas once each stage is proved.

The first term. If $\alpha_0 = 0$, the exponent string is empty. Equation (13) gives $1 + \tau(c_0)$, which is $d(c_0)$ for a dyadic coefficient and ω otherwise. This is precisely (17) with $e = 1$, or (15).

If $\alpha_0 > 0$, no minus is deleted and the block has length $1 + \omega \cdot \alpha_0 + \tau(c_0)$. The initial 1 is absorbed by the nonzero ω -multiple. We obtain $\omega \cdot \alpha_0 + (d(c_0) - 1)$ in the dyadic case, with $e = 0$, and $\omega \cdot (\alpha_0 + 1)$ in the non-dyadic case.

A successor term after a dyadic coefficient. Suppose the preceding index is β , the new one is $\alpha > \beta$, and the preceding coefficient is dyadic. Its initial normal form has length $\omega \cdot \beta + t$ for some finite t , by induction. Let $\eta = \text{otp}[\beta, \alpha)$, so that $\beta + \eta = \alpha$ and $\eta > 0$. No extra minus is deleted. Appending the new block gives

$$\begin{aligned} (\omega \cdot \beta + t) + (1 + \omega \cdot \eta + \tau(c_\alpha)) &= \omega \cdot \beta + \omega \cdot \eta + \tau(c_\alpha) \\ &= \omega \cdot \alpha + \tau(c_\alpha). \end{aligned}$$

The first equality uses that a finite ordinal followed by a nonzero ω -multiple is absorbed. This proves the formulas, with no endpoint correction in the dyadic last-coefficient case.

A successor term after a non-dyadic coefficient. The preceding segment has length $\omega \cdot (\beta + 1)$. Delete the first point of $[\beta, \alpha)$, leaving order type η' . Thus $1 + \eta' = \eta$ and $\beta + 1 + \eta' = \alpha$. If $\eta' > 0$, appending the new block gives

$$\omega \cdot (\beta + 1) + (1 + \omega \cdot \eta' + \tau(c_\alpha)) = \omega \cdot \alpha + \tau(c_\alpha).$$

Again the asserted formulas follow without correction.

If $\eta' = 0$, then $\alpha = \beta + 1$. There are no retained minus signs. Appending the new block gives

$$\omega \cdot \alpha + (1 + \tau(c_\alpha)).$$

For a dyadic last coefficient this is $\omega \cdot \alpha + d(c_\alpha)$, giving exactly $e = 1$. For a non-dyadic last coefficient it is $\omega \cdot (\alpha + 1)$, as required.

A term at a nonzero limit position. Suppose i is a nonzero limit ordinal and let $\delta = \sup_{j < i} \alpha_j$. The concatenation preceding term i has length $\sup_{j < i} P_j$. The bounds (18) and ordinary ordinal continuity give

$$\sup_{j < i} P_j = \omega \cdot \delta. \quad (19)$$

Indeed the preceding indices have no maximum and are cofinal in δ , so both bounding suprema are $\omega \cdot \delta$.

There is no additional successor deletion at a limit position. If $\delta < \alpha_i$, the retained interval is nonempty. The same absorption and concatenation calculation as above gives $\omega \cdot \alpha_i + \tau(c_i)$. If $\delta = \alpha_i$, the retained interval is empty, so the result is $\omega \cdot \alpha_i + (1 + \tau(c_i))$. The latter is the cofinal endpoint correction in (16).

A complete normal form with no last term. Its support has no maximum. Apply the same squeeze argument to all initial segments ending in a term. Equation (19), now with $\delta = \sigma$, gives (14). All steps preserve (18), completing the induction. $\square$

Corollary 5.2. *For $0 \neq x \in \mathcal{A}$, with $\sigma = \sup S(x)$,*

$$\omega \cdot \sigma \leq \ell(x) \leq \omega \cdot (\sigma + 1). \quad (20)$$

If $\ell(x)$ is infinite, then

$$\deg \ell(x) = 1 + \deg(\max\{1, \sigma\}). \quad (21)$$

Proof. The bounds follow term by term from Theorem 5.1. If $\sigma > 0$, the bounding ordinals have the same leading Cantor exponent, since $\deg(\sigma + 1) = \deg \sigma$. This gives $\deg \ell(x) = 1 + \deg \sigma$. If $\sigma = 0$, the number is a real constant; an infinite birthday then means a non-dyadic constant and $\ell(x) = \omega$, which also gives (21). $\square$

6 Products: a support estimate and a degree gap

Theorem 6.1 (Sharp support bound). *Let $0 \neq x, y \in \mathcal{A}$, and let $\alpha = \sup S(x)$, $\beta = \sup S(y)$. Then*

$$\ell(xy) \leq \omega \cdot ((\alpha \oplus \beta) + 1). \quad (22)$$

For every prescribed pair α, β , equality is attainable by some factors with those support suprema.

Proof. Every surviving product index is $a \oplus b$ with $a \leq \alpha$ and $b \leq \beta$. Monotonicity gives $a \oplus b \leq \alpha \oplus \beta$, so $\sup S(xy) \leq \alpha \oplus \beta$. Now apply (20). Coefficient cancellation only removes indices and cannot spoil this upper bound.

For sharpness choose $x = \frac{1}{3}\omega^{-\alpha}$ and $y = \omega^{-\beta}$. Their product is the monomial $\frac{1}{3}\omega^{-(\alpha \oplus \beta)}$, whose birthday is the right-hand side of (22) by (15). $\square$

The sharpness assertion fixes the support suprema, not the two input birthdays. These are different optimization problems. In particular it does not imply that the conjectured birthday-product bound is sharp for general non-real factors in this ring.

Proposition 6.2 (Strict degree gap). *Suppose $x, y \in \mathcal{A}$ have infinite birthdays. Put $d = \deg \ell(x)$, $e = \deg \ell(y)$, and $D = \max(d, e)$. Then*

$$\ell(xy) < \omega^{D+1} \leq \omega^{d \oplus e} \leq \ell(x) \otimes \ell(y). \quad (23)$$

In particular the conjectured inequality is strict for this pair.

Proof. The factors and their product are nonzero. Write $\alpha = \sup S(x)$, $\beta = \sup S(y)$ and set

$$r = \max(\deg \max\{1, \alpha\}, \deg \max\{1, \beta\}).$$

The ordinal $(\alpha \oplus \beta) + 1$ has leading exponent r . This is immediate by aligning Cantor forms, including the case $\alpha = \beta = 0$. Thus the upper bound in (22) has degree $1 + r$. On the other hand, (21) gives

$$D = \max(1 + \deg \max\{1, \alpha\}, 1 + \deg \max\{1, \beta\}) = 1 + r.$$

It follows that $\deg \ell(xy) \leq D$. Every ordinal of degree at most D is strictly below ω^{D+1} , proving the first inequality in (23).

Both d, e are positive because both input birthdays are infinite. Hence Lemma 2.2 gives $d \oplus e \geq D + 1$. Finally $\ell(x) \otimes \ell(y)$ has leading exponent $d \oplus e$ and positive leading coefficient, so it is at least $\omega^{d \oplus e}$. $\square$

7 The product conjecture and all equality cases on the ring

7.1 Finite factors

Lemma 7.1 (Dyadic factors). *For dyadic reals u, v , $\ell(uv) \leq \ell(u)\ell(v)$. Equality holds precisely when a factor is zero, a factor is ± 1 , or both factors are integers.*

Proof. The zero cases are immediate. For nonzero u, v , put $a = \lceil |u| \rceil$, $b = \lceil |v| \rceil$, and let k, l be the respective reduced denominator exponents. Thus $\ell(u) = a + k$ and $\ell(v) = b + l$. The product has denominator exponent at most $k + l$, and $\lceil |uv| \rceil \leq ab$. Hence

$$\ell(uv) \leq ab + k + l \leq (a + k)(b + l). \quad (24)$$

The difference in the second inequality is

$$(a + k)(b + l) - (ab + k + l) = l(a - 1) + k(b - 1) + kl.$$

If $k, l > 0$, this is positive. If $k > 0, l = 0$, it is positive unless $b = 1$, in which case the integer v is ± 1 . The remaining mixed case is symmetric. If $k = l = 0$, both numbers are integers and equality is immediate. Multiplication by ± 1 also preserves birthday, completing the classification. $\square$

Lemma 7.2 (A dyadic scalar cannot increase the leading birthday data). *Let $0 \neq q \in \mathbb{D}$ and $x \in \mathcal{A}$ have infinite birthday. The ordered pair $(\deg \ell(qx), \text{lc } \ell(qx))$ is lexicographically no larger than $(\deg \ell(x), \text{lc } \ell(x))$. Consequently, if $q \notin \{1, -1\}$, then*

$$\ell(qx) < \ell(q) \otimes \ell(x). \quad (25)$$

Proof. Multiplication by the nonzero constant q preserves the index support. If that support has no maximum, (14) gives exactly the same birthday.

Suppose it has maximum a . A dyadic coefficient remains dyadic after multiplication by q . If the last coefficient is dyadic, both birthdays have the form $\omega \cdot a + \text{finite}$. Since $\ell(x)$ is infinite, $a > 0$, and the finite tails do not change the leading Cantor data. The endpoint correction may change, but only within those finite tails.

If the last coefficient is non-dyadic, it may remain non-dyadic or become dyadic. In the former case both birthdays equal $\omega \cdot (a + 1)$. In the latter the new birthday is $\omega \cdot a + \text{finite} < \omega \cdot (a + 1)$, so its leading Cantor data cannot increase. This includes $a = 0$, when the new birthday may be finite.

Finally $m = \ell(q)$ is a positive integer, and $m \geq 2$ unless $q = \pm 1$. Natural multiplication by m preserves the leading exponent of $\ell(x)$ and multiplies its leading coefficient by m . The preceding leading-data comparison makes (25) strict. $\square$

Remark 7.3. One must not claim that a non-dyadic real coefficient remains non-dyadic under a dyadic scalar: $3 \cdot \frac{1}{3} = 1$. Nor is the entire birthday necessarily nonincreasing. For example,

$$\ell\left(\frac{1}{3} + \omega^{-1}\right) = \omega + 1, \quad \ell\left(3\left(\frac{1}{3} + \omega^{-1}\right)\right) = \omega + 2.$$

What is controlled in [Lemma 7.2](#) is the leading Cantor exponent and coefficient, not the finite tail.

7.2 Main theorem

Theorem 7.4 (Product birthdays on $\mathcal{A}$). *For every $x, y \in \mathcal{A}$,*

$$\boxed{\ell(xy) \leq \ell(x) \otimes \ell(y)}. \quad (26)$$

Equality holds if and only if at least one of the following holds:

- (i) $x = 0$ or $y = 0$;
- (ii) $x \in \{1, -1\}$ or $y \in \{1, -1\}$;
- (iii) x and y are both ordinary integers.

Proof. Zero factors are immediate. If both nonzero factors have finite birthdays, they are dyadic reals and [Lemma 7.1](#) gives the inequality and equality classification. If exactly one factor has finite birthday, it is a nonzero dyadic constant. Multiplication by ± 1 gives equality; every other such constant gives strict inequality by [Lemma 7.2](#). If both birthdays are infinite, [Proposition 6.2](#) gives strict inequality. These alternatives exhaust all pairs in $\mathcal{A}$, and each displayed equality case does indeed give equality. $\square$

Corollary 7.5 (Degree does not rise). *For nonzero $x, y \in \mathcal{A}$,*

$$\deg \ell(xy) \leq \max(\deg \ell(x), \deg \ell(y)). \quad (27)$$

Thus for every positive integer n , $\deg \ell(x^n) \leq \deg \ell(x)$.

Proof. When both birthdays are infinite this was proved in [Proposition 6.2](#). With one finite birthday use [Lemma 7.2](#); with both finite, all the degrees are zero. The power statement follows by induction. $\square$

The degree conclusion is much stronger than the degree estimate that would follow merely from (26). That weaker estimate permits $\deg \ell(x) \oplus \deg \ell(y)$, whereas here only their maximum is needed. This explains why equality is so rare within the ring.

8 A four-coefficient formula for finite supports

Suppose $x, y \in \mathcal{A} \setminus \{0\}$ have finite index supports, with maxima α, β . Put

$$\gamma = \alpha \oplus \beta, \quad p = c_\alpha d_\beta.$$

For an ordinal θ , let θ^- denote its immediate predecessor when it exists; it is undefined at zero and at a limit. An absent coefficient, including a coefficient at an undefined predecessor, is interpreted as zero. Define

$$q = c_{\alpha^-} d_\beta + c_\alpha d_{\beta^-}. \quad (28)$$

Proposition 8.1 (No full convolution is needed). *With the notation above, the product birthday is*

$$\ell(xy) = \begin{cases} \omega \cdot (\gamma + 1), & p \notin \mathbb{D}, \\ \omega \cdot \gamma + (d(p) - 1 + E), & p \in \mathbb{D}, \end{cases} \quad (29)$$

where

$$E = 1 \iff \gamma = 0 \text{ or } (\gamma \text{ is a successor ordinal and } q \notin \mathbb{D}).$$

After locating the two support maxima, at most the top and predecessor coefficient of each factor is needed: four coefficient lookups in total. The full product support need not be constructed.

Proof. Strict monotonicity of natural sum shows that any pair of indices other than (α, β) has sum strictly less than γ . Thus the product has maximum index exactly γ , with nonzero coefficient p .

If γ is a successor, a decomposition $a \oplus b = \gamma^-$, with $a \leq \alpha$, $b \leq \beta$, can only be (α^-, β) or (α, β^-) , when defined. To see this, compare the aligned Cantor coefficients. The target differs from γ only by subtracting one from the constant coefficient. A first decrease at a positive Cantor exponent in either input cannot be offset by the other input, which is also bounded by its corresponding maximum. Thus all positive-exponent coefficients must be unchanged, and exactly one constant coefficient decreases by one. The coefficient at γ^- is consequently q , including the possibility that it cancels to zero.

The product support is finite, so it cannot accumulate at a positive last index. Its endpoint correction is therefore precisely the one in the statement. Apply [Theorem 5.1](#). $\square$

Remark 8.2. The word “four” counts coefficient lookups, not bit operations. Ordinal-index arithmetic and exact coefficient multiplication still have costs, and locating a maximum in an unsorted input requires a scan. The claim is that a quadratic-size convolution can be avoided when only the birthday is requested. For infinite supports with a last limit index, the cofinal endpoint correction must also be considered; the proposition is deliberately restricted to finite supports.

9 Examples, cancellations, and transfinite endpoints

Put $\varepsilon = \omega^{-1}$, and define the normal-form sum $u = \sum_{n=1}^{\infty} \varepsilon^n$. The following values are direct applications of [Theorem 5.1](#). They also separate phenomena that would be easy to miss in a monomial-only investigation.

Number	Birthday	Controlling feature
$\varepsilon^m, 1 \leq m < \omega$	$\omega \cdot m$	Dyadic monomial
$3\varepsilon^2$	$\omega \cdot 2 + 2$	Finite coefficient tail
$\frac{1}{3}\varepsilon^2$	$\omega \cdot 3$	Non-dyadic last coefficient
$1 + \varepsilon$	ω	Earlier finite length absorbed
$\frac{1}{3} + \varepsilon$	$\omega + 1$	Successor endpoint correction
$\frac{1}{3}\varepsilon + \varepsilon^2$	$\omega \cdot 2 + 1$	Same correction above index 0
$\omega^{-\omega}$	ω^2	Transfinite monomial index
u	ω^2	Support has no maximum
$u + \omega^{-\omega}$	$\omega^2 + 1$	Cofinal support at last index
$u + 3\omega^{-\omega}$	$\omega^2 + 3$	Cofinal endpoint with longer tail
$u + \frac{1}{3}\omega^{-\omega}$	$\omega^2 + \omega$	Non-dyadic terminal coefficient

9.1 Why a last-index formula without correction fails

The tempting formula $\ell(x) = \omega \cdot a + d(c_a) - 1$ would give birthday ω for $\frac{1}{3} + \varepsilon$, instead of $\omega + 1$. It would also give ω^2 for $u + \omega^{-\omega}$, instead of $\omega^2 + 1$. These are counterexamples to that auxiliary formula, not counterexamples to Gonshor's conjecture. The two failures have different causes, reflected in the two clauses of (16).

The second example is genuinely transfinite. No finite list of smaller indices can be cofinal in ω . Testing arbitrarily many finite truncations would therefore not, by itself, establish the birthday at the new terminal index ω .

9.2 Unbounded power-series support has a fixed birthday

Every non-polynomial formal power series in ε , with real coefficients, has an infinite support contained in $\mathbb{N}$, hence cofinal in ω . Therefore

$$\ell\left(\sum_{n=0}^{\infty} c_n \varepsilon^n\right) = \omega^2 \quad \text{if infinitely many } c_n \neq 0. \quad (30)$$

This includes the coefficient sequences $c_n = 1$, $c_n = 1/n!$, and any other real sequence with infinitely many nonzero entries. No real-variable convergence hypothesis is involved.

An instructive cancellation uses

$$f = \sum_{n=0}^{\infty} \varepsilon^n, \quad g_- = 1 - \varepsilon, \quad g_+ = 1 + \varepsilon.$$

Coefficientwise multiplication gives

$$fg_- = 1, \quad fg_+ = 1 + 2 \sum_{n=1}^{\infty} \varepsilon^n.$$

The input birthdays are the same in the two products: $\ell(f) = \omega^2$ and $\ell(g_-) = \ell(g_+) = \omega$. The output birthdays are respectively 1 and ω^2 . Thus the product birthday is not determined by the two input birthdays, even in this very small part of $\mathcal{A}$.

9.3 Cofinal supports at arbitrary limits

Let λ be any nonzero limit ordinal and let $S \subset \lambda$ be cofinal, with no maximum. For arbitrary nonzero real coefficients,

$$\ell\left(\sum_{\alpha \in S} c_\alpha \omega^{-\alpha}\right) = \omega \cdot \lambda.$$

Appending a last term at index λ with dyadic coefficient c increases the birthday by $d(c)$, not by $d(c) - 1$. Appending a non-dyadic coefficient increases it by ω .

For $\lambda = \omega^\rho$, the original birthday is $\omega^{1+\rho}$. When $\rho = \omega$, this is ω^ω , not $\omega^{\omega+1}$. This is one concrete place where keeping ordinary left multiplication separate from natural multiplication is indispensable.

10 Set-sized local subrings and sharp birthday ceilings

10.1 A hierarchy of rings

For an ordinal $\rho > 0$, set $\kappa = \omega^\rho$ and define

$$\mathcal{A}_\kappa = \{x \in \mathcal{A} : S(x) \subseteq \kappa\}. \quad (31)$$

Every such object is encoded by a function $\kappa \rightarrow \mathbb{R}$, so this is a set, unlike the full class $\mathcal{A}$.

Proposition 10.1. *The set $\mathcal{A}_\kappa$ is a subring of $\mathcal{A}$, and*

$$\max\{\ell(x) : x \in \mathcal{A}_\kappa\} = \omega \cdot \kappa = \omega^{1+\rho}. \quad (32)$$

A nonzero element attains this maximum exactly when its index support is cofinal in κ .

Proof. An ordinal below ω^ρ has all Cantor exponents below ρ . Natural sum preserves this property, so the ring closure proof applies without leaving κ .

If a support has supremum $\sigma < \kappa$, its birthday is at most $\omega \cdot (\sigma + 1) < \omega \cdot \kappa$, since κ is a limit ordinal and left multiplication by ω is strictly increasing in its right argument. A cofinal support has no maximum and birthday $\omega \cdot \kappa$ by (14). Such an element exists: take coefficient 1 at every index below κ . Finally $\omega \cdot \omega^\rho = \omega^{1+\rho}$. $\square$

These ceilings are not obtained by restricting to numbers of birthday *strictly less* than a selected ordinal. They come from restrictions on normal-form indices. The resulting ring can contain elements whose birthday equals the stated ceiling, and products still remain inside.

10.2 Units and a formal-power-series interpretation

Proposition 10.2. *An element of $\mathcal{A}_\kappa$ is a unit in that ring if and only if its constant coefficient is nonzero. In particular $\mathcal{A}_\kappa$ is a local ring, with maximal ideal consisting of the elements having zero constant coefficient.*

Proof. The constant coefficient of a product is the product of the constant coefficients: the only solution to $\alpha \oplus \beta = 0$ is $\alpha = \beta = 0$. Thus a zero constant coefficient prevents invertibility.

Conversely write a series with nonzero constant coefficient as $c(1 + v)$, where $c \in \mathbb{R} \setminus \{0\}$ and v has zero constant coefficient. Consider the formal geometric inverse

$$c^{-1} \sum_{n=0}^{\infty} (-v)^n.$$

We verify local finiteness, rather than appealing to analytic convergence. For $\gamma = \sum_{j=1}^m \omega^{\rho_j} n_j$, put $\text{wt}(\gamma) = \sum_j n_j$, and $\text{wt}(0) = 0$. Natural sum adds these weights. Every positive ordinal has weight at least 1. Consequently a sum of n positive indices can equal γ only if $n \leq \text{wt}(\gamma)$. For each such n , distributing the finite Cantor coefficient list among n slots gives only finitely many ordered decompositions. Hence each coefficient of the geometric inverse is a finite sum over finitely many values of n .

All indices remain below κ , by natural-sum closure. Coefficientwise multiplication is therefore legitimate, and the finite geometric cancellation at each coefficient shows $(1 + v) \sum_{n \geq 0} (-v)^n = 1$. Thus the series is an inverse in $\mathcal{A}_\kappa$. The constant-coefficient map onto $\mathbb{R}$ has as kernel the stated ideal, and every element outside it is a unit. This proves that the ideal is maximal and unique. $\square$

To understand the algebra behind this proof, assign a formal variable X_ξ to $\omega^{-\omega^\xi}$, for each $\xi < \rho$. The Cantor normal form $\alpha = \sum_j \omega^{\xi_j} n_j$ corresponds to the ordinary monomial $\prod_j X_{\xi_j}^{n_j}$. Natural addition of indices becomes multiplication of these monomials. Thus $\mathcal{A}_\kappa$ is a coefficientwise formal-series ring in these variables. When there are infinitely many variables, this means arbitrary coefficient functions on the finite-support monomials; it should not be confused without further argument with a differently defined topological completion.

Every individual $x \in \mathcal{A}$ belongs to some $\mathcal{A}_\kappa$, by choosing ρ with $\sup S(x) < \omega^\rho$. This also avoids any need to interpret the proofs as set-indexed sums over a proper class.

11 Computation and reproducibility

11.1 What the programs implement

The companion module `code/ordinal_series.py` represents ordinals below ω^ω by finite tuples of nonnegative integers:

$$(a_0, a_1, \dots, a_k) \longleftrightarrow \omega^k a_k + \dots + \omega a_1 + a_0.$$

It distinguishes ordinary addition, natural addition, natural product, ordinary left multiplication by ω , and the order type of an ordinal interval. Coefficients are exact Python `Fraction` objects, including both dyadic and non-dyadic rational numbers.

There are three routes to a finite-support product birthday. The endpoint evaluator implements [Theorem 5.1](#). The block evaluator implements (13) directly, using shared-minus deletion and the extra non-dyadic deletion. The four-coefficient evaluator implements [Proposition 8.1](#) without constructing the product. The verifier compares all three, where applicable, against an explicitly constructed exact coefficient convolution.

11.2 Executed checks

The default run used seed 20260920. Its recorded result is **PASS**; detailed output is in `verification.json`.

Check	Number executed
Dyadic birthday versus explicit finite sign string	2,313
Ordinal interval concatenation identities	55
Endpoint versus block birthday, exhaustive series family	64,401
Exhaustive product pairs, unordered with repetition	76,636
Additional seeded random product pairs	20,000
Four-coefficient formula versus full convolution	96,636
Named finite-support examples	7

The exhaustive birthday family uses support size at most three from ten indices, including ω , $\omega + 1$, $\omega \cdot 2$, and ω^2 , with eight possible nonzero rational coefficients. The exhaustive product family has 391 members: support size at most two from five indices and six coefficient choices, including zero as the empty series. Every unordered pair with repetition is checked. Random tests use support size at most six and a seeded pool of 49 ordinal indices, with rational coefficients whose denominators can contain odd factors.

For each product pair the program checks the main inequality, the equality classification, the sharp support upper bound, the degree upper bound, the strict gap when both birthdays are infinite, and agreement with the four-coefficient formula. As an additional consistency check it verifies the standard additive natural-sum bound on the same inputs.

11.3 What these tests do not establish

No infinite support is represented or executed. The code treats ordinal indices with finite Cantor exponents, not arbitrary ordinals. The rational coefficient tests do not enumerate arbitrary real coefficients. Most importantly, agreement of the two birthday evaluators is not an independent formal verification of the classical sign-expansion theorem: both evaluators depend on that theorem, and both share the ordinal arithmetic implementation.

The purpose of the tests is narrower and useful: they expose off-by-one errors, ordinary/natural operation confusion, failure to handle non-dyadic coefficients, missed coefficient cancellation, and mistakes in the proposed equality condition. The transfinite proof in [Section 5](#), including its cofinal endpoint case, is the reason the mathematical statement covers infinite supports.

11.4 Reproducing the run

Only Python 3.10 or later and its standard library are needed:

```
python code/verify.py
python code/verify.py --seed 20260920 --random-pairs 20000
```

The default command writes `verification.json`. To preserve the shipped record, use a separate output file:

```
python code/verify.py --output verification-rerun.json
```

A different seed or a larger random sample can be selected explicitly. Failure raises an assertion with the offending pair or series. The code is research verification code, not a general-purpose surreal package.

12 Assessment of the attempt and the remaining general problem

12.1 What has actually been proved

The endpoint theorem gives exact birthdays on a ring of real-coefficient series with arbitrary set-sized ordinal-index support. It is not merely a calculation for powers ω^{-n} , a finite-support observation, or an estimate for individual monomials. The product theorem covers the whole ring, including cancellation and transfinite accumulation of support. Its strictness conclusion and its equality classification are consequences of a leading-Cantor-degree gap, not of numerical experimentation.

The associated four-coefficient formula makes a finite-support output birthday substantially easier to compute than the output normal form. The local-ring interpretation and the attained birthday ceilings show that the domain has familiar and nontrivial algebraic structure.

The main claims are deductions made in this investigation from classical sign-expansion theory. The literature search did not establish whether the exact endpoint formula or these specific deductions have appeared elsewhere. They should therefore be described as the results of this attempt, not as certified first discoveries.

12.2 Why the proof does not extend automatically

For a general negative surreal exponent, the sign string can contain plus signs. For example, $-\frac{1}{2}$ has sign string $-+$, whereas $-\alpha$ for an ordinal α has only minuses. In the former setting the exponent blocks are no longer uniformly ω -long. Reduced sign strings can have several retained regions rather than one terminal interval. The proof of (13) in its simplified form, and therefore the endpoint induction, no longer applies.

Allowing positive exponents changes the picture even more visibly. The numbers $x = y = \omega$ satisfy

$$\ell(xy) = \omega^2 = \ell(x) \otimes \ell(y),$$

although both birthdays are infinite. Thus the strictness theorem and equality classification proved for $\mathcal{A}$ cannot be generalized to all of $\mathbf{No}$ without a different statement. This example does not refute the general conjecture; it isolates the limitation of the stronger special-case result.

12.3 A concrete next extension

A natural next domain would allow exponents whose sign strings have one controlled plus block after an initial minus block. A successful extension would need a replacement for the single-interval calculation: it would track which exponent signs survive reduction, the ordinal sizes of their expanded blocks, and how these data behave under addition of exponents. The four-coefficient shortcut suggests separating the analysis of support endpoints from the analysis of coefficient tails, but it does not solve that new block-comparison problem.

The general conjecture remains a precise further target. The result of this attempt is a complete special-case proof with explicit boundaries, not a proof obtained by silently identifying all negative exponents with negative ordinals or by treating finite tests as transfinite evidence.

A A compact proof audit

The following checks are useful when reading or extending the argument.

Canonical objects. The quantity $\ell(x)$ is the length of the canonical sign expansion. Recursive form generation is never substituted for it.

Ordinal operations. Normal-form indices convolve by $\oplus$. The conjectured upper bound uses $\otimes$. Concatenated sign lengths use ordinary $+$ and ordinary left multiplication $\omega \cdot \alpha$. In $\omega^{1+\rho}$, the exponent addition is ordinary.

Coefficient types. Dyadic status is checked on the actual nonzero coefficient after all contributions at an index have been added. Products of non-dyadic reals can be dyadic, and a dyadic scalar can turn a non-dyadic real into a dyadic one. Both possibilities are included.

Limit stages. The limit argument takes the supremum of lengths of concatenated sign prefixes. It does not assume any general continuity of surreal addition or multiplication. A last index cofinal with its predecessors contributes an entire coefficient string, explaining the correction by one.

Set-sizedness and local finiteness. Every support is a set. The natural-sum fiber count (10) makes each convolution coefficient finite. For the geometric inverse, the finite Cantor weight bounds the number of factors that can contribute at a prescribed index.

Equality. The cases “both finite”, “one finite”, and “both infinite” are exhaustive because finite-birthday surreals are precisely dyadic reals. Strictness in the last case follows from a full leading-exponent gap; strictness in the mixed case follows from the leading coefficient.

B Source and artifact notes

The historical conjecture is documented by [Lán12] and the later first-person account [Ehr24]. The latter is the source used here for the reported scope of the van den Dries–Ehrlich monomial result. The original 2001 article was not used as a directly inspected proof source in preparing this draft.

The actual sign-expansion ingredient was checked in [BG22, Definition 2.20 and Theorem 2.21, printed p. 13]. Gonshor’s book is credited as the original source of that theorem; the version used in this derivation is the explicitly identified restatement. The normal-form background is as in [Ber20]. The distinction between forms and values, and the finite dyadic birthday formula, are also visible in [Rou18].

The archive contains this L^AT_EX source, its compiled PDF, the `references.bib` bibliography, two Python files, `verification.json`, a README, and build instructions. It does not redistribute the cited books or papers. The mathematical article is the primary result; the code provides reproducible finite checks with the limitations stated in Section 11.

References

- [Ber20] Alessandro Berarducci. *Surreal numbers, exponentiation and derivations*. 2020. arXiv:2008.06878v1, especially Sections 5, 6, and 8–10.
<https://arxiv.org/abs/2008.06878>.
- [BG22] Olivier Bournez and Quentin Guilmant. *Surreal fields stable under exponential and logarithmic functions*. 2022. arXiv:2201.08199v1, especially Definition 2.20 and Theorem 2.21, printed p. 13. <https://arxiv.org/abs/2201.08199>.
- [Ehr24] Philip Ehrlich. Answer to “Birthday of combinatorial game product”. *MathOverflow*, question 476824, 13 August 2024, edited the same day. Consulted 20 September 2026.
<https://mathoverflow.net/questions/476824>.
- [Gon86] Harry Gonshor. *An Introduction to the Theory of Surreal Numbers*. London Mathematical Society Lecture Note Series, vol. 110. Cambridge University Press, 1986. Original source credited for the conjecture and the normal-form sign-expansion theorems; the latter are used via [BG22].
- [Lán12] Lukáš Lánský. “Length of surreal product”. *Open Problem Garden*, 7 April 2012. Consulted 20 September 2026.
https://www.openproblemgarden.org/op/length_of_surreal_product.
- [Rou18] Matthew Roughan. *Surreal Birthdays and Their Arithmetic*. 2018. arXiv:1810.10373v2.
<https://arxiv.org/abs/1810.10373>.