

A MERGED RESEARCH CONTINUATION

Global Support Obstructions

Moving Divisors, Mittag–Leffler,
and a Picard Dichotomy

September 21, 2026

Abstract

We study the sheaf $\mathcal{H}_\Gamma$ of well-ordered Hahn series whose coefficients are ordinary holomorphic functions on a *common* ordinary domain, for a fixed set-sized surreal value group Γ , together with its nonnegative-support subring sheaf $\mathcal{H}_{\Gamma, \geq 0}$ and its positive-support additive subsheaf $\mathcal{H}_{\Gamma, +}$. The questions are global: when do infinitely many individually legitimate infinitesimal analytic prescriptions admit one coherent realization over the ordinary complex plane?

We prove a sharp moving-divisor theorem. A family of prescribed infinitesimal root clusters, encoded by monic cluster polynomials $P_c(u) = u^{m_c} + \sum_{j < m_c} a_{c,j} u^j$ over a closed discrete ordinary set A , is the exact zero divisor of one nonzero element of $\mathcal{H}_\Gamma(\mathbb{C})$ if and only if the *symmetric support* $E(P) = \bigcup_c \bigcup_{j < m_c} \text{supp}(a_{c,j})$ is well ordered. It is the support of the symmetric coefficients, not of separately labelled roots, that decides. Simple zeros at $n + t^{1/n}$ are therefore not realizable alone, while *all* roots of $(z - n)^n = t$ are, so a globally principal effective divisor can have a nonprincipal effective sub-divisor.

We prove the matching criteria for discrete Hahn principal parts, for deformed Hermite interpolation — whose correct target is a *support-restricted* product of the finite algebras $K_\Gamma[u]/(P_c)$, not their unrestricted product — for moving poles, and an exact additive Cousin criterion in terms of the singular support on a puncture-and-disk cover. We show that every ordinary-base stalk of $\mathcal{H}_\Gamma$ is a principal ideal domain, though generally not a local ring, which is what licenses classifying all tensor-invertible modules rather than only locally free rank-one sheaves. The resulting Picard group vanishes exactly when $\Gamma = \{0\}$ or Γ is order-isomorphic to $\mathbb{Z}$; for noncyclic Γ its complex dimension is at least $2^{\aleph_0}$, with equality when Γ is countable.

A second part carries the same coefficient sheaf over a second ordinary base: a fixed *compact* connected ordinary Riemann surface X of genus g . There the answers are different in both directions. A closed discrete cluster set is finite on this base, so its well-ordering obstruction disappears; in its place the Picard group splits canonically and completely,

$$\text{Pic}_{\text{lf}}(X, \mathcal{H}_\Gamma) \cong H^1(X, \underline{\Gamma}) \oplus \text{Pic}(X) \oplus (H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_\Gamma) \cong \Gamma^{2g} \oplus \text{Pic}(X) \oplus \mathfrak{m}_\Gamma^g,$$

the first summand being exactly the obstruction to a nonzero coherent meromorphic section — so there are line bundles of arbitrarily high degree with no such section at all. In the zero-valuation sector an explicit finite matrix $B = p_1 \delta (1 + h\delta)^{-1} i_0$ with entries in $\mathfrak{m}_\Gamma$ computes both cohomology groups and gives Riemann–Roch with no valuation-rank or cofinality hypothesis. A root-free Abel functional built from Newton power sums decides principality of a finite moving cluster over arbitrary, not necessarily divisible, value groups; on $y^2 = 1 + x^5$ the first symmetric displacement cancels while the obstruction does not. Finally, the inverse limit over all powers of one scale δ is split surjective with kernel exactly $H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \bigcap_n t^{n\delta} \mathcal{O}_\Gamma$, nonzero whenever $g > 0$ and the convex subgroup generated by δ is proper.

All proofs are support-controlled constructions in a fixed set-sized workspace, not convergence arguments in the fine surreal topology. *The compact results use compactness twice and do not specialize to the plane by setting $X = \mathbb{C}$ and $g = 0$* ; the two bases are kept apart throughout.

Status. This article merges three independent research continuations of one supplied manuscript; see [section 1.3](#). The principal global criteria, the restricted interpolation algebra, the exact singular-support obstruction and the Picard-group dichotomy over the plane, together with the compact-base classification, the finite obstruction matrix, the root-free Abel criterion and the one-scale comparison kernel, are proposed new results in the precise framework of that manuscript, and full proofs are given. The literature checks, conducted on 21 September 2026, were targeted rather than exhaustive; they did not locate these formulations, and they do not certify priority. No established named published conjecture is claimed solved. The proofs have not been refereed or machine checked, and there is no proof-assistant verification; the accompanying symbolic scripts validate displayed finite examples and are not machine verification of the general arguments.

Contents

1	The question: from local roots to global compatibility	1
1.1	The main results in concrete form	1
1.2	Conventions, and the coefficient ring that every theorem is stated over	2
1.3	Provenance, and relation to published work	3
2	The workspace and the support calculus	5
2.1	Fixed Hahn fields and infinitesimals	5
2.2	The Hahn–Neumann support lemma	5
3	The coefficient sheaf	6
3.1	Sections, evaluation and gluing	6
3.2	Units, positive logarithms, and the integral subring	8
4	The ordinary analytic input, with constructions	9
5	Local preparation and deformed division, with uniform support	10
5.1	Preparation by an exponent recursion	11
5.2	Deformed division and uniform remainders	12
6	The ordinary-base stalks, and what they buy	13
7	A sharp Hahn Mittag–Leffler theorem	15
7.1	Discrete principal-part data	15
7.2	Why this is not a summability condition over the points	16
8	The global theorem for moving surcomplex divisors	17
8.1	Cluster polynomials and the correct compatibility invariant	17
8.2	Uniqueness modulo global units	18
8.3	Root supports versus symmetric supports	19
9	Sharp examples and a nonfactorable divisor splitting	19
9.1	An impossible sequence of simple zeros	19
9.2	Complete clusters with unbounded multiplicities	19
9.3	A principal effective divisor with a nonprincipal sub-divisor	20
9.4	No repair by enlarging the value group	21
10	A support-restricted Chinese remainder theorem	21
10.1	The correct target algebra	21
10.2	A support-controlled right inverse	22
10.3	Bounded raw values need not be interpolable	23

11 A Mittag–Leffler theorem for moving poles	23
12 Near-identity automorphisms and interpolation of motions	24
13 The exact additive Cousin obstruction	25
13.1 A puncture-and-disk cover, and the singular support	26
13.2 An explicit infinite-dimensional family of obstructions	27
13.3 The divisor obstruction as a line-bundle class	28
14 The Picard-group dichotomy over the plane	29
14.1 Separating the ordinary and positive-support factors	29
14.2 Exactly which value groups give vanishing over the plane	30
14.3 The size of the nonzero plane Picard groups	31
14.4 Additive cohomology over the plane behaves differently	32
14.5 Why this does not contradict ordinary complex geometry	32
15 Moving divisors as nontrivial line bundles over the plane	32
15.1 What the existence and obstruction results say together	33
16 A second base: compact ordinary Riemann surfaces	34
16.1 Why the two global theories do not specialize to each other	34
16.2 What is kept, and what is renamed	34
17 The coefficient sheaf on a compact base, and Dolbeault comparison	35
17.1 The sheaves, and the one convention that carries over	35
17.2 The operator form of the support calculus	36
17.3 The coefficientwise resolution and the comparison theorem	37
18 The exact Picard classification over a compact base	38
18.1 Three independent components of a unit	38
18.2 A normal form with a finite support bound	39
18.3 The two bases side by side	40
19 A finite matrix for the zero-valuation sector	40
19.1 From holomorphic transitions to a global operator	40
19.2 The transferred matrix	41
19.3 Riemann–Roch, stable section counts, and integral torsion	42
20 Valuation monodromy is exactly the meromorphic obstruction	43
20.1 The meromorphic convention	43
20.2 Divisor classes occupy exactly the zero-valuation sector	44

21 A root-free Abel criterion for finite moving clusters	45
21.1 Finite cluster data and their ordinary reduction	45
21.2 Newton sums are the correct symmetric coordinates	45
21.3 The exact criterion	47
22 Compact examples: the sphere, an elliptic curve, and a nonlinear obstruction	48
22.1 The sphere, and a warning about the plane	48
22.2 An elliptic curve: exact displacement conservation	48
22.3 A genus-two cluster with zero first displacement and nonzero Abel class	49
23 The exact information lost by one-scale completion	49
23.1 Truncated coefficients and their Picard groups	50
23.2 The visible convex subgroup and the invisible ideal	50
23.3 A deformation beyond every integer-scale jet	52
23.4 Three tests, not one	52
23.5 Why arbitrary-rank support control is substantive	53
24 Interpretation, boundaries, and further questions	53
24.1 What has been established over the plane	53
24.2 What has been established over a compact base	54
24.3 What the conclusions do not say	54
24.4 Further precise directions	55
25 Conclusion	56
A Support and dependency audit	57
B Reproducible symbolic checks	59
C Provenance of the merged material	60
D Notation	62
References	62

1 The question: from local roots to global compatibility

The supplied manuscript [1] develops surcomplex analysis by separating fine-local power-series calculus on $\mathbf{SC} = \mathbf{No}[i]$ from Hahn-coherent holomorphy over an ordinary complex domain. In that middle layer a function on the infinitesimal thickening of an ordinary domain U is represented by

$$F(z) = \sum_{\gamma \in S} f_\gamma(z) t^\gamma, \quad f_\gamma \in \mathcal{O}(U), \quad S \subseteq \Gamma \text{ well ordered},$$

with $t^\gamma = \omega^{-\gamma}$ in the surcomplex interpretation. Its one-variable preparation theorem associates a finite monic polynomial to every infinitesimal zero cluster, and its root-lifting theorem counts the roots in a monad exactly. Those are starting points here, not new claims.

A question that local preparation does not answer is global. Suppose an ordinary discrete set has infinitely many points and a finite infinitesimal root configuration has been specified above each one. Can a single coherent entire datum realize all these configurations simultaneously? The classical Weierstrass theorem might suggest that ordinary discreteness suffices. It does not. A condition on the union of the *Hahn supports* is necessary, and — once the right invariant is extracted — sufficient.

A first illustration is the proposed list of simple zeros

$$z_n = n + t^{1/n}, \quad n \geq 2. \quad (1.1)$$

Every individual point is legitimate and every finite sublist is realizable, even by a polynomial. Nevertheless no Hahn-coherent entire datum has exactly these simple zeros and no others in the thickened plane. In contrast one can simultaneously realize, at every integer n , *all* roots of

$$(z - n)^n - t = 0. \quad (1.2)$$

These roots also involve the decreasing exponents $1/n$. The difference is that their symmetric cluster polynomial uses only the single exponent 1. That distinction is the content of the first main theorem, not an artifact of an overly strong sufficient hypothesis.

The same issue appears in the additive Cousin, or Mittag–Leffler, problem. Solving each complex coefficient separately is easy; assembling the solutions into one Hahn series may be impossible. Exponentiating the positive-support version of this obstruction produces nontrivial line bundles even though the ordinary base is the contractible Stein manifold $\mathbb{C}$.

1.1 The main results in concrete form

Fix a set-sized ordered additive subgroup $\Gamma \subseteq \mathbf{No}$ and a closed discrete $A \subset \mathbb{C}$. At each $c \in A$ prescribe a monic cluster polynomial

$$P_c(u) = u^{m_c} + \sum_{j < m_c} a_{c,j} u^j, \quad a_{c,j} \in \mathfrak{m}_\Gamma, \quad m_c \geq 1, \quad (1.3)$$

with no common upper bound on the multiplicities m_c and no separation assumption on the roots inside a cluster. The *symmetric support* is

$$E(P) = \bigcup_{c \in A} \bigcup_{j < m_c} \text{supp}(a_{c,j}) \subseteq \Gamma_{>0}, \quad (1.4)$$

and the family is *admissible* when $E(P)$ is well ordered. The principal conclusions are as follows. Items (1)–(5) are theorems over the ordinary base $\mathbb{C}$ and occupy [sections 2 to 15](#); item (6) is a second theory over a *compact* ordinary base and occupies [sections 16 to 23](#). Neither part implies the other; [section 16.1](#) says exactly why, and every statement names its base.

1. **Sharp moving-divisor realization** (theorem 8.2). The family (1.3) is the exact zero divisor of a nonzero $F \in \mathcal{H}_\Gamma(\mathbb{C})$ if and only if it is admissible; and then any prescribed ordinary entire f_0 with zero divisor $\sum_c m_c[c]$ can be retained as the exponent-zero coefficient of a realizer supported in $E(P)^*$.
2. **Sharp principal-part compatibility** (theorem 7.2). A discrete family of Hahn principal parts is the family of principal parts of a single section on the punctured plane if and only if the union of the principal-part exponents is well ordered. Relative to an admissible pole bound the same happens for moving poles (theorem 11.1).
3. **Support-restricted interpolation** (theorem 10.2 and corollary 10.3). For an admissible family the correct target of global interpolation is the *support-restricted* product $\mathcal{B}(P)$ of the finite algebras $K_\Gamma[u]/(P_c)$, not their unrestricted product, and $\mathcal{H}_\Gamma(\mathbb{C})/(G) \cong \mathcal{B}(P)$ for any divisor realizer G .
4. **An exact Cousin obstruction** (theorem 13.1). On a puncture-and-disk cover, additive gluing is possible exactly when the exponents carrying *nonremovable* ordinary coefficient singularities form a well-ordered set.
5. **Principal-ideal-domain stalks and a line-bundle dichotomy** (theorems 6.1, 14.3 and 14.4). Every ordinary-base stalk of $\mathcal{H}_\Gamma$ is a principal ideal domain, though nonlocal for $\Gamma \neq \{0\}$; consequently every tensor-invertible $\mathcal{H}_\Gamma$ -module is locally free of rank one, and

$$\mathrm{Pic}(\mathbb{C}, \mathcal{H}_\Gamma) = 0 \iff \Gamma = \{0\} \text{ or } \Gamma = \mathbb{Z}\delta \ (\delta > 0).$$

For noncyclic Γ one has $\dim_{\mathbb{C}} \mathrm{Pic}(\mathbb{C}, \mathcal{H}_\Gamma) \geq 2^{\aleph_0}$, with equality when Γ is countable; in particular $\dim_{\mathbb{C}} \mathrm{Pic}(\mathbb{C}, \mathcal{H}_{\mathbb{Q}}) = 2^{\aleph_0}$.

6. **A complete theory over a compact base** (theorems 17.7, 18.3, 19.2, 20.1, 21.5 and 23.3). For a compact connected ordinary Riemann surface X of genus g the cohomology of an ordinary bundle with Hahn coefficients is the ordinary cohomology tensored with the coefficient ring, the Picard group splits as $\Gamma^{2g} \oplus \mathrm{Pic}(X) \oplus \mathfrak{m}_\Gamma^g$, the first summand is exactly the obstruction to a nonzero coherent meromorphic section, the zero-valuation sector is computed by an explicit finite matrix over $\mathcal{O}_\Gamma$ with entries in $\mathfrak{m}_\Gamma$, a root-free Newton-sum Abel functional decides principality of a finite moving cluster, and the inverse limit over the powers of one scale has an exactly identified kernel.

These are global existence and obstruction theorems, not formal analogues of local identities. They distinguish finite from infinite collections of local data, cyclic from noncyclic value groups, and — across the two parts — a noncompact from a compact ordinary base.

1.2 Conventions, and the coefficient ring that every theorem is stated over

Several conventions are fixed once here and used without further comment.

One coefficient ring. *Every* numbered statement in this article is a statement about the sheaf

$$\mathcal{H}_\Gamma(U) = \mathcal{O}(U)((t^\Gamma)),$$

whose Hahn coefficients are ordinary holomorphic functions on the *common* ordinary domain U , or about its subsheaves $\mathcal{H}_{\Gamma, \geq 0}$ and $\mathcal{H}_{\Gamma, +}$, or about the ordinary-base stalks

$$R_c = (\mathcal{H}_\Gamma)_c = \varinjlim_{U \ni c} \mathcal{H}_\Gamma(U)$$

formed by shrinking ordinary neighbourhoods of an ordinary point c . *The ordinary base itself is part of the statement:* in sections 2 to 15 it is the plane $\mathbb{C}$ and $U \subseteq \mathbb{C}$, while in sections 16

to 23 the identical coefficient recipe is applied with U ranging over the opens of a fixed compact connected ordinary Riemann surface X (definition 17.1), the ordinary meromorphic version $\mathcal{M}_\Gamma$ being introduced there as well. Every numbered statement names its base, and section 16.1 records why the global theorems require their stated bases; local stalks agree in coordinate charts as explained in remark 18.5. Nothing here is asserted about, or transported to or from, a radius-free coefficient ring $\mathbb{C}\{z\}((t^\Gamma))$ whose Hahn coefficients are individually convergent germs with no common domain of convergence, or a formal coefficient ring $\mathbb{C}[[z]]((t^\Gamma))$. The common-domain requirement is used essentially and repeatedly: in the sheaf property (proposition 3.3), in the “no disk shrinking” clause of the divisor recursion (theorem 8.2), and in the uniform support bound of local preparation (theorem 5.1). The two rings that do occur are distinguished by name where it matters: the *fixed-disk ring* $\mathcal{O}(D)((t^\Gamma))$ of sections over one chosen ordinary disk D , in which preparation and division take place, and the *ordinary-base stalk* R_c , a filtered colimit of such rings, of which the principal-ideal-domain theorem speaks. The theorems are not interchangeable between the two: $\mathcal{O}(D)((t^\Gamma))$ is not claimed to be a principal ideal domain for a fixed D .

Reduction. The exponent-zero coefficient map

$$\text{red} : \mathcal{H}_{\Gamma, \geq 0} \longrightarrow \mathcal{O}, \quad \sum_{\gamma \geq 0} f_\gamma t^\gamma \longmapsto f_0,$$

is called the *reduction*. The two source manuscripts and the wider literature also call the induced operation the residue map, the standard part and the shadow; those are synonyms recorded once here and not used again. We write st only for the scalar standard part $\mathcal{O}_\Gamma \rightarrow \mathbb{C}$, $a \mapsto a_0$. For $\Gamma \neq 0$ this reduction does not extend to a ring map on $\mathcal{H}_\Gamma$; see remark 3.6.

Symbols. $\mathfrak{m}_\Gamma$ denotes the infinitesimals of the fixed Hahn field K_Γ and $\mathfrak{m}_{\mathbf{SC}}$ the infinitesimals of $\mathbf{SC}$; $\mathfrak{m}_\Gamma = \mathfrak{m}_{\mathbf{SC}} \cap K_\Gamma$. Cluster data are indexed by ordinary centres $c \in A$, with $u = z - c$ the local coordinate. The symmetric support of a cluster family is written $E(P)$ throughout; one of the sources writes $S_{\mathcal{D}}$ for the same set. The positive-support exponential and logarithm are written Exp_H and Log_H to keep them apart from the ordinary exponential e^h of a holomorphic function. The simultaneous deformed remainder map is $\mathcal{R}_P$. Line-bundle groups are written Pic_{lf} for locally free rank-one modules and Pic for tensor-invertible modules; corollary 6.3 proves these agree here, and remark 6.4 records exactly what that proof buys.

Set-sizedness. All supports, families, covers and recursions are sets. The value group Γ is fixed and set-sized, which is what makes the sheaves and their cohomology set-valued. Every finite tuple of surcomplex numbers occurring in an argument lies in some set-sized divisible workspace, obtained as the divisible hull of the group generated by the relevant normal-form supports; this never authorizes a proper-class sum or a proper-class polynomial ring.

1.3 Provenance, and relation to published work

This article merges three independent continuations, referred to below as *Manuscript A* (“Global Support Obstructions in Surcomplex Analysis: Moving Divisors, Mittag–Leffler Theory, and a Picard-Group Dichotomy”), *Manuscript B* (“Global Divisors and Cohomological Obstructions in Hahn-Coherent Surcomplex Analysis”) and *Manuscript C* (“Compact-Curve Geometry over Surcomplex Hahn Fields: Picard classification, valuation monodromy, finite cohomology models, and an exact Abel criterion”). Manuscripts A and B continue the same supplied manuscript [1] over the ordinary plane. The terminology $U^\sharp$, strong summability and Hahn coherence follows [1], as do the ordinary-base gluing mechanism, the injectivity and compatibility of evaluation, and the

one-variable preparation, division, root-lifting and pole-free-fraction arguments. These are credited, reproved here in the fixed- Γ form needed for infinitely many simultaneous prescriptions, and not claimed as new.

Where two continuations prove the same theorem, the merged text prints one proof; where one proves strictly more, the stronger statement is printed and the weaker one is retained as the statement that survives without the extra ingredient. Two such places are flagged explicitly in the text: [remark 6.4](#) on tensor-invertible versus locally free rank-one modules, and [remark 14.5](#) on the size of the Picard group. Attribution of the individually contributed results is recorded in [appendix C](#).

Manuscript C is a continuation of a different kind: it changes the ordinary base to a compact curve and was written against this repository rather than against [1], having inspected the documentation snapshot at commit `aa846271b4dcae2c055b216126a87210292ec19b` — specifically the documentation index and the README of this report — and made no change to it. It states that this report “explicitly emphasizes common ordinary domains and noncompact support obstructions” and “does not provide the compact-base classification or the fixed-scale Picard comparison proved here”, and [sections 16 to 23](#) are exactly that material. Two of its own boundaries are recorded here once and kept at their points of use. First, *it does not assume, and does not verify, this report’s unrefereed claims* — it names the principal-ideal stalk theorem [theorem 6.1](#) specifically — and it does not claim that every source manuscript or Lean file in the repository was individually audited; see [remark 18.5](#). Second, its novelty claim is the precise combination of arbitrary-rank support certificates, the finite matrix, the exact divisor criterion and the fixed-scale comparison kernel in the common-domain Hahn category: *the existence of a monodromy obstruction is not asserted to be a new general idea*. Logarithmic Picard theory and its tropicalization already study monodromy and divisorial representability [12], in a different geometric setting; here the base is a fixed smooth complex curve, the relevant group is H^1 of its ordinary topology, and no logarithmic structure, tropical dual graph or bounded-monodromy condition is imposed. The transferred finite matrix is a two-term instance of homological perturbation, classical and credited [5], and cohomological obstructions to deforming ordinary line-bundle cohomology have a substantial history [8]. The classical analytic input for the compact base is ordinary compact Riemann surface theory — Dolbeault, Hodge, Serre duality, Riemann–Roch, the Jacobian and Abel’s theorem [13]; these are used on ordinary complex curves only, and *no general surreal version of Hodge theory or Serre duality is proved or assumed*.

Normal forms, infinitesimal power-series evaluation and the underlying surreal field theory have established predecessors, including van den Dries–Ehrlich [17] and Gonshor [9]; the algebraic closure of Hahn fields with divisible value group is Poonen’s [15, Corollary 4]. The summability algebra is the classical Hahn–Neumann support theory [14]. Berarducci–Mantova’s surreal function theory [3] and Costin–Ehrlich’s integration theory [4] address different extension and summation questions; *we do not identify their function classes with the sheaf studied here*. Ehrlich–Kaplan construct canonical exponentials for surreal and surcomplex exponential fields [7]; *no novelty claim about merely defining a surcomplex exponential is made here, and no global exponential construction is claimed as a contribution*.

A terminological distinction matters. The Hahn-holomorphic and Hahn-meromorphic functions of Müller and Strohmaier are normally convergent generalized expansions near a point on a logarithmic cover, with a different analytic variable and admissibility condition [11, Definitions 2.3 and 2.5]. In our notation the Hahn parameter is an independent surreal scale and the coefficients are ordinary holomorphic functions on a common ordinary base; there is no ordinary convergence requirement across the Hahn exponents. *Similar terminology does not make the objects or the global support conditions identical, and no equivalence of the two categories is assumed*.

The classical analytic inputs are the Weierstrass product theorem, Mittag–Leffler’s theorem, entire Hermite interpolation, the scalar additive Cousin theorem and the vanishing of $H^1(\mathbb{C}, \mathcal{O})$ and $\text{Pic}(\mathbb{C}, \mathcal{O})$; see [2, 10, 6]. These are proved or cited in [section 4](#) and are kept strictly separate from the Hahn arguments. *The ordinary interpolation tools are classical, not new results of this article. The present proofs do not apply Cartan’s theorem to $\mathcal{H}_\Gamma$ as though it were a coherent $\mathcal{O}$ -module; see [section 14.5](#).* The cocycle and torsor interpretation of degree-one cohomology is standard [16, Tag 02FQ], and the local-stalk qualification for invertible sheaves is [16, Tag 09NT].

2 The workspace and the support calculus

2.1 Fixed Hahn fields and infinitesimals

Let Γ be a set-sized ordered additive subgroup of $\mathbf{No}$. Its complex Hahn field is

$$K_\Gamma = \mathbb{C}((t^\Gamma)) = \left\{ \sum_{\gamma \in S} a_\gamma t^\gamma : S \subseteq \Gamma \text{ well ordered, } a_\gamma \in \mathbb{C} \right\}, \quad (2.1)$$

with zero coefficients omitted. Conway normal forms embed this field in $\mathbf{SC}$ by $t^\gamma = \omega^{-\gamma}$. For $a \neq 0$ put $v(a) = \min \text{supp}(a)$, and $v(0) = +\infty$. Write

$$\mathfrak{m}_\Gamma = \{a \in K_\Gamma : v(a) > 0\}, \quad \mathcal{O}_\Gamma = \{a \in K_\Gamma : v(a) \geq 0\},$$

so that $0 \in \mathfrak{m}_\Gamma$. Elements of $\mathfrak{m}_\Gamma$ are infinitesimal in the surreal modulus and $\mathcal{O}_\Gamma$ consists of the finite elements of the workspace; the coefficient at exponent 0 is the standard part st of a finite element. We use algebraic closedness of $\mathbf{SC}$, and occasionally of K_Γ when Γ is divisible [15, Corollary 4]. No divisibility assumption enters the support, preparation, Mittag–Leffler, interpolation, Cousin or Picard theorems; roots of polynomials may always be discussed in $\mathbf{SC}$ or in a divisible Hahn enlargement.

A family of Hahn series is *strongly summable* when its total support is well ordered *and* only finitely many members contribute at each exponent. Addition at an exponent is then a finite sum. Both conditions are checked separately everywhere below.

2.2 The Hahn–Neumann support lemma

For $E \subseteq \Gamma_{>0}$ set $E^* = \{e_1 + \cdots + e_k : k \geq 0, e_i \in E\}$, the empty sum being zero.

Lemma 2.1 (Support calculus). *If S, T are well-ordered subsets of an ordered abelian group, then $S + T$ is well ordered and every element has only finitely many representations $s + t$ with $s \in S, t \in T$. If $E \subseteq \Gamma_{>0}$ is well ordered, then E^* is well ordered and every exponent is represented by only finitely many finite nondecreasing words in E ; in particular only finitely many word lengths occur at a fixed exponent. Consequently, a formal power series in finitely many positive-support Hahn elements is strongly summable, and more generally a family indexed by a term of a well-ordered set S together with a finite word in E is strongly summable whenever each such word contributes finitely many terms.*

Proof. The first two assertions are the Hahn–Neumann support lemma [14, 15]. For the last assertion the two requirements are separated. Well ordering follows by applying the first assertion to S and E^* . For a fixed output exponent there are finitely many pairs $(s, m) \in S \times E^*$ with $s + m$ equal

to it; for each m there are finitely many nondecreasing words in E summing to m , and each finite word has finitely many orderings. Hence there are finitely many contributions before cancellation, which is exactly strong summability. $\square$

The finiteness assertion refers to words in the *set of exponents*. It does not refer to an infinite family of spatial points carrying the same exponent, and a well-ordered support alone does not justify an arbitrary rearrangement when infinitely many summands contribute at one exponent. Every infinite operation performed below is audited against [lemma 2.1](#); the audit is collected in [appendix A](#).

Remark 2.2 (Three support cautions). First, strong summability is *not* a valuation limit. For $\Gamma = \mathbb{Q} + \mathbb{Q}\omega$ the strong sum of t^n over $n \geq 0$ is $(1 - t)^{-1}$, yet the tails never reach valuation ω . No argument below has the form “the valuation improves by a fixed positive amount at each iteration”; the recursions are well founded on a well-ordered exponent set, not contractive.

Second, “finitely many dependencies at a given exponent” does not mean “finitely many exponents below it”. The latter already fails below ω in $\mathbb{Q} + \mathbb{Q}\omega$. It is the former, and only the former, that legitimizes the finite-parameter replacements in the recursions of [theorems 5.1, 8.2 and 12.1](#) and [lemma 5.3](#).

Third, the phrase *positive perturbation* always refers to a well-ordered positive support, never merely to positive valuations of separately chosen terms: in a non-Archimedean ordered group $n\delta$ need not eventually exceed every positive element, and no proof below relies on that false implication. Relatedly, well ordering cannot be weakened to positivity, since $\{1/r : r \geq 1\}$ is positive but not well ordered — this is precisely the set that obstructs [\(1.1\)](#).

For a complex vector space V we write $V((t^\Gamma))$ for Hahn families with coefficients in V . Every complex-linear map $V \rightarrow W$ extends coefficientwise, without increasing support and without any continuity or norm hypothesis. This applies to differentiation, Taylor-jet extraction, ordinary interpolation operators and division by a fixed ordinary monomial, and it is used constantly.

3 The coefficient sheaf

3.1 Sections, evaluation and gluing

Definition 3.1 (Hahn-coherent coefficient sheaf). For connected nonempty ordinary open $U \subseteq \mathbb{C}$ define

$$\mathcal{H}_\Gamma(U) = \mathcal{O}(U)((t^\Gamma)) = \left\{ \sum_{\gamma \in S} f_\gamma t^\gamma : f_\gamma \in \mathcal{O}(U), S \subseteq \Gamma \text{ well ordered} \right\}. \quad (3.1)$$

For disconnected U take the product over connected components. Let $\mathcal{H}_{\Gamma,+}$ be given by requiring strictly positive exponents and $\mathcal{H}_{\Gamma,\geq 0}$ by requiring nonnegative exponents, componentwise; these are respectively an additive sheaf closed under multiplication and a sheaf of unital subrings of $\mathcal{H}_\Gamma$. The *support* of a datum is the set of exponents whose coefficient function is not identically zero — not the support of its value at a particular point. Derivatives differentiate coefficient functions and leave t^γ fixed. Because $\mathcal{O}(U)$ is a domain for connected U , the least exponent $v_H(F)$ satisfies the usual valuation identities for sums and products. The subscript $+$ refers to exponents, never to the sign of a complex coefficient.

Note the asymmetry that defines the whole theory: the coefficients are ordinary holomorphic functions on a common ordinary domain, while the Hahn variable is an independent surreal scale.

For an ordinary domain U write

$$U^\sharp = \{c + \epsilon : c \in U, \epsilon \in \mathfrak{m}_{\mathbf{SC}}\}, \quad \mu(c) = c + \mathfrak{m}_{\mathbf{SC}},$$

and define the evaluation of (3.1) by

$$F^\sharp(c + \epsilon) = \sum_{\gamma \in S} \sum_{k \geq 0} t^\gamma \frac{f_\gamma^{(k)}(c)}{k!} \epsilon^k. \quad (3.2)$$

Proposition 3.2 (Evaluation and identity). *Formula (3.2) is strongly summable, with support in $S + (\text{supp } \epsilon)^*$, and defines an injective algebra homomorphism from $\mathcal{H}_\Gamma(U)$ into functions on $U^\sharp$. It respects addition, multiplication, coefficientwise differentiation and formal Taylor expansion in infinitesimal displacements. A coherent datum vanishing on a nonempty ordinary open subset of connected U is zero.*

Proof. Let $E = \text{supp}(\epsilon)$. The total support lies in $S + E^*$, and a power ϵ^k contributes words of length k , so lemma 2.1 gives both well ordering and finite multiplicity at each exponent. The ordinary Taylor multiplication identities may therefore be regrouped, which proves the algebra assertion. Evaluation at an ordinary point is $\sum_\gamma f_\gamma(c) t^\gamma$, so uniqueness of Hahn normal forms and the ordinary identity theorem give injectivity and the last assertion. Taylor recentring uses the same support computation with the union of the finitely many displacement supports; the resulting local series has the coefficientwise derivatives, and its fine differentiability follows by factoring the remainder as the square of the increment times a locally bounded Hahn expression, as in the local evaluation argument of [1, §5]. $\square$

On the workspace thickening, where $\epsilon \in \mathfrak{m}_\Gamma$, the output lies in K_Γ . Allowing infinitesimals outside the workspace changes the evaluation domain, not the coefficient sheaf or its cohomology; roots are counted in the full monad in $\mathbf{SC}$ unless a field is explicitly fixed. The global constructions below are first equalities of coefficient data, and evaluation is applied only after those data have been assembled. *No sequence of partial Hahn sums is asserted to converge in the full fine surreal topology, and there is no claim that ordinary partial sums converge in that topology.*

For $F = f_0 + E$ with $E \in \mathcal{H}_{\Gamma,+}(U)$ and $f_0(U) \subseteq V$, composition with $H = \sum_\delta h_\delta t^\delta \in \mathcal{H}_\Gamma(V)$ is

$$H \circ F = \sum_\delta \sum_{k \geq 0} t^\delta \frac{h_\delta^{(k)} \circ f_0}{k!} E^k, \quad (3.3)$$

with support contained in $\text{supp } H + (\text{supp } E)^*$; evaluation turns (3.3) into actual composition. This formula is used in section 12.

Proposition 3.3 (Support-preserving sheaf property). *The prescriptions $\mathcal{H}_\Gamma$, $\mathcal{H}_{\Gamma,\geq 0}$, $\mathcal{H}_{\Gamma,+}$ are sheaves on the ordinary plane. On a connected base, compatible local sections glue to a section with a single well-ordered support.*

Proof. Refine a cover to connected opens. On an overlap, equality of Hahn data is equality of each coefficient function. A holomorphic coefficient that is nonzero on a connected open cannot vanish identically on a nonempty open subset, so two sections on intersecting connected opens with equal restrictions have the same exact support. The intersection graph of a cover of a connected space by connected opens is connected — otherwise its graph components would separate the space — hence all nonempty local descriptions share one support. Glue each holomorphic coefficient and retain that support; uniqueness is coefficientwise, and positivity and nonnegativity are preserved. On disconnected bases apply this to each component. This is the gluing mechanism of [1, §5]. $\square$

Notice what this does *not* say. Equality on overlaps supplies extra rigidity that an additive or multiplicative cocycle does not have. Arbitrary local sections need not glue after solving a correction problem: the corrections themselves may fail to have a common admissible support. That failure is the obstruction studied in [sections 13 and 14](#), and it is why arbitrary unions of supports are not permitted merely because a problem is stated locally.

3.2 Units, positive logarithms, and the integral subring

Proposition 3.4 (Exact unit criterion). *A nonzero $F = t^\lambda(f_\lambda + E) \in \mathcal{H}_\Gamma(U)$ on connected U , with $E \in \mathcal{H}_{\Gamma,+}(U)$, is a unit if and only if the ordinary function f_λ is nowhere zero on U . In that case*

$$F^{-1} = t^{-\lambda} f_\lambda^{-1} \sum_{k \geq 0} (-E/f_\lambda)^k, \quad (3.4)$$

and every unit has a unique factorization

$$F = t^\lambda h(1 + E'), \quad \lambda \in \Gamma, \quad h \in \mathcal{O}(U)^\times, \quad E' \in \mathcal{H}_{\Gamma,+}(U). \quad (3.5)$$

Proof. The valuation of a product is the sum of the leading exponents because $\mathcal{O}(U)$ is a domain; if $FG = 1$ the leading coefficient functions multiply to 1, so f_λ has no zeros. Conversely, dividing by the leading monomial and leading coefficient gives $1 + E/f_\lambda$ with positive support, and $\sum_{k \geq 0} (-E/f_\lambda)^k$ is strongly summable by [lemma 2.1](#) and inverts it. Uniqueness follows by comparing leading exponents and coefficients. $\square$

Proposition 3.5 (Positive-support exponential and logarithm). *The formulas*

$$\text{Exp}_H A = \sum_{k \geq 0} \frac{A^k}{k!}, \quad \text{Log}_H(1 + E) = \sum_{k \geq 1} \frac{(-1)^{k+1} E^k}{k} \quad (3.6)$$

define mutually inverse isomorphisms of abelian sheaves $(\mathcal{H}_{\Gamma,+}, +) \simeq (1 + \mathcal{H}_{\Gamma,+}, \cdot)$, with supports contained in $(\text{supp } A)^*$ and $(\text{supp } E)^* \setminus \{0\}$ respectively, the constant term of the exponential being treated separately. Consequently there are isomorphisms of abelian sheaves

$$\mathcal{H}_{\Gamma, \geq 0}^\times \simeq \mathcal{O}^\times \times \mathcal{H}_{\Gamma,+}, \quad (3.7)$$

$$\mathcal{H}_\Gamma^\times \simeq \underline{\Gamma} \times \mathcal{O}^\times \times \mathcal{H}_{\Gamma,+}, \quad (\lambda, h, A) \mapsto t^\lambda h \text{Exp}_H A, \quad (3.8)$$

where $\underline{\Gamma}$ is the locally constant sheaf with values in the additive group Γ .

Proof. All displayed terms have supports controlled by a positive monoid with finite word decompositions, so by [lemma 2.1](#) the formal characteristic-zero identities for exp and log may be evaluated and regrouped coefficientwise; they give the inverse identities and $\text{Exp}_H(A + B) = \text{Exp}_H(A) \text{Exp}_H(B)$ in the commutative ring $\mathcal{H}_\Gamma(U)$. For (3.8) apply [proposition 3.4](#) componentwise: on a connected patch every unit has the unique leading-term decomposition $t^\lambda f_\lambda \text{Exp}_H A$, multiplication adds λ , multiplies f_λ and adds A . A unit of $\mathcal{H}_{\Gamma, \geq 0}$ must have $\lambda = 0$, which gives (3.7). All constructions commute with restriction. $\square$

There is no selection of a global logarithm of an arbitrary ordinary holomorphic unit in this decomposition: that factor remains in $\mathcal{O}^\times$, and only the positive-support factor is logarithmized canonically. This logarithm is a logarithm of a positive-exponent perturbation of one; it has nothing to do with choosing a global branch of the ordinary logarithm, or with a global surcomplex exponential at infinite imaginary arguments. On $\mathbb{C}$ an ordinary nowhere-zero entire function is e^h for entire h determined up to $2\pi i\mathbb{Z}$, so a general global unit of $\mathcal{H}_\Gamma(\mathbb{C})$ is $t^\lambda e^h \text{Exp}_H A$.

Remark 3.6 (Reduction and the full coefficient sheaf). The reduction red of [section 1.2](#) is a surjective ring homomorphism $\mathcal{H}_{\Gamma, \geq 0} \rightarrow \mathcal{O}$ with kernel $\mathcal{H}_{\Gamma, +}$, so $\mathcal{H}_{\Gamma, \geq 0}/\mathcal{H}_{\Gamma, +} \cong \mathcal{O}$ as sheaves of rings. When $\Gamma \neq 0$ it does not extend to a ring homomorphism on $\mathcal{H}_{\Gamma}$: for $\delta \in \Gamma_{>0}$ the unit t^δ has reduction zero, whereas a unital ring map must send a unit to a unit. For $\Gamma = 0$, both sheaves are $\mathcal{O}$ and reduction is the identity. Below, a line bundle with transitions in $1 + \mathcal{H}_{\Gamma, +}$ has an integral model over $\mathcal{H}_{\Gamma, \geq 0}$ whose reduction is the trivial ordinary line bundle; *this statement does not assert a reduction map on the full ring $\mathcal{H}_{\Gamma}$, and reduction is always performed on the $\mathcal{H}_{\Gamma, \geq 0}$ model instead.*

4 The ordinary analytic input, with constructions

The following classical facts are the only places where analytic approximation occurs, and it occurs strictly at the level of one complex coefficient function. Including the constructions makes that separation visible. *These ordinary tools are classical and are not new results of this article.*

Lemma 4.1 (Scalar Weierstrass and Mittag–Leffler). *Let $A \subset \mathbb{C}$ be closed and discrete.*

1. *For positive integers m_c there is an entire f_0 whose zeros are precisely the points $c \in A$, with orders m_c .*
2. *For a finite Laurent principal part p_c prescribed at every $c \in A$ there is a meromorphic function on $\mathbb{C}$ with exactly those principal parts and no other poles.*

Proof. The finite case is immediate; in the infinite case enumerate the nonzero points as c_n with $|c_n| \rightarrow \infty$, treating a prescription at the origin by one separate factor or summand.

For (1) use the canonical factors $E_p(w) = (1 - w) \exp(w + w^2/2 + \cdots + w^p/p)$. On $|w| \leq 1/2$ the branch vanishing at the origin satisfies $\log E_p(w) = -\sum_{k>p} w^k/k$, so $|\log E_p(w)| \leq 2|w|^{p+1}$. Choose p_n with $m_{c_n} |\log E_{p_n}(z/c_n)| \leq 2^{-n}$ for $|z| \leq |c_n|/2$. The product of $E_{p_n}(z/c_n)^{m_{c_n}}$ then converges normally on compact sets once finitely many factors are separated, and off the specified zeros the tail is the exponential of a normally convergent sum, hence nonzero.

For (2) note that p_{c_n} is holomorphic on $|z| < |c_n|$. Choose a polynomial q_n approximating it within 2^{-n} on $|z| \leq |c_n|/2$ from its Taylor expansion there. Then $h = \sum_n (p_{c_n} - q_n)$ converges normally on compact subsets of $\mathbb{C} \setminus A$, and near any one point of A the remaining terms are holomorphic, so h has exactly the prescribed principal part. See also [\[10, Theorem 3.26\]](#) and [\[2\]](#). $\square$

Lemma 4.2 (Discrete Hermite interpolation). *For closed discrete $A \subset \mathbb{C}$ and finite $m_c \geq 1$ the simultaneous jet map*

$$R_0 : \mathcal{O}(\mathbb{C}) \longrightarrow V := \prod_{c \in A} \mathbb{C}[u]_{< m_c}, \quad R_0(f)_c = \sum_{j < m_c} \frac{f^{(j)}(c)}{j!} u^j, \quad (4.1)$$

is onto, and has a complex-linear right inverse $L : V \rightarrow \mathcal{O}(\mathbb{C})$.

Proof. Choose f_0 as in [lemma 4.1](#). Given $B_c \in \mathbb{C}[u]_{< m_c}$, take the principal part at c of $B_c(z-c)/f_0(z)$ and let h be a meromorphic function with exactly these principal parts. Then $f = f_0 h$ is entire and

$$f(c+u) - B_c(u) = f_0(c+u) \left(h(c+u) - \frac{B_c(u)}{f_0(c+u)} \right)$$

is holomorphically divisible by u^{m_c} , so $R_0 f = B$. A surjective linear map of complex vector spaces has a linear section, obtained by choosing a basis of the target and preimages of its members. $\square$

The operator L is not claimed to be continuous, unique or effective on arbitrary infinite input. What is used below is only the algebraic identity $R_0 L = \text{id}$, and the fact that extending L coefficientwise introduces no new Hahn exponent.

Lemma 4.3 (Scalar additive Cousin theorem on $\mathbb{C}$). *For any ordinary open cover (U_i) of $\mathbb{C}$ and any holomorphic additive cocycle b_{ij} there are $a_i \in \mathcal{O}(U_i)$ with $b_{ij} = a_i - a_j$. In particular $H^1(\mathbb{C}, \mathcal{O}) = 0$.*

Proof. A smooth partition of unity (ρ_k) subordinate to the cover, or to a locally finite refinement, gives smooth $v_i = \sum_k \rho_k b_{ik}$ with $v_i - v_j = b_{ij}$ for the convention $b_{ij} + b_{jk} = b_{ik}$. Their $\bar{\partial}v_i$ agree and define a global smooth $(0, 1)$ -form $a d\bar{z}$.

Every such form on $\mathbb{C}$ is $\bar{\partial}u$ for a global smooth u . Split $a = \sum_{n \geq 0} a_n$ by a locally finite smooth annular partition, each a_n compactly supported and, for large n , vanishing on $|z| < n - 1$. Its Cauchy transform

$$u_n(z) = \frac{1}{\pi} \int_{\mathbb{C}} \frac{a_n(\zeta)}{z - \zeta} dA(\zeta)$$

is smooth, satisfies $\bar{\partial}u_n = a_n$ and is holomorphic off $\text{supp } a_n$. For large n approximate the Taylor series of u_n at the origin by a polynomial p_n so that all derivatives of order at most n of $u_n - p_n$ are at most 2^{-n} on $|z| \leq n - 2$; no correction is needed for the finitely many small indices. Then $u = \sum_n (u_n - p_n)$ converges in every C^k norm on every compact set and satisfies $\bar{\partial}u = a$. Hence $a_i = v_i - u$ are holomorphic with the required differences. If a refinement was used, the local torsor sections still descend to a splitting on the original cover. The standard cocycle–torsor dictionary identifies this with $H^1(\mathbb{C}, \mathcal{O}) = 0$ [16, Tag 02FQ]. $\square$

We also use that every ordinary holomorphic line bundle on $\mathbb{C}$ is trivial. One route is the exponential sheaf sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathcal{O} \rightarrow \mathcal{O}^\times \rightarrow 1$: the preceding lemma gives $H^1(\mathbb{C}, \mathcal{O}) = 0$ and contractibility gives $H^2(\mathbb{C}, \mathbb{Z}) = 0$, hence $H^1(\mathbb{C}, \mathcal{O}^\times) = 0$; see [6]. Similarly $H^1(\mathbb{C}, \underline{G}) = 0$ for any constant abelian-group sheaf $\underline{G}$, in particular for $\underline{\Gamma}$. These are facts about the ordinary topological base, not about the fine topology of $\mathbf{SC}$.

Remark 4.4 (A support-safe implementation of the scalar step). When lemma 4.1(2) is applied once for every exponent γ of a well-ordered set S , the classical construction may be arranged as follows: exhaust $\mathbb{C}$ by closed disks whose boundary circles avoid A , group the finitely many new principal parts appearing at each stage, and subtract enough Taylor terms from each new rational principal part that its remaining tail is bounded by a prescribed summable positive real sequence on the preceding disk. The number of subtracted terms may depend arbitrarily on γ and on the spatial point. *None of these choices introduces a new Hahn exponent*, which is exactly why the construction preserves S , and no common real norm estimate across the coefficient functions is required. The same remark applies to lemmas 4.2 and 4.3.

5 Local preparation and deformed division, with uniform support

This section records the local input from [1] in the fixed- Γ form needed for infinitely many simultaneous prescriptions. The essential extra bookkeeping is that *one* positive support monoid controls all centres at once, even when the multiplicities are unbounded. Without that, local preparation would be a black box whose supports might depend uncontrollably on the centre, and no statement about an infinite family of centres could be made.

Throughout this section D is a fixed ordinary disk about the origin in the local coordinate u , and the ambient ring is the fixed-disk ring $\mathcal{H}_\Gamma(D) = \mathcal{O}(D)((t^\Gamma))$. For $h \in \mathcal{O}(D)$ put

$$R_m h = \sum_{j < m} \frac{h^{(j)}(0)}{j!} u^j, \quad D_m h = \frac{h - R_m h}{u^m},$$

so that both are holomorphic on the *same* disk, both are complex-linear, and

$$h = R_m h + u^m D_m h. \quad (5.1)$$

They act coefficientwise on Hahn data without increasing support.

5.1 Preparation by an exponent recursion

Theorem 5.1 (Uniform-support local preparation). *Let $F \in \mathcal{H}_\Gamma(U)$ be nonzero, with leading exponent $\lambda = v_H(F)$ and leading coefficient f_λ . Put*

$$T = (\text{supp } F - \lambda) \setminus \{0\} \subseteq \Gamma_{>0}, \quad M = T^*.$$

If f_λ has a zero of order $m \geq 1$ at $c \in U$, there is an ordinary disk D about the origin in $u = z - c$ on which

$$F(c + u) = t^\lambda q_0(u) P_c(u) Q_c(u), \quad (5.2)$$

where $q_0(u) = f_\lambda(c + u)/u^m$ is an ordinary holomorphic unit,

$$P_c(u) = u^m + \sum_{r < m} a_{c,r} u^r, \quad a_{c,r} \in \mathfrak{m}_\Gamma, \quad Q_c \in 1 + \mathcal{H}_{\Gamma,+}(D).$$

Every support $\text{supp}(a_{c,r})$ and $\text{supp}(Q_c - 1)$ lies in $M \setminus \{0\}$. The monoid M depends on F alone, not on c or m . The monic polynomial P_c and the normalized factor Q_c are unique.

Proof. Choose D so that q_0 has no zeros on it. Dividing by $t^\lambda q_0$ gives $u^m + E$ with $\text{supp } E \subseteq T$; division by a nonzero ordinary function changes no Hahn exponent, which is why the same T serves at every centre. Seek $P_c = u^m + A$ with $\deg_u A < m$ and $Q_c = 1 + B$ with positive supports. The equation is

$$E = A + u^m B + AB. \quad (5.3)$$

At $\nu \in M_{>0}$, after all smaller exponents have been treated, set

$$h_\nu = E_\nu - \sum_{\substack{\alpha+\beta=\nu \\ \alpha,\beta>0}} A_\alpha B_\beta, \quad A_\nu = R_m h_\nu, \quad B_\nu = D_m h_\nu. \quad (5.4)$$

The sum is finite by [lemma 2.1](#) and all its indices are strictly smaller than ν , so the recursion is legitimate and well founded on the well-ordered set $M_{>0}$. Identity (5.1) verifies (5.3) at each exponent, the coefficient functions are holomorphic on the same disk D , and the resulting supports lie in the fixed well-ordered M ; the finitely many coefficients of A have positive support and hence lie in $\mathfrak{m}_\Gamma$.

For uniqueness compare two normalized pairs and take the least exponent ν at which A or B differs, using the well-ordered union of the two candidate supports — so no common monoid need be assumed in advance. Products involving a positive exponent and a difference cannot contribute at ν , so $\Delta A_\nu + u^m \Delta B_\nu = 0$; applying R_m and D_m makes both terms zero, a contradiction. This is the preparation argument of [1, §7], with its centre-independent support bound made explicit. $\square$

Corollary 5.2 (All roots in a monad). *The zeros of $F^\sharp$ in the monad $\mu(c)$ are exactly c plus the roots of P_c , with their polynomial multiplicities, and their total multiplicity is $\text{ord}_c f_\lambda$. If $f_\lambda(c) \neq 0$ there are no zeros in that monad. When Γ is divisible all these roots lie in $c + \mathfrak{m}_\Gamma$.*

Proof. The evaluated factors q_0 and Q_c in (5.2) are nonzero throughout a small enough thickened disk: the relevant leading ordinary values are nonzero and Q_c is 1 plus an infinitesimal, and local nonvanishing units do not change zero orders. Every root ξ of a monic polynomial $u^m + \sum_{r < m} a_r u^r$ with infinitesimal lower coefficients is infinitesimal: otherwise $1/\xi$ is finite and

$$P_c(\xi)/\xi^m = 1 + \sum_{r < m} a_r \xi^{r-m}$$

is one plus an infinitesimal, hence nonzero. Algebraic closedness of **SC** then gives exactly m roots with multiplicity. If $f_\lambda(c) \neq 0$, evaluation at $c + \epsilon$ retains a nonzero leading value. The divisible case follows from algebraic closedness of K_Γ [15, Corollary 4]. $\square$

In particular any two monic infinitesimal cluster polynomials with the same surcomplex roots and multiplicities are equal. This elementary uniqueness is what identifies P_c intrinsically as the monic polynomial of the whole infinitesimal cluster rather than as a choice of labels for its roots, and it is used at every passage from a divisor to its coefficients.

5.2 Deformed division and uniform remainders

Lemma 5.3 (Deformed division). *Let $P = u^m + A$ with $\deg_u A < m$ and positive-exponent coefficients. For every $N \in \mathcal{H}_\Gamma(D)$ there are unique $J \in \mathcal{H}_\Gamma(D)$ and $R \in K_\Gamma[u]_{<m}$ with*

$$N = PJ + R. \quad (5.5)$$

If $S = \text{supp}(N)$ and E is a well-ordered positive set containing the supports of the coefficients of A , then J and R have support in $S + E^$. Explicitly, with $T(J) = D_m(AJ)$,*

$$J = \sum_{k \geq 0} (-T)^k D_m N, \quad R = R_m(N - AJ). \quad (5.6)$$

Proof. Applying D_m to (5.5) requires $(\text{id} + T)J = D_m N$, which (5.6) solves by the formal geometric identity; (5.1) then gives (5.5). Every application of T inserts one exponent from E together with a finite Taylor-division operation, so the support and finite-contribution assertions follow from lemma 2.1.

For uniqueness a difference satisfies $P \Delta J + \Delta R = 0$. If either is nonzero take the least exponent ν occurring in the two supports; multiplication by A cannot contribute at ν because it raises exponents and all earlier differences vanish, so $u^m \Delta J_\nu + \Delta R_\nu = 0$, and ordinary Taylor division forces both to vanish. $\square$

Write $\mathcal{R}_P N = R$ for the remainder; it is K_Γ -linear, and its correction to the ordinary Taylor remainder,

$$(\mathcal{R}_P - R_m)N = -R_m(AJ), \quad (5.7)$$

is supported in $S + E_{>0}^*$, every term containing at least one exponent from E . For a family (P_c) whose coefficient supports lie in the same E , all these conclusions hold with the same controlling set. Different m_c change the scalar operators but introduce no new exponent, and when the coefficients at all centres are viewed as one product vector the number of exponent decompositions is still finite; it is never necessary to sum over the centres.

Lemma 5.4 (Pole-free fractions). *Let $F, G \in \mathcal{H}_\Gamma(U)$ with $G \neq 0$. If the evaluated finite-meromorphic fraction F/G has no actual poles in $U^\sharp$, then after filling removable values it is a member of $\mathcal{H}_\Gamma(U)$.*

Proof. Where the leading coefficient of G is nonzero use its coherent inverse from [proposition 3.4](#). Near a leading zero, [theorem 5.1](#) writes G as a coherent unit times a monic P with infinitesimal roots; absorb the unit into F and divide, $N/P = J + R/P$ with $\deg R < \deg P$ by [lemma 5.3](#). Pole-freeness says R/P has no pole at any root of P , so P divides R over $\mathbf{SC}$, and the degree bound forces $R = 0$. This produces coherent representatives on ordinary neighbourhoods of every base point; they agree as fractions on overlaps and glue by [proposition 3.3](#). $\square$

Applying the lemma to both quotients shows that two nonzero coherent data with the same actual zero divisor differ by a coherent unit. That observation is used in [proposition 8.4](#), [corollary 10.3](#), and [theorem 11.1](#).

6 The ordinary-base stalks, and what they buy

Let $R_c = (\mathcal{H}_\Gamma)_c$ be the stalk at the *ordinary* base point c , formed by shrinking ordinary neighbourhoods. Such shrinking never separates two surcomplex points lying in the same monad. As fixed in [section 1.2](#), R_c is a filtered colimit of the fixed-disk rings $\mathcal{O}(D)((t^\Gamma))$ and the theorems below are about the colimit.

Theorem 6.1 (Principal-ideal-domain stalks). *For every $c \in \mathbb{C}$ the domain R_c is a principal ideal domain. Every nonzero element is associated to a unique monic polynomial in $u = z - c$ whose lower coefficients are infinitesimal. If $\Gamma \neq \{0\}$ then R_c is not a local ring. If Γ is divisible, its maximal ideals are precisely*

$$(u - \varepsilon), \quad \varepsilon \in \mathfrak{m}_\Gamma, \quad (6.1)$$

with residue fields K_Γ .

Proof. The ring is a domain because a nonzero coefficient function remains nonzero on smaller connected neighbourhoods. [Theorem 5.1](#) shows that every nonzero germ is a unit times a distinguished monic polynomial, a unit being represented by the polynomial 1; uniqueness follows from [corollary 5.2](#). Let $d(F)$ be the degree of that polynomial: a nonnegative integer vanishing exactly on units.

First take two nonzero germs F, G associated to $P, Q \in K_\Gamma[u]$ and let D be their monic greatest common divisor. The Euclidean algorithm in $K_\Gamma[u]$ gives $\mathcal{A}P + \mathcal{B}Q = D$ with $D \mid P$ and $D \mid Q$, so $(F, G) = (D)$ in R_c . Moreover D is distinguished: in an algebraically closed Hahn enlargement all its roots are roots of P , hence infinitesimal, and its lower coefficients are finite elementary symmetric sums of such roots. Thus $d(D) = \deg D$.

Now let I be any nonzero ideal and choose $F \in I \setminus \{0\}$ with $d(F)$ minimal. For every nonzero $G \in I$ the greatest common divisor just constructed lies in I and has degree at most $d(F)$, so by minimality its degree equals $d(F)$; the monic polynomial of F therefore divides that of G , and $F \mid G$ in R_c . Hence $I = (F)$. This treats arbitrary ideals, not merely finitely generated ones, and it uses only the Euclidean algorithm on *finite-degree* polynomials together with the least element of a nonempty set of nonnegative integers; no Noetherian property of an infinite coefficient ring is invoked.

If $\Gamma \neq 0$ choose $\delta \in \Gamma_{>0}$. Both u and $u - t^\delta$ are nonunits, their leading ordinary coefficient being u , while their difference t^δ is a unit; so the nonunits are not an ideal and R_c is not local.

If Γ is divisible, every distinguished polynomial factors into linear factors $u - \varepsilon$ with $\varepsilon \in \mathfrak{m}_\Gamma$, each a nonunit and irreducible by degree, so every nonzero maximal ideal of this principal ideal domain is generated by one of them. Conversely, evaluation at $c + \varepsilon$ is a surjection $R_c \rightarrow K_\Gamma$ because constants are available, and preparation with polynomial division identifies its kernel as $(u - \varepsilon)$. This proves (6.1) and the residue-field claim. $\square$

For $\Gamma = 0$ this is the familiar local ring of ordinary one-variable holomorphic germs. For nonzero Γ the extra maximal ideals record exactly the infinitesimally separated points that the ordinary base has collapsed into one point. *There is no contradiction with the fine-germ ring at a single surcomplex point considered in [1]: the two stalk constructions use different neighbourhoods.*

Proposition 6.2 (Non-locality, and the integral model). *If $\Gamma \neq 0$ the stalk $\mathcal{H}_{\Gamma,c} = R_c$ is not local. In contrast $\mathcal{H}_{\Gamma,\geq 0,c}$ is local, with residue field $\mathbb{C}$ and residue map*

$$\sum_{\gamma \geq 0} f_\gamma t^\gamma \mapsto f_0(c),$$

and as sheaves of rings $\mathcal{H}_{\Gamma,\geq 0}/\mathcal{H}_{\Gamma,+} \cong \mathcal{O}$.

Proof. Non-locality is the third paragraph of the proof of theorem 6.1. For $\mathcal{H}_{\Gamma,\geq 0}$, the displayed map is a surjective ring homomorphism. A germ outside its kernel has exponent-zero coefficient nonzero at c , hence nonzero on a smaller disk, and then the geometric inverse of (3.4) lies in $\mathcal{H}_{\Gamma,\geq 0}$ there; every germ in the kernel is a nonunit. So the kernel is the unique maximal ideal and the residue field is $\mathbb{C}$. Since $\text{red} : \mathcal{H}_{\Gamma,\geq 0} \rightarrow \mathcal{O}$ is already a surjective sheaf map with kernel $\mathcal{H}_{\Gamma,+}$ (remark 3.6), the final assertion follows. $\square$

This is the *integral model*: the nonnegative-support subsheaf is a genuinely locally ringed structure sheaf, and it is the object on which reduction to ordinary complex geometry is performed. The Laurent sheaf $\mathcal{H}_\Gamma$ is not.

Corollary 6.3 (Meaning of the Picard group over the plane). *For the ringed space $(\mathbb{C}, \mathcal{H}_\Gamma)$, every tensor-invertible sheaf of modules is locally free of rank one. Hence*

$$\text{Pic}(\mathbb{C}, \mathcal{H}_\Gamma) = \text{Pic}_{\text{lf}}(\mathbb{C}, \mathcal{H}_\Gamma) \simeq H^1(\mathbb{C}, \mathcal{H}_\Gamma^\times),$$

classified by unit-valued transition cocycles.

Proof. A tensor-invertible module over a commutative ring is a finitely generated projective module of rank one; at every stalk R_c it is free, by theorem 6.1. Let $\mathcal{L}$ have tensor inverse $\mathcal{M}$. Choose mutually dual generators at a stalk and represent them by local sections s of $\mathcal{L}$ and r of $\mathcal{M}$; after shrinking, their pairing is 1. At every nearby stalk both modules are free of rank one, and a pairing equal to 1 makes s a generator there. Hence multiplication by s is an isomorphism $\mathcal{H}_\Gamma \rightarrow \mathcal{L}$ on that neighbourhood. Local bases have unit transition functions, changing bases multiplies the cocycle by a coboundary, and gluing copies of $\mathcal{H}_\Gamma$ along a unit cocycle constructs the converse. Separately, for any sheaf of commutative rings $\mathcal{R}$ the same gluing gives

$$\text{Pic}_{\text{lf}}(\mathbb{C}, \mathcal{R}) = H^1(\mathbb{C}, \mathcal{R}^\times), \tag{6.2}$$

without any hypothesis on stalks. $\square$

Remark 6.4 (What the stalk theorem buys over the plane, and what is at stake without it). The stalk argument is not a formality. *For a general ringed space with nonlocal stalks, tensor-invertibility need not imply local generation by one section*; the distinction is explicitly noted in [16, Tag 09NT], and by [proposition 6.2](#) our stalks are exactly of that kind whenever $\Gamma \neq 0$. [Corollary 6.3](#) is what licenses dropping the qualification “locally free of rank one” from every Picard statement below.

One of the merged manuscripts declines that step. It proves [proposition 6.2](#), cites the same Stacks tag, and states its dichotomy only for $\mathrm{Pic}_{\mathrm{lf}}(\mathbb{C}, \mathcal{H}_\Gamma)$ and for $\mathrm{Pic}(\mathbb{C}, \mathcal{H}_{\Gamma, \geq 0})$ — writing that it *does not silently substitute the larger category of all tensor-invertible modules on an arbitrary ringed space*. We keep that distinction rather than delete it as superseded, and a third manuscript, written independently over a compact base, declines it again for the same reason and on the same authority: see [remark 18.5](#), where the compact classification is proved for $\mathrm{Pic}_{\mathrm{lf}}$ without a stalk hypothesis and the coordinate-chart consequence is recorded separately. That independent refusal is a confirmation of [proposition 6.2](#), not a competing claim. The honest summary is: (6.2) holds unconditionally; the identification of Pic with $\mathrm{Pic}_{\mathrm{lf}}$ on $(\mathbb{C}, \mathcal{H}_\Gamma)$ follows here from [theorem 6.1](#); and every theorem below stated for $\mathrm{Pic}(\mathbb{C}, \mathcal{H}_\Gamma)$ specializes verbatim to $\mathrm{Pic}_{\mathrm{lf}}(\mathbb{C}, \mathcal{H}_\Gamma)$ for a reader who declines [theorem 6.1](#).

Definition 6.5 (The line-bundle groups over the plane). A *Hahn-coherent line bundle* over the ordinary plane is a sheaf of $\mathcal{H}_\Gamma$ -modules locally isomorphic to $\mathcal{H}_\Gamma$; its class lies in $\mathrm{Pic}_{\mathrm{lf}}(\mathbb{C}, \mathcal{H}_\Gamma)$. We also use the ordinary Picard group of locally free rank-one $\mathcal{H}_{\Gamma, \geq 0}$ -modules; since $\mathcal{H}_{\Gamma, \geq 0}$ has local stalks by [proposition 6.2](#), that group coincides with its group of tensor-invertible modules and is written $\mathrm{Pic}(\mathbb{C}, \mathcal{H}_{\Gamma, \geq 0})$.

7 A sharp Hahn Mittag–Leffler theorem

7.1 Discrete principal-part data

Let $A = \{c_j : j \in J\} \subset \mathbb{C}$ be closed and discrete, with J finite or countably infinite. Choose pairwise disjoint ordinary disks U_j centred at c_j and set $U_0 = \mathbb{C} \setminus A$; the punctured plane U_0 is connected, the sets U_0, U_j cover $\mathbb{C}$, and the only nonempty overlaps between distinct members are the punctured disks $U_0 \cap U_j$. Prescribe, for each j ,

$$p_j(u) = \sum_{\gamma \in S_j} p_{j,\gamma}(u) t^\gamma, \quad p_{j,\gamma}(u) \in u^{-1}\mathbb{C}[u^{-1}], \quad u = z - c_j, \quad (7.1)$$

where $S_j \subseteq \Gamma$ is well ordered and zero principal parts are omitted. The pole order of $p_{j,\gamma}$ is finite but need not be bounded as j or γ varies. Each p_j is a legitimate section of $\mathcal{H}_\Gamma$ on its punctured disk.

Definition 7.1 (Coherent Mittag–Leffler solution). A *solution* for (7.1) is a section $B \in \mathcal{H}_\Gamma(U_0)$ such that $B - p_j$ extends to a section of $\mathcal{H}_\Gamma(U_j)$ for every j .

The definition imposes extension *coefficientwise across the ordinary centre*. It does not pretend that an arbitrary infinite family of meromorphic coefficients already has a finite moving-pole description on the omitted monad; the moving-pole version, where it does, is [theorem 11.1](#).

Theorem 7.2 (Uniform-support Mittag–Leffler criterion). *The principal-part data (7.1) have a coherent solution if and only if*

$$S := \bigcup_{j \in J} S_j \quad \text{is well ordered.} \quad (7.2)$$

When this holds a solution can be chosen with support contained in S , its coefficients being ordinary meromorphic functions on $\mathbb{C}$ with poles only in A . If $S \subseteq \Gamma_{>0}$, the solution and all holomorphic differences $B - p_j$ can be chosen with positive support. Any two solutions differ by a section of $\mathcal{H}_\Gamma(\mathbb{C})$.

Proof. Suppose $B = \sum_{\gamma \in T} b_\gamma t^\gamma$ is a solution. If some $\gamma \in S_j$ were absent from T , the coefficient of $B - p_j$ at γ would be the nonzero principal part $-p_{j,\gamma}$, which cannot extend holomorphically across c_j . Hence $S_j \subseteq T$ for every j , and a subset of a well-ordered set is well ordered, proving necessity of (7.2).

Conversely assume S well ordered. For each $\gamma \in S$ apply lemma 4.1(2) to the principal parts $p_{j,\gamma}$, taking zero where $\gamma \notin S_j$, and obtain an ordinary meromorphic b_γ on $\mathbb{C}$ with precisely those principal parts and no other poles. These choices are independent at different exponents and, by remark 4.4, introduce no new exponent. Then $B = \sum_{\gamma \in S} b_\gamma|_{U_0} t^\gamma$ is a section of $\mathcal{H}_\Gamma(U_0)$; every coefficient of $B - p_j$ extends holomorphically to U_j and the support is contained in S , so the extensions assemble into a valid section. If all exponents are positive so is the constructed solution.

Finally the difference of two solutions extends across each c_j , and the extensions agree on overlaps, so proposition 3.3 glues them to a section of $\mathcal{H}_\Gamma(\mathbb{C})$; no arbitrary union of the two solutions' local supports is taken in this step. $\square$

7.2 Why this is not a summability condition over the points

Condition (7.2) permits a fixed exponent at infinitely many points. For example $p_j = t/(z - j)$ is solvable: apply the ordinary Mittag–Leffler theorem once and multiply the result by t . The numerical family $(t, t, t, \dots)$ is not strongly summable, but no sum of that family is required — ordinary analytic summation across the spatial points takes place *within one coefficient function*.

In the opposite direction the data

$$p_j(z) = \frac{t^{1/j}}{z - j}, \quad j \geq 2, \quad (7.3)$$

are individually single-term Hahn sections and every finite subproblem is solvable, but their exponent union contains a strictly decreasing sequence and is not well ordered, so the infinite problem has no coherent solution. A common positive lower bound on valuations is insufficient: replacing $t^{1/j}$ by $t^{1+1/j}$ gives valuations greater than 1 and still violates (7.2). The obstruction concerns order structure throughout the support, not the smallest magnitude among the data.

Corollary 7.3 (Entire interpolation with an exact support criterion). *Let $a_j \in K_\Gamma$ be prescribed at a closed discrete set $\{c_j\}$. There exists $A \in \mathcal{H}_\Gamma(\mathbb{C})$ with $A(c_j) = a_j$ for all j if and only if $\bigcup_j \text{supp}(a_j)$ is well ordered; when that union is positive, A may be chosen in $\mathcal{H}_{\Gamma,+}(\mathbb{C})$. The same holds for finitely many prescribed derivatives at each point, the union being taken over all prescribed values.*

Proof. Necessity holds because evaluation at an ordinary point introduces no new exponent. For sufficiency interpolate the complex values, or the finite jets, at each exponent by lemma 4.2, and assemble over the given well-ordered union. $\square$

Corollary 7.3 is the undeformed case $P_c = u^{m_c}$ of the deformed interpolation theorem proved in section 10; it is recorded separately here because it is what section 12 needs, and because its proof requires nothing beyond lemma 4.2.

8 The global theorem for moving surcomplex divisors

8.1 Cluster polynomials and the correct compatibility invariant

Fix a closed discrete $A \subset \mathbb{C}$ and cluster data P_c as in (1.3). Every root of P_c in $\mathbf{SC}$ is infinitesimal, by the argument in corollary 5.2, and when Γ is divisible prescribing P_c is the same as prescribing an unordered finite multiset of elements of $\mathfrak{m}_\Gamma$. The zeros define an effective divisor on $\mathbb{C}^\sharp$: at the finite point $c + \xi$ take the multiplicity of ξ as a root of P_c . *Discreteness* means here that there are finitely many roots in each monad and only finitely many relevant monads above a compact ordinary set; fine-topological discreteness alone would be too weak.

Definition 8.1 (Symmetric support; admissibility). The symmetric support of the cluster data is $E(P)$ of (1.4), and the family is *admissible*, equivalently *uniformly supported*, when $E(P)$ is well ordered. The definition includes exact zeros $a_{c,j} = 0$, which contribute nothing to the union. For finitely many centres admissibility is automatic, a finite union of well-ordered sets being well ordered; for infinitely many centres it is a genuine restriction. The test is on the *symmetric coefficients*, not on separately chosen roots.

Theorem 8.2 (Sharp global moving-divisor criterion). *For cluster data (1.3) over a closed discrete $A \subset \mathbb{C}$, the following are equivalent.*

1. *The symmetric support $E(P)$ is well ordered.*
2. *There is a nonzero $F \in \mathcal{H}_\Gamma(\mathbb{C})$ whose complete zero divisor on $\mathbb{C}^\sharp$ consists exactly of the roots of $P_c(z - c)$ in the monads $\mu(c)$, $c \in A$, with their polynomial multiplicities.*
3. *For every ordinary entire f_0 whose zero divisor is $\sum_{c \in A} m_c[c]$ there is such an F with exponent-zero coefficient equal to f_0 , no negative exponents, and support contained in $M = E(P)^*$:*

$$F = f_0 + \sum_{\nu \in M_{>0}} f_\nu t^\nu \in \mathcal{H}_\Gamma(\mathbb{C}). \quad (8.1)$$

Equivalently, the prepared polynomial of F at every $c \in A$ is exactly P_c and its leading ordinary coefficient has no other zeros. On a fixed ordinary disk about each c the realizer factors as

$$F(c + u) = P_c(u)Q_c(u), \quad (Q_c)_0(u) = \frac{f_0(c + u)}{u^{m_c}}, \quad (8.2)$$

with Q_c a coherent unit supported in M . The ordinary function f_0 exists by the classical Weierstrass theorem, lemma 4.1(1).

Proof. Necessity, (2) $\Rightarrow$ (1). Let F be as in (2), with leading exponent λ and leading coefficient f_λ . By corollary 5.2 the zeros of f_λ are exactly the points of A , with orders m_c . Put $T = (\text{supp } F - \lambda) \setminus \{0\}$. At every c , theorem 5.1 constructs a monic polynomial whose lower coefficients have supports in $T^* \setminus \{0\}$; its roots and multiplicities are precisely those prescribed, so by the uniqueness after corollary 5.2 it equals P_c . Hence $E(P) \subseteq T^* \setminus \{0\}$, a well-ordered set, and every subset of a well-ordered set is well ordered. Multiplying F by an infinite or infinitesimal scalar cannot evade this: the normalization by t^λ has already removed that freedom.

Sufficiency, (1) $\Rightarrow$ (3). Assume $E = E(P)$ well ordered, put $M = E^*$, and fix f_0 as in (3). Shrink disjoint disks U_c so that $q_c(u) = f_0(c + u)/u^{m_c}$ is an ordinary holomorphic unit on each. On the puncture define

$$L_c(u) = \text{Log}_H \left(\frac{P_c(u)}{u^{m_c}} \right) = \sum_{k \geq 1} \frac{(-1)^{k+1}}{k} \left(\sum_{j < m_c} a_{c,j} u^{j-m_c} \right)^k. \quad (8.3)$$

Each L_c has support in $M \setminus \{0\}$, and at every exponent its coefficient is a *finite* polynomial in negative powers of u : finite decomposition in the positive monoid bounds the number of contributing exponent words by [lemma 2.1](#), and there are only finitely many indices j at each fixed c . Thus (8.3) is exactly a principal-part datum of the kind in [theorem 7.2](#), not an unrestricted Laurent germ.

Apply the positive-support half of [theorem 7.2](#) to all the L_c : it gives $B \in \mathcal{H}_{\Gamma,+}(U_0)$ with support in $M \setminus \{0\}$ such that $C_c = B - L_c \in \mathcal{H}_{\Gamma,+}(U_c)$. On the covering sets define

$$F_0 = f_0 \operatorname{Exp}_H B \text{ on } U_0, \quad F_c = q_c P_c \operatorname{Exp}_H C_c \text{ on } U_c. \quad (8.4)$$

The exponential law and $\operatorname{Exp}_H L_c = P_c/u^{m_c}$ show that these agree on every overlap, so by [proposition 3.3](#) they glue to $F \in \mathcal{H}_{\Gamma}(\mathbb{C})$, with support in M and exponent-zero coefficient f_0 . On $U_c^\sharp$ the factors q_c and $\operatorname{Exp}_H C_c$ are units, so the zeros and multiplicities there are exactly those of P_c , and (8.2) holds with $Q_c = q_c \operatorname{Exp}_H C_c$. On every other monad f_0 is nonzero, hence so is $F^\sharp$ by [corollary 5.2](#). Finally (3) $\Rightarrow$ (2) is immediate. $\square$

Remark 8.3 (A second, purely recursive construction of the realizer). Sufficiency can also be proved without logarithms or Mittag–Leffler, by a direct well-founded recursion on $M_{>0}$ which builds F and the local units $Q_c = \sum_{\nu \in M} q_{c,\nu} t^\nu$ simultaneously. Write $P_c = u^{m_c} + A_c$ and fix disks D_c on which $q_{c,0} = f_0(c+u)/u^{m_c}$ is nowhere zero. Suppose all coefficients below ν are chosen. The coefficient equation of (8.2) at ν is

$$f_\nu(c+u) = u^{m_c} q_{c,\nu}(u) + h_{c,\nu}(u), \quad h_{c,\nu} = \sum_{\substack{\alpha+\beta=\nu \\ \alpha>0, \beta\geq 0}} (A_c)_\alpha q_{c,\beta}, \quad (8.5)$$

where the correction $h_{c,\nu}$ is known: there are finitely many exponent pairs by [lemma 2.1](#) and every β is smaller than ν . Prescribe at every centre the finite ordinary jet $R_{m_c}(f_\nu(c+u)) = R_{m_c}h_{c,\nu}(u)$; [lemma 4.2](#) supplies one entire f_ν meeting all of these conditions at once, regardless of the number or the sizes of the jets. Then

$$q_{c,\nu}(u) = \frac{f_\nu(c+u) - h_{c,\nu}(u)}{u^{m_c}} \quad (8.6)$$

is holomorphic on the *same* disk D_c , since the numerator has the required vanishing order; no disk shrinking occurs during the recursion, which is where the common-domain hypothesis on the coefficient ring is used. All coefficients then live on the single well-ordered support M , and (8.5)–(8.6) prove the factorization exponent by exponent. This route is the one that generalizes in [section 10](#), since it produces the interpolation data directly.

8.2 Uniqueness modulo global units

Proposition 8.4 (The freedom in realization). *Two nonzero sections of $\mathcal{H}_{\Gamma}(\mathbb{C})$ have the same complete zero divisor on $\mathbb{C}^\sharp$ if and only if their quotient is a global unit. In particular all realizations in [theorem 8.2](#) differ by multiplication by a unit of $\mathcal{H}_{\Gamma}(\mathbb{C})$, that is, by*

$$t^\lambda e^h \operatorname{Exp}_H A, \quad \lambda \in \Gamma, \quad h \in \mathcal{O}(\mathbb{C}), \quad A \in \mathcal{H}_{\Gamma,+}(\mathbb{C}),$$

with h determined up to $2\pi i\mathbb{Z}$. When the exponent-zero leading coefficient is fixed to be the same f_0 , the unit is uniquely $\operatorname{Exp}_H A$ with $A \in \mathcal{H}_{\Gamma,+}(\mathbb{C})$, so the normalized realizers form a torsor under $1 + \mathcal{H}_{\Gamma,+}(\mathbb{C}) = \operatorname{Exp}_H \mathcal{H}_{\Gamma,+}(\mathbb{C})$.

Proof. One implication is [proposition 3.4](#). Conversely, both quotients are pole free, hence coherent by [lemma 5.4](#), and they are mutually inverse; so the quotient is a global unit. If both leading coefficients are f_0 at exponent zero, the ratio has leading monomial 1 and leading coefficient function 1, and [proposition 3.5](#) identifies it as $\text{Exp}_H A$. Multiplying any realizer by a unit changes no zero order, and [proposition 3.5](#) with $\mathcal{O}(\mathbb{C})^\times = e^{\mathcal{O}(\mathbb{C})}$ gives the displayed form of a general global unit. $\square$

8.3 Root supports versus symmetric supports

Suppose a cluster has roots $\varepsilon_{c,1}, \dots, \varepsilon_{c,m_c}$. Its coefficients are the elementary symmetric polynomials in those roots with alternating signs. A well-ordered union of all root supports is therefore *sufficient* for admissibility, because finite products use the positive monoid generated by that union. It is not *necessary*: symmetric cancellation can remove descending exponents from the coefficients even when they remain visible in every individual root.

This distinction is intrinsic, because a divisor is unordered. A root labelling, or the choice of one branch, can destroy precisely the cancellations that make a coherent global equation possible. The next section shows that this is not a marginal phenomenon.

9 Sharp examples and a nonfactorable divisor splitting

Throughout this section $t = \omega^{-1}$, and the displayed roots lie in $K_{\mathbb{Q}}$ even when their defining equations have integer Hahn exponents.

9.1 An impossible sequence of simple zeros

Example 9.1 (Individually valid motions with no exact global realization). For every integer $j \geq 2$ prescribe a single simple zero at $z_j = j + t^{1/j}$. The cluster polynomial is $P_j(u) = u - t^{1/j}$, so

$$E(P) = \{1/2, 1/3, 1/4, \dots\}$$

is not well ordered. By [theorem 8.2](#) there is no nonzero coherent entire datum having *exactly* these simple zeros and no other zeros on $\mathbb{C}^\sharp$. More generally the prescription $P_j(u) = u - a_j t^{1/j}$ with $a_j \in \mathbb{C}$ is realizable if and only if the sequence (a_j) has finite support, since an infinite subset of these decreasing exponents has no least element.

Every finite sub-divisor is realizable, even by a polynomial, so this is a genuinely infinite compatibility obstruction. It does *not* assert that the points z_j cannot all be among the zeros of a coherent entire datum; additional roots can restore the missing symmetric cancellations, as [example 9.2](#) shows.

9.2 Complete clusters with unbounded multiplicities

Example 9.2 (Uniform equations with nonuniform root scales). At the centre $j \geq 2$ prescribe

$$P_j(u) = u^j - t. \tag{9.1}$$

There is a single nonzero lower coefficient, with support $\{1\}$, so $E(P) = \{1\}$ and [theorem 8.2](#) produces a coherent entire datum with exactly the clusters

$$z_{j,k} = j + \zeta_j^k t^{1/j}, \quad 0 \leq k < j, \quad \zeta_j = e^{2\pi i/j}.$$

All roots are simple and the total multiplicity over the ordinary point j is j . The union of the root supports contains the descending sequence $(1/j)$, while the symmetric-coefficient support is the singleton $\{1\}$; indeed the defining datum can be chosen in $\mathcal{H}_{\mathbb{Z}, \geq 0}(\mathbb{C}) = \mathcal{O}(\mathbb{C})[[t]]$, although the roots require rational exponents and must be read in $K_{\mathbb{Q}}$ or in $\mathbf{SC}$.

This is an exact realization theorem for infinitely many clusters of unbounded size, not a finite-degree polynomial calculation in disguise. There is a relatively explicit construction. Let

$$f_0(z) = \prod_{j=2}^{\infty} \left[(1 - z/j) \exp(z/j + z^2/(2j^2)) \right]^j = \prod_{j=2}^{\infty} E_2(z/j)^j. \quad (9.2)$$

This ordinary Weierstrass product converges locally uniformly: on $|z| \leq R$ and $j > 2R$ the logarithm of its j th factor is bounded by $2j(R/j)^3 = 2R^3/j^2$. Its zeros are exactly the integers $j \geq 2$, with multiplicity j . On $U_0 = \mathbb{C} \setminus \{2, 3, \dots\}$ put

$$b_k(z) = -\frac{1}{k} \sum_{j=2}^{\infty} \frac{1}{(z-j)^{jk}}, \quad B = \sum_{k \geq 1} b_k(z) t^k. \quad (9.3)$$

For each fixed k the ordinary sum over j converges locally uniformly away from the indicated points: for large j on a compact set $|z-j| \geq j/2$, so the terms are bounded by $(2/j)^{jk}$. The principal part of b_k at j is exactly $-1/[k(z-j)^{jk}]$, so

$$B - \text{Log}_H(1 - t/(z-j)^j)$$

has holomorphic coefficients at j , and the section $f_0 \text{Exp}_H B$ on U_0 extends coherently to all of $\mathbb{C}$ by (8.4), with local polynomial $u^j - t$.

Equations (9.2) and (9.3) involve *ordinary* convergence inside single coefficient functions; the subsequent sum over k is an exact Hahn construction. These are two different operations, justified separately, and the proof never takes an infinite Hahn product of the cluster factors over j . An unqualified infinite product over the centres would obscure exactly that distinction.

9.3 A principal effective divisor with a nonprincipal sub-divisor

Corollary 9.3 (Failure of arbitrary global divisor splitting). *Let F be a coherent entire datum realizing [example 9.2](#), and select the root $j + t^{1/j}$ from each cluster. There is no factorization $F = GH$ by coherent entire data such that G has exactly the selected roots and H exactly the remaining roots, with the indicated multiplicities. Both prescribed sub-divisors are locally coherent, but neither is globally principal.*

Proof. The selected divisor is nonprincipal by [example 9.1](#). For the complement write $a = t^{1/j}$; its local polynomial is

$$Q_j(u) = \frac{u^j - a^j}{u - a} = u^{j-1} + au^{j-2} + \dots + a^{j-1},$$

whose coefficient of u^{j-2} is $t^{1/j}$, so its symmetric support union contains $\{1/j : j \geq 2\}$ and is not well ordered. In fact that union is $\{k/j : j \geq 2, 1 \leq k < j\} = \mathbb{Q} \cap (0, 1)$. The necessity half of [theorem 8.2](#) excludes a global realization. Local equations exist on each U_j and are units on the punctured overlaps, so the sub-divisors are locally coherent; a factorization with the prescribed zero divisors would contradict their nonprincipality. $\square$

This does *not* claim that F is irreducible, that every factorization of F fails, or that the global ring has any particular unique-factorization property. The assertion concerns this specific, intrinsically meaningful partition of one effective divisor.

9.4 No repair by enlarging the value group

The nonexistence in [example 9.1](#) persists after any set-sized enlargement of Γ inside **No**, and even for a putative section with an arbitrary set-sized surreal support: a larger ambient ordered group does not make the strictly decreasing sequence $1/2, 1/3, \dots$ well ordered, and the necessity proof forces that same sequence into one alleged support monoid. The obstruction is therefore also a nonexistence statement in the full set-supported Hahn-coherent category of [1]. By contrast, finite truncation of the *spatial* set removes the obstruction entirely. These two ways of enlarging or approximating the problem must not be confused.

10 A support-restricted Chinese remainder theorem

[Theorem 8.2](#) says when a divisor exists. It does not say what can be prescribed *on* that divisor. The answer is not the full product of the local finite algebras: the correct target is a support-restricted subalgebra, and the restriction is forced by the existence of a single coherent source datum rather than chosen as a convenient topology.

10.1 The correct target algebra

Assume the family $P = (P_c)_{c \in A}$ is admissible and set $E = E(P)$, $M = E^*$. Define

$$\mathcal{B}(P) = \left\{ (B_c)_{c \in A} : \begin{array}{l} B_c \in K_\Gamma[u]_{< m_c}, \\ \bigcup_{c \in A} \text{supp}(B_c) \text{ is well ordered} \end{array} \right\}, \quad (10.1)$$

where $\text{supp}(B_c)$ means the union of the supports of the finitely many polynomial coefficients of B_c . Multiplication is performed at each centre modulo P_c , taking the unique representative of degree $< m_c$ supplied by [lemma 5.3](#).

Proposition 10.1 (Restricted product algebra). *Formula (10.1) defines a K_Γ -algebra. For families supported in S and T the product is supported in $S + T + M$. The simultaneous deformed remainder map*

$$\mathcal{R}_P : \mathcal{H}_\Gamma(\mathbb{C}) \longrightarrow \mathcal{B}(P), \quad F \longmapsto (\mathcal{R}_{P_c}(F(c + u)))_{c \in A} \quad (10.2)$$

is a homomorphism of K_Γ -algebras.

Proof. Addition has support in the finite union of the two supports. For multiplication apply [lemma 5.3](#) to $B_c C_c$ at each centre: the input support lies in $S + T$ and all the divisions use the same positive well-ordered set E , so the output support lies in $S + T + M$, which is well ordered by [lemma 2.1](#). Uniqueness of division makes the ordinary quotient-ring identities valid. The same lemma shows that the image of (10.2) has a common well-ordered support, contained in $\text{supp}(F) + M$, and uniqueness makes $\mathcal{R}_P$ multiplicative and K_Γ -linear. $\square$

10.2 A support-controlled right inverse

Theorem 10.2 (Deformed Hermite interpolation). *For an admissible cluster family, prescribed polynomial remainders $B_c \in K_\Gamma[u]_{<m_c}$ are simultaneously realized by a member of $\mathcal{H}(\mathbb{C})$ if and only if their total coefficient support is well ordered. Thus (10.2) is surjective. If that support is S , a realizing datum can be chosen with support in $S + E^*$. Explicitly, let L be a complex-linear right inverse as in lemma 4.2, extended coefficientwise, let R_0 be the jet map (4.1) extended coefficientwise, and put $N = \mathcal{R}_P - R_0$. Then a K_Γ -linear right inverse of $\mathcal{R}_P$ is*

$$\mathcal{I}_P(B) = L \sum_{k \geq 0} (-NL)^k B. \quad (10.3)$$

Proof. Necessity is immediate from uniform deformed division: if F has support S_F then all its remainders lie in $S_F + E^*$ by lemma 5.3, a well-ordered set.

For sufficiency identify (10.1), as a vector space, with $V((t^\Gamma))$ for V the product in (4.1). A coefficient at one exponent is one element of that product; no sum over the centres is being taken. The scalar operator L extends to $V((t^\Gamma))$ without increasing support, and it is K_Γ -linear because multiplication by a fixed Hahn constant is finite convolution at each exponent.

By (5.6) and (5.7), every term of N inserts a nonempty finite word of exponents from E , so NL strictly raises the support by such words. In (10.3) a contribution from $(NL)^k$ is indexed by a term of S together with a word from E partitioned into k nonempty consecutive blocks. For a fixed output exponent lemma 2.1 leaves finitely many such words; each word has finite length and finitely many block partitions, so only finitely many k contribute. This is strong summability, not merely formal plausibility of a Neumann series, and the support lies in $S + E^*$.

Let $C = \sum_{k \geq 0} (-NL)^k B$, so that $(\text{id} + NL)C = B$. Since $R_0 L = \text{id}$,

$$\mathcal{R}_P(LC) = (R_0 + N)LC = (\text{id} + NL)C = B.$$

This proves existence, the support estimate and the stated right inverse. $\square$

The proof is valid for arbitrary ordered groups. It never assumes an Archimedean value group, a discrete valuation, a norm estimate for L , or a uniform bound on the multiplicities m_c .

Corollary 10.3 (Chinese remainder quotient). *Let G be any realizer of the admissible cluster divisor. Then*

$$\mathcal{H}_\Gamma(\mathbb{C})/(G) \cong \mathcal{B}(P) \quad (10.4)$$

as K_Γ -algebras, and the quotient map admits a noncanonical K_Γ -linear section.

Proof. Surjectivity and the linear section are theorem 10.2. If every local remainder of F vanishes then F vanishes to at least the multiplicity of every zero of G , so F/G has no actual pole and is coherent by lemma 5.4; hence the kernel is contained in (G) . The reverse inclusion follows from $G = P_c Q_c$ locally, (8.2). Apply the first isomorphism theorem. $\square$

This is an infinite Chinese remainder statement with a precisely identified image. *Replacing $\mathcal{B}(P)$ by the full product $\prod_c K_\Gamma[u]/(P_c)$ would be false, and the next example shows that the failure occurs already for prescribed values in $\{0, 1\}$.*

10.3 Bounded raw values need not be interpolable

Example 10.4 (Selecting one root in every growing cluster). Work in $\Gamma = \mathbb{Q}$ with the admissible family $P_n(u) = u^n - t$, $n \geq 1$, and write $s_n = t^{1/n}$. At the n roots of P_n prescribe the value 1 at s_n and the value 0 at each $\zeta_n^k s_n$, $1 \leq k < n$. The unique polynomial of degree $< n$ with those values is

$$B_n(u) = \frac{1}{n} \sum_{j=0}^{n-1} (u/s_n)^j = \frac{1}{n} \sum_{j=0}^{n-1} t^{-j/n} u^j, \quad (10.5)$$

since the finite geometric sum equals n at the root 1 of unity and 0 at the others. Its coefficient supports contain

$$-\frac{n-1}{n} = -1 + \frac{1}{n}, \quad n \geq 2,$$

an infinite strictly decreasing sequence, so [theorem 10.2](#) rules out a global coherent interpolant.

Every prescribed scalar value here was 0 or 1, and the underlying zero divisor is realizable even with integer Hahn exponents by [example 9.2](#). The obstruction appears only after converting point values into the intrinsic finite quotient coordinates. Every finite collection of these root-selection conditions is solvable. If repeated roots occur, values are replaced by finite jets at those roots and the same principle applies: solve the ordinary finite-dimensional polynomial Chinese remainder problem over a splitting field to obtain B_c , then test its global coefficient support. Raw jet coordinates at infinitesimally close points can conceal arbitrarily large negative powers of their separations.

11 A Mittag–Leffler theorem for moving poles

[Theorem 7.2](#) prescribes principal parts at the ordinary centres. We now prescribe them at the *moving* infinitesimal poles, relative to an admissible pole bound.

Keep an admissible family (P_c) . A prescribed moving principal part at c is a proper rational function

$$\frac{R_c(u)}{P_c(u)}, \quad R_c \in K_\Gamma[u]_{< m_c}. \quad (11.1)$$

Two local fractions have the same such principal part when their difference extends to a coherent holomorphic datum on a disk about the ordinary centre. By [lemma 5.3](#) every local coherent fraction whose poles are bounded by P_c has a unique representative of the form (11.1); this representative packages all the individual poles and residues of the cluster, including cancellations. A *global coherent meromorphic function* means here one fraction F/G with $F, G \in \mathcal{H}_\Gamma(\mathbb{C})$, $G \neq 0$, interpreted as finite-meromorphic germs with removable values filled. *We do not assume that an arbitrary componentwise meromorphic Hahn family belongs to this fraction field.*

Theorem 11.1 (Sharp moving-pole Mittag–Leffler theorem). *For an admissible pole bound (P_c) , the moving principal parts (11.1) are realized by a global coherent meromorphic fraction with no poles outside that bound if and only if*

$$S_R = \bigcup_{c \in A} \text{supp}(R_c) \quad (11.2)$$

is well ordered. Any two solutions differ by a member of $\mathcal{H}_\Gamma(\mathbb{C})$.

Proof. Choose the divisor realizer G of [theorem 8.2](#), so that on the disks D_c one has $G(c+u) = P_c(u)Q_c(u)$ with Q_c and Q_c^{-1} uniformly supported in $E(P)^*$ and with ordinary unit leading coefficient functions.

Suppose S_R is well ordered. Form the targets $B_c = \mathcal{R}_{P_c}(Q_c R_c)$; their total support lies in $S_R + E(P)^*$ by [proposition 10.1](#). By [theorem 10.2](#) choose $F \in \mathcal{H}_\Gamma(\mathbb{C})$ with those local remainders. Then $F - Q_c R_c$ is coherently divisible by P_c , and

$$\frac{F(c+u)}{G(c+u)} - \frac{R_c(u)}{P_c(u)} = \frac{F(c+u) - Q_c(u)R_c(u)}{P_c(u)Q_c(u)}$$

is holomorphic on the thickened disk. The fraction F/G has no other poles because the zeros of G are exactly the specified bound.

Conversely let a global fraction H have the required principal parts and no poles outside the bound. Then GH is pole free, hence equal to some $F \in \mathcal{H}_\Gamma(\mathbb{C})$ by [lemma 5.4](#), and locally $R_c = \mathcal{R}_{P_c}(F(c+u)/Q_c(u))$. Uniform inversion and deformed division put all these numerators in $\text{supp}(F) + E(P)^*$, so S_R is well ordered. Finally the difference of two solutions has no actual poles and is coherent by the same lemma. $\square$

The pole bound is part of the theorem. Without its admissibility one cannot assume that a single global denominator exists at all; [theorem 8.2](#) is the exact obstruction to that preliminary step.

Example 11.2 (Globally realizable poles with unbounded residue scales). Let $P_n(u) = u^n - t$ and prescribe $R_n(u) = 1$. Both the divisor support and the numerator support are admissible, so there is a global coherent meromorphic fraction locally congruent to $1/((z-n)^n - t)$ modulo coherent holomorphic data. At a root $\xi = \zeta_n^k s_n$ its residue is

$$\frac{1}{n\xi^{n-1}} = \frac{\zeta_n^k}{n} t^{-(n-1)/n}, \quad (11.3)$$

and the union of these individual residue supports is not well ordered. This does not prevent the global fraction from existing: its proper numerators are all equal to 1. For $n > 1$ the residues within the n th cluster sum to zero, while the $n = 1$ cluster contributes 1.

On an ordinary compact contour avoiding the leading denominator zeros, the moving-pole residue theorem of [1] applies to only finitely many clusters and returns the sum of their actual residues. *Neither the present theorem nor that contour theorem calls for an infinite sum of residues over the plane.* The example shows why the correct global input consists of proper rational cluster representatives rather than independently enumerated residue coefficients.

12 Near-identity automorphisms and interpolation of motions

For simple zero motions there is an alternative constructive mechanism, which also shows how rich the coherent category remains despite the obstruction theorems.

Theorem 12.1 (Global inverse for positive-support perturbations). *Let $U \subseteq \mathbb{C}$ be an ordinary domain and $E \in \mathcal{H}_{\Gamma,+}(U)$. The coherent map $G(z) = z + E(z)$ evaluates to an automorphism of $U^\sharp$. Its inverse is $H(z) = z + \mathcal{A}(z)$ for a unique $\mathcal{A} \in \mathcal{H}_{\Gamma,+}(U)$, and both compositions are identities as Hahn-coherent data. If $T = \text{supp } E$ then $\text{supp}(\mathcal{A}) \subseteq T^* \setminus \{0\}$.*

Proof. The standard part of $G(c + \epsilon)$ is c , so evaluation maps $U^\sharp$ into itself. To construct a right inverse, solve

$$\mathcal{A}(z) = -E(z + \mathcal{A}(z)) = - \sum_{\gamma \in T} \sum_{k \geq 0} \frac{e_\gamma^{(k)}(z)}{k!} t^\gamma \mathcal{A}(z)^k \quad (12.1)$$

by well-founded recursion on $T^* \setminus \{0\}$. At ν , the terms with $k \geq 1$ involve only coefficients of $\mathcal{A}$ at exponents strictly smaller than ν , because the external exponent γ is strictly positive, and the term with $k = 0$ is already known. Finite word decomposition (lemma 2.1) makes each coefficient formula finite, so the recursion defines holomorphic coefficients on the same U with the asserted support, and $G \circ H = \text{id}$ as coherent data by (3.3).

For cancellation, suppose two positive perturbations H_1, H_2 differ and let ν be the least exponent of their difference. In

$$G \circ H_1 - G \circ H_2 = (H_1 - H_2) + (E \circ H_1 - E \circ H_2)$$

the second summand has no term at ν , since every term containing a difference also carries an external positive exponent from E ; so the coefficient at ν of the displayed difference is nonzero, and left composition by G is injective in this class. Associativity of (3.3) then gives

$$G \circ (H \circ G) = (G \circ H) \circ G = G = G \circ \text{id},$$

whence $H \circ G = \text{id}$. The same least-exponent argument proves uniqueness among *all* positive-support inverses, not only those supported in T^* . $\square$

Corollary 12.2 (Exact interpolation of simple motions). *Let $\{c_j\}$ be closed discrete and prescribe $\delta_j \in \mathfrak{m}_\Gamma$. There is a coherent automorphism $G = \text{id} + E$ with $E \in \mathcal{H}_{\Gamma,+}(\mathbb{C})$ satisfying $G(c_j) = c_j + \delta_j$ for every j if and only if $\bigcup_j \text{supp}(\delta_j)$ is well ordered.*

Proof. Necessity follows by evaluating E at ordinary points. For sufficiency use corollary 7.3 to find $E \in \mathcal{H}_{\Gamma,+}(\mathbb{C})$ with $E(c_j) = \delta_j$ and apply theorem 12.1. $\square$

If f_0 has simple zeros exactly at the c_j , then $f_0 \circ G^{-1}$ realizes the moved simple divisor. This is consistent with, but strictly weaker than, theorem 8.2: a bijective change of coordinate cannot split a multiple zero into distinct roots, whereas the moving-divisor theorem permits precisely that splitting and allows arbitrarily large cluster degrees.

For example $G(z) = z + tz^2$ is a coherent automorphism of the finite surcomplex plane, although the same quadratic is not injective on all of $\mathbf{SC}$: its other algebraic preimage is generally infinite and lies outside $\mathbb{C}^\sharp$. Its inverse on $\mathbb{C}^\sharp$ is the positive-support branch

$$G^{-1}(z) = z - tz^2 + 2t^2z^3 - 5t^3z^4 + 14t^4z^5 - \dots,$$

whose coefficients are the Catalan numbers with alternating signs. This domain distinction is another reason not to identify coherent entire data on the finite plane with all-scale entire functions on the full class field.

13 The exact additive Cousin obstruction

The existence theorems above use a common support hypothesis and show it is sharp. We now identify the gluing obstruction directly, in a form that is invisible to every ordinary coefficientwise existence theorem.

13.1 A puncture-and-disk cover, and the singular support

Consider the ordinary cover

$$U_0 = \mathbb{C} \setminus \mathbb{N}_{>0}, \quad U_n = D(n, 1/4) \quad (n \geq 1). \quad (13.1)$$

The disks are disjoint, so the only nonempty pairwise intersections involving different patches are $W_n = U_0 \cap U_n = U_n \setminus \{n\}$, and there are no triple overlaps to check. Any family $b_n \in \mathcal{H}_\Gamma(W_n)$ is therefore additive Cousin data, with splitting convention

$$b_n = a_n - a_0 \quad \text{on } W_n. \quad (13.2)$$

Each b_n separately has well-ordered support; their union need not. Write $b_n = \sum_\gamma b_{n,\gamma} t^\gamma$, absent coefficients being zero, and define the *singular support*

$$\Sigma(b) = \{\gamma \in \Gamma : \text{some } b_{n,\gamma} \text{ has a nonremovable singularity at } n\}. \quad (13.3)$$

The singularity may be a pole or an ordinary essential singularity. This definition concerns ordinary coefficient functions; it is not about extending a coefficient's essential singularity into the infinitesimal monad.

Theorem 13.1 (Necessary and sufficient Cousin criterion). *For the cover (13.1), the data (b_n) admit a splitting (13.2) with $a_0 \in \mathcal{H}_\Gamma(U_0)$ and $a_n \in \mathcal{H}_\Gamma(U_n)$ if and only if $\Sigma(b)$ is well ordered. If all b_n have strictly positive support, the same condition is equivalent to a splitting with all $a_i \in \mathcal{H}_{\Gamma,+}(U_i)$.*

Proof. Necessity. If $b_{n,\gamma}$ has a nonremovable singularity at n then $(a_0)_\gamma$ cannot be identically zero on U_0 : otherwise $b_{n,\gamma} = (a_n)_\gamma$ would extend across n . Hence $\Sigma(b) \subseteq \text{supp}(a_0)$, which is well ordered.

Sufficiency. Fix $\gamma \in \Sigma(b)$. The scalar functions $b_{n,\gamma}$ form ordinary additive Cousin data, so lemma 4.3 gives $a_{0,\gamma} \in \mathcal{O}(U_0)$ such that every $b_{n,\gamma} + a_{0,\gamma}$ extends holomorphically across n . For $\gamma \notin \Sigma(b)$ set $a_{0,\gamma} = 0$, the corresponding $b_{n,\gamma}$ already extending. Let $a_{n,\gamma}$ be those extensions and assemble

$$a_0 = \sum_{\gamma \in \Sigma(b)} a_{0,\gamma} t^\gamma, \quad a_n = \sum_{\gamma} a_{n,\gamma} t^\gamma.$$

Then $\text{supp}(a_0) \subseteq \Sigma(b)$ and, for each fixed n , $\text{supp}(a_n) \subseteq \Sigma(b) \cup \text{supp}(b_n)$, a finite union of well-ordered sets. So every displayed expression is a legitimate coherent section and the coefficient equations give (13.2). If the input supports are positive then both sets in the last inclusion are positive, and the constructed splitting is positive. $\square$

A union of bad supports is therefore not, by itself, the obstruction. Exponents attached only to *removable* coefficient germs may stay on the separate disk patches; the exact obstruction is the support that must be carried by the single connected central patch.

Remark 13.2 (An explicit scalar solution on this cover). The scalar step can be written without sheaf cohomology. For fixed γ take the negative Laurent part p_n of $b_{n,\gamma}$ at n . Since its inner Laurent radius is zero, p_n is holomorphic on all of $\mathbb{C} \setminus \{n\}$ even when it is an infinite negative Laurent series. Choose a polynomial q_n with $|p_n - q_n| \leq 2^{-n}$ on $|z| \leq n/2$. Then

$$a_{0,\gamma} = - \sum_{n \geq 1} (p_n - q_n)$$

converges normally on compact subsets of U_0 and cancels every prescribed negative Laurent part, all other terms extending holomorphically at each centre. This is the correcting-polynomial proof of the required scalar Cousin step, and it applies verbatim when the negative part is an infinite series.

13.2 An explicit infinite-dimensional family of obstructions

Fix the discrete centres $c_j = j$, $j \geq 2$, with the cover of [section 7](#), and define complex vector spaces

$$\mathcal{V}_\Gamma = \prod_{j \geq 2} \mathfrak{m}_\Gamma, \quad (13.4)$$

$$\mathcal{W}_\Gamma = \left\{ (a_j) \in \mathcal{V}_\Gamma : \bigcup_{j \geq 2} \text{supp}(a_j) \text{ is well ordered} \right\}. \quad (13.5)$$

The second is a vector subspace, since finite unions of well-ordered sets are well ordered and addition can only remove exponents from such a union. *No Hahn summation over j is intended in the product (13.4).* An element $a = (a_j)$ determines an additive cocycle on the cover by

$$p_{0j}(z) = \frac{a_j}{z - j}, \quad p_{j0} = -p_{0j}. \quad (13.6)$$

Theorem 13.3 (A concrete obstruction subspace). *The construction (13.6) induces an injective complex-linear map*

$$\mathcal{V}_\Gamma / \mathcal{W}_\Gamma \hookrightarrow H^1(\mathbb{C}, \mathcal{H}_{\Gamma,+}). \quad (13.7)$$

Exponentiating the transition functions gives an injective group homomorphism

$$\mathcal{V}_\Gamma / \mathcal{W}_\Gamma \hookrightarrow \text{Pic}(\mathbb{C}, \mathcal{H}_\Gamma), \quad a \longmapsto \mathcal{L}(a), \quad (13.8)$$

where $\mathcal{L}(a)$ is glued by

$$g_{0j}(z) = \text{Exp}_H \left(\frac{a_j}{z - j} \right) \in 1 + \mathcal{H}_{\Gamma,+}(U_0 \cap U_j). \quad (13.9)$$

Its transitions lie in $1 + \mathcal{H}_{\Gamma,+}$, so $\mathcal{L}(a)$ has an integral model over $\mathcal{H}_{\Gamma,\geq 0}$ with trivial ordinary reduction; nevertheless $\mathcal{L}(a)$ is nontrivial exactly when $a \notin \mathcal{W}_\Gamma$.

Proof. An additive coboundary representation on this cover is a system $p_{0j} = B_0 - B_j$ with $B_0 \in \mathcal{H}_{\Gamma,+}(U_0)$ and $B_j \in \mathcal{H}_{\Gamma,+}(U_j)$, which is exactly the positive-support Mittag–Leffler problem for the principal parts $a_j/(z - j)$. By [theorem 7.2](#) it is solvable precisely when the support union is well ordered, so the kernel of the map from $\mathcal{V}_\Gamma$ is $\mathcal{W}_\Gamma$. Equivalently, the singular support of this cocycle is $\bigcup_j \text{supp}(a_j)$, and [theorem 13.1](#) gives the same conclusion.

This concerns the actual first cohomology class, not an artifact of one cover. An additive cocycle defines a sheaf torsor; if its class vanishes after refinement then the torsor has a global section, and expressing that section in the original local trivializations supplies a coboundary on the original cover [[16](#), Tag 02FQ]. No refinement can remove the obstruction without solving the stated Mittag–Leffler problem.

Nontriviality after exponentiation can also be checked directly. Suppose the line bundle with transitions (13.9) is trivial, so there are units $b_0 \in \mathcal{H}_\Gamma(U_0)^\times$, $b_j \in \mathcal{H}_\Gamma(U_j)^\times$ with $g_{0j} = b_0/b_j$ on the punctures. Write their leading factors as in (3.5). Since every g_{0j} has leading term 1, all leading exponents equal one λ and all leading ordinary coefficient functions agree on overlaps, so they glue to one nowhere-zero entire h . Dividing each b_i by $t^\lambda h|_{U_i}$ normalizes every unit into $1 + \mathcal{H}_{\Gamma,+}(U_i)$, and applying Log_H produces the forbidden additive coboundary unless $a \in \mathcal{W}_\Gamma$. Conversely, exponentiating an additive solution trivializes the bundle, and the exponential law makes tensor product correspond to addition of sequences, which gives (13.8). Finally the same transitions define a locally free rank-one $\mathcal{H}_{\Gamma,\geq 0}$ -module whose reduction is 1 on every patch, which is the integral-model statement of [remark 3.6](#). $\square$

Corollary 13.4 (First cohomology detected by descending scales). *Suppose Γ contains a strictly decreasing sequence $\gamma_1 > \gamma_2 > \dots > 0$. Then the cocycles*

$$b_n(a) = \frac{a_n t^{\gamma_n}}{z - n}, \quad a = (a_n) \in \mathbb{C}^{\mathbb{N}_{>0}}, \quad (13.10)$$

induce an injective complex-linear map

$$\mathbb{C}^{\mathbb{N}_{>0}} / \mathbb{C}^{(\mathbb{N}_{>0})} \hookrightarrow H^1(\mathbb{C}, \mathcal{H}_{\Gamma,+}), \quad (13.11)$$

where $\mathbb{C}^{(\mathbb{N}_{>0})}$ denotes the finitely supported sequences.

Proof. Each component of (13.10) is a legitimate positive Hahn section with one-element support, and its singular support is $\{\gamma_n : a_n \neq 0\}$. An infinite subset of a strictly decreasing sequence indexed by positive integers has no least element, so by [theorem 13.1](#) — or by the special case $a_j \mapsto a_j t^{\gamma_j}$ of [theorem 13.3](#) — the cocycle is a coboundary exactly when (a_n) is finitely supported. Linearity is immediate. $\square$

Corollary 13.5 (Size and local invisibility). *The space in (13.11) has complex dimension $2^{\aleph_0}$. Every individual class constructed there restricts to zero on every bounded ordinary disk, although globally it need not vanish.*

Proof. There are only continuum many complex sequences, which bounds the dimension above. For each $\lambda \in \mathbb{C}^\times$ consider $(\lambda^n)_{n \geq 1}$; any finite linear combination of such sequences with distinct λ that is eventually zero has all coefficients zero, by applying the Vandermonde determinant to a sufficiently late block of consecutive indices. These give continuum many independent vectors in the quotient. For the second assertion, only finitely many U_n meet a bounded disk, so the restricted cocycle has finite singular support and splits. *This statement is about these particular classes; it does not assert vanishing of the sheaf cohomology of every bounded disk for arbitrary data.* $\square$

Example 13.6 (An explicit nontrivial bundle over the ordinary plane). For $\Gamma = \mathbb{Q}$ take $a_j = t^{1/j}$. Then

$$g_{0j}(z) = \text{Exp}_H \left(\frac{t^{1/j}}{z - j} \right) = \sum_{k \geq 0} \frac{t^{k/j}}{k! (z - j)^k}$$

is a valid Hahn unit on every punctured disk, and these transitions define a nontrivial line bundle on $(\mathbb{C}, \mathcal{H}_{\mathbb{Q}})$. At each fixed j the support is a perfectly well-ordered arithmetic progression; nontriviality comes only from trying to trivialize all the transitions simultaneously, which would force the descending set $\{1/j : j \geq 2\}$ into one global support.

For the bundles of [theorem 13.3](#), restriction to any relatively compact ordinary domain is trivial: only finitely many centres lie in it, the finite principal-part problem is solvable, and principal parts at outside centres are already holomorphic there and are absorbed into the local gauges. A bundle can therefore be trivial on every disk of an ordinary exhaustion while admitting no compatible global trivialization.

13.3 The divisor obstruction as a line-bundle class

The cluster data (1.3) yield unit transitions

$$h_{0c} = P_c(z - c)/(z - c)^{m_c} \in 1 + \mathcal{H}_{\Gamma,+}(U_0 \cap U_c),$$

which define a line bundle even without a common support across c ; its logarithmic cocycle is L_c from (8.3).

Proposition 13.7 (The divisor and logarithmic obstructions coincide). *This line bundle is trivial if and only if the clusters are admissible; equivalently, exactly when the prescribed effective divisor is globally realizable as in theorem 8.2.*

Proof. The normalization argument in theorem 13.3 shows that triviality is equivalent to an additive Mittag–Leffler solution for the principal parts L_c , which by theorem 7.2 is equivalent to well ordering of $T = \bigcup_c \text{supp } L_c$. If $E(P)$ is well ordered then all the logarithmic supports lie in $E(P)^* \setminus \{0\}$. Conversely, if T is well ordered then exponentiating each L_c puts every coefficient support of P_c in T^* , so $E(P)$ is well ordered. Apply theorem 8.2 for the final equivalence. $\square$

In particular the effective sub-divisor of corollary 9.3 carries a genuine line-bundle obstruction; it is not merely the failure of a particular product formula. The transitions for a moved simple root are $1 - a_c/(z - c)$, whose logarithm has higher pole terms, and should not be confused with the simpler exponential transitions (13.9): both constructions detect the same well-ordering issue, but their cocycles are not asserted to be equal.

14 The Picard-group dichotomy over the plane

14.1 Separating the ordinary and positive-support factors

Proposition 14.1 (Reduction of the plane Picard group to positive supports). *There are natural isomorphisms of abelian groups*

$$\text{Pic}(\mathbb{C}, \mathcal{H}_{\Gamma, \geq 0}) \cong \text{Pic}(\mathbb{C}, \mathcal{H}_{\Gamma}) = \text{Pic}_{\text{lf}}(\mathbb{C}, \mathcal{H}_{\Gamma}) \cong H^1(\mathbb{C}, \mathcal{H}_{\Gamma, +}). \quad (14.1)$$

In particular all of these groups carry a complex vector-space structure in which tensor product corresponds to vector addition, and the map (13.8) is complex-linear for it. Every class has an integral model over $\mathcal{H}_{\Gamma, \geq 0}$ whose ordinary reduction is the trivial holomorphic line bundle.

Proof. By corollary 6.3 and (6.2), $\text{Pic}(\mathbb{C}, \mathcal{H}_{\Gamma}) = \text{Pic}_{\text{lf}}(\mathbb{C}, \mathcal{H}_{\Gamma}) = H^1(\mathbb{C}, \mathcal{H}_{\Gamma}^{\times})$ and $\text{Pic}(\mathbb{C}, \mathcal{H}_{\Gamma, \geq 0}) = H^1(\mathbb{C}, \mathcal{H}_{\Gamma, \geq 0}^{\times})$. Taking first cohomology in (3.7) and (3.8) of proposition 3.5 gives

$$\begin{aligned} H^1(\mathbb{C}, \mathcal{H}_{\Gamma}^{\times}) &\cong H^1(\mathbb{C}, \underline{\Gamma}) \times H^1(\mathbb{C}, \mathcal{O}^{\times}) \times H^1(\mathbb{C}, \mathcal{H}_{\Gamma, +}), \\ H^1(\mathbb{C}, \mathcal{H}_{\Gamma, \geq 0}^{\times}) &\cong H^1(\mathbb{C}, \mathcal{O}^{\times}) \times H^1(\mathbb{C}, \mathcal{H}_{\Gamma, +}). \end{aligned}$$

The factor $H^1(\mathbb{C}, \underline{\Gamma})$ vanishes because locally constant torsors over the simply connected, locally contractible plane have trivial monodromy and a global section. The factor $H^1(\mathbb{C}, \mathcal{O}^{\times})$ vanishes by the classical triviality of holomorphic line bundles on $\mathbb{C}$, derived in section 4 from $H^1(\mathbb{C}, \mathcal{O}) = 0$, $H^2(\mathbb{C}, \mathbb{Z}) = 0$ and the exponential sequence [6]. Only $H^1(\mathbb{C}, \mathcal{H}_{\Gamma, +})$ remains, and extension of scalars from $\mathcal{H}_{\Gamma, \geq 0}$ to $\mathcal{H}_{\Gamma}$ is the displayed isomorphism because both unit decompositions contain the same $\mathcal{H}_{\Gamma, +}$ factor. The last assertion is remark 3.6 together with the transitions being in $1 + \mathcal{H}_{\Gamma, +}$ after normalization. $\square$

This does not say that a contractible space can carry a nontrivial ordinary complex line bundle. It says that the coefficient sheaf is different: the obstruction lies in holomorphic dependence coupled

to a global well-ordered support, not in ordinary monodromy. When $\Gamma \neq 0$ there is no reduction ring map $\mathcal{H}_\Gamma \rightarrow \mathcal{O}$ sending positive monomials to zero, because those monomials are units in $\mathcal{H}_\Gamma$; reduction is always performed on the $\mathcal{H}_{\Gamma, \geq 0}$ model.

14.2 Exactly which value groups give vanishing over the plane

Lemma 14.2 (Descending positive sequences). *A nonzero ordered abelian group Γ has no infinite strictly decreasing sequence of positive elements if and only if it is order-isomorphic to $\mathbb{Z}$, that is, $\Gamma = \mathbb{Z}\delta$ for some $\delta > 0$.*

Proof. For $\Gamma = \mathbb{Z}\delta$ with $\delta > 0$ every positive element is a positive integer multiple of δ , so no such sequence exists. Conversely, if the positive cone has no least element, choose a smaller positive element repeatedly to construct the forbidden sequence. So there is a least positive δ . If $\Gamma \neq \mathbb{Z}\delta$, choose $\gamma > 0$ not an integer multiple of δ ; we claim $\gamma - n\delta > 0$ for every $n \geq 0$. Otherwise take the first n with $\gamma - n\delta \leq 0$; then $0 < \gamma - (n-1)\delta \leq \delta$, and minimality of δ forces equality, so $\gamma = n\delta$, a contradiction. Hence $\gamma, \gamma - \delta, \gamma - 2\delta, \dots$ is a strictly decreasing positive sequence, contradicting the hypothesis. $\square$

Theorem 14.3 (Picard-group vanishing dichotomy over the plane). *For every set-sized ordered additive subgroup $\Gamma \subseteq \mathbf{No}$,*

$$\mathrm{Pic}(\mathbb{C}, \mathcal{H}_{\Gamma, \geq 0}) = 0 \iff \mathrm{Pic}(\mathbb{C}, \mathcal{H}_\Gamma) = 0 \iff \Gamma = \{0\} \text{ or } \Gamma = \mathbb{Z}\delta \text{ for some } \delta > 0. \quad (14.2)$$

In every other case both groups, identified as in (14.1), contain an embedded complex vector space $\mathbb{C}^{\mathbb{N}}/\mathbb{C}^{(\mathbb{N})}$, and hence continuum many linearly independent classes, realized by the explicit transitions

$$g_{n0}(a) = \mathrm{Exp}_H\left(\frac{a_n t^{\gamma_n}}{z - n}\right) \quad \text{on } W_n, \quad \gamma_1 > \gamma_2 > \dots > 0. \quad (14.3)$$

Proof. By [proposition 14.1](#) it suffices to decide when $H^1(\mathbb{C}, \mathcal{H}_{\Gamma, +})$ vanishes. For $\Gamma = 0$ one has $\mathcal{H}_{\Gamma, +} = 0$ and $\mathcal{H}_\Gamma = \mathcal{O}$.

Suppose $\Gamma = \mathbb{Z}\delta$ with $\delta > 0$. On any ordinary open set a positive-support section is simply $\sum_{k \geq 1} a_k(z) t^{k\delta}$ with $a_k \in \mathcal{O}(U)$ and no condition beyond holomorphy, every subset of $\{k\delta : k \geq 1\}$ being well ordered; as an additive sheaf $\mathcal{H}_{\Gamma, +} = \prod_{k \geq 1} \mathcal{O} t^{k\delta}$. Given a positive-support additive cocycle on an ordinary cover, expand it at each $k\delta$; each coefficient is an ordinary holomorphic additive cocycle, so [lemma 4.3](#) splits it on the same cover, giving local cochains $b_{i,k}$. Their series $b_i = \sum_{k \geq 1} b_{i,k} t^{k\delta}$ are valid positive-support sections and solve the original cocycle. This argument works on the original cover and does not assume that infinite products commute with higher sheaf cohomology; alternatively the torsor interpretation supplies local coordinates in the original trivializations. Hence $H^1(\mathbb{C}, \mathcal{H}_{\Gamma, +}) = 0$.

If Γ is neither trivial nor cyclic, [lemma 14.2](#) supplies a strictly decreasing positive sequence (γ_n) , and the sequence $a_n = t^{\gamma_n}$ lies in $\mathcal{V}_\Gamma \setminus \mathcal{W}_\Gamma$, so its bundle is nontrivial by [theorem 13.3](#); [corollaries 13.4](#) and [13.5](#) supply the embedded vector space and its dimension, and exponentiation turns those additive cocycles into (14.3), legitimate on each overlap by positive-support summability. The unit decomposition (3.8) shows that no extra ordinary unit or monomial change of frame can trivialize these classes unless the underlying additive cocycle was already a coboundary. $\square$

The word *cyclic* cannot be replaced by “has a least positive element”. For example $\Gamma = \mathbb{Z} + \mathbb{Z}\omega \subseteq \mathbf{No}$ has least positive element 1 but is not cyclic, and $\gamma_n = \omega - n$ is positive and strictly decreasing,

so the transitions $\text{Exp}_H(t^{\omega-n}/(z-n))$ give a nontrivial line bundle. Higher-rank exponents beyond all multiples of the least positive element matter. Every nonzero divisible group, in particular $\Gamma = \mathbb{Q}$, also lies on the nontrivial side.

Over the plane version of the coefficientwise meromorphic sheaf $\mathcal{M}_\Gamma$ of [definition 17.1](#), these explicit classes, and every bundle of [theorem 13.3](#), become trivial: their transitions and inverses extend across each centre with meromorphic coefficients (nonabelian-support report, `docs/surcomplex/nonabelian-support/`, `nab:loc:prop:poles` and `nab:loc:cor:polestrivial`). This does not contradict (14.2), which concerns $\mathcal{H}_{\Gamma, \geq 0}$ and $\mathcal{H}_\Gamma$ only. Bundles of rank at least two that stay nontrivial over the meromorphic sheaf are constructed there (`nab:loc:thm:dichotomy`).

14.3 The size of the nonzero plane Picard groups

Theorem 14.4 (Continuum many independent classes over the plane, and the exact dimension). *If Γ is noncyclic then*

$$\dim_{\mathbb{C}} \text{Pic}(\mathbb{C}, \mathcal{H}_\Gamma) \geq 2^{\aleph_0}. \quad (14.4)$$

If Γ is countable and noncyclic, equality holds; in particular

$$\dim_{\mathbb{C}} \text{Pic}(\mathbb{C}, \mathcal{H}_{\mathbb{Q}}) = 2^{\aleph_0}. \quad (14.5)$$

Proof. Choose a strictly decreasing positive sequence $(\gamma_j)_{j \geq 2}$. There is a family $(I_x)_{x \in X}$ of infinite subsets of $\{2, 3, \dots\}$ with $|X| = 2^{\aleph_0}$ and all pairwise intersections finite: encode finite binary strings by integers and let I_x be the set of finite prefixes of an infinite binary sequence x , two distinct infinite sequences sharing only finitely many prefixes. For each x set $a_j^{(x)} = t^{\gamma_j}$ if $j \in I_x$ and 0 otherwise. Consider a finite nonzero complex-linear combination of the $a^{(x)}$ and pick one x with nonzero coefficient; infinitely many indices of I_x lie in none of the finitely many other selected sets, and at each such index the combination still has a nonzero coefficient at γ_j . Those exponents contain an infinite strictly decreasing subsequence, so the combination is not in $\mathcal{W}_\Gamma$. Hence the classes of the $a^{(x)}$ are linearly independent in $\mathcal{V}_\Gamma/\mathcal{W}_\Gamma$, and [theorem 13.3](#) and [proposition 14.1](#) give (14.4).

Assume now Γ countable. On any ordinary open set the number of holomorphic functions is at most $\mathfrak{c} = 2^{\aleph_0}$, since a function on each connected component is determined by its values on a countable dense set and there are at most countably many components. A section of $\mathcal{H}_{\Gamma,+}$ is specified by a subset of a countable group together with countably many such functions, so $|\mathcal{H}_{\Gamma,+}(U)| \leq \mathfrak{c}$ for every U . Every torsor can be trivialized on a countable cover selected from a fixed countable basis of the plane, and there are at most $\mathfrak{c}$ such covers; for each, a cocycle consists of countably many sections, so there are at most $\mathfrak{c}^{\aleph_0} = \mathfrak{c}$ cocycles. Hence $|H^1(\mathbb{C}, \mathcal{H}_{\Gamma,+})| \leq \mathfrak{c}$, and with the independent family already constructed the dimension is exactly $\mathfrak{c}$. $\square$

This determines vanishing for every value group and the exact dimension for countable noncyclic groups. It does *not* claim that (13.7) is surjective, and it supplies no canonical basis or normal form for an arbitrary line-bundle class; those are different classification questions.

Remark 14.5 (What survives without the stalk theorem). [Theorems 14.3](#) and [14.4](#) are stated for the unrestricted Picard group because [corollary 6.3](#), resting on [theorem 6.1](#), identifies it with Pic_{lf} . A reader who declines that identification — as one of the two merged manuscripts deliberately does, see [remark 6.4](#) — still obtains every statement above verbatim for $\text{Pic}_{\text{lf}}(\mathbb{C}, \mathcal{H}_\Gamma)$ and for $\text{Pic}(\mathbb{C}, \mathcal{H}_{\Gamma, \geq 0})$, since (6.2) and [proposition 14.1](#) need no hypothesis on stalks. What that manuscript asserts, and all it asserts, about the size in the nonvanishing cases is the *containment* of a copy of $\mathbb{C}^{\mathbb{N}}/\mathbb{C}^{(\mathbb{N})}$, i.e. [corollaries 13.4](#) and [13.5](#). The exact dimension in (14.5), with its countability hypothesis and the

cocycle-counting upper bound, is the sharper statement and is the one proved here. The two are not in conflict: the containment is what survives without the stalk theorem and without the counting argument, and it is the sharper statement for the purpose of exhibiting explicit independent classes, while (14.5) is the sharper statement for the purpose of measuring the group.

14.4 Additive cohomology over the plane behaves differently

Proposition 14.6 (The full additive sheaf over the plane is already obstructed in cyclic rank). *For every nonzero Γ the group $H^1(\mathbb{C}, \mathcal{H}_\Gamma)$ contains a copy of $\mathbb{C}^\mathbb{N}/\mathbb{C}^{(\mathbb{N})}$; in particular $H^1(\mathbb{C}, \mathcal{H}_\Gamma) \neq 0$ even when Γ is cyclic and the Picard group vanishes.*

Proof. Choose $\delta > 0$ in Γ and repeat the construction of corollary 13.4 with the exponents $\gamma_n = -n\delta$, which are strictly decreasing but not positive. The general, not necessarily positive, version of theorem 13.1 applies unchanged and gives precisely the same kernel: the cocycle $b_n(a) = a_n t^{-n\delta}/(z - n)$ is a coboundary exactly when (a_n) is finitely supported. The weaker assertion $H^1(\mathbb{C}, \mathcal{H}_\Gamma) \neq 0$ also follows directly from the necessity half of theorem 7.2 applied to $p_j = t^{-j\delta}/(z - j)$, since $\{-j\delta : j \geq 2\}$ is not well ordered. $\square$

There is no contradiction with theorem 14.3: Hahn exponentiation cannot be applied to negative-support cochains, so logarithms of units are governed by $\mathcal{H}_{\Gamma,+}$ and not by the full Laurent sheaf $\mathcal{H}_\Gamma$. The line-bundle group and the additive cohomology of $\mathcal{H}_\Gamma$ therefore vanish under different conditions, and only the former is cyclic-rank sensitive.

14.5 Why this does not contradict ordinary complex geometry

The ordinary plane is contractible and its ordinary holomorphic line bundles are trivial. The nontrivial bundles above are locally trivial for a *different* coefficient sheaf, and their obstruction is neither an ordinary winding number nor a topological hole: it is the impossibility of putting infinitely many forced exponents into one well-ordered coherent section on the central patch.

Nor is “Hahn-coherent” synonymous with “coherent $\mathcal{O}$ -module”. For $\Gamma \neq 0$ the sheaf $\mathcal{H}_\Gamma$ is not locally finitely generated over $\mathcal{O}$: at a point c , evaluation gives

$$R_c/(z - c)R_c \cong K_\Gamma,$$

which is infinite-dimensional over $\mathbb{C}$, while a finitely generated $\mathcal{O}_c$ -module would have a finite-dimensional quotient by its maximal ideal. The same computation for $\mathcal{H}_{\Gamma,+}$ gives the complex vector space of positive Hahn constants, which already contains the infinitely many independent elements $t^{k\delta}$ for any $\delta > 0$. *The adjective “Hahn-coherent” specifies a common-support condition, not ordinary module coherence, and the present proofs do not apply Cartan’s theorem to $\mathcal{H}_\Gamma$ as though it were a coherent $\mathcal{O}$ -module.*

15 Moving divisors as nontrivial line bundles over the plane

The realization theorem and the cohomology theorem describe one failure in two complementary languages. The following statement isolates the geometric form of that failure.

Proposition 15.1 (A plane divisor with trivial ordinary reduction but no global equation). *Let $\gamma_1 > \gamma_2 > \dots > 0$ in Γ . There is a locally principal effective divisor on $(\mathbb{C}, \mathcal{H}_{\Gamma, \geq 0})$ whose local*

surcomplex zeros are the simple points $n + t^{\gamma_n}$, $n \geq 1$, whose ordinary reduction is a principal ordinary divisor, and whose associated line bundle is nontrivial both over $\mathcal{H}_{\Gamma, \geq 0}$ and after extension of scalars to $\mathcal{H}_{\Gamma}$.

Proof. Choose an ordinary entire f_0 with simple zeros exactly at the positive integers. On the cover (13.1) take the local equations

$$h_0(z) = f_0(z) \text{ on } U_0, \quad h_n(z) = \frac{f_0(z)}{z - n} (z - n - t^{\gamma_n}) \text{ on } U_n,$$

noting that $f_0(z)/(z - n)$ is ordinary holomorphic and nowhere zero on U_n . On W_n the ratio is

$$g_{n0} = \frac{h_n}{h_0} = 1 - \frac{t^{\gamma_n}}{z - n}, \quad (15.1)$$

a unit of $\mathcal{H}_{\Gamma, \geq 0}(W_n)$ by [proposition 3.4](#). The equations are nonzerodivisors with unit ratios, so they define the asserted effective locally principal divisor and a rank-one line bundle, and reduction gives the ordinary equation f_0 on every patch.

Its positive logarithmic cocycle is

$$\text{Log}_H g_{n0} = - \sum_{k \geq 1} \frac{t^{k\gamma_n}}{k(z - n)^k}. \quad (15.2)$$

At the exponent γ_n the n th component has a genuine simple pole, so the singular support contains the whole descending sequence (γ_n) and is not well ordered. [Theorem 13.1](#) and [proposition 14.1](#) then show that the bundle is nontrivial in both categories. Equivalently, a trivializing unit frame v_i with $g_{n0} = v_n/v_0$ would make h_i/v_i glue to a global coherent datum with precisely the prescribed divisor, whose symmetric support is $\{\gamma_n : n \geq 1\}$; [theorem 8.2](#) rules that out. $\square$

This construction has three simultaneous features: all local equations are elementary, the ordinary reduction is globally principal, and the full coherent divisor is not principal. It therefore isolates a genuinely new support obstruction rather than importing a familiar nontrivial ordinary line bundle.

15.1 What the existence and obstruction results say together

There are four distinct levels of data in the examples.

Prescription	Intrinsic global datum	Outcome
One displaced simple root at each n	$u - t^{1/n}$	Not realizable: descending symmetric support (example 9.1).
All n roots of $u^n - t$ at each n	The single coefficient $-t$	Realizable, even with integer Hahn exponents (example 9.2).
Select one of those roots by a 0/1 value	Polynomial (10.5)	Not interpolable: descending negative remainder support (example 10.4).
Reciprocal principal part at each cluster	Numerator 1 over $u^n - t$	Realizable despite descending individual residue scales (example 11.2).

Neither ordinary discreteness, nor finite solvability, nor bounded raw values, nor the support of individually labelled roots or residues is the correct universal test. The appropriate finite-dimensional invariant must be extracted first — a monic cluster polynomial, a polynomial remainder, or a proper rational numerator — and only then is global support admissibility tested.

16 A second base: compact ordinary Riemann surfaces

Everything up to this point is a theorem about one ordinary base, the complex plane $\mathbb{C}$. The plane is noncompact, contractible and Stein, and each of those three properties is used somewhere above: contractibility kills $H^1(\mathbb{C}, \Gamma)$ and $H^1(\mathbb{C}, \mathcal{O}^\times)$ in [proposition 14.1](#), Steinness supplies [lemma 4.3](#), and noncompactness is what allows infinitely many prescribed centres in the first place — which is exactly where all the plane obstructions come from.

This part of the article changes exactly one thing: the ordinary base becomes a *fixed compact connected ordinary Riemann surface* X of genus g . The coefficient recipe, the value group, the monomial convention $t^\gamma = \omega^{-\gamma}$, the support calculus and the summability discipline are unchanged. What changes is the answer, and it changes in both directions at once. On a compact base a closed discrete set of centres is finite, so the plane’s well-ordering obstruction evaporates; in its place appear three genuinely new invariants that the plane cannot see, because on the plane two of them are the cohomology of a contractible space.

16.1 Why the two global theories do not specialize to each other

Neither part implies the other, and the failure is not symmetric.

The compact theorems do not specialize to the plane. The proof of the comparison theorem below ([theorem 17.7](#)) uses compactness *twice*, at two logically independent points: once to know that the ordinary harmonic spaces are finite-dimensional, which is what makes the transferred obstruction a *matrix*, and once to know that a globally defined smooth Hahn form has *one* well-ordered support ([lemma 17.2](#)), which is what makes a global coefficientwise Green operator legitimate at all. Local support bounds on infinitely many disjoint regions need not combine into a well-ordered set — indeed [example 9.1](#) and [corollary 13.4](#) are precisely the statement that they need not. So:

The compact-base results of [sections 17 to 23](#) must not be applied to the noncompact plane by replacing X with $\mathbb{C}$ and setting $g = 0$.

In a report that now proves both halves, this is not a courtesy sentence. [Example 22.1](#) below computes $\text{Pic}_{\text{lf}}(\mathbb{P}^1, \mathcal{H}_\Gamma) = \text{Pic}(\mathbb{P}^1) = \mathbb{Z}$ with no Abel obstruction whatsoever, and $\mathbb{P}^1$ is the compact genus-zero curve; [theorem 14.3](#) computes $\text{Pic}(\mathbb{C}, \mathcal{H}_\Gamma) \neq 0$ for every noncyclic Γ , and $\mathbb{C}$ is the noncompact genus-zero base. Read across bases, those two statements look like a contradiction. They are not: the first is a finite-cluster statement on a compact curve, the second is an infinite-cluster statement on a noncompact one, and the only thing they share is the coefficient sheaf.

The plane theorems do not specialize to the compact case. The sharp criteria of [theorems 7.2, 8.2 and 13.1](#) are about infinite families of prescriptions. On a compact base every coherent Cartier datum treated in [section 21](#) is supported on a *finite* set of ordinary points, so the union of finitely many well-ordered supports is automatically well ordered and the plane’s admissibility hypothesis is vacuous. The compact obstructions are therefore of a different kind: they are cohomological invariants of the base, not support pathologies.

16.2 What is kept, and what is renamed

No statement of [sections 1 to 15](#) is weakened, restated or reproved here; those are theorems about $(\mathbb{C}, \mathcal{H}_\Gamma)$ and they stand exactly as printed. Because the article now carries two bases, the Picard

material of [sections 14](#) and [15](#) names its base in every heading and every theorem title, and the compact statements below do the same. The *labels* of all earlier statements are unchanged.

Three pieces of the plane apparatus are used verbatim on the compact base, with proofs that never mentioned $\mathbb{C}$:

- the support calculus [lemma 2.1](#), together with its three negative companions in [remark 2.2](#);
- the positive-support exponential and logarithm [proposition 3.5](#), whose proof is a coefficientwise formal identity on any commutative Hahn coefficient algebra;
- the unit criterion [proposition 3.4](#) and the unit splitting [\(3.8\)](#), whose proof uses only that $\mathcal{O}(U)$ is an integral domain for connected U .

These are cited below rather than reproved. Everything else in this part is new material.

	Base $\mathbb{C}$ (sections 1 to 15)	Base compact X (sections 17 to 23)
Prescriptions	Infinitely many ordinary centres	Finitely many ordinary centres
Governing invariant	Well ordering of one global support	Ordinary cohomology of the base, coefficientwise
$H^1(\text{base}, \underline{\Gamma})$	0 (contractible)	Γ^{2g} , the valuation monodromy
Ordinary Picard factor	0 (Stein, contractible)	$\text{Pic}(X)$, nonzero already for $g = 0$
Infinitesimal factor	$H^1(\mathbb{C}, \mathcal{H}_{\Gamma,+})$, not computed in general	$H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_{\Gamma} \cong \mathfrak{m}_{\Gamma}^g$, computed exactly
Vanishing	Exactly for $\Gamma = 0$ or $\Gamma = \mathbb{Z}\delta$ (theorem 14.3)	Never for $g \geq 1$ and $\Gamma \neq 0$; for $g = 0$ the group is $\mathbb{Z}$

17 The coefficient sheaf on a compact base, and Dolbeault comparison

17.1 The sheaves, and the one convention that carries over

Throughout [sections 17 to 23](#), X is a fixed compact connected ordinary Riemann surface of genus g , with its ordinary complex topology, and $\Gamma \subseteq \mathbf{No}$ is the same fixed set-sized ordered group as before, with K_{Γ} , $\mathfrak{m}_{\Gamma}$, $\mathcal{O}_{\Gamma}$ as in [section 1.2](#). The source manuscript of this part writes R_{Γ} for the valuation ring $\mathcal{O}_{\Gamma} = \{a \in K_{\Gamma} : v(a) \geq 0\}$; we keep $\mathcal{O}_{\Gamma}$, which is a ring of *scalars* and is never the structure sheaf $\mathcal{O}_X$.

Definition 17.1 (The Hahn sheaves over a compact base). On an ordinary open $U \subseteq X$, a section of $\mathcal{H}_{\Gamma}$ is a family $(f_{\gamma})_{\gamma \in \Gamma}$ of ordinary holomorphic functions on U whose set of nonzero indices is *locally* contained in one well-ordered subset of Γ , all coefficients of a given local section being defined on the *same* ordinary open neighbourhood. The subsheaves $\mathcal{H}_{\Gamma, \geq 0}$ and $\mathcal{H}_{\Gamma, +}$ impose nonnegative and strictly positive support. The sheaf $\mathcal{M}_{\Gamma}$ is defined by the same recipe with ordinary *meromorphic* coefficients. We write $\mathcal{A}_{\Gamma}^{0,q}$ for the analogous sheaf of smooth Hahn $(0, q)$ -forms, with subscripts ≥ 0 and $+$ for its support restrictions. The symbols $\mathcal{H}_{\Gamma}$, $\mathcal{H}_{\Gamma, \geq 0}$, $\mathcal{H}_{\Gamma, +}$ are the same as in [definition 3.1](#); the base is named in every statement below.

The *local* form of the support condition is what makes this a presheaf with the gluing property on disconnected opens. On a connected nonempty U the ordinary identity theorem collapses it, exactly as in [proposition 3.3](#), to

$$\mathcal{H}_{\Gamma}(U) = \mathcal{O}_X(U)((t^{\Gamma})), \quad \mathcal{M}_{\Gamma}(U) = \mathcal{M}_X(U)((t^{\Gamma})), \quad (17.1)$$

because a nonzero holomorphic or meromorphic coefficient on a connected U cannot vanish identically on a nonempty open subset. On a disconnected open, different components may carry different supports.

Smooth coefficients satisfy no identity theorem, so (17.1) has no smooth analogue and the support of a smooth Hahn form is a genuinely local datum. Compactness is what repairs this.

Lemma 17.2 (Compact support uniformity). *A globally defined smooth Hahn form on a compact X has one well-ordered support. The same holds for forms valued in an ordinary finite-rank smooth bundle. Consequently every coefficientwise ordinary linear operator on such a global space is defined and support preserving.*

Proof. Choose a finite subcover of neighbourhoods carrying support bounds $S_1, \dots, S_N$. Every nonzero coefficient is nonzero on at least one member, so its exponent lies in $S_1 \cup \dots \cup S_N$. A finite union of well-ordered subsets of a linear order is well ordered: a nonempty subset meets some S_j , has a least element in each nonempty intersection, and the least of those finitely many elements is least in the subset. For a bundle use a finite ordinary trivializing cover; ordinary transition matrices introduce no exponent. $\square$

Remark 17.3 (Exactly what compactness has just bought). This is the point at which the two bases part company. On $\mathbb{C}$ the analogous statement is false, and its failure is the subject of sections 13 to 14: the cocycles of corollary 13.4 are built precisely so that local support bounds on infinitely many disjoint disks refuse to combine. Every later use of a *global* coefficientwise Green or Hodge operator on this compact base silently invokes lemma 17.2, and none of them is available over the plane.

17.2 The operator form of the support calculus

Lemma 2.1 controls formal power series in positive-support Hahn *elements*. The compact part needs the same control for a series of positive-support *operators* whose coefficients are ordinary linear maps on an infinite-dimensional space of smooth sections; those maps are not uniformly bounded in any sense, and no norm or convergence hypothesis is available or wanted.

Lemma 17.4 (Neumann inverse, operator form). *Let $E \subseteq \Gamma_{>0}$ be well ordered and let $A = \sum_{e \in E} t^e A_e$ act on Hahn families valued in a complex vector space, each A_e being an arbitrary complex-linear coefficient operator. Then*

$$(1 + A)^{-1} = \sum_{n \geq 0} (-A)^n \quad (17.2)$$

is a well-defined two-sided inverse: acting on a family with support in S , it produces a family with support in $S + E^$, and each output coefficient is a finite sum of terms $A_{e_1} \cdots A_{e_n}(x_s)$ with $s + e_1 + \dots + e_n$ the given exponent.*

Proof. By lemma 2.1 the set $S + E^*$ is well ordered and a fixed exponent has only finitely many representations $s + e_1 + \dots + e_n$, with only finitely many word lengths n occurring. Hence the displayed coefficient of $A^n x$ at a fixed exponent is a finite sum, and summing over n adds only finitely many further terms. Every coefficient of (17.2) is therefore a finite sum of ordinary operator compositions, and multiplying by $1 + A$ cancels the terms pairwise at each exponent. $\square$

Remark 17.5 (What a support certificate is, and what it refuses). A containment

$$\text{supp}(\text{output}) \subseteq S + E^* \quad (17.3)$$

is the only licence used below for an infinite construction. It proves that the result is *one* Hahn series, and it exhibits, for each fixed exponent, the finitely many ordinary terms that contribute there; ordinary differentiation, integration and Green operators are then applied to those finitely many terms only, so no exchange of a divergent analytic sum with an integral occurs anywhere.

The mechanism is finite decomposability of an *exact* exponent, not eventual domination. It need *not* be true that $n\delta$ eventually exceeds every element of $\Gamma_{>0}$: in $\Gamma = \mathbb{Q} + \mathbb{Q}\omega$ one has $n < \omega$ for every integer n . Neither [lemma 17.4](#) nor any proof in [sections 17 to 23](#) uses that false cofinality assertion, and none of them is a convergence argument in disguise. This is the third companion of [remark 2.2](#), restated here because the operators it now licenses are unbounded ones. Cofinality of a scale in Γ enters this part at exactly one place, [theorem 23.3](#), where it decides whether a one-scale formal test is faithful.

17.3 The coefficientwise resolution and the comparison theorem

Proposition 17.6 (Dolbeault resolution over a compact base). *There are fine resolutions of sheaves on X*

$$0 \longrightarrow \mathcal{H}_\Gamma \longrightarrow \mathcal{A}_\Gamma^{0,0} \xrightarrow{\bar{\partial}} \mathcal{A}_\Gamma^{0,1} \longrightarrow 0,$$

and the same for the nonnegative- and positive-support versions. For an ordinary holomorphic vector bundle V on X the analogous resolution computes the cohomology of V with Hahn coefficients.

Proof. The kernel statement is coefficientwise holomorphy. For local surjectivity take a coordinate disk on which the input coefficients are smooth, together with a fixed smaller disk, and one smooth cutoff supported in the first and equal to 1 on the second. The ordinary Cauchy–Green solution operator built from that *fixed* cutoff solves $\bar{\partial}$ for every coefficient on the same smaller disk, so no exponent-dependent shrinking is needed and the coefficientwise extension preserves support. An ordinary smooth partition of unity acts by multiplication in each coefficient, which gives fineness. The bundle statement is local in an ordinary holomorphic frame. On a curve there are no $(0, 2)$ -forms, so the resolution stops in degree one. $\square$

Fix ordinary Hermitian metrics, write $d = \bar{\partial}_V$, and let i_q include the ordinary harmonic representatives, p_q be ordinary harmonic projection and h the ordinary Hodge homotopy from degree one to degree zero, so that

$$hd = 1 - i_0 p_0, \quad dh = 1 - i_1 p_1, \quad p_0 h = 0, \quad h i_1 = 0, \quad p_q i_q = 1. \quad (17.4)$$

These are ordinary compact-curve Hodge theory [\[13\]](#). By [lemma 17.2](#) they remain valid after coefficientwise Hahn extension, with any of the three support restrictions.

Theorem 17.7 (Compact cohomology comparison). *Let X be compact of genus g and let V be an ordinary holomorphic vector bundle on X . For $q = 0, 1$,*

$$H^q(X, V_\Gamma) \cong H^q(X, V) \otimes_{\mathbb{C}} K_\Gamma, \quad (17.5)$$

$$H^q(X, V_{\Gamma, \geq 0}) \cong H^q(X, V) \otimes_{\mathbb{C}} \mathcal{O}_\Gamma, \quad (17.6)$$

$$H^q(X, V_{\Gamma, +}) \cong H^q(X, V) \otimes_{\mathbb{C}} \mathfrak{m}_\Gamma, \quad (17.7)$$

and all higher cohomology vanishes. In particular

$$H^1(X, \mathcal{H}_{\Gamma,+}) \cong H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_{\Gamma}, \quad H^0(X, \mathcal{H}_{\Gamma}) = K_{\Gamma}.$$

Proof. [Proposition 17.6](#) identifies sheaf cohomology with the cohomology of the global two-term complex. Extend (17.4) coefficientwise, which is legitimate by [lemma 17.2](#). The maps i and p then identify that cohomology with Hahn families of ordinary harmonic representatives, preserving the support restriction. Ordinary harmonic spaces on a compact curve are finite-dimensional, so such families are exactly the displayed tensor products — this is the one step where finite dimensionality is used, and it is also why the identification does not depend on the auxiliary metrics. The last assertion is that a holomorphic function on a compact connected Riemann surface is constant. $\square$

Remark 17.8 (Compactness is used twice, and the noncompact case is different). The proof needs two separate consequences of compactness: finite-dimensionality of the ordinary harmonic spaces, without which the transferred data below would not be a finite matrix, and a *single* global well-ordered support bound for smooth Hahn forms ([lemma 17.2](#)), without which the global coefficientwise Green operator is not even defined. Local bounds on infinitely many disjoint regions need not combine into a well-ordered set. *Theorem 17.7 must not be applied to the noncompact plane merely by replacing X by $\mathbb{C}$ and setting $g = 0$; corollaries 13.4 and 13.5 exhibit classes in $H^1(\mathbb{C}, \mathcal{H}_{\Gamma,+})$ that no such formula could produce, and $H^1(\mathbb{C}, \mathcal{O}) = 0$ by [lemma 4.3](#) while $H^1(\mathbb{C}, \mathcal{H}_{\Gamma,+}) \neq 0$ for every noncyclic Γ .*

18 The exact Picard classification over a compact base

18.1 Three independent components of a unit

Lemma 18.1 (Unit decomposition over a compact base). *On X there are canonical isomorphisms of abelian sheaves*

$$\mathcal{H}_{\Gamma}^{\times} \cong \underline{\Gamma} \times \mathcal{O}_X^{\times} \times \mathcal{H}_{\Gamma,+}, \quad (\gamma, a, h) \mapsto t^{\gamma} a \operatorname{Exp}_H(h), \quad \mathcal{H}_{\Gamma, \geq 0}^{\times} \cong \mathcal{O}_X^{\times} \times \mathcal{H}_{\Gamma,+}. \quad (18.1)$$

The splitting is relative to the fixed monomial section $\gamma \mapsto t^{\gamma}$ and the fixed inclusion of ordinary coefficients.

Proof. This is [propositions 3.4](#) and [3.5](#) and is not reproved: their proofs use only that $\mathcal{O}(U)$ is an integral domain on a connected U , that least exponents add under multiplication, and the coefficientwise formal identities for Exp_H and Log_H . None of that refers to the base. An integral unit has itself and its inverse of nonnegative support, which forces $\gamma = 0$. $\square$

Definition 18.2 (The three classes of a compact Hahn line bundle). A *Hahn line bundle* on X is a locally free rank-one $\mathcal{H}_{\Gamma}$ -module; its class lies in $\operatorname{Pic}_{\operatorname{lf}}(X, \mathcal{H}_{\Gamma}) = H^1(X, \mathcal{H}_{\Gamma}^{\times})$ and has three components under (18.1): the *valuation class* $\nu(L) \in H^1(X, \underline{\Gamma})$, the ordinary component $L_0 \in \operatorname{Pic}(X)$, and the *positive logarithmic class* $\xi(L) \in H^1(X, \mathcal{H}_{\Gamma,+})$. Its degree is $\deg L_0$. The ordinary component is a projection of unit cocycles; *it is not a reduction map on all of $\mathcal{H}_{\Gamma}$* , in accordance with [remark 3.6](#).

Theorem 18.3 (Exact Picard classification over a compact base). *For X compact connected of genus g there are canonical group isomorphisms*

$$\operatorname{Pic}_{\operatorname{lf}}(X, \mathcal{H}_{\Gamma}) \cong H^1(X, \underline{\Gamma}) \oplus \operatorname{Pic}(X) \oplus (H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_{\Gamma}), \quad (18.2)$$

$$\operatorname{Pic}_{\operatorname{lf}}(X, \mathcal{H}_{\Gamma, \geq 0}) \cong \operatorname{Pic}(X) \oplus (H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_{\Gamma}). \quad (18.3)$$

Moreover $H^1(X, \underline{\Gamma}) \cong \text{Hom}(H_1(X, \mathbb{Z}), \Gamma)$, so after choosing a homology basis and a basis of $H^1(X, \mathcal{O}_X)$,

$$\text{Pic}_{\text{lf}}(X, \mathcal{H}_{\Gamma}) \cong \Gamma^{2g} \oplus \text{Pic}(X) \oplus \mathfrak{m}_{\Gamma}^g. \quad (18.4)$$

Extension of scalars from $\mathcal{H}_{\Gamma, \geq 0}$ to $\mathcal{H}_{\Gamma}$ is injective on these groups, with image exactly the subgroup of zero valuation class.

Proof. For any ringed space, locally free rank-one modules are classified by unit-valued transition cocycles modulo changes of local frame; this is (6.2), which needs no hypothesis on stalks. Taking H^1 of the finite direct-product decomposition (18.1) gives (18.2) and (18.3), and theorem 17.7 identifies the last summand with $H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_{\Gamma}$. The ordinary unit term is the ordinary holomorphic Picard group $\text{Pic}(X)$.

A compact surface has a finite good cover, so constant-sheaf cohomology there agrees with singular cohomology with coefficients in Γ . In degree one the universal-coefficient sequence has zero extension term because $H_0(X, \mathbb{Z}) = \mathbb{Z}$ is free, which gives the Hom description. Finally $H_1(X, \mathbb{Z}) = \mathbb{Z}^{2g}$ and $\dim_{\mathbb{C}} H^1(X, \mathcal{O}_X) = g$ give (18.4), and the map induced by integral extension is the inclusion of the last two factors, the valuation factor being zero on integral units. $\square$

Remark 18.4 (What is *not* computed: the nonzero-valuation sector). Theorem 18.3 classifies the bundles. It does *not* compute their cohomology. Theorem 19.2 and corollary 19.4 below compute H^0 and H^1 only in the sector $\nu(L) = 0$, because the positive perturbation argument of proposition 19.1 needs a global ordinary component and a global scalar positive gauge, and neither exists in the required form when $\nu(L) \neq 0$. Theorem 20.1 proves that $H^0(X, L) = 0$ whenever $\nu(L) \neq 0$; that vanishing of sections determines neither $H^1(X, L)$ nor an Euler characteristic in that sector, and no such determination is claimed here. The plane part of this article has no analogous gap, because over $\mathbb{C}$ the valuation factor $H^1(\mathbb{C}, \underline{\Gamma})$ vanishes outright (proposition 14.1) and every class already lies in the zero-valuation sector. This asymmetry between the two bases is genuine and is not an artefact of the exposition.

18.2 A normal form with a finite support bound

If $\nu(L) = 0$, take a finite good trivializing cover and change local frames by monomials so as to remove the valuation cocycle. The remaining transitions are uniquely

$$g_{ij} = a_{ij} \text{Exp}_H(h_{ij}), \quad a_{ij} \in \mathcal{O}_X^{\times}(U_i \cap U_j), \quad h_{ij} \in \mathcal{H}_{\Gamma, +}(U_i \cap U_j), \quad (18.5)$$

the a_{ij} defining L_0 and the h_{ij} forming an additive cocycle. The cover being finite, a union of finitely many well-ordered sets gives one well-ordered $E \subseteq \Gamma_{>0}$ containing every $\text{supp } h_{ij}$; local smooth gauges and all later nonlinear operations are then certified in E^* by remark 17.5. *It is finiteness of the cover, not any property of Γ , that produces E* ; this is the compact counterpart of the plane's admissibility hypothesis, and it is why that hypothesis disappears here.

Remark 18.5 (Local stalk identification and the compact Picard group). The proof of theorem 18.3 uses only actual rank-one local trivializations and the unit sheaf, so it needs no hypothesis on stalks. The independently written compact source deliberately restricted its statements to Pic_{lf} and did not import the plane source's PID theorem. That is a statement about its proof dependencies, not an obstruction to comparing the stalks in the merged article.

Indeed, choose an ordinary holomorphic coordinate $u : U \rightarrow D$ about $x \in X$, with $u(x) = 0$. Composition with u^{-1} sends each holomorphic coefficient on U to one on D , introduces no Hahn

exponent and commutes with restriction. It therefore identifies $(\mathcal{H}_\Gamma)_x$ with the ordinary-base plane stalk R_0 . By [theorem 6.1](#) this stalk is a PID. The same argument works on any ordinary Riemann surface and uses no compactness.

Consequently the local dual-generator argument of [corollary 6.3](#) gives

$$\mathrm{Pic}(X, \mathcal{H}_\Gamma) = \mathrm{Pic}_{\mathrm{lf}}(X, \mathcal{H}_\Gamma).$$

The integral stalk is local by the same coordinate identification and [proposition 6.2](#), so the analogous equality holds for $\mathcal{H}_{\Gamma, \geq 0}$. Thus the compact classification also describes tensor-invertible modules. We retain $\mathrm{Pic}_{\mathrm{lf}}$ in its statements to make their original proof scope visible. This local transfer supplies neither the plane’s global vanishing dichotomy on X nor the compact cohomology comparison on the plane.

18.3 The two bases side by side

[Theorems 14.3](#) and [14.4](#) and [theorem 18.3](#) are two computations of the Picard group of *the same* coefficient sheaf over two different ordinary bases, and each is false over the other base. It is worth stating the contrast once, explicitly, since the two are easy to misquote across bases.

Over $\mathbb{C}$: $\mathrm{Pic}(\mathbb{C}, \mathcal{H}_\Gamma) = \mathrm{Pic}_{\mathrm{lf}}(\mathbb{C}, \mathcal{H}_\Gamma) \cong H^1(\mathbb{C}, \mathcal{H}_{\Gamma,+})$, which vanishes exactly when $\Gamma = \{0\}$ or $\Gamma = \mathbb{Z}\delta$, and otherwise has complex dimension at least $2^{\aleph_0}$, with equality for countable noncyclic Γ . The two vanishing factors $H^1(\mathbb{C}, \underline{\Gamma})$ and $H^1(\mathbb{C}, \mathcal{O}^\times)$ carry no information, and the surviving factor is *not* computed in closed form.

Over compact X : the three factors are identified, although individual factors can vanish, and the third is computed explicitly, $\Gamma^{2g} \oplus \mathrm{Pic}(X) \oplus \mathfrak{m}_\Gamma^g$. There is no well-ordering dichotomy at all: for $g \geq 1$ and $\Gamma \neq 0$ the group is never zero, and for $g = 0$ it is $\mathbb{Z}$ for every Γ , cyclic or not.

Neither statement is a special case of the other, and neither survives transport to the other base. The shared content includes the coefficient sheaf and the unit splitting ([18.1](#)). On the plane only the positive-support cohomology remains; the compact base can also contribute ordinary and valuation classes.

19 A finite matrix for the zero-valuation sector

19.1 From holomorphic transitions to a global operator

Proposition 19.1 (Positive Dolbeault normal form). *Let L be a Hahn line bundle on compact X with $\nu(L) = 0$ and transitions ([18.5](#)). There is a global positive-support scalar $(0, 1)$ -form α such that the smooth Hahn realization of L is identified with that of L_0 , and the holomorphic sections of L are the kernel of*

$$D = d + \delta, \quad d = \bar{\partial}_{L_0}, \quad \delta(s) = \alpha s. \quad (19.1)$$

Every coefficient of α may be taken ordinary harmonic, and the class of α is $\xi(L)$. If $\mathrm{supp} h_{ij} \subseteq E$ then $\mathrm{supp} \alpha \subseteq E$ and the smooth gauges have support in E^ .*

Proof. Fix an ordinary partition of unity (ρ_k) subordinate to the finite cover, with the convention $s_i = g_{ij}s_j$ for section coordinates, and set $b_i = \sum_k \rho_k h_{ik}$, each term extended smoothly by zero where its partition function vanishes. The cocycle identity gives $b_i - b_j = h_{ij}$, so $r_i = \mathrm{Exp}_H(-b_i)s_i$ has ordinary transitions a_{ij} . The holomorphic equation becomes

$$0 = \bar{\partial}s_i = \mathrm{Exp}_H(b_i)(\bar{\partial}r_i + (\bar{\partial}b_i)r_i),$$

and $\bar{\partial}(b_i - b_j) = 0$, so the forms $\bar{\partial}b_i$ glue to a global $(0, 1)$ -form α , the coefficientwise Čech–Dolbeault representative of $[h_{ij}]$. All operations preserve E except the exponential gauges, whose support lies in E^* by [proposition 3.5](#).

Every scalar $(0, 1)$ -form on a curve is $\bar{\partial}$ -closed, so ordinary Hodge splitting applied coefficientwise writes $\alpha = \alpha_{\text{harm}} + \bar{\partial}c$ with $\text{supp } c \subseteq E$; the further smooth change $r_i = \text{Exp}_H(-c)r'_i$ replaces α by its harmonic part. Both steps are legitimate on the global smooth spaces by [lemma 17.2](#). $\square$

The same construction applies to an integral Hahn line bundle using nonnegative-support coordinates. Local smooth solutions of $\bar{\partial}c = \alpha$ with the gauge $\text{Exp}_H(c)$ show that [\(19.1\)](#) is a fine Dolbeault resolution of L , whose global smooth terms are the Hahn extensions of the ordinary smooth terms of L_0 ; this follows from the finite trivializing cover together with the gauge support bound, and is *not* an assumption that a tensor product commutes with an infinite-dimensional space of sections.

19.2 The transferred matrix

Put

$$V_0 = H^0(X, L_0), \quad V_1 = H^1(X, L_0), \quad n_q = \dim_{\mathbb{C}} V_q,$$

and choose the ordinary Hodge maps [\(17.4\)](#) for L_0 , extended coefficientwise. Since $h\delta$ has positive support, [lemma 17.4](#) defines

$$T = (1 + h\delta)^{-1} = \sum_{n \geq 0} (-h\delta)^n, \quad B = p_1 \delta T i_0 : V_0 \otimes_{\mathbb{C}} K_{\Gamma} \longrightarrow V_1 \otimes_{\mathbb{C}} K_{\Gamma}, \quad (19.2)$$

equivalently

$$B = \sum_{n \geq 0} (-1)^n p_1 \delta (h\delta)^n i_0. \quad (19.3)$$

If $\text{supp } \alpha \subseteq E$ then every entry of B has support in $E^* \setminus \{0\}$; in particular B is an $n_1 \times n_0$ matrix over $\mathcal{O}_{\Gamma}$ with *all entries in $\mathfrak{m}_{\Gamma}$* . The inverse in [\(19.2\)](#) is a Hahn identity, not a limit: see [remark 17.5](#).

Theorem 19.2 (Finite cohomology and explicit representatives). *Let L be a Hahn line bundle on compact X with $\nu(L) = 0$, represented by [\(19.1\)](#). Then*

$$H^0(X, L) \cong \ker B, \quad H^1(X, L) \cong \text{coker } B. \quad (19.4)$$

The first isomorphism is realized by $v \mapsto T i_0 v$; in degree one a form w is represented by the finite vector

$$P_1 w = p_1 (1 + \delta h)^{-1} w = p_1 (w - \delta T h w), \quad (19.5)$$

taken modulo $\text{im } B$. For the normalized integral lattice $L_{\geq 0}$ the same statements hold with the finite complex

$$V_0 \otimes_{\mathbb{C}} \mathcal{O}_{\Gamma} \xrightarrow{B} V_1 \otimes_{\mathbb{C}} \mathcal{O}_{\Gamma}. \quad (19.6)$$

Every representative or correction built from input support S has support in $S + E^$. No valuation-rank and no cofinality hypothesis is required.*

Proof. We eliminate directly in the two-term complex, so that no infinite-dimensional perturbation theorem is invoked; every inverse expression is licensed first by [lemma 17.4](#), and only [\(17.4\)](#) is used.

Degree zero. Let $y = Ti_0v$, so $y + h\delta y = i_0v$. Applying d gives $dy + dh\delta y = 0$, and $dh = 1 - i_1p_1$ in degree one turns this into

$$Dy = i_1p_1\delta y = i_1Bv. \quad (19.7)$$

Also $p_0h = 0$ gives $p_0T = p_0$ and hence $p_0y = v$. So a vector in $\ker B$ determines a unique section. Conversely if $Ds = 0$, applying h and using $hd = 1 - i_0p_0$ gives $(1 + h\delta)s = i_0p_0s$, so $s = Ti_0p_0s$, and (19.7) gives $Bp_0s = 0$.

Degree one. For a degree-one form w put $y = Thw$; then $p_0y = 0$ because $p_0T = p_0$ and $p_0h = 0$. From $y + h\delta y = hw$,

$$h(w - Dy) = hw - (1 - i_0p_0)y - h\delta y = 0.$$

By $dh = 1 - i_1p_1$ a degree-one form killed by h is harmonic, so

$$w - DThw = i_1p_1(w - \delta Thw) = i_1P_1w, \quad (19.8)$$

and every class has a harmonic representative. The identity $1 - \delta Th = (1 + \delta h)^{-1}$ gives the second formula in (19.5).

Relations. Suppose $i_1b = Ds$. Applying h and using $hi_1 = 0$ gives $(1 + h\delta)s = i_0p_0s$, so $s = Ti_0v$ with $v = p_0s$, and (19.7) gives $b = Bv$; conversely every Bv is a boundary by the same identity. Hence degree-one cohomology is $\operatorname{coker} B$.

Every map used is coefficientwise linear or carries the positive Neumann support bound, so the argument runs unchanged with nonnegative supports and yields (19.6), together with the stated certificate. $\square$

Remark 19.3 (Relation to homological perturbation). The operators Ti_0 , P_1 and Th are the usual transferred inclusion, projection and homotopy of a two-term contraction, and B is its Schur complement. Homological perturbation and its deformation applications are classical [5, §2, especially Lemma 2.4][8], and *no discovery of that lemma is claimed*. Crainic uses $ip = 1 + dh + hd$; replacing his homotopy by $-h$ gives the sign convention (17.4) and the inverse $1 + h\delta$ here. What requires proof here is that the infinite expressions are admissible in an arbitrary-rank Hahn workspace and that the resulting global complex computes the common-domain sheaf cohomology of definition 17.1.

19.3 Riemann–Roch, stable section counts, and integral torsion

Corollary 19.4 (Riemann–Roch in the zero-valuation sector). *Let $\nu(L) = 0$ and $d = \deg L_0$. Both cohomology groups are finite-dimensional over K_Γ and, with $r = \operatorname{rank}_{K_\Gamma} B$,*

$$h_\Gamma^0(L) = n_0 - r, \quad h_\Gamma^1(L) = n_1 - r, \quad h_\Gamma^0(L) - h_\Gamma^1(L) = d + 1 - g.$$

In particular $h_\Gamma^q(L) \leq \dim_{\mathbb{C}} H^q(X, L_0)$. If $d > 2g - 2$ then $H^1(X, L) = 0$ and $h_\Gamma^0(L) = d + 1 - g$, and the integral section module is free of rank $d + 1 - g$ over $\mathcal{O}_\Gamma$, with basis the images under Ti_0 of an ordinary basis of $H^0(X, L_0)$. If $d < 0$ then $H^0(X, L) = 0$.

Proof. The kernel and cokernel of theorem 19.2 have the stated dimensions and their difference is $n_0 - n_1 = d + 1 - g$ by ordinary Riemann–Roch. Ordinary Serre duality gives $n_1 = 0$ for $d > 2g - 2$, so B has zero target and the integral conclusion follows from the same finite complex. For $d < 0$ one has $n_0 = 0$, so the kernel vanishes. $\square$

Proposition 19.5 (Degree-zero deformations of the trivial bundle). *Let $L_0 = \mathcal{O}_X$ with harmonic positive class $\xi \in H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_\Gamma$. Then $B : K_\Gamma \rightarrow K_\Gamma^g$ sends 1 to the coordinate vector of ξ . Hence $h_\Gamma^0(L) = 0$ and $h_\Gamma^1(L) = g - 1$ when $\xi \neq 0$, and 1 and g when $\xi = 0$. On an elliptic curve with $\xi = \lambda\eta$ for an ordinary harmonic generator $\eta \neq 0$ and $0 \neq \lambda \in \mathfrak{m}_\Gamma$, the integral groups are*

$$H^0(X, L_{\geq 0}) = 0, \quad H^1(X, L_{\geq 0}) \cong \mathcal{O}_\Gamma / \lambda \mathcal{O}_\Gamma. \quad (19.9)$$

Proof. The ordinary harmonic space in degree zero consists of constants, and α is harmonic, so $h\delta(1) = h\alpha = 0$, whence $T(1) = 1$ and $B(1) = p_1\alpha = \xi$. A nonzero map from a one-dimensional space has rank one. Over $\mathcal{O}_\Gamma$ the elliptic complex is multiplication by λ on $\mathcal{O}_\Gamma$, a domain, which gives (19.9). $\square$

The group (19.9) is genuine torsion of the *integral* model that is invisible over the field: by corollary 19.4 the same bundle has $H^0 = H^1 = 0$ over K_Γ . Passing to the field discards the torsion; reducing to ordinary coefficients discards the infinitesimal class. Those are two different operations and neither sees what the other sees.

Corollary 19.6 (Base extension does not remove an obstruction). *Let $\Gamma \subseteq \Gamma'$ be an inclusion of set-sized ordered groups using the same monomials. The induced map on compact Picard groups is injective, and for a valuation-zero bundle*

$$H^q(X, L) \otimes_{K_\Gamma} K_{\Gamma'} \cong H^q(X, L_{\Gamma'}), \quad q = 0, 1.$$

Proof. In (18.2) both $\mathfrak{m}_\Gamma \hookrightarrow \mathfrak{m}_{\Gamma'}$ and $\text{Hom}(H_1(X, \mathbb{Z}), \Gamma) \rightarrow \text{Hom}(H_1(X, \mathbb{Z}), \Gamma')$ are injective, and $\text{Pic}(X)$ is untouched. For cohomology use the same ordinary Hodge data: the matrix B is unchanged and merely viewed over a larger field, and kernels and cokernels of finite matrices commute with field extension. $\square$

This is the compact-base counterpart of section 9.4, and the two are proved by completely different means: over the plane, because a larger ambient ordered group cannot well-order a strictly decreasing sequence; over X , because a finite matrix keeps its rank. That both bases refuse the same repair is a fact about the coefficient sheaf, not a shared argument.

20 Valuation monodromy is exactly the meromorphic obstruction

20.1 The meromorphic convention

A *coherent meromorphic section* of a Hahn line bundle L on X is a section of $L \otimes_{\mathcal{H}_\Gamma} \mathcal{M}_\Gamma$. On a connected chart its coordinate is a Hahn series of ordinary meromorphic functions. Individual coefficients may have *different and unbounded* pole orders — no uniform bound is imposed — while the ordinary domain stays fixed. *This meromorphic sheaf is specified here and is not silently identified with any other sheaf of fractions or with a non-Archimedean analytic function field*; compare the corresponding refusal for the plane in section 11. Every nonzero section of $\mathcal{M}_\Gamma$ on a connected open is invertible: its leading coefficient is an invertible ordinary meromorphic function and the remaining factor is inverted by lemma 2.1.

Theorem 20.1 (Exact meromorphic-section criterion). *A Hahn line bundle L on compact connected X has a nonzero coherent meromorphic section if and only if*

$$\nu(L) = 0 \quad \text{in } H^1(X, \underline{\Gamma}). \quad (20.1)$$

If $\nu(L) \neq 0$ then $H^0(X, L) = 0$, whatever the degree of L .

Proof. Let s be a nonzero coherent meromorphic section and fix a finite connected good trivializing cover with $s_i = g_{ij}s_j$. No local coordinate can vanish identically: the coefficientwise identity theorem propagates a zero across any nonempty overlap and X is connected. So the local valuations $\beta_i = v(s_i)$ are defined, and the transition identity gives $v(g_{ij}) = \beta_i - \beta_j$, exhibiting the valuation cocycle as a coboundary.

Conversely let $\nu(L) = 0$. Fix an ordinary point $p \in X$ and $N \geq 0$ with $d + N > 2g - 2$ and $d + N + 1 - g > 0$. The twist $L(Np)$ still has zero valuation class, so by [corollary 19.4](#) it has a nonzero holomorphic section. The ordinary bundle $\mathcal{O}_X(Np)$ has its usual nonzero meromorphic trivialization, and using it converts that section into a nonzero coherent meromorphic section of L ; locally this multiplies a Hahn series by one ordinary meromorphic factor, preserving the common-domain and support conditions. A nonzero holomorphic section is in particular meromorphic, which gives the last assertion. $\square$

20.2 Divisor classes occupy exactly the zero-valuation sector

Define the sheaf of coherent meromorphic Cartier data by $\mathcal{D}_\Gamma = \mathcal{M}_\Gamma^\times / \mathcal{H}_\Gamma^\times$, put $\text{Div}_\Gamma^{\text{coh}}(X) = H^0(X, \mathcal{D}_\Gamma)$, and let $\text{Prin}_\Gamma^{\text{coh}}(X)$ be the image of $H^0(X, \mathcal{M}_\Gamma^\times)$. This convention includes the finite cluster data of [section 21](#), but *it does not assert that every section of $\mathcal{D}_\Gamma$ is a finite sum of moving points*.

Corollary 20.2 (The divisor-class sequence). *There is a split exact sequence*

$$0 \longrightarrow \frac{\text{Div}_\Gamma^{\text{coh}}(X)}{\text{Prin}_\Gamma^{\text{coh}}(X)} \longrightarrow \text{Pic}_{\text{lf}}(X, \mathcal{H}_\Gamma) \xrightarrow{\nu} H^1(X, \underline{\Gamma}) \longrightarrow 0, \quad (20.2)$$

so that

$$\frac{\text{Div}_\Gamma^{\text{coh}}(X)}{\text{Prin}_\Gamma^{\text{coh}}(X)} \cong \text{Pic}(X) \oplus (H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_\Gamma). \quad (20.3)$$

Proof. The exact sequence of multiplicative sheaves defining $\mathcal{D}_\Gamma$ embeds Cartier data modulo principal data into $H^1(X, \mathcal{H}_\Gamma^\times)$, with image the bundles that become trivial over $\mathcal{M}_\Gamma$, that is those carrying a nonzero coherent meromorphic section. By [theorem 20.1](#) that image is $\ker \nu$. The monomial map $\gamma \mapsto t^\gamma$ splits ν , and [theorem 18.3](#) identifies the kernel. $\square$

Example 20.3 (High degree without sections). Let $g \geq 1$ and $\Gamma \neq 0$, and choose a nonzero homomorphism $\rho : H_1(X, \mathbb{Z}) \rightarrow \Gamma$. A constant-sheaf cocycle for ρ gives a Hahn line bundle with monomial transitions $t^{\rho_{ij}}$. Tensoring with any ordinary degree- d line bundle leaves the valuation class equal to ρ , so by [theorem 20.1](#) the result has no nonzero coherent meromorphic section, and *a fortiori* no nonzero holomorphic section, for arbitrarily large d .

This does not contradict [corollary 19.4](#), whose hypothesis is $\nu(L) = 0$. Nor does it contradict Riemann–Roch for a proper algebraic curve over K_Γ : *the ordinary-base sheaf category is not thereby identified with that algebraic category*, and a purported equivalence preserving degree, ordinary twists and global sections would already fail on these examples. This is a boundary on the category, not a counterexample to anything classical, and it is the sharpest single obstruction proved in this part.

21 A root-free Abel criterion for finite moving clusters

21.1 Finite cluster data and their ordinary reduction

Fix a finite set $A \subset X$ of distinct ordinary points with pairwise disjoint coordinate disks, the coordinate at c being u_c with $u_c(c) = 0$. On each disk take two monic polynomials

$$P_c^\pm(u) = u^{m_c^\pm} + \sum_{r=0}^{m_c^\pm-1} a_{c,r}^\pm u^r, \quad a_{c,r}^\pm \in \mathfrak{m}_\Gamma, \quad (21.1)$$

an empty cluster having $m = 0$ and $P = 1$. Let D be the coherent Cartier datum given by P_c^+/P_c^- on the disks and by 1 off A ; this is legitimate because on a punctured disk each $P_c^\pm/u^{m_c^\pm}$ is a unit of $\mathcal{H}_\Gamma$ by [proposition 3.4](#). Its ordinary reduction is the ordinary divisor

$$D_0 = \sum_{c \in A} (m_c^+ - m_c^-) c, \quad (21.2)$$

and the valuation class of the associated Hahn line bundle is zero.

These are divisor data at infinitesimally displaced points, not merely values at the ordinary centres. *If Γ is divisible* then K_Γ is algebraically closed [[15](#), Corollary 4] and the roots of each nontrivial cluster polynomial lie in $\mathfrak{m}_\Gamma$ — a root of valuation $\beta \leq 0$ would give its leading term strictly smaller valuation than every lower term, which cannot cancel — so D may be read as a finite signed multiset of infinitesimal points. *The criterion below needs no roots and is stated without any divisibility hypothesis; without divisibility a cluster need not split in K_Γ , although its roots always exist in a divisible Hahn enlargement. Nothing below requires such an enlargement.*

21.2 Newton sums are the correct symmetric coordinates

For a monic $P(u) = u^m + c_1 u^{m-1} + \cdots + c_m$ define the Newton power sums $p_k(P)$, $k \geq 1$, by the integer recurrences

$$p_k + c_1 p_{k-1} + \cdots + c_{k-1} p_1 + k c_k = 0, \quad 1 \leq k \leq m, \quad (21.3)$$

$$p_k + c_1 p_{k-1} + \cdots + c_m p_{k-m} = 0, \quad k > m, \quad (21.4)$$

with all $p_k = 0$ for $P = 1$. These are universal symmetric polynomials with integer coefficients; *if* roots are chosen then $p_k(P) = \sum_j \varepsilon_j^k$ with multiplicity, but the recurrences do not choose any.

Remark 21.1 (The same insistence as [theorem 8.2](#), one step further). [Theorem 8.2](#) decides realizability by the symmetric support $E(P)$ of the cluster coefficients and not by the supports of separately labelled roots, and [section 9.4](#) and the discussion after it explain why a root labelling can destroy exactly the cancellations that make a global equation possible. The Newton power sums are that same insistence carried one step further: they are integer-coefficient functions of the symmetric coefficients, so the Abel criterion below is *stated and proved over arbitrary, not necessarily divisible, ordered groups*, and it yields a sharper support bound inside E^* than separately chosen fractional-exponent roots would.

Lemma 21.2 (Logarithmic principal-part identity). *Let E be the union of the supports of all nonleading coefficients in (21.1); it is a well-ordered subset of $\Gamma_{>0}$. On a punctured ordinary disk,*

$$\mathrm{Log}_H \frac{P(u)}{u^m} = - \sum_{k \geq 1} \frac{p_k(P)}{k u^k}. \quad (21.5)$$

The right-hand side is strongly summable as a family of meromorphic-coefficient Hahn series, its support lies in $E^* \setminus \{0\}$, and at each fixed Hahn exponent only finitely many negative powers of u occur.

Proof. E is a finite union of well-ordered positive supports, hence well ordered. Write $Q(u) = P(u)/u^m - 1 = \sum_{j=1}^m c_j u^{-j}$, so the left side is $\sum_{n \geq 1} (-1)^{n+1} Q(u)^n / n$. By lemma 2.1 a fixed exponent receives contributions from finitely many finite words in E , and each corresponding product is an ordinary polynomial in u^{-1} of finite degree; so the coefficient at that exponent has a finite principal part. For formal independent coefficients the identity

$$\log(1 + c_1 q + \cdots + c_m q^m) = - \sum_{k \geq 1} \frac{p_k(P)}{k} q^k$$

is the classical generating identity for Newton sums, obtained by logarithmic differentiation and comparison of coefficients using (21.3)–(21.4). Substituting the positive-support c_j and $q = u^{-1}$ is licensed by the finite-word argument just given, which also justifies regrouping by powers of u^{-1} . $\square$

Let η be an ordinary holomorphic differential on X , written at c as

$$\eta = \left(\sum_{n \geq 0} b_{c,n}(\eta) u_c^n \right) du_c, \quad (21.6)$$

the coefficients being the ordinary Taylor coefficients on that disk.

Definition 21.3 (The infinitesimal Abel functional). For the signed cluster datum D set

$$\mathcal{A}_D(\eta) = \sum_{c \in A} \sum_{k \geq 1} \frac{b_{c,k-1}(\eta)}{k} (p_k(P_c^+) - p_k(P_c^-)), \quad (21.7)$$

an element of $\mathfrak{m}_\Gamma$, linear in $\eta \in H^0(X, \Omega_X^1)$.

Proposition 21.4 (Well-definedness and a residue formula). *The sum (21.7) is strongly summable with support in $E^* \setminus \{0\}$. Putting*

$$\ell_c = \text{Log}_H \frac{P_c^+}{u_c^{m_c^+}} - \text{Log}_H \frac{P_c^-}{u_c^{m_c^-}},$$

one has

$$\mathcal{A}_D(\eta) = - \sum_{c \in A} \text{Res}_c(\ell_c \eta). \quad (21.8)$$

For split clusters it also equals the signed sum of local primitive displacements

$$\sum_{c,j} \int_0^{\varepsilon_{c,j}^+} \eta - \sum_{c,j} \int_0^{\varepsilon_{c,j}^-} \eta, \quad \int_0^\varepsilon \eta := \sum_{k \geq 1} \frac{b_{c,k-1}(\eta)}{k} \varepsilon^k. \quad (21.9)$$

The integrals in (21.9) are Taylor–Hahn substitutions, not fine-topological path integrals.

Proof. At a fixed Hahn exponent lemma 21.2 leaves only finitely many negative powers of u_c , so multiplying by the ordinary holomorphic Taylor series and extracting the residue uses finitely many Taylor coefficients; (21.5) then gives (21.8) and the summability claim. If roots are chosen, substituting $p_k = \sum_j \varepsilon_j^k$ and using lemma 2.1 on each of the finitely many primitive series gives (21.9). The symmetric formula supplies the sharper bound in E^* even when individual root supports would require fractional exponents. $\square$

21.3 The exact criterion

Theorem 21.5 (Support-controlled Abel criterion). *For the finite signed cluster datum D of (21.1) on compact X the following are equivalent.*

- (i) *There is $F \in \mathcal{M}_\Gamma(X)^\times$ such that $F/(P_c^+/P_c^-)$ is a unit of $\mathcal{H}_\Gamma$ on every cluster disk and F is a unit of $\mathcal{H}_\Gamma$ away from A .*
- (ii) *The ordinary divisor D_0 is principal, and $\mathcal{A}_D(\eta) = 0$ for every ordinary holomorphic differential η .*

It suffices to test a basis of the g -dimensional space $H^0(X, \Omega_X^1)$. When these hold, fix any ordinary meromorphic f_0 with $\text{div}(f_0) = D_0$; one may choose F with exponent-zero coefficient f_0 and support in E^ , and such an F is unique up to a nonzero constant of K_Γ . The criterion, and any nonzero obstruction, survive every ordered-group enlargement.*

Proof. Use the finite cover by the cluster disks together with $U_0 = X \setminus A$. The normalized bundle $\mathcal{O}(D) \otimes \mathcal{O}(-D_0)$ has positive logarithmic transition data ℓ_c on the punctured disks, zero valuation class and trivial ordinary component, so its infinitesimal class is some

$$\xi_D \in H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_\Gamma, \quad (21.10)$$

all coefficients of its logarithmic cocycle lying in E^* by lemma 21.2.

The ordinary Serre pairing between $H^1(X, \mathcal{O}_X)$ and $H^0(X, \Omega_X^1)$ may be computed by sums of residues of punctured-disk Čech representatives. Fix the convention under which the pairing of a Dolbeault class $[\alpha]$ with η is $(2\pi i)^{-1} \int_X \alpha \wedge \eta$ and the corresponding Čech representative has value $\sum_c \text{Res}_c(\ell_c \eta)$; the sign is verified by taking a cutoff equal to 1 on the inner and 0 on the outer circle of each annulus and applying Stokes' theorem there. That is ordinary complex analysis applied to one Hahn coefficient at a time. With proposition 21.4 it gives

$$\langle \xi_D, \eta \rangle = -\mathcal{A}_D(\eta). \quad (21.11)$$

Changing the Serre sign convention changes both sides by one overall sign and not the zero locus. The ordinary pairing is perfect, so its finite-dimensional coefficientwise extension is nondegenerate on $H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_\Gamma$, whence $\xi_D = 0$ if and only if $\mathcal{A}_D(\eta) = 0$ for every η .

If (i) holds the class of D vanishes; its ordinary component is $[D_0]$ and its positive component is ξ_D , so theorem 18.3 gives (ii). Conversely (ii) kills those two components while the valuation component is already zero, and corollary 20.2 then gives principality, which is (i).

For the constructive addendum, suppose $\xi_D = 0$. Every coefficient of the logarithmic Čech cocycle is then an ordinary coboundary on this fixed cover, and a support-preserving splitting is explicit: take a fixed partition of unity to get smooth b_i with $b_i - b_j = \ell_{ij}$, form the global exact $(0, 1)$ -form $\alpha = \bar{\partial} b_i$, and set $c = h\alpha$; then $k_i = b_i - c$ is holomorphic with $k_i - k_j = \ell_{ij}$. Applied coefficientwise this shrinks no domain and gives $\text{supp } k_i \subseteq E^* \setminus \{0\}$. Let f_i be the local defining function of D and let f_i^0 be $u_c^{m_c^+ - m_c^-}$ on a cluster disk and 1 on U_0 . The quotients $q_i = (f_i/f_i^0) \text{Exp}_H(-k_i)$ agree on overlaps, have meromorphic coefficients, exponent-zero coefficient 1 and support in E^* , so they define $q \in \mathcal{M}_\Gamma(X)$, and $F = f_0 q$ has the required local unit ratios and leading coefficient f_0 . Two such generators differ by a global unit of $\mathcal{H}_\Gamma$, which is a nonzero element of K_Γ by theorem 17.7. If the generators are normalized to have the same leading coefficient f_0 at exponent zero, their ratio lies in $1 + \mathfrak{m}_\Gamma$. Finally ordinary principality is unaffected by workspace enlargement and a nonzero Hahn coefficient of $\mathcal{A}_D$ stays nonzero, which is the last assertion. $\square$

Remark 21.6 (Ordinary periods versus infinitesimal displacement). The requirement that D_0 be principal is the ordinary Abel condition, degree zero together with the usual condition in the complex Jacobian [13]. The extra functional $\mathcal{A}_D$ has only *positive* Hahn exponents, while an ordinary nonzero period sits at exponent zero, so no period can cancel a positive-support obstruction. There is no unmentioned period quotient in (21.7): the infinitesimal condition is exact equality to zero.

Remark 21.7 (Coordinate independence, and what the theorem does not cover). For fixed moving-point data, changing the local coordinates changes the polynomial descriptions but not the sum of local primitive displacements of η ; equivalently the class ξ_D and its Serre pairing are intrinsic, so the vanishing criterion is coordinate independent. A polynomial such as $u^2 - s$ does describe a coordinate-dependent motion, and altering its coefficients without transporting the points is a change of divisor rather than a contradiction.

The theorem covers finite polynomial cluster Cartier data only. It does not assert that every section of $\mathcal{D}_\Gamma$ admits such a presentation, nor that an arbitrary meromorphic-coefficient Hahn series has a classical finite zero count.

22 Compact examples: the sphere, an elliptic curve, and a nonlinear obstruction

22.1 The sphere, and a warning about the plane

Example 22.1 (The sphere). If $X = \mathbb{P}^1(\mathbb{C})$ then $g = 0$ and [theorem 18.3](#) gives

$$\mathrm{Pic}_{\mathrm{lf}}(\mathbb{P}^1, \mathcal{H}_\Gamma) \cong \mathrm{Pic}(\mathbb{P}^1) \cong \mathbb{Z}.$$

There is neither valuation monodromy nor an infinitesimal Abel obstruction, and by [theorem 21.5](#) every finite signed cluster datum of total degree zero is principal: in standard affine coordinates the finite product of the defining polynomial ratios gives a rational generator, the factor at infinity accounting for the total degree. This explicit product uses coordinates $u_c = z - c$ at finite centres and $u_\infty = 1/z$ at infinity; for arbitrary chosen local coordinates the existence assertion follows from the Abel criterion, without identifying those coordinates with $z - c$.

This is not a global Mittag-Leffler assertion for the noncompact ordinary plane. On a compact base there are only finitely many prescribed clusters and the union of their supports is automatically well ordered, so the hypothesis of [theorem 8.2](#) is vacuous. Over $\mathbb{C}$ the same value group $\Gamma = \mathbb{Q}$ gives $\dim_{\mathbb{C}} \mathrm{Pic}(\mathbb{C}, \mathcal{H}_{\mathbb{Q}}) = 2^{N_0}$ by [theorem 14.4](#), and [example 9.1](#) exhibits an unrealizable divisor of exactly the kind that cannot occur here. Both statements are about genus zero and about the same sheaf; they differ only in compactness of the base, and neither may be quoted for the other.

22.2 An elliptic curve: exact displacement conservation

Example 22.2 (Elliptic displacement conservation). Let $X = \mathbb{C}/\Lambda$ with local coordinates induced by the uniformizing variable, so that dz generates $H^0(X, \Omega_X^1)$ and $b_{c,0} = 1$, $b_{c,n} = 0$ for $n > 0$. Then (21.7) reduces to

$$\mathcal{A}_D(dz) = \sum_c (p_1(P_c^+) - p_1(P_c^-)), \quad (22.1)$$

so the test is exact conservation of the sum of infinitesimal displacements, over and above the ordinary Abel condition.

For $P^+(u) = u - \varepsilon$ and $P^-(u) = u$ at one centre with $0 \neq \varepsilon \in \mathfrak{m}_\Gamma$, the ordinary divisor is zero while $\mathcal{A}_D(dz) = \varepsilon$, so a single moved point minus its original is not principal. By contrast $P^+(u) = u^2 - s$, $P^-(u) = u^2$ have $p_1 = 0$ for both clusters, and that signed divisor *is* principal in the uniformizing coordinate. The same curve with $\xi = \lambda\eta$ has no cohomology over K_Γ while its integral model has $H^1 \cong \mathcal{O}_\Gamma/\lambda\mathcal{O}_\Gamma$ by (19.9).

22.3 A genus-two cluster with zero first displacement and nonzero Abel class

Example 22.3 (Genus two, $y^2 = 1 + x^5$, balanced cluster). Consider the smooth projective completion of

$$y^2 = 1 + x^5. \quad (22.2)$$

The polynomial $1 + x^5$ has five simple roots, so the double cover of the x -sphere is branched at those five points and at infinity and Riemann–Hurwitz gives $g = 2$. A basis of holomorphic differentials is

$$\eta_0 = \frac{dx}{y}, \quad \eta_1 = \frac{x dx}{y}, \quad (22.3)$$

holomorphic at the finite branch points in their square-root parameters and at infinity in the parameter $x^{-1/2}$.

At $c = (0, 1)$ use $u = x$ and the branch $y = (1 + u^5)^{1/2}$. Fix $s = t^\gamma$ with $\gamma > 0$ and take the *balanced* cluster

$$P^+(u) = u^2 - s, \quad P^-(u) = u^2. \quad (22.4)$$

The ordinary divisor is zero, and in a divisible workspace the two moved roots are $\pm\sqrt{s}$, so the first symmetric displacement vanishes: $p_1(P^+) = p_1(P^-) = 0$. The power sums of P^+ are $p_{2r} = 2s^r$ and $p_{2r+1} = 0$, while all positive power sums of P^- vanish. Since $(1 + u^5)^{-1/2} = \sum_{n \geq 0} \binom{-1/2}{n} u^{5n}$, formula (21.7) gives the exact Hahn identities

$$\mathcal{A}_D(\eta_0) = \sum_{\ell \geq 0} \frac{\binom{-1/2}{2\ell+1}}{5\ell+3} s^{5\ell+3} = -\frac{s^3}{6} - \frac{5s^8}{128} - \cdots, \quad (22.5)$$

$$\mathcal{A}_D(\eta_1) = \sum_{\ell \geq 0} \frac{\binom{-1/2}{2\ell}}{5\ell+1} s^{5\ell+1} = s + \frac{s^6}{16} + \frac{35s^{11}}{1408} + \cdots. \quad (22.6)$$

The leading term s of (22.6) is nonzero, so [theorem 21.5](#) proves that the balanced cluster (22.4) is *not* principal: the first symmetric displacement cancels and a nonlinear Abel obstruction survives. All exponents lie in the monoid $\{n\gamma : n \geq 0\}$, and *no claim that $n\gamma$ is cofinal in Γ is needed or made*; compare [remark 17.5](#).

The displayed leading nonzero coefficient already certifies nonprincipality, and by [theorem 21.5](#) no enlargement of the value group removes it. The higher terms describe the full obstruction rather than an error term in any topology on $\mathbf{No}[i]$. *The series (22.5)–(22.6) are proved above; only their first few coefficients were confirmed symbolically by the accompanying script (appendix B).*

23 The exact information lost by one-scale completion

The integral sheaf is indispensable here. In $\mathcal{H}_\Gamma$ every t^δ is a unit, so a quotient by $t^{n\delta} \mathcal{H}_\Gamma$ would be the zero sheaf — the same point that [remark 3.6](#) makes about reduction. We therefore study

$\mathcal{H}_{\Gamma, \geq 0}/t^{n\delta}\mathcal{H}_{\Gamma, \geq 0}$ for $\delta \in \Gamma_{>0}$. Even its positive elements need not be nilpotent: if $0 < \varepsilon \ll \delta$ then $t^{k\varepsilon}$ survives for every integer k . Logarithms and exponentials remain available by strong summability, not by any false assumption that such series terminate.

23.1 Truncated coefficients and their Picard groups

Write

$$J_n = t^{n\delta}\mathcal{O}_\Gamma, \quad \mathcal{R}_n = \mathcal{H}_{\Gamma, \geq 0}/t^{n\delta}\mathcal{H}_{\Gamma, \geq 0}, \quad \mathfrak{m}_n = \mathfrak{m}_\Gamma/J_n,$$

noting $J_n \subseteq \mathfrak{m}_\Gamma$. Coefficients of $\mathcal{R}_n$ are represented uniquely by Hahn series with exponents in $[0, n\delta)$.

Lemma 23.1 (Truncated Picard classification). *For every positive integer n ,*

$$\mathrm{Pic}_{\mathrm{lf}}(X, \mathcal{R}_n) \cong \mathrm{Pic}(X) \oplus (H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_n), \quad (23.1)$$

and under this identification the reduction map from $\mathrm{Pic}_{\mathrm{lf}}(X, \mathcal{H}_{\Gamma, \geq 0})$ is the identity on $\mathrm{Pic}(X)$ and coefficient reduction on the second summand.

Proof. Cutting off exponents $\geq n\delta$ gives a canonical additive representative of a local quotient section; local representatives glue coefficientwise, and on a connected ordinary open the identity theorem makes their support well ordered, so the quotient sheaf really is the indicated sheaf of truncated Hahn coefficients rather than a pointwise quotient notation.

A truncated unit has invertible exponent-zero coefficient: necessity because the constant coefficients of two inverse representatives multiply to 1; sufficiency by dividing by that ordinary coefficient and using the geometric Hahn inverse of (3.4) before truncating. Positive exponential and logarithm respect J_n , because adding a nonnegative exponent to one that is at least $n\delta$ cannot return below $n\delta$, so they descend to inverse group maps on the positive truncated sheaf. This gives $\mathcal{R}_n^\times \cong \mathcal{O}_X^\times \times \mathcal{P}_n$ with $\mathcal{P}_n$ the positive truncated additive sheaf.

The proof of theorem 17.7 applies with exponents restricted to $(0, n\delta)$: the fixed local solution operators and the global Hodge maps preserve that restriction, and compactness again supplies one support bound through lemma 17.2. Hence $H^1(X, \mathcal{P}_n) = H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_n$, and taking H^1 of the unit decomposition gives the formula together with its compatibility with reduction. $\square$

23.2 The visible convex subgroup and the invisible ideal

Define

$$\Delta = \{\gamma \in \Gamma : |\gamma| \leq n\delta \text{ for some integer } n \geq 1\}, \quad I_\delta = \bigcap_{n \geq 1} t^{n\delta}\mathcal{O}_\Gamma. \quad (23.2)$$

Then Δ is the convex subgroup of Γ generated by δ . Write $R_\Delta = \mathbb{C}((t^\Delta))_{\geq 0}$ and $\mathfrak{m}_\Delta = \mathbb{C}((t^\Delta))_{>0}$. Truncation to exponents in Δ is a split surjective ring map

$$\mathcal{O}_\Gamma \longrightarrow R_\Delta, \quad (23.3)$$

whose kernel is exactly I_δ : a nonzero element lies in every J_n precisely when its least exponent exceeds every integer multiple of δ , and convexity of Δ then places all its other exponents outside Δ as well; conversely every positive exponent outside Δ exceeds every $n\delta$.

Lemma 23.2 (Inverse limit of the coefficient truncations). *There are canonical isomorphisms*

$$\varprojlim_n \mathcal{O}_\Gamma/J_n \cong R_\Delta, \quad \varprojlim_n \mathfrak{m}_\Gamma/J_n \cong \mathfrak{m}_\Delta. \quad (23.4)$$

Proof. A compatible family determines every coefficient with exponent in $\Delta_{\geq 0}$ by taking a large enough cutoff. Let S be the union of its nonzero indices; we check S is well ordered. Given a nonempty $T \subseteq S$ pick $\gamma_0 \in T$ and then n with $\gamma_0 < n\delta$. The nonempty set $T \cap [0, n\delta)$ lies in the well-ordered support of the n th truncation and so has a least element, which is least in all of T because every other exponent of T is at least $n\delta$. Hence the family defines an element of R_Δ . Restriction of a series in R_Δ is the inverse operation, and the maps respect addition and multiplication because a coefficient below $n\delta$ in a product of nonnegative series uses only exponents below that cutoff. Omitting the exponent-zero coefficient gives the positive version. $\square$

Theorem 23.3 (Exact one-scale Picard comparison). *Let X be compact of genus g , let $\delta \in \Gamma_{>0}$ and define Δ, I_δ by (23.2). The natural map*

$$\mathrm{Pic}_{\mathrm{lf}}(X, \mathcal{H}_{\Gamma, \geq 0}) \longrightarrow \varprojlim_n \mathrm{Pic}_{\mathrm{lf}}(X, \mathcal{H}_{\Gamma, \geq 0}/t^{n\delta} \mathcal{H}_{\Gamma, \geq 0}) \quad (23.5)$$

is split surjective, with

$$\varprojlim_n \mathrm{Pic}_{\mathrm{lf}}(X, \mathcal{H}_{\Gamma, \geq 0}/t^{n\delta} \mathcal{H}_{\Gamma, \geq 0}) \cong \mathrm{Pic}(X) \oplus (H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} \mathfrak{m}_\Delta), \quad (23.6)$$

$$\ker (23.5) \cong H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} I_\delta. \quad (23.7)$$

It is injective if and only if $g = 0$ or $\Delta = \Gamma$. Hence whenever $g > 0$ and the convex subgroup generated by δ is proper, all powers of that one scale miss genuine classes.

Proof. Apply lemma 23.1 at every level. The ordinary Picard factor has identity transition maps, and the other factor is a tensor product with the fixed finite-dimensional $H^1(X, \mathcal{O}_X)$; after choosing a basis it is a finite direct power, which commutes with inverse limits, so lemma 23.2 gives (23.6). Under the full integral classification (18.3) the map is the identity on $\mathrm{Pic}(X)$ and the convex-support cutoff $\mathfrak{m}_\Gamma \rightarrow \mathfrak{m}_\Delta$ on each infinitesimal coordinate; inclusion of Δ -supported coefficients splits it and its kernel is (23.7).

If $g = 0$ the kernel vanishes. For $g > 0$ it vanishes exactly when $I_\delta = 0$. If $\Delta = \Gamma$ no nonzero series has least exponent above every $n\delta$, so $I_\delta = 0$; if $\Delta \neq \Gamma$, choose a positive $\eta \notin \Delta$ and then $0 \neq t^\eta \in I_\delta$. $\square$

Remark 23.4 (What is, and is not, being completed). Theorem 23.3 computes an inverse limit of isomorphism classes of line bundles in the indicated integral quotient sheaves. It does not replace them by bundles over any fine surcomplex topology, and it does not infer a sheaf equivalence from a slogan about completion; the proof is an explicit compatible-coefficient calculation. Nor is the source Picard group given any complete Hausdorff topology a priori. What the kernel proves is that a topology built from the powers of a single scale would be nonseparated on the infinitesimal part whenever $g > 0$ and $\Delta \neq \Gamma$.

23.3 A deformation beyond every integer-scale jet

Example 23.5 (The t^ω witness). Take $\Gamma = \mathbb{Q} + \mathbb{Q}\omega \subset \mathbf{No}$ with its inherited order, $\delta = 1$, and a compact curve with $g \geq 1$. Then $\Delta = \mathbb{Q}$ and $t^\omega \in I_1$. Choose a nonzero $\zeta \in H^1(X, \mathcal{O}_X)$ represented on a finite cover by an ordinary additive cocycle h_{ij} . The transitions

$$g_{ij} = \text{Exp}_H(t^\omega h_{ij}) \quad (23.8)$$

define an integral line bundle of class $t^\omega \zeta \neq 0$, nontrivial over $\mathcal{H}_{\Gamma, \geq 0}$ and, by [theorem 18.3](#), over $\mathcal{H}_\Gamma$. Yet every transition reduces to 1 modulo $t^n \mathcal{H}_{\Gamma, \geq 0}$ for every positive integer n , so every one of those reductions is trivial. For an elliptic curve the integral cohomology is $H^1 \cong \mathcal{O}_\Gamma / t^\omega \mathcal{O}_\Gamma$ while the full Hahn cohomology vanishes in both degrees. *This is not a numerical failure to compute enough terms*: no integer multiple of the first scale reaches the obstruction, and a larger exponent scale is mathematically necessary. The same group $\Gamma = \mathbb{Q} + \mathbb{Q}\omega$ is the running witness of [remarks 2.2](#) and [17.5](#).

Corollary 23.6 (A set-indexed detection obstruction). *Let X be compact with $g \geq 1$ and let B be any set of positive surreal exponents. There exist a set-sized ordered subgroup $\Gamma \subset \mathbf{No}$ containing B and a nontrivial integral Hahn line bundle $L_{\geq 0}$ on X such that $L_{\geq 0} \bmod t^\beta \mathcal{H}_{\Gamma, \geq 0}$ is trivial for every $\beta \in B$; the same may be required for every positive integer multiple of every $\beta \in B$.*

Proof. The set of all those integer multiples is still a set, and every set of surreal numbers has a surreal strict upper bound, for instance by the usual set-sized cut construction. Choose a positive η strictly above that set and let Γ be the divisible additive subgroup generated by $B \cup \{\eta\}$, which is set-sized. Use the cocycle $\text{Exp}_H(t^\eta h_{ij})$ for a nonzero ordinary class $[h_{ij}]$: its Picard class is nonzero by [theorem 18.3](#), while all the stated reductions are trivial because $\eta > n\beta$ for every relevant n and β . *No cohomology of a proper-class space is formed anywhere in this construction*, in accordance with the set-sizedness convention of [section 1.2](#). $\square$

The order of the quantifiers in [corollary 23.6](#) matters: the workspace is chosen after the set of cutoffs. In a fixed nonzero Γ , all positive cutoffs together do detect every integral Picard class. The ordinary Picard component survives every reduction. If the positive component is nonzero, choose a basis of $H^1(X, \mathcal{O}_X)$ and a nonzero coordinate with least exponent $\gamma > 0$. Its coefficient at γ survives modulo t^β for any $\beta > \gamma$, for example $\beta = 2\gamma$, so [lemma 23.1](#) detects it. No single prescribed set of surreal cutoffs has this property uniformly over all enlarged workspaces.

23.4 Three tests, not one

The compact part separates three decisions that are easy to conflate. To decide whether a Hahn line bundle has any nonzero coherent meromorphic section, compute its valuation class ([theorem 20.1](#)). To decide whether a finite moving divisor is principal, compute its ordinary divisor class and its positive Abel functional ([theorem 21.5](#)). For $g > 0$, a one-scale formal test is faithful exactly when the scale is cofinal in the value group, that is when $\Delta = \Gamma$; for $g = 0$ it is always faithful ([theorem 23.3](#)).

None subsumes another. A nonzero valuation class prevents every meromorphic section, even in high degree ([example 20.3](#)). A nonzero infinitesimal class may admit many meromorphic sections while admitting no nowhere-vanishing holomorphic one. A nonzero infinitesimal class may also be invisible to every power of a smaller scale ([example 23.5](#)). And [example 22.3](#) shows that cancellation of the first displacement does not make the full infinitesimal Abel class vanish.

23.5 Why arbitrary-rank support control is substantive

In a rank-one complete valued field one proves an inverse formula by showing the terms approach zero rapidly. That argument is unavailable for general Γ : a term supported at ω is not reached by multiplying the scale 1 by ordinary integers. The proofs above use exact-exponent finiteness instead, which is why the same inverse matrix and residue formulas hold at ordinary small scales and at arbitrarily remote surreal scales while their detection by one chosen adic system may fail. Compactness supplies a second and independent form of control: a finite smooth cover with a uniform well-ordered support, and an ordinary Hodge projection with finite-dimensional image. Without the first, a partition-of-unity construction can create a forbidden union of supports; without the second, no finite matrix results. *Both are used explicitly, and neither is folded into a vague notion of analytic convergence.*

24 Interpretation, boundaries, and further questions

24.1 What has been established over the plane

The plane results are local-to-global theorems with exact obstructions. Ordinary analytic discreteness handles the positions of the standard parts; well ordering handles the distribution of surreal scales; neither condition can substitute for the other. The proofs succeed because ordinary Mittag–Leffler summation and ordinary Hermite interpolation introduce no new Hahn exponents, while the positive-support logarithm converts a nonlinear cluster equation into an additive principal-part problem — or, along the route of [remark 8.3](#), because a well-founded recursion on one fixed monoid can call the scalar interpolation lemma once per exponent.

Question	Exact positive result	Sharp boundary
Principal parts at a discrete set	Realizable exactly for a well-ordered union of principal-part supports	A descending exponent sequence prevents a coherent solution.
Finite moving clusters	Realizable exactly for a well-ordered union of symmetric coefficient supports	Individually well-defined roots need not form a principal divisor.
Prescribed data on a realizable divisor	The image is the support-restricted product $\mathcal{B}(P)$, with an explicit Neumann right inverse	Bounded 0/1 values can still be non-interpolable.
Additive gluing on the star cover	Splitting exactly when the singular support is well ordered	Removable coefficient germs impose no condition at all.
Ordinary-base local algebra	Every stalk is a principal ideal domain	It is nonlocal for every nonzero value group.
Simple infinitesimal motions	Uniform support gives a near-identity coherent automorphism	A bijection cannot split a multiple zero.
Line bundles over the plane	Vanishing exactly for trivial or cyclic value groups	Every noncyclic group gives continuum many independent classes.
Ordinary reduction	Positive-unit transition bundles have trivial integral reduction	Trivial reduction does not imply triviality over $\mathcal{H}_\Gamma$.

24.2 What has been established over a compact base

The compact results are classification and comparison theorems, and their mechanism is different: the base contributes finite-dimensional ordinary cohomology, compactness contributes one global well-ordered support, and the support calculus contributes an inverse that is an identity rather than a limit.

Question	Exact positive result	Sharp boundary
Cohomology of an ordinary bundle with Hahn coefficients	$H^q(X, V) \otimes_{\mathbb{C}}$ the coefficient ring, for $q = 0, 1$, with higher vanishing	Compactness used twice; false over the noncompact plane
Which rank-one modules exist	$\Gamma^{2g} \oplus \text{Pic}(X) \oplus \mathfrak{m}_{\Gamma}^g$ after choosing bases	The Pic_{lf} proof is independent of stalks; coordinate charts also identify Pic with Pic_{lf}
Nonzero coherent meromorphic section	Exists exactly when the valuation class vanishes	High degree with no section at all; H^1 in that sector is <i>not</i> computed
Cohomology when the valuation class vanishes	$\ker B$ and $\text{coker } B$ for a finite matrix over $\mathcal{O}_{\Gamma}$ with entries in $\mathfrak{m}_{\Gamma}$; Riemann–Roch	Integral torsion $\mathcal{O}_{\Gamma}/\lambda\mathcal{O}_{\Gamma}$ invisible over the field
Principality of a finite moving cluster	Ordinary principality <i>and</i> vanishing of a root-free Newton-sum Abel functional	Zero first displacement does not suffice: $y^2 = 1 + x^5$
What all powers of one scale see	Split surjection onto the limit, kernel $H^1(X, \mathcal{O}_X) \otimes_{\mathbb{C}} I_{\delta}$	Injective exactly for $g = 0$ or $\Delta = \Gamma$; prescribed cutoffs can be defeated after enlarging the workspace

24.3 What the conclusions do not say

All cohomology groups and line bundles here live over the ordinary complex plane or the specified compact Riemann surface, with its ordinary topology. *We have not assigned ordinary sheaf cohomology to the proper class of all surcomplex points*; each fixed coefficient group is a set, and every algebraic support and recursive construction is set-sized. *No ordinary real-parameter contour is claimed to be fine-continuous, and no transfinite support recursion is treated as a convergent fine-topological iteration.*

The divisor theorem concerns $\mathbb{C}^{\sharp}$, the finite surcomplex plane. It neither contradicts nor weakens the all-scale polynomial rigidity theorem of [1, §12]: a datum constructed by [theorem 8.2](#) has a coherent description at the ordinary scale and need not have coherent descriptions at every infinite surreal scale; and an infinite prescribed zero divisor cannot be the divisor of a polynomial, so these realizers cannot satisfy that stronger hypothesis. The word *entire* here means that every coefficient is ordinary entire and the evaluation is defined on $\mathbb{C}^{\sharp}$; it does not mean coherence at every surreal scale.

A Hahn principal-part solution on $\mathbb{C} \setminus A$ is not automatically a meromorphic function on every omitted monad with only finitely many poles. For moving divisors, finite polynomial cancellation supplies that extra structure and [theorem 11.1](#) is available; for the general [theorem 7.2](#) no such additional conclusion is claimed.

Our plane line-bundle classification is a vanishing/nonvanishing dichotomy plus a large explicit subspace, together with an exact dimension in the countable noncyclic case. It is not a computation of every element of $H^1(\mathbb{C}, \mathcal{H}_{\Gamma,+})$ for every noncyclic Γ . The exact Cousin criterion is for the

displayed puncture-and-disk cover; a similarly explicit criterion for arbitrary covers has not been established here. Nor does this article claim to settle canonical global exponentiation, sectorial summation, essential singularities in other surreal frameworks, or an all-scale transcendental theory. These limitations do not weaken the stated equivalences, whose proofs are complete within their hypotheses.

The compact part carries its own boundaries, and they are different ones.

The compact theorems are not statements about the plane. [Theorem 17.7](#) uses compactness twice — finite-dimensional ordinary harmonic spaces, and one global well-ordered support bound for smooth Hahn forms — and *it must not be applied to the noncompact plane by replacing X by $\mathbb{C}$ and setting $g = 0$* ([remark 17.8](#)). The global compact comparison does not follow from the plane results; local algebra does transfer through coordinate charts ([remark 18.5](#)).

Cohomology in the nonzero-valuation sector is not computed. [Theorem 19.2](#) and [corollary 19.4](#) apply only when $\nu(L) = 0$, because the positive perturbation argument of [proposition 19.1](#) has no global ordinary bundle and no global scalar positive gauge in the required form otherwise. [Theorem 20.1](#) proves $H^0(X, L) = 0$ when $\nu(L) \neq 0$, and *that vanishing determines neither $H^1(X, L)$ nor an Euler characteristic there*; see [remark 18.4](#). The plane part has no such gap, since over $\mathbb{C}$ the valuation factor vanishes identically.

The compact Abel theorem covers finite polynomial cluster Cartier data only. It is not asserted that every section of $\mathcal{D}_\Gamma$ admits such a presentation, nor that an arbitrary meromorphic-coefficient Hahn series has a classical finite zero count ([remark 21.7](#)). Divisibility of Γ makes K_Γ algebraically closed and guarantees that every cluster splits there. Without it an individual cluster may still split, and roots always exist in a divisible Hahn enlargement; the root-free criterion requires neither splitting nor enlargement.

No comparison functor is constructed to an algebraic curve over K_Γ , or to a rigid-analytic, Berkovich or logarithmic Picard functor. Such identifications would require comparison functors with stated hypotheses, and the valuation-monodromy examples of [example 20.3](#) already rule out the most naive degree- and section-preserving comparison. *The compact objects are not compact subsets of the fine surcomplex plane*: the base topology is ordinary complex topology, no fine-topological convergence occurs, and no general surreal version of Hodge theory or Serre duality is proved or assumed.

No uniform decision procedure is asserted for unrestricted surcomplex symbolic inputs. Each fixed Hahn coefficient uses finitely many ordinary operations, some of which are analytic solution operators rather than effective finite-dimensional calculations; a uniform algorithm over arbitrary symbolic coefficient families would need additional representability and equality assumptions that are not made here.

24.4 Further precise directions

The injections [\(13.7\)](#) and [\(13.11\)](#) leave a structural problem: characterize the cokernel, or find a useful normal form for an arbitrary class in $H^1(\mathbb{C}, \mathcal{K}_{\Gamma,+})$. The proof of [theorem 14.4](#) counts classes for countable Γ but supplies no such normal form, and for uncountable value groups the lower bound $2^{\aleph_0}$ need not be optimal; determining the dimension in terms of cardinal and order invariants of the positive cone would refine the dichotomy.

A second direction is the finite-rank vector-bundle theory: matrix-valued positive transition functions no longer reduce to an abelian logarithm identity, so a noncommutative support recursion must replace the rank-one argument.

A third direction was the same package over other ordinary bases. In the rank-one abelian case that direction is now taken up here, for a *compact* base, in [sections 16 to 23](#). That part is not a generalization of the plane part and does not contain it: compactness is load-bearing at two separate points ([remark 17.8](#)), and the plane’s obstructions arise precisely from the infinitely many centres a compact base cannot carry. *The general ordinary open Riemann surface, neither the plane nor compact, remains open here*: there the support recursions would have to be compared with scalar interpolation and with line-bundle obstructions on the base, and neither part of this article supplies that comparison. Two concrete questions the compact part leaves behind are the computation of H^1 for a pure monomial local system t^ρ with $\rho \neq 0$ — which would measure the cohomological cost of the extra Γ^{2g} factor, and for which the ordinary-base common-support condition rather than the abstract local system must enter — and a comparison of the zero-valuation sector with an algebraic or non-Archimedean analytic Picard functor for the same ordinary curve after scalar extension; [theorem 18.3](#) makes the source explicit but does not construct such a functor or prove it fully faithful.

Finally, the several-variable analogue. The present proofs use finite polynomial remainders in one distinguished local coordinate and isolated clusters indexed by a discrete ordinary set. Positive-dimensional analytic sets would require coherent parameter-dependent division and compatibility between local finite modules, and an exact global support criterion for higher-codimension analytic sets needs more than the one-variable logarithmic reduction. This article provides a test case and explicit obstructions that any such extension should preserve; it does not claim that result.

25 Conclusion

The globally coherent category has a precise compatibility boundary. Infinitely many infinitesimal data are compatible exactly when the appropriate coefficient supports can coexist inside one well-ordered set. For divisors the relevant coefficients are the symmetric coefficients of the local cluster polynomials; for additive principal parts they are the principal-part exponents themselves, and more precisely only those carrying nonremovable singularities; for prescribed data on a divisor they are the coefficients of the intrinsic polynomial remainders; for line bundles the positive logarithm exhibits the same issue as a first-cohomology class.

This yields a global Weierstrass theorem with an exact converse, a support-restricted Chinese remainder theorem, explicit principal divisors with nonprincipal effective sub-divisors, an exact Cousin criterion, and a complete vanishing dichotomy for the plane Picard group together with its exact dimension over countable noncyclic value groups. The phenomena already occur in the countable divisible workspace $\Gamma = \mathbb{Q}$, and they require neither proper-class summation nor any problematic notion of fine-topological convergence.

Over a compact ordinary base the same coefficient sheaf behaves differently and is classified completely. Finitely many clusters make the well-ordering question vacuous, and what remains is cohomological: a valuation monodromy in Γ^{2g} that is exactly the obstruction to a nonzero coherent meromorphic section, an ordinary line bundle, and an infinitesimal class in $\mathfrak{m}_\Gamma^g$ that a finite matrix over $\mathcal{O}_\Gamma$ turns into computable cohomology and a Riemann–Roch identity. Principality of a finite moving divisor is decided by ordinary principality together with a root-free Abel functional, and a genus-two witness shows that cancelling the first symmetric displacement is not enough. Finally, if $g > 0$, any prescribed set of surreal cutoffs can be defeated in a larger set-sized workspace. This quantifies over workspaces: in one fixed nonzero Γ , allowing all positive cutoffs detects every class, while powers of a single scale do so exactly when that scale is cofinal.

The two parts share their coefficient sheaf, their support calculus and their insistence that symmetric coordinates, not labelled roots, are what a global criterion may test. Their local algebra also agrees in coordinate charts; their global existence and classification theorems retain the stated base hypotheses.

A Support and dependency audit

The table records the operations at which infinite formal expressions occur, together with the set that controls them. A support in the right column is a *containing* support; cancellation may make the actual support smaller. Throughout, S is an input support and E a well-ordered positive set with $M = E^*$.

Construction	Controlling support and finiteness mechanism
Evaluation at $c + \epsilon$	$S + (\text{supp } \epsilon)^*$; words account for the Taylor powers, and finite exponent decompositions justify the regrouping.
Positive inverse, exponential, logarithm	M for positive well-ordered E ; finitely many exponent words contribute at each exponent.
Local preparation at any ordinary centre	T^* with T the relative positive support of the section; <i>the same monoid works at every centre</i> .
Deformed division	$S + E^*$, independently of the cluster multiplicity; each application of T inserts one exponent from E .
Global divisor construction	$E(P)^*$ throughout the additive solution, the exponentiation and the gluing; alternatively one scalar entire coefficient chosen per exponent.
Logarithm of a cluster polynomial	$E(P)^*$; finitely many coefficients at each centre keep every Hahn coefficient a finite principal part.
Interpolation Neumann series	$S + E(P)^*$; finite words and their finitely many block partitions bound the contributing powers of NL .
Moving Mittag–Leffler	$S_R + E(P)^*$ after multiplication by local units and deformed division.
Mittag–Leffler at ordinary centres	The union of the principal-part supports; ordinary analytic choices at a fixed exponent introduce no new exponent.
Cousin splitting	Central support $\Sigma(b)$; each disk support lies in $\Sigma(b) \cup \text{supp } b_n$.
Near-identity inverse	$(\text{supp } E)^*$; external positive exponents make the coefficient recursion triangular.
Cyclic-group additive coboundary	The universal set $\{\delta, 2\delta, \dots\}$; every coefficientwise solution is automatically admissible.
Nontriviality arguments	A forced infinite descending positive sequence would have to sit inside one legitimate support.
<i>Compact base (sections 16 to 23); E now comes from a <i>finite</i> cover, never from a hypothesis on Γ.</i>	
Global smooth Hahn forms on X	One well-ordered support from a finite subcover (lemma 17.2); false on a noncompact base.
Coefficientwise Green and Hodge operators	The same global support; a <i>fixed</i> cutoff solves $\bar{\partial}$ for every coefficient on one fixed smaller disk.
Operator Neumann inverse $(1 + A)^{-1}$	$S + E^*$ with unbounded coefficient operators; finitely many words, hence finitely many operator applications at each exponent.

Construction	Controlling support and finiteness mechanism
Positive Dolbeault normal form	E for α , E^* for the smooth gauges; the harmonic correction $\bar{\partial}c$ adds no exponent.
Transferred matrix $B = p_1\delta(1 + h\delta)^{-1}i_0$	$E^* \setminus \{0\}$; a finite matrix because ordinary harmonic spaces are finite-dimensional.
Logarithm of a cluster polynomial, Newton form	$E^* \setminus \{0\}$; finitely many negative powers of u at each fixed exponent.
Abel functional and its residue formula	$E^* \setminus \{0\}$; the residue at a fixed exponent uses finitely many ordinary Taylor coefficients.
One-scale inverse limit	Convexity of Δ makes a compatible family's union of supports well ordered.

Only the ordinary scalar constructions use analytic convergence: normal convergence of correcting-polynomial series, of Weierstrass products, and smooth convergence of corrected Cauchy transforms. In all Hahn constructions, well ordering *and* finite coefficient multiplicity are checked separately. The dependency chains are

support calculus $\implies$ units, logarithms, evaluation, preparation, division,
 preparation + scalar interpolation $\implies$ global divisor realization + deformed interpolation,
 these two constructions $\implies$ moving Mittag-Leffler,
 preparation $\implies$ principal-ideal-domain stalks,
 PID stalks $\implies$ tensor-invertible = rank-one locally free,
 scalar Cousin + exact singular-support criterion $\implies$ descending-scale classes,
 these classes + cyclic case $\implies$ the line-bundle dichotomy.

The connection between the last two chains is the explicit divisor-transition cocycle of [proposition 15.1](#). The principal-ideal-domain argument uses the Euclidean algorithm on finite-degree polynomials and the least element of a set of nonnegative integers; it does not invoke an unproved Noetherian property of an infinite coefficient ring. The plane Picard argument does not silently assume local stalks: local rank-one triviality is established separately in [corollary 6.3](#).

The compact part has its own chain, which shares only the first link:

support calculus $\implies$ operator Neumann inverse,
 compactness $\implies$ one global support + finite harmonic spaces $\implies$ compact comparison,
 unit splitting + compact comparison $\implies$ compact Picard classification $\implies$ divisor-class sequence,
 positive normal form + Hodge identities $\implies$ the finite matrix,
 the finite matrix $\implies$ Riemann-Roch + meromorphic criterion,
 Newton logarithm + ordinary Serre duality $\implies$ the Abel criterion,
 truncated unit splitting + convexity of Δ $\implies$ the one-scale kernel.

Potential failure points explicitly avoided there: no infinite-dimensional smooth Hahn space is replaced by an algebraic tensor product; compactness is established *before* a global Green operator is applied to a Hahn family; meromorphic coefficient pole orders are allowed to grow with the exponent while the ordinary domain is fixed; the exponential is used only on positive support; *no claim that*

$n\delta$ is cofinal is used unless $\Delta = \Gamma$ has already been imposed; the truncations of [section 23](#) are formed in the integral sheaf and not in the field sheaf, where they would be zero; and the classical results are used on ordinary complex curves only.

B Reproducible symbolic checks

The source archives supplied two plane verification scripts, maintained as `code/09-moving-divisors-picard-dichotomy.py` and `code/10-cohomological-obstructions.py`. They use exact symbolic algebra to check finite truncations and polynomial identities illustrating the proofs: a mixed principal-part logarithm and exponential through degree six in t ; cancellation for two finite root clusters; the two-sided inverse of $z + tz^2$ through degree seven, against the Catalan recurrence; complementary root-cluster factors for degrees two through ten; finite Lagrange interpolation of positive-support motions; a truncated instance of the preparation recursion; finite instances of the global divisor recursion and of local divisibility; root-selector idempotents; and finite residue identities. For the preparation check one takes

$$F(u) = u^2 + t(1 + u^3) + t^2(2u + u^4),$$

constructs A_n of degree less than two and polynomial B_n from $h_n = F_n - \sum_{i=1}^{n-1} A_i B_{n-i}$, $A_n = R_2 h_n$, $B_n = D_2 h_n$, and checks $(u^2 + A)(1 + B) = F$ through degree six — a finite example of the well-founded recursion in [theorem 5.1](#). For exponential coefficients the exact recurrence

$$e_0 = 1, \quad ne_n = \sum_{k=1}^n k b_k e_{n-k}, \quad \text{Exp}_H\left(\sum_{k \geq 1} b_k t^k\right) = \sum_{n \geq 0} e_n t^n,$$

avoids numerical error and makes every truncation order explicit.

For the compact part the source script is maintained as `code/11-compact-curve-picard-classification-verification.py`, an ordinary Python program using SymPy, whose recorded execution passed **53 of 53 exact checks** with no floating-point tolerances and no network access. It checks the Newton/logarithm generating identity for monic polynomials of degrees two and three through power nine; the displayed initial coefficients of the genus-two Abel functionals of [example 22.3](#), namely $-1/6$ at s^3 and $-5/128$ at s^8 for η_0 and 1 at s , $1/16$ at s^6 , $35/1408$ at s^{11} for η_1 ; and a purely algebraic three-dimensional two-term contraction model with

$$d = h = \text{diag}(0, 1, 1), \quad i = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad p = i^\top,$$

$$\delta = t \begin{pmatrix} 0 & 1 & 2 \\ 3 & 1 & 0 \\ 1 & -1 & 2 \end{pmatrix} + t^2 \begin{pmatrix} 1 & 0 & -1 \\ 0 & 2 & 1 \\ 1 & 0 & 0 \end{pmatrix},$$

for which the transferred 1×1 matrix is exactly

$$B(t) = \frac{t^2(2t^4 + 5t^3 + 4t^2 - 12t - 4)}{5t^3 + 4t^2 + 3t + 1}. \quad (\text{B.1})$$

The program verifies that this equals the Schur complement, verifies $DI_0 = iB$, $HD = 1 - I_0p$, $DH = 1 - iP_1$ and $P_1D = Bp$ with $I_0 = (1 + h\delta)^{-1}i$, $H = (1 + h\delta)^{-1}h$, $P_1 = p(1 + \delta h)^{-1}$, and

compares the first five positive coefficients with the truncated Neumann expansion (19.3). *This is an algebraic model, not an assertion that these matrices arise from a particular curve*; the leading order t^2 in (B.1) illustrates why a linearized obstruction need not be the whole transferred matrix. Only the first few coefficients of (22.5)–(22.6) were confirmed symbolically; the infinite formulas and their consequence for divisors are proved in section 22.

Remark B.1 (A defect in the shipped compact script, recorded). The compact script writes `verification_results.json` beside itself. In the original delivery this could overwrite the supplied results; after repository ingestion the historical results live under `data/`, but a rerun still overwrites any previous generated sibling file. Run a temporary copy to keep both the report tree and historical evidence untouched. Worse, its check helper raises an `AssertionError` on the first nonzero residual while the results file is written only after every check has passed. *The results file can therefore never structurally record a failure*: a failing run aborts before writing and leaves the previous all-pass file untouched. The recorded all-pass JSON is consequently evidence that some run completed, not evidence that the current source passes. Nothing in this article depends on that file; the checks it describes are reproducible by running the script, and in any case they certify only finite instances.

These tests are deliberately separate from the proofs. No finite symbolic computation establishes the nonexistence of a well ordering for an infinite descending sequence, the plane Picard-group dimension, the universality of the plane dichotomy, the infinite support lemma, the compact sheaf comparison, the compact Picard classification, the nonexistence of a meromorphic section, or the validity of an arbitrary-support transfinite recursion; those are proved in the main text. The scripts check signs, coefficients and finite algebraic instances. They are not a proof assistant, a machine verification of the general arguments, or a novelty checker, and do not establish Lean coverage. The repository’s maintained formalization index records the current coverage; the global sheaf and Picard results remain unformalized.

C Provenance of the merged material

Both source manuscripts prove lemmas 2.1 and 14.2, propositions 3.3 to 3.5, and theorems 5.1, 8.2 and 14.3, the two discriminating examples 9.1 and 9.2, and the realization torsor; they also both insist on the boundaries recorded in sections 1.3 and 14.5. Those are printed once.

Manuscript A is the base. Beyond the shared core it supplies theorem 6.1 and corollary 6.3; theorem 14.4, with the almost-disjoint prefix family and the cocycle-counting upper bound; the whole of section 12, including theorem 12.1 and corollary 12.2 and the Catalan inverse; corollary 9.3 with the explicit complementary factor and section 9.4; the series (9.3) of the explicit construction, whose Weierstrass product (9.2) both manuscripts print (the convergence estimate given here is Manuscript B’s); corollary 9.3’s nonprincipal effective sub-divisor, which Manuscript B does not treat; theorem 7.2 in the fixed-centre form, with corollary 7.3; and theorem 13.3, example 13.6, and proposition 13.7.

Manuscript C supplies the whole of the compact-base part, sections 16 to 23: lemmas 17.2 and 17.4, proposition 17.6, and theorem 17.7 with the compactness warning remark 17.8; the classification theorem 18.3 with the normal cocycle (18.5) and the nonzero-valuation gap remark 18.4; the finite matrix, propositions 19.1 and 19.5, theorem 19.2, and corollaries 19.4 and 19.6, including the elliptic integral torsion (19.9); the meromorphic criterion theorem 20.1 with corollary 20.2 and example 20.3; the root-free Abel package lemma 21.2, definition 21.3, proposition 21.4, and theorem 21.5 with remarks 21.6 and 21.7; the three compact examples examples 22.1 to 22.3; and the one-scale comparison lemmas 23.1 and 23.2 and theorem 23.3 with example 23.5 and corollary 23.6. It also

supplies the compact rows of [appendix A](#) and the compact half of [appendix B](#). Its original restriction to $\mathrm{Pic}_{\mathrm{lf}}$, without a stalk hypothesis, is recorded in [remark 18.5](#); the maintained review there adds the coordinate-chart consequence of the PID theorem in the merged text; its negative cofinality remark is kept as the second paragraph of [remark 17.5](#), alongside the third paragraph of [remark 2.2](#) which says the same thing for the plane. Manuscript C was written against this repository rather than against [\[1\]](#) and shares no proof with Manuscripts A and B beyond the support calculus and the unit splitting, both of which it proves for itself and which are printed here once, in the plane part.

Manuscript B supplies the constructive scalar section [section 4](#) ([lemmas 4.1 to 4.3](#)); [lemmas 5.3 and 5.4](#); the recursive route to the divisor theorem recorded in [remark 8.3](#); the whole of [section 10](#) — [proposition 10.1](#), [theorem 10.2](#), [corollary 10.3](#), and [example 10.4](#) — and with it the identification of the correct interpolation target as a support-restricted product; [theorem 11.1](#) and [example 11.2](#); the exact Cousin criterion [theorem 13.1](#) with the singular support and [remark 13.2](#); [corollaries 13.4 and 13.5](#) and [proposition 14.6](#) in the containment form; the positive half of [proposition 6.2](#), that the nonnegative-support subsheaf is local with residue field $\mathbb{C}$ — the integral model; and the $\mathrm{Pic}_{\mathrm{lf}}$ distinction with its Stacks citation, preserved in [remarks 6.4 and 14.5](#).

D Notation

Symbol	Meaning
$\mathbf{SC} = \mathbf{No}[i]$	Algebraically closed surcomplex class field.
t^γ	Conway monomial $\omega^{-\gamma}$; not an independently chosen exponential.
$K_\Gamma, \mathfrak{m}_\Gamma, \mathcal{O}_\Gamma$	Fixed complex Hahn field, its infinitesimals, its finite elements.
$\mathcal{H}_\Gamma$	Sheaf of Hahn data with ordinary holomorphic coefficients on a common ordinary base.
$\mathcal{H}_{\Gamma,+}, \mathcal{H}_{\Gamma,\geq 0}$	Positive-support additive subsheaf and nonnegative-support subring sheaf.
$U^\sharp, F^\sharp, \mu(c)$	Infinitesimal thickening, Taylor–Hahn evaluation, the monad of c .
E^*	Additive monoid generated by a positive well-ordered set E , including zero.
$P_c, E(P), M$	Monic cluster polynomial at c ; symmetric support; $M = E(P)^*$.
$\mathcal{R}_P, \mathcal{B}(P)$	Simultaneous deformed remainder map and its support-restricted target algebra.
$\Sigma(b)$	Singular support of additive Cousin data on the star cover.
R_c	Stalk of $\mathcal{H}_\Gamma$ at an ordinary base point — not a fine germ at one surcomplex point.
$\mathcal{V}_\Gamma/\mathcal{W}_\Gamma$	Infinitesimal sequences modulo uniformly supported sequences.
red, st	Reduction $\mathcal{H}_{\Gamma,\geq 0} \rightarrow \mathcal{O}$ (synonyms: residue, shadow, standard-part map) and the scalar standard part.
Pic, Pic _{lf}	Tensor-invertible modules; locally free rank-one modules. Equal on the plane by corollary 6.3 and on a Riemann surface by remark 18.5 ; compact statements retain Pic _{lf} .
<i>Compact base only</i> (sections 16 to 23).	
X, g	Fixed compact connected ordinary Riemann surface and its genus.
$\mathcal{M}_\Gamma, \mathcal{D}_\Gamma$	Hahn sheaf with ordinary meromorphic coefficients; its coherent Cartier quotient $\mathcal{M}_\Gamma^\times/\mathcal{H}_\Gamma^\times$.
$\mathcal{A}_\Gamma^{0,q}$	Smooth Hahn $(0, q)$ -forms, with ≥ 0 and $+$ support restrictions.
$\nu(L), L_0, \xi(L)$	Valuation class, ordinary component, positive logarithmic class of a Hahn line bundle.
$i_q, p_q, h, \delta, T, B$	Ordinary harmonic inclusion and projection, Hodge homotopy, multiplication by $\alpha, (1 + h\delta)^{-1}$, and the transferred matrix $p_1\delta Ti_0$.
$p_k(P), \mathcal{A}_D(\eta)$	Newton power sums of a monic cluster polynomial (defined by integer recurrences, not by roots); the infinitesimal Abel functional.
$\Delta, I_\delta, \mathcal{R}_n$	Convex subgroup generated by δ ; the invisible ideal $\bigcap_n t^{n\delta}\mathcal{O}_\Gamma$; the truncation $\mathcal{H}_{\Gamma,\geq 0}/t^{n\delta}\mathcal{H}_{\Gamma,\geq 0}$.
$\mathcal{O}_\Gamma$ (compact part)	The valuation ring of K_Γ , written R_Γ in the compact source manuscript; a ring of scalars, never the structure sheaf $\mathcal{O}_X$.

References

- [1] *Surcomplex Analysis: Infinitesimal Calculus, Hahn-Coherent Holomorphy, and Contour Theory*. User-supplied manuscript, September 21, 2026, file `surcomplex_analysis(3).tex`. The present article uses its §5 (coherent data, evaluation and support-preserving gluing), §7 (preparation, division, root lifting and pole-free fractions), its moving-pole residue theorem, and its distinction

- between finite-scale and all-scale coherence. Authorship and publication status were not supplied, and none is inferred from the filename.
- [2] Lars V. Ahlfors, *Complex Analysis*, 3rd edition, McGraw–Hill, 1979. Classical Weierstrass products, Mittag–Leffler theory, and elementary holomorphic function theory.
 - [3] Alessandro Berarducci and Vincenzo Mantova, “Transseries as germs of surreal functions,” *Transactions of the American Mathematical Society* **371** (2019), 3549–3592. [arXiv:1703.01995](#).
 - [4] Ovidiu Costin and Philip Ehrlich, “Integration on the surreals,” *Advances in Mathematics* **452** (2024), article 109823. [arXiv:2208.14331](#).
 - [5] Marius Crainic, “On the perturbation lemma, and deformations,” preprint, version 1 (2004), §2. [arXiv:math/0403266v1](#). The two-term transferred complex of [theorem 19.2](#) is an instance of this classical lemma; no discovery of the lemma is claimed here.
 - [6] Jean-Pierre Demailly, *Complex Analytic and Differential Geometry*, online book. Chapter V, §11 (holomorphic vector bundles) and Chapter IX (cohomological vanishing on Stein spaces). [Author’s full text](#), accessed September 21, 2026.
 - [7] Philip Ehrlich and Elliot Kaplan, “Surreal ordered exponential fields,” *The Journal of Symbolic Logic* **86** (2021), no. 3, 1066–1115. [doi:10.1017/jsl.2021.59](#); [arXiv:2002.07739](#). The surcomplex exponentiation discussion is in §11 of the arXiv version. Used to delimit prior work, not as an input to any proof here.
 - [8] Mark Green and Robert Lazarsfeld, “Higher obstructions to deforming cohomology groups of line bundles,” *Journal of the American Mathematical Society* **4** (1991), 87–103. [doi:10.1090/S0894-0347-1991-1076513-1](#). Cited to place the finite obstruction matrix in an existing tradition, not as an input to any proof here.
 - [9] Harry Gonshor, *An Introduction to the Theory of Surreal Numbers*, London Mathematical Society Lecture Note Series **110**, Cambridge University Press, 1986.
 - [10] Curtis T. McMullen, *Advanced Complex Analysis*, Harvard University Math 213a course notes, November 30, 2023. [Author’s notes](#). See §3 for ordinary entire and meromorphic functions, and Theorem 3.26 for Mittag–Leffler.
 - [11] Jörn Müller and Alexander Strohmaier, “The theory of Hahn-meromorphic functions, a holomorphic Fredholm theorem, and its applications,” *Analysis & PDE* **7** (2014), no. 3, 745–770. [doi:10.2140/apde.2014.7.745](#); [arXiv:1205.0236](#). Definitions 2.3 and 2.5 are the normally convergent generalized expansions from which the present coefficient sheaf is explicitly distinguished.
 - [12] Samouil Molcho and Jonathan Wise, “The logarithmic Picard group and its tropicalization,” [arXiv:1807.11364](#), version 5. The logarithmic-curve category and its bounded-monodromy condition are distinct from the ordinary-base Hahn category considered here; *no equivalence is assumed and no priority over that theory is claimed*. The Hahn–Tate report of this collection, [docs/surcomplex/hahn-tate-uniformization/](#), credits Section 5.1 of this paper as the precedent for its bounded-period quotient.
 - [13] Raghavan Narasimhan, *Compact Riemann Surfaces*, Lectures in Mathematics ETH Zürich, Birkhäuser, 1992. [doi:10.1007/978-3-0348-8617-8](#). The chapters on Dolbeault’s lemma, Hodge theory, Serre duality, Riemann–Roch, and the Jacobian and Abel’s theorem supply every ordinary compact-curve input of [sections 17 to 23](#).

- [14] B. H. Neumann, “On ordered division rings,” *Transactions of the American Mathematical Society* **66** (1949), 202–252. The support lemma underlying [lemma 2.1](#).
- [15] Bjorn Poonen, “Maximally complete fields,” *L’Enseignement Mathématique* (2) **39** (1993), 87–106. [Author’s full text](#). Corollary 4 supplies the algebraic-closure statement for Hahn fields with divisible value group.
- [16] The Stacks Project Authors, *The Stacks Project*. [Tag 02FQ](#) (abelian torsors and first cohomology); [Tag 09NT](#) (first cohomology and invertible sheaves). Accessed September 21, 2026. The local-stalk hypothesis in the latter is why [corollary 6.3](#) is proved rather than assumed, and why [remark 6.4](#) keeps the $\mathrm{Pic}_{\mathrm{lf}}$ distinction on record.
- [17] Lou van den Dries and Philip Ehrlich, “Fields of surreal numbers and exponentiation,” *Fundamenta Mathematicae* **167** (2001), 173–188. [doi:10.4064/fm167-2-3](#); erratum, *Fundamenta Mathematicae* **168** (2001), 295–297.