

OEIS RESEARCH REPORT

Fixed-composition excursions

D-finiteness, transcendence, and all-order
asymptotics for OEIS A215561

With an explicit correction for A215570
and asymptotic inversion

Merged from two independent manuscripts of the same day:

Fixed-composition excursions and *Balanced Late-Growing Permutations*; Parts IV and V
from a third and a fourth; Part VI from two more

Prepared for Vladimir Reshetnikov

October 1, 2026; Parts IV and V added October 2, 2026;

Part VI added October 5, 2026

Abstract

Let $A_r(n)$ count words containing n copies of each letter in $\{1, \dots, r\}$ whose every prefix has sum at most its average over all such words. We prove the fixed-row asymptotic conjecture recorded in OEIS A215561, identify its constant as a positive algebraic number, and establish a complete expansion in inverse powers of n . The central formula is

$$A_r(n) \sim \frac{\kappa_r}{rn} \frac{(rn)!}{(n!)^r}, \quad \kappa_r = \exp\left(\sum_{m \geq 1} \frac{\mathbb{P}(S_m = 0)}{m}\right),$$

where S_m is the centered, uniformly weighted walk associated with the alphabet. A finite kernel-root product computes κ_r .

We also prove that every fixed row is D-finite, while its ordinary generating function is transcendental for every $r \geq 3$. This separates recurrence existence from algebraicity and from the specific recurrence conjecture for A215570 (proved in Part IV). An explicit first correction for that sequence, a fixed-composition contact law, and an all-order Lambert- W inversion procedure are derived. Exact enumeration and symbolic checks accompany the proofs.

Part II adds the second manuscript's independent results: an elementary one-large-summand proof of the leading term for every finite labelled step set and rational balanced composition, a second all-orders route with a Gaussian-moment algorithm, second-order terms for $r = 2, 3$, a single giant primitive component with independent finite decorations and a heavy-tailed gap, eventual strict log-convexity, and an explicit rounding bound.

Part IV, from a third manuscript, proves the order-three recurrence that Kauers and Koutschan conjectured for A215570, by a positive Legendre-coefficient formula and a rational telescoping certificate, and computes $\alpha_{5,2}, \dots, \alpha_{5,8}$ exactly. Part V, from a fourth, adds the first correction $\alpha_{4,1}$ of A215562. Part VI, from two more, proves the main theorems a fifth and a sixth time and computes $\alpha_{5,2}$ without the recurrence.

Status. This is a newly written mathematical research manuscript, not a peer-reviewed publication or a Lean-verified development. Classical kernel and diagonal theorems are credited separately from the applications and deductions proved here. No claim of exhaustive bibliographic priority is made. Parts I and II do *not* prove the particular order-three recurrence conjecture of Kauers and Koutschan; Part IV (added October 2) does. The second manuscript describes itself as an “AI-assisted mathematical research draft. Unrefereed; not formally verified.”

Contents

1	Sources, merge decisions and conventions	6
1.1	Two independent manuscripts	6
1.2	What each source contributed	6
1.3	Where the merge chose	7
1.4	Repository context and re-derived material	8
1.5	Notation	9
I	Fixed-composition excursions	9
2	The problem, its significance, and the source audit	9
2.1	A natural multiset-permutation family	9
2.2	What is and is not resolved	11
2.3	Relationship to ProveIt	11
3	Main theorems	12
4	The exact excursion generating function	13
4.1	Encoding the prefix condition	13
4.2	The small-root formula, with a proof	14
4.3	Why D-finiteness follows	14
4.4	A concrete algebraic certificate for the fifth alphabet	15
5	The bridge logarithm and the algebraic constant	16
5.1	An exact identity retaining every composition	16
5.2	The critical value	17
6	Fixed-composition singularity analysis	18
6.1	Removing the two linear constraints	18
6.2	The moving critical point and its Hessian	18
6.3	A uniform square-root expansion	19
6.4	Why other phases do not contribute	20
6.5	Transfer and the Gaussian integral	20
6.6	Remainders and all orders	21
7	Explicit leading constants	22
8	Why every nontrivial higher row is transcendental	23
9	The first correction for the fifth row	24
9.1	A differential formula for the ratio correction	24

9.2	Exact ingredients for $r = 5$	25
10	A fixed-composition contact law	26
11	Inverting the asymptotic expansion	27
11.1	An exact Lambert core	27
11.2	Corrections and a remainder statement	28
11.3	Integer thresholds are a separate operation	28
12	Unequal but centered compositions	29
13	Reproducible checks and their scope	30
13.1	Exact finite enumeration	30
13.2	Asymptotic and inverse diagnostics	30
13.3	What the computations do not certify	31
II	The convolution route, general step sets and the shape of an excursion	31
14	Labelled steps and the log-excursion series	32
14.1	Labelled steps and composition	32
14.2	The log-excursion series	32
15	A directional convolution theorem	32
15.1	Critical multinomial weights	32
15.2	One large summand	33
16	The general balanced-composition formula	34
17	A second route to the complete expansion: lattice coordinates	35
17.1	Composition coordinates and the unique Fourier saddle	35
17.2	The critical point and the square-root expansion	36
17.3	An all-orders coefficient algorithm	37
18	First corrections by Gaussian moments	38
18.1	A coordinate-invariant first-correction formula	38
18.2	Computing the local kernel terms	39
18.3	Three evaluated cases	39
19	Primitive excursions and the giant component	40
19.1	A tight gap with an infinite-mean limit	41
20	Ratios, curvature and an explicit rounding bound	42
20.1	Ratios and eventual strict log-convexity	42

20.2	The inverse dominant sector	42
20.3	Discrete thresholds and the rounding boundary	42
21	Source 42's computations	43
21.1	Exact dynamic programming	43
21.2	Independent coefficient and algebra checks	43
21.3	Claims that remain outside the certificate	44
III	Research directions and conclusion	44
22	Further research questions and topics	44
22.1	Certify the explicit A215570 recurrence (resolved)	44
22.2	Find and explain the fourth-row recurrence	45
22.3	Automate all-order coefficient extraction	45
22.4	Determine sharp asymptotic error constants	46
22.5	Determine the exponentially small sectors	46
22.6	Let the alphabet grow	46
22.7	Resolve the mean size of the non-giant remainder	47
22.8	Approach the boundary of the composition simplex	47
22.9	Refine path statistics under exact composition	47
22.10	Formalize the exact-to-analytic interface	47
23	Conclusion	48
IV	The Kauers–Koutschan recurrence and the five-symbol row	48
24	Provenance, overlap and conventions of Part IV	49
24.1	A third manuscript	49
24.2	What the third manuscript re-proves	49
24.3	Conventions of Part IV	51
25	A positive coefficient formula for A215570	51
25.1	The sextic in compact form	51
25.2	Zero-step deletion and Lagrange inversion	52
25.3	A Legendre coefficient	52
26	Proof of Kauers–Koutschan Conjecture 15	53
26.1	A rational residue representation	54
26.2	The certificate identity	54
26.3	Exact generation, and what is not asserted	55

27 The exact coefficients $\alpha_{5,2}, \dots, \alpha_{5,8}$	55
27.1 A second route to the expansion, a third to $\alpha_{5,1}$	55
27.2 The triangular algorithm	56
28 Inversion, rounding and repository machinery	58
29 Source 75-03's computations	58
V Equal-content ballot words: the first correction for A215562	59
30 Provenance, overlap and conventions of Part V	60
30.1 A fourth manuscript	60
30.2 What the fourth manuscript re-proves	60
30.3 Conventions of Part V	61
31 The first correction for A215562	61
32 A second route to the global-arc bound	62
33 Literature pointers and further remarks	63
34 Source 77-44's computations and questions	64
VI Direct excursion proofs: every fixed alphabet, and the second correction of A215570 without the recurrence	64
35 Provenance, overlap and conventions of Part VI	65
35.1 Two bundle reports	65
35.2 What the two reports re-prove	66
35.3 Conventions of Part VI	68
36 Every fixed alphabet with the extreme steps unmarked (source 101-90)	69
36.1 Zero steps for odd alphabets	69
36.2 Extreme-unmarked coordinates and their Hessian	70
36.3 The full torus and the exact periodicity factor	71
36.4 The local saddle, the coefficient rule, and D-finiteness	72
36.5 Inversion for every row	74
37 The five-letter row with the zero steps deleted (source 101-86)	74
37.1 Two marks on the zero-free walk	74
37.2 The leading constant of the zero-free count	75
37.3 The coefficient rule in these coordinates	76

37.4	The second correction without the recurrence	76
37.5	Inversion, and monotonicity at every index	78
38	Computations, the lattice-peak theorem, stale claims and questions	79
38.1	Source 101-86 and the lattice-peak transfer theorem	79
38.2	The two packages	80
38.3	Stale and coordinate-dependent claims of the two reports	82
38.4	Further questions and research	82
A	A reproducible all-order construction	83
B	Package contents and build instructions	84
C	The full rational telescoping certificate (source 75-03)	85
C.1	Coefficients of $\mathcal{U}_n$	85
C.2	Coefficients of $\mathcal{V}_n$	86
C.3	How to verify the certificate without a guesser	86

1 Sources, merge decisions and conventions

1.1 Two independent manuscripts

This report merges two manuscripts of batch 73 of the repository’s incoming-reports intake (cluster O2). Both are dated October 1, 2026, both were prepared for Vladimir Reshetnikov, and both were placed in commit 6e193dd4f.

- **Source 61** (batch-73 manuscript 61): *Fixed-composition excursions. D-finiteness, transcendence, and all-order asymptotics for OEIS A215561. With an explicit correction for A215570 and asymptotic inversion* (a 1,505-line source and a 25-page PDF). Archive `oeis_balanced_excursions.zip`, arrival commit `aa43cc555`; it pins the repository at `40b17f2fc`.
- **Source 42** (batch-73 manuscript 42): *Balanced Late-Growing Permutations. Algebraic constants, D-finiteness, and complete asymptotic expansions. A proof of the fixed-row growth conjecture for OEIS A215561* (a 953-line source and a 22-page PDF, with the banner “AI-assisted mathematical research draft. Unrefereed; not formally verified”). Archive `late_growing_excursions.zip`, arrival commit `bdf1a1a73`; it pins the repository at `bc4d1fa2b`.

The two were written independently on the same day and share no prose: about 1.8 % of their 8-word shingles are common in each direction, all of them bibliography entries, OEIS rows and kernel polynomials. Source 42’s pin predates both arrivals. Source 61’s pin is later than source 42’s arrival and contains source 42’s archive as an unopened zip file, so source 61’s repository search (Section 2.3) could not have found it. Their 47 common exactly computed terms are equal, and their constants agree to the last printed digit for $r \leq 8$. Source 61 is the base text; Part II holds what only source 42 has.

(Addition of October 2, 2026.) A third independent manuscript on the same problem arrived in batch 75 (manuscript 03, *Fixed-Content Ballot Words*, here **source 75-03**; archive `OEIS_Fixed_Content_Ballot_Words.zip`, arrival commit `4b874cea0`, placed in `6ea60e367`, pin `29aca108e`). Part IV prints what it adds, chiefly a proof of the Kauers–Koutschan recurrence for A215570; Section 24 gives its provenance and Table 5 the results it re-proves, which are not printed again.

(Addition of October 2, 2026, batch 77.) A fourth independent manuscript, batch-77 manuscript 44, *Equal-content ballot words* (here **source 77-44**; archive `oeis-ballot-asymptotics-20261001.zip`, arrival commit `096ee7b87`, placed in `aa7345800`, no pin), proves the main theorems a fourth time and adds the first correction of A215562. Part V prints what it adds; Section 30 gives its provenance and Table 8 the results it re-proves.

(Addition of October 5, 2026, batch 101.) Two more independent reports on the same problem, bundle Reports 90 and 86 of a session bundle (here **sources 101-90** and **101-86**; archives `A215561_Fixed_Alphabet_Asymptotics.zip` and `A215570_Balanced_Ballot_Asymptotics.zip`, arrival commit `60f54ea06`, placed in `f7c612c72`, no pins), prove the main theorems a fifth and a sixth time, the second for $r = 5$ only. Part VI prints what they add; Section 35 gives their provenance and Table 9 the results they re-prove.

1.2 What each source contributed

Table 1 lists every result and where it is printed. “Both” means the two sources prove the result independently; it is then printed once, with source 61’s statement, and source 42’s proof is printed as well when it follows a different route.

Table 1: Results of the sources. The rows below the first inner rule are source 75-03's additions (Part IV), those below the second source 77-44's (Part V), those below the third sources 101-90's and 101-86's (Part VI); their re-proofs of other rows are listed in Tables 5, 8 and 9.

result	sources	printed as
fixed-row asymptotic, constant κ_r , all orders	both	Theorem 3.1; second routes Theorem 16.1, Section 17
D-finiteness of every row	both	Theorem 3.2
transcendence of every row with $r \geq 3$	61	Section 8
kernel product and bridge identity (classical)	both	Lemmas 4.1, 5.1, with second proof
root-product formula for κ_r	both	Proposition 5.2
integral formula for $\log \kappa_r$	61	(5.7)
sextic for $r = 5$	both	(4.5), Remark 4.2
Hessian determinant, normalization check	61	Lemma 6.1
$\alpha_{5,1} = -13/10 + 13\sqrt{5}/50$	both	Theorem 3.3; Sections 9, 18
$\alpha_{5,j} \in \mathbb{Q}(\sqrt{5})$ for all j	61	Theorem 3.3
second-order terms for $r = 2, 3$; $\alpha_{3,1}$ from the general formula	42	(18.6)
law of the number of arches	both	Theorem 10.1; Theorem 19.2 (general compositions)
giant primitive component, decoration tail	42	Section 19
eventual strict log-convexity	42	(20.2)
leading term for any finite labelled step set	42	Theorem 16.1
all orders and D-finiteness on centred rational rays	61	Corollary 12.1
Lambert- W inverse	both	Section 11
explicit rounding bound	42	(20.4)
constants: $r \leq 8$ (61), $r \leq 12$ (42)	both	Table 3
exact enumerations	both	Sections 13, 21
research directions: eight (61), ten (42)	both	Section 22
Kauers–Koutschan Conjecture 15: the A215570 recurrence	75-03	Theorem 26.1, Appendix C
positive Legendre formula for A215570	75-03	Theorem 25.2
exact $\alpha_{5,2}, \dots, \alpha_{5,8}$; second-order inverse for $r = 5$	75-03	Proposition 27.1, (28.1)
first correction $\alpha_{4,1}$ of A215562	77-44	Theorem 31.1
global mark arcs by an operator norm	77-44	Lemma 32.1
recurrence-free $\alpha_{5,2}$; constant of $A_5^o(n)$; $A_r(m+n) \geq A_r(m)A_r(n)$	101-86	Theorem 37.2, Propositions 37.1, 37.3
zero-step identity; extreme-unmarked Hessian, torus points	101-90	Propositions 36.1, 36.2, Lemma 36.3

1.3 Where the merge chose

Source 61 is the base: it carries the larger set of theorems about the OEIS family (transcendence, the field $\mathbb{Q}(\sqrt{5})$, all orders and D-finiteness on rational rays, the Hessian determinant, the integral formula, the $r = 5$ inverse numerics). Source 42's leading-term theorem has weaker analytic hypotheses (finite bridge mass, no saddle point) and a wider class of step sets, so the usual preference for the more general source would make it the base for that theorem alone; it is printed instead as the general theorem of Part II, with second-route status for the rows, because source 61 is stronger on its rays.

Source 42's proofs are kept where they follow a genuinely different route: the convolution

proof of the leading term (Theorem 16.1); the factorization proof of the bridge identity (after Lemma 5.1); the all-orders proof in lattice coordinates with its coefficient algorithm (Section 17); the first-correction formula (Section 18); the elimination for κ_4 (Section 7); the cutoff proof of the arch law (Theorem 19.2); and the second way of removing the auxiliary variable in the kernel lemma. Not printed a second time: source 42’s statements of the main theorem, of D-finiteness and of the root-product constant, its sextic (the negative of (4.5)), its kernel lemma and D-finiteness proof, and its Lambert- W core with first corrections, all of which coincide with source 61’s. The two constant tables are merged, the research directions are merged item by item, and source 61’s package appendix is rewritten for the shipped file names, with source 42’s instructions folded into it.

For source 75-03 (Part IV) the merge prints in full only what Parts I and II lack: the positive coefficient formula, the proof of the recurrence with its certificate (Appendix C), the coefficient algorithm and the exact $\alpha_{5,j}$. Its two-variable saddle is kept, briefly, as a second route to the $r = 5$ expansion, because it supplies $\alpha_{5,1}$ by a third contraction, and its resolvent derivation of the sextic as a remark. Everything else it proves is in Parts I and II already, mostly in a stronger form, and appears only as a row of Table 5, with two small increments stated in Section 24.2. Its research questions are merged into Section 22; its package description is folded into Appendix B.

For source 77-44 (Part V, October 2, 2026) the merge prints in full the first correction $\alpha_{4,1}$ with its ingredients, the operator-norm proof of the global-arc bound as a second route, its nested-logarithm inverse and its literature pointers; everything else is a row of Table 8. Its questions coincide with those of Section 22 and are listed in Section 34; its package is described in Appendix B.

For sources 101-90 and 101-86 (Part VI, October 5, 2026) the merge takes source 101-90, which treats every fixed alphabet, as the base and source 101-86 as its five-letter case. It prints in full what they add (the recurrence-free $\alpha_{5,2}$ with every intermediate value, the constant of the zero-free count, the supermultiplicativity of A_r , the zero-step identity, the extreme-unmarked determinant and torus count), prints their proofs of Theorems 3.1–3.3 as second routes, mapped in Table 10, and lists everything else as a row of Table 9. Their questions are merged into Section 22; their packages are described in Appendix B.

1.4 Repository context and re-derived material

The bridge–excursion identity (Lemma 5.1 and (14.3)) is the composition-marked case, with coefficient ring $\mathbb{Q}[x_1, \dots, x_r]$ and step symbol $\sum_i x_i u^{s_i}$, of Corollary `cor:excursion` of the collection’s report *Multiple-chain preorder polytopes: two proofs of the exponential formula* (SetTheory/Cardinals/docs/reports/enumerative-combinatorics/multiple-chain-exponential-formula), proved there over any commutative coefficient ring from its meander factorization; it is also classical [8]. Both sources’ proofs are kept as further routes.

The inversion of Section 11 and the rounding bound (20.4) re-derive, for this family, the apparatus of the chapter “Apparatus for inverting a rapidly growing function” (`p0:sec:top`) of the canonical transseries volume `Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/transseries_and_inversion.tex`: its exact-core and perturbed inversion (`p0:thm:perturbed-inversion`) and its staircase theorem `p0:thm:staircase` for integer thresholds. The separation clause of the staircase theorem and its failure for arbitrarily small error are formalized as the Lean theorems `Fabius.staircase_separation` and `Fabius.staircase_separation_fails` in `Analysis/FabiusFunction/Lean/FabiusFunction/StaircaseInversion.lean`. Neither source cites these theorems (source 61 cites the README of that volume, unchanged since its pin [14]); they are cited here, and no novelty is

claimed for the inversion or rounding statements. The same holds for source 75-03's inversion of A215570, which also does not cite them (Section 28). (2026-10-02 note: the same holds for source 77-44's inverse, Remark 33.1.) (2026-10-05 note: and for sources 101-90 and 101-86, Section 36.5.)

All three sources are AI-assisted and unrefereed. No Lean or Rocq development checks any statement of this report; the Lean theorems just named are general facts about ceilings and do not verify its asymptotic inputs, and its place in the research-report collection confers no formal status. Source 77-44 (Part V) says nothing about AI assistance; it is unrefereed and not formalized either. (2026-10-05 note: the same holds for sources 101-90 and 101-86, Part VI, whose receipts record only a pre-delivery approval by their producing pipeline.)

1.5 Notation

The two sources used different letters for the same objects and the same letters for different objects. This report fixes: $A_r(n)$ for the counts; κ_r for the constant and $\kappa(\mathbf{p})$ for a composition $\mathbf{p}$; $\alpha_{r,j}$ for the normalized coefficients of the expansion; b_j for the coefficients of its logarithm and μ_j for the coefficients of the inverse series; $N_r(Y)$ for the integer threshold (a bare N is always a length); $B(t; \mathbf{x})$ for the bridge generating function and $\mathcal{L}(\mathbf{x}) = \log \mathcal{E}(\mathbf{x})$ for the log-excursion (or log-bridge) series, whose marked specialization is the $\mathcal{L}(t; \mathbf{y})$ of Section 6 (the calligraphic $\mathcal{B}$ of (9.2) is a length correction, unrelated); H for the Hessian in marker coordinates and Q for the corresponding matrix in the lattice coordinates of Part II (not the kernel polynomial $Q_r(u)$); ν for the time period; $\mathcal{I}$ for the generating function of arches (primitive excursions); h for the depth of the most negative step; and V for the step variance. Table 2 lists every renamed symbol; no normalization changed (source 75-03's Q_r has the opposite sign; its other conventions are fixed in Section 24.3; source 77-44's step polynomial is r times P_r , Section 30.3). (2026-10-05 note: the symbols of sources 101-90 and 101-86 are renamed in Table 11, Section 35.3, which also lists six tempting false readings.) A few letters keep two meanings in separate places, where the context fixes them: q is the number of labelled steps in Part II, the tilted probabilities $q_i(\mathbf{y})$ in Section 6, and a root product in (4.5); X is a Gaussian vector in Part II and a discrete random vector in the proof of Lemma 6.1; Y is a target value in the inversions and Y_i a random step in Lemma 15.1; and the bridge weights b_k carry a composition index, the logarithmic coefficients b_j a scalar one.

Part I

Fixed-composition excursions

2 The problem, its significance, and the source audit

2.1 A natural multiset-permutation family

For a fixed integer $r \geq 2$, let $A_r(n)$ be the number of words $w = w_1 \cdots w_{rn}$ containing exactly n occurrences of each letter $1, \dots, r$ and satisfying

$$\sum_{j=1}^k w_j \leq \frac{r+1}{2}k \quad (0 \leq k \leq rn). \quad (2.1)$$

These are rows of the array A215561. The average on the right is exact: in a uniformly chosen multiset permutation, each position has expected letter $(r+1)/2$. This family in-

Table 2: Renamed symbols.

source 42	here	source 42	here
$T_r(n)$	$A_r(n)$	C, c_i (projection)	Ξ, ξ_i
$E_r, E(\mathbf{p}), E$	$\kappa_r, \kappa(\mathbf{p}), \kappa$	V (lattice basis)	$\mathbf{V}$
$B_r = \log E_r$	$\log \kappa_r$	$R(z)$	$P_*(z)$
B (bridge mass)	$\mathfrak{B}$	$\lambda(z), \lambda_m$	$\Psi(z), \Psi_m$
$\mathcal{B}$ (log-bridge)	$\mathcal{L}$	$E(t, z), E_*(z)$	$\widehat{\mathcal{E}}(t, z), \kappa_*(z)$
$d_j(r)$	$\alpha_{r,j}$	H_{an}, H_k	$\Upsilon_{\text{an}}, \Upsilon_k$
$q_j(r)$	b_j	G_j	$\mathcal{G}_j$
C_r	K_r	β, t_h (transfer)	γ, t_i
α_r, λ_r	β, λ	β_j	χ_j
$n_r(Y)$	$N_r(Y)$	$h = N^{-1/2}$	$\hbar$
$K_{r,J}$, truncation J	$\varpi_{r,M}, M$	η (exponent)	ε_1
δ (period), σ^2	ν, V	M_j, M_{12}	$\mathcal{M}_j, \mathcal{M}_{12}$
a (minimum step)	h	$a(z)$	$\mathbf{a}(z)$
$\alpha = (q+1)/2, a_0$	ς, η	ρ (nonprincipal roots)	ω
K_N (primitive factors)	J_N	$v_1, v_2, v_3; s$	$a_1, a_2, a_3; w$
J_N (giant component)	Ω_N	$h(\mathbf{k})$ (height)	$\text{ht}(\mathbf{k})$
U_N, V_N, U, V	$\mathcal{U}_N, \mathcal{V}_N, \mathcal{U}, \mathcal{V}$	$D(t)$	$\Theta(t)$
D (decoration length)	D_{dec}	U_j, V_j (roots)	$u_j, \tilde{u}_j$
$D(\ \cdot\)$	$D_{\text{KL}}(\ \cdot\)$	Y (sextic)	X
source 61	here	source 61	here
d_j (inverse series)	μ_j	H_0 (arches)	$\mathcal{I}$
$T_r(Y)$	$N_r(Y)$		
source 75-03	here	source 75-03	here
$\mathcal{E}_r, \mathcal{E}(\mathbf{p})$	$\kappa_r, \kappa(\mathbf{p})$	d_j (asymptotic)	$\alpha_{5,j}$
$E(\mathbf{x}), e(\mathbf{k})$	$\mathcal{E}, e_{\mathbf{k}}$	$H(\mathbf{x}), H_0$	$\mathcal{L}, \mathfrak{B}$
$h(\mathbf{k}), D = \frac{r+1}{2}$	$b_{\mathbf{k}}, \varsigma$	M (cutoff), ℓ (summands)	H, m
$K_N; q = 1 - \mathcal{E}^{-1}$	$J_N; \eta$	$Q_r(u)$	$-Q_r(u)$
z (length); b (depth)	$t; h$	$G_r(z); B(z)$	$F_r(z); B(t; \mathbf{x})$
Y (sextic); A, B, C, T	$X; A, B, C, \bar{c}$	d_n (zero-free count)	$A_5^{\varepsilon}(n)$
C_n (Legendre coefficient)	Λ_n	P_n (Legendre)	$\mathcal{P}_n$
$W; N, M; p, q, k$ (Lagrange)	$\mathcal{Y}; \mathbf{n}, \mathbf{n}+1; i, j, k$	$\tilde{a}_n, h_n$	$\tilde{A}_5(n), \mathfrak{h}_n$
R, S, U, V	$\mathfrak{R}, \mathfrak{S}, \mathfrak{U}, \mathfrak{V}$	$p_j, q_j, \lambda(n)$	$\mathfrak{p}_j, \mathfrak{q}_j, \mathfrak{c}(n)$
w_n, L_n	$\mathcal{W}_n, \mathcal{L}_n$	$V_j, D = t^2 - 1$	$\zeta_j, t^2 - 1$
$\mathcal{S}_0, \mathcal{S}_1; \mathcal{A}_n, \mathcal{B}_n$	$\mathcal{R}_0, \mathcal{R}_1; \mathcal{U}_n, \mathcal{V}_n$	$\alpha_j, \beta_j; R_m, S_m$	$\mathbf{u}_j, \mathbf{v}_j; \mathfrak{R}_{\sharp}, \mathfrak{S}_{\sharp}$
$\mathcal{P}(u, v); H$	$\Pi(u, v); H_{\Pi}$	$\rho; G = 1 + u_0$	$\varphi^{-2}; G_0$
$\eta_j; c_k; P_j(z)$	$\eta_j^{\Lambda}; \tilde{c}_k; \widehat{\mathfrak{p}}_j(z)$	$b_j, g_j; \kappa_j; (X, Y)$	$\ell_j^+, \ell_j^-; \mathcal{K}_j; (Z_1, Z_2)$
$L = 5 \log 5; y; N(y)$	$\lambda; Y; N_5(Y)$	D (log variable)	D_Y
source 77-44	here	source 77-44	here
$a_q(n), q$ (alphabet)	$A_r(n), r$	$r = \varphi - \sqrt{\varphi}$	$\varrho = \kappa_4/4$
$c_{q,j}$	$\alpha_{r,j}$	$E_*(\mathbf{1}), E_q$	κ_r
$\sigma_q^2; \Delta$	$V; \delta$	$D_{\text{uni}}, D_{\text{sad}}$	$\mathcal{B}, \mathcal{S}$
d_2, α (root jet)	D, c	$G = H^{-1}; \kappa_{ijk}$	$C; T_{abc}$
$c = -\min S_q; h = q - 2$	$h; d$	p_q (period)	ν
$\mathbf{w}; \mathbf{z} = \log \mathbf{w}$	$\mathbf{z}; \mathbf{y}$	$\eta = 1 - t/\rho$	x (with t rescaled)
$\mathcal{B}$ (bridge series)	$B(t; \mathbf{x})$	F, F_2 (sextic)	X, e
$P_q(u) = \sum_i u^{s_i}$	$rP_r(u)$	κ_q, β_q, C_q (inverse)	λ, β, K_r

sources 101-90 and 101-86: Table 11 (Part VI)

cludes Catalan numbers and a balanced Motzkin-word enumeration, but already the fourth and fifth rows lead to substantially more difficult coefficient extraction. The first rows are balanced versions of finite-step excursion enumeration, combining a ballot constraint with prescribed letter multiplicities; fixing the multiplicities is the significant extra condition, and the unrestricted-composition excursion sequence and a fixed-composition diagonal are different objects (source 42). Source 42 calls the counted words *late-growing permutations*, after the titles of the OEIS entries, and imposes (2.1) for $1 \leq k \leq rn$, which is equivalent.

The relevant sequence identifiers are

r	OEIS entry	Description of the row
2	A000108	Catalan numbers
3	A007004	Balanced three-letter excursions
4	A215562	Four-letter prefix-average words
5	A215570	Five-letter prefix-average words
6	A215571	Six-letter prefix-average words
7	A215593	Seven-letter prefix-average words

The array entry records Kotesovec’s 2016 conjecture that, for each fixed $r > 1$,

$$A_r(n) \sim c(r) \frac{r^{rn}}{\pi^{(r-1)/2} n^{(r+1)/2}}. \quad (2.2)$$

The fourth and fifth entries already supply explicit leading constants; those constants are credited to Kotesovec, rather than presented here as new discoveries [1, 2, 3].

2.2 What is and is not resolved

Kauers and Koutschan discuss A215570 and A215562 in Section 6.1 of their 2023 paper. Their finite transfer matrices grow with n , which prevents their argument from proving D-finiteness. They leave a particular order-three recurrence for A215570 conjectural and ask for a provably correct recurrence for A215562 [7]. We replace the growing matrix by a fixed algebraic generating function in composition-marking variables. (2026-10-02 note: Part IV proves that order-three recurrence, Theorem 26.1; the A215562 request remains open.)

The main results below establish the existence of polynomial-coefficient recurrences for *every* fixed row. They do not print a recurrence for the fourth row and do not certify the particular guessed operator for the fifth (Part IV now certifies it). Recurrence existence, an explicit annihilating operator, and minimality of that operator are three different assertions. Source 42 draws the same distinctions and divides the credit the same way: the kernel method and the bridge–excursion identity are classical ([8], Theorem 2 and Corollary 2), closure of D-finite series under diagonals is classical [9], and the contribution is their use with a directional coefficient theorem to resolve the fixed-composition growth assertion, determine its constants, and obtain the additional fixed-composition conclusions.

The source check used the live OEIS entries, the cited 2023 paper, the classical directed-path kernel theory, and targeted searches for the sequence identifiers. The fixed-row asymptotic was still labelled a conjecture in A215561 when checked. No proof of the precise results below was identified in that search. This is a bounded source audit, not a proof that no such result exists anywhere in the literature.

2.3 Relationship to ProveIt

The inspected ProveIt snapshot was

40b17f2fc79a10a6fc1a50ca9c0c3d6f77be69e5.

Its transseries-and-inversion documentation distinguishes exact cores, finite-order expansions, remainder transport, and discrete staircase inversion. We follow that organization in Section 11; the theorems of that volume which the inversion re-derives are named in Section 1.4. The current path inspected was

Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/README.md.

The document's own formalization inventory is deliberately narrower than its mathematical exposition [14]. No repository theorem is silently imported as a formal proof of our excursion results. Searches for A215561 and A215562 returned no repository matches; that limited check is not an audit of every file or of other identifiers for the same problem. (2026-10-01 placement note: manuscript 42, *Balanced Late-Growing Permutations*, delivered in bdf1a1a73 and present at the pin as a zipped archive, proves the same leading theorem independently; the two are merged in this report, Section 1.) Source 42 inspected the repository at its own pin bc4d1fa2b [15]; its targeted search did not locate an A215570 report, which was true then, and it states that neither that search nor its literature search is an exhaustive novelty certificate, and that it is mathematically independent of unverified repository theorems.

3 Main theorems

Put

$$g_r = \gcd(2, r-1), \quad s_i = \frac{2i - r - 1}{g_r}, \quad h = -s_1, \quad P_r(u) = \frac{1}{r} \sum_{i=1}^r u^{s_i}. \quad (3.1)$$

For odd $r = 2h + 1$, the steps are $-h, -h + 1, \dots, h$. For even r , they are the odd integers $-(r-1), -(r-3), \dots, r-1$. Let S_m be the sum of m independent steps chosen uniformly from this set.

Theorem 3.1 (All fixed rows: leading term and complete expansion). *For every fixed $r \geq 2$, the series*

$$\log \kappa_r = \sum_{m \geq 1} \frac{\mathbb{P}(S_m = 0)}{m} \quad (3.2)$$

converges, and κ_r is a positive real algebraic number. For each integer $M \geq 0$ there are real algebraic numbers $\alpha_{r,1}, \dots, \alpha_{r,M}$ such that

$$A_r(n) = K_r \frac{r^{rn}}{n^{(r+1)/2}} \left(1 + \sum_{j=1}^M \frac{\alpha_{r,j}}{n^j} + O_{r,M}(n^{-M-1}) \right), \quad (3.3)$$

where

$$K_r = \frac{\kappa_r}{\sqrt{r} (2\pi)^{(r-1)/2}}, \quad c(r) = \frac{\kappa_r}{\sqrt{r} 2^{(r-1)/2}}. \quad (3.4)$$

In particular,

$$\frac{A_r(n)}{(rn)!/(n!)^r} \sim \frac{\kappa_r}{rn}. \quad (3.5)$$

All coefficients in (3.3) are computable by finite algebraic operations on kernel expansions and Gaussian moments.

The constants implied by the O -notation are allowed to depend on r and M . Nothing in this statement is uniform when the alphabet size grows with n .

Theorem 3.2 (Algebraic ambient series; transcendental row series). *The multivariate generating function counting nonnegative excursions with steps (3.1), marking every step type independently, is algebraic over $\mathbb{Q}(x_1, \dots, x_r)$. Consequently*

$$F_r(z) = \sum_{n \geq 0} A_r(n) z^n$$

is D-finite for every fixed r . The function F_r is transcendental over $\mathbb{Q}(z)$ for every $r \geq 3$. For $r = 2$ it is the algebraic Catalan generating function.

Thus taking a diagonal preserves D-finiteness here, but does not preserve algebraicity. The transcendence assertion uses not just the growth exponent but, for even r , the power of π in the leading constant.

Theorem 3.3 (An explicit next term for A215570). *For the fifth row,*

$$A_5(n) = \frac{3\sqrt{5} - 5}{8\pi^2} \frac{5^{5n}}{n^3} \left(1 - \frac{13(5 - \sqrt{5})}{50n} + O(n^{-2}) \right). \quad (3.6)$$

Moreover every coefficient $\alpha_{5,j}$ in its full expansion belongs to $\mathbb{Q}(\sqrt{5})$.

The leading factor in (3.6) is the previously posted OEIS factor. The proof and the correction are developed below without assuming the conjectured recurrence.

Sources. Theorem 3.1 and the D-finiteness part of Theorem 3.2 are proved in both sources. Source 42 states them as its Theorems 1.1 and 1.2, with $1 < \kappa_r < \infty$; it proves the leading term without a saddle point (Theorem 16.1) and the complete expansion in lattice coordinates (Section 17). The transcendence part of Theorem 3.2 and the field statement of Theorem 3.3 are source 61's alone; the value of $\alpha_{5,1}$ was computed in both, by different contractions (Sections 9 and 18).

4 The exact excursion generating function

4.1 Encoding the prefix condition

Replacing a letter i by $r + 1 - i$ reverses the prefix inequality and preserves the multiset. It is therefore equivalent to count words whose centered partial sums, using steps s_i , are *nonnegative*. The total centered sum is zero. These are one-dimensional excursions with prescribed numbers of the individual steps.

For a count vector $\mathbf{k} = (k_1, \dots, k_r)$ write $|\mathbf{k}| = \sum_i k_i$ and $\mathbf{x}^{\mathbf{k}} = \prod_i x_i^{k_i}$. Let $e_{\mathbf{k}}$ count nonnegative excursions with that composition, and set

$$\mathcal{E}(\mathbf{x}) = \sum_{\mathbf{k} \geq 0} e_{\mathbf{k}} \mathbf{x}^{\mathbf{k}}. \quad (4.1)$$

The empty excursion contributes one. A nonzero coefficient requires $\sum_i s_i k_i = 0$, and

$$F_r(z) = \text{Diag } \mathcal{E}(z) := \sum_{n \geq 0} [x_1^n \cdots x_r^n] \mathcal{E}(\mathbf{x}) z^n. \quad (4.2)$$

In particular, the dimension of this ambient generating function is fixed when r is fixed.

4.2 The small-root formula, with a proof

Temporarily mark total length by t , and write

$$P_{\mathbf{x}}(u) = \sum_{i=1}^r x_i u^{s_i}.$$

The kernel equation is

$$u^h(1 - tP_{\mathbf{x}}(u)) = 0. \quad (4.3)$$

Of its $2h$ roots, exactly h tend to zero as $t \rightarrow 0$; these are called small roots. This description may be understood in a Puiseux extension with generic step weights.

Lemma 4.1 (Kernel product). *If $u_1(t), \dots, u_h(t)$ are the small roots, then*

$$\mathcal{E}(t\mathbf{x}) = \frac{(-1)^{h+1}}{tx_1} \prod_{j=1}^h u_j(t). \quad (4.4)$$

Consequently $\mathcal{E}$ is an algebraic multivariate formal power series.

Proof. Let $G_j(t)$ count nonnegative paths ending at height j , and put $G(t, u) = \sum_{j \geq 0} G_j(t)u^j$. Appending one step gives

$$(1 - tP_{\mathbf{x}}(u))G(t, u) = 1 - t \sum_{s_i < 0} x_i \sum_{j=0}^{-s_i-1} G_j(t)u^{j+s_i}.$$

After multiplication by u^h , the right-hand side is a monic polynomial $N(t, u)$ of degree h . Substitution of any small root is legitimate in the formal path equation and gives $N(t, u_j(t)) = 0$. Hence $N(t, u) = \prod_{j=1}^h (u - u_j(t))$. Its constant coefficient is $-tx_1 G_0(t)$: only the most negative step, followed from height zero, contributes to that coefficient. Equating constant coefficients proves (4.4), since $G_0(t) = \mathcal{E}(t\mathbf{x})$.

The roots are algebraic over $\mathbb{Q}(t, \mathbf{x})$, so their product is algebraic. An algebraic relation for $\mathcal{E}(t\mathbf{x})$ can be specialized to $t = 1$ after removing any common factor $t - 1$ from its coefficients. The specialization is valid coefficientwise in the $\mathbf{x}$ -adic topology and yields a nonzero polynomial relation for $\mathcal{E}(\mathbf{x})$. Alternatively (source 42), replace x_i by x_i/t in a polynomial relation for $\mathcal{E}(t\mathbf{x})$ and clear powers of t ; at least one coefficient in t is then a nonzero polynomial relation for $\mathcal{E}(\mathbf{x})$, and no relation that vanishes identically at $t = 1$ is ever specialized. $\square$

This is the classical directed-path kernel construction, included here to make the composition marking explicit; its general development is due to the directed-path literature, notably Banderier and Flajolet [8]. Source 42 gives the same proof for every finite list of labelled integer steps (Section 14).

4.3 Why D-finiteness follows

An algebraic multivariate formal series in characteristic zero is D-finite: its derivatives belong to its finite algebraic function field, so they span a finite-dimensional vector space over the rational function field. Lipshitz's diagonal theorem then implies that the complete diagonal (4.2) is D-finite [9].

Equivalently, for each fixed r there exist polynomials $q_0(n), \dots, q_d(n) \in \mathbb{Q}[n]$, not all zero, with

$$\sum_{j=0}^d q_j(n) A_r(n+j) = 0$$

for all sufficiently large n . This proves the D-finiteness portion of Theorem 3.2. Closure algorithms make this an effective existence result, but they do not imply a small order, a small degree, or a fast calculation of a useful recurrence. Source 42 proves the same statement by the same argument and notes why it does not conflict with the growing transfer matrices of [7]: a transfer matrix whose size increases with n does not have a fixed rational resolvent, but that observation is not an obstruction to a fixed *algebraic* multivariate series.

4.4 A concrete algebraic certificate for the fifth alphabet

For steps $-2, -1, 0, 1, 2$, give the weights the names a, b, c, d, e respectively. The ambient excursion series $X = \mathcal{E}(a, b, c, d, e)$ is the unique formal solution with constant coefficient one of

$$0 = a^3e^3X^6 + a^2e^2(c-1)X^5 + (abde - a^2e^2)X^4 + (ad^2 + b^2e + 2ae(1-c))X^3 + (bd - ae)X^2 + (c-1)X + 1. \quad (4.5)$$

Here the zero-step weight c is a variable, not the OEIS asymptotic constant.

To verify the equation, factor the quartic kernel as

$$e(u^2 - vu + q)(u^2 - wu + q') = eu^4 + du^3 + (c-1)u^2 + bu + a.$$

The two small roots have product $q = -aX$, while $q' = -1/(eX)$. Comparing the cubic and linear coefficients gives

$$v + w = -d/e, \quad vq' + wq = -b/e.$$

Solve these two equations for v, w , and substitute into $q + q' + vw = (c-1)/e$. Clearing denominators gives (4.5). This calculation is valid first for generic weights, and hence as a polynomial identity. At $a = b = c = d = e = 0$ the equation reduces to $1 - X = 0$, with nonzero derivative in X ; the formal implicit function theorem therefore selects a unique branch. The fifth row is the complete diagonal of this explicitly specified algebraic branch.

Remark 4.2 (Source 42's derivation). Source 42 obtains the same equation with the opposite overall sign, written in the abbreviations ae , bd and $ad^2 + b^2e$, by solving the coefficient equations explicitly. With the small-root factor $u^2 - vu + q$ and $q = -aX$ as above,

$$q' = -\frac{1}{eX}, \quad v = \frac{X(adX + b)}{1 - aeX^2}, \quad w = -\frac{d}{e} - v,$$

and substitution into the quadratic coefficient gives

$$[u^2] e(u^2 - vu + q)(u^2 - wu + q') - (c-1) = -\frac{S(X)}{X(aeX^2 - 1)^2},$$

where $S(X)$ is the right side of (4.5); source 42's program checks this identity exactly. The sextic is a finite algebraic input for creative telescoping. It is not a certificate that a guessed diagonal operator annihilates the diagonal; that further elimination, followed by initial-condition checking, remains a concrete computational target (Section 22). (2026-10-02 note: the guessed operator for A215570 is now proved by another route, which needs no elimination from the sextic: Theorem 26.1. A third derivation of the sextic is Remark 25.1.)

5 The bridge logarithm and the algebraic constant

5.1 An exact identity retaining every composition

Define the bridge generating function

$$B(t; \mathbf{x}) = \text{CT}_u \frac{1}{1 - tP_{\mathbf{x}}(u)}.$$

Unlike excursions, bridges have no sign restriction on their intermediate heights. Their composition coefficients are multinomial coefficients.

Lemma 5.1 (Bridge logarithm). *As formal power series,*

$$B(t; \mathbf{x}) = 1 + t \frac{d}{dt} \log \mathcal{E}(t\mathbf{x}), \quad (5.1)$$

and consequently

$$\log \mathcal{E}(\mathbf{x}) = \sum_{\substack{\mathbf{k} \neq 0 \\ \sum_i s_i k_i = 0}} \frac{(|\mathbf{k}| - 1)!}{\prod_i k_i!} \mathbf{x}^{\mathbf{k}}. \quad (5.2)$$

In particular,

$$[x_1^n \cdots x_r^n] \log \mathcal{E}(\mathbf{x}) = \frac{1}{rn} \frac{(rn)!}{(n!)^r} \quad (n \geq 1). \quad (5.3)$$

Proof. For small positive numerical weights and sufficiently small t , constant term extraction is a contour integral. Its residues at the small kernel roots give

$$B(t; \mathbf{x}) = \sum_{j=1}^h \frac{-1}{tu_j P'_{\mathbf{x}}(u_j)} = t \sum_{j=1}^h \frac{u'_j(t)}{u_j(t)}.$$

The second equality follows by differentiating $tP_{\mathbf{x}}(u_j(t)) = 1$. Taking the logarithmic derivative of (4.4) proves (5.1). The identity extends formally to arbitrary weights.

Expanding B in powers of t and then expanding $P_{\mathbf{x}}(u)^m$ gives the coefficient $m!/\prod k_i!$ precisely when $|\mathbf{k}| = m$ and $\sum s_i k_i = 0$. Integrating $(B - 1)/t$ supplies the factor $1/m$ and proves (5.2). Its balanced specialization gives (5.3). $\square$

The bridge–excursion relation is classical [8]. Its usefulness here is that the diagonal of the logarithm is *known exactly*. We can calibrate a multivariate saddle calculation against that coefficient and thereby avoid an unjustified guess about cyclic rotations or lattice periods.

Second proof of Lemma 5.1 (source 42). Expanding the logarithm gives

$$\text{CT}_u [-\log(1 - tP_{\mathbf{x}}(u))] = \sum_{m \geq 1} \frac{t^m}{m} [u^0] P_{\mathbf{x}}(u)^m,$$

and $[u^0] P_{\mathbf{x}}(u)^m$ is precisely the multinomial generating polynomial for zero-sum words of length m . So this constant term is the right side of (5.2) at $t\mathbf{x}$, and it remains to identify it with $\log \mathcal{E}(t\mathbf{x})$; (5.1) then follows by differentiation. Factor the kernel in an annulus between its small and large roots:

$$1 - tP_{\mathbf{x}}(u) = \Theta(t) \prod_{j=1}^h (1 - u_j/u) \prod_{j=1}^h (1 - u/\tilde{u}_j),$$

where $\tilde{u}_1, \dots, \tilde{u}_h$ are the large roots. The coefficient of u^{-h} gives $\Theta(t)(-1)^h \prod_j u_j = -tx_1$; by Lemma 4.1, $\Theta = 1/\mathcal{E}(t\mathbf{x})$. The logarithms of the small-root factors have only negative powers of u , and those of the large-root factors have only positive powers. Taking the Laurent constant term therefore gives $\text{CT}_u[-\log(1 - tP_{\mathbf{x}})] = -\log \Theta = \log \mathcal{E}(t\mathbf{x})$. One may first work at generic weights and then use formal coefficient identities to specialize them. All the series have the prescribed value at $t = 0$, fixing the logarithmic branch. $\square$

The identity is also the composition-marked case of a corollary proved in the collection's report on multiple-chain preorder polytopes over any commutative coefficient ring (Section 1.4).

5.2 The critical value

Set $\mathbf{p} = (1/r, \dots, 1/r)$. In the probability normalization,

$$B(t; \mathbf{p}) = \sum_{m \geq 0} \mathbb{P}(S_m = 0) t^m.$$

A finite-support, nondegenerate centered lattice walk has $\mathbb{P}(S_m = 0) = O(m^{-1/2})$. For completeness, this bound follows by Fourier inversion: the characteristic function has quadratic modulus decay near the finitely many points where its modulus is one, and has modulus bounded strictly below one on their complement. The integral near each such point is bounded by a Gaussian integral. Thus (3.2) converges. By monotone convergence in the nonnegative coefficient series,

$$\kappa_r = \lim_{t \uparrow 1} \mathcal{E}(t\mathbf{p}) = \mathcal{E}(\mathbf{p}) > 1. \quad (5.4)$$

Proposition 5.2 (Finite algebraic formula). *Let*

$$Q_r(u) = \sum_{i=1}^r u^{s_i+h} - ru^h. \quad (5.5)$$

The root $u = 1$ has multiplicity exactly two. Apart from that root, Q_r has $h - 1$ roots strictly inside the unit disk and $h - 1$ strictly outside, counting multiplicities. If the inside roots are $\alpha_1, \dots, \alpha_{h-1}$, then

$$\kappa_r = (-1)^{h+1} r \prod_{j=1}^{h-1} \alpha_j. \quad (5.6)$$

An empty product is one.

Proof. At $u = 1$, $P_r(1) = 1$, $P'_r(1) = 0$, and $P''_r(1) = \sum_i p_i s_i^2 > 0$. Hence the multiplicity is exactly two. If $|u| = 1$ and $P_r(u) = 1$, equality in the triangle inequality forces $u^{s_i} = 1$ for every i . The normalized step set has gcd one, so $u = 1$. There are no other roots on the unit circle.

For $0 < t < 1$, Rouché's theorem applied on $|u| = 1$ shows that $u^h(1 - tP_r(u))$ has exactly h roots inside. Near $t = 1$, its two roots near one split as $1 \pm c\sqrt{1-t} + O(1-t)$ with $c > 0$; one is inside and one outside. This leaves $h - 1$ strictly inside in the limit. Formula (5.6) is now the limit of (4.4). It also proves algebraicity. Positivity was already obtained from (5.4). $\square$

Internal multiple roots, should they occur, cause no difficulty: the product of a separated cluster is analytic even when individual root labels cannot be chosen analytically.

Source 42 proves the same formula as the limit at $t = 1$ of the small-root formula, with the finite value (3.2) identifying the branch and proving positivity. It counts the inside roots

without Rouché's theorem: Q_r is a reciprocal polynomial of degree $2h$ with a double root at $u = 1$ and no other root on the unit circle, so reciprocity leaves $h - 1$ roots inside the unit disk. The prescription is exact; numerical root selection is only one way to evaluate it, and certified algebraic root isolation can replace floating-point root selection wherever a numerical interval certificate is required.

For the symmetric step sets in this paper one also has the real integral

$$\log \kappa_r = -\frac{1}{2\pi} \int_0^{2\pi} \log(1 - P_r(e^{i\theta})) d\theta. \quad (5.7)$$

Indeed, expand first with a factor $t < 1$, integrate term by term, and let $t \uparrow 1$. The logarithmic singularity at $\theta = 0$ is integrable. The integrand is real because $P_r(e^{i\theta})$ is real and at most one. Equation (5.7) is an alternative numerical expression, not a replacement for the algebraic proof.

6 Fixed-composition singularity analysis

Source 42 proves the expansion of this section by the same strategy in lattice coordinates, with an explicit Gaussian-moment algorithm (Section 17), and proves its leading term without any saddle point (Theorem 16.1).

6.1 Removing the two linear constraints

Let $d = r - 2$ and mark only the first d step types in addition to total length. Use logarithmic markers $\mathbf{y} = (y_1, \dots, y_d)$ and set

$$P_{\mathbf{y}}(u) = \sum_{i=1}^r p_i e^{y_i^*} u^{s_i}, \quad y_i^* = \begin{cases} y_i & i \leq d, \\ 0 & i > d. \end{cases} \quad (6.1)$$

If total length is N and the first d counts are known, the remaining two counts are uniquely determined by length and total height zero, because $s_{r-1} \neq s_r$. At $N = rn$, extracting n occurrences of each marked type therefore extracts the complete balanced composition.

The relevant coefficient functional is

$$\mathcal{C}_N(U) = [t^N z_1^n \cdots z_d^n] U(tp_1 z_1, \dots, tp_d z_d, tp_{r-1}, tp_r), \quad N = rn. \quad (6.2)$$

For $d = 0$ the z extraction is absent. Thus

$$\mathcal{C}_N(\mathcal{E}) = r^{-N} A_r(n), \quad \mathcal{C}_N(\log \mathcal{E}) = r^{-N} \frac{(rn)!}{rn(n!)^r}.$$

6.2 The moving critical point and its Hessian

The analytic implicit function theorem gives a unique $\tau(\mathbf{y})$ near one solving

$$(u \partial_u) P_{\mathbf{y}}(u) \big|_{u=\tau(\mathbf{y})} = 0, \quad \tau(0) = 1. \quad (6.3)$$

The required derivative in $\log u$ is the positive variance

$$V = \sum_i p_i s_i^2.$$

Define

$$\begin{aligned}\rho(\mathbf{y}) &= P_{\mathbf{y}}(\tau(\mathbf{y}))^{-1}, \\ \psi(\mathbf{y}) &= \log P_{\mathbf{y}}(\tau(\mathbf{y})), \\ \Phi(\mathbf{y}) &= \psi(\mathbf{y}) - \sum_{a=1}^d p_a y_a.\end{aligned}\tag{6.4}$$

Then $\nabla\Phi(0) = 0$. Its Hessian is

$$H_{ab} = p_a \delta_{ab} - p_a p_b - \frac{p_a s_a p_b s_b}{V} \quad (1 \leq a, b \leq d).\tag{6.5}$$

Lemma 6.1 (Nondegenerate composition saddle). *The matrix H is positive definite, and, with $\delta = s_r - s_{r-1} = 2/g_r$,*

$$\det H = \frac{\delta^2}{V} \prod_{i=1}^r p_i.\tag{6.6}$$

For $d = 0$, the determinant convention is $\det H = 1$.

Proof. For numbers $v_1, \dots, v_d$, extend $f_i = v_i$ for $i \leq d$ by $f_{r-1} = f_r = 0$. Direct differentiation in (6.3) gives

$$v^\top H v = \text{Var}_p(f) - \frac{\text{Cov}_p(f, s)^2}{V}.$$

Cauchy–Schwarz makes this nonnegative. Equality would require $f_i = a + b s_i$ for every i . The two distinct steps with $f_i = 0$ force $a = b = 0$, hence $v = 0$.

For the determinant, let the random vector X have components $(\mathbf{1}_{\{i=1\}}, \dots, \mathbf{1}_{\{i=d\}}, s_i)$ with probability p_i . Its covariance matrix has Schur complement H in the last entry V , so its determinant is $V \det H$. A random vector supported on r affinely independent points in dimension $r - 1$ has covariance determinant equal to the product of its r probabilities times the square of the affine-simplex determinant. This identity follows by adjoining a row of ones to the support matrix and applying the Schur complement to its weighted Gram matrix. Here that affine determinant is, up to sign, $s_r - s_{r-1} = \delta$. This proves (6.6). $\square$

(2026-10-05 note: with the two extreme steps unmarked the same argument gives $\det H_{\text{ex}} = 4h^2 r^{-r}/V = (r-1)^2 \det H$, Proposition 36.2.)

6.3 A uniform square-root expansion

Put

$$q_i(\mathbf{y}) = \frac{p_i e^{y_i^* \tau(\mathbf{y})^{s_i}}}{P_{\mathbf{y}}(\tau(\mathbf{y}))}, \quad V(\mathbf{y}) = \sum_i q_i(\mathbf{y}) s_i^2.\tag{6.7}$$

The q_i sum to one and have mean step zero. Multiplying weights by τ^{s_i} does not change any excursion weight, so the critical value is

$$\kappa(\mathbf{y}) = \mathcal{E}(q(\mathbf{y})).$$

It is analytic near $\mathbf{y} = 0$ as a critical root-cluster product, and $\kappa(0) = \kappa_r$.

Let $x = 1 - t/\rho(\mathbf{y})$. The small root which reaches the critical point has the expansion

$$u_0(t, \mathbf{y}) = \tau(\mathbf{y}) \left(1 - c(\mathbf{y}) x^{1/2} + O(x)\right), \quad c(\mathbf{y}) = \sqrt{\frac{2}{V(\mathbf{y})}}.$$

All the other small roots are a separated cluster. Their product, and the factor $1/t$ in the kernel formula, are analytic at the critical point. Therefore, uniformly for complex $\mathbf{y}$ sufficiently near zero,

$$\mathcal{E}(t; \mathbf{y}) = \kappa(\mathbf{y}) \left(1 - c(\mathbf{y})x^{1/2} + O(x)\right), \quad (6.8)$$

$$\mathcal{L}(t; \mathbf{y}) := \log \mathcal{E}(t; \mathbf{y}) = \log \kappa(\mathbf{y}) - c(\mathbf{y})x^{1/2} + O(x). \quad (6.9)$$

Each expansion continues to arbitrary order in $x^{1/2}$. The notation $\mathcal{E}(t; \mathbf{y})$ means the marked specialization in (6.2), with $z_a = e^{y_a}$.

The square-root coefficients differ by exactly $\kappa(\mathbf{y})$. This analytic fact is the reason that the exact logarithmic reference coefficient in (5.3) identifies the fixed-composition leading constant.

6.4 Why other phases do not contribute

An asymptotic proof must control the full marker torus, not just a local expansion. Here this control is particularly simple.

Suppose $|t| = |u| = |z_a| = 1$ and a kernel root lies on $|u| = 1$. Equality in

$$\left| t \sum_i p_i z_i^* u^{s_i} \right| \leq 1$$

forces $t z_i^* u^{s_i} = 1$ for every i . The last two unmarked types imply $u^\delta = 1$. All step differences are multiples of δ , so then $z_a = 1$ for every marked type. If r is odd, $\delta = 1$ and $t = u = 1$. If r is even, $\delta = 2$ and the two possibilities are $(t, u) = (1, 1)$ and $(-1, -1)$.

Thus the only marker saddle is $\mathbf{z} = \mathbf{1}$. The only length period is

$$\nu = \begin{cases} 1 & r \text{ odd,} \\ 2 & r \text{ even,} \end{cases} \quad \text{so that } \nu = \delta. \quad (6.10)$$

In the even case every excursion has even length, and the two length singularities give identical contributions on the admissible subsequence $N = rn$. (2026-10-05 note: with the two *extreme* steps unmarked instead, there are $r - 1$ dominant mark points, each with ν time phases, Lemma 36.3; the count depends on the coordinates and the constant does not, Remark 36.4.)

More explicitly, on any compact part of the marker torus separated from $\mathbf{1}$, no kernel root can cross $|u| = 1$ for $|t| \leq 1$. By compactness this remains true for $|t| < 1 + \epsilon$ with some uniform $\epsilon > 0$. The small-root product is analytic there, even if roots within that cluster collide. It is nonzero away from $t = 0$, since the kernel has nonzero constant term for $t \neq 0$. The removable value at $t = 0$ is one. Consequently both $\mathcal{E}$ and its logarithm have uniform analytic continuations on that disk. Cauchy estimates make this part of the coefficient integral exponentially smaller than the main term.

6.5 Transfer and the Gaussian integral

For a half-integer γ which is not a nonnegative integer,

$$[t^N](1 - t/\rho)^\gamma = \rho^{-N} \frac{\Gamma(N - \gamma)}{\Gamma(-\gamma)\Gamma(N + 1)}. \quad (6.11)$$

The gamma quotient has a complete inverse-power expansion. Applying (6.11) to (6.8)–(6.9) gives local leading amplitudes

$$g_L(\mathbf{y}) = \frac{c(\mathbf{y})}{2\sqrt{\pi}}, \quad g_E(\mathbf{y}) = \kappa(\mathbf{y})g_L(\mathbf{y}), \quad (6.12)$$

multiplying $\rho(\mathbf{y})^{-N} N^{-3/2}$ and the period factor ν .

For marker extraction put $\mathbf{y} = i\boldsymbol{\theta}$. The exponential in the Cauchy integral is

$$\exp(N\Phi(i\boldsymbol{\theta})), \quad \Phi(i\boldsymbol{\theta}) = -\frac{1}{2}\boldsymbol{\theta}^\top H\boldsymbol{\theta} + O(|\boldsymbol{\theta}|^3).$$

The Hessian is positive definite. The leading Gaussian integral thus gives

$$\mathcal{C}_N(\mathcal{L}) \sim \frac{\nu g_L(0)}{(2\pi)^{d/2} \sqrt{\det H}} N^{-(d+3)/2}, \quad (6.13)$$

$$\mathcal{C}_N(\mathcal{E}) \sim \kappa_r \frac{\nu g_L(0)}{(2\pi)^{d/2} \sqrt{\det H}} N^{-(d+3)/2}. \quad (6.14)$$

The ratio is therefore κ_r . Substituting the exact logarithmic coefficient proves (3.5).

There is also a useful independent normalization check. From (6.6) and (6.10),

$$\frac{\nu g_L(0)}{(2\pi)^{d/2} \sqrt{\det H}} = \frac{\nu}{\delta} \frac{1}{(2\pi)^{(r-1)/2} \sqrt{\prod_i p_i}} = \frac{r^{r/2}}{(2\pi)^{(r-1)/2}}.$$

This is exactly the leading Stirling coefficient of $r^{-N} (rn)! / (rn(n!)^r)$ when $N = rn$. In particular, no hidden period-two factor is lost in the even alphabets.

6.6 Remainders and all orders

We spell out why the preceding calculation proves more than a leading heuristic. Fix any requested order M . The local kernel factorization provides Puiseux expansions to an arbitrarily high finite order, with analytic coefficient functions and uniform remainder estimates on a common dented neighborhood of the moving singularity. This follows from analytic factorization of the double root and analytic separation of the other root clusters. The same neighborhoods work for sufficiently small complex marker values.

After the length coefficient is extracted, Taylor-expand its analytic amplitudes and the phase Φ to a finite order larger than that needed for N^{-M-1} . In a neighborhood $|\boldsymbol{\theta}| \leq N^{-1/2+\varepsilon}$, substitute $\boldsymbol{\theta} = \mathbf{x}/\sqrt{N}$. Taylor's theorem and the positive-definite Gaussian majorant bound every discarded term by an integrable polynomial times a Gaussian, with the specified inverse power of N . One may first fix a sufficiently small $\varepsilon > 0$ depending on the requested truncation and then take sufficiently many Taylor terms. In the remainder of the small marker neighborhood, the real part of the phase is at most $-c|\boldsymbol{\theta}|^2$, giving an error smaller than every power of N^{-1} . The compact complement was handled by the analytic-continuation argument above.

Gaussian moments of odd total degree vanish. The half-integer powers in the length expansion differ by integers after coefficient transfer. Consequently only integer powers of N^{-1} remain after the common factor $N^{-(r+1)/2}$ is removed. This proves a full expansion with an $O(N^{-M-1})$ relative remainder after M corrections.

All finite Taylor jets involved are algebraic. Indeed, after replacing e^{y_a} by independent variables, the critical point solves polynomial kernel and derivative equations with algebraic base values. Exponential reparametrization has rational Taylor coefficients. Half-integer gamma factors contribute one common $\sqrt{\pi}$, and Gaussian moments contribute rational expressions in H^{-1} times a common determinant factor. After division by the leading coefficient, every correction is algebraic.

Finally, Stirling's expansion in (3.5), or directly (6.14), yields (3.3) and (3.4). This proves Theorem 3.1, including its all-order and algebraicity assertions.

Remark 6.2 (What “all orders” means here). Equation (3.3) determines the dominant exponential–power sector with a proved finite-order remainder for every fixed truncation. It does not assert convergence of the inverse-power series, identify every exponentially small contribution, or establish Borel summability. Those are separate questions. Source 42 adds that optimal truncation and a Gevrey bound are likewise not proved: the expansion is the dominant exponential sector of an asymptotic transseries, and its exponentially smaller sectors require additional global singularity analysis.

7 Explicit leading constants

The following values are obtained from the finite product, not from fitting a growth curve; Table 3 merges the tables of the two sources.

r	κ_r	$c(r)$ in (2.2)	$2\kappa_r - 1$
2	2.000000000000000	1.000000000000000	3.000000000000000
3	3.000000000000000	0.866025403784439	5.000000000000000
4	1.384057356943304	0.244669085661435	1.768114713886607
5	1.909830056250526	0.213525491562421	2.819660112501052
6	1.239095083832872	0.089423985025306	1.478190167665743
7	1.583896616939167	0.074832081265314	2.167793233878334
8	1.173744890362129	0.036679527823817	1.347489780724258
9	1.429003292368548	0.029770901924345	1.858006584737097
10	1.136489993032388	0.015882930501054	1.272979986064777
11	1.338806843936432	0.012614545363912	1.677613687872865
12	1.112404234469499	0.007095892609796	1.224808468938998

Table 3: Leading constants. The rows $r \leq 8$ of the first two columns were computed in both sources and agree to every digit either prints (source 61 printed 15 decimals, source 42 twelve); the rows $9 \leq r \leq 12$ and the last column, the limiting mean number of arches of Theorems 10.1 and 19.2, are source 42’s (its printed table stops at $r = 10$; its data file `data/42-late-growing-constants.csv` reaches $r = 12$, and those values are rounded here to 15 decimals). Decimal values are diagnostics, not interval-certified enclosures.

The printed decimals are numerical evaluations; the exact definition is (5.6). The evident even–odd pattern suggests a separate large-alphabet problem (source 42). None of the fixed- r proofs establishes a uniform two-parameter expansion.

For $r = 2, 3$, $h = 1$, so the product is empty and $\kappa_2 = 2$, $\kappa_3 = 3$. Directly,

$$A_2(n) = \frac{(2n)!}{n!(n+1)!}, \quad A_3(n) = \frac{(3n)!}{(n!)^3(n+1)}.$$

For the second formula, remove the n zero steps from a balanced Motzkin excursion, obtaining a Dyck path with n up and n down steps; reinsert the zero steps in $\binom{3n}{n}$ ways; source 42 derives both formulas the same way. These formulas give the checks

$$\alpha_{2,1} = -\frac{9}{8}, \quad \alpha_{3,1} = -\frac{11}{9}.$$

For the fourth row,

$$Q_4(u) = (u-1)^2(u^4 + 2u^3 + 4u^2 + 2u + 1).$$

The quartic is reciprocal. On dividing by u^2 and putting $v = u + u^{-1}$, it becomes $v^2 + 2v + 2 = 0$. Pair the two roots strictly inside the unit disk and multiply them. If $\varphi = (1 + \sqrt{5})/2$, the result is

$$\kappa_4 = 4(\varphi - \sqrt{\varphi}), \quad c(4) = \frac{\varphi - \sqrt{\varphi}}{\sqrt{2}}.$$

Substitution in (3.3) reproduces the formula credited in the OEIS entry to Kotesovec [2]. (2026-10-02 note: Part V gives the next coefficient, $\alpha_{4,1}$, Theorem 31.1.) Source 42 reaches the same value by elimination: write the quartic as $(u^2 - \sigma u + q)(u^2 - (\sigma/q)u + q^{-1})$, where q is the product of its two interior roots. Comparing cubic and quadratic coefficients gives $\sigma = -2q/(1 + q)$ and $q + q^{-1} + \sigma^2/q = 4$; eliminating σ gives $q^4 - 2q^3 - 2q^2 - 2q + 1 = 0$, and the root selected by $0 < q < 1$ has $q + q^{-1} = 1 + \sqrt{5} = 2\varphi$, so that $\kappa_4 = 4q$ is the value above.

For the fifth row,

$$Q_5(u) = (u - 1)^2(u^2 + 3u + 1).$$

The unique strict inside root is $\alpha = (-3 + \sqrt{5})/2$. Hence

$$\kappa_5 = \frac{5}{2}(3 - \sqrt{5}), \quad c(5) = \frac{3\sqrt{5} - 5}{8}.$$

This is the leading constant already recorded in A215570 [3]. As source 42 puts it, the claim for $r = 4, 5$ is a proof and a general prescription, not that these two radical expressions were previously absent. For reproducibility, two larger reduced kernels are

$$\begin{aligned} \frac{Q_6(u)}{(u - 1)^2} &= u^8 + 2u^7 + 4u^6 + 6u^5 + 9u^4 + 6u^3 + 4u^2 + 2u + 1, \\ \frac{Q_7(u)}{(u - 1)^2} &= u^4 + 3u^3 + 6u^2 + 3u + 1. \end{aligned}$$

Together with the inside-root selection, these give exact algebraic specifications of the constants for A215571 and A215593, whose checked entries did not give asymptotic formulas [4, 5].

8 Why every nontrivial higher row is transcendental

The exponent and constant in Theorem 3.1 imply a complete algebraicity classification for the row generating functions.

Proof of the transcendence assertion in Theorem 3.2. Set $R = r^{-r}$ and $G(x) = F_r(Rx)$. Its coefficients satisfy

$$[x^n]G(x) \sim K_r n^{-(r+1)/2}, \quad K_r > 0.$$

Assume for contradiction that G is algebraic over $\mathbb{Q}(x)$. Every ordinary derivative of G is then algebraic and, near the algebraic point $x = 1$, has a Puiseux expansion with algebraic coefficients.

First let $r = 2k + 1 \geq 3$. Then

$$G^{(k)}(x) \sim K_r \log \frac{1}{1 - x} \quad (x \uparrow 1).$$

To justify this Abelian estimate, its coefficient sequence is asymptotic to K_r/n up to the harmless derivative index shift. Compare it, outside any fixed finite initial segment, with $(K_r \pm \epsilon)/n$ and use $\sum_{n \geq 1} x^n/n = -\log(1 - x)$. No Puiseux series can have a leading logarithmic divergence. This is a contradiction.

Next let $r = 2k \geq 4$. Differentiation k times gives coefficients asymptotic to $K_r n^{-1/2}$. The same comparison argument and the Gaussian integral, or a Riemann-sum estimate with $x = e^{-t}$, yield

$$G^{(k)}(x) \sim K_r \sqrt{\pi} (1-x)^{-1/2}. \quad (8.1)$$

The leading Puiseux coefficient of an algebraic function over $\mathbb{Q}(x)$ at $x = 1$ must be algebraic. But

$$K_r \sqrt{\pi} = \frac{\kappa_r}{\sqrt{r} 2^{k-1/2}} \pi^{-(k-1)}$$

is transcendental: its prefactor is nonzero algebraic, and $k-1$ is a positive integer. If this number were algebraic, then π^{k-1} and hence π would be algebraic, contradicting Lindemann's theorem [13]. This contradicts (8.1).

Finally $F_2(z) = (1 - \sqrt{1-4z})/(2z)$ is algebraic, as expected. $\square$

The argument is not a generic claim that all diagonals are transcendental. It uses the particular positive leading asymptotic and its algebraic constant. It also explains why the even rows require an arithmetic argument in addition to their half-integer growth exponent.

9 The first correction for the fifth row

9.1 A differential formula for the ratio correction

This section makes the all-order construction concrete. Its calculation is independent of the conjectured A215570 recurrence (proved in Part IV, which uses the value of $\alpha_{5,1}$ found here as an initial datum).

Use the phase and marker notation from Section 6, and let $C = H^{-1}$. Repeated indices in the formulas below are summed from 1 to d . Put

$$\ell(\mathbf{y}) = \log \kappa(\mathbf{y}), \quad h_c(\mathbf{y}) = \log c(\mathbf{y}), \quad T_{abc} = \partial_a \partial_b \partial_c \Phi(0).$$

All derivatives without an argument are evaluated at zero. For an analytic amplitude g , the order- N^{-1} contribution in its Gaussian saddle integral is

$$-\frac{1}{2} C_{ab} g_{ab} + \frac{1}{2} g_a T_{bcd} C_{ab} C_{cd} + g \mathcal{J},$$

where $\mathcal{J}$ is independent of the amplitude. This follows by expanding $g(i\mathbf{x}/\sqrt{N})$ and the cubic and quartic terms of $N\Phi(i\mathbf{x}/\sqrt{N})$, then using Gaussian moments. The signs come from $i^2 = -1$; in particular the second derivative term is negative.

Subtracting the normalized correction for g_L from that for $g_E = \kappa g_L$ cancels $\mathcal{J}$ and gives

$$\mathcal{S} = -\frac{1}{2} C_{ab} (\ell_{ab} + \ell_a \ell_b) - C_{ab} \ell_a (h_c)_b + \frac{1}{2} \ell_a T_{bcd} C_{ab} C_{cd}. \quad (9.1)$$

There is a second correction from the length singularity itself. Write at $\mathbf{y} = 0$

$$\begin{aligned} \mathcal{E}(t) &= \kappa_r (1 - cx^{1/2} + Dx + E_3 x^{3/2} + O(x^2)), \\ \log \mathcal{E}(t) &= \log \kappa_r - cx^{1/2} + L_2 x + L_3 x^{3/2} + O(x^2), \quad x = 1 - t. \end{aligned}$$

Then $D = L_2 + c^2/2$. In the coefficient transfer formula the universal $3/(8N)$ term cancels between the two functions, and $\Gamma(-1/2)/\Gamma(-3/2) = -3/2$. It follows that the difference of their relative length corrections is

$$\mathcal{B} = -\frac{3}{2} L_2 - \frac{c^2}{4}. \quad (9.2)$$

Combining the two extractions,

$$A_r(n) = \frac{\kappa_r (rn)!}{rn (n!)^r} \left(1 + \frac{\Delta_r}{rn} + O(n^{-2}) \right), \quad \Delta_r = \mathcal{B} + \mathcal{S}. \quad (9.3)$$

Stirling's first correction then gives

$$\alpha_{r,1} = \frac{1/r - r}{12} + \frac{\Delta_r}{r}. \quad (9.4)$$

Equations (9.1)–(9.4) are also an implementable recipe for other fixed rows. (2026-10-02 note: source 77-44 evaluates it for $r = 4$, Theorem 31.1.)

9.2 Exact ingredients for $r = 5$

Here $s = (-2, -1, 0, 1, 2)$, $p_i = 1/5$, $d = 3$, and $V = 2$. The Hessian and inverse are

$$H = \begin{pmatrix} 2/25 & -2/25 & -1/25 \\ -2/25 & 7/50 & -1/25 \\ -1/25 & -1/25 & 4/25 \end{pmatrix}, \quad C = \begin{pmatrix} 130 & 90 & 55 \\ 90 & 70 & 40 \\ 55 & 40 & 30 \end{pmatrix}.$$

The large entries in C are a consequence of marking the first three letters while reconstructing the final two. They are not evidence of a degenerate saddle: H is strictly positive definite.

For compact formulas define the centered residual markers

$$R_a(i) = \mathbf{1}_{\{i=a\}} - p_a - \frac{p_a s_a}{V} s_i. \quad (9.5)$$

Then

$$T_{abc} = \sum_i p_i R_a(i) R_b(i) R_c(i), \quad (h_c)_a = -\frac{1}{2V} \sum_i p_i s_i^2 R_a(i). \quad (9.6)$$

The cubic identity is the third derivative of the profiled log moment generating function in (6.4); derivatives of the optimized height coordinate cancel because each residual marker is orthogonal to the height step.

There is an especially simple formula for $\kappa(\mathbf{q})$ for nearby centered five-step probabilities. Write

$$A = q_{-2}, \quad E = q_2, \quad B = q_1 + 2q_2 = q_{-1} + 2q_{-2}.$$

The equality for B is the zero-drift condition. The critical kernel factors as

$$\sum_{s=-2}^2 q_s u^{s+2} - u^2 = (u-1)^2 (Eu^2 + Bu + A),$$

so

$$\kappa(\mathbf{q}) = \frac{2}{B + \sqrt{B^2 - 4AE}}. \quad (9.7)$$

The square root is positive at the uniform probabilities and continued analytically nearby.

For transparency, let $v = \log \tau$. Differentiating the zero-drift equation and (6.7) gives

$$\begin{aligned} v_a &= -p_a s_a / V, & v_{ab} &= -V^{-1} \sum_i p_i s_i R_a(i) R_b(i), \\ (q_i)_a &= p_i R_a(i), & (q_i)_{ab} &= p_i (R_a(i) R_b(i) + s_i v_{ab} - H_{ab}). \end{aligned} \quad (9.8)$$

Equations (9.7)–(9.8) determine the needed derivatives using only rational arithmetic and $\sqrt{5}$. Their values are

$$(\ell_a) = \left(-\frac{3}{10} + \frac{\sqrt{5}}{10}, \quad \frac{1}{5} - \frac{\sqrt{5}}{10}, \quad \frac{1}{5} \right), \quad ((h_c)_a) = \left(-\frac{1}{10}, \frac{1}{20}, \frac{1}{10} \right),$$

and

$$(\ell_{ab}) = \begin{pmatrix} \frac{2}{25} - \frac{4\sqrt{5}}{125} & -\frac{2}{25} + \frac{11\sqrt{5}}{250} & -\frac{1}{25} \\ -\frac{2}{25} + \frac{11\sqrt{5}}{250} & \frac{7}{50} - \frac{17\sqrt{5}}{250} & -\frac{1}{25} \\ -\frac{1}{25} & -\frac{1}{25} & \frac{4}{25} \end{pmatrix}.$$

Substitution in (9.1), using (9.6), now gives the exact contraction

$$\mathcal{S} = -\frac{11}{4} + \sqrt{5}. \quad (9.9)$$

To obtain L_2 , note that the uniform law is symmetric. The logarithm of the critical small root has no x term: in the coordinate $u = e^w$, $P(e^w)$ is even in w , so $w = -c\sqrt{x} + O(x^{3/2})$. The other small root is $\alpha = (-3 + \sqrt{5})/2$. Differentiating its kernel equation yields

$$L_2 = 1 + \frac{1}{\alpha P'_5(\alpha)} = 1 - \frac{\sqrt{5}}{5}.$$

Since $c^2 = 2/V = 1$, equation (9.2) gives

$$\mathcal{B} = -\frac{7}{4} + \frac{3\sqrt{5}}{10}, \quad \Delta_5 = -\frac{9}{2} + \frac{13\sqrt{5}}{10}.$$

Finally (9.4) gives

$$\alpha_{5,1} = -\frac{13}{10} + \frac{13\sqrt{5}}{50} = -\frac{13(5 - \sqrt{5})}{50}.$$

This proves (3.6).

All higher finite jets at the uniform point stay in $\mathbb{Q}(\sqrt{5})$: the critical root-cluster data have that field, $c = 1$, and all implicit recursions have nonzero denominators in the same field. Gaussian and Stirling corrections introduce only rational operations. This proves the field assertion in Theorem 3.3.

Source 42 obtains $\alpha_{5,1}$ independently from a different, coordinate-invariant contraction formula (Section 18), and recovers $\alpha_{3,1} = -11/9$ from that general formula rather than from the closed form. In the multinomial normalization (9.3), $\Delta_5/5 = -9/10 + 13\sqrt{5}/50$, which is source 42's (18.7).

10 A fixed-composition contact law

The same saddle comparison provides distributional information, not only total enumeration. An *arch* is a nonempty excursion with no interior return to height zero. A zero step at height zero is itself an arch. Every nonempty excursion decomposes uniquely into a sequence of arches. Hence the arch generating function is

$$\mathcal{I}(\mathbf{x}) = 1 - \mathcal{E}(\mathbf{x})^{-1}.$$

Let J_n be the number of arches of a uniformly chosen balanced r -letter excursion, and let $C_n = J_n - 1$ be its number of interior returns to zero.

Theorem 10.1 (Contact limit under exact composition). *For each fixed $r \geq 2$ and $k \geq 0$,*

$$\mathbb{P}(C_n = k) \longrightarrow \frac{k+1}{\kappa_r^2} \left(1 - \frac{1}{\kappa_r}\right)^k. \quad (10.1)$$

The probability generating functions converge in a complex neighborhood of $v = 1$, and

$$\mathbb{E}C_n \longrightarrow 2(\kappa_r - 1). \quad (10.2)$$

Proof. For any analytic function f near κ_r , the leading square-root coefficient of $f(\mathcal{E})$ is $f'(\kappa_r)$ times that of $\mathcal{E}$. The same coefficient-extraction proof therefore gives the leading ratio $f'(\kappa_r)$, provided the composition is analytic on the necessary neighborhoods. For fixed k , apply this with $f(E) = (1 - E^{-1})^{k+1}$. It counts exactly $k+1$ arches, and the ratio is

$$f'(\kappa_r) = \frac{k+1}{\kappa_r^2} (1 - \kappa_r^{-1})^k.$$

These nonnegative limiting probabilities sum to one.

For the stronger generating-function statement, mark arches by v :

$$\mathcal{E}_v = \frac{1}{1 - v\mathcal{I}}.$$

At the critical weights, $\mathcal{I} = \eta := 1 - \kappa_r^{-1} < 1$. On the closed coefficient polydisk, nonnegative coefficients give $|\mathcal{I}| \leq \eta$. For $|v| < 1/\eta$ the denominator is consequently bounded away from zero; this remains true in a suitable local continuation. The preceding analysis is therefore uniform for v near one. Its derivative ratio is

$$\mathbb{E}(v^{J_n}) \longrightarrow \frac{v}{\kappa_r^2(1 - \eta v)^2}.$$

Divide by v and differentiate at one to obtain (10.2). □

A related negative-binomial contact law is classical when only length is conditioned [8]. The statement here imposes the complete step composition. Its proof explains why the same critical value survives this additional conditioning.

Source 42 proves the same limit law, stated for the number $J_n = C_n + 1$ of arches (its primitive factors) and for every centred rational composition, together with convergence of all fixed moments, by a cutoff argument that uses no singularity analysis (Theorem 19.2); its limiting mean is $\mathbb{E}J_n \rightarrow 2\kappa_r - 1$.

11 Inverting the asymptotic expansion

11.1 An exact Lambert core

For fixed r define

$$\lambda = r \log r, \quad \beta = \frac{r+1}{2}, \quad K = K_r.$$

Then the proved expansion has the form

$$A_r(n) \sim K e^{\lambda n} n^{-\beta} \left(1 + \frac{\alpha_1}{n} + \frac{\alpha_2}{n^2} + \cdots\right). \quad (11.1)$$

The large real solution of the *leading model* $Y = Ke^{\lambda x}x^{-\beta}$ is exactly

$$x_0(Y) = -\frac{\beta}{\lambda}W_{-1}\left(-\frac{\lambda}{\beta}\left(\frac{K}{Y}\right)^{1/\beta}\right). \quad (11.2)$$

The lower real branch W_{-1} is essential: it selects $x_0 \rightarrow \infty$. The argument lies in $(-e^{-1}, 0)$ for all sufficiently large Y . This exact-core organization parallels the inversion framework inspected in ProveIt [14]; no unverified asymptotic interpolation is needed. The exact core and its perturbation are the instance $(K, \lambda, \beta) = (K_r, r \log r, (r+1)/2)$ of the canonical volume's inversion apparatus (Section 1.4), and source 42 derives the same core and the same first corrections (Section 20); no novelty is claimed for them.

11.2 Corrections and a remainder statement

Write the formal logarithm of the unit correction as

$$\log\left(1 + \sum_{j \geq 1} \alpha_j x^{-j}\right) = \sum_{j \geq 1} b_j x^{-j}, \quad b_1 = \alpha_1, \quad b_2 = \alpha_2 - \frac{1}{2}\alpha_1^2.$$

Seek $x = x_0 + \sum_{j \geq 1} \mu_j x_0^{-j}$ in

$$\lambda(x - x_0) - \beta \log(x/x_0) + \sum_{j \geq 1} b_j x^{-j} = 0. \quad (11.3)$$

At every order the new coefficient occurs with nonzero multiplier λ , so the recursion is triangular. Its first terms are

$$x = x_0 - \frac{b_1}{\lambda x_0} - \frac{b_2/\lambda + \beta b_1/\lambda^2}{x_0^2} + O(x_0^{-3}). \quad (11.4)$$

This has a precise interpretation on the original sequence. Given any fixed order, evaluate the truncated inverse at $Y = A_r(n)$. The forward expansion has a controlled logarithmic residual, while the derivative of $\lambda x - \beta \log x$ is $\lambda - \beta/x$, bounded below by $\lambda/2$ for large x . The mean value theorem transports the residual to an error of the same inverse-power order in the recovered n . Thus the inversion is a theorem about the sequence values and their asymptotic inverse, rather than a claim that a canonical analytic interpolation of the integers has been constructed.

For the fifth row, (11.4) begins

$$n = x_0 + \frac{13(5 - \sqrt{5})}{250 \log 5} \frac{1}{x_0} + O(x_0^{-2}), \quad Y = A_5(n), \quad (11.5)$$

where $\beta = 3$, $\lambda = 5 \log 5$, and $K = (3\sqrt{5} - 5)/(8\pi^2)$ in (11.2). (2026-10-02 note: with $\alpha_{5,2}$ now known, the x_0^{-2} term is explicit, (28.1).)

11.3 Integer thresholds are a separate operation

For arbitrary real Y , define

$$N_r(Y) = \min\{n \geq 0 : A_r(n) \geq Y\}.$$

Appending a fixed balanced excursion block containing one of each step is an injection from the n th class into the $(n+1)$ st. Hence $A_r(n)$ is nondecreasing; its proved asymptotic also

gives $A_r(n+1)/A_r(n) \rightarrow r^r > 1$, so it is eventually strictly increasing. (2026-10-05 note: concatenation gives more, $A_r(m+n) \geq A_r(m)A_r(n)$, so for $r \geq 3$ the sequence increases strictly at every index, Proposition 37.3; source 101-86 states $A_5(n+1) \geq 35A_5(n)$.)

The continuous inverse predicts $N_r(Y)$ to bounded additive error. Taking a ceiling gives the exact threshold when an explicit inverse-error bound separates the predicted value from the relevant integers. Without that separation there is no universally valid exact rounding rule: Y can be arbitrarily close to a jump value $A_r(n)$. This is the discrete staircase issue emphasized in the repository's inversion documentation, and it must not be hidden in an $o(1)$ statement about a continuous model. In the canonical transseries volume it is the staircase theorem `p0:thm:staircase`, whose separation clause is formalized in Lean (Remark 20.1); source 42 gives an explicit separation bound, (20.4).

12 Unequal but centered compositions

The main construction is not restricted to equal multiplicities. Keep the step set (3.1), and let $p_1, \dots, p_r$ be fixed positive rational numbers with

$$\sum_i p_i = 1, \quad \sum_i p_i s_i = 0.$$

For admissible N , let $e(N\mathbf{p})$ count excursions with exactly Np_i occurrences of step i . Define $\kappa(\mathbf{p}) = \mathcal{E}(\mathbf{p})$ by the same critical kernel product, now using these probabilities.

Corollary 12.1 (Centered rational rays). *Along integers N for which all Np_i are integral,*

$$e(N\mathbf{p}) \sim \frac{\kappa(\mathbf{p})}{N} \frac{N!}{\prod_i (Np_i)!}. \quad (12.1)$$

There is a full inverse-power expansion in N with algebraic normalized coefficients. The coefficient series on every fixed positive integral ray is D-finite.

Proof. Replace the uniform probabilities everywhere in Section 6 by the specified p_i . Centering keeps the critical height coordinate at one. The Hessian proof, its determinant, and the marker-torus equality argument are unchanged. The exact coefficient of $\log \mathcal{E}$ is again the multinomial divided by N . Its ratio with the excursion coefficient gives (12.1), and the finite-jet argument gives all orders.

For the final statement, a fixed rational-ray extraction can be implemented by coefficient sections and a diagonal. Sections are finite root-of-unity filters after a monomial substitution; algebraicity is preserved by these operations, and the subsequent diagonal is D-finite. Equivalently one may use the multivariate D-finite residue closure for the fixed linear coefficient constraints. The dimensions and integer ray vector remain fixed. $\square$

This extension concerns excursions relative to the *same* height functional s_i . Centering ensures the final height is zero. Compositions with nonzero mean would describe a different endpoint problem and are not covered by this statement. Source 42 proves the leading term (12.1) for every finite list of labelled integer steps and every centred rational composition, by an elementary convolution argument (Theorem 16.1).

13 Reproducible checks and their scope

13.1 Exact finite enumeration

Source 61's program `code/61-balanced-verify.py` uses a direct composition recurrence. For a count vector $\mathbf{k}$, let $D(\mathbf{k})$ count nonnegative paths with that composition, regardless of final height. Then

$$D(\mathbf{0}) = 1, \quad D(\mathbf{k}) = 0 \text{ if } \sum_i s_i k_i < 0,$$

and for every other nonempty vector,

$$D(\mathbf{k}) = \sum_{i: k_i > 0} D(\mathbf{k} - \mathbf{e}_i).$$

The recurrence follows by deleting the last step. Its implementation stores a finite rectangular count-vector box in mixed-radix order, so every predecessor is already available. The diagonal gives $A_r(n)$.

The executed ranges were

r	Independently generated n	Rectangular states
2	0–24	625
3	0–24	15,625
4	0–18	130,321
5	0–16	1,419,857
6	0–6	117,649
7	0–4	78,125

The first two rows were compared with their factorial formulas. The other rows were compared with the available initial values in the corresponding OEIS entries. All comparisons passed. Source 42's independent dynamic program, with companion arrays for primitive words and returns, is described in Section 21; on the 47 indices that both programs computed ($r = 4, 5, 6, 7$ and $n \leq 18, 15, 6, 4$) the terms are equal.

An independent exact check of the bridge logarithm uses the identity $D_{\text{Euler}}\mathcal{E} = \mathcal{E}D_{\text{Euler}} \log \mathcal{E}$. Coefficientwise this is

$$|\mathbf{k}|e_{\mathbf{k}} = \sum_{\substack{0 < \mathbf{j} \leq \mathbf{k} \\ s: j=0}} \frac{|\mathbf{j}|!}{\prod_i j_i!} e_{\mathbf{k}-\mathbf{j}}. \quad (13.1)$$

Exact rational arithmetic in this identity agreed with the direct path recurrence for 120 nonempty balanced vectors across five test boxes. A separate height-based weighted enumeration verified the sextic (4.5) through length 24 at weights (2, 3, 5, 7, 11). The polynomial elimination proving that sextic is also reproduced in `code/61-balanced-derive_alpha5.py`.

13.2 Asymptotic and inverse diagnostics

For A_5 , write $L_5(n) = K_5 5^{5n} n^{-3}$ and $R_n = A_5(n)/L_5(n)$. The following values illustrate the independently derived first correction:

n	R_n	$n(R_n - 1)$	$n^2(R_n - 1 - \alpha_{5,1}/n)$
4	0.838539713679	-0.645841145284	0.291124722264
8	0.914884186065	-0.680926511482	0.301566514946
12	0.942234769960	-0.693182760477	0.305274784481
16	0.956285986890	-0.699424209762	0.307169857409
20	0.964839683966	-0.703206320687	0.308320103255
50	0.985752002111	-0.712399894449	0.311121570077

The predicted limit of the middle diagnostic is

$$\alpha_{5,1} = -0.7186223258500546789 \dots$$

The values for $n \leq 16$ were generated independently by the supplied program. The $n = 20, 50$ values were read from the published OEIS b-file; they were *not* independently enumerated here [6]. The last column is a numerical check of a second-order residual, not a claimed formula or a numerical certificate for $\alpha_{5,2}$. (2026-10-02 note: Part IV gives $\alpha_{5,2} = 36/25 - 63\sqrt{5}/125 = 0.31302\dots$ exactly, the limit of that column, Table 6; and its proved coefficient formula reproduces the $n = 20, 50$ rows, Theorem 25.2.) (2026-10-05 note: Part VI obtains the same $\alpha_{5,2}$ without the recurrence, by Gaussian moments, Theorem 37.2.)

At $Y = A_5(n)$, the Lambert core and its first correction give

n	$x_0(Y) - n$	$x_0(Y) - \alpha_{5,1}/(5 \log 5 x_0(Y)) - n$
8	$-1.1595249025 \cdot 10^{-2}$	$-4.1641746283 \cdot 10^{-4}$
16	$-5.6870504107 \cdot 10^{-3}$	$-1.0375143173 \cdot 10^{-4}$
50	$-1.7966866816 \cdot 10^{-3}$	$-1.0601877690 \cdot 10^{-5}$

The observed scales are consistent with the proved first- and second-order inverse errors (the last column is accounted for by the explicit x_0^{-2} term of (28.1)). Numerical agreement is a useful implementation check, not a substitute for the asymptotic proof.

13.3 What the computations do not certify

The package does not verify a Lean formalization, prove the guessed A215570 recurrence (source 75-03's program checks the certificate of Part IV, Section 29), find the minimal recurrence for any higher row, or supply interval-certified decimal enclosures for every kernel root. Its exact tests validate finite combinatorial and algebraic calculations. Its root computations and asymptotic tables are high-precision numerical diagnostics. The infinite statements are supported by the mathematical proofs above.

Part II

The convolution route, general step sets and the shape of an excursion

This Part prints the results of source 42 that source 61 does not contain, and source 42's proofs of shared results where they follow a genuinely different route. It uses the merged notation of Section 1.5; Table 2 lists every symbol of source 42 that was renamed. Statements proved in both sources are printed once, in Part I, and are only cited here.

14 Labelled steps and the log-excursion series

14.1 Labelled steps and composition

For a finite list of integer steps $\mathbf{s} = (s_1, \dots, s_q)$ having at least one negative and one positive entry, write

$$\mathcal{H} = \{\mathbf{k} \in \mathbb{N}^q : \mathbf{s} \cdot \mathbf{k} = 0\}, \quad |\mathbf{k}| = k_1 + \dots + k_q.$$

An excursion is a word in the *labelled* steps whose partial sums are nonnegative and whose final sum is zero. Distinct labels remain distinct even when their numerical steps coincide. Let $e_{\mathbf{k}}$ count excursions with composition $\mathbf{k}$, and set

$$\mathcal{E}(\mathbf{x}) = \sum_{\mathbf{k} \in \mathcal{H}} e_{\mathbf{k}} \mathbf{x}^{\mathbf{k}}, \quad e_{\mathbf{0}} = 1. \quad (14.1)$$

For the rows, $q = r$ and the steps are (3.1); the encoding of Section 4.1 turns the prefix condition into this nonnegative convention without changing the balanced composition, and (4.2) expresses $A_r(n)$ as a diagonal coefficient.

Lemma 4.1 holds for every such step list (source 42 states it in this generality): put $h = -\min_i s_i$, $h' = \max_i s_i$ and $x_- = \sum_{i:s_i=-h} x_i$, and replace x_1 by x_- in (4.4). The proof given there applies verbatim, because $u^h(1 - tP_{\mathbf{x}}(u))$ evaluated at $u = 0$ is $-tx_-$, and so do both arguments given there for removing the auxiliary variable. In particular $\mathcal{E}(\mathbf{x})$ is algebraic over $\mathbb{Q}(\mathbf{x})$. Generically the subset-product construction gives the finite degree bound $\binom{h+h'}{h}$, although the actual degree can be smaller.

14.2 The log-excursion series

Define

$$\mathcal{L}(\mathbf{x}) = \sum_{\substack{\mathbf{k} \in \mathcal{H} \\ |\mathbf{k}| > 0}} b_{\mathbf{k}}^0 \mathbf{x}^{\mathbf{k}}, \quad b_{\mathbf{k}}^0 = \frac{1}{|\mathbf{k}|} \binom{|\mathbf{k}|}{k_1, \dots, k_q}. \quad (14.2)$$

Lemma 5.1 and its proof, which uses only the kernel lemma, hold in this generality; in source 42's form the lemma reads

$$\mathcal{E}(\mathbf{x}) = \exp \mathcal{L}(\mathbf{x}), \quad \mathcal{L}(t\mathbf{x}) = \text{CT}_u [-\log(1 - tP_{\mathbf{x}}(u))]; \quad (14.3)$$

this is source 42's form of the bridge-excursion identity, and its own proof of it is printed after Lemma 5.1 as a second proof. The factor $1/|\mathbf{k}|$ is crucial. Replacing $\mathcal{L}$ by the unnormalized bridge series would give a false identity.

15 A directional convolution theorem

This section contains the elementary analytic mechanism behind the leading equivalent. It avoids assuming finite moments of the bridge-series mass; those moments generally diverge.

15.1 Critical multinomial weights

Fix positive rational numbers $p_1, \dots, p_q$ with

$$\sum_i p_i = 1, \quad \sum_i p_i s_i = 0, \quad \sum_i p_i s_i^2 > 0. \quad (15.1)$$

Let N tend to infinity through integers for which $\mathbf{k}_N = N\mathbf{p}$ is integral. Define

$$b_{\mathbf{k}} = b_{\mathbf{k}}^0 \mathbf{p}^{\mathbf{k}} \quad (\mathbf{k} \in \mathcal{H} \setminus \{0\}), \quad b_0 = 0, \quad \mathfrak{B} = \sum_{\mathbf{k}} b_{\mathbf{k}}.$$

Here and below sequences indexed off $\mathcal{H}$ are extended by zero. Put $\varsigma = (q+1)/2$.

Lemma 15.1 (Mass, bounds, and fixed shifts). *The mass $\mathfrak{B}$ is finite and positive. There are constants $c, C > 0$ such that*

$$b_{\mathbf{k}} \leq C|\mathbf{k}|^{-\varsigma} \quad (|\mathbf{k}| > 0), \quad b_{\mathbf{k}_N} \sim \frac{N^{-\varsigma}}{(2\pi)^{(q-1)/2} \sqrt{p_1 \cdots p_q}}. \quad (15.2)$$

For each fixed $\ell \in \mathcal{H}$,

$$\frac{b_{\mathbf{k}_N - \ell}}{b_{\mathbf{k}_N}} \longrightarrow 1. \quad (15.3)$$

The same conclusions along the target ray hold after any fixed admissible shift of that target.

Proof. Let Y take value s_i with probability p_i . Summing by length gives

$$\mathfrak{B} = \sum_{m \geq 1} \frac{\mathbb{P}(Y_1 + \cdots + Y_m = 0)}{m}.$$

The return probability is $O(m^{-1/2})$. One direct proof uses the Fourier integral of the characteristic function. Its modulus equals one at only finitely many points on a period, because the step distribution is nonconstant. Near each such point its modulus is at most $1 - c\theta^2$, after translating the point, and elsewhere it is bounded away from one. The Gaussian integral bounds the result by $Cm^{-1/2}$. The positive and negative steps and rational probabilities ensure a nonempty zero-sum word, so $\mathfrak{B} > 0$.

For the uniform pointwise bound in (15.2), the maximum point mass of a fixed q -category multinomial distribution is $O(m^{-(q-1)/2})$. To verify this directly, when every $k_i \geq p_i m/2$, Stirling's bounds give the stated power times $\exp(-mD_{\text{KL}}(\mathbf{k}/m \parallel \mathbf{p})) \leq 1$. Here $D_{\text{KL}}(\boldsymbol{\varpi} \parallel \mathbf{p}) = \sum_i \varpi_i \log(\varpi_i/p_i)$ is relative entropy, with $0 \log 0 = 0$. On the complementary closed subset of the probability simplex the relative entropy is bounded below by a positive constant. Factorial bounds, or the multinomial theorem followed by that entropy bound, give an exponentially small term, which is smaller than the required power. Dividing by m proves the first bound. Applying Stirling at $\mathbf{k}_N = N\mathbf{p}$ proves its asymptotic form.

Finally, with $L = |\ell|$, the ratio in (15.3) is exactly

$$\frac{(N-L-1)!}{(N-1)!} \prod_i \frac{(Np_i)!}{(Np_i - \ell_i)!} p_i^{-\ell_i},$$

which tends to one. Fixed shifts change none of the preceding power bounds along the ray. $\square$

15.2 One large summand

For sequences on $\mathbb{N}^q$, let $*$ denote additive convolution.

Theorem 15.2 (Directional one-large-summand principle). *For each fixed $m \geq 1$,*

$$\frac{(b^{*m})_{\mathbf{k}_N}}{b_{\mathbf{k}_N}} \longrightarrow m\mathfrak{B}^{m-1}. \quad (15.4)$$

Uniformly in $m \geq 1$ and all sufficiently large admissible N ,

$$\frac{(b^{*m})_{\mathbf{k}_N}}{b_{\mathbf{k}_N}} \leq C_0 m^{\varsigma+1} \mathfrak{B}^{m-1}. \quad (15.5)$$

Both assertions hold for a target with a fixed admissible shift.

Proof. In a decomposition $\mathbf{k}_N = \mathbf{k}^{(1)} + \cdots + \mathbf{k}^{(m)}$, some component has length at least N/m . Designate one such component. By Lemma 15.1 its weight is at most $C(N/m)^{-\varsigma}$. Summing the other $m-1$ components freely contributes at most $\mathfrak{B}^{m-1}$, and there are at most m choices for the designated index. The lower bound $b_{\mathbf{k}_N} \geq cN^{-\varsigma}$ proves (15.5). If $m > N$ the convolution vanishes, so no small-length exception is needed.

Now fix m and a cutoff H . Consider decompositions in which all but one component have length at most H . For $N > mH$ the large component is unique. The fixed-shift ratio (15.3) gives, after division by $b_{\mathbf{k}_N}$, the limit

$$m \left(\sum_{|\mathbf{k}| \leq H} b_{\mathbf{k}} \right)^{m-1}.$$

Every omitted decomposition has at least two components of length greater than H . Designate a largest one and one other such component. The same bound as before shows that their total normalized contribution is at most

$$C_m \mathfrak{B}^{m-2} \sum_{|\mathbf{k}| > H} b_{\mathbf{k}}$$

for $m \geq 2$. This tends to zero as $H \rightarrow \infty$. Taking first $N \rightarrow \infty$ and then $H \rightarrow \infty$ proves (15.4); $m = 1$ is immediate. The proof uses only finite total mass, not finite first moment. The shifted version follows from the shifted part of Lemma 15.1. $\square$

Corollary 15.3 (Entire functions of the bridge series). *For any fixed $z \in \mathbb{C}$,*

$$\frac{[\mathbf{x}^{\mathbf{k}_N}] \exp(z\mathcal{L}(\mathbf{p}\mathbf{x}))}{b_{\mathbf{k}_N}} \longrightarrow ze^{z\mathfrak{B}}. \quad (15.6)$$

In particular,

$$[\mathbf{x}^{\mathbf{k}_N}] \mathcal{E}(\mathbf{p}\mathbf{x}) \sim e^{\mathfrak{B}} b_{\mathbf{k}_N}.$$

Proof. For $N > 0$, expand the exponential and use (15.4) termwise. The absolute bounds

$$C_0 m^{\varsigma+1} \mathfrak{B}^{m-1} |z|^m / m!$$

are summable, so dominated convergence applies, also for negative or complex z . The limiting series is $ze^{z\mathfrak{B}}$. Tonelli's theorem applied to the nonnegative coefficient expansion also gives $\mathcal{E}(\mathbf{p}) = e^{\mathfrak{B}} < \infty$. $\square$

16 The general balanced-composition formula

Theorem 16.1 (Arbitrary rational balanced compositions). *Under (15.1), let $\kappa(\mathbf{p}) = \mathcal{E}(\mathbf{p}) = e^{\mathfrak{B}}$. Then*

$$e_{N\mathbf{p}} \sim \frac{\kappa(\mathbf{p})}{N} \binom{N}{Np_1, \dots, Np_q} \quad (16.1)$$

as $N \rightarrow \infty$ through admissible multiples. The constant $\kappa(\mathbf{p})$ is finite, positive, and algebraic.

Proof. Corollary 15.3 states $e_{N\mathbf{p}}\mathbf{p}^{N\mathbf{p}} \sim \kappa(\mathbf{p})b_{N\mathbf{p}}^0\mathbf{p}^{N\mathbf{p}}$. Cancel the common weight. Finiteness has already been proved. Algebraicity follows from the small-root formula at the rational critical weights, taking the finite limit from $t < 1$; equivalently take the product of its limiting algebraic roots. In particular, this argument does not need to infer algebraicity merely from convergence of a transcendental-looking exponential series. $\square$

For $p_i = 1/r$, $N = rn$, and the steps (3.1), Theorem 16.1 proves (3.5): this is source 42's proof of the leading term of Theorem 3.1, which uses no saddle point. Stirling gives

$$\frac{(rn)!}{(n!)^r} = \frac{\sqrt{r} r^{rn}}{(2\pi n)^{(r-1)/2}} \left(1 + \frac{r^{-1} - r}{12n} + O_r(n^{-2}) \right), \quad (16.2)$$

which proves the formula for $c(r)$ in (3.4) and hence the OEIS growth conjecture (2.2).

Theorem 16.1 and Corollary 12.1 meet on the step set (3.1) with centred rational probabilities, where they give the same leading term. Beyond that they are complementary: Theorem 16.1 covers every finite labelled list of integer steps with both signs and needs only the finite bridge mass, but gives the leading term only; Corollary 12.1 adds all orders and D-finiteness on its rays, for the rows' step set.

Remark 16.2 (A probabilistic interpretation). Among all balanced words, the probability of satisfying the prefix constraint is asymptotic to $\kappa_r/(rn)$. Thus κ_r measures the lattice correction to the naive inverse-length ballot probability. For $r = 2$ the exact probability is $1/(n+1)$; for $r = 3$ it is also $1/(n+1)$. The values $\kappa_2 = 2$ and $\kappa_3 = 3$ are consistent with both facts. There is no assertion that the analogous probability is exactly a rational function of n for larger r .

17 A second route to the complete expansion: lattice coordinates

Source 42 proves the all-orders part of Theorem 3.1 by the same strategy as Section 6 (Fourier extraction over the composition torus, a moving square-root singularity, coefficient transfer and Wick contraction), but in coordinates on the lattice of balanced compositions rather than marker coordinates, and with an explicit coefficient algorithm. Its formulation is printed here because Section 18 builds on it. Like Section 6, it uses local complex analysis, and the proof also specifies a finite coefficient procedure at every order.

17.1 Composition coordinates and the unique Fourier saddle

Fix r . Let

$$\Lambda = \{v \in \mathbb{Z}^r : \mathbf{1} \cdot v = \mathbf{s} \cdot v = 0\}, \quad d = r - 2.$$

Choose an integral basis of this lattice, with basis matrix $V \in \mathbb{Z}^{r \times d}$, and put $\Xi = (V^T V)^{-1} V^T$. Then

$$\Xi V = I_d, \quad \Xi \mathbf{1} = \Xi \mathbf{s} = 0. \quad (17.1)$$

Write $\xi_i \in \mathbb{Q}^d$ for the i th column of Ξ . At length $N = rn$, a zero-sum composition is uniquely $n\mathbf{1} + V\ell$ with $\ell \in \mathbb{Z}^d$, and $\Xi \mathbf{k} = \ell$. For $z \in \mathbb{C}^d$ set

$$P_z(u) = \frac{1}{r} \sum_{i=1}^r e^{\xi_i \cdot z} u^{s_i}, \quad \widehat{\mathcal{E}}(t, z) = \mathcal{E}\left(\frac{t}{r} e^{\xi_1 \cdot z}, \dots, \frac{t}{r} e^{\xi_r \cdot z}\right). \quad (17.2)$$

Fourier inversion, at precisely these lengths, gives

$$r^{-N} A_r(n) = \frac{1}{(2\pi)^d} \int_{[-\pi, \pi]^d} [t^N] \widehat{\mathcal{E}}(t, i\theta) d\theta. \quad (17.3)$$

For $d = 0$ the integral is absent.

The only saddle on this torus is zero, modulo its identified boundary. To see the needed strictness, suppose for some θ, ϕ that $|P_{i\theta}(e^{i\phi})| = 1$. Equality in the triangle inequality implies $\xi_i \cdot \theta + s_i \phi \equiv \gamma \pmod{2\pi}$ for all i . Pair this relation with any $v \in \Lambda$. It gives $\theta \cdot \Xi v \equiv 0$, so $\theta \in 2\pi\mathbb{Z}^d$. Hence outside a neighborhood of zero the maximum over ϕ is bounded strictly below one. The constant-term logarithm in (14.3) then converges uniformly for $|t| < 1 + \epsilon$, giving an exponentially small contribution to (17.3) there.

17.2 The critical point and the square-root expansion

There is a unique analytic function $\tau(z)$ near zero, with $\tau(0) = 1$, satisfying

$$u \frac{\partial P_z}{\partial u} \Big|_{u=\tau(z)} = 0.$$

This follows from the implicit-function theorem since

$$V = \frac{1}{r} \sum_i s_i^2 > 0.$$

Put $P_*(z) = P_z(\tau(z))$ and $\Psi(z) = \log P_*(z)$. The identities (17.1) imply

$$P_*(0) = 1, \quad \Psi'(0) = 0, \quad Q := \Psi''(0) = \frac{1}{r} \Xi \Xi^T > 0. \quad (17.4)$$

The notation Q in this Part is a positive matrix, not the kernel polynomial $Q_r(u)$, and not the Hessian H of Section 6, which is the corresponding matrix in marker coordinates.

At $t = P_*(z)^{-1}$, one small and one large root coalesce at $u = \tau(z)$. The second derivative in u is nonzero, so the analytic Morse lemma, or direct coefficient solving, gives a convergent Puiseux expansion in $(1 - tP_*(z))^{1/2}$. The product of the other small roots is analytic in t, z in this neighborhood. This remains true even if individual nonprincipal roots have an internal collision: the product of the roots inside a separating contour is analytic by symmetric root functions. Thus

$$\widehat{\mathcal{E}}(t, z) = \Upsilon_{\text{an}}(t, z) + \sum_{k \geq 0} \Upsilon_k(z) (1 - tP_*(z))^{k+1/2}, \quad (17.5)$$

locally, with analytic coefficients. Its leading coefficient is

$$\Upsilon_0(z) = -\kappa_*(z) \sqrt{\frac{2P_*(z)}{\tau(z)^2 P_z''(\tau(z))}}, \quad \kappa_*(z) = \widehat{\mathcal{E}}(P_*(z)^{-1}, z). \quad (17.6)$$

Here $\kappa_*(z)$ denotes the critical value on the continued small-root branch; it is not an assertion of absolute power-series convergence at every complex marked point. In particular $\Upsilon_0(0) = -\kappa_r \sqrt{2}/\sqrt{V}$.

The time period is $\nu = 1$ for odd r and $\nu = 2$ for even r . Indeed, odd r admits a zero step, while for even r all steps are odd and a two-step return is possible. At zero markings the only dominant singularities are the corresponding time rotations of one. This follows as well from the triangle-equality argument in the preceding subsection. At nearby complex markings their local contributions are the same on admissible lengths. Their separation from the remaining

singularities persists uniformly in a small neighborhood. Therefore coefficient transfer from (17.5) yields, for $N = rn$,

$$[t^N]\widehat{\mathcal{E}}(t, z) = \nu P_*(z)^N N^{-3/2} \left(\sum_{j=0}^M \frac{\mathcal{G}_j(z)}{N^j} + O(N^{-M-1}) \right), \quad (17.7)$$

uniformly locally in z , with

$$\mathcal{G}_0(z) = \frac{\kappa_*(z)}{\sqrt{2\pi}} \sqrt{\frac{P_*(z)}{\tau(z)^2 P''_z(\tau(z))}}. \quad (17.8)$$

Uniformity can be obtained by the same separated root contours and a common slit neighborhood for the coalescing pair. Outside those neighborhoods a coefficient contour can be pushed a fixed positive distance beyond the dominant radius. The analytic local terms have no singular contribution; the half-integral terms transfer by the gamma-ratio formula in the next subsection. These observations give the uniform remainder in (17.7), rather than merely a pointwise formal expansion.

17.3 An all-orders coefficient algorithm

For nonintegral γ define rational polynomials $t_i(\gamma)$ by

$$[t^N](1-t)^\gamma = \frac{N^{-\gamma-1}}{\Gamma(-\gamma)} \left(\sum_{i=0}^M \frac{t_i(\gamma)}{N^i} + O(N^{-M-1}) \right). \quad (17.9)$$

They are computed by expanding $\Gamma(N-\gamma)/\Gamma(N+1)$, as in (6.11); in particular $t_0 = 1$ and $t_1(\gamma) = \gamma(\gamma+1)/2$. Then

$$\mathcal{G}_j(z) = \sum_{k=0}^j \frac{\Upsilon_k(z)}{\Gamma(-k-1/2)} t_{j-k}(k+1/2). \quad (17.10)$$

Write $\Psi(z) = \frac{1}{2}z^\top Qz + \sum_{m \geq 3} \Psi_m(z)$, where Ψ_m is homogeneous of degree m . Let X be a centred d -dimensional Gaussian with covariance Q^{-1} . Form the formal series

$$\mathcal{A}(\hbar, X) = \left(\sum_{j \geq 0} \hbar^{2j} \mathcal{G}_j(i\hbar X) \right) \exp \left(\sum_{m \geq 3} \hbar^{m-2} \Psi_m(iX) \right). \quad (17.11)$$

Every coefficient needed below is a finite polynomial in X . Set

$$\chi_j = \mathbb{E}[\hbar^{2j}] \mathcal{A}(\hbar, X), \quad \alpha_{r,j} = \frac{\chi_j}{r^j \mathcal{G}_0(0)}. \quad (17.12)$$

Gaussian moments are computed exactly by pairings: an odd moment is zero, and an even moment is the sum, over pairings of its factors, of products of covariance entries. Thus (17.12) is a finite procedure, not an appeal to fitted coefficients.

Second proof of the all-orders part of Theorem 3.1 (source 42). Insert (17.7) into (17.3) and scale $\theta = X/\sqrt{N}$ in a neighborhood of the unique saddle. By (17.4),

$$N\Psi(iX/\sqrt{N}) = -\frac{1}{2}X^\top QX + \sum_{m \geq 3} N^{-(m-2)/2} \Psi_m(iX).$$

Taylor expansion of the analytic amplitudes gives exactly (17.11) with $\hbar = N^{-1/2}$. Terms of odd degree in $\hbar$ integrate to zero: in each such term the total degree in X is odd. The resulting expansion is

$$r^{-N} A_r(n) = \frac{\nu N^{-(d+3)/2}}{(2\pi)^{d/2} \sqrt{\det Q}} \left(\sum_{j=0}^M \chi_j N^{-j} + O(N^{-M-1}) \right).$$

To justify truncation, first restrict to $|X| \leq N^{\varepsilon_1}$ with sufficiently small fixed $\varepsilon_1 > 0$. Analytic Taylor remainders there are bounded by a chosen power of $N^{-1/2}$ times a polynomial in $|X|$ and a Gaussian envelope. Expand to enough extra orders to reach the prescribed M , and integrate that envelope. The part with $N^{\varepsilon_1} < |X|$ inside the fixed saddle neighborhood is exponentially small in $N^{2\varepsilon_1}$, since $\operatorname{Re} \Psi(i\theta) \leq -c|\theta|^2$. The complement of the saddle neighborhood was already exponentially controlled. This proves the stated remainder at every order.

The leading coefficient agrees with (3.5), already proved independently (Theorem 16.1); this also checks all lattice and time-period factors. Since $N = rn$ and $d + 3 = r + 1$, normalizing by the leading coefficient gives (3.3) and (17.12).

Finally, all local Taylor and Puiseux coefficients at zero are obtained by algebraic operations at algebraic roots of rational polynomials. Gamma values at the negative half-integers contribute only a common $1/\sqrt{\pi}$ times rational factors. The covariance is rational. Consequently $\chi_j/\mathcal{G}_0(0)$, and hence every $\alpha_{r,j}$, is algebraic. The finite steps just described establish effectiveness. $\square$

The scope of “complete” is the same as in the remark at the end of Section 6.

18 First corrections by Gaussian moments

18.1 A coordinate-invariant first-correction formula

The local calculation can use any rational full-rank $d \times r$ matrix Ξ annihilating $\mathbf{1}$ and $\mathbf{s}$, because an invertible linear change of local coordinates cancels from the normalized coefficients. In particular, consecutive second-difference rows $(1, -2, 1)$ are convenient. They need not be the globally normalized Fourier coordinates of Section 17; only the local ratios are being computed.

Put $L_i(x) = \xi_i \cdot x$ and define

$$\mathcal{M}_j(x) = \frac{1}{r} \sum_i L_i(x)^j, \quad \mathcal{M}_{12}(x) = \frac{1}{r} \sum_i s_i L_i(x)^2.$$

The critical phase has cubic and quartic terms

$$\Psi_3(x) = \frac{\mathcal{M}_3(x)}{6}, \quad \Psi_4(x) = \frac{\mathcal{M}_4(x) - 3\mathcal{M}_2(x)^2}{24} - \frac{\mathcal{M}_{12}(x)^2}{8V}. \quad (18.1)$$

Indeed, writing $v = \log \tau(z)$, the critical equation for $\log(r^{-1} \sum_i e^{\xi_i \cdot z + s_i v})$ gives $v = -\mathcal{M}_{12}(z)/(2V) + O(|z|^3)$. Substitution into its cumulant expansion proves (18.1).

Let $\mathbf{a}(z) = \mathcal{G}_0(z)/\mathcal{G}_0(0)$ and write $\mathbf{a}' = \nabla \mathbf{a}(0)$ and $\mathbf{a}'' = \nabla^2 \mathbf{a}(0)$. Equations (17.11)–(17.12) give

$$\begin{aligned} r \alpha_{r,1} &= \frac{\mathcal{G}_1(0)}{\mathcal{G}_0(0)} - \frac{1}{2} \operatorname{tr}(Q^{-1} \mathbf{a}'') \\ &\quad + \mathbb{E} \left[\frac{(\mathbf{a}' \cdot X) \mathcal{M}_3(X)}{6} + \Psi_4(X) - \frac{\mathcal{M}_3(X)^2}{72} \right]. \end{aligned} \quad (18.2)$$

The negative sign in the final term comes from $i^6 = -1$. Omitting it changes the answer.

18.2 Computing the local kernel terms

Here are explicit formulas sufficient to reproduce the first correction. At zero markings put $P = P_0$ and expand the principal small root at $t = 1 - w^2$ as $u = 1 + a_1 w + a_2 w^2 + a_3 w^3 + O(w^4)$, as in Appendix A. Solving $P(u) = 1/(1 - w^2)$ gives

$$\begin{aligned} a_1 &= -\sqrt{2/P''(1)}, & a_2 &= -\frac{P'''(1)a_1^2}{6P''(1)}, \\ a_3 &= \frac{1 - \frac{1}{2}P''(1)a_2^2 - \frac{1}{2}P'''(1)a_1^2a_2 - \frac{1}{24}P''''(1)a_1^4}{P''(1)a_1}. \end{aligned} \quad (18.3)$$

When the nonprincipal interior roots $\omega \in \mathcal{R}_r$ are simple, where $\mathcal{R}_r$ is the multiset of roots of $Q_r(u)/(u-1)^2$ in $|u| < 1$, their w^2 coefficients are $1/P'(\omega)$. Therefore

$$\frac{\mathcal{G}_1(0)}{\mathcal{G}_0(0)} = \frac{3}{8} - \frac{3}{2} \left(\frac{a_3}{a_1} + 1 + \sum_{\omega \in \mathcal{R}_r} \frac{1}{\omega P'(\omega)} \right). \quad (18.4)$$

For nonsimple nonprincipal roots, compute the derivative of their analytic product instead of differentiating the individual roots; the general algorithm in Section 17 is unaffected. The roots for $r = 3, 5$ used below are simple (or absent).

For the amplitude derivatives, use

$$\log \mathcal{G}_0(z) = \log \kappa_*(z) - \frac{1}{2} \log V(z) - \frac{1}{2} \log(2\pi), \quad V(z) = \left. \frac{\partial^2}{\partial v^2} \log P_z(e^v) \right|_{v=\log \tau(z)}.$$

The root product gives

$$\log \kappa_*(z) = \log P_*(z) + \log \tau(z) + \sum_{\omega} \log \omega(z) - \xi_- \cdot z + \text{constant}.$$

Here ξ_- is the column for the minimum step. The derivatives of a simple nonprincipal root follow by implicit differentiation of $P_z(\omega(z)) = P_*(z)$:

$$\begin{aligned} \omega_j &= -P_{z_j}(\omega)/P'(\omega), \\ (\log \omega)_{jk} &= \frac{Q_{jk} - P_{z_j z_k}(\omega) - P_{uz_j}(\omega)\omega_k - P_{uz_k}(\omega)\omega_j - P''(\omega)\omega_j\omega_k}{\omega P'(\omega)} - \frac{\omega_j\omega_k}{\omega^2}. \end{aligned} \quad (18.5)$$

All quantities on the right are evaluated at zero markings. In addition,

$$(\log \tau)_{jk} = -\frac{1}{rV} \sum_i s_i \xi_{i,j} \xi_{i,k}, \quad V_j = \frac{1}{r} \sum_i s_i^2 \xi_{i,j}, \quad V_{jk} = \frac{1}{r} \sum_i s_i^2 \xi_{i,j} \xi_{i,k} - V Q_{jk}.$$

The last identity uses the symmetry $\sum_i s_i^3 = 0$. These formulas make (18.2) an exact algebraic calculation.

18.3 Three evaluated cases

The two exact rows of Section 7 give the second-order terms

$$\begin{aligned} A_2(n) &= \frac{4^n}{\sqrt{\pi} n^{3/2}} \left(1 - \frac{9}{8n} + \frac{145}{128n^2} + O(n^{-3}) \right), \\ A_3(n) &= \frac{\sqrt{3} 27^n}{2\pi n^2} \left(1 - \frac{11}{9n} + \frac{101}{81n^2} + O(n^{-3}) \right). \end{aligned} \quad (18.6)$$

For $r = 5$, formulas (18.2)–(18.5) give the value of $\alpha_{5,1}$ in (3.6),

$$\alpha_{5,1} = -\frac{13}{10} + \frac{13\sqrt{5}}{50},$$

by a contraction different from the one of Section 9. In the multinomial normalization this is

$$\frac{A_5(n)}{\kappa_5(5n)!/(5n(n!)^5)} = 1 + \frac{-9/10 + 13\sqrt{5}/50}{n} + O(n^{-2}), \quad (18.7)$$

that is, $\Delta_5/5$ in (9.3). The shipped `code/42-late-growing-first_correction.py` carries out the rational Gaussian moments and algebraic root differentiation exactly for $r = 3, 5$. It obtains

$$\alpha_{3,1} = -11/9, \quad \alpha_{5,1} = -13/10 + 13\sqrt{5}/50 = -0.71862232585005467893\dots$$

The value of $\alpha_{3,1}$ is obtained here from the general formula and agrees with the value read off the closed form in Section 7. These values are not estimates from sequence data and do not use the conjectured recurrence.

19 Primitive excursions and the giant component

An excursion is *primitive* if it is nonempty and has no zero-height prefix before its endpoint; primitive excursions are the arches of Section 10. A one-step zero excursion is primitive. Every excursion factors uniquely at its returns to height zero. Let $\mathcal{I}(\mathbf{x})$ enumerate primitive excursions; then

$$\mathcal{E} = \frac{1}{1 - \mathcal{I}}, \quad \mathcal{I} = 1 - e^{-\mathcal{L}}. \quad (19.1)$$

We retain the general critical composition (15.1) in the first theorem and put $\kappa = \mathcal{E}(\mathbf{p})$.

Lemma 19.1 (Primitive weights). *If $i_{\mathbf{k}} = [\mathbf{x}^{\mathbf{k}}]\mathcal{I}(\mathbf{p}\mathbf{x})$, then*

$$0 \leq i_{\mathbf{k}} \leq b_{\mathbf{k}}, \quad \sum_{\mathbf{k}} i_{\mathbf{k}} = 1 - \kappa^{-1}, \quad i_{\mathbf{k}_N} \sim \kappa^{-1} b_{\mathbf{k}_N}.$$

The last assertion holds also after a fixed admissible shift.

Proof. A primitive excursion of length $m > 1$ has a unique minimum among its cyclic vertices before the endpoint. Its m rotations are distinct: a nontrivial word period would give an earlier zero return. No rotation class can contain two primitive excursions. Consequently m times the number of primitive words of a fixed composition is at most the number of all bridge words of that composition, proving $i_{\mathbf{k}} \leq b_{\mathbf{k}}$. Length one is immediate. The mass identity follows from (19.1). Finally apply Corollary 15.3 with $z = -1$ and use $\mathcal{I} = 1 - e^{-\mathcal{L}}$. $\square$

Theorem 19.2 (A size-biased geometric law). *Choose uniformly an excursion of composition $N\mathbf{p}$, and let J_N be its number of primitive factors. For each fixed $k \geq 1$,*

$$\mathbb{P}(J_N = k) \longrightarrow k\kappa^{-2}(1 - \kappa^{-1})^{k-1}. \quad (19.2)$$

All fixed moments converge. In particular,

$$\mathbb{E}J_N \longrightarrow 2\kappa - 1, \quad \mathbb{P}(J_N = 1) \longrightarrow \kappa^{-2}. \quad (19.3)$$

For the OEIS rows take $\kappa = \kappa_r$ and $N = rn$.

For the rows this is the law of Theorem 10.1, proved there by the analytic transfer of Section 6: J_N is the number J_n of arches, $C_n = J_n - 1$, and $2\kappa_r - 1 = 1 + 2(\kappa_r - 1)$. The proof below is source 42's second route, by the cutoff argument of Theorem 15.2; it also covers every centred rational composition and every fixed moment.

Proof. Repeat the cutoff proof of Theorem 15.2 for i in place of b . Lemma 19.1 supplies the same global bound and the local ratio $i_{\mathbf{k}_N - \ell} / b_{\mathbf{k}_N} \rightarrow \kappa^{-1}$. With $\eta = 1 - \kappa^{-1}$, it follows that

$$(i^{*k})_{\mathbf{k}_N} / b_{\mathbf{k}_N} \longrightarrow k\kappa^{-1}\eta^{k-1}.$$

Divide by the total excursion coefficient, asymptotic to $\kappa b_{\mathbf{k}_N}$. A uniform bound of the form $Ck^{\varsigma+1}\eta^{k-1}$ follows by designating a largest component. Since $\eta < 1$, it proves convergence of all fixed moments by dominated summation. The limiting probability generating function is

$$\frac{v}{\kappa^2(1 - \eta v)^2}. \quad (19.4)$$

Differentiating at $v = 1$ proves the mean formula. $\square$

Theorem 19.3 (One giant primitive component). *Write a uniform excursion of composition $N\mathbf{p}$ as $\mathcal{U}_N\Omega_N\mathcal{V}_N$, where Ω_N is a longest primitive component and ties are resolved by choosing the first one. Then $(\mathcal{U}_N, \mathcal{V}_N)$ converges in total variation to a pair $(\mathcal{U}, \mathcal{V})$ of independent finite excursions with*

$$\mathbb{P}(\mathcal{U} = u) = \frac{\mathbf{p}^{\text{comp}(u)}}{\kappa}, \quad \mathbb{P}(\mathcal{V} = v) = \frac{\mathbf{p}^{\text{comp}(v)}}{\kappa}. \quad (19.5)$$

In particular $N - |\Omega_N| = O_{\mathbb{P}}(1)$.

Proof. Fix two concrete finite excursions u, v and put $\ell = \text{comp}(u) + \text{comp}(v)$. For all sufficiently large N , an excursion of the form ujv with j primitive of composition $N\mathbf{p} - \ell$ automatically has j as its unique longest factor. Its probability is the ratio of that primitive count to $e_{N\mathbf{p}}$. The shifted primitive asymptotic gives the limit $\mathbf{p}^\ell / \kappa^2$, exactly the product in (19.5).

The sum of these limiting probabilities over all ordered pairs of finite excursions is $\mathcal{E}(\mathbf{p})^2 / \kappa^2 = 1$. Pointwise convergence on this countable space to a distribution of total mass one implies total-variation convergence: restrict first to a finite subset with limiting mass at least $1 - \epsilon$, then control its complement. Tightness of the two finite decoration lengths follows. It also shows that the probability of competing macroscopic factors tends to zero. $\square$

This limit is compatible with a nontrivial number of small factors: a giant *primitive* component need not mean that the whole excursion is primitive.

19.1 A tight gap with an infinite-mean limit

For the uniform alphabet, let $D_{\text{dec}} = |\mathcal{U}| + |\mathcal{V}|$ be the limiting decoration length. Its probability generating function is

$$\mathbb{E}t^{D_{\text{dec}}} = \left(\frac{\mathcal{E}(t/r, \dots, t/r)}{\kappa_r} \right)^2. \quad (19.6)$$

The square-root expansion at $t = 1$ gives

$$\mathbb{E}t^{D_{\text{dec}}} = 1 - \frac{2\sqrt{2}}{\sqrt{V}}\sqrt{1-t} + O(1-t).$$

Transfer and summation along the permitted time lattice yield

$$\mathbb{P}(D_{\text{dec}} \geq m) \sim \frac{2\sqrt{2}}{\sqrt{V\pi}} m^{-1/2}, \quad \mathbb{E}D_{\text{dec}} = \infty. \quad (19.7)$$

For even r , the support is even; the factor two in coefficient transfer cancels the half-density when the tail is summed. Thus (19.7) holds as written for unrestricted integer $m \rightarrow \infty$. Tightness of $N - |\Omega_N|$ does not imply convergence of its expectation to a finite constant. This is a useful warning against an otherwise tempting interpretation of the giant-component theorem.

20 Ratios, curvature and an explicit rounding bound

20.1 Ratios and eventual strict log-convexity

With $\beta = (r+1)/2$ and $\lambda = r \log r$ as in Section 11, the complete expansion gives

$$\frac{A_r(n+1)}{A_r(n)} = r^r \left(1 - \frac{\beta}{n} + \frac{\beta(\beta+1)/2 - \alpha_{r,1}}{n^2} + O_r(n^{-3}) \right). \quad (20.1)$$

Taking sufficiently many terms before taking second differences gives

$$\log A_r(n+1) - 2 \log A_r(n) + \log A_r(n-1) = \frac{\beta}{n^2} + O_r(n^{-3}) > 0 \quad (20.2)$$

for all sufficiently large n . Therefore every fixed row is eventually strictly log-convex, and its successive ratios eventually increase to r^r from below. This is not a claim of log-convexity at every finite index. A remainder controlled only by $o(1)$ would not suffice to infer the curvature; the higher-order expansion supplies the needed control.

20.2 The inverse dominant sector

Source 42 inverts the dominant sector exactly as Section 11 does: the large real solution of $Y = K_r e^{\lambda x} x^{-\beta}$ is (11.2), on the lower real branch (the principal branch would select the small solution, not the large inverse); the formula follows by raising the model equation to the power $-1/\beta$ and applying the defining equation $W e^W = z$. Its correction terms are those of (11.4), with $b_1 = \alpha_{r,1}$ and $b_2 = \alpha_{r,2} - \frac{1}{2}\alpha_{r,1}^2$, obtained by successive formal substitution or Newton correction; at every order the next coefficient is determined linearly, since the derivative of the leading logarithm is $\lambda - \beta/x$, bounded away from zero for large x . For orientation, writing $L_Y = \log Y$ gives

$$x(Y) = \frac{L_Y + \beta \log L_Y - \log K_r - \beta \log \lambda}{\lambda} + O_r \left(\frac{\log L_Y}{L_Y} \right). \quad (20.3)$$

For A215570 the first correction to the Lambert model is positive, because $\alpha_{5,1} < 0$; compare (11.5).

20.3 Discrete thresholds and the rounding boundary

Let $N_r(Y) = \min\{n : A_r(n) \geq Y\}$ for sufficiently large Y , as in Section 11.3. The sequence is eventually strictly increasing by (20.1), so this tail threshold is unambiguous. (2026-10-05 note: for $r \geq 3$ it is strictly increasing at every index, Proposition 37.3, so $N_r(Y)$ is unambiguous

for every Y .) Let $x_M(Y)$ solve the logarithmic expansion truncated after x^{-M} . There exists $\varpi_{r,M}$ such that, for all large Y ,

$$\left\lceil x_M(Y) - \varpi_{r,M}(\log Y)^{-M-1} \right\rceil \leq N_r(Y) \leq \left\lceil x_M(Y) + \varpi_{r,M}(\log Y)^{-M-1} \right\rceil. \quad (20.4)$$

This follows by comparing the logarithmic error at integer arguments with the derivative bounded away from zero. Away from the shrinking neighborhoods of integers the two bounds coincide. Near an integer, the error interval is necessary; a blanket claim that one may always take a ceiling of a truncated transseries would be false.

Remark 20.1 (Repository context). Inequality (20.4) is an instance of parts (2)–(3) of the staircase theorem `p0:thm:staircase` of the canonical transseries volume (Section 1.4), with $\eta = \varpi_{r,M}(\log Y)^{-M-1}$; its separation clause is formalized as the Lean theorem `Fabius.staircase_separation` and the failure of separation for arbitrarily small η as `Fabius.staircase_separation_fails`. Those are general statements about ceilings; they do not verify the asymptotic inputs of this report, and no novelty is claimed for the rounding statement.

21 Source 42’s computations

21.1 Exact dynamic programming

For an arbitrary composition $\mathbf{k}$, let $\text{ht}(\mathbf{k}) = \mathbf{s} \cdot \mathbf{k}$. Excursion-prefix counts satisfy

$$F(\mathbf{0}) = 1, \quad F(\mathbf{k}) = \begin{cases} 0, & \text{ht}(\mathbf{k}) < 0, \\ \sum_{i:k_i > 0} F(\mathbf{k} - \mathbf{e}_i), & \text{ht}(\mathbf{k}) \geq 0. \end{cases} \quad (21.1)$$

The desired value is $F(n\mathbf{1})$; this is the recurrence of Section 13, implemented independently. This recursion is exact: every admissible word is obtained uniquely by deleting its last letter. The implementation traverses a rectangular composition box in lexicographic order, which is topological for coordinate decrements. It uses integer arithmetic and no row recurrence conjecture.

Two companion arrays count primitive words and the sum of the numbers of returns. For the primitive array, do not propagate any nonempty zero-height state. For the moment array, add the current count whenever a newly reached state has height zero. This gives independent finite tests of both (19.2) and (19.3).

r	n	$A_r(n) / (\kappa_r(rn)! / (rn(n!)^r))$	Primitive probability	Mean returns
4	30	0.993523253212	0.527972597714	1.74565492347
5	15	0.979218847813	0.283110782306	2.70465628143
6	9	0.991084348462	0.662077457976	1.45242243648
7	5	0.971363174805	0.417632780048	2.04381815151

Table 4: Exact integer counts of source 42 evaluated against the proved limiting formulas. These modest-index ratios illustrate convergence but do not certify asymptotic error bounds. “Mean returns” is the mean number $\mathbb{E}J_N$ of primitive factors, whose limit is $2\kappa_r - 1$ (Table 3).

21.2 Independent coefficient and algebra checks

Source 42’s program performs the following finite checks.

- (a) The first seven $r = 4, 5$ values are checked against the displayed OEIS data. Full independently computed ranges are $0 \leq n \leq 30$ for $r = 4$, $0 \leq n \leq 15$ for $r = 5$, $0 \leq n \leq 9$ for $r = 6$, and $0 \leq n \leq 5$ for $r = 7$.
- (b) The bridge-excursion identity is checked coefficientwise in small multivariate boxes using the independent Euler identity (13.1). The tested boxes use $(r, \text{bound}) = (2, 5), (3, 4), (4, 3), (5, 2)$.
- (c) The sextic elimination identity is verified as an exact rational-function identity, including its specified branch at the origin.
- (d) Kernel polynomials are deflated exactly by $(u - 1)^2$. Their interior roots are evaluated at high precision for $2 \leq r \leq 12$ and the four radical constants are cross-checked. These decimal evaluations are not interval-certified root isolations.
- (e) The first-correction formula is evaluated symbolically for $r = 3, 5$ using exact Gaussian moments. The $r = 3$ result is independently checked against its elementary exact formula.

The shipped `data/42-late-growing-verification.txt` and `data/42-late-growing-first_correction.txt` record the successful runs. The CSV files retain exact counts and decimal diagnostics. All the supplied assertions passed in the recorded run. The full dynamic program stores arrays of size $(n + 1)^r$, so memory, rather than asymptotic arithmetic sophistication, is the main practical limitation of this deliberately simple independent verifier.

21.3 Claims that remain outside the certificate

The proofs are mathematical arguments in this report, not Lean or Rocq proofs. The scripts certify their stated finite computations, not every analytic estimate. No numerical fit is used to prove an asymptotic coefficient. No low-order recurrence is used as an assumption. No actual recurrence operator for $r = 4$ is supplied. No claim of exhaustive literature priority, uniformity in r , exponentially improved asymptotics, or an explicit finite log-convexity threshold is made.

Part III

Research directions and conclusion

22 Further research questions and topics

The research directions of the two sources are merged here: each subsection states source 61's question and then what source 42 adds; two subsections are source 42's alone. The eight questions of source 75-03 (Part IV) were added on October 2, 2026, as sentences marked with its name. The A215570 question of sources 61 and 42 is answered by source 75-03, and source 75-03's question on all orders for every row by Theorem 3.1. (2026-10-02 note: the four questions of source 77-44, Part V, coincide with subsections below; they are listed in Section 34.) (2026-10-05 note: the questions of sources 101-90 and 101-86, Part VI, coincide with subsections below and are credited there; Section 38.4 lists them as items 101.1–101.6.)

22.1 Certify the explicit A215570 recurrence (resolved)

The sextic (4.5), its specified formal branch, and a diagonal representation give a concrete input for creative telescoping. The next objective is to compute a certified annihilator and prove

that the order-three operator in Kauers–Koutschan Conjecture 15 is a right factor annihilating the same solution, with all initial conditions and singular indices checked. Merely matching additional terms would not accomplish this. A useful outcome would be a compact certificate independently checkable without a large symbolic-computation environment. Source 42 (its project 1) proposes the same route and asks for a machine-readable telescoping certificate and an independent checker: resolving the named conjecture, not just the existence of some recurrence, is the point.

Resolved (2026-10-02). Source 75-03 proves the conjecture (Theorem 26.1) with exactly such a compact certificate (Appendix C) and an exact checker, for every $n \geq 0$, so that no initial conditions or singular indices remain to be checked. Its route differs from the one proposed here: it does not eliminate from the sextic, but reduces the diagonal to the one-variable coefficient (25.7) and telescopes its Legendre residue representation. Minimality of the order-three operator is not claimed. The recurrence questions for A215562 and the higher rows remain open (next subsection).

22.2 Find and explain the fourth-row recurrence

A215562 is D-finite by the proof here, but its minimal recurrence and its computational complexity remain to be determined. The even-step parity and the larger kernel degree should be incorporated before elimination, not left for a generic telescoper to rediscover. What upper bounds on recurrence order and coefficient degree can be proved as functions of r ? Do the even and odd alphabets have qualitatively different minimal orders? Source 42 (project 2) notes that the generic small-root subset polynomial has degree at most $\binom{6}{3} = 20$ in this case, proposes exploiting the reciprocal and label-reversal symmetries before telescoping, and asks for a comparison of recurrence order, coefficient degree and bit complexity with direct composition dynamic programming. Source 75-03 asks for the same operator: eliminate the product of three small roots of the marked degree-six kernel and compute a diagonal telescoper, ideally with a compact rational certificate. The degree bound 20 suggests a larger problem than the five-symbol sextic but does not predict the minimal order; a smaller auxiliary coefficient representation, analogous to (25.7), would be especially valuable. (2026-10-05 note: source 101-90 lists “explicit minimal recurrences” among its open questions; its D-finiteness proof by an r -section, (36.17), is again existence only; item 101.1 of Section 38.4.)

22.3 Automate all-order coefficient extraction

The proof supplies finite algebraic operations, but a practical system should exploit residual-marker coordinates, symmetry, and Wick contractions to avoid dense high-dimensional tensors. An immediate target is exact $\alpha_{r,1}, \alpha_{r,2}, \alpha_{r,3}$ for $r = 4, 6, 7, 8$, together with machine-readable algebraic-number specifications. (2026-10-02 note: source 77-44 supplies $\alpha_{4,1}$ exactly in $\mathbb{Q}(\varrho)$, Theorem 31.1; $\alpha_{4,2}, \alpha_{4,3}$ and the rows $r = 6, 7, 8$ remain open.) For $r = 5$, the field $\mathbb{Q}(\sqrt{5})$ makes an especially clean regression suite. Source 42 (project 3) proposes the same engine built on (17.10) and (17.12), with exact algebraic numbers, branch-isolation certificates, symmetry reduction and regression tests from $r = 2, 3$, and targets $\alpha_{r,2}, \alpha_{r,3}$ for $r = 4, 5, 6$; its first-correction program is a prototype, not this engine. For $r = 5$ the target is met through $\alpha_{5,8}$ (Proposition 27.1), by the recurrence rather than by such an engine. Source 75-03 asks whether its transfer argument, which gives only a nonsharp $O(n^{-1/5})$ error, can be refined to all orders for every fixed alphabet; Theorem 3.1 already does so, so its question reduces to the targets above (it names an explicit, proved $1/n$ correction for $r = 4, 6, 7$; 2026-10-02 note: for $r = 4$ it is Theorem 31.1, for $r = 6, 7$ still open) and to a uniform version of the convolution method giving a better general error without the whole expansion. It also asks for the minimal polynomials

and degrees of the constants (5.6) and their dependence on parity, whether reciprocal symmetry reduces the generic subset-product degree bound substantially, and for rational isolating boxes of the inner roots that prove decimal intervals for κ_r . (2026-10-05 note: source 101-86 meets this target for $r = 5$ at second order *by the engine* rather than by the recurrence: Theorem 37.2 computes $\alpha_{5,2}$ by exact Gaussian moments, in agreement with Proposition 27.1. Source 101-90 states the rule for every r in marker coordinates, (36.16); evaluated at first order it reproduces $\alpha_{4,1}$ and $\alpha_{5,1}$ numerically, Remark 36.5. Its non-claim on the smallest field of the coefficients is item 101.5 of Section 38.4, the engine item 101.6. The values $\alpha_{4,2}$, $\alpha_{4,3}$ and the rows $r = 6, 7, 8$ remain open.)

22.4 Determine sharp asymptotic error constants

Turn the separated kernel contours and local Taylor bounds into explicit numerical inequalities. This would provide certified approximation intervals at finite n , certified threshold inversion near an integer boundary, and explicit eventual-log-convexity thresholds (source 42, project 4). Source 75-03 asks the same for the constant C of its rounded threshold for A215570 (Section 28): explicit finite- Y bounds that bracket exact threshold searches. (2026-10-05 note: sources 101-86 and 101-90 ask the same for their inverses, which assert “no numerical error constant or unconditional exact-ceiling rule”; item 101.3 of Section 38.4.)

22.5 Determine the exponentially small sectors

The dominant expansion does not identify subdominant critical points, connection constants, or Stokes phenomena. Once an explicit differential or recurrence operator is available, compare its formal solutions with complex kernel saddles. Which exponentially small contributions are actually present in the combinatorial solution? Can optimal truncation and rigorous exponentially improved error bounds be obtained without first computing a minimal recurrence? Source 42 (project 9) asks the same: identify non-dominant singularities of the algebraic kernel and the relevant complex composition saddles, which of them contribute after diagonal extraction, their exponential scales, and their Stokes transitions under parameter changes. This is what would be needed to replace the dominant-sector expansion by a more complete transseries. For A215570 the question is now concrete (source 75-03): the proved recurrence has normalized characteristic roots $1, 1, 2/27$ (Remark 27.2). Determine the decomposition of the actual solution into its dominant and exponentially smaller components, with connection constants, and whether and how optimally truncated dominant expansions detect the third mode. (2026-10-05 note: sources 101-86 and 101-90 list the smaller exponential sectors among their open questions; item 101.2 of Section 38.4.)

22.6 Let the alphabet grow

Our estimates keep r fixed. When $r = r(n)$ grows, kernel-root separation, Gaussian dimension, and constants in the remainder estimates all change. Determine useful uniform ranges, and analyze the large- r behavior of κ_r separately along even and odd alphabets. This would connect the fixed-row problem to the less regular column direction of A215561. Source 42 (project 5) adds that the parity split is real: odd alphabets include a zero step, while even alphabets have time period two, so fixed- r error constants cannot simply be reused. It asks for expansions of $\log \kappa_r$ separately for even and odd r , and for a range of joint growth $r = r(n)$ in which the fixed-row formula remains uniform. Source 75-03 asks for the same simultaneous limit and, separately, for the column $n = 1$ of the array, which is A147681; none of the fixed-row theorems settles that column problem. (2026-10-05 note: source 101-90 asks for

“uniform regimes in which the alphabet size grows”; in its coordinates the multiplicity $2h$ and $\det H_{\text{ex}} = 12(r-1)/(r^r(r+1))$ show the r -dependence explicitly; item 101.4 of Section 38.4.)

22.7 Resolve the mean size of the non-giant remainder

The limit gap D_{dec} has an $m^{-1/2}$ tail and infinite mean (19.7). Determine the growth of $\mathbb{E}(N - |\Omega_N|)$ and higher moments under exact composition conditioning, including their leading constants. Establish a transition law when the gap is a positive fraction of N and quantify the probability of two large components. Weak convergence alone does not answer these questions (source 42, project 6).

22.8 Approach the boundary of the composition simplex

In Corollary 12.1, positivity of all probabilities is essential for the full-dimensional saddle. If some p_i tend to zero, the determinant in (6.6) degenerates and the effective alphabet may change. Find uniform transition formulas interpolating between adjacent alphabet sizes and identify the scaling regimes in which a different asymptotic power appears. Source 42 (project 8) also asks at which scale the giant-component law of Theorem 19.3 changes; its Theorem 16.1 likewise assumes that every p_i is strictly positive. Source 75-03 asks the same when some type frequencies tend to zero with N : the multinomial covariance degenerates, the effective number of types can change and the power $(r+1)/2$ need not stay uniform; a useful theory would connect the full-dimensional regime to the lower-dimensional faces with explicit crossover scales.

22.9 Refine path statistics under exact composition

The contact law is only one consequence of the marked kernel. Natural next targets are its finite- n corrections, the maximum height, the area under the excursion, and functional convergence after diffusive scaling. These require additional arguments: neither the count asymptotic nor the contact law alone proves a Brownian-excursion limit or independence of other statistics. Source 42 (project 7) asks for the functional limit after diffusive height scaling jointly with the Boltzmann decorations of Theorem 19.3 and the primitive-factor count. Exact composition is a stronger conditioning than an endpoint constraint, and a satisfactory theorem should specify the covariance normalization and decide whether the limiting giant shape is asymptotically independent of the decorations. Source 75-03 asks for the same functional limit under exact content, expecting a Brownian-excursion-type limit with a content-determined variance, and names the height maximum, the area and the locations of short primitive components as observables; these need probability arguments beyond the scalar asymptotic.

22.10 Formalize the exact-to-analytic interface

A staged ProveIt development could first formalize the word-walk bijection, the finite-state recurrences, the bridge coefficient identity, and the formal algebraic kernel relation. A second stage would address algebraic-series diagonal closure and critical-root selection. The most substantial analytic stage is a reusable uniform marked square-root transfer theorem with a nondegenerate composition saddle and explicit lattice periods. Keeping these stages separate would prevent finite computations from being mistaken for proofs of analytic hypotheses. Source 42 (project 10) proposes to begin instead with its convolution route: the exact bridge coefficients, the factorial ratio lemma, the convolution cutoff bound and dominated exponential summation (Section 15) give the leading growth theorem with a relatively small analytic

interface, so that the substantial all-orders complex analysis need not be a prerequisite for a formally verified leading result. The kernel identity and the multivariate saddle estimates can be later modules. Source 75-03 proposes a formalization of the recurrence proof as a chain of small reusable certificates: content vectors, prefix validity, reversal and zero-step insertion; the marked kernel algebra and the Lagrange extraction; the coefficient identities and the polynomial certificate in a rational-function ring; the Legendre recurrence and derivative identities with the vanishing residue of a rational derivative; and, separately, the Fourier bounds and the composition-transfer theorem. These are distinct obligations: checking the printed certificate alone does not establish the counting semantics.

23 Conclusion

The fixed-row problem is governed by a finite one-dimensional step set, not by an intrinsically growing transfer matrix. Marking composition before taking a diagonal makes this structure visible. The resulting algebraic ambient series gives D-finiteness; its bridge logarithm gives an exact reference coefficient; and a marked critical-point calculation converts that coefficient into the conjectured row asymptotic with an algebraic constant.

The same analysis yields a complete dominant inverse-power expansion, a concrete correction for A215570, a contact distribution, and an asymptotic inversion procedure. A separate arithmetic argument shows that every row generating function above the Catalan case is transcendental. These results settle the stated fixed-row growth conjecture and the recurrence-existence issue while leaving the explicit recurrence certificates and exponentially small sectors as sharply specified next problems. (2026-10-02 note: Part IV supplies the certificate for A215570; the one for A215562 remains open.)

Source 42 reaches the leading formula by a different mechanism: along a strictly positive balanced composition ray, the coefficients of an exponential of the log-excursion series are dominated by one large summand, and the remaining summands contribute exactly the critical excursion value. The same mechanism, applied to the factorization into primitive excursions, explains the limiting number of returns and the single giant component with its heavy-tailed finite decorations. Certifying minimal recurrences, obtaining effective finite error bounds, and passing to growing alphabets are distinct next steps, not conclusions hidden inside the present theorems.

Source 75-03, finally, finds for the five-symbol row an exact representation much smaller than the content-state dynamic program: a positive one-variable coefficient with a Legendre-polynomial interpretation. It yields an explicit proof of the published order-three recurrence and, with the first correction, every dominant asymptotic coefficient and its inverse, while an explicit four-symbol recurrence, uniform growing-alphabet limits and the exponentially small sectors stay separate tasks.

Source 77-44 reaches the leading theorems a fourth time, by the same kernel, saddle and mark-extraction route, and contributes the first correction of the four-symbol row; the recurrence for that row is still open.

Sources 101-90 and 101-86 reach the leading theorems a fifth and a sixth time by direct marked-excursion extraction in two further coordinate systems, which show that the number of dominant torus points belongs to the coordinates, not to the problem. They add a recurrence-free second correction for A215570, the constant of its zero-free count, and the supermultiplicativity that makes every row with $r \geq 3$ strictly increasing at every index.

Part IV

The Kauers–Koutschan recurrence and the five-symbol row

Part IV was added on October 2, 2026, in batch 75 of ProveIt’s incoming-reports intake, from a third manuscript on the same problem (source 75-03), after the closing Part III. It proves what Parts I and II left open and Part III lists as an open question: the explicit order-three recurrence for A215570 conjectured by Kauers and Koutschan (Theorem 26.1). With that recurrence it computes the exact coefficients $\alpha_{5,2}, \dots, \alpha_{5,8}$ of (3.3) for $r = 5$ (Table 6). About two thirds of source 75-03 re-proves results of Parts I and II, some of them in a weaker form; those results are *not* printed again but listed, with the comparison, in Table 5. The rest of the report is unchanged apart from dated notes that point here, rows added to Tables 1 and 2, the merged research questions of Part III and the package description of Appendix B; no existing label or number was changed.

24 Provenance, overlap and conventions of Part IV

24.1 A third manuscript

Source 75-03 is batch-75 manuscript 03, *Fixed-Content Ballot Words. A proof of the A215570 recurrence, a fixed-row asymptotic theorem, and all-orders inversion*, dated October 1, 2026 and prepared for Vladimir Reshetnikov (a 917-line source with 87 labels and a 22-page PDF). Archive `OEIS_Fixed_Content_Ballot_Words.zip`, arrival commit `4b874cea0`, placed in commit `6ea60e367`; it pins the repository at `29aca108e` [16]. Its programs and data ship with the prefix `75-03-ballot-words-` (Section 29).

Source 75-03 was written independently of sources 61 and 42: about 1.8% of its 8-word shingles occur in this report (0.8% of the report’s occur in it), all of them preamble, bibliography and OEIS strings. Its pin (October 1, 19:03) predates this report’s placement (`6e193dd4f`, 19:16) by thirteen minutes and already contained the archives of sources 61 and 42 as unopened zip files. Its repository check (“its overview, combinatorics overview, and a targeted search for A215561”) could therefore not see this report, exactly as source 61’s search could not see source 42 (Section 2.3). It states that the check “was not an audit of every research artifact”. Its own status statement is kept: its results “have explicit mathematical proofs; the large recurrence identity additionally has an exact symbolic checker. They have not been independently peer-reviewed or formalized in Lean”; its source check of October 1, 2026 found the statements proved here still recorded as conjectural, but “it is not an exhaustive priority search”; and classical kernel, diagonal and saddle-point methods are credited rather than claimed.

Like sources 61 and 42, source 75-03 is AI-assisted and unrefereed, and no Lean or Rocq development checks any statement of this Part. Source 75-03 outlines a staged formalization (Section 22, last subsection); none exists.

24.2 What the third manuscript re-proves

Table 5 lists every result of source 75-03 that Parts I and II already contain, where it is printed, and how the two versions compare. Its main fixed-row theorem is a *third* proof of the leading term of Theorem 3.1, by the one-large-summand route of source 42 (Section 15)

made quantitative, and it is weaker than Theorem 3.1: a relative error $O(n^{-1/5})$ instead of a complete expansion. Source 75-03 presents it as an independent proof of the A215561 conjecture; in this report it is the third.

Table 5: Results of source 75-03 that Parts I and II already prove.
 “Same” means the same statement; the last column records where
 source 75-03 is weaker, stronger or merely different.

source 75-03	printed here	comparison
bridge factorization $\mathcal{E} = \exp \mathcal{L}$, algebraicity, small-root formula (its Thm. 2.1)	Lemmas 4.1, 5.1; (14.3)	same; the residue proof is source 61’s first proof
D-finiteness of every row (Cor. 2.2)	Theorem 3.2	same; its degree bounds $\binom{r-1}{(r-1)/2}$ (r odd) and $\binom{2r-2}{r-1}$ (r even), 6 for $r = 5$ and 20 for $r = 4$, are the bound $\binom{h+h'}{h}$ of Section 14
mass, central size, fixed shifts (Lemma 3.1)	Lemma 15.1	same, plus the uniform shift rate $1 + O((J + J^2)/N)$ and the tail bound $O(M^{-1/2})$ (Remark 24.1)
central composition transfer (Thm. 3.2), balanced excursions (Cor. 3.3)	Theorem 15.2, Corollary 15.3, Theorem 16.1	adds the rate $O(N^{-1/5})$ and F analytic on a disk of radius $> \mathfrak{B}$ instead of $F = \exp(z \cdot)$; Corollary 12.1 gives all orders for the rows’ step set
fixed-row asymptotic (Thm. 4.1)	Theorem 3.1	weaker: $O(n^{-1/5})$ against a complete expansion; same κ_r and $c(r)$
root-product formula (5.6) (its (4.5))	Proposition 5.2	same; its Q_r is $-Q_r$ here; inside roots counted by reciprocity, as source 42 does
$\kappa_4 = 4(\varphi - \sqrt{\varphi})$, $\kappa_5 = \frac{5}{2}(3 - \sqrt{5})$, A_2, A_3 closed forms, constants $r \leq 8$	Section 7, Table 3	same values to every printed digit; κ_4 ’s closed form is already printed there and credited in OEIS to Kotesovec [2]
marked sextic (Prop. 5.1)	(4.5)	the same polynomial term by term; a third derivation (Remark 25.1)
existence of the all-orders expansion for A215570, $\alpha_{5,j} \in \mathbb{Q}(\sqrt{5})$ (Thm. 7.1)	Theorems 3.1, 3.3	same; a second route by a two-variable saddle (Section 27.1)
$\alpha_{5,1} = -13/10 + 13\sqrt{5}/50$	(3.6)	same value; a third route, (27.3)
primitive-factor law, mean $2\kappa - 1$ (Thm. 8.1)	Theorems 10.1, 19.2	same, plus the limiting variance (24.2)
Lambert- W_{-1} anchor, first inverse corrections (Section 9)	(11.2), (11.4), (20.3)	same; for $r = 5$ the second-order coefficient is now explicit, (28.1)
rounded threshold $\lceil x_* \mp Cx_0^{-3} \rceil$	(20.4) with $M = 2$	same; an instance of <code>p0:thm:staircase</code>
monotonicity by appending a balanced block	Section 11.3	same argument

Two increments over Parts I and II are small enough to state here.

Remark 24.1 (A rate in the one-large-summand theorem (source 75-03)). In the notation of Section 15, with $q = r$ steps of the list (15.1), source 75-03 proves for every F analytic on a disk $|z| < R$ with $R > \mathfrak{B}$

$$\frac{[x^{k_N}]F(\mathcal{L}(px))}{b_{k_N}} = F'(\mathfrak{B}) + O(N^{-1/5}), \quad (24.1)$$

an additive estimate (not a relative equivalence when $F'(\mathfrak{B}) = 0$). The inputs are the two bounds of Lemma 15.1 with $\varsigma = (r + 1)/2$, the tail bound $\sum_{|k| > H} b_k = O(H^{-1/2})$ (from $\mathbb{P}(S_m = 0) = O(m^{-1/2})$), and the uniform shift ratio $b_{\mathbf{k}_N - \ell}/b_{\mathbf{k}_N} = 1 + O((L + L^2)/N)$ for $L = |\ell|$ with $L^2/N \rightarrow 0$. The cutoff proof of Theorem 15.2 is run with the cutoff $H = \lfloor N^{2/5} \rfloor$ and only for $m \leq N^{1/20}$ summands, so that $(mH)^2/N \leq N^{-1/10}$; the errors are $O(H^2/N + H^{-1/2}) = O(N^{-1/5})$, and the terms $m > N^{1/20}$ are exponentially small by (15.5) and Cauchy's estimate. With $F = \exp$ this gives the leading term of Theorem 16.1 with relative error $O(N^{-1/5})$ for every finite labelled step list, and for the rows the leading term of Theorem 3.1 with relative error $O(n^{-1/5})$. Source 75-03 calls the rate “deliberately nonsharp”; for the rows' step set it is superseded by Theorem 3.1 and Corollary 12.1.

The limiting probability generating function (19.4) is v times a negative binomial law of order two, so the number J_N of primitive factors (arches) has limiting variance

$$\text{Var } J_N \longrightarrow 2\kappa(\kappa - 1), \quad (24.2)$$

in the notation of Theorem 19.2 (for the rows also $\text{Var } C_n \rightarrow 2\kappa_r(\kappa_r - 1)$, since $C_n = J_n - 1$ in Theorem 10.1); convergence of the variance follows from the convergence of all fixed moments proved there. Source 75-03 states this variance with the mean $2\kappa - 1$, which for $r = 5$ is $14 - 5\sqrt{5}$, the last column of Table 3.

24.3 Conventions of Part IV

Part IV uses the merged notation of Section 1.5. Table 2 lists every symbol of source 75-03 that is renamed; no normalization changed except the sign of Q_r . In particular source 75-03's asymptotic coefficients d_j are the $\alpha_{5,j}$ of (3.3), its constant $\mathcal{E}_r$ is κ_r , and its excursion series Y of the sextic is the X of (4.5). Three readings are tempting and false:

- Λ_n (source 75-03's C_n , the Legendre coefficient of Theorem 25.2) is not the number C_n of interior returns of Section 10, not the functional $\mathcal{C}_N$ of (6.2), and not the lattice Λ of Section 17;
- $\mathcal{P}_n(t)$ is the Legendre polynomial with $\mathcal{P}_n(1) = 1$, not the step polynomial $P_r(u)$ of (3.1): $\mathcal{P}_5(t) \neq P_5(t)$;
- $A_5^o(n)$ (source 75-03's d_n) counts the zero-free excursions of (25.3); it is not an asymptotic coefficient.

The marking variables a, b, c, d, e of steps $-2, \dots, 2$ are those of (4.5); the letter a is never the sequence.

25 A positive coefficient formula for A215570

25.1 The sextic in compact form

Put

$$A = ae, \quad B = bd, \quad C = ad^2 + b^2e, \quad \bar{c} = 1 - c.$$

Then (4.5) reads

$$A^3X^6 - A^2\bar{c}X^5 + A(B - A)X^4 + (2A\bar{c} + C)X^3 + (B - A)X^2 - \bar{c}X + 1 = 0, \quad (25.1)$$

term by term the same polynomial; source 75-03 states it in this form.

Remark 25.1 (Third derivation of the sextic (source 75-03)). Let $e_1, \dots, e_4$ be the elementary symmetric functions of the roots of the kernel quartic $eu^4 + du^3 + (c-1)u^2 + bu + a$. The six pair products $u_i u_j$ ($i < j$) are the roots of the resolvent

$$R(v) = v^6 - e_2 v^5 + (e_1 e_3 - e_4) v^4 - (e_3^2 + e_1^2 e_4 - 2e_2 e_4) v^3 \\ + e_4(e_1 e_3 - e_4) v^2 - e_2 e_4^2 v + e_4^3,$$

by expansion of $\prod_{i < j} (v - u_i u_j)$. The two small roots have product $-aX$ (Lemma 4.1). Substituting $(e_1, e_2, e_3, e_4) = (-d/e, (c-1)/e, -b/e, a/e)$ and $v = -aX$ and multiplying by e^3/a^3 gives (25.1), as a polynomial identity; its X -derivative at the origin is -1 , which selects the formal branch. This is a third route beside the quadratic factorization of source 61 and the explicit coefficient solving of source 42 (Remark 4.2); the program `code/75-03-ballot-words-verify.py` checks the resolvent and its specialization symbolically.

With $U = AX^2$, (25.1) becomes $(1 - U)^2(1 + U - \bar{c}X) + (1 + U)BX^2 + CX^3 = 0$. Since the denominators below have constant term one, it is equivalent, in formal power series, to

$$X = 1 + cX + AX^2 + BX^2 \frac{1 + AX^2}{(1 - AX^2)^2} + CX^3 \frac{1}{(1 - AX^2)^2}. \quad (25.2)$$

25.2 Zero-step deletion and Lagrange inversion

Let $A_5^\circ(n)$ count nonnegative excursions with n copies of each of the steps $-2, -1, 1, 2$ and no zero step. Removing the n zero steps of an excursion counted by $A_5(n)$ and reinserting them gives

$$A_5(n) = \binom{5n}{n} A_5^\circ(n), \quad (25.3)$$

the same deletion that gives $A_3(n)$ in Section 7. Set $c = 0$ and give A, B, C the weights $2, 2, 3$. If $\mathbf{n} = 2i + 2j + 3k$, Lagrange inversion applied to $\mathcal{Y} = zX(z^2A, z^2B, z^3C)$, which by (25.2) satisfies $\mathcal{Y} = z(1 + A\mathcal{Y}^2 + B\mathcal{Y}^2(1 + A\mathcal{Y}^2)(1 - A\mathcal{Y}^2)^{-2} + C\mathcal{Y}^3(1 - A\mathcal{Y}^2)^{-2})$ for an auxiliary variable z , gives

$$[A^i B^j C^k]X = \frac{\mathbf{n}!}{j! k! (\mathbf{n} + 1 - j - k)!} [u^i] (1 + u)^{\mathbf{n}+1-k} (1 - u)^{-2(j+k)} : \quad (25.4)$$

select j copies of the B term and k of the C term in the $(\mathbf{n} + 1)$ st power of the Lagrange formula; the remaining powers of $1 + A\mathcal{Y}^2$ have total exponent $\mathbf{n} + 1 - k$, and extracting A^i gives (25.4). Equal powers of a, b, d, e force the two monomials of C to be used equally often, say ℓ times each; then $i = n - \ell$, $j = n - 2\ell$, $k = 2\ell$, $\mathbf{n} = 4n$, and $C^{2\ell}$ contributes $\binom{2\ell}{\ell}$. Hence

$$A_5^\circ(n) = \frac{(4n)!}{(3n + 1)!} \sum_{\ell=0}^{\lfloor n/2 \rfloor} \frac{[u^{n-\ell}] (1 + u)^{4n+1-2\ell} (1 - u)^{-2n}}{(n - 2\ell)! (\ell!)^2}. \quad (25.5)$$

25.3 A Legendre coefficient

Let $\mathcal{P}_n(t)$ be the Legendre polynomial, $\mathcal{P}_n(1) = 1$. Comparing coefficients of v^n in $((1 - v)^2 - 4zv^2)^{-1/2}$ gives

$$\sum_{\ell=0}^{\lfloor n/2 \rfloor} \frac{n! z^\ell}{(n - 2\ell)! (\ell!)^2} = (1 - 4z)^{n/2} \mathcal{P}_n((1 - 4z)^{-1/2}).$$

Substituting $z = u/(1 + u)^2$ in (25.5), for which $(1 - 4z)^{1/2} = (1 - u)/(1 + u)$, proves the following formula.

Theorem 25.2 (Positive coefficient formula (source 75-03)). *For $n \geq 0$ put*

$$\Lambda_n = [u^n](1+u)^{3n+1}(1-u)^{-n}\mathcal{P}_n\left(\frac{1+u}{1-u}\right) \quad (25.6)$$

$$= [u^n](1+u)^{3n+1}(1-u)^{-2n} \sum_{k=0}^n \binom{n}{k}^2 u^k. \quad (25.7)$$

Then

$$A_5(n) = \frac{(5n)!}{(n!)^2(3n+1)!} \Lambda_n. \quad (25.8)$$

Proof. (25.6) and (25.8) follow from (25.5) and (25.3). The two forms of Λ_n agree because $(1-u)^n \mathcal{P}_n((1+u)/(1-u)) = \sum_k \binom{n}{k}^2 u^k$, by the binomial expansion of the Legendre polynomial. $\square$

The first values are $\Lambda_0, \dots, \Lambda_5 = 1, 7, 104, 1935, 39854, 869736$, giving $A_5(0), A_5(1), A_5(2) = 1, 35, 18720$. Formula (25.7) is a finite positive coefficient computation, independent of the recurrence proved next. It agrees with the 17 independently enumerated terms of source 61, the 16 of source 42 and the 13 data terms of OEIS A215570; and at $n = 20$ and $n = 50$, where Section 13 used OEIS b-file values, its values reproduce the ratios recorded in `data/61-balanced-diagnostics.json` to all 25 recorded digits (checked at the intake, 2026-10-02).

26 Proof of Kauers–Koutschan Conjecture 15

Normalize

$$\tilde{A}_5(n) = \frac{(n!)^3((n+1)!)^2}{(5n)!} A_5(n) = \mathfrak{h}_n \Lambda_n, \quad \mathfrak{h}_n = \frac{n!((n+1)!)^2}{(3n+1)!}, \quad (26.1)$$

and put

$$\begin{aligned} \mathfrak{R}(n) &= 65n^3 + 593n^2 + 1772n + 1740, \\ \mathfrak{S}(n) &= 65n^3 + 398n^2 + 781n + 496, \\ \mathfrak{U}(n) &= 2015n^5 + 24428n^4 + 114387n^3 + 258294n^2 + 281088n + 118368, \\ \mathfrak{V}(n) &= 910n^5 + 11032n^4 + 52047n^3 + 119686n^2 + 134365n + 58980. \end{aligned}$$

Theorem 26.1 (Kauers–Koutschan Conjecture 15 (source 75-03)). *For every integer $n \geq 0$,*

$$\sum_{j=0}^3 \mathfrak{p}_j(n) \tilde{A}_5(n+j) = 0, \quad (26.2)$$

where

$$\begin{aligned} \mathfrak{p}_0(n) &= -2(n+1)(n+2)\mathfrak{R}(n), & \mathfrak{p}_1(n) &= \mathfrak{U}(n), \\ \mathfrak{p}_2(n) &= -4\mathfrak{V}(n), & \mathfrak{p}_3(n) &= 3(3n+8)(3n+10)\mathfrak{S}(n). \end{aligned} \quad (26.3)$$

Equivalently, with $K = \{1, \dots, 14\} \setminus \{5, 10\}$, the sequence satisfies the recurrence displayed in OEIS A215570 [3]:

$$\begin{aligned} &3(n+1)(n+2)^3(n+3)^4(n+4)^2(3n+8)(3n+10)\mathfrak{S}(n)A_5(n+3) \\ &- 20(n+1)(n+2)^3(n+3)^2 \prod_{k=11}^{14} (5n+k)\mathfrak{V}(n)A_5(n+2) \\ &+ 25(n+1)(n+2) \prod_{k \in K, k \geq 6} (5n+k)\mathfrak{U}(n)A_5(n+1) \\ &- 250 \prod_{k \in K} (5n+k)\mathfrak{R}(n)A_5(n) = 0. \end{aligned} \quad (26.4)$$

The form (26.4), of order three and degree 15, is the one Kauers and Koutschan guessed [7]; source 75-03 proves the normalized form (26.2) and calls its coefficients “the coefficients in Conjecture 15” without printing the unnormalized operator. The two are equivalent: with $w(n) = (n!)^3((n+1)!)^2/(5n)!$ and

$$\mathfrak{g}(n) = \frac{125 \prod_{k \in K} (5n+k)}{(n+1)(n+2)},$$

the coefficient of $A_5(n+j)$ in (26.4) is $\mathfrak{g}(n) \mathfrak{p}_j(n) w(n+j)/w(n)$ for $j = 0, 1, 2, 3$, so (26.4) is $\mathfrak{g}(n)/w(n)$ times (26.2), and $\mathfrak{g}(n) \neq 0$. This was checked symbolically at the intake (2026-10-02); the OEIS operator also annihilates the terms given by (25.8) for $0 \leq n \leq 59$.

26.1 A rational residue representation

The substitution $t = (1+u)/(1-u)$ in (25.6) gives

$$\Lambda_n = \text{Res}_{t=1} \mathcal{W}_n(t) \mathcal{P}_n(t), \quad \mathcal{W}_n(t) = \frac{2^{2n+2} t^{3n+1}}{(t-1)^{n+1} (t+1)^{n+2}}, \quad (26.5)$$

so that

$$\frac{\mathcal{W}_{n+j}}{\mathcal{W}_n} = \left(\frac{4t^3}{t^2-1} \right)^j, \quad \mathsf{L}_n(t) := \frac{\mathcal{W}'_n}{\mathcal{W}_n} = \frac{3n+1}{t} - \frac{n+1}{t-1} - \frac{n+2}{t+1}. \quad (26.6)$$

The factor $(t+1)$ in the denominator of $\mathcal{W}_n$ is essential. The coefficients

$$\begin{aligned} \mathfrak{q}_0(n) &= -18(n+1)(3n+2)(3n+4)(3n+5)(3n+7)\mathfrak{R}(n), \\ \mathfrak{q}_1(n) &= 3(n+2)(3n+5)(3n+7)\mathfrak{U}(n), \\ \mathfrak{q}_2(n) &= -4(n+2)(n+3)^2\mathfrak{V}(n), \\ \mathfrak{q}_3(n) &= (n+2)(n+3)^2(n+4)^2\mathfrak{S}(n) \end{aligned} \quad (26.7)$$

satisfy, by cancellation of factorials,

$$\mathfrak{q}_j(n) = \mathfrak{c}(n) \mathfrak{p}_j(n) \frac{\mathfrak{h}_{n+j}}{\mathfrak{h}_n}, \quad \mathfrak{c}(n) = \frac{9(3n+2)(3n+4)(3n+5)(3n+7)}{n+2}, \quad (26.8)$$

with all denominators nonzero for $n \geq 0$.

26.2 The certificate identity

Define vectors of rational functions

$$\zeta_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \zeta_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \zeta_{j+1} = \frac{(2(n+j)+1)t \zeta_j - (n+j) \zeta_{j-1}}{n+j+1} \quad (j = 1, 2).$$

By the Legendre recurrence, $\mathcal{P}_{n+j} = \zeta_{j,0} \mathcal{P}_n + \zeta_{j,1} \mathcal{P}_{n+1}$. Set

$$\begin{pmatrix} \mathcal{R}_0 \\ \mathcal{R}_1 \end{pmatrix} = \sum_{j=0}^3 \mathfrak{q}_j(n) \left(\frac{4t^3}{t^2-1} \right)^j \zeta_j. \quad (26.9)$$

The rational functions $\mathcal{U}_n(t), \mathcal{V}_n(t)$ printed in Appendix C satisfy the two identities in $\mathbb{Q}(n, t)$

$$\mathcal{R}_0 = \mathcal{U}'_n + \mathsf{L}_n \mathcal{U}_n - \frac{(n+1)t}{t^2-1} \mathcal{U}_n - \frac{n+1}{t^2-1} \mathcal{V}_n, \quad (26.10)$$

$$\mathcal{R}_1 = \mathcal{V}'_n + \mathsf{L}_n \mathcal{V}_n + \frac{n+1}{t^2-1} \mathcal{U}_n + \frac{(n+1)t}{t^2-1} \mathcal{V}_n. \quad (26.11)$$

This is an explicit finite certificate: it is verified by substituting the printed polynomials and clearing denominators. `code/75-03-ballot-words-verify.py` does exactly this with exact rational coefficients (`sympy.cancel` of both residuals over $\mathbb{Q}(n, t)$), with no numerical sampling.

Proof of Theorem 26.1. The Legendre derivative identities

$$\mathcal{P}'_n = \frac{n+1}{t^2-1}(\mathcal{P}_{n+1} - t\mathcal{P}_n), \quad \mathcal{P}'_{n+1} = \frac{n+1}{t^2-1}(t\mathcal{P}_{n+1} - \mathcal{P}_n)$$

show that (26.10)–(26.11) are precisely the coefficients of $\mathcal{P}_n$ and $\mathcal{P}_{n+1}$ in the scalar identity

$$\sum_{j=0}^3 \mathfrak{q}_j(n) \mathcal{W}_{n+j}(t) \mathcal{P}_{n+j}(t) = \frac{d}{dt} \left[\mathcal{W}_n(t) (\mathcal{U}_n(t) \mathcal{P}_n(t) + \mathcal{V}_n(t) \mathcal{P}_{n+1}(t)) \right]. \quad (26.12)$$

For each $n \geq 0$ every function here is rational in t ; the only parameter denominator of the certificate is $n+2$. The residue at a finite point of the derivative of a rational function is zero. Taking the residue at $t=1$ and using (26.5) gives $\sum_{j=0}^3 \mathfrak{q}_j(n) \Lambda_{n+j} = 0$. Substituting (26.8) and (26.1) and cancelling $\mathfrak{c}(n)/\mathfrak{h}_n$ yields (26.2) for every $n \geq 0$. $\square$

26.3 Exact generation, and what is not asserted

The leading coefficients $\mathfrak{p}_3(n)$ and $\mathfrak{q}_3(n)$ are positive for $n \geq 0$, so the recurrence and $\Lambda_0, \Lambda_1, \Lambda_2$ determine the sequence. Working with the integers Λ_n avoids the rational normalization; (25.8) recovers $A_5(n)$. Each new term costs a bounded number of arithmetic operations; this is not a constant bit-cost claim, since the integers have linearly growing bit length.

The order is three, as conjectured. No claim of minimality over every coefficient field is made. The normalized recurrence has the exact solution $3n+2$:

$$\sum_{j=0}^3 \mathfrak{p}_j(n) (3(n+j) + 2) = 0, \quad (26.13)$$

which is why a leading asymptotic alone does not determine every formal coefficient: a second asymptotic datum is needed (Section 27).

This settles Kauers and Koutschan's Conjecture 15 [7], listed as open in Section 22. Their request for a provably correct recurrence for A215562 ($r=4$) remains open. The route is not the creative telescoping of the sextic proposed there (Remark 4.2): zero-step deletion and Lagrange inversion reduce the diagonal to the one-variable coefficient (25.7), and the telescoping is done on its Legendre residue representation.

27 The exact coefficients $\alpha_{5,2}, \dots, \alpha_{5,8}$

27.1 A second route to the expansion, a third to $\alpha_{5,1}$

The existence of a complete expansion (3.3) for $r=5$ with $\alpha_{5,j} \in \mathbb{Q}(\sqrt{5})$ is Theorem 3.1 with Theorem 3.3. Source 75-03 proves it again from (25.7), by a saddle point in two variables instead of three marker variables. Since $\text{CT}_v(1+v)^n(1+u/v)^n = \sum_k \binom{n}{k}^2 u^k$,

$$\Lambda_n = \text{CT}_{u,v}(1+u) \Pi(u, v)^n, \quad \Pi(u, v) = \frac{(1+u)^3(1+v)(1+u/v)}{u(1-u)^2}. \quad (27.1)$$

The logarithmic gradient of Π vanishes at $(u_0, v_0) = (\varphi^{-4}, \varphi^{-2})$, $\varphi = (1 + \sqrt{5})/2$: the v equation gives $v^2 = u$, and with $u = t^2$, $v = t$ the u equation becomes $(t+1)(t^2 - 3t + 1) = 0$, whose root in $(0, 1)$ is $t = \varphi^{-2} = (3 - \sqrt{5})/2$. There

$$\Pi(u_0, v_0) = 27, \quad 1 + u_0 = 3\varphi^{-2}, \quad H_{\Pi} = \begin{pmatrix} 14/15 & -1/5 \\ -1/5 & 2/5 \end{pmatrix}, \quad \det H_{\Pi} = \frac{1}{3},$$

H_{Π} being the Hessian of $\log \Pi$ in logarithmic coordinates. On the torus $|u| = u_0$, $|v| = v_0$ one has $|\Pi| \leq 27$ with equality only at (u_0, v_0) (the triangle inequalities for $1 + u$ and $1 + v$ already force both phases to vanish, and $|1 - u| \geq 1 - u_0$). The standard saddle argument (restriction to angular distance $n^{-2/5}$, a Gaussian majorant inside and an $O(27^n e^{-cn^{1/5}})$ bound outside) gives, at every finite order,

$$\Lambda_n = \frac{3\sqrt{3}\varphi^{-2}}{2\pi} \frac{27^n}{n} \left(1 + \frac{\eta_1^\Lambda}{n} + \frac{\eta_2^\Lambda}{n^2} + \cdots \right), \quad \eta_j^\Lambda \in \mathbb{Q}(\sqrt{5}), \quad (27.2)$$

and with Stirling's series in (25.8) the expansion (3.3) for $r = 5$, with the constant K_5 of (3.6).

For the first correction put, for $j \geq 2$, $\ell_j^+(q) = (q\partial_q)^j \log(1+q)$ and $\ell_j^-(q) = (q\partial_q)^j [-\log(1-q)]$, and

$$\mathcal{K}_j(Z_1, Z_2) = (3\ell_j^+(u_0) + 2\ell_j^-(u_0))Z_1^j + \ell_j^+(\varphi^{-2})(Z_2^j + (Z_1 - Z_2)^j),$$

where (Z_1, Z_2) is a centred Gaussian vector with covariance $H_{\Pi}^{-1} = \begin{pmatrix} 6/5 & 3/5 \\ 3/5 & 14/5 \end{pmatrix}$. Taylor expansion of amplitude and exponent gives, with $G_0 = 1 + u_0$,

$$\eta_1^\Lambda = \frac{1}{G_0} \mathbb{E} \left[-\frac{u_0 Z_1^2}{2} + \frac{u_0 Z_1 \mathcal{K}_3}{6} + \frac{G_0 \mathcal{K}_4}{24} - \frac{G_0 \mathcal{K}_3^2}{72} \right] = -\frac{71}{90} + \frac{13\sqrt{5}}{50}. \quad (27.3)$$

The factorial multiplier in (25.8) has relative first correction $-23/45$, so $\alpha_{5,1} = \eta_1^\Lambda - 23/45 = -13/10 + 13\sqrt{5}/50$: the value of (3.6), here by a third contraction after Sections 9 and 18. The checker evaluates (27.3) exactly.

27.2 The triangular algorithm

By (3.3) and Stirling's series, $\tilde{A}_5(n)$ has a complete expansion

$$\tilde{A}_5(n) \sim n \sum_{k \geq 0} \tilde{c}_k n^{-k}, \quad \tilde{c}_0 = \varphi^{-2}, \quad \tilde{c}_1 = 1 - \frac{4\sqrt{5}}{25}, \quad (27.4)$$

where $\tilde{c}_0$ comes from K_5 and $\tilde{c}_1 = \varphi^{-2}(\eta_1^\Lambda + 17/9)$ from $\alpha_{5,1}$, $17/9$ being the first Stirling correction of $\mathfrak{h}_n$. Put $\hat{\mathfrak{p}}_j(z) = z^5 \mathfrak{p}_j(1/z)$. Substituting the expansion in (26.2) with $n = 1/z$ gives, for every $k \geq 2$,

$$[z^{k+2}] \sum_{j=0}^3 \hat{\mathfrak{p}}_j(z) \sum_{\ell=0}^k \tilde{c}_\ell z^\ell (1+jz)^{1-\ell} = 0. \quad (27.5)$$

For a trial term n^β the first nonzero coefficient is $[z^2] \sum_j \hat{\mathfrak{p}}_j(z) (1+jz)^\beta = 1625\beta(\beta-1)$, so the two free powers are n and 1 , and the coefficient of $\tilde{c}_k$ in (27.5) is $1625k(k-1) \neq 0$. Given $\tilde{c}_0$ and $\tilde{c}_1$, (27.5) is therefore triangular; for example $\tilde{c}_2 = 2\sqrt{5}/25$, $\tilde{c}_3 = -74\sqrt{5}/3125$, $\tilde{c}_4 = 814\sqrt{5}/78125$. Finally, as formal series in z ,

$$\sum_{k \geq 0} \alpha_{5,k} z^k = \frac{\sum_{k \geq 0} \tilde{c}_k z^k}{\varphi^{-2}(1+z)^2} \exp \left(\sum_{j \geq 1} \frac{B_{2j}(5^{1-2j} - 5)}{2j(2j-1)} z^{2j-1} \right), \quad (27.6)$$

with Bernoulli numbers B_{2j} : the exponential is Stirling's correction for $(5n)!/(n!)^5$, and $(1+z)^{-2}$ that of $(n+1)^{-2}$, since $A_5(n) = \tilde{A}_5(n)(5n)!/((n!)^5(n+1)^2)$.

Proposition 27.1 (Exact coefficients for A215570 (source 75-03)). *For every k , the coefficient $\alpha_{5,k}$ of (3.3) is the k th coefficient computed by (27.5) and (27.6) from (27.4). The values for $k \leq 8$ are those of Table 6.*

Proof. The expansion exists (Theorem 3.1, or Section 27.1); inserting it into (26.2), which holds for every n , gives a formal identity whose coefficients are (27.5). These determine $\tilde{c}_k$ for $k \geq 2$ uniquely from $\tilde{c}_0, \tilde{c}_1$, which are fixed by K_5 and $\alpha_{5,1}$ (Theorem 3.3). Stirling's series converts $\tilde{c}_k$ into $\alpha_{5,k}$ by (27.6). The table entries are the exact values recorded by `code/75-03-ballot-words-verify.py` in `data/75-03-ballot-words-verification_results.json`, regenerated identically at the intake. $\square$

Table 6: The exact coefficients of (3.3) for A215570 and of (27.4) (source 75-03). The values for $k = 0, 1$ were known (Theorem 3.3); those for $2 \leq k \leq 8$ are new here. Source 75-03 prints $k \leq 4$ in its text and $k \leq 8$ in its data file. Decimals are rounded.

k	$\alpha_{5,k}$	decimal	$\tilde{c}_k$
0	1	1	$\frac{3}{2} - \frac{\sqrt{5}}{2}$
1	$-\frac{13}{10} + \frac{13\sqrt{5}}{50}$	-0.71862232585	$1 - \frac{4\sqrt{5}}{25}$
2	$\frac{36}{25} - \frac{63\sqrt{5}}{125}$	0.31302173934	$\frac{2\sqrt{5}}{25}$
3	$-\frac{407}{250} + \frac{4283\sqrt{5}}{6250}$	-0.09566733638	$-\frac{74\sqrt{5}}{3125}$
4	$\frac{29387}{15625} - \frac{64554\sqrt{5}}{78125}$	0.03312470759	$\frac{814\sqrt{5}}{78125}$
5	$-\frac{1682397}{781250} + \frac{3745709\sqrt{5}}{3906250}$	-0.00929921333	$\frac{1918\sqrt{5}}{1953125}$
6	$\frac{23801122}{9765625} - \frac{52959909\sqrt{5}}{48828125}$	0.01195322150	$\frac{248974\sqrt{5}}{48828125}$
7	$-\frac{52826879}{19531250} + \frac{118935383\sqrt{5}}{97656250}$	0.01856723270	$\frac{1667798\sqrt{5}}{244140625}$
8	$\frac{735722961}{244140625} - \frac{1619559792\sqrt{5}}{1220703125}$	0.04683285829	$\frac{21982414\sqrt{5}}{1220703125}$

The value $\alpha_{5,2} = 0.31302\dots$ is the limit of the last column of the diagnostic table of Section 13, which source 61 did not claim as a formula: at $n = 50$ that column is 0.3111216, and $\alpha_{5,2} + \alpha_{5,3}/50 + \alpha_{5,4}/50^2 = 0.3111216$. (2026-10-05 note: source 101-86 computes $\alpha_{5,2}$ again without the recurrence, from the kernel and Gaussian moments alone, Theorem 37.2; the two derivations agree.) Table 7 shows source 75-03's numerical check of the truncations.

Table 7: Signed relative error $(K_5 5^{5n} n^{-3} \sum_{k \leq M} \alpha_{5,k} n^{-k})/A_5(n) - 1$ (source 75-03; rounded from its recorded results file, Section 29). Numerical illustrations, not interval-certified inequalities; the analytic argument, not this table, proves the expansion.

n	$M = 0$	$M = 1$	$M = 2$	$M = 4$	$M = 8$
20	$3.6442 \cdot 10^{-2}$	$-7.9889 \cdot 10^{-4}$	$1.2182 \cdot 10^{-5}$	$2.8011 \cdot 10^{-9}$	$-3.2337 \cdot 10^{-13}$
50	$1.4454 \cdot 10^{-2}$	$-1.2625 \cdot 10^{-4}$	$7.7105 \cdot 10^{-7}$	$2.9386 \cdot 10^{-11}$	$-7.3726 \cdot 10^{-17}$
100	$7.2066 \cdot 10^{-3}$	$-3.1432 \cdot 10^{-5}$	$9.6024 \cdot 10^{-8}$	$9.2439 \cdot 10^{-13}$	$-1.3794 \cdot 10^{-19}$
250	$2.8777 \cdot 10^{-3}$	$-5.0166 \cdot 10^{-6}$	$6.1318 \cdot 10^{-9}$	$9.5004 \cdot 10^{-15}$	$-3.5266 \cdot 10^{-23}$
1000	$7.1883 \cdot 10^{-4}$	$-3.1315 \cdot 10^{-7}$	$9.5703 \cdot 10^{-11}$	$9.2939 \cdot 10^{-18}$	$-1.3288 \cdot 10^{-28}$

Remark 27.2 (Scope (source 75-03)). Proposition 27.1 concerns the dominant exponential sector, as in the remark closing Section 6; it determines no exponentially small sector and no Stokes data. The leading characteristic polynomial of the normalized recurrence (26.2) is $65(\lambda - 1)^2(27\lambda - 2)$, here with λ a characteristic root, not $r \log r$. The double root 1

carries the two powers n and 1 above; the third root $2/27$ indicates a possible formal mode smaller by the factor $(2/27)^n$, whose contribution to the sequence is not determined here. The coefficients grow from $k = 6$ on, as expected of an asymptotic series; nothing is claimed about its summability.

28 Inversion, rounding and repository machinery

Source 75-03's inversion of A215570 is the instance $r = 5$ of Section 11 and of the rounding bound (20.4), and through them of the canonical transseries volume (Section 1.4): its Lambert anchor is (11.2), the large-solution case $b < 0$ of `p0:thm:lambert-core` (display `p0:eq:core-lambert-minus`, with $a = \lambda = 5 \log 5$ and $b = -\beta = -3$); its corrections are the first two coefficients of the Lambert-centred reversion `p0:thm:lambert-centered`, which is the explicit inverse `t2:thm:balanced-inverse` (display `t2:eq:balanced-first`) of the companion volume `Analysis/Transseries/docs/series-and-transseries/Combinatorial_Transseries_Inverses/Combinatorial_Transseries_Inverses.tex`; and its rounded threshold is parts (2)–(3) of `p0:thm:staircase` (Remark 20.1). Source 75-03 cites none of these, and no novelty is claimed for them. What is new is one value: with $\alpha_{5,2}$ known, (11.4) for $r = 5$ is explicit to second order,

$$n = x_0 + \frac{13(5 - \sqrt{5})}{50\lambda} \frac{1}{x_0} + \left(\frac{83\sqrt{5} - 213}{500\lambda} + \frac{39(5 - \sqrt{5})}{50\lambda^2} \right) \frac{1}{x_0^2} + O(x_0^{-3}), \quad (28.1)$$

for $Y = A_5(n)$, $x_0 = x_0(Y)$, $\lambda = 5 \log 5$; here $b_2 = \alpha_{5,2} - \frac{1}{2}\alpha_{5,1}^2 = (213 - 83\sqrt{5})/500$, and the two coefficients are $\mu_1 = 0.0893010312$ and $\mu_2 = 0.0264800992$. (2026-10-05 note: source 101-86 derives the same second-order inverse from its own, recurrence-free $\alpha_{5,2}$, Section 37.5.) The second term accounts for the residual column of the inverse diagnostic table in Section 13: μ_2/x_0^2 is $4.150 \cdot 10^{-4}$, $1.035 \cdot 10^{-4}$ and $1.059 \cdot 10^{-5}$ at $n = 8, 16, 50$, against the printed $-4.164 \cdot 10^{-4}$, $-1.038 \cdot 10^{-4}$, $-1.060 \cdot 10^{-5}$ (the column is $-\mu_2/x_0^2 + O(x_0^{-3})$). Source 75-03 states the rounded threshold

$$\lceil x_*(Y) - Cx_0^{-3} \rceil \leq N_5(Y) \leq \lceil x_*(Y) + Cx_0^{-3} \rceil,$$

with x_* the truncation of (28.1) after x_0^{-2} ; this is (20.4) with $M = 2$, since x_0^{-3} and $(\log Y)^{-3}$ are comparable. It also carries the reversion of the anchor one order beyond (20.3): with $D_Y = \log(Y/K_5) - 3 \log \lambda$, the number $u = \lambda x_0$ solves $u - 3 \log u = D_Y$, and

$$\lambda x_0 = D_Y + 3 \log D_Y + \frac{9 \log D_Y}{D_Y} + \frac{27 \log D_Y - \frac{27}{2}(\log D_Y)^2}{D_Y^2} + O\left(\frac{(\log D_Y)^3}{D_Y^3}\right), \quad (28.2)$$

the nested-logarithm expansion that follows `t2:thm:balanced-inverse` in the companion volume. Source 75-03 adds that for other rows the general theorem “currently supplies only the coarser $O(x^{-1/5})$ uncertainty”; in this report Theorem 3.1 gives every row a complete expansion, so Section 11 applies to every row at every order.

29 Source 75-03's computations

The shipped `code/75-03-ballot-words-verify.py`, with its integer certificate arrays in `code/75-03-ballot-words-certificate_data.py`, performs these exact checks:

- (a) the generic quartic pair-product resolvent of Remark 25.1 and its specialization to the sextic, as polynomial identities;

- (b) both certificate identities (26.10)–(26.11) identically in $\mathbb{Q}(n, t)$, the four relations (26.8), and the solution (26.13);
- (c) the Hessian H_{Π} and the Gaussian expectation (27.3) in $\mathbb{Q}(\sqrt{5})$;
- (d) agreement of the recurrence-generated Λ_n with (25.7) for $0 \leq n \leq 103$;
- (e) an independent content-state dynamic program, the recursion of Section 13 (no row recurrence is used), for $r = 2, 3, 4$ ($n \leq 10$), $r = 5$ ($n \leq 8$), $r = 6$ ($n \leq 5$) and $r = 7$ ($n \leq 4$), and the thirteen OEIS data terms of A215570.

It also generates Λ_n for $n \leq 1000$ by the recurrence, writes Λ_n and $A_5(n)$ for $n \leq 103$ to `data/75-03-ballot-words-a215570_terms.csv`, and records $\alpha_{5,0}, \dots, \alpha_{5,8}$, $\tilde{c}_0, \dots, \tilde{c}_8$, the constants of Table 3 for $r \leq 8$ (to 35 digits), the dynamic-programming rows and Table 7 in `data/75-03-ballot-words-verification_results.json`; its standard output is `data/75-03-ballot-words-verification_log.txt`. The recorded run used Python 3.13.5, SymPy 1.14.0 and mpmath 1.3.0 and took 3.6 s. The 53 dynamic-programming terms equal source 61’s enumerations (`data/61-balanced-exact_rows.json`). The intake reran the program on a copy (Python 3.14.4): all six checks passed, the CSV was regenerated byte for byte, and the JSON differs only in the recorded Python version and running time (and in its line endings: on Windows the program writes it with CRLF, while the shipped file has LF).

Source 75-03 marks the boundaries of these checks. The certificate check is finite symbolic algebra: it does not replace the derivation from words to (25.7), which is the proof above, and “finite agreement alone would not prove the conjecture”. Numerical errors are high-precision illustrations, not interval enclosures. “No Lean files are supplied as if they had been compiled.”

Part V

Equal-content ballot words: the first correction for A215562

Part V was added on October 2, 2026, in batch 77 of ProveIt’s incoming-reports intake, from a fourth manuscript on the same problem (source 77-44), after Part IV. Source 77-44 proves the leading constants of A215562 and A215570, every fixed order of their expansions, the fixed-row theorem for every alphabet and D-finiteness: a *fourth* independent proof of Theorems 3.1 and 3.2 (its D-finiteness part) and of the constructions of Sections 4–7, with identical constants and the same sextic. Those results are not printed again; Table 8 lists them. What it adds is one new value, the first correction $\alpha_{4,1}$ of A215562 (Theorem 31.1), which meets the target stated in Part III (Section 22); an operator-norm proof of the global-arc bound (Section 32), a second route to Section 6.4; the explicit Hessian for the fourth row; a nested-logarithm inverse for every row (Remark 33.1); and two literature pointers (Section 33). The rest of the report is unchanged apart from dated notes that point here, rows added to Tables 1 and 2, and the package description of Appendix B; no existing label or number was changed.

30 Provenance, overlap and conventions of Part V

30.1 A fourth manuscript

Source 77-44 is batch-77 manuscript 44, *Equal-content ballot words: leading constants and fixed-order expansions for OEIS A215562 and A215570*, dated October 1, 2026, author line “Research note with independently reviewed proofs” (a 594-line source with 18 labels and an 8-page PDF). Archive `oeis-ballot-asymptotics-20261001.zip`, arrival commit `096ee7b87`, placed in commit `aa7345800`. It names no ProveIt commit and records no repository search. Its programs and receipts ship with the prefix `77-44-bal-` (Section 34).

Source 77-44 was written without knowledge of this report: its receipts record the UTC date October 1, 2026, and its archive members are stamped 10:40–10:46 (no time zone), in any time zone before this report’s placement (`6e193dd4f`, October 2, 02:16 UTC). Its Theorem 1, Corollary 2, Lemmas 3 and 4, Proposition 5 and Theorem 6 re-prove Theorems 3.1 and 3.2 and Sections 4–7 of this report and are listed in Table 8, not reprinted. It presents the row-wise asymptotic recorded conjecturally in A215561 as a consequence of its own argument; in this report its proof is the fourth (after sources 61, 42 and 75-03). Its status statement is kept: “The proofs have been independently reviewed and the supporting computations reproduced. These are ordinary mathematical proofs, not machine-checked theorems or claims of publication priority. A complete exponentially improved transseries is not claimed.” The review it refers to is its own pre-delivery review (an approval receipt), not refereeing; the delivery says nothing about AI assistance. Its constants “exist for every fixed J ; this draft does not give numerical thresholds or optimized numerical remainder constants”. No Lean or Rocq development checks any statement of this Part.

Source 77-44 calls the Kauers–Koutschan recurrence for A215570 “guessed” and unproved, and lists “proving the particular guessed recurrence” among its future questions. (2026-10-02 note: the recurrence is proved in Part IV, Theorem 26.1; the A215562 recurrence remains open, Section 22.)

30.2 What the fourth manuscript re-proves

Table 8: Results of source 77-44 that Parts I, II and IV already prove. Statement numbers are source 77-44’s own (it numbers theorems, lemmas and propositions consecutively).

source 77-44	printed here	comparison
leading constants of A215562 and A215570, every fixed order (Thm. 1)	Theorem 3.1	same; its E_q is κ_r and its leading constant is K_r of (3.4)
every fixed row; $\det H = \Delta^2/(q^q \sigma^2)$; D-finiteness (Thm. 6)	Theorems 3.1, 3.2; Lemma 6.1, (6.6)	same (fourth proof); no transcendence statement
small-root formula, bridge identity $\mathcal{B} = 1 + t\partial_t \log \mathcal{E}$	Lemmas 4.1, 5.1	same; classical [8]
degree-six equation of the fully marked five-step series	(4.5)	the same polynomial term by term, under $(A, B, C, D, F_2, F) = (a, b, c, d, e, X)$; same derivation by small- and large-root quadratics (a fourth derivation, beside Remarks 4.2 and 25.1)

source 77-44	printed here	comparison
D-finiteness of A215562, A215570 (Cor. 2)	Theorem 3.2	same; it cites a different Lipshitz paper [10]
uniform square-root expansions, periods $p_4 = 2$, $p_5 = 1$ (Lemma 3); mark transfer (Prop. 5); lattice normalization check	Section 6, (6.10)	same; $p_q = \nu$
global mark arcs (Lemma 4)	Section 6, “Why other phases do not contribute”	same statement, different proof : Section 32
$E_*(1) = \frac{5}{2}(3 - \sqrt{5})$, 4ϱ	Section 7	same; its reciprocal factorization is source 42’s elimination
first-correction recipe $c_{q,1} = (D_{\text{uni}} + D_{\text{sad}})/q + (q^{-1} - q)/12$	(9.1)–(9.4)	the same formula: $D_{\text{sad}} = \mathcal{S}$, and $D_{\text{uni}} = -\frac{3}{2}d_2 + \frac{1}{2}\alpha^2 = \mathcal{B}$ because $d_2 = D = L_2 + c^2/2$ and $\alpha = c$
$c_{5,1} = 13(\sqrt{5} - 5)/50$, with D_{uni} , D_{sad}	(3.6), Section 9	same values; a fourth computation (after Sections 9, 18 and (27.3))
the higher-order algorithm	Section 6.6, Appendix A, Section 17	same
inverse models, threshold localization	Section 11, (20.4)	same; its general nested-logarithm form is Remark 33.1
exact diagonals through $n = 30$	Sections 13, 29	equal on every common index

30.3 Conventions of Part V

Part V uses the merged notation of Section 1.5; Table 2 lists every renamed symbol of source 77-44. Its alphabet size q is r here, and its normalized coefficients $c_{q,j}$ are the $\alpha_{r,j}$ of (3.3). One normalization differs: its step polynomial $P_q(u) = \sum_i u^{s_i}$ is r times the P_r of (3.1), so its dominant singularity is at $t = 1/q$, not $t = 1$. Three readings are tempting and false:

- $\varrho = \varphi - \sqrt{\varphi} = \kappa_4/4$ (source 77-44’s r) is not the alphabet size r ; it is the product q of the two interior roots in Section 7, the root in $(0, 1)$ of $\varrho^4 - 2\varrho^3 - 2\varrho^2 - 2\varrho + 1$.
- $c_{q,j}$ is not the OEIS-style constant $c(r)$ of (3.4), and source 77-44’s $c = -\min S_q$ is the depth h .
- Source 77-44’s $\mathcal{B}$ is the bridge generating function $B(t; \mathbf{x})$, not the length correction $\mathcal{B}$ of (9.2), which it calls D_{uni} .

31 The first correction for A215562

Theorem 31.1 (First correction of the fourth row (source 77-44)). *Let $\varphi = (1 + \sqrt{5})/2$ and $\varrho = \varphi - \sqrt{\varphi}$. Then*

$$A_4(n) = \frac{\varrho 256^n}{\sqrt{2} \pi^{3/2} n^{5/2}} \left(1 + \frac{\alpha_{4,1}}{n} + O(n^{-2}) \right), \quad (31.1)$$

$$\alpha_{4,1} = -\frac{37\varrho^3}{200} + \frac{11\varrho^2}{40} + \frac{13\varrho}{40} - \frac{259}{400} = -0.509784684867675\dots \quad (31.2)$$

The leading factor is $K_4 4^{4n} n^{-5/2}$ of (3.3), with $\kappa_4 = 4\varrho$; it is the formula credited to Kotesovec in the OEIS [2]. Every $\alpha_{4,j}$ lies in $\mathbb{Q}(\varrho)$, a quartic field, by the argument that gives $\alpha_{5,j} \in \mathbb{Q}(\sqrt{5})$ in Section 9; source 77-44 states that all its normalized coefficients are effectively computable real algebraic numbers.

Proof (source 77-44). Apply (9.4) with $r = 4$: $\alpha_{4,1} = (\mathcal{B} + \mathcal{S})/4 + (1/4 - 4)/12$. In source 77-44's form, the length part is

$$\mathcal{B} = -\frac{3}{2}d_2 + \frac{1}{2}\alpha^2, \quad \alpha^2 = \frac{2q}{P_q''(1)}, \quad \beta = -\frac{P_q'''(1)\alpha^2}{6P_q''(1)}, \quad d_2 = 1 + \beta + q \sum_{\substack{P_q(r_j)=q \\ |r_j|<1}} \frac{1}{r_j P_q'(r_j)},$$

where $u_1 = 1 - \alpha\eta^{1/2} + \beta\eta + \dots$ is the coalescing small root at unit marks, $\eta = 1 - t/\rho$, and d_2 is the coefficient of η in $\mathcal{E}/E_*$; for $q = 4$, $P_4(u) = u^{-3} + u^{-1} + u + u^3$, $P_4'(1) = 20$ and $\alpha^2 = 2/5 = 2/V$. The saddle part $\mathcal{S}$ is (9.1), written by source 77-44 as

$$\mathcal{S} = -\frac{1}{2} \sum_{ij} G_{ij}(g_{ij} + g_i g_j + 2(\partial_i \log b)g_j) + \frac{1}{2} \sum_{ijkl} g_i \kappa_{jkl} G_{ij} G_{kl},$$

with $G = H^{-1}$, $g = \log E_*(e^z)$ and $b = V(e^z)^{-1/2}$ in the logarithmic marks z of the first two letters. For $r = 4$ the marked steps are $-3, -1$, the unmarked ones $1, 3$, and

$$H_4 = \begin{pmatrix} 3/40 & -1/10 \\ -1/10 & 7/40 \end{pmatrix}, \quad \det H_4 = \frac{1}{320}. \quad (31.3)$$

Direct implicit differentiation of the kernel roots, in $\mathbb{Q}(\varrho)$ reduced modulo $\varrho^4 - 2\varrho^3 - 2\varrho^2 - 2\varrho + 1$, gives

$$\mathcal{B} = -\frac{6\varrho^3}{25} + \frac{3\varrho^2}{10} + \frac{3\varrho}{5} - \frac{47}{50}, \quad \mathcal{S} = -\frac{\varrho^3}{2} + \frac{4\varrho^2}{5} + \frac{7\varrho}{10} - \frac{2}{5}, \quad (31.4)$$

and (9.4) yields (31.2). $\square$

Source 77-44's exact script `code/77-44-bal-ballot_four_correction.py` implements this computation; its numerical-derivative check (`code/77-44-bal-verify_ballot_numerical_derivatives.py`) differentiates numerical implicit root solves directly, without the symbolic derivative formulas, and reproduces $\alpha_{4,1}$ and $\alpha_{5,1}$ to at least 54 decimal digits.

Remark 31.2 (Checks made at the merge). The intake checked that (31.4) with (9.4) gives exactly (31.2) ($(\mathcal{B} + \mathcal{S})/4 = -\frac{37}{200}\varrho^3 + \frac{11}{40}\varrho^2 + \frac{13}{40}\varrho - \frac{67}{200}$ and $(1/4 - 4)/12 = -\frac{5}{16}$), and that $\mathcal{B} \approx -0.706416072$ and $\mathcal{S} \approx -0.082722667$. Richardson extrapolation of $n(A_4(n)/(K_4 256^n n^{-5/2}) - 1)$ on the exact terms $n \leq 30$, which equal the enumerations of Section 13 on their common range, gives -0.5097844658 , -0.5097846437 and -0.5097846764 at orders 4, 6 and 8: agreement with (31.2) to eight digits. The same procedure reproduces $\alpha_{5,1}$ to thirteen digits. These are diagnostics, not part of the proof. $\det H_4 = 1/320$ and the five-letter matrix H_5 of source 77-44, with $\det H_5 = 1/6250$, are those of (6.6) and Section 9.

With $\alpha_{4,1}$ known, (11.4) is explicit to first order for the fourth row (this display was added at the merge):

$$n = x_0 + \frac{|\alpha_{4,1}|}{4 \log 4} \frac{1}{x_0} + O(x_0^{-2}), \quad \frac{|\alpha_{4,1}|}{4 \log 4} = 0.0919329795\dots, \quad Y = A_4(n), \quad (31.5)$$

with $x_0 = x_0(Y)$ the Lambert core (11.2) for $(K, \lambda, \beta) = (K_4, 4 \log 4, 5/2)$.

32 A second route to the global-arc bound

Section 6 (“Why other phases do not contribute”) shows that the marker torus contributes only near $z = \mathbf{1}$ by continuing the small-root product to a slightly larger disk away from that point. Source 77-44 proves the needed bound directly on coefficients.

Lemma 32.1 (Global mark arcs (source 77-44)). *Let $E_N(\mathbf{w})$ and $B_N(\mathbf{w})$ be the mark-weighted counts of nonnegative excursions and zero-height bridges of length N , with the first $r - 2$ step types marked by $\mathbf{w}$ and the unnormalized step polynomial $P(u; \mathbf{w})$. For $\mathbf{w}$ on the unit torus,*

$$|E_N(\mathbf{w})| \leq \left(\max_{|u|=1} |P(u; \mathbf{w})| \right)^N, \quad |B_N(\mathbf{w})| \leq \left(\max_{|u|=1} |P(u; \mathbf{w})| \right)^N.$$

On any closed subset of the torus excluding $\mathbf{1}$ the maximum is at most $r - \epsilon$ for some $\epsilon > 0$.

Proof (source 77-44). The bridge inequality follows from its Fourier constant-term integral. For excursions, let $L_{\mathbf{w}}$ be the bilateral Laurent operator on $\ell^2(\mathbb{Z})$ with the marked steps as its coefficients. Fourier transformation gives $\|L_{\mathbf{w}}\| = \max_{|u|=1} |P(u; \mathbf{w})|$. If Π projects onto heights $0, 1, \dots$, then $T = \Pi L_{\mathbf{w}} \Pi$ is the killed transfer operator and $E_N = \langle e_0, T^N e_0 \rangle$. Thus $|E_N| \leq \|T\|^N \leq \|L_{\mathbf{w}}\|^N$.

Equality in the triangle bound $|P(u; \mathbf{w})| \leq r$ requires all its unit-modulus summands to have the same phase. The two unmarked steps differ by δ (source 77-44's Δ), forcing $u^\delta = 1$: for odd r , $u = 1$; for even r all exponents are odd, so $u = \pm 1$ gives the same conclusion. Then every mark equals 1. Compactness gives a strict uniform gap away from that mark point. $\square$

Source 77-44 states the lemma for $r = 4, 5$ and extends it to every fixed row in its Theorem 6. With the uniform square-root expansions near $\mathbf{1}$ (Section 6), the lemma makes the mark integral outside a fixed neighbourhood of $\mathbf{1}$ exponentially smaller than the main term, which is what Section 6.4 provides by continuation. The operator bound needs no continuation in t and no analysis of colliding roots.

33 Literature pointers and further remarks

Source 77-44 adds two references to those of the other sources. Denisov and Wachtel [11] give a much more general local-limit framework for random walks in cones, compatible with the exponent; “its existence theorem alone does not replace the explicit constant calculation or all-order argument”. Banderier, Krattenthaler, Krinik, Kruchinin, Kruchinin, Nguyen and Wallner [12], Section 3.3 and Theorem 3.8, already give univariate basketball-walk asymptotics and explain the computation of all higher Puiseux orders; their statements fix length and endpoints, not equal multiplicities of every step, so the mark extraction and its global-arc control are the additional ingredients. In the five-letter case deletion and reinsertion of the zero steps reduces the four nonzero steps to basketball steps; this is the reduction (25.3) of Part IV. “These comparisons do not constitute a comprehensive publication-priority claim.” The OEIS leading constants “are prior statements, not new conjectures”.

Remark 33.1 (A nested-logarithm inverse for every row (source 77-44)). Write the expansion as $A_r(n) = K_r e^{\lambda n} n^{-\beta} (1 + \alpha_{r,1}/n + \dots)$ with $\lambda = r \log r$, $\beta = (r + 1)/2$, and fix J . Source 77-44 defines the *chosen analytic model* $F_J(x) = K_r e^{\lambda x} x^{-\beta} (1 + \sum_{j \leq J} \alpha_{r,j} x^{-j})$ for $x > X_J$, positive and strictly increasing for large X_J , so that $F_J^{-1}(A_r(n)) = n + O_J(n^{-J-1})$. With $L = \log Y$ and $B = \beta \log \lambda + \log K_r$ its compatible formal inverse begins

$$x = \frac{1}{\lambda} \left[L + \beta \log L - B + \frac{\beta^2 \log L - \beta B - \lambda \alpha_{r,1}}{L} + O\left(\frac{(\log L)^2}{L^2}\right) \right]. \quad (33.1)$$

This extends (20.3) by one order, for every row; for $r = 5$ it is consistent with (28.2) combined with (11.5). The discrete threshold is localized by rounding an interval of width $O_J(x^{-J-1})$ around $F_J^{-1}(Y)$; exact rounding requires separation from an integer at that accuracy, and “no

uniform exact-rounding claim is made”. As for the other sources, the core is the large-solution case $b < 0$ of `p0:thm:lambert-core` and the corrections are `p0:thm:perturbed-inversion` of the transseries volume (Section 1.4); the threshold step is `p0:thm:staircase`. No novelty is claimed.

The intake checked (33.1) by substituting it into $\lambda x - \beta \log x + \log K_r + \alpha_{r,1}/x = L$: the constant and the L^{-1} terms cancel.

34 Source 77-44’s computations and questions

Source 77-44 ships an exact checker, five derivation scripts and their receipts (Appendix B). `code/77-44-bal-verify_ballot_kernels.py` enumerates all count vectors in the box $0 \leq b_i \leq 3$, checks the bridge-excursion coefficient identity for both step sets and the degree-six equation for the five-step set with no nonzero residuals, and outputs the diagonals (1, 7, 403, 40350) and (1, 35, 18720, 19369350), which agree with the entries. `code/77-44-bal-verify_ballot_large_dp.py` recomputes the two diagonals through $n = 30$ by a rolling-array dynamic program; for five letters it removes and reinserts the zero steps. The first-correction scripts use exact symbolic arithmetic, and `code/77-44-bal-check.py` checks the correction formula identities, the generic degree-six identity, the covariance determinants for $q = 2, \dots, 16$ and 134 finite bridge identities. “These numerical checks, and numerical values from the OEIS, are diagnostics, not part of the proof.”

At placement the intake reran three of the scripts on copies: the kernel check, the dynamic program and the numerical-derivative check reproduced their receipts (modulo line endings), and the two first-correction scripts printed the values of (31.4) and of $\alpha_{5,1}$. The checker `code/77-44-bal-check.py` and the driver `code/77-44-bal-verify.py` did not finish within three minutes on a loaded machine and were not rerun; the delivered receipts record their passes. The diagonals through $n = 30$ equal `data/61-balanced-exact_rows.json` and `data/75-03-ballot-words-a215570_terms.csv` on every common index.

Source 77-44’s questions are those of Section 22: numerical effective remainder constants (“Determine sharp asymptotic error constants”); proving the particular guessed recurrence (resolved for A215570 by Part IV; open for A215562); “a genuinely exponentially improved transseries or a single selected exact analytic interpolation” (“Determine the exponentially small sectors”); and an alphabet size growing with n (“Let the alphabet grow”). “None of these is silently inferred” from its theorems. With $\alpha_{4,1}$, the target of Section 22 (“Automate all-order coefficient extraction”) is met for $r = 4$ at first order; $\alpha_{4,2}$, $\alpha_{4,3}$ and the rows $r = 6, 7, 8$ remain open.

Part VI

Direct excursion proofs: every fixed alphabet, and the second correction of A215570 without the recurrence

Part VI was added on October 5, 2026, in batch 101 of ProveIt’s incoming-reports intake, from two reports of one session bundle (bundle Reports 90 and 86, here sources 101-90 and 101-86), after Part V. Source 101-90 proves Theorem 3.1 and the D-finiteness part of Theorem 3.2

for every fixed alphabet, in marker coordinates that leave the two *extreme* steps unmarked; source 101-86 proves them, and Theorem 3.3, for the five-letter row after deleting the zero steps. They give a *fifth* and a *sixth* proof of results this report already has (after sources 61, 42, 75-03 and 77-44), and those results are not printed again: Table 9 lists them and Table 10 maps the second routes. What the two reports add is printed in full: source 101-86's computation of $\alpha_{5,2}$ *without* the recurrence of Part IV, with every intermediate value (Theorem 37.2); the leading constant of the zero-free count $A_5^o(n)$ (Proposition 37.1); the inequality $A_5(n+1) \geq 35A_5(n)$ for every n (Proposition 37.3, generalized at this write); source 101-90's zero-step identity for every odd alphabet (Proposition 36.1), its closed determinant (Proposition 36.2) and its account of the periodic torus points (Lemma 36.3). The rest of the report is unchanged apart from the title page and abstract, paragraphs added to Sections 1.1 and 1.3 and to the conclusion (Section 23), dated notes that point here, rows added to Tables 1 and 2, the merged research questions of Part III and the package description of Appendix B; no existing label or number was changed. Statements and proofs supplied by this write rather than by a source are marked [write]. An independent check of the write, made afterwards, is recorded at the end of Section 38.2.

35 Provenance, overlap and conventions of Part VI

35.1 Two bundle reports

The two reports arrived in commit 60f54ea06 as archives of a 177-archive session bundle and were placed in commit f7c612c72 (batch 101).

- **Source 101-90** (bundle Report 90): *Balanced ballot words over every fixed alphabet: exact constants, all algebraic orders, and inversion*, archive A215561_Fixed_Alphabet_Asymptotics.zip (248,562 bytes; a 474-line source with 24 labels and an 8-page US-letter PDF). It is the base of this Part, because it treats every fixed r .
- **Source 101-86** (bundle Report 86): *Balanced five-letter ballot words: an all-orders asymptotic and its inverse*, archive A215570_Balanced_Ballot_Asymptotics.zip (246,773 bytes; a 433-line source with 21 labels and a 7-page US-letter PDF). It is the case $r = 5$ in other coordinates, with one computation the others lack.

Both are dated October 1, 2026, with the author line “Research note”. Neither names a ProveIt commit or path, and neither records a repository search. Their PDFs were created at 09:37 (101-86) and 10:16 PDT (101-90), about nine hours before this report's first placement (6e193dd4f, 19:16 PDT), so neither could have seen any part of it. They come from one session (an identical build script; 4.7% and 4.2% of their 8-word shingles are common, mostly boilerplate); against this report only 1.3% (101-86) and 1.5% (101-90) of their shingles occur, so there is no textual copying. They describe their novelty against the OEIS and the literature only, in the same words: a bounded primary-source check “is a source-relative statement, not an exhaustive priority claim” (101-86) and “is not an exhaustive worldwide priority claim” (101-90).

Neither report says anything about AI assistance. Their receipts record a pre-delivery approval of the producing pipeline (in source 101-86's words, “Root and an independent research reviewer approved the complete proof and final transcription”) and an “independent” review file each; these are self-reported reviews, not refereeing (Section 38.2). Both are unrefereed, and no Lean or Rocq development checks any statement of this Part. Their programs and data ship with the prefixes 101-90-alphabet- and 101-86-five- (Section 38.2, Appendix B).

The two reports keep these limitations, which this Part keeps with them. Source 101-90: the alphabet is fixed and “uniformity when r grows is a separate problem”; D-finiteness is an existence statement, and “no particular guessed recurrence” or minimal order is identified; no smaller exponential sectors and no explicit numerical remainder constants; root products for larger r are “orientation only”; its code checks examples and is “not a general-purpose implementation”; and “no assertion that their smallest field is $\mathbb{Q}(\gamma_r)$ is needed”. Source 101-86: its expansion is a Poincaré expansion of the dominant exponential sector, “not a complete description of every smaller exponential contribution”; it does not use, and does not prove, the order-three recurrence; its inverse keeps its integer-rounding qualification, with “no numerical error constant or unconditional exact-ceiling rule” and no exponentially improved error; a single ceiling is certified only away from shrinking neighbourhoods of integers; its finite checks “supplement the analytic proof”; and the kernel method is classical [8].

35.2 What the two reports re-prove

Table 9 lists every result of the two reports that Parts I–V already contain, and Table 10 maps their proofs of Theorems 3.1–3.3 beside the earlier ones. They are second routes: the theorems were proved, four times, before the two reports were written, and no novelty is claimed for them here.

Table 9: Results of sources 101-90 and 101-86 that Parts I–V already prove. Equation names are the sources’ own labels.

source	printed here	comparison
101-90, Theorem 1: every fixed row, all orders, K_r , real algebraic $\alpha_{r,j}$; D-finiteness	Theorems 3.1, 3.2	same statement (its γ_r is κ_r); a fifth proof, Sections 36.2–36.4; no transcendence statement
101-90, eq:oeisconstant	$c(r)$ in (3.4)	same; its κ_r is $c(r)$
101-90, eq:rootconstant, eq:Q	Proposition 5.2	same; its Q_r is $-Q_r$; inside roots counted by reciprocity, as source 42 does; equal to Table 3 for $2 \leq r \leq 12$ to 15 digits (checked at the intake)
101-90, γ_4, γ_5	Section 7	same computation as sources 61 and 42 ($v = u + u^{-1}$, $v^3 - 2v = 4$, $p + p^{-1} = 2\varphi$); credited to Kotesovec [2, 3]
101-90, eq:product, eq:bridge	Lemmas 4.1, 5.1	same; classical [8]; its residue proof is source 61’s first proof
101-90, eq:puiseux, eq:local	(6.8), (6.12)	same, in its coordinates (Section 36.4)
101-90, eq:sigma, eq:det	Lemma 6.1	other coordinates; Proposition 36.2, with $\det H_{\text{ex}} = (r-1)^2 \det H$
101-90, eq:phases, eq:latticefactor	Section 6.4, (6.10)	coordinate variant: $r-1$ mark points instead of one (Lemma 36.3, Remark 36.4)
101-90, eq:transfer, eq:As, eq:recipe	Section 6.6, Appendix A, (17.10)–(17.12)	the same rule in marker coordinates (Section 36.4); evaluated at this write for $r = 4, 5$ (Remark 36.5)
101-90, $(\alpha_{2,1}, \alpha_{2,2}), (\alpha_{3,1}, \alpha_{3,2})$	(18.6)	same values (source 42)
101-90, D-finiteness by an r -section	Theorem 3.2; proof of Corollary 12.1	second route (Section 36.4)

source	printed here	comparison
101-90, <code>eq:W</code> , <code>eq:inverse</code> , <code>eq:round</code>	(11.2), (11.4), (20.4), (20.3)	same, for every row (Section 36.5); instances of <code>p0:thm:perturbed-inversion</code> and <code>p0:thm:staircase</code>
101-90, “eventually strictly increasing”	Section 11.3	same; superseded by Proposition 37.3
101-86, Theorem 1: leading term, all orders, $\alpha_{5,j} \in \mathbb{Q}(\sqrt{5})$	Theorems 3.1, 3.3	same; a sixth proof for $r = 5$ (Section 37)
101-86, $c_1 = 13(\sqrt{5} - 5)/50$	(3.6)	same value; a fifth computation (after Sections 9, 18, (27.3) and source 77-44)
101-86, $c_2 = (180 - 63\sqrt{5})/125$	Proposition 27.1, Table 6	same value, new route : no recurrence (Theorem 37.2)
101-86, <code>eq:algebraic</code>	(4.5)	–1 times (4.5) under $(x_1, \dots, x_5) = (a, \dots, e)$, term by term (source 42’s sign); derived by source 42’s pair sums and products (Remark 4.2), a fifth derivation
101-86, <code>eq:zeros</code>	(25.3)	same
101-86, <code>eq:four</code> : lattice factor 4	Section 6.4	coordinate variant (Section 37, Remark 38.1)
101-86, <code>eq:recipe-eq:cj</code>	Section 6.6, (17.10)–(17.12)	the same rule in its coordinates, with a Stirling factor for $\binom{5n}{n}$
101-86, <code>eq:W</code> , <code>eq:inverse</code>	(11.2), (28.1)	same; its ℓ_2 is b_2 ; derived from its own $\alpha_{5,2}$, so recurrence-free
101-86, <code>eq:rounding</code>	(20.4) with $r = 5$	same; an instance of <code>p0:thm:staircase</code>

Table 10: The proofs of Theorems 3.1–3.3 in this report. Sources 101-90 and 101-86 are second routes.

statement	earlier proofs	source 101-90	source 101-86
Theorem 3.1, leading term and constant	61 (Section 6); 42 (Theorem 16.1); 75-03 (rate $n^{-1/5}$); 77-44	extreme steps unmarked, Sections 36.2–36.3	$r = 5$ only: zero steps deleted, Proposition 37.1
Theorem 3.1, all orders, algebraic	61 (Section 6.6); 42 (Section 17); 75-03 ($r = 5$, Section 27.1); 77-44	coefficient rule (36.16)	$r = 5$: rule (37.8)
$\alpha_{r,j}$			
Theorem 3.2, D-finiteness	61, 42, 75-03, 77-44 (full diagonal)	r -section, then an ordinary diagonal (36.17)	full diagonal of the sextic
Theorem 3.2, transcendence ($r \geq 3$)	61 only	not claimed	not claimed
Theorem 3.3, $\alpha_{5,1}$	61, 42, 75-03, 77-44 (four contractions)	—	fifth contraction, Theorem 37.2
Theorem 3.3, $\alpha_{5,j} \in \mathbb{Q}(\sqrt{5})$	61; 75-03	—	by the rule (37.8)
$\alpha_{5,2}$ (beyond Theorem 3.3)	75-03 (from the recurrence, Proposition 27.1)	—	without the recurrence, Theorem 37.2

35.3 Conventions of Part VI

Part VI uses the merged notation of Section 1.5, and Table 11 lists every renamed symbol of the two reports. Two new coordinate systems appear, each with its own phase and Hessian: *extreme-unmarked* coordinates (source 101-90), which mark the $r - 2$ interior steps and leave $\pm h$ unmarked, carry the subscript “ex”; the *zero-free* coordinates of source 101-86, which mark two statistics of the four nonzero five-letter steps, carry the subscript “o”. Both reports use the unnormalized step polynomial $\sum_i u^{s_i}$, as source 77-44 does, so their critical times are $1/r$ and $1/4$, not 1. No other normalization changed.

Table 11: Renamed symbols of sources 101-90 and 101-86. A dash means that the source has no such object.

source 101-90	source 101-86	here	remark
$a_r(n)$	a_n	$A_r(n), A_5(n)$	
$\gamma_r = E_r(1/r)$	$E_c = 4\beta$	$\kappa_r; \kappa_4^\circ = 4\varphi^{-2}$	critical values
κ_r	—	$c(r)$	of (3.4)
g (gcd of $h + s$)	—	ν	time period (6.10); not g_r
z (height)	z (height)	u	
$u_j; a_j = \log u_j$	$u, v; a, b$	$z_j; y_j; z_A, z_D; y_A, y_D$	marks; log marks
—	A, D, H	M_2, D_2 , height	counts of a word
$q = r - 2$	—	$d = r - 2$	not Part II’s q
$\alpha_r = \frac{r+1}{2}$	3	β	of Section 11
—	$\beta = \frac{3-\sqrt{5}}{2}$	φ^{-2}	not that β
$c_{r,j}$	c_j	$\alpha_{r,j}, \alpha_{5,j}$	
$c = r \log r$	$c = 5 \log 5$	λ	not $c(r)$
$\sigma^2, \sigma^2(u)$	σ^2	$V, V(\mathbf{y})$	
$\Psi - \log r, \Psi_\ell$	$\Psi - \log 4, \Psi_\ell$	$\Phi_{\text{ex}}, \Phi_{\text{ex},\ell}; \Phi_{\circ}, \Phi_{\circ,\ell}$	phases, as Φ of (6.4)
$\Sigma; C$	Σ	$H_{\text{ex}}, C_{\text{ex}}, H_{\circ}$	not H , nor $C = H^{-1}$
$E(t, u), B(t, u)$	$E(t, u, v); \mathcal{E}$	$\mathcal{E}_{\text{ex}}, B_{\text{ex}}; \mathcal{E}_{\circ}, \mathcal{E}$	
$E_r(t), E_*(t)$	—	$\mathcal{E}_r(t), \mathcal{E}_r^\circ(t)$	length series
$E_c(u)$	E_c	$\kappa_{\text{ex}}(\mathbf{y}); \kappa_{\circ}(\mathbf{y})$	as $\kappa(\mathbf{y})$ of Section 6
$\rho(u), \tau(u)$	ρ, τ	$\rho_{\text{ex}}, \tau_{\text{ex}}; \rho_{\circ}, \tau_{\circ}$	$\rho_{\text{ex}}(0) = 1/r$
ζ_j	$\zeta = -\beta$	$\alpha_j; \zeta = -\varphi^{-2}$	inside roots
Q_r	—	$-Q_r$	(5.5)
$\omega; \alpha; \theta_j$	$\omega; \alpha; \theta, \phi$	$\phi_u; \phi_t; \theta_j, \theta_A, \theta_D$	torus angles
$A_s(u)$	$A_j(u, v)$	$\mathcal{G}_s$	of (17.10)
$e_h(u)$	e_h	$\mathbf{e}_h$	not the weight e of (4.5)
$T_j(\lambda)$	$T_j(\alpha)$	$t_j(\gamma)$	of (17.9)
$b_{r,j}$	b_j	$\mathbf{b}_{r,j}; \mathbf{b}_j^\circ$	not the b_j of Section 11
—	ℓ_2	b_2	of Section 11
—	d_j	$\mathfrak{d}_j$	not $d = r - 2$
—	B_n	$A_5^\circ(n)$	as in Part IV
—	$V_j; A, B, C$ (root jets)	$\mathcal{V}_j; \varkappa_1, \varkappa_2, \varkappa_3$	
$\nu_0, \nu; x; I_r(x)$	$\nu_0, \nu; y; N(y)$	$x_0, x; Y; N_r(Y)$	inversion
$\omega = e^{2\pi i/r}; F(T, u)$	—	$e^{2\pi i j/r}; F^{(r)}(T; \mathbf{z})$	not F_r

Six readings are tempting and false:

- Source 101-90’s κ_r is *not* the κ_r of this report: it is the OEIS-normalized $c(r)$ of (3.4), and its γ_r is this report’s κ_r . Copied verbatim, its display “ $\kappa_r = \gamma_r/(\sqrt{r} 2^{(r-1)/2})$ ” would read $\kappa_r = \kappa_r/(\sqrt{r} 2^{(r-1)/2})$.

- Source 101-90's g (the gcd of the integers $h + s$) is 1 for odd r and 2 for even r : it is the time period ν , with the *opposite* parity of $g_r = \gcd(2, r - 1)$ of (3.1), which is 2 for odd r ; indeed $g_r = 2/\nu$.
- Both reports use z for the height variable and u for marks; here u is the height and z (with $y = \log z$) the marks, as in Section 6.
- Source 101-86's β is $\varphi^{-2} = (3 - \sqrt{5})/2$, not the exponent $\beta = (r + 1)/2$ of Section 11, which source 101-90 calls α_r ; neither is a coefficient $\alpha_{r,j}$.
- The Gaussian-rule coefficients $\mathbf{b}_{r,j}, \mathbf{b}_j^\circ$ (the sources' $b_{r,j}, b_j$) are corrections in powers of the length N , not the logarithmic coefficients b_j of Section 11; source 101-86's ℓ_2 is that b_2 .
- The sources' ν_0, ν are the inverse variable (x_0, x here), not the time period ν ; and H_{ex}, H_o are Hessians in coordinates different from those of H in (6.5), with different determinants (Proposition 36.2).

36 Every fixed alphabet with the extreme steps unmarked (source 101-90)

36.1 Zero steps for odd alphabets

For odd r the step set (3.1) contains 0. Write $\mathcal{E}_r(t) = \mathcal{E}(t, \dots, t)$ for the length series of nonnegative excursions with unit step weights, and $\mathcal{E}_r^\circ(t)$ for the same series without the zero step; for a composition count, $A_r^\circ(n)$ counts zero-free nonnegative excursions with n copies of each nonzero step.

Proposition 36.1 (Zero-step insertion (source 101-90)). *For odd $r \geq 3$,*

$$\mathcal{E}_r(t) = \frac{1}{1-t} \mathcal{E}_r^\circ\left(\frac{t}{1-t}\right), \quad \kappa_r = \frac{r}{r-1} \mathcal{E}_r^\circ\left(\frac{1}{r-1}\right), \quad (36.1)$$

and [write], at balanced content,

$$A_r(n) = \binom{rn}{n} A_r^\circ(n) \quad (n \geq 0). \quad (36.2)$$

Proof [write; source 101-90 states the identity and the insertion factor]. A zero step changes no height. Deleting the zero steps of a nonnegative excursion of length $m + k$ with k zero steps therefore leaves a zero-free nonnegative excursion of length m , and conversely each of the $\binom{m+k}{k}$ ways of inserting k zero steps into a zero-free excursion of length m gives a nonnegative excursion; the two maps are inverse. Hence the length- N coefficient of $\mathcal{E}_r$ is $\sum_{m+k=N} \binom{m+k}{k} [t^m] \mathcal{E}_r^\circ$, and $\sum_k \binom{m+k}{k} t^k = (1-t)^{-m-1}$ gives the first identity. Both series have nonnegative coefficients, and $\kappa_r = \mathcal{E}_r(1/r)$ is finite by (5.4); letting $t \uparrow 1/r$, monotone convergence on both sides gives the second identity, with $t/(1-t) = 1/(r-1)$ and $1/(1-t) = r/(r-1)$ at $t = 1/r$. At balanced content the deletion is a bijection between the words counted by $A_r(n)$ and the pairs formed by a zero-free word counted by $A_r^\circ(n)$ and a set of n positions, among rn , for the zero steps; this is (36.2). $\square$

The cases $r = 3$ and $r = 5$ of (36.2) are the derivation of $A_3(n)$ in Section 7 and (25.3). The number $\kappa_{r-1}^\circ := \mathcal{E}_r^\circ(1/(r-1))$ is the critical excursion value of the uniform zero-free walk on the

$r - 1$ nonzero steps, so the second identity reads $\kappa_r/r = \kappa_{r-1}^\circ/(r-1)$. [write] This agrees with the leading terms: Theorem 16.1, applied to both walks, gives $A_r(n) \sim \frac{\kappa_r}{rn} \frac{(rn)!}{(n!)^r}$ and $A_r^\circ(n) \sim \frac{\kappa_{r-1}^\circ}{(r-1)n} \frac{((r-1)n)!}{(n!)^{r-1}}$, and (36.2) turns the second into the first exactly when $\kappa_r/r = \kappa_{r-1}^\circ/(r-1)$. For $r = 3$, $\kappa_2^\circ = 2$ (Dyck paths) and $\kappa_3 = 3$; for $r = 5$, $\kappa_4^\circ = 4\varphi^{-2}$ (Proposition 37.1) and $\kappa_5 = 5\varphi^{-2} = \frac{5}{2}(3 - \sqrt{5})$. Source 101-90 adds a warning that applies to its own proof: the marked extraction below retains the zero letter, so the factor $\binom{rn}{n}$ is already included and must not be inserted a second time.

36.2 Extreme-unmarked coordinates and their Hessian

Let $s_2 < \dots < s_{r-1}$ be the interior steps (all steps other than $\pm h$), $d = r - 2$, and give them marks $z_j = e^{y_j}$ ($2 \leq j \leq r - 1$), leaving the extreme steps $\pm h$ unmarked:

$$P_{\text{ex}}(u; \mathbf{y}) = u^{-h} + u^h + \sum_{j=2}^{r-1} e^{y_j} u^{s_j}. \quad (36.3)$$

Let $\mathcal{E}_{\text{ex}}(t; \mathbf{z})$ and $B_{\text{ex}}(t; \mathbf{z})$ count nonnegative excursions and bridges with these weights. At length $N = rn$, the extracted interior counts n and the two linear conditions (length rn , final height 0) force both extreme counts to be n , because $s_r - s_1 = 2h \neq 0$. Hence

$$A_r(n) = [t^{rn} z_2^n \cdots z_{r-1}^n] \mathcal{E}_{\text{ex}}(t; \mathbf{z}), \quad [t^{rn} z_2^n \cdots z_{r-1}^n] B_{\text{ex}}(t; \mathbf{z}) = \frac{(rn)!}{(n!)^r}. \quad (36.4)$$

Near $\mathbf{y} = 0$ let $\tau_{\text{ex}}(\mathbf{y})$ solve $\partial_u P_{\text{ex}} = 0$ with $\tau_{\text{ex}}(0) = 1$, and put $\rho_{\text{ex}} = 1/P_{\text{ex}}(\tau_{\text{ex}}; \mathbf{y})$ (so $\rho_{\text{ex}}(0) = 1/r$), $V(\mathbf{y}) = \tau_{\text{ex}}^2 \partial_u^2 P_{\text{ex}} / P_{\text{ex}}$ at $u = \tau_{\text{ex}}$, and

$$\Phi_{\text{ex}}(\mathbf{y}) = \log P_{\text{ex}}(\tau_{\text{ex}}(\mathbf{y}); \mathbf{y}) - \log r - \frac{1}{r} \sum_{j=2}^{r-1} y_j. \quad (36.5)$$

At equal marks

$$V = \frac{1}{r} \sum_i s_i^2 = \begin{cases} (r^2 - 1)/12, & r \text{ odd}, \\ (r^2 - 1)/3, & r \text{ even}, \end{cases} \quad (36.6)$$

since $\frac{2}{r} \sum_{k=1}^h k^2 = h(h+1)/3$ for $r = 2h + 1$, and $\frac{2}{r} \sum_{k=1}^{r/2} (2k-1)^2 = (r^2 - 1)/3$ for even r .

Proposition 36.2 (Hessian in extreme-unmarked coordinates (source 101-90)). Φ_{ex} is stationary at 0, and its Hessian there is

$$H_{\text{ex}} = C_{\text{ex}} - \frac{\mathbf{v}\mathbf{v}^\top}{V}, \quad C_{\text{ex}} = \frac{1}{r} I_d - \frac{1}{r^2} \mathbf{1}\mathbf{1}^\top, \quad v_j = \frac{s_j}{r}, \quad (36.7)$$

the covariance of the marked step indicators after elimination of the height under the uniform step law. It is positive definite, and

$$\det H_{\text{ex}} = \frac{4h^2}{r^r V} = \frac{12(r-1)}{r^r(r+1)} = (r-1)^2 \det H, \quad (36.8)$$

with $\det H$ of (6.6); for $r = 2$ the dimension is 0 and both sides are 1.

Proof (source 101-90). Positivity: otherwise a nonzero linear combination of the marked indicators would be an affine function of the height at every step; it vanishes at the two unmarked extreme steps, which have different heights, so the affine function is zero, and

then every coefficient vanishes. Determinant: C_{ex} has the eigenvalue $1/r$ on $\mathbf{1}^\perp$ and $2/r^2$ on $\mathbf{1}$, so $\det C_{\text{ex}} = 2/r^{r-1}$; the interior steps are symmetric, so $\mathbf{v} \perp \mathbf{1}$ and $C_{\text{ex}}^{-1}\mathbf{v} = r\mathbf{v}$; and $\sum_j s_j^2 = rV - 2h^2$. The matrix determinant lemma gives

$$\det H_{\text{ex}} = \det C_{\text{ex}} \left(1 - \frac{r|\mathbf{v}|^2}{V}\right) = \frac{2}{r^{r-1}} \cdot \frac{2h^2}{rV} = \frac{4h^2}{r^r V}.$$

With (36.6), $4h^2 = (r-1)^2$ and $V = (r^2-1)/12$ for odd r , and $4h^2 = 4(r-1)^2$ and $V = (r^2-1)/3$ for even r ; both give $12(r-1)/(r^r(r+1))$. $\square$

Second proof of (36.8), with the comparison [write]. Follow the proof of Lemma 6.1. Let X take the value $(\mathbf{1}_{\{i=2\}}, \dots, \mathbf{1}_{\{i=r-1\}}, s_i)$ with probability $1/r$. Its covariance matrix has Schur complement H_{ex} in the last entry V , so its determinant is $V \det H_{\text{ex}}$. Its r support points are affinely independent in dimension $r-1$, so that determinant is r^{-r} times the square of the affine-simplex determinant, $\det(\mathbf{1} \ x_i)_{i=1}^r$. Expanding along the d indicator columns, each of which has a single entry 1 (in its interior row), leaves $\pm \det(\mathbf{1} \ -h) = \pm 2h$ [corrected at the independent check; see the end of Section 38.2]. Hence $\det H_{\text{ex}} = 4h^2/(r^r V)$. The same argument with the top adjacent pair unmarked gives (6.6), with $\delta = s_r - s_{r-1}$ in place of $2h = s_r - s_1$. So $\det H_{\text{ex}}/\det H = (2h/\delta)^2$, and $2h/\delta = r-1$ in both parities ($2h = r-1$, $\delta = 1$ for odd r ; $2h = 2(r-1)$, $\delta = 2$ for even r). $\square$

For $r = 4$ and $r = 5$, $\det H_{\text{ex}} = 9/320$ and $8/3125$, against $\det H_4 = 1/320$ (Part V) and $\det H = 1/6250$ (Section 9). Source 101-90's program checks (36.8) exactly for $2 \leq r \leq 20$; the intake rechecked it for $2 \leq r \leq 12$.

36.3 The full torus and the exact periodicity factor

Lemma 36.3 (Dominant torus points (source 101-90)). *Let $|z_j| = |u| = 1$ and $|t| = 1/r$, and write $u = e^{i\phi_u}$, $t = e^{i\phi_t}/r$, $z_j = e^{i\theta_j}$. The kernel $u^h(1 - tP_{\text{ex}}(u; \mathbf{z}))$ vanishes on $|u| = 1$ exactly at*

$$\phi_u = \frac{\pi k}{h}, \quad \phi_t = \pi k, \quad \theta_j \equiv -(h + s_j)\phi_u \pmod{2\pi}, \quad 0 \leq k < 2h. \quad (36.9)$$

These are $2h/\nu = r-1$ distinct mark points, each with ν time phases. At each of them the complete local expansion of the extraction (36.4) equals that at $(t, \mathbf{z}) = (1/r, \mathbf{1})$ term by term, with phase factor one, so the total multiplicity is

$$\nu(r-1) = 2h. \quad (36.10)$$

Proof (source 101-90, with the gauge computation written out [write]). A zero requires equality in $|t \sum_i z_i u^{s_i}| \leq 1$ (with $z_1 = z_r = 1$), so $\phi_t + \theta_i + s_i \phi_u \equiv 0$ for every i . The two unmarked extreme steps give $\phi_t - h\phi_u \equiv \phi_t + h\phi_u \equiv 0$, hence $\phi_u = \pi k/h$ and $\phi_t = \pi k$, and then $\theta_j \equiv -\phi_t - s_j \phi_u = -(h + s_j)\phi_u$. Two values k, k' give the same marks if and only if $2h$ divides $(k - k')(h + s_j)$ for every interior j . The gcd of the integers $h + s_j$ is 1 for odd r (it contains $h + (1-h) = 1$) and 2 for even r (they are $2, 4, \dots, 2h-2$), which is ν ; so there are $2h/\nu$ mark points, each reached by ν values of k whose time phases differ by $\pi h \equiv \pi$ when $\nu = 2$ (for $r = 2$ there are no marks: one mark point with $\nu = 2$ time phases).

For the gauge identity, the weight of a step of type i at such a point is $e^{-i(\phi_t + s_i \phi_u)}$ (also for the extreme steps). Along a path of length N and final height 0 the product of these weights is $e^{-iN\phi_t}$, so $\mathcal{E}_{\text{ex}}(t; \mathbf{z}_k \mathbf{w}) = \mathcal{E}_{\text{ex}}(te^{-i\phi_t}; \mathbf{w})$ identically, and likewise for B_{ex} . In the Cauchy integral for (36.4) near $\mathbf{z}_k$ the factor $\mathbf{z}^{-n\mathbf{1}}$ contributes $e^{in \sum_j (\phi_t + s_j \phi_u)} = e^{ind\phi_t}$, since the interior steps sum to zero, and the time coefficient contributes $e^{-irn\phi_t}$. The total phase is

$e^{-2in\phi_t} = e^{-2\pi ink} = 1$. Since the identity is exact, every correction term is transported, not only the leading one. $\square$

Away from small neighbourhoods of these finitely many points, compactness gives $\epsilon > 0$ such that the kernel has no zero on $|u| = 1$ for $|t| \leq 1/r + \epsilon$; Rouché's theorem keeps h roots inside, and their product is analytic even through collisions inside that group (by contour power sums and Newton's identities), so the excursion series and the bridge constant-term contour are analytic in the larger disk, uniformly on the outer mark arcs, and Cauchy's bound makes those coefficients exponentially smaller. This is the argument of Section 6.4, in other coordinates. Near each dominant point the positive Hessian H_{ex} gives a Gaussian saddle in d variables, and the bridge saddle reproduces the multinomial coefficient exactly (source 101-90's normalization check):

$$\frac{2h}{(2\pi)^{(r-1)/2} \sqrt{V} \sqrt{\det H_{\text{ex}}}} r^N N^{-(r-1)/2} \Big|_{N=rn} = \sqrt{r} (2\pi n)^{-(r-1)/2} r^{rn}, \quad (36.11)$$

because $2h/(\sqrt{V} \sqrt{\det H_{\text{ex}}}) = r^{r/2}$ by (36.8). The excursion amplitude differs by $\kappa_r/N = \kappa_r/(rn)$, which proves (3.5) and (3.4) again, with all even/odd and zero-step normalizations.

Remark 36.4 (The count of dominant points depends on the coordinates [write]). Section 6.4 finds *one* dominant mark point with ν time phases; Lemma 36.3 finds $r-1$. Both are right: the count depends on which two steps are left unmarked, and the Gaussian determinant changes by exactly the compensating factor. Leave any two steps $s_a < s_b$ unmarked. The proof of Lemma 36.3 gives $\phi_t \equiv -s_a \phi_u$ and $(s_b - s_a) \phi_u \equiv 0$, hence exactly $s_b - s_a$ dominant pairs (mark point, time phase). They are distinct: two of them whose ϕ_u differ by $\Delta \phi_u$ have the same marks and time phase only if $s_i \Delta \phi_u \equiv 0$ for every step s_i , hence $\Delta \phi_u \equiv 0$, because the steps have gcd one. The gauge phase is $e^{-iN\phi_t + in \sum_{j \neq a,b} (\phi_t + s_j \phi_u)} = e^{in(s_a - s_b) \phi_u} = 1$ at $N = rn$, so all of them contribute equally. The proof of (36.8) gives $\det H_{a,b} = (s_b - s_a)^2 r^{-r} / V$. Therefore

$$\frac{\#\{\text{dominant pairs}\}}{\sqrt{\det H_{a,b}}} = \sqrt{r^r V} \quad \text{for every choice of } a < b, \quad (36.12)$$

which is $\nu/\sqrt{\det H}$ in the coordinates of Section 6 ($s_b - s_a = \delta = \nu$) and $2h/\sqrt{\det H_{\text{ex}}}$ here ($s_b - s_a = 2h$). The “ $r-1$ dominant mark points” of source 101-90's abstract are a property of its coordinates, not of the problem.

36.4 The local saddle, the coefficient rule, and D-finiteness

The weighted kernel formula [8, Theorem 2] and the bridge identity [8, Corollary 2] are Lemmas 4.1 and 5.1:

$$\mathcal{E}_{\text{ex}}(t; \mathbf{z}) = \frac{(-1)^{h-1}}{t} \prod_{j=1}^h u_j(t; \mathbf{z}), \quad B_{\text{ex}} = 1 + t \partial_t \log \mathcal{E}_{\text{ex}}, \quad (36.13)$$

with u_j the small roots of $u^h(1 - tP_{\text{ex}})$. Source 101-90 rechecks the bridge identity by the residue computation of source 61's first proof. The coalescing small root is $\tau_{\text{ex}} - \tau_{\text{ex}} \sqrt{2/V} \sqrt{1 - t/\rho_{\text{ex}}} + O(1 - t/\rho_{\text{ex}})$ and the product of the other inside roots is analytic, so

$$\mathcal{E}_{\text{ex}} = \kappa_{\text{ex}}(\mathbf{y}) \left(1 - \sqrt{2/V(\mathbf{y})} \sqrt{1 - t/\rho_{\text{ex}}(\mathbf{y})} + O(1 - t/\rho_{\text{ex}}) \right), \quad \kappa_{\text{ex}}(\mathbf{y}) = \mathcal{E}_{\text{ex}}(\rho_{\text{ex}}(\mathbf{y}); \mathbf{y}), \quad (36.14)$$

and the bridge identity gives the bridge singularity $1/\sqrt{2V(1 - t/\rho_{\text{ex}})}$. At one critical time the coefficient amplitudes are therefore $\rho_{\text{ex}}^{-N} N^{-3/2} \kappa_{\text{ex}} / \sqrt{2\pi V}$ for excursions and $\rho_{\text{ex}}^{-N} N^{-1/2} / \sqrt{2\pi V}$ for bridges, uniformly for complex marks near one, before the copies of (36.10) are summed.

All orders. The equal-mark critical kernel factors into the double factor $(u-1)^2$ and its inside and outside factors, which are pairwise coprime even if a stable group has repeated roots. Finite Hensel lifting (equivalently, the implicit equations for the coefficients of the three factors) produces their Taylor jets in time and marks and never divides by differences of roots in the same group; this is the analytic factorization of Section 6.6. With $t = \rho_{\text{ex}}(1-x^2)$ and the branch whose first displacement is $-\tau_{\text{ex}}\sqrt{2/V}x$, $\mathcal{E}_{\text{ex}} = \sum_{k \geq 0} \mathfrak{e}_k(\mathbf{y})(1-t/\rho_{\text{ex}})^{k/2}$, and uniform transfer gives the single-time expansion

$$E_N^{\text{loc}}(\mathbf{y}) = \rho_{\text{ex}}(\mathbf{y})^{-N} N^{-3/2} \left(\sum_{s=0}^J \mathcal{G}_s(\mathbf{y}) N^{-s} + O_{r,J}(N^{-J-1}) \right), \quad \mathcal{G}_s = \sum_{k=0}^s \frac{\mathfrak{e}_{2k+1}}{\Gamma(-k-\frac{1}{2})} t_{s-k} (k + \frac{1}{2}), \quad (36.15)$$

with t_j of (17.9); the time copies are summed afterwards. Let $\Phi_{\text{ex},\ell}$ be the homogeneous degree- ℓ part of Φ_{ex} and G a centred Gaussian vector with covariance H_{ex}^{-1} . Source 101-90's coefficient rule is

$$\mathbf{b}_{r,j} = [x^{2j}] \mathbb{E} \left[\sum_{s=0}^j x^{2s} \frac{\mathcal{G}_s(ixG)}{\mathcal{G}_0(0)} \exp \left(\sum_{\ell=3}^{2j+2} i^\ell x^{\ell-2} \Phi_{\text{ex},\ell}(G) \right) \right], \quad \alpha_{r,j} = r^{-j} \mathbf{b}_{r,j}, \quad (36.16)$$

where $\mathcal{G}_s(ixG)$ is the amplitude at the log marks $\mathbf{y} = ixG$, and the Gaussian expectation of a polynomial F is $[\exp(\frac{1}{2} \sum_{a,b} (H_{\text{ex}}^{-1})_{ab} \partial_a \partial_b) F](0)$. Every letter, including a zero step, is retained, so $N = rn$ is the only rescaling: no Stirling or zero-insertion factor is multiplied in. This is source 42's rule (17.11)–(17.12) ($\mathbf{b}_{r,j} = \chi_j / \mathcal{G}_0(0)$) in marker instead of lattice coordinates. Effective algebraicity follows as in Section 6.6: the root factors have algebraic coefficients, the phase jets and the covariance are rational, and the common $\sqrt{\pi}$ factors cancel. For the remainder, restrict the angular marks to radius $N^{-1/2+\varepsilon}$; Taylor remainders have a polynomial-times-Gaussian majorant, the rest of the local ball contributes $O(e^{-cN^{2\varepsilon}})$, the outer torus is exponentially small, and odd half-orders vanish under Gaussian integration; the gauge identity of Lemma 36.3 identifies all periodic contributions at every order, so no residue class of n is missed.

For $r = 2, 3$ the rule is checked by source 101-90's program from the Catalan and marked Motzkin kernels: $(\alpha_{2,1}, \alpha_{2,2}) = (-9/8, 145/128)$ and $(\alpha_{3,1}, \alpha_{3,2}) = (-11/9, 101/81)$, the values of (18.6) (source 42), and also those read off the closed forms $A_2(n) = \binom{2n}{n}/(n+1)$, $A_3(n) = (3n)!/((n!)^3(n+1))$.

Remark 36.5 (The rule at $r = 4, 5$, checked at this write [write]). Source 101-90's package exercises (36.16) only in dimension $d \leq 1$. At this write the rule was evaluated at $j = 1$ for $r = 2, 3, 4, 5$ in extreme-unmarked coordinates, numerically: kernel roots by 40-digit polynomial root finding, the coefficients in x by Cauchy integrals on small circles, and the Gaussian expectation by a four-point Gauss–Hermite product rule, which is exact for the polynomial of degree at most six that occurs. It reproduces $\alpha_{2,1}$ and $\alpha_{3,1}$, and

$$r^{-1} \mathbf{b}_{4,1} = -0.50978468486767535307\dots, \quad r^{-1} \mathbf{b}_{5,1} = -0.71862232585005467893\dots,$$

equal to $\alpha_{4,1}$ of Theorem 31.1 and to $\alpha_{5,1}$ of (3.6) to 33 and 34 digits. This is numerical evidence that the rule as printed (the factor r^{-j} , the multiplicity (36.10), the Hessian (36.7)) is normalized correctly in dimensions 2 and 3; it is not an interval certificate, and the proof of the rule is the one above.

D-finiteness by an r -section. By (36.13), $\mathcal{E}_{\text{ex}}$ is algebraic. To write the slope- r extraction (36.4) as an ordinary diagonal, source 101-90 first takes the r -section in time,

$$F^{(r)}(T; \mathbf{z}) = \sum_{N \geq 0} ([t^{rN}] \mathcal{E}_{\text{ex}}) T^N = \frac{1}{r} \sum_{j=0}^{r-1} \mathcal{E}_{\text{ex}}(e^{2\pi i j/r} T^{1/r}; \mathbf{z}), \quad (36.17)$$

a finite sum of algebraic functions with integer powers of T , hence algebraic over $\mathbb{Q}(T, \mathbf{z})$; then $A_r(n) = [T^n z_2^n \cdots z_{r-1}^n] F^{(r)}$ is an ordinary diagonal, D-finite by Lipshitz's theorem [9]. This is a second route to the D-finiteness of Theorem 3.2, which takes the full diagonal of the r -variable series (and which source 101-90 mentions as an alternative); the proof of Corollary 12.1 uses sections in the same way. It proves existence of a recurrence, not a particular or minimal one. Source 101-90 makes no transcendence statement.

36.5 Inversion for every row

With $\lambda = r \log r$, $\beta = (r+1)/2$ and $x_0(Y)$ the Lambert core (11.2), source 101-90 gives, for every r ,

$$x(Y) = x_0 - \frac{\alpha_{r,1}}{\lambda x_0 - \beta} - \frac{b_2}{x_0(\lambda x_0 - \beta)} + O_r(x_0^{-3}), \quad b_2 = \alpha_{r,2} - \frac{1}{2}\alpha_{r,1}^2, \quad (36.18)$$

checked by substitution into $\log K_r + \lambda x - \beta \log x + \alpha_{r,1}/x + b_2/x^2$, which leaves a residual $O_r(x_0^{-3})$; its program verifies this symbolically for arbitrary β . [write] Expanding $1/(\lambda x_0 - \beta) = (\lambda x_0)^{-1}(1 + \beta/(\lambda x_0) + O(x_0^{-2}))$ turns (36.18) into (11.4) exactly, so this is the inversion of Section 11; for $r = 5$ it is (28.1), and its nested-logarithm form is Remark 33.1. Further terms are recursive, each with the fixed-order error inherited from the forward expansion. For the threshold $N_r(Y) = \min\{n : A_r(n) \geq Y\}$ source 101-90 states $N_r(Y) = \lceil x_J(Y) + \epsilon_J(Y) \rceil$ with $\epsilon_J(Y) = O_{r,J}((\log Y)^{-J-1})$, where x_J is the large solution of the positive J -term approximation; ceilings of x_J plus and minus an error bound give rigorous bounds, a single ceiling is certified only when that bracket avoids the integers, and “no unqualified rule is asserted inside the boundary windows”. This is (20.4), an instance of `p0:thm:staircase` (Section 1.4), and its first consequence $N_r(Y) = (\log Y + \beta \log \log Y - \log K_r - \beta \log \lambda)/\lambda + O_r(1)$ is (20.3). No novelty is claimed.

37 The five-letter row with the zero steps deleted (source 101-86)

37.1 Two marks on the zero-free walk

Map a letter i to the step $3 - i$; the words counted by $A_5(n)$ become nonnegative excursions with n steps of each of $-2, \dots, 2$. Deleting the zero steps and choosing their positions gives (25.3), $A_5(n) = \binom{5n}{n} A_5^\circ(n)$, where the $\binom{5n}{n}$ choices include positions before the first and after the last nonzero step. Source 101-86 then marks two statistics of a zero-free excursion: the number M_2 of steps ± 2 (mark $z_A = e^{y_A}$) and the signed difference $D_2 = \#(+2) - \#(-2)$ (mark $z_D = e^{y_D}$). With

$$P_\circ(u; z_A, z_D) = u + u^{-1} + z_A(z_D u^2 + z_D^{-1} u^{-2}), \quad A_5^\circ(n) = [t^{4n} z_A^{2n} z_D^0] \mathcal{E}_\circ(t; z_A, z_D), \quad (37.1)$$

length $4n$, $M_2 = 2n$, $D_2 = 0$ and final height 0 force n copies of each nonzero step. The kernel formula [8, Theorem 2 and (20)] gives

$$\mathcal{E}_\circ = -\frac{u_1(t)u_2(t)}{tz_A/z_D}, \quad u^2 - t(z_A/z_D + u + u^3 + z_A z_D u^4) = 0, \quad (37.2)$$

with u_1, u_2 the small roots. Near $z_A = z_D = 1$ let $\tau_\circ$ solve $\partial_u P_\circ = 0$ with $\tau_\circ(1, 1) = 1$ (the implicit function theorem applies since $\partial_u^2 P_\circ(1; 1, 1) = 10$), and $\rho_\circ = 1/P_\circ(\tau_\circ)$. At equal marks

$$u^2 - \frac{1}{4}(1 + u + u^3 + u^4) = -\frac{1}{4}(u-1)^2(u^2 + 3u + 1),$$

so $\rho_o = 1/4$, the other small root is $\zeta = -\varphi^{-2}$ and the remaining root is $-\varphi^2$. The critical equation $u^3 \partial_u P_o = 0$ factors as $(u-1)(u+1)(2u^2+u+2) = 0$; $u = -1$ has $P_o = 0$ and no finite critical time, and $2u^2+u+2 = 0$ gives the only other finite critical value $t = -4/9$. The square-root singularity at ρ_o is therefore isolated uniformly for marks in a small complex neighbourhood of $(1, 1)$; source 101-86 takes “a fixed outer radius between $1/4$ and $4/9$, with a slit from the moving critical value” as the common transfer domain (a sketched step; the full argument of Section 6.6 covers it). With $V_o = \tau_o^2 \partial_u^2 P_o / P_o$ and $\kappa_o = -\tau_o \zeta / (\rho_o z_A / z_D)$,

$$V_o = \frac{5}{2}, \quad \kappa_o = 4\varphi^{-2}, \quad \mathcal{G}_0(1, 1) = \frac{4\varphi^{-2}}{\sqrt{5\pi}} \quad (37.3)$$

at equal marks, and (36.15) holds with ρ_o , τ_o , V_o , analytic amplitudes $\mathcal{G}_j(z_A, z_D)$, and positive branches at the equal-mark point.

The torus. For $|z_A| = |z_D| = |u| = 1$ and $|t| = 1/4$, with angles $\theta_A, \theta_D, \phi_u, \phi_t$, a kernel zero on $|u| = 1$ requires the four phases $\phi_t - \phi_u$, $\phi_t + \phi_u$, $\phi_t + \theta_A - \theta_D - 2\phi_u$, $\phi_t + \theta_A + \theta_D + 2\phi_u$ to vanish modulo 2π , so the only mark points are

$$(z_A, z_D) \in \{(1, 1), (-1, -1), (-1, 1), (1, -1)\}. \quad (37.4)$$

Away from them, compactness, Rouché’s theorem (exactly two roots inside) and the analytic product of those two roots (by contour power sums and Newton’s identities, through collisions) give an exponentially smaller contribution; the singularity of (37.2) at $t = 0$ is removable. For every zero-free excursion $N - M_2$ and $M_2 - D_2$ are even, so $\mathcal{E}_N(-z_A, z_D) = \mathcal{E}_N(z_A, -z_D) = (-1)^N \mathcal{E}_N(z_A, z_D)$ for the length- N coefficients; at $N = 4n$ and z_A -degree $2n$ the four neighbourhoods contribute equally with positive phase. Source 101-86 calls the resulting factor 4 the “lattice factor”. It is the index of the sublattice carrying (N, M_2, D_2) , a property of these coordinates (Remarks 36.4 and 38.1).

37.2 The leading constant of the zero-free count

Proposition 37.1 (The constant of $A_5^o(n)$ (source 101-86)). *Under the uniform law on the steps $-2, -1, 1, 2$, $\text{Var } M_2 = \frac{1}{4}$, $\text{Var } D_2 = \frac{1}{2}$, $\text{Var}(\text{height}) = \frac{5}{2}$, $\text{Cov}(D_2, \text{height}) = 1$, and the other covariances vanish (per step). The Hessian of $\Phi_o(y_A, y_D) = \log P_o(\tau_o; e^{y_A}, e^{y_D}) - \log 4 - y_A/2$ at 0 is therefore*

$$H_o = \text{diag}\left(\frac{1}{4}, \frac{1}{10}\right), \quad \det H_o = \frac{1}{40}, \quad (37.5)$$

and

$$A_5^o(n) = \frac{4\mathcal{G}_0(1, 1)}{2\pi\sqrt{\det H_o}} 4^{4n} (4n)^{-5/2} (1 + O(n^{-1})) = \frac{\varphi^{-2} 256^n}{\sqrt{2} \pi^{3/2} n^{5/2}} (1 + O(n^{-1})). \quad (37.6)$$

With

$$\binom{5n}{n} = \frac{\sqrt{5}}{2\sqrt{2\pi n}} \left(\frac{3125}{256}\right)^n \left(1 - \frac{7}{80n} + O(n^{-2})\right), \quad (37.7)$$

this gives the leading constant $K_5 = (3\sqrt{5} - 5)/(8\pi^2)$ of (3.6).

Proof. The covariances are elementary, and eliminating the height (the Schur complement in its entry $5/2$) changes only $\text{Var } D_2$, to $\frac{1}{2} - 1/\frac{5}{2} = \frac{1}{10}$; source 101-86 notes that this is also the implicit differentiation of $\partial_u P_o = 0$. The real part of the phase on angular marks has a strict Gaussian maximum, its quadratic bound controls the intermediate arcs and the torus argument above the rest, so the factor 4, the Gaussian integral in two variables and (37.3) give (37.6); Stirling’s formula gives (37.7). $\square$

[write] Three consistency checks. (i) For the uniform zero-free walk, Theorem 16.1 (source 42) gives $A_5^\circ(n) \sim \frac{\kappa_4^\circ}{4n} \frac{(4n)!}{(n!)^4}$, with κ_4° its critical excursion value; the small-root formula at the critical weights, as in Proposition 5.2, gives $\kappa_4^\circ = (-1)^3 \cdot 4 \cdot (-\varphi^{-2}) = 4\varphi^{-2} = \kappa_\circ$, and $(4n)!/(n!)^4 \sim 2 \cdot 256^n (2\pi n)^{-3/2}$ turns this into (37.6). (ii) Dividing $K_5 5^{5n} n^{-3}$ by (37.7) gives the same constant, $K_5 \cdot 2\sqrt{2\pi}/\sqrt{5} = \varphi^{-2}/(\sqrt{2}\pi^{3/2}) = 0.0485048217\dots$ (iii) Proposition 36.1 gives $\kappa_5 = \frac{5}{4}\kappa_4^\circ$. The ratio of $A_5^\circ(n)$ (source 101-86's shipped enumeration) to the right side of (37.6) is 0.93936, 0.96907, 0.97924 at $n = 10, 20, 30$. The constant of $A_5^\circ(n)$ was not printed before in this report (Part IV reaches $A_5^\circ(n)$ only through Λ_n). It has the shape of Theorem 31.1's leading term for A215562, with $\varrho = \kappa_4/4$ replaced by $\varphi^{-2} = \kappa_4^\circ/4$: both count balanced excursions of a four-step walk, and Theorem 16.1 gives every such count the form $\frac{\kappa}{4} 256^n / (\sqrt{2}\pi^{3/2} n^{5/2})$.

37.3 The coefficient rule in these coordinates

Let $\Phi_{\circ,\ell}$ be the homogeneous degree- ℓ part of $\Phi_\circ$ and $G = (G_A, G_D)$ a centred Gaussian with covariance $H_\circ^{-1} = \text{diag}(4, 10)$. With $\mathcal{E}_\circ = \sum_{k \geq 0} \mathfrak{e}_k(z_A, z_D)(1 - t/\rho_\circ)^{k/2}$, the amplitudes $\mathcal{G}_s$ are given by (36.15). Source 101-86 sets $\mathbf{b}_0^\circ = 1$ and

$$\mathbf{b}_j^\circ = [x^{2j}] \mathbb{E} \left[\sum_{s=0}^j x^{2s} \frac{\mathcal{G}_s(e^{ixG_A}, e^{ixG_D})}{\mathcal{G}_0(1, 1)} \exp \left(\sum_{\ell=3}^{2j+2} i^\ell x^{\ell-2} \Phi_{\circ,\ell}(G) \right) \right], \quad (37.8)$$

so that $A_5^\circ(n)$ is the right side of (37.6) times $1 + \sum_j \mathbf{b}_j^\circ (4n)^{-j}$. The binomial factor contributes

$$\sum_{j \geq 0} \mathfrak{d}_j w^j = \exp \left[\sum_{k \geq 1} \frac{B_{2k}}{2k(2k-1)} (5^{1-2k} - 1 - 4^{1-2k}) w^{2k-1} \right], \quad (37.9)$$

with Bernoulli numbers B_{2k} (Stirling's series for $\log(5n)! - \log n! - \log(4n)!$); so $\mathfrak{d}_1 = -7/80$ and $\mathfrak{d}_2 = \mathfrak{d}_1^2/2 = 49/12800$. Then

$$\alpha_{5,j} = \sum_{s=0}^j \mathfrak{d}_{j-s} \frac{\mathbf{b}_s^\circ}{4^s}. \quad (37.10)$$

Only the Taylor coefficients able to reach degree $2j$ are needed; the $\mathfrak{e}_k$ follow by finite algebraic substitution in the quartic kernel. All Gaussian moments are rational and all normalized root jets at the base point lie in $\mathbb{Q}(\sqrt{5})$, so (37.8)–(37.10) give $\alpha_{5,j} \in \mathbb{Q}(\sqrt{5})$ for every j , the field statement of Theorem 3.3 by a second route. For the remainder, source 101-86 retains finitely many terms of the uniform Puiseux transfer, Taylor-expands amplitudes and phase in a neighbourhood of radius $N^{-1/2+\epsilon}$, bounds the Taylor remainders by a polynomial times a strictly weaker Gaussian and the complement by $O(e^{-cN^{2\epsilon}})$ locally and exponentially small globally; “no exponentially improved error is asserted”.

37.4 The second correction without the recurrence

Theorem 37.2 (Two corrections for A215570 by Gaussian moments (source 101-86)). *The rule (37.8)–(37.10), evaluated in exact arithmetic in $\mathbb{Q}(\sqrt{5})$, gives*

$$\begin{aligned} \mathbf{b}_1^\circ &= -\frac{97}{20} + \frac{26\sqrt{5}}{25}, & \mathbf{b}_2^\circ &= \frac{681}{32} - \frac{77\sqrt{5}}{10}, \\ \alpha_{5,1} &= \frac{13(\sqrt{5}-5)}{50}, & \alpha_{5,2} &= \frac{180-63\sqrt{5}}{125} = \frac{36}{25} - \frac{63\sqrt{5}}{125}. \end{aligned} \quad (37.11)$$

The recurrence of Theorem 26.1 is not used.

The ingredients follow; they are source 101-86's Section 5 together with the complete jets its program records (`data/101-86-five-second_correction.json`), which the theorem needs and the manuscript only names. Write $w = \log \tau_o$. The critical equation and the step polynomial are

$$2 \sinh w + 4e^{y_A} \sinh(y_D + 2w) = 0, \quad P_o = 2 \cosh w + 2e^{y_A} \cosh(y_D + 2w),$$

and their solution begins

$$w = -\frac{2}{5}y_D - \frac{2}{25}y_A y_D + \frac{3}{125}y_A^2 y_D + \frac{1}{625}y_D^3 - \frac{1}{1875}y_A^3 y_D + \frac{7}{9375}y_A y_D^3 + O(|y|^5).$$

Phase. The homogeneous parts of Φ_o are

$$\begin{aligned} \Phi_{o,2} &= \frac{1}{8}y_A^2 + \frac{1}{20}y_D^2, & \Phi_{o,3} &= -\frac{3}{200}y_A y_D^2, & \Phi_{o,4} &= -\frac{1}{192}y_A^4 - \frac{1}{125}y_A^2 y_D^2 - \frac{41}{60000}y_D^4, \\ \Phi_{o,5} &= \frac{1}{100000}y_A y_D^2 (365y_A^2 + 82y_D^2), \\ \Phi_{o,6} &= \frac{1}{2880}y_A^6 + \frac{23}{37500}y_A^4 y_D^2 + \frac{217}{1200000}y_A^2 y_D^4 + \frac{293}{22500000}y_D^6; \end{aligned} \quad (37.12)$$

in derivatives, $\Phi_{aaa} = 0$, $\Phi_{abb} = -\frac{3}{100}$, $\Phi_{aaaa} = -\frac{1}{8}$, $\Phi_{aabb} = -\frac{4}{125}$, $\Phi_{bbbb} = -\frac{41}{2500}$ ($a = y_A$, $b = y_D$).

Amplitude. Source 101-86 computes the amplitude “particularly economically” from

$$\log \mathcal{G}_0 = \log P_o - y_A - \operatorname{arcosh}(c_0) - \frac{1}{2} \log V_o - \frac{1}{2} \log(2\pi), \quad c_0 = \cosh(y_D + 2w) + \frac{1}{2}e^{-y_A} \cosh w,$$

which follows from the remaining small root: $\tau_o \zeta z_D = -\exp[-\operatorname{arcosh}(c_0)]$ (at equal marks $c_0 = \frac{3}{2}$ and $e^{-\operatorname{arcosh}(3/2)} = \varphi^{-2}$). The normalized amplitude $\mathbf{a}_o = \mathcal{G}_0(e^{y_A}, e^{y_D})/\mathcal{G}_0(1, 1)$ has the homogeneous parts

$$\begin{aligned} [\mathbf{a}_o]_1 &= \left(\frac{\sqrt{5}}{5} - \frac{13}{20}\right)y_A, & [\mathbf{a}_o]_2 &= \left(\frac{367}{800} - \frac{17\sqrt{5}}{100}\right)y_A^2 + \left(\frac{59}{1000} - \frac{3\sqrt{5}}{125}\right)y_D^2, \\ [\mathbf{a}_o]_3 &= \frac{y_A}{240000} \left((25700\sqrt{5} - 57275)y_A^2 + (8496\sqrt{5} - 16908)y_D^2\right), \\ [\mathbf{a}_o]_4 &= \left(\frac{81821}{768000} - \frac{2423\sqrt{5}}{48000}\right)y_A^4 + \left(\frac{35273}{800000} - \frac{1353\sqrt{5}}{50000}\right)y_A^2 y_D^2 + \left(\frac{12517}{6000000} - \frac{23\sqrt{5}}{25000}\right)y_D^4, \end{aligned} \quad (37.13)$$

that is, $\mathcal{G}_{0,a}/\mathcal{G}_0 = 1/\sqrt{5} - 13/20$, $\mathcal{G}_{0,b}/\mathcal{G}_0 = 0$, $\mathcal{G}_{0,aa}/\mathcal{G}_0 = 367/400 - 17\sqrt{5}/50$ and $\mathcal{G}_{0,bb}/\mathcal{G}_0 = 59/500 - 6\sqrt{5}/125$.

Transfer at equal marks. Put $t = (1 - x^2)/4$ and $v_{\pm} = (-1 \pm \sqrt{9 + 4/t})/2$ (the values of $u + u^{-1}$ at the roots of the kernel). The small roots are $(v_+ - \sqrt{v_+^2 - 4})/2$ and $(v_- + \sqrt{v_-^2 - 4})/2$, and the odd Puiseux coefficients of (37.2) are

$$\mathbf{e}_1 = 4 - \frac{12\sqrt{5}}{5}, \quad \mathbf{e}_3 = 8 - \frac{536\sqrt{5}}{125}, \quad \mathbf{e}_5 = 12 - \frac{19108\sqrt{5}}{3125}. \quad (37.14)$$

By (36.15), with $t_1(\frac{1}{2}) = \frac{3}{8}$, $t_2(\frac{1}{2}) = \frac{25}{128}$, $t_1(\frac{3}{2}) = \frac{15}{8}$ and the gamma ratios $-\frac{3}{2}$, $\frac{15}{4}$,

$$\begin{aligned} \frac{\mathcal{G}_1}{\mathcal{G}_0}(1, 1) &= \frac{3}{8} - \frac{3\mathbf{e}_3}{2\mathbf{e}_1} = -\frac{381 - 48\sqrt{5}}{200}, \\ \frac{\mathcal{G}_2}{\mathcal{G}_0}(1, 1) &= \frac{25}{128} - \frac{45\mathbf{e}_3}{16\mathbf{e}_1} + \frac{15\mathbf{e}_5}{4\mathbf{e}_1} = \frac{53669}{16000} - \frac{411\sqrt{5}}{500}. \end{aligned} \quad (37.15)$$

The marked transfer correction. The second-order rule needs $\mathcal{G}_1/\mathcal{G}_0$ to degree two in the marks. With $\mathcal{V}_j = (u\partial_u)^j P_o/P_o$ at $u = \tau_o$, expand the coalescing root as $\tau_o \exp(\varkappa_1 x + \varkappa_2 x^2 + \varkappa_3 x^3 + \cdots)$; then

$$\varkappa_1^2 = \frac{2}{\mathcal{V}_2}, \quad \varkappa_2 = -\frac{\mathcal{V}_3}{3\mathcal{V}_2^2}, \quad \frac{\varkappa_3}{\varkappa_1} = \frac{1}{2} + \frac{5\mathcal{V}_3^2}{36\mathcal{V}_2^3} - \frac{\mathcal{V}_4}{12\mathcal{V}_2^2},$$

and implicit differentiation of $tP_\circ(\zeta(t)) = 1$ supplies the other small root, so that, as functions of the marks,

$$\frac{\mathfrak{e}_3}{\mathfrak{e}_1} = \frac{3}{2} + \frac{5\mathcal{V}_3^2}{36\mathcal{V}_2^3} - \frac{\mathcal{V}_4}{12\mathcal{V}_2^2} - \frac{\mathcal{V}_3}{3\mathcal{V}_2^2} + \frac{1}{3\mathcal{V}_2} + \frac{P_\circ(\tau_\circ)}{\zeta \partial_u P_\circ(\zeta)}. \quad (37.16)$$

This gives

$$\frac{\mathcal{G}_1}{\mathcal{G}_0} = -\frac{381}{200} + \frac{6\sqrt{5}}{25} + \left(\frac{33}{1000} + \frac{9\sqrt{5}}{125}\right)y_A + \left(\frac{3\sqrt{5}}{250} - \frac{171}{10000}\right)y_A^2 - \left(\frac{3\sqrt{5}}{250} + \frac{339}{50000}\right)y_D^2 + O(|y|^3). \quad (37.17)$$

First order. The Gaussian contraction at $j = 1$ is

$$\mathbf{b}_1^\circ = \frac{\mathcal{G}_1}{\mathcal{G}_0} - 2\frac{\mathcal{G}_{0,aa}}{\mathcal{G}_0} - 5\frac{\mathcal{G}_{0,bb}}{\mathcal{G}_0} - \frac{3}{5}\frac{\mathcal{G}_{0,a}}{\mathcal{G}_0} - \frac{91}{100} = -\frac{97}{20} + \frac{26\sqrt{5}}{25}, \quad (37.18)$$

where $-2\mathcal{G}_{0,aa}/\mathcal{G}_0 - 5\mathcal{G}_{0,bb}/\mathcal{G}_0$ is the amplitude Hessian against $H_\circ^{-1}$, $\mathbb{E}\Phi_{\circ,4}(G) = -\frac{31}{40}$ is the quartic phase term, $-\frac{1}{2}\mathbb{E}\Phi_{\circ,3}(G)^2 = -\frac{27}{200}$ the squared cubic term (their sum is $-\frac{91}{100}$), and $-\frac{3}{5}\mathcal{G}_{0,a}/\mathcal{G}_0$ the amplitude–cubic cross term. Then $\alpha_{5,1} = \mathbf{b}_1^\circ/4 - \frac{7}{80}$ by (37.10).

Second order. At $j = 2$, (37.8) needs the phase through degree six (37.12), the amplitude through degree four (37.13), $\mathcal{G}_1/\mathcal{G}_0$ through degree two (37.17) and $\mathcal{G}_2/\mathcal{G}_0$ at equal marks (37.15). In (37.8), $\mathcal{G}_s(e^{y_A}, e^{y_D})/\mathcal{G}_0(1, 1)$ is the product of the ratio $\mathcal{G}_s/\mathcal{G}_0$ with the normalized amplitude $\mathfrak{a}_\circ$ of (37.13); with the bare ratio (37.17) in its place the contraction would give $\frac{665871}{40000} - \frac{14621\sqrt{5}}{2500}$ instead of $\mathbf{b}_2^\circ$ [added at the independent check]. Gaussian integration, a finite exact computation that source 101-86's program performs and that neither the manuscript nor this Part prints term by term (every input is printed above), gives $\mathbf{b}_2^\circ$ of (37.11), and

$$\alpha_{5,2} = \frac{\mathbf{b}_2^\circ}{16} - \frac{7\mathbf{b}_1^\circ}{320} + \frac{49}{12800} = \frac{36}{25} - \frac{63\sqrt{5}}{125}.$$

“This is finite exact arithmetic using the displayed coefficient rule; the sequence recurrence is not involved.”

[write] The last step is checked by hand: $\frac{681}{512} + \frac{679}{6400} + \frac{49}{12800} = \frac{36}{25}$ and $-\frac{77}{160} - \frac{91}{4000} = -\frac{63}{125}$. The value equals $\alpha_{5,2}$ of Proposition 27.1, which source 75-03 obtained from the recurrence of Theorem 26.1 with $\alpha_{5,1}$ as an initial datum; the two derivations are independent (source 101-86 recomputes $\alpha_{5,1}$ as well), and their agreement checks both. At the intake, the exact terms $A_5(n)$ of Theorem 25.2 give $n^3(A_5(n)/(K_5 5^{5n} n^{-3}) - 1 - \alpha_{5,1}/n - \alpha_{5,2}/n^2) = -0.09501, -0.09534, -0.09550, -0.09556$ at $n = 50, 100, 200, 300$, approaching $\alpha_{5,3} = -0.0956673\dots$ of Table 6. Source 101-86's amplitude, phase and transfer jets were reconstructed by its second program in another way (Section 38.2).

37.5 Inversion, and monotonicity at every index

Source 101-86's inverse is the case $r = 5$ of Section 36.5: $x_0(Y) = -\frac{3}{\lambda}W_{-1}(-\frac{\lambda}{3}(K_5/Y)^{1/3})$, $\lambda = 5 \log 5$, and (36.18) with $b_2 = (213 - 83\sqrt{5})/500$ (its ℓ_2), which is (28.1); since its b_2 comes from Theorem 37.2, this second-order inverse is now also recurrence-free. Its rounded threshold $N_5(Y) = \lceil x_J(Y) + \epsilon_J(Y) \rceil$, $\epsilon_J = O_J((\log Y)^{-J-1})$, with the same qualification as in Section 36.5, is (20.4). New is the discrete input.

Proposition 37.3 (Supermultiplicativity (source 101-86 for $m = 1$, $r = 5$; general form [write])). *For every $r \geq 2$ and $m, n \geq 0$,*

$$A_r(m+n) \geq A_r(m)A_r(n); \quad \text{in particular} \quad A_5(n+1) \geq 35A_5(n). \quad (37.19)$$

For $r \geq 3$, $A_r(1) \geq 2$, so $A_r(n+1) \geq 2A_r(n) > A_r(n)$ for every $n \geq 0$; for $r = 2$ the Catalan numbers have $A_2(0) = A_2(1) = 1$ and increase strictly from $n = 1$ on.

Proof [write; source 101-86’s argument for one-copy words]. Let w have content $n\mathbf{1}$ and w' content $m\mathbf{1}$, both satisfying (2.1). The letters of w sum to exactly $\frac{r+1}{2}rn$, so a prefix of ww' of length $rn + k$ ($0 \leq k \leq rm$) has sum $\frac{r+1}{2}rn + \sum_{j \leq k} w'_j \leq \frac{r+1}{2}(rn + k)$; shorter prefixes are prefixes of w . So ww' is counted by $A_r(m + n)$, and w, w' are recovered from ww' by their lengths. For $r \geq 3$ the words $12 \cdots r$ and $213 \cdots r$ are counted by $A_r(1)$: their prefixes of length $k \geq 2$ have sum $k(k+1)/2 \leq \frac{r+1}{2}k$, and the first letter of the second is $2 \leq \frac{r+1}{2}$. Finally $A_r(n) \geq 1$ (the word $(12 \cdots r)^n$), and $A_5(1) = 35$. $\square$

Source 101-86 states only $A_5(n+1) \geq 35A_5(n)$, by concatenation with any one of the 35 valid one-copy words. Section 11.3 appends one fixed block and concludes that $A_r(n)$ is nondecreasing and *eventually* strictly increasing (also source 101-90, and Section 20); by Proposition 37.3 it is strictly increasing at every index for $r \geq 3$, with ratio at least $A_r(1)$ (the values $A_r(1) = 1, 3, 7, 35, 139, 1001$ for $r = 2, \dots, 7$), so the threshold $N_r(Y)$ needs no tail qualification. [write] By Fekete’s lemma the supermultiplicativity also gives $\lim A_r(n)^{1/n} = \sup_n A_r(n)^{1/n}$, which is r^r by Theorem 3.1; this adds nothing beyond the trivial bound $A_r(n) \leq (rn)!/(n!)^r \leq r^{rn}$. The enumerations of both packages satisfy (37.19) on their whole ranges ($r \leq 7$).

38 Computations, the lattice-peak theorem, stale claims and questions

38.1 Source 101-86 and the lattice-peak transfer theorem

Remark 38.1 (Not an instance; dated October 5, 2026). Part II of the collection’s report *Theta Asymptotics for High-Power Ballot Sums* (SetTheory/Cardinals/docs/reports/generating-functions-and-asymptotics/oeis-sequence-asymptotics/a357825-theta-ballot-power-sums, written in batch 98) proves a lattice-peak transfer theorem (tbs:lpt:thm:transfer, Theorem 15.1 there, and its multidimensional form tbs:lpt:thm:multi, Theorem 18.1): for power sums $S_n(r_n) = \sum_x C_{n,x}^{r_n}$ of a row on a moving lattice of mesh d_n , with growing exponents in the critical window $q_n = r_n d_n^2 / \sigma_n^2 \in K \subset (0, \infty)$, $S_n(r_n) / \Omega_n^{r_n} = \vartheta(q_n, \delta_n) + o(1)$, a theta factor that depends on the lattice phase δ_n . Source 101-86’s “lattice factor 4” is *not* an instance of it, for three reasons.

- (i) There is no power sum and no growing exponent: $A_5^\circ(n)$ is one coefficient, extracted by one Cauchy integral; in that theorem’s terms $r_n \equiv 1$.
- (ii) The factor counts the dominant points (37.4) of the mark torus, all of phase one by the parity identities; equivalently, it is the index of the sublattice that carries the support of (N, M_2, D_2) , the classical periodicity factor of a lattice local limit theorem. Read as a lattice sum of a Gaussian, the extraction has a fixed mesh and a variance growing like n , so $q \asymp 1/n \rightarrow 0$: the fine-lattice regime, where Proposition 16.1 there (tbs:lpt:prop:theta-regimes) gives $\vartheta(q, \delta) = \sqrt{2\pi/q} (1 + O(e^{-2\pi^2/q}))$, independent of the phase, and Remark 18.2 there (tbs:lpt:rem:variants) says to parametrize such a sublattice first, which turns the factor into its index. No theta oscillation arises, consistently with the complete, non-oscillating expansion of Theorem 3.1.
- (iii) The factor depends on the coordinates: source 101-90’s analogue is $2h = \nu(r-1)$ (Lemma 36.3), the coordinates of Section 6 give ν , and the combination (36.12) with the Gaussian determinant is the same in all of them. [write] For the zero-free walk with the steps $-2, -1$ marked and $1, 2$ unmarked, there is one dominant point and $\det = 1/640$; source 101-86 has four points and $\det H_o = 1/40 = 16/640$, and $4/\sqrt{1/40} = \sqrt{640}$.

The same holds for source 101-90. That report’s README lists source 101-86’s archive among later arrivals “not examined” for its write; this remark records the examination.

38.2 The two packages

Source 101-90 ships four programs and a driver (`code/101-90-alphabet-*`) with their recorded outputs (`data/101-90-alphabet-*`). `check_constants.py` checks the critical-multiplicity factorization and the inside-root count for $2 \leq r \leq 10$, the exact values $\kappa_2, \dots, \kappa_5$ and $c(4)$, $c(5)$, and records numerical root products, which it labels orientation for $r \geq 6$. `check_balanced_counts.py` enumerates content vectors directly for $r = 2, \dots, 7$ up to $n = 30, 20, 16, 9, 7, 5$, against the Catalan and $r = 3$ closed forms and OEIS initial terms. `check_lattice.py` checks the phase classes and gauge phases of Lemma 36.3, the determinant (36.8) and the bridge normalization exactly for $2 \leq r \leq 20$ (“PASS 19 alphabets”). `check_recipe_small.py` evaluates (36.16) at $j = 1, 2$ for $r = 2, 3$ and verifies (36.18) symbolically.

Source 101-86 ships six programs and a driver (`code/101-86-five-*`), their outputs and a review constants file (`data/101-86-five-*`). `check_kernel.py` checks the sextic over the coefficient box $0 \leq n_i \leq 3$ (1,024 positions, 112 nonzero excursion coefficients), the diagonal 1, 35, 18720, 19369350 and the zero-step removal, with $A_5^\circ(n) = 1, 7, 416, 42570$; `direct_terms.py` enumerates $A_5^\circ(n)$ for $n \leq 30$ by dynamic programming (467,976 states, no recurrence) and compares $\binom{5n}{n} A_5^\circ(n)$ with 13 OEIS terms; `check_jets.py` reconstructs the first-order jets and $\alpha_{5,1}$; `check_second_correction.py` all the second-order jets of Section 37.4 and $\alpha_{5,2}$ (about 100 s recorded); `check_auxiliary.py` the pair-product elimination of the sextic, the second inverse correction and b_2 ; and `audit-check_second_assembly_fast.py` (delivered as `checks/audit/check_second_assembly_fast.py`) reconstructs the implicit root, the phase, the amplitude (algebraically, without arcosh Taylor series) and the transfer jets by finite power-series convolution and assembles the Gaussian contraction independently.

What the second program does not do. Source 101-86’s README calls it the “final independent check”. It reads the main checker’s output `second_correction.json`: it takes the implicit root jet from there and only verifies it (the residual of the critical equation vanishes); it recomputes the phase, amplitude and transfer jets by another method and compares them with the recorded ones; and it then assembles b_2° and $\alpha_{5,2}$ from the recorded jets, which it has just found equal to its own. It is therefore a verification of the main computation, not a derivation independent of its output.

At the intake both suites were rerun on copies in their delivered layout (Python 3.14.4, SymPy 1.14.0): source 101-86’s six jobs passed (“All exact checks passed”; 445 s on a shared machine, against 131 s recorded) and source 101-90’s four (“All checks passed”; 24 s, against 12 s). Their regenerated outputs equal the shipped ones apart from line endings (three files each) or the recorded running time (three and two files). At this write the enumerated $A_5^\circ(n)$, $n \leq 30$, were found equal to $A_5(n)/\binom{5n}{n}$ from Theorem 25.2, and both enumerations to satisfy (37.19).

Both packages state their scope: finite checks “supplement the analytic proof” and use no guessed recurrence; source 101-90’s code “checks examples of that procedure; it is not a general-purpose implementation”. Their review files (`101-90-alphabet-structural_review.md`, `101-86-five-structural_review.md`) review drafts that were not delivered (`leading_comparison.md` and `all_orders_and_inverse.md`; `asymptotic_proof.md`), and their QA receipts record approvals by the producing pipeline. Source 101-86’s `check_receipt.json` has a key, `independent_portable_replay_passed`, that its driver does not write: it was added after the run. The report’s README discloses these points.

Independent check of the write (dated note, 5 October 2026). An independent adversarial check of Part VI, made by the intake after the write (013b32fca), reread every item that carries a proof or a printed value: Proposition 36.1 with its balanced form, Proposition 36.2 with both proofs, Lemma 36.3 with the gauge computation, Remark 36.4, the consistency checks after Proposition 37.1, Theorem 37.2 with every printed jet and the hand check after it, Proposition 37.3, and item (iii) of Remark 38.1. It found every item valid, no counterexample, and no gap in any proof chain; the hypotheses on r (odd, even, and the case $r = 2$) are stated correctly. Three refinements of wording were adopted where they stand; no statement or number changed. The second proof of (36.8) first expanded “along the interior rows”, but an interior row $(1, x_i)$ has up to three nonzero entries; the one-term expansion is along the d indicator columns, as the proof now says. Remark 36.4 now says why its $s_b - s_a$ dominant pairs are distinct: a coincidence would force $s_i \Delta \phi_u \equiv 0$ for every step, and the steps have gcd one. Section 37.4 now says that the rule (37.8) needs $\mathcal{G}_1(e^{y_A}, e^{y_D})/\mathcal{G}_0(1, 1)$, the product of the printed ratio (37.17) with the normalized amplitude $\mathbf{a}_0$. Contracted with the bare ratio, the printed jets give $\frac{665871}{40000} - \frac{14621\sqrt{5}}{2500} = 3.5694\dots$ instead of $\mathbf{b}_2^\circ = 4.0635\dots$; the check made exactly this slip on its first pass.

The check used neither the delivered programs nor the write’s. Its own counters are a dynamic program over content vectors, straight from the prefix condition (2.1), for $A_r(n)$ with $r = 2, \dots, 7$ and $n \leq 12, 9, 6, 5, 3, 3$, and a rolling dynamic program over the steps $\pm 1, \pm 2$ with every prefix height nonnegative, for $A_5^\circ(n)$ with $n \leq 80$; neither uses a recurrence. They agree with each other ($A_5(n) = \binom{5n}{n} A_5^\circ(n)$ for $n \leq 5$), and $\binom{5n}{n} A_5^\circ(n)$ equals Part IV’s formula (Theorem 25.2) for every $n \leq 80$.

For Section 36, the coefficient identity (36.1) holds exactly for $r = 3, 5, 7$ through length 24, and $\kappa_r/r = \kappa_{r-1}^\circ/(r-1)$ to 20 digits for $r = 3, 5, 7, 9$. Exact determinants for $r = 2, \dots, 15$ agree with both forms of (36.8) and with the ratio $(r-1)^2$, every H_{ex} is positive definite, and $\det H_{a,b} = (s_b - s_a)^2 r^{-r}/V$ holds for every pair $a < b$. A brute-force enumeration of the torus points with exact rational angles, for $r = 2, \dots, 13$, gives $r-1$ mark points with ν time phases each and gauge phase one for $n = 1, \dots, 5$, and, for every pair $a < b$, exactly $s_b - s_a$ distinct dominant pairs, all of gauge phase one; a grid search for $r = 4, 5$ finds no other ϕ_u . With the steps 1, 2 of the zero-free walk unmarked there is one dominant pair and the determinant is exactly $1/640$, as item (iii) of Remark 38.1 says.

For Section 37, the covariances and the Schur value $1/10$ behind H_0 were confirmed, the constant of check (ii) after Proposition 37.1 to 20 digits, $\mathfrak{d}_2 = 49/12800$ numerically, and the three ratios at $n = 10, 20, 30$ from the check’s own counts (0.93935647, 0.96906977, 0.97924208). Richardson extrapolation of its own $A_5^\circ(n)$, $n \leq 80$, gives $\alpha_{5,2} = 0.3130217393401059930098$, which agrees with (37.11) to 22 digits; $\alpha_{5,1}$, $\mathbf{b}_1^\circ$ and $\mathbf{b}_2^\circ$ agree to at least 20 digits, and $\alpha_{5,3} = -0.0956673363789441168$ agrees with $-\frac{407}{250} + \frac{4283\sqrt{5}}{6250}$ of Table 6 in all 19 digits. At 45 digits it then recomputed every intermediate jet from its own definitions: the critical point by Newton’s method, ζ as a kernel root, the amplitude as $\kappa/\sqrt{2\pi V}$, and $\mathbf{e}_1, \mathbf{e}_3, \mathbf{e}_5$ and $\mathcal{G}_1/\mathcal{G}_0$ as Cauchy integrals in x , with the coalescing root chosen by continuity, using neither the arcosh formula nor (37.16). The values $\mathbf{e}_1, \mathbf{e}_3, \mathbf{e}_5$, $\mathcal{G}_0(1, 1)$, $\mathcal{G}_1/\mathcal{G}_0(1, 1)$ and $\mathcal{G}_2/\mathcal{G}_0(1, 1)$ agree to 25 digits; the twelve coefficients of $\Phi_{0,2}, \dots, \Phi_{0,6}$, the eight of (37.13) and the four of (37.17) agree within $4.2 \cdot 10^{-30}$, $1.2 \cdot 10^{-28}$ and $6.1 \cdot 10^{-29}$. The derivative list, $\mathbb{E}\Phi_{0,4}(G) = -\frac{31}{40}$, $-\frac{1}{2}\mathbb{E}\Phi_{0,3}(G)^2 = -\frac{27}{200}$, the decomposition (37.18), the transfer constants and the gamma ratios were confirmed, and the formulas for $\varkappa_1, \varkappa_2, \varkappa_3$ and (37.16) were derived again by hand. An exact SymPy contraction of (37.8) with the printed jets and covariance diag(4, 10) gives exactly $\mathbf{b}_1^\circ$ and $\mathbf{b}_2^\circ$ of (37.11), so every input is indeed printed. The hand check after the theorem is correct. From the exact counts, $n^3(\dots) = -0.0950085$ at $n = 50$; with the next term $\alpha_{5,4}/n \approx 0.033/n$ fitted on $50 \leq n \leq 80$, the residual is predicted to be -0.09534 ,

$-0.09550, -0.09556$ at $n = 100, 200, 300$, as printed (those n lie beyond the check's own counts). Proposition 37.3 holds on all of its data, with $A_r(1) = 1, 3, 7, 35, 139, 1001$ for $r = 2, \dots, 7$, and $A_5(n+1) \geq 35A_5(n)$ for $n < 80$, with the smallest ratio, exactly 35, at $n = 0$. In passing it confirmed that expanding $1/(\lambda x_0 - \beta)$ turns (36.18) into (11.4), the normalization check (36.11), and the values 1, 35, 18720, 19369350 and $A_5^\circ(n) = 1, 7, 416, 42570$ of Section 38.2. This was a careful reading with numerical tests, not a formal verification or an external review.

38.3 Stale and coordinate-dependent claims of the two reports

Nothing in either report was found to be wrong: every formula checked agrees with Parts I–V and with the data. Three statements are stale against this report and are corrected here (5 October 2026).

- (a) Source 101-86 says that “the particular conjectured order-three recurrence is still unproved here” (its README; its abstract and its closing section say the proof does not use or prove it), and source 101-90 that “no particular guessed recurrence” is asserted. True of the reports, but the recurrence of Kauers and Koutschan is proved in Part IV (Theorem 26.1), for every $n \geq 0$.
- (b) Both report that a bounded source check “found no proof of this exact balanced-content equivalent” (101-86) or “no prior proof of this precise full fixed-alphabet statement” (101-90). As source-relative statements these are not false; as priority statements they are superseded: the reports were finished about nine hours before this report's first placement, and this report holds four proofs of the same day (sources 61, 42, 75-03, 77-44); the two reports' proofs are the fifth and the sixth.
- (c) Source 101-90's abstract “accounts explicitly for the $r-1$ dominant mark points”. Correct in its coordinates; the number is coordinate-dependent (Remark 36.4): Section 6.4 has one mark point.

Source 101-90's “the sequence is eventually strictly increasing” is true and is strengthened by Proposition 37.3.

38.4 Further questions and research

The open questions of the two reports coincide with questions of Section 22, where they are now credited (dated notes of October 5, 2026); none is answered by this Part. Each is listed with its source, its sketch and what is missing.

- 101.1 *Explicit minimal recurrences* (101-90; its D-finiteness is existence only, and 101-86 does not touch the recurrence). Merged into “Find and explain the fourth-row recurrence”. For $r = 5$ an order-three recurrence is proved (Part IV); its minimality, and every other row, remain open.
- 101.2 *Smaller exponential sectors* beyond the Poincaré expansion (101-86, 101-90). Merged into “Determine the exponentially small sectors”; for A215570 the candidate is the $(2/27)^n$ mode of Remark 27.2. Missing: connection constants and Stokes data.
- 101.3 *Effective error constants for the inversion* (101-86: “no numerical error constant or unconditional exact-ceiling rule”; 101-90: “effective numerical error bounds for inversion”). Merged into “Determine sharp asymptotic error constants”. Missing: explicit forms of the O_J in (36.18) and of ϵ_J , which would certify the single ceiling outside an explicit window.

- 101.4 *A growing alphabet* (101-90: “uniform regimes in which the alphabet size grows”). Merged into “Let the alphabet grow”. Missing: root separation and Gaussian bounds uniform in r ; the constants κ_r , $\det H_{\text{ex}} = 12(r-1)/(r^r(r+1))$ and the multiplicity $2h$ of Lemma 36.3 all move with r .
- 101.5 *The field of the coefficients* (from 101-90’s non-claim “no assertion that their smallest field is $\mathbb{Q}(\gamma_r)$ is needed”). Merged into “Automate all-order coefficient extraction”. Every known value lies in $\mathbb{Q}(\kappa_r)$: $\alpha_{5,j} \in \mathbb{Q}(\sqrt{5}) = \mathbb{Q}(\kappa_5)$ (Theorem 3.3) and $\alpha_{4,1} \in \mathbb{Q}(\varrho) = \mathbb{Q}(\kappa_4)$ (Theorem 31.1). Is $\alpha_{r,j} \in \mathbb{Q}(\kappa_r)$ for every r and j ? The proofs only give the field generated by the coefficients of the inside root factor, which may be larger for $r \geq 6$.
- 101.6 *An engine for every order* (101-86 shows that the Gaussian rule reaches $j = 2$ for $r = 5$ in exact arithmetic; 101-90’s rule is general). Merged into “Automate all-order coefficient extraction”. Missing: the values $\alpha_{4,2}$, $\alpha_{4,3}$ and the rows $r = 6, 7, 8$, which no package computes.

Two steps that the reports only sketch are *not* open, because the theorems they serve are proved in full elsewhere in this report: source 101-86’s common transfer domain (“a fixed outer radius between $1/4$ and $4/9$, with a slit”) and source 101-90’s “any other branch collision on the critical modulus is either one of the enumerated time copies or removable for the selected product” (Section 6.4 and Section 6.6, and the operator-norm bound of Lemma 32.1).

A A reproducible all-order construction

For clarity, the finite procedure implicit in the proof can be organized as follows. The input is a fixed r and a requested relative order M .

Critical algebraic data. Construct Q_r , remove $(u-1)^2$, and isolate the strict inside cluster. Represent its symmetric functions in an algebraic number field. Solve (6.3) recursively for a finite Taylor jet of $\tau(\mathbf{y})$; obtain ρ , Φ , and the tilted probabilities. Internal root collisions can be handled by the coefficients of the cluster factor instead of individual root labels.

Puiseux data. Set $t = \rho(\mathbf{y})(1-w^2)$. Substitute $u_0 = \tau(\mathbf{y})(1 + \sum_{j \geq 1} a_j(\mathbf{y})w^j)$ in the kernel, choosing $a_1 = -\sqrt{2/V(\mathbf{y})}$. The remaining coefficients are obtained recursively. Multiply by the analytic noncritical cluster product and the factor $1/t$ to form the needed Puiseux jet of $\mathcal{E}$ and, independently, of $\log \mathcal{E}$.

Length coefficients. For every odd power w^{2j+1} , use (6.11) and the gamma-ratio expansion. The even powers belong to the local analytic part. Retain enough terms for the requested order, and include the length-period factor ν .

Composition coefficients. Expand the phase about zero, substitute $\mathbf{y} = i\mathbf{x}/\sqrt{N}$, and multiply the amplitude jets. Replace each even monomial by its centered Gaussian moment with covariance H^{-1} ; discard odd monomials. Wick’s rule evaluates these moments as finite sums over pairings. The determinant and common π factor are kept outside the normalized series.

Calibration and inversion. Divide the resulting excursion expansion by the logarithmic reference expansion if desired. The exact coefficient (5.3) and Stirling’s series are an independent normalization check. Convert $N = rn$, compute the formal logarithm of the unit series, and use (11.3) for inverse coefficients.

Every stage is finite at fixed r, M . The procedure proves computability, not an optimal complexity bound or a claim that a complete general-purpose implementation is included in this package. The supplied exact symbolic implementation covers the first correction for $r = 5$. Source 42’s formulation of the same procedure in lattice coordinates is (17.10)–(17.12), its first-order specialization is (18.2), and its program covers the first correction for $r = 3$ and $r = 5$. For $r = 5$, source 75-03’s program (Part IV) replaces the saddle construction above the first order by the recurrence algorithm (27.5)–(27.6) and covers every order; it has recorded $\alpha_{5,j}$ for $j \leq 8$. (2026-10-05 note: source 101-90 states the procedure in marker coordinates with the extreme steps unmarked, (36.16), and its program covers $r = 2, 3$ to second order; source 101-86’s program carries it out for $r = 5$ to second order in exact arithmetic, without the recurrence, Theorem 37.2.)

B Package contents and build instructions

The report ships the programs and recorded outputs of both sources under prefixed names. Source 61 contributes `code/61-balanced-verify.py` and `code/61-balanced-derive_alpha5.py`, their outputs `data/61-balanced-*`, its source and claim provenance notes `61-balanced-SOURCES.md` and its build receipt `61-balanced-BUILD.md`. Source 42 contributes `code/42-late-growing-verify.py` and `code/42-late-growing-first_correction.py` and their outputs `data/42-late-growing-*`. The programs read and write their outputs under the delivered, unprefixed names in a `data` directory beside `code`, and both verification programs write a file named `verification.txt`; they must therefore be run on separate copies in the delivered layout. The report’s README gives the commands.

Source 75-03 (Part IV, added October 2, 2026) contributes `code/75-03-ballot-words-verify.py` and its certificate arrays `code/75-03-ballot-words-certificate_data.py`, the recorded standard output `data/75-03-ballot-words-verification_log.txt`, the exact results `data/75-03-ballot-words-verification_results.json`, and Λ_n and $A_5(n)$ for $n \leq 103$ in `data/75-03-ballot-words-a215570_terms.csv`. Its program imports `certificate_data` by that delivered name and writes its JSON and CSV beside itself under the delivered names; it must likewise be run on a copy under those names, never in this directory.

Source 77-44 (Part V, added October 2, 2026) contributes the driver `code/77-44-bal-verify.py`, the exact checker `code/77-44-bal-check.py`, the derivation scripts `code/77-44-bal-ballot_first_correction.py`, `code/77-44-bal-ballot_four_correction.py`, `code/77-44-bal-verify_ballot_kernels.py`, `code/77-44-bal-verify_ballot_large_dp.py` and `code/77-44-bal-verify_ballot_numerical_derivatives.py`, and the receipts `data/77-44-bal-*`. Its driver checks the hash of its unshipped manuscript, so it runs only in the delivered layout, rebuilt from the arrival archive; three of its scripts write their JSON receipts beside themselves. The README gives the commands.

Sources 101-90 and 101-86 (Part VI, added October 5, 2026) contribute, under the prefixes `101-90-alphabet-` and `101-86-five-`, their drivers `code/101-90-alphabet-run_checks.py` and `code/101-86-five-run_checks.py`, their check programs in `code/` (source 101-86’s review program as `code/101-86-five-audit-check_second_assembly_fast.py`), their recorded outputs, receipts and QA files in `data/`, and their review notes `101-90-alphabet-structural_review.md` and `101-86-five-structural_review.md` (Section 38.2). The

programs expect the delivered layout: each check program writes its JSON output beside itself in a directory `checks`, each driver writes `check_receipt.json` one level up, and source 101-86's review program, delivered in `checks/audit`, reads `../second_correction.json`. They must be run on a copy rebuilt in that layout, never in this directory; the README gives the commands.

The computations use arbitrary-precision integers, rational arithmetic, SymPy and mpmath (the shipped `data/61-balanced-requirements.txt`, whose content is identical to source 42's requirements file); they need no network access. Source 42's `-quick` switch reduces its dynamic-programming boxes; it changes the recorded finite ranges, not the mathematical results, and it overwrites the diagnostic data with the smaller run. Source URLs and the origin of the two externally supplied large-index values are recorded in `61-balanced-SOURCES.md`. No external paper is redistributed. The report is built with pdfL^AT_EX; its bibliography is embedded, so no BibTeX run is needed.

C The full rational telescoping certificate (source 75-03)

This appendix belongs to Part IV and gives every coefficient needed in (26.10)–(26.11). Put $m = n + 2$ (a local letter, not a length), and

$$\mathfrak{R}_{\sharp}(m) = 65m^3 + 203m^2 + 180m + 48, \quad \mathfrak{S}_{\sharp}(m) = 65m^3 + 8m^2 - 31m + 6,$$

which are $\mathfrak{R}(n)$ and $\mathfrak{S}(n)$ of Section 26 after the shift. Let

$$\mathcal{U}_n(t) = -\frac{2t}{m(t^2 - 1)^2} \sum_{j=0}^8 \mathfrak{u}_j(m)t^j, \quad \mathcal{V}_n(t) = \frac{2t^2}{m(t^2 - 1)^2} \sum_{j=0}^8 \mathfrak{v}_j(m)t^j. \quad (\text{C.1})$$

The coefficient polynomials $\mathfrak{u}_j, \mathfrak{v}_j$ (source 75-03's α_j, β_j , not the asymptotic coefficients) are printed below, split into powers of t so that no numerator term is omitted. The only parameter denominator is $m = n + 2$, harmless for $n \geq 0$.

C.1 Coefficients of $\mathcal{U}_n$

$$\begin{aligned} \mathfrak{u}_0 &= 9m(m-1)(3m-2)(3m-1)(3m+1)\mathfrak{R}_{\sharp}(m), \\ \mathfrak{u}_1 &= 3m(3m-2)(3m-1)(3m+1)\mathfrak{R}_{\sharp}(m), \\ \mathfrak{u}_2 &= -3m(3m-1)(3m+1)(28m^2 - 47m + 18)\mathfrak{R}_{\sharp}(m), \\ \mathfrak{u}_3 &= -6m(3m+1)(15m^2 - 18m + 5)\mathfrak{R}_{\sharp}(m), \\ \mathfrak{u}_4 &= (3m+1)(13195m^7 + 14169m^6 - 32535m^5 - 19461m^4 \\ &\quad + 15480m^3 + 5472m^2 - 1888m - 384), \\ \mathfrak{u}_5 &= 11765m^7 + 28943m^6 + 2891m^5 - 27207m^4 \\ &\quad - 12320m^3 + 5368m^2 + 2848m - 384, \\ \mathfrak{u}_6 &= -10920m^8 - 23899m^7 + 10511m^6 + 42923m^5 \\ &\quad + 12369m^4 - 16208m^3 - 5624m^2 + 2368m + 384, \\ \mathfrak{u}_7 &= -16(m-1)(m+1)(m+2)^2\mathfrak{S}_{\sharp}(m), \\ \mathfrak{u}_8 &= 32m(m-1)(m+1)(m+2)^2\mathfrak{S}_{\sharp}(m). \end{aligned}$$

C.2 Coefficients of $\mathcal{V}_n$

$$\begin{aligned}
v_0 &= -3m(m-1)(3m-2)(3m-1)(3m+1)\mathfrak{R}_\#(m), \\
v_1 &= -6m(m-1)(3m-1)(3m+1)\mathfrak{R}_\#(m), \\
v_2 &= -6m(3m+1)(1300m^6 + 3475m^5 + 1269m^4 \\
&\quad - 1895m^3 - 841m^2 + 244m + 48), \\
v_3 &= -2m(3m+1)(715m^5 + 2103m^4 + 1265m^3 \\
&\quad - 523m^2 - 168m + 208), \\
v_4 &= 43745m^8 + 126154m^7 + 67228m^6 - 57642m^5 \\
&\quad - 46597m^4 + 5216m^3 + 6760m^2 - 480m - 384, \\
v_5 &= 24(m+1)^2(325m^5 + 495m^4 - 330m^3 \\
&\quad - 202m^2 + 148m - 16), \\
v_6 &= -16(m+1)(1430m^7 + 3556m^6 + 471m^5 \\
&\quad - 2714m^4 - 307m^3 + 684m^2 - 72m - 24), \\
v_7 &= -16(m+1)(m+2)^2(2m-1)\mathfrak{S}_\#(m), \\
v_8 &= 32m(m+1)(m+2)^2(2m-1)\mathfrak{S}_\#(m).
\end{aligned}$$

C.3 How to verify the certificate without a guesser

Form $\mathcal{R}_0, \mathcal{R}_1$ from the vectors ζ_j of Section 26, insert (C.1) into the right sides of (26.10)–(26.11), subtract the left sides and multiply by common denominators in $\mathbb{Q}[n, t]$: every resulting polynomial coefficient is zero. The calculation involves no root choices, no limits and no sequence values. The shipped `code/75-03-ballot-words-certificate_data.py` stores the same u_j, v_j as integer coefficient arrays (in descending powers of m), and `code/75-03-ballot-words-verify.py` reduces the two residuals exactly to zero; the printed polynomials were expanded and compared with those arrays at the intake (2026-10-02), with no difference. The recurrence then needs only the residue representation (26.5) and the vanishing residue of a rational derivative.

References

- [1] The OEIS Foundation Inc., *A215561*, array of prefix-average multiset permutations (“Array read by antidiagonals: number of late-growing permutations”). Entry by Alois P. Heinz; fixed-row asymptotic conjecture by Vaclav Kotesovec, September 7, 2016. <https://oeis.org/A215561>. Checked October 1, 2026.
- [2] The OEIS Foundation Inc., *A215562*. Leading asymptotic formula by Vaclav Kotesovec, January 31, 2015. <https://oeis.org/A215562>. Checked October 1, 2026.
- [3] The OEIS Foundation Inc., *A215570*. Leading asymptotic formula by Vaclav Kotesovec, September 6, 2016; recurrence conjecture by Manuel Kauers and Christoph Koutschan, March 2, 2023. <https://oeis.org/A215570>. Checked October 1, 2026.
- [4] The OEIS Foundation Inc., *A215571*. <https://oeis.org/A215571>. Checked October 1, 2026.

- [5] The OEIS Foundation Inc., *A215593*. <https://oeis.org/A215593>. Checked October 1, 2026.
- [6] Manuel Kauers and Christoph Koutschan, with terms 0–47 credited in the OEIS entry to Vaclav Kotesovec, *Table of A215570*, $0 \leq n \leq 50$. <https://oeis.org/A215570/b215570.txt>. Checked October 1, 2026.
- [7] M. Kauers and C. Koutschan, *Some D-finite and some possibly D-finite sequences in the OEIS*, Journal of Integer Sequences **26** (2023), Article 23.4.5, especially Section 6.1 and Conjecture 15, pp. 29–30. <https://cs.uwaterloo.ca/journals/JIS/VOL26/Koutschan/kout4.pdf>. Also <https://cs.uwaterloo.ca/journals/JIS/VOL26/Koutschan/kout4.html> and arXiv:2303.02793.
- [8] C. Banderier and P. Flajolet, *Basic analytic combinatorics of directed lattice paths*, Theoretical Computer Science **281** (2002), 37–80. DOI: [10.1016/S0304-3975\(02\)00007-5](https://doi.org/10.1016/S0304-3975(02)00007-5). https://www-lipn.univ-paris13.fr/~banderier/Papers/tcs_banderier_flajolet_2002.pdf. Source 42 cites in particular Theorem 2, Corollary 2 and the singularity analysis of Section 3.
- [9] L. Lipshitz, *The diagonal of a D-finite power series is D-finite*, Journal of Algebra **113** (1988), 373–378. DOI: [10.1016/0021-8693\(88\)90166-4](https://doi.org/10.1016/0021-8693(88)90166-4).
- [10] L. Lipshitz, *D-finite power series*, Journal of Algebra **122** (1989), 353–373. DOI: [10.1016/0021-8693\(89\)90222-6](https://doi.org/10.1016/0021-8693(89)90222-6). Cited by source 77-44 for the diagonal closure.
- [11] D. Denisov and V. Wachtel, *Random walks in cones*, Annals of Probability **43** (2015), 992–1044. <https://arxiv.org/abs/1110.1254>. Cited by source 77-44.
- [12] C. Banderier, C. Krattenthaler, A. Krinik, D. Kruchinin, V. Kruchinin, D. T. Nguyen and M. Wallner, *Explicit formulas for enumeration of lattice paths: basketball and the kernel method*, arXiv:1609.06473, version 2 (2017). <https://arxiv.org/pdf/1609.06473>. Cited by source 77-44, Section 3.3 and Theorem 3.8.
- [13] F. Lindemann, *Ueber die Zahl π* , Mathematische Annalen **20** (1882), 213–225. DOI: [10.1007/BF01446522](https://doi.org/10.1007/BF01446522).
- [14] V. Reshetnikov, *ProveIt*. Repository snapshot: 40b17f2fc79a10a6fc1a50ca9c0c3d6f77be69e5.
Inspected source: the transseries-and-inversion README, at the path identified in Section 2.3. https://github.com/VladimirReshetnikov/ProveIt/tree/40b17f2fc79a10a6fc1a50ca9c0c3d6f77be69e5/Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion. Accessed through the GitHub connector on October 1, 2026.
- [15] V. Reshetnikov, *ProveIt*, GitHub repository, inspected by source 42 at commit bc4d1fa2b849b644461e3b8c597d77aa1c54b9fe, October 1, 2026. <https://github.com/VladimirReshetnikov/ProveIt>. The research-report catalogue distinguishes report claims from refereed or machine-checked results.
- [16] V. Reshetnikov, *ProveIt*, repository root README and Combinatorics/README.md, inspected by source 75-03 at commit 29aca108ed25d714f31b1316512777fbbdc8e006, October 1, 2026, through the GitHub connector. <https://github.com/VladimirReshetnikov/ProveIt/blob/29aca108ed25d714f31b1316512777fbbdc8e006/Combinatorics/README.md>.