

Exact asymptotics and inversion for column convex permutominoes

Research report for OEIS A196275

2 October 2026

Abstract

An exact Bernstein-basis representation converts the known triangular counting recurrence for column-convex permutominoes into moments of a fixed positive self-adjoint operator. Its scalar resolvent is elementary. This gives a closed form for the factorial-normalized generating function, proves the previously conjectured factorial-exponential equivalent, and identifies both numerical constants exactly. The secondary term is an inverse-logarithmic spectral-edge contribution, beginning $1/(2n \log^2 n)$ after factorial normalization. Explicit coefficient functions give an all-orders mixed inverse-power and inverse-logarithmic expansion with sector-level error bounds. A natural spectral interpolation admits a convergent hierarchy of exponentially small inverse corrections about the exact Gamma-envelope inverse. Separately, an elementary global bound shows that rounding the exact envelope inverse recovers every positive integer index from its sequence value. Proofs are analytical; computation is used only for independent validation. A bounded literature search found no prior occurrence of this solution, but does not establish global priority.

[write] **Status and trust boundary, added 5 October 2026, batch 101.** This is bundle Report 96, filed in ProveIt as a single-source research report (provenance in the note at the end of Section 1). It is AI-assisted (its PDF metadata name the author “Research report prepared with OpenAI”), unrefereed, and not formalized. Its one external input is Tomás’s triangular recurrence (1)–(3), which the report takes from [2] and does not re-derive; ProveIt’s intake checked only that it reproduces all 200 terms of the OEIS b-file. The statements about the literature (that the factorial-exponential equivalent was an open conjecture, and that no prior solution exists) are the source’s bounded search of 2 October 2026; the intake did not check them. The numerical checks of Section 11 are not interval certificates.

Contents

1	The counting problem and main results	3
2	An exact Bernstein transfer operator	6
3	The scalar resolvent and its singularities	6
4	The complete spectral measure	8
5	The logarithmic second sector	8
6	A mixed all orders expansion with controlled sectors	9
7	The natural interpolation and ordinary inverse scale	11
8	Convergent exponentially small inverse sectors	12
9	Exact integer recovery and threshold inversion	14

10 Non D finiteness and non P recursiveness	15
11 Reproducible validation	16
12 Literature scope and further research	18

1 The counting problem and main results

A permutomino is a lattice polyomino whose boundary encodes a pair of permutations. Here size n means n rows and n columns, not area. Column-convexity requires each column to be connected. Transposition gives the same count for row-convex objects. Rotations and reflections are not identified in the convention used here.

Let a_n , $n \geq 1$, denote this count, OEIS A196275 [3]. It begins

$$1, 4, 22, 152, 1262, 12232, 135544, 1690080, 23417928, 356958816.$$

The input to this report is the recurrence of Tomás [2, Section 5.2], after changing the index from the number of reflex vertices to size:

$$P_{1,1} = 1, \quad P_{n,p} = 0 \quad \text{outside } 1 \leq p \leq n, \quad (1)$$

$$P_{n,p} = (n+1-p)P_{n-1,p} + \sum_{j=1}^{n+1-p} P_{n-1,j}, \quad n \geq 2, \quad 1 \leq p \leq n, \quad (2)$$

$$a_n = \sum_{p=1}^n P_{n,p}. \quad (3)$$

Set

$$b_0 = \frac{1}{2}, \quad b_n = \frac{a_n}{(n+1)!} \quad (n \geq 1), \quad B(z) = \sum_{n \geq 0} b_n z^n. \quad (4)$$

The artificial value $b_0 = 1/2$ is a normalization, not a count of size-zero objects. Thus B is a factorial-normalized (shifted exponential) generating function. The conventional positive-index exponential generating function is $(zB(z))' - 1/2 = \sum_{n \geq 1} a_n z^n / n!$.

Theorem 1.1 (Exact generating function and spectral decomposition). *For the recurrence (1)–(3),*

$$B(z) = \frac{-(2-z)\log(1-z)}{2z\{2-z+\log(1-z)\}}. \quad (5)$$

Let W_0 be the principal real Lambert function, and put

$$\rho = 1 - W_0(e^{-1}), \quad h = \rho^{-1}, \quad k = \frac{(2h-1)(h-1)}{2}. \quad (6)$$

For every integer $n \geq 0$,

$$b_n = kh^n + r_n, \quad r_n = \frac{1}{2} \int_0^1 t^n w(t) dt, \quad (7)$$

where

$$w(t) = \frac{(2t-1)^2}{\{2t-1-t\log(t/(1-t))\}^2 + \pi^2 t^2}, \quad 0 < t < 1. \quad (8)$$

In particular,

$$a_n \sim k(n+1)!h^n, \quad r_n \sim \frac{1}{2n(\log n)^2}. \quad (9)$$

The constants are

$$h = 1.38593327599819425386062181488251586064513279657058 \dots, \\ k = 0.34191113152179550788392501698972761198399013707239 \dots$$

The first equivalent in (9) proves the numerical conjecture of Beaton, Disanto, Guttman and Rinaldi [1, Section 3.3]. Their exploratory fit $r_n \approx en^g$, with $g \approx -1.272$, is replaced by the second equivalent and its logarithmic corrections. Their paper explicitly expressed uncertainty about the secondary exponent; it was not an established fractional-power law.

[write] **Added 5 October 2026, batch 101.** The status of the 2011 conjecture is the source’s reading of [1], [4] and [5] (Section 12); ProveIt’s intake did not consult these papers or search the literature, so neither the conjectural status nor the absence of a prior proof is checked here (Question 7). The OEIS entry, read by the intake on 5 October 2026, states no formula and no asymptotic for A196275.

The subsequent sections prove the theorem, resolve the secondary sector to all orders, and give two distinct inversion results. The exact interpolation is made explicit wherever continuous inverse terms are discussed.

[write] **Added 5 October 2026, batch 101.** *Provenance.* This report is Report 96 of the session bundle that ProveIt’s docs/incoming intake processed as batch 101: the archive A196275_Exact_Asymptotics_and_Inversion_Source.zip (wrapper directory a196275/, 18 files, 1,611,688 bytes), arrival commit 60f54ea06, placed in f7c612c72. The text is the delivered source/permutomino_asymptotics_standalone.tex, which is the delivered modular source permutomino_asymptotics.tex with its generated fragment validation_summary.tex inlined (the lines of Section 11 from the comment “Generated by render_validation_tables.py” on); the modular edition and the delivered 14-page PDF are not shipped. The package names no ProveIt commit, path or report, so the report has no pin; its title page names no person, and its PDF metadata give the author as “Research report prepared with OpenAI”. It is a single-source report, so no merge choice was made. The write prefixed the 59 delivered labels with `ccp:`, added labels for three sections, the further-questions subsection, its seven questions and one proposition, and added the notes marked [write], the subsection “Further questions and research” (Section 12.1), Proposition 8.2 and the profile bound in Question 2, both with proofs; no statement, proof or number of the source was changed. Theorem numbers follow the section counter (Theorem 1.1 is Theorem 1.1, and so on).

Repository. At the time of writing no other file of the repository mentions A196275, permutominoes or Tomás’s recurrence, and nothing in this report re-proves a result of another report. Where methods meet:

- `generating-functions-and-asymptotics/oeis-sequence-asymptotics/a126764-lconvex-polyominoes` (prefix `lcp:`) counts a different class, L-convex polyominoes by area, and shares no question with this report. Both reports invert a *specified* interpolation and recover the integer threshold by a ceiling (its Section 9.2 and `lcp:ao:cor:brackets`); there the rounding brackets carry asymptotic existence constants without a finite starting index, and batch 101’s Report 95 (its Part III, written separately) finds that eventual rounding of a Lambert model of A126764 is wrong for $2 \leq n \leq 555$. Here Theorem 9.1 gives an explicit rounding rule valid for every $n \geq 1$. Its Lambert branch is W_{-1} , here W_0 .
- `generating-functions-and-asymptotics/oeis-sequence-asymptotics/a113227-powered-catalan` (prefix `pcn:`) also writes counts as moments of a positive measure (`pcn:eq:moments`) and reads the measure off a Stieltjes-type transform (`pcn:eq:Fmeasure`); there the measure is discrete, here it is one atom plus the density w of (18).
- The transseries-and-inversion volume (`Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/`) states the inversion facts used in Sections 7 and 9 in general form; credits are in the notes there.

No statement of this report is formalized in Lean or Rocq, and its place in the collection confers no formal status.

Notation. Many letters carry more than one local meaning. The table fixes them; no symbol was renamed.

Symbol	Meanings (where)	Not to be read as
n, r	size n = number of rows = number of columns; Tomás’s index $r = n - 1$ counts reflex vertices [2]	area; Tomás’s “ r vertices” (a harmless typo in his prose for r reflex vertices)

Symbol	Meanings (where)	Not to be read as
a_n, b_n	$a_n = \text{A196275}$ (offset 1); $b_n = a_n/(n+1)!$ with the artificial $b_0 = 1/2$ (4)	b_0 a count; $a_n/n!$
h, k	$h = 1/(1 - W_0(e^{-1}))$, $k = (2h-1)(h-1)/2$ (6); k is also a summation index in (30), (31) and the coefficient label of $C_{j,k}$	H, K
H	the Laplace weight $H(u) = w(e^{-u})$ (22)	the constant h
w	the density $w(t)$ (8); in the proof of Theorem 10.1, the covering coordinate $w = \log(1-z)$	the same object
B	the series $B(z)$ (4); Bernstein polynomials $B_{j,m}$ (Lemma 2.1); Bernoulli numbers B_{2j} (37)	
W	Lambert W , branches W_0 and W_j ; in Section 7.1 also the number $W = W_0(hY/e)$	
ℓ	$\ell(\lambda) = \int_0^1 dt/(\lambda - t)$ (14); a summation index in the proof of Theorem 6.1	
m	the scalar resolvent $m(\lambda)$ (16); an index in $a_{j,m}$ (27) and the sector index of Theorem 8.1	
c	$c = 2\lambda - 1$ (Section 3); the coefficients c_j (25); small constants c, c' in the proof of Theorem 8.1	
D, d	$D(\lambda)$ (Section 3) and $D(z) = 2 - z + \log(1-z)$ with coefficients d_j (Section 11); $d(\rho)$ in the proof of Proposition 3.2; d_0, d_1 in Section 7.1	$D(z)$ as $D(\lambda)$ at $\lambda = 1/z$ (they differ by a factor λ)
ρ	$\rho = 1/h$ (6); in the proof of Proposition 3.2 the variable $1/\lambda$	
q	the analytic function $q(u)$ (23); $q = 1 - \rho$ in the proof of Proposition 3.2; $q = F''(X_*)$ in Theorem 8.1; $q_0 = \frac{1}{2} \log(2\pi X_0)$ (Section 7.1)	
p	the column index in $P_{n,p}$; $p = F'(X_*)$ in Theorem 8.1; $p_0 = \log(hX_0)$ (Section 7.1)	
N	$N = n + 1$ (20); the threshold $N(y)$ (Corollary 9.2)	
A, a	$A = 1 - \gamma$ (20); $A_0(u) = (2 - e^u)^2$ (22); the interpolation $\mathcal{A}(x)$ (32); Taylor coefficients $a_{j,m}$ (27)	$a_{j,m}$ related to the counts a_n
S	the spectral ratio $S(X)$ (34); in the proof of Theorem 6.1, $S = L - 1 - \log v - i\pi$	
$F, \mathcal{F}, \mathcal{E}$	$F(X)$ the log-envelope (33); $\mathcal{F}_j(L)$ the coefficient functions (28); $\mathcal{E}(x) = k\Gamma(x+2)h^x$ the Gamma envelope (44)	the shipped renderer and verifier README's F , which is $\mathcal{E}$, not $\log \mathcal{E}$
u, v	$u = \lambda - 1 + x$, $v = \lambda - x$ (Lemma 3.1); the Laplace variable u and $v = Nu$ (Sections 5–6); the inverse shift u (Section 8)	
x, X	$x \in [0, 1]$ in Sections 2–3; the real size variable x and $X = x + 1$ (Sections 7–8); X_* the envelope inverse (36), $X(y)$ the interpolation inverse, X_0 the Lambert approximation	X_* and $X(y)$ the same
e	in “ $r_n \approx en^g$ ” (Section 1) a fitted constant of [1]	Euler's number e

What is not claimed (collected from the source): the numerical tests are supporting checks, not the proof, and nothing is interval-certified or formalized; the literature search was bounded and “does not establish global priority”; Theorem 8.1 concerns the specified interpolation only, and other functions agreeing with a_n at the integers can have different exponentially small inverse terms; the rounding of Theorem 9.1 applies to exact sequence values, not to arbitrary y , and forward rounding of $\mathcal{E}(n)$ already fails at $n = 4$; Theorem 6.1 claims no convergence of the inverse-logarithmic series, and fixed-order sums need not improve monotonically at modest n ; $b_0 = 1/2$ is artificial; the coefficients $C_{0,5}$ and $C_{0,6}$ printed in the verifier's README are derived from (31) by the source and not attributed to an external publication; third-party PDFs are not redistributed and no OEIS entry was edited. Open questions are collected in Section 12.1.

2 An exact Bernstein transfer operator

Define the bounded operator on $L^2[0, 1]$

$$(Tf)(x) = (1-x)f(x) + \int_0^{1-x} f(y) dy. \quad (10)$$

It is important that this is an exact encoding of the discrete recurrence, rather than a continuum approximation.

Lemma 2.1 (Bernstein embedding). *For $B_{j,m}(x) = \binom{m}{j} x^j (1-x)^{m-j}$, one has*

$$TB_{j,n-1} = \frac{n-j}{n} B_{j,n} + \frac{1}{n} \sum_{i=0}^{n-j-1} B_{i,n}. \quad (11)$$

Consequently,

$$T^n 1 = \frac{2}{n!} \sum_{p=1}^n P_{n,p} B_{p-1,n}, \quad n \geq 1, \quad b_n = \frac{1}{2} \langle 1, T^n 1 \rangle, \quad n \geq 0. \quad (12)$$

Proof. Multiplication gives $(1-x)B_{j,n-1} = (n-j)B_{j,n}/n$. The integrated Bernstein identity is

$$\int_0^s B_{j,n-1}(y) dy = \frac{1}{n} \sum_{i=j+1}^n B_{i,n}(s).$$

Taking $s = 1-x$ and using $B_{i,n}(1-x) = B_{n-i,n}(x)$ proves (11). Since $T1 = 2B_{0,1}$, induction shows that the coefficient of $B_{p-1,n}$ in $T^n 1$ is $2P_{n,p}/n!$: the first term gives $(n+1-p)P_{n-1,p}$ and the sum gives exactly the second term of (2). Finally, $\int_0^1 B_{j,n}(x) dx = 1/(n+1)$ proves (12), including the separately defined $n = 0$ case. $\square$

Proposition 2.2 (Positivity and cyclicity). *The operator T is positive and self-adjoint, with essential spectrum $[0, 1]$. The vector 1 is cyclic.*

Proof. The integral kernel $\mathbf{1}_{x+y \leq 1}$ is real and symmetric and lies in $L^2([0, 1]^2)$, so the integral part is self-adjoint and compact. Moreover,

$$\langle f, Tf \rangle = \frac{1}{2} \iint_{x+y \leq 1} |f(x) + f(y)|^2 dx dy \geq 0. \quad (13)$$

The multiplication part has essential spectrum $[0, 1]$, unchanged by the compact perturbation. For cyclicity, T raises the degree of a nonzero degree- d polynomial by one: the leading multiplier on x^d is

$$-1 + \frac{(-1)^{d+1}}{d+1} \neq 0.$$

Thus $1, T1, T^2 1, \dots$ span all polynomials, which are dense in $L^2[0, 1]$. $\square$

3 The scalar resolvent and its singularities

For real $\lambda > \|T\|$, define

$$\ell(\lambda) = \log \frac{\lambda}{\lambda-1}, \quad c = 2\lambda - 1, \quad D(\lambda) = c - \lambda \ell(\lambda).$$

The logarithm is continued to $\mathbb{C} \setminus [0, 1]$ by

$$\ell(\lambda) = \int_0^1 \frac{dt}{\lambda - t}. \quad (14)$$

Lemma 3.1 (Explicit resolvent vector). *Writing $u = \lambda - 1 + x$ and $v = \lambda - x$,*

$$f_\lambda(x) = \frac{c/u - \log(u/v)}{D(\lambda)} \quad (15)$$

satisfies $(\lambda - T)f_\lambda = 1$. Hence

$$m(\lambda) := \langle 1, (\lambda - T)^{-1}1 \rangle = \frac{(2\lambda - 1)\ell(\lambda)}{2\lambda - 1 - \lambda\ell(\lambda)}. \quad (16)$$

Proof. Let $g(x) = c/u - \log(u/v)$. Direct differentiation and reflection yield

$$ug'(x) + g(x) + g(1 - x) = 0.$$

The derivative of $(\lambda - T)f_\lambda$ therefore vanishes. At $x = 1$, the integral in T is zero and

$$\lambda f_\lambda(1) = \frac{c - \lambda\ell(\lambda)}{D(\lambda)} = 1.$$

This proves the vector identity. The logarithm in (15) is antisymmetric under $x \mapsto 1 - x$, so its integral is zero, whereas $\int_0^1 c/u \, dx = c\ell(\lambda)$. This proves (16). $\square$

By (12), $B(z) = m(1/z)/(2z)$ near zero, giving (5). The apparent singularity at $z = 0$ is removable.

Proposition 3.2 (The unique outlying eigenvalue). *The only zero of D off $[0, 1]$ is a simple real zero $h > 1$, defined in (6). It is the unique simple dominant eigenvalue of T . The residue of m there is $2k$.*

Proof. There are no nonreal zeros. Indeed,

$$\frac{D(\lambda)}{\lambda} = 2 - \frac{1}{\lambda} - \int_0^1 \frac{dt}{\lambda - t}.$$

For $\lambda = a + ib$ with $b \neq 0$, its imaginary part equals

$$b \left(\frac{1}{|\lambda|^2} + \int_0^1 \frac{dt}{|\lambda - t|^2} \right),$$

which has the strict sign of b . There are no negative zeros, since each subtracted reciprocal term in the last expression is positive when $\lambda < 0$. For $\lambda > 1$, put $\rho = 1/\lambda$. The zero condition becomes

$$d(\rho) := 2 - \rho + \log(1 - \rho) = 0.$$

The function d decreases strictly from 2 to $-\infty$ on $(0, 1)$, so it has exactly one zero. Setting $q = 1 - \rho$ gives $qe^q = e^{-1}$ and hence (6). At the root,

$$\ell(h) = 2 - \frac{1}{h}, \quad D'(h) = \frac{2h - 1}{h(h - 1)}.$$

Therefore

$$\text{Res}_{\lambda=h} m(\lambda) = (2h - 1)(h - 1) = 2k.$$

The pole is an atom of the scalar spectral measure. Cyclicity in Proposition 2.2 identifies the full operator spectrum with the support of this measure. In particular the eigenspace at the pole has dimension one. The remaining spectral support is $[0, 1]$, as verified explicitly next. $\square$

4 The complete spectral measure

For $0 < t < 1$, the upper-half-plane boundary value is

$$\ell(t + i0) = \log \frac{t}{1-t} - i\pi.$$

Substitution in (16) gives

$$-\frac{1}{\pi} \operatorname{Im} m(t + i0) = w(t), \quad (17)$$

with w as in (8). The density is nonnegative and continuous in the open interval; its zero at $t = 1/2$ is harmless. At the endpoints,

$$w(t) \longrightarrow 1 \quad (t \downarrow 0), \quad w(t) \sim \frac{1}{\{\log(1/(1-t)) - 1\}^2 + \pi^2} \quad (t \uparrow 1).$$

Proof of the spectral identity in Theorem 1.1. Stieltjes inversion applied to the scalar spectral measure of 1 gives the density (17) on $(0, 1)$. On every compact subinterval the boundary values are smooth, so there is no additional singular measure there. The endpoint atoms vanish because

$$\lim_{\lambda \rightarrow 0} \lambda m(\lambda) = 0, \quad \lim_{\lambda \rightarrow 1} (\lambda - 1)m(\lambda) = 0,$$

with the limits taken off the interval. Proposition 3.2 supplies the only atom outside the interval, of mass $2k$ at h . Thus

$$d\mu(t) = 2k \delta_h(dt) + w(t) \mathbf{1}_{(0,1)}(t) dt. \quad (18)$$

Its total mass is $\|1\|^2 = 1$. Taking moments and using (12) proves (7), and in particular

$$\int_0^1 w(t) dt = 1 - 2k, \quad 0 < r_n \leq \frac{1}{2} - k. \quad (19)$$

Since $h > 1$, this already proves $a_n \sim k(n+1)!h^n$. $\square$

The same formula also shows that $(r_n)_{n \geq 0}$ is a Hausdorff moment sequence, hence completely monotone in the discrete sense. For example $r_n > r_{n+1} > 0$, and every alternating finite difference has the appropriate nonnegative sign. These exact restrictions are useful numerical consistency checks.

5 The logarithmic second sector

Write

$$N = n + 1, \quad L = \log N, \quad A = 1 - \gamma. \quad (20)$$

The substitution $t = e^{-u}$ in (7) gives the exact Laplace integral

$$r_n = \frac{1}{2} \int_0^\infty e^{-Nu} H(u) du, \quad (21)$$

where

$$H(u) = \frac{A_0(u)}{\{\log u + 1 + q(u)\}^2 + \pi^2}, \quad (22)$$

$$\begin{aligned} A_0(u) &= (2 - e^u)^2, \quad q(u) = 1 - e^u + \log \frac{e^u - 1}{u} \\ &= -\frac{u}{2} - \frac{11u^2}{24} - \frac{u^3}{6} - \frac{121u^4}{2880} + O(u^5). \end{aligned} \quad (23)$$

Both A_0 and q are analytic at zero. The function $H(u) = w(e^{-u})$ is bounded on $(0, \infty)$.

Theorem 5.1 (Inverse logarithmic expansion). *For each fixed $K \geq 0$,*

$$r_n = \frac{1}{2NL^2} \left(\sum_{j=0}^K \frac{c_j}{L^j} + O_K(L^{-K-1}) \right), \quad (24)$$

where

$$c_j = \frac{1}{\pi} \operatorname{Im} \left[(\partial_\alpha + 1 + i\pi)^{j+1} \Gamma(\alpha + 1) \right]_{\alpha=0}. \quad (25)$$

The first coefficients are

$$\begin{aligned} c_0 &= 1, & c_1 &= 2A, \\ c_2 &= 3A^2 - \pi^2/2, & c_3 &= 4A^3 - 2\pi^2 A - 8\zeta(3), \\ c_4 &= 5A^4 - 5\pi^2 A^2 - 40A\zeta(3) + \pi^4/12. \end{aligned}$$

Proof. Scale $u = v/N$. At leading order in $1/N$, (21) becomes

$$\frac{1}{2\pi N} \operatorname{Im} \int_0^\infty \frac{e^{-v}}{L - 1 - \log v - i\pi} dv. \quad (26)$$

The replacement has error $O(N^{-2}L^{-2})$, a special case of the stronger theorem below. Expanding the reciprocal in $1/L$ gives (25), since

$$\int_0^\infty e^{-v} (\log v)^j dv = \Gamma^{(j)}(1).$$

For rigor, split at $v = e^{-\varepsilon L}$ and $v = e^{\varepsilon L}$, with any fixed $0 < \varepsilon < 1/2$. On the middle interval a finite geometric expansion has a uniformly controlled remainder, which is bounded after integration by the next logarithmic moment. The small- v tail is exponentially small in L and the large- v tail is superexponentially small. This proves every fixed-order assertion in (24). The displayed coefficients follow from the standard derivatives of Γ at 1. $\square$

[write] **Added 5 October 2026, batch 101.** The rigour paragraph above is condensed: it names the split points and the size of each tail but writes out neither the remainder bound of the finite geometric expansion on the middle interval nor the comparison of the truncated Gamma moments with the full ones. The intake checked the steps and found them correct as sketched; a written-out version is part of Question 6. The verifier's README (`verification-README.md`) prints two further coefficients, $C_{0,5} = c_5$ and $C_{0,6} = c_6$, obtained from (31) by the source's symbolic code; the intake confirmed numerically that they equal (25) at $j = 5, 6$.

Thus the effective exponent of the residual tends to -1 with a leading drift $-2/\log n$. A finite-range fit can consequently resemble a more negative fractional exponent. There is no algebraic $1/n$ correction multiplying the dominant term kh^n in b_n : that contribution is the exact isolated pole, while all remaining moments come from $[0, 1]$.

6 A mixed all orders expansion with controlled sectors

A finite truncation in inverse logarithms cannot resolve the next power of $1/N$. To preserve this distinction, define actual coefficient functions, rather than only a double formal series.

Put

$$a_{j,m} = [u^j] A_0(u) q(u)^m, \quad 0 \leq m \leq j, \quad (27)$$

and

$$\mathcal{F}_j(L) = \frac{1}{\pi} \operatorname{Im} \sum_{m=0}^j a_{j,m} \int_0^\infty \frac{e^{-v} v^j}{(L-1-\log v - i\pi)^{m+1}} dv. \quad (28)$$

The integrals converge for every real L .

Theorem 6.1 (Mixed sector calculus). *For each fixed $J \geq 0$,*

$$r_n = \frac{1}{2} \sum_{j=0}^J N^{-j-1} \mathcal{F}_j(\log N) + O_J(N^{-J-2}(\log N)^{-2}). \quad (29)$$

Each coefficient function has an all-orders expansion

$$\mathcal{F}_j(L) \sim \sum_{k \geq 0} C_{j,k} L^{-k-2}, \quad (30)$$

where

$$C_{j,k} = \frac{1}{\pi} \operatorname{Im} \sum_{m=0}^{\min(j,k)} a_{j,m} \binom{k+1}{m} \left[(\partial_\alpha + 1 + i\pi)^{k+1-m} \Gamma(\alpha+1) \right]_{\alpha=j}. \quad (31)$$

In particular $C_{0,k} = c_k$ and

$$\mathcal{F}_1(L) = -2L^{-2} + (-9 + 4\gamma)L^{-3} + O(L^{-4}).$$

Proof. Let $S = L - 1 - \log v - i\pi$. After $u = v/N$, the kernel in (21) is

$$\frac{1}{\pi} \operatorname{Im} \frac{A_0(v/N)}{S - q(v/N)}.$$

The Taylor coefficients in v/N are precisely

$$\sum_{m=0}^j a_{j,m} S^{-m-1}.$$

Fix $0 < \eta < 1$ and a small $\delta > 0$. On $v \leq N^\eta$, $|S| \geq (1-\eta)L - 1$, and $q(v/N)$ is uniformly small. Taylor's theorem and a finite geometric expansion give an imaginary-part remainder $O_J((v/N)^{J+1}L^{-2})$. The second denominator power comes from taking the imaginary part of a rational function with real Taylor coefficients. Integrating against e^{-v} and including the factor $1/(2N)$ proves the stated error there. The range $N^\eta < v < N\delta$ is exponentially small in N^η ; the remaining $u \geq \delta$ tail is $O(e^{-N\delta})$, since H is bounded. Extending each coefficient integral to infinity changes only exponentially small tails. This proves (29).

Finally expand

$$S^{-m-1} = \sum_{\ell \geq 0} \binom{m+\ell}{m} (1 + \log v + i\pi)^\ell L^{-m-\ell-1}$$

to any finite order. The same tail argument as in Theorem 5.1 justifies termwise integration. Taking the imaginary part eliminates the zero-power term, and collecting the coefficient of L^{-k-2} gives (31). $\square$

Theorem 6.1 is the precise meaning of the mixed expansion in this report. It provides an explicit algorithm for every coefficient and a remainder at the level of each inverse-power sector. It does not claim convergence of the separate inverse-logarithmic series.

[write] **Added 5 October 2026, batch 101.** The proof of Theorem 6.1 is also condensed. Two steps are asserted rather than written out: that the Taylor remainder of $A_0(v/N)/(S - q(v/N))$ has *imaginary part* $O_J((v/N)^{J+1}L^{-2})$ on $v \leq N^\eta$ (one power of L better than its modulus, because $\text{Im } S = -\pi$ is bounded while $|S| \gtrsim L$), and the termwise integration of the expansion of S^{-m-1} , which is referred to the tail argument of Theorem 5.1. The intake rederived $\mathcal{F}_1(L) = -2L^{-2} + (-9 + 4\gamma)L^{-3} + O(L^{-4})$ by hand. Writing these steps out is part of Question 6.

7 The natural interpolation and ordinary inverse scale

The spectral measure supplies a specific interpolation:

$$\mathcal{A}(x) = \Gamma(x+2) \left[kh^x + \frac{1}{2} \int_0^1 t^x w(t) dt \right], \quad x > -1. \quad (32)$$

It agrees with a_n at every positive integer and is real analytic. In fact it is strictly increasing on all of $[1, \infty)$. To prove this, write

$$b(x) = \frac{1}{2} \int t^x d\mu(t).$$

The second derivative of $\log b(x)$ is the variance of $\log t$ under the tilted positive measure, hence is strictly positive. Since $b(0) = b(1) = 1/2$, strict convexity implies $(\log b)'(1) > 0$ and therefore $b'(x) > 0$ for $x \geq 1$. The Gamma factor is also strictly increasing there. Thus for every $y \geq 1$, the interpolation inverse exists and the integer threshold is exactly $\lceil \mathcal{A}^{-1}(y) \rceil$. This assertion uses the specified interpolation, not an envelope approximation.

Set $X = x + 1$ and define

$$F(X) = \log \Gamma(X+1) + X \log h + \log(k/h), \quad (33)$$

$$I(X) = \int_0^\infty e^{-Xu} H(u) du, \quad S(X) = \frac{h}{2k} h^{-X} I(X). \quad (34)$$

Then, exactly,

$$\mathcal{A}(X-1) = e^{F(X)}(1 + S(X)). \quad (35)$$

The function I is analytic for $\Re X > 0$; its differentiated asymptotic expansions follow by inserting powers of u into the Laplace integral.

For a target y tending to infinity, let $X_* = X_*(y)$ be the *exact* Gamma-envelope inverse

$$F(X_*) = \log y. \quad (36)$$

This definition will be retained when exponentially small corrections are resolved.

7.1 Lambert W and the Stirling expansion

Let

$$Y = \log(yh/k), \quad W = W_0(hY/e), \quad X_0 = Y/W, \quad p_0 = \log(hX_0) = W + 1.$$

The leading equation is $Y = X_0(\log(hX_0) - 1)$. Stirling's expansion gives

$$Y = X_*(\log(hX_*) - 1) + \frac{1}{2} \log(2\pi X_*) + \sum_{j \geq 1} \frac{B_{2j}}{2j(2j-1)X_*^{2j-1}}, \quad (37)$$

as an asymptotic expansion, where B_{2j} are Bernoulli numbers.

For clarity, the first two corrections are explicit. Put $q_0 = \frac{1}{2} \log(2\pi X_0)$ and $d_0 = -q_0/p_0$. Then

$$X_* = X_0 - \frac{q_0}{p_0} - \frac{d_0^2/2 + d_0/2 + 1/12}{X_0 p_0} + O\left(\frac{1}{X_0^2 p_0}\right). \quad (38)$$

Indeed substitute $X_* = X_0 + d_0 + d_1$ into (37), expand through X_0^{-1} , and set the successive residuals to zero. The unremoved residual is $O(X_0^{-2})$, while the derivative is asymptotic to p_0 . Repeating this coefficient cancellation with further Stirling terms yields arbitrary ordinary inverse order. For real positive arguments, the usual remainder bounds for Stirling's formula justify each fixed truncation [7, Section 5.11].

A finite truncation of (38) has an algebraic error much larger than the spectral correction. Therefore the next section is centered on (36), not on a finitely truncated Lambert approximation.

[write] **Added 5 October 2026, batch 101: instances of repository results.** The inversion of this section is an instance of the general apparatus of ProveIt's transseries-and-inversion volume (`Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/transseries_and_inversion.tex`), and no novelty is claimed for the method. The leading equation $Y = X_0(\log(hX_0) - 1)$ is `p0:prop:factorial-core` with $\kappa = 1$ and $d = \log h - 1$: its $v = W_0(Ye^d) = W_0(hY/e)$ is the W above, its solution $\exp(v - d) = e^{W+1}/h$ equals Y/W because $We^W = hY/e$, and its slope $\kappa(v+1) = W+1$ is p_0 . The threshold identity $N(y) = \lceil \mathcal{A}^{-1}(y) \rceil$ stated before (33) is part (1) of `p0:thm:staircase` for the admissible interpolation $\mathcal{A}$ on $[1, \infty)$; the source supplies the monotonicity proof that makes $\mathcal{A}$ admissible.

8 Convergent exponentially small inverse sectors

Let $X(y)$ denote the eventual inverse in (35). Define, near $u = 0$,

$$G(u) = F(X_* + u) - F(X_*), \quad Q(u) = \frac{uS(X_* + u)}{1 - e^{-G(u)}}, \quad (39)$$

with the removable value $Q(0) = S(X_*)/F'(X_*)$.

Theorem 8.1 (All exponential inverse sectors). *For y sufficiently large, the following series converges to the inverse of the interpolation (32):*

$$X(y) = X_* + \sum_{m \geq 1} \frac{(-1)^m}{m!} \left[\frac{d^{m-1}}{du^{m-1}} Q(u)^m \right]_{u=0}. \quad (40)$$

For each fixed M , its remainder after $m = M$ is

$$O_M \left(\frac{S(X_*)^{M+1}}{F'(X_*)} \right). \quad (41)$$

Writing $p = F'(X_*)$, $q = F''(X_*)$, and evaluating S, S' at X_* , the first two sectors are

$$X - X_* = -\frac{S}{p} + \frac{SS'}{p^2} + \frac{S^2}{2p} - \frac{qS^2}{2p^3} + O(S^3/p). \quad (42)$$

For each fixed m , the leading term of the m th sector is

$$\frac{(-1)^m}{m} \left(\frac{h}{2k} \right)^m \frac{h^{-mX_*}}{X_*^m (\log X_*)^{2m} \log(hX_*)}. \quad (43)$$

All its mixed inverse-power and inverse-logarithmic corrections are generated by (31), Gamma derivatives, and the finite differentiation in (40).

Proof. Replace S in (35) by εS . Relative to the envelope solution, the inverse equation is

$$e^{G(u)}(1 + \varepsilon S(X_* + u)) = 1,$$

equivalently $u = -\varepsilon Q(u)$. Lagrange inversion gives (40) as the Taylor series in ε at zero.

To justify convergence at $\varepsilon = 1$, choose a sufficiently small fixed $c > 0$ and the disk $|u| \leq c/F'(X_*)$. The function $G(u)$ is asymptotically $F'(X_*)u$ uniformly on this disk; hence $1 - e^{-G(u)}$ has only its simple zero at zero there. Moreover the Laplace representation gives

$$|S(X_* + u)| \leq CS(X_*), \quad |Q(u)| \leq CS(X_*)/F'(X_*).$$

For example, bound the modulus of the Laplace transform by $I(X_* - |\Re u|)$ and use its asymptotic equivalent to compare it with $I(X_*)$. Rouché's theorem on the disk now gives a unique root of $u + \varepsilon Q(u) = 0$ for $|\varepsilon| < c'/S(X_*)$. That root is simple and analytic in ε . This disk contains 1 for all sufficiently large X_* . Cauchy's coefficient bounds on a fixed fraction of this radius give (41).

Expanding the equation to order ε^2 proves (42). Finally $F' \sim \log(hX_*)$ and

$$S(X_*) \sim \frac{h}{2k} \frac{h^{-X_*}}{X_*(\log X_*)^2}.$$

On the local scale $1/F'$, S is asymptotically constant and F is asymptotically linear. The resulting inverse shift is exactly $-\log(1 + \varepsilon S)/F'$, whose m th coefficient is $(-1)^m S^m/(mF')$. This proves (43); derivatives of S and higher derivatives of F give the lower orders through the exact formula. $\square$

[write] **Added 5 October 2026, batch 101: the disk estimate written out.** The convergence paragraph of the proof above asserts the bound $|S(X_* + u)| \leq CS(X_*)$ on the disk, with the hint “bound the modulus of the Laplace transform by $I(X_* - |\Re u|)$ and use its asymptotic equivalent”. The proposition below proves the bound with an elementary log-derivative estimate in place of the asymptotic equivalent, and writes out the Rouché and Cauchy steps with explicit radii. It is a proof by the write, not by the source; the radii $1/(4p)$ and $1/(8p)$ are one admissible choice of the source's $c/F'(X_*)$.

Proposition 8.2 ([write] Disk estimate for Theorem 8.1). *Write $p = F'(X_*)$, $S_* = S(X_*)$ and $\Delta = \{u \in \mathbb{C} : |u| \leq 1/(4p)\}$. There is X_1 such that for every $X_* \geq X_1$:*

1. $|S(X_* + u)| \leq 2h^{1/4}S_*$ for $u \in \Delta$;
2. on Δ , $1 - e^{-G(u)}$ vanishes only at $u = 0$, where its zero is simple, and $|1 - e^{-G(u)}| \geq 0.45p|u|$; hence Q is analytic on Δ and $|Q(u)| \leq KS_*/p$ there, with $K = 5h^{1/4}$;
3. for $|\varepsilon| < R := 1/(8KS_*)$ the equation $u + \varepsilon Q(u) = 0$ has exactly one root $u(\varepsilon)$ in $|u| < 1/(8p)$; it is analytic in ε , real for real ε , and its Taylor coefficients at $\varepsilon = 0$ are the terms of (40);
4. if moreover $R \geq 2$, then $X(y) = X_* + u(1)$ and the remainder of (40) after $m = M$ is at most $\frac{1}{4p}(8KS_*)^{M+1}$, which is (41).

Since $S_* \rightarrow 0$ as $X_* \rightarrow \infty$ (below), $R \geq 2$ holds for all large X_* , that is, for all large y .

Proof. Facts about H and I . The density w is continuous on $(0, 1)$ with finite limits at both ends (Section 4), so $0 \leq H \leq M_H$ for a constant M_H ; and $H > 0$ except where w vanishes, at the single point $u = \log 2$, so $I(X) > 0$ for real $X > 0$. For $0 < u \leq 1$ one has $1 < (e^u - 1)/u \leq e - 1$, hence $1 - e < q(u) \leq \log(e - 1)$ and $|1 + q(u)| \leq 2$; also $A_0(u) \geq \frac{1}{2}$ for $u \leq \frac{1}{4}$. If $0 < u \leq e^{-7}$, then $\Lambda = \log(1/u) \geq 7$ and $(\Lambda + 2)^2 + \pi^2 \leq 2\Lambda^2$ (true for $\Lambda \geq 2 + \sqrt{8 + \pi^2}$), so (22) gives

$$H(u) \geq \frac{1}{4\log^2(1/u)} \quad (0 < u \leq e^{-7}).$$

For real $t > 0$ the integrals defining $I(t)$ and $-I'(t) = \int_0^\infty se^{-ts}H(s)ds$ converge, because H is bounded, and I is decreasing. For $t \geq e^7$ the interval $[1/(2t), 1/t]$ lies in $(0, e^{-7}]$, and $\log(1/s) \leq \log(2t)$ on it, so

$$I(t) \geq \int_{1/(2t)}^{1/t} e^{-ts}H(s)ds \geq \frac{1}{8et\log^2(2t)}, \quad -I'(t) \leq M_H \int_0^\infty se^{-ts}ds = \frac{M_H}{t^2}.$$

Hence $\mu(t) := -I'(t)/I(t) \leq 8eM_H \log^2(2t)/t$, which decreases for $t \geq e^7$. For real $X \geq e^7 + \frac{1}{4}$,

$$\log \frac{I(X - \frac{1}{4})}{I(X)} = \int_{X-1/4}^X \mu(t) dt \leq \frac{2eM_H \log^2(2X)}{X - \frac{1}{4}} \rightarrow 0,$$

so $I(X - \frac{1}{4}) \leq 2I(X)$ for all $X \geq X_2$, say. Also $I(X) \leq M_H/X$, so $S(X) = \frac{h}{2k} h^{-X} I(X) \rightarrow 0$ as $X \rightarrow \infty$.

(1). Let $X_* \geq 3$. Then $p = \psi(X_* + 1) + \log h \geq \psi(4) + \log h > 1$, so $|u| \leq \frac{1}{4}$ on Δ . For complex $Z = X_* + u$, $|h^{-Z}| = h^{-X_* - \Re u} \leq h^{1/4} h^{-X_*}$, and $|I(Z)| \leq \int_0^\infty e^{-(X_* + \Re u)s} H(s) ds = I(X_* + \Re u) \leq I(X_* - \frac{1}{4})$. If also $X_* \geq X_2$, then $|I(Z)| \leq 2I(X_*)$, and (1) follows from (34).

(2). F is analytic for $\Re Z > -1$ with $F''(Z) = \psi'(Z + 1) = \sum_{j \geq 0} (Z + 1 + j)^{-2}$. For $\Re Z \geq X_* - \frac{1}{4} \geq 1$, $|F''(Z)| \leq \sum_{j \geq 0} (\Re Z + 1 + j)^{-2} \leq 1/\Re Z \leq 2/X_*$. Integrating twice along the segment from 0 to $u \in \Delta$, $|G(u) - pu| \leq |u|^2/X_* \leq |u|/(4pX_*) \leq p|u|/10$, the last step because $4p^2X_* \geq 12 > 10$. Thus $0.9p|u| \leq |G(u)| \leq 1.1p|u| \leq \frac{1.1}{4} < \frac{1}{2}$. For $|G| \leq 1$, $|1 - e^{-G} - G| \leq |G|^2 \sum_{j \geq 2} 1/j! \leq |G|^2$, so $|1 - e^{-G(u)}| \geq |G| - |G|^2 \geq |G|/2 \geq 0.45p|u|$. This vanishes only at $u = 0$, and the zero is simple because the derivative there is $G'(0) = p \neq 0$. So Q is analytic on Δ (its value at 0 is the removable one), and by (1), $|Q(u)| \leq |u| \cdot 2h^{1/4} S_*/(0.45p|u|) \leq 5h^{1/4} S_*/p$ for $u \neq 0$, hence also at $u = 0$.

(3). On the circle $|u| = r := 1/(8p)$, which lies inside Δ , $|\varepsilon Q(u)| \leq |\varepsilon| K S_*/p < r$ whenever $|\varepsilon| < R$. By Rouché's theorem $u + \varepsilon Q(u)$ has as many zeros in $|u| < r$ as u , namely one. The root $u(\varepsilon) = \frac{1}{2\pi i} \oint_{|u|=r} u \frac{1+\varepsilon Q'(u)}{u+\varepsilon Q(u)} du$ is analytic in ε for $|\varepsilon| < R$. The function Q is real on real u (there G and S are real), so for real ε the conjugate of the root is again a root in the disk, and uniqueness makes it real. With $\phi = -Q$, analytic on Δ , the root solves $u = \varepsilon \phi(u)$, and the Lagrange inversion theorem gives its Taylor coefficients as $\frac{1}{m!} \left[\frac{d^{m-1}}{du^{m-1}} \phi(u)^m \right]_{u=0}$, which are the terms of (40).

(4). If $R \geq 2$, then 1 lies in the disk of analyticity and $u(1)$ is real with $|u(1)| < 1/(8p)$. At $\varepsilon = 1$ the equation $u + Q(u) = 0$ with $u \neq 0$ is, after multiplying by $(1 - e^{-G(u)})/u$, the same as $S(X_* + u) = e^{-G(u)} - 1$, that is $e^{F(X_* + u)}(1 + S(X_* + u)) = e^{F(X_*)} = y$ by (36); and $u = 0$ is not a root, since $Q(0) = S_*/p \neq 0$. By (35) the real number $X_* + u(1)$ therefore solves $\mathcal{A}(X - 1) = y$. Since $X_* + u(1) - 1 > 3 - \frac{1}{4} - 1 \geq 1$ and $\mathcal{A}$ is strictly increasing on $[1, \infty)$ (Section 7), $X_* + u(1) = X(y)$. Finally $|u(\varepsilon)| < r$ on $|\varepsilon| < R$, so Cauchy's estimate gives $|u_m| \leq r R^{-m}$ for the m th Taylor coefficient, and for $R \geq 2$, $\sum_{m > M} |u_m| \leq 2r R^{-M-1} = \frac{1}{4p} (8K S_*)^{M+1}$. Take $X_1 \geq \max(3, X_2)$ so large that also $8K S(X_*) \leq \frac{1}{2}$ for $X_* \geq X_1$ (possible because $S(X) \rightarrow 0$); the constant K is absolute, so this bound is $O_M(S_*^{M+1}/p)$. $\square$

This is an interpolation-specific inverse theorem. Arbitrary functions agreeing with a_n at integers can have different exponentially small inverse terms. The spectral interpolation is distinguished here by the exact positive moment representation.

[write] **Added 5 October 2026, batch 101.** The interpolation dependence just stated is the general phenomenon of `p0:rem:three-differ` in the transseries-and-inversion volume: distinct admissible interpolations agree at the integers and have different inverses between them. The convergence step of the proof of Theorem 8.1 is written out in Proposition 8.2.

9 Exact integer recovery and threshold inversion

The next conclusion is stronger for integer inputs and requires no asymptotic onset. Define the envelope in the original size variable by

$$\mathcal{E}(x) = k\Gamma(x + 2)h^x, \quad x \geq 1. \quad (44)$$

It is strictly increasing.

Theorem 9.1 (Global rounding corollary). *For every integer $n \geq 1$,*

$$\mathcal{E}(n) < a_n < \mathcal{E}(n + 1/2). \quad (45)$$

Consequently, if $z = \mathcal{E}^{-1}(a_n)$ is the exact real inverse, then

$$n < z < n + 1/2, \quad \lfloor z \rfloor = n, \quad \text{nearestinteger}(z) = n. \quad (46)$$

Proof. The defining equation for h gives $4/3 < h < 3/2$. Thus $k > 5/18$ and $1/2 - k < 2/9$. By (19), for every $n \geq 1$,

$$0 < \frac{a_n}{\mathcal{E}(n)} - 1 = \frac{r_n}{kh^n} \leq \frac{1/2 - k}{kh^n} < \frac{3}{5}.$$

The ratio $\Gamma(x+1/2)/\Gamma(x)$ increases with $x > 0$, since its logarithmic derivative is $\psi(x+1/2) - \psi(x) > 0$. Therefore

$$\frac{\mathcal{E}(n+1/2)}{\mathcal{E}(n)} > \sqrt{\frac{4}{3}} \frac{\Gamma(7/2)}{\Gamma(3)} = \frac{15}{8} \sqrt{\frac{\pi}{3}} > \frac{15}{8} > \frac{8}{5}.$$

Combining the two bounds proves (45). Strict monotonicity gives (46). $\square$

For completeness, $4/3 < h < 3/2$ follows by evaluating $d(\rho) = 2 - \rho + \log(1 - \rho)$ at $\rho = 3/4$ and $2/3$: $d(3/4) = 5/4 - \log 4 < 0$ and $d(2/3) = 4/3 - \log 3 > 0$.

Corollary 9.2 (At most one exact threshold comparison). *For $y \geq \mathcal{E}(1)$ set $z = \mathcal{E}^{-1}(y)$ and let*

$$N(y) = \min\{n \geq 1 : a_n \geq y\}.$$

Then

$$\max\{1, \lfloor z - 1/2 \rfloor + 1\} \leq N(y) \leq \lceil z \rceil. \quad (47)$$

The interval contains at most two integers. If it contains two, comparing y with the sequence value at the smaller index determines $N(y)$.

Proof. If $n \leq z - 1/2$, then $a_n < \mathcal{E}(n + 1/2) \leq y$, so such an n cannot attain the threshold. If $n \geq z$, then $a_n > \mathcal{E}(n) \geq y$, so $\lceil z \rceil$ suffices. The bounds differ by at most one. The original recurrence makes (a_n) strictly increasing, or this follows directly from (45), since $\mathcal{E}(n + 1/2) < \mathcal{E}(n + 1)$. $\square$

Values $y < \mathcal{E}(1)$ have $N(y) = 1$. The simple rounding statement in Theorem 9.1 applies to exact sequence inputs; it must not be used as an arbitrary- y threshold rule. A numerical implementation should certify the envelope inverse interval before rounding near an integer or half-integer boundary.

[write] Added 5 October 2026, batch 101: relation to repository results. In the language of `p0:thm:staircase` (transseries-and-inversion volume), Theorem 9.1 supplies, for every $n \geq 1$, an approximation $x_* = \mathcal{E}^{-1}(a_n)$ of the interpolated inverse $\mathcal{A}^{-1}(a_n) = n$ with error $\eta < \frac{1}{2}$, which is the hypothesis of part (3) of that theorem. For arbitrary y , Corollary 9.2 replaces the separation condition of part (2), which can fail near integers, by a bracket containing at most two integers that one exact comparison resolves. What is specific here is that the error bound is explicit and global, from $n = 1$, by the elementary inequality (45). Compare `a126764-1convex-polyominoes`, whose rounding brackets (`lcp:ao:cor:brackets`) carry asymptotic existence constants without a finite starting index (note at the end of Section 1).

10 Non D finiteness and non P recursiveness

The closed form also determines the holonomic status of this counting sequence.

Theorem 10.1. *The function B is not D-finite over $\mathbb{C}(z)$. Neither (b_n) nor (a_n) is P-recursive over $\mathbb{C}[n]$.*

Proof. A D-finite function satisfies a nonzero linear differential equation with polynomial coefficients. All its analytic continuations can have finite-plane singularities only among the finitely many zeros of the leading polynomial.

Now use the logarithmic covering coordinate $w = \log(1 - z)$, so $z = 1 - e^w$. On this surface,

$$B(1 - e^w) = \frac{-(1 + e^w)w}{2(1 - e^w)(1 + e^w + w)}.$$

For every integer branch index j of Lambert W [7, Section 4.13], the denominator has a zero

$$w_j = -1 - W_j(e^{-1}), \quad z_j = 1 - W_j(e^{-1}).$$

These are infinitely many distinct projected points. At each such point, the numerator reduces to $w_j^2 \neq 0$, the derivative of $1 + e^w + w$ is $1 + W_j(e^{-1}) \neq 0$, and $dz/dw = -W_j(e^{-1}) \neq 0$. Also $1 - e^{w_j} \neq 0$. Thus each is a genuine simple pole on an analytic continuation of the original germ.

To make the continuation argument precise, paths in the w -plane can be chosen from a small nonzero point near $w = 0$ to each w_j , avoiding the discrete poles and the preimages of any proposed finite ODE singular set. Projection gives a continuation path in the z -plane. Choose a z_j outside that proposed finite set. An ODE solution must remain analytic at this ordinary point, contradicting the genuine pole. Therefore B is not D-finite.

The standard coefficient equivalence between D-finite series and P-recursive sequences then excludes P-recursive-ness of (b_n) . If (a_n) satisfied a polynomial-coefficient recurrence, substitute $a_{n+j} = (n+j+1)!b_{n+j}$ and divide by $(n+1)!$. Every factorial ratio is a polynomial in n , giving a P-recurrence for (b_n) , a contradiction. Finite initial modifications have no effect on this argument. $\square$

11 Reproducible validation

The source bundle includes a standalone Python validation program using exact rational arithmetic and high-precision quadrature. These checks support the algebra and indexing; the proofs above do not depend on data fitting.

11.1 An independent scalar recurrence from the generating function

Write $D(z) = 2 - z + \log(1 - z) = \sum d_j z^j$ and let the numerator of (5) be $\sum \nu_j z^j$. Then

$$d_0 = 2, \quad d_1 = -2, \quad d_j = -1/j \quad (j \geq 2), \quad \nu_n = \frac{1}{n+1} - \frac{1}{2n} \quad (n \geq 1).$$

Consequently

$$2b_n - 2b_{n-1} - \sum_{j=2}^n \frac{b_{n-j}}{j} = \frac{1}{n+1} - \frac{1}{2n}, \quad n \geq 1. \quad (48)$$

Starting from $b_0 = 1/2$, exact rational arithmetic verifies agreement between (48) and the original integer triangle through the configured validation limit. The program also checks the initial OEIS terms and the total spectral mass.

11.2 Residual and inverse checks

Residuals should be evaluated from the positive integral, or by exact integer counting with sufficient guard digits. Subtracting kh^n from b_n in ordinary floating point loses the secondary term rapidly.

The validation computes both independently and compares them. It also evaluates the inverse of the Gamma envelope at exact sequence values and checks the all-index half-step bound.

Independent code uses the integer triangle through $n = 1000$, with prefix sums in each row, and rational formal division of the stated generating function through $n = 250$. All coefficients agree exactly. The freshly retrieved OEIS b-file contains indices $1 \leq n \leq 200$; all 200 counts agree exactly. The script uses explicit exception-based checks, not Python `assert` statements. The source manifest records retrieval and SHA-256 hashes.

For $r_n = b_n - kh^n$, direct subtraction requires precision growing linearly with n . The table uses a scaled positive Laplace integral and compares it against exact counts with sufficiently increased working precision. The reported discrepancies are numerical observations, not certified errors. The first row uses the normalization $b_0 = 1/2$.

n	r_n from positive integral	Relative discrepancy
0	0.1580888684782	1.16×10^{-90}
1	0.02613398538975	6.76×10^{-93}
4	0.00518039343134	2.39×10^{-92}
100	$1.895125258712 \times 10^{-4}$	8.8×10^{-92}
1000	$1.020228640383 \times 10^{-5}$	8.12×10^{-92}

The total continuous mass is numerically 0.316177736956408984232149966021, agreeing with $1 - 2k$. Independent direct t -quadratures check the first few moments; 65- and 90-digit scaled quadratures agree to over 55 relative digits in the reported precision-stability tests. These are consistency checks only.

Let $R_n = 2(n+1)\log^2(n+1)r_n$, and let P_K be the logarithmic polynomial inside the square brackets truncated after L^{-K} . The following values are empirical evaluations of the exact positive integral and those polynomials. A fixed-order asymptotic expansion does not promise monotone improvement as more terms are added at modest n .

n	R_n	P_3	P_6
1000	0.974902401719	0.976684812798	0.976581527058
10^6	1.03111428094	1.03146229717	1.03110591297
10^{12}	1.02397377958	1.02400377395	1.02397369004
10^{24}	1.013754047	1.0137561217	1.01375404629
10^{96}	1.00373360566	1.00373361425	1.00373360566

The derived coefficients $C_{0,0}, \dots, C_{0,4}$ and $C_{1,0}, C_{1,1}$ were checked by exact symbolic algebra. The supplemental JSON also records $C_{j,k}$ for $0 \leq j \leq 2$, $0 \leq k \leq 6$ and mixed-sector tests.

For the exact Gamma envelope $\mathcal{E}(x) = k\Gamma(x+2)h^x$, the numerical inverse shifts below are positive. Every index $1 \leq n \leq 200$ passed the finite check $0 < \mathcal{E}^{-1}(a_n) - n < 1/2$. The all-index statement follows from the analytical bound in the text, not from this finite check.

n	$\mathcal{E}^{-1}(a_n) - n$
1	0.0426892402655284
4	0.00201614717486298
10	$1.04718232840828 \times 10^{-4}$
100	$7.50239646203492 \times 10^{-19}$

Forward rounding is a different operation and already fails at $n = 4$: $\mathcal{E}(4) = 151.378352788239\dots$, while $a_4 = 152$. Thus rounding $\mathcal{E}(4)$ to its nearest integer gives 151. The forward absolute discrepancy subsequently grows, despite the exponentially small relative discrepancy.

For the natural interpolation, direct high-precision solution about selected exact envelope inverses $X_* = 5, 10, 25, 50, 100$ validates the signs and scale of the first two inverse sectors. At $X_* = 100$ the inverse shift is approximately $-1.0551771299966891351 \times 10^{-18}$; including the second sector reduces the observed relative error from 2.25×10^{-18} to 6.44×10^{-36} . These diagnostics use 138 working decimal digits.

A cautionary test at $n = 500$ illustrates why fixed precision is unsuitable: 50-digit subtraction gives approximately -4.02×10^{21} instead of the positive residual $2.4464646945801111 \times 10^{-5}$. No negative residual, secondary exponent, or inverse correction should be inferred from such a cancellation-dominated computation.

[write] **Added 5 October 2026, batch 101: shipped files and intake reruns.** The validation program and its records are shipped beside this report: `code/verify_a196275.py`, `code/render_validation_tables.py`, `data/verification_results.json` and its python `-O` counterpart, `data/run_manifest.json`, `data/source_manifest.json`, the frozen b-file `data/oeis_b196275.txt` and the renderer’s raw output `data/validation_summary.tex` (identical to the text above except that it writes F for $\mathcal{E}$). The two exact-count tables (1,263,683 bytes each, byte-identical) are not shipped; the report’s README says how to regenerate them or retrieve them from the arrival commit. The b-file is OEIS data (table credited by OEIS to V. Kotěšovec, computed by A. J. Guttmann) under CC BY-SA 4.0, not under the repository’s MIT-0 licence. On copies of the delivered layout, the intake verified all 17 hashes of the delivered `manifest.json` and all 9 of `run_manifest.json`, reran the full normal-mode verifier (Python 3.14.4, SymPy 1.14.0, mpmath 1.3.0: all checks passed, in about 9 minutes on a loaded machine, not the “minute or two” the verifier’s README states), and reran the renderer; the outputs agree with the shipped ones except for line endings, the output file name and the Python version. The python `-O` run was not repeated. Independent intake scripts rechecked the triangle against the b-file ($n \leq 60$), the generating function in exact rationals, $\int_0^1 w = 1 - 2k$, the moments at $n = 0, 1, 2, 5, 10$, Theorem 9.1 for $1 \leq n \leq 200$, and the closed forms of $c_0, \dots, c_6$; all passed.

12 Literature scope and further research

The 2011 paper [1] obtains the original functional equation and gives the numerical conjecture addressed here. Tomás [2] provides the simpler triangular recurrence used as the exact input. Rinaldi and Socci [4] still describe the unrestricted column-convex equivalent as conjectural; their exact formula concerns the directed subclass. Duchi, Rinaldi and Socci [5] likewise retain the conjectural statement. Duchi’s later work [6] concerns doubly convex permutominoes and square permutations, rather than resolving this singly convex class.

The literature search on 2 October 2026 included the identifier A196275, the exact class name and row-convex or vertically-convex variants, grid-ogon terminology, the decimal growth constant, and combinations with Bernstein polynomials, transfer and integral operators, resolvents, spectra, and Lambert W . The checked author bibliography still linked the 2011 problem. No prior instance of the exact solution above was located. This is a bounded-search result, not a claim that all published, unpublished, or non-indexed work has been exhausted. Source dates were distinguished from search-engine crawl dates.

Several further directions follow naturally.

1. **Combinatorial explanation of the normalized generating function.** The Bernstein identity proves the formula analytically. A direct bijective or labeled-combinatorial derivation of the denominator $2 - z + \log(1 - z)$ could explain its simple structure more transparently.
2. **Refined shape statistics.** The spectral projection at h provides an explicit polynomial-profile

limit after Bernstein smoothing. Turning that into local asymptotics for the original coefficients $P_{n,p}$, particularly near the boundaries, requires a separate coefficient analysis.

3. **Uniform secondary asymptotics.** The present hierarchy is valid to every fixed order. Optimal truncation, late-term growth, and exponentially improved estimates within the logarithmic sector are additional questions.
4. **Efficient certified inversion.** The global half-step bound reduces threshold recovery to at most one exact count comparison. Interval arithmetic for the Gamma-envelope inverse could turn this into a robust implementation with explicit precision targets.
5. **Related counting recurrences.** The decisive feature was that multiplication and reflected cumulative sums close exactly in the Bernstein basis. Other growing triangular recurrences with this structure may admit fixed positive-operator models and similarly explicit moment asymptotics.

The established conclusions are the exact generating function and measure, non-P-recursiveness, the proved leading equivalent, the full mixed secondary expansion, the specified interpolation's convergent inverse sectors, and the global integer-recovery theorem. These should be distinguished from the further questions just listed.

12.1 Further questions and research

[write] **Added 5 October 2026, batch 101.** This subsection collects every statement of the source that it does not prove, following Vladimir's standing rule of 4 October 2026: each is stated as an open question with its source, the argument offered for it, and what is missing. Items 1–5 are the five directions listed above (Report 96, Section 12); item 6 collects the steps the source argues only in condensed form; item 7 the external inputs the intake did not check. The intake found no claim of the source false. The convergence step of Theorem 8.1, which the source also argues briefly, is now written out in Proposition 8.2.

1. **A combinatorial reason for the denominator.** Is there a bijective or labelled-combinatorial derivation of (5), in particular of the denominator $2 - z + \log(1 - z)$? The source derives it analytically, through the Bernstein embedding (Lemma 2.1) and the resolvent (Lemma 3.1). A remark of the write, not an answer: (48) has the renewal form $b_n = b_{n-1} + \sum_{j=2}^n b_{n-j}/(2j) + \frac{1}{2}\nu_n$, so a combinatorial model would have to explain the step weights 1 and $1/(2j)$, $j \geq 2$, and the inhomogeneous term $\nu_n = 1/(n+1) - 1/(2n)$. Missing: any such model.
2. **Shape statistics.** What are the local asymptotics of the triangle $P_{n,p}$ of (2), especially for p near 1 or near n ? The source says that the spectral projection at h gives a smoothed profile limit. The write makes this explicit (proof below): with

$$g_h(x) = \frac{2h-1}{h-1+x} - \log \frac{h-1+x}{h-x}, \quad 0 \leq x \leq 1,$$

one has $Tg_h = hg_h$, and for every $n \geq 1$, in $L^2[0, 1]$,

$$\left\| \frac{2}{n!} \sum_{p=1}^n P_{n,p} B_{p-1,n} - h^n \frac{\langle 1, g_h \rangle}{\|g_h\|^2} g_h \right\| \leq 1.$$

Proof. For real $\lambda > h$, Lemma 3.1 gives $(\lambda - T)g_\lambda = D(\lambda) \cdot 1$ with $g_\lambda = c/u - \log(u/v)$ the numerator of (15). For $\lambda \geq h > 1$ both $u = \lambda - 1 + x$ and $v = \lambda - x$ are at least $h - 1 > 0$ on $[0, 1]$, so $g_\lambda \rightarrow g_h$ uniformly as $\lambda \downarrow h$; T is bounded and $D(h) = 0$, so $(h - T)g_h = 0$. Also $g_h \neq 0$, since $\int_0^1 g_h = (2h - 1)\ell(h) > 0$ (proof of Lemma 3.1). By Proposition 3.2 and the measure (18), the spectrum of T is $\{h\} \cup [0, 1]$ with h a simple isolated eigenvalue. Let P be the orthogonal projection onto $\mathbb{C}g_h$. The self-adjoint operator T leaves $g_h^\perp$ invariant; the spectrum of

its restriction lies in that of T and does not contain the isolated point h , which would otherwise be an eigenvalue with an eigenvector orthogonal to g_h . Hence the restriction has norm at most 1, and $T^n 1 = h^n P 1 + T^n(1 - P)1$ with $\|T^n(1 - P)1\| \leq \|(1 - P)1\| \leq 1$. Lemma 2.1 identifies $T^n 1$ with the Bernstein sum. $\square$ (Consistently, $\|P 1\|^2 = 2k$ is the atom of (18).) Missing: passing from this L^2 statement about Bernstein-smoothed rows to the coefficients $P_{n,p}$ themselves, which needs a separate coefficient analysis, and the boundary behaviour.

3. **Uniform secondary asymptotics.** What are the optimal truncation, the late-term growth of c_j and $C_{j,k}$, and exponentially improved estimates within the logarithmic sector? Theorems 5.1 and 6.1 hold to every fixed order and claim no convergence. *Evidence* (the verifier’s README): at $n = 1000$ the normalized residual $2N \log^2 N r_n$ is $0.97490240\dots$, the sum through L^{-3} is $0.97668481\dots$ and the sum through L^{-4} is $0.96752333\dots$, so adding a term can worsen the approximation at modest n . Missing: any bound on the coefficients uniform in their order.
4. **Certified inversion.** Can interval arithmetic for the inverse of $\mathcal{E}$ turn Theorem 9.1 and Corollary 9.2 into an implementation with explicit precision targets? The theorem reduces threshold recovery to at most one exact comparison; the shipped checks (Section 11) evaluate $\mathcal{E}^{-1}(a_n)$ in arbitrary but uncertified precision, for $n \leq 200$. Missing: rigorous enclosures of h , k , $\log \Gamma$ and the inverse.
5. **Other triangular recurrences.** Which other growing triangular recurrences close exactly in the Bernstein basis, as (2) does under multiplication by $1 - x$ and the reflected cumulative sum, and do they admit fixed positive-operator models with explicit moment asymptotics? The source names the mechanism and no second example. Missing: a characterization, or further instances.
6. **Written-out tail estimates.** The proofs of Theorems 5.1 and 6.1 name their splits and the order of each piece but leave the bounds implicit (notes after each theorem). The intake checked them and believes them correct. Missing: explicit remainder bounds for the finite geometric expansions on the middle ranges, the gain of one power of L from taking imaginary parts, and the termwise integration of the expansion of S^{-m-1} . The corresponding step of Theorem 8.1 is done (Proposition 8.2).
7. **External inputs.** (a) The source states that [1] conjectured the equivalent $a_n \sim k(n+1)!h^n$ numerically, that [4] and [5] still call it conjectural, and that its bounded search of 2 October 2026 found no prior solution. The intake read none of these papers and made no search, so these statements are the source’s. (b) The recurrence (1)–(3) is proved by Tomás [2], not here; the intake checked only that it reproduces the 200 terms of the OEIS b-file. Missing: a reading of the four cited papers against the source’s summaries, and an independent literature search.

References

- [1] N. R. Beaton, F. Disanto, A. J. Guttmann, and S. Rinaldi. *On the enumeration of column-convex permutominoes*. DMTCS Proceedings AO (FPSAC 2011), 111–122. doi:10.46298/dmtcs.2895. Author PDF.
- [2] A. P. Tomás. *On the Enumeration of Permutominoes*. FCT 2015, Section 5.2. doi:10.1007/978-3-319-22177-9_4. Author PDF.
- [3] OEIS Foundation. *A196275: Number of column-convex permutominoes of size n*. <https://oeis.org/A196275>. Accessed 2 October 2026.
- [4] S. Rinaldi and D. Socci. *About Half Permutations*. Electronic Journal of Combinatorics 21(1) (2014), P1.35. <https://www.combinatorics.org/ojs/index.php/eljc/article/download/v21i1p35/pdf/>.
- [5] E. Duchi, S. Rinaldi, and D. Socci. *3-dimensional polygons determined by permutations*. Journal of Combinatorics 9(1) (2018), 57–94. https://www.irif.fr/~duchi/JOC_09_01_A05.pdf.
- [6] E. Duchi. *A code for square permutations and convex permutominoes* (2019). <https://arxiv.org/abs/1904.02691>.

- [7] NIST Digital Library of Mathematical Functions. Sections 4.13 and 5.11, *Lambert W Function* and *Asymptotic Expansions*. <https://dlmf.nist.gov/4.13> and <https://dlmf.nist.gov/5.11>.