

Uniform rooted trees and binary partitions

Ratio nonconvergence and all fixed inverse log orders

Report215

4 October 2026

Abstract

Let $a_n = A003238(n)$ be the uniform rooted-tree count and $S_n = \sum_{j=1}^n a_j = A003318(n)$. We prove that neither $a_{n+1}/A018819(n)$ nor $S_{n+1}/A000123(n)$ converges. A positive dyadic identity, a cut positive operator, and exact directed-integer evaluation of 2^{24} recurrence coefficients give a contradiction to a hypothetical common radial limit. A positive Abelian argument transfers this contradiction to both ratios without using coefficient asymptotics. A separate midpoint argument certifies actual profile oscillation greater than 0.00001839189788, also a lower bound for each sequence ratio's limit-superior minus limit-inferior. Separately, an analytic dyadically periodic amplitude and an integrated positive-prefix bound yield individual and summatory expansions to every fixed inverse-logarithmic order, with relative remainder $O_J((\log n)^{-J-1})$. A finite Gaussian-moment prescription supplies every correction. Smooth inverse centers have uncertainty $O_J(x/(\log x)^{J+2})$, which need not shrink absolutely. Erdős and Loxton already established periodic summatory leading behavior and binary-prefix approximation in 1979; the cumulative ratio question appears explicitly on their page 328. Our source comparison is bounded, not a worldwide priority claim. The reproducibility archive contains the analytic proof, standard-library exact arithmetic, compact receipts, and deterministic build instructions.

[write] **Status and trust boundary, added 7 October 2026, batch 112.** This is bundle Report 215 of an external AI-assisted research session, filed in ProveIt as the single-source research report **a003238-uniform-trees-binary-partitions** (provenance, the sources as the write read them, what was checked, relation to the repository, the collected non-claims and the reading conventions are in Section 1.1). It is unrefereed and not formalized. Its author line and PDF author field are empty, and the title block reads “Report215”: it names no person or tool. *What rests on what.* Theorem 1.1, which answers in the negative the question Erdős and Loxton left open in 1979, and Theorem 7.1 are *computer-assisted*: their proofs reduce, by the analytic arguments of Sections 3–7, to finite inequalities that Python programs certify in exact integer arithmetic and 256-bit directed interval arithmetic, including the exact computation of 2^{24} recurrence coefficients. The write reran every step of that certificate except the 2^{24} -coefficient radial sieve (about 1.6 GiB of memory), whose receipt is taken as delivered. Theorem 1.2 and Theorem 12.1 have conventional proofs that use no computation. The write adds Remarks 1.3 and 12.2 (the OEIS entries quoted, and Cloitre’s 2004 conjecture in A003238 refuted with proof; the inverses against the transseries volume) and dated notes, among them a confirmation of the Erdős–Loxton citations from the paper itself. No statement, proof or number of the source was changed.

Contents

1 The exact sequences and the results

3

1.1	Provenance, sources and reading conventions	5
2	What the primary sources already establish	7
3	Positive radial identities and a global bound	8
4	The cut operator and every analytic tail	10
4.1	Power majorants for the operator	10
4.2	The exterior terms	11
4.3	Finite recursion and its omitted terms	12
5	The finite contradiction	12
6	Abelian transfer to the two sequence ratios	13
7	A separate quantitative oscillation corollary	14
8	The total profile is holomorphic	15
9	Saddle parameters and the finite correction rule	16
10	An integrated bound for the whole circle	17
10.1	Short prefixes	18
10.2	Long prefixes at intermediate angles	18
10.3	The outer angles and dyadic secondary peaks	18
11	Local expansion and its remainder	19
12	Inverse centers and integer thresholds	20
12.1	Explicit leading inverse scales	21
13	Exact arithmetic and the reproduction contract	22
13.1	Arithmetic rules	22
13.2	Checks and their limits	22
13.3	Rebuilding	23
14	Further directions	23

1 The exact sequences and the results

All logarithms are natural. Define $a_1 = 1$ and, for $n \geq 1$,

$$a_{n+1} = \sum_{d|n} a_d, \quad S_n = \sum_{j=1}^n a_j, \quad S_0 = 0. \quad (1)$$

These are OEIS A003238 and A003318 with index starting at one [1, 2]. The first terms are

$$(a_1, \dots, a_{10}) = (1, 1, 2, 3, 5, 6, 10, 11, 16, 19), \quad (S_1, \dots, S_{10}) = (1, 2, 4, 7, 12, 18, 28, 39, 55, 74).$$

The recurrence can serve as the definition throughout this report, so no alternative convention for planted roots enters the calculations. In the usual tree interpretation, a uniform rooted tree is built recursively from a root and a positive number of isomorphic copies of a smaller uniform rooted tree, together with the one-vertex tree. A tree with $n + 1$ vertices has k copies of a d -vertex tree attached to its root, where $kd = n$; this gives (1). Equivalently, vertices at a common distance from the root have equal numbers of children.

Write

$$q(n) = a_{n+1}, \quad Q(n) = \sum_{j=0}^n q(j) = S_{n+1} \quad (n \geq 0). \quad (2)$$

The numbers $q(n)$ count partitions of n into positive parts arranged in a divisibility chain, repetitions allowed, including the empty partition when $n = 0$. To see the recurrence directly, divide all parts by the smallest part d and remove one resulting part equal to one. This gives a chain partition of $n/d - 1$ and hence $q(n) = \sum_{d|n} q(n/d - 1)$. Relabeling divisors yields (1).

Let β_n count partitions of n into powers of two, with $\beta_0 = 1$, and put $B_n = \sum_{j=0}^n \beta_j$. Then

$$\mathcal{B}(z) = \sum_{n \geq 0} \beta_n z^n = \prod_{j \geq 0} (1 - z^{2^j})^{-1}, \quad \beta_n = \text{A018819}(n), \quad B_n = \text{A000123}(n). \quad (3)$$

These conventions agree with the displayed binary entries [3]. In particular $B_n = \beta_{2n}$; A000123 is not the individual binary-partition count at the same argument. This identity follows either from the product or from $\beta_{2n+1} = \beta_{2n}$ and $\beta_{2n} = \beta_{2n-1} + \beta_n$.

Theorem 1.1 (Computer-assisted ratio nonconvergence). *Neither of the positive sequence ratios*

$$\frac{a_{n+1}}{\beta_n} = \frac{\text{A003238}(n+1)}{\text{A018819}(n)}, \quad \frac{S_{n+1}}{B_n} = \frac{\text{A003318}(n+1)}{\text{A000123}(n)}$$

has a limit as $n \rightarrow \infty$, even as an extended real number.

The proof in Sections 3–6 is independent of the all-order saddle theorem. Its computational component evaluates exact integer coefficients and includes every analytic truncation error. The two displayed intervals later in the report constrain a *hypothetical common constant*; they do not enclose two actual values of the limiting profile.

For the asymptotic result, introduce

$$\begin{aligned} F(z) &= \sum_{n \geq 1} a_n z^n, \quad f(t) = F(e^{-t}), \quad B(t) = \mathcal{B}(e^{-t}), \quad h = \log 2, \quad \rho = \frac{\log(3/2)}{h}, \\ G(t) &= \frac{e^t f(t)}{B(t)}, \quad H(t) = \frac{e^{-t} f(t)}{B(t)} = e^{-2t} G(t). \end{aligned} \quad (4)$$

The radial $B(t)$ is distinct from the cumulative sequence B_n . For $k \geq 1$ put

$$b_k(t) = (-1)^k \frac{d^k}{dt^k} \log B(t). \quad (5)$$

There is a unique $t = t_N > 0$ satisfying $b_1(t) = N$. For coefficients indexed by n , always use $N = n + 1$, and write $L = \log(1/t)$ and $v = b_2(t)$.

Theorem 1.2 (Every fixed inverse-logarithmic order). *There is a positive function $\mathcal{P}$ on $(0, \infty)$, holomorphic in a complex sector about this ray, satisfying $\mathcal{P}(2t) = \mathcal{P}(t)$. It is nonconstant by Theorem 1.1's radial certificate. For every fixed integer $J \geq 0$,*

$$a_n = \frac{e^{(n+1)t} B(t)}{\sqrt{2\pi v}} \left\{ \sum_{r=0}^J \mathcal{A}_r[\mathcal{P}](t) + O_J(L^{-J-1}) \right\}, \quad (6)$$

$$S_n = \frac{e^{(n+1)t} B(t)}{\sqrt{2\pi v}} \left\{ \sum_{r=0}^J \mathcal{A}_r[\mathcal{P}/t](t) + O_J(t^{-1} L^{-J-1}) \right\}. \quad (7)$$

Here $t = t_{n+1}$, $\mathcal{A}_0[P] = P$, and the finite prescription (50) defines all corrections. For each fixed r ,

$$\mathcal{A}_r[\mathcal{P}](t) = O_r(L^{-r}), \quad \mathcal{A}_r[\mathcal{P}/t](t) = O_r(t^{-1} L^{-r}).$$

The errors in (6) and (7) are relative errors of order L^{-J-1} after dividing by their respective leading brackets. In particular,

$$S_n \sim \frac{a_n}{t_{n+1}} \sim \frac{hn}{\log n} a_n. \quad (8)$$

The notation $\mathcal{A}_r[\mathcal{P}/t]$ means to differentiate the complete amplitude $t \mapsto \mathcal{P}(t)/t$ in the prescription. It does not mean $\mathcal{A}_r[\mathcal{P}]/t$. The same saddle is used for both formulas, rather than hiding a shift in a newly chosen saddle.

All orders in this report are fixed before $n \rightarrow \infty$. No convergence of the infinite correction series, effective onset, growing-order uniformity, or expansion in every power of $1/n$ is claimed. The implicit constants are existential, except in the separate finite nonconvergence certificate. Section 12 states the precise inverse consequences and the uncertainty attached to integer thresholds.

Remark 1.3 ([write] The OEIS entries quoted, and a refuted conjecture, 7 October 2026). The write read the four entries in the internal format on 7 October 2026. *A003238* (revision #133, 30 May 2026; N. J. A. Sloane), “Number of rooted trees with n vertices in which vertices at the same level have the same degree.”, offset 1, b-file $n \leq 10000$ (Robert Israel). Its formula section contains

“Conjecture: $\log(a(n))$ is asymptotic to $c \cdot \log(n)^2$ where $0.4 < c < 0.5$ - __Benoit Cloitre__, Apr 13 2004”,

and a comment by Andrei Zabolotskii (22 January 2017) reads: “According to the MathOverflow link, $\log(a(n)) \sim \log(n)^2 / \log(4)$, and a more precise asymptotic expansion is similar to that of *A018819* and hence *A000123*, so the conjecture in the Formula section is partly correct.” *A003318* (#60, 30 May 2026), “ $a(n+1) = 1 + a(\text{floor}(n/1)) + a(\text{floor}(n/2)) + \dots + a(\text{floor}(n/n))$.”, offset 1, b-file $n \leq 1000$; *A003238*'s formula section states that its partial sums are *A003318*. *A018819* (#219, 3 August 2026), “Binary partition function: number of partitions of n into powers of 2.”, offset 0. *A000123* (#324, 23 July 2026), “Number of binary partitions: number of partitions of $2n$ into powers of 2.”, offset 0.

Cloitre's conjecture is false as stated. The asymptotic form is right, but the constant is $c = 1/(2\log 2) = 0.72134752\dots$, outside $(0.4, 0.5)$. Proof: by (71), $\log a_n = L^2/(2h) + O(L)$, and by (49), $L = \log n + O(\log \log n)$; hence $\log a_n / \log^2 n \rightarrow 1/(2h) = 1/(2\log 2)$. (Independently of this report, the same follows from Erdős–Loxton's Lemma 1, $\beta_n \leq a_{n+1} \leq C\beta_n$, and the classical $\log \beta_n \sim \log^2 n / (2\log 2)$ of Mahler and de Bruijn; it is also the value $1/\log 4$ in Zabolotskii's comment.) The finite data explain the conjecture: with the write's own sieve, $\log a_n / \log^2 n$ is 0.486, 0.479, 0.489 and 0.493 at $n = 10^2, 10^3, 10^4$ and $2 \cdot 10^4$, inside $(0.4, 0.5)$, and approaches 0.7213 only at the speed of the $O(L)$ and $\log \log n$ terms. The entry's word “partly correct” refers to this situation. No OEIS edit is part of this work.

Checks. The write recomputed a_n by the ascending divisor sieve of Section 5 for $n \leq 20000$, S_n , β_n from its recurrence, and B_n : all 10000 b-file terms of A003238, all 1000 of A003318, all b-file terms of A018819 with $n \leq 1000$ and of A000123 with $n \leq 20000$ agree, as do the first ten values printed above, the identity $B_n = \beta_{2n}$ for $n \leq 20000$, and the 48 terms of each shipped fixture (data/code-b003238.txt, data/code-b003318.txt).

1.1 Provenance, sources and reading conventions

[write] **Added 7 October 2026, batch 112.** *Provenance.* This report is Report 215 of the session bundle that ProveIt's docs/incoming intake processed as batch 112: the archive `Report215-reproducibility.zip` (601,832 bytes, 37 files in a wrapper directory `Report215/`), arrival commit 60f54ea06, placed in f79c9bef1. The text is the delivered `Report215.tex` (794 lines, dated 4 October 2026), shipped as `article.tex`. The programs are shipped under `code/` (the root `reproduce.py` and `test_package_guards.py` beside them), the receipts as `data/code-*.json`, the root `PACKAGE_GUARD_CHECKS.json` and `requirements.txt` under `data/`, the two 48-term OEIS fixtures as `data/code-b003238.txt` and `data/code-b003318.txt`, and the source ledger `SOURCES.txt`, the inventory `SOURCE_FILES.txt` and the code note `code/README.txt` (as `code-README.txt`) at the top level. Not shipped, and retrievable from the arrival commit: the delivered PDF, the pure checksum manifest `MANIFEST.json` (36 entries, verified at the write) and the delivered `README.txt`, replaced by the report `README`. The package records no ProveIt commit; it carries no “prepared for private review” line and no personal data. It is a single-source report, so no merge choice was made. The write prefixed the delivered labels with `urt:` and updated the references to them; it added this subsection, the notes marked [write] and Remarks 1.3 and 12.2, each the last statement of its section, and inserted nothing before a delivered display or statement, so every theorem, section and equation of the source keeps its delivered number.

The sources, as the write read them (7 October 2026): OEIS A003238, A003318, A018819 and A000123 in the internal format with their b-files (Remark 1.3); Erdős–Loxton [4] (the published PDF, freely served by Cambridge University Press), Section 3: Lemma 1, Theorem 3 on p. 324 and the last paragraph of Section 3 on p. 328 (the note at the end of Section 2); the transseries volume `Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/transseries_and_inversion.tex` (p0:def:model, p0:def:three-inverses, p0:thm:staircase, plt:def:lw-monomial-log-datum, plt:thm:lw-template). Not read by the write: the other cited works; the source's account of them (Section 2) is reported, not confirmed.

What was checked at intake. The batch-112 dossier read the manuscript, found no demonstrably wrong claim, and reran the delivered suite on a copy except the 2^{24} -coefficient radial sieve, whose regression values it confirmed with its own sieve to 10^6 ; the only failures were Windows artifacts (carriage returns change the hashes of regenerated receipts). *What the write checked.* It read every proof and found no error. On a fresh copy it reran, with Python 3.14.4, `certify_global_bound.py` (60 s), `operator_constants.py`, `certify_operator.py` (203 s), `combine_certificate.py`, `verify_oscillation.py`, `verify_finite_models.py`, `verify_exact.py`, `verify_saddle_engine.py` and `verify_arithmetic.py`: the global-bound, operator-constant and operator-value receipts are byte-identical to the delivered ones after removing carriage returns, the oscillation, exact and saddle receipts are equal as JSON (key order differs),

and the combined nonconstancy receipt differs only in the SHA-256 hashes it records for its inputs, which change with the line ends. `certify_radial_values.py` was not run (about 1.6 GiB; the machine is shared), so the radial enclosures enter `combine_certificate.py` as delivered. It confirmed against the receipts that the printed intervals I_1, I_2 of Proposition 5.1 are outward roundings of the 256-bit endpoints, that the exact gap (37) is 0.00001847273026448250..., that the global bound's upper endpoint is the printed 12.790169983612993410799254364259, and that the oscillation bound's exact interval is the one printed in Section 7; it recomputed in exact rationals $2g/(R_1 + R_2) = 821010233977/44639777749800000$ and the displayed margin, $C_4 = 499968$, $x_* < 1$ in (18), $3^{200} < 2^{317}$, $Mt_1/2 = 36$ and $Mt_2/2 = 48$; and it rederived the leading inverse constants $2/(e\sqrt{h})$ and $1/(e\sqrt{h})$ of (73) from (71)–(72) and (49). Finite computations are observations, except the certificate steps, which are the source's computer-assisted proof.

Relation to the repository. Before batch 112 the repository named A003238 and A000123 only as targets rejected by the source audits of two other reports (`a047874-long-increasing-subsequences` and `a126348-stable-hilbert-series`, in the same directory). The nearest reports are in the same directory: `a055779-labeled-fat-trees`, `a242375-many-color-rooted-trees` and `a244407-high-outdegree-rooted-trees` (batch 112; other tree families). They share no result with this report, so no reciprocal note is proposed. No Lean or Rocq file treats these sequences; no statement of this report is formalized, and its place in the collection confers no formal status. The inverses are compared with the transseries volume in Remark 12.2.

[write] **What is not claimed, collected from the source.** The periodic summatory leading asymptotic, the binary comparison and the binary-prefix approximation are Erdős and Loxton's; dyadic profiles and all-order saddle polynomials are not new methods (Mahler, de Bruijn, Dumas–Flajolet, Banderier–Hwang–Ravelomanana–Zacharovas credited); the source comparison is bounded, with Pennington, Kato–McLeod and the final 2026 version of the distinct-chain-partition preprint not read, and no worldwide priority claim. The conditional intervals I_1, I_2 constrain a hypothetical common constant: they are not enclosures of $\mathcal{P}(t_1)$, $\mathcal{P}(t_2)$, and their gap is not an oscillation bound (that is the separate Theorem 7.1); rounding that bound up to 0.0000183919 is not justified. All orders are fixed; no convergence of the correction series, effective onset, growing-order uniformity or expansion in powers of $1/n$; the asymptotic constants are existential; the inverse error is relative and its absolute width need not shrink, and no exact eventual ceiling formula is given; the 48-term fixtures are not full b-files; finite model and saddle-engine checks validate algebra, not the analytic remainder; byte identity of the PDF and ZIP only with the recorded toolchain. *The write adds:* its checks are exact or floating computations; it did not rerun the radial sieve; its transseries remark claims no novelty for any inversion.

Reading conventions (letters the source uses in more than one sense, with the tempting false readings; no symbol was renamed).

Symbol	Meanings
$B, \mathcal{B}, \mathcal{B}$	$B(t) = \mathcal{B}(e^{-t})$ the radial binary product; $\mathcal{B}_n = \text{A000123}(n)$ the cumulative binary count; B_3 the alternating exterior expression (28). <i>False reading:</i> $B(t)$ is not $\mathcal{B}_n$ (the source says so). Erdős–Loxton’s $B(x)$ is $\mathcal{B}_{\lfloor x \rfloor}$.
Q, q	$q(n) = a_{n+1}$ and $Q(n) = S_{n+1}$; q and Q also Erdős–Loxton’s $q(n)$, $Q(x)$; q_* an additive error in (18); q a factor parameter in Section 13.
D	$D(a)$ the power-majorant constants; $D(z)$ the positive prefix remainder (52); D an even degree in (50); $D = (0, u_0]$ the operator domain in Section 7.
A	$\mathcal{A}_r[P]$ the correction prescription; $A_j(z)$ the binary prefix products; $A_3 = I - T + T^2 - T^3$.
H, G, R	$H(t) = e^{-t} f(t)/B(t)$, $G = e^{2t} H$, $R(t) = H(t) - H(2t)$; R_i the ratios U_3/S_3 in Theorem 7.1.
S	$S_n = \text{A003318}(n)$; $S_3 = 1 - T1 + T^21 - T^31$. <i>False reading:</i> S_3 is not the third partial sum.
M, C	$M = 2^{24}$ the coefficient cutoff; M a running bound in the proof of Proposition 3.2; $M = \sup \mathcal{P}$ in Theorem 7.1; C the hypothetical common constant and generic constants C_r, C_b, C_J .
L, t, v	$L = \log(1/t)$, $t = t_N$ the saddle with $b_1(t) = N$, $v = b_2(t)$.
h, ρ, σ	$h = \log 2$; $\rho = \log(3/2)/\log 2$; $\sigma = \rho - d_0$.
c	Cloitre’s conjectured constant (Remark 1.3); generic constants c, c_ε .

2 What the primary sources already establish

Erdős and Loxton [4], Section 3, study exactly $q(n)$ and $Q(x) = S_{\lfloor x \rfloor + 1}$. Their Lemma 1 gives binary comparison $\beta_n \leq q(n) \leq C\beta_n$. Their Lemma 2 constructs positive binary-recurrence approximants with uniform multiplicative control. Their Theorem 3, page 324, establishes a periodic summatory logarithmic asymptotic. In their notation its form is

$$\begin{aligned} \log Q(x) = & \frac{\log^2(x/\log x)}{2h} + \left(\frac{1}{2} + \frac{1 + \log h}{h} \right) \log x \\ & - \left(1 + \frac{\log h}{h} \right) \log \log x + V \left(\frac{\log x - \log \log x}{h} \right) + o(1), \end{aligned} \quad (9)$$

where V is one-periodic. Their final paragraph of Section 3, page 328, explicitly leaves the convergence of $Q(x)/\sum_{j \leq x} \beta_j$ unresolved. Theorem 1.1 gives a negative answer for these exact conventions. Equation (9), the existence of a periodic leading shape, and binary-prefix approximation are prior results, not conclusions first proposed here.

Gol’dberg and Livshits [5] identify the size of a minimal universal tree with the relevant uniform-tree partial sum and establish its logarithmic leading growth. Gati, Harary, and Robinson [6], Section 3, connect the chain-partition and tree descriptions and reproduce the Erdős–Loxton summatory formula. Harary and Robinson [7], equation (36), give the matching generating equation in a planted convention with one extra root. The recurrence (1) fixes the convention used here.

The methods also have substantial predecessors. Mahler [8] and de Bruijn [9] treat binary or power-partition growth and periodic corrections. Dumas and Flajolet [10] give complete periodic coefficient expansions for a cyclotomic Mahler-product family and analyze multiple saddles. Banderier, Hwang, Ravelomanana, and Zacharovas [11] obtain inverse-logarithmic expansions with periodic coefficients for their binomially weighted recurrence. Those exact hypotheses do not identify their objects with (1). These works nevertheless preclude treating generic dyadic profiles or all-order saddle polynomials as new methods.

The source comparison for this report additionally inspected the cited works of Chung–Coppersmith–Graham, Kochkarev, Read, and the 2025 distinct-chain-partition preprint, together with the actual nearby ProveIt source listed in `SOURCES.txt`. In the inspected material, no theorem supplied either the present ratio nonconvergence or the complete fixed-order individual and summatory expansion for this exact recurrence. This is a bounded finding about those sources. It does not exclude uninspected later work, including a final 2026 version of the distinct-chain-partition paper or full original results of Pennington and Kato–McLeod that were not independently read. The proposed mathematical sharpening is the exact-target coefficient and inverse theorem with its quantified remainder, and the separately certified nonconvergence result. No worldwide priority assertion is made.

[write] The Erdős–Loxton citations, checked 7 October 2026. The write read the published paper [4]. Section 3 defines $q(n)$ by (8), the binary-partition count $b(n)$ by (9), $B(x) = \sum_{n \leq x} b(n) = b(2[x])$ and the sum function $Q(x)$ of q ; Lemma 1 is $b(n) \leq q(n) \leq cb(n)$; Theorem 3 is on p. 324; and the last paragraph of Section 3, on p. 328, says that their methods could bound the oscillation of $Q(x)/B(x)$, but “we have not been able to decide whether or not $Q(x)/B(x)$ approaches a limit”. With $B(n) = \mathbf{B}_n$ and $Q(n) = S_{n+1}$, this is exactly the second ratio of Theorem 1.1, so the citations above are accurate and the theorem answers that question in the negative, by a computer-assisted proof. Erdős and Loxton anticipated upper bounds for the oscillation; Theorem 7.1 gives a lower bound.

3 Positive radial identities and a global bound

Since a chain partition is an ordinary partition, $q(n) \leq p(n)$, where p is the ordinary partition function. Thus F is analytic in $|z| < 1$. Summing (1) gives the normally convergent identity

$$F(z) = \frac{z}{1-z} \left(1 + \sum_{k \geq 2} F(z^k) \right). \quad (10)$$

For example, direct expansion of the right side assigns to z^{n+1} the sum of a_d over all divisors d of n . The binary product has

$$B(t) = \frac{B(2t)}{1 - e^{-t}}. \quad (11)$$

Combining (10) and (11) yields

$$G(t) = \frac{1}{B(2t)} + e^{-2t} G(2t) + \sum_{k \geq 3} \frac{e^{-kt} B(kt)}{B(2t)} G(kt), \quad (12)$$

$$H(t) - H(2t) = R(t) := \frac{e^{-2t}}{B(2t)} \left(1 + \sum_{k \geq 3} f(kt) \right) > 0. \quad (13)$$

Lemma 3.1 (Product-ratio majorant). *For $k \geq 3$ and $kt \leq 1$, let $\lambda_k = \log_2 k - 1$. Then*

$$\frac{B(kt)}{B(2t)} \leq e^2 (kt)^{\lambda_k}. \quad (14)$$

For $0 < t \leq 1/16$,

$$W(t) := \sum_{k \geq 3} \frac{e^{-kt} B(kt)}{B(2t)} \leq 15t^\rho + 24t, \quad \frac{1}{B(2t)} \leq 8t^2. \quad (15)$$

Proof. For $u > 0$,

$$\frac{1 - e^{-2u}}{1 - e^{-ku}} \leq \frac{2}{k} e^{ku}.$$

Multiply through the last dyadic $u = 2^j t$ for which $ku \leq 1$. The sum of the retained ku is at most 2, and if J is the last index, then $(2/k)^{J+1} \leq (kt)^{\lambda_k}$. Every subsequent factor ratio is at most one. This proves (14).

The $k = 3$ term is at most $e^2 3^\rho t^\rho < 15t^\rho$. One elementary verification is $\rho < 3/5$, $3^{3/5} < 2$, and $e^2 < 15/2$. For $4 \leq k \leq 7$, monotonicity of B and cancellation in (11) give a ratio at most $1 - e^{-2t} \leq 2t$, so the total is at most $8t$. For $k \geq 8$ the first two canceled factors give a ratio at most $8t^2$; its exponential sum is at most $(e^t - 1)^{-1} \leq 1/t$. Thus $16t$ already bounds the terms with $k \geq 4$; $24t$ is a convenient enlargement. Finally, retain the first two factors of $1/B(2t)$. $\square$

Proposition 3.2 (Certified uniform bound). *For every real $t > 0$, $0 < G(t) < 14$ and $0 < H(t) < 14$.*

Proof. The finite portion and its tails are specified here so that the computation is a finite proof obligation. From

$$\log \prod_{m \geq 1} (1 - e^{-ms})^{-1} = \sum_{r \geq 1} \frac{1}{r(e^{rs} - 1)} < \frac{2}{s}$$

and positivity, $p(r) \leq \exp(2\sqrt{2r}) \leq \exp(3\sqrt{r+1})$. Hence $a_n \leq \exp(3\sqrt{n})$. For $t \geq 1/16$, concavity of the square root at $n = 10000$ bounds the omitted polynomial tail by

$$\sum_{n > 10000} a_n e^{-nt} < \frac{e^{-325}}{e^{19/400} - 1} < 33e^{-325} < 2^{-319}. \quad (16)$$

The last inequality already follows from $e > 2$ and $33 < 64$.

Compute a_n through 10000. On each interval $u \leq t \leq v$ in the grid $u_0 = 1/16$, $u_{i+1} = \min(4, 65u_i/64)$, use the monotonicity of $e^t f(t) = \sum a_n e^{-(n-1)t}$ and of $B(t)$ to obtain

$$G(t) \leq \frac{e^u f(u)}{B(v)}.$$

Use (16) for $f(u)$. For $t \geq 4$, use $G(t) \leq e^4 f(4)$ and $B(t) \geq 1$. Directed evaluation certifies $G < 4$ throughout $[1/16, \infty)$.

Suppose $G \leq M$ has been covered for $t \geq b$, and extend to $a \leq t \leq b = 2a$. Every kt with $k \geq 2$ is already covered. Each ratio $B(kt)/B(2t)$ increases with t : differentiate individual factor ratios and use the decrease of $x/(e^{xu} - 1)$ in x . Therefore (12) is bounded by

$$\frac{1}{B(2b)} + (e^{-2a} + W_{a,b}) M, \quad (17)$$

where the finite upper bound is

$$W_{a,b} = \sum_{k=3}^{63} \frac{e^{-ka} B(kb)}{B(2b)} + \frac{B(64b)e^{-64a}}{B(2b)(1 - e^{-a})}.$$

Take the maximum of M and (17) at each step. After 22 such bands, the lower endpoint is $t_* = 2^{-26}$. For the remaining bands, (15) bounds the accumulated multiplicative exponent and additive error by

$$x_* = 45(2/3)^{26} + 48 \cdot 2^{-26}, \quad q_* = (32/3)2^{-52}. \quad (18)$$

Indeed $t_*^\rho = (2/3)^{26}$ and summing the successive geometric scales gives these constants. If M_* is the finite-band bound, every remaining band has bound at most

$$(M_* + q_*)e^{x_*} \leq \frac{M_* + q_*}{1 - x_*}.$$

The executable `certify_global_bound.py` verifies $x_* < 1$ and gives the outward upper endpoint

$$12.790169983612993410799254364259 < 14.$$

The arithmetic rules and replay procedure appear in Section 13. Finally $H = e^{-2t}G$. $\square$

Equations (13) and (15) now give $R(t) = O(t^\rho)$. In detail $f(kt) = e^{-kt}B(kt)G(kt)$ and $G < 14$, so $R \leq 1/B(2t) + 14W(t)$. Consequently

$$\mathcal{P}(t) := \lim_{m \rightarrow \infty} H(t/2^m) = H(t) + \sum_{j \geq 1} R(t/2^j) \quad (19)$$

converges uniformly on compact real intervals, is continuous, and satisfies $\mathcal{P}(2t) = \mathcal{P}(t)$ and

$$H(t) = \mathcal{P}(t) + O(t^\rho). \quad (20)$$

It is bounded below by a positive constant: $H(t) \geq H(2t)$ lets one move every small t into a fixed compact dyadic annulus, where H has a positive minimum. The same compact-annulus argument and (19) give a positive lower bound for $\mathcal{P}$. This construction alone does not prove nonconstancy.

4 The cut operator and every analytic tail

Set $u_0 = 2^{-10}$. For bounded nonnegative v on $(0, u_0]$, define

$$(Tv)(t) = \sum_{j \geq 1} \sum_{\substack{k \geq 3 \\ kt/2^j \leq u_0}} \frac{e^{(k-2)t/2^j} B(kt/2^j)}{B(2t/2^j)} v(kt/2^j). \quad (21)$$

The exponent in this kernel is positive. It comes from substituting $f(ku) = e^{ku}B(ku)H(ku)$ into (13); changing its sign would change the certificate. The domain includes equality $ku = u_0$.

Let $b(t)$ be the remaining positive part of $\sum_{j \geq 1} R(t/2^j)$, comprising the inhomogeneous 1 and all terms with $kt/2^j > u_0$. If $H(t)$ had a limit C as $t \downarrow 0$, telescoping (13) would give

$$H + TH + b = C \quad \text{on } (0, u_0]. \quad (22)$$

The same identity follows from the hypothesis that $\mathcal{P}$ is the constant C . The next estimates establish convergence of the operator sums, including the omitted exterior terms.

4.1 Power majorants for the operator

If $a \geq 0$ and $a + \rho < 9$, Lemma 3.1 gives

$$T(t^a) \leq D(a)t^{a+\rho}, \quad D(a) = \frac{e^{2+u_0}3^{10}}{2^{a+\rho}-1} \sum_{k \geq 3} k^{a+\rho-10}. \quad (23)$$

To verify the exponent and constant, put $u = t/2^j$. Since $ku \leq u_0$ and $\lambda_k \geq \rho$,

$$\begin{aligned} e^{(k-2)u} \frac{B(ku)}{B(2u)} (ku)^a &\leq e^{2+u_0} k^{a+\lambda_k} u^{a+\lambda_k} \\ &\leq e^{2+u_0} k^{a+\rho} u_0^{\lambda_k-\rho} u^{a+\rho} \\ &= e^{2+u_0} 3^{10} k^{a+\rho-10} u^{a+\rho}. \end{aligned}$$

Summing $j \geq 1$ gives the denominator in (23). The condition $a + \rho < 9$ is exactly what makes the k -series convergent.

The directed computation verifies

$$\frac{a}{D(a) <} \left| \begin{array}{cccc} 0 & \rho & 2\rho & 3\rho \\ 31 & 24 & 24 & 28 \end{array} \right| \quad \frac{a}{D(a) <} \left| \begin{array}{ccc} 7 & 7+\rho & 7+2\rho \\ 434 & 852 & 3306 \end{array} \right| \quad (24)$$

These are finite checks with explicit tails, not decimal estimates for infinite series. Use $\rho < 117/200$, certified by $3^{200} < 2^{317}$, to increase each numerator power. The denominators remain exact:

$$2^{(r+1)\rho} = (3/2)^{r+1}, \quad 2^{7+(r+1)\rho} = 128(3/2)^{r+1}.$$

For the sum over k , evaluate $3 \leq k \leq 31$ and use, for $p > 1$,

$$\sum_{k \geq K} k^{-p} \leq K^{-p} + \frac{K^{1-p}}{p-1}, \quad K = 32. \quad (25)$$

Rational powers are enclosed by the directed logarithm and exponential rules of Section 13.

Put $d = (31, 24, 24, 28)$, $C_0 = 1$, and $C_r = \prod_{i=0}^{r-1} d_i$. In particular

$$T^r 1(t) \leq C_r t^{r\rho} \quad (0 \leq r \leq 4), \quad C_4 = 499968. \quad (26)$$

4.2 The exterior terms

For $0 < u \leq u_0$, Proposition 3.2 gives

$$\sum_{ku > u_0} f(ku) \leq \frac{14B(u_0)}{1 - e^{-u}} \leq \frac{28B(u_0)}{u}.$$

The inhomogeneous 1 is absorbed into $29B(u_0)/u$. Moreover

$$\frac{1}{B(2u)} \leq \prod_{\ell=1}^8 (1 - e^{-2^\ell u}) \leq 2^{36} u^8.$$

Drop the factor $e^{-2u} \leq 1$ and sum over $j \geq 1$ to obtain

$$0 \leq b(t) \leq C_b t^7, \quad C_b = \frac{29B(u_0)2^{36}}{127}. \quad (27)$$

The denominator 127 is $2^7 - 1$. Applying the last three constants in (24) bounds the alternating expression

$$B_3 = b - Tb + T^2b - T^3b \quad (28)$$

by $|B_3(t)| \leq \varepsilon_b(t)$, where

$$\varepsilon_b(t) = C_b t^7 \left(1 + 434t^\rho + 434 \cdot 852t^{2\rho} + 434 \cdot 852 \cdot 3306t^{3\rho} \right). \quad (29)$$

There is no assumption that B_3 is positive.

4.3 Finite recursion and its omitted terms

To enclose $T^r 1$ for $r = 1, 2, 3$, retain $3 \leq k \leq 15$ and $1 \leq j \leq J_r$, where

$$(J_1, J_2, J_3) = (35, 20, 15). \quad (30)$$

Enforce the exact condition $ku \leq u_0$ in every summand. Every inner interval is multiplied by a positive kernel, and all omitted positive terms receive an upper endpoint contribution.

For the omitted j -tail, (23) gives

$$C_r t^{r\rho} (2/3)^{rJ_r}. \quad (31)$$

For $k \geq 16$, use

$$\frac{C_{r-1} e^{2+u_0} 3^{10}}{(3/2)^r - 1} t^{r\rho} \sum_{k \geq 16} k^{r\rho-10}, \quad (32)$$

and bound the last series by (25), with $K = 16$ and $p = 10 - 117r/200$. Adding (31) and (32) double-counts their overlap, which is harmless. An inner recursion may be stopped with the full interval $[0, C_r t^{r\rho}]$. The code uses the rational enlargement

$$C_r m^r (2/3)^{er} \geq C_r (m/2^e)^{r\rho}, \quad t = m/2^e,$$

when this upper bound is below 10^{-12} . This is an enclosure, not a fitted pruning error. All additions and products are directed.

5 The finite contradiction

Four substitutions in (22) yield

$$H = CS_3 - B_3 + T^4 H, \quad S_3 = 1 - T1 + T^2 1 - T^3 1. \quad (33)$$

Since $0 < H < 14$ and T is positive, any common constant C must obey, whenever $S_3(t) > 0$,

$$\frac{H(t) - 14T^4 1(t) - \varepsilon_b(t)}{S_3(t)} \leq C \leq \frac{H(t) + \varepsilon_b(t)}{S_3(t)}. \quad (34)$$

The positivity of S_3 is explicitly certified at both test points.

Set

$$t_1 = \frac{9}{2^{21}}, \quad t_2 = \frac{3}{2^{19}}, \quad M = 2^{24}. \quad (35)$$

The program computes every a_n , $1 \leq n \leq M$, as an arbitrary-precision integer using the ascending divisor sieve. At step d , it adds a_d to each entry a_{1+kd} , so each value is final before it is used. There are no floating-point coefficients. Horner evaluation encloses $\sum_{n=1}^M a_n e^{-nt}$, and directed product evaluation encloses $B(t)$.

The omitted radial tail is bounded by

$$0 \leq H(t) - \frac{e^{-t}}{B(t)} \sum_{n=1}^M a_n e^{-nt} \leq \frac{14e^{-Mt/2}}{1 - e^{-t/2}}. \quad (36)$$

Indeed positivity gives $\sum_{n>M} a_n e^{-nt} \leq e^{-Mt/2} f(t/2)$. Apply $f(t/2) < 14e^{-t/2} B(t/2)$ and the exact relation $B(t/2)/B(t) = (1 - e^{-t/2})^{-1}$; dropping $e^{-3t/2}$ enlarges the result. The two exponents are exactly $Mt_1/2 = 36$ and $Mt_2/2 = 48$. This estimate is used as an interval endpoint, not just quoted as small.

Proposition 5.1 (Directed conditional intervals). *Under either hypothesis $\lim_{t \downarrow 0} H(t) = C$ or $\mathcal{P} \equiv C$, the same real number C belongs to both intervals below. Endpoints are rounded outward:*

$$I_1 = [4.9933847050426907, 4.9933867654594445],$$

$$I_2 = [4.9933622876055544, 4.9933662323124263].$$

The full 256-bit integer endpoints are included in the accompanying receipt. Their exact separation is

$$\frac{2138996031241830049592557078436499944663290918102101242885845234882473289}{2^{256}} > 0. \quad (37)$$

Proof. Compute the radial enclosures by (36), the operator enclosures by (30)–(32), and the T^4 and exterior bounds by (26) and (29). Substitute the resulting intervals in (34), using outward division by the certified positive S_3 . The compact receipts retain every integer endpoint; `combine_certificate.py` computes the intervals and computes the first lower endpoint minus the second upper endpoint; the replay compares it with the pinned positive numerator in (37) divided by 2^{256} . The reproduction procedure regenerates the underlying 2^{24} coefficients and all three operator powers before comparing receipts. $\square$

The numerator in (37) is an exact integer, not a digitized plot or a decimal-fit residual. To avoid a common misinterpretation, the intervals I_1, I_2 are *conditional constraints on a single constant*. They are not pointwise enclosures for $\mathcal{P}(t_1)$ and $\mathcal{P}(t_2)$. Their gap is not asserted to be a lower bound on the actual oscillation of $\mathcal{P}$. They prove that H has no radial limit and that $\mathcal{P}$ is nonconstant.

6 Abelian transfer to the two sequence ratios

Lemma 6.1 (Positive Abelian averaging). *Let $u_n \geq 0$, $v_n > 0$, and suppose $V(z) = \sum v_n z^n \rightarrow \infty$ as $z \uparrow 1$. If $u_n/v_n \rightarrow C < \infty$, then*

$$\frac{\sum u_n z^n}{\sum v_n z^n} \rightarrow C.$$

If instead $u_n/v_n \rightarrow +\infty$, the generating-function ratio tends to $+\infty$.

Proof. For finite C , split $\sum (u_n - Cv_n)z^n$ at an index beyond which $|u_n - Cv_n| \leq \varepsilon v_n$. The fixed initial part divided by $V(z)$ tends to zero and the tail contributes at most ε . For an infinite limit, use $u_n \geq Av_n$ beyond a fixed index and then let the arbitrary A increase. $\square$

The generating functions of $q(n)$ and $Q(n)$ are respectively $F(z)/z$ and $F(z)/(z(1-z))$. Those of β_n and B_n are $\mathcal{B}(z)$ and $\mathcal{B}(z)/(1-z)$. Both generating-function ratios are therefore

$$\frac{F(z)}{z\mathcal{B}(z)} = G(-\log z). \quad (38)$$

Both denominators diverge because $\beta_n \geq 1$. If either sequence ratio in Theorem 1.1 converged to a finite C , Lemma 6.1 would force $G(t) \rightarrow C$ and hence $H(t) = e^{-2t}G(t) \rightarrow C$. Proposition 5.1 rules this out. The global bound $G < 14$ rules out a ratio tending to $+\infty$ by the same lemma; nonnegativity excludes $-\infty$. This proves Theorem 1.1. No phase-density statement, Tauberian theorem, saddle expansion, or coefficient equivalent is used.

For the original real-variable summatory question, $Q(x)$ and its binary comparator depend on $\lfloor x \rfloor$. The integer nonconvergence therefore gives the same negative answer for real $x \rightarrow \infty$.

7 A separate quantitative oscillation corollary

The conditional intervals also yield an actual oscillation lower bound, but only after the additional argument below. They remain conditional constraints on a common constant and are never reinterpreted as values of $\mathcal{P}$.

Theorem 7.1 (Actual oscillation lower bound). *Let $\omega = \sup_{t>0} \mathcal{P}(t) - \inf_{t>0} \mathcal{P}(t)$. Then*

$$\omega > 0.00001839189788. \quad (39)$$

For each of the two ratios in Theorem 1.1, its limit superior minus its limit inferior is at least ω , and therefore exceeds the same displayed decimal.

Proof. On $D = (0, u_0]$, T is a bounded positive operator on bounded real functions: $T1(t) \leq 31t^\rho \leq 31u_0^\rho$. Also b is bounded by (27). Without assuming constancy, the exact telescoping identity is

$$\mathcal{P} = H + TH + b.$$

Put $A_3 = I - T + T^2 - T^3$, $S_3 = A_31$, and $U_3 = (I + T + T^2 + T^3)1$. Four substitutions, or $A_3(I + T) = I - T^4$, give

$$H = A_3\mathcal{P} - B_3 + T^4H. \quad (40)$$

Every argument at which the operator evaluates a function remains in D .

Let $m = \inf_D \mathcal{P}$, $M = \sup_D \mathcal{P}$, $C = (M + m)/2$, and $e = \mathcal{P} - C$. These numbers are finite, since $0 < \mathcal{P} \leq 14$. Dyadic periodicity makes the range on D equal to the full positive-axis range, so $M - m = \omega$. Positivity gives

$$|A_3e(t)| \leq \frac{\omega}{2}U_3(t).$$

At either test point, define the auxiliary quantity

$$X(t) = \frac{H(t) + B_3(t) - T^4H(t)}{S_3(t)} = C + \frac{A_3e(t)}{S_3(t)}.$$

The already established bounds place $X(t_i)$ in the full fixed-point interval used to print I_i in Proposition 5.1. This is an unconditional enclosure of X , which is a different quantity from $\mathcal{P}(t_i)$. If $g > 0$ is the exact interval gap (37) and $R_i = U_3(t_i)/S_3(t_i)$, then

$$g \leq X(t_1) - X(t_2) \leq \frac{\omega}{2}(R_1 + R_2).$$

Thus $\omega \geq 2g/(R_1 + R_2)$.

To bound R_i from above, sum the stored upper endpoints for $T^r1(t_i)$, $0 \leq r \leq 3$, and divide by the stored strictly positive lower endpoint for $S_3(t_i)$. Independent exact rational reconstruction certifies

$$\begin{aligned} g &> 0.0000184727302644825, \\ R_1 &< 1.004024051499, \quad R_2 < 1.004765947242. \end{aligned}$$

Consequently

$$\omega > \frac{821010233977}{44639777749800000} > 0.00001839189788. \quad (41)$$

Thus

$$\frac{B(t)}{|B(w)|} \leq C_{y_0} t^{-d_0}, \quad d_0 = \frac{\log(1 + y_0^2)}{2h}. \quad (42)$$

Choose y_0 small enough that $\sigma := \rho - d_0 > 0$. Positivity gives $|f(w)| \leq f(t)$. Apply (42) at $2t$ to the exact expression for R in (13) to obtain

$$|R(w)| \leq C t^\sigma. \quad (43)$$

The series (19) therefore converges normally in the open sector $|\Im w| < y_0 \operatorname{Re} w$, defines a holomorphic $\mathcal{P}$, and satisfies $\mathcal{P}(2w) = \mathcal{P}(w)$. On every narrower sector,

$$H(w) = \mathcal{P}(w) + O(|w|^\sigma), \quad H^{(k)}(w) - \mathcal{P}^{(k)}(w) = O_k(|w|^{\sigma-k}) \quad (44)$$

for every fixed k , by Cauchy estimates on disks of radius proportional to $|w|$. Multiplicative periodicity and compactness of a dyadic annulus imply

$$|t^k \mathcal{P}^{(k)}(t)| \leq C_k \quad (t > 0). \quad (45)$$

The exact normalization for subsequent transfer is

$$f(w) = e^w B(w) (\mathcal{P}(w) + O(|w|^\sigma)). \quad (46)$$

This sector estimate does not by itself control a coefficient circle. That missing step is proved in Section 10.

9 Saddle parameters and the finite correction rule

Differentiating the absolutely convergent product gives

$$b_k(t) = \sum_{j \geq 0} 2^{kj} \sum_{r \geq 1} r^{k-1} e^{-r 2^j t}. \quad (47)$$

For every fixed $k \geq 1$,

$$t^k b_k(t) = (k-1)! \frac{L}{h} + O_k(1). \quad (48)$$

Indeed $u^k \sum_{r \geq 1} r^{k-1} e^{-ru} = (k-1)! + O_k(u)$ as $u \downarrow 0$. Sum this over $u = 2^j t \leq 1$, whose error sum is bounded, and bound the remaining dyadic tail by exponential decay. The small- u estimate follows by differentiating $1/(e^u - 1) = u^{-1} + O(1)$ a fixed number of times, or by its convergent analytic expansion near zero after removing the pole.

In particular $b'_1(t) = -b_2(t) < 0$, $b_1(0+) = \infty$, and $b_1(\infty) = 0$, proving the existence and uniqueness of t_N . At $N = n + 1$,

$$L = \log N - \log \log N + \log h + o(1), \quad t \sim \frac{\log N}{hN}, \quad v = b_2(t) \sim \frac{L}{ht^2}. \quad (49)$$

For a differentiable amplitude P , define $\mathcal{A}_r[P]$ by summing over all nonnegative integers $\ell, m_3, \dots, m_{2r+2}$ with

$$\ell + \sum_{k=3}^{2r+2} (k-2)m_k = 2r.$$

Put $D = \ell + \sum km_k$, which is even. Then

$$\mathcal{A}_r[P](t) = \sum \frac{(-1)^{\ell+r+\sum m_k} (D-1)!! P^{(\ell)}(t)}{\ell! v^{\ell/2}} \prod_{k=3}^{2r+2} \frac{1}{m_k!} \left(\frac{b_k(t)}{k! v^{k/2}} \right)^{m_k}. \quad (50)$$

The sum is finite and $(-1)!! = 1$. The full exponent of v in any term is the integer $D/2$, even though the factored display uses half-powers. Thus (50) is directly evaluable with exact rational jets.

The first correction is

$$\mathcal{A}_1[P] = P \left(\frac{b_4}{8v^2} - \frac{5b_3^2}{24v^3} \right) - \frac{P'b_3}{2v^2} - \frac{P''}{2v}. \quad (51)$$

Using (45) and (48), a term of grade $2r$ has size $O_r(L^{-r})$ for $P = \mathcal{P}$. For $P = \mathcal{P}/t$, it has size $O_r(t^{-1}L^{-r})$.

In any fixed truncation, one may replace $\mathcal{P}$ by H in the entire prescription without changing the remainders in Theorem 1.2; (44) makes the resulting difference power-small in t . This is a way to retain extra radial information without knowing Fourier coefficients of the limiting profile. It does not turn a finite radial computation into a certified pointwise profile evaluation unless its own tail has been bounded.

10 An integrated bound for the whole circle

Define

$$D(z) = \frac{z}{1-z} \left(1 + \sum_{k \geq 3} F(z^k) \right), \quad A_j(z) = \prod_{\ell=0}^{j-1} \frac{z^{2^\ell}}{1-z^{2^\ell}}, \quad A_0 = 1. \quad (52)$$

Iterating the binary part of (10) gives the normally convergent positive-prefix expansion

$$F(z) = \sum_{j \geq 0} A_j(z) D(z^{2^j}). \quad (53)$$

For clarity, the remainder after j steps is $A_j(z)F(z^{2^j})$. On each compact disk $|z| \leq r < 1$, the powers r^{2^j} and $F(r^{2^j})$ decay fast enough to make this remainder tend uniformly to zero. Every coefficient in D is nonnegative.

At $r = e^{-t}$, let

$$w_j = \frac{A_j(r)D(r^{2^j})}{F(r)}, \quad w_j \geq 0, \quad \sum_{j \geq 0} w_j = 1.$$

For small u , (13) gives

$$\frac{D(e^{-u})}{f(u)} = \frac{R(u)}{H(u)} = O(u^\rho).$$

The cancellation of the binary factors and powers is exact:

$$\frac{A_j(r)f(2^j t)}{f(t)} = \frac{H(2^j t)}{H(t)}, \quad w_j = \frac{R(2^j t)}{H(t)}. \quad (54)$$

Put $K = \lfloor L/(2h) \rfloor$. Boundedness above and below of H near zero then gives

$$\sum_{j < K} w_j = O(t^{\rho/2}). \quad (55)$$

For $u > 0$ define

$$\chi(u, \theta) = \frac{1 - e^{-u}}{|1 - e^{-u+i\theta}|}, \quad \Xi_K(t, \theta) = \prod_{\ell=0}^{K-1} \chi(2^\ell t, 2^\ell \theta).$$

The function D contains one geometric denominator beyond the j prefix factors; its remaining coefficients are nonnegative. Therefore the exact bound is

$$\frac{|F(re^{i\theta})|}{F(r)} \leq \sum_{j \geq 0} w_j \Xi_{j+1}(t, \theta). \quad (56)$$

This is a bound for the actual F , not an assumption that F is an Euler product.

10.1 Short prefixes

The first factor has

$$\int_{-\pi}^{\pi} \chi(t, \theta) d\theta \leq Ct(1 + L). \quad (57)$$

For example, $|1 - e^{-t+i\theta}|^2 = (1 - e^{-t})^2 + 4e^{-t} \sin^2(\theta/2)$ and $\sin(|\theta|/2) \geq |\theta|/\pi$ on this interval imply the estimate by elementary integration. Since every $\chi \leq 1$, (55) and (57) bound the integral of the terms $j < K$ in (56) by

$$O(t^{1+\rho/2}(1 + L)). \quad (58)$$

10.2 Long prefixes at intermediate angles

For $j \geq K$, $\Xi_{j+1} \leq \Xi_K$. If $|\theta| \leq \sqrt{t}$, then $2^K \leq t^{-1/2}$ makes every retained radial argument and angle bounded by one. The same exact factor formula gives

$$\Xi_K(t, \theta) \leq [1 + c(\theta/t)^2]^{-K/2} \quad (59)$$

with an absolute $c > 0$. Set $\theta_0 = tL^{-2/5}$. Integrating (59), for instance after $y = \theta/t$, gives

$$\int_{\theta_0 \leq |\theta| \leq \sqrt{t}} \Xi_K(t, \theta) d\theta \leq C \frac{t}{\sqrt{L}} \exp(-cL^{1/5}). \quad (60)$$

One can split $|y| \leq 1$ from $|y| \geq 1$: in the first region use $\log(1 + cy^2) \geq c'y^2$; in the second keep a fixed integrable power and an exponentially small factor in K . This justifies the Gaussian-scale prefactor and does not rely on a pointwise estimate times the entire circle length.

10.3 The outer angles and dyadic secondary peaks

For $\sqrt{t} \leq |\theta| \leq \pi$, the first factor obeys $\chi(t, \theta) \leq C\sqrt{t}$. The next two factors have an L^1 product of order t . In fact

$$\int_{-\pi}^{\pi} \chi(u, \theta)^2 d\theta = 2\pi \tanh(u/2), \quad (61)$$

by the geometric-series Poisson kernel. Periodic changes of variable and Cauchy-Schwarz give

$$\int_{-\pi}^{\pi} \chi(2t, 2\theta) \chi(4t, 4\theta) d\theta \leq 2\pi \sqrt{\tanh(t) \tanh(2t)} \leq Ct.$$

For sufficiently small t , $K \geq 3$; discard the remaining factors, which are at most one. Thus

$$\int_{\sqrt{t} \leq |\theta| \leq \pi} \Xi_K(t, \theta) d\theta \leq Ct^{3/2}. \quad (62)$$

This integrated estimate controls the possible secondary dyadic peaks. Bounding their height alone would not be enough at the central coefficient scale.

Combining (58), (60), and (62), and dividing by the central angular volume $1/\sqrt{v} \asymp t/\sqrt{L}$, shows that the omitted circle $|\theta| > \theta_0$ contributes a relative error

$$O\left(L^{3/2}t^{3/2} + \sqrt{L}t^{1/2} + \exp(-cL^{1/5})\right). \quad (63)$$

It is smaller than every fixed inverse power of L . For the partial-sum generating function $F(z)/(1-z)$, the normalized additional factor on the same circle is precisely $\chi(t, \theta) \leq 1$. The same whole-circle estimate is therefore valid at its own radial scale.

11 Local expansion and its remainder

Cauchy's formula has integrand $f(t - i\theta)e^{n(t-i\theta)}$. On $|\theta| \leq \theta_0$, substitute (46) to obtain the principal integrand

$$B(t - i\theta)e^{(n+1)(t-i\theta)}\mathcal{P}(t - i\theta).$$

The power-small error in (46) integrates to a relative $O(t^\sigma)$ after the local Gaussian bound below. The factor $n + 1$ is the reason for the saddle convention $b_1(t) = n + 1$.

Put $\theta = x/\sqrt{v}$. The saddle cancels the linear phase, leaving

$$\exp\left(-\frac{x^2}{2} + \sum_{k \geq 3} \frac{b_k(t)(ix)^k}{k!v^{k/2}}\right). \quad (64)$$

Taylor-expand $\mathcal{P}(t - ix/\sqrt{v})$. Assign grade ℓ to its ℓ th derivative factor and grade $k - 2$ to the k th cumulant in (64). By (48) and (45), a grade- g monomial is $O(L^{-g/2})$ times a fixed polynomial in x . Odd grades integrate to zero over the symmetric interval. The Gaussian integral for an even total degree D is $(D - 1)!!$, with the powers of i and the amplitude sign giving exactly (50).

Here is a fixed-order remainder justification. On the central interval,

$$|x| \leq CL^{1/10}, \quad \left| \sum_{k \geq 3} \frac{b_k(t)(ix)^k}{k!v^{k/2}} \right| \leq C \frac{|x|^3}{\sqrt{L}} = o(x^2).$$

For the last estimate use analyticity of $\log B(w)$ on $|w - t| \leq t/2$ and the derivative bounds $b_k = O_k(Lt^{-k})$; $|\theta|/t = L^{-2/5}$ tends to zero. Fix J . Expand the amplitude and phase through enough derivatives to retain all terms of grade at most $2J + 1$. Taylor's theorem on a smaller disk, followed by the exponential remainder formula, bounds the omitted integrand by

$$C_J L^{-J-1} (1 + |x|^{m_J}) e^{-cx^2} \quad (65)$$

for a fixed integer m_J . More explicitly, the amplitude remainder after degree $2J + 1$ is $O(L^{-J-1}|x|^{2J+2})$; the logarithmic-phase remainder after cumulant degree $2J + 3$ is $O(L^{-J-1}|x|^{2J+4})$. Expand the exponential through power $2J + 1$ of the retained nonquadratic phase polynomial. Its

tail is $O_J(L^{-J-1}|x|^{6J+6})$, since that polynomial is $O_J(L^{-1/2}|x|^3) = o(1)$ uniformly on this arc. The factor from the logarithmic Taylor remainder contributes $O_J(L^{-J-1}|x|^{2J+4})$. In the resulting finite product, discard monomials of grade at least $2J+2$; each is $O_J(L^{-J-1})$ times a fixed polynomial. All these errors are absorbed by a fixed part of the Gaussian. The retained terms of odd grade integrate to zero; the next even grade is $2J+2$.

Integration of (65) gives $O_J(L^{-J-1})$. Extending every retained Gaussian polynomial integral to the real line costs $O_J(\exp(-cL^{1/5}))$. The whole-circle error (63) and the sector error $O(t^\sigma)$ are both smaller than every fixed inverse power. This proves (6) with its claimed remainder.

For partial sums,

$$\sum_{n \geq 0} S_n z^n = \frac{F(z)}{1-z}.$$

On the central sector the amplitude becomes $\mathcal{P}(w)/(1 - e^{-w})$. Replace it by $\mathcal{P}(w)/w$, since $w/(1 - e^{-w}) = 1 + O(w)$ with the corresponding fixed derivative estimates. This changes the bracket by a power-small relative error. The derivatives of $\mathcal{P}(t)/t$ are $O_k(t^{-k-1})$, so the same grade calculation gives the bracket remainder $O_J(t^{-1}L^{-J-1})$. The normalized whole-circle bound was already checked after (63). This proves (7) and completes Theorem 1.2. Taking $J = 0$ and using (49) gives (8).

12 Inverse centers and integer thresholds

For real Y tending to infinity, define

$$\nu_a(Y) = \min\{n \geq 1 : a_n \geq Y\}, \quad \nu_S(Y) = \min\{n \geq 1 : S_n \geq Y\}.$$

The recurrence gives $a_{n+1} \geq a_n + 1$ for $n \geq 2$, because both divisors 1 and n occur. Thus the counts are eventually strictly increasing and unbounded; S_n is strictly increasing from the beginning.

For real x large, set $b_1(t) = x + 1$ and define the smooth truncations

$$f_{a,J}(x) = \frac{e^{(x+1)t} B(t)}{\sqrt{2\pi b_2(t)}} \sum_{r=0}^J \mathcal{A}_r[\mathcal{P}](t), \quad (66)$$

$$f_{S,J}(x) = \frac{e^{(x+1)t} B(t)}{\sqrt{2\pi b_2(t)}} \sum_{r=0}^J \mathcal{A}_r[\mathcal{P}/t](t). \quad (67)$$

These are positive and eventually strictly increasing. To prove this, differentiate the approximants themselves, not the unknown integer-index error in the asymptotic theorem. Since $t' = -1/b_2(t)$, the derivative of the Legendre term $(x+1)t + \log B(t)$ is exactly t . The logarithmic derivative of the variance factor is $O(t/L)$. The positive lower bound for $\mathcal{P}$, its derivative bounds, and the fixed correction formulas show that the remaining logarithmic derivative is also $O_J(t/L)$. The factor $1/t$ in the summatory amplitude has the same bound. Hence

$$\frac{d}{dx} \log f_{\star,J}(x) = t(1 + O_J(L^{-1})), \quad \star \in \{a, S\}. \quad (68)$$

Theorem 12.1 (Fixed-order inverse uncertainty). *Let $x_{\star,J}(Y)$ be the eventually unique real solution of $f_{\star,J}(x) = Y$. For each fixed $J \geq 0$, there are constants C_J and an onset such that, for either $\star = a$ or S and all sufficiently large Y ,*

$$\left[x_{\star,J} - \frac{C_J x_{\star,J}}{(\log x_{\star,J})^{J+2}} \right] \leq \nu_{\star}(Y) \leq \left[x_{\star,J} + \frac{C_J x_{\star,J}}{(\log x_{\star,J})^{J+2}} \right]. \quad (69)$$

In particular this is an inverse approximation at every fixed relative logarithmic order. Its absolute bracket width need not tend to zero.

Proof. Theorem 1.2 says $a_n/f_{a,J}(n) = 1 + O_J((\log n)^{-J-1})$, and the same holds for S_n and $f_{S,J}$. Integrating (68) over an interval of length $\delta = C_J x / (\log x)^{J+2}$ changes $\log f_{*,J}$ by a constant multiple of

$$t\delta \asymp \frac{\log x}{x} \frac{C_J x}{(\log x)^{J+2}} = C_J (\log x)^{-J-1}.$$

Choose C_J larger than the uniform error constant. Compare the exact count at $\lfloor x - \delta \rfloor$ and $\lceil x + \delta \rceil$, both asymptotic to x . The first is less than Y and the second is at least Y . Monotonicity then controls every earlier index and gives (69). Rounding at the upper endpoint only increases the approximant, so no unproved exact-ceiling inference is involved. $\square$

12.1 Explicit leading inverse scales

Elementary dyadic product estimates give

$$\log B(t) = \frac{L^2}{2h} + \frac{L}{2} + O(1), \quad tb_1(t) = \frac{L}{h} + O(1). \quad (70)$$

For the first estimate, use $-\log(1 - e^{-u}) = -\log u + O(u)$ through $2^j t \leq 1$. The sum of the $O(u)$ errors and the large- u tail are bounded. If $m = \lfloor L/h \rfloor$, the main sum is $(m+1)L - hm(m+1)/2 = L^2/(2h) + L/2 + O(1)$. The second estimate follows similarly from $u/(e^u - 1) = 1 + O(u)$.

Using the bounded positive amplitude and the variance estimate in the leading saddle formula gives

$$\log a_n = \frac{L^2}{2h} + \left(\frac{1}{h} - \frac{1}{2}\right)L - \frac{1}{2}\log L + O(1), \quad (71)$$

$$\log S_n = \frac{L^2}{2h} + \left(\frac{1}{h} + \frac{1}{2}\right)L - \frac{1}{2}\log L + O(1). \quad (72)$$

The bounded terms include the classical dyadic fluctuation. Set $y = \log Y$. Solving these equations gives respectively

$$L = \sqrt{2hy} - 1 + \frac{h}{2} + o(1), \quad L = \sqrt{2hy} - 1 - \frac{h}{2} + o(1).$$

The errors here are $O(\log y / \sqrt{y}) = o(1)$, so exponentiation preserves the leading constants. Since $n \sim (L/h)e^L$,

$$\begin{aligned} \nu_a(Y) &\sim \frac{2}{e\sqrt{h}} \sqrt{\log Y} \exp(\sqrt{2h \log Y}), \\ \nu_S(Y) &\sim \frac{1}{e\sqrt{h}} \sqrt{\log Y} \exp(\sqrt{2h \log Y}). \end{aligned} \quad (73)$$

The factor two distinguishes exact-size counts from their cumulative count. The inverse techniques are standard; the exact recurrence and the fixed-order error control determine their application here.

Remark 12.2 ([write] The inverses and the transseries volume, 7 October 2026). Compared statement by statement with the repository's transseries volume [12]; its symbols are not used here.

1. Theorem 12.1 brackets the integer staircases ν_a, ν_S (the volume's N_* of `p0: def: three-inverses(1)`; a_n is eventually strictly increasing and S_n strictly increasing)

around the solutions $x_{*,J}$ of $f_{*,J}(x) = Y$. The smooth approximants $f_{*,J}$ are not interpolations of the sequences, so `p0:thm:staircase(1)` does not apply to them; the bracket is proved directly at the integers from Theorem 1.2 and the slope (68). It is an analogue of `p0:thm:staircase(2)`, not an instance, with a relative (not absolute) width.

2. The centres $x_{*,J}$ are implicit inverses of explicit smooth functions; no formal inverse series is printed. The growth $\log a_n \sim \log^2 n / (2 \log 2)$ is not of the exponential–power type of `p0:def:model` ($\log a_n$ is neither an^α nor comparable to it), and $\log a_n$ as a function of $\log n$ has a quadratic leading balance, which is not the monomial–logarithmic form $\rho Z + \mu \log Z$ that `plt:def:lw-monomial-log-datum` requires; after the substitution $x = \exp \sqrt{2h} \log Y$ the leading balance changes, and the write did not examine whether the inverse then becomes an instance of `plt:thm:lw-template`. The centres and the leading scales (73) are therefore not shown to be instances of a theorem of the volume.

13 Exact arithmetic and the reproduction contract

The archive is deliberately small. It contains this L^AT_EX source and PDF, standard-library Python code, short explicitly labeled OEIS prefix fixtures, exact semantic receipts, a source ledger, an inventory, and a SHA-256 manifest. It contains no third-party papers, private audit files, full coefficient arrays, or floating-point diagnostics. The paper proves the infinite analytic estimates; the scripts certify the explicitly finite inequalities and interval endpoints.

13.1 Arithmetic rules

All interval endpoints are integers on the lattice $2^{-256}\mathbb{Z}$. Conversion from a rational number uses floor and ceiling. Addition and subtraction are exact at the endpoint level; multiplication and division use all four corners and outward floor/ceiling, with division rejecting an interval containing zero. Nonnegative integer powers use interval multiplication.

To enclose e^{-x} for rational $x \geq 0$, halve x until it lies in $[0, 1/16]$, use the alternating rational Taylor series with its next-term enclosure, and square back using directed arithmetic. Negative x uses reciprocal enclosure. To enclose logarithms, reduce a positive rational argument by powers of two to $[1, 2]$ and use

$$\log x = 2 \sum_{j \geq 0} \frac{z^{2j+1}}{2j+1}, \quad z = \frac{x-1}{x+1}.$$

After M_0 terms the nonnegative tail is at most $2z^{2M_0+1}/((2M_0+1)(1-z^2))$. Rational powers use this logarithm followed by monotone exponential enclosure.

The binary product stops when its next factor parameter has upper bound $q \leq 2/2^{256}$. The remaining logarithmic tail is at most $q/(1-q)^2 < 4/2^{256}$, so the omitted product lies in $[1, 1+8/2^{256}]$. Horner evaluation of nonnegative polynomials is directed at every multiplication. No floating-point quantity affects an endpoint or pass/fail decision.

13.2 Checks and their limits

The finite model check independently compares divisor-by-divisor and sieve implementations through 10000, checks 29 direct chain-partition cases, 999 summatory floor recurrences, 5001 binary cumulative identities, and 9998 strict-increase cases. It compares 207 displayed OEIS terms across the four

entries. The optional named `b003238.txt` and `b003318.txt` files contain only 48 displayed terms each; their labels and receipts do not claim a full b-file comparison.

The finite saddle engine enumerates the multiplicities in (50). A separate implementation builds the formal exponential by a graded differential recurrence and integrates exact Gaussian moments. It compares orders zero through four on 16 rational jet inputs, for 80 exact cases. These tests validate the finite algebra, not the analytic remainder theorem.

All essential guards are explicit exceptions, not removable `assert` statements. Deliberately corrupted receipts test precision, row count, test points, positive denominator, nonnegative error, endpoint orientation, disjointness, ordering, and status. The package-level checks test source inventory, hashes, unsafe paths, duplicate JSON keys, and pre-write output safety. Normal and optimized Python runs must generate identical semantic receipts. Runtime and memory metadata are excluded from those receipts.

13.3 Rebuilding

From the extracted package, run

```
python3 reproduce.py --out /absolute/new/output-directory
```

The output directory must not exist and must be outside the source tree. The runner verifies the manifest, makes an isolated copy, runs the full exact certificate sequentially, compares every newly generated receipt with its supplied baseline, runs the supplemental and deliberate-failure checks in normal and optimized modes, rebuilds the PDF with a private TeX format until its bytes stabilize, and writes the deterministic ZIP. The complete arithmetic sequence is

```
certify_global_bound.py
operator_constants.py
certify_radial_values.py
certify_operator.py
combine_certificate.py
```

The radial step genuinely computes all 2^{24} coefficients each time. It needs roughly 1.6 GiB of memory; the development replay took about 140 seconds for that step and about 90 seconds for the operator step. These are planning estimates, not guarantees or inputs. Serialize memory-heavy replays.

Python uses only its standard library. PDF reproduction additionally requires the documented pdfTeX/L^AT_EX and Latin Modern toolchain. The runner disables shell escape, fixes the build epoch and archive metadata, uses an explicit source inventory, and excludes runtime logs from the archive. Exact PDF and ZIP byte identity is promised only with the same TeX and font toolchain; the mathematical receipts are independently deterministic. See `README.txt` for normal and optimized replay commands and comparison against the original ZIP.

14 Further directions

Certified profile values and sharper oscillation bounds. Theorem 7.1 gives a genuine oscillation lower bound through a midpoint argument. Directly enclosing $H(t) + \sum_{j \geq 1} R(t/2^j)$ with

sufficiently sharp correlated tails could additionally yield true pointwise intervals for $\mathcal{P}(t)$, sharpen that lower bound, and supply an upper bound. The original conditional interval gap by itself is still not an oscillation bound.

Effective asymptotic onsets. The fixed-order theorem has existential constants. Making the sector constants, integrated-circle constants, and local Taylor constants explicit would turn the inverse brackets into usable certified numerical thresholds. It would also quantify how large n must be before a selected correction improves a given tolerance.

Fourier information and the total amplitude. Writing $\mathcal{P}(t) = \Phi(\log_2 t)$ gives an analytic one-periodic function on a strip. Its Fourier coefficients may be very small. A rigorous coefficient enclosure would illuminate the size and phase of the fluctuation and could improve the positive-operator certificate. No numerical Fourier coefficient is asserted here.

Smaller certificates. The cut u_0 , the two test points, the recursion cutoffs, and the 2^{24} polynomial cutoff were chosen to give a comfortable certified separation. Sharper power majorants, a more efficient exterior treatment, or a different positive iteration may reduce the exact coefficient workload. Such changes need a new full proof of every altered tail and a fresh exact replay.

Beyond fixed inverse-logarithmic order. The terms suppressed as powers of t include information from the full all-integer dilation forcing. Resolving them would require a finer expansion than Theorem 1.2, together with corresponding control of other circle regions. A growing-order or exponentially improved expansion is a separate problem, not an automatic corollary of the finite Gaussian prescription.

Continued source comparison. The full original Pennington and Kato–McLeod theorem scopes, a final 2026 version of the distinct-chain-partition work, and later citations of Erdős–Loxton are natural further checks. A newly found exact-target theorem could change the historical framing without changing the finite certificate. The current conclusions rest on the displayed proof and bounded source comparison.

[write] **Added 7 October 2026, under Vladimir’s standing rule of 4 October 2026.** Every open question of the source is in this section, with its sketch and what is missing: certified pointwise profile values and an upper bound for the oscillation; effective onsets for Theorem 1.2 and Theorem 12.1; Fourier coefficients of the profile; smaller certificates; the terms below every fixed inverse-logarithmic order; and the continued source comparison (the full original results of Pennington and of Kato–McLeod, a final 2026 version of the distinct-chain-partition preprint, later citations of Erdős–Loxton), none of which the write read. Also open, from the non-claims: an exact eventual ceiling rule and growing-order uniformity. One claim outside the source is refuted: Cloitre’s 2004 conjecture in A003238 that $0.4 < c < 0.5$ (Remark 1.3, with proof; $c = 1/(2 \log 2)$). No claim of the source was found to be false.

[write] **Independent check of the write, 7 October 2026.** An independent adversarial check read A003238 (#133), A003318 (#60), A018819 (#219) and A000123 (#324) again. Every quotation of Remark 1.3 is verbatim. It also re-read Section 3 of Erdős–Loxton in the published scan: (8), (9), $B(x) = b(2[x])$, Lemma 1, Theorem 3 on p. 324, and the open question on p. 328. These are exactly as the note at the end of Section 2 reports. That paper says only that its methods “could be used to give bounds for the oscillation”. Reading these as upper bounds, as the note does, is an inference: a positive lower bound would already have decided the question.

- *The refutation of Cloitre’s conjecture.* The refutation stands; the derivation from (71) and (49) was redone. It is also independent of the logarithm’s base: in base b the constant is $\log b/(2 \log 2)$, which is $\frac{1}{2}$ in base 2 and so still outside the open interval. A floating divisor sieve to 2^{25} continues the write’s data: $\log a_n/\log^2 n$ is 0.503, 0.518, 0.531 and 0.537 at $n = 10^5, 10^6, 10^7$ and 2^{25} . It already leaves $(0.4, 0.5)$ near $n = 10^5$; finite data prove nothing asymptotic, but they agree with the proof.
- *OEIS data.* The exact sieve agrees with all b-file terms the write compared and with both 48-term fixtures. $B_n = \beta_{2n}$ holds for $n \leq 20000$.
- *The radial sieve, independently.* The write did not rerun it. The check computed $a_{2^{24}}$ exactly by the same ascending sieve in eight word-size prime moduli and the Chinese remainder theorem; its own code, without big integers, gives exactly the receipt’s endpoint

$$a_{2^{24}} = 13242676671038590346167600227591139744796090194799488130189680393.$$

A floating sum of $a_n e^{-nt}$ over $n \leq 2^{24}$ reproduces the receipt’s finite radial values H at t_1 and t_2 to relative $2 \cdot 10^{-13}$. The actual tail between 2^{24} and 2^{25} is $2.2 \cdot 10^{-13}$ at t_1 , well below the bound $1.51 \cdot 10^{-9}$ of (36), and invisible at t_2 .

- *The operator.* A direct floating evaluation of $T1(t_i)$ from (21), with no code of the package, gives 0.0020079643... and 0.0023772839.... Both lie inside the certified enclosures.
- *Exact reconstruction from the receipts.* From the stored radial, operator, T^4 and exterior endpoints, the check rebuilt S_3, I_1, I_2 , the gap (37), R_1, R_2 and $2g/(R_1 + R_2)$ in exact rationals, with its own code. Everything agrees with the receipts and with the printed decimals, the rational $821010233977/4463977749800000$ and the margin of Section 7. The logic of the oscillation corollary and of the Abelian transfer was re-read.
- *Reruns and leading terms.* The delivered certificate chain was rerun on a copy (all steps but the radial sieve). The leading inverse constants of (73) were re-derived by hand.

The provenance figures (601,832 bytes, 37 files, 794 lines, 21 pages, 36 manifest entries), the byte identity of the 33 staged files, the README listing and the numbering (95 labels unchanged, 3 added) are confirmed. No claim of the write was found wrong, and nothing needed correcting.

References

- [1] OEIS Foundation, *A003238*, uniform rooted-tree counts, <https://oeis.org/A003238>, inspected 4 October 2026.
- [2] OEIS Foundation, *A003318*, partial sums of *A003238*, <https://oeis.org/A003318>, inspected 4 October 2026.
- [3] OEIS Foundation, *A018819* and *A000123*, <https://oeis.org/A018819> and <https://oeis.org/A000123>, inspected 4 October 2026.
- [4] P. Erdős and J. H. Loxton, Some problems in partitio numerorum, *J. Austral. Math. Soc. Ser. A* **27** (1979), 319–331. <https://doi.org/10.1017/S144678870001243X>.
- [5] M. K. Gol’dberg and E. M. Livshits, On minimal universal trees, *Math. Notes* **4** (1968), 713–717; Russian original in *Mat. Zametki* **4**. <https://www.mathnet.ru/eng/mzm9458>.
- [6] G. Gati, F. Harary, and R. W. Robinson, Line colored trees with extendable automorphisms, *Acta Math. Sci.* **2** (1982), 105–110. <https://oeis.org/A003238/a003238.pdf>.

- [7] F. Harary and R. W. Robinson, The number of achiral trees, *J. Reine Angew. Math.* **278/279** (1975), 322–335. <https://doi.org/10.1515/crll.1975.278-279.322>.
- [8] K. Mahler, On a special functional equation, *J. London Math. Soc.* **15** (1940), 115–123. Primary archive: <https://carmamaths.org/resources/mahler/docs/068.pdf>.
- [9] N. G. de Bruijn, On Mahler’s partition problem, *Proc. Section of Sciences, Koninklijke Nederlandse Akademie van Wetenschappen* **51**(6) (1948), 659–669. Primary scan: <https://pure.tue.nl/ws/portalfiles/portal/4342300/597482.pdf>.
- [10] P. Dumas and P. Flajolet, Asymptotique des récurrences mahlériennes: le cas cyclotomique, *J. Théor. Nombres Bordeaux* **8** (1996), 1–30. <https://doi.org/10.5802/jtnb.154>.
- [11] C. Banderier, H.-K. Hwang, V. Ravelomanana, and V. Zacharovas, Analysis of an exhaustive search algorithm in random graphs and the $n^{c \log n}$ -asymptotics, *SIAM J. Discrete Math.* **28** (2014), 342–371. Preprint and versions: <https://arxiv.org/abs/1207.6549>.
- [12] ProveIt, *Transseries and inversion*,
 Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/
 transseries_and_inversion.tex; Part 0 (p0:def:model, p0:def:three-inverses,
 p0:thm:staircase), plt:def:lw-monomial-log-datum, plt:thm:lw-template. [Added in
 the write.]