

The First- κ Coefficients

Omitted Types, Singular Compression, and Completion in Surreal and Surcomplex Hahn Fields

Research manuscript prepared for Vladimir Reshetnikov
OpenAI ChatGPT

22 September 2026

Abstract

Let κ be any uncountable cardinal, possibly singular, let $\Gamma \neq 0$ be a divisible ordered abelian group, and let $K_{\kappa,k} \subseteq k((t^\Gamma))$ consist of the Hahn series having fewer than κ nonzero coefficients, with $k = \mathbb{R}$ or $\mathbb{C}$. We give an explicit description of the one-variable types over this field that are realized in its full Hahn extension but omitted in the smaller field. The invariant is exactly the first κ nonzero terms of the series: these terms determine its complete ordered-field type in the real case and its complete valued-field type in the complex case. We compute the least number of formulas needed to witness omission of each such type and obtain $\text{cf}(\kappa)$, including at singular cardinals. In the pure complex field language, by contrast, all the missing elements have the same type and the corresponding number is $|K_{\kappa,\mathbb{C}}|$.

The same coefficient invariant gives the exact approximation-value cut, the least cardinality of an empty nest of closed valuation balls, and a concrete description of the valuation completion. Whenever $K_{\kappa,k}$ is proper, it is complete if and only if $\text{cf}(\Gamma) \neq \text{cf}(\kappa)$; it is never spherically complete, and its first empty ball nests have cardinality $\text{cf}(\kappa)$. We also compute the precise valuation loss of every linear retraction from $K + Kx$ to K . Explicit Conway normal forms give surreal examples with countable omitted types at $\kappa = \aleph_\omega$, as well as complete fields exhibiting the same countable ball obstruction.

The constructions, general approximation-type theory, and quantifier-elimination inputs are established mathematics. The proposed contribution is the joint first- κ classification and its exact cardinal, topological, and linear consequences. Full proofs are supplied. This is an unrefereed research contribution with a limited priority audit, not a claimed solution of a named longstanding conjecture or a proof-assistant-certified result.

This single-source report is built from one manuscript, pinned at repository commit 465a54b. [Section 1.4](#) records its provenance, its notation conventions against the collection's, and its relation to neighbouring reports; [Section 11.6](#) collects its non-claims.

Conventions. All Hahn fields and parameter sets are sets in ZFC. All topological assertions use the intrinsic Hahn valuation topology, not the subspace topology inherited from the full class **No**. The support bound is a *cardinality* bound, not a bound on the magnitude of the exponents.

Contents

1	The problem and the contribution	4
1.1	Standing setup	4
1.2	A guide to the main results	4
1.3	What is established, and what is proposed here?	5
1.4	This report in the collection	5
2	Hahn arithmetic and singular support bounds	7
2.1	The support conventions	7
2.2	A small exponent group contains all finite algebraic data	7
2.3	The imported logical tools	8
3	The first-κ approximation theorem	8
4	Complete one-types and their fibers	9
4.1	The complex valued-field language	9
4.2	The real ordered-field language	10
5	The exact number of formulas needed for omission	11
5.1	Formula-by-formula stabilization	11
5.2	What forgetting the valuation does	12
5.3	Singular compression along elementary chains	12
6	The precise spherical-completeness threshold	13
7	The valuation completion and its cofinality criterion	14
7.1	An explicit completion inside the full Hahn field	14
7.2	The group-theoretic existence calculation	15
8	An exact two-dimensional linear obstruction	16
9	Explicit examples at regular and singular cardinals	17
9.1	A concrete family of exponent groups	17
9.2	A singular bound with a countable missing Cauchy limit	18
9.3	The same countable obstruction in a complete field	18
9.4	A completion strictly between the two support fields	18
9.5	A regular bound with a genuinely uncountable obstruction	19
9.6	The rank-one boundary case	19
10	Transfer to actual surreal and surcomplex numbers	19
10.1	The normal-form realization	19
10.2	The explicit groups already live in the surreal field	20
10.3	The topology and language boundaries	20
11	Scope, provenance, and a formalization route	20
11.1	Repository boundary	20
11.2	Literature and novelty ledger	21
11.3	A feasible proof-assistant decomposition	21
11.4	Further questions, not conclusions	22
11.5	Corrections to the manuscript's repository statements	22
11.6	What is not claimed	23
12	Conclusion	23
A	Proof-risk checklist and finite verification	24

References**24**

1 The problem and the contribution

A series can fail to belong to a subfield for very different reasons. It may require new coefficients, a larger value group, or too many terms. The last obstruction is especially natural for surreal numbers: Conway normal form gives each individual surreal a set-sized, reverse-well-ordered sequence of monomials, but no fixed cardinal bounds the normal forms of all surreals. A support restriction therefore provides a set-theoretic way to organize exact surreal arithmetic.

This paper asks a more precise question than whether the support-restricted field is proper:

What can first-order formulas over a cardinally support-restricted field detect about a missing series, and how many formulas are needed to force its absence from the field?

The answer will retain the coefficients of an initial segment, not merely the order type or cofinality of the support. Its cardinal consequence is that a singular support bound compresses a large amount of missing information into only $\text{cf}(\kappa)$ conditions.

1.1 Standing setup

Fix a nonzero divisible ordered abelian group Γ , an uncountable cardinal κ , and write $\mu = \text{cf}(\kappa)$. Cardinals are identified with their initial ordinals. For $k \in \{\mathbb{R}, \mathbb{C}\}$, put

$$F_k = k((t^\Gamma)), \quad K_{\kappa,k} = \{f \in F_k : |\text{supp } f| < \kappa\}. \quad (1.1)$$

The support is well ordered in the increasing order of Γ , and

$$v(f) = \min \text{supp } f \quad (f \neq 0), \quad v(0) = \infty.$$

Thus larger values mean smaller elements. For real coefficients, the sign is the sign of the coefficient at the least exponent. We reserve $\text{cf}(\Gamma)$ for the least cardinality of a cofinal subset of Γ toward $+\infty$.

If $x \in F_k \setminus K_{\kappa,k}$, enumerate its support increasingly as

$$x = \sum_{\alpha < \lambda} c_\alpha t^{s_\alpha}, \quad c_\alpha \neq 0, \quad s_\alpha < s_\beta \quad (\alpha < \beta), \quad \lambda \geq \kappa. \quad (1.2)$$

Its *first- κ prefix* is

$$T_\kappa(x) = \sum_{\alpha < \kappa} c_\alpha t^{s_\alpha}. \quad (1.3)$$

This series itself is not in $K_{\kappa,k}$. Every proper initial segment of it is.

1.2 A guide to the main results

The following statements are proved separately below. The real types use the language of ordered rings; the complex valued types use the language of rings with the binary relation $a \mid_v b$ meaning $v(a) \leq v(b)$. All parameters belong to the indicated smaller field.

Theorem 1.1 (Main theorem package). *In the standing setup, the following hold.*

(i) $K_{\kappa,\mathbb{R}}$ is real closed and $K_{\kappa,\mathbb{C}}$ is algebraically closed, even when κ is singular. Their inclusions in the full Hahn fields are elementary in the stated languages.

(ii) For missing elements x, y in the same full Hahn field,

$$\text{tp}(x/K_{\kappa,k}) = \text{tp}(y/K_{\kappa,k}) \iff T_\kappa(x) = T_\kappa(y).$$

(iii) For each such type p , the least size of a subset of p with no realization in $K_{\kappa,k}$ is exactly μ .

(iv) If $K_{\kappa,k} \neq F_k$, every nest of fewer than μ closed valuation balls meets, and some nest of μ such balls does not.

(v) *The completion is the subfield of F_k consisting of those f for which*

$$|\text{supp } f \cap (-\infty, \gamma)| < \kappa \quad (\gamma \in \Gamma).$$

If $K_{\kappa,k} \neq F_k$, this is a proper extension of $K_{\kappa,k}$ exactly when $\text{cf}(\Gamma) = \mu$.

The pure complex field language gives a deliberately contrasting answer: there is only one missing one-variable type, and its omission number is $|K_{\kappa,\mathbb{C}}|$ rather than μ ; see [Theorem 5.4](#). The linear application in [Theorem 8.1](#) provides an exact bound for every retraction $K + Kx \rightarrow K$, not merely an existence or nonexistence assertion.

1.3 What is established, and what is proposed here?

Cardinally bounded Hahn fields are not new. Kuhlmann and Shelah use κ -bounded Hahn fields for regular uncountable κ in their construction of exponential-logarithmic fields [1]. Related bounded constructions appear in the work of Berarducci, Kuhlmann, Mantova, and Matusinski on omega-maps [2]. We do not claim the construction, Hahn arithmetic, or its familiar regular-cardinal instances as new.

General approximation-type theory already describes immediate extensions by nests of balls. Kuhlmann’s account includes their relation to first-order one-types [3]; the passage from polynomial data to complete types uses classical quantifier elimination [5, 6, 7]. The contribution proposed here is an explicit *cardinal-prefix calculation* for the fields in (1.1), together with its sharp singular-cardinal and completion consequences. In particular, the exact omission number is proved for arbitrary formulas, not only for the displayed ball conditions.

The manuscript’s repository comparison was made at its pinned revision; see [Section 11](#). Its report on transcendence over bounded support concerns order-bounded supports and their fraction field, not the cardinal restriction in (1.1). Its foundations and Hahn-vector-space reports supply a natural context, but the present proofs do not depend on a new, unaudited theorem from those manuscripts. A limited search did not identify the entire theorem package in the material reviewed. This is a scoped novelty assessment, not a guarantee that no equivalent statement exists. The comparison with the collection as it now stands, including countable-support analogues of [Theorems 7.1](#) and [7.4](#) in two reports that the manuscript’s comparison did not mention, is in [Section 1.4](#).

1.4 This report in the collection

Provenance. This report is built from one manuscript, *The First- κ Coefficients: Omitted Types, Singular Compression, and Completion in Surreal and Surcomplex Hahn Fields*, dated 22 September 2026 (21 pages, author line “OpenAI ChatGPT”), delivered as the archive `surreal_first_kappa_types` (item 04 of batch 22) and placed as raw files in commit 7b5f934. The report was assembled in commit 68e2960. Its repository comparison is pinned at commit 465a54b479a1ee842cbf7db1689a7d2f6bfe25e1. It is not a merge: no other manuscript proves its results, and the nearest material in the collection is cross-referenced below, not merged. The statements, proofs, limitations and priority caveats are the manuscript’s. During assembly the following changed.

- Every label received the prefix `fkc`; all 72 labels of the manuscript are kept, and no statement was renumbered or removed.
- Added: this subsection; [Remark 7.6](#) (the countable analogues in other reports); [Remark 9.6](#) (the groups Γ_θ elsewhere in the collection); a pointer after the rank-one boundary case; the corrections of [Section 11.5](#); the collected non-claims of [Section 11.6](#); bibliography entries for collection reports; and the file paths of [Appendix A](#), which now name this directory’s layout.
- The program code/`verify.py`, the script code/`build.sh`, the recorded run data/`verification.json`, the build record data/`build_audit.json` and the file `research_audit.md` are the delivered files, unchanged. The delivered member README and PDF are not shipped; data/`build_audit.json` describes the delivered 21-page PDF, not the PDF of this report.
- Cross-references are now printed with the right names. All environments share one counter, and in the delivered build every reference to a proposition, lemma, corollary or example was printed as “theorem”, and the reference to the appendix as “section”. The preamble now declares the names; no statement changed.
- The family `surcomplex/` was chosen over `surreal/` because the results are stated uniformly for $k = \mathbb{R}$ and $k = \mathbb{C}$; this was the only placement choice.

Notation. No symbol of the manuscript was renamed. Its symbols are fully subscripted and local to this report, and the following conventions fix them against the collection’s notation guide (docs/NOTATION.md) and neighbouring reports.

- $F_k = k((t^\Gamma))$ denotes the full Hahn field for *both* $k = \mathbb{R}$ and $k = \mathbb{C}$. The guide writes $F_\Gamma = \mathbb{R}((t^\Gamma))$ for the real workspace and $K_\Gamma = \mathbb{C}((t^\Gamma))$ for the complex one. Thus $F_\mathbb{R}$ here is the guide’s F_Γ , and $F_\mathbb{C}$ here is the guide’s K_Γ . The subscript Γ is suppressed because Γ is fixed throughout.
- $K_{\kappa,k}$ is the *cardinally bounded subfield* of (1.1), real or complex. It is never the guide’s full complex workspace K_Γ . From Section 8 on, K and F abbreviate $K_{\kappa,k}$ and F_k .
- $L_{\kappa,k}$ is the completion of (7.1); D_x is the approximation cut of (3.1), not the quaternion algebra D_Γ of the surquaternions report; $E_x = K + Kx$ is a two-dimensional valued space, neither the algebraic scalar extension $E_\Gamma(V)$ of the three-duals report nor the ring of entire functions of the entire-functions report.
- κ is an uncountable cardinal and $\mu = \text{cf}(\kappa)$ is a regular cardinal. Neither is a measure, a residue field (the entire-functions report writes κ_n for residue fields) or a valuation gain (the expanding-dynamics report writes $\kappa = v(q)$).
- I_D in (3.3) is an additive subgroup and not an ideal; it is unrelated to the error ideal I_κ of the expanding-dynamics report. $T_\kappa(x)$ is an external invariant, not an operation of any language used here.
- $\text{cf}(\Gamma)$ is cofinality toward $+\infty$; $v = \min \text{supp}$, larger values mean smaller elements, and $t^\gamma \mapsto \omega^{-j(\gamma)}$. These agree with the guide.

Four words carry more than one meaning, and each tempting false reading is excluded here.

- *Bounded.* A support bound is a bound on the *cardinality* of the support. In the report on transcendence over bounded support, “bounded support” means bounded above in the exponent order, and that report says explicitly that it is not a cardinal restriction.
- *Complete.* A *complete type* is a maximal consistent set of formulas; a *complete field* is complete for Cauchy nets of the valuation uniformity; a *spherically complete field* has nonempty intersection for every nest of closed balls. Theorem 7.4 shows that the last two differ: a proper $K_{\kappa,k}$ is never spherically complete but can be complete.
- *Closure.* Theorem 7.1 concerns topological closure inside F_k ; Proposition 2.2 concerns real and algebraic closedness.
- *Omitted.* An omitted type here is a one-type over the set-sized field $K_{\kappa,k}$, realized in F_k and not in $K_{\kappa,k}$. It is not an omitted cut of a proper-class field.

Neighbouring reports. The comparisons below were checked against the current collection. None of them is used in a proof, and none proves or refutes a statement here.

- foundations-and-computation/surreal-fields-across-universes (prefix univ:) [10] studies saturation and omitted cuts of the old proper-class field $\mathbf{No}^M$ inside $\mathbf{No}^N$ for inner models $M \subseteq N$. It shares the vocabulary of omitted types, saturation and cofinality spectra, and it too contrasts languages, but its objects are different: it has no Hahn field, no support bound and no completion, and its topology is the fine order topology of proper classes. This report answers none of its questions. It was added after this manuscript’s pin.
- surreal/transcendence-over-bounded-support (prefix bst:) [11] works over the order-bounded base $\{f : \text{supp } f \text{ bounded above}\}$ and its fraction field. It explicitly excludes cardinal support bounds (its definition of the base, labelled bst:eq:base, and its list of conventions). Its corollary bst:cor:optimal is consistent with this report: its proof uses the fact that a support of cardinality less than $\text{cf}(\Gamma)$ is bounded above. This report answers none of its questions and is not a new version of its independence theorem.
- surcomplex/three-duals-of-hahn-vector-spaces (prefix duals:) [12] proves the vector-space analogue with “finite coefficient rank” in place of “fewer than κ support terms”; see Remark 7.6. Its full-Hahn spherical completeness, duals:prop:spherical, is the gluing argument that Lemma 6.1 refines by a cardinal bound.
- surcomplex/rank-one-berkovich (bare labels) [13] proves spherical completeness of $\mathbb{C}((t^\mathbb{R}))$ (its label prop:spherical) and describes the closure of the finite-support series in $\mathbb{C}((t^\mathbb{Q}))$; see Remark 7.6 and the rank-one boundary case in Section 9.

- `surcomplex/entire-functions-at-arbitrary-rank` (prefix `ent:`) [14] uses the group $\Gamma_\infty = \bigoplus_{j \geq 0} \mathbb{Q}e_j$ with $e_j = \omega^j$ and the group Γ_{ω_1} . These are the groups Γ_ω and Γ_{ω_1} of (9.1), with the same embedding in **No**; see Remark 9.6.
- `foundations-and-computation/foundations` (prefix `found:`) [15] distinguishes the intrinsic Hahn-workspace topology from the full fine topology (`found:prop:twotopologies`) and proves that small subsets and small-index nets are discrete in the latter (`found:thm:discrete`). Section 10 relies on that distinction.
- `surcomplex/hidden-negative-hermitian-directions` (prefix `hnd:`) [16] asks, in an unlabelled question, which support conditions on a support-family subfield $E \subseteq F_\Gamma$ give order density and which support invariants bound the degree of approximants to an omitted valuation ball. $K_{\kappa, \mathbb{R}}$ is such a subfield, but this report computes approximation cuts and empty ball nests, not order density or degrees of approximants, so that question is not answered here.

2 Hahn arithmetic and singular support bounds

2.1 The support conventions

A Hahn series is a function $f : \Gamma \rightarrow k$ whose nonzero set is well ordered. Addition is coefficientwise, and multiplication is the Hahn convolution. The well-ordering and finite-contribution properties needed for multiplication are the standard Hahn–Neumann support facts. No ordinary convergence of finite partial sums is implicit in this notation.

For $\gamma \in \Gamma$, define

$$f_{<\gamma} = \sum_{g < \gamma} f_g t^g, \quad f_{\leq \gamma} = \sum_{g \leq \gamma} f_g t^g.$$

Subseries are Hahn series because a subset of a well-ordered set is well ordered. The essential valuation identity is

$$v(f - g) \geq \gamma \iff f_h = g_h \text{ for every } h < \gamma. \quad (2.1)$$

Strict inequality means agreement at every $h \leq \gamma$. When $v(f) \neq v(g)$,

$$v(f + g) = \min\{v(f), v(g)\}. \quad (2.2)$$

Both formulas follow by looking at the least exponent with nonzero coefficient.

2.2 A small exponent group contains all finite algebraic data

Lemma 2.1 (Small-workspace lemma). *Let $A \subseteq F_k$ be finite and suppose each element of A has support of cardinality less than κ . There is a divisible ordered subgroup $H \subseteq \Gamma$ with $|H| < \kappa$ such that*

$$A \subseteq k((t^H)) \subseteq K_{\kappa, k}.$$

Proof. Let S be the union of the supports of the finitely many elements of A . Then $|S| < \kappa$. Take its rational linear span H in Γ . Since ordered abelian groups are torsion free and Γ is divisible, this span is a well-defined divisible subgroup. Every element of H is a finite rational linear combination of members of S , so

$$|H| \leq \max\{|S|, \aleph_0\} < \kappa.$$

The last strict inequality uses uncountability, not regularity, of κ . Every support contained in H has cardinality at most $|H|$, which proves the second inclusion; the first is immediate from the choice of S . $\square$

Proposition 2.2 (Algebraic closedness and real closedness). *$K_{\kappa, k}$ is a subfield of F_k . Moreover, $K_{\kappa, \mathbb{C}}$ is algebraically closed, $K_{\kappa, \mathbb{R}}$ is real closed, and*

$$K_{\kappa, \mathbb{R}}[i] = K_{\kappa, \mathbb{C}}.$$

The valuation of $K_{\kappa, k}$ has value group Γ and residue field k , so $F_k/K_{\kappa, k}$ is immediate.

Proof. For two inputs, apply [Lemma 2.1](#). Their sum, product, and, for a nonzero input, inverse lie in the Hahn field $k((t^H))$. This proves the subfield assertion without any union of κ many supports.

We use the classical theorem that a full Hahn field with algebraically closed coefficient field and divisible exponent group is algebraically closed; see [4, Corollary 4]. Given a polynomial with coefficients in $K_{\kappa, \mathbb{C}}$, the small-workspace lemma puts all its coefficients in $\mathbb{C}((t^H))$. The polynomial splits there, and every root has support of size at most $|H| < \kappa$. Thus it splits in $K_{\kappa, \mathbb{C}}$.

Complex coefficients split uniquely into real and imaginary parts. Each resulting support is a subset of the original support, while the union of two supports of size less than κ still has size less than κ . Hence $K_{\kappa, \mathbb{R}}[i] = K_{\kappa, \mathbb{C}}$. The field $K_{\kappa, \mathbb{R}}$ is ordered by its leading coefficient, and its quadratic extension by i is algebraically closed. By the real-closed-field criterion, $K_{\kappa, \mathbb{R}}$ is real closed. Equivalently, one can apply the usual real Hahn theorem inside each small H .

Finally, every monomial t^γ and every constant from k has finite support. Thus all values in Γ and all residues in k already occur in $K_{\kappa, k}$. They are also the value group and residue field of F_k , proving immediacy. $\square$

Remark 2.3 (Why singular cardinals cause no algebraic failure). A countable union of sets of size less than a singular κ can have size κ . That fact does not invalidate [Proposition 2.2](#): a fixed polynomial has only finitely many coefficients, and all its roots can be found in *one* exponent group of size less than κ . The relevant bound is not an uncontrolled union along a root construction. This distinction is central later: algebraic equations stay in one small workspace, whereas an infinite type may use parameters from many increasingly large workspaces.

2.3 The imported logical tools

We use the following standard quantifier-elimination statements. Real closed fields eliminate quantifiers in the ordered-ring language [5]. Algebraically closed fields eliminate quantifiers in the ring language. Algebraically closed nontrivially valued fields eliminate quantifiers in the ring language expanded by valuation divisibility [6, 7]. The last language has no named coefficient extraction, truncation operation, exponential, or cross-section.

It follows that $K_{\kappa, \mathbb{R}} \leq F_{\mathbb{R}}$ as ordered fields and $K_{\kappa, \mathbb{C}} \leq F_{\mathbb{C}}$ both as pure fields and as valued fields. Here $\leq$ means elementary substructure. Each valuation is nontrivial because $\Gamma \neq 0$.

We will only need the following concrete consequences. A formula in one variable over a real closed field is equivalent to a Boolean combination of polynomial sign conditions. A valued-field formula in one variable over an algebraically closed valued field is equivalent to a Boolean combination of polynomial equations and comparisons $v(P(X)) \leq v(Q(X))$. Over an algebraically closed field, a one-variable definable set is finite or cofinite. These are background inputs, not claims of new quantifier elimination.

3 The first- κ approximation theorem

For a missing element x as in (1.2), define

$$S_\kappa(x) = \{s_\alpha : \alpha < \kappa\}, \quad D_x = \{g \in \Gamma : g \leq s_\alpha \text{ for some } \alpha < \kappa\}. \quad (3.1)$$

The set D_x is an initial segment of Γ with no greatest element. It can be all of Γ or a proper initial segment. Do not replace D_x by a presumed supremum in Γ : that supremum need not exist.

Theorem 3.1 (Exact approximation values). *For every $x \in F_k \setminus K_{\kappa, k}$,*

$$A(x/K_{\kappa, k}) := \{v(x - a) : a \in K_{\kappa, k}\} = D_x, \quad \text{cf}(D_x) = \text{cf}(\kappa) = \mu. \quad (3.2)$$

In particular, x has no best approximation in $K_{\kappa, k}$ in the sense of a largest value of $v(x - a)$.

Proof. Fix $a \in K_{\kappa, k}$. It cannot have the same nonzero coefficient as x at every s_α for $\alpha < \kappa$, because that would place κ distinct elements in $\text{supp } a$. Choose $\alpha < \kappa$ where those coefficients differ. The least exponent where x and a differ is at most s_α . Thus $v(x - a) \in D_x$.

Conversely, suppose $g \in D_x$ and choose $\alpha < \kappa$ with $g \leq s_\alpha$. The support of $x_{<g}$ is contained in $\{s_\beta : \beta < \alpha\}$, so it has size less than κ . Choose $b \in k$ different from the coefficient x_g , and put

$$a = x_{<g} + bt^g.$$

This lies in $K_{\kappa,k}$, agrees with x below g , and differs at g . Hence $v(x - a) = g$, proving the equality.

Choose a strictly increasing cofinal family $(\alpha_i)_{i < \mu}$ in the ordinal κ . Then $(s_{\alpha_i})_{i < \mu}$ is cofinal in D_x , so $\text{cf}(D_x) \leq \mu$. For the reverse bound, suppose $E \subseteq D_x$ is cofinal and $|E| < \mu$. For each $e \in E$, choose $\beta_e < \kappa$ with $e \leq s_{\beta_e}$. Since $|E| < \text{cf}(\kappa)$, these indices are bounded below κ . A later s_β exceeds every member of E , contradicting cofinality. Thus $\text{cf}(D_x) = \mu$.

Finally, for every s_α in the initial segment there is a later s_β , because the infinite initial ordinal κ is a limit ordinal. Therefore D_x has no greatest element, which proves the final assertion. $\square$

Definition 3.2 (The invisible tail subgroup). For an initial segment $D \subseteq \Gamma$, set

$$I_D = \{0\} \cup \{h \in F_k \setminus \{0\} : v(h) > d \text{ for all } d \in D\}. \quad (3.3)$$

This is an additive subgroup. No assertion that it is an ideal of the field F_k is intended. If $D = \Gamma$, then $I_D = \{0\}$.

Lemma 3.3 (Prefix fibers). For missing elements $x, y \in F_k \setminus K_{\kappa,k}$, the following are equivalent:

$$T_\kappa(x) = T_\kappa(y), \quad x - y \in I_{D_x}, \quad y \in T_\kappa(x) + I_{D_x}.$$

Every element of $T_\kappa(x) + I_{D_x}$ is missing from $K_{\kappa,k}$.

Proof. Equal prefixes mean that the two series agree at every exponent at most s_α , for every $\alpha < \kappa$. Any nonzero difference therefore has value larger than all those exponents, hence larger than all of D_x . Conversely, such a difference changes none of the first κ nonzero coefficients and introduces no additional term before them. It cannot change the first- κ prefix. The difference $x - T_\kappa(x)$ itself belongs to I_{D_x} , giving the third formulation. Each series in that coset still contains all κ nonzero terms of the prefix, so it cannot lie in $K_{\kappa,k}$. $\square$

Remark 3.4 (A cut without coefficients is insufficient). Two series may have identical sets D_x but different first- κ prefixes. For example, changing a single coefficient in the prefix does not change D_x . Approximation values forget the centers of the balls; the full first-order type will remember the coefficients through those centers. The distinction between D_x and $T_\kappa(x)$ prevents an erroneous classification solely by cofinality or valuation rank.

4 Complete one-types and their fibers

A complete one-type over K is the collection of all formulas $\varphi(X)$ with parameters in K that hold of an element in an elementary extension. We only classify the types realized in the specified full Hahn extension F_k . Other elementary extensions can add new residue elements or values, and their types are outside this classification.

4.1 The complex valued-field language

Theorem 4.1 (First- κ classification: valued case). For $x, y \in F_{\mathbb{C}} \setminus K_{\kappa, \mathbb{C}}$,

$$\text{tp}_{\text{ACVF}}(x/K_{\kappa, \mathbb{C}}) = \text{tp}_{\text{ACVF}}(y/K_{\kappa, \mathbb{C}}) \iff T_\kappa(x) = T_\kappa(y). \quad (4.1)$$

The set of realizations in $F_{\mathbb{C}}$ of the type of x is exactly

$$T_\kappa(x) + I_{D_x}.$$

Proof. Suppose first that the prefixes agree. By Lemma 3.3, either $x = y$ or $v(x - y)$ is greater than every element of D_x . For each $a \in K_{\kappa, \mathbb{C}}$, Theorem 3.1 gives $v(x - a) \in D_x$, and therefore

$$v(y - a) = v(x - a), \quad \text{lc}(y - a) = \text{lc}(x - a). \quad (4.2)$$

Indeed, the two differences differ only at strictly higher valuation.

Every nonzero polynomial over $K_{\kappa, \mathbb{C}}$ factors as

$$P(X) = b \prod_{j=1}^n (X - a_j), \quad b \in K_{\kappa, \mathbb{C}}^\times, \quad a_j \in K_{\kappa, \mathbb{C}},$$

with multiplicities retained. Neither x nor y is a root, since $K_{\kappa, \mathbb{C}}$ is algebraically closed and they do not belong to it. Formula (4.2) implies

$$v(P(x)) = v(b) + \sum_j v(x - a_j) = v(b) + \sum_j v(y - a_j) = v(P(y)).$$

The zero polynomial is handled separately, with value ∞ . Thus x and y satisfy the same polynomial equations and the same comparisons of polynomial valuations. Quantifier elimination gives equality of their complete valued-field types.

Conversely, suppose the types agree. For every $\alpha < \kappa$, the element

$$b_\alpha = x_{\leq s_\alpha}$$

has fewer than κ support terms and lies in $K_{\kappa, \mathbb{C}}$. The following is a formula over $K_{\kappa, \mathbb{C}}$, using t^{s_α} as a parameter to name its value:

$$v(X - b_\alpha) > v(t^{s_\alpha}). \quad (4.3)$$

It holds of x , hence of y . The two series consequently agree at every exponent at most s_α , for all $\alpha < \kappa$. Their prefixes agree. The assertion about the entire realization set follows from Lemma 3.3, which also excludes realizations in $K_{\kappa, \mathbb{C}}$. $\square$

Corollary 4.2 (Rational-function data). *If x, y have the same first- κ prefix, the isomorphism*

$$K_{\kappa, \mathbb{C}}(x) \longrightarrow K_{\kappa, \mathbb{C}}(y), \quad x \longmapsto y,$$

fixing $K_{\kappa, \mathbb{C}}$ preserves both valuation and leading coefficient of every nonzero rational function. In the real case, the corresponding isomorphism of rational function fields is order preserving.

Proof. The missing elements are transcendental over their respective base fields. For complex coefficients, factor numerator and denominator into linear factors and apply (4.2); leading coefficients multiply and divide. In the real case, the same identity shows that $x - a$ and $y - a$ have the same sign for every a in the base. Factor real polynomials into real linear factors and monic irreducible quadratic factors, the latter positive everywhere. Signs of nonzero rational functions are therefore preserved. $\square$

4.2 The real ordered-field language

Theorem 4.3 (First- κ classification: ordered case). *For $x, y \in F_{\mathbb{R}} \setminus K_{\kappa, \mathbb{R}}$,*

$$\text{tp}_{\text{RCF}}(x/K_{\kappa, \mathbb{R}}) = \text{tp}_{\text{RCF}}(y/K_{\kappa, \mathbb{R}}) \iff T_\kappa(x) = T_\kappa(y). \quad (4.4)$$

The realizations in $F_{\mathbb{R}}$ of the type of x are exactly $T_\kappa(x) + I_{D_x}$.

Proof. Equal prefixes imply equal signs of $x - a$ and $y - a$ for all $a \in K_{\kappa, \mathbb{R}}$, by the leading-coefficient argument in (4.2). A nonzero real polynomial factors into linear factors and positive monic irreducible quadratic factors, together with a nonzero scalar. Thus its values at x and y have the same sign. Quantifier elimination for real closed fields now gives equal types.

For the reverse direction, set $b_\alpha = x_{\leq s_\alpha} \in K_{\kappa, \mathbb{R}}$. The ordered-ring formula

$$b_\alpha - t^{s_\alpha} < X < b_\alpha + t^{s_\alpha} \quad (4.5)$$

holds of x : its difference from b_α has valuation strictly greater than s_α and is therefore smaller in absolute value than the positive monomial t^{s_α} . Equal types imply that (4.5) holds of y as well.

The inequality $|y - b_\alpha| < t^{s_\alpha}$ forces agreement of y and b_α at all exponents strictly below s_α . If there were a least disagreement $g < s_\alpha$, its nonzero real leading coefficient times t^g would dominate t^{s_α} in absolute value. For any fixed $\beta < \kappa$, choose α strictly later than β . Agreement below s_α includes the coefficient at s_β and all earlier exponents. Consequently the first- κ prefixes agree. The realization-set description follows as before. $\square$

Corollary 4.4 (Counting and identifying the missing types). *In either language of Theorems 4.1 and 4.3, the missing one-types realized in F_k are in bijection with the Hahn series whose support has order type exactly κ . The bijection sends a type to its first- κ prefix. If $K_{\kappa,k} \neq F_k$, there are at least 2^κ such types.*

Proof. Each missing element has exactly one prefix of order type κ , and each such prefix is itself a missing element. The two type-classification theorems show that this correspondence is well defined and bijective. Fix an increasing support $(s_\alpha)_{\alpha < \kappa}$. Choosing each coefficient independently from $\{1, 2\}$ gives 2^κ different prefixes and therefore 2^κ types. $\square$

Remark 4.5 (Types are not asserted to be automorphism orbits in F_k). Equality of types guarantees conjugacy in a sufficiently homogeneous elementary extension. It does not, by itself, produce an automorphism of the particular full Hahn field F_k taking x to y . No homogeneity theorem for F_k , and no extension of a partial automorphism to the full class **No**, is used here.

5 The exact number of formulas needed for omission

Definition 5.1 (Omission number of a type). Let $M \leq N$ and let $p = \text{tp}(x/M)$ be omitted in M . Define

$$\text{om}_M(p) = \min\{|\Phi| : \Phi \subseteq p \text{ has no realization in } M\}. \quad (5.1)$$

This minimum exists because p itself is a set of formulas omitted in M . Each formula is finitary, but different formulas may use different parameters. This invariant is not the saturation cardinal of the entire structure.

5.1 Formula-by-formula stabilization

Choose a strictly increasing cofinal sequence $(\alpha_i)_{i < \mu}$ in κ , and put

$$a_i = x_{< s_{\alpha_i}} \in K_{\kappa,k}. \quad (5.2)$$

Then

$$v(x - a_i) = s_{\alpha_i}, \quad v(a_j - a_i) = s_{\alpha_i} \quad (i < j). \quad (5.3)$$

The second equality uses the nonzero coefficient c_{α_i} . Thus (a_i) is a strictly pseudo-Cauchy family. It need not be a Cauchy net: its error values may stay bounded in Γ .

Lemma 5.2 (Eventual elementary agreement). *In the real ordered-field or complex valued-field language, for every formula $\varphi(X)$ over $K_{\kappa,k}$ there is $i_0 < \mu$ such that*

$$F_k \models \varphi(a_i) \iff F_k \models \varphi(x) \quad (i_0 \leq i < \mu).$$

Proof. First consider a fixed $b \in K_{\kappa,k}$. By Theorem 3.1, $r = v(x - b)$ belongs to D_x . For all sufficiently large i , $s_{\alpha_i} > r$. Hence

$$v(a_i - b) = v(x - b), \quad \text{lc}(a_i - b) = \text{lc}(x - b).$$

In particular, $a_i \neq b$ eventually.

For a finite collection of nonzero complex polynomials over $K_{\kappa,\mathbb{C}}$, take their finitely many roots, with multiplicity ignored for this choice of a bound. Choose i_0 making the preceding identities hold for every root. For all $i \geq i_0$, each polynomial is nonzero at a_i and has the same value as at x . Consequently every Boolean combination of their polynomial equations and valuation comparisons stabilizes. Quantifier elimination reduces any fixed valued-field formula to such a finite Boolean combination.

Over $K_{\kappa,\mathbb{R}}$, instead factor the finitely many polynomials from a quantifier-free ordered formula into linear and positive irreducible quadratic factors. Their linear factors have stable signs by the first paragraph, and the quadratic signs are already constant. All polynomial sign conditions stabilize, and real quantifier elimination proves the assertion. Constant and identically zero polynomials require no stabilization condition. $\square$

Theorem 5.3 (Exact omission number). *For each $x \in F_\kappa \setminus K_{\kappa,k}$, in the ordered real or valued complex language,*

$$\boxed{\text{om}_{K_{\kappa,k}}(\text{tp}(x/K_{\kappa,k})) = \text{cf}(\kappa).} \quad (5.4)$$

More precisely, every family of fewer than μ formulas from the type is realized by a sufficiently late truncation a_i , whereas an explicitly given family of μ ball or interval formulas is omitted.

Proof. For the lower bound, let Φ be a subset of the type of cardinality less than μ . For each $\varphi \in \Phi$, choose a stabilization index i_φ from Lemma 5.2. The cardinal $\mu = \text{cf}(\kappa)$ is regular, so fewer than μ such indices are bounded by some $i_* < \mu$. Then a_i realizes every formula in Φ for all $i \geq i_*$. Elementarity lets us view this realization either in F_κ or in $K_{\kappa,k}$.

For the upper bound in the complex valued case, use the μ formulas

$$v(X - a_i) \geq v(t^{s_{\alpha_i}}) \quad (i < \mu). \quad (5.5)$$

They all hold of x . A common realization $a \in K_{\kappa,\mathbb{C}}$ would agree with x at every exponent below each s_{α_i} . Cofinality of the indices then forces agreement at all first κ nonzero terms, placing at least κ elements in $\text{supp } a$. This is impossible.

For the upper bound in the real case, put $b_i = x_{\leq s_{\alpha_i}}$ and use

$$b_i - t^{s_{\alpha_i}} < X < b_i + t^{s_{\alpha_i}} \quad (i < \mu). \quad (5.6)$$

These formulas belong to the type of x . As in the proof of Theorem 4.3, a common realization must agree with x below every s_{α_i} and hence on all first κ terms. Again it cannot lie in $K_{\kappa,\mathbb{R}}$. This proves the upper bound and the equality. $\square$

5.2 What forgetting the valuation does

Theorem 5.4 (Pure-field collapse). *Assume $K_{\kappa,\mathbb{C}} \neq F_{\mathbb{C}}$ and work in the pure ring language. All elements of $F_{\mathbb{C}} \setminus K_{\kappa,\mathbb{C}}$ realize the same complete one-type p_{gen} over $K_{\kappa,\mathbb{C}}$. Its omission number is*

$$\text{om}_{K_{\kappa,\mathbb{C}}}(p_{\text{gen}}) = |K_{\kappa,\mathbb{C}}|. \quad (5.7)$$

Proof. Since $K_{\kappa,\mathbb{C}}$ is algebraically closed, every element of $F_{\mathbb{C}} \setminus K_{\kappa,\mathbb{C}}$ is transcendental over it. Such an element satisfies precisely the generic one-type over the algebraically closed field: all nonzero polynomials over $K_{\kappa,\mathbb{C}}$ are nonzero there. Quantifier elimination makes this a complete type.

Every one-variable formula from the generic type defines a cofinite subset of $K_{\kappa,\mathbb{C}}$. Indeed, the definable subset is finite or cofinite, and a finite set definable over an algebraically closed base cannot contain a transcendental element in an elementary extension. Fewer than $|K_{\kappa,\mathbb{C}}|$ finite exceptional sets cannot cover the infinite field $K_{\kappa,\mathbb{C}}$. This remains true if $|K_{\kappa,\mathbb{C}}|$ is singular: for an infinite cardinal $\nu < |K_{\kappa,\mathbb{C}}|$, a union of ν finite sets has cardinality at most ν . Therefore fewer than $|K_{\kappa,\mathbb{C}}|$ formulas from p_{gen} have a common realization in $K_{\kappa,\mathbb{C}}$.

Conversely, the family $\{X \neq a : a \in K_{\kappa,\mathbb{C}}\}$ belongs to the generic type and has no realization in $K_{\kappa,\mathbb{C}}$. Its cardinality is $|K_{\kappa,\mathbb{C}}|$, proving the equality. $\square$

In particular, when κ is singular and $K_{\kappa,\mathbb{C}} \neq F_{\mathbb{C}}$, the same individual missing element has omission number $\text{cf}(\kappa) < \kappa \leq |K_{\kappa,\mathbb{C}}|$ in the valued language and omission number $|K_{\kappa,\mathbb{C}}|$ in the pure field language. This is a quantitative distinction between two descriptions of a surcomplex workspace, not a change in its underlying arithmetic.

5.3 Singular compression along elementary chains

Corollary 5.5 (Restriction of types is truncation of prefixes). *Let $\kappa < \lambda$ be uncountable cardinals, with the same k and Γ . Then*

$$K_{\kappa,k} \leq K_{\lambda,k} \leq F_k$$

in the real ordered or complex valued language. If $x \notin K_{\lambda,k}$, restriction of its type from $K_{\lambda,k}$ to $K_{\kappa,k}$ corresponds to the map

$$T_\lambda(x) \mapsto T_\kappa(x).$$

The two omission numbers are respectively $\text{cf}(\lambda)$ and $\text{cf}(\kappa)$.

Proof. The inclusions are elementary by [Proposition 2.2](#) and quantifier elimination. The assertions about prefixes and omission numbers follow from [Theorems 4.1, 4.3 and 5.3](#) applied at the two cardinals. $\square$

For example, moving from the bound $\aleph_1$ to the larger bound $\aleph_\omega$ can *decrease* the omission number from $\aleph_1$ to $\aleph_0$. There is no conflict with compactness: each parameter in the larger field is a single object but may encode a much longer initial segment. Countably many such parameters can encode $\aleph_\omega$ support positions. At the smaller bound, no analogous countable family of parameters can do so for these types.

6 The precise spherical-completeness threshold

A closed valuation ball in a valued field K is

$$B(a, \gamma) = \{z \in K : v(z - a) \geq \gamma\}, \quad a \in K, \quad \gamma \in \Gamma.$$

A *nest* is a nonempty set of such balls linearly ordered by inclusion. Spherical completeness means that every nest has nonempty intersection. The radii here are values in Γ , not necessarily real numbers; nests need not be sequences.

Lemma 6.1 (Coefficient gluing with a cardinal bound). *Let $\{B(a_j, \gamma_j) : j \in J\}$ be a nest of balls in $K_{\kappa, k}$. If $|J| < \text{cf}(\kappa)$, its intersection contains an element of $K_{\kappa, k}$.*

Proof. For any $i, j \in J$, nesting implies

$$v(a_i - a_j) \geq \min\{\gamma_i, \gamma_j\}.$$

Consequently, for a fixed $g \in \Gamma$, all the coefficients $(a_j)_g$ with $g < \gamma_j$ agree. Define f_g to be this common coefficient when such a j exists, and zero otherwise.

We first check that $\text{supp } f$ is well ordered. Let $U \subseteq \text{supp } f$ be nonempty, and choose $g_0 \in U$. There is j_0 with $g_0 < \gamma_{j_0}$, and on all exponents below γ_{j_0} the coefficients of f agree with those of a_{j_0} . Thus the nonempty set $U \cap (-\infty, g_0]$ is a subset of the well-ordered support of a_{j_0} and has a least element. That element is also least in U . Hence f is a Hahn series.

Moreover,

$$\text{supp } f \subseteq \bigcup_{j \in J} \text{supp } a_j.$$

The union on the right has size less than κ : a union of fewer than $\text{cf}(\kappa)$ sets, each of cardinality less than κ , still has cardinality less than κ . Therefore $f \in K_{\kappa, k}$. For each j , it agrees with a_j at every exponent below γ_j , so [\(2.1\)](#) gives $f \in B(a_j, \gamma_j)$. $\square$

The gluing argument without the final cardinal bound is the standard proof of spherical completeness for full Hahn fields; compare [\[4, Lemma 4\]](#). The extra point here is exactly where the union of supports remains below κ .

Theorem 6.2 (First empty nests). *If $K_{\kappa, k} \neq F_k$, the least cardinality of an empty nest of closed valuation balls in $K_{\kappa, k}$ is $\mu = \text{cf}(\kappa)$. In particular, $K_{\kappa, k}$ is not spherically complete.*

Proof. The lower bound is [Lemma 6.1](#). For the upper bound, choose a missing x and use the approximants from [\(5.2\)](#). Let

$$B_i = B(a_i, s_{\alpha_i}) \quad (i < \mu).$$

For $i < j$, the center difference has value s_{α_i} , and the radius value s_{α_j} is larger. The ultrametric inequality therefore gives $B_j \subseteq B_i$. The inclusion is strict: $a_i \notin B_j$. A common point would satisfy all formulas [\(5.5\)](#), whose common realization in $K_{\kappa, k}$ was shown impossible. Thus this is an empty nest of cardinality exactly μ . $\square$

Corollary 6.3 (Its intersection in the full Hahn field). *The balls just constructed, interpreted in F_k , have intersection*

$$\bigcap_{i < \mu} B_{F_k}(a_i, s_{\alpha_i}) = T_\kappa(x) + I_{D_x}.$$

Thus the same coset is a fiber of the type map and a full-field intersection of a canonical omitted ball nest.

Proof. The ball conditions say precisely that all coefficients below the cofinal thresholds s_{α_i} agree with those of x . Their union forces exactly the first- κ prefix and leaves precisely the tails in I_{D_x} free. $\square$

Remark 6.4 (Not a full saturation theorem). The ball theorem does not say that every type over fewer than μ parameters is realized. Our lower bound in [Theorem 5.3](#) applies to types realized in the fixed full Hahn extension. Other types may demand new values or residues. A field can have all the small ball intersections established here while failing a different saturation condition. Likewise, the pure-field omission number in [Theorem 5.4](#) is not contradicted by the smaller valued-field ball obstruction.

7 The valuation completion and its cofinality criterion

Completeness and spherical completeness differ because a Cauchy net must reach every valuation scale, whereas the radii of a nest can remain confined to a proper initial segment of the value group. In the present fields this distinction is determined exactly by the first- κ support.

7.1 An explicit completion inside the full Hahn field

Define

$$L_{\kappa,k} = \{f \in F_k : |\text{supp } f \cap (-\infty, \gamma)| < \kappa \text{ for every } \gamma \in \Gamma\}. \quad (7.1)$$

The condition is local in the *exponent order*, not a condition of local finiteness in an ordinary real variable.

Theorem 7.1 (Completion by locally small support). *The closure of $K_{\kappa,k}$ in F_k is $L_{\kappa,k}$. This is a complete valued subfield of F_k and, with the given inclusion of $K_{\kappa,k}$, is its Hausdorff valuation completion.*

Proof. Suppose f is in the closure. Given $\gamma \in \Gamma$, there is $a \in K_{\kappa,k}$ with $v(f - a) \geq \gamma$. By [\(2.1\)](#), the support of f below γ is contained in the support of a , so it has size less than κ . Thus $f \in L_{\kappa,k}$.

Conversely, if $f \in L_{\kappa,k}$, then $f_{<\gamma} \in K_{\kappa,k}$ for every γ . Moreover,

$$v(f - f_{<\gamma}) \geq \gamma.$$

Direct Γ by its increasing order. These truncations converge to f : given a target value β , choose $\gamma > \beta$. Hence f lies in the closure of $K_{\kappa,k}$.

The full Hahn field is complete; this follows from its classical spherical completeness, or directly from coefficient stabilization in a Cauchy net. For clarity, in a Cauchy net all coefficients up to any fixed exponent eventually agree simultaneously. Their eventual values define a Hahn series: any nonempty part of its proposed support has a least element by comparison below one selected exponent with a sufficiently late member of the net. The same stabilization proves convergence to that series.

The closure of a subfield in a topological field is a subfield: addition and multiplication are continuous, and inversion is continuous at each nonzero point. A closed subspace of the complete Hahn uniform space is complete. Since $K_{\kappa,k}$ is dense in its closure, this closure is its Hausdorff uniform completion, with the induced field and valuation structures. $\square$

Proposition 7.2 (Exactly which supports are added). *For $f \in F_k \setminus K_{\kappa,k}$, the following are equivalent:*

- (i) $f \in L_{\kappa,k}$;
- (ii) $\text{supp } f$ has order type exactly κ and is cofinal in Γ ;
- (iii) $D_f = \Gamma$.

In particular, every new completion element satisfies $T_\kappa(f) = f$.

Proof. Write the support increasingly as $(s_\alpha)_{\alpha < \lambda}$ with $\lambda \geq \kappa$. If $\lambda > \kappa$, the exponent s_κ exists, and the support strictly below it already has cardinality κ . This violates [\(7.1\)](#). Hence membership in $L_{\kappa,k}$ forces $\lambda = \kappa$.

If that support were not cofinal in Γ , choose γ strictly above it. Then the support below γ has size κ , again a contradiction. This proves (i) $\Rightarrow$ (ii).

If the support has order type κ and is cofinal, then for each γ there is $\alpha < \kappa$ with $\gamma \leq s_\alpha$. Its support below γ is contained in the earlier α positions, so it has size less than κ . Thus (ii) $\Rightarrow$ (i). Condition (ii) clearly implies (iii). Conversely, (iii) says that the first κ support positions are already cofinal in Γ ; there can be no later exponent s_κ above them all. The full support therefore has exactly order type κ , proving (iii) $\Rightarrow$ (ii). $\square$

7.2 The group-theoretic existence calculation

Lemma 7.3 (Long cofinal supports). *Suppose Γ contains a well-ordered subset of cardinality κ . Then it contains a cofinal well-ordered subset of order type κ if and only if*

$$\text{cf}(\Gamma) = \text{cf}(\kappa). \quad (7.2)$$

Proof. If a cofinal increasing sequence of order type κ exists, its cofinality is $\text{cf}(\kappa)$, and cofinal subsets have the same cofinality as the ambient linear order. More explicitly, a cofinal subsequence of κ indices gives the upper bound, and a smaller cofinal subset of Γ would, by choosing sequence entries above its members, give a smaller cofinal subset of κ . This proves necessity.

For sufficiency, put $\mu = \text{cf}(\Gamma) = \text{cf}(\kappa)$. Take an increasing sequence $(s_\alpha)_{\alpha < \kappa}$ in Γ . If it is already cofinal, there is nothing to prove. Otherwise, translating it by $-s_0$ gives a sequence $(u_\alpha)_{\alpha < \kappa}$ in an interval $[0, h)$ for some $h > 0$.

If κ is regular, then $\mu = \kappa$, and the definition of $\text{cf}(\Gamma)$ gives a strictly increasing cofinal sequence of length κ . Such a sequence can be built by choosing, at each stage, a point above the fewer than $\text{cf}(\Gamma)$ previous choices and above a prescribed cofinal seed.

Suppose now that κ is singular. Choose increasing infinite cardinals $\kappa_i < \kappa$ for $i < \mu$ with supremum κ . Construct a cofinal increasing sequence $(q_i)_{i < \mu}$ in Γ such that

$$q_j > q_i + h \quad (i < j < \mu).$$

At stage j , fewer than $\text{cf}(\Gamma)$ previously selected values $q_i + h$ have been specified, so they are bounded. Choose the next value above them and above the next member of a fixed cofinal seed.

Now concatenate the blocks

$$\{q_i + u_\alpha : \alpha < \kappa_i\} \quad (i < \mu).$$

Different blocks are strictly separated, and their union is cofinal because it contains every q_i . Its order type is the ordinal sum $\sum_{i < \mu} \kappa_i$. This sum equals the initial ordinal κ . Indeed, every proper initial collection of blocks has cardinality less than κ , since it uses fewer than $\text{cf}(\kappa)$ cardinals below κ ; the same is true after adding a proper initial part of the next block. Thus every proper initial segment of the concatenation has cardinality less than κ , while the whole concatenation has cardinality κ . Its order type is therefore exactly κ . $\square$

Theorem 7.4 (Completeness dichotomy). *The following exhaustive alternatives hold.*

- (i) *If Γ has no well-ordered subset of cardinality κ , then $K_{\kappa, \kappa} = F_\kappa$, so $K_{\kappa, \kappa}$ is complete and spherically complete.*
- (ii) *If Γ has such a subset, then $K_{\kappa, \kappa} \neq F_\kappa$ and*

$$\boxed{K_{\kappa, \kappa} \text{ is complete} \iff \text{cf}(\Gamma) \neq \text{cf}(\kappa).} \quad (7.3)$$

In this second alternative, the least size of an empty closed-ball nest is always $\text{cf}(\kappa)$, regardless of completeness.

Proof. A missing Hahn series yields a well-ordered support containing its first κ positions. Conversely, an increasing support of order type κ supports a Hahn series with all coefficients equal to one, which is missing from $K_{\kappa, \kappa}$. This proves the criterion for properness.

By Theorem 7.1 and Proposition 7.2, a proper $K_{\kappa, \kappa}$ fails to be complete exactly when Γ admits a cofinal increasing sequence of order type κ . By Lemma 7.3, under the properness hypothesis this is equivalent to equality of the two cofinalities. The final assertion is Theorem 6.2. $\square$

Remark 7.5 (The completion need not be the full Hahn field). A new completion element is required to have cofinal support of order type exactly κ . A series with κ support positions confined below a fixed exponent is never added by completion. Thus the completion is not obtained simply by replacing $< \kappa$ by $\leq \kappa$, nor by replacing κ with κ^+ . Example 9.4 gives an explicit strict chain $K_{\kappa,k} \subsetneq L_{\kappa,k} \subsetneq F_k$.

Remark 7.6 (The countable analogue elsewhere in the collection; added during assembly). The case $\kappa = \aleph_0$ is excluded here, since finite-support series do not form a field (Appendix A). Two other reports prove the countable analogue of Theorems 7.1 and 7.4 for other objects. In the report on three duals of Hahn vector spaces, the completion of the algebraic scalar extension $E_\Gamma(V)$ of an infinite-dimensional k -vector space V consists of the vectors with finite coefficient rank below every $\gamma \in \Gamma$ (duals:thm:completion), and it is strictly larger than $E_\Gamma(V)$ exactly when $\text{cf}(\Gamma) = \aleph_0$ (duals:thm:cofinality-completion). This is the shape of (7.1) and (7.3) with “finite coefficient rank” in place of “fewer than κ support terms” and $\aleph_0 = \text{cf}(\aleph_0)$ in place of $\text{cf}(\kappa)$. In the rank-one Berkovich report, the metric closure of the finite-support series in $\mathbb{C}((t^{\mathbb{Q}}))$ consists of the series whose support meets every $(-\infty, B]$ in a finite set, which is (7.1) with κ replaced by $\aleph_0$. Neither result is used here, and neither implies nor is implied by the theorems above: the objects there are a vector space over the full Hahn field and the ring of finite-support series, and the bound is countable.

8 An exact two-dimensional linear obstruction

The same approximation cut has a direct functional interpretation. Fix either coefficient field and abbreviate $K = K_{\kappa,k}$, $F = F_k$. For $x \in F \setminus K$, let

$$E_x = K + Kx \subseteq F$$

with the valuation inherited from F . The vectors $1, x$ are linearly independent over K , so E_x is two dimensional. This is a Γ -valued linear space, not an asserted real-normed Banach space.

Every K -linear retraction $P : E_x \rightarrow K$ of the inclusion $K \subseteq E_x$ has the form

$$P_c(a + bx) = a + bc \quad (a, b \in K) \quad (8.1)$$

for a unique $c \in K$. For $\delta \in \Gamma_{\geq 0}$, say that it has *valuation loss at most* δ when

$$v(P_c z) \geq v(z) - \delta \quad (z \in E_x). \quad (8.2)$$

Loss zero is the nonexpanding, or contractive, case in valuation notation.

Theorem 8.1 (Exact retraction-loss formula). *Put $r_c = v(x - c) \in D_x$. Then*

$$P_c \text{ satisfies (8.2)} \iff d \leq r_c + \delta \text{ for every } d \in D_x. \quad (8.3)$$

Consequently no P_c has loss zero. A continuous retraction exists if and only if D_x is bounded above in Γ , equivalently $x \notin L_{\kappa,k}$.

Proof. Vectors in K satisfy (8.2) automatically. A vector $a + bx$ with $b \neq 0$ is $b(x - d)$ for $d = -a/b \in K$. By linearity and multiplicativity of the valuation, it suffices to check

$$v(c - d) \geq v(x - d) - \delta \quad (d \in K). \quad (8.4)$$

Write $r_d = v(x - d)$. If $r_d < r_c$, then (2.2) gives $v(c - d) = r_d$, so (8.4) holds. If $r_d = r_c$, the ultrametric inequality gives $v(c - d) \geq r_c$, so it also holds. If $r_d > r_c$, the exact value is $v(c - d) = r_c$, and (8.4) is equivalent to $r_d \leq r_c + \delta$. As the set of all r_d is D_x , this is exactly (8.3).

There is no zero-loss retraction because D_x has no greatest element. If D_x is bounded above by b , then for any c the nonnegative value $\delta = b - r_c$ is an admissible loss. Such a bound implies continuity directly.

Conversely, any continuous K -linear map $P : E_x \rightarrow K$ has some uniform valuation-loss bound. Indeed, continuity at zero supplies $\beta \in \Gamma$ such that $v(z) > \beta$ implies $v(Pz) > 0$. Choose $\eta > 0$. For $z \neq 0$, multiplication by the scalar

$$q = t^{\beta + \eta - v(z)} \in K$$

makes $v(qz) = \beta + \eta > \beta$. If $Pz \neq 0$, linearity yields

$$v(Pz) > v(z) - \beta - \eta.$$

The same lower bound is automatic if $Pz = 0$. Thus $\delta = \max\{0, \beta + \eta\}$ is a loss bound. Applying (8.3) to a continuous retraction forces D_x to be bounded above. Finally, an initial segment of a linear order is unbounded exactly when it is the whole order, and Proposition 7.2 identifies $D_x = \Gamma$ with $x \in L_{\kappa,k}$. $\square$

Corollary 8.2 (A zero-or-full continuous-dual dichotomy). *Let $E'_x = \text{Hom}_{K,\text{cont}}(E_x, K)$. Then*

$$\dim_K E'_x = \begin{cases} 0, & x \in L_{\kappa,k} \setminus K, \\ 2, & x \in F_k \setminus L_{\kappa,k}. \end{cases}$$

In the second case every algebraic linear functional is continuous, although no retraction has valuation loss zero.

Proof. If $x \in L_{\kappa,k} \setminus K$, then K is dense in E_x : approximate x by elements of K and multiply and translate to approximate $a + bx$. If a continuous functional ℓ had $\ell(1) \neq 0$, dividing by $\ell(1)$ would produce a continuous retraction, contrary to Theorem 8.1. Hence ℓ vanishes on the dense subspace K and therefore vanishes everywhere, since the target is Hausdorff.

Suppose instead that D_x is bounded. Fix $c \in K$ and a loss bound δ for P_c . The coefficient functional $Q(a + bx) = b$ is continuous as well. For $z = a + bx$, the identity $z - P_c z = b(x - c)$ gives, whenever $b \neq 0$,

$$v(b) = v(z - P_c z) - r_c \geq \min\{v(z), v(P_c z)\} - r_c \geq v(z) - \delta - r_c.$$

Replacing the right-hand loss by a larger nonnegative value if necessary gives a uniform continuity bound. Since P_c and Q span the algebraic dual of E_x , every linear functional is continuous and the dual dimension is two. $\square$

This application does not claim a new general Hahn–Banach theorem. The mechanism is the elementary best-approximation obstruction: contractivity of P_c would require c to approximate x at least as well as every other base-field element. The contribution within the present package is that the full set of permitted losses, and the continuous-dual dichotomy, are explicitly calculated from the first- κ support cut.

9 Explicit examples at regular and singular cardinals

9.1 A concrete family of exponent groups

For an infinite limit ordinal θ , let

$$\Gamma_\theta = \bigoplus_{\alpha < \theta} \mathbb{Q}e_\alpha \tag{9.1}$$

with finite supports and the order determined by the *largest* nonzero coordinate. Thus $0 < e_\alpha \ll e_\beta$ for $\alpha < \beta$: every positive integral multiple of e_α is less than e_β . This is a divisible ordered abelian group.

Lemma 9.1. *For an infinite limit ordinal θ ,*

$$\text{cf}(\Gamma_\theta) = \text{cf}(\theta).$$

Moreover, $(e_\alpha)_{\alpha < \theta}$ is increasing and cofinal in Γ_θ .

Proof. Every group element has finite support. An e_β with index larger than all indices in that support exceeds the group element, proving cofinality of the displayed sequence. Taking a cofinal subset of indices gives $\text{cf}(\Gamma_\theta) \leq \text{cf}(\theta)$. Conversely, a set of fewer than $\text{cf}(\theta)$ group elements uses a bounded set of coordinate indices: it is a union of fewer than $\text{cf}(\theta)$ finite sets. Some later e_β exceeds them all, so the set cannot be cofinal. This proves the reverse inequality. $\square$

9.2 A singular bound with a countable missing Cauchy limit

Example 9.2. Let $\kappa = \aleph_\omega$, $\Gamma = \Gamma_\kappa$, and take either $k = \mathbb{R}$ or $\mathbb{C}$. Define

$$x_\kappa = \sum_{\alpha < \kappa} t^{e_\alpha}, \quad a_n = \sum_{\alpha < \aleph_n} t^{e_\alpha} \quad (n < \omega). \quad (9.2)$$

Each a_n belongs to $K_{\kappa,k}$, whereas x_κ does not. Furthermore,

$$v(x_\kappa - a_n) = e_{\aleph_n}, \quad v(a_m - a_n) = e_{\aleph_n} \quad (n < m).$$

These values are cofinal in Γ_κ , so (a_n) is a Cauchy sequence with limit x_κ in the full Hahn field and no limit in $K_{\kappa,k}$.

All algebraic equations appropriate to real closedness or algebraic closedness can still be solved in the smaller field, by [Proposition 2.2](#). The omitted type of x_κ in the ordered real or valued complex language needs exactly countably many formulas. Its realization set in this particular full Hahn field is a singleton, because $D_{x_\kappa} = \Gamma_\kappa$ and $I_{D_{x_\kappa}} = \{0\}$. This does not mean the type is isolated by a single formula: [Theorem 5.3](#) proves the opposite.

For the two-dimensional valued space $K + Kx_\kappa$, the continuous K -linear dual is zero. Thus even two algebraic coordinates need not supply continuous coordinate functionals over this incomplete base field with its inherited valuation.

In fact, the completion in [Example 9.2](#) is the entire full Hahn field. Every bounded interval in Γ_κ has cardinality less than κ . To see this, choose an index $\beta < \kappa$ at least as large as all coordinate indices in its two endpoints. A group element with a nonzero coordinate beyond β lies strictly above both endpoints or strictly below both, according to the sign of its last coordinate. The interval is consequently contained in the rational span of $\{e_\alpha : \alpha \leq \beta\}$, a set of size less than κ . Since a nonempty Hahn support has a least exponent, each initial slice of it is contained in such a bounded interval. Thus every Hahn series has locally small support, and [Theorem 7.1](#) gives $L_{\kappa,k} = F_k$.

9.3 The same countable obstruction in a complete field

Example 9.3. Keep $\kappa = \aleph_\omega$ but enlarge the exponent group to Γ_{κ^+} . The same series x_κ in (9.2) now has support bounded above by e_κ . Since

$$\text{cf}(\Gamma_{\kappa^+}) = \kappa^+ \neq \omega = \text{cf}(\kappa),$$

[Theorem 7.4](#) says that $K_{\kappa,k}$ is complete. Nevertheless, the countable nest

$$B(a_n, e_{\aleph_n}) \quad (n < \omega)$$

has empty intersection in $K_{\kappa,k}$. Its radii never reach the scale e_κ , so it is not a Cauchy obstruction. In particular, the sequence (a_n) itself is not Cauchy in this larger valuation topology.

For this example, all retractions $K + Kx_\kappa \rightarrow K$ are continuous but none is contractive. Moreover,

$$x_\kappa \quad \text{and} \quad x_\kappa + t^{e_\kappa}$$

are distinct elements with exactly the same ordered real type or complex valued type over $K_{\kappa,k}$. The coefficient at e_κ is part of the invisible tail. Its addition changes the full support order type from κ to $\kappa + 1$ but leaves the first- κ prefix unchanged.

9.4 A completion strictly between the two support fields

Example 9.4. Let $\kappa = \aleph_\omega$ and take the ordered rational vector space

$$\Gamma^* = \Gamma_\kappa \oplus \mathbb{Q}u,$$

where the u coordinate dominates every coordinate of Γ_κ . Its cofinality is ω , since $(nu)_{n \geq 1}$ is cofinal. In $F_k = k((t^{\Gamma^*}))$, the series $x_\kappa = \sum_{\alpha < \kappa} t^{e_\alpha}$ has its entire support below u , so it does not belong to the completion of $K_{\kappa,k}$. On the other hand,

$$y = \sum_{n < \omega} \sum_{\alpha < \aleph_n} t^{nu + e_\alpha}$$

has a cofinal well-ordered support of order type $\sum_{n < \omega} \aleph_n = \aleph_\omega$. The blocks are strictly separated because every e_α is less than u . Therefore $y \in L_{\kappa,k} \setminus K_{\kappa,k}$ by [Proposition 7.2](#), whereas $x_\kappa \in F_k \setminus L_{\kappa,k}$. This proves the strict chain

$$K_{\kappa,k} \subsetneq L_{\kappa,k} \subsetneq F_k.$$

9.5 A regular bound with a genuinely uncountable obstruction

Example 9.5. Take $\kappa = \aleph_1$ and $\Gamma = \Gamma_{\omega_1}$. Every countable nest of closed balls in $K_{\aleph_1,k}$ meets, but a nest of cardinality $\aleph_1$ can be empty. The field is not complete, because $\text{cf}(\Gamma) = \aleph_1 = \text{cf}(\kappa)$, and

$$x = \sum_{\alpha < \omega_1} t^{e_\alpha}$$

is a missing limit of its ω_1 -indexed truncation net. Its ordered or valued type has omission number $\aleph_1$.

Here every Cauchy sequence is eventually constant. To prove this directly, consider the countable set of finite values $v(z_m - z_n)$ of all nonzero pairwise differences in a given sequence. Since $\text{cf}(\Gamma) > \omega$, this set is bounded above by some $\beta \in \Gamma$. The Cauchy condition at scale β forces every sufficiently late pairwise difference to be zero. Thus sequential completeness does not imply completeness for the valuation uniformity in this example. The missing net cannot be replaced by a countable Cauchy sequence.

9.6 The rank-one boundary case

If Γ is a subgroup of $(\mathbb{R}, +)$, every well-ordered subset of Γ is countable. One proof is to assign to each nonfinal element of a well-ordered subset a rational number strictly between it and its successor; the chosen intervals are disjoint. There are only countably many rationals. Consequently, for every uncountable κ ,

$$K_{\kappa,k} = k((t^\Gamma)).$$

The nontrivial cardinal-support phenomena in this paper therefore require value groups beyond the Archimedean case. Merely replacing the real coefficients by complex ones in a rank-one Hahn field does not create these missing types. This is consistent with the rank-one Berkovich report, where the proper closure of the finite-support series in $\mathbb{C}((t^\mathbb{Q}))$ occurs only at the excluded countable bound ([Remark 7.6](#)).

Remark 9.6 (The same groups elsewhere in the collection; added during assembly). The construction (9.1), together with the embedding $e_\alpha \mapsto \omega^\alpha$ of [Section 10](#), is the one used in the report on entire functions at arbitrary rank. Its group $\Gamma_\infty = \bigoplus_{j \geq 0} \mathbb{Q}e_j$ with $e_j = \omega^j$ (`ent : eq : concrete-group`) is Γ_ω here, and its group Γ_{ω_1} (`ent : eq : Gamma-omegaone`) is the group of [Example 9.5](#). Both reports compute the same cofinalities, $\aleph_0$ and $\aleph_1$. The questions asked about these groups are different: there, entire functions and their rings; here, cardinally bounded subfields.

10 Transfer to actual surreal and surcomplex numbers

10.1 The normal-form realization

Let $j : \Gamma \rightarrow (\mathbf{No}, +)$ be a specified increasing additive embedding of the set-sized exponent group. Conway normal-form arithmetic gives the field embedding

$$\iota_j : \mathbb{R}((t^\Gamma)) \longrightarrow \mathbf{No}, \quad \sum_{\gamma \in S} r_\gamma t^\gamma \mapsto \sum_{\gamma \in S} r_\gamma \omega^{-j(\gamma)}. \quad (10.1)$$

The exponent set on the right is reverse well ordered, since S is well ordered and j is increasing. The minus sign is essential: the increasing-support convention for t^Γ corresponds to decreasing exponents of ω . Existence, uniqueness, and arithmetic of these Conway normal forms are the standard surreal normal-form theorem [8].

The map is injective, order preserving, and preserves the support cardinality and its indexing order type after reversing the exponent convention. For complex coefficients, use the coordinatewise embedding

$$\mathbb{C}((t^\Gamma)) \cong \mathbb{R}((t^\Gamma))[i] \longrightarrow \mathbf{No}[i]. \quad (10.2)$$

Its two real coordinates are the real and imaginary coefficient series. Their union is exactly the support of the original complex series, so the cardinal restriction transfers without change.

Consequently every theorem in this paper applies to the actual subfields $\iota_j(K_{\kappa, \mathbb{R}}) \subseteq \mathbf{No}$ and their quadratic complexifications, when furnished with the transported valued or ordered structures. No proper-class quantification is required for the one-types: their base fields and ambient full Hahn workspaces remain sets.

10.2 The explicit groups already live in the surreal field

For the groups Γ_θ in (9.1), define

$$j\left(\sum_{\alpha \in A} q_\alpha e_\alpha\right) = \sum_{\alpha \in A} q_\alpha \omega^\alpha, \quad A \subseteq \theta \text{ finite.}$$

Finite Conway normal forms make this an additive embedding, and the largest exponent controls the sign, so it has exactly the required order. Thus e_α can be represented concretely by the surreal monomial ω^α .

Under this embedding, the missing series in (9.2) is the actual surreal

$$X_\kappa = \sum_{\alpha < \kappa} \omega^{-\omega^\alpha}. \quad (10.3)$$

For $\kappa = \aleph_\omega$, its approximants are

$$A_n = \sum_{\alpha < \aleph_n} \omega^{-\omega^\alpha}.$$

In the workspace with exponent group $j(\Gamma_\kappa)$, these are a Cauchy sequence in the intrinsic valuation topology and have no limit in the support-restricted subfield. In the workspace with exponent group $j(\Gamma_{\kappa^+})$, the same displayed approximants are not Cauchy, and the support-restricted field is complete. The smaller scale $\omega^{-\omega^\kappa}$ now exists in the workspace and detects the failure of the Cauchy condition.

This is a precise example of why a surreal limit assertion must specify which scales are permitted in its neighborhoods. The formulas for the numbers alone do not determine a topology on an arbitrarily chosen subfield.

10.3 The topology and language boundaries

The topological statements here concern the valuation taking its finite values in the chosen group Γ . They are not statements about the topology induced from the order topology of the full class $\mathbf{No}$. In the latter setting, a set of nonzero differences has a positive surreal lower bound for its absolute values, so every set-sized subset is discrete. The repository's foundations treatment explicitly distinguishes this full fine topology from an intrinsic Hahn-workspace topology [9] (found:prop:twotopologies in [15]). In particular, convergence of the displayed sequence in one workspace does not contradict the eventual-constancy results for small-index nets in the full fine topology.

Similarly, the support-restricted subfields are not asserted to be birthday-initial segments. No closure under the full surreal exponential, no compatibility with the Berarducci–Mantova derivation, and no definability of the prefix operator in the field language is claimed. The prefix is an external invariant describing the types; the individual tests used in the proofs are ordinary finitary formulas with allowed parameters.

Naming complex conjugation would create another language. Its fixed field is the real axis, and positivity on that fixed field is definable by being a nonzero square. Thus the pure-field collapse from Theorem 5.4 cannot be transferred without qualification to a surcomplex field with named conjugation. We have not classified all one-types in that expanded language: an arbitrary complex element corresponds to a pair of real coordinates, whose joint algebraic relations require a tuple analysis.

11 Scope, provenance, and a formalization route

11.1 Repository boundary

The repository was examined at the pinned commit

465a54b479a1ee842cbf7db1689a7d2f6bfe25e1.

The root overview, documentation catalogue, the docs/new directory, and the research audit of the closest bounded-support transcendence report were read through the GitHub connector. The catalogue at that revision describes 44 reports and explicitly distinguishes mathematical arguments, finite checks, and Lean coverage [9]. No complete audit of every article, historical version, or Lean declaration is claimed.

Three comparisons matter. The report on *transcendence over bounded support* uses supports bounded in exponent order and the fraction field they generate; our base is instead the already-real-closed or algebraically closed field of all supports of cardinality less than κ . The foundations report treats universes, localization, and the full fine topology; our theorems concern exact one-types and intrinsic completion in a fixed set-sized localization. The Hahn-vector-space report studies strong maps and several duals; [Corollary 8.2](#) instead uses the inherited valuation on a two-dimensional space over a potentially incomplete cardinally bounded base field. These differences are substantive, not merely changes of notation.

A targeted repository search returned no indexed matches for one relevant term, but the index was not treated as an exhaustive content audit. The comparisons above rely on the retrieved catalogue and the closest report’s stated base field and scope, not on that negative search.

11.2 Literature and novelty ledger

The mathematical dependency boundary can be summarized without claiming publication priority for standard ingredients.

Ingredient or conclusion	Status in this manuscript
Hahn fields; normal-form realization	Classical input; not new [4, 8].
Regular κ -bounded constructions	Prior literature; not new [1, 2].
Quantifier elimination	Classical logical input; not new [5, 6, 7].
Approximation types and ball conditions	General framework already established [3].
First- κ type fibers; exact omission number	Explicit calculation proved here, including singular κ ; proposed contribution.
Completion and first empty nests	Sharp cardinal-support classification proved here; gluing mechanism is classical.
Retraction loss and dual dichotomy	Direct consequences of the exact approximation cut; no general new Hahn–Banach theorem claimed.

Kuhlmann’s general approximation-type account is the closest logical precedent, especially its discussion of one-types and its ball formulas [3, Section 4.2]. Merely using such ball conditions would not establish novelty. Here the cardinal computation supplies the explicit prefix parameterization, proves the lower bound for *arbitrary* formulas, and connects it to a completion criterion for both regular and singular bounds. These are the statements proposed for further expert review.

The search was targeted rather than exhaustive. It included cardinally bounded Hahn fields, approximation types, singular-cardinal support bounds, completion, and spherical completeness. Some broad queries returned irrelevant results; those were not counted as evidence of absence. No exhaustive MathSciNet or zbMATH priority check, citation-network review, or survey of unpublished manuscripts was performed. Accordingly, the warranted claim is that the combined statements were not identified in the reviewed material. A later identification of an earlier equivalent theorem would change the novelty assessment, not replace the proofs provided here.

11.3 A feasible proof-assistant decomposition

No Lean file or Lean build accompanies this manuscript. A formalization would be most naturally divided into the following interfaces rather than beginning with proper-class model theory.

Coefficient layer. Work with a set-sized ordered exponent type and a Hahn-series type. Formalize strict and non-strict truncations, first disagreement, the cardinality of finite unions of supports, and the small-workspace lemma. The argument at singular κ needs the countable cardinal bound on the rational span, not regularity.

Cardinal and support layer. Represent the first- κ prefix by an order isomorphism from a well-ordered support to its ordinal order type. Prove the exact approximation set, prefix fibers, coefficient gluing, and the long-cofinal-support lemma. These conclusions do not depend on quantifier elimination. A useful interface theorem is that fewer than $\text{cf}(\kappa)$ subsets of size less than κ have union of size less than κ .

Algebra and topology layer. Import full-Hahn real or algebraic closedness with its hypotheses explicitly checked. Then formalize completion as closure inside the full Hahn field, the classification of newly added supports, and the retraction-loss trichotomy. This already certifies the main valuation and linear conclusions without a theory-of-types library.

Logical layer. State the quantifier-elimination interfaces for RCF and ACVF and prove eventual agreement first for atomic polynomial data, then for Boolean combinations, and finally for formulas. This isolates the only model-theoretic import in the omission-number lower bound. An implementation that assumes these interfaces proves a conditional theorem; it must not be described as an unconditional Lean verification until those imports are discharged.

11.4 Further questions, not conclusions

The one-variable invariant is complete precisely because over an algebraically closed base every polynomial splits into linear factors, while real polynomials add only positive quadratic factors. In several variables there is no analogous factorization into conditions on single coordinates. A natural next question is whether finite tuples in a full Hahn extension admit an effective prefix description that also records cancellations in polynomial combinations. The one-type theorem alone does not answer it.

A second question concerns richer languages: named truncation, named conjugation, a surreal exponential, or a derivation. Each can make additional relations visible, and closure of the base field under the new operations must be established before even formulating an elementary-substructure comparison. The present language-sensitive examples supply test cases, not a claim that the same classification survives those expansions.

11.5 Corrections to the manuscript’s repository statements

The statements above describe the repository at the manuscript’s pin 465a54b, which is kept as provenance; the pinned URLs in [9] are also kept. During assembly they were checked against the current collection, with the following results.

- (1) *Catalogue size.* At the pin the catalogue described 44 reports, as stated. Since then batch 21 added `surreal-fields-across-universes` and `birthday-cutoffs-and-hereditary-sets`, and batch 22 added this report and `single-dilation-hahn-support`; the collection had 48 report directories when this report was written.
- (2) *Universes.* At the pin, universes were treated in the foundations report. The collection now also has a report on surreal fields across set-theoretic universes, which studies saturation and omitted cuts of proper-class surreal fields of inner models. Its objects differ from the set-sized fields here; see [Section 1.4](#).
- (3) *The negative search.* The indexed search that returned no matches was for the word “spherical” (`research_audit.md`). The negative result was wrong: at the pin the collection already stated spherical completeness of full Hahn fields, in the rank-one Berkovich report (its `prop:spherical`) and in the three-duals report (`duals:prop:spherical`). The manuscript did not rely on that search, as it says, and both statements are the classical full-Hahn result that it credits to Poonen.
- (4) *The Hahn-vector-space comparison was incomplete.* Besides strong maps and duals, the three-duals report already contained, at the pin, the completion formula `duals:thm:completion` and the cofinality criterion `duals:thm:cofinality-completion`, and the rank-one Berkovich report the closure of the finite-support

series in $\mathbb{C}((t^{\mathbb{Q}}))$. These are countable analogues of [Theorems 7.1](#) and [7.4](#) ([Remark 7.6](#)). They are not cardinally bounded fields and do not anticipate the uncountable, and in particular the singular, case treated here, but they belong in the priority context of the completion results.

- (5) *Statements found accurate.* The description of the bounded-support report (order-bounded supports and their fraction field, explicitly not a cardinal restriction), the foundations report’s separation of the fine and intrinsic topologies (`found:prop:twotopologies`), and the absence from the collection of any study of cardinally bounded Hahn fields, their types or their completions were confirmed.

11.6 What is not claimed

The manuscript states its limitations where they arise. They are collected here, with the one limitation added during assembly; none is withdrawn.

- N1. κ is uncountable; the finite-support case $\kappa = \aleph_0$ is excluded ([Appendix A](#)).
- N2. Established mathematics is credited, not claimed: cardinally bounded Hahn fields (Kuhlmann–Shelah for regular κ ; Berarducci–Kuhlmann–Mantova–Matusinski), Hahn arithmetic and closedness, Conway normal forms, general approximation-type theory including its ball formulas (F.-V. Kuhlmann, Proposition 4.4 and Section 4.2), full-Hahn ball gluing (Poonen), and quantifier elimination for real closed and algebraically closed valued fields (Tarski, Robinson, Yin).
- N3. The contribution is proposed only: the joint first- κ classification, the exact omission number and the completion and linear consequences. The combined statements were not identified in the reviewed material; the priority search was targeted, not exhaustive (no MathSciNet or zbMATH search), and expert review may find equivalents or a simpler general formulation.
- N4. No named longstanding conjecture is settled. The repository comparison was not an exhaustive audit, and a negative index search was not treated as evidence (it was in fact a false negative; [Section 11.5](#)). There is no Lean build or certification.
- N5. Only types realized in the fixed full Hahn extension F_k are classified. No saturation theorem and no automorphism homogeneity is proved, and types are not asserted to be automorphism orbits of F_k .
- N6. All topology is intrinsic to the set-sized workspace, not the subspace topology from **No**; support bounds are cardinalities, not order bounds.
- N7. The subfields are not asserted to be birthday-initial. No closure under the surreal exponential, no compatibility with the Berarducci–Mantova derivation, and no definability of the prefix operator is claimed.
- N8. Languages with named conjugation, truncation, exponential or derivation are not classified, nor are types of several-variable tuples; these remain questions.
- N9. No new general Hahn–Banach theorem is claimed. The zero continuous dual is over an incomplete base in the inherited topology and does not contradict finite-dimensional functional analysis.
- N10. This is not a new version of the independence theorem of the report on transcendence over bounded support.
- N11. The 9,622 finite assertions of `code/verify.py` test finite local identities only. They do not verify the transfinite theorems, quantifier elimination, novelty or any Lean implementation.
- N12. Added during assembly: the comparisons with other reports in [Section 1.4](#) and [Remarks 7.6](#) and [9.6](#) are editorial. They prove no new theorem, and no statement here is derived from those reports.

12 Conclusion

The obstruction created by a cardinal support bound has a precise shape. A missing Hahn series is first-order visible through its first κ nonzero terms, but the least number of formulas needed to force that missing prefix is only $\text{cf}(\kappa)$. Its approximation values are the downward closure of those first support positions. Whether this cut reaches all of Γ decides

both membership in the completion and the existence of continuous retractions from the associated two-dimensional valued space.

This gives a unified explanation of three superficially different effects: singular-cardinal compression of omitted types, complete fields with small empty ball nests, and failure of continuous coordinate functionals over an incomplete base. All three occur in explicitly specified surreal and surcomplex subfields. Their formulation requires neither a proper-class first-order structure nor an unspecified notion of surreal convergence.

A Proof-risk checklist and finite verification

Cardinal bounds. The entire discussion assumes $\kappa > \aleph_0$. At $\kappa = \aleph_0$, finite-support series are generally not a field: $(1 - t^g)^{-1} = \sum_{n \geq 0} t^{ng}$ for $g > 0$ has infinite support. Regularity is not assumed for κ ; it is used only for $\mu = \text{cf}(\kappa)$, which is regular. The small-workspace and small-union arguments are separate and use different bounds.

Prefix versus full support. A missing series can have support order type larger than κ . Its type still remembers only the first κ positions. By contrast, a newly added completion element must have support order type *exactly* κ , not merely cardinality κ . The example $x_\kappa + t^{e_\kappa}$ shows why these statements differ.

Valuation inequalities. A closed ball of radius value γ fixes coefficients *below* γ . A strict valuation inequality fixes coefficients *at and below* γ . An ordered inequality $|z| < t^\gamma$ forces $v(z) \geq \gamma$, not necessarily $v(z) > \gamma$. The real-type proofs use later cofinal thresholds to recover the coefficient at a given position; they do not make the stronger, false implication.

Cauchy versus pseudo-Cauchy. The identity $v(a_j - a_i) = s_{\alpha_i}$ proves a pseudo-Cauchy condition. It gives a Cauchy net only when those values are cofinal in the whole group. This is why the countable family in the complete-field example is not a missing Cauchy limit. The topology induced from full **No** is a third, separate matter.

Scope of quantifier elimination. The type theorems use exactly the stated ordered or valued language. In the pure complex ring language the answer changes, as proved. A monomial appearing as a parameter in a formula is not a named global cross-section symbol. No quantifier-elimination result for a language with a prefix operation is assumed.

Computational checks. The companion code/verify.py uses exact rational coefficients in finite Laurent-polynomial supports over $\mathbb{Z}^2$ with lexicographic order. It checks first-disagreement identities, invariance of polynomial leading data under sufficiently high tails, nested-ball coefficient agreement, and the local trichotomy underlying the retraction-loss formula. A fixed seed makes its checks reproducible, and it writes their assertion counts to a file verification.json in its own directory; the recorded run is data/verification.json (9,622 assertions, all passed). These checks test finite algebraic mechanisms only. They do not prove any infinite-support, cardinality, completeness, first-order, or novelty statement. In particular, a finite support cutoff is *not* treated as a field model of an uncountable cardinal cutoff.

The manuscript was delivered with its TeX source, compiled PDF, build script, build record and a separate research-audit file. This report's directory contains the source and PDF of this report, the delivered script code/build.sh, the delivered build record data/build_audit.json (which describes the delivered 21-page PDF) and research_audit.md. The script was written for the delivery's flat layout; the README of this directory explains how to rerun it on a copy. The written proofs, finite checks, and document-layout checks are distinct forms of evidence.

References

- [1] S. Kuhlmann and S. Shelah, *κ -bounded Exponential-Logarithmic Power Series Fields*, arXiv:math/0512220v1 (2005). <https://arxiv.org/abs/math/0512220>.

- [2] A. Berarducci, S. Kuhlmann, V. Mantova, and M. Matusinski, *Exponential fields and Conway’s omega-map*, Proceedings of the American Mathematical Society **151** (2023), 2655–2669. DOI: [10.1090/proc/14577](https://doi.org/10.1090/proc/14577). Preprint: <https://arxiv.org/abs/1810.03029>.
- [3] F.-V. Kuhlmann, *Approximation types describing extensions of valuations to rational function fields*, arXiv:2111.10529v1 (2021), especially Sections 2.4–2.6 and 4.2. <https://arxiv.org/abs/2111.10529>.
- [4] B. Poonen, *Maximally complete fields*, L’Enseignement Mathématique **39** (1993), 87–106, especially Lemma 4 and Corollary 4. Author’s paper hosted at Stanford: <https://math.stanford.edu/~conrad/Perfseminar/refs/poonencomplete.pdf>.
- [5] A. Tarski, *A Decision Method for Elementary Algebra and Geometry*, prepared for publication with the assistance of J. C. C. McKinsey, RAND Corporation, R-109 (1951). <https://www.rand.org/pubs/reports/R109.html>.
- [6] A. Robinson, *Complete Theories*, North-Holland, Amsterdam (1956). The one-sorted valued-field quantifier-elimination attribution is also recorded in the introduction of [7].
- [7] Y. Yin, *Quantifier elimination and minimality conditions in algebraically closed valued fields*, arXiv:1006.1393v1 (2010). <https://arxiv.org/abs/1006.1393>. The introduction distinguishes the classical one-sorted divisibility language from the richer languages studied in that paper.
- [8] H. Gonshor, *An Introduction to the Theory of Surreal Numbers*, London Mathematical Society Lecture Note Series **110**, Cambridge University Press (1986). Classical reference for Conway normal forms and their arithmetic; the complete book was not freshly audited for this manuscript.
- [9] V. Reshetnikov and contributors, *Surreal*, GitHub repository, revision 465a54b479a1ee842cbf7db1689a7d2f6bfe25e1. Documentation catalogue: <https://github.com/VladimirReshetnikov/Surreal/blob/465a54b479a1ee842cbf7db1689a7d2f6bfe25e1/docs/README.md>. Closest report’s audit: https://github.com/VladimirReshetnikov/Surreal/blob/465a54b479a1ee842cbf7db1689a7d2f6bfe25e1/docs/surreal/transcendence-over-bounded-support/research_audit.md. See Section 11 for the actual inspection boundary.
- [10] Surreal collection, *Surreal Fields Across Set-Theoretic Universes*, report docs/foundations-and-computation/surreal-fields-across-universes (label prefix univ:). Added after the manuscript’s pin.
- [11] Surreal collection, *Bounded-Support Hahn Arithmetic*, report docs/surreal/transcendence-over-bounded-support (label prefix bst:).
- [12] Surreal collection, *Three Duals at Surreal Scales*, report docs/surcomplex/three-duals-of-hahn-vector-spaces (label prefix duals:).
- [13] Surreal collection, *Rank-One Non-Archimedean Analytic Geometry inside the Surcomplex Numbers*, report docs/surcomplex/rank-one-berkovich (bare labels).
- [14] Surreal collection, *Cofinality, Factorization and Scalar Extension for Entire Hahn Functions at Arbitrary Rank*, report docs/surcomplex/entire-functions-at-arbitrary-rank (label prefix ent:).
- [15] Surreal collection, *Foundations for Surreal and Surcomplex Mathematics*, report docs/foundations-and-computation/foundations (label prefix found:).
- [16] Surreal collection, *Hidden Negative Directions in Surcomplex Linear Algebra*, report docs/surcomplex/hidden-negative-hermitian-directions (label prefix hnd:).