

Finite Separators and Discriminant Factorizations for Sextic Block Resolvents

A sharp 196-point guarantee and a bounded replacement
for ProveIt’s separating-invariant search

Prepared for Vladimir Reshetnikov

September 28, 2026

ProveIt research-report collection, continuing the formal project Algebra/PolynomialFormulas. Batches 40–41 intake, batch 41, manuscript 03 (archive ProveIt_Finite_Sextic_Separators); pinned to ProveIt e2b1f016a. AI-assisted, unrefereed, not formalized. Notes marked **[write]** were added when the manuscript was written into the collection; see [sections 1.4](#) and [1.5](#).

Abstract

The sextic radical-solvability development in ProveIt uses terminating but unbounded search to specialize two block-partition descriptors. We replace that search, at the level of an ordinary mathematical proof, by a single explicit parameter range. For every separable sextic in characteristic zero, one of the integers $1, \dots, 196$ separates all fifteen pair partitions and all ten triple partitions simultaneously. The underlying descriptors compress to quadratics and linears. Their collisions are controlled by three invariant polynomials Q_f, U_f, L_f , with degrees at most 30, 120, 45, and we prove

$$\text{disc}_Z R_{2,f} = \text{disc}(f)^6 Q_f^6 U_f^2, \quad \text{disc}_Z R_{3,f} = \text{disc}(f)^3 L_f^2.$$

For the generic sextic, the three factors are irreducible and have the stated exact degrees. Consequently 196 is optimal for a guarantee valid for every set of distinct nonzero complex test parameters, although optimality of the initial integer interval is not established. A split integer example gives a lower bound of eight for that latter problem. We derive a coefficient-only primitive-recursive decision architecture, a height-only separator requiring no search, a precise repository integration plan, and further research questions. Exact symbolic checks and a modular Bézout certificate accompany the proofs.

Status and scope. This is an unrefereed research article with complete written proofs and reproducible exact checks. The new results have not been formalized or run in Lean. General decidability, including polynomial-time decidability of solvability by radicals, is classical; it is not claimed as a new discovery. The specific separator and collision-factorization results are proposed contributions relative to the inspected repository. Historical priority is not certified by the literature check.

Contents

1	The repository question and the contribution	2
2	Notation and complete partition descriptors	6
3	Compression to quadratics and linears	7
4	The two types of pair-partition collision	8
5	Triple collisions as exchanges of two roots	9
6	Three invariant obstruction polynomials	10
7	Exact discriminant factorizations	11
8	Generic irreducibility of the three components	12
9	The finite separator theorem and its sharpness	14
10	Why the separated resolvents decide an irreducible sextic	16
11	A bounded coefficient-only decision architecture	17
12	A separator determined by coefficient height alone	19
13	Exact checks, certificates, and their limits	19
14	A precise route back into ProveIt	21
15	A reusable general finite-separation principle	22
16	Further research questions and priorities	22
17	Conclusion	24
A	Reproducibility and package contents	24
B	Source and claim ledger	25
	References	26

1 The repository question and the contribution

The starting point is a concrete implementation boundary, not an assertion that sextic solvability is an unsolved problem. In the pinned ProveIt snapshot, `SexticRadicalDecision.lean` proves a total recursive criterion for integer sextics. Its introductory comment explicitly does not claim primitive recursiveness, because the separating-invariant search uses unbounded minimization. The supporting file `SexticIrreducibleDecision.lean` constructs pair and triple parameters using `Nat.find`. The file `SexticSeparatingInvariants.lean` proves that suitable parameters exist by representing each block by its root polynomial and each partition by the polynomial of those block polynomials [4, 5, 6, 7].

That is a particularly useful research target: the existing semantic architecture already identifies the right invariant and its correctness condition. What is missing from this route is a small explicit bound and a sufficiently transparent explanation of the exceptional specializations. Our answer is stronger than merely bounding an enumeration of pairs of natural numbers. A one-dimensional curve works for every distinct root tuple, its failure set has three explicit components, and its generic number of failures can be determined exactly.

1.1 What is classical, inherited, and proved here

The equivalence between radical solvability and solvability of the Galois group is classical. So is the sextic criterion using the two imprimitive wreath-product groups of degrees fifteen and ten in the corresponding coset actions; see, for example, Boswell and Glasser [2]. Landau and Miller proved polynomial-time decidability of radical solvability in 1985 [1]. Thus even a primitive-recursive decision for sextics is not, as an abstract existence theorem, new.

The partition descriptors and the strategy of separating their specialized values are inherited from the inspected ProveIt source. More broadly, the construction and use of relative invariants are central to computational Galois theory; Fieker and Klüners discuss general methods and the practical importance of structured invariants [3].

The claims established in this article are the explicit compression, the special curve $(u, v) = (t, 2t^2)$, the 196-point uniform bound, the three-factor description of its obstruction, the exact discriminant identities, and generic irreducibility and sharpness. Their proofs are given below rather than inferred from computations. The bounded decision architecture is an application to the repository’s mathematical gap, not a replacement claim for the stronger classical complexity theorem.

The literature check consulted the primary sources just cited and the pinned source files. It does not establish that no equivalent identity has appeared elsewhere under different coordinates or resolvent conventions. Accordingly, “contribution” here means a result developed and proved for this report, with a demonstrated use relative to this repository; it does not mean a certified first-in-literature result.

1.2 Main results at a glance

The root tuple is always assumed distinct. The polynomials h_P and k_T are defined explicitly in [section 3](#); their resolvents have degrees fifteen and ten in Z .

Uniform separator. Among any 196 distinct nonzero elements of a characteristic-zero field containing the roots, there is a parameter t for which both maps $P \mapsto h_P(t)$ and $T \mapsto k_T(t)$ are injective. In particular, testing $1, \dots, 196$ always succeeds.

Exact obstruction. There is a nonzero polynomial $E_f = Q_f U_f L_f$ of degree at most 195, with coefficients polynomially expressible in the sextic coefficients, such that $E_f(t) \neq 0$ is exactly simultaneous separation. Its three factors have generic degrees 30, 120, 45 and are generically irreducible.

Sharpness and implementation. The number 196 is optimal for arbitrary complex test sets. For the initial integer interval the proven bounds are $8 \leq N_{\text{int}} \leq 196$. The explicit finite range removes unbounded minimization from a coefficient-only sextic decision architecture.

[write] Scope of the last sentence. The finite range removes unbounded minimization from the architecture of [section 11](#), on paper. It removes nothing from the Lean development yet: none of this report is formalized, and the Lean search it bounds is still an unbounded `Nat.find`; see [section 1.5](#).

1.3 [write] Conventions and notation

No symbol of the delivered text was renamed and no normalization was changed. The table lists the letters that carry more than one meaning in the text, in the Lean development it continues, or in the sibling report [specialization-safe-radical-solvers \[13\]](#); where a false reading is tempting it is printed beside the true one.

Symbol	Meaning here, and readings to avoid
t	The one parameter on the curve $(u, v) = (t, 2t^2)$ in descriptor space. Lean's parameter is a pair $x: \text{Fin } 2 \rightarrow \mathbb{N}$, with X0 the outer (block) variable v and X1 the root variable u . <i>True:</i> $x(0) = 2t^2, x(1) = t$. <i>False:</i> $x(0) = t, x(1) = 2t^2$.
n (Lean)	The <code>Nat.find</code> index of the Lean search is a <code>Primcodable</code> code, decoded to a pair by <code>parameterAt</code> ; it is not t . The bound 196 bounds t , not the code.
$p_B(u), p_b, p$	p_B is the root polynomial of a block B of any size, equation (2.3) ; $p_b = \alpha_i \alpha_j$ is the product of a root pair b , equation (3.1) ; $p = 1,000,003$ is the prime of equation (9.3) .
$e_j, e_j(B), e$	e_j are the elementary symmetric functions of all six roots, $e_j(B)$ those of a block; e, e' are root pairs. In lemma 5.1 the letters $a, \dots, f$ are root values, so there f is not the sextic and e is not a pair.
$A_P, B_P, C_P; a_T, b_T; A, B$	Pair and triple coefficients, equations (3.1) and (3.4) . The bare A, B are the coefficients of $h_P - h_Q$ as a linear polynomial in one root in the proof of lemma 8.2 , and the Bézout cofactors in equation (9.3) .
$R_{2,f}, R_{3,f}, R$	The compressed resolvents equation (7.1) , of degrees 15 and 10 in Z , are built from the values $h_P(t), k_T(t)$; Lean's evaluated resolvents are built from the descriptor values $D_P(t, 2t^2) = K_2(t) - 2t^2 h_P(t)$, an affine image, so the two are separable at the same $t \neq 0$. $R = 1 + H$ is a root bound (section 11). Neither is the sibling report's subset-sum resolvent $R_{p,k,f}$.
$h_P(t)$	The pair invariant equation (3.1) , quadratic in t ; not the sibling report's subset compression $h_A(u) = \sum_r u^{r-1} e_r(x_A)$.
K, K_2, K_3	K is the field; K_2, K_3 are the partition-independent parts in equations (3.3) and (3.6) .
H, H_R	H is the largest absolute value of the lower coefficients of the monic sextic (sections 11 and 12); in the proof of lemma 8.2 H is a putative common factor; H_R is the analogue of H for a resolvent.
Q, U, L, E and Q_f, U_f, L_f, E_f	Universal polynomials over $\mathbb{Q}[e_1, \dots, e_6, t]$ (section 8) and their specializations to one sextic; $\bar{E}_f$ is E_f reduced modulo p .
Degree	Unqualified degrees 30, 120, 45, 195 are degrees in t . The homogeneous degrees 30, 240, 90, 360 of the remark after proposition 6.2 count roots and t together.

Symbol	Meaning here, and readings to avoid
196 versus N_{int}	196 is sharp for <i>arbitrary</i> sets of distinct nonzero complex test values (theorem 9.2). For the initial integer interval only $8 \leq N_{\text{int}} \leq 196$ is proved (proposition 9.3). <i>False reading</i> : “some sextic defeats all of $1, \dots, 195$ ”.

1.4 [write] Provenance of this report

This report is built from **one** manuscript: batch 41, manuscript 03 of ProveIt’s incoming-reports intake (written together with batch 40), delivered as the archive `ProveIt_Finite_Sextic_Separators` (11 files, including a 24-page Letter PDF) in commit `9754e8360`, pinned to ProveIt commit `e2b1f016a94102663f12b970e6dd434229f90d01`, and placed in commit `eaf787d50`. It is filed in ProveIt’s research-report collection beside `specialization-safe-radical-solvers` [[13](#)], with which it shares no theorem, and continues the formal project `Algebra/PolynomialFormulas`. That project is unchanged between the pin and the placement commit, so every repository statement of the delivered text was checked against the tree it describes. Every statement, proof, table, example, remark, question and limitation of the delivered text is printed. No merge choice arose.

The write changed only the following. Every label carries the prefix `sbr:`. The title-page provenance line, [sections 1.3 to 1.6](#), the unnumbered [write] notes, and the bibliography entries [[11](#)], [[12](#)] and [[13](#)] were added, and the title page no longer duplicates the PDF anchor of page 1. Two corrections of form: the title-page spacing was tightened so that the added line fits, and the environments sharing the theorem counter now use alias counters, because in the delivered build every cross-reference to a lemma, proposition or corollary printed the word “theorem”. The notes use no theorem counter, so every theorem, equation and section number is that of the delivered PDF. The delivered text names the delivery layout: its files `checks/X` are shipped as `data/X`, its `build.sh` as `code/build.sh` and its `requirements.txt` as `data/requirements.txt`; the delivered PDF, README and checksum file `SHA256SUMS` are not shipped (the checksums were verified, 10 of 10, at placement). [Section A](#) carries a note with commands that work with the shipped names.

Relation to the sibling report. The manuscript does not know `specialization-safe-radical-solvers` [[13](#)]. That report’s separating repair compresses the k -subsets of the roots of a prime-degree polynomial by $h_A(u)$ and bounds the bad parameters by adding the degrees of all pairwise differences (1191 values suffice for septic triples, not claimed sharp); its question on smaller deterministic separating families asks for a better number for $(n, k) = (7, 3)$ by counting collision orbits under the Galois action. The present report carries out that kind of orbit count for a different family, the sextic block partitions, and reaches a sharp bound. It answers no question of the sibling report.

1.5 [write] Formal status in ProveIt

None of this report’s statements is formalized. No `sbr:` label has a Lean or Rocq counterpart. In particular the compression ([propositions 3.1](#) and [3.2](#)), the collision factorizations, the exact failure test [equation \(6.6\)](#), the 196-point bound ([theorem 9.1](#)), the discriminant identities, generic irreducibility, sharpness, the N_{int} bracket, the bounded architecture ([theorem 11.1](#)) and the height separator have written proofs only. The modules proposed in [section 14](#) (`SexticCompressedDescriptors.lean`, `SexticCollisionFactors.lean`, `SexticBoundedSeparatingSearch.lean`) do not exist. Placement beside a Lean and Rocq development confers no formal status.

What ProveIt does prove in Lean. All in namespace `LeanProofs.PolynomialFormulas` under `Algebra/PolynomialFormulas/Lean/PolynomialFormulas/`, at the pin and unchanged since; line numbers are those of the declarations.

- `SexticSeparatingInvariants.lean` [7]: `rootPolynomial_injective` (l. 115), `pairDescriptor_injective` and `tripleDescriptor_injective` (l. 422, 428), which together are [lemma 2.1](#); `bivariateDescriptor` (l. 618), a product of X_0 minus block polynomials in X_1 , and `pairBivariateDescriptor_injective`, `tripleBivariateDescriptor_injective` (l. 668, 677); `pairDescriptorValue`, `tripleDescriptorValue` (l. 790, 794), the evaluation at $x: \text{Fin } 2 \rightarrow \mathbb{N}$; `exists_nat_eval_injective` (l. 1048), which separates a finite injective family of multivariate polynomials over a characteristic-zero domain by one natural-number point (through `exists_nat_eval_ne_zero`, l. 1019, the combinatorial Nullstellensatz); and its sextic instances `exists_pairDescriptor_separating_evaluation`, `exists_tripleDescriptor_separating_evaluation` (l. 1074, 1083) and `exists_pairDescriptorValue_injective`, `exists_tripleDescriptorValue_injective` (l. 1092, 1102). These statements are purely existential. The Nullstellensatz proof finds the point in a finite grid, but no bound is stated or exported, and nothing restricts the point to a curve.
- `SexticSeparatingSearch.lean` [11]: the Boolean tests `pairSeparatesB`, `tripleSeparatesB` (l. 536, 539), through the collision products of `pairCollisionPolynomial` (l. 114) and `collisionProduct` (l. 318); the unbounded searches `pairSeparatingCode`, `tripleSeparatingCode` (l. 604, 608; `Nat.find` at l. 606, 610) over codes decoded by `parameterAt` (l. 517); the parameters `pairSeparatingParameter`, `tripleSeparatingParameter` (l. 612, 616); and their correctness `pairSeparatingParameter_injective`, `tripleSeparatingParameter_injective` (l. 620, 629). This is the search this report bounds. *The manuscript does not cite this file*: its citation [6] is `SexticIrreducibleDecision.lean`.
- `SexticIrreducibleDecision.lean` [6]: the guarded total searches `pairTotalCode`, `tripleTotalCode` (l. 87, 90; `Nat.find` at l. 88, 91), total on every monic sextic because a bounded reducibility test returns code 0 on reducible inputs, and `pairTotalCode_computable` (l. 101).
- `SexticRadicalDecision.lean` [5]: the public results `sexticRadicalDecision_correct` (l. 34), `allRootsRadical_computablePred` (l. 49) and `has_verified_sextic_radical_turing_machine` (l. 56). Its docstring (l. 8–9) calls the criterion “not claimed primitive recursive, because the separating-invariant search uses unbounded minimization”.

The Rocq route (`Algebra/PolynomialFormulas/Coq/SexticMuRec*.v`) also selects the separating parameters “by genuine unbounded minimization”, as the project README states; the manuscript mentions it only in passing and does not analyse it.

What formalizing the bound would give (a formalization target, not a claim). A Lean proof of [theorem 9.1](#) for the root tuple of an irreducible monic sextic, together with the affine relations [equations \(3.3\)](#) and [\(3.6\)](#) and the convention $x = (2t^2, t)$ of the first table row, would make the Lean search provably bounded: by minimality of `Nat.find`, `pairSeparatingCode` would be at most the largest `encodeParameter` of the points $(2t^2, t)$, $1 \leq t \leq 151$, and `tripleSeparatingCode` the same with $t \leq 46$; one common $t \leq 196$ serves both. Equivalently the search could be replaced by a fixed scan of $t = 1, \dots, 196$. That bounds the parameter selector only. The Lean pipeline has two further unbounded searches: `SexticReducibleDecision.linearTotalCode` (l. 72, `Nat.find` at l. 73), which the manuscript notes in [section 14](#); and `SexticSparseSymmetricSearch.elementaryCode` (l. 634, `Nat.find` at l. 637) [12], the certificate search behind `elementarySparse` (l. 671), which produces the elementary-symmetric formulas `pairElementarySparse`, `tripleElementarySparse` (l. 1133, 1138) for the resolvent coefficients and `pairCollisionElementary`, `tripleCollisionElementary` (`SexticSeparatingSearch.lean`, l. 282, 286) for the collision products. The manuscript does not mention this search. Its inputs are the parameter and a coefficient index, not the sextic, so over a finite parameter range it searches for finitely many

fixed certificates; a **Primrec** proof must still bound or replace it. This is the condition, in [section 14](#), that the coefficient evaluators themselves have primitive-recursive proofs.

1.6 [write] What is not claimed

The delivered text states each of the following, and each is kept where the text states it. The article is unrefereed. Its new results were not formalized or run in Lean or Rocq, the repository was not rebuilt or modified, and nothing was benchmarked against production Galois software. Decidability of radical solvability, even in polynomial time, is classical (Landau and Miller), as is the sextic block criterion; the contributions are relative to the inspected repository and the consulted sources, and no exhaustive priority search was made. The Python files are an exact verifier and a root-level prototype, not a coefficient-only solver and not an extraction of a formal proof; the expanded universal coefficient tables are specified but not supplied. The value 196 is sharp for arbitrary complex test sets only; the initial integer interval is bracketed, not determined. The fixed-parameter counterexamples are reducible and say nothing about irreducible rational sextics. [Theorem 11.1](#) is a mathematical realization, not a changed Lean function or an accepted **Primrec** proof, and it claims no polynomial bit complexity. The height separator is a termination certificate or comparison construction, not a recommendation for practical evaluation. The Kronecker-curve principle ([proposition 15.1](#)) is a standard monomial-encoding argument, not a novelty claim.

At the write the two scripts were rerun on a copy with SymPy 1.14.0; both recorded outputs were reproduced exactly, apart from line endings. The write also checked, by a separate script, the Lean convention of the first table row against the definitions (300 random integer root tuples, each at $t = 1, \dots, 11$, no disagreement between Lean-style descriptor values and h_P, k_T), the first separator 8 of $(-3, 0, 1, 2, 3, 4)$, and, over $\mathbb{Z}$, that E_f for the roots [equation \(9.1\)](#) has degree 195, a nonzero constant term and no repeated factor. None of this is an independent proof review.

2 Notation and complete partition descriptors

Let K be a field of characteristic zero, let $\alpha_1, \dots, \alpha_6 \in K$ be distinct, and put

$$f(X) = \prod_{i=1}^6 (X - \alpha_i) = X^6 - e_1 X^5 + e_2 X^4 - e_3 X^3 + e_4 X^2 - e_5 X + e_6. \quad (2.1)$$

If the coefficients originally lie in a smaller field, take K to contain a splitting field. Our arguments are algebraic and do not depend on an ordering, numerical approximation, or choice of complex branches.

Write $[6] = \{1, \dots, 6\}$ and $\mathcal{E} = \binom{[6]}{2}$. Let $\mathcal{P}_2$ be the set of partitions of $[6]$ into three unordered two-element blocks, and let $\mathcal{P}_3$ be the set of partitions into two unordered three-element blocks. Their cardinalities are

$$|\mathcal{P}_2| = \frac{6!}{(2!)^3 3!} = 15, \quad |\mathcal{P}_3| = \frac{\binom{6}{3}}{2} = 10. \quad (2.2)$$

We call members of $\mathcal{P}_2$ *matchings*. When two matchings have no common block, we call them *block-disjoint*; they still use the same six roots.

For a block $B \subseteq [6]$, define its root polynomial and the complete partition descriptor by

$$p_B(u) = \prod_{i \in B} (u - \alpha_i), \quad D_{\mathcal{B}}(u, v) = \prod_{B \in \mathcal{B}} (v - p_B(u)). \quad (2.3)$$

These are the ordinary polynomial counterparts of the descriptors in `SexticSeparatingInvariants.lean` [7]. The distinction between the variables u and v matters: u enters the root polynomials, whereas v records the unordered collection of blocks.

Lemma 2.1 (A descriptor determines the partition). *For distinct roots, the map $\mathcal{B} \mapsto D_{\mathcal{B}}(u, v)$ is injective on either partition family.*

Proof. First, the map $B \mapsto p_B(u)$ is injective. Indeed, the roots of p_B among $\alpha_1, \dots, \alpha_6$ are precisely the roots indexed by B . Next regard $D_{\mathcal{B}}$ as a monic polynomial in v over the field $K(u)$. Its distinct linear factors are $v - p_B(u)$. Equality of two such products implies equality of their sets of roots in $K(u)$, so it implies equality of the sets of block polynomials. Their injectivity recovers the blocks and hence the partition. This proof also explains why repeated sextic roots must be excluded at this stage. $\square$

The descriptors therefore solve the information-loss problem before specialization. The remaining task is to evaluate them at a pair (u, v) without causing two distinct descriptors to become equal. Arbitrary fixed evaluations need not preserve injectivity. We next expose the unusually small amount of partition-dependent data in the sextic case.

3 Compression to quadratics and linears

3.1 Pair partitions

For a pair $b = \{i, j\}$, write $s_b = \alpha_i + \alpha_j$ and $p_b = \alpha_i \alpha_j$. For a matching P , set

$$\begin{aligned} A_P &= \sum_{b \in P} p_b, \\ B_P &= \sum_{\substack{b, c \in P \\ b \neq c}} s_b p_c, \\ C_P &= \sum_{\{b, c\} \in \binom{P}{2}} p_b p_c, \end{aligned} \quad h_P(t) = A_P t^2 + B_P t - C_P. \quad (3.1)$$

The sum defining B_P is over ordered pairs of different blocks, so it has six summands before expanding the s_b .

Proposition 3.1 (Pair descriptor expansion). *For every matching P ,*

$$\begin{aligned} D_P(u, v) &= v^3 - (3u^2 - e_1 u + A_P) v^2 \\ &\quad + (3u^4 - 2e_1 u^3 + (e_2 + A_P) u^2 - B_P u + C_P) v - f(u). \end{aligned} \quad (3.2)$$

Consequently, if $K_2(t) = 2t^6 + 2e_2 t^4 - f(t)$, then

$$D_P(t, 2t^2) = K_2(t) - 2t^2 h_P(t). \quad (3.3)$$

The map $P \mapsto h_P(t)$ is injective as a map into $K[t]$.

Proof. Each block polynomial is $u^2 - s_b u + p_b$. Their sum is $3u^2 - e_1 u + A_P$. The sum of their pairwise products has leading coefficients 3, $-2e_1$, and its coefficient of u^2 is $2A_P + \sum_{b < c} s_b s_c$. Since the global e_2 is $A_P + \sum_{b < c} s_b s_c$, that coefficient is $e_2 + A_P$. The remaining two coefficients are $-B_P$ and C_P . The product of all three block polynomials is $f(u)$. Expanding the monic cubic in v gives [equation \(3.2\)](#); substituting $u = t, v = 2t^2$ gives [equation \(3.3\)](#).

If two h polynomials agree, their coefficients give equality of A, B, C . The displayed descriptor formula then gives equal descriptors. Apply [lemma 2.1](#). $\square$

3.2 Triple partitions

For a three-element block B , let $e_j(B)$ denote the elementary symmetric polynomial of degree j in its three roots. For $T \in \mathcal{P}_3$, define

$$a_T = \sum_{B \in T} e_2(B), \quad b_T = \sum_{B \in T} e_3(B), \quad k_T(t) = a_T t - b_T. \quad (3.4)$$

Proposition 3.2 (Triple descriptor expansion). *For every triple partition T ,*

$$D_T(u, v) = v^2 - (2u^3 - e_1 u^2 + a_T u - b_T) v + f(u). \quad (3.5)$$

On $(u, v) = (t, 2t^2)$ this has the form

$$D_T(t, 2t^2) = K_3(t) - 2t^2 k_T(t), \quad K_3(t) = 4t^4 - 4t^5 + 2e_1 t^4 + f(t). \quad (3.6)$$

The map $T \mapsto k_T(t)$ into $K[t]$ is injective.

Proof. The sum of the two monic cubic block polynomials is $2u^3 - e_1 u^2 + a_T u - b_T$, and their product is $f(u)$. Expanding their quadratic descriptor proves the formula. Equality of k polynomials gives equality of the two descriptor-dependent coefficients, and then [lemma 2.1](#) proves injectivity. $\square$

In particular every difference $h_P - h_Q$ for $P \neq Q$ is a nonzero polynomial of degree at most two; every difference $k_T - k_{T'}$ is nonzero of degree at most one. A first root-counting bound would therefore allow $2\binom{15}{2} + \binom{10}{2} = 255$ bad parameters. The next two sections remove repeated collision factors and reduce this to 195.

Remark 3.3 (The zero parameter and the source coordinate order). The compressed families make sense and may separate at $t = 0$, but the original descriptors on the curve all collapse within each family there, because the multiplier $2t^2$ vanishes. We use positive integer tests when transferring results to the repository. In its `Fin2` convention, `X0` is the outer descriptor variable v and `X1` is u . Thus the required parameter is $x(0) = 2t^2$, $x(1) = t$, not the reverse [7].

[write] **Checked against the Lean definitions.** In `SexticSeparatingInvariants.lean`, `bivariateRootPolynomial` (l. 614) is the product of `X1` minus the roots, and `bivariateDescriptor` (l. 618) the product of `X0` minus those block polynomials, so `X0` = v and `X1` = u as stated; `pairDescriptorValue` (l. 790) evaluates `Xi` at $x(i)$. The Lean search itself ranges over all pairs x , not over the curve; the curve is a subset of its search space. See [section 1.5](#).

4 The two types of pair-partition collision

Two different matchings can share at most one block, since two common pairs force the remaining pair to agree. There are exactly 45 unordered pairs of matchings sharing a block: choose the common pair in 15 ways, then choose two of the three matchings on the four remaining indices. The other $\binom{15}{2} - 45 = 60$ pairs are block-disjoint. Their union is a single alternating six-cycle, since a cycle of length two would be a common block.

For a root pair $e = \{i, j\}$, define

$$q_e(t) = t^2 + s_e t - p_e = t^2 + (\alpha_i + \alpha_j)t - \alpha_i \alpha_j. \quad (4.1)$$

Lemma 4.1 (Common-block collision factor). *If distinct matchings P, Q share the pair e , then*

$$h_P(t) - h_Q(t) = (A_P - A_Q)q_e(t), \quad A_P - A_Q \neq 0. \quad (4.2)$$

Proof. Let the four remaining roots be a, b, c, d . Their two-block pairing affects A through one of

$$z_1 = ab + cd, \quad z_2 = ac + bd, \quad z_3 = ad + bc.$$

The products of the two remaining pair products are always $abcd$. The contribution to B involving only those two blocks is always the third elementary symmetric function of a, b, c, d . Therefore changing the pairing changes B by s_e times the change in A , and changes C by p_e times the change in A . Substitution in [equation \(3.1\)](#) proves the factorization.

The three differences are

$$z_1 - z_2 = (a - d)(b - c), \quad z_1 - z_3 = (a - c)(b - d), \quad z_2 - z_3 = (a - b)(c - d). \quad (4.3)$$

Distinctness of the roots makes every one nonzero. $\square$

Thus the 45 shared-block comparisons do not contribute 90 unrelated exceptional parameters: they contribute roots of only fifteen quadratics. The 60 block-disjoint comparisons each still contribute a quadratic. There are at most $2(15 + 60) = 150$ pair-family failures. Some of these quadratics can have lower degree or share roots after specializing the sextic, so this is a uniform upper bound rather than a claim about every input.

5 Triple collisions as exchanges of two roots

For disjoint root pairs e, e' , set

$$\ell_{e,e'}(t) = (s_e - s_{e'})t - (p_e - p_{e'}). \quad (5.1)$$

Interchanging e, e' changes only the sign. There are $\binom{6}{4} \cdot 3 = 45$ unordered choices of two disjoint pairs.

Lemma 5.1 (Exchange identity). *For the triple partitions*

$$T = \{\{a, b, c\}, \{d, e, f\}\}, \quad T' = \{\{a, b, d\}, \{c, e, f\}\},$$

where letters now denote root values,

$$k_T(t) - k_{T'}(t) = (c - d)((a + b - e - f)t - (ab - ef)). \quad (5.2)$$

Every unordered pair of distinct triple partitions occurs exactly once in this manner, up to the irrelevant orientations, from an unordered pair of disjoint root pairs $\{\{a, b\}, \{c, d\}\}$.

Proof. Subtracting the internal pair sums gives $a_T - a_{T'} = (c - d)(a + b - e - f)$, while subtracting the two triple products gives $b_T - b_{T'} = (c - d)(ab - ef)$. This proves the formula.

For the combinatorial assertion, two distinct partitions can be oriented so that one block from each has two elements in common. Call these a, b ; call the exchanged elements c, d ; the other two elements are e, f . Conversely, two disjoint pairs and their two complementary elements specify exactly the two partitions in the display. Simultaneously switching the chosen blocks or reversing the exchange does not change the unordered data. This establishes the bijection, with 45 elements on either side. $\square$

Lemma 5.2 (No identically vanishing exchange factor). *For disjoint pairs of distinct roots, $\ell_{e,e'}$ is not the zero polynomial.*

Proof. If it vanished identically, the two pair sums and the two pair products would agree. The two monic quadratics $X^2 - s_e X + p_e$ and $X^2 - s_{e'} X + p_{e'}$ would agree, contradicting their disjoint root sets. $\square$

The triple-family failure set is therefore the union of the roots of 45 nonzero linears, and has cardinality at most 45. A nonzero constant factor simply contributes no root.

6 Three invariant obstruction polynomials

Fix total orders on the matchings and on the root pairs, solely to orient differences in products. Define

$$Q_f(t) = \prod_{e \in \mathcal{E}} q_e(t), \quad (6.1)$$

$$U_f(t) = \prod_{\substack{P < Q \\ P \cap Q = \emptyset}} (h_P(t) - h_Q(t)), \quad (6.2)$$

$$L_f(t) = \prod_{\substack{e < e' \\ e \cap e' = \emptyset}} \ell_{e,e'}(t), \quad (6.3)$$

$$E_f(t) = Q_f(t)U_f(t)L_f(t). \quad (6.4)$$

Here $P \cap Q = \emptyset$ refers to their block collections. Reversing a chosen orientation can multiply U_f or L_f by -1 , which changes neither the zero set nor any squared identity below.

It is not automatic that an oriented product of differences is symmetric; the ordinary Vandermonde product is the familiar warning. The following simple sign argument proves what is needed here.

Lemma 6.1 (Graph-product sign test). *Let a finite group act on a finite graph's vertices, permuting quantities w_v . Form an oriented product $\prod_{\{v,w\} \in E} (w_v - w_w)$, with one orientation for each edge. For a group element τ of order two, its sign on this product is $(-1)^r$, where r is the number of edges whose endpoints are interchanged by τ .*

Proof. Edges exchanged with other edges occur in two-element orbits. The two orientation signs in each such orbit multiply to $+1$, since applying τ twice fixes each oriented difference. A fixed edge contributes $+1$ if its endpoints are individually fixed, and -1 if they are interchanged. Multiplying these contributions gives the claim. $\square$

Proposition 6.2 (Symmetry, integrality, and exact failure test). *The expressions Q_f, U_f, L_f are symmetric polynomials with integer coefficients in the six roots and t . They are nonzero for every distinct root tuple and satisfy*

$$\deg Q_f = 30, \quad \deg U_f \leq 120, \quad \deg L_f \leq 45. \quad (6.5)$$

Their coefficients are universal integer polynomials in $e_1, \dots, e_6$. For every $t \in K$,

$$E_f(t) \neq 0 \iff (P \mapsto h_P(t)) \text{ and } (T \mapsto k_T(t)) \text{ are both injective.} \quad (6.6)$$

Proof. The product Q_f is visibly symmetric. For U_f , use the graph with matchings as vertices and block-disjoint pairs as edges. A transposition of two root labels sends any matching either to itself or to a matching sharing its untouched third block. Hence it interchanges the endpoints of no edge in this graph. By [lemma 6.1](#), it fixes U_f . Root transpositions generate S_6 , so U_f is symmetric.

For L_f use the graph on two-element subsets, joined when they are disjoint, with vertex quantity $s_e t - p_e$. A root transposition sends a pair either to itself or to a pair sharing one root. It therefore reverses no disjointness edge, so the same argument proves symmetry of L_f . The fundamental theorem of symmetric polynomials over $\mathbb{Z}[t]$ now yields the integer coefficient expressions in the e_i .

Every factor of Q_f is monic quadratic. Every factor of U_f is nonzero by [proposition 3.1](#), and every factor of L_f is nonzero by [lemma 5.2](#). This proves nonvanishing and the degree bounds. Finally, [lemma 4.1](#) accounts for all shared-block collisions, [equation \(6.2\)](#) accounts for all block-disjoint ones, and [lemma 5.1](#) accounts for all triple collisions. Their scalar root-difference factors are nonzero. This proves [equation \(6.6\)](#). $\square$

Remark 6.3 (Homogeneity and scaling). Assign weight one to every root and to t . Then $h_P, k_T, q_e, \ell_{e,e'}$ have respective total degrees 4, 3, 2, 2. Consequently Q_f, U_f, L_f, E_f have total degrees 30, 240, 90, 360. Scaling all roots and t by c multiplies $E_f(t)$ by c^{360} . These are homogeneous degrees in the roots and parameter together, not the parameter degrees in [equation \(6.5\)](#).

7 Exact discriminant factorizations

Define the compressed resolvents

$$R_{2,f}(Z, t) = \prod_{P \in \mathcal{P}_2} (Z - h_P(t)), \quad R_{3,f}(Z, t) = \prod_{T \in \mathcal{P}_3} (Z - k_T(t)). \quad (7.1)$$

Their coefficients are symmetric integer polynomials in the roots, so they are coefficient-only constructions just as E_f is. Let

$$\Delta_f = \text{disc}(f) = \prod_{1 \leq i < j \leq 6} (\alpha_i - \alpha_j)^2. \quad (7.2)$$

For a monic polynomial with roots $z_1, \dots, z_m$, we use the convention $\text{disc} = \prod_{i < j} (z_i - z_j)^2$.

Theorem 7.1 (Discriminant identities). *As polynomial identities in the roots and t ,*

$$\text{disc}_Z R_{2,f}(Z, t) = \Delta_f^6 Q_f(t)^6 U_f(t)^2, \quad (7.3)$$

$$\text{disc}_Z R_{3,f}(Z, t) = \Delta_f^3 L_f(t)^2. \quad (7.4)$$

In particular the formulas remain valid at nonseparating parameter values.

Proof. The pair-resolvent discriminant is the product of all 105 squared matching differences. The 60 block-disjoint differences give U_f^2 directly. Fix a common pair e . The three unordered comparisons between matchings containing e give, by [lemma 4.1](#),

$$q_e(t)^6 \prod_{1 \leq i < j \leq 3} (z_i - z_j)^2.$$

By [equation \(4.3\)](#), the second factor is exactly the discriminant of the monic quartic whose roots are the other four roots. Multiplying over the fifteen choices of e gives Q_f^6 times the product of these fifteen quartic discriminants. A fixed root difference $(\alpha_i - \alpha_j)^2$ occurs in precisely six of those quartics: choose the excluded pair among the four other roots. Their product is therefore Δ_f^6 , proving [equation \(7.3\)](#).

For triples, [lemma 5.1](#) gives one squared linear factor for each unordered pair of disjoint root pairs, producing L_f^2 . The two leftover roots are the exchanged roots. For a fixed exchanged pair there are three pairings of the other four roots, so its squared difference appears three times. The product of the root-difference factors is Δ_f^3 . This proves [equation \(7.4\)](#). No division was used, so these are polynomial identities, not merely identities on the open set of good parameters. $\square$

Corollary 7.2 (Discriminant square classes). *Over any characteristic-zero coefficient field, the pair-resolvent discriminant is a square. If both f and the triple resolvent are separable, the triple-resolvent discriminant has the same square class as Δ_f .*

Proof. The two formulas read

$$\text{disc}_Z R_{2,f} = (\Delta_f^3 Q_f^3 U_f)^2, \quad \text{disc}_Z R_{3,f} = \Delta_f (\Delta_f L_f)^2.$$

In the second assertion all displayed factors are nonzero. $\square$

There is a useful independent permutation explanation. A transposition of root labels fixes three of the fifteen matchings and interchanges the other twelve in six pairs; it acts evenly. It fixes four triple partitions and interchanges the other six in three pairs; it acts oddly. Thus the fifteen-point action has trivial sign character and the ten-point action has the ordinary six-point sign character. The explicit factorizations identify much more than this parity information: they also isolate every collision factor and its multiplicity.

8 Generic irreducibility of the three components

The degree bound is not merely the result of an inefficient factor count. Generically all its factors are needed. In this section take $x_1, \dots, x_6, t$ to be algebraically independent over $\mathbb{Q}$ and put $S = \mathbb{Q}[x_1, \dots, x_6, t]$. Its symmetric subring is

$$S^{S_6} = \mathbb{Q}[e_1, \dots, e_6, t].$$

The symbols Q, U, L now denote the universal polynomials in this ring, prior to any specialization of the sextic coefficients.

8.1 Irreducibility of the individual root-coordinate factors

Lemma 8.1 (Primitive linear-polynomial test). *Let R be a unique factorization domain. If $A, B \in R$ are coprime and $A \neq 0$, then $Ay + B$ is irreducible in $R[y]$.*

Proof. The polynomial is primitive, and is linear, hence irreducible, over the fraction field of R . Gauss's lemma gives irreducibility over R . $\square$

Lemma 8.2 (The three kinds of factor are irreducible). *In S , each q_e , each $\ell_{e,e'}$ for disjoint pairs, and each $h_P - h_Q$ for block-disjoint matchings is irreducible.*

Proof. For q_e , rename its two roots a, b and write

$$q_e = a(t - b) + t(t + b).$$

The coefficients as a linear polynomial in a are coprime: substituting $t = b$ in the second gives $2b^2 \neq 0$. Apply [lemma 8.1](#). For a linear exchange factor, rename its disjoint pairs $\{a, b\}$ and $\{c, d\}$. Then

$$\ell = a(t - b) + (b - c - d)t + cd.$$

Substituting $t = b$ in its constant coefficient gives $(b - c)(b - d) \neq 0$, so the same test applies. Adjoining unused independent variables preserves these irreducibilities.

For the third type, all block-disjoint matching pairs are related by a root permutation, because their union is an alternating six-cycle. It suffices to consider

$$P = (ab)(cd)(ef), \quad Q = (ac)(be)(df).$$

Write $h_P - h_Q = aA + B$, where $A, B \in \mathbb{Q}[b, c, d, e, f, t]$. This is linear in a because every monomial in [equation \(3.1\)](#) uses each root at most once. It is homogeneous of total degree four; hence A is homogeneous of degree three and B of degree four. At $t = 0$, direct expansion gives

$$\begin{aligned} A_0 &= -bcd + bce - bef + cdf, \\ B_0 &= def(b - c). \end{aligned} \tag{8.1}$$

The polynomial A_0 is divisible by none of $d, e, f, b - c$. Indeed its respective substitutions at zero or at $b = c$ are

$$be(c - f), \quad cd(f - b), \quad bc(e - d), \quad c(e - d)(c - f),$$

all nonzero. Therefore $\gcd(A_0, B_0) = 1$.

Suppose an irreducible nonconstant polynomial H divided both A and B . A factor of a homogeneous polynomial can be chosen homogeneous, so H is homogeneous. If $H|_{t=0}$ is nonzero, it has the same positive total degree and divides both A_0 and B_0 , a contradiction. If it is zero, then t divides H , so irreducibility makes H associated to t ; but $A_0 \neq 0$. Thus A, B are coprime. Apply [lemma 8.1](#) with variable a to finish the proof. $\square$

The homogeneity in the last argument is essential: without it, coprime specializations need not imply coprimality of the original polynomials. Here it prevents a nonconstant common homogeneous factor from specializing to a nonzero constant.

8.2 Descent from root coordinates to coefficients

Lemma 8.3 (One-orbit product lemma). *Let a finite group act on a polynomial ring over a field. Suppose a finite collection of pairwise nonassociate irreducible polynomials forms a single orbit up to units, and its product F is invariant. Then F is irreducible in the invariant subring.*

Proof. A factorization $F = AB$ in the invariant subring is also a factorization in the ambient unique factorization domain. The irreducible factors of A therefore form a subset of the factors of F . Invariance makes this subset stable under the group. Transitivity forces it to be empty or the whole set. In the first case A is a unit, and in the second B is a unit. $\square$

Theorem 8.4 (Generic three-factor theorem). *The universal polynomials Q, U, L are pairwise nonassociate irreducible polynomials in $\mathbb{Q}[e_1, \dots, e_6, t]$, and are irreducible in $\mathbb{Q}(e_1, \dots, e_6)[t]$. Their degrees in t are respectively 30, 120, 45. Their product E is squarefree over the generic coefficient field and has degree 195 and nonzero constant coefficient.*

Proof. The individual factors are irreducible by [lemma 8.2](#). The symmetric group is transitive on root pairs, on unordered pairs of disjoint root pairs, and on unordered pairs of block-disjoint matchings. In each collection the individual factors are pairwise nonassociate. For q_e , this follows from its monic leading term and its two root indices. For $\ell_{e,e'}$, the coefficient of t specifies the two pairs with opposite signs, up to their interchange. For $h_P - h_Q$, the coefficient of t^2 is $A_P - A_Q$: its six distinct root-product monomials, with coefficients $+1$ or -1 , specify the two block-disjoint matchings up to interchange. Thus no further associations occur.

Each product is invariant by [proposition 6.2](#). The one-orbit product lemma proves irreducibility in the symmetric subring. The leading coefficient in t of each individual q is one, of each individual h difference is a nonzero $A_P - A_Q$, and of each individual ℓ is a nonzero difference of pair sums. The ambient ring is a domain, so the three parameter degrees are exactly 30, 120, 45. They are consequently pairwise nonassociate. Their irreducibility and positive parameter degree also imply that they are primitive over $\mathbb{Q}[e_1, \dots, e_6]$. Gauss's lemma proves irreducibility over its fraction field.

In characteristic zero every irreducible polynomial of positive degree is separable. The three are pairwise coprime there, so E is squarefree and has degree 195. Its constant coefficient is nonzero: every $q_e(0)$ and $\ell_{e,e'}(0)$ is a nonzero polynomial, and every block-disjoint h difference at zero is nonzero by the representative calculation [equation \(8.1\)](#) and root permutation. $\square$

This theorem is a statement about the generic coefficient field, not about every numerical sextic. After specialization a factor can become reducible, lose degree, or share roots with another factor. None of these possibilities threatens the uniform separator theorem; they only reduce the number of bad parameters. On the open coefficient locus $\Delta_f \neq 0$, the three polynomials describe the three irreducible families of bad specializations over $\mathbb{Q}$. The constructive factor proofs also work after extending the constant field to any characteristic-zero field.

9 The finite separator theorem and its sharpness

Theorem 9.1 (Uniform simultaneous separation). *For any six distinct roots in a characteristic-zero field, at most 195 parameters cause a collision in either compressed family. Therefore any 196 distinct nonzero test parameters contain a simultaneous separator. The integers $1, \dots, 196$ always contain one, and this same parameter separates both original descriptors on $(u, v) = (t, 2t^2)$.*

Proof. By [proposition 6.2](#), the bad parameters are exactly the roots of the nonzero polynomial E_f , which has degree at most $30 + 120 + 45 = 195$. A nonzero polynomial over a field has at most its degree many distinct roots. Characteristic zero makes the positive integers $1, \dots, 196$ distinct and nonzero. Finally use [equations \(3.3\) and \(3.6\)](#), whose common multiplier is nonzero at these parameters. $\square$

The same proof gives 151 test values for the pair family alone and 46 for the triple family alone. A common parameter is preferable when a single uniform interface is wanted; separate searches can stop sooner on some inputs. There is no requirement that the least good pair parameter agree with the least good triple parameter.

Theorem 9.2 (Sharpness for arbitrary test sets). *The constant 196 in the assertion about every set of distinct nonzero complex test parameters cannot be lowered. There are separable sextics with exactly 195 distinct nonzero bad parameters. Such a sextic can even be chosen split with integer roots.*

Proof. Over the generic coefficient field, [theorem 8.4](#) gives a degree-195 squarefree polynomial with nonzero constant coefficient. Thus its leading coefficient, constant coefficient, and discriminant in t are all nonzero. Multiply these coefficient polynomials by Δ_f and substitute the elementary symmetric functions of six independent root variables. The result is still a nonzero polynomial in those variables, because the symmetric-polynomial substitution is injective.

The integer lattice is Zariski dense in affine six-space: a nonzero polynomial over $\mathbb{Q}$ cannot vanish on all integer tuples, as follows by induction on the number of variables and the one-variable root bound. Choose an integer tuple where the displayed product is nonzero. Its roots are distinct, and its E_f has degree 195, no zero root, and no repeated root. Over $\mathbb{C}$ it has exactly 195 distinct nonzero roots. These are all bad by [equation \(6.6\)](#); taking them as the test set disproves any 195-point guarantee of the stated kind. $\square$

9.1 An explicit sharpness certificate

The existence proof above is entirely algebraic. The accompanying code also gives a concrete witness with roots

$$1, 4, 10, 23, 51, 109. \tag{9.1}$$

For the orientation used in the verifier, reduction modulo $p = 1,000,003$ gives

$$\deg \bar{E}_f = 195, \quad \bar{E}_f(0) = 521291, \quad \text{lc}(\bar{E}_f) = 670728. \tag{9.2}$$

The file `checks/sharpness_certificate.json`¹ stores the coefficients of $\overline{E}_f$ and polynomials A, B satisfying the exact identity

$$A\overline{E}_f + B\overline{E}'_f = 1 \quad \text{in } \mathbb{F}_p[t]. \quad (9.3)$$

The verifier checks primality of p by trial division and checks this identity by finite polynomial arithmetic. Degree preservation, the nonzero constant coefficient, and [equation \(9.3\)](#) imply that the integer polynomial E_f has degree 195 and 195 distinct nonzero complex roots. This is a reproducible computational certificate, not a Lean-checked one. It is supplementary: [theorem 9.2](#) does not rely on its execution.

9.2 The initial integer interval is a different problem

Define N_{int} to be the least positive integer N such that $1, \dots, N$ contains a simultaneous separator for every distinct complex root tuple. The sharpness proof using complex roots of E_f does *not* prove $N_{\text{int}} = 196$.

Proposition 9.3 (An eight-point lower bound). *For this initial-interval problem,*

$$8 \leq N_{\text{int}} \leq 196. \quad (9.4)$$

Proof. The upper bound is [theorem 9.1](#). For the lower bound use the distinct integer roots $(-3, 0, 1, 2, 3, 4)$, whose polynomial is

$$X^6 - 7X^5 + 5X^4 + 55X^3 - 126X^2 + 72X.$$

Direct substitution in [equations \(3.1\)](#) and [\(3.4\)](#) gives the following exact numbers of distinct values:

t	1	2	3	4	5	6	7	8
$ \{h_P(t)\} $	12	13	13	14	15	15	15	15
$ \{k_T(t)\} $	10	8	9	10	9	8	9	10

Thus none of $1, \dots, 7$ works. For additional explicit checks, the sorted pair values at $t = 8$ are

$$\begin{aligned} &-1128, -1044, -870, -792, -724, -486, -432, -424, \\ &72, 90, 192, 396, 776, 816, 936, \end{aligned}$$

and the sorted triple values are

$$-32, -28, -22, -20, -14, 1, 32, 46, 92, 160.$$

They are all distinct within their respective families. □

Proposition 9.4 (No universal single positive parameter). *For each fixed positive integer t_0 , there is a split separable integer sextic for which t_0 fails to separate the pair family.*

Proof. Take roots $0, t_0, 2t_0, 3t_0, 4t_0, 5t_0$. For the common pair $\{2t_0, 3t_0\}$, its quadratic satisfies $q(t_0) = t_0^2 + 5t_0^2 - 6t_0^2 = 0$. Any two different matchings containing that pair collide by [lemma 4.1](#). □

The example is reducible. It does not establish that a fixed parameter must fail on some irreducible rational sextic; that is a separate question.

¹[write] Shipped as `data/sharpness_certificate.json`; the delivered `checks/` directory is `data/` here.

10 Why the separated resolvents decide an irreducible sextic

The group-theoretic criterion in this section is classical. We include its proof to show exactly what the new separator supplies and what it does not. It is the criterion underlying the degree-fifteen and degree-ten sextic resolvents in the literature and the repository [2, 6]. Let $f \in \mathbb{Q}[X]$ now be irreducible of degree six, let M be its splitting field, and view $G = \text{Gal}(M/\mathbb{Q})$ as a transitive subgroup of S_6 .

Lemma 10.1 (Solvable primitive groups have prime-power degree). *A finite solvable group acting faithfully and primitively on a set with more than one element has degree p^r for some prime p .*

Proof. Choose a minimal nontrivial normal subgroup N of G . Because G is solvable, N is elementary abelian. Here is the standard short justification: a last nontrivial term of the derived series of N is an abelian characteristic subgroup of N , hence normal in G , so minimality makes N abelian. Its primary components are characteristic, so only one prime p occurs. The characteristic subgroup of elements killed by p is nontrivial and hence all of N .

The orbits of a normal subgroup form a G -invariant partition. Primitivity and faithfulness force N to be transitive: otherwise its orbits are singletons, making N act trivially. An abelian transitive permutation group is regular. Indeed, an element fixing one point fixes every point by commuting with all elements that carry it to another point. Faithfulness then makes the stabilizer trivial. Thus the degree is $|N| = p^r$. $\square$

Proposition 10.2 (The two block systems are complete). *For a transitive subgroup $G \leq S_6$, the following are equivalent: G is solvable; G is imprimitive; G preserves a partition in $\mathcal{P}_2$ or in $\mathcal{P}_3$.*

Proof. A solvable primitive group cannot have degree six by lemma 10.1. Thus solvability implies imprimitivity. For a transitive action, the blocks in a nontrivial block system have a common size properly dividing six, so that size is two or three. This proves the next implication.

The full stabilizer of a pair partition is $S_2 \wr S_3$, an extension of S_2^3 by S_3 . The full stabilizer of a triple partition is $S_3 \wr S_2$, an extension of S_3^2 by S_2 . Both are solvable, since finite direct products and extensions of solvable groups are solvable. Any subgroup preserving the corresponding partition is a subgroup of its stabilizer and is therefore solvable. $\square$

Theorem 10.3 (Separated rational-root criterion). *Let $f \in \mathbb{Q}[X]$ be an irreducible monic sextic, and let $t \in \mathbb{Q}$ be a simultaneous separator. Then f is solvable by radicals if and only if at least one of $R_{2,f}(Z, t)$ and $R_{3,f}(Z, t)$ has a rational root.*

Proof. For a matching P and $\sigma \in G$, coefficient equivariance gives $\sigma(h_P(t)) = h_{\sigma P}(t)$, since t is rational. If P is fixed by G , then $h_P(t)$ is fixed by G and lies in $\mathbb{Q}$. Conversely, if $h_P(t)$ is rational, then $h_{\sigma P}(t) = h_P(t)$ for every σ , and separation gives $\sigma P = P$. A rational root of the pair resolvent is one of its explicit values, so it is equivalent to a G -fixed pair partition. The same proof applies to k_T and the triple resolvent. Now use proposition 10.2 and the classical equivalence between solvability by radicals and solvability of the Galois group. $\square$

Separation is not a cosmetic condition in the converse. Without it, $h_{\sigma P}(t) = h_P(t)$ would not imply $\sigma P = P$. A specialized value could be fixed while its underlying partition moves among colliding partitions. This is precisely the logical failure prevented by the nonvanishing certificate $E_f(t) \neq 0$.

For a monic integer sextic, both resolvents are monic integer polynomials in Z . Any rational root is therefore an integer, so the final condition can be decided using a finite integer interval. No splitting field or numerical root approximation needs to occur in the returned-data algorithm.

11 A bounded coefficient-only decision architecture

11.1 Computing the invariant coefficients without roots

A root-coordinate definition can hide a computational oracle unless the coefficient bridge is made explicit. Here the bridge is the constructive fundamental theorem of symmetric polynomials. For each fixed power of Z in [equation \(7.1\)](#), its coefficient is a fixed symmetric polynomial in six roots with coefficients in $\mathbb{Z}[t]$. It can be reduced to a fixed polynomial in $e_1, \dots, e_6, t$ using integer arithmetic only.

For completeness, the reduction is finite in a particularly elementary sense. Order root monomials lexicographically with $x_1 > \dots > x_6$. The leading exponent vector of a symmetric polynomial has the form $\lambda_1 \geq \dots \geq \lambda_6$. The product

$$e_1^{\lambda_1 - \lambda_2} e_2^{\lambda_2 - \lambda_3} \dots e_5^{\lambda_5 - \lambda_6} e_6^{\lambda_6} \quad (11.1)$$

has exactly that leading root monomial, with coefficient one. Subtract its multiple by the leading coefficient, which is a polynomial in t . The leading monomial strictly decreases and the total root degree never increases. There are only finitely many root monomials within the original degree bound, so the process terminates with an expression over $\mathbb{Z}[e_1, \dots, e_6, t]$. This proof gives a finite symbolic construction, not a choice of an unspecified algebraic root.

For the coefficient of Z^{15-j} in $R_{2,f}$, the root degree is at most $4j$ and parameter degree at most $2j$. For the coefficient of Z^{10-j} in $R_{3,f}$, they are at most $3j$ and j . Hence degrees at most 60 and 30 in the root variables suffice for the two complete coefficient tables. These are fixed bounds, independent of the input height. The analogous bound for E is 360 in the root variables and 195 in t .

The article does not distribute expanded versions of these potentially large universal tables. Their construction and their evaluation are mathematically specified by the fixed finite products and the reduction above. The supplied executable code is instead a verifier on exact root tuples. This distinction is important when assessing what has and has not been implemented.

11.2 Monicization and bounded factor search

Let the input be an integer polynomial $g(X) = a_6 X^6 + \dots + a_0$, with $a_6 \neq 0$. Define

$$f(Y) = a_6^5 g(Y/a_6) = Y^6 + \sum_{i=0}^5 a_i a_6^{5-i} Y^i. \quad (11.2)$$

This is monic integral, and its roots are a_6 times those of g . Multiplication and division by the nonzero rational a_6 preserve the radical closure of $\mathbb{Q}$, so this transformation preserves all-root radical solvability. The repository already provides this semantic bridge [8].

Write H for the maximum absolute value of the six lower coefficients of f , and let $R = 1 + H$. Every complex root of f has absolute value at most R . For example, if $|z| > 1 + H$, the geometric sum estimate gives

$$H \sum_{i=0}^5 |z|^i < |z|^6,$$

so the leading term cannot be canceled by the others. If a monic degree- d integer polynomial divides f , its roots are a submultiset of these roots, and the coefficient at distance j below its leading term has absolute value at most

$$\binom{d}{j} R^j. \quad (11.3)$$

Gauss's lemma implies that rational reducibility of a monic integer polynomial has a monic integer factor. Any proper factorization of a sextic has a factor of degree at most three. Thus enumeration of the finite coefficient boxes in [equation \(11.3\)](#), for $d = 1, 2, 3$, completely decides reducibility.

A degree-two or degree-three factor leaves a quotient of degree at most four; then every root is radical. A degree-one factor leaves a quintic, whose radical solvability must actually be tested. In particular, one must not accept every reducible sextic: a nonsolvable quintic multiplied by a linear factor is the basic counterexample. The existing primitive-recursive quintic decision can be reused [\[4, 10\]](#). Repeated factors are handled by these reducible branches; the irreducible branch is automatically separable in characteristic zero.

11.3 The algorithm

The following is a mathematical specification of a finite coefficient algorithm. A default return on malformed input makes it a total Boolean on seven encoded integers; the target predicate includes exact degree six.

Bounded sextic radical criterion.

1. Reject if $a_6 = 0$. Otherwise form the monic integer polynomial f in [equation \(11.2\)](#) and the root bound R .
2. Search the explicit boxes for a monic factor of degree two or three. If one exists, accept.
3. Search the explicit interval for a linear factor. If one exists, choose the first in a fixed order, form its quintic quotient, and return the existing primitive-recursive quintic criterion.
4. The remaining input is irreducible. For $t = 1, \dots, 196$, evaluate the universal coefficient polynomials of the two compressed resolvents. Choose the first t for which both are squarefree, or, equivalently, for which $E_f(t) \neq 0$.
5. Accept exactly when one of the two resulting monic integer resolvents has an integer root. For each resolvent, test all integers in its Cauchy interval $[-(1 + H_R), 1 + H_R]$, where H_R is the maximum absolute value of its lower coefficients.

Squarefreeness in step 4 can be tested by a rational polynomial gcd with the derivative, by a fixed-size resultant determinant, or by the obstruction value itself. For an especially direct primitive-recursive implementation, the fixed-size determinant avoids any concern about a hidden unbounded gcd loop. The resolvent degrees are always fifteen and ten.

Theorem 11.1 (Bounded architecture and correctness). *The coefficient procedure above has a primitive-recursive Boolean realization and accepts exactly the integer inputs of degree six whose complex roots are all expressible by radicals over $\mathbb{Q}$.*

Proof. Monicization is integer arithmetic with fixed exponents. The factor boxes have primitive-recursive bounds, and polynomial division for their fixed degrees is a finite coefficient computation. The correctness of their branches follows from [equation \(11.3\)](#) and the degree arguments above. The quintic subroutine is the established primitive-recursive criterion, not an unbounded search for radical expressions.

On the irreducible branch, [theorem 9.1](#) guarantees that step 4 finds a parameter. Each coefficient is the evaluation of a fixed integer polynomial. Evaluation, fixed-size determinants, a loop of length 196, and bounded selection are primitive recursive under the usual effective encodings of integers and finite tuples. Step 5 is a bounded integer search, and is complete because the resolvents are monic and the Cauchy bound includes all their integer roots. Its correctness is [theorem 10.3](#). Composing these primitive-recursive operations proves the claim. $\square$

This theorem concerns a mathematical realization. It does not assert that the existing Lean function has already been changed, that a `Primrec` proof has been accepted by the kernel, or that the attached verifier is this coefficient solver. A source-level integration must additionally supply the coefficient-evaluation lemmas and bounded-selector code discussed in [section 14](#).

Nor does the naive factor and integer-root enumeration above claim polynomial bit complexity. Its bounds depend polynomially on coefficient *magnitudes*, which can still give exponential running time in their bit lengths. The stronger general polynomial-time theorem is classical [1]; our objective is a small, transparent invariant search with a tractable verification boundary.

12 A separator determined by coefficient height alone

The finite search has only 196 candidates, but it is also possible to give one coefficient-dependent parameter that is guaranteed to work without any separation test. It is useful as a termination certificate or a comparison construction, not as a recommendation for practical evaluation.

For a polynomial $P(t) = \sum c_i t^i$ over $\mathbb{C}$, write $\|P\|_1 = \sum |c_i|$. This norm is submultiplicative. Let $f \in \mathbb{Z}[X]$ be monic and separable, put $R = 1 + H$ as above, and use $|\alpha_i| \leq R$. Then

$$\begin{aligned} \|q_e\|_1 &\leq 1 + 2R + R^2 \leq 4R^2, \\ \|h_P - h_Q\|_1 &\leq 6R^2 + 24R^3 + 6R^4 \leq 36R^4, \\ \|\ell_{e,e'}\|_1 &\leq 4R + 2R^2 \leq 6R^2. \end{aligned} \tag{12.1}$$

For the middle estimate, $|A_P| \leq 3R^2$, $|B_P| \leq 12R^3$, and $|C_P| \leq 3R^4$; subtracting two such quantities doubles each bound. Multiplying the bounds for fifteen, sixty, and forty-five factors gives

$$\|E_f\|_1 \leq 4^{15} 6^{165} R^{360} =: M(H). \tag{12.2}$$

Theorem 12.1 (A search-free height separator). *The positive integer*

$$t_* = 4^{15} 6^{165} (1 + H)^{360} + 2 \tag{12.3}$$

separates both compressed families and both original descriptors on the curve.

Proof. The nonzero polynomial E_f has integer coefficients, so its leading coefficient has absolute value at least one. Its Cauchy root bound is at most $1 + M(H)$ by [equation \(12.2\)](#). Hence $E_f(t_*) \neq 0$. Apply [equation \(6.6\)](#) and the nonzero-multiplier identities. $\square$

Although t_* is numerically enormous, its bit length is $O(1 + \log(1 + H))$ with a large absolute constant. Its role is to show that even separator selection itself can be replaced by an explicit arithmetic expression. The first successful candidate in $1, \dots, 196$ will generally be much smaller and avoid unnecessary coefficient growth.

13 Exact checks, certificates, and their limits

The attached computations are designed to test local algebra, combinatorial counts, discriminant multiplicities, symmetry signs, and explicit examples. They are not used as evidence that an unproved general statement should be accepted. Each general theorem above has a written argument, including the irreducibility and arbitrary-test-set sharpness results.

13.1 Local symbolic identities

The script `code/verify.py` uses SymPy to expand and check four universal identities: the pair and triple descriptor expansions, the common-block factorization, and the triple exchange factorization. These checks involve independent root symbols, rather than approximate roots of sample polynomials. They therefore test the precise signs and coefficients in [equations \(3.2\), \(3.5\), \(4.2\) and \(5.2\)](#).

13.2 Integer arithmetic checks

The same script enumerates the fifteen matchings and ten triple partitions without a stored table. It checks the $45 + 60$ decomposition of matching comparisons and the 45 disjoint-root-pair comparisons. For each root transposition it checks that no edge of the block-disjoint matching graph is reversed, matching the sign argument in [lemma 6.1](#).

For four distinct integer root tuples and every $t = 0, \dots, 16$, the script checks both discriminant identities using exact integer products, and verifies the compressed-to-descriptor identities and the exact obstruction test. This yields 68 parameter cases, each testing both resolvent discriminants. It also checks invariance of the three factors under all fifteen root transpositions for those four tuples, giving 60 exact symmetry cases.

It exhausts all $\binom{11}{6} = 462$ six-element subsets of $\{-3, -2, \dots, 7\}$ and computes their first positive simultaneous separator. The resulting distribution is

First separator	1	2	3	4	5	6	7	8
Number of tuples	93	83	89	125	48	15	8	1

These finite data do not support a claim that eight works universally. They provide reproducible examples, including the lower-bound witness in [proposition 9.3](#). Twenty scaled examples additionally check the failure of fixed parameters described in [proposition 9.4](#).

13.3 A modular certificate for the sharp example

The independent standard-library script `code/sharpness_certificate.py` constructs E_f modulo 1,000,003 for [equation \(9.1\)](#). It uses only its 120 explicit factors, computes an extended Euclidean identity with its derivative, and checks [equation \(9.3\)](#). The certificate stores every coefficient in ascending order, so an independent implementation can verify the identity by multiplication and addition without repeating the Euclidean algorithm.

Because reduction preserves the degree and gives gcd one, the characteristic-zero polynomial is squarefree. Equivalently, the resultant with its derivative is nonzero modulo the prime, hence is a nonzero integer. This proves that the example reaches 195 distinct nonzero complex failures for this parameterization. It does not imply that any of those failures, let alone the first 195 positive integers, are all integers or all real.

13.4 Verification status

Both scripts were executed successfully for this report. The package records the exact results, the complete modular identity, and the scripts needed to reproduce them. There was no Lean or Rocq execution for the new results, no rebuild of the repository, and no benchmark against production Galois software. The code supplied is an exact verifier and root-level prototype, not a completed coefficient-only radical solver.

14 A precise route back into ProveIt

All repository statements here refer to snapshot `e2b1f016a94102663f12b970e6dd434229f90d01`. The relevant source directory is `Algebra/PolynomialFormulas/Lean/PolynomialFormulas/`. A subsequent repository version may have changed its interfaces or already closed some of these obligations.

14.1 Reusable existing boundaries

The source already provides descriptor injectivity and evaluated-resolvent separability equivalences in `SexticSeparatingInvariants.lean`; the irreducible decision combines the chosen parameters with the Galois criterion in `SexticIrreducibleDecision.lean`. The final all-roots predicate and both computable and primitive-recursive criterion interfaces are in `SexticRadicalComputability.lean`. Monicization is separated cleanly into `SexticRadicalSemantics.lean` [7, 6, 9, 8].

The reducible branch also contains a smaller search issue: `linearTotalCode` uses `Nat.find`, even though a finite symmetric integer-code bound is already present in `linearWitnessB`. A complete primitive-recursive integration must replace or bound that selector as well, not stop after fixing the pair and triple search [10].

[write] Two files this section does not name. The unguarded pair and triple searches, `pairSeparatingCode` and `tripleSeparatingCode`, are in `SexticSeparatingSearch.lean` [11]; `SexticIrreducibleDecision.lean` wraps them in the total searches `pairTotalCode` and `tripleTotalCode`. A third unbounded search, `elementaryCode` in `SexticSparseSymmetricSearch.lean` [12], finds the elementary-symmetric certificates for the resolvent coefficients and the collision products. Declarations and lines are in [section 1.5](#).

14.2 Suggested module sequence

A useful first module, tentatively `SexticCompressedDescriptors.lean`, would define A, B, C, a, b, h, k and prove [equations \(3.2\) and \(3.5\)](#). These are finite ring identities, followed by short applications of the existing descriptor injectivity theorem. Its main public result should identify the current parameter $x(0) = 2t^2, x(1) = t$ with the compressed values through the common nonzero affine transformation.

A second module, `SexticCollisionFactors.lean`, would formalize the shared-block/exchange identities and the 45/60/45 finite partition counts. It can obtain the 195 bad-value bound directly from a product over the finite factors, without first formalizing generic irreducibility or the full discriminant formulas. Thus the shortest route to the bounded selector is substantially smaller than the whole article.

A third module, `SexticBoundedSeparatingSearch.lean`, would prove that one of $1, \dots, 196$ works and implement a fixed-length list scan. Its return specification should establish simultaneous injectivity, followed by the existing resolvent correctness bridge. A total fallback value can be present in the definition, but its unreachable status on separable inputs must be proved rather than assumed.

The computational module should then prove primitive recursiveness of each fixed universal coefficient evaluation, of the squarefreeness test, and of the bounded selector. One legitimate shortcut is to specialize at each of the 196 constant parameters and unroll the finite search, provided the coefficient evaluators themselves have primitive-recursive proofs. Merely replacing `Nat.find` by a bounded loop around a function known only to be computable would not establish `Primrec`.

Finally, replace the linear-factor selector by its existing bounded interval, reuse the primitive-recursive quintic branch, and compose the final criterion. The independent Rocq route can use the same mathematical bound and compact collision products, but no claim of a completed port is made here.

14.3 Separate proof products from optional refinements

The exact discriminant identities, generic irreducibility, and the 195-failure sharpness example deserve their own modules. They make the mathematical story stronger, but they are not prerequisites for a correct bounded sextic decision. Keeping them separate would preserve a small trusted dependency chain for the computational endpoint.

A sensible audit should report four distinct endpoints: the universal mathematical separator theorem; the correctness of the bounded returned parameter; a primitive-recursive coefficient function; and the final all-roots radical predicate. Exact sample evaluations are a fifth product, not a substitute for any of the first four.

15 A reusable general finite-separation principle

The sextic construction is special in its low degrees and factorization, but the idea of converting polynomial descriptors to a finite hitting set has an elementary general form. The following statement explains why an existence proof by avoiding finitely many polynomial equalities often admits an explicit bounded counterpart.

Proposition 15.1 (Kronecker-curve separation). *Let $F_1(u, v), \dots, F_m(u, v)$ be distinct polynomials over a characteristic-zero field, each having u -degree at most a and v -degree at most b . Put*

$$G_i(t) = F_i(t, t^{a+1}), \quad D = a + b(a + 1).$$

Then the G_i are distinct, have degree at most D , and among any $D \binom{m}{2} + 1$ distinct parameter values there is one at which all their values are different.

Proof. A monomial $u^i v^j$ becomes $t^{i+(a+1)j}$. For $0 \leq i \leq a$, the exponent uniquely determines i, j by division with remainder by $a + 1$. Thus the substitution is injective on the stated polynomial space and preserves nonzero differences. The product of the $\binom{m}{2}$ differences is nonzero of degree at most $D \binom{m}{2}$, and the root bound applies. $\square$

This is a standard monomial-encoding argument, not a novelty claim. Its contrast with the sextic result is instructive. A general curve must preserve all descriptor coefficients. Here we first subtract the partition-independent part, use the coefficient relations to compress the family, and then recognize common collision factors. The result is a quadratic curve, a low-degree obstruction, and an exact generic factor count rather than a large undifferentiated degree estimate.

For other resolvent actions the promising order of work is therefore: identify a complete descriptor; isolate its varying coefficients; find a low-degree curve faithful to those coefficients; organize collision pairs by group orbits; and only then use a root bound. The graph-product sign test and the one-orbit product lemma can be reused independently of the sextic setting.

16 Further research questions and priorities

The results leave several concrete problems. They fall into three groups: sharp arithmetic separation, the geometry of the obstruction, and certified implementation. The distinction between the solved arbitrary-set bound and the unresolved integer-interval bound should be retained in all follow-up work.

Question 16.1 (The optimal initial integer interval). Determine N_{int} , currently bracketed by $8 \leq N_{\text{int}} \leq 196$ in this report. A proof that eight suffices would require new control over simultaneous integer roots of the three factors, not just an improvement in generic complex root counting. Conversely, construct a distinct root tuple for which $1, \dots, 8$ all fail.

A practical first experiment is an exact search over larger integer and rational root sets, recording which factor accounts for each failed parameter. The combinatorial pattern of overlaps may be more informative than the maximum alone. Any proposed universal improvement should be proved independently of a finite sample.

Question 16.2 (Short fixed test sets not forming an interval). What is the smallest cardinality of a fixed set $S \subset \mathbb{Z}_{>0}$ that separates every distinct complex root tuple? The sharpness theorem for arbitrary input-dependent complex test sets does not answer this question. Can a sparse integer set outperform an initial interval substantially?

Question 16.3 (Irreducible rational inputs). Do the best uniform bounds decrease when f is required to be irreducible over $\mathbb{Q}$? For a prescribed positive integer t_0 , can one construct an irreducible rational sextic with $E_f(t_0) = 0$? The split counterexamples in this article deliberately do not settle this.

A route through rational points on the three collision hypersurfaces would have to preserve irreducibility of the sextic, rather than merely produce a generic point in root coordinates.

Question 16.4 (Alternative low-degree curves). Can a different scalar compression or a different curve in descriptor space lower the maximum number of exceptional parameters below 195? The proof of sharpness applies to the particular h, k parameterization, not to every possible separating invariant.

The common-block factorization suggests looking for a curve that collapses more repeated comparisons without making an entire family identical. Any such proposal must still distinguish all partitions for every distinct root tuple, including special symmetric configurations.

Question 16.5 (Geometry of the three components). Describe the intersections, singularities, and branching of the three irreducible hypersurfaces $Q = 0$, $U = 0$, and $L = 0$ over the separable-sextic coefficient locus. Which specializations cause degree loss, repeated parameter roots, or intersections between the components?

The generic irreducibility theorem reduces this from an unspecified collision locus to three explicit orbit products. Their normalizations and projection monodromy are natural next invariants.

Question 16.6 (Galois groups of the obstruction factors). Determine the Galois groups over $\mathbb{Q}(e_1, \dots, e_6)$ of the degree-30, degree-120, and degree-45 parameter polynomials. Their factorizations into quadratics and linears over the generic sextic splitting field provide strong constraints, but do not by themselves determine the full groups.

Question 16.7 (Compact coefficient circuits). Construct certified straight-line programs for the two resolvents or for $E_f(t)$ that avoid expanding large symmetric-polynomial tables. Compare Newton identities, traces in finite algebras, and modular reconstruction, with exact operation counts and coefficient-height estimates.

A useful first target is not the entire final solver but an independently checkable evaluator for the obstruction at a supplied integer parameter.

Question 16.8 (End-to-end formal primitive recursiveness). Implement the module sequence in [section 14](#), close the bounded linear-selector issue, and obtain a kernel-checked `Primrec` theorem for the complete coefficient criterion. Port the same bounded search to the independent Rocq development.

The decisive artifact would be a verified executable Boolean and its all-complex-roots specification, not only a classical existence theorem for a suitable natural number.

Question 16.9 (Modular certificates and practical performance). Can a small collection of finite-field evaluations certify good parameters and rational-root decisions substantially faster than direct integer expansion? Establish explicit reconstruction bounds and fallback conditions so that modular success is a certificate rather than a probabilistic guess.

Question 16.10 (Positive characteristic and field size). Determine the precise characteristics and finite-field sizes for which analogues hold. The compressed formulas and many collision identities survive algebraically, but the multiplier $2t^2$, the distinctness of integer parameters, and generic irreducibility arguments must be revisited. Extension-field parameters may be necessary.

Question 16.11 (Higher-degree block systems). For partitions into equal-sized blocks in higher degrees, classify collision orbits and search for compressed descriptors with similarly small invariant products. Which actions admit a graph-product factorization and irreducible orbit factors of manageable degree?

Question 16.12 (From a decision to explicit radical expressions). Use a certified rational resolvent root to recover a fixed block system and then construct branch-compatible radical expressions. A Boolean solvability certificate is not yet an explicit all-roots formula; the recovery and radical-tower interfaces require separate proofs.

17 Conclusion

The repository’s unbounded separating search can be replaced by a mathematically explicit finite mechanism. The central reason is not a general computability theorem, but a small amount of partition-dependent information: three coefficients for pair partitions, two for triple partitions, and a quadratic curve that preserves them. The resulting collision factors have enough structure to yield an exact discriminant factorization and a generic irreducibility theorem.

This resolves the bounded-search obligation at the written-proof level and identifies a precise next formalization target. It also settles a sharpness question for arbitrary complex test sets while leaving the more arithmetic positive-integer problem genuinely open. Keeping these scopes separate is necessary both for mathematical accuracy and for a useful research program.

[write] Status of the obligation in ProveIt. In the Lean and Rocq developments the obligation stays open: the separating parameters are still found by unbounded minimization, and formalizing [theorem 9.1](#) would bound only the parameter selector ([section 1.5](#)).

A Reproducibility and package contents

The archive contains the self-contained source `article.tex`, its compiled `article.pdf`, `README.md`, a build script, two Python scripts, the exact verification report, and the modular Bézout certificate. Bibliographic entries are embedded in the TeX source; no separate bibliography tool is required. A checksum file records the packaged artifacts.

Compile the document by running

```
./build.sh
```

from the package directory. The script uses `pdflatex` three times and writes the PDF in the same directory. The source uses commonly available TeX Live packages, including `newtx`, `microtype`, and `tcolorbox`.

Run the checks with

```
python code/verify.py
python code/sharpness_certificate.py
```

The first script needs SymPy for its four local symbolic identities. The second uses only the Python standard library and the combinatorial and integer-arithmetic functions defined in `verify.py`; importing that module does not import SymPy or run its test suite. The certificate polynomials use ascending coefficient order. The default scripts use assertions for test failures and should be run without Python’s optimization flag.

[write] The shipped layout. This appendix describes the delivered archive. In the collection the report directory holds `article.tex`, a build of it as `article.pdf`, a report `README.md` (not the delivered one), `code/verify.py`, `code/sharpness_certificate.py`, `code/build.sh`, and `data/verification.json`, `data/sharpness_certificate.json`, `data/environment.json`, `data/requirements.txt`; the delivered PDF, `README` and checksum file are not shipped. `code/build.sh` changes to its own directory, where there is no `article.tex`; build instead with `latexmk-pdf` from a copy of the report directory. Each script writes its output into a directory `checks/` next to the `code/` directory that contains it, so run them on a copy of `code/`, never in place, and compare the new `checks/X` with `data/X`; the `README` gives the commands.

B Source and claim ledger

The following ledger separates historical dependencies, inspected source status, written results, and executed checks.

Claim or artifact	Basis and status
General radical-solvability decidability	Classical; Landau–Miller proves polynomial-time decidability. Not a new result of this article.
Sextic pair/triple group criterion	Classical; proved again in section 10 to expose the exact role of separation.
Complete block descriptors	Inherited from the inspected repository; injectivity reproved in lemma 2.1 .
Quadratic/linear compression	Written proof in section 3 ; local identities checked symbolically.
Three invariant factors and discriminants	Written proofs in sections 6 and 7 ; exact integer tests supplied.
Generic irreducibility and 196 sharpness	Written proofs in sections 8 and 9 ; explicit sharp example additionally has a modular certificate.
Initial interval optimum	Not determined; this report proves only $8 \leq N_{\text{int}} \leq 196$.
Primitive-recursive coefficient criterion	Mathematical construction and proof in section 11 ; not an executed Lean development or a packaged extracted solver.
Executable checks	Both attached scripts executed successfully; exact outputs in <code>checks/</code> (shipped as <code>data/</code> ; [write]). No floating-point evidence is used.
Historical novelty	Specific contributions are relative to the inspected repository and consulted sources; no exhaustive priority certification.

References

- [1] Susan Landau and Gary L. Miller, *Solvability by radicals is in polynomial time*, Journal of Computer and System Sciences **30**(2) (1985), 179–208. DOI: [10.1016/0022-0000\(85\)90013-3](https://doi.org/10.1016/0022-0000(85)90013-3). Author’s bibliographic record: <https://www.cs.cmu.edu/~glmiller/Publications/b2hd-LaMi85.html>.
- [2] C. Boswell and M. L. Glasser, *Solvable sextic equations*, 2005 preprint, arXiv:math-ph/0504001. <https://arxiv.org/abs/math-ph/0504001>. The two imprimitive subgroup types are described on page 3.
- [3] Claus Fieker and Jürgen Klüners, *Computation of Galois groups of rational polynomials*, LMS Journal of Computation and Mathematics **17** (2014), 141–158. DOI: [10.1112/S1461157013000302](https://doi.org/10.1112/S1461157013000302). <https://arxiv.org/abs/1211.3588>.
- [4] Vladimir Reshetnikov, *ProveIt: Polynomial formulas and the Abel–Ruffini obstruction*, repository documentation, snapshot e2b1f016a94102663f12b970e6dd434229f90d01, inspected September 28, 2026. <https://github.com/VladimirReshetnikov/ProveIt/blob/e2b1f016a94102663f12b970e6dd434229f90d01/Algebra/PolynomialFormulas/README.md>.
- [5] Vladimir Reshetnikov, *SexticRadicalDecision.lean*, same snapshot. <https://github.com/VladimirReshetnikov/ProveIt/blob/e2b1f016a94102663f12b970e6dd434229f90d01/Algebra/PolynomialFormulas/Lean/PolynomialFormulas/SexticRadicalDecision.lean>.
- [6] Vladimir Reshetnikov, *SexticIrreducibleDecision.lean*, same snapshot. <https://github.com/VladimirReshetnikov/ProveIt/blob/e2b1f016a94102663f12b970e6dd434229f90d01/Algebra/PolynomialFormulas/Lean/PolynomialFormulas/SexticIrreducibleDecision.lean>.
- [7] Vladimir Reshetnikov, *SexticSeparatingInvariants.lean*, same snapshot. <https://github.com/VladimirReshetnikov/ProveIt/blob/e2b1f016a94102663f12b970e6dd434229f90d01/Algebra/PolynomialFormulas/Lean/PolynomialFormulas/SexticSeparatingInvariants.lean>.
- [8] Vladimir Reshetnikov, *SexticRadicalSemantics.lean*, same snapshot. <https://github.com/VladimirReshetnikov/ProveIt/blob/e2b1f016a94102663f12b970e6dd434229f90d01/Algebra/PolynomialFormulas/Lean/PolynomialFormulas/SexticRadicalSemantics.lean>.
- [9] Vladimir Reshetnikov, *SexticRadicalComputability.lean*, same snapshot. <https://github.com/VladimirReshetnikov/ProveIt/blob/e2b1f016a94102663f12b970e6dd434229f90d01/Algebra/PolynomialFormulas/Lean/PolynomialFormulas/SexticRadicalComputability.lean>.
- [10] Vladimir Reshetnikov, *SexticReducibleDecision.lean*, same snapshot. <https://github.com/VladimirReshetnikov/ProveIt/blob/e2b1f016a94102663f12b970e6dd434229f90d01/Algebra/PolynomialFormulas/Lean/PolynomialFormulas/SexticReducibleDecision.lean>.
- [11] Vladimir Reshetnikov, *SexticSeparatingSearch.lean*, same snapshot [write; not cited by the manuscript]. <https://github.com/VladimirReshetnikov/ProveIt/blob/e2b1f016a94102663f12b970e6dd434229f90d01/Algebra/PolynomialFormulas/Lean/PolynomialFormulas/SexticSeparatingSearch.lean>.
- [12] Vladimir Reshetnikov, *SexticSparseSymmetricSearch.lean*, same snapshot [write; not cited by the manuscript]. <https://github.com/VladimirReshetnikov/ProveIt/blob/e2b1f016a94102663f12b970e6dd434229f90d01/Algebra/PolynomialFormulas/Lean/PolynomialFormulas/SexticSparseSymmetricSearch.lean>.

- [13] *Specialization-Safe Radical Solvers: Exact Bad Characteristics for Subset-Sum Resolvents, a Fourier–Kummer Atlas, and a Global Obstruction*, ProveIt research-report collection, batch 36 [write; sibling report, not cited by the manuscript]. `SetTheory/Cardinals/docs/reports/galois-theory-and-radicals/specialization-safe-radical-solvers/`.