Polynomial Centering of the Inverse Pell Trace

Ten positive unknowns: 89 operations/degree 210 and 90/206

Research note, 26 October 2026; prepared with AI assistance

1 Result and the late comparison

The complete literal polynomials here use ten independently strictly positive integer unknowns, with operation/degree pairs

89/210	and	90/206.

All binary additions, subtractions, scalar multiplications, and squares are charged; leaves are integers. The bounded late source172 at ProveIt `09764940c289c493aa573530eda86dd75f319787` supplies87/214. The new pairs exchange two or three gates for four or eight degrees. They are not simultaneous improvements or global minimum claims.

The earlier scaled-chart87/228 and88/224 artifacts remain supplied as verified method controls, but source172 strictly dominates them in operation/degree. They are no longer headline frontier points. Source172's free quotient $Y = s$, scale $\lambda = q$, and index-three-compatible transport proof are credited to that source and its retained predecessors. It does not already perform the polynomial centering used here.

The whole historical radix67 tuple remains little-endian. Each new front also has a separately calibrated whole Start-marker4177 specialization: 98,219 raw bytes, largest port196,790 bits, minimal unsigned big-endian. This is the pinned tuple at the earlier5274bc751 snapshot, not a global storage record: the bounded late source174 supplies a distinct4127 encoding at97,046 bytes. No ports or controller certificates are mixed.

The result is an ordinary proof composition at the explicit retained native/refutation boundary. It is not a consistency decision, a claimed zero, a new formal-kernel object, or an encoded internal-ZFC replay.

2 The complete free-chart equations

The ten independent inputs are

$$J, F, \alpha, i, T, w, s, n, \eta, \zeta \in \mathbb{Z}_{>0}.$$

For the selected entire fixed tuple β, K_0, M_C, M_F, define

$$\begin{gathered}
h = \beta J, \quad q = h + 1, \quad C = q - F - \alpha, \\
X = qw, \quad Y = s, \quad \lambda = q, \quad k = \eta + \zeta, \quad c = kY + \eta, \\
A = qY(X + 1) + 1, \quad \Delta = A^2 - 1, \quad H = 4A - 5, \\
v = Xq^2Y^2, \quad \ell = vk, \quad E = 1 + \ell(\ell + k), \\
R = [h(q - F) + \alpha](q^2 - 1) + (M_C + qM_F)J, \\
B = (K_0 + w)C - F, \quad U = ic^2, \quad a = \Delta U^2 E, \\
F_0 = 1 + a, \quad G = EF_0, \quad f = a + F_0, \quad Q = 4\Delta a F_0, \\
b = cT - R, \quad V = fb - c, \quad K = X(3c - 2X) + 2, \\
e_6 = 6(k - R - 1) - \ell(k^2 - 4), \quad D = 6q^7, \quad M = hDH.
\end{gathered}$$

Choose one polynomial center,

$$\widehat{g} = AUE \quad (90/206), \qquad \widehat{g} = (A + 1)UE \quad (89/210).$$

Then set

$$Z = hD\{H[n - V(bB + \widehat{g})] + K\} - H[DB + h(e_6 - 6q^4 s)], \tag{1}$$

$$P = Q(Z^2 - M^2 GV^2) + (M^2 G - Z^2). \tag{2}$$

The equation is $P = 0$. All named quantities after the ten inputs are polynomial abbreviations, not further witnesses. There are no rational leaves, divisions, recurrences, binomials, or variable-exponent gates.

For proof only let $L = bV$, $W = 1 + hL$, $t = B/h$, $\kappa = K/H$, $e^* = e_6/6$, $\rho = e^*/q^7$, $\sigma = Y/q^3$, and $z = Z/M$. Before asserting integrality one has the exact rational identity

$$z = n + \kappa + \sigma - Wt - \rho - \widehat{g}V. \tag{3}$$

If Z_0 is the uncentered source172 numerator, then $Z = Z_0 - M\widehat{g}V$ identically.

3 Soundness, including the auxiliary index-three branch

The complete source172 proof is retained in `late172/parent/`. Its root extraction and signed all-index folding use the final norm, not the specific carrier numerator. At a zero they give positive integral τ, f_0, g, y and $\epsilon = \pm 1$ with

$$E = \tau^2, \quad F_0 = f_0^2, \quad g = \tau f_0, \quad z = \epsilon g y, \quad QV^2 - (Q-1)y^2 = 1,$$
$$f^2 - \Delta I^2 = 1, \quad I = 2Ug, \quad c^2 \mid I, \quad c = \psi_A(p), \quad p \geq 1 \text{ odd}.$$

Since $c \geq 3$, $p \geq 3$ and $c \geq c_3 = 4\Delta + 3 > 4q^4$, hence $K > 0$. This works even at $Y = 1$: no assumption $q \mid Y$, exclusion of $p = 3$ using that assumption, or exclusion of auxiliary index3 is imported.

The source's trace-linear argument gives nonzero b, V of the same sign, and

$$L = |b||V| > 0, \qquad y < \tfrac{3}{2}|V|, \qquad f > 6g, \qquad |b| > 4\Delta f - 1.$$

Although the source displayed the weaker consequence $|b| > 4g$, its actual preceding bound gives, using $\Delta \geq 3$ and integral $g \geq 1$,

$$|b| > 24\Delta g - 1 \geq 72g - 1 > 8g. \tag{4}$$

For $g_0 = AUE$, the identity $g_0^2 - G = U^2E^2 - E$ gives $0 < g_0 < 2g$. For $g_+ = (A+1)UE$,

$$g_+^2/G < (A+1)^2/\Delta = (A+1)/(A-1) < 2,$$

so also $0 < g_+ < 2g$. Therefore either center satisfies

$$gy + \widehat{g}|V| < \tfrac{7}{2}g|V| < \tfrac{7}{16}L < L/2. \tag{5}$$

Since $k \geq 2$, $e^* \leq k - R - 1 < b$, hence $\rho < |b|/q^7$. If $B \leq -1$, then $Wt < -L$. Positive κ, σ and (3) give

$$n < L/2 - L + |b|/q^7 < 0,$$

a contradiction. Thus $B \geq 0, C > 0$, before assuming any sign of R. The whole-tuple mask bound now gives $3q < R < q^4 - 2$. As $c > 2R$, $b, V > 0$. The unchanged normalized-rank and signed auxiliary-index theorem yields $p = R$, followed by R odd, q even, and J odd.

4 Authentication, projection, and positive completeness

Because $Z - Z_0$ is a multiple of $M = hDH$, every numerator reduction modulo D and H in source172 remains unchanged. First $k = 2\psi_{1+2v}(m)$ with $m \geq 2$, and literal cancellation of6 gives $q^4 \mid e^*$. The full recurrence and cubic-difference identity

$$3[e^* - (2m - R - 1)] = 4v^2(24Z_m - A_m S_m)$$

with $v = q^3 w s^2$ give $q^4 \mid [e^* - (2m - R - 1)]$ at every prime, including3. Monotonicity gives $m < R$, hence $m = (R+1)/2$. No q^7 divisibility is inferred yet.

Reduction modulo H gives $H \mid 6hK$. The ratio theorem is applied in its actual uniform family $\lambda = q, Y \geq 1$, giving $Y > X^r/3$ for $R = 2r + 1, r \geq 48$. The universal projector and whole-product height bound then yield $X = 2^R$ without canceling an unproved factor of6h. Now q is a power of2; rank growth gives $q^6 \mid X$, hence $q^8 \mid v$. Only at this point does the cubic identity give $q^7 \mid e^*$. Thus ρ, κ are integral. Since $\widehat{g}V$ is integral on every input, the remaining carrier yields $q^3 \mid Y$ and $h \mid B$, so $t \geq 0$ is integral. The retained fixed-minus parity, uniform half-binomial rounding, and fresh native reconstruction at $Y(X + 1) + 2$ then give the actual native words and a finite syntactic ZFC refutation.

Conversely the separately calibrated parent construction supplies a positive zero with the free chart $Y = s, \lambda = q$ and $V > 0$. Keep all inputs except n and define

$$n_{\text{new}} = n_{\text{old}} + \widehat{g}V > 0.$$

The center is independent of n, and exactly

$$n_{\text{new}} - V(bB + \widehat{g}) = n_{\text{old}} - bVB.$$

Thus the numerator and final polynomial value are preserved. No new cofinal seed or Pell-branch choice is needed. This is a positive map on constructed parent zeros; soundness is independent because an arbitrary inverse shift need not stay positive.

5 Literal costs and all highest-degree terms

The source uses87 gates. Replacing its three gates for $L = bV$, LB, $n - LB$ by four for bB, $bB + \widehat{g}$, $V(bB + \widehat{g})$ and $n - V(bB + \widehat{g})$ adds one. For AUE, computing $A = (A - 1) + 1$ and $A(UE)$ adds two more, giving90. For $(A + 1)UE$, $A + 1$ is already live, so only one more gate is needed, giving89.

The exact cancellation is

$$Z^2 - M^2GV^2 = Z_0^2 - 2M\widehat{g}VZ_0 + M^2(\widehat{g}^2 - G)V^2,$$

where

$$(AUE)^2 - G = U^2E^2 - E, \qquad ((A + 1)UE)^2 - G = 2(A + 1)U^2E^2 - E.$$

The actual free-chart degrees are

$$(\deg Q, \deg M, \deg G, \deg V, \deg Z_0, \deg U, \deg E, \deg \widehat{g}) = (72, 12, 46, 36, 54, 5, 14, 23).$$

For the first center, the three inside terms have degrees108,125,134, so multiplication by Q gives206. For the second, the last is138, giving210. The final correction has degree142 in both cases.

Put $q_0 = \beta J$, $k_0 = \eta + \zeta$. The complete leading forms are

$$[P_A]_{206} = 9 \cdot 2^{10} q_0^{82} w^{24} s^{56} i^{10} k_0^{32} (q_0 - F)^2,$$
$$[P_{A+1}]_{210} = 9 \cdot 2^{11} q_0^{84} w^{25} s^{57} i^{10} k_0^{32} (q_0 - F)^2.$$

Each has99 terms and is nonzero for actual $\beta > 0$. Here the leading term of b is $-q_0^3(q_0 - F)$; the scaled chart's T-dependent tie must not be copied. Independent exact cuts cover every actual gate, prove the cancellation, and check all99 terms. All180 or178 alternative gate-operation corruptions are rejected for each complete tuple.

6 Whole-tuple calibration, recovery, and checks

The source172 closure was reconstructed at the bounded late pin using59 exact PLACEMENT aliases; all67 inherited dependency pins match. Its builder and full508-mutation deep checker pass. Normal/optimized checks for the four new graphs also agree. The historical tuple and its full finite recipe remain unchanged; the entire4177 tuple, actual physical support and Pcs are separately reconstructed.

For4177, there are980 physical positions but977 nonzero weights. Marker1048 has zero weight and remains physical; the separate kernel monomial at1048 copies Start selector0. Every one of the3297 coefficient slots and all four integers are checked, with $M_F = M_{F,\text{native}} + \beta$ exactly once. The4177-specific Pcs uses primes4177/224113, orders23/87, spacing2001, and nonzero H residues9/2620/158694 modulo47/4177/224113.

The free-chart endpoint is exactly $X = 2^R, q^3 \mid Y, C > 0$ and $(K_0 + w)C - F = (q-1)t$ with integral $t \geq 0$. It reconstructs fresh source witnesses at $Y(X+1)+2$, decoded by the4177 native proof. Conversely that proof provides fresh canonical outer data to which the source172 $\lambda = q$ construction and our positive centering map apply. Neither old words nor old controller certificates are substituted.

The frozen4177 gzip blobs are unchanged. The inherited page builder has the explicitly recorded runtime OS-header03/ff difference; raw integer reconstruction, stored hashes, new graph bytechecks and Pcs pass. We do not claim that the runtime-sensitive inherited compressed bytecheck passed.

At each whole retained calibration, ZFC proves absence of a strictly positive zero equivalent to Con(ZFC). Finite checks are separate from the unbounded Pell and native theorems. This is an ordinary composed result, not a new end-to-end formalization or a consistency proof. Full proofs, original attribution, licenses and finite recipes accompany every numerical specialization.

A Complete historical coefficient recipe

In this appendix K denotes the port K_0 in the new equations. Every symbol introduced here is a fixed constant, independent of the ten quantified variables. The formulas and 436-character seed are the retained indexed BFS943 radix67 specification; they reconstruct the exact supplied payloads, with $\beta = 2^{590069} - 1$.

The four fixed integers. Every index, set, sum, product and floor below defines a constant, not another unknown. Put

$$\rho = 2^{67}, \quad B = \rho^{8807}, \quad \beta = B - 1, \quad G_m(x) = \sum_{j=0}^{m-1} x^j.$$

Define

$$\begin{aligned}\mathcal{E} = {} & \{0, \dots, 942\} \cup \{983, \dots, 992\} \cup \{1003, \dots, 1011\} \\ & \cup \{1013, \dots, 1031\} \cup \{1063, 1064, 1095, 2195, 3296\}.\end{aligned}$$

Then

$$\begin{aligned}M_C &= B - 1 - \sum_{e \in \mathcal{E} \setminus \{1\}} \rho^e - 2\rho^{1063}, \\ M_F &= B - 1 + 4 + (2^{51} - 2)(\rho^{4393} + \rho^{5495}) \\ &\quad + \rho^{1013} G_{10}(\rho) + \rho^{1065} G_9(\rho) + \rho^{2195} + \rho^{3296} \\ &\quad + (2^{65} - 1)\rho^{8792} G_{13}(\rho) + (2^{19} - 1)\rho^{8805}, \\ K &= \rho^{30} + \rho^{62} + \rho^{1100} + \rho^{2201} + \rho^{8849} \\ &\quad + (\rho^{4393} + \rho^{5495})\Phi(1/\rho),\end{aligned}$$

where

$$\begin{aligned}\Phi(x) &= (1 + 2^{51}) G_{943}(x) - 2^{10} \sum_{i=0}^{942} v(t_i) x^i \\ &\quad + \sum_{(b,e,m) \in \mathcal{C}} 2^b x^e G_m(2x), \\ \mathcal{C} = {} & \{(10, 1023, 8), (18, 1013, 9), (27, 1003, 8), \\ & (35, 983, 9), (44, 1031, 1), (45, 1022, 1), (46, 1011, 1), \\ & (47, 992, 1), (48, 1095, 1), (49, 2195, 1), (50, 3296, 1)\}.\end{aligned}$$

The rational intermediate in K cancels exactly; all four results are positive integers.

The 943 indexed fixed tiles. Use the actions u_j, v_j, d_j decoded below. Let

$$\begin{gathered}h(q, b) = 2(q+1) + b, \quad \mathcal{R} = \{u_j : d_j = 1,\ 0 \leq j < 430\}, \\ \mathcal{L} = \{u_j : d_j = 0,\ 0 \leq j < 430\}, \\ r_q = 6 + |\mathcal{R} \cap \{0, \dots, q-1\}|, \quad l_q = 205 + |\mathcal{L} \cap \{0, \dots, q-1\}|.\end{gathered}$$

An interval ending at -1 is empty. The table defines t_i for $0 \le i \le 942$; all index ranges are fixed:

index i	tile $t_i = (w, n, e, s)$
$0, 1$	$(2, 435, 3, 1),\ (5, 0, 0, 0)$
$j + 2\ (j = 2q + b < 430)$	$((1 - d_j) l_{u_j}, h(q, b), d_j r_{u_j}, v_j)$
$432 + b\ (b = 0, 1)$	$(0, h(215, b), 0, h(215, b))$
$434 + b\ (b = 0, 1)$	$(0, b, 0, b)$
$424 + 2r_q + b\ (q \in \mathcal{R}, b = 0, 1)$	$(r_q, b, 0, h(q, b))$
$424 + 2l_q + b\ (q \in \mathcal{L}, b = 0, 1)$	$(0, b, l_q, h(q, b))$
$924 + j\ (0 \le j < 4)$	$(0, j + 430 \lfloor j/2 \rfloor, 0, 438)$
928	$(2, 434, 2, 0)$
929	$(3, 436, 4, 4)$
930	$(4, 436, 4, 0)$
$497 + t\ (t = 434, 435, 436)$	$(1, 438, 1, t)$
$934+3a+b-6\lfloor a/4 \rfloor$ $a \in \{0,1,4\},\ b=0,1,2$	$(a, 437, b, 437)$

For $t = (w, n, e, s)$ put

$$v(t) = w + 2^8 n + 2^{17} e + 2^{25} s + 127\, 2^{34}.$$

A closed decoder for the 430 actions. Put $D_b(A, k) = \lfloor A/b^k \rfloor \bmod b$, where $a \bmod b = a - b \lfloor a/b \rfloor$. For $0 \le q < 215$, $b = 0, 1$, $j = 2q + b$:

$$N = \mathcal{A} \bmod 2^{636}, \quad T_0 = \lfloor \mathcal{A}/2^{636} \rfloor \bmod 2^{1675},$$
$$E_0 = \lfloor \mathcal{A}/2^{2311} \rfloor,$$
$$\nu_j = D_2(N, j), \quad \sigma_j = \sum_{k=0}^{j-1} \nu_k,$$
$$m_q = D_2(N, 430 + q), \quad \tau_q = \sum_{k=0}^{q-1} m_k,$$
$$F_q = 14(1 - m_q) + m_q D_{16}(E_0, \tau_q),$$
$$u_j = \nu_j(1 + \sigma_j) + (1 - \nu_j) D_{216}(T_0, j - \sigma_j),$$
$$v_j = D_2(F_q, 2b), \quad d_j = D_2(F_q, 2b + 1).$$

Empty sums are zero. Here $(v_0, d_0, u_0) = (1, 1, 1)$; the constructor tile uses this actual first target, including its state renaming.

The integer $\mathcal{A}$. The alphabet, in order of digit values $0, \ldots, 61$, is

```
0123456789ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz.
```

Concatenate the following 436 characters as digits $e_0, \ldots, e_{435}$; put $\mathcal{A} = \sum_{j=0}^{435} e_j 62^{435-j}$.

```
BxEnhvy6NipzkWmROo0GsA3uK3twWLxscLq20UTumy5Cdu461LwD6g7RDNzF8Y
iYDFTpB8ehAypd7d6OsOL4APamnJeNXQypUezCmW9n4oXP6XfgmiWuAWZ8ZJ7U
OP7gu4kVdlGZqZHULQMar1dXNeJW4i7oNVI8usp8BlovjjZbwPCovZI328fRUq
4q92R81s5GjWQaQfGhkq9UYMK3rRMaBmsOIBkJqc2cj2VRIZ6mjzkBIOkW9SjZ
FzeA6J5u4j3sBL3SdgwXzM9mHvQivu8EJ1TBTpRhHDqZIbmYgwQ8VqTAFotl6F
We1YC554GKrVHsA8oUGKSA6F6k5KREBSDOTgd7bFALY84LLbpw574BQyvdoHrl
OOe35uwDf15DpT6VJWZBjSbHPpPAINYTteOeUxnTFTKBlGj8Am6sb1gLldh8WZ
8P
```

B Complete Start-marker4177 coefficient recipe

Here K denotes the fixed port K_0, not an unknown. These are the whole coherent big-endian Start-marker4177 ports, not the historical radix67 tuple. **The four fixed integers.** Every index, set, sum, product and floor below defines a constant, not another unknown. Put

$$\rho = 2^{47}, \quad B = \rho^{4177}, \quad \beta = B - 1,$$
$$G_m(x) = \sum_{j=0}^{m-1} x^j, \quad \Lambda = \rho^{2059} + \rho^{3108}.$$

$$\begin{aligned} M_C &= \beta + \rho - (\rho+3)\rho^{990} \\ &\quad - G_{943}(\rho) - \Psi(\rho, 1), \\ M_F &= B + 3 + (2^{44} - 2)\Lambda \\ &\quad + \rho^{973} G_9(\rho) + \rho^{1012} G_8(\rho) + \rho^{1048} \\ &\quad + 2^{29}\rho^{4175} - \rho^{4157}, \\ K &= \rho^{20} + \rho^{30} + \rho^{1048} + \rho^{4187} \\ &\quad + \Lambda\Phi(1/\rho), \end{aligned}$$

where

$$\begin{aligned} \Phi(x) &= \sum_{i=0}^{942} [1 + 2^{10}(1 + w_i + 2^8 n_i \\ &\quad + 2^{17} p_i + 2^{25} s_i)] x^i + 2^{10}\Psi(x, 2), \\ \Psi(x, t) &= x^{992}(x^{10} + t^{17}) G_8(tx) \\ &\quad + t^8 x^{943}(x^{30} + t^{17}) G_9(tx) \\ &\quad + t^{34}(2 - t) x^{1048}. \end{aligned}$$

The rational intermediate in K cancels exactly; all four results are positive integers.

The 943 indexed fixed tiles. Use the actions u_j, v_j, d_j decoded below. For $d = 0, 1$, let

$$h(q, b) = 2q + b + 2, \quad \mathcal{S}_d = \{u_j : d_j = d,\ 0 \le j < 430\},$$
$$r_{d,q} = 205 - 199d + |\mathcal{S}_d \cap \{0, \dots, q-1\}|.$$

An interval ending at -1 is empty. The table defines t_i for $0 \le i \le 942$; all index ranges are fixed:

index i	tile $t_i = (w_i, n_i, p_i, s_i)$
$0, 1$	$(2, 435, 3, 1),\ (5, 0, 0, 0)$
$j + 2\ (j = 2q + b < 430)$	$((1 - d_j) r_{0,u_j}, h(q, b), d_j r_{1,u_j}, v_j)$
$434 - 2a + b\ (a, b = 0, 1)$	$(0, 432a + b, 0, 432a + b)$
$\substack{424 + 2r_{d,q} + b \\ q \in \mathcal{S}_d,\ d,b = 0,1}$	$(dr_{d,q}, b, (1 - d) r_{d,q}, h(q, b))$
$924 + 2a + b\ (a, b = 0, 1)$	$(0, 432a + b, 0, 438)$
928	$(2, 434, 2, 0)$
929	$(3, 436, 4, 4)$
930	$(4, 436, 4, 0)$
$497 + t\ (t = 434, 435, 436)$	$(1, 438, 1, t)$
$934 + 3a + b\ (a, b = 0, 1, 2)$	$(a^2, 437, b, 437)$

A closed decoder for the 430 actions. Put $D_b(A, k) = \lfloor A/b^k \rfloor \bmod b$, where $a \bmod b = a - b\lfloor a/b \rfloor$. For

$0 \le q < 215$, $b = 0, 1$, $j = 2q + b$:

$$N = \mathcal{A} \bmod 2^{636}, \quad T_0 = \left\lfloor \mathcal{A}/2^{636} \right\rfloor \bmod 2^{1675},$$
$$E_0 = \left\lfloor \mathcal{A}/2^{2311} \right\rfloor,$$
$$\nu_j = D_2(N, j), \quad \sigma_j = \sum_{k=0}^{j-1} \nu_k,$$
$$m_q = D_2(N, 430 + q), \quad \tau_q = \sum_{k=0}^{q-1} m_k,$$
$$F_q = 14 + m_q(D_{16}(E_0, \tau_q) - 14),$$
$$u_j = \nu_j(1 + \sigma_j) + (1 - \nu_j)D_{216}(T_0, j - \sigma_j),$$
$$v_j = D_2(F_q, 2b), \quad d_j = D_2(F_q, 2b + 1).$$

Empty sums are zero. Here $(v_0, d_0, u_0) = (1, 1, 1)$; the constructor tile uses this actual first target, including its state renaming.

The integer $\mathcal{A}$. The alphabet, in order of digit values $0, \ldots, 61$, is:

```
0123456789ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz.
```

Concatenate the following 436 characters as digits $e_0, \ldots, e_{435}$; put $\mathcal{A} = \sum_{j=0}^{435} e_j 62^{435-j}$.

```
BxEnhvy6NipzkWmROo0GsA3uK3twWLxscLq2OUTumy5Cdu461LwD6g7RDNzF8Y
iYDFTpB8ehAypd7d6OsOL4APamnJeNXQypUezCmW9n4oXP6XfgmiWuAWZ8ZJ7U
OP7gu4kVdlGZqZHULQMar1dXNeJW4i7oNVI8usp8BlovjjZbwPCovZI328fRUq
4q92R81s5GjWQaQfGhkq9UYMK3rRMaBmsOIBkJqc2cj2VRIZ6mjzkBIOkW9SjZ
FzeA6J5u4j3sBL3SdgwXzM9mHvQivu8EJ1TBTpRhHDqZIbmYgwQ8VqTAFotl6F
We1YC554GKrVHsA8oUGKSA6F6k5KREBSD0Tgd7bFALY84LLbpw574BQyvdoHrl
0Oe35uwDf15DpT6VJWZBjSbHPpPAINYTte0eUxnTFTKBlGj8Am6sb1gLldh8WZ
8P
```