

Holonomic Rigidity for Entire Hahn Functions

A sharp order-unit criterion for dilations,
with uniform exclusion bounds,
and the exact boundary of nonlinear differential rigidity

Merged research report · 22–23 September 2026

Abstract

Fix a nonzero set-sized ordered abelian group Γ and the complex Hahn field $K_\Gamma = \mathbb{C}((t^\Gamma))$. *Entire* means strong Hahn summability of the evaluation family at every point of this one field. We prove that every strongly entire D-finite power series is a polynomial, in one or in finitely many variables, and that a fixed differential operator already excludes every nonterminating formal solution on an explicit exterior valuation region, with a common degree bound.

For dilations there is an exact alternative. If $q \in K_\Gamma^\times$ has infinite multiplicative order, a nonpolynomial strongly entire solution of a nonzero linear q -difference equation with polynomial coefficients exists if and only if $v(q) \neq 0$ and $|v(q)|$ is an order unit of Γ . A nonzero valuation is not enough: its integer multiples must be cofinal among all scales. The proof covers units whose residues are roots of unity, and coefficients and parameters with arbitrary well-ordered Hahn supports. A partial theta series proves sharpness, and its exact strong evaluation domain is computed for an arbitrary Hahn parameter. Quantitatively, the recurrence supplies an exclusion bound that is attained on its own boundary, and a coarsened exclusion bound valid for every noncofinal dilation. A mixed differential–dilation extension is proved for nonzero noncofinal dilation valuations.

The common mechanism is an elementary finite-step escape chain: bounded valuation costs in a coefficient recurrence force a subsequence of evaluated leading exponents that never increases. This contradicts strong summability without replacing an arbitrary-rank valuation by a real-valued norm. In an explicit surreal workspace with countable cofinality and no order unit, the series $\sum_{n \geq 0} \omega^{-\omega^n} z^n$ is strongly entire yet satisfies no nonzero linear differential equation and no nonzero linear dilation equation for any nontorsion parameter of that workspace.

Over $k((t^\Gamma))$, k any field of characteristic zero, later sources show: first-order algebraic differential equations have only polynomial strongly entire solutions; without an order unit every nonpolynomial strongly entire series is differentially transcendental, jointly with its dilates by parameters whose quotients are not roots of unity; with one, a Chebyshev-descended theta series has minimal differential order three; for every Γ no such series satisfies an equation of order ≤ 2 , or of order three and jet degree ≤ 3 ; linear mixed rigidity holds at unit dilations; a dominant argument $f(P(z))$, $\deg P \geq 2$, forces polynomiality for every Γ ; and unit-root recurrences have finitely generated coefficient algebras. Divisibility of Γ is *not* assumed, except in three threshold constructions that state it (Section 1.4).

Status. An AI-assisted research report; priority and independent refereeing are not certified. Lean covers support and escape lemmas, exponential and partial-theta domains, polynomial-orbit valuations, constant-point avoidance, inward stability and torsion covariance. The full classifications and the theorem packages of the ten later parts remain pending. The variable derivative is not the Berarducci–Mantova derivation on surreal coefficients.

Contents

1	The question and the principal results	3
2	Hahn support and what evaluation means	16
3	The finite-step escape mechanism	19
4	Valuations on integer and geometric orbits	21
5	Differential rigidity	24
6	Generic lines and several-variable rigidity	27
7	Dilation equations and the necessity of a cofinal scale	29
8	Partial theta series: sharp witness and exact domain	31
9	Torsion parameters and why they are excluded	34
10	An extension to mixed differential–dilation equations	35
11	Surreal workspaces that separate the theories	36
12	Coefficient fields, degenerate value groups, characteristic	39
13	Quantitative certificates and symbolic representation	39
14	Nonlinear equations: finite initial polynomials	41
15	First-order algebraic differential equations	46
16	Higher order and positive-weight Euler equations	51
17	Nonlinear consequences, hypotheses and predecessors	55
18	The thirteen finite check suites	58
19	Originality assessment and questions left open	71
20	All-order rigidity without an order unit: setting and valuations	94
21	The coefficient-field theorem and its consequences	101
22	Explicit families, surreal realizations and boundaries	108

23 Theta descent: an entire series with an equation of order three	116
24 An equation of order three, and none of lower order	123
25 The exact boundary, surreal consequences and scope	131
26 Second-order rigidity: finite endpoint spectra at exterior breaks	137
27 Order three: endpoint gaps, theta breaks and scope	145
28 Unit-dilation rigidity: valuations of Hahn exponential polynomials	150
29 Mixed rigidity and the relative-scale dichotomy	157
30 Order three through cubic jet degree: first polars on the twisted cubic	166
31 Exact classifications beyond the first-polar test, and scope	173
32 Independent theta scales and unbounded differential order	177
33 Continuum fields, integer sampling, a third route to order three, and scope	184
34 Differential freedom at a single scale: exact relation ideals	188
35 Entire, surreal and omnific realizations of single-scale freedom	194
36 Polynomial-composition rigidity: profiles and the weighted theorem	201
37 Omnific coefficients, universality and boundaries of composition rigidity	208
38 Hahn-valued recurrences: observables and Noetherian compression	213
39 Surreal recurrences: leading scales, omnific coding, matrix powers and mixed operators	219
40 Conclusion	227
A Notation	229
B Proof-dependency and hypothesis audit	233
C Reproducing the document and the finite checks	236
References	239

1 The question and the principal results

1.1 Why the question is scale sensitive

A familiar way to specify a special function is to give a finite linear differential equation. Another is to give a finite relation between its values at $z, qz, q^2z, \dots$. Over the complex numbers both descriptions produce nonpolynomial entire functions. Over a surreal Hahn workspace, “entire” asks for evaluation not only at ordinary complex constants but at every infinite and infinitesimal scale available in that workspace. In a Hahn field an infinite expression is legal only when its supports satisfy an order condition and a finiteness condition, and an equation can force either condition to fail at a sufficiently large argument. The question of this report is what a finite linear equation can accommodate under that stronger global requirement. Sections 14–17 ask the same of a finite *nonlinear* algebraic differential equation, and answer it in order one; Sections 20–22 answer it in every order when the value group has no order unit; and Sections 23–25 show that with an order unit the answer changes: an explicit nonpolynomial strongly entire series satisfies an algebraic differential equation of order three. Sections 26–27 show that three is the least such order: for every value group, no nonpolynomial strongly entire series satisfies an equation of order at most two. Sections 28–29 return to linear equations that mix derivatives and dilations, and settle them by the relative scale of the dilations; Sections 30–31 exclude equations of order three and total degree at most three in the derivatives. Sections 32–33 show, for $\Gamma = \mathbb{R}$, that minimal orders are unbounded and that the differentially algebraic strongly entire series generate a field of transcendence degree $2^{\aleph_0}$; Sections 34–35 give, at a single order-unit scale, explicit series with coefficients in $\mathbb{Q}(t^\mu)$ that satisfy no algebraic differential equation at all, with their whole relation ideals. Sections 36–37 replace the dilation by a polynomial argument of degree at least two, and there rigidity holds for every value group, with explicit degree bounds. Sections 38–39 turn from functions to the ordinary linear recurrences with Hahn or surreal values that underlie all these proofs, and read their individual coefficients: tame, in one finitely generated algebra, when the characteristic roots are units, and arbitrary for a first-order omnific recurrence.

The answer depends decisively on which operator is used, and for dilation equations it depends further on whether the valuation of the particular parameter reaches every scale of the value group.

1.2 Conventions

We work with a fixed ordered abelian group Γ and

$$K_\Gamma = \mathbb{C}((t^\Gamma)).$$

An element of K_Γ is a formal sum $a = \sum_\gamma a_\gamma t^\gamma$ with $a_\gamma \in \mathbb{C}$ whose support $\text{supp}(a) = \{\gamma : a_\gamma \neq 0\}$ is a well-ordered set. For $a \neq 0$ put

$$v(a) = \min \text{supp}(a), \quad \text{lc}(a) = a_{v(a)} \in \mathbb{C}^\times, \quad \text{res}(a) = \text{lc}(a) \text{ when } v(a) = 0,$$

and put $v(0) = +\infty$ only where convenient; this symbol is never subtracted. Products and sums are the usual Hahn operations, with

$$v(ab) = v(a) + v(b), \quad v(a + b) \geq \min\{v(a), v(b)\}, \tag{1.1}$$

with equality in the second law when the two valuations differ. The coefficient field $\mathbb{C}$ is trivially valued, so every nonzero ordinary integer and every factorial has valuation zero. Throughout, $\mathbb{N} = \{0, 1, 2, \dots\}$, and $|\lambda|$ means $\max(\lambda, -\lambda)$ in the ordered group.

Convention 1.1 (Standing hypothesis on the value group). Γ is a nonzero set-sized ordered abelian group. Divisibility is *not* part of this standing hypothesis. Every statement below that needs divisibility says so in its own statement; those statements are collected in Section 1.4.

Definition 1.2 (Strong summability, evaluation domain, entireness). A set-indexed family $(b_i)_{i \in I}$ in K_Γ is *strongly Hahn summable* if $\bigcup_i \text{supp}(b_i)$ is well ordered and each $\gamma \in \Gamma$ lies in the supports of only finitely many members. Its sum is then defined coefficientwise, by a finite sum in $\mathbb{C}$ at each exponent. For $f = \sum_{n \geq 0} a_n z^n \in K_\Gamma[[z]]$ let $\text{Dom}_{\text{str}}(f)$ be the set of $x \in K_\Gamma$ for which $(a_n x^n)_{n \geq 0}$ is strongly summable, and let

$$\mathcal{E}_\Gamma = \{f : \text{Dom}_{\text{str}}(f) = K_\Gamma\}.$$

In d variables the family to be tested is $(a_\alpha x^\alpha)_{\alpha \in \mathbb{N}^d}$.

The word *entire* always means Definition 1.2 relative to one fixed workspace. It does not mean that a complex coefficient function is entire, it does not mean convergence in the fine topology inherited from the full surreal class, and it is never a claim about all of $\mathbf{No}[i]$. The definition also concerns the family *before* cancellation between different indices: one cannot first cancel a nonsummable family and then declare it summable, and convergence of a numerical coefficient series does not license infinitely many contributions at one Hahn exponent.

Definition 1.3 (Order unit). A positive $\mu \in \Gamma$ is an *order unit* if for every $\gamma \in \Gamma$ there is $N \in \mathbb{N}$ with $|\gamma| \leq N\mu$; equivalently, if the positive integer multiples of μ are cofinal in Γ . Zero is not an order unit.

Definition 1.4 (The two operator classes). Let D_z be the formal derivative in z , so $D_z(a) = 0$ for every $a \in K_\Gamma$, and for $q \in K_\Gamma^\times$ let $\sigma_q f(z) = f(qz)$. A series $f \in K_\Gamma[[z]]$ is *D-finite over $K_\Gamma(z)$* if

$$\sum_{r=0}^R p_r(z) D_z^r f(z) = 0, \quad p_r \in K_\Gamma[z], \quad (p_0, \dots, p_R) \neq 0, \quad (1.2)$$

and *linearly q-difference finite* if

$$\sum_{\ell=0}^r p_\ell(z) f(q^\ell z) = 0, \quad p_\ell \in K_\Gamma[z], \quad (p_0, \dots, p_r) \neq 0. \quad (1.3)$$

These are formal identities in $K_\Gamma[[z]]$, not assertions about limits of partial sums. Rational coefficients give the same notions after denominators are cleared.

We say “linearly q -difference finite” rather than silently identifying several meanings of “ q -holonomic”. In the sequence literature q is often an indeterminate with coefficients in $\mathbb{Q}(q)$; here q is a *fixed* Hahn element, possibly with infinitely many terms. The differential terminology is the usual characteristic-zero one [1]. *Nontorsion* and “of infinite multiplicative order” are used interchangeably.

1.3 The principal results

Theorem A (Differential rigidity, with a uniform exterior obstruction). *Let Γ be as in Convention 1.1 and let $0 \neq L = \sum_{r,k} c_{r,k} z^k D_z^r$ be a nonzero linear differential operator with coefficients in $K_\Gamma[z]$. There are $B_L \in \Gamma$ with $B_L \geq 0$ and $N_L \in \mathbb{N}$, both determined by the finitely many coefficients of L , such that every formal solution of $Lf = 0$ that is strongly evaluable at even one $x \neq 0$ with $v(x) \leq -B_L$ has degree less than N_L . Consequently*

$$\{f \in \mathcal{E}_\Gamma : f \text{ is } D\text{-finite over } K_\Gamma(z)\} = K_\Gamma[z].$$

Theorem B (Sharp dilation criterion). *Let Γ be as in Convention 1.1 and let $q \in K_\Gamma^\times$ have infinite multiplicative order. The following are equivalent.*

- (i) *There is a nonpolynomial $f \in \mathcal{E}_\Gamma$ that is linearly q -difference finite over $K_\Gamma(z)$.*
- (ii) *$v(q) \neq 0$ and $|v(q)|$ is an order unit of Γ .*

When (ii) fails, every fixed equation (1.3) has a uniform exterior obstruction and a uniform polynomial degree bound, exactly as in Theorem A. When (ii) holds, an explicit partial theta series with parameter q or q^{-1} is a nonpolynomial entire solution.

Theorem C (Several-variable rigidity). *Let Γ be as in Convention 1.1 and let $F \in K_\Gamma[[z_1, \dots, z_d]]$ be strongly Hahn summable at every point of K_Γ^d . If the mixed partial derivatives of F span a finite-dimensional vector space over $K_\Gamma(z_1, \dots, z_d)$, then F is a polynomial.*

The fourth principal theorem comes from the third source manuscript and is proved in Sections 14–15. It allows an arbitrary coefficient field of characteristic zero; see Convention 14.1, which also explains why that generality is not transferred to Theorems A–C.

Theorem D (First-order nonlinear rigidity). *Let Γ be as in Convention 1.1, let k be any field of characteristic zero, and let $\mathbb{K} = k((t^\Gamma))$, with strong evaluation and entireness defined as in Definition 1.2 with k in place of $\mathbb{C}$. For every nonzero $P \in \mathbb{K}[z, Y_0, Y_1]$, every strongly entire formal solution $f \in \mathbb{K}[[z]]$ of*

$$P(z, f, D_z f) = 0$$

is a polynomial. More strongly, if a nonpolynomial formal solution exists, there is $\delta_0 \in \Gamma_{>0}$, depending on the equation and on finitely many coefficients of that solution, such that $t^{-\delta} \notin \text{Dom}_{\text{str}}(f)$ for every $\delta \geq \delta_0$. Consequently, for every nonpolynomial strongly entire f , the series f and $D_z f$ are algebraically independent over $\mathbb{K}(z)$.

Theorem D has no separant, irreducibility or solved-form hypothesis on P . Its exclusion bound depends on the solution, and Theorem 15.10 shows that this cannot be avoided, in contrast with the operator-only bound of Theorem A. By itself it does not say that a nonpolynomial entire function is differentially transcendental in all orders: independence of f and $D_z f$ does not exclude a relation involving $D_z^2 f$. Theorem E below says this when Γ has no order unit, and Theorem H shows that the hypothesis cannot be dropped: with an order unit a nonpolynomial strongly entire series can satisfy a relation of order three. Theorem I below extends Theorem D to order two for every Γ , so that three is the least order; Question 19.1, once re-scoped to order exactly two, is now re-scoped to the classification of the series of minimal order three.

The last three principal theorems come from sources 11 and 12 and are proved in Sections 20–22, over the field $\mathbb{K} = k((t^\Gamma))$ of Convention 14.1 for an arbitrary field k of characteristic zero; $\mathcal{E}_\Gamma(k)$ is its set of strongly entire series, $\text{Frac } \mathcal{E}_\Gamma(k)$ its fraction field inside $\mathbb{K}((z))$, and $\mathcal{J}_{\lambda,j} f = (D_z^j f)(\lambda z)$ (Convention 20.1).

Theorem E (All-order rigidity without an order unit). *Let Γ be as in Convention 1.1, and suppose that Γ has no order unit.*

- (a) (Sources 11 and 12.) *Every nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ is differentially transcendental over $\mathbb{K}(z)$: the series $f, D_z f, D_z^2 f, \dots$ are algebraically independent over $\mathbb{K}(z)$. Equivalently, every strongly entire solution of a nonzero algebraic differential equation $P(z, f, D_z f, \dots, D_z^r f) = 0$ over $\mathbb{K}(z)$, of any finite order and with any residue corner, is a polynomial.*

- (b) (Source 12.) *Let $\Lambda \subseteq \mathbb{K}^\times$ be such that λ/λ' is not a root of unity for distinct $\lambda, \lambda' \in \Lambda$. For every nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ the family $\{\mathcal{J}_{\lambda,j}f : \lambda \in \Lambda, j \geq 0\}$ is algebraically independent over $\mathbb{K}(z)$. In particular, for every $q \in \mathbb{K}^\times$ of infinite multiplicative order, including $v(q) = 0$, the series $(D_z^j f)(q^i z)$, $i \in \mathbb{Z}, j \geq 0$, are jointly algebraically independent over $\mathbb{K}(z)$.*
- (c) (Source 11.) *The elements of $\text{Frac } \mathcal{E}_\Gamma(k)$ that are differentially algebraic over $\mathbb{K}(z)$ are exactly the elements of $\mathbb{K}(z)$.*

For the coefficient-value rank $\text{cvr}(f)$, the dimension of the rational span of the values of the nonzero Taylor coefficients (20.4), there is a test for a single series that needs neither entireness nor a hypothesis on Γ .

Theorem F (Coefficient-value-rank criterion; source 12). *Let Γ be as in Convention 1.1 and $f \in \mathbb{K}[[z]]$. If $\text{cvr}(f)$ is infinite, the family of Theorem E (b) is algebraically independent over $\mathbb{K}(z)$ for every Λ as there.*

Theorem G (Order-unit dichotomy for mixed jets; source 12). *For Γ as in Convention 1.1, the following are equivalent.*

- (i) Γ has an order unit.
- (ii) Some nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ has all its coefficients in a subfield $L \supseteq k$ of $\mathbb{K}$ with $\text{trdeg}_k L < \infty$.
- (iii) For some nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ and some $q \in \mathbb{K}^\times$ of infinite order, the family $\{(D_z^j f)(q^i z) : i \in \mathbb{Z}, j \geq 0\}$ is algebraically dependent over $\mathbb{K}(z)$.
- (iv) (Added by the merge of source 13.) Some nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ is differentially algebraic over $\mathbb{K}(z)$.

If μ is an order unit, the partial theta series Θ_p of (8.1) with $p = t^\mu$ is a witness for (ii) and (iii): its coefficients lie in $k(p)$, and $\Theta_p(z) = 1 + z\Theta_p(pz)$. The series $\mathcal{T}_p$ of Theorem H is a witness for (ii)–(iv), and for (iii) with every q of infinite order.

As proved by source 12, Theorem G concerned the *mixed* conclusion: its witness Θ_p satisfies a dilation equation, and conditions (i)–(iii) do not say that an order unit produces a nonpolynomial strongly entire *differentially* algebraic series. Clause (iv) says this; it is added by the merge from Theorem H below, and its proof is in Section 25.1. Theorem E answers Question 19.1, as it then stood, in every order for groups without an order unit, including the equations with vanishing residue corner, and its part (b) answers the same case of Question 19.4. Both sources prove part (a) by one argument, printed once: a finite relation, linearized along its solution, puts all Taylor coefficients into one finitely generated field (Theorem 21.2); such a field has a value group of finite rational rank, which lies in a proper convex subgroup when there is no order unit, while a nonpolynomial entire series needs coefficient values at cofinally many scales. Part (b) adds dilations through the classical Skolem–Mahler–Lech theorem. Source 11 also proves a coarsening descent for arbitrary coefficient fields (Theorem 21.11), corollaries for rational systems and for Painlevé I in $\text{Frac } \mathcal{E}_\Gamma(k)$, a joint-independence criterion, and, over $\mathbb{C}((t^\Gamma))$, a continuum-sized family of strongly entire series with jointly independent jets, so that the fraction field has differential transcendence degree $2^{\aleph_0}$ (Theorem 22.6). The series $\sum_{n \geq 0} \omega^{-\omega^n} z^n$ satisfies no algebraic differential equation of any order (Theorem 22.1).

The eighth principal theorem comes from source 13 and is proved in Sections 23–25, over $\mathbb{K} = k((t^\Gamma))$ for an arbitrary field k of characteristic zero (Convention 23.1). For $\mu > 0$ in Γ put

$p = t^\mu$, let C_n be the monic Chebyshev polynomials, $C_0 = 2$, $C_1 = z$, $C_{n+1} = zC_n - C_{n-1}$, so that $C_n(u + u^{-1}) = u^n + u^{-n}$, and put

$$\mathcal{T}_p(z) = 1 + \sum_{n \geq 1} p^{n^2} C_n(z). \quad (1.4)$$

Theorem H (Exact boundary of differential algebraicity; source 13). *For Γ as in Convention 1.1 and every field k of characteristic zero, the following are equivalent.*

- (i) Γ has an order unit.
- (ii) Some nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ is differentially algebraic over $\mathbb{K}(z)$.
- (iii) Some nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ satisfies a nonzero algebraic differential equation of order three over $\mathbb{K}(z)$, while $f, D_z f, D_z^2 f$ are algebraically independent over $\mathbb{K}(z)$ and $D_z^3 f$ has degree two over $\mathbb{K}(z)(f, D_z f, D_z^2 f)$.

If μ is an order unit, $\mathcal{T}_p$ is a witness for (iii); it is not D -finite over $\mathbb{K}(z)$. For every $\mu > 0$ the strong domain of $\mathcal{T}_p$ is $\{0\} \cup \{x : v(x) \geq -N\mu \text{ for some } N \in \mathbb{N}\}$, so $\mathcal{T}_p \in \mathcal{E}_\Gamma(k)$ exactly when μ is an order unit.

Theorem H answers Question 19.1 as it stood after the merge of sources 11 and 12: with an order unit a nonpolynomial entire series can satisfy an algebraic differential equation of order at least two, a differentially algebraic element of $\text{Frac } \mathcal{E}_\Gamma(k)$ need not be rational, and the absence of an order unit is necessary for differential algebraic rigidity. With Theorem D, the least order of such an equation over all examples was then two or three, and the question was re-scoped to order two; Theorem I below (source 14) shows that it is three (Section 19). The equation of $\mathcal{T}_p$ is explicit (Theorem 24.1); its residue corner polynomial vanishes identically while its full corner polynomial is a nonzero constant (Proposition 24.2). The direction (ii) $\Rightarrow$ (i) is Theorem E (a), which source 13 proves again; so Theorem E (a) and (c) hold exactly when Γ has no order unit (Corollary 25.2). The series $\mathcal{T}_p$ is a Chebyshev descent of the bilateral theta series $\sum_{n \in \mathbb{Z}} p^{n^2} u^n$ along $z = u + u^{-1}$; the classical triple product gives all its zeros, and in the surreal specialization $p = \omega^{-1}$ none of them is an omnific integer (Corollary 25.3).

The ninth principal theorem comes from source 14 and is proved in Sections 26–27, again over $\mathbb{K} = k((t^\Gamma))$ for an arbitrary field k of characteristic zero (Convention 26.1).

Theorem I (Second-order rigidity and the order-three threshold; source 14). *Let Γ be as in Convention 1.1 and k any field of characteristic zero.*

- (a) Every $f \in \mathcal{E}_\Gamma(k)$ that satisfies a nonzero algebraic differential equation $P(z, f, D_z f, D_z^2 f) = 0$ with $P \in \mathbb{K}(z)[Y_0, Y_1, Y_2]$ is a polynomial. Equivalently, for every nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ the series $f, D_z f$ and $D_z^2 f$ are algebraically independent over $\mathbb{K}(z)$.
- (b) If Γ has an order unit, the least order of a nonzero algebraic differential equation over $\mathbb{K}(z)$ satisfied by a nonpolynomial element of $\mathcal{E}_\Gamma(k)$ is exactly three. If Γ has no order unit, no nonpolynomial element of $\mathcal{E}_\Gamma(k)$ satisfies one of any order.

Part (a) needs no divisibility, rank, algebraic closedness or order-unit hypothesis, and no condition on the residue corner. Its polynomiality conclusion contains those of Theorem D and of Theorem 16.1 in orders at most two, but not their explicit exclusion bounds, and it holds in particular for equations with $\chi_P = 0$, such as (16.1). The new ingredient is the lower bound three; the witness of order three and the no-order-unit direction in part (b) are Theorems H and E (a). The proof keeps the whole

initial polynomial at the infinitely many scales where it is not a monomial, reduces the equation there to one fixed homogeneous polynomial over k , and shows with logarithmic Euler derivatives that both endpoints of every such initial polynomial lie in a finite set, while a nonpolynomial entire series has such scales with arbitrarily large endpoints (Section 26.3). At order three an extra integer, the gap next to an endpoint, survives (Theorem 27.2); for the equation of $\mathcal{T}_p$ it is always one, the initial polynomials of every nonpolynomial entire solution are eventually $bX^m(1+aX)$, and the coefficient values of such a solution are eventually strictly convex (Theorems 27.4 and 27.5). Consequences include rational two-state systems and quotients with a polynomial denominator (Corollaries 26.10 and 26.11).

The tenth principal theorem comes from source 18 and is proved in Sections 28–29, over $\mathbb{K} = k((t^\Gamma))$ for an arbitrary field k of characteristic zero (Convention 28.1). Its operators are the finite sums

$$L = \sum_{\ell=1}^m \sum_{r=0}^R p_{\ell,r}(z) D_z^r \sigma_{\lambda_\ell}, \quad p_{\ell,r} \in \mathbb{K}[z] \text{ not all zero,}$$

for multipliers $\lambda_1, \dots, \lambda_m \in \mathbb{K}^\times$.

Theorem J (Mixed equations: a relative-scale criterion; source 18). *Let Γ be as in Convention 1.1, k any field of characteristic zero, and $\lambda_1, \dots, \lambda_m \in \mathbb{K}^\times$ with no quotient λ_i/λ_j , $i \neq j$, a root of unity.*

- (a) *If no $|v(\lambda_i/\lambda_j)|$ is an order unit of Γ , then for every such L and every $\bar{e} \in \mathbb{N}$ there are $N \in \mathbb{N}$ and $B \in \Gamma_{\geq 0}$ such that every formal solution of $Lf = \mathbf{h}$, with $\mathbf{h} \in \mathbb{K}[z]$ of degree at most $\bar{e}$, that is strongly evaluable at one $x \neq 0$ with $v(x) < -B$ has degree less than N . In particular every strongly entire solution is a polynomial.*
- (b) *Some nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ satisfies $Lf = 0$ for some such L if and only if $|v(\lambda_i/\lambda_j)|$ is an order unit for some i, j ; then two of the multipliers and derivatives of order at most one suffice.*

In particular, for every Γ and every $q \in \mathbb{K}^\times$ of infinite multiplicative order with $v(q) = 0$, every strongly entire solution of a nonzero equation (10.1) with coefficients in $\mathbb{K}[z]$ is a polynomial.

The last assertion answers Question 19.4 positively, for every Γ (Corollary 29.6). Part (a) contains Theorem A, the negative direction of Theorem B and Theorem 10.2, now over every characteristic-zero k (Remark 29.5); part (b) answers the existence half of Question 19.6 for a prescribed finite set of multipliers, but not for prescribed operators or systems. The new ingredient is a valuation form of the Skolem–Mahler–Lech theorem: an exponential polynomial $\sum_j R_j(n)\lambda_j^n$ with arbitrary Hahn coefficients has eventually affine valuations on finitely many arithmetic progressions, and eventually periodic ones at unit valuation (Theorems 28.5 and 28.10); for $v(P(n, q^n))$ this is the estimate that Question 19.4 named (Corollary 28.7).

The eleventh principal theorem comes from source 19 and is proved in Sections 30–31, over $\mathbb{K} = k((t^\Gamma))$ for an arbitrary field k of characteristic zero (Convention 30.1); source 19 also re-derives Theorem I(a) independently, and that proof is printed once, in Section 26.

Theorem K (Order three through cubic jet degree; source 19). *Let Γ be as in Convention 1.1 and k any field of characteristic zero. Every $f \in \mathcal{E}_\Gamma(k)$ satisfying $P(z, f, D_z f, D_z^2 f, D_z^3 f) = 0$ for a nonzero $P \in \mathbb{K}(z)[Y_0, Y_1, Y_2, Y_3]$ of total degree at most three in $Y_0, \dots, Y_3$ is a polynomial. More generally, the same holds for every equation of order at most three whose exterior form, or each of whose irreducible factors over k , either takes a nonzero value on the monomial jet curve $(1, T, T^2, T^3)$ or has a nonzero first polar along it.*

The degree in z is unrestricted. At order three a nonmonomial initial polynomial at a far break must satisfy the two endpoint conditions of Theorem 27.2; when the first polar is nonzero these say that the exponent next to each end lies on a fixed side of it, which fails at one of the two ends far out (Proposition 30.6), and a secant argument shows that a nonzero cubic form vanishing on the twisted cubic has a nonzero first polar (Lemma 30.7). Beyond degree three the test can fail: the quartic H_3 of order three and the quadratic Q_4 of order four have exactly the monomials and the doubling binomials $az^n + bz^{2n}$ as formal solutions (Theorems 31.1 and 31.2), and a sparse strongly entire series satisfies both exterior equations at every scale without satisfying either equation (Example 31.3). With an order unit, $\mathcal{T}_p$ therefore satisfies no equation of order three and jet degree below four (Corollary 31.4). Consequences include external surcomplex and Gaussian-omnific constants and algebraic two-state systems (Corollary 30.12, Theorem 30.16).

The twelfth principal theorem comes from source 16 and is proved in Sections 32–33, over $\mathbb{K} = k((t^{\mathbb{R}}))$ with $k = \mathbb{R}$ or $\mathbb{C}$ (Convention 32.1). For a real $\mu > 0$, $\mathcal{T}_{t^\mu}$ is (1.4) with $p = t^\mu$, and $\text{dord}(f)$ is the least r with $f, D_z f, \dots, D_z^r f$ algebraically dependent over $\mathbb{K}(z)$.

Theorem L (Independent theta scales and unbounded differential order; source 16). *Let $\Gamma = \mathbb{R}$ and $k = \mathbb{R}$ or $\mathbb{C}$.*

- (a) *If $\mu_1, \dots, \mu_N > 0$ and $\mu_1^{-1}, \dots, \mu_N^{-1}$ are linearly independent over $\mathbb{Q}$, then $\mathcal{T}_{t^{\mu_1}}, \dots, \mathcal{T}_{t^{\mu_N}}$ are algebraically independent over $\mathbb{K}(z)$; there is such a family of cardinality $2^{\aleph_0}$, each member of minimal order three.*
- (b) *For every $N \geq 1$ and $\mu_i = 1/\sqrt{p_i}$ with distinct primes p_i , some $g = \sum_{i=1}^N (\sum_{j < N} c_{ij} z^j) \mathcal{T}_{t^{\mu_i}}$ with $c_{ij} \in \{0, 1, \dots, N\}$, one of at most $(N+1)^{N^2}$ candidates, is strongly entire, differentially algebraic over $\mathbb{K}(z)$, and satisfies $N \leq \text{dord}(g) \leq 3N$.*
- (c) *The differential subfield $\mathcal{D}_{\text{DA}}$ of $\text{Frac } \mathcal{E}_\Gamma(k)$ generated over $\mathbb{K}(z)$ by all differentially algebraic strongly entire series has $\text{trdeg}_{\mathbb{K}(z)} \mathcal{D}_{\text{DA}} = 2^{\aleph_0}$ and $\text{dtrdeg}_{\mathbb{K}(z)} \mathcal{D}_{\text{DA}} = 0$; it is not finitely generated as a differential field.*

Part (a) rests on the exact Gauss profile $\gamma_\delta(\mathcal{T}_{t^\mu}) = -\delta^2/(4\mu) + O(1)$ (Theorem 32.3): distinct monomials of a relation have distinct quadratic growth. Part (b) combines it with a lemma of pure differential algebra, bounded polynomial jet compression (Theorem 32.9). With Theorem I the minimal orders of nonpolynomial differentially algebraic strongly entire series over $\mathbb{K}$ form an unbounded set of integers at least three containing three (Remark 32.18). Source 16 also gives a valuation proof of the minimal order three of $\mathcal{T}_p$ (Section 33.3), a third route beside those of sources 13 and 14.

The thirteenth principal theorem comes from source 17 and is proved in Sections 34–35 (Convention 34.1). Let $(b_n)_{n \geq 2}$ be positive integers with $b_n/b_{n-1} \rightarrow \infty$, for instance $n!$, and for $A \subseteq \mathbb{N}_{\geq 2}$ put $\mathcal{G}_A(p, z) = \sum_{n \in A} p^{b_n} z^n$.

Theorem M (Differential freedom at a single scale; source 17). *Let k be a field of characteristic zero.*

- (a) *For every finite family $A_1, \dots, A_m$, after subtracting explicit polynomials that account for the finitely many indices with patterns $(\mathbf{1}_{A_i}(n))_i$ occurring finitely often, the differential relation ideal of $(\mathcal{G}_{A_i})_i$ over $k(p, z)$, for $\vartheta = zD_z$ and $\vartheta_p = p d/dp$, is generated by the rational linear relations among the patterns that occur infinitely often; if every A_i has infinitely many indices belonging to it alone, all mixed jets are algebraically independent.*

- (b) If Γ has an order unit μ , $\mathbb{K} = k((t^\Gamma))$ and $p = t^\mu$, every $\mathcal{G}_A$ with A infinite is a nonpolynomial element of $\mathcal{E}_\Gamma(k)$ that satisfies no nonzero algebraic differential equation over $\mathbb{K}(z)$, and the D_z -relations of a finite family over $\mathbb{K}(z)$ are exactly those of (a).
- (c) For $\mathbb{K} = \mathbb{R}((t^\mathbb{Q}))$ or $\mathbb{C}((t^\mathbb{Q}))$, $\text{dtrdeg}_{\mathbb{K}(z)} \text{Frac } \mathcal{E}_\Gamma(k) = 2^{\aleph_0}$, with a witnessing family whose Taylor coefficients all lie in $\mathbb{Q}(t)$.

The coefficient values $\mathbf{b}_n \mu$ span $\mathbb{Q}\mu$, so neither Theorem E (which needs no order unit) nor Theorem F (which needs infinite coefficient-value rank) applies: the freedom comes from the support, through a height-separation band and polarization of the top-degree part of a relation (Section 34.2). With Theorems H and L, an order-unit field thus carries strongly entire series of order three, of arbitrarily large order (for $\Gamma = \mathbb{R}$), and of no finite order. Source 17 also proves the independence of the Berarducci–Mantova jets of the numbers $\sum_{n \in A} \omega^{-\mathbf{b}_n}$ over a bounded-support field, the exact common strong domain $\{x : v(x) \geq -m \text{ for some } m\}$ of these series in $\mathbf{No}[i]$, and independent omnific codes with factorial floor profiles (Section 35). Differential transcendence of one such lacunary series is classical (Grönwall, via Ostrowski [36]).

The fourteenth principal theorem comes from source 15 and is proved in Sections 36–37 (Convention 36.1). For monomials $Y^{\mathbf{a}}$ in variables attached to arguments of degrees $e_1, \dots, e_{\bar{m}}$, the *composition weight* is $\text{cw}(\mathbf{a}) = \sum_\nu a_\nu e_\nu$.

Theorem N (Polynomial-composition rigidity; source 15). *Let k be a field of characteristic zero and $\mathbb{K} = k((t^\Gamma))$. Let $P \in \mathbb{K}[z]$ have degree $d \geq 2$, let $P_1, \dots, P_{\bar{m}} \in \mathbb{K}[z]$ be nonconstant of degrees $e_\nu < d$, let $L_1, \dots, L_{\bar{s}} (\bar{s} \geq 1)$ and $L'_1, \dots, L'_{\bar{m}}$ be nonzero linear differential operators with coefficients in $\mathbb{K}[z]$, let $0 \neq c_* \in \mathbb{K}[z]$, and let $\mathbf{G} \in \mathbb{K}[z][Y_1, \dots, Y_{\bar{m}}]$ have only monomials of composition weight $\text{cw}(\mathbf{a}) < \bar{s}d$. Then every $f \in \mathcal{E}_\Gamma(k)$ with*

$$c_*(z) \prod_{\ell=1}^{\bar{s}} L_\ell(f(P(z))) = \mathbf{G}(z, L'_1(f(P_1(z))), \dots, L'_{\bar{m}}(f(P_{\bar{m}}(z))))$$

is a polynomial, of degree at most an explicit integer B_{cmp} computed from the degrees, the operator shifts and the integer roots of the exact indicial polynomials $I_{L_\ell}(dT)$ (Theorem 36.12). In particular, for every Γ , every strongly entire solution of a linear equation $\sum_{j=0}^m L_j(f(P_j(z))) = g(z)$ with $L_m \neq 0$ and a unique argument P_m of largest degree $d_m \geq 2$ is a polynomial.

The lower arguments may be dilations λz . So, unlike Theorems B and J, no order unit and no condition on scales enters: a strictly dominant argument of degree at least two defeats the partial theta mechanism (Remark 36.10). The proof reads how a polynomial argument rescales the Gauss data of Section 14.3 and compares scales by an integer convexity inequality. Source 15 also proves a matrix version, describes the solution modules of linear equations with integer data and omnific coefficients, and shows that every affine hypersurface over $\mathbb{Q}$ is the entire-solution locus of one equation of this class, so omnific solvability is undecidable inside it (Theorem 37.3, Corollary 37.5).

The fifteenth principal theorem comes from source 20 and is proved in Sections 38–39 (Convention 38.1). A *coefficient-algebraic condition* on a vector of Hahn series is a polynomial equation over k in finitely many of their coefficients (Definition 38.7).

Theorem O (Coefficient observables of recurrences; source 20). *Let k be a field of characteristic zero and $\mathbb{K} = k((t^\Gamma))$.*

- (a) *Let $\mathbf{y}_n$ be a finite vector of exponential polynomials $\sum_j R_j(n) \lambda_j^n$, $R_j \in \mathbb{K}[U]$, whose bases λ_j have valuation zero and residues $\zeta_1, \dots, \zeta_m$. Every coefficient sequence $n \mapsto [t^\gamma]$ of every coordinate lies in the one finitely generated k -algebra $k[n, \zeta_1^n, \dots, \zeta_m^n]$ of sequences.*

- (b) *For every set-sized family of coefficient-algebraic conditions, a finite subfamily is equivalent to the whole family at every point of the orbit $(\mathbf{y}_n)_{n \geq 0}$, and the set of n at which $\mathbf{y}_n$ satisfies the family is a finite union of full residue classes modulo $\mathbf{M}_{\text{tor}}$, the exponent of the torsion subgroup of $\langle \zeta_1, \dots, \zeta_m \rangle$, and a finite set; one $\mathbf{M}_{\text{tor}}$ serves for every family.*
- (c) *Without units this fails completely: for every real sequence $(\mathbf{b}_m)_{m \geq 0}$ there is a purely infinite omnific integer $\mathbf{w}$ with $[\omega^\omega](\omega^n \mathbf{w}) = \mathbf{b}_n$ for every ordinary n , so the first-order omnific recurrence $y_{n+1} = \omega y_n$ realizes every subset of $\mathbb{N}$ as the nonzero set of one fixed Conway coefficient.*

Part (a) rests on the support envelope of Lemma 28.4; part (b) adds Hilbert’s basis theorem and a Skolem–Mahler–Lech period valid for all monomials in the residues (Lemma 38.4), for which the period of source 18’s valuation theorem is too small (Remark 38.5). The element in part (c) is a stream already in the omnific-notations report (Remark 39.6); new is its reading as a recurrence observable, which excludes a coefficientwise Skolem–Mahler–Lech theorem for omnific recurrences (Corollary 39.4), while the leading exponent of every recurrence over $\mathbf{No}[i]$, $\mathbf{Oz}$ or $\mathbf{Oz}[i]$ stays eventually affine on finitely many progressions (Theorem 39.1). Source 20 also re-derives, independently of source 18, the valuation theorem for Hahn exponential polynomials and the unit-dilation case of Theorem J, printed once; its exact finite-group criterion is a special case of that theorem (Section 39.4).

For the linear theorems, the hypotheses separate two obstructions. Differential equations have bounded valuation costs after passing to their coefficient recurrence. Dilation equations can have costs increasing linearly with the index, but only along the single scale $v(q)$; a scale that is not an order unit cannot reach all of Γ , even if its multiples are unbounded within their own Archimedean component. Strong entireness detects that failure.

Beyond the first three, the report proves: a mixed differential–dilation rigidity theorem at every nonzero noncofinal dilation scale (Theorem 10.2, extended by Theorem J to unit dilations and to finite sets of multipliers); a recurrence exclusion bound that is *attained*, boundary included, by an explicit formal exponential (Corollary 3.5 and Proposition 5.3); a coarsened exclusion bound valid at every nonzero noncofinal dilation valuation (Theorem 7.4); and the exact strong evaluation domain of the partial theta series for an arbitrary Hahn parameter, including arbitrary tails (Theorem 8.1).

1.4 Where divisibility of Γ is used

The core arguments — the escape mechanism, the orbit valuation lemmas, the two conversions, the theta domain theorem, and all of Theorems A–C — use only Convention 1.1. Divisibility of Γ enters in exactly three threshold constructions, with the hypothesis stated where it is used.

- (1) The refined recurrence threshold B_{rec} of (3.4), whose defining maximum divides a valuation difference by the jump length j . Corollary 3.5 states it under divisibility; Corollary 3.4 is the divisibility-free predecessor, with a coarser threshold and the same qualitative conclusion.
- (2) The refined coarsened threshold $\overline{B}_{\text{rec}}$ of (7.4), for the same reason, in the quotient group. Theorem 7.4 states the divisibility-free form and the divisible refinement separately.
- (3) The worked sparse threshold in Example 13.1, an instance of (1).

The torsion eigenspace equivalence in Proposition 9.1 needs no divisibility: inward stability of strong evaluation replaces the root-extraction argument of the source manuscripts.

The nonlinear part, Sections 14–17, uses no divisibility of Γ anywhere: the degree cutoff (15.11) divides ordinary integers in $\mathbb{Q}$, and its exclusion bounds are chosen from finitely many strict inequalities with positive integer multipliers.

The coefficient-field part, Sections 20–22, uses no divisibility either. It works in $\Gamma \otimes_{\mathbb{Z}} \mathbb{Q}$ only to measure rational rank, and the principal convex bound of Lemma 20.7 uses integer multiples only.

The theta-descent part uses no divisibility. The order-three-threshold part, Sections 26–27, states its theorems for Γ as in Convention 1.1; to find scales at which the initial polynomial is not a monomial it divides in the divisible hull $\Gamma \otimes_{\mathbb{Z}} \mathbb{Q}$, after extending the field by Lemma 26.2, and transfers the polynomiality and coefficient conclusions back to Γ , in which the hull is cofinal (Lemmas 26.2 and 26.4).

The unit-dilation part, Sections 28–29, uses no divisibility: its cost bounds are formed from finitely many elements and a noncofinality witness, and its jump-sensitive comparison multiplies by the jump instead of dividing by it. The Newton-rigidity part, Sections 30–31, passes to $\Gamma \otimes_{\mathbb{Z}} \mathbb{Q}$ exactly as the order-three-threshold part does, through Lemma 26.2, and transfers its conclusions back.

The theta-hierarchy part works in $\Gamma = \mathbb{R}$, which is divisible, and states it; the single-scale part uses no divisibility, and neither does the polynomial-composition part, whose degree bounds are floors of ratios of ordinary integers and whose convexity inequality multiplies by positive integers. The recurrence part uses no divisibility either: its periods are exponents of finite groups of residues, and its cost bounds are those of the unit-dilation part.

No theorem in this report states a conclusion under Convention 1.1 while resting on a proof step that divides by an integer in Γ ; the divisions of the order-three-threshold and Newton-rigidity parts take place in $\Gamma \otimes_{\mathbb{Z}} \mathbb{Q}$. The division-free predecessors in (1) and (2) are weaker in the constant only, not in the conclusion.

1.5 Position relative to the collection and the literature

Provenance. This report is built from thirteen manuscripts, numbered as they arrived in the collection. Sources 08 and 09 were written independently on 22 September 2026 and both prove the differential rigidity theorem and the order-unit dilation criterion (Theorems A and B); source 08 is the base, because it assumes only $\Gamma \neq 0$ while source 09 assumed divisibility, and every proof taken from 09 was re-read for a hidden division (Section 1.4). Together they supply Sections 1–13. Source 10, delivered later the same day, supplies the nonlinear part, Sections 14–17 and Theorem D. The repository snapshot consulted by sources 08 and 09 is

a3124af79f66b8b9c196d76b4cbc5ac3938907c4,

and the one consulted by source 10 is

048b72cf7cbfc8ab246e4f73788c10460cb3f6e0.

At the second pin this report consisted of sources 08 and 09 alone and its text was identical to the text into which source 10 was merged, so source 10’s references to “Question 15.1” and “Theorems A–C” of this report were accurate there; that question is now Question 19.1, re-scoped in Section 19. The choices the merge made (renamed symbols, results printed once, the placement of the nonlinear part after the linear theory) are recorded in Section 14.1.

Sources 11 and 12, delivered together in a later batch and written independently, are titled *Coarsening Descent and Nonlinear Differential Rigidity for Entire Hahn–Surcomplex Functions*

and *Coefficient-Field Rigidity at Unbounded Surreal Scales*. Both answer the no-order-unit case of Question 19.1 by the same argument, and together they form the coefficient-field part, Sections 20–22, with Theorems E–G. Source 12 is the base of that part: it adds dilations, answers the no-order-unit case of Question 19.4, and gives the order-unit dichotomy. Source 11 adds coarsening descent, rationality in the fraction field, the rational-system and Painlevé corollaries, the joint-independence criterion and the continuum family. Both consulted the snapshot

465a54b479a1ee842cbf7db1689a7d2f6bfe25e1,

at which this report was exactly Sections 1–19 and the conclusion, so their references to Questions 19.1 and 19.4 and to the README were accurate. The coefficient-field part is placed after Section 19 so that the numbering of Sections 1–19, of Questions 19.1–19.6 and of Theorems A–D, which the shipped audit files cite, is unchanged; the new questions are Questions 19.7–19.9. The merge’s choices for this part are recorded in Section 20.1.

Source 13, *Theta Descent and the Exact Boundary of Differential Rigidity*, delivered in a later batch, answers the order-unit case of Question 19.1 by an explicit series of minimal differential order three, and so shows that the order unit is the exact boundary. It forms the theta-descent part, Sections 23–25, with Theorem H, and adds clause (iv) to Theorem G. It consulted the snapshot

4cdeaec43ff1692ed2ad9f5bdf761a41872975a5,

at which this report was exactly the five-source text, so its references to Question 19.1 and Theorem E were accurate. The new part is placed after Section 22, so that the numbering of Sections 1–22, of Questions 19.1–19.9 and of Theorems A–G is unchanged; its one new question is Question 19.10. The merge’s choices for this part, including two steps it writes out, are recorded in Section 23.1.

Source 14, *The Order-Three Threshold for Entire Hahn–Surcomplex Functions*, delivered in a later batch, answers the order-two question that source 13 left open: no nonpolynomial strongly entire series satisfies an algebraic differential equation of order at most two, for every value group. It forms the order-three-threshold part, Sections 26–27, with Theorem I. It consulted the snapshot

b895e8672990e8b5a56f97dd7246f9dd5f86c771,

at which this report’s text was still the five-source text and source 13 was known to it only as a manuscript; its description of this report was accurate there. The new part is placed after Section 25, so that the numbering of Sections 1–25, of Questions 19.1–19.10 and of Theorems A–H is unchanged; the conclusion moves to the end. Its new questions are Questions 19.11–19.18. The merge’s choices for this part are recorded in Section 26.1.

Sources 18 and 19 arrived together in a later batch (batch 31, manuscripts 05 and 08), written independently of each other. Source 18, *Unit-Dilation Rigidity at Surreal Scales*, answers Question 19.4 positively for every value group and proves the relative-scale criterion of Theorem J; it forms the unit-dilation part, Sections 28–29. It consulted the snapshot

bcac55ae6fd3f2b354e568ba1d2e94d496a2cc59.

Source 19, *Low-Order Nonlinear Rigidity at Surreal Scales*, re-derives Theorem I (a) independently, printed once, and proves Theorem K; it forms the Newton-rigidity part, Sections 30–31. It consulted the snapshot

3d40856953f4bb0f5d069e45a3b4bab6eda33040.

At both snapshots this report was still the five-source text, so their descriptions of it were accurate there; what has since become stale is recorded in Sections 28.1 and 30.1. The two parts are placed after Section 27, so that the numbering of Sections 1–27, of Questions 19.1–19.18 and of Theorems A–I is unchanged; the conclusion moves to the end. Their principal theorems are Theorems J and K, and their new questions are Questions 19.19–19.29 and 19.30–19.35.

Sources 16 and 17 (batch 31, manuscripts 02 and 03), written independently, were merged next. Source 16, *Independent Theta Families and Unbounded Differential Order in Hahn–Surcomplex Analysis*, forms the theta-hierarchy part, Sections 32–33, with Theorem L; it records the snapshot

0fffc26c670c15c7db451ed899f98d4384f9df1e,

at which this report’s text was the five-source one and source 13’s audits were placed. Source 17, *Differential Freedom at a Single Surreal Scale*, forms the single-scale part, Sections 34–35, with Theorem M; it consulted `bcac55a`, the pin of source 18. What has since become stale in them is recorded in Sections 32.1 and 34.1. The two parts follow Section 31, so that the numbering of Sections 1–31, of Questions 19.1–19.35 and of Theorems A–K is unchanged; the conclusion moves to the end. Their new questions are Questions 19.36–19.44 and 19.45–19.54.

Source 15 (batch 31, manuscript 01), *Polynomial-Composition Rigidity at Surreal Scales*, was merged last of its batch. It forms the polynomial-composition part, Sections 36–37, with Theorem N; it consulted `bcac55a`, the pin of source 18, at which this report was the five-source text. What has since become stale in it, and one statement that was already incorrect there, are recorded in Section 36.1. The part follows Section 35, so that the numbering of Sections 1–35, of Questions 19.1–19.54 and of Theorems A–M is unchanged; the conclusion moves to the end. Its new questions are Questions 19.55–19.62.

Source 20 (batch 33, manuscript 01), *Valuation Rigidity and Coefficient Universality in Surreal Recurrences*, forms the recurrence part, Sections 38–39, with Theorem O. It consulted the snapshot

efc5446229dbf61a97bdd5edae91dba20af9d131,

at which this report was the seven-source text (sources 08–14) and sources 15–19 were placed but not yet merged; its statement that the mixed unit-valuation case was left open was accurate there and is stale now, and its re-derivations of source 18’s results are printed once (Section 38.1). The part follows Section 37, so that the numbering of Sections 1–37, of Questions 19.1–19.62 and of Theorems A–N is unchanged; the conclusion moves to the end. Its new questions are Questions 19.63–19.70.

The companion entire-function report. The collection already contains a merged report on cofinality, factorization and scalar extension for entire Hahn functions at arbitrary rank [12]. It fixes the same class $\mathcal{E}_\Gamma$ over the same kind of workspace, classifies the ring structure of $\mathcal{E}_\Gamma$, and computes the exact strong evaluation domain after an enlargement of the value group. None of its ring-theoretic conclusions is used to prove anything here: the present proofs start from Hahn support arithmetic. Two points of contact matter.

First, we do not reprove the order-unit coarsening machinery. That report’s Lemma 9.1, labelled `ent:lem:coarsening`, states that if $h > 0$ is an order unit of Γ there is an order-preserving homomorphism $\pi : \Gamma \rightarrow \mathbb{R}$ with $\pi(h) = 1$ whose kernel is the convex subgroup $H = \{\gamma : n|\gamma| < h \text{ for all } n \geq 1\}$, so that $|a|_\pi = \exp(-\pi(v(a)))$ is a nontrivial real-valued non-Archimedean absolute value inducing the intrinsic valuation topology. That is exactly the sense in which sources 08 and 09 of this report use the phrase “coarsening”, and we cite it rather than repeat it; see Remark 2.6.

Second, and this is easy to get wrong: *the two order-unit dichotomies are not the same theorem* — the companion report’s order unit decides Hermite interpolation and the Bézout property of the ring $\mathcal{E}_\Gamma$ (its Theorem 9.3), whereas the order unit here decides which annihilating operators a nonpolynomial member of $\mathcal{E}_\Gamma$ can satisfy. Neither dichotomy implies the other, and the coincidence of the invariant is not a coincidence of statement. In particular the existence of nonpolynomial entire functions at countable cofinality, and the fact that $\Gamma_\infty = \bigoplus_{j \geq 0} \mathbb{Q}\omega^j$ has countable cofinality and no order unit, are background from that report [12], not targets of this one.

The collection’s differential-equations report uses a derivation on surreal *scalars* [13]; here D_z kills every scalar and differentiates only the formal variable, and no theorem about that scalar derivation is restated as ordinary D-finite rigidity. The rank-one analytic report [14] already contains the quadratic-exponent example $\sum t^{n^2} z^n$; our partial theta witness is explicitly classical and is not counted as a new function.

External literature. Stanley’s D-finite/P-recursive correspondence is the standard algebraic starting point [1]. We make the coefficient conversion explicit because the valuations of the resulting recurrence, rather than ordinary coefficient growth, drive the proof. Ramis treats growth of ordinary complex entire solutions of linear q -difference equations and also studies simultaneous differential and difference constraints [2]; the arbitrary-rank strong summability condition here is a different global condition, and none of its order-unit conclusions is inferred from an ordinary complex growth estimate. Garoufalidis proves eventual quadratic quasi-polynomial degree behaviour for q -holonomic sequences in a rational-function and Puiseux setting [3]; that is an important antecedent to the idea that dilation equations constrain coefficient scales, we do not claim to originate it, and we do not strengthen its quadratic quasi-polynomial conclusion. Di Vizio’s ultrametric Maillet–Malgrange theorem treats Gevrey behaviour of formal solutions of nonlinear ultrametric q -difference equations, including the valuation-zero case with Diophantine conditions [4]; we make no nonlinear generalization of it. The partial theta series belongs to the classical partial-theta literature [5], and Higman’s finite-word theorem is a standard external ingredient in the positive-support monoid lemma [6].

Terminology needs care in two further directions. Hahn meromorphic functions in the analytic setting of Müller and Strohmaier involve a convergence framework different from the formal strong evaluation used here [7]. The Berarducci–Mantova derivation differentiates surreal coefficients themselves and is not D_z [8]. Work on operation-closed surreal subfields gives context for set-sized workspaces [9], and the Conway normal-form conventions used in Section 2.4 follow current surreal field work [10]; closure under a scalar derivation is assumed in no theorem below.

The inspections of sources 08 and 09 were targeted: repository documentation, the report catalogue, the entire-function report, and the primary sources named above. Neither was a line-by-line audit of every repository manuscript or an exhaustive bibliography search, and a repository content search for `holonomic` returning no matches is not proof of absence. Source 10’s repository and literature comparison, equally targeted, is recorded in Section 17.4, that of sources 11 and 12 in Section 22.6, that of source 13 in Section 25.5, that of source 14 in Section 27.5, those of sources 18 and 19 in Sections 29.7 and 31.4, and those of sources 16 and 17 in Sections 33.4 and 35.6, that of source 15 in Section 37.5, and that of source 20 in Section 39.6. Section 19 and Sections 22.7, 25.6, 27.6, 29.8, 31.5, 33.5, 35.7, 37.6 and 39.7 record exactly what is and is not claimed.

2 Hahn support and what evaluation means

2.1 Support calculus

Lemma 2.1 (Hahn–Neumann support calculus and the positive monoid). *Let A, B be well-ordered subsets of an ordered abelian group.*

- (1) *$A + B$ is well ordered, and each γ has only finitely many representations $\gamma = a + b$ with $(a, b) \in A \times B$.*
- (2) *If S is a well-ordered subset of $\Gamma_{>0}$, the additive monoid*

$$M(S) = \{s_1 + \cdots + s_k : k \geq 0, s_i \in S\}$$

is well ordered, and each of its elements has only finitely many representations as a finite word in letters from S .

- (3) *Multiplication by a fixed Hahn element preserves strong summability, and multiplication by a nonzero one preserves and reflects it.*

Proof. An infinite sequence in a well-ordered set has an infinite nondecreasing subsequence. Given a strictly decreasing sequence $a_n + b_n$, pass first to a subsequence with a_n nondecreasing and then to one with b_n nondecreasing, a contradiction. If infinitely many distinct pairs sum to a fixed γ , the distinct first coordinates contain an increasing sequence, forcing a decreasing sequence of second coordinates. This is part (1).

For part (2), order finite words over S by subsequence embedding with coordinatewise increase. Higman’s theorem says that words over a well-quasi-ordered alphabet are well quasi ordered [6]. An embedding cannot decrease the sum, and increases it strictly unless the two words are identical, since inserted letters are positive and a strict coordinate increase is positive. A descending sequence of sums therefore gives a bad sequence of words, and so do infinitely many distinct words with one fixed sum. Both are impossible.

For part (3), let A be the union of the supports of a strongly summable family and B the support of the multiplier. The new supports lie in $A + B$. Part (1) leaves finitely many admissible pairs at a specified exponent, and for each first coordinate only finitely many family members contribute. For a nonzero multiplier apply the same to its inverse. $\square$

In part (2) words carry their order; a fixed finite multiset has only finitely many permutations, so the commutative monoid version follows. The argument does not assume that S is bounded below by an order unit, and remains valid at arbitrary valuation rank.

Lemma 2.2 (The nonincreasing leading-exponent obstruction). *A family of nonzero Hahn elements possessing an infinite subsequence whose leading exponents are nonincreasing is not strongly summable.*

Proof. If that sequence of leading exponents takes infinitely many distinct values it has an infinite strictly decreasing subsequence, so the union of the supports is not well ordered. Otherwise it is eventually constant, and one exponent lies in the supports of infinitely many members, so local finiteness fails. $\square$

The second alternative is essential and is what makes the exclusion boundary below a *closed* region. Merely excluding a descending support does not establish strong summability: the family $(1 \cdot z^n)_{n \geq 0}$ at $z = 1$ has support union $\{0\}$, yet every summand contributes at exponent zero.

Lemma 2.3 (Two sufficient certificates). *Let $(b_n)_{n \geq 0}$ be a family in K_Γ .*

- (a) *If $v(b_n)$ tends cofinally to $+\infty$, meaning that for each $\beta \in \Gamma$ all but finitely many nonzero b_n satisfy $v(b_n) > \beta$, then (b_n) is strongly summable.*
- (b) *If $\text{supp}(b_n) \subseteq h_n + M(S)$ for a well-ordered $S \subseteq \Gamma_{>0}$, where the set of values of h_n is well ordered and each value is taken by only finitely many indices, then (b_n) is strongly summable.*

Moreover, grouping a strongly summable family into finite blocks preserves strong summability and its sum.

Proof. For (a) fix β . Only finitely many supports meet $(-\infty, \beta]$, and a finite union of well-ordered subsets of a linear order is well ordered. A descending sequence in the full union would lie below its first member, hence inside such a finite union. The same finiteness bounds the multiplicity of an exponent. For (b) use parts (1) and (2) of Lemma 2.1: a fixed target exponent admits finitely many pairs $(h, m) \in \{h_n\} \times M(S)$ with $h + m$ equal to it, and finitely many indices carry each such h ; actual supports can only be smaller than the indicated sets. For the last assertion, each block support is contained in the union of the old supports and the blocks contributing at any exponent form a finite set, while coefficientwise associativity is finite. $\square$

Criterion (a) and criterion (b) are genuinely different tools and both are used below. Criterion (a) is the shorter route whenever the term valuations really do escape every bound. Criterion (b) is what survives at the boundary, where the term valuations increase strictly but remain trapped below a fixed element of Γ ; Theorem 8.1 is exactly such a case, and no leading-valuation argument alone can decide it.

Lemma 2.4 (Inward stability of strong evaluation). *Let $f = \sum_n a_n z^n \in K_\Gamma[[z]]$, and let $x, y \in K_\Gamma^\times$ with $v(y) \geq v(x)$. If $x \in \text{Dom}_{\text{str}}(f)$, then $y \in \text{Dom}_{\text{str}}(f)$. Thus membership of a nonzero argument in $\text{Dom}_{\text{str}}(f)$ depends only on its valuation, and entireness can be tested on monomial arguments.*

Proof. Put $b_n = a_n x^n$ and $u = y/x$, so (b_n) is strongly summable and $v(u) = \eta \geq 0$. Write $u = c t^\eta(1 + w)$, where $c \in \mathbb{C}^\times$ and w is zero or has positive valuation. Let S be $\text{supp}(w)$, with η adjoined when $\eta > 0$, and put $M = M(S)$. This is a well-ordered positive alphabet, so Lemma 2.1 makes M well ordered. Every finite power u^n has support in M .

Let $A = \bigcup_n \text{supp}(b_n)$. The supports of $a_n y^n = b_n u^n$ lie in $A + M$, a well-ordered set. For a fixed exponent γ there are only finitely many pairs $(\alpha, \mu) \in A \times M$ with $\alpha + \mu = \gamma$. For each such α , strong summability of (b_n) leaves only finitely many indices n . Hence γ occurs in only finitely many supports of $b_n u^n$, proving strong summability. Equal valuations allow the argument in both directions; compare an arbitrary nonzero y with $t^{v(y)}$ to obtain the monomial test. Evaluation at zero is always finite. $\square$

The statement fixes the full coefficient series a_n . It does not say that their leading valuations alone determine the domain: higher coefficient tails can still matter at a boundary scale.

2.2 Positive valuation is not cofinal decay

For $\beta > 0$ write

$$\Delta_\beta = \{\gamma \in \Gamma : |\gamma| \leq N\beta \text{ for some } N \in \mathbb{N}\}, \quad (2.1)$$

the smallest convex subgroup containing β . It is all of Γ exactly when β is an order unit. If it is proper, there is $\gamma > 0$ outside it, and then

$$\gamma > N\beta \quad \text{for every ordinary } N \in \mathbb{N}, \quad (2.2)$$

since a positive element bounded by some $N\beta$ would lie in Δ_β .

Remark 2.5 (Sufficiency is not necessity). Let $0 < \epsilon < H$ in Γ with $n\epsilon < H$ for every ordinary integer n ; for instance $\Gamma = \mathbb{Q}H \oplus \mathbb{Q}\epsilon$ ordered by the highest nonzero coordinate. The family $(t^{n\epsilon})_{n \geq 0}$ is strongly summable, since its support is an increasing sequence and each exponent occurs once, and its sum is the Hahn element $(1 - t^\epsilon)^{-1}$. Yet its valuations do not tend cofinally to $+\infty$. This is the most elementary warning against identifying positive valuation with topological nilpotence, and it is why criterion (b) of Lemma 2.3 is needed at all. No appeal to convergence of partial sums at the H scale is involved.

Remark 2.6 (The real-valued coarsening is imported, not reproved). When Γ does have an order unit h , the companion report [12, Lemma 9.1, `ent:lem:coarsening`] supplies an order-preserving $\pi : \Gamma \rightarrow \mathbb{R}$ with $\pi(h) = 1$ and kernel the convex subgroup $H = \{\gamma : n|\gamma| < h \text{ for all } n \geq 1\}$, hence a nontrivial real-valued non-Archimedean absolute value $|a|_\pi = \exp(-\pi(v(a)))$ inducing the intrinsic topology. We cite that lemma and do not reprove it. No proof in this report uses it: every argument below is carried out in Γ itself, which is what makes the arguments valid when no order unit exists. The coarsening is mentioned twice, as an optional rank-one picture of the order-unit case (Remark 8.3) and for the warning of Section 1.5 that the interpolation/Bézout dichotomy of that report and the annihilating-operator dichotomy of this one share an invariant but are different theorems.

2.3 The entire class and its closure properties

Polynomials are entire, because all the evaluation families are finite. An ordinary complex entire Taylor series need not be entire in the sense of Definition 1.2: at a monomial argument $x = t^\gamma$ the family

$$\left(\frac{x^n}{n!}\right)_{n \geq 0} \quad (2.3)$$

is strongly summable exactly when $\gamma > 0$, since $\gamma = 0$ puts infinitely many contributions at exponent zero and $\gamma < 0$ gives a descending sequence. Proposition 5.3 computes the exact domain of the corresponding family with an arbitrary Hahn argument.

Proposition 2.7 (Closure). $\mathcal{E}_\Gamma$ is a K_Γ -algebra, $\sigma_q(\mathcal{E}_\Gamma) \subseteq \mathcal{E}_\Gamma$ for every $q \in K_\Gamma^\times$, and $D_z(\mathcal{E}_\Gamma) \subseteq \mathcal{E}_\Gamma$.

Proof. For addition, the evaluation supports lie in the union of two well-ordered sets with finite exponent multiplicities. For multiplication, the doubly indexed product family is strongly summable by Lemma 2.1 (1) and (3), and grouping by ordinary total degree preserves strong summability by the last clause of Lemma 2.3. Dilation is evaluation at qx . For $x \neq 0$ the derivative family is

$$(n+1)a_{n+1}x^n = x^{-1} \cdot (n+1)a_{n+1}x^{n+1}.$$

Discarding one term and multiplying each term by a nonzero ordinary integer enlarges neither supports nor multiplicities, and multiplication by x^{-1} preserves summability by Lemma 2.1 (3). At $x = 0$ only the constant term contributes. Iterate. $\square$

Proposition 2.8 (Faithfulness of evaluation). *If $f \in \mathcal{E}_\Gamma$ vanishes at infinitely many distinct ordinary complex arguments, then $f = 0$ as a formal series. Consequently a linear equation with rational-function coefficients that holds pointwise wherever its coefficients are defined is also a formal equation in the sense of Definition 1.4.*

Proof. Strong evaluation at 1 says that the coefficient family (a_n) has a well-ordered union of supports and is locally finite at every exponent. Hence for each γ

$$P_\gamma(T) = \sum_n (a_n)_\gamma T^n \in \mathbb{C}[T]$$

is a polynomial, and its value at an ordinary complex c is the coefficient of t^γ in $f(c)$. If f vanishes at infinitely many such c , every P_γ has infinitely many roots and is zero, so every a_n is zero. For the last assertion, clear denominators; the left-hand side is in $\mathcal{E}_\Gamma$ by Proposition 2.7, and nonzero polynomial denominators exclude at most finitely many ordinary complex arguments. $\square$

The nonexistence proofs may therefore be run on formal coefficients without weakening their reading as functional obstructions, and no interchange of infinite limits is needed anywhere.

The theorems below use only the valuation laws (1.1), the existence of every monomial t^γ , triviality of the valuation on $\mathbb{C}^\times$, and the support facts of this section. They do not require algebraic closedness, spherical completeness, Weierstrass preparation, or an embedding of Γ into $\mathbb{R}$.

2.4 Embedding the workspace into the surcomplex numbers

If Γ is an additive ordered subgroup of $\mathbf{No}$, the Conway normal-form map

$$\sum_{\gamma \in S} c_\gamma t^\gamma \mapsto \sum_{\gamma \in S} c_\gamma \omega^{-\gamma} \tag{2.4}$$

identifies the real-coefficient Hahn field with a subfield of $\mathbf{No}$ and its complexification with a subfield of $\mathbf{No}[i]$; a well-ordered S becomes a reverse-well-ordered set of Conway exponents. This viewpoint and its set-sized support discipline are standard [9, 10]. Here t^γ is a formal monomial and $\omega^{-\gamma}$ a Conway monomial; the notation does not silently identify every formal power with a choice of surreal exponentiation.

The results may therefore be read twice. Abstractly they concern Hahn fields and need no surreal construction. Concretely they restrict the exact linear descriptions available for surcomplex-valued power series on specified surreal scales. Formula (2.4) is the only bridge used, and no global surcomplex exponential is invoked.

3 The finite-step escape mechanism

The following isolates the mechanism used throughout. It needs no formula for the recurrence coefficients, no asymptotic slope for the valuations, and no assumption that cancellations are rare.

Lemma 3.1 (Escape chain). *Suppose a sequence (a_n) in K_Γ satisfies, for every $n \geq N$,*

$$c_0(n)a_n + \sum_{j=1}^s c_j(n)a_{n+j} = 0, \quad c_0(n) \neq 0, \quad (3.1)$$

and suppose $a_{n_0} \neq 0$ for some $n_0 \geq N$. Then $s \geq 1$ and there is an infinite sequence $n_0 < n_1 < n_2 < \dots$ with $j_r := n_{r+1} - n_r \in \{1, \dots, s\}$, with $a_{n_r} \neq 0$ and $c_{j_r}(n_r) \neq 0$ for every r , and with

$$v(a_{n_{r+1}}) - v(a_{n_r}) \leq v(c_0(n_r)) - v(c_{j_r}(n_r)). \quad (3.2)$$

Proof. Let $n \geq N$ with $a_n \neq 0$. The leftmost term of (3.1) is nonzero, so $s \geq 1$ and at least one later term is nonzero. If every nonzero later term had valuation strictly larger than $v(c_0(n)a_n)$, the leading monomial of the leftmost term could not cancel, contradicting (1.1). So some $j \in \{1, \dots, s\}$ has $c_j(n)a_{n+j} \neq 0$ and

$$v(c_j(n)a_{n+j}) \leq v(c_0(n)a_n),$$

which is (3.2). Taking the least eligible j makes the recursion definite. The successor coefficient is nonzero and its index is again at least N , so the construction never stops. $\square$

Remark 3.2 (The direction of the inequality matters). Solving the recurrence for its *last* coefficient typically yields a *lower* bound on a valuation, which obstructs nothing. The comparison above is made from the first nonzero coefficient forward and produces a later coefficient whose valuation is controlled from above. Cancellation among the other terms can raise the valuation of their sum, but cannot make every term on the right have valuation strictly larger than the single nonzero term on the left. Zeros and arbitrarily large cancellations elsewhere therefore do not invalidate the chain; the chain records only that finite inequality, repeatedly.

Theorem 3.3 (Evaluation exclusion criterion). *In the situation of Lemma 3.1, let $x \in K_\Gamma^\times$ and suppose that for every $n \geq N$ and every $j \in \{1, \dots, s\}$ with $c_j(n) \neq 0$,*

$$v(c_0(n)) - v(c_j(n)) + jv(x) \leq 0. \quad (3.3)$$

If $a_{n_0} \neq 0$ for some $n_0 \geq N$, then $(a_n x^n)_{n \geq 0}$ is not strongly summable. Equivalently: if $(a_n x^n)_{n \geq 0}$ is strongly summable, then $a_n = 0$ for every $n \geq N$.

Proof. Build the chain of Lemma 3.1 from n_0 and add $n_{r+1}v(x)$ to each side of (3.2), using $n_{r+1} = n_r + j_r$:

$$v(a_{n_{r+1}}x^{n_{r+1}}) \leq v(a_{n_r}x^{n_r}) + v(c_0(n_r)) - v(c_{j_r}(n_r)) + j_r v(x) \leq v(a_{n_r}x^{n_r}).$$

The evaluated leading exponents along the chain are therefore nonincreasing, and Lemma 2.2 applies. $\square$

Note that (3.3) is a weak inequality. This is the difference between an open and a closed excluded region, and it is available only because Lemma 2.2 also rules out a constant leading exponent. Section 5.2 exhibits an equation whose excluded region is exactly the closed one, and Section 8.5 exhibits a coarsened comparison where the boundary genuinely must be left in.

Corollary 3.4 (Uniform bounded cost; no divisibility). *Suppose in (3.1) there is a single $B \in \Gamma$ with $v(c_0(n)) - v(c_j(n)) \leq B$ for all $n \geq N$ and all $j \geq 1$ with $c_j(n) \neq 0$. Put $B^+ = \max(B, 0) \in \Gamma$. Then for every $x \neq 0$ with $v(x) \leq -B^+$, strong summability of $(a_n x^n)_n$ forces $a_n = 0$ for all $n \geq N$. Such x exist: $x = t^{-B^+}$, and $x = t^{-\delta}$ for every $\delta \geq B^+$.*

Proof. Since $B^+ \geq 0$ we have $v(x) \leq 0$, so $jv(x) \leq v(x) \leq -B^+ \leq -B$ for $j \geq 1$. Hence $v(c_0(n)) - v(c_j(n)) + jv(x) \leq B - B = 0$, and Theorem 3.3 applies. Every ordered abelian group admits a maximum of finitely many elements and the monomial t^{-B^+} exists, so no cofinal cyclic subgroup is needed. $\square$

The absence of a cofinality requirement here is exactly why the mechanism proves rigidity in groups of very large rank. It is also why the refined bound below, which divides by j , is a refinement of the constant and never of the conclusion.

Corollary 3.5 (Refined periodic threshold; requires Γ divisible). *Assume in addition that Γ is divisible. Assume each slot of (3.1) is either identically zero for $n \geq N$ or nonzero for all $n \geq N$ with periodic valuation there of common period $m \geq 1$; enlarge N past any initial nonperiodic range first. Let $\alpha_{j,r}$ be the valuation of slot j on the residue class $r \bmod m$, the identically zero slots being omitted. Put*

$$B_{\text{rec}} = \max_{\substack{0 \leq r < m \\ 1 \leq j \leq s, c_j \neq 0}} \frac{\alpha_{0,r} - \alpha_{j,r}}{j} \in \Gamma. \quad (3.4)$$

If there is no active forward slot $j \geq 1$, set $B_{\text{rec}} = 0$; the recurrence then forces $a_n = 0$ for all $n \geq N$ directly. Then for every $x \neq 0$ with $v(x) \leq -B_{\text{rec}}$, strong summability of $(a_n x^n)_n$ forces $a_n = 0$ for all $n \geq N$. In particular every solution with $a_n \neq 0$ for some $n \geq N$ — so in particular every nonpolynomial solution — satisfies

$$\text{Dom}_{\text{str}}(f) \subseteq \{0\} \cup \{x \in K_\Gamma^\times : v(x) > -B_{\text{rec}}\}, \quad f = \sum_n a_n z^n. \quad (3.5)$$

Proof. The case with no forward slot was handled in the statement. Otherwise divisibility makes each quotient $(\alpha_{0,r} - \alpha_{j,r})/j$ an element of Γ , so B_{rec} is defined. If $v(x) \leq -B_{\text{rec}}$ then $jv(x) \leq -j(\alpha_{0,r} - \alpha_{j,r})/j = -(\alpha_{0,r} - \alpha_{j,r})$, because multiplying a weak inequality by the positive integer j preserves it. Hence (3.3) holds and Theorem 3.3 applies. $\square$

The inequality in (3.5) is strict on the right: at $v(x) = -B_{\text{rec}}$ the chain may keep a constant leading exponent rather than descend, which is precisely why both halves of Definition 1.2 are needed. The bound is an *exclusion* bound. It does not assert that every point on the other side lies in the domain.

4 Valuations on integer and geometric orbits

4.1 Ordinary integers cannot create unbounded valuation loss

Lemma 4.1 (Integer evaluation). *Let $0 \neq P(X) = \sum_k p_k X^k \in K_\Gamma[X]$ and put $\beta(P) = \min_{p_k \neq 0} v(p_k)$. For all but finitely many nonnegative ordinary integers n ,*

$$P(n) \neq 0 \quad \text{and} \quad v(P(n)) = \beta(P). \quad (4.1)$$

Proof. The polynomial $t^{-\beta(P)}P$ has coefficients of nonnegative valuation and at least one of valuation zero, so its coefficientwise reduction

$$\overline{P}(X) = \sum_{v(p_k) = \beta(P)} \text{lc}(p_k) X^k \in \mathbb{C}[X]$$

is nonzero and has finitely many roots. The ordinary integers have distinct images in $\mathbb{C}$. At every integer n with $\overline{P}(n) \neq 0$, the Hahn coefficient of $t^{-\beta(P)}P(n)$ at exponent zero is nonzero, and no smaller exponent occurs. $\square$

Residue characteristic zero does real work here. The proof does not apply to p -adic integer inputs, where positive integers can have arbitrarily large valuation. Conversely no Diophantine approximation over $\mathbb{C}$ is relevant to v : every nonzero complex number, however small in ordinary modulus, has valuation zero.

4.2 A geometric orbit of nonzero valuation

Lemma 4.2 (Eventually affine valuation). *Let $q \in K_\Gamma^\times$ with $\lambda = v(q) \neq 0$ and let $0 \neq P(X) = \sum_k p_k X^k \in K_\Gamma[X]$. There are an index k_P with $p_{k_P} \neq 0$, the value $\beta_P = v(p_{k_P})$, and $N_P \in \mathbb{N}$, such that*

$$v(P(q^n)) = \beta_P + k_P n \lambda \quad (n \geq N_P). \quad (4.2)$$

In particular $P(q^n)$ is eventually nonzero.

Proof. The nonzero terms have valuations $v(p_k) + kn\lambda$. For two distinct indices the difference $v(p_k) - v(p_l) + (k - l)n\lambda$ is strictly monotone in n , so its sign is eventually constant and it vanishes for at most one integer, whether or not its values are cofinal in any convex subgroup. Finitely many pairs give one common threshold beyond which every pairwise comparison is strict and fixed, hence a unique minimizing index k_P . Its term cannot cancel against terms of strictly larger valuation. $\square$

Remark 4.3 (A rank-one shortcut that fails). The minimizing index need not be the smallest when $v(q) > 0$. In $\Gamma = \mathbb{Q}H \oplus \mathbb{Q}\epsilon$ with H dominating every $n\epsilon$, take $q = t^\epsilon$ and $P(X) = X + t^{-H}$: the constant term has smaller valuation than q^n for every n , however large. Conversely with $P(X) = 1 + t^{-H}X$ the nonconstant term is the minimum for every n . The same phenomenon in the other sign convention: in $\Gamma = \mathbb{Q}e_0 \oplus \mathbb{Q}e_1$ with $e_1 \gg e_0 > 0$, $q = t^{e_0}$ and $P(X) = t^{e_1} + X$ give $v(P(q^n)) = ne_0$ for every $n \geq 0$, so the nonzero constant coefficient never becomes the least-valued term. The proof above uses the eventual ordering of finitely many affine functions, not the Archimedean assertion that $n\epsilon$ eventually exceeds H , and not the rank-one slogan that q^n eventually beats every coefficient ratio.

4.3 Valuation-zero dilations, including residues that are roots of unity

Distinct powers of a valuation-zero q all have the same valuation, so the previous lemma says nothing. Polynomial evaluation nevertheless has only a finite periodic valuation pattern.

Theorem 4.4 (Unit-orbit valuations). *Let $q \in K_\Gamma^\times$ have $v(q) = 0$ and infinite multiplicative order, and let $0 \neq P \in K_\Gamma[X]$. Then $v(P(q^n))$ is eventually finite and periodic. More precisely, with $\zeta = \text{res}(q) \in \mathbb{C}^\times$:*

- (a) *if ζ is not a root of unity, $v(P(q^n))$ is eventually constant, equal to the minimum valuation of the coefficients of P ;*
- (b) *if ζ has exact order h , then $v(P(q^n))$ is eventually constant on each residue class modulo h , so its eventual period divides h .*

Proof. For (a) write $P(X) = \sum_k p_k X^k$ and $\beta = \min_k v(p_k)$. The coefficient of $P(q^n)$ at exponent β is $\overline{P}(\zeta^n)$ with $\overline{P}(X) = \sum_{v(p_k)=\beta} \text{lc}(p_k) X^k$, a nonzero complex polynomial. The points ζ^n are distinct, so this vanishes only finitely often, and no exponent below β occurs.

For (b) write

$$q = \zeta(1 + u), \quad \eta = v(u) > 0, \quad d = \text{lc}(u).$$

Here $u = q/\zeta - 1$ has positive valuation because q/ζ has valuation zero and residue 1, and $u \neq 0$ because otherwise $q = \zeta$ would have finite order. For every positive ordinary integer n the finite binomial expansion gives

$$y_n := (1 + u)^n - 1, \quad v(y_n) = \eta, \quad \text{lc}(y_n) = nd, \quad (4.3)$$

the leading coefficient being nonzero in characteristic zero. There is no convergence assertion: for each fixed n only finitely many powers occur.

Fix $r \in \{0, \dots, h-1\}$. For $n \equiv r \pmod{h}$ we have $\zeta^n = \zeta^r$, hence $q^n = \zeta^r(1 + y_n)$. Expand the nonzero polynomial

$$P(\zeta^r(1 + Y)) = \sum_{k=0}^{\deg P} b_{r,k} Y^k \quad (4.4)$$

and set

$$\beta_r = \min_{b_{r,k} \neq 0} (v(b_{r,k}) + k\eta), \quad J_r = \{k : v(b_{r,k}) + k\eta = \beta_r\}. \quad (4.5)$$

Then $P(q^n) = \sum_k b_{r,k} y_n^k$, and its coefficient at exponent β_r is the value at n of

$$R_r(T) = \sum_{k \in J_r} \text{lc}(b_{r,k}) d^k T^k \in \mathbb{C}[T]. \quad (4.6)$$

This polynomial is nonzero: its terms have distinct degrees and all the indicated coefficients are nonzero. Outside its finitely many integer roots the coefficient at β_r survives, so $v(P(q^n)) = \beta_r$ eventually on that class. Finitely many classes give one finite exceptional range. $\square$

Remark 4.5 (A second normalization of the same proof). Sources 08 and 09 both prove Theorem 4.4 (b), by two slightly different normalizations, and we record the other one because it is occasionally more convenient. Instead of factoring out the complex root of unity, put $q^h = 1 + u$ directly, with $\eta = v(u) > 0$ and $u \neq 0$ for the same reason. Write $n = r + hk$ and expand the polynomial $Q_r(Y) = P(q^r Y)$ as $Q_r(1 + V) = \sum_k b_{r,k} V^k$. The finite binomial expansion gives $V_k = (1 + u)^k - 1$ with $v(V_k) = \eta$ and $\text{lc}(V_k) = k \text{lc}(u)$, and $P(q^n) = Q_r(1 + V_k)$. The minimum construction (4.5) and the corresponding nonzero residue polynomial (4.6), now in the variable k rather than n , finish the argument. Neither version sums a binomial series to a noninteger exponent; every expansion is finite, and arbitrary supports in q and in the coefficients of P enter only through the leading valuation and leading coefficient of finitely many Hahn elements.

Example 4.6 (A genuine two-phase pattern). Let $q = -(1 + t^\eta)$ with $\eta > 0$. Then q is nontorsion, $\text{res}(q) = -1$ has order two, and for $P(X) = X - 1$ and $n \geq 1$

$$v(P(q^n)) = v(q^n - 1) = \begin{cases} 0, & n \text{ odd}, \\ \eta, & n \text{ even}. \end{cases}$$

For odd n the residue is -2 ; for even positive n the leading term is nt^η . The valuation sequence is periodic, not the dilation itself, and a claim of eventual *constancy* without splitting residue classes would be false. The escape argument uses the bounded valuation costs; it does not replace q by its residue.

Example 4.7 (Finite exceptional indices really occur). For $q = 1 + t^\eta$ and $P(X) = X - 1 - 7t^\eta$, the leading coefficient of $P(q^n)$ at exponent η is $n - 7$, so $v(P(q^n)) = \eta$ for positive $n \neq 7$ while $v(P(q^7)) = 2\eta$. With $P(X) = X - 1 - 3t^\eta$ the exceptional index is $n = 3$, where $P(q^3) = 3t^{2\eta} + t^{3\eta}$. The exceptions are harmless for the escape argument, since it needs only an eventual range, but they cannot be ignored in a statement asserted for all indices. This is why Lemma 4.1 and Theorem 4.4 both allow finitely many exceptions even when $\text{res}(q) = 1$.

For an ordinary complex q with $|q|_\mathbb{C} \neq 1$, the complex modulus does not alter these conclusions: as an element of the Hahn coefficient field, q still has valuation zero. The ordinary modulus and the scale valuation answer different questions.

5 Differential rigidity

5.1 From an operator to a forward recurrence

A formal series is D-finite exactly when its derivatives span a finite-dimensional space over $K_\Gamma(z)$. The D-finite/P-recursive equivalence is classical [1, Theorem 1.5]; the following records the exact finite conversion, because the valuations of the resulting recurrence, not ordinary coefficient growth, are what the proof consumes.

Lemma 5.1 (Differential coefficient recurrence). *Write*

$$L = \sum_{r=0}^R \sum_{k=0}^m c_{r,k} z^k D_z^r, \quad c_{r,k} \in K_\Gamma, \quad (5.1)$$

with at least one $c_{r,k} \neq 0$, and let $f = \sum_{n \geq 0} a_n z^n$. For $n \geq m$ the coefficient of z^n in Lf is

$$\sum_{s=-m}^R Q_s(n) a_{n+s}, \quad Q_s(X) = \sum_{r-k=s} c_{r,k} (X+s)_r, \quad (5.2)$$

where $(Y)_r = Y(Y-1)\cdots(Y-r+1)$ and $(Y)_0 = 1$. At least one Q_s is nonzero. Writing s_0 for the least active shift and $u = n + s_0$, every formal solution of $Lf = 0$ satisfies, for all sufficiently large u ,

$$\sum_{j=0}^s P_j(u) a_{u+j} = 0, \quad P_j(X) = Q_{s_0+j}(X - s_0), \quad P_0 \neq 0, \quad (5.3)$$

with $s = R - s_0$. If $s = 0$, every formal solution is a polynomial.

Proof. The operator $z^k D_z^r$ carries the term of index $n + r - k$ to degree n and multiplies it by $(n+s)_r$ with $s = r - k$, which gives (5.2). For a fixed shift s the distinct derivative orders r contribute polynomials of distinct degrees in X , so a nonempty group containing a nonzero coefficient cannot cancel identically and some Q_s is nonzero. Reindexing gives (5.3). If $s = 0$ the recurrence reads $P_0(u) a_u = 0$ with $P_0 \neq 0$, so by Lemma 4.1 all but finitely many a_u vanish. $\square$

Proposition 5.2 (Uniform obstruction for P-recursive coefficients). *Suppose (a_n) satisfies (5.3) for all $n \geq N_1$, with $P_j \in K_\Gamma[X]$ and $P_0 \neq 0$. Let $\beta_j = \beta(P_j)$ for the nonzero P_j and put*

$$B = \max(\{0\} \cup \{\beta_0 - \beta_j : j \geq 1, P_j \neq 0\}) \geq 0.$$

There is $N \geq N_1$, beyond the finitely many exceptional integers of Lemma 4.1 for the finitely many nonzero P_j , such that for every $x \neq 0$ with $v(x) \leq -B$, strong summability of $(a_n x^n)_n$ forces $a_n = 0$ for all $n \geq N$. If Γ is divisible one may replace B by the smaller threshold

$$B_{\text{rec}} = \max\left(\left\{\frac{\beta_0 - \beta_i}{j} : j \geq 1, P_j \neq 0\right\}\right), \quad (5.4)$$

which is (3.4) with period $m = 1$. If the displayed set is empty, take $B_{\text{rec}} = 0$; then every solution terminates beyond N without an evaluation hypothesis.

Proof. Discard the identically zero P_j . By Lemma 4.1 there is N beyond which $v(P_j(n)) = \beta_j$ for all the finitely many nonzero P_j simultaneously. Apply Corollary 3.4 with this B , or Corollary 3.5 with $m = 1$ when Γ is divisible. If no $j \geq 1$ remains, the recurrence forces the same vanishing directly. $\square$

These are certified exterior obstructions, not a computation of the exact evaluation domain of each solution. Individual solutions can have stricter restrictions or can terminate much earlier.

Proof of Theorem A. Lemma 5.1 converts L into (5.3). Take $B_L = B$ from Proposition 5.2 and take N_L to be an positive integer beyond the recurrence's starting index and beyond every nonnegative integer root of the finitely many residue polynomials $\overline{P_j}$. If a formal solution is strongly evaluable at one $x \neq 0$ with $v(x) \leq -B_L$, then $a_n = 0$ for all $n \geq N_L$, so its degree is less than N_L . An entire solution is in particular strongly evaluable at the available monomial t^{-B_L} , hence is a polynomial. Conversely a polynomial is entire and is annihilated by a sufficiently high derivative, which gives the displayed equality. $\square$

The closed region $v(x) \leq -B_L$, rather than an open one, comes from the weak inequality in Theorem 3.3, hence from Lemma 2.2. A polynomial solution of degree at least N_L would in any case contradict the recurrence at its last nonzero coefficient, since $P_0(n)$ no longer vanishes there.

Theorem A is stronger than merely excluding a globally extended exponential. The operator coefficients may themselves be arbitrary Hahn series carrying infinite surreal scales. There are nevertheless only finitely many of them, so their leading valuations give a finite obstruction that one additional exterior scale defeats.

5.2 An exact boundary, not only a nonexistence result

Proposition 5.3 (Formal exponential domain). *For $\eta \in \Gamma$ let*

$$E_\eta(z) = \sum_{n \geq 0} \frac{t^{n\eta}}{n!} z^n.$$

Then $D_z E_\eta = t^\eta E_\eta$ and

$$\text{Dom}_{\text{str}}(E_\eta) = \{0\} \cup \{x \in K_\Gamma^\times : v(x) > -\eta\}. \quad (5.5)$$

Proof. The differential identity is a formal coefficient comparison. Put $y = t^\eta x$ for $x \neq 0$, so the n th evaluation term is $y^n/n!$.

If $v(y) > 0$, write $y = ct^{v(y)}(1+u)$ with $c \in \mathbb{C}^\times$ and u zero or of positive valuation, and put $S = \text{supp}(u) \subseteq \Gamma_{>0}$. Since $(1+u)^n$ is a finite integer power, $\text{supp}(y^n) \subseteq h_n + M(S)$ with $h_n = nv(y)$,

and (h_n) is strictly increasing, so its values are well ordered and each is taken once. Lemma 2.3 (b) makes the family strongly summable; the factor $1/n!$ is a nonzero complex scalar and changes no support.

If $v(y) < 0$ the leading exponents $nv(y)$ strictly decrease. If $v(y) = 0$, then $y^n/n!$ has the nonzero coefficient $\text{lc}(y)^n/n!$ at exponent zero for every n , so local finiteness fails. These cases exhaust $x \neq 0$, and $v(y) > 0$ is $v(x) > -\eta$. $\square$

The recurrence of E_η is $(n+1)a_{n+1} - t^\eta a_n = 0$, with one active shift $j = 1$, so $\alpha_0 = \eta$, $\alpha_1 = 0$, and (3.4) gives $B_{\text{rec}} = \eta$ with no division by an integer at all. Comparing with (5.5): the excluded region $v(x) \leq -\eta$ of Corollary 3.5 is here *exactly* the complement of the domain, boundary included. The general exclusion bound can therefore be attained, and the closed form of the boundary is not an artefact of a lossy estimate. This example is the reason for keeping Lemma 2.2 rather than the strictly descending version alone.

There is no conflict with any scalar exponential on a surreal field: E_η is an ordinary power series in z , not a total exponential evaluated by an independent construction. Taking $\eta = 0$ recovers the elementary observation (2.3).

5.3 Algebraic and system consequences

Corollary 5.4 (Algebraic entire series). *An element of $\mathcal{E}_\Gamma$ algebraic over $K_\Gamma(z)$ belongs to $K_\Gamma[z]$.*

Proof. Let $F = K_\Gamma(z, f)$ be the finite extension generated by the algebraic series. In characteristic zero its minimal polynomial is separable, and differentiating that polynomial expresses $D_z f$ inside F , because the derivative with respect to the algebraic variable is nonzero at f and hence invertible in F . So D_z preserves F , all derivatives of f lie in the finite-dimensional $K_\Gamma(z)$ -space F , and f is D-finite. Apply Theorem A. $\square$

This is not offered as a new algebraic rigidity theorem; it records that the linear result subsumes the algebraic case without any factorization theorem for $\mathcal{E}_\Gamma$.

Corollary 5.5 (Rational differential systems). *If a finite vector Y of strongly entire formal series satisfies $D_z Y = A(z)Y$ with $A \in M_r(K_\Gamma(z))$, every component of Y is a polynomial. The same holds for an inhomogeneous system with a rational forcing vector, provided all its solution components are strongly entire.*

Proof. The $K_\Gamma(z)$ -span of the components is stable under D_z , so the successive derivatives of each component span a finite-dimensional space. For $Y' = AY + b$ append the constant component 1 and use $\begin{pmatrix} A & b \\ 0 & 0 \end{pmatrix}$. $\square$

Proposition 5.6 (Rational inhomogeneous equations). *The polynomiality conclusions of this report remain valid when the right-hand side of a linear equation is a rational function of z , provided the corresponding homogeneous hypothesis on L or on q holds.*

Proof. Clear denominators in the coefficients and in the right-hand side. The result has a polynomial right-hand side, so the coefficient recurrence is homogeneous for all sufficiently large indices, and the escape statements were formulated for eventual recurrences. The left-hand operator is still nonzero, having been multiplied by a nonzero polynomial. $\square$

This applies, for instance, to the inhomogeneous first-order theta identity (8.2) when the parameter valuation is noncofinal: it is not necessary to homogenize it in order to see the obstruction.

Example 5.7 (A sparse recurrence does not escape the obstruction). The formal Airy-type equation $f'' - zf = 0$ has the recurrence

$$a_{n+3} = \frac{a_n}{(n+3)(n+2)} \quad (n \geq 0), \quad a_2 = 0.$$

With $a_0 = 1$ and $a_1 = 0$ there are infinitely many nonzero coefficients, at indices $3n$, all of valuation zero. At $x = t^{-\delta}$ their leading exponents are $-3n\delta$, strictly decreasing for every $\delta > 0$. Gaps between nonzero coefficients do not help: the escape chain follows only the surviving congruence class.

Remark 5.8 (What transcendence has actually been proved). Every nonpolynomial member of $\mathcal{E}_\Gamma$ is transcendental over $K_\Gamma(z)$ and satisfies no nonzero *linear* differential equation over $K_\Gamma(z)$, and no finite family $f, D_z f, \dots, D_z^R f$ is linearly dependent over $K_\Gamma(z)$. This does not imply differential transcendence in the nonlinear sense. A relation

$$P(z, f, D_z f, \dots, D_z^R f) = 0$$

involving products of f and its derivatives lies outside the present proofs unless it can separately be reduced to a finite linear system. The distinction is not cosmetic: the escape mechanism exploits a finite *linear* cancellation in which one nonzero coefficient forces a later nonzero coefficient, whereas nonlinear coefficient equations involve convolutions of many earlier coefficients and supply no such forward step without further work.

That further work is the nonlinear part of this report (Sections 14–17, from source 10). It does not repair the escape chain; it replaces the recurrence by a finite initial polynomial at one large scale. Every relation of order $R \leq 1$ is thereby excluded (Theorem D), and so is every relation of higher order whose residue corner polynomial is nonzero (Theorem 16.1). Relations of order at least two with a vanishing residue corner remain outside the proofs of Sections 14–17. When Γ has no order unit they are excluded in every order by a third, different argument, which bounds the field generated by the Taylor coefficients (Theorem E, sources 11 and 12). With an order unit they are not always excluded: the series $\mathcal{T}_p$ of Theorem H (source 13) is a nonpolynomial strongly entire solution of a relation of order three with vanishing residue corner. Relations of order two are excluded for every Γ by a fourth argument, through initial polynomials at infinitely many scales (Theorem I, source 14), so three is the least order.

6 Generic lines and several-variable rigidity

A single differential equation in several variables is not enough to force polynomiality: it could ignore most of the variables. The correct hypothesis is finite-dimensionality of the span of *all* mixed partial derivatives, which is what Theorem C assumes.

Lemma 6.1 (Avoiding countably many polynomial conditions). *For any countable family of nonzero polynomials in $K_\Gamma[z_1, \dots, z_d]$ there is $c \in \mathbb{C}^d$ at which none of them vanishes.*

Proof. For one polynomial H , take the least Hahn exponent among its finitely many nonzero coefficients; the coefficient at that exponent is a nonzero $\overline{H} \in \mathbb{C}[z_1, \dots, z_d]$, and $\overline{H}(c) \neq 0$ implies

$H(c) \neq 0$. The coefficients of the countably many polynomials $\overline{H}$ form a countable subset of $\mathbb{C}$, and the field they generate over $\mathbb{Q}$ is countable. Choose $c_1, \dots, c_d$ algebraically independent over that field, successively outside the algebraic closure of a countable field, which remains countable while $\mathbb{C}$ is uncountable. $\square$

The same countable-field device is used in the companion report held in `docs/surcomplex/entire-functions-at-arbitrary-rank/ (ent:mv:thm:exceptional)` to show that the set of polynomial directions of a nonpolynomial series is not all of projective space.

Proof of Theorem C. Suppose F is not a polynomial and write it by total degree,

$$F(z) = \sum_{n \geq 0} F_n(z), \quad F_n \text{ homogeneous of degree } n, \quad (6.1)$$

so infinitely many F_n are nonzero. Let $g_1, \dots, g_\sigma$ be a basis of the finite-dimensional $K_\Gamma(z_1, \dots, z_d)$ -span of all mixed derivatives of F , taken to be mixed derivatives themselves with $g_1 = F$. The span is stable under differentiation, so there are rational matrices A_i with

$$\partial_i g = A_i(z)g \quad (1 \leq i \leq d). \quad (6.2)$$

These matrices have finitely many nonzero denominator polynomials; choose one nonzero homogeneous component of each. Apply Lemma 6.1 to every nonzero F_n , to those finitely many chosen components, and to the coordinate polynomials z_i , obtaining $c \in \mathbb{C}^d$ with $F_n(c) \neq 0$ whenever $F_n \neq 0$ and with no denominator becoming the zero polynomial under $z = wc$. Then

$$h(w) = F(wc) = \sum_{n \geq 0} F_n(c)w^n$$

is nonpolynomial.

For each $x \in K_\Gamma$ the family $(a_\alpha c^\alpha x^{|\alpha|})_{\alpha \in \mathbb{N}^d}$ is strongly summable, by entireness of F at $(c_1 x, \dots, c_d x)$. Group it into the finite blocks of fixed total degree; by the last clause of Lemma 2.3, $\sum_n F_n(c)x^n$ is strongly summable, so $h \in \mathcal{E}_\Gamma$.

The formal chain rule, applied after clearing denominators in (6.2), gives

$$\frac{d}{dw} g(wc) = \left(\sum_{i=1}^d c_i A_i(wc) \right) g(wc). \quad (6.3)$$

Every denominator here is a nonzero element of $K_\Gamma[w]$. It may vanish at $w = 0$, which is harmless: read the identity in the Laurent series field $K_\Gamma((w))$, with matrices in $M_\sigma(K_\Gamma(w))$. The finite $K_\Gamma(w)$ -span of the components of $g(wc)$ is preserved by d/dw , so its first component h is D-finite over $K_\Gamma(w)$, and Theorem A forces h to be a polynomial. This contradicts its construction. $\square$

Corollary 6.2 (Multivariate algebraic rigidity). *If a strongly entire $F \in K_\Gamma[[z_1, \dots, z_d]]$ is algebraic over $K_\Gamma(z_1, \dots, z_d)$, it is a polynomial.*

Proof. Each partial derivation extends to the finite separable extension generated by F , by differentiation of its minimal polynomial, so all mixed partial derivatives lie in one finite-dimensional space. Apply Theorem C. $\square$

The generic line is an existence device, not a promised effective choice of ordinary complex constants. The proof does not say that one prescribed line detects every nonpolynomial multivariate series; the line is chosen only after the countably many nonzero homogeneous parts and the finitely many rational connection matrices are known.

7 Dilation equations and the necessity of a cofinal scale

7.1 Exact coefficient conversion

Lemma 7.1 (Dilation coefficient recurrence). *Let $q \in K_\Gamma^\times$ have infinite multiplicative order and let $f = \sum a_n z^n$ satisfy (1.3). Write $p_\ell(z) = \sum_k p_{\ell,k} z^k$ and $Q_k(X) = \sum_\ell p_{\ell,k} X^\ell$, let d and e be the largest and smallest k with $Q_k \neq 0$, and put $s = d - e$. Then for every $n \geq 0$*

$$\sum_{h=0}^s R_h(q^n) a_{n+h} = 0, \quad R_h(X) = Q_{d-h}(q^h X), \quad (7.1)$$

with $R_0 = Q_d \neq 0$ and $R_s = Q_e(q^s \cdot) \neq 0$. Each nonzero R_h satisfies $R_h(q^n) \neq 0$ for all but finitely many n . If $s = 0$, every formal solution is a polynomial.

Proof. The coefficient of z^{n+d} in $\sum_\ell p_\ell(z) f(q^\ell z)$ is $\sum_{k=e}^d Q_k(q^{n+d-k}) a_{n+d-k}$; substituting $h = d - k$ and using $Q_{d-h}(q^{n+h}) = Q_{d-h}(q^h \cdot q^n)$ gives (7.1). Since q has infinite order the points q^n are distinct, and a nonzero univariate polynomial vanishes at only finitely many of them; this already gives eventual nonvanishing, independently of the finer valuation results. If $s = 0$ the relation reads $R_0(q^n) a_n = 0$ for every n , so all but finitely many a_n vanish. $\square$

The powers q^h in (7.1) matter: dropping them changes the recurrence and can change its valuation constants. Equivalently, taking $m = \max_\ell \deg p_\ell$ and reading off the coefficient of z^{n+m} gives $\sum_j P_j(q^n) a_{n+j} = 0$ with $P_j(X) = \sum_\ell p_{\ell, m-j} q^{\ell j} X^\ell$ and $P_0 = Q_m \neq 0$; both conversions are checked on finite exact examples by the programs of Section 18. If $m = 0$ the equation already reads $P_0(q^n) a_n = 0$ and all solutions are polynomials with no entireness assumption at all, so we may assume $s \geq 1$ below.

7.2 The valuation-zero case

Corollary 7.2 (Every unit dilation is globally rigid). *Let $q \in K_\Gamma^\times$ have $v(q) = 0$ and infinite multiplicative order. Every linearly q -difference finite member of $\mathcal{E}_\Gamma$ is a polynomial, and each fixed equation (1.3) has a uniform exterior obstruction and a uniform degree bound. If Γ is in addition divisible, every nonpolynomial formal solution has its domain bounded as in (3.5) by the constant (3.4) formed from the eventual residue-class valuations of (7.1).*

Proof. Apply Theorem 4.4 simultaneously to the finitely many nonzero R_h of Lemma 7.1. Their valuations take only finitely many values after a common starting index, with common period one or the exact order h_0 of $\text{res}(q)$. All differences $v(R_0(q^n)) - v(R_h(q^n))$ are therefore bounded above by a single element of Γ , and Corollary 3.4 gives the exterior obstruction and, via the residue polynomials, the degree bound. When Γ is divisible, Corollary 3.5 applies with $m = 1$ if the residue has infinite order, and $m = h_0$ otherwise. In both cases take the starting index beyond the exceptional range. $\square$

This includes parameters arbitrarily close to a root of unity without being one. The finite exceptional cancellations of Theorem 4.4 can move the starting index but do not remove the bound.

7.3 Nonzero but noncofinal valuation

Proposition 7.3 (Noncofinal dilation scales force polynomiality). *Let $q \in K_\Gamma^\times$ with $\beta = v(q) \neq 0$ and suppose $|\beta|$ is not an order unit of Γ . Every linearly q -difference finite member of $\mathcal{E}_\Gamma$ is a polynomial, with a uniform exterior obstruction and a uniform degree bound for each fixed equation. Divisibility of Γ is not used.*

Proof. By Lemma 4.2 applied to the finitely many nonzero R_h of (7.1), there is a common index beyond which

$$v(R_h(q^n)) = \alpha_h + k_h n \beta, \quad k_h \in \mathbb{N}.$$

Put $V = |\beta| > 0$, choose $B \in \Gamma$ with $B \geq 0$ and $B \geq |\alpha_0 - \alpha_h|$ for all relevant h , and put $M = \max_h |k_0 - k_h| \in \mathbb{N}$. Then for $h \geq 1$

$$v(R_0(q^n)) - v(R_h(q^n)) = \alpha_0 - \alpha_h + (k_0 - k_h)n\beta \leq B + MnV. \quad (7.2)$$

Because $|\beta|$ is not an order unit, (2.2) supplies $\gamma > 0$ with $\gamma > NV$ for every $N \in \mathbb{N}$; in particular $MnV < \gamma$ for all n , trivially so if $M = 0$. Hence the right side of (7.2) is bounded above by the single element $B + \gamma$. Corollary 3.4 with this bound excludes every $x \neq 0$ with $v(x) \leq -(B + \gamma)$, and the explicit witness $x = t^{-(B+\gamma)}$ is available. Since an entire solution is evaluable there, it satisfies $a_n = 0$ beyond the common starting index, hence is a polynomial of degree less than that index. Nothing in this argument divides an element of Γ by an integer. $\square$

The proof covers two different higher-rank situations. The group may have no order unit at all, so every nonzero dilation valuation is noncofinal; alternatively the group may have an order unit while the chosen q lives on a smaller scale. Global q -difference finiteness depends on the valuation of that particular dilation, not only on the group. The argument also requires no asymptotic formula for $v(a_n)$ and no eventually quadratic coefficient valuation: a single infinite escape chain suffices, which is why cancellations and sparse zero coefficients do not force a separate Skolem–Mahler–Lech analysis.

7.4 An intrinsic coarsened exclusion bound

The previous proof produces more than one bad point. Let $\beta = v(q) \neq 0$, let $\Delta = \Delta_{|\beta|}$ be the convex subgroup (2.1), and assume $\Delta \neq \Gamma$. Let $\pi : \Gamma \rightarrow \Gamma/\Delta$ be the ordered quotient and put $w = \pi \circ v$. With the eventual constants α_h of the previous proof, put

$$\overline{B}_{\text{rec}}^\flat = \max(\{0\} \cup \{\pi(\alpha_0) - \pi(\alpha_h) : 1 \leq h \leq s, R_h \neq 0\}), \quad (7.3)$$

and, when Γ is divisible so that Γ/Δ is a divisible ordered group,

$$\overline{B}_{\text{rec}} = \max_{1 \leq h \leq s, R_h \neq 0} \frac{\pi(\alpha_0) - \pi(\alpha_h)}{h}. \quad (7.4)$$

Theorem 7.4 (Coarsened exclusion). *Let $f = \sum a_n z^n$ be a nonpolynomial formal solution of (1.3) with $v(q) \neq 0$ and $\Delta = \Delta_{|v(q)|}$ proper.*

- (a) *Under Convention 1.1 alone, every $x \neq 0$ with $-w(x) > \overline{B}_{\text{rec}}^\flat$ lies outside $\text{Dom}_{\text{str}}(f)$.*
- (b) *If Γ is in addition divisible, every $x \neq 0$ with $-w(x) > \overline{B}_{\text{rec}}$ lies outside $\text{Dom}_{\text{str}}(f)$.*

Proof. Coarsening by Δ kills $n\beta$, since $n\beta \in \Delta$ for every n , so the projected valuations of the recurrence coefficients are the constants $\pi(\alpha_h)$. In case (b), $-w(x) > \overline{B}_{\text{rec}}$ gives $hw(x) < -(\pi(\alpha_0) - \pi(\alpha_h))$ for every relevant $h \geq 1$. In case (a) we have $w(x) < -\overline{B}_{\text{rec}}^b \leq 0$, hence $hw(x) \leq w(x) < -\overline{B}_{\text{rec}}^b \leq -(\pi(\alpha_0) - \pi(\alpha_h))$. Either way the projected leading valuation along an escape chain strictly decreases at every step. A strict decrease in the ordered quotient is a strict decrease in Γ , so the chain produces an infinite strictly descending sequence in the union of the supports. $\square$

The boundary is deliberately *not* excluded here. Equal leading values in the quotient can correspond to a strictly increasing sequence in Γ , and hence to a strongly summable family; Section 8.5 exhibits exactly that. Unlike Corollary 3.5, a coarsened comparison alone cannot decide local finiteness at an exponent of Γ .

8 Partial theta series: sharp witness and exact domain

8.1 The classical identities

For $p \in K_\Gamma^\times$ with $\beta = v(p) > 0$ set

$$\Theta_p(z) = \sum_{n \geq 0} p^{\binom{n}{2}} z^n = \sum_{n \geq 0} p^{n(n-1)/2} z^n. \quad (8.1)$$

This is a standard partial theta series up to a harmless sign convention [5]. None of its coefficients vanishes, so it is not a polynomial. The coefficient identity $p^{n(n-1)/2} = p^{n-1} p^{(n-1)(n-2)/2}$ for $n \geq 1$ gives

$$\Theta_p(z) = 1 + z\Theta_p(pz), \quad (8.2)$$

and applying $\sigma_p - 1$ to $(1 - z\sigma_p)\Theta_p = 1$, with $\sigma_p z = pz\sigma_p$, yields the homogeneous equation

$$\Theta_p(z) - (1 + z)\Theta_p(pz) + pz\Theta_p(p^2z) = 0, \quad (8.3)$$

which is also obtained by substituting pz for z in (8.2) and subtracting the two inhomogeneous identities. The existence of these identities is not the new part. What matters is precisely where the series is strongly evaluable, for an arbitrary-rank value group and an arbitrary Hahn parameter.

8.2 Exact domain with arbitrary coefficient tails

Theorem 8.1 (Exact partial theta domain). *Let Γ be as in Convention 1.1 and let $p \in K_\Gamma^\times$ with $\beta = v(p) > 0$, and let Δ_β be as in (2.1). Then*

$$\begin{aligned} \text{Dom}_{\text{str}}(\Theta_p) &= \{0\} \cup \{x \in K_\Gamma^\times : v(x) \geq -N\beta \text{ for some } N \in \mathbb{N}\} \\ &= \{0\} \cup \{x \in K_\Gamma^\times : (v \bmod \Delta_\beta)(x) \geq 0\}. \end{aligned} \quad (8.4)$$

In particular $\Theta_p \in \mathcal{E}_\Gamma$ if and only if β is an order unit of Γ . No monomial hypothesis on p or on x is needed, and Γ need not be divisible.

Proof. Evaluation at 0 is finite. For $x \neq 0$ put $\delta = v(x)$ and write

$$p = ct^\beta(1+u), \quad x = dt^\delta(1+w), \quad c, d \in \mathbb{C}^\times,$$

where each of u, w is zero or has strictly positive valuation. Let $S = \text{supp}(u) \cup \text{supp}(w) \subseteq \Gamma_{>0}$ and $M = M(S)$. The n th evaluation term is

$$p^{(n)}x^n = c^{(n)}d^n t^{h_n} (1+u)^{(n)}(1+w)^n, \quad h_n = \binom{n}{2}\beta + n\delta,$$

and the last two factors are finite integer powers, so

$$\text{supp}(p^{(n)}x^n) \subseteq h_n + M, \quad v(p^{(n)}x^n) = h_n. \quad (8.5)$$

Suppose first that $\delta \geq -N\beta$ for some $N \in \mathbb{N}$. Then

$$h_{n+1} - h_n = n\beta + \delta \geq (n - N)\beta > 0 \quad (n > N),$$

so (h_n) is eventually strictly increasing. Its set of values is therefore well ordered, including the finite initial segment, and each value is taken by finitely many indices. Lemma 2.3(b) applied to (8.5) proves strong summability.

If no such N exists, then $\delta < -N\beta$ for every N , so $h_{n+1} - h_n = n\beta + \delta < 0$ for every n . By (8.5) the leading exponents are exactly h_n , and they form a strictly decreasing sequence, so strong summability fails by Lemma 2.2.

Finally, $\delta \geq -N\beta$ for some N says exactly that the image of δ in Γ/Δ_β is nonnegative: a nonnegative coset either is positive, whence $\delta > 0$, or is zero, whence $\delta \in \Delta_\beta$ and $\delta \geq -N\beta$ for some N ; a negative coset lies below every element of Δ_β . The last assertion is the case $\Delta_\beta = \Gamma$. $\square$

This proof controls the whole support, not only the leading valuation. Leading valuation formulas by themselves do not in general determine a Hahn evaluation domain, because hidden tails can matter; here the common positive monoid M is the extra certificate that makes the formula exact. Note also that the entire calculation is carried out in Γ : nothing is divided by an integer, and no real-valued surrogate for v is introduced.

Remark 8.2 (A second, shorter proof of the order-unit half). When $\mu = \beta$ is an order unit, entireness also follows from the cofinal-valuation criterion Lemma 2.3(a) alone, and the routes of sources 08 and 09 are both kept here because they are genuinely different. For $x \neq 0$ with $\alpha = v(x)$, the n th evaluated term has valuation

$$\frac{n(n-1)}{2}\mu + n\alpha. \quad (8.6)$$

Given a target $\beta' \in \Gamma$, choose ordinary nonnegative integers M', B' with $\alpha \geq -M'\mu$ and $\beta' \leq B'\mu$, which is possible exactly because μ is an order unit. Then (8.6) is at least $(\frac{n(n-1)}{2} - M'n)\mu$, which exceeds $B'\mu$ for all sufficiently large ordinary n . So the term valuations tend cofinally to $+\infty$ and Lemma 2.3(a) applies. Only leading term valuations are used, so unrestricted higher Hahn tails in p do not invalidate the conclusion. This shorter route is available only in the order-unit case: at the boundary $v(x) = -N\beta$ with β noncofinal, the term valuations increase strictly but stay trapped below a fixed element of Γ , so criterion (a) says nothing and the monoid certificate is essential.

Remark 8.3 (The order-unit case has an optional rank-one picture). When β is an order unit, the real-valued coarsening $\pi : \Gamma \rightarrow \mathbb{R}$ of [12, Lemma 9.1], normalized by $\pi(\beta) = 1$, gives the same field K_Γ a rank-one absolute value. For $x \neq 0$, its theta terms have coarsened valuations $\binom{n}{2} + n\pi(v(x))$, tending to $+\infty$. This is an optional normed-field picture of the order-unit case, not a map that replaces the Hahn exponents by their images in $\mathbb{R}$.

Such a coefficientwise replacement need not define a Hahn series. In $\mathbb{Q}H \oplus \mathbb{Q}\epsilon$ with H dominating all $n\epsilon$, the coarsening normalized by H kills ϵ . The valid series $\sum_{n \geq 0} t^{n\epsilon}$ would then place infinitely many coefficients at exponent zero. Coarsening a valuation retains the field and changes the value map; it is not an embedding into a complex-coefficient Hahn field with smaller exponent group. No proof here requires this optional coarsening.

8.3 Completion of the sharp criterion

Proof of Theorem B. Assume (i) and suppose (ii) fails. If $v(q) = 0$, Corollary 7.2 makes every entire solution a polynomial. If $v(q) \neq 0$ and $|v(q)|$ is not an order unit, Proposition 7.3 does the same. Either way (i) fails, which proves (i) $\Rightarrow$ (ii) and supplies the stated uniform obstruction and degree bound.

Assume (ii). If $v(q) > 0$ is an order unit take $p = q$: by Theorem 8.1, $\Theta_q \in \mathcal{E}_\Gamma \setminus K_\Gamma[z]$, and (8.3) is a nonzero linear q -difference equation with polynomial coefficients. If $v(q) < 0$ and $-v(q)$ is an order unit take $p = q^{-1}$, so $v(p) = -v(q) > 0$ is an order unit and $\Theta_{q^{-1}} \in \mathcal{E}_\Gamma$ by the same theorem. Substituting q^2z for z in (8.3), with $p = q^{-1}$ so that $pz \mapsto qz$ and $p^2z \mapsto z$, gives

$$\Theta_{q^{-1}}(q^2z) - (1 + q^2z)\Theta_{q^{-1}}(qz) + qz\Theta_{q^{-1}}(z) = 0, \quad (8.7)$$

which uses only nonnegative powers of σ_q and has polynomial coefficients that are not all zero. Both signs of $v(q)$ are covered. $\square$

The partial theta function is used as an established type of construction, not advertised as a newly discovered special function. Its role is precise: it realizes the positive side of a necessary and sufficient criterion, including nonmonomial q and arbitrary valuation rank.

8.4 Two scales in the same rank-two field

Example 8.4 (One field, opposite answers). Let $\Gamma = \mathbb{Q}e_0 \oplus \mathbb{Q}e_1$ with $e_1 \gg e_0 > 0$, ordered by the highest nonzero coordinate. Then e_1 is an order unit and e_0 is not.

For $q = t^{e_0}$, no nonpolynomial strongly entire series satisfies a nonzero linear q -difference equation. The series Θ_q itself is not entire: by Theorem 8.1 its exact domain is the valuation ring obtained by forgetting the e_0 coordinate, which contains every x with $v(x) = -Ne_0$ and excludes $x = t^{-e_1}$.

For $q = t^{e_1}$, Θ_q is strongly entire, though the value group still has rank two. The two conclusions occur in the *same* Hahn field: the distinction is the dilation scale, not the rank of the field.

Under $e_0 = 1$, $e_1 = \omega$ and $t^\gamma = \omega^{-\gamma}$, the first dilation is multiplication by ω^{-1} and the second by $\omega^{-\omega}$. Quadratic multiples of the first valuation never reach the second scale; quadratic multiples of the second dominate every scale in this workspace. A fuller table for the same group, including unit dilations and the torsion exception, appears in Section 11.3.

8.5 The coarse boundary really can be included

For the theta series the forward recurrence is $a_{n+1} = p^n a_n$, read off from (8.2), whose inhomogeneity affects only the index $n = 0$ and is removed by Proposition 5.6. Its two slots have valuations $n\beta$ and 0, both of which project to 0 modulo Δ_β , so the projected gap is zero and $\overline{B}_{\text{rec}} = \overline{B}_{\text{rec}}^b = 0$. Theorem 7.4 therefore excludes the points of negative coarsened valuation, while Theorem 8.1

includes *all* points of zero coarsened valuation. The strict inequality in the coarsened exclusion theorem cannot be replaced by a weak one in general. This is the exact mirror image of Section 5.2, where the uncoarsened bound was attained with its boundary excluded from the domain. The two boundaries behave differently because one comparison is made in Γ and the other in a quotient that has forgotten the scale doing the work.

9 Torsion parameters and why they are excluded

If $q^h = 1$ then $\sigma_q^h - 1$ annihilates *every* formal power series, so the unrestricted definition of linear q -difference finiteness is vacuous for such a parameter. This is not a technical inconvenience removable by a better proof; it is a genuine counterexample to dropping the nontorsion hypothesis in Theorem B.

The difficulty persists even after excluding operators with zero action. For $h > 1$ and

$$f(z) = z^r g(z^h),$$

one has $f(qz) = q^r f(z)$, and the operator $\sigma_q - q^r$ is not zero on all formal series when $h > 1$. A nonpolynomial entire g produces a nonpolynomial entire f , since evaluation of f at x is evaluation of g at x^h followed by multiplication by x^r . Finite-order dilations therefore have a real periodic-support exception and are not a limiting case of Theorem B. In particular “the residue of q is a root of unity” and “ q is a root of unity” are fundamentally different hypotheses: Example 4.6 is covered by the theorem and this section is not.

There is nevertheless a meaningful eigenspace statement.

Proposition 9.1 (Torsion covariance). *Let $q \in \mathbb{C}^\times$ have exact finite order m and let $0 \leq r < m$.*

- (a) *A formal series satisfies $f(qz) = q^r f(z)$ if and only if $f(z) = z^r G(z^m)$ for a unique $G \in K_\Gamma[[z]]$.*
- (b) *If $G \in \mathcal{E}_\Gamma$ then $f = z^r G(z^m) \in \mathcal{E}_\Gamma$.*
- (c) *The converse of (b) also holds, so $f \in \mathcal{E}_\Gamma$ if and only if $G \in \mathcal{E}_\Gamma$, without divisibility of Γ .*

Moreover, every root of unity of K_Γ lies in $\mathbb{C}$, so the hypothesis $q \in \mathbb{C}^\times$ loses no torsion element of the Hahn field. All parts use only Convention 1.1.

Proof. For (a), coefficient comparison gives $(q^n - q^r)a_n = 0$, so only degrees congruent to r modulo m occur, and G is read off uniquely.

For (b), evaluation of $z^r G(z^m)$ at x is x^r times the family for G at x^m , reindexed; multiplication by the fixed element x^r preserves strong summability by Lemma 2.1 (3), and at $x = 0$ the evaluation is finite.

For (c), suppose $f \in \mathcal{E}_\Gamma$ and let $y \in K_\Gamma^\times$. Set $\delta = v(y)$, $\gamma = \min(\delta, 0)$, and $x = t^\gamma$. Since $m \geq 1$, we have $v(x^m) = m\gamma \leq \delta$. Strong evaluation of f at x , restricted to the degrees $r + mn$ and multiplied by x^{-r} , gives strong evaluation of G at x^m . Lemma 2.4 then gives strong evaluation at y . At $y = 0$ evaluation is finite. No root of y is extracted.

Finally, let $\zeta \in K_\Gamma$ be a root of unity. Then $v(\zeta) = 0$; dividing by $\text{res}(\zeta)$ gives $1 + u$ with u zero or of positive valuation, and $(1 + u)^{m'} = 1$ for some $m' \geq 1$. If $u \neq 0$ then $(1 + u)^{m'} - 1$ has valuation $v(u)$ and leading coefficient $m' \text{lc}(u) \neq 0$ in characteristic zero, a contradiction. So $u = 0$ and $\zeta = \text{res}(\zeta) \in \mathbb{C}$. $\square$

The converse uses only that the values $m\gamma$ are coinital in Γ , as the explicit choice $\gamma = \min(\delta, 0)$ shows. Actual m th roots need not exist in the original Hahn field; this stronger algebraic requirement is unnecessary for the entireness equivalence.

10 An extension to mixed differential–dilation equations

For a nonzero noncofinal dilation valuation, derivatives do not repair the missing-scale problem. The extension is useful when a symbolic system allows both differentiation and rescaling of the argument.

Lemma 10.1 (Polynomial dependence on the index as well as the dilation). *Let $q \in K_\Gamma^\times$ with $\lambda = v(q) \neq 0$ and let $0 \neq P(U, X) \in K_\Gamma[U, X]$. There are $\beta \in \Gamma$, $k \in \mathbb{N}$ and $N \in \mathbb{N}$ with*

$$v(P(n, q^n)) = \beta + kn\lambda \quad (n \geq N).$$

Proof. Write $P(U, X) = \sum_k C_k(U)X^k$. By Lemma 4.1 each nonzero $C_k(n)$ has the fixed valuation $\beta_k = \beta(C_k)$ for all sufficiently large n . The finitely many summands then have valuations $\beta_k + kn\lambda$, and the pairwise-order argument of Lemma 4.2 produces a unique eventual minimizing index, whose term cannot cancel. $\square$

Theorem 10.2 (Mixed rigidity at a noncofinal nonzero scale). *Let Γ be as in Convention 1.1, let $q \in K_\Gamma^\times$ with $v(q) \neq 0$, and suppose $|v(q)|$ is not an order unit of Γ . If a strongly entire f satisfies a nonzero equation*

$$\sum_{r,\ell} p_{r,\ell}(z) D_z^r \sigma_q^\ell f(z) = 0, \quad p_{r,\ell} \in K_\Gamma[z], \quad (10.1)$$

where the sum is finite and the coefficients are not all zero, then f is a polynomial. Each fixed equation has a uniform exterior obstruction and a uniform degree bound.

Proof. Expand $p_{r,\ell}(z) = \sum_k c_{r,\ell,k} z^k$. With $s = r - k$, the term of coefficient index $n + s$ contributes to degree n the quantity $c_{r,\ell,k}(n + s) q^{\ell(n+s)} a_{n+s}$, so the coefficient recurrence has shift polynomials

$$P_s(U, X) = \sum_{r-k=s} \sum_{\ell} c_{r,\ell,k} q^{\ell s} (U + s) X^\ell, \quad (10.2)$$

evaluated at $(U, X) = (n, q^n)$. For fixed s and ℓ , distinct r give distinct degrees in U , so a nonempty group of nonzero coefficients cannot cancel identically and some P_s is nonzero. Reindex from the smallest active shift; replacing U by $U - s_0$ and X by $q^{-s_0} X$ preserves polynomiality and nonzeroness, so the leftmost coefficient is $A_0(n) = P'_0(n, q^n)$ for a nonzero bivariate P'_0 , and Lemma 10.1 makes it eventually nonzero.

Lemma 10.1 gives eventual affine valuations for all the nonzero coefficients, so their differences obey a bound $B + Mn|v(q)|$ exactly as in (7.2). Since $|v(q)|$ is not an order unit, (2.2) bounds this by one element of Γ , and Corollary 3.4 finishes the proof. $\square$

The hypothesis $v(q) \neq 0$ is essential to *this* proof. For $v(q) = 0$ the separate pure differential and pure dilation conclusions have been proved above, but they are not a proof of the mixed conclusion. When Γ has no order unit, the mixed conclusion at unit valuation, for nonlinear relations as well, is Theorem E (b) of source 12, proved by a different method. For every Γ , with or without an order unit, the linear mixed conclusion at unit valuation is Corollary 29.6 (source 18), which answers Question 19.4: Lemma 10.1 is replaced there by the eventually periodic valuations of $P(n, q^n)$ (Corollary 28.7), and the theorem above is a special case of Theorem 29.4.

11 Surreal workspaces that separate the theories

The results so far are stated in set-sized Hahn fields, which keeps the foundations transparent and allows a literal surreal reading without quantifying over a proper class as though it were a set.

11.1 The workspace without an order unit

Take

$$\Gamma_\infty = \bigoplus_{j \geq 0} \mathbb{Q}e_j, \quad 0 < e_0 \ll e_1 \ll e_2 \ll \cdots, \quad (11.1)$$

ordered so that the highest index with a nonzero coordinate determines the sign, equivalently $e_{j+1} > Ne_j > 0$ for all $j \geq 0$ and $N \geq 1$. Every element has only finitely many nonzero coordinates. One may identify e_j with the surreal ω^j .

The group has no order unit: if $\gamma > 0$ has largest active index j , then $e_{j+1} > n\gamma$ for every ordinary n . The sequence (e_n) is nevertheless cofinal in the whole group, so this is a countable cofinality that cannot be generated by repeatedly adding one positive element. This group and this distinction appear in the companion entire-function report [12] and are background here, not a new observation.

Corollary 11.1 (Universal nontorsion dilation rigidity). *Suppose Γ has no order unit. Then every nonpolynomial $f \in \mathcal{E}_\Gamma$ satisfies, simultaneously for all nontorsion $q \in K_\Gamma^\times$ and all $r \geq 0$,*

$$f(z), f(qz), \dots, f(q^r z) \quad \text{are linearly independent over } K_\Gamma(z).$$

Such fields can contain nonpolynomial strongly entire series, as (11.1) and Theorem 11.2 show.

Proof. A linear dependence over $K_\Gamma(z)$ becomes, after clearing denominators, a nonzero equation (1.3). Since Γ has no order unit, no $|v(q)|$ is an order unit, so Theorem B forces f to be a polynomial. $\square$

Theorem 11.2 (An explicit universally nonholonomic entire series). *In K_{Γ_∞} the series*

$$F(z) = \sum_{n \geq 0} t^{e_n} z^n = \sum_{n \geq 0} \omega^{-\omega^n} z^n \quad (11.2)$$

is strongly entire and nonpolynomial. It satisfies no nonzero linear differential equation over $K_{\Gamma_\infty}(z)$ and, for every nontorsion $q \in K_{\Gamma_\infty}^\times$, no nonzero polynomial-coefficient linear q -difference equation. For every q of nonzero valuation it satisfies no nonzero mixed equation (10.1) either.

Proof. Fix $x \neq 0$ and write $x = ct^\delta(1+w)$ with $c \in \mathbb{C}^\times$ and w zero or of positive valuation. Choose M with δ involving only the coordinates $e_0, \dots, e_M$. The n th evaluation term has leading exponent $h_n = e_n + n\delta$ and support contained in $h_n + M(\text{supp } w)$. For $n > M$ the highest coordinate of h_n is e_n with coefficient 1, so $h_n > 0$, and the highest coordinate of

$$h_{n+1} - h_n = e_{n+1} - e_n + \delta$$

is e_{n+1} with coefficient 1, so (h_n) is eventually strictly increasing. Lemma 2.3(b) proves strong summability; at $x = 0$ only the constant term remains. All coefficients are nonzero, so F is nonpolynomial.

Alternatively, for any $\beta \in \Gamma_\infty$ choose j larger than every active coordinate index of δ and of β ; for $n > j$ the largest active coordinate of $h_n - \beta$ is n with coefficient 1, so $h_n > \beta$, and Lemma 2.3 (a) applies for arbitrary w . Cofinal leading valuations control the whole support, since every exponent of a Hahn term is at least its leading exponent.

The differential conclusion is Theorem A. Since Γ_∞ has no order unit, Theorem B — equivalently Corollary 11.1 — excludes every nontorsion dilation equation, and Theorem 10.2 excludes the mixed equations at every q with $v(q) \neq 0$. Rational coefficient equations reduce to the polynomial-coefficient case by clearing denominators. $\square$

The universal quantifier over q matters. The parameter may be chosen after F is fixed and may itself have infinite support; each individual $v(q)$ still has finitely many coordinates in Γ_∞ , hence is noncofinal. So the theorem excludes a linear relation for *each* such parameter, not merely for an advance list of simple dilations. The series has a simple explicit coefficient rule, and this is a statement about finite linear annihilating equations, not a claim that the rule is uncomputable or lacks a finite description.

11.2 Why the same formula is not entire on all surcomplex numbers

Let $e_* = \omega^\omega$, which dominates every integer multiple of every positive element of Γ_∞ , and enlarge the group to $\Gamma_\infty + \mathbb{Q}e_*$. At the surcomplex point $x = \omega^{e_*} = t^{-e_*}$ the evaluated leading exponents are $e_n - ne_*$, whose successive differences $e_{n+1} - e_n - e_*$ are negative. The defining family is therefore not strongly summable, by Lemma 2.2.

This explicit obstruction prevents the fixed-workspace theorem from being misread as an assertion of a nonpolynomial power series entire on all of $\mathbf{No}[i]$. The example illustrates two independent restrictions: a workspace can be large enough that no single dilation scale is cofinal, yet small enough to admit a countable sequence of coefficient scales escaping all its arguments, and enlarging it further can destroy the very entireness on which the theorem operates. The companion report studies this kind of boundary in general [12]; the calculation here is included only to state the domain of the present example unambiguously, and no new universal scalar-extension theorem is claimed.

11.3 Two classification tables

Let $\Gamma = \mathbb{Q}H \oplus \mathbb{Q}\epsilon$ with H dominating every $n\epsilon$, so that H is an order unit and ϵ is not.

Dilation q	Its valuation	Nonpolynomial entire solution of a nonzero linear q -difference equation?
t^H	H	Yes: Θ_{t^H} , by (8.3).
t^{-H}	$-H$	Yes: Θ_{t^H} , by (8.7).
t^ϵ	ϵ	No: positive, but noncofinal.
$1 + t^H$	0	No: an infinite-order unit.
2	0	No: complex size does not change the Hahn valuation.
-1	0	Outside the theorem: a torsion parameter, with periodic-support exceptions.

Workspace group	Parameter	Solution exists?	Reason
$\mathbb{Q}$	$v(q) \neq 0$	Yes	Every positive nonzero scale is an order unit.
$\mathbb{Q}$	$v(q) = 0$, nontorsion	No	Unit-orbit valuations are eventually periodic.
$\mathbb{Q}e_0 \oplus \mathbb{Q}e_1$, $e_1 \gg e_0$	$v(q) = e_0$	No	A lower scale is not cofinal.
the same group	$v(q) = e_1$	Yes	The top scale is an order unit.
$\bigoplus_{n \geq 0} \mathbb{Q}e_n$	any nontorsion q	No	No scale is an order unit, although F is entire.

In every row, nonpolynomial strongly entire D-finite series are impossible. Nonpolynomial strongly entire differentially algebraic series exist exactly in the rows whose group has an order unit, that is, in all rows except the last row of the second table (Theorem H), and their least differential order is then three (Theorem I); this does not depend on the parameter. Allowing derivatives beside the powers of σ_q changes no answer in either table: a nonzero mixed equation in D_z and powers of σ_q has a nonpolynomial strongly entire solution exactly in the rows answered “yes” (Corollary 29.11, source 18); the torsion row stays outside that corollary, and there $\sigma_{-1} - 1$ already has nonpolynomial entire solutions (Section 9). Torsion parameters are deliberately absent from the second table: their annihilating operator may be an identity on every series. Finite valuation rank alone is therefore not the criterion, and the distinction persists for nonmonomial dilations, because the theorem uses the valuation rather than a requirement that q be a monomial.

Corollary 11.3 (Detecting order units by functional equations). *For a nonzero set-sized ordered abelian group Γ the following are equivalent:*

- (i) Γ has an order unit;
- (ii) some nontorsion $q \in K_\Gamma^\times$ admits a nonpolynomial strongly entire solution of a nonzero polynomial-coefficient linear q -difference equation.

Proof. Theorem B gives (ii) $\Rightarrow$ (i). For the converse take $q = t^\mu$ for an order unit μ ; then $v(q) = \mu$ is an order unit and q is nontorsion. $\square$

This recovers a property of the ordered group from a global functional equation existence question. Theorem H recovers the same property from algebraic differential equations, with no dilation. The workspace (11.1) shows why the existence of nonpolynomial entire functions is by itself strictly weaker than the existence of an order unit.

11.4 Several meanings of transcendence, kept apart

For the explicit F the established conclusions are: transcendence over $K_{\Gamma_\infty}(z)$ by Corollary 5.4; linear differential independence of every finite family of derivatives; and linear independence along every single nontorsion geometric dilation orbit. These are substantial restrictions, and they do *not* imply that no polynomial relation $P(z, F, D_z F, \dots, D_z^R F) = 0$ exists. Nonlinear differential equations lie outside these proofs, and a relation involving $F(q_1 z)$ and $F(q_2 z)$ for multiplicatively independent parameters is not covered merely by applying the single- q theorem twice.

The nonlinear part adds one more conclusion by a separate proof: F and $D_z F$ are algebraically independent over $K_{\Gamma_\infty}(z)$ (Corollary 17.2), so every nonlinear relation with $R \leq 1$ is excluded. The methods of that part exclude relations with $R \geq 2$ only when their residue corner polynomial is nonzero, and give no statement about several dilations. Because Γ_∞ has no order unit, the coefficient-field part removes both limits for this F : Theorem 22.1 (source 12) makes all $(D_z^j F)(\lambda z)$, for dilations λ with pairwise quotients that are not roots of unity, algebraically independent over

$K_{\Gamma_\infty}(z)$. In particular F satisfies no algebraic differential equation of any order, and no nonzero relation, linear or not, links $F(q_1 z)$ and $F(q_2 z)$ for multiplicatively independent q_1, q_2 (Corollary 22.8, a merge observation).

12 Coefficient fields, degenerate value groups, characteristic

The one-variable proofs work over any characteristic-zero coefficient field with trivial valuation: the root-of-unity argument uses only finitely many ordinary polynomial roots and needs no algebraic closedness, and the integer-evaluation lemma needs only residue characteristic zero. All theorems also hold over $\mathbb{R}((t^\Gamma))$, the generic-line proof of Theorem C using uncountability of $\mathbb{R}$ in place of uncountability of $\mathbb{C}$. The results therefore concern surreal as well as surcomplex workspaces. More generally the core versions work over $k((t^\Gamma))$ for any characteristic-zero field k , and, as Section 1.4 records, without divisibility of Γ .

No statement here is a theorem for arbitrary valued fields of positive residue characteristic. In that setting Lemma 4.1 fails and the valuations of near-torsion geometric powers can behave differently, so any comparison with the ultrametric literature must keep the coefficient field and the residue characteristic visible. Example 17.3, from source 10, shows that characteristic zero is needed even for the single linear equation $D_z f = 0$: over $\mathbb{F}_p((t^\mathbb{Q}))$ a nonpolynomial strongly entire series has zero derivative.

The nonlinear part (Convention 14.1) is proved by its source over an arbitrary field of characteristic zero. That is a statement about Sections 14–17 only; the generality of the linear theorems is governed by the first paragraph of this section and its own argument. The same holds for the coefficient-field part (Convention 20.1), whose sources also work over an arbitrary field of characteristic zero.

We assumed $\Gamma \neq 0$ in order to speak of positive exterior scales. When $\Gamma = 0$ the field is $\mathbb{C}$ with the trivial valuation, strong evaluation at $x = 1$ already permits only finitely many nonzero coefficients because every nonzero term has support $\{0\}$, and entire series are polynomials for that immediate support reason.

13 Quantitative certificates and symbolic representation

13.1 What one can extract from an annihilating equation

The classification has a direct representational consequence. A system whose nonpolynomial special functions are represented only by finite linear differential equations cannot cover the nonpolynomial strong entire functions of a Hahn workspace. Allowing linear dilation equations enlarges the class only when a cofinal dilation scale is available. In the no-order-unit example, an exact coefficient rule such as (11.2) lies outside both pure equation formats.

This does not rule out exact symbolic computation with such functions. It specifies what a faithful representation must keep separate: the coefficient-generation rule, the exponent group, the strong evaluation domain, and the type of finite equation being asserted. A field-independent label such as “entire theta function” discards relevant information; for Θ_p the proof certificate is not merely $v(p) > 0$, it includes that $v(p)$ is an order unit of the selected group.

For a differential operator the reduction yields a concrete certificate. Compute the finite shift polynomials Q_s of (5.2) and shift the least active index to zero. For each resulting P_j record its least coefficient valuation β_j and the corresponding nonzero complex residue polynomial $\bar{P}_j$. Record

an integer N beyond their nonnegative integer roots and beyond the recurrence's starting index. Record the threshold B of Proposition 5.2, or (5.4) when Γ is divisible. The escape statements then certify that every formal solution strongly evaluable at some $x \neq 0$ with $v(x) \leq -B$ has degree less than N .

For a valuation-zero dilation the same plan uses (7.1): if $\text{res}(q)$ has infinite order use the minimum coefficient valuation, and if it has order h use (4.4)–(4.6) on each residue class. For a nonzero-valuation dilation, finitely many affine comparisons select the eventual winner of each $R_h(q^n)$; project their constants modulo $\Delta_{|v(q)|}$ to obtain the coarsened bound (7.3), or (7.4) when Γ is divisible. When the parameter scale is noncofinal one can instead display $\gamma \gg |v(q)|$ and the explicit bad argument $t^{-(B+\gamma)}$ of Proposition 7.3.

These are mathematical certificate schemas relative to exact coefficient data and order comparisons. They are *not* a terminating algorithm for arbitrary encoded surreals. Deciding whether an arbitrary complex residue is a root of unity, comparing arbitrary surreal values, extracting arbitrary supports, or deciding whether an abstract scale is cofinal can require information not effectively available in a chosen representation, and none of this has been made computable by the proofs.

13.2 A valuation recurrence with zero coefficients allowed

Example 13.1 (Jump lengths, not just jumps). Suppose a sequence that is not eventually zero satisfies

$$t^A a_n + (n-2)t^B a_{n+1} + t^C a_{n+3} = 0 \quad (n \geq 0),$$

with $A, B, C \in \Gamma$. The middle coefficient vanishes at $n = 2$, a finite exceptional index; beyond it the three valuations are A, B, C . The escape chain may move forward by one or by three, and no assumption that all a_n are nonzero is needed. Corollary 3.4 excludes the region $v(x) \leq -\max\{A - B, A - C, 0\}$ under Convention 1.1 alone. If Γ is *divisible*, Corollary 3.5 gives the sharper excluded region

$$v(x) \leq -\max\left\{A - B, \frac{A - C}{3}\right\}.$$

The maximum must be divided by the actual jump length: replacing every jump by one need not preserve the intended threshold, especially when the valuation gaps have different signs. This is a worked instance of item (1) of Section 1.4.

13.3 Current Lean coverage and remaining dependency order

The ledger `docs/FORMALIZATION.md` records the formalized escape mechanism in `Surreal/HahnSeries/EscapeChain.lean`: the nonincreasing-order obstruction, escape step and infinite chain, evaluation exclusion, and the uniform bounded-cost corollary. It also covers the univariate strong-domain definitions and the formal exponential identity with its exact domain and attained exclusion boundary. Part (2) of Lemma 2.1 is mapped separately to `Surreal/HahnSeries/NeumannWords.lean`.

`Surreal/HahnSeries/PartialThetaDomain.lean` covers both sufficient criteria of Lemma 2.3, the scale-subgroup facts, the exact partial-theta domain and the theta identities. It also proves (ii) $\Rightarrow$ (i) of Theorem B and (i) $\Rightarrow$ (ii) of Corollary 11.3; the converse implications remain pending.

`Surreal/HahnSeries/UnitOrbitPolynomials.lean` covers integer valuation stabilization, eventually affine geometric-orbit valuations, the unit-orbit theorem and the bivariate index–dilation

lemma. `Surreal/Algebra/GenericConstantPoint.lean` proves Lemma 6.1, including its least-exponent reduction, for uncountable constants. `Surreal/HahnSeries/TorsionCovariance.lean` covers inward stability, all of Proposition 9.1, and the preservation half of Lemma 2.1 (3). These results require no divisibility of the value group.

The remaining dependencies include the refined periodic threshold, part (1) and the reflection half of part (3) of Lemma 2.1, finite-block grouping in Lemma 2.3, Proposition 2.7, and the several-variable strong-domain definition. The operator-to-recurrence conversions and full differential and dilation classifications remain pending. For the several-variable classification, constant-point avoidance is proved; generic-line restriction, restriction of a rational differential system, and the resulting D-finiteness argument are pending.

The nonlinear rigidity theorems in Sections 14–17, the coefficient-field theorem package in Sections 20–22, the theta-descent part in Sections 23–25, the order-three-threshold part in Sections 26–27, and the unit-dilation, Newton-rigidity, theta-hierarchy, single-scale, polynomial-composition and recurrence parts in Sections 28–39, whose suggested formalization orders are recorded in Sections 29.6, 33.3 and 37.4 and Questions 19.35, 19.53 and 19.29 (with its batch-33 status), remain unformalized; the shared support and inward-stability prerequisites above do not prove them. For the coefficient-field package source 12 suggests the order: valuation-rank inequality, proper-convex-subgroup obstruction, formal coefficient recurrence, all-order differential corollary; the mixed statements would also need a formal theorem implying the zero-set lemma for nondegenerate exponential polynomials. For the order-three-threshold part source 14 suggests two nearly independent layers: finite algebra (dehomogenization, the logarithmic Euler identities, constants of rational differentiation in characteristic zero and the two endpoint evaluations, which give the finite endpoint lemma without surreal infrastructure), then the Hahn layer (the all-scale criterion, transport through the divisible hull, finite active-index sets and the estimate (26.4)). The build and axiom audit cover only imported declarations recorded in the ledger. The Python checks are finite checks of formulas and examples; they do not establish any remaining infinite statement.

14 Nonlinear equations: finite initial polynomials

This section and the next three contain the material of source 10. They answer the first-order case of Question 19.1, which sources 08 and 09 had left open, together with a nondegenerate higher-order class and positive-weight Euler equations in several variables. The method is not the escape chain of Section 3. The warning in Remark 5.8 stands: a nonlinear equation gives coefficient convolutions, not a bounded-step linear recurrence. Instead, a whole scaled series is reduced to a finite polynomial at one large scale, and an exact leading-term computation gives the contradiction.

14.1 Coefficient field, conventions and the merge’s choices

Convention 14.1 (Coefficient field of the nonlinear part). In Sections 14–17, k is an arbitrary field of characteristic zero with the trivial valuation, Γ is as in Convention 1.1, and

$$\mathbb{K} = k((t^\Gamma)).$$

The valuation v , leading coefficient lc , residue res , strong summability, $\text{Dom}_{\text{str}}(f)$ and entireness are defined exactly as in Section 1.2 and Definition 1.2, with k in place of $\mathbb{C}$, except that, following source 10, $\text{res}(a)$ is defined for every a of nonnegative valuation as its coefficient at exponent zero, so that $\text{res}(a) = 0$ when $v(a) > 0$. Write $\mathcal{E}_\Gamma(k)$ for the strongly entire series in $\mathbb{K}[[z]]$. In

several variables, strong entireness of $f = \sum_{\alpha} a_{\alpha} x^{\alpha}$ means strong summability of the *joint* family $(a_{\alpha} b^{\alpha})_{\alpha \in \mathbb{N}^m}$ at every $b \in \mathbb{K}^m$. For $k = \mathbb{C}$ one has $\mathbb{K} = K_{\Gamma}$ and $\mathcal{E}_{\Gamma}(k) = \mathcal{E}_{\Gamma}$. The derivative D_z kills $\mathbb{K}$, and $f' = D_z f$; no scalar surreal derivation, analytic continuation or fine-topology limit is involved.

Why the coefficient field differs. Source 10 proves every statement of the nonlinear part over an arbitrary characteristic-zero field k , and it uses neither algebraic nor real closedness. Its hypothesis on the coefficient field is therefore weaker than the hypothesis $k = \mathbb{C}$ under which Theorems A–C and the rest of Sections 1–13 are stated. This generality is recorded for the nonlinear part only. It is *not* transferred to the linear theorems: they remain stated over $\mathbb{C}$, and their possible extension to other coefficient fields is the separate remark of Section 12, resting on its own argument. In the other direction, the nonlinear part borrows from Section 2 only facts about well-ordered subsets of Γ and finite coefficient sums (Lemmas 2.1 and 2.3), whose proofs do not use any property of $\mathbb{C}$; source 10 proves the facts it needs over k itself, except the positive-support lemma used in Theorem 15.10, which it imports as standard. The group Γ is not assumed divisible, and no order unit, no countable cofinal subset and no real-valued norm is used anywhere in the nonlinear part.

Renamed symbols. Several symbols of source 10 would collide with symbols of Sections 1–13; they are renamed here, in the new material only.

Source 10	Here	Reason
$D = \max_{\nu} d_{\nu}, W$	d_P, w_P	D_z is the derivative.
H_P (residue corner polynomial)	χ_P	H is an element of Γ in Sections 2–11.
B_P (degree cutoff, in $\mathbb{Q}$)	κ_P	B, B_L, B_{rec} are elements of Γ .
B (set of right sides)	$\mathcal{R}$	The same.
r, r_0 (scale, “radius”)	δ, δ_0	r indexes derivative orders in (1.2); $t^{-\delta}$ is the notation of Corollary 3.4.
h_r, p_r (Gauss value, initial polynomial)	$\gamma_{\delta}, \phi_{\delta}$	h is a torsion order, p_r an operator coefficient.
M (an integer bound)	n_{*}	$M(S)$ is a support monoid.
E_q (Euler operator)	ϑ_q	E_{η} is the formal exponential of Proposition 5.3.
Theorems A, B, C	Theorems D, 16.5, 16.1	Theorems A–C of this report are different theorems.

The shipped audit files `10-nonlinear-rigidity-PROOF_AUDIT.md` and `10-nonlinear-rigidity-SOURCES_AND_SCOPE.md` keep the source’s notation and numbering.

Words with two meanings. *Escape* keeps the meaning of Section 3 (a chain through a linear recurrence); source 10’s lemma “escape of active degree” is a different statement and is called Lemma 14.5 here. An *exclusion bound* is, as in Section 3, an element $\delta_0 \in \Gamma$ such that evaluation fails at the arguments $t^{-\delta}$ with $\delta \geq \delta_0$; the source’s word “radius” is not used. A *corner* is always the finite set of terms of Section 14.4, never a full differential Newton polygon. The *residue polynomials* $\bar{P}$ and R_r of Section 4 are unrelated to the residue corner polynomial χ_P .

Results printed once. Source 10 reproves several facts that Sections 2 and 11 already contain. They are printed once and cited: the support calculus inside Lemma 14.2 (Lemma 2.1 (1),(3) and Proposition 2.7); its “cofinal escape of coefficient minima” lemma, which is Lemma 2.3 (a); the

positive-support geometric-series lemma, which source 10 imports and Lemma 2.1 (2) proves; the surreal embedding, which is Section 2.4 with the same monomial convention $t^\gamma \mapsto \omega^{-\gamma}$; strong entireness of the series (11.2), which is Theorem 11.2; and the remark on $\Gamma = 0$, which is in Section 12. Everything else in source 10 is printed in Sections 14–17, with its proofs.

14.2 The coefficient-family algebra

For a finite tuple $X = (X_1, \dots, X_m)$ of indeterminates put

$$\mathcal{S}_m = \left\{ F = \sum_{\alpha} b_{\alpha} X^{\alpha} \in \mathbb{K}[[X]] : (b_{\alpha})_{\alpha} \text{ is strongly summable} \right\},$$

and let $\mathcal{S}_m^+$ be the subset whose coefficients all have nonnegative valuation. The condition concerns the family of coefficients, not convergence in the X -adic topology.

Lemma 14.2 (Hahn-family algebra and its residue). *$\mathcal{S}_m$ is a $\mathbb{K}$ -subalgebra of $\mathbb{K}[[X]]$, stable under the formal partial derivatives and under the Euler operators $\sum_{\nu} q_{\nu} X_{\nu} \partial_{\nu}$ with $q_{\nu} \in k$. Coefficientwise residue is a ring homomorphism*

$$\rho : \mathcal{S}_m^+ \longrightarrow k[X], \quad \rho\left(\sum_{\alpha} b_{\alpha} X^{\alpha}\right) = \sum_{\alpha} \text{res}(b_{\alpha}) X^{\alpha},$$

whose values are polynomials, not arbitrary formal power series; ρ commutes with these differential operators.

Proof. Closure under sums, scalar multiples and products is the support calculation of Lemma 2.1 (1) and (3), as in the proof of Proposition 2.7. For strongly summable (b_{α}) and (c_{ϵ}) a fixed exponent has finitely many representations as a sum of two support exponents, each carried by finitely many indices, so the double family $(b_{\alpha} c_{\epsilon})$ is strongly summable, and regrouping by $\alpha + \epsilon$ gives the Cauchy product: regrouping can remove supported exponents by cancellation, but cannot introduce new ones or new contributors at a fixed exponent. A scalar is a one-member Hahn family, so the same argument gives scalar multiplication; a finite union handles addition. Source 10 proves this over k by the same two-coordinate sequence argument; like the Hahn-field construction itself, that argument uses ordinary set-theoretic choice. A formal derivative or an Euler operator with coefficients in k reindexes a subfamily and multiplies its members by elements of the trivially valued field k ; it creates no new support exponent. For $F \in \mathcal{S}_m^+$ the exponent zero lies in only finitely many coefficient supports, so $\rho(F)$ is a polynomial. Each X -coefficient of a product is a finite sum of products of coefficients, the residue is additive and multiplicative on elements of nonnegative valuation, and the local finiteness just proved justifies every interchange; so ρ is a ring homomorphism. Differentiation commutes with residue coefficientwise. $\square$

For $0 \neq F = \sum_{\alpha} b_{\alpha} X^{\alpha} \in \mathcal{S}_m$ put

$$\gamma(F) = \min_{b_{\alpha} \neq 0} v(b_{\alpha}), \quad \text{in}(F) = \rho(t^{-\gamma(F)} F).$$

The minimum exists because it is the least element of the nonempty well-ordered union of supports. Then $t^{-\gamma(F)} F \in \mathcal{S}_m^+$ and the *initial polynomial* $\text{in}(F)$ is nonzero.

Corollary 14.3 (Gauss multiplicativity). *For nonzero $F, G \in \mathcal{S}_m$,*

$$\gamma(FG) = \gamma(F) + \gamma(G), \quad \text{in}(FG) = \text{in}(F) \text{in}(G).$$

Proof. Normalize both factors and apply Lemma 14.2. The product of the two nonzero residues is nonzero because $k[X]$ is an integral domain, so some coefficient of $t^{-\gamma(F)-\gamma(G)}FG$ has valuation exactly zero, and none has negative valuation. This proves the asserted minimum. Taking the residue of that normalization then gives the initial-polynomial identity. $\square$

14.3 Scaled initial polynomials

Let $f = \sum_{n \geq 0} a_n z^n \in \mathbb{K}[[z]]$ and $\delta \in \Gamma$ with $t^{-\delta} \in \text{Dom}_{\text{str}}(f)$. Then

$$F_\delta(X) = f(t^{-\delta}X) = \sum_{n \geq 0} a_n t^{-n\delta} X^n \in \mathcal{S}_1.$$

For $f \neq 0$ put

$$\gamma_\delta = \min_{a_n \neq 0} (v(a_n) - n\delta), \quad g_\delta = t^{-\gamma_\delta} F_\delta, \quad \phi_\delta = \rho(g_\delta), \quad N_\delta = \deg \phi_\delta. \quad (14.1)$$

The *active* indices are the n attaining γ_δ ; there are finitely many, N_δ is the largest, and $A_\delta = \text{lc}(\phi_\delta) \in k^\times$. More explicitly,

$$\phi_\delta(X) = \sum_{v(a_n) - n\delta = \gamma_\delta} \text{lc}(a_n) X^n.$$

Here only nonzero a_n are included. Each active index contributes at the same Hahn exponent γ_δ in the scaled coefficient family, so local finiteness makes this sum finite. Distinct indices remain distinct powers of X ; their nonzero leading coefficients cannot cancel as a polynomial.

The element γ_δ is a *coefficient Gauss value*. It need not equal $v(f(t^{-\delta}))$: $\phi_\delta(1)$ can vanish although $\phi_\delta \neq 0$. The proofs below use identities in $\mathcal{S}_1$ and in $k[X]$ only, never an identification of these two quantities, and never replace the initial polynomial by its value at 1. For example, with $\eta > 0$ the finite family $F(X) = 1 - X + t^\eta X^2$ has $\gamma(F) = 0$ and $\text{in}(F) = 1 - X$, but $F(1) = t^\eta$ has valuation $\eta > 0$.

Lemma 14.4 (Exact derivative scaling). *For every $j \geq 0$,*

$$(D_z^j f)(t^{-\delta}X) = t^{\gamma_\delta + j\delta} g_\delta^{(j)}(X), \quad \rho(g_\delta^{(j)}) = \phi_\delta^{(j)}. \quad (14.2)$$

If $N_\delta \geq j$, then $\phi_\delta^{(j)}$ has degree $N_\delta - j$ and leading coefficient $A_\delta(N_\delta)_j$.

Proof. The scaling identity is a formal coefficient computation: $D_X^j F_\delta(X) = t^{-j\delta} (D_z^j f)(t^{-\delta}X)$. The residue identity is Lemma 14.2. The falling factorial is a nonzero element of k for $N_\delta \geq j$ because k has characteristic zero. $\square$

Lemma 14.5 (Large active degrees). *Let f be nonpolynomial, let $n_* \in \mathbb{N}$, and choose $m > n_*$ with $a_m \neq 0$. There is $\delta_1 \in \Gamma_{>0}$ such that, whenever $\delta \geq \delta_1$ and $t^{-\delta} \in \text{Dom}_{\text{str}}(f)$, every active index exceeds n_* . Moreover $\gamma_\delta \leq v(a_m) - m\delta$.*

Proof. Choose $\delta_1 > 0$ with

$$(m - n)\delta_1 > v(a_m) - v(a_n) \quad (0 \leq n \leq n_*, a_n \neq 0). \quad (14.3)$$

There are finitely many inequalities, and they need no division in Γ : take δ_1 positive and larger than the absolute values of all right-hand sides, since each multiplier $m - n$ is a positive integer. Such an

element exists: add any positive element of the nonzero ordered group to the maximum of zero and those finitely many absolute values. This also covers an empty list of low-index coefficients. The inequalities persist for $\delta \geq \delta_1$ and say that each such low-index term has scaled valuation $v(a_n) - n\delta$ strictly larger than that of the m th term. The last assertion is the definition of γ_δ . $\square$

This is not a limit $N_\delta \rightarrow \infty$ along a countable cofinal sequence, which need not exist; it concerns a final segment of $\Gamma_{>0}$ and only those scales at which strong evaluation is available. It is also unrelated to the escape chain of Lemma 3.1: no recurrence is used.

14.4 The finite corner of a differential polynomial

Let P be a nonzero differential polynomial of order at most s , written in combined-monomial form

$$P(z, Y_0, \dots, Y_s) = \sum_{\nu \in J} c_\nu z^{k_\nu} \prod_{j=0}^s Y_j^{m_{\nu j}}, \quad (14.4)$$

with J finite, every $c_\nu \in \mathbb{K}^\times$, and identical monomials already combined. Write $P[f] = P(z, f, D_z f, \dots, D_z^s f)$. Define the integers

$$d_\nu = \sum_{j=0}^s m_{\nu j}, \quad w_\nu = k_\nu - \sum_{j=0}^s j m_{\nu j}, \quad d_P = \max_{\nu} d_\nu, \quad w_P = \max_{d_\nu = d_P} w_\nu, \quad (14.5)$$

and

$$\mathcal{C}_P = \{\nu : d_\nu = d_P, w_\nu = w_P\}, \quad \beta_P = \min_{\nu \in \mathcal{C}_P} v(c_\nu), \quad \bar{c}_\nu = \text{res}(t^{-\beta_P} c_\nu) \in k \quad (\nu \in \mathcal{C}_P).$$

Thus (d_P, w_P) is chosen lexicographically: first the largest total differential degree, then the largest shift w . This is a finite corner selection, not the construction of a differential Newton polygon. The *residue corner polynomial* and the *full corner polynomial* are

$$\chi_P(T) = \sum_{\nu \in \mathcal{C}_P} \bar{c}_\nu \prod_{j=0}^s (T)_j^{m_{\nu j}} \in k[T], \quad I_P(T) = \sum_{\nu \in \mathcal{C}_P} c_\nu \prod_{j=0}^s (T)_j^{m_{\nu j}} \in \mathbb{K}[T]. \quad (14.6)$$

In general χ_P can vanish although some $\bar{c}_\nu \neq 0$, because different products of falling factorials can be linearly dependent. If $\chi_P \neq 0$, then $I_P \neq 0$, and

$$I_P(n) = 0 \implies \chi_P(n) = 0 \quad (n \in \mathbb{N}), \quad (14.7)$$

because $t^{-\beta_P} I_P$ has coefficients of nonnegative valuation and residue commutes with substitution of an ordinary integer.

Lemma 14.6 (The exact corner identity). *Let $\phi \in k[X]$ have degree $N \geq s$ and leading coefficient $A \neq 0$, and put*

$$\Omega_\phi(X) = \sum_{\nu \in \mathcal{C}_P} \bar{c}_\nu X^{k_\nu} \prod_{j=0}^s (\phi^{(j)}(X))^{m_{\nu j}}.$$

Then $\deg \Omega_\phi \leq d_P N + w_P$ and

$$[X^{d_P N + w_P}] \Omega_\phi = A^{d_P} \chi_P(N). \quad (14.8)$$

In particular $\chi_P(N) \neq 0$ implies $\Omega_\phi \neq 0$.

Proof. Each corner summand has degree at most $k_\nu + \sum_j m_{\nu j}(N - j) = d_P N + w_P$. This common degree is a nonnegative integer because $N \geq s \geq j$ and $k_\nu, m_{\nu j} \geq 0$, even if $w_P < 0$. The coefficient of the summand in that degree is $\bar{c}_\nu A^{d_P} \prod_j (N)_j^{m_{\nu j}}$. Summing gives the formula. $\square$

Only finitely many nonnegative integers are roots of a nonzero χ_P , even when k is not algebraically closed. The active degree is an ordinary integer, not an element of Γ , and no root extraction or division in Γ occurs in this step.

15 First-order algebraic differential equations

15.1 A finite certificate excluding all large scales

The following theorem carries the proof of Theorem D and is also a practical audit object: its scale inequalities use only finitely many coefficients of the solution. It applies to equations of any order.

Theorem 15.1 (Nonlinear finite exclusion certificate). *Let P be as in (14.4) with $d_P \geq 1$ and $\chi_P \neq 0$, and let $f = \sum_n a_n z^n$ be a nonpolynomial formal solution of $P[f] = 0$. Choose an integer $n_* \geq s$ at least as large as every nonnegative integer root of χ_P , and choose $m > n_*$ with $a_m \neq 0$ and*

$$(d_P - d_\nu)m + w_P - w_\nu > 0 \quad (d_\nu < d_P). \quad (15.1)$$

Every $\delta \in \Gamma_{>0}$ satisfying the finitely many strict inequalities

$$(m - n)\delta > v(a_m) - v(a_n) \quad (0 \leq n \leq n_*, a_n \neq 0), \quad (15.2)$$

$$(w_P - w_\nu)\delta > \beta_P - v(c_\nu) \quad (d_\nu = d_P, w_\nu < w_P), \quad (15.3)$$

$$((d_P - d_\nu)m + w_P - w_\nu)\delta > \beta_P - v(c_\nu) + (d_P - d_\nu)v(a_m) \quad (d_\nu < d_P) \quad (15.4)$$

has $t^{-\delta} \notin \text{Dom}_{\text{str}}(f)$. Such m and δ exist, and the inequalities persist when δ increases. Consequently $\text{Dom}_{\text{str}}(f)$ omits $t^{-\delta}$ for every δ in a final segment of $\Gamma_{>0}$, and every strongly entire solution of $P[f] = 0$ is a polynomial.

Proof. An integer n_* exists because $\chi_P \neq 0$. For $d_\nu < d_P$ the coefficient $d_P - d_\nu$ of m in (15.1) is a positive integer, so all sufficiently large m satisfy it, and a nonpolynomial f has nonzero coefficients of arbitrarily large index. Every multiplier on the left of (15.2)–(15.4) is then a positive integer, so any δ that is positive and larger than the absolute values of all right-hand sides satisfies them, as does every larger δ .

Suppose $t^{-\delta} \in \text{Dom}_{\text{str}}(f)$ for such a δ , and form $\gamma = \gamma_\delta$, $g = g_\delta$, $\phi = \phi_\delta$, $N = N_\delta$ as in (14.1). Each inequality in (15.2) puts that low-index term strictly above the m th term in scaled valuation. Thus no index at most n_* is active, by the comparison in the proof of Lemma 14.5, and

$$N > n_*, \quad \gamma \leq v(a_m) - m\delta. \quad (15.5)$$

All derivatives of g lie in $\mathcal{S}_1^+$. Substituting $z = t^{-\delta}X$, a ring homomorphism $\mathbb{K}[[z]] \rightarrow \mathbb{K}[[X]]$, Lemma 14.4 turns the ν th summand of $P[f]$ into

$$c_\nu t^{d_\nu \gamma - w_\nu \delta} X^{k_\nu} \prod_j (g^{(j)})^{m_{\nu j}}.$$

Multiply the whole identity by $t^{-\lambda}$ with $\lambda = \beta_P + d_P\gamma - w_P\delta$. The scalar in front of the ν th product then has valuation

$$v(c_\nu) - \beta_P + (d_\nu - d_P)\gamma + (w_P - w_\nu)\delta. \quad (15.6)$$

For $\nu \in \mathcal{C}_P$ this is $v(c_\nu) - \beta_P \geq 0$. For $d_\nu = d_P$ and $w_\nu < w_P$ it is positive by (15.3). For $d_\nu < d_P$, multiplying the bound on γ in (15.5) by the negative integer $d_\nu - d_P$ reverses it and gives the lower bound

$$v(c_\nu) - \beta_P - (d_P - d_\nu)v(a_m) + ((d_P - d_\nu)m + w_P - w_\nu)\delta > 0$$

by (15.4). Every polynomial product of the $g^{(j)}$ has coefficients of nonnegative valuation. Multiplication by a positive-valuation scalar makes its residue zero, including when the product itself is zero. Thus the normalized identity lies in $\mathcal{S}_1^+$, and only corner summands can survive the residue map. Applying ρ to the formal identity gives

$$0 = \sum_{\nu \in \mathcal{C}_P} \bar{c}_\nu X^{k_\nu} \prod_j (\phi^{(j)})^{m_{\nu j}} = \Omega_\phi(X). \quad (15.7)$$

But $N > n_* \geq s$ and $\chi_P(N) \neq 0$, so by Lemma 14.6 the coefficient of $X^{d_P N + w_P}$ in Ω_ϕ is $\text{lc}(\phi)^{d_P} \chi_P(N) \neq 0$, a contradiction. The last assertion follows because a strongly entire series is evaluable at every monomial argument. $\square$

An explicit, division-free exclusion bound. Let $\mathcal{R}$ be the finite set of right-hand sides of (15.2)–(15.4); an empty list contributes nothing. For any $\epsilon \in \Gamma_{>0}$,

$$\delta_0 = \epsilon + \max(\{0\} \cup \{|b| : b \in \mathcal{R}\}) \quad (15.8)$$

satisfies all of them, since their multipliers are positive integers. This is an explicit certificate; it is not claimed to be optimal. The proof uses the full supports to form the initial polynomial but only finitely many leading valuations to choose δ_0 . Leading valuations alone do *not* in general decide strong evaluation at a boundary argument, and the certificate has no converse.

For $k = \mathbb{C}$, Lemma 2.4 (inward stability) turns the monomial conclusion into the exterior form used in Theorem A: a nonpolynomial formal solution is then not strongly evaluable at any $x \neq 0$ with $v(x) \leq -\delta_0$. This combination of the two statements is made by the merge; source 10 states only the monomial form.

Corollary 15.2 (Cofinal scales suffice). *If $\chi_P \neq 0$, a formal solution of $P[f] = 0$ strongly evaluable at $t^{-\delta}$ for a cofinal set of $\delta \in \Gamma_{>0}$ is a polynomial.*

Proof. If $d_P = 0$, then P is a nonzero polynomial in z alone and has no formal solution. Otherwise a cofinal subset meets the final segment excluded by Theorem 15.1 for a nonpolynomial solution. $\square$

Dependency audit. The only infinite-support step is Lemma 14.2, with the existence and finiteness of a leading coefficient layer; the rest is finite polynomial algebra. The proof makes none of the following replacements: a higher-rank group by a real-valued norm; strong summation by convergence of finite subsums; a nonsummable family by a cancelled expression; the residue polynomial by its value at 1; a nonlinear coefficient recurrence by a linear one. The three critical checkpoints are the minimum in (14.1), the order reversal when (15.5) is multiplied by $d_\nu - d_P < 0$, and the nonvanishing of $\chi_P(N)$.

15.2 Why the corner never degenerates in order one; proof of Theorem D

Write a first-order equation in combined-monomial form as

$$P(z, Y, Y') = \sum_{k,i,j} c_{kij} z^k Y^i (Y')^j. \quad (15.9)$$

For a term, $d = i + j$ and $w = k - j$. At the corner, fixing j forces $i = d_P - j$ and $k = w_P + j$, so there is at most one corner term for each j , and (14.6) becomes

$$\chi_P(T) = \sum_{\substack{i+j=d_P \\ k-j=w_P}} \bar{c}_{kij} T^j, \quad I_P(T) = \sum_{\substack{i+j=d_P \\ k-j=w_P}} c_{kij} T^j. \quad (15.10)$$

At least one $\bar{c}_{kij}$ is nonzero by the choice of β_P , and its power T^j cannot cancel against another summand. Hence $\chi_P \neq 0$.

Proof of Theorem D. If $d_P = 0$, then P is a nonzero polynomial in z alone and has no formal solutions. Otherwise the preceding observation and Theorem 15.1 give polynomiality and the exclusion statement, with δ_0 as in (15.8). A nonzero algebraic relation between f and $D_z f$ over $\mathbb{K}(z)$ becomes a nonzero polynomial of the form (15.9) after denominators are cleared. Multiplication by a common nonzero denominator preserves both the nonzero relation and its vanishing in $\mathbb{K}((z))$, even if that denominator vanishes at $z = 0$. Thus the polynomiality conclusion contradicts nonpolynomiality of f , proving the independence statement. $\square$

The uniqueness of a corner term for fixed j is a structural fact of order one. It is exactly what fails when several derivative orders can contribute the same falling-factorial polynomial (Section 16.2).

15.3 Finite candidate degrees and the entire-solution locus

Once polynomiality is known, ordinary polynomial degree gives a sharper finite candidate set that uses the *full* coefficients. Assume $d_P \geq 1$ and define the nonnegative rational number

$$\kappa_P = \max\left(\{0\} \cup \left\{ \frac{w_\nu - w_P}{d_P - d_\nu} : d_\nu < d_P \right\}\right) \in \mathbb{Q}. \quad (15.11)$$

This divides ordinary integers in $\mathbb{Q}$, not elements of Γ .

Theorem 15.3 (Finite candidate degrees). *For the first-order polynomial (15.9) with $d_P \geq 1$, every nonconstant strongly entire solution has degree in the finite set*

$$\mathcal{D}_P = \{n \in \mathbb{Z}_{>0} : n \leq \kappa_P\} \cup \{n \in \mathbb{Z}_{>0} : I_P(n) = 0\}. \quad (15.12)$$

The constant solutions are exactly the $c \in \mathbb{K}$ with $P(z, c, 0) = 0$ as a polynomial in z . The set (15.12) is necessary; it does not promise that each candidate occurs.

Proof. Polynomiality is Theorem D. Let $f = az^n + \dots$ with $n \geq 1$ and $a \neq 0$. A term of (15.9) has degree $k + in + j(n-1) = dn + w$ and leading coefficient $c_{kij} a^d n^j$. If $n > \kappa_P$, every term with $d < d_P$ has degree below $d_P n + w_P$, and so do the terms with $d = d_P$, $w < w_P$. So the coefficient of $z^{d_P n + w_P}$ in $P[f]$ is $a^{d_P} I_P(n)$, which must vanish. I_P is nonzero by (15.10), so its set of ordinary integer roots is finite. The assertion on constants is immediate by substitution. $\square$

Remark 15.4 (What “finite” does and does not compute). The set of ordinary integer roots of I_P is well defined and finite for arbitrary Hahn coefficients, but its existence is not a uniform algorithm for arbitrary coefficient names: extracting a leading exponent, deciding coefficient equality, or testing an arbitrary complex coefficient for exact integrality can need information that no numerical approximation supplies. For rational coefficients, or other exact representations with the needed operations, (15.12) is an actual finite algebraic search. The shipped program uses only exact rational data.

Replacing $I_P(n) = 0$ by $\chi_P(n) = 0$ gives a coarser necessary set, by (14.7). Keeping I_P can be genuinely sharper.

Example 15.5 (A resonance visible only in the first layer). Let $c = t^\eta$ with $\eta > 0$, $N \in \mathbb{Z}_{>0}$, and $P = zY' - (N + c)Y$. Then $\chi_P(T) = T - N$ but $I_P(T) = T - N - c$ has no ordinary nonnegative integer root. The only formal solution in $\mathbb{K}[[z]]$ is zero, since coefficient comparison gives $(n - N - c)a_n = 0$ for every n . A residue resonance alone is not a polynomial solution.

Corollary 15.6 (An algebraic entire-solution locus). *For a first-order equation P with $d_P \geq 1$ put $n_P = \max(\{0\} \cup \mathcal{D}_P)$. There is an explicitly defined affine algebraic set $\mathcal{V}_P \subseteq \mathbb{A}_{\mathbb{K}}^{n_P+1}$ whose $\mathbb{K}$ -points are in bijection with all strongly entire formal solutions of $P[f] = 0$. The same equations parametrize all strongly entire solutions over every Hahn-field extension $\mathbb{K} \hookrightarrow \mathbb{L} = k'((t^{\Gamma}))$ induced by an embedding of coefficient fields $k \hookrightarrow k'$ and an embedding of ordered groups $\Gamma \hookrightarrow \Gamma'$.*

Proof. With indeterminates $u_0, \dots, u_{n_P}$ put $F_u(z) = \sum_{n \leq n_P} u_n z^n$ and expand $P(z, F_u, F'_u) = \sum_{\ell} Q_{\ell}(u) z^{\ell}$ with $Q_{\ell} \in \mathbb{K}[u_0, \dots, u_{n_P}]$. Let $\mathcal{V}_P$ be the common zero locus of these finitely many Q_{ℓ} . Its points are exactly the polynomial solutions of degree at most n_P , each of which is strongly entire, and Theorem 15.3 shows there are no others. Under the stated extension the monomial degrees and κ_P are unchanged, and for an ordinary integer n the element $I_P(n)$ vanishes in $\mathbb{L}$ if and only if it vanishes in $\mathbb{K}$. Theorem D applies over $\mathbb{L}$, whose coefficient field again has characteristic zero and whose group is again nonzero, so the same degree bound and equations remain exhaustive there. $\square$

This is a finite-dimensional algebraic parametrization, not a claim that the number of solutions is finite, and not a moduli statement over coefficient rings with nilpotents. The extension class is exactly the one stated.

15.4 Worked first-order equations

All identities use the formal derivative D_z . For $k = \mathbb{C}$ they are statements about surcomplex Hahn workspaces; for $k = \mathbb{R}$ the real versions follow with the non-real constant roots omitted. Consider

$$D_z f = f^2 + p(z), \quad p \in \mathbb{K}[z]. \quad (15.13)$$

The corner is the term $-Y^2$, so $d_P = 2$, $w_P = 0$ and $I_P = -1$. If p has degree $e \geq 0$, then $\kappa_P = \max(0, e/2)$, so a nonconstant entire solution has degree at most $\lfloor e/2 \rfloor$. More precisely, a solution of degree $n \geq 1$ requires $\deg p = 2n$ and the leading coefficient of p to be minus the square of that of f , since the top term of f^2 cannot be cancelled by $D_z f$. These conditions are necessary only; the remaining coefficient equations must still hold.

Proposition 15.7 (Two exact Riccati classifications). *Over $\mathbb{K} = \mathbb{C}((t^{\Gamma}))$, the strongly entire solutions of $D_z f = f^2 + 1$ are exactly $f = i$ and $f = -i$, and the only strongly entire solution of $D_z f = f^2 + 1 - z^2$ is $f = z$.*

Proof. In the first equation the degree bound excludes nonconstant solutions, and a constant must satisfy $c^2 + 1 = 0$; in the field $\mathbb{K}$, which contains i , the roots are $\pm i$. In the second the degree bound is one and constants are impossible. For $f = az + b$ coefficient comparison gives $a^2 = 1$, $2ab = 0$ and $a = b^2 + 1$, so $b = 0$ and $a = 1$, and $f = z$ is verified directly. $\square$

The first equation is also `diff:cor:riccati` of the collection's differential-equations report [13], with the same two solutions, but there $\partial u = 1 + u^2$ is solved in $\mathbf{No}[i]$ for a derivation on surreal *scalars*. The two statements concern different derivations and different objects, and neither implies the other.

Example 15.8 (The degree cutoff is attained). For every integer $m \geq 1$ the equation $D_z f = f^2 + mz^{m-1} - z^{2m}$ has the strongly entire solution $f = z^m$. Here $I_P = -1$ and $\kappa_P = m$ exactly, so the low-degree part of (15.12) cannot in general be replaced by $n < \kappa_P$.

Example 15.9 (Arbitrarily large resonant degrees). For $N \geq 1$ the equation

$$zf D_z f - Nf^2 = 0 \quad (15.14)$$

has $d_P = 2$, $w_P = 0$ and $I_P(T) = T - N$. Since $\mathbb{K}[[z]]$ is an integral domain, a nonzero solution satisfies $zD_z f = Nf$, and coefficient comparison gives exactly $f = cz^N$ with $c \in \mathbb{K}$ (including $c = 0$). The exceptional integer root is an actual degree, and no degree bound can depend only on the positions of the monomials of P : the coefficient N matters.

15.5 The exclusion bound cannot be uniform

Theorem A has an exclusion bound depending only on the operator. For nonlinear equations this is false, already for one fixed equation of first order.

Theorem 15.10 (A fixed Riccati equation defeats every uniform bound). *For $\lambda \in \Gamma$ let $c = t^\lambda$ and*

$$f_\lambda(z) = \frac{c}{1 - cz} = \sum_{n \geq 0} c^{n+1} z^n \in \mathbb{K}[[z]].$$

Then $D_z f_\lambda = f_\lambda^2$, f_λ is nonpolynomial, and

$$\text{Dom}_{\text{str}}(f_\lambda) = \{0\} \cup \{x \in \mathbb{K}^\times : v(x) + \lambda > 0\}. \quad (15.15)$$

Consequently, for every $\delta \in \Gamma$ some nonpolynomial solution of the fixed equation $D_z f = f^2$ is strongly evaluable at $t^{-\delta}$, and no exclusion bound depending on the equation alone excludes all its nonpolynomial solutions.

Proof. The derivative identity is a coefficient comparison. For $x \neq 0$ put $u = cx$, so the n th evaluation term is cu^n . If $v(u) > 0$, write $u = bt^{v(u)}(1 + w)$ with $b \in k^\times$ and w zero or of positive valuation; then $\text{supp}(u^n) \subseteq nv(u) + M(\text{supp } w)$, the leading exponents $nv(u)$ increase strictly, and Lemma 2.3 (b), resting on the positive-support monoid of Lemma 2.1 (2), gives strong summability; multiplication by c preserves it. (Source 10 cites the Hahn–Neumann positive-support lemma here instead of proving it; no Archimedean convergence of $nv(u)$ is used.) If $v(u) = 0$, every cu^n has the nonzero coefficient $\text{lc}(u)^n$ at exponent λ , so local finiteness fails. If $v(u) < 0$, the leading exponents $\lambda + nv(u)$ strictly decrease. At $x = 0$ only one member is nonzero. This proves (15.15). Given δ , choose $\lambda = \delta + \epsilon$ with any $\epsilon \in \Gamma_{>0}$, available because $\Gamma \neq 0$; no order unit is required. Then $v(t^{-\delta}) + \lambda = \lambda - \delta = \epsilon > 0$. $\square$

There is no conflict with Theorem 15.1, whose finite data, and hence whose bound, may depend on the solution; for f_λ the exact boundary moves with λ . At the boundary $v(x) = -\lambda$ geometric-series algebra cannot legalize infinitely many contributions at one exponent. Nor is there a conflict with Theorem A: each f_λ also satisfies the linear equation $(1 - t^\lambda z)D_z f - t^\lambda f = 0$, whose operator depends on λ , exactly as the operator $D_z - t^\eta$ of the formal exponential of Proposition 5.3 depends on η . (This comparison is the merge's; it is a direct computation.)

16 Higher order and positive-weight Euler equations

16.1 The nondegenerate higher-order class

Theorem 16.1 (Higher-order corner criterion and its boundary). *Let P be a nonzero differential polynomial (14.4) of any finite order with $d_P \geq 1$ and $\chi_P \neq 0$. Then every strongly entire formal solution of $P[f] = 0$ is a polynomial, and every nonpolynomial formal solution fails strong evaluation at $t^{-\delta}$ for all δ in a final segment of $\Gamma_{>0}$, with the finite certificate of Theorem 15.1. The condition $\chi_P \neq 0$ holds automatically in order one, but not in order two: a fixed second-order equation can have polynomial solutions of every degree (Proposition 16.4).*

Proof. The first two assertions are Theorem 15.1; the third is Section 15.2 and Proposition 16.4. $\square$

Proposition 16.2 (Higher-order degree candidates). *Suppose $d_P \geq 1$ and $\chi_P \neq 0$ for an equation of order at most s . Every nonzero strongly entire solution has degree in the finite set*

$$\{0, \dots, s-1\} \cup \{n \in \mathbb{N} : n \geq s, n \leq \kappa_P\} \cup \{n \in \mathbb{N} : n \geq s, I_P(n) = 0\},$$

with κ_P as in (15.11). When $s = 0$ the first set is empty. Constant solutions can always be checked by substitution.

Proof. Polynomiality is Theorem 15.1. For degree $n \geq s$ every derivative leading-term formula of Lemma 14.6 applies, and if also $n > \kappa_P$ the coefficient of $z^{d_P n + w_P}$ in $P[f]$ is $\text{lc}(f)^{d_P} I_P(n)$, which must vanish; $I_P \neq 0$ because $\chi_P \neq 0$. Degrees below s , where derivative leading terms can vanish, are listed separately. $\square$

Proposition 16.3 (The first two Painlevé polynomial equations). *In characteristic zero, the equation $D_z^2 f = 6f^2 + z$ has no strongly entire formal solution, and the equation $D_z^2 f = 2f^3 + zf + \alpha$ with $\alpha \in \mathbb{K}$ has a strongly entire solution exactly when $\alpha = 0$, the only one then being $f = 0$.*

Proof. The first equation has $d_P = 2$, $w_P = 0$, $\chi_P = -6$; the second has $d_P = 3$, $w_P = 0$, $\chi_P = -2$. Theorem 15.1 makes every entire solution a polynomial. In the first equation a polynomial of degree $n \geq 1$ gives f^2 of degree $2n$, larger than $\deg D_z^2 f$ (when nonzero) and than $\deg z = 1$, so no cancellation is possible, and a constant cannot cancel z . In the second, for $n \geq 1$ one has $3n > n + 1$ and $3n > \deg D_z^2 f$, so $2f^3$ cannot cancel; a constant c must satisfy $2c^3 + zc + \alpha = 0$ identically, which forces $c = 0$ and $\alpha = 0$, and substitution proves sufficiency. $\square$

The names only identify the displayed equations; no classification of their classical meromorphic solutions is imported. The examples illustrate a class beyond first order: a unique term of greatest total differential degree has a nonzero corner whenever its falling-factorial product is nonzero. In particular an equation $D_z^s f = Q(z, f)$ with Q of degree at least two in f has a nonzero corner and hence only polynomial strongly entire solutions; the coefficient of the top power of f may be a nonconstant polynomial in z , whose highest z -term then selects the corner.

16.2 A vanishing corner

Consider

$$z^2 f D_z^2 f + z f D_z f - z^2 (D_z f)^2 = 0. \quad (16.1)$$

All three terms have $d = 2$ and $w = 0$, and the residue corner polynomial is

$$T(T - 1) + T - T^2 = 0, \quad (16.2)$$

an identity in $k[T]$, not an exceptional integer resonance.

Proposition 16.4 (Complete formal classification of the example). *The formal solutions of (16.1) in $\mathbb{K}[[z]]$ are exactly 0 and cz^n with $c \in \mathbb{K}^\times$ and $n \in \mathbb{N}$. All are strongly entire, and their degrees are unbounded for this single equation.*

Proof. With $\vartheta = zD_z$, equation (16.1) reads $f \vartheta^2 f - (\vartheta f)^2 = 0$. For $f \neq 0$ work in the Laurent series field $\mathbb{K}((z))$ and divide by f^2 : $\vartheta(\vartheta f/f) = 0$. The constants of ϑ in $\mathbb{K}((z))$ are exactly $\mathbb{K}$, since $\vartheta \sum b_n z^n = \sum n b_n z^n$ and characteristic zero kills no nonzero integer. Hence $\vartheta f/f = c_0 \in \mathbb{K}$; if the first nonzero term of f has exponent n , comparing lowest terms gives $c_0 = n$. Then $\vartheta f = n f$ kills every coefficient except the n th. Substitution verifies the monomials and zero. $\square$

The example has two messages. A universal higher-order degree bound cannot be obtained by declaring χ_P nonzero: it may vanish identically, and a fixed equation can then have unbounded degrees. And the example is *not* a nonpolynomial entire solution, so it does not refute polynomial rigidity in higher order; it locates a failure of this proof and of a stronger uniform conclusion.

The corner is also not an invariant of the solution set. Multiplying an equation by another differential polynomial, or passing to a differential consequence, can change the corner data. For a concrete example put

$$Q = z^2 Y_0 Y_2 + z Y_0 Y_1 - z^2 Y_1^2, \quad P = Y_1, \quad \tilde{P} = Y_1(1 + zQ).$$

For every formal f , the factor $1 + zQ[f]$ has constant coefficient one and is a unit of $\mathbb{K}[[z]]$. Hence $P[f] = 0$ and $\tilde{P}[f] = 0$ have exactly the same solutions, the constants. But $\chi_P(T) = T$, whereas the degree-three corner of $\tilde{P}$ gives $\chi_{\tilde{P}}(T) = T(T(T - 1) + T - T^2) = 0$. The criterion $\chi_P \neq 0$ is a sufficient condition on the displayed combined polynomial: when $\chi_P = 0$ another equation satisfied by the same function may have a nonzero corner, or a logarithmic-derivative reduction as in Proposition 16.4 may be available. A zero corner cannot be repaired by retaining only monomial degrees, since the residue coefficients and their exact cancellation are essential. The methods of this section do not decide whether a higher-order equation with vanishing residue corner has nonpolynomial strongly entire solutions. When Γ has no order unit, Theorem E (sources 11 and 12) decides it negatively in every order; Example 21.6 computes the nonzero linearization multiplier of (16.1) along its solutions. With an order unit the answer is different: the cleared third-order equation of source 13 has residue corner $\chi_P = 0$ and the nonpolynomial strongly entire solution $\mathcal{T}_p$ (Theorem H); its full corner polynomial I_P is a nonzero constant of positive valuation, so the hypothesis $\chi_P \neq 0$ of Theorems 15.1 and 16.1 cannot be weakened to $I_P \neq 0$ (Proposition 24.2 and Remark 24.3). In order two the vanishing-corner case is decided negatively for every Γ by source 14 (Theorem I(a)): its endpoint lemma uses the whole initial polynomial at infinitely many scales, and for (16.1) it permits no nonmonomial initial polynomial at all (Example 26.7).

16.3 Positive-weight Euler equations in several variables

Fix positive integers $q = (q_1, \dots, q_m)$, write $\text{wt}(\alpha) = q \cdot \alpha$ for $\alpha \in \mathbb{N}^m$, and let

$$\vartheta_q = \sum_{\nu=1}^m q_\nu x_\nu \frac{\partial}{\partial x_\nu}, \quad \vartheta_q x^\alpha = \text{wt}(\alpha) x^\alpha.$$

For $N \geq 0$, a multi-index of weight at most N satisfies $0 \leq \alpha_\nu \leq \lfloor N/q_\nu \rfloor$ in each coordinate. Thus at most $\prod_\nu (1 + \lfloor N/q_\nu \rfloor)$ multi-indices have that bound. This finiteness, with the eigenvalue identity, is the essential feature of positive weights. Write a nonzero equation in combined-monomial form

$$P(x, Y, Z) = \sum_{\nu \in J} c_\nu x^{\alpha_\nu} Y^{i_\nu} Z^{j_\nu}, \quad (16.3)$$

to be read as $P(x, f, \vartheta_q f) = 0$, and set $d_\nu = i_\nu + j_\nu$, $w_\nu = \text{wt}(\alpha_\nu)$, with d_P , w_P , $\mathcal{C}_P$, β_P and $\bar{c}_\nu$ as in Section 14.4. The corner polynomials now keep the spatial variables:

$$\chi(T, X) = \sum_{\nu \in \mathcal{C}_P} \bar{c}_\nu X^{\alpha_\nu} T^{j_\nu}, \quad I(T, X) = \sum_{\nu \in \mathcal{C}_P} c_\nu X^{\alpha_\nu} T^{j_\nu}. \quad (16.4)$$

Both are nonzero: for fixed (α_ν, j_ν) the condition $d_\nu = d_P$ forces $i_\nu = d_P - j_\nu$, so no two combined corner terms give the same monomial in (X, T) . Only finitely many integers n make $\chi(n, X)$ the zero polynomial, since they must all be roots of one nonzero X -coefficient of χ , and likewise for I .

Theorem 16.5 (Positive-weight Euler rigidity). *Let $f \in \mathbb{K}[[x_1, \dots, x_m]]$ satisfy $P(x, f, \vartheta_q f) = 0$ for a nonzero $P \in \mathbb{K}[x_1, \dots, x_m, Y, Z]$. If f is nonpolynomial, it fails joint strong evaluation at $(t^{-q_1\delta}, \dots, t^{-q_m\delta})$ for every sufficiently large $\delta \in \Gamma_{>0}$. Consequently, if f is strongly entire, or merely jointly strongly evaluable at these points for a cofinal set of $\delta \in \Gamma_{>0}$, then f is a polynomial. Every fixed equation has a finite set of possible positive weighted degrees (Corollary 16.6).*

Proof. If $d_P = 0$, then P is a nonzero polynomial in x alone and there is no solution; assume $d_P \geq 1$. Choose $n_* \geq 0$ with $\chi(n, X) \neq 0$ for every integer $n > n_*$. Since $f = \sum_\alpha a_\alpha x^\alpha$ is nonpolynomial and bounded-weight sets are finite, choose $a_\mu \neq 0$ of weight $n_\mu = \text{wt}(\mu) > n_*$ so large that $(d_P - d_\nu)n_\mu + w_P - w_\nu > 0$ for $d_\nu < d_P$. Take $\delta > 0$ satisfying the finitely many inequalities

$$\begin{aligned} (n_\mu - \text{wt}(\alpha))\delta &> v(a_\mu) - v(a_\alpha) & (\text{wt}(\alpha) \leq n_*, a_\alpha \neq 0), \\ (w_P - w_\nu)\delta &> \beta_P - v(c_\nu) & (d_\nu = d_P, w_\nu < w_P), \\ ((d_P - d_\nu)n_\mu + w_P - w_\nu)\delta &> \beta_P - v(c_\nu) + (d_P - d_\nu)v(a_\mu) & (d_\nu < d_P); \end{aligned}$$

as before they can be met in every nonzero ordered group and persist on a final segment. Suppose the joint family is strongly summable at such a scale, and put

$$F(X) = f(t^{-q_1\delta} X_1, \dots, t^{-q_m\delta} X_m) \in \mathcal{S}_m, \quad \gamma = \gamma(F), \quad g = t^{-\gamma} F, \quad \phi = \rho(g).$$

Then $\gamma \leq v(a_\mu) - n_\mu\delta$, and the first inequalities show that ϕ has no monomial of weight at most n_* ; its greatest weight N exceeds n_* . Let $\phi_N \neq 0$ be its weight- N homogeneous part. The Euler operator commutes with the scaling, so $(\vartheta_q f)(t^{-q_1\delta} X_1, \dots, t^{-q_m\delta} X_m) = t^\gamma \vartheta_q g$. Normalizing the substituted equation by $t^{-\beta_P - d_P\gamma + w_P\delta}$, the scalar computation (15.6) shows that only corner terms survive the residue, whence

$$0 = \sum_{\nu \in \mathcal{C}_P} \bar{c}_\nu X^{\alpha_\nu} \phi^{i_\nu} (\vartheta_q \phi)^{j_\nu}. \quad (16.5)$$

The largest possible weight on the right is $d_P N + w_P$. Since the weight- N part of $\vartheta_q \phi$ is $N\phi_N$, the homogeneous part of that weight is

$$\phi_N(X)^{d_P} \chi(N, X), \quad (16.6)$$

a product of two nonzero polynomials, hence nonzero. This contradicts (16.5). $\square$

The proof uses *joint* strong evaluation, not a scalar expression obtained by first setting $x_1 = \dots = x_m$ and cancelling: a factor $x_1 - x_2$ can make a diagonal scalar expression zero while the ungrouped monomial family is not summable. Keeping $X_1, \dots, X_m$ in (16.6) avoids that error.

Corollary 16.6 (Weighted-degree candidates). *With κ_P defined by (15.11) from the Euler data d_ν, w_ν , every nonconstant strongly entire solution of $P(x, f, \vartheta_q f) = 0$ has greatest weight in the finite set*

$$\{n \in \mathbb{Z}_{>0} : n \leq \kappa_P\} \cup \{n \in \mathbb{Z}_{>0} : I(n, X) \equiv 0\}.$$

For a fixed equation all strongly entire solutions are therefore described by finitely many polynomial equations in the coefficients of a bounded-weight polynomial ansatz.

Proof. Polynomiality is Theorem 16.5. For a polynomial of greatest weight $N > \kappa_P$, the terms of lower differential degree have smaller total weight, and the top-weight part of the equation is $f_N^{d_P} I(N, X)$, which must vanish; so $I(N, X) = 0$. The finite-root argument after (16.4) applies to I , and a bounded-weight ansatz has finitely many coefficients. $\square$

Example 16.7 (An exactly described solution space). For $N \geq 0$ the solutions of $\vartheta_q f = Nf$ in $\mathbb{K}[[x]]$ are exactly the weighted homogeneous polynomials of weight N , since coefficient comparison gives $(\text{wt}(\alpha) - N)a_\alpha = 0$. For $q = (1, 2)$ this space has dimension $\lfloor N/2 \rfloor + 1$ over $\mathbb{K}$, with basis $x_1^{N-2j} x_2^j$, $0 \leq j \leq \lfloor N/2 \rfloor$. The nonlinear equation $f(\vartheta_q f - Nf) = 0$ has the same solutions, because $\mathbb{K}[[x]]$ is an integral domain.

Why positive weights are needed. Let $\mathbb{K} = \mathbb{C}((t^{\mathbb{Q}}))$ and $G(u) = \sum_{n \geq 0} t^{n^2} u^n$, which is strongly entire (Corollary 17.1). The function $f(x, y) = G(y)$ is nonpolynomial and strongly entire but satisfies $x\partial_x f = 0$, so a zero weight invalidates the theorem. For the mixed weights $(1, -1)$, the nonpolynomial strongly entire function $f(x, y) = G(xy)$ satisfies $(x\partial_x - y\partial_y)f = 0$; at a fixed point its nonzero joint evaluation terms are exactly $t^{n^2} (xy)^n$, which are strongly summable. Positivity is therefore not a cosmetic substitute for nonzero weights, and a general first-order partial differential equation need not have the rigidity of a positive Euler operator.

Relation to Theorem C. Theorem C assumes that *all* mixed partial derivatives span a finite-dimensional space, and Section 6 notes that a single differential equation in several variables cannot force polynomiality in general, since it may ignore variables; the zero-weight example above is an instance. Theorem 16.5 instead assumes one algebraic relation between f and $\vartheta_q f$ with all weights positive, with no finite-dimensional-span hypothesis. These are different input criteria and different proof routes. Within the strongly entire class, each criterion forces a polynomial, which then satisfies both criteria: its mixed derivatives span a finite-dimensional space, and the relation $Y - f(x) = 0$ is an algebraic Euler relation.

17 Nonlinear consequences, hypotheses and predecessors

17.1 Two-jet independence of explicit series

With the surreal embedding of Section 2.4, for instance $\mathbb{C}((\omega^{-\mathbb{Q}}))$ is a set-sized surcomplex workspace to which Theorem D applies, and its coefficients and admissible arguments are actual elements of $\mathbf{No}[i]$. The independence conclusions, however, concern *formal functions*: each value $f(x)$ lies in $\mathbb{K}$ and is of course not transcendental over $\mathbb{K}$.

Corollary 17.1 (Two independent jets of the quadratic-exponent series). *Over $\mathbb{K} = \mathbb{C}((t^{\mathbb{Q}}))$ let $G(z) = \sum_{n \geq 0} t^{n^2} z^n$. Then G is nonpolynomial and strongly entire, and G and $D_z G$ are algebraically independent over $\mathbb{K}(z)$.*

Proof. For $x \neq 0$ with $v(x) = \sigma \in \mathbb{Q}$ the n th evaluated member has valuation $n^2 + n\sigma$, which eventually exceeds every rational bound, so Lemma 2.3 (a) gives strong summability even when x has an arbitrary infinite Hahn tail; at $x = 0$ the family is finite. All coefficients are nonzero. Theorem D gives the independence. $\square$

The series is not new: it is the rank-one example of the rank-one-Berkovich report [14], Example `found:ex:internal` of the foundations report [15], and a rescaled partial theta series: $G(z) = \Theta_{t^2}(tz)$ in the notation of (8.1). The contribution of source 10 is the nonlinear two-jet consequence only.

Corollary 17.2 (An example without an order unit). *In K_{Γ_∞} , with Γ_∞ as in (11.1), the strongly entire series $F = \sum_{n \geq 0} t^{e_n} z^n = \sum_{n \geq 0} \omega^{-\omega^n} z^n$ of (11.2) satisfies: F and $D_z F$ are algebraically independent over $K_{\Gamma_\infty}(z)$.*

Proof. Strong entireness is Theorem 11.2; source 10 proves it again by the cofinal-valuation route recorded in the second paragraph of that proof. Apply Theorem D. $\square$

F and its linear rigidity are in Theorem 11.2; the new statement is the two-jet strengthening. It answers the first-order case of the last sentence of Question 19.1 as posed by sources 08 and 09 (recorded in its status paragraph) with no coefficient restriction, and says nothing about relations of order two or more; Theorem 22.1 (source 12) answers that sentence in every order, again with no coefficient restriction.

Extension of scalars versus preservation of entireness. The degree classification is stable under the Hahn-field extensions of Corollary 15.6, but entireness of a nonpolynomial series need not be. Enlarge $\mathbb{Q}$ to $\mathbb{Q} + \mathbb{Q}\omega$ with its surreal order and view G over the larger field. At $z = t^{-\omega}$ the member exponents are $n^2 - n\omega$, with successive differences $(2n + 1) - \omega < 0$, so the support union contains an infinite descending sequence and the evaluation is not strong. The same computation appears in the rank-one-Berkovich report [14], and for F in Section 11.2. No theorem here claims that G or F is entire on the proper class $\mathbf{No}[i]$; hypotheses and derivative are relative to a specified set-sized workspace.

17.2 Which hypotheses are used

Example 17.3 (Positive characteristic destroys the conclusion). Let p be prime and $\mathbb{K} = \mathbb{F}_p((t^{\mathbb{Q}}))$. The nonpolynomial series $U(z) = \sum_{n \geq 0} t^{n^2} z^{pn}$ is strongly entire: at a point of valuation σ its

member valuations are $n^2 + pn\sigma$, and Lemma 2.3 (a) applies, its proof being independent of the coefficient field. Yet $D_z U = 0$.

So the conclusion fails in positive characteristic even for one linear equation. The characteristic-zero requirements of the proof are substantive: ordinary integer eigenvalues stay distinct, nonzero falling factorials stay nonzero, and a nonzero polynomial cannot vanish at all ordinary integers.

The hypothesis $\Gamma \neq 0$ is needed for the scale argument; the degenerate case $\Gamma = 0$ is treated in Section 12. Under ordinary complex analytic summation, by contrast, $\exp z$ is a nonpolynomial entire function with $f' = f$; these are different notions of infinite sum, and the classical exponential does not contradict the Hahn theorem. The coefficient field may be $\mathbb{R}$, $\mathbb{C}$ or any field of characteristic zero; neither algebraic nor real closedness is used, no completeness of a real-valued norm is used, and Γ needs neither an order unit nor a countable cofinal subset. Strong summability supplies exactly the finite initial layer the argument needs.

17.3 An exact finite workflow

For first-order input P the reduction is as follows. Combine like monomials and remove zero coefficients. If no Y or Y' remains, decide the polynomial identity directly. Otherwise compute d_P , w_P , I_P and κ_P from the finite term list, find the positive integer roots of I_P in the chosen exact representation of the coefficients, take the largest candidate degree, substitute a generic polynomial of that degree, and solve the resulting finite algebraic system. This procedure is complete for *entire* solutions whenever its finite coefficient operations and algebraic solving are available. It is not a general algorithm for arbitrary surreal names, it does not enumerate the local formal solutions, and it does not decide arbitrary strong-support membership. A negative answer after an exhaustive exact solve is a genuine nonexistence certificate for entire solutions; failure of a numerical solver is not.

For a nonpolynomial candidate formal solution a second workflow builds the certificate (15.2)–(15.4) from one sufficiently high nonzero coefficient and finitely many low-index ones; a claimed certificate must supply their exact leading valuations. The shipped program checks finite algebraic identities and sample certificates; it does not decide infinitary summability.

17.4 Predecessors, repository comparison and priority

Inherited ideas. Hahn fields, the positive-support summation lemma, Conway normal forms, Gauss initial forms, Euler eigenvalues and indicial-polynomial reasoning are established ideas and are not relabelled as new. The organizing method is related in spirit to maximum-term and central-index arguments in the Wiman–Valiron tradition, surveyed by Hayman [18]; here the finite residue polynomial gives an exact identity rather than an analytic approximation, and no real-valued growth estimate is imported. Source 10 consulted Hayman’s survey through its publisher metadata and abstract only, and uses no theorem from its full text.

Newton-polygon methods for nonlinear equations have a substantial history. Cano and Fortuny Ayuso construct initial and indicial polynomials for nonlinear q -difference equations and obtain support and growth results for generalized-series solutions [19]. Their operator and setting are not those of a formal z -series with arbitrary-rank Hahn *coefficients* evaluated strongly at every workspace scale. The conceptual precedent is important, and no invention of differential Newton polygons is claimed.

Non-Archimedean value distribution already restricts nonlinear functional and differential equations. Hu and Luan work over a complete algebraically closed field of characteristic zero with a nontrivial non-Archimedean absolute value and prove Clunie-, Malmquist- and Mokhon’ko-type results [20]; source 10 consulted the parsed text of that paper for its field hypotheses and Theorems 1.1, 1.3 and 1.5, reports that a remote screenshot service failed so that no visual inspection of the PDF is claimed, and reproduces no figure or table from it. Hu and Yang’s monograph [21] is a broader predecessor; only its publisher metadata and description were checked, so no claim is made that all rank-one special cases were previously unknown. Giansiracusa and Mereta develop differential enhancements of non-Archimedean norms and a tropicalization of differential equations [22]; the finite corner here records only the data this obstruction needs, and no equivalence to their framework and no new tropical fundamental theorem is asserted. Mantova and Matusinski’s survey [17] was consulted for normal forms.

Repository comparison at source 10’s pin. Source 10 compared its results with the collection at 048b72c; the comparison, with the corrections needed now, is as follows.

Material	Existing result or scope	Difference in the nonlinear part
This report, Theorem A	Strongly entire D-finite series are polynomials; the nonlinear case was the open Question 15.1, now Question 19.1.	All first-order algebraic equations, a higher-order corner criterion, and solution-dependent finite certificates.
This report, Theorem C	Finite-dimensional span of all mixed derivatives forces polynomiality.	One algebraic relation in f and $\vartheta_q f$ suffices for positive weights; no finite-dimensional span.
Tail-spans report [16]	Particular Galois-supported constructions have all-order differential independence, for scalar or fixed-domain analytic derivations over specified smaller base fields.	A two-jet theorem for every nonpolynomial strongly entire function over its own full Hahn field.
Documentation catalogue docs/README.md	Distinguishes workspaces, strong summability, coefficient rings and scalar derivations.	Those distinctions are kept; no global proper-class analytic claim is added.

The tail-span examples are neither weakened nor rediscovered: that report proves stronger all-order independence for particular constructions under different coefficient and function-class hypotheses, while the present universal statement assumes strong entireness and concludes only two-jet independence. (The coefficient-field part, from sources 11 and 12, now concludes all-order independence for groups without an order unit; its comparison with the tail-span report is in Section 22.2.) Source 10 cited the quadratic-exponent series as occurring in this report and in the catalogue; within the collection it occurs as the rank-one-Berkovich example [14], as `found:ex:internal` [15], and as a case of (8.1). This is the targeted comparison recorded at the pin, not an absence claim about every report in the current collection. The Riccati statement `diff:cor:riccati` of the differential-equations report concerns a scalar derivation (compare Proposition 15.7).

Source 10’s repository audit covered the root README, the catalogue, contents listings, this report’s definitions, theorems and nonlinear-question passages, and the README files of the tail-span, analytic-geometry and product-birthday reports (the last two for context only). It was a targeted comparison, not a line-by-line search of every report, Lean file, archive, branch or the Git history, and no absence claim rests on a truncated preview.

Priority and proof status. Source 10’s literature search, made on 22 September 2026, did not locate the exact all-rank strong-family first-order theorem, the finite exclusion certificate, or the positive-weight Euler formulation. Some searches were poorly targeted, and no exhaustive MathSciNet, zbMATH, full-book or citation-network review was made, so priority is provisional. The safest description is a proposed new theorem package and a proved first-order answer to a repository question, which is not a named historical conjecture; it is not a certified first discovery of nonlinear non-Archimedean rigidity. The written proofs are the evidence. The finite checks of Section 18 establish no arbitrary-support theorem; the main nonlinear rigidity theorems remain unmapped in the current implementation ledger. No independent human referee report accompanies the delivered source package.

18 The thirteen finite check suites

Each source manuscript shipped an independent exact-arithmetic suite. The suites of sources 08 and 09 were run on copies and passed, and neither supersedes the other: they cover overlapping but distinct finite facts of the linear theory, and they are kept side by side in `code/` for that reason. The suite of source 10 checks finite facts of the nonlinear part only; it was also run on a copy for this merge and reproduced its recorded counts exactly. The suites of sources 11 and 12 check finite facts of the coefficient-field part; both were run on copies for this merge and reproduced their records exactly, apart from the recorded interpreter and package versions. The suite of source 13 checks finite facts of the theta-descent part; it was run on a copy for this merge and reproduced its record apart from the interpreter version and the elapsed time. The suite of source 14 checks finite facts of the order-three-threshold part; it was run on a copy for this merge and reproduced its record apart from the interpreter version and the elapsed time. The suites of sources 18 and 19 check finite facts of the unit-dilation and Newton-rigidity parts; both were run on copies for this merge, source 18’s reproduced its record exactly and source 19’s apart from the recorded interpreter version. The suites of sources 16 and 17 check finite facts of the theta-hierarchy and single-scale parts; both were run on copies for this merge, source 16’s reproduced its record apart from the elapsed time and source 17’s apart from the recorded interpreter version. The suite of source 15 checks finite facts of the polynomial-composition part; it was run on a copy for this merge and reproduced its record apart from the recorded interpreter version. The suite of source 20 checks finite facts of the recurrence part; it was run on a copy for this merge and reproduced its record exactly, up to line endings. The suites of sources 08, 09, 10, 12, 16, 18 and 20 use only the Python standard library, integers, and `fractions.Fraction`; the suites of sources 11, 13, 14, 15, 17 and 19 use exact symbolic and rational arithmetic in SymPy, the only third-party package any suite needs. None uses floating-point mathematics or network access; those of sources 13, 14 and 16 record their elapsed time.

code/08-finite-recurrences-verify.py — 3,705 checks, 0 failures. It emphasises the operator-to-recurrence conversions in all three formats and the higher-rank ordered-group arithmetic.

Finite calculation	Checks
Differential operator to coefficient recurrence	648
Dilation operator to coefficient recurrence	336
Mixed operator to coefficient recurrence	276
Unit residues, parity, and finite exceptional cancellation	400
Partial theta identities and finite quadratic estimates	552
Higher-rank lexicographic valuation examples	1,292
Sparse Airy recurrence and finite escape edges	201
Total	3,705

The operator tests compare the grouped recurrence formulas against direct finite polynomial differentiation, dilation and multiplication; the theta tests compare Laurent monomial exponents exactly, including the q^{-1} -to- q conversion of (8.7); the unit tests retain the exceptional index $n = 7$ rather than silently checking only generic values; and the ordered-group checks use lexicographic integer tuples, not a real-valued surrogate norm. This script takes a *required* `--output` path and exits nonzero without one, which is deliberate: it prevents a re-run from overwriting the delivered record.

code/09-entire-hahn-holonomic-verification.py — 2,043 checks, 0 failures. It emphasises the boundary behaviour of the exclusion bounds and the exact theta domain.

Finite calculation	Checks
Theta homogeneous equation, positive and negative valuation	92
Theta first-order coefficient recurrence	45
Dilation and differential forward recurrence conversions	72
Unit orbit: parity, exceptional cancellation, residue polynomial	504
Rank-two noncofinal affine minimum	200
Theta excluded upper-scale argument	200
Theta included coarse boundary	170
Theta cofinal-scale growth	120
Finite-coordinate entire witness growth	480
Sparse recurrence exact identity	50
Escape boundary: constant leading exponent, strict exterior descent	50
Formal exponential recurrence	60
Total	2,043

Two of these rows exist only in this suite and check the two boundary phenomena that the report insists are different: “theta included coarse boundary” checks that a zero coarse valuation coexists with positive growth in the original lower-scale coordinate, which is Section 8.5, and “escape boundary” checks the constant-leading-exponent case that Lemma 2.2 rules out, which is what makes the excluded region in Corollary 3.5 closed. Its `--output` argument is optional. Seeds are recorded in the shipped records.

code/10-nonlinear-rigidity-verify.py — 2,113 checks, 0 failures. It checks the finite algebra of Sections 14–17.

Finite calculation	Checks
Finite Hahn initial layers and exclusion inequalities	450
Automatic first-order nondegeneracy of the corner	240
Univariate falling-factorial corner identities, orders 0 to 3	480
Lexicographic higher-rank ordered-group exclusion bounds	500
Weighted Euler homogeneous-part identities	360
Worked examples: Riccati, degree cutoff, resonance, Painlevé, vanishing corner, Euler solution space	83
Total	2,113

The group names in the record `data/10-nonlinear-rigidity-verification.json` are `finite_hahn_certificate`, `first_order_nondegeneracy`, `higher_order_corner`, `ordered_group_radius`, `weighted_euler` and `worked_examples`, in the order of the table; “radius” there is source 10’s word for an exclusion bound. The default seed is 20260922. The `--output` argument is required, and the program refuses to overwrite an existing file. The vanishing-corner example is tested for the monomial degrees $0, \dots, 20$ only. The counts are program checks, not independently proved theorems.

code/11-coarsening-differential-rigidity-verify.py — 8,673 cases in 8 groups, 0 failures. It checks finite facts of source 11’s material and needs SymPy (`data/11-coarsening-differential-rigidity-requirements.txt` pins `sympy>=1.12,<2`; the record was made with Python 3.13.5 and SymPy 1.14.0, the merge’s rerun with Python 3.14.4 and SymPy 1.14.0).

Group in <code>data/11-coarsening-differential-rigidity-verification.json</code>	Cases
<code>singular_tail_coefficient_isolation</code> : Taylor identity (21.4), orders 1–3	12
<code>vanishing_corner_indicial_polynomial</code> : (21.6) for $0 \leq m \leq 9$	10
<code>painleve_local_formal_solutions</code> : 24 coefficient equations, three initial conditions	72
<code>common_factor_reduction_models</code> : numerator and denominator reduction	3
<code>binary_prefix_coding</code> : code ranges and all pairs of length-seven branches	8,136
<code>finite_reverse_lexicographic_tests</code> : growth and loss after an exterior scale	434
<code>generator_value_cancellation</code> : Remark 20.8	1
<code>boundary_equations</code> : rational Riccati solution, characteristic p	5
Total	8,673

Most of the count is prefix-pair comparisons; it is not a count of independent theorems. The `--output` argument is optional and defaults to `verification.json` in the current directory, so a bare run can overwrite a file of that name; use an explicit new path.

code/12-coefficient-field-rigidity-verify.py — 9,308 checks, 0 failures. It checks finite facts of source 12’s material with the standard library only.

Category in <code>data/12-coefficient-field-rigidity-verification.json</code>	Checks
coefficient recovery; formal solution coefficients	387; 637
linearized recurrence identity; mixed derivative multiplier scaling	389; 59
derivative-before-dilation convention (20.1)	405
degenerate Euler multiplier (21.6)	1,500
exponential-polynomial annihilator	257
distinct monomial exponent vectors	256
exterior descending leading values; successive Archimedean levels	462; 1,400
partial-theta exponent identity; partial-theta finite escape bounds	200; 2,706
positive-characteristic derivative obstruction	500
torsion even-function identity	150
Total	9,308

The record also lists thirteen formal recurrence cases with their offsets and sampled zero multipliers; the sampled zeros are not an effective Skolem–Mahler–Lech bound. The `--output` argument is required and an existing file is refused unless `--force` is given.

code/13-theta-descent-verify.py — 1,622 checks, 0 failures. It checks finite facts of the theta-descent part with exact rational arithmetic in SymPy (`data/13-theta-descent-requirements.txt` pins `sympy>=1.12,<2`; the record was made with Python 3.13.5 and SymPy 1.14.0 at precision p^{128} , the merge’s rerun with Python 3.14.4 and SymPy 1.14.0).

Group in <code>data/13-theta-descent-verification.json</code>	Checks
Chebyshev closed coefficients (23.3); Chebyshev reciprocal substitution (23.2)	30; 30
Taylor coefficient extraction: the coefficients of C_n entering (23.4)–(23.5)	495
Theta shift exponents (23.10)	81
Jacobi product p -coefficients (23.12)	128
Third-order ODE p -coefficients (24.7)	128
Exact jet identities: (24.8) and the system (24.18)	6
Root-factor cancellations; root and rounding leading terms (25.2)–(25.4)	16; 48
Finite local-linearization leading terms (23.15)	560
Lexicographic exclusion checks (Example 25.5)	100
Total	1,622

The record’s group names and its field `q_precision_exclusive` use source 13’s letter q for the nome called p here. The product and the cleared equation have zero residual in every p -coefficient below p^{128} , each checked as an exact polynomial in the variable; the Lambert constants use the nome p^2 . The `--output` argument is required and an existing file is overwritten, so pass a new path; `--precision` defaults to 64, and the record uses 128. For this merge the identities used by Lemmas 24.5 and 24.7 and Proposition 24.2, the expansions of Section 24.1, the rational forms of Theorems 24.1 and 24.10, and the cleared equation and product modulo p^{50} were also checked with an independent SymPy script, which is not shipped; it also confirms that with the nome p in place of p^2 the residual of the cleared equation is nonzero.

code/14-order-three-threshold-verify.py — 600 checks, 0 failures. It checks finite facts of the order-three-threshold part with exact rational and polynomial arithmetic in SymPy (**data/14-order-three-threshold-requirements.txt** pins `sympy>=1.13,<2`; the record was made with Python 3.13.5 and SymPy 1.14.0, the merge’s rerun with Python 3.14.4 and SymPy 1.14.0).

Group in data/14-order-three-threshold-verification.json	Checks
symbolic_jets: the coefficient $(z^2 - 4)^3 Y_0^4$ of Y_3^2 ; after (27.3), z -degree six, the form (27.4) and its residue (27.5); the binomial gap identity $(g^2 - 1)v^2$; a second-order endpoint example	6
endpoint_cancellation: removal of V^{i_0} , $0 \leq i_0 \leq 4$, and the resulting endpoint polynomial, as in Lemma 26.8	10
finite_initial_classification: all 242 nonzero polynomials of degree at most four with coefficients in $\{-1, 0, 1\}$ against Theorem 27.4	242
chebyshev: identities (23.2) and (23.3)	72
newton_corners: at $\delta_n = (2n + 1)\mu$, $0 \leq n \leq 80$, the active indices $\{n, n + 1\}$ and the value $-n(n + 1)\mu$ for $\mathcal{T}_p$	162
positive_characteristic: derivative zero of $\sum z^{pn}$ modulo $\mathfrak{p} \in \{2, 3, 5, 7, 11\}$	5
theta_product_truncation; theta_ode_truncation: the product (23.12) and the cleared equation (24.7) through p^{47}	48; 48
theta_coefficient_values: (23.6)	7
Total	600

The record’s group names use source 14’s word “corner” for a break and its letter q for the nome called p here; its field **cutoff_exclusive** is 48, so every p -coefficient below p^{48} of the product and of the cleared equation is compared as an exact polynomial in z . The initial-form enumeration admits 26 of the 242 polynomials and is not the proof of the classification. The script takes no arguments: it writes **data/verification.json** in the parent of its own directory. In this layout that is a new file beside the delivered record **data/14-order-three-threshold-verification.json**, which it does not touch, but a second run overwrites the first; run it on a copy.

code/18-unit-dilation-verify_finite.py — **16,579 checks, 0 failures.** It checks finite facts of the unit-dilation part with the standard library only (exact integers and `fractions.Fraction`; fixed seed 20260923; Python 3.10 or later, as the source’s README states).

Category in <code>data/18-unit-dilation-finite_verification.json</code>	Checks
binomial_tail : the tails of Example 29.14, $n \leq 40$, $d \leq 13$	975
parity_valuation : Example 4.6, $1 \leq n \leq 100$	200
theta_identity ; differentiated_theta : the coefficients of (8.2) and (29.7) for five rational parameters	205; 200
two_prescribed_dilations : the coefficients of (29.8) for rational multipliers	600
operator_output_formula ; reindexed_shift_formula : (29.2) and (29.4)–(29.5) against direct application, 160 random operators	3,187; 3,187
finite_independence_matrix : nonzero confluent Vandermonde determinants	12
frobenius_boundary : $\binom{\mathfrak{p}^a}{i} \equiv 0 \pmod{\mathfrak{p}}$ for $0 < i < \mathfrak{p}^a$, $\mathfrak{p} \leq 7$, $a \leq 3$	12
escape_inequality : $B + j\delta < 0$ for $\delta = -B - g$, small integers	1,440
unbounded_polynomial_degree : the coefficients of $(\sigma_2 - 2^a)z^a$, $a, n \leq 80$	6,561
Total	16,579

Rational parameters check finite coefficient identities, not entireness. Two categories check only elementary integer arithmetic: the 6,561 checks of **unbounded_polynomial_degree** compare an expression that is zero by construction, and the 1,440 checks of **escape_inequality** verify $B + j\delta < 0$ for small integers; together they are 8,001 of the 16,579, which is why the count is no measure of proof coverage, as the source says. The `--output` argument is optional: without it the program only prints its record; with it, it writes the given path, creating directories and overwriting an existing file, so pass a new path. The record names no interpreter version.

code/19-newton-rigidity-verify.py — **259 checks, 0 failures.** It checks finite facts of the Newton-rigidity part with exact polynomial arithmetic in SymPy (`data/19-newton-rigidity-requirements.txt` pins `sympy==1.14.0`; the record was made with Python 3.13.5 and SymPy 1.14.0, the merge’s rerun with Python 3.14.4 and SymPy 1.14.0).

Groups in <code>data/19-newton-rigidity-verification.json</code>	Checks
<code>conic_homogeneous_rank, conic_kernel_dimension</code> : the conic kernel of Remark 30.2, degrees 1–9	18
<code>twisted_cubic_gradient_injectivity</code> : Lemma 30.7 as a rank computation, degrees 1–3	3
<code>first_polar_double_factor, first_polar_remaining_degree</code> : (30.2) on a basis of curve-vanishing forms	26
<code>secant_jacobian, cubic_polar_example</code> : (30.4) and Example 30.10	2
<code>conic_lowest_cross_term</code> : the coefficient $a_m a_n (n - m)^2$ of Remark 30.2	60
<code>quartic_resultant, quartic_singular_on_twisted_cubic</code> : (31.2) and $\text{Pol}_{H_3} = 0$	5
<code>quartic_binomials, fourth_order_binomials</code> : doubling binomials solve H_3 and Q_4 , $n \leq 12$	72
<code>fourth_order_pair_kernel, hidden_residual_z5</code> : (31.4) and (31.6)	2
<code>quadratic_newton_transition, doubling_newton_transition</code> : finite windows of the breaks of G_1 and F_{sp}	51
<code>first_polar_lowest_coefficient</code> : (30.3) for four forms	20
Total	259

The script takes no arguments and always writes `data/verification.json` in the parent of its own directory, recording the interpreter and SymPy versions. In this layout that is the same file that source 14’s script writes, a new file beside both delivered records, overwritten by each run; run it on a copy. The source’s `code/19-newton-rigidity-build.py` was written for the delivery layout: it typesets an `article.tex` in its own directory `code/`, where there is none, and its `--verify` option calls a `code/verify.py` relative to that directory, which is not present. The shipped `data/19-newton-rigidity-build_audit.json` describes the 24-page source manuscript, whose source and PDF, hashed there, are not shipped.

code/16-theta-hierarchy-verify.py — exact comparisons, status PASS. It checks finite facts of the theta-hierarchy part with the standard library only, in $\mathbb{Z}[z][p]/(p^{128})$ with exact rationals; its record lists the comparisons by kind and gives no single total.

Field in <code>data/16-theta-hierarchy-verification.json</code>	Count
Chebyshev recurrence against the closed formula (23.3)	30
leading Taylor terms of the truncated descent (Proposition 23.3)	12
Jacobi product (23.12) against the descent, p -coefficients below p^{128}	128
cleared order-three identity (source 16's Tate form), p -coefficients below p^{128}	128
Gauss profile of Theorem 32.3 and its bounds, rational cases	852
dominance in Example 32.6, exact $\mathbb{Q}(\sqrt{2})$ comparisons	65
unique squarefree wedge assignment, $N = 1, \dots, 8$	8
two-function constant-derivative model of the grid: candidates, nonzero jet determinants	81; 50

The last row is an illustrative finite specialization, not the compression theorem, and a vanishing residual modulo p^{128} checks normalization and finite algebra, not an infinite identity. The script's `--output` defaults to `verification.json` in the current directory and overwrites it; `--precision` defaults to 128, the value of the record; the record names the elapsed time (0.496s) but no interpreter version. The shipped `code/16-theta-hierarchy-Makefile` names the delivery paths `article.tex` and `verify.py`: its default target would run `latexmk` on whatever `article.tex` is in the current directory, and its `verify` target calls a `verify.py` that is not present under that name. `data/16-theta-hierarchy-build_audit.json` describes the 25-page source manuscript, which is not shipped.

code/17-single-scale-verify.py — 8,608 checks in 21 groups, 0 failures. It checks finite facts of the single-scale part with exact integer and sparse polynomial arithmetic and SymPy (data/17-single-scale-requirements.txt pins `sympy==1.14.0`; the record was made with Python 3.13.5 and SymPy 1.14.0, the merge's rerun with Python 3.14.4 and SymPy 1.14.0).

Groups in <code>data/17-single-scale-verification_results.json</code>	Checks
<code>height_separation</code> : Lemma 34.3 in finite windows	7,198
<code>coding_scale_cosets</code> ; <code>commuting_jets</code> ; <code>Euler_conversion</code>	432; 256; 144
<code>intrinsic_monomials</code> ; <code>intrinsic_triangular</code> : (35.1), Lemma 35.5	153; 9
<code>finite_valuation_domain</code> ; <code>exterior_domain_obstruction</code> : Theorem 35.10	124; 16
<code>polarization</code> ; <code>polarization_orbits</code> ; <code>polarization_cancellation</code> : Lemma 34.5, Example 34.6	53; 53; 1
<code>binary_prefixes</code> ; <code>private_prefixes</code>	46; 32
<code>floor_cutoffs</code> ; <code>floor_examples</code> : Theorem 35.18	45; 5
<code>omnific_code_support</code> ; <code>jet_block_rank</code> ; <code>finite_patterns</code>	16; 12; 6
<code>coefficient_witnesses</code> ; <code>necessary_threshold</code> ; <code>polynomial_height_obstruction</code>	5; 1; 1
Total	8,608

Most of the count is one lemma checked in many windows; it is not a count of theorems. The record also lists three sparse coefficient witnesses (for instance the coefficient 1 of $p^{403200}z^{17}$ in $\mathcal{G}\partial^2\mathcal{G} - (\partial\mathcal{G})^2$) and the floor examples. The `--output` argument is optional and defaults to

`verification_results.json` *beside the script*, here in `code/`, overwriting any file of that name; pass a new path. The source's `code/17-single-scale-build.py` looks for an `article.tex` in its own directory `code/`, where there is none, and stops with an error before writing anything; in the delivery layout it wrote into `.build/` and replaced `article.pdf` and `build_audit.json` in place. `data/17-single-scale-build_audit.json` and `data/17-single-scale-visual_audit.json` describe the 24-page source manuscript, which is not shipped.

code/15-polynomial-composition-verify.py — 2,369 checks in 12 groups, 0 failures.

It checks finite facts of the polynomial-composition part with SymPy and exact rational Hahn polynomials over value groups $\mathbb{Z}^2$ (lexicographic), with the fixed seed 20260923 (`data/15-polynomial-composition-requirements.txt` pins `sympy==1.14.0`; the record was made with Python 3.13.5 and SymPy 1.14.0, the merge's rerun with Python 3.14.4 and SymPy 1.14.0).

Groups in <code>data/15-polynomial-composition-verification.json</code>	Checks
<code>universal_encoding</code> : Theorem 37.3	648
<code>indicial_leading_terms</code> ; <code>exact_resonance</code> : Lemma 36.8, Theorem 36.12	445; 208
<code>integer_convexity</code> : Lemma 36.4	360
<code>euler_projectors</code> : (37.1)	228
<code>smith_certificates</code> : Theorem 37.2	150
<code>rank_two_operator_profiles</code> ; <code>rank_two_pullback_profiles</code> ; <code>rank_two_gauss_p</code>	90; 70; 35
<code>roducts</code> : Propositions 36.7, 36.5, Corollary 14.3	
<code>boundary_coefficient_checks</code> : Section 37.4	75
<code>sharp_degree_family</code> ; <code>worked_equations</code> : Section 37.3	48; 12
Total	2,369

The record also lists the degree bounds 1, 3, 1 and 2 of the four worked equations of Section 37.3 with a bound. The `--output` argument is optional and defaults to `verification.json` in the current directory, overwriting any file of that name; pass a new path. The source's `code/15-polynomial-composition-Makefile` was written for its delivery layout: its default target runs `pdflatex` three times on an `article.tex` in the current directory, rebuilding `article.pdf` in place, its `verify` target calls a `verify.py` there, and its `clean` target deletes auxiliary files with `rm -f`; do not run it here. `data/15-polynomial-composition-BUILD_REPORT.json` describes the 24-page source manuscript, which is not shipped.

code/20-recurrences-verify.py — **4,605 checks in 12 families, 0 failures.** It checks finite facts of the recurrence part with the standard library only (exact integers and `fractions.Fraction`, finite Laurent polynomials in t , lexicographic pairs for exponents $a\omega + b$; fixed seed 20260923; Python 3.10 or later, as the source states).

Family in <code>data/20-recurrences-verification.txt</code>	Checks
unit coefficient formula: (38.1) against direct expansion, three cases, $n \leq 18$, exponents -3 to 24	1,596
residual parity valuation: Example 4.6, $1 \leq n \leq 80$	80
principal-unit cancellation: the case $d = 2$ of Example 29.14, $n \leq 44$	45
unbounded coordinate transients: $\binom{n}{d} = 0$ exactly for $n < d$ (Example 38.10), $d \leq 18$, $n \leq 44$	810
higher-rank affine comparison: $\min((1, 0), (0, n)) = (0, n)$, $n \leq 200$	201
omnific coefficient coding: Theorem 39.3 for a rational sequence and for the squares, finite windows	281
Catalan algebraic identity; Catalan power coefficients; declining omnific support: Example 39.2	40; 88; 88
partial-theta operator: the coefficients of (8.3) as Laurent polynomials in the parameter, $n \leq 100$	101
mixed-operator coefficient indexing: (29.2) against direct application, 35 random operators with multipliers 1, 2, 3	875
positive-characteristic boundary: Example 39.12 for $\mathfrak{p} \leq 7$, $n \leq 100$	400
Total	4,605

Integer multipliers check coefficient indexing, not entireness, and a finite truncation of λ_- or of $\mathfrak{w}_b$ checks only the coefficients that the omitted tail cannot reach. Four families check facts that hold by construction: the 810 binomial-vanishing checks, the 201 and 88 lexicographic comparisons, and the 90 positivity checks inside the coding family; together they are 1,189 of the 4,605, so the count is no measure of proof coverage, as the source says. The script takes no arguments and prints its record to standard output only; the delivered record is that output (the merge’s rerun, under Python 3.14.4, matched it exactly after normalizing the Windows line endings of the redirected output). The source’s `code/20-recurrences-build.sh` was written for its delivery layout: it changes to its own directory, runs `pdflatex` three times on an `article.tex` there and then `python3 verify.py > verification.txt`, overwriting that record by redirection. Here its directory is `code/`, which has neither file: run in this layout it stops at the first `pdflatex` call and leaves `texput.log` in `code/`; do not run it here. Source 20 shipped no build record, and its 22-page manuscript is not shipped.

What the checks do not do. No suite proves any infinite theorem. No finite prefix can prove strong summability of an infinite family, cofinality of (e_n) , noncofinality of $(n\epsilon)$, the existence of an infinite escape path, the eventual orbit statements, the generic-line theorem, or the absence of every possible annihilating equation. No infinite Hahn sum is evaluated, or numerically approximated, by any of the programs. In particular a finite truncation of the explicit series (11.2) is a polynomial and is D-finite, so no finite computation can witness its nonholonomicity; for the same reason no finite computation can witness the two-jet independence of Corollaries 17.1 and 17.2. Source 10’s suite in particular does not test arbitrary infinite supports, cofinality, all Hahn elements, the nonexistence of all annihilating equations, or historical novelty. The suites of sources 11 and 12 do not prove the coefficient-field theorem in general, rationality descent for infinite series, the Skolem–Mahler–Lech

theorem, infinite support well ordering, cofinality, or any algebraic independence statement, and no finite computation can witness the independence of all jets of F . Source 13's suite does not verify strong summability of infinite supports, minimal differential order or algebraic independence, the no-order-unit obstruction, completeness of the infinite zero set, or priority, as its record states; a finite truncation of $\mathcal{T}_p$ is a polynomial. Source 14's suite does not establish the quantifiers over arbitrary Hahn supports, ordered groups, scales or formal solutions, and its enumeration of initial polynomials is not the classification proof. Source 18's suite does not prove well ordering of arbitrary supports, strong summability at all field elements, the Skolem–Mahler–Lech theorem, eventual bounds for exceptional zeros, the nonexistence of an annihilator, or novelty; source 19's suite certifies no infinite-support argument, no cofinality statement, nothing over arbitrary fields, and no novelty, and a finite truncation of F_{sp} or of G_1 is a polynomial. Source 16's suite proves no summability of infinite supports, no algebraic independence of the full functions, not the compression theorem, and no minimal order or cardinality. Source 17's suite tests finite windows of the band lemma and finite examples; it quantifies over no supports, equations or continuum families. Source 15's suite does not establish summability of arbitrary infinite supports, the weighted theorem, the universal encoding for all polynomials, or novelty; its Hahn profiles are computed for polynomials over $\mathbb{Z}^2$ only. Source 20's suite does not prove the Hahn support lemma, the Skolem–Mahler–Lech theorem, the infinite-support minimum argument, Noetherian compression or the proper-class localization; its coding checks are finite windows. What the suites can detect is sign, indexing, conversion and finite valuation-comparison mistakes. Everything else rests on the written arguments.

The mathematical dependency chain is short.

Conclusion	Load-bearing ingredients
Differential rigidity	Integer evaluation lemma; finite shift conversion; escape chain and exclusion criterion.
Unit-dilation rigidity	Residue polynomial; finite binomial leading terms; escape chain and exclusion criterion.
Noncofinal nonunit rigidity	Eventual order of finitely many affine values; noncofinality witness; escape chain and exclusion criterion.
Coarsened exclusion	The same affine values, projected to $\Gamma/\Delta_{ v(q) }$; strictness of the projected descent.
Positive dilation existence	Support monoid certificate and the eventual strict increase of h_n ; or, in the order-unit case, quadratic leading valuations and the cofinal-valuation criterion.
Several-variable rigidity	Countable polynomial avoidance; finite-block regrouping; rational differential system on a generic line.
First-order nonlinear rigidity	Hahn-family algebra and its residue homomorphism; large active degrees; exact corner identity; uniqueness of a corner term for each power of Y' .
Higher-order corner criterion	The same, with $\chi_P \neq 0$ assumed.
Positive-weight Euler rigidity	The same algebra in m variables; finite bounded-weight sets; the eigenvalue identity of ϑ_q .
All-order rigidity without an order unit	Minimal relation and its linearization; one finitely generated coefficient field; valuation-rank inequality; principal convex bound; exterior obstruction, or coarsened reduction.
Mixed-jet independence	The same, with an exponential-polynomial multiplier and the imported Skolem–Mahler–Lech theorem.

Conclusion	Load-bearing ingredients
Rationality in $\text{Frac } \mathcal{E}_\Gamma(k)$	Finite coefficient field; common coarsened reduction of numerator and denominator; rationality descent.
Continuum family	Joint-independence criterion; independent scale monomials; binary-prefix codes; a separate cardinality bound.
Exact domain of $\mathcal{T}_p$	Chebyshev coefficients; quadratic positive-support certificate; descending leading exponents.
Order-three equation for $\mathcal{T}_p$	Classical triple product; elliptic pole cancellation; formal identity over $\mathbb{Q}$; ring maps to $\mathbb{K}$.
No equation of order below three	Laurent ring and substitution homomorphism; theta scaling law; fixed-field, additive-difference and u -adic obstructions; $\psi_3 \neq 0$.
Degree two in the third derivative	Three-jet independence; odd pole order of a cubic in the transcendental $\mathcal{A}$.
No linear equation for $\mathcal{T}_p$	Transcendence of ψ_1 over the elliptic field; highest power in normalized derivatives.
Omnific consequences	Linear-factor product; normal forms; local product linearization.
Second-order rigidity (Theorem I)	All-scale criterion; cofinal divisible extension; local finiteness and late breaks; exterior homogeneous reduction by jet degree, then z -degree; nonzero dehomogenization; removal of V^{i_0} ; nonconstant logarithmic Euler derivative in characteristic zero; the finite endpoint lemma.
Minimum order three	Second-order rigidity; the entire witness $\mathcal{T}_p$ and its equation of order three.
Third-order gap certificate	Homogeneous logarithmic substitution; lowest transverse degree; expansions at 0 and ∞ .
Theta initial forms and eventual convexity	Residue form of the exterior equation; rational Riccati reduction; constants of rational differentiation; exclusion of skipped and triply tied indices; local finiteness.
Periodic-affine valuations (Theorems 28.5, 28.10)	Common support envelope; one period from the residues; the imported Skolem–Mahler–Lech theorem through Lemma 21.1; the least coefficient sequence that is not an identity; eventual order of finitely many affine functions.
Relative-scale rigidity and dichotomy (Theorem J)	Exact shift polynomials; periodic-affine valuations; noncofinality witness; escape chain and exclusion criterion; the differentiated theta identity and Theorem 8.1.
Third order through cubic jet degree (Theorem K)	Late breaks and the exterior form (as for Theorem I); bottom and top first-polar coefficients; the double factor; the eventual side of integral roots through a $\mathbb{Q}$ -linear functional; the secant Jacobian.
Exact classifications of $\mathbf{H}_3$ and $\mathbf{Q}_4$	Resultant identity and a common-root parameter; constants of ϑ ; the pair kernel (31.4).
Independent theta scales (Theorem L (a))	Coefficient values μn^2 ; exact Gauss profile; Gauss multiplicativity; eventually affine Gauss values of polynomials; a unique dominant summand.

Conclusion	Load-bearing ingredients
Unbounded order (Theorem L (b))	Kähler criterion; invertible coefficient-to-jet change; unique squarefree wedge term; finite-grid nonvanishing; separant argument for finite composita; the order-three equation of $\mathcal{T}_p$.
The field $\mathcal{D}_{\text{DA}}$	Hamel basis (choice); countable well-ordered real supports; finite composita.
Valuation route to order three	Laurent ring, shift $\mathbf{S}$, Gauss value of radius one, eventually constant values of polynomials at integers, quadratic growth of iterates.
Rapid-height freedom (Theorem M)	Height separation of maximal-degree multisets; multicolour polarization; hereditary Zariski density of $(n, \mathbf{b}_n)$; full-row-rank pattern change; finite coefficient minors or constant extension.
Exact domain in $\mathbf{No}[i]$, codes, floors	Leading values $\mathbf{b}_n + n\gamma$; positive support monoid; disjoint real cosets of powers of M_Ω ; discreteness of $\mathbf{Oz}$.
Polynomial-composition rigidity (Theorem N)	Exact pullback profile; exact operator profile at large active degrees; Gauss multiplicativity; integer convexity; superlinear escape.
Degree bounds and coefficient loci	Exact leading term through I_L ; finite integer resonance set; comparison of polynomial degrees.
Omnific modules, universality, undecidability	Smith normal form and the constant-term retraction; Euler projectors; the MRDP theorem and Sun's theorem (imported).
Observable algebra (Theorem 38.2)	Common support envelope; finite binomial expansion with $\binom{U}{i} \in k[U]$ in characteristic zero.
Noetherian compression (Theorem 38.8)	Observable algebra; Hilbert's basis theorem; one torsion period for all monomials in the residues; the imported Skolem–Mahler–Lech theorem through Lemma 21.1.
Surreal recurrence profiles, matrix powers	Algebraic closedness of $\mathbf{No}[i]$; localization to a set-sized workspace; Jordan form of a finite companion matrix; periodic-affine valuations; exterior powers.
Omnific coding (Theorem 39.3)	The normal-form support definition and monomial multiplication only; no Skolem–Mahler–Lech.

All supports are sets. All operators are finite sums. All integer indices are ordinary nonnegative integers. The coefficient valuation is trivial in characteristic zero; the coefficient field is $\mathbb{C}$ in Sections 1–13 and any characteristic-zero field in the nonlinear, coefficient-field, theta-descent, order-three-threshold, unit-dilation, Newton-rigidity, polynomial-composition and recurrence parts. The coefficient-field, unit-dilation and recurrence parts import one external recurrence theorem without proof, the Skolem–Mahler–Lech theorem, used only for relations involving at least two distinct dilations or multipliers, or exponential polynomials with at least two bases. The polynomial-composition part imports the MRDP theorem and Sun's theorem, used only for Corollaries 37.5 and 37.6. The equations use the external formal variable derivative. None of the proofs asserts convergence in the fine topology of $\mathbf{No}[i]$, substitutes a real-valued norm for an arbitrary-rank valuation, or transports a fixed-workspace evaluation to every surcomplex argument.

19 Originality assessment and questions left open

19.1 Established background and proposed contribution

The D-finite/P-recursive correspondence [1], ordinary properties of Hahn fields, the Hahn–Neumann support lemmas and Higman’s finite-word theorem [6], finite polynomial root bounds, the Conway normal-form identification [9, 10], and partial theta series with their elementary functional identity [5] are established material. The companion report’s cofinality and scalar-extension work, including the group Γ_∞ and the distinction between order units and countable cofinality, is prior project material [12]. The rank-one quadratic-exponent entire example is already in the collection [14]. We claim to originate none of these, and we do not claim to have disproved any published theorem about a different analytic function class.

The proposed research contribution is their combination into the sharp arbitrary-rank dilation classification, Theorem B: in particular the necessity argument for all noncofinal scales, the unit case with root-of-unity residue, the uniform exterior obstruction applying to every solution of a fixed equation, the quantitative exclusion bounds with their two different boundary behaviours, and the exact theta evaluation domain for an arbitrary Hahn parameter. The mixed noncofinal extension (Theorem 10.2), generic-line rigidity (Theorem C), and the explicit series excluded from every nontorsion pure dilation equation (Theorem 11.2) are further consequences proved here.

The nonlinear part (source 10) inherits Gauss initial forms, Euler eigenvalues and indicial-polynomial reasoning, and is related to central-index, Newton-polygon, non-Archimedean value-distribution and tropical differential methods (Section 17.4). Its proposed contribution is the precise arbitrary-rank strong-family formulation: Theorem D with its finite exclusion certificate (Theorem 15.1), the candidate degrees and the entire-solution locus, the higher-order corner criterion, the positive-weight Euler theorem, and the non-uniformity of the nonlinear exclusion bound. Its priority is provisional in the sense recorded there.

The coefficient-field part (sources 11 and 12) inherits classical coefficient recurrences, the finite-generation principle, the valuation-rank inequality and the Skolem–Mahler–Lech theorem (Section 22.6). Its proposed contributions are the all-order no-order-unit rigidity with its joint-dilation form (Theorem E), the coefficient-value-rank criterion and the order-unit dichotomy (Theorems F and G), the meromorphic coarsening descent (Theorem 21.11) and the full-constant-field continuum family (Theorem 22.6); its priority is equally provisional.

The theta-descent part (source 13) inherits the Chebyshev identity, Jacobi’s triple product and the theta-elliptic differential identity, all classical, the bilateral theta domain of the Hahn–Tate report, and the no-order-unit direction of Theorem E (Section 25.5). Its proposed contributions are the support-controlled descent to an entire series with its exact domain at every scale, the exact boundary of Theorem H with the minimal-order certificate, the differential field and non-D-finiteness, and the surreal and omnific consequences; its priority is equally provisional.

The order-three-threshold part (source 14) inherits the theta material of source 13, the no-order-unit direction of Theorem E, the Gauss initial polynomials of source 10 and classical logarithmic derivatives (Section 27.5). Its proposed contributions are second-order rigidity for every value group (Theorem I (a)) with its finite endpoint lemma, the exact least order three, the third-order endpoint-gap obstruction, the classification of theta initial polynomials and eventual Newton convexity; its priority is equally provisional.

The unit-dilation part (source 18) inherits the support calculus, the escape mechanism, the partial theta series and the imported Skolem–Mahler–Lech theorem (Section 29.7). Its proposed

contributions are the periodic-affine valuation theorem for Hahn exponential polynomials, the bivariate unit-orbit estimate and the answer to Question 19.4, and the relative-scale dichotomy of Theorem J; its priority is equally provisional.

The Newton-rigidity part (source 19) inherits the all-scale criterion, the corner and break techniques, and second-order rigidity, which it re-derives independently (Section 31.4). Its proposed contributions are third-order rigidity through cubic jet degree (Theorem K) with the first-polar, secant and eventual-side lemmas, the constant-extension and algebraic two-state consequences, and the exact classifications of two boundary equations; its priority is equally provisional.

The theta-hierarchy part (source 16) inherits source 13's seed, equation and minimal order three and the classical theta product and Tate identity (Section 33.4). Its proposed contributions are independence across scales, bounded polynomial jet compression, unbounded minimal order, the transcendence degrees of the differentially algebraic entire field for $\Gamma = \mathbb{R}$, integer sampling and a valuation route to order three; its priority is equally provisional.

The single-scale part (source 17) inherits Grönwall's single-series gap criterion, the Berarducci–Mantova derivation, the omnific integers and the bounded-support descent (Section 35.6). Its proposed contributions are the exact differential relation ideals of rapid-height families, their differential transcendence over the full Hahn field at one order-unit scale, and the surreal, domain and omnific realizations; its priority is equally provisional.

The recurrence part (source 20) inherits the support lemma, the Skolem–Mahler–Lech theorem, Hilbert's basis theorem, the escape mechanism and the partial theta series, and re-derives source 18's valuation theorem (Section 39.6). Its proposed contributions are the finitely generated algebra of coefficient observables, Noetherian compression of coefficient conditions with one universal residual period, the recurrence profiles over $\mathbf{No}[i]$, $\mathbf{Oz}$ and $\mathbf{Oz}[i]$, the reading of a known omnific stream as a universal coefficient observable (Theorem O), and determinantal profiles of matrix powers; its priority is equally provisional.

19.2 A result-by-result attribution ledger

Result or ingredient	Status in this report
Hahn arithmetic, strong support sums, surreal normal forms	Standard background; references and support proofs supplied.
Higman's finite-word theorem	External standard ingredient; alternatively import the classical Hahn–Neumann support lemma directly.
D-finite / polynomial recurrence conversion	Classical; credited to Stanley and derived explicitly for Hahn coefficients.
Partial theta series and functional identity	Classical; not a novelty claim.
Escape chain and exclusion criterion, with the local-finiteness boundary	Elementary support-theoretic lemma and quantitative formulation developed here.
Eventual affine and periodic orbit valuations	Proved here for arbitrary Hahn coefficients; the finite residue-polynomial treatment covers near-torsion units.
Exact fixed- q order-unit classification	Principal proposed contribution: arbitrary rank, arbitrary Hahn parameter, both signs of $v(q)$.

Result or ingredient	Status in this report
Exact theta evaluation domain	Support-controlled sharpness theorem; related in spirit to the companion report's scalar-extension results, proved independently here.
Uniform exterior obstruction and common degree bound	Proved here; stronger than eventual termination of each solution separately.
Universal single-dilation nonholonomicity on no-order-unit workspaces	Consequence of the classification, with an explicit entire witness.
Mixed differential–dilation rigidity at noncofinal nonzero scales	Proved here; the unit-valuation mixed case is answered without an order unit by source 12 (Theorem E), and for every Γ , for linear equations, by source 18 (Corollary 29.6).
Several-variable rigidity and multivariate algebraic rigidity	Proved here; the generic line is an existence device.
Gauss initial forms, Euler eigenvalues, indicial polynomials, central-index and Newton-polygon ideas	Established (source 10); credited to Hayman, Cano and Fortuny Ayuso, Hu–Luan, Hu–Yang and Giansiracusa–Mereta as predecessors in different settings.
First-order nonlinear rigidity (Theorem D), finite exclusion certificate, candidate degrees and entire-solution locus	Proposed contribution of source 10, over any characteristic-zero coefficient field; priority provisional.
Higher-order corner criterion and positive-weight Euler rigidity	Proposed by source 10; sufficient conditions only; the vanishing-corner case is settled without an order unit by sources 11 and 12, and with one source 13's equation of order three has vanishing residue corner and a nonpolynomial entire solution.
Solution-dependence of the nonlinear exclusion bound	Proved by source 10 with the explicit Riccati family of Theorem 15.10.
Two-jet independence of $\sum t^{n^2} z^n$ and of (11.2)	New consequences (source 10); both series are prior collection material.
Classical coefficient recurrences, finite generation, valuation-rank inequality, Skolem–Mahler–Lech	Established (sources 11 and 12); credited to Hurwitz, Laohakosol–Kongsakorn–Ubolsri, Bell and Lech; the last imported without proof.
All-order rigidity without an order unit, with dilations (Theorem E)	Proposed by sources 11 and 12 (part (b) and the dilations by source 12, part (c) by source 11); priority provisional.
Coefficient-value-rank criterion, order-unit dichotomy, rank-one family	Proposed by source 12; sufficient conditions and one equivalence, as stated.
Coarsening descent, rationality in $\text{Frac } \mathcal{E}_\Gamma(k)$, systems, Painlevé I in $\text{Frac } \mathcal{E}_\Gamma(k)$, joint-independence criterion, continuum family	Proposed by source 11; the polynomial half of the descent is the entire-function report's obstruction, and neither the cardinal nor the almost-disjoint device is new.
All-order independence of (11.2)	New consequence (sources 11 and 12); the series is prior material.

Result or ingredient	Status in this report
Several multiplicatively independent dilations (Corollary 22.8)	The merge’s observation from source 12’s theorem; not claimed by source 12.
Mixed jets in the fraction field (Corollary 22.9)	Consequence added here from source 12’s coefficient theorem and source 11’s descent; neither source states it.
Chebyshev identity, Jacobi triple product, theta-elliptic differential identity	Classical (source 13); DLMF, Sebbar and Ohyama credited; no novelty claim for third-order theta relations.
Bilateral theta domain	The Hahn–Tate report’s theorem, specialized; reproved by source 13, not claimed anew.
Exact domain of $\mathcal{T}_p$, product and zeros, local valuations, omnific rounding	Proposed by source 13, derived from the classical product.
Exact boundary (Theorem H), minimal order three, differential field, non-D-finiteness	Proposed by source 13; the no-order-unit direction is Theorem E (a); priority provisional.
Substitution homomorphism, global proof of $\psi_3 \neq 0$, corner data of the cleared equation, clause (iv) of Theorem G, Corollary 25.2	Added by the merge from source 13’s results; the first two write out steps source 13 used.
Second-order rigidity for every Γ and exact least order three (Theorem I), endpoint lemma, endpoint–gap obstruction, theta initial forms, eventual Newton convexity, two-state systems and polynomial-denominator quotients	Proposed by source 14; the witness of order three is source 13’s and the no-order-unit direction is Theorem E (a); priority provisional.
Cofinal scalar extension, all-scale criterion, surreal full-class boundary	Lemma 26.2 and Proposition 27.6 are source 14’s; the criterion is Proposition 20.2, printed once.
Exterior form as the residue corner in Euler jets; three jets of G and Θ_p	Added by the merge (Remark 26.6, Corollary 26.13).
Periodic-affine valuations of Hahn exponential polynomials, the bivariate unit-orbit estimate, mixed rigidity for noncofinal relative scales, the relative-scale dichotomy (Theorem J)	Proposed by source 18; the Skolem–Mahler–Lech theorem is imported; pure differential, cyclic pure dilation and nonzero noncofinal mixed rigidity are this report’s earlier results, now special cases; priority provisional.
Finite-word lemma with proof	The standard minimal-bad-sequence argument, printed from source 18 (Lemma 28.2); not new.
Second-order rigidity, re-derived	Source 19’s Theorem A is Theorem I (a) of source 14, found independently the same day; printed once, no priority between them asserted.

Result or ingredient	Status in this report
Third order through cubic jet degree (Theorem K), first-polar and factor criteria, secant and eventual-side lemmas, constant-extension, omnific and algebraic two-state consequences, exact classifications of H_3 and Q_4 , the sparse series	Proposed by source 19; priority provisional.
First polar inside the endpoint-gap obstruction, a cubic outside Corollary 27.3, jet degree of the theta equation, comparison of the two-state results	Added by the merge (Remark 30.11, Corollary 31.4, Section 30.3).
Named published open conjecture	None claimed solved; the nonlinear question, answered in order one, without an order unit in all orders, with an order unit by an example of order three, and in order two for every Γ , is a question of this report, not a historical conjecture; so is the mixed unit-valuation question answered by source 18.
Seed, equation and minimal order three of $\mathcal{T}_p$ reused by source 16; theta product, Tate identity	Source 13's and classical; credited, not claimed.
Independence across scales, exact Gauss profile, noncancellation certificate, jet compression, unbounded order, trdeg and dtrdeg of $\mathcal{D}_{DA}$, integer sampling, valuation route to order three	Proposed by source 16, for $\Gamma = \mathbb{R}$; priority provisional.
Single-series gap differential transcendence; normal forms, Berarducci–Mantova derivation, Oz , bounded-support descent	Classical or prior (Grönwall via Ostrowski; [8]; [38]); not claimed.
Exact relation ideals of rapid-height families, pure D_z freedom over the full Hahn field, Hermite data, ∂_{BM} -jets over the bounded-support field, exact domain in $\mathbf{No}[z]$, omnific codes and floors	Proposed by source 17; priority provisional.
Identity of the Tate and Weierstrass seed equations, the known part of the order spectrum, three fields of entire series, three routes to order three, placement of single-scale freedom	Added by the merge (Section 32.1, Remarks 32.18, 33.4, 33.10, 35.2).
Complex entire Mahler lemma, Mahler's method, normal forms and the constant-term retraction, Smith normal forms, the MRDP theorem and Sun's theorem	Classical or prior ([42, 43, 37, 44, 45]; [41]); credited, not claimed.

Result or ingredient	Status in this report
Strict composition-weight rigidity for every Γ (Theorem N), exact-resonance degree bounds, extension-stable coefficient loci, the matrix version, linear omnific solution modules, the universal Euler-projector encoding and its undecidability consequences	Proposed by source 15; priority provisional.
Removal of the order-unit dichotomy by a dominant argument of degree at least two, identification of source 15's indicial polynomials, added Mahler references	Added by the merge (Remark 36.10, Sections 36.1 and 37.5).
Support envelope, unit-root and periodic-affine valuations, mixed unit rigidity, re-derived; exact finite-group criterion	Source 20 found these independently of source 18; printed once as source 18's (Section 38.1); its finite-group criterion and its convex-scale theorem are special cases of Theorem J, printed as Corollaries 39.9 and 39.8.
The omnific stream $\sum_m \mathbf{b}_m \omega^{\omega-m}$	The omnific-notations report's (<code>onot:eq:guardbasic</code>); credited by the merge, not claimed.
One finitely generated observable algebra, Noetherian compression with one residual period, recurrence profiles over $\mathbf{No}[i]$, $\mathbf{Oz}$ and $\mathbf{Oz}[i]$, the decreasing omnific recurrence, the coding reading and the failure of a coefficientwise Skolem–Mahler–Lech theorem, determinantal profiles	Proposed by source 20 (Theorem O); priority provisional.
Two periods, finite generation in the convex-hull criterion, the stream credit, the exact positive-characteristic formula beside the unit-dilation part	Added by the merge (Remarks 38.5, 39.10, 39.6, Example 39.12).
Priority, peer review, formal verification	Priority and independent refereeing are not certified. Lean covers support and escape lemmas, exponential and partial-theta domains, polynomial-orbit valuations, constant-point avoidance, inward stability and torsion covariance. The full classifications and the theorem packages of the nonlinear, coefficient-field, theta-descent, order-three-threshold, unit-dilation, Newton-rigidity, theta-hierarchy, single-scale, polynomial-composition and recurrence parts remain pending.

The targeted literature checks of sources 08 and 09 located the standard D-finite framework, classical complex q -difference growth theory, q -holonomic degree growth in a rational-function setting, and ultrametric q -difference theory, but did not establish whether an identical arbitrary-rank strong-Hahn formulation appears elsewhere. Several broad keyword searches returned irrelevant

material and were not treated as evidence of absence; a repository content search for `holonomic` returning no matches is likewise not proof that no sentence anywhere in the repository can be reformulated into one of these results. Accordingly this is a proposed original result with proofs, not a certified claim of bibliographic priority. Independent mathematical review is still needed; machine-checked scope is the partial coverage recorded in Section 13 and the formalization ledger. The corresponding records for the nonlinear, coefficient-field, theta-descent, order-three-threshold, unit-dilation, Newton-rigidity, theta-hierarchy, single-scale, polynomial-composition and recurrence parts, with their own search limits, are Sections 17.4, 22.6, 25.5, 27.5, 29.7, 31.4, 33.4, 35.6, 37.5 and 39.6.

19.3 What the nonlinear part does not claim

Every limitation stated by source 10, in its manuscript or its two audit files, is kept; they are collected here. They describe source 10's results; where the coefficient-field or theta-descent part goes further, a parenthesis says so. The limitations of sources 11 and 12 are listed separately as (C1)–(C26) in Section 22.7, those of source 13 as (T1)–(T19) in Section 25.6, those of source 14 as (O1)–(O18) in Section 27.6, those of source 18 as (U1)–(U15) in Section 29.8, and those of source 19 as (R1)–(R16) in Section 31.5, those of source 16 as (Th1)–(Th16) in Section 33.5, those of source 17 as (F1)–(F19) in Section 35.7, and those of source 15 as (P1)–(P18) in Section 37.6.

- (N1) Only the first-order case of Question 19.1 is answered, not the unrestricted higher-order question. (Sources 11 and 12 answer the case without an order unit in every order; source 13 shows that with an order unit a nonpolynomial strongly entire series can satisfy an equation of order three; source 14 answers order two for every Γ , and source 19 order three with total jet degree at most three.)
- (N2) Theorem D does not say that a nonpolynomial entire function is differentially transcendental in all orders; independence of f and $D_z f$ does not exclude a relation involving $D_z^2 f$. (Theorem E (a) says it when Γ has no order unit, and Theorem H shows it fails with one; Theorem I adds $D_z^2 f$ to the independent jets for every Γ .)
- (N3) The criterion $\chi_P \neq 0$ is sufficient, not a complete invariant of a solution set, and the methods of source 10 do not decide equations of order at least two with vanishing residue corner. Proposition 16.4 is not a nonpolynomial entire solution and does not refute higher-order rigidity. (Source 13's order-three equation, with $\chi_P = 0$, does have one when Γ has an order unit; Proposition 24.2. Source 14 decides order two negatively for every Γ , Theorem I, and source 19 order three in jet degree at most three, Theorem K.)
- (N4) The corner is a finite selection, not a full differential Newton polygon; no invention of differential Newton polygons, no equivalence to tropical differential algebra, and no new tropical fundamental theorem is claimed.
- (N5) The exclusion bound (15.8) is not claimed optimal; the certificate has no converse; leading valuations alone do not decide evaluation at a boundary argument.
- (N6) Candidate degree sets are necessary, not realized; their finiteness is not a uniform algorithm for arbitrary coefficient names, and the workflow of Section 17.3 is not an algorithm for arbitrary surreal names, does not enumerate local formal solutions, does not decide strong-support membership, and accepts no numerical solver failure as a certificate.

- (N7) Corollary 15.6 does not say that the number of solutions is finite, is not a moduli statement over rings with nilpotents, and covers only the extension class it states.
- (N8) No theorem says that G , F or any nonpolynomial series is entire on the proper class $\mathbf{No}[i]$; entireness and the derivative are relative to one set-sized workspace.
- (N9) The independence statements concern formal functions; no statement concerns algebraic independence of a value $f(x)$ from its own ambient field.
- (N10) D_z kills all Hahn coefficients and is not the Berarducci–Mantova derivation; no scalar-derivation result is claimed.
- (N11) The series $\sum t^{n^2} z^n$ and (11.2), and the linear rigidity of the latter, are prior material; only their nonlinear two-jet consequences are new. The tail-span report’s all-order independence results are neither weakened nor rediscovered.
- (N12) Hahn fields, the positive-support summation lemma, Conway normal forms, Gauss initial forms, Euler eigenvalues and indicial-polynomial reasoning are established ideas, not relabelled as new; the positive-support lemma is imported by source 10, and proved in this report by Lemma 2.1 (2).
- (N13) The precise all-rank statements are proposed contributions; historical priority is not certified. The literature search of 22 September 2026 was targeted and not exhaustive; Hayman’s survey and the Hu–Yang monograph were consulted through metadata only, and no claim is made that rank-one special cases were previously unknown.
- (N14) The repository comparison at 048b72c was targeted: no line-by-line audit of every report, Lean file, archive, branch or the Git history, and no absence claim rests on a truncated preview.
- (N15) The answered question is a question of this report, not a named historical conjecture.
- (N16) Source 10 supplied no Lean formalization or independent referee report for the nonlinear part. The 2,113 finite checks are program checks, not theorems; they do not test infinite supports, cofinality, all Hahn elements, nonexistence of annihilating equations or novelty, and no infinite surreal sum is numerically approximated. A successful PDF build is not a proof check.
- (N17) The sequence arguments behind the support calculus use ordinary set-theoretic choice.

19.4 Further mathematical questions

Question 19.1 (Differentially algebraic entire series of minimal order three). Suppose Γ has an order unit. By Theorem I no nonpolynomial $f \in \mathcal{E}_\Gamma$ satisfies a nonzero algebraic differential equation over $K_\Gamma(z)$ of order at most two, and by Theorem H order three occurs. Which nonpolynomial strongly entire series have minimal differential order three? Are they all accounted for by support-controlled algebraic descents of elliptic or theta constructions, or are there fundamentally different families? More broadly, to what extent do finite algebraic descents of theta or abelian functions, combined with elementary operations preserving strong entireness, account for the nonpolynomial strongly entire series that are differentially algebraic over $K_\Gamma(z)$? The same questions are open over $\mathbb{K} = k((t^\Gamma))$ for every field k of characteristic zero.

Status. As posed by sources 08 and 09, the question asked about nonlinear algebraic differential equations of every order, for every Γ , and ended by asking under which hypotheses on Γ linear rigidity can be strengthened to differential algebraic rigidity in all orders, and whether the explicit series

(11.2) avoids all equations of order at least two under a natural coefficient restriction. Source 10 answers its first-order case negatively, for every nonzero Γ and every characteristic-zero coefficient field (Theorem D): in order one no hypothesis on Γ beyond $\Gamma \neq 0$ is needed, and (11.2) avoids every first-order equation with no coefficient restriction (Corollary 17.2). It also settles the higher-order equations with $\chi_P \neq 0$ (Theorem 16.1) and the relations between f and a positive-weight Euler derivative (Theorem 16.5). After source 10, order at least two with a vanishing residue corner stayed open. Proposition 16.4 shows that such equations exist and can have polynomial solutions of unbounded degree, so the study cannot rely on a uniform degree bound for all polynomial solutions of an arbitrary higher-order equation; it is not a nonpolynomial entire solution and gives no negative answer. Source 10 suggests as a next step classifying the polynomial solutions of a vanishing-corner residue equation beyond their leading monomial and determining how those constraints propagate between scales.

Sources 11 and 12 then answered the question, negatively, in every order for every Γ without an order unit, whatever the residue corner and for every characteristic-zero coefficient field (Theorem E (a)); source 11 extends the answer to quotients of entire series (Theorem E (c)). Their method needs no degree bound: along the solutions of (16.1) its linearization multiplier is $(T - m)^2 \neq 0$ (Example 21.6). The last sentence of the question as posed by sources 08 and 09 is answered positively for (11.2) with no coefficient restriction (Theorem 22.1, source 12; source 11 for $f_* = F - \omega^{-1}$). Its sentence on hypotheses is answered in one direction: the absence of an order unit suffices. The question was therefore re-scoped to groups with an order unit, in the wording recorded below, which merges source 11's Question 9.1 and source 12's Question 10.1; both sources leave it open and give no counterexample. There a finitely generated coefficient field can have values whose convex hull is all of Γ , and the argument of Section 21 stops. The partial theta series Θ_p and G are not counterexamples: Θ_p satisfies a dilation equation, not a known algebraic differential one, and only the first-order consequence is known for G (Corollary 17.1). Theorem G shows that the order unit is exactly the boundary of the *mixed* statement; whether it is also the boundary of the purely differential one was, at that stage, the question.

As re-scoped after sources 11 and 12, the question read: *Suppose Γ has an order unit. Can a nonpolynomial $f \in \mathcal{E}_\Gamma$ satisfy a nonzero algebraic differential equation $P(z, f, D_z f, \dots, D_z^s f) = 0$ over $K_\Gamma(z)$ of order $s \geq 2$? By Theorems D and 16.1, such an f satisfies no nonzero equation of order at most one and no equation with $\chi_P \neq 0$; equivalently, can a nonpolynomial strongly entire function satisfy an equation whose residue corner polynomial vanishes identically, without satisfying any other equation to which Theorem 15.1 applies? More strongly, is every element of $\text{Frac } \mathcal{E}_\Gamma$ that is differentially algebraic over $K_\Gamma(z)$ rational? Is the absence of an order unit necessary for differential algebraic rigidity in all orders, or can linear rigidity be strengthened to it for every nonzero Γ ?* Source 13 (the theta-descent part) answers every sentence of it, for every characteristic-zero coefficient field, including $k = \mathbb{C}$. With an order unit μ and $p = t^\mu$, the series $\mathcal{T}_p$ of (1.4) is nonpolynomial and strongly entire and satisfies an explicit equation of order three and none of lower order (Theorems H, 24.1 and 24.10), so the answer to the first sentence is yes. Its equation has $\chi_P = 0$, and $\mathcal{T}_p$ satisfies no equation with $\chi_P \neq 0$ (Proposition 24.2, Remark 24.3), so the answer to the second is also yes. $\mathcal{T}_p$ is a differentially algebraic element of $\text{Frac } \mathcal{E}_\Gamma$ that is not rational, so the rationality question has a negative answer. The absence of an order unit is necessary: linear rigidity (Theorem A) cannot be strengthened to differential algebraic rigidity for any Γ with an order unit (Corollary 25.2). Source 13 leaves open whether order exactly two occurs and asks for a classification (its Questions 11.1 and 11.2); the question was re-scoped to these, in the wording recorded in the next paragraph. Neither Θ_p nor G is decided: whether either is differentially algebraic stays open.

As re-scoped after source 13, the question read: *Suppose Γ has an order unit. Can a nonpolynomial $f \in \mathcal{E}_\Gamma$ satisfy a nonzero algebraic differential equation $P(z, f, D_z f, \dots, D_z^s f) = 0$ over $K_\Gamma(z)$ of order $s = 2$, that is, are f , $D_z f$ and $D_z^2 f$ algebraically dependent over $K_\Gamma(z)$ for some nonpolynomial $f \in \mathcal{E}_\Gamma$? By Theorem D no equation of order at most one is possible, and by Theorem H order three occurs, so the least order over all examples is two or three. More broadly, to what extent do finite algebraic descents of theta or abelian functions, combined with elementary operations preserving strong entireness, account for the nonpolynomial strongly entire series that are differentially algebraic over $K_\Gamma(z)$? The same questions are open over $\mathbb{K} = k((t^\Gamma))$ for every field k of characteristic zero.* Source 14 (the order-three-threshold part) answers the first sentence negatively, for every Γ , with or without an order unit, and for every characteristic-zero coefficient field, including $k = \mathbb{C}$: a strongly entire solution of a nonzero equation of order at most two is a polynomial (Theorem I(a)), even when its residue corner vanishes. So the least order is exactly three (Theorem I(b)). Source 14 was written against the repository at [b895e86](#), before source 13 was merged, and answers source 13's question `q:order2`, which is the same question (Section 26.1). The classification sentence stays open; source 14 poses it again as its Research question 1, which is merged here, and the question is re-scoped to it, as printed above. For Θ_p and G the answer gives three independent jets (Corollary 26.13); whether either satisfies an equation of order at least three stays open. The eight further questions of source 14 are Questions 19.11–19.18 below.

Status (batch 31). Source 19 re-derives the order-two answer independently (Section 30.1) and narrows the classification from below: a series of minimal order three satisfies no equation of order three and total jet degree at most three (Theorem K), and the exterior form of every equation of order three that it satisfies has vanishing first polar (Corollary 31.4); for $\mathcal{T}_p$ the least such jet degree is four, five or six. Source 19's own Question 12.1, the order-unit case of this question in all orders, is answered by source 13, and its abstract's “unresolved” is stale against the current text. The classification sentence stays open, and the question keeps its wording.

Status (batch 31, sources 16 and 17). Source 16 shows, for $\Gamma = \mathbb{R}$, that minimal orders are unbounded: polynomially weighted sums of theta descents at reciprocal-independent scales have orders in $[N, 3N]$ for every N , and the differentially algebraic strongly entire series generate a field of transcendence degree $2^{\aleph_0}$ and differential transcendence degree zero (Theorem L). These are finite algebraic combinations of theta descents under operations preserving strong entireness, as the last sentence of the question contemplates; so the minimal orders form an unbounded set containing three (Remark 32.18), and which orders above three occur is Question 19.36. Source 17 shows that the same order-unit fields carry explicit nonpolynomial strongly entire series with coefficients in $\mathbb{Q}(t^\mu)$ satisfying no algebraic differential equation (Theorem M); its Question 12.1, this question in the form posed before source 13, is answered by Theorem H. Neither source decides Θ_p or G (Question 19.45). The question keeps its wording.

The escape proof does not answer this, in any order. A nonlinear differential relation produces convolution sums in the coefficients rather than a bounded-step linear recurrence, and treating it as though it had the form (3.1) would be an invalid extension. The first-order answer uses the different route of Section 14, the no-order-unit answer the third route of Section 21, and the order-two answer a fourth, initial polynomials at infinitely many scales (Section 26.3).

Question 19.2 (Beyond a positive Euler operator). Which polynomial vector fields V have the property that every strongly entire f satisfying a nonzero equation $P(x, f, Vf) = 0$ is a polynomial?

Positive integer Euler weights give one family, and zero and mixed weights give explicit counterexamples (Section 16.3). A useful extension would need both a finite-dimensional filtration and

a sufficiently nondegenerate action on its highest layers; diagonalization alone is not enough if infinitely many monomials share a bounded eigenvalue. (Source 10.)

Question 19.3 (Effective representations and minimal certificates). For which exact Hahn coefficient representations can the candidate degree set and an optimal excluded valuation cut be computed uniformly?

The bound (15.8) is always a valid choice from exact finite valuation data but can be far from minimal, and Theorem 15.10 shows that an optimal cut must in general depend on solution data. An algorithmic refinement must state how it receives those data and how it decides coefficient cancellation. (Source 10; the linear analogue is Question 19.5.)

Question 19.4 (Mixed operators at unit valuation, with an order unit). Suppose Γ has an order unit. For a nontorsion q with $v(q) = 0$, must every strongly entire solution of a nonzero mixed equation (10.1) be a polynomial?

As posed by sources 08 and 09 the question concerned every Γ . The unresolved point in the escape method is uniform control of $v(P(n, q^n))$ for a general bivariate P in the unit case. The pure q -difference proof controls $P(q^n)$ and the differential proof controls $P(n)$; neither statement by itself handles their combination. This is a question suggested by the argument, not asserted to be a previously catalogued open problem.

Status. When Γ has no order unit, source 12 answers it positively and more strongly: by Theorem E (b) with $\Lambda = \{q^i : i \in \mathbb{Z}\}$, a nonpolynomial $f \in \mathcal{E}_\Gamma$ satisfies no nonzero algebraic relation, linear or not, among the $(D_z^j f)(q^i z)$, and a nonzero equation (10.1) is such a relation (Section 22.5). The method bounds the coefficient field instead of the valuations $v(P(n, q^n))$. The question is re-scoped to groups with an order unit, where it stays open: the witness of Theorem G uses a parameter of nonzero valuation, and source 13's series $\mathcal{T}_p$, which gives nonlinear mixed relations for every nontorsion q (Corollary 25.2), is not D-finite, and source 13 says nothing about linear mixed equations.

Status (batch 31). Source 18 answers the question positively, for every Γ with or without an order unit and over every characteristic-zero coefficient field: for nontorsion q with $v(q) = 0$ every strongly entire solution of a nonzero equation (10.1), also with polynomial or rational forcing, is a polynomial, with a uniform degree bound and exterior obstruction for each equation (Corollary 29.6, Theorem J). The obstruction named above, uniform control of $v(P(n, q^n))$, is supplied by Corollary 28.7: the valuation is eventually periodic, with period dividing the order of $\text{res}(q)$ when that is a root of unity. The question is answered in the form posed; the nonlinear analogue is settled negatively by Corollary 25.2, as recorded before.

Status (batch 33). Source 20 re-derives the answer independently, for multipliers in a torsion-free group, through the same valuation theorem for Hahn exponential polynomials (Section 39.4); it was written against the text before source 18 was merged, when the question was still open there.

Question 19.5 (Optimal exterior regions and exact solution domains). For a fixed differential or rigid dilation operator, can one compute the largest common strong evaluation domain of its formal solution space from finite valued operator data, rather than only a sufficient exclusion bound? Can the coarse exclusion bound of Theorem 7.4 be supplemented by support hypotheses that give the exact domain?

The thresholds supplied here are uniform and explicit once the valuation certificates are available, but need not be optimal, and distinct solutions may have cancellations or terminating tails that change their individual domains. Proposition 5.3 shows that the uncoarsened bound can be exactly attained, and Theorem 8.1 exhibits one sufficient pattern for an exact answer: a common positive support monoid together with a controlled sequence of leading exponents. A full answer should retain arbitrary Hahn tails rather than use only a numerical radius.

Question 19.6 (Several independent dilations). When Γ has an order unit, what is the existence criterion for a nonpolynomial strongly entire solution constrained by equations using several multiplicatively independent dilations? Is an appropriate cofinality condition on their joint valuation data sufficient, or do compatibility relations impose additional obstructions? In the no-order-unit workspace (11.1), can one classify the finite linear relations among $F(q_1z), \dots, F(q_s z)$ when some quotient q_i/q_j is a root of unity?

Status. As posed by sources 08 and 09 the first two sentences concerned every Γ , and the last asked about parameters that are not powers of a common element. In a group without an order unit the merge observes (Corollary 22.8, from source 12's Theorem E (b); source 12 does not claim it) that no nonpolynomial strongly entire series satisfies any nonzero equation, linear or not, in finitely many derivatives at dilations from the group generated by multiplicatively independent parameters, and that $F(q_1z), \dots, F(q_s z)$ satisfy only the trivial linear relation whenever no quotient q_i/q_j is a root of unity, whether or not the q_i are powers of a common element. The question is re-scoped to what this leaves: groups with an order unit, and parameters with a root-of-unity quotient.

Status (batch 31). For linear equations source 18 answers the existence half of the first sentence, for every Γ , in the following form. For a finite set of multipliers with no root-of-unity quotient, some nonzero mixed linear equation supported in the set has a nonpolynomial strongly entire solution if and only if $|v(\lambda_i/\lambda_j)|$ is an order unit for some pair; two multipliers and first derivatives then suffice, and otherwise every such equation, with polynomial forcing, has only polynomial entire solutions (Theorems 29.4 and 29.9). So the appropriate cofinality condition is the relative one, and a common absolute scale is irrelevant. This is a partial answer (Remark 29.10): it concerns the existence of *some* equation, not the solvability of a prescribed operator (Question 19.19) or of simultaneous equations (Question 19.20), and it keeps the exclusion of root-of-unity quotients (Question 19.21). The question stays open in those respects, and its last sentence, on root-of-unity quotients in Γ_∞ , is untouched.

Status (batch 31, source 15). If an equation also contains a nonzero operator applied to $f(P(z))$ with $\deg P \geq 2$ strictly above the degrees of all other arguments, every strongly entire solution is a polynomial, for every Γ and whatever the dilations (Corollary 36.9, Remark 36.10). The question concerns arguments of degree one and is otherwise unaffected.

The single-dilation theorem identifies the exact obstruction when only one scale is repeatedly added. With several dilation directions, distinct monomials can have equal valuation slopes and their leading coefficients can produce nontrivial recurrence phenomena of their own. A meaningful formulation must also quotient out identities caused by repeated parameters and any genuine algebraic relations among them. No multidilation classification is implied by Theorem B.

The next three questions come from sources 11 and 12.

Question 19.7 (An intrinsic descent criterion; source 11). For a fixed Hahn field $\mathbb{K}$ and a subfield $L \subseteq \mathbb{K}$, can the equality $\text{Frac } \mathcal{E}_\Gamma(k) \cap L((z)) = L(z)$ be characterized solely in terms of the valued extension $\mathbb{K}/L$?

Containment of $v(L^\times)$ in a proper convex subgroup is sufficient (Theorem 21.11) and applies to many infinitely generated fields; it is not claimed necessary. A criterion using rational rank alone will not capture all cases: what matters to the reduction is whether the coefficient values reach every Archimedean scale of Γ .

Question 19.8 (A finite-rank replacement; source 12). Which additional properties of a finitely generated coefficient field, of its valuation, or of the recurrence (21.5) force a linear upper bound on the relevant nonzero coefficient values, and so exclude strong entireness even when those values are cofinal?

Bounding the multiplier $\Xi(n)$ is not enough: a nonlinear recurrence can involve cancellations among many products of earlier coefficients, and an extension must control them or replace them by another invariant. An answer would bear on Question 19.1. The series $\mathcal{T}_p$ of Theorem H is differentially algebraic and strongly entire, with a finitely generated coefficient field and coefficient values $m^2\mu$ that are cofinal and not linearly bounded; with an order unit, any such property must therefore fail for it (Section 25.2).

Question 19.9 (Torsion blocks; source 12). When some dilations have root-of-unity quotients, can the coefficient-field conclusion of Theorem 21.2 (b) be recovered separately on the finitely many residue classes of Taylor indices that survive the resonances?

The even-series example of Section 22.4 shows that no unconditional version for all coefficients exists; a blockwise theorem would need explicit conditions on which residue classes are constrained, not only on the nonzero part of the multiplier.

The next question comes from source 13 (its Question 11.3).

Question 19.10 (Effective strong domains for differential solutions). For an explicitly represented formal solution of an algebraic differential equation, can its maximal strong domain be recovered from a finite differential system and coefficient data?

For $\mathcal{T}_p$ the answer is the coarsened valuation ring $\mathcal{O}_{\Delta_\mu}$ (Theorem 23.5), whose boundary depends on the convex subgroup generated by one scale. General solutions can have much more complicated supports. An algorithm needs a representation of those supports and a decision procedure for the relevant coefficient cancellations; inspecting the leading differential monomials cannot suffice in the vanishing-corner case. Compare Questions 19.3 and 19.5.

The next eight questions come from source 14 (its Research questions 2–9; its Research question 1 is merged into Question 19.1).

Question 19.11 (Lifting admissible initial polynomials; source 14). Given an equation of order three and an unbounded sequence of initial polynomials satisfying its exterior form, what compatibility conditions guarantee that they are the initial polynomials of one strongly entire solution?

A criterion would have to control both coefficient lifting and global support: local formal solutions at separate scales need not assemble into one series. $\mathcal{T}_p$ is a fully worked positive case, with the initial polynomials $X^n(1+X)$ at its breaks.

Status (batch 31). Source 19 gives a concrete negative instance (Example 31.3): the initial polynomials of the strongly entire series F_{sp} satisfy the exterior forms of H_3 and of Q_4 at every scale, yet F_{sp} solves neither equation, the first failure being the coefficient $-126 t^{17}$ of z^5 in $Q_4(F_{\text{sp}})$. So agreement of initial polynomials at every break is not a sufficient compatibility condition, even when those polynomials come from one entire series. Question 19.30 asks for the higher-layer conditions; the question stays open.

Question 19.12 (The entire solutions of the theta equation; source 14). For the constants (24.2), determine all strongly entire solutions of (24.7). Which are scalar multiples, coordinate transforms or other members of a theta family, and which come from singular branches of the cleared equation?

Theorem 27.5 forces eventually full coefficient support and strictly increasing consecutive coefficient differences. The next step would use the lower exterior terms, discarded by the exterior form, to constrain those differences and the leading coefficients; the cleared form may have exceptional solutions not represented in logarithmic coordinates, which need separate treatment. Since (24.8) is homogeneous in the jets, every constant multiple of $\mathcal{T}_p$ is a solution.

Question 19.13 (Effective third-order exclusion; source 14). For which coefficient fields and equations can boundedness of the nonnegative-integral endpoint projection of $\Upsilon_{\mathfrak{h}_P}(T, G) = 0$ be decided? Can a finite library of classifications of residue polynomials extend the certificate of Corollary 27.3?

At order two the construction is finite algebra once the leading coefficient values are accessible: dehomogenize, substitute, remove a power of V and inspect the remainder on $V = 0$. Order three introduces a genuine two-integer constraint. Any algorithmic claim must include computability of the Hahn coefficients of the input and of their exact leading terms; compare Question 19.3.

Status (batch 31). Source 19 extends the certificate in one direction: when the exterior form has a nonzero first polar, that is $i_0 = 1$, the two endpoint conditions cannot both hold far out, whatever the upper-endpoint set of Corollary 27.3 is (Proposition 30.6, Remark 30.11); the test applies factor by factor (Corollary 30.9), and it settles every equation of order three and total jet degree at most three (Theorem K). The quartic H_3 , with $i_0 = 2$ and $\Upsilon = (G + T)(2G - T)$, shows that the test fails in degree four; its formal solutions are classified exactly (Theorem 31.1). The question stays open for $i_0 \geq 2$.

Question 19.14 (Higher-order break invariants; source 14). Can the expansion transverse to constant logarithmic derivatives be organized into a hierarchy of gap and multiplicity spectra for orders four and above, with rigidity theorems at each level?

At higher orders several logarithmic derivatives vanish to the same first local order, with leading ratios determined by powers of an endpoint gap; secondary terms could record further support spacings. When the resulting constraints are complete is open.

Status (batch 31). Source 19's quadratic Q_4 of order four is a first worked case: its cross coefficient is the pair kernel $(m - n)^2(m - 2n)(n - 2m)$, and its formal solutions are exactly the monomials and the doubling binomials (Theorem 31.2); Question 19.31 asks for the forms with such sparse solutions in general.

Question 19.15 (Meromorphic second-order solutions; source 14). Beyond polynomial denominators, which algebraic differential equations of order two have nonrational quotients of strongly entire series as solutions? Can the break method bound the density or the valuation distribution of their poles?

Elliptic meromorphic functions suggest that polynomiality cannot simply be extended to every quotient of entire series when an order unit exists; the comparison involves separate Newton data for numerator and denominator, including cancellation and common zeros. Without an order unit, Theorem E (c) already makes every differentially algebraic element of $\text{Frac } \mathcal{E}_\Gamma(k)$ rational.

Source 19 asks the same question (its Question 12.4), for equations of order at most two and elements of $\text{Frac } \mathcal{E}_\Gamma(k)$ in the order-unit case, and notes that the initial object of a quotient is rational rather than polynomial, so the first- and last-support arguments need replacement.

Question 19.16 (Partial differential systems; source 14). Is there a useful several-variable analogue of three-jet independence for series strongly summable at every point of $\mathbb{K}^d$? Which finite-dimensional rational differential systems force polynomiality of all strongly entire observables?

Restriction to lines may settle special cases, as for Theorem C, but algebraic relations among several partial derivatives do not in general reduce to a relation among the first two derivatives on every line; a several-variable analysis of initial forms is needed.

Source 19 asks the same question (its Question 12.6) for a positive-weight Euler operator: a leading weighted component is then a homogeneous polynomial rather than a multiple of one monomial, the conic factorization suggests a route, and diagonal specialization can hide cancellation, so a multivariable break statement should come first.

Question 19.17 (Positive residue characteristic; source 14). Can a replacement for second-order rigidity be formulated with Hasse or divided derivatives, after excluding series that are constant for the Frobenius and specifying the behaviour of integer valuations?

The example of Section 27.4 and Example 17.3 show that ordinary derivatives cannot suffice. A replacement must distinguish the characteristic of the field, that of the residue field and the notion of summation.

Source 19 asks the same question (its Question 12.8), adding that neither the conic cross coefficient nor the double factor of the first polar transfers verbatim, and that equal characteristic zero, mixed characteristic and positive characteristic must be distinguished; source 18's version for the unit theorem is Question 19.26.

Question 19.18 (Omnific approximation of differential zero sets; source 14). For explicitly represented entire solutions of minimal order three, which zero sets admit uniform approximation by omnific integers, and how do endpoint gaps constrain the valuations of the rounding errors?

The zeros of $\mathcal{T}_{\omega-1}$ have exact rounding errors $\omega^{-(2m-1)}$ (Corollary 25.3). For other entire solutions the coefficient valuations are not enough: residue directions and higher terms can move individual zeros.

The next eleven questions come from source 18 (its Research questions 11.2–11.12; its Research question 11.1 is answered in the current text, Remark 29.19).

Question 19.19 (Prescribed operators at a cofinal quotient; source 18). For a specified mixed operator L whose multipliers cross the top valuation scale, which conditions characterize the existence of a nonpolynomial strongly entire solution of $Lf = 0$?

Theorem 29.9 lets one choose L and does not answer this. A first target is a two-multiplier operator of differential order one: its recurrence has explicit valuation slopes, and one must decide whether compatible solutions have coefficient valuations escaping fast enough, not merely whether a formal solution exists.

Question 19.20 (Simultaneous equations; source 18). Can a nonpolynomial strongly entire series satisfy two independently prescribed mixed equations whose cofinal dilation parameters are multiplicatively independent? Which compatibility conditions on their recurrences are decisive?

The witness for one pair of multipliers need not satisfy an equation for another pair. This sharpens Question 19.6; a starting point is to compare the two induced recurrences before seeking a general difference–differential Galois theory.

Question 19.21 (A torsion-block criterion for linear mixed rigidity; source 18). When some multiplier quotients are roots of unity, can Theorem 29.4 be replaced by a finite block criterion deciding which residue classes of Taylor indices are forced to terminate?

Theorem 28.10 already allows identically zero progressions; the difficulty is that recurrence jumps can move between classes on which the leftmost coefficient vanishes. A finite directed graph of residue classes with nonvanishing shift data on its edges is a plausible tool, but every surviving infinite path needs an escape inequality; checking classes in isolation is not enough. Question 19.9 asks the analogous question for the coefficient-field conclusion.

Status (batch 33). Source 20 poses the same question (its Question 12.2) for a finitely generated dilation group with torsion: decompose formal series by degree modulo the torsion order, account for the sector shifts caused by derivatives and polynomial coefficients, and distinguish a genuinely constrained sector from the free even series of Section 39.5. It stays open.

Question 19.22 (Several variables without an ellipticity shortcut; source 18). For jointly strongly entire series in several variables, which finite systems of partial differential–dilation equations admit an analogous exact relative-scale obstruction?

A single partial derivative may annihilate an arbitrary function of the other variables, so the one-variable statement does not extend verbatim. The hypothesis should constrain all escaping multi-indices, perhaps through finitely many positive-weight directions or a rank condition; restriction to a generic line, as for Theorem C, would need the operator system and joint strong summability to be preserved. Compare Question 19.16.

Status (batch 33). Source 20’s Question 12.12, on finite linear substitutions in several variables and matrix dilations, is the neighbouring Question 19.70.

Question 19.23 (Optimal exterior cuts for mixed operators; source 18). Given the exact periodic-affine valuation data of a mixed operator, can one compute the largest common strong evaluation domain of its formal solutions?

The jump-sensitive inequalities of Theorem 3.3 are sufficient; optimality would require knowing which recurrence edges actual solutions realize and how their nonleading supports behave, and valuation data alone may not decide it. A first problem is an operator with monomial coefficients and a fixed finite jump graph. This is the mixed form of Question 19.5.

Question 19.24 (Effective coefficient representations; source 18). For which finitely specified Hahn subfields can the support envelope, the first coefficient sequence that is not an identity, and all exceptional indices in Theorem 28.10 be computed?

A first target is Laurent or Puiseux polynomial data with finite-order residue quotients, where the zero tests reduce to polynomial calculations once a support bound is proved; arbitrary residue bases need a separate exact zero-set analysis. No universal algorithm follows from the qualitative theorem.

Status (batch 33). Source 20 poses the same question (its Question 12.1) for Hahn inputs with algebraic coefficients and finitely generated, explicitly represented supports, and asks to separate

the search for the first coefficient sequence that is not an identity from the location of the zeros of that exponential polynomial, which the existence proof does not eliminate. It adds a target of the same kind: Theorem 38.8 gives a finite equivalent subfamily of a set-sized family of coefficient conditions only existentially (Section 39.5). Both stay open.

Question 19.25 (Valuation profiles beyond Hahn fields; source 18). Which equicharacteristic-zero valued fields have the periodic-affine property of Theorem 28.10 for exponential polynomials, without a specified Hahn expansion?

A support-preserving embedding into a Hahn field would transfer the result when available, but finding sufficient hypotheses for it is part of the problem; an intrinsic proof needs substitutes for the well-ordered set of candidate coefficients and for the least coefficient sequence that is not an identity. No embedding theorem beyond the explicit surreal workspaces is asserted.

Status (batch 33). Source 20's Question 12.5 is the same question: which embedding or completion hypotheses preserve the conclusion without unjustified summability assumptions, so that the genuinely Hahn-theoretic part of the theorem is isolated. It stays open.

Question 19.26 (A positive-characteristic replacement for the unit theorem; source 18). Can Frobenius-sensitive index sets and a modified notion of differential operator give a useful replacement for Theorem 28.5 and Corollary 29.6 in characteristic $\mathfrak{p}$?

Ordinary derivatives cannot suffice (Section 29.5). Hasse derivatives and automatic zero-set patterns [33] are natural ingredients, but unbounded valuations along $\mathfrak{p}$ -power indices must be treated explicitly. Compare Question 19.17.

Question 19.27 (Scalar extension of the obstruction; source 18). For an ordered-group extension $\Gamma \subseteq \Gamma'$, which entire solution spaces survive the passage from $k((t^\Gamma))$ to $k((t^{\Gamma'}))$?

Rigidity survives whenever the relative scales stay noncofinal; a cofinal quotient in Γ may stop being cofinal in Γ' , and its theta witness then loses entireness. An exact domain theorem should track full support certificates, not only the extension of the coefficient field; the enlargement of G_μ in Section 29.4 is a first example.

Question 19.28 (Arithmetic restrictions on omnific operators; source 18). Do operators with coefficients in an omnific subring admit stronger bounds on polynomial solutions, or more effective valuation certificates, than general Hahn-coefficient operators?

The theorems allow such coefficients but use neither their one-sided supports nor their integer constant terms. A first task is to study finite-support omnific coefficients and determine which cancellation profiles in Theorem 28.10 occur; any improvement must account for $\sigma_2 - 2^a$ (Example 29.17), which already has ordinary integer coefficients.

Status (batch 33). Source 20 continues the question on the side of sequences rather than operators. The leading exponent of a recurrence over $\mathbf{Oz}$ or $\mathbf{Oz}[i]$ is eventually affine on finitely many progressions, two sufficing when the characteristic roots are real (Theorem 39.1); for a real-surreal unit-root recurrence, support conditions are decided by finitely many coefficients along the orbit, with hitting sets periodic modulo M_{tor} up to a finite set (Corollary 38.9); but one fixed Conway coefficient of the purely infinite orbit $y_{n+1} = \omega y_n$ can be any real sequence (Theorem 39.3). Full omnific membership is Question 19.64. The question about operators stays open.

Question 19.29 (A proof-assistant implementation with an explicit trust boundary; source 18). Can the finite-order-residue case and the two-multiplier witness be formalized first, and the general theorem once the Skolem–Mahler–Lech input is available in verified form or as an explicit hypothesis?

This separates a support and order component, a finite algebraic component and a number-theoretic zero-set component, and keeps finite testing, imported theorems and kernel-checked proof apart.

Status (batch 33). Source 20’s Question 12.11 proposes the same separation for its chain: formalize Theorem 38.2 with Lemma 28.4, the argument by the least coefficient sequence that is not an identity, and the escape lemma, keeping the Skolem–Mahler–Lech theorem as a declared imported theorem until it is available; the repository’s support and recurrence-escape infrastructure would need a theorem-label audit before being reused as checked coverage. No Lean work is added.

The next six questions come from source 19 (its Questions 12.2, 12.3, 12.5, 12.7, 12.9 and 12.10; its Question 12.1 is answered by Theorem H, and its Questions 12.4, 12.6 and 12.8 are Questions 19.15, 19.16 and 19.17).

Question 19.30 (Higher residue layers and differential prolongation; source 19). Is there a finite collection of higher-layer compatibility conditions that excludes every nonpolynomial entire lift of an exterior form whose initial polynomial solutions are not eventually monomials?

Example 31.3 is a concrete test: all initial polynomials of F_{sp} satisfy both boundary equations, while the coefficient (31.6) detects a later obstruction. In the quartic case, differentiating a common-root parameter produced the decisive identity (31.3); a general version might combine breaks with differential elimination and separants. This refines Question 19.11.

Question 19.31 (Jet-curve ideals and sparse polynomial solutions; source 19). For the rational normal curve $\mathbf{m}_r(T) = (1, T, \dots, T^r)$, classify the homogeneous forms $\mathfrak{h}$ for which $\mathfrak{h}(\phi, \vartheta_X \phi, \dots, \vartheta_X^r \phi) = 0$ has nonmonomial polynomial solutions of arbitrarily large lowest degree.

Relevant invariants include the multiplicity of vanishing along the curve, first and higher polars and secant varieties. The conic’s principal ideal is unusually rigid, and $\mathbf{H}_3$ shows that singularity along the twisted cubic allows unbounded doubling supports. An effective criterion on $\mathfrak{h}$ is the useful outcome, not merely a classification of its zero set. Compare Questions 19.13 and 19.14.

Question 19.32 (Systems in three or more variables; source 19). Which strongly entire solutions of finite-dimensional rational or algebraic first-order systems must be polynomial? In particular, with an order unit, do three states suffice to produce a nonpolynomial strongly entire observable?

The planar result follows from three-jet independence. Without an order unit every number of states is covered (Corollary 21.16); with one, four states suffice to produce $\mathcal{T}_p$ (Section 30.3), and three are open. Dimension three would need higher-order independence or a bound on the degree of an eliminated relation, since Theorem K applies only to relations of total degree at most three and such a bound cannot be assumed from the dimension. (The second sentence is the merge’s re-scoping.)

Question 19.33 (Effective certificates and bounds; source 19). Given a finitely described differential equation and coefficient profile, can one compute an exterior non-entireness certificate for every nonpolynomial formal solution?

The corner inequalities are finite and explicit but can depend on a high nonzero coefficient of the particular solution, and uniform degree bounds are impossible in general by (16.1). A realistic target is a solution-dependent certificate with a finite exceptional-degree analysis where the monomial test does not vanish identically. Compare Questions 19.3 and 19.13.

Question 19.34 (Omnific normalization with presentations; source 19). For finitely presented surcomplex constants, when can the multiplier ω^ρ of Lemma 30.14 be computed with a useful bound on the complexity of ρ ?

Existence uses only a set-sized bound on supports; effective normalization needs representations of those supports and an algorithm bounding them, without confusing coefficient clearance with effective global surreal computation.

Question 19.35 (Formal verification and independent review; source 19). Can the conic and first-polar arguments be checked in Lean using only the repository’s stated Hahn interfaces, with every constant extension and universe change explicit?

The finite algebra (conic kernel, bottom and top first-polar coefficients, double factor, eventual side, secant Jacobian) is small enough to isolate first; the break construction, the sign-sensitive inequality of the exterior reduction and the assembly would follow, and finite descent should precede the transfer to $\mathbf{No}[i]$. Independent review of the break construction and the eventual-side lemma should precede broader claims.

The next nine questions come from source 16 (its Questions 11.1 and 11.3–11.10; its Question 11.2 is answered by Theorem I, Remark 32.18).

Question 19.36 (The exact order spectrum; source 16). Which integers occur as $\text{dord}(f)$ for nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ differentially algebraic over $\mathbb{K}(z)$? Does every integer $r \geq 3$ occur?

Theorem L gives unboundedness and realizes the dimensions d_N ; neither identifies the spectrum, which would need control of derivative interactions beyond leading Gauss growth. By Theorem I the spectrum lies in $\{3, 4, \dots\}$, and it contains 3 (Remark 32.18, merge). Compare Question 19.1.

Question 19.37 (Joint jet independence; source 16). For scales with $\mathbb{Q}$ -independent reciprocals, are the $3N$ series $\mathcal{T}_{t^{\mu_i}}, D_z \mathcal{T}_{t^{\mu_i}}, D_z^2 \mathcal{T}_{t^{\mu_i}}, 1 \leq i \leq N$, algebraically independent over $\mathbb{K}(z)$, that is, is $d_N = 3N$?

A positive answer would give exact orders $3N$ through Theorem 32.19. The one-seed difference argument does not apply scale by scale: dilating the shared variable changes all seeds at once.

Question 19.38 (Commensurable scales; source 16). Classify the algebraic relations among $\mathcal{T}_{t^{\mu_1}}, \dots, \mathcal{T}_{t^{\mu_N}}$ when the reciprocals are $\mathbb{Q}$ -dependent. Is there a criterion by theta transformations or isogenies describing the exceptional configurations exactly?

The present condition is sufficient, not necessary: its failure lets two monomials of a relation share their quadratic growth, and produces no relation. A classification must use the bounded and linear terms or further elliptic structure.

Question 19.39 (Efficient compression; source 16). Can the bound $(N+1)^{N^2}$ be reduced substantially, can simpler weights be prescribed for the theta family, and can a good candidate be certified from a finite computable presentation of the relevant Kähler differentials?

The universal argument ignores the structure of theta functions and is probably wasteful for them; an effective version needs a certified representation of jet relations, not a numerical search.

Question 19.40 (Smaller exponent groups; source 16). Over $\mathbb{C}((t^{\mathbb{Q}}))$, $\mathbb{C}((t^{\mathbb{Z}}))$ or another rank-one group, are the minimal orders of differentially algebraic strongly entire series still unbounded? Can large independent families be produced without $\mathbb{Q}$ -independent real growth constants?

The mechanism needs rationally independent reciprocal scales; embedding a smaller field in $\mathbb{K}$ does not answer the question, since the base field and the coefficient field of the constructed series change.

Question 19.41 (Beyond quadratic profiles; source 16). Which differentially algebraic strongly entire series have profiles $\gamma_{\delta}(f) = -c\delta^{\varpi} + o(\delta^{\varpi})$ with $\varpi \neq 2$, and can a Legendre-transform analysis give further independent families and sharper order bounds?

Theorem 32.4 is ready for any $\varpi > 1$; what is missing are differentially algebraic seeds. Fast coefficient decay alone gives entire series, not differential algebraicity (compare Theorem 35.1).

Question 19.42 (Arbitrary value groups; source 16). Which version of the hierarchy survives in higher-rank Hahn fields with an order unit, and what replaces the real-valued growth comparison? Can a finite dominance certificate be stated in the ordered group without embedding a subgroup in $\mathbb{R}$?

A generalization should isolate the dominating Archimedean quotient and check that coefficient values in lower convex layers do not spoil the comparison.

Question 19.43 (Intrinsic derivations and global domains; source 16). Is there an order hierarchy for an intrinsic surreal derivation that moves the coefficients? Separately, which identities survive beyond the real-exponent workspace without inadmissible strong sums?

These are distinct problems, one changing the derivation and the other the domain; neither is solved by renaming t .

Question 19.44 (Omnific sampling beyond the ordinary integers; source 16). For which explicit infinite subsets of $\mathbf{Oz}$ does a nonzero entire series of a fixed Hahn workspace have only finitely many zeros? Can the residue argument of Theorem 33.5 be replaced by finitely many residue tests at varying leading scales?

Theorem 33.5 uses distinct ordinary residues; general omnific inputs can share a residue or leave the workspace, and any extension must fix both scales and domain.

The next ten questions come from source 17 (its Questions 12.2–12.11; its Question 12.1 is answered in this report, Remark 35.20).

Question 19.45 (Pure differential freedom below rapid ratios; source 17). Which height sequences make every infinite subseries $\sum_{n \in A} p^{b_n} z^n$ differentially transcendental over $\mathbb{K}(z)$ for D_z ?

Quadratic heights are the instructive case: the series are entire, the mixed theorem is obstructed by Proposition 35.19, and for $A = \mathbb{N}_{\geq 2}$ the series is G_{μ} without its first two terms, a rescaled partial theta series, whose differential algebraicity is the open point recorded after Corollary 26.13 and in Question 19.1. Fixed-ratio exponential heights also fall outside the band argument.

Question 19.46 (An exact mixed-height criterion; source 17). Is mixed differential freedom characterized by a combination of bounded-degree additive separation of the $\mathbf{b}_n$ and Zariski density of the points $(n, \mathbf{b}_n)$?

The hypotheses are sufficient, not necessary; a useful theorem should distinguish failure of the proof from an actual equation.

Question 19.47 (Infinite coefficient alphabets; source 17). For pattern vectors $\mathbf{v}_n$ taking infinitely many values, what replaces the finite pattern matrix?

$\mathbf{v}_n = (1, n)$ shows that the invariant must see differential shifts; one candidate is the eventual linear geometry of the vectors $n^a \mathbf{b}_n^b \mathbf{v}_n$, uniformly in the derivative orders.

Question 19.48 (Effective relation reconstruction; source 17). Given a computable finite-alphabet stream and an oracle for its eventual patterns, can the generators of Theorem 34.13 and small coefficient witnesses be produced with practical complexity bounds?

The factorial threshold is explicit but far from economical; finite-index certificates may be much smaller.

Question 19.49 (Larger differential base fields; source 17). Which logarithmic, exponential or differentially algebraic extensions of the bounded-support base of Corollary 35.8 preserve the independence of all intrinsic jets?

The endpoint is a field containing the witnesses; a sharp intermediate boundary would connect support geometry with differential closure.

Question 19.50 (Intrinsic differentiation of the codes; source 17). For a given coding scale M_Ω , is there a natural differential base containing its intrinsic derivatives over which the $\mathfrak{z}_A$ remain differentially independent?

Multiplication by a differential unit preserves independence over a suitable base, but enlarging the base may destroy the original independence.

Question 19.51 (Infinite interpolation; source 17). Which infinite discrete collections of prescribed jets admit a continuum of independent strongly entire solutions in one workspace?

The supports forced by interpolation, not only the linear equations, must be controlled.

Question 19.52 (Specialization and chain rules; source 17). For which substitutions $z = z(\omega)$ can mixed formal independence be transferred to intrinsic numerical independence over a specified smaller field?

Both derivations must be tracked, and specializations that collapse the external variable, such as ordinary constants, excluded.

Question 19.53 (Formalization of the finite core; source 17). Can the band lemma, multicolour polarization, the finite-pattern kernel theorem and the constant-extension minor lemma be formalized in Lean independently of surreal numbers, with the Hahn and Conway interpretation as a separate layer?

Question 19.54 (Arithmetic of the codes; source 17). What can be said about divisibility, irreducibility or ideal membership of the $\mathfrak{z}_A$ in $\mathbf{Oz}$?

Algebraic independence over $\mathcal{F}^{\text{bd}}(M_\Omega)$ is not a factorization theorem; positive support and zero constant term interact with omnific divisibility in ways the field-theoretic result does not determine.

The next eight questions come from source 15 (its Questions 12.1–12.8), in the notation of Section 36; they are research questions formulated there, not claims of literature-wide open status.

Question 19.55 (Critical composition weight; source 15). Classify the equations (36.5) in which some competing monomials have weight exactly $\bar{s}d$. Can the leading coefficient, the indicial factors and the next face of the coefficient Newton polygon replace the strict inequality in significant subclasses?

The proof fails at one identified place: the coefficient of $\gamma_\sigma(f)$ in (36.7) becomes zero, and the remaining expression must be controlled rather than the strict argument reused. The elementary equation $f(z^d) = f(z)^d$ is classified in Example 37.7; its polynomial perturbations and multi-term critical identities are the natural next tests.

Question 19.56 (Several top products; source 15). If the largest-degree pullback occurs in a differential polynomial rather than in a single product of nonzero linear differential transforms, which explicit nonvanishing conditions on its indicial or initial polynomial still force an exact top Gauss profile?

The corner-cancellation issues for ordinary nonlinear equations are those of Sections 15–31; the new ingredient is their interaction with distinct polynomial arguments. The question is not the unrestricted higher-order problem.

Question 19.57 (Matrix differential pullbacks; source 15). Does Theorem 36.15 extend to matrix differential operators whose maximal-shift indicial matrix has nonzero determinant? What is the criterion when different rows have different shifts?

A promising route introduces row and column shifts, normalizes a matrix initial form and tracks the integer roots of its determinant; lower-shift couplings must be accounted for, and entrywise use of the scalar result does not suffice.

Question 19.58 (Several variables; source 15). For a polynomial map $P : \mathbb{K}^r \rightarrow \mathbb{K}^r$, can a positive weight vector and a strictly expanding Newton-polytope face replace the scalar degree d ? Which finite-map or noncancellation assumptions make the pullback of the active initial polynomial injective?

The scalar proof uses the injective substitution $X \mapsto \bar{c}X^d$; multivariable leading forms can satisfy algebraic relations, a genuine obstruction.

Question 19.59 (Meromorphic solutions; source 15). For a quotient of strongly entire series satisfying a differential–Mahler equation, when does the conclusion improve from a finite coefficient reduction to rationality? Can pole divisors and pullback degrees replace the single Gauss profile?

Classical Mahler theory gives a rationality precedent [42]; arbitrary rank and strong summation need a separate divisor and support argument.

Question 19.60 (Minimal universal encoders; source 15). What are the least differential order, polynomial degree bound and number of pullback arguments of an undecidable omnific-coefficient subclass? Can the Euler projectors be replaced by a smaller sparse operator without losing an explicit bound?

The bound ten of Corollary 37.6 is inherited from a known integer encoding and not claimed optimal; a lower bound would have to separate analytic rigidity from arithmetic universality.

Question 19.61 (Arithmetic subclasses beyond linear equations; source 15). Which nonlinear subclasses have decidable omnific solvability despite Corollary 37.5? Natural targets are quadratic coefficient loci of fixed dimension, equations with a prescribed rational parametrization, and cases in which positivity forces ordinary coefficients.

A general algorithm must avoid the universal encoder explicitly; a finite degree bound alone is not such a condition.

Question 19.62 (Formal verification and certified solvers; source 15). Can the scalar weighted theorem and the universal encoder be formalized with the repository’s Hahn summability and polynomial infrastructure? Can a solver emit a compact certificate of operator shifts, indicial polynomials, a degree bound and a verified finite coefficient system?

The dependency order proposed by source 15 is in Section 37.4. Its suggested research order: first the matrix-indicial criterion, then critical-weight equations with one lower argument; the several-variable and meromorphic problems need further geometric infrastructure. The universal encoder supplies adversarial tests for symbolic solvers: a program claiming to decide all nonlinear omnific equations of this class must be incorrect, however well it computes degree bounds on examples.

The next eight questions come from source 20 (its Questions 12.3, 12.4, 12.6–12.10 and 12.12, the fourth re-scoped by the merge), in the notation of Section 38; its Questions 12.1, 12.2, 12.5 and 12.11 are recorded in the statuses of Questions 19.24, 19.21, 19.25 and 19.29.

Question 19.63 (Nonunit coefficient tests without arbitrary coding; source 20). Which restrictions on the initial support or on the observed coefficient cut restore eventual periodicity of coefficient tests for recurrences whose roots are not units?

Finite support, supports in a fixed Archimedean subgroup and one-sided valuation hypotheses are natural starting points. Theorem 39.3 shows that arbitrary reverse well-ordered omnific support cannot be allowed without further restrictions.

Question 19.64 (Full omnific-membership sets; source 20). For real-surreal y_n whose characteristic roots are units, Section 38.4 controls the absence of negative Conway exponents. Which further hypotheses make the integer constant-term condition, and so the set $\{n : y_n \in \mathbf{Oz}\}$, effectively classifiable?

This is the intersection of a support-theoretic problem with an arithmetic integrality problem for an exponential polynomial, not another application of coefficientwise algebraicity.

Question 19.65 (Multivariate time; source 20). For $\mathbf{n} \in \mathbb{N}^{\bar{r}}$, what are the coefficients and valuations of $\sum_j R_j(\mathbf{n}) \prod_i \lambda_{ij}^{n_i}$?

The coefficient-algebra reduction still suggests finitely generated observable algebras, but one-variable arithmetic progressions must be replaced by a finer description of lattice subsets; polynomial conditions in several integer variables warn against a naive semilinearity assertion.

Question 19.66 (Nonlinear mixed equations at unit dilation; source 20, re-scoped). Suppose Γ has an order unit and $q \in \mathbb{K}^\times$ is a nontorsion unit. What are the least differential order and the least degree of a nonzero algebraic relation over $\mathbb{K}(z)$ among finitely many $(D_z^r f)(q^\ell z)$, involving at least two powers of q , that has a nonpolynomial strongly entire solution?

As posed by source 20 the question asked for the smallest nonlinear differential–dilation classes admitting such solutions with an order unit, measured by differential order and nonlinear degree, noting that its linear theorem does not extrapolate to nonlinear equations. In the current text unrestricted nonlinear rigidity is already refuted (Remark 29.19): $\mathcal{T}_p$ satisfies a relation of order three, of degree six, with no dilation, and for relations without dilations the least order is three (Theorem I) and the least degree at order three is at least four (Theorem K); which series attain order three is Question 19.1. The merge re-scopes source 20’s question to what these do not decide: whether a unit dilation lowers the order or the degree needed.

Question 19.67 (Derivations of the coefficient field; source 20). Replace D_z , which fixes $\mathbb{K}$, by an operator that also uses a specified series derivation of $\mathbb{K}$, such as a compatible restriction of the Berarducci–Mantova derivation. Which valuation-cost estimates survive?

Derivatives of coefficients introduce new support dynamics that the present normal-form computation does not cover.

Question 19.68 (Definability of the coding observation; source 20). In which expansions of $(\mathbf{No}, \mathbf{Oz})$ can the coefficient map $[\omega^\omega]$ and the orbit $n \mapsto \omega^n \mathbf{v}$ be defined uniformly?

The coding theorem becomes a model-theoretic interpretation only after these definability questions are answered; the normal-form identity alone interprets no second-order arithmetic in the pure ring.

Question 19.69 (Uniform complexity bounds for finite observations; source 20). For a prescribed finite set of coefficient coordinates and bounded polynomial degrees, can one bound the dimension, the recurrence order, or the number of possible exceptional zeros of the resulting observation sequences?

Such bounds can be useful even when the last exceptional index is not effectively known; by Example 38.10 any bound must depend on the complexity of the observation.

Question 19.70 (Beyond scalar commuting dilations; source 20). What replaces the valuation theory for finite linear substitutions in several variables and for finite-dimensional matrix dilations?

The one-variable proof relies on scalar exponential-polynomial coefficients; noncommuting substitutions may need an orbit theorem for matrix products rather than a direct reuse of the Skolem–Mahler–Lech theorem, and simultaneously triangularizable cases are a tractable first step. Compare Questions 19.22 and 19.58.

These questions are directions suggested by the proved statements, not claims that they were already posed in the published literature, and their formulations are deliberately narrower than a general theory of nonlinear surcomplex differential equations.

20 All-order rigidity without an order unit: setting and valuations

This section and the next two contain the material of sources 11 and 12, two manuscripts delivered together after the rest of this report was written. They answer Question 19.1 in every differential order when Γ has no order unit, including the equations with vanishing residue corner that Sections 14–17 leave open, and source 12 answers the same case of Question 19.4. The method is neither the escape chain of Section 3 nor the finite initial polynomial of Section 14. A finite relation forces all

Taylor coefficients of a solution into one finitely generated field (Theorem 21.2); such a field has a value group of finite rational rank (Lemma 20.6), which is trapped in a proper convex subgroup when Γ has no order unit (Lemma 20.7); and strong entireness needs coefficient values at cofinally many scales (Proposition 20.2). The warning of Remark 5.8 stands: nothing here turns a nonlinear equation into a bounded-step linear recurrence with controlled valuations. The argument uses only that one scalar multiplier is eventually nonzero, and no valuation of it.

20.1 Sources, conventions and the merge's choices

Convention 20.1 (Coefficient field of the coefficient-field part). In Sections 20–22, k is an arbitrary field of characteristic zero with the trivial valuation, Γ is as in Convention 1.1, $\mathbb{K} = k((t^\Gamma))$, and v , strong summability, $\text{Dom}_{\text{str}}(f)$ and $\mathcal{E}_\Gamma(k)$ are as in Convention 14.1. Write $\text{Frac } \mathcal{E}_\Gamma(k)$ for the fraction field of $\mathcal{E}_\Gamma(k)$ inside the formal Laurent field $\mathbb{K}((z))$; the word *meromorphic* in this part means membership of $\text{Frac } \mathcal{E}_\Gamma(k)$, and nothing else. The derivative D_z kills $\mathbb{K}$ and extends to $\mathbb{K}((z))$ by the quotient rule. A series u is *differentially algebraic* over a differential subfield $B \subseteq \mathbb{K}((z))$ if $P(u, D_z u, \dots, D_z^r u) = 0$ for some finite r and some nonzero $P \in B[Y_0, \dots, Y_r]$, and *differentially transcendental* otherwise. For $\lambda \in \mathbb{K}^\times$ and $j \geq 0$ put

$$\mathcal{J}_{\lambda,j} f = (D_z^j f)(\lambda z) = \sum_{n \geq j} a_n(n)_j \lambda^{n-j} z^{n-j} \quad \left(f = \sum_n a_n z^n \right), \quad (20.1)$$

differentiation *before* dilation. The same notation is used for $u \in \mathbb{K}((z))$, using formal Laurent substitution. Since $D_z^j \sigma_\lambda f = \lambda^j \sigma_\lambda D_z^j f$, the operator $D_z^r \sigma_q^\ell$ of (10.1), which dilates first, differs from $\mathcal{J}_{q^\ell,r}$ by the nonzero scalar $q^{\ell r}$. This changes no algebraic independence statement, but recurrence coefficients must use one convention throughout. An infinite family is algebraically independent when every finite subfamily is. For a subgroup $\Delta' \subseteq \Gamma$, $\text{rrk } \Delta' = \dim_{\mathbb{Q}}(\Delta' \otimes_{\mathbb{Z}} \mathbb{Q})$ is its rational rank; ordered groups are torsion free, so they embed in their divisible hulls, and no divisibility of Γ is assumed.

For $k = \mathbb{C}$ one has $\mathbb{K} = K_\Gamma$ and $\mathcal{E}_\Gamma(k) = \mathcal{E}_\Gamma$. As for the nonlinear part, the generality in k is recorded for this part only and is *not* transferred to Theorems A–C.

Two sources, one spine. Sources 11 and 12 were written independently and prove the same five steps.

- (1) Along an actual solution, a minimal relation is linearized. The evaluated partial derivatives have an offset θ and a nonzero *linearization multiplier* Ξ , the quadratic remainder has z -order at least $2(n-r) > n + \theta$, and so all Taylor coefficients lie in one field generated by finitely many of them and finitely many scalars (source 11's Theorem 4.1 and Corollary 4.3; source 12's Theorems 4.1 and 5.2).
- (2) The valuation-rank inequality (source 11's Lemma 5.1, source 12's Lemma 3.1).
- (3) A subgroup of finite rational rank lies in a principal convex subgroup Δ_β , which is proper when there is no order unit (source 11's Lemma 5.2, source 12's Lemma 3.2).
- (4) An exterior obstruction at $t^{-\eta}$ with $\eta > \Delta_\beta$ (source 11's Proposition 5.4, source 12's Proposition 3.4).
- (5) The same worked example: along the solutions az^m of the vanishing-corner equation (16.1) the linearization multiplier is a nonzero multiple of $(T-m)^2$.

Both assume only characteristic zero, $\Gamma \neq 0$ set-sized, no divisibility, and no order unit where that is used; neither has weaker hypotheses than the other. The spine is printed once, with both sources credited at each statement. **Source 12 is the base** of this part: its all-order theorem with dilations (Theorem E (b) here) contains, for the single dilation 1, source 11's statement about entire functions; it answers the no-order-unit case of Question 19.4; and its order-unit dichotomy (Theorem G) gives the exact boundary of the mixed statement. Source 11 contributes what source 12 lacks: coarsening descent for arbitrary, not necessarily finitely generated, coefficient fields (Theorem 21.11); rationality of every differentially algebraic element of $\text{Frac } \mathcal{E}_\Gamma(k)$ (Theorem E (c)); the rational-system and Painlevé corollaries; the joint-independence criterion (Theorem 22.2); and a continuum-sized independent family (Theorem 22.6). Two of source 11's proofs differ genuinely from source 12's and are kept as marked second routes: its choice of a minimal relation (Remark 21.3) and its proof of polynomiality by coarsened reduction (proof of Theorem E). Every proof taken from source 11 was re-read against the statement it is attached to: source 11 has no stronger standing hypothesis than source 12; its cardinal equality uses $k = \mathbb{C}$, while its lower bound holds for every k ; and its coefficient principle is relative to a Laurent field $k_0((z))$, a form kept because the joint-independence criterion needs it.

Renamed symbols. The following symbols of the two sources would collide with symbols of Sections 1–19 or of docs/NOTATION.md. They are renamed here, in the new material only.

Source 11	Source 12	Here	Reason
D	D	D_z	D is a scalar derivation in the notation guide; Definition 1.4.
$K, \mathcal{E}_\Gamma$	$K, \mathcal{E}$	$\mathbb{K}, \mathcal{E}_\Gamma(k)$	Convention 14.1; $\mathcal{E}_\Gamma$ is over $\mathbb{C}$.
$\mathcal{M}_\Gamma = \text{Frac } \mathcal{E}_\Gamma$	—	$\text{Frac } \mathcal{E}_\Gamma(k)$	$M(S)$ is a support monoid.
H, H_δ	H, H_h	Δ, Δ_β	(2.1); $H \in \Gamma$ in Section 11.3.
$w, \mathcal{O}_w, \mathbf{m}_w, L, \text{res}$	w_H	$v_\Delta, \mathcal{O}_\Delta, \mathbf{m}_\Delta, k_\Delta, \text{red}_\Delta$	w_P is a corner shift; res is the residue at v .
$s, A_j, C_j, Q(T)$	$\ell, S_{ij}, c_{ij}, B(T)$	$\theta, \Pi_{ij}, \xi_{ij}, \Xi(T)$	s is an order in Question 19.1, ℓ a dilation power in (10.1), B a threshold, Q_s a shift polynomial.
F (a subfield)	L, F	L, K_0, k_0	F is the series (11.2).
$F_\alpha, \sigma, c(\sigma)$	—	$\mathcal{F}_\alpha, b, \text{code}(b)$	F as above; σ_q is a dilation.
B_i	—	$k_{J,i}$	B is a threshold.
$\Gamma_\infty = \bigoplus_{m \geq 1} \mathbb{Q}e_m$	$\bigoplus_{n \geq 0} \mathbb{Q}e_n$	(11.1)	One group, indexed from 0.
f_*	F_ω	F	See below.
Θ over $\mathbb{Q}$	$\Theta_u, u = t^h$	G, Θ_p	Corollary 17.1; (8.1).
—	$\rho(f)$	$\text{cvr}(f)$	ρ is the residue of Lemma 14.2.
—	E_h	E_η	Proposition 5.3.
—	F_τ	R_τ	F as above.
—	s, λ, μ	m, λ, λ'	s is an order; μ a dilation valuation.
Theorems 1.1–1.3	Theorems A, B, C	Theorems 21.11, E, 22.6; E (b), F, G	Theorems A–D of this report are different theorems.

The shipped audit files 11-coarsening-differential-rigidity-* and 12-coefficient-field-rigidity-SOURCE_AUDIT.md keep the sources' notation and numbering.

Words with two meanings. *No order unit* is not a synonym of *non-Archimedean*, *higher rank* or *infinite rational rank* (Section 21.4). *Rank one* means Archimedean; the *rational rank* rrk is

a $\mathbb{Q}$ -dimension. A *dilation* is always the argument dilation $f(z) \mapsto f(\lambda z)$ with $\lambda \in \mathbb{K}^\times$, never an automorphism of the scalar field. *Residue* keeps its meaning at v (res, ρ); the residue at a coarsened valuation is written red_Δ . The *linearization multiplier* Ξ depends on a solution; it is not the residue corner polynomial χ_P or the full corner polynomial I_P of (14.6), which depend on the equation alone, and not a shift polynomial Q_s or P_s of Sections 5–10. Source 12’s lemma “escape of leading values is sufficient” is Lemma 2.3 (a), not the escape chain.

Results printed once. Sources 11 and 12 reprove several facts that the report already contains. They are printed once and cited: the definition of strong summability (Definition 1.2); the sufficiency half of the all-scale criterion (Lemma 2.3 (a)); closure of the entire class under derivatives and dilations (Proposition 2.7, whose proof uses no property of $\mathbb{C}$); the surreal normal-form embedding (Section 2.4, with the same convention $t^\gamma \mapsto \omega^{-\gamma}$); the group Γ_∞ and strong entireness of F (Theorem 11.2); the failure of that entireness after a dominating scale is adjoined (Section 11.2); the domain of the formal exponential (Proposition 5.3); and the failure in positive characteristic (Example 17.3). Everything else in the two sources is printed in Sections 20–22, with proofs.

Where each numbered result went. Source 11: Theorems 1.1–1.3 are Theorem 21.11, Theorem E (a),(c) and Theorem 22.6; Lemma 2.2 is Proposition 20.2, Remark 2.3 the paragraph after it, Proposition 2.4 and Lemma 2.5 are Proposition 20.4 and Lemma 20.5; Section 3 is Section 21.2; Theorem 4.1 is Theorem 21.2 (a) with Remark 21.3, Remark 4.2 is Remark 21.4, Corollary 4.3 is Corollary 21.8, Example 4.4 is Example 21.6, and Example 4.5 follows it; Lemmas 5.1–5.2, Remark 5.3 and Proposition 5.4 are Lemmas 20.6–20.7, Remark 20.8 and Proposition 20.9; Corollaries 6.1–6.3 and Example 6.4 are Corollaries 21.15–21.17 and the paragraph after the last; Section 7 is Section 22.2; Section 8 is Section 22.3; Questions 9.1 and 9.3 are Questions 19.1 (re-scoped) and 19.7, and Example 9.2 is discussed in Section 21.4. Source 12: Theorems A–C are Theorems E (a),(b), F and G; Proposition 2.3, Corollary 2.4 and Proposition 2.5 are Proposition 20.2, Corollary 20.3 and Proposition 20.4; Lemmas 3.1–3.2, Remark 3.3, Proposition 3.4 and Corollaries 3.5–3.6 are Lemmas 20.6–20.7, Remark 20.8, Proposition 20.9 and Corollaries 20.10–20.11; Theorems 4.1 and 5.2 are Theorem 21.2 (a) and (b), Remarks 4.2 and 5.3 are Remarks 21.4 and 21.5, Corollary 4.3 is Theorem E (a), Corollary 4.4 is Corollary 21.18, Lemma 5.1 is Lemma 21.1 and Corollary 5.4 is Corollary 21.14; Proposition 6.1 and Theorem 6.2 are Proposition 21.19 and Theorem 21.20; Theorem 7.1 and Example 7.2 are Theorem 22.1 and the paragraph after it; Propositions 8.1 and 8.2 are Proposition 22.7 and the whole-class failure of Section 22.3; Section 9 is Section 22.4; Questions 10.1–10.3 are Questions 19.1 (re-scoped), 19.8 and 19.9.

The explicit series, and an indexing discrepancy. Source 12’s F_ω is exactly F of (11.2), over the same group indexed from 0. Source 11 indexes the group from 1, maps e_m to ω^m , and writes $f_* = \sum_{n \geq 1} \omega^{-\omega^n} z^n$, calling it “this exact entire example” of the report. That is off by a constant: the term $n = 0$ of F is $t^{e_0} = \omega^{-1}$, so $F = f_* + \omega^{-1}$. The discrepancy is harmless, because F and f_* differ by an element of the field, their derivatives agree, and a polynomial $P(Y_0, Y_1, \dots)$ vanishes on the jets of F exactly when $P(Y_0 + \omega^{-1}, Y_1, \dots)$ vanishes on those of f_* . It is disclosed here and not corrected in the shipped audit files. Source 11’s branch family uses only codes at least 2, so it is printed over the group (11.1) unchanged.

20.2 Strong entireness at every scale

Proposition 20.2 (All-scale coefficient criterion; sources 11 and 12). *For $f = \sum_n a_n z^n \in \mathbb{K}[[z]]$ the following are equivalent:*

- (i) $f \in \mathcal{E}_\Gamma(k)$;
- (ii) the family $(a_n t^{n\gamma})_{n \geq 0}$ is strongly summable for every $\gamma \in \Gamma$;
- (iii) for all $\gamma, \beta \in \Gamma$, only finitely many n with $a_n \neq 0$ satisfy $v(a_n) + n\gamma \leq \beta$.

Proof. (i) $\Rightarrow$ (ii) is immediate. For (ii) $\Rightarrow$ (iii) fix γ and suppose that infinitely many nonzero coefficients have $A_n := v(a_n) + n\gamma \leq \beta$. Strong summability at t^γ gives a common lower bound α for all A_n , the least element of the union of the supports. Choose any $\eta > 0$ in Γ and set $\epsilon = \max(\beta - \alpha, 0) + \eta$. Then $\epsilon > 0$ and $\epsilon > \beta - \alpha$, without division or an order-unit assumption. For two selected indices $n' < n$,

$$(A_n - n\epsilon) - (A_{n'} - n'\epsilon) \leq \beta - \alpha - (n - n')\epsilon < 0,$$

so at the more exterior argument $t^{\gamma-\epsilon}$ the leading exponents of the selected summands form a strictly decreasing sequence, contradicting well-ordered support. For (iii) $\Rightarrow$ (i), a nonzero x with $v(x) = \gamma$ gives summands $a_n x^n$ of valuation $v(a_n) + n\gamma$, and Lemma 2.3 (a), whose proof uses no property of $\mathbb{C}$, applies; at $x = 0$ only one summand is nonzero. $\square$

The sufficiency half is Lemma 2.3 (a). Source 11 describes the criterion as part of the collection's support framework and source 12 as part of its entire-function programme [12]; both prove it, and it is printed here once, with the necessity half, which Sections 1–13 did not need. The necessity proof uses a second, more exterior argument: strong summability at one argument does not by itself give cofinal growth. In $k((t^\mathbb{Q}))$ the family $(t^{n/(n+1)})_{n \geq 1}$ is strongly summable although its valuations are bounded (source 12), and Remark 2.5 gives a higher-rank version. Replacing the all-scale hypothesis by evaluation on one ordinary complex disk would therefore be an error (source 11).

Corollary 20.3 (When the statements are nonvacuous; source 12). *A nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ exists if and only if Γ has countable cofinality.*

Proof. If f is nonpolynomial, (iii) with $\gamma = 0$ makes the values of its infinitely many nonzero coefficients a countable cofinal subset; a nonzero ordered abelian group has no largest element, so its cofinality is not finite. Conversely choose a nondecreasing cofinal sequence of positive δ_n , $n \geq 1$, and put $a_0 = 1$, $a_n = t^{n\delta_n}$. Given γ, β , eventually $\delta_n + \gamma > \max(\beta, 0)$, so $n(\delta_n + \gamma) > \beta$, and Proposition 20.2 applies. $\square$

The equivalence is background from the entire-function report [12] (Section 1.5); source 12's proof is printed because the next sections use it. Theorems E and G therefore have nonpolynomial instances exactly for groups of countable cofinality; in uncountable cofinality they remain true but are vacuous.

Proposition 20.4 (Differential algebra structure; source 11). *$\mathcal{E}_\Gamma(k)$ is a differential $\mathbb{K}$ -subalgebra of $\mathbb{K}[[z]]$, stable under every dilation σ_λ , $\lambda \in \mathbb{K}^\times$. Its fraction field $\text{Frac } \mathcal{E}_\Gamma(k)$ is a differential subfield of $\mathbb{K}((z))$.*

Proof. Closure under sums, products, D_z and σ_λ is Proposition 2.7, whose proof uses no property of $\mathbb{C}$. Through Proposition 20.2 it can also be read off the weighted values: for a product, if

$A_i = v(a_i) + i\gamma$ and $B_j = v(b_j) + j\gamma$ have lower bounds m_A, m_B , then $A_i + B_j \leq \beta$ forces $A_i \leq \beta - m_B$ and $B_j \leq \beta - m_A$, which only finitely many pairs satisfy. If a nonzero product coefficient at degree n has weighted value at most β , one of its finitely many nonzero terms $a_i b_j$, $i + j = n$, must have weighted value at most β . Thus only finitely many degrees can occur, even when some terms cancel. A zero factor is immediate. For $\mathcal{J}_{\lambda,j} f$ the coefficient of z^n is $a_{n+j}(n+j)_j \lambda^n$, whose weighted value at γ is that of a_{n+j} at $v(\lambda) + \gamma$ minus the constant $j(v(\lambda) + \gamma)$. Here the falling factorial is a nonzero element of the trivially valued prime field, so it contributes no valuation (source 12). The quotient rule extends D_z to $\text{Frac } \mathcal{E}_\Gamma(k)$, and every nonzero power series is invertible in $\mathbb{K}((z))$, so $\text{Frac } \mathcal{E}_\Gamma(k)$ embeds there. $\square$

Lemma 20.5 (A minimum coefficient value; source 11). *If $0 \neq h = \sum_n h_n z^n \in \mathcal{E}_\Gamma(k)$, some h_j satisfies $v(h_j) \leq v(h_n)$ for every nonzero h_n .*

Proof. Take any nonzero coefficient value β . By Proposition 20.2 with $\gamma = 0$ only finitely many coefficients have value at most β ; the least of them is the minimum. $\square$

20.3 Value groups of finitely generated fields

Lemma 20.6 (Valuation-rank inequality; sources 11 and 12). *Let $k_0 \subseteq L$ be fields and v a valuation on L , with values in an ordered abelian group, trivial on $k_0^\times$. If $x_1, \dots, x_d \in L^\times$ have values linearly independent over $\mathbb{Q}$, they are algebraically independent over k_0 . Consequently*

$$\text{rrk } v(L^\times) \leq \text{trdeg}_{k_0} L. \quad (20.2)$$

Proof. For distinct $\alpha, \alpha' \in \mathbb{N}^d$ the monomials x^α and $x^{\alpha'}$ have different values, since equality would be an integer relation among the $v(x_i)$. A nonzero coefficient from k_0 has value zero. So the finitely many nonzero terms of a nonzero polynomial evaluated at the x_i have pairwise distinct values, and the unique term of least value cannot be cancelled. For the rank inequality, choose a rational basis of $v(L^\times) \otimes_{\mathbb{Z}} \mathbb{Q}$ from the images of elements of $v(L^\times)$, and lift those values to elements of $L^\times$. Every polynomial uses only finitely many of the lifts, so the preceding argument proves that the whole lifted family is algebraically independent. Its cardinality is the rational rank, giving the inequality also when either rank is infinite. $\square$

Source 12 states the lemma for $k \subseteq L \subseteq \mathbb{K}$, source 11 for $k_0 = \mathbb{Q}$ and any valuation trivial on $\mathbb{Q}$; both cases are used below.

Lemma 20.7 (A principal convex bound; sources 11 and 12). *Let $\Delta' \subseteq \Gamma$ be a subgroup of finite rational rank. Either $\Delta' = 0$, or $\Delta' \subseteq \Delta_\beta$ for some $\beta > 0$, where Δ_β is the convex subgroup (2.1); in the second case β can be chosen in Δ' , and Δ_β is then the smallest convex subgroup containing Δ' . The subgroup Δ_β is proper exactly when β is not an order unit, so it is proper for every β when Γ has no order unit.*

Proof. Choose $\delta_1, \dots, \delta_d \in \Delta'$ whose images form a basis of $\Delta' \otimes \mathbb{Q}$ and put $\beta = \sum_i |\delta_i| > 0$, an element of Δ' . For $\delta \in \Delta'$, clearing denominators gives $m\delta = \sum_i n_i \delta_i$ with a positive integer m and integers n_i , so $|\delta| \leq m|\delta| \leq (\sum_i |n_i|)\beta$. Hence $\Delta' \subseteq \Delta_\beta$, and Δ_β , the smallest convex subgroup containing $\beta \in \Delta'$, lies in every convex subgroup containing Δ' . The last assertion is the definition of an order unit. $\square$

No division in Γ occurs. Source 11 takes $\delta = \eta + \sum_i |\delta_i|$ with an arbitrary $\eta > 0$, which covers $\Delta' = 0$ by the same formula.

Remark 20.8 (Field values, not generator values; sources 11 and 12). The bound (20.2) concerns the value group of the whole field. It would be false to use the values of a chosen list of generators: for $0 < \alpha < \beta$ with α, β rationally independent, both generators of $\mathbb{Q}(t^\alpha, t^\alpha + t^\beta)$ have value α , but their difference has value β (source 11). Conversely, one generator $b \in \mathbb{K}$ may have infinitely many rationally independent exponents in its support without contradicting $\text{rrk } v(k(b)^\times) \leq 1$: the field $k(b)$ contains rational functions of b , not the truncations of its Hahn expansion (source 12). This is why the coefficients of the equations below may be arbitrary Hahn elements.

For a convex subgroup $\Delta \subseteq \Gamma$ the quotient Γ/Δ is an ordered abelian group, and

$$v_\Delta(a) = v(a) + \Delta \in \Gamma/\Delta \quad (a \in \mathbb{K}^\times), \quad v_\Delta(0) = +\infty,$$

is a valuation on the same field $\mathbb{K}$, a coarsening of v , as in Theorem 7.4; it is not a coefficientwise map to a smaller Hahn field. Write $\mathcal{O}_\Delta = \{a : v_\Delta(a) \geq 0\}$, $\mathfrak{m}_\Delta = \{a : v_\Delta(a) > 0\}$, $k_\Delta = \mathcal{O}_\Delta/\mathfrak{m}_\Delta$ and $\text{red}_\Delta : \mathcal{O}_\Delta \rightarrow k_\Delta$. No splitting of $\Gamma \rightarrow \Gamma/\Delta$ and no identification of k_Δ with a Hahn field is used. Write $\eta > \Delta$ when $\eta > \delta$ for every $\delta \in \Delta$; by convexity every positive element outside Δ satisfies this, and $v_\Delta(x) < 0$ means exactly $v(x) = -\eta$ with $\eta > \Delta$.

Proposition 20.9 (Coarsened exterior obstruction; sources 11 and 12). *Suppose all nonzero coefficients of a nonpolynomial $f = \sum_n a_n z^n \in \mathbb{K}[[z]]$ have values in a proper convex subgroup Δ . Then*

$$\text{Dom}_{\text{str}}(f) \subseteq \{0\} \cup \{x \in \mathbb{K}^\times : v_\Delta(x) \geq 0\}; \quad (20.3)$$

in particular $t^{-\eta} \notin \text{Dom}_{\text{str}}(f)$ for every $\eta > \Delta$.

Proof. Let $v(x) = -\eta$ with $\eta > \Delta$. For nonzero coefficients at indices $m > n$, $v(a_m x^m) - v(a_n x^n) = (v(a_m) - v(a_n)) - (m - n)\eta < 0$, because the first difference lies in Δ . The leading exponents of the infinitely many nonzero summands therefore strictly decrease, and Lemma 2.2 applies. $\square$

Source 11 states this for $x = t^{-\eta}$ and coefficients in a field whose values lie in Δ ; source 12 states (20.3). The inclusion is necessary, not sufficient: points with $v_\Delta(x) \geq 0$ need not lie in $\text{Dom}_{\text{str}}(f)$. For $k = \mathbb{C}$ it is consistent with inward stability (Lemma 2.4).

Corollary 20.10 (Finite transcendence degree; source 12). *Assume Γ has no order unit. If all coefficients of $f \in \mathcal{E}_\Gamma(k)$ lie in a subfield $L \supseteq k$ of $\mathbb{K}$ with $\text{trdeg}_k L < \infty$, then $f \in \mathbb{K}[z]$.*

Proof. By Lemmas 20.6 and 20.7, $v(L^\times)$ lies in $\{0\}$ or in some Δ_β , and both are proper. Proposition 20.9 excludes a nonpolynomial f . $\square$

For $f = \sum_n a_n z^n$ define the *coefficient-value rank*

$$\text{cvr}(f) = \dim_{\mathbb{Q}} \left(\text{span}_{\mathbb{Q}} \{v(a_n) \otimes 1 : a_n \neq 0\} \right). \quad (20.4)$$

The span is taken in $\Gamma \otimes_{\mathbb{Z}} \mathbb{Q}$, and $\text{cvr}(f)$ is its dimension. It is the rational rank of the values of the Taylor coefficients, not the number of support exponents of any single coefficient.

Corollary 20.11 (Necessary infinite coefficient-value rank; source 12). *If Γ has no order unit and $f \in \mathcal{E}_\Gamma(k)$ is not a polynomial, then $\text{cvr}(f) = \aleph_0$ and $\text{trdeg}_k k(a_0, a_1, \dots) = \aleph_0$.*

Proof. If $\text{cvr}(f)$ were finite, the group generated by the coefficient values would lie in a proper Δ_β by Lemma 20.7, contradicting Proposition 20.9. There are countably many coefficients, which bounds the rank; Lemma 20.6 gives the lower bound for the transcendence degree and countable generation the upper bound. $\square$

21 The coefficient-field theorem and its consequences

21.1 One finitely generated field for every finite relation

Coefficient recurrences for algebraic differential equations go back to Hurwitz [23], and Lemma 3 of Laohakosol, Kongsakorn and Ubolsri [24] isolates a high coefficient through a polynomial in its index; Krattenthaler and Rivoal treat the arithmetic of such coefficients [25], and Ait El Manssour, Sattelberger and Teguia Tabugua the general framework of D-algebraic functions [26]. Only the field-theoretic consequence is needed here. Both sources supply a self-contained formal linearization proof over arbitrary characteristic-zero fields that allows singular solutions, and neither advertises the finite-generation principle or the valuation-rank inequality as a discovery.

For several dilations one classical theorem is imported and not proved: the *Skolem–Mahler–Lech theorem*, that the zero set of a linear recurrence sequence over a field of characteristic zero is a finite set together with finitely many infinite arithmetic progressions. Source 12 uses it in the general-field form of Bell’s Theorem 1.1 [27], credited there to Lech [28]; no generalized dynamical form of the theorem is used.

Lemma 21.1 (Nondegenerate exponential polynomials; source 12). *Let K_0 be a field of characteristic zero and let $\lambda_1, \dots, \lambda_m \in K_0^\times$ with $\lambda_i/\lambda_{i'}$ not a root of unity for $i \neq i'$. If $p_1, \dots, p_m \in K_0[T]$ are not all zero, then*

$$\Xi(n) = \sum_{i=1}^m \lambda_i^n p_i(n) \quad (21.1)$$

is nonzero for all sufficiently large $n \in \mathbb{N}$.

Proof. Discard the zero p_i , and let E be the shift $(Eu)(n) = u(n+1)$. The sequence $\lambda_i^n p_i(n)$ is annihilated by $(E - \lambda_i)^{\deg p_i + 1}$, so the product of these annihilators kills Ξ : they commute because they are polynomials in the same shift E . Thus Ξ satisfies a linear recurrence with constant coefficients in K_0 , and by the imported theorem its zero set is a finite set together with finitely many infinite arithmetic progressions. No infinite progression can occur. If $\Xi(a + bu) = 0$ for all $u \geq 0$, with $b \geq 1$, then $\sum_i (\lambda_i^b)^u \lambda_i^a p_i(a + bu) = 0$ for all u , and the bases $\alpha_i = \lambda_i^b$ are pairwise distinct by the root-of-unity hypothesis. Polynomial-exponential sequences with pairwise distinct nonzero bases are linearly independent: to isolate the base α_i , apply $\prod_{i' \neq i} (E - \alpha_{i'})^{d_{i'} + 1}$, where $d_{i'}$ is the degree of the polynomial at $\alpha_{i'}$; this kills the other terms, and on $\alpha_i^u p_i(u)$ each factor keeps the degree and multiplies the leading coefficient by $\alpha_i - \alpha_{i'} \neq 0$, leaving a nonzero power times a nonzero polynomial, which cannot vanish at every natural number in characteristic zero. Hence every $p_i(a + bT)$, and so every p_i , is zero, a contradiction. $\square$

For $m = 1$ the lemma is the finiteness of the roots of a nonzero polynomial, since $\lambda_1^n \neq 0$. When only the $j = 0$ jets occur and $\lambda_i = q^{m_i}$ with distinct integers m_i and a nontorsion q , the multiplier is a nonzero Laurent polynomial evaluated at the distinct elements q^n , again without the imported theorem. The general mixed-dilation argument uses the imported theorem.

Theorem 21.2 (Finite coefficient field; sources 11 and 12). *Let $k_0 \subseteq K_0$ be fields of characteristic zero and $u = \sum_{n \geq 0} a_n z^n \in K_0[[z]]$, with D_z killing K_0 .*

- (a) (Sources 11 and 12.) *If u is differentially algebraic over $k_0((z))$, there is $N \geq 0$ with $a_n \in k_0(a_0, \dots, a_{N-1})$ for every n .*
- (b) (Source 12.) *Let $\lambda_1, \dots, \lambda_m \in K_0^\times$ have pairwise quotients that are not roots of unity. If finitely many of the series $\mathcal{J}_{\lambda_i, j} u$ are algebraically dependent over $K_0(z)$, then all a_n lie in one subfield $L \subseteq K_0$ finitely generated over k_0 , generated by the λ_i , the finitely many coefficients of a suitable polynomial relation, and finitely many a_n .*

Proof. Treat both parts at once, first choosing a relation involving derivative orders at most r . In case (a) put $m = 1$, $\lambda_1 = 1$ and $R = k_0[[z]]$, and multiply the relation by a sufficiently large power of z to remove the finitely many Laurent poles of its coefficients. In case (b) put $R = K_0[z]$ and clear a common polynomial denominator of its finitely many rational-function coefficients. In either case there is a nonzero polynomial over R in variables Y_{ij} , $1 \leq i \leq m$, $0 \leq j \leq r$, vanishing at $Y_{ij} = \mathcal{J}_{\lambda_i, j} u$. Among these relations choose P of least total degree in the Y_{ij} . Its degree is positive, since a nonzero element of R does not vanish in $K_0[[z]]$. Put

$$\Pi_{ij} = \frac{\partial P}{\partial Y_{ij}}((\mathcal{J}_{\lambda_i, j} u)_{i, j}) \in K_0[[z]].$$

A nonzero polynomial partial has nonzero evaluation, by minimality of the total degree, and characteristic zero gives at least one. Put

$$\theta = \min_{\Pi_{ij} \neq 0} (\text{ord}_z \Pi_{ij} - j), \quad \xi_{ij} = [z^{\theta+j}] \Pi_{ij}, \quad (21.2)$$

with $\xi_{ij} = 0$ when $\Pi_{ij} = 0$ or $\theta + j < 0$, so $\theta \geq -r$ and some $\xi_{ij} \neq 0$, and put

$$\Xi(n) = \sum_{i, j} \xi_{ij} \lambda_i^{n-j} (n)_{\underline{j}} = \sum_i \lambda_i^n p_i(n), \quad p_i(T) = \sum_j \xi_{ij} \lambda_i^{-j} (T)_{\underline{j}}. \quad (21.3)$$

Falling factorials of distinct degrees are linearly independent, so $p_i \neq 0$ whenever some $\xi_{ij} \neq 0$, and Lemma 21.1 makes $\Xi(n) \neq 0$ for all large n . In case (a), $\Xi(T) = \sum_j \xi_{1j} (T)_{\underline{j}}$ is a nonzero polynomial, and the lemma is not needed.

Let $T_n = \sum_{m' < n} a_{m'} z^{m'}$ and $g_n = u - T_n$. By (20.1), $\text{ord}_z \mathcal{J}_{\lambda_i, j} g_n \geq n - j \geq n - r$ for $n \geq r$. Taylor expansion of P at the jets of u gives

$$P((\mathcal{J}_{\lambda_i, j} T_n)_{i, j}) = - \sum_{i, j} \Pi_{ij} \mathcal{J}_{\lambda_i, j} g_n + R_n, \quad \text{ord}_z R_n \geq 2(n - r), \quad (21.4)$$

because every term of R_n has at least two factors $\mathcal{J}_{\lambda_i, j} g_n$ and its other factors have nonnegative order; this holds even when the evaluated partials vanish to high order. For $n > \theta + 2r$ one has $2(n - r) > n + \theta$, so R_n contributes nothing in degree $n + \theta$. For each nonzero Π_{ij} the coefficient contributed by $a_{m'}$ to degree $n + \theta$ is

$$a_{m'} (m')_{\underline{j}} \lambda_i^{m'-j} [z^{n+\theta-m'+j}] \Pi_{ij}, \quad m' \geq n.$$

If $m' > n$, the displayed exponent is below $\theta + j$ and hence below $\text{ord}_z \Pi_{ij}$, or is negative, so this coefficient is zero. At $m' = n$ it is $a_n (n)_{\underline{j}} \lambda_i^{n-j} \xi_{ij}$. Terms with $\Pi_{ij} = 0$ contribute nothing. Thus only the stated multiplier of a_n remains:

$$a_n \Xi(n) = -[z^{n+\theta}] P((\mathcal{J}_{\lambda_i, j} T_n)_{i, j}). \quad (21.5)$$

Put $N_0 = \theta + 2r \geq r$. Let $L_0 = k_0$ in case (a); in case (b), let L_0 be generated over k_0 by the λ_i and the finitely many scalar coefficients of P . Replacing u by T_{N_0+1} changes every jet of order at most r only in degrees at least $N_0 + 1 - r = \theta + r + 1$. Polynomial substitution with coefficients in R preserves this congruence. Since ξ_{ij} is a coefficient in degree $\theta + j \leq \theta + r$, it is determined by that truncation; thus $\xi_{ij} \in L_0(a_0, \dots, a_{N_0})$.

Choose $N > N_0$ so that $\Xi(n) \neq 0$ for every $n \geq N$, and put $L = L_0(a_0, \dots, a_{N-1})$. Its generators contain the required prefix, so $\Xi(n) \in L^\times$ for $n \geq N$. If all $a_{m'}$ with $m' < n$ already lie in L , then the right side of (21.5) lies in L : in case (a) every coefficient of the finitely many power series occurring as coefficients of P lies in k_0 , and in case (b) all scalar coefficients of P lie in L_0 . Division by $\Xi(n)$ gives $a_n \in L$, completing the induction. In case (a), $L = k_0(a_0, \dots, a_{N-1})$; in case (b), L is finitely generated over k_0 . $\square$

Remark 21.3 (Second route: source 11's minimal relation). For part (a), source 11 chooses a relation of least order r and then least degree in Y_r . Its separant $\partial P / \partial Y_r$ is a nonzero polynomial in characteristic zero, and its evaluation is nonzero, since otherwise it would be a relation of smaller order, or of the same order and smaller degree in Y_r . It then defines the offset and multiplier from the nonzero evaluated partials exactly as in (21.2). Source 12's least total degree makes *every* nonzero partial evaluate to a nonzero series. The proof needs only one nonzero evaluated partial, so either choice works; they are the two sources' routes to the same recurrence.

Remark 21.4 (No nonsingular initial condition; sources 11 and 12). The theorem assumes no evaluated partial nonzero at $z = 0$, no equation solved for the highest derivative, no irreducibility and no nonzero residue corner. A nonzero evaluated partial is used as a formal series, not through its constant term; the offset θ absorbs its vanishing order, and the finitely many natural zeros of Ξ are placed in the initial segment. No implicit-function theorem at an ordinary point is invoked.

Remark 21.5 (Existential, not an algorithm; source 12). The invoked form of Skolem–Mahler–Lech supplies no effective bound on the last exceptional index, and no effective equality, leading-exponent access or computable representation of the coefficients of P is assumed. Part (b) is an existence statement. In part (a) the multiplier is explicitly a polynomial, but computational claims would still have to say how its scalar coefficients are represented.

Example 21.6 (Linearization along the vanishing-corner example; sources 11 and 12). For the equation (16.1), $P = z^2 Y_0 Y_2 + z Y_0 Y_1 - z^2 Y_1^2$, whose residue corner polynomial vanishes identically, take the solution $u = az^m$ with $a \neq 0$ and $m \geq 0$ (Proposition 16.4). Writing Π_j for Π_{1j} , the evaluated partials are

$$\Pi_0 = am^2 z^m, \quad \Pi_1 = a(1 - 2m)z^{m+1}, \quad \Pi_2 = az^{m+2},$$

so $\theta = m$ (for $m = 0$ after discarding $\Pi_0 = 0$) and

$$\Xi(T) = a(m^2 + (1 - 2m)T + T(T - 1)) = a(T - m)^2 \neq 0. \quad (21.6)$$

Source 12 takes $a = 1$; source 11 carries a . The multiplier is nonzero although $\chi_P = 0$: it is computed along a solution, χ_P from the equation alone, and its only exceptional index $n = m$ moves with the solution. This is why the argument needs no degree bound for polynomial solutions, which this equation does not have. The displayed P is not of least degree for az^m , which also satisfies $zD_z u - mu = 0$; the computation shows that the recurrence (21.5) can be run for this P directly, since one of its evaluated partials is nonzero. Neither source infers the general theorem from the example.

Example 21.7 (The initial segment is not bounded by the order; source 11). For an integer $N \geq 0$, the equation $zD_z u - Nu = 0$ has the solution az^N with arbitrary a , and $\Xi(T) = T - N$. Over a field not containing a , the initial segment must reach degree N , so the differential order alone does not bound its length.

Corollary 21.8 (Finitely many scalar parameters; source 11). *Let K_0 be a field of characteristic zero. If $u \in K_0((z))$ is differentially algebraic over $K_0(z)$, all its Laurent coefficients lie in a subfield of K_0 finitely generated over $\mathbb{Q}$.*

Proof. Clear denominators in a relation; its finitely many scalar coefficients $c_1, \dots, c_q$ generate $k_0 = \mathbb{Q}(c_1, \dots, c_q)$. Write $u = z^{-m'}y$ with a power series y , taking $m' \geq 0$. Leibniz's formula expresses the jets of u through order r as a triangular $k_0(z)$ -linear combination of the jets of y through that order, with nonzero diagonal $z^{-m'}$. This invertible substitution preserves nonzero polynomial relations, so y is differentially algebraic over $k_0(z) \subseteq k_0((z))$. Theorem 21.2 (a) puts the coefficients of y , hence of u , in k_0 with finitely many initial coefficients adjoined. $\square$

21.2 Coarsening descent

This subsection is source 11's. Its polynomial half is a coefficient-field form of the entire-function report's obstruction for noncofinal extensions of the workspace [12]; the fraction-field half is the part that needs a new argument.

Lemma 21.9 (Polynomial reduction of an entire series; source 11). *Let $\Delta \subsetneq \Gamma$ be convex. If $g = \sum_n g_n z^n \in \mathcal{E}_\Gamma(k)$ has all coefficients in $\mathcal{O}_\Delta$, its coefficientwise reduction $\bar{g} = \sum_n \text{red}_\Delta(g_n) z^n$ is a polynomial in $k_\Delta[z]$. Every nonzero $h \in \mathcal{E}_\Gamma(k)$ has a scalar multiple h/c with coefficients in $\mathcal{O}_\Delta$ and nonzero reduction.*

Proof. Because Δ is proper, choose a positive element η outside it; convexity gives $\eta > \Delta$. By Proposition 20.2 with $\gamma = 0$, eventually $v(g_n) > \eta$, so $v_\Delta(g_n) > 0$ and $\text{red}_\Delta(g_n) = 0$. For the second assertion take $c = h_j$ as in Lemma 20.5: all coefficients of h/c have nonnegative v , and the j th is 1. $\square$

Lemma 21.10 (Rationality needs no new constants; source 11). *For any extension of fields $L \subseteq L'$, $L'(z) \cap L((z)) = L(z)$ inside $L'((z))$.*

Proof. After multiplying by a power of z , let $u = \sum_{n \geq 0} u_n z^n \in L[[z]]$ lie in $L'(z)$. Being regular at zero, $u = p/q$ with $p, q \in L'[z]$ and $q(0) = 1$; put $d = \deg q$ and take $N \geq \max(d, \deg p)$. Then $u_n + q_1 u_{n-1} + \dots + q_d u_{n-d} = 0$ for $n > N$. This is a linear system with coefficients in L in the finitely many unknowns $q_1, \dots, q_d$. The augmented rows lie in L^{d+1} , so finitely many of them span all the rows. Solving that finite subsystem is therefore equivalent to solving the whole system. Gaussian elimination uses coefficients in L , and cannot produce an inconsistent row because a solution exists in L' . Setting its free variables to zero gives a solution $b_1, \dots, b_d \in L$. Then $\tilde{q} = 1 + \sum_{j=1}^d b_j z^j \in L[z]$ satisfies $[z^n](\tilde{q}u) = 0$ for all $n > N$, so $\tilde{q}u \in L[z]$ and $u \in L(z)$. This includes $d = 0$, when there are no unknowns and u is already a polynomial. $\square$

Theorem 21.11 (Coarsening descent; source 11). *Let $L \subseteq \mathbb{K}$ be a subfield and $\Delta \subsetneq \Gamma$ a convex subgroup with $v(L^\times) \subseteq \Delta$. Then, inside $\mathbb{K}((z))$,*

$$\mathcal{E}_\Gamma(k) \cap L[[z]] = L[z], \quad \text{Frac } \mathcal{E}_\Gamma(k) \cap L((z)) = L(z). \quad (21.7)$$

Proof. Let $u = \sum_n u_n z^n \in \mathcal{E}_\Gamma(k) \cap L[[z]]$. Every nonzero u_n has value in Δ , below any fixed $\eta > \Delta$, so by Proposition 20.2 only finitely many u_n are nonzero.

Next let $u \in \text{Frac } \mathcal{E}_\Gamma(k) \cap L[[z]]$, say $u = g/h$ with $g, h \in \mathcal{E}_\Gamma(k)$ and $h \neq 0$. Normalize h by c as in Lemma 21.9. Since v_Δ vanishes on $L^\times$, red_Δ embeds L as a subfield $\bar{L} \subseteq k_\Delta$: every nonzero element of L is a unit of $\mathcal{O}_\Delta$, and so has nonzero residue. In the formal identity

$$g/c = (h/c)u \quad (21.8)$$

every coefficient on the right is a *finite* sum of products of elements of $\mathcal{O}_\Delta$, so g/c has coefficients in $\mathcal{O}_\Delta$; both g/c and h/c lie in $\mathcal{E}_\Gamma(k)$. By Lemma 21.9 their reductions are polynomials, the second nonzero, and reducing (21.8) coefficientwise gives $\overline{h/c} \bar{u} = \overline{g/c}$ in $k_\Delta[[z]]$. So $\bar{u} \in k_\Delta(z) \cap \bar{L}[[z]]$, and Lemma 21.10 gives $\bar{u} \in \bar{L}(z)$. Write $\bar{u} = \bar{p}/\bar{q}$ with $\bar{p}, \bar{q} \in \bar{L}[z]$ and $\bar{q} \neq 0$. Lift their finitely many coefficients through the isomorphism $L \rightarrow \bar{L}$ to $p, q \in L[z]$, with $q \neq 0$. Every coefficient of $qu - p$ belongs to L and has zero residue; injectivity on L therefore gives $qu = p$, hence $u \in L(z)$. This lifts a rational identity over the embedded field, not arbitrary residue coefficients from k_Δ . Finally, for $u \in \text{Frac } \mathcal{E}_\Gamma(k) \cap L((z))$ choose $m' \geq 0$ so that $z^{m'}u \in L[[z]]$, apply the power-series case, and divide by $z^{m'}$. The inclusions $\supseteq$ hold because all polynomials over L are strongly entire and their quotients belong to $\text{Frac } \mathcal{E}_\Gamma(k)$. $\square$

Remark 21.12 (Numerator and denominator together; source 11). The coefficients of g and h need not lie in L , and neither need be a polynomial. The proof works because their *common* coarsened reduction gives a polynomial equation for the quotient; asserting that $g, h \in L[[z]]$ would be unjustified.

Example 21.13 (The formal exponential; source 11). With $L = k$ and $\Delta = \{0\}$, Theorem 21.11 gives $\text{Frac } \mathcal{E}_\Gamma(k) \cap k((z)) = k(z)$ for every nonzero Γ . The formal series $\sum_n z^n/n!$ is not rational: if $r = p/q \neq 0$ with $p, q \in k[z]$ satisfied $D_z r = r$, then $p'q - pq' = pq$. The left side has degree at most $\deg p + \deg q - 1$ (or is zero), while the right side has degree $\deg p + \deg q$. This is impossible, so the formal exponential is not a quotient of strongly entire series. This says nothing about the complex exponential, which belongs to a different analytic category.

21.3 Proofs of Theorems E–G and further consequences

Proof of Theorem F (source 12). A dependence uses finitely many $\lambda \in \Lambda$ and derivative orders. Theorem 21.2 (b) with $k_0 = k$ and $K_0 = \mathbb{K}$ puts every a_n in a subfield $L \subseteq \mathbb{K}$ finitely generated over k . By Lemma 20.6, $v(L^\times)$ has finite rational rank, and it contains every coefficient value, contradicting $\text{cvr}(f) = \infty$. $\square$

Proof of Theorem E. (b) (Source 12.) By Corollary 20.11, $\text{cvr}(f) = \aleph_0$, and Theorem F applies. For q of infinite order and $\Lambda = \{q^i : i \in \mathbb{Z}\}$, a quotient $q^{i-i'}$ is a root of unity only if $i = i'$.

(a) (Sources 11 and 12.) Take $\Lambda = \{1\}$ in (b). This case needs only Theorem 21.2 (a), with k_0 generated over k by the finitely many scalar coefficients of a relation, and Corollary 20.10; no Skolem–Mahler–Lech. An algebraic dependence among $f, D_z f, \dots, D_z^r f$ becomes, after clearing denominators, a nonzero equation $P(z, f, \dots, D_z^r f) = 0$ with $P \in \mathbb{K}[z, Y]$, and conversely. *Second route* (source 11): a differentially algebraic $f \in \mathcal{E}_\Gamma(k)$ has, by Corollary 21.8, its coefficients in a field L finitely generated over $\mathbb{Q}$; by Lemmas 20.6 (with $k_0 = \mathbb{Q}$) and 20.7, $v(L^\times)$ lies in a proper convex subgroup; and the polynomial half of Theorem 21.11 gives $f \in L[z]$.

(c) (Source 11.) Let $u \in \text{Frac } \mathcal{E}_\Gamma(k)$ be differentially algebraic over $\mathbb{K}(z)$. As in the second route, its Laurent coefficients lie in a field L with $v(L^\times)$ in a proper convex subgroup Δ , and the

fraction-field half of Theorem 21.11 gives $u \in L(z) \subseteq \mathbb{K}(z)$. Conversely $\mathbb{K}(z) \subseteq \text{Frac } \mathcal{E}_\Gamma(k)$, since polynomials are entire, and rational functions are algebraic, hence differentially algebraic, over $\mathbb{K}(z)$. $\square$

Corollary 21.14 (A free differential algebra of dilates; source 12). *Under the hypotheses of Theorem E (b), the homomorphism from the differential polynomial algebra $\mathbb{K}(z)\{X_\lambda : \lambda \in \Lambda\}$, with $D_z(z) = 1$ and $D_z(\mathbb{K}) = 0$, to $\mathbb{K}((z))$ sending X_λ to $f(\lambda z)$ is injective. For $\Lambda = \{q^i : i \in \mathbb{Z}\}$ this gives countably many differentially algebraically independent functions.*

Proof. $D_z^j X_\lambda$ goes to $\lambda^j \mathcal{J}_{\lambda,j} f$, and multiplying members of an algebraically independent family by nonzero elements of the base field keeps it independent. $\square$

This is stronger than differential transcendence of each dilate separately: for $q = 2$, no nonzero polynomial over $\mathbb{K}(z)$ relates $f(z)$, $D_z f(z)$, $f(2z)$, $(D_z^2 f)(4z)$ and any further finite selection from the family.

Corollary 21.15 (Relative differential-algebraic closedness; source 11). *If Γ has no order unit, no element of $\text{Frac } \mathcal{E}_\Gamma(k) \setminus \mathbb{K}(z)$ is differentially algebraic over $\mathbb{K}(z)$.*

This is Theorem E (c) restated. It does not say that $\mathbb{K}(z)$ is differentially closed; it describes only the extension $\text{Frac } \mathcal{E}_\Gamma(k)/\mathbb{K}(z)$.

Corollary 21.16 (Finite rational differential systems; source 11). *Assume Γ has no order unit, and let $u_1, \dots, u_{m'} \in \text{Frac } \mathcal{E}_\Gamma(k)$ satisfy $D_z u_i = R_i(z, u_1, \dots, u_{m'})$ with $R_i \in \mathbb{K}(z)(Y_1, \dots, Y_{m'})$, each denominator nonzero after substitution. Then every u_i lies in $\mathbb{K}(z)$, and every strongly entire u_i is a polynomial.*

Proof. The field $\mathbb{K}(z)(u_1, \dots, u_{m'})$ is stable under D_z : the equations put each $D_z u_i$ in the field, and the quotient rule handles every rational expression in the u_i . Its transcendence degree over $\mathbb{K}(z)$ is at most m' , so the $m' + 1$ elements $u_i, D_z u_i, \dots, D_z^{m'} u_i$ are algebraically dependent. Apply Theorem E (c), and (a) for the entire ones. $\square$

Corollary 5.5 is the linear case for every $\Gamma \neq 0$; this corollary allows nonlinear rational systems and quotients of entire series, but needs the absence of an order unit.

Corollary 21.17 (Painlevé I in the fraction field; source 11). *If Γ has no order unit, the equation $D_z^2 u = 6u^2 + z$ has no solution in $\text{Frac } \mathcal{E}_\Gamma(k)$.*

Proof. A solution would be rational by Theorem E (c), and $u = 0$ fails. A nonzero rational u has a leading term cz^d at infinity. If $d \geq 1$, then $6u^2$ has degree $2d$, above the degree 1 of z and the degree at most $d - 2$ of $D_z^2 u$. If $d \leq 0$, then z has degree 1 while u^2 and $D_z^2 u$ have degree at most 0. No cancellation is possible. $\square$

Proposition 16.3 already excludes strongly entire solutions for every $\Gamma \neq 0$; the corollary extends this to quotients of entire series when there is no order unit. It is not a statement about local formal solutions, which exist for all $a_0, a_1 \in \mathbb{K}$ through $(n + 1)(n + 2)a_{n+2} = 6 \sum_{j=0}^n a_j a_{n-j} + \mathbf{1}_{n=1}$, nor about classical complex solutions. The conclusion cannot be strengthened from rational to constant: for every $c \in \mathbb{K}$, $u = 1/(z - c) \in \text{Frac } \mathcal{E}_\Gamma(k)$ satisfies $D_z u = -u^2$, and at $c = 0$ it has a pole at the origin, which is why the Laurent form of Theorem 21.11 is needed (source 11).

Corollary 21.18 (A solution-dependent exterior certificate; source 12). *Let $f \in \mathbb{K}[[z]]$ be a nonpolynomial formal series, differentially algebraic over $\mathbb{K}(z)$, and let $L \supseteq k$ be a field finitely generated over k that contains its coefficients, as supplied by Theorem 21.2 (a) with k_0 generated over k by the scalar coefficients of a relation. If the smallest convex subgroup Δ containing $v(L^\times)$ is proper, every x with $v_\Delta(x) < 0$ lies outside $\text{Dom}_{\text{str}}(f)$. If Γ has no order unit, Δ is always proper.*

Proof. Lemmas 20.6 and 20.7 make $\Delta = \{0\}$ or $\Delta = \Delta_\beta$ with $\beta \in v(L^\times)$; apply Proposition 20.9. $\square$

The certificate is structural, not a uniform algorithm for arbitrary Hahn names: it can need the values of rational functions of the finitely many generators of L , which their raw supports do not supply. It differs from Theorem 15.1, which needs $\chi_P \neq 0$ and finitely many leading valuations of the solution but no hypothesis on convex subgroups.

21.4 The order-unit boundary

Proposition 21.19 (A one-generator coefficient field; source 12). *Let $\mu > 0$ be an order unit of Γ and $p = t^\mu$. The partial theta series Θ_p of (8.1) is strongly entire and nonpolynomial, satisfies $\Theta_p(z) - z\Theta_p(pz) - 1 = 0$, has all its coefficients in $k(p)$, and p has infinite multiplicative order.*

Proof. All coefficients are nonzero and $v(p^{m'}) = m'\mu \neq 0$ for $m' \neq 0$. Given γ, β , choose positive integers A, B with $\gamma \geq -A\mu$ and $\beta \leq B\mu$; then $v(p^{n(n-1)/2}) + n\gamma \geq (n(n-1)/2 - An)\mu > \beta$ for large n , and Proposition 20.2 applies. The functional identity is (8.2). $\square$

For $k = \mathbb{C}$ this is also a case of Theorem 8.1; the short proof above works over any k .

Proof of Theorem G (source 12). Proposition 21.19 gives (i) $\Rightarrow$ (ii) and (i) $\Rightarrow$ (iii): the identity $\Theta_p(z) - z\Theta_p(pz) - 1 = 0$ is the nonzero polynomial $Y_0 - zY_1 - 1$ vanishing at $(\mathcal{J}_{1,0}\Theta_p, \mathcal{J}_{p,0}\Theta_p)$. If (ii) holds without an order unit, Corollary 20.10 contradicts nonpolynomiality, so (ii) $\Rightarrow$ (i). A dependence as in (iii) involves finitely many powers of q and derivative orders, and Theorem 21.2 (b) puts every coefficient in a field finitely generated over k , so (iii) $\Rightarrow$ (ii). $\square$

Theorem B and Corollary 11.3 detect an order unit by linear dilation equations with a cofinal parameter; Theorem G detects it by coefficient fields and by mixed jets, with the same partial theta witness. As proved by source 12, neither said that an order unit produces a nonpolynomial strongly entire *differentially* algebraic series: the witness's equation contains a dilation. That was Question 19.1 before source 13; clause (iv), proved in Section 25.1 from Theorem H, now says it, with the witness $\mathcal{T}_p$.

What “no order unit” does and does not mean (sources 11 and 12). It is not a synonym of non-Archimedean or higher rank: $\mathbb{Q} \oplus \mathbb{Q}$ ordered lexicographically, with the first coordinate dominant, has two Archimedean classes, and $(1, 0)$ is an order unit; so has every finite lexicographic product. An Archimedean group, $\mathbb{R}$ for instance, can have infinite rational rank and an order unit, and an infinite-rank group has an order unit when it has a largest Archimedean class. The increasing direct sum (11.1) has none. The proofs use exactly that every Δ_β is proper. With an order unit a finitely generated coefficient field can already have values whose convex hull is all of Γ ; then Lemma 20.7 forces no proper coarsening, and the argument stops for that definite reason. The series G of Corollary 17.1 over $\mathbb{C}((t^\mathbb{Q}))$ is the example of source 11: strongly entire, with coefficients

in $\mathbb{Q}(t)$ whose value group $\mathbb{Z}$ has convex hull $\mathbb{Q}$. The collection already contains G and its first-order consequence, and neither source makes a new all-order claim for it.

Theorem 21.20 (An explicit rank-one family; source 12). *Let $k = \mathbb{C}$, $\Gamma = \mathbb{R}$, and let $\tau > 1$ be a transcendental real number. The series $R_\tau(z) = \sum_{n \geq 0} t^{\tau^n} z^n$ is strongly entire over $\mathbb{C}((t^\mathbb{R}))$, and for every $\Lambda \subseteq \mathbb{C}((t^\mathbb{R}))^\times$ with pairwise quotients that are not roots of unity, all $(D_z^j R_\tau)(\lambda z)$ are algebraically independent over $\mathbb{C}((t^\mathbb{R}))(z)$.*

Proof. For real γ , $\tau^n + n\gamma \rightarrow +\infty$, so Proposition 20.2 gives strong entireness. A nontrivial rational relation among distinct τ^n would be a nonzero polynomial equation for τ over $\mathbb{Q}$, so $\text{cvr}(R_\tau) = \infty$, and Theorem F applies. $\square$

The group $\mathbb{R}$ has an order unit, so Theorem E does not apply; the theorem uses the infinite rational rank of $\mathbb{R}$ through Theorem F. It says nothing about a series whose coefficient values span only $\mathbb{Q}\mu$, such as Θ_p or G , and it does not settle the purely differential problem over $\mathbb{C}((t^\mathbb{Q}))$. (Source 17's lacunary series $\mathcal{G}_A$ have coefficient values spanning $\mathbb{Q}\mu$ and are differentially transcendental over every order-unit field, including $\mathbb{C}((t^\mathbb{Q}))$, Theorem M; Θ_p and G remain undecided.) Through (2.4) its surreal realization is $\sum_{n \geq 0} \omega^{-\tau^n} z^n$ on the image of $\mathbb{C}((t^\mathbb{R}))$ in $\mathbf{No}[i]$, with independence over the full scalar field of that workspace.

22 Explicit families, surreal realizations and boundaries

22.1 The explicit series in all orders

Theorem 22.1 (Full mixed-jet independence of the explicit series; source 12). *Let F be the strongly entire series (11.2) over K_{Γ_∞} . For every $\Lambda \subseteq K_{\Gamma_\infty}^\times$ with pairwise quotients that are not roots of unity, the family $\{(D_z^j F)(\lambda z) : \lambda \in \Lambda, j \geq 0\}$ is algebraically independent over $K_{\Gamma_\infty}(z)$. In particular F satisfies no nonzero algebraic differential equation of any order over $K_{\Gamma_\infty}(z)$.*

Proof. Strong entireness is Theorem 11.2; source 12 reproves it by the highest-coordinate argument recorded there. The values e_n are rationally independent, so $\text{cvr}(F) = \infty$ and Theorem F applies; Theorem E gives the same, since Γ_∞ has no order unit. $\square$

Source 11 derives the purely differential part for $f_* = F - \omega^{-1}$ from Theorem E (a); by the remark on indexing in Section 20.1 this is the same statement. The theorem upgrades the linear rigidity of Theorem 11.2 and the two-jet independence of Corollary 17.2; the series itself is not new. For example, all formal series $(D_z^j F)(q^i z)$, $i \in \mathbb{Z}$, $j \geq 0$, for any one of $q = 2$, $q = 1 + t^{e_0}$ or $q = t^{e_0}$ are jointly algebraically independent over $K_{\Gamma_\infty}(z)$: the first two parameters have valuation zero, and the third has a positive valuation that is not an order unit (source 12).

22.2 Joint independence and a continuum-sized family

This subsection is source 11's.

Theorem 22.2 (Infinite independent coefficient supply; source 11). *Let $k_0 \subseteq K_0$ be fields of characteristic zero and $f_i = \sum_{n \geq 0} a_{i,n} z^n \in K_0[[z]]$ for $i \in I$. For every nonempty finite $J \subseteq I$ and $i \in J$ put $k_{J,i} = k_0(a_{i',n} : i' \in J \setminus \{i\}, n \geq 0)$, and suppose*

$$\text{trdeg}_{k_{J,i}} k_{J,i}(a_{i,n} : n \geq 0) = \infty \quad (22.1)$$

for all such J and i . Then the jets $D_z^j f_i$, $i \in I$, $j \geq 0$, are algebraically independent over $K_0(z)$.

Proof. Otherwise choose an inclusion-minimal nonempty finite J whose members have algebraically dependent jets, and fix $i \in J$. By minimality the jets of the other members are algebraically independent over $K_0(z)$. Take a nonzero relation, clear its denominators in z , and let $c_1, \dots, c_q \in K_0$ be its scalar coefficients. As a polynomial in the jets of f_i its coefficients are polynomials in z and in the jets of the other members, and they do not all vanish after substitution, because that substitution is injective by the independence just noted. So f_i satisfies a nonzero differential equation with coefficients in $k_{J,i}(c_1, \dots, c_q)[[z]]$, and Theorem 21.2 (a) puts all $a_{i,n}$ in $k_{J,i}(c_1, \dots, c_q, a_{i,0}, \dots, a_{i,N-1})$, of transcendence degree at most $q + N$ over $k_{J,i}$, contradicting (22.1). $\square$

The criterion does not require disjoint coefficient fields; finite overlaps are harmless. It assumes nothing about the Hahn supports of constants in a relation, which may be arbitrary elements of K_0 .

Lemma 22.3 (Independent scale monomials; source 11). *The monomials $t^{e_0}, t^{e_1}, \dots$ in $k((t^{\Gamma_\infty}))$ are algebraically independent over k .*

Proof. A polynomial in finitely many of them is a finite sum of monomials $t^{\sum_j m_j e_j}$; distinct exponent vectors give distinct elements of Γ_∞ , so no cancellation occurs. $\square$

For a binary word $b = (b_1, \dots, b_n)$ with $n \geq 1$ put

$$\text{code}(b) = 2^n + \sum_{j=1}^n b_j 2^{n-j} \in [2^n, 2^{n+1} - 1]. \quad (22.2)$$

Distinct words have distinct codes, codes strictly increase along an infinite branch, and two distinct branches share only finitely many prefixes, so their code sets meet in a finite set. The almost-disjoint prefix device is standard and also appears, with a different coefficient construction, in the tail-span report [16]. For $\alpha \in \{0, 1\}^{\mathbb{N}_{>0}}$ put

$$\mathcal{F}_\alpha(z) = \sum_{n \geq 1} t^{e_{\text{code}(\alpha|n)}} z^n, \quad \alpha|n = (\alpha_1, \dots, \alpha_n). \quad (22.3)$$

Each nonzero coefficient is a Hahn monomial with coefficient 1 in the prime field.

Proposition 22.4 (Source 11). *Every $\mathcal{F}_\alpha$ is strongly entire over $k((t^{\Gamma_\infty}))$.*

Proof. Fix $\gamma, \beta \in \Gamma_\infty$, both supported on $\{e_0, \dots, e_{n_*}\}$. Once $\text{code}(\alpha|n) > n_*$, the largest nonzero coordinate of $e_{\text{code}(\alpha|n)} + n\gamma - \beta$ is at index $\text{code}(\alpha|n)$ and equals 1, so it is positive. Proposition 20.2 applies. $\square$

Proposition 22.5 (Source 11). *Over $K = k((t^{\Gamma_\infty}))$, all $D_z^j \mathcal{F}_\alpha$, with $\alpha \in \{0, 1\}^{\mathbb{N}_{>0}}$ and $j \geq 0$, are algebraically independent over $K(z)$.*

Proof. Given finitely many distinct branches and one of them, take the largest common-prefix length with the other branches, or 0 if there are none. Beyond that length every code of the chosen branch occurs on none of the others. So the chosen series has infinitely many coefficient monomials t^{e_c} indexed by codes absent from all other coefficient fields, and by Lemma 22.3 they are algebraically independent over the field generated by the other branches' monomials. Indeed, a putative polynomial relation uses only finitely many private monomials and rational functions of finitely many other monomials. Clearing denominators would contradict the algebraic independence of that finite combined family. Condition (22.1) holds with $k_0 = k$, and Theorem 22.2 applies. $\square$

Theorem 22.6 (Maximal differential transcendence in one workspace; source 11). *Let $k = \mathbb{C}$ and $K_{\Gamma_\infty} = \mathbb{C}((t^{\Gamma_\infty}))$. The family $(\mathcal{F}_\alpha)_{\alpha \in \{0,1\}^{\mathbb{N}_{>0}}}$ of strongly entire series has all its jets jointly algebraically independent over $K_{\Gamma_\infty}(z)$, and*

$$\text{dtrdeg}_{K_{\Gamma_\infty}(z)} \text{Frac } \mathcal{E}_{\Gamma_\infty} = 2^{\aleph_0}. \quad (22.4)$$

Proof. Propositions 22.4 and 22.5 give a differentially independent family indexed by the $2^{\aleph_0}$ branches, so the differential transcendence degree is at least $2^{\aleph_0}$. Conversely Γ_∞ is countable and $|\mathbb{C}| = 2^{\aleph_0}$, so $|K_{\Gamma_\infty}| \leq |\mathbb{C}^{\Gamma_\infty}| = 2^{\aleph_0}$ and $|K_{\Gamma_\infty}[[z]]| = 2^{\aleph_0}$; no differentially independent subset of the fraction field can be larger. $\square$

The lower bound holds for every k ; the equality is stated, as by source 11, for $k = \mathbb{C}$. Equivalently, the free differential polynomial algebra $K_{\Gamma_\infty}(z)\{X_\alpha\}$ embeds in the fraction field by $X_\alpha \mapsto \mathcal{F}_\alpha$. This is cardinal maximality only: the family is not claimed maximal under inclusion, nor to be a differential transcendence basis.

The tail-span report already constructs continuum-sized differentially independent families [16], using algebraic coefficient extensions and Galois sign changes over smaller fields, and its analytic construction lives on a fixed ordinary disk rather than being strongly entire at every Hahn scale. Here the coefficients are exponent monomials, the functions are strongly entire on one full Hahn field, and the independence is over that whole field as field of constants. Neither the cardinal $2^{\aleph_0}$ nor the almost-disjoint combinatorics is claimed new.

22.3 Surreal realization, scalar extension, whole-class failure

Through (2.4), with $e_j \mapsto \omega^j$, the branch family becomes

$$\mathcal{F}_\alpha(z) = \sum_{n \geq 1} \omega^{-\omega^{\text{code}(\alpha|n)}} z^n,$$

and at every argument of the embedded workspace its evaluation is an actual surcomplex number. Everything is set-sized: the value group is countable, the field has the cardinality of the continuum, and no class-sized sum or unrestricted class quantification is used (source 11).

Proposition 22.7 (Independence survives extension of constants; source 12). *Let $E/\mathbb{K}$ be a field extension and (g_α) a family in $\mathbb{K}[[z]]$ algebraically independent over $\mathbb{K}(z)$. Under the coefficientwise inclusion $\mathbb{K}[[z]] \subseteq E[[z]]$ the family is algebraically independent over $E(z)$.*

Proof. A relation over $E(z)$ among finitely many g_α gives, after clearing denominators, a nonzero $Q \in E[z, X_1, \dots, X_{m'}]$ vanishing at them. Choose a $\mathbb{K}$ -basis $b_1, \dots, b_d$ of the $\mathbb{K}$ -span of its coefficients and write $Q = \sum_\nu b_\nu Q_\nu$ with $Q_\nu \in \mathbb{K}[z, X]$, some $Q_\nu \neq 0$. Each $Q_\nu(z, g)$ has coefficients in $\mathbb{K}$, so linear independence of the b_ν over $\mathbb{K}$ forces every $Q_\nu(z, g) = 0$, and a nonzero Q_ν is a relation over $\mathbb{K}(z)$. $\square$

The proposition concerns fixed formal functions and extensions of the constant field. It adjoins no new functional variable, says nothing about evaluation domains, and does not cover new dilation parameters taken from the extension. Entireness behaves quite differently. Adjoin a dominating level, $\Gamma' = \Gamma_\infty \oplus \mathbb{Q}e_*$ with $e_* > \Gamma_\infty$, realized by $e_* = \omega^\omega$; the new group has the order unit e_* . At $x = t^{-e_*} = \omega^{\omega^\omega}$ the evaluation family of F is not strongly summable (Section 11.2; source 12 repeats this computation), and neither is that of any $\mathcal{F}_\alpha$, whose leading exponents $e_{\text{code}(\alpha|n)} - ne_*$

strictly decrease since $e_{\text{code}(\alpha|n+1)} - e_{\text{code}(\alpha|n)} - e_* < 0$ (source 11). So the formal independences survive the larger field by Proposition 22.7, while entireness does not: two different kinds of stability. Nothing in this part is entire on the proper class $\mathbf{No}[i]$.

The derivative D_z kills every scalar, including every $\omega^{-\gamma}$, and $D_z(z) = 1$. The Berarducci–Mantova derivation acts on surreal scalars and is a different operation [8]. The tail-span report’s all-order independence results for particular constructions, other base fields and other derivations [16] are not restated here, and neither kind of statement is transferred to the other’s derivation. The independence statements concern formal functions: at a point $x \in \mathbb{K}$ every value $f(x)$ lies in $\mathbb{K}$ and cannot be transcendental over it, and at $z = 0$ the j th jet has the value $j!a_j \in \mathbb{K}$ (sources 11 and 12).

22.4 Boundary examples

Positive characteristic (sources 11 and 12). If k has characteristic $p > 0$, the series $\sum_{n \geq 0} t^{e_n} z^{pn}$ over $k((t^{\Gamma_\infty}))$ is strongly entire, by the highest-coordinate argument and Lemma 2.3 (a), whose proof does not use the characteristic, yet has D_z -derivative zero. This is the phenomenon of Example 17.3 in the no-order-unit workspace; characteristic zero is essential even for order-one linear equations, and the finite coefficient-field principle fails there. No claim about alternative derivations in positive characteristic is made.

Torsion-related dilations (source 12). Let $\zeta \in k$ be a root of unity of order $m' \geq 2$ and, over $k((t^{\Gamma_\infty}))$, let $g(z) = F(z^{m'}) = \sum_n t^{e_n} z^{m'n}$. The same highest-coordinate argument makes g strongly entire, and $\text{cvr}(g) = \infty$, yet $g(\zeta z) - g(z) = 0$; for $\zeta = -1$, an even series relates its dilates by 1 and -1 . The non-root-of-unity condition therefore cannot be dropped from Theorems E (b) and F; compare Section 9.

Local evaluation is not entireness (source 12). The formal exponential E_η of Proposition 5.3, whose proof works over any k , satisfies $D_z E_\eta = t^\eta E_\eta$, has its coefficients in $k(t^\eta)$, and is strongly evaluable exactly on $\{0\} \cup \{v(x) > -\eta\}$. In higher rank the leading exponents $nv(y)$ of its evaluation need not escape cofinally when $v(y) > 0$, which is why the positive-support lemma, not the cofinal criterion, decides that domain (Remark 2.5).

Finite coefficient-value rank is not a converse (source 12). Theorem F is a sufficient condition. A series with $\text{cvr}(f) < \infty$ may still be differentially transcendental, and finite $\text{cvr}(f)$ does not by itself put the coefficients in a finitely generated field, since residues and tails can carry further algebraic complexity. The implications proved are

$$\text{a mixed relation} \implies \text{a finitely generated coefficient field} \implies \text{cvr}(f) < \infty,$$

not their converses.

An explicit example, added here, is

$$f(z) = F(z) + \frac{1}{1-z} = \sum_{n \geq 0} (1 + t^{e_n}) z^n \quad \text{over } K_{\Gamma_\infty}.$$

Every coefficient has value zero, so $\text{cvr}(f) = 0$, while subtracting 1 recovers the algebraically independent monomials of Lemma 22.3; its coefficient field has infinite transcendence degree. This

f is not strongly entire, by Proposition 20.2 with $\gamma = \beta = 0$. Nevertheless

$$(D_z^j f)(\lambda z) = (D_z^j F)(\lambda z) + \frac{j!}{(1 - \lambda z)^{j+1}}.$$

For every finite selection of jets the added terms belong to $K_{\Gamma_\infty}(z)$, so translation by them is a polynomial-algebra automorphism. The full mixed independence of Theorem 22.1 therefore holds for f with the same restriction on the dilation quotients, despite its zero coefficient-value rank.

22.5 Consequences for the questions of Section 19

Question 19.1. Theorem E (a) (sources 11 and 12) answers it for every Γ without an order unit, in every order: no nonpolynomial strongly entire series satisfies a nonzero algebraic differential equation over $\mathbb{K}(z)$, whatever its residue corner, for every characteristic-zero k , without divisibility, with arbitrary Hahn coefficients in the equation. Theorem E (c) (source 11) extends this to quotients of entire series. The last sentence of the question as posed by sources 08 and 09 is answered for F with no coefficient restriction (Theorem 22.1, source 12; source 11 for f_*). The question's sentence about hypotheses on Γ is answered in one direction: the absence of an order unit suffices. With an order unit it stayed open after sources 11 and 12 and was re-scoped to that case; source 13 has since answered it there (Theorem H), source 14 has excluded order two for every Γ (Theorem I), and it is now re-scoped to the classification of series of minimal order three in Section 19.

Question 19.4. Theorem E (b) with $\Lambda = \{q^i : i \in \mathbb{Z}\}$ (source 12) excludes, for every nontorsion q including those with $v(q) = 0$, every nonzero algebraic relation among the $\mathcal{J}_{q^i, j} f$ of a nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ when Γ has no order unit. Since $D_z^r \sigma_q^\ell f = q^{\ell r} \mathcal{J}_{q^\ell, r} f$, a nonzero equation (10.1) is such a relation, and so are nonlinear mixed relations. With an order unit the unit-valuation case stayed open after sources 11 and 12; the witness of Theorem G uses the parameter t^μ of nonzero valuation and says nothing there. Source 18 has since answered it for linear equations and every Γ (Corollary 29.6). Theorem 10.2 and Theorem E (b) do not contain each other: the first covers every $\Gamma \neq 0$ but only nonzero noncofinal $v(q)$, the second every nontorsion q but only groups without an order unit. Theorem 29.4 contains the first and, for linear relations, the second's cyclic case, and is contained in neither (Remark 29.5).

Question 19.6: a merge observation. Source 12 does not claim this question. The following consequence of its theorem was checked for this merge.

Corollary 22.8 (Several multiplicatively independent dilations; merge observation). *Suppose Γ has no order unit, and let $q_1, \dots, q_s \in \mathbb{K}^\times$ be multiplicatively independent, that is, $q_1^{m_1} \cdots q_s^{m_s} = 1$ with $m_i \in \mathbb{Z}$ only for $m_1 = \cdots = m_s = 0$. Let Λ be the subgroup of $\mathbb{K}^\times$ they generate. Then for every nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ the family $\{\mathcal{J}_{\lambda, j} f : \lambda \in \Lambda, j \geq 0\}$ is algebraically independent over $\mathbb{K}(z)$. In particular no nonpolynomial strongly entire series satisfies a nonzero equation, linear or not, in finitely many $(D_z^j f)(q_1^{m_1} \cdots q_s^{m_s} z)$. For the series F of (11.2), the only linear relation over $K_{\Gamma_\infty}(z)$ among $F(\lambda_1 z), \dots, F(\lambda_{s'} z)$ is the trivial one whenever $\lambda_1, \dots, \lambda_{s'} \in K_{\Gamma_\infty}^\times$ have pairwise quotients that are not roots of unity, for instance when they are distinct elements of a group generated by multiplicatively independent parameters.*

Proof. Λ is free abelian on the q_i , hence torsion free. For distinct $\lambda, \lambda' \in \Lambda$ the quotient λ/λ' is a nonidentity element of Λ , so no positive power of it is 1: it is not a root of unity. Apply Theorem E (b), and Theorem 22.1 for the last clause. $\square$

So the first sentence of Question 19.6 has, in a group without an order unit, the answer that no nonpolynomial strongly entire solution exists, and its last sentence has the answer that only the trivial relation occurs whenever no quotient of parameters is a root of unity. The existence criterion in groups with an order unit, and relations among parameters with a root-of-unity quotient, remained open, and the question was re-scoped accordingly in Section 19; source 18 has since given the existence criterion for some linear equation supported in a prescribed finite set of multipliers (Theorem 29.9), recorded in the status of the question.

Mixed jets in the fraction field. The coefficient theorem of source 12 and the fraction-field descent of source 11 also give the following consequence. Neither source states it; it is added here by combining their results.

Corollary 22.9 (Mixed rigidity in the fraction field). *Suppose Γ has no order unit, and let $u \in \text{Frac } \mathcal{E}_\Gamma(k) \setminus \mathbb{K}(z)$. If $\Lambda \subseteq \mathbb{K}^\times$ has no root-of-unity quotient between distinct elements, the Laurent series $\{(D_z^j u)(\lambda z) : \lambda \in \Lambda, j \geq 0\}$ are algebraically independent over $\mathbb{K}(z)$.*

Proof. Choose $m' \geq 0$ so that $y = z^{m'} u \in \mathbb{K}[[z]]$; also $y \in \text{Frac } \mathcal{E}_\Gamma(k)$. If the stated family were dependent, a nonzero relation would involve finitely many parameters and derivative orders. Adjoin the lower orders through the largest order r used, and for each parameter use Leibniz's formula, with differentiation before dilation:

$$(D_z^j u)(\lambda z) = \sum_{\ell=0}^j \binom{j}{\ell} (-m')_{j-\ell} (\lambda z)^{-m'-j+\ell} (D_z^\ell y)(\lambda z).$$

For each λ this is a triangular change of jet variables over $\mathbb{K}(z)$ with nonzero diagonal $(\lambda z)^{-m'}$. The combined change is therefore an automorphism of the finite polynomial algebra of jet variables, so it carries the relation to a nonzero relation among the jets of y . Clear its rational denominators and apply Theorem 21.2 (b), with $k_0 = k$ and $K_0 = \mathbb{K}$. All coefficients of y lie in a field $L \subseteq \mathbb{K}$ finitely generated over k . Lemmas 20.6 and 20.7 put $v(L^\times)$ in a proper convex subgroup. Since $y \in \text{Frac } \mathcal{E}_\Gamma(k) \cap L[[z]]$, the fraction-field part of Theorem 21.11 gives $y \in L(z)$. Thus $u = z^{-m'} y \in \mathbb{K}(z)$, a contradiction. $\square$

The excluded class is the rational functions: a nonpolynomial rational function still satisfies a polynomial relation over $\mathbb{K}(z)$. Taking $\Lambda = \{1\}$ recovers the purely differential statement in Theorem E (c). This consequence uses the same characteristic-zero and quotient hypotheses as the mixed coefficient theorem. The no-order-unit hypothesis supplies the proper convex subgroup needed for descent.

22.6 Predecessors, repository comparison and priority

Literature. Source 11 consulted the NUMDAM bibliographic page of Hurwitz's paper [23] and its attribution in [24], without a complete fresh reading of Hurwitz's proof; it read the parsed text of Laohakosol–Kongsakorn–Ubolsri and viewed its pages 404, 408 and 409, where Lemma 3 isolates a high coefficient through a polynomial in its index; it consulted the abstract of Krattenthaler–Rivoal [25] as arithmetic context, not as an imported theorem, and the framework of [26] for context, using none of its software. Source 12 read the general characteristic-zero form of the Skolem–Mahler–Lech theorem in Bell's Theorem 1.1 [27], imports only that classical theorem and not Bell's affine-variety generalization, and checked Lech's paper [28] through publisher metadata

only, without reconstructing its proof. Both consulted Hu–Luan [20], whose setting is an algebraically closed field of characteristic zero, complete for a nontrivial real-valued non-Archimedean absolute value; its hypotheses are not transferred to arbitrary-rank Hahn fields, its results are neither inferred nor replaced, and no rank-one question is claimed resolved. For normal forms source 11 used Mantova–Matusinski, Section 2.2 [17], and source 12 Bournez–Guilmant, Corollary 2.7 [39], with the same convention $t^a \mapsto \omega^{-a}$ as (2.4).

Repository comparison at the pin. Both sources compared their results with the collection at 465a54b, at which this report consisted of Sections 1–19 and the conclusion and was unchanged up to the merge. Their statements about it were checked and are accurate as of that pin: Question 19.1 left order at least two with vanishing residue corner open, and the README named $z^2ff'' + zff' - z^2(D_zf)^2 = 0$; Question 19.4 is the unit-valuation mixed case (source 12’s phrase “including unit-valued parameters” refers to exactly that); the source 10 audit file still says “Question 15.1”, as source 11 notes; the report already contained the sufficiency half of the all-scale criterion, the series F and its domain warning, first-order independence and the polynomial Painlevé exclusions. The entire-function report [12] contains the polynomial intersection obstruction $\mathcal{E}_\Delta \cap K_\Gamma[[Z]] = K_\Gamma[Z]$ for noncofinal extensions, coarsened support methods and the distinction between countable cofinality and order units; the tail-span report [16] contains continuum-sized families built with binary-prefix almost-disjoint sets. Since the pin, the statements of this report that called the vanishing-corner case open have been re-scoped as recorded in Section 19.

Source 11 inspected the root README, the catalogue, directory listings, this report’s README, the source 10 audit, and the READMEs of the entire-function and tail-span reports; the `docs/new` listing then contained only a README. Source 12 inspected the root README, the catalogue, this report’s README (rereading its lines 131–182) and the tail-span README, and identified the foundations and entire-function reports through the catalogue. Neither was a line-by-line audit of the manuscripts, Lean declarations, archived deliveries, branches or history.

Priority. Both literature searches were made on 22 September 2026, were targeted, and included no exhaustive MathSciNet, zbMATH, Zentralblatt, full-book or citation-network review; some searches returned irrelevant or inaccessible material, and no failed search is used as evidence of absence. The inspected sources did not show the meromorphic descent theorem with its no-order-unit consequence, the full-constant-field entire branch family, the all-order no-order-unit and joint-dilation conclusions, the coefficient-value-rank criterion or the order-unit dichotomy. These are proposed contributions with proofs, not certified historical firsts, and the answered questions are questions of this report, not named published conjectures.

22.7 What the coefficient-field part does not claim

Every limitation stated by source 11 or source 12, in its manuscript, README or audit files, is kept. The tag says which source states it.

- (C1) [11, 12] The purely differential case with an order unit is not settled, and no counterexample is given. Partial theta, Θ_p or G , is not a counterexample to purely differential rigidity: its equation contains a dilation, and no new all-order claim is made for it. (Source 13 settles the case with the series $\mathcal{T}_p$ of Theorem H; Θ_p and G remain undecided.)

- (C2) [11, 12] The delivered sources supply no Lean formalization of this theorem package or independent referee report. Source 11 performed no Lean build of the repository. The finite checks provide no computer-algebra certification of an infinite statement.
- (C3) [11, 12] Historical priority is not certified; the literature searches were targeted, with the access limits recorded in Section 22.6, and a failed search is not evidence.
- (C4) [11, 12] The repository comparison was targeted, not line by line; there is no blanket claim that no equivalent result occurs elsewhere in the collection.
- (C5) [11, 12] Not claimed new: classical coefficient recurrences (Hurwitz, Laohakosol–Kongsakorn–Ubolsri; Krattenthaler–Rivoal as context), the finite-generation principle, the valuation-rank inequality, Hahn algebra and Hahn–Neumann support, Conway normal forms, the Skolem–Mahler–Lech theorem, partial theta series, the all-scale criterion, the polynomial half of the descent theorem, continuum cardinality and almost-disjoint combinatorics, and the series F (or f_*) with its two-jet consequence.
- (C6) [11, 12] *Entire* means strong evaluation on one set-sized field, not on the class $\mathbf{No}[i]$ and not ordinary convergence of constant coefficients; entireness fails after an exterior scale is adjoined, with the explicit witness $x = \omega^{\omega^{\omega}}$.
- (C7) [11, 12] D_z is not the Berarducci–Mantova derivation; nothing is said about scalar surreal differential equations, and nothing is transferred to the tail-span report’s derivations.
- (C8) [11, 12] Independence concerns formal functions, not point values.
- (C9) [11] Containment of $v(L^\times)$ in a proper convex subgroup is sufficient for descent, not claimed necessary (Question 19.7); a criterion using rational rank alone will not suffice.
- (C10) [11] Corollary 21.15 does not say that $\mathbb{K}(z)$ is differentially closed.
- (C11) [11] The continuum family is cardinally maximal only; it is not maximal under inclusion and not a differential transcendence basis.
- (C12) [11] Corollary 21.17 concerns $\text{Frac } \mathcal{E}_\Gamma(k)$; local formal solutions exist, and nothing is said about classical complex solutions.
- (C13) [11] “Meromorphic” means membership of $\text{Frac } \mathcal{E}_\Gamma(k)$, not a sheaf and not a fine-topology notion.
- (C14) [11, 12] Characteristic zero is essential, and no claim is made about alternative derivations in positive characteristic.
- (C15) [11, 12] There is no uniform algorithm for extracting a convex subgroup from Hahn names; Corollary 21.18 is structural and Theorem 21.2 (b) existential.
- (C16) [11, 12] Hu–Luan’s rank-one hypotheses are not transferred; Hu–Luan is context only, and no rank-one question is claimed resolved.
- (C17) [11, 12] The finite checks prove no arbitrary-support, continuum, recurrence zero-set, cofinality or transcendence statement.
- (C18) [11] No font files or third-party texts are distributed.
- (C19) [12] The answered questions are questions of this research programme, not named historical or published conjectures.

- (C20) [12] The Skolem–Mahler–Lech theorem is imported through Bell’s Theorem 1.1; Bell’s affine generalization is not used, Lech’s proof was not reconstructed, and the zero multipliers sampled by the checks are not an effective bound.
- (C21) [12] Theorem F is sufficient only; finite $\text{cvr}(f)$ does not give a finitely generated coefficient field.
- (C22) [12] Proposition 22.7 covers extension of constants only, not new dilation parameters from the extension.
- (C23) [12] Theorem 21.20 does not settle the purely differential problem over $\mathbb{C}((t^{\mathbb{Q}}))$.
- (C24) [12] Dilations with a root-of-unity quotient are excluded (Question 19.9).
- (C25) [12] In groups of uncountable cofinality there is no nonpolynomial strongly entire series, so statements requiring one are vacuous (Corollary 20.3). The coefficient-value-rank criterion for arbitrary formal series has no such restriction.
- (C26) [merge] Corollary 22.8 is a consequence assembled from source 12. Corollary 22.9 is an additional consequence of sources 11 and 12 together. Neither is attributed to the delivered sources as a stated result, and no historical priority is claimed for either deduction.

23 Theta descent: an entire series with an equation of order three

This section and the next two contain the material of source 13, delivered after the coefficient-field part had been merged. They answer the order-unit case of Question 19.1 as it stood after that merge: when Γ has an order unit, a nonpolynomial strongly entire series satisfies a nonzero algebraic differential equation of order three and none of lower order (Theorem H). With Theorem E (a) this makes the order unit the exact boundary of differential algebraic rigidity. The ingredients are classical: a bilateral theta series, Jacobi’s triple product and the theta–elliptic differential identity. What source 13 adds is their descent to an ordinary power series by the substitution $z = u + u^{-1}$, the support control that makes the descended series strongly entire exactly when its scale is an order unit, and a difference-field proof that no equation of order below three exists. The method is neither the escape chain, nor the finite initial polynomial, nor the finite coefficient field; the last is used only for the converse direction, which is Theorem E (a).

23.1 Source, conventions and the merge’s choices

Convention 23.1 (Setting of the theta-descent part). In Sections 23–25, k is an arbitrary field of characteristic zero with the trivial valuation, Γ is as in Convention 1.1, $\mathbb{K} = k((t^{\Gamma}))$, and v , lc , res , strong summability, $\text{Dom}_{\text{str}}(f)$, $\mathcal{E}_{\Gamma}(k)$, $\text{Frac } \mathcal{E}_{\Gamma}(k)$ and D_z are as in Conventions 14.1 and 20.1. Throughout, $\mu \in \Gamma$ is positive, $p = t^{\mu}$, and Δ_{μ} is the principal convex subgroup (2.1); μ is an order unit only where this is stated, and then $\Delta_{\mu} = \Gamma$. The power-series variable is z , while x and y denote elements of $\mathbb{K}$, and u is the variable of the Laurent series of Section 24.5. Write $\vartheta = zD_z$, as in the proof of Proposition 16.4, and $\vartheta_u = uD_u$. The elliptic constants below use the nome p^2 , not p .

For $k = \mathbb{C}$ one has $\mathbb{K} = K_{\Gamma}$ and $\mathcal{E}_{\Gamma}(k) = \mathcal{E}_{\Gamma}$. As for the two previous parts, the generality in k is recorded for this part only.

The source. Source 13 is *Theta Descent and the Exact Boundary of Differential Rigidity* (21 pages, 23 September 2026). It consulted the repository at

4cdeaec43ff1692ed2ad9f5bdf761a41872975a5,

at which this report was exactly the five-source text into which it is merged here; its references to Question 19.1 (`hol:q:nonlinear`), to its statement and status passage, to Theorem E and to the coefficient-field part were accurate there. It assumes characteristic zero, $\Gamma \neq 0$ set-sized, and, where stated, that the scale μ is an order unit. It uses no divisibility, no rank hypothesis, no real or algebraic closedness and no countable cofinal subset. Its standing hypotheses are therefore those of the coefficient-field part, and every proof taken from it was re-read against the statement it is attached to.

Renamed symbols. The following symbols of source 13 would collide with symbols of the earlier parts. They are renamed here, in the new material only.

Source 13	Here	Reason
variable x	z	Definition 1.4; x denotes points. (Source 13's $\mathbb{K} = k((t^\Gamma))$ and $\mathcal{E}_\Gamma(k)$ are this report's.)
$h, q = t^h$	$\mu, p = t^\mu$	h is a torsion order and q a dilation parameter; μ and $p = t^\mu$ are the notation of Theorem G.
$H_h, \mathcal{D}_h$	$\Delta_\mu, \mathcal{O}_{\Delta_\mu}$	(2.1); $\mathcal{D}_P$ is a candidate set (15.12); $\mathcal{O}_\Delta$ as in Section 20.3.
F_q	$\mathcal{T}_p$	F is the series (11.2).
Θ_q (bilateral)	Ψ_p	Θ_p is the partial theta series (8.1).
r_m, P_q, j_m, e, η	$x_m, \mathcal{P}_p, 2m-1, y, \varpi$	r indexes derivative orders, P is a differential polynomial, e_n and η are elements of Γ .
$w_\gamma(F_q)$	γ_δ	The coefficient Gauss value (14.1), with $\delta = -\gamma$.
$Q = q^2, s_r(Q), c$	$p^2, \mathcal{Q}_r(p^2), \mathfrak{c}$	Q_s is a shift polynomial, s an order, c_ν a corner coefficient.
A_f, H_f, J_f, B_f	$\mathcal{A}[f], \mathcal{H}[f], \mathcal{K}[f],$ $\mathcal{B}[f]$	A_δ is a leading coefficient, $H \in \Gamma$, J indexes monomials in (14.4), B is a threshold, $\mathcal{J}_{\lambda,j}$ a jet.
$S_Q(w), W(z), a, b$	$\mathcal{Z}(w), \mathcal{W}(y), \mathbf{a}_2, \mathbf{a}_4$	z is the power-series variable; a_n are Taylor coefficients.
$\delta = u \, d/du, s = u - u^{-1}$	$\vartheta_u, \tilde{z}$	δ is an element of Γ ; s is an order.
σ, τ	S, S_a	σ_q is an argument dilation; τ is the parameter of R_τ .
$\mathcal{A}, \mathcal{M} = \text{Frac } \mathcal{A}$	$\mathcal{U}_\mu, \text{Frac } \mathcal{U}_\mu$	$\mathcal{A}[f]$ as above; $M(S)$ is a support monoid.
L, A, V (in u); Z	$\psi_1, \psi_2, \psi_3; V$	L is a coefficient field; z as above.
$\ell, B = A', R(T), \mu = q^{-1}u^{-1}$	$\mathcal{L}, \mathcal{A}_1, \Phi(T),$ written out	ℓ is a dilation power; R_τ is a series; μ is the scale.
$\vartheta = x \, d/dx; \mathbf{No}(i)$	$\vartheta = zD_z; \mathbf{No}[i]$	The same operator; not ϑ_q . The collection writes $\mathbf{No}[i]$.
Theorem 1.2	Theorem H	Theorems A–G of this report are different theorems.

The integer subscripts of the classical invariants g_2, g_3 distinguish them from the scaled series g_δ of (14.1). The shipped audit files `13-theta-descent-PROOF_AUDIT.md` and `13-theta-descent-SOURCES_AND_SCOPE.md` keep the source's notation, including F_q, Θ, q, h, L, A and σ .

Words with two meanings. The *order* of an algebraic differential equation is the highest derivative occurring in it; an *order unit* is as in Definition 1.3; the *multiplicative order* of q is as in Theorem B; and ord_u is the u -adic order of a rational function. The *minimal order* of a series is the least order of a nonzero algebraic differential equation over $\mathbb{K}(z)$ that it satisfies. *Descent* in this part means the Chebyshev substitution $z = u + u^{-1}$, not the coarsening descent of Theorem 21.11. Three theta series occur and are not interchangeable: the partial series Θ_p of (8.1), which satisfies a dilation equation; the bilateral series Ψ_p of (23.8); and the Hahn–Tate report’s series $\sum_{n \in \mathbb{Z}} (-1)^n q^{n(n-1)/2} u^n$ [40], of which $\Psi_p(u)$ is the value at the nome p^2 and the argument $-pu$. A *corner* keeps the meaning of Section 14.4: source 13’s “vanishing corner” is the vanishing of the residue corner polynomial χ_P , and the full corner polynomial I_P need not vanish with it (Proposition 24.2). A *zero* of $\mathcal{T}_p$ is a point $x \in \text{Dom}_{\text{str}}(\mathcal{T}_p)$ with $\mathcal{T}_p(x) = 0$.

Results printed once. Source 13 reproves several facts that the report already contains. They are printed once and cited: the definitions of an order unit and of strong summability (Definitions 1.3 and 1.2); its cofinal-valuation lemma, which is Lemma 2.3(a); the positive-support lemma, which is Lemma 2.1(2); the surreal normal-form embedding (2.4); the strong domain of the bilateral theta series, which is the Hahn–Tate report’s theta-domain theorem [40, `tate:thm:theta`] after the substitution just described; and its whole Section 7, the no-order-unit direction, which is Theorem E(a) with Theorem 21.2(a), Lemmas 20.6 and 20.7 and Proposition 20.9; its Euler-derivative presentation of that argument is recorded in Remark 25.1. For $k = \mathbb{C}$ the non-D-finiteness of $\mathcal{T}_p$ is also a case of Theorem A; source 13’s separate proof, valid for every k , is kept (Proposition 24.11). Everything else in source 13 is printed in Sections 23–25, with its proofs.

Where each numbered result went. Theorem 1.2 is Theorem H; Lemma 2.3 is Lemma 23.4; Lemma 3.1, Proposition 3.2 and Theorem 3.3 are Lemma 23.2, Proposition 23.3 and Theorem 23.5; Theorem 4.1 and Proposition 4.2 are Theorem 23.6 and Proposition 23.7; Theorem 5.1 is Theorem 24.1, and Remark 5.2 is Remark 24.3; Lemma 6.1, Theorem 6.2, Corollary 6.3 and Theorem 6.4 are Lemma 24.6, Theorem 24.8, Corollary 24.9 and Theorem 24.10, and Section 6.5 is Proposition 24.11; Lemmas 7.1–7.2 and Theorem 7.3 are printed once as above; Corollary 8.1 is Corollary 25.3; Examples 9.1–9.3 and Corollary 9.4 are Examples 25.4–25.6 and Corollary 25.7; Questions 11.1 and 11.2 are merged into the re-scoped Question 19.1, and Question 11.3 is Question 19.10; Appendix A is (24.8) and the expansions after (24.2); Appendix B is merged into the dependency table of Section 18.

Two steps written out, and other choices. The review of source 13 for this merge found its main line correct and two steps used but not written. Both are supplied here, marked as the merge’s. First, the minimal-order proof substitutes $z = u + u^{-1}$ into strongly entire series and uses that this is a ring homomorphism into a Laurent ring, compatible with the derivations; Lemma 24.5 states and proves this. Second, source 13 proved $\partial_u \psi_2 \neq 0$ from “formal meromorphic germs” at the simple zero $u_0 = -p$ of Ψ_p , a local expansion inside $\text{Frac} \mathcal{U}_\mu$ that it did not define; Lemma 24.7 replaces this by the global identity $\Psi_p^3 \partial_u \psi_2 = p(u - u^{-1}) + O(p^2)$. The passage from the complex elliptic identity to $\mathbb{K}$ is written as three ring maps (Section 24.3), which is the content of source 13’s “support-controlled specialization”. The merge also computes the corner data of the cleared equation (Proposition 24.2), adds clause (iv) to Theorem G, and records the consequences for Theorem E (Corollary 25.2). Source 13’s theorems, proofs and non-claims are otherwise printed as delivered, in the renamed notation.

23.2 Chebyshev descent and the exact Taylor coefficients

The monic Chebyshev polynomials $C_n \in \mathbb{Z}[z]$ are defined by

$$C_0 = 2, \quad C_1 = z, \quad C_{n+1} = zC_n - C_{n-1} \quad (n \geq 1), \quad (23.1)$$

and $\mathcal{T}_p$ is the series (1.4). In ordinary notation $C_n(z) = 2T_n(z/2)$; the normalization $C_0 = 2$ is why the constant term of $\mathcal{T}_p$ is written separately.

Lemma 23.2 (Chebyshev identities; source 13). *For $n \geq 1$, C_n is monic of degree n and*

$$C_n(u + u^{-1}) = u^n + u^{-n}, \quad (23.2)$$

$$C_n(z) = \sum_{j=0}^{\lfloor n/2 \rfloor} (-1)^j \frac{n}{n-j} \binom{n-j}{j} z^{n-2j}. \quad (23.3)$$

All the coefficients in the second formula are integers.

Proof. The first identity follows from the initial values and the same three-term recurrence on both sides. The recurrence also proves monicity and integrality. Substituting the displayed coefficients into the recurrence and using Pascal's identity proves (23.3); the initial cases are $C_1 = z$ and $C_2 = z^2 - 2$. $\square$

Thus

$$\mathcal{T}_p(z) = 1 + pz + p^4(z^2 - 2) + p^9(z^3 - 3z) + p^{16}(z^4 - 4z^2 + 2) + p^{25}(z^5 - 5z^3 + 5z) + \cdots$$

Proposition 23.3 (Exact Taylor coefficients; source 13). *For every $\mu > 0$ write $\mathcal{T}_p = \sum_{m \geq 0} a_m z^m$. Then*

$$a_0 = 1 + 2 \sum_{j \geq 1} (-1)^j p^{4j^2}, \quad (23.4)$$

$$a_m = \sum_{j \geq 0} (-1)^j \frac{m+2j}{m+j} \binom{m+j}{j} p^{(m+2j)^2} \quad (m \geq 1), \quad (23.5)$$

and

$$v(a_0) = 0, \quad v(a_m) = m^2 \mu, \quad \text{lc}(a_m) = 1 \quad (m \geq 1). \quad (23.6)$$

No coefficient vanishes, so $\mathcal{T}_p$ is not a polynomial.

Proof. For each m collect the terms $n = m + 2j$ of (23.3). Their exponents $(m + 2j)^2 \mu$ increase strictly, so the family is strongly summable for every $\mu > 0$ (Lemma 2.3 (b) with $S = \{\mu\}$) and each exponent carries one term. For $m \geq 1$ the first coefficient is one and every later term has strictly larger valuation. The constant term follows from $C_{2j}(0) = 2(-1)^j$ and $C_{2j+1}(0) = 0$. $\square$

23.3 The exact strong domain

Lemma 23.4 (A quadratic support certificate; source 13). *Let $\mu > 0$ and $x \in \mathbb{K}$ with $x = 0$ or $v(x) \geq -N\mu$ for some $N \in \mathbb{N}$. The family*

$$(p^{n^2} x^{n-2j})_{n \geq 1, 0 \leq j \leq \lfloor n/2 \rfloor}$$

is strongly summable, and so is every family obtained from it by multiplying members by ordinary integers or omitting members.

Proof. For $x = 0$ only the members with $n - 2j = 0$ are nonzero, and their exponents $n^2\mu$ are distinct. Otherwise write

$$x = at^{-N\mu+\beta}(1+\varepsilon_x), \quad a \in k^\times, \quad \beta \geq 0, \quad \varepsilon_x = 0 \text{ or } v(\varepsilon_x) > 0.$$

The support of a member lies in $[n^2 - N(n - 2j)]\mu + (n - 2j)\beta + M(\text{supp } \varepsilon_x)$. For $n > N$ the integer $c_{n,j} = n^2 - N(n - 2j) \geq n(n - N)$ is positive and tends to infinity with n , uniformly in j . Apply Lemma 2.1 (2) to the well-ordered positive alphabet $S = \{\mu\} \cup (\{\beta\} \setminus \{0\}) \cup \text{supp } \varepsilon_x$. After the binomial expansion of $(1 + \varepsilon_x)^{n-2j}$, every support element of a member with $n > N$ is represented by a word with at least $c_{n,j}$ letters μ . A fixed exponent has only finitely many words, so the number of their letters μ is bounded, which bounds n and hence j ; each of the finitely many remaining members is one Hahn element. The members with $n \leq N$ are finitely many. The same argument shows that the union of the supports is well ordered. Multiplication by ordinary integers and omission enlarge neither supports nor multiplicities. No cofinality of $(n\mu)$ is assumed. $\square$

Theorem 23.5 (Exact strong domain; source 13). *For every $\mu > 0$, not necessarily an order unit,*

$$\text{Dom}_{\text{str}}(\mathcal{T}_p) = \mathcal{O}_{\Delta_\mu} = \{0\} \cup \{x \in \mathbb{K}^\times : v(x) \geq -N\mu \text{ for some } N \in \mathbb{N}\}, \quad (23.7)$$

the valuation ring of the coarsened valuation v_{Δ_μ} . Consequently $\mathcal{T}_p \in \mathcal{E}_\Gamma(k)$ if and only if μ is an order unit of Γ .

Proof. For $x \in \mathcal{O}_{\Delta_\mu}$, Lemma 23.4 applies to the full expansion $(p^{n^2}c_{n,j}x^{n-2j})_{n,j}$, with $c_{n,j}$ the integers of (23.3). Regrouping it by powers of x gives exactly $(a_mx^m)_{m \geq 0}$. Here and below we use the standard associativity of strong sums: if a strongly summable family is partitioned into arbitrary subfamilies, each subfamily is strongly summable, and the family of their sums is strongly summable with the same sum, because the supports of the partial sums lie in the union of the original supports and only finitely many subfamilies contain a member contributing at a fixed exponent.

If $x \notin \mathcal{O}_{\Delta_\mu}$, put $\gamma = v(x)$, so $\gamma < -N\mu$ for every N . By (23.6) the members with $m \geq 1$ have leading exponents $m^2\mu + m\gamma$, whose consecutive differences $(2m + 1)\mu + \gamma$ are negative. They strictly decrease, and Lemma 2.2 excludes strong summability.

A coset in Γ/Δ_μ is nonnegative exactly when its representative lies in Δ_μ or is positive and outside it, that is, when it is bounded below by some $-N\mu$; this identifies the valuation ring. It equals $\mathbb{K}$ exactly when every negative value is so bounded, that is, when $\Delta_\mu = \Gamma$. $\square$

When μ is an order unit there is the shorter route: for fixed γ the values $m^2\mu + m\gamma$ escape cofinally, and Lemma 2.3 (a) applies. The support proof is needed for the noncofinal boundary, where strong summability and convergence in the full valuation topology differ (Remark 2.5). When μ is not an order unit, the coefficient values of $\mathcal{T}_p$ lie in the proper convex subgroup Δ_μ , and Theorem 23.5 shows that the necessary inclusion of Proposition 20.9 is then an equality for this series.

23.4 The bilateral series, the product and the zeros

Put

$$\Psi_p(u) = \sum_{n \in \mathbb{Z}} p^{n^2} u^n. \quad (23.8)$$

At $u \in \mathbb{K}^\times$ this family is strongly summable exactly when $v(u) \in \Delta_\mu$: both tails are covered by the argument of Lemma 23.4 when $|v(u)| \leq N\mu$, and otherwise one of the sequences $n^2\mu \pm nv(u)$ strictly

decreases. Since $\Psi_p(u) = \sum_n (-1)^n (p^2)^{n(n-1)/2} (-pu)^n$, this is the Hahn–Tate theta domain [40, `tate:thm:theta`] at the nome p^2 and the argument $-pu$, printed once; source 13 describes it as the symmetric specialization of that domain. On this domain, and as identities of formal series in Section 24.3, reindexing gives

$$\Psi_p(u^{-1}) = \Psi_p(u), \quad (23.9)$$

$$\Psi_p(p^2 u) = p^{-1} u^{-1} \Psi_p(u), \quad (23.10)$$

$$\mathcal{T}_p(u + u^{-1}) = \Psi_p(u). \quad (23.11)$$

For (23.10) the exponent identity is $n^2 + 2n = (n+1)^2 - 1$. For (23.11), $x = u + u^{-1}$ has $v(x) \geq -|v(u)|$, so Lemma 23.4 applies at x ; grouping the full expansion by n gives $(p^{n^2} C_n(x))_n = (p^{n^2} (u^n + u^{-n}))_n$ by (23.2), and grouping it by powers of x gives the Taylor family. No square root or inverse cosine is used to define $\mathcal{T}_p$, and no branch is chosen.

Jacobi’s triple product, in the normalization of DLMF 20.5.9 [50], is

$$\sum_{n \in \mathbb{Z}} q^{n^2} u^n = \prod_{m \geq 1} (1 - q^{2m})(1 + q^{2m-1} u)(1 + q^{2m-1} u^{-1}).$$

It is classical, not a new theorem about theta functions. Pairing the reciprocal factors gives the following form.

Theorem 23.6 (Product and complete zero set; source 13). *Let $\mu > 0$ and $\mathcal{P}_p = \prod_{m \geq 1} (1 - p^{2m})$. For every $x \in \mathcal{O}_{\Delta_\mu}$,*

$$\mathcal{T}_p(x) = \mathcal{P}_p \prod_{m \geq 1} (1 + p^{2m-1} x + p^{4m-2}), \quad (23.12)$$

where the products are strong products, that is, strong sums of their finite-subset expansions. The scalar $\mathcal{P}_p$ is a unit of valuation zero and leading coefficient one. The zeros of $\mathcal{T}_p$ in its domain are exactly

$$x_m = -(p^{2m-1} + p^{-(2m-1)}) \quad (m = 1, 2, \dots), \quad (23.13)$$

and every zero is simple. The conclusion is already split over $k((t^\Gamma))$; no algebraic closedness is used.

Proof. First let p be a formal variable. At each p -degree both sides of the triple product have Laurent-polynomial coefficients in u , and the identity holds in $\mathbb{Q}[u, u^{-1}][[p]]$. Since $(1 + p^{2m-1} u)(1 + p^{2m-1} u^{-1}) = 1 + p^{2m-1}(u + u^{-1}) + p^{4m-2}$ and the substitution $z \mapsto u + u^{-1}$ is injective on $\mathbb{Q}[z]$ (the leading power u^d of a polynomial of degree d cannot cancel), the paired identity $1 + \sum_{n \geq 1} p^{n^2} C_n(z) = \mathcal{P}_p \prod_m (1 + p^{2m-1} z + p^{4m-2})$ holds in $\mathbb{Q}[z][[p]]$.

Now let $p = t^\mu$ and $v(x) \geq -N\mu$. For $2m-1 > N$ both $p^{2m-1}x$ and p^{4m-2} have positive valuation, and their supports, with the factor labels retained, lie in a positive support monoid generated by μ and the support of the normalized x as in Lemma 23.4; every nonempty choice from a late factor uses at least one letter μ , so a fixed exponent bounds the number and the indices of the late factors involved. Lemma 2.1 (2) therefore makes the finite-subset expansion of the tail product strongly summable, and the finitely many early factors multiply it by a finite family. The same holds for $\mathcal{P}_p$. Expand both sides of the paired identity at x into families of terms $e p^k x^j$ with $e \in \mathbb{Z}$: the Taylor side is strongly summable by Lemma 23.4, the product side by the argument just given. Grouping either family by the formal monomial $p^k z^j$ gives the same family, by the identity in $\mathbb{Q}[z][[p]]$, so the two sums agree. The tail product lies in $1 + \mathfrak{m}_v$ and is nonzero.

Each factor of (23.12) equals $p^{2m-1}(x-x_m)$, and the roots have the distinct valuations $-(2m-1)\mu$. At a fixed point all sufficiently late factors form a nonzero unit product, so a zero comes from one of finitely many early factors; there are no other zeros. For simplicity, split off the m th factor $\Pi_m(z) = 1 + p^{2m-1}z + p^{4m-2}$: the formal identity gives $\mathcal{T}_p = \Pi_m \cdot G_m$ in $\mathbb{K}[[z]]$, with G_m the image of the complementary product, which is strongly evaluable on $\mathcal{O}_{\Delta_\mu}$ by the same argument. Differentiating and evaluating at the nonzero point x_m , where Π_m vanishes, gives $(D_z \mathcal{T}_p)(x_m) = p^{2m-1}G_m(x_m)$, a nonzero strong product. $\square$

For $k = \mathbb{R}$ every zero is real and negative; for $k = \mathbb{C}$ there are no further nonreal zeros. These conclusions come from the displayed factorization, not from a claim that non-Archimedean entire functions split in general.

Valuations at the zeros. Write $\mathcal{T}'_p = D_z \mathcal{T}_p$. At $x = x_m$ the factors with $k < m$ have valuation $-2(m-k)\mu$ and leading coefficient -1 , and those with $k > m$ have valuation zero and leading coefficient one. Hence

$$v(\mathcal{T}'_p(x_m)) = (-m^2 + 3m - 1)\mu, \quad \text{lc}(\mathcal{T}'_p(x_m)) = (-1)^{m-1}, \quad (23.14)$$

which is also a direct certificate that the derivative does not vanish.

Proposition 23.7 (Exact linearization on the nearest-root ball; source 13). *If $y \neq 0$ and $v(y) > -(2m-1)\mu$, then*

$$\mathcal{T}_p(x_m + y) = \mathcal{T}'_p(x_m) y (1 + \varpi), \quad \varpi = 0 \text{ or } v(\varpi) > 0. \quad (23.15)$$

Therefore $v(\mathcal{T}_p(x_m + y)) = (-m^2 + 3m - 1)\mu + v(y)$.

Proof. For $k \neq m$ the valuation of $x_m - x_k$ is the smaller of the two root valuations and is at most $-(2m-1)\mu$, so $v(y/(x_m - x_k)) > 0$. Dividing (23.12) at $x_m + y$ by $\mathcal{T}'_p(x_m) y$ gives

$$\frac{\mathcal{T}_p(x_m + y)}{\mathcal{T}'_p(x_m) y} = \prod_{k \neq m} \left(1 + \frac{y}{x_m - x_k}\right).$$

The finitely many early factors lie in $1 + \mathfrak{m}_v$. In the tail, $1/(x_m - x_k)$ has valuation $(2k-1)\mu$; expanding after normalizing its leading monomial gives the same positive support certificate as for the original product. The strong product therefore lies in $1 + \mathfrak{m}_v$. $\square$

Newton profile. In the notation (14.1), at a scale δ with $t^{-\delta} \in \text{Dom}_{\text{str}}(\mathcal{T}_p)$, that is $\delta \leq N\mu$ for some N , the coefficient Gauss value is $\gamma_\delta = \min_{n \geq 0} (n^2\mu - n\delta)$. The minimum exists because the members with $n > N$ exceed the member $n = 0$, and consecutive differences are $(2n+1)\mu - \delta$. There is one active index n when $(2n-1)\mu < \delta < (2n+1)\mu$ (with the evident change for $n = 0$), and two, n and $n+1$, when $\delta = (2n+1)\mu$; the root x_{n+1} has exactly the valuation $-(2n+1)\mu$. The quadratic Newton profile and the linear-factor product give the same complete shell classification, without a general Newton-polygon theorem.

24 An equation of order three, and none of lower order

24.1 The cleared third-order equation

For $r \in \{1, 3, 5\}$ put

$$\mathcal{Q}_r(p^2) = \sum_{n \geq 1} \frac{n^r p^{2n}}{1 - p^{2n}} = \sum_{N \geq 1} \left(\sum_{d|N} d^r \right) p^{2N}, \quad (24.1)$$

and

$$\mathfrak{c} = \frac{1}{12} - 2\mathcal{Q}_1(p^2), \quad g_2 = \frac{1}{12} + 20\mathcal{Q}_3(p^2), \quad g_3 = -\frac{1}{216} + \frac{7}{3}\mathcal{Q}_5(p^2). \quad (24.2)$$

The powers are supported on positive multiples of 2μ , and each exponent has finitely many divisors, so the series exist for every $\mu > 0$. They begin

$$\begin{aligned} \mathfrak{c} &= \frac{1}{12} - 2p^2 - 6p^4 - 8p^6 - 14p^8 + O(p^{10}), \\ g_2 &= \frac{1}{12} + 20p^2 + 180p^4 + 560p^6 + 1460p^8 + O(p^{10}), \\ g_3 &= -\frac{1}{216} + \frac{7}{3}p^2 + 77p^4 + \frac{1708}{3}p^6 + \frac{7399}{3}p^8 + O(p^{10}), \end{aligned}$$

where $O(p^{10})$ denotes a formal series divisible by p^{10} , not a numerical error bound. Put

$$\Phi(T) = 4(\mathfrak{c} - T)^3 - g_2(\mathfrak{c} - T) - g_3. \quad (24.3)$$

For a nonzero element f of a differential field extending $\mathbb{K}(z)$ with derivation D_z , put

$$\mathcal{A}[f] = (z^2 - 4) \left(\frac{D_z^2 f}{f} - \frac{(D_z f)^2}{f^2} \right) + z \frac{D_z f}{f}, \quad (24.4)$$

$$\mathcal{H}[f] = (z^2 - 4)(f D_z^2 f - (D_z f)^2) + z f D_z f, \quad (24.5)$$

$$\mathcal{K}[f] = f D_z \mathcal{H}[f] - 2 D_z f \mathcal{H}[f], \quad \mathcal{B}[f] = \mathfrak{c} f^2 - \mathcal{H}[f], \quad (24.6)$$

so that $\mathcal{A}[f] = \mathcal{H}[f]/f^2$ and $D_z \mathcal{A}[f] = \mathcal{K}[f]/f^3$. These are rational differential expressions; no logarithm of a field element is used.

Theorem 24.1 (Cleared third-order equation; source 13). *For every $\mu > 0$ the series $f = \mathcal{T}_p$ satisfies, in $\mathbb{K}[[z]]$,*

$$(z^2 - 4)\mathcal{K}[f]^2 - 4\mathcal{B}[f]^3 + g_2\mathcal{B}[f]f^4 + g_3f^6 = 0. \quad (24.7)$$

Its left side is $P(z, f, D_z f, D_z^2 f, D_z^3 f)$ for the nonzero polynomial

$$P = (z^2 - 4)K^2 - 4B^3 + g_2BY_0^4 + g_3Y_0^6 \in \mathbb{K}[z, Y_0, Y_1, Y_2, Y_3], \quad (24.8)$$

where $H = (z^2 - 4)(Y_0Y_2 - Y_1^2) + zY_0Y_1$, $H_1 = (z^2 - 4)(Y_0Y_3 - Y_1Y_2) + 2z(Y_0Y_2 - Y_1^2) + Y_0Y_1 + z(Y_1^2 + Y_0Y_2)$ (the total derivative of H), $K = Y_0H_1 - 2Y_1H$ and $B = \mathfrak{c}Y_0^2 - H$. The polynomial P is homogeneous of degree six in $Y_0, \dots, Y_3$, and its coefficient of Y_3^2 is $(z^2 - 4)^3Y_0^4$, so no values of $\mathfrak{c}$, g_2 , g_3 make it zero. In $\mathbb{K}((z))$ the equivalent rational form is

$$(z^2 - 4)(D_z \mathcal{A}[f])^2 = \Phi(\mathcal{A}[f]). \quad (24.9)$$

Both are formal identities. The cleared one can be evaluated at every point of $\text{Dom}_{\text{str}}(\mathcal{T}_p)$, including the zeros of f and $z = \pm 2$, since evaluation is a ring homomorphism on series evaluable at a common point and no division occurs.

The proof occupies the next two subsections. The coefficient of $D_z^3 f$ in $\mathcal{K}[f]$ is $(z^2 - 4)f^2$, which gives the leading coefficient in Y_3 .

24.2 The classical elliptic identity

Source 13 supplies the normalization calculation because a sign or nome convention changes all three constants. Theta-elliptic differential identities are classical [51, 55], and Tate coordinates give another route to the same normalization [53, 40].

In this subsection only, p is an ordinary real number with $0 < p < 1$, and $\Psi(u) = \sum_{n \in \mathbb{Z}} p^{n^2} u^n$ for $u \in \mathbb{C}^\times$. Let $\psi_1 = \vartheta_u \Psi / \Psi$ and $\psi_2 = \vartheta_u \psi_1$. Differentiating the product away from its zeros gives

$$\psi_2 = -\mathcal{Z}(-pu), \quad \mathcal{Z}(w) = \sum_{n \in \mathbb{Z}} \frac{p^{2n} w}{(1 - p^{2n} w)^2}. \quad (24.10)$$

For negative n the summand can be rewritten through $w/(1-w)^2 = w^{-1}/(1-w^{-1})^2$, so this is a normally convergent meromorphic series on $\mathbb{C}^\times$, invariant under $w \mapsto p^2 w$ and $w \mapsto w^{-1}$. Put

$$\mathcal{W}(y) = \mathcal{Z}(e^y) - 2\mathcal{Q}_1(p^2) + \frac{1}{12}.$$

It is meromorphic with periods $2\pi i$ and $\log p^2$, and its poles are exactly the points of the period lattice, all double. Near $y = 0$ the summand $n = 0$ is

$$\frac{e^y}{(1 - e^y)^2} = y^{-2} - \frac{1}{12} + \frac{y^2}{240} - \frac{y^4}{6048} + O(y^6),$$

and pairing the summands n and $-n$ for $n \geq 1$ gives $2 \sum_{d \geq 1} dp^{2nd} \cosh(dy)$. Consequently

$$\mathcal{W}(y) = y^{-2} + \mathfrak{a}_2 y^2 + \mathfrak{a}_4 y^4 + O(y^6), \quad \mathfrak{a}_2 = \frac{1}{240} + \mathcal{Q}_3(p^2), \quad \mathfrak{a}_4 = -\frac{1}{6048} + \frac{1}{12} \mathcal{Q}_5(p^2). \quad (24.11)$$

The elliptic function $(\mathcal{W}')^2 - 4\mathcal{W}^3 + 20\mathfrak{a}_2 \mathcal{W} + 28\mathfrak{a}_4$ has no pole: its y^{-6} , y^{-2} and constant terms cancel by (24.11), and periodicity treats every lattice point. It is bounded on a compact period parallelogram, hence constant, and its limit at 0 is zero. As $20\mathfrak{a}_2 = g_2$ and $28\mathfrak{a}_4 = g_3$,

$$(\mathcal{W}')^2 = 4\mathcal{W}^3 - g_2 \mathcal{W} - g_3.$$

With $e^y = -pu$ one has $\vartheta_u = d/dy$ and $\mathcal{W} = \mathfrak{c} - \psi_2$, so, writing $\psi_3 = \vartheta_u \psi_2$,

$$\psi_3^2 = \Phi(\psi_2) = 4(\mathfrak{c} - \psi_2)^3 - g_2(\mathfrak{c} - \psi_2) - g_3. \quad (24.12)$$

The proof uses only the classical product, a Laurent expansion and the constancy of an elliptic function without poles.

24.3 From the complex identity to the Hahn field

Now let p be an indeterminate again, and work in the ring $\mathbb{Q}[u, u^{-1}][[p]]$ of power series in p whose coefficients are Laurent polynomials in u . The series $\Psi = \sum_n p^{n^2} u^n = 1 + p(u + u^{-1}) + \dots$ is a unit there, so ψ_1, ψ_2, ψ_3 belong to it, and $\mathfrak{c}, g_2, g_3 \in \mathbb{Q}[[p]]$. For u in the annulus $\frac{1}{2} \leq |u| \leq 2$ and $0 < p < \epsilon_0$, with ϵ_0 small, all these formal expansions converge absolutely to the complex functions of Section 24.2: $|\Psi - 1| < \frac{1}{2}$, so $1/\Psi$ is a convergent geometric series. Hence, for each such u , the power series in p obtained from $\psi_3^2 - \Phi(\psi_2)$ vanishes on $(0, \epsilon_0)$, so all its coefficients vanish there; each coefficient is a Laurent polynomial vanishing on an annulus, hence zero. Thus (24.12) is an identity in $\mathbb{Q}[u, u^{-1}][[p]]$, over $\mathbb{Q}$, and not merely for particular complex nomes.

From u to z . The map $\iota : \mathbb{Q}[z][[p]] \rightarrow \mathbb{Q}[u, u^{-1}][[p]]$, $\sum_k P_k(z)p^k \mapsto \sum_k P_k(u + u^{-1})p^k$, is an injective ring homomorphism. With $\check{z} = u - u^{-1}$,

$$\vartheta_u(u + u^{-1}) = \check{z}, \quad \vartheta_u \check{z} = u + u^{-1}, \quad \check{z}^2 = \iota(z^2 - 4), \quad \vartheta_u(\iota g) = \check{z} \iota(D_z g), \quad (24.13)$$

the last by the chain rule, coefficientwise in p . The formal series $\mathcal{T} = 1 + \sum_{n \geq 1} p^{n^2} C_n(z) \in \mathbb{Q}[z][[p]]$ has $\iota(\mathcal{T}) = \Psi$ by (23.2). From (24.13) one computes $\iota(\mathcal{H}[\mathcal{T}]) = \Psi \vartheta_u^2 \Psi - (\vartheta_u \Psi)^2 = \Psi^2 \psi_2$, then $\check{z} \iota(\mathcal{K}[\mathcal{T}]) = \Psi^3 \psi_3$ and $\iota(\mathcal{B}[\mathcal{T}]) = \Psi^2(\mathfrak{c} - \psi_2)$. Hence ι maps the left side of (24.7) for $\mathcal{T}$ to $\Psi^6(\psi_3^2 - \Phi(\psi_2)) = 0$, and injectivity gives (24.7) in $\mathbb{Q}[z][[p]]$.

From $\mathbb{Q}[[p]]$ to $\mathbb{K}$. For every $\mu > 0$ the map $s_\mu : \mathbb{Q}[[p]] \rightarrow \mathbb{K}$, $\sum_k c_k p^k \mapsto \sum_k c_k t^{k\mu}$, is an injective ring homomorphism: the support lies in the well-ordered set $\{k\mu\}$, each exponent comes from one k , and the Cauchy product is preserved. Applied coefficientwise it is a ring homomorphism $\mathbb{Q}[[p]][[z]] \rightarrow \mathbb{K}[[z]]$ commuting with D_z , and $\mathbb{Q}[z][[p]] \subseteq \mathbb{Q}[[p]][[z]]$ as subrings of $\mathbb{Q}[[p, z]]$. It maps the formal $\mathcal{T}$ to $\mathcal{T}_p$, whose coefficients are (23.4)–(23.5), and the formal constants to (24.2). This proves (24.7) in $\mathbb{K}[[z]]$ for every $\mu > 0$. Dividing by f^6 in $\mathbb{K}((z))$ gives (24.9).

Inverses are taken only in $\mathbb{Q}[u, u^{-1}][[p]]$, where Ψ is a unit, and only the cleared identity is specialized. This avoids the incorrect claim that the p -adic geometric expansion of $1/\Psi_p(u)$ converges at every Hahn argument, a point source 13 makes explicitly; in a Hahn function field the inverse is taken as a fraction.

24.4 The corner of the cleared equation

Proposition 24.2 (Corner data of the cleared equation; merge). *Let P be the polynomial (24.8) with the constants (24.2), written in the combined-monomial form (14.4). In the notation of Section 14.4, $d_P = 6$, $w_P = 0$, $\beta_P = 0$, and*

$$I_P(T) = -\Phi(0) = -(4\mathfrak{c}^3 - g_2\mathfrak{c} - g_3), \quad \chi_P(T) = 0.$$

The constant $4\mathfrak{c}^3 - g_2\mathfrak{c} - g_3$ equals $-4p^2 - 48p^4 - 144p^6 - 448p^8 - 600p^{10} + O(p^{12})$; so I_P is a nonzero constant of valuation 2μ and leading coefficient 4, while the residue corner polynomial vanishes identically.

Proof. Every monomial of P has total degree six in the Y_j , so $d_P = 6$. For the weight $w_\nu = k_\nu - \sum_j j m_{\nu j}$ of (14.5), every term of H has weight at most 0, with weight-zero part $H_0 = z^2(Y_0 Y_2 - Y_1^2) + z Y_0 Y_1$ (the terms $-4(Y_0 Y_2 - Y_1^2)$ have weight -2). The total derivative lowers the weight of each term by exactly one, so the terms of H_1 have weight at most -1 , with weight -1 part the total derivative H'_0 of H_0 . Hence every monomial of P has weight at most 0, and its weight-zero part is

$$P_0 = z^2(Y_0 H'_0 - 2Y_1 H_0)^2 - 4(\mathfrak{c} Y_0^2 - H_0)^3 + g_2(\mathfrak{c} Y_0^2 - H_0) Y_0^4 + g_3 Y_0^6.$$

The substitution $z \mapsto 1$, $Y_j \mapsto (T)_j$ is a ring homomorphism $\mathbb{K}[z, Y] \rightarrow \mathbb{K}[T]$ sending a monomial of weight 0 to the product of falling factorials in (14.6). It sends H_0 to $T(T-1) - T^2 + T = 0$ and $H'_0 = z^2(Y_0 Y_3 - Y_1 Y_2) + 2z(Y_0 Y_2 - Y_1^2) + Y_0 Y_1 + z(Y_1^2 + Y_0 Y_2)$ to

$$T(T-1)(T-2) - T^2(T-1) + 2(T(T-1) - T^2) + T + T^2 + T(T-1) = 0,$$

so P_0 goes to $-4\mathfrak{c}^3 + g_2\mathfrak{c} + g_3 = -\Phi(0)$. From $\mathfrak{c} = \frac{1}{12} - 2p^2 + O(p^4)$, $g_2 = \frac{1}{12} + 20p^2 + O(p^4)$ and $g_3 = -\frac{1}{216} + \frac{7}{3}p^2 + O(p^4)$ one finds $4\mathfrak{c}^3 = \frac{1}{432} - \frac{1}{6}p^2 + O(p^4)$ and $g_2\mathfrak{c} = \frac{1}{144} + \frac{3}{2}p^2 + O(p^4)$, so $4\mathfrak{c}^3 - g_2\mathfrak{c} - g_3 = -4p^2 + O(p^4)$; the further terms were computed with SymPy. In particular the

coefficient of the weight-zero monomial Y_0^6 in P is $-\Phi(0) \neq 0$, so $P_0 \neq 0$, $w_P = 0$, and $I_P = -\Phi(0)$. The monomial $z^6 Y_0^4 Y_3^2$ of P_0 has coefficient 1, and every coefficient of P lies in $\mathbb{Z}[\mathfrak{c}, g_2, g_3] \subseteq s_\mu(\mathbb{Q}[[p]])$, which has nonnegative valuation; so $\beta_P = 0$. Since res is a ring homomorphism on elements of nonnegative valuation and the falling-factorial products have integer coefficients, χ_P is obtained by applying res to the coefficients of I_P , and $\text{res}(-\Phi(0)) = 0$ because $v(\Phi(0)) = 2\mu > 0$. $\square$

The corner consists of 19 monomials (SymPy). The computation has a transparent form: the weight-zero part H_0 of $\mathcal{H}$ is exactly the left side of the vanishing-corner equation (16.1), whose residue corner polynomial is the identity (16.2).

Remark 24.3 (The vanishing corner; source 13 and merge). The corner criterion of Theorem 16.1 is a sufficient rigidity criterion, not a theorem for every higher-order equation, and the cleared equation necessarily lies outside it. Source 13 records the cancellation directly: on a trial monomial $f = z^N$ the top-degree part of $\mathcal{A}[f]$ vanishes, because $z^2(-N/z^2) + z(N/z) = 0$, so retaining only separate derivative degrees and ignoring their cancellation would miss the example. Proposition 24.2 makes this exact and adds three consequences. (1) The equation is consistent with Theorems 15.1 and 16.1: they require $\chi_P \neq 0$, and here $\chi_P = 0$. (2) When μ is an order unit, $\mathcal{T}_p$ satisfies no nonzero equation with $d_P \geq 1$ and $\chi_P \neq 0$ at all (and a nonzero P with $d_P = 0$ is a nonzero element of $\mathbb{K}[z]$, which no series satisfies), since by Theorem 16.1 such an equation has only polynomial strongly entire solutions; so a nonpolynomial strongly entire series can satisfy an equation whose residue corner vanishes identically without satisfying any equation to which Theorem 15.1 applies. (3) The hypothesis $\chi_P \neq 0$ of those theorems cannot be weakened to $I_P \neq 0$: here the full corner polynomial is a nonzero constant of positive valuation, while the residue corner vanishes. The containment (14.7) and Proposition 16.2, which assume $\chi_P \neq 0$, are unaffected. None of this contradicts the first-order or D-finite rigidity theorems.

24.5 The Laurent ring and the substitution $z = u + u^{-1}$

In this subsection and the next two, μ is an order unit. This supplies a common Laurent function field for all nonzero arguments; the lower bound concerns formal functions, not the algebraic independence of values in their own ambient field.

Let $\mathcal{U}_\mu$ be the set of formal Laurent series $b = \sum_{m \in \mathbb{Z}} b_m u^m$ with $b_m \in \mathbb{K}$ such that, for all $\gamma, \beta \in \Gamma$, only finitely many m with $b_m \neq 0$ satisfy $v(b_m) + m\gamma \leq \beta$. This is source 13's condition " $v(b_m) + m\gamma \rightarrow +\infty$ as $|m| \rightarrow \infty$ ", written in the form of Proposition 20.2 (iii). For each γ the weighted values $v(b_m) + m\gamma$ of the nonzero coefficients then have a least element, attained by finitely many m .

Lemma 24.4 (The Laurent ring; source 13). (a) $\mathcal{U}_\mu$ is an integral domain containing $\mathbb{K}[u, u^{-1}]$, with the product $(\sum b_m u^m)(\sum c_m u^m) = \sum_m (\sum_{m_1+m_2=m} b_{m_1} c_{m_2}) u^m$, the inner sums being strong sums.

(b) $\vartheta_u(\sum b_m u^m) = \sum m b_m u^m$ is a derivation of $\mathcal{U}_\mu$ vanishing on $\mathbb{K}$, and $S(\sum b_m u^m) = \sum b_m p^{2m} u^m$ is a ring automorphism fixing $\mathbb{K}$ and commuting with ϑ_u . Both extend to $\text{Frac} \mathcal{U}_\mu$.

(c) Ψ_p and all $\vartheta_u^j \Psi_p$ belong to $\mathcal{U}_\mu$, and $\Psi_p \neq 0$.

Proof. (a) With $\gamma = 0$ the coefficient values of b and c are bounded below and escape cofinally, so for fixed m the family $(b_{m_1} c_{m-m_1})_{m_1}$ is strongly summable by Lemma 2.3 (a). For fixed γ , if the product coefficient $d_m \neq 0$ has $v(d_m) + m\gamma \leq \beta$, some term has $(v(b_{m_1}) + m_1\gamma) + (v(c_{m_2}) + m_2\gamma) \leq \beta$,

and the lower bounds of the two weighted families leave finitely many pairs, as in the proof of Proposition 20.4; so the product lies in $\mathcal{U}_\mu$. Commutativity, associativity and distributivity follow from the associativity of strong sums applied to the families $(b_{m_1}c_{m_2}e_{m_3})$. For integrality, let w_b and w_c be the least coefficient values of nonzero b and c and $\text{in}(b) = \sum_{v(b_m)=w_b} \text{lc}(b_m)u^m \in k[u, u^{-1}]$, a nonzero Laurent polynomial. The coefficient of $t^{w_b+w_c}u^m$ in bc is the coefficient of u^m in $\text{in}(b)\text{in}(c)$, because every term has value at least $w_b + w_c$. Since $k[u, u^{-1}]$ is a domain, $bc \neq 0$.

(b) Multiplying coefficients by ordinary integers preserves the defining condition, and Leibniz's rule holds coefficientwise since $m = m_1 + m_2$. For $\mathbf{S}$, $v(b_m p^{2m}) + m\gamma = v(b_m) + m(\gamma + 2\mu)$, the inverse uses p^{-2m} , and $p^{2m} = p^{2m_1}p^{2m_2}$ with Lemma 2.1 (3) gives multiplicativity. Both operators act diagonally on u^m , so they commute. An automorphism and a derivation of a domain extend to its fraction field.

(c) If $|\gamma| \leq N\mu$ and $\beta \leq B\mu$, then $v(p^{n^2}) + n\gamma \geq (n^2 - N|n|)\mu > \beta$ for all but finitely many n ; multiplying coefficients by n^j changes no valuation. $\square$

Lemma 24.5 (Substitution of $u + u^{-1}$; merge). *For $f = \sum_n a_n z^n \in \mathcal{E}_\Gamma(k)$ and $m \in \mathbb{Z}$ the family $(a_n \binom{n}{(n-m)/2})_{n \geq |m|, n \equiv m \pmod{2}}$ is strongly summable. Write b_m for its sum and $f^\natural = \sum_m b_m u^m$. Then:*

- (a) $f^\natural \in \mathcal{U}_\mu$;
- (b) $f \mapsto f^\natural$ is a $\mathbb{K}$ -algebra homomorphism $\mathcal{E}_\Gamma(k) \rightarrow \mathcal{U}_\mu$, equal to the substitution $z \mapsto u + u^{-1}$ on $\mathbb{K}[z]$;
- (c) $\vartheta_u(f^\natural) = \check{z}(D_z f)^\natural$, where $\check{z} = u - u^{-1}$;
- (d) $\mathcal{T}_p^\natural = \Psi_p$.

Proof. Write $\beta_{n,m} = \binom{n}{(n-m)/2}$ when $n \geq |m|$ and $n \equiv m \pmod{2}$, and $\beta_{n,m} = 0$ otherwise; this is the coefficient of u^m in $(u + u^{-1})^n$. By Proposition 20.2 (iii) with $\gamma = 0$, the coefficient values of f escape cofinally, and the nonzero integers $\beta_{n,m}$ have value zero, so Lemma 2.3 (a) gives strong summability.

(a) Fix γ, β and put $\gamma' = -|\gamma|$. If $b_m \neq 0$ and $v(b_m) + m\gamma \leq \beta$, some $n \geq |m|$ with $a_n \neq 0$ has $v(a_n) + m\gamma \leq \beta$, and then $v(a_n) + n\gamma' \leq v(a_n) + m\gamma \leq \beta$ because $n\gamma' \leq |m|\gamma' \leq m\gamma$. By Proposition 20.2 (iii) at γ' only finitely many n qualify, and each serves only the m with $|m| \leq n$.

(b) Linearity over $\mathbb{K}$ holds because strong sums are linear. For multiplicativity fix m and consider the family $(a_i c_j \beta_{i,m_1} \beta_{j,m_2})$ indexed by (i, j, m_1, m_2) with $m_1 + m_2 = m$, where $g = \sum c_j z^j \in \mathcal{E}_\Gamma(k)$. The double family $(a_i c_j)$ is strongly summable, as in the proof of Lemma 14.2; each pair (i, j) carries finitely many (m_1, m_2) , and nonzero integer factors change neither supports nor multiplicities. Grouping by (m_1, m_2) and then by m_1 gives the coefficient of u^m in $f^\natural g^\natural$. Grouping by (i, j) gives $a_i c_j \beta_{i+j,m}$, because $(u + u^{-1})^i (u + u^{-1})^j = (u + u^{-1})^{i+j}$, and grouping those by $n = i + j$ gives $(\sum_{i+j=n} a_i c_j) \beta_{n,m}$, whose sum is the coefficient of u^m in $(fg)^\natural$. By associativity of strong sums the two totals agree. On polynomials all families are finite.

(c) Since $\vartheta_u(u + u^{-1})^n = n(u + u^{-1})^{n-1}(u - u^{-1})$, one has $m\beta_{n,m} = n(\beta_{n-1,m-1} - \beta_{n-1,m+1})$. The coefficient of u^m in $\vartheta_u(f^\natural)$ is $mb_m = \sum_n a_n m\beta_{n,m}$, and that of $\check{z}(D_z f)^\natural$ is $\sum_n n a_n (\beta_{n-1,m-1} - \beta_{n-1,m+1})$; they agree termwise, and $D_z f \in \mathcal{E}_\Gamma(k)$ by Proposition 20.4.

(d) With $c_{N,j}$ the integers of (23.3), the coefficient of u^m in $\mathcal{T}_p^\natural$ is the sum of the family $(p^{N^2} c_{N,j} \beta_{N-2j,m})_{N,j}$, grouped by $n = N - 2j$ (the constant term included with $N = 0$). Its exponents $N^2\mu$ each come from finitely many j , so it is strongly summable. Grouping instead by

N gives p^{N^2} times the coefficient of u^m in $C_N(u + u^{-1}) = u^N + u^{-N}$, which is nonzero only for $N = |m|$. The sum is p^{m^2} . $\square$

Injectivity of $f \mapsto f^\natural$ is not needed. By (c) and Proposition 20.4, $(D_z^r f)^\natural = (\check{z}^{-1} \vartheta_u)^r f^\natural$ in $\text{Frac} \mathcal{U}_\mu$, which equals $\check{z}^{-r} \vartheta_u^r f^\natural$ plus a combination of lower $\vartheta_u^j f^\natural$ with coefficients in $\mathbb{Q}(u)$. For $\mathcal{T}_p$,

$$\Psi_p = \mathcal{T}_p^\natural, \quad \vartheta_u \Psi_p = \check{z} (D_z \mathcal{T}_p)^\natural, \quad \vartheta_u^2 \Psi_p = \check{z}^2 (D_z^2 \mathcal{T}_p)^\natural + (u + u^{-1}) (D_z \mathcal{T}_p)^\natural. \quad (24.14)$$

24.6 Three independent theta quantities

Lemma 24.6 (Three rational-function obstructions; source 13). *Let E be a field of characteristic zero and $a \in E^\times$ not a root of unity, and let S_a be the automorphism of $E(u)$ with $S_a u = au$ fixing E .*

- (a) *If $S_a r = r$, then $r \in E$.*
- (b) *For $d \in E^\times$ there is no $r \in E(u)$ with $S_a r - r = d$.*
- (c) *On $E(u, V)$, with V transcendental, extend S_a by $S_a V = V - 1$. Then S_a preserves the u -adic order ord_u of every nonzero element.*

Proof. Expand a rational function as a formal Laurent series at $u = 0$. For (a), each coefficient of degree $n \neq 0$ is multiplied by $a^n \neq 1$, so invariance leaves only degree zero. For (b), the coefficient of u^0 in $r(au) - r(u)$ is zero, also when r has a pole at 0. For (c), a lowest nonzero coefficient in $E(V)((u))$ is sent to the nonzero coefficient obtained by $V \mapsto V - 1$, multiplied by a nonzero power of a ; its degree is unchanged. $\square$

In $\text{Frac} \mathcal{U}_\mu$ put $\Psi = \Psi_p$, $\psi_1 = \vartheta_u \Psi / \Psi$, $\psi_2 = \vartheta_u \psi_1$ and $\psi_3 = \vartheta_u \psi_2$; the field $\mathbb{K}(u)$ lies in it and is preserved by S . By (23.10),

$$S\Psi = p^{-1}u^{-1}\Psi, \quad S\psi_1 = \psi_1 - 1, \quad S\psi_2 = \psi_2. \quad (24.15)$$

Lemma 24.7 (Transfer, and a global certificate for $\psi_3 \neq 0$; merge). (a) *For every polynomial Π in four variables with coefficients in $\mathbb{Q}[[p]]$ and every $m \in \mathbb{Z}$, the coefficient of u^m in*

$$\Pi(\Psi, \vartheta_u \Psi, \vartheta_u^2 \Psi, \vartheta_u^3 \Psi),$$

computed in $\mathcal{U}_\mu$ with the coefficients of Π mapped by s_μ , is the image under s_μ of the same coefficient computed in $\mathbb{Q}[u, u^{-1}][[p]]$. In particular $\psi_3^2 = \Phi(\psi_2)$ in $\text{Frac} \mathcal{U}_\mu$.

- (b) *In $\mathcal{U}_\mu$,*

$$\Psi^3 \psi_3 = \mathcal{N} := \Psi^2 \vartheta_u^3 \Psi - 3\Psi \vartheta_u \Psi \vartheta_u^2 \Psi + 2(\vartheta_u \Psi)^3,$$

and the coefficient of u in $\mathcal{N}$ has valuation μ and leading coefficient 1. In particular $\psi_3 \neq 0$.

Proof. (a) The four series $\vartheta_u^j \Psi = \sum_n n^j p^{n^2} u^n$ lie in $\mathcal{U}_\mu$ (Lemma 24.4). By associativity of strong sums, the coefficient of u^m on the $\mathcal{U}_\mu$ side is the strong sum of the family of terms $e t^{(k+n_1^2+\dots+n_r^2)\mu}$, $e \in \mathbb{Z}$, indexed by a monomial of Π of degree r , a term ep^k of its coefficient, and a tuple $(n_1, \dots, n_r)$ with $n_1 + \dots + n_r = m$. The exponent $K\mu$ is carried by the finitely many members with $k + \sum n_i^2 = K$, whose integer coefficients add up to the coefficient of $u^m p^K$ on the formal side. This is the assertion. Applied to the cleared form of (24.12), $\Psi^6(\psi_3^2 - \Phi(\psi_2))$, which is a polynomial in the four series

with coefficients in $\mathbb{Z}[\mathfrak{c}, g_2, g_3]$ and vanishes in $\mathbb{Q}[u, u^{-1}][[p]]$ (Section 24.3), it gives the identity in $\text{Frac}\mathcal{U}_\mu$, because $\Psi \neq 0$ there.

(b) The identity is the quotient rule applied twice. By (a), the coefficient of u in $\mathcal{N}$ is the image under the injective ring homomorphism s_μ of the coefficient of u in the same expression computed in $\mathbb{Q}[u, u^{-1}][[p]]$. There $\Psi = 1 + p(u + u^{-1}) + O(p^4)$, $\vartheta_u \Psi = \vartheta_u^3 \Psi = p(u - u^{-1}) + O(p^4)$ and $\vartheta_u^2 \Psi = p(u + u^{-1}) + O(p^4)$, so

$$\mathcal{N} = p(u - u^{-1}) + O(p^2).$$

The coefficient of u is $p + O(p^2)$; SymPy gives $p - 8p^3 + 20p^5 - 70p^9 + 64p^{11} + O(p^{12})$. Its image $t^\mu + \dots$ is nonzero, so $\mathcal{N} \neq 0$, and since $\mathcal{U}_\mu$ is a domain and $\Psi \neq 0$, $\psi_3 = \mathcal{N}/\Psi^3 \neq 0$. $\square$

Source 13's proof of this step expanded ψ_1 at the simple zero $u_0 = -p$ of Ψ as $u_0/(u - u_0) + O(1)$ and read off $\psi_3 = 2u_0^3/(u - u_0)^3 + O((u - u_0)^{-2})$; the lemma replaces that local argument, which used expansions in $\text{Frac}\mathcal{U}_\mu$ that were not defined. Neither route uses a transcendence theorem for complex theta values.

Theorem 24.8 (Three independent theta quantities; source 13). *Let μ be an order unit. The elements Ψ, ψ_1, ψ_2 of $\text{Frac}\mathcal{U}_\mu$ are algebraically independent over $\mathbb{K}(u)$. Equivalently, $\Psi, \vartheta_u \Psi, \vartheta_u^2 \Psi$ are algebraically independent over $\mathbb{K}(u)$.*

Proof. ψ_2 is transcendental over $\mathbb{K}(u)$. Otherwise let $\pi(T)$ be its monic minimal polynomial. Since $S\psi_2 = \psi_2$ and S preserves $\mathbb{K}(u)$, applying S to the coefficients gives the same monic minimal polynomial, so by Lemma 24.6 (a), with $E = \mathbb{K}$ and $a = p^2$, all its coefficients lie in $\mathbb{K}$. Applying ϑ_u to $\pi(\psi_2) = 0$ gives $\pi'(\psi_2)\psi_3 = 0$. By separability in characteristic zero $\pi'(\psi_2) \neq 0$, contradicting Lemma 24.7.

ψ_1 is transcendental over $\mathbb{K}(u, \psi_2)$. This field is $E(u)$ with $E = \mathbb{K}(\psi_2)$ fixed by S . Suppose the monic minimal polynomial of ψ_1 over it has degree $d \geq 1$ and begins $T^d + bT^{d-1} + \dots$. Applying S and then replacing T by $T - 1$ gives another monic polynomial vanishing at ψ_1 , because $S\psi_1 = \psi_1 - 1$; its next coefficient is $Sb - d$. Uniqueness gives $Sb - b = d$, contradicting Lemma 24.6 (b).

Ψ is transcendental over $\mathbb{K}(u, \psi_2, \psi_1)$. Let its hypothetical monic minimal polynomial have degree d and constant coefficient b_0 , nonzero because $\Psi \neq 0$. Applying S , substituting $p^{-1}u^{-1}T$ for T and dividing by $(p^{-1}u^{-1})^d$ gives another monic polynomial vanishing at Ψ , so $Sb_0 = (p^{-1}u^{-1})^d b_0$. By the first two steps the field is the rational function field $\mathbb{K}(\psi_2)(u, \psi_1)$, on which S acts as in Lemma 24.6 (c) with $V = \psi_1$. The left side has the u -adic order of b_0 , the right side that order minus d , which is impossible.

Finally $\vartheta_u \Psi = \psi_1 \Psi$ and $\vartheta_u^2 \Psi = (\psi_2 + \psi_1^2) \Psi$, and these formulas and their rational inverses identify the fields generated by the two triples. $\square$

Corollary 24.9 (Exact minimal order; source 13). *Let μ be an order unit. Then $\mathcal{T}_p, D_z \mathcal{T}_p$ and $D_z^2 \mathcal{T}_p$ are algebraically independent over $\mathbb{K}(z)$, and the least order of a nonzero algebraic differential equation for $\mathcal{T}_p$ over $\mathbb{K}(z)$ is three.*

Proof. Suppose that a nonzero polynomial P in Y_0, Y_1, Y_2 with coefficients in $\mathbb{K}[z]$, obtained after clearing denominators, satisfies $P(z, \mathcal{T}_p, D_z \mathcal{T}_p, D_z^2 \mathcal{T}_p) = 0$. Apply Lemma 24.5 (b): the image of this identity in $\mathcal{U}_\mu$ is $P^\natural(\Psi, (D_z \mathcal{T}_p)^\natural, (D_z^2 \mathcal{T}_p)^\natural) = 0$, where $P^\natural$ has the coefficients of P with $u + u^{-1}$ substituted for z . By (24.14) the three u -jets are obtained from $\Psi, \vartheta_u \Psi, \vartheta_u^2 \Psi$ by a triangular linear change with diagonal $1, \tilde{z}^{-1}, \tilde{z}^{-2}$, invertible over $\mathbb{K}(u)$ because $\tilde{z} \neq 0$. The polynomial $P^\natural$ is nonzero,

because $z \mapsto u + u^{-1}$ is injective on $\mathbb{K}[z]$, so this is a nonzero relation among $\Psi, \vartheta_u \Psi, \vartheta_u^2 \Psi$ over $\mathbb{K}(u)$, contradicting Theorem 24.8. This proves the lower bound; Theorem 24.1 gives the upper bound. $\square$

24.7 The differential field and non-D-finiteness

Let μ be an order unit and write, in $\mathbb{K}((z))$,

$$f = \mathcal{T}_p, \quad \mathcal{L} = D_z f / f, \quad \mathcal{A} = \mathcal{A}[f], \quad \mathcal{A}_1 = D_z \mathcal{A}.$$

The first three are algebraically independent over $\mathbb{K}(z)$, being an invertible rational change of $f, D_z f, D_z^2 f$.

Theorem 24.10 (Differential-field presentation; source 13). *Let μ be an order unit. The differential field generated by $\mathcal{T}_p$ over $\mathbb{K}(z)$ is*

$$\mathbb{K}(z)\langle \mathcal{T}_p \rangle = \mathbb{K}(z)(f, \mathcal{L}, \mathcal{A}, \mathcal{A}_1), \quad (z^2 - 4)\mathcal{A}_1^2 = \Phi(\mathcal{A}). \quad (24.16)$$

It has transcendence degree three over $\mathbb{K}(z)$ and degree two over the rational function field $\mathbb{K}(z)(f, \mathcal{L}, \mathcal{A})$. Its derivation is

$$D_z f = \mathcal{L}f, \quad D_z \mathcal{L} = \frac{\mathcal{A} - z\mathcal{L}}{z^2 - 4}, \quad (24.17)$$

$$D_z \mathcal{A} = \mathcal{A}_1, \quad D_z \mathcal{A}_1 = \frac{-12(\mathfrak{c} - \mathcal{A})^2 + g_2 - 2z\mathcal{A}_1}{2(z^2 - 4)}. \quad (24.18)$$

Consequently the minimal polynomial of $D_z^3 \mathcal{T}_p$ over $\mathbb{K}(z)(\mathcal{T}_p, D_z \mathcal{T}_p, D_z^2 \mathcal{T}_p)$ has degree two, and (24.7) is that minimal relation up to a nonzero factor.

Proof. Regard $\Phi(\mathcal{A})/(z^2 - 4)$ as a rational function of the transcendental $\mathcal{A}$ over $\mathbb{K}(z)(f, \mathcal{L})$. Its order at $\mathcal{A} = \infty$ is -3 , an odd integer, so it is not a square, and adjoining $\mathcal{A}_1$ is a quadratic extension. In particular $\mathcal{A}_1 \neq 0$, as also follows from $\Phi \neq 0$ and the transcendence of $\mathcal{A}$.

The formulas for $D_z f$ and $D_z \mathcal{L}$ follow from the definitions, since $\mathcal{A} = (z^2 - 4)D_z \mathcal{L} + z\mathcal{L}$. Differentiate $(z^2 - 4)\mathcal{A}_1^2 = \Phi(\mathcal{A})$ and use $D_z \mathcal{A} = \mathcal{A}_1$ and $\Phi'(T) = -12(\mathfrak{c} - T)^2 + g_2$; cancelling $\mathcal{A}_1 \neq 0$ gives (24.18). So all derivatives of the four generators remain in the field, and conversely the generators are rational in $f, D_z f, D_z^2 f, D_z^3 f$; the field is the differential field generated by f . Finally $\mathcal{A}_1 = \mathcal{K}[f]/f^3$ is affine linear in $D_z^3 f$ with nonzero leading coefficient $(z^2 - 4)/f$, so the degree-two assertion for $\mathcal{A}_1$ is the same assertion for $D_z^3 f$, and clearing powers of f introduces only a nonzero factor. $\square$

The poles at $z = \pm 2$ of this rational system describe a coordinate chart, not poles of the entire f ; the globally meaningful equation is the cleared identity (24.7). The cubic in the elliptic coordinate $\mathfrak{c} - \mathcal{A}$ accounts for the algebraic part of the differential field, while $\mathcal{L}$ and f carry the logarithmic and multiplicative theta information.

Proposition 24.11 (Non-D-finiteness; source 13). *Let μ be an order unit. Then $\mathcal{T}_p$ satisfies no nonzero linear differential equation over $\mathbb{K}(z)$.*

Proof. Minimal algebraic order three does not by itself exclude a linear equation of higher order, so a separate argument is needed. In the u -coordinate, (24.12) holds in $\text{Frac } \mathcal{U}_\mu$, $\psi_3^2 = \Phi(\psi_2)$, and, after differentiation and cancellation of $\psi_3 \neq 0$, $\vartheta_u \psi_3 = \frac{1}{2}\Phi'(\psi_2)$ (Lemma 24.7). Hence $\mathbb{K}(\psi_2, \psi_3)$ is closed

under ϑ_u and algebraic over $\mathbb{K}(\psi_2)$, and by Theorem 24.8 ψ_1 is transcendental over $\mathbb{K}(u, \psi_2, \psi_3)$. Inductively,

$$\frac{\vartheta_u^n \Psi}{\Psi} = \psi_1^n + \text{a polynomial of degree less than } n \text{ in } \psi_1 \text{ over } \mathbb{K}(\psi_2, \psi_3) :$$

apply ϑ_u , add multiplication by ψ_1 , and use $\vartheta_u \psi_1 = \psi_2$. A nontrivial $\mathbb{K}(u)$ -linear combination of the $\vartheta_u^n \Psi$, divided by Ψ , has a nonzero highest power of the transcendental ψ_1 , so it does not vanish. A nonzero linear operator $\sum_{r \leq R} \rho_r(z) D_z^r$ with $\rho_R \neq 0$ becomes, through Lemma 24.5 and the triangular formula after it, a $\mathbb{K}(u)$ -linear combination of $\vartheta_u^r \Psi$, $r \leq R$, whose coefficient of $\vartheta_u^R \Psi$ is $\rho_R(u + u^{-1}) \tilde{z}^{-R} \neq 0$. $\square$

For $k = \mathbb{C}$ the proposition is also a case of Theorem A, since $\mathcal{T}_p$ is then a nonpolynomial element of $\mathcal{E}_\Gamma$; the argument above works for every k and uses no escape chain.

25 The exact boundary, surreal consequences and scope

25.1 Proof of Theorem H, and Theorems E and G

Proof of Theorem H. (ii) $\Rightarrow$ (i) is the contrapositive of Theorem E (a) (sources 11 and 12; source 13 proves it again, Remark 25.1). For (i) $\Rightarrow$ (iii) let μ be an order unit and $p = t^\mu$. Theorem 23.5 gives $\mathcal{T}_p \in \mathcal{E}_\Gamma(k)$ and Proposition 23.3 nonpolynomiality; Theorem 24.1, Corollary 24.9 and Theorem 24.10 give an equation of order three, algebraic independence of $\mathcal{T}_p, D_z \mathcal{T}_p, D_z^2 \mathcal{T}_p$ and degree two of $D_z^3 \mathcal{T}_p$ over the field they generate. (iii) $\Rightarrow$ (ii) is immediate. Non-D-finiteness is Proposition 24.11. Finally $\mathcal{T}_p$ is not in $\mathbb{K}(z)$, since it is transcendental over $\mathbb{K}(z)$. $\square$

Remark 25.1 (Source 13's proof of the no-order-unit direction). Section 7 of source 13 proves (ii) $\Rightarrow$ (i) self-contained, so that its classification does not rest on a citation. It is the argument of Theorem E (a) and is printed once there. Source 13 writes a relation with the Euler operator $\vartheta = zD_z$, which is a triangular rational change of the ordinary derivatives, chooses it of least total degree in the jets, and linearizes along the solution: if $\pi_j = \partial P / \partial Y_j$ evaluated at $(f, \vartheta f, \dots)$, θ' is the least z -order of a nonzero π_j and $I(T) = \sum_j [z^{\theta'}] \pi_j T^j$, then, because ϑ does not lower z -order, $I(n)a_n = -[z^{n+\theta'}]P(z, f_{<n}, \vartheta f_{<n}, \dots)$ for $n > \theta'$. With the Euler operator the linearization multiplier is a polynomial in the index. Its Lemmas 7.1 and 7.2 are Theorem 21.2 (a) and Lemmas 20.6–20.7, and its Theorem 7.3, via the exterior obstruction of Proposition 20.9, is Theorem E (a). Source 13 credits this direction to the coefficient-field part and claims no new finite-generation principle.

Proof of Theorem G (iv). (i) $\Leftrightarrow$ (iv) is (i) $\Leftrightarrow$ (ii) of Theorem H; with the proof of Theorem G given in Section 21.4, all four conditions are equivalent. $\square$

Corollary 25.2 (Theorem E is sharp; merge). *For Γ as in Convention 1.1:*

- (a) every nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ is differentially transcendental over $\mathbb{K}(z)$ if and only if Γ has no order unit;
- (b) the elements of $\text{Frac } \mathcal{E}_\Gamma(k)$ that are differentially algebraic over $\mathbb{K}(z)$ are exactly those of $\mathbb{K}(z)$ if and only if Γ has no order unit.

If Γ has an order unit μ , then for $f = \mathcal{T}_p$ and every $q \in \mathbb{K}^\times$ of infinite multiplicative order, including $v(q) = 0$, the family of Theorem G (iii) is algebraically dependent over $\mathbb{K}(z)$.

Proof. (a) is Theorem H, (i) $\Leftrightarrow$ (ii). For (b), Theorem E(c) gives the equality without an order unit; with one, $\mathcal{T}_p \in \mathcal{E}_\Gamma(k) \subseteq \text{Frac } \mathcal{E}_\Gamma(k)$ is differentially algebraic and not in $\mathbb{K}(z)$. The last assertion holds because the subfamily with $i = 0$ is already dependent (Theorem 24.1). $\square$

25.2 Consistency with the rest of the report

The new part contradicts no earlier statement; the points of contact are these.

- *Theorem E.* Without an order unit $\mathcal{T}_p$ is not entire: $\text{Dom}_{\text{str}}(\mathcal{T}_p) = \mathcal{O}_{\Delta_\mu} \neq \mathbb{K}$ (Theorem 23.5). Theorem E(a) is thus not contradicted, and by Corollary 25.2 its hypothesis is necessary.
- *Theorem D and the corner criterion.* The minimal order is three, not one; and $\chi_P = 0$ for the cleared equation, as Theorem 16.1 requires of an equation with a nonpolynomial entire solution, while $I_P \neq 0$ (Proposition 24.2, Remark 24.3).
- *Theorem F.* The coefficient values $m^2\mu$ span $\mathbb{Q}\mu$, so $\text{cvr}(\mathcal{T}_p) = 1$ and Theorem F does not apply; it is a sufficient condition only (Section 22.4). Theorem 21.20 says nothing about series whose coefficient values span $\mathbb{Q}\mu$, and $\mathcal{T}_p$ is such a series with an algebraic differential relation.
- *Theorem G.* $\mathcal{T}_p$ is a second witness for (ii) and (iii), and a stronger one for (iii): its jets are already dependent at $i = 0$, so (iii) holds for every q of infinite order, including $v(q) = 0$. Its coefficients lie in a field finitely generated over k , by Theorem 21.2(a) with $k_0 = k(\mathfrak{c}, g_2, g_3)$, as (ii) requires.
- *Theorem 21.11.* The coefficient values lie in $\mathbb{Z}\mu$, whose smallest containing convex subgroup Δ_μ is all of Γ exactly when μ is an order unit; the descent theorem needs a proper convex subgroup and does not apply.
- *The partial theta series.* Θ_p and G (Proposition 21.19, Corollary 17.1) are untouched: source 13 explains why a bilateral descent is used instead, since the dilation equation of Θ_p is not an algebraic differential equation. Whether Θ_p or G is differentially algebraic stays open.
- *Question 19.4.* It concerns *linear* mixed equations at unit valuation. $\mathcal{T}_p$ is not D-finite, and source 13 says nothing about linear mixed equations, so the question stayed open after source 13; source 18 has since answered it positively (Corollary 29.6), and the nonlinear analogue is settled by Corollary 25.2.
- *Question 19.8.* $\mathcal{T}_p$ is differentially algebraic and strongly entire, with a finitely generated coefficient field whose values $m^2\mu$ are cofinal and grow faster than any linear bound. With an order unit, therefore, no property shared by the coefficient fields of all differentially algebraic series can force a linear bound. The question asks for additional properties and stays open.
- *The Hahn–Tate report.* Ψ_p is that report’s theta series at the nome p^2 and the argument $-pu$; its domain and product (`tate:thm:theta`, `tate:prop:theta-product`, the latter for $0 \leq v(u) < v(q)$) are specialized here, not claimed anew [40].

25.3 Surreal, surcomplex and omnific consequences

Through (2.4) take $\Gamma = \mathbb{R}$ and $\mu = 1$. Then $\mathcal{T}_t$ is an actual nonpolynomial strongly entire function on the set-sized surcomplex field $\mathbb{C}((\omega^{-\mathbb{R}})) \subseteq \mathbf{No}[i]$, with coefficients in the smaller field $\mathbb{Q}((\omega^{-1}))$; the same holds on $\mathbb{C}((\omega^{-\mathbb{Z}}))$ and $\mathbb{C}((\omega^{-\mathbb{Q}}))$, and algebraic closedness of the workspace is not needed.

The derivative D_z still kills the coefficients, including $p = \omega^{-1}$, so this is a statement about functions whose values and coefficients are surreal or surcomplex, not about the scalar derivation of **No**.

Not entire on the class. With $p = \omega^{-1}$ the same coefficient formula can be evaluated at every surreal or surcomplex point whose leading exponent is bounded above by an ordinary integer; the support proof is pointwise and uses only set-sized supports. It fails at $x = \omega^\omega$: the leading exponent of $a_m x^m$ in Conway's convention is $m\omega - m^2$, and these increase strictly, so they are not the decreasing well-ordered support of a strong surreal sum. The example is therefore not strongly entire on **No** or **No** $[i]$, as for F in Section 11.2. More generally, enlarging Γ to Γ' while keeping $p = t^\mu$ changes the domain exactly to $\{x : v(x) \geq -N\mu \text{ for some } N\}$ in the larger field: an extension can turn an order unit into a non-order-unit and destroy entireness without changing a coefficient or the equation.

Omnific integers. The omnific integers are $\mathbf{Oz} = \mathbb{Z} \oplus \Pi$, where Π consists of the surreals with purely positive Conway exponents [41, `odg:def:rings`]: a nonzero term with negative exponent excludes membership. $\mathbf{Oz}$ is discretely ordered with 1 the least positive element, and a surreal that is not an omnific integer lies between two consecutive ones, its omnific floor and ceiling; for the numbers below this is read off their normal forms.

Corollary 25.3 (Zeros next to omnific integers; source 13). *For $p = \omega^{-1}$ the zeros of $\mathcal{T}_p$ in its strong domain are $x_m = -\omega^{2m-1} - \omega^{-(2m-1)}$, $m \geq 1$. None is an omnific integer, and*

$$\lfloor x_m \rfloor_{\mathbf{Oz}} = -\omega^{2m-1} - 1, \quad \lceil x_m \rceil_{\mathbf{Oz}} = -\omega^{2m-1}. \quad (25.1)$$

So $\mathcal{T}_{\omega^{-1}}(x) = 0$ has no omnific-integer solution in its strong domain, although it has infinitely many simple real surreal solutions there.

Proof. The zeros are those of Theorem 23.6. The last summand of x_m has negative exponent and nonzero coefficient, so $x_m \notin \mathbf{Oz}$. Since $0 < \omega^{-(2m-1)} < 1$, x_m lies strictly between the two consecutive omnific integers of (25.1), and there is no other zero. $\square$

This concerns a strongly evaluated infinite-series equation, not a polynomial Diophantine statement: the differential equation of $\mathcal{T}_p$ is polynomial in its jets, but $\mathcal{T}_p(x) = 0$ is not a polynomial equation in x .

Rounding can magnify an infinitesimal error. Let $x_m^+ = -\omega^{2m-1}$ and $x_m^- = x_m^+ - 1$, the ceiling and floor of x_m . Then $x_m^+ - x_m = \omega^{-(2m-1)}$ is infinitesimal and $x_m^- - x_m = -1 + \omega^{-(2m-1)}$ is finite with leading coefficient -1 , and both satisfy the hypothesis of Proposition 23.7. With the valuation normalized by $v(\omega^{-1}) = 1$,

$$v(\mathcal{T}_p'(x_m)) = -m^2 + 3m - 1, \quad \text{lc}(\mathcal{T}_p'(x_m)) = (-1)^{m-1}, \quad (25.2)$$

$$v(\mathcal{T}_p(x_m^+)) = -m^2 + 5m - 2, \quad \text{lc}(\mathcal{T}_p(x_m^+)) = (-1)^{m-1}, \quad (25.3)$$

$$v(\mathcal{T}_p(x_m^-)) = -m^2 + 3m - 1, \quad \text{lc}(\mathcal{T}_p(x_m^-)) = (-1)^m. \quad (25.4)$$

m	$v(\mathcal{T}'_p(x_m))$	$v(\mathcal{T}_p(x_m^+))$	$v(x_m^+ - x_m)$
1	1	2	1
2	1	4	3
3	-1	4	5
4	-5	2	7
5	-11	-2	9
6	-19	-8	11

For every $m \geq 5$, $\mathcal{T}_p(x_m^+)$ is infinite although x_m^+ differs from a genuine zero by an infinitesimal. This violates no local continuity: the derivative at these increasingly large roots is itself large. The exact local formula records both scales; an argument based only on the small input difference would lose the dominant effect.

25.4 Examples

Example 25.4 (The simplest discrete field; source 13). For $\mathbb{K} = \mathbb{Q}((t))$ and $p = t$, the series $1 + tz + t^4(z^2 - 2) + t^9(z^3 - 3z) + \cdots$ is strongly entire, nonpolynomial, differentially algebraic of minimal order three and not D-finite. It needs no uncountable constant field, no real closure and no divisible exponents.

Example 25.5 (One group, two scales; source 13). Let $\Gamma = \mathbb{Z} \oplus_{\text{lex}} \mathbb{Z}$ with the first coordinate dominant. The scale $\mu = (1, 0)$ gives a strongly entire witness; $\mu = (0, 1)$ does not: its strong domain is the valuation ring for the projection to the first coordinate, and $t^{(-1, 0)}$ is excluded. The existence classification quantifies over the scale; it does not say that every positive μ works.

Example 25.6 (Countable cofinality without an order unit; source 13). Order $\Gamma = \bigoplus_{j \geq 0} \mathbb{Z}e_j$ by the largest nonzero coordinate. The sequence (e_j) is cofinal, but every fixed $\mu > 0$, and every ordinary multiple of it, is dominated by some e_N . By Theorem H, or Theorem E(a), no nonpolynomial strongly entire series is differentially algebraic. This does not exclude nonpolynomial strongly entire series; it says that any such series has no algebraic differential relation over the full coefficient field. The group is the integer analogue of (11.1), over which the series F of (11.2) is strongly entire.

Corollary 25.7 (Stability under affine changes; source 13). *Let μ be an order unit, $a \in \mathbb{K}^\times$ and $b \in \mathbb{K}$. Then $g(z) = \mathcal{T}_p(az + b)$ is strongly entire and has minimal order three, and its zeros are exactly the $(x_m - b)/a$, all simple.*

Proof. For a fixed input, $v(ax + b)$ is bounded below by a multiple of $-\mu$, and the quadratic support bound, expanded binomially, justifies the Taylor coefficients of g and their evaluation. By the regrouping argument of Lemma 24.5 (b), $f \mapsto f(az + b)$ is a $\mathbb{K}$ -algebra endomorphism of $\mathcal{E}_\Gamma(k)$ with inverse $f \mapsto f((z - b)/a)$, and $D_z^j(f(az + b)) = a^j(D_z^j f)(az + b)$. A nonzero affine substitution is an automorphism of $\mathbb{K}(z)$ and scales the successive derivatives by nonzero powers of a , so it preserves the algebraic independence of the first three jets and the order of an equation. The zeros follow by substitution. $\square$

There are therefore many explicit entire nonlinear solutions, not one isolated series. The argument classifies neither all differentially algebraic entire series nor the possible minimal orders. (For $\Gamma = \mathbb{R}$ the minimal orders are unbounded, by source 16's Theorem L (b).) Order two, which source 13 left open for other series, is excluded for every series by Theorem I (source 14); the classification is Question 19.1.

25.5 Predecessors, repository comparison and priority

Literature. Source 13 consulted the NIST DLMF product and elliptic sections [50, 51]; the Tate-curve lecture notes hosted by M. Woodbury [53], whose §16 (printed page 32, the Tate equation and invariant differential) it inspected as a PDF image and whose normalization agrees with the constants (24.2), the hosting attribution not being a claim that Woodbury wrote every part; the publisher abstract and bibliography of Sebbar’s paper [55], whose abstract explicitly discusses classical third-order theta differential relations, without its full text; the publication record of Ohyama’s paper [54], not used in place of a proof; and publication records of Mantova–Matusinski [17] for normal forms. Several broad searches returned irrelevant results and are not evidence of absence; no exhaustive MathSciNet or zbMATH review, survey of book treatments or citation-network review was made. In particular no claim is made that Sebbar or Ohyama lacks a related minimal-order theorem.

Repository comparison at the pin. Source 13 read the repository at 4cdeaec through a connector: the root and documentation inventories, this report’s README, its nonlinear corner criterion and limitations, the vanishing-corner example, the first-order limitation, its attribution and dependency sections, and the question and status passage of Question 19.1 (source lines 4135–4202 of the article; its audit file gives the read range as about 4125–4290); and the README of the Hahn–Tate report, not its full article. Its statements were checked for this merge and are accurate at the pin: Question 19.1 asked for existence with an order unit, for rationality of differentially algebraic elements of $\text{Frac } \mathcal{E}_T$, and whether the absence of an order unit is necessary; its status settled the case without an order unit and gave no counterexample with one; and the Hahn–Tate README contains the bilateral theta domain on U_q . A repository content search that returned no matches was followed by direct file reads and not used as proof of absence. This was a targeted comparison, not an audit of every file, archived manuscript, branch or commit. Since the pin the statements of this report that called the order-unit case open have been updated, and Question 19.1 has been re-scoped, as recorded in Section 19.

Priority. The triple product, the Chebyshev identity and the theta–elliptic identity are classical, and no novelty is claimed for a third-order theta relation; the counterexample is a classical function, not a new theta identity. The no-order-unit direction is Theorem E (a), and the bilateral theta domain and uniformization belong to the Hahn–Tate report. The proposed contribution of source 13 is the support-controlled descent to an entire series, its exact domain at every scale, the existence classification of Theorem H, the minimal-order certificate with the differential field and non-D-finiteness, and the stated surreal and omnific consequences. Whether the arbitrary-rank formulation, the descent as an entire-function counterexample or the three-stage minimal-order proof already occurs in the literature needs a more exhaustive bibliographic review; the answered question is a question of this report, not a named historical conjecture.

25.6 What the theta-descent part does not claim

Every limitation stated by source 13, in its manuscript, README, audit files or verification record, is kept.

- (T1) No Lean formalization, no independent referee report and no certified bibliographic priority accompany source 13; its compiled and visually inspected PDF is no evidence of mathematical correctness.

- (T2) The classical theta, triple-product, Chebyshev and elliptic identities are not new, and no novelty is claimed for third-order theta differential relations; the counterexample is a classical function.
- (T3) The no-order-unit direction is prior repository material (Theorem E (a)), re-proved by source 13 and not claimed.
- (T4) The bilateral theta domain and the Hahn–Tate uniformization are not claimed anew.
- (T5) $\mathcal{T}_p$ is not strongly entire on $\mathbf{No}$ or $\mathbf{No}[i]$; it fails at $x = \omega^\omega$. Entireness is relative to one specified Hahn field, and an extension of the value group can destroy it.
- (T6) D_z kills every Hahn coefficient, including $p = \omega^{-1}$, and is not the Berarducci–Mantova derivation.
- (T7) The finite checks do not verify strong summability of infinite or arbitrary supports, transcendence, algebraic independence or minimal order, the no-order-unit obstruction or the classification, completeness of the infinite zero set, novelty, or correctness in general; they are not a formal verification.
- (T8) No classification of strongly entire differentially algebraic series is given, and whether minimal order two occurs for some other series is open; a lower-order equation for $\mathcal{T}_p$ itself is excluded, which is a different assertion. (Source 14 has since excluded order two for every series, Theorem I; the non-claim is kept as source 13’s statement, and the classification stays open.) The geometric route suggested by Theorem 24.10 is a research direction, not a theorem.
- (T9) “No omnific zero” concerns the strong domain only; it is not a polynomial Diophantine statement, and no values are assigned outside the domain.
- (T10) The local valuation formulas are not uniform across the roots; a small input error does not imply a small output error.
- (T11) No transcendence theorem for numerical theta values is used or claimed; the independence statements, and all fields of meromorphic functions in the proofs, concern formal functions, not values at a point.
- (T12) The minimal-order, differential-field and non-D-finiteness arguments assume that μ is an order unit. The parameter is the monomial $p = t^\mu$; no result is generalized to a parameter with an arbitrary higher-order tail.
- (T13) Sebbar’s paper and Ohyama’s paper were not read in full, so no absence claim is made about them; broad searches are not evidence of absence, and no exhaustive MathSciNet or zbMATH review was made.
- (T14) The repository comparison at `4cdeaec` was targeted, not a full audit of every file, archived manuscript, branch or commit; the Hahn–Tate report was compared through its README.
- (T15) The answered question is a question of this report; no named historical conjecture is claimed solved.
- (T16) The existence classification quantifies over the scale: not every positive μ gives an entire series (Example 25.5).
- (T17) Without an order unit the conclusion excludes differentially algebraic nonpolynomial entire series only, not nonpolynomial entire series (Example 25.6).
- (T18) The poles of the rational system at $z = \pm 2$ are those of a coordinate chart; the globally meaningful equation is the cleared one.

- (T19) *[merge]* Lemmas 24.5 and 24.7, Proposition 24.2, clause (iv) of Theorem G, Corollary 25.2 and the consistency statements of Section 25.2 are the merge’s. The first two write out steps that source 13 used; no historical priority is claimed for any of them.

26 Second-order rigidity: finite endpoint spectra at exterior breaks

This section and the next contain the material of source 14, delivered after the theta-descent part had been merged. They answer the question left open by source 13 and re-scoped in Section 19 after it: no nonpolynomial strongly entire series satisfies a nonzero algebraic differential equation of order at most two, for every value group and every coefficient field of characteristic zero (Theorem I(a)). With Theorem H this makes the least order exactly three when Γ has an order unit (Theorem I(b)). The method is the finite initial polynomial of Sections 14–17, applied not at one scale but at infinitely many scales where the initial polynomial is not a monomial, and read through logarithmic Euler derivatives; it succeeds exactly where the corner criterion of Theorem 16.1 is silent, since an equation of order two may have $\chi_P = 0$ (Proposition 16.4). It is neither the escape chain, nor the finite coefficient field, nor a difference-field argument.

26.1 Source, conventions and the merge’s choices

Convention 26.1 (Setting of the order-three-threshold part). In Sections 26–27, k is an arbitrary field of characteristic zero with the trivial valuation, Γ is as in Convention 1.1, $\mathbb{K} = k((t^\Gamma))$, and v , lc , res , strong summability, $\text{Dom}_{\text{str}}(f)$, $\mathcal{E}_\Gamma(k)$, $\text{Frac } \mathcal{E}_\Gamma(k)$, D_z , the coefficient family algebra $\mathcal{S}_1$ with its residue homomorphism ρ , and the Gauss data γ_δ , ϕ_δ , N_δ of (14.1) are as in Conventions 14.1 and 20.1 and Section 14.3. Write $\vartheta = zD_z$, as in Convention 23.1, and $\vartheta_X = X d/dX$ on $k(X)$. Where a statement needs a divisible group it says so; it is then applied to the divisible hull $\Gamma_{\mathbb{Q}} = \Gamma \otimes_{\mathbb{Z}} \mathbb{Q}$ and $\mathbb{K}_{\mathbb{Q}} = k((t^{\Gamma_{\mathbb{Q}}}))$ (Lemma 26.2). U, V, W, T and G are indeterminates of polynomial rings over k in this part, except that in Corollary 26.13 G is the series of Corollary 17.1; $E_0, \dots, E_s$ are Euler-jet variables.

For $k = \mathbb{C}$ one has $\mathbb{K} = K_\Gamma$ and $\mathcal{E}_\Gamma(k) = \mathcal{E}_\Gamma$. As for the three previous parts, the generality in k is recorded for this part only.

The source. Source 14 is *The Order-Three Threshold for Entire Hahn–Surcomplex Functions: Second-order rigidity, Newton-corner spectra, and sharp theta examples* (26 pages, 23 September 2026). It consulted the repository at

b895e8672990e8b5a56f97dd7246f9dd5f86c771.

At that pin the article and README of this report were exactly the five-source text (sources 08–12); source 13’s audit files, programs and records had been placed in this directory but its manuscript had not yet been merged. Source 14 read this report’s README there, and took source 13 from the author’s own file library as an unpublished manuscript, with its baseline `4cdeaec` and its question `q:order2`, “whether a different example could have order two”. Its statements about this report (first-order nonlinear rigidity and the all-order obstruction without an order unit) were accurate at the pin. Since then source 13 has been merged as Sections 23–25, and `q:order2` is its Question 11.1, merged into Question 19.1; the question source 14 answers is therefore the one recorded there, and source 14’s description of source 13 as a private project source that is not a published paper

describes the state at its pin. Source 14 assumes characteristic zero and an arbitrary ordered abelian group; it passes to the divisible hull by a cofinal extension and uses no divisibility, rank, algebraic or real closedness or order-unit hypothesis in its main theorem. Its standing hypotheses are therefore those of the coefficient-field and theta-descent parts, and every proof taken from it was re-read against the statement it is attached to.

Renamed symbols. The following symbols of source 14 would collide with symbols of the earlier parts. They are renamed here, in the new material only.

Source 14	Here	Reason
$D = d/dz, f', f''$	$D_z, D_z f, D_z^2 f$	The notation of Convention 14.1. (Source 14's $\mathbb{K} = k((t^\Gamma))$, $\mathcal{E}_\Gamma(k)$ and $\vartheta = zD_z$ are this report's.)
radius r ; $\alpha_f(r)$; ϕ_r ; $I_f(r)$; $m_f(r)$, $N_f(r)$ Newton corner	scale δ ; γ_δ ; ϕ_δ ; Act_δ ; $\underline{N}_\delta$, N_δ break (Definition before Lemma 26.3)	The Gauss data (14.1); r indexes derivative orders, and I_P is the full corner polynomial. A <i>corner</i> is the finite term selection of Section 14.4.
restricted ring $\mathcal{O}\langle X \rangle_v$, red order bound ℓ ; blocks $P_{d,j}$; μ , M , C H (exterior homogeneous polynomial)	$\mathcal{S}_1^+$, ρ s ; $P_{d,w}$; β , n_1 , C_1 $\mathfrak{h}_P$	Lemma 14.2; red_Δ is a coarsened residue. ℓ is a dilation power; μ is a scale; $M(S)$ is a support monoid. $H \in \Gamma$, and $\mathcal{H}[f]$ is a theta expression.
$Q(U, V)$, $Q(U, V, W)$; s , R ; Q_s S_H	$\mathfrak{q}$; i_0 , $\mathfrak{r}$; $\mathfrak{q}_{i_0}$ $\text{Sp}(\mathfrak{h})$	Q_s is a shift polynomial, s an order, R an order bound. S is a support alphabet.
$u = \vartheta_X \phi / \phi$, V_ϕ , W_ϕ ; $h = W/V$, r $\chi_H(T, G)$	$\mathfrak{u}_\phi$, $\mathfrak{v}_\phi$, $\mathfrak{w}_\phi$; $\mathbf{k}$, $\mathbf{r}$ $\Upsilon_{\mathfrak{h}}(T, G)$	u is the Laurent variable of Section 24.5. χ_P is the residue corner polynomial.
upper gap g , lower gap e h , $q = t^h$, $Q = q^2$, $s_r(Q)$, c	$\mathbf{g}^+$, $\mathbf{g}^-$ μ , $p = t^\mu$, p^2 , $\mathcal{Q}_r(p^2)$, $\mathfrak{c}$	g_2, g_3 are theta invariants, $e_n \in \Gamma$. As for source 13 (Section 23.1).
F_q ; H_f , J_f , B_f $r_n = v(a_{n+1}) - v(a_n)$ $F = g/p$; characteristic p	$\mathcal{T}_p$; $\mathcal{H}[f]$, $\mathcal{K}[f]$, $\mathcal{B}[f]$ δ_n $F_0 = g/p$; $\mathfrak{p}$	(1.4), (24.5), (24.6). It is a break scale. F is the series (11.2); $p = t^\mu$.
differential field L ; state variables u, v Stirling numbers (implicit)	$\mathcal{L} \subseteq \mathfrak{F}$; u_1, u_2 $\varsigma_{j,i}$	L is a coefficient subfield, $M(S)$ a support monoid. s is an order.
$\Gamma_{\mathbb{Q}}$; $\mathbf{No}(i)$ Theorems 1.1 and 1.3	$\Gamma_{\mathbb{Q}}$; $\mathbf{No}[i]$ Theorem I (a) and (b)	The collection writes $\mathbf{No}[i]$. Theorems A–H of this report are different theorems.

The shipped audit file `14-order-three-threshold-SOURCE_AUDIT.md` keeps the source's notation.

Words with two meanings. A *break* is a scale δ at which the initial polynomial ϕ_δ is not a monomial; source 14 calls it a Newton corner. A *corner* keeps the meaning of Section 14.4, and the residue corner polynomial χ_P of a differential polynomial is not the endpoint-gap polynomial $\Upsilon_{\mathfrak{h}}$ of a homogeneous one; Remark 26.6 gives their exact relation. *Exterior* means, as elsewhere in this report, large δ , that is arguments $t^{-\delta}$ of large absolute valuation. The *endpoint spectrum* $\text{Sp}(\mathfrak{h})$ is a finite set of nonnegative integers, not the spectrum of an operator. *Order* and *minimal order* are

as in Section 23.1: the order of an equation is the highest derivative in it. A *gap* is a difference of exponents of ϕ at one end. u_ϕ, v_ϕ, w_ϕ are rational functions of X attached to a polynomial ϕ , while U, V, W are indeterminates.

Results printed once. Source 14 reproves several facts that the report already contains. They are printed once and cited: the definitions of an order unit and of strong entireness (Definitions 1.3 and 1.2); its all-scale coefficient criterion (Lemma 2.2), which is Proposition 20.2, proved there by the same two-scale argument; the restricted-series ring and its reduction homomorphism, which are $\mathcal{S}_1^+$ and ρ of Lemma 14.2, with the commutation of Euler derivatives and reduction in Lemma 14.4; the first assertion of its Lemma 3.2, which is Lemma 14.5; its Proposition 5.2, whose equation $z^2(ff'' - (f')^2) + zff' = 0$ is (16.1) and whose classification is Proposition 16.4; its Corollary 6.4 on Painlevé I and II, which is Proposition 16.3 (source 14's second route is recorded in Remark 26.12); its Section 8, the theta example, whose entireness, coefficients, cleared equation and derivation are Theorem 23.5, Proposition 23.3, Theorem 24.1 and Sections 24.2–24.3, except that its minimality argument is a genuinely different route and is kept (Remark 26.14); its Section 10, the no-order-unit direction, which is Theorem E (a) in the Euler presentation already recorded in Remark 25.1 (its Lemmas 10.1–10.2 and Theorem 10.3 are Theorem 21.2 (a), Lemmas 20.6–20.7 and Proposition 20.9); its Examples 10.4 and 10.5, which are Example 25.6 and, over $\mathbb{R} \oplus_{\text{lex}} \mathbb{R}$ instead of $\mathbb{Z} \oplus_{\text{lex}} \mathbb{Z}$, Example 25.5; the surreal embedding (2.4); the surreal witness, product, zeros and omnific rounding of its Section 11.3, which are Section 25.3, Theorem 23.6 and Corollary 25.3; its Example 12.1 in positive characteristic, which is Example 17.3 with the scale 1 of $\mathbb{Q}$ replaced by an order unit of Γ ; and the $\Gamma = 0$ remark, which is in Section 12. Source 14 credits the theta material and the no-order-unit direction as prior and claims neither.

Where each numbered result went. Theorem 1.1 is Theorem I(a), and Theorem 1.3 is Theorem I(b); Lemma 2.4 is Lemma 26.2; Lemma 3.1 is Lemma 26.3; the second assertion of Lemma 3.2 is Lemma 26.4; Lemma 4.1 and Remark 4.2 are Lemma 26.5 and Example 26.7; Lemma 5.1 is Lemma 26.8; Corollaries 6.1–6.3 are Corollaries 26.9–26.11; Definition 7.1, Theorem 7.2 and Corollary 7.3 are Definition 27.1, Theorem 27.2 and Corollary 27.3; Theorems 9.1 and 9.2 are Theorems 27.4 and 27.5; Proposition 11.1 is Proposition 27.6; Research question 1 is merged into the re-scoped Question 19.1, and Research questions 2–9 are Questions 19.11–19.18; Appendix A is (24.8) with (27.3); Appendix B and the proof checkpoints of Section 14.1 are merged into the dependency table of Section 18.

The merge's choices. The review of source 14 for this merge found its proofs correct. Three points are written out here and marked as the merge's. First, the existence of a late break (Lemma 26.4) is proved by naming the break scale, the least of finitely many quotients, rather than by the phrase “a change of the minimum occurs at an equality”. Second, the exterior form $\mathfrak{h}_P$ of Lemma 26.5 is identified with the residue corner of Section 14.4 written in Euler jets, and its values on monomials are χ_P (Remark 26.6); this is what connects source 14's endpoint lemma to the corner criterion of source 10. Third, Corollary 26.9 and Corollary 26.10 are stated inside an explicit differential field containing $\mathbb{K}((z))$, which is how source 14's phrase “every $f \in L \cap \mathcal{E}_\Gamma(k)$ ” must be read, and Corollary 26.13 applies Theorem I to the series G and Θ_p of earlier parts. Source 14's theorems, proofs and non-claims are otherwise printed as delivered, in the renamed notation.

26.2 Breaks and the exterior form

Lemma 26.2 (Cofinal scalar extension; source 14). *Let $\Gamma \subseteq \Lambda$ be an extension of ordered abelian groups with Γ cofinal in Λ . Every $f \in \mathcal{E}_\Gamma(k)$ is strongly entire over $k((t^\Lambda))$. In particular this holds for $\Lambda = \Gamma_\mathbb{Q}$, in which Γ is cofinal.*

Proof. Write $f = \sum_n a_n z^n$ and use Proposition 20.2 (iii) over Λ . Given $\gamma, \beta \in \Lambda$, choose $\gamma' \in \Gamma$ with $\gamma' \leq \gamma$ and $\beta' \in \Gamma$ with $\beta' \geq \beta$; since Γ is cofinal in the group Λ , it is also coinital. If $n \geq 0$ and $v(a_n) + n\gamma \leq \beta$, then $v(a_n) + n\gamma' \leq \beta'$, which holds for only finitely many n with $a_n \neq 0$. For $\Lambda = \Gamma_\mathbb{Q}$, the torsion-free group Γ embeds in $\Gamma \otimes_\mathbb{Z} \mathbb{Q}$, and an element g/m with $g \in \Gamma$ and $m \geq 1$ satisfies $|g/m| \leq |g|$. $\square$

A series is a polynomial over $\mathbb{K}_\mathbb{Q}$ exactly when it is one over $\mathbb{K}$, and a polynomial relation over $\mathbb{K}(z)$ remains a nonzero relation over $\mathbb{K}_\mathbb{Q}(z)$. So a polynomiality conclusion proved over $\mathbb{K}_\mathbb{Q}$ descends to $\mathbb{K}$. This is the only use of divisibility in the part.

Let $f = \sum_n a_n z^n \in \mathcal{E}_\Gamma(k)$ be nonzero. For every $\delta \in \Gamma$ the Gauss value γ_δ and the initial polynomial ϕ_δ of (14.1) exist. Put

$$\text{Act}_\delta = \{n : a_n \neq 0, v(a_n) - n\delta = \gamma_\delta\}, \quad \underline{N}_\delta = \min \text{Act}_\delta, \quad N_\delta = \max \text{Act}_\delta, \quad (26.1)$$

so that $\underline{N}_\delta = \text{ord}_X \phi_\delta$, the X -adic order, and $N_\delta = \deg \phi_\delta$. A scale δ is a *break* of f if $\underline{N}_\delta < N_\delta$, that is, if ϕ_δ is not a monomial. Lemma 14.2 gives $\rho(t^{-\gamma_\delta}(\vartheta^j f)(t^{-\delta} X)) = \vartheta_X^j \phi_\delta$ for every j , because $\vartheta^j f$ scales like f with each coefficient multiplied by the integer n^j , and Euler operators commute with ρ . A derivative on the right may vanish; no equality of individual valuations is asserted.

Lemma 26.3 (Local finiteness of active indices; source 14). *Let $f \in \mathcal{E}_\Gamma(k)$ be nonzero and $\delta_1 \leq \delta_2$ in Γ . Only finitely many indices lie in Act_δ for some $\delta \in [\delta_1, \delta_2]$. If $\delta < \delta'$, $n \in \text{Act}_\delta$ and $n' \in \text{Act}_{\delta'}$, then $n \leq n'$.*

Proof. Fix $a_j \neq 0$. On the interval its weighted value $v(a_j) - j\delta$ is at most $C = v(a_j) - j\delta_1$. By Proposition 20.2 (iii) at $\gamma = -\delta_2$ all but finitely many nonzero a_n have $v(a_n) - n\delta_2 > C$, and then $v(a_n) - n\delta \geq v(a_n) - n\delta_2 > C$ for $\delta \leq \delta_2$, so such n is never active. For the monotonicity, the two minimality inequalities $v(a_n) - n\delta \leq v(a_{n'}) - n'\delta$ and $v(a_{n'}) - n'\delta' \leq v(a_n) - n\delta'$ add to $(n' - n)(\delta' - \delta) \geq 0$. $\square$

Lemma 26.4 (Late breaks; source 14). *Let Γ be divisible and $f \in \mathcal{E}_\Gamma(k)$ nonpolynomial. For every $\delta_* \in \Gamma$ and every $B \in \mathbb{N}$ there is a break $\delta \geq \delta_*$ of f with $\underline{N}_\delta > B$.*

Proof. By Lemma 14.5 there is δ_B such that every active index at every $\delta \geq \delta_B$ exceeds B . Let $\delta_1 = \max(\delta_*, \delta_B)$. If δ_1 is a break we are done; otherwise $\text{Act}_{\delta_1} = \{N\}$. Lemma 14.5 again gives $\delta_2 > \delta_1$ at which every active index exceeds N . By Lemma 26.3 the set F of indices active somewhere on $[\delta_1, \delta_2]$ is finite, and on that interval γ_δ is the minimum of the finitely many affine functions $v(a_n) - n\delta$, $n \in F$. For $n \in F$ with $n > N$ put

$$\delta^{(n)} = \frac{v(a_n) - v(a_N)}{n - N} \in \Gamma,$$

using divisibility; then $v(a_N) - N\delta \leq v(a_n) - n\delta$ exactly when $\delta \leq \delta^{(n)}$. Since N is active at δ_1 , every $\delta^{(n)} \geq \delta_1$, and since some $n > N$ in F is active at δ_2 while N is not, $\delta^{(n)} < \delta_2$ for that n . Let δ^* be the least $\delta^{(n)}$ (the merge names this scale). For $n \in F$ with $n < N$, activity of N at δ_1 and

$\delta^* \geq \delta_1$ give $v(a_N) - N\delta^* \leq v(a_n) - n\delta^*$; for $n > N$ the same holds because $\delta^* \leq \delta^{(n)}$, with equality for an index attaining δ^* . Since the least weighted value on the interval is attained in F , N and that index are both active at δ^* , which is a break, and $\delta^* \geq \delta_1 \geq \delta_B$ makes every active index there exceed B . $\square$

The proof uses only finite lower envelopes on bounded intervals: no completeness of Γ , no real-valued norm, no countable cofinal sequence and no order unit.

Let $P \in \mathbb{K}[z, Y_0, \dots, Y_s]$ be a nonzero differential polynomial of order at most s . Since $z^j D_z^j = \vartheta(\vartheta - 1) \cdots (\vartheta - j + 1)$, the substitution

$$Y_j \mapsto z^{-j} \sum_{i=0}^j \varsigma_{j,i} E_i \quad (0 \leq j \leq s), \quad (26.2)$$

with $\varsigma_{j,i}$ the signed Stirling numbers of the first kind, so that $\sum_i \varsigma_{j,i} T^i = (T)_j$, rewrites $P(z, f, D_z f, \dots)$ as a Laurent polynomial in z in the Euler jets $E_i = \vartheta^i f$. After multiplication by a power of z it becomes $\tilde{P}(z, f, \vartheta f, \dots, \vartheta^s f)$ with $0 \neq \tilde{P} \in \mathbb{K}[z, E_0, \dots, E_s]$; the change is invertible and preserves total jet degree. Decompose

$$\tilde{P} = \sum_{d,w} z^w P_{d,w}(E_0, \dots, E_s),$$

with $P_{d,w}$ homogeneous of total degree d in the jets, choose the largest d occurring, then the largest w with $P_{d,w} \neq 0$, let β be the least valuation of a coefficient of $P_{d,w}$, and put

$$\mathfrak{h}_P = \text{res}(t^{-\beta} P_{d,w}) \in k[Y_0, \dots, Y_s], \quad (26.3)$$

applying res to coefficients and renaming E_i as Y_i . It is a nonzero homogeneous polynomial, the *exterior form* of P .

Lemma 26.5 (Exterior homogeneous reduction; source 14). *Let $f \in \mathcal{E}_\Gamma(k)$ be nonpolynomial with $P(z, f, D_z f, \dots, D_z^s f) = 0$ for a nonzero P of order at most s . Then $\mathfrak{h}_P$ has positive total degree, and there is $\delta_0 \in \Gamma_{>0}$ such that for every $\delta \geq \delta_0$*

$$\mathfrak{h}_P(\phi_\delta, \vartheta_X \phi_\delta, \dots, \vartheta_X^s \phi_\delta) = 0 \quad \text{in } k[X].$$

No divisibility of Γ is used.

Proof. If $d = 0$, then $\tilde{P}$ is a nonzero element of $\mathbb{K}[z]$, which cannot vanish as a series; so $d \geq 1$. Because f is nonpolynomial there is n_1 with $a_{n_1} \neq 0$ and $(d - d')n_1 + w - w' > 0$ for every block $P_{d',w'} \neq 0$ with $d' < d$; put $C_1 = v(a_{n_1})$, so that $\gamma_\delta \leq C_1 - n_1\delta$ for every δ .

Evaluate at $z = t^{-\delta} X$ and normalize by $t^{\beta + d\gamma_\delta - w\delta}$. Each factor $(\vartheta^i f)(t^{-\delta} X)$ equals t^{γ_δ} times an element of $\mathcal{S}_1^+$, so a monomial of a block $z^{w'} P_{d',w'}$ with coefficient $b \neq 0$ becomes $t^{v(b) - w'\delta + d'\gamma_\delta}$ times an element of $\mathcal{S}_1^+$. Relative to the normalization its exponent is at least

$$v(b) - \beta - (d - d')\gamma_\delta + (w - w')\delta \geq v(b) - \beta - (d - d')C_1 + ((d - d')n_1 + w - w')\delta. \quad (26.4)$$

For $d' < d$ the coefficient of δ is a positive integer by the choice of n_1 ; for $d' = d$ and $(d', w') \neq (d, w)$ it is $w - w' > 0$. There are finitely many monomials, so the right side is positive for all $\delta \geq \delta_0$ once $\delta_0 > 0$ exceeds the absolute values of the finitely many constants; no division occurs. The monomials of $P_{d,w}$ itself with coefficients of value above β also lose their residue. Applying the ring homomorphism ρ of Lemma 14.2 to the normalized identity $0 = \tilde{P}(\dots)$ leaves $X^w \mathfrak{h}_P(\phi_\delta, \vartheta_X \phi_\delta, \dots) = 0$, and X^w cancels in $k[X]$. $\square$

Remark 26.6 (The exterior form is the residue corner in Euler jets; merge). Write P in the combined-monomial form (14.4). By (26.2) the monomial $c_\nu z^{k_\nu} \prod_j Y_j^{m_{\nu j}}$ becomes $c_\nu z^{w_\nu} \prod_j (\sum_i \varsigma_{j,i} E_i)^{m_{\nu j}}$, with the weight w_ν of (14.5). The substitution $Y_j \mapsto \sum_i \varsigma_{j,i} Y_i$ is a unitriangular, hence invertible, linear change of variables over $\mathbb{Z}$ and preserves total degree, so the selection of Lemma 26.5 picks $d = d_P$ and the z -power of weight w_P : its block is the image of $\sum_{\nu \in \mathcal{C}_P} c_\nu \prod_j Y_j^{m_{\nu j}}$. Its least coefficient value is β_P , because the image of the nonzero residue vector $(\bar{c}_\nu)$ under the invertible integral change is nonzero. Hence

$$\mathfrak{h}_P(Y_0, \dots, Y_s) = \sum_{\nu \in \mathcal{C}_P} \bar{c}_\nu \prod_{j=0}^s \left(\sum_{i=0}^j \varsigma_{j,i} Y_i \right)^{m_{\nu j}}, \quad \mathfrak{h}_P(1, T, T^2, \dots, T^s) = \chi_P(T),$$

the second because $\sum_i \varsigma_{j,i} T^i = (T)_{\underline{j}}$. On a monomial $\phi = X^N$ one has $\vartheta_X^i X^N = N^i X^N$, so $\mathfrak{h}_P(X^N, \vartheta_X X^N, \dots) = \chi_P(N) X^{d_P N}$: the monomial test of the exterior form is the residue corner polynomial, and the corner criterion of Theorem 16.1 uses exactly this test. When $\chi_P = 0$ the exterior form vanishes on every monomial but is still a nonzero polynomial, and source 14 keeps the whole initial polynomial at breaks. For the equation (16.1), $\mathfrak{h}_P = Y_0 Y_2 - Y_1^2$; for the cleared theta equation (24.7), $\beta_P = 0$ and the dehomogenized exterior form is (27.5) below, consistent with Proposition 24.2.

Example 26.7 (The cancellation that must not be ignored; source 14). For $\mathfrak{h} = Y_0 Y_2 - Y_1^2$ one has $\mathfrak{h}(X^N, \vartheta_X X^N, \vartheta_X^2 X^N) = 0$ for every N , so substituting a trial monomial and assuming that the resulting polynomial in N is nonzero proves nothing. This $\mathfrak{h}$ is the exterior form of (16.1), whose formal solutions are exactly 0 and the monomials $c z^n$ (Proposition 16.4, which is source 14's Proposition 5.2). Lemma 26.8 below gives $\text{Sp}(\mathfrak{h}) = \emptyset$ for it: no nonmonomial initial polynomial is permitted at any exterior break, which is stronger than the identically vanishing monomial test and consistent with unbounded monomial solutions. Polynomiality is therefore not a uniform degree bound: a fixed equation can have polynomial solutions of every degree.

26.3 The endpoint lemma and the proof of Theorem I (a)

For a nonzero $\phi \in k[X]$ put $\mathfrak{u}_\phi = \vartheta_X \phi / \phi$, $\mathfrak{v}_\phi = \vartheta_X \mathfrak{u}_\phi$ and $\mathfrak{w}_\phi = \vartheta_X \mathfrak{v}_\phi$ in $k(X)$. Then

$$\frac{\vartheta_X^2 \phi}{\phi} = \mathfrak{u}_\phi^2 + \mathfrak{v}_\phi, \quad \frac{\vartheta_X^3 \phi}{\phi} = \mathfrak{u}_\phi^3 + 3\mathfrak{u}_\phi \mathfrak{v}_\phi + \mathfrak{w}_\phi.$$

Lemma 26.8 (Finite endpoint spectrum; source 14). *Let $0 \neq \mathfrak{h} \in k[Y_0, Y_1, Y_2]$ be homogeneous. There is a finite set $\text{Sp}(\mathfrak{h}) \subseteq \mathbb{N}$ such that every nonmonomial $\phi \in k[X]$ with $\mathfrak{h}(\phi, \vartheta_X \phi, \vartheta_X^2 \phi) = 0$ has both its X -adic order $\text{ord}_X \phi$ and $\deg \phi$ in $\text{Sp}(\mathfrak{h})$.*

Proof. Put

$$\mathfrak{q}(U, V) = \mathfrak{h}(1, U, U^2 + V). \quad (26.5)$$

It is nonzero: setting $Y_0 = 1$ is injective on homogeneous polynomials of a fixed degree, since the missing power of Y_0 is determined by the other exponents, and $Y_2 \mapsto U^2 + V$ is a triangular automorphism. Let V^{i_0} be the largest power of V dividing $\mathfrak{q}$ and write

$$\mathfrak{q} = V^{i_0} \mathfrak{r}, \quad \mathfrak{r}(U, 0) \neq 0. \quad (26.6)$$

By homogeneity $\mathfrak{q}(\mathfrak{u}_\phi, \mathfrak{v}_\phi) = 0$. The function $\mathfrak{u}_\phi$ is not constant: $\vartheta_X \phi = b\phi$ with $b \in k$ forces $n = b$ for every exponent n of ϕ , and in characteristic zero there is at most one such exponent. The

constants of d/dX in $k(X)$ are k , so $v_\phi \neq 0$, and cancelling $v_\phi^{i_0}$ in the field $k(X)$ gives $\mathfrak{r}(u_\phi, v_\phi) = 0$. With $m = \text{ord}_X \phi$ and $N = \deg \phi$, the expansions at 0 and at ∞ are

$$u_\phi = m + O(X), \quad v_\phi = O(X); \quad u_\phi = N + O(X^{-1}), \quad v_\phi = O(X^{-1}),$$

and their constant terms give $\mathfrak{r}(m, 0) = \mathfrak{r}(N, 0) = 0$. The nonzero polynomial $\mathfrak{r}(T, 0)$ has finitely many roots in k , and distinct nonnegative integers are distinct in k ; let $\text{Sp}(\mathfrak{h})$ be its set of roots in $\mathbb{N}$. $\square$

Proof of Theorem I(a). Let $f \in \mathcal{E}_\Gamma(k)$ satisfy a nonzero relation of order at most two over $\mathbb{K}(z)$; clearing denominators gives a nonzero $P \in \mathbb{K}[z, Y_0, Y_1, Y_2]$. Suppose f is nonpolynomial. By Lemma 26.2, f is strongly entire over $\mathbb{K}_\mathbb{Q}$ and satisfies the same relation there, so we may assume Γ divisible. Lemma 26.5 gives $\mathfrak{h}_P \neq 0$ and δ_0 . Choose $B \in \mathbb{N}$ larger than every element of $\text{Sp}(\mathfrak{h}_P)$ ($B = 0$ if it is empty). Lemma 26.4 gives a break $\delta \geq \delta_0$ with $N_\delta > B$; its initial polynomial is nonmonomial, satisfies $\mathfrak{h}_P(\phi_\delta, \vartheta_X \phi_\delta, \vartheta_X^2 \phi_\delta) = 0$, and has $\text{ord}_X \phi_\delta = N_\delta \notin \text{Sp}(\mathfrak{h}_P)$, contradicting Lemma 26.8. So f is a polynomial over $\mathbb{K}_\mathbb{Q}$, hence over $\mathbb{K}$. The equivalent formulation is the same statement for the algebraic dependence of $f, D_z f, D_z^2 f$ over $\mathbb{K}(z)$. $\square$

The proof uses characteristic zero twice: nonzero integer multipliers keep valuation zero under differentiation, and infinitely many degree indices stay distinct in the residue field. The finite endpoint set is what the corner criterion lacks when $\chi_P = 0$: at order two the degenerate part of an equation leaves only the finitely many roots of $\mathfrak{r}(T, 0)$ for both endpoints of every nonmonomial initial polynomial, while a nonpolynomial entire series has breaks with arbitrarily large endpoints.

26.4 Consequences and the proof of Theorem I(b)

Corollary 26.9 (Three-jet transcendence; source 14). *For nonpolynomial $f \in \mathcal{E}_\Gamma(k)$,*

$$\text{trdeg}_{\mathbb{K}(z)} \mathbb{K}(z)(f, D_z f, D_z^2 f) = 3.$$

More generally, let $\mathfrak{F}$ be a differential field extension of $\mathbb{K}((z))$ whose derivation extends D_z , and $\mathfrak{L} \subseteq \mathfrak{F}$ a differential subfield containing $\mathbb{K}(z)$ with $\text{trdeg}_{\mathbb{K}(z)} \mathfrak{L} \leq 2$. Every $f \in \mathfrak{L} \cap \mathcal{E}_\Gamma(k)$ is a polynomial.

Proof. The first assertion is Theorem I(a). For the second, $f, D_z f, D_z^2 f \in \mathfrak{L}$ are algebraically dependent over $\mathbb{K}(z)$ in $\mathfrak{F}$, and a polynomial relation among elements of $\mathbb{K}((z))$ holds in $\mathfrak{F}$ exactly when it holds in $\mathbb{K}((z))$. $\square$

Corollary 26.10 (Rational two-state systems; source 14). *Let $\mathfrak{F}$ be as in Corollary 26.9 and let $u_1, u_2 \in \mathfrak{F}$ satisfy $D_z u_1 = R_1(z, u_1, u_2)$ and $D_z u_2 = R_2(z, u_1, u_2)$ with $R_1, R_2 \in \mathbb{K}(z, Y_1, Y_2)$, all denominators nonzero after substitution. Every element of $\mathbb{K}(z, u_1, u_2)$ that lies in $\mathcal{E}_\Gamma(k)$ is a polynomial.*

Proof. The system makes $\mathbb{K}(z, u_1, u_2)$ a differential field, by the quotient rule, of transcendence degree at most two over $\mathbb{K}(z)$; apply Corollary 26.9. $\square$

The state variables themselves need not be entire; the corollary restricts the strongly entire observables of the field. A nonpolynomial strongly entire function therefore cannot be generated by a rational nonautonomous system with two state variables over the full Hahn coefficient field, for

every Γ . Corollary 21.16 (source 11) allows any number of states and quotients of entire series but needs the absence of an order unit, and Corollary 5.5 is the linear case. With an order unit, $\mathcal{T}_p$ is an observable of the rational system with the four states $f, \mathcal{L}, \mathcal{A}, \mathcal{A}_1$ of Theorem 24.10, so there the number two cannot be replaced by four; three states are not decided here (an observation of the merge). Arbitrary transcendental right-hand sides remove the finite differential-field hypothesis and are outside the corollary.

Corollary 26.11 (Polynomial-denominator meromorphic rigidity; source 14). *Let $g \in \mathcal{E}_\Gamma(k)$, $0 \neq p \in \mathbb{K}[z]$ and $F_0 = g/p \in \mathbb{K}((z))$. If F_0 satisfies a nonzero algebraic differential equation of order at most two over $\mathbb{K}(z)$, then $F_0 \in \mathbb{K}(z)$.*

Proof. The relations $g = pF_0$, $D_z g = D_z p F_0 + p D_z F_0$ and $D_z^2 g = D_z^2 p F_0 + 2D_z p D_z F_0 + p D_z^2 F_0$ form an invertible triangular linear change of jets over $\mathbb{K}(z)$, so g satisfies a nonzero relation of order at most two, and Theorem I(a) makes it a polynomial. $\square$

This concerns quotients with a polynomial denominator only; it does not assert that every object called meromorphic with finitely many poles in a Hahn analytic category has such a representation. For the fraction field $\text{Frac } \mathcal{E}_\Gamma(k)$ without an order unit, Theorem E(c) gives rationality in every order.

Remark 26.12 (Painlevé I and II: a second route; source 14). Source 14's Corollary 6.4 is Proposition 16.3, with the same degree comparison for polynomials. Source 10 reaches polynomiality through the nonzero corners $\chi_P = -6$ and $\chi_P = -2$; source 14 reaches it through Theorem I(a), since both equations have order two. The equations of DLMF §32.2 [52] are read as formal equations over $\mathbb{K}$; nothing is said about local formal solutions, which exist, or about classical complex Painlevé transcendents.

Corollary 26.13 (The explicit series of earlier parts; merge). *The series $G = \sum_{n \geq 0} t^{n^2} z^n$ over $\mathbb{C}((t^\mathbb{Q}))$ of Corollary 17.1, and, for an order unit μ and $p = t^\mu$, the partial theta series $\Theta_p \in \mathcal{E}_\Gamma(k)$ of Proposition 21.19, have three algebraically independent jets $f, D_z f, D_z^2 f$ over the respective $\mathbb{K}(z)$.*

Proof. Both are nonpolynomial and strongly entire (Corollary 17.1, Proposition 21.19); apply Theorem I(a). $\square$

Whether G or Θ_p satisfies an algebraic differential equation of order at least three is not decided; the corollary only raises the known lower bound from two jets (Corollary 17.1) to three.

Proof of Theorem I(b). Let μ be an order unit and $p = t^\mu$. By Theorem 23.5 and Proposition 23.3, $\mathcal{T}_p$ is a nonpolynomial element of $\mathcal{E}_\Gamma(k)$, and by Theorem 24.1 it satisfies a nonzero algebraic differential equation of order three. By part (a) no nonpolynomial element of $\mathcal{E}_\Gamma(k)$ satisfies one of order at most two; an equation of order zero is included, being of order at most two. So the least order is three. Without an order unit, Theorem E(a) excludes every finite order. $\square$

When $\Gamma = 0$ every strongly entire series is a polynomial, by strong summability at $x = 1$ (Section 12); source 14 includes this case in its statement, and Convention 1.1 excludes it here.

Remark 26.14 (A second route to minimal order three for $\mathcal{T}_p$; source 14). The lower bound of Corollary 24.9, that $\mathcal{T}_p, D_z \mathcal{T}_p, D_z^2 \mathcal{T}_p$ are algebraically independent over $\mathbb{K}(z)$ when μ is an order unit, is also a case of Theorem I(a), since $\mathcal{T}_p$ is nonpolynomial and strongly entire. Source 14 proves minimality this way, “without the separate difference-field minimality argument”. The two routes

are genuinely different: source 13's argument uses the Laurent ring, the substitution homomorphism and three difference-field obstructions for $u \mapsto p^2u$, and also yields the degree-two statement of Theorem 24.10 and non-D-finiteness (Proposition 24.11), which Theorem I does not give; source 14's argument uses no property of $\mathcal{T}_p$ beyond nonpolynomial strong entireness. Both are kept. The strict inequality of orders also shows again that $\mathcal{T}_p$ satisfies no equation to which Theorem 15.1 applies (Remark 24.3), but not conversely.

27 Order three: endpoint gaps, theta breaks and scope

27.1 The endpoint-gap polynomial

At order three the endpoint lemma has a precise analogue with one more integer parameter; this is what an equation of order three can use, and the theta equation uses it.

Definition 27.1 (Endpoint gaps; source 14). For a nonmonomial $\phi \in k[X]$ with highest exponent N and second-highest exponent $N - \mathbf{g}^+$, the positive integer $\mathbf{g}^+$ is its *upper gap*; with lowest exponent m and second-lowest exponent $m + \mathbf{g}^-$, the positive integer $\mathbf{g}^-$ is its *lower gap*. The pairs $(N, \mathbf{g}^+)$ and $(m, \mathbf{g}^-)$ are its endpoint-gap pairs.

Theorem 27.2 (Third-order endpoint-gap obstruction; source 14). *Let $0 \neq \mathfrak{h} \in k[Y_0, Y_1, Y_2, Y_3]$ be homogeneous and*

$$\mathfrak{q}(U, V, W) = \mathfrak{h}(1, U, U^2 + V, U^3 + 3UV + W). \quad (27.1)$$

Let i_0 be the least total degree in (V, W) of a nonzero term of $\mathfrak{q}$ and $\mathfrak{q}_{i_0}$ the sum of the terms of that degree. Then

$$\Upsilon_{\mathfrak{h}}(T, G) = \mathfrak{q}_{i_0}(T, 1, -G) \in k[T, G] \quad (27.2)$$

is nonzero, and every nonmonomial $\phi \in k[X]$ with $\mathfrak{h}(\phi, \vartheta_X \phi, \vartheta_X^2 \phi, \vartheta_X^3 \phi) = 0$ satisfies $\Upsilon_{\mathfrak{h}}(N, \mathbf{g}^+) = 0$ and $\Upsilon_{\mathfrak{h}}(m, -\mathbf{g}^-) = 0$ for its endpoint-gap pairs.

Proof. As in Lemma 26.8, $\mathfrak{q} \neq 0$. The polynomial $\mathfrak{q}_{i_0}$ is homogeneous of degree i_0 in (V, W) , and $V = 1, W = -G$ is injective on such polynomials, so $\Upsilon_{\mathfrak{h}} \neq 0$. By homogeneity and the identities before Lemma 26.8, $\mathfrak{q}(\mathbf{u}_\phi, \mathbf{v}_\phi, \mathbf{w}_\phi) = 0$. At infinity write $\phi = aX^N(1 + bX^{-\mathbf{g}^+} + \dots)$ with $ab \neq 0$; then, in $k((X^{-1}))$,

$$\mathbf{u}_\phi = N - \mathbf{g}^+ bX^{-\mathbf{g}^+} + \dots, \quad \mathbf{v}_\phi = (\mathbf{g}^+)^2 bX^{-\mathbf{g}^+} + \dots, \quad \mathbf{w}_\phi / \mathbf{v}_\phi = -\mathbf{g}^+ + O(X^{-1}).$$

Writing each homogeneous part as $\mathfrak{q}_i(\mathbf{u}_\phi, \mathbf{v}_\phi, \mathbf{w}_\phi) = \mathbf{v}_\phi^i \mathfrak{q}_i(\mathbf{u}_\phi, 1, \mathbf{w}_\phi / \mathbf{v}_\phi)$, the parts with $i > i_0$ have X^{-1} -order above $i_0 \mathbf{g}^+$, and the coefficient of $X^{-i_0 \mathbf{g}^+}$ is $((\mathbf{g}^+)^2 b)^{i_0} \mathfrak{q}_{i_0}(N, 1, -\mathbf{g}^+)$; replacing $\mathbf{u}_\phi$ by N changes only higher orders. So $\Upsilon_{\mathfrak{h}}(N, \mathbf{g}^+) = 0$. At 0 write $\phi = aX^m(1 + bX^{\mathbf{g}^-} + \dots)$: then $\mathbf{u}_\phi \rightarrow m$, $\mathbf{v}_\phi$ has order $\mathbf{g}^-$ and $\mathbf{w}_\phi / \mathbf{v}_\phi \rightarrow \mathbf{g}^-$, and the same argument gives $\mathfrak{q}_{i_0}(m, 1, \mathbf{g}^-) = \Upsilon_{\mathfrak{h}}(m, -\mathbf{g}^-) = 0$. $\square$

At order two the lowest transverse part is $A(U)V^{i_0}$, whose specialization is $A(T)$: no gap variable remains, which is Lemma 26.8. At order three a polynomial such as $G^2 - 1$ permits arbitrarily large endpoints while fixing the gap to one; this is what happens for the theta equation (Section 27.2).

Corollary 27.3 (A third-order rigidity certificate; source 14). *Let P have order at most three, with exterior form $\mathfrak{h}_P$ as in Lemma 26.5. If the set*

$$\{N \in \mathbb{N} : \Upsilon_{\mathfrak{h}_P}(N, \mathbf{g}) = 0 \text{ for some integer } 1 \leq \mathbf{g} \leq N\}$$

is bounded, every strongly entire solution of $P[f] = 0$ is a polynomial.

Proof. Pass to $\Gamma_{\mathbb{Q}}$ as in the proof of Theorem I(a). A nonpolynomial entire solution has breaks $\delta \geq \delta_0$ with $\underline{N}_{\delta}$, hence N_{δ} , arbitrarily large (Lemma 26.4), and the upper endpoint-gap pair $(N_{\delta}, \mathbf{g}^+)$ of ϕ_{δ} has $1 \leq \mathbf{g}^+ \leq N_{\delta}$; Theorem 27.2 and Lemma 26.5 contradict the boundedness. $\square$

This is a sufficient certificate, not a decision algorithm for third-order equations: deciding the integral zero set of a bivariate polynomial is a separate arithmetic problem, and an unbounded zero set of $\Upsilon_{\mathfrak{h}_P}$ is necessary for escaping the obstruction, not sufficient for the existence of an entire solution.

27.2 The exterior form of the theta equation

Let P be the polynomial (24.8) with the constants (24.2); it is homogeneous of degree six in the jets. In Euler jets E_j the substitution (26.2) reads

$$Y_0 = E_0, \quad Y_1 = E_1/z, \quad Y_2 = (E_2 - E_1)/z^2, \quad Y_3 = (E_3 - 3E_2 + 2E_1)/z^3, \quad (27.3)$$

and after multiplication by z^6 the highest z -power is six. Its block, dehomogenized by $E_0 = 1$, $E_1 = U$, $E_2 = U^2 + V$, $E_3 = U^3 + 3UV + W$, is

$$W^2 - 4(\mathfrak{c} - V)^3 + g_2(\mathfrak{c} - V) + g_3 = W^2 + 4V^3 - 12\mathfrak{c}V^2 + (12\mathfrak{c}^2 - g_2)V - 4\mathfrak{c}^3 + \mathfrak{c}g_2 + g_3. \quad (27.4)$$

Directly: in Euler jets the top part of $\mathcal{H}[f]$ is $f\vartheta^2 f - (\vartheta f)^2$, that is $E_0 E_2 - E_1^2$, which dehomogenizes to V ; the top part of $z\mathcal{K}[f]$ is $f\vartheta\mathcal{H}_0 - 2\vartheta f\mathcal{H}_0$ with $\mathcal{H}_0$ that part, which dehomogenizes to W ; so $\mathcal{B}[f]$ contributes $\mathfrak{c} - V$ and $(z^2 - 4)\mathcal{K}[f]^2$ contributes W^2 . The coefficient of W^2 is 1 and all coefficients lie in $\mathbb{Z}[\mathfrak{c}, g_2, g_3]$, of nonnegative valuation, so $\beta = 0$ in (26.3). With the residues $\text{res } \mathfrak{c} = \frac{1}{12}$, $\text{res } g_2 = \frac{1}{12}$ and $\text{res } g_3 = -\frac{1}{216}$ the constant and linear terms cancel and the dehomogenized exterior form is

$$W^2 - V^2 + 4V^3. \quad (27.5)$$

Its value at $V = W = 0$ is 0, which is $\chi_P = 0$ in Proposition 24.2, by Remark 26.6. Its transverse part of least degree is $W^2 - V^2$, so Theorem 27.2 gives $\Upsilon(T, G) = G^2 - 1$: every exterior upper and lower gap of a solution is one. The Euler-jet conversion and (27.5) are checked symbolically by source 14's program.

27.3 Theta breaks and eventual Newton convexity

Theorem 27.4 (Polynomial solutions of the theta exterior equation; source 14). *Let $0 \neq \phi \in k[X]$. Then $\mathbf{w}_{\phi}^2 - \mathbf{v}_{\phi}^2 + 4\mathbf{v}_{\phi}^3 = 0$ if and only if $\phi = bX^m$ or $\phi = bX^m(1 + aX)$ with $m \in \mathbb{N}$ and $a, b \in k^{\times}$.*

Proof. If $\mathbf{v}_{\phi} = 0$, then $\mathbf{u}_{\phi}$ is constant and ϕ is a monomial, as in the proof of Lemma 26.8. Suppose $\mathbf{v}_{\phi} \neq 0$ and put $\mathbf{k} = \mathbf{w}_{\phi}/\mathbf{v}_{\phi}$. The equation becomes $\mathbf{k}^2 = 1 - 4\mathbf{v}_{\phi}$. If $\mathbf{k} = 0$ then $\mathbf{v}_{\phi} = \frac{1}{4}$, impossible because the constant term of $\vartheta_X \mathbf{u}_{\phi}$ in its Laurent expansion at 0 is zero. Applying ϑ_X and using $\mathbf{w}_{\phi} = \mathbf{k}\mathbf{v}_{\phi}$ gives $2\mathbf{k}\vartheta_X \mathbf{k} = -4\mathbf{k}\mathbf{v}_{\phi}$, so $\vartheta_X \mathbf{k} = (\mathbf{k}^2 - 1)/2$; and $\mathbf{k} \neq \pm 1$, since either would give $\mathbf{v}_{\phi} = 0$. For $\mathbf{r} = (\mathbf{k} - 1)/(\mathbf{k} + 1)$ a direct computation gives $\vartheta_X \mathbf{r} = \mathbf{r}$, so $d(\mathbf{r}/X)/dX = 0$ and $\mathbf{r} = b_1 X$ with $b_1 \in k^{\times}$. With $a = -b_1$,

$$\mathbf{k} = \frac{1 - aX}{1 + aX}, \quad \mathbf{v}_{\phi} = \frac{aX}{(1 + aX)^2}, \quad \vartheta_X \left(\mathbf{u}_{\phi} - \frac{aX}{1 + aX} \right) = 0.$$

The constant is $m = \text{ord}_X \phi$, by evaluation at 0, so $\phi/(X^m(1+aX))$ has zero logarithmic derivative and is a nonzero constant. Conversely a monomial has $\mathbf{v}_\phi = \mathbf{w}_\phi = 0$, and for $\phi = bX^m(1+aX)$

$$\mathbf{u}_\phi = m + \frac{aX}{1+aX}, \quad \mathbf{v}_\phi = \frac{aX}{(1+aX)^2}, \quad \mathbf{w}_\phi = \frac{aX(1-aX)}{(1+aX)^3},$$

which satisfy the equation. $\square$

For a binomial with gap $\mathbf{g}$, $\phi = bX^m(1+aX^{\mathbf{g}})$, the same computation with $aX^{\mathbf{g}}$ in place of aX gives $\mathbf{w}_\phi^2 - \mathbf{v}_\phi^2 + 4\mathbf{v}_\phi^3 = (\mathbf{g}^2 - 1)\mathbf{v}_\phi^2$: gap one is accepted and every larger binomial gap is rejected (source 14, Appendix A).

Theorem 27.5 (Eventual strict Newton convexity; source 14). *Let $\mathbf{c}, g_2, g_3 \in \mathbb{K}$ have nonnegative valuation and the residues $\frac{1}{12}, \frac{1}{12}, -\frac{1}{216}$, and let a nonpolynomial $f = \sum_n a_n z^n \in \mathcal{E}_\Gamma(k)$ satisfy the cleared equation (24.7) with these constants. Then:*

- (i) *after passing to $\Gamma_{\mathbb{Q}}$, every break $\delta \geq \delta_0$ of f has initial polynomial $bX^m(1+aX)$ with $a, b \in k^\times$;*
- (ii) *$a_n \neq 0$ for every sufficiently large n ;*
- (iii) *the differences $\delta_n = v(a_{n+1}) - v(a_n) \in \Gamma$ are strictly increasing for all sufficiently large n and tend cofinally to $+\infty$ in Γ .*

Proof. Pass to $\Gamma_{\mathbb{Q}}$ (Lemma 26.2). The exterior form of the equation, dehomogenized, is (27.5), by the computation of Section 27.2, which uses only the residues and the nonnegative valuations of the constants; so, beyond the δ_0 of Lemma 26.5, every initial polynomial satisfies $\mathbf{w}^2 - \mathbf{v}^2 + 4\mathbf{v}^3 = 0$, and Theorem 27.4 excludes the monomial case at a break, proving (i). Beyond δ_0 , therefore, a break has exactly two active indices n and $n+1$, with no skipped index and no three-way tie, and between breaks the active index is single. By Lemma 26.3 only finitely many indices are active on a bounded interval, and a change of the active index happens only at a break (as in the proof of Lemma 26.4); so the active index increases by exactly one at each break, and by Lemma 14.5 it exceeds every bound. Every sufficiently large integer is therefore active at some scale, and its coefficient is nonzero, proving (ii). The break at which n is replaced by $n+1$ is the solution of $v(a_n) - n\delta = v(a_{n+1}) - (n+1)\delta$, namely $\delta = \delta_n$, which lies in Γ without any division. These scales increase by the monotonicity in Lemma 26.3, strictly because a repeated break would carry three active indices, and they are cofinal because an upper bound would put infinitely many active indices in one bounded interval. Nonvanishing of coefficients, inequalities between their values and cofinality in $\Gamma_{\mathbb{Q}}$ descend to Γ , since Γ is cofinal in $\Gamma_{\mathbb{Q}}$. $\square$

For $f = \mathcal{T}_p$ with μ an order unit, (23.6) gives the realization $\delta_n = (2n+1)\mu$, $\gamma_{\delta_n} = -n(n+1)\mu$ and $\phi_{\delta_n} = X^n(1+X)$ for $n \geq 0$: at δ_n the weighted value of degree j is $(j^2 - (2n+1)j)\mu$, least exactly at $j = n, n+1$. This is the Newton profile of Section 23.4, and it makes the order-three escape visible without analytic asymptotics: unbounded degrees coexist with the fixed gap one. Theorem 27.5 is a necessary condition on solutions, not a classification of the entire solutions of the theta equation: it does not force $v(a_n) = n^2\mu$, does not determine coefficients from their values, and does not assert that every eventually strictly convex value sequence occurs.

27.4 Surreal consequences and boundaries

Through (2.4), Theorem I applies to every set-sized surreal or surcomplex field $k((\omega^{-\Gamma}))$ with $\Gamma \subseteq \mathbf{No}$ and $k = \mathbb{R}$ or $\mathbb{C}$: there no nonpolynomial strongly entire series satisfies an algebraic

differential equation of order at most two, whatever the rank of Γ , and algebraic closedness of the coefficient field is not needed. For $\Gamma = \mathbb{Q}$ or $\mathbb{R}$ and $p = \omega^{-1}$ the series $\mathcal{T}_p$ of Section 25.3 has three algebraically independent jets over the field of rational functions and an explicit equation of order three. The following proposition, from source 14, states once for every series what Section 11.2, Section 25.3 and Corollary 17.1 show for particular ones.

Proposition 27.6 (No nonpolynomial series is entire on the class; source 14). *A power series $\sum_n a_n z^n$ with coefficients in $\mathbf{No}[i]$ that is strongly summable at every point of $\mathbf{No}[i]$ has only finitely many nonzero coefficients.*

Proof. Suppose infinitely many a_n are nonzero, and let $v(a)$ be the negative of the largest Conway exponent of $a \neq 0$. The values $v(a_n)$ of the nonzero coefficients and their pairwise differences form a set of surreals, so there is a positive surreal γ larger than the absolute value of every such difference. At $x = \omega^\gamma$ the leading values of the terms are $v(a_n) - n\gamma$, and along increasing indices with $a_n \neq 0$ they decrease strictly, since $v(a_{n'}) - v(a_n) < \gamma \leq (n' - n)\gamma$. They cannot lie in a well-ordered union of Hahn supports, contradicting strong summability at x . $\square$

This is a size-and-domain obstruction, not an instance of second-order rigidity: entireness is always relative to one set-sized field, $\mathcal{T}_p$ is entire on its field because μ is an order unit there, and enlarging to every surreal scale destroys that property. The zeros of $\mathcal{T}_p$ and their omnific floors and ceilings are those of Theorem 23.6 and Corollary 25.3, inherited; source 14's new contribution there is the universal lower bound for the function that produces them and the necessary Newton geometry of Theorem 27.5 for every other entire solution of its equation.

Boundaries recorded by source 14. In characteristic $\mathfrak{p} > 0$ and with an order unit μ , the series $\sum_{n \geq 0} t^{n^2 \mu} z^{pn}$ is nonpolynomial and strongly entire with derivative zero, so Theorem I fails already in order one; this is Example 17.3 with $\mathbb{Q}$ replaced by any group with an order unit. Mixed-characteristic $\mathfrak{p}$ -adic fields are not covered merely because they have characteristic zero: their residue characteristic and integer valuations differ from the hypotheses used here. The ordinary exponential $\sum z^n/n!$ solves $D_z f = f$ formally but is not strongly summable at $z = 1$, since infinitely many terms have exponent zero; allowing ordinary convergence inside a fixed coefficient layer would change the function class and need a different theorem. The proof classifies neither functions patched from local formulas nor analytic identities in the fine topology of $\mathbf{No}[i]$. It passes only to a cofinal divisible extension and never reduces the group to rank one; the no-order-unit argument uses finite rational rank of a coefficient field, not of the ambient group, and a group of high rank can have an order unit while one of countable cofinality can lack one (Examples 25.5 and 25.6).

27.5 Predecessors, repository comparison and priority

Literature. Source 14 consulted NIST DLMF §20.5 for the triple product and logarithmic derivatives [50], §23.3 for the Weierstrass differential equation [51], and §32.2 for the Painlevé equations [52]; the Mantova–Matusinski survey [17] for normal forms, not as a source for the second-order theorem; and the publisher abstract of Sebbar's paper [55], which explicitly treats classical third-order theta relations, without its full text. It also consulted the Wikipedia article on surreal numbers for orientation only. General searches for second-order Hahn and non-Archimedean rigidity returned unrelated results, including Hahn difference operators; these are not evidence of absence. No exhaustive MathSciNet or zbMATH review and no full subscription-literature review was made. For this merge the DLMF sections and the Mantova–Matusinski survey were checked,

and the bibliographic data of Sebbar’s paper were checked against its Crossref record; its content was not checked by this merge, and nothing in this part rests on it.

Repository comparison at the pin. Source 14 read the repository at `b895e86` through a connector: the root README and tree, `docs/README.md`, this report’s README and neighbouring report inventories. The formalization ledger did not return text, so no assertion about it is made, and the new proof is not represented as covered by the Lean build. It took source 13 from the author’s file library (`article(20260923-193210).tex`, baseline `4cdeaec`). Its statements were checked for this merge and are accurate at the pin, as recorded in Section 26.1. This was a targeted comparison, not an audit of every file, branch or archived draft.

Priority. The triple product, the theta–elliptic identity and third-order theta relations are classical; the Chebyshev descent of the theta series, its equation, zeros and omnific rounding are source 13’s; the no-order-unit obstruction and the finite-generation principle are the coefficient-field part’s. The proposed contribution of source 14 is universal second-order rigidity for every ordered value group and every characteristic-zero coefficient field (Theorem I(a)), with the endpoint lemma at degenerate breaks, the resulting exact minimum order (Theorem I(b)), the third-order endpoint–gap obstruction, the classification of theta initial polynomials and eventual Newton convexity, and the stated consequences for two-state systems and polynomial-denominator quotients. Whether an equivalent result exists in the literature needs a more exhaustive review; the answered question is a question of this report, not a named classical conjecture.

27.6 What the order-three-threshold part does not claim

Every limitation stated by source 14, in its manuscript, README, audit file or verification record, is kept.

- (O1) No Lean formalization, no independent referee report and no certified bibliographic priority accompany source 14; no new Lean file is presented as verified, and the repository’s build and axiom audit do not cover this part. Independent review of the exterior reduction and the endpoint lemma is the most useful next check.
- (O2) Only the question of this report is resolved; no named classical conjecture is claimed solved.
- (O3) The repository comparison at `b895e86` was targeted; the formalization ledger was not read, and no audit of every branch or archived draft was made.
- (O4) Entireness is relative to one specified Hahn field; $\mathcal{T}_p$ is not strongly entire on $\mathbf{No}[i]$, and no nonpolynomial series is (Proposition 27.6).
- (O5) D_z acts on the external variable and kills every Hahn coefficient; it is not the Berarducci–Mantova derivation.
- (O6) The theta product and the elliptic identity are classical, and the theta descent with its equation, zeros and omnific rounding and the no-order-unit obstruction are prior; source 14 reproves them for completeness and claims none. “No omnific zero” concerns the strong domain only.
- (O7) Theorem 27.5 is necessary only: it gives no sufficiency and no classification of entire solutions, does not force $v(a_n) = n^2\mu$, does not determine coefficients from their values, and does not assert that every eventually strictly convex sequence occurs.

- (O8) Corollary 27.3 is a sufficient certificate, not a decision algorithm for third-order equations; an unbounded zero set of $\Upsilon_{\mathfrak{h}_p}$ is necessary, not sufficient, for an entire solution.
- (O9) Corollary 26.11 concerns quotients $g/\mathfrak{p}$ with a polynomial denominator only; no global division theorem in a Hahn analytic category is asserted.
- (O10) Corollary 26.10 concerns rational systems; arbitrary transcendental right-hand sides are outside it.
- (O11) Nothing is claimed about local formal solutions or about classical complex Painlevé functions; the all-scale requirement, not local existence theory, does the work.
- (O12) Positive characteristic is excluded, with a counterexample, and mixed-characteristic $\mathfrak{p}$ -adic fields are not covered.
- (O13) Theorem I gives no uniform degree bound for polynomial solutions of a fixed equation (Example 26.7).
- (O14) The proof does not reduce every group to rank one; it classifies neither functions patched from local formulas nor identities in the fine topology, and a different summation convention is a different function class.
- (O15) The 600 finite checks, with the theta product and cleared equation compared modulo p^{48} (source 14's q^{48}), do not certify the quantifiers over arbitrary supports, value groups, scales or formal solutions; the classification of Theorem 27.4 is proved for rational functions, not extrapolated from the finite enumeration, and no floating-point agreement is used as evidence.
- (O16) Sebbar's paper was seen through its publisher abstract only, and broad searches are not evidence of absence; no exhaustive MathSciNet or zbMATH review was made.
- (O17) The research questions of source 14 (Questions 19.1 and 19.11–19.18) are not asserted solved.
- (O18) *[merge]* Remark 26.6, Corollary 26.13, the named break scale in the proof of Lemma 26.4, the reading of Corollaries 26.9 and 26.10 inside a differential field containing $\mathbb{K}((z))$, and the four-state observation after Corollary 26.10 are the merge's; no historical priority is claimed for them.

28 Unit-dilation rigidity: valuations of Hahn exponential polynomials

This section and the next contain the material of source 18, delivered after the order-three-threshold part had been merged. They answer Question 19.4 positively for every value group: for a nontorsion q with $v(q) = 0$, every strongly entire solution of a nonzero mixed equation (10.1) is a polynomial (Corollary 29.6). More generally, for a finite set of multipliers with no root-of-unity quotient, a nonpolynomial strongly entire solution of *some* nonzero mixed linear equation exists exactly when the absolute valuation of one quotient is an order unit (Theorem J). The missing ingredient named in the status of Question 19.4, uniform control of $v(P(n, q^n))$ for a bivariate P at unit valuation, is Corollary 28.7; it comes from a valuation form of the Skolem–Mahler–Lech theorem for exponential polynomials with arbitrary Hahn coefficients (Theorems 28.5 and 28.10). The recurrence is then closed by the escape chain of Section 3; neither the finite coefficient field of the coefficient-field part nor any nonlinear method is used.

28.1 Source, conventions and the merge’s choices

Convention 28.1 (Setting of the unit-dilation part). In Sections 28–29, k is an arbitrary field of characteristic zero with the trivial valuation, Γ is as in Convention 1.1, $\mathbb{K} = k((t^\Gamma))$, and v , lc , res , strong summability, $\text{Dom}_{\text{str}}(f)$, $\mathcal{E}_\Gamma(k)$, D_z and $\sigma_\lambda f(z) = f(\lambda z)$ are as in Conventions 14.1 and 20.1. A *unit* is an element of valuation zero, not merely a nonzero element. Every $\lambda \in \mathbb{K}$ with $v(\lambda) = 0$ has a unique expression

$$\lambda = \zeta(1 + \tau), \quad \zeta = \text{res}(\lambda) \in k^\times, \quad \tau = 0 \text{ or } v(\tau) > 0. \quad (28.1)$$

The multipliers of a mixed operator are $\lambda_1, \dots, \lambda_m \in \mathbb{K}^\times$, as the dilation parameters of Theorem E (b); q is a single dilation parameter, as in Definition 1.4.

For $k = \mathbb{C}$ one has $\mathbb{K} = K_\Gamma$ and $\mathcal{E}_\Gamma(k) = \mathcal{E}_\Gamma$. As for the four previous parts, the generality in k is recorded for this part; unlike them, this part also contains Theorem A, the negative direction of Theorem B and Theorem 10.2 as special cases over every such k (Remark 29.5).

The source. Source 18 is *Unit-Dilation Rigidity at Surreal Scales: A valuation form of Skolem–Mahler–Lech and an exact relative-scale criterion for mixed differential–dilation equations* (26 pages, A4, 23 September 2026), delivered as the archive `Surreal_Unit_Dilation_Rigidity` (batch 31, manuscript 05). It consulted the repository at

bcac55ae6fd3f2b354e568ba1d2e94d496a2cc59,

where this report’s `article.tex` is the blob 86fc4925b4c3997b984894e6b4756e327f27a4c6; its shipped file `18-unit-dilation-SOURCES_AND_SCOPE.md` lists the line ranges it read. At that pin this report was the five-source text (sources 08–12): neither the theta-descent nor the order-three-threshold part had been merged. Source 18’s description of this report was accurate there: Question 19.4 was open with an order unit, the mixed theorem at unit valuation was not asserted, and “higher-order nonlinear equations with an order unit” were open. The last point is stale now. Theorem H and Theorem I have since settled that question up to the classification of Question 19.1, and source 18’s first research question is answered negatively in the current text (Remark 29.19). Source 18 claims neither the pure differential theorem, the cyclic pure-dilation criterion, the mixed theorem at nonzero noncofinal valuation, the partial theta series, nor the escape mechanism; it credits them to this report.

Renamed symbols. The following symbols of source 18 would collide with symbols of the earlier parts. They are renamed here, in the new material only.

Source 18	Here	Reason
$K = k((t^\Gamma))$, $\mathcal{E}_\Gamma(k)$, D	$\mathbb{K}$, $\mathcal{E}_\Gamma(k)$, D_z	K_Γ is $\mathbb{C}((t^\Gamma))$ in this report.
multipliers $u_1, \dots, u_m$;	$\lambda_1, \dots, \lambda_m$;	u is the Laurent variable of Section 24.5; $\epsilon \in \Gamma$
$u = c(1 + \epsilon)$	$\lambda = \zeta(1 + \tau)$	below; c is a coefficient.
$A(n) = \sum_j P_j(n)u_j^n$; P_j	$\mathcal{Y}(n) = \sum_j R_j(n)\lambda_j^n$;	P is a differential polynomial; R_h plays the same
	R_j	role in Lemma 7.1.
$A_j(n)$ (recurrence	$c_j(n)$, $R_{j,\ell}$	As in Lemma 3.1.
coefficients), $Q_{j,\ell}$		
$Q_{j,\gamma}$ (coefficient	$r_{j,\gamma}$; Σ	T is an indeterminate.
polynomials); envelope T		

Source 18	Here	Reason
forward shift S	Sh	S is a support alphabet, as here.
period M ; residue class r	M ; b	$M(S)$ is the positive monoid; r is a derivative order.
slopes $\lambda_r = v(u_j)$	ϖ_b	λ now names a multiplier.
relative scale μ	$\text{rs}(\lambda_1, \dots, \lambda_m)$	μ is an order unit or a positive scale, with $p = t^\mu$.
noncofinality witness η	γ of (2.2)	η is $v(\tau)$ in Theorem 4.4.
operator coefficients $c_{\ell,r,k}$, z^k ; output index m	$\varkappa_{\ell,r,i}$, z^i ; $\bar{n}$	k is the coefficient field; m counts multipliers.
right side h , $\deg h \leq e$; binomial index h	$\mathbf{h}$, $\deg \mathbf{h} \leq \bar{e}$; i	h is the order of a root of unity, as in Theorem 4.4.
Θ_q , q with $v(q)$ an order unit	Θ_p , p	(8.1); q is kept for a dilation parameter.
order unit H ; H_* ; $F = \sum t^{n^2 H} z^n$; $G = \sum t^{n^2 H} z^{hn}$	μ ; e_* ; G_μ ; $G_\mu(z^h)$	F is (11.2); G is the series of Corollary 17.1, which is G_1 over $\mathbb{C}((t^\mathbb{Q}))$.
Δ (maximal proper convex subgroup)	Δ^{top}	Δ_β is the least convex subgroup containing β .
field F of its Definition 3.1	K_0	F is (11.2); K_0 as in Theorem 21.2.
Theorems 4.2, 4.7, 6.4, 7.2	Theorems 28.5, 28.10, 29.4, 29.9	Theorems A–N of this report are different theorems; Theorem J collects the last two.

The shipped file `18-unit-dilation-SOURCES_AND_SCOPE.md` keeps the source's notation.

Words with two meanings. A *unit* is a valuation-zero element (Convention 28.1), and a *unit dilation* is σ_q with $v(q) = 0$. An *order unit* is Definition 1.3; the two words are unrelated. The *relative scale* of a list of multipliers is $\text{rs}(\lambda_1, \dots, \lambda_m) = \max_{i,j} |v(\lambda_i) - v(\lambda_j)|$ (Definition 29.3); it is *noncofinal* if it is 0 or not an order unit. *Nondegenerate* describes an exponential polynomial none of whose base quotients is a root of unity; it has nothing to do with the nondegenerate corners of Section 16. A *root-of-unity quotient* concerns the actual multipliers λ_i/λ_j , not their residues; the residues may be roots of unity (Example 4.6).

Results printed once. Source 18 reproves several facts the report already contains. They are printed once and cited: its definition of strong summability and entireness (Definition 1.2); its Lemma 2.2 (sums of well-ordered sets) and Lemma 2.4 (the positive monoid), which are Lemma 2.1 (1) and (2), except that its proof of the finite-word lemma is printed below as Lemma 28.2, since this report imports that lemma from Higman; its Lemmas 2.5 and 2.6, which are Lemma 2.3 (a) and Lemma 2.2; the invariance of $\mathcal{E}_\Gamma(k)$ under σ_λ and D_z , which is Proposition 2.7; its Lemma 3.2 and Corollary 3.4 (polynomial–exponential independence, finitely many zeros of a nonzero nondegenerate exponential polynomial, also on every arithmetic progression), which are Lemma 21.1 and its proof; its Theorem 3.3, the classical Skolem–Mahler–Lech theorem, imported there as here (Section 21.1); its Example 4.8, which is the third example of Remark 4.3; its Section 5 (Theorem 5.1, Corollary 5.2, Proposition 5.3), which is Lemma 3.1, Theorem 3.3 and Corollary 3.4, the last with the closed region $v(x) \leq -B^+$ where source 18 states the open one $v(x) < -B$; the entireness half of its Lemma 7.1, which is Theorem 8.1 by the route of Remark 8.2; the embedding (2.4); its parity example (Section 9.1), which is Example 4.6; its torsion example (Section 9.5), which is the periodic-support construction of Section 9; its positive-characteristic series, which is Example 17.3; and its remark on $\Gamma = 0$, which is in Section 12.

Where each numbered result went. Lemma 2.3 is Lemma 28.2; Lemma 3.5 is Lemma 28.3; Lemma 4.1, Theorem 4.2, Remark 4.3, Corollary 4.4 and Remark 4.5 are Lemma 28.4, Theorem 28.5, Remark 28.6, Corollary 28.7 and Remark 28.8; Lemma 4.6 and Theorem 4.7 are Lemma 28.9 and Theorem 28.10; Lemma 6.1, Corollary 6.2, Definition 6.3, Theorem 6.4, Corollaries 6.5 and 6.6 are Lemma 29.1, Corollary 29.2, Definition 29.3, Theorem 29.4 and Corollaries 29.6 and 29.7; Lemma 7.1, Theorem 7.2, Remark 7.3, Corollary 7.4 and Proposition 7.5 are Lemma 29.8, Theorem 29.9, Remark 29.10, Corollary 29.11 and Proposition 29.12; Corollary 8.1 is Corollary 29.13; Proposition 10.1 is Proposition 29.18; Research question 11.1 is recorded in Remark 29.19, and Research questions 11.2–11.12 are Questions 19.19–19.29. Theorem J in the introduction combines Theorems 6.4 and 7.2.

The merge’s choices. The review of source 18 for this merge found its proofs correct. The merge adds four points, each marked. First, Remark 29.5 records that Theorem 29.4 contains Theorem A, the negative direction of Theorem B and Theorem 10.2, over every characteristic-zero k , and compares it with Theorem E(b). Second, the exterior region of Theorem 29.4 is closed, by Corollary 3.4. Third, Remark 29.19 records the answer in the current text to source 18’s nonlinear question. Fourth, Corollary 29.11 is read beside Theorem B: adding derivatives creates no new existence case. Source 18’s theorems, proofs, examples and non-claims are otherwise printed as delivered, in the renamed notation.

28.2 Finite words and exponential polynomials

Lemma 2.1 (2) imports the finite-word lemma from Higman [6]. Source 18 proves the case it needs, by the usual minimal-bad-sequence argument; the proof is printed here because it makes the positive-monoid lemma self-contained. It is standard, not new.

Lemma 28.2 (Finite-word lemma for a well-ordered alphabet; source 18). *Let S be well ordered, and order finite words over S by subsequence embedding with coordinatewise increase of letters. Every infinite sequence of words contains an earlier word that embeds in a later one.*

Proof. Suppose a bad sequence exists, that is, one with no such pair. Choose a bad sequence $w_0, w_1, \dots$ so that, at each position, w_n has the least possible length among the choices that allow a bad continuation of the prefix already chosen. No word is empty. Write $w_n = w'_n a_n$ with a_n its last letter, and choose $i_0 < i_1 < \dots$ with $a_{i_0} \leq a_{i_1} \leq \dots$, which is possible in a well-ordered set. By the minimality at position i_0 the sequence $w_0, \dots, w_{i_0-1}, w'_{i_0}, w'_{i_1}, \dots$ is not bad. An embedding between two words of the prefix contradicts badness of the original sequence; an embedding of a prefix word into some w'_{i_j} is an embedding into w_{i_j} ; and an embedding $w'_{i_j} \hookrightarrow w'_{i_l}$ with $j < l$ extends, by $a_{i_j} \leq a_{i_l}$, to $w_{i_j} \hookrightarrow w_{i_l}$. Every case is a contradiction. $\square$

With it, the proof of Lemma 2.1 (2) needs no import: a strictly decreasing sequence of word sums would give an embedding of an earlier word into a later one, which cannot decrease the sum; and if infinitely many distinct nondecreasingly sorted words had one sum, an embedding between two of them would force equal lengths, since inserted letters are positive, and then equal letters, since the sums agree.

An *exponential polynomial* over a field K_0 of characteristic zero is a sequence $\mathcal{V}(n) = \sum_{j=1}^m R_j(n) \lambda_j^n$ with $R_j \in K_0[U]$ and $\lambda_j \in K_0^\times$; it is *nondegenerate* if the bases are distinct and no quotient λ_i/λ_j , $i \neq j$, is a root of unity. A constant-coefficient recurrence annihilates it (a product of powers of

Sh $-\lambda_j$, with Sh the forward shift), so the Skolem–Mahler–Lech theorem applies over k and over $\mathbb{K}$. Source 18 imports it in the characteristic-zero form of Lech [28] and Bell’s account [29]; the report imports it through Bell’s Theorem 1.1 [27]. Its consequence used below is Lemma 21.1: a nonzero nondegenerate exponential polynomial has only finitely many zeros, and so has its restriction to every arithmetic progression, the restriction being again nonzero and nondegenerate; the proof of that lemma also shows that polynomial–exponential sequences with distinct bases are linearly independent on every tail.

Lemma 28.3 (One period removes all residue torsion; source 18). *For $\zeta_1, \dots, \zeta_m \in k^\times$ let M be the least common multiple of the orders of those quotients ζ_i/ζ_j that are roots of unity ($M = 1$ if there are none), or any multiple of it. After equal bases are combined, every sequence*

$$l \mapsto \sum_j r_j(Ml + b) \zeta_j^{Ml+b}, \quad r_j \in k[U],$$

is either identically zero or nondegenerate. The same M serves for every choice of the polynomials r_j and every b .

Proof. The bases in l are ζ_j^M . If $(\zeta_i/\zeta_j)^M$ is a root of unity, so is ζ_i/ζ_j , whose order then divides M ; so ζ_i^M and ζ_j^M are equal or have a nontorsion quotient. Combine equal bases. $\square$

The uniformity in the last sentence is essential below. There may be infinitely many Hahn coefficient sequences to consider, but they all use the same finite list of residue bases: the proof never takes a maximum of infinitely many unrelated exceptional zero sets or periods.

28.3 The valuation theorem

Lemma 28.4 (A common support envelope for unit orbits; source 18). *Let $R_j \in \mathbb{K}[U]$ and let $\lambda_j = \zeta_j(1 + \tau_j)$ be units, $1 \leq j \leq m$. There is one well-ordered set $\Sigma \subseteq \Gamma$ containing the support of*

$$\mathcal{Y}(n) = \sum_j R_j(n) \lambda_j^n$$

for every $n \geq 0$, and for each $\gamma \in \Sigma$ there are polynomials $r_{j,\gamma} \in k[U]$ with

$$[t^\gamma] \mathcal{Y}(n) = \sum_j \zeta_j^n r_{j,\gamma}(n) \quad (n \geq 0). \quad (28.2)$$

Proof. Let C be the union of the supports of the coefficients of all R_j and S the union of the supports of the nonzero τ_j : a finite union of well-ordered sets, and a well-ordered subset of $\Gamma_{>0}$. If all R_j vanish take $\Sigma = \emptyset$; otherwise $\Sigma = C + M(S)$, well ordered by Lemma 2.1. For $n \geq 0$, $(1 + \tau_j)^n = \sum_{i=0}^n \binom{n}{i} \tau_j^i$, and every resulting support lies in Σ . Fix $\gamma \in \Sigma$ and write $R_j(U) = \sum_a \rho_{j,a} U^a$. The coefficient in question is

$$\sum_{j,a,i} \zeta_j^n n^a \binom{n}{i} [t^\gamma] (\rho_{j,a} \tau_j^i). \quad (28.3)$$

Although i is displayed as unbounded, only finitely many i and finitely many support decompositions contribute at the fixed γ , by Lemma 2.1 (1) and (2), and the finite list does not depend on n . Each $U^a \binom{n}{i}$ is a polynomial over k , since $i!$ is invertible, and $\binom{n}{i} = 0$ for $i > n$, so extending the sum in i changes no value at a nonnegative integer. Collecting the finitely many polynomials gives (28.2). $\square$

No bound on $\deg r_{j,\gamma}$ uniform in γ is asserted, and none need hold. The period of Lemma 28.3, by contrast, does not depend on those degrees.

Theorem 28.5 (Eventual periodicity at unit valuation; source 18). *In Lemma 28.4, choose M for the residues ζ_j by Lemma 28.3. For each $b \in \{0, \dots, M-1\}$ exactly one of the following holds:*

- (i) $\mathcal{Y}(Ml + b) = 0$ for every $l \geq 0$;
- (ii) there is $\beta_b \in \Gamma$ with $v(\mathcal{Y}(Ml + b)) = \beta_b$ for all sufficiently large l .

If the representation is nonzero and no quotient λ_i/λ_j of the actual multipliers is a root of unity, only (ii) occurs; then $\mathcal{Y}(n) \neq 0$ for all large n , and its eventual valuations lie in a finite subset of Γ .

Proof. Fix b . For every $\gamma \in \Sigma$ the restriction of (28.2) to $n = Ml + b$ is the zero sequence or a nondegenerate exponential polynomial (Lemma 28.3), and in the second case it has only finitely many zeros (Lemma 21.1). If every such coefficient sequence is identically zero, (i) holds. Otherwise the set of $\gamma \in \Sigma$ for which $l \mapsto [t^\gamma]\mathcal{Y}(Ml + b)$ is not identically zero has a least element β_b . Every coefficient below β_b vanishes *identically*, at every index of the progression, not merely eventually; at β_b there are finitely many exceptional zeros, and beyond them β_b is exactly the least exponent of $\mathcal{Y}(Ml + b)$.

For the last assertion, the restriction of $\mathcal{Y}$ to a progression has the bases λ_j^M and the nonzero polynomial factors $\lambda_j^b R_j(MU + b)$. The hypothesis on the actual quotients makes the bases distinct, and the independence statement in the proof of Lemma 21.1, applied over $\mathbb{K}$, excludes an identically zero restriction. The finitely many progressions have one common tail. $\square$

Remark 28.6 (Why a naive coefficientwise argument would be invalid; source 18). It would not be enough to say that every coefficient sequence is eventually stable and then to take a maximum of infinitely many exceptional indices: such a maximum need not exist. The proof selects the least coefficient sequence that is not an identity; all earlier ones vanish at *every* index of the progression, and only one finite exceptional set is used on each progression.

Corollary 28.7 (The bivariate unit-orbit estimate; source 18). *Let $q \in \mathbb{K}^\times$ have $v(q) = 0$ and infinite multiplicative order. For every nonzero $P(U, X) \in \mathbb{K}[U, X]$, $P(n, q^n)$ is eventually nonzero and $v(P(n, q^n))$ is eventually periodic. If $\text{res}(q)$ is not a root of unity the valuation is eventually constant; if $\text{res}(q)$ has finite order h , a period dividing h suffices.*

Proof. Write $P(U, X) = \sum_j P_j(U)X^j$. The multipliers q^j with $P_j \neq 0$ have no root-of-unity quotient, since q is nontorsion, and their residues are powers of $\text{res}(q)$. If $\text{res}(q)$ is not a root of unity no nontrivial residue quotient is one, and $M = 1$; if it has order h , $M = h$ is admissible in Lemma 28.3. Apply Theorem 28.5. $\square$

This is the estimate named in the status of Question 19.4. It extends Theorem 4.4, which treats $P(q^n)$ for a univariate P , to the bivariate orbit $P(n, q^n)$; the univariate case of part (b) there has the elementary proof printed there, and the parity example of Example 4.6 shows that periodicity, not constancy, is the right conclusion.

Remark 28.8 (An elementary subcase; source 18). When $\text{res}(q)$ has finite order h , every coefficient sequence on a class modulo h in the preceding proof is an ordinary polynomial in the new index times a nonzero constant, and the only zero-set input needed is the finiteness of the roots of a nonzero polynomial. The full Skolem–Mahler–Lech theorem is needed for general residue bases, not for this subcase. The actual q must still be nontorsion, to exclude a zero progression.

Lemma 28.9 (Eventually ordered affine functions; source 18). *For finitely many functions $n \mapsto \beta_i + n\varpi_i$ into an ordered abelian group, the pairwise comparisons stabilize as n runs through $\mathbb{N}$. If the slopes ϖ_i are pairwise distinct, all comparisons are eventually strict and one index is the unique eventual minimum. The same holds along any fixed arithmetic progression.*

Proof. A difference $\beta + n\varpi$ has constant sign if $\varpi = 0$, and is strictly monotone, crossing zero at most once, otherwise. Choose one tail on which all finitely many comparisons have stabilized. On a progression the slope becomes a positive integer multiple and only the intercept changes. $\square$

This is the argument inside Lemma 4.2, stated for an arbitrary finite list; like that lemma it uses no Archimedean comparison.

Theorem 28.10 (Periodic-affine valuations of Hahn exponential polynomials; source 18). *Let $R_j \in \mathbb{K}[U]$, $\lambda_j \in \mathbb{K}^\times$ and $\mathcal{Y}(n) = \sum_j R_j(n)\lambda_j^n$. There is $M \geq 1$ such that for each class b modulo M either*

$$\mathcal{Y}(n) = 0 \quad \text{for every } n \geq 0 \text{ with } n \equiv b \pmod{M}, \quad (28.4)$$

or there are $\beta_b \in \Gamma$ and $\varpi_b \in \{v(\lambda_1), \dots, v(\lambda_m)\}$ with

$$v(\mathcal{Y}(n)) = \beta_b + n\varpi_b \quad \text{for all sufficiently large } n \equiv b \pmod{M}. \quad (28.5)$$

If the representation is nonzero and no quotient λ_i/λ_j is a root of unity, no zero progression occurs; in particular $\mathcal{Y}(n)$ is eventually nonzero. M may be chosen from the residues of the normalized multipliers $t^{-v(\lambda_j)}\lambda_j$ alone, independently of the coefficients of the R_j .

Proof. Put $\hat{\lambda}_j = t^{-v(\lambda_j)}\lambda_j$, a unit, and group the indices with equal valuation:

$$\mathcal{Y}(n) = \sum_{\varpi} t^{n\varpi} \mathcal{Y}_{\varpi}(n), \quad \mathcal{Y}_{\varpi}(n) = \sum_{j: v(\lambda_j)=\varpi} R_j(n)\hat{\lambda}_j^n.$$

Choose one period M for all residues $\text{res}(\hat{\lambda}_j)$ by Lemma 28.3; it is admissible for each group. On a class b , Theorem 28.5 makes every $\mathcal{Y}_{\varpi}$ either identically zero there or of eventually constant valuation $\beta_{\varpi,b}$. If all groups vanish identically on the class, (28.4) holds. Otherwise discard the zero groups; the rest have, on a common tail of the class, the valuations $n\varpi + \beta_{\varpi,b}$ with pairwise distinct slopes, and Lemma 28.9 gives a unique least one on a further tail. A unique least-valuation summand cannot cancel, which is (28.5). The assertion without zero progressions follows from the independence statement of Lemma 21.1 over $\mathbb{K}$ on a progression, as in Theorem 28.5. $\square$

The smallest slope need not win eventually: in the third example of Remark 4.3, which is source 18's Example 4.8, the intercept e_1 dominates every multiple of the slope e_0 . The proof uses the eventual winner of finitely many affine functions, not the rule “take the smallest slope”.

What is, and is not, effective (source 18). The theorem is an existence statement for a finite valuation description. Given arbitrary Hahn coefficients it provides no algorithm for the first coefficient sequence that is not an identity, for the eventual winning indices, or for the last exceptional zero. In the finite-order residue case the relevant zero sets are polynomial roots, but selecting the coefficients still needs exact support information. No decision procedure for arbitrary surreal inputs follows.

29 Mixed rigidity and the relative-scale dichotomy

29.1 Operator normal form and its recurrence

The operators are D_z and σ_λ , with $D_z \sigma_\lambda = \lambda \sigma_\lambda D_z$. Write a finite operator as

$$L = \sum_{\ell=1}^m \sum_{r=0}^R p_{\ell,r}(z) D_z^r \sigma_{\lambda_\ell}, \quad p_{\ell,r}(z) = \sum_{i=0}^{d_L} \varkappa_{\ell,r,i} z^i \in \mathbb{K}[z], \quad (29.1)$$

with $\lambda_\ell \neq 0$ and some $\varkappa_{\ell,r,i} \neq 0$; $D_z^r \sigma_\lambda$ is kept in that order, not silently replaced by $\sigma_\lambda D_z^r$. Since $D_z^r \sigma_\lambda = \lambda^r \sigma_\lambda D_z^r$, the terms are the jets $\lambda^r \mathcal{J}_{\lambda,r} f$ of (20.1), and (10.1) is the case $\lambda_\ell = q^\ell$.

Lemma 29.1 (Exact shift polynomials; source 18). *Let $f = \sum_n a_n z^n$. For an output degree $\bar{n} \geq d_L$, the term $\varkappa_{\ell,r,i} z^i D_z^r \sigma_{\lambda_\ell} f$ contributes*

$$\varkappa_{\ell,r,i} (\bar{n} + r - i)_{\underline{r}} \lambda_\ell^{\bar{n}+r-i} a_{\bar{n}+r-i} \quad (29.2)$$

to $[z^{\bar{n}}] Lf$. Let s_0 be the least active shift $r - i$ and put $n = \bar{n} + s_0$. After grouping equal coefficient indices, the high coefficients of $Lf = \mathbf{h}$ read

$$c_0(n) a_n + \sum_{j=1}^J c_j(n) a_{n+j} = [z^{n-s_0}] \mathbf{h}, \quad (29.3)$$

with

$$c_j(n) = \sum_{\ell=1}^m \lambda_\ell^n R_{j,\ell}(n), \quad (29.4)$$

$$R_{j,\ell}(U) = \lambda_\ell^j \sum_{r-i=s_0+j} \varkappa_{\ell,r,i} (U+j)_{\underline{r}}. \quad (29.5)$$

The active j run from 0 to a finite J ; for each active j some $R_{j,\ell}$ is nonzero, in particular some $R_{0,\ell}$. The right side vanishes once $\bar{n} > \deg \mathbf{h}$.

Proof. The monomial $a_N z^N$ becomes $a_N \lambda_\ell^N (N)_{\underline{r}} z^{N-r}$ under $D_z^r \sigma_{\lambda_\ell}$, and multiplication by z^i gives degree $N - r + i$; setting it equal to $\bar{n}$ gives (29.2), and $\bar{n} \geq d_L$ ensures $N \geq r$. With $r - i = s_0 + j$ one has $N = n + j$, which gives (29.4)–(29.5). For fixed j and ℓ distinct contributing r give the monic polynomials $(U+j)_{\underline{r}}$ of distinct degrees, so a nonempty combination with nonzero coefficients is nonzero, and for fixed r and j there is only one i . $\square$

For $\lambda_\ell = q^\ell$ this is the conversion (10.2) of Theorem 10.2, grouped by multiplier instead of by powers of X .

Corollary 29.2 (No hidden zero operator; source 18). *If the λ_ℓ are distinct, a normal form (29.1) with some nonzero coefficient is a nonzero operator on $\mathbb{K}[[z]]$.*

Proof. At the least active shift, (29.4) is a nonzero polynomial–exponential representation with distinct bases, so $c_0(n) \neq 0$ for arbitrarily large n by the independence statement of Lemma 21.1; for such n the coefficient of degree $n - s_0$ in $L(z^n)$ is $c_0(n) \neq 0$. $\square$

29.2 Relative scales and rigidity

Definition 29.3 (Relative scale; source 18). For a nonempty finite list of multipliers put

$$\text{rs}(\lambda_1, \dots, \lambda_m) = \max_{i,j} |v(\lambda_i) - v(\lambda_j)| \in \Gamma_{\geq 0}.$$

It is 0 for one multiplier. It is *noncofinal* if it is 0 or is not an order unit.

If $\text{rs} > 0$ is not an order unit, (2.2) supplies $\gamma > 0$ with $n \text{rs} < \gamma$ for every $n \geq 0$. For a finite list, noncofinality of every $|v(\lambda_i/\lambda_j)|$ is equivalent to noncofinality of their maximum.

Theorem 29.4 (Rigidity for noncofinal relative scales; source 18). *Suppose no quotient λ_i/λ_j , $i \neq j$, is a root of unity and $\text{rs}(\lambda_1, \dots, \lambda_m)$ is noncofinal. Let L be a nonzero normal form (29.1). For every $\bar{e} \in \mathbb{N}$ there are $N \geq 1$ and $B \in \Gamma_{\geq 0}$ such that, for every $\mathbf{h} \in \mathbb{K}[z]$ with $\deg \mathbf{h} \leq \bar{e}$ and every formal solution of $Lf = \mathbf{h}$:*

- (i) *strong evaluation at one $x \neq 0$ with $v(x) < -B$ forces $\deg f < N$;*
- (ii) *every strongly entire solution is a polynomial of degree less than N ;*
- (iii) *every nonpolynomial formal solution fails strong evaluation throughout that common exterior region.*

The polynomial conclusion also holds for rational operator coefficients and a rational right side, read as an identity of formal Laurent series. By Corollary 3.4 the region in (i) and (iii) may be taken closed, $v(x) \leq -B$ (merge).

Proof. Apply Lemma 29.1. For each active j , $c_j(n)$ is a nonzero exponential polynomial with bases among the λ_ℓ , so Theorem 28.10 gives one period M and one tail on which, for $n \equiv b \pmod{M}$,

$$v(c_j(n)) = \beta_{j,b} + n\varpi_{j,b}, \quad \varpi_{j,b} \in \{v(\lambda_1), \dots, v(\lambda_m)\}, \quad (29.6)$$

and all these $c_j(n)$, in particular $c_0(n)$, are nonzero there. Inactive j give identically zero coefficients and are discarded; if no $j \geq 1$ is active the recurrence terminates directly. Otherwise let $B_0 \geq 0$ bound the finitely many differences $\beta_{0,b} - \beta_{j,b}$, $j \geq 1$. Then

$$v(c_0(n)) - v(c_j(n)) = \beta_{0,b} - \beta_{j,b} + n(\varpi_{0,b} - \varpi_{j,b}) \leq B_0 + n \text{rs}(\lambda_1, \dots, \lambda_m).$$

If the relative scale is 0 put $B = B_0$; otherwise put $B = B_0 + \gamma$ with γ from (2.2). This is a uniform cost bound. Enlarge N past the finite valuation exceptions and so that $n - s_0 \geq d_L$ and $n - s_0 > \bar{e}$. Lemma 3.1 with Theorem 3.3 and Corollary 3.4 proves (i)–(iii): a nonzero a_n with $n \geq N$ starts an escape chain whose evaluated leading exponents do not increase at such x . For rational data, multiply the equation on the left by a common nonzero polynomial denominator: the operator stays a nonzero normal form, the right side becomes a polynomial, and no denominator is differentiated. $\square$

Remark 29.5 (What the theorem contains; merge). With $m = 1$ and $\lambda_1 = 1$ the relative scale is 0, and Theorem 29.4 is Theorem A, with its uniform exterior obstruction and degree bound, over every characteristic-zero k . With $R = 0$ and $\lambda_\ell = q^\ell$ it is the negative direction of Theorem B: the relative scale is a multiple of $|v(q)|$, and is noncofinal exactly when (ii) there fails. With $\lambda_\ell = q^\ell$ and $v(q) \neq 0$ noncofinal it is Theorem 10.2. The proofs of this report reach those three theorems through Theorem 4.4, Lemma 4.2 and Lemma 10.1; source 18 reaches all of them, and the unit case,

through Theorem 28.10. The comparison with Theorem E (b) is this: that theorem excludes every algebraic relation, linear or not, among the mixed jets when Γ has no order unit, which is then the case of every list of multipliers; Theorem 29.4 excludes only linear relations, but for every Γ with a noncofinal relative scale, with a uniform degree bound and exterior region. Neither contains the other.

Corollary 29.6 (Mixed equations at unit valuation; source 18). *Let $q \in \mathbb{K}^\times$ be nontorsion with $v(q) = 0$. Every strongly entire solution of a nonzero equation $\sum_{r,\ell} p_{r,\ell}(z) D_z^r \sigma_q^\ell f(z) = h(z)$ with $p_{r,\ell}, h \in \mathbb{K}[z]$ is a polynomial, for every Γ . Rational coefficients are allowed as in Theorem 29.4, and each fixed equation has a uniform degree bound and a uniform exterior obstruction for its nonpolynomial formal solutions.*

Proof. The distinct multipliers q^ℓ that occur have nontorsion quotients and valuation zero, so the relative scale is 0; apply Theorem 29.4. Negative powers of q change nothing. $\square$

For $h = 0$ and $k = \mathbb{C}$ this is Question 19.4, answered positively, with no order-unit hypothesis. The decisive estimate is Corollary 28.7; no coefficient-field theorem and no nonlinear method is used.

Corollary 29.7 (Mixed linear independence; source 18). *Let $f \in \mathcal{E}_\Gamma(k)$ be nonpolynomial. For every finite list of multipliers with no root-of-unity quotient and a noncofinal relative scale, the family $\{D_z^r \sigma_{\lambda_j} f : 1 \leq j \leq m, r \geq 0\}$ is linearly independent over $\mathbb{K}(z)$; equivalently, so is $\{\mathcal{J}_{\lambda_j, r} f\}_{j,r}$.*

Proof. Clearing denominators in a nontrivial relation gives a nonzero normal form annihilating f , and Theorem 29.4 makes f a polynomial. The nonzero scalars λ_j^r do not affect independence. $\square$

This is linear independence of formal functions over a field of rational functions. It is not algebraic independence of values at a point, and it does not exclude polynomial relations among these functions; for those see Theorem E (b) without an order unit and Corollary 25.2 with one.

29.3 The dichotomy for a finite set of multipliers

Lemma 29.8 (A mixed equation for partial theta; source 18). *If $p \in \mathbb{K}^\times$ and $v(p) > 0$ is an order unit, then Θ_p of (8.1) is a nonpolynomial element of $\mathcal{E}_\Gamma(k)$ and satisfies*

$$D_z \Theta_p - \sigma_p \Theta_p - z D_z \sigma_p \Theta_p = 0. \quad (29.7)$$

Proof. Entireness and nonpolynomiality are Theorem 8.1, whose order-unit half source 18 proves by the route of Remark 8.2. Differentiate (8.2), $\Theta_p(z) = 1 + z \Theta_p(pz)$, with p constant for D_z : the product rule gives $D_z \Theta_p = \sigma_p \Theta_p + z D_z(\sigma_p \Theta_p)$. $\square$

Theorem 29.9 (Exact relative-scale dichotomy; source 18). *Let $\lambda_1, \dots, \lambda_m \in \mathbb{K}^\times$ with no quotient λ_i/λ_j , $i \neq j$, a root of unity. The following are equivalent.*

- (i) *There are a nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ and a nonzero operator (29.1) with $Lf = 0$.*
- (ii) *$|v(\lambda_i/\lambda_j)|$ is an order unit for some i, j .*
- (iii) *There are such f and L using only two of the λ_ℓ and derivatives of order at most one.*

If (ii) fails, every nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ has the independence property of Corollary 29.7 for the whole list.

Proof. If (ii) fails the relative scale is noncofinal and Theorem 29.4 excludes (i). Assume (ii) and order the pair so that $p = \lambda_j/\lambda_i$ has positive order-unit valuation. Put $f(z) = \Theta_p(z/\lambda_i)$, nonpolynomial and strongly entire since dilation preserves $\mathcal{E}_\Gamma(k)$. Then $\sigma_{\lambda_i}f = \Theta_p$ and $\sigma_{\lambda_j}f = \sigma_p\Theta_p$, and (29.7) becomes

$$(D_z\sigma_{\lambda_i} - \sigma_{\lambda_j} - zD_z\sigma_{\lambda_j})f = 0, \quad (29.8)$$

a nonzero normal form by Corollary 29.2. This is (iii), and (iii) $\Rightarrow$ (i) is immediate. $\square$

Proof of Theorem J. Part (a) is Theorem 29.4, part (b) is Theorem 29.9, and the last assertion is Corollary 29.6. $\square$

Remark 29.10 (The quantifiers matter; source 18). Theorem 29.9 classifies the existence of *at least one* nonzero mixed equation supported in a prescribed finite set of multipliers. It does not say that every operator using a cofinal quotient has a nonpolynomial entire solution, it does not decide the solvability of a specified equation, and it does not classify simultaneous solutions of several independently prescribed equations. The positive construction may give zero coefficients to all but two of the listed multipliers. This is the precise sense in which it bears on Question 19.6.

Corollary 29.11 (Cyclic mixed equations; source 18). *Let $q \in \mathbb{K}^\times$ be nontorsion. A nonpolynomial strongly entire solution of some nonzero polynomial-coefficient equation in D_z and nonnegative powers of σ_q exists if and only if $v(q) \neq 0$ and $|v(q)|$ is an order unit. In the negative case every such equation, also with polynomial forcing, has only polynomial strongly entire solutions. The same holds with negative powers of σ_q admitted.*

Proof. The valuation differences of a finite list of powers of q are integer multiples of $v(q)$, and a nonzero integer multiple of a positive element is an order unit exactly when the element is. If $v(q) = 0$ or $|v(q)|$ is noncofinal, Theorem 29.4 applies to every finite list. In the positive case apply Theorem 29.9 to $\{1, q\}$; its equation uses only σ_1 and σ_q , whatever the sign of $v(q)$. $\square$

With derivatives removed, the existence half is Theorem B, whose witness is a pure dilation equation. So adding finite derivative orders creates *no new existence case* (merge's reading): the criterion of Theorem B is also the criterion for mixed equations in D_z and σ_q , and the unit case missing before this part is now included.

Why a common absolute scale does not matter (source 18). Replacing every λ_j by $c\lambda_j$ with one $c \neq 0$ gives $\sum p_{j,r}D_z^r\sigma_{c\lambda_j}f = \sum p_{j,r}D_z^r\sigma_{\lambda_j}(\sigma_cf)$, and σ_c is a bijection of $\mathcal{E}_\Gamma(k)$ preserving polynomiality, so the existence problem is unchanged. An absolute-scale criterion would therefore be wrong: one multiplier of arbitrarily large valuation gives an ordinary differential equation for the entire series $\sigma_\lambda f$ and cannot by itself produce a nonpolynomial strongly entire solution.

Proposition 29.12 (The top Archimedean scale; source 18). *Suppose Γ has an order unit and put $\Delta^{\text{top}} = \{\gamma \in \Gamma : \gamma = 0 \text{ or } |\gamma| \text{ is not an order unit}\}$. Then Δ^{top} is the unique maximal proper convex subgroup of Γ and $\Gamma/\Delta^{\text{top}}$ is Archimedean. The negative case of Theorem 29.9 holds exactly when all $v(\lambda_j)$ have the same image in $\Gamma/\Delta^{\text{top}}$.*

Proof. For $\gamma_1, \gamma_2 \in \Delta^{\text{top}}$ not both zero, $|\gamma_1 + \gamma_2| \leq 2\rho$ with $\rho = \max(|\gamma_1|, |\gamma_2|)$ not an order unit, and 2ρ is then not one either; negatives and convexity are immediate. An order unit lies outside, so the subgroup is proper, and a convex subgroup containing an order unit is Γ , since every element lies between two integer multiples of it; so every proper convex subgroup lies in Δ^{top} . For positive

classes $\bar{\xi}_1, \bar{\xi}_2$ of the quotient a positive representative ξ_1 is an order unit, and $n\xi_1 > \xi_2$ for some n . Finally $v(\lambda_i) - v(\lambda_j) \in \Delta^{\text{top}}$ says exactly that its absolute value is not an order unit. $\square$

These ordered-group facts are elementary background, not a new classification. For an order unit μ , Δ^{top} is the kernel $\{\gamma : n|\gamma| < \mu \text{ for all } n \geq 1\}$ of the real-valued coarsening cited in Remark 2.6. The interpretation: under the no-root-of-unity restriction, a nonpolynomial strongly entire mixed relation exists exactly when the multipliers are separated at the top Archimedean scale.

29.4 Surreal, surcomplex and omnific coefficients

Through (2.4) every statement above is a statement about series with surreal or surcomplex coefficients in one fixed set-sized workspace $K_\Gamma \subseteq \mathbf{No}[i]$, with $\Gamma \subseteq \mathbf{No}$.

Corollary 29.13 (A surcomplex no-annihilator theorem; source 18). *Let $\Gamma \subseteq \mathbf{No}$ be a nonzero set-sized subgroup, $f \in K_\Gamma[[z]]$ nonpolynomial and strongly entire on K_Γ , and $q \in K_\Gamma^\times$ nontorsion with $v(q) = 0$. There is no nonzero finite relation $\sum_{r,\ell} R_{r,\ell}(z)(D_z^r f)(q^\ell z) = 0$ with $R_{r,\ell} \in K_\Gamma(z)$. The same holds for every finite list of multipliers with a noncofinal relative scale and no root-of-unity quotient.*

Proof. Corollary 29.7; the factors $q^{r\ell}$ relating $D_z^r \sigma_q^\ell f$ to $(D_z^r f)(q^\ell z)$ are absorbed in the rational functions. $\square$

An example with an order unit (source 18). Let $\Gamma = \mathbb{Q}\mu \oplus \mathbb{Q}\epsilon$ with $\mu > n\epsilon > 0$ for every $n \geq 1$, realized by $\mu = \omega$, $\epsilon = 1$. Then μ is an order unit and ϵ is not. Put

$$G_\mu(z) = \sum_{n \geq 0} t^{n^2\mu} z^n = \sum_{n \geq 0} \omega^{-n^2\omega} z^n, \quad q = 1 + t^\epsilon = 1 + \omega^{-1}.$$

The valuations $n^2\mu + nv(x)$ are cofinal for every $x \neq 0$, so G_μ is nonpolynomial and strongly entire by Lemma 2.3 (a). The unit q is nontorsion, since $(1 + t^\epsilon)^n - 1$ has leading term nt^ϵ . By Corollary 29.7, all finite collections of derivatives of $G_\mu(z)$, $G_\mu(qz)$, $G_\mu(q^2z)$, $\dots$ are linearly independent over $K_\Gamma(z)$. This lies in the order-unit regime, where Theorem E (b) does not apply. At a different dilation scale there is a relation: $G_\mu(z) = \Theta_p(t^\mu z)$ with $p = t^{2\mu}$, so G_μ satisfies the dilation equation (8.3) rescaled; excluding every unit-dilation mixed equation does not exclude one at a cofinal nonzero dilation valuation.

Omnific coefficients (source 18). Under (2.4), omnific coefficients in the workspace have Hahn support in $\Gamma_{\leq 0}$, with an integer coefficient at exponent 0; Gaussian omnific coefficients have a Gaussian-integer constant term. The proofs place no boundedness or positivity condition on operator coefficients, so coefficients in $\mathbf{Oz} \cap K_\Gamma$ or $\mathbf{Oz}[i] \cap K_\Gamma$ are allowed, in the real or complex workspace; infinite omnific coefficients do not evade the theorem. Every set of surreal coefficients lies in some set-sized workspace: the union of their normal-form supports is a set, the group it generates is set sized, and one nonzero exponent may be adjoined if needed. This chooses a common workspace; it does *not* give strong entireness there, which remains a hypothesis and can be lost on enlarging the group. No result on prime omnific integers, their integral closure, automorphisms or Diophantine definability is claimed; the arithmetic connection is only that full, possibly infinite normal forms are legitimate coefficients.

Enlargement (source 18). Enlarge the group of the example by e_* with $e_* > n\mu$ for every n , for instance $e_* = \omega^2$. At $x = t^{-e_*}$ the n th term of G_μ has valuation $n^2\mu - ne_*$, with successive differences $(2n+1)\mu - e_* < 0$, so the evaluation family is not strongly summable (Lemma 2.2): G_μ is entire on the original workspace but not on this enlargement, let alone on $\mathbf{No}[i]$. This is the computation of Section 11.2 and after Corollary 17.2, and no nonpolynomial series is entire on $\mathbf{No}[i]$ (Proposition 27.6).

29.5 Examples and necessary boundaries

Periodic valuations from a torsion residue. Source 18's first example, $q = -(1 + t^\epsilon)$ with $v(q^n - 1) = 0$ for odd n and $= \epsilon$ for even $n \geq 1$, is Example 4.6: even for one nontorsion parameter, Theorem 28.5 needs periodicity, not eventual constancy. At $n = 0$ the value is zero, an isolated exceptional index.

Example 29.14 (Cancellation below a prescribed coefficient; source 18). For $q = 1 + t^\epsilon$ and an integer $d \geq 1$ put $\mathcal{Y}_d(n) = q^n - \sum_{i=0}^{d-1} \binom{n}{i} t^{i\epsilon}$, an exponential polynomial with the bases q and 1. For $n \geq d$ its leading term is $\binom{n}{d} t^{d\epsilon}$, so $v(\mathcal{Y}_d(n)) = d\epsilon$; for $n < d$ it is 0. All coefficient sequences below $d\epsilon$ vanish identically. This is the first-nonidentity-coefficient selection of Theorem 28.5 at work: inspecting the residue of $\mathcal{Y}_d(n)$ would never find the valuation.

Example 29.15 (A large common absolute scale is not enough; source 18). In the group of the example of Section 29.4 take $\lambda_1 = t^\mu$, $\lambda_2 = t^\mu(1 + t^\epsilon)$ and $\lambda_3 = 2t^\mu(1 + t^{2\epsilon})$. All have the cofinal valuation μ and their relative valuations are zero. Their quotients are nontorsion: a quotient with residue 2 or $\frac{1}{2}$ is not a root of unity in characteristic zero, and $1 + t^\epsilon$ is nontorsion by its leading term. So no nonpolynomial strongly entire series satisfies a nonzero mixed equation using this list; the common cofinal factor is irrelevant.

Example 29.16 (Crossing the cofinality boundary; source 18). In the same group, for $\lambda_1 = 1$ and $\lambda_2 = t^\epsilon$ the quotient valuation is positive but noncofinal, and all mixed entire solutions are polynomials. Replacing λ_2 by t^μ changes the answer: Θ_{t^μ} is nonpolynomial and entire and satisfies

$$D_z \Theta_{t^\mu}(z) - \Theta_{t^\mu}(t^\mu z) - z D_z(\Theta_{t^\mu}(t^\mu z)) = 0.$$

So “nonzero valuation” is not the boundary, even for a first-order equation with two multipliers; compare Example 8.4 for pure dilations.

Actual root-of-unity quotients cannot be ignored (source 18). If k contains a root of unity $\zeta \neq 1$ of order h and μ is an order unit, then $G_\mu(z^h) = \sum_n t^{n^2\mu} z^{hn}$ is nonpolynomial and strongly entire, and $(\sigma_\zeta - 1)G_\mu(z^h) = 0$, although $\sigma_\zeta - 1$ is not zero on $\mathbb{K}[[z]]$ (it sends z to $(\zeta - 1)z$) and the relative scale of $1, \zeta$ is zero. This is the periodic-support construction of Proposition 9.1 (a) read as a mixed equation. The hypothesis on root-of-unity quotients in Theorems 29.4 and 29.9 is therefore a real restriction, not a device to avoid a syntactically zero operator; the mechanism is an unconstrained residue class of Taylor indices.

Positive characteristic breaks both steps (source 18). Over $\mathbb{F}_p((t))$ the unit $q = 1 + t$ satisfies $q^{p^a} - 1 = t^{p^a}$ for $a \geq 0$, so $v(q^{p^a} - 1)$ is unbounded and the finite-range conclusion of Theorem 28.5 fails. The series of Example 17.3 is nonpolynomial and strongly entire with derivative zero, while D_z is not zero ($D_z z = 1$). So characteristic zero is essential to the coefficient-polynomial argument and to differential rigidity; the positive-characteristic zero-set theory [33] is a different subject.

Example 29.17 (No degree bound from the shape of the operator; source 18). For $q = 2 \in k$ and every $a \in \mathbb{N}$, $L_a = \sigma_2 - 2^a$ has the strongly entire solution z^a . Each L_a has differential order zero, two dilation terms and constant coefficients, yet the degrees of their polynomial solutions are unbounded. So the degree bound of Theorem 29.4 depends on the exact operator data, not only on the number of terms, derivative orders and coefficient degrees; compare Proposition 16.4 for a single nonlinear equation.

The trivial group $\Gamma = 0$, where every strongly entire series is a polynomial by evaluation at $z = 1$, is excluded to keep the scale language meaningful; it is elementary, not an uncovered counterexample (Section 12).

29.6 Certificates and computation

For a fixed polynomial operator L the proof of Theorem 29.4 reduces to finite data *once the eventual valuation descriptions are established*: the coefficients $\varkappa_{\ell,r,i}$, the least active shift s_0 , the polynomials $R_{j,\ell}$ of (29.5), one period M , one common tail index N , the pairs $(\beta_{j,b}, \varpi_{j,b})$ of (29.6), either $rs = 0$ or a witness γ for (2.2), and a bound on the finitely many intercept differences. These give B , hence the common bad argument $t^{-B-\rho'}$ for any $\rho' > 0$. The support envelope and the first-nonidentity coefficient are part of the certificate's *proof*, not information recoverable from a long finite prefix: a late exceptional zero of an exponential polynomial can invalidate an unjustified choice of N .

Proposition 29.18 (The entire solution space is finite; source 18). *Under Theorem 29.4 the strongly entire solutions of $Lf = 0$ form a finite-dimensional $\mathbb{K}$ -vector space inside $\mathbb{K}[z]_{<N}$; those of $Lf = h$ form the empty set or an affine translate of it, inside the same space. Once a valid N is supplied, all of them are determined by a finite linear system over $\mathbb{K}$.*

Proof. Every entire solution lies in $\mathbb{K}[z]_{<N}$, and every polynomial is entire. If the coefficients of L have degree at most d_L , then L maps $\mathbb{K}[z]_{<N}$ into polynomials of degree at most $N - 1 + d_L$, so $Lf = h$ is finitely many linear equations in the N unknown coefficients. No algorithm for arbitrary encoded Hahn elements is assumed. $\square$

A formalization route proposed by source 18. The report's Lean ledger records modules for escape chains, positive-support words, partial-theta domains and unit-orbit polynomials, which source 18 names as natural reuse points; no Lean build was run and no declaration is added. The proposed order is: the operator-to-recurrence conversion and polynomial–exponential independence; the common support envelope; the first-nonidentity coefficient argument relative to a clearly named zero-set theorem; the relative-scale cost bound and the two-dilation theta witness. The Skolem–Mahler–Lech input should either be proved in the library or appear openly as a hypothesis of a conditional theorem, never as a hidden axiom of a result advertised as kernel checked. The finite-order-residue case (Remark 28.8) needs only root finiteness and is a smaller first target that still includes nontorsion units such as $1 + \omega^{-1}$ in an order-unit workspace.

Remark 29.19 (Source 18's nonlinear question; merge). Source 18's Research question 11.1 asks: in a workspace with an order unit and for a nontorsion unit q , must every strongly entire solution of a nonzero algebraic relation over $\mathbb{K}(z)$ among finitely many $(D_z^r f)(q^\ell z)$ be a polynomial, after genuine algebraic identities among the displayed expressions are excluded? At its pin that was open; in the current text the answer is no. With an order unit μ and $p = t^\mu$, the nonpolynomial strongly entire series $\mathcal{T}_p$ satisfies a nonzero algebraic relation among $f, D_z f, D_z^2 f, D_z^3 f$, the terms with $\ell = 0$ (Theorem 24.1). It is no identity among the displayed expressions: it is a nonzero polynomial in the

jet variables, and it fails for $f = z$, where its coefficient of z^2 is $64 - 192\mathfrak{c}$, of residue 48. So the family of Theorem G (iii) is dependent for every nontorsion q , including $v(q) = 0$ (Corollary 25.2). Source 18's suggested first test case, a second-order differential polynomial without dilations whose leading residue expression vanishes identically, is decided by Theorem I (a): every strongly entire solution of an equation of order at most two is a polynomial. What remains open with an order unit is the classification of Question 19.1; the question is therefore not printed as an open question.

29.7 Predecessors, repository comparison and priority

Literature. Source 18 imports the Skolem–Mahler–Lech theorem in its characteristic-zero form, citing Lech [28] as the primary source (bibliographic metadata verified) and Bell's account [29], whose accessible text it read, including the first two pages; it credits Neumann [30] for the support calculus, whose needed lemma it proves; Gonshor [32] for the normal-form embedding (publisher metadata checked, the monograph not reread); Fuchs–Heintze [31] as related work on growth of recurrences over one-variable function fields, whose nondegeneracy hypothesis and valuation-bound problem differ and which is not an input; and Derksen [33] as background for a positive-characteristic direction. It consulted the Wikipedia article on surreal numbers for orientation only. Its literature searches were targeted, not an exhaustive priority audit. For this merge the bibliographic data of these references were checked; their content was not rechecked, and no proof here rests on Fuchs–Heintze or Derksen.

Repository comparison at the pin. Source 18 read this report's `article.tex` at `bcac55a` in the line ranges listed in its scope file, including the opening statements, the mixed noncofinal theorem and its limitation, the certificate and formalization notes and the questions, comparing against `hol:thm:mixed`, `hol:q:mixedunit`, `hol:q:severaldilations` and `hol:q:nonlinear`; it also read the repository overview and selected openings of other reports. Some responses were truncated, and no absence claim rests on their unreturned parts. There was no line-by-line audit of the repository, the companion archives, the Lean modules or the history. Its statements about this report were accurate at the pin, as recorded in Section 28.1.

Priority. Pure differential rigidity, the cyclic pure-dilation criterion, mixed rigidity at a nonzero noncofinal valuation, the partial theta series, the escape mechanism and the stronger nonlinear statements without an order unit are this report's earlier material and are not claimed by source 18. Its proposed contributions are the periodic-affine valuation theorem for Hahn exponential polynomials with arbitrary well-ordered supports (Theorems 28.5 and 28.10), the bivariate unit-orbit estimate and the answer to Question 19.4, with uniform degree bounds and polynomial forcing (Theorem 29.4, Corollary 29.6), and the relative-scale dichotomy for a finite set of multipliers with its two-multiplier witness (Theorem 29.9). Historical novelty of the valuation theorem is not certified, and no rank-one special case is claimed to be new; the answered question is a question of this report, not a named historical conjecture.

Batch 33 (merge). Source 20 found the support envelope, the unit and periodic-affine valuation theorems, the operator reduction and the mixed unit-valuation corollary independently, on the same day, against the text before this part was merged. They are printed once, here, and its narrower criteria are Corollaries 39.8 and 39.9 (Section 39.4); no priority between the two sources is asserted.

29.8 What the unit-dilation part does not claim

Every limitation stated by source 18, in its manuscript, README, scope file or verification record, is kept.

- (U1) No independent referee report and no Lean formalization accompany source 18; no Lean build was run, no declaration was added, and the repository's axiom audit does not cover this part. A PDF build and a finite test log are not refereeing.
- (U2) The Skolem–Mahler–Lech theorem is imported, not proved anew; no effective bound on exceptional zeros is given.
- (U3) Historical novelty of the valuation theorem and of its full scope is not certified; no rank-one special case is claimed to be historically new, and independence from all earlier rank-one results is not claimed. The answered question is a question of this report at the pin `bcac55a`, not a named historical conjecture.
- (U4) No nonlinear rigidity is claimed: source 18 solves neither the higher-order nonlinear order-unit question nor its own nonlinear question (Remark 29.19 records that the latter is answered negatively in the current text by source 13's series).
- (U5) No unconditional torsion-block theorem is claimed: the hypothesis that no quotient of actual multipliers is a root of unity is essential (Section 29.5), and a finite block criterion is open (Question 19.21).
- (U6) Theorem 29.9 concerns the existence of *some* equation supported in a prescribed set of multipliers; it does not decide prescribed operators or simultaneous systems (Remark 29.10).
- (U7) The proofs are existence proofs: no algorithm finds the first nonidentity coefficient, the eventual winners or the last exceptional zero, no effective zero set is computed, and no decision procedure for arbitrary surreal inputs follows.
- (U8) The exterior regions and degree bounds are uniform for a fixed operator but are not claimed to be exact domains, and they depend on the exact operator data (Example 29.17).
- (U9) Entireness is relative to one set-sized Hahn field: nothing is entire on the proper class $\mathbf{No}[i]$, and entireness can be lost on enlarging the group. D_z kills every coefficient and is not the Berarducci–Mantova derivation; no statement concerns that derivation or power series entire on all of $\mathbf{No}[i]$.
- (U10) Omnific integers enter only as permissible coefficients: no factorization, integral-closure, automorphism or definability result is claimed.
- (U11) Linear independence of formal functions is not algebraic independence of values at a point, and does not exclude polynomial relations.
- (U12) Positive characteristic is excluded, with counterexamples to both the valuation theorem and differential rigidity.
- (U13) The 16,579 finite assertions check finite identities and examples. They do not prove well ordering of arbitrary supports, strong summability at all field elements, the Skolem–Mahler–Lech theorem, finite exceptional sets, the nonexistence of an annihilator, or novelty, and their number is not a measure of proof coverage.

- (U14) The repository comparison was partial and targeted: selected line ranges of this report at the pin, the overview and selected openings of other reports; no audit of all reports, archives, Lean modules or history, and no absence claim from truncated responses. The literature search was targeted.
- (U15) *[merge]* Remarks 29.5 and 29.19, the closed exterior region in Theorem 29.4, and the reading of Corollary 29.11 beside Theorem B are the merge’s; no historical priority is claimed for them.

30 Order three through cubic jet degree: first polars on the twisted cubic

This section and the next contain the material of source 19, delivered after the order-three-threshold part had been merged. Its first theorem, second-order rigidity for every value group and every differential degree, is Theorem I(a), which source 19 found independently of source 14; it is printed once, in Section 26. The new material starts one order higher. At order three the endpoint–gap obstruction of Theorem 27.2 constrains an exterior initial polynomial at both ends; source 19 shows that when the exterior form has a nonzero *first polar* along the twisted cubic of monomial Euler jets, the two constraints push the next exponent to opposite sides of the endpoint and cannot both hold far out, and that every nonzero cubic form vanishing on that curve has a nonzero first polar. Hence no nonpolynomial strongly entire series satisfies an equation of order three and total jet degree at most three (Theorem K). Source 19 adds external constants and Gaussian-omnific coefficients, algebraic two-state systems, exact classifications of two boundary equations beyond the first-polar test, and a sparse strongly entire series whose initial polynomials satisfy both boundary equations at every scale although the series satisfies neither.

30.1 Source, conventions and the merge’s choices

Convention 30.1 (Setting of the Newton-rigidity part). In Sections 30–31, k , Γ , $\mathbb{K}$, v , res , $\mathcal{E}_\Gamma(k)$, D_z , $\vartheta = zD_z$, ϑ_X , the Gauss data γ_δ , ϕ_δ , Act_δ , $\underline{N}_\delta$, N_δ , breaks, the exterior form $\mathfrak{h}_P$ of (26.3), its dehomogenization $\mathfrak{q}$ and the integer i_0 are as in Convention 26.1 and Section 26.2. For a homogeneous $\mathfrak{h} \in k[Y_0, \dots, Y_r]$ write $\mathbf{m}_r(T) = (1, T, \dots, T^r)$ for the *monomial jet curve* (a conic for $r = 2$, the twisted cubic for $r = 3$) and $\chi_{\mathfrak{h}}(T) = \mathfrak{h}(\mathbf{m}_r(T))$; by Remark 26.6, $\chi_{\mathfrak{h}_P} = \chi_P$. A constant extension is a field extension $\mathbb{L}/\mathbb{K}$, with D_z extended to $\mathbb{L}((z))$ by killing $\mathbb{L}$; entireness always refers to $\mathbb{K}$.

For $k = \mathbb{C}$ one has $\mathbb{K} = K_\Gamma$ and $\mathcal{E}_\Gamma(k) = \mathcal{E}_\Gamma$; the generality in k is recorded for this part only.

The source. Source 19 is *Low-Order Nonlinear Rigidity at Surreal Scales: Newton transitions, Euler-jet geometry, and omnific coefficient descent* (24 pages, US letter, 23 September 2026), delivered as the archive `surreal_newton_rigidity` (batch 31, manuscript 08). It consulted the repository at

3d40856953f4bb0f5d069e45a3b4bab6eda33040,

through a connector: the root README (partially), `docs/README.md`, this report’s directory and README, the complete files `10-nonlinear-rigidity-PROOF_AUDIT.md` and `11-coarsening-differential-rigidity-SOURCES_AND_SCOPE.md`, and the first hundred lines of `article.tex`;

the formalization ledger returned no content, and the repository was not built. At that pin this report was the five-source text (sources 08–12): neither source 13 nor source 14 had been merged. Source 19’s description was accurate there: first-order rigidity for every value group, all-order rigidity without an order unit, and the nonlinear question, Question 19.1 (formerly 15.1), open with an order unit. It is stale now in three places. Its principal claim, the unrestricted second-order case, is Theorem I(a), merged from source 14 before source 19 arrived; the two were written on the same day, each against a text of this report that did not contain the other, and neither cites the other. Its abstract’s “the unrestricted all-order, order-unit case is not resolved” and its Question 12.1 (“is every strongly entire differentially algebraic series a polynomial?” with an order unit) are answered negatively by Theorem H: $\mathcal{T}_p$ is a nonpolynomial strongly entire series with an equation of order three. Its statement that its own second-order theorem is the principal proposed contribution is therefore recorded as an independent re-derivation.

Renamed symbols. The following symbols of source 19 would collide with symbols of the earlier parts. They are renamed here, in the new material only.

Source 19	Here	Reason
$\partial = d/dz$; $E = z d/dz$, E_X	D_z ; ϑ , ϑ_X	The notation of Conventions 14.1 and 23.1; E_i are Euler-jet variables. (Source 19’s $K_\Gamma = k((t^\Gamma))$ and $\mathcal{E}_\Gamma(k)$ are $\mathbb{K}$ and $\mathcal{E}_\Gamma(k)$.)
α_n , I_δ , F_δ	$v(a_n)$, Act_δ , $t^{-\gamma_\delta} f(t^{-\delta} X)$	F is (11.2); I_P is the full corner polynomial.
Newton transition	break	As in Section 26.1.
D, W, μ ; residue corner H	d, w, β ; $\mathfrak{h}_P$	Lemma 26.5; μ is an order unit, $H \in \Gamma$.
$c_2(T)$, $c_3(T)$, $c_r(T)$	$\mathbf{m}_2$, $\mathbf{m}_3$, $\mathbf{m}_r$	$\mathfrak{c}$ is a theta constant, c_ν corner coefficients.
$S = Y_0 Y_2 - Y_1^2$; $H = S^e Q$	$\mathfrak{s}$; $\mathfrak{h} = \mathfrak{s}^e \mathfrak{h}_1$	S is a support alphabet; Q a cofactor.
χ_H ; first polar A_H , a , b	$\chi_{\mathfrak{h}}$; $\text{Pol}_{\mathfrak{h}}$, $a_{\mathfrak{h}}$, $b_{\mathfrak{h}}$	χ_P is the residue corner polynomial, $\mathcal{A}$ a theta expression, $A(U)$ a transverse part.
H_3 , Q_4 , $B(m, n)$	$\mathbf{H}_3$, $\mathbf{Q}_4$, $\mathbf{B}(m, n)$	H , Q_s , B and $\mathcal{B}$ have other meanings.
$A(N)$, $B(N)$, N ; $U = E f / f$	$\mathbf{A}_1(\mathbf{n})$, $\mathbf{A}_2(\mathbf{n})$, $\mathbf{n}$; $\mathbf{u}_f$	N is an index; $\mathbf{u}_\phi$ as in Section 26.3.
L/K_Γ (constants); $F = L(z, f, g)$, F_1 ; $q(T)$	$\mathbb{L}/\mathbb{K}$; $\mathfrak{M}$, $\mathfrak{M}_1$; $\mathbf{m}(T)$	L is a coefficient subfield; q a dilation.
u (order unit); c_n ; $\Theta(z) = 1 + t^u z \Theta(t^{2u} z)$	μ ; b_n ; $G_\mu(z) = \sum t^{n^2 \mu} z^n$	u is the Laurent variable; source 19’s Θ is $\Theta_{p^2}(pz)$ with $p = t^\mu$, not Θ_p of (8.1).
$F = \sum t^{e_n} z^n$	F	The same series (11.2).
field F of Section 10; $p = a_m z^m + a_{2m} z^{2m}$	K_0 ; $\mathbf{d}_0$	F is (11.2), as in the row above; K_0 as in Theorem 21.2; $p = t^\mu$.
characteristic p	$\mathfrak{p}$	$p = t^\mu$.
Theorems A, B, C	Theorem I(a), Theorem K, Corollary 30.12 and Theorem 30.16	Theorems A–N of this report are different theorems.

The shipped audit files 19-newton-rigidity-PROOF_AUDIT.md and 19-newton-rigidity-SOURCES_AND_SCOPE.md keep the source’s notation.

Words with two meanings. Source 19’s *residue corner* is the exterior form $\mathfrak{h}_P$, not the residue corner polynomial χ_P ; its *Newton transition* is a break. The *first polar* $\text{Pol}_{\mathfrak{h}}$ is a polynomial in two variables attached to a homogeneous form, not the full corner polynomial I_P . *Degree* of an equation

means its total degree in the jet variables $Y_0, \dots, Y_r$ (source 19: differential degree); the degree in z is unrestricted. *Order* is the highest derivative, as in Section 23.1.

Results printed once. Source 19’s Theorem A, its principal claim, is Theorem I (a), with identical hypotheses (any k of characteristic zero, any Γ , any degree, any residue corner), and its proof is in substance source 14’s: Theorem 3.2 (arbitrarily exterior nonmonomial initial polynomials) is Lemma 26.4, with the break found by a crossing slope in a finite window below one prescribed scale rather than on an interval; Proposition 4.1 (exterior corner reduction) is Lemma 26.5, with the same sign-sensitive inequality (26.4); and its Lemma 5.1 and Proposition 5.3 are Lemma 26.8 in homogeneous coordinates (Remark 30.2). Also printed once: its all-scale criterion (Proposition 2.2), which is Proposition 20.2; the closure of $\mathcal{E}_\Gamma(k)$ (Corollary 2.3), which is Proposition 2.7 with $\vartheta = zD_z$; the cofinal scalar extension (Lemma 2.4), which is Lemma 26.2; the escape of the active support (Lemma 3.1), which is Lemma 14.5; its Remark 5.4, whose equation $z^2ff'' + zff' - z^2f'^2 = 0$ is (16.1), with the classification of Proposition 16.4; the finite linear descent (Lemma 7.1), which is Proposition 22.7, whose basis proof also preserves the total degree; the full-class obstruction (Proposition 7.6), which is Proposition 27.6; the embedding (2.4); and its examples in characteristic $\mathfrak{p}$ and with $\sum z^n/n!$ (Section 9.3), which are Example 17.3, the positive-characteristic paragraph of Section 27.4 and (2.3). The quadratic profile of its Example 3.4 (for G_1 at $\delta = 2n + 1$ the initial polynomial is $X^n + X^{n+1}$) is the Newton profile recorded after Theorem 27.5 for $\mathcal{T}_p$.

Where each numbered result went. Theorem A is Theorem I (a), and Theorem B is Theorem K; Definition 6.1 and Lemmas 6.2, 6.3 are Definition 30.3 and Lemmas 30.4, 30.5; Proposition 6.4, Lemma 6.5, Remark 6.6, Corollary 6.7 and Example 6.8 are Proposition 30.6, Lemma 30.7, Remark 30.8, Corollary 30.9 and Example 30.10; Corollaries 7.2, 7.3, Lemma 7.4 and Corollary 7.5 are Corollaries 30.12, 30.13, Lemma 30.14 and Corollary 30.15; Theorem 8.1 and Corollary 8.2 are Theorem 30.16 and Corollary 30.17; Proposition 9.1 is Proposition 31.5; Theorems 10.1 and 10.2 are Theorems 31.1 and 31.2, and Section 10.3 is Example 31.3; Question 12.1 is answered (Section 30.1); Questions 12.4, 12.6 and 12.8 are Questions 19.15, 19.16 and 19.17, asked by source 14 in the same form, with status notes recording source 19’s versions; Questions 12.2, 12.3, 12.5, 12.7, 12.9 and 12.10 are Questions 19.30–19.35.

The merge’s choices. The review of source 19 for this merge found its proofs correct; the identities it states were rechecked with an independent SymPy script, which is not shipped. The merge adds four points, each marked. First, Remark 30.11 places the first-polar criterion inside the endpoint–gap obstruction of Theorem 27.2: for $\chi = 0$ the first polar is nonzero exactly when $i_0 = 1$, the two linear conditions of source 19 are the two endpoint conditions of Theorem 27.2, and an explicit cubic shows that Corollary 27.3, which uses the upper endpoint only, does not contain Theorem K. Second, Corollary 31.4 applies Theorem K to $\mathcal{T}_p$ and to the theta exterior form. Third, Example 31.3 is read as a negative instance for Question 19.11. Fourth, the difference between Theorem 30.16 and Corollary 26.10 is stated. Source 19’s theorems, proofs, examples and non-claims are otherwise printed as delivered, in the renamed notation.

Remark 30.2 (Source 19’s route to Theorem I (a); source 19). Source 19 proves the endpoint lemma in homogeneous coordinates. Modulo $\mathfrak{s} = Y_0Y_2 - Y_1^2$ the monomials $Y_0^{d-b}Y_1^b$ ($0 \leq b \leq d$) and $Y_1^{d-c}Y_2^c$ ($1 \leq c \leq d$) span degree d , and their images $1, T, \dots, T^{2d}$ under $Y \mapsto \mathbf{m}_2(T)$ are independent; so a homogeneous $\mathfrak{h}$ with $\chi_{\mathfrak{h}} = 0$ is divisible by $\mathfrak{s}$, and every nonzero homogeneous $\mathfrak{h}$ is uniquely $\mathfrak{s}^e \mathfrak{h}_1$ with $\chi_{\mathfrak{h}_1} \neq 0$. Since $\mathfrak{s}(1, U, U^2 + V) = V$, this e is the i_0 of (26.6) and $\mathfrak{h}_1(1, U, U^2 + V) = \mathfrak{r}$, so the

endpoint spectrum $\text{Sp}(\mathfrak{h})$ is the set of nonnegative integral roots of $\chi_{\mathfrak{h}_1} = \mathfrak{r}(T, 0)$. Source 19 then argues at the top degree only: if $\deg \phi$ exceeds these roots, the coefficient of degree $(\deg \mathfrak{h}_1) \deg \phi$ in $\mathfrak{h}_1(\phi, \vartheta_X \phi, \vartheta_X^2 \phi)$ is $\text{lc}(\phi)^{\deg \mathfrak{h}_1} \chi_{\mathfrak{h}_1}(\deg \phi) \neq 0$, so there is no such ϕ if $e = 0$, and $\mathfrak{s}(\phi, \vartheta_X \phi, \vartheta_X^2 \phi) = 0$ if $e > 0$; and a nonzero polynomial with $\mathfrak{s}(\phi, \dots) = 0$ is a monomial, because for its first two exponents $m < n$ the coefficient of X^{m+n} in $\phi \vartheta_X^2 \phi - (\vartheta_X \phi)^2$ is $a_m a_n (n - m)^2 \neq 0$. Source 14 uses logarithmic Euler derivatives at both endpoints instead. The two routes prove the same lemma; the variant is recorded, not printed as a separate theorem.

30.2 The first polar and third-order rigidity

Definition 30.3 (First polar along the twisted cubic; source 19). For a homogeneous $\mathfrak{h} \in k[Y_0, Y_1, Y_2, Y_3]$ of degree $d \geq 1$ put

$$\text{Pol}_{\mathfrak{h}}(T, U) = \sum_{j=0}^3 \frac{\partial \mathfrak{h}}{\partial Y_j}(\mathbf{m}_3(T)) U^j. \quad (30.1)$$

Lemma 30.4 (The double factor; source 19). *If $\chi_{\mathfrak{h}} = 0$, there are $a_{\mathfrak{h}}, b_{\mathfrak{h}} \in k[T]$ with*

$$\text{Pol}_{\mathfrak{h}}(T, U) = (U - T)^2(a_{\mathfrak{h}}(T)U + b_{\mathfrak{h}}(T)), \quad (30.2)$$

not both zero if $\text{Pol}_{\mathfrak{h}} \neq 0$.

Proof. Euler's identity gives $\text{Pol}_{\mathfrak{h}}(T, T) = d\chi_{\mathfrak{h}}(T) = 0$, and differentiating $\chi_{\mathfrak{h}}(T) = 0$ gives $\partial_U \text{Pol}_{\mathfrak{h}}(T, U)|_{U=T} = 0$. Division in $k[T][U]$ gives the factor $(U - T)^2$, and the U -degree is at most three. $\square$

Lemma 30.5 (An eventual side for integral roots; source 19). *Let $a, b \in k[T]$ be not both zero. There is $B \in \mathbb{N}$ such that for integers $m > B$ every integral solution $n \neq m$ of $a(m)n + b(m) = 0$ lies on one fixed side of m : always $n > m$, or always $n < m$, the side not depending on m . There may be no solutions.*

Proof. If $a = 0$, then $b(m) \neq 0$ for all but finitely many m , and eventually there are no solutions. Let $a \neq 0$ and $C(T) = Ta(T) + b(T)$. If $C = 0$, then away from the roots of a the only root is $n = m$, which is excluded. Let $a, C \neq 0$ and let $V \subseteq k$ be the finite-dimensional $\mathbb{Q}$ -span of their coefficients. There is a $\mathbb{Q}$ -linear $\ell : V \rightarrow \mathbb{Q}$ for which neither coefficientwise image $a_*, C_* \in \mathbb{Q}[T]$ is zero: the functionals killing all coefficients of a , or all of C , form two proper subspaces of V^* , and a vector space over the infinite field $\mathbb{Q}$ is not the union of two proper subspaces. Applying ℓ to $a(m)(n - m) + C(m) = 0$ gives $a_*(m)(n - m) + C_*(m) = 0$. For large m both $a_*(m)$ and $C_*(m)$ have fixed nonzero signs, so $n - m = -C_*(m)/a_*(m)$ has a fixed strict sign. This is a statement about integers, not an ordering of k . $\square$

Proposition 30.6 (The first-polar criterion; source 19). *Let $0 \neq \mathfrak{h} \in k[Y_0, Y_1, Y_2, Y_3]$ be homogeneous, and suppose $\chi_{\mathfrak{h}} \neq 0$, or $\chi_{\mathfrak{h}} = 0$ and $\text{Pol}_{\mathfrak{h}} \neq 0$. There is $B_{\mathfrak{h}} \in \mathbb{N}$ such that every nonzero $\phi \in k[X]$ with*

$$\mathfrak{h}(\phi, \vartheta_X \phi, \vartheta_X^2 \phi, \vartheta_X^3 \phi) = 0, \quad \text{ord}_X \phi > B_{\mathfrak{h}},$$

is a monomial.

Proof. If $\chi_{\mathfrak{h}} \neq 0$, the top coefficient of the expression is $\text{lc}(\phi)^d \chi_{\mathfrak{h}}(\deg \phi) \neq 0$ once $\deg \phi$ exceeds the integral roots of $\chi_{\mathfrak{h}}$, and there is no such solution. Let $\chi_{\mathfrak{h}} = 0 \neq \text{Pol}_{\mathfrak{h}}$ and let ϕ be nonmonomial, with first two exponents $m < n$, last two $\ell < M$, and coefficients a_j . Start with all d factors at degree m ; one replacement by n gives exactly the first polar, and two replacements, or one by a larger exponent, raise the degree further. So the coefficient of $X^{(d-1)m+n}$ is

$$a_m^{d-1} a_n \text{Pol}_{\mathfrak{h}}(m, n), \quad (30.3)$$

and symmetrically the coefficient of $X^{(d-1)M+\ell}$ is $a_M^{d-1} a_{\ell} \text{Pol}_{\mathfrak{h}}(M, \ell)$. Both vanish; by (30.2) and characteristic zero, $a_{\mathfrak{h}}(m)n + b_{\mathfrak{h}}(m) = 0$ and $a_{\mathfrak{h}}(M)\ell + b_{\mathfrak{h}}(M) = 0$. Once m , and so $M \geq m$, exceeds the bound of Lemma 30.5, these are incompatible: $n > m$ but $\ell < M$, on opposite sides. $\square$

Lemma 30.7 (No cubic is singular along the twisted cubic; source 19). *If a homogeneous $\mathfrak{h} \in k[Y_0, Y_1, Y_2, Y_3]$ of positive degree at most three has all first partial derivatives zero on $\mathbf{m}_3(T)$, then $\mathfrak{h} = 0$.*

Proof. Extend k to an algebraic closure. By Euler's identity also $\chi_{\mathfrak{h}} = 0$, since the degree is nonzero in k . Restrict $\mathfrak{h}$ to secants: $G(A, B) = \mathfrak{h}(A\mathbf{m}_3(s) + B\mathbf{m}_3(t))$ is homogeneous of degree at most three in (A, B) , and its value and first transverse derivative vanish at $B = 0$, so $B^2 \mid G$; likewise $A^2 \mid G$; hence $G = 0$. The secant map $(A, B, s, t) \mapsto (A + B, As + Bt, As^2 + Bt^2, As^3 + Bt^3)$ has Jacobian determinant

$$-AB(s - t)^4 \neq 0, \quad (30.4)$$

so in characteristic zero its coordinates are algebraically independent (a minimal-degree relation, differentiated, would have all partial derivatives zero by the invertible Jacobian), and no nonzero polynomial vanishes on its image. So $\mathfrak{h} = 0$. $\square$

Remark 30.8 (The exact use of the lemma; source 19). $\text{Pol}_{\mathfrak{h}} = 0$ says that every coefficient in U of (30.1) vanishes, that is, every first partial vanishes on the curve. So a nonzero homogeneous $\mathfrak{h}$ of degree at most three with $\chi_{\mathfrak{h}} = 0$ has $\text{Pol}_{\mathfrak{h}} \neq 0$.

Proof of Theorem K. Suppose a nonpolynomial $f \in \mathcal{E}_{\Gamma}(k)$ satisfies a nonzero equation of order at most three and total jet degree at most three. Pass to $\mathbb{K}_{\mathbb{Q}}$ (Lemma 26.2) and form the exterior form $\mathfrak{h}_P$, nonzero homogeneous of degree between one and three, since the Euler substitution (26.2) preserves total jet degree (Lemma 26.5). If $\chi_{\mathfrak{h}_P} = 0$, Lemma 30.7 gives $\text{Pol}_{\mathfrak{h}_P} \neq 0$. Either way Proposition 30.6 applies: beyond the δ_0 of Lemma 26.5 and the scale of Lemma 14.5 beyond which every active index exceeds $B_{\mathfrak{h}_P}$, every initial polynomial is a monomial, while Lemma 26.4 gives a break there. For the general assertion, apply Proposition 30.6 to $\mathfrak{h}_P$, or Corollary 30.9 to its irreducible factors, in the same argument. $\square$

Corollary 30.9 (A factor criterion beyond cubic degree; source 19). *For an equation of order at most three, factor $\mathfrak{h}_P$ into homogeneous irreducible factors over k . If each factor $\mathfrak{g}$ has $\chi_{\mathfrak{g}} \neq 0$ or $\text{Pol}_{\mathfrak{g}} \neq 0$, every strongly entire solution is a polynomial.*

Proof. Take the largest of the bounds of Proposition 30.6 for the finitely many factors. Since $k[X]$ is a domain, an initial polynomial solving the product equation solves one factor equation, and the argument above applies. Factoring first removes repeated factors, whose unfactored first polar can vanish identically. $\square$

Example 30.10 (Two ends forcing incompatible next exponents; source 19). For $\mathfrak{h} = Y_0^2 Y_3 - 4Y_0 Y_1 Y_2 + 3Y_1^3$ one has $\chi_{\mathfrak{h}} = 0$ and $\text{Pol}_{\mathfrak{h}}(T, U) = (U - T)^2(U - 2T)$. A nonmonomial polynomial solution with lowest exponent m would need its next exponent to be $2m$, and at its largest exponent M the next smaller one would also have to be $2M$, which is impossible. This case shows the two-end argument without the rational functional of Lemma 30.5.

Remark 30.11 (Relation to the endpoint-gap obstruction; merge). Let $\mathfrak{h}$ be homogeneous in $Y_0, \dots, Y_3$ with $\chi_{\mathfrak{h}} = 0$, and write $\mathfrak{h}_j$ for $\partial \mathfrak{h} / \partial Y_j$ evaluated on $\mathbf{m}_3(T)$. In (27.1) the linear part of $\mathfrak{q}$ in (V, W) is $V(\mathfrak{h}_2 + 3T\mathfrak{h}_3) + W\mathfrak{h}_3$ at $U = T$, and the relations $\mathfrak{h}_0 + T\mathfrak{h}_1 + T^2\mathfrak{h}_2 + T^3\mathfrak{h}_3 = 0$ (Euler) and $\mathfrak{h}_1 + 2T\mathfrak{h}_2 + 3T^2\mathfrak{h}_3 = 0$ (derivative of $\chi_{\mathfrak{h}}$) show that it vanishes exactly when all $\mathfrak{h}_j$ vanish. So, when $\chi_{\mathfrak{h}} = 0$, $\text{Pol}_{\mathfrak{h}} \neq 0$ exactly when $i_0 = 1$, and then, comparing coefficients in (30.2), $a_{\mathfrak{h}} = \mathfrak{h}_3$, $b_{\mathfrak{h}} = \mathfrak{h}_2 + 2T\mathfrak{h}_3$ and

$$\Upsilon_{\mathfrak{h}}(T, G) = a_{\mathfrak{h}}(T)(T - G) + b_{\mathfrak{h}}(T).$$

The conditions $\Upsilon_{\mathfrak{h}}(N, \mathbf{g}^+) = 0$ and $\Upsilon_{\mathfrak{h}}(m, -\mathbf{g}^-) = 0$ of Theorem 27.2 are then exactly source 19's $a_{\mathfrak{h}}(N)\ell + b_{\mathfrak{h}}(N) = 0$ with $\ell = N - \mathbf{g}^+$, and $a_{\mathfrak{h}}(m)n + b_{\mathfrak{h}}(m) = 0$ with $n = m + \mathbf{g}^-$. The new ingredients of Theorem K are therefore Lemma 30.5, which makes one of the two endpoint conditions eventually unsatisfiable, and Lemma 30.7, which forces $i_0 \leq 1$ in degree at most three. Corollary 27.3 tests the upper endpoint only and does not contain the theorem. For the cubic

$$\mathfrak{h} = 2Y_0^2 Y_3 - 5Y_0 Y_1 Y_2 + 3Y_1^3, \quad \chi_{\mathfrak{h}} = 0, \quad \text{Pol}_{\mathfrak{h}} = (U - T)^2(2U - T), \quad \Upsilon_{\mathfrak{h}} = T - 2G,$$

the upper condition allows the gap $\mathbf{g}^+ = N/2$ for every even N , so the set in Corollary 27.3 is unbounded, while the lower condition asks for the next exponent $m/2 < m$ and has no solution; the equation $2f^2 \vartheta^3 f - 5f \vartheta f \vartheta^2 f + 3(\vartheta f)^3 = 0$ is covered by Theorem K and not by that corollary. For Example 30.10, $\Upsilon_{\mathfrak{h}} = -(G + T)$ and the upper condition alone suffices. These identities were checked with SymPy.

30.3 External constants and planar systems

Corollary 30.12 (External constants; source 19). *For nonpolynomial $f \in \mathcal{E}_{\Gamma}(k)$ and every field extension $\mathbb{L}/\mathbb{K}$, the series $f, D_z f, D_z^2 f$ are algebraically independent over $\mathbb{L}(z)$, and $f, D_z f, D_z^2 f, D_z^3 f$ satisfy no nonzero polynomial relation of total degree at most three over $\mathbb{L}(z)$.*

Proof. A relation over $\mathbb{L}(z)$ gives one over $\mathbb{K}(z)$ of no larger total degree: in the proof of Proposition 22.7 each Q_{ν} has total degree at most that of Q (source 19 uses a $\mathbb{K}$ -linear functional on the finite span of the coefficients instead of a basis). Apply Theorems I(a) and K. $\square$

This is stronger than reapplying the analytic theorems over a larger Hahn field, which would need entireness there and can fail; the finite algebraic descent needs no analytic hypothesis.

Corollary 30.13 (External surcomplex coefficients; source 19). *Let $\Gamma \subseteq \mathbf{No}$ be a set-sized subgroup and $f \in \mathbb{C}((t^{\Gamma}))[[z]]$ nonpolynomial and strongly entire on that workspace. No nonzero polynomial relation with finitely many coefficients in $\mathbf{No}[i](z)$ holds among $f, D_z f, D_z^2 f$, and none of total degree at most three among $f, \dots, D_z^3 f$.*

Proof. The workspace and the finitely many coefficients of a relation lie in a set-sized subfield of $\mathbf{No}[i]$; apply Corollary 30.12. No proper-class polynomial ring or tensor product is used. $\square$

Lemma 30.14 (Finite monomial clearance; source 19). *For finitely many $c_1, \dots, c_s \in \mathbf{No}[i]$ there is a surreal $\rho > 0$ with $\omega^\rho c_j \in \mathbf{Oz}[i]$ for all j ; the real and imaginary parts of these products have constant term 0.*

Proof. The union of the normal-form supports of the real and imaginary parts is a set; take $\rho > 0$ larger than the negative of each of its exponents. Translating by ρ makes every exponent positive and preserves the order of the support, which is the normal-form description of $\mathbf{Oz}$ with constant term 0, as in `odg:def:rings` [41]. If the union is empty any $\rho > 0$ works. $\square$

Corollary 30.15 (Omnific differential-relation obstruction; source 19). *For every nonpolynomial strongly entire f in a surcomplex Hahn workspace, $P(z, f, D_z f, D_z^2 f) \neq 0$ for every nonzero $P \in \mathbf{Oz}[i][z, Y_0, Y_1, Y_2]$, and the same holds through $D_z^3 f$ when the total jet degree is at most three. Conversely every surcomplex polynomial differential relation with finitely many coefficients becomes a Gaussian-omnific one after clearing z -denominators and multiplying by one monomial ω^ρ .*

Proof. The obstruction is a case of Corollary 30.13; the conversion is Lemma 30.14, and multiplication by a nonzero scalar preserves order and degree. $\square$

Monomial clearance is elementary normal-form algebra, not a new arithmetic theorem. The consequence concerns finitely many coefficients of formal relations: it says nothing about factorization of omnific integers and is not algebraic independence of a value $f(x)$ from the field containing it.

Theorem 30.16 (Algebraic two-state systems; source 19). *Let $\mathbb{L}/\mathbb{K}$ be a field extension and $f, g \in \mathcal{E}_\Gamma(k)$. If $D_z f$ and $D_z g$ are algebraic over $\mathfrak{M} = \mathbb{L}(z, f, g) \subseteq \mathbb{L}((z))$, then $f, g \in \mathbb{K}[z]$.*

Proof. $\mathfrak{M}_1 = \mathfrak{M}(D_z f, D_z g)$ is finite algebraic over $\mathfrak{M}$, so $\text{trdeg}_{\mathbb{L}(z)} \mathfrak{M}_1 \leq 2$, and derivatives of elements of $\mathfrak{M}$ lie in $\mathfrak{M}_1$. Let $m(T) \in \mathfrak{M}[T]$ be the minimal polynomial of $D_z f$; it is separable in characteristic zero, so $m'(D_z f) \neq 0$. Differentiating $m(D_z f) = 0$ gives $D_z^2 f = -m^D(D_z f)/m'(D_z f) \in \mathfrak{M}_1$, with m^D obtained by differentiating the coefficients. Likewise $D_z^2 g \in \mathfrak{M}_1$. So $f, D_z f, D_z^2 f$ lie in a field of transcendence degree at most two over $\mathbb{L}(z)$ and are dependent, and Corollary 30.12 makes f a polynomial; likewise g . $\square$

The separant is nonzero because it comes from a minimal polynomial, not from an arbitrary singular presentation of an implicit equation, and $\mathfrak{M}_1$ is in fact closed under D_z .

Corollary 30.17 (Rational planar systems; source 19). *Every strongly entire pair f, g with $D_z f = R_1(z, f, g)$ and $D_z g = R_2(z, f, g)$, $R_1, R_2 \in \mathbb{L}(z, Y_1, Y_2)$, denominators nonzero after substitution, is polynomial. The coefficients may be external surcomplex constants.*

These statements assert neither the existence of polynomial solutions, nor finiteness of their number, nor a uniform bound on their degrees; the strongly entire solution locus contains no genuinely transcendental pair, and (16.1) has polynomial solutions of unbounded degree.

Comparison with Corollary 26.10 (merge). The two statements overlap without containing each other. Corollary 26.10 (source 14) needs a *rational* system over $\mathbb{K}(z)$ but lets the states u_1, u_2 be arbitrary elements of a differential field, not necessarily entire, and concludes for every entire element of $\mathbb{K}(z, u_1, u_2)$. Theorem 30.16 (source 19) needs the states f, g themselves to be entire but allows *algebraic* dependence of $D_z f, D_z g$ over $\mathbb{L}(z, f, g)$ and external constants $\mathbb{L} \supseteq \mathbb{K}$. Both rest on three-jet independence (Corollary 26.9), the second through Proposition 22.7. With an order unit,

$\mathcal{T}_p$ is an observable of a four-state rational system (Theorem 24.10), so neither statement extends to four states; three states are open (Question 19.32).

31 Exact classifications beyond the first-polar test, and scope

The first-polar criterion is sufficient, not necessary. The following two equations show its boundary without producing a counterexample to rigidity. In Sections 31.1 and 31.2, K_0 is any field of characteristic zero, $f \in K_0[[z]]$, and $\vartheta = z d/dz$; no Hahn or entire hypothesis is used.

31.1 A quartic equation of order three

Put

$$H_3 = 2Y_0^2Y_3^2 - 13Y_0Y_1Y_2Y_3 + 9Y_0Y_2^3 + 9Y_1^3Y_3 - 7Y_1^2Y_2^2. \quad (31.1)$$

All its first partial derivatives vanish on $\mathbf{m}_3(T)$, so $\text{Pol}_{H_3} = 0$ and Proposition 30.6 does not apply; and H_3 annihilates the Euler jets of $az^n + bz^{2n}$ for all a, b, n , which have arbitrarily large lowest degree.

Theorem 31.1 (Exact formal solutions of the quartic; source 19). *The formal power series with $H_3(f, \vartheta f, \vartheta^2 f, \vartheta^3 f) = 0$ are exactly 0, the monomials az^m ($m \in \mathbb{N}$) and the binomials $az^n + bz^{2n}$ with $n \geq 1$ and $ab \neq 0$.*

Proof. Write $Y_j = \vartheta^j f$ and, in an auxiliary variable n ,

$$A_1(n) = 2Y_0n^2 - 3Y_1n + Y_2, \quad A_2(n) = 2Y_1n^2 - 3Y_2n + Y_3.$$

Expanding the Sylvester determinant gives

$$\text{Res}_n(A_1, A_2) = 2H_3. \quad (31.2)$$

Let $f \neq 0$ and put $\mathfrak{s}_f = f\vartheta^2 f - (\vartheta f)^2$. If $\mathfrak{s}_f = 0$, the first-two-exponents argument of Remark 30.2, valid for formal series, or Proposition 16.4, makes f a monomial. Let $\mathfrak{s}_f \neq 0$. By (31.2), A_1 and A_2 have a common root in an algebraic closure of $K_0((z))$ (A_1 is genuinely quadratic since $f \neq 0$; a degree drop in A_2 does not affect this). The combination $Y_0A_2 - Y_1A_1 = -3\mathfrak{s}_fn + (Y_0Y_3 - Y_1Y_2)$ shows that the root is $n = (f\vartheta^3 f - \vartheta f \vartheta^2 f)/(3\mathfrak{s}_f) \in K_0((z))$. Applying ϑ to $A_1(n) = 0$, with $\vartheta Y_j = Y_{j+1}$,

$$A_2(n) + (4fn - 3\vartheta f) \vartheta n = 0, \quad (31.3)$$

so $\vartheta n = 0$ or $4fn = 3\vartheta f$. In the first case $n \in K_0$, the constants of ϑ on $K_0((z))$, and the coefficients of $A_1(n) = 0$ read $(j-n)(j-2n)a_j = 0$: at most two coefficients are nonzero, and if two, their degrees are n and $2n$ with n a positive integer. In the second case substitution gives $8f\vartheta^2 f - 9(\vartheta f)^2 = 0$, that is $8\vartheta u_f - u_f^2 = 0$ for $u_f = \vartheta f/f$. With $m = \text{ord}_z f$, $u_f = m + O(z)$ and the constant term forces $m = 0$; if $u_f \neq 0$, its first positive degree r gives a nonzero term of degree r in $8\vartheta u_f$ while u_f^2 starts at degree $2r$. So $u_f = 0$, f is constant and $\mathfrak{s}_f = 0$, a contradiction. Substitution verifies the listed solutions. $\square$

31.2 A quadratic equation of order four

Put $Q_4 = -2Y_0Y_4 + 9Y_1Y_3 - 7Y_2^2$. For distinct monomial degrees m, n its cross coefficient is governed by

$$B(m, n) = -2(m^4 + n^4) + 9(mn^3 + nm^3) - 14m^2n^2 = (m - n)^2(m - 2n)(n - 2m). \quad (31.4)$$

Theorem 31.2 (Exact formal solutions of the order-four quadratic; source 19). *The solutions in $K_0[[z]]$ of $-2f\vartheta^4f + 9\vartheta f\vartheta^3f - 7(\vartheta^2f)^2 = 0$ are exactly those of Theorem 31.1: zero, monomials and doubling binomials.*

Proof. Monomials are solutions since $-2 + 9 - 7 = 0$. If f has two terms or more, let $m < n$ be its first two exponents; the coefficient at z^{m+n} is $a_ma_nB(m, n)$, so (31.4) forces $n = 2m$ and $m > 0$. For $d_0 = a_mz^m + a_{2m}z^{2m}$ the same factorization gives $Q_4(d_0) = 0$. If $f - d_0 \neq 0$ has first exponent $r > 2m$, the coefficient at z^{m+r} of $Q_4(f)$ is $a_ma_rB(m, r)$: the other cross terms start later and the square of $f - d_0$ starts at $2r > m + r$. No factor of $B(m, r)$ vanishes for $r > 2m$. $\square$

Theorems 31.1 and 31.2 are exact positive results, not counterexamples to polynomial rigidity. They locate an obstruction to a *proof technique*: their exterior equations admit nonmonomial polynomial solutions of arbitrarily large lowest degree. In the language of Theorem 27.2 (merge), H_3 has $i_0 = 2$ and $\Upsilon_{H_3}(T, G) = (G + T)(2G - T)$, which accepts the upper gap $N/2$ and the lower gap m of a doubling binomial at every scale; Q_4 has order four, outside that theorem.

Example 31.3 (Valid initial equations at every scale, and no identity; source 19). Over a rank-one workspace with the order unit 1, for instance $\mathbb{K} = k((t^{\mathbb{Q}}))$, put

$$F_{\text{sp}}(z) = \sum_{j \geq 0} t^{4^j} z^{2^j}. \quad (31.5)$$

It is strongly entire, since $4^j - 2^j\delta \rightarrow +\infty$ for every δ . At the break $\delta = 3 \cdot 2^j$ the active degrees are 2^j and 2^{j+1} and $\phi_\delta = X^{2^j} + X^{2^{j+1}}$; between breaks the initial polynomial is a monomial. So every initial polynomial satisfies both $H_3 = 0$ and $Q_4 = 0$, yet F_{sp} satisfies neither: only the degrees 1 and 4 contribute to z^5 in $Q_4(F_{\text{sp}})$, and

$$[z^5] Q_4(F_{\text{sp}}) = B(1, 4)t^{1+16} = -126t^{17} \neq 0, \quad (31.6)$$

while Theorem 31.1 excludes $H_3(F_{\text{sp}}) = 0$. Vanishing leading terms at all scales therefore do not yield an actual differential identity: higher support layers and the compatibility of successive breaks carry information that no single initial equation sees. For Question 19.11 this is a concrete negative instance (merge's reading): an unbounded sequence of initial polynomials, each satisfying the exterior form of an equation, need not be the initial polynomials of any solution of that equation, although here they are those of one strongly entire series.

Corollary 31.4 (The theta equation and its jet degree; merge). *Suppose Γ has an order unit μ and $p = t^\mu$. Every nonzero algebraic differential equation of order at most three over $\mathbb{K}(z)$ satisfied by T_p has order exactly three and total jet degree at least four; the equation of Theorem 24.1 has degree six, so the least such degree is four, five or six. The exterior form (27.5) of that equation has $i_0 = 2$ and vanishing first polar, as Proposition 30.6 requires of every equation of order at most three with a nonpolynomial strongly entire solution.*

Proof. $\mathcal{T}_p$ is nonpolynomial and strongly entire (Theorem H); Theorem I(a) excludes order at most two and Theorem K total degree at most three. The equation (24.8) is homogeneous of degree six in the jets. The least transverse degree of $W^2 - V^2 + 4V^3$ is two, and with $\chi = 0$ the first polar vanishes exactly when $i_0 \geq 2$ (Remark 30.11); if an equation of order at most three had an exterior form with $\chi \neq 0$ or a nonzero first polar, all its strongly entire solutions would be polynomials, by the proof of Theorem K. $\square$

Which of four, five and six is the least degree for $\mathcal{T}_p$ is not decided here: it depends on whether (24.8) is irreducible over $\mathbb{K}(z)$, which was not checked.

31.3 Examples and the role of the hypotheses

Proposition 31.5 (A large order-unit family; source 19). *Let $\mu > 0$ be an order unit of Γ and let each $b_n \in \mathbb{K}$ be zero or of valuation zero. Then $f(z) = \sum_{n \geq 0} b_n t^{n^2 \mu} z^n$ is strongly entire. If infinitely many b_n are nonzero, its first three jets are algebraically independent over every constant extension $\mathbb{L}(z)$, and its first four jets satisfy no polynomial relation of total degree at most three over $\mathbb{L}(z)$.*

Proof. Given $\delta, \beta \in \Gamma$ choose positive integers r, s with $\delta \leq r\mu$ and $\beta \leq s\mu$; then $n^2\mu - n\delta \geq (n^2 - rn)\mu > \beta$ for large n . Apply Proposition 20.2 and Corollary 30.12. $\square$

With $\Gamma = \mathbb{Q} \subseteq \mathbf{No}$ and $\mu = 1$, the surcomplex reading is $\sum_n b_n \omega^{-n^2} z^n$ on $\mathbb{C}((\omega^{-\mathbb{Q}}))$, and the coefficients b_n , among zero and the units, are arbitrary: the conclusion is not about one named special function. For $b_n = 1$ the series is G_μ , with $G_\mu(z) = 1 + t^\mu z G_\mu(t^{2\mu} z)$, that is $G_\mu(z) = \Theta_{p^2}(pz)$ for $p = t^\mu$; this dilation example is prior material (Corollary 17.1, Section 29.4), and three independent jets for G_1 are Corollary 26.13. It illustrates why differential and dilation equations have different rigidity thresholds.

A higher-rank comparison (source 19). Over Γ_∞ of (11.1), which has no order unit, the series F of (11.2) is strongly entire, and Theorems I and K apply to it; but Theorem 22.1 is already stronger for it, in every order. The comparison explains the division of labour and adds no second novelty claim. If Γ has no countable cofinal subset, every strongly entire series is a polynomial, since by Proposition 20.2 at $\delta = 0$ the countably many nonzero coefficient values would be cofinal; the order-unit examples are not affected by this size obstruction.

The literature boundary (source 19). Non-Archimedean functional-equation theory [20] is context: its absolute-value hypotheses and particular equations are not transferred, and no theorem is imported from it. The general D-algebraic framework [26] is context; the low-order obstructions here use strong entireness in addition.

31.4 Predecessors, repository comparison and priority

Literature. Source 19 consulted the Mantova–Matusinski survey [17], Section 2.2, for normal forms and the ω -map and to distinguish scalar surreal derivations from D_z ; Hu–Luan [20] for comparison with non-Archimedean functional equations; and Ait El Manssour, Sattelberger and Tegua Tabugua [26] for general differential-algebraic context. No theorem of these is used in its proofs. It opened the Wikipedia article on surreal numbers for orientation only and located related coefficient-arithmetic literature not used as a dependency. Several targeted queries returned

irrelevant results, which support no absence claim; no exhaustive MathSciNet, zbMATH, book, thesis or citation-network audit was made.

Repository comparison at the pin. Source 19 read the files listed in Section 30.1 at 3d40856 through a connector; a code search for the question label returned nothing, and the question was located through the scope documents; a container clone failed for lack of network access. This was not a line-by-line audit of the 91-page report as it then stood, of the 57 reports then catalogued, of archived submissions, history or Lean declarations, and directory inventories are not used to prove absence. Its statements were accurate at the pin; the stale ones are recorded in Section 30.1.

Priority. Second-order rigidity is Theorem I(a), merged from source 14; source 19's proof is an independent re-derivation written the same day against a text of this report that did not contain source 14, and no priority between the two is asserted. The all-scale criterion, the corner technique, the no-order-unit results and the partial-theta witness are credited as prior. The proposed contributions of source 19 that are new to this report are third-order rigidity through cubic jet degree (Theorem K) with the first-polar and factor criteria, the secant lemma and the eventual-side lemma; the external-constant, Gaussian-omnific and algebraic two-state consequences; and the exact classifications of H_3 and Q_4 with the sparse series. Whether equivalent results exist in the literature needs a more exhaustive review, and no named conjecture is claimed solved.

31.5 What the Newton-rigidity part does not claim

Every limitation stated by source 19, in its manuscript, README, audit files or verification record, is kept.

- (R1) No all-order rigidity with an order unit is claimed (and none holds: Theorem H); the unrestricted order-unit question was not resolved by source 19.
- (R2) No meromorphic order-unit analogue is proved: for quotients of entire series with an order unit nothing beyond Corollary 26.11 is claimed.
- (R3) All-order differential independence is not inferred from three-jet independence.
- (R4) No independence statement concerns point values: the relations are formal, and $f(x) \in \mathbb{K}$ for every x .
- (R5) No omnific factorization or arithmetic structure theorem is claimed; the omnific consequence concerns finitely many coefficients of formal relations.
- (R6) No uniform degree bound for polynomial solutions of a fixed equation is claimed; (16.1) has solutions of every degree, and the systems results assert neither existence, nor finiteness, nor bounded degrees of polynomial solutions.
- (R7) Entireness is relative to one Hahn field; nothing is claimed about entireness on the proper class $\mathbf{No}[i]$, where no nonpolynomial series is entire (Proposition 27.6).
- (R8) D_z acts on the external variable and kills every scalar; it is not the Berarducci–Mantova derivation, and no theorem about that derivation is claimed.
- (R9) External constants change the coefficients allowed in a relation, not the domain of strong summation.

- (R10) The first-polar criterion and the factor criterion are sufficient, not necessary (Theorems 31.1 and 31.2).
- (R11) H_3 , Q_4 and F_{sp} are not counterexamples to polynomial rigidity; an obstruction to a proof technique is not a counterexample.
- (R12) The 259 finite assertions check polynomial identities and finite Newton windows; they certify nothing about infinite supports, cofinality, arbitrary fields or novelty, and a large count is no substitute for the written proofs.
- (R13) No Lean proof or checked Lean theorem is supplied, the repository was not built by source 19, its formalization claims are not certificates for these results, and no independent referee report or historical priority is claimed.
- (R14) The repository and literature comparisons were targeted, not exhaustive, as recorded in Section 31.4.
- (R15) Positive characteristic and mixed characteristic are excluded: the proofs use $v(n) = 0$ for nonzero integers, and the theorems are not statements about $\mathfrak{p}$ -adic analytic fields.
- (R16) *[merge]* Remark 30.11 with its cubic, Corollary 31.4, the reading of Example 31.3 for Question 19.11, and the comparison of Theorem 30.16 with Corollary 26.10 are the merge's; no historical priority is claimed for them.

32 Independent theta scales and unbounded differential order

This section and the next contain the material of source 16, delivered in batch 31 with sources 15, 17, 18 and 19, and merged after the order-three-threshold part. They take source 13's descended theta series as given and ask how many differentially algebraic strongly entire series there are, and how large their minimal orders can be. Over $\mathbb{K} = k((t^{\mathbb{R}}))$ with $k = \mathbb{R}$ or $\mathbb{C}$ the answer is: the descents $\mathcal{T}_{t^\mu}$ for scales μ with $\mathbb{Q}$ -independent reciprocals are algebraically independent, a family of cardinality $2^{\aleph_0}$; polynomially weighted sums of them have arbitrarily large minimal differential order; and the field generated by all differentially algebraic strongly entire series has transcendence degree $2^{\aleph_0}$ and differential transcendence degree zero (Theorem L). The mechanism is the exact quadratic Gauss profile of $\mathcal{T}_{t^\mu}$ together with a finite-grid compression lemma of differential algebra. Source 16 also gives a third route to the minimal order three of $\mathcal{T}_p$ (Section 33.3).

32.1 Source, conventions and the merge's choices

Convention 32.1 (Setting of the theta-hierarchy part). In Sections 32–33, unless a statement says otherwise, $\Gamma = \mathbb{R}$, k is $\mathbb{R}$ or $\mathbb{C}$, $\mathbb{K} = k((t^{\mathbb{R}}))$, and $\mathcal{E}_\Gamma(k)$, $\text{Frac } \mathcal{E}_\Gamma(k)$, v , res , D_z and the Gauss value $\gamma_\delta(f) = \min_n(v(a_n) - n\delta)$ of (14.1), here for real δ and extended multiplicatively to $\text{Frac } \mathcal{E}_\Gamma(k)$, are as in Conventions 14.1 and 20.1. For a real $\mu > 0$ the series $\mathcal{T}_{t^\mu}$ is (1.4) with $p = t^\mu$; every positive real is an order unit of $\mathbb{R}$. The abstract results of Section 32.3 hold in any differential field of characteristic zero; there ∂ is the derivation and d the universal derivation of Kähler differentials, which are unrelated to the partial derivatives ∂_i of Section 6.

The source. Source 16 is *Independent Theta Families and Unbounded Differential Order in Hahn–Surcomplex Analysis: Valuation growth, bounded polynomial jet compression, and a continuum-sized*

differential-algebraic function field (25 pages, A4, 23 September 2026), delivered as the archive `Surreal_Theta_Hierarchy` (batch 31, manuscript 02). It records the reference snapshot

`0fffc26c670c15c7db451ed899f98d4384f9df1e`,

taken while the repository was changing. At that snapshot this report’s article and README were the five-source text (sources 08–12), and source 13’s two audit files had been placed in this directory without its manuscript being merged; source 16 read those audits and took the seed, its equation and its minimal order three from them, crediting all three to source 13. Its description was accurate at the snapshot; its statement that “the repository explicitly leaves the order-two alternative unresolved” and its Question 11.2 are stale now, answered by Theorem I. Source 16 works over $\Gamma = \mathbb{R}$ only, with $k = \mathbb{C}$ and, by the same proofs, $k = \mathbb{R}$; its asymptotics are real-valued, and no statement of this part is transferred to other value groups.

Renamed symbols. The following symbols of source 16 would collide with symbols of the earlier parts. They are renamed here, in the new material only.

Source 16	Here	Reason
function variable X ; $\partial = d/dX$; $\mathbb{F} = \mathbb{K}(X)$	z ; D_z ; $\mathbb{K}(z)$	The notation of Convention 14.1; X is the variable of initial polynomials.
$\mathbb{K} = \mathbb{C}((t^{\mathbb{R}}))$, $\mathbb{K}_{\mathbb{R}} = \mathbb{R}((t^{\mathbb{R}}))$; $\mathcal{E}_{\mathbb{K}}$	$\mathbb{K}$ with $k = \mathbb{C}, \mathbb{R}$; $\mathcal{E}_{\Gamma}(k)$	The same objects; only the real-coefficient field is renamed.
scale h ; $q = t^h$; $Q = q^2$; s_k	μ ; $p = t^{\mu}$; p^2 ; $\mathcal{Q}_k(p^2)$	As in Section 23.1; q is a dilation parameter.
F_h	$\mathcal{T}_{t^{\mu}}$	The series (1.4); F is (11.2).
Laurent variable z ; $\Theta = \sum q^{n^2} z^n$; $D = z d/dz$	u ; Ψ_p ; ϑ_u	Section 24.5; Θ_p is the partial theta series.
$G = D\Theta/\Theta$; $H = -DG - 2s_1$; $Y = z - z^{-1}$	ψ_1 ; $x = -\psi_2 - 2\mathcal{Q}_1(p^2)$; $\tilde{z}$	G, H have other meanings; x is a Tate x -coordinate.
$\Delta = X^2 - 4$; A, B of its (3.3)	$z^2 - 4$; $\mathcal{B}[f] - \frac{1}{12}f^2$, $\mathcal{H}[f]$	Δ_{μ} is a convex subgroup; (24.5), (24.6).
a_4, a_6 (Tate)	kept	Related to g_2, g_3 in Section 32.1.
radius parameter r ; w_r	δ ; γ_{δ}	The Gauss data of Section 14.3; r is a derivative order.
$\text{ord}_{\partial}(f)$	$\text{dord}(f)$	ord_z is the z -adic order.
$\mathfrak{c} = 2^{\aleph_0}$	$2^{\aleph_0}$	$\mathfrak{c}$ is a theta constant.
$F \subseteq L$, s , a_i , $p_i(s)$, c_{ij} (compression)	$\mathfrak{K}_0 \subseteq \mathfrak{K}_1$, $\mathfrak{s}$, y_i , $P_i(\mathfrak{s})$, $\mathfrak{c}_{ij}$	L is a coefficient subfield, a_n Taylor coefficients, $p = t^{\mu}$.
W , U_{ir} , λ_i	W , U_{ir} , ε_i	W, U are indeterminates, λ a multiplier.
primes p_i	p_i	$p = t^{\mu}$.
$\mathbf{m}$, $p_{\mathbf{m}}$, $A_{\mathbf{m}}$, $B_{\mathbf{m}}$, $\delta_{\mathbf{m}}$, $b_{\mathbf{m}}$, $c_{\mathbf{m}}$, R_0 , R	$\mathbf{e}$, $P_{\mathbf{e}}$, $\kappa_{\mathbf{e}}$, $\kappa'_{\mathbf{e}}$, $\kappa_* - \kappa_{\mathbf{e}}$, $\bar{d}_{\mathbf{e}}$, $\bar{\alpha}_{\mathbf{e}}$, $\delta^{(0)}$, δ^{cert}	Radii are scales δ here.
Hamel set B	$\mathfrak{B}$	B is a threshold.
$\mathcal{A}$, $\mathcal{D}$	$\mathcal{A}_{\text{DA}}$, $\mathcal{D}_{\text{DA}}$	$\mathcal{A}[f]$ is a theta expression.
G , $\mu = w_0(G)$, $\bar{G}$ (sampling)	g , $\gamma_0(g)$, ϕ_0	ϕ_0 is the initial polynomial at the scale 0.
Laurent-entire ring $\mathcal{L}$; w ; σ	$\mathcal{U}_{\mu}$; $\gamma^{\mathcal{U}}$; $\mathbf{S}$	Section 24.5.
Theorems 1.1–1.3	Theorem L	Theorems A–N of this report are different theorems.

The shipped files `16-theta-hierarchy-PROOF_AUDIT.md` and `16-theta-hierarchy-SOURCES_AND_SCOPE.md` keep the source’s notation.

Words with two meanings. The *minimal differential order* of f , written $\text{dord}(f)$, is the least r with $f, D_z f, \dots, D_z^r f$ algebraically dependent over $\mathbb{K}(z)$; it is the least order of Section 23.1, not the degree of an equation and not the growth order of a complex entire function. A *radius* $t^{-\delta}$ is a Hahn monomial, not a real radius. *Compression* is the finite-grid lemma of Section 32.3, not a primitive-element theorem. *Independent* without qualifier means algebraically independent over $\mathbb{K}(z)$ *without* derivatives; differential independence is stated as such.

Results printed once. Source 16 reproves several facts already in the report. They are printed once and cited: its escape lemma (Lemma 2.1), which is Lemma 2.3 (a); its definition of entireness (Definition 2.2), which is Definition 1.2 in the form of Proposition 20.2; closure and evaluation (Proposition 2.3), which is Proposition 2.7; Gauss multiplicativity (Lemma 2.5), which is Corollary 14.3; the coefficients of the seed (Proposition 3.1), which are Proposition 23.3; its product (3.4), which is (23.12); its explicit equation (Proposition 3.2), which is (24.7) (see below); its Proposition 9.1, the failure at ω^ω , which is in Section 25.3; its Laurent ring and shift (Appendix A.1), which are Lemma 24.4 and (23.10); its Lemma A.1, whose proof is that of Lemma 4.1 over any valued field of residue characteristic zero; and the embedding (2.4).

Source 16 states the seed equation in the Tate normalization, with the constants $a_4 = -5\mathcal{Q}_3(p^2)$, $a_6 = -(5\mathcal{Q}_3(p^2) + 7\mathcal{Q}_5(p^2))/12$ and $A = \mathcal{B}[f] - f^2/12$:

$$(z^2 - 4)(f D_z A - 2D_z f A)^2 = 4A^3 + A^2 f^2 + 4a_4 A f^4 + 4a_6 f^6.$$

Since $f D_z A - 2D_z f A = -\mathcal{K}[f]$, $g_2 = 1/12 - 4a_4$ and $g_3 = -1/216 + a_4/3 - 4a_6$, this is (24.7) as a polynomial identity in z and the jets (the merge's check, with SymPy), and its leading coefficient $(z^2 - 4)^3 f^4$ in $(D_z^3 f)^2$ agrees with Theorem 24.1. The Tate cubic $y^2 + xy = x^3 + a_4 x + a_6$ is the Weierstrass form after the shift $x \mapsto x - 1/12$; source 16 imports it from the course notes [53], as source 13 did.

Where each numbered result went. Theorems 1.1–1.3 are Theorem L (a)–(c); Lemma 2.6 is Lemma 32.2; Theorems 4.1, 4.2, Proposition 4.4 and Example 4.5 are Theorems 32.3, 32.4, Proposition 32.5 and Example 32.6; Lemmas 5.1, 5.2, Theorem 5.3, Remark 5.4, Example 5.5 and Corollary 5.6 are Lemmas 32.7, 32.8, Theorem 32.9, Remark 32.10, Example 32.11 and Corollary 32.12; Definition 6.1, Lemma 6.2, Corollaries 6.3, 6.4, Remark 6.5 and Theorem 6.6 are Definition 32.13, Lemma 32.14, Corollaries 32.15, 32.16, Remark 32.17 and Theorem 32.19; Lemmas 7.1, 7.3 and Corollary 7.2 are Lemmas 33.1, 33.3 and Corollary 33.2; Theorem 8.1 and Corollaries 8.2, 8.3 are Theorem 33.5 and Corollaries 33.6, 33.7; Theorem A.2 and Corollary A.3 are Theorem 33.8 and Corollary 33.9; Questions 11.1 and 11.3–11.10 are Questions 19.36–19.44, and Question 11.2 is answered (Remark 32.18); Appendix B is merged into the non-claims of Section 33.5 and the dependency table of Section 18.

The merge's choices. The review of source 16 for this merge found its proofs correct. The merge adds four points, each marked. First, the identity of the two seed equations above. Second, Remark 32.18: with Theorems I and K, the minimal orders of nonpolynomial differentially algebraic strongly entire series over $\mathbb{K}$ form a subset of $\{3, 4, \dots\}$ that contains 3 and is unbounded, and the compressed series of Theorem L (b) have order at least $\max(N, 3)$. Third, Remark 33.4 compares Theorem L (c) with Theorem 22.6 and with source 17's Corollary 35.3. Fourth, Remark 33.10 compares the three routes to minimal order three. Source 16's theorems, proofs, examples and non-claims are otherwise printed as delivered, in the renamed notation.

32.2 Gauss profiles and independence across scales

Lemma 32.2 (Polynomial growth; source 16). *If $P = \sum_{i=0}^d c_i z^i \in \mathbb{K}[z]$ with $c_d \neq 0$, then $\gamma_\delta(P) = v(c_d) - d\delta$ for all sufficiently large real δ ; for a nonzero rational function γ_δ is eventually an affine function of δ .*

Proof. For each $i < d$ with $c_i \neq 0$ the top term has strictly smaller weighted value once $\delta > (v(c_d) - v(c_i))/(d - i)$; there are finitely many such inequalities. Subtract for numerator and denominator. $\square$

Theorem 32.3 (Exact quadratic profile; source 16). *For real $\mu > 0$ and $\delta \geq 0$,*

$$\gamma_\delta(\mathcal{T}_{t^\mu}) = \min_{n \geq 0} (\mu n^2 - \delta n) = -\frac{\delta^2}{4\mu} + \mu \operatorname{dist}\left(\frac{\delta}{2\mu}, \mathbb{Z}_{\geq 0}\right)^2,$$

so $-\delta^2/(4\mu) \leq \gamma_\delta(\mathcal{T}_{t^\mu}) \leq -\delta^2/(4\mu) + \mu/4$; for $\delta < 0$, $\gamma_\delta(\mathcal{T}_{t^\mu}) = 0$.

Proof. By (23.6), $v(a_n) = \mu n^2$. Complete the square, $\mu n^2 - \delta n = \mu(n - \delta/(2\mu))^2 - \delta^2/(4\mu)$, and use that a nonnegative real lies within $\frac{1}{2}$ of a nonnegative integer. For $\delta < 0$ the expression is positive for $n > 0$ and zero at $n = 0$. $\square$

The corners are at $\delta = (2n + 1)\mu$, where n and $n + 1$ are active: these are the breaks δ_n of Theorem 27.5, with $\gamma_{\delta_n} = -n(n + 1)\mu$. The bounded oscillation lets distinct growth constants be compared uniformly, without special sequences of scales.

Theorem 32.4 (Superlinear growth separation; source 16). *Let $(f_i)_{i \in I}$ be nonzero elements of $\mathcal{E}_\Gamma(k)$ and $\varpi > 1$ real, and suppose $\gamma_\delta(f_i) = -c_i \delta^\varpi + o(\delta^\varpi)$ as $\delta \rightarrow +\infty$, with positive reals c_i linearly independent over $\mathbb{Q}$. Then $(f_i)_{i \in I}$ is algebraically independent over $\mathbb{K}(z)$.*

Proof. A relation involves finitely many f_i ; clear denominators to get $\sum_{\mathbf{e} \in E} P_{\mathbf{e}}(z) \prod_{i=1}^N f_i^{e_i} = 0$ with $E \subset \mathbb{Z}_{\geq 0}^N$ finite and every $P_{\mathbf{e}} \neq 0$. By Corollary 14.3 and Lemma 32.2 the summand indexed by $\mathbf{e}$ has $\gamma_\delta = -(\sum_i e_i c_i) \delta^\varpi + o(\delta^\varpi) + O(\delta)$. The numbers $\sum_i e_i c_i$ are pairwise distinct, by the $\mathbb{Q}$ -independence, so one is largest, and since $\varpi > 1$ its summand has strictly the least Gauss value for all large δ ; a finite sum with a unique least term is nonzero. $\square$

Proof of Theorem L(a). Apply Theorem 32.4 with $\varpi = 2$ and $c_i = 1/(4\mu_i)$, using the bounds of Theorem 32.3. Each $\mathcal{T}_{t^{\mu_i}}$ is a nonpolynomial element of $\mathcal{E}_\Gamma(k)$ of minimal order three (Theorem H); the cardinality assertion is Corollary 33.2. $\square$

The coefficients of a relation may be arbitrary elements of $\mathbb{K}$: a fixed polynomial in z has finitely many coefficients, each with one valuation, so Lemma 32.2 still gives only linear growth. That is why the quadratic separation survives the whole Hahn coefficient field.

Proposition 32.5 (An explicit noncancellation threshold; source 16). *In a relation as in the proof of Theorem 32.4 with $f_i = \mathcal{T}_{t^{\mu_i}}$, put $\kappa_{\mathbf{e}} = \frac{1}{4} \sum_i e_i / \mu_i$, $\kappa'_{\mathbf{e}} = \frac{1}{4} \sum_i e_i \mu_i$, $d_{\mathbf{e}} = \deg P_{\mathbf{e}}$ and $\alpha_{\mathbf{e}} = v(\operatorname{lc} P_{\mathbf{e}})$, let $\mathbf{e}_*$ be the unique maximizer of $\kappa_{\mathbf{e}}$, write $\kappa_*, d_*, \alpha_*, \kappa'_*$ for its values, and let $\delta^{(0)}$ be the largest of 0 and the numbers $(\alpha_{\mathbf{e}} - v(c_{\mathbf{e},i})) / (d_{\mathbf{e}} - i)$ over all coefficients $c_{\mathbf{e},i} \neq 0$ of $P_{\mathbf{e}} = \sum_i c_{\mathbf{e},i} z^i$ with $i < d_{\mathbf{e}}$. For $\mathbf{e} \neq \mathbf{e}_*$ put $\bar{d}_{\mathbf{e}} = |d_* - d_{\mathbf{e}}|$ and $\bar{\alpha}_{\mathbf{e}} = |\alpha_{\mathbf{e}} - \alpha_*| + \kappa'_*$. Then every real $\delta \geq \delta^{\operatorname{cert}}$, where*

$$\delta^{\operatorname{cert}} = 1 + \max\left(\{\delta^{(0)}\} \cup \left\{ \frac{2\bar{d}_{\mathbf{e}}}{\kappa_* - \kappa_{\mathbf{e}}}, \sqrt{\frac{2\bar{\alpha}_{\mathbf{e}}}{\kappa_* - \kappa_{\mathbf{e}}}} : \mathbf{e} \neq \mathbf{e}_* \right\}\right),$$

makes the $\mathbf{e}_*$ summand the unique one of least Gauss value; in particular the expression is nonzero.

Proof. For $\delta > \delta^{(0)}$ the summand $\mathbf{e}$ has Gauss value $-\kappa_{\mathbf{e}}\delta^2 - d_{\mathbf{e}}\delta + \alpha_{\mathbf{e}} + \epsilon_{\mathbf{e}}(\delta)$ with $0 \leq \epsilon_{\mathbf{e}}(\delta) \leq \kappa'_{\mathbf{e}}$, by Theorem 32.3. Its difference from the $\mathbf{e}_*$ value is at least $(\kappa_* - \kappa_{\mathbf{e}})\delta^2 - \bar{d}_{\mathbf{e}}\delta - \bar{\alpha}_{\mathbf{e}}$, and the two conditions in δ^{cert} make each of the last two terms smaller than half of the first. A single summand needs only $\delta^{(0)}$. $\square$

This certifies a specified polynomial expression; it is not a decision procedure for differential relations.

Example 32.6 (Two explicit scales; source 16). Take $\mu_1 = 1$, $\mu_2 = 1/\sqrt{2}$ and $P = Y_1^2 + (z+t^{-7})Y_2 + 1$. The growth constants of the three summands are $\frac{1}{2}$, $\frac{\sqrt{2}}{4}$ and 0, so the Y_1^2 summand dominates; $\delta^{(0)} = 7$, and $\delta \geq 16$ is a valid certificate (the exact value is about 14.7). So $\mathcal{T}_t^2 + (z+t^{-7})\mathcal{T}_{t^{1/\sqrt{2}}} + 1 \neq 0$, with the infinite coefficient t^{-7} allowed.

32.3 Bounded polynomial jet compression

The valuation argument yields independent *functions*, not one function of high order; a plain sum can lose information. The following is pure differential algebra, related in purpose to differential primitive-element theorems [34, 35] but different in conclusion: it keeps a prescribed number of independent jets, with an explicit grid of integer weights, and does not claim a generator of the whole extension.

Lemma 32.7 (Kähler criterion; source 16). *Let $\mathfrak{K}_0 \subseteq \mathfrak{K}_1$ be fields of characteristic zero and $y_1, \dots, y_N \in \mathfrak{K}_1$. They are algebraically independent over $\mathfrak{K}_0$ if and only if $dy_1 \wedge \dots \wedge dy_N \neq 0$ in $\bigwedge_{\mathfrak{K}_1}^N \Omega_{\mathfrak{K}_1/\mathfrak{K}_0}$.*

Proof. If they are independent, extend them to a transcendence basis; the coordinate derivations of the rational function field extend to $\mathfrak{K}_1$ by separability and give a dual system to the dy_i . Conversely, if they are dependent, every y outside a maximal independent subfamily is algebraic over it, and differentiating its minimal polynomial (nonzero derivative by separability) makes dy a combination of the others, so the wedge is zero. $\square$

Lemma 32.8 (Finite-grid nonvanishing; source 16). *A nonzero polynomial in M variables over a field of characteristic zero, of degree at most N in each variable, is nonzero at some point of $\{0, 1, \dots, N\}^M$.*

Proof. For $M = 1$ this is the root bound. Otherwise write the polynomial in the last variable; vanishing on the grid makes every specialization of the first $M - 1$ variables have $N + 1$ roots, so every coefficient vanishes on the smaller grid, and induction applies. $\square$

Theorem 32.9 (Bounded polynomial jet compression; source 16). *Let $(\mathfrak{K}_0, \partial) \subseteq (\mathfrak{K}_1, \partial)$ be differential fields of characteristic zero, $\mathbf{s} \in \mathfrak{K}_0$ with $\partial \mathbf{s} = 1$, and $y_1, \dots, y_N \in \mathfrak{K}_1$ algebraically independent over $\mathfrak{K}_0$. There are $P_i(\mathbf{s}) = \sum_{j=0}^{N-1} c_{ij} \mathbf{s}^j$ with $c_{ij} \in \{0, 1, \dots, N\}$ such that for $g = \sum_i P_i(\mathbf{s}) y_i$ the elements $g, \partial g, \dots, \partial^{N-1} g$ are algebraically independent over $\mathfrak{K}_0$. At most $(N + 1)^{N^2}$ coefficient arrays need be considered.*

Proof. Treat the c_{ij} first as indeterminates, constant for ∂ and killed by d , which is relative to $\mathfrak{K}_0$. The wedge

$$W(c) = dg \wedge d(\partial g) \wedge \cdots \wedge d(\partial^{N-1}g) \in \left(\bigwedge_{\mathfrak{K}_1}^N \Omega_{\mathfrak{K}_1/\mathfrak{K}_0} \right)[c]$$

has total degree N , each factor being linear in the parameters. Replace them by the jet parameters $U_{ir} = \partial^r P_i(s)$, $0 \leq r < N$; this change is invertible over $\mathfrak{K}_0$, triangular with the factorials $0!, \dots, (N-1)!$ on the diagonal, so it suffices to show $W \neq 0$ in the new variables. By Leibniz,

$$d(\partial^j g) = \sum_{i=1}^N \sum_{r=0}^j \binom{j}{r} U_{ir} d(\partial^{j-r} y_i),$$

with no differentials of the U_{ir} , whose specialized values lie in $\mathfrak{K}_0$. Specialize $U_{ir} = \varepsilon_i$ if $r = i - 1$ and 0 otherwise, with new indeterminates ε_i . The j th factor then involves only $\varepsilon_1, \dots, \varepsilon_{j+1}$, so the squarefree monomial $\varepsilon_1 \cdots \varepsilon_N$ arises only when the j th factor supplies $\varepsilon_{j+1} dy_{j+1}$, with binomial coefficient one; its coefficient is $dy_1 \wedge \cdots \wedge dy_N \neq 0$ by Lemma 32.7. So W is a nonzero polynomial. A $\mathfrak{K}_1$ -linear functional on the finite span of its coefficients not killing a nonzero one gives a nonzero scalar polynomial of degree at most N in each of the N^2 variables; Lemma 32.8 gives a point of $\{0, \dots, N\}^{N^2}$ where it, hence W , is nonzero, and Lemma 32.7 concludes. $\square$

Remark 32.10 (Why the formal jet specialization is legitimate; source 16). The specialization $U_{ir} = \varepsilon_i \mathbf{1}_{r=i-1}$ only proves that a polynomial is nonzero; it is not a claim that one polynomial in s has independently prescribed derivatives. Invertibility of the coefficient-to-jet change over $\mathfrak{K}_0$ is what makes the test valid. The final specialization is in the original coefficients, where the grid gives actual integer polynomials.

Example 32.11 (The first two cases; source 16). For $N = 1$ the grid contains $g = y_1$. For $N = 2$, among the 81 candidates $(c_{10} + c_{11}s)y_1 + (c_{20} + c_{21}s)y_2$ with $c_{ij} \in \{0, 1, 2\}$ at least one has $g, \partial g$ independent; an unweighted sum $y_1 + y_2$ is not asserted to work. The grid is uniform: it works for every independent pair in the stated setting.

Corollary 32.12 (An entire-preserving version; source 16). *If $y_1, \dots, y_N \in \mathcal{E}_\Gamma(k)$ are algebraically independent over $\mathbb{K}(z)$, one of the candidates of Theorem 32.9 with $s = z$ is strongly entire and has its first N jets algebraically independent over $\mathbb{K}(z)$; with real Hahn coefficients, so does the candidate.*

Proof. Apply the theorem in $\text{Frac } \mathcal{E}_\Gamma(k)$; polynomial weights and finite sums preserve $\mathcal{E}_\Gamma(k)$ (Proposition 2.7), and integer weights preserve the real subfield. $\square$

32.4 Minimal differential order and the unbounded hierarchy

Definition 32.13 (Minimal differential order; source 16). For a differential field extension $\mathfrak{K}_0 \subseteq \mathfrak{K}_1$ and $f \in \mathfrak{K}_1$ differentially algebraic over $\mathfrak{K}_0$, put $\text{dord}(f/\mathfrak{K}_0) = \min\{r \geq 0 : f, \partial f, \dots, \partial^r f \text{ are algebraically dependent over } \mathfrak{K}_0\}$; write $\text{dord}(f)$ when $\mathfrak{K}_0 = \mathbb{K}(z)$, and $\mathfrak{K}_0\langle f \rangle$ for the differential field generated by f .

A rational function has order zero. For a nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ the number $\text{dord}(f)$ is the least order of Theorem I and Section 23.1.

Lemma 32.14 (Order equals transcendence degree; source 16). *In characteristic zero, if f is differentially algebraic over $\mathfrak{K}_0$ and $r = \text{dord}(f/\mathfrak{K}_0)$, then $\mathfrak{K}_0\langle f \rangle = \mathfrak{K}_0(f, \partial f, \dots, \partial^r f)$ and its transcendence degree over $\mathfrak{K}_0$ is r .*

Proof. $f, \dots, \partial^{r-1}f$ are independent and $\partial^r f$ is algebraic over the field they generate; differentiate its minimal polynomial, whose derivative at $\partial^r f$ is nonzero by separability, and solve for $\partial^{r+1}f$ inside $\mathfrak{K}_0(f, \dots, \partial^r f)$, which is therefore closed under ∂ . This is the argument of Theorem 30.16 and of the finite differential fields of Theorem 24.10; it is standard. $\square$

Corollary 32.15 (Finite composita; source 16). *If $f_1, \dots, f_N$ are differentially algebraic over $\mathfrak{K}_0$ of orders $r_1, \dots, r_N$, then $\mathfrak{K}_0\langle f_1, \dots, f_N \rangle$ is a finitely generated field extension of transcendence degree at most $r_1 + \dots + r_N$, and each of its elements is differentially algebraic of order at most that transcendence degree.*

Proof. The compositum of the fields of Lemma 32.14 is closed under ∂ ; any $d+1$ jets of an element of a field of transcendence degree d are dependent. $\square$

Proof of Theorem L(b). Take $\mu_i = 1/\sqrt{p_i}$ for distinct primes p_i . The $\mathcal{T}_{t^{\mu_i}}$ are independent by part (a), and Corollary 32.12 gives an entire candidate g with N independent jets, so $\text{dord}(g) \geq N$. Each seed has an equation of order three (Theorem 24.1), so by Corollary 32.15 g is differentially algebraic with $\text{dord}(g) \leq 3N$; being of order at least $N \geq 1$ it is not rational, hence not a polynomial. $\square$

The square roots of distinct primes are linearly independent over $\mathbb{Q}$: in the multiquadratic field $\mathbb{Q}(\sqrt{p_1}, \dots, \sqrt{p_N})$ the sign of any one square root can be changed while the others are fixed, and applying such a change to a linear relation and subtracting isolates a coefficient. No nonconstructive choice is needed for finite N .

Corollary 32.16 (No finite upper bound; source 16). *For every $M \geq 0$ some strongly entire differentially algebraic series over $\mathbb{K}$, for $k = \mathbb{R}$ and for $k = \mathbb{C}$, has minimal differential order greater than M .*

Remark 32.17 (An important limitation; source 16). The interval $[N, 3N]$ does not show that every integer in it occurs; $N = 2$ does not produce an entire series of order two, and the argument does not prove that the first three jets of all the seeds are jointly independent.

Remark 32.18 (Order two, and the known part of the spectrum; merge). Source 16's Question 11.2, whether a nonpolynomial strongly entire series of minimal order two exists, is answered negatively by Theorem I(a), for every Γ . Hence over $\mathbb{K}$ the minimal orders of nonpolynomial differentially algebraic strongly entire series form a subset of $\{3, 4, 5, \dots\}$ that contains 3 (Theorem H) and is unbounded (Corollary 32.16); the compressed series of Theorem L(b) have order at least $\max(N, 3)$, and by Theorem K every equation of order three that such a series satisfies has total jet degree at least four. The dimension d_N of Theorem 32.19 is at least $\max(N, 3)$. Which orders above three occur stays open (Question 19.36).

Theorem 32.19 (Exact order at the compositum dimension; source 16). *For scales with $\mathbb{Q}$ -independent reciprocals let $\mathfrak{L}_N = \mathbb{K}(z)\langle \mathcal{T}_{t^{\mu_1}}, \dots, \mathcal{T}_{t^{\mu_N}} \rangle$ and $d_N = \text{trdeg}_{\mathbb{K}(z)} \mathfrak{L}_N$. Then $N \leq d_N \leq 3N$, and there is $u_N \in \mathfrak{L}_N \cap \mathcal{E}_\Gamma(k)$ with $\text{dord}(u_N) = d_N$: a bounded polynomial combination, with weights of degree below d_N and coefficients in $\{0, \dots, d_N\}$, of a transcendence basis chosen from $\{\mathcal{T}_{t^{\mu_i}}, D_z \mathcal{T}_{t^{\mu_i}}, D_z^2 \mathcal{T}_{t^{\mu_i}}\}_{i \leq N}$. Moreover $\mathfrak{L}_N$ is a finite algebraic extension of $\mathbb{K}(z)\langle u_N \rangle$.*

Proof. Each third derivative is algebraic over the field of the displayed finite list (Theorem 24.1), so by Lemma 32.14 $\mathfrak{L}_N$ is finite over that field and a maximal independent subfamily of the list is a transcendence basis, of size d_N . Compress it (Theorem 32.9) inside $\mathfrak{L}_N$: the result has d_N

independent jets and, by Corollary 32.15, order at most d_N . The two fields have the same finite transcendence degree, and $\mathfrak{L}_N$ is finitely generated over the smaller one. $\square$

This realizes an intrinsically defined dimension as an exact order without computing d_N , and does not identify u_N as a differential primitive element.

33 Continuum fields, integer sampling, a third route to order three, and scope

33.1 Continuum many independent differentially algebraic series

Lemma 33.1 (A positive Hamel set; source 16). *There is $\mathfrak{V} \subset \mathbb{R}_{>0}$ of cardinality $2^{\aleph_0}$ linearly independent over $\mathbb{Q}$.*

Proof. Take a Hamel basis of $\mathbb{R}$ over $\mathbb{Q}$ and replace negative elements by their negatives; an infinite-dimensional space over a countable field has cardinality equal to its dimension, so the basis has size $2^{\aleph_0}$. $\square$

Corollary 33.2 (The independent family; source 16). *$\{\mathcal{T}_{t^{1/b}} : b \in \mathfrak{V}\}$ is algebraically independent over $\mathbb{K}(z)$, consists of nonpolynomial strongly entire series of minimal order three, and has cardinality $2^{\aleph_0}$.*

Proof. Every finite subfamily satisfies Theorem L (a). $\square$

The finite examples use explicit prime scales; the uncountable family uses the axiom of choice through a Hamel basis, and no continuum hypothesis or large-cardinal axiom.

Let $\mathcal{A}_{\text{DA}} \subseteq \mathcal{E}_{\Gamma}(k)$ be the set of strongly entire series that are differentially algebraic over $\mathbb{K}(z)$. By Proposition 2.7 and Corollary 32.15 it is a differential subring containing $\mathbb{K}[z]$, and its fraction field is the differential field $\mathcal{D}_{\text{DA}} \subseteq \text{Frac } \mathcal{E}_{\Gamma}(k)$ of Theorem L (c).

Lemma 33.3 (Cardinality; source 16). $|\mathbb{K}| = |\mathcal{E}_{\Gamma}(k)| = 2^{\aleph_0}$.

Proof. A well-ordered subset of $\mathbb{R}$ is countable (choose a rational between each element and its successor), so there are at most $(2^{\aleph_0})^{\aleph_0}$ supports and coefficient assignments; a power series is a countable sequence of elements of $\mathbb{K}$. Constants give the lower bounds. $\square$

Proof of Theorem L (c). Corollary 33.2 gives the lower bound $2^{\aleph_0}$ and Lemma 33.3 the upper bound for the transcendence degree. Each element of $\mathcal{D}_{\text{DA}}$ lies in a differential field generated by finitely many elements of $\mathcal{A}_{\text{DA}}$, so is differentially algebraic (Corollary 32.15); the differential transcendence degree is zero. Finitely many differential generators would give finite transcendence degree. $\square$

Algebraic independence concerns relations without derivatives; differential algebraicity allows relations involving derivatives of the same element. The family is as large as possible in the first sense and consists of constrained elements in the second; this difference is what compression uses.

Remark 33.4 (Three fields of entire series; merge). Theorem 22.6 gives $\text{dtrdeg } \text{Frac } \mathcal{E}_{\Gamma}(k) = 2^{\aleph_0}$ over $\mathbb{C}((t^{\Gamma^\infty}))$, a group without an order unit, where every nonpolynomial entire series is differentially transcendental. Over $\mathbb{K} = k((t^{\mathbb{R}}))$, which has an order unit, Theorem L (c) shows that the

differentially algebraic part $\mathcal{D}_{\text{DA}}$ of $\text{Frac } \mathcal{E}_\Gamma(k)$ is already of transcendence degree $2^{\aleph_0}$, and source 17's family (Theorem M with $p = t$) shows that $\text{Frac } \mathcal{E}_\Gamma(k)$ itself has $\text{dtrdeg}_{\mathbb{K}(z)} \text{Frac } \mathcal{E}_\Gamma(k) = 2^{\aleph_0}$ there too: its lower bound applies because 1 is an order unit of $\mathbb{R}$, and Lemma 33.3 gives the upper bound. Source 17 states this for $k((t^{\mathbb{Q}}))$ (Corollary 35.3); the case of $k((t^{\mathbb{R}}))$ is the merge's.

33.2 Sampling on ordinary integers

Theorem 33.5 (Residue-polynomial sampling; source 16). *Let $0 \neq g = \sum_n a_n z^n \in \mathcal{E}_\Gamma(k)$ and let $\phi_0 \in k[z]$ be its initial polynomial at the scale 0, $\phi_0 = \sum_n \text{res}(t^{-\gamma_0(g)} a_n) z^n$ (14.1). Then ϕ_0 is a nonzero polynomial, the zeros of g in $k \subseteq \mathbb{K}$ are zeros of ϕ_0 , and in particular $\#\{m \in \mathbb{Z} : g(m) = 0\} \leq \deg \phi_0$.*

Proof. By entireness at $\delta = 0$ only finitely many coefficients have the least value $\gamma_0(g)$, and at least one does. For $c \in k$, $v(c) \geq 0$, and the residue of the normalized value $t^{-\gamma_0(g)} g(c)$ is $\phi_0(c)$: terms of positive valuation contribute nothing, and finitely many of valuation zero remain. So $g(c) = 0$ forces $\phi_0(c) = 0$. $\square$

This is a quantitative form of Proposition 2.8, which says that a strongly entire series vanishing at infinitely many ordinary complex arguments is zero; here the number of such zeros is bounded by the degree of one initial polynomial.

Corollary 33.6 (No fixed relation on infinitely many integers; source 16). *For scales with $\mathbb{Q}$ -independent reciprocals and a nonzero $P \in \mathbb{K}[z, Y_1, \dots, Y_N]$, $P(m, \mathcal{T}_{t^{\mu_1}}(m), \dots, \mathcal{T}_{t^{\mu_N}}(m)) \neq 0$ for all but finitely many integers m ; for finitely many polynomials the exceptional set is common.*

Proof. $P(z, \mathcal{T}_{t^{\mu_1}}, \dots)$ is a nonzero element of $\mathcal{E}_\Gamma(k)$ (Theorem L (a), Proposition 2.7); apply Theorem 33.5. $\square$

Corollary 33.7 (Jet-sampling rigidity; source 16). *For a series g of Theorem L (b) and every nonzero $P \in \mathbb{K}[z, Y_0, \dots, Y_{N-1}]$, $P(m, g(m), D_z g(m), \dots, D_z^{N-1} g(m)) \neq 0$ for all but finitely many integers m .*

These statements concern a *fixed* polynomial evaluated at varying integers. They do not say that the values at one point are algebraically independent over $\mathbb{K}$: every value lies in $\mathbb{K}$, and $Y - \mathcal{T}_{t^\mu}(m)$ vanishes at the point m . Ordinary integers are omnific integers, so the corollaries give a concrete omnific test set against functional identities; they are not a classification of Diophantine equations over $\mathbf{Oz}$ and use no accumulation of integers in a surreal topology.

33.3 A third route to minimal order three

Source 16 reproves, for $\Gamma = \mathbb{R}$ and $k = \mathbb{C}$, the lower bound of Corollary 24.9, by a valuation argument. Let $p = t^\mu$ with real $\mu > 0$, let $\mathcal{U}_\mu$ and $\mathbf{S}$ be as in Lemma 24.4 (source 16's Laurent-entire ring is exactly $\mathcal{U}_\mu$ for $\Gamma = \mathbb{R}$), and let $\gamma^\mathcal{U}$ be the Gauss valuation of radius one, $\gamma^\mathcal{U}(\sum b_m u^m) = \min_m v(b_m)$; it is multiplicative on $\mathcal{U}_\mu$, since the normalized reduction of a product is the product of two nonzero Laurent polynomials, and extends to $\text{Frac } \mathcal{U}_\mu$, with residue characteristic zero and $\gamma^\mathcal{U}(u) = 0$. With $\Psi = \Psi_p$, $\psi_1 = \vartheta_u \Psi / \Psi$ and $\mathbf{x} = -\vartheta_u \psi_1 - 2\mathcal{Q}_1(p^2)$, (23.10) gives

$$\mathbf{S}^m \Psi = p^{-m^2} u^{-m} \Psi, \quad \mathbf{S} \psi_1 = \psi_1 - 1, \quad \mathbf{S} \mathbf{x} = \mathbf{x},$$

and $\gamma^{\mathcal{U}}(\Psi) = 0$, $x = -p(u + u^{-1}) + \text{terms of Gauss value at least } 2\mu$, so $\vartheta_u x = -p(u - u^{-1}) + \dots \neq 0$ and x is transcendental over $\mathbb{K}$ (an element algebraic over the constants $\mathbb{K}$ of ϑ_u has zero derivative). For a nonzero $R \in \text{Frac}\mathcal{U}_\mu[T]$, $\gamma^{\mathcal{U}}(R(m))$ is constant for all but finitely many $m \in \mathbb{N}$, by the proof of Lemma 4.1 (source 16's Lemma A.1).

Theorem 33.8 (Three theta coordinates; source 16). *Ψ , ψ_1 and x are algebraically independent over $\mathbb{K}(u)$.*

Proof. A relation can be written $\sum_{i,j} u^j \Psi^i P_{ij}(\psi_1, x) = 0$ with finitely many distinct pairs (i, j) , $i \geq 0$, and nonzero $P_{ij} \in \mathbb{K}[U, V]$. For each, $P_{ij}(\psi_1 - T, x)$ is a nonzero polynomial in T : its leading coefficient is, up to sign, the leading coefficient in U of $P_{ij}(U, x)$, a nonzero polynomial in the transcendental x . Apply S^m : the (i, j) summand becomes $p^{-im^2+2jm} u^{j-im} \Psi^i P_{ij}(\psi_1 - m, x)$, of Gauss value $-i\mu m^2 + 2j\mu m + \beta_{ij}$ for all large m , with β_{ij} fixed. The pair with the largest i and, among those, the smallest j has the uniquely least value for all large m , so the transformed sum is nonzero, a contradiction. $\square$

Corollary 33.9 (Minimal order three; source 16, credited conclusion). *$\mathcal{T}_{t^\mu}$, $D_z \mathcal{T}_{t^\mu}$ and $D_z^2 \mathcal{T}_{t^\mu}$ are algebraically independent over $\mathbb{K}(z)$, and $\text{dord}(\mathcal{T}_{t^\mu}) = 3$.*

Proof. Substitute $z = u + u^{-1}$, $\tilde{z} = u - u^{-1}$ (Lemma 24.5). With $f = \mathcal{T}_{t^\mu}$ one has $\Psi = f^\natural$, $\psi_1 = \tilde{z}(D_z f)^\natural / f^\natural$ and $x = -\tilde{z}^2(D_z^2 f / f - (D_z f / f)^2)^\natural - (u + u^{-1})(D_z f / f)^\natural - 2Q_1(p^2)$, by (24.14), an invertible rational change of the three coordinates over $\mathbb{K}(u)$; Theorem 33.8 excludes every relation, and Theorem 24.1 supplies one at the next jet. $\square$

The argument separates finitely many terms by the growth of iterates of a difference automorphism; it concerns functions, not values of theta functions.

Remark 33.10 (Three routes to minimal order three; merge). The lower bound of Corollary 24.9 now has three proofs. Source 13's (Theorem 24.8) uses the same shift S and the same three coordinates, and three field-theoretic obstructions on rational functions: fixed elements are constants, an additive difference equation has no rational solution, and S preserves u -adic order (Lemma 24.6), with the global certificate $\psi_3 \neq 0$ of Lemma 24.7. Source 16's replaces those three steps by one valuation argument: iterating S makes the Gauss values of the summands quadratic in the number of iterations, and a unique least one survives; the needed nonvanishing is $\vartheta_u x \neq 0$, read off from the leading term. Source 14's (Remark 26.14) uses no property of $\mathcal{T}_p$ beyond nonpolynomial strong entireness. Source 16's route is stated for $\Gamma = \mathbb{R}$ and $k = \mathbb{C}$ only, as delivered; the other two hold for every Γ with an order unit and every k of characteristic zero. All three are kept.

The surreal boundary and the derivation (source 16). Through $t^a \mapsto \omega^{-a}$ the field $\mathbb{R}((t^\mathbb{R}))$ is the surreal subfield with real normal-form exponents, and $p = t^\mu$ becomes $\omega^{-\mu}$; every real $\mu > 0$ is cofinal in $\mathbb{R}$, which is why quadratic coefficient values make these series entire on that field. At ω^ω the evaluation family of $\mathcal{T}_{t^\mu}$ is not strongly summable (Section 25.3), so nothing here is entire on **No** or **No**[i]. The derivative D_z fixes $\mathbb{K}$; an intrinsic surreal derivation such as that of Berarducci and Mantova [8] need not fix p , and replacing D_z by it changes the equations and invalidates the order calculations. Values at omnific inputs need not be omnific; the only omnific consequence claimed is the sampling statement above.

What the bounded grid makes effective (source 16). For given N the scales $1/\sqrt{p_i}$, the coefficient range and the degree bound are explicit, so all $(N+1)^{N^2}$ candidates can be listed. The proof guarantees that one has N independent jets but supplies no way to recognize it from finite truncations: W lives in a module of Kähler differentials of a function field without an effective presentation here, and Proposition 32.5 handles polynomial expressions in the seeds, not mixed jet expressions.

A formalization plan proposed by source 16. The most reusable lemma is Theorem 32.9, with no surreal input: the Kähler criterion, the invertible coefficient-to-jet matrix, the unique squarefree coefficient and grid nonvanishing. The analytic part needs the entire-series ring with its Gauss valuation, the coefficient values of the seed, a finite dominant-term lemma and the separant argument; the theta equation can first be a named imported theorem. No Lean code or certificate accompanies source 16.

33.4 Predecessors, repository comparison and priority

Literature. Source 16 consulted DLMF §20.5 and §23.6 [50, 51]; the Tate equation, coefficients and invariant differential on printed page 32 of the course notes hosted by M. Woodbury [53], inspected as an image (it attributes the notes to a course of J. Ellenberg; the hosting attribution does not identify the author of every part); Pogudin’s primitive-element papers [34, 35] for context; and the Mantova–Matusinski survey [17] for normal forms and intrinsic derivations. It consulted the Wikipedia article on surreal numbers for orientation only. No exhaustive MathSciNet or zbMATH search, citation-network review or referee report was made, and irrelevant keyword searches were not taken as evidence of novelty. A close variant of the compression argument, or a classical antecedent of the hierarchy in differential algebra or non-Archimedean function theory, may exist. For this merge the Pogudin preprints’ bibliographic data were checked; their content was not.

Repository comparison at the snapshot. Source 16 read, through a connector, the root and `docs/` inventories, this report’s directory and README and source 13’s two audit files, at a recorded snapshot `0fffc26` taken while the repository was changing; this is not an assertion that every read was pinned to that commit, nor an audit of all reports, archives, branches or commits. A search returning no match was not treated as evidence of absence. Its statements were accurate at the snapshot, as recorded in Section 32.1; its first research direction, theta descent as a counterexample to order-unit rigidity, was dropped when source 13’s audit showed it was already done.

Priority. The seed, its coefficient values, its equation and its minimal order three are source 13’s; the theta product and the Tate identity are classical and imported. The proposed contributions of source 16 are independence across reciprocal-independent scales with the exact profile and the finite-radius certificate, bounded polynomial jet compression, unbounded minimal order with the bounds $[N, 3N]$ and the exact-dimension refinement, the transcendence degrees of $\mathcal{D}_{\text{DA}}$, integer-sampling rigidity, and the valuation route to minimal order three; priority is provisional.

33.5 What the theta-hierarchy part does not claim

Every limitation stated by source 16, in its manuscript, README, audit files or verification record, is kept.

- (Th1) Not every candidate of Theorem L (b) is claimed to work, and no algorithm recognizes a working candidate.
- (Th2) The bound $3N$ is not shown sharp, d_N is not computed, and the full order spectrum is not determined; not every integer in $[N, 3N]$ is claimed to occur.
- (Th3) Order two is not decided by the case $N = 2$ (it is decided, negatively, by Theorem I, not by source 16).
- (Th4) Joint independence of all $3N$ seed jets is not proved.
- (Th5) The finite checks do not decide differential dependence, prove summability of infinite supports, algebraic independence of the full functions, the compression theorem, minimal orders or cardinalities.
- (Th6) Nothing is entire on the full class **No** or **No** $[i]$.
- (Th7) No point values are claimed algebraically independent over their own coefficient field.
- (Th8) The seed, its equation and its minimal order three are credited as prior (source 13); the theta product and the Tate identity are imported classical inputs.
- (Th9) Priority is provisional: a primitive-element-type antecedent of the compression lemma, or of the hierarchy, may exist.
- (Th10) No independent referee report and no Lean formalization.
- (Th11) The repository comparison was targeted, at a snapshot recorded while the repository was changing.
- (Th12) The continuum family uses choice (a Hamel basis), not the continuum hypothesis or a large-cardinal axiom.
- (Th13) The asymptotics are real-valued: the part is proved for $\Gamma = \mathbb{R}$ only, and the third route to order three for $\Gamma = \mathbb{R}$, $k = \mathbb{C}$.
- (Th14) No named historical conjecture is claimed solved, and no complete classification.
- (Th15) The derivative is the external D_z ; nothing is claimed for an intrinsic surreal derivation.
- (Th16) *[merge]* The identification of the two seed equations, Remarks 32.18, 33.4 and 33.10 are the merge's; no historical priority is claimed for them.

34 Differential freedom at a single scale: exact relation ideals

This section and the next contain the material of source 17, delivered in batch 31 with sources 15, 16, 18 and 19. The coefficient-field part explains differential transcendence without an order unit by coefficients whose values reach every Archimedean level; Theorem F does it for a single series by infinite coefficient-value rank. Source 17 supplies the opposite mechanism, *support*: the lacunary series $\mathcal{G}_A(p, z) = \sum_{n \in A} p^{\mathbf{b}_n} z^n$, with heights $\mathbf{b}_n$ whose successive ratios tend to infinity, have all Taylor coefficients in $\mathbb{Q}(p)$ and coefficient values spanning $\mathbb{Q}\mu$ when $p = t^\mu$, yet every infinite subseries is strongly entire, nonpolynomial and differentially transcendental over the full field $\mathbb{K}(z)$ whenever μ is an order unit (Theorem M). More strongly, the whole differential relation ideal of every finite family is determined: after finite polynomial corrections it is generated by the linear relations among

the membership patterns that occur infinitely often. The same finite core gives numerical surreals with independent Berarducci–Mantova jets, an exact strong-evaluation domain in $\mathbf{No}[i]$, omnific codes and floor profiles.

34.1 Source, conventions and the merge’s choices

Convention 34.1 (Setting of the single-scale part). In Sections 34–35, k is a field of characteristic zero and $n \geq 2$ throughout. In the formal sections p and z are algebraically independent indeterminates, with the commuting derivations $\vartheta_p = p \, d/dp$ and $\vartheta = z \, D_z$ on $k(p, z)$, $\vartheta_p z = 0$ and $\vartheta p = 0$; over a larger coefficient field $E(z)$ the derivation D_z kills all of E . In the Hahn sections Γ has an order unit μ , $\mathbb{K} = k((t^\Gamma))$, $p = t^\mu$, and v , $\mathcal{E}_\Gamma(k)$, $\text{Frac } \mathcal{E}_\Gamma(k)$ and strong summability are as in Conventions 14.1 and 20.1. A family of derivatives is *differentially algebraically independent* if every finite list of distinct iterated derivatives is algebraically independent over the stated base.

The source. Source 17 is *Differential Freedom at a Single Surreal Scale: Factorial-height entire functions, exact relation ideals, and omnific codes* (24 pages, A4, 23 September 2026), delivered as the archive `single_scale_surreal_research` (batch 31, manuscript 03). It consulted the repository at

bcac55ae6fd3f2b354e568ba1d2e94d496a2cc59,

the pin of source 18, at which this report was the five-source text (sources 08–12). It read this report’s directory, catalogue entry and the source 12 audit, and the READMEs of the tail-span [16] and bounded-support [38] reports. Its description of this report was accurate there. Its statement that “the general question whether every differentially algebraic strongly entire function in an order-unit Hahn field is a polynomial remains open” and its Question 12.1 are stale now: they are answered negatively by Theorem H, with least order three by Theorem I (Remark 35.20). Source 17 does not claim that question; its results concern explicit families.

Renamed symbols. The following symbols of source 17 would collide with symbols of the earlier parts. They are renamed here, in the new material only.

Source 17	Here	Reason
formal variable q ; $q = t^\eta$; order unit η	p ; $p = t^\mu$; μ	q is a dilation parameter; $p = t^\mu$ as in Section 23.1; $\eta = v(\tau)$ in Theorem 4.4.
$\delta = q \, d/dq$; $\Theta = z \, d/dz$; $D = d/dz$	ϑ_p ; ϑ ; D_z	δ is a scale; Θ_p is the partial theta series.
$S_A(q)$; $F_A(q, z)$; heights b_n , $\mathbf{b}$	$\mathcal{N}_A(p)$; $\mathcal{G}_A(p, z)$; $\mathbf{b}_n$	$\mathcal{S}$ is the coefficient-family algebra, F is (11.2), $\mathcal{F}_\alpha$ the branch family; b is a derivative order.
$K = k((t^\Gamma))$, $\mathcal{E}(K)$ (cofinally entire)	$\mathbb{K}$, $\mathcal{E}_\Gamma(k)$	The same class (Proposition 20.2).
patterns v_n , T_v , $\mathcal{P}_\infty$, E_0 ; $p(q, z)$, $s(q)$; V , W , d ; λ_j ; L	$\mathbf{v}_n$, $A_{\mathbf{v}}$, $\mathcal{P}_\infty$, A_{fin} ; $\mathbf{c}(p, z)$, $\mathbf{c}_0(p)$; $\mathbf{V}$, its column span, d_{pat} ; ℓ_j ; $\mathbf{L}$	$p = t^\mu$; V , W are indeterminates; λ is a multiplier; L an operator.
D , H , T , U , k , B_T (Lemma 3.1)	$\bar{d}$, $\bar{h}$, $\mathbf{T}$, $\mathbf{U}$, n_* , $\mathbf{b}_T$	D_z is the derivative, $H \in \Gamma$, T an indeterminate, k the coefficient field.
L , Q , J , $U_{i,h}$, $X_{i,h}$, W_Q (polarization)	$\mathfrak{K}$, $\mathfrak{P}$, $\mathcal{J}$, $\varepsilon_{i,h}$, $\mathbf{x}_{i,h}$, $\mathbf{W}_{\mathfrak{P}}$	Q_s , Q_r and W have other meanings.

Source 17	Here	Reason
$\mathcal{D}$ (domain in $\mathbf{No}[i]$)	$\mathcal{O}_{\langle 1 \rangle}$	The valuation ring of the coarsening by the convex hull of $\mathbb{Z}$, as $\mathcal{O}_{\Delta_\mu}$ in Theorem 23.5.
$\mathcal{B}_\Gamma(k), \mathcal{F}_\Gamma(k); H = \mathbb{R}((t^\mathbb{R}))$	$\mathcal{B}_\Gamma^{\text{bd}}(k), \mathcal{F}_\Gamma^{\text{bd}}(k); \mathbb{R}((t^\mathbb{R}))$	$\mathcal{B}[f]$ and $\mathcal{F}_\alpha$ have other meanings.
$\xi_A = \sum \omega^{-b_n}; \partial_{\text{BM}}$	$\mathfrak{f}_A; \partial_{\text{BM}}$	ξ_{ij} are leading coefficients; the tail-span report's ξ_A are different numbers.
$\Lambda, M = \omega^\Lambda; Z_A; I_A(a)$	$\Omega, M_\Omega = \omega^\Omega; \mathfrak{z}_A; \mathfrak{l}_A(a)$	Λ is a set of dilations, I_P the full corner polynomial.
nodes a_i , multiplicities m_i , values $c_{i,j}, M, H_\alpha, P, G_\alpha$ (Hermite)	$x_i, e_i, y_{ij}, \bar{e}, R_\alpha, R_0, \mathcal{G}_\alpha^*$	a_n are Taylor coefficients.
polynomial height $h(n)$	$\mathfrak{h}(n)$	h is a root-of-unity order.
$\mathfrak{c} = 2^{\aleph_0}$	$2^{\aleph_0}$	$\mathfrak{c}$ is a theta constant.
Theorems 4.2, 4.3, 5.1, 7.2	Theorems 34.9, 34.10, 34.13, 35.1	Theorem M collects them; Theorems A–N of this report are different theorems.

The shipped files `17-single-scale-PROOF_AUDIT.md` and `17-single-scale-SOURCES_AND_SCOPE.md` keep the source's notation.

Words with two meanings. A *rapid height sequence* is Definition 34.2; it has nothing to do with the scale μ . A *pattern* is the membership vector $\mathbf{v}_n$ of an index. *Cofinally entire* (source 17) is exactly $\mathcal{E}_\Gamma(k)$. *Differential independence* of the numerical series $\mathbf{N}_A$ refers to ϑ_p , or to ∂_{BM} after the substitution $p = \omega^{-1}$, never to D_z ; that of $\mathcal{G}_A$ refers to D_z (pure) or to ϑ, ϑ_p (mixed), over the base stated each time. The *support* mechanism of this part is not the *coefficient* mechanism of Sections 20–22 or of the tail-span report.

Results printed once. Source 17's definition of the entire class (Definition 2.2) is $\mathcal{E}_\Gamma(k)$ in the form of Proposition 20.2; its Lemma 7.1 (the entire ring) is Proposition 2.7; its Euler–Stirling conversion (2.1) is (26.2); its Hahn–Neumann support facts are Lemma 2.1; its constant-extension step (Lemma 6.1 and Proposition 6.2) is the argument of Proposition 22.7, over $\mathbb{Q}(p)$ instead of $\mathbb{K}$, applied to the family of all jets, source 17's proof via an invertible finite coefficient minor being a second route to the linear case; its normal-form embedding is (2.4); its bounded-support descent (Lemma 8.3) is the integer-lattice case of `bst:thm:descent` [38], reproved below for completeness; its binary-prefix family is the almost-disjoint device already used for $\mathcal{F}_\alpha$ (Section 22.2); and its floor profile (Theorem 10.4) is a case of `odg:thm:floor` [41], printed with its short proof.

Where each numbered result went. Definition 2.1 is Definition 34.2; Lemma 3.1, Corollary 3.2, Lemma 3.3, Example 3.4 and Lemma 3.5 are Lemma 34.3, Corollary 34.4, Lemma 34.5, Example 34.6 and Lemma 34.7; Definition 4.1, Theorems 4.2, 4.3, Corollary 4.4 and Remark 4.5 are Definition 34.8, Theorems 34.9, 34.10, Corollary 34.11 and Remark 34.12; Theorem 5.1, Corollary 5.2, Examples 5.3, 5.4 and Corollary 5.5 are Theorem 34.13, Corollary 34.14, Examples 34.15, 34.16 and Corollary 34.17; Theorem 6.3 and Warning 6.4 are Theorem 34.18 and Remark 34.19; Theorem 7.2, Corollary 7.3 and Theorem 7.4 are Theorem 35.1, Corollary 35.3 and Theorem 35.4; Lemma 8.1, Theorem 8.2, Lemma 8.3, Corollary 8.4 and Warning 8.5 are Lemma 35.5, Theorem 35.6, Lemma 35.7, Corollary 35.8 and Remark 35.9; Theorem 9.1, Propositions 9.2, 9.3 and Warning 9.4 are Theorem 35.10, Propositions 35.11, 35.13 and Remark 35.14; Lemma 10.1, Theorem 10.2, Warn-

ing 10.3 and Theorem 10.4 are Lemma 35.15, Theorem 35.16, Remark 35.17 and Theorem 35.18; Proposition 11.1 is Proposition 35.19; Warning 1.1 is merged into Section 35.6; Question 12.1 is answered (Remark 35.20) and Questions 12.2–12.11 are Questions 19.45–19.54; Appendices A and B are merged into the dependency table and suite description of Section 18, and Appendix C into Section 35.6.

The merge’s choices. The review of source 17 for this merge found its proofs correct. The merge adds three points, each marked. First, Remark 35.20 records the answer to source 17’s universal question. Second, Remark 35.2 places Theorem 35.1 beside Theorems F, G, H and L, and records that it fills the gap noted after Theorem 21.20. Third, Remark 35.12 compares the common domain of Theorem 35.10 with that of $\mathcal{T}_p$. Source 17’s theorems, proofs, examples and non-claims are otherwise printed as delivered, in the renamed notation.

34.2 Rapid heights and the finite separation mechanism

Definition 34.2 (Rapid heights; source 17). A *rapid height sequence* is a strictly increasing sequence of positive integers $(\mathbf{b}_n)_{n \geq 2}$ with $\mathbf{b}_n/\mathbf{b}_{n-1} \rightarrow +\infty$, for instance $\mathbf{b}_n = n!$. For $A \subseteq \mathbb{N}_{\geq 2}$ put

$$\mathbf{N}_A(p) = \sum_{n \in A} p^{\mathbf{b}_n} \in k[[p]], \quad \mathcal{G}_A(p, z) = \sum_{n \in A} p^{\mathbf{b}_n} z^n \in k[[p]][[z]].$$

Then $\vartheta_p^b \mathbf{N}_A = \sum_{n \in A} \mathbf{b}_n^b p^{\mathbf{b}_n}$ and $\vartheta^a \vartheta_p^b \mathcal{G}_A = \sum_{n \in A} n^a \mathbf{b}_n^b p^{\mathbf{b}_n} z^n$. By (26.2) and its analogue in p , finite-jet independence and its transcendence degree are the same for Euler and ordinary derivatives, the changes being triangular and invertible over $k(p, z)$.

Lemma 34.3 (Height separation; source 17). *Fix rapid heights and integers $\bar{d} \geq 1$, $\bar{h} \geq 0$. For all large N : if $\mathbf{T}$ is a multiset of exactly $\bar{d}$ indices, all at least N , and $\mathbf{U} \neq \mathbf{T}$ a multiset of at most $\bar{d}$ indices, then $|\sum_{n \in \mathbf{T}} \mathbf{b}_n - \sum_{n \in \mathbf{U}} \mathbf{b}_n| > \bar{h}$. For $\mathbf{b}_n = n!$ it suffices that $N \geq \max(3, 4\bar{d})$ and $N!/2 > \bar{h}$.*

Proof. If the multiplicities agree at every index at least N , then $\mathbf{U}$ already contains all $\bar{d}$ entries of $\mathbf{T}$ and has no room for more, so $\mathbf{U} = \mathbf{T}$. Hence there is a largest differing index $n_* \geq N$. There the multiplicity difference is at least one in absolute value, the lower indices contribute at most $2\bar{d}\mathbf{b}_{n_*-1}$, and higher indices cancel, so the difference is at least $\mathbf{b}_{n_*} - 2\bar{d}\mathbf{b}_{n_*-1}$. Choose N with $\mathbf{b}_n > 4\bar{d}\mathbf{b}_{n-1}$ for $n \geq N$ and $\mathbf{b}_N/2 > \bar{h}$. For $n!$ and $n \geq 4\bar{d}$ the bound is at least $n!/2$. $\square$

That $\mathbf{T}$ has *exactly* the maximal degree is essential: it stops a competitor from matching every high index and adding low ones.

Corollary 34.4 (Isolated bands; source 17). *With $\mathbf{b}_{\mathbf{T}} = \sum_{n \in \mathbf{T}} \mathbf{b}_n$, the integer interval $[\mathbf{b}_{\mathbf{T}}, \mathbf{b}_{\mathbf{T}} + \bar{h}]$ is disjoint from $[\mathbf{b}_{\mathbf{U}}, \mathbf{b}_{\mathbf{U}} + \bar{h}]$ for every different $\mathbf{U}$ of size at most $\bar{d}$.*

Lemma 34.5 (Multicolour polarization; source 17). *Let $\mathfrak{K}$ have characteristic zero, $\mathcal{J} \subseteq \mathbb{N}^s$ be finite, and $\mathfrak{P} \neq 0$ be a polynomial in variables $Y_{i,\alpha}$ ($1 \leq i \leq m$, $\alpha \in \mathcal{J}$), multihomogeneous of degree d_i in the variables of colour i . For each colour take d_i slots with scalars $\varepsilon_{i,h}$ and s -tuples $\mathbf{x}_{i,h}$, and substitute $Y_{i,\alpha} = \sum_h \varepsilon_{i,h} \mathbf{x}_{i,h}^\alpha$. The coefficient $\mathbf{W}_{\mathfrak{P}}$ of $\prod_{i,h} \varepsilon_{i,h}$ is a nonzero polynomial in the slot tuples.*

Proof. A monomial of $\mathfrak{P}$ contributes the sum over all assignments of its exponent vectors, colour by colour, to the labelled slots, with positive integer multiplicities. Different monomials differ in

the multiset of exponent vectors of some colour, so their orbits of slot monomials are disjoint, and characteristic zero keeps the multiplicities nonzero. $\square$

When the indices put in the slots are distinct, $W_{\mathfrak{P}}$ is exactly the coefficient obtained by taking each once in the series product; distinctness is enforced below, and no repeated-index factorial is divided out.

Example 34.6 (A cancellation-sensitive case; source 17). For $\mathfrak{P} = Y_0Y_2 - Y_1^2$, $W_{\mathfrak{P}}(x_1, x_2) = (x_1 - x_2)^2$: a single repeated index gives zero, two distinct indices detect it. This is the conic $\mathfrak{s}$ of Remark 30.2, whose cross coefficient $(n - m)^2$ drives the same phenomenon there.

Lemma 34.7 (Hereditary density; source 17). *For rapid heights, on every infinite set of indices a nonzero polynomial over a field of characteristic zero vanishes at only finitely many of the points $\mathbf{b}_n \in \mathbb{A}^1$, $n \in \mathbb{A}^1$, and $(n, \mathbf{b}_n) \in \mathbb{A}^2$.*

Proof. The one-dimensional cases follow from distinctness. Eventually $\mathbf{b}_n \geq 2\mathbf{b}_{n-1}$, so $\mathbf{b}_n$ dominates every power of n ; for $0 \neq P = \sum_{j \leq e} P_j(X)Y^j \in \mathbb{Q}[X, Y]$ with $P_e \neq 0$ the term $P_e(n)\mathbf{b}_n^e$ dominates the others in absolute value for large n . For general coefficients write $P = \sum_j e_j P_j$ over a $\mathbb{Q}$ -basis (e_j) of the span of its coefficients; some $P_j \neq 0$, and $P(n, \mathbf{b}_n) = 0$ forces $P_j(n, \mathbf{b}_n) = 0$. $\square$

Products of such sets are Zariski dense by induction on the number of tuples, and an infinite index set splits into any finite number of disjoint infinite sets, so distinct indices can be chosen for all slots, arbitrarily far out, with $W_{\mathfrak{P}} \neq 0$; this holds over $\mathfrak{K} = k(p)$.

34.3 Joint independence

Definition 34.8 (Finite private tails; source 17). A family $\mathcal{A}$ of infinite subsets of $\mathbb{N}_{\geq 2}$ has the *finite private-tail property* if for distinct $A_1, \dots, A_m \in \mathcal{A}$ every $A_i \setminus \bigcup_{j \neq i} A_j$ is infinite; pairwise almost-disjoint families have it.

Theorem 34.9 (Numerical all-order independence; source 17). *For rapid heights and a family $\mathcal{A}$ with finite private tails, the elements $\vartheta_p^b \mathbf{N}_A$, $A \in \mathcal{A}$, $b \geq 0$, are algebraically independent over $k(p)$.*

Proof. A relation gives, after clearing denominators, a nonzero $P \in k[p][Y_{i,j}]$ of top total degree $\bar{d}$ and p -degree $\bar{h}$; $\bar{d} = 0$ is impossible since p is transcendental over k . Take a nonzero multihomogeneous part $\mathfrak{P}$ of degree $\bar{d}$ with colour degrees d_i , take N from Lemma 34.3, split each colour's private indices beyond N into d_i infinite sets, and by Lemmas 34.5 and 34.7 over $k(p)$, with slot weights $\mathbf{b}_n$, choose $\bar{d}$ distinct private indices beyond N with polarized coefficient $C(p) \neq 0$ in $k[p]$, of degree at most $\bar{h}$. Let $\mathbf{T}$ be this multiset. Every contribution to P evaluated comes from an index multiset of size at most $\bar{d}$ followed by a shift in $[0, \bar{h}]$; by Corollary 34.4 only $\mathbf{T}$ reaches the band $[\mathbf{b}_{\mathbf{T}}, \mathbf{b}_{\mathbf{T}} + \bar{h}]$, the private indices force the colour degrees d_i , and the contribution there is $p^{\mathbf{b}_{\mathbf{T}}} C(p) \neq 0$. $\square$

Theorem 34.10 (Mixed differential independence; source 17). *For the same heights and family, all mixed jets $\vartheta^a \vartheta_p^b \mathcal{G}_A$, $A \in \mathcal{A}$, $a, b \geq 0$, are algebraically independent over $k(p, z)$, and so are all ordinary mixed partial derivatives.*

Proof. From a nonzero $P \in k[p, z][Y_{i,a,b}]$ of top degree $\bar{d}$ and p -degree $\bar{h}$, choose z^{j_0} and a colour multidegree of total degree $\bar{d}$ with nonzero part $\mathfrak{P} \in k[p][Y_{i,a,b}]$, polarize with the two-coordinate weights $(n, \mathbf{b}_n)$, and choose $\bar{d}$ distinct private target indices beyond the separation threshold with

polarized weight $C(p) \neq 0$. With $\mathbf{b}_T = \sum_T \mathbf{b}_n$ and $M = j_0 + \sum_T n$, the band $[\mathbf{b}_T, \mathbf{b}_T + \bar{h}]$ of the coefficient of z^M receives only the target multiset, which forces the colours and the power z^{j_0} , and equals $p^{\mathbf{b}_T} C(p)$. Each z -coefficient is a polynomial in p , so the argument takes place in an ordinary polynomial ring; the triangular changes give the ordinary derivatives. $\square$

Corollary 34.11 (Every infinite subseries; source 17). *For infinite A , $\mathbf{N}_A$ is differentially transcendental over $k(p)$, and $\mathcal{G}_A$ satisfies no nonzero algebraic partial differential equation in ϑ, ϑ_p over $k(p, z)$, in particular no algebraic ordinary differential equation in D_z over $k(p, z)$.*

Remark 34.12 (Why nonlinear equations are handled; source 17). The selected contribution uses the top *total* degree of the whole relation, not its highest derivative; polarization handles cancellation among nonlinear monomials, and the band lemma excludes every lower-degree competitor at once. No first-order, separant, linearization or residue hypothesis is imposed. For a single ordinary series the conclusion is classical (Section 35.6).

34.4 The exact relation ideal

Fix arbitrary $A_1, \dots, A_m \subseteq \mathbb{N}_{\geq 2}$. For a nonzero pattern $\mathbf{v} \in \{0, 1\}^m$ let $A_{\mathbf{v}} = \{n : \mathbf{v}_n = \mathbf{v}\}$, where $\mathbf{v}_n = (\mathbf{1}_{A_1}(n), \dots, \mathbf{1}_{A_m}(n))$, and $\mathcal{P}_{\infty} = \{\mathbf{v} \neq 0 : A_{\mathbf{v}} \text{ infinite}\}$. The union A_{fin} of the finite nonzero pattern classes is finite; put $\mathbf{c}(p, z) = \sum_{n \in A_{\text{fin}}} \mathbf{v}_n p^{b_n} z^n$ and $\mathbf{c}_0(p) = \sum_{n \in A_{\text{fin}}} \mathbf{v}_n p^{b_n}$, let $\mathbf{V}$ be the matrix with the columns $\mathbf{v} \in \mathcal{P}_{\infty}$, and $d_{\text{pat}} = \text{rank } \mathbf{V}$, the same over every field of characteristic zero. With $\mathcal{G} = (\mathcal{G}_{A_i})_i$ and $\mathbf{N} = (\mathbf{N}_{A_i})_i$,

$$\mathcal{G} = \mathbf{c} + \mathbf{V}(\mathcal{G}_{A_{\mathbf{v}}})_{\mathbf{v} \in \mathcal{P}_{\infty}}, \quad \mathbf{N} = \mathbf{c}_0 + \mathbf{V}(\mathbf{N}_{A_{\mathbf{v}}})_{\mathbf{v} \in \mathcal{P}_{\infty}}, \quad (34.1)$$

where the $A_{\mathbf{v}}$ are pairwise disjoint and infinite, so Theorems 34.9 and 34.10 apply to them.

Theorem 34.13 (Complete relation classification; source 17). *Let $\ell_1, \dots, \ell_{m-d_{\text{pat}}} \in \mathbb{Q}^m$ be a basis of the left kernel of $\mathbf{V}$. The kernel of evaluation of the differential polynomial ring over $k(p)$ at $\mathbf{N}$ is the differential ideal generated by $\ell_j \cdot (Y - \mathbf{c}_0(p))$; the kernel over $k(p, z)$ at $\mathcal{G}$, with the derivations ϑ, ϑ_p , is the differential ideal generated by $\ell_j \cdot (Y - \mathbf{c}(p, z))$. There are no further nonlinear relations; for $d_{\text{pat}} = 0$ all functions are the corrections, and for $d_{\text{pat}} = m$ the family is differentially algebraically independent.*

Proof. Choose $\mathbf{L} \in \text{GL}_m(\mathbb{Q})$ whose last $m - d_{\text{pat}}$ rows span the left kernel. After the affine change $X = \mathbf{L}(Y - \mathbf{c})$ the last coordinates vanish and the first d_{pat} are linear forms of full row rank in the pattern functions, acting separately on each block of their independent jets; a full-row-rank substitution is injective on polynomial rings (a right inverse of the matrix gives a left inverse). So the kernel is generated by the jets of the last coordinates. The numerical case is the same with $\mathbf{c}_0$ and ϑ_p . $\square$

Corollary 34.14 (Exact finite-jet degrees; source 17). *For every finite family,*

$$\begin{aligned} \text{trdeg}_{k(p)} k(p)(\vartheta_p^b \mathbf{N}_{A_i} : i \leq m, b \leq s) &= (s+1)d_{\text{pat}}, \\ \text{trdeg}_{k(p,z)} k(p,z)(\vartheta^a \vartheta_p^b \mathcal{G}_{A_i} : i \leq m, a \leq r, b \leq s) &= (r+1)(s+1)d_{\text{pat}}, \end{aligned}$$

and the same with ordinary derivatives in the corresponding rectangles.

Equivalently, d_{pat} is the rank of the indicator classes of the A_i in $\mathbb{Q}^{\mathbb{N}_{\geq 2}}/\mathbb{Q}^{(\mathbb{N}_{\geq 2})}$: a rational combination of indicators is eventually zero exactly when it annihilates every recurring pattern. Up to finite corrections, the differential dependence geometry of $A \mapsto \mathcal{G}_A$ is ordinary linear dependence modulo finite sets.

Example 34.15 (Union and intersection; source 17). $\mathcal{G}_{A \cup B} + \mathcal{G}_{A \cap B} = \mathcal{G}_A + \mathcal{G}_B$, and likewise for $\mathbf{N}$. If $A \setminus B$, $B \setminus A$ and $A \cap B$ are infinite, the four series have pattern rank three, and this identity with its derivatives generates their whole relation ideal; the family of all infinite subsets is not independent.

Example 34.16 (Finite changes; source 17). If $B = A \cup \{n_0\}$ with $n_0 \notin A$, then $\mathcal{G}_B - \mathcal{G}_A = p^{b_{n_0}} z^{n_0}$: the ideal keeps this affine correction, and after it there is exactly one differential generator when A is infinite.

Corollary 34.17 (Finite alphabets; source 17). *The classification holds when the patterns $\mathbf{v}_n$ range over a fixed finite subset of k^m , with the recurring vectors, the corrections and k -linear rank.*

Finiteness of the alphabet matters: for $\mathbf{v}_n = (1, n)$ the second series is ϑ of the first, and no finite-pattern description applies.

34.5 The full scalar field

Theorem 34.18 (All scalars may enter the equation; source 17). *Let E have characteristic zero and contain an element p transcendental over its prime field, and regard $\mathcal{G}_A$ in $E[[z]]$. For rapid heights, the pure D_z relation ideal over $E(z)$ of every finite family is exactly the ideal generated by the $\ell_j \cdot (Y - \mathbf{c}(p, z))$ and their D_z -derivatives. In particular every infinite subseries is differentially transcendental over $E(z)$, and a finite-private-tail family is jointly so.*

Proof. The pattern functions have coefficients in $\mathbb{Q}(p) \subseteq E$ and all their D_z -jets are algebraically independent over $\mathbb{Q}(p)(z)$ (Theorem 34.10 with $k = \mathbb{Q}$ and the triangular change); the proof of Proposition 22.7, with $\mathbb{Q}(p)$ in place of $\mathbb{K}$, carries this to $E(z)$. Source 17 proves this step by writing a relation as an E -linear relation among $\mathbb{Q}(p)$ -linearly independent series $z^j \prod (D_z^a f_i)^{e_{i,a}}$ and choosing an invertible finite minor of their coefficient matrix over $\mathbb{Q}(p)$, whose determinant stays nonzero in E . The full-row-rank argument of Theorem 34.13 then gives the ideal. $\square$

Remark 34.19 (What constant extension does not say; source 17). The theorem freezes all of E when differentiating in the external variable z . It does not make the numbers $\mathbf{N}_A(p)$ independent over $E = k((p))$, which contains them, and it equips E with no derivation extending ϑ_p : the mixed theorem has base $k(p, z)$ only. A finite relation with coefficients in a proper class, such as $\mathbf{No}[i]$, involves finitely many coefficients, which with p generate a set-sized field to which the theorem applies.

35 Entire, surreal and omnific realizations of single-scale freedom

35.1 Strongly entire series at one cofinal scale

Theorem 35.1 (Entire differential freedom; source 17). *Let Γ have an order unit μ , k have characteristic zero, $\mathbb{K} = k((t^\Gamma))$ and $p = t^\mu$. Every $\mathcal{G}_A$ with rapid heights lies in $\mathcal{E}_\Gamma(k)$; for each finite family its pure differential relations over $\mathbb{K}(z)$ are those of Theorem 34.18; in particular every infinite subseries is a nonpolynomial strongly entire series satisfying no algebraic ordinary differential equation over $\mathbb{K}(z)$ of any order.*

Proof. For $n \in A$ the coefficient value is $\mathbf{b}_n \mu$. For $\gamma \in \Gamma$ choose m with $\gamma \geq -m\mu$; then $\mathbf{b}_n \mu + n\gamma \geq (\mathbf{b}_n - mn)\mu$, and $\mathbf{b}_n/n \rightarrow \infty$ with μ an order unit makes this cofinal, so $\mathcal{G}_A \in \mathcal{E}_\Gamma(k)$ by Proposition 20.2. The monomial p is transcendental over the prime field; apply Theorem 34.18 with $E = \mathbb{K}$. $\square$

Remark 35.2 (Where the theorem sits; merge). The coefficient values $\mathbf{b}_n \mu$ of $\mathcal{G}_A$ span $\mathbb{Q}\mu$, so its coefficient-value rank is one and Theorem F does not apply; the remark after Theorem 21.20 noted that that theorem “says nothing about a series whose coefficient values span only $\mathbb{Q}\mu$ ” and “does not settle the purely differential problem over $\mathbb{C}((t^\mathbb{Q}))$ ”. Theorem 35.1 supplies, over $\mathbb{C}((t^\mathbb{Q}))$ and every order-unit field, explicit nonpolynomial strongly entire series of coefficient-value rank one that are differentially transcendental over $\mathbb{K}(z)$. Their coefficients lie in $\mathbb{Q}(p)$, a field of transcendence degree one over $\mathbb{Q}$, so they are further witnesses for condition (ii) of Theorem G: a finitely generated coefficient field is necessary for differential algebraicity (Theorem 21.2 (a)) but, with an order unit, not sufficient. Beside Theorem H and Theorem L, the same fields therefore carry nonpolynomial strongly entire series of every minimal order in a set that contains 3 and is unbounded (for $\Gamma = \mathbb{R}$), and series of no finite order. They do not decide Θ_p or G , whose heights n^2 are not rapid (Proposition 35.19, Question 19.45).

For a binary sequence $\alpha = (\alpha_1, \alpha_2, \dots)$ put $n_\ell(\alpha) = 2^\ell + \sum_{j \leq \ell} \alpha_j 2^{\ell-j}$ and $A_\alpha = \{n_\ell(\alpha) : \ell \geq 1\}$. Each $n_\ell(\alpha)$ is 1 followed by the first ℓ digits of α in binary, different lengths lie in disjoint dyadic intervals, and $A_\alpha \cap A_\beta$ is finite for $\alpha \neq \beta$; so the family has finite private tails. Each A_α is computable relative to α ; uniform computability of all members is not claimed.

Corollary 35.3 (Maximal differential transcendence degree; source 17). *For $\mathbb{K} = \mathbb{R}((t^\mathbb{Q}))$ or $\mathbb{C}((t^\mathbb{Q}))$, with $p = t = \omega^{-1}$ in the surreal realization, $\text{dtrdeg}_{\mathbb{K}(z)} \text{Frac } \mathcal{E}_\Gamma(k) = 2^{\aleph_0}$, with the lower bound supplied by $(\mathcal{G}_{A_\alpha})_{\alpha \in 2^\mathbb{N}}$, all of whose coefficients lie in $\mathbb{Q}(t)$.*

Proof. Joint differential independence gives the lower bound; $|\mathbb{K}[[z]]| = (2^{\aleph_0})^{\aleph_0}$ gives the upper bound. $\square$

This is the order-unit counterpart of Theorem 22.6; for $\mathbb{K} = k((t^\mathbb{R}))$ see Remark 33.4. A family of maximal cardinality is not claimed to be a differential transcendence basis.

Theorem 35.4 (Freedom inside every finite-jet fibre; source 17). *In either field of Corollary 35.3, fix distinct nodes $x_1, \dots, x_s \in \mathbb{K}$, multiplicities $e_i \geq 1$ and values $y_{ij} \in \mathbb{K}$, $0 \leq j < e_i$. There are $2^{\aleph_0}$ jointly differentially algebraically independent $\mathcal{G}_\alpha^* \in \mathcal{E}_\Gamma(k)$ with $D_z^j \mathcal{G}_\alpha^*(x_i) = y_{ij}$ for all i, j, α .*

Proof. With $\bar{e} = \sum e_i$, the map $\mathbb{K}[z]_{<\bar{e}} \rightarrow \prod_i \mathbb{K}^{e_i}$, $R \mapsto (D_z^j R(x_i))$, is injective, a polynomial in its kernel being divisible by $\prod (z - x_i)^{e_i}$, hence bijective. Let R_α interpolate the jets of $\mathcal{G}_{A_\alpha}$ and R_0 the values y_{ij} , and put $\mathcal{G}_\alpha^* = \mathcal{G}_{A_\alpha} - R_\alpha + R_0$. For finitely many α the translation is an invertible change of variables over $\mathbb{K}(z)$. $\square$

35.2 Numerical surreals and the intrinsic derivation

The normalized Berarducci–Mantova derivation ∂_{BM} has real constants, is strongly additive and satisfies $\partial_{\text{BM}} \omega = 1$ [8]. On the integer-exponent Laurent field with $p = \omega^{-1}$ it therefore acts as

$$\partial_{\text{BM}} \left(\sum_{n \geq n_0} c_n p^n \right) = - \sum_{n \geq n_0} n c_n p^{n+1} = -p \vartheta_p \left(\sum_{n \geq n_0} c_n p^n \right), \quad (35.1)$$

first on monomials and then by strong additivity. It is not D_z : it moves p .

Lemma 35.5 (Triangular intrinsic jets; source 17). *On that field $\partial_{\text{BM}}^j = (-1)^j p^j \vartheta_p(\vartheta_p + 1) \cdots (\vartheta_p + j - 1)$, so intrinsic and Euler jets of order at most r are related by an invertible triangular matrix over $\mathbb{R}(p)$.*

Proof. $\vartheta_p(p^j g) = p^j(\vartheta_p + j)g$; induct from $\partial_{\text{BM}} = -p\vartheta_p$. $\square$

Theorem 35.6 (Rational-coefficient surreal differential freedom; source 17). *Let $\mathfrak{x}_A = \sum_{n \in A} \omega^{-b_n} \in \mathbb{R}((\omega^{-1})) \subseteq \mathbf{No}$. For a finite-private-tail family of infinite sets, all $\partial_{\text{BM}}^j \mathfrak{x}_A$ are jointly algebraically independent over $\mathbb{R}(\omega)$; for arbitrary finite families the relation ideal is that of Theorem 34.13 with the correction $\mathbf{c}_0(\omega^{-1})$ and ∂_{BM} in place of ϑ_p . In particular $\text{dtrdeg}_{\mathbb{R}(\omega)} \mathbb{R}((\omega^{-1})) = 2^{\aleph_0}$ for ∂_{BM} , and the analogous statement holds over $\mathbb{C}(\omega)$.*

Proof. Theorem 34.9 and the classification with $k = \mathbb{R}$, then Lemma 35.5; each nonzero $\mathfrak{x}_A$ is a positive infinitesimal; the binary-prefix family and the cardinality of the Laurent field give the transcendence degree. $\square$

Let $\mathcal{B}_{\Gamma}^{\text{bd}}(k)$ be the ring of $f \in k((t^{\Gamma}))$ with support bounded above, and $\mathcal{F}_{\Gamma}^{\text{bd}}(k)$ its fraction field; it contains $k[p, p^{-1}]$ for $p = t^{\mu}$.

Lemma 35.7 (Bounded-support descent; source 17, after `bst:thm:descent`). *Let μ be an order unit. If $g_1, \dots, g_N \in k((p))$ are linearly independent over $k(p)$, they are linearly independent over $\mathcal{F}_{\Gamma}^{\text{bd}}(k)$ in $k((t^{\Gamma}))$.*

Proof. From $\sum_j b_j g_j = 0$ with $b_j \in \mathcal{B}_{\Gamma}^{\text{bd}}(k)$, project to a coset $\gamma + \mathbb{Z}\mu$: the support of b_j there is bounded below and above, hence finite because μ is an order unit, so the projection is $t^{\gamma} P_{j,\gamma}(p)$ with a Laurent polynomial $P_{j,\gamma}$. Multiplication by $g_j \in k((p))$ preserves cosets, so $\sum_j P_{j,\gamma}(p) g_j = 0$ and every $P_{j,\gamma} = 0$. $\square$

Corollary 35.8 (All intrinsic jets over the bounded-support field; source 17). *For $p = \omega^{-1}$ and $\mathcal{F}^{\text{bd}} = \mathcal{F}_{\mathbb{R}}^{\text{bd}}(\mathbb{R}) \subseteq \mathbb{R}((t^{\mathbb{R}})) \subseteq \mathbf{No}$, the jets $\partial_{\text{BM}}^j \mathfrak{x}_A$ of a finite-private-tail family are algebraically independent over $\mathcal{F}^{\text{bd}}$, also with complex coefficients; for an arbitrary finite family the ideal of each finite jet block is the extension of its ideal over $\mathbb{R}(p)$.*

Proof. Jet monomials lie in $\mathbb{R}((p))$ and are linearly independent over $\mathbb{R}(p)$ by Theorem 35.6; apply Lemma 35.7, after the affine change of Theorem 34.13 in the dependent case. $\square$

Remark 35.9 (An absolute base-field boundary; source 17). The witnesses $\mathfrak{x}_A$ and their jets lie in $\mathbb{R}((p))$, so they cannot be independent over a base containing them, for instance $\mathbb{R}((p))$ or $\mathbf{No}$; the external-variable scalar theorem does not contradict this. After a noncofinal enlargement of the value group all integer supports can become bounded, and the bounded-support base then absorbs every witness.

35.3 An exact domain in the full class

With $t = \omega^{-1}$ and valuations of surcomplex normal forms in that convention, put $\mathcal{O}_{\langle 1 \rangle} = \{0\} \cup \{x \in \mathbf{No}[i]^{\times} : v(x) \geq -m \text{ for some } m \in \mathbb{N}\}$, equivalently the x with $|x| < \omega^m$ for some m .

Theorem 35.10 (Exact common domain; source 17). *For rapid heights and infinite A , the family $(\omega^{-b_n} x^n)_{n \in A}$ is strongly summable exactly when $x \in \mathcal{O}_{\langle 1 \rangle}$, and every termwise D_z -derivative has the same domain; the domain depends neither on A nor on the heights.*

Proof. Write $x = ct^\gamma(1+w)$ with $v(w) > 0$. The n th term has leading value $\beta_n = \mathbf{b}_n + n\gamma$. If $\gamma \geq -m$, the increments of $\mathbf{b}_n$ tend to infinity, so β_n increases eventually, and the supports lie in $\beta_n + M(\text{supp } w)$, so Lemma 2.3 (b) gives strong summability, with $(1+w)^n$ a finite expansion. If $\gamma < -m$ for every m , then for $n' > n$, $\beta_{n'} - \beta_n = (\mathbf{b}_{n'} - \mathbf{b}_n) + (n' - n)\gamma < 0$, an ordinary integer plus a negative infinite element, so the leading exponents decrease strictly and Lemma 2.2 excludes summability. A j th derivative shifts β_n by $-j\gamma$ and multiplies by nonzero integers. $\square$

Proposition 35.11 (The domain is a valuation ring; source 17). *$\mathcal{O}_{\langle 1 \rangle}$ is a valuation subring of $\mathbf{No}[i]$, with units the elements whose valuations are bounded in absolute value by an integer and maximal ideal $\{0\}$ together with the elements of positive infinite valuation.*

Proof. Integer lower bounds are preserved by sums and add under products; for $x \neq 0$ one of x, x^{-1} lies in the ring, both exactly when $v(x)$ is bounded on both sides. $\square$

Remark 35.12 (The same shape as the domain of $\mathcal{T}_p$; merge). For $p = \omega^{-1}$ the strong domain of $\mathcal{T}_p$ in its field is $\mathcal{O}_{\Delta_\mu}$ (Theorem 23.5), the valuation ring of the coarsening by the convex subgroup generated by the scale; $\mathcal{O}_{\langle 1 \rangle}$ is the same construction in the whole class $\mathbf{No}[i]$. Both are proper, as they must be (Proposition 27.6): ω^ω is outside both.

Proposition 35.13 (Faithfulness on the domain; source 17). *Every nonzero finite polynomial expression in z and the D_z -jets of a finite-private-tail family $(\mathcal{G}_A)$ with $p = \omega^{-1}$, with coefficients in $\mathbf{No}[i]$, is nonzero at some $x \in \mathcal{O}_{\langle 1 \rangle}$, which can be taken of arbitrarily large positive valuation; so the formal independence over $\mathbf{No}[i](z)$ excludes identities on the common domain.*

Proof. The expression $g = \sum g_n z^n$ is nonzero by Theorem 34.18, in the set-sized field of its coefficients and p . The jets of $\mathcal{G}_A$ have coefficients of nonnegative value, so $v(g_n) \geq L$ for one L . With m least such that $g_m \neq 0$ and a positive $\rho > v(g_m) - L$, the term of degree m is the unique least one at $x = t^\rho$, since $v(g_n) + n\rho \geq L + n\rho > v(g_m) + m\rho$ for $n > m$; evaluation and regrouping are legitimate by Theorem 35.10. $\square$

Remark 35.14 (Three noninterchangeable operations; source 17). Strong summation on $\mathcal{O}_{\langle 1 \rangle}$, cofinal convergence in a fixed order-unit workspace and convergence in the fine topology of the full class are different notions, and the third is not inferred from the first two. The derivative in Proposition 35.13 is the termwise D_z , not ∂_{BM} after a substitution $z = z(\omega)$. For instance ω^a with real a lies in $\mathcal{O}_{\langle 1 \rangle}$, and ω^ω does not: the series are entire on $\mathbb{C}((t^\mathbb{Q}))$ and not on $\mathbf{No}[i]$.

35.4 Omnific codes and floor profiles

Let Ω be a positive infinite surreal, $M_\Omega = \omega^\Omega$, and

$$\mathfrak{z}_A = M_\Omega \mathfrak{x}_A = \sum_{n \in A} \omega^{\Omega - \mathbf{b}_n}.$$

All exponents are positive and the constant term is zero, so $\mathfrak{z}_A \in \mathbf{Oz}$, a positive purely infinite omnific integer for infinite A [41].

Lemma 35.15 (The coding scale is transcendental; source 17). *M_Ω is transcendental over $\mathbb{R}((t^\mathbb{R}))$, and the support cosets of $M_\Omega^j h$ and $M_\Omega^{j'} h'$, $h, h' \in \mathbb{R}((t^\mathbb{R}))$, are disjoint for $j \neq j'$.*

Proof. $j\Omega - r = j'\Omega - r'$ with real r, r' would put $(j - j')\Omega$ in $\mathbb{R}$. $\square$

Theorem 35.16 (Independent omnific codes; source 17). *For a finite-private-tail family of infinite sets, the $\mathfrak{z}_A$ are algebraically independent over $\mathcal{F}^{\text{bd}}(M_\Omega)$, hence over $\mathbb{R}(\omega, M_\Omega)$, and the binary-prefix family gives $2^{\aleph_0}$ of them. For an arbitrary finite family the algebraic relation ideal over $\mathcal{F}^{\text{bd}}(M_\Omega)$ is generated by $\ell_j \cdot (Y - M_\Omega c_0(p))$, and its transcendence degree is d_{pat} .*

Proof. Corollary 35.8 at order zero gives independence of the pattern numbers over $\mathcal{F}^{\text{bd}}$ inside $\mathbb{R}((t^{\mathbb{R}}))$; adjoining M_Ω preserves it by Lemma 35.15, comparing powers of M_Ω ; multiplication by M_Ω is an invertible substitution. Multiply (34.1) by M_Ω for the general case. $\square$

Remark 35.17 (What the codes do not give; source 17). This is an *algebraic* theorem: intrinsic differential independence of the $\mathfrak{z}_A$ would require tracking the derivatives of M_Ω . No irreducibility, primality or divisibility statement is made, and multiplying by M_Ω is not the forbidden evaluation $\mathcal{G}_A(M_\Omega)$ outside $\mathcal{O}_{(1)}$.

For factorial heights and an ordinary real $a \geq 0$, Theorem 35.10 gives $\mathcal{G}_A(\omega^{-1}, \omega^a) = \sum_{n \in A} \omega^{an - n!}$; put $\mathfrak{l}_A(a) = \sum_{n \in A, (n-1)! \leq a} \omega^{an - n!}$.

Theorem 35.18 (Factorial floor profile; source 17). *For infinite $A \subseteq \mathbb{N}_{\geq 2}$ and real $a \geq 0$, $[\mathcal{G}_A(\omega^{-1}, \omega^a)]_{\mathbf{Oz}} = \mathfrak{l}_A(a)$, and $0 < \mathcal{G}_A(\omega^{-1}, \omega^a) - \mathfrak{l}_A(a) < 1$; the cutoffs are the $(n-1)!$ with $n \in A$.*

Proof. $an - n! \geq 0$ exactly when $(n-1)! \leq a$, for finitely many n ; at most one exponent is zero, so $\mathfrak{l}_A(a)$ has an integer constant term 0 or 1 and lies in $\mathbf{Oz}$. The rest has negative real exponents and coefficients one, a positive infinitesimal. A positive nonzero omnific integer is at least 1. This is the case of `odg:thm:floor` [41] in which the real part is 0 or 1 and the infinitesimal part is positive. $\square$

For $A = \mathbb{N}_{\geq 2}$ the floors at $a = 0, 1, 2, 3, 6$ are 0, 1, $\omega^2 + 1$, $\omega^4 + \omega^3$ and $\omega^{12} + \omega^{10} + 1$.

35.5 The role of the growth hypothesis

Rapid ratios isolate a maximal-degree target against every bounded-degree competitor, and make the points $(n, \mathbf{b}_n)$ eventually avoid every algebraic curve; $\mathbf{b}_n = n!$, $(n!)^2$ and 2^{n^2} qualify.

Proposition 35.19 (A polynomial-height obstruction; source 17). *If $\mathbf{b}_n = \mathbf{h}(n)$ for an integer-valued polynomial $\mathbf{h}$ on the chosen indices, then $\vartheta_p \mathcal{G}_A = \mathbf{h}(\vartheta) \mathcal{G}_A$ for every A ; for $\mathbf{b}_n = n^2$, $\vartheta_p \mathcal{G}_A = \vartheta^2 \mathcal{G}_A$.*

Proof. Both sides multiply the coefficient of $p^{\mathbf{b}_n} z^n$ by $\mathbf{h}(n)$. $\square$

This concerns *mixed* derivations; it gives no pure D_z equation over a Hahn coefficient field. With $\mathbf{b}_n = n^2$ and $p = t^\mu$, $\mathcal{G}_{\mathbb{N}_{\geq 2}}$ is, up to its first two terms, the series G_μ of Section 29.4, which is strongly entire at an order unit and whose differential algebraicity is not decided (Question 19.45).

For a proposed relation of top degree $\bar{d}$ and p -degree $\bar{h}$ the factorial threshold $N \geq \max(3, 4\bar{d})$, $N!/2 > \bar{h}$ is explicit, and polarization reduces nonvanishing to a polynomial test at far-out points. This is a finite certificate given effective access to enough private indices, not a decision procedure: whether a computable set is infinite, or a pattern recurs, may be undecidable. Intrinsic differentiation of $\mathcal{G}_A(\omega^{-1}, z(\omega))$ mixes coefficient and chain-rule terms, so formal D_z independence with frozen scalars cannot be substituted into it; no single surreal number is transcendental over all of $\mathbf{No}$. The proofs use finite polynomials, countable sequences and $(2^{\aleph_0})^{\aleph_0} = 2^{\aleph_0}$; no large-cardinal hypothesis, CH or GCH is needed.

Remark 35.20 (Source 17’s universal question; merge). Source 17’s Question 12.1 asks whether, in a characteristic-zero order-unit Hahn field, every strongly entire solution of an algebraic ordinary differential equation over $\mathbb{K}(z)$ must be a polynomial. At its pin that was Question 19.1 as then posed; it is answered negatively by $\mathcal{T}_p$ (Theorem H), with least order three (Theorem I), and it is not printed as an open question. What remains is the classification of Question 19.1 and its companions.

35.6 Predecessors, repository comparison and priority

Literature. Differential transcendence of a single ordinary power series with unbounded ratios of successive nonzero exponents is classical: Ostrowski’s 1920 paper, Section 5 just before Theorem 11, attributes the criterion to Grönwall [36]. Source 17 read Ostrowski in the English translation of Hansen, Stone and Wolfson; the Grönwall original was not inspected. So differential transcendence of an individual factorial-gap series over $\mathbb{C}(p)$ is not new; the unified proof is given because it also controls joint mixed jets and the whole relation ideal. Source 17 imports the normal-form identification and the Berarducci–Mantova derivation [8, 39] and the normal-form definition of **Oz** [37]; it consulted the Wikipedia article on surreal numbers for orientation only, and located general lacunary-series literature not used as a premise. No exhaustive MathSciNet, zbMATH, thesis, historical-language or citation-network search was made. For this merge the bibliographic data of the Ostrowski translation and of the L’Innocente–Mantova preprint were checked; their content was not.

Repository comparison at the pin. Source 17 read, at bcac55a, the root README, recursive tree and catalogue, this report’s directory, catalogue entry and source 12’s audit, and the READMEs of the tail-span and bounded-support reports; README claims were recorded as claims, not verified proofs, and this was not a line-by-line audit of the 57 catalogued reports, their histories or the Lean declarations. The tail-span report already has an exact cofinite-span relation theory and numerical ∂_{BM} jets for different witnesses (its `tail:thm:surreal`, `tail:eq:BMtriangular` and `tail:rem:indexfamily` [16]); source 17’s mechanism is support, not coefficients, and its numerical base differs. The bounded-support report already has factorial-support algebraic independence, the witnesses $\eta_A = \sum_n c_A(n)\omega^{-n!}$, algebraically independent over the bounded-support field, and the descent theorem, and states that it proves no differential independence [38]; source 17 continues it differentially. These overlaps are recorded, not hidden.

Priority. Not claimed new: single-series gap differential transcendence; normal forms, the Berarducci–Mantova derivation and **Oz**; lacunarity, polarization, row rank, almost-disjoint families and the cardinality of a continuum family; cofinite-span invariants as an idea; factorial-support algebraic independence and bounded-support descent; existence of differentially transcendental surreal numbers in general. The proposed contributions are the complete ordinary and mixed relation ideals for finite-alphabet rapid-height families with exact jet degrees, pure D_z independence over the full scalar Hahn field for strongly entire series with coefficients in $\mathbb{Q}(p)$, arbitrary finite Hermite data in a continuum family, ∂_{BM} -jet independence over the bounded-support field, the exact common domain in **No**[i], and the omnific codes and floors; priority is provisional, and a correct theorem can be a rediscovery.

35.7 What the single-scale part does not claim

Every limitation stated by source 17, in its manuscript, README, audit files or verification records, is kept.

- (F1) Differential transcendence of one factorial-gap series is classical (Grönwall, via Ostrowski 1920) and not new; the original was not inspected.
- (F2) Normal forms, the Berarducci–Mantova derivation, **Oz**, lacunarity, polarization, almost-disjoint families, cofinite spans and bounded-support descent are not claimed new; nor is the existence of differentially transcendental surreal numbers.
- (F3) The universal order-unit problem is not solved by source 17 (it is answered in this report by source 13; Remark 35.20).
- (F4) The codes carry no primality, irreducibility or divisibility statement, and their independence is algebraic, not intrinsic differential.
- (F5) The mixed theorem has base $k(p, z)$ with its two named derivations only.
- (F6) The pure z theorem freezes E ; the witnesses are not transcendental over $k((p))$, $\mathbb{R}((t^{\mathbb{R}}))$ or **No**.
- (F7) Only finite alphabets are classified; $\mathbf{v}_n = (1, n)$ shows the obstruction.
- (F8) The continuum family is not a differential transcendence basis and is not uniformly computable.
- (F9) The Hermite data are finite.
- (F10) The domain theorem uses strong summation, not the fine topology; ∂_{BM} is not D_z after a substitution.
- (F11) Bounded-support descent needs the integer scale to be cofinal.
- (F12) Rapid ratios are sufficient, not necessary; polynomial heights fail the mixed theorem.
- (F13) The finite witness is not a decision procedure.
- (F14) No large cardinals, CH or GCH are used.
- (F15) No independent referee report and no Lean formalization; the 8,608 checks are regression checks, and their number is not a count of theorems.
- (F16) Byte-for-byte reproducibility of the PDF is not claimed.
- (F17) The repository comparison covered three reports' documentation, not every report or Lean declaration; the literature search was targeted.
- (F18) The floor theorem is proved only for ordinary real $a \geq 0$.
- (F19) *[merge]* Remarks 35.2, 35.12 and 35.20 and the comparison with Proposition 22.7 are the merge's; no historical priority is claimed for them.

36 Polynomial-composition rigidity: profiles and the weighted theorem

This section and the next contain the material of source 15, delivered in batch 31 with sources 16–19 and merged last. The dilation theory of Sections 7–10 and the mixed criterion of Theorem J concern arguments λz of degree one, and there an order unit decides everything. Source 15 replaces the dominant argument by a polynomial $P(z)$ of degree $d \geq 2$. Then no hypothesis on Γ is needed at all: a product of $\bar{s}$ nonzero linear differential transforms of $f(P(z))$ cannot be balanced by any polynomial in differential transforms of $f(P_\nu(z))$ with $\deg P_\nu < d$ whose monomials have *composition weight* below $\bar{s}d$, unless f is a polynomial (Theorem N). The proof keeps the Gauss data of Section 14.3, reads how a polynomial argument rescales them, and compares scales through an integer convexity inequality. An explicit degree bound retains all exact indicial resonances; the linear case with ordinary integer data has a complete omnific solution module; and every affine hypersurface over $\mathbb{Q}$ is the entire-solution locus of one such equation, which makes omnific solvability undecidable inside this rigid class.

36.1 Source, conventions and the merge’s choices

Convention 36.1 (Setting of the polynomial-composition part). In Sections 36–37, k is an arbitrary field of characteristic zero with the trivial valuation, Γ is as in Convention 1.1, $\mathbb{K} = k((t^\Gamma))$, and v , lc , res , strong summability, $\mathcal{E}_\Gamma(k)$, D_z and the Gauss data $\gamma_\delta(f)$, $\text{Act}_\delta(f)$, $\phi_\delta(f)$ of a nonzero $f \in \mathcal{E}_\Gamma(k)$ are as in Conventions 14.1 and 26.1, (14.1) and (26.1); we put $\gamma_\delta(0) = +\infty$ and $\vartheta = zD_z$. For nonzero $b \in \mathbb{K}[z]$, $\text{lc}_z(b) \in \mathbb{K}^\times$ is its leading coefficient in z ; lc remains the leading Hahn coefficient. “For large δ ” quantifies over a final segment of Γ and presupposes no countable cofinal sequence. A nonzero linear differential operator is written $L = \sum_{r=0}^{R_L} b_r(z)D_z^r$ with $b_r \in \mathbb{K}[z]$, the derivative acting on what stands to its right. D_z kills $\mathbb{K}$ and is not the Berarducci–Mantova derivation.

For $k = \mathbb{C}$ one has $\mathbb{K} = K_\Gamma$ and $\mathcal{E}_\Gamma(k) = \mathcal{E}_\Gamma$. As for the parts before it, the generality in k is recorded for this part.

The source. Source 15 is *Polynomial-Composition Rigidity at Surreal Scales: Weighted nonlinear equations, finite degree bounds, and omnific Diophantine universality* (24 pages, US letter, 23 September 2026), delivered as the archive `Surreal_Polynomial_Composition_Rigidity` (batch 31, manuscript 01). It consulted the repository at

bcac55ae6fd3f2b354e568ba1d2e94d496a2cc59

through a connector (a container clone failed), reading the root README, the catalogue, this report’s README and source 10’s proof audit, an omnific Diophantine audit, and the READMEs of the quotient and lattice reports; its shipped file `15-polynomial-composition-SOURCE_AUDIT.md` lists them. At that pin this report was the five-source text (sources 08–12). Source 15’s description of the project, linear rigidity, dilations, first-order nonlinear rigidity and “a higher-order theorem in fields whose value group has no order unit”, was accurate there; the nonlinear picture has since been completed by Theorems H, I and K, which it does not use. Its statement that “a keyword search for Mahler material returned no matches” was not correct at its pin: the single-dilation report already discussed Mahler difference algebra and cited Favrejon and Roques on linear Mahler equations with Hahn-series solutions [46, 47], and the computable-surreals report cited the same paper; this report mentioned only the Skolem–Mahler–Lech theorem. The comparison is corrected in Section 37.5.

Renamed symbols. The following symbols of source 15 would collide with symbols of the earlier parts. They are renamed here, in the new material only. The most important row is the first: source 15’s two indicial polynomials carry hol’s names in *reverse*.

Source 15	Here	Reason
exact indicial χ_L ; residue indicial $\bar{\chi}_L$	I_L ; χ_L	For the linear differential polynomial $P = L[Y]$ they are exactly I_P and χ_P of (14.6): <i>source 15’s χ_L is this report’s I_L, not its χ_L.</i>
shift h_L ; α_L	w_L ; β_L	The corner weight w_P and value β_P of (14.5) for $P = L[Y]$.
radius ρ ; height $M_f(\rho)$; $I_f(\rho)$; $\text{in}_\rho(f)$	scale δ ; $-\gamma_\delta(f)$; $\text{Act}_\delta(f)$; $\phi_\delta(f)$	The Gauss data; source 15’s height is the <i>negative</i> of the Gauss value, so its inequalities are reversed here.
reference radius R ; $\mathcal{T}_\rho$; σ (Proposition 2.4)	δ_{ref} ; $\mathcal{E}^{(\delta)}$; δ'	R is an order bound, $\mathcal{T}_\rho$ the theta descent, σ a dilation.
$\kappa = v(\text{lc}_z P)$	$v(\text{lc}_z P)$	κ_P is the degree cutoff (15.11).
leading coefficient $\text{lc}(P)$ of a polynomial	$\text{lc}_z(P)$	lc is the leading Hahn coefficient of Section 1.2.
$A(z)$; $Q(z, Y)$; q lower arguments; s top factors; T_ν	$c_*(z)$; $\mathbf{G}(z, Y)$; $\bar{m}$; $\bar{s}$; L'_ν	$A[f]$, Q_s , q (a dilation), s (an order) and T (an indeterminate) have other meanings.
composition weight $w(\mathbf{a})$; H , b ; H_{top} , $U_{\mathbf{a}}$; $h_{\mathbf{a}}$, $b_{\mathbf{a}}$	$\text{cw}(\mathbf{a})$; w_{top} , β_{top} ; γ_{top} , $\gamma_{\mathbf{a}}$; $w_{\mathbf{a}}$, $\beta_{\mathbf{a}}$	w_P is a corner weight and wt an Euler weight; $H \in \Gamma$.
resonance set $\mathcal{R}$; bound B	Rsn ; B_{cmp}	$\mathcal{R}$ is a set of right sides in Section 15; B a threshold.
$D = \mathbb{Z}$ or $\mathbb{Z}[i]$; $R_{D, \Gamma}$; R	$\mathfrak{o}$; $\mathfrak{D}_{\mathfrak{o}, \Gamma}$; $\mathfrak{R}$	D_z is the derivative.
$E_z = zD$; projectors S_j ; $L_{d, N}$	ϑ ; Pr_j ; $\text{Ann}_{d, N}$	E_η is the formal exponential, S a shift.
$q = \max \mathbf{a} $, $T = \max \sum j a_j$ (encoder)	$\deg p$, $\text{wd}(p)$	As above.
theta F ; even series H ; char- p series H ; rank-infinity G	Θ_p ; f_{ev} ; U of Example 17.3; G (kept)	(8.1); F is (11.2).
“Mahler equation”	kept, with its meaning	Here it means equations in $f(z^{b^j})$, argument substitution; not the exponent dilation S_d of the autonomous-dilation report [48], and not the scalar dilation of the single-dilation report.
Theorems 5.1, 6.1, 7.1, 8.2, 9.1	Theorem N and Theorems 36.12, 36.15, 37.2, 37.3	Theorems A–N of this report are different theorems.

The shipped audit file 15-polynomial-composition-SOURCE_AUDIT.md keeps the source’s notation.

Words with two meanings. A *pullback* is $f \mapsto f \circ P$ for a polynomial P ; a *dilation* is the degree-one case $P = \lambda z$. The *composition weight* of a monomial is Definition (36.4); it is neither the corner weight w_P nor the Euler weight wt . An *exact* resonance is an ordinary integer root of I_L , a *residue* resonance one of χ_L ; Example 15.5 shows they differ.

Results printed once. Source 15’s Definition 2.1 and Lemma 2.2 are Definition 1.2 and Lemma 2.3 (a); its Theorem 2.3, the exact entire-support criterion, is Proposition 20.2, proved

there by the same two-scale argument; its Proposition 2.4 is Proposition 2.7 together with closure under polynomial composition, which is contained in `ent:prop:operations` of the entire-functions report [12] (entire composition) and is also proved directly below; its Lemma 3.1 is Corollary 14.3 with the derivative estimate of Lemma 36.6 for $L = D_z^r$; its Proposition 3.2 is the residue argument of Theorem 33.5, over any k (compare Proposition 2.8); the first half of its Lemma 3.3 is Lemma 14.5; its Remark 3.6, that the Gauss value is not $v(f(t^{-\delta}))$, is the remark after (14.1); its Remark 4.4, $L = zD_z - N + \varepsilon$, is Example 15.5; the conformal embedding in its Section 8.1 is (2.4); the partial theta series, the characteristic- p series, the rank-infinity series and the failure on the full Conway field in its Section 11 are (8.1), Example 17.3, the series F of Theorem 11.2 (source 15 uses $\bigoplus_{n \geq 1} \mathbb{R}e_n$, this report Γ_∞ of (11.1)), Section 11.2 and Proposition 27.6; and its remark that uncountable cofinality forces $\mathcal{E}_\Gamma(k) = \mathbb{K}[z]$ is the remark in Section 31.3.

Where each numbered result went. Proposition 3.5 (pullback profile), Lemma 3.4 (integer convexity) and the second half of Lemma 3.3 are Proposition 36.5, Lemma 36.4 and Lemma 36.3; Lemmas 4.1, 4.3 and Proposition 4.2 are Lemmas 36.6, 36.8 and Proposition 36.7; Theorem 5.1 and Corollaries 5.2, 5.3 are Theorem N (with its proof in Section 36.4) and Corollaries 36.9, 36.11; Theorem 6.1, Corollary 6.2 and Proposition 6.3 are Theorem 36.12, Corollary 36.13 and Proposition 36.14; Theorem 7.1 is Theorem 36.15; Proposition 8.1 and Theorem 8.2 are Proposition 37.1 and Theorem 37.2; Theorem 9.1, Remark 9.2 and Corollaries 9.3, 9.4 are Theorem 37.3, Remark 37.4 and Corollaries 37.5, 37.6; the examples of Sections 10 and 11 are in Sections 37.3 and 37.4; Questions 12.1–12.8 are Questions 19.55–19.62; Section 13 and the appendices are merged into Sections 37.5, 37.6 and 18.

The merge's choices. The review of source 15 for this merge found its proofs correct; the worked equations were rechecked with SymPy. The merge adds three points, each marked. First, Remark 36.10 states the relation to Theorems B and J, to Theorem 10.2 and to Corollary 11.3: with a strictly dominant argument of degree at least two the order-unit dichotomy disappears. Second, the Gauss data are those of this report, with source 15's signs reversed, and its two indicial polynomials are identified with I_P and χ_P . Third, the comparison with the Mahler and omnific literature of the collection is completed (Section 37.5). Source 15's theorems, proofs, examples and non-claims are otherwise printed as delivered, in the renamed notation.

36.2 Pullback profiles

For $\delta \in \Gamma$ let $\mathcal{E}^{(\delta)}$ be the set of $\sum_n b_n z^n \in \mathbb{K}[[z]]$ with $v(b_n) - n\delta \rightarrow +\infty$ cofinally. After $z = t^{-\delta}X$ its elements have coefficients whose values tend cofinally to infinity; it is closed under products (finite convolutions, with the two minima normalized to zero), sums and D_z (the multiplier n has value zero), and a family (H_n) in it whose minimal weighted coefficient values tend cofinally to infinity has a coefficientwise sum in it, which may be regrouped. By Proposition 20.2, $\mathcal{E}_\Gamma(k) = \bigcap_\delta \mathcal{E}^{(\delta)}$.

Proposition 36.2 (Polynomial composition; source 15). *For $f = \sum a_n z^n \in \mathcal{E}_\Gamma(k)$ and $P \in \mathbb{K}[z]$, the series $f \circ P = \sum_n a_n P^n$ is defined coefficientwise by strongly summable families and lies in $\mathcal{E}_\Gamma(k)$; $D_z(f \circ P)$ may be computed termwise. P need not vanish at 0.*

Proof. Fix δ and put $\delta' = \max_j(j\delta - v([z^j]P))$. Every coefficient of $a_n P^n$, weighted at δ , has value at least $v(a_n) - n\delta'$, and these tend cofinally to infinity by Proposition 20.2; so $\sum a_n P^n$ exists in $\mathcal{E}^{(\delta)}$, with coefficients independent of δ . It lies in every $\mathcal{E}^{(\delta)}$. The same bounds justify termwise differentiation. $\square$

Ordinary formal substitution into a power series needs $P(0) = 0$; strong entireness supplies the convergence that replaces it.

Lemma 36.3 (Superlinear escape; source 15). *If $f \in \mathcal{E}_\Gamma(k)$ is not a polynomial, then for every integer C , every positive integer d and every fixed $\kappa \in \Gamma$, $\gamma_{d\delta-\kappa}(f) + C\delta \rightarrow -\infty$ as $\delta \rightarrow +\infty$, that is, it is eventually below every element of Γ .*

Proof. Choose m with $md > C$ and $a_m \neq 0$. Then $\gamma_{d\delta-\kappa}(f) + C\delta \leq v(a_m) + m\kappa - (md - C)\delta$, and positive integer multiples of an unbounded variable are cofinal, whether or not Γ has an order unit. $\square$

Lemma 36.4 (Integer convexity; source 15). *Let $d > e > 0$ be integers and $\sigma, \tau, \delta_{\text{ref}} \in \Gamma$ with $d\tau \leq e\sigma + (d - e)\delta_{\text{ref}}$. Then for every nonzero $f \in \mathcal{E}_\Gamma(k)$*

$$d\gamma_\tau(f) \geq e\gamma_\sigma(f) + (d - e)\gamma_{\delta_{\text{ref}}}(f). \quad (36.1)$$

Proof. For $a_n \neq 0$, $n \geq 0$ gives $d(v(a_n) - n\tau) \geq e(v(a_n) - n\sigma) + (d - e)(v(a_n) - n\delta_{\text{ref}})$, which is at least the right side of (36.1); take the minimum. No division in Γ occurs. $\square$

Proposition 36.5 (Exact pullback profile; source 15). *Let $P = cz^d + \dots$ have degree $d \geq 1$. For all large δ , depending on P only, and every nonzero $f \in \mathcal{E}_\Gamma(k)$,*

$$\gamma_\delta(f \circ P) = \gamma_{d\delta-v(c)}(f), \quad \text{Act}_\delta(f \circ P) = d \text{Act}_{d\delta-v(c)}(f), \quad \phi_\delta(f \circ P)(X) = \phi_{d\delta-v(c)}(f)(\bar{c}X^d), \quad (36.2)$$

with $\bar{c} = \text{res}(t^{-v(c)}c)$.

Proof. For large δ the top term uniquely attains the Gauss value of P : $v(c) - d\delta < v([z^j]P) - j\delta$ for $j < d$. With $\sigma = d\delta - v(c)$, the polynomial $t^\sigma P(t^{-\delta}X)$ has integral coefficients and residue $\bar{c}X^d$, and Proposition 36.2 with reduction gives $\text{res}(t^{-\gamma_\sigma(f)}f(P(t^{-\delta}X))) = \phi_\sigma(f)(\bar{c}X^d)$, a nonzero polynomial with support $d \text{Act}_\sigma(f)$. $\square$

36.3 Operators and their two indicial polynomials

For $L = \sum_r b_r(z)D_z^r \neq 0$ put

$$w_L = \max_{b_r \neq 0} (\deg b_r - r), \quad I_L(T) = \sum_{\deg b_r - r = w_L} \text{lc}_z(b_r)(T)_T, \quad \beta_L = \min_{\deg b_r - r = w_L} v(\text{lc}_z b_r), \quad (36.3)$$

and $\chi_L(T) = \sum \text{res}(t^{-\beta_L} \text{lc}_z b_r)(T)_T$ over the r with $\deg b_r - r = w_L$ and $v(\text{lc}_z b_r) = \beta_L$. For the linear differential polynomial $L[Y]$ these are its corner weight w_P , full corner polynomial I_P , corner value β_P and residue corner polynomial χ_P ((14.5), (14.6)). Both I_L and χ_L are nonzero, since distinct falling factorials are independent, and χ_L has finitely many roots in $\mathbb{N}$.

Lemma 36.6 (Uniform operator bound; source 15). *For all large δ , depending on L only, and every $u \in \mathcal{E}_\Gamma(k)$, $\gamma_\delta(Lu) \geq \gamma_\delta(u) - w_L\delta + \beta_L$.*

Proof. The term $c_{rj}z^jD_z^r u$, $c_{rj} = [z^j]b_r \neq 0$, has Gauss value at least $\gamma_\delta(u) - (j - r)\delta + v(c_{rj})$. For $j - r < w_L$ this exceeds the bound for large δ ; at $j - r = w_L$, $j = \deg b_r$ and $v(c_{rj}) \geq \beta_L$. $\square$

Proposition 36.7 (Exact eventual operator profile; source 15). *If $u \in \mathcal{E}_\Gamma(k)$ is not a polynomial, then for all large δ*

$$\gamma_\delta(Lu) = \gamma_\delta(u) - w_L\delta + \beta_L, \quad \phi_\delta(Lu) = \sum_{n \in \text{Act}_\delta(u)} [X^n] \phi_\delta(u) \chi_L(n) X^{n+w_L};$$

in particular $Lu \neq 0$.

Proof. Take δ beyond Lemma 36.6 and so large that every active index exceeds R_L and every root of χ_L in $\mathbb{N}$ (Lemma 14.5). Normalizing at the proposed value, the terms of smaller shift and the top-shift terms with coefficient value above β_L have positive value, and at input degree n the surviving terms have the residue multiplier $\chi_L(n) \neq 0$; inactive coefficients contribute nothing. The exponents $n + w_L$ are distinct and nonnegative. $\square$

This is the corner identity of Section 14.4 for the linear polynomial $L[Y]$, applied at every exterior scale.

Lemma 36.8 (Exact polynomial leading term; source 15). *For a polynomial u of degree n with leading coefficient a , $\deg Lu \leq n + w_L$; if $I_L(n) \neq 0$ equality holds with leading coefficient $aI_L(n)$, and if $I_L(n) = 0$ the degree is smaller or $Lu = 0$.*

Proof. The coefficient at degree $n + w_L$ is the top-shift contribution of az^n , namely $aI_L(n)$; everything else has smaller degree. If $n + w_L < 0$ every term vanishes, and the falling factorials give $I_L(n) = 0$. $\square$

A residue resonance $\chi_L(n) = 0$ need not be an exact one (Example 15.5): residue resonance controls the exterior proof, exact resonance the degree bounds.

36.4 The weighted composition theorem

Fix P of degree $d \geq 2$, nonzero operators $L_1, \dots, L_{\bar{s}}$ ($\bar{s} \geq 1$), a nonzero $c_* \in \mathbb{K}[z]$, and for $1 \leq \nu \leq \bar{m}$ ($\bar{m} \geq 0$) nonconstant P_ν of degree $e_\nu < d$ and nonzero operators L'_ν . Let $\mathbf{G}(z, Y_1, \dots, Y_{\bar{m}}) = \sum_{\mathbf{a} \in \mathcal{A}} c_{\mathbf{a}}(z) Y^{\mathbf{a}}$ in collected form, with nonzero $c_{\mathbf{a}} \in \mathbb{K}[z]$ ($\mathcal{A} = \emptyset$ if $\mathbf{G} = 0$), and put

$$\text{cw}(\mathbf{a}) = \sum_{\nu=1}^{\bar{m}} a_\nu e_\nu. \quad (36.4)$$

The equation is

$$c_*(z) \prod_{\ell=1}^{\bar{s}} L_\ell(f(P(z))) = \mathbf{G}(z, L'_1(f(P_1(z))), \dots, L'_{\bar{m}}(f(P_{\bar{m}}(z)))). \quad (36.5)$$

Proof of Theorem N. Suppose f is not a polynomial. Put $\sigma = d\delta - v(\text{lc}_z P)$ and $\sigma_\nu = e_\nu\delta - v(\text{lc}_z P_\nu)$, and take δ large for the finitely many profiles below. By Proposition 36.5 and Lemma 14.5, $f \circ P$ is not a polynomial, so Proposition 36.7 applies to each $L_\ell(f \circ P)$, and by Corollary 14.3 the left side has the exact Gauss value

$$\gamma_{\text{top}}(\delta) = \bar{s} \gamma_\sigma(f) - w_{\text{top}}\delta + \beta_{\text{top}}, \quad w_{\text{top}} = \deg c_* + \sum_{\ell} w_{L_\ell}, \quad \beta_{\text{top}} = v(\text{lc}_z c_*) + \sum_{\ell} \beta_{L_\ell}. \quad (36.6)$$

Choose one δ_{ref} with $(d - e_\nu)\delta_{\text{ref}} \geq e_\nu v(\text{lc}_z P) - d v(\text{lc}_z P_\nu)$ for every ν ; then $d\sigma_\nu \leq e_\nu\sigma + (d - e_\nu)\delta_{\text{ref}}$ identically in δ , and Lemma 36.4 gives $d\gamma_{\sigma_\nu}(f) \geq e_\nu\gamma_\sigma(f) + (d - e_\nu)\gamma_{\delta_{\text{ref}}}(f)$. With $w_{\mathbf{a}} = \deg c_{\mathbf{a}} + \sum_\nu a_\nu w_{L'_\nu}$ and $\beta_{\mathbf{a}} = v(\text{lc}_z c_{\mathbf{a}}) + \sum_\nu a_\nu \beta_{L'_\nu}$, the monomial $\mathbf{a}$ has, by Lemma 36.6 and Proposition 36.5, Gauss value at least $\gamma_{\mathbf{a}}(\delta) = \sum_\nu a_\nu \gamma_{\sigma_\nu}(f) - w_{\mathbf{a}}\delta + \beta_{\mathbf{a}}$, also when a factor vanishes. Hence, with $|\mathbf{a}| = \sum a_\nu$,

$$d(\gamma_{\mathbf{a}} - \gamma_{\text{top}}) \geq -(\bar{s}d - \text{cw}(\mathbf{a}))\gamma_\sigma(f) + d(w_{\text{top}} - w_{\mathbf{a}})\delta + (d|\mathbf{a}| - \text{cw}(\mathbf{a}))\gamma_{\delta_{\text{ref}}}(f) + d(\beta_{\mathbf{a}} - \beta_{\text{top}}). \quad (36.7)$$

The coefficient $\bar{s}d - \text{cw}(\mathbf{a})$ is a positive integer, so by Lemma 36.3 the right side is eventually above every element of Γ . On a common final segment, then, every monomial of the right side has strictly larger Gauss value than the left side, which is impossible for an identity; if $G = 0$ the nonzero left side already contradicts it. $\square$

Corollary 36.9 (Mixed differential–Mahler equations; source 15). *Let $P_0, \dots, P_m \in \mathbb{K}[z]$ be nonconstant with $d_m = \deg P_m > \max_{j < m} \deg P_j$ and $d_m \geq 2$, let $L_0, \dots, L_m$ be linear differential operators over $\mathbb{K}[z]$ with $L_m \neq 0$, and $g \in \mathbb{K}[z]$. Every strongly entire solution of*

$$\sum_{j=0}^m L_j(f(P_j(z))) = g(z) \quad (36.8)$$

is a polynomial; so is every solution with rational coefficients, after a common denominator is cleared.

Proof. Move the lower terms to the right: $\bar{s} = 1$, $P = P_m$, $c_* = 1$, a linear G whose monomials have weights $\deg P_j < d_m$, and a forcing term of weight zero. Clearing denominators multiplies the identity by a polynomial and moves no coefficient through a derivative. $\square$

With $P_j = z^{b_j}$, $b \geq 2$, every strongly entire solution of a linear Mahler equation, or of one whose pullbacks are multiplied by finite-order differential operators, is a polynomial. The pure complex-analytic antecedent is Bell, Coons and Rowland’s [42, Lemma 6]; the arbitrary-rank mixed statement is source 15’s.

Remark 36.10 (Degree at least two removes the order-unit dichotomy; merge). The lower arguments in Corollary 36.9 may be dilations λz , of degree one. So every linear equation in D_z , finitely many dilations σ_λ , and at least one polynomial argument of degree $d \geq 2$ strictly above the degrees of all other arguments has only polynomial strongly entire solutions, *for every* Γ , with or without an order unit and whatever the valuations of the dilation parameters. By contrast, Theorem B, Theorem J and Corollary 11.3 concern equations all of whose arguments have degree one; there a nonpolynomial solution exists exactly when some (relative) dilation valuation is an order unit, and the partial theta series is the witness. A dominant argument of degree at least two therefore detects nothing about order units. Theorem 10.2 and Corollary 29.6 are the degree-one mixed statements. The hypothesis of a *strictly* dominant degree is used: Section 37.4 shows that equal degrees of the top arguments, including degree one, allow nonpolynomial solutions. Nor does Theorem N say anything about algebraic differential equations without a pullback, which are Theorems D, H, I and K.

Corollary 36.11 (A necessary Newton-weight condition; source 15). *If $f \in \mathcal{E}_\Gamma(k)$ is not a polynomial, every relation of the form (36.5) that it satisfies has a monomial of weight at least $\bar{s}d$.*

This restricts possible relations; it is not an algebraic or differential independence statement, and relations with equal weights, or with cancellation among several top-weight expressions, are not classified.

36.5 Degree bounds and finite coefficient loci

Under the hypotheses of Theorem N let $\text{Rsn} = \{N \in \mathbb{N} : I_{L_\ell}(dN) = 0 \text{ for some } \ell\}$, a finite set, and

$$B_{\text{cmp}} = \max\left(\{0\} \cup \text{Rsn} \cup \left\{\left\lfloor \frac{w_{\mathbf{a}} - w_{\text{top}}}{\bar{s}d - \text{cw}(\mathbf{a})} \right\rfloor : \mathbf{a} \in \mathcal{A}\right\}\right) \in \mathbb{N}, \quad (36.9)$$

a ratio of ordinary integers, with no division in Γ .

Theorem 36.12 (Resonant degree bound; source 15). *Every nonzero strongly entire solution of (36.5) has degree at most B_{cmp} ; the zero series is tested directly, and the bound need not be attained.*

Proof. The solution is a polynomial. If its degree N exceeds B_{cmp} , then $N \notin \text{Rsn}$, and by Lemma 36.8 the left side has degree $\bar{s}dN + w_{\text{top}}$ with leading coefficient $\text{lc}_z(c_*)a^{\bar{s}}\text{lc}_z(P)^{\bar{s}N} \prod_{\ell} I_{L_\ell}(dN) \neq 0$, while the monomial $\mathbf{a}$ has degree at most $\text{cw}(\mathbf{a})N + w_{\mathbf{a}}$, which the definition of B_{cmp} makes smaller. $\square$

Corollary 36.13 (Linear bound; source 15). *For (36.8), with $w_j = w_{L_j}$, $d = d_m$ and $\text{Rsn}_m = \{N \in \mathbb{N} : I_{L_m}(dN) = 0\}$, every nonzero entire solution has degree at most*

$$B_{\text{lin}} = \max\left(\{0\} \cup \text{Rsn}_m \cup \left\{\left\lfloor \frac{w_j - w_m}{d - \deg P_j} \right\rfloor : j < m, L_j \neq 0\right\} \cup \left\{\left\lfloor \frac{\deg g - w_m}{d} \right\rfloor : g \neq 0\right\}\right).$$

Substituting $f = u_0 + u_1z + \cdots + u_Bz^B$ with $B = B_{\text{cmp}}$ and comparing coefficients gives finitely many polynomial equations in the u_j over the field generated by the data, whose solutions in $\mathbb{K}$ are exactly the entire solutions; linear equations give a linear system. This is the composition analogue of Theorem 15.3 and Corollary 15.6. It is not a decision procedure: one must represent the data and decide equalities, find the integer roots in Rsn , and solve the finite system. For rational data the first two are effective; for arbitrary named surreal coefficients no algorithm is asserted, and over the omnific integers even integer data can be undecidable (Corollary 37.5).

Proposition 36.14 (Extension-stable coefficient description; source 15). *For an injective field map $\mathbb{K} \hookrightarrow \mathbb{K}'$ between Hahn fields of the standing type, with the finite data transported, the same bound and the same finite coefficient equations describe the entire solutions over $\mathbb{K}'$; no entire solution of higher degree appears.*

Proof. Degrees, shifts and the vanishing of $I_{L_\ell}(dN)$ are preserved and reflected, so B_{cmp} is unchanged; apply Theorems N and 36.12 over $\mathbb{K}'$. Polynomials are entire. $\square$

This concerns the solutions of a fixed equation, which are polynomials; an arbitrary infinite series can lose entireness under extension (Section 37.4).

36.6 Vector equations

For $F = (f_1, \dots, f_r) \in \mathcal{E}_\Gamma(k)^r$ put $\gamma_\delta(F) = \min_i \gamma_\delta(f_i)$; Lemmas 36.4 and 36.3 hold for it, minimizing over components and indices.

Theorem 36.15 (Matrix polynomial-composition rigidity; source 15). *Let $P_0, \dots, P_m$ be non-constant, $d_m = \deg P_m > \max_{j < m} \deg P_j =: \max d_j$, $d_m \geq 2$, $A_0, \dots, A_m \in \text{Mat}_r(\mathbb{K}[z])$ with $\det A_m \neq 0$, and $b \in \mathbb{K}[z]^r$. Every $F \in \mathcal{E}_\Gamma(k)^r$ with $\sum_{j=0}^m A_j(z)F(P_j(z)) = b(z)$ is a polynomial vector. With $a = \det A_m$, $C_j = \text{adj}(A_m)A_j$, $c = \text{adj}(A_m)b$ and entry degrees, the largest component degree of a nonzero solution is at most*

$$B_{\text{mat}} = \max\left(\{0\} \cup \left\{\left\lfloor \frac{\deg C_j - \deg a}{d_m - d_j} \right\rfloor : C_j \neq 0\right\} \cup \left\{\left\lfloor \frac{\deg c - \deg a}{d_m} \right\rfloor : c \neq 0\right\}\right).$$

Proof. Multiplying by the adjugate gives $aF(P_m) = c - \sum_{j < m} C_j F(P_j)$. If a component is nonpolynomial, the left side has Gauss value $\gamma_{d_m \delta - v(\text{lc}_z P_m)}(F) - \deg(a)\delta + v(\text{lc}_z a)$ for large δ (componentwise, by Proposition 36.5), while the right terms are bounded below by $\gamma_{d_j \delta - v(\text{lc}_z P_j)}(F)$ plus a fixed integer multiple of δ and a fixed element of Γ ; the argument of Theorem N with weights $d_j < d_m$ makes every right term strictly larger in value. For the degree bound compare $\deg a + d_m N$ with $\deg c$ and $\deg C_j + d_j N$. $\square$

Nonsingularity cannot be dropped: with $A_m = \text{diag}(1, 0)$ and all other terms zero, every $(0, f)$ is a solution. Matrices of differential operators are not treated (Question 19.57).

37 Omnific coefficients, universality and boundaries of composition rigidity

37.1 Omnific coefficients and linear solution modules

For $(\mathfrak{o}, k) = (\mathbb{Z}, \mathbb{R})$ or $(\mathbb{Z}[i], \mathbb{C})$ put $\mathfrak{D}_{\mathfrak{o}, \Gamma} = \mathfrak{o} \oplus \Pi_{k, \Gamma}$, with $\Pi_{k, \Gamma} = \{\sum_{\gamma < 0} a_\gamma t^\gamma\}$ (well-ordered supports) and the constant-term retraction $\text{ct} : \mathfrak{D}_{\mathfrak{o}, \Gamma} \rightarrow \mathfrak{o}$, a ring homomorphism because $\Pi_{k, \Gamma}$ is closed under sums, under products and under multiplication by $\mathfrak{o}$, all supports staying negative. Through (2.4), allowing all of $\mathbf{No}$ as exponents (each support a set) gives $\mathbf{Oz} = \mathbb{Z} \oplus \Pi$ and $\mathbf{Oz}[i]$, with the same retraction [41, 37]; these are not new.

Proposition 37.1 (An elementary coefficient restriction; source 15). *A strongly entire series with all coefficients in $\mathfrak{D}_{\mathfrak{o}, \Gamma}$ is a polynomial, with no equation.*

Proof. Nonzero coefficients have value at most 0; infinitely many would violate Proposition 20.2 at $\gamma = \beta = 0$. $\square$

So for omnific coefficients the analytic content of Theorem N is automatic; what an equation adds is its degree bound and the structure of its coefficient locus.

Theorem 37.2 (Linear omnific solution modules; source 15). *Consider (36.8) with all arguments, operator coefficients and forcing in $\mathfrak{o}[z]$, $\mathfrak{o} = \mathbb{Z}$ or $\mathbb{Z}[i]$, and let $\mathfrak{R}$ be $\mathfrak{D}_{\mathfrak{o}, \Gamma}$, or $\mathbf{Oz}$, or $\mathbf{Oz}[i]$, accordingly. An effective procedure on the data either certifies that there is no polynomial solution in $\mathfrak{R}[z]$ or gives $f_0, h_1, \dots, h_r \in \mathfrak{o}[z]$ such that the solutions are exactly $f_0 + \sum_i u_i h_i$, $u_i \in \mathfrak{R}$, with unique parameters. Existence over $\mathfrak{R}$ is equivalent to existence over $\mathfrak{o}$, and for a fixed Hahn workspace these are also exactly its strongly entire solutions with coefficients in $\mathfrak{R}$.*

Proof. The degree argument of Lemma 36.8 and Corollary 36.13 is algebraic and applies to polynomials over $\mathbf{No}$ and $\mathbf{No}[i]$; the integer roots of $I_{L_m}(d_m T)$ are computable. Substituting a polynomial of degree B_{lin} gives $Mu = b$ over $\mathfrak{o}$. Take a Smith normal form $UMV = S$ over the principal ideal domain $\mathfrak{o}$, and put $u = Vy$, $b' = Ub$. The equations are $s_i y_i = b'_i$ and zero rows. A solution over $\mathfrak{R}$ gives one over $\mathfrak{o}$ by ct , and conversely; so divisibility and zero-row tests decide existence. Since $\mathfrak{R}$ is a domain, the homogeneous diagonal equations force $y_i = 0$ for $s_i \neq 0$, the other y_i are free, and V gives the basis. The last assertion is Proposition 37.1. $\square$

For $\mathfrak{o} = \mathbb{Z}$ and $\mathfrak{R} = \mathbf{Oz}$ the linear algebra is `odg:thm:smith` and `odg:thm:linear` of the omnific Diophantine report [41], applied to the finite system produced by the degree bound; the Gaussian case is the same argument over $\mathbb{Z}[i]$. The proof uses neither that $\mathbf{Oz}$ is a principal ideal domain nor any factorization theory.

37.2 Universality and undecidability

Fix $N \geq 0$. For $0 \leq j \leq N$ put

$$\text{Pr}_j = N! \prod_{0 \leq k \leq N, k \neq j} \frac{\vartheta - k}{j - k} \in \mathbb{Z}[z] \langle D_z \rangle; \quad (37.1)$$

the denominator $(-1)^{N-j} j! (N-j)!$ divides $N!$, and $\vartheta^i = \sum_j S(i, j) z^j D_z^j$ with Stirling numbers $S(i, j)$ of the second kind. Each Pr_j has shift zero and $\text{Pr}_j f = N! u_j z^j$ for $f = \sum_{j \leq N} u_j z^j$: interpolation in the Euler eigenvalue.

Theorem 37.3 (Universal hypersurface encoding; source 15). *For every nonzero $p = \sum_{\mathbf{a}} p_{\mathbf{a}} U^{\mathbf{a}} \in \mathbb{Z}[U_0, \dots, U_N]$ put $\text{wd}(p) = \max\{\sum_j j a_j : p_{\mathbf{a}} \neq 0\}$, $d = \deg p + \text{wd}(p) + 2$ and $\text{Ann}_{d,N} = \prod_{k=0}^N (\vartheta - dk)$. The equation*

$$\text{Ann}_{d,N}(f(z^d)) = \sum_{\mathbf{a}} p_{\mathbf{a}} (N!)^{\deg p - |\mathbf{a}|} z^{\text{wd}(p) - \sum_j j a_j} \prod_{j=0}^N (\text{Pr}_j f(z))^a_j \quad (37.2)$$

has integer data and satisfies the hypothesis of Theorem N; over every Hahn field of the standing type its entire solutions are exactly $f = \sum_{j \leq N} u_j z^j$ with $p(u_0, \dots, u_N) = 0$, and its bound (36.9) is exactly N .

Proof. Here $\bar{s} = 1$ and every lower argument is z , so $\text{cw}(\mathbf{a}) = |\mathbf{a}| \leq \deg p < d$, and the solutions are polynomials. $\text{Ann}_{d,N}$ has shift 0 and $I(T) = \prod_k (T - dk)$, so $\text{Rsn} = \{0, \dots, N\}$; the monomial shifts are at most $\text{wd}(p)$ and $d - |\mathbf{a}| \geq \text{wd}(p) + 2$, so every fraction in (36.9) has floor 0 and $B_{\text{cmp}} = N$. For f of degree at most N the left side vanishes term by term and the right side is $(N!)^{\deg p} z^{\text{wd}(p)} p(u_0, \dots, u_N)$, zero exactly when $p(u) = 0$ in characteristic zero. $\square$

The margin $d = \deg p + \text{wd}(p) + 2$, rather than merely $d > \max(\deg p, \text{wd}(p))$, is what makes the bound exactly N in every case.

Remark 37.4 (What universality says; source 15). After clearing denominators every affine hypersurface over $\mathbb{Q}$ arises. A finite degree certificate therefore implies neither finitely many solutions, nor a smooth solution set, nor a simple classification: the coefficient loci carry the full complexity of affine algebraic geometry.

Corollary 37.5 (Omnific undecidability inside the rigid class; source 15). *No algorithm decides, from the integer data of an equation of Theorem N, whether it has a polynomial solution with coefficients in $\mathbf{Oz}$; nor, in any fixed real Hahn workspace, whether it has a strongly entire solution with coefficients in $\mathfrak{D}_{\mathbb{Z},\Gamma}$.*

Proof. By Theorem 37.3 and the retraction ct , the equation for p has such a solution exactly when p has a zero in $\mathbb{Z}^{N+1}$; the Davis–Putnam–Robinson–Matiyasevich theorem [44] makes that undecidable. $\square$

Undecidability of polynomial equations over $\mathbf{Oz}$ itself is `odg:cor:H10` of the omnific Diophantine report [41]; what is new is the encoding into a rigid functional equation with a prescribed degree bound.

Corollary 37.6 (A fixed degree bound; source 15). *Using Sun’s undecidability theorem in eleven integer unknowns [45], the undecidability persists for an effectively given subclass in which every entire solution has degree at most ten, the top operator has order eleven and every lower operator order at most ten.*

Proof. Take $N = 10$ in Theorem 37.3; d varies with p , but the resonances and orders do not. $\square$

This applies known Diophantine undecidability; it claims no optimal variable count. The linear ordinary-data subclass is decidable (Theorem 37.2), while the nonlinear subclass is already undecidable at a fixed degree bound.

37.3 Worked equations

Every degree attains the linear bound (source 15). For $d \geq 2$ and $N \geq 0$, $f(z^d) = z^{(d-1)N}f(z)$ has $B_{\text{lin}} = N$; comparing largest and smallest degrees gives $\deg f = \text{ord}_z f = N$, so the entire solutions are exactly cz^N .

A free omnific parameter (source 15). $f(z^2) - f(z) = z^2 - z$ has bound one and entire solutions $z + c$, $c \in \mathbb{K}$; its omnific solutions are $z + c$, $c \in \mathbf{Oz}$, the simplest module of Theorem 37.2.

Resonances that cannot be discarded (source 15). $(\vartheta - dN)f(z^d) = 0$ has the solutions cz^N ; a bound ignoring I_L would leave only constants. For

$$(zD_z - 6)f(z^2 + z) + (z^2D_z + 1)f(z) = -3z^5 - 3z^4 - 2z^3 - 6z^2 - 8z - 5$$

the top resonance is $N = 3$ while the nonresonant estimate is two; for $f = a_0 + \cdots + a_3z^3$ the left side is $-3a_3z^5 + (-2a_2 - 3a_3)z^4 + (-4a_2 - 2a_3)z^3 + (-3a_1 - 3a_2)z^2 - 4a_1z - 5a_0$, and the unique solution is $f = z^3 + 2z + 1$.

Nonlinear weighted equations (source 15). In $f(z^2)^2 = f(z)^3 + (D_z f(z))^2 + z^4 - z^3 - z^2 - 3z - 1$ the top weight is four, the right monomials have weights three and two with shifts 0 and -2 , and the forcing has weight 0 and degree four, so $B_{\text{cmp}} = 1$; $f = az + b$ gives $a^2 = 1$, $a^3 = 1$, $2ab = 3a^2b - 1$, hence the unique solution $f = z + 1$. Likewise $f(z^3) = f(z)^2 + z^6 - z^4 - z^3 - 3z^2 - 2z$ has bound two and unique solution $z^2 + z + 1$. The total degree three of the right side of the first equation exceeds the two factors on the left; the weight, not the degree, is what matters. These solutions were rechecked with SymPy for this merge.

A Diophantine encoder (source 15). For $N = 1$ and $p = U_0^2 + U_1^2 - m$, $m \in \mathbb{Z}$: $\deg p = \text{wd}(p) = 2$, $d = 6$, $\text{Pr}_0 = 1 - \vartheta$, $\text{Pr}_1 = \vartheta$, and the equation is $\vartheta(\vartheta - 6)f(z^6) = z^2((1 - \vartheta)f)^2 + (\vartheta f)^2 - mz^2$. Every entire solution is $a + bz$ with $a^2 + b^2 = m$. For $m = -1$ there is no surreal solution; for $m = 1$ there are surreal and surcomplex families, and the real omnific solutions are exactly ± 1 and $\pm z$, since $a^2 + b^2 = 1$ forces a, b finite, hence ordinary integers (an elementary observation not needed for universality).

37.4 Boundaries

Degree-one arguments are different (source 15). Over $\Gamma = \mathbb{R}$ with $p = t^\gamma$, $\gamma > 0$, the partial theta series Θ_p of (8.1) is strongly entire and satisfies $\Theta_p(z) = 1 + z\Theta_p(pz)$: both arguments have degree one, and Corollary 36.9 does not apply. This is the report's own witness (Theorem 8.1), used here only to delimit the theorem. Equal degrees also allow cancellation at higher degree: for a nonpolynomial strongly entire even series f_{ev} , for instance $\sum_n t^{n^2} z^{2n}$ over $\mathbb{R}((t^\mathbb{R}))$, one has $f_{\text{ev}}(z^2) - f_{\text{ev}}(-z^2) = 0$.

Characteristic zero (source 15). With the prime p and the series U of Example 17.3, also $D_z(U(z^{p+1})) = 0$: infinitely many indices share a residue, and Proposition 36.7 fails.

Rank, cofinality and enlargement (source 15). Over $\bigoplus_{n \geq 1} \mathbb{R}e_n$ ($0 < e_1 \ll e_2 \ll \dots$), with no order unit, $G = \sum_n t^{e_n} z^n$ is strongly entire (as F in Theorem 11.2), so the theorem has nonpolynomial inputs to exclude at infinite rank; with uncountable cofinality every entire series is a polynomial and only the degree bounds and encodings remain informative. Adjoining $h > ne_j$ for all n, j destroys the entireness of G (Section 11.2), which does not contradict Proposition 36.14, and no nonpolynomial series is strongly evaluable on all of **No** (Proposition 27.6).

Example 37.7 (A critical-weight equation; source 15). For $d \geq 2$ the formal solutions over a field of characteristic zero of $f(z^d) = f(z)^d$, a weight-critical case ($\text{cw} = \bar{s}d = d$), are exactly 0 and cz^N with $N \geq 0$ and $c^{d-1} = 1$. If cz^N is the lowest term, degree dN gives $c = c^d$; writing $f = cz^N U$ with $U(0) = 1$, $U(z^d) = U(z)^d$, and a first nonconstant term bz^r of U would give 0 on the left at degree r and $db \neq 0$ on the right. This elementary classification is the function-variable twin of `adr:lem:monomial` of the autonomous-dilation report [48], where the exponent dilation S_d replaces $z \mapsto z^d$.

So the strict inequality is not necessary for every equation. It is sufficient only: the case $\text{cw} = \bar{s}d$ may still be rigid, and several top-weight terms may carry further obstructions; no nonpolynomial characteristic-zero counterexample at the boundary is asserted. Algebraic differential equations without pullbacks, meromorphic solutions and several variables are separate questions.

The formalization order proposed by source 15. Restricted-series tails; finite initial polynomials; integer convexity; polynomial pullback; the operator residue profile; weighted domination; the polynomial degree bound; Euler projectors; arithmetic reduction. The infinite support argument and the finite algebraic certificate should stay separate, and no Python check replaces the former. For linear equations with rational or Gaussian rational data source 15 describes a certificate: the arguments and operator coefficients, the degrees, shifts and I_{L_j} , the full integer-root set of $I_{L_m}(d_m T)$ and B_{lin} , and the coefficient matrix with a row-reduction certificate, or, for integer or Gaussian-integer solutions, a unimodular Smith certificate, since a rational nullspace basis need not be a lattice basis; for non-effective data this is a certificate format, not an algorithm.

37.5 Predecessors, repository comparison and priority

Literature. Bell, Coons and Rowland prove that a complex entire Mahler function is a polynomial [42, Lemma 6]; source 15 inspected the page with that lemma, and does not claim the complex-analytic statement. Nishioka’s monograph [43] is the standard reference for Mahler’s method (publisher metadata and scope only). The Hahn and omnific background is L’Innocente–Mantova [37]; the undecidability inputs are Matiyasevich [44] (metadata and abstract) and Sun [45] (its abstract, eleven unknowns). Source 15 also looked at the abstract of Bell and Coons’s transcendence tests for Mahler functions and the Wikipedia article on surreal numbers, for orientation only. Its searches were not exhaustive. Two relevant references are missing from source 15 and are added by this merge: Faverjon and Roques treat linear Mahler equations with Hahn-series solutions algorithmically [46], as the single-dilation report records [47]; and the autonomous-dilation report [48] records, through Nishioka–Nishioka, Mahler’s 1983 criterion for convergent power-series solutions of $P(f(z), f(z^d)) = 0$ with constant coefficients. Neither concerns strong entireness over an arbitrary-rank Hahn field, and neither is used here; their content was not rechecked by this merge.

Repository comparison at the pin. Source 15 read seven files at `bcac55a`, listed in its audit, not the 91-page text of this report as it then stood, and compared no report theorem by theorem among the 57 then catalogued. Its keyword-search claim is corrected in Section 36.1. Its statements about this report were accurate at the pin. The universal quotient theorem of the quotient report and the curve and lattice classifications of the omnific reports [49, 41] are not reproduced. The autonomous-dilation report, which now discusses autonomous Mahler equations for the exponent dilation, was not yet in the repository at the pin.

Priority. Not claimed: the complex entire-Mahler lemma, Mahler’s method, normal forms and the constant-term retraction, Smith normal forms, MRDP and Sun’s theorem, the support calculus and Gauss data of this report, and the elementary critical classification. The proposed contributions are strict composition-weight rigidity over arbitrary-rank Hahn fields, its exact-resonance degree bound and extension-stable coefficient loci, the matrix version, the linear omnific module theorem as a consequence, and the universal Euler-projector encoding with its undecidability consequences; priority is provisional, and the word “breakthrough”, which the source considered, is expressly disclaimed.

37.6 What the polynomial-composition part does not claim

Every limitation stated by source 15, in its manuscript, audit file or verification record, is kept.

- (P1) No independent referee report, no Lean formalization, no repository build (a container clone failed), and no certified priority; the label “breakthrough” is disclaimed.
- (P2) Bell–Coons–Rowland’s Lemma 6, Nishioka, the MRDP theorem and Sun’s theorem are credited inputs; the variable count is not optimal, and the bound ten is inherited.
- (P3) Entireness is relative to one set-sized Hahn field; nothing is claimed about unrestricted surreal analytic functions or entireness on **No**.
- (P4) D_z is the function-variable derivative, not the Berarducci–Mantova derivation.
- (P5) The general higher-order algebraic differential rigidity problem is not addressed (in this report it is Theorems H, I and K).
- (P6) Polynomiality of entire series with omnific coefficients is elementary (Proposition 37.1).
- (P7) The condition $cw < \bar{s}d$ is sufficient, not necessary; the critical case $cw = \bar{s}d$ and several top products are open.
- (P8) Corollary 36.11 is not an independence statement.
- (P9) The finite coefficient locus is not a decision procedure, and no algorithm is given for arbitrary named surreal coefficients.
- (P10) The matrix theorem needs a nonsingular dominant matrix and does not treat matrix differential operators.
- (P11) Several variables and meromorphic solutions are not treated.
- (P12) Proposition 36.14 concerns fixed equations only.
- (P13) **Oz** is not claimed to be a principal ideal domain.

- (P14) The classification of $f(z^d) = f(z)^d$ is elementary.
- (P15) The 2,369 finite assertions do not cover infinite supports, all ranks or universal statements, and do not certify novelty.
- (P16) The repository comparison was limited to seven files, and the 57-report catalogue was not compared theorem by theorem; its statement that the repository contained no Mahler material was incorrect at the pin.
- (P17) The universal quotient theorem and the curve and lattice results of the omnific reports are not reproduced.
- (P18) *[merge]* Remark 36.10, the identification of source 15's indicial polynomials with I_P and χ_P , and the added Mahler references are the merge's; no historical priority is claimed for them.

38 Hahn-valued recurrences: observables and Noetherian compression

This section and the next contain the material of source 20, delivered in batch 33 after the fourteen principal theorems above had been merged. Its subject is the *ordinary* constant-coefficient recurrence $y_{n+d} = a_{d-1}y_{n+d-1} + \cdots + a_0y_n$ whose coefficients and values lie in a Hahn field, or in $\mathbf{No}[i]$; its index n is always an ordinary natural number. A single surreal term can carry infinitely many real coefficients, and source 20 separates two kinds of information in such a sequence: the leading scale, which is eventually affine on finitely many progressions, and a prescribed lower Hahn or Conway coefficient, which is tame when the characteristic roots are units and can be arbitrary otherwise.

Much of source 20's valuation theory was found independently of source 18 and is printed once, in Sections 28–29. What is new here is the *coefficient* side. All coefficient sequences of a unit-root exponential polynomial lie in one finitely generated algebra of sequences (Theorem 38.2); hence every set-sized family of polynomial conditions on finitely many coefficients at a time reduces, along the orbit, to a finite subfamily, and its hitting set consists of full progressions modulo one period and a finite set (Theorem 38.8). In the opposite direction one fixed Conway coefficient of a first-order omnific recurrence can reproduce any real sequence (Theorem 39.3). The recurrence statements over $\mathbf{No}[i]$, $\mathbf{Oz}$ and $\mathbf{Oz}[i]$, an omnific recurrence whose degree decreases forever, and determinantal profiles of matrix powers complete the part. Source 20's Sections 9–10 on mixed operators re-derive special cases of Theorem J and are recorded in Section 39.4.

38.1 Source, conventions and the merge's choices

Convention 38.1 (Setting of the recurrence part). In Sections 38–39, k is an arbitrary field of characteristic zero with the trivial valuation, Γ is as in Convention 1.1, $\mathbb{K} = k((t^\Gamma))$, and v , lc , res , strong summability, $\mathcal{E}_\Gamma(k)$, D_z and σ_λ are as in Convention 28.1: D_z kills $\mathbb{K}$ and is not the Berarducci–Mantova derivation, and entineness is relative to this one set-sized field. Sequences are indexed by ordinary $n \in \mathbb{N}$. A unit (an element of valuation zero) is written $\lambda = \zeta(1 + \tau)$ as in (28.1). An *exponential polynomial* is $\mathcal{Y}(n) = \sum_{j=1}^m R_j(n)\lambda_j^n$ with $R_j \in \mathbb{K}[U]$ and $\lambda_j \in \mathbb{K}^\times$, as in Section 28.2; it is *unit-root* if every λ_j is a unit. For units $\lambda_1, \dots, \lambda_m$ with residues $\zeta_1, \dots, \zeta_m$ put $\langle \zeta \rangle = \langle \zeta_1, \dots, \zeta_m \rangle \leq k^\times$, a finitely generated abelian group, whose torsion subgroup is therefore finite, and let M_{tor} be the exponent of that torsion subgroup ($M_{\text{tor}} = 1$ when it is trivial).

For $k = \mathbb{C}$ one has $\mathbb{K} = K_\Gamma$ and $\mathcal{E}_\Gamma(k) = \mathcal{E}_\Gamma$. The statements of Section 38 hold verbatim for $\Gamma = 0$, where $\mathbb{K} = k$; Convention 1.1 matters only where scales or entireness enter.

The source. Source 20 is *Valuation Rigidity and Coefficient Universality in Surreal Recurrences: A mixed differential–dilation theorem and the limits of coefficientwise predictability* (22 pages, US letter, 23 September 2026), delivered as the archive `surreal_recurrences_research` (batch 33, manuscript 01; local number 20 in this directory). It consulted the repository at

efc5446229dbf61a97bdd5edae91dba20af9d131,

where this report’s `article.tex` is the blob `9fd5bd460a19448b9078a4ef7f012e647f545e59`; its shipped file `20-recurrences-SOURCE_AUDIT.md` lists the line ranges it read. At that pin this report was the seven-source text (sources 08–14, 137 pages). Sources 15–19 had been placed in this directory, with their audit and code files, in `9d28e28`, an ancestor of the pin, but none of them had yet been merged into the text; source 18 was merged later the same day, in `2c4debb`. Source 20’s description of this report was accurate at the pin: the paragraph after Theorem 10.2 did leave the mixed unit-valuation case unclassified when Γ has an order unit, and pointed to Question 19.4. That statement is *stale* now: source 18 answers Question 19.4 positively for every Γ (Corollary 29.6) and proves the relative-scale criterion of Theorem J, which contains source 20’s Theorems 10.1, 10.3 and 10.4 (Section 39.4). So source 20’s description of its Section 10 as closing “the repository gap”, its Corollary 10.2 (“the case explicitly left open”), and its Remark 10.5 are printed here as an independent second derivation, not as current contributions of this report. Source 20 claims neither the pure differential or dilation theorems, the escape mechanism, the order-unit definition, nor the partial theta series; it credits them to this report.

Renamed symbols. The following symbols of source 20 would collide with symbols of the earlier parts. They are renamed here, in the new material only.

Source 20	Here	Reason
$K = k((t^\Gamma))$, $\mathcal{E}(K)$	$\mathbb{K}$, $\mathcal{E}_\Gamma(k)$	K_Γ is $\mathbb{C}((t^\Gamma))$ and $\mathcal{E}_\Gamma$ its entire series in this report.
unit roots $q_i = c_i(1 + \varepsilon_i)$; $u_n = \sum_i P_i(n)q_i^n$	$\lambda_j = \zeta_j(1 + \tau_j)$; $\mathcal{Y}(n) = \sum_j R_j(n)\lambda_j^n$	q is a dilation parameter; the unit-dilation notation of Convention 28.1 is kept.
envelope $W = A + M(S)$; $R_{i,\gamma}$	$\Sigma = C + M(S)$; $r_{j,\gamma}$	As in Lemma 28.4; R_j are the polynomial factors.
$C = \langle c_i \rangle$; period $M = \exp \operatorname{Tor}(C)$; class r	$\langle \zeta \rangle$; M_{tor} ; b	$\mathbb{C}$ is the complex field; M is source 18’s smaller period (Lemma 28.3, Remark 38.5); $M(S)$ is the positive monoid; r is a derivative order.
$\mathcal{A} = k[n, c_i^n]$; $R_0 = k[T, X_1, \dots, X_s]$	$\mathfrak{A}$; $k[U, X_1, \dots, X_m]$	$\mathcal{A}[f]$ is a differential expression of the theta-descent part; U is the polynomial variable of $k[U]$.
forward shift E	Sh	As in Section 28.2.
recurrence terms u_n	y_n	u is the Laurent variable of Section 24.5; y_n is otherwise local to the proof of Theorem 4.4.
vector $x \in K^d$, set X , $T \subseteq \Gamma$, $F \in k[X_1, \dots, X_h]$	$\mathbf{x} \in \mathbb{K}^{\bar{d}}$, $\mathcal{X}$, $\mathbf{T}$, $F \in k[Z_1, \dots, Z_h]$	x is an evaluation point, d a degree, T an indeterminate.
purely infinite ideal $\mathcal{J}$	Π	$\mathbf{Oz} = \mathbb{Z} \oplus \Pi$ as in Corollary 25.3; $\mathcal{J}_{\lambda,j}$ is a jet.
sequence b ; $A_b = \sum b_m \omega^{\omega-m}$; set $S \subseteq \mathbb{N}$	$\mathbf{b} = (\mathbf{b}_m)$; $\mathbf{w}_\mathbf{b}$; A	b is a residue class; S is a support alphabet.

Source 20	Here	Reason
Catalan root λ ; $C(s)$, C_k	λ_- ; $\text{Cat}(X)$, Cat_k	C_n are the Chebyshev polynomials of (23.1).
matrix A ; $\delta_j(n)$	$\mathbf{A}$; $\text{mv}_j(n)$	δ is a scale.
dilation group G ; convex hull H of $v(G)$	$\mathfrak{G}$; $\Delta_{\mathfrak{G}}$	G is the series of Corollary 17.1; convex subgroups are Δ here.
operator $\sum c_{\lambda,j,k} z^k D_z^j \sigma_{\lambda}$; $A_s(n)$, $B_j(n)$	$\sum \kappa_{\ell,r,i} z^i D_z^r \sigma_{\lambda_{\ell}}$; $c_j(n)$	As in (29.1) and (29.4).
witness $\eta \notin H$; η in examples	γ ; ϵ	$\eta = v(\tau)$ in Theorem 4.4.
Θ_Q ; characteristic p	Θ_p ; $\mathfrak{p}$	(8.1).
Theorems 4.1, 5.1, 5.5, 6.2, 7.1, 7.3, 10.1, 10.3, 10.4	see below	Theorems A–O of this report are different theorems.

The shipped audit file `20-recurrences-SOURCE_AUDIT.md` keeps the source’s notation.

Words with two meanings. A *unit* is a valuation-zero element (a “valuation unit” in source 20); it is neither an *order unit* (Definition 1.3) nor a unit of the ring $\mathbf{Oz}$, whose only units are ± 1 . *Coefficient* means a Hahn coefficient $[t^{\gamma}]x$, a Conway coefficient $[\omega^{\alpha}]x$ (the same number as $[t^{-\alpha}]x$ under (2.4)), a Taylor coefficient a_n , or a recurrence coefficient $c_j(n)$; a *coefficient observable* is a sequence $n \mapsto [t^{\gamma}]y_n$ for one fixed γ . *Compression* here means *Noetherian compression*, a finite subfamily of coefficient conditions equivalent to the whole family along an orbit (Theorem 38.8); it is unrelated to the bounded polynomial *jet* compression of Theorem 32.9, a statement of differential algebra. Two *periods* occur: source 18’s M suffices for valuations, while the hitting sets of Theorem 38.8 need M_{tor} , which M divides (Remark 38.5); the tempting reading “the valuation period is also the period of every coefficient test” is false. Finally $\deg_{\omega}$ is the greatest exponent of a Conway normal form, neither the degree of a polynomial nor a birthday.

Results printed once. Source 20 reproves several facts the report already contains. They are printed once and cited. Its definitions of strong summability, of $\mathcal{E}(K)$, of valuation units and of order units are Definitions 1.2 and 1.3 and Convention 28.1; its surreal translation $t^{\gamma} \mapsto \omega^{-\gamma}$ is (2.4), and its localization of finitely many surreal data to a set-sized Hahn field is the paragraph on omnific coefficients in Section 29.4. Its Lemma 3.1 (support lemma) is Lemma 2.1; its Lemma 3.2 and Corollary 3.4 are Lemma 21.1 and its proof; its Theorem 3.3, the classical Skolem–Mahler–Lech theorem, is imported there as here. Its Theorem 4.1 (a)–(b) is Lemma 28.4, proved by the same finite binomial expansion and support count; its Remark 4.2 is the remark after that lemma, and its Example 4.3, $v((1+t^{\epsilon})^n - 1 - nt^{\epsilon}) = 2\epsilon$ for $n \geq 2$ with $y_0 = y_1 = 0$, is the case $d = 2$ of Example 29.14. Its Theorem 5.1 is Theorem 28.5 with the admissible period M_{tor} in place of M (Remark 38.5), and its Corollary 5.2 is the last clause of that theorem together with its finitely many exceptional indices; its Example 5.3 is Example 4.6; its Lemma 5.4 is Lemma 28.9; its Theorem 5.5 is Theorem 28.10, which states the zero alternative for every index of the class where source 20 states it beyond a threshold; its Remark 5.6 is the third example of Remark 4.3. Its Lemma 9.1 is Lemma 29.1 with Corollary 29.2, and its Lemma 9.2 is Lemma 3.1 with Corollary 3.4, with the same closed exterior region $v(x) \leq -B$; its Theorem 10.1 is Theorem 29.4 for relative scale 0, and its Corollary 10.2 is Corollary 29.6. The entireness of its theta witness is Theorem 8.1, by the route of Remark 8.2, and its identities are (8.2) and (8.3). Its torsion example (Section 11.1) is the case $h = 2$ of the paragraph on root-of-unity quotients in Section 29.5 with Proposition 9.1, and its rank-two example (Section 11.2) is Example 29.16 with Example 8.4.

Where each numbered result went. Theorem 4.1 (c) is Theorem 38.2; the period paragraph (38.3) of its Section 3 is Lemma 38.4; Corollary 5.7 is Corollary 38.6; Definition 6.1, Theorem 6.2, Corollary 6.3, Example 6.4 and Remark 6.5 are Definition 38.7, Theorem 38.8, Corollary 38.9, Example 38.10 and Remark 38.11; Theorem 7.1, Example 7.2, Theorem 7.3, Corollary 7.4 and Remark 7.5 are Theorem 39.1, Example 39.2, Theorem 39.3, Corollary 39.4 and Remark 39.5; Corollary 8.1 is Corollary 39.7; Theorems 10.3 and 10.4 are Corollaries 39.8 and 39.9; Remark 10.5 is Remark 39.11; Section 11 is Section 39.5, with its positive-characteristic formula as Example 39.12; Questions 12.1, 12.2, 12.5 and 12.11 are recorded in the statuses of Questions 19.24, 19.21, 19.25 and 19.29, and Questions 12.3, 12.4, 12.6–12.10 and 12.12 are Questions 19.63–19.70, Question 12.7 re-scoped; the appendix and the audit file are merged into Sections 39.6, 39.7 and 18. Theorem O in the introduction collects Theorems 38.2, 38.8 and 39.3.

The merge's choices. The review of source 20 for this merge found its proofs correct; its worked examples were rechecked by hand and by its own suite. The merge adds five points, each marked. First, Remark 38.5 compares the two periods and shows that source 18's cannot replace source 20's for hitting sets. Second, Section 39.4 identifies source 20's Theorems 10.3 and 10.4 as special cases of Theorems 29.4 and 29.9, notes through Example 29.15 that the inclusion is strict, and shows in Remark 39.10 that finite generation cannot be dropped from the convex-hull form of the criterion. Third, Remark 39.6 credits the omnific-notations report, which already contains the element of the coding theorem. Fourth, Example 39.12 keeps source 20's exact positive-characteristic valuation formula, of which Section 29.5 records a subsequence. Fifth, the statuses of the related questions of source 18 are updated and source 20's Question 12.7 is re-scoped (Question 19.66). Source 20's theorems, proofs, examples and non-claims are otherwise printed as delivered, in the renamed notation.

38.2 One finitely generated algebra of coefficient observables

The support envelope and the coefficient polynomials of Lemma 28.4 give more than a valuation theorem: they say where every coefficient sequence lives.

Theorem 38.2 (One finitely generated observable algebra; source 20). *Let $\mathcal{Y}^{(1)}, \dots, \mathcal{Y}^{(\bar{d})}$ be unit-root exponential polynomials over $\mathbb{K}$, let $\lambda_1, \dots, \lambda_m$ list all their bases and $\zeta_1, \dots, \zeta_m$ the residues. There is one well-ordered $\Sigma \subseteq \Gamma$ containing $\text{supp } \mathcal{Y}^{(i)}(n)$ for every i and every $n \geq 0$. For every i and every $\gamma \in \Gamma$ there are $r_{j,\gamma}^{(i)} \in k[U]$ with*

$$[t^\gamma] \mathcal{Y}^{(i)}(n) = \sum_{j=1}^m \zeta_j^n r_{j,\gamma}^{(i)}(n) \quad (n \geq 0), \quad (38.1)$$

and so every coefficient observable $n \mapsto [t^\gamma] \mathcal{Y}^{(i)}(n)$ lies in the one finitely generated k -algebra

$$\mathfrak{A} = k[n, \zeta_1^n, \dots, \zeta_m^n] \subseteq k^{\mathbb{N}}, \quad (38.2)$$

the image of the evaluation homomorphism $\text{ev} : k[U, X_1, \dots, X_m] \rightarrow k^{\mathbb{N}}$, $U \mapsto (n)_n$, $X_j \mapsto (\zeta_j^n)_n$, sequences being multiplied pointwise.

Proof. Lemma 28.4, applied to each $\mathcal{Y}^{(i)}$ with its own bases (the others entering with zero polynomial factor), gives a well-ordered envelope $\Sigma^{(i)}$ and the identity (38.1) for $\gamma \in \Sigma^{(i)}$; for $\gamma \notin \Sigma^{(i)}$ take $r_{j,\gamma}^{(i)} = 0$. The finite union $\Sigma = \bigcup_i \Sigma^{(i)}$ is well ordered. The identity holds for every $n \geq 0$, small

n included: the polynomials $\binom{U}{i}$ vanish at the integers $n < i$, and characteristic zero makes them polynomials over k . Each summand $\zeta_j^n r_{j,\gamma}^{(i)}(n)$ is $\text{ev}(X_j r_{j,\gamma}^{(i)}(U))$. No sum over n is asserted to be strongly summable. $\square$

Remark 38.3 (Finitely generated, not finite-dimensional; source 20). The algebra $\mathfrak{A}$ has $m + 1$ generators, but the observables do not lie in a finite-dimensional subspace: already $[t^d](1+t)^n = \binom{n}{d}$ has degree d in n , so the degrees of the $r_{j,\gamma}$ are unbounded in γ , as recorded after Lemma 28.4. The theorem gives finitely many algebra generators for all coefficient sequences at once, not a finite basis; and it is the finite generation, not a degree bound, that the next subsection uses.

For $q = 1 + t^\epsilon$ with $\epsilon > 0$ the sequence $y_n = q^n - 1 - nt^\epsilon = \sum_{i=2}^n \binom{n}{i} t^{i\epsilon}$ vanishes for $n \leq 1$ and has valuation 2ϵ for $n \geq 2$; replacing q by its residue 1 would lose the whole sequence. This is source 20's Example 4.3, the case $d = 2$ of Example 29.14: the observables keep every principal-unit correction while each fixed coefficient is an ordinary exponential polynomial over k .

38.3 One period for all coefficient tests, and split recurrences

Source 20's Section 3 fixes the period

$$M_{\text{tor}} = \exp \text{Tor} \langle \zeta \rangle, \quad (38.3)$$

the exponent of the finite torsion subgroup of $\langle \zeta \rangle$, and observes that it serves for all polynomial degrees and all finite products of the residues at once.

Lemma 38.4 (One period for all monomials in the residues; source 20). *Let $\zeta^{\alpha_1}, \dots, \zeta^{\alpha_s}$ be monomials $\zeta^\alpha = \prod_j \zeta_j^{\alpha_j}$, $\alpha_i \in \mathbb{Z}^m$, let $\varrho_1, \dots, \varrho_s \in k[U]$ and $b \in \mathbb{N}$. The sequence*

$$l \mapsto \sum_{i=1}^s \varrho_i(M_{\text{tor}}l + b) (\zeta^{\alpha_i})^{M_{\text{tor}}l + b}$$

is either identically zero or has only finitely many zeros. The same M_{tor} serves for every such list, every choice of degrees and every b ; the finite zero sets need not have a common bound.

Proof. The bases in l are the elements $(\zeta^{\alpha_i})^{M_{\text{tor}}}$; combine equal ones. If $(\zeta^{\alpha_i}/\zeta^{\alpha_{i'}})^{M_{\text{tor}}}$ is a root of unity, then $\zeta^{\alpha_i}/\zeta^{\alpha_{i'}} \in \langle \zeta \rangle$ is a torsion element, its order divides M_{tor} , and the two bases are equal. So after combination the distinct bases have no root-of-unity quotient, and Lemma 21.1 gives the alternative. $\square$

Remark 38.5 (Two periods; merge). Source 18's period M of Lemma 28.3, the least common multiple of the orders of those quotients ζ_i/ζ_j that are roots of unity, divides M_{tor} , since each such quotient lies in the torsion subgroup of $\langle \zeta \rangle$. Both are admissible in Theorem 28.5, which with M_{tor} is source 20's Theorem 5.1. For hitting sets the smaller period does not suffice. For $m = 1$ and $\mathcal{Y}(n) = (-1)^n$ there is no quotient, so $M = 1$, and indeed $v(\mathcal{Y}(n)) = 0$ for every n ; but $M_{\text{tor}} = 2$, and the single condition $[t^0]x - 1 = 0$ holds exactly for even n . Valuations see only quotients of residues, because the least coefficient that is not an identity is a combination with those residues as bases; coefficient tests see arbitrary monomials in them, which is why Theorem 38.8 is stated with M_{tor} .

Corollary 38.6 (Recurrences with split characteristic polynomial; source 20). *Let (y_n) satisfy a constant-coefficient recurrence $y_{n+d} = a_{d-1}y_{n+d-1} + \dots + a_0y_n$ over $\mathbb{K}$ whose characteristic polynomial*

splits in $\mathbb{K}$. There are $M \geq 1$ and $N \geq 0$ such that on each class b modulo M either $y_n = 0$ for all $n \geq N$ in the class, or

$$v(y_n) = \beta_b + n\varpi_b \quad (n \geq N, n \equiv b \pmod{M}),$$

with $\beta_b \in \Gamma$ and ϖ_b the valuation of a nonzero characteristic root. The leading coefficient on each nonzero class is eventually an exponential polynomial over k in $(n - b)/M$, and M may be chosen from the residues of the normalized nonzero roots $t^{-v(\lambda)}\lambda$ alone. A zero characteristic root affects only a finite initial segment.

Proof. The Jordan decomposition of the companion matrix over $\mathbb{K}$ gives, for all n at least the multiplicity of the root 0, a representation $y_n = \sum_j R_j(n)\lambda_j^n$ over the distinct nonzero roots, with $\deg R_j$ below the multiplicity of λ_j ; nilpotent blocks vanish after finitely many steps. Apply Theorem 28.10 to the exponential polynomial defined by this representation for all n , which agrees with y_n on that tail, and take the larger threshold. The statement on leading coefficients is the one in Theorem 28.5, applied to the winning group of roots. $\square$

38.4 Noetherian compression of infinitely many coefficient tests

Definition 38.7 (Coefficient-algebraic conditions; source 20). A *coefficient-algebraic condition* on $\mathbf{x} = (x^{(1)}, \dots, x^{(\bar{d})}) \in \mathbb{K}^{\bar{d}}$ is an equation

$$F([t^{\gamma_1}]x^{(i_1)}, \dots, [t^{\gamma_h}]x^{(i_h)}) = 0, \quad F \in k[Z_1, \dots, Z_h],$$

in finitely many coefficient coordinates. A *coefficient-algebraic set* $\mathcal{X} \subseteq \mathbb{K}^{\bar{d}}$ is the common zero set of an arbitrary set-sized family $(F_a)_{a \in I}$ of such conditions.

The family may be infinite, but each member is an ordinary polynomial in finitely many ordinary coefficients. For a prescribed $T \subseteq \Gamma$ the support condition $\text{supp}(x) \subseteq T$ is the family $[t^\gamma]x = 0$, $\gamma \notin T$. The condition that an arbitrary real coefficient be an *integer* is not of this form.

Theorem 38.8 (Noetherian compression and one residual period; source 20). *Let $\mathbf{y}_n = (\mathcal{Y}^{(1)}(n), \dots, \mathcal{Y}^{(\bar{d})}(n))$ be a finite vector of unit-root exponential polynomials, let $\zeta_1, \dots, \zeta_m$ be the residues of all their bases, and let M_{tor} be as in (38.3). For every coefficient-algebraic set $\mathcal{X} \subseteq \mathbb{K}^{\bar{d}}$, defined by $(F_a)_{a \in I}$:*

- (a) *there is a finite $I_0 \subseteq I$ such that, for every $n \geq 0$, $\mathbf{y}_n \in \mathcal{X}$ if and only if F_a vanishes at $\mathbf{y}_n$ for all $a \in I_0$;*
- (b) *for each class b modulo M_{tor} , either $\mathbf{y}_{M_{\text{tor}}l+b} \in \mathcal{X}$ for every $l \geq 0$, or this holds for only finitely many l ;*
- (c) *consequently $\{n \geq 0 : \mathbf{y}_n \in \mathcal{X}\}$ is a finite union of full residue classes modulo M_{tor} and a finite set.*

The same M_{tor} serves for every $\mathcal{X}$; the finite subfamily I_0 and the exceptional set depend on $\mathcal{X}$.

Proof. By Theorem 38.2 every coordinate observable is ev of a polynomial in $k[U, X_1, \dots, X_m]$. For each $a \in I$ substitute such representatives for the finitely many coordinates in F_a ; this gives $\hat{F}_a \in k[U, X_1, \dots, X_m]$ whose value at $(n, \zeta_1^n, \dots, \zeta_m^n)$ is exactly F_a evaluated at $\mathbf{y}_n$. The ring $k[U, X_1, \dots, X_m]$ is Noetherian (Hilbert's basis theorem), so the ideal generated by all $\hat{F}_a$ is

generated by finitely many elements, each a finite combination of finitely many $\widehat{F}_a$; collect those indices into I_0 . A point (n, ζ^n) at which the $\widehat{F}_a$, $a \in I_0$, vanish kills the whole ideal, hence every $\widehat{F}_a$. This is (a), for every n and not only eventually.

Each $\widehat{F}_a(n, \zeta_1^n, \dots, \zeta_m^n)$ is a finite sum of terms $\varrho(n)(\zeta^\alpha)^n$ with $\alpha \in \mathbb{N}^m$. On a class modulo $\mathbf{M}_{\text{tor}}$ it is identically zero or has finitely many zeros by Lemma 38.4. If every condition vanishes identically on the class, every orbit point of the class lies in $\mathcal{X}$; otherwise a single condition that is not an identity there already leaves only finitely many hits. This is (b), and (c) follows by taking the finite union over classes. $\square$

Corollary 38.9 (Support and tail recurrence; source 20). *For a unit-root exponential polynomial $\mathcal{Y}$ and any $\mathbf{T} \subseteq \Gamma$, the set $\{n : \text{supp } \mathcal{Y}(n) \subseteq \mathbf{T}\}$ is a finite union of full classes modulo $\mathbf{M}_{\text{tor}}$ and a finite set; if it meets a class infinitely often, it contains the whole class. Along the orbit, the vanishing of all coefficients outside $\mathbf{T}$ is equivalent to the vanishing of finitely many of them.*

Proof. Apply Theorem 38.8 to the family $[t^\gamma]x = 0$, $\gamma \notin \mathbf{T}$; part (a) chooses a finite subset of exactly these equations. $\square$

Under (2.4), for a real-surreal unit-root sequence, the absence of an infinitesimal tail ($\text{supp} \subseteq \Gamma_{\leq 0}$), membership in the purely infinite support space ($\text{supp} \subseteq \Gamma_{< 0}$) and the vanishing of a prescribed truncation are instances of Corollary 38.9. Full membership in $\mathbf{Oz}$ also requires the constant coefficient to be an ordinary integer, an arithmetic condition on an exponential polynomial that is not a Zariski-closed coefficient condition and is not disposed of here (Question 19.64).

Example 38.10 (No common transient bound; source 20). For $\mathcal{Y}(n) = (1+t)^n$ the single condition $[t^d]x = 0$ holds exactly for $n = 0, \dots, d-1$. The period is $\mathbf{M}_{\text{tor}} = 1$ for every d , but the last exceptional index tends to infinity with d . A universal residual period does not give a uniform transient length for all coefficient tests on one orbit.

Remark 38.11 (Why infinitely many conditions leave no hidden gap; source 20). It would be invalid to take the union of infinitely many finite exceptional sets and call it finite, and the proof does not do so. On each class either every condition is an identity, or one condition that is not an identity already leaves finitely many possible hits; separately, Noetherianity supplies the finite subfamily. These are two independent safeguards against that inference, as in Remark 28.6 for valuations.

39 Surreal recurrences: leading scales, omnific coding, matrix powers and mixed operators

39.1 Leading Conway exponents

For a nonzero x in a workspace $K_\Gamma \subseteq \mathbf{No}[i]$ of (2.4), put $\deg_\omega(x) = -v(x)$: the greatest exponent of its Conway normal form, the real and imaginary normal forms being combined into one complex-coefficient normal form. For finitely many surreal or surcomplex numbers such a workspace exists: the group generated by the union of their supports is a set (Section 29.4). So every statement below about a fixed finite recurrence takes place in one set-sized Hahn field, and no proper-class product, sum or determinant is formed.

Theorem 39.1 (Surreal and surcomplex recurrence profiles; source 20). *Let (y_n) satisfy a constant-coefficient recurrence of finite order over $\mathbf{No}[i]$, with initial values in $\mathbf{No}[i]$. There are $\mathbf{M} \geq 1$ and*

$N \geq 0$ such that for each class b modulo $\mathbf{M}$ either $y_n = 0$ for all $n \geq N$ in the class, or

$$\deg_\omega(y_n) = n \mathbf{d}_b + \mathbf{e}_b \quad (n \geq N, n \equiv b \pmod{\mathbf{M}}), \quad (39.1)$$

with $\mathbf{d}_b, \mathbf{e}_b \in \mathbf{No}$, $\mathbf{d}_b$ the degree of a nonzero characteristic root. This applies in particular to every recurrence over $\mathbf{Oz}$ or $\mathbf{Oz}[i]$ with initial values in the same ring. If every nonzero characteristic root lies in $\mathbf{No}$, $\mathbf{M} = 2$ suffices; if every one is positive, $\mathbf{M} = 1$ suffices.

Proof. $\mathbf{No}[i]$ is algebraically closed, so the characteristic polynomial splits. Its finitely many roots, the coefficients of a representation $y_n = \sum_j R_j(n) \lambda_j^n$ on the tail after the nilpotent segment, and the recurrence data lie in one set-sized Hahn field $\mathbb{C}((t^\Gamma))$ with $\Gamma \leq \mathbf{No}$, as above. Corollary 38.6 applies there, and negating the valuation gives (39.1). For roots in $\mathbf{No}$ the normalized leading coefficients lie in $\mathbb{R}^\times$, whose only nontrivial torsion element is -1 ; so $\mathbf{M}_{\text{tor}}$ of their residues divides two, and it is an admissible period (Remark 38.5). Positive roots have positive leading coefficients, which generate a torsion-free subgroup of $\mathbb{R}_{>0}$. Closure of $\mathbf{Oz}$ and $\mathbf{Oz}[i]$ under ring operations keeps the terms in those rings. $\square$

Example 39.2 (An intercept larger than every ordinary time; source 20). Let $\lambda_- = \frac{1}{2}(\omega - \sqrt{\omega^2 - 4})$, the infinitesimal root of $X^2 - \omega X + 1$. With $t = \omega^{-1}$,

$$\lambda_- = t \text{Cat}(t^2), \quad \text{Cat}(X) = \sum_{k \geq 0} \text{Cat}_k X^k, \quad \text{Cat}_k = \frac{1}{k+1} \binom{2k}{k},$$

because $\text{Cat}(X) = 1 + X \text{Cat}(X)^2$. Put $y_n = \omega^\omega \lambda_-^n$. Then

$$y_{n+2} = \omega y_{n+1} - y_n, \quad y_n \in \Pi, \quad \deg_\omega(y_n) = \omega - n. \quad (39.2)$$

For $n \geq 1$, λ_-^n has positive integer coefficients, supported at the Conway exponents $-(n + 2k)$, $k \geq 0$, with $[t^{n+2k}] \lambda_-^n = \frac{n}{n+2k} \binom{n+2k}{k}$; multiplication by ω^ω moves them to $\omega - n - 2k > 0$, a reverse well-ordered support. Also $y_0 = \omega^\omega \in \Pi$, so all initial values and all later terms are omnific, and the recurrence follows from $\lambda_-^2 = \omega \lambda_- - 1$. For every ordinary n , y_n is positive and infinite, and $y_{n+1} < y_n$ since $0 < \lambda_- < 1$. Its degree decreases with slope -1 but never reaches zero at an ordinary time. This agrees with Theorem 39.1 ($\mathbf{d} = -1$, $\mathbf{e} = \omega$) and contradicts any argument that silently treats $\omega - n$ as though n were cofinal in the surreal exponents.

39.2 One fixed coefficient can encode any sequence

For $x \in \mathbf{No}[i]$ and $\alpha \in \mathbf{No}$ let $[\omega^\alpha]x$ be the coefficient of ω^α in the normal form of x .

Theorem 39.3 (A first-order omnific recurrence encodes any sequence; source 20). *For every real sequence $\mathbf{b} = (\mathbf{b}_m)_{m \geq 0}$ there is $\mathbf{w}_\mathbf{b} \in \Pi$ such that*

$$y_0 = \mathbf{w}_\mathbf{b}, \quad y_{n+1} = \omega y_n, \quad y_n \in \Pi, \quad [\omega^\omega] y_n = \mathbf{b}_n$$

for every ordinary $n \geq 0$. The same holds for complex sequences, with $\mathbf{w}_\mathbf{b}, y_n \in \Pi + i\Pi \subseteq \mathbf{Oz}[i]$. For every $A \subseteq \mathbb{N}$ one can therefore arrange $\{n : [\omega^\omega] y_n \neq 0\} = A$; when A is nonempty and $\mathbf{b}$ takes the values 0 and 1, every y_n is positive.

Proof. Put

$$\mathfrak{w}_{\mathbf{b}} = \sum_{m \geq 0} \mathbf{b}_m \omega^{\omega-m}, \quad (39.3)$$

zero terms omitted. The exponents $\omega - m$ strictly decrease and are all positive; every nonempty subset of them has a greatest element, the one of least index. So the support is a reverse well-ordered set, of order type at most ω in its decreasing enumeration, and (39.3) is a purely infinite omnific integer. Multiplication by the monomial ω^n translates exponents with no convolution: $y_n = \omega^n \mathfrak{w}_{\mathbf{b}} = \sum_m \mathbf{b}_m \omega^{\omega+n-m}$. All exponents stay positive, n and m being ordinary, and exactly the term $m = n$ has exponent ω . The complex case is the same coefficientwise, or follows by separating real and imaginary parts. For the last assertion take $\mathbf{b}$ the indicator of A ; the leading coefficient is then 1. $\square$

Corollary 39.4 (No coefficientwise Skolem–Mahler–Lech theorem for omnific recurrences; source 20). *It is false that all fixed-Conway-coefficient zero sets, or nonzero sets, of finite omnific recurrences are eventually periodic. Already the fixed multiplier ω , the fixed observed exponent ω and a varying purely infinite initial value realize every subset of $\mathbb{N}$.*

Remark 39.5 (No conflict with the universal quotient; source 20). The map $x \mapsto [\omega^\omega]x$ is additive on $\mathbf{Oz}$ but not multiplicative: it kills $\omega^{\omega/2}$ and sends its square to 1. So Theorem 39.3 does not contradict the theorem of the set-sized-quotient report that every additive and multiplicative map from $\mathbf{Oz}$ to a set-sized ring factors through the constant term (`osq:thm:universal` [49]); every y_n above has constant term zero. *Related (batch 34)*. The set-sized target in that theorem is needed: the same report constructs unital ring homomorphisms $\text{sp}_a : \mathbf{Oz} \rightarrow \mathbf{No}[i]$, $a > 0$, into the proper-class field $\mathbf{No}[i]$ with $\text{sp}_a(\omega^a) = -1$ (`osq:cr:thm:residually`), so they do not factor through the constant term. Neither result bears on Theorem 39.3, whose observable is additive and not multiplicative.

Remark 39.6 (The element is a known stream; merge). The number $\mathfrak{w}_{\mathbf{b}}$ is, with $a_n = \mathbf{b}_n$, the series `onot:eq:guardbasic` of the omnific-notations report [58], which already observes that $\sum_n a_n \omega^{\omega-n} = \omega^\omega \sum_n a_n (\omega^{-1})^n$ has positive exponents and a valid support for every real sequence, and that an algorithm for each a_n need not decide whether all a_n vanish. Source 20 does not cite it. The element is therefore not new; what source 20 adds is its reading as a recurrence observable, the orbit under multiplication by ω read at one fixed exponent, and Corollary 39.4. The independent omnific codes of Theorem 35.16 encode subsets of $\mathbb{N}$ differently, through algebraic independence, not a coefficient observation.

The exact distinction between the two sides (source 20). In Theorem 38.8 the roots are units, so the supports of all iterates lie in one well-ordered envelope, and a fixed coefficient receives only finitely many binomial contributions. In Theorem 39.3 multiplication by ω has nonzero valuation and translates the support, and the fixed observation reads a new stored coefficient at each step. This movement of support, not nonlinear arithmetic, permits universality. The leading exponent stays tame: if $\mathbf{b} \neq 0$ and m_0 is its first nonzero index, then $\deg_\omega(y_n) = \omega + n - m_0$, as Theorem 39.1 requires. The exact leading-scale theorem and unrestricted lower coefficients coexist in the simplest recurrence.

39.3 Determinantal valuation profiles

Corollary 39.7 (Determinantal valuation profiles; source 20). *Let $\mathbf{A} \in \mathrm{GL}_{\bar{d}}(\mathbf{No}[i])$, and for $1 \leq j \leq \bar{d}$ put*

$$\mathrm{mv}_j(n) = \min\{v(\det \mathbf{B}) : \mathbf{B} \text{ a } j \times j \text{ minor of } \mathbf{A}^n, \det \mathbf{B} \neq 0\}, \quad \mathrm{mv}_0(n) = 0,$$

in a workspace containing the entries of $\mathbf{A}$. There is one finite period such that every $\mathrm{mv}_j(n)$ and every difference $\mathrm{mv}_j(n) - \mathrm{mv}_{j-1}(n)$ is eventually affine on each residue class.

Proof. The $j \times j$ minors of $\mathbf{A}^n$ are the entries of $(\bigwedge^j \mathbf{A})^n$, and each entry satisfies the constant-coefficient recurrence given by the characteristic polynomial of the finite matrix $\bigwedge^j \mathbf{A}$. By Theorem 39.1, in its valuation form, each is eventually zero or of eventually affine valuation on every class of a finite period; refine the finitely many periods to one and take a common threshold. Invertibility of $\mathbf{A}^n$ gives a nonzero minor of each size at every n , so on each class some minor is active. The minimum of finitely many eventually affine functions is eventually one of them (Lemma 28.9), and differences of eventually affine functions are eventually affine. $\square$

Over a valuation ring these minima are the sums of the first j invariant-factor valuations of a finite matrix, after a common monomial rescaling if necessary; such a finite diagonal reduction is obtained directly by choosing an entry of least valuation, clearing its row and column by divisibility in the valuation ring, and repeating. The corollary gives exact eventual scale profiles of these finite-dimensional invariants. It asserts neither a real-valued Lyapunov limit nor eventual periodicity of all matrix coefficients (source 20).

39.4 Mixed operators: an independent derivation of part of Theorem J

Source 20's Sections 9–10 treat finite operators (29.1) whose multipliers lie in a torsion-free subgroup $\mathfrak{G} \leq \mathbb{K}^\times$. Distinct elements of such a group have no root-of-unity quotient, so the hypothesis of Theorems 29.4 and 29.9 holds, and source 18 needs less: faithfulness of the normal form needs only distinct multipliers (Corollary 29.2), and rigidity only a noncofinal relative scale. Source 20's reduction to a recurrence, its escape lemma with the closed region, and its unit-valuation theorem are printed once (Section 38.1); it found them independently of source 18, whose text it had not seen. Its remaining two theorems are special cases, printed here with its route.

Corollary 39.8 (Mixed rigidity below a proper convex scale; source 20). *Let $\mathfrak{G} \leq \mathbb{K}^\times$ be torsion-free and L a nonzero normal form (29.1) with distinct multipliers in $\mathfrak{G}$. If their valuations lie in a proper convex subgroup $\Delta < \Gamma$, then L has a uniform exterior obstruction and degree bound as in Theorem 29.4; in particular every strongly entire solution of $Lf = 0$ is a polynomial.*

Proof. The differences $v(\lambda_i) - v(\lambda_j)$ lie in Δ , and a nonzero element of a proper convex subgroup is not an order unit, since a convex subgroup containing an order unit is Γ . So the relative scale is noncofinal and Theorem 29.4 applies. Source 20's route avoids the relative scale: by Theorem 28.10 the recurrence coefficients have valuations $\beta_{j,b} + n\varpi_{j,b}$ with slopes in Δ ; for any $\gamma > 0$ outside Δ , convexity gives $n(\varpi_{0,b} - \varpi_{j,b}) < \gamma$ for every $n \geq 0$, whatever the sign of the difference, so the costs are bounded by $C + \gamma$ with C the largest intercept difference, and Corollary 3.4 concludes. The contradiction occurs in Γ itself; no coarsened summability is used. $\square$

The converse inclusion fails (merge): the multipliers of Example 29.15 have the cofinal common valuation μ , which lies in no proper convex subgroup, yet Theorem 29.4 excludes every nonpolynomial entire solution, because only relative valuations matter. Source 18's theorem is strictly stronger than source 20's Theorem 10.3.

Corollary 39.9 (Exact finite-group criterion; source 20). *Let $\mathfrak{G} \leq \mathbb{K}^\times$ be finitely generated and torsion-free, and let $\Delta_{\mathfrak{G}}$ be the convex subgroup of Γ generated by $v(\mathfrak{G})$. The following are equivalent.*

- (i) *Some nonpolynomial $f \in \mathcal{E}_\Gamma(k)$ is annihilated by a nonzero finite operator $\sum_{\lambda,r} p_{\lambda,r}(z) D_z^r \sigma_\lambda$ with multipliers $\lambda \in \mathfrak{G}$ and $p_{\lambda,r} \in \mathbb{K}[z]$.*
- (ii) $\Delta_{\mathfrak{G}} = \Gamma$.
- (iii) *Some $p \in \mathfrak{G}$ has positive order-unit valuation.*

When (iii) holds, Θ_p of (8.1) is a witness for (i): it satisfies (8.3), with the multipliers $1, p, p^2 \in \mathfrak{G}$ and no derivative.

Proof. (i) $\Leftrightarrow$ (iii): a finite list of distinct elements of $\mathfrak{G}$ has no root-of-unity quotient, so by Theorem 29.9, (i) holds if and only if $|v(\lambda_i/\lambda_j)|$ is an order unit for some $\lambda_i, \lambda_j \in \mathfrak{G}$; as $\lambda_i/\lambda_j \in \mathfrak{G}$ and $1 \in \mathfrak{G}$, this is (iii), after inverting if the valuation is negative. For the witness, Θ_p is a nonpolynomial element of $\mathcal{E}_\Gamma(k)$ (Lemma 29.8), and the operator $\sigma_1 - (1+z)\sigma_p + pz\sigma_{p^2}$ of (8.3) is a nonzero normal form (Corollary 29.2). (ii) $\Leftrightarrow$ (iii), source 20's argument: for generators $q_1, \dots, q_s$ of $\mathfrak{G}$ the convex subgroup generated by $v(\mathfrak{G})$ is $\{\gamma : |\gamma| \leq n\eta \text{ for some } n \geq 1\}$ with $\eta = \max_i |v(q_i)|$ (zero if every $v(q_i) = 0$). If it is the nonzero group Γ , then $\eta > 0$ is an order unit, and a generator or its inverse has valuation η ; conversely an order-unit valuation forces $\Delta_{\mathfrak{G}} = \Gamma$. Source 20 proves (i) $\Rightarrow$ (ii) by Corollary 39.8 and (iii) $\Rightarrow$ (i) by the direct estimate $h_{n+1} - h_n = n\beta + \delta > 0$ for $n > A$, where $h_n = \binom{n}{2}\beta + n\delta$, $\beta = v(p)$, $\delta = v(x)$ and $|\delta| \leq A\beta$, followed by the support count of Remark 8.2. $\square$

When Γ has an order unit, (ii) says that $v(\mathfrak{G})$ is not contained in the maximal proper convex subgroup Δ^{top} of Proposition 29.12; that proposition is the reformulation of the negative case for an arbitrary finite list.

Remark 39.10 (Finite generation cannot be dropped from (ii); merge). The equivalence of (i) and (iii) uses no finite generation. The equivalence with (ii) does. In Γ_∞ of (11.1), which has no order unit, the torsion-free group $\mathfrak{G}$ generated by the monomials t^{ω^j} , $j \geq 0$, has cofinal valuations, so $\Delta_{\mathfrak{G}} = \Gamma_\infty$; but no element of Γ_∞ is an order unit, and by Theorem 29.9, or already by Theorem E (b), no nonpolynomial entire series satisfies a nonzero mixed equation with multipliers in $\mathfrak{G}$.

Remark 39.11 (Which part source 20 counted as new; source 20). Source 20 recorded that the partial-theta witness and the one-parameter order-unit boundary were already in this report, and counted as its own contribution the general exponential-polynomial valuation theorem and its use for the mixed unit case and the finite multibase criterion; it also stressed that it does not infer a mixed assertion by combining two pure ones. The first sentence is accurate. The second was accurate against the text at its pin and is stale against the current one: source 18, placed before that pin and merged after it, proves the valuation theorem (Theorems 28.5 and 28.10) and the relative-scale criterion, of which source 20's Theorems 10.1, 10.3 and 10.4 are special cases. The two sources are independent derivations of the same day; no priority between them is asserted.

39.5 Boundary examples

Torsion is a genuine obstruction (source 20). Over $\mathbb{K} = k((t^{\mathbb{Q}}))$ let $p = t$, so that Θ_p is strongly entire, and put $f(z) = \Theta_p(z^2)$. Then f is nonpolynomial and strongly entire, and $f(-z) - f(z) = 0$; the operator $\sigma_{-1} - 1$ is not zero on all formal series, it merely kills every even series. So excluding only the formally zero identity $\sigma_{-1}^2 - 1$ would not repair the theorems; a theory with torsion must track residue-degree sectors and the operator's action on them. This is the case $h = 2$ of the paragraph on root-of-unity quotients in Section 29.5 and of Proposition 9.1. Residual torsion, by contrast, is allowed throughout: $-(1 + t)$ has infinite order and is covered by Corollary 29.6.

Two scales inside one rank-two workspace (source 20). Let $\Gamma = \mathbb{Q}e_0 \oplus \mathbb{Q}e_1$ with $0 < e_0 \ll e_1$. Dilation by t^{e_0} , together with any finite family of unit dilations generating a torsion-free group, stays in the proper convex subgroup $\mathbb{Q}e_0$, and by Corollary 39.8 supports no nonpolynomial entire solution of a nonzero mixed equation; dilation by t^{e_1} admits the partial theta witness, e_1 being an order unit of the same group. So valuation rank alone is not the classification parameter, as in Examples 8.4 and 29.16.

Example 39.12 (Characteristic zero cannot be omitted; source 20). In $\mathbb{F}_p((t))$, for every $n \geq 1$,

$$v((1 + t)^n - 1) = \mathfrak{p}^{v_{\mathfrak{p}}(n)}.$$

Indeed, writing $n = \mathfrak{p}^a m'$ with $\mathfrak{p} \nmid m'$, $(1 + t)^n = (1 + t^{\mathfrak{p}^a})^{m'}$, whose first nonconstant term is $m't^{\mathfrak{p}^a} \neq 0$. These valuations have no eventually periodic finite range although $1 + t$ is a unit; the binomial-polynomial argument over k and the characteristic-zero zero-set theorem are both essential. The paragraph on positive characteristic in Section 29.5 records the subsequence $n = \mathfrak{p}^a$; the formula gives every n (merge's comparison). Positive-characteristic zero sets need a different theory [33].

Leading exponents are not birthdays or signs (source 20). $\deg_{\omega}$ records a leading normal-form exponent, not the birthday of the surreal, and nothing here bears on Gonshor's product-birthday problem. An eventually affine valuation does not give an eventually periodic sign: ordinary real recurrence sequences have valuation zero whenever nonzero, while their signs can oscillate with complex phases. No sign classification is used or claimed.

Effectivity and the classical Skolem problem (source 20). The proofs give the finite exceptional sets existentially, and an arbitrary coefficient exponential polynomial invokes the Skolem–Mahler–Lech theorem. No algorithm locating all exceptional zeros is claimed; a general one would solve the ordinary Skolem problem, by applying it to constant Hahn series. Noetherian compression likewise gives the existence of a finite subfamily, not an algorithm finding it from an arbitrary infinitely presented family. The coding theorem is deliberately non-effective when $\mathbf{b}$ is noncomputable: its initial value already contains the sequence. It gives no finite effective representation of every subset of $\mathbb{N}$, and it does not show that coefficient extraction is definable in the pure ring language of $\mathbf{Oz}$.

39.6 Predecessors, repository comparison and priority

Literature. Source 20 imports Conway normal forms and the Hahn-field viewpoint as established foundations [8]; the Hahn–Neumann support lemma, credited historically to Neumann [30] (not

audited as a historical paper) and consulted in Bagayoko, Krapp, Kuhlmann, Panazzolo and Serra [56] (version 2), for positive-support summability; and the classical Skolem–Mahler–Lech theorem [28, 29], whose Bell account it read and inspected visually. Fuchs and Heintze [31] obtain effective valuation-growth bounds for recurrences over one-variable function fields under nondegeneracy assumptions; source 20’s exact eventual-affine statement allows arbitrary Hahn supports and ordered value groups but sacrifices a general effective exceptional-index bound, and does not improve their constants. Krapp, Kuhlmann and Serra [57] study generalized recurrence relations indexed by exponent groups and their relation to fraction fields: there the recurrence index is an exponent, here it is an ordinary natural number and only the values lie in a Hahn field; the setups are distinct and no theorem of theirs is imported. Derksen [33] is context for positive characteristic. Formal summability is used in the support-sensitive sense of the Hahn literature, not as numerical convergence. The Wikipedia article on surreal numbers was read for orientation only. For this merge the bibliographic data were checked; their content was not rechecked, and no proof here rests on Fuchs–Heintze, Krapp–Kuhlmann–Serra or Derksen.

Repository comparison at the pin. Source 20 read, at `efc5446`, the repository README and catalogue, this report’s `article.tex` in the line ranges listed in its audit (1330–1500, the unit-orbit examples and differential recurrences; 2020–2150, the sharp dilation criterion, partial theta and torsion; 2150–2228, Lemma 10.1, Theorem 10.2 and the limitation after it), and the README of the set-sized-quotient report [49], whose universal quotient theorem it uses only to explain the absence of a contradiction (Remark 39.5), without auditing that report. It did not read the placed audit file of source 18, nor the omnific-notations report (Remark 39.6). The comparison was targeted, not a proof-by-proof audit of all reports, and the absence of a matching theorem in it is not a proof of novelty. Its statements about this report were accurate at the pin, as recorded in Section 38.1.

Priority. Source 20 separates four groups. *Imported foundations:* Hahn multiplication and positive-support summability, Conway normal forms, real closedness of **No**, the Skolem–Mahler–Lech theorem and Hilbert’s basis theorem. *Reproved repository mechanisms:* finite-step recurrence escape, the noncofinal-scale obstruction and the partial-theta sharpness construction. *Candidate-original package:* the uniform coefficient algebra, coefficient-constraint compression, arbitrary-rank exact valuation profiles, their mixed unit and multibase application, and the juxtaposed omnific coefficient-universality theorem. *Not claimed:* independent priority certification, formal verification, effective Skolem decision procedures, arbitrary nonlinear mixed rigidity, and a classification of full omnific membership sets for unrestricted recurrences. Against the current text, the valuation profiles and the mixed application are source 18’s, found independently (Remark 39.11), and the coding element is the omnific-notations report’s (Remark 39.6); what remains proposed by source 20 is the coefficient algebra, Noetherian compression with its universal period, the recurrence statements over **No**[i], **Oz** and **Oz**[i] with the decreasing omnific example, the coding reading with Corollary 39.4, and the determinantal profiles. “Candidate-original” means developed and proved at manuscript level with no matching result found in a targeted comparison; it does not establish priority.

39.7 What the recurrence part does not claim

Every limitation stated by source 20, in its manuscript, README, audit file or verification record, is kept.

- (V1) Source 20 is an AI-assisted, unrefereed research draft with manuscript arguments, not an independently certified breakthrough. No independent referee report and no Lean formalization accompany it; no Lean build was run, no declaration was added, and the repository’s axiom audit does not cover this part. Independent correctness and worldwide priority are not certified.
- (V2) The Skolem–Mahler–Lech theorem, the Hahn–Neumann support lemma and Hilbert’s basis theorem are imported. No effective bound for exceptional zeros and no algorithm locating them is given; such an algorithm would solve the ordinary Skolem problem, which is not claimed solved.
- (V3) Noetherian compression gives the existence of a finite subfamily only, not an algorithm finding it from an infinitely presented family.
- (V4) The degrees of the coefficient polynomials $r_{j,\gamma}$ are unbounded, and a universal residual period gives no common transient bound for all coefficient tests on one orbit (Example 38.10).
- (V5) No nonlinear mixed rigidity is asserted; source 20’s nonlinear question is re-scoped here (Question 19.66).
- (V6) No classification of full omnific membership sets: the integer constant-term condition is not a coefficient-algebraic condition and is not handled (Question 19.64).
- (V7) The coding theorem is storage of a sequence in an infinite initial value, not an algorithm that generates information; it is non-effective for noncomputable sequences, gives no finite effective representation of subsets of $\mathbb{N}$, and shows neither that coefficient extraction is definable in the pure ring language of **Oz** nor any interpretation of second-order arithmetic.
- (V8) Entireness is relative to one set-sized Hahn field; a series entire in one workspace need not be entire after enlargement, and nothing concerns simultaneous entireness on all of **No**[i]. Statements over **No**[i] concern each finite recurrence localized to a set-sized Hahn field.
- (V9) D_z fixes the coefficients and is not the Berarducci–Mantova derivation; derivations of the coefficient field are not treated (Question 19.67).
- (V10) Actual torsion among the dilation parameters and positive characteristic are excluded, with counterexamples (Section 39.5).
- (V11) Nothing is claimed about real-valued Lyapunov limits or eventual periodicity of the coefficients of matrix powers.
- (V12) $\deg_\omega$ is neither a birthday nor a sign; Gonshor’s product-birthday problem is not addressed, and no sign classification is claimed.
- (V13) The effective constants of Fuchs and Heintze are not improved, and their setting differs; the setup of Krapp, Kuhlmann and Serra is distinct.
- (V14) Neumann’s paper is credited historically, not audited; the set-sized-quotient report was not audited, and its theorem is used only to explain the absence of a contradiction.
- (V15) The 4,605 finite checks concern finite identities and examples. They prove neither the support lemma, the Skolem–Mahler–Lech theorem, the infinite-support minimum argument nor the proper-class localization, and a PDF build concerns document integrity, not truth. Their number measures no proof coverage: 1,189 of them compare elementary integer or lexicographic facts that hold by construction (Section 18).

- (V16) The repository comparison at `efc5446` was targeted, not exhaustive; its description of this report is stale in the respects recorded in Section 38.1, and its search was not a literature priority audit.
- (V17) *[merge]* Remarks 38.5, 39.6 and 39.10, the identification of source 20's Theorems 10.3 and 10.4 as special cases of Theorem J, the comparison in Example 39.12, and the re-scoping of Question 19.66 are the merge's; no historical priority is claimed for them.

40 Conclusion

Strong entireness over a surreal Hahn workspace imposes a global scale requirement that a finite linear differential equation cannot meet unless its solution terminates. A dilation equation can meet that requirement, but exactly when its valuation reaches every scale through integer multiples. The decisive distinction is therefore not between Archimedean and non-Archimedean fields, nor between rank one and higher rank; it is between *a cofinal dilation scale* and *a scale trapped below another one*.

The obstruction is finite and robust. A linear recurrence forces a later nonzero coefficient with a controlled valuation, and a large enough argument turns an infinite chain of such coefficients into a nonincreasing sequence of leading exponents. Descending supports and infinite reuse of one exponent are two distinct ways in which the evaluation fails, and keeping them separate is what makes one exclusion boundary closed and the other open. The mechanism works with arbitrary Hahn coefficient tails, arbitrarily high valuation rank, and sparse coefficient sequences, without replacing strong summability by topological convergence and without any hidden completeness assumption.

The countable-rank example makes the separation concrete. The series $\sum_{n \geq 0} \omega^{-\omega^n} z^n$ is evaluable at every point of its full workspace, yet no single nontorsion dilation in that workspace supplies a finite linear equation for it. That gap between existence and finite linear describability is the main mathematical contribution proposed here.

The nonlinear part pushes the same global requirement past linear equations by a different mechanism. A finite algebraic relation between a function and its first derivative is incompatible with nonpolynomial strong entireness over any nontrivial characteristic-zero Hahn workspace: at one large admissible scale the whole series leaves a finite initial polynomial, whose top coefficient is controlled by a corner polynomial that is automatically nonzero in order one. In higher order an exact cancellation of that corner is a genuine obstruction to that method, and a fixed nonlinear equation can then have polynomial solutions of unbounded degree.

The coefficient-field part removes the order restriction when the value group has no order unit, by a third mechanism. A finite algebraic relation, linearized along its own solution, confines every Taylor coefficient to one finitely generated field, and such a field cannot supply independent valuations at cofinally many Archimedean levels; a nonpolynomial entire series needs them. So every nonpolynomial strongly entire series is then differentially transcendental over the full field of constants, jointly with its dilates, and every differentially algebraic quotient of entire series is rational, while one explicit workspace still carries continuum many entire series with jointly independent jets. With an order unit the mixed statement fails.

The theta-descent part shows that the purely differential statement fails too, so the order unit is the exact boundary. A bilateral theta series, descended along $z = u + u^{-1}$ by the Chebyshev identity, becomes an ordinary power series that is strongly entire exactly when its scale is an order

unit, and the classical elliptic identity gives it an explicit algebraic differential equation of order three. Its residue corner vanishes, as it must. The shift of the theta series by $u \mapsto p^2u$ then excludes every equation of lower order, by three elementary difference-field arguments.

The order-three-threshold part shows that no other series does better. At the infinitely many scales where a nonpolynomial entire series has a nonmonomial initial polynomial, a fixed equation of order at most two reduces to one homogeneous polynomial over the residue field, and logarithmic Euler derivatives confine both endpoints of those initial polynomials to a finite set, even when the leading-monomial test vanishes identically. So order three is the least possible for every value group with an order unit. At order three one further integer, the gap next to an endpoint, survives, and the theta equation uses it with gap one. Which series attain order three is the open part of Question 19.1.

The unit-dilation part closes the linear theory. Mixing derivatives with dilations adds no new way to escape: the coefficients of a mixed recurrence are exponential polynomials with Hahn coefficients, whose valuations are eventually affine on finitely many progressions, and so for every value group a nonpolynomial entire solution of some mixed linear equation exists exactly when two multipliers are separated at the top Archimedean scale. In particular a unit dilation, however close to a root of unity its residue, adds nothing to derivatives.

The Newton-rigidity part adds a first result at order three. When the exterior form of an equation has a nonzero first polar along the twisted cubic of monomial jets, the two ends of a far initial polynomial demand next exponents on opposite sides, and every cubic form vanishing on that curve has such a polar; so order three with jet degree at most three is also rigid. In degree four the test fails, the exact solutions of two boundary equations are doubling binomials, and a sparse entire series shows that initial equations valid at every scale need not lift to an identity.

The theta-hierarchy and single-scale parts measure what order three leaves room for. Theta descents at scales with independent reciprocals have distinct quadratic Gauss growth, so over $\mathbb{R}((t^{\mathbb{R}}))$ they are algebraically independent, and polynomially weighted sums of them reach every large differential order; yet every one is differentially algebraic. At the other extreme a single scale already carries lacunary series with rational Taylor coefficients that satisfy no algebraic differential equation at all, because a maximal-degree band of heights cannot be cancelled.

The polynomial-composition part shows how much of the order-unit phenomenon belongs to degree one. A polynomial argument of degree $d \geq 2$ multiplies the scale by d , so a superlinearly growing Gauss profile cannot be balanced by arguments of lower degree unless their composition weight reaches the top one; then every strongly entire solution is a polynomial, for every value group, with a degree bound that keeps the exact indicial resonances. The finite coefficient locus that remains can be as complicated as any affine hypersurface over $\mathbb{Q}$, and its omnific points are undecidable in general.

The recurrence part reads the ordinary recurrences behind all these proofs one coefficient at a time. When the characteristic roots are units, every iterate lies in one well-ordered support envelope and each fixed coefficient is an ordinary exponential polynomial over the residue field; all of them lie in one finitely generated algebra, so any set-sized family of coefficient conditions reduces along the orbit to finitely many, with hitting sets periodic modulo one period up to a finite set. When a root moves the support, as ω does, one fixed coefficient can store any real sequence, although the leading exponent stays eventually affine.

A Notation

Symbol	Meaning
Γ	A fixed nonzero set-sized ordered abelian group; divisible only where stated.
K_Γ	The complex Hahn field $\mathbb{C}((t^\Gamma))$.
$v(a), \text{lc}(a), \text{res}(a)$	Least support exponent, its coefficient, and the residue when $v(a) = 0$.
$\mathcal{E}_\Gamma, \text{Dom}_{\text{str}}(f)$	Series strongly evaluable at every point of K_Γ ; the exact set where a series is strongly evaluable.
D_z	Formal derivative in z , zero on every element of K_Γ .
σ_q	Argument dilation $f(z) \mapsto f(qz)$.
$M(S)$	The additive monoid generated by a well-ordered $S \subseteq \Gamma_{>0}$.
Δ_β	The smallest convex subgroup containing $\beta > 0$, (2.1).
B, B^+, B_{rec}	Exclusion thresholds in Γ ; B_{rec} divides by the jump length and needs Γ divisible.
$\overline{B}_{\text{rec}}^\flat, \overline{B}_{\text{rec}}$	Their coarsened analogues in $\Gamma/\Delta_{ v(q) }$.
μ, β	Usually a positive dilation valuation; an order unit only when stated.
$(X)_r$	The ordinary falling factorial polynomial of degree r .
Θ_p	The partial theta series (8.1).
Γ_∞	The increasing-rank direct sum (11.1).
<i>Nonlinear part, Sections 14–17</i>	
$k, \mathbb{K}, \mathcal{E}_\Gamma(k)$	Any characteristic-zero field; $k((t^\Gamma))$; its strongly entire series (Convention 14.1).
$\mathcal{S}_m, \mathcal{S}_m^+, \rho$	Series with strongly summable coefficient family; those with coefficients of nonnegative valuation; coefficientwise residue.
$\gamma_\delta, \phi_\delta, N_\delta$	Gauss value, initial polynomial and top active degree at the scale $t^{-\delta}$, (14.1).
$d_P, w_P, \mathcal{C}_P, \beta_P$	Corner data of a differential polynomial, (14.5).
χ_P, I_P	Residue and full corner polynomials, (14.6).
κ_P	The rational degree cutoff (15.11).
δ_0	A solution-dependent exclusion bound, (15.8).
ϑ_q, wt	Positive-weight Euler operator and weight.
<i>Coefficient-field part, Sections 20–22</i>	
$\text{Frac } \mathcal{E}_\Gamma(k)$	The fraction field of $\mathcal{E}_\Gamma(k)$ in $\mathbb{K}((z))$ (“meromorphic”).
$\mathcal{J}_{\lambda,j}f$	$(D_z^j f)(\lambda z)$, differentiation before dilation, (20.1).
$\text{rrk}, \text{trdeg}, \text{dtrdeg}$	Rational rank; transcendence degree; differential transcendence degree.
$\text{cvr}(f)$	Coefficient-value rank (20.4).
$\Delta, v_\Delta, k_\Delta, \text{red}_\Delta$	A convex subgroup, the coarsened valuation $v + \Delta$, its residue field and residue map.
$\Pi_{ij}, \theta, \xi_{ij}, \Xi$	Evaluated partials, offset, leading coefficients and linearization multiplier, (21.2)–(21.3).
$L, k_0 \subseteq K_0$	A coefficient subfield; a generic pair of characteristic-zero fields.
$\mathcal{F}_\alpha, \text{code}, R_\tau$	The branch family (22.3), its binary code, the rank-one family.

Symbol	Meaning
<i>Theta-descent part, Sections 23–25</i>	
μ, p, Δ_μ	A positive scale (an order unit only where stated), $p = t^\mu$, the convex subgroup (2.1).
$C_n, \mathcal{T}_p$	Monic Chebyshev polynomials (23.1); the descended series (1.4).
$\Psi_p, u, \tilde{z}$	The bilateral theta series (23.8); its variable; $u = u^{-1}$.
$\mathcal{O}_{\Delta_\mu}, x_m, \mathcal{P}_p$	The strong domain of $\mathcal{T}_p$; its zeros (23.13); $\prod_m (1 - p^{2m})$.
$\mathcal{Q}_r, \mathfrak{c}, g_2, g_3, \Phi$	Lambert series in the nome p^2 and the constants (24.1)–(24.3).
$\mathcal{A}[f], \mathcal{H}[f], \mathcal{K}[f], \mathcal{B}[f]$	Differential expressions (24.4)–(24.6).
$\mathcal{U}_\mu, f^\natural, \vartheta_u, \mathcal{S}$	The Laurent ring, the substitution $z = u + u^{-1}$, uD_u , and $u \mapsto p^2u$.
$\psi_1, \psi_2, \psi_3; \mathcal{L}, \mathcal{A}_1$	$\vartheta_u \Psi / \Psi$ and its two ϑ_u -derivatives; $D_z f / f$ and $D_z \mathcal{A}[f]$.
<i>Order-three-threshold part, Sections 26–27</i>	
$\Gamma_{\mathbb{Q}}, \mathbb{K}_{\mathbb{Q}}$	The divisible hull $\Gamma \otimes_{\mathbb{Z}} \mathbb{Q}$ and $k((t^{\Gamma_{\mathbb{Q}}}))$.
$\text{Act}_\delta, \underline{N}_\delta$	Active indices at the scale δ and the least of them (26.1); a <i>break</i> is a scale with $\underline{N}_\delta < N_\delta$.
$\vartheta_X; \mathfrak{u}_\phi, \mathfrak{v}_\phi, \mathfrak{w}_\phi$	XD_X ; $\vartheta_X \phi / \phi$ and its two ϑ_X -derivatives.
$\mathfrak{h}_P$	The exterior form of P (26.3).
$\mathfrak{q}, i_0, \mathfrak{r}, \text{Sp}(\mathfrak{h})$	Dehomogenization, the power of V removed, the remainder, the finite endpoint spectrum (Lemma 26.8).
$\Upsilon_{\mathfrak{h}}, \mathfrak{g}^\pm, \delta_n$	Endpoint-gap polynomial (27.2); upper and lower gaps; the break scale $v(a_{n+1}) - v(a_n)$.

Symbol	Meaning
<i>Unit-dilation part, Sections 28–29</i>	
$\lambda_1, \dots, \lambda_m; \zeta, \tau$	Multipliers of a mixed operator; residue and tail of a unit (28.1).
$\mathcal{Y}(n), R_j; \Sigma, r_{j,\gamma}$	An exponential polynomial $\sum_j R_j(n)\lambda_j^n$; its support envelope and coefficient polynomials (28.2).
$M, b; \beta_b, \varpi_b$	Period and residue class; eventual intercept and slope (28.5).
Sh	The forward shift of sequences.
$\varkappa_{\ell,r,i}, d_L, s_0, c_j(n), R_{j,\ell}$	Coefficients of (29.1), their z -degree bound, the least active shift, the recurrence coefficients (29.4) and shift polynomials (29.5).
$h, \bar{e}, \bar{n}$	A polynomial right side, a bound on its degree, an output degree.
$\text{rs}(\lambda_1, \dots, \lambda_m)$	The relative scale (Definition 29.3).
Δ^{top}	The maximal proper convex subgroup when Γ has an order unit (Proposition 29.12).
G_μ	$\sum_n t^{n^2\mu} z^n = \Theta_{p^2}(pz)$, $p = t^\mu$; G_1 over $\mathbb{C}((t^\mathbb{Q}))$ is the series G of Corollary 17.1.
<i>Newton-rigidity part, Sections 30–31</i>	
$\mathbf{m}_r(T); \chi_{\mathfrak{h}}$	The monomial jet curve $(1, T, \dots, T^r)$; $\mathfrak{h}(\mathbf{m}_r(T))$, so that $\chi_{\mathfrak{h}_P} = \chi_P$.
$\mathfrak{s}, \mathfrak{h}_1$	$Y_0 Y_2 - Y_1^2$ and the cofactor in $\mathfrak{h} = \mathfrak{s}^e \mathfrak{h}_1$ (Remark 30.2).
$\text{Pol}_{\mathfrak{h}}, a_{\mathfrak{h}}, b_{\mathfrak{h}}$	The first polar (30.1) and its factorization (30.2).
$H_3, Q_4, B(m, n)$	The quartic of order three, the quadratic of order four and its pair kernel (31.1)–(31.4).
A_1, A_2, n, u_f	The auxiliary quadratics and common-root parameter of Theorem 31.1; $\vartheta f/f$.
F_{sp}	The sparse series (31.5).
$\mathbb{L}; \mathfrak{M}, \mathfrak{M}_1, \mathfrak{m}$	A field of external constants; the fields and minimal polynomial of Theorem 30.16.

Symbol	Meaning
<i>Theta-hierarchy part, Sections 32–33</i>	
$\mathcal{T}_{t^\mu}$	(1.4) with $p = t^\mu$, $\mu > 0$ real.
$\text{dord}(f)$	The least r with $f, \dots, D_z^r f$ dependent (Definition 32.13).
$\mathfrak{K}_0 \subseteq \mathfrak{K}_1, \partial, \text{d}, \mathfrak{s}$	Differential fields, their derivation, the universal derivation, an element with $\partial \mathfrak{s} = 1$.
$P_i, c_{ij}, W, U_{ir}, \varepsilon_i$	Compression weights, grid coefficients, the wedge, jet parameters, formal parameters (Theorem 32.9).
$\kappa_{\mathbf{e}}, \kappa'_{\mathbf{e}}, \delta^{(0)}, \delta^{\text{cert}}$	Growth constants and thresholds of Proposition 32.5.
$d_N, \mathfrak{L}_N, u_N$	The compositum and its exact-order element (Theorem 32.19).
$\mathfrak{V}; \mathcal{A}_{\text{DA}}, \mathcal{D}_{\text{DA}}$	A $\mathbb{Q}$ -independent set of positive reals; the differentially algebraic entire series and their fraction field.
$x, \gamma^{\mathcal{U}}$	$-\partial_u \psi_1 - 2\mathcal{Q}_1(p^2)$; the radius-one Gauss valuation on $\mathcal{U}_\mu$.
<i>Single-scale part, Sections 34–35</i>	
$\mathbf{b}_n; \mathbf{N}_A, \mathcal{G}_A$	Rapid heights; the series of Definition 34.2.
∂_p	$p \text{d}/\text{d}p$.
$\mathbf{v}_n, A_{\mathbf{v}}, A_{\text{fin}}, \mathbf{c}, \mathbf{c}_0, \mathbf{V}, d_{\text{pat}}, \ell_j$	Patterns, pattern classes, finite corrections, pattern matrix and rank, kernel basis (Section 34.4).
$\bar{d}, \bar{h}, \mathbf{T}, \mathbf{b}_{\mathbf{T}}; \mathfrak{P}, W_{\mathfrak{P}}$	Top degree, p -degree, target multiset, band start; multihomogeneous part and its polarization.
$\mathfrak{x}_A, \partial_{\text{BM}}$	$\sum_{n \in A} \omega^{-\mathbf{b}_n}$; the Berarducci–Mantova derivation.
$\mathcal{B}_{\Gamma}^{\text{bd}}(k), \mathcal{F}_{\Gamma}^{\text{bd}}(k)$	Series with support bounded above; its fraction field.
$\mathcal{O}_{\langle 1 \rangle}$	$\{0\} \cup \{x \in \mathbf{No}[i]^\times : v(x) \geq -m\}$.
$M_{\Omega}, \mathfrak{z}_A, \mathfrak{l}_A(a)$	ω^{Ω} ; the omnific codes; the floor profile.
<i>Polynomial-composition part, Sections 36–37</i>	
$\text{lc}_z(b)$	Leading coefficient in z of $0 \neq b \in \mathbb{K}[z]$.
$\mathcal{E}^{(\delta)}$	Series with $v(b_n) - n\delta \rightarrow +\infty$ cofinally.
$w_L, \beta_L, I_L, \chi_L$	Shift, corner value, exact and residue indicial polynomials of an operator (36.3).
δ_{ref}	Reference scale of Lemma 36.4.
$\bar{s}, \bar{m}, c_*, \mathbf{G}, L_{\ell}, L'_{\nu}, P_{\nu}, e_{\nu}$	Data of (36.5).
$\text{cw}(\mathbf{a})$	Composition weight (36.4).
$w_{\text{top}}, \beta_{\text{top}}, w_{\mathbf{a}}, \beta_{\mathbf{a}}, \gamma_{\text{top}}, \gamma_{\mathbf{a}}$	Shifts, corner values and Gauss bounds in the proof of Theorem N.
$\text{Rsn}, B_{\text{cmp}}, B_{\text{lin}}, B_{\text{mat}}$	Resonances and degree bounds (Section 36.5, Theorem 36.15).
$\mathfrak{o}, \mathfrak{O}_{\mathfrak{o}, \Gamma}, \Pi_{k, \Gamma}, \text{ct}, \mathfrak{R}$	$\mathbb{Z}$ or $\mathbb{Z}[i]$; omnific coefficient ring; its negative part; constant term; a coefficient ring of Theorem 37.2.
$\text{Pr}_j, \text{Ann}_{d, N}, \text{wd}(p)$	Euler projectors, annihilator and weighted degree of the encoder.

Symbol	Meaning
<i>Recurrence part, Sections 38–39</i>	
y_n	A recurrence sequence with values in $\mathbb{K}$ or $\mathbf{No}[i]$.
$\langle \zeta \rangle, \mathbf{M}_{\text{tor}}$	The group generated by the residues of unit bases; the exponent of its torsion subgroup (38.3) (divisible by $\mathbf{M}$).
$\mathfrak{A}, \text{ev}$	The observable algebra $k[n, \zeta_1^n, \dots, \zeta_m^n]$ of (38.2); the evaluation map onto it from $k[U, X_1, \dots, X_m]$.
$\mathcal{X}, F_a, \mathbf{T}$	A coefficient-algebraic set, its defining conditions, a prescribed support set (Definition 38.7).
$\deg_\omega; \mathbf{d}_b, \mathbf{e}_b$	$-v$, the greatest Conway exponent; slope and intercept in (39.1).
$\Pi; [\omega^\alpha]x$	The purely infinite ideal, $\mathbf{Oz} = \mathbb{Z} \oplus \Pi$; a Conway coefficient.
$\mathbf{b}, \mathbf{w}_b$	A real sequence and its omnific code (39.3).
λ_-, Cat_k	The infinitesimal root of $X^2 - \omega X + 1$; Catalan numbers.
$\mathbf{A}, \text{mv}_j(n)$	An invertible matrix; the least valuation of a nonzero $j \times j$ minor of $\mathbf{A}^n$.
$\mathfrak{G}, \Delta_{\mathfrak{G}}$	A torsion-free group of dilation parameters; the convex subgroup generated by $v(\mathfrak{G})$.

B Proof-dependency and hypothesis audit

The logical dependencies are short. Differential rigidity uses the integer-evaluation lemma, the coefficient conversion, and the escape chain with its exclusion criterion. Dilation necessity uses the geometric-orbit lemma or the unit-orbit theorem, the dilation conversion, and the same criterion. Dilation sufficiency uses only the partial theta identity and the support certificate. The mixed extension uses the bivariate orbit lemma. The several-variable theorem uses countable polynomial avoidance, finite-block regrouping and the one-variable theorem. The infinite-rank application uses a direct strong-entireness proof and the completed classification. The nonlinear part uses none of these linear arguments: first-order rigidity uses the Hahn-family algebra (with the support calculus of Section 2), the large-active-degree lemma, the corner identity and the order-one nondegeneracy of the corner; the higher-order criterion replaces the last step by the hypothesis $\chi_P \neq 0$; the Euler theorem uses the same algebra in several variables with finite bounded-weight sets. The coefficient-field part uses neither the escape chain nor the corner: its all-order theorem uses the linearized coefficient recurrence, the valuation-rank inequality, the principal convex bound and the exterior obstruction (or the coarsened reduction), with the cofinal criterion of Proposition 20.2; its mixed statements add the imported Skolem–Mahler–Lech theorem. The theta-descent part uses none of these for its existence direction: the exact domain uses the quadratic support certificate and the nonincreasing-exponent obstruction; the equation of order three uses the classical triple product and elliptic identity, proved as a formal identity over $\mathbb{Q}$ and transferred by ring homomorphisms; the lower bound uses the Laurent ring, the substitution homomorphism and three difference-field obstructions. Its necessity direction is Theorem E (a). The order-three-threshold part uses the all-scale criterion, a cofinal extension to the divisible hull, the Hahn-family algebra and its residue homomorphism, local finiteness of active indices and late breaks, the exterior reduction and the finite endpoint lemma; its least-order statement adds the witness $\mathcal{T}_p$ and Theorem E (a), and its theta results the exterior form of the cleared equation and a rational Riccati reduction. The unit-dilation part uses the support calculus, the imported Skolem–Mahler–Lech theorem through Lemma 21.1, a common support envelope and period, the least coefficient sequence that is not an identity, the eventual order of finitely many affine functions, the escape chain with its exclusion criterion, and, for existence, the partial theta identity and domain. The Newton-rigidity part uses the machinery

of the order-three-threshold part up to the exterior form, then the first polar with its double factor, the eventual side of integral roots through a $\mathbb{Q}$ -linear functional and the secant Jacobian; its exact classifications use only formal power series over a field of characteristic zero. The theta-hierarchy part uses the coefficient values of $\mathcal{T}_p$, Gauss multiplicativity, a unique dominant summand, the Kähler criterion with a finite grid, the separant argument and, for the continuum, a Hamel basis; its route to order three uses the Laurent ring and the shift with a Gauss valuation. The single-scale part uses height separation, polarization, Zariski density and linear algebra over $k(p, z)$, constant extension, and, for its realizations, the positive support monoid, the Berarducci–Mantova derivation and the normal-form definition of $\mathbf{Oz}$. The polynomial-composition part uses the entire-support criterion, Gauss multiplicativity and large active degrees, then the exact pullback and operator profiles, integer convexity and superlinear escape; its degree bounds use exact leading terms, its module theorem the Smith normal form and the constant-term retraction, and its encoding Euler projectors. The recurrence part uses the support envelope of the unit-dilation part and finite binomial expansions for its observable algebra, Hilbert’s basis theorem and a torsion period with the imported Skolem–Mahler–Lech theorem for compression, the algebraic closedness of $\mathbf{No}[i]$, a Jordan form and the periodic-affine valuations for its surreal and matrix profiles, and only normal-form support arithmetic for the coding theorem.

No unproved ring-theoretic conclusion from another report in this collection is used. The real-valued coarsening of [12, Lemma 9.1] is cited and not reproved, and no proof here depends on it. Higman’s theorem is an external standard ingredient; alternatively one may import the classical Hahn–Neumann support lemma directly. The surreal interpretation imports the normal-form field embedding, while every substantive function statement is already proved in the abstract Hahn field. The coefficient-field part imports one further external theorem without proof, the Skolem–Mahler–Lech theorem in the form of Bell’s Theorem 1.1 [27], and only its statements with several dilations use it. The theta-descent part imports the classical Jacobi triple product [50] and the constancy of an elliptic function without poles; it proves the elliptic identity it needs. The unit-dilation part imports the Skolem–Mahler–Lech theorem [28, 29] for exponential polynomials with at least two bases, and proves the finite-word lemma that Lemma 2.1 imports from Higman (Lemma 28.2). The Newton-rigidity part imports nothing beyond the Jacobian criterion for algebraic independence in characteristic zero, which it proves in one line. The theta-hierarchy part imports the theta product and the Tate cubic identity, as source 13 did, and uses the axiom of choice for a Hamel basis. The single-scale part imports the normalized Berarducci–Mantova derivation [8], the normal-form identification and definition of $\mathbf{Oz}$, and cites Grönwall’s single-series criterion without using it. The polynomial-composition part imports the Davis–Putnam–Robinson–Matiyasevich theorem [44] and Sun’s eleven-unknown theorem [45] for its two undecidability corollaries, and cites the complex entire-Mahler lemma [42, Lemma 6] without using it. The recurrence part imports the Skolem–Mahler–Lech theorem [28, 29], Hilbert’s basis theorem, the Hahn–Neumann support lemma [30, 56] and the algebraic closedness of $\mathbf{No}[i]$; it cites the set-sized-quotient theorem [49] only to explain the absence of a contradiction.

The potentially dangerous hypotheses, collected for review:

- Supports and the value group are sets; all operators are finite sums. Power-series degrees and derivative orders are ordinary nonnegative integers; dilation powers may be negative, as may the linearization offset θ . Formal Laurent series have only finitely many negative powers.
- $\Gamma \neq 0$, set-sized, ordered abelian. *Divisibility is not assumed* except in Corollary 3.5, the refined threshold (5.4), Theorem 7.4 (b), and Example 13.1, each of which states it, and in Lemma 26.4,

which is applied, in the order-three-threshold and Newton-rigidity parts, only after the cofinal extension of Lemma 26.2 to the divisible hull.

- Series are ordinary $\mathbb{N}$ -indexed power series in one variable, except in the several-variable Sections 6 and 16.3, and in the formal Laurent fraction field $\text{Frac } \mathcal{E}_\Gamma(k)$. The main workspace rigidity equations have rational-function coefficients in the variable, with arbitrary Hahn scalar coefficients. The relative coefficient-field theorem (Theorem 21.2 (a)) also allows coefficients in $k_0((z))$; that generality is used in the joint-independence criterion.
- The residue field is characteristic zero and trivially valued. Algebraic closedness is not used. It is $\mathbb{C}$ in Sections 1–13 and an arbitrary characteristic-zero field in the nonlinear, coefficient-field, theta-descent, order-three-threshold, unit-dilation, Newton-rigidity, single-scale, polynomial-composition and recurrence parts (with $\mathbb{R}$ or $\mathbb{C}$ for their omnific rings), and $\mathbb{R}$ or $\mathbb{C}$ in the theta-hierarchy part, which also fixes $\Gamma = \mathbb{R}$; the latter generality is not transferred to the statements of the linear theorems (Convention 14.1), although the unit-dilation part contains Theorem A, the negative direction of Theorem B and Theorem 10.2 over every such field (Remark 29.5). Characteristic zero is essential (Example 17.3).
- The single parameter q in the main dilation classification is nonzero and nontorsion. In the mixed-jet statements every λ is nonzero, and distinct parameters have non-root-of-unity quotients; an individual parameter may be 1; the same holds for the multipliers of the unit-dilation part. Section 9 and Section 29.5 explain the failure when this quotient condition is dropped.
- The derivative is d/dz , not a scalar surreal derivation.
- Entireness is always relative to one fixed workspace, never to $\mathbf{No}[z]$. Strong summability requires both well-ordering and local finiteness.
- The conclusions of Sections 1–13 concern *linear* equations. For every $\Gamma \neq 0$, nonlinear differential equations are treated only in order one, in higher order under $\chi_P \neq 0$, and for positive-weight Euler relations (the nonlinear part), in order two (the order-three-threshold part), and in order three with total jet degree at most three (the Newton-rigidity part); for explicit lacunary series all orders are excluded at one order-unit scale (the single-scale part), while for $\Gamma = \mathbb{R}$ minimal orders are unbounded (the theta-hierarchy part). Linear mixed equations in derivatives and finitely many dilations are treated for every Γ (the unit-dilation part), and so are equations in which one polynomial argument of degree at least two strictly dominates the composition weights (the polynomial-composition part). The recurrence part concerns ordinary constant-coefficient recurrences indexed by $\mathbb{N}$, not equations for functions, apart from its re-derived mixed statements. Nonlinear differential transcendence in all orders, and algebraic independence of derivatives at dilations with pairwise quotients that are not roots of unity, are proved only when Γ has no order unit, or for a single series with infinite coefficient-value rank (the coefficient-field part). No universal all-order rigidity theorem is claimed for groups with an order unit, and none can hold: for them the theta-descent part gives a nonpolynomial entire series with an algebraic differential equation of order three, and three is then the least order. No general mixed-jet independence theorem is claimed when a quotient is a root of unity.

Two boundary behaviours deserve separate emphasis because they point in opposite directions. The most delicate equality is in Corollary 3.5: weakly decreasing leading exponents already obstruct strong summability, so the excluded region is closed, and Proposition 5.3 shows the bound is attained. The most delicate strictness is in Theorem 7.4: equal *coarse* values need not obstruct summability in Γ , so that excluded region must be open, and Theorem 8.1 with Section 8.5 shows why.

Deliberate exclusions, recorded so that no reader supplies them by analogy: finite-order q is excluded from the main classification and has genuine periodic-support counterexamples; linear mixed operators at unit valuation are classified for every Γ (Corollary 29.6), but multiplier sets with a root-of-unity quotient, prescribed operators at a cofinal quotient and simultaneous systems are not. For groups with an order unit, the universal nonlinear rigidity results cover orders one and two, order three in total jet degree at most three, the nondegenerate higher-order class and positive-weight Euler relations; the infinite-coefficient-value-rank criterion gives additional all-order results for individual series, including Theorem 21.20, while Theorem H shows that universal differential algebraic rigidity fails for them, with least order three by Theorem I; which series have minimal order three is not decided. The nonlinear part keeps the non-claims of Section 19.3, the no-order-unit results keep those of Section 22.7, the theta-descent part keeps those of Section 25.6, the order-three-threshold part those of Section 27.6, the unit-dilation part those of Section 29.8, the Newton-rigidity part those of Section 31.5, the theta-hierarchy part those of Section 33.5, the single-scale part those of Section 35.7, and the polynomial-composition part those of Section 37.6 (critical weights, several top products and matrix differential pullbacks are not treated), and the recurrence part those of Section 39.7 (no effective exceptional zeros, no classification of omnific membership, and a coding theorem that stores rather than computes); no complete algorithm for arbitrary surreal comparisons or cofinality decisions is claimed; the finite checks prove no infinite theorem. The full classifications and the nonlinear, coefficient-field, theta-descent, order-three-threshold, unit-dilation, Newton-rigidity, theta-hierarchy, single-scale, polynomial-composition and recurrence theorem packages remain without Lean verification or independent refereeing. The formalization ledger records the proved support and escape lemmas, exponential and partial-theta domains, polynomial-orbit valuations, constant-point avoidance, inward stability and torsion covariance, including the partial-theta directions of the dilation classification.

C Reproducing the document and the finite checks

The directory contains this standalone L^AT_EX source, its compiled PDF, a README, twenty source audit documents at the top level, twenty-two files under `code/` (the thirteen verification programs of the sources merged here, the build scripts of sources 10, 11, 13, 17, 19 and 20, and the `Makefiles` of sources 12, 15 and 16) and thirty-two under `data/` (the thirteen recorded outputs, the source build, layout and visual reports, and the requirements files of sources 11, 13, 14, 15, 17 and 19), all belonging to the thirteen sources merged into this text. No source document from the inspected repository is redistributed, and no source manuscript is shipped. The bibliography is embedded: no BibTeX run, external graphics, shell escape, or repository checkout is required, and a conventional TeX Live or MiKTeX installation with the named packages suffices.

```
latexmk -pdf -interaction=nonstopmode -halt-on-error article.tex
python code/08-finite-recurrences-verify.py --output rerun-08.txt
python code/09-entire-hahn-holonomic-verification.py --output rerun-09.txt
python code/10-nonlinear-rigidity-verify.py --output rerun-10.json
pip install -r data/11-coarsening-differential-rigidity-requirements.txt
python code/11-coarsening-differential-rigidity-verify.py --output rerun-11.json
python code/12-coefficient-field-rigidity-verify.py --output rerun-12.json
pip install -r data/13-theta-descent-requirements.txt
python code/13-theta-descent-verify.py --precision 128 --output rerun-13.json
pip install -r data/14-order-three-threshold-requirements.txt
```

```
python code/14-order-three-threshold-verify.py
python code/18-unit-dilation-verify_finite.py --output rerun-18.json
pip install -r data/19-newton-rigidity-requirements.txt
python code/19-newton-rigidity-verify.py
python code/16-theta-hierarchy-verify.py --precision 128 --output rerun-16.json
pip install -r data/17-single-scale-requirements.txt
python code/17-single-scale-verify.py --output rerun-17.json
pip install -r data/15-polynomial-composition-requirements.txt
python code/15-polynomial-composition-verify.py --output rerun-15.json
python code/20-recurrences-verify.py > rerun-20.txt
```

The `--output` argument is required by the scripts of sources 08, 10, 12 and 13, so that a rerun need not overwrite the record distributed with the report, and optional for those of sources 09 and 11; the scripts of sources 10 and 12 refuse an existing output path (source 12's unless `--force` is given), source 13's overwrites one, and source 11's writes `verification.json` in the current directory when no path is given. Source 14's takes no arguments and always writes `data/verification.json` next to its own `code/` directory: here a new file beside the delivered record, overwritten by the next run, so it should be run on a copy. Source 19's script behaves the same way and writes the same file, `data/verification.json`; source 18's writes only the path given by `--output`, overwriting it, and otherwise only prints. Source 16's writes `verification.json` in the current directory unless `--output` is given, and source 17's writes `verification_results.json` beside itself in `code/`; both overwrite, so pass new paths. Source 15's, like source 16's, writes `verification.json` in the current directory unless `--output` is given, overwriting it. Source 20's takes no arguments and only prints its record, so redirect it to a new file. All use exact integer and rational arithmetic, those of sources 11, 13, 14, 15, 17 and 19 through SymPy, and exit with a nonzero status if a check fails. The build script `code/10-nonlinear-rigidity-build.py` was written for source 10's own manuscript: run here it would typeset this report, leave auxiliary files in the directory and write `data/build_report.json`. The shipped `data/10-nonlinear-rigidity-build_report.json` and `data/10-nonlinear-rigidity-layout_audit.json` describe the 23-page source manuscript, not this report; use `latexmk` as above, on a copy. Likewise `code/11-coarsening-differential-rigidity-build.py` looks for an `article.tex` in `code/` and fails there, and `code/12-coefficient-field-rigidity-Makefile` names the delivery paths `article.tex` and `verify.py` and would rebuild `article.pdf` in place; the shipped `data/11-coarsening-differential-rigidity-build_audit.json` and `data/12-coefficient-field-rigidity-build_report.json` describe the 20- and 23-page source manuscripts, whose PDFs and sources, hashed in the first, are not shipped. Source 13's `code/13-theta-descent-build.py` typesets an `article.tex` in its own directory `code/`, where there is none, and its `--verify` option would call a `verify.py` there; run here it creates `code/build/`, writes a failed first-pass log into it and exits with an error. In its original delivery layout it writes only into `build/`. The shipped `data/13-theta-descent-build_report.json` describes the 21-page source manuscript, whose source and PDF, hashed there, are not shipped. Likewise `data/14-order-three-threshold-build_report.json` describes source 14's 26-page A4 manuscript, whose source and PDF, hashed there, are not shipped; source 14 shipped no build script. Source 18 shipped none either and no build record. Source 19's `code/19-newton-rigidity-build.py` looks for an `article.tex` in `code/` and fails there, and its `--verify` option would call a `code/verify.py` relative to `code/`, which does not exist; `data/19-newton-rigidity-build_audit.json` describes the 24-page source manuscript, whose source and PDF, hashed there, are not shipped. Source 16's `code/16-theta-hierarchy-Makefile` and source 17's `code/17-single-scale-build.py` were written for their delivery layouts (Section 18); do not use them here, and the build and layout

records of both describe their 25- and 24-page manuscripts, which are not shipped. The same holds for source 15's `code/15-polynomial-composition-Makefile`, which would rebuild an `article.pdf` in the current directory and whose `clean` target deletes auxiliary files there, and for `data/15-polynomial-composition-BUILD_REPORT.json`, which describes its 24-page manuscript. Source 20's `code/20-recurrences-build.sh` runs `pdflatex` on an `article.tex` in its own directory and then overwrites `verification.txt` there by redirection; here it stops at the first `pdflatex` call and leaves `texput.log` in `code/` (Section 18), so do not run it.

References

- [1] R. P. Stanley, *Differentiably finite power series*, European Journal of Combinatorics **1** (1980), no. 2, 175–188. Author-hosted copy: <https://math.mit.edu/~rstan/pubs/pubfiles/45.pdf>. Theorem 1.5 (printed page 176) gives the D-finite/P-recursive equivalence. The finite coefficient conversion is reproved in this report for its stated characteristic-zero Hahn coefficient field.
- [2] J.-P. Ramis, *About the growth of entire functions solutions of linear algebraic q -difference equations*, Annales de la Faculté des Sciences de Toulouse, Mathématiques, Série 6, **1** (1992), no. 1, 53–94. https://www.numdam.org/item/AFST_1992_6_1_1_53_0/.
- [3] S. Garoufalidis, *The degree of a q -holonomic sequence is a quadratic quasi-polynomial*, Electronic Journal of Combinatorics **18** (2011), no. 2, Paper P4. Inspected manuscript: arXiv:1005.4580v4. <https://arxiv.org/abs/1005.4580>.
- [4] L. Di Vizio, *An ultrametric version of the Maillet–Malgrange theorem for nonlinear q -difference equations*, Proceedings of the American Mathematical Society **136** (2008), no. 8, 2803–2814. Inspected manuscript: arXiv:0709.2464v2. <https://arxiv.org/abs/0709.2464>.
- [5] G. E. Andrews and S. O. Warnaar, *The product of partial theta functions*. Author-hosted manuscript; the definition on its first page is the series used here, up to a sign convention. <https://people.smp.uq.edu.au/OleWarnaar/pubs/Partial-thetas.pdf>.
- [6] G. Higman, *Ordering by divisibility in abstract algebras*, Proceedings of the London Mathematical Society (3) **2** (1952), 326–336. <https://doi.org/10.1112/plms/s3-2.1.326>.
- [7] J. Müller and A. Strohmaier, *The theory of Hahn meromorphic functions, a holomorphic Fredholm theorem and its applications*, Analysis & PDE **7** (2014), no. 3, 745–770. <https://arxiv.org/abs/1205.0236>. <https://doi.org/10.2140/apde.2014.7.745>.
- [8] A. Berarducci and V. Mantova, *Surreal numbers, derivations and transseries*, Journal of the European Mathematical Society **20** (2018), 339–390. <https://arxiv.org/abs/1503.00315>. <https://doi.org/10.4171/JEMS/769>.
- [9] O. Bournez and Q. Guilmant, *Surreal fields stable under exponential, logarithmic, derivative and anti-derivative functions*, arXiv:2211.08396 (2022). <https://arxiv.org/abs/2211.08396>.
- [10] E. Kaplan, L. S. Krapp and M. Serra, *Decomposing the automorphism group of the surreal numbers*, arXiv:2509.22374v3. Used for the Hahn-field and surreal normal-form conventions, not for its automorphism results. <https://arxiv.org/html/2509.22374v3>.
- [11] VladimirReshetnikov/Surreal project, *Repository documentation and research-report catalogue*, snapshot a3124af79f66b8b9c196d76b4cbc5ac3938907c4, consulted 22 September 2026. <https://github.com/VladimirReshetnikov/Surreal>. The inspection used the root README and docs/README.md at this commit; these are project documentation, not independent published validation.
- [12] VladimirReshetnikov/Surreal project, *Cofinality, factorization and scalar extension for entire Hahn functions at arbitrary rank*, docs/surcomplex/entire-functions-at-arbitrary-rank/, same pinned snapshot. <https://github.com/VladimirReshetnikov/Surreal/tree/a3124af79f66b8b9c196d76b4cbc5ac3938907c4/docs/surcomplex/entire-functions-at-arbitrary-rank>. Its Lemma 9.1, labelled

- `ent:lem:coarsening`, is the real-valued coarsening cited in Remark 2.6; its Theorem 9.3 is the Bézout statement referred to in Section 1.5. Source 11 inspected its README at 465a54b for the polynomial intersection obstruction for noncofinal extensions and for coarsened support methods.
- [13] VladimirReshetnikov/Surreal project, *Differential algebra and differential equations over the surreal and surcomplex numbers*, docs/surcomplex/differential-equations/, same pinned snapshot. <https://github.com/VladimirReshetnikov/Surreal/tree/a3124af79f66b8b9c196d76b4cbc5ac3938907c4/docs/surcomplex/differential-equations>. Cited to distinguish the scalar surreal derivation from D_z .
 - [14] VladimirReshetnikov/Surreal project, *Rank-one non-Archimedean analytic geometry inside the surcomplex numbers*, docs/surcomplex/rank-one-berkovich/, same pinned snapshot. <https://github.com/VladimirReshetnikov/Surreal/tree/a3124af79f66b8b9c196d76b4cbc5ac3938907c4/docs/surcomplex/rank-one-berkovich>. Cited for the pre-existing quadratic-exponent rank-one example.
 - [15] VladimirReshetnikov/Surreal project, *Foundations for Surreal and Surcomplex Mathematics*, docs/foundations-and-computation/foundations/. Its Example `found:ex:internal` is the series $\sum t^{n^2} z^n$ over $\mathbb{C}((t^{\mathbb{Q}}))$; cited by this merge, not by source 10.
 - [16] VladimirReshetnikov/Surreal project, *Tail-spans and differential transcendence in surreal and surcomplex Hahn fields*, docs/surreal/tail-spans-and-differential-transcendence/, at source 10's pin 048b72cf7cbfc8ab246e4f73788c10460cb3f6e0. <https://github.com/VladimirReshetnikov/Surreal/tree/048b72cf7cbfc8ab246e4f73788c10460cb3f6e0/docs/surreal/tail-spans-and-differential-transcendence>. Its README was inspected by source 10 for function-class and derivative boundaries, and again by sources 11 and 12 at their pin 465a54b, for its continuum-sized families and its all-order results.
 - [17] V. Mantova and M. Matusinski, *Surreal numbers with derivation, Hardy fields and transseries: a survey*, arXiv:1608.03413, inspected version 2, Section 2 on normal forms (Section 2.2 by source 11). <https://arxiv.org/html/1608.03413v2>. Published in *Contemporary Mathematics* **697** (2017), 265–290; source 13 consulted the publication records and the preprint for normal forms, and source 14 the preprint for normal-form and Hahn-field background.
 - [18] W. K. Hayman, *The local growth of power series: a survey of the Wiman–Valiron method*, Canadian Mathematical Bulletin **17** (1974), no. 3, 317–358. <https://doi.org/10.4153/CMB-1974-064-0>. Publisher metadata and abstract consulted by source 10.
 - [19] J. Cano and P. Fortuny Ayuso, *Power series solutions of non-linear q -difference equations and the Newton–Puiseux polygon*, Qualitative Theory of Dynamical Systems **21** (2022), article 123. <https://doi.org/10.1007/s12346-022-00656-0>. Introduction and Section 2 (initial and indicial polynomials) consulted by source 10.
 - [20] P.-C. Hu and Y.-Z. Luan, *Non-Archimedean meromorphic solutions of functional equations*, arXiv:1311.5291v1 (2013). <https://arxiv.org/abs/1311.5291>. Field hypotheses and Theorems 1.1, 1.3 and 1.5 consulted by source 10; introduction and field hypotheses by sources 11 and 12, as context only.

- [21] P.-C. Hu and C.-C. Yang, *Meromorphic Functions over Non-Archimedean Fields*, Mathematics and Its Applications, Springer, Dordrecht, 2000. <https://doi.org/10.1007/978-94-015-9415-8>. Publisher metadata and description only; the book was not audited.
- [22] J. Giansiracusa and S. Mereta, *A general framework for tropical differential equations*, manuscripta mathematica **173** (2024), 1273–1304. <https://doi.org/10.1007/s00229-023-01492-5>.
- [23] A. Hurwitz, *Sur le développement des fonctions satisfaisant à une équation différentielle algébrique*, Annales scientifiques de l'École Normale Supérieure (3) **6** (1889), 327–332. <https://doi.org/10.24033/asens.326>. Bibliographic page and the attribution in [24] consulted by source 11.
- [24] V. Laohakosol, K. Kongsakorn and P. Ubolsri, *Coefficients of differentially algebraic series*, Journal of the Australian Mathematical Society, Series A **48** (1990), 402–412. <https://doi.org/10.1017/S1446788700029955>. Lemma 3 consulted by source 11.
- [25] C. Krattenthaler and T. Rivoal, *Arithmetic properties of the Taylor coefficients of differentially algebraic power series*, arXiv:2502.09259v1 (2025). <https://arxiv.org/abs/2502.09259>. Abstract consulted by source 11, as context only.
- [26] R. Ait El Manssour, A.-L. Sattelberger and B. Teguia Tabuguia, *D-algebraic functions*, arXiv:2301.02512v3. <https://arxiv.org/abs/2301.02512>. Framework consulted by source 11 for context.
- [27] J. P. Bell, *A generalised Skolem–Mahler–Lech theorem for affine varieties*, arXiv:math/0501309v2 (2007). <https://arxiv.org/abs/math/0501309>. Its Theorem 1.1, the classical theorem over a field of characteristic zero, is the form imported by source 12.
- [28] C. Lech, *A note on recurring series*, Arkiv för Matematik **2** (1953), 417–421. <https://doi.org/10.1007/BF02590997>. Publisher metadata checked by source 12; the proof was not reconstructed. Cited by source 18 as the primary source of the characteristic-zero theorem, with its metadata verified, and by source 20 for the same theorem.
- [29] J. P. Bell, *The Skolem–Mahler–Lech theorem*, Documenta Mathematica, Extra Volume Mahler Selecta (2019), 173–178. <https://ems.press/content/book-chapter-files/27398>. Read by source 18, including the first two pages, for the precise characteristic-zero statement and its exponential-polynomial formulation; a different text from [27]. Source 20 read and visually inspected the same PDF.
- [30] B. H. Neumann, *On ordered division rings*, Transactions of the American Mathematical Society **66** (1949), 202–252. <https://doi.org/10.1090/S0002-9947-1949-0032593-5>. Credited by source 18 for the support calculus, whose needed lemma it proves, and by source 20 as the historical source of the support lemma, not audited as a historical paper.
- [31] C. Fuchs and S. Heintze, *On the growth of linear recurrences in function fields*, Bulletin of the Australian Mathematical Society **104** (2021), 11–20. <https://doi.org/10.1017/S0004972720001094>. Author version: <https://arxiv.org/abs/2006.11074>. Related work consulted by source 18 through the author version and publisher metadata; not an input. Cited by source 20 as related work whose effective constants it does not improve.

- [32] H. Gonshor, *An Introduction to the Theory of Surreal Numbers*, London Mathematical Society Lecture Note Series 110, Cambridge University Press, 1986.
<https://doi.org/10.1017/CBO9780511629143>. Normal-form background for source 18; publisher metadata checked, the monograph not reread.
- [33] H. Derksen, *A Skolem–Mahler–Lech theorem in positive characteristic and finite automata*, *Inventiones Mathematicae* **168** (2007), 175–224. Author version:
<https://arxiv.org/abs/math/0510583>. Background for source 18’s positive-characteristic direction, and for source 20’s; not an input to any proof.
- [34] G. Pogudin, *The primitive element theorem for differential fields with zero derivation on the ground field*, arXiv:1409.3847. <https://arxiv.org/abs/1409.3847>. Consulted by source 16 as context for jet compression.
- [35] G. Pogudin, *A primitive element theorem for fields with commuting derivations and automorphisms*, arXiv:1812.11375v3. <https://arxiv.org/abs/1812.11375>. Consulted by source 16 (HTML text) as context; not a substitute for Theorem 32.9.
- [36] A. Ostrowski, *Über Dirichletsche Reihen und algebraische Differentialgleichungen*, *Mathematische Zeitschrift* **8** (1920), 241–298. Read by source 17 in the English translation by E. C. Hansen, Y. Stone and J. Wolfson, arXiv:2211.02088v1 (2022), <https://arxiv.org/abs/2211.02088>; the passage before Theorem 11 in Section 5 attributes the gap criterion to Grönwall, whose paper was not inspected.
- [37] S. L’Innocente and V. Mantova, *A factorisation theory for generalised power series and omnific integers*, arXiv:1710.07304v5 (2024). <https://arxiv.org/abs/1710.07304>. Used by source 17 for the normal-form definition of **Oz** only.
- [38] VladimirReshetnikov/Surreal project, *Bounded-Support Hahn Arithmetic: cofinal descent, optimal support thresholds and maximal transcendence*, `docs/surreal/transcendence-over-bounded-support/`. Its README was read by source 17 at `bcac55a`; its `bst:thm:descent` contains the descent of Lemma 35.7.
- [39] O. Bournez and Q. Guilmant, *Surreal fields stable under exponential and logarithmic functions*, arXiv:2201.08199v1 (2022). <https://arxiv.org/abs/2201.08199>. Its Corollary 2.7, the normal-form identification, consulted by source 12; a different paper from [9].
- [40] VladimirReshetnikov/Surreal project, *Hahn–Tate Uniformization at Arbitrary Valuation Rank*, `docs/surcomplex/hahn-tate-uniformization/`, at source 13’s pin `4cdeaec43ff1692ed2ad9f5bdf761a41872975a5`.
<https://github.com/VladimirReshetnikov/Surreal/tree/4cdeaec43ff1692ed2ad9f5bdf761a41872975a5/docs/surcomplex/hahn-tate-uniformization>. Its theta series $\sum_{n \in \mathbb{Z}} (-1)^n q^{n(n-1)/2} u^n$, its theta domain (`tate:thm:theta`) and its product and zeros (`tate:prop:theta-product`) are cited in the theta-descent part; source 13 inspected its README only.
- [41] VladimirReshetnikov/Surreal project, *Omnific Integers and Omnific–Diophantine Geometry*, `docs/surreal/omnific-diophantine-geometry/`. Its definition `odg:def:rings` of **Oz** = $\mathbb{Z} \oplus \Pi$ is cited by this merge for the omnific integers of Corollary 25.3.
- [42] J. P. Bell, M. Coons and E. Rowland, *The rational–transcendental dichotomy of Mahler functions*, *J. Integer Seq.* **16** (2013), Article 13.2.10; arXiv:1210.2070.

- <https://arxiv.org/abs/1210.2070>. Its Lemma 6 is the complex entire-Mahler lemma credited by source 15.
- [43] K. Nishioka, *Mahler Functions and Transcendence*, Lecture Notes in Mathematics 1631, Springer, 1996. <https://doi.org/10.1007/BFb0093672>. Cited by source 15 from publisher metadata only.
 - [44] Yu. V. Matiyasevich, *Diophantine representation of enumerable predicates*, Math. USSR-Izv. **5** (1971), 1–28. <https://doi.org/10.1070/IM1971v005n01ABEH001004>.
 - [45] Z.-W. Sun, *Further results on Hilbert’s Tenth Problem*, Sci. China Math. **64** (2021), 281–306; arXiv:1704.03504v7. <https://arxiv.org/abs/1704.03504>. Cited by source 15 only for undecidability in eleven integer unknowns.
 - [46] C. Faverjon and J. Roques, *Hahn series and Mahler equations: algorithmic aspects*, J. London Math. Soc. (2) **110** (2024), e12945; arXiv:2412.04928. <https://doi.org/10.1112/jlms.12945>. Added by this merge from the single-dilation report’s bibliography; not read for this merge and not used.
 - [47] VladimirReshetnikov/Surreal project, *A Single Dilation Recovers Hahn Support*, docs/surcomplex/single-dilation-hahn-support/. At bcac55a it already discussed Mahler difference algebra and cited [46].
 - [48] VladimirReshetnikov/Surreal project, *Dilation Rigidity of Surreal Numbers and Omnific Integers*, docs/surcomplex/autonomous-dilation-relations/. Its adr:lem:monomial is the exponent-dilation twin of Example 37.7, and its README records Mahler’s 1983 criterion through Nishioka–Nishioka.
 - [49] VladimirReshetnikov/Surreal project, *Set-Sized Quotients of the Omnific Integers*, docs/surreal/set-sized-quotients-of-omnific-integers/, and *Omnific Groups and Lattices*, docs/surreal/omnific-groups-and-lattices/. Their READMEs were read by source 15 at bcac55a; nothing from them is used. Source 20 read the first README at efc5446 and cites its universal quotient theorem, osq:thm:universal, only in Remark 39.5, without auditing that report.
 - [50] NIST Digital Library of Mathematical Functions, Chapter 20, §20.5, especially equations 20.5.3 and 20.5.9 (Jacobi’s triple product), and §20.5(ii) (logarithmic derivatives). <https://dlmf.nist.gov/20.5>. Accessed by source 13 on 23 September 2026.
 - [51] NIST Digital Library of Mathematical Functions, Chapter 23, §23.6 (relations to theta functions) and §23.8 (trigonometric series and products). <https://dlmf.nist.gov/23.6>; <https://dlmf.nist.gov/23.8>. Accessed by source 13 on 23 September 2026. Source 14 consulted §23.3 (differential equations of the Weierstrass function), <https://dlmf.nist.gov/23.3>, on the same day.
 - [52] NIST Digital Library of Mathematical Functions, Chapter 32, §32.2 (differential equations; the definitions of Painlevé I and II). <https://dlmf.nist.gov/32.2>. Accessed by source 14 on 23 September 2026.
 - [53] *Elliptic Curves and Modular Forms*, lecture notes hosted by M. Woodbury, 2010, §16, printed page 32: the Tate equation and invariant differential. <https://www.mi.uni-koeln.de/~woodbury/research/ecnotes.pdf>. Inspected by source 13 as a PDF image; the hosting attribution is not a claim of authorship of every part.

- [54] Y. Ohyama, *Differential relations of theta functions*, Osaka Journal of Mathematics **32** (1995), 431–450. <https://projecteuclid.org/journals/osaka-journal-of-mathematics/volume-32/issue-2/Differential-relations-of-theta-functions/ojm/1200786061.full>. Publication record consulted by source 13; not used in place of a proof.
- [55] A. Sebbar, *Finite and infinite order differential relations for theta functions*, Milan Journal of Mathematics **84** (2016), 317–347. <https://doi.org/10.1007/s00032-016-0261-6>. Publisher abstract and bibliography consulted by source 13, and the publisher abstract by source 14; full text not inspected by either. For the merge of source 14 the bibliographic data were checked against the Crossref record (volume 84, issue 2); the content was not verified.
- [56] V. Bagayoko, L. S. Krapp, S. Kuhlmann, D. Panazzolo and M. Serra, *Automorphisms and derivations on algebras endowed with formal infinite sums*, arXiv:2403.05827, version 2. <https://arxiv.org/abs/2403.05827>. Consulted by source 20 for positive-support summability.
- [57] L. S. Krapp, S. Kuhlmann and M. Serra, *Generalised power series determined by linear recurrence relations*, arXiv:2206.04126; published in Journal of Algebra (2025). <https://arxiv.org/abs/2206.04126>. Cited by source 20 to distinguish exponent-indexed recurrences from the ordinary-indexed recurrences of the recurrence part; nothing is imported. A different paper, and a different author list, from [10].
- [58] VladimirReshetnikov/Surreal project, *Notations for Omnific Integers*, [docs/foundations-and-computation/omnific-notations/](https://github.com/VladimirReshetnikov/Surreal/blob/main/docs/foundations-and-computation/omnific-notations/). Its display `onot:eq:guardbasic` is the element of Theorem 39.3; cited by this merge, not by source 20.