

A parity factorization and sharp stabilization for Cigler's Hankel polynomials

A proposed proof of Conjecture 16 in arXiv:2111.14492v3

Part II (added 29 September 2026): rectangular Schur structure
and sharp recurrences

Part III (added 30 September 2026): cyclotomic pole collapse
and exact minimal recurrences at complex parameters

Research draft prepared for Vladimir Reshetnikov

20 September 2026 (Part I); 29 September 2026 (Part II); 30 September 2026 (Part III)

Abstract

We study the shifted Hankel determinants of Cigler's polynomial extension $c_n(t)$ of the middle binomial coefficients. An even-odd decomposition factors every normalized determinant into two principal minors of powers of simple tridiagonal matrices. This gives a proof of polynomiality, strict coefficient positivity, reciprocity, and the exact degree. An elementary alternant evaluation of the two factors proves the stable-coefficient assertions of Conjecture 16. More precisely, if $a = \lfloor k/2 \rfloor$, $b = \lfloor (k-1)/2 \rfloor$, and $\mathcal{H}_k(n; t) = (-1)^k \binom{n}{2} t^{-\binom{n}{2}} \det(c_{k+i+j}(t))_{0 \leq i, j < n}$, then for $k \geq 2$,

$$\mathcal{H}_k(n; t) = \frac{1}{(1-t)^a (1-t^2)^{ab}} - \binom{n+a}{a-1} t^{n+1} + O(t^{n+2}).$$

Thus the conjectured stable range is optimal, and its first defect has a closed formula. We also obtain fixed-size determinant evaluations, an explicit nonminimal recurrence, and convergence on compact subsets of the open unit disk. Exact integer and polynomial verification programs accompany the article. A stronger rectangular-Schur identity suggested by computation is separated from the proved results and is not used in any proof. [Added 29 September 2026, batch 56: Part II proves this identity and the exact minimal recurrence.] [Added 30 September 2026, batch 66: Part III determines the reduced denominator and the minimal recurrence at every complex parameter, including every root of unity.]

Status and scope. This is an AI-assisted research draft with a full mathematical argument, not an independently refereed or proof-assistant-checked result. The target is Conjecture 16 in Section 6 of Cigler's paper, not Conjectures 13–15 in Section 5, which are already represented in the supplied manifest. A targeted literature search did not locate a later solution of this specific conjecture; that is not a guarantee of priority.

Contents

Part I. A parity factorization and sharp stabilization	6
1 The problem and the main result	6
1.1 Why this problem is not a duplicate	6
1.2 Definitions and conventions	6
1.3 Proof architecture	7
2 Two auxiliary moment sequences	7
2.1 Tridiagonal realizations	8
3 Hankel determinants as principal minors	8
4 The parity factorization	9
5 Strict positivity, degree, and reciprocity	10
6 A small alternant evaluation	11
6.1 The Christoffel determinant identity	11
6.2 The two explicit numerators	11
7 Sharp stable coefficients	13
8 Examples and checks on the statement	14
8.1 The first three shifts	14
8.2 A first-defect example	14
8.3 Signs and exceptional parameters	15
9 Algorithms and further consequences	15
9.1 Exact computation at fixed shift	15
9.2 A concrete, nonminimal recurrence	15
9.3 Actual convergence, not just formal stabilization	16
10 A stronger identity suggested by the computations	17
11 Reproducibility and proof audit	17
11.1 What was checked computationally	17
11.2 Logical dependencies and possible failure points	18
11.3 Literature status and exclusions	18
A The removable specialization at $t = 1$	18
B How to run the package	20

Part II. Rectangular Schur structure and sharp recurrences	21
12 The selected question and the results	21
12.1 Provenance of Part II	21
12.2 The manuscript's title, abstract and status	22
12.3 Notation shared with Part I	23
12.4 Provenance and what is added	24
12.5 Definitions	25
12.6 Main statements	25
13 Laurent moments and the four parity factors	26
13.1 Two elementary Gram identities	26
13.2 The moments	26
13.3 Block elimination	27
14 The missing Toeplitz factorization	28
15 Proof of the Schur identity and coefficient consequences	29
15.1 A tableau model and reciprocal coefficients	30
16 A confluent spectral theorem for rectangles	30
16.1 Statement with an explicit leading coefficient	31
16.2 The confluent bialternant	31
16.3 Degree loss forced by confluent rows	32
16.4 Evaluation of the unique leading term	32
16.5 Why the recurrence is minimal	33
17 Exact recurrences for Cigler's family	33
17.1 Exact computation without a large Hankel matrix	34
18 Generating numerators, negative indices, and a corrected sign	34
18.1 Negative-width continuation	35
18.2 The exact numerator theorem	35
18.3 What this says about the printed Conjecture 18	36
18.4 The period-four signed normalization	36
19 Sharp convergence and all real specializations	37
19.1 The leading nonconstant spectral term	37
19.2 The exceptional real parameters	38
20 Reproducibility, proof dependencies, and formalization	39
20.1 What the accompanying program actually checks	39
20.2 Dependency and trust boundaries	40

20.3 A finite-algebra formalization plan	40
21 Further research questions and topics	41
21.1 Full coefficient structure in Cigler's Conjecture 17	41
21.2 Complex roots of unity and spectral cancellation	41
21.3 Coefficient shape for even shifts	42
21.4 A two-parameter Hankel deformation	42
21.5 A uniform crossover as the poles merge	42
21.6 Broader noncollision criteria and certified algorithms	42
22 Conclusion	42
C Small-shift formulas and independent checks	43
D Package contents and reproduction	43
Part III. Cyclotomic pole collapse and exact minimal recurrences at complex parameters	45
23 The problem, its source, and the mathematical interface	45
23.1 Provenance of Part III	45
23.2 The manuscript's title, abstract and status	47
23.3 Notation shared with Parts I and II	47
23.4 Which question is being resolved?	49
23.5 Cigler's moments and the normalization	50
23.6 What is new here, and what is background?	50
24 The exact cyclotomic classification	50
24.1 Minimal recurrence and pole multiplicity	50
24.2 Small examples	51
24.3 Outline of the proof	52
25 Confluent rectangular characters and their spectral terms	52
25.1 The distinct-variable subset identity	52
25.2 Specialization to the three-cluster alphabet	53
26 Why there is at most one cancellation class	54
26.1 The concavity restriction	54
26.2 Reflection about the canonical center	54
27 A general first subleading coefficient for confluent Schur modes	55
28 Ordinary cyclotomic cancellations stop after one order	56

29 Exact fourth-root product formulas	58
29.1 The odd-shift parity factorization	58
29.2 Binomial and Catalan moment determinants	59
29.3 Evaluation of the two Gram factors at a fourth root	59
30 The exceptional cancellation is exactly three orders	60
30.1 The relevant discrete Fourier coefficient	60
30.2 Root power sums through degree five	60
31 Consequences and explicit forms	62
31.1 Every complex parameter, not just roots of unity	62
31.2 Compact formulas at primitive fourth roots	62
31.3 Quasipolynomial periods and exact reconstruction	63
31.4 A quadratic recurrence-order law	63
32 Verification, dependency audit, and formalization prospects	64
32.1 What the included computations check	64
32.2 Logical dependencies	64
32.3 A practical proof-assistant route	65
33 Further research questions	65
33.1 More than one middle variable	65
33.2 A general bound on cancellation depth	65
33.3 Higher coefficients of a canonical spectral mode	66
33.4 Why do fourth roots force the additional cancellation?	66
33.5 Detuning a root of unity	66
33.6 Numerators and zeros at cyclotomic parameters	66
33.7 Original, unnormalized Hankel determinants	66
33.8 Joint growth of the shift and cyclotomic order	66
33.9 Certified recurrence generation	67
34 Conclusion	67
E An elementary beta determinant and the product evaluations	67
F A minimal exponent algorithm	68
G Package and reproduction	69

1 The problem and the main result

1.1 Why this problem is not a duplicate

The supplied manifest [3] catalogues a report entitled *Product formulas for ballot-polynomial Hankel determinants*, concerning Conjectures 13–15 in Section 5 of Cigler's paper [1]. Those conjectures concern the moment family denoted there by $a_n(t)$. Here the family is $c_n(t)$ from equation (6), and the target is *Conjecture 16 in Section 6*, on pages 21–22 of version 3. The relationship is methodological—polynomial moment sequences and their Hankel determinants—but the input moments and the numbered problem differ. We do not assume that any proof catalogued in the manifest is correct.

The source continues to present this statement as a conjecture in the version located during the search. The arXiv record identifies version 3 as dated 30 December 2021 [1]. The bounded status audit in Section 11 explains what was and was not established about subsequent literature.

1.2 Definitions and conventions

A binomial coefficient $\binom{r}{j}$ is zero unless $0 \leq j \leq r$. Set $c_0(t) = 1$, and for $r \geq 1$ define

$$c_r(t) = \sum_{j=1}^r \binom{\lfloor (r-1)/2 \rfloor}{\lfloor (j-1)/2 \rfloor} \binom{\lfloor r/2 \rfloor}{\lfloor j/2 \rfloor} t^{j-1}. \quad (1.1)$$

This is Cigler's equation (6), with its zero $j = 0$ term omitted so that no negative power of t is displayed. For example,

$$\begin{array}{lll} c_0 = 1, & c_1 = 1, & c_2 = 1 + t, \\ c_3 = 1 + t + t^2, & c_4 = 1 + 2t + 2t^2 + t^3, & c_5 = 1 + 2t + 4t^2 + 2t^3 + t^4. \end{array}$$

For $k, n \geq 0$, write

$$D_k(n; t) = \det(c_{k+i+j}(t))_{0 \leq i, j < n}, \quad D_k(0; t) = 1. \quad (1.2)$$

For $k \geq 1$ initially define, in $\mathbb{Q}(t)$,

$$\mathcal{H}_k(n; t) = (-1)^{k \binom{n}{2}} \frac{D_k(n; t)}{t^{\binom{n}{2}}}. \quad (1.3)$$

Polynomiality, including a well-defined value at $t = 0$, will be proved. It is important not to substitute $t = 0$ into the quotient before this step.

Let

$$a = \left\lfloor \frac{k}{2} \right\rfloor, \quad b = \left\lfloor \frac{k-1}{2} \right\rfloor, \quad \mathcal{F}_k(t) = \frac{1}{(1-t)^a (1-t^2)^{ab}}. \quad (1.4)$$

An expression $O(t^r)$ in a formal-series identity means an element of $t^r \mathbb{Q}[[t]]$; it has no analytic meaning unless one is explicitly stated.

Theorem 1.1 (Conjecture 16, with a sharp first defect). *For every $k \geq 1$ and $n \geq 0$, $\mathcal{H}_k(n; t)$ is a polynomial in $\mathbb{Z}[t]$ of exact degree $(k-1)n$. Its constant and leading coefficients are both 1, every coefficient between them is strictly positive, and*

$$\mathcal{H}_k(n; t) = t^{(k-1)n} \mathcal{H}_k(n; t^{-1}). \quad (1.5)$$

For $k = 1$, $\mathcal{H}_1(n; t) = 1$. For $k \geq 2$,

$$\mathcal{H}_k(n; t) = \mathcal{F}_k(t) - \binom{n+a}{a-1} t^{n+1} + O(t^{n+2}). \quad (1.6)$$

Consequently,

$$[t^j]\mathcal{H}_k(n; t) = [t^j]\mathcal{F}_k(t) \quad (0 \leq j \leq n), \quad (1.7)$$

and $n + 1$ is the first degree of disagreement for every $k \geq 2$.

For $k = 2a$, equation (1.7) has denominator $(1 - t)^a(1 - t^2)^{a^2 - a}$; for $k = 2a + 1$ it has denominator $(1 - t)^a(1 - t^2)^{a^2}$. These are precisely the two stable-coefficient assertions in the source. The first-defect formula (1.6) is a refinement, not part of the original statement.

1.3 Proof architecture

The proof has three independent ingredients. First, the relation $c_{2r+2} = (1 + t)c_{2r+1}$ turns a Hankel matrix into a two-by-two block matrix whose determinant factors by row operations. Second, the resulting factors are principal minors of powers of coefficientwise totally nonnegative tridiagonal matrices. This establishes positivity and degree without any alternant or Schur identity. Finally, small alternant determinants give the first omitted terms of the two factors and hence the exact stable range. All determinant identities needed below are proved in the article.

[Added 29 September 2026, batch 56: Sections 1–11 and Appendices A and B are this report as of 20 September 2026 (Part I). Part II (Sections 12–22 and Appendices C and D; provenance in Section 12.1) is a later manuscript written against this text. It proves the experimental formula of Section 10, determines the exact minimal recurrence that replaces the bound of Proposition 9.1, and re-derives the parity factorization by a second route. It changes no statement of Part I.]

2 Two auxiliary moment sequences

Put $B = 1 + t^2$ and $s = 1 + t$, so that $s^2 = B + 2t$. For a Laurent polynomial in z , CT_z denotes its constant coefficient. Define

$$Q(z) = B + t(z + z^{-1}) = (1 + tz)(1 + t/z), \quad u_r = \text{CT}_z(1 + z)Q(z)^r, \quad v_r = \text{CT}_z(1 - z^2)Q(z)^r. \quad (2.1)$$

Both u_r and v_r belong to $\mathbb{Z}[t]$.

Lemma 2.1 (The pairing relation). *For every $r \geq 0$,*

$$c_{2r+1} = u_r, \quad c_{2r+2} = su_r, \quad s^2u_r - u_{r+1} = tv_r. \quad (2.2)$$

Proof. Expanding $Q(z)^r = (1 + tz)^r(1 + t/z)^r$ gives

$$u_r = \sum_{j=0}^r \binom{r}{j}^2 t^{2j} + \sum_{j=0}^{r-1} \binom{r}{j} \binom{r}{j+1} t^{2j+1}. \quad (2.3)$$

Separating the even and odd indices of (1.1) gives $c_{2r+1} = u_r$. The coefficients of t^{2j} and t^{2j+1} in $(1 + t)u_r$ are respectively

$$\binom{r}{j} \left(\binom{r}{j} + \binom{r}{j-1} \right), \quad \binom{r}{j} \left(\binom{r}{j} + \binom{r}{j+1} \right).$$

Pascal's identity identifies these with the coefficients of c_{2r+2} . For the last assertion, observe that

$$(s^2 - Q(z))(1 + z) = t(1 + z - z^2 - z^{-1}).$$

Since $Q(z) = Q(z^{-1})$, the constant terms of $zQ(z)^r$ and $z^{-1}Q(z)^r$ agree. The unwanted two terms cancel, leaving tv_r . $\square$

2.1 Tridiagonal realizations

Let $e_0, e_1, \dots$ be the standard basis, let $Se_j = e_{j+1}$ be the unilateral shift, and let $E_{00} = e_0 e_0^\top$. For $\varepsilon = 0, 1$ put

$$M_\varepsilon = BI + t(S + S^\top) + \varepsilon t E_{00}. \quad (2.4)$$

Thus every off-diagonal entry adjacent to the diagonal is t , every diagonal entry is $1 + t^2$, and for M_1 the first diagonal entry has the additional summand t . These are infinite banded matrices: every entry of a fixed power and every minor used below is a finite calculation.

Lemma 2.2 (Moment realization). *For $r \geq 0$,*

$$u_r = (M_1^r)_{00}, \quad v_r = (M_0^r)_{00}. \quad (2.5)$$

Proof. For the first identity use Laurent polynomials $f_h = z^h + z^{-h-1}$, $h \geq 0$. Multiplication by $z + z^{-1}$ sends f_0 to $f_0 + f_1$ and sends f_h , $h > 0$, to $f_{h-1} + f_{h+1}$. Also $\text{CT } f_h$ is 1 for $h = 0$ and 0 otherwise. Multiplication by $Q(z)$ in this basis therefore has matrix M_1 , and $\text{CT } Q(z)^r f_0 = (M_1^r)_{00}$. Replacing z by z^{-1} gives the constant term defining u_r .

For the second identity use $g_h = z^h - z^{-h-2}$. Multiplication by $z + z^{-1}$ sends g_0 to g_1 and, for $h > 0$, sends g_h to $g_{h-1} + g_{h+1}$. Again $\text{CT } g_h = \delta_{h0}$. This gives M_0 and $\text{CT } Q(z)^r (1 - z^{-2}) = v_r$. $\square$

3 Hankel determinants as principal minors

For a matrix A , $A[m]$ means its principal submatrix on indices $0, \dots, m-1$, and $\det A[0] = 1$. Define

$$\mathcal{U}_a(m; t) = \det(M_1^a[m]), \quad \mathcal{V}_a(m; t) = \det(M_0^a[m]) \quad (a, m \geq 0). \quad (3.1)$$

Here $M_\varepsilon^a[m]$ means $(M_\varepsilon^a)[m]$, *not* $(M_\varepsilon[m])^a$. Paths may leave the first m indices and return before the a multiplications are complete.

Lemma 3.1 (The auxiliary Gram identity). *For $a, m \geq 0$,*

$$\det(u_{a+i+j})_{0 \leq i, j < m} = t^{m(m-1)} \mathcal{U}_a(m; t), \quad (3.2)$$

$$\det(v_{a+i+j})_{0 \leq i, j < m} = t^{m(m-1)} \mathcal{V}_a(m; t). \quad (3.3)$$

Proof. We prove both statements together. Write $M = M_\varepsilon$ and define a moment functional $L(X^r) = e_0^\top M^r e_0$. Let

$$p_0 = 1, \quad p_1 = X - B - \varepsilon t, \quad p_{j+1} = (X - B)p_j - t^2 p_{j-1} \quad (j \geq 1).$$

Induction gives $p_j(M)e_0 = t^j e_j$. Consequently,

$$L(p_i p_j) = t^{i+j} \delta_{ij}, \quad L(X^a p_i p_j) = t^{i+j} (M^a)_{ij}. \quad (3.4)$$

The change from $1, X, \dots, X^{m-1}$ to the monic polynomials $p_0, \dots, p_{m-1}$ is unit triangular. Changing both bases in the Hankel Gram matrix therefore preserves its determinant. Taking the factors t^i and t^j out of its rows and columns gives $t^{2 \sum_{i=0}^{m-1} i} = t^{m(m-1)}$, as required. This argument is an identity over $\mathbb{Z}[t]$ and does not require $t \neq 0$. $\square$

The normalized auxiliary quantities are thus polynomials before we use any rational evaluation of them. In particular $\mathcal{U}_0(m) = \mathcal{V}_0(m) = 1$ and $\mathcal{U}_a(0) = \mathcal{V}_a(0) = 1$.

4 The parity factorization

Theorem 4.1 (Four parity cases). *For $a, m \geq 0$,*

$$\mathcal{H}_{2a+1}(2m; t) = \mathcal{U}_a(m; t) \mathcal{V}_a(m; t), \quad (4.1)$$

$$\mathcal{H}_{2a+1}(2m+1; t) = \mathcal{U}_a(m+1; t) \mathcal{V}_a(m; t). \quad (4.2)$$

For $a \geq 1$ and $m \geq 0$,

$$\mathcal{H}_{2a}(2m; t) = \mathcal{U}_a(m; t) \mathcal{V}_{a-1}(m; t), \quad (4.3)$$

$$\mathcal{H}_{2a}(2m+1; t) = (1+t) \mathcal{U}_{a-1}(m+1; t) \mathcal{V}_a(m; t). \quad (4.4)$$

In particular the normalization in (1.3) is polynomial.

Proof. Reorder both rows and columns of the Hankel matrix by putting the even indices first. Applying the same permutation on both sides contributes no sign. Write $U_a(r)$ for the unnormalized matrix $(u_{a+i+j})_{0 \leq i, j < r}$ and $V_a(r)$ for the analogous matrix of v 's.

For an odd shift $2a+1$, the four blocks, with the appropriate rectangular sizes, are

$$\begin{pmatrix} (u_{a+i+j}) & (su_{a+i+j}) \\ (su_{a+i+j}) & (u_{a+i+j+1}) \end{pmatrix}.$$

Subtract s times the corresponding even row from each odd row. The lower left block becomes zero and the lower right block becomes $(-tv_{a+i+j})$, by Lemma 2.1. The upper left block has size m when $n = 2m$ and size $m+1$ when $n = 2m+1$. Therefore

$$D_{2a+1}(2m) = (-t)^m \det U_a(m) \det V_a(m),$$

$$D_{2a+1}(2m+1) = (-t)^m \det U_a(m+1) \det V_a(m).$$

Using Lemma 3.1, the total powers of t are respectively $2m^2 - m = \binom{2m}{2}$ and $2m^2 + m = \binom{2m+1}{2}$. Also $(-1)^m = (-1)^{\binom{n}{2}}$ for both $n = 2m$ and $n = 2m+1$. This proves the two odd-shift formulas.

For shift $2a$ and size $2m$, the reordered matrix is

$$\begin{pmatrix} sU_{a-1}(m) & U_a(m) \\ U_a(m) & sU_a(m) \end{pmatrix}.$$

Work temporarily in $\mathbb{Q}(t)$ and subtract s^{-1} times each bottom row from the matching top row. The upper right block becomes zero, and the upper left block is $(t/s)V_{a-1}(m)$. Hence

$$D_{2a}(2m) = t^m \det V_{a-1}(m) \det U_a(m).$$

For size $2m+1$, the even block has $m+1$ rows. Subtract s^{-1} times even row $i+1$ from odd row i , for $0 \leq i < m$. The lower left block vanishes; the lower right block becomes $(t/s)V_a(m)$, while the upper left block is $sU_{a-1}(m+1)$. Thus

$$D_{2a}(2m+1) = st^m \det U_{a-1}(m+1) \det V_a(m).$$

The Gram identities give exactly the same two normalization exponents as before. These are polynomial identities after denominators are cleared, so the temporary use of $1/s$ causes no exception at $t = -1$. $\square$

5 Strict positivity, degree, and reciprocity

For polynomials P, Q , write $P \succeq Q$ if $P - Q$ has nonnegative integer coefficients. A matrix is *coefficientwise totally nonnegative* if every finite minor, with row and column indices in increasing order, is a polynomial with nonnegative coefficients.

Lemma 5.1 (Total nonnegativity of the auxiliary matrices). *Both M_0 and M_1 , and all of their nonnegative integer powers, are coefficientwise totally nonnegative.*

Proof. Since $S^\top S = I$ for the unilateral shift,

$$M_0 = (I + tS^\top)(I + tS). \quad (5.1)$$

The two bidiagonal factors are totally nonnegative: a nonzero minor has an order-preserving matching of its selected rows and columns and has a nonnegative monomial weight. This can also be checked by expanding at the first selected row; the bidiagonal zero pattern prevents a negatively signed nonzero matching.

The Cauchy–Binet formula proves total nonnegativity of their product. Although the matrices are infinite, each of these sums is finite because the factors are banded. Next $M_1 = M_0 + tE_{00}$. A minor not selecting both row 0 and column 0 is unchanged. A minor selecting both acquires t times the minor obtained by deleting its first row and first column. The cofactor sign is positive because the selected indices are increasing. Thus adding this corner entry preserves coefficientwise total nonnegativity. Repeated Cauchy–Binet proves the assertion for powers. $\square$

Proposition 5.2 (Support and reciprocity of the factors). *For $a, m \geq 0$, $\mathcal{U}_a(m)$ and $\mathcal{V}_a(m)$ are reciprocal polynomials of exact degree $2am$, with constant and leading coefficients 1. For $a, m \geq 1$, every coefficient of $\mathcal{U}_a(m)$ is positive, whereas $\mathcal{V}_a(m)$ has a positive coefficient at every even degree between 0 and $2am$ and no odd-degree terms. More quantitatively,*

$$\mathcal{U}_a(m; t) \succeq (1 + t + \cdots + t^{2m})^a, \quad (5.2)$$

$$\mathcal{V}_a(m; t) \succeq (1 + t^2 + \cdots + t^{2m})^a. \quad (5.3)$$

Proof. The matrices satisfy $M_\varepsilon(t^{-1}) = t^{-2}M_\varepsilon(t)$. Taking the a th power and an m -row principal minor gives

$$\mathcal{U}_a(m; t) = t^{2am}\mathcal{U}_a(m; t^{-1}), \quad \mathcal{V}_a(m; t) = t^{2am}\mathcal{V}_a(m; t^{-1}).$$

At $t = 0$, the matrices are I , so both determinants have constant term 1. Their exact degrees and leading coefficients follow.

The tridiagonal determinant recurrence, with $B = 1 + t^2$, gives

$$\det M_1[m] = 1 + t + \cdots + t^{2m}, \quad \det M_0[m] = 1 + t^2 + \cdots + t^{2m}.$$

For example the second sequence has initial values $1, B$ and recurrence $P_m = BP_{m-1} - t^2P_{m-2}$; the first has initial values $1, B + t$ and the same recurrence. The displayed finite sums satisfy these recurrences.

Expand the principal minor of M_ε^a by repeated Cauchy–Binet. All terms are coefficientwise nonnegative. Selecting the same index set $\{0, \dots, m-1\}$ at every intermediate stage supplies the term $(\det M_\varepsilon[m])^a$, proving the two lower bounds. Finally $M_0(-t) = JM_0(t)J$, where $J = \text{diag}(1, -1, 1, -1, \dots)$. Its principal minors of powers are unchanged, so $\mathcal{V}_a(m)$ is even. $\square$

Proof of the positivity and reciprocity part of Theorem 1.1. Apply Theorem 4.1. The degrees in its four lines are

$$4am, \quad 2a(2m+1), \quad 2m(2a-1), \quad (2a-1)(2m+1),$$

which equal $(k-1)n$ in each case. All factors are reciprocal, including $1+t$, and their constant terms are 1.

For strict positivity, the full consecutive support of a nonconstant $\mathcal{U}$ factor overlaps the translates arising from the even support of a $\mathcal{V}$ factor. More explicitly, an interval $\{0, \dots, L\}$ with $L \geq 1$, added to $\{0, 2, \dots, 2R\}$, covers every integer from 0 to $L+2R$. The only exceptional-looking even-shift case is $a=1$, n odd: there the factor $1+t$ supplies precisely the needed interval of length one. The cases $n=0$ and $k=1$ give the constant polynomial 1 directly. $\square$

6 A small alternant evaluation

Positivity did not require symmetric-function identities. The stabilization proof will also be elementary: the relevant formulas are quotients of explicit alternating determinants. This section supplies their derivation, including the classical moment determinant identity that is often used for this purpose; compare the general framework in [2].

6.1 The Christoffel determinant identity

Lemma 6.1 (Moment modification). *Let L be a moment functional over a characteristic-zero field, with monic orthogonal polynomials p_j of nonzero norms $h_j = L(p_j^2)$. Put $H_m = \det(L(X^{i+j}))_{0 \leq i,j < m}$. For independent parameters $\eta_1, \dots, \eta_a$,*

$$\frac{\det(L(X^{i+j} \prod_{r=1}^a (X - \eta_r)))_{0 \leq i,j < m}}{H_m} = (-1)^{am} \frac{\det(p_{m+j-1}(\eta_i))_{1 \leq i,j \leq a}}{\Delta(\eta)}, \quad (6.1)$$

where $\Delta(\eta) = \prod_{i < j} (\eta_j - \eta_i)$.

Proof. Form an $(m+a)$ -square matrix whose first m rows, indexed by r , are $L(X^r X^j)$ for $0 \leq j < m+a$, and whose last a rows are η_i^j . Change the column polynomials from the monomials to $p_0, \dots, p_{m+a-1}$. The change is unit triangular. Orthogonality makes the upper right block zero, and the determinant of the upper left block is $\prod_{r=0}^{m-1} h_r = H_m$. Thus this determinant is $H_m \det(p_{m+j-1}(\eta_i))$.

Alternatively change the column polynomials to $1, X, \dots, X^{a-1}, P, XP, \dots, X^{m-1}P$, where $P = \prod_r (X - \eta_r)$. This is again a monic unit-triangular change. Move the last a rows to the top, contributing $(-1)^{am}$. The top left block is a Vandermonde matrix and the top right block is zero. The bottom right block is the modified moment matrix on the left of (6.1). Equating the two evaluations proves the identity. $\square$

6.2 The two explicit numerators

For $a, m \geq 0$ and $\delta \in \{0, 1\}$, define

$$N_{a,m}^\delta(t) = \det \left(\binom{j-1}{i-1} t^{j-i} - \binom{2m+2a-j+\delta}{i-1} t^{2m+2a-j+\delta-i+1} \right)_{1 \leq i,j \leq a}. \quad (6.2)$$

The first summand is interpreted as zero when $j < i$, without evaluating its negative power. An empty determinant is 1.

Theorem 6.2 (Fixed-size evaluations). *For all $a, m \geq 0$,*

$$\mathcal{U}_a(m; t) = \frac{N_{a,m}^0(t)}{(1-t)^a(1-t^2)^{\binom{a}{2}}}, \quad (6.3)$$

$$\mathcal{V}_a(m; t) = \frac{N_{a,m}^1(t)}{(1-t^2)^{\binom{a+1}{2}}}. \quad (6.4)$$

These identities hold in $\mathbb{Q}(t)$ and determine the polynomials already constructed in (3.1); apparent singularities at $t = 1$ or $t = -1$ are removable. The determinant orders are a , independently of m .

Proof. Work first over $\mathbb{Q}(t)$. Let $C_0(y) = 1$, $C_1(y) = y$, and $C_{j+1}(y) = yC_j(y) - C_{j-1}(y)$; also put $C_{-1} = 0$. The orthogonal polynomials in Lemma 3.1 are

$$p_j^\varepsilon(X) = t^j \left[C_j \left(\frac{X-B}{t} \right) - \varepsilon C_{j-1} \left(\frac{X-B}{t} \right) \right].$$

Define monic polynomials $A_j^\varepsilon(y) = C_j(y) + \varepsilon C_{j-1}(y)$ and put

$$\eta_i = B - t(x_i + x_i^{-1}).$$

Then $p_j^\varepsilon(\eta_i) = (-t)^j A_j^\varepsilon(x_i + x_i^{-1})$. The elementary recurrence gives

$$A_j^0(x + x^{-1}) = \frac{x^{j+1} - x^{-j-1}}{x - x^{-1}}, \quad (6.5)$$

$$A_j^1(x + x^{-1}) = \frac{x^{j+1/2} - x^{-j-1/2}}{x^{1/2} - x^{-1/2}} = \sum_{r=-j}^j x^r. \quad (6.6)$$

The half powers in the middle expression cancel; the expression on the right is a Laurent polynomial.

Since the A_j^ε are monic in y , their first a evaluation columns have determinant $\Delta(y)$, where $y_i = x_i + x_i^{-1}$. Apply Lemma 6.1, cancel the column powers of $-t$ and the Vandermonde scale, and then specialize every η_i to zero. Choosing every $x_i = t$ gives

$$\det(M_\varepsilon^a[m]) = t^{am} \frac{\det(A_{m+j-1}^\varepsilon(x_i + x_i^{-1}))}{\det(A_{j-1}^\varepsilon(x_i + x_i^{-1}))} \Big|_{x_1=\dots=x_a=t}. \quad (6.7)$$

The equal-parameter value is the limit of an alternating quotient, not a quotient of two separately substituted zero determinants.

Reverse all columns in the numerator and denominator of this quotient and use (6.5)–(6.6). For $\delta = 1 - \varepsilon$, multiplication of row i by the appropriate power of x_i gives

$$\left(\prod_{i=1}^a x_i \right)^m \frac{\det(A_{m+j-1}^\varepsilon(x_i + x_i^{-1}))}{\det(A_{j-1}^\varepsilon(x_i + x_i^{-1}))} = \frac{\det(x_i^{j-1} - x_i^{2m+2a-j+\delta})}{\det(x_i^{j-1} - x_i^{2a-j+\delta})}. \quad (6.8)$$

At $x_i = t$, the prefactor on the left is exactly t^{am} .

The denominator on the right has the factorization

$$\det(x_i^{j-1} - x_i^{2a-j+\delta}) = \Delta(x) \prod_{i=1}^a (1 - x_i^{1+\delta}) \prod_{i < j} (1 - x_i x_j). \quad (6.9)$$

Here is a direct verification. Alternation supplies $\Delta(x)$. At $x_i = 1$ a row vanishes, and for $\delta = 1$ it also vanishes at $x_i = -1$. At $x_j = x_i^{-1}$ the two rows are proportional, with proportionality

factor $-x_i^{-(2a-1+\delta)}$, so every $1 - x_i x_j$ divides the determinant. These factors have exactly its total degree. Its lowest homogeneous term is $\Delta(x)$, which fixes the multiplicative constant as 1.

Finally, for polynomials f_j , Taylor expansion gives the confluent identity

$$\frac{\det(f_j(x_i))}{\Delta(x)} \Big|_{x_1=\dots=x_a=t} = \det \left(\frac{f_j^{(i-1)}(t)}{(i-1)!} \right)_{1 \leq i, j \leq a}. \quad (6.10)$$

Indeed the lowest alternating term in the Taylor expansion around t uses the distinct exponents $0, 1, \dots, a-1$. Apply (6.10) to the numerator in (6.8). It gives exactly (6.2). The remaining factors in (6.9), after equal specialization, give the denominators in (6.3) and (6.4). $\square$

Remark 6.3. The quotients before equal specialization are familiar rectangular orthogonal and symplectic character alternants. No character theory, branching formula, or bounded Schur-sum identity is being imported: the moment identity, the recurrence, the denominator factorization, and the confluent limit above give a complete determinant proof.

7 Sharp stable coefficients

Lemma 7.1 (First nonconstant numerator term). *For $a \geq 1$, $m \geq 0$, and $\delta \in \{0, 1\}$, put $r = 2m + 1 + \delta$. Then*

$$N_{a,m}^\delta(t) = 1 - \binom{a+r-1}{a-1} t^r + O(t^{r+1}). \quad (7.1)$$

Proof. Before equal specialization, expand $\det(x_i^{j-1} - x_i^{2m+2a-j+\delta})$ by choosing either the low or the high monomial in each column. If the high monomial is selected in column j , the total degree increases, relative to $\Delta(x)$, by

$$2m + 2a - 2j + 1 + \delta = r + 2(a - j).$$

The all-low choice is $\Delta(x)$. The unique choice of smallest positive additional degree is to select only the high monomial in column a ; its sign is negative and its additional degree is r . Every other choice has strictly larger additional degree.

After division by $\Delta(x)$ and equal specialization, the coefficient of this first correction can be computed directly from (6.10). The column exponents are $0, 1, \dots, a-2, a-1+r$. The first $a-1$ columns are triangular with unit diagonal, so the remaining determinant is $\binom{a-1+r}{a-1} t^r$. This proves the assertion, including $a = 1$. $\square$

Define the formal series

$$U_a^\infty(t) = \frac{1}{(1-t)^a (1-t^2)^{\binom{a}{2}}}, \quad V_a^\infty(t) = \frac{1}{(1-t^2)^{\binom{a+1}{2}}}. \quad (7.2)$$

Their constant terms are 1. For $a = 0$ both are exactly 1. Theorem 6.2 and Lemma 7.1 imply, for $a \geq 1$,

$$\mathcal{U}_a(m; t) = U_a^\infty(t) - \binom{2m+a}{a-1} t^{2m+1} + O(t^{2m+2}), \quad (7.3)$$

$$\mathcal{V}_a(m; t) = V_a^\infty(t) - \binom{2m+a+1}{a-1} t^{2m+2} + O(t^{2m+3}). \quad (7.4)$$

Multiplying by a series with constant term 1 does not change the first nonzero correction, which is why the same binomial coefficient appears in these formulas.

Completion of Theorem 1.1. For $n = 2m$ the earliest correction in both (4.1) and (4.3) comes from $\mathcal{U}_a(m)$, at degree $2m + 1 = n + 1$. Its magnitude is $\binom{2m+a}{a-1} = \binom{n+a}{a-1}$. The products of the corresponding limiting series are respectively

$$U_a^\infty V_a^\infty = \frac{1}{(1-t)^a(1-t^2)^{a^2}}, \quad U_a^\infty V_{a-1}^\infty = \frac{1}{(1-t)^a(1-t^2)^{a(a-1)}}.$$

For $n = 2m + 1$ the earliest correction in both (4.2) and (4.4) comes from $\mathcal{V}_a(m)$, at degree $2m + 2 = n + 1$, with magnitude $\binom{2m+a+1}{a-1} = \binom{n+a}{a-1}$. For the last parity case the limiting product simplifies as

$$(1+t)U_{a-1}^\infty V_a^\infty = \frac{1+t}{(1-t)^{a-1}(1-t^2)^{a^2-a+1}} = \frac{1}{(1-t)^a(1-t^2)^{a(a-1)}}.$$

All multiplying factors have constant term 1, so they do not alter the first correction coefficient. When $a = 1$, any factor with index $a - 1 = 0$ is identically 1, and the same argument applies. The positive binomial coefficient proves optimality of the range. The case $k = 1$ was already obtained directly from the factorization. $\square$

Corollary 7.2 (Coefficient extraction without a Hankel determinant). *For $k \geq 2$, $0 \leq j \leq n$, and a, b as in (1.4),*

$$[t^j]\mathcal{H}_k(n; t) = \sum_{\ell=0}^{\lfloor j/2 \rfloor} W(ab, \ell) W(a, j-2\ell), \quad (7.5)$$

where $W(q, r) = \binom{q+r-1}{r}$ for $q \geq 1$, and $W(0, r) = 1$ for $r = 0$ and 0 otherwise.

Proof. Expand $(1-t)^{-a}$ and $(1-t^2)^{-ab}$ and use (1.7). The convention for $W(0, r)$ handles $k = 2$ without an artificial negative upper binomial index. $\square$

8 Examples and checks on the statement

8.1 The first three shifts

The factorization immediately gives

$$\mathcal{H}_1(n; t) = 1, \quad \mathcal{H}_2(n; t) = 1 + t + \cdots + t^n. \quad (8.1)$$

For $k = 3$, insert $\mathcal{U}_1(m) = 1 + t + \cdots + t^{2m}$ and $\mathcal{V}_1(m) = 1 + t^2 + \cdots + t^{2m}$ in the two parity cases. Both simplify to

$$\mathcal{H}_3(n; t) = \frac{(1-t^{n+1})(1-t^{n+2})}{(1-t)(1-t^2)}. \quad (8.2)$$

These small-shift formulas are already printed in Cigler's paper; they serve as indexing and sign checks, not as new results.

8.2 A first-defect example

For $k = 4$, the limiting series is

$$\mathcal{F}_4(t) = \frac{1}{(1-t)^2(1-t^2)^2} = 1 + 2t + 5t^2 + 8t^3 + 14t^4 + \cdots.$$

At $n = 2$ the determinant gives

$$\mathcal{H}_4(2; t) = 1 + 2t + 5t^2 + 4t^3 + 5t^4 + 2t^5 + t^6.$$

The first three coefficients agree. At degree $3 = n + 1$ the deficit is $8 - 4 = 4 = \binom{2+2}{2-1}$, exactly as predicted. At $n = 3$,

$$\mathcal{H}_4(3; t) = 1 + 2t + 5t^2 + 8t^3 + 9t^4 + 9t^5 + 8t^6 + 5t^7 + 2t^8 + t^9,$$

and the first deficit is $14 - 9 = 5 = \binom{3+2}{2-1}$.

These examples also delimit the result. The coefficients 5, 4, 5 in $\mathcal{H}_4(2; t)$ have a strict valley. Thus positivity and reciprocity do *not* imply unimodality here. No unimodality or log-concavity assertion is included in Theorem 1.1.

8.3 Signs and exceptional parameters

For real $t > 0$, strict coefficient positivity gives the exact sign

$$\operatorname{sgn} D_k(n; t) = (-1)^{k \binom{n}{2}}. \quad (8.3)$$

At $t = 0$ the normalized polynomial has value 1, although the original Hankel determinant vanishes when $n \geq 2$. At $t = 1$ the source moments are the middle binomial coefficients. At $t = -1$ the factor $1 + t$ in (4.4) shows that every positive even-shift, odd-size determinant vanishes. These statements follow from polynomial identities; no argument uses an undefined specialization of a rational formula.

9 Algorithms and further consequences

9.1 Exact computation at fixed shift

Theorem 6.2 evaluates the factors through determinants of order at most $\lfloor k/2 \rfloor$, even when n is very large. A practical algorithm is to determine the parity of k, n , compute the one or two numerators in (6.2), divide by their proved exact denominators, and multiply according to Theorem 4.1. For $t = \pm 1$, use the principal-minor definition or take the polynomial limit instead of evaluating a zero denominator.

The reduction is in determinant *order*, not in bit complexity. The entries contain powers whose exponents grow with n , and an expanded answer has degree $(k - 1)n$. Once entries are constructed, ordinary dense elimination uses a cubic number of field operations in the small determinant order; large integers and polynomial output still carry their usual costs. For only the first $j \leq n$ coefficients, (7.5) avoids both determinant constructions.

9.2 A concrete, nonminimal recurrence

Proposition 9.1 (Rationality at each fixed shift). *For $k \geq 2$, set*

$$d_k = \begin{cases} (a - 1)^2, & k = 2a, \\ a(a - 1), & k = 2a + 1. \end{cases}$$

The series $\sum_{n \geq 0} \mathcal{H}_k(n; t)x^n$ is rational over $\mathbb{Q}(t)$, with a denominator dividing

$$\prod_{q=0}^{k-1} (1 - t^{2q}x^2)^{d_{k+1}}. \quad (9.1)$$

Equivalently, with E denoting forward shift in n ,

$$\prod_{q=0}^{k-1} (E^2 - t^{2q})^{d_{k+1}} \mathcal{H}_k(n; t) = 0. \quad (9.2)$$

The bound is not asserted to be minimal and is not a resolution of all the more precise denominator and numerator assertions in Conjecture 18.

Proof. Expand (6.2) column by column. A term selecting q high columns has the form $P(m; t)t^{2qm}$, where P is a polynomial in m with coefficients in $\mathbb{Q}(t)$. Its degree in m is at most $\sum_{i=1}^a (i-1) = \binom{a}{2}$, because the binomial in row i has degree at most $i-1$. The denominators in Theorem 6.2 do not depend on m . Thus each auxiliary factor is a finite sum of $m^j(t^{2q})^m$, with $0 \leq q \leq a$ and $j \leq \binom{a}{2}$.

The products in the two parity subsequences have $0 \leq q \leq k-1$. Their degree bounds are $2\binom{a}{2} = a(a-1)$ in the odd-shift case and $\binom{a}{2} + \binom{a-1}{2} = (a-1)^2$ in the even-shift case. Replacing m by $m+1$ changes neither bound. A polynomial of degree at most d times λ^m is annihilated by $(E_m - \lambda)^{d+1}$, as follows by taking successive finite differences. Interleaving the even and odd subsequences replaces E_m by E^2 , proving the claim. $\square$

[Added 29 September 2026, batch 56: in the notation of this proposition, the exact reduced denominator is $\prod_{q=0}^{k-1} (1 - t^q x)^{1 + \lfloor q(k-1-q)/2 \rfloor}$, with no cancellation over $\mathbb{Q}(t)$ or at any real $t \notin \{0, \pm 1\}$ (Theorem 12.2; Part II writes z for the variable x). Part II also gives the numerator degree and reciprocity sign (Theorem 18.2), the signed denominator for odd k (Theorem 18.3), and the reduced denominators at $t = 0, \pm 1$ (Theorem 19.2). Cigler's denominators (84) and (86) are minimal; the positive sign printed in his (85) is false when a is even, the correct sign being $(-1)^{a+1}$.]

For the unadjusted normalized determinant $d_k(n; t) = t^{-\binom{n}{2}} D_k(n; t)$, the even- k recurrence is unchanged. For odd k , the period-four sign replaces $1 - t^{2q}x^2$ in the displayed coarse denominator by $1 + t^{2q}x^2$.

9.3 Actual convergence, not just formal stabilization

Proposition 9.2 (Uniform convergence inside the unit disk). *Fix $k \geq 2$ and $0 < r < 1$. There is a constant $C_{k,r}$ such that, for all $n \geq 0$,*

$$\sup_{|t| \leq r} |\mathcal{H}_k(n; t) - \mathcal{F}_k(t)| \leq C_{k,r} (n+1) \binom{a}{2} r^{n+1}. \quad (9.3)$$

In particular the polynomials converge to $\mathcal{F}_k$ locally uniformly on $|t| < 1$.

Proof. In the expansion of $N_{a,m}^\delta - 1$, every nonzero term has degree at least $2m + 1 + \delta$ in t . There are finitely many determinant terms for fixed a , and each coefficient is bounded by a constant times $(m+1)\binom{a}{2}$, by the row-degree estimate just used. It follows, uniformly on $|t| \leq r$, that

$$N_{a,m}^\delta(t) - 1 = O_{a,r}((m+1)\binom{a}{2}r^{2m+1+\delta}).$$

The two fixed denominators in Theorem 6.2 are bounded away from zero on this disk. Hence the factors converge uniformly to (7.2), with these bounds, and in particular remain uniformly bounded as m varies. Subtract the limiting products in the four parity formulas. The earliest error has exponent $n+1$, exactly as in Section 7; the largest polynomial exponent is $\binom{a}{2}$. The finitely many small cases can be absorbed in $C_{k,r}$. $\square$

[Added 29 September 2026, batch 56: for fixed $0 < t < 1$ the exact rate is $\mathcal{F}_k(t) - \mathcal{H}_k(n; t) \sim n^{a-1}t^{n+1}/((a-1)!(1-t)^{b+1}(1-t^2)^{(a-1)b})$, with relative error $O(1/n)$ locally uniformly on $(0, 1)$ (Theorem 19.1).]

10 A stronger identity suggested by the computations

The discovery phase suggested a compact formula not needed for any theorem above. We record it as a separate conjecture so that a promising observation is not silently promoted to a proof.

For an alphabet $X = (x_1, \dots, x_k)$, define $h_j(X)$ by $\sum_{j \geq 0} h_j(X) z^j = \prod_{i=1}^k (1 - x_i z)^{-1}$, with $h_j = 0$ for $j < 0$. The rectangular Schur polynomial is

$$s_{(n^a)}(X) = \det(h_{n-i+j}(X))_{1 \leq i, j \leq a}.$$

Conjecture 10.1 (Experimental rectangular-Schur formula). With $a = \lfloor k/2 \rfloor$ and $b = \lfloor (k-1)/2 \rfloor$,

$$\mathcal{H}_k(n; t) = s_{(n^a)}(\underbrace{1, \dots, 1}_a, t, \underbrace{t^2, \dots, t^2}_b). \quad (10.1)$$

The conjecture agrees with all 1,248 exact evaluations in the range recorded in Section 11.1. For instance, at $k = 4, n = 2$, its right-hand side is $h_2^2 - h_1 h_3$ in the alphabet $(1, 1, t, t^2)$, giving the polynomial displayed in Section 8.

A proof of (10.1) would give an attractive unified symmetric-function interpretation. It may be approachable through factorizations of rectangular Schur polynomials into classical-character alternants. This article does not assert such a factorization without proof. The proof of Conjecture 16 instead uses the parity factors, coefficientwise total nonnegativity, and the fully derived alternant formulas. The larger conjecture can be deleted from this article without changing any proved statement.

[Added 29 September 2026, batch 56: Conjecture 10.1 is proved in Part II for all $k \geq 1, n \geq 0$ as an identity in $\mathbb{Z}[t]$ (Theorem 12.1), through a Toeplitz factorization with a boundary parameter (Theorem 14.1). None of Part I's proofs is changed.]

11 Reproducibility and proof audit

11.1 What was checked computationally

The accompanying program `code/verify.py` uses exact integers and exact polynomials in $\mathbb{Z}[t]$. It does not infer symbolic identities from floating-point samples or from interpolation. The direct moment routine implements the source definition (1.1); the principal minor routine independently multiplies the banded matrices; and the small determinant routine implements (6.2).

Check	Range	Cases
Original Hankel versus parity formula	$1 \leq k \leq 12, 0 \leq n \leq 12$, eight integer parameters	1,248
Auxiliary corner versus alternant	$0 \leq a \leq 7, 0 \leq m \leq 8$, both factors, six parameters	864
Three-way polynomial equality; degree, positivity, reciprocity, stability, first defect	$1 \leq k \leq 10, 0 \leq n \leq 8$, in $\mathbb{Z}[t]$	90
Auxiliary first numerator term	$1 \leq a \leq 6, 0 \leq m \leq 6$, both numerators, in $\mathbb{Z}[t]$	84
Experimental Schur formula only	$1 \leq k \leq 12, 0 \leq n \leq 12$, eight integer parameters	1,248

The eight parameters are $-3, -2, -1, 0, 1, 2, 3, 5$. The six-parameter alternant test omits ± 1 , where its displayed denominator vanishes; these parameters are nevertheless covered by the original-versus-corner checks. All checks passed with Python 3.13.5 and SymPy 1.14.0. The execution record is in `data/verification.json`, with the console transcript in `data/verification.log` and coefficient arrays in `data/sample_polynomials.json`. A

supplementary script checks the specialization products in Appendix A in 234 cases and the recurrence in Proposition 9.1 in 126 cases; its exact ranges and results are recorded in `data/additional_verification.json`. These finite checks supplement the all-index proofs; they are not a substitute for them.

11.2 Logical dependencies and possible failure points

The main theorem depends only on the binomial definition, the Laurent basis calculation, elementary block row operations, Cauchy–Binet, monic orthogonal-polynomial recurrences, and alternating determinants. The proof of the moment-modification identity is included rather than used as an unexplained import. The exact role of each delicate point is as follows.

The odd/even row reorder is applied to both rows and columns, so there is no permutation sign. The only sign remaining is $(-1)^{\lfloor n/2 \rfloor}$ for odd shifts, which equals the prescribed $(-1)^{\binom{n}{2}}$. The temporary division by $1+t$ in the even-shift factorization is removed before specialization. The auxiliary normalization is justified by an identity in $\mathbb{Z}[t]$, not by dividing a determinant whose vanishing has not been established. Equal-variable alternants are evaluated after cancelling the Vandermonde, not by a 0/0 substitution. Finally, strict positivity is obtained from explicit Cauchy–Binet summands; nonnegativity alone would not have been sufficient.

11.3 Literature status and exclusions

The source pages and their mathematical notation were inspected directly. Searches by the paper identifier, its title, and the exact numbered conjecture did not locate a later proof of this specific statement. A search hit for Manuel Kauers's *A Proof of Conjecture 16* [4] was investigated separately: indexed material identifies its target as the sequence A339987, not Cigler's $c_n(t)$ conjecture. It was therefore not counted as a solution of the present problem. The full talk PDF was not successfully fetched, so this disambiguation rests on the indexed author material rather than a complete reading of the talk.

The article makes no absolute priority claim. It also does not claim to settle all parts of Cigler's Conjectures 17 and 18, or the separate Schur formula (10.1). What is claimed mathematically is the full argument for Conjecture 16 and the additional results explicitly proved here. Independent expert checking remains appropriate before publication or addition to a collection of established theorems.

[Added 29 September 2026, batch 56: Part II proves the Schur formula and corrected forms of the generating-function assertions of Conjecture 18; Conjecture 17 remains open (Section 21).]

A The removable specialization at $t = 1$

The factors have convenient product evaluations at 1:

$$\mathcal{U}_a(m; 1) = \prod_{r=1}^a \frac{2m+2r-1}{2r-1} \prod_{1 \leq r < s \leq a} \frac{2m+r+s-1}{r+s-1}, \quad (\text{A.1})$$

$$\mathcal{V}_a(m; 1) = \prod_{r=1}^a \frac{m+r}{r} \prod_{1 \leq r < s \leq a} \frac{2m+r+s}{r+s}. \quad (\text{A.2})$$

Empty products have value 1. To see these formulas directly from the proved alternants, set $x_i = e^{z_i}$ before taking the equal-parameter limit in (6.7). For $\mathcal{V}$, the numerator contains $\sinh((m+r)z_i)$ and the denominator $\sinh(rz_i)$, with $1 \leq r \leq a$. For $\mathcal{U}$, replace r by $r - \frac{1}{2}$. The

lowest nonzero alternating term of $\det(\sinh(\rho_j z_i))$ is a common nonzero constant times

$$\left(\prod_i z_i\right) \Delta(z_1^2, \dots, z_a^2) \left(\prod_j \rho_j\right) \Delta(\rho_1^2, \dots, \rho_a^2).$$

This follows by selecting the powers $1, 3, \dots, 2a - 1$ in the Taylor series. In the ratio, the differences between the ρ_j cancel, whereas the sums give the second products in (A.1)–(A.2). These specialize the new parity formula to the classical middle-binomial Hankel setting already treated in the source; the unweighted setting itself is not claimed as new.

B How to run the package

The archive contains the article source and PDF, three Python files, exact verification output, example coefficient arrays, and a status/source audit. The commands, from the archive's root directory, are

```
python -m pip install -r requirements.txt
python code/verify.py
python code/verify_additional.py
pdflatex -interaction=nonstopmode -halt-on-error article.tex
pdflatex -interaction=nonstopmode -halt-on-error article.tex
```

The `Makefile` provides `make verify` and `make pdf`. The verification program stops at the first failed assertion or inexact division. It records the mathematical test ranges and software versions, not just a generic success message. Its stronger Schur checks are labelled `EXPERIMENTAL` in both the code and the output.

Part II. Rectangular Schur structure and sharp recurrences

Sections 1–11 and Appendices A and B form the original report (Part I) of 20 September 2026, unchanged except for dated pointers to this addition, marked [Added 29 September 2026, batch 56: ...]. Part II was added on 29 September 2026 in batch 56 of ProveIt’s incoming-report intake, from one manuscript written against the pinned state of Part I. It proves Part I’s experimental rectangular-Schur formula (Conjecture 10.1) and replaces the nonminimal recurrence of Proposition 9.1 by the exact minimal one. Its Sections 12–22 and Appendices C and D are the manuscript’s Sections 1–11 and Appendices A and B.

12 The selected question and the results

12.1 Provenance of Part II

Part II is batch-56 manuscript 01, *Rectangular Schur Structure and Sharp Recurrences for Cigler’s Hankel Polynomials: A proof of the repository’s Schur conjecture, exact pole multiplicities, and corrected generating-function reciprocity*, an AI-assisted research draft “prepared for Vladimir Reshetnikov” and dated 29 September 2026 (22 pages, A4, as delivered). It arrived as the archive `ProveIt_Cigler_Schur_Recurrences.zip` (inner directory `cigler_schur_research/`, main file `article.tex`) in ProveIt commit `e47ed8547` and was placed in `ae718a441`. It was written against ProveIt commit `e9c09b125` (in full `e9c09b12549ea4e3bea7cebc52b450b761f6f581`), at which Part I’s `article.tex` was the Git blob `f33543af1f` and the `article.tex` of the sibling report `cigler-conjecture-8-schur` [5] the blob `9c984baeeb`. No commit touched this report between `dc54c3cb3` (which brought the research-report collection into ProveIt) and the placement, so every statement the manuscript makes about “the repository’s report” or “the inspected ProveIt report” refers to Part I as printed above. The manuscript read Part I at the pinned commit through a GitHub connector, not from a library copy (its source audit, `02-schur-recurrences-sources.md`); Part I’s file is `SetTheory/Cardinals/docs/reports/hankel-determinants/catalan-and-ballot/cigler-conjecture-16-parity/article.tex`.

Its code and records are shipped beside Part I’s files with the prefix `02-schur-recurrences-` (Part I’s unprefixed files count as 01): the verifier `code/02-schur-recurrences-verify.py` and its Makefile `code/02-schur-recurrences-Makefile`; under `data/` the recorded run `02-schur-recurrences-verification.json` with its console log `02-schur-recurrences-verification.log`, the coefficient arrays `02-schur-recurrences-sample_polynomials.json`, the requirement pin `02-schur-recurrences-requirements.txt` and a hand-written layout record of the delivered PDF, `02-schur-recurrences-pdf_checks.json`; and in the report root its research-status record `02-schur-recurrences-STATUS.md` and its source audit `02-schur-recurrences-sources.md`. These files are byte-identical to the delivery, so their text still uses the delivery names. The manuscript, its PDF and its README are not shipped; they survive in the arrival commit. The report README maps every delivery name to its shipped path, and Appendix D explains why the shipped Makefile must not be run.

The manuscript contributes, in the order printed here: a proof of Part I’s Conjecture 10.1 for all $k \geq 1$ and $n \geq 0$ (Theorem 12.1), through a Toeplitz factorization with a boundary parameter (Theorem 14.1); a second derivation of Part I’s parity factorization (Proposition 13.1) and of Part I’s shape statements (Corollary 15.1); parity-wise unimodality for odd shifts (Proposition 15.2); a confluent spectral theorem for rectangular Schur polynomials with explicit nonzero leading coefficients (Theorem 16.1); the exact minimal recurrence of Cigler’s family (Theorem 12.2 and Corollary 17.1); the degree and reciprocity sign of the generating numerator (Lemma 18.1,

Theorem 18.2), with a correction of the sign printed in Cigler’s equation (85); the period-four signed normalization (Theorem 18.3); a sharp fixed-parameter convergence equivalent (Theorem 19.1); the reduced denominators at every real parameter (Theorem 19.2); its finite checks, proof-dependency audit and formalization plan (Section 20); six research topics (Section 21); a conclusion; and small-shift examples and package notes (Appendices C and D). Every result, proof, example, remark, limitation and question of the manuscript is printed.

Where the merge had to choose.

- *Duplicated material.* Section 13 re-derives Part I’s Sections 2–4 (Gram identities in the bases P_i , R_i and block elimination, in place of Part I’s tridiagonal principal minors); Corollary 15.1 recovers the shape statements of Theorem 1.1 through tableaux and is not claimed as new; equation (19.4) restates (1.6); Theorem 19.1 sharpens Proposition 9.2 for real $0 < t < 1$; the definitions (12.1)–(12.3) are those of (1.1), (1.2) and (1.3). The manuscript re-derives this material so that its argument does not rest on Part I; it is printed in both parts, and Part II’s proofs are second routes.
- *Numbering.* The manuscript’s Section n is Section $n + 11$ here, and its Theorem, Lemma, Proposition, Corollary and equation $n.m$ are $(n + 11).m$; its Appendices A and B are Appendices C and D. Its title block, abstract and status box are printed in Section 12.2, and this subsection and the next two precede its Section 1.1, which is Section 12.4 here. Its table of contents is merged into the report’s.
- *Names.* Every label of Part II carries the prefix **rs:**. The manuscript’s macro for $\mathcal{H}$ is replaced by Part I’s $\backslash\mathcal{H}$; both print the same glyph for the same object. Its bibliography key CK (Ciucu–Krattenthaler) is printed as CiucuK, because Part I’s CK is Cigler–Krattenthaler, a different paper; its entry for Part I is replaced by internal references, and its entry for Cigler’s paper is merged with Part I’s (the same version 3). No printed symbol is renamed and no normalization is changed. Symbols that mean different things in the two parts are listed in Section 12.3.
- *Added text.* Dated notes were added in Part I (title, abstract, Section 1.3, after Propositions 9.1 and 9.2, at the end of Section 10 and in Section 11.3), and four bracketed notes in Part II: on the pinned paths (Section 12.4), on Cigler’s printed page 24 (after (18.13); an observation of this merge, not of the manuscript), on the shipped names and a rerun (Section 20), and on the delivered commands (Appendix D). All are marked [Added 29 September 2026, batch 56: ...].

12.2 The manuscript’s title, abstract and status

The manuscript’s title block reads *Rectangular Schur Structure and Sharp Recurrences for Cigler’s Hankel Polynomials*, with the subtitle *A proof of the repository’s Schur conjecture, exact pole multiplicities, and corrected generating-function reciprocity*, the author line “Research draft prepared for Vladimir Reshetnikov” and the date 29 September 2026. Its abstract:

A report in the ProveIt repository leaves explicitly unproved a rectangular-Schur formula for normalized Hankel determinants of Cigler’s third polynomial extension of the middle binomial coefficients. We prove that formula for all shifts and orders. Writing $a = \lfloor k/2 \rfloor$ and $b = \lfloor (k - 1)/2 \rfloor$, the identity is

$$\mathcal{H}_k(n; t) = s_{(n^a)}((1)^{[a]}, t, (t^2)^{[b]}).$$

An elementary Toeplitz factorization supplies the missing bridge from the repository’s parity factors to the Schur determinant. We then prove a confluent spectral theorem for rectangular Schur polynomials: each allocation of rows among repeated-variable clusters has an explicit nonzero leading coefficient. For Cigler’s family this determines the *minimal* generic recurrence,

with characteristic polynomial

$$\prod_{q=0}^{k-1} (E - t^q)^{1 + \lfloor q(k-1-q)/2 \rfloor}.$$

We determine the exact generating-numerator degree and reciprocity sign, correct a sign in the printed Conjecture 18, obtain a sharp fixed-parameter convergence equivalent, and classify the reduced denominators at every real parameter. Exact finite checks and a focused further-research program accompany the all-index arguments.

Its status box:

Research status. The conclusions are mathematical proof claims, with the arguments supplied below. They have not been independently refereed or checked in Lean or Rocq. The unresolved repository formula is identified at a fixed commit; the general Schur, character, and confluence machinery is classical and is credited. No exhaustive historical-priority claim is made. The literal entirety of Cigler's printed Conjecture 18 is *not* asserted: one of its signs is false, and the corrected statements are given precisely.

12.3 Notation shared with Part I

Part II keeps the manuscript's notation. The two parts agree on $c_r(t)$, $D_k(n; t)$, $\mathcal{H}_k(n; t)$, a , b , $s = 1 + t$, u_r , v_r , the Laurent polynomial $Q(z) = Q_t(z) = (1 + tz)(1 + t/z)$, h_j , $s_{(n^a)}$ and the forward shift E . The following letters mean different things in the two parts; the right-hand column gives the tempting false reading where one exists.

Symbol	Part I	Part II	Do not read as
$\mathcal{U}, \mathcal{V}$	$\mathcal{U}_a(m; t) = \det(M_1^a[m])$, $\mathcal{V}_a(m; t) = \det(M_0^a[m])$: power a first, size m second	$\mathcal{U}_m(q)$, $\mathcal{V}_m(q)$: Gram determinants of size m of a symbol q , (13.1)–(13.2); $U_h(m) = \mathcal{U}_m(Q_t^h)$, $V_h(m) = \mathcal{V}_m(Q_t^h)$	$\mathcal{U}_m(q)$ as $\mathcal{U}_a(m; t)$ with $a = m$. The true dictionary is $\mathcal{U}_a(m; t) =$ $U_a(m) = \mathcal{U}_m(Q_t^a)$ and $\mathcal{V}_a(m; t) = V_a(m) =$ $\mathcal{V}_m(Q_t^a)$
italic U, V	$U_a(r)$, $V_a(r)$: unnormalized moment matrices in the proof of Theorem 4.1; U_a^∞ , V_a^∞ : limit series (7.2)	$U_h(m)$, $V_h(m)$: normalized determinants; the unnormalized matrices are U_r'	Part II's $U_h(m)$ as a matrix
$\mathcal{F}_k, F_k$	$\mathcal{F}_k(t)$ (1.4)	$F_k(t)$ (19.2), the same function	—
d	d_k : coarse exponent bound of Proposition 9.1; $d_k(n; t) = t^{-\binom{n}{2}} D_k(n; t)$	$d_k(n, t)$: the same unadjusted normalization (Cigler's); d_r , d_q , $d_\pm$: polynomial degrees in n	d_q as Part I's d_k
N	$N_{a,m}^\delta(t)$: numerator determinants (6.2)	N_k : order of the minimal recurrence (12.8)	—
δ	$\delta \in \{0, 1\}$ in $N_{a,m}^\delta$	$\delta_k = \lfloor k^2/4 \rfloor$ (18.1)	—
C	$C_j(y)$: Chebyshev-type polynomials in the proof of Theorem 6.2; $C_{k,r}$: constant of Proposition 9.2	$C_k = \binom{k}{2}$; $C = K - A$ in Section 16; $C_{2m}(u)$, $C_{2m+1}(u)$: normalizing scalars	—
B	$B = 1 + t^2$	$B(m, r)$: factorial ratio (16.3)	—
M	M_ε : tridiagonal matrices (2.4)	$M = k - 1$ (12.4)	—

Symbol	Part I	Part II	Do not read as
x, z	z : Laurent variable; x : generating variable of Proposition 9.1	z : Laurent variable (Sections 13–15) and generating variable (Sections 12 and 16–19, Appendix C); $x = z + z^{-1}$ in Sections 13–14	Cigler’s generating variable x (quoted in the note after (18.13)) as Part II’s $x = z + z^{-1}$
e	e_j : standard basis vectors	$e_{k,q}$: pole multiplicities; $e_j(X)$: elementary symmetric polynomials; (e_j) : bialternant exponents	—
h	— (Part I indexes the parity factors by a)	index of the parity factors in Section 13; derivative orders in Section 16	h as the complete homogeneous h_j
$A, \mathcal{A}$	$A[m]$: principal submatrix; A_j^ε : monic polynomials	A : height of the rectangle in Section 16; $\mathcal{A}_k$: generating numerator	—
J	$J = \text{diag}(1, -1, 1, \dots)$	the involution $Jf(z) = zf(z^{-1})$	—
L	the moment functional L	L_r : leading coefficients (16.5); LC: leading coefficient	—

12.4 Provenance and what is added

The source problem is Section 6 of Johann Cigler’s paper [1], version 3, dated 30 December 2021. Its moment family is denoted $c_n(t)$; it must not be confused with the paper’s other families $a_n(t)$ and $b_n(t)$. In ProveIt, the report *A parity factorization and sharp stabilization for Cigler’s Hankel polynomials* (Part I of this report) proves a parity factorization, coefficient positivity, reciprocity, stable edge coefficients, and a first-defect formula. Its Section 10 explicitly records an experimental rectangular-Schur identity as a conjecture, not a proof dependency. Its recurrence bound is explicitly nonminimal.

The inspected repository commit is

e9c09b12549ea4e3bea7cebc52b450b761f6f581.

The source article’s Git blob identifier is f33543af1f846c24511dc3004fb3542c2a20c50f. The selected conjecture belongs to the directory `cigler-conjecture-16-parity`; full paths are in the bibliography and the package’s source audit. The related repository report on *Conjecture 8* [5] uses a Laurent–Gram/Schur method for the *different* $b_n(t)$ moments. It is relevant methodological background, not a proof of the identity selected here.

[Added 29 September 2026, batch 56: in this report the bibliography entry for Part I is replaced by internal references; the pinned commit, blob identifiers and paths are recorded in Section 12.1, and the package’s source audit is shipped as `02-schur-recurrences-sources.md`.]

Our principal addition is the missing all-index Schur identity. Beyond that identity, the essential new calculation relative to the inspected reports is a noncancellation theorem for confluent spectral terms. It proves exact pole multiplicities, rather than merely giving a denominator that might cancel. The resulting numerator reciprocity also distinguishes a genuine statement from a sign error in the original conjecture.

Rectangular Schur factorizations at reciprocal alphabets are established mathematics; see Ciucu–Krattenthaler [6]. Recurrence existence for stretched Schur polynomials is also established; see Alexandersson [7]. Repeated-variable bialternants are treated explicitly by González-Serrano–Maximenko [8]. We use and explain these kinds of classical tools, rather than advertising

their existence as a discovery. The specific identity, exact multiplicities, constants, and source corrections are the objects proved here. A bounded literature search does not certify that no equivalent formulation has appeared elsewhere.

12.5 Definitions

For $r \geq 1$, put

$$c_r(t) = \sum_{j=1}^r \binom{\lfloor (r-1)/2 \rfloor}{\lfloor (j-1)/2 \rfloor} \binom{\lfloor r/2 \rfloor}{\lfloor j/2 \rfloor} t^{j-1}, \quad c_0(t) = 1. \quad (12.1)$$

A binomial coefficient is zero when its lower index is outside the usual range. These are Cigler's equation (6), with the zero $j = 0$ summand omitted. For a positive shift k and order $n \geq 0$, define

$$D_k(n; t) = \det(c_{k+i+j}(t))_{0 \leq i, j < n}, \quad (12.2)$$

$$\mathcal{H}_k(n; t) = (-1)^k \binom{n}{2} t^{-\binom{n}{2}} D_k(n; t). \quad (12.3)$$

Empty determinants are one. Initially (12.3) is read in $\mathbb{Q}(t)$; polynomiality and specialization at $t = 0$ are conclusions. Throughout, write

$$a = \lfloor k/2 \rfloor, \quad b = \lfloor (k-1)/2 \rfloor, \quad M = k-1, \quad \mathcal{X}_k = ((1)^{[a]}, t, (t^2)^{[b]}). \quad (12.4)$$

Here $x^{[r]}$ means r copies of x , including no copies when $r = 0$. The alphabet $\mathcal{X}_k$ has exactly k entries.

For an alphabet X , define $h_j(X)$ and $e_j(X)$ by

$$\sum_{j \geq 0} h_j(X) w^j = \prod_{x \in X} (1 - xw)^{-1}, \quad \sum_{j \geq 0} e_j(X) w^j = \prod_{x \in X} (1 + xw). \quad (12.5)$$

Subscripts below zero give zero. The Schur polynomial of the rectangle with a rows of length n has the classical Jacobi–Trudi descriptions

$$s_{(n^a)}(X) = \det(h_{n-i+j}(X))_{1 \leq i, j \leq a} = \det(e_{a-i+j}(X))_{1 \leq i, j \leq n}. \quad (12.6)$$

These identities, the tableau interpretation, and the bialternant formula are standard Schur identities [9, 8]. Empty rectangles have value one. They are the only symmetric-function identifications needed for the main determinant proof.

12.6 Main statements

Theorem 12.1 (The repository's rectangular-Schur conjecture). *For all $k \geq 1$ and $n \geq 0$, there is an identity in $\mathbb{Z}[t]$*

$$\boxed{D_k(n; t) = (-1)^k \binom{n}{2} t^{\binom{n}{2}} s_{(n^a)}(\mathcal{X}_k).} \quad (12.7)$$

Consequently $\mathcal{H}_k(n; t) = s_{(n^a)}(\mathcal{X}_k)$, with no exceptional value of t .

Theorem 12.2 (Exact generic recurrence). *For $k \geq 1$, set*

$$e_{k,q} = 1 + \left\lfloor \frac{q(k-1-q)}{2} \right\rfloor, \quad N_k = \sum_{q=0}^{k-1} e_{k,q}. \quad (12.8)$$

Over $\mathbb{Q}(t)$, the reduced denominator of $\mathcal{G}_k(z, t) = \sum_{n \geq 0} \mathcal{H}_k(n; t) z^n$, normalized to have constant term one, is exactly

$$\boxed{\mathcal{D}_k(z, t) = \prod_{q=0}^{k-1} (1 - t^q z)^{e_{k,q}}.} \quad (12.9)$$

Equivalently, the monic minimal constant-coefficient recurrence operator is $\prod_q (E - t^q)^{e_{k,q}}$, where $Ef(n) = f(n+1)$. There is no cancellation in (12.9). The same statement holds at each real $t \notin \{0, 1, -1\}$.

The precise numerator and its sign are given in Theorem 18.2; the period-four signed normalization used by Cigler is treated in Theorem 18.3. Theorem 19.2 completes the real-parameter classification. The general mechanism that makes the recurrences minimal is Theorem 16.1, whose explicit leading coefficient also gives the sharp asymptotic equivalent in Theorem 19.1.

13 Laurent moments and the four parity factors

This section rederives the relevant part of Part I (Sections 2–4), so the new argument does not depend on trusting an unverified report.

13.1 Two elementary Gram identities

Let $q(z) = q(z^{-1})$ be a symmetric Laurent polynomial, and put $q_j = [z^j]q(z)$, so $q_{-j} = q_j$. Define

$$\mathcal{U}_m(q) = \det(q_{i-j} + q_{i+j+1})_{0 \leq i, j < m}, \quad (13.1)$$

$$\mathcal{V}_m(q) = \det(q_{i-j} - q_{i+j+2})_{0 \leq i, j < m}. \quad (13.2)$$

Both empty determinants are one. Set $x = z + z^{-1}$ and $y = z - z^{-1}$. For $i \geq 0$, the Laurent expressions

$$P_i(x) = \frac{z^{i+1} + z^{-i}}{z + 1}, \quad R_i(x) = \frac{z^{i+1} - z^{-i-1}}{z - z^{-1}} \quad (13.3)$$

are monic polynomials in x of degree i . Indeed, $P_0 = R_0 = 1$, $P_1 = x - 1$, $R_1 = x$, and each sequence satisfies $F_{i+1} = xF_i - F_{i-1}$. A direct multiplication gives

$$\frac{1}{2} \text{CT}_z q(z)(x+2)P_i(x)P_j(x) = q_{i-j} + q_{i+j+1}, \quad (13.4)$$

$$\frac{1}{2} \text{CT}_z q(z)(4-x^2)R_i(x)R_j(x) = q_{i-j} - q_{i+j+2}. \quad (13.5)$$

For example, $(x+2) = (z+1)^2/z$, and the product on the first line has four monomials before the constant term is taken. Symmetry of q merges them in pairs. For the second line use $4-x^2 = -y^2$. Thus $\mathcal{U}_m$ and $\mathcal{V}_m$ are ordinary polynomial Gram determinants for the two displayed constant-term weights. Monicity means that replacing P_i or R_i by x^i does not change either determinant.

13.2 The moments

Put

$$Q_t(z) = 1 + t^2 + t(z + z^{-1}) = (1 + tz)(1 + t/z), \quad s = 1 + t, \quad (13.6)$$

and define

$$u_r = \text{CT}_z (1+z)Q_t(z)^r, \quad v_r = \text{CT}_z (1-z^2)Q_t(z)^r. \quad (13.7)$$

Symmetry allows the alternative weights $(x+2)/2$ and $(4-x^2)/2$, respectively. Expansion of the two binomials in Q_t^r yields

$$u_r = \sum_{j=0}^r \binom{r}{j}^2 t^{2j} + \sum_{j=0}^{r-1} \binom{r}{j} \binom{r}{j+1} t^{2j+1}. \quad (13.8)$$

Separating parity in (12.1) and applying Pascal's identity gives

$$c_{2r+1} = u_r, \quad c_{2r+2} = su_r. \quad (13.9)$$

Moreover, $(s^2 - Q_t)(1 + z) = t(1 + z - z^2 - z^{-1})$. The constant terms of zQ_t^r and $z^{-1}Q_t^r$ agree, so

$$s^2 u_r - u_{r+1} = t v_r. \quad (13.10)$$

For short, write

$$U_h(m) = \mathcal{U}_m(Q_t^h), \quad V_h(m) = \mathcal{V}_m(Q_t^h) \quad (h, m \geq 0). \quad (13.11)$$

Since $Q_t = 1 + t^2 + tx$, its i th power has leading coefficient t^i as a polynomial in x . Applying the monic basis changes in (13.4)–(13.5) therefore gives

$$\det(u_{h+i+j})_{0 \leq i, j < m} = t^{m(m-1)} U_h(m), \quad (13.12)$$

$$\det(v_{h+i+j})_{0 \leq i, j < m} = t^{m(m-1)} V_h(m). \quad (13.13)$$

The exponent is twice $\sum_{i=0}^{m-1} i$. These are polynomial identities, not assertions obtained by substituting zero into an undefined quotient.

13.3 Block elimination

Proposition 13.1 (Parity formulas). *For $h, m \geq 0$,*

$$\mathcal{H}_{2h+1}(2m; t) = U_h(m) V_h(m), \quad (13.14)$$

$$\mathcal{H}_{2h+1}(2m+1; t) = U_h(m+1) V_h(m). \quad (13.15)$$

For $h \geq 1$ and $m \geq 0$,

$$\mathcal{H}_{2h}(2m; t) = U_h(m) V_{h-1}(m), \quad (13.16)$$

$$\mathcal{H}_{2h}(2m+1; t) = (1+t) U_{h-1}(m+1) V_h(m). \quad (13.17)$$

Proof. Reorder both rows and columns of the Hankel matrix with the even indices first. The permutation signs cancel. For an odd shift $2h+1$, the four blocks have entries u_{h+i+j} , su_{h+i+j} , su_{h+i+j} , and $u_{h+i+j+1}$. Subtract s times even row i from odd row i . The lower-left block becomes zero; the lower-right block becomes $-t(v_{h+i+j})$. Its size is m for both orders $2m$ and $2m+1$; the upper-left block has size m or $m+1$. Equations (13.12)–(13.13) give the normalization powers $2m^2 - m = \binom{2m}{2}$ and $2m^2 + m = \binom{2m+1}{2}$. The remaining sign $(-1)^m$ equals $(-1)^{\binom{n}{2}}$.

For an even shift $2h$ and even order, abbreviate the unnormalized moment matrices by $U'_r = (u_{r+i+j})_{0 \leq i, j < m}$. The reordered matrix is

$$\begin{pmatrix} sU'_{h-1} & U'_h \\ U'_h & sU'_h \end{pmatrix}.$$

Over $\mathbb{Q}(t)$, subtract s^{-1} times each lower row from the corresponding upper row. The upper-right block vanishes and the upper-left becomes $(t/s)(v_{h-1+i+j})$. The determinant is consequently $t^m \det(v_{h-1+i+j}) \det(u_{h+i+j})$.

For odd order $2m+1$, subtract s^{-1} times even row $i+1$ from odd row i , for $0 \leq i < m$. The lower-left block vanishes; the lower-right is $(t/s)(v_{h+i+j})$, and the upper-left is $s(u_{h-1+i+j})$ of size $m+1$. The remaining scalar factor is st^m . The same two normalization powers give (13.16) and (13.17). All final expressions are polynomials, so the temporary use of s^{-1} introduces no exception at $t = -1$. $\square$

14 The missing Toeplitz factorization

For a Laurent polynomial f , let

$$T_n(f) = \det([z^{j-i}]f(z))_{0 \leq i, j < n}. \quad (14.1)$$

The following two formulas are the bridge needed by the repository's experimental conjecture. We prove them by elementary finite-dimensional linear algebra, rather than requiring a classical-group character factorization as a black box.

Theorem 14.1 (Two factorizations with a boundary parameter). *For every symmetric Laurent polynomial q and every $m \geq 0$,*

$$T_{2m}((1+z)(1+u/z)q) = \mathcal{U}_m(Q_u q) \mathcal{V}_m(q), \quad (14.2)$$

$$T_{2m+1}((1+z)(1+u/z)q) = (1+u) \mathcal{U}_{m+1}(q) \mathcal{V}_m(Q_u q), \quad (14.3)$$

where $Q_u = 1 + u^2 + u(z + z^{-1})$. These are polynomial identities in u and the coefficients of q . In particular,

$$T_n((1+z)q) = \mathcal{U}_{\lceil n/2 \rceil}(q) \mathcal{V}_{\lfloor n/2 \rfloor}(q). \quad (14.4)$$

Proof. It is enough first to work over a characteristic-zero rational-function field, with $u \neq 0, 1, -1$ and independent coefficients for q . For Laurent polynomials define

$$\langle f, g \rangle_q = \text{CT}_z q(z) f(z) g(z^{-1}).$$

This form is symmetric. In consecutive monomial bases for a space V , the pairing of the left vectors $(1+z)z^i$ with the right vectors $(1+uz)z^j$ is the Toeplitz matrix of $(1+z)(1+u/z)q$. Translating all exponents by the same integer leaves its entries unchanged. We choose centered exponent ranges to expose the symmetry.

Even order. For $n = 2m$, take

$$V = \text{span}\{z^{-m}, \dots, z^{m-1}\}, \quad W = \text{span}\{z^{-m}, \dots, z^m\}.$$

Every element of W has a unique decomposition $p(x) + yr(x)$ with $\deg p \leq m$ and $\deg r \leq m-1$. The left space $(1+z)V$ is exactly the subspace of W vanishing at $z = -1$. Thus it has basis

$$(x+2)x^i \ (0 \leq i < m), \quad yx^i \ (0 \leq i < m).$$

Put

$$z_0 = -u^{-1}, \quad x_0 = -(u + u^{-1}), \quad y_0 = u - u^{-1}.$$

The right space $(1+uz)V$ is the kernel of evaluation at z_0 and has basis

$$(x - x_0)x^j \ (0 \leq j < m), \quad yx^j - y_0x_0^j \ (0 \leq j < m).$$

Linear independence follows separately from the symmetric and antisymmetric parts; their total dimensions are $2m$. Symmetric Laurent polynomials are orthogonal to antisymmetric ones for $\langle \cdot, \cdot \rangle_q$. The lower-left block is therefore zero. The two diagonal blocks are the moment matrices for the weights

$$(x+2)(x-x_0)q \quad \text{and} \quad (4-x^2)q.$$

Because $u(x-x_0) = Q_u$, their determinants are, up to factors depending only on u and m , $\mathcal{U}_m(Q_u q)$ and $\mathcal{V}_m(q)$ by (13.4)–(13.5).

Every basis change just made is independent of q . Consequently there is a nonzero rational function $C_{2m}(u)$, independent of q , such that the left side of (14.2) equals $C_{2m}(u) \mathcal{U}_m(Q_u q) \mathcal{V}_m(q)$. We determine this scalar below. This observation avoids hiding any powers of two, powers of u , or basis orientation signs.

Odd order. For $n = 2m + 1$, take

$$V = \text{span}\{z^{-m}, \dots, z^m\}, \quad W = \text{span}\{z^{-m}, \dots, z^{m+1}\}.$$

The involution $Jf(z) = zf(z^{-1})$ is an isometry of the pairing. Its two eigenspaces are orthogonal, and every element of W is uniquely

$$(1+z)p(x) + (z-1)r(x), \quad \deg p, \deg r \leq m.$$

A basis of the left space is

$$(1+z)x^i \ (0 \leq i \leq m), \quad (z-1)(x+2)x^i \ (0 \leq i < m).$$

This follows by writing the symmetric and antisymmetric parts of V and using $(1+z)y = (z-1)(x+2)$. Set $\gamma = (u-1)/(u+1)$. A basis of the right evaluation kernel is

$$(1+z)x^j + \gamma(z-1)x_0^j \ (0 \leq j \leq m), \quad (z-1)(x-x_0)x^j \ (0 \leq j < m).$$

Indeed $(1+z_0) + \gamma(z_0-1) = 0$, and dimensions and independence again follow from the two eigenspaces. The upper-right block is zero. The diagonal blocks now have weights

$$(x+2)q \quad \text{and} \quad (4-x^2)(x-x_0)q,$$

because $(z-1)(z^{-1}-1) = 2-x$. Their determinants are proportional to $\mathcal{U}_{m+1}(q)\mathcal{V}_m(Q_u q)$, with a scalar $C_{2m+1}(u)$ independent of q .

Normalization and specialization. For $q = 1$, both $\mathcal{U}_m(1)$ and $\mathcal{V}_m(1)$ equal one. Tridiagonal continuant recurrences give

$$\begin{aligned} T_n((1+z)(1+u/z)) &= 1 + u + \dots + u^n, \\ \mathcal{U}_m(Q_u) &= 1 + u + \dots + u^{2m}, \\ \mathcal{V}_m(Q_u) &= 1 + u^2 + \dots + u^{2m}. \end{aligned}$$

For example, all three recurrences follow by expanding the last row; the first has diagonal $1+u$ and off-diagonal product u , while the other two have diagonal $1+u^2$, off-diagonal product u^2 , and, for $\mathcal{U}$, an extra u in the first diagonal position. Their initial values fix the displayed sums. They imply $C_{2m} = 1$ and $C_{2m+1} = 1+u$. Both sides of the asserted identities are polynomials. Equality on the generic parameter field proves them identically, including $u = 0, 1, -1$ and singular choices of q . Substitution $u = 0$ gives (14.4). $\square$

15 Proof of the Schur identity and coefficient consequences

Proof of Theorem 12.1. Write $E_k(w) = \prod_{x \in \mathcal{X}_k} (1+xw)$ and, temporarily over $\mathbb{Q}(t)$, put

$$f_k(z) = t^a z^{-a} E_k(z/t). \quad (15.1)$$

Its coefficient at z^{j-i} is $t^{i-j} e_{a+j-i}(\mathcal{X}_k)$. The row factors t^i and column factors t^{-j} cancel in the determinant, so dual Jacobi–Trudi gives

$$T_n(f_k) = s_{(n^a)}(\mathcal{X}_k). \quad (15.2)$$

The point of the chosen alphabet is the following exact factorization:

$$f_k(z) = \begin{cases} (1+z)Q_t(z)^a, & k = 2a+1, \\ (1+z)(1+t/z)Q_t(z)^{a-1}, & k = 2a. \end{cases} \quad (15.3)$$

For instance, each paired factor contributes $tz^{-1}(1+z/t)(1+tz) = Q_t(z)$. For odd k , (14.4) yields exactly (13.14)–(13.15). For even k , use (14.2)–(14.3) with $u = t$ and $q = Q_t^{a-1}$; these yield (13.16)–(13.17). Thus $T_n(f_k) = \mathcal{H}_k(n; t)$ in all four cases. The resulting equality (12.7) is between polynomials in $\mathbb{Z}[t]$, so it holds there and at every specialization. The case $k = 1$ gives the empty-height Schur polynomial one. $\square$

15.1 A tableau model and reciprocal coefficients

The identity says that $\mathcal{H}_k(n; t)$ counts semistandard tableaux of shape (n^a) , with letters in an ordered alphabet of k symbols whose weights are 0 for the first a letters, 1 for the next, and 2 for the last b . The exponent of t is the sum of those weights. Thus the normalization in (12.3), including its sign, turns the original Hankel determinant into a genuine nonnegative generating polynomial.

Corollary 15.1 (Previously known shape properties, recovered). *The polynomial $\mathcal{H}_k(n; t)$ has exact degree $(k-1)n$, constant and leading coefficients one, positive coefficients at every intermediate degree, and*

$$\mathcal{H}_k(n; t^{-1}) = t^{-(k-1)n} \mathcal{H}_k(n; t). \quad (15.4)$$

Proof. The all-zero-weight tableau is unique: every column must contain the first a letters in order. Thus the constant term is one. For odd k , the inverse alphabet is a permutation of $t^{-2}\mathcal{X}_k$. Homogeneity of Schur degree an proves (15.4). For even $k = 2a$, use the rectangular complement identity

$$s_{(n^{K-A})}(X) = \left(\prod X\right)^n s_{(n^A)}(X^{-1}) \quad (|X| = K). \quad (15.5)$$

It follows immediately by reversing the exponent columns in the bialternant, or by complementing columns of a rectangular tableau. Here $A = K - A = a$ and $\prod \mathcal{X}_k = t^{k-1}$. This proves reciprocity and the assertion about the top degree and coefficient.

For full support, regard a column as an increasing a -subset of $\{1, \dots, k\}$. There is a saturated componentwise chain from the first a letters to the last a letters, increasing a single letter by one at each step. Each step changes the column weight by zero or one, so the chain contains columns of all weights $0, 1, \dots, k-1$. For a desired total weight between 0 and $(k-1)n$, use some initial zero-weight columns, at most one intermediate-weight column from this chain, and some final maximum-weight columns. The columns are componentwise increasing, hence form a semistandard tableau. The empty cases are immediate. $\square$

These shape properties are not claimed as new relative to Part I (Theorem 1.1). The Schur identity gives a unified interpretation and an alternative proof. Ordinary unimodality is false: already

$$\mathcal{H}_4(2; t) = 1 + 2t + 5t^2 + 4t^3 + 5t^4 + 2t^5 + t^6.$$

A different, weaker coefficient inequality does follow for odd shifts.

Proposition 15.2 (Parity-wise unimodality for odd shifts). *If $k = 2a + 1$ and $\mathcal{H}_k(n; t) = \sum_j c_j t^j$, then $c_j = c_{2an-j}$ and $c_j \leq c_{j+2}$ whenever $0 \leq j \leq an - 1$. Thus the coefficients in each fixed parity class are unimodal.*

Proof. The eigenvalue alphabet $((t^{-1})^{[a]}, 1, (t)^{[a]})$ is the character alphabet of the $\mathrm{SL}_2(\mathbb{C})$ module $(\mathbb{C}^2)^{\oplus a} \oplus \mathbb{C}$. The classical Schur-module character theorem [9] identifies $t^{-an} \mathcal{H}_{2a+1}(n; t)$ with the character of the Schur functor of shape (n^a) on that module. Finite-dimensional SL_2 modules in characteristic zero are sums of irreducibles, whose characters are $t^r + t^{r-2} + \dots + t^{-r}$. Each such string is symmetric and has nondecreasing coefficients in steps of two toward the center. Their sum has the same properties. The representation-theoretic facts are imported classical results; they are not needed elsewhere in this article. $\square$

16 A confluent spectral theorem for rectangles

The next theorem separates recurrence existence from recurrence minimality. A determinant expansion easily produces exponential terms, but cancellation can remove their highest powers

of n . We identify a unique extremal confluent minor and evaluate its leading coefficient, which prevents that cancellation.

16.1 Statement with an explicit leading coefficient

Let $\xi_1, \dots, \xi_\ell$ be distinct nonzero elements of a field of characteristic zero. Give them positive multiplicities $m_1, \dots, m_\ell$, put $K = \sum_i m_i$, choose $1 \leq A < K$, and put $C = K - A$. For each allocation

$$0 \leq r_i \leq m_i, \quad \sum_i r_i = A, \quad (16.1)$$

define

$$s_i = m_i - r_i, \quad \lambda_r = \prod_i \xi_i^{r_i}, \quad d_r = \sum_i r_i s_i, \quad (16.2)$$

and use the factorial ratio

$$B(m, r) = \frac{\prod_{h=0}^{r-1} h!}{\prod_{h=m-r}^{m-1} h!}. \quad (16.3)$$

An empty product is one, so $B(m, 0) = B(m, m) = 1$.

Theorem 16.1 (Cluster allocation and exact polynomial degree). *There are polynomials $P_r(n)$ such that, for every $n \geq 0$,*

$$s_{(n^A)}((\xi_1)^{[m_1]}, \dots, (\xi_\ell)^{[m_\ell]}) = \sum_r P_r(n) \lambda_r^n. \quad (16.4)$$

The canonical polynomials obtained by the confluent Laplace expansion have exact degree d_r . Their leading coefficients are

$$L_r = (-1)^{\sum_{i < j} r_i s_j} \frac{\prod_i B(m_i, r_i) \xi_i^{C r_i - r_i s_i}}{\prod_{i < j} (\xi_j - \xi_i)^{s_i r_j + r_i s_j}}. \quad (16.5)$$

In particular every L_r is nonzero. If the products λ_r are pairwise distinct, the reduced generating function denominator is exactly

$$\prod_r (1 - \lambda_r z)^{d_r + 1}, \quad (16.6)$$

and the corresponding characteristic polynomial is the minimal recurrence.

Distinctness of the nodes is not the same as distinctness of the products. The first hypothesis suffices for the degree and leading-coefficient statements for the canonical summands. The second is essential to the last assertion: summands with a common product must otherwise be combined.

16.2 The confluent bialternant

For distinct variables $X = (x_1, \dots, x_K)$, the rectangular bialternant is

$$s_{(n^A)}(X) = \frac{\det(x_i^{e_j})_{1 \leq i, j \leq K}}{\prod_{i < j} (x_j - x_i)}, \quad (e_j) = (0, 1, \dots, C-1, n+C, \dots, n+K-1). \quad (16.7)$$

To coalesce m_i variables at ξ_i , replace their rows by the normalized derivative rows of orders $0, 1, \dots, m_i - 1$. Taylor expansion and multilinearity show that the within-cluster Vandermonde factor cancels between numerator and denominator. Thus the numerator matrix has rows (i, h) and entries

$$\binom{e_j}{h} \xi_i^{e_j - h}, \quad 0 \leq h < m_i, \quad (16.8)$$

and its denominator is

$$\Delta_m(\xi) = \prod_{i < j} (\xi_j - \xi_i)^{m_i m_j}. \quad (16.9)$$

Rows are ordered by increasing cluster label, then increasing derivative order. This convention fixes all signs. This limiting identity is a standard repeated-variable bialternant; [8] supplies general treatments. The Taylor argument just described explains the factorial normalization used here.

Expand the numerator along its last A columns. Suppose these columns select r_i rows from cluster i , with derivative-order set R_i . The remaining C columns contain no n . From every selected cluster row in the last block one can extract ξ_i^n , leaving a polynomial in n . Grouping by $(r_1, \dots, r_\ell)$ proves the existence of (16.4). The main issue is its exact degree.

16.3 Degree loss forced by confluent rows

Write $D^h f = f^{(h)}/h!$ and $g_j(x) = x^{C+j}$ for $0 \leq j < A$. Normalized Leibniz expansion gives

$$D^h(x^n g_j(x))|_{x=\xi_i} = \xi_i^n \sum_{p=0}^h \binom{n}{h-p} \xi_i^{p-h} D^p g_j(\xi_i). \quad (16.10)$$

For a fixed cluster, the r_i selected rows are linear combinations of the rows $(D^p g_j(\xi_i))_j$. In a nonzero determinant contribution, the selected p 's in that cluster must be distinct. Cauchy–Binet makes this statement literal: the coefficient matrix is block diagonal by cluster, so each nonzero minor selects exactly r_i derivative rows from cluster i . The sum of their orders is at least $\binom{r_i}{2}$. Hence the degree of the high-column determinant, after removing λ_r^n , is at most

$$\sum_i \left(\sum_{h \in R_i} h - \binom{r_i}{2} \right). \quad (16.11)$$

Among all r_i -subsets of $\{0, \dots, m_i - 1\}$, the sum of orders is uniquely maximized by

$$R_i = \{s_i, s_i + 1, \dots, m_i - 1\}. \quad (16.12)$$

For that set the contribution to (16.11) is $r_i s_i$. Every other row selection has strictly lower total degree. This proves that only one Laplace row selection can contribute to n^{dr} . It remains to show that its coefficient is nonzero and to calculate it.

16.4 Evaluation of the unique leading term

For the choice (16.12), the lowest possible derivative orders in (16.10) are $p = 0, \dots, r_i - 1$. Since $\binom{n}{h-p}$ has leading coefficient $1/(h-p)!$, the local leading coefficient determinant is

$$\det \left(\frac{1}{(s_i + h - p)!} \right)_{0 \leq h, p < r_i} = B(m_i, r_i), \quad (16.13)$$

where a negative factorial is interpreted as giving entry zero. To verify this identity, multiply row h by $(s_i + h)!$. The entry in column p becomes the monic falling-factorial polynomial $(s_i + h)_p$ of degree p . Its determinant is the Vandermonde of $s_i, s_i + 1, \dots, s_i + r_i - 1$, namely $\prod_{h=0}^{r_i-1} h!$. Division by the row factors proves (16.13). The powers of ξ_i in (16.10) supply $\xi_i^{-r_i s_i}$.

The low-column minor uses derivative orders $0, \dots, s_i - 1$ in each cluster, so its determinant is $\prod_{i < j} (\xi_j - \xi_i)^{s_i s_j}$. The high-column derivative matrix with orders $0, \dots, r_i - 1$ is the confluent

matrix of $g_j(x) = x^C x^j$. Multiplication by x^C induces, within each cluster, a lower-triangular derivative change with diagonal ξ_i^C . Its determinant is therefore

$$\left(\prod_i \xi_i^{Cr_i} \right) \prod_{i < j} (\xi_j - \xi_i)^{r_i r_j}.$$

In the full Laplace expansion the selected high rows follow the low rows within each cluster. Moving all low rows before all high rows crosses $r_i s_j$ pairs for each $i < j$, so the sign is $(-1)^{\sum_{i < j} r_i s_j}$. Finally, divide by (16.9). The cross exponent becomes $m_i m_j - s_i s_j - r_i r_j = s_i r_j + r_i s_j$, giving exactly (16.5). None of its factors vanishes. This proves the exact degree.

16.5 Why the recurrence is minimal

For a nonzero polynomial P of degree d and nonzero λ ,

$$(E - \lambda)(P(n)\lambda^n) = \lambda^{n+1}(P(n+1) - P(n)).$$

Thus $(E - \lambda)^{d+1}$ kills this sequence, whereas the d th power does not. If $\mu \neq \lambda$, applying $E - \mu$ preserves its polynomial degree and multiplies its leading coefficient by $\lambda - \mu$. These observations prove independence of the spaces $\{P(n)\lambda^n\}$ for distinct λ : apply operators killing all but one of a putative relation. Equivalently, $\sum_{n \geq 0} P(n)(\lambda z)^n$ has a pole of exact order $d+1$ at $z = \lambda^{-1}$; its top pole coefficient is a nonzero multiple of $d! \text{LC}(P)$. Distinct poles cannot cancel one another. Applying this to (16.4) proves (16.6) and finishes Theorem 16.1.

17 Exact recurrences for Cigler's family

Proof of Theorem 12.2. For $k \geq 2$, specialize Theorem 16.1 to cluster nodes $(1, t, t^2)$ and multiplicities $(a, 1, b)$, omitting a zero-multiplicity cluster. The height is $A = a$. An allocation satisfies $r_1 + r_2 + r_3 = a$ and $r_2 \in \{0, 1\}$. Its product is t^q , where $q = r_2 + 2r_3$. For each feasible q the allocation is unique:

$$r_3 = \lfloor q/2 \rfloor, \quad r_2 = q - 2\lfloor q/2 \rfloor, \quad r_1 = a - \lfloor q/2 \rfloor. \quad (17.1)$$

The feasible values are exactly $0 \leq q \leq k-1$. Uniqueness is crucial: the algebraic relation between the three nodes creates no collision between different allocations because the middle cluster has multiplicity one.

The exact degree is

$$d_q = r_1(a - r_1) + r_3(b - r_3) = \left\lfloor \frac{q(k-1-q)}{2} \right\rfloor. \quad (17.2)$$

For example, if $q = 2h$ the first expression is $h(a + b - 2h)$. If $q = 2h + 1$, it is $(a - h - 1)(h + 1) + h(b - h)$; substituting $b = a$ or $a - 1$ gives the displayed floor in either parity of k . Thus

$$\mathcal{H}_k(n; t) = \sum_{q=0}^{k-1} P_{k,q}(n; t) t^{q^n}, \quad \deg_n P_{k,q} = e_{k,q} - 1, \quad (17.3)$$

with nonzero leading coefficients supplied by (16.5). Over $\mathbb{Q}(t)$ the products t^q are distinct; Theorem 16.1 proves minimality. At a real parameter outside $\{0, 1, -1\}$, the nodes $1, t, t^2$ and the products t^q remain distinct and nonzero, and all the same leading coefficients remain nonzero. The case $k = 1$ is simply the constant sequence one. $\square$

Corollary 17.1 (Exact recurrence orders). *For $a \geq 1$,*

$$N_{2a} = 2a + \frac{a(a-1)(2a-1)}{3}, \quad (17.4)$$

$$N_{2a+1} = 2a + 1 + \frac{2a(a^2-1)}{3}. \quad (17.5)$$

Also $N_1 = 1$.

Proof. Sum (12.8). If $k = 2a$, the product $q(2a-1-q)$ is always even, so there is no rounding. If $k = 2a+1$, subtract $1/2$ for each odd q from $q(2a-q)/2$; there are a such values. The usual sums of integers and squares give the formulas. $\square$

k	Pole multiplicities $(e_{k,0}, \dots, e_{k,k-1})$	N_k	$N_k - k$
1	(1)	1	0
2	(1, 1)	2	0
3	(1, 1, 1)	3	0
4	(1, 2, 2, 1)	6	2
5	(1, 2, 3, 2, 1)	9	4
6	(1, 3, 4, 4, 3, 1)	16	10
7	(1, 3, 5, 5, 5, 3, 1)	23	16
8	(1, 4, 6, 7, 7, 6, 4, 1)	36	28

This is stronger than recurrence existence and stronger than the repository's coarse product of powers of $E^2 - t^{2q}$. The proof identifies every needed root and every needed multiplicity. It also explains why a numerical fit at $t = 1$ would be misleading: that parameter merges all the roots, producing a very different reduced recurrence.

17.1 Exact computation without a large Hankel matrix

For fixed k , Jacobi–Trudi computes $\mathcal{H}_k(n; t)$ as an $a \times a$ determinant, not an $n \times n$ determinant. Its entries are generated by

$$\sum_{j \geq 0} h_j(\mathcal{X}_k) w^j = \frac{1}{(1-w)^a (1-tw)(1-t^2w)^b}. \quad (17.6)$$

For many successive orders, one can instead compute an initial segment and use the exact order- N_k recurrence. For an isolated very large order, recurrence exponentiation gives an alternative whose field-operation count is logarithmic in n for fixed k . None of these statements removes output size or integer bit complexity: an expanded polynomial still has degree $(k-1)n$, and integer specializations may have linearly growing bit length. The polynomial $P_{k,q}(n; t)$ can be obtained directly from the finite confluent Laplace expansion, without guessing it by interpolation.

18 Generating numerators, negative indices, and a corrected sign

Set

$$\delta_k = a(k-a) = \lfloor k^2/4 \rfloor, \quad C_k = \binom{k}{2}, \quad R_k = N_k - k. \quad (18.1)$$

Here δ_k is not a Hankel determinant. It is the degree of the dimension polynomial that appears at $t = 1$.

18.1 Negative-width continuation

The exponential-polynomial expression (17.3) defines $\mathcal{H}_k(n; t)$ for every integer n , over $\mathbb{Q}(t)$. This is an algebraic continuation in the width index, not a combinatorial interpretation of a negative rectangle.

Lemma 18.1 (A zero gap and negative-index reciprocity). *For $k \geq 2$,*

$$\mathcal{H}_k(-j; t) = 0 \quad (1 \leq j \leq k-1), \quad (18.2)$$

$$\mathcal{H}_k(-k-r; t) = (-1)^{\delta_k} t^{-C_k-(k-1)r} \mathcal{H}_k(r; t) \quad (r \geq 0). \quad (18.3)$$

For $k = 1$, the second identity holds and the zero range is empty.

Proof. Use the bialternant (16.7), initially for distinct nonzero variables. Its exponent columns are $0, \dots, K-A-1$ and $n+K-A, \dots, n+K-1$. If $n = -j$ with $1 \leq j \leq K-1$, the two intervals overlap, so two columns coincide. If $n = -K-r$, the second interval is $-A-r, \dots, -r-1$. Move that block before the first block, with sign $(-1)^{A(K-A)}$, then multiply each variable row by x_i^{A+r} . The resulting bialternant is that of $s_{(rK-A)}(X)$. Therefore the continued value is

$$(-1)^{A(K-A)} (\prod X)^{-A-r} s_{(rK-A)}(X) = (-1)^{A(K-A)} (\prod X)^{-A} s_{(rA)}(X^{-1}),$$

where the last step is (15.5). The identities survive confluence. On $\mathcal{X}_k$, one has $(\prod \mathcal{X}_k)^a = t^{C_k}$, and (15.4) gives the remaining factor. Since both the bialternant and the spectral expansion agree for all $n \geq 0$, their polynomial-exponential extensions agree identically. $\square$

18.2 The exact numerator theorem

Theorem 18.2 (Degree and corrected reciprocity). *The reduced generating function has the form*

$$\mathcal{G}_k(z, t) = \frac{\mathcal{A}_k(z, t)}{\mathcal{Q}_k(z, t)}, \quad \mathcal{A}_k \in \mathbb{Z}[t, z], \quad \mathcal{A}_k(0, t) = 1. \quad (18.4)$$

Its numerator has exact z -degree $R_k = N_k - k$, with leading coefficient

$$[z^{R_k}] \mathcal{A}_k = (-1)^{N_k + \delta_k + 1} t^{(k-1)R_k/2}, \quad (18.5)$$

and satisfies

$$t^{(k-1)R_k/2} z^{R_k} \mathcal{A}_k\left(\frac{1}{t^{k-1}z}, t\right) = (-1)^{N_k + \delta_k + 1} \mathcal{A}_k(z, t). \quad (18.6)$$

The sign is $(-1)^{a+1}$ when $k = 2a$ and is $+1$ when k is odd.

Proof. Every exponential-polynomial sequence $F(n)$ has the following expansion of its ordinary rational generating function at infinity:

$$\sum_{n \geq 0} F(n) z^n = - \sum_{j \geq 1} F(-j) z^{-j}. \quad (18.7)$$

This is a statement about expansions of one rational function at two places, not an equality between two convergent series on a common annulus. For $F(n) = \lambda^n$ it is the two expansions of $(1 - \lambda z)^{-1}$. Applying $(z d/dz)^h$ proves it for $n^h \lambda^n$, hence in general.

Lemma 18.1 shows that the first nonzero term at infinity is $-\mathcal{H}_k(-k; t) z^{-k} = (-1)^{\delta_k + 1} t^{-C_k} z^{-k}$. The denominator has degree N_k , so the numerator has degree $N_k - k$. Its leading coefficient is obtained by multiplying that term by the leading coefficient $(-1)^{N_k} t^{\sum_q q e_{k,q}}$ of $\mathcal{Q}_k$. Symmetry

$e_{k,q} = e_{k,k-1-q}$ gives $\sum_q q e_{k,q} = (k-1)N_k/2$, proving (18.5). The recurrence makes the numerator a finite convolution of polynomials in $\mathbb{Z}[t]$, so it lies in $\mathbb{Z}[t, z]$.

Substitute $1/(t^{k-1}z)$ in the expansion at infinity and use (18.3). Since $k(k-1) = 2C_k$, this gives

$$\mathcal{G}_k\left(\frac{1}{t^{k-1}z}, t\right) = (-1)^{\delta_k+1} t^{C_k} z^k \mathcal{G}_k(z, t). \quad (18.8)$$

On the other hand,

$$\mathcal{Q}_k\left(\frac{1}{t^{k-1}z}, t\right) = (-1)^{N_k} t^{-(k-1)N_k/2} z^{-N_k} \mathcal{Q}_k(z, t).$$

Combining these two formulas proves (18.6). For even $k = 2a$, N_k is even and $\delta_k = a^2$. For odd k , N_k is odd and $\delta_k = a(a+1)$ is even. These parity facts give the stated signs. They also show explicitly that all displayed half-exponents of t are integers. $\square$

18.3 What this says about the printed Conjecture 18

For even shift $k = 2a$, Cigler's $d_{2a}(n, t)$ is exactly our $\mathcal{H}_{2a}(n; t)$. The denominator exponent in his equation (84) is

$$aj - \frac{(j-1)(j+2)}{2} = 1 + \frac{j(2a-1-j)}{2} = e_{2a,j}.$$

Thus the present theorem proves that proposed denominator and proves it is minimal. Its numerator degree is

$$R_{2a} = \frac{a(a-1)(2a-1)}{3}.$$

However, the positive-sign reciprocity printed in equation (85) is false when a is even. The corrected sign is $(-1)^{a+1}$. This is not a subtle large-index discrepancy: the same source prints

$$\mathcal{G}_4(z, t) = \frac{1 - t^3 z^2}{(1-z)(1-tz)^2(1-t^2 z^2)(1-t^3 z)}. \quad (18.9)$$

For its numerator, $t^3 z^2 \mathcal{A}_4(1/(t^3 z), t) = t^3 z^2 - 1 = -\mathcal{A}_4(z, t)$. This directly contradicts the positive sign in the printed equation. We therefore prove a corrected formulation, not the literal whole of Conjecture 18. The original source, printed page 24, was inspected directly for this comparison [1].

18.4 The period-four signed normalization

The source uses $d_k(n, t) = t^{-\binom{n}{2}} D_k(n; t)$. For odd k this is $\sigma_n \mathcal{H}_k(n; t)$, where

$$\sigma_n = (-1)^{\binom{n}{2}} = \frac{1-i}{2} i^n + \frac{1+i}{2} (-i)^n. \quad (18.10)$$

The two normalizations must not be interchanged in a generating function.

Theorem 18.3 (Exact signed denominator and numerator). *Let $k = 2a + 1$ and write $\widehat{\mathcal{G}}_k(z, t) = \sum_{n \geq 0} d_k(n, t) z^n$. Its reduced denominator over $\mathbb{Q}(t)$ is exactly*

$$\widehat{\mathcal{D}}_k(z, t) = \prod_{q=0}^{k-1} (1 + t^{2q} z^2)^{e_{k,q}}. \quad (18.11)$$

The numerator $\widehat{\mathcal{A}}_k$ lies in $\mathbb{Z}[t, z]$, has constant term one and exact degree $\widehat{R}_k = 2N_k - k$, and satisfies

$$t^{(k-1)\widehat{R}_k/2} z^{\widehat{R}_k} \widehat{\mathcal{A}}_k\left(\frac{1}{t^{k-1}z}, t\right) = (-1)^a \widehat{\mathcal{A}}_k(z, t). \quad (18.12)$$

Proof. Equation (18.10) changes each spectral root t^q into the two roots it^q and $-it^q$, with nonzero coefficients. They are all distinct over $\mathbb{Q}(t, i)$, so the minimality argument doubles the recurrence order and gives (18.11). The product has rational coefficients, and minimality over the extension implies minimality over $\mathbb{Q}(t)$. For negative indices, $\sigma_{-k-r} = (-1)^{a+1}\sigma_r$. Since δ_k is even, the proof of (18.8) becomes

$$\widehat{\mathcal{G}}_k\left(\frac{1}{t^{k-1}z}, t\right) = (-1)^a t^{C_k} z^k \widehat{\mathcal{G}}_k(z, t).$$

The gap of $k-1$ zero negative indices gives degree $2N_k - k$ as before. The quadratic denominator transforms without a sign, with factor $t^{-(k-1)N_k} z^{-2N_k}$. This proves (18.12) and the remaining assertions. $\square$

As a useful normalization check, the first nontrivial odd shift gives

$$\widehat{\mathcal{G}}_3(z, t) = \frac{1 + (1+t+t^2)z - t(1+t+t^2)z^2 - t^3z^3}{(1+z^2)(1+t^2z^2)(1+t^4z^2)}. \quad (18.13)$$

It has $d_3(1, t) = c_3(t) = 1 + t + t^2$, as the moment definition requires. The cubic term and the sign in (18.13) cannot be omitted. This provides a second direct check on the printed small-shift generating functions, independent of the general numerator proof.

[Added 29 September 2026, batch 56. Observation of this merge, not of the manuscript; it concerns the external source only. The generating function for d_3 printed on p. 24 of [1], with numerator $1 + (1+t)^2x + t^2x^2$ over $(1+x^2)(1+t^2x^2)(1+t^4x^2)$, is incorrect: its coefficient of x is $(1+t)^2$, whereas $d_3(1, t) = c_3(t) = 1 + t + t^2$ (at $t = 2$: 9 against 7). The numerator of (18.13) factors as $(1-tz)(1+(1+t)^2z+t^2z^2)$, so the printed numerator omits the factor $1-tx$. Equation (18.13) agrees with the determinants for $n < 10$ at $t = 2, 3, -5/2$, and so do the printed generating functions for d_1, d_2 and d_4 . In the same source, equation (87), for the shift $2k+1$, has the sign $(-1)^k$ and the argument $1/(t^{2k}x)$ of Theorem 18.3 with $a = k$, but its exponents of x and t , $\rho(k) = \frac{2k(k-1)(2k-1)}{3} - 2(k-1)^2 + 1$ and $k\rho(k)$, hold only with $\rho(k+1) = (4k^3 + 2k + 3)/3 = \widehat{R}_{2k+1}$ in place of $\rho(k)$: for the shift 3 the printed exponent is $\rho(1) = 1$, while the numerator (18.13) has degree 3. So (87) mixes the indices of two shifts; its sign is right. Both readings were checked on a rendering of the printed page of arXiv:2111.14492v3 and by exact computation during the intake.]

19 Sharp convergence and all real specializations

19.1 The leading nonconstant spectral term

Theorem 19.1 (Sharp fixed-parameter convergence equivalent). *Fix $k \geq 2$ and $0 < t < 1$. Then, as $n \rightarrow \infty$,*

$$\frac{1}{(1-t)^a(1-t^2)^{ab}} - \mathcal{H}_k(n; t) \sim \frac{n^{a-1}t^{n+1}}{(a-1)!(1-t)^{b+1}(1-t^2)^{(a-1)b}}. \quad (19.1)$$

The equivalent is uniform for t in each compact subinterval of $(0, 1)$; more precisely, its relative error is $O(1/n)$ on such a subinterval.

Proof. At $q = 0$, the allocation is $(a, 0, 0)$ and $d_0 = 0$. Formula (16.5) consequently gives the whole constant spectral term, not merely a leading approximation:

$$P_{k,0}(n; t) = F_k(t) := \frac{1}{(1-t)^a(1-t^2)^{ab}}. \quad (19.2)$$

At $q = 1$, the allocation is $(a - 1, 1, 0)$, of degree $a - 1$. Here $B(a, a - 1) = 1/(a - 1)!$. Substitution into (16.5), followed by cancellation of the factors $t^2 - t = t(t - 1)$, yields

$$[n^{a-1}]P_{k,1}(n; t) = -\frac{t}{(a-1)!(1-t)^{b+1}(1-t^2)^{(a-1)b}}. \quad (19.3)$$

All remaining terms in (17.3) contain t^{qn} with $q \geq 2$. For fixed k , their polynomial degrees are bounded and their rational coefficients are bounded on any compact subinterval of $(0, 1)$. They are exponentially smaller than $n^{a-1}t^n$ there. The lower coefficients of $P_{k,1}$ give relative error $O(1/n)$; when $a = 1$ there are no such coefficients. Equations (19.2)–(19.3) prove the assertion. $\square$

The repository already identifies F_k as the limiting series and proves the formal first defect

$$\mathcal{H}_k(n; t) = F_k(t) - \binom{n+a}{a-1} t^{n+1} + O(t^{n+2}) \quad \text{in } \mathbb{Q}[[t]] \quad (19.4)$$

for fixed n (Part I, Theorem 1.1). The limit in (19.1) is a different statement: t is fixed and n grows, so all powers beyond t^{n+1} can affect the prefactor. The rational factor in (19.1) accounts for those contributions. In particular, a formal first-defect formula alone would not justify this asymptotic equivalent. For example,

$$F_4(1/2) - \mathcal{H}_4(n; 1/2) \sim \frac{8}{3} n 2^{-n}.$$

No uniform assertion near $t = 1$ is made; the pole collision there requires a different scaling analysis.

19.2 The exceptional real parameters

Theorem 19.2 (Complete real-parameter denominator classification). *For $\sum_{n \geq 0} \mathcal{H}_k(n; t)z^n$, the reduced denominator at every real parameter is as follows, always normalized to constant term one:*

parameter	reduced denominator
$t \notin \{0, 1, -1\}$	$\prod_{q=0}^{k-1} (1 - t^q z)^{e_{k,q}}$
$t = 0$	$1 - z$
$t = 1$	$(1 - z)^{\delta_k + 1}$
$t = -1, k = 2a$	$(1 - z^2)^{a(a-1)+1}$
$t = -1, k = 2a + 1$	$(1 - z)^{a^2+1}(1 + z)^{a^2}.$

(19.5)

All exponents are exact, not just upper bounds. In particular the last line includes $k = 1$ with $a = 0$.

Proof. The generic real case was proved in Theorem 12.2. At zero, the alphabet contains a ones and zeros. Only the tableau using all a one-valued letters in each column contributes, so $\mathcal{H}_k(n; 0) = 1$.

At one, every alphabet entry is one. The confluent bialternant reduces to the elementary dimension product

$$\mathcal{H}_k(n; 1) = \prod_{i=1}^a \prod_{j=a+1}^k \frac{n+j-i}{j-i}. \quad (19.6)$$

One can derive this directly: the determinant of the normalized derivative matrix at one is the Vandermonde of its exponent columns divided by $\prod_{h=0}^{k-1} h!$. The within-block factors cancel,

leaving precisely the cross-block ratios in (19.6). The product is a nonzero polynomial in n of exact degree δ_k . Its generating denominator is therefore $(1 - z)^{\delta_k + 1}$.

At minus one, the alphabet is 1 repeated $k - 1$ times together with a single -1 . For $k \geq 2$, apply Theorem 16.1 with these two nodes. There are only two allocations: the -1 cluster contributes zero or one selected row. The spectral roots are 1 and -1 , and the polynomial degrees are respectively

$$d_+ = a(k - 1 - a) = ab, \quad d_- = (a - 1)(k - a) = (a - 1)(b + 1).$$

Their leading coefficients are nonzero by (16.5). If $k = 2a$, both degrees are $a(a - 1)$; if $k = 2a + 1$, they are a^2 and $a^2 - 1$. This proves the two last lines. For $k = 1$ the sequence is one directly. $\square$

The determinant values at $t = 1$ and the Catalan-related simplifications at $t = -1$ are related to the source's specializations ([1]; Part I, Section 8 and Appendix A). The conclusion here is an exact reduced-recurrence classification at those specializations, including the proof that the surviving poles really occur. It would be incorrect to substitute $t = 1$ into the generic product and call the resulting multiplicity N_k minimal: the actual multiplicity is $\delta_k + 1$.

20 Reproducibility, proof dependencies, and formalization

20.1 What the accompanying program actually checks

The package contains `code/verify.py`, a runnable verifier. It uses fraction-free Bareiss elimination with an exact-division check at every step. The original binomial moments and the Schur side are constructed independently. For the latter, complete homogeneous polynomials are computed from (17.6), then an $a \times a$ Jacobi–Trudi determinant is evaluated. All numerical checks use integers or rational numbers. SymPy is used only for exact polynomial-ring checks.

The run supplied in `data/verification.json` and `data/verification.log` used Python 3.13.5 and SymPy 1.14.0. All of the following checks passed:

Check	Range or method	Cases
Original Hankel vs. Schur	$1 \leq k \leq 12, 0 \leq n \leq 12; t = -3, -2, -1, 1, 2, 3, 5$	1,092
Universal Toeplitz factorization	12 signed symmetric symbols; $0 \leq n \leq 12; u = -3, -2, -1, 0, 1, 2, 3$	1,092
Ordinary recurrence and numerator	$1 \leq k \leq 12; t = -2, 2, 3$; sequence checked through $2N_k + 8$	36
Period-four signed recurrence	Odd k from 1 to 11; $t = -2, 2, 3$	18
Numerator reciprocity coefficients	All coefficients of both tested numerator families	2,052
Confluent leading coefficients	Five cluster-multiplicity families at prime nodes; also Cigler $2 \leq k \leq 10, t = 2$	143
Exceptional real parameters	$1 \leq k \leq 12, t = 0, 1, -1$	36
Exact polynomial Hankel–Schur identity	$1 \leq k \leq 8, 0 \leq n \leq 6$ in $\mathbb{Z}[t]$	56

The polynomial checks also test reciprocity, positive support, and the odd-shift parity-wise coefficient inequalities. Their actual coefficient arrays are saved in `data/sample_polynomials.json`. The universal Toeplitz test uses a reproducible pseudorandom seed, 260929, and includes negative coefficients, so it is not restricted to positive moment forms. The JSON record lists every cluster family and range explicitly.

[Added 29 September 2026, batch 56: in this report the verifier is shipped as `code/02-schur-recurrences-verify.py` and its records as `data/02-schur-recurrences-verification`.

json, data/02-schur-recurrences-verification.log and data/02-schur-recurrences-sample_polynomials.json; code/verify.py and data/verification.json are Part I's files (Section 11.1). Run the verifier with an explicit `-output` outside `data/`: by default it writes `data/verification.json`, and it always writes `sample_polynomials.json` next to its output, so a run without `-output` would overwrite Part I's records. A rerun on a copy on 29 September 2026 (Python 3.13.5, SymPy 1.14.0) passed every check and reproduced the recorded files up to line endings (the script writes CRLF on Windows) and the field `elapsed_seconds`. For the exceptional parameters the recorded run checks that the denominators of Theorem 19.2 annihilate the sequences; their minimality is proved in the text, and an independent Berlekamp–Massey computation during the intake found exactly those minimal denominators for $k \leq 10$ at $t = 0, \pm 1$, and exactly $\mathcal{Q}_k$ for $k \leq 9$ at $t = 2, 3, -2, 1/2$.]

There is a useful independent exact test of (16.5). For distinct spectral roots λ_j with degrees d_j , the leading coefficient of P_j can be extracted directly from the sequence F by

$$\text{LC}(P_j) = \frac{\left[\prod_{\ell \neq j} (E - \lambda_\ell)^{d_\ell+1} (E - \lambda_j)^{d_j} F \right] (0)}{\lambda_j^{d_j} d_j! \prod_{\ell \neq j} (\lambda_j - \lambda_\ell)^{d_\ell+1}}. \quad (20.1)$$

The verifier constructs the left-side sequence from Jacobi–Trudi, applies the operator using exact arithmetic, and compares the result with the factorial product (16.5). This checks the signs, the powers of the cluster nodes, and the cross-difference exponents independently of an implementation of the confluent determinant itself.

Finite checks are not a proof of an infinite family. In particular they cannot establish the all-index minimality theorem or historical novelty. Those depend on the mathematical arguments in Sections 14 and 16. No theorem prover was run for this article.

20.2 Dependency and trust boundaries

The main identity depends on the binomial definition, constant terms, finite-dimensional basis changes, block elimination, and dual Jacobi–Trudi. The exact recurrence additionally uses the bialternant, its Taylor confluence, Cauchy–Binet, and finite differences. The numerator theorem uses only those results and an algebraic expansion at infinity. The exceptional-parameter and asymptotic theorems depend on the same explicit leading coefficients. Only Proposition 15.2 imports Schur-module characters and complete reducibility for SL_2 .

Several points are especially important for an independent audit. The Toeplitz pairing is bilinear, not a positive Hermitian inner product; no positivity assumption is used. The odd-order pairing uses the involution $Jf = zf(z^{-1})$, not the ordinary inversion involution. The normalizing constants in the factorization are explicitly fixed at $q = 1$. In the confluence calculation, normalized derivatives remove hidden factorials, and only the highest derivative-order row sets contribute to the maximal degree. Minimality is asserted only after spectral products have been checked for distinctness. Finally, generating functions for $\mathcal{H}_k$ and d_k are kept separate throughout.

20.3 A finite-algebra formalization plan

A Lean implementation can be divided into three layers. First formalize symmetric Laurent coefficient sequences and the two Gram matrices, using finite support; prove the pairing identities and the parameterized Toeplitz factorization. A formal proof can compute basis-change determinants directly or prove their independence from the symbol and use the normalization

specialization as above. All exceptional parameters should be handled by polynomial identity rather than separate division arguments.

The second layer identifies the Toeplitz matrix with the elementary Jacobi–Trudi matrix. Depending on available symmetric-function support, one can initially define the rectangular Schur object by this determinant and separately prove its equality with the complete-homogeneous determinant. This already yields the Hankel identity without a representation-theoretic library.

The third layer formalizes normalized derivative matrices, the factorial Vandermonde determinant (16.13), and degree bounds for Laplace selections. The unique-maximizer argument is finite combinatorics; it is the most important piece to keep inside the trusted proof boundary. The final recurrence follows from explicit shift-operator lemmas. The Python code may generate examples and candidate identities but is not a trusted replacement for any of these steps. No uncompiled Lean skeleton is included as if it were a completed formalization.

21 Further research questions and topics

The following questions are proposed extensions of the results in this article. Except where the original Cigler conjecture is named explicitly, they are not claims about the status of a separately published conjecture.

21.1 Full coefficient structure in Cigler's Conjecture 17

The spectral polynomials $P_{k,q}(n;t)$ have exact degrees in n , but the present paper does not identify all of their coefficient signs and factorizations after a common denominator is cleared. The numerator polynomials in the source's Conjecture 17 organize related information in a more refined way. A useful next target is to express those polynomials as flagged tableaux or minors with manifest sign, and to determine exactly which of the printed degree and reciprocity claims survive normalization. The confluent Laplace expansion gives a finite formula from which to start; it is not itself a proof of every Conjecture 17 assertion.

[Added 30 September 2026, batch 66: Part III's question in Section 33.6 asks for the reduced numerator at cyclotomic parameters; it is related to this topic, not part of it.]

21.2 Complex roots of unity and spectral cancellation

At a complex root of unity, the roots t^q can coincide even when $1, t, t^2$ are distinct. The exact reduced multiplicity is the degree of a sum of canonical $P_{k,q}$'s in one residue class, plus one, unless that sum vanishes. Determine a closed formula for these multiplicities as a function of k , the root order, and the residue class. The real points $0, 1, -1$ are settled by Theorem 19.2; simply substituting into the generic product does not solve the complex problem. Schur root-of-unity identities and cyclic-sieving methods may give a structural approach, but cancellation must be checked, not assumed absent.

[Added 30 September 2026, batch 66: answered in Part III. Theorem 24.1 gives these multiplicities in closed form as a function of k , the root order $\ell \geq 3$ and the residue class, including empty residue classes and factors that disappear: at most one residue class can lose order relative to the maximum-degree baseline: one order in general, three for $\ell = 4$, $k \equiv 1 \pmod{4}$, $k \geq 9$, and the whole simple factor for $\ell = 4$, $k = 5$. Corollary 31.1 completes the classification at every complex parameter.]

21.3 Coefficient shape for even shifts

Odd shifts have an SL_2 interpretation that proves parity-wise unimodality. Does the analogous step-two inequality hold for every even shift and every order? If not, what is the first structural obstruction, and what weaker inequalities characterize these tableau weights? The ordinary unimodality assertion is already false, so one should not promote a few unimodal examples into a general conjecture. A satisfactory result would explain the difference between the equal outer multiplicities of odd shifts and the unequal outer multiplicities of even shifts.

21.4 A two-parameter Hankel deformation

The Schur alphabet $((1)^{[a]}, u, (v)^{[b]})$ makes sense for independent u, v . The present moment construction lies on $v = u^2$. Find a natural two-parameter moment family whose normalized shifted Hankel determinants realize the full alphabet, or prove that a specified class of finite-period Jacobi recurrences cannot do so. The parameterized Toeplitz factorization offers a starting point, but the compatibility of all shifts within one moment sequence is a genuine additional condition.

21.5 A uniform crossover as the poles merge

Theorem 19.1 applies for fixed $t < 1$, while at $t = 1$ the answer grows like n^{δ_k} . In the scaling $t = e^{-s/n}$, study

$$n^{-\delta_k} \mathcal{H}_k(n; e^{-s/n})$$

for fixed k and s in bounded complex sets. Determine the limiting function, uniform error bounds, and an all-orders expansion. The finite confluent bialternant should allow a systematic derivation, but the separate spectral coefficients become singular and their cancellations must be organized before taking the limit. This is a natural meeting point of the repository's determinant and asymptotic-analysis programs.

21.6 Broader noncollision criteria and certified algorithms

The middle multiplicity one in $(a, 1, b)$ makes $q = r_2 + 2r_3$ determine the whole allocation. For alphabets with more clusters, classify linear relations among their exponents that still yield a unique allocation, or yield a sum whose leading coefficient can be proved nonzero. Such a criterion would turn Theorem 16.1 into a reusable exact minimal-recurrence algorithm for specialized rectangular Schur families. A formalized implementation should output both the recurrence and a certificate for every pole multiplicity, rather than an annihilator alone.

[Added 30 September 2026, batch 66: Part III bears on this topic without answering it. Its Theorem 27.1 gives the first centered subleading coefficient of every canonical confluent mode for arbitrary cluster multiplicities, the tool its proof uses when two allocations of Cigler's alphabet share a product at a root of unity; the case of a middle cluster of multiplicity at least two and certified recurrence generation are its questions in Sections 33.1 and 33.9.]

22 Conclusion

The experimental Schur formula in the inspected ProveIt report admits a finite algebraic proof. Its decisive step is the equality of the original Hankel parity factors with the factors of a particular Toeplitz determinant. The resulting Schur structure supports a sharper conclusion than rationality: all generic recurrence roots and multiplicities are exact, with explicitly nonzero confluent leading coefficients. Negative-width continuation then fixes the numerator degree and

reciprocity sign, including a necessary correction to the original printed conjecture. The same calculation gives a sharp convergence equivalent and all reduced denominators at real parameters. The remaining questions concern finer coefficient structure, complex spectral collisions, and uniform pole-merger asymptotics, rather than an unproved bridge in the central determinant identity.

[Added 30 September 2026, batch 66: the complex spectral collisions are classified in Part III (Theorem 24.1 and Corollary 31.1).]

C Small-shift formulas and independent checks

The first three sign-adjusted sequences are

$$\begin{aligned}\mathcal{H}_1(n; t) &= 1, \\ \mathcal{H}_2(n; t) &= 1 + t + \cdots + t^n, \\ \mathcal{H}_3(n; t) &= h_n(1, t, t^2) = \frac{(1 - t^{n+1})(1 - t^{n+2})}{(1 - t)(1 - t^2)}.\end{aligned}$$

The displayed quotient is interpreted by its polynomial continuation at $t = 1, -1$. Their ordinary generating functions are respectively

$$\frac{1}{1 - z}, \quad \frac{1}{(1 - z)(1 - tz)}, \quad \frac{1}{(1 - z)(1 - tz)(1 - t^2z)}.$$

For shift four, (18.9) has numerator degree two, in agreement with $N_4 - 4 = 2$. At $t = -1$ it becomes

$$\sum_{n \geq 0} \mathcal{H}_4(n; -1) z^n = \frac{1 + z^2}{(1 - z^2)^3},$$

so $\mathcal{H}_4(2m; -1) = (m + 1)^2$ and $\mathcal{H}_4(2m + 1; -1) = 0$. At $t = 1$, the dimension product gives

$$\mathcal{H}_4(n; 1) = \frac{(n + 1)(n + 2)^2(n + 3)}{12}.$$

These two specializations make visible the difference between a generic recurrence, a pole-merger specialization, and a parity-supported sequence.

D Package contents and reproduction

The archive contains the present `article.tex` and its compiled `article.pdf`, the verifier, exact JSON records, a console log, `README.md`, `STATUS.md`, `sources.md`, and a small `Makefile`. To reproduce the checks and the PDF, run

```
python -m pip install -r requirements.txt
python code/verify.py
latexmk -pdf -interaction=nonstopmode -halt-on-error article.tex
```

The recorded run completed all listed checks. Run time is environment-dependent and is not a complexity claim. The package does not include a Lean proof or assert independent referee approval.

[Added 29 September 2026, batch 56: this report ships the verifier, the JSON records (including `data/pdf_checks.json`, a hand-written layout record of the delivered 22-page PDF), the console

log, STATUS.md, sources.md, requirements.txt and the Makefile under the prefixed names listed in Section 12.1. The archive's article.tex, article.pdf and README.md are not shipped: Part II and the report's article.pdf replace them. The commands above are the delivery's and must not be used from this report's root: there `python code/verify.py` runs Part I's verifier, which rewrites Part I's records in data/, and `latexmk` rebuilds the whole report in place. The shipped `code/02-schur-recurrences-Makefile` runs exactly these commands (its `clean` target runs `latexmk -c article.tex`), so it must not be run from the report root, and from any other directory its paths do not resolve. Part II's checks are reproduced, from the report root, by

```
uv run --no-project --with sympy==1.14.0 python \  
    code/02-schur-recurrences-verify.py --output <scratch>/verification.json
```

with <scratch> a directory outside the report.]

Part III. Cyclotomic pole collapse and exact minimal recurrences at complex parameters

Part III was added on 30 September 2026 in batch 66 of ProveIt’s incoming-report intake, from one manuscript written against the pinned state of Parts I and II. It answers Part II’s research question “Complex roots of unity and spectral cancellation” (Section 21.2) completely: it determines the reduced generating denominator and the minimal recurrence of $\mathcal{H}_k(n; t)$ at every complex parameter, where Part II’s Theorems 12.2 and 19.2 cover $\mathbb{Q}(t)$ and the real line. Its Sections 23–34 and Appendices E–G are the manuscript’s Sections 1–12 and Appendices A–C. Parts I and II are unchanged except for dated pointers to this addition, marked [Added 30 September 2026, batch 66: ...].

23 The problem, its source, and the mathematical interface

23.1 Provenance of Part III

Part III is batch-66 manuscript 02, *Cyclotomic Pole Collapse and Exact Minimal Recurrences for Cigler’s Hankel Polynomials: A complete complex-parameter classification and an exceptional fourth-root cancellation law*, an AI-assisted research manuscript dated 30 September 2026 (22 pages, A4, as delivered). It arrived as the archive `ProveIt_Cyclotomic_Pole_Collapse.zip` (inner directory `ProveIt_Cyclotomic_Pole_Collapse/`, main file `article.tex`) in ProveIt commit `9ad899cbe` and was placed in `4fee1cd07`. It was written against ProveIt commit `ff6969475` (in full `ff6969475bad8f61a33268418aaf23d23fcd1fb`), at which this report’s `article.tex` was the Git blob `07c41161857e`: Parts I and II exactly as printed above, since no commit touched this report between Part II’s write commit `3a06260c4` and the placement. Every statement the manuscript makes about “the relevant ProveIt report”, “the inspected repository” or “Part II” therefore refers to Parts I and II of the present report, and the numbers it cites (Section 21.2; Theorems 12.1, 16.1 and 19.2) are this report’s own. The manuscript read the report at the pinned commit through a GitHub connector (its source audit, `03-cyclotomic-poles-SOURCES.md`). Its author line, “Research draft prepared for Vladimir Reshetnikov”, names no assistant.

Its code and records are shipped beside the earlier parts’ files with the prefix `03-cyclotomic-poles-` (Part I’s unprefix files count as `01`, Part II’s `02-schur-recurrences-` files as `02`):

Shipped as	Content
<code>code/03-cyclotomic-poles-verify.py</code>	the exact verifier
<code>code/03-cyclotomic-poles-modular.py</code>	its finite-field helper module
<code>code/03-cyclotomic-poles-Makefile</code>	the delivered Makefile (not to be run)
<code>data/03-cyclotomic-poles-verification.json</code>	the recorded run
<code>data/03-cyclotomic-poles-verification.log</code>	its console log
<code>data/03-cyclotomic-poles-modular_checks.json</code>	the 240 finite-field records
<code>data/03-cyclotomic-poles-requirements.txt</code>	the requirement pin
<code>data/03-cyclotomic-poles-pdf_checks.json</code>	build and layout record of the delivered PDF
<code>03-cyclotomic-poles-CLAIM_STATUS.md</code>	claim status and limitations
<code>03-cyclotomic-poles-SOURCES.md</code>	repository pin and literature audit

These files are byte-identical to the delivery, so their text still uses the delivery names. The manuscript, its PDF and its README are not shipped; they survive in the arrival commit `9ad899cbe`. The report README maps every delivery name to its shipped path, and Appendix G explains why the shipped Makefile must not be run and how to rerun the verifier without touching Part I’s records.

The manuscript contributes, in the order printed here: the complete classification of the reduced denominators and minimal recurrences at every primitive root of unity of order $\ell \geq 3$ (Theorem 24.1), with small examples, among them $\mathcal{Q}_{9,i}(z) = (1 - z)^9(1 + z)^4(1 + z^2)^8$ (24.8); a restatement of Part II's confluent leading-coefficient theorem with its specialization to Cigler's alphabet (Lemma 25.1, (25.7)) and the minimality lemma for exponential polynomials (Lemma 25.2); the concavity restriction to one tied residue (Lemma 26.1) and a mode-by-mode negative-index reflection (Lemma 26.2); a general formula for the first centered subleading coefficient of a canonical confluent rectangular-Schur mode with arbitrary cluster multiplicities (Theorem 27.1); the one-order loss outside the exceptional family (Lemma 28.1, Proposition 28.2); exact fourth-root product formulas for $k = 4m + 1$ (Theorem 29.1); the exact three-order loss in that family (Theorem 30.1); four corollaries (Corollaries 31.1–31.4: every complex parameter, compact fourth-root forms, quasipolynomial periods, and the quadratic recurrence-order law $N_{k,\ell} = \ell(k-1)^2/8 + O_\ell(1)$); its finite checks, dependency audit and formalization route (Section 32); nine research questions (Section 33); a conclusion; an elementary beta-moment determinant (Appendix E); an integer exponent algorithm (Appendix F); and package notes (Appendix G). Every result, proof, example, remark, limitation and question of the manuscript is printed.

Relation to other reports. The report `shifted-catalan-hankel-polynomials` in this directory classifies, in its Part III, the cyclotomic resonances of one repeated root for weighted Catalan moments, a different family; there too at most one residue class loses an order. The two manuscripts were written independently and neither uses the other; this parallel is an observation of this merge. The sibling report `cigler-conjecture-8-schur` remarks that specialized parameters can merge the poles of its two-cluster family $b_n(t)$; Part III does not treat that family.

Where the merge had to choose.

- *Duplicated material.* Lemma 25.1 is Part II's Theorem 16.1, with nodes α_i in place of ξ_i and L clusters in place of ℓ , and (25.4) is (16.5) with the cross factors normalized; the manuscript credits the argument to Part II. (25.6), (24.2) and (25.7) restate (17.1), (17.2) and (17.3); the Gram determinants (29.7) are Part II's (13.1)–(13.2), and the odd-shift identity (29.8) is the odd part of Part II's Proposition 13.1 (the manuscript calls it “not an independent new claim”); (26.3) is the negative-index continuation behind Part II's Lemma 18.1, of which Lemma 26.2 is a mode-by-mode refinement. In Corollary 31.1 the cases $t \in \{0, 1, -1\}$ and real t are Part II's Theorems 19.2 and 12.2 (agreement checked during the intake); complex t not a root of unity extends Theorem 12.2 from the real line, and roots of unity are new. The manuscript re-derives this material so that its argument is self-contained; it is printed in both parts, and Part III's proofs are second routes. The interface (23.4) is Part II's Theorem 12.1; Part III uses it as an input and does not reprove it.
- *Numbering.* The manuscript's Section n is Section $n + 22$ here, and its Theorem, Lemma, Proposition, Corollary, Remark and equation $n.m$ are $(n + 22).m$; its Appendices A–C are Appendices E–G. This subsection and the next two precede its Section 1.1, which is Section 23.4 here. Its table of contents is merged into the report's. Its claim-status file numbers the results as delivered (Theorem 2.1 is Theorem 24.1, and so on).
- *Names.* Every label of Part III carries the prefix `cp:`; four of the manuscript's labels (`thm:main`, `sec:main`, `sec:consequences`, `eq:UV`) would otherwise collide with Part I's. The manuscript's macros for $\mathcal{H}$, $\mathbb{Q}$, $\mathbb{Z}$, $\mathbb{N}$, $\mathbb{C}$ are replaced by the report's, with the same glyphs. Three glyphs change: the manuscript's calligraphic $\mathcal{Q}$ (reduced denominators) and $\mathcal{G}$ (generating function) are printed as Part II's script $\mathcal{Q}$ and $\mathcal{G}$, which denote the same kinds of object; and the report's repetition macro prints $(x)^{[r]}$ with parentheses where the manuscript printed $x^{[r]}$. Its bibliography entry for Parts I and II is replaced by

internal references; its entries for Cigler's paper and for González-Serrano–Maximenko are merged with Part II's (the same papers; the manuscript adds the page range 2613–2647 of the latter), and Krattenthaler's survey and Chern–Shi's paper are added. A layout-only \Needspace before Theorem 24.1 is dropped. No normalization is changed. Symbols that mean different things in Part III and the earlier parts are listed in Section 23.3.

- *Added text.* Dated notes were added on the title page, in the abstract, in Part II's research-topic Sections 21.1, 21.2 and 21.6 (which received new labels `cp:sec:rs-...`) and in Part II's conclusion; and six bracketed notes in Part III: on the pinned paths and cited numbers (Section 23.4), on the duplicated cluster lemma (after Lemma 25.1), on Part II's real-parameter results (after Corollary 31.1), on the shipped names and a rerun (Section 32), on the relation of the research questions to Part II's (Section 33), and on the delivered commands (Appendix G). All are marked [Added 30 September 2026, batch 66: ...]. The research questions received labels `cp:q:...`.

23.2 The manuscript's title, abstract and status

The manuscript's title block reads *Cyclotomic Pole Collapse and Exact Minimal Recurrences for Cigler's Hankel Polynomials*, with the subtitle *A complete complex-parameter classification and an exceptional fourth-root cancellation law*, and the date 30 September 2026. Its abstract:

A ProveIt research report identifies normalized shifted Hankel determinants of Cigler's middle-binomial polynomial family with rectangular Schur polynomials, and determines their minimal recurrence away from complex root-of-unity collisions. It explicitly leaves the latter cancellations unclassified. We give the missing classification for every shift and every root of unity. At a primitive root of order $\ell \geq 3$, the possible spectral degrees are $\lfloor q(k-1-q)/2 \rfloor$. In each residue class modulo ℓ the largest one survives, except possibly in a single residue class. An explicit parity criterion decides whether cancellation occurs. When it does, the pole order usually drops by one; precisely for $\ell = 4$, $k \equiv 1 \pmod{4}$, it drops by three, with a small-shift disappearance exception.

The proof combines a reflection identity with a general explicit formula for the first subleading coefficient of a confluent rectangular-Schur spectral polynomial. A fourth-root product evaluation and a fifth elementary-symmetric coefficient calculation resolve the exceptional case. We obtain minimal recurrences at every complex parameter, exact fourth-root quasipolynomials, and a quadratic rather than cubic recurrence-order law at fixed cyclotomic order. All-index proofs, reproducible exact checks, a dependency audit, and further research questions are included.

Its status box:

Status. The article supplies mathematical proofs, not a Lean or Rocq formalization. The Hankel–Schur identification is an explicitly cited input from the inspected repository; the spectral classification is proved here. The general confluent and orthogonal-polynomial machinery is classical. Neither external peer review nor exhaustive historical priority is claimed. The target is the repository's stated cyclotomic gap, not the entirety of Cigler's Conjectures 17 or 18.

23.3 Notation shared with Parts I and II

Part III keeps the manuscript's notation. It agrees with Part II on $c_r(t)$, $D_k(n;t)$, $\mathcal{H}_k(n;t)$, a , b , $M = k - 1$, the forward shift E , $Q_t(z) = (1 + tz)(1 + t/z)$, the Gram determinants $\mathcal{U}_N(q)$, $\mathcal{V}_N(q)$ and $U_h(N)$, $V_h(N)$, the bases P_j , R_j , the allocation data r_i , $s_i = m_i - r_i$, K , A , $C = K - A$, $B(m, r)$, λ_r , L_r , the degrees d_q and the canonical polynomials $P_{k,q}(n;t)$. Its alphabet $X_k(t)$ is Part II's $\mathcal{X}_k$. The following letters mean different things in Part III and the earlier parts, or within Part III; the right-hand column gives the tempting false reading.

Symbol	Parts I and II	Part III	Do not read as
$\mathcal{Q}$	$\mathcal{Q}_k(z, t)$: the generic reduced denominator (12.9) over $\mathbb{Q}(t)$	$\mathcal{Q}_{k,\zeta}(z)$, $\mathcal{Q}_{k,t}(z)$: the reduced denominator at the specialized parameter (24.7), (31.1)	$\mathcal{Q}_{k,\zeta}(z)$ as $\mathcal{Q}_k(z, \zeta)$, which adds the colliding multiplicities and is not reduced
e	$e_{k,q} = 1 + d_q$, indexed by the eigenvalue exponent q (12.8); $e_j(X)$: elementary symmetric polynomials	e_ρ , e_ρ^0 : exponents indexed by a residue ρ modulo ℓ (24.3); $e_j^{(r)}$: elementary symmetric functions of root multisets (Section 30)	e_ρ^0 as $e_{k,\rho}$, or as a sum of the $e_{k,q}$ with $q \equiv \rho$; it is their maximum
N	$N_k = \sum_q e_{k,q}$: generic order	$N_{k,\ell} = \sum_\rho e_\rho$: order at a root of order ℓ	—
ℓ, L	ℓ : the number of distinct nodes $\xi_1, \dots, \xi_\ell$ in Theorem 16.1; L : Part I's moment functional	ℓ : the exact order of the root ζ ; L : the number of nodes $\alpha_1, \dots, \alpha_L$ (Section 25), an integer variable in $\mathcal{H}_{4m+1}(4L + r; i)$ (Sections 29 and 30), and a matrix size (Appendix E)	Part III's ℓ as a number of clusters
δ	$\delta \in \{0, 1\}$ (Part I); $\delta_k = \lfloor k^2/4 \rfloor$ (18.1)	$\delta = 2m^2$ for $k = 4m + 1$ (Sections 29–31); $\eta = 2m(m + 1)$ for $k = 4m + 3$	δ as δ_k : for $k = 4m + 1$, $\delta_k = 4m^2 + 2m$
D	$D_k(n; t)$: the Hankel determinant	also $D_k(n; t)$; and the scalar tie degree $D = \lfloor (M^2 - \ell^2)/8 \rfloor$ (24.4); D_N^0 , $D_L(A, B)$: moment determinants (Section 29, Appendix E)	the scalar D as a determinant
U, V	$U_a(r)$, $V_a(r)$: matrices (Part I); $U_h(m)$, $V_h(m)$: normalized Gram determinants (Part II)	$U(t) = (1 + t)/(1 - t)$, $V(t) = (1 + t^2)/(1 - t^2)$ (28.1) (Section 28); $U_h(N)$, $V_h(N)$ as in Part II and their rescalings U_N^0 , V_N^0 (Section 29)	$U(t)$ as a Gram determinant
x	Part I: generating variable; Part II: $x = z + z^{-1}$	$x = n + k/2$ (26.4); $n = x - K/2$ in Theorem 27.1; $x = z + z^{-1}$ in Section 29; $x = n + 2m + 1/2$ in Section 30	—
c	$c_r(t)$: Cigler's moments	also $c_r(t)$; c_r , $c_q(t)$: centered first coefficients (27.2), (28.2); $c = 2m + 1/2$ (Section 30); $c \geq 2$: a middle multiplicity (Section 33.1)	—
A, T	A : the height of a rectangle (Part II); $\mathcal{A}_k$: generating numerator	also the height A ; $A_s(L)$, $T_s(L)$: binomial and Catalan moment determinants (29.9); $A_\rho(n)$, $A_j(n)$: grouped polynomials; A , B : beta parameters (Appendix E); $T_s(b, j)$: triangular power sums (30.5)	$T_s(L)$ as $T_s(b, j)$
B	$B = 1 + t^2$ (Part I); $B(m, r)$ (Part II)	$B(m, r)$ as in Part II; B_i (27.3); B : the beta function	—
C	$C_k = \binom{k}{2}$; $C = K - A$ (Part II); $C_j(y)$ (Part I)	$C = K - A$; $C_j = \binom{2j}{j}/(j + 1)$: Catalan numbers (Section 29)	—
F, G	$\mathcal{F}_k$, F_k : the limit function; $\mathcal{G}_k$: the generating function	$F_r(n)$, $G_r(x)$: residue polynomials at $t = i$ (Section 30); $\mathcal{G}_k$ as in Part II	F_r as the limit function F_k
α, β	—	α_i : cluster nodes (Part II's ξ_i); $\alpha_s(L)$, $\beta_s(L)$: product polynomials (29.1)–(29.2)	$\alpha_s(L)$ as a node

Symbol	Parts I and II	Part III	Do not read as
h	the index of the parity factors; derivative orders; h_j	also the parity-factor index ($h = 2m$ in Section 29); h_i , h : selected sums (Section 27); $h = M - 2q$ (Section 28)	—
p, P	P_i, R_i : monic bases; $P_{k,q}$: canonical polynomials	the same; also $p_N(x)$: monic orthogonal polynomials, $p_j^{(r)}$: power sums, P_{bad} : the exceptional phase polynomial	—
R	R_i : monic bases (Part II)	R_j the same; R_i : derivative-order sets (proof of Lemma 25.1); $\mathcal{R}_j$: root multisets; $R_{k,\zeta}(E)$: minimal recurrence	—
ε, J	ε (Part I); $J = \text{diag}(1, -1, \dots)$ (Part I), $Jf(z) = zf(z^{-1})$ (Part II)	$\varepsilon = \pm 1$ (26.5); ε : a splitting parameter, $\mathcal{J}$: an alternant functional (Section 27)	—

23.4 Which question is being resolved?

The relevant ProveIt report is *A parity factorization and sharp stabilization for Cigler's Hankel polynomials*, with its Part II, *Rectangular Schur structure and sharp recurrences*: Parts I and II of the present report. At the inspected commit, Part II proves a rectangular-Schur identity, exact generic recurrence multiplicities, and the real-parameter classification. Its Section 21.2 explicitly leaves spectral cancellations at complex roots of unity unclassified. That is the question resolved below.

This distinction is important. Simply specializing the generic denominator does not determine a minimal recurrence. Different exponential modes acquire the same eigenvalue, their polynomial coefficients must be added, and the highest terms of that sum can cancel. A bound on an annihilating recurrence therefore does not answer the question.

The source snapshot is ProveIt commit

ff6969475bad8f61a33268418aaf23d23fdcd1fb.

The inspected manuscript is

SetTheory/Cardinals/docs/reports/hankel-determinants/catalan-and-ballot/
cigler-conjecture-16-parity/article.tex.

Its Git blob is 07c41161857ed17854b2208b66418e9d73b956d5. The companion status document also records the cyclotomic limitation. These are research manuscripts, not proof-assistant certificates. Their mathematical inputs are isolated below so that the present results can be read independently of the repository's other projects.

[Added 30 September 2026, batch 66: the inspected manuscript is this report's `article.tex` at the pin, that is, Parts I and II as printed above (Section 23.1); the companion status document is shipped as `02-schur-recurrences-STATUS.md`, whose line 51 records the limitation. The manuscript's bibliography entry for the report is replaced by internal references; the numbers it cites, Section 21.2 and Theorems 12.1, 16.1 and 19.2, are Section 21.2 and Theorems 12.1, 16.1 and 19.2 of this report.]

23.5 Cigler's moments and the normalization

A binomial coefficient is zero outside its ordinary range. Following Cigler [1], define

$$c_0(t) = 1, \quad c_r(t) = \sum_{j=1}^r \binom{\lfloor (r-1)/2 \rfloor}{\lfloor (j-1)/2 \rfloor} \binom{\lfloor r/2 \rfloor}{\lfloor j/2 \rfloor} t^{j-1} \quad (r \geq 1). \quad (23.1)$$

For instance, $c_1 = 1$, $c_2 = 1 + t$, $c_3 = 1 + t + t^2$, and $c_4 = 1 + 2t + 2t^2 + t^3$. Put

$$D_k(n; t) = \det(c_{k+i+j}(t))_{0 \leq i, j < n}, \quad D_k(0; t) = 1. \quad (23.2)$$

Throughout the article $k \geq 1$ and

$$a = \lfloor k/2 \rfloor, \quad b = \lfloor (k-1)/2 \rfloor, \quad M = k-1, \quad X_k(t) = ((1)^{[a]}, t, (t^2)^{[b]}). \quad (23.3)$$

Here $(x)^{[r]}$ denotes r repetitions of x , not an exponent. The explicit interface supplied by Part II (Theorem 12.1) is

$$\mathcal{H}_k(n; t) := s_{(n^a)}(X_k(t)) = (-1)^{k \binom{n}{2}} t^{-\binom{n}{2}} D_k(n; t). \quad (23.4)$$

The identity means that the quotient extends to the polynomial on the left, including at $t = 0$. It is not permission to evaluate an undefined quotient. The empty-height Schur polynomial gives $\mathcal{H}_1(n; t) = 1$.

Our spectral theorems are unconditional statements about the explicitly defined Schur sequence on the left of (23.4). Their application to the original Hankel determinants uses exactly the displayed identity. The repository proves that identity through a Laurent-moment parity factorization and a Toeplitz factorization; it is not assumed merely because finite numerical values agree. For the fourth-root analysis we also use its odd-shift parity identity, which is rederived in Section 29.

23.6 What is new here, and what is background?

The principal addition is the all- k , all- ℓ minimality theorem in Section 24. Two useful intermediate results are the general centered subleading-coefficient formula in Section 27 and the four explicit fourth-root product formulas in Section 29. The five-coefficient calculation in Section 30 proves, rather than extrapolates, the exceptional loss of three pole orders.

Schur polynomials, the bialternant formula, confluence of repeated variables, Christoffel transformations, and beta-moment Hankel determinants are existing mathematics [8, 10]. We supply the particular calculations needed here, and do not present these general techniques as discoveries. The repository's Schur identity and generic leading-coefficient theorem are likewise not claimed as additions of this article.

A bounded literature check included Cigler's original paper, repeated-variable bialternant work [8], determinant-calculus sources [10], and Chern-Shi's August 2026 paper [11]. The latter treats shifted Catalan-like arrays indexed by a fixed meander height, a different parameter family from the three-cluster rectangular character studied here. No matching all-shift cyclotomic pole classification was identified in that check. This is a description of the check, not a proof of worldwide priority.

24 The exact cyclotomic classification

24.1 Minimal recurrence and pole multiplicity

Let E be the forward shift: $(Ef)(n) = f(n+1)$. A monic polynomial $R(E)$ is a constant-coefficient recurrence if $R(E)f(n) = 0$ for every $n \geq 0$. The minimal one divides every such

recurrence. The denominator of

$$\mathcal{G}_k(z, t) = \sum_{n \geq 0} \mathcal{H}_k(n; t) z^n \quad (24.1)$$

is always normalized to have constant term one and to be relatively prime to its numerator. An exponent zero means that a factor is omitted.

For $0 \leq q \leq M$, define

$$d_q = \left\lfloor \frac{q(M-q)}{2} \right\rfloor. \quad (24.2)$$

At a primitive ℓ th root ζ , $\ell \geq 3$, define the baseline exponents

$$e_\rho^0 = \begin{cases} 1 + \max\{d_q : 0 \leq q \leq M, q \equiv \rho \pmod{\ell}\}, & \text{if this set is nonempty,} \\ 0, & \text{otherwise,} \end{cases} \quad 0 \leq \rho < \ell. \quad (24.3)$$

They use a *maximum*, not a sum. The following theorem gives the exact correction to that maximum.

Theorem 24.1 (Complete cyclotomic pole classification). *Let $k \geq 1$ and let ζ have exact order $\ell \geq 3$. Start with $e_\rho = e_\rho^0$ from (24.3).*

If $\ell > M$ or $\ell \not\equiv M \pmod{2}$, no correction is required. Otherwise put

$$q_* = (M - \ell)/2, \quad \rho_* = q_* \bmod \ell, \quad D = \left\lfloor \frac{M^2 - \ell^2}{8} \right\rfloor. \quad (24.4)$$

Only e_{ρ_} can change. The cancellation criterion is*

$$\begin{array}{l|l} k = 2a & a + D \text{ is odd} \\ k = 2a + 1 & D \text{ is even.} \end{array} \quad (24.5)$$

If the criterion fails, leave $e_{\rho_} = D + 1$. If it holds, set*

$$e_{\rho_*} = \begin{cases} \max(0, D - 2), & \ell = 4 \text{ and } k \equiv 1 \pmod{4}, \\ D, & \text{otherwise.} \end{cases} \quad (24.6)$$

Then the reduced denominator and the monic minimal recurrence are exactly

$$\boxed{\mathcal{Q}_{k,\zeta}(z) = \prod_{\rho=0}^{\ell-1} (1 - \zeta^\rho z)^{e_\rho}, \quad R_{k,\zeta}(E) = \prod_{\rho=0}^{\ell-1} (E - \zeta^\rho)^{e_\rho}.} \quad (24.7)$$

The exponent list depends on k and ℓ , not on the choice of primitive root of that order.

Remark 24.2 (What “loss of three” means). In the exceptional family write $k = 4m + 1$, $m \geq 1$. Then $D = 2m^2 - 2$. For $m \geq 2$ the exponent drops from $D + 1$ to $D - 2$, exactly three orders. For $m = 1$, the baseline exponent is one and the factor disappears; negative exponents are not introduced.

24.2 Small examples

The exponent vectors below are ordered by phases $1, \zeta, \dots, \zeta^{\ell-1}$.

k	ℓ	baseline e^0	exact e	outcome
4	3	(1, 2, 2)	(1, 2, 2)	tie, no cancellation
5	4	(1, 2, 3, 2)	(0, 2, 3, 2)	disappearance
6	3	(4, 3, 4)	(4, 2, 4)	one-order loss
6	5	(1, 3, 4, 4, 3)	(0, 3, 4, 4, 3)	disappearance
7	4	(5, 3, 5, 5)	(5, 2, 5, 5)	one-order loss
9	4	(9, 8, 7, 8)	(9, 8, 4, 8)	three-order loss

In particular,

$$\mathcal{Q}_{9,i}(z) = (1 - z)^9(1 + z)^4(1 + z^2)^8. \quad (24.8)$$

The maximum-degree rule alone would incorrectly give exponent seven at $1 + z$. Direct substitution into the generic denominator would be still less accurate because it would add all colliding multiplicities.

24.3 Outline of the proof

The proof has three separable parts. First, strict concavity shows that only one residue can have two maximal-degree contributions. Second, a negative-index reflection identity decides whether their leading terms cancel. The explicit first subleading coefficient proves that cancellation stops immediately, except at primitive fourth roots in one congruence class of k . Third, exact fourth-root product formulas show that two further coefficients cancel there, and that the next permitted coefficient is nonzero. Each ingredient is proved in the following sections.

25 Confluent rectangular characters and their spectral terms

25.1 The distinct-variable subset identity

For K distinct nonzero variables $x_1, \dots, x_K$, a rectangle of height A , and $C = K - A$, the bialternant gives

$$s_{(n^A)}(X) = \sum_{\substack{I \subseteq \{1, \dots, K\} \\ |I|=A}} \left(\prod_{i \in I} x_i \right)^n \prod_{\substack{i \in I \\ j \notin I}} (1 - x_j/x_i)^{-1}. \quad (25.1)$$

Indeed, use exponent columns $0, 1, \dots, C - 1, n + C, \dots, n + K - 1$, and expand along the last A columns. The Vandermonde factors internal to I and its complement cancel. The cross factors and the powers x_i^C give precisely (25.1). This also defines an exponential-polynomial extension $\tilde{s}_A(n; X)$ for all integer n ; no negative partition is being asserted to exist.

Now coalesce m_i variables at distinct nonzero nodes α_i , $1 \leq i \leq L$, where $\sum_i m_i = K$. An allocation is a vector $\mathbf{r}$ with $0 \leq r_i \leq m_i$ and $\sum_i r_i = A$. Write

$$s_i = m_i - r_i, \quad d_i = r_i s_i, \quad d = \sum_i d_i, \quad \lambda_{\mathbf{r}} = \prod_i \alpha_i^{r_i}, \quad B(m, r) = \frac{\prod_{h=0}^{r-1} h!}{\prod_{h=m-r}^{m-1} h!}. \quad (25.2)$$

Empty products equal one, in particular $B(m, 0) = B(m, m) = 1$.

Lemma 25.1 (Canonical cluster polynomials). *There are canonically grouped polynomials $P_{\mathbf{r}}(n)$ such that*

$$s_{(n^A)}((\alpha_1)^{[m_1]}, \dots, (\alpha_L)^{[m_L]}) = \sum_{\mathbf{r}} \lambda_{\mathbf{r}}^n P_{\mathbf{r}}(n). \quad (25.3)$$

They have exact degree d and nonzero leading coefficient

$$L_{\mathbf{r}} = \prod_i B(m_i, r_i) \prod_{i \neq j} (1 - \alpha_j / \alpha_i)^{-r_i s_j}. \quad (25.4)$$

The grouping is by allocations, even when two products $\lambda_{\mathbf{r}}$ coincide.

Proof. The normalized confluent row of order h at α_i in an exponent column e is $\binom{e}{h} \alpha_i^{e-h}$. The denominator is $\prod_{i < j} (\alpha_j - \alpha_i)^{m_i m_j}$. Expand the numerator along its A high columns. Selecting r_i rows in cluster i extracts the factor $\alpha_i^{n r_i}$, so the remaining coefficient is polynomial in n .

Suppose the selected derivative orders in this cluster form R_i . The normalized Leibniz rule for $x^n g(x)$ gives a degree bound $\sum_{h \in R_i} h - \binom{r_i}{2}$: distinct derivative rows of g are required in a nonzero determinant, and their smallest possible total order is $\binom{r_i}{2}$. The unique maximizing selection is $R_i = \{s_i, \dots, m_i - 1\}$, giving degree $r_i s_i$. Its local leading determinant is

$$\det \left(\frac{1}{(s_i + h - p)!} \right)_{0 \leq h, p < r_i} = B(m_i, r_i), \quad (25.5)$$

where negative factorials mean zero entries. Multiply row h by $(s_i + h)!$; the columns become monic falling-factorial polynomials at $s_i, s_i + 1, \dots, s_i + r_i - 1$, proving (25.5).

The remaining low and high confluent Vandermondes give

$$(-1)^{\sum_{i < j} r_i s_j} \frac{\prod_i B(m_i, r_i) \alpha_i^{C r_i - r_i s_i}}{\prod_{i < j} (\alpha_j - \alpha_i)^{s_i r_j + r_i s_j}},$$

which is (25.4). All factors are nonzero. This proof is the confluent leading-term argument already used in Part II (Theorem 16.1); general repeated-variable bialternants are treated in [8]. $\square$

[Added 30 September 2026, batch 66: Lemma 25.1 is Part II's Theorem 16.1, whose nodes $\xi_1, \dots, \xi_\ell$ are written $\alpha_1, \dots, \alpha_L$ here, and (25.4) is (16.5), written with normalized cross factors; the proof above condenses Part II's argument, as it says. Part II's remark after that theorem already notes that summands with a common product must be combined; Sections 26–30 carry out that combination at roots of unity.]

25.2 Specialization to the three-cluster alphabet

For $t \neq 0, \pm 1$, the nodes $1, t, t^2$ are distinct. In (25.2) take multiplicities $(a, 1, b)$ and height $A = a$. There is exactly one allocation for each $q = 0, \dots, M$:

$$r_0 = a - \lceil q/2 \rceil, \quad r_1 = q \bmod 2, \quad r_2 = \lfloor q/2 \rfloor. \quad (25.6)$$

Its eigenvalue is t^q . Direct substitution gives degree d_q in (24.2). Hence

$$\mathcal{H}_k(n; t) = \sum_{q=0}^M t^{qn} P_{k,q}(n; t), \quad \deg_n P_{k,q} = d_q. \quad (25.7)$$

Each canonical $P_{k,q}$ is a rational function of t with polynomial coefficients in n ; its only possible parameter poles are $0, 1, -1$. Thus (25.7) specializes regularly at every root of unity of order at least three. Confluence of *nodes* is not needed there: only eigenvalue products collide.

Lemma 25.2 (Minimality for exponential polynomials). *If $f(n) = \sum_{\lambda} \lambda^n A_{\lambda}(n)$, with distinct nonzero λ and nonzero polynomials A_{λ} , then its minimal recurrence is $\prod_{\lambda} (E - \lambda)^{1 + \deg A_{\lambda}}$. Its reduced generating denominator is $\prod_{\lambda} (1 - \lambda z)^{1 + \deg A_{\lambda}}$.*

Proof. The operator $E - \lambda$ lowers the polynomial degree in the λ mode by one, whereas $E - \mu$, $\mu \neq \lambda$, preserves it with nonzero leading multiplier $\lambda - \mu$. Applying operators that kill all but one mode proves their linear independence and the stated minimality. Equivalently, a degree- d polynomial times λ^n has a pole of exact order $d + 1$ at $z = \lambda^{-1}$. $\square$

For a root ζ , therefore, the entire question is the exact degree of

$$A_\rho(n) = \sum_{q \equiv \rho \pmod{\ell}} P_{k,q}(n; \zeta). \quad (25.8)$$

If this polynomial is zero, its phase is absent.

26 Why there is at most one cancellation class

26.1 The concavity restriction

Lemma 26.1 (Unique possible tie). *Among the q in a fixed residue class modulo $\ell \geq 3$, the maximum of d_q is unique except possibly for a pair $q, M - q$. A two-way maximum occurs in exactly one residue if and only if $\ell \leq M$ and $\ell \equiv M \pmod{2}$. Its two indices are q_* and $q_* + \ell$ from (24.4). Every other index in that residue, when present, has degree at most $D - \ell^2$.*

Proof. If $d_q = d_p$, the integers $q(M - q)$ and $p(M - p)$ differ by at most one. Their difference is $(q - p)(M - q - p)$. For different indices in one residue, $|q - p| \geq \ell \geq 3$, so necessarily $q + p = M$. If two symmetric maxima were separated by more than ℓ , an intermediate index of the same residue would be strictly closer to $M/2$ and have larger degree. Thus $p - q = \ell$, proving the criterion and uniqueness. The next possible indices are at distance $3\ell/2$ from $M/2$, rather than $\ell/2$. The quadratic values divided by two decrease by exactly ℓ^2 ; taking floors preserves this integral difference. $\square$

A unique maximum cannot cancel because its leading coefficient in (25.4) is nonzero. The lemma already proves all uncorrected entries of Theorem 24.1.

26.2 Reflection about the canonical center

Lemma 26.2 (Negative-index reflection). *The canonical polynomials of (25.7) satisfy*

$$P_{k,M-q}(n; t) = (-1)^a t^{a(M-2q)} P_{k,q}(-n - k; t), \quad k = 2a, \quad (26.1)$$

$$P_{k,M-q}(n; t) = t^{k(a-q)} P_{k,q}(-n - k; t), \quad k = 2a + 1. \quad (26.2)$$

These are identities of rational functions in t and polynomials in n .

Proof. Complementing the subsets in (25.1) gives

$$\tilde{s}_A(-n - K; X) = (-1)^{A(K-A)} \left(\prod X \right)^{-n-A} s_{(n^{K-A})}(X). \quad (26.3)$$

For example, each cross denominator changes sign on complementation; there are $A(K - A)$ such signs. This argument works for the exponential-polynomial extension, so there is no appeal to a Schur polynomial indexed by a negative partition.

For even k , $K - A = A = a$ and $\prod X_k = t^M$. Comparing distinct powers t^{qn} over $\mathbb{Q}(t)$ in (26.3) gives (26.1). For odd k , $\prod X_k = t^k$ and $X_k^{-1} = t^{-2} X_k$ as multisets. The rectangular complement identity $s_{(n^{K-A})}(X) = (\prod X)^n s_{(n^A)}(X^{-1})$ therefore yields $s_{(n^{a+1})}(X_k) = t^n s_{(n^a)}(X_k)$. Equation (26.2) follows. The identities continue to every $t \neq 0, \pm 1$ because the canonical coefficients are regular there. $\square$

At a tied residue let $q = q_*$ and introduce the centered variable

$$x = n + k/2. \quad (26.4)$$

If $p(x) = P_{k,q}(x - k/2; \zeta)$, the sum of the two maximal-degree polynomials is

$$p(x) + \varepsilon p(-x), \quad \varepsilon = \begin{cases} (-1)^a, & k \text{ even,} \\ -1, & k \text{ odd.} \end{cases} \quad (26.5)$$

For the odd case, $a - q_* = \ell/2$ and $\zeta^{k\ell/2} = (-1)^k = -1$. Its leading coefficient is multiplied by $1 + \varepsilon(-1)^D$, giving exactly (24.5). If $D = 0$ and the criterion holds, the constant pair disappears, and there are no lower terms in that residue by Lemma 26.1. We henceforth consider $D > 0$.

27 A general first subleading coefficient for confluent Schur modes

The leading coefficient alone cannot distinguish an ordinary cancellation from the fourth-root exception. We now calculate one further coefficient for arbitrary cluster multiplicities. This calculation is useful beyond Cigler's family.

Theorem 27.1 (Centered first-coefficient formula). *Use the notation of Lemma 25.1, and suppose $d \geq 1$. Then*

$$P_r(x - K/2) = L_r(x^d + c_r x^{d-1} + O(x^{d-2})), \quad (27.1)$$

where

$$c_r = \frac{1}{2} \sum_{i < j} [d_i(2r_j - m_j) - d_j(2r_i - m_i)] \frac{\alpha_i + \alpha_j}{\alpha_i - \alpha_j}. \quad (27.2)$$

Here $O(x^u)$ denotes a polynomial of degree at most u ; if $u < 0$, that polynomial is zero. Equivalently, before centering,

$$[n^{d-1}]P_r(n) = L_r \sum_i d_i B_i, \quad B_i = \frac{m_i}{2} + \sum_{j \neq i} \frac{r_j - s_j \alpha_j / \alpha_i}{1 - \alpha_j / \alpha_i}. \quad (27.3)$$

Proof. Split the i th cluster formally into $\alpha_i \exp(\epsilon y_{i,1}), \dots, \alpha_i \exp(\epsilon y_{i,m_i})$, with $\sum_u y_{i,u} = 0$. The y 's may be chosen distinct and generic. In (25.1), group subsets that select r_i variables in each cluster. Such a grouped expression has a removable singularity as $\epsilon \rightarrow 0$: antisymmetrizing within each cluster cancels its internal Vandermonde, while the cross-cluster denominators remain nonzero. Its limit is exactly the canonical $P_r(n)$ times λ_r^n .

For a selected set I_i let $h_i = \sum_{u \in I_i} y_{i,u}$ and $h = \sum_i h_i$. Consider the finite alternant functional

$$\mathcal{J}(f) = \sum_{|I_i|=r_i} \frac{f(I_1, \dots, I_L)}{\prod_i \prod_{u \in I_i, v \notin I_i} (y_{i,u} - y_{i,v})}. \quad (27.4)$$

For a polynomial separately symmetric in selected and unselected variables, possible poles cancel in pairs under an interchange of two variables. Thus the result is polynomial and symmetric in each whole cluster. If the input is homogeneous of degree less than d , its output is zero; at degree d it is a constant. This also follows directly by multiplying by the cluster Vandermondes: the alternating numerator must be divisible by each of them.

The functional factors by cluster. The same leading confluent determinant (25.5), applied to the exponential subset sum, gives

$$\mathcal{J}\left(\frac{h^d}{d!}\right) = \prod_i B(m_i, r_i), \quad \mathcal{J}\left(\frac{h_i h^{d-1}}{(d-1)!}\right) = d_i \prod_j B(m_j, r_j). \quad (27.5)$$

For clarity, the one-cluster identity in the first formula is $\mathcal{J}_i(h_i^{d_i}/d_i!) = B(m_i, r_i)$; it is the coefficient of n^{d_i} in the coalescing bialternant. In the product functional every cluster must receive degree at least d_i . At total degree d , it receives exactly that degree. The multinomial theorem proves both displayed formulas, with multiplier d_i in the second. This also covers $d_i = 0$.

For selected u and unselected v in one cluster,

$$\frac{1}{1 - e^{\epsilon(v-u)}} = \frac{1}{\epsilon(u-v)} \left(1 + \frac{\epsilon}{2}(u-v) + O(\epsilon^2)\right). \quad (27.6)$$

The total linear correction in cluster i is $m_i h_i/2$ because $\sum_{u \in I_i, v \notin I_i} (y_{i,u} - y_{i,v}) = m_i h_i$. For two different clusters put $z_{ij} = \alpha_j/\alpha_i$. Their cross factor expands as

$$\frac{1}{1 - z_{ij} e^{\epsilon(v-u)}} = \frac{1}{1 - z_{ij}} \left(1 + \epsilon \frac{z_{ij}(v-u)}{1 - z_{ij}} + O(\epsilon^2)\right). \quad (27.7)$$

Summing $v - u$ over selected variables in i and unselected variables in j gives $-r_i h_j - s_j h_i$. After both orientations of each pair are collected, the coefficient of h_i is exactly B_i in (27.3). The exponential factor contributes $e^{\epsilon n h}$.

After extracting ϵ^{-d} and the constant cross-factor product, the coefficient of n^d in the finite limit comes from $h^d/d!$. The coefficient of n^{d-1} comes only from the linear correction $\sum_i B_i h_i$ times $h^{d-1}/(d-1)!$. Equation (27.5) proves (27.3). No higher-order expansion term can contribute to n^{d-1} .

Replacing n by $x - K/2$ subtracts $dK/2$ from the normalized subleading coefficient. Finally,

$$B_i - K/2 = \frac{1}{2} \sum_{j \neq i} (2r_j - m_j) \frac{\alpha_i + \alpha_j}{\alpha_i - \alpha_j}.$$

Antisymmetrization in i, j yields (27.2). □

Remark 27.2 (A useful consistency check). For a single cluster the sum in (27.2) is empty. Thus a rectangular Weyl-dimension polynomial is centered at $-K/2$ with zero first subleading coefficient. The formula detects the additional asymmetry introduced by separated nodes, rather than an arbitrary choice of origin.

28 Ordinary cyclotomic cancellations stop after one order

Set

$$U(t) = \frac{1+t}{1-t}, \quad V(t) = \frac{1+t^2}{1-t^2}. \quad (28.1)$$

For the allocation (25.6), put $q = 2r + \sigma$, $\sigma \in \{0, 1\}$, and

$$d_0 = (a - r - \sigma)(r + \sigma), \quad d_2 = r(b - r), \quad u_0 = a - 2r - 2\sigma, \quad u_2 = 2r - b.$$

The middle cluster has $d_1 = 0$. Equation (27.2) becomes

$$c_q(t) = \frac{1}{2} [(2\sigma - 1)(d_0 - d_2)U(t) + (d_0 u_2 - d_2 u_0)V(t)]. \quad (28.2)$$

Lemma 28.1 (Nonvanishing of the centered coefficient). *Suppose a tied maximal pair has common degree $D > 0$ and satisfies the cancellation criterion (24.5). Then $c_q(\zeta) \neq 0$, except when $\ell = 4$ and $k \equiv 1 \pmod{4}$.*

Proof. Write $\zeta = e^{i\theta}$. We repeatedly use

$$\frac{V(\zeta)}{U(\zeta)} = \frac{\cos \theta}{1 + \cos \theta}. \quad (28.3)$$

All denominators here are legitimate because $\ell \geq 3$. If the cosine of the argument of a root of unity is rational, then $2 \cos \theta = \zeta + \zeta^{-1}$ is a rational algebraic integer and hence an integer. The only nonreal possibilities are $-1/2, 0, 1/2$, corresponding to orders 3, 4, 6.

Even $k = 2a$. Here M and the tied separation ℓ are odd. Choose the even index $q = 2r$ in the symmetric pair, and let $h = M - 2q = \pm \ell$. Then $D = r(M - q)$, and (28.2) simplifies to

$$c_q = -\frac{r}{2}U + \frac{r - Dh}{4}V. \quad (28.4)$$

Since $D > 0$, $r > 0$. Vanishing would imply $\cos \theta = -2r/(r + Dh)$; a zero denominator cannot solve the equation. The order is odd, so the only possibility is $\ell = 3$ and cosine $-1/2$, giving $Dh = 3r$. If $h = -3$ this contradicts positivity. If $h = 3$, it would require $M - q = 1$, whereas $M - q = q + 3 \geq 3$. Thus c_q cannot vanish.

Odd $k = 2a + 1$, even tied index. Take the smaller tied index $q = q_*$ and put $v = a - q = \ell/2 > 0$. When q is even, (28.2) reduces to

$$c_q = -\frac{Dv}{2}V. \quad (28.5)$$

It vanishes exactly when $\zeta^2 = -1$, that is, $\ell = 4$. Then $q = a - 2$ is even, so a is even and $k \equiv 1 \pmod{4}$.

Odd $k = 2a + 1$, odd tied index. For odd q , the corresponding simplification is

$$c_q = \frac{v}{2}(U - (D + 1)V). \quad (28.6)$$

Vanishing would force $\cos \theta = 1/D$. The only admissible rational cosine is $1/2$, hence $\ell = 6$, $v = 3$, and $D = 2$. But $D = \lfloor (a^2 - 9)/2 \rfloor = 2$ would give $13 \leq a^2 < 15$, impossible for an integer a . $\square$

Proposition 28.2 (One-order cancellation outside the exceptional family). *Under the hypotheses of Lemma 28.1, excluding its exception, $\deg A_{\rho_*} = D - 1$. Thus the exact exponent is D .*

Proof. In (26.5), cancellation of the degree- D term means $\varepsilon(-1)^D = -1$. The degree- $(D - 1)$ coefficient is therefore $2L_q c_q$, which is nonzero by Lemma 28.1. All remaining terms in the residue have degree at most $D - \ell^2 < D - 1$. They cannot alter it. The $D = 0$ case was dealt with immediately after (26.5). $\square$

29 Exact fourth-root product formulas

The remaining family is $k = 4m + 1$, $m \geq 1$, at $t = i$ or $-i$. We first evaluate its four residue polynomials completely.

Define, for $s \geq 1$,

$$\alpha_s(L) = \prod_{1 \leq u \leq v \leq s-1} \frac{2L + u + v - 1}{u + v - 1}, \quad (29.1)$$

$$\beta_s(L) = \prod_{1 \leq u \leq v \leq s-1} \frac{2L + u + v}{u + v}. \quad (29.2)$$

They are polynomials in L with rational coefficients; $\alpha_1 = \beta_1 = 1$.

Theorem 29.1 (Fourth-root quasipolynomials). *For all integers $m \geq 1$ and $L \geq 0$,*

$$\mathcal{H}_{4m+1}(4L; i) = \alpha_{m+1}(L) \beta_m(L)^2 \beta_{m+1}(L), \quad (29.3)$$

$$\mathcal{H}_{4m+1}(4L + 1; i) = (-1)^m 2 \alpha_m(L + 1) \beta_m(L) \beta_{m+1}(L)^2, \quad (29.4)$$

$$\mathcal{H}_{4m+1}(4L + 2; i) = 2 \alpha_m(L + 1) \beta_m(L + 1) \beta_{m+1}(L)^2, \quad (29.5)$$

$$\mathcal{H}_{4m+1}(4L + 3; i) = (-1)^m \alpha_{m+1}(L + 1) \beta_m(L + 1)^2 \beta_{m+1}(L). \quad (29.6)$$

The same values hold at $-i$. Each residue polynomial has degree $\delta = 2m^2$.

29.1 The odd-shift parity factorization

Here is the specific Hankel reduction used in the proof. Put

$$Q_t(z) = (1 + tz)(1 + t/z) = 1 + t^2 + t(z + z^{-1}), \quad x = z + z^{-1},$$

and, for a symmetric Laurent polynomial $q(z) = q(z^{-1})$, write $q_j = [z^j]q(z)$ and define

$$\mathcal{U}_N(q) = \det(q_{i-j} + q_{i+j+1})_{i,j < N}, \quad \mathcal{V}_N(q) = \det(q_{i-j} - q_{i+j+2})_{i,j < N}. \quad (29.7)$$

Both empty determinants are one. They equal the polynomial moment Gram determinants with weights $q(x)(x + 2)/2$ and $q(x)(4 - x^2)/2$ under CT_z . To check this, use the monic bases

$$P_j(x) = \frac{z^{j+1} + z^{-j}}{z + 1}, \quad R_j(x) = \frac{z^{j+1} - z^{-j-1}}{z - z^{-1}}.$$

Both satisfy the monic recurrence $F_{j+1} = xF_j - F_{j-1}$. Multiplication gives the entries in (29.7); replacing these monic bases by powers of x does not change the determinants.

Write $U_h(N) = \mathcal{U}_N(Q_t^h)$ and $V_h(N) = \mathcal{V}_N(Q_t^h)$. Then

$$\mathcal{H}_{2h+1}(2N; t) = U_h(N) V_h(N), \quad \mathcal{H}_{2h+1}(2N + 1; t) = U_h(N + 1) V_h(N). \quad (29.8)$$

For completeness, these identities follow directly from (23.1). Let $u_j = \text{CT}(1 + z)Q_t^j$ and $v_j = \text{CT}(1 - z^2)Q_t^j$. Binomial expansion gives $c_{2j+1} = u_j$, $c_{2j+2} = (1 + t)u_j$ and $(1 + t)^2 u_j - u_{j+1} = t v_j$. Reorder a Hankel matrix of odd shift with even indices first and subtract $(1 + t)$ times each corresponding even row from the odd row. Its diagonal blocks are a u -moment block and a $-tv$ -moment block. Since $Q_t = 1 + t^2 + tx$, changing powers of Q_t to powers of x in a size- N Gram determinant contributes $t^{N(N-1)}$. For even order the total power is $N + 2N(N - 1) = \binom{2N}{2}$; for odd order it is $N + N(N + 1) + N(N - 1) = \binom{2N+1}{2}$. The signs also cancel the normalization in (23.4). This proves (29.8); it is the odd part of the repository's parity factorization, not an independent new claim.

29.2 Binomial and Catalan moment determinants

Let $C_j = \binom{2j}{j}/(j+1)$ and, for $s \geq 1$, define

$$A_s(L) = \det \left(\binom{2(s+i+j)}{s+i+j} \right)_{i,j < L}, \quad T_s(L) = \det (C_{s+i+j})_{i,j < L}. \quad (29.9)$$

The following elementary evaluations will also specify every normalization in the calculation:

$$A_s(L) = 2^L \alpha_s(L), \quad T_s(L) = \beta_s(L), \quad \frac{T_s(L)}{A_s(L)} = \frac{2^L (1/2)_L}{(s+L)_L}. \quad (29.10)$$

Here $(u)_L = u(u+1) \cdots (u+L-1)$. A proof from beta moments, including an elementary determinant proof rather than an appeal to a Selberg integral, is given in Appendix E.

29.3 Evaluation of the two Gram factors at a fourth root

At $t = i$, $Q_t = ix$. For $h = 2m$, remove the scalar $(-1)^{mN}$ from both size- N Gram determinants, and denote the remaining ones by U_N^0, V_N^0 . Their weights are respectively $x^{2m}(x+2)/2$ and $x^{2m}(4-x^2)/2$. Because $\text{CT } x^{2j} = \binom{2j}{j}$ and odd moments vanish,

$$\frac{1}{2} \text{CT } x^{2j}(4-x^2) = C_j.$$

Separating even and odd monomials therefore gives

$$V_{2L}^0 = T_m(L)T_{m+1}(L), \quad V_{2L+1}^0 = T_m(L+1)T_{m+1}(L). \quad (29.11)$$

Let D_N^0 be the moment determinant for the even weight x^{2m} alone, and let $p_N(x)$ be its monic orthogonal polynomial. The Christoffel determinant identity at -2 , together with $p_N(-x) = (-1)^N p_N(x)$, gives

$$U_N^0 = 2^{-N} D_N^0 p_N(2). \quad (29.12)$$

Under $y = x^2/4$, the even polynomials correspond to beta weight $y^{m-1/2}(1-y)^{-1/2}$ and the odd polynomials to $y^{m+1/2}(1-y)^{-1/2}$. The endpoint ratio of beta determinants gives

$$p_{2L}(2) = \frac{4^L (1/2)_L}{(m+L)_L}, \quad p_{2L+1}(2) = \frac{2 \cdot 4^L (1/2)_L}{(m+L+1)_L}. \quad (29.13)$$

For instance, a monic polynomial π_L for beta weight with parameters (A, B) satisfies $\pi_L(1) = (B)_L / (A+B+L-1)_L$; this follows by applying the Christoffel identity to $(1-y)$ and taking the quotient of the two beta determinants in Appendix E. Thus no unspecified Jacobi-polynomial normalization is used.

Even-odd separation gives $D_{2L}^0 = A_m(L)A_{m+1}(L)$ and $D_{2L+1}^0 = A_m(L+1)A_{m+1}(L)$. Combining (29.10)–(29.13) yields

$$U_{2L}^0 = 2^{-L} T_m(L) A_{m+1}(L), \quad U_{2L+1}^0 = 2^{-L} A_m(L+1) T_{m+1}(L). \quad (29.14)$$

Proof of Theorem 29.1. Insert (29.14) and (29.11) into (29.8), with $h = 2m$. At even order the scalar factors multiply to one; at odd order they multiply to $(-1)^m$. Substitution of (29.10) gives all four displayed formulas, including the factors two. Their degrees sum to $2m^2$. Their values are real, so conjugation gives the identical formulas at $-i$. $\square$

30 The exceptional cancellation is exactly three orders

Let $F_r(n)$ be the polynomial obtained from (29.3)–(29.6) by substituting $L = (n - r)/4$. Thus $F_r(n) = \mathcal{H}_{4m+1}(n; i)$ on the residue $n \equiv r \pmod{4}$. Remove the odd-residue signs by setting

$$G_0 = F_0, \quad G_2 = F_2, \quad G_1 = (-1)^m F_1, \quad G_3 = (-1)^m F_3.$$

In this section these polynomials are expressed in the centered variable $x = n + 2m + 1/2$. All four have degree $\delta = 2m^2$ and the same positive leading coefficient

$$\Lambda_m = \frac{2^{-2m^2}}{\prod_{1 \leq u \leq v \leq m} (u+v-1) \prod_{1 \leq u \leq v \leq m-1} (u+v)^2 \prod_{1 \leq u \leq v \leq m} (u+v)}. \quad (30.1)$$

For G_1 , the ratio to the displayed leading coefficient is $2(m)_m/(m+1)_m = 1$. Reflection gives

$$G_3(x) = G_0(-x), \quad G_2(x) = G_1(-x). \quad (30.2)$$

One can verify these identities from the factors. Alternatively, (26.3) at $k = 4m + 1, t = i$ reads $\tilde{\mathcal{H}}_k(-n - k; i) = (-1)^m \tilde{\mathcal{H}}_k(n; i)$ for integer n ; it swaps residues 0, 3 and 1, 2. Since each residue agrees with a polynomial at infinitely many integers, this proves (30.2) as a polynomial identity.

30.1 The relevant discrete Fourier coefficient

For any period-four polynomial representation, $F_r(n) = \sum_{j=0}^3 i^{jr} A_j(n)$, so $A_j(n) = \frac{1}{4} \sum_{r=0}^3 i^{-jr} F_r(n)$. The potentially exceptional phase is $\lambda_{\text{bad}} = (-1)^{m-1}$. In either parity of m , its grouped polynomial is

$$P_{\text{bad}}(x) = \frac{1}{4}(G_0 - G_1 + G_2 - G_3) = \frac{1}{2}(\text{odd } G_0 - \text{odd } G_1), \quad (30.3)$$

where $\text{odd } G = (G(x) - G(-x))/2$. Thus three consecutive candidate odd-degree coefficients can be inspected without expanding the full high-degree products.

30.2 Root power sums through degree five

Write $G_j(x) = \Lambda_m \prod_{u \in \mathcal{R}_j} (x + u)$ for $j = 0, 1$. Let $c = 2m + 1/2$ and let a triangular set of size s mean $1 \leq u \leq v \leq s$. The two offset multisets are

$$\begin{aligned} \mathcal{R}_0 : \quad & 2(u+v) - 2 - c \quad (s = m), \\ & 2(u+v) - c \quad (s = m-1), \text{ twice}, \\ & 2(u+v) - c \quad (s = m); \\ \mathcal{R}_1 : \quad & 2(u+v) + 1 - c \quad (s = m-1), \\ & 2(u+v) - 1 - c \quad (s = m-1), \\ & 2(u+v) - 1 - c \quad (s = m), \text{ twice}. \end{aligned}$$

Put $p_j^{(r)} = \sum_{u \in \mathcal{R}_r} u^j$, $\bar{p}_j = (p_j^{(0)} + p_j^{(1)})/2$, and $\Delta p_j = p_j^{(0)} - p_j^{(1)}$. Direct summation gives

j	$\bar{p}_j$	Δp_j
1	m	0
2	$m^2(8m^2 + 1)/6$	$2m$
3	$m(16m^2 - 9)/4$	$3m^2$
4	not needed	$m(16m^2 - 11)$
5	not needed	$\frac{5}{2}m^2(8m^2 - 5)$.

(30.4)

These are finite polynomial identities, not asymptotic approximations. A reproducible way to check every entry is to use

$$T_s(b, j) = \sum_{v=1}^s \sum_{u=1}^v (2(u+v) + b - c)^j = \sum_{h=0}^j \binom{j}{h} 2^h (b - c)^{j-h} \sum_{v=1}^s \sum_{r=0}^h \binom{h}{r} v^{h-r} \sum_{u=1}^v u^r. \quad (30.5)$$

The two power sums are respectively $T_m(-2, j) + 2T_{m-1}(0, j) + T_m(0, j)$ and $T_{m-1}(1, j) + T_{m-1}(-1, j) + 2T_m(-1, j)$. The usual finite sums of powers through degree six evaluate the right side of (30.5) and give (30.4) for every integer $m \geq 1$. The included symbolic verifier repeats this polynomial calculation with m left as a symbol.

Let $e_j^{(r)}$ be the elementary symmetric functions of the respective multisets. Newton's identities yield

$$\Delta e_1 = 0, \quad \Delta e_3 = 0, \quad \boxed{\Delta e_5 = -m^2(m^2 - 1)}. \quad (30.6)$$

For an explicit algebra check, $e_3 = (p_1^3 - 3p_1p_2 + 2p_3)/6$, while

$$120e_5 = p_1^5 - 10p_1^3p_2 + 15p_1p_2^2 + 20p_1^2p_3 - 20p_2p_3 - 30p_1p_4 + 24p_5.$$

Subtracting between the two multisets, the terms containing $\bar{p}_2$ cancel. What remains is

$$120\Delta e_5 = 40m^4 - 40m\bar{p}_3 + 30m^2 = -120m^2(m^2 - 1),$$

which proves the last identity without relying on an unevaluated computer output.

Theorem 30.1 (Exact exceptional degree). *For $m \geq 2$, the polynomial of phase $(-1)^{m-1}$ in $\mathcal{H}_{4m+1}(n; i)$ has degree $2m^2 - 5$ and leading term*

$$-\frac{\Lambda_m}{2} m^2 (m^2 - 1) x^{2m^2-5}, \quad x = n + 2m + 1/2. \quad (30.7)$$

For $m = 1$ that phase polynomial is identically zero.

Proof. Since δ is even, the odd parts of G_0, G_1 have their first three possible degrees $\delta-1, \delta-3, \delta-5$. Equation (30.6), inserted into (30.3), cancels the first two and gives the nonzero third coefficient (30.7). For $m = 1$, the polynomials have degree two, and their odd linear terms agree, so (30.3) vanishes. $\square$

Completion of Theorem 24.1. Lemma 26.1 settles every unique-maximum residue. Reflection (26.5) settles the leading cancellation criterion in the single possible tied residue. Proposition 28.2 gives an exact one-degree loss except at the stated fourth roots, including the separately treated constant disappearance case. In the exceptional family $D = 2m^2 - 2$; Theorem 30.1 gives exponent $1 + (2m^2 - 5) = D - 2$ for $m \geq 2$ and zero for $m = 1$. Conjugation treats $-i$. Finally Lemma 25.2 converts these exact grouped degrees to the reduced denominator and minimal recurrence. This proves every case of the theorem. $\square$

31 Consequences and explicit forms

31.1 Every complex parameter, not just roots of unity

Corollary 31.1 (Complete complex-parameter classification). *For $k = 1$ the reduced denominator is $1 - z$ for all t . For $k \geq 2$:*

$$\mathcal{Q}_{k,t}(z) = \begin{cases} 1 - z, & t = 0, \\ (1 - z)^{a(k-a)+1}, & t = 1, \\ (1 - z)^{ab+1}(1 + z)^{(a-1)(b+1)+1}, & t = -1, \\ \prod_{q=0}^M (1 - t^q z)^{d_q+1}, & t \neq 0 \text{ is not a root of unity}, \\ \prod_{\rho=0}^{\ell-1} (1 - t^\rho z)^{e_\rho}, & t \text{ has order } \ell \geq 3, \end{cases} \quad (31.1)$$

where the last exponents are given in Theorem 24.1.

Proof. At $t = 0$ the Schur alphabet has exactly a nonzero entries, all one, and the rectangular tableau is unique. At $t = 1$ the one-cluster formula in Lemma 25.1 has degree $a(k - a)$. At $t = -1$ there are two nodes: 1 of multiplicity $k - 1$ and -1 of multiplicity one. The allocations select respectively a or $a - 1$ entries from the first cluster. Their degrees are ab and $(a - 1)(b + 1)$, and their distinct phases have nonzero leading coefficients. Away from roots of unity all products t^q are distinct, so (25.7) and Lemma 25.2 apply directly. The remaining case is the main theorem. The first three cases and the generic real case agree with the preceding repository report. $\square$

[Added 30 September 2026, batch 66: the preceding repository report is Part II; the cases $t = 0, \pm 1$ are its Theorem 19.2, and the generic case is its Theorem 12.2, stated there over $\mathbb{Q}(t)$ and at real $t \notin \{0, \pm 1\}$. The line for complex t that is not a root of unity extends that theorem to the complex plane.]

31.2 Compact formulas at primitive fourth roots

Corollary 31.2 (Odd shifts at $t = i$). *For $m \geq 1$, put $\delta = 2m^2$. Then*

$$\mathcal{Q}_{4m+1,i}(z) = (1 - (-1)^m z)^{\delta+1} (1 + (-1)^m z)^{\max(0, \delta-4)} (1 + z^2)^\delta. \quad (31.2)$$

For $m \geq 0$, put $\eta = 2m(m + 1)$. Then

$$\mathcal{Q}_{4m+3,i}(z) = (1 - z^2)^{\eta+1} (1 - (-1)^m i z)^{\eta+1} (1 + (-1)^m i z)^{\max(0, \eta-2)}. \quad (31.3)$$

At $-i$ take complex conjugates.

Proof. For $k = 4m + 1$, the index nearest the center in the central residue is $q = 2m$, of degree δ . The two adjacent residues have maximal degree $\delta - 1$, and the opposite real phase has the exceptional exponent $\max(0, \delta - 4)$. For $k = 4m + 3$, the two real phases and the central imaginary phase have maximal degree η ; the opposite imaginary phase has degree $\eta - 3$ after ordinary cancellation. The small case $m = 0$ has no such phase and is covered by the maximum with zero. $\square$

For example, Theorem 29.1 at $m = 1$ gives particularly simple integer values:

$$\begin{aligned} \mathcal{H}_5(4L; i) &= (2L + 1)(L + 1), & \mathcal{H}_5(4L + 1; i) &= -2(L + 1)^2, \\ \mathcal{H}_5(4L + 2; i) &= 2(L + 1)^2, & \mathcal{H}_5(4L + 3; i) &= -(2L + 3)(L + 1). \end{aligned}$$

The phase $+1$ cancels identically, not merely in its highest coefficient. The denominator is $(1+z)^3(1+z^2)^2$.

31.3 Quasipolynomial periods and exact reconstruction

Corollary 31.3 (Quasipolynomial structure). *At a root of unity of order $\ell \geq 3$, $\mathcal{H}_k(n; \zeta)$ is a quasipolynomial of period dividing ℓ . Its least quasipolynomial period is*

$$\text{lcm}\{\text{ord}(\zeta^\rho) : e_\rho > 0\}. \quad (31.4)$$

On each residue it is a polynomial of degree at most $\max_\rho(e_\rho - 1)$.

Proof. Each surviving mode is a polynomial times $\zeta^{\rho n}$, which is periodic in n . Conversely a quasipolynomial of period s has only s th roots of unity as its exponential phases, by finite Fourier inversion on its residue polynomials. Independence in Lemma 25.2 forces every surviving phase to have order dividing s . $\square$

This gives an exact reconstruction procedure, not just an existence result. Compute the exponents from the main theorem, then either compute the canonical confluent polynomials and combine their equal phases, or compute sufficient initial values by Jacobi–Trudi and interpolate the bounded-degree polynomial on each residue. The proof supplies the a priori bound that makes such interpolation a certified finite task in exact arithmetic. The Berlekamp–Massey experiments included with this article serve as independent checks, rather than as the source of that bound.

For the unnormalized determinants $D_k(n; \zeta)$, the multiplier in (23.4) is a nonzero periodic quadratic phase, of period dividing 4ℓ . Thus they too are quasipolynomials, of period dividing 4ℓ . Their *minimal* recurrences need a further Fourier combination and are not asserted to equal (24.7).

31.4 A quadratic recurrence-order law

Corollary 31.4 (Fixed cyclotomic order). *Let $N_{k,\ell} = \sum_\rho e_\rho$ and fix $\ell \geq 3$. For $M = k - 1 \geq \ell$,*

$$\frac{\ell(M^2 - \ell^2)}{8} - 3 \leq N_{k,\ell} \leq \frac{\ell M^2}{8} + \ell. \quad (31.5)$$

In particular

$$N_{k,\ell} = \frac{\ell(k-1)^2}{8} + O_\ell(1) \quad (k \rightarrow \infty). \quad (31.6)$$

Proof. In each residue choose an index nearest to $M/2$. Its distance is at most $\ell/2$, and $M \geq \ell$ ensures that a nearest choice lies in $[0, M]$. Since $q(M-q)/2 = M^2/8 - (q-M/2)^2/2$, the baseline exponent lies between $(M^2 - \ell^2)/8$ and $M^2/8 + 1$. Sum over residues and use the fact that the total correction is at most three. $\square$

Generically the order is $\sum_{q=0}^{k-1} (1 + d_q) = k^3/12 + O(k^2)$. Therefore a fixed cyclotomic order changes the growth from cubic in the shift to quadratic. This is a statement about recurrence order, not a bit-complexity assertion: large exact coefficients and initial-value computation still have costs.

32 Verification, dependency audit, and formalization prospects

32.1 What the included computations check

The archive contains an executable Python verifier, exact output records, and a source audit. The program uses integer arithmetic, rational arithmetic, Gaussian-rational arithmetic, symbolic polynomial arithmetic, and finite fields; it never estimates pole orders from floating-point roots.

The independent checks comprise the following finite tasks.

1. Canonical confluent polynomials for $2 \leq k \leq 7$ at $t = 2, 3$, constructed by Laplace expansion rather than by the claimed coefficient formulas. All 54 modes are checked for degree, leading coefficient, centered first coefficient, and reflection. An additional 22 modes with other cluster multiplicities check the general coefficient formula.
2. The four product formulas for $1 \leq m \leq 4$, $0 \leq n < 20$, checked against Jacobi–Trudi over $\mathbb{Q}(i)$: 80 exact values.
3. The power-sum and Newton identities in (30.4)–(30.6), checked with m left symbolic. This is polynomial identity verification, not a list of numerical samples.
4. Finite-field recurrence reconstruction for $2 \leq k \leq 17$ and $3 \leq \ell \leq 17$: 240 cases. For each order a prime $p > 10^6$ with $p \equiv 1 \pmod{\ell}$ is used, and a primitive ℓ th root is formed in $\mathbb{F}_p$. At least twice the baseline denominator degree plus eight sequence terms are computed. Berlekamp–Massey output is factored into the predicted linear factors and compared with the exponent list.
5. A separate finite check compares the original normalized Hankel formula with Schur evaluation at rational and Gaussian-rational parameters. This includes $1 \leq k \leq 8$, $0 \leq n \leq 5$, and $t = 2, -2, i$: 144 comparisons. The generated JSON report records these ranges and counts.

The finite-field test is a check of specialized finite sequences, not an independent characteristic-zero proof and not a theorem for untested parameters. Bad reduction could in principle conceal or create a cancellation; the all-index arguments above, not the modular experiments, establish the characteristic-zero classification. No claim is made that this script certifies the repository as a whole.

[Added 30 September 2026, batch 66: in this report the verifier is shipped as `code/03-cyclotomic-poles-verify.py`, its helper module as `code/03-cyclotomic-poles-modular.py`, and its records as `data/03-cyclotomic-poles-verification.json` (with the console log `data/03-cyclotomic-poles-verification.log`) and `data/03-cyclotomic-poles-modular_checks.json`. The file `data/verification.json` is Part I's record (Section 11.1). The verifier has no output option: it always writes `verification.json` and `modular_checks.json` into the `data/` directory beside its own `code/` directory, and it imports its helper under the delivered name `modular`. It must therefore be run only on a copy with the delivered layout (Appendix G). Run that way on 30 September 2026 (Python 3.13.5, SymPy 1.14.0), it passed every check; its `modular_checks.json` equals the record after CRLF→LF (the script writes CRLF on Windows), and its `verification.json` differs only in the fields `utc` and `elapsed_seconds`. During the intake the pole exponents were also recomputed from scratch for $3 \leq \ell \leq 8$, $1 \leq k \leq 15$ (Jacobi–Trudi in exact $\mathbb{Z}[x]/\Phi_\ell(x)$, lowering each phase exponent while the operator still annihilates): all 90 cases agree with Theorem 24.1.]

32.2 Logical dependencies

The new spectral argument depends on a short, explicit list of ingredients. The bialternant identity and Vandermonde confluence give the canonical polynomials. Reflection and the elementary concavity lemma restrict their collisions. The first-coefficient theorem resolves every ordinary

collision. The fourth-root Gram reduction, beta determinant calculation, and Newton identities resolve the only exception. Finally exponential-polynomial independence proves minimality, not just annihilation.

The spectral theorem can be formalized without first constructing the whole Hankel-moment development: take $s_{(n^a)}(X_k(t))$ as the initial definition. To transfer it to the original moments one needs exactly (23.4). The fourth-root proof as written uses the same identification together with the directly verified odd-shift block elimination in Section 29. All assumptions are thus explicit; no numerical recurrence is promoted to an all-index theorem.

32.3 A practical proof-assistant route

A realistic first stage is entirely algebraic: formalize exponential polynomials, their annihilators, and exact minimality for distinct nonzero phases. Next prove the quadratic maximum lemma and implement the integer exponent function from Theorem 24.1. These steps do not require analysis or infinite series.

The main algebraic effort is then the canonical confluent determinant and its first two coefficients. The proof in Section 27 can be translated into finite polynomial identities by clearing all differences $\alpha_i - \alpha_j$. Formal power series in ϵ are one route; a finite Taylor-jet ring truncated after degree $d + 1$ avoids an analytic limit. The divisibility of alternating polynomials by Vandermondes should be kept as a reusable lemma.

For the exceptional case, formalize the beta determinant in the factorial specializations actually used, rather than importing real integration. The triangular power sums and Newton identities are elementary polynomial identities and are suitable for ring normalization with generated, independently checkable coefficient tables. Only after those components are in place is it useful to attach the original Hankel–Schur interface. No Lean or Rocq implementation of these new results is included or claimed.

33 Further research questions

The classification suggests several concrete follow-up problems. The questions below are not used in the proofs.

[Added 30 September 2026, batch 66: none of these questions repeats one of Part II’s. Section 33.1 and Section 33.9 continue Part II’s topic “Broader noncollision criteria and certified algorithms” (Section 21.6), on which Theorem 27.1 bears; Section 33.6 is related to Part II’s Section 21.1 on Cigler’s Conjecture 17, as it says; Section 33.5 moves the parameter near a root of unity, whereas Part II’s “uniform crossover as the poles merge” moves it to $t = 1$.]

33.1 More than one middle variable

Replace the alphabet $(1^{[a]}, t, (t^2)^{[b]})$ by $(1^{[a]}, t^{[c]}, (t^2)^{[b]})$ with $c \geq 2$. Several allocations can then have the same product even before a root-of-unity specialization. Determine their combined degrees and the exact minimal recurrence. The first-coefficient formula remains available, but the unique-allocation mechanism (25.6) disappears. A useful first target is $c = 2$, where one can enumerate the competing allocations explicitly.

33.2 A general bound on cancellation depth

For arbitrary repeated-node rectangular Schur polynomials, can the maximal degree lost when products collide be bounded in terms of the number of clusters, rather than the total multiplicity?

The present family has loss at most three, and its exceptional loss is detected by a fifth symmetric coefficient. A structural theorem would explain when a finite number of Taylor coefficients suffices to establish minimality.

33.3 Higher coefficients of a canonical spectral mode

Extend Theorem 27.1 to the second and third centered coefficients in closed cluster notation. Expanding the finite functional proof further introduces quadratic and cubic characteristic polynomials in the selected sums. Such formulas could replace special-purpose product evaluations and make multi-cluster collision tests systematic.

33.4 Why do fourth roots force the additional cancellation?

The proof here is an explicit algebraic one: the first and third elementary symmetric differences vanish and the fifth does not. Is there a character- theoretic or geometric explanation that predicts this pattern before calculation? In particular, does a symmetry of the relevant rectangular Schur module identify the missing low-order terms in the bad phase? Any such explanation should recover the coefficient $-\Lambda_m m^2(m^2 - 1)/2$, not merely predict vanishing.

33.5 Detuning a root of unity

Investigate the transition when $t = \zeta e^{u/n}$ with u in a bounded complex set. Distinct generic phases then remain separated on a scale visible to the polynomial prefactors, while at $u = 0$ they cancel. Uniform asymptotic formulas should distinguish the ordinary one-order transition from the exceptional three-order one. This is a separate moving-parameter problem; fixed- t recurrence minimality alone does not solve it.

33.6 Numerators and zeros at cyclotomic parameters

The denominator is now known exactly. Determine the reduced numerator, its reciprocity, and its zero set at roots of unity. Which of its coefficients admit an enumerative or signed representation? This problem is related to, but should not be conflated with, the full coefficient assertions in Cigler's Conjecture 17 and the corrected generic numerator claims in the repository.

33.7 Original, unnormalized Hankel determinants

Classify the minimal recurrence of $D_k(n; \zeta)$ itself, including the periodic quadratic phase in (23.4). Its Fourier transform can mix the surviving $\mathcal{H}$ phases, so the answer need not be a simple relabeling of the denominator obtained here. Exact Gauss-sum identities may provide a useful description of this additional mixing.

33.8 Joint growth of the shift and cyclotomic order

Corollary 31.4 fixes ℓ . Determine a uniform recurrence-order asymptotic when ℓ/k approaches a nonzero limit, and describe the transition between the quadratic and cubic regimes. The finite maximum formula reduces this to a lattice-point problem with a small cancellation correction, making this an accessible next theorem rather than an unstructured asymptotic conjecture.

33.9 Certified recurrence generation

Produce a proof-producing implementation that accepts k and a cyclotomic order and returns the exact denominator, its minimality certificate, and a certified evaluator. A first practical deliverable is the exponent function plus a checked proof of the concavity and reflection reductions; the later confluence and quartic certificates can be attached modularly.

34 Conclusion

The distinction between an annihilating denominator and a reduced denominator is decisive at roots of unity. For Cigler's normalized Hankel polynomials, strict concavity reduces all possible leading cancellations to one residue. Reflection gives an exact parity test, a general centered coefficient formula settles every ordinary case, and an explicit fourth-root product calculation proves the unique three-order exception. The resulting theorem classifies the minimal recurrence at every complex parameter and closes the precise cyclotomic gap identified in the inspected ProveIt report.

The further questions concern genuinely additional work: larger middle multiplicity, deeper multi-cluster collisions, moving-parameter transitions, numerators, and proof-assistant integration. They are not needed to complete the classification proved here.

E An elementary beta determinant and the product evaluations

This appendix supplies the normalizations used in (29.10) and (29.13). Although these evaluations belong to classical determinant and orthogonal-polynomial theory [10], the required determinant can be proved directly.

Lemma E.1 (Beta-moment determinant). *For positive real A, B and $L \geq 0$,*

$$\det(B(A + i + j, B))_{0 \leq i, j < L} = \prod_{j=0}^{L-1} \frac{j! \Gamma(A + j) \Gamma(B + j)}{\Gamma(A + B + L + j - 1)}. \quad (\text{E.1})$$

Proof. Set $X_i = A + i$. Factor $B(X_i, B)$ from row i . The remaining entry in column j is $(X_i)_j / (X_i + B)_j$. For arbitrary distinct X_i one has

$$\det \left(\frac{(X_i)_j}{(X_i + B)_j} \right)_{i, j < L} = \frac{\prod_{j=0}^{L-1} (B)_j \prod_{i < r} (X_r - X_i)}{\prod_{i=0}^{L-1} (X_i + B)_{L-1}}. \quad (\text{E.2})$$

To prove this, multiply row i by its denominator on the right. Column j becomes the polynomial $f_j(X) = (X)_j (X + B + j)_{L-1-j}$ of degree $L - 1$. Its evaluation determinant is its coefficient determinant times the Vandermonde. Evaluate at $X_i = -i$: the matrix is lower triangular, with diagonal $(-1)^i i! (B)_{L-1-i}$. The Vandermonde at these points is $(-1)^{\binom{L}{2}} \prod_i i!$. Their quotient is $\prod_j (B)_j$, proving (E.2). Substitute $X_i = A + i$, use $\prod_{i < r} (r - i) = \prod_j j!$, and simplify the gamma factors. The empty determinant is immediate. $\square$

The binomial and Catalan moments have beta representations

$$\binom{2j}{j} = \frac{4^j}{\pi} B(j + 1/2, 1/2), \quad C_j = \frac{2 \cdot 4^j}{\pi} B(j + 1/2, 3/2). \quad (\text{E.3})$$

They follow from the gamma recurrence and the half-integer formula. Applying (E.1) after factoring the powers of four gives

$$A_s(L) = 4^{Ls+L(L-1)} \pi^{-L} \prod_{j=0}^{L-1} \frac{j! \Gamma(s+j+1/2) \Gamma(j+1/2)}{\Gamma(s+L+j)}, \quad (\text{E.4})$$

$$T_s(L) = 4^{Ls+L(L-1)} (2/\pi)^L \prod_{j=0}^{L-1} \frac{j! \Gamma(s+j+1/2) \Gamma(j+3/2)}{\Gamma(s+L+j+1)}. \quad (\text{E.5})$$

Their quotient immediately yields the third formula in (29.10). For an elementary check of the first two, the factorial products give $A_1(L) = 2^L$, $T_1(L) = 1$, and

$$\begin{aligned} \frac{A_{s+1}(L)}{A_s(L)} &= 4^L \frac{(s+1/2)_L}{(s+L)_L} = \frac{(2L+s)_s}{(s)_s}, \\ \frac{T_{s+1}(L)}{T_s(L)} &= 4^L \frac{(s+1/2)_L}{(s+L+1)_L} = \frac{(2L+s+1)_s}{(s+1)_s}. \end{aligned}$$

The right sides are exactly the successive triangular factors in (29.1) and (29.2). Induction on s proves (29.10) for all $s \geq 1$, $L \geq 0$.

Finally, the Christoffel determinant identity follows by appending the row $1, y, \dots, y^L$ to the moment matrix: the resulting determinant divided by the size- L moment determinant is the monic orthogonal polynomial. Putting $y = 1$ and subtracting adjacent columns identifies its value with the determinant ratio for weights $(1-y)w$ and w . For beta weight, (E.1) gives

$$\pi_L(1) = \frac{D_L(A, B+1)}{D_L(A, B)} = \frac{(B)_L}{(A+B+L-1)_L}. \quad (\text{E.6})$$

Scaling $y = x^2/4$ proves (29.13). Positivity of the beta weights ensures all the moment determinants used to define these monic polynomials are nonzero. The final identities themselves are polynomial or rational identities and need no limiting interpretation.

F A minimal exponent algorithm

The following pseudocode is an exact integer algorithm for Theorem 24.1. It does not need a numerical approximation to a root of unity.

```
cyclotomic_exponents(k, ell):
  require k >= 1 and ell >= 3
  M = k - 1
  a = floor(k/2)
  e = array of ell zeros
  for q in 0..M:
    e[q mod ell] = max(e[q mod ell], 1 + floor(q*(M-q)/2))
  if ell > M or (ell-M) is odd:
    return e
  q = (M-ell)/2
  D = floor((M*M-ell*ell)/8)
  cancels = (a+D is odd) if k is even else (D is even)
  if cancels:
    e[q mod ell] = max(0, D-2) if ell == 4 and k mod 4 == 1 else D
  return e
```

A direct implementation uses $O(k + \ell)$ elementary integer operations and $O(\ell)$ storage for a dense output array. These are arithmetic-operation counts; bit costs depend on the sizes of the integer indices. A sparse map avoids allocating ℓ entries when ℓ is large compared with k . The coefficient field for the actual recurrence is $\mathbb{Q}(\zeta)$, and its factorized form is usually preferable to expanding it.

G Package and reproduction

The package has a standalone `article.tex`, its compiled `article.pdf`, a `README.md`, a claim-status file, a source audit, and code and data directories. Compile the paper with

```
pdflatex -interaction=nonstopmode -halt-on-error article.tex
pdflatex -interaction=nonstopmode -halt-on-error article.tex
```

Run the exact verification suite from the package root with

```
python code/verify.py
```

It writes `data/verification.json` and `data/modular_checks.json`. The delivered record reports the versions and counts actually used. Successful finite checks are explicitly separated from the all-index mathematical proofs. No network access, proprietary computer algebra system, or repository checkout is required to rerun them.

[Added 30 September 2026, batch 66: this report ships the verifier, its helper module, the JSON records, the console log, the claim-status file, the source audit, `requirements.txt` and the Makefile under the prefixed names listed in Section 23.1. The archive's `article.tex`, `article.pdf` and `README.md` are not shipped: Part III and the report's `article.pdf` replace them. The commands above are the delivery's and must not be used from this report's root: there `python code/verify.py` runs Part I's verifier, which rewrites Part I's records in `data/`, and `pdflatex` rebuilds the whole report in place. The shipped `code/03-cyclotomic-poles-Makefile` runs exactly these commands (`make check` is `python code/verify.py`; its `clean` target deletes `article.aux`, `article.log`, `article.out` and `article.toc`), so it must not be run from the report root, and from `code/` its paths do not resolve. Part III's checks are reproduced on a copy with the delivered layout:

```
W=<scratch>; mkdir -p "$W/code"
cp code/03-cyclotomic-poles-verify.py "$W/code/verify.py"
cp code/03-cyclotomic-poles-modular.py "$W/code/modular.py"
uv run --no-project --with sympy==1.14.0 python "$W/code/verify.py"
```

with `<scratch>` a directory outside the report; the outputs land in `<scratch>/data/`.]

References

- [1] Johann Cigler, *Hankel determinants of middle binomial coefficients and conjectures for some polynomial extensions and modifications*, arXiv:2111.14492v3, 30 December 2021. The present target is Conjecture 16, pp. 21–22; the input moments are equation (6), p. 3. Part II uses especially Section 6 and printed pages 22–24, including equations (84)–(87) of Conjecture 18. <https://arxiv.org/abs/2111.14492v3>.
- [2] Johann Cigler and Christian Krattenthaler, *Hankel determinants of linear combinations of moments of orthogonal polynomials*, arXiv:2003.01676, 2020. <https://arxiv.org/abs/2003.01676>.

- [3] Vladimir Reshetnikov, *A manifest of docs/reports: Seventy-one independent mathematical research packages, classified by subject*, supplied file `manifest.tex`, dated 20 September 2026. See the entry *Product formulas for ballot-polynomial Hankel determinants under Catalan and ballot moments*. The manifest records report claims, not independent proof verification.
- [4] Manuel Kauers, *A Proof of Conjecture 16*, author-hosted lecture material, <https://www.algebra.uni-linz.ac.at/people/mkauers/publications/kauers26a.pdf>. Search-index excerpts consulted 20 September 2026 identify the target as A339987. The complete PDF was not successfully retrieved; this reference is used only to disambiguate the numbered problem, not as a mathematical proof source.
- [5] ProveIt research report, *Rectangular Schur Polynomials and Cigler's Conjecture 8*, 20 September 2026. Repository VladimirReshetnikov/ProveIt, commit `e9c09b12549ea4e3bea7cebc52b450b761f6f581`, file `SetTheory/Cardinals/docs/reports/hankel-determinants/catalan-and-ballot/cigler-conjecture-8-schur/article.tex`. This report concerns the different moment family $b_n(t)$.
- [6] M. Ciucu and C. Krattenthaler, *A factorization theorem for classical group characters, with applications to plane partitions and rhombus tilings*, in *Advances in Combinatorial Mathematics: Proceedings of the Waterloo Workshop in Computer Algebra 2008*, Springer, 2010, pp. 39–60. <https://arxiv.org/abs/0812.1251>.
- [7] P. Alexandersson, *Stretched skew Schur polynomials are recurrent*, Journal of Combinatorial Theory, Series A **122** (2014), 1–8. DOI: [10.1016/j.jcta.2013.09.009](https://arxiv.org/abs/1210.0377). <https://arxiv.org/abs/1210.0377>.
- [8] L. A. González-Serrano and E. A. Maximenko, *Bialternant formula for Schur polynomials with repeating variables*, Linear and Multilinear Algebra (2025). DOI: [10.1080/03081087.2025.2464639](https://arxiv.org/abs/2312.15680). <https://arxiv.org/abs/2312.15680>.
- [9] W. Fulton, *Young Tableaux: With Applications to Representation Theory and Geometry*, London Mathematical Society Student Texts 35, Cambridge University Press, 1997. Used for classical Schur identities and Schur-module characters.
- [10] C. Krattenthaler, *Advanced determinant calculus*, Séminaire Lotharingien de Combinatoire **42** (1999), Article B42q. <https://arxiv.org/abs/math/9902004>.
- [11] S. Chern and W. Shi, *Hankel determinants of Catalan-like sequences*, arXiv:2608.27208v1, 27 August 2026. <https://arxiv.org/abs/2608.27208>.