

Finite Residue Obstructions and Logarithmic-Depth Promotion in Hahn Transseries

An explicit Fredholm reduction for small
Euler-type differential perturbations

Research prepared for Vladimir Reshetnikov

29 September 2026

Abstract

At a fixed logarithmic depth, a formal differential equation can fail to have a solution even when all its nonresonant blocks are explicitly invertible. We give a concrete obstruction calculus for this failure. In the real Hahn field generated by $x, \log x, \dots, \log_n x$, the image of the m th derivative is characterized by exactly m residue moments. For an Euler-type operator $P(xD)$ perturbed by differential terms whose coefficients uniformly lower the exponent of x , all solvability conditions reduce to a finite real matrix. Its entries involve only finitely many perturbation steps, with an explicit cutoff determined by the real indicial roots and the support decrease. Its rank measures exactly the loss of homogeneous solutions and the remaining inhomogeneous obstructions at the original depth.

Adjoining one further logarithm removes every obstruction in this class. More precisely, if P has s distinct real roots, a particular solution has degree at most s in the new logarithm, and every homogeneous solution has degree at most $s - 1$. These bounds depend on the number of distinct roots, not their multiplicities. A family $\prod_{j=0}^{s-1} (xD - j) + (x \log x \cdots \log_n x)^{-1}$ attains both bounds and has a single Jordan-shift obstruction matrix. For its second-order member we construct actual real solutions and give an explicit error bound for the forced $\log \log x$ block, including a signed factorial-series remainder. The paper provides conventional proofs, finite exact verification scripts, a repository crosswalk, and a program of further questions.

Scope and status. This is an unrefereed, self-contained research manuscript, not a Lean-verified development. Classical Hahn support calculus, block inversion, and perturbative elimination are acknowledged explicitly. The new contribution proposed here is their concrete residue-matrix and sharp logarithmic-degree synthesis for the stated operator class. No exhaustive priority claim, general transseries solvability theorem, or general analytic summability theorem is asserted.

Contents

1	Research target, repository context, and main conclusions	3
2	The finite-depth Hahn model	4
3	A residue exact sequence and normalized integration	6
4	Higher derivatives and residue moments	8
5	The unperturbed Euler operator and its finite defect	9
6	Finite-dimensional reduction of a Hahn differential equation	10
7	One new logarithm suffices, with a sharp degree bound	13
8	A sharp family with an exact Jordan-shift obstruction	15
9	A second-order equation: formal resonance and real error bounds	17
10	Proof boundaries, checks, and formalization	21
11	Further research questions and concrete next targets	22
12	Conclusion	25
A	A compact abstract form of the reduction	26
B	Support facts used in the constructions	26
C	Package contents and reproducibility	27

Reading guide

For the main classification theorem, start with theorem 6.2 and the finite cutoff in theorem 6.3. The logarithmic-depth and degree theorem is theorem 7.2; its sharpness examples are theorems 8.1 and 8.2 and corollary 8.3.

The logical route is: normalized integration and residue moments (sections 3 and 4); the finite splitting of the Euler operator (section 5); perturbative elimination and support bounds (section 6); one-logarithm promotion and sharp examples (sections 7 and 8). These formal arguments do not require an analytic realization theorem.

The real-variable result can be read separately in section 9. Its endpoint is the explicit inequality (9.17), proved by a Volterra integral argument. Verification scope and the proposed Lean implementation are discussed in section 10; ten further research directions are developed in section 11.

1 Research target, repository context, and main conclusions

1.1 The question being answered

Formal antidifferentiation is often described by saying that one divides a monomial by its exponent, adding a logarithm at resonance. This description is useful but incomplete for a differential equation. Several resonant blocks may interact, the interaction may be mediated by infinitely many smaller terms, and it is not automatic that the resulting support is admissible. Nor does a formal solution by itself provide a real function with a certified error estimate.

This article isolates a class in which these questions have exact answers. The coefficients may be arbitrary real Hahn series of finite logarithmic depth; their supports need not be finitely generated. The essential restriction is that every perturbing coefficient decreases the *outer exponent of x* by a common positive amount. Inside a fixed x -block, arbitrary real powers of the logarithms remain available.

The central questions are as follows. Which forcing terms can be solved without increasing logarithmic depth? Can infinitely supported perturbations be reduced to finite obstruction data? How many new logarithms, and how large a power of the new logarithm, are necessary? These are formulated and resolved below for a precise class, rather than presented as a solution of an unspecified broad conjecture.

1.2 Relation to ProveIt

The inspected ProveIt development separates the transseries library from the Fabius-function library. Its current transseries directory contains asymptotic scales, well-based supports, polynomial-logarithmic blocks, and inversion material [1, 2]. The actual module `TransseriesBlockAntiderivative.lean` proves an explicit inverse for $\partial_L - c$ on polynomials when $c \neq 0$, and proves polynomial antidifferentiation at $c = 0$ [3]. Its finite inverse is

$$(\partial_L - c)^{-1}f = - \sum_{k=0}^{\deg f} c^{-k-1} \partial_L^k f. \quad (1.1)$$

The documentation carefully distinguishes these results from the remaining construction of a concrete Laurent/Hahn ambient and associated uniqueness and evaluation statements [2].

We extend the mathematical setting from polynomial blocks to well-based Hahn blocks, replace the finite sum in (1.1) by a strongly summable operator series when needed, and retain the resonant defects instead of hiding them in an enlarged ambient field. This produces a finite matrix that decides whether the enlargement is actually necessary. We do not claim that the existing Lean results already prove the theorems of this article, and we have not modified or rebuilt the repository.

The provenance snapshot used for the repository references is

8813fb3a2cdd1702d3ea7f7ca5d63ae710630bef.

The transseries destination is `Analysis/Transseries/`, not the historical Fabius documentation directory. The investigation was targeted at the relevant documentation and modules; it was not an exhaustive audit of every source in the repository.

1.3 Classical ingredients and the proposed contribution

Hahn fields, strong series derivations, and general integration criteria have an established theory. In particular, Kuhlmann and Matusinski study strong derivations of Hardy type and

give a general surjectivity criterion [4]. The one-residue integration result below is an explicit finite-depth specialization of that surrounding theory, not a claim to have first discovered formal integration. Van der Hoeven's operator theory already supplies powerful implicit-function methods for generalized series [5], and his monograph develops a broad transseries differential algebra [6]. Our geometric operator inverses are deliberately elementary instances of support-contracting methods.

Likewise, passing from an invertible complement to a finite effective operator is standard perturbative elimination; it is related in structure to homological perturbation methods [7]. We prove the needed algebra directly. What is proposed here as a useful research extension is the *explicit combination* of residue moments, a sharp finite perturbation cutoff, exact fixed-depth kernel/cokernel dimensions, a distinct-root bound for the power of one newly adjoined logarithm, and a family attaining the bounds. The literature check establishes relevant precedents, not priority for every formulation in this synthesis.

1.4 Roadmap of the results

Write $E = xD$ and let P be a nonzero real polynomial. Let r be the sum of the multiplicities of its real roots and s their number. For

$$\mathcal{L} = P(E) + R, \quad R = \sum_{j=0}^{\deg P} a_j E^j, \quad d_0(a_j) \leq -\varepsilon < 0, \quad (1.2)$$

the conclusions are:

- (i) A canonically normalized block construction gives a finite matrix M and a surjective defect map b such that $\mathcal{L}u = f$ at depth n if and only if $Mc = b(f)$ is solvable. Both kernel and cokernel have dimension $r - \text{rank } M$.
- (ii) If Δ is the difference between the largest and smallest real roots, the matrix uses at most $\lfloor \Delta/\varepsilon \rfloor$ perturbation factors. This is an exact cutoff, not an asymptotic approximation.
- (iii) In the next logarithmic field there is a particular solution polynomial of degree at most s in the new logarithm; all homogeneous solutions are polynomial of degree at most $s - 1$. Explicit examples attain these bounds.

The formal theorems are in sections 6 to 8. An actual differential equation with a quantitative real-variable certificate is treated in section 9.

2 The finite-depth Hahn model

2.1 Monomials, support, and summability

For $n \geq 0$, introduce formal symbols

$$\ell_0 = x, \quad \ell_1 = \log x, \quad \dots, \quad \ell_n = \log_n x.$$

These names specify the intended differential structure, not an assertion that every infinite formal sum converges after real evaluation. For $a \in \mathbb{R}^{n+1}$ put

$$\ell^a = \prod_{j=0}^n \ell_j^{a_j}.$$

Order exponents lexicographically, with a_0 the most significant coordinate. Larger exponent vectors mean larger monomials. Define

$$\mathcal{H}_n = \left\{ \sum_{a \in S} c_a \ell^a : c_a \in \mathbb{R}, S \subset \mathbb{R}^{n+1} \text{ is reverse well-ordered} \right\}. \quad (2.1)$$

Reverse well-ordered means that every nonempty subset has a greatest element. Coefficients may vanish; support always means the set of nonzero coefficients. We use the large-to-small convention throughout. Reversing the exponent order gives the usual valuation convention for Hahn series.

A family $(f_i)_{i \in I}$ is *summable* if the union of its supports is reverse well-ordered and each monomial occurs in only finitely many members. Its sum is then defined coefficientwise. A linear operator is *strong* if it preserves summable families and commutes with their sums.

We use the standard Hahn support lemma: finite unions and finite sums of reverse well-ordered subsets of an ordered abelian group are reverse well-ordered, and a fixed exponent has only finitely many decompositions as a sum of elements from two fixed such supports. The positive-support version of Neumann's lemma also makes geometric inverses of small Hahn series legitimate. These facts give the usual field operations in (2.1); they are part of the classical support calculus [5, 6]. The arguments below specify the additional support bounds needed for every operator series used in the paper.

2.2 Outer blocks

For $n \geq 1$, let

$$L = \ell_1, \quad \mathcal{B}_n = \mathbb{R}((L^{\mathbb{R}} \ell_2^{\mathbb{R}} \cdots \ell_n^{\mathbb{R}})).$$

Thus $\mathcal{B}_n$ is a copy of $\mathcal{H}_{n-1}$ whose first variable is L . Each element of $\mathcal{H}_n$ has a unique block decomposition

$$f = \sum_{\alpha} x^{\alpha} f_{\alpha}, \quad f_{\alpha} \in \mathcal{B}_n. \quad (2.2)$$

The set of nonzero outer exponents is reverse well-ordered, and each block has reverse well-ordered lower support. Conversely these two properties imply reverse well-ordering of the full lexicographic support. Indeed, to find the greatest element of a nonempty subset, first choose its greatest outer exponent and then the greatest element of that fibre. The same argument proves the projection and fibre assertions in the other direction.

For $f \neq 0$, define

$$d_0(f) = \max\{\alpha : f_{\alpha} \neq 0\}, \quad d_0(0) = -\infty.$$

Then $d_0(fg) = d_0(f) + d_0(g)$ for nonzero products, and $d_0(f + g) \leq \max(d_0(f), d_0(g))$. These statements concern support only; they are not estimates for real-valued functions.

2.3 The strong derivative

Define on monomials

$$D\ell^a = \sum_{j=0}^n a_j \ell^{a - \mathbf{1}_{\{0, \dots, j\}}}. \quad (2.3)$$

In particular, $D\ell_0 = 1$ and

$$D\ell_j = (\ell_0 \ell_1 \cdots \ell_{j-1})^{-1} \quad (j \geq 1).$$

There are only $n+1$ possible support shifts in (2.3). Their finite union is admissible, and a fixed output monomial receives at most $n+1$ input contributions. Consequently D extends strongly to $\mathcal{H}_n$. The Leibniz rule holds on monomials and then on arbitrary series by summability of Hahn products.

Let $\delta = d/dL$ denote the same derivative construction on $\mathcal{B}_n$. Directly from (2.3),

$$D(x^\alpha A) = x^{\alpha-1}(\alpha + \delta)A, \quad (2.4)$$

$$E(x^\alpha A) = x^\alpha(\alpha + \delta)A, \quad E = xD. \quad (2.5)$$

Every term of δA has its L -exponent lowered by one. This elementary fact replaces analytic norm estimates inside a block.

Lemma 2.1 (Formal functional calculus inside a block). *For $F(z) = \sum_{k \geq 0} F_k z^k \in \mathbb{R}[[z]]$, the operator $F(\delta)A = \sum_{k \geq 0} F_k \delta^k A$ is well-defined and strong on $\mathcal{B}_n$. The substitution respects sums and products of formal power series. If $F(0) \neq 0$, $F(\delta)$ is invertible with inverse $(1/F)(\delta)$.*

Proof. If the largest L -exponent of A is B , then that of $\delta^k A$ is at most $B - k$. At any fixed L -exponent, only finitely many k can contribute. The union of supports is reverse well-ordered: above any fixed lower bound for the L -exponent only finitely many translated supports are involved. The same reasoning applies to a summable family because its union has a single upper bound B . It justifies rearranging the double sums in multiplication of formal operator series. Substituting the formal identity $F(z)(1/F(z)) = 1$ proves the inverse assertion. $\square$

Remark 2.2. This argument requires no positive lower bound on $|F(0)|$ as parameters vary from one outer block to another. Small denominators may matter for analytic realization, but do not invalidate the formal block construction. An analytic estimate must therefore not be inferred from this lemma.

3 A residue exact sequence and normalized integration

Put

$$\rho_n = (x\ell_1 \cdots \ell_n)^{-1}, \quad \text{Res}_n(f) = [\rho_n]f.$$

The coefficient of the constant monomial is denoted by $[1]f$.

Theorem 3.1 (The fixed-depth integration defect). *There is a unique strong linear operator $I_n : \mathcal{H}_n \rightarrow \mathcal{H}_n$ satisfying*

$$DI_n f = f - \text{Res}_n(f)\rho_n, \quad [1]I_n f = 0, \quad (3.1)$$

$$I_n Df = f - [1]f. \quad (3.2)$$

In particular,

$$0 \longrightarrow \mathbb{R} \longrightarrow \mathcal{H}_n \xrightarrow{D} \mathcal{H}_n \xrightarrow{\text{Res}_n} \mathbb{R} \longrightarrow 0 \quad (3.3)$$

is exact. A primitive of f lies in $\mathcal{H}_n$ exactly when $\text{Res}_n(f) = 0$. In $\mathcal{H}_{n+1}$ a primitive is

$$J_n f = I_n f + \text{Res}_n(f)\ell_{n+1}. \quad (3.4)$$

All primitives differ by a real constant.

Proof. For $n = 0$, integrate x^a as $x^{a+1}/(a+1)$ when $a \neq -1$, and send x^{-1} to zero. The exceptional coefficient is exactly Res_0 . Translation of support proves strongness, and the asserted identities follow termwise.

Suppose the result has been proved for the lower field $\mathcal{B}_n$. In (2.2), write $I^{(L)}$ for its normalized integration operator. Define the primitive of a block with $\alpha \neq -1$ by

$$I_n(x^\alpha A) = x^{\alpha+1} \sum_{k \geq 0} \frac{(-1)^k}{(\alpha+1)^{k+1}} \delta^k A. \quad (3.5)$$

The sum is the inverse of $\alpha + 1 + \delta$ from lemma 2.1. At $\alpha = -1$ define

$$I_n(x^{-1} A) = I^{(L)} A.$$

The missing lower residue is the coefficient of $\sigma_n = (L\ell_2 \cdots \ell_n)^{-1}$ in A , and its outer counterpart is $\rho_n = x^{-1}\sigma_n$. This proves the first identity in (3.2).

Each input outer exponent is shifted to exactly one possible output outer exponent, namely $\alpha + 1$. Within a nonresonant block the construction is strong by lemma 2.1, and within the exceptional block it is strong by induction. The projection/fibre criterion therefore proves strongness on the full field. No constant is produced in a nonresonant block, and the exceptional block has been normalized to have zero constant.

To compute the kernel of D , apply (2.5) to a putative constant. A block with outer exponent $\alpha \neq 0$ must vanish because $\alpha + \delta$ is invertible. The remaining outer-zero block lies in the kernel of δ , which is $\mathbb{R}$ by induction. Thus $\ker D = \mathbb{R}$. Directly from the monomial derivative, $\text{Res}_n(Df) = 0$: every possible input that could produce ρ_n has zero exponent in the coordinate being differentiated, and hence contributes zero. Hence $I_n Df$ and f differ by a constant; normalization identifies it as $[1]f$.

The functional Res_n is onto because $\text{Res}_n \rho_n = 1$, proving exactness. Finally $D\ell_{n+1} = \rho_n$. The formula (3.4) and the kernel calculation at depth $n+1$ prove the last assertions. Uniqueness of I_n follows from its differential identity and zero-constant normalization. $\square$

Editorial note (ProveIt, 2026-09-29). Part of theorem 3.1 is already Theorem `plt:thm:ext-tower-strict` of the canonical volume (`Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/transseries_and_inversion.tex`), which this article does not cite. That theorem works in $\mathcal{H}_r(\mathbb{K}, \Gamma)$ with $\mathbb{Z} \subseteq \Gamma \subseteq \mathbb{R}$ and the monomial $\mathbf{n}_r = L_0 L_1 \cdots L_r$, $L_0 = x$, $L_{j+1} = \log L_j$, so that $\mathbf{n}_n^{-1} = \rho_n$. Its equation `plt:eq:ext-residue`, $[\mathbf{n}_r^{-1}] \partial_r F = 0$ for all $F \in \mathcal{H}_r$, is the identity $\text{Res}_n \circ D = 0$, and its consequence that $L_{r+1} \notin \mathcal{H}_r$ is the statement here that ℓ_{n+1} is a primitive of ρ_n outside $\mathcal{H}_n$. Theorem 3.1 re-derives both for real coefficients and strengthens them to the exactness of (3.3): every f with $\text{Res}_n(f) = 0$ has a primitive in $\mathcal{H}_n$, given by the explicit operator I_n ; theorem 4.1 extends the criterion to D^m . Neither result bears on the canonical volume's open Remark `plt:rmk:ext-open-depth`, which concerns compositional inverses rather than differential operators.

Corollary 3.2 (Residue integration by parts). *For $f, g \in \mathcal{H}_n$,*

$$\text{Res}_n(fDg) = -\text{Res}_n(gDf). \quad (3.6)$$

Proof. Apply Res_n to $D(fg)$ and use theorem 3.1. $\square$

Example 3.3. At depth one, $\rho_1 = 1/(x \log x)$. Thus this monomial has no primitive in $\mathcal{H}_1$ and has primitive $\log \log x$ in $\mathcal{H}_2$. In contrast, $x^{-1}(\log x)^{-2}$ integrates inside $\mathcal{H}_1$ to $-(\log x)^{-1}$. The issue is a single exact coefficient, not merely the presence of a negative logarithmic power.

4 Higher derivatives and residue moments

Theorem 4.1 (The complete obstruction to an m th primitive). *For every integer $m \geq 1$,*

$$f \in D^m \mathcal{H}_n \iff \text{Res}_n(x^k f) = 0 \quad (0 \leq k < m). \quad (4.1)$$

The kernel is $\mathbb{R}[x]_{<m}$, and the cokernel has dimension m over $\mathbb{R}$. With $T = \ell_{n+1}$, a particular m th primitive is

$$Y_m(f) = \frac{1}{(m-1)!} \sum_{k=0}^{m-1} (-1)^k \binom{m-1}{k} x^{m-1-k} J_n(x^k f). \quad (4.2)$$

It has the form $U_m(f) + TV_m(f)$, with $U_m(f) \in \mathcal{H}_n$ and

$$V_m(f) = \frac{1}{(m-1)!} \sum_{k=0}^{m-1} (-1)^k \binom{m-1}{k} x^{m-1-k} \text{Res}_n(x^k f). \quad (4.3)$$

Thus an arbitrary number of repeated integrations requires at most the first power of one additional logarithm.

Proof. If $f = D^m u$, repeated use of (3.6) gives $\text{Res}_n(x^k D^m u) = (-1)^m \text{Res}_n(u D^m x^k) = 0$ for $k < m$.

We verify the Green formula. For $m = 1$, it is $Y_1 = J_n$, and $DY_1(f) = f$. For $m > 1$, differentiating a primitive factor in (4.2) contributes

$$\frac{x^{m-1} f}{(m-1)!} \sum_{k=0}^{m-1} (-1)^k \binom{m-1}{k} = 0.$$

Differentiating the power of x instead gives $Y_{m-1}(f)$, because

$$(m-1-k) \binom{m-1}{k} = (m-1) \binom{m-2}{k}.$$

Therefore $D^m Y_m(f) = f$. Substitution of (3.4) yields (4.3). If the moments vanish, $Y_m(f) \in \mathcal{H}_n$, proving sufficiency in (4.1).

For the kernel, induct on m . If $D^m u = 0$, then $D^{m-1} u$ is constant; subtract a suitable multiple of x^{m-1} and use the induction hypothesis. Finally, the moment functionals are independent: their values on $x^{-j} \rho_n$, $0 \leq j < m$, form the identity matrix. This gives cokernel dimension m . The distinct powers of x in (4.3) also show that $V_m(f) = 0$ exactly when all moments vanish. $\square$

Remark 4.2 (Why successive naive integrations obscure the answer). One may integrate m times and introduce a logarithm at each stage. Formula (4.2) proves that this is unnecessary. The entire defect is already a finite polynomial of residue moments multiplying one new logarithm. This uniform control is the reason that repeated indicial roots will not increase the logarithmic-degree bound in section 7.

4.1 Integration of polynomials in the new logarithm

We need a slightly stronger statement for later perturbative steps.

Lemma 4.3 (One-step degree increase). *On $\mathcal{H}_n[T] \subset \mathcal{H}_{n+1}$, with $T = \ell_{n+1}$, the derivative is surjective. It admits a linear right inverse that raises T -degree by at most one and is strong on each subspace of bounded T -degree. For every $m \geq 1$, D^m also admits such a right inverse, still raising T -degree by at most one, independently of m .*

Proof. Let the highest-degree term of a polynomial forcing be $a_d T^d$. The derivative of

$$I_n(a_d)T^d + \frac{\text{Res}_n(a_d)}{d+1}T^{d+1}$$

is

$$a_d T^d + d\rho_n I_n(a_d)T^{d-1}.$$

Subtract the second term from the remaining forcing and repeat at the next lower degree. The process ends after finitely many polynomial degrees, is linear, and uses only strong operations. Its degree increase is at most one. Apply the Green formula (4.2), now using this right inverse for each polynomial forcing $x^k f$, to obtain a right inverse of D^m with the same degree bound. The verification of that formula used only $DJ = \text{id}$, so applies unchanged. $\square$

5 The unperturbed Euler operator and its finite defect

Throughout the rest of the formal analysis let $n \geq 1$. Write the distinct real roots of a nonzero polynomial $P \in \mathbb{R}[z]$ as

$$\lambda_1 < \cdots < \lambda_s,$$

with multiplicities $m_1, \dots, m_s$. Set $r = \sum_{i=1}^s m_i$; if there are no real roots, $s = r = 0$. Nonreal roots are allowed. They do not generate real-power homogeneous monomials in the present field.

For a root λ , factor

$$P(\lambda + z) = z^{m_\lambda} Q_\lambda(z), \quad Q_\lambda(0) \neq 0. \quad (5.1)$$

On $x^\alpha \mathcal{B}_n$, the operator $P_0 = P(E)$ acts as $P(\alpha + \delta)$.

5.1 Kernel coordinates and residue coordinates

Let $V = \mathbb{R}^r$, with coordinates indexed by pairs (λ, k) where $0 \leq k < m_\lambda$. Define

$$Kc = \sum_{\lambda} \sum_{k=0}^{m_\lambda-1} c_{\lambda,k} x^\lambda L^k, \quad (5.2)$$

$$(\Lambda f)_{\lambda,k} = [x^\lambda L^k \ell_2^0 \cdots \ell_n^0] f, \quad (5.3)$$

$$(\Pi f)_{\lambda,k} = \text{Res}^{(L)}(L^k f_\lambda), \quad (5.4)$$

$$J e_{\lambda,k} = x^\lambda L^{-k} \sigma_n, \quad \sigma_n = (L \ell_2 \cdots \ell_n)^{-1}. \quad (5.5)$$

Here $\text{Res}^{(L)}$ is the residue functional of the lower field. The maps K, J have different purposes: K represents homogeneous solutions, whereas J represents obstructions. Direct coefficient extraction gives

$$\Lambda K = \text{id}_V, \quad \Pi J = \text{id}_V. \quad (5.6)$$

Theorem 5.1 (Explicit splitting of $P(E)$). *The kernel of P_0 is KV , and its image is $\ker \Pi$. There is a unique normalized strong linear operator $G : \mathcal{H}_n \rightarrow \mathcal{H}_n$ such that*

$$P_0 G = \text{id} - J \Pi, \quad G P_0 = \text{id} - K \Lambda, \quad (5.7)$$

$$\Lambda G = 0, \quad G J = 0. \quad (5.8)$$

It preserves each outer x -exponent block. Thus both the kernel and cokernel of P_0 have dimension r .

Proof. If $P(\alpha) \neq 0$, lemma 2.1 gives the inverse

$$[P(\alpha + \delta)]^{-1} = \frac{1}{P(\alpha + z)} \Big|_{z=\delta}. \quad (5.9)$$

At a real root λ , (5.1) gives the product $\delta^{m_\lambda} Q_\lambda(\delta)$. The second factor is invertible and commutes with δ . Therefore it maps $\text{im } \delta^{m_\lambda}$ onto itself: both it and its inverse commute with δ^{m_λ} . By theorem 4.1, the image is characterized by

$$\text{Res}^{(L)}(L^k f_\lambda) = 0 \quad (0 \leq k < m_\lambda).$$

The kernel is the space of polynomials in L of degree less than m_λ . Indeed, this is the kernel of δ^{m_λ} , and $Q_\lambda(\delta)$ is an invertible endomorphism of that finite polynomial space.

For arbitrary f , the series $f - J\Pi f$ satisfies all image conditions. Solve $P_0 u = f - J\Pi f$ blockwise using (5.9), and at a resonant block using $Q_\lambda(\delta)^{-1}$ and (4.2). Subtract $K\Lambda u$ to normalize. The result is the unique Gf with $\Lambda Gf = 0$. It is strong because the lower integration, residue, and functional-calculus operations are strong; there are only finitely many exceptional outer blocks.

This proves $P_0 G = \text{id} - J\Pi$. Applied to $P_0 f$, the same normalized uniqueness gives $GP_0 f = f - K\Lambda f$. If $f = Je$, then $f - J\Pi f = 0$, so $GJe = 0$. The construction never changes an outer exponent, proving the final support assertion. $\square$

Remark 5.2. The residue coordinates in (5.5) do not need factors of $Q_\lambda(\delta)^{-1}$. Such factors would give an alternative basis of obstruction functionals, but the commutation argument in the proof shows that the unweighted moment test already characterizes the image. This simplifies the matrix in the next section.

6 Finite-dimensional reduction of a Hahn differential equation

6.1 The support-contracting hypothesis

Let $d = \deg P$ and let

$$R = \sum_{j=0}^d a_j E^j, \quad a_j \in \mathcal{H}_n. \quad (6.1)$$

Suppose that a fixed $\varepsilon > 0$ satisfies

$$a_j \neq 0 \implies d_0(a_j) \leq -\varepsilon. \quad (6.2)$$

The derivatives act on the argument to their right, not on a_j . Since E preserves an outer block, multiplication gives

$$d_0(Rf) \leq d_0(f) - \varepsilon, \quad d_0(GRf) \leq d_0(f) - \varepsilon. \quad (6.3)$$

No restriction is imposed on logarithmic exponents inside the coefficient blocks. Taking $a_d \neq 0$ is permitted; the leading differential coefficient is still a nonzero perturbation of the leading constant coefficient of P .

Lemma 6.1 (The geometric inverse). *The operator*

$$\mathcal{N} = (\text{id} + GR)^{-1} = \sum_{q \geq 0} (-GR)^q \quad (6.4)$$

is well-defined and strong on $\mathcal{H}_n$. It is a two-sided inverse.

Proof. For a fixed f with $d_0(f) = A$, the q th term has outer exponent at most $A - q\varepsilon$. At any prescribed outer exponent only finitely many values of q are possible. The union of the outer supports is reverse well-ordered because their maxima tend to $-\infty$, and each of its fibres is a finite union of admissible supports. The argument is uniform on a summable family, using the greatest outer exponent of its union. Thus the series is strong. The finite geometric identity has remainder $(-GR)^{N+1}f$; every coefficient of that remainder vanishes for sufficiently large N , proving both inverse identities. $\square$

Define the *resonance matrix* and the *effective forcing* by

$$M = \Pi R \mathcal{N} K : V \longrightarrow V, \quad (6.5)$$

$$b(f) = \Pi f - \Pi R \mathcal{N} G f. \quad (6.6)$$

All compositions in these formulas are now defined, including their infinite operator series.

Theorem 6.2 (Finite-residue Fredholm alternative). *Under (6.2), the equation*

$$(P(E) + R)u = f \quad (6.7)$$

has a solution in $\mathcal{H}_n$ if and only if

$$Mc = b(f) \quad (6.8)$$

has a solution in $V = \mathbb{R}^r$. All solutions are given by

$$u = \mathcal{N}(Kc + Gf), \quad Mc = b(f). \quad (6.9)$$

The parameter is exactly $c = \Lambda u$. Moreover,

$$\ker(P(E) + R) \cong \ker M, \quad (6.10)$$

$$\text{coker}(P(E) + R) \cong V / \text{im } M, \quad (6.11)$$

$$\dim \ker(P(E) + R) = \dim \text{coker}(P(E) + R) = r - \text{rank } M. \quad (6.12)$$

These are algebraic dimensions; no Banach-space Fredholm assertion is intended.

Proof. Apply G to (6.7) and use (5.8):

$$u - K\Lambda u + GRu = Gf.$$

With $c = \Lambda u$, the invertibility from lemma 6.1 gives (6.9). Applying Π to the original equation gives $\Pi Ru = \Pi f$, which after substitution is precisely (6.8).

Conversely, suppose $Mc = b(f)$ and define u by (6.9). Since $\Lambda G = 0$, the equation $(\text{id} + GR)u = Kc + Gf$ implies $\Lambda u = c$. The residual $h = (P_0 + R)u - f$ then satisfies $Gh = 0$. The matrix equation says $\Pi h = 0$. The identity $P_0 G = \text{id} - J\Pi$ now gives $h = P_0 Gh + J\Pi h = 0$.

For the cokernel, first note that $bJ = \text{id}_V$, because $GJ = 0$. Thus b is surjective. For arbitrary u ,

$$\mathcal{N}G(P_0 + R)u = u - \mathcal{N}K\Lambda u,$$

and hence

$$b((P_0 + R)u) = \Pi R \mathcal{N} K \Lambda u = M\Lambda u.$$

The solvability equivalence already proved says that $b(f) \in \text{im } M$ exactly when $f \in \text{im}(P_0 + R)$. Consequently b induces the asserted cokernel isomorphism. Formula (6.9), together with $\Lambda u = c$, proves the kernel isomorphism. Rank-nullity in V completes the proof. $\square$

6.2 An exact finite perturbation cutoff

The solution reconstruction usually is an infinite Hahn series. The matrix itself is not an infinite perturbation problem.

Theorem 6.3 (Finite cutoff and triangular structure). *Suppose $r > 0$, and put*

$$\lambda_{\min} = \lambda_1, \quad \lambda_{\max} = \lambda_s, \quad \Delta = \lambda_{\max} - \lambda_{\min}, \quad Q = \left\lfloor \frac{\Delta}{\varepsilon} \right\rfloor.$$

Then

$$M = \sum_{q=1}^Q (-1)^{q-1} \Pi R (GR)^{q-1} K. \quad (6.13)$$

An empty sum is zero. A matrix entry can be nonzero only from a source root λ to a strictly smaller target root μ . Thus M is strictly block triangular and $M^s = 0$.

For $f \neq 0$ with $A = d_0(f)$, the correction in $b(f)$ similarly terminates after at most

$$Q_f = \max \left(0, \left\lfloor \frac{A - \lambda_{\min}}{\varepsilon} \right\rfloor \right) \quad (6.14)$$

perturbation factors.

Proof. A term starting in the source block x^λ and containing q factors of R can reach only outer exponents at most $\lambda - q\varepsilon$. The maps G preserve outer exponents, and Π reads only the real root blocks. If $q\varepsilon > \Delta$, no target root is reachable from any source root. This proves (6.13). Every nonzero term also has target strictly below source, proving the triangular assertion. A product of s such strict root decreases is impossible, so $M^s = 0$.

For the forcing correction replace the source bound $\lambda_{\max}$ by A ; Gf has no greater outer exponent than f . This gives (6.14). $\square$

Corollary 6.4 (A coefficient-support selection rule). *Let $S \subset (0, \infty)$ be the set of all outer drops $-\beta$ for which some coefficient a_j has a nonzero x^β block. A matrix block from λ to μ vanishes unless*

$$\lambda - \mu \in S + S + \cdots + S \quad \text{for some positive finite number of summands.} \quad (6.15)$$

In particular, if no positive difference of real roots lies in this positive additive semigroup, then $M = 0$.

Proof. Each perturbation factor contributes one outer drop from S , while all intervening operators preserve the outer exponent. The total drop along a contributing term must equal $\lambda - \mu$. $\square$

Remark 6.5 (Multiplicity and remaining dimension). Strict block triangularity implies at least $\max_i m_i$ homogeneous solutions at the original depth whenever there are real roots. Indeed, at any chosen root level i , the columns up to that level have image contained in the strictly lower row blocks, while the remaining columns span a space of dimension at most the total multiplicity above that level. Thus $\text{rank } M \leq r - m_i$ for every i . If there is only one distinct real root, $M = 0$, although not every forcing is solvable: the r residue-moment obstructions remain.

6.3 A rigorous support truncation certificate

Proposition 6.6. *For $y = Kc + Gf \neq 0$, let*

$$u_N = \sum_{q=0}^N (-GR)^q y, \quad u = \mathcal{N}y.$$

Then

$$d_0(u - u_N) \leq d_0(y) - (N + 1)\varepsilon. \quad (6.16)$$

Hence all coefficients with outer exponent greater than the right-hand side are exact after N steps.

Proof. Every term in the omitted tail contains at least $N + 1$ support decreases. Cancellation can only reduce, not increase, the greatest remaining outer exponent. $\square$

The bound (6.16) is useful for symbolic computation, but it is not a numerical error bound. It controls the outer exponent only and does not bound the real sizes or growth of the coefficient blocks. The analytic certificate in section 9 is obtained by a separate argument.

6.4 What is and is not finite

Equation (6.13) is finite in *perturbation depth*. Individual coefficient blocks may still be infinitely supported, and arbitrary real Hahn coefficients need not be given by computable data. Therefore the theorem is not an unconditional finite-time algorithm on all set-theoretically possible inputs.

For an effective representation with exact coefficient extraction, exact outer-exponent arithmetic, and the requisite block operators, the formula becomes an algorithm. For finite rational supports, the included verification program illustrates part of this regime. For arbitrary noncomputable real supports, it remains a mathematical finite-dimensional characterization. This distinction is essential: finite-dimensional output does not by itself make its entries computable.

7 One new logarithm suffices, with a sharp degree bound

Let $T = \ell_{n+1}$. The ring $\mathcal{H}_n[T]$ is naturally a differential subring of $\mathcal{H}_{n+1}$. Its elements have only finitely many nonnegative integer powers of T , although their coefficients can be arbitrary elements of $\mathcal{H}_n$.

Lemma 7.1 (A blockwise right inverse after promotion). *There is a linear right inverse $\tilde{G}$ of $P_0 = P(E)$ on $\mathcal{H}_n[T]$ with the following properties:*

- (a) *it preserves the outer x -exponent;*
- (b) *at a nonresonant outer exponent it does not increase T -degree;*
- (c) *at a resonant outer exponent it raises T -degree by at most one, regardless of the root multiplicity;*
- (d) *it is strong on every family with uniformly bounded T -degree.*

It may be normalized by subtracting its kernel coordinates.

Proof. On the lower polynomial ring $\mathcal{B}_n[T]$, the derivative still lowers the L -exponent by one, including the term $\delta T = \sigma_n$. Thus the functional calculus of lemma 2.1 remains valid on each bounded-degree polynomial space. Differentiation never increases T -degree.

At a nonresonant exponent use (5.9). At a root λ , use (5.1), first a right inverse of δ^{m_λ} from lemma 4.3 in the lower variable L , and then $Q_\lambda(\delta)^{-1}$. This raises degree by at most one, not by m_λ . Strongness and preservation of the outer block follow from the construction. Subtracting the finitely many coordinates in the kernel does not affect any of these bounds. $\square$

Theorem 7.2 (Logarithmic-depth promotion). *Let $\mathcal{L} = P(E) + R$ satisfy (6.2), and let s and r be as in section 5.*

For every $f \in \mathcal{H}_n$, there is a solution of $\mathcal{L}u = f$ in $\mathcal{H}_n[T]$ with

$$\deg_T u \leq s. \quad (7.1)$$

The homogeneous solution space in $\mathcal{H}_{n+1}$ has dimension r , and all its elements lie in $\mathcal{H}_n[T]$. If $s \geq 1$, they satisfy

$$\deg_T u \leq s - 1. \quad (7.2)$$

If $s = 0$, the homogeneous kernel is zero and every forcing has a unique solution already in $\mathcal{H}_n$.

Consequently the matrix test of theorem 6.2 gives the exact answer to whether depth n suffices. Whenever it fails, exactly one further logarithmic level suffices. For homogeneous solutions, the number of dimensions missing at depth n is precisely rank M .

Proof. Define successively the polynomial terms

$$u = \sum_{q \geq 0} (-\tilde{G}R)^q \tilde{G}f. \quad (7.3)$$

We first prove a uniform degree bound, before summing in the polynomial ring. Every application of R lowers the outer exponent by at least ε . The operators $\tilde{G}$ preserve it. Along any contributing sequence of blocks, the outer exponents at successive applications of $\tilde{G}$ therefore strictly decrease. Such a sequence can visit each real root at most once. By lemma 7.1, only a visit to a root can increase the T -degree, and each visit increases it by at most one. Starting from f of T -degree zero, all terms in (7.3) have degree at most s .

This path assertion is coefficientwise legitimate. At a fixed perturbation length, the Hahn product and strong block operators give summable contributions. At a fixed final outer exponent, only finitely many perturbation lengths can occur, by the same outer bound as in lemma 6.1. Thus the sum is well-defined in $\mathcal{H}_{n+1}$, and its uniform degree bound puts it in $\mathcal{H}_n[T]$. It satisfies $u + \tilde{G}Ru = \tilde{G}f$; applying P_0 proves $\mathcal{L}u = f$.

For a kernel seed Kc , set

$$u_c = \sum_{q \geq 0} (-\tilde{G}R)^q Kc. \quad (7.4)$$

Applying P_0 gives $\mathcal{L}u_c = 0$. A seed in the root block λ begins with T -degree zero and undergoes a perturbation before any use of $\tilde{G}$. It can therefore encounter only roots strictly smaller than λ . There are at most $s - 1$ such roots, proving (7.2). The leading outer block of the solution from $x^\lambda L^k$ is exactly that seed, so the r resulting solutions are linearly independent.

It remains to exclude additional homogeneous solutions in the full field $\mathcal{H}_{n+1}$. Let $v \neq 0$ solve $\mathcal{L}v = 0$, and let α be its greatest outer exponent. Since Rv is strictly lower, the leading block v_α satisfies $P(\alpha + \delta)v_\alpha = 0$ in the lower field with one more logarithmic variable. The same kernel proof as in theorem 5.1 shows that α is a real root and v_α is a polynomial in L of degree less than its multiplicity. Subtract the corresponding linear combination of the

solutions (7.4). The greatest possible remaining outer exponent is a strictly lower real root. Repeating removes the solution after at most s root levels. Thus the constructed r solutions span the entire kernel in $\mathcal{H}_{n+1}$.

When there are no real roots, G is an actual inverse of P_0 , so theorem 6.2 gives the stated unique fixed-depth solution. Finally, a failed matrix test means no solution belongs to $\mathcal{H}_n$, whereas (7.3) supplies one in the next field. The dimension statement follows by subtracting (6.12) from r . $\square$

7.1 Distinct roots, not multiplicity

The distinction in (7.1)–(7.2) is significant. For example, when $P(z) = (z - \lambda)^m$, there is only one root level, however large m is. All m homogeneous solutions remain at the original logarithmic depth after an outer-small perturbation. A forcing may still need the new logarithm, but at most its first power. Repeated differentiation produces multiple residue moments, while increased degree in the *new* logarithm is caused by traversing different resonant outer exponents.

This also explains why the two statements “one new logarithm suffices” and “the correction is linear in that logarithm” are not equivalent. The former is true throughout our class; the latter is false as soon as a long chain of distinct resonances is present.

8 A sharp family with an exact Jordan-shift obstruction

For $n \geq 1$ and an integer $s \geq 1$, set

$$P_s(z) = \prod_{j=0}^{s-1} (z - j), \quad \mathcal{L}_{n,s} = P_s(E) + \rho_n. \quad (8.1)$$

Here the last term is multiplication by $\rho_n = x^{-1}\sigma_n$. The uniform support drop is $\varepsilon = 1$, and all roots are simple.

Theorem 8.1 (Maximal loss of fixed-depth homogeneous solutions). *In the bases $Ke_j = x^j$ and $Je_j = x^j\sigma_n$, $0 \leq j < s$, the matrix of theorem 6.2 is exactly*

$$Me_0 = 0, \quad Me_j = e_{j-1} \quad (1 \leq j < s). \quad (8.2)$$

Consequently

$$\text{rank } M = s - 1, \quad \dim \ker_{\mathcal{H}_n} \mathcal{L}_{n,s} = \dim \text{coker}_{\mathcal{H}_n} \mathcal{L}_{n,s} = 1,$$

whereas $\dim \ker_{\mathcal{H}_{n+1}} \mathcal{L}_{n,s} = s$.

Proof. For $j \geq 1$, multiplication by ρ_n gives $RKe_j = Je_{j-1}$. Since $GJ = 0$, every higher perturbation correction starting from that vector vanishes before the next R . The first term gives $\Pi RKe_j = e_{j-1}$. Starting from e_0 , the first perturbation has outer exponent -1 , and all later terms have still smaller outer exponent. No root block can be reached, so the column is zero. This proves (8.2). The dimension conclusions follow from theorems 6.2 and 7.2. $\square$

Theorem 8.2 (Sharp homogeneous logarithmic degree). *There is a homogeneous solution of (8.1) whose leading outer block is x^{s-1} . In its block of outer exponent $s - 1 - k$, $0 \leq k < s$, the coefficient of the highest power T^k is the nonzero real number*

$$c_0 = 1, \quad c_k = -\frac{c_{k-1}}{kP'_s(s-1-k)} = \frac{(-1)^k}{k! \prod_{h=1}^k P'_s(s-1-h)}. \quad (8.3)$$

In particular, the bound $s - 1$ in (7.2) is attained. No homogeneous solution with that leading block can have smaller T -degree.

Proof. Use (7.4) with seed x^{s-1} . Write the successive root blocks as $x^{s-1-k}A_k$. Since R shifts the outer exponent by exactly -1 , the root-block recursion is

$$P_s(s-1-k+\delta)A_k = -\sigma_n A_{k-1}, \quad A_0 = 1. \quad (8.4)$$

At $k \geq 1$, factor the left operator as $\delta Q(\delta)$ with $Q(0) = P'_s(s-1-k)$. Suppose A_{k-1} has top T -coefficient c_{k-1} , a real constant. The top contribution to a primitive of $-c_{k-1}\sigma_n T^{k-1}$ is $-c_{k-1}T^k/k$, since $\delta T = \sigma_n$. Terms of smaller T -degree can produce degree at most $k-1$.

Applying $Q(\delta)^{-1}$ divides the coefficient of T^k by $Q(0)$. Every positive power of δ differentiates the constant coefficient times T^k and lowers its T -degree. This proves the recurrence and shows that the top coefficient remains a real constant. It is nonzero because the roots are simple. Below the last root, all blocks are nonresonant, so no further degree increase occurs.

Two solutions with the same leading block differ by homogeneous solutions seeded at smaller roots. Such a solution can traverse at most $s-2$ lower resonances, so cannot cancel the nonzero T^{s-1} coefficient at the last root. This proves the last assertion. $\square$

Using

$$P'_s(j) = (-1)^{s-1-j} j!(s-1-j)!, \quad (8.5)$$

the last coefficients for the first six cases are

s	Necessary homogeneous degree	c_{s-1} in the x^0 block
1	0	1
2	1	1
3	2	$-1/4$
4	3	$-1/144$
5	4	$1/82944$
6	5	$1/1194393600$

These rational identities were also checked by the included script, independently of the proof by recurrence.

Corollary 8.3 (Sharp inhomogeneous degree). *For the forcing*

$$f = x^{s-1}\sigma_n, \quad (8.6)$$

every solution of $\mathcal{L}_{n,s}u = f$ in $\mathcal{H}_{n+1}$ has T -degree exactly s . The coefficient of T^s in the outer-zero block is

$$C_s = \frac{(-1)^{s-1+s(s-1)/2}}{s! \left(\prod_{j=0}^{s-1} j! \right)^2}. \quad (8.7)$$

There is no solution in $\mathcal{H}_n$.

Proof. At the highest root, the forcing produces the top term $T/P'_s(s-1)$. The same recursion as in (8.4) then increases degree at each of the remaining $s-1$ roots. The top coefficient at the last root is

$$\frac{(-1)^{s-1}}{s! \prod_{j=0}^{s-1} P'_s(j)},$$

which equals (8.7) by (8.5). Every homogeneous solution in the entire next-depth field has degree at most $s - 1$ by theorem 7.2, so the coefficient cannot be canceled by choosing another particular solution.

The original-depth obstruction can also be read directly from the matrix. Here $f = Je_{s-1}$, so $b(f) = e_{s-1}$, which is not in the range of the Jordan shift (8.2). $\square$

8.1 Consequences for proposed closure principles

The family gives a counterexample to the statement that outer-small perturbations of an Euler equation preserve all its homogeneous solutions in a fixed logarithmic field. The loss can be $s - 1$ out of s dimensions, even though the perturbation is only multiplication by one monomial. It also refutes a uniform bound on the degree of the newly needed logarithm that is independent of the number of distinct resonances.

By contrast, it does *not* refute closure after one logarithmic-depth promotion: that is exactly what theorem 7.2 proves. The distinction between depth and degree is the structural point of these examples.

9 A second-order equation: formal resonance and real error bounds

We now pass from a formal theorem to an actual real-variable calculation. This section realizes the decisive first logarithmic block of one example, not every arbitrary Hahn series in the preceding sections.

Take $n = 1$ and $s = 2$. With $L = \log x$ and $T = \log L$, the equation is

$$\left(E(E - 1) + \frac{1}{xL}\right)y = 0, \quad \text{equivalently} \quad y'' + \frac{y}{x^3 \log x} = 0. \quad (9.1)$$

Its fixed-depth matrix is $Me_1 = e_0$, $Me_0 = 0$. Exactly one homogeneous dimension remains in $\mathcal{H}_1$, and the growing solution seeded by x requires T .

9.1 The exact formal outer-zero block

The next block after x solves

$$\delta(\delta - 1)A_0 = -\frac{1}{L}. \quad (9.2)$$

Define the formal factorial series

$$\widehat{S}(L) = \sum_{k=0}^{\infty} (-1)^k k! L^{-k-1}. \quad (9.3)$$

It is a legitimate Hahn series, though divergent as a numerical power series for every fixed positive L . Termwise differentiation gives

$$(\delta - 1)\widehat{S} = -\frac{1}{L}. \quad (9.4)$$

Since $\delta T = 1/L$, one obtains

$$\delta(\delta - 1)(T + \widehat{S}) = -\frac{1}{L}.$$

The zero-constant choice for the outer-zero block is therefore

$$y_{\text{grow}} = x + T + \widehat{S}(L) + \text{blocks with outer exponent at most } -1. \quad (9.5)$$

All lower blocks are polynomial of degree at most one in T by theorem 7.2.

For comparison, the solution seeded by 1 has the first correction

$$y_{\text{bd}} = 1 + x^{-1}v(L) + \text{lower outer blocks}, \quad (9.6)$$

$$\begin{aligned} v(L) &= - \sum_{k \geq 0} (-1)^k k! (1 - 2^{-k-1}) L^{-k-1} \\ &= -\frac{1}{2L} + \frac{3}{4L^2} - \frac{7}{4L^3} + \cdots. \end{aligned} \quad (9.7)$$

Indeed, the block equation is $(\delta - 1)(\delta - 2)v = -1/L$, and

$$\frac{1}{(z-1)(z-2)} = \sum_{k \geq 0} (1 - 2^{-k-1}) z^k.$$

Neither (9.5) nor (9.7) is yet a real convergence assertion.

9.2 An exact sum and a signed factorial remainder

For $L > 0$, define

$$S(L) = \int_0^\infty \frac{e^{-Lt}}{1+t} dt. \quad (9.8)$$

Differentiation under the integral, justified by exponential domination on compact positive L -intervals, gives

$$S'(L) - S(L) = -\frac{1}{L}, \quad 0 < S(L) < \frac{1}{L}. \quad (9.9)$$

Thus S is an exact real counterpart of the formal block $\widehat{S}$.

Proposition 9.1 (Signed remainder). *For every integer $N \geq 0$ and every $L > 0$,*

$$0 < (-1)^N \left(S(L) - \sum_{k=0}^{N-1} (-1)^k k! L^{-k-1} \right) < N! L^{-N-1}. \quad (9.10)$$

The empty sum for $N = 0$ is zero.

Proof. Use the finite geometric identity

$$\frac{1}{1+t} = \sum_{k=0}^{N-1} (-1)^k t^k + \frac{(-1)^N t^N}{1+t}$$

in (9.8). Each polynomial integral is $k! L^{-k-1}$. The remaining positive integral is strictly between zero and $\int_0^\infty e^{-Lt} t^N dt = N! L^{-N-1}$. $\square$

The Borel transform in the convention $\sum a_k L^{-k-1} \mapsto \sum a_k t^k / k!$ is $1/(1+t)$. Formula (9.8) is its Laplace integral on the positive axis. This establishes ordinary Borel summability of this single block in that direction. It does not establish Borel summability of the entire Hahn solution or arbitrary coefficient fields.

9.3 Volterra construction of two real solutions

Let $X \geq e^2$, and on $[X, \infty)$ put

$$q(x) = \frac{1}{x^3 \log x}, \quad (Vg)(x) = \int_x^\infty (t-x)q(t)g(t) dt.$$

For bounded continuous g ,

$$\|Vg\|_{[x, \infty)} \leq \eta_x \|g\|_{[x, \infty)}, \quad \eta_x = \frac{1}{2x \log x} < 1. \quad (9.11)$$

Here the norm is the supremum on the indicated tail. To prove this, use $\log t \geq \log x$ and

$$\int_x^\infty \frac{t-x}{t^3} dt = \frac{1}{2x}.$$

The Neumann series for $(\text{id} + V)^{-1}$ consequently converges in the usual supremum norm. Also $(Vg)'' = qg$ for continuous bounded g , by differentiating the integral twice.

The bounded solution

$$y_0 = (\text{id} + V)^{-1} 1 \quad (9.12)$$

satisfies $y_0'' = -qy_0$ and

$$|y_0(x) - 1| \leq \frac{\eta_x}{1 - \eta_x}. \quad (9.13)$$

Thus $y_0(x) \rightarrow 1$. The tail estimate follows by restricting the same Volterra equation to $[x, \infty)$ and applying (9.11).

For the growing solution, define

$$z(x) = x + \log \log x + S(\log x) = x + T + S(L). \quad (9.14)$$

A direct calculation from (9.9) gives

$$z'' + qz = q(T + S(L)). \quad (9.15)$$

The function $T + S(L)$ is unbounded but its Volterra transform $h = V(T + S(L))$ is bounded. Set

$$w = -(\text{id} + V)^{-1} h, \quad y_1 = z + w. \quad (9.16)$$

Then $w = -V(T + S(L)) - Vw$, so $w'' = -q(T + S(L) + w)$. Combining with (9.15) proves that y_1 solves (9.1) exactly.

Theorem 9.2 (An explicit analytic certificate). *For every $x \geq e^2$, the real solution y_1 constructed in (9.16) satisfies*

$$|y_1(x) - x - \log \log x - S(\log x)| \leq \frac{\frac{\log \log x}{2x \log x} + \frac{5}{4x(\log x)^2}}{1 - \frac{1}{2x \log x}}. \quad (9.17)$$

The solutions y_0, y_1 are linearly independent. For every fixed $N \geq 0$,

$$y_1(x) = x + \log \log x + \sum_{k=0}^{N-1} (-1)^k k! (\log x)^{-k-1} + \mathcal{E}_N(x), \quad (9.18)$$

$$|\mathcal{E}_N(x)| \leq N! (\log x)^{-N-1} + \frac{\frac{\log \log x}{2x \log x} + \frac{5}{4x(\log x)^2}}{1 - \frac{1}{2x \log x}}. \quad (9.19)$$

Proof. The integrand defining h is positive for $x \geq e^2$, and h is decreasing. Substitute $t = xv$, with $L = \log x$ and $T = \log L$. For $v \geq 1$,

$$\log \log(xv) = \log(L + \log v) \leq T + \frac{\log v}{L}, \quad S(\log(xv)) \leq \frac{1}{L}.$$

Consequently

$$\begin{aligned} h(x) &\leq \frac{1}{xL} \int_1^\infty \frac{v-1}{v^3} \left(T + \frac{\log v + 1}{L} \right) dv \\ &= \frac{T}{2xL} + \frac{5}{4xL^2}, \end{aligned}$$

where

$$\int_1^\infty \frac{v-1}{v^3} dv = \frac{1}{2}, \quad \int_1^\infty \frac{v-1}{v^3} \log v dv = \frac{3}{4}.$$

Because h is decreasing, its norm on $[x, \infty)$ is $h(x)$. The inverse bound from (9.11) therefore gives $|w(x)| \leq h(x)/(1 - \eta_x)$, proving (9.17).

We have $y_0(x) \rightarrow 1$, while $y_1(x)/x \rightarrow 1$, so the solutions cannot be scalar multiples. The integral equations make them C^2 solutions, and the ordinary uniqueness theorem for a second-order linear equation shows that they constitute a basis on the interval. Finally combine (9.17) with proposition 9.1 to obtain (9.19). $\square$

The normalization implicit here is

$$y_0(x) \rightarrow 1, \quad y_1(x) - x - \log \log x \rightarrow 0.$$

Adding a multiple of y_0 to y_1 would change the second limit. This selects a definite real growing solution, rather than merely its leading asymptotic equivalence.

9.4 Numerical illustrations and their interpretation

The accompanying script evaluates S , z , and the right-hand side of (9.17) at high working precision. It records the following approximate values:

x	$S(\log x)$	$z(x)$	Bound in (9.17)
10	0.323897895933	11.1579303412	$4.26125667 \cdot 10^{-2}$
100	0.182974349963	101.710153976	$2.24996864 \cdot 10^{-3}$
10^4	0.0987515406201	10002.3190783	$1.35270483 \cdot 10^{-5}$

The table entries are rounded illustrations, not outward-rounded interval certificates. The exact formula (9.17), proved above, is the certificate. No numerical integration of the differential equation is needed to justify it.

For fixed N , the second term on the right of (9.19) is smaller than every fixed inverse power of $\log x$. This realizes the whole displayed factorial expansion within the decisive outer-zero block. We do not claim here that the construction has been matched term by term to every smaller outer block of the full formal solution.

10 Proof boundaries, checks, and formalization

10.1 What the proofs establish

The formal statements are algebraic statements about explicitly defined Hahn fields and strong operators. Their infinite sums are justified by two distinct mechanisms: inside a block, δ lowers the L -exponent by one; between blocks, R lowers the x -exponent by a uniform ε . Residue extraction accounts for the finite exceptional part. Neither mechanism uses a hidden analytic convergence assumption.

The analytic theorem concerns a concrete real equation and is established separately by a supremum-norm contraction. It does not follow merely by substituting a real number into an arbitrary formal Hahn series. The factorial block is divergent as an ordinary power series, but its explicit Laplace integral has a signed remainder bound.

No claim of Lean compilation, mechanized checking of the new theorems, or proof-assistant coverage is made. The repository modules motivated and supplied the mathematical starting point; the proofs in this article stand as conventional proofs requiring normal mathematical review.

10.2 Finite exact verification supplied with the article

The package contains `verification/verify.py` and its recorded output `verification/results.json`. The run completed with all checks passing. Its scope is intentionally narrower than theorem verification.

The exact symbolic part uses rational arithmetic to check 212 block identities. A finite block is represented as a dictionary of coefficients of $L^b T^c$, with integer b and nonnegative integer c . Differentiation uses

$$\delta(L^b T^c) = bL^{b-1}T^c + cL^{b-1}T^{c-1}.$$

The program tests resonant and nonresonant inverses, repeated-root cases, and the sharp-family recursion for $1 \leq s \leq 6$. It retains guard terms down to L -exponent -28 and checks residuals only at exponents at least -10 . The gap prevents a discarded lower term from being mistaken for a checked identity at the visible cutoff.

A second exact test constructs 24 finite-dimensional split-operator examples, with deterministic random seed 20260929. It checks the geometric inverse, finite matrix, reconstruction formula, and equality of kernel/cokernel dimensions using exact SymPy matrices. These examples validate the algebraic arrangement and signs; they do not replace the infinite-support proof.

The numerical part uses `mpmath` for the integral block and remainder illustrations. It is not an interval-arithmetic verification. Its values are reproducible at the script's chosen working precision. The symbolic dependency versions used in the delivered run are recorded in the package README.

10.3 A staged Lean implementation plan

The following separation would keep a future formalization auditable.

Stage 1: the concrete ambient and strong derivative. Construct the ordered exponent group $\mathbb{R}^{n+1}$ with lexicographic order and its order-dual support convention. Establish block

decomposition, coefficient extraction, and the finite-shift derivative. This stage needs a concrete Hahn ambient, not only polynomial identities. The repository's well-based support lemmas and its existing polynomial block API are relevant starting points [2, 3].

Stage 2: normalized integration and residue moments. Formalize the lower-exponent contracting functional calculus, define I_n , and prove (3.2). Then derive residue integration by parts and the Green formula. The moment theorem should be proved abstractly for any differential algebra with $Dx = 1$, constants $\mathbb{R}$, and the stated residue splitting; the Hahn construction supplies an instance.

Stage 3: a generic finite-defect perturbation theorem. Separate theorem 6.2 from the concrete field. Its proof only uses the identities (5.8), a geometric inverse, and finite-dimensional linear algebra. This reusable algebraic theorem can be formalized before all transseries details are available. Its hypotheses must explicitly include the sum/existence conditions rather than assume them implicitly.

Stage 4: support bounds and logarithmic promotion. Instantiate the abstract theorem using the primary-exponent decrease. Prove the exact cutoff, the degree-controlled right inverse on $\mathcal{H}_n[T]$, and the strictly descending root-path argument. This is the part that turns a general perturbative method into a sharp fixed-depth theorem.

Stage 5: examples and analytic certification. The Jordan-shift family is a compact integration test of the formal API. The Volterra theorem belongs in a separate real-analysis development with explicit differentiation-under-the-integral and norm bounds. Formal symbolic truncation and analytic error bounds should remain different theorem types.

This plan does not presume that a full class of real germs realizing arbitrary Hahn series can be formalized as a consequence. Such an evaluation theory would require additional hypotheses and constructions.

11 Further research questions and concrete next targets

The following are proposed questions arising from the proved theorems. They are not presented as a catalog of already established open problems in the literature. Each indicates what the present argument does not settle.

11.1 The exact minimal power of the promoted logarithm

The finite matrix decides whether degree zero in T is possible; the root-path argument gives the universal upper bound s . It does not in general compute the least possible positive degree. Can one construct a finite sequence of residue matrices whose successive ranks recover the full filtration by T -degree of homogeneous and particular solutions?

A concrete first target is the subclass with finitely many rational outer drops and finite Laurent-log coefficient blocks. One would seek an algorithm that outputs the smallest d such that a forcing has a solution in $\mathcal{H}_n[T]_{\leq d}$, together with an obstruction certificate for degree $d - 1$. The sharp family should yield the complete sequence $0, 1, \dots, s$.

Editorial note (ProveIt, 2026-09-29). The later package `Analysis/Transseries/docs/series-and-transseries/Hahn_Fuchsian_Resonance_Analytic_Normalization/` answers the

homogeneous half of this question in a neighbouring setting, without citing it (it knew this article only through its README). For first-order systems $xY' = (A_0 + A_+(x))Y$ over the log-free Hahn field with $L = \log x$ adjoined, with constant coefficient matrices, real spectrum and well-ordered positive exponents, the solutions of degree at most d in L form a space of dimension $\dim \ker B^{d+1}$ for one finite nilpotent matrix $B = N + \sum_{\rho} R_{\rho}$ (`hahn_fuchsian.tex`, Theorem `thm:logs`), whose entries depend polynomially on finitely many input coefficients (Theorem `thm:ancestry`). That is depth 0, where the outer-block construction of this article, which needs $n \geq 1$, does not apply; particular solutions of forced equations are not treated there. At depth $n \geq 1$, with the new logarithm T and with forcing, the question is answered for this article's whole class by the later package `Analysis/Transseries/docs/series-and-transseries/Exact_Logarithmic_Degree_Smith_Invariants/` (batch 50; not reviewed): the Smith exponents of a lifted residue series satisfy $\sum \nu_i = r$, $0 \leq \nu_i \leq s$, give the whole filtration, and are decided by block Toeplitz ranks of a finite jet, with rational obstruction certificates; the Jordan-shift family of theorem 8.1 is its single-block case.

11.2 Cancellation-sensitive resonance graphs

The semigroup condition (6.15) detects impossible matrix entries but does not prove that a permitted path contributes nontrivially. Different derivative orders and intermediate blocks may cancel. Can a weighted directed graph of residue transfers characterize the leading nonvanishing path and sharpen both the cutoff and the logarithmic-degree bound?

For monomial coefficients one can first classify which edge weights depend only on $P'(\lambda)$ and which require higher residue moments. A useful theorem would distinguish structural zeros from coefficient-dependent cancellation without expanding all lower logarithmic terms.

Editorial note (ProveIt, 2026-09-29). For operators $P(E) + \sigma \sum_{\eta} x^{-\eta} q_{\eta}(E)$ with simple real roots, the later package `Analysis/Transseries/docs/series-and-transseries/Exact_Logarithmic_Degree_Smith_Invariants/` (batch 50; not reviewed) shows that the relevant weighted graph is the matrix $D_P^{-1}N$, $N_{ij} = q_{\lambda_j - \lambda_i}(\lambda_j)$, whose Jordan block sizes are the degrees; a fourth-order family has the same unweighted graph and different degrees. The general case and the sharpening of the cutoff remain open.

11.3 Perturbations small only in logarithms

The hypothesis $d_0(a_j) \leq -\varepsilon$ excludes coefficients such as $1/\log x$, whose outer exponent is zero. In that setting a perturbation can stay in one outer root block indefinitely. Can a multi-level version of the reduction, contracting successively in x , $\log x$, and deeper logarithms, still yield finite obstruction data?

The present proof cannot simply replace “outer-small” by “small” in the full lexicographic order. Its finite cutoff and its rule that a root is visited at most once would no longer apply. A precise next step is a classification of single-root operators with coefficients supported in negative powers of L , separating those needing only T from those requiring exponentials or further logarithmic depth.

11.4 Parameter-dependent root collisions

When two distinct real roots coalesce, s decreases but a multiplicity increases. The degree theorem predicts less growth in the new logarithm, while the natural coefficient formulas may become singular in the root difference. Can one construct a parameter-uniform block basis

and obstruction matrix that pass continuously, or in a controlled confluence sense, through such a collision?

A test case is $P_\tau(z) = z(z - \tau)$ with a monomial outer-small perturbation. The fixed-parameter theorem is already proved here; the unresolved target is a uniform representation and an interpretation of the divergent coefficients as $\tau \rightarrow 0$.

11.5 Matrix systems and nonreal indicial roots

For a matrix polynomial $P(E)$, kernels at a root may contain Jordan chains before any perturbation. A matrix-valued residue complex should replace the scalar moments. Which invariant controls the new-logarithm degree: the number of distinct root levels, the lengths of internal Jordan chains, or a combination?

Nonreal scalar roots raise a different issue. The present real monomial field deliberately omits $x^{i\beta}$ and oscillation. An extension with complex powers would need a careful support convention rather than an artificial total ordering of oscillatory functions. A first goal is an algebraic version for a fixed finite collection of imaginary exponents, without claiming an ordered Hardy-field interpretation.

Editorial note (ProveIt, 2026-09-29). For first-order systems at depth 0 with real spectrum (the setting of the previous note), the later package `Analysis/Transseries/docs/series-and-transseries/Hahn_Fuchsian_Resonance_Analytic_Normalization/` identifies the invariant: the new-logarithm degree of a fundamental matrix is exactly $\nu(B) - 1 \leq m - 1$, where $\nu(B)$ is the nilpotency index of $B = N + \sum_\rho R_\rho$, which combines the internal Jordan chains (through N) with the passage between root levels (through the resonant blocks R_ρ) (`hahn_fuchsian.tex`, Theorem `thm:logs`; the bound is attained in Example `ex:chain`, the analogue of the Jordan-shift family of theorem 8.1). When $N = 0$, B is strictly block-triangular over the s distinct eigenvalue levels, so the degree is at most $s - 1$, the homogeneous bound of theorem 7.2; that package does not state this consequence. Matrix polynomials $P(E)$ of higher order, depth $n \geq 1$, and nonreal roots are not treated there.

11.6 Nonlinear equations and an effective obstruction variety

Consider $P(E)u + Ru + F(u) = f$, with a nonlinear term that is suitably small in a support filtration. Eliminating the infinite-dimensional complement may leave a finite nonlinear equation in the kernel coordinates. Can the linear matrix M be replaced by a convergent formal obstruction map, with explicit support bounds for its Taylor coefficients?

The existing generalized-series implicit-function theory is a natural framework [5]. The specific next theorem would retain resonance coordinates rather than assume an everywhere invertible linearization, and would state which nonlinearities preserve a finite logarithmic-depth bound.

11.7 Effective coefficient representations

For arbitrary real Hahn coefficients, coefficient extraction and equality need not be computable. What practical representation makes the finite-depth cutoff of theorem 6.3 an actual terminating symbolic algorithm?

A useful restricted target combines rational exponent lattices, computable well-based supports, and coefficient blocks generated by a finite differential-algebraic grammar. One should specify the exact oracle assumptions and prove complexity in the number of reachable root differences, rather than describe the finite matrix as automatically computable.

11.8 Analytic realization beyond the displayed block

The real construction in section 9 certifies the decisive outer-zero block, not every subsequent block. Can one show that the same Volterra solutions have the entire formal Hahn expansion, with explicit truncation bounds simultaneously in powers of x^{-1} and $(\log x)^{-1}$?

For a broader class, impose analytic or Gevrey bounds on the coefficient blocks and ask whether the finite residue matrix identifies precisely which logarithmic terms occur in sectorial actual solutions. A particularly useful result would translate (6.16) into an analytic residual bound without silently treating formal support order as numerical size.

11.9 Other ordered exponent groups

Our outer exponents are real, and the uniform ε gives a simple Archimedean cutoff. A more general ordered group may have positive infinitesimals or no useful integer-valued depth bound. Which version of “strict decrease” guarantees summability of the geometric operator inverse and finiteness of its projection to the resonant part?

One should separate two demands: existence of the full formal inverse and termination of the finite-dimensional obstruction calculation. They coincide under the present real uniform-drop hypothesis, but need not coincide in a non-Archimedean exponent group.

Editorial note (ProveIt, 2026-09-29). The later package `Analysis/Transseries/docs/series-and-transseries/Reversion_Beyond_Archimedean_Valuations/` draws the analogous separation for compositional reversion over an arbitrary ordered exponent group: strong Hahn convergence always holds, uniform valuation convergence holds exactly when the multiples of the least support exponent are cofinal, and every finite coefficient request factors through a finite-rank quotient. It concerns reversion of near-identity maps, not the differential operators of this article, so this question remains open.

11.10 A reusable proof-assistant library for finite defects

The identities $P_0G = \text{id} - J\Pi$ and $GP_0 = \text{id} - K\Lambda$ package an infinite-dimensional problem into a finite kernel/cokernel interface. Can a library expose this interface independently of the Hahn implementation and reuse it for difference equations, summation, or sequence inversion?

The immediate target is a formally checked version of theorem 6.2 with explicit summability hypotheses and then one concrete Hahn instance. This would be a meaningful extension of the repository’s existing block-antiderivative results, rather than a new count of unrelated declarations.

12 Conclusion

At finite logarithmic depth, the obstruction to solving a differential equation need not be an uncontrolled infinite tail. For outer-small perturbations of $P(xD)$, it is a finite residue matrix, and only finitely many perturbation depths can affect that matrix. Its rank measures the exact discrepancy between solvability at the original depth and solvability after promotion.

One more logarithm always suffices in this class, but the degree in that logarithm records the chain of distinct resonant outer exponents. The Jordan-shift family proves that arbitrarily high powers can be forced while only one new logarithmic level is needed. The second-order member demonstrates that the formal obstruction can also be accompanied by actual real solutions and a quantitative bound.

The outcome is a specific extension of the repository's polynomial-block calculus: a concrete Hahn ambient, explicit residue moments, finite-dimensional obstruction data, sharp depth/degree control, and a separate analytic certificate. The broader nonlinear, parameter-uniform, and general analytic-realization questions remain distinct research tasks.

A A compact abstract form of the reduction

For reuse outside transseries, let X be a real vector space, V finite-dimensional, and suppose linear maps

$$P_0, G : X \rightarrow X, \quad K, J : V \rightarrow X, \quad \Lambda, \Pi : X \rightarrow V$$

satisfy

$$\begin{aligned} P_0 K &= 0, & \Pi P_0 &= 0, & \Lambda K &= \Pi J = \text{id}, \\ P_0 G &= \text{id} - J\Pi, & G P_0 &= \text{id} - K\Lambda, & \Lambda G &= GJ = 0. \end{aligned}$$

For any $R : X \rightarrow X$ such that $\text{id} + GR$ is invertible, define $\mathcal{N}$, M , and b as in (6.4) and (6.6), interpreting $\mathcal{N}$ simply as the inverse if no infinite sum is involved. The proof of theorem 6.2 gives

$$\ker(P_0 + R) \cong \ker M, \quad \text{coker}(P_0 + R) \cong \text{coker } M.$$

No norm, topology, or differential structure is needed for this algebraic statement. The Hahn-specific work is to construct the splitting maps, establish the geometric inverse, and show that its finite-dimensional projection terminates. This decomposition is also why the finite-dimensional exact tests in the supplied script are relevant but cannot certify the Hahn support arguments.

B Support facts used in the constructions

We spell out the elementary local-finiteness principle used repeatedly. Suppose $(S_q)_{q \geq 0}$ are reverse well-ordered subsets of $\mathbb{R}^{n+1}$ in lexicographic order and every first coordinate in S_q is at most $A - q\varepsilon$, where $\varepsilon > 0$. Then $S = \bigcup_q S_q$ is reverse well-ordered. To see this, choose one element a of a nonempty subset of S . Only finitely many S_q can contain an element with first coordinate at least a_0 . Each of their intersections with the subset has a greatest element, and the greatest of those finitely many choices is greatest in the whole subset. Every fixed exponent lies in only finitely many S_q as well.

If each S_q is itself the support union of a summable family, the finite-contribution condition at a fixed exponent then proves summability of the doubly indexed family. The same argument applies inside an outer block with L in place of x and $\varepsilon = 1$. It proves the rearrangement assertions for all geometric and formal functional-calculus sums used in the paper.

For ordinary Hahn multiplication, finite sums of reverse well-ordered supports are reverse well-ordered, and each resulting exponent has finitely many decompositions. One way to check the obstruction to an infinite bad sequence is to use the following elementary fact: a sequence in a reverse well-ordered total order has an infinite nonincreasing subsequence. If infinitely many indices are at least as large as all their successors, choose those; otherwise construct a forbidden infinite increasing subsequence. Given an increasing sequence of sums, first pass to a nonincreasing subsequence in one summand support and then in the other, obtaining a contradiction. Given infinitely many distinct decompositions of one fixed exponent, a strictly decreasing subsequence in one coordinate would force a strictly increasing one in the other. This proves the required local finiteness. Finite unions are immediate.

These arguments explain why none of the infinite operator expressions in the article is being justified merely by suggestive notation or by an unproved interchange of infinite sums.

C Package contents and reproducibility

The delivered archive contains the editable source `residue_fredholm.tex`, the compiled article `residue_fredholm.pdf`, a build/readme file, the verification script, and its recorded JSON output. The document uses standard L^AT_EX packages and can be built with

```
pdflatex -interaction=nonstopmode -halt-on-error residue_fredholm.tex
pdflatex -interaction=nonstopmode -halt-on-error residue_fredholm.tex
pdflatex -interaction=nonstopmode -halt-on-error residue_fredholm.tex
python verification/verify.py
```

The verification script uses rational arithmetic for symbolic checks and requires `sympy` and `mpmath`. The exact output counts and numerical illustration values described in section 10 refer to the run included with this package. They do not assert a proof-assistant verification of the mathematical manuscript.

References

- [1] Vladimir Reshetnikov, *ProveIt: Transseries project README*, repository snapshot 8813fb3a2cdd1702d3ea7f7ca5d63ae710630bef, inspected 29 September 2026. <https://github.com/VladimirReshetnikov/ProveIt/blob/8813fb3a2cdd1702d3ea7f7ca5d63ae710630bef/Analysis/Transseries/README.md>.
- [2] Vladimir Reshetnikov, *ProveIt: Series and transseries documentation and canonical-volume status*, same snapshot and access date. <https://github.com/VladimirReshetnikov/ProveIt/blob/8813fb3a2cdd1702d3ea7f7ca5d63ae710630bef/Analysis/Transseries/docs/series-and-transseries/README.md>. See also the adjacent *Transseries_And_Inversion/README.md* for the declaration-level crosswalk and explicit scope boundaries.
- [3] Vladimir Reshetnikov, *ProveIt: TransseriesBlockAntiderivative.lean*, same snapshot and access date. Inspected file blob: 1a4b5e335b5429a267bcfa4c0c9b35aadf9be527. <https://github.com/VladimirReshetnikov/ProveIt/blob/8813fb3a2cdd1702d3ea7f7ca5d63ae710630bef/Analysis/Transseries/Lean/Transseries/TransseriesBlockAntiderivative.lean>.
- [4] Salma Kuhlmann and Mickaël Matusinski, *Hardy type derivations on generalised series fields*, *Journal of Algebra* **351** (2012), 185–203. DOI: 10.1016/j.jalgebra.2011.11.024. Preprint: arXiv:0903.2197.
- [5] Joris van der Hoeven, *Operators on generalized power series*, *Illinois Journal of Mathematics* **45** (2001), 1161–1190. DOI: 10.1215/ijm/1258138061. Author’s text: <https://www.texmacs.org/joris/noeth/noeth.html>.
- [6] Joris van der Hoeven, *Transseries and Real Differential Algebra*, *Lecture Notes in Mathematics* **1888**, Springer, 2006. DOI: 10.1007/3-540-35590-1.
- [7] Matthew Hogancamp, *Homological perturbation theory with curvature*, preprint, 2019. arXiv:1912.03843. Cited for the broader perturbative framework, not as a source for the particular Hahn residue and degree formulas proved here.