

Shifted Catalan Hankel Polynomials

Exact coefficients, sharp recurrences,
and a denominator conjecture

A research note prepared with OpenAI ChatGPT

19 September 2026

With Part II (28 September 2026): root collisions and sharp recurrences for arbitrary polynomial multipliers, and Parts III–VI (29 September 2026): cyclotomic resonances, two-endpoint confluence, interior root collisions, and growing shifts

Abstract

For the Catalan numbers C_n , consider the size- N determinant $D_N^{(m)}(a, b) = \det(aC_{i+j+m} + bC_{i+j+m+1})_{0 \leq i, j < N}$. We give a self-contained proof of the denominator-exponent conjecture stated in Paul Barry's 2020 note on this family. With $d = \binom{m}{2}$, the generating function is $P_m(t)/(1 - (a + 2b)t + b^2t^2)^{d+1}$. The denominator is minimal whenever $b \neq 0$, $a \neq 0$, and $a \neq -4b$. We determine every exceptional case, prove that $\deg P_m = (m-1)^2$ for $b \neq 0$, and establish a reciprocal symmetry of the numerator. An explicit positive-integer coefficient formula also identifies and repairs a one-unit shift in the source's separate Conjecture 2. The essential degree reduction is a Vandermonde cancellation in confluent Christoffel minors. It yields an exact sum of two exponentials with polynomial coefficients, a finite full asymptotic expansion, and an explicit first correction. Independent integer and rational computations, including direct determinants and polynomial-identity certificates at fixed sizes, accompany the proofs.

Addition of 28 September 2026 (batch 39). Part II, from a later manuscript, replaces $x^m(a + bx)$ by an arbitrary nonzero multiplier $q(x) = \gamma x^m(x-4)^\ell \prod_i (x - c_i)^{t_i}$. Each exponential sector has a polynomial coefficient of exact degree $\binom{m}{2} + \binom{\ell+1}{2} + \sum_i k_i(t_i - k_i)$ with a nonzero product leading coefficient, so off resonance the minimal recurrence order is $\prod_i (t_i + 1) \left(\binom{m}{2} + \binom{\ell+1}{2} + 1 + \frac{1}{6} \sum_i t_i(t_i - 1) \right)$; a secondary Hankel determinant detects exactly when it drops, and a certified finite-prefix algorithm handles the resonances.

Addition of 29 September 2026 (batch 42). Part III gives every characteristic multiplicity for one repeated root $(x - c)^t$, $c = 2 + z + z^{-1} \notin \{0, 4\}$; it depends on the sign of z^L , not only on t and $L = \text{ord}(z^2)$.

Addition of 29 September 2026 (batch 53). Part IV lets roots approach the endpoints 0 and 4 on the scale N^{-2} : the normalized determinant is an even convergent series in $1/(N + \deg q/2)$, and outward displacements are negligible exactly when they are $o(N^{-2})$.

Addition of 29 September 2026 (batch 55). Part V lets roots collide away from 0 and 4 on the scale N^{-1} . *Addition of 29 September 2026 (batch 57).* Part VI lets the shift m grow with N .

Main result. For all nonexceptional complex parameters,

$$\text{minimal constant-coefficient recurrence order} = m(m-1) + 2.$$

The proof is uniform in the determinant size and the Catalan shift.

Status and scope. This is an unrefereed, AI-assisted mathematical manuscript, not a formalized proof. Its theorems are supported by explicit arguments, not inferred from finite tests. The target is a conjecture in a specified source version; no claim of first discovery or exhaustive verification of present-day open status is made. Classical Christoffel formulas and the known general rationality theory are credited, rather than presented as new.

Reproducibility. The accompanying archive contains this L^AT_EX source, the PDF, a standard-library-only Python implementation, an independently organized verification suite, and exact machine-readable data. No external mathematical software is needed to rerun the tests.

Contents

Part I. Shifted linear multipliers	10
1 The problem and the contribution	10
1.1 A notation guide	10
2 Statements of the main theorems	11
2.1 The scale of the improvement	12
3 Moments, orthogonality, and the coefficient formula	12
3.1 Catalan moments and a normalized Jacobi polynomial	12
3.2 A characteristic-polynomial determinant identity	13
3.3 A proof of the shifted Catalan product	13
3.4 Completion of the coefficient proof	14
3.5 Real roots and coefficient inequalities	14
4 A fixed-size confluent determinant	15
5 The degree cancellation and its first correction	16
6 Two exponentials and the minimal recurrence	17
6.1 Why the identity extends to all parameters	18
6.2 Proof of the exceptional classification	18
7 The numerator degree and reciprocal symmetry	19
7.1 An explicit finite prescription for the numerator	20
8 Exact exponential expansions and asymptotics	20
9 Worked examples and integer sequences	21
9.1 Small shifts	21
9.2 The specialization $a = b = 1$	22
9.3 Rational characteristic roots: $a = 1, b = 2$	22
9.4 An endpoint example	23
10 Algorithms, exact experiments, and reproducibility	23
10.1 Implementation choices	23
10.2 Verification that was actually run	23
10.3 How to reproduce the package	24
11 What has been resolved, and what has not	24
Part II. Root collisions: arbitrary polynomial multipliers	26
12 Provenance and conventions of Part II	26
12.1 One manuscript, one addition	26
12.2 Reading conventions	27
12.3 Notation of Part II	27
12.4 Where the merge had to choose	29
13 The research question and its provenance	29
13.1 A concrete extension of the ProveIt report	29
13.2 What is classical, and what is being added	30
14 Definitions and principal results	30
14.1 The weighted determinant and its spectral parameters	30
14.2 The exact multiplicity law	31

14.3	An exact detector for exceptional parameters	32
15	Catalan orthogonality and the fixed-size determinant	33
15.1	The orthogonal polynomials	33
15.2	Christoffel's determinant, including confluence	33
16	Local jet structure	34
16.1	The two branches away from the endpoints	34
16.2	The two endpoint jets	35
17	The determinant degree calculus	36
17.1	Polynomial shifts and confluent exponential rows	36
17.2	Three local leading determinants	36
18	Canonical sectors and the proof of sharpness	37
18.1	A definition that still makes sense at resonance	37
18.2	The leading coefficient in product form	37
19	Minimal recurrences, order, and the secondary factorization	39
19.1	Exponential polynomials and minimality	39
19.2	Comparison with the classical exponential bound	39
19.3	Proof of the secondary Hankel factorization	40
20	Resonance and a root-free exact algorithm	41
20.1	The exact cancellation rule	41
20.2	Why a finite prefix can be a proof	41
20.3	No factorization into individual roots is required	42
20.4	Algebraic rank-drop loci	42
21	Worked examples and checks against the repository	43
21.1	Recovering the shifted linear multiplier	43
21.2	A repeated fourth power with order fifteen	43
21.3	Resonance can cancel an entire branch	44
21.4	A nine-value proof of a resonant fourth-power formula	44
21.5	An irreducible rational input	45
22	Exact asymptotics and stability of the leading sector	45
23	Verification, implementation, and formalization status	46
23.1	Independent exact calculations	46
23.2	The implementation interface	47
23.3	A feasible formalization route	48
24	Further research questions and proposed next steps	48
24.1	Closed classification of cyclotomic resonances	48
24.2	Pairwise resonance and cancellation order	48
24.3	Sector polynomials and symmetric functions	49
24.4	Uniform endpoint and root-collision limits	49
24.5	Other eventually constant Jacobi recurrences	49
24.6	Rational multipliers and finite poles	50
24.7	Coefficient-space equations and arithmetic reductions	50
24.8	Fast certified algorithms and formal verification	50
25	Conclusion	50
26	Normalization and sign audit	50
27	Reproducing the package	51

Part III. Cyclotomic resonances of one repeated root	53
28 Provenance and conventions of Part III	53
28.1 One manuscript, one addition	53
28.2 Reading conventions	54
28.3 Notation of Part III	55
28.4 What Part III answers, and how it corrects the question	56
28.5 Where the merge had to choose	57
29 The repository question and the scope of the answer	58
29.1 What is already known	58
29.2 Why the problem is not solved by identifying equal rates	59
30 Definitions and the complete classification	59
30.1 Immediate generating-function and scaling consequences	60
31 Catalan orthogonality and the confluent determinant	61
31.1 A convenient monic orthonormal family	61
31.2 The determinant formula and its sign	61
32 An exact subset expansion for every sector	62
33 The first two sector coefficients and reflection	63
33.1 A signed finite Vandermonde identity	63
33.2 Leading coefficient and the decisive correction	64
33.3 An exact reflection of the whole sector polynomial	65
34 Why the noncentral cancellation has exactly one defect	65
34.1 Minimal annihilators of exponential polynomials	65
34.2 The geometry of a residue class	66
34.3 The tied group and the nonvanishing next term	66
35 Closed orders, exceptional parameters, and reflection	67
35.1 Summing the grouped degrees	67
35.2 An exact finite exceptional locus	67
35.3 The missing sign datum and reflection of the root	68
35.4 Quadratic quasipolynomial growth in the multiplicity	68
36 The central exception and the known product	69
36.1 Reduction to middle-binomial moments	69
36.2 A self-contained condensation proof	69
36.3 Minimality and why the one-defect law excludes this point	70
37 A root-free minimal-recurrence algorithm	70
37.1 Detecting a resonance by a short scalar recurrence	71
37.2 Linear and quadratic factors in the original coefficient field	71
38 Worked examples and a complete rational-parameter corollary	72
38.1 Two cubic multipliers with the same cyclotomic order	72
38.2 A noncentral loss below a positive degree	73
38.3 All rational nonendpoint roots	73
39 Exact verification and reproducibility	74
39.1 Arithmetic models and independent routes	74
39.2 Parameter ranges and the recorded run	74
39.3 Rebuilding the article and the audit	75
40 Further research questions and proposed next steps	76
40.1 Several distinct repeated roots	76
40.2 Endpoint factors coupled to a cyclotomic interior root	76

40.3	Higher centered coefficients and a symmetric-function formula	77
40.4	Uniform limits near a disappearing characteristic factor	77
40.5	Degrees and zeros on individual size progressions	77
40.6	Other eventually constant Jacobi recurrences	77
40.7	Effective arithmetic and reduction modulo primes	77
40.8	A focused formalization in Lean	77
41	Conclusion	78
42	Normalization, boundary, and proof audit	78
43	Source and package concordance	79
Part IV.	Two-endpoint confluence	80
44	Provenance and conventions of Part IV	80
44.1	One manuscript, one addition	80
44.2	Reading conventions	81
44.3	Notation of Part IV	82
44.4	What Part IV answers, and what stays open	83
44.5	Where the merge had to choose	84
45	The repository question and the contribution	85
45.1	The endpoint moving-root question	85
45.2	What is classical, and what is established here	86
45.3	A guide to the proof	86
46	Notation and the two-endpoint model	86
47	Main results	87
47.1	An exact even expansion	87
47.2	The complete first correction	87
47.3	The endpoint value	88
47.4	A sharp scale criterion	88
48	The Catalan moment functional and Christoffel reduction	89
48.1	Orthogonal polynomials with unit norms	89
48.2	A proof of the fixed-size determinant identity	89
49	The centered hyperbolic alternant	90
49.1	The quadratic generator	90
49.2	The all-orders meaning	90
50	Proof of the even, collision-uniform expansion	91
50.1	An exact normalized formula	91
50.2	Removal of the apparent singularities	91
50.3	An explicit remainder bound	92
51	Derivation of the first interaction coefficient	92
51.1	The five relevant minors	92
51.2	Reduction to ordinary kernel derivatives	93
51.3	The geometric denominator correction	93
52	Positive Schur expansions of the outward profiles	94
52.1	A coefficientwise-positive formula	94
52.2	Consequences and a useful relative error bound	95
53	Proof of the sharp outward critical-scale theorem	95
53.1	Monotonicity at every finite size	95

53.2	Finite scaled coordinates	95
53.3	Necessity and escape from the critical window	95
54	The original scaling and other coordinate choices	96
54.1	An explicit answer to the repository's question	96
54.2	Exact spatial displacements also have an even expansion	96
55	Worked examples and checks on the constants	97
55.1	One root at the left endpoint	97
55.2	One root at the right endpoint	97
55.3	One root at each endpoint	97
55.4	Two coincident roots at an endpoint	97
55.5	A genuinely confluent two-cluster numerical example	98
56	Computation, reproducibility, and the verification boundary	98
56.1	Two independent determinant paths	98
56.2	Executed checks	98
56.3	Computing higher coefficients	99
56.4	A formalization route	99
57	Further research questions and proposed next steps	99
57.1	Nonendpoint collisions on the reciprocal-size scale	99
57.2	Endpoint clusters in the presence of fixed interior roots	100
57.3	Higher interaction operators	100
57.4	Multiplicities growing with determinant size	100
57.5	Other Jacobi endpoint exponents	100
57.6	Zeros of inward confluent profiles	100
57.7	Degeneration of finite recurrences	101
57.8	Sharper analytic bounds and stable numerical evaluation	101
57.9	Rational modifications and Cauchy-transform blocks	101
57.10A	machine-checked coefficient and confluence library	101
58	Conclusion	101
59	A sign and normalization audit	101
60	Archive contents and reproduction	102
Part V.	Interior root collisions	104
61	Provenance and conventions of Part V	104
61.1	One manuscript, one addition	104
61.2	Reading conventions	106
61.3	Notation of Part V	106
61.4	What Part V answers, and what stays open	108
61.5	Where the merge had to choose	109
62	The repository question and the contribution	110
62.1	A specific unresolved continuation	110
62.2	What is established here, and what is classical	111
62.3	Why the answer has to retain phases	111
63	Catalan normalization and the classical determinant reduction	111
63.1	The moment measure and its polynomials	111
63.2	Vandermonde convention and collisions	112
64	The one-cluster theorem	113
64.1	Coordinates and universal profiles	113

65 Proof by centered exponential alternants	114
65.1 The centered Vandermonde identity	114
65.2 Divisibility in the inverse size and in the coordinates	115
65.3 Recovery of the determinant and the leading term	115
66 The first correction and all-order construction	116
66.1 An exact finite coefficient algorithm	116
67 An explicit collision-uniform remainder	117
68 Several separated nonendpoint clusters	118
69 Inside the moment interval: dominant sectors and parity	119
70 Sine-kernel duality and a positive Gram representation	119
70.1 A bilinear Christoffel identity	119
71 Two equal-multiplicity roots: the complete first-order law	121
71.1 Identification with the classical oscillatory Jacobi determinant	122
72 The classical profile at large real frequency	123
73 Finite-size zeros and convergent phase-dependent transport	124
73.1 A genuinely convergent expansion, not only an asymptotic one	125
74 Explicit examples and a sharp failure of a phase-free correction	125
74.1 Two simple roots: an exact formula	125
74.2 Two roots of multiplicity two	126
74.3 A repeated root at the collision center	127
75 Reproducible checks and their evidentiary status	127
75.1 What the verification program does	127
75.2 First-correction data	128
75.3 Zero-transport data	128
75.4 Reproduction and computational limitations	128
76 What is resolved and what is not	129
77 Further research questions and proposed topics	130
77.1 A uniform bridge from interior collisions to endpoint confluence	130
77.2 Mixed endpoint and interior clusters	130
77.3 Moving zero indices and mesoscopic collision coordinates	130
77.4 Zeros of the transport coefficient and suppressed drift	130
77.5 Higher correction operators in a compact basis	130
77.6 Growing degree and conditioning	130
77.7 Eventually constant Jacobi recurrences	131
77.8 Multiple phase tori and resonant zero geometry	131
77.9 Exact certificates, interval numerics, and formalization	131
78 Conclusion	131
79 A proof-dependency map and implementation audit	131
80 Normalization checklist	132
Part VI. Growing shifts	133
81 Provenance and conventions of Part VI	133
81.1 One manuscript, one addition	133
81.2 Reading conventions	135

81.3	Notation of Part VI	136
81.4	What Part VI answers, and what stays open	137
81.5	Where the merge had to choose	138
82	The repository question and the contribution	139
82.1	What is classical and what is being added	139
83	Normalization and a map of the main results	139
84	An exact Jacobi reduction, with normalization checked	140
85	Inverse-zero weights and the exact concentration index	142
85.1	Positive weights from orthogonality	142
85.2	Riccati recurrence	142
86	A uniform bound on the largest inverse zero	143
87	The first critical scale and exact error control	144
88	A size-biased inverse-zero law	144
89	An all-orders hierarchy of critical scales	146
90	Resumming the hierarchy: the full nonlinear scale	147
90.1	Mean and variance on the nonlinear scale	148
91	The second critical scale: displacement of the nominal Poisson law	149
92	A sharp prefactor in proportional growth	150
92.1	The limiting saddle and the coefficient rate function	151
93	Boundary regimes and the meaning in the original variables	152
93.1	What happens when only one parameter grows?	152
93.2	Physical sizes of the perturbation	152
94	Reproducible checks and computational methods	152
94.1	Two thresholds on a common sequence	153
95	Proof dependencies and limitations	154
96	Further research questions	154
96.1	A sharp extreme-zero theorem	154
96.2	Finite-size corrections to the inverse-zero measure	154
96.3	A uniform bridge from fixed to growing multiplicity	155
96.4	Several moving factors and their interactions	155
96.5	Both endpoint exponents growing	155
96.6	Complex perturbations at the first zero edge	155
96.7	Growing truncation order	155
96.8	A direct combinatorial coefficient model	155
96.9	Certified numerical evaluation	155
96.10A	staged formalization	156
97	Conclusion	156
98	A compact formula sheet	156
99	Reproduction instructions	157
A	The precise correction to the printed conjecture	158
B	Compact reference and archive conventions	159

References**160**

1 The problem and the contribution

Let

$$C_n = \frac{1}{n+1} \binom{2n}{n} \quad (n \in \mathbb{N}_0) \quad (1.1)$$

be the Catalan numbers. We use determinant size, rather than the last row index, throughout:

$$D_N^{(m)}(a, b) = \det_{0 \leq i, j < N} (aC_{i+j+m} + bC_{i+j+m+1}), \quad D_0^{(m)} = 1. \quad (1.2)$$

Here $m, N \in \mathbb{N}_0$, and the parameters may be indeterminates or complex numbers. Define the ordinary generating function

$$F_m(t; a, b) = \sum_{N \geq 0} D_N^{(m)}(a, b) t^N. \quad (1.3)$$

All generating-function identities below first make sense as identities of formal power series. Assertions about poles concern the associated rational functions.

Barry's preprint [1], version 1, explicitly conjectures on page 4 that the exponent of the quadratic generating-function denominator is $\binom{m}{2} + 1$. The same page contains a more complicated coefficient conjecture. These are distinct statements: the denominator claim is correct in the uniform sense proved here, whereas the printed coefficient formula uses a shift one smaller than its accompanying determinant. Appendix A makes the translation precise.

There is substantial prior theory. Dougherty, French, Saderholm, and Qian [2] studied recurrences for linear combinations of Catalan numbers. Krattenthaler [3] gives a detailed account of the Christoffel-type determinant identity, confluence, and constant-coefficient recurrences; his Section 8 also records Elouafi's prior proof of general Catalan recurrence assertions. For a multiplier of degree r , that general construction supplies an order at most 2^r . Consequently, rationality by itself is *not* the research claim here. For the special multiplier $x^m(a + bx)$, the objective is the exact, much smaller order, together with its sharp parameter classification.

The article separates two useful calculations. Jacobi orthogonality immediately exposes the individual coefficients. A confluent determinant, on the other hand, exposes the dependence on N and explains the precise recurrence order. Their independent implementations provide a useful check against indexing and normalization errors.

1.1 A notation guide

We shall repeatedly use

$$d = \binom{m}{2}, \quad e = d + 1, \quad K = (m - 1)^2, \quad q(t) = 1 - (a + 2b)t + b^2t^2. \quad (1.4)$$

The shifted pure Catalan determinant and its leading coefficient are

$$H_N^{(m)} = \det_{0 \leq i, j < N} (C_{i+j+m}) = \prod_{1 \leq j \leq i < m} \frac{2N + i + j}{i + j}, \quad (1.5)$$

$$h_m = \prod_{1 \leq j \leq i < m} \frac{2}{i + j}. \quad (1.6)$$

Every empty product and empty determinant equals 1. In particular, $H_N^{(0)} = H_N^{(1)} = 1$ and $h_0 = h_1 = 1$. We prove (1.5) rather than assuming it. As a polynomial in N , $H_N^{(m)}$ has degree d and leading coefficient h_m . The notation $(u)_k = u(u + 1) \cdots (u + k - 1)$ denotes a rising factorial, with $(u)_0 = 1$.

2 Statements of the main theorems

Theorem 2.1 (Explicit coefficients). *For every $N, m \in \mathbb{N}_0$,*

$$D_N^{(m)}(a, b) = \sum_{k=0}^N c_{N,m,k} a^k b^{N-k}, \quad c_{N,m,k} = H_N^{(m+1)} \binom{N}{k} \frac{(N+m+1)_k}{4^k (m+\frac{1}{2})_k}. \quad (2.1)$$

All the coefficients $c_{N,m,k}$ are positive integers. Equivalently,

$$c_{N,m,k} = H_N^{(m+1)} \binom{N}{k} \frac{(N+m+k)!(2m)!(m+k)!}{(N+m)!(2m+2k)!m!}. \quad (2.2)$$

They can be generated without evaluating any determinant:

$$c_{N,m,0} = H_N^{(m+1)}, \quad \frac{c_{N,m,k+1}}{c_{N,m,k}} = \frac{(N-k)(N+m+1+k)}{2(k+1)(2m+2k+1)}. \quad (2.3)$$

Theorem 2.2 (Uniform rational form and sharp denominator). *For every $m \in \mathbb{N}_0$ there is a polynomial $P_m \in \mathbb{Z}[a, b, t]$ such that*

$$F_m(t; a, b) = \frac{P_m(t; a, b)}{q(t)^{d+1}}. \quad (2.4)$$

It satisfies $P_m(0; a, b) = 1$ and $\deg_t P_m \leq K$. If $b \neq 0$, then $\deg_t P_m = K$ and

$$P_m(t; a, b) = s_m (bt)^K P_m\left(\frac{1}{b^2 t}; a, b\right), \quad s_m = (-1)^{(m-1)(m-2)/2}. \quad (2.5)$$

In particular, $[t^K]P_m = s_m b^K$. If $b \neq 0$, $a \neq 0$, and $a \neq -4b$, the representation (2.4) is reduced over $\mathbb{C}[t]$. The sequence has minimal constant-coefficient recurrence characteristic polynomial

$$(X^2 - (a+2b)X + b^2)^{d+1}, \quad (2.6)$$

and hence minimal order $2d+2 = m(m-1)+2$.

Here “minimal” refers even to recurrences allowed to start after a finite initial segment. Thus minimality is not an artifact of the empty-determinant convention. The polynomial numerator in (2.4) remains the *uniform, possibly unreduced* numerator at exceptional parameters.

Theorem 2.3 (Every exceptional denominator). *For fixed complex a, b , the reduced denominator of F_m , normalized to have constant term 1, is exactly the following:*

Parameter condition	Reduced denominator
$a = b = 0$	1
$b = 0, a \neq 0$	$(1 - at)^{\binom{m}{2}+1}$
$a = 0, b \neq 0$	$(1 - bt)^{\binom{m+1}{2}+1}$
$a = -4b, b \neq 0$	$(1 + bt)^{\binom{m}{2}+2}$
$b \neq 0, a(a+4b) \neq 0$	$q(t)^{\binom{m}{2}+1}$

Theorem 2.4 (Finite asymptotic expansion). *Let $a, b > 0$ be fixed and put*

$$r = \frac{a/b + 2 + \sqrt{(a/b)(a/b+4)}}{2} > 1, \quad \lambda_+ = br, \quad \lambda_- = b/r. \quad (2.7)$$

There are explicit degree- d polynomials $B_+(N), B_-(N)$ such that

$$D_N^{(m)}(a, b) = B_+(N)\lambda_+^N + B_-(N)\lambda_-^N \quad (N \geq 0). \quad (2.8)$$

In particular,

$$D_N^{(m)}(a, b) = A_m N^d \lambda_+^N \left(1 + \frac{d \left(\frac{m}{2} - \frac{1}{r-1} \right)}{N} + O(N^{-2}) + O(r^{-2N}) \right), \quad (2.9)$$

$$A_m = h_m \frac{r^{m+1}}{(r-1)^m(r+1)}. \quad (2.10)$$

The entire algebraic expansion terminates after N^{-d} ; the second exponential is also multiplied by a polynomial of degree d . For $d = 0$, there are no algebraic correction terms at all.

2.1 The scale of the improvement

For a degree- $(m+1)$ multiplier the general 2^{m+1} construction need not be minimal. The theorem determines the actual order in this family. The first values are:

m	Exponent e	Exact order $2e$	Numerator degree K	2^{m+1}
0	1	2	1	2
1	1	2	0	4
2	2	4	1	8
3	4	8	4	16
4	7	14	9	32
5	11	22	16	64
6	16	32	25	128
8	29	58	49	512
10	46	92	81	2048

The reduction is not obtained by fitting a shorter recurrence to a long initial segment. It comes from a cancellation of polynomial degree in a family of minors, proved in Section 5.

3 Moments, orthogonality, and the coefficient formula

3.1 Catalan moments and a normalized Jacobi polynomial

For $m \in \mathbb{N}_0$ define a positive measure on $(0, 4)$ by

$$d\mu_m(x) = \frac{1}{2\pi} x^{m-1/2} (4-x)^{1/2} dx. \quad (3.1)$$

The beta integral, after $x = 4t$, gives

$$\int_0^4 x^n d\mu_m(x) = \frac{4^{n+m+1}}{2\pi} B(n+m+\frac{1}{2}, \frac{3}{2}) = C_{n+m}. \quad (3.2)$$

Consequently every moment matrix is positive definite, so its determinant is nonzero and its monic orthogonal polynomials are well-defined.

Lemma 3.1 (The monic polynomial). *The degree- N monic orthogonal polynomial for μ_m is*

$$p_N^{(m)}(x) = (-1)^N \frac{4^N (m + \frac{1}{2})_N}{(N + m + 1)_N} {}_2F_1 \left(\begin{matrix} -N, N + m + 1 \\ m + \frac{1}{2} \end{matrix}; x/4 \right). \quad (3.3)$$

The hypergeometric series terminates after $N + 1$ terms.

Proof. Write $\alpha = m - \frac{1}{2}$, $\beta = \frac{1}{2}$, so both are greater than -1 . A Rodrigues representation on $0 < t < 1$ is

$$\frac{t^{-\alpha}(1-t)^{-\beta}}{(\alpha+1)_N} \frac{d^N}{dt^N} \left(t^{N+\alpha}(1-t)^{N+\beta} \right) = {}_2F_1 \left(\begin{matrix} -N, N + \alpha + \beta + 1 \\ \alpha + 1 \end{matrix}; t \right). \quad (3.4)$$

Leibniz's rule gives the polynomial on the right, including its normalization at $t = 0$. Multiplying the left side by $t^j(1-t)^\beta$ and integrating by parts N times yields zero for $0 \leq j < N$. Every boundary term vanishes: after at most $N - 1$ derivatives, the endpoint exponents that can occur are at least $\alpha + 1 > 0$ and $\beta + 1 > 0$. The final derivative of t^j is zero.

The coefficient of t^N on the right is $(-1)^N (N + \alpha + \beta + 1)_N / (\alpha + 1)_N$. Replacing t by $x/4$ and normalizing this leading coefficient gives (3.3). This also identifies the polynomial with the standard Jacobi family; see the conventional hypergeometric representation in DLMF [4], Eq. 18.5.7. $\square$

3.2 A characteristic-polynomial determinant identity

The following elementary identity will be used twice.

Lemma 3.2. *Let v_j be the moments of a positive measure, let $H_N = \det(v_{i+j})_{0 \leq i, j < N}$, and let p_N be its monic orthogonal polynomial. Then*

$$\det_{0 \leq i, j < N} (v_{i+j+1} - z v_{i+j}) = (-1)^N H_N p_N(z). \quad (3.5)$$

Proof. The monic polynomial is represented by

$$p_N(z) = \frac{1}{H_N} \det \begin{pmatrix} v_0 & v_1 & \cdots & v_N \\ v_1 & v_2 & \cdots & v_{N+1} \\ \vdots & \vdots & & \vdots \\ v_{N-1} & v_N & \cdots & v_{2N-1} \\ 1 & z & \cdots & z^N \end{pmatrix}.$$

Its leading coefficient is one, and integrating it against any power z^j , $j < N$, creates a repeated row. Thus it is indeed p_N . Starting with the last column and moving left, replace col_j by $\text{col}_j - z \text{col}_{j-1}$. The last row becomes $(1, 0, \dots, 0)$. Expansion along it gives (3.5), with sign $(-1)^N$. $\square$

Applying the lemma to μ_m and $z = -a/b$ gives, for $b \neq 0$,

$$D_N^{(m)}(a, b) = (-b)^N H_N^{(m)} p_N^{(m)}(-a/b). \quad (3.6)$$

3.3 A proof of the shifted Catalan product

For $m = 0$, (3.3) reduces to

$$p_n^{(0)}(x) = \sum_{j=0}^n (-1)^{n-j} \binom{n+j}{2j} x^j. \quad (3.7)$$

Its initial values and recurrence are

$$p_0^{(0)} = 1, \quad p_1^{(0)} = x - 1, \quad p_{n+2}^{(0)} = (x - 2)p_{n+1}^{(0)} - p_n^{(0)}. \quad (3.8)$$

These follow directly by comparing the binomial coefficients in (3.7). If η_n denotes the squared norm of $p_n^{(0)}$, the coefficient 1 of $p_{n-1}^{(0)}$ in $x p_n^{(0)}$ implies $\eta_n = \eta_{n-1}$. Indeed, the two ways of writing $\langle x p_n^{(0)}, p_{n-1}^{(0)} \rangle$ give η_{n-1} and η_n , respectively. Since $\eta_0 = C_0 = 1$, all these norms equal 1. Changing from monomials to monic orthogonal polynomials is unitriangular, and therefore $H_N^{(0)} = \prod_{n=0}^{N-1} \eta_n = 1$.

Next set $z = 0$ in (3.5) and use (3.3). We obtain

$$\frac{H_N^{(m+1)}}{H_N^{(m)}} = (-1)^N p_N^{(m)}(0) = \frac{4^N (m + \frac{1}{2})_N}{(N + m + 1)_N} = \prod_{j=1}^m \frac{2N + m + j}{m + j}. \quad (3.9)$$

For completeness, the middle expression equals

$$\frac{(2m + 2N)! m! (N + m)!}{(m + N)! (2m)! (2N + m)!} = \frac{(2N + 2m)! m!}{(2m)! (2N + m)!},$$

which is the last product. Induction on m , beginning with $H_N^{(0)} = 1$, proves (1.5).

3.4 Completion of the coefficient proof

Insert (3.3) into (3.6). The constant factor is $H_N^{(m+1)}$ by (3.9). In each summand, $(-N)_k (-1)^k / k! = \binom{N}{k}$. This gives exactly (2.1). The formula is a polynomial identity for $b \neq 0$, so it holds for $b = 0$ as well. The duplication identity for a rising half-integer factorial, or direct cancellation, gives (2.2). Taking adjacent coefficient ratios gives (2.3).

All factors in (2.1) are positive rational numbers. The left side is a determinant with entries in $\mathbb{Z}[a, b]$, so its coefficients are integers. This proves positivity and integrality simultaneously and completes Theorem 2.1. In particular,

$$c_{N,m,0} = H_N^{(m+1)}, \quad c_{N,m,N} = H_N^{(m)}. \quad (3.10)$$

3.5 Real roots and coefficient inequalities

Corollary 3.3. *For $N \geq 1$, the polynomial $a \mapsto D_N^{(m)}(a, 1)$ has exactly N simple real roots, all in $(-4, 0)$. For $N \geq 2$, its normalized coefficients $u_k = c_{N,m,k} / \binom{N}{k}$ satisfy*

$$u_k^2 > u_{k-1} u_{k+1} \quad (1 \leq k < N). \quad (3.11)$$

Proof. A degree- N orthogonal polynomial for a strictly positive density on $(0, 4)$ has N simple roots in that interval. One elementary proof is to multiply it by the product of its distinct sign-changing roots in $(0, 4)$. If there were fewer than N such roots, that product would have degree below N , while the integral of its product with the orthogonal polynomial would have constant nonzero sign. This contradicts orthogonality. Equation (3.6) now negates the roots.

Alternatively, the coefficient inequality follows without any root theory. Theorem 2.1 gives

$$\frac{u_{k+1}}{u_k} = \frac{N + m + 1 + k}{4(m + \frac{1}{2} + k)}.$$

This is strictly decreasing in k , because the numerator's constant term exceeds the denominator's unscaled constant term by $N + 1/2$. The decrease is equivalent to (3.11). $\square$

4 A fixed-size confluent determinant

The explicit coefficient formula grows in length with N . To understand the recurrence order, we need a representation whose determinant size depends only on m . Define

$$R_n(y) = (-1)^n p_n^{(0)}(-y) = \sum_{i=0}^n \binom{n+i}{2i} y^i. \quad (4.1)$$

Thus

$$R_0(y) = 1, \quad R_1(y) = y + 1, \quad R_{n+2}(y) = (y + 2)R_{n+1}(y) - R_n(y). \quad (4.2)$$

Extend R_n to all integer indices using this recurrence. Since $R_{-1} = R_0 = 1$, it also satisfies the reflection identity

$$R_{-n-1}(y) = R_n(y) \quad (n \in \mathbb{Z}). \quad (4.3)$$

The reversed form of the polynomials is convenient because it eliminates unnecessary signs from Christoffel's identity.

Lemma 4.1 (Christoffel determinant in the present normalization). *For pairwise distinct $y_1, \dots, y_r$,*

$$\det_{0 \leq i, j < N} \int_0^4 x^{i+j} \prod_{\ell=1}^r (x + y_\ell) d\mu_0(x) = \frac{\det[R_{N+j}(y_i)]_{\substack{1 \leq i \leq r \\ 0 \leq j < r}}}{\prod_{1 \leq i < j \leq r} (y_j - y_i)}. \quad (4.4)$$

The quotient extends polynomially when some y_i coincide.

Proof. Here is a proof that also verifies the normalization. Write $\Delta(x_1, \dots, x_N) = \det[x_i^{j-1}]_{1 \leq i, j \leq N}$. Expanding two determinants and integrating termwise gives the identity

$$\frac{1}{N!} \int \cdots \int \det[f_i(x_j)] \det[g_i(x_j)] \prod_j d\mu_0(x_j) = \det \left[\int f_i(x) g_j(x) d\mu_0(x) \right].$$

No analytic interchange difficulty occurs here: the sums are finite and all polynomials are integrable on a bounded interval.

For external variables $z_1, \dots, z_r$ we have

$$\Delta(x, z) = \Delta(x) \Delta(z) \prod_{i=1}^N \prod_{\ell=1}^r (z_\ell - x_i).$$

In the determinant defining $\Delta(x, z)$, replace monomials by the monic polynomials $p_j^{(0)}$; the determinant is unchanged. Expand along its first N rows and integrate against $\Delta(x)$. By orthogonality, every choice of columns vanishes except columns of degrees $0, \dots, N-1$. That surviving integral equals $N! H_N^{(0)} = N!$. The remaining minor is $\det[p_{N+j}^{(0)}(z_i)]$. Dividing by $\Delta(z)$ yields the identity for the multiplier $\prod_\ell (z_\ell - x)$.

Finally put $z_\ell = -y_\ell$ and use $p_{N+j}^{(0)}(-y) = (-1)^{N+j} R_{N+j}(y)$. The signs from the multiplier, column factors, and Vandermonde cancel, giving (4.4). The numerator is alternating in the y_i , hence divisible by their Vandermonde as a polynomial. This proves the extension statement. This is the classical fixed-size identity discussed in [3]; its inclusion here is a normalization proof, not a novelty claim. $\square$

Take $r = m + 1$, let the first m variables tend to zero, and put the last one equal to $y = a/b \neq 0$. Dividing by the Vandermonde and taking the limit replaces those first rows by Taylor coefficients. Since $[y^i] R_{N+j}(y) = \binom{N+j+i}{2i}$, we obtain:

Proposition 4.2 (Confluent representation). *If $ab \neq 0$, then*

$$D_N^{(m)}(a, b) = b^N y^{-m} \det \begin{pmatrix} \binom{N+j+i}{2i} & (0 \leq i < m, 0 \leq j \leq m) \\ R_{N+j}(y) & (0 \leq j \leq m) \end{pmatrix}, \quad y = a/b. \quad (4.5)$$

The upper block has m rows and the lower block has one row. For $m = 0$ the formula simply reads $D_N^{(0)} = b^N R_N(a/b)$.

Define the $m \times m$ minors

$$M_{m,\ell}(N) = \det_{\substack{0 \leq i < m \\ j \in J_\ell}} \binom{N+j+i}{2i}, \quad J_\ell = \{0, 1, \dots, m\} \setminus \{\ell\}, \quad (4.6)$$

where columns are in increasing order and $0 \leq \ell \leq m$. Expansion of (4.5) along its last row gives

$$D_N^{(m)}(a, b) = b^N y^{-m} \sum_{\ell=0}^m (-1)^{m+\ell} M_{m,\ell}(N) R_{N+\ell}(y). \quad (4.7)$$

Although a naive degree bound for $M_{m,\ell}$ is $m(m-1) = 2d$, its actual degree is only d . This is the key point.

5 The degree cancellation and its first correction

Lemma 5.1 (Sharp minor degree). *For $0 \leq \ell \leq m$, $M_{m,\ell}(N)$ is a polynomial of degree $d = \binom{m}{2}$ with leading coefficient*

$$[N^d] M_{m,\ell}(N) = h_m \binom{m}{\ell}. \quad (5.1)$$

For $d \geq 1$, the next coefficient is

$$[N^{d-1}] M_{m,\ell}(N) = h_m \binom{m}{\ell} \left(\frac{d(m+2)}{2} - \frac{(m-1)\ell}{2} \right). \quad (5.2)$$

Proof. Let $f_i(x) = \binom{x+i}{2i}$, so f_i has degree $2i$ and leading coefficient $1/(2i)!$. The determinant $\det[f_i(x_j)]_{0 \leq i, j < m}$ is alternating in its m independent variables. It is divisible by $\Delta(x) = \prod_{p < q} (x_q - x_p)$, whose degree is d . The determinant has total degree at most $\sum_i 2i = 2d$; its quotient by Δ therefore has total degree at most d . Substituting $x_j = N + j$ for $j \in J_\ell$ makes Δ a nonzero constant. Hence the degree in N is at most d .

To obtain the leading coefficient, expand $f_i(N + j)$ in Taylor powers of j . The first possible nonzero determinant uses distinct Taylor orders $0, 1, \dots, m-1$; these have the least possible sum d . The coefficient of N^d is

$$\frac{\det[\binom{2i}{r}]_{0 \leq i, r < m}}{\prod_{i=0}^{m-1} (2i)!} \Delta(J_\ell) = \frac{2^d \Delta(J_\ell)}{\prod_{i=0}^{m-1} (2i)!}. \quad (5.3)$$

To see the second equality, regard $\binom{2i}{r}$ as a polynomial of degree r in i with leading coefficient $2^r/r!$ and evaluate its determinant at $i = 0, \dots, m-1$. The Vandermonde is $\prod_{r=1}^{m-1} r!$, so these factorials cancel. Removing ℓ from $0, \dots, m$ also gives

$$\Delta(J_\ell) = \binom{m}{\ell} \prod_{r=1}^{m-1} r!. \quad (5.4)$$

Since

$$h_m = \frac{2^d \prod_{r=1}^{m-1} r!}{\prod_{i=0}^{m-1} (2i)!} \quad (m \geq 1),$$

we obtain (5.1). It is nonzero, proving the degree assertion. The case $m = 0$ is the empty determinant and is immediate.

For the next coefficient, observe the symmetric factorization

$$f_i(x) = \frac{1}{(2i)!} \prod_{r=1}^i \left(\left(x + \frac{1}{2}\right)^2 - \left(r - \frac{1}{2}\right)^2 \right). \quad (5.5)$$

Thus $f_i(x) = (x + \frac{1}{2})^{2i} / (2i)!$ plus terms of degree at most $2i - 2$. Replacing one or more rows by those lower-degree terms reduces the total degree by at least two. The same alternating argument then bounds their degree after translation by $d - 2$. The leading two terms are consequently obtained from

$$\det \left[\frac{(N + j + \frac{1}{2})^{2i}}{(2i)!} \right]_{0 \leq i < m, j \in J_\ell} = \frac{\prod_{\substack{p < q \\ p, q \in J_\ell}} (q - p)(2N + p + q + 1)}{\prod_{i=0}^{m-1} (2i)!}. \quad (5.6)$$

The relative coefficient of N^{d-1} is

$$\frac{1}{2} \sum_{\substack{p < q \\ p, q \in J_\ell}} (p + q + 1) = \frac{1}{2} \left((m-1) \left(\frac{m(m+1)}{2} - \ell \right) + d \right),$$

which simplifies to (5.2). If $d = 1$, the putative remainder of degree $d - 2 = -1$ is identically zero. $\square$

It is useful to collect the minors in a polynomial in an auxiliary variable r :

$$Q_m(N; r) = \sum_{\ell=0}^m (-1)^{m+\ell} M_{m,\ell}(N) r^\ell. \quad (5.7)$$

By the binomial theorem,

$$[N^d] Q_m(N; r) = h_m(r - 1)^m. \quad (5.8)$$

When $r \neq 1$ and $d \geq 1$, the relative next coefficient is

$$\frac{[N^{d-1}] Q_m(N; r)}{[N^d] Q_m(N; r)} = \frac{d(m+2)}{2} - d \frac{r}{r-1} = d \left(\frac{m}{2} - \frac{1}{r-1} \right). \quad (5.9)$$

Indeed, applying $r \, d/dr$ to $(r - 1)^m$ sums the terms weighted by ℓ in (5.2). The exact value d , rather than an upper bound $2d$, is what makes the denominator exponent sharp.

6 Two exponentials and the minimal recurrence

Suppose first that $b \neq 0$ and $a(a + 4b) \neq 0$. Set $y = a/b$ and choose one ordering of the two roots

$$r_{\pm} = \frac{y + 2 \pm \sqrt{y(y+4)}}{2}, \quad r_+ r_- = 1. \quad (6.1)$$

They are distinct and nonzero. Solving the recurrence (4.2) with its two initial values gives

$$R_n(y) = \frac{(r_+ - 1)r_+^n - (r_- - 1)r_-^n}{r_+ - r_-}. \quad (6.2)$$

Write

$$\lambda_{\pm} = br_{\pm} = \frac{a + 2b \pm \sqrt{a(a+4b)}}{2}. \quad (6.3)$$

Substitution in (4.7) gives the exact identity

$$D_N^{(m)}(a, b) = \frac{(r_+ - 1)Q_m(N; r_+) \lambda_+^N - (r_- - 1)Q_m(N; r_-) \lambda_-^N}{y^m(r_+ - r_-)}. \quad (6.4)$$

In particular, both polynomial coefficients in this two-exponential representation have degree *exactly* d : neither r_+ nor r_- can equal 1, since $y \neq 0$.

Lemma 6.1 (Minimal annihilator of an exponential polynomial). *If $\lambda_1, \dots, \lambda_s$ are distinct nonzero complex numbers and the nonzero polynomials $A_i(N)$ have degrees d_i , then $u_N = \sum_i A_i(N) \lambda_i^N$ has minimal constant-coefficient annihilator*

$$\prod_{i=1}^s (E - \lambda_i)^{d_i+1}, \quad Eu_N = u_{N+1}. \quad (6.5)$$

Its generating function has poles of orders $d_i + 1$ at $t = 1/\lambda_i$.

Proof. On $A(N)\lambda^N$, the operator $E - \lambda$ is $\lambda^{N+1}(A(N+1) - A(N))$; it lowers the polynomial degree by exactly one until the result vanishes. For $\mu \neq \lambda$, $E - \mu$ preserves that degree, multiplying the leading coefficient by $\lambda - \mu$. The generalized eigenspaces for distinct eigenvalues are independent: polynomial projectors from the Chinese remainder identity isolate each one. Thus every annihilator needs the stated multiplicity at every eigenvalue. This argument remains valid after any finite shift in N .

Equivalently, applying $(z d/dz)^k$ to $1/(1-z)$ shows that $\sum_{N \geq 0} N^k z^N$ has a pole of exact order $k+1$ at $z=1$, with nonzero leading pole coefficient $k!$. Distinct pole locations cannot cancel. This proves the generating-function assertion too. $\square$

Applying the lemma to (6.4) proves the minimal characteristic polynomial (2.6). It also proves that $q^{d+1}F_m$ is a polynomial of degree at most $2d+1 = 2e-1$ for these parameters. Section 7 sharpens its degree to K .

6.1 Why the identity extends to all parameters

Each $D_N^{(m)}$ is a polynomial in a, b with integer coefficients. The coefficients of $q^{d+1}F_m$ are therefore such polynomials too. Every coefficient that vanishes for all $b \neq 0$ and $a(a+4b) \neq 0$ vanishes identically: a polynomial cannot be nonzero and vanish on that nonempty open set. Hence the uniform rational identity, and its coefficient integrality, extend to every a, b , even where its displayed denominator is not reduced. This argument is coefficientwise in a formal series and does not require a convergence limit in t .

6.2 Proof of the exceptional classification

When $b = 0$,

$$D_N^{(m)}(a, 0) = a^N H_N^{(m)}. \quad (6.6)$$

For $a \neq 0$, the polynomial $H_N^{(m)}$ has degree d , so Lemma 6.1 gives denominator $(1-at)^{d+1}$. When $a = 0$ and $b \neq 0$,

$$D_N^{(m)}(0, b) = b^N H_N^{(m+1)}, \quad (6.7)$$

whose polynomial degree is $\binom{m+1}{2}$.

At the remaining collision $a = -4b$, with $b \neq 0$, the recurrence gives

$$R_n(-4) = (-1)^n (2n+1). \quad (6.8)$$

Inserting this into (4.7) and tracking the signs produces

$$D_N^{(m)}(-4b, b) = (-b)^N E_m(N), \quad (6.9)$$

$$E_m(N) = 4^{-m} \sum_{\ell=0}^m (2N + 2\ell + 1) M_{m,\ell}(N). \quad (6.10)$$

By (5.1), its degree is exactly $d + 1$, with

$$[N^{d+1}]E_m(N) = 4^{-m} 2h_m \sum_{\ell=0}^m \binom{m}{\ell} = 2^{1-m} h_m \neq 0. \quad (6.11)$$

Thus the reduced denominator is $(1 + bt)^{d+2}$. Finally, if $a = b = 0$, every nonempty determinant vanishes and $F_m = 1$. Together with the nonexceptional result, this proves Theorem 2.3.

The two root-collision lines have different pole orders. This is not a contradiction: $x = 0$ and $x = 4$ are inequivalent endpoints of the weight $x^{m-1/2}(4-x)^{1/2}$. In particular, blindly substituting $a = 0$ into a generic minimal-order statement would be incorrect.

7 The numerator degree and reciprocal symmetry

A rational function can have a denominator of large degree and a surprisingly short numerator. Here the explanation is a string of zero values in the natural bilateral extension of the determinant.

For the moment retain the nonexceptional hypotheses. Define D_N for all $N \in \mathbb{Z}$ by the right side of (4.5). The upper rows are interpreted through their polynomial binomial coefficients; this agrees with the Taylor coefficients of R_n at zero, including negative n , because

$$\binom{-n-1+i}{2i} = \binom{n+i}{2i}.$$

It also agrees with the unique bilateral extension supplied by (6.4).

Lemma 7.1 (Reflection and a zero interval). *Let $\sigma_m = (-1)^{\binom{m+1}{2}}$. Then*

$$D_{-N-m-1} = \sigma_m b^{-2N-m-1} D_N \quad (N \in \mathbb{Z}), \quad (7.1)$$

$$D_{-1} = D_{-2} = \cdots = D_{-m} = 0. \quad (7.2)$$

The second statement is empty for $m = 0$.

Proof. The index change $N \mapsto -N - m - 1$, followed by $R_{-n-1} = R_n$, reverses all $m + 1$ columns of the determinant. The sign of that reversal is σ_m . Comparing its prefactor b^N on the two sides gives (7.1). If $N = -s$ with $1 \leq s \leq m$, columns $j = s - 1$ and $j = s$ correspond to R_{-1} and R_0 , respectively, which are the same polynomial. Their values and all needed Taylor coefficients coincide, so the determinant vanishes. $\square$

Lemma 7.2 (The expansion at infinity). *For a sequence of exponential-polynomial form with nonzero characteristic roots and its bilateral extension, its proper rational generating function satisfies*

$$F(t) = - \sum_{r \geq 1} D_{-r} t^{-r} \quad (7.3)$$

as a formal Laurent expansion at infinity.

Proof. For $D_N = \lambda^N$ the assertion is simply $(1 - \lambda t)^{-1} = - \sum_{r \geq 1} \lambda^{-r} t^{-r}$. Apply powers of $t \, d/dt$ to obtain the assertion for $D_N = N^k \lambda^N$, and then use linearity. $\square$

Combining (7.2), (7.1), and $D_0 = 1$ yields

$$F_m(t) = -\sigma_m b^{-m-1} t^{-m-1} + O(t^{-m-2}) \quad (t \rightarrow \infty). \quad (7.4)$$

Since q^e has degree $2e$ and leading coefficient b^{2e} , the numerator has degree

$$2e - m - 1 = m(m - 1) + 2 - m - 1 = (m - 1)^2 = K \quad (7.5)$$

and leading coefficient $-\sigma_m b^{2e-m-1} = s_m b^K$. The identity $s_m = -\sigma_m$ follows because the difference of the two exponents $\binom{m+1}{2}$ and $(m - 1)(m - 2)/2$ is $2m - 1$, an odd integer. This calculation proves the sharp degree assertion, including $m = 0$.

The full reflection identity yields more than the leading term. Putting $1/(b^2 t)$ into (7.3) and reindexing with (7.1), we find

$$F_m\left(\frac{1}{b^2 t}\right) = s_m b^{m+1} t^{m+1} F_m(t). \quad (7.6)$$

But $q(1/(b^2 t)) = q(t)/(b^2 t^2)$. Substitution into (7.6) gives exactly (2.5). After denominators in b are cleared, each of these numerator identities is polynomial in a, b . Thus they extend to the exceptional parameters; for $b \neq 0$ the leading coefficient remains nonzero. This completes the proof of Theorem 2.2.

7.1 An explicit finite prescription for the numerator

No uncomputed infinite series is needed to determine P_m . Expand

$$q(t)^e = \sum_{j=0}^{2e} q_j t^j. \quad (7.7)$$

For example, its coefficients have the finite expression

$$q_j = \sum_{v=\max(0, j-e)}^{\lfloor j/2 \rfloor} \frac{e!}{(j-2v)! v! (e-j+v)!} (-(a+2b))^{j-2v} b^{2v}. \quad (7.8)$$

Then

$$P_m(t) = \sum_{\ell=0}^K \left(\sum_{j=0}^{\min(\ell, 2e)} q_j D_{\ell-j}^{(m)}(a, b) \right) t^\ell, \quad (7.9)$$

where every determinant in the right side is given by Theorem 2.1. Equations (7.8) and (7.9) therefore constitute a finite sum/product formula for every numerator coefficient. All coefficients of $q^e F_m$ after degree K vanish.

The usual recurrence is

$$\sum_{j=0}^{2e} q_j D_{N-j}^{(m)}(a, b) = 0 \quad (N \geq 2e). \quad (7.10)$$

In fact the truncated convolution already vanishes for $N > K$ when negative-index terms are taken as zero for the purpose of multiplying ordinary power series. Those artificial zeros should not be confused with the distinct bilateral extension used in the proof above.

8 Exact exponential expansions and asymptotics

Equation (6.4) supplies every coefficient of the two polynomials in Theorem 2.4. Explicitly,

$$B_+(N) = \frac{(r_+ - 1)Q_m(N; r_+)}{y^m(r_+ - r_-)}, \quad B_-(N) = -\frac{(r_- - 1)Q_m(N; r_-)}{y^m(r_+ - r_-)}. \quad (8.1)$$

These formulas are valid for arbitrary nonexceptional complex parameters; the assumption $a, b > 0$ is only needed to select a strictly dominant real exponential for the asymptotic statement.

For $a, b > 0$, write $r = r_+$, so $r_- = r^{-1}$ and

$$y = r + r^{-1} - 2 = \frac{(r-1)^2}{r}. \quad (8.2)$$

Using (5.8), the coefficient of N^d in B_+ is

$$\frac{h_m(r-1)^{m+1}}{y^m(r-r^{-1})} = h_m \frac{r^{m+1}}{(r-1)^m(r+1)} = A_m.$$

Its relative next coefficient is exactly (5.9). Also $\lambda_-/\lambda_+ = r^{-2}$. Division of (6.4) by $A_m N^d \lambda_+^N$ proves (2.9).

One may write the complete answer in the finite form

$$D_N^{(m)} = \lambda_+^N \sum_{j=0}^d \beta_j^+ N^{d-j} + \lambda_-^N \sum_{j=0}^d \beta_j^- N^{d-j}, \quad (8.3)$$

where

$$\beta_j^+ = \frac{r_+ - 1}{y^m(r_+ - r_-)} [N^{d-j}] Q_m(N; r_+), \quad \beta_j^- = -\frac{r_- - 1}{y^m(r_+ - r_-)} [N^{d-j}] Q_m(N; r_-). \quad (8.4)$$

Thus the “full asymptotic expansion” is particularly strong here: it is an exact expression, with no uncomputed remainder once both finite sectors are retained. No assertion of asymptotic convergence or summability is needed. The leading coefficient ratio of the two sectors is

$$\frac{\beta_0^-}{\beta_0^+} = (-1)^m r^{-(m+1)}. \quad (8.5)$$

These estimates are for fixed m, a, b . They are not claimed to be uniform when $a \rightarrow 0$, when the two roots coalesce, or when m grows with N . In such regimes the exact finite formula remains valid away from the collision itself, but a different uniform asymptotic analysis would be required. At a collision, use the exact exceptional formulas from Section 6 instead.

[Added 29 September 2026, batch 57: for this family with real $a/b \geq 0$, limits with m growing with N are treated in Part VI. With $v = m + \frac{1}{2}$, $B = N + m + 1$ and $\tau = (a/b)NB/(4v)$, the ratio $D_N^{(m)}(a, b)/D_N^{(m)}(0, b)$ tends to 1 exactly when $\tau \rightarrow 0$, for every relative growth of N and m (Corollary 84.2); when $N, m \rightarrow \infty$ its logarithm is governed by $\beta = (1 + c(v + \frac{3}{2}))/(v + 1)$, $c = v/(NB)$, with critical scales $\tau \asymp \beta^{-1/2}, \beta^{-2/3}$ and β^{-1} (Theorem 83.1). Coalescing roots and $a/b < 0$ are not treated there.]

9 Worked examples and integer sequences

9.1 Small shifts

The first uniform generating functions are

$$F_0(t) = \frac{1 - bt}{q(t)}, \quad F_1(t) = \frac{1}{q(t)}, \quad (9.1)$$

$$F_2(t) = \frac{1 + bt}{q(t)^2}, \quad F_3(t) = \frac{(1 - b^2 t^2)(1 + (a + 6b)t + b^2 t^2)}{q(t)^4}. \quad (9.2)$$

They can be obtained directly from (7.9). Notice that $m = 0$ is a genuine boundary case: its numerator has degree 1, not 0. For $m = 3$, the numerator is anti-reciprocal; for $m = 2$, it is reciprocal. This is the pattern described uniformly by s_m .

For $m = 2$, put $A = a + 2b$. The recurrence becomes

$$D_N = 2AD_{N-1} - (A^2 + 2b^2)D_{N-2} + 2Ab^2D_{N-3} - b^4D_{N-4} \quad (N \geq 4). \quad (9.3)$$

The initial polynomials are

$$\begin{aligned} D_0 &= 1, \\ D_1 &= 2a + 5b, \\ D_2 &= 3a^2 + 14ab + 14b^2, \\ D_3 &= 4a^3 + 27a^2b + 54ab^2 + 30b^3. \end{aligned}$$

The recurrence is minimal outside the exceptional lines; displaying only a valid order-four relation would not by itself prove that fact.

9.2 The specialization $a = b = 1$

The first sequences, all beginning with the empty determinant, are

$m \backslash N$	0	1	2	3	4	5	6
0	1	2	5	13	34	89	233
1	1	3	8	21	55	144	377
2	1	7	31	115	390	1254	3893
3	1	19	170	1075	5580	25529	107036

The first two are F_{2N+1} and F_{2N+2} , respectively, as follows from their common recurrence $u_{N+2} = 3u_{N+1} - u_N$ and initial values. They connect the family to the odd and even Fibonacci subsequences in OEIS A001519 and A001906 [6]; Catalan numbers themselves are A000108 [5]. The higher rows and their coefficient triangles are natural OEIS-style objects, but the archive does not assert that any particular row is absent from OEIS.

For every fixed shift, the dominant exponential is

$$\lambda_+ = \frac{3 + \sqrt{5}}{2},$$

while its polynomial prefactor has degree $\binom{m}{2}$. Thus increasing the shift changes the power of N , not the exponential growth constant. For $m = 2$, the first relative asymptotic correction equals $1 - 1/(r - 1) = (3 - \sqrt{5})/2$ with $r = (3 + \sqrt{5})/2$.

9.3 Rational characteristic roots: $a = 1, b = 2$

This specialization gives $r_+ = 2, r_- = 1/2$ and $\lambda_+ = 4, \lambda_- = 1$. Therefore the exact formulas have rational polynomial coefficients and no radicals:

$$D_N^{(0)}(1, 2) = \frac{2 \cdot 4^N + 1}{3}, \quad (9.4)$$

$$D_N^{(1)}(1, 2) = \frac{4 \cdot 4^N - 1}{3}, \quad (9.5)$$

$$D_N^{(2)}(1, 2) = \frac{8N 4^N + N + 3}{3}, \quad (9.6)$$

$$D_N^{(3)}(1, 2) = \frac{(32N^3 + 48N^2 + 16N + 96)4^N - (2N^3 + 21N^2 + 73N + 78)}{18}. \quad (9.7)$$

For example, the $m = 2$ sequence starts 1, 12, 87, 514, 2733, 13656, ... Its dominant polynomial is $(8/3)N$ with no constant term. This vanishing agrees with the first-correction formula $d(m/2 - 1/(r - 1)) = 1(1 - 1) = 0$; the entire correction is the smaller exponential sector $(N + 3)/3$. For $m = 3$, the relative coefficient of N^{-1} in the dominant sector is $3/2$, again exactly as predicted.

9.4 An endpoint example

For $m = 2$, (6.10) simplifies to

$$E_2(N) = \frac{(N + 1)(N + 2)}{2}.$$

Hence

$$D_N^{(2)}(-4b, b) = (-b)^N \frac{(N + 1)(N + 2)}{2}, \quad F_2(t; -4b, b) = \frac{1}{(1 + bt)^3}. \quad (9.8)$$

This also follows by cancelling $1 + bt$ from $(1 + bt)/(1 + bt)^4$, the uniform form of F_2 . At the other collision, $a = 0$, one instead obtains $F_2(t; 0, b) = (1 + bt)/(1 - bt)^4$. These explicit cases illustrate why the reduced denominator needs a separate classification.

10 Algorithms, exact experiments, and reproducibility

10.1 Implementation choices

The file `code/catalan_hankel.py` implements the coefficient formula, pure shifted product, homogeneous polynomial evaluation, fixed-size determinant, and uniform and reduced denominators. All numerical parameters in its public computation interface are integers; internal fractions use Python's exact `Fraction`. The mathematical theorems have the broader complex-parameter scope stated above. No floating-point arithmetic enters the tests.

For a coefficient row, first compute the triangular product $H_N^{(m+1)}$ and then apply (2.3) successively. The number of arithmetic operations is $O(m^2 + N)$; this is an *arithmetic-operation count*, not a bit-complexity claim. Every division in the update is checked for exactness. Homogeneous Horner evaluation avoids division by a or b , so zero parameters are handled without special numerical limits.

The independent determinant path uses a fraction-free Bareiss algorithm on the original matrix of Catalan moments, including pivoting when a diagonal pivot vanishes. This matters for signed parameters: an implementation tested only on positive definite examples can conceal pivot failures. The source-indexing formula is implemented separately, not obtained by calling the corrected coefficient routine.

For many consecutive values at fixed m , it is also possible to use (7.10) after producing the first $2e$ values. For very large isolated indices, powering the companion matrix or reducing X^N modulo (2.6) provides standard recurrence-based methods. These methods are consequences of the theorem; the supplied reference implementation prioritizes transparent exact verification rather than optimized large-integer throughput.

10.2 Verification that was actually run

The included `data/verification.json` records PASS for all checks in the following table. The report is generated by the test program, not manually written.

Check	Count
Direct determinant versus coefficient evaluation	1,183
Additional direct signed/degenerate parameter evaluations	810
Printed source formula versus corrected shift	990
Confluent fixed-size determinant evaluations	364
Minor polynomials exactly interpolated and checked	36
Generating-function/recurrence coefficient identities	2,816
Numerator reciprocal coefficient identities	604
Nonexceptional numerator–denominator gcd checks	44
Exceptional reduced-denominator checks	44
Exact two-exponential evaluations	168
Strict normalized log-concavity inequalities	5,655

The first 1,183 evaluations cover every pair $0 \leq N, m \leq 12$, using $a = 0, 1, \dots, N$ and $b = 1$. At each fixed (N, m) , both sides are polynomials of degree at most N in a . Equality at $N + 1$ distinct points certifies that particular polynomial identity exactly. Thus this part certifies 169 full coefficient polynomials, not merely 169 evaluations. It still does not prove the result for untested values of N or m ; the proof in Section 3 does that.

For $0 \leq m \leq 7$, the minor tests interpolate using $2d + 2$ points, more than enough for the elementary degree bound $2d$. They verify the drop to degree d , both leading coefficients from Lemma 5.1, and additional negative-index values. The recurrence checks include $0 \leq m \leq 10$ and four nonexceptional parameter pairs, while separate tests examine every exceptional class. The exact two-exponential tests use $(a, b) = (1, 2)$, $m \leq 7$, and $0 \leq N \leq 20$. These tests check different parts of the derivation, but are not represented as formal proof certificates for arbitrary parameters.

10.3 How to reproduce the package

From the archive's top directory, run:

```
python3 code/verify.py --out data
python3 code/catalan_hankel.py 8 3 --a 2 --b 3 --direct
latexmk -pdf -interaction=nonstopmode -halt-on-error article.tex
```

Use Python 3.10 or later and do not pass -0 , because the verification suite deliberately uses assertions. The script writes coefficient rows, generating functions, minor polynomials, exact exponential-sector polynomials, and integer sequence tables in JSON or CSV. Rational polynomial coefficients are stored as strings such as "8/3", in ascending power order, to avoid any loss of exactness when imported into other systems. The included README documents every file and its indexing convention. [Added 28 September 2026, batch 39: the first command regenerates the shipped files in data/; in the repository, run it on a copy (see the report README). Part II's separate suite, and why its delivered commands must not be run in this directory, are described in Section 27.]

11 What has been resolved, and what has not

The all-shift denominator formula is proved here, with a stronger minimality assertion and a complete description of the degenerate parameters. The coefficient expression is likewise proved for all N, m ; comparison with the exact formula printed in the target source requires the explicit indexing correction in the next appendix. The reciprocal numerator and finite exponential expansions are proved from the same structural determinant calculation.

These are substantive mathematical conclusions about the selected conjectural family. They are not evidence that the Christoffel identity, Jacobi representation, Catalan shifted product, or general rationality

are new. A literature search was performed against the source title, arXiv identifier, denominator phrase, and related recurrence work. The cited version still labels the sharp exponent conjectural, and the retrieved sources did not supply an explicit statement of this full sharp classification. That is a bounded search result, not a proof that no earlier equivalent theorem exists. Before a priority claim or submission, the exact formulas should be checked against the wider orthogonal-polynomial and determinant literature and by a human referee.

Several natural extensions remain separate from the theorem. One may seek a direct combinatorial interpretation of the individual positive integers $c_{N,m,k}$, or a product formula for the minors $M_{m,\ell}(N)$ beyond their first two coefficients. Multiple distinct linear factors in the multiplier lead to more exponential sectors and new collision patterns. A joint limit with growing m or a parameter approaching $a = 0$ would require uniform, rather than fixed-parameter, asymptotics. None of these extensions is assumed in the present proofs.

[Added 28 September 2026, batch 39: the preceding paragraph describes the report at the state fbba58593 against which Part II was written. The extension to several distinct linear factors, and to repeated factors, is treated in Part II: for every nonzero polynomial multiplier the exponential sectors, their exact degrees and nonzero leading coefficients (Theorems 14.2 and 18.1), the minimal order off resonance (14.12), an exact detector of the parameters where it drops (Theorem 14.5), and a certified finite algorithm on that resonance locus (Theorems 20.1 and 20.2). Part II gives no short closed classification of the resonant cancellations; it poses that as a research direction (Section 24). The other extensions named here remain open: Part II does not address a combinatorial interpretation of $c_{N,m,k}$, a product formula for the minors $M_{m,\ell}(N)$, or joint limits with growing m ; and its asymptotics (Section 22) are for fixed parameters, so a parameter approaching $a = 0$ (a root reaching the endpoint 0) is among the uniform endpoint limits it poses, not solves.]

[Added 29 September 2026, batch 42: for a single repeated root away from the endpoints, Part III gives that classification (Theorems 30.1, 30.4 and 37.1); resonant cancellations among several distinct roots, or with endpoint factors, remain unclassified.]

[Added 29 September 2026, batch 53: for fixed m the limit $a \rightarrow 0$ is treated in Part IV. For $b \neq 0$, $x^m(a + bx)$ is b times $x^m(x + a/b)$, a cluster of $m + 1$ roots at the endpoint 0; for real $a/b \geq 0$ the ratio $D_N^{(m)}(a, b)/D_N^{(m)}(0, b)$ tends to 1 exactly when $a/b = o(N^{-2})$, to an explicit profile when $N^2 a/b$ converges, and to infinity when $N^2 a/b \rightarrow \infty$ (Theorem 47.4 and Section 44.4). Joint limits with growing m remain open.]

[Added 29 September 2026, batch 57: joint limits with growing m are treated in Part VI for this one-factor family with $a/b \geq 0$ (Theorems 83.1, 88.2, 90.1 and 92.1); for fixed m its criterion reduces to Part IV's $N^2 a/b \rightarrow 0$. A combinatorial interpretation of $c_{N,m,k}$ and a product formula for $M_{m,\ell}(N)$ remain open.]

Part II. Root collisions: arbitrary polynomial multipliers

Sections 1–11 and Appendices A–B form the original report (Part I) of 19 September 2026, unchanged except for dated pointers to this addition, a line on the title page, contents entries for the two parts, and two bibliography entries used only by Part II. Part II was added on 28 September 2026 in batch 39 of ProveIt’s incoming-report intake, from one manuscript written against the pinned state of Part I. It answers the extension that Part I’s Section 11 named but left open, “multiple distinct linear factors in the multiplier” and their “new collision patterns”: for every nonzero polynomial multiplier it determines the exponential sectors, their exact polynomial degrees and leading coefficients, the minimal recurrence off resonance, and a finite algorithm on the resonance locus. Section 12 records the provenance and fixes the notation of the addition.

12 Provenance and conventions of Part II

12.1 One manuscript, one addition

Part II is batch-39 manuscript 03, *Root Collisions and Sharp Recurrences for Polynomially Weighted Catalan Hankel Determinants*, a proof-focused research article prepared with OpenAI ChatGPT for Vladimir Reshetnikov’s ProveIt research program and dated 28 September 2026 (23 pages as delivered). It arrived as the archive `ProveIt_Catalan_Root_Collisions.zip` (inner directory `ProveIt_Catalan_Root_Collisions/`, main file `article.tex`) in ProveIt commit `74f7f5bdb` and was placed in `e2b1f016a`. It was written against ProveIt commit `fbba58593dc0622aa914972896150d4848f935b5`, reading Part I’s `article.tex` and `README.md` at that commit (consulted 28 September 2026 through the pinned repository tree). Part I’s files are unchanged between that commit and the placement commit, so every statement the manuscript makes about “the repository report” refers to Part I as printed above.

The manuscript’s code and data are shipped beside Part I’s files with the prefix `02-root-collisions-` (Part I’s unprefix files count as 01), and its provenance notes, delivered as `notes/provenance.md`, as `02-root-collisions-provenance.md` at the report root. Its manuscript, delivery README and PDF are not shipped; they survive in the arrival commit. Section 27 lists the shipped files.

The manuscript contributes, in the order printed here: its research question and a statement of what is classical (Section 13); the definitions and principal results, namely the exact sector-degree law, the minimal recurrence off resonance and the secondary-Hankel factorization (Section 14); the orthogonal polynomials and the confluent Christoffel identity in its normalization (Section 15); the local jets at interior points and at both endpoints (Section 16); the determinant degree calculus (Section 17); the canonical sectors and the exact leading coefficient (Section 18); minimality, the order formula and the secondary factorization (Section 19); resonance, finite-prefix certification and a root-free algorithm (Section 20); worked examples, including the recovery of Part I’s theorem (Section 21); exact asymptotics (Section 22); its verification record, implementation interface and a formalization route (Section 23); eight research directions (Section 24); a conclusion; a normalization and sign audit (Section 26); and its reproduction instructions (Section 27). Every result, proof, example, remark, limitation and question of the manuscript is printed. Where it re-derives something Part I already proves, the overlap is marked in the text (Section 12.4).

The manuscript’s own abstract reads:

For a nonzero polynomial q , let $H_N(q)$ be the size- N Hankel determinant of the Catalan moments multiplied by q . General constant-coefficient recurrences for this family are classical. We determine their sharp multiplicities. Write $q(x) = \gamma x^m (x-4)^\ell \prod_i (x-c_i)^{t_i}$, with distinct $c_i \notin \{0, 4\}$, and put $c_i = 2 + z_i + z_i^{-1}$. The branch-labelled exponential with rate $(-1)^{\ell+\sum t_i} \gamma \prod_i z_i^{2k_i-t_i}$ has a polynomial

coefficient whose degree is exactly $\binom{m}{2} + \binom{\ell+1}{2} + \sum_i k_i(t_i - k_i)$. A nonvanishing confluent-Vandermonde product gives its leading coefficient. Consequently, when the rates are distinct, the minimal recurrence order is $\prod_i (t_i + 1) \left(\binom{m}{2} + \binom{\ell+1}{2} + 1 + \frac{1}{6} \sum_i t_i(t_i - 1) \right)$. We prove an exact secondary-Hankel factorization that detects every loss of this maximal order, and give a finite, root-free algorithm determining the minimal recurrence at all rational or algebraic specializations. For one repeated nonendpoint root the generic order is cubic in the multiplicity, rather than the classical exponential bound. The example $q = (2x - 9)^4$ has order 15, while resonant fourth powers can have order 5. The proofs extend the repository's linear-multiplier result to arbitrary root-multiplicity patterns, including both endpoints, and are accompanied by exact-arithmetic verification code.

Status (manuscript 03). This is an unrefereed, AI-assisted mathematical manuscript. The universal results are proved in the text; the finite computations are an independent audit, not a substitute for those arguments. No Lean formalization or claim of first discovery is made. General polynomial-multiplier rationality, the Christoffel formula and its confluent form are classical and are not presented as new; the proposed contribution is the exact branch degree, its nonvanishing leading constant, the resulting minimality classification off resonance, and their assembled structural and algorithmic consequences. No novelty is claimed for the secondary-Hankel identity of an arbitrary exponential polynomial (Lemma 19.2), an elementary confluent-Vandermonde calculation. The literature comparison was targeted (the sources cited in Section 13) and does not establish that the theorem has never appeared in another notation; Schur-function, Toeplitz/Hankel determinant and orthogonal-polynomial literature may contain equivalent specializations, and expert review and a broader priority search would be needed before any first-discovery claim. No current global open-problem status is asserted for anything, including the eight research directions of Section 24. The theorems are over characteristic zero and are not to be transported unchanged to small positive characteristic. Nothing in Part II is formalized in any proof assistant, and its placement beside a Lean project confers no formal status. (Sources: the manuscript and its provenance notes, 02-root-collisions-provenance.md.)

12.2 Reading conventions

Inside Sections 13–27, “this article” and “the present article” mean manuscript 03 as printed in Part II, “the repository report” and “the ProveIt report” mean Part I at the pin fbba58593, and “the archive” or “the package” means the delivered package, whose code and data are shipped under the prefixed names listed in Section 27. The manuscript’s two appendices are printed as Sections 26 and 27, so that Part I’s appendices keep their letters. The manuscript cites Krattenthaler’s Ramanujan Journal paper in its preprint form arXiv:2101.04225v5 (10 September 2021), and points to its Theorem 1, Proposition 5 and Section 8, Corollary 9; that is the same paper as Part I’s reference [3]. The manuscript cites Barry’s note [1] for the denominator-exponent assertion in its introduction and does not reassert the printed coefficient conjecture without the index correction of Appendix A.

12.3 Notation of Part II

Part II uses the manuscript’s symbols, which differ from Part I’s in many places. Within Sections 13–27 every symbol has its Part II meaning, except where a sentence names Part I or cites one of its equations. Part I’s parameters a, b appear only where Part I’s multiplier $x^m(a + bx)$ is discussed: the first subsection of Section 13 and the first subsection of Section 21. Elsewhere a and b are local letters (roots, indices, block counts, a leading coefficient). Table 1 lists the symbols that a reader of Part I is likely to misread. One symbol was renamed: the manuscript’s maximal order R_0 is written $\mathcal{R}_0$, because Part I’s $R_0(y) = 1$ is the first of its polynomials (4.1). No normalization changed.

Table 1: Symbols of Part II. The last column gives Part I's meaning of the same letter, or the tempting false reading.

Symbol	Meaning in Part II	Not to be confused with
$q(x)$; subscripts in $\Delta_q, \chi_q, \Psi_q, B_q$	the nonzero multiplier polynomial (14.2)	Part I's $q(t) = 1 - (a + 2b)t + b^2 t^2$ (1.4), the quadratic denominator; Part II's generating-function variable is u , not t .
$H_N(q)$	the weighted Hankel determinant (14.1)	Part I's $H_N^{(m)} = H_N(x^m)$ and $D_N^{(m)}(a, b) = H_N(x^m(a + bx))$ are special cases; in Theorem 20.2, $h = (h_N)$ is an arbitrary sequence, not Part I's h_m .
m	multiplicity of the root 0 of q	agrees with Part I's shift m when $q = x^m(a + bx)$.
ℓ	multiplicity of the root 4 of q	Part I's column index in $M_{m,\ell}$ and Q_m .
$s; s(z), s_i$	number of distinct roots outside $\{0, 4\}$; the scalar $1/(z - z^{-1})$ of (16.2)	Part I's sign s_m in (2.5), and the source shift s of Appendix A.
c_i, z_i, t_i	roots outside $\{0, 4\}$, spectral parameters $c_i = 2 + z_i + z_i^{-1}$, multiplicities	Part I's coefficients $c_{N,m,k}$ and its variable t ; for $q = x^m(a + bx)$, $-z^{\pm 1}$ are Part I's $r_{\pm}$.
T, D	$\sum_i t_i$ and $\deg q$	Part I's $T_{n,k,s}$ (Appendix A) and its determinant $D_N^{(m)}$.
$\mathcal{K}, \mathbf{k}, \kappa$	index set (14.4), a label in it, and $\binom{m}{2} + \binom{\ell+1}{2}$	Part I's numerator degree $K = (m - 1)^2$.
λ_k	rate of the sector $\mathbf{k}$ (14.6)	for $q = x^m(a + bx)$ the two rates are Part I's $\lambda_{\pm}$ (6.3).
d_k, r_k	exact sector degree (14.7) and $d_k + 1$	Part I's $d = \binom{m}{2}$ and $e = d + 1$ (the linear case), its roots $r_{\pm}$ and the variable r of $Q_m(N; r)$.
$\mathcal{R}_0$ (renamed from R_0)	the maximal recurrence order (14.12)	Part I's polynomial $R_0(y) = 1$ (4.1); Part II's plain R is another order or size (an assumed bound in Theorem 20.2, a matrix size in Lemma 19.2, the root-free bound (20.6)).
$P_k(N), L_k$	sector polynomial in the size N (18.4), and its leading coefficient	Part I's numerator $P_m(t; a, b)$, a polynomial in t ; Part I's sector polynomials are $B_{\pm}(N)$.
$A_{\pm}(z), A_{r,\pm}(n; z)$	branch amplitudes and jet polynomials (16.4)	Part I's asymptotic constant A_m (2.10).
$B_0(m), B_4(\ell), B_z(k); B_i; B_q(u)$	local leading determinants (18.7), (17.5); blocks in Lemma 19.2; the numerator in (19.3)	Part I's $B_{\pm}(N)$ of Theorem 2.4.
$M_N; M$	the $D \times D$ Christoffel matrix (15.7); $\max_{\mathbf{k}} d_{\mathbf{k}}$ in Section 22	Part I's minors $M_{m,\ell}(N)$ (4.6).
E	forward shift, $(Ef)(N) = f(N + 1)$	Part I's shift E in Lemma 6.1 and its endpoint polynomial $E_m(N)$ (6.10).
Λ, Q_{λ}	set of distinct rates; grouped sector polynomial (20.1)	Part I's minor generating polynomial $Q_m(N; r)$ (5.7).
$(a_u, v_u)_{u \leq b}; (w_a, u_a)_{a \leq b}; b_t$	the distinct roots of q , their multiplicities and number b (Section 15); nodes and block sizes in Lemma 17.1; degree of the square-free factor f_t	Part I's parameters a, b , which Part II uses only in the first subsections of Sections 13 and 21.
p_n, μ	monic orthogonal polynomials (15.1) and the Catalan measure	they are Part I's $p_n^{(0)}$ and μ_0 ; Part I's $p_N^{(m)}$ and μ_m with $m \geq 1$ are different.

Symbol	Meaning in Part II	Not to be confused with
$\Delta_q, V, \mathcal{V}$	cross-root product (15.6); Vandermonde (matrix); confluent product (18.6)	Part I's Vandermonde $\Delta(x)$.
C (proof of Theorem 18.1)	a matrix of coefficient polynomials	the Catalan numbers C_n .

12.4 Where the merge had to choose

1. *Placement.* Part II is printed after Part I's Section 11 and before Part I's appendices, so that no section, equation, theorem or table number of Part I moves (checked against the `.aux` file of a build of the committed text). Part II's sections continue the numbering (Sections 12 to 27); the manuscript's Sections 1–13 are Sections 13–25, and its Appendices A and B are Sections 26 and 27.
2. *Labels.* Every label of Part II carries the prefix `rc::`; without it, the manuscript's `sec:main`, `thm:main`, `eq:R`, `eq:lambda`, `eq:measure`, `lem:Christoffel`, `eq:Christoffel` and several others would have collided with Part I's.
3. *Overlaps with Part I.* The manuscript re-derives, without assuming Part I correct, several things Part I already proves. Printed with a pointer and kept: the orthogonality of the p_n and $\det(C_{i+j}) = 1$ (a second route; Section 15); the Christoffel identity (Part I's Lemma 4.1 is its special case $\gamma = 1$, with Part I's normalization R_n , and Proposition 4.2 its confluent case at the root 0); the binomial determinant (17.3) (as in (5.3)); the minimal-annihilator lemma (Lemma 19.1 restates Lemma 6.1; the manuscript's proof is kept as a second proof); and the recovery of Part I's sharp order and exceptional orders (a second proof by a different route, Section 21). No statement of Part I was changed.
4. *Bibliography.* The manuscript's references to Barry and to Krattenthaler's Ramanujan Journal paper are Part I's [1, 3]; its versions and pointers are recorded in Section 12.2. Its references to Cigler and Krattenthaler and to Krattenthaler's Uvarov-formula preprint are added to the common bibliography. Its reference to the pinned repository report is replaced by Part I itself and the pin recorded in Section 12.1.
5. *File names.* Sections 23 and 27 give the shipped names beside the delivered ones. The manuscript's reproduction commands are printed as delivered but must not be run in place here; Section 27 gives a recipe on a scratch copy.
6. *Front matter.* The manuscript's title page, abstract, status paragraph and archive paragraph are replaced by Section 12.1, which prints the abstract and the status. The archive paragraph is not reprinted because the delivered source and PDF are not shipped; the shipped code and data are listed in Section 27.
7. *Notation.* The manuscript's symbols are kept, with the single rename $R_0 \rightarrow \mathcal{R}_0$ and the scoping convention and Table 1 of Section 12.3.

13 The research question and its provenance

13.1 A concrete extension of the ProveIt report

The starting point is the report *Shifted Catalan Hankel Polynomials: Exact coefficients, sharp recurrences, and a denominator conjecture*, dated 19 September 2026, in ProveIt, printed above as Part I (§12.1 records the pinned snapshot). For

$$D_N^{(m)}(a, b) = \det_{0 \leq i, j < N} (aC_{i+j+m} + bC_{i+j+m+1}), \quad C_n = \frac{1}{n+1} \binom{2n}{n},$$

it proves the sharp characteristic polynomial

$$(X^2 - (a + 2b)X + b^2)^{\binom{m}{2}+1} \quad (13.1)$$

when $b \neq 0$ and $a(a + 4b) \neq 0$ (this is (2.6) of Theorem 2.2). Its concluding discussion (Section 11) explicitly leaves multiple linear factors and their collision patterns for further work. That is the extension addressed here.

The repository was inspected at commit

fbba58593dc0622aa914972896150d4848f935b5.

The relevant source is under

SetTheory/Cardinals/docs/reports/hankel-determinants/
catalan-and-ballot/shifted-catalan-hankel-polynomials/article.tex.

The present proof does not assume that an unrefereed repository manuscript is correct. We rederive the determinant identity and the local degree calculus, then recover (13.1) as a specialization.

The question is not merely whether another rational generating function exists. It is this:

Given every root and its multiplicity in a polynomial multiplier, which exponential terms actually occur, what is the exact polynomial degree of each term, and precisely when does the minimal recurrence order drop?

A repeated root is not equivalent to several independently chosen roots. The associated determinant forces cancellations within a repeated block. Those cancellations are the main subject of this article.

13.2 What is classical, and what is being added

The Christoffel determinant formula and its confluent form are classical; Krattenthaler's treatment [3] gives several proofs and explicitly handles coalescing evaluation points. Its Section 8, Corollary 9, supplies a constant-coefficient recurrence of order at most 2^D for a multiplier of degree D , including Catalan moments. That account also credits Elouafi's earlier proof of the Catalan recurrence assertion. General rationality is therefore not a new conclusion of this article.

Barry's note [1] motivates the shifted linear combination and its quadratic-denominator exponent. The earlier work of Cigler and Krattenthaler [7] gives further context for polynomial linear combinations of moments. Our comparison is with these particular sources and the pinned repository report, not with an exhaustive classification of the literature.

The additions proved below are an exact degree law for every repeated-root block, its nonzero leading constant with endpoint factors included, the resulting sharp recurrence and a factorization detecting its exceptional locus. A root-free finite procedure handles resonances without assuming that a generic formula remains minimal after specialization. The underlying determinant techniques are familiar; the full multiplicity law and its assembled consequences are the proposed research contribution. An equivalent earlier formulation in the wider orthogonal-polynomial or symmetric-function literature has not been ruled out.

14 Definitions and principal results

14.1 The weighted determinant and its spectral parameters

For $q(x) = \sum_{r=0}^D q_r x^r \in \mathbb{C}[x] \setminus \{0\}$, define

$$H_N(q) = \det_{0 \leq i, j < N} \left(\sum_{r=0}^D q_r C_{i+j+r} \right), \quad N \in \mathbb{N}_0, \quad H_0(q) = 1. \quad (14.1)$$

Write its factorization as

$$q(x) = \gamma x^m (x - 4)^\ell \prod_{i=1}^s (x - c_i)^{t_i}, \quad (14.2)$$

where $\gamma \neq 0$, $m, \ell \in \mathbb{N}_0$, $t_i \geq 1$, and the c_i are pairwise distinct and avoid 0, 4. Thus

$$T = \sum_{i=1}^s t_i, \quad D = m + \ell + T.$$

An empty product is 1. Constant multipliers are included by taking $m = \ell = s = 0$. The identically zero multiplier is excluded; its sequence would simply be 1, 0, 0, ...

For each i choose a solution of

$$c_i = 2 + z_i + z_i^{-1}. \quad (14.3)$$

Then $z_i \in \mathbb{C}^\times \setminus \{1, -1\}$. Replacing z_i by z_i^{-1} only relabels an index below. No final recurrence depends on this choice. Define

$$\mathcal{K} = \prod_{i=1}^s \{0, 1, \dots, t_i\}, \quad (14.4)$$

$$\kappa = \binom{m}{2} + \binom{\ell+1}{2}, \quad (14.5)$$

$$\lambda_{\mathbf{k}} = (-1)^{\ell+T} \gamma \prod_{i=1}^s z_i^{2k_i - t_i}, \quad (14.6)$$

$$d_{\mathbf{k}} = \kappa + \sum_{i=1}^s k_i(t_i - k_i), \quad r_{\mathbf{k}} = d_{\mathbf{k}} + 1. \quad (14.7)$$

When $s = 0$, the index set $\mathcal{K}$ has one element, the empty tuple.

Definition 14.1 (Resonance). The parameters are *nonresonant* if the values $\lambda_{\mathbf{k}}$, $\mathbf{k} \in \mathcal{K}$, are pairwise distinct. Otherwise there is a *resonance*. Equivalently, a resonance is an identity

$$\prod_i z_i^{2(k_i - j_i)} = 1 \quad \text{for some distinct } \mathbf{k}, \mathbf{j} \in \mathcal{K}. \quad (14.8)$$

This is different from a repeated root of q : repeated roots have already been recorded by the t_i .

14.2 The exact multiplicity law

Theorem 14.2 (Branch degrees and minimal recurrence). *For every multiplier (14.2), there are canonically branch-labelled polynomials $P_{\mathbf{k}} \in \mathbb{C}[N]$ such that*

$$H_N(q) = \sum_{\mathbf{k} \in \mathcal{K}} P_{\mathbf{k}}(N) \lambda_{\mathbf{k}}^N \quad (N \geq 0), \quad (14.9)$$

and

$$\deg P_{\mathbf{k}} = d_{\mathbf{k}} \quad \text{for every } \mathbf{k}. \quad (14.10)$$

Their leading coefficients are given by the nonzero product in Theorem 18.1. These individual assertions remain valid at resonances, when different branch labels can have the same numerical rate.

If the parameters are nonresonant, the minimal characteristic polynomial of the sequence is

$$\chi_q(X) = \prod_{\mathbf{k} \in \mathcal{K}} (X - \lambda_{\mathbf{k}})^{d_{\mathbf{k}}+1}. \quad (14.11)$$

Its degree is

$$\mathcal{R}_0 = \prod_{i=1}^s (t_i + 1) \left(\kappa + 1 + \frac{1}{6} \sum_{i=1}^s t_i(t_i - 1) \right). \quad (14.12)$$

For arbitrary parameters, (14.11) is still an annihilating polynomial of degree $\mathcal{R}_0$. The true minimal order equals $\mathcal{R}_0$ if and only if the parameters are nonresonant.

A characteristic polynomial $g(X) = \sum_{j=0}^r g_j X^j$, with $g_r = 1$, means the recurrence

$$\sum_{j=0}^r g_j H_{N+j}(q) = 0 \quad \text{for every } N \geq 0. \quad (14.13)$$

Thus the assertions concern the entire sequence, including its initial values, not only an eventual recurrence. A polynomial of minimal degree with this property is unique.

Corollary 14.3 (One repeated nonendpoint root). *Let $q(x) = \gamma(x-c)^t$, where $c \notin \{0, 4\}$ and $c = 2+z+z^{-1}$. If $z^{2h} \neq 1$ for $1 \leq h \leq t$, then*

$$\text{ord}(H(q)) = t + 1 + \binom{t+1}{3}. \quad (14.14)$$

The exponential with rate $(-1)^t \gamma z^{2k-t}$ has polynomial degree $k(t-k)$. In particular the first orders, for $t = 1, \dots, 8$, are

$$2, 4, 8, 15, 26, 42, 64, 93.$$

The order in (14.14) is asymptotic to $t^3/6$, in contrast to the universal bound 2^t . The first strict improvement occurs at $t = 4$. This is a uniform family of exact minimality results, rather than an isolated fitted recurrence.

Corollary 14.4 (Simple factors and endpoints). *If every $t_i = 1$ and the parameters are nonresonant, the minimal order is*

$$2^s \left(\binom{m}{2} + \binom{\ell+1}{2} + 1 \right). \quad (14.15)$$

If $s = 0$, no nonresonance hypothesis is needed: $H_N(q)$ is $((-1)^\ell \gamma)^N$ times a polynomial of degree exactly κ . The minimal polynomial is $(X - (-1)^\ell \gamma)^{\kappa+1}$.

The asymmetry between 0 and 4 is essential. A multiplicity m at 0 contributes $\binom{m}{2}$ degrees, whereas a multiplicity ℓ at 4 contributes $\binom{\ell+1}{2}$. Treating both endpoints as a generic collision of two equally weighted branches gives the wrong answer.

14.3 An exact detector for exceptional parameters

Put $L_k = \text{lc}(P_k)$, and choose any ordering of $\mathcal{K}$.

Theorem 14.5 (Secondary Hankel factorization). *With $\mathcal{R}_0 = \sum_k r_k$, one has*

$$\begin{aligned} & \det_{0 \leq u, v < \mathcal{R}_0} (H_{u+v}(q)) \\ &= (-1)^{\sum_k \binom{r_k}{2}} \prod_k \left((d_k! L_k)^{r_k} \lambda_k^{r_k(r_k-1)} \right) \prod_{k < j} (\lambda_j - \lambda_k)^{2r_k r_j}. \end{aligned} \quad (14.16)$$

This identity holds at resonant as well as nonresonant parameters. All factors outside the final product are nonzero. Consequently this secondary determinant vanishes exactly on the resonance locus.

The formula is stronger than a statement that a generic matrix has full rank. It identifies every source of rank loss and its multiplicity in spectral coordinates. Sections 19 and 20 prove this assertion and make it algorithmic.

15 Catalan orthogonality and the fixed-size determinant

This section proves the classical infrastructure with our normalization. It is included to make the signs, initial index and confluent factorials fully auditable. The general identities are not claimed as new.

15.1 The orthogonal polynomials

Define monic polynomials by

$$p_0(x) = 1, \quad p_1(x) = x - 1, \quad p_{n+1}(x) = (x - 2)p_n(x) - p_{n-1}(x) \quad (n \geq 1). \quad (15.1)$$

The Catalan probability measure is

$$d\mu(x) = \frac{1}{2\pi} \sqrt{\frac{4-x}{x}} \mathbf{1}_{(0,4)}(x) dx. \quad (15.2)$$

Substitution $x = 4u$ and the beta integral give

$$\int x^n d\mu(x) = \frac{2 \cdot 4^n}{\pi} B\left(n + \frac{1}{2}, \frac{3}{2}\right) = \frac{(2n)!}{n!(n+1)!} = C_n.$$

There is also an elementary trigonometric verification of the orthogonality. For $x = 2 + 2 \cos \theta$, $0 < \theta < \pi$, the recurrence gives

$$p_n(x) = \frac{\sin((n + \frac{1}{2})\theta)}{\sin(\theta/2)}, \quad d\mu = \frac{1 - \cos \theta}{\pi} d\theta. \quad (15.3)$$

The product-to-sum formula then yields

$$\int p_n(x) p_j(x) d\mu(x) = \delta_{n,j}. \quad (15.4)$$

Indeed, after cancellation of $\sin^2(\theta/2)$ the integral is $\frac{2}{\pi} \int_0^\pi \sin((n + \frac{1}{2})\theta) \sin((j + \frac{1}{2})\theta) d\theta$. The change from monomials to monic p_n is triangular with determinant 1. Hence (15.4) also proves

$$\det_{0 \leq i, j < N} (C_{i+j}) = 1. \quad (15.5)$$

Relation to Part I. The measure μ is Part I's μ_0 of (3.1), and p_n is Part I's $p_n^{(0)}$, with the same recurrence (3.8). Part I proves (15.5), $H_N^{(0)} = 1$, from the constant squared norms of the $p_n^{(0)}$ (Section 3); the trigonometric orthogonality above is a second route, kept as the manuscript printed it.

15.2 Christoffel's determinant, including confluence

List the distinct roots in (14.2) in the order $0, 4, c_1, \dots, c_s$, omitting endpoints whose multiplicities are zero. Write this list as $(a_1, \nu_1), \dots, (a_b, \nu_b)$ and set

$$\Delta_q = \prod_{1 \leq u < v \leq b} (a_v - a_u)^{\nu_u \nu_v}. \quad (15.6)$$

It is nonzero. Let M_N have rows indexed by (a_u, r) , $0 \leq r < \nu_u$, and columns indexed by $j = 0, \dots, D-1$, with entries

$$(M_N)_{(a_u, r), j} = \frac{p_{N+j}^{(r)}(a_u)}{r!}. \quad (15.7)$$

Lemma 15.1 (Confluent Christoffel identity). *For all $N \geq 0$,*

$$H_N(q) = (-1)^{ND} \gamma^N \frac{\det M_N}{\Delta_q}. \quad (15.8)$$

Proof. First suppose all roots $a_1, \dots, a_D$ are distinct. For an ordered list y , write $V(y) = \prod_{i < j} (y_j - y_i)$. Expanding two determinants under the integral and using the product measure proves the elementary integration identity

$$\int \det(f_i(x_j))_{i,j=1}^N \det(g_i(x_j))_{i,j=1}^N \prod_j d\mu(x_j) = N! \det \left(\int f_i(x) g_j(x) d\mu(x) \right)_{i,j=1}^N.$$

Taking monomials with the multiplier included gives

$$H_N(q) = \frac{1}{N!} \int V(x_1, \dots, x_N)^2 \prod_{i=1}^N q(x_i) d\mu(x_i).$$

Since

$$V(x, a) = V(x) V(a) \prod_{i,u} (a_u - x_i),$$

we obtain

$$H_N(q) = \frac{(-1)^{ND} \gamma^N}{N! V(a)} \int V(x) V(x, a) \prod_i d\mu(x_i). \quad (15.9)$$

Replace the monomials in both Vandermonde determinants by the monic polynomials p_j . Expand $V(x, a)$ along its first N rows. On integration against $V(x)$, orthogonality kills every selected set of columns except $0, 1, \dots, N-1$. That surviving integral is $N!$, and its complementary minor is $(p_{N+j}(a_u))_{u,j}$. The Laplace sign is positive because the selected columns are the first N . This proves (15.8) for distinct roots.

Now let ν distinct roots tend to a common value a . Taylor expansion shows that division by their internal Vandermonde replaces their successive evaluation rows by the rows $p_{N+j}^{(r)}(a)/r!$, $r = 0, \dots, \nu-1$. Cross-block differences tend to the product (15.6). Applying this separately to every block gives the stated formula. Equivalently, both sides are polynomial identities in the root variables after the removable Vandermonde factors are canceled. Thus the limiting argument does not assume that any weighted Hankel determinant is nonzero. At $N = 0$, both the empty moment determinant and the normalized confluent Vandermonde equal 1. $\square$

The important computational consequence is that the determinant on the right has size D , independent of N . The important theoretical consequence is different: all dependence on N has been localized in jets of a two-branch orthogonal-polynomial sequence.

Relation to Part I. Lemma 4.1 of Part I is the case $\gamma = 1$, roots $a_u = -y_u$, in the reversed normalization $R_n(y) = (-1)^n p_n(-y)$ of (4.1); its proof follows the same integration-and-Laplace route for distinct roots. Part I then takes the confluent limit only for the m -fold root at 0 (Proposition 4.2). Lemma 15.1 adds confluence at every root, both endpoints, and the leading coefficient γ with its sign, so it is printed in full.

16 Local jet structure

16.1 The two branches away from the endpoints

The characteristic equation of (15.1), at $x = 2 + z + z^{-1}$, has roots z, z^{-1} . Matching the first two values gives

$$p_n(2 + z + z^{-1}) = \frac{z^{n+1} - z^{-n}}{z - 1} = A_+(z)z^n + A_-(z)z^{-n}, \quad (16.1)$$

where

$$A_+(z) = \frac{z}{z-1}, \quad A_-(z) = -\frac{1}{z-1}, \quad s(z) = \frac{1}{z-z^{-1}}. \quad (16.2)$$

The letter $s(z)$ is a scalar function, not the number s of distinct nonendpoint roots. For clarity we often write $s_i = s(z_i)$. The inverse function theorem applies locally because $z \neq \pm 1$, and

$$\frac{d}{dx} \log z(x) = s(z), \quad \frac{d}{dx} = z s(z) \frac{d}{dz}. \quad (16.3)$$

Lemma 16.1 (Offendpoint jets). *For $c = 2 + z + z^{-1} \notin \{0, 4\}$ and $r \geq 0$,*

$$\frac{p_n^{(r)}(c)}{r!} = z^n A_{r,+}(n; z) + z^{-n} A_{r,-}(n; z), \quad (16.4)$$

where $A_{r,\pm}(n; z)$ are polynomials of degree exactly r in n . Their leading coefficients are

$$[n^r] A_{r,\pm}(n; z) = \frac{A_{\pm}(z)(\pm s(z))^r}{r!}. \quad (16.5)$$

They can be constructed recursively from $A_{0,\pm} = A_{\pm}$ by

$$A_{r+1,\pm}(n; z) = \frac{zs(z) \partial_z A_{r,\pm}(n; z) \pm ns(z) A_{r,\pm}(n; z)}{r+1}. \quad (16.6)$$

Proof. Differentiate (16.4) once and use (16.3). The normalization by $(r+1)!$ gives (16.6). The derivative term does not increase the degree in n , while multiplication by $\pm ns(z)$ increases it by one. Induction gives (16.5). All its factors are nonzero, so the degree is exact. $\square$

16.2 The two endpoint jets

From (15.1),

$$\sum_{n \geq 0} p_n(x) u^n = \frac{1+u}{1-(x-2)u+u^2}. \quad (16.7)$$

Differentiation in x , normalized by $r!$, gives

$$\sum_{n \geq 0} \frac{p_n^{(r)}(x)}{r!} u^n = \frac{(1+u)u^r}{(1-(x-2)u+u^2)^{r+1}}. \quad (16.8)$$

Lemma 16.2 (Endpoint degrees). *For all $n, r \geq 0$,*

$$\frac{p_n^{(r)}(0)}{r!} = (-1)^{n-r} \binom{n+r}{2r}, \quad (16.9)$$

$$\frac{p_n^{(r)}(4)}{r!} = \frac{2n+1}{2r+1} \binom{n+r}{2r}. \quad (16.10)$$

After the common factor $(-1)^n$ is removed at 0, the polynomial in n has degree $2r$ and leading coefficient $(-1)^r/(2r)!$. At 4 the degree is $2r+1$, with leading coefficient $2/(2r+1)!$.

Proof. At $x = 0$, (16.8) becomes $u^r/(1+u)^{2r+1}$, which gives (16.9). At $x = 4$ it becomes $(1+u)u^r/(1-u)^{2r+2}$. The coefficient is

$$\binom{n+r+1}{2r+1} + \binom{n+r}{2r+1} = \frac{2n+1}{2r+1} \binom{n+r}{2r}.$$

The assertions on leading coefficients follow immediately. For $n < r$ the binomial coefficient is zero, as it must be for a derivative whose order exceeds the polynomial degree. $\square$

The same quadratic recurrence has two different endpoint amplitudes: $p_n(0) = (-1)^n$ but $p_n(4) = 2n+1$. This single extra degree at 4 is what ultimately changes $\binom{\ell}{2}$ into $\binom{\ell+1}{2}$.

17 The determinant degree calculus

17.1 Polynomial shifts and confluent exponential rows

The next two identities isolate the cancellation responsible for the sharp order. If f is a polynomial of degree d with leading coefficient a , then

$$[j^h]f(N+j) \text{ has degree at most } d-h \text{ in } N, \quad [N^{d-h}j^h]f(N+j) = a \binom{d}{h}. \quad (17.1)$$

An expression with $h > d$ is interpreted as zero.

For distinct nonzero numbers $w_1, \dots, w_b$ and positive integers $u_1, \dots, u_b$ summing to D , form the matrix

$$V_{(a,h),j} = w_a^j j^h, \quad 0 \leq h < u_a, \quad 0 \leq j < D.$$

The convention $0^0 = 1$ is used only in the polynomial value j^0 .

Lemma 17.1 (Confluent exponential Vandermonde). *In the indicated block order,*

$$\det V = \prod_{a=1}^b \left(w_a^{\binom{u_a}{2}} \prod_{h=0}^{u_a-1} h! \right) \prod_{a < b} (w_b - w_a)^{u_a u_b}. \quad (17.2)$$

Proof. Start with the ordinary Vandermonde determinant and coalesce u_a variables to each w_a . The matrix of ordinary, unnormalized derivatives of w^j then has determinant $\prod_a \prod_{h < u_a} h! \prod_{a < b} (w_b - w_a)^{u_a u_b}$. Within a block replace ∂_w^h by $(w \partial_w)^h$. This is a triangular change of rows: the coefficient of ∂_w^h is w^h , and all other terms have smaller derivative order. The determinant is multiplied by $\prod_a w_a^{0+1+\dots+(u_a-1)}$. Since $(w \partial_w)^h w^j = j^h w^j$, the resulting matrix is V . $\square$

17.2 Three local leading determinants

For an endpoint block of size u , the leading matrices obtained from (17.1) have entries proportional to $\binom{2r}{h}$ or $\binom{2r+1}{h}$, $0 \leq r, h < u$. As a polynomial in its upper argument, $\binom{x}{h}$ has leading coefficient $1/h!$. Consequently

$$\det_{0 \leq r, h < u} \binom{2r}{h} = \det_{0 \leq r, h < u} \binom{2r+1}{h} = 2^{\binom{u}{2}}. \quad (17.3)$$

Here the Vandermonde in $0, 2, \dots, 2u-2$ or $1, 3, \dots, 2u-1$ contributes $2^{\binom{u}{2}} \prod_{h < u} h!$, canceling the column leading denominators. (The first equality with $u = m$ is the determinant evaluated in (5.3) of Part I, by the same argument.)

For the nonendpoint calculation let $t = k + h$, and write $A_+ = A_+(z)$, $A_- = A_-(z)$ and $s = s(z)$. Consider the $t \times t$ matrix with rows $r = 0, \dots, t-1$, followed first by k plus columns and then by h minus columns:

$$B_{r,(+,j)} = \frac{A_+ s^r}{r!} \binom{r}{j} \quad (0 \leq j < k), \quad B_{r,(-,j)} = \frac{A_- (-s)^r}{r!} \binom{r}{j} \quad (0 \leq j < h). \quad (17.4)$$

Lemma 17.2 (The mixed-block determinant). *The determinant in (17.4) equals*

$$B_z(k) = \frac{A_+^k A_-^h s^{\binom{k}{2}} (-s)^{\binom{h}{2}} (-2s)^{kh}}{\prod_{r=0}^{t-1} r!}, \quad h = t - k, \quad (17.5)$$

and is nonzero.

Proof. Remove the row factors $1/r!$ and the column factors $A_{\pm}$. Use

$$w^r \binom{r}{j} = w^j \frac{1}{j!} \left(\frac{d}{dw} \right)^j w^r.$$

The remaining normalized-derivative matrix is the transpose of a confluent Vandermonde at $w = s$ and $w = -s$. Its determinant is $(-2s)^{kh}$. The column powers supply $s^{\binom{k}{2}}(-s)^{\binom{h}{2}}$, proving (17.5). Every factor is nonzero because $z \notin \{0, 1, -1\}$. $\square$

This lemma is the point at which a mere degree bound becomes an exact degree theorem. It sums all competing assignments of the derivative rows to the two branches, including their determinant signs. Ignoring those signs, or arguing that one assignment has a nonzero leading term, would not establish noncancellation of the whole sector.

18 Canonical sectors and the proof of sharpness

18.1 A definition that still makes sense at resonance

Set

$$f_{0,r}(n) = (-1)^r \binom{n+r}{2r}, \quad f_{4,r}(n) = \frac{2n+1}{2r+1} \binom{n+r}{2r}.$$

Introduce independent marker variables $y_1, \dots, y_s$. Let $\tilde{M}(N; \mathbf{y})$ be the $D \times D$ matrix whose column j contains the following rows, in the same root order as before:

$$(-1)^j f_{0,r}(N+j), \quad 0 \leq r < m, \quad (18.1)$$

$$f_{4,r}(N+j), \quad 0 \leq r < \ell, \quad (18.2)$$

$$y_i z_i^j A_{r,+}(N+j; z_i) + z_i^{-j} A_{r,-}(N+j; z_i), \quad 0 \leq r < t_i. \quad (18.3)$$

Its determinant is a polynomial in N and the markers. Define

$$P_{\mathbf{k}}(N) = \frac{1}{\Delta_q} [y_1^{k_1} \cdots y_s^{k_s}] \det \tilde{M}(N; \mathbf{y}). \quad (18.4)$$

This definition does not require distinct numerical rates. It is the meaning of “canonically branch-labelled” in Theorem 14.2.

In the original matrix M_N , remove $(-1)^N$ from each of its m zero-endpoint rows and z_i^{-N} from each of the t_i rows at c_i . The remaining matrix is (18.1)–(18.3) with $y_i = z_i^{2N}$. Therefore (15.8) gives

$$H_N(q) = \left((-1)^{D+m} \gamma \prod_i z_i^{-t_i} \right)^N \sum_{\mathbf{k}} P_{\mathbf{k}}(N) \prod_i z_i^{2N k_i}.$$

Since $D+m$ has the parity of $\ell + T$, this proves (14.9), with the rates in (14.6).

18.2 The leading coefficient in product form

For each label $\mathbf{k}$, form an ordered list of spectral blocks

$$\mathcal{W}_{\mathbf{k}} = ((-1, m), (1, \ell), (z_1, k_1), (z_1^{-1}, t_1 - k_1), \dots, (z_s, k_s), (z_s^{-1}, t_s - k_s)), \quad (18.5)$$

omitting all blocks of size zero. Its nodes are pairwise distinct: an equality between a node from c_i and a node from c_j would imply $c_i = c_j$, and none is an endpoint node ± 1 . For any such ordered list write

$$\mathcal{V}((w_a, u_a)_{a=1}^b) = \prod_a \left(w_a^{\binom{u_a}{2}} \prod_{h=0}^{u_a-1} h! \right) \prod_{a < b} (w_b - w_a)^{u_a u_b}. \quad (18.6)$$

Finally define

$$B_0(m) = \frac{(-2)^{\binom{m}{2}}}{\prod_{r=0}^{m-1} (2r)!}, \quad B_4(\ell) = \frac{2^{\binom{\ell+1}{2}}}{\prod_{r=0}^{\ell-1} (2r+1)!}. \quad (18.7)$$

Both equal 1 at multiplicity zero.

Theorem 18.1 (Exact leading coefficient). *For P_k defined by (18.4),*

$$[N^{d_k}]P_k(N) = \frac{B_0(m)B_4(\ell) \prod_{i=1}^s B_{z_i}(k_i)}{\Delta_q} \mathcal{V}(\mathcal{W}_k) \neq 0. \quad (18.8)$$

Here $B_{z_i}(k_i)$ is (17.5) with $t = t_i$. Moreover P_k has no term of degree greater than d_k .

Proof. Expand each row of $\tilde{M}$ in the finite collection of basis rows

$$(w^j j^h)_{j=0}^{D-1}. \quad (18.9)$$

Only the finitely many h up to the largest relevant polynomial degree are needed. This factors $\tilde{M} = C(N; \mathbf{y})V$, with C a matrix of coefficient polynomials and V the matrix of basis rows. Cauchy–Binet expresses its determinant as a sum over selections of D basis rows.

A nonzero coefficient minor must select exactly m rows associated with node -1 , exactly ℓ with node 1 , and exactly t_i rows from the two nodes z_i, z_i^{-1} for each i . Otherwise a root block of C , whose support is confined to its own nodes, cannot have full rank. The coefficient of $\mathbf{y}^k$ selects precisely k_i rows at z_i and $t_i - k_i$ rows at z_i^{-1} .

For a fixed node with u selected basis rows, their h indices must be distinct. Their sum is therefore at least $0 + 1 + \dots + (u-1) = \binom{u}{2}$, with equality only for the set $\{0, \dots, u-1\}$. By (17.1), selecting such an index h subtracts h from the maximum possible degree in N of its coefficient. Thus the zero-endpoint block contributes at most

$$\sum_{r=0}^{m-1} 2r - \binom{m}{2} = \binom{m}{2},$$

the four-endpoint block contributes at most

$$\sum_{r=0}^{\ell-1} (2r+1) - \binom{\ell}{2} = \binom{\ell+1}{2},$$

and the i th nonendpoint block contributes at most

$$\sum_{r=0}^{t_i-1} r - \binom{k_i}{2} - \binom{t_i - k_i}{2} = k_i(t_i - k_i).$$

The sum of these bounds is d_k .

There is exactly one selection of basis rows that can attain this total degree: at every node take $h = 0, \dots, u-1$. The associated basis determinant is (18.6) for the list $\mathcal{W}_k$, by Lemma 17.1. In its coefficient minor, the root blocks separate. For the endpoint 0, the leading row factors from Lemma 16.2, followed by (17.3), give $B_0(m)$. At 4 they give $B_4(\ell)$. At c_i the leading coefficient matrix is exactly (17.4), so Lemma 17.2 gives $B_{z_i}(k_i)$. Dividing by Δ_q proves (18.8).

Every factor in (18.8) is nonzero. In particular, the unique maximal-degree basis selection cannot vanish or be canceled by another selection, because every other selection has strictly smaller degree. This proves both exactness and noncancellation. $\square$

Remark 18.2 (Why the naive degree is too large). For a block of multiplicity t , multiplying row degrees without taking the determinant into account suggests $\binom{t}{2}$ degrees. The k plus-branch rows consume $\binom{k}{2}$ degrees, and the $t - k$ minus rows consume $\binom{t-k}{2}$. The surviving degree is their difference, $k(t - k)$. The consumption is forced by linear dependence of equal low-degree basis rows; it is not an accidental cancellation at special numerical parameters.

19 Minimal recurrences, order, and the secondary factorization

19.1 Exponential polynomials and minimality

Let $(Ef)(N) = f(N + 1)$ denote forward shift.

Lemma 19.1 (Minimal annihilators). *Suppose $\lambda_1, \dots, \lambda_b$ are distinct nonzero complex numbers and Q_i are nonzero polynomials of degrees d_i . Then $f(N) = \sum_i Q_i(N)\lambda_i^N$ has minimal characteristic polynomial*

$$\prod_{i=1}^b (X - \lambda_i)^{d_i+1}. \quad (19.1)$$

Its ordinary generating function has reduced denominator $\prod_i (1 - \lambda_i u)^{d_i+1}$.

This is Lemma 6.1 of Part I, restated in Part II's notation (Part I writes E for the shift and t for the generating-function variable). The manuscript proves it again; its proof isolates each block by annihilating the others instead of using Chinese-remainder projectors, and is kept as a second route.

Second proof (manuscript 03). On a term $Q(N)\lambda^N$, the operator $E - \lambda$ gives $\lambda^{N+1}(Q(N+1) - Q(N))$. It lowers the degree by exactly one until the polynomial is zero. In contrast, $E - \mu$, for $\mu \neq \lambda$, preserves the degree and multiplies the leading coefficient by $\lambda - \mu$.

These observations also prove independence of distinct exponential blocks. To isolate one block in a purported zero sum, annihilate all other blocks by their respective powers of $E - \lambda_i$. The selected block remains nonzero unless it was zero initially. If a polynomial g has a zero of multiplicity e at λ_i , then its action on that block is nonzero for $e \leq d_i$ and zero for $e > d_i$. Hence each factor in (19.1) is necessary and their product suffices.

For the generating function, apply powers of $u d/du$ to $(1 - \lambda_i u)^{-1}$. A degree- d_i polynomial multiplier yields a pole of order exactly $d_i + 1$ at $u = \lambda_i^{-1}$. Distinct poles cannot cancel. This gives the stated reduced denominator. $\square$

Applying the lemma to Theorem 18.1 proves (14.11) under nonresonance. Without that hypothesis its labelled product still annihilates every term. Summing its exponents uses

$$\sum_{k=0}^t k(t-k) = \frac{t(t+1)(t-1)}{6}, \quad (19.2)$$

and gives (14.12). If two or more labels have the same rate, their sum has degree at most their largest degree, strictly less in order contribution than the sum of their separate $d + 1$ values. Thus any resonance strictly lowers the order below $\mathcal{R}_0$. This completes the proof of Theorem 14.2 and its corollaries.

In the nonresonant case the ordinary generating function is therefore

$$\sum_{N \geq 0} H_N(q)u^N = \frac{B_q(u)}{\prod_k (1 - \lambda_k u)^{d_k+1}}, \quad \deg B_q < \mathcal{R}_0, \quad B_q(0) = 1, \quad (19.3)$$

and the displayed fraction is reduced. A limiting specialization of this numerator and denominator need not remain reduced.

19.2 Comparison with the classical exponential bound

There is a useful elementary consistency check on the sharp formula. For $0 \leq k \leq t$,

$$1 + k(t-k) \leq \binom{t}{k}. \quad (19.4)$$

To see this, identify a k -subset of $\{1, \dots, t\}$ with a partition inside a $k \times (t - k)$ rectangle. Every size from 0 to $k(t - k)$ is attained: fill complete rows and then at most one partial row. There are therefore at least $k(t - k) + 1$ different partitions. Also

$$1 + \binom{m}{2} \leq 2^m, \quad 1 + \binom{\ell + 1}{2} = 1 + \ell + \binom{\ell}{2} \leq 2^\ell.$$

For nonnegative numbers a_j , $1 + \sum_j a_j \leq \prod_j (1 + a_j)$. Combining these inequalities, label by label and then summing, proves

$$\mathcal{R}_0 \leq 2^{m+\ell} \prod_i \sum_{k_i=0}^{t_i} \binom{t_i}{k_i} = 2^D. \quad (19.5)$$

Thus the order bound here is compatible with the classical one, while retaining the determinant cancellations that the larger bound ignores. For a single repeated root, summing $1 + k(t - k)$ gives $t + 1 + \binom{t+1}{3}$. Strictness for $t \geq 4$ already follows from $\binom{t}{2} > 1 + 2(t - 2)$ at $k = 2$.

19.3 Proof of the secondary Hankel factorization

We prove a general identity, then substitute the explicit leading coefficients of our determinant sequence.

Lemma 19.2 (A finite Hankel determinant of exponential polynomials). *Suppose $f(N) = \sum_{i=1}^b Q_i(N) \lambda_i^N$, with $\deg Q_i = d_i$, leading coefficient a_i , and $r_i = d_i + 1$. Put $R = \sum_i r_i$. The rates may coincide. Then*

$$\det_{0 \leq u, v < R} f(u + v) = (-1)^{\sum_i \binom{r_i}{2}} \prod_i (d_i! a_i)^{r_i} \lambda_i^{r_i(r_i-1)} \prod_{i < j} (\lambda_j - \lambda_i)^{2r_i r_j}. \quad (19.6)$$

Proof. Write $Q_i(N) = \sum_{h=0}^{d_i} a_{i,h} N^h$. Let $U_{u,(i,h)} = \lambda_i^u u^h$, $0 \leq h \leq d_i$, and let B be block diagonal, with i th block

$$(B_i)_{h,j} = \begin{cases} \binom{h+j}{h} a_{i,h+j}, & h+j \leq d_i, \\ 0, & h+j > d_i. \end{cases}$$

The binomial expansion of $Q_i(u + v)$ gives the exact factorization $(f(u + v)) = U B U^\top$. Each B_i is anti-triangular, so

$$\det B_i = (-1)^{\binom{r_i}{2}} a_i^{r_i} \prod_{h=0}^{d_i} \binom{d_i}{h}.$$

Lemma 17.1 gives $\det U$. The factorials simplify using

$$\left(\prod_{h=0}^{d_i} h! \right)^2 \prod_{h=0}^{d_i} \binom{d_i}{h} = (d_i!)^{r_i}.$$

Taking determinants proves (19.6) for distinct rates. The same identity holds for coincident rates by polynomial continuation in the λ_i , or directly because U then has repeated columns and both sides vanish. $\square$

Theorem 14.5 follows on taking the canonical labels $\mathbf{k}$ as the blocks of Lemma 19.2 and inserting (18.8). All its non-resonance factors are nonzero throughout the admissible root-multiplicity stratum. In particular, no hidden amplitude-zero locus can lower the maximal recurrence order.

Suppose, locally, exactly two rates meet, their difference vanishes to first order along a parameter curve, and no root itself reaches another root or an endpoint. Formula (14.16) then shows that the secondary determinant vanishes to exactly $2r_k r_j$ orders along that curve. This is a quantitative statement about rank loss, not just a test that some exceptional parameter exists.

20 Resonance and a root-free exact algorithm

20.1 The exact cancellation rule

Let Λ be the set of distinct numerical values among the rates. For $\lambda \in \Lambda$ define

$$Q_\lambda(N) = \sum_{k:\lambda_k=\lambda} P_k(N). \quad (20.1)$$

Some of these sums may be identically zero even though every summand has a nonzero leading coefficient.

Theorem 20.1 (All-parameter minimality rule). *At every admissible parameter choice, the minimal characteristic polynomial is*

$$\chi_q(X) = \prod_{\substack{\lambda \in \Lambda \\ Q_\lambda \neq 0}} (X - \lambda)^{1+\deg Q_\lambda}. \quad (20.2)$$

Before computing any cancellations, a valid smaller annihilator than the labelled product is

$$\Psi_q(X) = \prod_{\lambda \in \Lambda} (X - \lambda)^{1+\max\{d_k:\lambda_k=\lambda\}}. \quad (20.3)$$

Proof. Group (14.9) by numerical rate, discard zero polynomials, and apply Lemma 19.1. The degree of a sum is at most the maximum degree of its summands, proving (20.3). $\square$

Equation (20.2) is exact, but it is not a short classification of every possible resonant cancellation. Such a classification would require a more detailed description of the polynomials Q_λ . The next theorem instead gives a terminating algebraic procedure for each exact input, without constructing any spectral roots.

[Added 29 September 2026, batch 42: for a single repeated root $q = \gamma(x - c)^t$, $c \notin \{0, 4\}$, Part III describes the grouped polynomials exactly (Section 34): after grouping, at most one group loses further multiplicity, and exactly one, except at $c = 2$ (Theorems 30.1 and 30.4).]

20.2 Why a finite prefix can be a proof

Theorem 20.2 (Certified finite-prefix recovery). *Let $h = (h_N)_{N \geq 0}$ be a nonzero sequence over a characteristic-zero field, known to satisfy a monic constant-coefficient recurrence of order at most R from $N = 0$ onward. Its minimal recurrence can be recovered from $h_0, \dots, h_{2R-1}$ as follows. For $r = 1, \dots, R$, solve*

$$h_{n+r} + \sum_{j=0}^{r-1} g_j h_{n+j} = 0, \quad 0 \leq n < R. \quad (20.4)$$

The first soluble system gives the unique minimal monic characteristic polynomial $X^r + \sum_{j < r} g_j X^j$.

Proof. Let $A(E)h = 0$ be a known-to-exist monic annihilator of order R , padding a lower-order one by a power of X if necessary. For any candidate g the residual sequence $u = g(E)h$ also satisfies $A(E)u = 0$, because polynomials in shift commute. If (20.4) holds, its first R values are zero. The monic recurrence then forces every subsequent value to be zero by induction. Thus every soluble system gives a genuine all-index annihilator, not merely a fitted prefix.

The actual minimal annihilator gives a solution for its order, so the first successful order is minimal. Two different monic annihilators of that degree would differ by a nonzero annihilator of smaller degree, which is impossible. The largest index used by the algorithm is $R - 1 + R = 2R - 1$. $\square$

The prior order bound is indispensable. Without it, an arbitrarily long matching prefix does not certify a recurrence for an arbitrary sequence. For (14.1), Theorem 14.2 supplies exactly the bound needed.

20.3 No factorization into individual roots is required

Suppose $q \in \mathbb{Q}[x] \setminus \{0\}$. Strip its powers of x and $x - 4$, and perform square-free decomposition on the remainder:

$$q(x) = \gamma x^m (x - 4)^\ell \prod_{t \geq 1} f_t(x)^t, \quad (20.5)$$

where the nonconstant f_t are monic, square-free and pairwise coprime. Let $b_t = \deg f_t$. Ordinary polynomial gcds with the derivative produce this decomposition; irreducible factorization and numerical root finding are unnecessary. The bound becomes

$$R = \prod_{t \geq 1} (t + 1)^{b_t} \left(\kappa + 1 + \frac{1}{6} \sum_{t \geq 1} b_t t(t - 1) \right). \quad (20.6)$$

Compute the first $2R$ determinants directly from their rational moment matrices and apply Theorem 20.2. This determines the true minimal polynomial even when the roots are complex, multiplicatively dependent, or not conveniently expressible by radicals.

All recovered coefficients lie in $\mathbb{Q}$. More generally, if the input coefficients lie in an exactly computable algebraic number field, the same linear systems work over that field. Solvability over its algebraic closure is equivalent to solvability over the field itself, by Gaussian elimination; uniqueness of the minimal monic polynomial gives descent. For arbitrary unspecified complex numbers, “algorithm” presupposes an exact representation with equality and field operations available. We do not assert an effective equality test for arbitrary complex oracles.

The transparent root-free implementation uses $O(R^4)$ field operations for generating the determinants and searching all orders with naive Gaussian elimination, apart from polynomial preprocessing and moment generation. This is an arithmetic-operation estimate, not a bit-complexity claim. The included fixed-size Christoffel path is much faster when the roots are already given exactly.

20.4 Algebraic rank-drop loci

The finite-prefix proof also turns exceptional orders into explicit polynomial conditions. Fix a degree $D \geq 0$ and consider nonzero polynomial multipliers of exactly that degree. The uniform all-index bound $S = 2^D$ follows from (19.5). For $1 \leq r < S$, form

$$\mathcal{H}_{S,r}(q) = (H_{n+j}(q))_{0 \leq n < S, 0 \leq j \leq r}. \quad (20.7)$$

Then

$$\text{ord}(H(q)) \leq r \iff \text{rank } \mathcal{H}_{S,r}(q) \leq r. \quad (20.8)$$

For the forward direction, the recurrence gives a column dependence. For the reverse direction, take a nonzero dependence, normalize its highest nonzero coefficient, and apply the residual argument of Theorem 20.2. Its degree is at most r .

Every entry $H_n(q)$ is a polynomial in the coefficients of q . Consequently the order-at-most- r locus is defined, within the degree- D coefficient space, by the vanishing of all $(r + 1)$ -minors of (20.7). It is relatively Zariski closed. On a fixed root-multiplicity stratum, S may be replaced by the sharper $\mathcal{R}_0$. Theorem 14.5 is an explicit factorization of the maximal-rank test on that stratum; lower-rank tests can involve further cancellation conditions after rates have already collided.

21 Worked examples and checks against the repository

21.1 Recovering the shifted linear multiplier

Take $q(x) = x^m(a + bx)$, with $b \neq 0$ and $a(a + 4b) \neq 0$. Then $\gamma = b$, $\ell = 0$, $s = 1$, $t_1 = 1$ and $c_1 = -a/b$. The two rates are $-bz$ and $-bz^{-1}$, so their sum and product are

$$-b(z + z^{-1}) = -b(c_1 - 2) = a + 2b, \quad (-bz)(-bz^{-1}) = b^2.$$

Their two degrees are both $\binom{m}{2}$. Theorem 14.2 gives exactly (13.1). At $a = 0$ the multiplier is bx^{m+1} , whose order is $\binom{m+1}{2} + 1$. At $a = -4b$ it is $bx^m(x - 4)$, whose order is $\binom{m}{2} + 2$. These endpoint results follow from the same theorem, without taking a singular limit of the two nonendpoint amplitudes.

Relation to Part I. In this subsection a and b are Part I's parameters, $q(x) = x^m(a + bx)$ is the multiplier of $D_N^{(m)}(a, b) = H_N(q)$, and $-z^{\pm 1}$ are Part I's roots $r_{\pm}$ of (6.1), so the two rates are Part I's $\lambda_{\pm}$ of (6.3). The paragraph is thus a second proof, by a different route, of the minimality in Theorem 2.2 and of the rows $a = 0$, $a = -4b$ and $a(a + 4b) \neq 0$ of Theorem 2.3; the row $b = 0$, $a \neq 0$ is the case $s = 0$ of Corollary 14.4 (an observation of this merge). It does not reprove Part I's coefficient formula, numerator degree, reciprocity or asymptotic constant. One further consistency check (an observation of this merge, not of the manuscript): for $q = x^m$, Theorem 18.1 gives the leading coefficient $B_0(m) (-1)^{\binom{m}{2}} \prod_{h < m} h! = 2^{\binom{m}{2}} \prod_{h < m} h! / \prod_{r < m} (2r)!$, which is Part I's h_m of (1.6), the leading coefficient of $H_N^{(m)}$.

21.2 A repeated fourth power with order fifteen

For

$$q(x) = (2x - 9)^4 = 16(x - \frac{9}{2})^4, \quad (21.1)$$

choose $z = 2$. The five rates, their exact degrees and their leading coefficients are as follows.

k	λ_k	d_k	$\text{lc}(P_k)$
0	1	0	1/729
1	4	3	8/81
2	16	4	-64/27
3	64	3	-2048/81
4	256	0	65536/729

The exact minimal characteristic polynomial is

$$\chi_q(X) = (X - 1)(X - 4)^4(X - 16)^5(X - 64)^4(X - 256). \quad (21.2)$$

The full expansion $H_N = \sum_{k=0}^4 P_k(N) \lambda_k^N$ is

$$\begin{aligned} P_0(N) &= \frac{1}{729}, \\ P_1(N) &= \frac{24N^3 + 204N^2 + 572N + 520}{243}, \\ P_2(N) &= -\frac{1728N^4 + 13824N^3 + 43344N^2 + 62784N + 35648}{729}, \\ P_3(N) &= -\frac{6144N^3 + 21504N^2 + 23552N + 10240}{243}, \\ P_4(N) &= \frac{65536}{729}. \end{aligned} \quad (21.3)$$

These polynomials may be obtained by (18.4), or by exact interpolation after the theorem has established their degrees and nonvanishing. The first determinant values are

$$1, \quad 3401, \quad 2480745, \quad 1026904425, \quad 327789036905, \quad 92503701028201.$$

No polynomial of degree 14 can annihilate this sequence: all five rates are distinct and every displayed top coefficient is nonzero. This nonexistence assertion is proved for the entire sequence, not inferred from the displayed six values.

21.3 Resonance can cancel an entire branch

For $q(x) = (x - 2)^2$, choose $z = i$. The three canonical sectors are

$$P_0(N) = \frac{i}{4}, \quad P_1(N) = N + 1, \quad P_2(N) = -\frac{i}{4}, \quad (21.4)$$

with rates $-1, 1, -1$. The two nonzero constant sectors cancel exactly, so

$$H_N((x - 2)^2) = N + 1, \quad \chi_q(X) = (X - 1)^2.$$

For completeness, the leading-coefficient formula gives the two constants and the coefficient 1 of N in P_1 . The condition $H_0 = 1$ then fixes its constant term. This already proves (21.4). Taking the maximum degree separately for each collided rate, without checking cancellation, would retain an unnecessary $X + 1$ factor.

A similar phenomenon need not involve repeated roots of q . Take

$$q(x) = (x - \frac{9}{2})(x + \frac{1}{2}), \quad z_1 = 2, \quad z_2 = -2. \quad (21.5)$$

The four canonical constants, in order $(0, 0)$, $(0, 1)$, $(1, 0)$, $(1, 1)$, are $-1/15, -1/3, 1/3, 16/15$; their rates are $-1/4, -1, -1, -4$. Hence

$$H_N(q) = \frac{16}{15}(-4)^N - \frac{1}{15}(-\frac{1}{4})^N, \quad \chi_q(X) = X^2 + \frac{17}{4}X + 1. \quad (21.6)$$

The multiplicity profile permits generic order 4, the grouped annihilator has order 3, and exact cancellation lowers the true order to 2. Thus root multiplicities alone cannot determine the order on the resonance locus.

21.4 A nine-value proof of a resonant fourth-power formula

For $q = (x - 2)^4$ and $z = i$, the degree law and grouped annihilator give

$$(E - 1)^5(E + 1)^4H = 0. \quad (21.7)$$

The exact closed formula is

$$H_N((x - 2)^4) = \frac{(N + 1)(N + 2)^2(N + 3)}{12}. \quad (21.8)$$

Here is a finite proof using the already established all-index bound, rather than an extrapolation from data. The right side is a polynomial of degree 4, so it satisfies (21.7). The first nine values of both sides are

N	0	1	2	3	4	5	6	7	8
H_N	1	6	20	50	105	196	336	540	825

These determinant values follow from the 4×4 formula (15.8). They can be reproduced using only integers: write $u_{n,r} = p_n^{(r)}(2)/r!$, start with

$$(u_{0,r})_{r=0}^3 = (1, 0, 0, 0), \quad (u_{1,r})_{r=0}^3 = (1, 1, 0, 0),$$

and apply

$$u_{n+1,r} = u_{n,r-1} - u_{n-1,r}, \quad u_{n,-1} = 0.$$

Then $H_N = \det(u_{N+j,r})_{0 \leq r, j < 4}$. The difference between the two sides of (21.8) satisfies a monic recurrence of order 9 and has its first nine values zero. Induction proves that it is identically zero. Since the product polynomial has degree 4 and nonzero leading coefficient $1/12$, its minimal polynomial is $(X - 1)^5$. The generic order 15 has dropped to 5.

[Added 29 September 2026, batch 42: (21.4) and (21.8) are the cases $t = 2, 4$ of a known product of Cigler for every t , printed with a condensation proof as Theorem 30.4 in Part III.]

21.5 An irreducible rational input

No spectral representation is needed to run the exact algorithm. For $q = x^2 + 1$, the square-free profile is $b_1 = 2$, so $R = 4$. The root-free calculation gives

$$\chi_q(X) = X^4 - 5X^3 + 4X^2 - 5X + 1. \quad (21.9)$$

For $q = (x^2 + 1)^2$ the profile is $b_2 = 2$, and the order is 15. The archive records its full characteristic polynomial and exact input coefficients. These examples illustrate the intended use of the algorithm: its input is a rational coefficient list, not approximate complex roots or guessed multiplicative relations among them.

22 Exact asymptotics and stability of the leading sector

The expansion (14.9) is finite and exact. Consequently it contains more information than a leading asymptotic equivalence. Suppose a label $\mathbf{k}_*$ has strictly larger rate modulus than every other label. Put $\lambda_* = \lambda_{\mathbf{k}_*}$, $d_* = d_{\mathbf{k}_*}$, and let $M = \max_{\mathbf{k}} d_{\mathbf{k}}$. Choose $\rho < |\lambda_*|$ no smaller than the other rate moduli. Then

$$H_N(q) = \lambda_*^N P_{\mathbf{k}_*}(N) + O(\rho^N N^M). \quad (22.1)$$

When there are no other labels the error term is zero. In particular,

$$H_N(q) \sim L_{\mathbf{k}_*} N^{d_*} \lambda_*^N.$$

Every algebraic correction in inverse powers of N can be read from the finite polynomial $P_{\mathbf{k}_*}$; no infinite asymptotic series is needed for fixed parameters.

Corollary 22.1 (Repeated roots outside the interval). *Suppose all nonendpoint roots satisfy $c_i > 4$, and choose $z_i > 1$. Then the unique largest-modulus sector is $k_i = t_i$ for every i . Thus*

$$H_N(q) \sim L_{(t_1, \dots, t_s)} N^\kappa \left((-1)^{\ell+T} \gamma \prod_i z_i^{t_i} \right)^N. \quad (22.2)$$

No multiplicative-independence assumption is needed for this conclusion.

Proof. Reducing any k_i multiplies the modulus by a factor $z_i^{-2} < 1$. Hence the maximal sector is unique even if some of the smaller sectors resonate. At $k_i = t_i$ all $k_i(t_i - k_i)$ vanish, leaving degree κ . Its leading coefficient is nonzero by Theorem 18.1. $\square$

This separates two effects that are easy to conflate. Repeating a root outside the support changes the exponential growth rate and can greatly increase the recurrence order. It does *not* increase the power of N in the leading asymptotic term; that power comes entirely from the endpoint multiplicities. The repeated offendpoint roots instead create higher polynomial degrees in intermediate, subdominant sectors. For (21.1), for instance,

$$H_N((2x - 9)^4) \sim \frac{65536}{729} 256^N,$$

even though its 16^N sector has degree 4 and its recurrence has order 15.

The estimate is for fixed parameters. As roots approach 0, 4 or each other, the spectral gaps and the constants need not be uniform. A double-scaling limit cannot be justified by simply using (22.1) with moving parameters. That is one of the separate research directions listed below.

[Added 29 September 2026, batch 53: for roots approaching 0 or 4 on the scale N^{-2} , Part IV proves the uniform double-scaling limit directly (Theorems 47.1 and 47.4), without using (22.1); roots approaching each other away from the endpoints are not covered.]

[Added 29 September 2026, batch 55: for roots approaching each other at fixed nonendpoint bases on the scale N^{-1} , Part V proves the uniform double-scaling expansion directly (Theorems 64.1, 67.1 and 68.1), without using (22.1); limits uniform up to the endpoints, or as two distinct bases merge, are not covered.]

23 Verification, implementation, and formalization status

23.1 Independent exact calculations

The archive contains a reference implementation in Python 3.10 or later, using only the standard library. Every numerical calculation uses integers or exact rational numbers. Gaussian elimination uses pivoting; there is no assumption that a signed or resonant moment matrix is positive definite.

The implementation has two determinant paths. The first constructs the original matrix in (14.1) from Catalan numbers. The second computes jets from the polynomial recurrence (15.1) and uses the fixed-size Christoffel determinant. The paths share the low-level elimination routine, but neither uses the explicit sector leading constants. A third calculation interpolates sector polynomials at distinct rational spectral parameters and checks their degrees, leading constants and extra values. The secondary-Hankel factorization is tested by yet another determinant, built from the already computed sequence values.

The following are the actual results recorded by `code/verify.py` (shipped here as `code/02-root-collisions-verify.py`; its recorded run is `data/02-root-collisions-verification.json`, with console text `data/02-root-collisions-verification.txt`). Counts include simple and boundary instances; they measure executed identities, not independent universal theorems.

Executed check	Count
Endpoint-jet identities	304
Original moment determinant versus Christoffel determinant	1,554
Summed sector orders versus the multiplicity formula	12
Secondary-Hankel product factorizations, including resonances	15
Exact sector degrees and leading coefficients	48
Full exponential-polynomial evaluations	328
Nonresonant recurrence residuals	200
Nonresonant minimal-recurrence recoveries	9
Resonant recurrence residuals	204
Resonant minimal-recurrence recoveries	12
Inter-root resonance recoveries	3
Root-free multiplicity profiles	111
Root-free recurrence recoveries	3
Central-root fourth-power product evaluations	40
Total; all passed	2,843

The original-matrix comparisons cover 222 multiplier/scale choices at sizes $N = 0, \dots, 6$. They include roots below the support, in its interior, at both endpoints and above the support; multiplicities up to five occur. The sector checks use twelve nonresonant patterns, including mixed endpoint and repeated-root blocks, with recurrence orders up to 26. The central fourth-power formula is checked through $N = 39$, although its finite proof in Section 21 needs only the first nine values.

The distinction between evidence and proof matters. A successful interpolation for twelve parameter choices does not prove Theorem 18.1. Its proof is the unique maximal-degree Cauchy–Binet selection and the three nonzero local determinants. Conversely, once a universal order bound is available, the finite certificate for a *particular* specialization is a rigorous proof by Theorem 20.2. The article uses both kinds of computation but does not conflate them.

23.2 The implementation interface

The module `code/catalan_collisions.py` provides the following principal entry points. (It is shipped here as `code/02-root-collisions-catalan_collisions.py`; §27 explains how to import it.) Polynomial arrays always use ascending powers.

Routine	Purpose
<code>direct_hankel</code>	Original moment determinant for rational coefficients.
<code>christoffel_values</code>	Fixed-size evaluation with rational roots and multiplicities.
<code>sectors</code>	Rates, exact degrees and leading constants for rational z_i .
<code>secondary_hankel_formula</code>	The product in (14.16).
<code>fit_sectors</code>	Exact polynomial interpolation when rates are distinct.
<code>multiplicity_profile</code>	Endpoint multiplicities and square-free root counts.
<code>minimal_recurrence</code>	Recovery from a prefix, given a proved order bound.
<code>recurrence_from_coefficients</code>	Root-free minimal recurrence for rational q .

The last routine is intentionally transparent, not optimized for large bounds. Its optional `max_bound` argument prevents an unexpected large computation. The mathematical theorem permits all complex roots; the reference implementation uses rational inputs, and does not purport to implement a general algebraic-number package.

23.3 A feasible formalization route

No Lean source has been compiled or checked for this article. The repository location does not confer formal status on the present proofs. A formalization should separate the argument into small algebraic layers.

First formalize normalized jets of (15.1) over a characteristic-zero field and their generating functions. Next prove the confluent Vandermonde identity, including the Euler-derivative row change, and formalize the polynomial-shift coefficient bound (17.1). The decisive reusable lemma is the maximal-degree Cauchy–Binet argument: it tracks one distinct list of derivative orders per spectral node and identifies the unique smallest such list.

The orthogonality and integral proof of Christoffel may be formalized analytically, but an algebraic alternative is attractive. One may define the moment functional by $L(x^n) = C_n$, prove its Gram identities for p_n directly from generating functions, and prove Christoffel by determinant multilinearity. This would avoid importing measure theory into the core recurrence theorem.

Finally, the shift-operator minimality lemma, the UBU^T secondary factorization and the finite-prefix recovery theorem are pure finite linear algebra over a field. They should be reusable outside Catalan determinants. An audit must distinguish imported classical facts from newly checked lemmas and must not replace a missing confluence proof with an admitted theorem hidden inside a computation.

24 Further research questions and proposed next steps

The questions below are precise continuations of this article. They are not all asserted to be unsolved in the published literature; a broader literature check is part of each proposed project.

24.1 Closed classification of cyclotomic resonances

For a single repeated root, classify the minimal recurrence as an explicit function of t and the order of z^2 when z is a root of unity other than ± 1 . The degree law reduces the task to finitely many grouped polynomials, but the cancellations can be substantial. The examples at $z = i$ show that counting distinct rates is not enough. A first target is a uniform closed form for $H_N((x-2)^t)$ for every t , together with a direct explanation of the parity effects, rather than a separate finite certificate for each t .

[Added 29 September 2026, batch 42: answered by Part III for one repeated root away from the endpoints, with a correction to the question itself: t and the order L of z^2 do not determine the minimal recurrence, and the sign of z^L is also needed (for $t = 3$, $c = 1$ and $c = 3$ both have $L = 3$ but orders 7 and 6). Theorem 30.1 gives every characteristic multiplicity for $c \neq 2$ in terms of t , L and z^L , and Theorem 37.1 computes it from c without roots. The uniform central formula is a known product of Cigler, credited and re-proved in Theorem 30.4; the parity effects come from the single tied pair of Lemma 34.2. The endpoint cases $c = 0, 4$ are outside Part III.]

24.2 Pairwise resonance and cancellation order

Suppose exactly two labelled rates agree and all other rates are separate. Determine when their leading coefficients cancel, and how many further coefficients can cancel. Theorem 18.1 makes the first test explicit, while (16.6) determines all later tests. The aim is a structural criterion in the root parameters, not just the output of Gaussian elimination. The distinct-root example (21.5) is a small initial case.

[Added 29 September 2026, batch 42: partly answered. Within the rate groups of one repeated root, Part III's Section 34 gives the structural criterion: only the tied pair $k_{\pm} = (t \mp L)/2$ can cancel at the leading order, it cancels exactly when the sign ρ of (34.5) is -1 , and then exactly one further degree is lost unless $c = 2$. Collisions between distinct roots, including the example above, remain open.]

24.3 Sector polynomials and symmetric functions

The degree $k(t-k)$ is the area of a $k \times (t-k)$ rectangle, and the proof uses a mixed confluent Vandermonde. Seek a Schur-function or factorial-Schur description of the entire P_k , not only its leading coefficient. Useful outputs would be exact factorizations, coefficient sign rules for real roots outside $[0, 4]$, and formulas that remain stable as two parameters approach one another. Such a description could also identify an earlier equivalent formulation of the multiplicity law.

24.4 Uniform endpoint and root-collision limits

A concrete endpoint scaling is

$$c_N = 2 + 2 \cosh(\tau/N), \quad q_N(x) = (x - c_N)^t.$$

Determine the limit, locally uniformly in τ , of

$$\frac{H_N(q_N)}{(-1)^{tN} N^{t(t+1)/2}},$$

including its value at $\tau = 0$. The analogous left-endpoint scaling $c_N = 2 - 2 \cosh(\tau/N)$ should use the exponent $t(t-1)/2$. For $t = 1$ the branch formula already reduces these limits to elementary hyperbolic functions. The general problem asks for a uniform confluent determinant limit and error bounds. A second version lets two distinct nonendpoint roots approach at a rate of order N^{-1} .

[Added 29 September 2026, batch 53: the endpoint half is answered by Part IV for any fixed numbers m and ℓ of roots approaching 0 and 4 together, with arbitrary internal collisions: with $\nu = N + (m + \ell)/2$ the normalized determinant is a convergent series in ν^{-2} , locally uniform in the scaled coordinates (Theorem 47.1); its leading term factorizes into confluent cosine and sine determinants, its first correction is explicit (Theorem 47.2), and in the scaling posed here the limits are $F_t^+(\tau)$ at the right endpoint and $F_t^-(\tau)$ at the left, with the stated values at $\tau = 0$ and an N^{-1} term that is only the centring $N \mapsto N + t/2$ (Corollary 54.1); an explicit error bound is Theorem 50.2. Relative to the fixed-endpoint determinant the ratio tends to 1 exactly when every outward displacement is $o(N^{-2})$ (Theorem 47.4). The nonendpoint collisions of the second version remain open (Section 57.1).]

[Added 29 September 2026, batch 55: the nonendpoint collisions of the second version are answered by Part V for any fixed total degree d and any base $c = 2 + 2 \cosh w \notin \{0, 4\}$, with arbitrary collisions inside a cluster and several separated clusters. With $\nu = N + d/2$ and roots $2 + 2 \cosh(w + u_i/\nu)$, H_N is an exact finite sum of sectors $e^{(2k-d)\nu w} \nu^{k(d-k)} F_k(1/\nu)$ whose amplitudes are convergent power series in $1/\nu$, locally uniform through every collision (Theorem 64.1); the first coefficient is explicit (Theorem 66.1), an explicit remainder bound is Theorem 67.1, and separated clusters are Theorem 68.1. For two simple roots the determinant is explicit (Example 74.1). For two roots of multiplicity t at $2 + 2 \cos(\theta \pm \tau/(N+t))$ the normalized limit is an oscillatory Jacobi Hankel determinant and the ν^{-1} term keeps the phase $\sin(2\nu\theta)$ (Theorem 71.1); the value at $\tau = 0$ is (74.7). Unlike Part IV's, the expansion is not even. Limits uniform as the base approaches 0 or 4 remain open (Section 77.1).]

24.5 Other eventually constant Jacobi recurrences

Krattenthaler's recurrence theorem covers many moment families beyond Catalan numbers [3]. Extend the exact degree analysis to a three-term recurrence whose coefficients are eventually constant. Away from branch points, the main new issue is a zero of one branch amplitude caused by the finite initial perturbation. Classify how its vanishing order changes the jet degrees and the noncancellation matrix. This would distinguish universal collision behavior from behavior tied to the Catalan boundary condition $p_1 = x - 1$.

24.6 Rational multipliers and finite poles

Replace $q(x)$ by a rational function whose poles avoid the support. The determinant identities underlying Uvarov transformations provide a starting point [8]. The target is an exact analogue of the sector-degree formula when evaluation and Cauchy-transform blocks occur together. One must first specify which normalizing factors are removed and which recurrences hold from the initial index rather than only eventually.

24.7 Coefficient-space equations and arithmetic reductions

Eliminate the spectral variables from (14.16) to obtain small defining equations for maximal-order failure directly in the coefficients of q . The finite minors in (20.7) are correct but can be very large. Seek sharper degrees, irreducible components and efficient membership tests. For monic integral multipliers, a related problem is to classify primes at which the minimal recurrence drops after reduction. The factorials in the leading constants show why characteristic-zero noncancellation cannot simply be reused unchanged in small characteristic.

24.8 Fast certified algorithms and formal verification

Combine the proved order bound with fast minimal-recurrence algorithms, but reduce the cost of obtaining the initial determinants as well. A useful target is an output-sensitive exact algorithm with bit-complexity bounds in D , the coefficient heights and the final recurrence order. Separately, formalize the maximal-degree determinant lemma and the finite-prefix certification theorem first: these are the shortest reusable path from the present article to a machine-checked result.

25 Conclusion

The selected repository extension has a uniform answer at the level of exponential sectors. A root of multiplicity t_i away from 0, 4 contributes the exact degree $k_i(t_i - k_i)$ to its k_i th branch sector. The two endpoints contribute different triangular numbers. The leading coefficient is a nonzero product, so distinct sectors cannot disappear at nonresonant parameters. This proves the sharp order formula (14.12), with cubic rather than exponential complexity for a single repeated root.

The secondary-Hankel factorization identifies the maximal-order exceptional locus exactly. Resonances may cause further cancellations, but they do not obstruct an exact answer for a specified rational or algebraic multiplier: a proved order bound turns finitely many determinant values into a complete minimal-recurrence certificate. The remaining research questions concern structural descriptions of those cancellations, uniform collision limits, broader moment families and formalization, not gaps in the stated multiplicity proof.

26 Normalization and sign audit

There are four easy ways to obtain an incorrect statement in this problem. This section (Appendix A of the manuscript) records the conventions that prevent them.

Size versus last index. Throughout, N is the determinant size. In a convention that indexes rows by $0 \leq i, j \leq n$, one must substitute $N = n + 1$. The initial term $H_0 = 1$ is part of the recurrence sequence used here.

The Christoffel sign. The product in the multiplier is $\prod(x - a_u)$, whereas the mixed Vandermonde supplies $\prod(a_u - x)$. This contributes $(-1)^{ND}$. The m zero-endpoint rows add $(-1)^{mN}$ when their factors are removed. The remaining sign is therefore $(-1)^{(\ell+T)N}$, exactly the sign in (14.6).

Normalized versus unnormalized jets. The rows of M_N are $p_{N+j}^{(r)}(a)/r!$. Thus Δ_q contains only cross-root differences. Extra products of $r!$ in its denominator would double-count confluence. The factorials appearing later in B_0, B_4, B_z and $\mathcal{V}$ have separate origins and must not be removed by changing the jet convention halfway through the proof.

Two different kinds of collision. A collision of roots of q changes the factorization profile and must be treated by normalized jets. A collision of rates λ_k can occur even when all roots of q are distinct, and must be treated by adding their polynomial coefficients. These are logically different operations. The first creates the degree law; the second can reduce the actual recurrence order.

As a final check, $H_N(uq) = u^N H_N(q)$. The determinant on the left of (14.16) therefore scales by $u^{\mathcal{R}_0(\mathcal{R}_0-1)}$. On its right, the rate powers and rate differences contribute

$$\sum_k r_k(r_k - 1) + 2 \sum_{k < j} r_k r_j = \mathcal{R}_0(\mathcal{R}_0 - 1),$$

while the sector leading coefficients do not depend on the overall scale γ . The scaling agrees exactly.

27 Reproducing the package

This section is Appendix B of the manuscript, adapted to the shipped file names. As delivered, the package was reproduced from its top directory by

```
python3 code/verify.py --out data
latexmk -pdf -interaction=nonstopmode -halt-on-error article.tex
```

The first command regenerates the recorded numerical outputs; the second builds the PDF. There are no third-party Python dependencies. *In this report these commands must not be run in place.* Here `code/verify.py` and `data/` belong to Part I, so the first command would run Part I's suite and rewrite Part I's data; the shipped verifier `code/02-root-collisions-verify.py` still imports its sibling module under the delivered name `catalan_collisions`, which is not importable under the shipped name, and its default output directory `data` would overwrite Part I's `data/verification.json` and `data/verification.txt`. The delivered Makefile, shipped as `code/02-root-collisions-Makefile`, has the same defect: its `check` target runs `code/verify.py --outdata`, which here is Part I's verifier. Restore the delivered names in a scratch copy instead (POSIX shell, run from the report directory):

```
W=/path/to/scratch/root-collisions
mkdir -p "$W/code" "$W/recorded"
for f in code/02-root-collisions-*.py; do
  cp "$f" "$W/code/${f#code/02-root-collisions-}"; done
for f in data/02-root-collisions-*; do
  cp "$f" "$W/recorded/${f#data/02-root-collisions-}"; done
cd "$W" && python3 code/verify.py --out data
```

The run writes seven files to `$W/data`, which can be compared with the recorded copies in `$W/recorded`; on 28 September 2026 this recipe passed all 2,843 checks, and the seven files agreed with the recorded ones apart from Windows line endings. The following root-free example, run in the same scratch directory, needs no spectral factorization:

```
import sys
```

```

sys.path.insert(0, "code")
from catalan_collisions import recurrence_from_coefficients

chi, bound = recurrence_from_coefficients([1, 0, 1], max_bound=16)
print(bound) # 4
print(chi)    # ascending coefficients: 1, -5, 4, -5, 1

```

The main data files are shipped in `data/` with the prefix `02-root-collisions-` (delivered without it):

- `verification.json`, `verification.txt`: the recorded run of Section 23;
- `sectors.json`: the twelve nonresonant patterns, with sector polynomials, leading constants and characteristic polynomials;
- `resonances.json`: exact minimal recurrences at resonant parameters;
- `rational_input_examples.json`: root-free rational-coefficient examples;
- `central_root_certificate.json`: the nine-value certificate of (21.8);
- `fourth_power_sequence.csv`: exact values of $H_N((2x - 9)^4)$.

The code is shipped in `code/` with the same prefix (`verify.py`, `catalan_collisions.py` and the delivered root Makefile), and the provenance notes, delivered as `notes/provenance.md`, as `02-root-collisions-provenance.md` at the report root. Rational values are stored as exact strings, for example `"8/81"`. Arrays representing polynomials use increasing powers. Sector polynomials are polynomials in the determinant size N , not generating-function numerators.

Part III. Cyclotomic resonances of one repeated root

Part III was added on 29 September 2026 in batch 42 of ProveIt's incoming-report intake, from one manuscript written against the pinned state of Parts I and II. It answers Part II's research question "Closed classification of cyclotomic resonances" (Section 24.1) for a single repeated root away from the endpoints: at every parameter it determines the minimal recurrence, every characteristic multiplicity, and a root-free factorization, and it shows that the question's proposed parameters, the multiplicity t and the order of z^2 , do not determine the answer. Parts I and II are unchanged except for dated pointers to this addition (Sections 11, 20, 21 and 24), a line on the title page (whose top space was reduced by 3 mm so that it still fits on one page) and a paragraph of the abstract, a contents entry, one bibliography entry used only by Part III and a clause in the closing note on source dates, and two new labels on Part II's research subsections. Section 28 records the provenance and fixes the notation of the addition.

28 Provenance and conventions of Part III

28.1 One manuscript, one addition

Part III is batch-42 manuscript 01, *A One-Defect Law for Cyclotomic Catalan Hankel Recurrences*, subtitled *Complete single-root classification, a necessary sign parameter, and the central exception*, a research article prepared with OpenAI ChatGPT for Vladimir Reshetnikov's ProveIt research program and dated 29 September 2026 (22 A4 pages as delivered). It arrived as the archive `ProveIt_Cyclotomic_Catalan_Recurrences.zip` (inner directory `cyclotomic_catalan/`, main file `article.tex`) in ProveIt commit 8315d24e3 and was placed in 3609d0473. It was written against ProveIt commit 9754e83603e223811b7515eb892f22c847144593, reading this report's combined `article.tex` (Git blob b29ea42fc5bba69d23c3b036bee25357e202f356), its provenance notes and the directory listing at that commit; its source notes add that the standalone Part II manuscript was read as well. The report's files are unchanged between the pin and the placement commit (the same blob), so every statement the manuscript makes about "the repository report" or "the prior report" refers to Parts I and II as printed above.

The manuscript's code and data are shipped beside the earlier files with the prefix `03-cyclotomic-` (Part I's unprefix files count as 01, Part II's as 02), and its source and comparison notes, delivered as `SOURCES.md`, as `03-cyclotomic-SOURCES.md` at the report root. Its manuscript, delivery README and PDF are not shipped; they survive in the arrival commit. Section 39 lists the shipped files.

The manuscript contributes, in the order printed here: the repository question, what is already known, and why grouping equal rates is not enough (Section 29); the definitions, the complete noncentral classification with its elementary sign test, and the central theorem (Section 30); a self-contained derivation of the confluent Christoffel identity in its normalization (Section 31); an exact subset expansion of every canonical sector (Section 32); the leading coefficient, the first centred correction and an exact reflection of the sector polynomials (Section 33); the one-defect proof (Section 34); closed orders, the exceptional locus and its count, the reflection $c \mapsto 4 - c$ and quadratic growth in the multiplicity (Section 35); the central exception via Cigler's known product, with a self-contained condensation proof (Section 36); a root-free factorization (Section 37); worked examples and the classification of all rational roots (Section 38); its verification record and rebuild instructions (Section 39); eight research directions (Section 40); a conclusion; a normalization and proof audit (Section 42); and a source and package concordance (Section 43). Every result, proof, example, remark, limitation and question of the manuscript is printed. Where it re-derives something Parts I or II already prove, the overlap is marked (Section 28.5).

The manuscript's own abstract reads:

Let $H_N^{(t)}(c)$ be the size- N Hankel determinant of the Catalan moments multiplied by $(x - c)^t$, with $c \neq 0, 4$. Write $c = 2 + z + z^{-1}$ and let L be the order of z^2 , possibly infinite. The nonresonant minimal order $t + 1 + \binom{t+1}{3}$ is already established in the ProveIt report that motivates this work. We determine the minimal recurrence at every remaining specialization.

For $3 \leq L \leq t$, put $\eta = \mathbf{1}_{t \equiv L \pmod{2}}$. The exact order is

$$R(t, c) = \frac{L(3t^2 - L^2 + 13 - 3\eta)}{12} - \delta, \quad \delta \in \{0, 1\}.$$

The defect occurs precisely when $\eta = 1$ and $(-1)^{L(t+1)/2} z^{Lt} = -1$. We determine the multiplicity of every characteristic root, not only their total. The proof isolates the first two coefficients of each canonical exponential sector using an exact signed-Vandermonde identity. A collided sector can lose just one degree away from $c = 2$; at $c = 2$ an entire group can disappear. The latter case is governed by a product formula already proved by Cigler for middle-binomial Hankel determinants, which we credit and rederive with the required normalization.

The answer cannot depend on t and L alone: for $t = 3$, $c = 1$ and $c = 3$ both have $L = 3$, but their minimal orders are 7 and 6. Consequences include a complete rational-parameter classification, an exact count of exceptional parameters, an eventual quadratic quasipolynomial law in t , and a root-free factorization computed with $O(t)$ scalar field operations. A standard-library Python audit supplies 22,124 successful exact checks, including independent determinant computations in cyclotomic integer rings.

Status (manuscript 01). This is an unrefereed, AI-assisted manuscript with ordinary mathematical proofs, not a Lean formalization. The proposed extension is the noncentral cancellation classification and its coefficient mechanism: the explicit first centred sector coefficient, the noncentral one-defect theorem, the minimal characteristic polynomial at every cyclotomic specialization and the resulting sign-sensitive order laws, with the exceptional-parameter count, the reflection law and the rational-input classification as corollaries. Classical determinant identities (the Christoffel formula and its confluent form, Andréief's identity, Desnanot–Jacobi condensation), general rationality and the order-at-most- 2^t construction [3], Part II's generic degree law and finite recurrence-recovery algorithm, and Cigler's central product [9] are not presented as new; the signed Vandermonde evaluation and the shift-operator minimality lemma are elementary uses of established techniques, not standalone priority claims. First-discovery priority has not been established: a specialized equivalent result may exist in Wronskian, orthogonal-polynomial or symmetric-function notation, the targeted searches that returned unrelated results do not establish absence from the literature, and independent expert review and a broader priority audit are still appropriate. The comparison with this report is targeted, not a proof audit of every report in ProveIt. The manuscript does not solve multi-root resonance, does not claim uniform collision asymptotics, and does not transfer its characteristic-zero minimality theorem unchanged to positive characteristic; the endpoints $c = 0, 4$ are outside its theorems. Its 22,124 exact assertions are finite supporting checks, not a universal proof. No proof assistant was run, nothing in Part III is formalized, and its placement beside a Lean project confers no formal status. (Sources: the manuscript, its delivery README and its source notes, 03-cyclotomic-SOURCES.md.)

28.2 Reading conventions

Inside Sections 29–43, “this article” and “the present paper” mean manuscript 01 as printed in Part III; “the repository report”, “the ProveIt report”, “the prior report” and “the preceding repository work” mean Parts I and II at the pin 9754e8360; and “the older root-collisions manuscript” is Part II. The manuscript cites Part II's standalone section numbers: its Section 12.1 and Section 12.2 are Sections 24.1 and 24.2 here (“Closed classification of cyclotomic resonances” and “Pairwise resonance and cancellation order”). “The archive” or “the package” means the delivered package, whose code and data are shipped under the prefixed names listed in Section 39. The manuscript's two appendices are printed as Sections 42 and 43, so that Part I's appendices keep their letters. The manuscript cites Krattenthaler's Ramanujan

Journal paper through arXiv:2101.04225, version 5 of 10 September 2021, and points to its Theorem 1, its confluence discussion and Section 8; that is Part I’s reference [3]. Its Barry reference is Part I’s [1], cited as background only. Bracketed paragraphs beginning “Merge note” are this merge’s, not the manuscript’s.

28.3 Notation of Part III

Part III keeps the manuscript’s symbols. Within Sections 29–43 every symbol has its Part III meaning, except in merge notes and where a sentence names Part I or II or cites one of their equations. The single multiplier is $(x - c)^t$ with one root $c \notin \{0, 4\}$; in Part II’s notation this is $q = \gamma x^m (x - 4)^\ell \prod_i (x - c_i)^{t_i}$ with $\gamma = 1$, $m = \ell = 0$, $s = 1$, $t_1 = t$, $c_1 = c$, $z_1 = z$, and the manuscript’s $H_N^{(t)}(c)$ is Part II’s $H_N((x - c)^t)$ (14.1). Table 2 lists the symbols that a reader of Parts I and II is likely to misread, and the notions that need care. No symbol was renamed and no normalization changed: the manuscript’s rates λ_k are Part II’s λ_k for one root, and its canonical sector polynomials $P_{t,k}$ have Part II’s normalization (Section 28.5).

Table 2: Symbols and notions of Part III. The last column gives the meaning of the same letter in Part I or II, or the tempting false reading.

Symbol or notion	Meaning in Part III	Not to be confused with
$H_N^{(t)}(c)$	$H_N((x - c)^t)$, the superscript being the multiplicity t (30.1)	Part I’s $H_N^{(m)} = \det(C_{i+j+m})$, whose superscript is the shift; formally $H_N^{(t)}(0) = H_N^{(t)}$, but $c = 0$ is outside the theorems.
<i>resonance</i> ; <i>exceptional</i> ; the “resonant range”	for one root, Part II’s resonance (two equal rates) happens exactly when $L \leq t$, and Corollary 35.1 shows that this is exactly when $R(t, c) < R_{\text{gen}}(t)$ (“exceptional”); the manuscript’s “resonant range” $3 \leq L \leq t$ omits the central case $L = 2$, $c = 2$	reading “resonant range” as all resonances: the central root is resonant for $t \geq 2$ but is treated separately (Theorem 30.4).
<i>one defect</i> ; δ	further loss <i>after</i> grouping equal rates, 0 or 1; the baseline $R_{\text{grp}} = \sum_r (d_r + 1)$ is the degree of Part II’s grouped annihilator Ψ_q (20.3)	a loss of at most one from the generic order $R_{\text{gen}}(t)$, which is false (Remark 30.2).
t, c, z, γ	multiplicity, root, spectral parameter $c = 2 + z + z^{-1}$, scale	Part II’s t_i, c_i, z_i, γ (agreeing for one root); Part I’s generating-function variable t and coefficients $c_{N,m,k}$.
$L = \text{ord}(z^2)$ $\lambda_k = (-1)^t z^{2k-t}$	the order of z^2 , possibly infinite rate of the k th sector (30.4)	Part II’s sector leading coefficients L_k . equal to Part II’s λ_k for one root with $\gamma = 1$; Part I’s $\lambda_\pm$.
$R(t, c), \chi_{t,c}, R_{\text{gen}}(t), R_{\text{grp}}$	minimal order, minimal polynomial, generic order (29.1) (Part II’s $\mathcal{R}_0$ for one root, (14.14)), grouped order (35.2)	Part I’s polynomials $R_n(y)$ (4.1); Part II’s plain R (a bound or a size).
$I_r, d_r, e_r, G_r(N)$	residue classes of sector indices, their maximal degree, exponents, and grouped sector polynomials (34.2)	Part I’s $d = \binom{m}{2}$ and $e = d + 1$; Part II’s d_k ; G_r is Part II’s Q_λ (20.1).
$\eta, k_\pm, r_*, d_*, \rho, \lambda_*$	parity indicator, tied indices, their class, tied degree, leading ratio, defect root	—
$T = \binom{t}{2}$ (Section 32 onward)	a triangular number	Part II’s $T = \sum_i t_i$ (which is t here); Part I’s $T_{n,k,s}$.
$A_u = \prod_{j < u} j!, K_t(z)$	superfactorial and the common sector factor (32.1)	Part II’s amplitudes $A_\pm(z)$; Part I’s constant A_m and $K = (m - 1)^2$; Part II’s index set $\mathcal{K}$.

Symbol or notion	Meaning in Part III	Not to be confused with
$J, S, S^c, \sigma(S), \Delta(S)$	$\{0, \dots, t-1\}$, a subset, its complement, its sum, its Vandermonde magnitude (32.2)	$\sigma = z^L$ without argument (Section 37); S_a in (37.4); Part II's $S = 2^D$ and Δ_q ; $\Delta_N(x)$ in Section 31 is a Vandermonde.
$F_{t,k}, P_{t,k}(N; z)$	raw and canonical sector polynomials (32.3), (32.4)	$P_{t,k}$ is Part II's P_k for one root; Part I's numerator $P_m(t)$.
q	a multiplier in (31.5), but an indeterminate in Lemma 33.1 and $q = z^2$ in the proof of Theorem 33.2	Part II's multiplier $q(x)$; Part I's denominator $q(t)$.
$h = t - k, d = kh, w = N + t/2$	complementary index, sector degree, centred size variable	Part I's h_m and $d = \binom{m}{2}$; Part II's nodes w_a .
$\ell_{t,k}(z), b_{t,k}(z), Q_{t,k}(w)$	leading coefficient (33.3), first centred correction (33.7), remainder of degree at most $d - 2$	Part II's multiplicity ℓ of the root 4; Part I's parameter b ; $Q_a(X)$ below; Part II's Q_λ and Part I's $Q_m(N; r)$.
$B(S), \bar{B}; B_{a,b}(N)$	offset sum and its weighted mean (33.5); the rectangular product (30.12) with integers a, b	Part I's $B_\pm(N)$ and parameters a, b ; Part II's B_0, B_4, B_z, B_i and $B_q(u)$.
$\mu; \mu_n; \alpha$	the Catalan measure; centred moments (36.2); the arcsine measure	Part I's shifted measures μ_m (μ is Part I's μ_0 and Part II's μ); Part III's μ_n is a number, not a measure.
$D_t(N); h_N^{(t)}$	Cigler's middle-binomial Hankel determinant; $\det(\mu_{i+j+t})$ (Section 36)	Part I's $D_N^{(m)}(a, b)$ and h_m ; Part II's $D = \deg q$.
$s_N = (-1)^{N(N+1)/2}$	a period-four sign	Part I's sign s_m ; Part II's s and $s(z)$.
$v, U_a, S_a, \sigma, Q_a(X), E_a$	$c - 2$; Chebyshev values; $z^a + z^{-a}$; z^L ; quadratic factors and their exponents (Section 37)	Part II's matrix U in UBU^T ; the shift operator E (as in Part II); Part I's $E_m(N)$.
$M, \Phi_M, \varphi, \mathcal{E}_t$	primitive order of z in the audit ring (so $L = M/\gcd(M, 2)$), cyclotomic polynomial, Euler's function, the exceptional set (35.3)	Part II's M_N and M ; Part I's $M_{m,\ell}$; E the shift.

28.4 What Part III answers, and how it corrects the question

Part II's first research question (Section 24.1) asked, for a single repeated root, for the minimal recurrence “as an explicit function of t and the order of z^2 ”, with a uniform closed form for $H_N((x-2)^t)$ and an explanation of the parity effects. Part III answers it for one repeated root away from the endpoints: Theorem 30.1 gives every characteristic multiplicity for $c \neq 2$, Theorem 30.4 the central root (a known product of Cigler, credited), and Theorem 37.1 a root-free factorization. It also shows that the question's framing was incomplete: t and $L = \text{ord}(z^2)$ do *not* determine the answer in odd parity, and the sign of z^L is a necessary further datum. The smallest instance is $t = L = 3$, where $c = 1$ and $c = 3$ have minimal orders 7 and 6 (Corollary 35.2 and (38.1)–(38.2)). The corrected statement of the answer is therefore a function of t, L and $z^L \in \{1, -1\}$, and the parity effects are explained by the tied pair of Lemma 34.2 and the sign ρ of (34.5). Part II's second question (“Pairwise resonance and cancellation order”) is answered only for the tied groups of one repeated root (Section 34); collisions between distinct roots, and endpoint factors, remain open (Section 40).

Consistency with Part II's data (an observation of this merge, checked by a separate exact computation not shipped with the report). Part II's `data/02-root-collisions-resonances.json` records twelve single-root cases, $c = 1, 2, 3$ with $t = 1, \dots, 4$, each with its minimal order, full characteristic polynomial and initial values. All twelve characteristic polynomials, not only the orders, coincide with those given by Theorems 30.1, 30.4 and 37.1, and the orders coincide with the rows $t \leq 4$ of `data/03-cyclotomic-orders_c1_c2_c3.csv`: 0 mismatches. Seven of the twelve lie below the generic order ($c = 1, 3$ at $t = 3, 4$ and $c = 2$ at $t = 2, 3, 4$); the other five ($t = 1$ for all three roots, and

$t = 2$ for $c = 1, 3$) have $L > t$, so they are nonresonant by Corollary 35.1 and have the generic order, although that file is described as recording resonant parameters. The case $(x - 2)^3$ of `data/02-root-collisions-rational_input_examples.json`, with characteristic polynomial $(X^2 + 1)^3$, is the case $m = 1$ of (30.16). The central products at $t = 2, 4$ are $B_{1,1}(N) = N + 1$ and $B_{2,2}(N) = (N + 1)(N + 2)^2(N + 3)/12$, which are (21.4) and (21.8); the manuscript's residue-class formulas (38.3) and the table for $c = 1$ in Section 38 agree with direct determinant evaluation.

28.5 Where the merge had to choose

1. *Placement.* Part III is printed after Part II and before Part I's appendices, so that no section, equation, theorem or table number of Parts I and II moves (checked against the `.aux` file of a build of the committed text). Its sections continue the numbering (Sections 28 to 43); the manuscript's Sections 1–13 are Sections 29–41, and its Appendices A and B are Sections 42 and 43.
2. *Labels.* Every label of Part III carries the prefix `cyc:`; without it, eight of the manuscript's labels, `sec:main`, `sec:examples`, `thm:main`, `lem:Christoffel`, `lem:minimal`, `eq:Christoffel`, `eq:lambda` and `eq:measure`, would have collided with Part I's bare labels. Two labels were added to Part II's research subsections so that they can be cited; no existing label was changed.
3. *Overlaps with Parts I and II.* The manuscript re-derives, without assuming the repository report correct, several things already proved there. Printed with a pointer and kept: the Catalan measure, orthogonality and $\det(C_{i+j}) = 1$ (a third route, after Part I's Section 3 and Part II's Section 15); the confluent Christoffel identity (Lemma 31.1 is Part II's (15.8) for $q = (x - c)^t$, and Part I's Lemma 4.1 and Proposition 4.2 are other special cases); the minimal-annihilator lemma (Lemma 34.1 restates Part I's Lemma 6.1 and Part II's Lemma 19.1, with a third proof and the generating-function statement); the exact sector degree $k(t - k)$ and the generic order (Part II's Theorem 14.2 and Corollary 14.3, re-proved here by the subset expansion and the explicit leading coefficient (33.3), a different route from Part II's Cauchy–Binet argument); and the central examples at $t = 2, 4$, which the central product of Theorem 30.4 contains. As an observation of this merge, the leading coefficients (33.3) reproduce all five leading coefficients of Part II's order-fifteen example (the table before (21.2), at $z = 2$) and the two constants of (21.4): the two sector normalizations agree. No statement of Parts I or II was changed.
4. *A third route to half of the central product* (an observation of this merge). The neighbouring report `cigler-conjecture-8-schur` (same directory) proves, for Cigler's polynomials $b_m(t)$ with $b_m(1) = \binom{m}{\lfloor m/2 \rfloor}$, an even-shift Hankel–Schur identity (its Theorem 3.1) and, at $t = 1$, the dimension product $\prod_{i,j \leq k} (n + i + j - 1)/(i + j - 1)$ (its Proposition 7.2). Together they give $D_{2k}(n) = B_{k,k}(n)$, the even case of Cigler's product used in Section 36, by a Schur-function route; the odd case is not covered there. That report and its two siblings concern conjectures of the same Cigler paper [9]; Part III uses its proved Theorem 1, not a conjecture.
5. *Bibliography.* The manuscript's references to Krattenthaler and Barry are Part I's [3, 1]; its reference to Cigler's middle-binomial paper is added to the common bibliography as [9]. Its reference to the pinned repository report is replaced by Parts I and II themselves and the pin recorded in Section 28.1; the sentence that cited it now points to Section 28.1.
6. *File names.* Section 39 gives the shipped names beside the delivered ones. The manuscript's rebuild commands are printed as delivered but must not be run in place here; a recipe on a scratch copy follows them.
7. *Front matter.* The manuscript's title page, abstract, status paragraph and package paragraph are replaced by Section 28.1, which prints the abstract and the status. The package paragraph is not reprinted because the delivered source and PDF are not shipped; the shipped code and data are listed in Section 39.
8. *Notation.* The manuscript's symbols are kept, with the scoping convention and Table 2 of Section 28.3; nothing was renamed. One evident index slip in (31.7) is corrected, with a merge note after Lemma 31.1.

29 The repository question and the scope of the answer

The relevant ProveIt package is *Shifted Catalan Hankel Polynomials*, whose Part II incorporates *Root Collisions and Sharp Recurrences for Polynomially Weighted Catalan Hankel Determinants* (Parts I and II above; the manuscript’s bibliography entry for this repository report is replaced by Section 28.1). The selected source was checked against repository commit

9754e83603e223811b7515eb892f22c847144593.

Its directory is

SetTheory/Cardinals/docs/reports/hankel-determinants/catalan-and-ballot/
shifted-catalan-hankel-polynomials/.

The combined `article.tex` has Git blob identifier `b29ea42fc5bba69d23c3b036bee25357e202f356`.

For a single nonendpoint root of multiplicity t , that report gives a canonical expansion into $t + 1$ exponential sectors, of respective polynomial degrees $k(t - k)$ for $0 \leq k \leq t$. Distinct exponential rates therefore give the exact order

$$R_{\text{gen}}(t) = \sum_{k=0}^t (k(t - k) + 1) = t + 1 + \binom{t + 1}{3}. \quad (29.1)$$

At resonances it provides a finite exact algorithm, but its research agenda asks for a structural classification when z is a root of unity. It also asks for a uniform central-root formula and an explanation of parity, beyond its separately certified examples at multiplicities two and four. These are the local research questions addressed here.

[Merge note: in this report the canonical expansion and its degrees are Part II’s Theorem 14.2, the generic order (29.1) is its Corollary 14.3, the finite exact algorithm is Theorems 20.1 and 20.2, the research agenda is Section 24, and the certified central examples at multiplicities two and four are (21.4) and (21.8).]

There are two distinct conclusions. First, the noncentral cyclotomic problem admits a closed answer: at most one collided rate has a leading cancellation, and that cancellation loses exactly one multiplicity. Second, the suggested input “multiplicity and order of z^2 ” is insufficient in odd parity. The sign of z^L is necessary. Thus the result both completes a classification and corrects the parameter list needed to state it.

29.1 What is already known

The Christoffel determinant formula, its confluent version, and general constant-coefficient recurrences for polynomially modified Catalan moments are classical. Krattenthaler [3] treats these identities and, in Section 8, gives the familiar order-at-most- 2^t recurrence construction. Barry’s note [1] is part of the background for the repository’s shifted-linear-multiplier work. Neither rationality alone nor the existence of an exponential expansion is a novelty claim here.

The central product used in Section 36 is also not new. Cigler’s Theorem 1, equation (23), in [9] evaluates the shifted Hankel determinants of $\binom{n}{\lfloor n/2 \rfloor}$. A translation and a row–column sign change give exactly the $c = 2$ product required here. His proof also uses condensation. Recognizing this prior result closes the repository’s central subproblem without assigning it new priority.

The additional content is the explicit first centered sector coefficient, the noncentral one-defect theorem, the minimal characteristic polynomial at every cyclotomic specialization, and the resulting sign-sensitive order laws. These statements are not present in the selected repository source. A broader literature equivalence in different orthogonal-polynomial or symmetric-function notation has not been ruled out. The comparison is not a proof audit of every report in ProveIt.

29.2 Why the problem is not solved by identifying equal rates

An exponential polynomial $\sum_k P_k(N) \lambda_k^N$ must first be grouped by equal numerical values of λ_k . The largest degree within one group is only an upper bound for the degree of its sum. Equal leading coefficients can cancel, and symmetry can make the whole sum zero. The prior report's central square already illustrates the latter phenomenon [(21.4)].

The main proof computes one more coefficient than the degree law. That additional coefficient is sufficient everywhere except at the single central value $c = 2$. This is what turns a finite procedure for each example into a uniform classification.

30 Definitions and the complete classification

Write $\mathbb{N}_0 = \{0, 1, 2, \dots\}$ and

$$C_n = \frac{1}{n+1} \binom{2n}{n}, \quad H_N^{(t)}(c) = \det_{0 \leq i, j < N} \left(\sum_{a=0}^t \binom{t}{a} (-c)^{t-a} C_{i+j+a} \right), \quad H_0^{(t)}(c) = 1. \quad (30.1)$$

Throughout the main classification, $t \geq 1$ and $c \in \mathbb{C} \setminus \{0, 4\}$. Choose a root z of

$$z + z^{-1} = c - 2. \quad (30.2)$$

Then $z \neq 0, 1, -1$. The two choices are reciprocal and give the same answers after relabeling.

A monic polynomial $g(X) = \sum_{j=0}^R g_j X^j$ is an annihilator when

$$\sum_{j=0}^R g_j H_{N+j}^{(t)}(c) = 0 \quad \text{for every } N \geq 0. \quad (30.3)$$

The least possible degree is denoted $R(t, c)$, and the monic polynomial of that degree by $\chi_{t,c}$. All characteristic roots occurring below are nonzero; consequently the same minimal polynomial is necessary even for recurrences required only after a finite initial segment.

Let $L = \text{ord}(z^2) \in \{2, 3, \dots\} \cup \{\infty\}$. Define

$$\lambda_k = (-1)^t z^{2k-t} \quad (0 \leq k \leq t). \quad (30.4)$$

If $L < \infty$, equality of these rates means exactly equality of the indices modulo L .

Theorem 30.1 (Minimal recurrence at every noncentral parameter). *Assume $c \notin \{0, 2, 4\}$.*

If $L > t$, including $L = \infty$, then

$$\chi_{t,c}(X) = \prod_{k=0}^t (X - \lambda_k)^{k(t-k)+1}, \quad R(t, c) = R_{\text{gen}}(t). \quad (30.5)$$

If $3 \leq L \leq t$, define, for $0 \leq r < L$,

$$I_r = \{k : 0 \leq k \leq t, k \equiv r \pmod{L}\}, \quad d_r = \max_{k \in I_r} k(t-k), \quad \eta = \mathbf{1}_{t \equiv L \pmod{2}}. \quad (30.6)$$

When $\eta = 1$, also put

$$k_- = (t-L)/2, \quad k_+ = (t+L)/2, \quad r_* \equiv k_- \pmod{L}, \quad 0 \leq r_* < L, \quad (30.7)$$

and set

$$\delta = \begin{cases} 1, & \eta = 1 \text{ and } (-1)^{L(t+1)/2} z^{Lt} = -1, \\ 0, & \text{otherwise.} \end{cases} \quad (30.8)$$

The exponent $L(t+1)/2$ is an integer whenever it is used. Then

$$\chi_{t,c}(X) = \prod_{r=0}^{L-1} (X - \lambda_r)^{e_r}, \quad e_r = d_r + 1 - \delta \mathbf{1}_{r=r_*}. \quad (30.9)$$

When $\eta = 0$, the subtracted term means zero and no r_* is needed. A factor with exponent zero is omitted. In particular,

$$R(t, c) = \frac{L(3t^2 - L^2 + 13 - 3\eta)}{12} - \delta. \quad (30.10)$$

Remark 30.2 (The meaning of one defect). The baseline in (30.10) is $R_{\text{grp}} = \sum_r (d_r + 1)$, not the larger generic order $R_{\text{gen}}(t)$. Merging many rates can already cause a large decrease from R_{gen} . The theorem says that *further cancellation after this grouping* decreases the order by either zero or one.

Corollary 30.3 (An elementary version of the defect test). *In the resonant range $3 \leq L \leq t$, the defect is one exactly in one of the following two disjoint cases:*

$$\begin{aligned} t, L \text{ even}, \quad L &\equiv 2 \pmod{4}; \\ t, L \text{ odd}, \quad z^L &= (-1)^{(t-1)/2}. \end{aligned} \quad (30.11)$$

There is no defect when t and L have different parity.

The proof of the theorem occupies Sections 31–35. The remaining parameter is governed by a known product.

Theorem 30.4 (Central root; Cigler's product with Catalan normalization). *For $a, b \in \mathbb{N}_0$ put*

$$B_{a,b}(N) = \prod_{i=1}^a \prod_{j=1}^b \frac{N+i+j-1}{i+j-1}. \quad (30.12)$$

Every empty product is one. For $m \in \mathbb{N}_0$,

$$H_N^{(2m)}(2) = B_{m,m}(N), \quad (30.13)$$

$$H_N^{(2m+1)}(2) = (-1)^{N(N+1)/2} B_{m,m+1}(N). \quad (30.14)$$

Their minimal polynomials are

$$\chi_{2m,2}(X) = (X-1)^{m^2+1}, \quad (30.15)$$

$$\chi_{2m+1,2}(X) = (X^2+1)^{m(m+1)+1}. \quad (30.16)$$

The even formula includes the trivial multiplier $t = 0$.

Together these results classify every single repeated nonendpoint root, not only rational roots or a bounded list of multiplicities. The endpoints $c = 0, 4$ are intentionally outside the theorem; they require endpoint jets and have different triangular degree laws.

30.1 Immediate generating-function and scaling consequences

If the distinct characteristic roots and their exponents are (λ_r, e_r) , the reduced denominator of $\sum_{N \geq 0} H_N^{(t)}(c) u^N$ is exactly

$$\prod_r (1 - \lambda_r u)^{e_r}. \quad (30.17)$$

There is a numerator of degree less than $R(t, c)$ and constant term one. Thus the theorem determines all poles and their exact orders as well as the recurrence. For the multiplier $\gamma(x-c)^t$, $\gamma \neq 0$, all determinants are multiplied by γ^N . Its monic minimal polynomial is $\gamma^{R(t,c)} \chi_{t,c}(X/\gamma)$.

31 Catalan orthogonality and the confluent determinant

We derive the needed determinant identity rather than assuming the correctness of an unrefereed repository lemma. These are classical arguments; their role here is normalization and self-containment.

31.1 A convenient monic orthonormal family

Define

$$d\mu(x) = \frac{1}{2\pi} \sqrt{\frac{4-x}{x}} dx \quad (0 < x < 4). \quad (31.1)$$

Its moments are C_n . For example, the substitution $x = 4v$ gives a beta integral; integration by parts gives successive moment ratio $2(2n+1)/(n+2)$, with zeroth moment one, exactly the Catalan ratio.

Let

$$p_0(x) = 1, \quad p_1(x) = x - 1, \quad p_{n+1}(x) = (x-2)p_n(x) - p_{n-1}(x) \quad (n \geq 1). \quad (31.2)$$

All p_n are monic. At $x = 2 + 2\cos\theta$, $0 < \theta < \pi$,

$$p_n(x) = \frac{\sin((n+\frac{1}{2})\theta)}{\sin(\theta/2)}, \quad d\mu = \frac{2}{\pi} \sin^2(\theta/2) d\theta, \quad (31.3)$$

where the orientation of the integral is reversed as necessary. The elementary sine orthogonality integral therefore gives

$$\int_0^4 p_n(x) p_m(x) d\mu(x) = \mathbf{1}_{n=m}. \quad (31.4)$$

A triangular monic change of polynomial basis has determinant one. Consequently the unweighted size- N Catalan Hankel determinant is one.

31.2 The determinant formula and its sign

For variables $x_1, \dots, x_N$ set $\Delta_N(x) = \prod_{i < j} (x_j - x_i)$. Expansion of two determinants and integration term by term gives

$$\det \left(\int x^{i+j} q(x) d\mu(x) \right)_{0 \leq i, j < N} = \frac{1}{N!} \int \Delta_N(x)^2 \prod_{j=1}^N q(x_j) d\mu(x_j). \quad (31.5)$$

This is the usual Andréief identity in the particular form needed here. It follows directly because the double sum over permutations contains $N!$ equal copies of the determinant on the left.

Take distinct $a_1, \dots, a_t$ and $q(x) = \prod_{i=1}^t (x - a_i)$. The mixed Vandermonde obeys

$$\Delta_{N+t}(x, a) = \Delta_N(x) \Delta_t(a) \prod_{j=1}^N \prod_{i=1}^t (a_i - x_j). \quad (31.6)$$

Replace its monomial columns by the monic $p_0, \dots, p_{N+t-1}$ without changing the determinant. In its Laplace expansion along the first N rows, integrate against $\det(p_{i-1}(x_j))_{j,i=1}^N$. Orthogonality kills every choice of N columns other than $p_0, \dots, p_{N-1}$. That choice contributes $N!$ times the remaining evaluation determinant, with positive Laplace sign. Combining (31.5) and (31.6) yields

$$\det \left(\int x^{i+j} \prod_{u=1}^t (x - a_u) d\mu(x) \right) = (-1)^{Nt} \frac{\det(p_{N+j}(a_i))_{1 \leq i \leq t, 0 \leq j < t}}{\Delta_t(a)}. \quad (31.7)$$

The sign is caused by $x - a$ in the multiplier and $a - x$ in the mixed Vandermonde. The left side is polynomial in all a_i , so the quotient has a well-defined confluent limit.

Lemma 31.1 (Confluent Christoffel identity). *For every $N, t \in \mathbb{N}_0$ and every $c \in \mathbb{C}$,*

$$H_N^{(t)}(c) = (-1)^{Nt} \det_{0 \leq r, j < t} \left(\frac{p_{N+j}^{(r)}(c)}{r!} \right). \quad (31.8)$$

Proof. In (31.7), let $a_i = c + \varepsilon_i$ and expand each evaluation row by Taylor's formula. The least possible distinct Taylor orders are $0, 1, \dots, t-1$. Their determinant is $\Delta_t(\varepsilon)$ times the determinant of the normalized jets $p_{N+j}^{(r)}(c)/r!$. All higher Taylor terms vanish after division by the Vandermonde and passage to the common limit. This proves the formula, including the empty determinant cases. $\square$

No division by $H_N^{(t)}(c)$ occurred. Some of those weighted determinants can be zero; the proof and the classification remain valid.

[Merge note: Lemma 31.1 is Part II's (15.8) for $q = (x - c)^t$, derived here a second time; Part I's Lemma 4.1 and Proposition 4.2 are further special cases, and the measure and orthogonality above repeat Part I's Section 3 and Part II's Section 15. The recursion (31.2) is Part II's (15.1). In (31.7) the manuscript prints the multiplier as $\prod_{a=1}^t (x - a_a)$; the product index is written u here (as in Part II), since i is a row index.]

32 An exact subset expansion for every sector

Fix $t \geq 1$ and abbreviate

$$T = \binom{t}{2}, \quad A_u = \prod_{j=0}^{u-1} j!, \quad K_t(z) = \frac{(z/(z^2 - 1))^T}{(z - 1)^t A_t}. \quad (32.1)$$

For a subset $S \subseteq J := \{0, \dots, t-1\}$ define

$$\sigma(S) = \sum_{j \in S} j, \quad \Delta(S) = \prod_{\substack{i, j \in S \\ i < j}} (j - i), \quad S^c = J \setminus S. \quad (32.2)$$

Empty products equal one.

Theorem 32.1 (Canonical sector formula). *For $0 \leq k \leq t$, put $h = t - k$ and define*

$$F_{t,k}(N; z) = \sum_{\substack{S \subseteq J \\ |S|=k}} (-1)^{T+h-\sigma(S)} z^{2\sigma(S)+k-T} \Delta(S) \Delta(S^c) \prod_{\substack{i \in S \\ j \in S^c}} (2N + i + j + 1), \quad (32.3)$$

$$P_{t,k}(N; z) = K_t(z) F_{t,k}(N; z). \quad (32.4)$$

Then the all-index identity

$$H_N^{(t)}(c) = \sum_{k=0}^t P_{t,k}(N; z) \lambda_k^N \quad (N \geq 0) \quad (32.5)$$

holds. Each $P_{t,k}$ is polynomial in N of degree at most $k(t - k)$. The definition remains meaningful when the numerical rates coincide.

Proof. Locally write $z = e^u$ and $x = 2 + 2 \cosh u$. Formula (31.2) gives

$$p_n(x) = \frac{z^{n+1} - z^{-n}}{z - 1} = g(u) (e^{(n+1/2)u} - e^{-(n+1/2)u}), \quad g(u) = \frac{1}{z^{1/2} - z^{-1/2}}. \quad (32.6)$$

Also $du/dx = z/(z^2 - 1)$. For a Wronskian of t functions, change of variable multiplies its determinant by $(du/dx)^T$: row r has leading derivative term $(du/dx)^r \partial_u^r$, and all other terms are combinations of earlier

rows. A common factor g in all functions similarly multiplies the Wronskian by g^t , by the Leibniz rule and a triangular row transformation. The normalization of row r by $r!$ contributes A_t^{-1} .

Set $v_j = N + j + \frac{1}{2}$. After these row transformations the Wronskian in (31.8) has columns

$$(v_j^r e^{v_j u} - (-v_j)^r e^{-v_j u})_{r=0}^{t-1}.$$

Expand each column by its two summands. Let S be the plus columns and S^c the minus columns. The exponential factor is $z^{(2k-t)N+2\sigma(S)-T+(2k-t)/2}$. The remaining determinant is the Vandermonde in the signed v_j . Within each sign block its magnitude is $\Delta(S)\Delta(S^c)$; between the blocks it contributes $\prod_{i \in S, j \in S^c} (v_i + v_j)$.

There are h signs from choosing the minus summand, $\binom{h}{2}$ from pairs of minus columns, and one for each plus column preceding a minus column. The number of the latter pairs is $k(t-1) - \sigma(S) - \binom{k}{2}$. Their total is exactly $T + h - \sigma(S)$. Finally $g(u)^t = z^{t/2}/(z-1)^t$ cancels all half-integral powers, leaving $z^{2\sigma(S)+k-T}$ outside $z^{(2k-t)N}$. The sign $(-1)^{Nt}$ in the Christoffel identity is absorbed into λ_k^N . This proves (32.3)–(32.5). There are kh linear factors in each product, giving the stated bound. $\square$

Remark 32.2 (No branch choice in the final formula). The auxiliary local logarithm and square root in the proof disappear from (32.3). Only integer powers of z remain. Since the result is a rational identity in z , valid off $0, \pm 1$, no global logarithm or continuation convention is required.

This expression is useful computationally: $F_{t,k}$ has Laurent polynomial coefficients in z with integer coefficients. In a cyclotomic integer ring it can be evaluated without any division. The common nonzero factor $K_t(z)$ has no effect on degrees or on cancellation among sectors.

33 The first two sector coefficients and reflection

The exact degree law alone is inherited from the preceding repository work. The additional coefficient calculated here distinguishes a one-degree loss from an arbitrarily long cancellation. [Merge note: the degree law is Part II's Theorem 14.2; Theorem 33.2 below re-proves it for one root, and its leading coefficient (33.3) is the one-root case of Part II's Theorem 18.1 (Section 28.5).]

33.1 A signed finite Vandermonde identity

Lemma 33.1. *For $h = t - k$ and an indeterminate q ,*

$$W_{t,k}(q) := \sum_{\substack{S \subseteq J \\ |S|=k}} (-1)^{\sigma(S)} \Delta(S) \Delta(S^c) q^{\sigma(S)} = (-1)^{\binom{k}{2}} A_k A_h q^{\binom{k}{2}} (1-q)^{kh}. \quad (33.1)$$

Proof. Take the $t \times t$ matrix with rows indexed by $j = 0, \dots, t-1$. Its first k columns are $j^a q^j$, $0 \leq a < k$, and its last h columns are j^b , $0 \leq b < h$. The convention $j^0 = 1$ includes $j = 0$. Laplace expansion along the first k columns gives the left side of (33.1) multiplied by $(-1)^{\binom{k}{2}}$.

To compute the same determinant, start with columns of derivatives of w^j at the two nodes $q, 1$. The ordinary confluent Vandermonde is $A_k A_h (1-q)^{kh}$. Replacing ∂_w^a by $(w \partial_w)^a$ is triangular in the derivative order and multiplies the determinant by $q^{0+1+\dots+(k-1)}$ in the first block and one in the second. Its columns are exactly those of our matrix. Equating the two evaluations proves the identity. $\square$

For $q \neq 0, 1$, logarithmic differentiation gives

$$\frac{q W'_{t,k}(q)}{W_{t,k}(q)} = \binom{k}{2} - \frac{khq}{1-q}. \quad (33.2)$$

This is an algebraic weighted mean of $\sigma(S)$, not a positivity or probability assertion. The weights have signs and can be complex.

33.2 Leading coefficient and the decisive correction

Theorem 33.2 (Exact degree and first centered correction). *Let $h = t - k$, $d = kh$, and $w = N + t/2$. The leading coefficient of $P_{t,k}$ is*

$$\ell_{t,k}(z) = (-1)^{h(h+1)/2} \frac{2^{kh} A_k A_h z^{k^2}}{A_t (z-1)^t (z^2-1)^{\binom{k}{2} + \binom{h}{2}}}, \quad \ell_{t,k}(z) \neq 0. \quad (33.3)$$

In particular $\deg_N P_{t,k} = kh$. If $d \geq 1$, then

$$P_{t,k}(N; z) = \ell_{t,k}(z) \left(w^d + \frac{kh(h-k)(z^2+1)}{4(z^2-1)} w^{d-1} \right) + Q_{t,k}(w), \quad \deg Q_{t,k} \leq d-2. \quad (33.4)$$

A polynomial of degree at most -1 means zero. If $d = 0$, the sector is the constant $\ell_{t,k}(z)$.

Proof. Every product in (32.3) has leading coefficient 2^d . Its weighted sum is, by (33.1),

$$[N^d]F_{t,k} = (-1)^{T+h+\binom{k}{2}} 2^d A_k A_h z^{k^2-T} (1-z^2)^d.$$

Multiply by $K_t(z)$. Since $T = \binom{k}{2} + \binom{h}{2} + kh$, collecting the signs and powers gives exactly (33.3). All factors are nonzero under the standing hypotheses, proving the exact degree.

For the next coefficient, rewrite the mixed factor as

$$2N + i + j + 1 = 2w + i + j + 1 - t.$$

For a fixed subset S , the sum of its d offsets equals

$$\begin{aligned} B(S) &= \sum_{i \in S, j \in S^c} (i + j + 1 - t) \\ &= (h-k)\sigma(S) + kT - kh(t-1). \end{aligned} \quad (33.5)$$

Indeed $\sum_{j \in S^c} j = T - \sigma(S)$, which proves the second line by direct summation. The relative coefficient of w^{d-1} in this product is $B(S)/2$.

The weighted sum uses the same weights as $W_{t,k}(z^2)$. By (33.2), its mean offset is

$$\begin{aligned} \bar{B} &= (h-k) \left(\binom{k}{2} - \frac{khq}{1-q} \right) + kT - kh(t-1) \\ &= \frac{kh(h-k)(q+1)}{2(q-1)}, \quad q = z^2. \end{aligned} \quad (33.6)$$

Dividing by two gives the coefficient in (33.4). Only lower powers remain, proving the assertion. $\square$

It is useful to isolate the correction as

$$b_{t,k}(z) = \frac{k(t-k)(t-2k)(z^2+1)}{4(z^2-1)}. \quad (33.7)$$

Then $b_{t,t-k} = -b_{t,k}$. Except for the trivial factors $k = 0, t$ or $k = t/2$, its only zero away from the endpoints is $z^2 = -1$, namely $c = 2$.

33.3 An exact reflection of the whole sector polynomial

Proposition 33.3 (Centered sector reflection). *For every $0 \leq k \leq t$, with $h = t - k$,*

$$P_{t,h}(N; z) = (-1)^{t(t+1)/2} z^{(h-k)t} P_{t,k}(-N - t; z). \quad (33.8)$$

Consequently,

$$\frac{\ell_{t,h}(z)}{\ell_{t,k}(z)} = (-1)^{(h-k)(t+1)/2} z^{(h-k)t}. \quad (33.9)$$

The exponent in the sign is always an integer.

Proof. In the subset formula replace S by the complement of $\{t - 1 - j : j \in S\}$. This new subset has size h and sum $T - k(t - 1) + \sigma(S)$. Its two Vandermonde magnitudes are unchanged, its power of z gains $(h - k)t$, and replacing N by $-N - t$ changes the sign of every one of the kh mixed linear factors. The remaining sign simplifies to $(-1)^{t(t+1)/2}$. This is a termwise bijection of the two sums and proves (33.8). Alternatively the same reflection comes from $p_{-n-1} = -p_n$ and reversal of the columns of the Wronskian.

Comparison of the leading coefficients gives $(-1)^{t(t+1)/2+kh} z^{(h-k)t}$. Its sign equals the one in (33.9); the difference of the two exponents is even. One can also obtain (33.9) immediately by dividing the products in (33.3). $\square$

The center $-t/2$ is therefore natural, rather than a convenient numerical fit. At a colliding symmetric pair, the two leading terms are either equal or opposite, and the first centered corrections have opposite relative signs. That is the algebraic source of the classification.

34 Why the noncentral cancellation has exactly one defect

34.1 Minimal annihilators of exponential polynomials

Lemma 34.1. *Suppose $f(N) = \sum_{j=1}^b Q_j(N) \alpha_j^N$, where the α_j are distinct and nonzero and each Q_j is a nonzero polynomial of degree d_j . The minimal annihilator of f is*

$$\prod_{j=1}^b (X - \alpha_j)^{d_j+1}. \quad (34.1)$$

The same conclusion holds for eventual annihilators. Its generating function has reduced denominator $\prod_j (1 - \alpha_j u)^{d_j+1}$.

Proof. On $Q(N) \alpha^N$, the operator $E - \alpha$, where $(Ef)(N) = f(N + 1)$, gives $\alpha^{N+1}(Q(N + 1) - Q(N))$. It lowers the polynomial degree by exactly one until the term becomes zero. In contrast, $E - \beta$ with $\beta \neq \alpha$ preserves the degree and has a nonzero leading coefficient. Distinct exponential blocks are independent: applying the annihilators of all but one block isolates a nonzero polynomial multiple of the remaining exponential. A nonzero polynomial cannot vanish at every sufficiently large integer. It follows that any annihilator, even an eventual one, must contain every factor in (34.1) to the displayed multiplicity. Their product clearly suffices.

For generating functions, $(u d/du)^a (1 - \alpha u)^{-1}$ is the series of $N^a \alpha^N$. A polynomial multiplier of degree d therefore gives a pole of exact order $d + 1$ at $u = \alpha^{-1}$. Distinct poles cannot cancel. This proves the denominator assertion. $\square$

[Merge note: Lemma 34.1 restates Part I's Lemma 6.1 and Part II's Lemma 19.1; the manuscript's proof is kept as a third proof.]

The nonresonant case of Theorem 30.1 now follows from Theorems 32.1 and 33.2. It remains to determine the degree of each grouped polynomial

$$G_r(N) = \sum_{k \in I_r} P_{t,k}(N; z). \quad (34.2)$$

34.2 The geometry of a residue class

Lemma 34.2 (The unique possible top tie). *Assume $2 \leq L \leq t$. In every class I_r , the maximum of $k(t - k)$ has a unique maximizer except possibly in one class. A tied class exists exactly when $t \equiv L \pmod{2}$. Its two maximizers are k_-, k_+ from (30.7); they are consecutive members of that class, separated by L , and*

$$d_* := k_-(t - k_-) = k_+(t - k_+) = (t^2 - L^2)/4. \quad (34.3)$$

Any other index in that class has degree at most $d_* - 2L^2$.

Proof. The identity

$$k(t - k) = \frac{t^2 - (2k - t)^2}{4} \quad (34.4)$$

shows that one must choose the members of the arithmetic progression I_r nearest $t/2$. There are at most two. A tie requires two consecutive members to be symmetric about $t/2$, so their difference is L and their sum is t . They are consequently the two indices in (30.7), and exist exactly under the parity condition. Their values $2k - t$ are $-L, L$. Any further index in that same class has $|2k - t| \geq 3L$, so (34.4) decreases its degree by at least $((3L)^2 - L^2)/4 = 2L^2$. $\square$

A group with a unique maximizer has degree d_r , since its unique leading coefficient is nonzero by (33.3). Thus the only remaining issue is the one tied group.

34.3 The tied group and the nonvanishing next term

Assume the parity condition and use $k = k_-$, $h = k_+$, so $h - k = L$. By (33.9), the ratio of their leading coefficients is

$$\rho = (-1)^{L(t+1)/2} z^{Lt}. \quad (34.5)$$

Since $(z^L)^2 = 1$, this is either 1 or -1 . When $\rho = 1$, the leading coefficient of G_{r_*} is $2\ell_{t,k} \neq 0$ and there is no defect.

Suppose $\rho = -1$. If $d_* = 0$, then $t = L$, the two indices are $0, t$, and they are the only indices in their class. Their sectors are opposite constants, so $G_{r_*} = 0$. The class contributes no characteristic factor rather than the naive exponent one.

If $d_* > 0$, apply (33.4) in the centered variable $w = N + t/2$. The leading terms cancel, while the coefficient of w^{d_*-1} in their sum is

$$\ell_{t,k} b_{t,k} + (-\ell_{t,k})(-b_{t,k}) = \frac{\ell_{t,k} d_* L (z^2 + 1)}{2(z^2 - 1)}. \quad (34.6)$$

It is nonzero when $c \neq 2$. The lower members of the same residue class cannot affect this coefficient, by Lemma 34.2. Therefore $\deg G_{r_*} = d_* - 1$ exactly.

We have now proved all the exponents in (30.9), including the disappearing constant case. Applying Lemma 34.1 proves their minimality and the generating-function assertion. The total order formula will follow from a finite sum of squares.

Corollary 34.3 (Only a real characteristic root can lose multiplicity). *Whenever the noncentral defect occurs, its characteristic root is*

$$\lambda_* = (-1)^t z^{-L} = (-1)^t z^L \in \{1, -1\}. \quad (34.7)$$

No other characteristic root loses multiplicity beyond grouping.

Proof. Substitute $2k_- - t = -L$ into (30.4) and use $z^{2L} = 1$. The degree proof above excludes every other group. $\square$

35 Closed orders, exceptional parameters, and reflection

35.1 Summing the grouped degrees

For each residue r , select a nearest centered value $a_r = 2k_- - t$ at a maximizer. In a tie either choice gives the same square. When t and L have different parity, their squares are the squares of the complete list

$$-(L-1), -(L-3), \dots, L-3, L-1.$$

When the parities agree, they are the squares of

$$-L, -L+2, \dots, L-4, L-2.$$

The finite interval $[-t, t]$ contains all chosen representatives because $t \geq L$. Summing squares in these arithmetic progressions gives

$$\sum_{r=0}^{L-1} a_r^2 = \frac{L(L^2-1)}{3} + L\eta. \quad (35.1)$$

Using (34.4),

$$\begin{aligned} R_{\text{grp}}(t, L) &= \sum_{r=0}^{L-1} (d_r + 1) = L + \frac{Lt^2 - \sum_r a_r^2}{4} \\ &= \frac{L(3t^2 - L^2 + 13 - 3\eta)}{12}. \end{aligned} \quad (35.2)$$

The loss established in the preceding section is exactly δ , proving (30.10).

For completeness, Corollary 30.3 follows without any complex arithmetic. If t, L are even, $z^{Lt} = 1$ and the parity of $L(t+1)/2$ is that of $L/2$. If both are odd, $z^{Lt} = z^L$ and the condition $\rho = -1$ is equivalent to $z^L = (-1)^{(t-1)/2}$. This proves all the sign cases.

35.2 An exact finite exceptional locus

Corollary 35.1 (Every nongeneric nonendpoint parameter). *For fixed $t \geq 1$, the parameters $c \neq 0, 4$ at which $R(t, c) < R_{\text{gen}}(t)$ are exactly*

$$\mathcal{E}_t = \left\{ 2 + 2 \cos \frac{\pi a}{L} : 2 \leq L \leq t, \quad 1 \leq a < L, \quad \gcd(a, L) = 1 \right\}. \quad (35.3)$$

In particular all exceptional parameters are real and lie in $(0, 4)$, and

$$|\mathcal{E}_t| = \sum_{L=2}^t \varphi(L), \quad (35.4)$$

where $\varphi(L)$ is the number of integers $1 \leq a < L$ coprime to L . At the parameter represented by a/L , one has $z^L = (-1)^a$.

Proof. A collision of two rates means $z^{2(k-j)} = 1$ with $1 \leq |k-j| \leq t$. This is equivalent to finite $L \leq t$. Then z is a root of unity, so $c = 2 + z + z^{-1}$ is real in $(0, 4)$. Choose the reciprocal of z as needed to write $z = e^{\pi i a/L}$ with $1 \leq a < L$ and $\gcd(a, L) = 1$. Conversely any such parameter has exactly the stated order L .

At a collision, grouping replaces at least two positive contributions $d_k + 1$ by at most their maximum, strictly decreasing the total generic order. Further cancellations cannot undo that decrease. In the absence of collisions, the nonzero leading coefficients give the full generic order, including $t = 1, c = 2$. Finally $\cos \theta$ is injective on $(0, \pi)$, and the fractions a/L in (35.3) are reduced. No two listed parameters coincide. Counting them proves (35.4). $\square$

Thus the algebraic position of a root, not its numerical proximity to another parameter, determines whether an exact recurrence drops order. There is no open interval of exceptional values.

35.3 The missing sign datum and reflection of the root

Corollary 35.2 (Order under $c \mapsto 4 - c$). *For $c \neq 0, 2, 4$, the orders at c and $4 - c$ differ exactly when $L \leq t$ and L, t are both odd. In that case they differ by one:*

$$R(t, 4 - c) - R(t, c) = 2\delta - 1. \quad (35.5)$$

In every other case they are equal.

Proof. The replacement $z \mapsto -z$ sends c to $4 - c$ without changing z^2 or L . For odd L it changes the sign of z^L ; for even L it does not. Corollary 30.3 shows that the two defects are complementary precisely when t, L are odd and $L \leq t$. Their grouped orders are identical. In the nonresonant case both orders are generic. $\square$

The example $t = L = 3, z = e^{2\pi i/3}$ versus $z = e^{\pi i/3}$, is the smallest instance. The two input roots are $c = 1$ and $c = 3$. The orders 7 and 6 show that t and $\text{ord}(z^2)$ alone cannot parametrize the answer. The additional sign is invariant under the allowed choice $z \leftrightarrow z^{-1}$, so it is genuine information about c , not an artifact of a branch convention.

35.4 Quadratic quasipolynomial growth in the multiplicity

Corollary 35.3. *Fix a noncentral cyclotomic parameter with $L \geq 3$. For $t \geq L$, $R(t, c)$ is a quadratic quasipolynomial in t , with period dividing four and leading coefficient $L/4$. In particular,*

$$R(t, c) = \frac{L}{4}t^2 + O_L(1), \quad (35.6)$$

whereas the nonresonant order is $t^3/6 + O(t)$.

Proof. In (30.10), η depends only on parity and δ depends only on t modulo four once L and z^L are fixed. The remaining term is $Lt^2/4$ plus a constant depending on L . Equation (29.1) gives the comparison. $\square$

Here the asymptotic variable is the root multiplicity t , not the Hankel size N . This distinction matters: the determinant itself is an exact quasipolynomial in N at a cyclotomic parameter, but its recurrence order is a different function.

36 The central exception and the known product

36.1 Reduction to middle-binomial moments

Set $y = x - 2$ in (31.1). Then

$$d\mu = (1 - y/2) d\alpha(y), \quad d\alpha(y) = \frac{dy}{\pi\sqrt{4 - y^2}}, \quad -2 < y < 2. \quad (36.1)$$

The arcsine moments are zero in odd degree and $\int y^{2r} d\alpha = \binom{2r}{r}$, as follows from $y = 2 \cos \theta$. Consequently the centered Catalan moments are

$$\mu_n := \int (x - 2)^n d\mu(x) = (-1)^n \binom{n}{\lfloor n/2 \rfloor}. \quad (36.2)$$

Indeed the odd moment is $-\frac{1}{2} \binom{2r+2}{r+1} = -\binom{2r+1}{r}$.

A monic translation of the polynomial basis leaves the determinant unchanged, giving $H_N^{(t)}(2) = \det(\mu_{i+j+t})$. If

$$D_t(N) = \det_{0 \leq i, j < N} \left(\binom{i+j+t}{\lfloor (i+j+t)/2 \rfloor} \right),$$

row and column signs yield

$$H_N^{(t)}(2) = (-1)^{Nt} D_t(N). \quad (36.3)$$

Cigler's Theorem 1 [9] states $D_t(N) = (-1)^t \binom{N}{2} B_{\lfloor t/2 \rfloor, \lceil t/2 \rceil}(N)$. Equation (36.3) is therefore precisely (30.13)–(30.14). In particular, the uniform central formula requested by the repository [Section 24.1] is already a result established in a 2021 preprint after this change of normalization.

36.2 A self-contained condensation proof

We include the proof to keep the classification independent of an external determinant evaluation. It uses the same classical condensation mechanism as Cigler's argument, not a new method. Write $h_N^{(t)} = \det(\mu_{i+j+t})$ and set $h_0^{(t)} = 1$. The Desnanot–Jacobi determinant identity gives, for $N \geq 2$,

$$h_N^{(t)} h_{N-2}^{(t+2)} = h_{N-1}^{(t)} h_{N-1}^{(t+2)} - (h_{N-1}^{(t+1)})^2. \quad (36.4)$$

It is a polynomial determinant identity and does not require any minor to be nonzero.

We verify the proposed products in (30.13) and (30.14), and then induct on N simultaneously for all $t \geq 0$. At $N = 0$ the products equal one. At $N = 1$, cancellation in (30.12) gives

$$B_{a,b}(1) = \prod_{i=1}^a \frac{i+b}{i} = \binom{a+b}{a}, \quad (36.5)$$

so the products give exactly the moments (36.2).

For the induction put $n = N - 1 \geq 1$. Four ratios suffice. Direct cancellation of the rectangular products yields

$$\frac{B_{m,m}(n) B_{m+1,m+1}(n)}{B_{m,m+1}(n)^2} = \frac{n+2m+1}{2m+1}, \quad (36.6)$$

$$\frac{B_{m,m}(n+1) B_{m+1,m+1}(n-1)}{B_{m,m+1}(n)^2} = \frac{n}{2m+1}, \quad (36.7)$$

$$\frac{B_{m,m+1}(n) B_{m+1,m+2}(n)}{B_{m+1,m+1}(n)^2} = \frac{n+2m+2}{2(n+m+1)}, \quad (36.8)$$

$$\frac{B_{m,m+1}(n+1) B_{m+1,m+2}(n-1)}{B_{m+1,m+1}(n)^2} = \frac{n}{2(n+m+1)}. \quad (36.9)$$

For example, adding the last column of a rectangle gives

$$\frac{B_{a,b+1}(n)}{B_{a,b}(n)} = \prod_{i=1}^a \frac{n+i+b}{i+b}, \quad (36.10)$$

and increasing n telescopes within each row:

$$\frac{B_{a,b}(n+1)}{B_{a,b}(n)} = \prod_{i=1}^a \frac{n+i+b}{n+i}. \quad (36.11)$$

Applying (36.10) and (36.11), or their inverses, to the four ratios cancels all interior factors and gives the displayed right sides. These identities also cover $m = 0$.

For even $t = 2m$, divide the product version of (36.4) by $B_{m,m+1}(n)^2$. The squared odd sign is one, and the identity becomes

$$\frac{n}{2m+1} = \frac{n+2m+1}{2m+1} - 1.$$

For odd $t = 2m+1$, write $s_N = (-1)^{N(N+1)/2}$. One has $s_N s_{N-2} = -1$ and $s_{N-1}^2 = 1$. Division by $B_{m+1,m+1}(n)^2$ reduces the same identity to

$$-\frac{n}{2(n+m+1)} = \frac{n+2m+2}{2(n+m+1)} - 1.$$

Thus the proposed products satisfy the condensation equation.

By the induction hypothesis, the actual denominator $h_{N-2}^{(t+2)}$ in solving (36.4) for $h_N^{(t)}$ is the proposed signed product, which is nonzero at nonnegative integer $N-2$. Solving is therefore legitimate. It proves the product at size N for every t , completing the induction. The nonzero denominator is a conclusion established earlier in this induction, not an assumption about arbitrary weighted Hankel minors.

36.3 Minimality and why the one-defect law excludes this point

The product $B_{a,b}(N)$ has degree ab and nonzero leading coefficient. The even product therefore has minimal polynomial $(X-1)^{m^2+1}$ by Lemma 34.1. In the odd case,

$$s_N = \frac{1+i}{2} i^N + \frac{1-i}{2} (-i)^N. \quad (36.12)$$

Both coefficients are nonzero. Multiplication by $B_{m,m+1}(N)$ gives both characteristic roots $i, -i$ with multiplicity $m(m+1)+1$, proving Theorem 30.4 in full.

For $t = 2m \geq 2$, the two grouped rates at $z = i$ are $1, -1$. Their uncorrected exponents would be $m^2 + 1$ and m^2 , so $R_{\text{grp}} = 2m^2 + 1$. The exact product has only the rate 1: the entire -1 group vanishes, and the actual order is $m^2 + 1$. The additional loss is m^2 , not one. Formula (34.6) correctly signals the problem because $z^2 + 1 = 0$ here.

For odd t , $L = 2$ has different parity from t , so there is no tied maximum at all. The unique-maximum argument already gives the correct odd central order. The product evaluation provides much more: all values and the exact period-four sign.

37 A root-free minimal-recurrence algorithm

The classification need not be implemented by factoring the quadratic for z , recognizing a root of unity, or computing Hankel determinants. There is a direct exact algorithm over the coefficient field of c .

37.1 Detecting a resonance by a short scalar recurrence

Put $v = c - 2$ and define

$$U_0 = 1, \quad U_1 = v, \quad U_{a+1} = vU_a - U_{a-1}. \quad (37.1)$$

These are scalar values of the Chebyshev polynomials of the second kind, $U_a(v/2)$; we suppress the argument to emphasize computation. From the recurrence,

$$U_a = \frac{z^{a+1} - z^{-a-1}}{z - z^{-1}}. \quad (37.2)$$

Since $z \neq \pm 1$, the first zero $U_{L-1} = 0$ occurs exactly at $L = \text{ord}(z^2)$, if that order is finite. Moreover,

$$\sigma := z^L = -U_{L-2} \in \{1, -1\}. \quad (37.3)$$

Indeed $U_L = (z^{L+1} - z^{-L-1})/(z - z^{-1}) = z^L$, while the recurrence at the zero gives $U_L = -U_{L-2}$.

Testing $U_1, \dots, U_{t-1}$ therefore decides whether the recurrence is resonant at the given multiplicity. There is no need to decide whether z has some much larger finite order: all $L > t$ have the same nonresonant formula. This requires $O(t)$ exact field operations and zero tests.

37.2 Linear and quadratic factors in the original coefficient field

Define another scalar sequence

$$S_0 = 2, \quad S_1 = v, \quad S_{a+1} = vS_a - S_{a-1}, \quad S_a = z^a + z^{-a}, \quad (37.4)$$

and set

$$Q_a(X) = X^2 - (-1)^t S_a X + 1, \quad E_a = \frac{t^2 - a^2}{4} + 1 \quad (a \equiv t \pmod{2}). \quad (37.5)$$

These factors have coefficients in $\mathbb{Q}(c)$, or in any characteristic-zero coefficient field containing c .

Theorem 37.1 (Root-free factorization). *For $c \neq 0, 4$, test (37.1) through U_{t-1} . If no zero occurs, then*

$$\chi_{t,c}(X) = (X - 1)^{\mathbf{1}_{2|t}(t^2/4+1)} \prod_{\substack{1 \leq a \leq t \\ a \equiv t \pmod{2}}} Q_a(X)^{E_a}. \quad (37.6)$$

The first factor is omitted when t is odd. If the first zero occurs at $L - 1$ with $L = 2$, use (30.15)–(30.16). If $3 \leq L \leq t$, compute σ by (37.3) and δ by (30.8), replacing z^{Lt} by σ^t . Then

$$\chi_{t,c}(X) = (X - 1)^{\mathbf{1}_{2|t}(t^2/4+1)} \prod_{\substack{1 \leq a < L \\ a \equiv t \pmod{2}}} Q_a(X)^{E_a} \cdot (X - (-1)^t \sigma)^{\eta(E_L - \delta)}. \quad (37.7)$$

The last factor is omitted when $\eta = 0$ or its exponent is zero; E_L is only used when $\eta = 1$. The nonconstant displayed factors have pairwise disjoint roots, and each displayed quadratic has distinct roots.

Proof. In the nonresonant case, pair the sectors $k, t - k$ and put $a = |2k - t|$. Their rates are $(-1)^t z^a$ and $(-1)^t z^{-a}$, whose quadratic is (37.5). Their common exponent is E_a . When t is even, the unpaired sector $k = t/2$ has rate one and exponent $t^2/4 + 1$. Distinctness follows from $L > t$.

In the resonant case, use the centered representatives in Section 35. The pairs $\pm a$ with $0 < a < L$ give exactly the quadratics in (37.7). The possible center $a = 0$ gives the first linear factor. The tied endpoints $a = \pm L$ occur exactly when $\eta = 1$ and give the single rate $(-1)^t \sigma$, with its one-defect exponent. No two different interior pairs coincide: equality of rates would force their corresponding k indices to agree modulo L , contrary to the choice of distinct centered representatives. An interior quadratic cannot have a double root because that would require $z^{2a} = 1$ with $0 < a < L$. If both linear factors occur, L is even, whence $z^L = -1$; they are therefore also distinct. Theorem 30.1 supplies minimality. $\square$

The theorem outputs $O(t)$ linear or quadratic factors with integer exponents after $O(t)$ scalar field operations. This is an arithmetic operation count for the *factored* answer; it is not a bit-complexity bound and not the cost of expanding a degree- $\Theta(t^3)$ polynomial. Exact equality tests in the input field are essential. Approximate floating-point tests cannot reliably detect an exact resonance.

This improves the task-specific route considerably over building a long prefix of large Hankel determinants and solving for an unknown recurrence. It does not replace the repository's more general algorithm for a multiplier with several different roots [Theorem 20.2 with the bound (20.6)].

38 Worked examples and a complete rational-parameter corollary

38.1 Two cubic multipliers with the same cyclotomic order

For $t = 3, c = 1$, choose $z = e^{2\pi i/3}$. Then $L = 3, z^L = 1$, and $\delta = 0$. The tied constant sectors retain the rate -1 . For $t = 3, c = 3$, choose $z = e^{\pi i/3}$. Then $L = 3, z^L = -1$, and $\delta = 1$. The tied constant sectors at rate 1 cancel entirely. The root-free factors are particularly simple:

$$\chi_{3,1}(X) = (X + 1)(X^2 - X + 1)^3, \quad R(3, 1) = 7, \quad (38.1)$$

$$\chi_{3,3}(X) = (X^2 + X + 1)^3, \quad R(3, 3) = 6. \quad (38.2)$$

For both inputs the generic order would be eight.

There are concise formulas for the determinants themselves. For $n \geq 0$,

$$\begin{aligned} H_{3n}^{(3)}(3) &= 2n + 1, \\ H_{3n+1}^{(3)}(3) &= -(n + 1)(18n + 13), \\ H_{3n+2}^{(3)}(3) &= (n + 1)(18n + 23). \end{aligned} \quad (38.3)$$

For $c = 1$, the six formulas are

r	$H_{6n+r}^{(3)}(1)$	$n = 0$	$n = 1$	$n = 2$
0	$(4n + 1)^2$	1	25	81
1	$(2n + 1)(4n + 1)$	1	15	45
2	$-(2n + 1)(4n + 3)$	-3	-21	-55
3	$-(4n + 3)^2$	-9	-49	-121
4	$-2(n + 1)(4n + 3)$	-6	-28	-66
5	$2(n + 1)(4n + 5)$	10	36	78

These are exact identities, not extrapolations. The sector theorem shows that each displayed progression is a polynomial of degree at most two: the rates have periods three and six, respectively, and all sector degrees are at most two. The first three values in each progression determine its quadratic. Evaluating them from (31.8) gives the displayed formulas. The accompanying code additionally verifies each identity through $N = 59$.

For reference, their first twelve values are

$$\begin{aligned} H_N^{(3)}(1) : & \quad 1, 1, -3, -9, -6, 10, 25, 15, -21, -49, -28, 36, \dots, \\ H_N^{(3)}(3) : & \quad 1, -13, 23, 3, -62, 82, 5, -147, 177, 7, -268, 308, \dots \end{aligned}$$

The difference between their minimal orders persists even though both are low-degree quasipolynomial sequences.

38.2 A noncentral loss below a positive degree

Take $c = 2 + \sqrt{3}$, $z = e^{\pi i/6}$, $L = 6$, and $t = 8$. The tied indices are 1, 7, of degree seven, at rate -1 . Since $L \equiv 2 \pmod{4}$, their leading terms cancel. Here $z^2 + 1 \neq 0$, so (34.6) proves that the grouped polynomial has degree six, not less. The full answer is

$$\chi_{8,2+\sqrt{3}}(X) = (X-1)^{17}(X+1)^7(X^2+X+1)^{13}(X^2-X+1)^{16}. \quad (38.4)$$

Its order is 82, compared with grouped order 83 and generic order 93. This example illustrates the genuinely positive-degree part of the one-defect theorem, rather than just cancellation of two constants. The same recurrence holds at $c = 2 - \sqrt{3}$.

38.3 All rational nonendpoint roots

Corollary 38.1. *If $c \in \mathbb{Q} \setminus \{0, 1, 2, 3, 4\}$, then $R(t, c) = R_{\text{gen}}(t)$ for every $t \geq 1$. For $c = 1$ and $c = 3$, $R(1, c) = 2$. For $t \geq 2$ their orders are*

$$\begin{array}{c|cc} & c = 1 & c = 3 \\ \hline t \text{ even} & 3t^2/4 + 1 & 3t^2/4 + 1 \\ t \text{ odd} & (3t^2 + 1)/4 - \mathbf{1}_{t \equiv 1 \pmod{4}} & (3t^2 + 1)/4 - \mathbf{1}_{t \equiv 3 \pmod{4}}. \end{array} \quad (38.5)$$

For $c = 2$, the exact orders are

$$R(t, 2) = \begin{cases} t^2/4 + 1, & t \text{ even}, \\ (t^2 + 3)/2, & t \text{ odd}. \end{cases} \quad (38.6)$$

Proof. If z is a root of unity, $z + z^{-1}$ is an algebraic integer. A rational algebraic integer is an integer: substitution of a reduced fraction into its monic integer equation forces its denominator to be one. Also $-2 < z + z^{-1} < 2$ when $c \neq 0, 4$ is real and cyclotomic. Thus the only rational possibilities are $c = 1, 2, 3$. Every other rational nonendpoint input is nonresonant. At $c = 1, 3$ use $L = 3$ and $z^L = 1, -1$, respectively, in (30.10). The case $t = 1$ precedes the resonant range and is stated separately. The central formula is Theorem 30.4. $\square$

t	$R(t, 1)$	$R(t, 2)$	$R(t, 3)$	$R_{\text{gen}}(t)$
1	2	2	2	2
2	4	2	4	4
3	7	6	6	8
4	13	5	13	15
5	18	14	19	26
6	28	10	28	42
7	37	26	36	64
8	49	17	49	93
9	60	42	61	130
10	76	26	76	176
11	91	62	90	232
12	109	37	109	299

All entries follow from the proved formulas. The machine-readable version extends through $t = 20$.

39 Exact verification and reproducibility

The universal results above follow from the displayed proofs. The verification suite is an independent finite audit of formulas, signs, normalizations, and algorithmic implementations. No assertion of universality is inferred merely from the number of successful tests.

39.1 Arithmetic models and independent routes

The program `code/verify.py` (shipped here as `code/03-cyclotomic-verify.py`) uses Python's standard library only. Cyclotomic arithmetic takes place in

$$\mathbb{Z}[z]/(\Phi_M(z)), \quad (39.1)$$

with coefficient vectors in the usual basis of degree less than $\deg \Phi_M$. The monic polynomials Φ_M are constructed by exact successive division of $x^M - 1$ by the previously constructed factors for its proper divisors. All arithmetic is integral; comparisons do not use tolerances or approximate complex roots. Negative powers of z are reduced using $z^M = 1$.

The raw sector formula (32.3) is evaluated as an exact polynomial in N . Its degree, leading coefficient, first centered correction, and reflected polynomial are compared with the theorems. The sectors are then added according to equal rates, and their actual polynomial degrees are compared with every predicted characteristic multiplicity. The root-free algorithm is tested separately: it does not access the cyclotomic order or the spectral generator when constructing its factors from c . Its factors are checked against the actual grouped rates and multiplicities.

The original small Hankel determinants in the same rings are computed by the Leibniz permutation formula, not by the sector determinant. To compare with the rational sector expression, the test clears the common denominator. Explicitly it checks

$$H_N^{(t)}(c)(z^2 - 1)^T(z - 1)^t A_t = (-1)^{Nt} z^T \sum_{k=0}^t F_{t,k}(N; z) z^{(2k-t)N}. \quad (39.2)$$

Thus no inversion in a number field is needed even in this independent comparison.

For integer c , a second route evaluates the original moment matrix by pivoted fraction-free Bareiss elimination and compares it with the $t \times t$ normalized-jet determinant. The jets are generated from (31.2): if $u_{n,r} = p_n^{(r)}(c)/r!$, then

$$u_{n+1,r} = (c - 2)u_{n,r} + u_{n,r-1} - u_{n-1,r}, \quad u_{n,-1} = 0, \quad (39.3)$$

with $u_{0,0} = 1$, $u_{1,0} = c - 1$, $u_{1,1} = 1$, and other initial jets zero. This is independent of the subset expansion.

39.2 Parameter ranges and the recorded run

The principal cyclotomic grid uses primitive orders $3 \leq M \leq 24$ and root multiplicities $1 \leq t \leq 12$. A computation in $\mathbb{Z}[z]/\Phi_M$ is exact at a primitive root and respects all its Galois conjugates, rather than sampling a floating-point angle. The original cyclotomic determinant comparisons use $M \leq 14$, $t \leq 8$, and sizes $N = 0, \dots, 4$.

The integer determinant comparison uses $c \in \{-2, 0, 1, 2, 3, 4, 5\}$, $t = 1, \dots, 10$, and $N = 0, \dots, 10$. The endpoint values here only audit the classical determinant identity; they are not applications of the nonendpoint classification. The central product is checked for $t = 0, \dots, 16$ and $N = 0, \dots, 16$. The scalar sign and square-sum identities are checked on an additional grid with $M \leq 60$ and $t \leq 100$.

Exact assertion category	Checks passed
Canonical sector degrees	1,980
Product leading coefficients	1,980
First centered corrections	1,452
Full sector reflections	1,980
Grouped minimal multiplicities	1,569
Closed total-order formula	264
Root-free detection, factor degrees, and multiplicities	2,097
Independent cyclotomic Hankel determinants	480
Original integer matrices versus jet determinants	770
Central product evaluations	289
Four condensation ratio identities	252
Centered residue-class square sums	4,495
Simplified defect signs	4,396
Worked residue-class formulas	120
Total	22,124

The run completed successfully. The individual counts and the parameter-by-parameter recurrence orders are stored as JSON, and the rational examples are also stored as CSV. A check is one explicit assertion as counted in the program; the number is not a measure of proof strength or a substitute for peer review.

[Merge note: the recorded run is shipped as `data/03-cyclotomic-verification_summary.json` (status PASS and the twenty counts behind the table; the table's condensation row adds four counts of 63, and its root-free row adds $264 + 264 + 1,569$) and `data/03-cyclotomic-verification_run.txt` (the console transcript); the parameter-by-parameter orders, the 264 principal cases with their multiplicities by residue class, as `data/03-cyclotomic-cyclotomic_orders.json`; and the rational examples, the orders at $c = 1, 2, 3$ and the generic order for $t = 1, \dots, 20$, as `data/03-cyclotomic-orders_c1_c2_c3.csv` (all-CRLF as written by Python's csv module). All four were delivered in `data/` without the prefix.]

39.3 Rebuilding the article and the audit

From the package directory, run

```
python3 code/verify.py
pdflatex -interaction=nonstopmode -halt-on-error article.tex
pdflatex -interaction=nonstopmode -halt-on-error article.tex
```

The first command regenerates the data files. A smaller main grid is available with `-quick`; it is explicitly marked as such in its output. The supplied full-run transcript is `data/verification_run.txt`. The $\text{\LaTeX}$ source uses ordinary TeX Live packages, including `newtxtext` and `newtxmath`. The source bibliography is embedded, so no separate bibliography processor or network access is needed.

In this report these commands must not be run in place. Here `code/verify.py` is Part I's verifier, and the shipped `code/03-cyclotomic-verify.py` writes by default into the report's `data/`, adding the three unprefixed files `verification_summary.json`, `cyclotomic_orders.json` and `orders_c1_c2_c3.csv` beside the shipped ones; the transcript is its console output, not a file it writes. The delivered Makefile, shipped as `code/03-cyclotomic-Makefile`, has the same defect when run from the report directory: its `verify` target runs Part I's `code/verify.py`, which rewrites Part I's data files; its `quick` target passes that verifier options it does not know, and its `pdf` and `clean` targets build and clean the

merged `article.tex` in place. The article is built as described in the report's README. Restore the delivered layout in a scratch copy instead (POSIX shell, run from the report directory):

```
W=/path/to/scratch/cyclotomic
mkdir -p "$W/code" "$W/recorded"
cp code/03-cyclotomic-verify.py "$W/code/verify.py"
for f in data/03-cyclotomic-*; do
  cp "$f" "$W/recorded/${f#data/03-cyclotomic-}"; done
cd "$W" && python3 code/verify.py --output data > verification_run.txt
```

The run writes three files to `$W/data` and the transcript to `$W/verification_run.txt`; compare them with the copies in `$W/recorded`. Run this way on 29 September 2026 (Python 3.14.4, about 30 s), it passed; the CSV was byte-identical and the two JSON files and the transcript agreed with the recorded ones apart from Windows line endings. The quick grid is `python3 code/verify.py -quick -output data_quick`. The delivery README's example of the root-free routine, run in the same scratch directory, reproduces (38.4):

```
import sys
sys.path.insert(0, "code")
from verify import CyclotomicRing, root_free_factors

ring = CyclotomicRing(12)
c = ring.add(ring.integer(2), ring.add(ring.zp(1), ring.zp(-1)))
L, factors = root_free_factors(8, c, ring) # c = 2 + sqrt(3)
assert L == 6
assert sum((len(p) - 1) * e for p, e in factors) == 82
```

It returns pairs (polynomial coefficients in ascending order, exponent), and reads neither the spectral generator nor the cyclotomic order. The subset expansion in the program is deliberately exponential in t and serves testing; the root-free routine is the compact algorithm.

40 Further research questions and proposed next steps

The following are precise continuations of the present results. They are research tasks relative to this article; they are not all asserted to be globally unsolved problems in the published literature.

40.1 Several distinct repeated roots

For $q(x) = \prod_i (x - c_i)^{t_i}$, equal rates are described by multiplicative relations among several z_i . The degree function is $\sum_i k_i(t_i - k_i)$, so maximizing labels lie on a higher-dimensional lattice rather than a single arithmetic progression. Can one classify when several maximal labels cancel, and bound the additional loss in terms of the dimension or shape of that maximizing set? The one-dimensional proof shows exactly which ingredient is missing: a tractable sum of leading and first surviving coefficients over all maximal labels.

40.2 Endpoint factors coupled to a cyclotomic interior root

Investigate $x^a(x-4)^b(x-c)^t$. The repository's degree law already [Theorem 14.2] adds fixed endpoint triangular contributions. The open structural issue is how those endpoint blocks alter the ratio of two tied interior leading coefficients and the first centered correction. Determine which endpoint multiplicities preserve a one-defect law and which produce higher-order cancellation.

40.3 Higher centered coefficients and a symmetric-function formula

The first correction arose from a logarithmic derivative of a signed Vandermonde. Higher corrections involve elementary symmetric functions of the mixed offsets $i + j + 1 - t$. Find a closed differential or symmetric-function expression for all centered coefficients of $P_{t,k}$. A particularly useful target is a formula exposing their parity under (33.8) and their factors at $z^2 = -1$. The goal is structural information, not merely a larger interpolation.

40.4 Uniform limits near a disappearing characteristic factor

The present formulas are exact at fixed parameters. They do not provide a uniform large- N approximation as c approaches a cyclotomic value, especially the center where whole sectors vanish. Determine the correct scaling of $c - c_0$ with N near a defect and identify the transition between the generic and specialized exponential-polynomial expansions. The exact subset formula is a finite starting point with no asymptotic error to hide.

40.5 Degrees and zeros on individual size progressions

At a cyclotomic parameter, restrict $H_N^{(t)}(c)$ to one residue class of N modulo a common rate period. Different characteristic rates now become the same, so additional cancellations can occur: (38.3) has a linear progression alongside two quadratic ones. Classify the exact degree and leading coefficient on each progression, and determine which progressions can vanish identically. This would lead to a precise description of the zero set of the Hankel determinant at cyclotomic parameters.

40.6 Other eventually constant Jacobi recurrences

Replace (31.2) by a family with a different finite initial Jacobi segment and the same constant tail. The exponential branches survive, but their amplitudes change. Determine a condition on those amplitudes under which only one extra multiplicity can be lost at a single-root cyclotomic resonance. Such a theorem would separate the universal lattice mechanism from the special Catalan boundary condition.

40.7 Effective arithmetic and reduction modulo primes

The root-free algorithm uses exact characteristic-zero equality. Determine a bit-complexity bound for algebraic inputs, including the cost of field arithmetic and coefficient growth, and study which primes preserve the characteristic-zero minimal polynomial on reduction. In positive characteristic, finite differences may lower polynomial degree abnormally and factorial factors may vanish, so neither the degree lemma nor the minimality theorem transports unchanged.

40.8 A focused formalization in Lean

The shortest reusable formalization targets are the exponential-block minimality lemma, the nearest-representative and square-sum lemmas, and the purely algebraic implication from the two sector coefficients to the one-defect law. These can be separated from integration and orthogonal polynomials. A second stage would formalize the signed confluent Vandermonde and the Christoffel identity, replacing the coefficient hypotheses by proved formulas. This staged plan must keep imported hypotheses visible rather than labeling the entire analytic argument formalized after only its final combinatorics has been checked.

41 Conclusion

The single repeated nonendpoint-root problem admits a complete sign-sensitive answer. When the rates are distinct, their nonzero sector leading coefficients give the known cubic generic order. At a cyclotomic resonance, the parabolic sector-degree function has at most one tied maximizing pair. Its leading ratio is a sign; when that sign is negative, the first centered correction proves that the loss is exactly one, except at the central root. The latter is an entire-group cancellation controlled by Cigler's known product.

This yields the minimal characteristic polynomial at every such parameter, not merely an annihilating bound. It also explains why t and the order of z^2 alone do not suffice, counts the exceptional parameters, and gives a root-free factorization computed directly from c by two short scalar recurrences. The proofs and exact audit are provided for independent checking. They establish the claims made here within ordinary mathematics; they do not establish priority or machine-checked formal status.

42 Normalization, boundary, and proof audit

This section is Appendix A of the manuscript.

Size rather than last index. N is the matrix size. An indexing convention with rows $0, \dots, n$ requires $N = n + 1$. The value at $N = 0$ is part of the sequence and equals one.

Multiplier sign. The multiplier is $(x - c)^t$, not $(c - x)^t$. This is the source of $(-1)^{Nt}$ in the Christoffel determinant and $(-1)^t$ in every characteristic root. Reversing the multiplier sign would scale the entire sequence by $(-1)^{Nt}$.

Jet factorials. The determinant uses $p_n^{(r)}(c)/r!$. Accordingly A_t appears once, in K_t , and not a second time in the confluence denominator. The subset Vandermondes use actual index differences, with their products of factorials kept explicitly.

Two different orders of a root of unity. The theorem uses $L = \text{ord}(z^2)$, while the audit ring uses a primitive M th root z , so $L = M/\text{gcd}(M, 2)$. For odd L , z can have order L or $2L$. These give opposite signs of z^L and can give different recurrence orders. This distinction is essential.

Tied constants. When $t = L$ and the defect sign is negative, $d_* = 0$. There is no degree -1 nonzero polynomial; the whole constant group is zero and its characteristic exponent is zero. This case is not covered by dividing the correction by d_* and must be handled separately, as in Section 34.

What the computation does not prove. Finite success at $t \leq 12$ does not prove an unbounded-multiplicity claim. The general mechanism is the explicit coefficient identity and the maximizing-pair lemma. Likewise, inspecting a finite set of sources does not prove historical novelty.

No unsupported nonvanishing assumptions. The Christoffel derivation divides only by a Vandermonde before confluence, not by weighted Hankel minors. The central condensation proof divides by a smaller minor only after its nonzero product formula has been established by induction. Minimality uses the nonzero coefficients actually calculated, not genericity by analogy.

43 Source and package concordance

This section is Appendix B of the manuscript.

The older root-collisions manuscript is now Part II of the repository article cited in Section 29. Its standalone Section 12.1 is the cyclotomic-classification question answered here; its Section 12.2 asks for structural cancellation criteria, addressed here for the single repeated root. The generic degree law and finite recurrence-recovery algorithm belong to that prior work. Its certified central fourth-power example is subsumed by the already known general product.

[Merge note: the standalone Sections 12.1 and 12.2 are Sections 24.1 and 24.2 here; the generic degree law and the recovery algorithm are Theorems 14.2 and 20.2; the certified central fourth-power example is (21.8).]

The present paper does not claim a general solution of multi-root resonance, uniform collision asymptotics, or positive-characteristic minimality. Those remain separate tasks. No code was added to the ProveIt repository and no Lean compiler was run for this article. [Merge note: this describes the manuscript at its pin. Its code and data have since been placed in this report, byte-identical, under the prefix 03-cyclotomic- (placement commit 3609d0473); nothing has been formalized.]

The archive contains `article.tex`, `article.pdf`, `README.md`, `SOURCES.md`, `Makefile`, `code/verify.py`, and the generated data directory. The JSON records distinguish actual grouped minimal multiplicities from generic order bounds. No external source PDFs or font files are included.

[Merge note: of these, `article.tex`, `article.pdf` and `README.md` are not shipped here; they survive in the arrival commit 8315d24e3. `SOURCES.md`, `Makefile`, `code/verify.py` and the four data files are shipped with the prefix 03-cyclotomic- (the `Makefile` in `code/`), as listed in Section 39.]

Part IV. Two-endpoint confluence

Part IV was added on 29 September 2026 in batch 53 of ProveIt's incoming-report intake, from one manuscript written against the pinned state of Parts I–III. It answers the endpoint half of Part II's research question “Uniform endpoint and root-collision limits” (Section 24.4): for any fixed numbers of roots approaching 0 and 4, with arbitrary collisions inside each cluster, it proves an exact even convergent expansion of the normalized determinant, its complete first correction, and a sharp outward scale. The question's other half, collisions of roots away from the endpoints, remains open. Parts I–III are unchanged except for dated pointers to this addition (Sections 11, 22 and 24.4), a line on the title page (whose vertical spaces were reduced by 12 mm in all so that it still fits on one page) and a paragraph of the abstract, a contents entry, three bibliography entries used only by Part IV (with a label column widened for the longer list), a clause in the closing note on source dates, and two new labels, on Part II's Section 24.4 and Part III's Section 40.2. Section 44 records the provenance and fixes the notation of the addition.

[Added 29 September 2026, batch 55: the other half, collisions of roots away from the endpoints, is answered by Part V for fixed degree and fixed nonendpoint bases (Section 61.4); limits uniform up to the endpoints remain open.]

44 Provenance and conventions of Part IV

44.1 One manuscript, one addition

Part IV is batch-53 manuscript 02, *Two-Endpoint Confluence for Catalan Hankel Determinants*, subtitled *An exact even finite-size expansion, the first interaction law, and a sharp critical scale*, an AI-assisted research article prepared for Vladimir Reshetnikov and dated 29 September 2026 (20 A4 pages as delivered). Unlike the sources of Parts I–III, it does not name the assistant that prepared it. It arrived as the archive `Two_Endpoint_Catalan_Confluence.zip` (inner directory `Two_Endpoint_Catalan_Confluence/`, main file `article.tex`) in ProveIt commit 148a906aa and was placed in 1dc874990. It was written against ProveIt commit 1ee53d57de253d16cdaad79d1f54bbd95d76d682, reading this report's `article.tex` (Git blob e5a4d81408c80f0385ae66c26fbd11bba7dfdb09) and README through the GitHub connector; its source notes name Part II's Sections 22 and 24.4, and Part III, as the relevant material. The report's `article.tex` is unchanged between the pin and the placement commit (the same blob), so every statement the manuscript makes about “the repository”, “the source” or “the report” refers to Parts I–III as printed above.

The manuscript's program and recorded outputs are shipped beside the earlier files with the prefix `04-endpoint-confluence-` (Part I's unprefix files count as 01, Part II's as 02, Part III's as 03), its dependency file under `data/`, and its source notes and proof audit, delivered as `SOURCES.md` and `PROOF_AUDIT.md`, as `04-endpoint-confluence-SOURCES.md` and `04-endpoint-confluence-PROOF_AUDIT.md` at the report root. Its manuscript, delivery README and PDF are not shipped; they survive in the arrival commit. Section 60 lists the shipped files.

The manuscript contributes, in the order printed here: the repository question, what is classical, and a guide to the proof (Section 45); the two-endpoint model and its confluent cosine and sine kernels (Section 46); the four main statements, namely the exact even expansion, the first interaction law, the endpoint values and the sharp outward scale (Section 47); a self-contained Christoffel reduction (Section 48); the exact centred hyperbolic alternant that generates every coefficient (Section 49); the proof of the expansion, with an explicit remainder bound (Section 50); the derivation of the first correction (Section 51); positive Schur expansions of the outward profiles (Section 52); the proof of the sharp scale (Section 53); the answer in the question's own uncentred scaling, and exact spatial displacements (Section 54); worked examples (Section 55); its computations and verification boundary (Section 56); ten research directions (Section 57); a conclusion; a sign and normalization audit (Section 59); and its archive

and reproduction notes (Section 60). Every result, proof, example, remark, limitation and question of the manuscript is printed. Where it re-derives something Parts I–III already prove, the overlap is marked (Section 44.5).

The manuscript’s own abstract reads:

The Catalan-Hankel report in *ProveIt* asks for uniform limits when a repeated root approaches either endpoint of the Catalan moment interval. We solve that endpoint problem for arbitrary fixed multiplicities and, more generally, for two simultaneous clusters with arbitrary internal collisions. For total multiplier degree d , the effective size is $\nu = N + d/2$. After removing the endpoint sign and the power $\nu^{\binom{m}{2} + \binom{\ell+1}{2}}$, the determinant is the restriction of a holomorphic *even* function of $1/\nu$. Its expansion is convergent, locally uniform in all scaled root coordinates, and has an explicit Cauchy remainder bound. The leading coefficient factors into two confluent cosine/sine determinants. We derive the complete first correction, including a universal mixed derivative with coefficient $-1/4$. A centered hyperbolic Vandermonde identity generates every higher coefficient. Positive Schur expansions of the outward profiles then prove an exact scale criterion: the determinant is asymptotic to its fixed-endpoint value if and only if all outward displacements are $o(N^{-2})$; if the largest scaled displacement tends to infinity, the ratio diverges. Written proofs are accompanied by 810 exact-arithmetic and high-precision checks, clearly separated by their evidentiary status.

Status (manuscript 02). This is an unrefereed, AI-assisted manuscript with conventional written proofs, not a Lean or Rocq formalization. It resolves the endpoint portion of Part II’s moving-root question, not its nonendpoint collision problem, and it does not resolve hard-edge universality for arbitrary weights or limits with multiplicities that grow with N : the multiplicities m and ℓ are fixed, and the constants and the analytic radius are not claimed to be uniform in them. The if-and-only-if scale criterion is restricted to outward, nonnegative displacements. Classical identities are credited, not presented as new: the fixed-size Christoffel reduction and its confluence (Krattenthaler [3], re-proved in the present normalization), characteristic-polynomial determinants, integrable kernels and hard-edge limits from random matrix theory [10, 11], and the Mehler–Heine phenomenon of which the one-variable cosine and sine limits are elementary half-integer cases [12]. The proposed contribution is the explicit combined package: the simultaneous two-endpoint, collision-uniform theorem, the centre $N + d/2$, the convergent expansion without odd powers, the first interaction operator, the all-orders coefficient generator, the explicit error bound and the sharp outward criterion. The literature check was bounded, not exhaustive, and worldwide priority for every identity or corollary is not established; no claim of settling a famous external conjecture is made. The proofs do not assume the report’s sector-degree or resonance theorems. The 810 recorded checks (710 exact, 100 at 100 digits) count executions, not universal theorems; the high-precision checks are diagnostics, not outward-rounded interval certificates, and the shipped program verifies the leading term and the first correction; it is not an arbitrary-order coefficient engine. No proof assistant was run, nothing in Part IV is formalized, and its placement beside a Lean project confers no formal status. (Sources: the manuscript, its delivery README, and its source notes and proof audit, 04-endpoint-confluence-SOURCES.md and 04-endpoint-confluence-PROOF_AUDIT.md.)

44.2 Reading conventions

Inside Sections 45–60, “this article”, “the present article” and “here” mean manuscript 02 as printed in Part IV; “the repository”, “the repository’s question”, “the source” and “the report” mean Parts I–III at the pin 1ee53d57d. The manuscript cites this report’s own section numbers, which did not change: its “Part II, Section 24.4” is Section 24.4 and its “Section 22” is Section 22. “The archive” means the delivered package, whose program and data are shipped under the prefixed names listed in Section 60. The manuscript’s two appendices are printed as Sections 59 and 60, so that Part I’s appendices keep their letters. Its Sections 1–14 are Sections 45–58, so a manuscript number $n.j$ of a section, theorem, lemma, corollary or equation is printed as $(n + 44).j$ (for example, its Theorem 3.1 is Theorem 47.1), and its Appendices A and B are Sections 59 and 60. The manuscript cites Krattenthaler’s Ramanujan Journal paper through

arXiv:2101.04225v5, pointing to its Theorem 1 and “the confluent discussion in Section 5”; that is Part I’s reference [3]. Its DLMF reference is Section 18.11(ii), not the Section 18.5 of Part I’s [4], and is listed separately as [12]. Bracketed paragraphs beginning “Merge note” are this merge’s, not the manuscript’s.

44.3 Notation of Part IV

Part IV keeps the manuscript’s symbols. Within Sections 45–60 every symbol has its Part IV meaning, except in merge notes and where a sentence names Part I, II or III or cites one of their equations. The multiplier has m roots near 0 and ℓ near 4 and no other roots; with all of them at the endpoints it is Part II’s $q = \gamma x^m (x - 4)^\ell \prod_i (x - c_i)^{t_i}$ with $\gamma = 1$ and $s = 0$, and $H_N(q)$ is Part II’s (14.1). Table 3 lists the symbols that a reader of Parts I–III is likely to misread. No symbol was renamed and no normalization changed. Two readings need particular care: the letter t is the variable of the kernels in Sections 46 to 53, but the multiplicity of one repeated root in (45.2) and Section 54, as in Part II’s question; and the derivative columns of the kernels (46.8)–(46.9) are *not* divided by factorials, whereas repeated-root rows are (Section 59).

Table 3: Symbols and notions of Part IV. The last column gives the meaning of the same letter in Part I, II or III, or the tempting false reading.

Symbol or notion	Meaning in Part IV	Not to be confused with
$H_N(q)$	the weighted determinant (45.1), Part II’s (14.1)	Part I’s $H_N^{(m)} = H_N(x^m)$, whose superscript is a shift; Part III’s $H_N^{(t)}(c) = H_N((x - c)^t)$.
$m, \ell, d = m + \ell$	numbers of roots near 0 and near 4; the degree of the multiplier	Part II’s multiplicities m, ℓ of the exact roots 0, 4 (the same when the roots sit there); d is Part II’s $D = \deg q$ here, not Part I’s $d = \binom{m}{2}$ nor Part II’s sector degrees d_k .
b_r, ρ, κ	$\binom{r}{2}, b_m + b_\ell, b_m + \binom{\ell+1}{2}$ (46.1)	κ is Part II’s κ (14.5); b_r is not Part I’s parameter b ; ρ is not Part II’s modulus bound in (22.1) nor Part III’s leading ratio (34.5).
$\nu = N + d/2, h = 1/\nu$	centred size and its inverse; the expansion is in h^2	Part II’s block multiplicities ν_u ; Part I’s moments ν_{i+j} ; Part III’s $h = t - k$ (compare its centred size $w = N + t/2$).
t, s	variables of C, S and of the kernels; but t is a multiplicity in (45.2) and Section 54	Part I’s generating-function variable t ; Part II’s number s of interior roots; Part III’s multiplicity t .
$C(t, y), S(t, y); C, S$ in (55.1)	$\cos(t\sqrt{y})$ and $\sin(t\sqrt{y})/\sqrt{y}$ as entire series (46.2)–(46.3); abbreviations of $C(1, y), S(1, y)$	the Catalan numbers C_n (always subscripted); Part II’s $S = 2^D$ and matrix C ; Part III’s subsets S .
$y_i, v_j; c_i^\pm; q_N$	scaled coordinates (negative real means outward); the moving roots (46.5); the size-dependent multiplier	Part II’s fixed interior roots c_i and fixed multiplier q .
$Z_N, Z(h; \mathbf{y}, \mathbf{v}), A_j$	normalized determinant (46.7), its even interpolation, its coefficients	Part II’s amplitudes $A_\pm(z)$; Part III’s superfactorials A_u ; Part I’s constant A_m .
$K_m^-, K_\ell^+; K_n^\varepsilon$	confluent cosine and sine kernels (46.8), (46.9); $\varepsilon = 0, 1$ selects them in Theorem 52.1	Part I’s $K = (m - 1)^2$; Part III’s common sector factor $K_t(z)$.
$\mathcal{A}_{m,\ell}$	endpoint value $A_0(\mathbf{0}, \mathbf{0})$ (47.5)	it equals Part II’s leading coefficient for $q = x^m(x - 4)^\ell$ (Section 44.4).

Symbol or notion	Meaning in Part IV	Not to be confused with
$a, b; \mathbf{a}, \mathbf{b}$	b_m and $\binom{\ell+1}{2}$ in Corollary 47.3; limits of scaled displacements in Theorem 47.4; a fixed positive number in Section 53	Part I's parameters a, b of $x^m(a + bx)$, which appear in the dated note of Section 11 and in Section 44.4.
$\epsilon_{i,N}, \eta_{j,N}; \varepsilon$	outward displacements at 0 and 4; the kernel type 0 or 1	Part III's parity indicator η .
R_N, M_N	ratio to the fixed-endpoint determinant (47.8); N^2 times the largest displacement	Part II's Christoffel matrix M_N (15.7) and bound R ; Part III's minimal order $R(t, c)$.
$M(h; \mathbf{y}, \mathbf{v}), Q, E(h; \mathbf{u}, \mathbf{w})$	the matrices (50.3), (49.1) and the quotient (50.5)	Part II's M_N ; Part I's $E_m(N)$ and the shift E ; the multiplier q .
$\mathcal{R}, \mathcal{R}_2$	hyperbolic generator (49.3) (constant term 1) and its h^2 coefficient (49.5)	Part II's maximal order $\mathcal{R}_0$; Part IV has no $\mathcal{R}_0$.
$\alpha_j; u_i, w_a; U, W; Y, V$	centred exponents $j - (d - 1)/2$; auxiliary variables and their sums; coordinate sums $\sum y_i, \sum v_j$	Part III's arcsine measure α and values U_a ; Part II's nodes w_a ; Part II's matrix V .
ϕ_h, c_h, s_h, D_h	the even entire functions of Section 50	Part III's sign s_N ; Part I's $D_N^{(m)}(a, b)$.
$\mathcal{D}_\pm$	Euler operators (47.4)	—
J_0, J_1, J_2, J_{11}	kernel minors with raised derivative orders (Section 51)	Part III's index set J .
k_j, λ, s_λ	strictly increasing exponents, their partition (52.1), the Schur polynomial	Part II's labels $\mathbf{k}$ and rates $\lambda_{\mathbf{k}}$; Part III's rates $\lambda_{\mathbf{k}}$; Part I's $\lambda_\pm$.
$R, r, T, B_{m,\ell,R}$	coordinate radius, disk radius, bound and constant of Theorem 50.2	Part III's $T = \binom{t}{2}$ and product $B_{a,b}(N)$; Part II's $T = \sum_i t_i$ and B_0, B_4 .
$F_t^\pm, \tau_t, \kappa_t, f_\varepsilon$	repeated-root profiles (54.1); κ_t is $\binom{t}{2}$ at 0 and $\binom{t+1}{2}$ at 4; $f_0 = C, f_1 = S$	—
$p_n, \mu; a_{n,r}$	monic orthogonal polynomials (48.2) and the Catalan measure (48.1); Taylor jets of p_n	the same as Part II's p_n, μ (Part I's $p_n^{(0)}, \mu_0$); Part III's centred moments μ_n .
Δ	the Vandermonde $\prod_{i < j} (u_j - u_i)$	Part II's Δ_q , with multiplicities; Part III's $\Delta(S)$.
$\text{sinc}, \text{sinhc}, (x)_\ell$	$\sin z/z, \sinh z/z$ (both 1 at 0), the falling factorial	—

44.4 What Part IV answers, and what stays open

Part II's Section 24.4 asked, for $c_N = 2 \pm 2 \cosh(\tau/N)$ and $q_N = (x - c_N)^t$, for the locally uniform limit of $H_N(q_N)$ normalized by $(-1)^{tN} N^{t(t+1)/2}$ at the right endpoint and by $N^{t(t-1)/2}$ at the left, with its value at $\tau = 0$, a uniform confluent determinant limit and error bounds; and, in a second version, for two distinct nonendpoint roots approaching each other at rate N^{-1} . Part IV answers the first version. Corollary 54.1 gives the limits $F_t^+(\tau)$ and $F_t^-(\tau)$, entire and even in τ , their values at $\tau = 0$ (54.5), and the N^{-1} term, which is only the centring $N \mapsto N + t/2$; Theorem 50.2 gives an explicit remainder bound. It does so inside a larger statement: m roots near 0 and ℓ near 4 at once, any collision pattern within each cluster, coordinates in compact complex sets (Theorems 47.1 and 47.2), and a sharp outward scale (Theorem 47.4). For $t = 1$ the question noted that the limits are elementary hyperbolic functions; they are $F_1^-(\tau) = \cosh \tau$ and $F_1^+(\tau) = 2 \sinh \tau/\tau$ (Section 55). The second version, collisions away from the endpoints, is *not* answered: it is Part IV's first research question (Section 57.1), and Section 24.4 stays open for it.

[Added 29 September 2026, batch 55: Part V answers the second version, and with it Section 57.1, for fixed degree and fixed nonendpoint bases (Section 61.4).]

The same results settle, for endpoint roots, the caveat that closes Part II's Section 22: its fixed-parameter estimate cannot be used with moving parameters, and Part IV proves the double-scaling limit directly, on the scale N^{-2} , without using that estimate. They also treat the limit $a \rightarrow 0$ that Part I's Section 11

names, at fixed m . For $b \neq 0$ and $\varepsilon = a/b$, $D_N^{(m)}(a, b) = b^N H_N(x^m(x + \varepsilon))$, a cluster of $m + 1$ roots at the endpoint 0, m of them fixed there. For real $\varepsilon \geq 0$ the root $-\varepsilon$ is an outward displacement, so by Theorem 47.4, with $\ell = 0$ and $m + 1$ roots, $D_N^{(m)}(a, b)/D_N^{(m)}(0, b)$ tends to 1 exactly when $\varepsilon = o(N^{-2})$, to the profile ratio (47.10) when $N^2\varepsilon$ converges, and to infinity when $N^2\varepsilon \rightarrow \infty$. Joint limits with m growing with N remain open (Section 57, “Multiplicities growing with determinant size”). Part III’s research directions on endpoint factors coupled to a cyclotomic interior root (Section 40.2) and on uniform limits near a disappearing characteristic factor concern interior roots and are not addressed; Part IV’s second research question (endpoint clusters in the presence of fixed interior roots) is close to the first of them.

[Added 29 September 2026, batch 57: for the cluster $x^m(x + \varepsilon)$ of $m + 1$ roots at 0 with one outward displacement $\varepsilon \geq 0$, Part VI treats m growing with N at every relative rate (Section 81.4); see the note under Section 57.4.]

Consistency with Parts II and III (observations of this merge, checked by a separate exact computation not shipped with the report).

1. The endpoint value (47.5) is Part II’s leading coefficient (18.8) for $q = x^m(x - 4)^\ell$ ($s = 0$, $\gamma = 1$): there $\mathcal{W} = ((-1, m), (1, \ell))$, $\Delta_q = 4^{m\ell}$ and $\mathcal{V}(\mathcal{W}) = (-1)^{b_m} 2^{m\ell} \prod_{h < m} h! \prod_{h < \ell} h!$, and $B_0(m)B_4(\ell)\mathcal{V}(\mathcal{W})/\Delta_q$ simplifies to $\mathcal{A}_{m,\ell}$; exact arithmetic agrees for $m, \ell \leq 3$. The sign $(-1)^{N\ell}$ removed in (46.7) is Part II’s rate $(-1)^\ell$. So Corollary 47.3 is a second route to the case $s = 0$ of Theorem 18.1, and it adds the next coefficient. More is true at exact endpoints: Part II’s single sector polynomial $P(N)$ of degree κ satisfies $\nu^{-\kappa}P = Z(1/\nu)$, and a polynomial in h that agrees with the even series $Z(h)$ of Theorem 47.1 at infinitely many points accumulating at 0 has only even powers; hence $P = \sum_j A_j(\mathbf{0}, \mathbf{0})\nu^{\kappa-2j}$ contains only powers of $\nu = N + d/2$ of the parity of κ . For example $H_N((x - 4)^2) = \nu(4\nu^2 - 1)/3$ with $\nu = N + 1$, and $H_N(x^2(x - 4)^2) = \nu^2(\nu^2 - 1)/12$ with $\nu = N + 2$.
2. The product (55.4) for $H_N(x^2(x - 4)^2)$ is the same polynomial as Part II’s (21.8) for $H_N((x - 2)^4)$, which is Part III’s central product $B_{2,2}(N)$; the coincidence of the two determinants is observed (both formulas are proved, and they agree for $N \leq 7$ by direct evaluation), not explained here.
3. The small determinants of Section 59 ($H_N(x)$, $H_N(x - 4)$, $H_N(x(x - 4))$, $H_N(x^2)$, $H_N((x - 4)^2)$) agree with direct evaluation for $N \leq 7$, and the table of Section 55 agrees with the case both-double recorded in data/04-endpoint-confluence-verification.json.

44.5 Where the merge had to choose

1. *Placement.* Part IV is printed after Part III and before Part I’s appendices, so that no section, equation, theorem or table number of Parts I–III moves (checked against the .aux file of a build of the committed text). Its sections continue the numbering (Sections 44 to 60), with the concordance of Section 44.2.
2. *Labels.* Every label of Part IV carries the prefix tec::; without it, seven of the manuscript’s labels, thm:main, lem:Christoffel, eq:Christoffel, eq:measure, eq:R, eq:M and eq:Q, would have collided with Part I’s bare labels. Labels were added to the manuscript’s sixteen sections and to its first research subsection, and to Part II’s Section 24.4 and Part III’s Section 40.2 so that they can be cited; no existing label was changed.
3. *Overlaps with Parts I–III.* The manuscript re-derives, without assuming the report correct, several things already proved there. Printed with a pointer and kept: the determinant (45.1) (Part II’s (14.1)); the Catalan measure, the monic orthogonal polynomials, their trigonometric form (48.3) and $H_N(1) = 1$ (a fourth route, after Part I’s Section 3, Part II’s Section 15, whose (15.3) is (48.3), and Part III’s Section 31); the Gram integral identity (48.6) and the Christoffel identity (Lemma 48.1 is Part II’s Lemma 15.1 for $\gamma = 1$, and Part III’s Lemma 31.1 and Part I’s Lemma 4.1 are other special cases); the endpoint jets (56.1), which are Part II’s (16.9) and (16.10) (the signs $(-1)^{n+r}$ and $(-1)^{n-r}$ agree); and the endpoint value of Corollary 47.3 (Section 44.4). Everything else in Part IV (the evenness

and convergence, the first interaction law, the alternant generator, the remainder bound, the Schur positivity and the sharp scale) has no counterpart in Parts I–III, which are fixed-parameter throughout. No statement of Parts I–III was changed.

4. *Bibliography.* The manuscript's Krattenthaler reference is Part I's [3]. Its references to Strahov–Fyodorov, Akemann–Fyodorov and DLMF Section 18.11(ii) are added to the common bibliography as [10], [11] and [12]. Its reference to the pinned repository report is replaced by Parts I–III themselves and the pin recorded in Section 44.1; the sentence that cited it (Section 45) now points there, and gains a pointer to Section 22 and a merge note. The manuscript's references to Part II's Section 24.4 are printed as cross-references to Section 24.4; the numbers are unchanged.
5. *File names.* Sections 56 and 60 give the shipped names beside the delivered ones in merge notes. The manuscript's reproduction command is printed as delivered but must not be run in place here, because the program would overwrite Part I's recorded outputs; a recipe on a scratch copy follows it.
6. *Front matter.* The manuscript's title page, abstract, scope box and contents are replaced by Section 44.1, which prints the abstract and the status.
7. *Notation and macros.* The manuscript's symbols are kept, with the scoping convention and Table 3 of Section 44.3; nothing was renamed. Its macros `\Del`, `\bv`, `\ba`, `\bb`, `\bu`, `\bw`, `\sinc`, `\sinhc` and `\fall` were added to the preamble with the manuscript's definitions; its `\by` and `\CC` are the report's, and its `\repo` is spelled out. No mathematical text of the manuscript was changed.

45 The repository question and the contribution

45.1 The endpoint moving-root question

For $C_n = \binom{2n}{n}/(n+1)$ and $q(x) = \sum_r q_r x^r$, put

$$H_N(q) = \det \left(\sum_r q_r C_{i+j+r} \right)_{0 \leq i, j < N}, \quad H_0(q) = 1. \quad (45.1)$$

The report *Shifted Catalan Hankel Polynomials* in *ProveIt* contains three parts: a shifted linear-multiplier analysis, an extension to arbitrary polynomial multipliers, and a single-root cyclotomic classification. Its Part II, Section 24.4, asks for the locally uniform limit associated with

$$c_N = 2 + 2 \cosh(\tau/N), \quad q_N(x) = (x - c_N)^t, \quad \frac{H_N(q_N)}{(-1)^{tN} N^{t(t+1)/2}}, \quad (45.2)$$

and for the corresponding left-endpoint limit with exponent $t(t-1)/2$. It specifically requests confluent determinant limits and error bounds. The same subsection separately asks about collisions away from the endpoints; that separate problem is not solved here.

The source was read at repository commit 1ee53d57de253d16cdaad79d1f54bbd95d76d682. The mathematical source file had blob e5a4d81408c80f0385ae66c26fbd11bba7dfdb09. The path, relevant sections, and comparison are recorded in Section 44.1 and the accompanying SOURCES.md. In particular, the source's fixed-parameter asymptotics (Section 22) expressly warn that they cannot simply be substituted into a moving-root limit.

[Merge note: the manuscript's SOURCES.md is shipped as 04-endpoint-confluence-SOURCES.md; its bibliography item for the pinned report is replaced by Parts I–III themselves and the pin recorded in Section 44.1. Section 24.4 of Part II and Section 22 of Part II, as the manuscript cites them, are Sections 24.4 and 22 here: the numbers did not change.]

The results here address a larger fixed-degree problem. Any number m of roots may approach 0, any number ℓ may approach 4, and roots inside either cluster may coincide in any pattern. The approach may

be from outside the interval, inside it, or in complex directions. Uniformity includes all these collisions on compact sets of the scaled coordinates.

45.2 What is classical, and what is established here

The reduction of a polynomially modified moment determinant to a fixed-size determinant of orthogonal polynomials is classical. Krattenthaler gives a detailed account, four proofs, a confluent version, and recurrence applications [3]. We use that identity, with a self-contained proof for the present normalization. The broader appearance of characteristic-polynomial determinants, integrable kernels, and hard-edge limits is likewise established in random matrix theory; see Strahov–Fyodorov and Akemann–Fyodorov [10, 11]. The one-variable cosine and sine limits are elementary half-integer cases of the Mehler–Heine phenomenon [12].

The contribution is the following explicit package for the repository’s problem: a simultaneous two-endpoint, collision-uniform theorem; the exact center $N + d/2$; a genuinely convergent expansion containing no odd powers; a closed first interaction operator; an all-orders coefficient generator; an elementary, explicit error bound; and a necessary-and-sufficient outward stability criterion. These conclusions are proved directly, without assuming the repository’s newer sector-degree or resonance claims. A bounded literature check does not establish priority for every identity or corollary in this package.

45.3 A guide to the proof

The decisive algebra is not a delicate large determinant estimate. Centering the columns turns an exponential alternant into a product of hyperbolic sines within each cluster and hyperbolic cosines between the clusters. After the forced zero is divided out, that product is even in the inverse size. A fixed-dimensional Christoffel formula transports this identity to Catalan determinants. The first correction follows from the quadratic term of the product; positivity of the limiting profiles then settles the sharp scale question.

46 Notation and the two-endpoint model

Fix nonnegative integers m, ℓ with $d = m + \ell \geq 1$. Write

$$b_r = \binom{r}{2}, \quad \rho = b_m + b_\ell, \quad \kappa = b_m + \binom{\ell+1}{2} = \rho + \ell, \quad \nu = N + \frac{d}{2}, \quad h = \nu^{-1}. \quad (46.1)$$

For a tuple $\mathbf{u}$, let $\Delta(\mathbf{u}) = \prod_{i < j} (u_j - u_i)$; empty products and empty determinants are 1.

All square-root expressions below denote entire power series in the squared variable, so no branch choice is part of the definition. Set

$$C(t, y) = \cos(t\sqrt{y}) = \sum_{k \geq 0} \frac{(-1)^k t^{2k} y^k}{(2k)!}, \quad (46.2)$$

$$S(t, y) = \frac{\sin(t\sqrt{y})}{\sqrt{y}} = \sum_{k \geq 0} \frac{(-1)^k t^{2k+1} y^k}{(2k+1)!}. \quad (46.3)$$

In particular $S(t, 0) = t$. For $\mathbf{y} \in \mathbb{C}^m$ and $\mathbf{v} \in \mathbb{C}^\ell$, define the moving roots and multiplier by

$$c_i^-(h) = 2 - 2\cos(h\sqrt{y_i}), \quad 1 \leq i \leq m, \quad (46.4)$$

$$c_j^+(h) = 2 + 2\cos(h\sqrt{v_j}), \quad 1 \leq j \leq \ell, \quad (46.5)$$

$$q_N(x) = \prod_{i=1}^m (x - c_i^-(h)) \prod_{j=1}^\ell (x - c_j^+(h)). \quad (46.6)$$

Thus $c_i^- = h^2 y_i + O(h^4)$ and $c_j^+ = 4 - h^2 v_j + O(h^4)$. Negative real coordinates mean outward motion. The normalized determinant is

$$Z_N(\mathbf{y}, \mathbf{v}) = (-1)^{N\ell} v^{-\kappa} H_N(q_N). \quad (46.7)$$

Define the two entire, symmetric kernel functions by

$$K_m^-(t; \mathbf{y}) = (-1)^{b_m} \frac{\det[\partial_t^r C(t, y_i)]_{1 \leq i \leq m, 0 \leq r < m}}{\Delta(\mathbf{y})}, \quad (46.8)$$

$$K_\ell^+(s; \mathbf{v}) = (-1)^{b_\ell} \frac{\det[\partial_s^r S(s, v_j)]_{1 \leq j \leq \ell, 0 \leq r < \ell}}{\Delta(\mathbf{v})}. \quad (46.9)$$

The quotients are defined at collisions by analytic continuation. More concretely, a block of r repeated coordinates z is replaced by rows of normalized derivatives $(1/a!) \partial_z^a$, $0 \leq a < r$; the remaining denominator is the product of cross-block differences to the products of their multiplicities. Set $K_0^\pm = 1$.

47 Main results

47.1 An exact even expansion

Theorem 47.1 (Two-endpoint confluence). *There are entire functions $A_j(\mathbf{y}, \mathbf{v})$, symmetric separately in each cluster, such that for all sufficiently large N ,*

$$Z_N(\mathbf{y}, \mathbf{v}) = \sum_{j=0}^{\infty} \frac{A_j(\mathbf{y}, \mathbf{v})}{v^{2j}}. \quad (47.1)$$

The series converges locally uniformly in the root coordinates and uniformly for $|h|$ in a sufficiently small closed disk. It is the Taylor series of an even holomorphic function $Z(h; \mathbf{y}, \mathbf{v})$ with $Z_N = Z(v^{-1}; \mathbf{y}, \mathbf{v})$. In particular,

$$A_0(\mathbf{y}, \mathbf{v}) = 2^{\ell-m\ell} K_m^-(1; \mathbf{y}) K_\ell^+(1; \mathbf{v}). \quad (47.2)$$

All statements include repeated roots, zero coordinates, and mixed collision patterns. For every fixed J the remainder after A_J is $O(v^{-2J-2})$ locally uniformly, including after any fixed number of derivatives in the coordinates.

The theorem is stronger than the existence of a formal asymptotic series. It constructs an analytic function of the inverse size. The absence of odd powers is exact, not a cancellation observed through a finite order. The restriction to fixed m, ℓ is important: the constants and the analytic radius are not claimed to be uniform in growing multiplicities.

47.2 The complete first correction

In the next statement every kernel and its derivatives are evaluated at $t = s = 1$. Put $Y = \sum_i y_i$ and $V = \sum_j v_j$.

Theorem 47.2 (First interaction law). *The coefficient of v^{-2} in (47.1) is*

$$A_1 = 2^{\ell-m\ell} \left\{ \frac{(m+3\ell+1)Y + (\ell+3m-1)V}{24} K_m^- K_\ell^+ - \frac{(\partial_t^2 K_m^-) K_\ell^+ + K_m^- (\partial_s^2 K_\ell^+)}{24} - \frac{(\partial_t K_m^-)(\partial_s K_\ell^+)}{4} \right\}. \quad (47.3)$$

This identity is entire in every coordinate; it does not divide by the leading profile, so it remains valid at zeros of A_0 .

At leading order the clusters decouple, up to the scalar $2^{\ell-m\ell}$. The first correction has a mixed derivative whose coefficient is always $-1/4$. It is therefore incorrect, in general, to multiply two one-endpoint finite-size expansions and expect the simultaneous two-endpoint correction.

An equivalent form uses the Euler operators

$$\mathcal{D}_- = b_m + 2 \sum_i y_i \partial_{y_i}, \quad \mathcal{D}_+ = \binom{\ell+1}{2} + 2 \sum_j v_j \partial_{v_j}. \quad (47.4)$$

On the product of the kernels, replace ∂_t by $\mathcal{D}_-$, ∂_t^2 by $\mathcal{D}_-(\mathcal{D}_- - 1)$, and analogously for s . This eliminates auxiliary variables from the correction formula.

47.3 The endpoint value

Corollary 47.3. *At $y = \mathbf{0}$, $v = \mathbf{0}$,*

$$A_0(\mathbf{0}, \mathbf{0}) = \mathcal{A}_{m,\ell} := 2^{\ell-m\ell+b_m+b_\ell} \prod_{r=0}^{m-1} \frac{r!}{(2r)!} \prod_{r=0}^{\ell-1} \frac{r!}{(2r+1)!} > 0. \quad (47.5)$$

Writing $a = b_m$ and $b = \binom{\ell+1}{2}$, one also has

$$\frac{A_1(\mathbf{0}, \mathbf{0})}{\mathcal{A}_{m,\ell}} = -\frac{a(a-1) + b(b-1) + 6ab}{24}. \quad (47.6)$$

47.4 A sharp scale criterion

For nonnegative real outward displacements define

$$\tilde{q}_N(x) = \prod_{i=1}^m (x + \epsilon_{i,N}) \prod_{j=1}^{\ell} (x - 4 - \eta_{j,N}), \quad (47.7)$$

$$R_N = \frac{H_N(\tilde{q}_N)}{H_N(x^m(x-4)^\ell)}, \quad M_N = N^2 \max\{\epsilon_{i,N}, \eta_{j,N}\}. \quad (47.8)$$

The denominator is nonzero and R_N is positive.

Theorem 47.4 (Exact outward stability threshold). *With fixed m, ℓ as above,*

$$R_N \longrightarrow 1 \iff M_N \longrightarrow 0. \quad (47.9)$$

If $N^2\epsilon_{i,N} \rightarrow a_i$ and $N^2\eta_{j,N} \rightarrow b_j$ with finite nonnegative limits, then

$$R_N \longrightarrow \frac{K_m^-(1; -\mathbf{a})}{K_m^-(1; \mathbf{0})} \frac{K_\ell^+(1; -\mathbf{b})}{K_\ell^+(1; \mathbf{0})}. \quad (47.10)$$

This limit exceeds 1 if at least one limit coordinate is positive. If $M_N \rightarrow \infty$, then $R_N \rightarrow \infty$.

The equivalence is restricted to outward, nonnegative displacements. Inside the interval, oscillations and zeros prevent the same monotonicity argument and may invalidate such an equivalence.

48 The Catalan moment functional and Christoffel reduction

48.1 Orthogonal polynomials with unit norms

The Catalan numbers are moments of the probability measure

$$d\mu(x) = \frac{1}{2\pi} \sqrt{\frac{4-x}{x}} dx, \quad 0 < x < 4. \quad (48.1)$$

Indeed, $x = 4u$ gives $\int x^k d\mu = (2/\pi)4^k B(k + 1/2, 3/2) = C_k$. Define monic polynomials by

$$p_0 = 1, \quad p_1 = x - 1, \quad p_{n+1} = (x - 2)p_n - p_{n-1} \quad (n \geq 1). \quad (48.2)$$

Their elementary trigonometric expressions are

$$p_n(2 + 2 \cos \theta) = \frac{\sin((n + 1/2)\theta)}{\sin(\theta/2)}, \quad (48.3)$$

$$p_n(2 - 2 \cos \theta) = (-1)^n \frac{\cos((n + 1/2)\theta)}{\cos(\theta/2)}. \quad (48.4)$$

The identities follow by checking $n = 0, 1$ and the recurrence; the apparent singularities at special angles are removable where appropriate.

Under $x = 2 + 2 \cos \theta$, the measure is $(2/\pi) \sin^2(\theta/2) d\theta$ on $0 < \theta < \pi$. Thus the functions in (48.3) are orthonormal, since the half-integer sine functions have squared integral $\pi/2$ and distinct ones are orthogonal. In particular $\int p_n p_k d\mu = \delta_{nk}$ and $H_N(1) = 1$. These facts fix all normalizing constants used below.

48.2 A proof of the fixed-size determinant identity

Lemma 48.1 (Christoffel identity). For $q(x) = \prod_{a=1}^d (x - c_a)$,

$$H_N(q) = (-1)^{Nd} \frac{\det[p_{N+j}(c_i)]_{1 \leq i \leq d, 0 \leq j < d}}{\Delta(c_1, \dots, c_d)}. \quad (48.5)$$

The quotient is a symmetric polynomial in the c_a and hence extends to all collisions by normalized derivative rows.

Proof. Expanding two determinants and integrating term by term yields the Gram integral identity

$$H_N(q) = \frac{1}{N!} \int_{(0,4)^N} \Delta(x_1, \dots, x_N)^2 \prod_{a=1}^N q(x_a) d\mu(x_a). \quad (48.6)$$

The same identity holds for complex q , by finite multilinearity. The enlarged Vandermonde in the ordered tuple $(x_1, \dots, x_N, c_1, \dots, c_d)$ is

$$\Delta(\mathbf{x})\Delta(\mathbf{c}) \prod_{a,i} (c_i - x_a).$$

Replace its monomial columns by the monic columns $p_0, p_1, \dots$; the determinant does not change. Multiply by $\Delta(\mathbf{x}) = \det[p_{j-1}(x_i)]_{1 \leq i, j \leq N}$ and integrate. Laplace expansion on the first N rows followed by orthogonality kills every choice of N columns except the first N . The surviving integral is $N!$, and the remaining determinant is $\det[p_{N+j}(c_i)]$. Comparing with (48.6) gives (48.5), including the sign $(-1)^{Nd}$. An alternating polynomial is divisible by the Vandermonde. Taylor expansion of rows in a colliding block proves the normalized derivative rule and its factorials. $\square$

This proof is included to make the present article self-contained, not as a claim of a new Christoffel identity. The general identity and its confluence are explicitly treated in [3].

49 The centered hyperbolic alternant

The following finite determinant is the algebraic engine of the article. Let $\alpha_j = j - (d - 1)/2$, $0 \leq j < d$. Introduce independent auxiliary variables $u_1, \dots, u_m, w_1, \dots, w_\ell$ and the $d \times d$ matrix

$$E(h; \mathbf{u}, \mathbf{w}) = \begin{bmatrix} [(-1)^j e^{\alpha_j h u_i}]_{i,j} \\ [e^{\alpha_j h w_a}]_{a,j} \end{bmatrix}. \quad (49.1)$$

We use $\operatorname{sinhc} z = \sinh(z)/z$, with value 1 at zero.

Lemma 49.1 (Exact centered alternant). *One has the identity of entire functions*

$$\det E = (-1)^{b_m} 2^{m\ell} h^\rho \Delta(\mathbf{u}) \Delta(\mathbf{w}) \mathcal{R}(h; \mathbf{u}, \mathbf{w}), \quad (49.2)$$

where

$$\mathcal{R} = \prod_{i < i'} \operatorname{sinhc} \left(\frac{h}{2} (u_{i'} - u_i) \right) \prod_{a < a'} \operatorname{sinhc} \left(\frac{h}{2} (w_{a'} - w_a) \right) \prod_{i,a} \cosh \left(\frac{h}{2} (w_a - u_i) \right). \quad (49.3)$$

Consequently $\mathcal{R}$ is even in h , symmetric within each tuple, and its constant term is 1.

Proof. Factor $e^{-h(d-1)u_i/2}$ from each lower row and $e^{-h(d-1)w_a/2}$ from each upper row. The remaining determinant is the Vandermonde at nodes $-e^{hu_i}$ and e^{hw_a} . Within the first cluster its factors are $-2e^{h(u_i+u_{i'})/2} \sinh(h(u_{i'} - u_i)/2)$; within the second cluster they are the analogous positive factors. Between clusters they are $2e^{h(u_i+w_a)/2} \cosh(h(w_a - u_i)/2)$. Every variable occurs in exactly $d - 1$ exponential factors, so the centering cancels all exponentials. Extract h times each within-cluster difference. There are ρ such differences and $m\ell$ cross pairs. This gives (49.2), including its sign and power of 2. $\square$

49.1 The quadratic generator

The first nonconstant term is

$$\mathcal{R} = 1 + h^2 \mathcal{R}_2 + O(h^4), \quad (49.4)$$

where, writing $U = \sum_i u_i$ and $W = \sum_a w_a$,

$$24\mathcal{R}_2 = (m + 3\ell) \sum_i u_i^2 - U^2 + (\ell + 3m) \sum_a w_a^2 - W^2 - 6UW. \quad (49.5)$$

Indeed, a within-cluster pair contributes its squared difference divided by 24, and a cross pair contributes its squared difference divided by 8. Summing the differences gives the displayed formula.

49.2 The all-orders meaning

Expand the right side of (49.3) to any prescribed order. The coefficient of h^{2j} is a finite symmetric polynomial with rational coefficients. In the alternant on the left, expanding each row in powers of h identifies the coefficients with minors of $[(-1)^j \alpha_j^r / r!]$ and $[\alpha_j^s / s!]$. Equivalently, expand $\mathcal{R}$ in the two Schur bases; a Schur partition records exactly which derivative orders replace the consecutive orders $0, \dots, m - 1$ or $0, \dots, \ell - 1$.

This supplies an all-orders coefficient algorithm. It can also be read as an identity of constant-coefficient differential operators: replace u_i, w_a by derivatives in separate row variables and apply the result to a product of the functions C and S . Each finite Taylor coefficient uses only finitely many derivatives. Convergence is justified below by holomorphy, rather than by a purely formal interchange of infinite operators.

50 Proof of the even, collision-uniform expansion

50.1 An exact normalized formula

Put

$$\phi_h(y) = \frac{2 - 2 \cos(h\sqrt{y})}{h^2}, \quad \phi_0(y) = y, \quad (50.1)$$

$$c_h(y) = \cos(h\sqrt{y}/2), \quad s_h(y) = \text{sinc}(h\sqrt{y}/2). \quad (50.2)$$

All three functions are entire in (h, y) and even in h . Define

$$M(h; \mathbf{y}, \mathbf{v}) = \begin{bmatrix} [(-1)^j C(1 + \alpha_j h, y_i)]_{i,j} \\ [S(1 + \alpha_j h, v_a)]_{a,j} \end{bmatrix}. \quad (50.3)$$

Using (48.3)–(48.4) in (48.5) yields

$$Z(h; \mathbf{y}, \mathbf{v}) = \frac{2^\ell (-1)^{b_\ell} h^{-\rho} \det M}{\prod_i c_h(y_i) \prod_a s_h(v_a) \Delta(\phi_h(\mathbf{y})) \Delta(\phi_h(\mathbf{v})) \prod_{i,a} [4 - h^2(\phi_h(y_i) + \phi_h(v_a))]} \quad (50.4)$$

At $h = (N + d/2)^{-1}$ the right side equals Z_N .

For completeness, the powers and signs are as follows. The m lower rows supply $(-1)^{Nm}$, which combines with the Christoffel sign to leave $(-1)^{N\ell}$. The upper rows each supply $2/h$. The lower within-cluster Vandermonde is $h^{2b_m} \Delta(\phi_h(\mathbf{y}))$; the upper one is $(-1)^{b_\ell} h^{2b_\ell} \Delta(\phi_h(\mathbf{v}))$. Multiplication by h^κ leaves $h^{-\rho}$, as in (50.4). No additional factorials belong in this formula.

50.2 Removal of the apparent singularities

Lemma 50.1. *The function*

$$Q(h; \mathbf{y}, \mathbf{v}) = \frac{\det M(h; \mathbf{y}, \mathbf{v})}{h^\rho \Delta(\mathbf{y}) \Delta(\mathbf{v})} \quad (50.5)$$

is entire and even in h , and entire in every coordinate.

Proof. Taylor-expand each row in h . In a nonzero determinant term the selected derivative orders within one cluster must be distinct, since equal orders would give proportional rows in the corresponding coefficient matrix. Their smallest possible sum is b_m for the lower cluster and b_ℓ for the upper cluster. Thus the determinant vanishes to order at least ρ . Its quotient by h^ρ is entire, jointly with the coordinates. Alternation in the coordinates of either cluster then removes the two Vandermondes. One may justify joint removal by successive analytic divided differences, which also gives the stated confluent rule.

For parity, reverse the d columns. Since $\alpha_{d-1-j} = -\alpha_j$, one obtains

$$\det M(-h) = (-1)^{b_d+m(d-1)} \det M(h) = (-1)^\rho \det M(h).$$

The last equality follows because $b_d + m(d-1) - \rho = m(m-1) + 2m\ell$ is even. Dividing by h^ρ proves evenness. Alternatively, every odd coefficient vanishes in (49.2). $\square$

Write

$$D_h(\mathbf{y}) = \prod_{i < i'} \frac{\phi_h(y_{i'}) - \phi_h(y_i)}{y_{i'} - y_i}, \quad (50.6)$$

with derivatives at coincidences, and similarly for $\mathbf{v}$. Then (50.4) is $2^\ell (-1)^{b_\ell} Q$ divided by $\prod c_h \prod s_h D_h(\mathbf{y}) D_h(\mathbf{v})$ and the cross factors. At $h = 0$ these denominator factors are respectively 1, 1, 1, 1 and $4^{m\ell}$. They remain nonzero on a sufficiently small disk, uniformly on any compact coordinate set. This proves holomorphy and convergence of the even Taylor series.

The least-order term in the row expansion uses precisely the derivative orders $0, \dots, m-1$ and $0, \dots, \ell-1$. By the constant term of (49.2), its coefficient is

$$(-1)^{b_m} 2^{m\ell} \det[\partial_i^r C(1, y_i)] \det[\partial_s^r S(1, v_a)].$$

Substitution in (50.4) gives (47.2). All Taylor coefficients are entire in the coordinates, because every denominator has a nonzero constant in h and every numerator coefficient is an entire alternating quotient. Uniform convergence of differentiated series follows from Cauchy's estimates on a slightly larger compact set. This proves Theorem 47.1.

50.3 An explicit remainder bound

Theorem 50.2 (A conservative effective bound). *Suppose $|y_i|, |v_a| \leq R$, $R \geq 0$. Let*

$$r = \frac{1}{4\sqrt{R+1}}, \quad T = 1 + \frac{d-1}{2}r, \quad (50.7)$$

$$B_{m,\ell,R} = 2^{\ell+d+\rho} d! r^{-\rho} T^d \exp(dT\sqrt{R+1}). \quad (50.8)$$

Then Z is holomorphic on a neighborhood of $|h| \leq r$ and has modulus at most $B_{m,\ell,R}$ on $|h| = r$. For $|h| < r$,

$$\left| Z(h) - \sum_{j=0}^J A_j h^{2j} \right| \leq B_{m,\ell,R} \frac{(|h|/r)^{2J+2}}{1 - (|h|/r)^2}. \quad (50.9)$$

Proof. Here $|h\sqrt{y}| \leq 1/4$. The power-series bounds give $|c_h(y) - 1| < 1/2$, $|s_h(y) - 1| < 1/2$, and

$$|\phi'_h(y) - 1| = |\text{sinc}(h\sqrt{y}) - 1| \leq \frac{\sinh(1/4)}{1/4} - 1 < \frac{1}{50}.$$

A divided difference of ϕ_h is an integral of its derivative on the straight segment between its nodes. It consequently has distance less than $1/50$ from 1. Also $|\phi_h(y)| \leq (51/50)R$, so each cross factor has modulus greater than 3. In particular all denominator factors in the rewritten version of (50.4) are nonzero; for an upper bound their inverse product is at most $2^{d+\rho}$.

To bound the determinant divided by the coordinate Vandermondes, replace rows by successive divided differences within each cluster. Every such row entry is bounded by the supremum of its original entire row function on $|y| \leq R+1$, by Cauchy's derivative estimate and the integral formula for divided differences. On that larger disk, $|1 + \alpha_j h| \leq T$, and both C and S have modulus at most $T \exp(T\sqrt{R+1})$. The determinant is therefore bounded by $d!$ times the d th power of this number. On $|h| = r$, division by h^ρ costs $r^{-\rho}$. The factor 2^ℓ in (50.4) gives (50.8). Finally apply Cauchy's coefficient bound to the even Taylor series and sum its geometric tail. $\square$

The constants deliberately favor transparency over sharpness. They are explicit analytic bounds, whereas the numerical tests later in the article are not interval certificates.

51 Derivation of the first interaction coefficient

51.1 The five relevant minors

For a cluster of size n , denote the determinant with column derivative orders $0, \dots, n-1$ by J_0 . Let J_1, J_2, J_{11} use, respectively,

$$(0, \dots, n-2, n), \quad (0, \dots, n-2, n+1), \quad (0, \dots, n-3, n-1, n).$$

Set $J_{11} = 0$ if $n < 2$. The symbols refer to either the C or S cluster. They may be divided by the same Vandermonde and multiplied by the same sign as the base kernel.

The coefficient of degree two in the centered alternant gives the following ratios to the base coefficient. The row coefficients use $\alpha_j^r/r!$, so the factorials in this table are already included.

Derivative selection	Coefficient ratio
Lower J_2 , upper J_0	$(m + 3\ell - 1)/24$
Lower J_{11} , upper J_0	$-(m + 3\ell + 1)/24$
Lower J_0 , upper J_2	$(\ell + 3m - 1)/24$
Lower J_0 , upper J_{11}	$-(\ell + 3m + 1)/24$
Lower J_1 , upper J_1	$-1/4$

A selection involving a nonexistent cluster is simply absent. To verify the table directly, use $s_{(2)} - s_{(1,1)} = \sum u_i^2$ and $s_{(2)} + s_{(1,1)} = (\sum u_i)^2$ in (49.5). The term $UW = s_{(1)}(\mathbf{u})s_{(1)}(\mathbf{w})$ gives the mixed coefficient. There are no degree-one selections.

51.2 Reduction to ordinary kernel derivatives

Every row function satisfies $\partial_t^2 f(t, y) = -y f(t, y)$. Differentiating J_0 column by column leaves only the last column, hence

$$J'_0 = J_1, \quad J''_0 = J_2 + J_{11}, \quad J_2 - J_{11} = -\left(\sum y_i\right)J_0. \quad (51.1)$$

For the last identity, sum the determinants obtained by applying two derivatives to one column. The differential equation says that this sum is the coefficient of a diagonal row multiplication by $-y_i$, hence $-(\sum y_i)J_0$. All terms except the last two columns vanish; the next-to-last term is $-J_{11}$ because its two last columns are reversed. These identities are polynomial determinant identities and hold even where $J_0 = 0$.

It follows from the table that the normalized numerator correction is

$$-\frac{(K_m^-)''K_\ell^+ + K_m^-(K_\ell^+)''}{24} - \frac{(m + 3\ell)Y + (\ell + 3m)V}{24}K_m^-K_\ell^+ - \frac{(K_m^-)'(K_\ell^+)'}{4}. \quad (51.2)$$

51.3 The geometric denominator correction

The remaining factors must not be omitted. Their expansions are

$$\begin{aligned} \prod_i c_h(y_i)^{-1} &= 1 + \frac{h^2 Y}{8} + O(h^4), \\ \prod_a s_h(v_a)^{-1} &= 1 + \frac{h^2 V}{24} + O(h^4), \\ D_h(\mathbf{y})^{-1} &= 1 + \frac{h^2(m-1)Y}{12} + O(h^4), \\ D_h(\mathbf{v})^{-1} &= 1 + \frac{h^2(\ell-1)V}{12} + O(h^4), \\ \prod_{i,a} [4 - h^2(\phi_h(y_i) + \phi_h(v_a))]^{-1} &= 4^{-m\ell} \left(1 + \frac{h^2(\ell Y + mV)}{4} + O(h^4) \right). \end{aligned}$$

For example $\phi_h(y) = y - h^2 y^2/12 + O(h^4)$ proves the two divided Vandermonde expansions even at collisions. Adding these contributions to (51.2) gives exactly (47.3). This proves Theorem 47.2.

Finally, rescaling the kernel determinants gives

$$K_m^-(t; \mathbf{y}) = t^{b_m} K_m^-(1; t^2 \mathbf{y}), \quad (51.3)$$

$$K_\ell^+(s; \mathbf{v}) = s^{\binom{\ell+1}{2}} K_\ell^+(1; s^2 \mathbf{v}). \quad (51.4)$$

The identities first follow for nonzero t, s by extracting powers from rows, columns and Vandermondes, then extend to zero by holomorphy. Differentiation at 1 proves the Euler-operator form.

52 Positive Schur expansions of the outward profiles

52.1 A coefficientwise-positive formula

For a strictly increasing tuple $0 \leq k_0 < \dots < k_{n-1}$, define the partition

$$\lambda_i = k_{n-i} - (n - i), \quad 1 \leq i \leq n. \quad (52.1)$$

Let s_λ denote the Schur polynomial, equivalently the quotient $\det[a_i^{k_j}]/\Delta(\mathbf{a})$ in this ordering.

Theorem 52.1 (Positive outward expansion). *For $\varepsilon = 0, 1$, write K_n^ε for K_n^- and K_n^+ , respectively. Then*

$$K_n^\varepsilon(1; -\mathbf{a}) = 2^{b_n} \sum_{0 \leq k_0 < \dots < k_{n-1}} \frac{\Delta(k_0, \dots, k_{n-1})}{\prod_{j=0}^{n-1} (2k_j + \varepsilon)!} s_\lambda(\mathbf{a}). \quad (52.2)$$

The series converges locally uniformly on $\mathbb{C}^n$ and all its coefficients in the variables a_i are nonnegative real numbers. Its constant term is

$$K_n^\varepsilon(1; \mathbf{0}) = 2^{b_n} \prod_{j=0}^{n-1} \frac{j!}{(2j + \varepsilon)!}. \quad (52.3)$$

For $n \geq 1$, the coefficient of $a_1 + \dots + a_n$ is strictly positive.

Proof. At negative coordinates, the series for the row functions has positive coefficients in a . Apply Cauchy–Binet to its power-series factorization. For chosen powers $k_0 < \dots < k_{n-1}$, the second determinant has entries

$$\frac{(2k_i + \varepsilon)_r}{(2k_i + \varepsilon)!}, \quad 0 \leq i, r < n.$$

The numerator is the Vandermonde in $2k_i + \varepsilon$, because the falling factorials are monic polynomials of their degrees. It equals $2^{b_n} \Delta(\mathbf{k})$. The sign in the definition of the kernel cancels the sign of $\Delta(-\mathbf{a})$, leaving (52.2).

This computation may first be carried out for truncated series. Their entire row functions converge with all derivatives on compact sets, so the determinant divided differences converge there as well. Alternatively, on a positive diagonal $a_i = A$ the series is monotone and bounded by the confluent determinant limit; its positive coefficients then majorize the series on $|a_i| \leq A$. This gives local uniform convergence. Schur polynomials have nonnegative coefficients: the branching identity

$$s_\lambda(a_1, \dots, a_n) = \sum_{\mu \text{ interlaces } \lambda} a_n^{|\lambda| - |\mu|} s_\mu(a_1, \dots, a_{n-1})$$

proves this by induction, starting with a monomial in one variable. The identity follows from the bialternant formula after clearing its Vandermonde. The smallest tuple is $k_j = j$, which gives the constant term since $\Delta(0, 1, \dots, n-1) = \prod_j j!$. The tuple $(0, 1, \dots, n-2, n)$ gives $\lambda = (1)$ and a strictly positive coefficient. For $n = 1$ the same assertion uses $k_0 = 1$. $\square$

52.2 Consequences and a useful relative error bound

For nonnegative $\mathbf{a}$,

$$K_n^\varepsilon(1; -\mathbf{a}) \geq K_n^\varepsilon(1; \mathbf{0}) > 0. \quad (52.4)$$

The inequality is strict if at least one coordinate is positive, and every coordinate derivative of the profile is nonnegative. In particular, $A_0(-\mathbf{a}, -\mathbf{b}) \geq \mathcal{A}_{m,\ell}$. Thus the absolute error in Theorem 50.2, divided by $\mathcal{A}_{m,\ell}$, is a valid relative-to-leading-term bound on an outward compact set.

Formula (52.3) and the homogeneity laws immediately give (47.5) and (47.6). This proves Corollary 47.3. The two different triangular exponents are now visible directly: the cosine determinant contributes b_m , whereas the sine-divided-by-root determinant contributes $b_\ell + \ell$.

53 Proof of the sharp outward critical-scale theorem

53.1 Monotonicity at every finite size

Multiply (48.6) by $(-1)^{N\ell}$ for the outward multiplier. Its integrand contains

$$\prod_{a=1}^N \left\{ \prod_i (x_a + \epsilon_{i,N}) \prod_j (4 + \eta_{j,N} - x_a) \right\}.$$

Every factor is positive for $0 < x_a < 4$. The integral is positive, and increasing any displacement increases the integrand pointwise. Consequently $R_N \geq 1$ and it is nondecreasing in every displacement. No assumption of distinct roots is required.

53.2 Finite scaled coordinates

For an outward displacement $\epsilon \geq 0$, define its angular coordinate by

$$a(h, \epsilon) = h^{-2} \operatorname{arcosh}(1 + \epsilon/2)^2. \quad (53.1)$$

Then $2 - 2 \cos(h\sqrt{-a}) = -\epsilon$. Since $\operatorname{arcosh}(1 + \epsilon/2)^2 = \epsilon + O(\epsilon^2)$ as $\epsilon \rightarrow 0$, bounded $N^2\epsilon$ implies $a(h, \epsilon) - N^2\epsilon \rightarrow 0$. The same coordinate works for $4 + \eta$. Apply Theorem 47.1 to these bounded moving coordinates and divide by its value at zero. This proves (47.10), including $R_N \rightarrow 1$ when $M_N \rightarrow 0$.

53.3 Necessity and escape from the critical window

Suppose M_N does not tend to zero. Along a subsequence, one fixed coordinate has N^2 times its displacement at least some $a > 0$; there is a fixed coordinate after passing to a further subsequence because d is fixed. By finite-size monotonicity, R_N is bounded below by the ratio with that displacement replaced by a/N^2 and all other displacements set to zero. Its limit is strictly greater than 1 by Theorem 52.1. Hence R_N cannot tend to 1.

Now suppose $M_N \rightarrow \infty$. Fix $A > 0$. For all sufficiently large N , some displacement is at least A/N^2 . The same comparison bounds R_N below by the minimum of the finitely many one-coordinate comparison ratios. For each location, its limiting Schur series has a positive linear coefficient in A , so the minimum of those finitely many limits tends to infinity as $A \rightarrow \infty$. Given any desired lower bound, choose A first and then N sufficiently large. This proves $R_N \rightarrow \infty$ and completes Theorem 47.4.

The divergence conclusion is not a large-coordinate use of the compact asymptotic theorem. It follows by comparison with arbitrarily large *fixed* critical coordinates, which is why no unjustified uniformity at infinity is needed.

54 The original scaling and other coordinate choices

54.1 An explicit answer to the repository's question

Let $\tau_t = (-\tau^2, \dots, -\tau^2) \in \mathbb{C}^t$. Define

$$F_t^-(\tau) = K_t^-(1; \tau_t), \quad F_t^+(\tau) = 2^t K_t^+(1; \tau_t). \quad (54.1)$$

Both functions are entire in τ and even. At repeated coordinates they are the explicit $t \times t$ determinants

$$K_t^\varepsilon(1; (z, \dots, z)) = (-1)^{b_t} \det \left[\frac{1}{i!} \partial_z^i \partial_s^j f_\varepsilon(s, z) \right]_{s=1} \Big|_{0 \leq i, j < t}, \quad (54.2)$$

where $f_0 = C$ and $f_1 = S$.

Corollary 54.1 (Original uncentered formulation). *Locally uniformly for $\tau \in \mathbb{C}$,*

$$\frac{H_N((x - (2 - 2 \cosh(\tau/N)))^t)}{N^{t(t-1)/2}} = F_t^-(\tau) + O(N^{-1}), \quad (54.3)$$

$$\frac{H_N((x - (2 + 2 \cosh(\tau/N)))^t)}{(-1)^t N^{t(t+1)/2}} = F_t^+(\tau) + O(N^{-1}). \quad (54.4)$$

The endpoint values are

$$F_t^-(0) = 2^{b_t} \prod_{j=0}^{t-1} \frac{j!}{(2j)!}, \quad F_t^+(0) = 2^{t+b_t} \prod_{j=0}^{t-1} \frac{j!}{(2j+1)!}. \quad (54.5)$$

If κ_t is the corresponding triangular exponent, the first uncentered correction is explicitly

$$F_t^\pm(\tau) + \frac{t}{2N} \left(\kappa_t + \tau \frac{d}{d\tau} \right) F_t^\pm(\tau) + O(N^{-2}). \quad (54.6)$$

Proof. Put $\nu = N + t/2$. In the centered theorem the coordinates are $-(\nu/N)^2 \tau^2$, and the normalization changes by $(\nu/N)^{\kappa_t}$. Expand this analytic rescaling. The centered finite-size correction itself starts at ν^{-2} , so the entire N^{-1} term is exactly the coordinate and normalization change. All expansions are uniform on compact sets. Formula (54.5) follows from (52.3). $\square$

Thus the $O(N^{-1})$ term in the originally posed coordinates is real, but it is a centering effect. Replacing N by $N + t/2$ in both the angular scale and the normalization removes it and every later odd inverse power.

54.2 Exact spatial displacements also have an even expansion

Suppose the desired roots are $h^2 y_i$ and $4 - h^2 v_a$, rather than (46.5). Inverting ϕ_h near zero gives

$$\widehat{y} = y + \frac{h^2 y^2}{12} + O(h^4), \quad \phi_h(\widehat{y}) = y.$$

The inverse is analytic in h^2 and locally uniform in y . Therefore the exact spatial-coordinate model also has an even convergent expansion. Its first correction is

$$A_1^{\text{spatial}} = A_1 + \frac{1}{12} \left(\sum_i y_i^2 \partial_{y_i} + \sum_a v_a^2 \partial_{v_a} \right) A_0. \quad (54.7)$$

This demonstrates that evenness is not confined to a trigonometric notation for the roots. The center of the determinant size, however, is essential to the stated form.

55 Worked examples and checks on the constants

55.1 One root at the left endpoint

For $m = 1$, $\ell = 0$, $\nu = N + 1/2$ and $\kappa = 0$, the exact formula is

$$Z_N(y) = \frac{\cos \sqrt{y}}{\cos(h\sqrt{y}/2)}.$$

Hence $A_0 = \cos \sqrt{y}$ and $A_1 = (y/8) \cos \sqrt{y}$. At $y = -\tau^2$ this is $\cosh \tau$ divided by $\cosh(h\tau/2)$, and the limiting profile is $F_1^-(\tau) = \cosh \tau$. This also makes the sharp N^{-2} threshold visible without determinants.

55.2 One root at the right endpoint

For $m = 0$, $\ell = 1$,

$$Z_N(v) = \frac{2 \sin \sqrt{v}/\sqrt{v}}{\text{sinc}(h\sqrt{v}/2)}.$$

Thus $A_0 = 2 \sin \sqrt{v}/\sqrt{v}$ and $A_1 = \nu A_0/24$. The outward profile is $F_1^+(\tau) = 2 \sinh \tau/\tau$, with value 2 at zero. These two one-root examples verify both boundary normalizations.

55.3 One root at each endpoint

For $m = \ell = 1$, write $C = C(1, y)$ and $S = S(1, v)$. Then

$$A_0 = CS, \quad A_1 = \left(\frac{y}{4} + \frac{v}{6}\right) CS - \frac{1}{4} C_t(1, y) S_t(1, v). \quad (55.1)$$

The mixed term generally does not vanish. At $y = v = 0$ the exact unsigned endpoint determinant is $N + 1$, and the centered normalization with $\nu = N + 1$ is identically 1; accordingly $A_1(0, 0) = 0$.

55.4 Two coincident roots at an endpoint

The two repeated-root profiles from (54.2) simplify to

$$F_2^-(\tau) = \frac{1}{2} + \frac{\sinh(2\tau)}{4\tau}, \quad (55.2)$$

$$F_2^+(\tau) = \frac{\sinh(2\tau) - 2\tau}{\tau^3}. \quad (55.3)$$

The singularities at zero are removable; the values there are 1 and $4/3$, respectively. To derive them, evaluate the 2×2 determinant in (54.2), or differentiate the two elementary hyperbolic row functions before taking the coincident limit.

For example, with two roots at each endpoint, all exactly at the endpoints, one has

$$H_N(x^2(x-4)^2) = \frac{(N+1)(N+2)^2(N+3)}{12}. \quad (55.4)$$

This follows by inserting the endpoint jets in the four-row Christoffel determinant and factoring the resulting polynomial. Writing $\nu = N + 2$ gives $Z_N = (1 - h^2)/12$, agreeing with both (47.5) and (47.6).

55.5 A genuinely confluent two-cluster numerical example

Take $m = \ell = 2$, $\mathbf{y} = (-1, -1)$ and $\mathbf{v} = (-2, -2)$. The first two coefficients, computed from the confluent kernels, are

$$A_0 = 0.1741166983662596749147659 \dots, \quad A_1 = -0.7183984032902893460064576 \dots$$

The following are high-precision evaluations, not interval enclosures. Here $h = 1/(N + 2)$.

N	$ Z_N - A_0 $	$ Z_N - A_0 - h^2 A_1 $	$(Z_N - A_0 - h^2 A_1)/h^4$
16	$2.2027646 \cdot 10^{-3}$	$1.4514422 \cdot 10^{-5}$	1.52366601
32	$6.2030784 \cdot 10^{-4}$	$1.1440675 \cdot 10^{-6}$	1.52885861
64	$1.6484093 \cdot 10^{-4}$	$8.0651949 \cdot 10^{-8}$	1.53034944
128	$4.2503422 \cdot 10^{-5}$	$5.3595798 \cdot 10^{-9}$	1.53074959

The corrected residual displays the predicted fourth-order behavior. The proof of the error order is Theorem 47.1, not the table.

56 Computation, reproducibility, and the verification boundary

56.1 Two independent determinant paths

The accompanying code/verify.py (shipped as code/04-endpoint-confluence-verify.py) uses exact rational arithmetic for its algebraic tests. One path forms the original Catalan moment matrix from (45.1). A second path uses the fixed-size confluent Christoffel determinant. The latter obtains Taylor jets from (48.2): if $p_n(c + u) = \sum_r a_{n,r} u^r$, then

$$a_{n+1,r} = (c - 2)a_{n,r} + a_{n,r-1} - a_{n-1,r}, \quad a_{n,-1} = 0.$$

No trigonometric root extraction is needed for these exact tests. The endpoint jet identities checked separately are

$$\frac{p_n^{(r)}(0)}{r!} = (-1)^{n+r} \binom{n+r}{2r}, \quad \frac{p_n^{(r)}(4)}{r!} = \frac{2n+1}{2r+1} \binom{n+r}{2r}, \quad (56.1)$$

with zero values when $r > n$. Both follow from the recurrence, or the trigonometric formulas expanded at the endpoints.

The code also evaluates the normalized kernel jets by the convergent entire series

$$\frac{1}{q!} \partial_y^q \partial_t^r f_\varepsilon(t, y) = \sum_{\substack{k \geq q \\ 2k + \varepsilon \geq r}} \frac{(-1)^k \binom{k}{q} y^{k-q} t^{2k+\varepsilon-r}}{(2k + \varepsilon - r)!}. \quad (56.2)$$

This handles zero coordinates and repeated roots without dividing by small root differences. High-precision tests use 100 decimal digits. They include inward roots, complex roots, exact endpoints, and mixed repeated-root blocks of total degree up to eight in the endpoint coefficient checks.

56.2 Executed checks

The run recorded in data/verification.json and data/verification.txt completed the following checks.

[Merge note: those two records are shipped as data/04-endpoint-confluence-verification.json and data/04-endpoint-confluence-verification.txt. In this report the unprefix data/verification.json and data/verification.txt are Part I's recorded outputs; see Section 60 before rerunning anything.]

Check	Count
Exact original determinants versus confluent Christoffel determinants	104
Exact endpoint Taylor-jet identities	378
Exact leading constants of centered coefficient minors	35
Exact vanishing of degree-one centered minors	60
Exact second-order one-cluster minor ratios	108
Exact mixed minor ratios $-1/4$	25
High-precision first-correction residual checks	40
High-precision analytic evenness checks	6
Analytic continuation formula versus Christoffel evaluation	6
High-precision endpoint leading and correction constants	48
Total; all passed	810

The first six rows comprise 710 exact-arithmetic checks. The last four rows comprise 100 high-precision diagnostic checks. A check count measures executed identities, not distinct universal results. In particular, finite tests of the table of minors do not replace the all-parameter proof from (49.2).

56.3 Computing higher coefficients

For a specified order J , expand the finite product (49.3) through degree $2J$. Convert its separately symmetric coefficients to Schur polynomials, or directly to the coefficient minors of the centered exponential matrix. Replace the corresponding alternants by kernel determinants with the indicated increased derivative orders. Finally multiply by the truncated reciprocal series of c_h, s_h, D_h and the cross factors in (50.4). Only finite operations are needed for each J .

This is a constructive mathematical algorithm, but the shipped script is a verification implementation for the leading and first correction, not a fully optimized arbitrary-order engine. The bound in Theorem 50.2 can be applied once the desired exact coefficients or certified enclosures for them have been computed. The present high-precision implementation does not perform outward-rounded interval arithmetic.

56.4 A formalization route

No Lean or Rocq file is claimed for this article. A feasible formalization would begin with the polynomial Vandermonde identity underlying (49.2), the first coefficient table over $\mathbb{Q}$, and (51.1). Formal power series can express the exact parity and coefficient algorithm independently of analysis. A second layer would prove the Catalan moment functional and the Christoffel identity. Only the final layer needs holomorphic divided differences, uniform convergence, and the Cauchy remainder estimate. Keeping those layers separate would make the new algebra reusable for other moment families.

57 Further research questions and proposed next steps

The following are research directions, not assertions that every item is globally open in the published literature.

57.1 Nonendpoint collisions on the reciprocal-size scale

The unsolved second portion of the repository's question lets two nonendpoint roots approach at order N^{-1} . Construct a uniform counterpart of the present theorem with oscillatory sector amplitudes retained. A useful first target is two equal-multiplicity roots with spectral coordinates $ze^{\pm\tau/N}$ and $z \neq \pm 1$, allowing z on the unit circle. Here different exponential sectors can compete; the endpoint factorization alone does not settle their cancellation.

[Added 29 September 2026, batch 55: answered by Part V (Theorems 64.1 and 71.1), in the centred coordinates $2 + 2 \cos(\theta \pm \tau/(N + t))$, with $z = e^{i\theta}$ on the unit circle and the oscillatory sectors retained; off the circle one sector dominates exponentially. In the coordinates $ze^{\pm\tau/N}$ proposed here the N^{-1} term gains a non-oscillatory summand (Remark 61.1, an observation of this merge).]

57.2 Endpoint clusters in the presence of fixed interior roots

Add a fixed polynomial multiplier whose roots avoid 0 and 4. Determine the correct effective size and the influence of its boundary amplitudes on evenness. A finite perturbation of the Jacobi recurrence may alter the amplitude without changing the local oscillatory scale. An exact criterion for disappearance of odd powers would separate structural symmetry from a choice of normalization.

57.3 Higher interaction operators

The quadratic generator produces the universal mixed term $-1/4$. Compute the quartic and sextic operators in a minimal basis of Euler operators and symmetric coordinate polynomials. Determine whether the logarithm of the coefficient generator yields a particularly simple connected-interaction expansion. Its elementary factors already show that only even orders can occur, but do not automatically give the most compact differential representation.

57.4 Multiplicities growing with determinant size

The theorem keeps m, ℓ fixed. Find a regime $m = m_N, \ell = \ell_N$ in which the factorized leading profile or the first correction remains uniform. The explicit bound here is deliberately too coarse to locate a sharp growth threshold. Such a threshold should account simultaneously for the number of pair interactions, growth of the confluent kernels, and the shrinking domain on which an effective relative error is small.

[Added 29 September 2026, batch 57: partly answered by Part VI, for $\ell = 0$ and the cluster $x^m(x + \varepsilon)$ with one outward displacement $\varepsilon \geq 0$: the ratio to $H_N(x^{m+1})$ tends to 1 exactly when $\varepsilon N(N + m + 1)/(4m + 2) \rightarrow 0$, for every growth of m (Corollary 84.2). For fixed m this is the threshold $N^2\varepsilon \rightarrow 0$ of Theorem 47.4, and when $N^2\varepsilon \rightarrow y$ the limit is ${}_0F_1(; m + \frac{1}{2}; y/4)$ (93.1); for $N, m \rightarrow \infty$ the fixed- m profile is replaced by e^τ with the corrections of Theorem 89.1, on the critical scales of Theorem 83.1. Several displaced roots, $\ell > 0$, the first correction in the sense of Theorem 47.1 and an approximation uniform in $1 \leq m \leq N$ (Section 96.3) remain open (Section 81.4).]

57.5 Other Jacobi endpoint exponents

Replace the Catalan weight by $x^\alpha(4 - x)^\beta$ with $\alpha, \beta > -1$. Determine which effective size optimizes convergence, and whether an exact even analytic interpolation survives beyond the half-integer trigonometric cases. The appropriate limiting kernels are Bessel-type, so the classical hard-edge theory supplies context; the specific objective is an explicit collision-uniform coefficient and remainder calculus analogous to the one proved here.

57.6 Zeros of inward confluent profiles

For positive scaled coordinates, the kernel determinants can vanish. Classify real zeros of the one-parameter repeated-root profiles and establish simplicity or interlacing where true. Holomorphic convergence and a nonzero derivative at a limiting zero would then give quantitative locations of zeros of the finite Catalan determinant. The absolute first-correction formula remains meaningful there even when a relative formula would be singular.

57.7 Degeneration of finite recurrences

For each fixed multiplier the repository gives a finite exponential-sector recurrence description. Explain how those recurrences degenerate under the present moving-root limit and recover differential equations for the profile determinants. A rigorous bridge between coalescing characteristic roots and the centered even series would connect the repository's algebraic and asymptotic descriptions rather than treating them separately.

57.8 Sharper analytic bounds and stable numerical evaluation

Locate the nearest genuine singularities of the analytic interpolation in h , distinguishing removable singularities from denominator zeros. Use that geometry to improve (50.9) and select a practical truncation order. Develop interval implementations using divided differences or jets so that collisions are handled without loss of significance. This would convert the available analytic error estimates into end-to-end numerical certificates.

57.9 Rational modifications and Cauchy-transform blocks

Allow poles of a rational multiplier away from the integration interval, with selected poles approaching the endpoints on the same scale. Characteristic-polynomial ratio identities suggest determinant blocks involving Cauchy transforms [10, 11]. Determine which sign, centering, and evenness statements survive, and what replaces the monotonicity argument for the sharp outward stability criterion.

57.10 A machine-checked coefficient and confluence library

Formalize the centered alternant and the second-order interaction formula first, then implement a verified finite coefficient generator. A reusable library should state its divided-difference confluence conventions explicitly and track every factorial. The final target is a checked interface connecting exact rational algebra, analytic remainder bounds, and certified evaluations of the confluent kernels.

58 Conclusion

The repository's endpoint moving-root question has an explicit, collision-uniform answer for every fixed multiplicity, with a simultaneous two-endpoint extension. The principal structural fact is exact evenness in $(N + d/2)^{-1}$ after the natural sign and power are removed. The two endpoints decouple in the leading term but interact at the next order through a computable differential operator. Positive Schur coefficients turn the asymptotic profile into a sharp necessary-and-sufficient criterion for stability under outward perturbations.

The scope remains precise: the work proves the endpoint results stated here, not a general solution of all root collisions, all recurrence resonances, or hard-edge universality for arbitrary weights. The result is a conventional mathematical manuscript with explicit tests; neither formal verification nor worldwide priority is asserted.

59 A sign and normalization audit

This section is Appendix A of the manuscript.

The following elementary checks are useful when porting the formulas. For one multiplier factor, (48.5) says $H_N(x - c) = (-1)^N p_N(c)$. Thus

$$H_N(x) = 1, \quad H_N(x - 4) = (-1)^N (2N + 1).$$

For two endpoint factors the same formula and $p_n(0) = (-1)^n$, $p_n(4) = 2n + 1$ give $H_N(x(x - 4)) = (-1)^N(N + 1)$. For two repeated lower factors the derivative determinant gives $H_N(x^2) = N + 1$; for two upper factors it gives

$$H_N((x - 4)^2) = \frac{(N + 1)(2N + 1)(2N + 3)}{3}.$$

These identities include $N = 0$. They fix the asymmetric triangular powers and the factor 2^ℓ at the right endpoint. They also verify that derivatives in a repeated-root block must be divided by their factorials, whereas the derivative columns in (46.8) and (46.9) are *not* factorial-normalized.

At a tuple of all zero coordinates, the kernel coefficient matrix has entries $(-1)^i(2i + \varepsilon)_j / (2i + \varepsilon)!$. Its determinant has sign $(-1)^{b_n}$ and nonzero magnitude $2^{b_n} \prod i! / (2i + \varepsilon)!$, exactly as used in (52.3). This is a direct independent derivation of the endpoint constant without evaluating a Selberg integral.

60 Archive contents and reproduction

This section is Appendix B of the manuscript.

The archive contains the TeX source and compiled PDF, `code/verify.py`, its recorded data, a reproduction README, source notes, and a dependency file. After installing `requirements.txt`, run `python code/verify.py`; the script writes to `data/` regardless of the working directory. Exact tests use the standard library; `mpmath` is used only for numerical diagnostics. Rebuild with `latexmk -pdf article.tex`, or repeated `pdflatex` runs. No external images or custom font files are required.

[Merge note: of these, the TeX source, the 20-page PDF and the reproduction README are not shipped here; they survive in the arrival commit 148a906aa. The program, its two recorded outputs, the source notes, the proof audit and the dependency file are shipped byte-identical with the prefix

`04-endpoint-confluence-`:

- `code/04-endpoint-confluence-verify.py` (delivered as `code/verify.py`);
- `data/04-endpoint-confluence-verification.json` (delivered as `data/verification.json`);
- `data/04-endpoint-confluence-verification.txt` (delivered as `data/verification.txt`);
- `data/04-endpoint-confluence-requirements.txt` (delivered as `requirements.txt`; it pins `mpmath==1.3.0`);
- `04-endpoint-confluence-SOURCES.md` and `04-endpoint-confluence-PROOF_AUDIT.md` at the report root (delivered as `SOURCES.md` and `PROOF_AUDIT.md`).

The command above must *not* be run in this directory. Here `code/verify.py` is Part I's verifier, and the shipped Part IV program, which takes no arguments, always writes `verification.json` and `verification.txt` into the `data/` directory beside its own `code/` directory: run in place as `code/04-endpoint-confluence-verify.py`, it would overwrite Part I's recorded `data/verification.json` and `data/verification.txt`. On Windows it also writes CRLF line endings, whereas the recorded files are LF. Restore the delivered layout in a scratch copy instead (a POSIX shell, from the report directory):]

```
W=/path/to/scratch/endpoint-confluence
mkdir -p "$W/code" "$W/recorded"
cp code/04-endpoint-confluence-verify.py "$W/code/verify.py"
for f in data/04-endpoint-confluence-verification.*; do
  cp "$f" "$W/recorded/${f#data/04-endpoint-confluence-}"; done
cd "$W" && python3 code/verify.py # writes $W/data; compare with $W/recorded
```

[Run this way on 29 September 2026 with Python 3.13.5 and mpmath 1.3.0, it printed TOTAL: 810 in about two seconds, and both outputs equalled the recorded ones apart from Windows line endings. The checks are Python assertions, so do not pass -0.]

Part V. Interior root collisions

Part V was added on 29 September 2026 in batch 55 of ProveIt's incoming-report intake, from one manuscript written against the pinned README of Parts I–III and the standalone manuscripts of Parts II and IV. It answers the half of Part II's research question “Uniform endpoint and root-collision limits” (Section 24.4) that Part IV left open, and with it Part IV's first research question (Section 57.1): for any fixed number of roots colliding, on the scale N^{-1} , at fixed bases away from 0 and 4, with arbitrary collisions inside each cluster and several separated clusters, it proves an exact finite sector expansion with amplitudes convergent in $1/(N + d/2)$ uniformly through every collision, an explicit first correction and remainder bound, the phase-dependent law of two equal-multiplicity roots inside $(0, 4)$, and a convergent expansion of the moving finite-size zeros. Limits uniform up to the endpoints remain open. Parts I–IV are unchanged except for dated pointers to this addition (Sections 22 and 24.4, the introduction of Part IV, and Sections 44.4 and 57.1), a shortened line on the title page (whose vertical spaces were reduced by a further 7 mm so that it still fits on one page) and a paragraph of the abstract, a contents entry, one bibliography entry used only by Part V (and a clause in the note of another), a clause in the closing note on source dates, and three new labels, on Part II's Section 24.5, Part III's Section 40.4 and Part IV's Section 57.2, so that they can be cited. Section 61 records the provenance and fixes the notation of the addition.

61 Provenance and conventions of Part V

61.1 One manuscript, one addition

Part V is batch-55 manuscript 03, *Interior Root Collisions in Catalan Hankel Determinants*, subtitled *Convergent sector expansions, multiple clusters, and phase-dependent zero motion*, an AI-assisted research article prepared for Vladimir Reshetnikov and dated 29 September 2026 (24 A4 pages as delivered). Its visible byline reads “Prepared for Vladimir Reshetnikov / AI-assisted mathematical research” and names no assistant; the delivered PDF's metadata gives the author as “OpenAI ChatGPT; prepared for Vladimir Reshetnikov”. It arrived as the archive `ProveIt_Interior_Collision_Research.zip` (inner directory `ProveIt_Interior_Collision_Research/`, main file `article.tex`) in ProveIt commit 2d4919838 and was placed in 26473dfa0.

It was written against ProveIt commit e1afd75e35a4de734d5aa47aec5cdb917b82ed3e (the record of batch 53 in the incoming-report README), and it records no blob. Through the GitHub connector it read only this report's README there, which described three parts: at that commit this report's `article.tex` is still Git blob e5a4d81408c80f0385ae66c26fbd11bba7dfdb09, Parts I–III, the blob Part IV's own pin recorded; Part IV's text was written into it afterwards, in e300500cb. By its source notes, the manuscript took Part II's question from the standalone Part II manuscript and Part IV's scope and question from the standalone batch-53 manuscript, both read from Vladimir's file library; that manuscript's “Section 13.1” is Section 57.1 here. The body of Section 24.4 is unchanged between the pin and the placement commit apart from its label and the dated batch-53 note after it, and Section 57.1 as printed here is the passage the manuscript paraphrases (checked for this merge). So the manuscript's “repository” and “inspected repository reports” are Parts I–III as printed above, its “root-collision report” is Part II, and its “endpoint continuation” and “later manuscript” are Part IV.

The manuscript's program and recorded outputs are shipped beside the earlier files with the prefix 05-interior-collision- (Part I's unprefixed files count as 01, and Parts II–IV have 02– to 04–), its dependency file under `data/`, and its source audit and proof status, delivered as `SOURCES.md` and `PROOF_STATUS.md`, at the report root:

- `code/05-interior-collision-verify.py` (delivered as `code/verify.py`);
- `code/05-interior-collision-Makefile` (delivered at the package root as `Makefile`; it must not be run in this directory, Section 75);

- `data/05-interior-collision-verification.json` and the four CSV files `data/05-interior-collision-finite_size_corrections.csv`, `-zero_transport.csv`, `-jacobi_parity_audit.csv` and `-multiple_clusters.csv` (delivered under `data/` without the prefix; the CSV files have CRLF line endings, as delivered);
- `data/05-interior-collision-requirements.txt` (delivered as `requirements.txt`; it pins `mpmath==1.3.0`);
- `05-interior-collision-SOURCES.md` and `05-interior-collision-PROOF_STATUS.md`.

All are byte-identical to the delivery. The manuscript, its PDF, its delivery README and its checksum list are not shipped; they survive in the arrival commit (the checksum list was verified at placement and retired).

The manuscript contributes, in the order printed here: the repository question, what is classical, and why phases must be retained (Section 62); the Catalan normalization and a self-contained Christoffel reduction (Section 63); the one-cluster theorem, an exact sector expansion with convergent collision-uniform amplitudes, and the values and symmetries of its universal profile (Section 64); its proof by centred exponential alternants (Section 65); the first-correction operator and an all-orders coefficient construction (Section 66); an explicit Cauchy remainder (Section 67); several separated clusters (Section 68); the dominant sectors inside the moment interval (Section 69); sine-kernel duality and Gram positivity (Section 70); the complete first-order law of two equal-multiplicity roots and its identification with the oscillatory Jacobi (kissing-polynomial) determinant (Section 71); the profile at large real frequency (Section 72); the transport of finite-size zeros and its convergent Laurent-phase expansion (Section 73); explicit examples, including periodic and quasiperiodic scaled zero motion (Section 74); its checks and their evidentiary status (Section 75); what is resolved and what is not (Section 76); nine research directions (Section 77); a conclusion; a proof-dependency map (Section 79); and a normalization checklist (Section 80). Every result, proof, example, remark, limitation and question of the manuscript is printed. Where it re-derives something Parts I–IV already prove, the overlap is marked (Section 61.5).

The manuscript's own abstract reads:

The Catalan-Hankel reports in PROVEIt ask for uniform limits when roots of a polynomial multiplier collide away from the endpoints of the moment interval. We resolve this fixed-degree, reciprocal-size collision problem, including arbitrary internal coincidences and several separated clusters. With effective size $\nu = N + d/2$, the determinant has an exact finite sum of exponential sectors whose amplitudes are holomorphic in ν^{-1} . We give every leading amplitude, a closed first-correction operator, an all-orders coefficient construction, and an explicit compact-uniform Cauchy remainder bound. For two equal-multiplicity roots colliding inside $(0, 4)$, the leading limit is a classical oscillatory Jacobi Hankel determinant, but its first correction retains the phase $\sin(2\nu\theta)$. At each simple limiting zero we obtain a convergent phase-dependent expansion of the finite-size zero, not merely a formal asymptotic series. A completely explicit two-simple-root case exhibits periodic or quasiperiodic scaled zero motion. Classical Christoffel, sine-kernel, and kissing-polynomial results are credited and rederived where needed. The written proofs are accompanied by 392 exact-arithmetic and 126 high-precision checks.

Its title page carried this scope box:

Scope and status. The resolved question is the nonendpoint moving-root problem explicitly proposed in the repository and its endpoint continuation. This is an unrefereed manuscript, not a Lean or Rocq formalization. The contribution is the proved uniform finite-size and zero-transport package; no claim of a first discovery of classical bulk limits, kissing-polynomial parity, or worldwide priority for every formula is made.

Status (manuscript 03). This is an unrefereed, AI-assisted manuscript with conventional written proofs, not a Lean or Rocq formalization and not an interval-certified computation. It resolves the nonendpoint moving-root question of Sections 24.4 and 57.1 for fixed multiplier degree and fixed nonendpoint bases. Classical ingredients are credited, not presented as new: the Catalan moment measure and its orthogonal polynomials, the Christoffel polynomial-modification identity, its bilinear kernel variant and divided-difference confluence (Krattenthaler [3]; re-proved in the present normalization), determinants of orthogonal-polynomial kernels and bulk (sine-kernel) universality for characteristic-polynomial correlations (Strahov–Fyodorov [10], context only), and the oscillatory Jacobi Hankel determinant of the kissing-polynomial literature with its even-degree nondegeneracy, odd-degree degeneracy and large-frequency analysis (Celsus–Deaño–Huybrechs–Iserles [13]), whose versions are independently proved here in the needed normalization; the centred exponential Vandermonde is an elementary repackaging of the ordinary one. The claimed contribution relative to the inspected reports is the collision-uniform convergent amplitude construction, its explicit first coefficient and compact Cauchy bound, the separated-cluster constants, and the convergent phase-dependent finite-size zero expansion. The literature audit was bounded: the full text of the kissing-polynomial paper could not be retrieved, so no theorem-by-theorem comparison with it was made, and worldwide priority for every identity, coefficient or corollary is not established. Not claimed: uniformity as a base approaches an endpoint or another base; growing degree, unbounded scaled coordinates varying with N , or zero indices growing with N ; a classification of nonsimple zeros or of all recurrence resonances; constant-coefficient recurrences for a moving multiplier q_N ; validation of every theorem or file of the repository. No theorem of the unrefereed source reports is used as an unproved premise. The 518 recorded assertions (392 exact, 126 at 100 digits) are finite audits; the CSV experiments are not certificates, and numerical root decimals are not certified isolating intervals. No proof assistant was run, nothing in Part V is formalized, and its placement beside a Lean project confers no formal status. (Sources: the manuscript, its delivery README, and its source audit and proof status, 05-interior-collision-SOURCES.md and 05-interior-collision-PROOF_STATUS.md.)

61.2 Reading conventions

Inside Sections 62–80, “this paper”, “the present manuscript” and “here” mean manuscript 03 as printed in Part V; “the repository”, “the inspected repository reports” and “the report” mean Parts I–III at the pin e1afd75e3 (its “root-collision report” is Part II), and “the endpoint continuation” means Part IV. The manuscript cites Part II’s Section 24.4 by this report’s number, which did not change (Section 24.4); its citation of the standalone Part IV manuscript’s Section 13.1 is printed as a cross-reference to Section 57.1. “The package” means the delivered archive, whose files are shipped under the prefixed names listed in Section 61.1. The manuscript’s two appendices are printed as Sections 79 and 80, so that Part I’s appendices keep their letters. Its Sections 1–17 are Sections 62–78, so a manuscript number $n.j$ of a section, theorem, lemma, proposition, corollary, example, remark or equation is printed as $(n + 61).j$ (for example, its Theorem 3.1 is Theorem 64.1 and its Theorem 10.1 is Theorem 71.1), and its Appendices A and B are Sections 79 and 80; its two data tables are Tables 5 and 6. The shipped 05-interior-collision-PROOF_STATUS.md and 05-interior-collision-SOURCES.md use the manuscript’s numbers. The manuscript cites Krattenthaler’s Ramanujan Journal paper (with arXiv:2101.04225v5) and the Strahov–Fyodorov paper; they are Part I’s reference [3] and Part IV’s reference [10]. Bracketed paragraphs beginning “Merge note” are this merge’s, not the manuscript’s.

61.3 Notation of Part V

Part V keeps the manuscript’s symbols. Within Sections 62–80 every symbol has its Part V meaning, except in merge notes and where a sentence names Part I, II, III or IV or cites one of their equations. The multiplier q_N has only moving roots; with all of them at one fixed base it is Part II’s $q = \gamma \prod_i (x - c_i)^{t_i}$ with $\gamma = 1$, $m = \ell = 0$, and $H_N(q)$ is Part II’s (14.1). Table 4 lists the symbols that a reader of Parts I–IV

is likely to misread. One symbol was renamed: the manuscript's $\text{shc } z = \sinh z/z$ is printed $\text{sinhc } z$, Part IV's name for the same function; and the manuscript's upright Re is printed $\Re$. No normalization changed. Four readings need particular care: $D = \binom{d}{2}$ is *not* Part II's $D = \deg q$, which is Part V's d ; the spectral base w is not Part III's centred size $w = N + t/2$; the letter t is a formal branch-counting variable in (65.3) and in the proof of Theorem 68.1 (there t_a), but the multiplicity of each root of a symmetric pair from Section 71 on; and Z_N in (73.1) is normalized by $A_t(\theta)v^{t^2}$, not like Part IV's Z_N (46.7).

Table 4: Symbols and notions of Part V. The last column gives the meaning of the same letter in Part I, II, III or IV, or the tempting false reading.

Symbol or notion	Meaning in Part V	Not to be confused with
$H_N(q), p_n, \mu$	the weighted determinant (62.1); the monic orthonormal polynomials (63.2) and the Catalan measure (63.1)	the same as Part II's (14.1), p_n, μ and Part IV's; Part I's $H_N^{(m)} = H_N(x^m)$ (a shift) and Part III's $H_N^{(t)}(c) = H_N((x-c)^t)$ (a multiplicity).
$d, k, r = d - k, D, e_k$	degree of the multiplier; sector index; $\binom{d}{2}$; $\binom{k}{2} + \binom{r}{2}$ (64.3)	d is Part IV's d and Part II's $D = \deg q$; D here is <i>not</i> Part II's D ; Part I's $d = \binom{m}{2}$; Part II's sector degrees d_k and labels k .
$w, c = 2 + 2 \cosh w$; θ with $w = i\theta$	spectral coordinate of the base, $w \notin \pi i\mathbb{Z}$ (64.1); the base angle inside $(0, 4)$	$z = e^w$ is Part II's and Part III's z ($c = 2 + z + z^{-1}$); Part III's centred size $w = N + t/2$; Part II's nodes w_a ; Part IV's auxiliary w_a .
$\nu = N + d/2, h = 1/\nu$	centred size and its inverse ($\nu = N + t$ for a pair, $d = 2t$); the expansion is in h , not h^2	Part IV's ν (same formula, but even expansions there); Part II's block multiplicities ν_u ; Part I's moments ν_{i+j} ; Part III's $h = t - k$.
$u_i, U = \sum u_i, \mathcal{E}$	scaled coordinates (64.2), their sum, the Euler operator (66.1)	Part IV's auxiliary u_i , sum U and Euler operators $\mathcal{D}_\pm$; Part III's values U_a ; the shift E .
$S_{d,k}(\mathbf{u}), B_{d,k}(w)$	universal profile (64.4); nonzero base constant (64.6)	Part IV's $S(t, y)$; Part II's $S = 2^D$; Part III's subsets S ; Part II's $B_0(m)$, $B_4(\ell)$; Part III's product $B_{a,b}(N)$; Part IV's $B_{m,\ell,R}$.
$F_k(h; \mathbf{u}, w), f_{k,j}; R_k, L(h; \mathbf{u}, w)$	sector amplitudes and Taylor coefficients (64.8); numerator and denominator of (65.5)	Part IV's profiles $F_t^\pm$; Part III's $L = \text{ord}(z^2)$ and order $R(t, c)$; Part II's bound R ; L_j below.
$\alpha_j, \Delta, \text{sinhc}$	centred exponents $j - (d - 1)/2$ (65.1); the Vandermonde $\prod_{i < j} (v_j - v_i)$ (63.5); $\sinh z/z$	the same as Part IV's; Part III's arcsine measure α ; Part II's Δ_q , with multiplicities.
ϵ, ρ, R, M_k	radius data and bound of Theorem 67.1	Part IV's ρ, R, r of Theorem 50.2; Part III's ratio ρ (34.5).
$s, d_a, k_a, r_a, \mathbf{k}, T_{ab}$	number of clusters, their sizes, sector vector, cross factor (68.3)	Part II's s (number of interior roots) and $T = \sum t_i$; Part III's $T = \binom{t}{2}$; Part IV's bound T .
$K_M(x, y)$	kernel $\sum_{\ell \leq M} p_\ell(x)p_\ell(y)$ (70.1)	Part I's $K = (m - 1)^2$; Part III's $K_t(z)$; Part IV's kernels K_m^-, K_ℓ^+ .
$t; (i\tau)^t$	multiplicity of each root of the pair (71.1), as in Part III; but a formal variable in (65.3); in a tuple, $(i\tau)^t$ means t repeated entries, not a power	Part IV's kernel variable t ; Part I's generating-function variable.
$\tau, \tau_*, \tau_N, \delta_N$	collision coordinate on the scale $1/\nu$; a simple profile zero; the moving finite-size zero and its displacement	Part II's and Part IV's τ in $2 \pm 2 \cosh(\tau/N)$ (a hyperbolic coordinate: Section 57.1's $ze^{\pm\tau/N}$ is Part V's τ times $-i$, Remark 61.1); Part IV's τ_t .

Symbol or notion	Meaning in Part V	Not to be confused with
$s_t, A_t(\theta), \Phi_t, \Psi_t$	constant (71.2), normalization (71.3), leading and adjacent profiles (71.4)–(71.5)	Part III's sign s_N ; Part II's amplitudes $A_{\pm}(z)$; Part III's A_u ; Part IV's A_j ; Part I's A_m ; Part II's annihilator Ψ_q .
$m_j(\tau), Z_t; m$ in $d = 2m, t = 2m + 1$	oscillatory moments (71.8) and the constant (71.11); half the degree in Sections 69 and 72	Part I's shift m ; Part II's and Part IV's m (roots at 0); Part III's centred moments μ_n .
$Z_N(\tau), \mathcal{Z}(h, \zeta, \tau), \zeta, b_\ell$	normalized determinant (73.1), its interpolation (73.3) in the phase $\zeta = e^{2i\nu\theta}$, the zero coefficients (73.5)	Part IV's Z_N (46.7) and $Z(h; \mathbf{y}, \mathbf{v})$; Part I's parameter b ; Part IV's $b_r = \binom{r}{2}$.
L_j, E, Q_t	$\prod_{\ell \leq j} \ell!(\ell - 1)!$ (72.1); exponent and constant of Proposition 72.2	$L(h; \mathbf{u}, w)$ above; Part III's L ; Part I's $E_m(N)$; Part III's quadratic factors Q_a .
$\mathbf{x}, \mathbf{y}$	tuples of Theorem 70.2; integration variables	Part IV's coordinates $\mathbf{y}, \mathbf{v}$.

61.4 What Part V answers, and what stays open

Part II's Section 24.4 asked, besides the endpoint limits that Part IV supplied, for a second version in which two distinct nonendpoint roots approach each other at rate N^{-1} , with a uniform confluent limit and error bounds; Part IV's Section 57.1 repeated it, asking for oscillatory sector amplitudes to be retained and proposing as a first target two equal-multiplicity roots with spectral coordinates $ze^{\pm\tau/N}$, $z \neq \pm 1$, z on the unit circle allowed. Part V answers both, for fixed total degree and fixed nonendpoint bases, inside a larger statement: any number of roots in a cluster, any collision pattern (Theorem 64.1), with the explicit first coefficient (Theorem 66.1), an explicit remainder bound (Theorem 67.1) and several separated clusters (Theorem 68.1). For two simple roots the determinant is explicit (Example 74.1). For two roots of multiplicity t on the unit circle the limit is Φ_t , an oscillatory Jacobi Hankel determinant, and the ν^{-1} term keeps the phase $\sin(2\nu\theta)$ (Theorem 71.1); the value at $\tau = 0$ is (74.7); off the unit circle one sector dominates exponentially (the paragraph after Theorem 64.1). Unlike Part IV's, the expansion is not even (the remark closing Section 66). The same theorems settle, for collisions away from the endpoints, the caveat that closes Part II's Section 22: the double-scaling limit is proved directly, without (22.1).

The coordinates differ from those proposed in Section 57.1. The manuscript states its pair law only in centred coordinates; the following translation is this merge's.

Remark 61.1 (The coordinates of Section 57.1; an observation of this merge). Fix $t \geq 1$ and $0 < \theta < \pi$, and put $\nu = N + t$. For the uncentred pair $q_N = (x - 2 - 2\cos(\theta + \tau/N))^t (x - 2 - 2\cos(\theta - \tau/N))^t$, uniformly on compact sets of complex τ ,

$$\frac{H_N(q_N)}{A_t(\theta)\nu^{t^2}} = \Phi_t(\tau) + \frac{1}{\nu} \left[t\tau\Phi'_t(\tau) + (-1)^t \frac{\sin(2\nu\theta)}{\sin\theta} \Psi_t(\tau) \right] + O(\nu^{-2}).$$

Indeed $\theta \pm \tau/N = \theta \pm \tau'/\nu$ with $\tau' = \tau\nu/N = \tau + t\tau/\nu + O(\nu^{-2})$, and Theorem 71.1 holds uniformly with derivatives on a slightly larger compact set, so Taylor expansion of $\Phi_t(\tau')$ at τ gives the extra summand, while $\Psi_t(\tau')/\nu$ changes only at order ν^{-2} . With $z = e^{i\theta}$, the spectral coordinates $ze^{\pm\sigma/N}$ of Section 57.1 give the roots $2 + 2\cosh(i\theta \pm \sigma/N) = 2 + 2\cos(\theta \mp i\sigma/N)$, that is, $\tau = i\sigma$ (the profiles are even, so the sign is immaterial). So in the coordinates proposed there the ν^{-1} term has the non-oscillatory summand $t\tau\Phi'_t(\tau)$ besides the oscillatory one. As in Part IV's Corollary 54.1, where the N^{-1} term is only the centring, the extra summand is produced by the centring alone, here of the coordinate. Normalizing by N^{t^2} instead of ν^{t^2} would add a further $t^3\Phi_t(\tau)/\nu$. (Checked numerically for this merge against direct determinants for $t = 1, \theta = 1.1, \tau = 0.8$ and $N = 20, 40, 80$: ν times the residual of Theorem 71.1's terms stays near $t\tau\Phi'_t(\tau) \approx -0.65$, and ν^2 times the residual after the extra summand stays bounded; the computation is not shipped.)

Not answered. Nothing in Part V is uniform as a base approaches 0 or 4; the bridge to Part IV is Part V's own first research question (Section 77.1), so Section 24.4 as a whole, with limits uniform across the whole spectrum, is still open there. Part IV's second research question, endpoint clusters with fixed interior roots (Section 57.2), is generalized to moving interior clusters in Part V's Section 77.2, not answered. Part III's research directions on endpoint factors coupled to a cyclotomic interior root (Section 40.2) and on uniform limits near a disappearing characteristic factor (Section 40.4) are not discussed by the manuscript; its theorems hold at every nonendpoint base, cyclotomic ones included, but whether they describe the transition asked for there is not examined. Part V's Section 77.7 restates, with a sharper target, Part II's research question on other eventually constant Jacobi recurrences (Section 24.5), as the manuscript says.

Consistency with Parts II–IV (observations of this merge).

1. At $\mathbf{u} = \mathbf{0}$, one root of multiplicity d at $c = 2 + z + z^{-1}$, $z = e^w$, the sector k of Theorem 64.1 is $((-1)^d z^{2k-d})^N$ times $z^{(2k-d)d/2} \nu^{k(d-k)} F_k(\nu^{-1}; \mathbf{0}, w)$, whose leading amplitude $f_{k,0} = B_{d,k}(w) S_{d,k}(\mathbf{0})$ is nonzero by (64.10): the rate $(-1)^d z^{2k-d}$ and the growth $N^{k(d-k)}$ are the rate and the exact sector degree of Part II's Corollary 14.3. (The leading constants were not compared symbolically with Part II's Theorem 18.1 by this merge.)
2. At the central base $\theta = \pi/2$ ($c = 2$), (74.7) has no ν^{-1} term, since $\sin(\nu\pi) = 0$ for the integer $\nu = N + t$. This agrees with Part III's central product, $H_N((x-2)^{2t}) = B_{t,t}(N) = \prod_{i,j=1}^t (N+i+j-1)/(i+j-1)$ (Theorem 30.4): its factors are $\nu + (i+j-1-t)$, symmetric about ν , so it is a polynomial in ν of the parity of t^2 , and its leading coefficient $\prod_{j<t} j!^2 / \prod_{j<2t} j!$ equals $A_t(\pi/2)$ by (71.2)–(71.3). For $t = 1, 2$ it is ν and $\nu^2(\nu^2 - 1)/12$.
3. Lemma 63.1 is Part II's Lemma 15.1 for $\gamma = 1$ and Part IV's Lemma 48.1, of which Part III's Lemma 31.1 and Part I's Lemma 4.1 are special cases; with the Catalan measure, the monic orthonormal polynomials and their trigonometric form (63.3) (Part II's (15.3), Part IV's (48.3)) it is a fifth self-contained route. Lemma 65.1 and the sinh c quotients of (65.4) use the centring $\alpha_j = j - (d-1)/2$ of Part IV's exact centred alternant (Lemma 49.1), which the manuscript does not cite.
4. Tables 5 and 6 are rounded subsets (rows $N = 40, 160, 320$) of the shipped CSV files `finite_size_corrections.csv` and `zero_transport.csv` (prefixed `05-interior-collision-` under `data/`), which also have rows $N = 80$; a rerun reproduces both files byte for byte (Section 75).

61.5 Where the merge had to choose

1. *Placement.* Part V is printed after Part IV and before Part I's appendices, so that no section, equation, theorem or table number of Parts I–IV moves (checked against the `.aux` file of a build of the committed text). Its sections continue the numbering (Sections 61 to 80), with the concordance of Section 61.2.
2. *Labels.* Every label of Part V carries the prefix `ic:`; without it, six of the manuscript's labels, `thm:main`, `sec:main`, `sec:examples`, `eq:measure`, `eq:R` and `eq:Jacobi`, would have collided with Part I's bare labels. Labels were added to the manuscript's four unlabelled sections (its Section 2, its conclusion and its two appendices) and to three of its research subsections, and to Part II's Section 24.5, Part III's Section 40.4 and Part IV's Section 57.2 so that they can be cited; no existing label was changed.
3. *Overlaps with Parts I–IV.* The manuscript re-derives, without assuming the report correct, the Catalan measure, the orthogonal polynomials and the Christoffel identity (item 3 above). Printed with a pointer and kept. Everything else (the collision-uniform sector expansion at nonendpoint bases, its first correction, bound and multi-cluster extension, the pair law and the zero transport) has no counterpart in Parts I–III, which are fixed-parameter, nor in Part IV, which is at the endpoints. No statement of Parts I–IV was changed.
4. *Bibliography.* The manuscript's Krattenthaler and Strahov–Fyodorov references are [3] and [10]; its kissing-polynomial reference is added to the common bibliography as [13]. Its three references to the

pinned repository report, the standalone Part II manuscript and the standalone Part IV manuscript are replaced by Parts I–IV themselves and the pin recorded in Section 61.1; the sentences that cited them now read “(Parts I–II above)”, “(Part IV above)” after the Part IV title, “its Section 57.1” for “its Section 13.1”, “(Part IV, Theorem 47.1)” and “(Part II, Section 24.5)”; “Part II, Section 24.4” is printed as a cross-reference to Section 24.4, with the same number. The manuscript’s bibliography entry for the pinned report called it “Parts I–III” although the manuscript also relies on Part IV through the standalone manuscript; the replacement makes this moot.

5. *File names.* Merge notes in Sections 62, 75 and 79 give the shipped names beside the delivered ones. The manuscript’s reproduction commands are printed as delivered but must not be run in place here: the program’s default output directory would overwrite Part I’s recorded outputs, and the delivered Makefile, which the placement commit shipped as `code/05-interior-collision-Makefile`, acts on Part I’s verifier and on this report’s `article.tex`; a recipe on a scratch copy follows them.
6. *Front matter.* The manuscript’s title page, abstract, scope box and contents are replaced by Section 61.1, which prints the abstract, the scope box and the status.
7. *Notation and macros.* The manuscript’s symbols are kept, with the scoping convention and Table 4 of Section 61.3. Its `\shc` is printed with the report’s `\sinhc` (the same function), and its `\Rea` (upright “Re”) as `\Re`. Its `\dd` is `\,`, `\mathrm d`, the report’s is `\mathrm d`, so every `\dd` of Part V is printed `\,`, `\dd`, with the manuscript’s spacing. Its macros `\RR`, `\bx` and `\EE` were added to the preamble with the manuscript’s definitions (`\EE` is $\mathcal{E}$, not the report’s shift E); its unused `\HH` (the report’s `\HH` takes two arguments) was dropped; its `\repo` is printed `PROVEIT`; the others are the report’s with the same definitions. Its `tcloft`, `tocloft` and `xurl` packages are not needed. No mathematical text of the manuscript was changed.

62 The repository question and the contribution

62.1 A specific unresolved continuation

Let $C_n = \binom{2n}{n}/(n+1)$ be the Catalan numbers. For a polynomial $q(x) = \sum_r q_r x^r$, define

$$H_N(q) = \det \left[\sum_r q_r C_{i+j+r} \right]_{0 \leq i, j < N}, \quad H_0(q) = 1. \quad (62.1)$$

The report *Shifted Catalan Hankel Polynomials* in `PROVEIT` includes an analysis of fixed polynomial multipliers, their exponential sectors, and their sharp constant-coefficient recurrences (Parts I–II above). Part II, Section 24.4, proposes uniform limits when roots move with N . Its nonendpoint version asks what happens when two roots approach one another at order N^{-1} . Fixed-parameter asymptotics alone do not answer this question: their separate coefficients can become singular as the roots collide, while the determinant itself remains regular.

The later manuscript *Two-Endpoint Confluence for Catalan Hankel Determinants* (Part IV above) addresses the endpoint part of that program, but explicitly leaves the nonendpoint problem open in its Section 57.1. It suggests spectral coordinates $ze^{\pm\tau/N}$, with $z \neq \pm 1$, and asks for uniformity with the oscillatory sectors retained. This is the problem solved here. We treat all fixed multiplicities, arbitrary collision patterns within a cluster, and multiple separated nonendpoint clusters, rather than just two simple roots.

The repository README used for scope comparison was inspected at commit

e1afd75e35a4de734d5aa47aec5cdb917b82ed3e.

The exact question and predecessor scope were also checked in the standalone manuscripts available in the user’s file library. The source audit is recorded in `SOURCES.md`. This is a targeted continuation, not an assertion that every file or every incoming report in a changing repository has been exhaustively compared.

[Merge note: the commit above is the pin recorded in Section 61.1; at that commit this report's `article.tex` held Parts I–III only. The “standalone manuscripts” are those of Parts II and IV, and `SOURCES.md` is shipped as `05-interior-collision-SOURCES.md`.]

62.2 What is established here, and what is classical

Our first principal result is Theorem 64.1. It replaces singular fixed-root formulas by an exact finite sum of sectors with convergent analytic amplitudes. The centering $\nu = N + d/2$ is forced by the Christoffel columns. Theorem 66.1 evaluates the complete first coefficient using an Euler differential operator, while Theorem 67.1 supplies an explicit absolute remainder. Theorem 68.1 extends the construction to several separated clusters. These statements include collisions and coordinate derivatives.

The second principal result concerns zeros. For a symmetric pair of multiplicity- t roots, Theorem 71.1 identifies the first phase-dependent correction. Theorems 73.1 and 73.2 then give the zero displacement and a convergent expansion whose coefficients are Laurent polynomials in the oscillatory phase. Example 74.1 shows explicitly why an ordinary N -independent first coefficient can fail to exist.

Several ingredients are classical and are not presented as new. Krattenthaler gives the polynomial-modification determinant identity, its confluent interpretation, and recurrence applications [3]. Determinants of orthogonal-polynomial kernels and bulk universality for characteristic-polynomial correlations are established topics; see Strahov–Fyodorov [10]. The oscillatory moment determinant with weight $e^{i\omega x}$ on $[-1, 1]$ is the subject of the kissing-polynomial literature, including even-degree nondegeneracy, odd-degree degeneracy, and large-frequency asymptotics [13]. We explicitly identify our limiting profile with that family. Its parity and sine-kernel interpretations are context and tools, not claims of a newly discovered special function or a newly solved classical parity problem.

The bounded literature audit did not establish worldwide novelty for all finite-size coefficients. The precise claim is that the following proofs solve the stated repository continuation and give an explicit, self-contained package that is not supplied by the inspected fixed-parameter and endpoint reports. No unverified theorem from those reports is needed in the proofs below.

62.3 Why the answer has to retain phases

Two effects coexist. Within a collision cluster the Vandermonde in the root locations has small factors, producing powers of N after cancellation. Independently, every orthogonal polynomial has two spectral branches. If the base root is inside $(0, 4)$, both branches have the same absolute size and their phases cannot be discarded. For an even total multiplicity there is one largest power of N , so the leading limit can be phase-free; adjacent sectors are only one power smaller and reappear in the first correction. For odd total multiplicity two sectors already tie at leading order. The exact analytic decomposition keeps all of this information without dividing by a potentially zero limiting profile.

63 Catalan normalization and the classical determinant reduction

63.1 The moment measure and its polynomials

Put

$$d\mu(x) = \frac{1}{2\pi} \sqrt{\frac{4-x}{x}} \mathbf{1}_{(0,4)}(x) dx. \quad (63.1)$$

A beta integral, or the substitution $x = 4s$, gives $\int x^n d\mu(x) = C_n$. Let

$$p_0(x) = 1, \quad p_1(x) = x - 1, \quad p_{n+1}(x) = (x - 2)p_n(x) - p_{n-1}(x) \quad (n \geq 1). \quad (63.2)$$

These polynomials are monic and have squared norm 1. For completeness, write $x = 2 + 2 \cos \theta$, $0 < \theta < \pi$. Then

$$d\mu = \frac{2}{\pi} \sin^2(\theta/2) d\theta, \quad p_n(2 + 2 \cos \theta) = \frac{\sin((n + \frac{1}{2})\theta)}{\sin(\theta/2)}. \quad (63.3)$$

Orthogonality and the norms follow immediately from the orthogonality of the half-integer sine functions on $(0, \pi)$. Analytic continuation gives

$$p_n(2 + 2 \cosh w) = \frac{e^{(n+1/2)w} - e^{-(n+1/2)w}}{2 \sinh(w/2)}. \quad (63.4)$$

At singular spectral coordinates the right side is interpreted by continuity, but the main theorems keep the base away from them.

63.2 Vandermonde convention and collisions

Throughout,

$$\Delta(v_1, \dots, v_d) = \prod_{i < j} (v_j - v_i). \quad (63.5)$$

An empty product or determinant is 1. The quotient of an alternating holomorphic function by Δ extends holomorphically across all coincidences. For a determinant of row evaluations, the extension at a block of m repeated nodes v is obtained by replacing its rows with

$$\frac{1}{a!} \partial_v^a (\text{row at } v), \quad 0 \leq a < m, \quad (63.6)$$

and retaining the cross-block Vandermonde factors. We use these *normalized* derivatives everywhere. Omitting the factorials would change the constants in several theorems.

Lemma 63.1 (Christoffel reduction, in the present normalization). *For $q(x) = \prod_{i=1}^d (x - c_i)$,*

$$H_N(q) = (-1)^{Nd} \frac{\det[p_{N+j}(c_i)]_{1 \leq i \leq d, 0 \leq j < d}}{\Delta(c_1, \dots, c_d)}. \quad (63.7)$$

The formula includes arbitrary repeated roots by (63.6).

Proof. Andréief's determinant integration identity, obtained by expanding two determinants and integrating each term, gives

$$H_N(q) = \frac{1}{N!} \int_{(0,4)^N} \Delta(\mathbf{x})^2 \prod_{a=1}^N q(x_a) \prod_{a=1}^N d\mu(x_a).$$

For distinct c_i ,

$$\Delta(x_1, \dots, x_N, c_1, \dots, c_d) = \Delta(\mathbf{x}) \Delta(\mathbf{c}) \prod_{a,i} (c_i - x_a).$$

Replace the monomial columns in this augmented Vandermonde by the monic polynomials $p_0, \dots, p_{N+d-1}$; its determinant is unchanged. Multiply by $\Delta(\mathbf{x})$, likewise written using $p_0, \dots, p_{N-1}$, and integrate. Orthogonality forces the first N columns to pair with the integration rows and contributes $N!$, since all squared norms are 1. The surviving minor is $\det[p_{N+j}(c_i)]$. The product $\prod_{a,i} (c_i - x_a) = (-1)^{Nd} \prod_a q(x_a)$ supplies the sign. This proves (63.7). Polynomial continuity, equivalently the holomorphic divided-difference construction, proves the confluent case. $\square$

64 The one-cluster theorem

64.1 Coordinates and universal profiles

Fix an integer $d \geq 1$ and a spectral coordinate

$$w \in \mathbb{C} \setminus \pi i \mathbb{Z}. \quad (64.1)$$

Equivalently, the base point $c = 2 + 2 \cosh w$ is neither 0 nor 4. Set

$$\nu = N + \frac{d}{2}, \quad h = \nu^{-1}, \quad c_i(h) = 2 + 2 \cosh(w + hu_i), \quad q_N(x) = \prod_{i=1}^d (x - c_i(h)). \quad (64.2)$$

Thus $c_i(h) = c + 2 \sinh(w)hu_i + O(h^2)$ uniformly on compact coordinate sets. The nonzero derivative $2 \sinh w$ shows that this describes the ordinary reciprocal-size collision scale in the root variable as well.

For $0 \leq k \leq d$, write

$$r = d - k, \quad D = \binom{d}{2}, \quad e_k = \binom{k}{2} + \binom{r}{2} = D - kr. \quad (64.3)$$

The universal profile is

$$S_{d,k}(\mathbf{u}) = \sum_{\substack{A \subseteq [d] \\ |A|=k}} \frac{\exp(\sum_{a \in A} u_a - \sum_{b \notin A} u_b)}{\prod_{a \in A, b \notin A} (u_a - u_b)}. \quad (64.4)$$

The individual summands need distinct coordinates, but their sum does not. Indeed, Laplace expansion yields

$$S_{d,k}(\mathbf{u}) = (-1)^{kr} \frac{\det \left[u_i^j e^{u_i} \ (0 \leq j < k) \mid u_i^j e^{-u_i} \ (0 \leq j < r) \right]_{1 \leq i \leq d}}{\Delta(\mathbf{u})}. \quad (64.5)$$

The vertical bar separates blocks of columns. Consequently $S_{d,k}$ is an entire symmetric function of all coordinates, including when $k = 0$ or $r = 0$. In particular $S_{d,0} = e^{-\sum u_i}$ and $S_{d,d} = e^{\sum u_i}$.

Define the nonzero base constant

$$B_{d,k}(w) = \frac{(-1)^{r(r+1)/2}}{(2 \sinh(w/2))^d (2 \sinh w)^{e_k}}. \quad (64.6)$$

Theorem 64.1 (Convergent collision-sector expansion). *There are functions $F_k(h; \mathbf{u}, w)$, $0 \leq k \leq d$, holomorphic near $h = 0$ locally uniformly in $\mathbf{u} \in \mathbb{C}^d$ and in nonendpoint w , for which the following identity is exact for all sufficiently large N :*

$$H_N(q_N) = (-1)^{Nd} \sum_{k=0}^d e^{(2k-d)\nu w} \nu^{k(d-k)} F_k(\nu^{-1}; \mathbf{u}, w). \quad (64.7)$$

Each amplitude is symmetric in $\mathbf{u}$ and has a genuinely convergent Taylor series

$$F_k(h; \mathbf{u}, w) = \sum_{j=0}^{\infty} f_{k,j}(\mathbf{u}, w) h^j, \quad f_{k,0} = B_{d,k}(w) S_{d,k}(\mathbf{u}). \quad (64.8)$$

The coefficient functions are entire in $\mathbf{u}$. On every compact coordinate set the expansion and any fixed number of coordinate derivatives converge uniformly for sufficiently small $|h|$. After retaining $j \leq J$, the absolute remainder in (64.7) is bounded by a constant times

$$\sum_{k=0}^d |e^{(2k-d)\nu w}| \nu^{k(d-k)-J-1}. \quad (64.9)$$

All these assertions include arbitrary internal collisions.

The theorem is deliberately stated with an absolute remainder. A relative estimate obtained by dividing by a leading profile would be false or meaningless at its zeros. Outside the moment interval a single spectral direction may dominate exponentially; on the unit circle the retained sectors can interfere. Formula (64.7) handles both cases.

Proposition 64.2 (Values and elementary symmetries of the profile). *With $r = d - k$,*

$$S_{d,k}(\mathbf{0}) = 2^{kr} \frac{\prod_{j=0}^{k-1} j! \prod_{j=0}^{r-1} j!}{\prod_{j=0}^{d-1} j!} > 0, \quad (64.10)$$

$$S_{d,k}(\mathbf{u} + a\mathbf{1}) = e^{(2k-d)a} S_{d,k}(\mathbf{u}), \quad (64.11)$$

$$S_{d,d-k}(\mathbf{u}) = S_{d,k}(-\mathbf{u}). \quad (64.12)$$

Proof. The last two identities follow directly from (64.4) on the distinct-coordinate locus, and then everywhere by analyticity. For (64.10), use (64.5) with the normalized derivatives of orders $0, \dots, d-1$ at zero. Before dividing the rows by their factorials, the determinant is the confluent Vandermonde at the two spectral values $+1$ and -1 : it equals $(-1)^{kr} \prod_{j < k} j! \prod_{j < r} j!$. The factor $(-1)^{kr}$ in (64.5) cancels its sign. Division by $\prod_{j < d} j!$ proves the formula. The confluent Vandermonde identity itself follows by letting k distinct spectral parameters tend to $+1$ and r tend to -1 in the ordinary exponential Wronskian. $\square$

65 Proof by centered exponential alternants

65.1 The centered Vandermonde identity

Let

$$\alpha_j = j - \frac{d-1}{2}, \quad 0 \leq j < d. \quad (65.1)$$

Then $N + j + \frac{1}{2} = \nu + \alpha_j$. This equality explains the effective size in (64.2); it is not an empirically fitted shift.

Lemma 65.1 (Centered exponential Vandermonde). *For all $\nu_1, \dots, \nu_d \in \mathbb{C}$,*

$$\det[e^{\alpha_j \nu_i}]_{i,j} = \prod_{i < j} 2 \sinh\left(\frac{\nu_j - \nu_i}{2}\right). \quad (65.2)$$

Proof. Factor $e^{-(d-1)\nu_i/2}$ from each row and use the ordinary Vandermonde in e^{ν_i} . In each factor $e^{\nu_j} - e^{\nu_i} = e^{(\nu_i + \nu_j)/2} 2 \sinh((\nu_j - \nu_i)/2)$, the exponential prefactors multiply to the inverse of the row factors. $\square$

Put $\text{sinhc } z = \sinh z / z$, with $\text{sinhc } 0 = 1$. Define

$$R_k(h; \mathbf{u}, w) = [t^k] \det \left[t e^{u_i + \alpha_j(w + hu_i)} - e^{-u_i - \alpha_j(w + hu_i)} \right]_{i,j}, \quad (65.3)$$

$$\begin{aligned} L(h; \mathbf{u}, w) &= \prod_i 2 \sinh\left(\frac{w + hu_i}{2}\right) \\ &\times \prod_{i < j} \left[2 \sinh\left(w + \frac{h(u_i + u_j)}{2}\right) \text{sinhc}\left(\frac{h(u_j - u_i)}{2}\right) \right]. \end{aligned} \quad (65.4)$$

Here $[t^k]$ denotes the coefficient of t^k in a polynomial of degree at most d . The promised exact analytic amplitude is

$$F_k(h; \mathbf{u}, w) = \frac{R_k(h; \mathbf{u}, w)}{h^{ek} \Delta(\mathbf{u}) L(h; \mathbf{u}, w)}, \quad (65.5)$$

with removable singularities filled in as proved next.

65.2 Divisibility in the inverse size and in the coordinates

Lemma 65.2. R_k is divisible by $h^{e_k} \Delta(\mathbf{u})$ as a holomorphic function of $(h, \mathbf{u}, w)$. The quotient is entire in $h, \mathbf{u}, w$.

Proof. Expand the determinant by choosing the positive branch in a set A of k rows and the negative branch in its complement B . Apart from the row factors $e^{\sum_A u - \sum_B u}$ and a sign, the determinant is (65.2) with

$$v_i = \begin{cases} w + hu_i, & i \in A, \\ -w - hu_i, & i \in B. \end{cases}$$

Every pair within A or within B contributes a factor divisible by h . There are exactly e_k such pairs. Thus every summand, and hence R_k , is divisible by h^{e_k} in the ring of entire functions.

After dividing by this common power, the sum is alternating in all u_i , because the original determinant has that property. It vanishes on every hyperplane $u_i = u_j$. Holomorphic division by a linear coordinate difference is elementary: in coordinates $a = u_j - u_i$, a holomorphic function vanishing at $a = 0$ is a times a holomorphic function, as follows from its Taylor series. Divide successively by all the differences in Δ . Each subsequent vanishing continues to hold away from the already divided hyperplanes and therefore on their intersections by continuity. This proves simultaneous divisibility, including at $h = 0$ and arbitrary multiple collisions. $\square$

At $h = 0$ the denominator that remains is

$$L(0; \mathbf{u}, w) = (2 \sinh(w/2))^d (2 \sinh w)^D \neq 0. \quad (65.6)$$

Therefore (65.5) is holomorphic near $h = 0$ locally uniformly in all coordinates. The location of this neighborhood can depend on a compact coordinate set and on the distance of w from the excluded spectral points.

65.3 Recovery of the determinant and the leading term

Factor the row denominators in (63.4) from the Christoffel matrix and use $\nu h = 1$. A row-sign choice with k positive branches has the factor $e^{(2k-d)\nu w}$ and the remaining coefficient is exactly R_k . Also,

$$c_j(h) - c_i(h) = 2 \sinh \left(w + \frac{h(u_i + u_j)}{2} \right) h(u_j - u_i) \operatorname{sinhc} \left(\frac{h(u_j - u_i)}{2} \right). \quad (65.7)$$

Consequently the row denominators times $\Delta(\mathbf{c}(h))$ equal $h^D \Delta(\mathbf{u}) L(h; \mathbf{u}, w)$. Since $D - e_k = kr$, substitution in (63.7) gives (64.7) exactly.

For distinct coordinates the centered identity also gives the useful formula

$$\begin{aligned} F_k(h; \mathbf{u}, w) &= (-1)^{r(r+1)/2} \sum_{|A|=k} \frac{e^{\sum_A u - \sum_B u}}{\prod_i 2 \sinh \left(\frac{w + hu_i}{2} \right)} \\ &\quad \times \frac{1}{\prod_{\{i,j\} \subset A \text{ or } B} 2 \sinh \left(w + \frac{h(u_i + u_j)}{2} \right)} \\ &\quad \times \frac{1}{\prod_{a \in A, b \in B} (u_a - u_b) \operatorname{sinhc} \left(\frac{h(u_a - u_b)}{2} \right)}. \end{aligned} \quad (65.8)$$

Here $B = [d] \setminus A$, and unordered within-group pairs occur once. The sign includes the r negative row choices and the $\binom{r}{2}$ reversed within-negative pairs; conversion of cross factors to the convention in (64.4) gives the displayed common sign. At $h = 0$, (65.8) becomes $B_{d,k}(w)S_{d,k}(\mathbf{u})$. The identity extends across collisions by Lemma 65.2. Taylor's theorem for the analytic amplitudes now proves convergence, derivative-uniformity, and the remainder order asserted in Theorem 64.1.

66 The first correction and all-order construction

Set $U = \sum_{i=1}^d u_i$ and let

$$\mathcal{E} = \sum_{i=1}^d u_i \partial_{u_i}. \quad (66.1)$$

Theorem 66.1 (First-correction operator). *The first Taylor coefficient in (64.8) is*

$$f_{k,1}(\mathbf{u}, w) = B_{d,k}(w) \left[- \left(\frac{1}{2} \coth \frac{w}{2} + \frac{d-2}{4} \coth w \right) U S_{d,k}(\mathbf{u}) - \frac{2k-d}{4} \coth w (\mathcal{E} + k(d-k)) S_{d,k}(\mathbf{u}) \right]. \quad (66.2)$$

In particular, for $d = 2m$, $k = m$, and $U = 0$, the first coefficient of the central amplitude vanishes identically.

Proof. Work first with distinct coordinates. The factors $\sinh c$ in (65.8) are even in h and have no linear contribution. The logarithmic derivative of a summand's other reciprocal factors at zero is

$$-\frac{1}{2} \coth(w/2)U - \frac{1}{2} \coth w \sum_{\{i,j\} \subset A \text{ or } B} (u_i + u_j).$$

Write $V_A = \sum_A u - \sum_B u$. A direct count gives

$$\sum_{\{i,j\} \subset A \text{ or } B} (u_i + u_j) = \frac{d-2}{2}U + \frac{2k-d}{2}V_A. \quad (66.3)$$

The cross denominator in (64.4) is homogeneous of degree kr . Hence, applying $\mathcal{E}$ to each summand,

$$\sum_{|A|=k} \frac{V_A e^{V_A}}{\prod_{a \in A, b \in B} (u_a - u_b)} = (\mathcal{E} + kr) S_{d,k}(\mathbf{u}). \quad (66.4)$$

Substituting (66.3) and (66.4) proves (66.2) on the distinct-coordinate set. Both sides are entire in $\mathbf{u}$, so the formula holds everywhere. $\square$

66.1 An exact finite coefficient algorithm

Formula (65.5) gives every coefficient without estimating a large $N \times N$ determinant. For fixed d, k, J , expand each entry in (65.3) through order $e_k + J$ in h , extract $[t^k]$, divide by $h^{e_k} \Delta(\mathbf{u})$, and multiply by the reciprocal Taylor series of L through order J . All operations are finite. At repeated coordinates use normalized row jets before taking the determinant. The divisibility lemma guarantees that the apparent negative powers disappear.

Equivalently, away from coordinate collisions one may expand the finite subset formula (65.8); analyticity then uniquely specifies the result at collisions. This is often a convenient symbolic route for low orders, but the divided-difference or confluent-determinant route is preferable numerically when coordinates are close. The theorem is an all-orders construction, not a claim that the supplied verification program is an optimized arbitrary-order symbolic engine.

Remark 66.2 (A caution about evenness). For endpoint clusters, a suitable normalization may have an exactly even inverse-size expansion (Part IV, Theorem 47.1). There is no analogous blanket statement here. Even the central sector has a linear term when $U \neq 0$, and when $U = 0$ the neighboring sectors usually contribute at order ν^{-1} . Centering removes an artificial shift but does not remove actual phase-dependent corrections.

67 An explicit collision-uniform remainder

The compact-uniform statement in Theorem 64.1 is already enough for many applications. The following deliberately conservative bound makes the analytic radius and Taylor remainder explicit.

Fix $R \geq 0$ and a nonendpoint w , and set

$$\epsilon = \min \left\{ \frac{1}{4}, |\sinh(w/2)|e^{-(|\Re w|+1)/2}, \frac{1}{2} |\sinh w|e^{-(|\Re w|+1)} \right\}, \quad \rho = \frac{\epsilon}{R+1}. \quad (67.1)$$

For $e = e_k$, define

$$M_k = \frac{\rho^{-e} d^{d/2} \left[2(R+1) \exp \left(R+1 + \frac{d-1}{2} (|\Re w| + \epsilon) \right) \right]^d}{|\sinh(w/2)|^d (|\sinh w|/2)^D}. \quad (67.2)$$

The constants are intended as rigorous bounds, not as sharp estimates.

Theorem 67.1 (Explicit Cauchy remainder). For $|u_i| \leq R$, $|h| < \rho$, and $J \geq 0$,

$$\left| F_k(h; \mathbf{u}, w) - \sum_{j=0}^J f_{k,j}(\mathbf{u}, w) h^j \right| \leq M_k \frac{(|h|/\rho)^{J+1}}{1 - |h|/\rho}. \quad (67.3)$$

The estimate includes all coordinate collisions. Multiplication by the sector weights in (64.7) and summation yields an explicit absolute error for $H_N(q_N)$ whenever $\nu^{-1} < \rho$.

Proof. For $|h| \leq \rho$ and $|u_i| \leq R$, the elementary bound $|\cosh z| \leq e^{|\Re z|}$ and integration along a line segment show

$$\left| 2 \sinh \frac{w + hu_i}{2} \right| \geq |\sinh(w/2)|, \quad \left| 2 \sinh \left(w + \frac{h(u_i + u_j)}{2} \right) \right| \geq |\sinh w|.$$

Indeed, the change in the first expression from $2 \sinh(w/2)$ is at most $\epsilon e^{(|\Re w|+1)/2}$, and the change in the second from $2 \sinh w$ is at most $2\epsilon e^{|\Re w|+1}$. The definition of ϵ gives the inequalities. Also $|z| \leq 1/4$ implies $|\sinh z - 1| < 1/2$ by its power series. Therefore

$$|L(h; \mathbf{u}, w)| \geq |\sinh(w/2)|^d (|\sinh w|/2)^D. \quad (67.4)$$

It remains to bound R_k/Δ without dividing by small coordinate differences. For each column in (65.3), replace the evaluation rows successively by Newton divided differences at u_1 , then u_1, u_2 , and so on. The resulting determinant is the original determinant divided by $\Delta(\mathbf{u})$. A divided difference of an entire function g has the contour representation

$$g[u_1, \dots, u_a] = \frac{1}{2\pi i} \int_{|z|=R+1} \frac{g(z)}{\prod_{i=1}^a (z - u_i)} dz.$$

Every denominator on this contour has modulus at least 1. For $|t| = 1$, $|h| = \rho$, and $|z| = R+1$, each original entry of (65.3) has modulus at most

$$2 \exp \left(R+1 + \frac{d-1}{2} (|\Re w| + \epsilon) \right).$$

The contour bound multiplies this by at most $R + 1$. Hadamard's inequality therefore bounds the divided-difference determinant by the numerator in (67.2) without its factor ρ^{-e} . Cauchy's coefficient formula in t gives the same bound for its k th coefficient. Division by h^e on $|h| = \rho$, followed by (67.4), proves $|F_k| \leq M_k$ on that circle. The removable singularities were already established in Lemma 65.2. Cauchy's coefficient bound $|f_{k,j}| \leq M_k \rho^{-j}$ and summation of a geometric series prove (67.3). $\square$

For coordinate derivatives, apply the same theorem on a slightly larger polydisk and then Cauchy's derivative formula on the smaller one. This provides the derivative-uniform estimates used below, including near zeros. The constants necessarily deteriorate as w approaches an endpoint spectral value, and no endpoint-uniform conclusion is hidden in this theorem.

68 Several separated nonendpoint clusters

Let there be s clusters with fixed sizes $d_a \geq 1$, total size $d = \sum_a d_a$, and base spectral values $w_a \notin \pi i \mathbb{Z}$. Assume

$$c_a = 2 + 2 \cosh w_a \quad \text{are pairwise distinct.} \quad (68.1)$$

With the *global* effective size $\nu = N + d/2$, define

$$c_{a,i,N} = 2 + 2 \cosh \left(w_a + \frac{u_{a,i}}{\nu} \right), \quad 1 \leq i \leq d_a, \quad q_N(x) = \prod_{a,i} (x - c_{a,i,N}). \quad (68.2)$$

For a sector vector $\mathbf{k} = (k_1, \dots, k_s)$, put $r_a = d_a - k_a$, $n_{a,+} = k_a$, and $n_{a,-} = r_a$. Define the nonzero cross-cluster factor

$$T_{ab}(k_a, k_b) = \frac{\prod_{\sigma, \tau \in \{+1, -1\}} \left[2 \sinh \frac{\tau w_b - \sigma w_a}{2} \right]^{n_{a,\sigma} n_{b,\tau}}}{(c_b - c_a)^{d_a d_b}}. \quad (68.3)$$

There is no ambiguity from an ordering of roots inside a cluster; the cluster order in the numerator and denominator is the same.

Theorem 68.1 (Multiple-cluster confluence). *There are analytic amplitudes $F_{\mathbf{k}}(h; \mathbf{u}, \mathbf{w})$ such that, exactly for all sufficiently large N ,*

$$H_N(q_N) = (-1)^{Nd} \sum_{0 \leq k_a \leq d_a} e^{\nu \sum_a (2k_a - d_a) w_a} \nu^{\sum_a k_a r_a} F_{\mathbf{k}}(\nu^{-1}; \mathbf{u}, \mathbf{w}). \quad (68.4)$$

They are symmetric in each cluster's coordinates, have convergent Taylor series locally uniformly including every internal collision, and satisfy

$$F_{\mathbf{k}}(0; \mathbf{u}, \mathbf{w}) = \prod_a [B_{d_a, k_a}(w_a) S_{d_a, k_a}(\mathbf{u}_a)] \prod_{a < b} T_{ab}(k_a, k_b). \quad (68.5)$$

The same uniformity holds after any fixed number of coordinate derivatives. On compact sets of pairwise separated nonendpoint bases, Taylor truncation gives the analogue of (64.9), with $k(d - k)$ replaced by $\sum_a k_a r_a$ and the corresponding exponential weight in (68.4).

Proof. Use the same centered columns $\alpha_j = j - (d - 1)/2$ in the full $d \times d$ Christoffel determinant. Introduce one branch-counting variable t_a for each cluster and extract $\prod_a t_a^{k_a}$. For a row-sign choice, apply Lemma 65.1 to all d signed spectral coordinates at once.

Within cluster a , equal-sign pairs force $h^{\binom{k_a}{2} + \binom{r_a}{2}}$. The root Vandermonde within that cluster has $h^{\binom{d_a}{2}}$, leaving $h^{-k_a r_a}$. The coefficient is alternating in each cluster's coordinates and hence divisible by

$\prod_a \Delta(\mathbf{u}_a)$ even when the coordinates collide. Every cross-cluster factor is nonzero at $h = 0$, by (68.1): a zero of $\sinh((\tau w_b - \sigma w_a)/2)$ would imply $\cosh w_a = \cosh w_b$. The same is true of $c_b - c_a$.

At $h = 0$, the cross factors depend only on the numbers of positive and negative rows in each cluster, not on which coordinate carries which sign. The sum over row-sign choices therefore factors into one S_{d_a, k_a} for each cluster. The remaining within-cluster constants are exactly B_{d_a, k_a} , and the cross constants are (68.3). This proves (68.5). Analyticity of the quotients and nonvanishing of their residual denominators prove the convergent expansion and the uniformity assertions as in the one-cluster case. $\square$

Remark 68.2 (Separation is a genuine hypothesis). The theorem permits arbitrary collisions *inside* each prescribed cluster. It does not assert uniformity while two different base values c_a, c_b approach one another. Such a regime changes the cluster structure and must be recentered as a larger cluster or analyzed with an additional scaling parameter. Nor may one replace the global $N + d/2$ by a different local size for each cluster without correspondingly changing the amplitudes.

69 Inside the moment interval: dominant sectors and parity

Take $w = i\theta$ with $0 < \theta < \pi$. The base root is in $(0, 4)$, and all the factors $e^{(2k-d)\nu w}$ have modulus 1. The dominant powers are determined by the elementary concave quadratic $k(d - k)$.

Corollary 69.1 (Even and odd total multiplicity). *For $d = 2m$,*

$$\frac{H_N(q_N)}{\nu^{m^2}} = B_{2m, m} S_{2m, m} + \frac{1}{\nu} [f_{m, 1} + e^{2i\nu\theta} B_{2m, m+1} S_{2m, m+1} + e^{-2i\nu\theta} B_{2m, m-1} S_{2m, m-1}] + O(\nu^{-2}). \quad (69.1)$$

All B and f are evaluated at $i\theta$, and all profiles at $\mathbf{u}$. For $d = 2m + 1$, the two leading sectors give

$$\frac{(-1)^N H_N(q_N)}{\nu^{m(m+1)}} = e^{-i\nu\theta} B_{d, m} S_{d, m} + e^{i\nu\theta} B_{d, m+1} S_{d, m+1} + O(\nu^{-1}). \quad (69.2)$$

The errors are compact-uniform with coordinate derivatives. In (69.2) the first corrections come only from these two leading amplitudes; the next pair of sectors starts two powers lower.

Proof. For $d = 2m$, put $k = m + j$; then $k(d - k) = m^2 - j^2$. For $d = 2m + 1$, the maximum $m(m + 1)$ occurs exactly at $k = m, m + 1$, and the next value is smaller by 2. Substitute these identities in Theorem 64.1. $\square$

The odd case need not have an N -independent limit even at leading order. The even case usually does have one, but it need not have an N -independent first correction. This distinction is essential in using bulk limits to predict finite determinant zeros.

70 Sine-kernel duality and a positive Gram representation

This section identifies classical structure inside the universal profile. The proofs also provide an independent route to its positivity properties and an independent numerical check of its normalization.

70.1 A bilinear Christoffel identity

Define

$$K_M(x, y) = \sum_{\ell=0}^M p_\ell(x) p_\ell(y). \quad (70.1)$$

Lemma 70.1 (Bilinear kernel formula). *For two tuples $\mathbf{a}, \mathbf{b}$ of length t ,*

$$H_N \left(\prod_{i=1}^t (x - a_i)(x - b_i) \right) = \frac{\det[K_{N+t-1}(a_i, b_j)]_{i,j=1}^t}{\Delta(\mathbf{a})\Delta(\mathbf{b})}. \quad (70.2)$$

The quotient extends to repeated nodes by normalized derivatives.

Proof. Multiply the two augmented Vandermondes

$$\Delta(x_1, \dots, x_N, a_1, \dots, a_t) \quad \text{and} \quad \Delta(x_1, \dots, x_N, b_1, \dots, b_t),$$

divide by $\Delta(\mathbf{a})\Delta(\mathbf{b})$, and integrate against $\prod_{a=1}^N d\mu(x_a)/N!$. The product of the two sign factors is 1, so this is the left side of (70.2). Rewrite both determinants in the orthonormal basis $p_0, \dots, p_{N+t-1}$. Expanding the last t rows and integrating the first N rows forces identical complementary column sets, with unit norm factors. The sum over the remaining t -element column sets is, by Cauchy–Binet, $\det K_{N+t-1}(a_i, b_j)$. The identical expansion signs occur twice and cancel. Continuity proves confluence. $\square$

The finite kernel has a particularly transparent bulk expression. With $\nu = N + t$, summing the products of sines in (63.3) gives

$$\begin{aligned} & K_{\nu-1}(2 + 2 \cos \theta_a, 2 + 2 \cos \theta_b) \\ &= \frac{1}{4 \sin(\theta_a/2) \sin(\theta_b/2)} \left[\frac{\sin(\nu(\theta_a - \theta_b))}{\sin((\theta_a - \theta_b)/2)} - \frac{\sin(\nu(\theta_a + \theta_b))}{\sin((\theta_a + \theta_b)/2)} \right]. \end{aligned} \quad (70.3)$$

Every coincident argument is interpreted by continuity. In particular, for $\theta_a = \theta + x/\nu$ and $\theta_b = \theta + y/\nu$,

$$\nu^{-1} K_{\nu-1} \rightarrow \frac{\text{sinc}(x - y)}{2 \sin^2(\theta/2)}, \quad \text{sinc } z = \frac{\sin z}{z}, \quad \text{sinc } 0 = 1, \quad (70.4)$$

locally uniformly in complex x, y , with all fixed derivatives. The second term of (70.3) is bounded on compact coordinate sets and is precisely one source of the oscillatory finite-size correction.

Theorem 70.2 (Bilinear universal profile). *For two complex tuples $\mathbf{x}, \mathbf{y}$ of length t ,*

$$S_{2t,t}(ix_1, \dots, ix_t, iy_1, \dots, iy_t) = 2^t \frac{\det[\text{sinc}(x_i - y_j)]_{i,j=1}^t}{\Delta(\mathbf{x})\Delta(\mathbf{y})}. \quad (70.5)$$

Both sides are entire with their confluent interpretations. If $\mathbf{x} = \mathbf{y}$ is real, the quotient on the right is strictly positive, including after any collisions within the tuple.

Proof. Apply Lemma 70.1 with $a_i = 2 + 2 \cos(\theta + x_i/\nu)$ and $b_j = 2 + 2 \cos(\theta + y_j/\nu)$. Use (70.4) and

$$\Delta(\mathbf{a})\Delta(\mathbf{b}) = \left(\frac{-2 \sin \theta}{\nu} \right)^{t(t-1)} \Delta(\mathbf{x})\Delta(\mathbf{y})(1 + O(\nu^{-1})).$$

Compare the resulting leading term with the unique central sector in Theorem 64.1, noting that

$$B_{2t,t}(i\theta) = \frac{1}{(2 \sin(\theta/2))^{2t} (2 \sin \theta)^{t(t-1)}}. \quad (70.6)$$

The constants differ by the factor 2^t in (70.5). This proves the formula first for distinct coordinates; derivative-uniformity or analytic continuation proves it at collisions.

For real distinct x_i , use

$$\text{sinc}(x_i - x_j) = \frac{1}{2} \int_{-1}^1 e^{ix_i s} \overline{e^{ix_j s}} ds.$$

The matrix is the Gram matrix of linearly independent exponentials in $L^2([-1, 1], ds/2)$ and is positive definite. Division by $\Delta(x)^2 > 0$ preserves positivity. At repeated nodes, its normalized confluent matrix is the Gram matrix of functions $(is)^a e^{ixs}/a!$ attached to the distinct node blocks. These exponential-polynomials are still linearly independent: an identity on an interval extends analytically, and repeated application of the operators $\partial_s - ix$ separates the distinct frequencies and their polynomial degrees. The confluent Gram determinant is therefore strictly positive. The remaining cross-block Vandermonde factors are squares of nonzero real numbers. This proves strict positivity at every collision pattern. $\square$

71 Two equal-multiplicity roots: the complete first-order law

Fix an integer $t \geq 1$ and $0 < \theta < \pi$. For $\nu = N + t$ put

$$c_{\pm, N} = 2 + 2 \cos \left(\theta \pm \frac{\tau}{\nu} \right), \quad q_{N, t, \tau}(x) = (x - c_{+, N})^t (x - c_{-, N})^t. \quad (71.1)$$

Let

$$s_t = 2^{t^2} \frac{\left(\prod_{j=0}^{t-1} j! \right)^2}{\prod_{j=0}^{2t-1} j!}, \quad (71.2)$$

$$A_t(\theta) = \frac{s_t}{(2 \sin(\theta/2))^{2t} (2 \sin \theta)^{t(t-1)}} > 0, \quad (71.3)$$

$$\Phi_t(\tau) = \frac{S_{2t, t}((i\tau)^t, (-i\tau)^t)}{s_t}, \quad (71.4)$$

$$\Psi_t(\tau) = \frac{S_{2t, t+1}((i\tau)^t, (-i\tau)^t)}{s_t}. \quad (71.5)$$

The notation $(i\tau)^t$ in a tuple means t repeated coordinates, not a power. Both profiles are entire even functions of τ , real-valued on the real axis. Their values at zero are

$$\Phi_t(0) = 1, \quad \Psi_t(0) = \frac{t}{2}. \quad (71.6)$$

These follow from Proposition 64.2. Reality follows by complex conjugation and permutation of the two repeated blocks.

Theorem 71.1 (Symmetric collision and its phase correction). *Uniformly on compact subsets of complex τ , with any fixed number of derivatives,*

$$\frac{H_N(q_{N, t, \tau})}{A_t(\theta)\nu^{t^2}} = \Phi_t(\tau) + \frac{(-1)^t \sin(2\nu\theta)}{\nu \sin \theta} \Psi_t(\tau) + O(\nu^{-2}). \quad (71.7)$$

In particular the leading limit exists without selecting a phase subsequence, although its first correction generally oscillates with N .

Proof. Use the one-cluster theorem with $d = 2t$, $w = i\theta$, and coordinate multiset $\mathbf{u} = ((i\tau)^t, (-i\tau)^t)$. Its sum U is zero, so Theorem 66.1 makes the first central amplitude correction vanish. The adjacent sectors $k = t \pm 1$ are exactly one power below the central sector. By reflection and symmetry of the coordinate multiset,

$$S_{2t, t-1}(\mathbf{u}) = S_{2t, t+1}(\mathbf{u}).$$

A direct calculation from (64.6) gives

$$\frac{B_{2t, t+1}(w)}{B_{2t, t}(w)} = \frac{(-1)^t}{2 \sinh w}, \quad \frac{B_{2t, t-1}(w)}{B_{2t, t}(w)} = \frac{(-1)^{t+1}}{2 \sinh w}.$$

Their phase-weighted sum is $(-1)^t \sin(2\nu\theta)/\sin \theta$. Division by $A_t = B_{2t, t}(i\theta)s_t$ proves (71.7). The residual central Taylor series begins at order ν^{-2} , the adjacent amplitude corrections also start at that order after their sector weight, and all other sectors are at least four powers below the center. The uniform remainder follows from Theorem 64.1. $\square$

71.1 Identification with the classical oscillatory Jacobi determinant

For $j \geq 0$ define

$$m_j(\tau) = \frac{1}{2} \int_{-1}^1 y^j e^{2i\tau y} dy. \quad (71.8)$$

At $\tau = 0$, $m_j(0) = 0$ for odd j and $1/(j+1)$ for even j .

Proposition 71.2 (Jacobi Hankel duality). *The collision profile has the identities*

$$\Phi_t(\tau) = \frac{\det[m_{i+j}(\tau)]_{0 \leq i, j < t}}{\det[m_{i+j}(0)]_{0 \leq i, j < t}}, \quad (71.9)$$

$$= \frac{1}{Z_t} \int_{[-1, 1]^t} e^{2i\tau \sum_j y_j} \Delta(\mathbf{y})^2 d\mathbf{y}, \quad (71.10)$$

$$Z_t = 2^{t^2} \prod_{j=0}^{t-1} \frac{(j!)^2 (j+1)!}{(t+j)!}. \quad (71.11)$$

Thus Φ_t is a normalized version of the Hankel determinant studied in the kissing-polynomial literature, with frequency $\omega = 2\tau$.

Proof. In Theorem 70.2, let all x_i tend to τ and all y_j to $-\tau$. A confluent entry is

$$\frac{1}{a!b!} \partial_x^a \partial_y^b \operatorname{sinc}(x-y) \Big|_{x=\tau, y=-\tau} = \frac{i^a (-i)^b}{a!b!} m_{a+b}(\tau).$$

The row and column phase factors multiply to 1, and the factorial factors cancel upon division by the same expression at zero. This proves (71.9). Andréief gives (71.10) and $Z_t = 2^t t! \det[m_{i+j}(0)]$.

For the product evaluation, the monic Legendre polynomial of degree j has squared norm

$$\frac{2}{2j+1} \frac{2^{2j}}{\binom{2j}{j}^2}$$

with respect to dy on $[-1, 1]$. This follows from the Rodrigues formula by j integrations by parts and the leading coefficient $2^{-j} \binom{2j}{j}$ of the usual Legendre polynomial. Hence $Z_t = t!$ times the product of these monic squared norms. Simplifying the factorials gives (71.11). $\square$

Corollary 71.3 (Even multiplicities have no real profile zeros). *For every $m \geq 1$ and every real τ ,*

$$\Phi_{2m}(\tau) > 0. \quad (71.12)$$

For real roots in (71.1), the corresponding finite $H_N(q_{N,2m,\tau})$ is also strictly positive for every N .

Proof. Take $t = 2m$ and let the real tuple $\mathbf{x}$ of length t consist of m copies of τ and m copies of $-\tau$. The coordinate multiset $(i\mathbf{x}, i\mathbf{x})$ is the one defining Φ_t . Strict positivity follows from the confluent Gram statement in Theorem 70.2 and $s_t > 0$. For the finite determinant, $q_{N,2m,\tau}$ is the square of a nonzero real polynomial. The modified measure $q d\mu$ is positive and has support containing an interval apart from finitely many points. Its moment Gram matrix is positive definite. $\square$

This elementary proof does not make the positivity statement new. The same nondegeneracy phenomenon is part of the established analysis of oscillatory orthogonal polynomials [13]. Its role here is to separate genuine moving zeros from a spurious prediction obtained by truncating a finite-size expansion outside its uniform regime.

72 The classical profile at large real frequency

We give a self-contained asymptotic argument for the odd profile zeros needed in the next section. The subject and its parity phenomenon are classical [13]; the finite- N transport theorem, rather than a new claim about the existence of kissing points, is the application. The proof is elementary here because the integrand in (71.10), apart from its exponential, is a polynomial.

Put

$$L_j = \prod_{\ell=1}^j \ell! (\ell-1)!, \quad L_0 = 1. \quad (72.1)$$

Lemma 72.1 (Vertex contribution). *For $\tau \neq 0$, the integral in (71.10) is an exact finite sum of exponentials in τ with Laurent-polynomial coefficients. For a class of vertices with j coordinates at $+1$ and $r = t - j$ at -1 , its leading contribution to the normalized integral is*

$$\binom{t}{j} \frac{2^{2jr} L_j L_r}{Z_t} e^{2i\tau(j-r)} (2i\tau)^{-j^2} (-2i\tau)^{-r^2} \left[1 + \frac{jr(r-j)}{2i\tau} + O(\tau^{-2}) \right]. \quad (72.2)$$

The remainder has the corresponding differentiated asymptotics along the positive real axis.

Proof. Repeated integration by parts in each variable of a polynomial times $e^{2i\tau \Sigma y}$ terminates. The answer is a finite sum of boundary jets at the vertices of the cube, with powers of $(2i\tau)^{-1}$. Thus no unproved steepest-descent interchange is required.

To compute the first nonzero jets at one vertex, write the j upper coordinates as $1 - a_i$ and the r lower coordinates as $-1 + b_i$. The within-group Vandermonde squares are $\Delta(\mathbf{a})^2$ and $\Delta(\mathbf{b})^2$, of degrees $j(j-1)$ and $r(r-1)$. The cross part is

$$\prod_{i=1}^j \prod_{\ell=1}^r (2 - a_i - b_\ell)^2 = 2^{2jr} \left[1 - r \sum_i a_i - j \sum_\ell b_\ell + \text{terms of degree at least 2} \right]. \quad (72.3)$$

The boundary-jet functional on a monomial of degree a at an upper endpoint is $a!/(2i\tau)^{a+1}$, and at a lower endpoint is $a!/(-2i\tau)^{a+1}$. Their sums can be computed using the convergent Laplace integral

$$\int_{[0,\infty)^j} e^{-\Sigma v_i} \Delta(\mathbf{v})^2 d\mathbf{v} = j! \det[(a+b)!]_{0 \leq a, b < j} = L_j.$$

For the determinant evaluation, factor $a!$ and $b!$ from its rows and columns; the remaining Pascal-type determinant $\det[\binom{a+b}{a}]$ is 1, by the factorization into a lower Pascal matrix and its transpose. Scaling the Laplace parameter shows that the same integral with an extra Σv_i is $j^2 L_j$.

These identities give the leading powers j^2 and r^2 in (72.2). The linear part of (72.3) contributes

$$-\frac{rj^2}{2i\tau} - \frac{jr^2}{-2i\tau} = \frac{jr(r-j)}{2i\tau}.$$

There are $\binom{t}{j}$ vertices of this type. All remaining terms are at least two powers smaller within that vertex contribution. Because the integration-by-parts expansion is finite, its differentiated remainder bounds follow directly. $\square$

Proposition 72.2 (Parity asymptotics and large odd zeros). *As $\tau \rightarrow +\infty$,*

$$\Phi_{2m}(\tau) = \frac{\binom{2m}{m} L_m^2}{Z_{2m}} \tau^{-2m^2} [1 + O(\tau^{-2})]. \quad (72.4)$$

For $t = 2m + 1$, set $E = (t^2 + 1)/2$ and $Q_t = \binom{t}{m} L_m L_{m+1} / Z_t$. Then

$$\Phi_t(\tau) = (-1)^m Q_t \tau^{-E} \left[\sin(2\tau) + \frac{m(m+1)}{2\tau} \cos(2\tau) + O(\tau^{-2}) \right]. \quad (72.5)$$

The normalized bracket has the corresponding derivative remainder. In particular, for each sufficiently large integer j there is exactly one simple positive real zero near $j\pi/2$, and

$$\tau_j = \frac{j\pi}{2} - \frac{m(m+1)}{2\pi j} + O(j^{-2}). \quad (72.6)$$

These account for all sufficiently large positive real zeros.

Proof. The smallest exponent $j^2 + (t-j)^2$ in Lemma 72.1 occurs at $j = m$ for $t = 2m$, and at $j = m, m+1$ for $t = 2m+1$. In the even case, the linear correction vanishes, the next vertex class is two powers smaller, and the powers of 2 and i in the central contribution cancel. This proves (72.4).

In the odd case the two leading vertices are complex conjugates. Each has the factor $Q_t/2$, and their phases combine into $(-1)^m \sin(2\tau)$. Their opposite linear corrections combine into $(-1)^m m(m+1) \cos(2\tau)/(2\tau)$. The next vertex class is four powers smaller, proving (72.5) and its derivative form.

Let the normalized bracket be $g(\tau)$. Its estimate implies that any large zero lies within $O(1/\tau)$ of an integer multiple of $\pi/2$. On each such neighborhood, $g'(\tau) = 2 \cos(2\tau) + O(1/\tau)$ is bounded away from zero for sufficiently large τ . The values on the two sides of a sufficiently large fixed multiple of $1/\tau$ have opposite signs. Hence there is precisely one real zero in the neighborhood, and it is simple. Writing $\tau = j\pi/2 + \delta$ in (72.5) gives $2\delta + m(m+1)/(2(j\pi/2)) = O(j^{-2})$, which is (72.6). The localization argument rules out other large real zeros. $\square$

Remark 72.3 (Order of limits). The assertion above has fixed t and $\tau \rightarrow \infty$. The collision expansion has fixed t and bounded τ as $N \rightarrow \infty$. They may be used successively to choose any particular sufficiently large simple profile zero and then transport that fixed zero as N grows. They do not, without further estimates, give a theorem in which the zero index j grows with N .

73 Finite-size zeros and convergent phase-dependent transport

Write

$$Z_N(\tau) = \frac{H_N(q_{N,t,\tau})}{A_t(\theta)v^{t^2}}, \quad v = N + t. \quad (73.1)$$

The preceding sections provide simple profile zeros for every odd t . The next two results apply at any simple zero, without requiring its large-frequency approximation.

Theorem 73.1 (First displacement of a real zero). *Suppose $\tau_* \in \mathbb{R}$ satisfies $\Phi_t(\tau_*) = 0$ and $\Phi'_t(\tau_*) \neq 0$. For all sufficiently large N , there is exactly one zero τ_N of Z_N in a fixed sufficiently small complex disk about τ_* . That zero is real and simple, and*

$$\tau_N = \tau_* + \frac{(-1)^{t+1}}{v} \frac{\sin(2v\theta)}{\sin \theta} \frac{\Psi_t(\tau_*)}{\Phi'_t(\tau_*)} + O(v^{-2}). \quad (73.2)$$

The error is uniform in the phase $e^{2iv\theta}$ for fixed t, θ, τ_* .

Proof. Choose a small closed disk centered at τ_* containing no other zero of Φ_t . On its boundary $|\Phi_t|$ has a positive minimum. The derivative-uniform expansion in Theorem 71.1 gives $Z_N = \Phi_t + O(v^{-1})$. Rouché's theorem therefore gives one zero in the disk, counted with multiplicity. It is simple. Since Z_N is real on the real axis, nonreal zeros come in conjugate pairs; uniqueness in a conjugation-invariant disk makes this zero real.

The nonvanishing derivative and the $O(v^{-1})$ perturbation first give $\delta_N = \tau_N - \tau_* = O(v^{-1})$. Taylor expansion of (71.7) at τ_* now yields

$$0 = \Phi'_t(\tau_*)\delta_N + \frac{(-1)^t}{v} \frac{\sin(2v\theta)}{\sin \theta} \Psi_t(\tau_*) + O(v^{-2}).$$

Solving this equation proves (73.2). Every bound used was uniform for the phase on the unit circle. $\square$

73.1 A genuinely convergent expansion, not only an asymptotic one

The fast phase can be treated as an independent variable. Write $\mathbf{u}(\tau) = ((i\tau)^t, (-i\tau)^t)$ and define

$$\mathcal{Z}(h, \zeta, \tau) = \frac{1}{A_t(\theta)} \sum_{j=-t}^t h^{j^2} \zeta^j F_{t+j}(h; \mathbf{u}(\tau), i\theta), \quad (73.3)$$

where the amplitudes are those for $d = 2t$. Then

$$\mathcal{Z}_N(\tau) = \mathcal{Z}\left(v^{-1}, e^{2iv\theta}, \tau\right), \quad \mathcal{Z}(0, \zeta, \tau) = \Phi_t(\tau). \quad (73.4)$$

The function is holomorphic for h small, τ in any fixed compact neighborhood, and ζ in an annulus about the unit circle. It is not asserted to extend to $\zeta = 0$.

Theorem 73.2 (Convergent Laurent-phase zero expansion). *Under the hypotheses of Theorem 73.1, there are Laurent polynomials $b_\ell(\zeta)$, $\ell \geq 1$, whose exponents lie between $-\ell$ and ℓ , such that*

$$\tau_N = \tau_* + \sum_{\ell=1}^{\infty} v^{-\ell} b_\ell(e^{2iv\theta}). \quad (73.5)$$

The series is convergent for all sufficiently large N , uniformly in the phase on the unit circle. More generally it is the restriction of a holomorphic root function $\tau(h, \zeta) = \tau_ + \sum_{\ell \geq 1} b_\ell(\zeta) h^\ell$ on a product of a disk in h and an annulus in ζ . Its first coefficient is*

$$b_1(\zeta) = \frac{(-1)^{t+1}}{2i \sin \theta} (\zeta - \zeta^{-1}) \frac{\Psi_t(\tau_*)}{\Phi'_t(\tau_*)}. \quad (73.6)$$

Proof. At every point $(0, \zeta, \tau_*)$, $\partial_\tau \mathcal{Z} = \Phi'_t(\tau_*) \neq 0$. Fix a closed annulus contained in $\mathbb{C} \setminus \{0\}$ with the unit circle in its interior. Uniform Rouché estimates on its compact set of phases, or the holomorphic implicit-function theorem followed by compactness, give a single root in a common small disk about τ_* for all sufficiently small $|h|$. The roots obtained locally agree by uniqueness. They form a jointly holomorphic function $\tau(h, \zeta)$, whose Taylor expansion in h converges uniformly on a smaller product domain.

It remains to prove the finite Laurent support rather than merely holomorphy in ζ . The coefficient of h^a in (73.3) is a Laurent polynomial with $|j| \leq a$, because a sector of frequency j first appears at h^{j^2} and $j^2 \geq |j|$. At order h^ℓ in the identity $\mathcal{Z}(h, \zeta, \tau(h, \zeta)) = 0$, the only term containing the new unknown b_ℓ is $\Phi'_t(\tau_*) b_\ell$. All other terms are finite products of earlier coefficients and derivatives of the coefficient functions of $\mathcal{Z}$. By induction, their Laurent degrees are bounded by the sum of their h -orders, namely ℓ . Division by the nonzero scalar $\Phi'_t(\tau_*)$ proves the claim. The coefficient at order one is exactly (73.6), by Theorem 71.1. $\square$

The recursive proof is also an all-orders zero-coefficient algorithm. It uses differentiation and finite Laurent-polynomial arithmetic after the amplitude coefficients have been computed. The statement is stronger than saying that each phase subsequence has a formal series: it gives one analytic root function valid on a whole annulus of phases. It is also weaker, and more accurate, than asserting a phase-independent power series in $1/N$.

74 Explicit examples and a sharp failure of a phase-free correction

74.1 Two simple roots: an exact formula

Example 74.1. Take $t = 1$ and $v = N + 1$. Lemma 70.1, now a 1×1 identity, and (70.3) give

$$H_N(q_{N,1,\tau}) = \frac{\frac{\sin(2\tau)}{\sin(\tau/v)} - \frac{\sin(2v\theta)}{\sin \theta}}{4 \sin((\theta + \tau/v)/2) \sin((\theta - \tau/v)/2)}. \quad (74.1)$$

At $\tau = 0$ the first quotient is 2ν . On every fixed compact τ -set the denominator is nonzero for all sufficiently large N . The profiles are

$$\Phi_1(\tau) = \frac{\sin(2\tau)}{2\tau}, \quad \Psi_1(\tau) = \frac{1}{2}, \quad A_1(\theta) = \frac{1}{2\sin^2(\theta/2)}. \quad (74.2)$$

For a fixed integer $j \geq 1$, the limiting zero $\tau_* = j\pi/2$ is simple, and

$$\tau_{N,j} = \tau_* + \frac{(-1)^j \tau_*}{2\nu \sin \theta} \sin(2\nu\theta) + O(\nu^{-2}). \quad (74.3)$$

Indeed, away from removable singularities, the exact zero equation is

$$\sin(2\tau) = \frac{\sin(2\nu\theta)}{\sin \theta} \sin(\tau/\nu). \quad (74.4)$$

Expanding it at $j\pi/2$ reproduces (74.3) independently of the general theorem.

Corollary 74.2 (Periodic and quasiperiodic scaled zero motion). *For the fixed zero in Example 74.1, if θ/π is irrational, the set of subsequential limits of $\nu(\tau_{N,j} - \tau_*)$ is the full interval*

$$\left[-\frac{\tau_*}{2\sin \theta}, \frac{\tau_*}{2\sin \theta} \right]. \quad (74.5)$$

If θ/π is rational, there are finitely many phase limits, obtained from its periodic phase sequence. At $\theta = \pi/2$, the fixed zeros $j\pi/2$ are exact finite-size zeros for all sufficiently large N .

Proof. When θ/π is irrational, the sequence $e^{2i(N+1)\theta}$ is dense on the unit circle, so its sine has precisely $[-1, 1]$ as its set of subsequential limits. This standard rotation fact also follows from the pigeonhole approximation argument: integer multiples of an irrational angle have arbitrarily small nonzero steps modulo 2π , whose successive multiples form arbitrarily fine nets. Formula (74.3) then gives (74.5); its scaled error tends to zero. For rational θ/π , the phase is periodic. At $\theta = \pi/2$, $\sin(2\nu\theta) = \sin(\nu\pi) = 0$ since $\nu = N + 1$ is an integer. Equation (74.4) then has the claimed exact roots, with nonzero denominator for large N . $\square$

Thus the first zero displacement need not approach any one constant after multiplication by N . Replacing the phase-dependent correction by an ordinary coefficient would lose an entire interval of possible scaled behavior. This is already visible in degree two, and the general Laurent-phase theorem explains its higher-multiplicity analogue.

74.2 Two roots of multiplicity two

For $t = 2$, the 2×2 determinant in (71.9) simplifies to

$$\Phi_2(\tau) = \frac{3(4\tau^2 - \sin^2(2\tau))}{16\tau^4}, \quad \Phi_2(0) = 1. \quad (74.6)$$

For real nonzero τ , $|\sin(2\tau)| < 2|\tau|$, so the expression is strictly positive, agreeing with Corollary 71.3. Its first finite-size correction is nevertheless oscillatory:

$$Z_N(\tau) = \Phi_2(\tau) + \frac{\sin(2(N+2)\theta)}{(N+2)\sin \theta} \Psi_2(\tau) + O((N+2)^{-2}).$$

Positivity of the leading profile is not a justification for suppressing that correction when accurate values, rather than only existence of a limit, are needed.

74.3 A repeated root at the collision center

At $\tau = 0$, all $2t$ roots coincide, and Theorem 71.1 gives the particularly simple law

$$\frac{H_N((x - (2 + 2 \cos \theta))^2)^{2t}}{A_t(\theta)(N + t)^{t^2}} = 1 + \frac{(-1)^t t}{2(N + t)} \frac{\sin(2(N + t)\theta)}{\sin \theta} + O((N + t)^{-2}). \quad (74.7)$$

This overlaps the fixed-root setting, but here it appears as the regular center of an entire collision family, with a uniform neighborhood and all coordinate derivatives. The limit has not been obtained by substituting coincident roots into separately singular coefficients.

75 Reproducible checks and their evidentiary status

75.1 What the verification program does

The accompanying code/`verify.py` implements the Catalan moment determinant, confluent Christoffel determinants, the universal profile, the exact finite-sector amplitude, the first correction, and the oscillatory Jacobi moment determinant. Its exact tests use Python's `Fraction`; numerical tests use `mpmath` at 100 decimal digits. All 518 assertions passed in the recorded run:

Exact-arithmetic assertions	Count
Moment determinant versus confluent Christoffel, $N = 0, \dots, 6$	63
Centered Vandermonde identities, dimensions $1, \dots, 7$	7
Leading sector coefficients at full confluence, $d \leq 7$	35
First sector coefficients at full confluence, $d \leq 7$	35
Forced lower-order zeros in the confluent sector numerator	252
Exact subtotal	392

The 126 numerical assertions comprise 35 profile values at zero, 25 profile/Jacobi duality checks through $t = 5$, 30 exact finite-sector identity comparisons, 14 first-correction comparisons, four bilinear sine-kernel comparisons, twelve evaluations of the exact two-root formula, and six positive samples of even profiles.

[Merge note: the program is shipped as `code/05-interior-collision-verify.py`; in this report the unprefix code/`verify.py` is Part I's verifier. The recorded run is `data/05-interior-collision-verification.json`, and the four CSV files named below are shipped as `data/05-interior-collision-finite_size_corrections.csv`, `data/05-interior-collision-zero_transport.csv`, `data/05-interior-collision-multiple_clusters.csv` and `data/05-interior-collision-jacobi_parity_audit.csv`.]

The exact coefficient audit uses rational spectral value $z = e^w = 4$ and all-collided coordinates, dimensions $d \leq 7$. The numerical sector audit also includes distinct coordinates, real and complex off-circle bases, and an interior base on the unit circle. None of these finite parameter sets is a substitute for the all-parameter divisibility and analyticity proofs.

The largest normalized discrepancy in the numerical finite-sector identity comparisons was approximately 2.16×10^{-90} ; for the Jacobi duality it was 2.68×10^{-92} . The numerical derivative check of the first coefficient used a centered difference with $h = 10^{-6}$, and its largest normalized discrepancy was 6.62×10^{-13} , consistent with the deliberately finite $O(h^2)$ differentiation error. These are high-precision consistency checks, *not* outward-rounded interval enclosures or proof-assistant certificates.

75.2 First-correction data

Table 5 uses $\theta = 1.1$, $\tau = 0.8$. The leading error is $|Z_N - \Phi_t|$; the corrected error subtracts the explicitly oscillatory ν^{-1} term in (71.7). The CSV files retain more digits and the scaled residuals.

t	N	Leading error	Corrected error	ν^2 corrected error
1	40	1.0518×10^{-2}	2.5363×10^{-4}	0.4263
1	160	2.4831×10^{-3}	1.6630×10^{-5}	0.4311
1	320	1.0651×10^{-3}	4.1914×10^{-6}	0.4319
2	40	1.9496×10^{-2}	1.9676×10^{-4}	0.3471
2	160	5.2145×10^{-3}	1.6170×10^{-5}	0.4244
2	320	2.6644×10^{-3}	4.8569×10^{-6}	0.5036
3	40	1.2476×10^{-2}	2.4082×10^{-3}	4.4527
3	160	3.5713×10^{-3}	1.6426×10^{-4}	4.3643
3	320	2.2341×10^{-3}	4.0235×10^{-5}	4.1977

Table 5: Computed finite-size errors for the symmetric collision law. The bounded scaled residual is consistent with the proved absolute $O(\nu^{-2})$ error; the displayed samples do not prove that order.

75.3 Zero-transport data

For $t = 3$, a numerically located profile zero is

$$\tau_* = 6.201688214318496432280709 \dots$$

The displayed decimal is numerical, not a certified isolating interval. The theorem applies to any simple zero; Proposition 72.2 independently supplies infinitely many such zeros for odd t . Table 6 compares the observed root displacement with the prediction $\delta_N^{(1)}$ in (73.2), again at $\theta = 1.1$.

t	N	Finite zero τ_N	Predicted shift $\delta_N^{(1)}$	$\nu^2 \tau_N - \tau_* - \delta_N^{(1)} $
1	40	1.5540579439	-0.0169195592	0.3046
1	160	1.5668796236	-0.0039265155	0.2543
1	320	1.5691184590	-0.0016796655	0.1852
3	40	6.1748653119	-0.0279145143	2.0184
3	160	6.1923244738	-0.0094465544	2.2003
3	320	6.1956309248	-0.0060828969	2.6716

Table 6: Numerical zero transport. For $t = 1$, $\tau_* = \pi/2$. The nonmonotone scaled error is compatible with higher phase-dependent coefficients; monotonicity is not asserted.

The package additionally records a two-separated-cluster experiment and audit values for the classical large-frequency profile expansion. These experiments are labeled separately from the 518 asserted checks. Their complete parameters are in the code, and their outputs are in `data/multiple_clusters.csv` and `data/jacobi_parity_audit.csv`.

75.4 Reproduction and computational limitations

From the package directory, run

```
python -m pip install -r requirements.txt
python code/verify.py --out data
pdflatex -interaction=nonstopmode -halt-on-error article.tex
pdflatex -interaction=nonstopmode -halt-on-error article.tex
```

[Merge note: these commands assume the delivered package, which is not the layout here, and must *not* be run in this directory. The manuscript `article.tex` and the package `requirements.txt` are not at the report root (the manuscript is printed as Part V, and the dependency file is shipped as `data/05-interior-collision-requirements.txt`); here `code/verify.py` is Part I's verifier, and the shipped Part V program, run from this directory without `--out`, would write `verification.json` into `data/`, overwriting Part I's recorded `data/verification.json`, and would add four unprefixed CSV files. The delivered Makefile, shipped as `code/05-interior-collision-Makefile`, runs `python code/verify.py --out data` (Part I's verifier, here) and `pdflatex` on `article.tex`, and its clean target deletes `article.aux`, `.log`, `.out`, `.toc`, `.fls` and `.fdb_latexmk`; do not run it from this directory either. Restore the delivered layout in a scratch copy instead (a POSIX shell, from the report directory):]

```
W=/path/to/scratch/interior-collision
mkdir -p "$W/code" "$W/recorded"
cp code/05-interior-collision-verify.py "$W/code/verify.py"
for f in data/05-interior-collision-*; do
  cp "$f" "$W/recorded/${f#data/05-interior-collision-}"; done
cd "$W" && uv run --no-project --with mpmath==1.3.0 \
  python code/verify.py --out data # compare $W/data with $W/recorded
```

[Run this way on 29 September 2026 with Python 3.13.5 and `mpmath` 1.3.0, it passed all 518 assertions in a few seconds; the four CSV files were byte-identical to the recorded ones (CRLF, as recorded, because the `csv` module ends rows with CRLF on every system), and `verification.json` equalled the recorded file apart from `elapsed_seconds` and Windows line endings. The checks raise exceptions explicitly, so `-O` does not disable them.] The supplied determinant routines are small reference implementations. A degree- d Christoffel calculation replaces the original $N \times N$ determinant by a $d \times d$ determinant, with polynomial jets generated by recurrence. At close but nonidentical nodes, cancellation can still require extra precision; exact coincidence should use confluent jets rather than tiny numerical separations. The Fourier coefficient extraction used in the numerical amplitude routine is a finite root-of-unity evaluation at high precision, not an interval algorithm. A production implementation should adapt its working precision and, when certification is required, use exact or outward-rounded arithmetic.

76 What is resolved and what is not

The reciprocal-size collision problem in the inspected repository reports is resolved for fixed multiplier degree and fixed nonendpoint bases. The answer is a convergent analytic sector decomposition, not only the existence of a leading limit. Arbitrary multiplicities and coordinate collisions are included, and several separated clusters are covered with explicit cross constants. For symmetric equal-multiplicity interior collisions, the phase-dependent first correction and the convergent motion of simple zeros are obtained in closed form.

This does not amount to a solution of every nearby problem. The estimates are not uniform as a base approaches 0 or 4, as the degree grows with N , or as the scaled coordinate grows without bound with N . Multiple or nonsimple profile zeros require another local analysis. The results do not classify every resonance of the repository's fixed-parameter recurrence problem. They do not assert that a moving sequence $H_N(q_N)$ itself satisfies the same constant-coefficient recurrence as a sequence with a fixed multiplier q .

The leading sine-kernel and oscillatory Jacobi structures, together with the classical kissing-polynomial parity mechanism, are credited prior mathematics. The new work asserted relative to the inspected repository is the explicit collision-uniform analytic package and its finite-size phase consequences. The manuscript's proof status and its historical priority status are separate: detailed proofs are provided, while an exhaustive worldwide priority determination has not been made.

77 Further research questions and proposed topics

These are proposed continuations of the present and repository results, not a claim that every formulation below is an open problem throughout the published literature.

77.1 A uniform bridge from interior collisions to endpoint confluence

Let $\theta = \theta_N \rightarrow 0$ or π while an interior cluster collides. The factors $\sin \theta$ in the bulk constants diverge, and the natural root scale changes from N^{-1} to N^{-2} . Construct one matched analytic description that recovers this paper when $N\theta_N$ is large and the endpoint continuation when $N\theta_N$ is bounded. A precise first target is $\theta_N = \lambda/\nu$ with a fixed-degree symmetric pair, retaining the collision coordinate τ and controlling the joint (λ, τ) confluence.

77.2 Mixed endpoint and interior clusters

Allow a fixed number of roots to approach 0 and 4, while the remaining roots form one or more interior clusters. The centered Vandermonde proof already identifies the types of within- and cross-cluster factors, but endpoint branch cancellations alter their valuations. Derive the correct common normalization and determine exactly which amplitudes retain even inverse-size expansions. An explicit first mixed interaction operator would connect the two existing continuation theories.

77.3 Moving zero indices and mesoscopic collision coordinates

Theorem 73.1 fixes a simple zero before letting N grow. Determine the largest regime $j = j_N \rightarrow \infty$ for which a uniform version of (73.2) holds at the j th odd profile zero. This requires balancing the large-frequency decay of Φ'_t with the growth of the collision remainder, not simply substituting (72.6) into a compact-uniform estimate. Even the exact $t = 1$ equation offers a tractable test case for sharp thresholds.

77.4 Zeros of the transport coefficient and suppressed drift

At a simple zero of Φ_t , the first phase correction disappears when Ψ_t also vanishes, or when the sampled phase has zero sine. Classify the former condition and its relation to the determinant identities of oscillatory orthogonal polynomials. Determine whether the leading nonzero displacement then has a universal second-order form. This is a question about the pair (Φ_t, Ψ_t) and finite-size transport; basic kissing-polynomial nondegeneracy should be compared with the existing literature before assigning novelty to a subproblem.

77.5 Higher correction operators in a compact basis

Theorem 66.1 uses only multiplication by U and an Euler operator. At quadratic and higher order, express the coefficients in a minimal basis of symmetric differential operators acting on $S_{d,k}$. The logarithm of the finite product in (65.8) gives a systematic generator, but the most economical collision-regular form is not supplied here. Such a form could make higher zero coefficients practical without large determinant expansions.

77.6 Growing degree and conditioning

Let $d = d_N$ grow. The explicit Cauchy constant in (67.2) is too coarse to reveal a sharp threshold for uniform convergence of a fixed number of leading sectors. Separate analytic radius loss from numerical conditioning and from factorial growth of the profile. A useful target is a regime in which relative errors

are controlled on compact sets avoiding profile zeros, together with an absolute result on the full compact set.

77.7 Eventually constant Jacobi recurrences

Replace the Catalan polynomials by a family whose three-term recurrence is eventually constant. Its spectral expression generally has two nonconstant branch amplitudes. A zero of one branch amplitude at the base can change the sector valuation. Classify those changes and derive an analogue of (66.2) that separates the universal centered Vandermonde part from the finite initial perturbation. This continues a separate question already stated in the repository's root-collision report (Part II, Section 24.5).

77.8 Multiple phase tori and resonant zero geometry

In Theorem 68.1, interior base angles produce several phase variables. Develop a convergent zero or critical-point expansion over their phase torus, including resonant rational relations among the angles. For one scalar simple zero, the implicit-function argument extends readily, but collisions of several zeros or several constraints may produce nontrivial singular phase loci. Determining those loci is a more substantial target than merely repeating the one-phase proof.

77.9 Exact certificates, interval numerics, and formalization

The shortest formalization path is algebraic: the centered Vandermonde identity, row-sign sector extraction, forced valuations, and the first-coefficient identity. Formal power series can encode these steps before complex analysis is introduced. A second layer would formalize the Catalan moment functional and the Christoffel reduction. The final layer needs holomorphic divided differences, Cauchy estimates, and the implicit-function argument. In parallel, an interval implementation could certify specific profile-zero isolations and quantitative finite- N displacement bounds. The present package does neither of those certification tasks and makes no machine-checked proof claim.

78 Conclusion

Interior root collisions have a regular description once their spectral branches are retained and the Christoffel columns are centered. Same-sign pairs account for the forced inverse-size zeros, alternating division removes every coordinate collision, and the residual amplitudes are holomorphic. This yields an exact convergent sector expansion with explicit leading terms and first correction, and it extends to several separated clusters.

For a symmetric pair inside the Catalan interval, the leading profile is classical, but its finite-size behavior contains information that the leading bulk limit suppresses. Adjacent sectors produce a $\sin(2\nu\theta)/\nu$ correction. At a simple profile zero, that correction becomes a calculable zero displacement and, more strongly, a convergent Laurent-phase expansion. The exact two-simple-root example shows both periodic cancellation and quasiperiodic scaled motion. These results answer the nonendpoint continuation explicitly posed in the repository, with the endpoint, growing-degree, and moving-zero-index regimes left as clearly separated research directions.

79 A proof-dependency map and implementation audit

This section is Appendix A of the manuscript.

The principal dependency chain is

Layer	Ingredients and output
Catalan normalization	The moment measure, orthonormal recurrence, and classical Christoffel identity reduce the problem to fixed dimension.
Centered algebra	The exponential Vandermonde and row-sign expansion prove forced powers and analytic amplitude formulas.
Uniform analysis	Alternating division and nonvanishing residual factors prove collision-uniform convergence and Cauchy remainders.
Coefficient calculus	Logarithmic differentiation gives the first operator and the finite construction of higher coefficients.
Profile structure	The bilinear kernel identity supplies sine-kernel and oscillatory Jacobi representations; Gram positivity and finite integration by parts give the needed profile information.
Zero transport	Rouché and the holomorphic implicit-function theorem turn derivative-uniform convergence into convergent phase-dependent zero expansions.

The universal proofs do not depend on any computed decimal value or any assertion in `verification.json`. The program checks independent routes wherever possible: moment matrices against confluent polynomial jets, profile alternants against oscillatory moment matrices, and sector amplitudes against direct Christoffel evaluations. The first-coefficient identity is checked both by exact full-confluence polynomials and by numerical derivatives at distinct coordinates. The multi-cluster and zero tables are experiments supporting interpretation, not additional universal proof certificates.

[Merge note: `verification.json` is shipped as `data/05-interior-collision-verification.json`.]

80 Normalization checklist

This section is Appendix B of the manuscript.

For reuse, the following conventions should be preserved together. The determinant size is N , not $N + 1$; $H_0 = 1$. The Catalan polynomials are monic with $p_1 = x - 1$, not $x - 2$, and have squared norm 1. The Vandermonde is $\prod_{i < j} (u_j - u_i)$, whereas the subset profile has cross denominator $\prod_{a \in A, b \notin A} (u_a - u_b)$. The sign $(-1)^{k(d-k)}$ in its alternant representation reconciles those conventions. Confluent rows use $1/a!$ times the a th derivative. The effective size is the global $N + d/2$. For the paired kernel identity, the final polynomial index is $N + t - 1$, while the effective size is $N + t$. The oscillatory Jacobi frequency is 2τ , not τ . These points explain the signs, the factors of 2, and the first-order phase in the main formulas.

Part VI. Growing shifts

Part VI was added on 29 September 2026 in batch 57 of ProveIt’s incoming-report intake, from one manuscript written against this report as it stood with Parts I–IV. It takes up the regime that Part I (Sections 8 and 11) and Part IV (Section 44.4) left open: the shift m growing with the determinant size N . For the one-factor family $D_N^{(m)}(a, 1)$ with $a \geq 0$ and arbitrary relative growth of N and m , it proves an exact criterion for when the factor $x + a$ is invisible, a uniform bound on the largest inverse Jacobi zero, a size-biased inverse-zero law equal to a shifted Catalan (Marchenko–Pastur) law, an all-orders hierarchy of critical scales, an algebraic free-energy limit, the exact total-variation transition away from the nominal Poisson law, and a uniform saddle prefactor in proportional growth. Several moving factors, both endpoint exponents growing, and a sharp extreme-zero theorem remain open. Parts I–V are unchanged except for dated pointers to this addition (Sections 8 and 11, Section 44.4 and Section 57.4), a longer line on the title page (now set in a smaller font, with the title page’s vertical spaces reduced by a further 7 mm, so that the page still fits), a sentence run into the last paragraph of the abstract, a contents entry, three bibliography entries used only by Part VI, a clause in the closing note on source dates, one new label on Part IV’s Section 57.4 so that it can be cited, and the `graphicx` package, which Part VI’s two figures need. Section 81 records the provenance and fixes the notation of the addition.

81 Provenance and conventions of Part VI

81.1 One manuscript, one addition

Part VI is batch-57 manuscript 02, *A Catalan Law Behind Growing Shifts*, subtitled *Three critical scales, an all-orders hierarchy, and uniform Hankel asymptotics*, an AI-assisted research article dated 29 September 2026 (21 A4 pages as delivered). Its byline reads “AI-assisted research prepared for Vladimir Reshetnikov / In continuation of the ProveIt repository”, and its PDF metadata gives the same author string; it names no assistant. It arrived as the archive `ProveIt_Catalan_Growing_Shifts.zip` (inner directory `Catalan_Growing_Shifts/`, main file `article.tex`) in ProveIt commit `6ad01e9c8` and was placed in `70b39943a`.

It was written against a ProveIt commit (a fix commit for the batch-54 Fabius-tree arrivals), and it records two Git blobs there, both checked for this merge:

commit	1085b506d65e207a05b7e9c861bb1fe88432fe38
this report’s <code>article.tex</code>	b23831b8f20efab08342b11bb4939809d0a8047e
this report’s <code>README.md</code>	ce3436af04afdb64c683d9d36be3f0b45c6ccf5e

At that commit the article held Parts I–IV; Part V was written into it afterwards, in `b2aed2d23`, and that is the only change to the report between the pin and the placement commit. Every passage the manuscript continues (Part I’s Sections 8 and 11, Part I’s Theorem 2.1, Part IV’s Section 44.4 and its research questions in Section 57) is textually unchanged between the pin and the placement commit, only shifted in the source, apart from the dated notes that Part V added to Sections 44.4 and 57.1 and its label on Section 57.2; so the manuscript’s “repository”, “combined source manuscript” and “inspected reports” are Parts I–IV as printed above, its “Part I” is Part I, and its “later endpoint-confluence part” is Part IV. Its source audit also names Part V’s shipped source notes, `05-interior-collision-SOURCES.md`, which were already in the repository at the pin (placed in `26473dfa0`); it read them only to avoid treating that fixed-degree direction as new. The line numbers the source audit gives for “the combined source” refer to the article at the pin, not to the present file.

The manuscript’s program, its figure script and its recorded outputs are shipped beside the earlier files with the prefix `06-growing-shifts-` (Part I’s unprefix files count as 01, and Parts II–V have 02–to 05–), its dependency file under `data/`, its delivered `Makefile` under `code/`, and its source audit,

proof audit and build check, delivered as SOURCES.md, PROOF_AUDIT.md and BUILD_CHECK.md, at the report root:

- code/06-growing-shifts-verify.py and code/06-growing-shifts-make_figures.py (delivered as code/verify.py and code/make_figures.py; the second imports the first by its delivered name, Section 99);
- code/06-growing-shifts-Makefile (delivered at the package root as Makefile; it must not be run in this directory);
- data/06-growing-shifts-verification_summary.json and the four CSV files data/06-growing-shifts-critical_scales.csv, -spectral_moments.csv, -free_energy.csv and -saddle_prefactor.csv (delivered under data/ without the prefix; the CSV files have CRLF line endings, as delivered);
- data/06-growing-shifts-requirements.txt (delivered as requirements.txt; it states the unpinned ranges numpy>=1.24, scipy>=1.10, matplotlib>=3.7);
- 06-growing-shifts-SOURCES.md, 06-growing-shifts-PROOF_AUDIT.md and 06-growing-shifts-BUILD_CHECK.md (the last describes the delivered 21-page PDF, not this report);
- figures/06-growing-shifts-determinant_crossover.pdf and figures/06-growing-shifts-free_energy.pdf, Figures 1 and 2.

All but the two figures are byte-identical to the delivery. The delivered figure PDFs embedded their DejaVu Sans text as Type 3 fonts, which the rest of this report does not use; the shipped figures were regenerated for this merge by the delivered code/make_figures.py, unchanged, on a copy of the delivery with Matplotlib 3.10.8 (the version recorded in the delivered figures), NumPy 2.3.5 and SciPy 1.17.0 and a local matplotlibrc setting pdf.fonttype: 42, so that they embed TrueType fonts. They plot the same curves but are not byte-identical to the deliveries. The manuscript, its PDF, its delivery README, its delivered figures and its checksum list are not shipped; they survive in the arrival commit (the checksum list was verified at placement and retired).

The manuscript contributes, in the order printed here: the repository question and what is classical (Section 82); the normalization, the concentration index β and the main asymptotic picture (Section 83); an exact Jacobi reduction, the normalized coefficients and an exact invisibility criterion valid for every growth of m (Section 84); positive inverse-zero weights and a Riccati recurrence for all their moments (Section 85); a uniform Catalan majorant and the bound $W \leq 8\beta$ on the largest inverse zero (Section 86); the first (Gaussian) critical scale with explicit error bounds (Section 87); the size-biased inverse-zero law (Section 88); the all-orders hierarchy of critical scales (Section 89); the algebraic free energy and a nonlinear central limit theorem (Section 90); the nominal-Poisson transition (Section 91); the saddle prefactor in proportional growth (Section 92); boundary regimes and the physical sizes of the perturbation (Section 93); its checks (Section 94); proof dependencies and limitations (Section 95); ten research directions (Section 96); a conclusion; a formula sheet (Section 98); and its reproduction instructions (Section 99). Every result, proof, remark, limitation and question of the manuscript is printed. Where it re-derives something Parts I–V already prove, the overlap is marked (Section 81.5).

The manuscript's own abstract reads:

Consider the shifted Catalan Hankel determinant $D_N^{(m)}(a, 1) = \det(aC_{i+j+m} + C_{i+j+m+1})_{0 \leq i, j < N}$. The repository's fixed-shift formulas do not give uniform asymptotics when m grows with N . We develop such a theory for a single moving linear factor, allowing *arbitrary relative growth* of N and m . An exact normalization reduces the determinant to a polynomial $\mathcal{R}(\tau)$ whose negative zeros define positive weights of total mass one. An explicit quadratic index β controls these weights, and an elementary Riccati argument proves that their maximum is at most 8β .

A size-biased inverse-zero measure converges to a shifted and scaled Marchenko–Pastur law; its moments are binomial transforms of the Catalan numbers. This single law determines an infinite

hierarchy of renormalized limits at $\tau \asymp \beta^{-(r-1)/r}$. In particular, the approximation $\mathcal{R}(\tau) \sim e^\tau$ breaks down at $\beta^{-1/2}$, whereas the determinant's coefficient distribution separates from $\text{Pois}(\tau)$ at $\beta^{-2/3}$. At $\tau \asymp \beta^{-1}$, the hierarchy resums to an explicit algebraic free-energy law, valid also when m/N tends to zero or infinity. A separate saddle calculation supplies a uniform leading prefactor when m/N remains in a compact subset of $(0, \infty)$.

All principal claims have written proofs. Exact-arithmetic audits and floating-point diagnostics accompany the article. The underlying Jacobi, Christoffel, and Poisson-approximation machinery is classical; the claimed advance is relative to the inspected repository questions. Worldwide priority is not established, and no Lean formalization is claimed.

Its title page carried this scope line:

Scope. One moving linear factor; nonnegative real perturbations for probability and sharp-order statements; complex compact limits where stated. No claim is made about arbitrary growing-degree multipliers.

Status (manuscript 02). This is an unrefereed, AI-assisted manuscript with conventional written proofs, not a Lean or Rocq formalization and not an interval-certified computation. It treats one additional linear factor $x + a$ beside a growing endpoint power x^m , with $a/b \geq 0$ for its probabilistic and sharp-order statements, and complex limits only where stated. Classical ingredients are credited, not presented as new: the Catalan moment measure, the one-factor Christoffel identity (Krattenthaler [3]; re-derived in the present normalization), the finite Jacobi hypergeometric representation (DLMF [4]), large-parameter Jacobi-zero asymptotics (Dette–Studden [14], context only), the Barbour–Hall Bernoulli-sum Poisson bound [15], the only Poisson approximation estimate used, and mod-Poisson and higher-order Chen–Stein approximation (Méliot–Nikeghbali–Visentin [16], context only). The fixed- $m {}_0F_1$ limit is a standard special-function phenomenon, not claimed as new, and the coefficient formula (84.5) is Part I's. The claimed contribution relative to the inspected reports is the integrated calculation: the exact normalization and concentration index, the uniform inverse-zero bound, the size-biased Catalan law in every relative-growth regime, the all-orders hierarchy, the distinct determinant and nominal-Poisson thresholds, and the algebraic resummation with a proportional-growth prefactor. Equivalent formulations may exist in the orthogonal-polynomial literature; the bounded literature check does not settle first-discovery priority, and requested PDF fetches of the Krattenthaler and Dette–Studden papers did not succeed (only their abstracts were read). Not claimed: several moving factors or arbitrary growing-degree multipliers; both endpoint exponents growing; negative perturbations, across which the positive-axis statements need not hold; an extreme-zero theorem (the constant 8 is not claimed sharp); uniformity in the hierarchy order q ; a saddle prefactor uniform as m/N or t approaches 0 or ∞ ; exhaustive inspection of the repository. The recorded assertions (270 direct determinant comparisons, 420 moment parameter pairs with 5,040 moment equalities and 5,040 majorant checks, 60 limiting-recurrence checks, 16 floating-point Jacobi-root parameter pairs) are finite audits, and the CSV tables are floating-point diagnostics, not outward-rounded certificates. No proof assistant was run, nothing in Part VI is formalized, and its placement beside a Lean project confers no formal status. (Sources: the manuscript, its delivery README, and its source and proof audits, 06-growing-shifts-SOURCES.md and 06-growing-shifts-PROOF_AUDIT.md.)

81.2 Reading conventions

Inside Sections 82–99, “this article” and “here” mean manuscript 02 as printed in Part VI; “the repository”, “the inspected reports” and “the combined source manuscript” mean Parts I–IV at the pin 1085b506d. “The package” and “the delivered archive” mean the delivered archive, whose files are shipped under the prefixed names listed in Section 81.1. The manuscript's two appendices are printed as Sections 98 and 99, so that Part I's appendices keep their letters. Its Sections 1–16 are Sections 82–97, so a manuscript number $n.j$ of a section, theorem, proposition, lemma, corollary, definition, remark or equation is printed as $(n + 81).j$ (for example, its Theorem 2.1 is Theorem 83.1 and its Theorem 7.2 is Theorem 88.2),

and its Appendices A and B are Sections 98 and 99; its two figures are Figures 1 and 2. The shipped 06-growing-shifts-PROOF_AUDIT.md and 06-growing-shifts-SOURCES.md use the manuscript's numbers. The manuscript cites Krattenthaler's Ramanujan Journal paper (with arXiv:2101.04225) and DLMF equation 18.5.7; they are Part I's references [3] and [4]. Bracketed paragraphs beginning "Merge note" are this merge's, not the manuscript's.

81.3 Notation of Part VI

Part VI keeps the manuscript's symbols. Within Sections 82–99 every symbol has its Part VI meaning, except in merge notes and where a sentence names Part I, II, III, IV or V or cites one of their equations. The determinants are Part I's: $D_N^{(m)}(a, b)$ and $H_N^{(m)}$ are Part I's (1.2) and (1.5), a and b are Part I's parameters, and the manuscript works at $b = 1$ except in Section 93.2, where it restores $b > 0$ (there a is replaced by a/b). In Part IV's terms, $D_N^{(m)}(a, b) = b^N H_N(x^m(x + \varepsilon))$ with $\varepsilon = a/b$: a cluster of $m + 1$ roots at the endpoint 0, m of them fixed there and one displaced outward when $\varepsilon > 0$. Table 7 lists the symbols that a reader of Parts I–V is likely to misread. No printed symbol was renamed, but two macros were: the manuscript's calligraphic $\mathcal{R}$ for the normalized determinant (83.4) is typed `\Rgs` in the source (it prints the same $\mathcal{R}$), so that no bare `\R` stands beside Part II's $\mathcal{R}_0$; and the manuscript's expectation $\mathbb{E}$ is typed `\Egs`, because the report's `\E` is Part II's sans-serif shift E. No normalization changed. Five readings need particular care: β is the concentration index (83.5), not a Jacobi endpoint exponent (which is $\frac{1}{2}$ here) nor Part I's $\beta_j^\pm$; $c = v/(NB)$ is not Part I's coefficient $c_{N,m,k}$ nor the base c of Parts II, III and V; $\nu_{N,m}$ and ν_θ are probability measures, not Part IV's and Part V's centred size ν ; b is Part I's parameter in Section 93.2 but $(N + m + 1)/N$ in Section 92 (a clash inside the manuscript); and t is $\beta\tau$ in Sections 83 and 90 but τ/N in Section 92 (also inside the manuscript).

Table 7: Symbols and notions of Part VI. The last column gives the meaning of the same letter in Part I, II, III, IV or V, or the tempting false reading.

Symbol or notion	Meaning in Part VI	Not to be confused with
ν, B, c, μ	$m + \frac{1}{2}$, $N + m + 1$, $v/(NB)$ and the scalar $NB/(4\nu)$ (83.3); $a = \tau/\mu$	Part IV's coordinates ν_j ; Part III's rectangular product $B_{a,b}(N)$; Part I's $c_{N,m,k}$ and the base c of Parts II, III, V; Part I's measure μ_m and the Catalan measure μ of Parts II, IV, V.
$\mathcal{R}(\tau) = \mathcal{R}_{N,m}(\tau)$	the normalized determinant (83.4), a polynomial of degree N in τ	Part II's maximal order $\mathcal{R}_0$; Part III's order $R(t, c)$; Part IV's ratio R_N and hyperbolic generator $\mathcal{R}$; Part I's polynomials $R_n(y)$.
τ, s, u, t	the normalized perturbation; the scaled variables $\tau\sqrt{\beta}$, $\tau\beta^{(q-1)/q}$ and $\beta\tau$ (but $t = \tau/N$ in Section 92)	Part IV's kernel variable t and Part V's collision coordinate τ ; Part II's number s of interior roots; Part V's coordinates u_i ; Parts III and V's multiplicity t .
$\beta, \theta_{N,m}, \theta, \gamma$	the concentration index and the shape (83.5), its limit, and $1 - \theta$	Part I's $\beta_j^\pm$; the Jacobi exponents $m - \frac{1}{2}$, $\frac{1}{2}$; Part V's base angle θ ; Part II's leading coefficient γ .
x_j, w_j, W, S_r	the zeros of $p_N^{(m)}$, the inverse-zero weights $1/(\mu x_j)$, their maximum and power sums (85.1)	Part V's spectral base w and profiles $S_{d,k}$; Part IV's $S(t, y)$; Part III's $w = N + t/2$.
$L = \mathcal{R}'/\mathcal{R}, L_\theta, M(z)$	the logarithmic derivative, its limit (90.3), the limiting moment series (88.8)	Part III's $L = \text{ord}(z^2)$; Part V's $L(h; u, w)$; Part I's minors $M_{m,\ell}$; Part II's Christoffel matrix M_N .
$\omega, Y, \text{MP}_1, C(z)$	the Catalan measure (84.1) (Part I's μ_0), a random variable with that law, its name, the Catalan generating function	Part IV's $Y = \sum_i y_i$; Part IV's kernel $C(t, y)$.

Symbol or notion	Meaning in Part VI	Not to be confused with
$\nu_{N,m}, \nu_\theta, d_r(\theta), D_r$	the size-biased inverse-zero measure (88.1), its limit (88.3), the limiting moments (88.5), and S_r/β^{r-1}	Part IV's and Part V's centred size ν ; Part I's $d = \binom{m}{2}$; Part V's d ; Part I's $D_N^{(m)}$.
$\mathcal{F}_\theta, Q_\theta, V_\theta$	the free energy (90.1), the limiting mean and variance of Corollary 90.2	Part I's generating function $F_m(t; a, b)$; Part IV's profiles $F_t^\pm$; Part III's quadratic factors Q_a .
$K_\tau, p_j, \lambda, \delta$	the coefficient distribution, a sum of Bernoulli variables with parameters p_j (90.8); its mean and $\tau - \lambda$	Part I's $K = (m-1)^2$; Part IV's kernels $K^\pm$; the orthogonal polynomials p_n (and $p_N^{(m)}$ in Section 84); Part II's rates λ_k ; Part IV's partitions λ .
$\Phi, N(\cdot, 1), \sigma$	the standard normal distribution function; a normal law; a limit parameter in $[0, \infty]$	Part V's profiles Φ_t ; the determinant size N ; Part III's subset sums $\sigma(S)$.
$r, b, q, \ell, h(x), f, \mathcal{A}$	in Section 92: $(m + \frac{1}{2})/N, (N + m + 1)/N$, the saddle point, an auxiliary in (92.5), $-f''$, the rate (92.2), the prefactor (92.6)	the moment index r elsewhere in Part VI; Part I's parameter b ; Part I's $q(t)$ and Part II's multiplier q ; the multiplicity ℓ of the root 4 (Parts II, IV); Part I's h_m (1.6), Part IV's and Part V's $h = 1/\nu$; Part IV's constant $\mathcal{A}_{m,\ell}$.
$\rho, b_\rho, b_*, T_k, A_{N,m}(k)$	the limit of m/N ; $\lim N\beta$; an auxiliary; the k th term of $\mathcal{R}$; its normalized coefficient (84.4)	Part IV's $\rho = \binom{m}{2} + \binom{\ell}{2}$; Part IV's A_j ; Part V's $A_t(\theta)$.
$d_{TV}, \text{Pois}, \mathcal{L}, \mathbb{E}, \mathbb{P}, \text{Var}$	total variation (half the ℓ^1 distance), a Poisson law, the law of a random variable, expectation, probability, variance	the shift E of Part II.

81.4 What Part VI answers, and what stays open

Part I's status paragraph (Section 8) restricts its estimates to fixed m, a, b , “not claimed to be uniform . . . when m grows with N ”, and its Section 11 names joint limits with growing m as requiring uniform asymptotics; its dated notes of batches 39 and 53 there left them open. Part IV treats $a \rightarrow 0$ at fixed m (Theorem 47.4 and Section 44.4) and repeats that “joint limits with m growing with N remain open”, and its research question “Multiplicities growing with determinant size” (Section 57.4) asks for a regime $m = m_N$, $\ell = \ell_N$ in which the factorized leading profile or the first correction remains uniform. Part VI answers the growing- m question for the one-factor family with real $a/b \geq 0$, at every relative growth of N and m : the ratio $D_N^{(m)}(a, b)/D_N^{(m)}(0, b)$ tends to 1 exactly when $\tau = (a/b)NB/(4\nu) \rightarrow 0$ (Corollary 84.2); when $N, m \rightarrow \infty$ its logarithm is governed by β , with the critical scales $\beta^{-1/2}$, $\beta^{-2/3}$ and β^{-1} (Theorem 83.1), the size-biased inverse-zero law (Theorem 88.2), every fixed level of the hierarchy (Theorem 89.1), the algebraic free energy (Theorem 90.1) and, in proportional growth, a uniform prefactor (Theorem 92.1). In Part IV's terms this is $\ell = 0$ and the cluster $x^m(x + \varepsilon)$, $\varepsilon = a/b \geq 0$, with one displaced root. The manuscript does not cite Part IV's Section 57.4; the identification is this merge's.

Not answered. Part IV's Section 57.4 stays open for several displaced roots, for $\ell > 0$, and for the first correction in the sense of Part IV's Theorem 47.1; a single approximation uniform for $1 \leq m \leq N$ bridging the fixed- m and growing- m regimes is Part VI's own question (Section 96.3). Part VI's Section 96.4 poses several moving factors, and its Section 96.5, both endpoint exponents growing, is related to Part IV's question “Other Jacobi endpoint exponents” but not the same. Coalescing roots and $a/b < 0$ (Part I's other caveats) are not treated. A combinatorial interpretation of Part I's $c_{N,m,k}$ and a product formula for its minors $M_{m,\ell}(N)$ remain open; Part VI's Section 96.8 asks for a combinatorial coefficient model in the same spirit. Part V's “Not claimed” (growing degree, or zero indices growing with N) concerns interior roots and is neither answered nor contradicted.

Consistency with Parts I and IV (observations of this merge).

1. (84.5) is Part I's (2.1) at $b = 1$, as the manuscript says: (84.4) is the statement $A_{N,m}(k)\mu^k/k! = \binom{N}{k}(B)_k/(4^k(v)_k) = c_{N,m,k}/H_N^{(m+1)}$.
2. For fixed m , Corollary 84.2 reduces to Part IV's threshold: $\tau = \varepsilon N(N+m+1)/(4m+2) \rightarrow 0$ exactly when $N^2\varepsilon \rightarrow 0$ (Theorem 47.4 with $\ell = 0$ and $m+1$ roots at 0). If $N^2\varepsilon \rightarrow y$, then $v\tau \rightarrow y/4$ and (93.1) gives the limit ${}_0F_1(; m + \frac{1}{2}; y/4)$, which must therefore be Part IV's profile ratio (47.10) at the coordinates $(0, \dots, 0, y)$. By hand for $m = 0$ and $m = 1$, Part IV's kernel quotient (46.8) gives $\cosh \sqrt{y}$ and $\sinh \sqrt{y}/\sqrt{y}$, which are ${}_0F_1(; \frac{1}{2}; y/4)$ and ${}_0F_1(; \frac{3}{2}; y/4)$. For $m \rightarrow \infty$ with τ bounded, the limit is e^τ instead (Section 93.1), with the corrections of Theorem 89.1.
3. The table of Section 94.1 is a rounded copy of the shipped data/`06-growing-shifts-critical_scales.csv` (checked for this merge).

81.5 Where the merge had to choose

1. *Placement.* Part VI is printed after Part V and before Part I's appendices, so that no section, equation, theorem, figure or table number of Parts I–V moves (checked against the `.aux` file of a build of the committed text). Its sections continue the numbering (Sections 81 to 99), with the concordance of Section 81.2. The manuscript's `\clearpage` before its Appendix B was dropped.
2. *Labels.* Every label of Part VI carries the prefix `gs:`; without it two of the manuscript's labels, `eq:measure` and `eq:h`, would have collided with Part I's bare labels. Labels were added to the manuscript's conclusion, its two appendices, its second figure, four of its subsections and its ten research subsections, and to Part IV's Section 57.4 so that it can be cited; no existing label was changed.
3. *Overlaps with Parts I–V.* The manuscript re-derives, without assuming the report correct, the Catalan measure, the one-factor Christoffel identity, the Jacobi hypergeometric representation and Part I's coefficient formula, which it credits (merge note after (84.5)). Printed with a pointer and kept. Everything else has no counterpart in Parts I–V, all of which keep the multiplier degree and the endpoint multiplicities fixed. No statement of Parts I–V was changed.
4. *Bibliography.* The manuscript's Krattenthaler and DLMF references are [3] and [4]; its Dette–Studden, Barbour–Hall and Méliot–Nikeghbali–Visentin references are added to the common bibliography as [14], [15] and [16]. Its two references to the pinned repository report and to its README and “continuation source notes” are replaced by Parts I–IV and the pin recorded in Section 81.1; the two sentences that cited them now read “(Parts I and IV above, and this report's README at the pin)” and “(Part I, Theorem 2.1)”. The first entry described the pinned article's “Part I, explicit coefficients and the fixed-parameter scope of its asymptotics; Part I's growing-shift agenda; Part IV's fixed multiplicity limitations”; the second said that those files “distinguish existing fixed-multiplicity endpoint and nonendpoint work from the growing-shift question”. Both pointed to `SOURCES.md`, shipped as `06-growing-shifts-SOURCES.md`.
5. *File names and figures.* Merge notes in Sections 94 and 99 give the shipped names beside the delivered ones. The manuscript's reproduction commands are printed as delivered but must not be run in place here: the program's default output directory is the report's shared `data/`, the figure script imports the program by its delivered name, and the delivered Makefile acts on Part I's verifier and on this report's `article.tex`; a recipe on a scratch copy follows them. The two figures are the regenerated ones (Section 81.1), included from `figures/`; the `graphicx` package was added to the preamble for them.
6. *Front matter.* The manuscript's title page, abstract, scope line and contents are replaced by Section 81.1, which prints the abstract, the scope line and the status.
7. *Notation and macros.* The manuscript's symbols are kept, with the scoping convention and Table 7 of Section 81.3. Its $\mathbb{R}$ ($\mathcal{R}$) is typed `\Rgs` and its $\mathbb{E}$ ($\mathbb{E}$) `\Egs`; both print as in the manuscript. Its d is `\, \mathit{d}`, the report's is `\mathit{d}`, so every d of Part VI is printed `\, \mathrm{d}`, with the manuscript's spacing. Its macros `\PP`, `\Pois`, `\Var`, `\TV`, `\MP`, `\Law` and `\calF` were added to the

preamble with the manuscript's definitions; its \RR and \CC are the report's, with the same definitions; its unused \NN (the report's \NN is $\mathbb{N}_0$), \supp, \Cat and \onehalf were dropped. No mathematical text of the manuscript was changed.

82 The repository question and the contribution

The relevant ProveIt project is *Shifted Catalan Hankel Polynomials*, inside the Catalan-and-ballot Hankel-determinant reports. Its Part I gives an exact coefficient formula and sharp recurrences for $x^m(a + bx)$, but explicitly restricts its asymptotic estimates to fixed m, a, b . Its research agenda asks for joint limits with growing m . The later endpoint-confluence part fixes its endpoint multiplicities and explicitly does not make its constants uniform in them (Parts I and IV above, and this report's README at the pin). The snapshot inspected here is

1085b506d65e207a05b7e9c861bb1fe88432fe38.

The combined source manuscript has blob b23831b8f20efab08342b11bb4939809d0a8047e.

We address the following precise version of the growing-shift question:

How does $D_N^{(m)}(a, 1)/D_N^{(m)}(0, 1)$ behave when $N, m \rightarrow \infty$ with no restriction on their ratio, and how far can its simplest exponential approximation be trusted?

The determinant also defines a natural probability distribution by normalizing its nonnegative coefficients at $a > 0$. We ask the corresponding question for that distribution, rather than assuming that accuracy of the normalizing constant and accuracy of the probability law have the same threshold.

The answer is organized by an exact parameter β , not by N alone. The three most visible scales are

$$\beta^{-1/2}, \quad \beta^{-2/3}, \quad \beta^{-1}. \quad (82.1)$$

The first is a relative-error threshold for the determinant. The second is a *mean-displacement threshold* for comparison with the nominal law $\text{Pois}(\tau)$; a Poisson law with the corrected mean remains accurate much longer. The third is a genuinely nonlinear regime for the logarithm of the determinant. Between the first and third lies a full hierarchy of exponents $(r - 1)/r$.

82.1 What is classical and what is being added

The Christoffel determinant reduction is classical; a modern detailed account is Krattenthaler [3]. Jacobi hypergeometric formulas are recorded in DLMF [4]. Large-parameter zero asymptotics for Jacobi polynomials already have an extensive literature; Dette–Studden [14] is particularly relevant to regimes in which parameters outgrow the degree. Bernoulli sums and higher-order Poisson corrections belong to the established Chen–Stein and mod-Poisson frameworks [15, 16]. None of these broad mechanisms is claimed as new.

The addition relative to the inspected reports is the following integrated calculation: the exact normalization and concentration index; a uniform inverse-zero bound; the explicit size-biased Catalan law in every relative growth regime; the associated all-orders critical hierarchy; the distinct sharp determinant and nominal-Poisson thresholds; and an algebraic resummation with a proportional-growth prefactor. Equivalent formulations may exist in the orthogonal-polynomial literature. The bounded literature check conducted for this article does not settle first-discovery priority.

83 Normalization and a map of the main results

Let

$$C_k = \frac{1}{k+1} \binom{2k}{k}, \quad H_N^{(m)} = \det(C_{i+j+m})_{0 \leq i, j < N}, \quad (83.1)$$

and

$$D_N^{(m)}(a, b) = \det(aC_{i+j+m} + bC_{i+j+m+1})_{0 \leq i, j < N}. \quad (83.2)$$

Throughout the article, $N \geq 1$ and $m \geq 0$ are integers. Set

$$v = m + \frac{1}{2}, \quad B = N + m + 1, \quad c = \frac{v}{NB}, \quad \mu = \frac{NB}{4v} = \frac{1}{4c}, \quad (83.3)$$

and define

$$\mathcal{R}_{N,m}(\tau) = \frac{D_N^{(m)}(\tau/\mu, 1)}{H_N^{(m+1)}}. \quad (83.4)$$

Thus the original perturbation is $a = \tau/\mu$, not $a = \tau$. The concentration and shape parameters are

$$\beta = \frac{1 + c(v + 3/2)}{v + 1}, \quad \theta_{N,m} = \frac{1}{(v + 1)\beta}. \quad (83.5)$$

In particular, $0 < \theta_{N,m} < 1$.

Theorem 83.1 (Main asymptotic picture). *Suppose $N, m \rightarrow \infty$. Then $\beta \rightarrow 0$ and the following hold.*

- (i) *For $\tau \geq 0$, $\mathcal{R}(\tau)/e^\tau \rightarrow 1$ if and only if $\beta\tau^2 \rightarrow 0$. If $\tau\sqrt{\beta} \rightarrow s \in [0, \infty)$, then $\mathcal{R}(\tau)e^{-\tau} \rightarrow e^{-s^2/2}$. If $\tau\sqrt{\beta} \rightarrow \infty$, this ratio tends to zero.*
- (ii) *Define K_τ by $\mathbb{P}(K_\tau = k) = [z^k]\mathcal{R}(z)\tau^k/\mathcal{R}(\tau)$. If $\beta\tau \rightarrow 0$ and $\beta\tau^{3/2} \rightarrow \sigma \in [0, \infty]$, then*

$$d_{\text{TV}}(\mathcal{L}(K_\tau), \text{Pois}(\tau)) \rightarrow 2\Phi(\sigma/2) - 1, \quad (83.6)$$

with the right side interpreted as 1 at $\sigma = \infty$.

- (iii) *If $\theta_{N,m} \rightarrow \theta \in [0, 1]$, then, for every $t \geq 0$,*

$$\beta \log \mathcal{R}(t/\beta) \rightarrow \mathcal{F}_\theta(t). \quad (83.7)$$

The convergence is uniform on compact real t -intervals. The derivative $L_\theta = \mathcal{F}'_\theta$ is the positive solution of

$$\theta t L_\theta(t)^2 + L_\theta(t) = \frac{1}{1 + (1 - \theta)t}, \quad \mathcal{F}_\theta(0) = 0. \quad (83.8)$$

The proof is distributed across Sections 85–91. The more informative result is the inverse-zero law in Theorem 88.2, which also yields every intermediate critical scale in Theorem 89.1.

84 An exact Jacobi reduction, with normalization checked

The Catalan numbers are the moments of

$$d\omega(x) = \frac{1}{2\pi} \sqrt{\frac{4-x}{x}} \mathbf{1}_{(0,4)}(x) dx. \quad (84.1)$$

Indeed, $x = 4u$ reduces the k th moment to a beta integral and gives $4^k \Gamma(k + 1/2) / (\sqrt{\pi} \Gamma(k + 2)) = C_k$. Consequently, the Hankel matrix for $H_N^{(m)}$ is the Gram matrix of $1, x, \dots, x^{N-1}$ for $x^m d\omega(x)$. It is positive definite.

Let $p_N^{(m)}$ be the monic degree- N orthogonal polynomial for this positive measure. The one-factor Christoffel identity gives

$$D_N^{(m)}(a, 1) = (-1)^N H_N^{(m)} p_N^{(m)}(-a). \quad (84.2)$$

For completeness, form the usual moment determinant with first N rows $(C_{i+j+m})_{j=0}^N$ and last row $(1, y, \dots, y^N)$. Dividing by $H_N^{(m)}$ gives a monic polynomial orthogonal to degrees below N . Equivalently, integrating $\prod_{j=1}^N (y - x_j)$ against the squared Vandermonde Gram integral gives that same polynomial. Substituting $y = -a$ proves (84.2), including its sign. This is the standard identity, rederived only to fix conventions [3].

Under $y = 1 - x/2$, the weight is proportional to $(1 - y)^{m-1/2}(1 + y)^{1/2}$ on $(-1, 1)$. Thus the quotient of endpoint values is a Jacobi quotient. The finite hypergeometric representation [4, 18.5.7] yields

$$\frac{D_N^{(m)}(a, 1)}{H_N^{(m+1)}} = {}_2F_1\left(\begin{matrix} -N, B \\ v \end{matrix}; -a/4\right). \quad (84.3)$$

All normalizing factors cancel because the quotient equals one at $a = 0$.

Proposition 84.1 (Exact normalized coefficients). *One has*

$$\mathcal{R}(\tau) = \sum_{k=0}^N A_{N,m}(k) \frac{\tau^k}{k!}, \quad A_{N,m}(k) = \prod_{j=0}^{k-1} \frac{(1 - j/N)(1 + j/B)}{1 + j/v}. \quad (84.4)$$

Every $A(k)$ is positive for $0 \leq k \leq N$, and $A(0) = A(1) = 1$. Moreover $A(k) \leq 1$.

Proof. Expand (84.3) and use $a = 4c\tau$. For $0 \leq j < N$,

$$(1 - j/N)(1 + j/B) = 1 - j(1/N - 1/B) - j^2/(NB) \leq 1,$$

where $B > N$. The denominator is at least one. □

The unnormalized coefficient formula is therefore

$$D_N^{(m)}(a, 1) = H_N^{(m+1)} \sum_{k=0}^N \binom{N}{k} \frac{(B)_k}{4^k (v)_k} a^k. \quad (84.5)$$

This is the repository's existing coefficient theorem, not a new coefficient formula (Part I, Theorem 2.1). It is useful here because it is exact for every N, m and remains meaningful when fixed- m asymptotics do not. [Merge note: (84.5) is Part I's (2.1) at $b = 1$, since $B = N + m + 1$ and $v = m + \frac{1}{2}$; the normalization (83.4) divides by Part I's $c_{N,m,0} = H_N^{(m+1)}$. The measure ω of (84.1) is Part I's μ_0 (3.1), and $x^m d\omega(x) = d\mu_m(x)$; the monic polynomial $p_N^{(m)}$ is Part I's (3.3), whose value at $x = -a$ gives (84.3); and (84.2) is the one-factor Christoffel identity for μ_m , of which Part I's Lemma 4.1 (for μ_0 and several factors) and its successors in Parts II, IV and V are relatives. The manuscript re-derives all three without assuming Part I.]

Corollary 84.2 (An exact invisibility criterion). *For every N, m and $\tau \geq 0$,*

$$1 + \tau \leq \mathcal{R}(\tau) \leq e^\tau. \quad (84.6)$$

Along any parameter sequence with $a \geq 0$,

$$\frac{D_N^{(m)}(a, 1)}{H_N^{(m+1)}} \longrightarrow 1 \iff \frac{aN(N + m + 1)}{4(m + 1/2)} \longrightarrow 0. \quad (84.7)$$

No condition that m be fixed or tend to infinity is needed.

Proof. Use the constant and linear coefficients for the lower bound and $A(k) \leq 1$ for the upper bound. Apply the squeeze theorem in one direction and $\mathcal{R} \geq 1 + \tau$ in the other. □

85 Inverse-zero weights and the exact concentration index

85.1 Positive weights from orthogonality

A monic orthogonal polynomial of degree N for a positive measure on $(0, 4)$ has N simple zeros in that interval. One elementary proof counts its sign changes there. If it had fewer than N , multiply it by the polynomial whose roots are precisely the sign-changing zeros. The product would have constant nonzero sign almost everywhere, contradicting orthogonality to that polynomial of degree less than N . The degree bound then forces every zero to be simple and internal.

Write these zeros as $x_1, \dots, x_N$, and put

$$w_j = \frac{1}{\mu x_j} > 0, \quad S_r = \sum_{j=1}^N w_j^r, \quad W = \max_j w_j. \quad (85.1)$$

Equations (84.2) and (83.4) imply

$$\mathcal{R}(\tau) = \prod_{j=1}^N (1 + \tau w_j), \quad S_1 = 1. \quad (85.2)$$

In particular, $W \leq 1$. The weights describe a deterministic zero configuration; they are not random eigenvalues.

85.2 Riccati recurrence

The hypergeometric differential equation transforms into

$$\tau(1 + c\tau)\mathcal{R}'' + [v + (v + 3/2)c\tau]\mathcal{R}' - v\mathcal{R} = 0. \quad (85.3)$$

Set $L = \mathcal{R}'/\mathcal{R}$. Near the origin,

$$L(\tau) = \sum_{k \geq 0} (-1)^k S_{k+1} \tau^k. \quad (85.4)$$

Substitution in (85.3) gives the following identity.

Proposition 85.1 (All inverse-zero moments). *For every $k \geq 1$,*

$$(v + k)S_{k+1} = \sum_{i=1}^k S_i S_{k+1-i} + c(k + v + 1/2)S_k - c \sum_{i=1}^{k-1} S_i S_{k-i}, \quad S_1 = 1. \quad (85.5)$$

In particular,

$$S_2 = \beta, \quad S_3 = \frac{2\beta + c(v + 5/2)\beta - c}{v + 2}. \quad (85.6)$$

Proof. The coefficient of τ^k in $\tau(1 + c\tau)(L' + L^2) + [v + (v + 3/2)c\tau]L - v$ is zero. Multiply that coefficient by $(-1)^{k-1}$ and collect the S_{k+1} terms. The derivative contributions combine to $v + k$; the coefficient multiplying cS_k is $k - 1 + v + 3/2$. This gives (85.5). The cases $k = 1, 2$ give (85.6). $\square$

This recurrence is rational and requires no zero computation. Computing $S_1, \dots, S_r$ costs $O(r^2)$ rational arithmetic operations; this is an arithmetic-operation count, not a bit-complexity claim.

Lemma 85.2 (When the concentration index vanishes). *One has*

$$\max \left\{ \frac{1}{N}, \frac{1}{v+1} \right\} \leq \beta \leq \frac{1}{v+1} + \frac{4}{3N}. \quad (85.7)$$

Consequently, $\beta \rightarrow 0$ if and only if $N \rightarrow \infty$ and $m \rightarrow \infty$. If both tend to infinity, then

$$\beta \sim \frac{1}{N} + \frac{1}{m+1/2} - \frac{1}{N+m+1}. \quad (85.8)$$

If $m/N \rightarrow \rho \in [0, \infty]$, then

$$\theta_{N,m} \rightarrow \theta(\rho) = \frac{1+\rho}{1+\rho+\rho^2}, \quad (85.9)$$

with $\theta(0) = 1$ and $\theta(\infty) = 0$.

Proof. Cauchy–Schwarz and $S_1 = 1$ give $S_2 \geq 1/N$. The exact formula for β gives the other lower bound. Since $c < 1/N$ and $(v+3/2)/(v+1) \leq 4/3$ for $v \geq 1/2$, the upper bound follows. For (85.8), put $b_* = 1/N + 1/v - 1/B$. Direct algebra gives

$$\beta = \frac{vb_* + c}{v+1}, \quad b_* - c = \frac{1}{v} + \frac{1}{2NB}.$$

Thus $(b_* - \beta)/b_* \rightarrow 0$. Finally $\theta_{N,m} = [1 + v(v+3/2)/(NB)]^{-1}$, which gives the asserted limits. $\square$

86 A uniform bound on the largest inverse zero

Control of S_2 alone is not sufficient for the sharp critical limits: one weight of order $\sqrt{\beta}$ could survive on the first critical scale. The next theorem excludes that possibility uniformly, even when m/N has no limit.

Theorem 86.1 (Uniform inverse-zero majorant). *For every integer $r \geq 1$,*

$$S_r \leq C_{r-1}(2\beta)^{r-1}. \quad (86.1)$$

Consequently,

$$W \leq \min\{1, 8\beta\}. \quad (86.2)$$

Proof. Because $c < 1/N \leq \beta$, $1/(v+k) \leq 1/(v+1) \leq \beta$, and

$$\frac{c(k+v+1/2)}{v+k} \leq 2\beta,$$

we may discard the negative last term of (85.5) and obtain

$$S_{k+1} \leq \beta \sum_{i=1}^k S_i S_{k+1-i} + 2\beta S_k. \quad (86.3)$$

The bounds for $r = 1, 2$ follow from $S_1 = 1$ and $S_2 = \beta$. Suppose the bounds hold through $r = k$, where $k \geq 2$. Using the Catalan convolution gives

$$\sum_{i=1}^k S_i S_{k+1-i} \leq (2\beta)^{k-1} \sum_{i=1}^k C_{i-1} C_{k-i} = (2\beta)^{k-1} C_k.$$

Also $C_k \geq 2C_{k-1}$ for $k \geq 2$. Equation (86.3) is therefore at most $C_k(2\beta)^k$, completing the induction. Finally $W^r \leq S_r$ and $C_{r-1}^{1/r} \rightarrow 4$. Taking r th roots and passing to the limit proves $W \leq 8\beta$. $\square$

The constant eight is a transparent bound, not a claim of optimality. The limiting measure below suggests the much more precise edge $W/\beta \rightarrow 1 + 3\theta$, but weak convergence alone does not exclude outliers. We leave that strengthening explicitly open here.

87 The first critical scale and exact error control

Theorem 87.1 (Finite-parameter logarithmic bounds). *For every $\tau \geq 0$,*

$$\tau - \frac{\beta\tau^2}{2} \leq \log \mathcal{R}(\tau) \leq \tau - \frac{\beta\tau^2}{2(1+W\tau)}, \quad (87.1)$$

$$0 \leq \log \mathcal{R}(\tau) - \tau + \frac{\beta\tau^2}{2} \leq \frac{S_3\tau^3}{3} \leq \frac{8}{3}\beta^2\tau^3. \quad (87.2)$$

For $8\beta|z| < 1$, with the logarithm normalized by $\log \mathcal{R}(0) = 0$,

$$\left| \log \mathcal{R}(z) - z + \frac{\beta z^2}{2} \right| \leq \frac{8\beta^2|z|^3}{3(1-8\beta|z|)}. \quad (87.3)$$

Proof. For $x \geq 0$,

$$\frac{x^2}{2(1+x)} \leq x - \log(1+x) \leq \frac{x^2}{2},$$

by integrating $u/(1+u)$ from zero to x . Also $0 \leq \log(1+x) - x + x^2/2 \leq x^3/3$. Sum these inequalities at $x = \tau w_j$, using $S_2 = \beta$ and $S_3 \leq WS_2 \leq 8\beta^2$. For complex z , expand every $\log(1 + zw_j)$ in its convergent Taylor series. Since $S_r \leq \beta W^{r-2}$ for $r \geq 2$, the tail from $r = 3$ is bounded by a geometric series. The disk is zero-free because $|z|W < 1$. $\square$

Corollary 87.2 (Sharp Gaussian relative correction). *As $N, m \rightarrow \infty$,*

$$e^{-z/\sqrt{\beta}} \mathcal{R}(z/\sqrt{\beta}) \rightarrow e^{-z^2/2} \quad (87.4)$$

locally uniformly for $z \in \mathbb{C}$. On the nonnegative real axis, $\mathcal{R}(\tau)/e^\tau \rightarrow 1$ exactly when $\tau\sqrt{\beta} \rightarrow 0$, and it tends to zero when $\tau\sqrt{\beta} \rightarrow \infty$.

Proof. For bounded z , the right side of (87.3) after scaling is $O(\sqrt{\beta})$, proving (87.4). The ratio $e^{-\tau} \mathcal{R}(\tau)$ decreases with $\tau \geq 0$, since its logarithmic derivative is

$$-\sum_j \frac{\tau w_j^2}{1 + \tau w_j} < 0 \quad (\tau > 0).$$

The bounded-scale limit and monotonicity show necessity of the little- o condition and give the limit zero beyond every fixed critical multiple. Sufficiency also follows directly from (87.1). $\square$

88 A size-biased inverse-zero law

Definition 88.1. The normalized inverse-zero measure is

$$\nu_{N,m} = \sum_{j=1}^N w_j \delta_{w_j/\beta}. \quad (88.1)$$

It is a probability measure, because $S_1 = 1$.

The weighting by w_j is essential: this is not the uniform empirical measure $N^{-1} \sum \delta_{w_j/\beta}$. Its $(r-1)$ st moment is S_r/β^{r-1} , so the exact recurrence directly determines its limit.

Let Y have the Marchenko–Pastur law of parameter one, defined here by

$$\mathbb{P}(Y \in dy) = \frac{1}{2\pi} \sqrt{\frac{4-y}{y}} \mathbf{1}_{(0,4)}(y) dy. \quad (88.2)$$




Figure 1: The first critical scale, with $N = m$. The horizontal variable is $s = \tau\sqrt{\beta}$. The dashed curve is the proved limit, not a fitted curve. Finite-size curves are floating-point diagnostics.

This is numerically the Catalan moment measure (84.1); the name only identifies the standard law. Its moments are $\mathbb{E}Y^j = C_j$ by the same beta-integral computation. For $0 \leq \theta \leq 1$, put

$$\nu_\theta = \mathcal{L}(1 - \theta + \theta Y). \quad (88.3)$$

Thus $\nu_0 = \delta_1$, while $\nu_1 = \text{MP}_1$.

Theorem 88.2 (The shifted Catalan law). *If $N, m \rightarrow \infty$ and $\theta_{N,m} \rightarrow \theta$, then*

$$\nu_{N,m} \Rightarrow \nu_\theta. \quad (88.4)$$

More explicitly, for each fixed integer $r \geq 1$,

$$\frac{S_r}{\beta^{r-1}} \longrightarrow d_r(\theta), \quad d_r(\theta) = \sum_{j=0}^{r-1} \binom{r-1}{j} C_j \theta^j (1-\theta)^{r-1-j}. \quad (88.5)$$

All measures $\nu_{N,m}$ are supported in $[0, 8]$.

Proof. Write $D_r = S_r/\beta^{r-1}$. The majorant theorem bounds every fixed D_r uniformly. From (83.5),

$$\frac{1}{\nu\beta} \rightarrow \theta, \quad \frac{c}{\beta} \rightarrow 1 - \theta. \quad (88.6)$$

Indeed the first differs from $\theta_{N,m}$ by a factor $(\nu+1)/\nu$, and the second differs from $1 - \theta_{N,m}$ by a factor $(\nu+1)/(\nu+3/2)$. Divide (85.5) by $(\nu+k)\beta^k$. Induction on k yields limits satisfying, with $\gamma = 1 - \theta$,

$$d_{k+1} = \theta \sum_{i=1}^k d_i d_{k+1-i} + \gamma d_k - \theta \gamma \sum_{i=1}^{k-1} d_i d_{k-i}, \quad d_1 = 1. \quad (88.7)$$

This induction works unchanged at $\theta = 0$ and $\theta = 1$.

Let $M(z) = \sum_{k \geq 1} d_k z^{k-1}$. Multiplication of (88.7) by z^k and summation gives

$$M(z) = \frac{1}{1 - \gamma z} + \theta z M(z)^2. \quad (88.8)$$

If $C(z) = \sum C_j z^j$ is the Catalan generating function, the unique formal solution with constant term one is

$$M(z) = \frac{1}{1 - \gamma z} C\left(\frac{\theta z}{1 - \gamma z}\right). \quad (88.9)$$

Expanding the denominator gives exactly (88.5); equivalently, these are the moments of $\gamma + \theta Y$. Finally $W/\beta \leq 8$ makes the measures uniformly compactly supported. Polynomials approximate continuous functions on $[0, 8]$, so convergence of all moments implies the asserted weak convergence. $\square$

The first coefficients, displayed without a change of normalization, are

$$\begin{aligned} d_1 &= 1, & d_2 &= 1, & d_3 &= 1 + \theta^2, & (88.10) \\ d_4 &= 1 + 3\theta^2 + \theta^3, & d_5 &= 1 + 6\theta^2 + 4\theta^3 + 3\theta^4. \end{aligned}$$

This identifies the same quadratic index as universal at second order, while the aspect ratio first becomes visible in the cubic correction.

Remark 88.3 (What weak convergence does not say). For $\theta > 0$, the support of ν_θ is $[1 - \theta, 1 + 3\theta]$. The theorem gives convergence of weighted mass and of fixed moments. It does not prove convergence of the largest atom's location, a rate uniform in r , or an asymptotic for an individual Jacobi zero. Those are stronger statements.

89 An all-orders hierarchy of critical scales

The product representation makes the relation with asymptotic series especially transparent:

$$\log \mathcal{R}(z) = \sum_{r \geq 1} \frac{(-1)^{r+1}}{r} S_r z^r, \quad |z| < 1/W. \quad (89.1)$$

The coefficient S_r is of order β^{r-1} for fixed r . Consequently, its first order-one contribution occurs at $z \asymp \beta^{-(r-1)/r}$.

Theorem 89.1 (Every fixed correction level). *Assume the hypotheses of Theorem 88.2. Fix an integer $q \geq 2$ and put $z_\beta = u\beta^{-(q-1)/q}$. Then*

$$\mathcal{R}(z_\beta) \exp\left\{-\sum_{r=1}^{q-1} \frac{(-1)^{r+1}}{r} S_r z_\beta^r\right\} \longrightarrow \exp\left\{\frac{(-1)^{q+1}}{q} d_q(\theta) u^q\right\} \quad (89.2)$$

locally uniformly for $u \in \mathbb{C}$.

Proof. For $W|z| < 1$, the tail beyond degree q satisfies

$$\left| \log \mathcal{R}(z) - \sum_{r=1}^q \frac{(-1)^{r+1}}{r} S_r z^r \right| \leq \frac{\beta W^{q-1} |z|^{q+1}}{(q+1)(1 - W|z|)}. \quad (89.3)$$

Use $S_r \leq \beta W^{r-2}$ and sum the geometric majorant. At $z = z_\beta$, $W|z| = O(\beta^{1/q})$ and the bound is $O(\beta^{1/q})$ on compact u -sets. The remaining degree- q term converges by (88.5). Exponentiation proves the statement. $\square$

For $q = 2$ this is the Gaussian correction. For $q = 3$,

$$\mathcal{R}(u\beta^{-2/3}) \exp\{-u\beta^{-2/3} + u^2\beta^{-1/3}/2\} \longrightarrow \exp\{(1 + \theta^2)u^3/3\}. \quad (89.4)$$

For $q = 4$, the limiting fourth term is $\exp\{-(1 + 3\theta^2 + \theta^3)u^4/4\}$ after subtracting the first three *exact* terms.

Remark 89.2 (A necessary precision rule). The lower-order S_r in (89.2) cannot in general be replaced by $\beta^{r-1}d_r(\theta)$ without an additional rate hypothesis. Those lower-order contributions diverge on the q th scale. An $o(1)$ relative error in a coefficient can then become an order-one or unbounded error in the exponent. The theorem deliberately keeps the finite-parameter moments exact. It asserts fixed- q limits, not a result with $q \rightarrow \infty$.

90 Resumming the hierarchy: the full nonlinear scale

Define for $t \geq 0$

$$\mathcal{F}_\theta(t) = \int \frac{\log(1 + tx)}{x} d\nu_\theta(x), \quad (90.1)$$

where the integrand at $x = 0$ is defined as t .

Theorem 90.1 (Uniform nonlinear logarithmic limit). *Under the hypotheses of Theorem 88.2,*

$$\beta \log \mathcal{R}(t/\beta) \longrightarrow \mathcal{F}_\theta(t) \quad (90.2)$$

uniformly on compact subsets of $[0, \infty)$, together with every fixed number of derivatives in t . Its derivative is

$$L_\theta(t) = \frac{2}{(1 + (1 - \theta)t) \left(1 + \sqrt{1 + \frac{4\theta t}{1 + (1 - \theta)t}}\right)}. \quad (90.3)$$

The formula includes $t = 0$ and $\theta = 0$ without singularities.

Proof. The exact finite-parameter identity is

$$\beta \log \mathcal{R}(t/\beta) = \int \frac{\log(1 + tx)}{x} d\nu_{N,m}(x). \quad (90.4)$$

On $[0, 8]$ the integrand, continued at $x = 0$, and all its t -derivatives are continuous and uniformly bounded for t in a fixed nonnegative compact interval. Weak convergence and compactness give the claimed uniform limits, including derivatives.

Differentiating (90.1) gives $L_\theta(t) = \int (1 + tx)^{-1} d\nu_\theta(x) = M(-t)$ near zero. Equation (88.8) therefore becomes (83.8). Its positive solution is (90.3); the identity extends to all $t \geq 0$, either by the integral representation or by analytic continuation along that ray. $\square$

At the two extreme aspect ratios the integral is elementary:

$$\mathcal{F}_0(t) = \log(1 + t), \quad (90.5)$$

$$\mathcal{F}_1(t) = \sqrt{1 + 4t} - 1 - \log\left(\frac{1 + \sqrt{1 + 4t}}{2}\right). \quad (90.6)$$

The Taylor series begins

$$\mathcal{F}_\theta(t) = t - \frac{t^2}{2} + \frac{1 + \theta^2}{3}t^3 - \frac{1 + 3\theta^2 + \theta^3}{4}t^4 + \dots. \quad (90.7)$$

It is convergent at least for $|t| < (1 + 3\theta)^{-1}$, with the convention that the radius at $\theta = 0$ is one. Thus the hierarchy is not merely a list of unrelated corrections: its limiting coefficients belong to one analytic function.




Figure 2: The limiting logarithm at the full scale. The middle curve has $m/N \rightarrow 1$, hence $\theta = 2/3$. The endpoints correspond to $m/N \rightarrow \infty$ and $m/N \rightarrow 0$, always with both $m, N \rightarrow \infty$. This is a smooth crossover, not a claim of a singular phase transition at finite positive t .

90.1 Mean and variance on the nonlinear scale

Let K_τ be the coefficient distribution from Theorem 83.1. Its probability generating function is

$$\mathbb{E}_z^{K_\tau} = \frac{\mathcal{R}(\tau z)}{\mathcal{R}(\tau)} = \prod_{j=1}^N (1 - p_j + p_j z), \quad p_j = \frac{\tau w_j}{1 + \tau w_j}. \quad (90.8)$$

Hence K_τ is a sum of independent Bernoulli variables with these parameters. In particular,

$$\mathbb{E} K_\tau = \tau \frac{\mathcal{R}'(\tau)}{\mathcal{R}(\tau)}, \quad \text{Var } K_\tau = \tau \frac{d}{d\tau} \mathbb{E} K_\tau. \quad (90.9)$$

Corollary 90.2 (Nonlinear coefficient fluctuations). *For fixed $t > 0$, under the hypotheses of Theorem 90.1,*

$$\beta \mathbb{E} K_{t/\beta} \rightarrow Q_\theta(t) := t L_\theta(t), \quad (90.10)$$

$$\beta \text{Var } K_{t/\beta} \rightarrow V_\theta(t) := t \int \frac{1}{(1 + tx)^2} d\nu_\theta(x) > 0. \quad (90.11)$$

Moreover

$$\frac{K_{t/\beta} - \mathbb{E} K_{t/\beta}}{\sqrt{\text{Var } K_{t/\beta}}} \Rightarrow N(0, 1). \quad (90.12)$$

Proof. The limits follow from (90.4) by differentiation or directly from the Bernoulli sums. The limiting variance is positive; thus the unscaled variance diverges. For independent centered Bernoulli summands, the sum of third absolute moments is at most their total variance. After division by the variance to the power $3/2$, this tends to zero. The usual characteristic-function proof of the Lyapunov central limit theorem applies: the quadratic term tends to $-u^2/2$ and the total third-order remainder tends to zero. $\square$

Only exact centering is used here. Replacing the mean by $Q_\theta(t)/\beta$ requires an error estimate stronger than the weak convergence in Theorem 88.2.

91 The second critical scale: displacement of the nominal Poisson law

Write $\lambda = \mathbb{E}K_\tau$. From (90.8),

$$\lambda = \tau \sum_j \frac{w_j}{1 + \tau w_j}, \quad \delta := \tau - \lambda = \tau^2 \sum_j \frac{w_j^2}{1 + \tau w_j}. \quad (91.1)$$

Thus

$$\frac{\beta\tau^2}{1 + W\tau} \leq \delta \leq \beta\tau^2, \quad \lambda \geq \frac{\tau}{1 + W\tau}. \quad (91.2)$$

The shift δ becomes visible against a Poisson standard deviation $\sqrt{\tau}$ when $\beta\tau^{3/2}$ becomes order one.

We use the standard Bernoulli-sum Poisson bound [15]:

$$d_{TV} \left(\mathcal{L} \left(\sum_j \text{Bern}(p_j) \right), \text{Pois}(\lambda) \right) \leq \frac{1 - e^{-\lambda}}{\lambda} \sum_j p_j^2, \quad \lambda = \sum_j p_j. \quad (91.3)$$

The convention for d_{TV} is one half the ℓ^1 distance. This external classical inequality is the only Poisson approximation estimate used below; its provenance is not absorbed into the new claims.

Theorem 91.1 (Quantitative corrected-mean approximation). *For $\tau > 0$,*

$$d_{TV}(\mathcal{L}(K_\tau), \text{Pois}(\lambda)) \leq \beta\tau(1 + 8\beta\tau). \quad (91.4)$$

Also

$$d_{TV}(\mathcal{L}(K_\tau), \text{Pois}(\tau)) \leq \beta\tau(1 + 8\beta\tau) + \beta\tau^{3/2}. \quad (91.5)$$

Bounds exceeding one may of course be replaced by one.

Proof. One has $\sum p_j^2 \leq \tau^2\beta$. Apply (91.3), then $\lambda \geq \tau/(1 + W\tau)$ and $W \leq 8\beta$. For two Poisson laws, their Hellinger affinity equals $\exp[-(\sqrt{\lambda} - \sqrt{\tau})^2/2]$. Cauchy–Schwarz gives

$$d_{TV}(\text{Pois}(\lambda), \text{Pois}(\tau)) \leq \sqrt{1 - e^{-(\sqrt{\lambda} - \sqrt{\tau})^2}} \leq |\sqrt{\lambda} - \sqrt{\tau}| \leq \frac{\delta}{\sqrt{\tau}}.$$

Use $\delta \leq \beta\tau^2$ and the triangle inequality. □

Theorem 91.2 (Exact transition profile). *Let $N, m \rightarrow \infty$, $\tau \geq 0$, and $\beta\tau \rightarrow 0$. If $\beta\tau^{3/2} \rightarrow \sigma \in [0, \infty]$, then (83.6) holds. If additionally $\tau \rightarrow \infty$ and $\sigma < \infty$, then*

$$\frac{K_\tau - \tau}{\sqrt{\tau}} \Rightarrow N(-\sigma, 1). \quad (91.6)$$

Proof. By (91.4), K_τ is asymptotically indistinguishable in total variation from $\text{Pois}(\lambda)$. Equations (91.2) give $\lambda/\tau \rightarrow 1$ and $\delta/\sqrt{\tau} \sim \beta\tau^{3/2}$ whenever the latter is nonzero. If $\sigma = 0$, (91.5) already proves the result.

For $0 < \sigma < \infty$, necessarily $\tau \rightarrow \infty$. The likelihood ratio of $\text{Pois}(\lambda)$ to $\text{Pois}(\tau)$ is $e^\delta (\lambda/\tau)^k$, decreasing in k . Its crossing point is

$$k_* = \frac{\delta}{\log(\tau/\lambda)} = \tau - \frac{\delta}{2} + O(\delta^2/\tau).$$

The total variation distance is the difference of the two probabilities of $k \leq \lfloor k_* \rfloor$. The Poisson central limit theorem gives limits $\Phi(\sigma/2)$ and $\Phi(-\sigma/2)$, respectively, proving (83.6). The same theorem, $\lambda/\tau \rightarrow 1$, and $\delta/\sqrt{\tau} \rightarrow \sigma$ give (91.6).

If $\sigma = \infty$, the two means are separated by much more than $\sqrt{\tau}$. At the midpoint of the two means, Chebyshev's inequality shows that the corresponding Poisson probabilities tend to one and zero. Their total variation distance tends to one; the corrected-mean error still tends to zero. This completes every case. □

Remark 91.3 (Do not conflate the two thresholds). At $\tau = s\beta^{-1/2}$ the determinant has a nontrivial factor $e^{-s^2/2}$, yet the nominal Poisson error tends to zero. At $\tau = u\beta^{-2/3}$, that nominal error tends to $2\Phi(u^{3/2}/2) - 1$, but the corrected law $\text{Pois}(\lambda)$ is still accurate. The second threshold is not a failure of all Poisson approximation. Higher-order Bernoulli corrections are part of the broader mod-Poisson literature [16]; the point here is their exact scale and constants for this determinant family.

92 A sharp prefactor in proportional growth

The inverse-zero method controls the scaled logarithm for every aspect ratio. When m/N stays bounded away from zero and infinity, the exact coefficient formula yields a stronger multiplicative asymptotic. This section gives a second route to the nonlinear regime.

Set

$$r = \frac{m + 1/2}{N}, \quad b = \frac{N + m + 1}{N}, \quad \tau = Nt. \quad (92.1)$$

Here $b = 1 + r + 1/(2N)$; retaining this finite- N difference matters for the prefactor. For $0 < x < 1$, define

$$f(x; r, b, t) = x \log(t/x) - (1 - x) \log(1 - x) + (b + x) \log(1 + x/b) - (r + x) \log(1 + x/r), \quad (92.2)$$

continued continuously at the endpoints. Its second derivative is

$$-f''(x) = h(x) := \frac{1}{x} + \frac{1}{1 - x} + \frac{1}{r + x} - \frac{1}{b + x} > 0. \quad (92.3)$$

The derivative tends to $+\infty$ at zero and $-\infty$ at one. There is therefore a unique maximum $q \in (0, 1)$, characterized by

$$tr(1 - q)(b + q) = qb(r + q). \quad (92.4)$$

It is the positive root of a quadratic. A cancellation-resistant formula is

$$q = \frac{2rtb}{\ell + \sqrt{\ell^2 + 4(rt + b)rtb}}, \quad \ell = br - rt(1 - b). \quad (92.5)$$

Finally define

$$\mathcal{A}(r, b, t) = \sqrt{\frac{b(r + q)}{q(1 - q)r(b + q)h(q)}}. \quad (92.6)$$

Theorem 92.1 (Uniform leading prefactor). *Let m/N and t range over fixed compact subsets of $(0, \infty)$. Then, uniformly over these parameters,*

$$\mathcal{R}_{N,m}(Nt) = \mathcal{A}(r, b, t) \exp\{Nf(q; r, b, t)\} (1 + O(N^{-1})). \quad (92.7)$$

For the coefficient distribution at $\tau = Nt$,

$$\frac{K_{Nt} - Nq}{\sqrt{N/h(q)}} \Rightarrow N(0, 1), \quad (92.8)$$

uniformly on such compact parameter sets.

Proof. Let $T_k = A(k)(Nt)^k/k!$, with $x = k/N$. The exact gamma-function expression for T_k is

$$T_k = \frac{(Nt)^k}{k!} \frac{N!}{(N - k)!N^k} \frac{\Gamma(Nb + k)}{\Gamma(Nb)(Nb)^k} \frac{\Gamma(Nr)(Nr)^k}{\Gamma(Nr + k)}. \quad (92.9)$$

For x in any compact subinterval of $(0, 1)$, Stirling's formula, including its square-root factors, gives uniformly

$$T_k = \frac{1}{\sqrt{2\pi N}} \sqrt{\frac{b(r+x)}{x(1-x)r(b+x)}} e^{Nf(x;r,b,t)} (1 + O(N^{-1})). \quad (92.10)$$

The factorial ratio contributes $(1-x)^{-1/2}$; the two rising-factorial ratios contribute $(b/(b+x))^{1/2}$ and $((r+x)/r)^{1/2}$. These factors account for all terms in (92.6).

On the stated compact parameter sets, q stays uniformly inside $(0, 1)$, and $h(q)$ is bounded above and below by positive constants. Strict concavity gives a uniform exponential gap outside any fixed neighborhood of q . Endpoint versions of the rough Stirling bounds, with $O(\log N)$ in the logarithm, show that the sum of the terms outside that neighborhood is exponentially smaller than $e^{Nf(q)}$.

Inside it, put $k = Nq + y\sqrt{N}$ and Taylor-expand f and the smooth amplitude in (92.10). The leading exponential is $e^{-h(q)y^2/2}$. Terms of order $N^{-1/2}$ are odd polynomials in y times that Gaussian, and their integrals vanish. Gaussian tails permit termwise summation after truncation at $|y| \leq N^{1/12}$; the omitted tails are exponentially small in a positive power of N . The Gaussian sums on a mesh of width $N^{-1/2}$ differ from their integrals by exponentially small errors, uniformly in the fractional part of Nq (for example, by the Fourier series of a periodized Gaussian). Thus the relative error is $O(N^{-1})$, not merely $O(N^{-1/2})$.

The leading sum multiplies (92.10) at $x = q$ by $\sqrt{2\pi N/h(q)}$. This is (92.7). Normalizing the same local expansion, with the exponentially small tails, proves (92.8). $\square$

92.1 The limiting saddle and the coefficient rate function

If $m/N \rightarrow \rho \in (0, \infty)$, then $r \rightarrow \rho$, $b \rightarrow 1 + \rho$, and (92.4) simplifies to

$$q(q + \rho) = \frac{\rho(1 + \rho)t}{1 + \rho + \rho t}. \quad (92.11)$$

Writing $f_{\rho,t}(x) = f(x; \rho, 1 + \rho, t)$, we obtain

$$\frac{1}{N} \log \mathcal{R}(Nt) \rightarrow f_{\rho,t}(q). \quad (92.12)$$

For any integers k_N with $k_N/N \rightarrow x \in [0, 1]$,

$$\frac{1}{N} \log \mathbb{P}(K_{Nt} = k_N) \rightarrow f_{\rho,t}(x) - f_{\rho,t}(q). \quad (92.13)$$

The endpoint cases follow directly from (92.9) using the continuous extension of f ; the interior cases use (92.10). Thus the nonnegative function $f_{\rho,t}(q) - f_{\rho,t}(x)$ is the pointwise coefficient rate function.

There is an exact consistency check between the two nonlinear descriptions. Put

$$b_\rho = \frac{1 + \rho + \rho^2}{\rho(1 + \rho)}, \quad \theta = \frac{1 + \rho}{1 + \rho + \rho^2}. \quad (92.14)$$

Then $N\beta \rightarrow b_\rho$, and the two proven limits imply

$$b_\rho f_{\rho,t}(q) = \mathcal{F}_\theta(b_\rho t). \quad (92.15)$$

It can also be checked by differentiating both sides, using (92.11) and (83.8), and matching the value zero at $t = 0$.

93 Boundary regimes and the meaning in the original variables

93.1 What happens when only one parameter grows?

The requirement $N, m \rightarrow \infty$ is genuine, not a convenience. From (84.4), dominated coefficient convergence gives locally uniformly in $\tau \in \mathbb{C}$:

$$\mathcal{R}_{N,m}(\tau) \longrightarrow {}_0F_1(; m + 1/2; (m + 1/2)\tau) \quad \text{if } m \text{ is fixed and } N \rightarrow \infty, \quad (93.1)$$

$$\mathcal{R}_{N,m}(\tau) \longrightarrow (1 + \tau/N)^N \quad \text{if } N \text{ is fixed and } m \rightarrow \infty. \quad (93.2)$$

In the first line, each finite coefficient tends to $v^k/(v)_k$; in the second it tends to N^k/N^k and the series is finite. The bound $|A(k)\tau^k/k!| \leq |\tau|^k/k!$ supplies domination. The fixed- m function is a standard hypergeometric/Bessel limit; no novelty is claimed for that special-function phenomenon.

If $N, m \rightarrow \infty$, the same argument gives $\mathcal{R}(\tau) \rightarrow e^\tau$ locally uniformly. Conversely, convergence to e^τ locally uniformly forces $S_2 = \beta \rightarrow 0$ by comparing second derivatives at zero, hence forces both parameters to diverge. Even convergence at a single fixed positive τ is enough: the upper bound in (87.1), using $W \leq 1$, forces $\beta \rightarrow 0$.

93.2 Physical sizes of the perturbation

Remember $a = \tau/\mu$. For $1 \ll m \ll N$, $\mu \sim N^2/(4m)$ and $\beta \sim 1/m$; for $m \gg N \gg 1$, $\mu \sim N/4$ and $\beta \sim 1/N$. The following table records *orders of magnitude*; constants depend on the aspect ratio in the proportional regime.

Regime	First $O(1)$ response	Relative exponential threshold	Nominal Poisson threshold
$1 \ll m \ll N$	m/N^2	$m^{3/2}/N^2$	$m^{5/3}/N^2$
$m \asymp N \rightarrow \infty$	N^{-1}	$N^{-1/2}$	$N^{-1/3}$
$m \gg N \gg 1$	N^{-1}	$N^{-1/2}$	$N^{-1/3}$

The full nonlinear scale has $a \asymp m^2/N^2$ in the first regime and $a \asymp 1$ in the other two. The usual fixed-multiplicity N^{-2} scale therefore does not remain the correct first-response scale when the multiplicity grows. Corollary 84.2 gives the exact criterion, without relying on this table.

For $b > 0$, factor b from every row:

$$D_N^{(m)}(a, b) = b^N D_N^{(m)}(a/b, 1). \quad (93.3)$$

All conclusions translate by replacing a with a/b . The probabilistic statements require $a/b \geq 0$; complex versions retain only the analytically stated conclusions, not positivity or ordering assertions.

94 Reproducible checks and computational methods

The package includes `code/verify.py`. Its exact layer uses rational arithmetic and its numerical layer uses NumPy and SciPy. The latter does not perform outward rounding and is *not* an interval certificate. The all-parameter results rest on the proofs above, not on the size of the finite test grid.

Executed audit	Count
Direct rational Hankel determinants versus the coefficient formula ($1 \leq N \leq 6$, $0 \leq m \leq 8$, five rational a values)	270
Parameter pairs comparing logarithmic-series extraction with the Riccati recurrence, through moment 12	420
Individual exact moment equalities within those pairs	5,040
Individual exact Catalan-majorant inequalities on the same grid	5,040
Limiting-moment recurrence checks at five rational shapes, through twelve recurrence steps	60
Floating-point Jacobi-root parameter pairs	16

The 5,040 moment equalities are components of the 420 parameter-pair checks, not 5,040 additional independently selected parameter pairs. All assertions passed in the delivered run. The environment and detailed counts are recorded in `data/verification_summary.json`.

[Merge note: in this report the program is shipped as `code/06-growing-shifts-verify.py` and the summary as `data/06-growing-shifts-verification_summary.json`; see Section 81.1.]

For positive τ , stable numerical evaluation uses the term ratio

$$\frac{T_{k+1}}{T_k} = \frac{\tau}{k+1} \frac{(1 - k/N)(1 + k/B)}{1 + k/v}. \quad (94.1)$$

The program accumulates logarithms of this ratio, applies log-sum-exp, and then normalizes the terms. No ill-conditioned Hankel determinant or approximate zero is needed for those tables. The exact moment recurrence is used separately, so its checks are independent of floating-point Jacobi-root finding.

94.1 Two thresholds on a common sequence

For $N = m$, the following values use the exact β in (83.5). The determinant column has $\tau = \beta^{-1/2}$; the last column has $\tau = \beta^{-2/3}$.

$N = m$	β	$e^{-\tau} \mathcal{R}(\tau)$	$d_{\text{TV}}(\mathcal{L}(K_\tau), \text{Pois}(\tau))$
64	$2.3139313 \cdot 10^{-2}$	0.64422016	0.31341650
256	$5.8404126 \cdot 10^{-3}$	0.62704465	0.33303790
1024	$1.4636534 \cdot 10^{-3}$	0.61723707	0.34952143
4096	$3.6613646 \cdot 10^{-4}$	0.61200022	0.36109809
16384	$9.1548078 \cdot 10^{-5}$	0.60929502	0.36871668
Limit	0	$e^{-1/2} = 0.60653066$	$2\Phi(1/2) - 1 = 0.38292492$

The nominal-Poisson convergence is visibly slower. The finite values are not advertised as close enough to establish the limit numerically; Theorem 91.2 proves it independently.

The files `spectral_moments.csv`, `free_energy.csv`, and `saddle_prefactor.csv` give additional checks of the Catalan moment profile, the algebraic free energy, and the multiplicative saddle formula. For the last, the program records both the relative error and N times that error, rather than only a visually agreeable plot.

[Merge note: the table of Section 94.1 is a rounded copy of the columns `beta`, `det_ratio` and `tv` of the shipped file `data/06-growing-shifts-critical_scales.csv`, and the three files named here are shipped under `data/` with the same prefix. All four are floating-point diagnostics.]

95 Proof dependencies and limitations

The dependency structure is short enough to audit explicitly:

Result	Required ingredients
Exact normalized polynomial	Catalan beta integral; one-factor Christoffel identity; Jacobi finite series
Moment recurrence and $W \leq 8\beta$	Hypergeometric ODE; coefficient comparison; Catalan convolution
Inverse-zero limit law	Moment recurrence; compact moment convergence
Critical hierarchy and free energy	Product representation; logarithmic Taylor bounds; inverse-zero law
Nominal-Poisson transition	Bernoulli factorization; classical bound (91.3); elementary Poisson comparison
Proportional-growth prefactor	Exact coefficients; uniform Stirling bounds; one-dimensional discrete Laplace expansion

No unrefereed asymptotic assertion from another repository article is needed as an unproved premise. The already-known coefficient identity is rederived in the needed normalization. The exact numerical tests are consistent with every result, but they do not replace any analytic step.

There are several important boundaries. The general theorems concern one extra factor $x + a$ with a growing fixed-endpoint power x^m . They do not settle moving clusters of growing cardinality, simultaneous arbitrary shifts at both endpoints, or arbitrary polynomial multipliers. The sharp-order and probability results are for $\tau \geq 0$. Negative perturbations can meet zeros, so analogous ratio criteria need not hold. The critical hierarchy fixes its order q before taking limits. The saddle prefactor is uniform only when the aspect ratio and t stay in positive compact sets; it is not asserted uniform as those compact sets approach zero or infinity.

Finally, this is an AI-assisted, unrefereed manuscript. No formal verification in Lean or Rocq has been performed. The repository was read, not modified. The literature audit establishes relevant precedents, not worldwide novelty of each corollary.

96 Further research questions

96.1 A sharp extreme-zero theorem

Does $W/\beta \rightarrow 1 + 3\theta$ hold under exactly the hypotheses of Theorem 88.2, including the extreme ratios $\theta = 0, 1$? An affirmative answer would replace the coarse constant eight by the correct asymptotic edge. A proof needs an outlier-exclusion argument, not just fixed-moment convergence. Quantitative Jacobi zero bounds or recurrence-matrix comparisons are natural routes.

96.2 Finite-size corrections to the inverse-zero measure

The exact Riccati recurrence gives every fixed moment with rational coefficients in N, m . Determine a signed-measure or resolvent expansion around $\nu_{\theta_{N,m}}$, retaining separate N^{-1} and m^{-1} effects. This would produce a multiplicative correction to Theorem 90.1 beyond the proportional-growth case and give rates for exact centering in the nonlinear central limit theorem.

96.3 A uniform bridge from fixed to growing multiplicity

The fixed- m limit is ${}_0F_1(; \nu; \nu\tau)$, whereas growing m gives the hierarchy above. Construct a single two-parameter approximation with a remainder uniform for $1 \leq m \leq N$, not merely an iterated limit. The exact coefficient product suggests a route, but uniform control near its large-parameter transition requires more than dominated convergence.

96.4 Several moving factors and their interactions

For fixed $s \geq 2$, study $H_N(x^m \prod_{j=1}^s (x + a_j))/H_N(x^{m+s})$ as $N, m \rightarrow \infty$. The one-factor theory does not imply a product of one-factor answers: Christoffel determinants can contain mixed corrections. Determine the first interaction term and whether a matrix-valued counterpart of ν_θ governs the joint hierarchy. Then let s grow as well.

96.5 Both endpoint exponents growing

Replace the weight by $x^m(4-x)^\ell d\omega(x)$ with $m, \ell \rightarrow \infty$. Its Jacobi differential equation still provides a rational Riccati recurrence, but the shape space now has more than one parameter. Identify the corresponding normalized inverse-zero law and the sharp small-perturbation scales without assuming $m, \ell \asymp N$.

96.6 Complex perturbations at the first zero edge

The current complex limits stay far inside a shrinking zero-free disk in the original perturbation variable. Near the full negative scale, $1 + tx$ can vanish on the limiting support. Determine the local Airy or Bessel regimes, the zero spacing, and the analytic continuation of the free energy there. Positive-axis estimates cannot simply be continued through a zero.

96.7 Growing truncation order

Theorem 89.1 fixes q . Determine the largest $q = q(N, m)$ for which exact moment subtraction remains uniformly useful, and compare it with direct evaluation of the algebraic limit. This asks for moment asymptotics uniform in the order, a problem not solved by the current fixed-order induction.

96.8 A direct combinatorial coefficient model

The Bernoulli representation uses generally algebraic Jacobi zeros. Find a combinatorial model for the normalized coefficients in which the shape parameter and the mean displacement are visible without those zeros. A positive integer interpretation of the repository's coefficient triangle would be especially useful for this purpose.

96.9 Certified numerical evaluation

Turn (94.1) into an interval algorithm with rigorous finite tail bounds, and combine it with rational moment subtraction. The delivered program is a diagnostic implementation, not that certified algorithm. An efficient certificate should avoid forming large Hankel matrices and should report bit complexity, not only operation counts.

96.10 A staged formalization

A practical Lean project could first formalize the finite hypergeometric polynomial, its ODE as a coefficient identity, and the rational recurrence. The next stage is the positive-root factorization and the Catalan majorant. The logarithmic estimates and the compact moment limit can then be separated from the external Poisson theorem. This is a proposed formalization architecture, not a claim that the necessary code has already been written or checked.

97 Conclusion

The growing-shift question has a useful answer in a variable not visible in fixed- m recurrence formulas: the exact second inverse-zero moment β . It identifies when both size parameters are genuinely large, controls the largest inverse zero uniformly, and sets the critical scales. The corresponding size-biased zero measure converges to a shifted Catalan-moment law, which supplies every fixed-order logarithmic correction and an algebraic resummation at the nonlinear scale.

The main distinction is operational as well as mathematical. A determinant can already differ from its exponential approximation by a fixed factor while its coefficient law is still close to the nominal Poisson law. Later, the nominal mean becomes detectably wrong while a corrected-mean Poisson approximation remains valid. Only at the full scale does the algebraic nonlinear law replace the perturbative description. These statements resolve the specified growing-shift continuation of the repository problem, while leaving the stronger multi-factor, edge, quantitative, and formalization questions explicitly open.

98 A compact formula sheet

This section is Appendix A of the manuscript.

$$\begin{aligned} v &= m + \frac{1}{2}, & B &= N + m + 1, & c &= \frac{v}{NB}, & \mu &= \frac{NB}{4v}, \\ \mathcal{R}(\tau) &= {}_2F_1(-N, B; v; -c\tau), & \beta &= \frac{1 + c(v + 3/2)}{v + 1}, & \theta &= \frac{1}{(v + 1)\beta}. \end{aligned}$$

The moments start with $S_1 = 1$, $S_2 = \beta$ and satisfy (85.5). If $S_r \sim \beta^{r-1} d_r(\theta)$, then

$$d_r(\theta) = \mathbb{E}(1 - \theta + \theta Y)^{r-1}, \quad Y \sim \text{MP}_1.$$

The three central comparisons are

$$\begin{aligned} \tau = s\beta^{-1/2} : & \quad e^{-\tau} \mathcal{R}(\tau) \rightarrow e^{-s^2/2}, \\ \tau = u\beta^{-2/3} : & \quad d_{\text{TV}}(\mathcal{L}(K_\tau), \text{Pois}(\tau)) \rightarrow 2\Phi(u^{3/2}/2) - 1, \\ \tau = t\beta^{-1} : & \quad \beta \log \mathcal{R}(\tau) \rightarrow \mathcal{F}_\theta(t). \end{aligned}$$

For positive real τ , the first finite correction has the explicit error certificate

$$0 \leq \log \mathcal{R}(\tau) - \tau + \frac{1}{2}\beta\tau^2 \leq \frac{8}{3}\beta^2\tau^3.$$

This is a proved analytic bound. Floating-point evaluation of its entries still needs outward rounding before it becomes a numerical interval certificate.

99 Reproduction instructions

This section is Appendix B of the manuscript.

The delivered archive has one top-level directory, `Catalan_Growing_Shifts`. From that directory, run

```
python code/verify.py --output data
python code/make_figures.py
pdflatex -interaction=nonstopmode -halt-on-error article.tex
pdflatex -interaction=nonstopmode -halt-on-error article.tex
```

The exact audit uses only the Python standard library. The full script also imports NumPy and SciPy for diagnostics; figure generation uses Matplotlib. The delivered run used Python 3.13.5, NumPy 2.3.5 and SciPy 1.17.0. These are recorded environment versions, not a requirement to install those exact releases. No external data download is needed.

[Merge note: in this report the delivered files are shipped as listed in Section 81.1, and the commands above must *not* be run in this directory. Here `code/verify.py` is Part I's verifier, and `-output data` would write into the report's shared `data/`; the shipped `code/06-growing-shifts-make_figures.py` does `from verify import ...`, which here would find Part I's verifier and fail, and it writes unprefixes figure files into the `figures/` directory beside its own `code/` directory, here the report's; the delivered Makefile, shipped as `code/06-growing-shifts-Makefile`, runs the same commands and `pdflatex` on this report's `article.tex` in place. The program's CSV writer ends rows with CRLF on every system, which is why the four shipped CSV files are CRLF; its JSON output has CRLF line endings on Windows only. Restore the delivered layout in a scratch copy instead (a POSIX shell, from the report directory):]

```
W=/path/to/scratch/growing-shifts
mkdir -p "$W/code" "$W/recorded" "$W/out"
for f in verify make_figures; do
  cp "code/06-growing-shifts-$f.py" "$W/code/$f.py"; done
for f in data/06-growing-shifts-*; do
  cp "$f" "$W/recorded/${f#data/06-growing-shifts-}"; done
cd "$W" && uv run --no-project --with numpy==2.3.5 --with scipy==1.17.0 \
  python code/verify.py --output out # compare $W/out with $W/recorded
printf 'pdf.fonttype: 42\n' > matplotlibrc # TrueType, not Type 3, fonts
uv run --no-project --with matplotlib==3.10.8 --with numpy==2.3.5 \
  --with scipy==1.17.0 python code/make_figures.py # writes $W/figures
```

[Run this way on 29 September 2026 with Python 3.13.5, NumPy 2.3.5 and SciPy 1.17.0, the program passed every assertion in a few seconds and recorded the counts of Section 94; `free_energy.csv` was byte-identical to the recorded file, the other three CSV files differed only in the last digits of floating-point entries (at most $9.3 \cdot 10^{-10}$, in the cancellation-limited column `n_times_error` of `saddle_prefactor.csv`), and `verification_summary.json` was identical after removal of carriage returns. The figure step reproduced the two shipped figures except for the creation date in their metadata. The manuscript's appendix lists two `pdflatex` runs; its delivered README and Makefile list three.]

A The precise correction to the printed conjecture

This appendix distinguishes two kinds of index. The source uses a size- $(n + 1)$ determinant. We write $N = n + 1$. To avoid silently identifying incompatible shift conventions, write s for the source's symbol in its displayed coefficient formula, and reserve m for our actual Catalan shift.

For $s \geq 2$ and $0 \leq k \leq n + 1$, the coefficient expression printed in Conjecture 2 of [1] can be written as

$$T_{n,k,s} = \frac{C_s}{2s-2} \binom{s+k-2}{s-2} \binom{n+k+2s-2}{2k+2s-3} \times \prod_{j=0}^{\lfloor (s-1)/2 \rfloor - 1} \frac{\binom{2n+2s-2j-1}{2s-4j-5}}{\binom{2s-2j-1}{2s-4j-5}} \prod_{j=0}^{s-3} \frac{2s-j-2}{n+k+2s-j-2}. \quad (\text{A.1})$$

Only the natural, nonambiguous range $s \geq 2$ is needed here. Empty products have value 1.

Proposition A.1 (Correct shift). *For $N \geq 1$, $s \geq 2$, and $0 \leq k \leq N$,*

$$T_{N-1,k,s} = c_{N,s-1,k}. \quad (\text{A.2})$$

Consequently, the polynomial using $T_{N-1,k,s}$ is $D_N^{(s-1)}(a, b)$, not $D_N^{(s)}(a, b)$. Equivalently, to use that printed expression for an actual shift $m \geq 1$, replace its shift symbol by $s = m + 1$.

Proof. Taking a ratio of successive terms in (A.1) and cancelling factorials gives

$$\frac{T_{n,k+1,s}}{T_{n,k,s}} = \frac{(n+1-k)(n+k+s+1)}{2(k+1)(2k+2s-1)}. \quad (\text{A.3})$$

This agrees with (2.3) after $N = n + 1$ and $m = s - 1$. It remains to match the initial coefficient. Putting $n = N - 1$, $k = 0$, and simplifying the factorials gives

$$T_{N-1,0,s} = C_s \frac{(N+s-1)!}{N!s!} \prod_{j=0}^{p-1} \frac{\binom{2N+2s-2j-3}{2s-4j-5}}{\binom{2s-2j-1}{2s-4j-5}}, \quad p = \left\lfloor \frac{s-1}{2} \right\rfloor. \quad (\text{A.4})$$

At $N = 1$ this is $C_s = H_1^{(s)}$. Its ratio at successive N is

$$\begin{aligned} \frac{T_{N,0,s}}{T_{N-1,0,s}} &= \frac{N+s}{N+1} \prod_{j=0}^{p-1} \frac{(2N+2s-2j-1)(2N+2s-2j-2)}{(2N+2j+4)(2N+2j+3)} \\ &= \frac{(2N+2s)!(2N+1)!}{(2N+s)!(2N+s+1)!}. \end{aligned} \quad (\text{A.5})$$

In the product, the numerator factors form a consecutive interval of integers, as do the denominator factors; the last equality holds for either parity of s . The shifted-product formula gives the identical ratio

$$\frac{H_{N+1}^{(s)}}{H_N^{(s)}} = \prod_{i=1}^{s-1} \frac{(2N+2i+1)(2N+2i+2)}{(2N+i+1)(2N+i+2)} = \frac{(2N+2s)!(2N+1)!}{(2N+s)!(2N+s+1)!}.$$

Induction in N therefore proves $T_{N-1,0,s} = H_N^{(s)}$. The ratio (A.3) then proves (A.2) for every k . $\square$

The discrepancy is visible in the smallest nontrivial instance. With $N = 1$ and $s = 2$, the printed expression is

$$T_{0,0,2}b + T_{0,1,2}a = 2b + a,$$

whereas the determinant with the stated shift $s = 2$ is

$$D_1^{(2)}(a, b) = aC_2 + bC_3 = 2a + 5b.$$

At $a = b = 1$, these are 3 and 7. The expression $a + 2b$ is exactly $D_1^{(1)}(a, b)$, so the correction is systematic rather than an isolated numerical exception. The original PDF page was inspected as well as its HTML rendering; this is not an artifact of an HTML conversion. The denominator-exponent statement is separate and uses the actual shift m correctly.

B Compact reference and archive conventions

Symbol	Meaning
N	Determinant size, with $D_0 = 1$
m	Actual Catalan shift in $aC_{i+j+m} + bC_{i+j+m+1}$
$H_N^{(m)}$	Pure shifted Catalan determinant
d, e, K	$\binom{m}{2}$, $\binom{m}{2} + 1$, and $(m - 1)^2$
P_m	Uniform numerator, not automatically reduced at exceptions
q	$1 - (a + 2b)t + b^2t^2$
$M_{m,\ell}$	Minor with column ℓ deleted from the binomial block
$Q_m(N; r)$	Signed generating polynomial of those minors
$r_{\pm}$	Roots of $r^2 - (a/b + 2)r + 1$
$\lambda_{\pm}$	$br_{\pm}$, the characteristic roots in determinant size
s in Appendix A	Source shift symbol, equal to actual shift plus one

[Added 28 September 2026, batch 39: this table covers Part I. Part II reuses several of these letters with other meanings, among them q , s , ℓ , d , K , P , M and r ; its symbols are listed in Table 1, and its data files in Section 27.]

The JSON files use ascending polynomial powers. In `coefficient_rows.json`, position k is the coefficient of $a^k b^{N-k}$. In `minor_polynomials.json`, position j is the coefficient of N^j . In `two_branch_polynomials.json`, the arrays `P_plus` and `P_minus` describe the polynomial multipliers of 4^N and 1^N for $(a, b) = (1, 2)$; they are not the ordinary-generating-function numerator P_m .

The source audit records exact URLs, source versions, and search limitations. The tests establish reproducibility of the computation, not priority or formal verification. The archive intentionally contains no third-party article PDFs, no font files, no compiler auxiliaries, and no Python bytecode.

References

- [1] P. Barry, *Notes on the Hankel transform of linear combinations of consecutive pairs of Catalan numbers*, arXiv:2011.10827v1 (21 November 2020), 29 pp. The denominator conjecture and Conjecture 2 are on p. 4. <https://arxiv.org/abs/2011.10827>.
- [2] M. Dougherty, C. French, B. Saderholm, and W. Qian, Hankel transforms of linear combinations of Catalan numbers, *Journal of Integer Sequences* **14** (2011), Article 11.5.1. <https://cs.uwaterloo.ca/journals/JIS/VOL14/French/french2.html>.
- [3] C. Krattenthaler, Hankel determinants of linear combinations of moments of orthogonal polynomials, II, *The Ramanujan Journal* **61** (2023), 597–627; published online 22 November 2021. <https://doi.org/10.1007/s11139-021-00514-8>. See the determinant formula, the confluence discussion, and Section 8 for existing general recurrence theory and its earlier sources.
- [4] NIST Digital Library of Mathematical Functions, Section 18.5, Eq. 18.5.7, Jacobi polynomial hypergeometric representation. <https://dlmf.nist.gov/18.5.E7>.
- [5] OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, A000108 (Catalan numbers). <https://oeis.org/A000108>.
- [6] OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, A001519 and A001906 (Fibonacci subsequences). <https://oeis.org/A001519>; <https://oeis.org/A001906>.
- [7] J. Cigler and C. Krattenthaler, Hankel determinants of linear combinations of moments of orthogonal polynomials, *International Journal of Number Theory* **17** (2021), 341–369. Preprint arXiv:2003.01676. <https://arxiv.org/abs/2003.01676>. Cited in Part II.
- [8] C. Krattenthaler, A determinant identity for moments of orthogonal polynomials that implies Uvarov’s formula for the orthogonal polynomials of rationally related densities, preprint arXiv:2103.03969. <https://arxiv.org/abs/2103.03969>. Cited in Part II as a starting point for the rational-multiplier research direction, not as an input to its proofs.
- [9] J. Cigler, Hankel determinants of middle binomial coefficients and conjectures for some polynomial extensions and modifications, preprint arXiv:2111.14492 (2021). <https://arxiv.org/abs/2111.14492>. Cited in Part III: its Theorem 1, equation (23), supplies the known central product after (36.3); its proof also uses condensation. Part III’s manuscript does not state which arXiv version it read.
- [10] E. Strahov and Y. V. Fyodorov, Universal results for correlations of characteristic polynomials: Riemann–Hilbert approach, *Communications in Mathematical Physics* **241** (2003), 343–382. <https://arxiv.org/abs/math-ph/0210010>. Cited in Parts IV and V for context, not as an input to their proofs.
- [11] G. Akemann and Y. V. Fyodorov, Universal random matrix correlations of ratios of characteristic polynomials at the spectral edges, *Nuclear Physics B* **664** (2003), 457–476. <https://arxiv.org/abs/hep-th/0304095>. Cited in Part IV for hard-edge context, not as an input to its proofs.
- [12] NIST Digital Library of Mathematical Functions, Section 18.11(ii), Formulas of Mehler–Heine type. <https://dlmf.nist.gov/18.11>. Cited in Part IV; consulted by its manuscript on 29 September 2026.
- [13] A. F. Celsus, A. Deaño, D. Huybrechs, and A. Iserles, The kissing polynomials and their Hankel determinants, *Transactions of Mathematics and Its Applications* **6** (2022), article tnab005. <https://doi.org/10.1093/imatrm/tnab005>; preprint arXiv:1504.07297, <https://arxiv.org/abs/1504.07297>. Cited in Part V: the oscillatory Jacobi determinant, its parity phenomena and its large-frequency analysis are prior literature, not novelty claims there.

- [14] H. Dette and W. J. Studden, *Some new asymptotic properties for the zeros of Jacobi, Laguerre and Hermite polynomials*, arXiv:math/9406224 (1994). <https://arxiv.org/abs/math/9406224>. Cited in Part VI for the established large-parameter zero-asymptotic context; no unverified specialized theorem from this paper is invoked.
- [15] A. D. Barbour and P. Hall, *On the rate of Poisson convergence*, Mathematical Proceedings of the Cambridge Philosophical Society **95** (1984), 473–480. <https://doi.org/10.1017/S0305004100061806>. Cited in Part VI for the Bernoulli-sum Poisson bound (91.3).
- [16] P.-L. Méliot, A. Nikeghbali, and G. Visentin, *Mod-Poisson approximation schemes and higher-order Chen–Stein inequalities*, arXiv:2210.13818 (2022). <https://arxiv.org/abs/2210.13818>. Cited in Part VI as prior context for higher-order Bernoulli-sum approximation, not as a novelty claim for that general mechanism.

Online sources checked on 19 September 2026 (Part I) and 28 September 2026 (Part II); Part III's manuscript is dated 29 September 2026, and its source notes do not date its checks; Part IV's manuscript and its source notes are dated 29 September 2026, the day it consulted DLMF; Part V's and Part VI's manuscripts and source audits are dated 29 September 2026 (both audits give the Pacific date). All theorem numbering in the body refers to the present manuscript, unless a source is explicitly named.