

Birthday Cutoffs and Hereditary Sets

The set-theoretic content of bounded surreal arithmetic:
bi-interpretations, elementary inclusions, axiom spectra,
outer models, and surcomplex orientation

September 22, 2026

Abstract

Expand the ordered field of surreal numbers of birthday below an ordinal λ by its birthday function. The global connection of this kind of structure with set theory is the subject of a bi-interpretation theorem announced by Chen, Hamkins and Yang. This report, merged from five manuscripts that prove overlapping versions of one local theorem, asks *which set universe is recovered from a particular birthday cutoff, and what that recovery implies*.

For an epsilon number λ let $\kappa(\lambda)$ be the least cardinal not below λ . One parameter-free interpretation recovers exactly $H_{\kappa(\lambda)}$. At cardinal cutoffs, singular ones included, this is a uniform bi-interpretation with H_λ ; at noncardinal cutoffs it is a bi-interpretation with $(H_{\kappa(\lambda)}, \in, \lambda)$, and the cutoff is recovered by a definable collector. Thus the field of birthdays below ε_0 already recovers all hereditarily countable sets. Small arithmetic certificates justify the reverse interpretation without Replacement in H_κ , and a second route codes graphs modulo bisimulation. The forward interpretation extends to the ring cutoffs, including ω .

Consequences within one universe: for epsilon numbers $\lambda < \eta$ the inclusion of birthday structures is elementary exactly when both are cardinals and $H_\lambda \prec H_\eta$; elementary embeddings correspond to those of the hereditary universes; one sentence detects noncardinal cutoffs and another the strong-limit cardinals; birthday eternity is Replacement at cardinal cutoffs, and with fiber packing it selects the worldly cardinals; in ZF the graph coding reaches exactly the sets with well-orderable transitive closure. Across universes: old arithmetic is absolute, a bounded structure is unchanged exactly when no bounded subset is added, the first new birthday is the least ordinal with a new subset and is a ground-model cardinal, explicit formulas detect new subsets, and, assuming a measurable cardinal, Prikry forcing leaves the enriched field unchanged while singularizing its cutoff. Under GBC the full structure has no nontrivial elementary self-embedding. For surcomplex numbers with conjugation and coordinate birthday, each real elementary embedding has exactly two lifts, and one nonreal parameter, but no set of real parameters, defines an orientation.

Two later parts work below ZFC. Over Zermelo set theory without Replacement, bounding the birthdays of definable surreal families is ordinal bounding (with a hierarchy axiom, Replacement), and kernel models with atoms have exact Collection spectra; over ZF, universality of the surreals for free fields on all sets is exactly the first Kinna–Wagner principle.

Built from five manuscripts written independently in relation to the `VladimirReshetnikov/Surreal` project, all pinned to commit `4f26455`, and merged after placement commit `d4e71b7`; Parts VI–VII (batch 89) add two more, pinned to `2b7b388`, placed by `23adb85`.

An AI-assisted research draft with written proofs and finite regression checks; not refereed; not a claim of any Lean formalization. The global bi-interpretation is prior announced work of Chen–Hamkins–Yang.

Contents

I	The cutoff theorem	1
1	The question, the results, and their provenance	1
2	Foundations and the two ordinal languages	9
3	Small certificates for surreal arithmetic	11
4	Definable sign access and a bounded pairing map	14
5	Pointed relation codes and the exact interpreted universe	16
6	Bi-interpretation and recovery of the cutoff	19
7	Second route: graphs modulo bisimulation	22
8	Ring cutoffs and the finite boundary	24
II	Consequences inside one universe	24
9	The exact elementary-inclusion spectrum	25
10	Birthday growth and set-existence axioms	28
11	Worldly cutoffs and separation examples	31
12	Model-theoretic consequences	34
13	The exact Choice boundary of ordinal graph coding	36
III	Outer models	37
14	Old arithmetic and cross-model coherence	37
15	The preservation criterion and the first new birthday	39
16	Detecting new subsets	41
17	Prikry forcing: an unchanged field with a singularized cutoff	43
IV	The full class and the surcomplex numbers	44
18	The full class: the announced theorem and Kunen rigidity	45
19	Surcomplex numbers and the orientation obstruction	47
V	Scope, questions and provenance	50
20	What the theorems do and do not imply	51
21	Open questions, re-scoped	54

22	Provenance and verification	55
23	Conclusion	58
VI	Replacement-free surreal cuts, definable families and atom-support spectra	58
24	Source 10: question, results, notation and relation to Part II	58
25	The base theories and the meaning of a family	63
26	Sign sequences and bounded containers	64
27	Actual set cuts: a Replacement-free simplicity proof	66
28	The equivalence theorem for definable surreal families	67
29	When the test becomes full Replacement	69
30	An explicit omitted cut in a rank-initial universe	69
31	Urelements: rank bounds versus atom-base bounds	70
32	Cardinal-support models and their basic axioms	71
33	The exact Collection spectrum	72
34	Naming an enumeration: exact failure of Replacement	74
35	Why pure surreal structures cannot see the support obstruction	75
36	A formalization route for ProveIt	76
37	Executable finite checks and reproducibility	78
38	Further research questions and proposed projects	79
39	Conclusion of source 10, ledger and provenance	81
VII	Surreal universality without choice	83
40	Source 11: question, results, notation and relation to the report	83
41	Foundations, notation, and the surreal interface	87
42	The exact coding strength of a surreal coordinate set	88
43	Double monomials and free-algebra universality	89
44	An optimal set-sized real-closed target	91
45	Prescribed order is a different universality problem	93
46	A convex-resolution criterion for ordered embeddings	95
47	Orbit factorization of polynomial representations	95

48 Sharp support bounds in two permutation models	96
49 An ordered field with all finite stages embeddable but no global embedding	99
50 Proof audit and relation to the existing repository	101
51 Further research questions and concrete next targets	103
52 Reproducibility and limits of the computational evidence	105
53 Conclusion of source 11, checklist and provenance	105
A Formula expansion and size checks	107
B An essential negative regression	109
C A small concrete membership code	109

Part I

The cutoff theorem

1 The question, the results, and their provenance

The field structure of the surreal numbers and their construction history carry very different information. Real-closed-field quantifier elimination makes all inclusions between real closed subfields elementary in the ordered field language. A birthday function remembers much more: together with the order it reads individual sign coordinates, and thereby encodes subsets, relations and well-founded membership graphs. The issue addressed here is not merely whether this coding is possible globally, but how its strength changes when only birthdays below one ordinal are allowed, and what the resulting dictionary says about elementary inclusions, set-existence axioms, outer models of set theory, and the surcomplex numbers.

1.1 What is already known, and credited

The global theorem is prior work. Chen, Hamkins, and Yang announced that the surreal ordered field with birthday order is bi-interpretable with the set-theoretic universe, and described a first-order theory of surreal arithmetic intended to be bi-interpretable with ZFC. Hamkins’s September and November 2025 slides give the global coding by pointed well-founded extensional relations, together with birthday-growth principles; the March 2026 workshop announcement and slides repeat the global theorem and label the axiomatic work as still being checked [6, 7, 8]. This report does *not* claim that theorem, its pointed-graph coding idea, or the association of birthday growth with set-existence axioms as new. The collection’s **foundations-and-computation/foundations** report already records the March 2026 announcement, at slide level and with its status boundary, in its subsection **found:sub:announcement**, and lists it among its non-claims as work in progress that is not a proved dependency of that report. Of the five manuscripts merged here, sources 02, 06 and 09 credit the announcement prominently. Sources 05 and 07 do not cite it; source 07 presented the full-class bi-interpretation as a proposed new contribution. That presentation is corrected here: [theorem 18.1](#) states the announced theorem with its credit, and the argument printed there, taken from source 07, is a written proof of the announced theorem, not a new result.

Other prior inputs. If λ is an epsilon number, $\mathbf{No}_{<\lambda}$ is a real closed subfield. The original van den Dries–Ehrlich paper must be used together with its erratum [3, 4]; Bournez–Guilmant state the corrected ring, field and real-closedness results in their Theorems 1.1–1.2 [5]. The collection’s foundations report states the same classical fact in its subsection **found:sub:cutoffs**, including that ε_0 is already an epsilon number, so that no large cardinal is needed. The sign construction and recursive arithmetic are classical [1, 2]. Mostowski collapse, well-founded recursion, hereditary-size universes and forcing are standard [13]; quantifier elimination for real closed and algebraically closed fields is standard [14, 15]. The general obstruction to parameter-free bi-interpretation caused by complex conjugation, and the role of a named imaginary unit, were explained by Jeřábek in 2021 [9]; every use of that obstruction below ([section 19](#)) is credited to him. Bi-interpretation methodology follows Freire–Hamkins [10]. The Prikry forcing facts are imported from Poveda–Rinot–Sinapova [11], and the class form of Kunen’s inconsistency from Hamkins–Kirmayer–Perlmutter [12]. Rangel–Mariano’s algebraic set theory of surreal numbers and Bertram’s realization of numbers as sets of

ordinals are separate programs with different languages and aims; nothing here settles or refutes them [16, 17].

1.2 The principal statements

We use $\text{TC}(\{x\})$ for the transitive closure containing x and all its iterated members, and write

$$H_\kappa = \{x : |\text{TC}(\{x\})| < \kappa\}.$$

For an epsilon number λ (an ordinal with $\omega^\lambda = \lambda$ in ordinary ordinal exponentiation) put

$$\mathcal{B}_\lambda = (\mathbf{No}_{<\lambda}; 0, 1, +, -, \cdot, <, \mathbf{b}),$$

where $\mathbf{b}(x)$ is the birthday of x , regarded as the all-plus surreal ordinal of that length. “Uniform” means that fixed finite formulas work at every cutoff in the stated class, not that a separate language is chosen for each cardinal. The source numbers in brackets are explained in [section 1.3](#).

Theorem 1.1 (Cardinal rounding and cutoff recovery; [theorem 6.3](#)). *Let λ be an epsilon number and let*

$$\kappa(\lambda) = \min\{\kappa : \kappa \text{ is a cardinal and } \lambda \leq \kappa\}.$$

There is one parameter-free interpretation $\mathcal{I}$ such that

$$\mathcal{I}(\mathcal{B}_\lambda) \cong (H_{\kappa(\lambda)}, \in).$$

If λ is a cardinal, $\mathcal{B}_\lambda$ and $(H_\lambda, \in)$ are uniformly bi-interpretable. If λ is not a cardinal, $\mathcal{B}_\lambda$ and $(H_{\kappa(\lambda)}, \in, \lambda)$ are uniformly bi-interpretable, and the named ordinal λ is definable, as an interpreted set, in $\mathcal{B}_\lambda$ itself.

This is the most general form in the cluster, and it is source 09’s. The other four sources prove cases of it with their own codes:

- source 02: the forward interpretation onto $H_{\kappa(\lambda)}$ at every epsilon number, one way only (it explicitly claims no reverse interpretation);
- source 05: the bi-interpretation at uncountable *regular* cardinals, with graphs modulo bisimulation ([section 7](#));
- sources 06 and 07: the bi-interpretation at every uncountable cardinal, singular ones included; source 07 also proves the forward interpretation onto $H_{\kappa(\lambda)}$ at every epsilon number, which it calls a *cardinality plateau* and states forward only.

This is a cardinal-rounding statement, not a rank-cutoff statement: for example $\kappa(\varepsilon_0) = \omega_1$, whereas V_{ε_0} is not the recovered universe. At a singular cardinal κ the recovered structure is still H_κ ; no regularity assumption is hidden in the theorem.

Theorem 1.2 (Elementary-inclusion spectrum; [theorem 9.3](#)). *For epsilon numbers $\lambda < \eta$,*

$$\mathcal{B}_\lambda \prec \mathcal{B}_\eta \iff \lambda, \eta \text{ are cardinals and } H_\lambda \prec H_\eta.$$

The ordered-field reducts, in contrast, always form an elementary inclusion.

Source 09 proves the equivalence at all epsilon numbers. Source 02 proves the forward implication and left the converse as its first open question; that question is answered by source 09 ([remark 9.4](#)). Sources 05 (regular cardinals), 06 and 07 (uncountable cardinals) prove the equivalence between cardinal cutoffs, and 05 and 06 add the correspondence of elementary embeddings ([theorem 9.10](#)).

Theorem 1.3 (Growth spectrum; [theorem 11.2](#)). *For an epsilon-number cutoff, birthday eternity Et fails at every noncardinal cutoff. At an uncountable cardinal κ ,*

$$\mathcal{B}_\kappa \models \text{Et} \iff H_\kappa \models \text{Replacement}.$$

Fiber packing Pack, and also the translated Power Set sentence $\text{Pow}^\mathcal{I}$, hold exactly at the uncountable strong-limit cardinal cutoffs. Consequently

$$\mathcal{B}_\lambda \models \text{Et} + \text{Pack} \iff \lambda \text{ is a cardinal and } V_\lambda \models \text{ZFC}.$$

The eternity, packing and worldly statements are source 09's. The translated Power Set sentence is source 02's, at every epsilon number, and source 06's at uncountable cardinals ([theorem 10.7](#)).

Theorem 1.4 (Extension rigidity at a strong-limit cutoff; [theorem 16.4](#)). *Let $M \subseteq N$ be transitive models of ZFC with the same ordinals, and let κ be an uncountable cardinal in both, strong limit in M . Then*

$$\mathcal{B}_\kappa^M \prec \mathcal{B}_\kappa^N \iff \mathcal{B}_\kappa^M = \mathcal{B}_\kappa^N \iff \mathcal{P}^M(\alpha) = \mathcal{P}^N(\alpha) \quad (\alpha < \kappa).$$

Without the strong-limit hypothesis, a new subset of $\alpha < \kappa$ still prevents elementarity whenever $(2^{|\alpha|})^M < \kappa$.

The detector is sharp for the old-power-set *parameter* ([proposition 16.2](#)), not a claim that every other possible detector needs that much space. The report also proves:

- the forward interpretation at ring cutoffs, including $\mathbf{No}_{<\omega}$ onto the hereditarily finite sets ([proposition 8.1](#), source 02);
- one sentence true exactly at noncardinal epsilon cutoffs ([theorem 6.2](#), sources 02 and 09);
- a separation of $\mathcal{B}_{\omega^-}$ from $\mathcal{B}_{\omega^+}$ by one Replacement instance ([theorem 11.5](#), source 06);
- the exact range of the graph coding over ZF without Choice ([theorem 13.2](#), source 06);
- the transfer of elementary equivalence, the interpretation of full second-order arithmetic, the independence property of the bit formula, an omitted parameter-free type, and nondefinability of birthday in the field reduct ([section 12](#), sources 05, 06, 07);
- the exact preservation criterion across outer models, the first new birthday, three explicit new-subset detectors, and the full-class statement $\mathcal{B}^M \prec \mathcal{B}^N \iff M = N$ ([Part III](#), sources 02, 05, 06, 07, 09);
- conditional on a measurable cardinal, an unchanged enriched field whose cutoff changes cofinality, the preserved $H_\kappa = V_\kappa^M$, and the exact onset of new elements above the cutoff ([section 17](#), sources 02 and 06);
- under GBC, no nontrivial class elementary self-embedding of the full birthday structure, by transfer of Kunen's inconsistency ([theorem 18.3](#), source 07);
- for surcomplex numbers with conjugation and coordinate birthday: exactly two elementary lifts of each real elementary embedding (source 06), the exact parameter threshold for a definable orientation (source 06), the mutual-interpretation versus bi-interpretation distinction (sources 05, 06, 09, after Jeřábek), and, under GBC, the elementary self-embeddings of the full structure (source 07) ([section 19](#)).
- in Part VI (source 10, added in batch 89), over Zermelo set theory with Foundation and without Replacement or Choice: simplest separators of set cuts without Replacement, the equivalence of six definable-family schemes with ordinal bounding and, over an explicit hierarchy axiom, with full Replacement and Collection (the weak-base form of [theorem 10.4](#)); the exact Collection spectra of kernel urelement models; and a language-sensitive separation that pure surreal structures cannot see ([theorems 24.1 to 24.3](#));

- in Part VII (source 11, added in batch 89), over ZF: a set injects into **No** iff it injects into the power set of an ordinal; universality of **No** and **Oz** for free algebras on all sets is exactly the first Kinna–Wagner principle; optimal set-sized real-closed targets; sharp transcendence bounds in two permutation models and, in ZFA, an ordered field with no surreal embedding (theorems 43.5, 44.3 and 49.1).

1.3 Five manuscripts, one report

This report is built from five manuscripts, written independently on September 22, 2026, all pinned to repository commit 4f2645599121fa872c7104995e47f86f0382351f and placed in the collection by commit d4e71b7. Their numbers are those of the placement batch and are local to this directory.

Source	Manuscript	Contribution to this report
09	<i>The Set-Theoretic Content of Bounded Surreal Arithmetic: hereditary-size universes, coherent bi-interpretations, axiom spectra, and extension rigidity</i>	The base. Its text is the spine of Part I (sections 2 to 6), the elementary-inclusion classification (section 9), the eternity, packing and worldly calibration (section 10), the cross-model coherence, detector and extension rigidity (section 16), the orientation obstruction (theorem 19.4), and most of section 20. Files prefixed 09-bounded-arithmetic-.
02	<i>What a Birthday Cutoff Remembers: hereditary set universes, elementary obstructions, and forcing invisibility</i>	The one-way forward theorem at all epsilon numbers (a special case of the base), ring cutoffs (section 8), the collector sentence (printed once with 09's), the translated Power Set sentence at every epsilon number, the preservation criterion at epsilon cutoffs and its countable corollary, the Prikry invisibility theorem with its preserved H_κ and the onset above the cutoff (section 17), the concrete example in Appendix C. Files prefixed 02-birthday-cutoffs-, and its source audit 02-birthday-cutoffs-SOURCE_AUDIT.md.
05	<i>Birthdays Recover Sets: bi-interpretability, elementary embeddings, and forcing in surreal and surcomplex fields</i>	The second route by codes modulo bisimulation (section 7), the explicit embedding formula and critical-point corollary, the separation of $\mathcal{B}_{\omega_1}$ and $\mathcal{B}_{\omega_2}$, nondefinability of birthday, the explicit missing-row detector New, the full-class statement $\mathcal{B}^M \prec \mathcal{B}^N \iff M = N$, the first new birthday for arbitrary same-ordinal outer models, the countably closed example, and the surcomplex forcing contrast. Files prefixed 05-birthdays-recover-sets-.

Source	Manuscript	Contribution to this report
06	<i>What Birthday-Enriched Surreal Fields Remember: hereditary-size interpretations, choice, forcing, and surcomplex orientation</i>	The bi-interpretation at singular cardinals (printed once with the base), the second route by prefix-pair arithmetic tables (section 3.4), definitional equivalence with $(\mathbf{No}_{<\kappa}; <, \mathbf{b}, G_\omega)$, transfer of elementary equivalence, the $\sqsupset_\omega$ Replacement instance, the ZF range theorem, the four-way preservation criterion, the Prikry corollaries, the two complex lifts, and the orientation threshold. Files prefixed 06-hereditary-size-orientation-.
07	<i>The Birthday Expansion of the Surreal Field: recovering set theory, detecting forcing, and rigidity of surcomplex embeddings</i>	The written proof of the announced full-class theorem (theorem 18.1), the radix pairing, the ε_0 localization (printed once as a case of the base), recovery of models of ZFC, stage agreement at every ordinal, the two-parameter outer-model witness, the independence property and omitted type, the o-minimal nondefinability, Kunen transfer (theorem 18.3) and the surcomplex elementary self-embeddings. Files prefixed 07-birthday-expansion-.

Why one report. All five prove the same spine: the ordered field of surreals below a cutoff, expanded by birthday, interprets the hereditary sets H_κ by sign bits read from order and birthday, a bounded ordinal pairing polynomial, and pointed well-founded relation codes, and in most versions is bi-interpretable with them. They differ in the cutoff class and in whether the result is one-way or two-way. Source 09 has the widest cutoff class (all epsilon numbers, which include all uncountable cardinals by lemma 2.2) and the strongest conclusion (two-way, with the correct pointed target), so it is the base.

Printed once. The prefix and bit formulas (all five), the bounded pairing (all five, in four variants compared in remark 4.4), the rooted collapse and exact range (02, 06, 07, 09), the arithmetic certificates (05, 06, 07, 09), the bi-interpretation at cardinal cutoffs (05 regular; 06, 07, 09 all), the ε_0 case (02, 07, 09), the collector sentence (02, 09), the elementary-inclusion classification (02 one direction; 05, 06, 07, 09), the translated Power Set threshold (02, 06, with 09's version inside H_κ), absoluteness of old arithmetic (02, 05, 07, 09), the preservation criterion (02, 06, 07, 09), the first new birthday (02, 05, 07), the outer-model non-elementarity (05, 07, 09), the Prikry invisibility (02, 06), rigidity of the bounded structures (05, 06, 07, 09), the surcomplex real-axis recovery (all five) and the two-element automorphism group (05, 06, 07, 09). Each is printed once and credited to every source that proves it.

Kept twice, as different proofs. The reverse arithmetic has two routes: source 09's option-graph certificates built from finite signed words (theorem 3.3), and the prefix-pair tables of sources 05, 06 and 07 (section 3.4), whose single- μ size bound is source 06's argument at singular cutoffs. The coding of sets has two routes: source 09's extensional rooted codes with isomorphism as equality (section 5), and source 05's codes modulo bisimulation (section 7). The first new birthday has two

existence arguments: the new-subset lemma of 05 and 07 for any outer model, and the generic filter of 02 for set forcing (remark 15.5). The new-subset detectors are three different formulas (section 16).

Where the merge had to choose. (i) Hypotheses: whenever a result is proved by several sources under different hypotheses, the weakest proved hypothesis is printed, and the proof printed is one that uses only it; where this attaches a source’s argument to a weaker hypothesis than the source stated, the fact is said at the point of use (remarks 3.7 and 16.7). (ii) The first new birthday is stated for arbitrary same-ordinal transitive outer models (05, 07), with 02’s set-forcing theorem as a special case. (iii) The full-class bi-interpretation is printed as the announced theorem of Chen–Hamkins–Yang, not as a result of source 07. (iv) Symbols were renamed as listed in section 1.4.

Two later parts. [Added 3 October 2026, batch 89.] Parts VI and VII print two further manuscripts, written independently of the five above and of each other, both pinned to commit 2b7b388ba, delivered in commit 7c0f2d9f9 and placed by commit 23adb85f9. Source 10 (batch-89 manuscript 03, *Surreal Cuts, Replacement, and Atom-Support Spectra*; files 10-cuts-replacement-, labels hset:zr:) is Part VI; source 11 (batch-89 manuscript 04, *Surreal Universality Without Choice*; files 11-choiceless-universality-, labels hset:kw:) is Part VII. The numbers 10 and 11 continue this directory’s local sequence after 09. Each part keeps its own setting, hypotheses and notation convention and ends with its non-claims and provenance (sections 39 and 53). Nothing in Parts I–V was renumbered; dated notes in Parts II and V record where the new parts bear on them.

1.4 Conventions and renamed symbols

Convention 1.5 (One meaning per symbol). The following conventions hold throughout, including in material taken from sources that used other letters.

Notion	Here	In the sources
Birthday of x , as a surreal ordinal	$b(x)$	$\mathbf{b}$ (02); b (05, 07, 09); b for the ordinal and $\mathbf{b}$ for its surreal (06)
Cutoff carrier	$\mathbf{No}_{<\lambda}$	F_λ (02), S_κ (05, 06), $\mathbf{No}_{<\kappa}$ (07), K_λ (09)
Birthday structure	$\mathcal{B}_\lambda$; full class $\mathcal{B}$	$\mathfrak{B}_\lambda$ (02), $\mathcal{S}_\kappa$ (05), $\mathcal{A}_\kappa$ (06), $\mathcal{B}_\kappa$ (07, 09)
Cardinal ceiling	$\kappa(\lambda)$	$\theta(\lambda)$, the least <i>uncountable</i> cardinal $\geq \lambda$ (07); equal for epsilon λ
Pairing length	$\Lambda(\theta) = \theta^2 + \theta$	B_θ (02); $L(a, d)$ (05); $\tau(\delta)$ (06); q^2 (07)
Code	$C = (\delta, r, t)$: domain, relation word, root	(θ, d, e, r) (02); (λ, c, r) with λ a node count (05); (δ, e, r) (06); (q, d, e, r) (07)
Interpreted $=, \in$	$\sim, \in_{\mathcal{I}}$	$\mathbf{Eq}, \mathbf{Mem}$ (02); $\equiv, \in^*$ (06); $\simeq, \in_*$ (07)
Ordinal code	$\mathbf{OC}(a)$	$\mathbf{OC}$ (02), $\mathbf{Sync}$ (05), $\mathbf{OrdCode}$ (06), $\mathbf{OCode}$ (07), $O(a)$ (09)
Collector	$\mathbf{Coll}(C)$; sentence $\exists C \mathbf{Coll}(C)$	$\mathbf{CollectOrd}$ (02), $\mathbf{Coll}$ (09)

Notion	Here	In the sources
Replacement instance at $\beth_\omega$	$\text{Rep}_{\beth}$	$\text{Collect}_{\beth}$ (06)
Translated Power Set	$\text{Pow}^{\mathcal{I}}$	Power (02), $\text{Pow}^{\#}$ (06)
Eternity, fiber packing	Et , Pack	E , P (09)
Conjugation	c	c (02, 06); overline (05, 07); σ (09)
Coordinate birthday	$\text{cb}(x + iy) = \max(\text{b } x, \text{b } y)$	β (02, 05, 09); B (06, 07)
Surcomplex structures	$\mathcal{C}_\lambda$ (no i named), $\mathcal{C}_\lambda^i$ (i named); full $\mathcal{C}$, $\mathcal{C}^i$	$\mathfrak{K}_\lambda$ with i (02); $\mathcal{C}_\kappa, \mathcal{C}_\kappa^-$ (05); $\mathcal{K}_\kappa$ (06); $\mathcal{C}_b, \mathcal{C}_b^i$ (07); $\mathcal{C}_\lambda^\pm$ (09)
First new birthday	$\beta = \beta(M, N)$	δ (02, 05, 07)
New-subset formula of 05	$\text{New}(m, \theta, \nu)$	$\text{New}(c, \theta, \lambda)$, λ an enumeration length (05)
Eternity bound	ϱ	β (09)

Three collisions with the collection’s shared notation are resolved as follows (`docs/NOTATION.md`). The collection reserves F_Γ and K_Γ for the real and complex Hahn workspaces and K for a set-sized *complex* field; sources 02 and 09 used F_λ and K_λ for birthday cutoffs (09’s K_λ a *real* field), so this report writes $\mathbf{No}_{<\lambda}$ and $\mathbf{No}_{<\lambda}[i]$ instead. The collection’s H_α is a subgroup of a value group (bounded multiples of α); here H_κ is always the universe of sets of hereditary size below κ , and no value group occurs. The collection’s $\mathcal{O}$ is a valuation ring, hence the ordinal code is written OC. The letter β is reserved for the first new birthday of an outer model, as in the sibling report `foundations-and-computation/surreal-fields-across-universes`; that report uses δ for the first new *ordinal-sequence length*, a notion not named here, while here δ is only the domain ordinal of a code. Sources 02, 05 and 07 wrote δ for the first new birthday.

Two further words have two meanings in the sources. “Collect” named a sentence collecting the literal ordinals in source 02 and a Replacement instance in source 06; the two are $\exists C \text{ Coll}(C)$ and $\text{Rep}_{\beth}$ here. A sans-serif P was fiber packing in source 09 and Power Set in sources 02 and 06; they are Pack and $\text{Pow}^{\mathcal{I}}$ here, and Pack is *not* the Power Set axiom. Birthday precedence and prefix simplicity are different relations (`found:rem:simplicity` of the foundations report); prefix is defined below from order and birthday, and is never identified with precedence.

[Added 3 October 2026, batch 89: Parts VI and VII keep these conventions and add their own (`conventions 24.4` and `40.1`), where their sources’ H , SG , M_κ , S_κ , K_κ , C_α , β and δ are renamed.]

1.5 Relation to the collection and corrected repository statements

Foundations. The foundations report records the Chen–Hamkins–Yang announcement (`found:sub:announcement`), the epsilon-number cutoffs (`found:sub:cutoffs`) and the distinction between simplicity and birthday precedence (`found:rem:simplicity`). Its inventory notes that the field language has no birthday symbol; [proposition 12.5](#) supplies the standard nondefinability proof. None of the interpretation theorems of this report was in the collection before it.

Surcomplex automorphisms. The report `surcomplex/surcomplex-field-automorphisms` proves $\text{Aut}(K/F) = \{\text{id}, c\}$ for $K = F(i)$ with F real closed (`saut:thm:axis`) and, in its section `saut:sec:exponential`, subsection “Simplicity restores the canonical surreal numbers”, that a bijection preserving numerical order and prefix simplicity is the identity, so that a real-axis-preserving automorphism whose real restriction preserves simplicity is the identity or conjugation.

Since order and birthday define prefix (lemma 4.1), the bounded automorphism statements of sources 05, 06, 07 and 09 are consequences of those results; they are printed once, as lemma 19.3 and theorem 19.4, and credited there. What is new relative to that report is only source 07’s classification of the class elementary self-embeddings of the full structure, which are not assumed surjective (theorem 19.7), and source 06’s two-lift classification of elementary embeddings between different cutoffs (theorem 19.5). No question named in that report is answered or refuted; in particular its question whether $\{\text{id}, c\}$ is a proper subgroup of its group Aut_L of field automorphisms commuting with $L(z) = \log|z|$ (equivalently, whether a nontrivial exponential automorphism of **No** exists) is untouched.

Product birthdays. No argument here uses a bound $b(xy) \leq b(x) \otimes b(y)$. That bound is Gonshor’s product-birthday conjecture; the collection’s `surreal/gonshor-product-birthdays` report proves it on a specified ring of ordinal-support series and states that the general conjecture is not settled. Sources 05, 06, 07 and 09 all say that they avoid it, and the size estimates below use only field closure of the epsilon-number cutoffs.

Sibling report. The report `foundations-and-computation/surreal-fields-across-universes`, placed in the same batch, studies the external saturation of the *pure* ordered field $\mathbf{No}^M$ inside an outer universe, without a birthday symbol. Its setting meets this report’s outer-model part (Part III): the absoluteness of old arithmetic, the first new birthday, and the forcing examples. The two languages give opposite phenomena: the pure-field inclusion stays elementary but can lose saturation, while the birthday expansion loses elementarity (remark 15.8).

Other reports. The `foundations-and-computation/computable-surreals` report concerns countable effective subfields and is a different program; the undecidability remarks of section 12 are standard. No report of the collection is contradicted by this one.

Repository statements of the sources, corrected. All five sources compared themselves with the collection at the pin `4f2645599121fa872c7104995e47f86f0382351f`, which is 32 commits before the placement commit `d4e71b7`.

- Sources 05 and 07 report that the catalogue at the pin described forty catalogued reports and four newly placed drafts, and that the formalization ledger counted forty-four main texts. This was true at the pin. Before this batch was placed the catalogue listed 44 research reports in five families.
- Sources 05 and 07 report that no matching reconstruction theorem was located in the material they inspected, and source 07 that the uniform bi-interpretation had not been located in earlier published work or in the repository. Source 05 read only the first 230 lines of the foundations report and source 07 its introduction; the Chen–Hamkins–Yang announcement is recorded in that report at the pin and now. The full-class statement is therefore credited to the announcement (theorem 18.1), and the local refinements of 05 and 07 carry the same priority caveat as those of 02, 06 and 09.
- Sources 05 and 06 cite line ranges and section structure of the surcomplex automorphisms report at the pin. That report has since been rewritten; its subsection “Simplicity restores the canonical surreal numbers” still exists, now in its section `saut:sec:exponential`, and the relative result is its `saut:thm:axis`.

- Source 06 reports that a GitHub search for “Hamkins” returned no indexed matches. That was a search-index observation; a direct search of the repository finds the announcement in the foundations report.
- Sources 02 and 07 report that the incoming-report directory contained only its README at the pin; that was true at the pin.

Source 02’s own source audit, kept byte-identical as `02-birthday-cutoffs-SOURCE_AUDIT.md`, contains no statement about the repository that is false; its closing description of the elementary-inclusion criterion as “necessary, not claimed sufficient” describes source 02 and is superseded inside this report by [theorem 9.3](#).

2 Foundations and the two ordinal languages

2.1 Ambient theory and class conventions

The proofs take place in ZFC, except in [section 13](#) (ZF) and in [section 18.2](#) (GBC, stated there). All cutoff structures are sets. The statements about the full surreal class are definable-class statements, understood one first-order formula at a time. Neither a set of all surreals nor an internally available truth predicate for the entire universe is assumed. Quotient interpretations use definable equivalence relations; the Mostowski collapse supplies set representatives when we compare them with H_κ . Global Choice is not used outside [section 18.2](#).

When a transitive model M is explicitly mentioned, its surreals are the sign sequences belonging to M , and cardinalities with superscript M are computed there. A statement proved internally in an arbitrary model of ZFC does not assert that the model’s well-founded relations are externally well founded. The outer-model comparison theorems explicitly use transitive models.

2.2 The carrier and the birthday function

A surreal is a function $s : \alpha \longrightarrow \{-, +\}$ for an ordinal α . Comparison is lexicographic, with termination between $-$ and $+$. The all-plus sequence of length α is the embedded ordinal α . Throughout, $b(s)$ is the *surreal ordinal* corresponding to the length of s . Thus it is a function on the surreal carrier, not an extra ordinal sort. Set-theoretic lengths and their surreal embeddings are distinguished by context.

For an ordinal λ put $\mathbf{No}_{<\lambda} = \{x \in \mathbf{No} : b(x) < \lambda\}$, a set. For an epsilon number λ define

$$\mathcal{B}_\lambda = (\mathbf{No}_{<\lambda}; 0, 1, +, -, \cdot, <, b), \quad \mathcal{B} = (\mathbf{No}; 0, 1, +, -, \cdot, <, b).$$

The language includes no exponentiation, support predicate, membership predicate, or second-order quantifier.

Lemma 2.1 (Birthday order and birthday function). *On each $\mathcal{B}_\lambda$, and on $\mathcal{B}$, the birthday function and the birthday-precedence relation $x \prec_b y \iff b(x) < b(y)$ are definitionally interconvertible. So is the equal-birthday relation $E_b(x, y) \iff b(x) = b(y)$. Moreover, the embedded ordinals are exactly the fixed points of b .*

Proof. Given the function, precedence and equal birthday are immediate. Conversely, equality of birthdays is expressed by neither $x \prec_b y$ nor $y \prec_b x$. In the class of sign sequences of length α , the

largest in numerical order is the all-plus sequence: any other sequence of that length has a first minus sign and is smaller. Therefore $b(x)$ is the unique largest member of x 's equal-birthday class,

$$y = b(x) \iff E_b(x, y) \wedge \forall z (E_b(x, z) \rightarrow z \leq y),$$

and this uses only E_b and the order. A sign sequence is fixed by this function exactly when it is all plus. Each cutoff contains the all-plus representative of each of its birthdays, so the definitions work in every $\mathcal{B}_\lambda$. $\square$

The interconvertibility with precedence is proved in sources 02, 06 and 09; the sufficiency of the coarser equal-birthday relation is source 07's observation. In particular the results below concern a definitional variant of the birthday-precedence language of the Chen–Hamkins–Yang announcement, not an essentially stronger coding apparatus. We abbreviate $\text{Ord}(a)$ by $b(a) = a$. Quantification over embedded ordinals in $\mathbf{No}_{<\lambda}$ is precisely quantification over $\alpha < \lambda$.

2.3 Natural arithmetic, not ordinary ordinal arithmetic

For ordinal arguments, surreal addition and multiplication are the Hessenberg operations

$$\alpha + \gamma = \alpha \oplus \gamma, \quad \alpha\gamma = \alpha \otimes \gamma.$$

They are strictly increasing in each positive argument where appropriate, and $\alpha + 1$ agrees with ordinary ordinal successor. We use ordinary ordinal exponentiation only in metamathematical bounds and in explicitly set-theoretic constructions. It is not silently identified with a field operation.

Lemma 2.2 (Closure and cardinal estimates). *Every uncountable cardinal is an epsilon number. For infinite ordinals α, γ , natural sums and products have cardinality at most $\max(\aleph_0, |\alpha|, |\gamma|)$. In particular, if κ is an uncountable cardinal and $\alpha, \gamma < \kappa$, their natural sum and product are below κ .*

Proof. In Cantor normal form, a natural sum joins finitely many exponent terms; a natural product uses finitely many natural sums of their exponents. The cardinality estimate follows from the corresponding estimate for ω^ξ : for infinite ξ , its cardinality is at most $\max(\aleph_0, |\xi|)$. One can see the latter by representing ordinals below ω^ξ by finite Cantor-normal-form lists with exponents below ξ and natural-number coefficients. These observations also cover finite arguments after adjoining $\aleph_0$ to the bound.

If κ is uncountable and initial, $\omega^\alpha < \kappa$ for every $\alpha < \kappa$. Continuity gives $\omega^\kappa \leq \kappa$. The opposite inequality follows from $\omega^\alpha \geq \alpha$ and taking suprema. Hence $\omega^\kappa = \kappa$. Neither step uses regularity of κ . $\square$

All five sources prove the first assertion; van den Dries and Ehrlich already note that every uncountable cardinal is an epsilon number [3], as source 06 records. For a general epsilon number, closure of natural arithmetic below λ also follows from the classical field-cutoff theorem. This simple closure is all the ordinal arithmetic needed by the relation codes.

2.4 Hereditary size rather than rank

Lemma 2.3 (Basic hereditary-size facts). *Let κ be an uncountable cardinal. A sign sequence of length α belongs to H_κ exactly when $\alpha < \kappa$. Every subset of an ordinal $\alpha < \kappa$ belongs to H_κ . Also $H_\kappa \subseteq V_\kappa$, and $\text{Ord} \cap H_\kappa = \kappa$.*

Proof. The transitive closure of the usual set-coded graph of a sign sequence has cardinality at most $\max(\aleph_0, |\alpha|)$, and it contains enough of the domain ordinal to have cardinality at least $|\alpha|$ when α is infinite. The same estimate applies to a subset of α , and $\text{TC}(\{\alpha\}) = \alpha + 1$.

If $|\text{TC}(\{x\})| \leq \mu$ for an infinite $\mu < \kappa$, the membership relation on this transitive closure is a well-founded relation of size at most μ . Its rank function takes values below μ^+ . This follows by well-founded induction, using regularity of the successor cardinal μ^+ : the supremum of at most μ ordinals below μ^+ remains below μ^+ . Thus $\text{rank}(x) < \mu^+ \leq \kappa$, and $x \in V_\kappa$. $\square$

The statement that every subset of α belongs to H_κ does *not* say that the collection $\mathcal{P}(\alpha)$ is an element of H_κ . This distinction drives the growth results. Three notions must not be interchanged: a family whose individual members are small, a family that is a small set, and a family that belongs to H_κ . For example, all subsets of ω belong to H_{ω_1} , but their collection does not (source 06).

3 Small certificates for surreal arithmetic

A reverse interpretation cannot simply say “perform the usual set-theoretic recursion inside H_κ .” At a singular cutoff, Replacement may fail in that structure. We instead prove that each individual arithmetic computation has a certificate of the same hereditary cardinal scale as its inputs. Two routes are given: source 09’s option-graph certificates ([theorem 3.3](#)) and the prefix-pair tables of sources 05, 06 and 07 ([section 3.4](#)).

3.1 Legal cuts can be checked using prefixes

For sets of sign sequences $L < R$, let $\{L \mid R\}$ denote their simplest separator. The classical simplicity theorem supplies this separator and the bound

$$b(\{L \mid R\}) \leq \sup\{b(a) + 1 : a \in L \cup R\}. \quad (3.1)$$

The supremum of the empty set is zero. These statements can equivalently be proved by the sign-by-sign cut construction [2].

Lemma 3.1 (Prefix test for a cut value). *A sign sequence z equals $\{L \mid R\}$ if and only if it separates L and R and no proper sign prefix of z separates them.*

Proof. The forward implication is immediate. For the converse, suppose y is a separator shorter than z . If y is a prefix of z , the stated test fails. If it is not, their longest common prefix lies numerically between them and is a proper prefix of z . The set of separators is convex, so this common prefix is another separator. Again the test fails. Thus z has minimum birthday among separators, and uniqueness of the simplest separator gives the result. $\square$

The check in this lemma quantifies only over the displayed option sets and the prefixes of z . In set coding it is a bounded check: one does not quantify over all sign sequences of earlier birthdays.

3.2 Well-founded option graphs of small size

A labeled option graph has a set D of vertices, left and right successor relations, and a rank label that strictly decreases along every option edge. A valuation of such a graph assigns a sign sequence to each vertex so that its value passes [lemma 3.1](#) for the values at its options. When the graph is numeric, this valuation exists and is unique.

Lemma 3.2 (Small numeric graphs have small certificates). *Suppose μ is infinite and a numeric, well-founded option graph has at most μ vertices, with its structural data of hereditary size at most μ . Then it has a rank-labeled valuation certificate of hereditary size at most μ .*

Proof. The graph rank of every vertex is below μ^+ , by the successor-cardinal argument in lemma 2.3. Applying (3.1) by induction shows that a vertex value has birthday at most that vertex's graph rank. Thus each rank label and each sign value has hereditary size at most μ . There are at most μ vertices and at most $\mu^2 = \mu$ edges, so the union of all these data still has hereditary size at most μ . The existence construction here takes place in the ambient ZFC universe, not in a possibly deficient H_κ . $\square$

3.3 The computation graph for addition and multiplication

Take all sign prefixes of the inputs, including the inputs themselves. With

$$\mu = \max(\aleph_0, |b(x)|, |b(y)|),$$

this prefix family and its data have hereditary size at most μ . For addition, use finite signed words of prefix atoms $a(a)$; an option moves one atom to one of its sign-prefix options, reversing left and right for a negative atom. The root word is $a(x) + a(y)$. This is the usual disjunctive-sum construction of the numeric game for $x + y$.

For multiplication, use finite signed words of pair atoms $p(a, a')$, with a, a' prefixes of the respective inputs. An option of a positive pair atom is a signed three-term word

$$p(a^\circ, a') + p(a, a'^\circ) - p(a^\circ, a'^\circ). \quad (3.2)$$

It is a left option when (a°, a'°) is left–left or right–right, and a right option for the two mixed choices. Negative atoms exchange the sides and negate the replacement word. An option of a word replaces one atom by one of these words. The root is $p(x, y)$. This spells out Conway's multiplication recursion without first constructing a large universe of games.

There are at most μ possible finite words. These graphs are well founded. For addition, use the natural sum, over atoms of the word, of $\omega^{b(a)}$. For multiplication put

$$\rho(a, a') = b(a) \oplus b(a')$$

and use the natural sum of $\omega^{\rho(a, a')}$ over the pair atoms. Every pair atom in (3.2) has strictly smaller ρ than the replaced atom. A finite sum of lower powers of ω is smaller than the replaced power, so each move strictly decreases the displayed measure. The standard numeric-game arithmetic theorem identifies the values of these graphs with $x + y$ and xy ; their options are exactly the classical recursive options, including the signs in the negative summands [1].

Theorem 3.3 (Hereditary-size arithmetic certificates). *There are fixed first-order set-theoretic formulas defining surreal addition, negation, and multiplication on sign sequences such that, for inputs of hereditary size at most an infinite μ , a correct computation has a certificate of hereditary size at most μ . Verification of a supplied certificate is absolute between transitive sets containing its data. Consequently, for every uncountable cardinal κ , these formulas define in H_κ exactly the external surreal operations on its sign sequences. No assumption $H_\kappa \models \text{Replacement}$ is required.*

Proof. For $+$ and multiplication, existentially quantify a set of word vertices containing the root and closed under the explicit option rule, its exact option relations, an ordinal rank labeling decreasing on

edges, and a valuation by sign sequences passing the prefix cut test at every vertex. The node labels specify the input prefixes and the finite signed words; all checks of option generation quantify over those input prefix families and over entries of the finite words. Exactness requires both that each generated option is present and that every recorded edge is generated by the rule. The predicates for sign sequence, prefix, numerical comparison, finite word, and decreasing ordinal rank have bounded verifications on the supplied sets. In particular, a supplied prefix family is not accepted merely because its members are prefixes: for each ordinal $\xi \leq \text{dom}(s)$ it must contain the unique restriction $s \upharpoonright \xi$, and every recorded prefix must be one of these restrictions. The same completeness check is imposed on the prefix families of vertex values used by the cut test. These are bounded checks against $\text{dom}(s) + 1$ and the supplied families. A word length is a finite ordinal; this is checked by requiring every ordinal at most that length to be zero or a successor. Thus a missing prefix or a nonstandard-length word cannot evade an option requirement. The verifier is consequently a fixed bounded set-theoretic predicate, after including these domains and prefix families as certificate data. Their cardinality and hereditary-size bounds remain μ .

Any valid certificate is sound: external well-founded induction on its rank labels identifies each vertex value with the value of its generated numeric game. This also proves uniqueness of the root value. A certificate exists by the preceding graph construction and [lemma 3.2](#), with hereditary size at most μ . Negation simply reverses all signs and has the same size estimate.

For inputs in H_κ , choose the infinite bound $\mu < \kappa$ given above. The external certificate belongs to H_κ ; conversely any certificate there is externally sound by bounded absoluteness. Hence the existential formulas have the same intended values. The uses of cardinal regularity were at μ^+ , not at κ . $\square$

Remark 3.4 (Scope of the certificate theorem). This is not a proof that every class-recursive surreal operation is absolute in every hereditary-size structure. It proves the specific finite-signature claim needed here. Inversion is definable from multiplication by $xy = 1$; its existence in the epsilon-number cutoffs, and real closedness, are supplied by the classical cutoff theorem. No normal-form birthday estimates or unproved sharp product-birthday conjecture enter the argument ([section 1.5](#)).

3.4 Second route: prefix-pair tables

Sources 05, 06 and 07 certify arithmetic by a single table indexed by pairs of input prefixes. Source 05 states it at regular cardinals; source 06 isolates the single-cardinal size bound that makes it work at every uncountable cardinal, singular ones included, and source 07 uses the same bound. We print this route with 06's bound.

For a sign sequence x write $\text{Pf}_x = \{x \upharpoonright \alpha : \alpha \leq \text{b}(x)\}$ for its prefix set, including x , and let L_x and R_x be its proper prefixes below and above x ; then $x = \{L_x \mid R_x\}$. Say that w is the *simplest separator* of two families when every left member is below w , every right member is above w , and no sign sequence of smaller length has the same property. For any candidate $w \in H_\kappa$, all sign sequences shorter than w also lie in H_κ ([lemma 2.3](#)); so this minimality test is absolute between H_κ and V whenever the families are in H_κ . Uniqueness of a least-birthday separator follows from signs: two distinct separators of the same least birthday would have a common prefix at their first difference strictly between them, still separating and of smaller birthday.

Lemma 3.5 (Hereditary size of arithmetic tables). *Let κ be an uncountable cardinal, let $x, y \in \mathbf{No}_{<\kappa}$ and set*

$$\mu = \max\{\aleph_0, |\text{b}(x)|, |\text{b}(y)|\} < \kappa.$$

The functions on $\text{Pf}_x \times \text{Pf}_y$ given by $T_{x,y}(u, v) = u + v$ and $U_{x,y}(u, v) = uv$ have graphs of hereditary size at most μ ; in particular they belong to H_κ .

Proof. There are at most μ pairs of prefixes. Every prefix belongs to $\mathbf{No}_{<\mu^+}$, which is a field by lemma 2.2 and the classical cutoff theorem. Therefore every value in either table has birthday below μ^+ and hereditary size at most μ . A set of at most μ objects, each of hereditary size at most μ , has hereditary size at most μ ; the finite layers needed for ordered pairs and triples do not change this.

This argument is carried out in the ambient ZFC universe to establish the existence of a particular witness in H_κ . It does not invoke Replacement inside H_κ , and it uses a *single* infinite cardinal μ , not a union of sizes approaching a singular κ . It also works when $\mu^+ = \kappa$. $\square$

Theorem 3.6 (Table definitions of the operations; sources 05, 06, 07). *Let $\text{Add}(x, y, z)$ say that there is a function T on $\text{Pf}_x \times \text{Pf}_y$ with $T(x, y) = z$ such that every entry obeys*

$$T(u, v) = \{T(u^L, v), T(u, v^L) \mid T(u^R, v), T(u, v^R)\} \quad (u^L \in L_u, v^L \in L_v, u^R \in R_u, v^R \in R_v),$$

the braces meaning the simplest-separator predicate. Let $\text{Mult}(x, y, z)$ say that there is a function U on $\text{Pf}_x \times \text{Pf}_y$ with $U(x, y) = z$ such that $U(u, v)$ is the simplest separator of the left options $U(u', v) + U(u, v') - U(u', v')$ with $(u', v') \in L_u \times L_v \cup R_u \times R_v$ and the right options of the same form with $(u', v') \in L_u \times R_v \cup R_u \times L_v$, where each sum and difference is certified by Add and sign reversal. Then for every uncountable cardinal κ and all $x, y, z \in \mathbf{No}_{<\kappa}$,

$$H_\kappa \models \text{Add}(x, y, z) \iff z = x + y, \quad H_\kappa \models \text{Mult}(x, y, z) \iff z = xy.$$

Proof. The actual tables are in H_κ by lemma 3.5 and satisfy the canonical Conway recursions, proving existence. Conversely, any accepted table is identified with the actual one by external well-founded induction on the pair of prefix lengths, ordered so that decreasing either coordinate decreases the pair: every option entry is earlier, the simplest-separator test is absolute, and the separator is unique. For multiplication, addition has already been identified. Each invocation of Add has its own hereditarily small witness; first-order satisfaction does not require that a single set collect the witnesses to all invocations. $\square$

The two routes certify the same operations; the table route uses fewer auxiliary objects, the option-graph route makes the verifier an explicitly bounded predicate. Either suffices for the reverse interpretation of section 6.

Remark 3.7 (Regularity in source 05). Source 05 states its size bookkeeping and its table lemma for uncountable regular κ and declines to infer a singular-cardinal theorem. Its table argument uses only the field closure of $\mathbf{No}_{<\mu^+}$ and the single bound μ , exactly as in lemma 3.5; regularity of κ is not used there. The singular case is sources 06 and 07's theorem, and the base's.

4 Definable sign access and a bounded pairing map

4.1 Recovering the sign tree from the two orders

For x, y, z let $\text{Between}(z; x, y)$ say that z lies in the closed numerical interval with endpoints x and y . Define

$$\text{Pre}(x, y) \iff \forall z ((\text{Between}(z; x, y) \wedge z \neq x) \longrightarrow b(x) < b(z)). \quad (4.1)$$

Lemma 4.1 (Definable prefix relation). *In every $\mathcal{B}_\lambda$, and in $\mathcal{B}$, (4.1) holds exactly when x is a sign prefix of y , allowing equality.*

Proof. If x is a prefix of y , every other point between them must continue the sign sequence of x ; a point diverging earlier would lie outside that interval. Its birthday is therefore larger.

If x is not a prefix of y and y is a proper prefix of x , the endpoint $z = y$ violates the formula. In the remaining case the longest common prefix of x, y lies strictly between them and has birthday smaller than that of x . This prefix belongs to the same cutoff. Finally, for $x = y$ the formula is vacuous, as required. $\square$

Remark 4.2 (Four prefix formulas). Each source defines prefix from $<$ and b by its own formula; all are correct. Source 02 requires $b(x) \leq b(y)$ and that every z with $b(z) < b(x)$ compares with x as with y ; source 06 adds the symmetric comparison. Sources 05 and 07 require $b(x) \leq b(y)$ and that no z strictly between x and y has $b(z) \leq b(x)$ (05) or $b(z) < b(x)$ (07). In the betweenness form the length conjunct cannot be dropped: for $x = 1$ and $y = 0$ no surreal of birthday at most 1 lies strictly between them, but 1 is not a prefix of 0 (source 05, which checks this case in its finite suite).

Define the bit predicate by

$$\begin{aligned} \text{Bit}(s, \alpha) \iff & \text{Ord}(\alpha) \wedge \alpha < b(s) \wedge \\ & \exists t (\text{Pre}(t, s) \wedge b(t) = \alpha \wedge t < s). \end{aligned} \quad (4.2)$$

The unique prefix of length α lies below s precisely when the sign of s at position α is plus. Thus

$$\{\alpha < \delta : \text{Bit}(s, \alpha)\}, \quad b(s) = \delta, \quad (4.3)$$

runs through *all* subsets of δ , once each, for every $\delta < \lambda$. This uses a first-order quantifier over numbers of one birthday, not a second-order quantifier in the language. The exact length matters: trailing minus signs are not discarded, so the code has an explicit domain.

4.2 Pairing without reconstructing ordinary ordinal multiplication

For a positive embedded ordinal θ , put

$$p_\theta(\xi, \zeta) = \theta\xi + \zeta, \quad \Lambda(\theta) = \theta^2 + \theta. \quad (4.4)$$

All operations in (4.4) are field operations, hence natural ordinal operations on these arguments.

Lemma 4.3 (Bounded pairing). *The map p_θ injects $\theta \times \theta$ into $\Lambda(\theta)$. For every epsilon number λ and every $0 < \theta < \lambda$, $\Lambda(\theta) < \lambda$. For infinite θ , $|\Lambda(\theta)| = |\theta|$.*

Proof. Suppose $\xi < \xi' < \theta$ and $\zeta, \zeta' < \theta$. Since $\xi + 1 \leq \xi'$ in ordinal order, positivity and distributivity give

$$\theta\xi + \zeta < \theta\xi + \theta = \theta(\xi + 1) \leq \theta\xi' \leq \theta\xi' + \zeta'.$$

For fixed ξ , cancellation gives injectivity in ζ . Also $\theta\xi + \zeta < \theta^2 + \theta$. Closure below λ follows from its being a field cutoff. The cardinal estimate is lemma 2.2, together with $\theta \leq \Lambda(\theta)$. $\square$

Source 02 uses the same pairing and bound (its B_θ). Consequently a sign sequence of birthday $\Lambda(\theta)$ codes a binary relation on θ . To make this code canonical for a given relation, require every

plus position to be in the image of p_θ ; unused positions are minus. This last requirement is the first-order formula

$$\forall \rho < \Lambda(\theta) (\text{Bit}(r, \rho) \longrightarrow \exists \xi, \zeta < \theta [\rho = p_\theta(\xi, \zeta)]).$$

Here and below bounded ordinal quantifiers include the predicate Ord .

For a graph between two different ordinals $\delta, \epsilon < \lambda$, use $\theta = \delta + \epsilon + 1$, which is above both. Coding the graph in the $\delta \times \epsilon$ rectangle of this square again stays below λ . The construction requires no definable global pairing on the entire class of ordinals, and no reconstruction of ordinary ordinal arithmetic.

Remark 4.4 (Four pairings in the sources). The other sources pack relations differently; each variant works with the codes below after the obvious changes. Source 05 uses the rectangle map $p_a(i, j) = aj + i$ for $i < a$, $j < d$, with length $a(d + 1)$, which is the same idea with the roles of the coordinates exchanged. Source 06 uses the polynomial of [lemma 4.5](#), and source 07 the radix pairing of [lemma 4.6](#).

Lemma 4.5 (A one-polynomial pairing; source 06). *On ordinal surreals put $\varpi(\alpha, \gamma) = (\alpha + \gamma)^2 + \alpha$. Then ϖ is injective on pairs of ordinals, increasing in each coordinate, and maps $\kappa \times \kappa$ into κ for every uncountable cardinal κ .*

Proof. Put $\sigma = \alpha + \gamma$. Since $0 \leq \alpha \leq \sigma$, $\sigma^2 \leq \varpi(\alpha, \gamma) \leq \sigma^2 + \sigma$. If $\sigma < \sigma'$ are ordinals, then $\sigma + 1 \leq \sigma'$ and $\sigma^2 + \sigma < (\sigma + 1)^2 \leq \sigma'^2$. The intervals belonging to different sums are therefore disjoint. Equal values force equal sums, then equal first coordinates by subtraction, and finally equal second coordinates. Monotonicity follows from positivity and monotonicity of addition and squaring on nonnegative elements, and the bound is [lemma 2.2](#). $\square$

Lemma 4.6 (Admissible radices; source 07). *Let $\text{Rad}(q)$ say that q is an ordinal, $q > 1$, and $a + a' < q$ for all ordinals $a, a' < q$. The ordinals satisfying Rad are exactly $q = \omega^\rho$ with $\rho > 0$; for each of them $p_q(a, a') = qa + a'$ injects $q \times q$ into q^2 . Both statements are absolute between $\mathcal{B}$ and every initial cutoff containing the ordinals in question. Below an epsilon number λ the admissible radices are cofinal and $q^2 < \lambda$ for each of them.*

Proof. For $q = \omega^\rho$ every ordinal below q has Cantor normal form with exponents below ρ , and natural addition introduces no new exponent. If $q > 1$ is not a single monomial with coefficient one, split its normal form into two nonzero proper parts (splitting a coefficient if needed) to get $a, a' < q$ with $a \oplus a' = q$; $q = 1$ is excluded. For $a < q$ the exponents of $q \otimes a$ are all at least ρ , those of $a' < q$ are below ρ , and cancellation of natural addition recovers a and a' ; moreover $qa + a' < q(a + 1) \leq q^2$. For $\xi < \lambda$ the radix $\omega^{\xi+1}$ exceeds ξ and lies below λ , and $(\omega^\rho)^2 = \omega^{\rho \oplus \rho} < \lambda$ since $\lambda = \omega^\lambda$ is closed under natural addition below itself. All ordinals below an argument of an initial cutoff are present there, so the quantified test is unchanged. $\square$

5 Pointed relation codes and the exact interpreted universe

5.1 The definable domain of codes

A prospective code is a triple

$$C = (\delta, r, t), \quad 0 < \delta < \lambda, \quad t < \delta, \quad \text{b}(r) = \Lambda(\delta),$$

with δ, t embedded ordinals and with unused bits of r equal to minus. Write

$$E_C(i, j) \iff i, j < \delta \wedge \text{Bit}(r, p_\delta(i, j)).$$

The orientation is *child belongs to parent*: $i E_C j$ will mean that the object at node i is a member of the object at node j .

Require extensionality,

$$\forall i, j < \delta [\forall k < \delta (E_C(k, i) \leftrightarrow E_C(k, j)) \longrightarrow i = j], \quad (5.1)$$

and well-foundedness in the following form:

$$\begin{aligned} \forall u [b(u) = \delta \longrightarrow ((\exists j < \delta \text{Bit}(u, j)) \longrightarrow \\ \exists j < \delta [\text{Bit}(u, j) \wedge \forall i < \delta (\text{Bit}(u, i) \longrightarrow \neg E_C(i, j))]]). \end{aligned} \quad (5.2)$$

By (4.3), this is *actual* well-foundedness of the set relation, not just well-foundedness against a restricted collection of subsets.

Finally require generation by the root:

$$\begin{aligned} \forall u [b(u) = \delta \longrightarrow (\text{Bit}(u, t) \wedge \forall i, j < \delta [\text{Bit}(u, j) \wedge E_C(i, j) \longrightarrow \text{Bit}(u, i)] \\ \longrightarrow \forall i < \delta \text{Bit}(u, i))]. \end{aligned} \quad (5.3)$$

Thus no proper downward-closed subset of the graph contains the root. Let $\text{Code}(C)$ be the conjunction of these conditions. The triple is an ordinary finite-dimensional interpretation domain in the single-sorted structure $\mathcal{B}_\lambda$. Source 02 uses four-tuples with an extra domain word that selects a subset of θ as nodes, and source 07 four-tuples with a radix; in both, a “clean” or “no junk” condition plays the role of the unused-bit convention.

Lemma 5.1 (Rooted collapse). *Every valid code has a unique Mostowski collapse $\pi_C : \delta \longrightarrow T_C$ onto a transitive set. If $x_C = \pi_C(t)$, then*

$$T_C = \text{TC}(\{x_C\}), \quad |\delta| = |\text{TC}(\{x_C\})|.$$

Proof. Define by well-founded recursion

$$\pi_C(j) = \{\pi_C(i) : i E_C j\}.$$

Extensionality and well-founded induction show that π_C is injective: if a collision existed, choose one minimizing the maximum of the two node ranks; comparison of predecessor values gives equal predecessor sets, and extensionality identifies the nodes. Its image is transitive, and the displayed recursion gives an isomorphism. Uniqueness follows by well-founded induction as well; this is the usual Mostowski-collapse argument.

The nodes reachable from t by finitely many downward edges form a downward-closed subset containing t . By (5.3) this is all of δ . Under the collapse, these are exactly x_C and its iterated members. Thus the image is $\text{TC}(\{x_C\})$, giving the cardinal equality. $\square$

Root generation is not needed merely to encode arbitrary sets. It is useful here because it makes the hereditary-size cost exact rather than just an upper bound polluted by unrelated graph components.

5.2 Equality and membership are definable

For codes $C = (\delta, r, t)$ and $D = (\epsilon, s, u)$, define $C \sim D$ when a coded bijection $f : \delta \rightarrow \epsilon$ preserves E , sends t to u , and reflects E . Such a graph is expressed by a sign sequence in a square of width $\delta + \epsilon + 1$, as in [lemma 4.3](#). The familiar function, bijection, and edge-preservation clauses use only bounded ordinal quantifiers and bits. Therefore $\sim$ is a single first-order relation.

For membership, let $C \in_{\mathcal{I}} D$ mean that there is a coded injection $f : \delta \rightarrow \epsilon$ such that

$$E_D(f(t), u), \tag{5.4}$$

$$E_C(i, j) \longleftrightarrow E_D(f(i), f(j)) \quad (i, j < \delta), \tag{5.5}$$

$$E_D(v, f(j)) \longrightarrow \exists i < \delta [E_C(i, j) \wedge f(i) = v] \quad (j < \delta, 0 \leq v < \epsilon). \tag{5.6}$$

In (5.6), the quantifier on v simply ranges over the embedded ordinals below ϵ . The clause says that the image contains *all* predecessors of its nodes, not merely some of them. Sources 06 and 07 use the same downward-closed image clause; source 02 instead defines membership through an isomorphism of the root with the cone of a predecessor, using a definable reachability relation.

Lemma 5.2 (Correctness of equality and membership). *For valid codes,*

$$C \sim D \iff x_C = x_D, \quad C \in_{\mathcal{I}} D \iff x_C \in x_D.$$

Proof. A root-preserving isomorphism gives equal collapsed roots by uniqueness of collapse. Conversely, if the roots agree, the two images are the same transitive closure by [lemma 5.1](#), and $\pi_D^{-1}\pi_C$ gives the required isomorphism. Its graph has size at most $\max(|\delta|, |\epsilon|)$ and is codable within the same cutoff.

For membership, (5.5) and (5.6), followed by induction along E_C , imply $\pi_D(f(j)) = \pi_C(j)$ for all j . The root condition then gives $x_C \in x_D$. Conversely, if $x_C \in x_D$, the inclusion $\text{TC}(\{x_C\}) \subseteq \text{TC}(\{x_D\})$ transports through the collapse maps to an injection satisfying all three clauses. Its image is predecessor-closed, so (5.6) really does hold. The graph is again small enough to be coded below the cutoff. $\square$

The predecessor-full clause cannot be omitted; Appendix B gives the counterexample, which the finite suites of sources 06, 07 and 09 check.

Let $\mathcal{I}(\mathcal{B}_\lambda)$ be the quotient of valid codes by $\sim$, with the membership relation just defined. All quantifiers in its interpreted formulas are replaced by quantifiers over triples satisfying Code. No quotient class is assumed to be an element of the original structure.

5.3 Cardinal rounding

Theorem 5.3 (Exact collapse range). *For every epsilon number λ , the map $[C] \mapsto x_C$ is an isomorphism*

$$\mathcal{I}(\mathcal{B}_\lambda) \cong (H_{\kappa(\lambda)}, \in).$$

More precisely, the ordinals δ supporting rooted codes for a given set x are exactly those with

$$0 < \delta < \lambda, \quad |\delta| = |\text{TC}(\{x\})|. \tag{5.7}$$

Proof. Every code has $\delta < \lambda \leq \kappa(\lambda)$. Hence its collapse has hereditary size $|\delta| < \kappa(\lambda)$, by [lemma 5.1](#). This proves one inclusion of the range.

Conversely, suppose $x \in H_{\kappa(\lambda)}$. Its nonempty transitive closure has some cardinal $\nu < \kappa(\lambda)$. By the definition of cardinal ceiling, $\nu < \lambda$: if λ is a cardinal this is immediate; otherwise $\nu \leq |\lambda| < \lambda$. Choose a bijection from ν onto $\text{TC}(\{x\})$, transfer membership to ν , and designate the node for x . The resulting graph is extensional, well founded, and generated by that node. It has a sign code of length $\Lambda(\nu) < \lambda$. Thus x occurs in the range.

For any other $\delta < \lambda$ equipotent to this transitive closure, the same construction works with a bijection from δ . Necessity in (5.7) was already proved by rooted collapse. Correctness of both relations is lemma 5.2. $\square$

Choice is used here only to well-order one transitive closure at a time; the interpretation keeps all valid codes and never selects one code per set. Source 02 proves this theorem one way, at every epsilon number, as a four-dimensional interpretation; source 07 proves the same range at every epsilon number (its cardinality plateaux, with $\theta(\lambda)$ in place of $\kappa(\lambda)$), also stating it forward only; sources 05 and 06 prove it at regular, respectively all uncountable, cardinals.

Remark 5.4 (Why the bound is sharp; source 02). No set of hereditary size $\kappa(\lambda)$ can be represented by these codes: its collapse would require at least $\kappa(\lambda)$ distinct nodes, while each domain is an ordinal below $\lambda \leq \kappa(\lambda)$. This obstruction is intrinsic to the stated interpretation. It is not a claim that no different interpretation in $\mathcal{B}_\lambda$ could represent a larger structure.

Example 5.5 (The empty set and a first nonempty code). The empty set uses one node, no edges, and root zero. With $\delta = 1$, the canonical relation string has birthday $\Lambda(1) = 2$ and both signs minus. Thus a code for the empty set is $(1, --, 0)$, not the surreal zero itself. For $\{\emptyset\}$ take $\delta = 2$, root 1, and the single edge $0E1$. The relation string has length 6 and its only plus is at $p_2(0, 1) = 1$. Numeric values and the sets they code must not be identified. A three-node example with two labelings is in Appendix C.

Corollary 5.6 (Countable cutoffs recover all countable sets; sources 02, 07, 09). *For every countable epsilon number λ , in particular ε_0 , $\mathcal{I}(\mathcal{B}_\lambda) \cong (H_{\omega_1}, \in)$.*

Proof. A countable epsilon number is strictly larger than ω and strictly below ω_1 , so its cardinal ceiling is ω_1 . Concretely, every hereditarily countable set has a rooted code on a finite ordinal or on ω , and $\Lambda(\omega) = \omega^2 + \omega < \varepsilon_0$. $\square$

The recovered sets include countable ordinals far larger than ε_0 . Those larger ordinals are recovered as *collapsed relations*, not as all-plus scalar sign sequences in the cutoff: every countable ordinal α has a membership graph labeled by natural numbers, whose relation word has birthday below $\omega^2 + \omega$. Labels need not preserve ordinal order; the collapse recovers that order from the edges. One should therefore not infer a low set-theoretic rank of the interpreted objects from a low birthday of the numerical codes (source 07).

6 Bi-interpretation and recovery of the cutoff

6.1 The ordinal and sign-graph bridges

For an actual embedded ordinal $a < \lambda$, there is a distinguished code $\text{OC}(a)$ with domain $a + 1$, root a , and edge relation iEj iff $i < j$. Its sign relation code is uniquely determined by the unused-bit convention. The formula defining $\text{OC}(a)$ is uniform, and its collapse is the von Neumann ordinal a . Because λ is a limit and a ring cutoff, both $a + 1$ and $\Lambda(a + 1)$ are below λ .

Definition 6.1 (Visible interpreted ordinals). For a set code C define

$$\text{VisOrd}(C) \iff \exists a [\text{Ord}(a) \wedge C \sim \text{OC}(a)].$$

Under the collapse isomorphism, these are exactly the ordinals below λ , not necessarily all ordinals of $H_{\kappa(\lambda)}$.

There is also a definable bridge $\text{SignGraph}(s, C)$ saying that C represents the set-coded sign function of the scalar s : its domain is $\text{OC}(\text{b}(s))$, its values are the interpreted ordinals 0 and 1, and at each actual $\alpha < \text{b}(s)$ it has value 1 exactly when $\text{Bit}(s, \alpha)$ holds. Application of a set-coded function and the Kuratowski ordered-pair construction are ordinary fixed formulas in the interpreted membership language. The ordinal bridge supplies their arguments. Thus SignGraph is a fixed first-order formula, not an external assignment operation.

For each s there is exactly one equivalence class of codes C with $\text{SignGraph}(s, C)$; conversely, every sign function whose length is below λ arises this way. Existence follows from the hereditary-size estimate for sign graphs and [theorem 5.3](#).

6.2 A first-order test for a cardinal cutoff

Define the *visible-ordinal collector* formula

$$\text{Coll}(C) \iff \text{Code}(C) \wedge \forall D [\text{Code}(D) \longrightarrow (D \in_{\mathcal{I}} C \leftrightarrow \text{VisOrd}(D))]. \quad (6.1)$$

It says that one interpreted set collects all the scalar ordinals. This is not the assertion that the scalar carrier itself is a set inside itself.

Theorem 6.2 (The cutoff is a set exactly in the noncardinal case; sources 02, 09). *For an epsilon number λ ,*

$$\mathcal{B}_\lambda \models \exists C \text{ Coll}(C) \iff \lambda \text{ is not a cardinal.}$$

When it exists, the collector is unique modulo $\sim$ and its collapse is exactly the ordinal λ .

Proof. The elements specified in (6.1) are exactly all ordinals $a < \lambda$. In the ambient set universe the unique set with precisely those elements is λ . Hence a collector exists precisely when $\lambda \in H_{\kappa(\lambda)}$, equivalently $|\lambda| < \kappa(\lambda)$, since $\text{TC}(\{\lambda\}) = \lambda + 1$. For a cardinal λ the right side fails; otherwise $\kappa(\lambda) = (|\lambda|)^+$ and it holds. Uniqueness follows from extensionality of the interpreted universe. $\square$

Source 02's sentence CollectOrd is this sentence written with its own codes. The theorem recovers the *ordinal value* of a noncardinal cutoff as an interpreted object. It is not a construction of a surreal number with birthday λ inside $\mathbf{No}_{<\lambda}$. This collector is the additional information needed to recover a noncardinal cutoff after interpreting its hereditary-size universe. Merely saying that $\mathcal{B}_\lambda$ interprets $H_{\kappa(\lambda)}$ would lose it.

6.3 The reverse interpretation

If $\lambda = \kappa$ is a cardinal, interpret surreals in H_κ as all set-coded functions from an ordinal into $\{0, 1\}$. By [lemma 2.3](#), these are exactly the sign sequences of length below κ . Define the order by first disagreement, birthday by the all-one function on the domain ordinal, and arithmetic by the certificate formulas of [theorem 3.3](#) (or the tables of [theorem 3.6](#)). Call this interpretation $\mathcal{J}$.

If λ is not a cardinal, work instead in the pointed structure $(H_{\kappa(\lambda)}, \in, \lambda)$. Use the same definitions, with the extra domain restriction that the sign length is below the named ordinal λ . The certificate verifier may use auxiliary sign sequences anywhere in $H_{\kappa(\lambda)}$; it is not required to carry out a large recursion inside the smaller interpreted sign domain. The root outputs are below λ by classical cutoff closure. This gives a uniform interpretation $\mathcal{J}_{\text{cut}}$ in the language with one named cutoff. Without the named cutoff, interpreting sign sequences in H_{ω_1} gives $\mathbf{No}_{<\omega_1}$, not $\mathbf{No}_{<\varepsilon_0}$; this is why source 07 stated its plateau theorem forward only.

Theorem 6.3 (Uniform local bi-interpretation). *The interpretations just constructed prove all assertions of [theorem 1.1](#). The maps witnessing the two bi-interpretations are definable, not merely external bijections.*

Proof. Start on the set side. A code in the interpreted surreal structure gives an extensional well-founded relation on an ordinal. Its Mostowski-collapse root defines the map back to the original set. The graph of the collapse belongs to the same H_κ : its domain, range, and all values have hereditary size bounded by a fixed infinite cardinal below κ . Consequently this correspondence is defined by a first-order formula in H_κ , even when Replacement is not an axiom of that structure. The collapse equation is a bounded verification of the supplied function. Every set has a code, and equality and membership are respected by [lemma 5.2](#). In the pointed case, the collector collapses to the named cutoff, so the correspondence preserves that constant as well.

Start on the surreal side. The formula $\text{SignGraph}(s, C)$ identifies s with the sign function representing it in the copy of the set universe. In the cardinal case all sign functions in the interpreted H_λ occur. In the noncardinal case, the collector defines λ , and the restricted reverse interpretation uses exactly lengths below that ordinal. Thus SignGraph is a definable bijection to the twice-interpreted surreal structure. It preserves numerical order and birthday directly from the sign rules, and it preserves arithmetic by the certificate theorem. This proves the required definable isomorphisms in both directions. $\square$

At cardinal cutoffs sources 06 and 07 give the same two round trips with their own codes (06 by a formula Eval for the collapse and a sign-function bridge; 07 by an ordinal-code comparison and the interpreted sign name $\langle b(x), A_x \rangle$), and source 05 gives them with graphs modulo bisimulation ([section 7](#)).

Corollary 6.4 (The first epsilon cutoff and hereditary countability). *The structure $\mathcal{B}_{\varepsilon_0}$ is parameter-free bi-interpretable with $(H_{\omega_1}, \in)$.*

Proof. The theorem first gives the pointed structure $(H_{\omega_1}, \in, \varepsilon_0)$. The ordinal ε_0 is parameter-free definable in H_{ω_1} as the least nonzero fixed point of ordinary $\alpha \mapsto \omega^\alpha$. For any countable arguments, the functions witnessing ordinal arithmetic and the relevant well-orders are countable sets, hence belong to H_{ω_1} . Their well-founded recursive verification is absolute. The least-fixed-point specification therefore picks out the actual ε_0 there. Naming it is a definitional expansion, which may be removed. $\square$

The same argument applies to any noncardinal epsilon number that is parameter-free definable, with its correct ordinal value, in the recovered hereditary-size universe. It is not an assertion that every ordinal can be named parameter-free.

Remark 6.5 (Questions of the sources answered here). Source 02 claimed only a one-way interpretation and warned that a two-way theorem “should specify whether the target carries a predicate for that segment or a constant for λ ”. [Theorem 6.3](#), source 09’s, is exactly such a theorem, with a constant;

at ε_0 it is even unpointed. Source 06's question on noncardinal cutoffs, source 07's question on recovering a noncardinal cutoff, and the noncardinal half of source 05's question on optimal cutoffs are answered for epsilon-number cutoffs by the same theorem; the singular-cardinal half of 05's question is answered by sources 06 and 07. The questions stay open for cutoffs that are not epsilon numbers (section 21).

6.4 Coherence of the interpretation

Proposition 6.6 (Coherence under cutoff inclusions). *If $\lambda \leq \eta$ are epsilon numbers, the inclusion of scalar cutoffs induces, under $\mathcal{I}$, the actual inclusion*

$$H_{\kappa(\lambda)} \longrightarrow H_{\kappa(\eta)}.$$

A code keeps the same collapsed value, and all sign-graph bridges agree. For cardinal cutoffs the reverse interpretations also commute with the actual inclusions.

Proof. For a code already below λ , the larger structure has no additional sign sequences of birthday δ for any $\delta < \lambda$: both contain *all* sequences of each such length in the same ambient universe. Thus the subset tests in well-foundedness and root generation are unchanged. Equality and membership agree by their collapse characterizations; any needed witness graph has a code below λ as proved above. The collapse is defined from the same relation, so its value is unchanged. The sign-graph bridge refers to the same prefixes and bits. Finally the reverse arithmetic formulas compute the same external operations. $\square$

Sources 02 (at epsilon numbers, forward), 06 and 07 (at cardinal cutoffs, both directions) prove the same compatibility. If two noncardinal cutoffs have the same cardinal ceiling, the induced inclusion on recovered *unpointed* set universes is the identity. Their recovered distinguished cutoffs nevertheless differ. This is precisely why elementary inclusion will usually fail.

7 Second route: graphs modulo bisimulation

Source 05 codes sets by well-founded graphs that need not be extensional or generated by their root, and takes *bisimilarity* as equality. This avoids any extensionality or canonical-enumeration requirement: two different terminal nodes both represent $\emptyset$, and a disjoint component below no root changes nothing. We write its codes with the pairing of lemma 4.3; source 05 uses its rectangle map (remark 4.4), which changes nothing below. Source 05 proved the resulting bi-interpretation at uncountable regular cardinals; the argument below uses regularity nowhere (the certificates of section 3 are the only arithmetic input), and we state it at the cardinal cutoffs of theorem 1.1, where the base already proves the conclusion by the first route. The value of the second route is its different equality test.

A *graph code* is a triple $G = (\nu, e, t)$ with $\nu > 0$ and $t < \nu$ embedded ordinals and $b(e) = \Lambda(\nu)$; its relation is $E_G(i, j) \iff i, j < \nu \wedge \text{Bit}(e, p_\nu(i, j))$, again oriented from a member to a set. Let $\text{WF}(G)$ be the well-foundedness clause (5.2). No extensionality or root generation is required. The *decoration* of G is the unique function with

$$v_G(j) = \{v_G(i) : i E_G j\}, \quad \text{dec}(G) = v_G(t);$$

ordinary well-founded recursion suffices, and the Mostowski collapse is its extensional special case.

Proposition 7.1 (Exactly the hereditary sets; source 05). *For an uncountable cardinal κ , the decorations of graph codes in $\mathcal{B}_\kappa$ are exactly the members of H_κ .*

Proof. Every member of $\text{TC}(\{v_G(j)\})$ is a value at a node reachable by a finite predecessor chain, so it lies in the range of v_G , of cardinality at most $|\nu| < \kappa$. Conversely, pull membership on a transitive set containing $a \in H_\kappa$ back to an ordinal $\nu < \kappa$ along a bijection and root it at the label of a ; induction gives decoration a . $\square$

For graph codes $G = (\nu, e, t)$ and $G' = (\nu', e', t')$, a relation $Q \subseteq \nu \times \nu'$ is a *bisimulation* when every $(i, j) \in Q$ satisfies

$$\begin{aligned} i' E_G i &\implies \exists j' (j' E_{G'} j \wedge (i', j') \in Q), \\ j' E_{G'} j &\implies \exists i' (i' E_G i \wedge (i', j') \in Q). \end{aligned}$$

Code Q by one sign sequence q in a square of width $\nu + \nu' + 1$, and let $\text{Bis}(G, G', q)$ be the resulting first-order formula. Define

$$G \approx G' \iff \exists q [\text{Bis}(G, G', q) \wedge (t, t') \in Q_q], \quad G \in_\approx G' \iff \exists j < \nu' [E_{G'}(j, t') \wedge G \approx G'[j]],$$

where $G'[j] = (\nu', e', j)$ is the same graph rerooted at j .

Theorem 7.2 (Equality is bisimilarity; source 05). *For well-founded graph codes, $G \approx G' \iff \text{dec}(G) = \text{dec}(G')$ and $G \in_\approx G' \iff \text{dec}(G) \in \text{dec}(G')$. Consequently the graph codes modulo $\approx$, with $\in_\approx$, form a parameter-free interpretation of $(H_\kappa, \in)$ in $\mathcal{B}_\kappa$.*

Proof. Suppose Q is a bisimulation relating the roots. By well-founded induction on i along E_G , every j with $(i, j) \in Q$ satisfies $v_G(i) = v_{G'}(j)$: a member $v_G(i')$ of $v_G(i)$ is matched by the forth clause with a predecessor j' of j and equals $v_{G'}(j')$ by induction, and the back clause gives the other inclusion. Conversely, if the root values agree, the set relation $Q = \{(i, j) : v_G(i) = v_{G'}(j)\}$ is a bisimulation relating the roots, because equal decorated sets supply the forth and back witnesses without selecting unique matches; it has a code in the same cutoff. Membership follows by applying this to the rerooted graph. Hence $\approx$ is an equivalence relation, a conclusion established semantically rather than presumed from the shape of the definition, and [proposition 7.1](#) gives surjectivity. $\square$

Example 7.3 (Duplicates and irrelevant components). One terminal node represents $\emptyset$. A graph with two terminal nodes and a root above both represents $\{\emptyset\}$, as does a graph with only one terminal child. Their rooted graphs are not isomorphic, but their roots are bisimilar. Graph isomorphism is therefore *not* the right equality for these codes; source 05's finite suite exercises exactly this case.

The round trips of source 05 are the analogues of those in [theorem 6.3](#). The set-side map is the decoration relation “there is a function v on ν satisfying the recursion with $v(t) = a$ ”, whose witness has hereditary size at most $\max(\aleph_0, |\nu|)$. On the surreal side, for an ordinal α let $\text{OC}(\alpha)$ be the graph on $\alpha + 1$ with $i E j \iff i < j$ and root α , so that $G \approx \text{OC}(\alpha)$ iff $\text{dec}(G) = \alpha$; and for a surreal x let A_x be the interpreted set whose members are the classes $\text{OC}(\gamma)$ for the ordinals $\gamma < \text{b}(x)$ with $\text{Bit}(x, \gamma)$. The number x is then matched with the interpreted sign name $\langle \text{b}(x), A_x \rangle$, which externally decodes to $\langle \text{b}(x), \{\gamma : x(\gamma) = +\} \rangle$. Source 07 uses the same sign-name round trip with extensional codes.

Remark 7.4 (Internal versions and models of ZFC; sources 05, 07). All constructions above are theorems of ZFC and can be carried out inside any model $M \models \text{ZFC}$, with M 's own ordinals, subsets and well-founded relations; for the full class $\mathbf{No}^M$ they recover M itself. For a nonstandard M this is an internal statement: graph well-foundedness is M 's notion, not external well-foundedness. For set models M, N of ZFC, writing $\mathcal{B}^M$ and $\mathcal{B}^N$ for the internal birthday structures regarded externally, source 07 derives

$$M \equiv N \iff \mathcal{B}^M \equiv \mathcal{B}^N, \quad M \cong N \iff \mathcal{B}^M \cong \mathcal{B}^N,$$

by transporting sentences and isomorphisms through the uniform interpretations and their definable composites. These full-class and internal statements are instances of the announced Chen–Hamkins–Yang theorem ([theorem 18.1](#)), not results of sources 05 or 07.

8 Ring cutoffs and the finite boundary

The classical ring statement is weaker than the field statement: the infinite ring cutoffs $\mathbf{No}_{<\lambda}$ are exactly those with $\lambda = \omega^{\omega^\alpha}$ [5, Theorems 1.1–1.2]. Source 02 observes that the forward interpretation needs only ring operations.

Proposition 8.1 (Ring version; source 02). *The interpretation $\mathcal{I}$ and its exact image remain valid for every infinite ordinal λ for which $\mathbf{No}_{<\lambda}$ is a subring of $\mathbf{No}$, when $\mathcal{B}_\lambda$ is regarded as a birthday-enriched ordered ring. Its image is $H_{\kappa(\lambda)}$, where now $\kappa(\lambda)$ is the least infinite cardinal $\geq \lambda$. In particular*

$$\mathcal{I}(\mathbf{No}_{<\omega}; +, -, \cdot, <, b) \cong (H_\omega, \in),$$

the hereditarily finite sets, and $\mathcal{I}(\mathcal{B}_{\omega^\omega}) \cong (H_{\omega_1}, \in)$.

Proof. Every formula of the interpretation uses only the ordered-ring operations and birthday. The bounds $\Lambda(\theta)$ and $\theta + \epsilon + 1$ remain in the cutoff by ring closure, and an infinite ring cutoff is a limit ordinal, so the ordinal codes also fit. All later coding and collapse arguments use only these facts and the cardinal-ceiling calculation. For $\lambda = \omega$ every domain is finite, all codes for its subsets and finite isomorphisms have finite length, and the represented transitive closures are exactly the finite ones. $\square$

Real closedness matters for the comparison with decidable field theories and for the surcomplex transfer, but it is not what enables the set coding. The decisive ingredients are complete sign layers, definable reading of those signs, and enough ordinal arithmetic to fit relation codes below the cutoff. Only the forward interpretation is claimed at ring cutoffs that are not epsilon numbers. At ω the cutoff is the dyadic ring, not a field, and it would be incorrect to reuse any real-closed-field argument there. Arbitrary ordinal cutoffs that are not ring cutoffs are not classified.

Part II

Consequences inside one universe

9 The exact elementary-inclusion spectrum

The elementary-inclusion result is stronger than the observation that a birthday expansion is not governed by real-closed-field quantifier elimination. It identifies exactly which initial inclusions are possible.

Lemma 9.1 (Noncardinal lower cutoffs never include elementarily; sources 02, 09). *If $\lambda < \eta$ are epsilon numbers and λ is not a cardinal, then $\mathcal{B}_\lambda \not\prec \mathcal{B}_\eta$.*

Proof. Choose a code C in $\mathcal{B}_\lambda$ for its visible-ordinal collector. Its collapse is λ , and it satisfies $\text{Coll}(C)$ there. By coherence it still collapses to λ in $\mathcal{B}_\eta$. But λ itself is now a scalar ordinal, since $\lambda < \eta$. Its interpreted ordinal code is visible, and is not a member of the set coded by C , because $\lambda \notin \lambda$. Thus $\text{Coll}(C)$ is false in the larger structure. This single formula, with the three entries of C as parameters, witnesses failure of elementarity. $\square$

Source 02’s witness is the weaker formula “the coded set contains every literal ordinal”, which is also true in $\mathcal{B}_\lambda$ and false in $\mathcal{B}_\eta$ for the same code. As source 02 observes, $\mathcal{B}_{\varepsilon_0}$ and $\mathcal{B}_{\varepsilon_1}$ interpret the same H_{ω_1} , yet their inclusion is not elementary: interpretation does not discard the additional first-order information of the original structure.

Lemma 9.2 (A cardinal lower cutoff cannot include elementarily into a noncardinal cutoff). *If λ is a cardinal and η is not, then $\mathcal{B}_\lambda \not\prec \mathcal{B}_\eta$.*

Proof. The parameter-free sentence $\exists C \text{ Coll}(C)$ is false in the first structure and true in the second, by [theorem 6.2](#). $\square$

Theorem 9.3 (Classification of elementary initial inclusions). *For epsilon numbers $\lambda < \eta$,*

$$\mathcal{B}_\lambda \prec \mathcal{B}_\eta \iff \lambda, \eta \text{ are cardinals and } H_\lambda \prec H_\eta.$$

Proof. The preceding two lemmas force both cutoffs to be cardinals; they are uncountable, since no countable cardinal is an epsilon number. For cardinal cutoffs, an elementary inclusion of the surreal structures induces an elementary inclusion of their interpreted set structures. By [proposition 6.6](#) this is the actual inclusion $H_\lambda \subseteq H_\eta$. Conversely, an elementary inclusion of those set structures induces an elementary inclusion of the uniformly interpreted sign structures. The reverse interpretation and its definable sign-graph isomorphism identify this with the original inclusion $\mathcal{B}_\lambda \subseteq \mathcal{B}_\eta$. $\square$

Remark 9.4 (Source 02’s open question, answered). Source 02 proved the forward implication (its necessary conditions) and asked whether they could be sharpened to a sufficient criterion, noting that any proof would have to control the reverse interpretation locally. Source 09’s [theorem 9.3](#) is that criterion: the reverse interpretation of [theorem 6.3](#) and the coherence of [proposition 6.6](#) give the converse. Between cardinal cutoffs the equivalence is also proved by source 05 (regular cardinals) and sources 06 and 07 (all uncountable cardinals).

Corollary 9.5 (Two kinds of elementary extension). *The ordered field inclusion $\mathbf{No}_{<\lambda} \subseteq \mathbf{No}_{<\eta}$ is elementary for all epsilon numbers $\lambda < \eta$. Adding birthday changes its elementary status according to [theorem 9.3](#). In particular,*

$$\mathbf{No}_{<\varepsilon_0} \prec \mathbf{No}_{<\omega_1}, \quad \mathcal{B}_{\varepsilon_0} \not\prec \mathcal{B}_{\omega_1}.$$

Proof. Both fields are real closed, so ordered-field quantifier elimination gives the first assertion. The birthday assertions are instances of the theorem. $\square$

Despite the last failure, both structures in that example are separately bi-interpretable with H_{ω_1} . There is no contradiction: a bi-interpretation does not make *every particular inclusion* elementary. The smaller structure defines an interpreted cutoff set; that formula changes its meaning under the displayed inclusion.

Example 9.6 (Two field-equivalent cutoffs separated by one sentence; sources 05, 07). Let χ say that every set admits an injection into ω (defined as the least infinite ordinal). Then $H_{\omega_1} \models \chi$ and $H_{\omega_2} \models \neg\chi$, since $\omega_1 \in H_{\omega_2}$. Hence the parameter-free sentence $\chi^{\mathcal{I}}$ gives $\mathcal{B}_{\omega_1} \not\equiv \mathcal{B}_{\omega_2}$, although their ordered-field reducts are elementarily equivalent and the field inclusion is elementary.

Corollary 9.7 (Transfer of elementary equivalence; source 06). *For uncountable cardinals κ, λ , $\mathcal{B}_\kappa \equiv \mathcal{B}_\lambda \iff H_\kappa \equiv H_\lambda$. The complete theories $\text{Th}(H_\kappa, \in)$ and $\text{Th}(\mathcal{B}_\kappa)$ are computably many-one reducible to each other by translations independent of κ (source 05, at regular cardinals).*

Proof. Translate sentences through the uniform interpretations in both directions; the translations are effective syntactic operations. $\square$

Remark 9.8 (Bi-interpretation versus elementary equivalence; source 05). Distinct membership structures can be bi-interpretable without being isomorphic or elementarily equivalent in their own language; Freire and Hamkins give relative-consistency instances involving H_{ω_1} and H_{ω_2} [10]. [Example 9.6](#) does not contradict this: the theorem here supplies one uniform interpretation between each cutoff and its own hereditary-size structure, and does not classify all interpretations between two different hereditary-size structures.

Corollary 9.9 (Comparison with the full class). *For an epsilon number λ , interpreted formula by formula,*

$$\mathcal{B}_\lambda \prec \mathcal{B} \iff \lambda \text{ is a cardinal and } H_\lambda \prec V.$$

Proof. The same collector obstruction excludes a noncardinal lower cutoff. For a cardinal lower cutoff, the uniform interpretation and reverse sign-graph interpretation extend to definable classes ([theorem 18.1](#)). The induced set map is $H_\lambda \subseteq V$, so the elementary equivalence follows for each formula. This uses no internal satisfaction class for V . $\square$

The condition $H_\lambda \prec H_\eta$ in [theorem 9.3](#) is not replaced here by a conjectural large-cardinal characterization. It is an exact reduction of one elementary-inclusion problem to another familiar one, together with the complete exclusion of every noncardinal initial cutoff. If both cutoffs are strongly inaccessible, $H_\rho = V_\rho$ and the condition reads $V_\lambda \prec V_\eta$ (source 05).

9.1 Elementary embeddings

Sources 05 (regular cardinals) and 06 (all uncountable cardinals) prove that the uniform bi-interpretation also transfers elementary embeddings that are not inclusions. We state the theorem at uncountable cardinals and give source 05’s explicit formula. Write $\text{name}(x) = \langle b(x), A_x \rangle$ with $A_x = \{\alpha < b(x) : x(\alpha) = +\}$ for the set-theoretic sign name of x .

Theorem 9.10 (Elementary-embedding correspondence; sources 05, 06). *For uncountable cardinals κ, λ there is a canonical bijection*

$$\text{Emb}_{\text{el}}((H_\kappa, \in), (H_\lambda, \in)) \longleftrightarrow \text{Emb}_{\text{el}}(\mathcal{B}_\kappa, \mathcal{B}_\lambda),$$

respecting identities and composition. For $j : H_\kappa \rightarrow H_\lambda$ the corresponding map e_j is given by

$$\text{name}(e_j(x)) = j(\text{name}(x)) = \langle j(b(x)), j(A_x) \rangle,$$

and for $e : \mathcal{B}_\kappa \rightarrow \mathcal{B}_\lambda$ the corresponding j_e is $j_e(x_C) = x_{e(C)}$, with e applied coordinatewise to codes. In particular $e_j(\alpha) = j(\alpha)$ on ordinals and $e_j(x)(j(\alpha)) = x(\alpha)$ for $\alpha < b(x)$.

Proof. Sign names, their order, birthday and arithmetic are uniformly defined in the membership language (section 6), so an elementary j maps sign names to sign names and preserves every translated formula; this proves elementarity of e_j . Since the code domain and $\sim$ are parameter-free, an elementary e maps codes to codes and equivalent codes to equivalent codes, so j_e is well defined, and applying $\mathcal{I}$ to each membership formula proves it elementary. The definable round-trip isomorphisms commute with elementary maps, because an elementary map preserves their defining formulas and their unique outputs modulo the equivalence relations; hence $e_{j_e} = e$ and $j_{e_j} = j$, and composition and identities are respected. The name of an ordinal α is $\langle \alpha, \alpha \rangle$, which gives $e_j(\alpha) = j(\alpha)$, and elementarity of Bit gives the last identity. $\square$

Remark 9.11 (Do not replace $j(A)$ by the pointwise image). For a nontrivial elementary set embedding, $j(A)$ need not equal $j^{\text{“}}A$. The formula uses the actual image *set* $j(A)$: the sign sequence $e_j(x)$ has domain $j(b(x))$ and may have coordinates outside $j^{\text{“}}b(x)$, which a pointwise transport of signs would leave undetermined.

Corollary 9.12 (Critical ordinals and the identity case; source 05). *Corresponding embeddings move exactly the same ordinals; when a least moved ordinal exists it is the same for both. An elementary embedding $\mathcal{B}_\kappa \rightarrow \mathcal{B}_\lambda$ fixing every ordinal below κ is the canonical inclusion.*

Proof. If e fixes all ordinals, then $b(e(x)) = e(b(x)) = b(x)$, and elementarity of the bit formula gives the same signs below that birthday. $\square$

The correspondence is a classification by reduction, not an existence theorem for such embeddings: it proves no new large-cardinal consistency statement and does not refute rank-into-rank hypotheses. A proper canonical inclusion moves no ordinal, so “proper embedding” must not be confused with “embedding having a critical point”. Every automorphism of $\mathcal{B}_\kappa$ is the identity (lemma 19.3); that statement overlaps the collection’s existing simplicity rigidity and is not counted as new.

10 Birthday growth and set-existence axioms

10.1 Birthday eternity

Definition 10.1 (The eternity scheme Et; source 09). For every formula $\varphi(a, x, \vec{p})$ in the birthday-field language, include the assertion that if γ is an embedded ordinal and

$$\forall a < \gamma \exists! x \varphi(a, x, \vec{p}),$$

then some embedded ordinal ϱ satisfies

$$\forall a < \gamma \forall x (\varphi(a, x, \vec{p}) \longrightarrow b(x) < \varrho).$$

The tuple $\vec{p}$ consists of arbitrary parameters from the structure.

This is a precisely specified version of birthday boundedness for definable ordinal-indexed functions. The analogy with the eternity principle in the Chen–Hamkins–Yang announcements is intentional; the cutoff classification below is proved directly, rather than imported from an unexpanded axiom scheme in the slides.

Theorem 10.2 (Eternity forces cardinality of an epsilon cutoff). *If λ is a noncardinal epsilon number, then $\mathcal{B}_\lambda \not\models \text{Et}$.*

Proof. Put $\mu = |\lambda|$, so $\mu < \lambda$ and $\kappa(\lambda) = \mu^+$. Choose a bijection $f : \mu \rightarrow \lambda$. Its graph has hereditary size at most μ , so it belongs to H_{μ^+} and has a code in $\mathcal{B}_\lambda$. Fix one such code as a parameter. Using the interpreted graph of f and the ordinal bridge, a single formula now defines a scalar map on the actual ordinal μ : at $a < \mu$ its value is the embedded ordinal $f(a) < \lambda$. Its birthdays exhaust λ , hence have no bound below λ . This violates the corresponding instance of Et. $\square$

At a singular cardinal κ , an external cofinal map into κ generally has hereditary size κ and need not belong to H_κ . Thus the same argument cannot simply be repeated with $\mu = \text{cf}(\kappa)$. Whether an internally definable cofinal map exists is the point of the next theorem.

10.2 The axioms that all hereditary-size universes already satisfy

Lemma 10.3 (The fixed background axioms). *For every uncountable cardinal κ , H_κ satisfies Extensionality, Empty Set, Pairing, Union, Infinity, Foundation, every Separation instance, and Choice. Moreover the actual transitive closure of each of its elements belongs to H_κ , and its cardinality is correctly computed there.*

Proof. Transitivity gives Extensionality and Foundation. Empty Set, Pairing, Union, and Infinity preserve hereditary size below κ . For any fixed formula and parameters, the externally defined subset of a set $a \in H_\kappa$ satisfying that formula in the set structure H_κ has hereditary size bounded by that of a (up to finitely many additional objects). It therefore belongs to H_κ , proving Separation, including formulas with unbounded quantifiers in that structure.

An ambient choice function for a family $a \in H_\kappa$ has its domain and chosen values inside the transitive closure of a , and its graph has the same infinite hereditary-cardinality bound. It belongs to H_κ . This proves Choice for set families, not a class-choice principle.

Writing $T = \text{TC}(\{x\})$, the transitive closure of T adds at most the set T to its already transitive data, so $T \in H_\kappa$. Its description as the least transitive set containing x as an element is correct

internally: the actual least such set is available there. A bijection between T and its least equipotent ordinal has hereditary size below κ , and any bijection asserted internally is externally a bijection. Minimality among ordinals is therefore also computed correctly. $\square$

10.3 Replacement is exactly eternity at cardinal cutoffs

Theorem 10.4 (Eternity–Replacement equivalence). *For every uncountable cardinal κ ,*

$$\mathcal{B}_\kappa \models \text{Et} \iff H_\kappa \models \text{every instance of Replacement.}$$

Proof. Assume first Replacement in H_κ . A definable scalar function on an ordinal $\gamma < \kappa$ corresponds under the bi-interpretation to a set-theoretically definable function assigning to $a < \gamma$ its unique sign graph. Replacement collects those sign graphs in one set of H_κ . The union of their domains, with one added successor for each length, is bounded below κ : otherwise the transitive closure of the collected set would contain at least κ ordinals. More explicitly,

$$\varrho = \sup\{\text{dom}(s) + 1 : s \text{ is an image sign graph}\} < \kappa$$

bounds their birthdays strictly. This proves the corresponding eternity instance; the empty-domain case is vacuous.

Conversely, assume eternity and consider a formula defining, in H_κ , a function $F : a \rightarrow H_\kappa$, where $a \in H_\kappa$ and the formula has fixed parameters. Choose a bijection $e : \delta \rightarrow a$ with $\delta < \kappa$. Its graph belongs to H_κ . For each $i < \delta$, put externally

$$\nu_i = |\text{TC}(\{F(e(i))\})|,$$

regarded as an initial ordinal. By [lemma 10.3](#), this is a uniquely defined ordinal of H_κ , with the correct external value. The definition of F , the parameter e , and the hereditary-cardinality formula translate into a definable scalar map $i \mapsto \nu_i$ in $\mathcal{B}_\kappa$. Here the ordinal bridge is available for every ordinal below κ . No definable choice of a code for $F(e(i))$ is required: its *hereditary cardinal* is unique.

Eternity gives a $\varrho < \kappa$ bounding all these ordinals. Ambient Replacement forms the actual image set $A = F[a]$, and its transitive closure has cardinality at most

$$\max(\aleph_0, |\delta|, |\varrho|) < \kappa.$$

Indeed it is contained, up to the image set itself, in the union of $|\delta|$ transitive closures, each of cardinality at most $\max(\aleph_0, |\varrho|)$. Thus $A \in H_\kappa$, proving the desired Replacement instance in that structure. This proof uses a fixed cardinal bound below κ ; it never assumes that an arbitrary union of fewer than κ small sets is small. $\square$

Corollary 10.5. *If κ is uncountable and regular, then $\mathcal{B}_\kappa \models \text{Et}$.*

Proof. Any externally given function on an ordinal below κ has fewer than κ birthdays in its image. Regularity bounds them below κ . This argument applies in particular to definable functions. $\square$

This corollary genuinely uses regularity; the equivalence of the theorem does not.

[Added 3 October 2026, batch 89: the same principle on a weak base. Source 10 ([Part VI](#)) proves over Zermelo set theory with Foundation, without Replacement or Choice, that bounding the birthdays of definable set-indexed families of surreals, even strictly inside $(-1, 0)$, is equivalent

to ordinal bounding, and, over an explicit hierarchy axiom, to full Replacement and Collection (theorems 28.2 and 29.1). That is not this theorem: it is an internal scheme equivalence over a weaker base theory, uses only the order of sign sequences, and does not apply to H_κ unless H_κ satisfies that base theory. Remark 24.5 compares the two and records where they combine.]

10.4 Power Set and packing a birthday fiber

Lemma 10.6 (Power Set in a hereditary-size universe; sources 02, 06, 09). *For every infinite cardinal κ ,*

$$H_\kappa \models \text{Power Set} \iff \kappa \text{ is strong limit } (2^\nu < \kappa \text{ for all cardinals } \nu < \kappa).$$

For an individual $\alpha < \kappa$, the actual power set $\mathcal{P}(\alpha)$ is an element of H_κ exactly when $2^{|\alpha|} < \kappa$. Regularity is not required.

Proof. Every subset of a set $a \in H_\kappa$ is already in H_κ , and the internal subset relation is the actual one by transitivity. So an internal power set of a , if it exists, is the full actual power set $\mathcal{P}(a)$. Its hereditary size is $2^{|\alpha|}$ up to harmless finite or countable bounds. This proves necessity (apply it to the cardinals $\nu < \kappa$) and the individual assertion. Conversely, for $x \in H_\kappa$ with κ uncountable, the transitive closure of $\mathcal{P}(x)$ has size at most $\max(\aleph_0, |\text{TC}(\{x\})|, 2^{|x|})$, which is below a strong-limit κ . If $\kappa = \omega$, power sets of hereditarily finite sets are hereditarily finite. $\square$

Theorem 10.7 (The translated Power Set sentence; sources 02, 06). *Let $\text{Pow}^\mathcal{I}$ be the $\mathcal{I}$ -translation of the Power Set axiom $\forall a \exists p \forall x (x \in p \leftrightarrow \forall y (y \in x \rightarrow y \in a))$; it is one fixed parameter-free sentence of the birthday-field language, not a scheme. For every epsilon number λ ,*

$$\mathcal{B}_\lambda \models \text{Pow}^\mathcal{I} \iff \lambda \text{ is a strong-limit cardinal.}$$

Among uncountable regular cutoffs it holds exactly at the strongly inaccessible ones.

Proof. By theorem 5.3 and lemma 10.6, $\text{Pow}^\mathcal{I}$ holds exactly when $\kappa(\lambda)$ is strong limit. If λ is not a cardinal, write $\nu = |\lambda|$; then $\kappa(\lambda) = \nu^+$ and Cantor's theorem gives $2^\nu \geq \nu^+$, so the condition fails. If λ is a cardinal, its ceiling is itself. Adding regularity yields strong inaccessibility. $\square$

Source 02 proves this at every epsilon number, source 06 at uncountable cardinals (its $\text{Pow}^\sharp$). It would be incorrect to remove the words “among uncountable regular cutoffs”: a singular strong-limit cardinal such as $\beth_\omega$ also satisfies $\text{Pow}^\mathcal{I}$, and section 17 shows that the same enriched field can have a regular cutoff in one universe and a singular one in another. The sentence detects the strong-limit property, not regularity, Replacement, or full ZFC.

Definition 10.8 (The fiber-packing principle Pack; source 09). For every embedded ordinal α , there exist a positive embedded ordinal θ and a scalar r , with

$$\chi = \theta + \alpha + 1, \quad \text{b}(r) = \Lambda(\chi),$$

such that

$$\forall u [\text{b}(u) = \alpha \longrightarrow \exists \xi < \theta \forall \zeta < \alpha (\text{Bit}(u, \zeta) \leftrightarrow \text{Bit}(r, p_\chi(\xi, \zeta)))] \quad (10.1)$$

This is one first-order sentence, using the previously defined bit and pairing formulas; it is *not* the Power Set axiom. Its rows list every sign string of birthday α ; repetitions are allowed. Unused coordinates can be ignored. It does not use ordinary ordinal concatenation or a primitive power-set operation.

Theorem 10.9 (Exact packing threshold). *In $\mathcal{B}_\lambda$, the instance of (10.1) for a particular $\alpha < \lambda$ holds exactly when*

$$2^{|\alpha|} < \kappa(\lambda), \quad \text{equivalently, } 2^{|\alpha|} < \lambda,$$

where the cardinal on the left is compared as an ordinal in the second inequality. Consequently Pack holds exactly when λ is an uncountable strong-limit cardinal.

Proof. A packing table with $\theta < \lambda$ rows gives a surjection from θ onto $\mathcal{P}(\alpha)$, so $2^{|\alpha|} \leq |\theta| < \kappa(\lambda)$. Conversely, if that power-set cardinal is $\nu < \kappa(\lambda)$, then $\nu < \lambda$ by cardinal rounding. Choose an enumeration of the fiber by $\theta = \nu$ and encode its table in the square of width $\chi = \theta + \alpha + 1$. The square code stays below λ by closure, so the required r exists. The equivalence of the two inequalities follows because no cardinal lies in $[\lambda, \kappa(\lambda))$.

For a cardinal λ , validity for all $\alpha < \lambda$ is exactly the strong-limit property. If λ is not a cardinal, let $\mu = |\lambda| < \lambda$ and use $\alpha = \mu$. Cantor's theorem gives $2^\mu \geq \mu^+ = \kappa(\lambda)$, so this instance fails. $\square$

Notice that for a noncardinal cutoff λ , the recovered $H_{\kappa(\lambda)}$ and the scalar domain do not have the same ordinal height. Assertions about *all* interpreted ordinals and assertions about only scalar ordinals must be kept separate. The failure at $\alpha = |\lambda|$ is enough to settle the full packing principle.

Cutoff λ	Recovered universe	$\exists C \text{ Coll}(C)$	$\text{Pow}^{\mathcal{T}}, \text{Pack}$
Countable epsilon number	H_{ω_1}	true	false
Noncardinal, $ \lambda = \nu$	H_{ν^+}	true	false
Uncountable successor cardinal κ	H_κ	false	false
Singular strong-limit cardinal κ	H_κ	false	true
Strongly inaccessible κ	$H_\kappa = V_\kappa$	false	true

The table is source 02's, with the packing column added from source 09.

11 Worldly cutoffs and separation examples

11.1 The exact conjunction

Call an uncountable cardinal κ *worldly* here when $V_\kappa \models \text{ZFC}$. This definition does not impose regularity. We need the following familiar set-theoretic fact, for which a proof is included to fix the precise meaning.

Lemma 11.1 (When hereditary size becomes cumulative rank). *For an uncountable cardinal κ ,*

$$H_\kappa \models \text{ZFC} \iff H_\kappa = V_\kappa \text{ and } V_\kappa \models \text{ZFC}.$$

Moreover, $V_\kappa \models \text{ZFC}$ by itself implies $H_\kappa = V_\kappa$.

Proof. Suppose $H_\kappa \models \text{ZFC}$. Its internal cumulative hierarchy exists to every ordinal $\alpha < \kappa$. Every subset of each of its sets belongs to H_κ , so its internal power sets are actual power sets. Induction on α shows that its internal V_α is the actual V_α ; the limit step is ordinary union over the actual smaller ordinals. Thus every V_α for $\alpha < \kappa$ belongs to H_κ , and transitivity gives $V_\kappa \subseteq H_\kappa$. The reverse inclusion is lemma 2.3.

Conversely, assume $V_\kappa \models \text{ZFC}$. For every $\alpha < \kappa$, that model can well-order V_α and supply a bijection with some ordinal $\delta < \kappa$. These are actual sets and actual bijections, because the model is transitive. Hence $|V_\alpha| < \kappa$. The transitive closure of any $x \in V_\kappa$ is contained in some such V_α , and therefore has size below κ . Again $H_\kappa = V_\kappa$ follows. The remaining implication is immediate from equality. $\square$

Theorem 11.2 (Worldly-cutoff characterization). *For an epsilon number λ ,*

$$\begin{aligned} \mathcal{B}_\lambda \models \text{Et} + \text{Pack} &\iff \lambda \text{ is a cardinal and } H_\lambda \models \text{ZFC} \\ &\iff \lambda \text{ is a worldly cardinal.} \end{aligned}$$

For regular uncountable λ , this holds exactly when λ is strongly inaccessible.

Proof. Either growth principle excludes noncardinal epsilon cutoffs. At cardinal cutoffs, [theorem 10.4](#) and [lemmas 10.3](#) and [10.6](#) identify the two missing ZFC requirements as Replacement and Power Set. The equivalence with the rank formulation is [lemma 11.1](#). For regular uncountable cardinals, eternity is automatic and packing is exactly strong limit, giving the usual inaccessible condition. $\square$

This proves [theorem 1.3](#). It does not identify the two formulas with every formulation of a theory called SA. It is an exact theorem about the explicit scheme and sentence defined here.

11.2 The two growth principles do different work

Example 11.3 (Eternity without fiber packing). At $\kappa = \omega_1$, regularity gives eternity. The birthday- ω fiber has $2^{\aleph_0} \geq \aleph_1$ members, so no ordinal below ω_1 can index a packing of it. Hence

$$\mathcal{B}_{\omega_1} \models \text{Et}, \quad \mathcal{B}_{\omega_1} \not\models \text{Pack}.$$

In the decoded structure, every individual real is present, but the set of all reals is not an element of H_{ω_1} .

Example 11.4 (Fiber packing without eternity). Let $\kappa = \beth_\omega = \sup_{n < \omega} \beth_n$, with $\beth_0 = \aleph_0$ and $\beth_{n+1} = 2^{\beth_n}$. This is a strong-limit cardinal: for every $\mu < \kappa$ some $\beth_n$ is at least μ , and then $2^\mu \leq \beth_{n+1} < \kappa$. Thus packing holds.

In H_κ , each finite initial segment of the beth recursion is present and correctly computed. The necessary finite sequence of power sets, cardinals, and bijections has hereditary size below κ , and Power Set is correct there by [lemma 10.6](#). A single formula therefore defines $n \mapsto \beth_n$ on the actual ω . Transporting this formula through the ordinal bridge gives a definable scalar function whose birthdays are cofinal in κ . Consequently

$$\mathcal{B}_{\beth_\omega} \models \text{Pack}, \quad \mathcal{B}_{\beth_\omega} \not\models \text{Et}.$$

The infinite sequence of all these ordinals is precisely what cannot be collected as an element of H_κ .

Source 06 isolates the same phenomenon as one explicit sentence, and uses it to separate $\beth_\omega$ from its successor. Let $\vartheta(n, \gamma)$ be the membership formula saying that a finite sequence of length $n + 1$ begins with ω , ends with γ , and at each step takes the least ordinal equinumerous with the power set of the previous entry; and let

$$\text{Rep}_\beth : [\forall n \in \omega \exists! \gamma \vartheta(n, \gamma)] \longrightarrow \exists y \forall \gamma [\gamma \in y \leftrightarrow \exists n \in \omega \vartheta(n, \gamma)],$$

a single Replacement instance.

Theorem 11.5 (A singular Replacement boundary; source 06). *With $\kappa = \beth_\omega$, $H_\kappa \not\models \text{Rep}_\beth$ and $H_{\kappa^+} \models \text{Rep}_\beth$, the antecedent being true in both. Consequently*

$$\mathcal{B}_{\beth_\omega} \not\prec \mathcal{B}_{\beth_\omega^+},$$

although $\mathcal{B}_{\beth_\omega} \models \text{Pow}^\mathcal{I}$ and the ordered-field inclusion is elementary.

Proof. Every finite beth computation, with the power sets and bijections at its finitely many stages, has hereditary size below κ and belongs to both structures, where it is correctly computed; so $\vartheta(n, \gamma)$ picks out the actual $\beth_n$ in both. A set collecting all $\beth_n$ has every ordinal below κ in its transitive closure, so it is not in H_κ ; the actual set $\{\beth_n : n < \omega\}$ has hereditary size κ and lies in H_{κ^+} . Transfer by [theorem 9.3](#) (or the translation $\text{Rep}_\beth^\mathcal{I}$) gives the failure of elementarity, and [theorem 10.7](#) the Power Set assertion. $\square$

The theorem gives one singular example with failed Replacement. It does *not* show that every singular κ fails Replacement in H_κ : an externally cofinal sequence can fail to be definable over H_κ or fail to be an element of it. Nor does internal ZFC imply external inaccessibility; [corollary 17.2](#) gives a sharp example.

Example 11.6 (A noncardinal field cutoff). Both principles fail at ε_0 . Packing fails at the ordinal ω . Eternity fails by [theorem 10.2](#); one may also use the definable finite power-tower sequence cofinal in ε_0 , with ordinary ordinal exponentiation computed in the decoded H_{ω_1} . Nevertheless the field is real closed and its birthday expansion is bi-interpretable with hereditary countability.

These are actual structures exhibiting independence of the two displayed growth requirements within this family. They are not an assertion of independence from an unspecified list of all the other axioms of SA.

11.3 Singular cutoffs satisfying both principles

Proposition 11.7 (A conditional source of singular worldly cutoffs). *If Ω is strongly inaccessible, there is a cardinal $\kappa < \Omega$ of cofinality ω such that*

$$V_\kappa \prec V_\Omega.$$

In particular, κ is worldly and $\mathcal{B}_\kappa$ satisfies both growth principles.

Proof. Enumerate all existential formulas in the language of set theory. Construct strictly increasing infinite cardinals $\alpha_n < \Omega$. Given α_n , consider the first n formulas and all their finite parameter tuples from V_{α_n} . Whenever V_Ω has a witness, choose one. There are fewer than Ω such requirements, since $|V_{\alpha_n}| < \Omega$. Regularity bounds their witness ranks below Ω . Choose a cardinal $\alpha_{n+1} < \Omega$ above that bound and above α_n .

Set $\kappa = \sup_n \alpha_n$. Regularity of Ω gives $\kappa < \Omega$; it is a cardinal of cofinality ω . Every finite tuple in V_κ appears in one stage, and every existential formula is treated at all sufficiently late stages. Thus every existential statement true in V_Ω with parameters in V_κ has a witness in V_κ . The Tarski–Vaught criterion gives elementarity. Since V_Ω satisfies ZFC, so does V_κ . Apply [theorem 11.2](#). $\square$

The reflection construction is standard set-theoretic reasoning, included as an illustration rather than as a novelty claim, and it needs the stated inaccessible. It demonstrates why replacing “worldly” by “inaccessible” in [theorem 11.2](#) would be wrong. No inaccessible cardinal is assumed in the unconditional cutoff theorems.

12 Model-theoretic consequences

12.1 Full second-order arithmetic and undecidability

Proposition 12.1 (The standard natural numbers and all their subsets; sources 06, 07, 09). *In $\mathcal{B}$ and in every $\mathcal{B}_\lambda$ (λ an epsilon number) there is a parameter-free interpretation of the two-sorted structure $(\mathbb{N}, \mathcal{P}(\mathbb{N}); 0, 1, +, \cdot, <, \in)$ with its full actual subset sort. The number sort consists of the ordinals below the least nonzero limit ordinal w (externally $w = \omega$), the subset sort of the s with $b(s) = w$, and membership is $\text{Bit}(s, n)$.*

Proof. “ q is a nonzero limit ordinal” is $\text{Ord}(q) \wedge q > 0 \wedge \forall a < q \exists d (a < d < q)$ with ordinal guards, and w is its least solution. The ordinals below w are the standard finite ordinals, and natural arithmetic on them is ordinary arithmetic. The fiber $b(s) = \omega$ is in bijection with all subsets of ω by (4.3); no restriction to definable subsets is imposed. $\square$

When working inside a model M , the subset sort is $\mathcal{P}(\omega)^M$, as it must be. Sources 06 and 07 state the proposition, and the next two results, at uncountable cardinals and source 09 at ε_0 ; the proofs use only $\omega < \lambda$, which holds at every epsilon number. This is an ordinary first-order interpretation of a two-sorted structure, not a replacement of first-order by second-order syntax. It does not make set theory definable in the pure ordered-field reduct, where the birthday structure is absent.

Corollary 12.2 (Undecidability; sources 05, 06, 07, 09). *True first-order arithmetic reduces effectively to the complete theory of every $\mathcal{B}_\lambda$, and formula by formula to $\mathcal{B}$. Consequently these complete theories are neither decidable nor computably enumerable.*

Proof. Relativize arithmetic quantifiers to the number sort. If the target theory were computably enumerable, the inverse image would enumerate true arithmetic; a complete computably enumerable theory is decidable, and true arithmetic is not. $\square$

These are logical-complexity consequences, not new algorithms. A code can contain an arbitrary noncomputable subset of an ordinal, and first-order definability in the birthday expansion must not be confused with decidability from a finite symbolic representation.

12.2 Independence and an omitted type

Proposition 12.3 (Independence property and a countable omitted type; source 07). *The formula $\text{Bit}(s, n)$ has the independence property in every $\mathcal{B}_\lambda$. The parameter-free partial type $\{x < w \wedge \text{Ord}(x)\} \cup \{x > n : n \in \mathbb{N}\}$, with each standard n named by its definition, is finitely satisfiable but omitted.*

Proof. For distinct finite ordinals $n_1, \dots, n_k$ and any $S \subseteq \{1, \dots, k\}$, a length- ω sign code positive exactly at the n_j with $j \in S$ realizes the pattern. Finite parts of the type are realized by large finite ordinals; a realization of the whole type would be a standard natural number above all standard natural numbers. $\square$

Remark 12.4 (Saturation of the pure field, and the sibling report). For regular uncountable κ , the pure real closed field $\mathbf{No}_{<\kappa}$ realizes every one-variable type over fewer than κ parameters: by quantifier elimination a nonalgebraic type over the real closed subfield generated by the parameters

is a cut, regularity bounds the birthdays of its fewer than κ endpoints, and (3.1) supplies a separator. Source 07 records this as the usual saturation property of the pure field, not a new result. Naming birthday nevertheless produces the omitted type of [proposition 12.3](#). External saturation of the pure field $\mathbf{No}^M$ inside an outer universe is the subject of the sibling report [foundations-and-computation/surreal-fields-across-universes](#); this report does not study it.

[Added 2 October 2026, batch 80: the order-theoretic case of this remark, which uses neither the field structure nor uncountability, is proved and used in the report [foundations-and-computation/surreal-well-orders](#) (batch-80 manuscripts 08, 09, 11 and 12). Its Lemma 7.2 (`swo:sw:stageeta`, its source 11's) shows, by the birthday bound (3.1), that for every regular infinite κ , $\kappa = \omega$ included, $\mathbf{No}_{<\kappa}$ fills every cut $A < B$ with $|A \cup B| < \kappa$, even avoiding any prescribed set of fewer than κ points; it calls this property η_κ . Its Note 7.4 (`swo:sk:prop:cutoffeta`, its source 08's Proposition 6.1) states it for regular κ with $2^{<\kappa} = \kappa$ and cites this remark. Its Proposition 12.1 (`swo:sw:permutationeta`) shows that the lexicographic order of the well-orderings of length κ of an η_κ -order of cardinality κ is again η_κ ; with Lemma 7.2 this applies to $\mathbf{No}_{<\kappa}$ when $2^{<\kappa} = \kappa$. No result here uses that report.]

12.3 Birthday is not visible in the field reduct

Proposition 12.5 (Nondefinability of birthday; sources 05, 07). *On $\mathbf{No}_{<\lambda}$, for λ an epsilon number, neither birthday nor prefix simplicity is definable, even with parameters, in the ordered-field reduct, nor in any o-minimal expansion of it. The same holds formula by formula on $\mathbf{No}$.*

Proof. If birthday were definable, its fixed-point set, the embedded ordinals, would be definable. It is infinite and contains no interval: between ordinals $\alpha < \gamma$ lies $\alpha + \frac{1}{2}$, which is not an ordinal. An o-minimal structure (in particular a real closed field, by quantifier elimination) has only finite unions of points and intervals as definable subsets of the line. If prefix simplicity were definable, the ordinals would be defined by “every proper prefix of x lies below x ”, with the same contradiction. $\square$

Source 05 states the proposition at regular cardinals and source 07 at uncountable cardinals; the proof uses only real closedness (o-minimality). The collection's foundations report notes that the field language has no birthday symbol; this proposition is the corresponding standard nondefinability proof, new to the collection but not a novelty claim.

12.4 How little arithmetic is needed initially

Let $G_\varpi(\alpha, \gamma, \rho)$ be the ternary relation on $\mathbf{No}_{<\kappa}$ saying that α, γ, ρ are ordinals and $\rho = \varpi(\alpha, \gamma)$, with ϖ the polynomial of [lemma 4.5](#).

Corollary 12.6 (Reconstruction from ordinal pairing; source 06). *For every uncountable cardinal κ , all operations and constants of $\mathcal{B}_\kappa$ are uniformly definable in $(\mathbf{No}_{<\kappa}; <, b, G_\varpi)$. Hence the two structures are definitionally equivalent.*

Proof. Run the coding of [section 5](#) with relation words of length $\varpi(\delta, \delta) + 1$ whose plus positions are the $\varpi(u, v)$, as source 06 does; ordinal successor is definable from the ordinal order. The code formulas then use field arithmetic only to evaluate ϖ on ordinals, which G_ϖ supplies, so this reduct still interprets H_κ . Translate the arithmetic definitions of [section 3](#) through that interpretation

and use the surreal-side round trip to put their graphs on the original carrier. The converse is the polynomial formula for G_ω . $\square$

No minimal-language claim is made: the corollary does not remove birthday data or show that arithmetic is definable from order and prefix simplicity alone. It is partial progress on the minimal-signature questions of sources 02, 05 and 07, which stay open (section 21).

13 The exact Choice boundary of ordinal graph coding

This section, source 06's, works in ZF . It considers the full class of sign sequences and the graph interpretation of section 5 without a cutoff. All its defining formulas remain first-order formulas about individual sets or surreals, and class assertions are read formula by formula.

Definition 13.1. Let $\text{HWO} = \{x : \text{TC}(\{x\}) \text{ admits a well-ordering}\}$. One well-ordering of the whole transitive closure is required, with no definability condition such as the one in HOD.

Theorem 13.2 (Choice-sensitive range; source 06). *Over ZF , the pointed ordinal-graph interpretation has quotient structure isomorphic to $(\text{HWO}, \in)$. Its external collapse map is surjective onto V if and only if the Axiom of Choice holds.*

Proof. The Mostowski collapse of a valid code (δ, r, t) is injective with domain the ordinal δ , and its range is $\text{TC}(\{x_C\})$ by root generation; transporting the ordinal order well-orders that closure. Conversely, a well-order of $\text{TC}(\{x\})$ gives a bijection from an ordinal, along which membership pulls back to a valid code for x ; no simultaneous choice for a family is used. Equality and membership remain correct in ZF : all subsets of a fixed ordinal exist and have characteristic words, so the well-foundedness test still quantifies over every subset, and Mostowski collapse is a theorem of ZF . If Choice holds every transitive closure is well-orderable. Conversely, surjectivity well-orders every $\text{TC}(\{x\})$ and hence every $x \subseteq \text{TC}(\{x\})$, and the well-ordering principle implies Choice. $\square$

Proposition 13.3 (Internal Choice is a different assertion; source 06). *In ZF , HWO is transitive, contains every ordinal and every subset of an ordinal, and internally satisfies the statement that every set can be well-ordered. If $\mathcal{P}(\omega)$ is not well-orderable in the ambient universe, $(\text{HWO}, \in)$ fails the Power Set axiom at ω .*

Proof. If $y \in x \in \text{HWO}$ then $\text{TC}(\{y\}) \subseteq \text{TC}(\{x\})$ is well-orderable. For $a \subseteq \alpha$, $\text{TC}(\{a\}) \subseteq \{a\} \cup \alpha$. For $x \in \text{HWO}$ the restriction to x of a well-order of $\text{TC}(\{x\})$, and every subset of x , lie in HWO, since finite products and finite subsets of a well-orderable set are well-orderable in ZF ; so the internal well-order has a witness. Every subset of ω is in HWO, so an internal power set of ω would be the actual $\mathcal{P}(\omega)$, which is then well-orderable, contrary to the hypothesis. $\square$

The surjectivity statement of theorem 13.2 is an external comparison of ranges, not the value of the internal Choice sentence. The theorem is about the specified ordinal-graph construction; no claim is made that every possible interpretation of sets in surreals has this boundary. Global Choice is unnecessary here and for the local ZFC bi-interpretations: existence of a code is proved object by object and the quotient identifies all codes.

Remark 13.4 (Source 09's fourth target, answered). Source 09 listed “the choice-free analogue” among its targets and observed that ordinal-indexed relations naturally recover the sets with well-orderable transitive closures. Theorem 13.2, source 06's, is that analogue for the full class; source 07 records

the same range in a remark without a separate proof. Whether a richer coding signature can escape the boundary, and a choice-free version of the *cutoff* theorems, remain open.

[Added 2 October 2026, batch 80: a class-level counterpart. The report `foundations-and-computation/surreal-well-orders` proves in its Theorem 13.1 (`swo:cf:choice`, its source 11's) that over GB without set choice, Global Choice holds if and only if **No** has a class well-order, if and only if it has a set-like one, if and only if there is a class bijection $\text{Ord} \rightarrow \mathbf{No}$. From a class well-order of the sign sequences, the proof well-orders every V_α by set-valued recursion, coding a subset of V_β by a sign sequence along the well-order of V_β already built. Its Remark 13.4 (`swo:rem:answered-choice`) compares this with the ordinal-graph coding of [theorem 13.2](#) and notes that the theorem here concerns a different statement and is unchanged. The two open problems just named are not affected.]

[Added 3 October 2026, batch 89: an injection counterpart over ZF. Source 11 ([Part VII](#)) proves that a set X injects into **No** iff it injects into $\mathcal{P}(\alpha)$ for some ordinal α ([theorem 42.1](#)), so that universality of **No** for free fields on all sets is exactly the first Kinna–Wagner principle ([theorem 43.5](#)). The sets injecting into **No** form a wider class than HWO: $\mathcal{P}(\omega)$ always injects into **No** ([remark 40.3](#)). An injection names elements; it does not recover a set as the coding here does. [Theorem 13.2](#) and the two open problems of [remark 13.4](#) are unchanged.]

Part III

Outer models

Throughout this part, $M \subseteq N$ are transitive models of ZFC with the same ordinals. They may be set models viewed from an ambient metatheory, or class models with formulas interpreted one at a time. Superscripts indicate the model in which a construction is performed, and all references to M -cardinality are internal to M . The intended examples are set-forcing extensions, but no result below assumes forcing unless it says so.

14 Old arithmetic and cross-model coherence

Lemma 14.1 (An old simplest separator stays simplest; sources 02, 05, 07). *If $L, R \in M$ are separated sets of old sign sequences and $w \in M$ is their simplest separator computed in M , then w is their simplest separator in N .*

Proof. Comparisons of old sign sequences are absolute, so w still separates. Suppose $z \in N$ separates and is shorter. If z is a prefix of w it is a restriction of w to an ordinal, hence old, contradicting minimality in M . Otherwise z and w first disagree before z ends; their common prefix p is a restriction of w , hence old, lies strictly between z and w , and is a shorter separator by convexity of the set of separators, with all its comparisons with L, R holding in M . This again contradicts minimality. $\square$

Source 02 gives a constructive variant: build the separator sign by sign, appending $+$ if some $\ell \in L$ is at least the current word and $-$ otherwise, and taking unions at limits; every test involves only the given sets and the previous prefix, so the stopping place and every sign agree in M and N .

Proposition 14.2 (Absolute old field structure; sources 02, 05, 07). *The inclusion of old sign sequences $\mathbf{No}^M \subseteq \mathbf{No}^N$ preserves and reflects order and preserves birthday, constants, negation, addition, multiplication and inverses. Its ordered-field reduct is an elementary inclusion, and so is the pure-field inclusion $\mathbf{No}^M[i] \subseteq \mathbf{No}^N[i]$. The same holds for $\mathbf{No}_{<\lambda}^M \subseteq \mathbf{No}_{<\lambda}^N$ at every common epsilon number λ .*

Proof. Order, length, constants and sign reversal are absolute. For addition induct on pairs of old prefixes: all prefixes of an old number are old, the options are identified by induction, and lemma 14.1 identifies their simplest separator. Multiplication follows by the same induction using addition, and inverses by uniqueness. Both real fields are real closed and both complex fields algebraically closed of characteristic zero, so quantifier elimination makes the inclusions elementary [14, 15]. For proper classes the equivalences are applied one formula at a time. $\square$

This is the usual absoluteness of the canonical construction, not an assertion about arbitrary chosen presentations of isomorphic fields. It does not say that the outer universe cannot distinguish the inner one once extra structure is named: the rest of this part shows that birthday does.

Proposition 14.3 (Cross-model coherence; source 09). *Let κ be an uncountable cardinal of both models. There are actual inclusions*

$$\mathcal{B}_\kappa^M \subseteq \mathcal{B}_\kappa^N, \quad H_\kappa^M \subseteq H_\kappa^N,$$

compatible with the interpretations and their definable isomorphisms. Consequently

$$\mathcal{B}_\kappa^M \prec \mathcal{B}_\kappa^N \iff H_\kappa^M \prec H_\kappa^N.$$

Proof. A sign sequence of M remains the same sign sequence in N ; order and birthday are absolute. A small arithmetic certificate existing in M remains valid in N , so the field operations agree. An old set of hereditary size below κ comes with an old bijection witnessing that bound; since κ is still a cardinal, it belongs to H_κ^N .

For code validity, one must not claim that the two models have the same subsets of the coding ordinal. Instead, a relation well founded in M has an ordinal rank function in M , which remains decreasing in N . Its well-foundedness therefore persists despite the extra subsets in N . Root generation says that every node is connected to the root by a finite downward path, as established in lemma 5.1; these paths are absolute. Extensionality is bounded and absolute. Failure of any of the code conditions in M also persists, using its old witness. Thus validity on old codes is unchanged.

The old Mostowski collapse still satisfies its bounded defining equations in N and hence is the same collapse. Equality and membership of decoded old sets are unchanged. Witnesses to their equality and membership can be constructed in M from those collapses, so their required graph codes already exist there. The reverse sign-graph interpretation likewise sends old objects to their actual images. Applying each interpretation to an elementary inclusion now proves the equivalence. $\square$

Unlike two cutoffs in a single universe, $\mathcal{B}_\kappa^M$ and $\mathcal{B}_\kappa^N$ may have different birthday fibers even at the same small ordinal. Their ordered field reducts nevertheless always include elementarily (proposition 14.2).

15 The preservation criterion and the first new birthday

Theorem 15.1 (Exact preservation criterion; sources 02, 06, 07, 09). *For every common ordinal γ ,*

$$(\mathbf{No}_{<\gamma})^M = (\mathbf{No}_{<\gamma})^N \iff \forall \alpha < \gamma \ P(\alpha)^M = P(\alpha)^N. \quad (15.1)$$

If $\gamma = \lambda$ is an epsilon number, these conditions are equivalent to the literal equality $\mathcal{B}_\lambda^M = \mathcal{B}_\lambda^N$ of the canonically presented structures, including all operations and the birthday function. If $\gamma = \kappa$ is an uncountable cardinal in both models, they are also equivalent to $H_\kappa^M = H_\kappa^N$.

Proof. Taking plus positions is an absolute bijection between sign sequences of length α and subsets of α ; this gives (15.1). Once the carriers agree, proposition 14.2 gives equality of all operations; conversely, equal structures have equal carriers.

For H_κ : a subset of $\alpha < \kappa$ has hereditary size below κ in either model, so $H_\kappa^M = H_\kappa^N$ implies that no bounded subset is new. Conversely assume no new bounded subsets. Old sets of hereditary size below κ stay so in N (proposition 14.3). For $x \in H_\kappa^N$ take in N a valid code (δ, r, t) with $\delta < \kappa$. Its relation word has length $\Lambda(\delta) < \kappa$, so it is old, as are δ and t . The coded relation is well founded in N , hence in M (an old nonempty subset without a minimal element would also be one in N); extensionality is absolute; the collapse computed in M satisfies its recursion in N , so it is the collapse there. Hence $x \in M$, with hereditary size below κ witnessed in M . $\square$

Source 07 proves (15.1) for every common ordinal, source 02 the epsilon-number clause, source 06 the cardinal clause with H_κ , and source 09 the structure equality inside theorem 16.4. The right side of (15.1) must *not* be replaced by “no new short sequences of ordinals”: a short sequence with unbounded range can be new while every bounded subset below a large cutoff stays old, which is exactly what the Prikry example of section 17 exploits. Nor is “adds no real” identified with “adds no surreal” (example 15.7).

Corollary 15.2 (Countable cutoffs detect exactly new reals; source 02). *If λ is a countable epsilon number, then $\mathcal{B}_\lambda^M = \mathcal{B}_\lambda^N \iff \mathcal{P}(\omega)^M = \mathcal{P}(\omega)^N$.*

Proof. For infinite $\alpha < \lambda$ a bijection between ω and α lies in M and remains one in N , so a new subset of α gives a new subset of ω ; finite ordinals have no new subsets; and $\omega < \lambda$. $\square$

Lemma 15.3 (New sets imply a new ordinal subset; sources 05, 07). *If $M \subsetneq N$, then N has a subset of some ordinal that is not in M .*

Proof. Suppose not, and let $a \in N$. In N enumerate $\text{TC}(\{a\})$ by an ordinal and code the pulled-back membership relation by a subset of an ordinal through an old injective pairing (for example the natural-arithmetic rectangle map). By hypothesis the code, hence the relation, is in M . Well-founded in N , it is well founded in M ; its decoration computed in M is the decoration in N by uniqueness of recursion, and its root value is a . So $a \in M$, and $M = N$. $\square$

Theorem 15.4 (First new birthday; sources 02, 05, 07). *Let $M \subsetneq N$ and*

$$\beta = \beta(M, N) = \min\{\alpha : \mathcal{P}(\alpha)^N \neq \mathcal{P}(\alpha)^M\}.$$

Then β exists and:

- (i) *every surreal of N of birthday below β belongs to M ;*

- (ii) N has a surreal of birthday exactly β not in M , all of whose proper prefixes are in M ; thus β is the least birthday of an element of $\mathbf{No}^N \setminus \mathbf{No}^M$;
- (iii) β is an infinite cardinal of M .

Proof. β exists by lemma 15.3. A sign sequence of birthday α is determined by α and its plus set, which is old when $\alpha < \beta$; this proves (i), and a new subset of β gives the new sign of (ii), whose proper prefixes are old by (i). No finite ordinal has a new subset. If β were not an M -cardinal, M would have a bijection $f : \nu \rightarrow \beta$ with $\nu < \beta$; for a new $A \subseteq \beta$ the pullback $f^{-1}[A]$ would be old by minimality, and then $A = f[f^{-1}[A]]$ would be old. $\square$

Remark 15.5 (Source 02's version). Source 02 proved the theorem for a set-forcing extension $N = M[G] \neq M$, obtaining existence of β from the generic filter: a ground-model well-order of the forcing set turns G into a new subset of an ordinal. Sources 05 and 07 remove the forcing hypothesis by lemma 15.3. No claim that all cardinals are preserved is needed. The theorem does not assert that β is regular, and it does not identify the signs of birthday at most ω with real numbers.

Example 15.6 (Adding a real). If $N = M[g]$ adds a Cohen subset g of ω , no finite subset is new and g is, so $\beta = \omega$. The sign with plus set g is a new surreal of birthday ω , and each of its finite prefixes is old. Such an extension changes $\mathbf{No}_{<\kappa}$ for every common uncountable cardinal κ .

Example 15.7 (No new reals, but new surreals; sources 05, 07). Force over M with countable partial functions from ω_1^M to 2, ordered by extension. The union of a descending ω -chain is a condition, so the forcing is countably closed; the successive-decision argument shows that it adds no new countable sequence of ordinals, hence no subset of any old countable ordinal, and preserves ω_1 . The generic union differs from every old subset of ω_1^M by a density argument [13]. Hence $\beta = \omega_1^M$: all countable-birthday surreals are unchanged, there are new signs of birthday ω_1^M , and the full birthday inclusion still fails elementarity (theorem 16.5). Agreement on the reals, or even on every countable sign sequence, does not exhaust what the birthday expansion remembers.

Remark 15.8 (The sibling report). The sibling report `foundations-and-computation/surreal-fields-across-universes` works in the same setting of transitive $M \subseteq N$ with the same ordinals and uses the same letter β for the first new birthday; it writes δ for the first new ordinal-sequence length, which is not named here. It studies the external saturation of the *pure* ordered field $\mathbf{No}^M$ inside N , which is elementary in $\mathbf{No}^N$ by proposition 14.2; the present report studies the birthday expansion, whose inclusion is non-elementary as soon as $M \neq N$ (theorem 16.5). The first-new-birthday theorem and the absoluteness of old arithmetic are shared ground; the countably closed and Prikry examples appear in both.

Remark 15.9 (Gap pairs of the cutoff fields; batch 37). Added in batch 37 (28 September 2026). Part VIII of the sibling report computes the external gaps of this report's cutoff fields as *pure* ordered fields (no birthday symbol). Let $M \subseteq N$ have the same ordinals and the same cofinalities, and let θ be a common uncountable regular cardinal. By its Lemma 26.1 (`univ:gs:lem:boundary`), the gap pairs of $(\mathbf{No}_{<\theta})^M$ in N are exactly the symmetric pairs (μ, μ) , for μ the N -cofinality of the length of a fresh sign (a new sign sequence all of whose proper initial segments are old) of length below θ , together with the boundary pairs (μ, θ) and (θ, μ) for every infinite regular $\mu \leq \theta$, which come from old signs of length θ . Pairs of the first kind, all with $\mu < \theta$, are absent exactly when $(\mathbf{No}_{<\theta})^M = (\mathbf{No}_{<\theta})^N$, that is, by theorem 15.1, when no subset of an ordinal below θ is new. With $M = N$ this shows that the internal field $\mathbf{No}_{<\theta}$ already has asymmetric gaps such as (ω, θ) . This is consistent with remark 12.4, since each such gap has a side of cofinality θ . The sibling's Theorem 26.2 and Corollary 26.3 (`univ:gs:thm:cutoff`, `univ:gs:cor:setfields`) use these fields,

at $\theta = \aleph_{\omega+2}$, for $2^{\aleph_0}$ pairwise nonisomorphic real closed fields of one cardinality with one saturation spectrum.

[Added 3 October 2026, batch 81: the case $M = N$ holds for every infinite cardinal. Source 17 of the report `foundations-and-computation/surreal-well-orders` computes the gap pairs of the ordered set $\mathbf{No}_{<\theta}$ for every infinite cardinal θ , including singular θ and $\theta = \omega$: with $r = \text{cf } \theta$ they are (r, r) and (ρ, r) , (r, ρ) for every infinite regular $\rho < \theta$, and deleting fewer than $2^{<\theta}$ points does not change them (its Theorem 29.4, `swo:gs:thm:alphabet-gaps`, with the same witnesses, ρ pluses followed by θ minuses). At uncountable regular θ this is the case $M = N$ of `univ:gs:lem:boundary` above, which that report records with a pointer.]

16 Detecting new subsets

16.1 The old-power-set detector at a cutoff

Theorem 16.1 (Uniform detector for an added subset; source 09). *Let κ be an uncountable cardinal of both models. Suppose $\alpha < \kappa$, N has a subset of α not in M , and*

$$\nu = (2^{|\alpha|})^M < \kappa.$$

Then $\mathcal{B}_\kappa^M \not\equiv \mathcal{B}_\kappa^N$. One fixed interpreted formula witnesses the failure using parameters coding α and $A = \mathcal{P}^M(\alpha)$.

Proof. The cardinal bound gives $A \in H_\kappa^M$. In the membership language consider

$$\exists y [\forall v \in y (v \in \alpha) \wedge y \notin A]. \quad (16.1)$$

This is false in H_κ^M , because all its subsets of α lie in A . It is true in H_κ^N : any newly added subset of α has hereditary size below the still-cardinal κ and is not an element of the old set A . Translate (16.1) through $\mathcal{I}$ and choose old codes for the two parameters. Cross-model coherence gives the claimed failure in the birthday structures. $\square$

The membership-language formula (16.1) is Σ_1 : its matrix has only bounded quantifiers. Its fully expanded birthday-field translation need *not* be existential or Σ_1 in that different language, because code validity includes the bit-quantified well-foundedness test. No quantifier-complexity claim is made across that translation.

Proposition 16.2 (Sharpness of the detector's parameter cost; source 09). *The old parameter $\mathcal{P}^M(\alpha)$ belongs to H_κ^M exactly when $(2^{|\alpha|})^M < \kappa$. In the rooted coding scheme, every code for that parameter has node-domain cardinality $|\text{TC}(\{\mathcal{P}^M(\alpha)\})|^M$, which equals the power-set cardinal for infinite α . In that infinite case, a code of birthday cardinality no larger than that cardinal can be chosen.*

Proof. The first statement is lemma 10.6 inside M . The exact node-domain cardinality is (5.7). For infinite α , the transitive closure consists of the power set, its subsets, and the ordinal data beneath α , giving cardinality $2^{|\alpha|}$. Choosing its initial ordinal as the node domain gives relation birthday $\Lambda(\nu)$, of cardinality ν by lemma 4.3. $\square$

This is not a theorem that elementarity must hold when the bound fails. Another formula can detect a change at a smaller cutoff. It says exactly when this particular, universally applicable old-power-set witness is available.

Example 16.3 (A conditional continuum-sized illustration). Suppose M satisfies CH, N preserves ω_2^M , and N adds a real. Then the old set of reals has hereditary size ω_1^M , so the detector works at the cutoff ω_2^M . That old set is not an element of $H_{\omega_1^M}^M$, so the same parameter cannot be used at the cutoff ω_1^M . No claim of elementarity at the smaller cutoff follows.

16.2 Rigidity under extensions at a strong limit

Theorem 16.4 (Strong-limit extension rigidity; source 09). *Let κ be an uncountable cardinal of both models, strong limit in M . Then*

$$\begin{aligned} \mathcal{B}_\kappa^M \prec \mathcal{B}_\kappa^N &\iff \mathcal{B}_\kappa^M = \mathcal{B}_\kappa^N \\ &\iff \mathcal{P}^M(\alpha) = \mathcal{P}^N(\alpha) \quad \text{for every } \alpha < \kappa. \end{aligned}$$

Proof. If some subset of an $\alpha < \kappa$ is new, the strong-limit property makes its old power-set parameter small enough, and [theorem 16.1](#) excludes elementarity. Conversely, if all these power sets agree, the structures are equal by [theorem 15.1](#), hence the inclusion is elementary. Equality of the scalar domains forces equality of the corresponding plus-sets, proving the remaining direction. $\square$

This proves [theorem 1.4](#). It applies, for example, to any forcing extension satisfying the stated cardinal-preservation hypotheses, but uses no particular forcing construction. It is a theorem about *standard cutoff structures at a fixed common cardinal*, not a prohibition on abstract elementary extensions supplied by model theory. Source 02 explicitly declined to claim that a proper enlargement at a fixed cutoff is non-elementary merely because it is proper; the theorem above and [theorem 16.1](#) give sufficient conditions, and nothing more is claimed. Their ordered field reducts still include elementarily even when the birthday expansion fails the detector; a set-theoretic change can be invisible to ordered-field formulas yet visible to one birthday-field formula with old parameters.

16.3 The full class, and an explicit row-search formula

For the full classes write $\mathcal{B}^M = (\mathbf{No}^M; 0, 1, +, -, \cdot, <, b)$ and likewise $\mathcal{B}^N$; by [proposition 14.2](#) the old signs form a substructure.

Theorem 16.5 (An outer universe is detectable; sources 05, 07). $\mathcal{B}^M \prec \mathcal{B}^N \iff M = N$. *When $M \neq N$ the failure is witnessed with old parameters: by the $\mathcal{I}$ -translation of (16.1) with codes for α and $\mathcal{P}(\alpha)^M$, for any α with a new subset (source 07), and by the formula New of [theorem 16.6](#) (source 05).*

Proof. If $M \neq N$, [lemma 15.3](#) gives an ordinal α with a new subset. Old valid codes remain valid in N and decode to the same sets (as in [proposition 14.3](#), without a cardinal bound), and (16.1) is false in M and true in N with those parameters. $\square$

Source 05's detector reads the birthday language directly. Fix a nonzero ordinal θ and in M an enumeration $(A_j)_{j < \nu}$, $\nu > 0$, of $\mathcal{P}(\theta)^M$ (a surjection suffices), and code it by one old sign m of birthday $\Lambda(\theta + \nu + 1)$ whose bit at $p_{\theta+\nu+1}(j, \xi)$ is plus exactly when $\xi \in A_j$. Let

$$\text{New}(m, \theta, \nu) : \quad \exists x \left[b(x) = \theta \wedge \forall j < \nu \exists \xi < \theta \neg (\text{Bit}(x, \xi) \leftrightarrow \text{Bit}(m, p_{\theta+\nu+1}(j, \xi))) \right],$$

together with the ordinal and length conditions on θ, ν, m ; it asks for a row of length θ different from every old row.

Theorem 16.6 (An explicit new-subset detector; source 05). *With such parameters, $\mathcal{B}^M \models \neg \text{New}(m, \theta, \nu)$, and $\mathcal{B}^N \models \text{New}(m, \theta, \nu)$ if and only if $\mathcal{P}(\theta)^N \neq \mathcal{P}(\theta)^M$.*

Proof. Every sign of birthday θ in M has an old plus set, which is some row A_j . The old sign m has the same bits in N , so it still has exactly the old rows. A new subset of θ gives a witness in N ; conversely a witness has a plus set differing from every old subset, hence new. $\square$

Source 05 writes the matrix with its rectangle map and length $\theta(\nu + 1)$; the formula is the same up to that pairing. The enumeration is a set in M and the matrix one sign of set length; no class of ground-model signs is named. The formula is not claimed to be existential after the abbreviations are expanded, since the prefix and bit predicates have quantifiers of their own. The detector branches off after the pairing lemma and does not need the arithmetic certificates.

Remark 16.7 (The bounded version). At a cutoff the matrix must fit. Source 05 states: if a common ordinal κ is an uncountable regular cardinal in both models, the old matrix has $\theta, \nu < \kappa$ and length below κ , and N has a new subset of θ , then $\mathcal{B}_\kappa^M \not\models \mathcal{B}_\kappa^N$, witnessed by New . Its proof uses regularity only through closure of natural arithmetic below κ , which holds at every uncountable cardinal (lemma 2.2); so the conclusion holds whenever κ is an uncountable cardinal of both models and $(2^{|\theta|})^M < \kappa$, the hypothesis of theorem 16.1. Without a bound on the old enumeration the corollary cannot be inferred from $\theta < \kappa$ alone, because the set of old subsets of a small ordinal may have cardinality at least κ .

Remark 16.8 (The conclusions use parameters). These theorems say that the *canonical inclusion* is not elementary. They do not by themselves show that the two structures have different parameter-free complete theories, nor do they exclude unrelated embeddings or isomorphisms. If the extension changes a parameter-free set-theoretic sentence, its translation does distinguish the theories (sources 05, 07).

17 Prikry forcing: an unchanged field with a singularized cutoff

17.1 The imported forcing theorem

We use the classical ordinary Prikry forcing associated with a normal measure U on a measurable cardinal κ . Conditions consist of a finite increasing stem $s \subset \kappa$ and a measure-one set $A \in U$ above that stem; an extension may lengthen the stem using points of A and shrink A , a direct extension only shrinks A . Its standard properties are: the generic stem union is cofinal of order type ω in κ ; the forcing adds no bounded subsets of κ ; and it preserves cardinals. These are imported from the exposition of Poveda–Rinot–Sinapova, especially Lemma 2.10 and Section 3.1 [11], and are not reproved in full. For orientation, source 02 sketches the bounded-subset consequence of the Prikry property: decide a name for a subset of $\alpha < \kappa$ by successive direct extensions, using that intersections of fewer than κ measure-one sets have measure one; bounded subset preservation keeps the cardinals below κ and κ itself, and the κ^+ -chain condition the cardinals above.

Theorem 17.1 (Forcing invisibility, conditional; sources 02, 06). *Suppose a transitive model M of ZFC has a measurable cardinal κ , and let $N = M[G]$ be an ordinary Prikry extension at κ . Then*

$$\mathcal{B}_\kappa^M = \mathcal{B}_\kappa^N, \quad H_\kappa^M = H_\kappa^N = V_\kappa^M,$$

with the same carrier, functions and relations, although

$$M \models \text{cf}(\kappa) = \kappa, \quad N \models \text{cf}(\kappa) = \omega.$$

The common membership structure satisfies ZFC in both universes.

Proof. In M the measurable κ is strongly inaccessible, hence an epsilon number by lemma 2.2; it remains a cardinal, hence an epsilon number, in N , although singular there. Prikry forcing adds no subset of any $\alpha < \kappa$, so theorem 15.1 gives both equalities of structures. Since κ is inaccessible in M , $H_\kappa^M = V_\kappa^M$ (sets in V_κ have small transitive closures by strong limit and regularity, and a well-founded relation on fewer than κ nodes has rank below κ), and $V_\kappa^M \models \text{ZFC}$. The generic sequence witnesses $\text{cf}^N(\kappa) = \omega$. Truth of each fixed first-order formula in one unchanged set structure is absolute between transitive models, by induction on formulas. $\square$

Corollary 17.2 (No uniform regularity detector; sources 02, 06). *Relative to the existence of such a pair of models, no first-order sentence in the language of $\mathcal{B}_\kappa$ holds, in every universe and at every uncountable cardinal cutoff, exactly when the external cutoff is regular, or exactly when it is strongly inaccessible. More strongly, no invariant determined by the isomorphism type of the enriched structure alone determines the external cofinality of its cutoff in a universe-independent way.*

Proof. Such a sentence or invariant would take different values on the two identical structures of theorem 17.1. $\square$

This is a relative-consistency separation using a measurable cardinal: no measurable cardinal is constructed or proved to exist, and ZFC is not claimed to prove the existence of such a pair of universes. The statement concerns fixed first-order sentences; unrestricted second-order satisfaction is *not* asserted to be unchanged, since the family of external subsets of the carrier changes. The equality established in N is with the *old* V_κ^M ; it is not claimed that $H_\kappa^N = V_\kappa^N$, and the rank hierarchy of the extension can contain new objects. The corollary also shows why diagnosing singularity by failure of some unspecified Replacement instance is unreliable: the small universe keeps every axiom of ZFC while the ambient universe acquires an unbounded sequence invisible to it.

Proposition 17.3 (Where the missing sequence lives; sources 02, 06). *In the situation of theorem 17.1, the first new birthday is exactly $\beta(M, N) = \kappa$. The Prikry sequence $h : \omega \rightarrow \kappa$, and its range, have hereditary size κ , so they lie in $H_{\kappa^+}^N \setminus H_\kappa^N$; any faithful pointed code for them needs at least κ nodes. For every epsilon number λ with $\kappa < \lambda < \kappa^+$, of which there are some, the carrier $\mathbf{No}_{<\lambda}$ changes and its interpreted universe H_{κ^+} contains the new sequence.*

Proof. No subset below κ is new, while the range of h is a new subset of κ (already by regularity of κ in M); theorem 15.4 gives $\beta = \kappa$, and every $\lambda > \kappa$ contains that sign layer. The values of h are cofinal ordinals below κ , so the transitive closure of h contains all of κ and has size κ . For $\kappa < \lambda < \kappa^+$ the cardinal ceiling is κ^+ in both models. Iterating $\alpha \mapsto \omega^\alpha$ ω times from above κ gives a fixed point of cardinality κ , an epsilon number in the interval. $\square$

The proposition distinguishes the *appearance of new elements* from a *change of the complete first-order theory*; it proves the former and does not assert the latter for every enlarged cutoff. By contrast, an extension adding a real changes $\mathbf{No}_{<\kappa}$ at every common uncountable cardinal (example 15.6); the preservation criterion distinguishes the two kinds of extension without referring to the names of their forcing notions. The surcomplex enrichment is likewise unchanged in the Prikry situation (theorem 19.8).

Part IV

The full class and the surcomplex numbers

18 The full class: the announced theorem and Kunen rigidity

18.1 The announced global bi-interpretation

Theorem 18.1 (Chen–Hamkins–Yang, announced [6, 7, 8]). *In ZFC there are parameter-free first-order interpretations $\mathcal{I}$ and $\mathcal{J}$ such that, formula by formula, $\mathcal{I}(\mathcal{B})$ is definably identified with $(V, \in)$, $\mathcal{J}(V, \in)$ with $\mathcal{B}$, and both composites have definable isomorphisms to the original structures.*

Status and credit. This is the global theorem announced by Chen, Hamkins and Yang, stated there for the surreal field with birthday order; by lemma 2.1 that language and the birthday-function language are definitionally equivalent. Their slides describe the coding of sets by pointed well-founded extensional relations and then by sign sequences, and mark the work as still being checked; the collection’s foundations report records the announcement in `found:sub:announcement`. The written argument below is source 07’s; source 05 gives the same statement with its bisimulation codes, and internally in every model of ZFC (remark 7.4). It is printed as *a proof of the announced theorem*, not as a new result, and it is not an audit of an unpublished manuscript of Chen–Hamkins–Yang or of their proposed axiomatization.

Proof (source 07). Use the codes of section 5 without a cutoff. Every set has a valid code: choose an injection of $\text{TC}(\{a\})$ into an ordinal (one application of Choice) and code the transported membership relation. The correctness of $\sim$ and $\in_{\mathcal{I}}$ (lemma 5.2) and of the collapse is proved for set-sized graphs and applies to each code. The reverse interpretation takes sign names $\langle \alpha, A \rangle$ with $A \subseteq \alpha$ and defines the arithmetic by the certificates of section 3, which now exist without a size restriction. The two composites are the collapse relation and the sign-name relation $x \mapsto \langle b(x), A_x \rangle$ of sections 6 and 7; both are single formulas, and their correctness is proved one formula at a time, with no truth predicate for V and no class of chosen representatives. $\square$

The local refinements proved in Part I were stated for V by the announcement; the cutoff theorems, the noncardinal pointed version, and the local certificates at singular cutoffs are what sources 02, 05, 06, 07 and 09 propose to add, with the priority caveats of section 20. Source 09 already noted that its refinements may overlap unpublished details of the Chen–Hamkins–Yang project.

18.2 Transfer of Kunen’s inconsistency

An automorphism and a non-surjective elementary self-embedding must not be conflated. For automorphisms of the full class the argument of lemma 19.3 applies (the ordinals form a definable well-ordered class). It fails for a non-surjective elementary self-embedding j , whose restriction to the ordinals need not be an automorphism, and which satisfies $b(j(x)) = j(b(x))$ rather than the externally birthday-fixed equation $b(j(x)) = b(x)$. Literature on birthday-fixed surreal morphisms, such as the structures of Rangel–Mariano [16], must be compared with this distinction in mind (source 07).

Lemma 18.2 (Transferring an elementary embedding; source 07). *Let $j : \mathcal{B} \rightarrow \mathcal{B}$ be a class elementary embedding. For each set a choose any code C with $x_C = a$ and put $\hat{j}(a) = x_{j(C)}$, with j applied coordinatewise. This is a well-defined class elementary embedding $(V, \in) \rightarrow (V, \in)$, and $\hat{j}$ is the identity only if j is.*

Proof. Elementarity preserves validity and $\sim$, so the value does not depend on the code, and $\hat{j}$ is a class defined from j without choosing representatives. For each set-theoretic formula and codes for its parameters, [theorem 18.1](#) and elementarity of j give $\varphi(\vec{a}) \iff \varphi(\hat{j}(\vec{a}))$. The definable sign-name composite is natural under elementary maps; if $\hat{j}$ is the identity, it identifies $j(x)$ and x with the same interpreted sign name, so $j(x) = x$. $\square$

The class-theoretic hypothesis. The classical Kunen inconsistency rules out a nonidentity elementary embedding $V \rightarrow V$ under Choice, in the form that permits the embedding as a class parameter in Separation and Replacement (Hamkins–Kirmayer–Perlmutter [12]). We work, as source 07 does, in GBC for convenience; global choice is stronger than needed. The class $\hat{j}$ is definable from j and inherits availability as a class parameter. Elementarity may be read formula by formula; for a single class formulation one may use Δ_0 -elementarity with membership-cofinality, which holds because $\hat{j}(\alpha) \geq \alpha$ and $\hat{j}(V_\alpha) = V_{\hat{j}(\alpha)}$.

Theorem 18.3 (No nontrivial class elementary self-embedding; source 07). *In GBC, or under the class-parameter Separation and Replacement hypotheses just described, every class elementary self-embedding of $\mathcal{B}$ is the identity.*

Proof. A nonidentity j would give a nonidentity elementary $\hat{j} : V \rightarrow V$ by [lemma 18.2](#), contradicting the cited Kunen theorem. $\square$

The set-theoretic input is the classical Kunen inconsistency; this is not a new proof of it. The theorem concerns the full birthday-expanded class. It does not assert that the pure surreal ordered field has no nontrivial embeddings or automorphisms; it does not classify the elementary self-embeddings of the bounded $\mathcal{B}_\kappa$ (which [theorem 9.10](#) reduces to those of H_κ); and it does not prohibit external self-maps of a countable nonstandard model’s internal surreal structure, which need not be classes of that model with the required Separation and Replacement properties.

[Added 4 October 2026, batch 89: how much the class-parameter hypothesis carries in a setting with atoms. In the urelement kernel models (M_κ^{ker} of [Part VI](#), written $M_\kappa(A)$ there) of the research-report collection’s [SetTheory/Cardinals/docs/reports/ordinals-and-order-types/naming-elementary-embeddings](#) (batch-89 manuscript 05), every injection of the atoms lifts to an elementary self-embedding that fixes every pure set and is amenable (its restriction to each set is a set), and these lifts are exactly the elementary self-embeddings with transitive range (its Theorems 3.2 and 3.4, `nee:thm:lifts`, `nee:thm:initial`). Naming finitely many such lifts keeps Replacement in the expanded language exactly under a condition on the components of the named atom injections (its Theorem 5.1, `nee:thm:profile`). This does not touch [theorem 18.3](#): those embeddings fix every pure set, so every ordinal and every surreal number, while the theorem concerns embeddings of the pure class $\mathcal{B}$, which [lemma 18.2](#) turns into embeddings of V . That report is AI-assisted, unrefereed and not formalized.]

19 Surcomplex numbers and the orientation obstruction

19.1 The enriched surcomplex structures

For an epsilon number λ , $\mathbf{No}_{<\lambda}[i]$ is algebraically closed because $\mathbf{No}_{<\lambda}$ is real closed. Let $c(x+iy) = x - iy$ be conjugation and

$$\text{cb}(x+iy) = \max(\text{b}(x), \text{b}(y)),$$

the *coordinate birthday*, whose value is an embedded real ordinal. Put

$$\mathcal{C}_\lambda = (\mathbf{No}_{<\lambda}[i]; 0, 1, +, -, \cdot, c, \text{cb}), \quad \mathcal{C}_\lambda^i = (\mathcal{C}_\lambda, i),$$

and $\mathcal{C}, \mathcal{C}^i$ for the full classes on $\mathbf{No}[i]$. The coordinate birthday is a stated convention, not the birthday of a combinatorial-game representative of a complex object, and not asserted to be a pre-existing universal definition of a “surcomplex birthday”. Conjugation, cb and i are explicit extra structure; none is asserted to be definable in the pure algebraically closed field, whose inclusions are elementary in the pure language [14].

Lemma 19.1 (Independence of orientation; source 06). *cb is unchanged if i is replaced by $-i$, and its restriction to $\mathbf{No}_{<\lambda}$ is b.*

Proof. The replacement changes coordinates (x, y) to $(x, -y)$, and negation reverses every sign without changing length, so $\text{b}(-y) = \text{b}(y)$; also $\text{b}(0) = 0$. $\square$

Proposition 19.2 (Recovery of the real birthday structure; all five sources). *Both $\mathcal{C}_\lambda$ and $\mathcal{C}_\lambda^i$ interpret $\mathcal{B}_\lambda$ without parameters on the fixed field of c , with order*

$$x < y \iff \exists t [c(t) = t \wedge t \neq 0 \wedge t^2 = y - x]$$

for real x, y . The structure $\mathcal{C}_\lambda^i$ is uniformly bi-interpretable with $\mathcal{B}_\lambda$, hence with $(H_\lambda, \in)$ at cardinal cutoffs and with $(H_{\kappa(\lambda)}, \in, \lambda)$ otherwise; the same holds for $\mathcal{C}^i, \mathcal{B}$ and $(V, \in)$ formula by formula.

Proof. The fixed field is $\mathbf{No}_{<\lambda}$, real closed, where positivity is being a nonzero square; field operations restrict, and cb restricts to b. Conversely interpret complex numbers as pairs of reals with $(x, y)(u, v) = (xu - yv, xv + yu)$, $c(x, y) = (x, -y)$, $i = (0, 1)$ and $\text{cb}(x, y) = (\max(\text{b } x, \text{b } y), 0)$. With i named, $(x, y) \mapsto x + iy$ is definable with inverse

$$z \mapsto \left(\frac{z + c(z)}{2}, \frac{z - c(z)}{2i} \right),$$

and the real composite is $x \mapsto (x, 0)$. These are the definable isomorphisms of a bi-interpretation; compose with [theorem 6.3](#) and [theorem 18.1](#). $\square$

Thus cardinal rounding, recovery of a noncardinal cutoff, and all the set-theoretic distinctions above transfer to the oriented surcomplex structures, and adjoining i creates no new set-theoretic size requirement. The real-axis recovery is standard quadratic-extension algebra; its role here is to identify which enrichment carries the foundational information. In the Prikry situation of [theorem 17.1](#), $\mathcal{C}_\kappa$ and $\mathcal{C}_\kappa^i$ are also unchanged, because both coordinates and all displayed operations are; so adjoining i and retaining conjugation does not repair the inability to recover the ambient cofinality (sources 02, 06).

19.2 Rigidity and the two-element automorphism group

Lemma 19.3 (Rigidity of a standard birthday cutoff; sources 05, 06, 07, 09). *Every automorphism of $\mathcal{B}_\lambda$ is the identity, and every automorphism of a transitive structure $(H_\kappa, \in)$, or $(H_\kappa, \in, \lambda)$, is the identity.*

Proof. The embedded ordinals are definable and numerically well ordered, so any order automorphism of this segment fixes every ordinal by transfinite induction; the automorphism therefore fixes all birthdays, preserves the definable bit predicate at each fixed coordinate, and fixes every sign sequence. For H_κ , external well-founded induction gives $h(x) = \{h(y) : y \in x\} = x$; naming an ordinal does not change this. $\square$

The first assertion is a case of the collection’s existing simplicity rigidity: the report `surcomplex/surcomplex-field-automorphisms` shows, in its section `saut:sec:exponential` (subsection “Simplicity restores the canonical surreal numbers”), that a bijection preserving numerical order and prefix simplicity is the identity, and prefix is definable from order and birthday (lemma 4.1). It is printed here once, not as a new result. Rigidity is not pointwise definability: a rigid structure can have elements that are not individually definable without parameters.

Theorem 19.4 (The orientation obstruction; sources 05, 06, 09, after Jeřábek). *For every epsilon number λ , $\text{Aut}(\mathcal{C}_\lambda) = \{\text{id}, c\}$. Consequently $\mathcal{C}_\lambda$ is parameter-free mutually interpretable with, but not parameter-free bi-interpretable with, $(H_\lambda, \in)$ at cardinal cutoffs, or $(H_{\kappa(\lambda)}, \in, \lambda)$ otherwise; after naming i the bi-interpretation holds (proposition 19.2).*

Proof. An automorphism preserves the fixed field of c , its definable order and birthday, so its restriction is the identity by lemma 19.3; it sends i to a root of $X^2 + 1$, so it is the identity or conjugation, as in `saut:thm:axis` ($\text{Aut}(F(i)/F) = \{\text{id}, c\}$ for real closed F). Both preserve c and cb (lemma 19.1).

A parameter-free bi-interpretation induces mutually inverse homomorphisms between the automorphism groups: an automorphism preserves the definable interpretation domain and equivalence relation, induces an automorphism of the interpreted structure, and the definable round-trip isomorphism recovers it. The set structure is rigid (lemma 19.3), whereas conjugation is a nonidentity automorphism of $\mathcal{C}_\lambda$. This excludes *every* parameter-free bi-interpretation, not only the coordinate one. Mutual interpretation holds: recover the real birthday structure inside $\mathcal{C}_\lambda$ and then its set universe; in the other direction use the reverse surreal interpretation and the pair field, omitting the constant i . $\square$

The general real/complex obstruction, that real and complex fields with conjugation are mutually interpretable but need a named imaginary unit for a parameter-free bi-interpretation, was explained by Jeřábek in 2021 [9]; only source 06 credited it, and the credit applies to the versions in sources 05 and 09 as well. The group computation is a consequence of classical rigidity and of `saut:thm:axis`, not a new classification of the unrestricted field automorphisms of $\mathbf{No}[i]$ studied in the automorphisms report. The point here is the distinction between mutual interpretability and bi-interpretability, and why an unnamed imaginary unit cannot be treated as harmless notation in the latter claim. The two roots i and $-i$ have the same parameter-free type in $\mathcal{C}_\lambda$; interpreting a rigid universe does not force the interpreting structure to be rigid, and it is the definability of both round trips that transmits rigidity (source 06).

19.3 Two lifts of each real embedding

Theorem 19.5 (Two lifts, and only two; source 06). *Let κ, λ be uncountable cardinals. Every elementary embedding $f : \mathcal{B}_\kappa \rightarrow \mathcal{B}_\lambda$ has exactly two elementary extensions $\mathcal{C}_\kappa \rightarrow \mathcal{C}_\lambda$,*

$$\widehat{f}_\epsilon(x + iy) = f(x) + \epsilon i f(y), \quad \epsilon \in \{1, -1\},$$

relative to reference roots of -1 , and every elementary embedding $\mathcal{C}_\kappa \rightarrow \mathcal{C}_\lambda$ is one of these. After naming the reference imaginary units only the positive lift remains. Hence

$$\text{Emb}_{\text{el}}(\mathcal{C}_\kappa, \mathcal{C}_\lambda) \simeq \text{Emb}_{\text{el}}(H_\kappa, H_\lambda) \times \{1, -1\}, \quad \text{Emb}_{\text{el}}(\mathcal{C}_\kappa^i, \mathcal{C}_\lambda^i) \simeq \text{Emb}_{\text{el}}(H_\kappa, H_\lambda),$$

the signs multiplying under composition, and $\mathcal{C}_\kappa^i \prec \mathcal{C}_\lambda^i \iff H_\kappa \prec H_\lambda$ for $\kappa < \lambda$ (source 05 proved the last equivalence, and the positive-lift correspondence, at regular cardinals).

Proof. The displayed maps are field embeddings commuting with c and preserving cb , because f preserves order and birthdays, and they differ on i . For elementarity, name i and translate each complex formula into $\mathcal{B}_\kappa$ by coordinates, as in [proposition 19.2](#); elementarity of f gives elementarity of $\widehat{f}_1$, and composing with c gives the other lift in the language without i . Conversely an elementary embedding of the complex structures restricts to an elementary embedding of the definable real structures and sends i to $\pm i$, and the field laws force the displayed form. Combine with [theorem 9.10](#). $\square$

At $\kappa = \lambda$ with $f = \text{id}$ this gives $\text{Aut}(\mathcal{C}_\kappa) = \{\text{id}, c\}$ again. The statement concerns the specified birthday-and-conjugation enrichment; it is emphatically not a classification of automorphisms of the pure algebraically closed field, which has far more symmetry (see the automorphisms report). It is a specialization combining the local theorem with coordinate complexification, not a new general Galois principle.

19.4 The exact parameter threshold for orientation

Theorem 19.6 (Orientation threshold; source 06). *Let κ be an uncountable cardinal and $A \subseteq \text{No}_{<\kappa}[i]$ any set of parameters. The following are equivalent:*

- (i) *some root of $X^2 + 1$ is definable in $\mathcal{C}_\kappa$ over A ;*
- (ii) *a linear order of the whole carrier is definable in $\mathcal{C}_\kappa$ over A ;*
- (iii) *A contains a nonreal element.*

In particular neither an imaginary unit nor a linear order is definable over any collection of real parameters, and one nonreal parameter suffices for both.

Proof. If all parameters are real, conjugation fixes them and exchanges the two roots, so neither root is definable; and a definable linear order $\triangleleft$ would be preserved by c , while for nonreal z exactly one of $z \triangleleft c(z)$ and $c(z) \triangleleft z$ holds and c reverses it. If $z \in A$ is nonreal, $v = z - c(z)$ is nonzero and purely imaginary, $-v^2$ is real and positive, and

$$j = \frac{z - c(z)}{\sqrt{-(z - c(z))^2}},$$

with the positive square root of the definable real field, is definable from z and satisfies $j^2 = -1$. The definable coordinates $(\frac{w+c(w)}{2}, \frac{w-c(w)}{2j})$, ordered lexicographically, give a definable linear order of the carrier; it is not a field order. $\square$

Each defining formula uses finitely many parameters; allowing arbitrarily many real parameters does not help because conjugation fixes them all. The proof uses only that $\mathbf{No}_{<\kappa}$ is real closed. A nonreal parameter removes the twofold obstruction to the coordinate round trip, but if it is kept as a named constant, its two real coordinates must be kept in the real interpretation and the corresponding sets in the H_κ interpretation. It is *not* inferred that an arbitrary expansion $(\mathcal{C}_\kappa, z)$ is bi-interpretable with the *unexpanded* H_κ ; naming a root of -1 avoids the issue because its coordinates $(0, \pm 1)$ are already definable.

19.5 The full class, under GBC

Theorem 19.7 (Surcomplex elementary self-embeddings; source 07). *In the class setting of theorem 18.3, the class elementary self-embeddings of $\mathcal{C}$ are exactly the identity and conjugation, and those of $\mathcal{C}^i$ only the identity. In particular every such self-embedding is surjective.*

Proof. An elementary self-embedding of $\mathcal{C}$ restricts to one of the definable real structure $\mathcal{B}$, which is the identity by theorem 18.3; it sends i to $\pm i$, and the coordinates are fixed, so it is the identity or conjugation. Both are automorphisms. Naming i excludes conjugation. $\square$

This is the one surcomplex result of this report that goes beyond the automorphism statements of the automorphisms report: it concerns elementary self-embeddings that are not assumed surjective, of the full structure, under GBC. With lemma 19.3 in place of Kunen’s theorem the same argument gives only the two-element *automorphism* group of the bounded structures. Nothing is asserted about unrestricted field embeddings, and the specified conjugation and coordinate birthday are not claimed to be recoverable from the pure surcomplex field.

19.6 Outer models on the surcomplex side

Theorem 19.8 (The surcomplex outer-model contrast; sources 05, 07). *For transitive $M \subseteq N$ with the same ordinals, the inclusion of the full surcomplex fields is elementary in the pure field language, and also after naming i and conjugation. After additionally naming cb , with or without i , it is elementary exactly when $M = N$. If $M \neq N$, the least coordinate birthday of a new surcomplex number is $\beta(M, N)$.*

Proof. The pure-field statement is proposition 14.2. With i and c named, translate every formula into real coordinates; it becomes an ordered-field formula, so the inclusion is elementary. With cb named the real axis with its birthday is definable, so an elementary inclusion would restrict to $\mathcal{B}^M \prec \mathcal{B}^N$, which forces $M = N$ by theorem 16.5. If $cb(x + iy) < \beta$, both coordinates are old; a new real sign of birthday β is a new surcomplex number of coordinate birthday β . $\square$

The result concerns field and coordinate structure, not complex analysis: it involves no contour integral, topology or total surcomplex exponential, and is not a consequence of an analytic continuation or a valuation argument.

Part V

Scope, questions and provenance

20 What the theorems do and do not imply

20.1 Logical strength versus field strength

Already $\mathcal{B}_{\varepsilon_0}$ recovers H_{ω_1} . Within that membership structure, the natural numbers and all subsets of the natural numbers form the usual full two-sorted structure of second-order arithmetic ([proposition 12.1](#)). The birthday-field interpretation thus includes quantification over all actual subsets of ω , not merely over computable subsets, because the numbers coding those subsets are first-order elements. This does not make set theory definable in the pure ordered-field reduct, where birthday is not definable ([proposition 12.5](#)), and it does not give a decision procedure ([corollary 12.2](#)).

20.2 Four statements that must not be conflated

For a cutoff λ the following can have very different answers:

Question	Exact issue in this report
Is $\mathbf{No}_{<\lambda}$ a real closed field?	Classical epsilon-number cutoff theorem.
Which sets can be decoded?	Exactly $H_{\kappa(\lambda)}$, by rooted relation codes.
Does the decoded universe satisfy ZFC?	At cardinal cutoffs, exactly the worldly condition.
Is an initial inclusion elementary?	Both cutoffs must be cardinals, followed by elementarity of their hereditary-size universes.

An algebraic closure theorem alone does not answer any of the last three questions. Conversely, a transitive set model satisfying ZFC need not arise from a regular inaccessible cutoff ([proposition 11.7](#)), and the same enriched field can sit below a regular and a singular cutoff in two universes ([theorem 17.1](#)). Hereditary size, not rank, is what the birthday structure measures: the recovered universe is H_κ , which in general differs from V_κ ; the two agree, for example, at every worldly κ ([lemma 11.1](#)).

20.3 Choice and quotients

Ordinary Choice is used to enumerate an individual transitive closure and, in some proofs, to enumerate a particular fiber or choose a particular bijection. The interpretation does not choose one representative simultaneously for every set. It keeps all valid codes and quotients by a definable equivalence relation. This is why Global Choice is unnecessary outside [section 18.2](#).

The exact cardinal-rounding conclusion is stated in ZFC. Without Choice, not every transitive closure is well-orderable, and a code on an ordinal cannot represent every set; [theorem 13.2](#) identifies the exact range of the full-class coding in ZF. No choice-free version of the cutoff theorems is claimed, and nothing in the ZFC proofs justifies silently deleting Choice.

20.4 Set models, nonstandard models, and class truth

At a fixed cutoff, the structures in the main theorems are actual set structures with ordinary first-order satisfaction; their interpretations are not hypothetical syntactic models satisfying only a fragment of surreal arithmetic. A nonstandard model of ZFC can internally perform the same constructions, but its coded well-foundedness and its internal cardinal ceiling must be read in that model; an externally ill-founded model is not covered by an external Mostowski-collapse claim merely because it internally believes its code is well founded. Arbitrary nonstandard models of a proposed surreal-arithmetic axiom system need a separate analysis.

For the full class **No**, one applies the translation to a fixed formula and fixed set parameters. Statements about “all formulas” are metatheoretic schemes. We do not form a set of all class functions, a proper-class membership quotient, or a truth predicate for V inside ZFC. The cutoff results do not require a class theory, a Grothendieck universe, or a large-cardinal existence axiom; GBC is used only in [section 18.2](#) and [theorem 19.7](#), a measurable cardinal only in [section 17](#), and an inaccessible only in [proposition 11.7](#).

20.5 Why uncountability and the epsilon hypothesis matter

For the hereditary-size bi-interpretation at cardinal cutoffs, uncountability ensures that countably sized computation certificates and the ordinary infinite set ω remain below the cutoff. At ω the finite-birthday surreals form the dyadic ring, not a field, and only the forward interpretation onto the hereditarily finite sets is claimed ([proposition 8.1](#)). For noncardinal cutoffs, the epsilon hypothesis supplies closure of the bounded pairing terms and field operations. Arbitrary ordinal sign cutoffs that fail those closure conditions are not classified; the full birthday-field statement would then need a different language or partial operations. The reconstruction also applies to the canonical birthday function only, and uses the presence of every sign of each length below the cutoff: a small subfield with an inherited birthday may omit most subset codes, and no such generalization is asserted (source 05). For outer models all ordinals are assumed common; models of different heights are a different problem.

20.6 Non-claims of the sources

Every limitation, disclaimer and priority caveat stated by a source is kept. Most appear at their point of use above; this subsection lists them by source.

Across the report. The proofs are written mathematical arguments. They have not been independently refereed and have not been formalized in Lean; no Lean source is supplied. The finite programs check finite analogues only. No named longstanding conjecture is resolved. Priority is not certified by any of the five targeted literature and repository searches; “not found” is a bounded search result, not proof of absence.

[Added 3 October 2026, batch 89: this subsection lists the non-claims of sources 02, 05, 06, 07 and 09. Those of sources 10 and 11 are collected in [sections 39.4](#) and [53.3](#).]

Source 02. The full-class interpretation is Chen–Hamkins–Yang’s prior announced work; its contributions are the bounded refinement and its consequences, with priority not certified. Interpreting $H_{\kappa(\lambda)}$ is not interpreting the ambient universe and does not assert $H_{\kappa(\lambda)} \models \text{ZFC}$. The translated Power Set sentence is one axiom; it does not characterize Replacement, regularity or

full ZFC. Its own elementary-inclusion conditions were necessary only (the converse is now [theorem 9.3](#)). The results concern concrete externally well-founded structures. Sharpness is intrinsic to the interpretation. The collector recovers an ordinal value, not a surreal of birthday λ . Prikry forcing is imported; no measurable cardinal is constructed; the statement is relative consistency and concerns fixed first-order sentences, not second-order satisfaction. The preservation criterion must not be replaced by “no new short sequences”, and a proper enlargement at a fixed cutoff is not claimed non-elementary merely because it is proper. The onset proposition concerns new elements, not a change of complete theory. In the surcomplex part i , c and the coordinate birthday are explicit structure, not definable in the pure algebraically closed field, and no new automorphism classification is claimed. Only one-way interpretation was claimed at ring cutoffs and by source 02 generally. The Rangel–Mariano program is not settled. The finite tests do not prove the transfinite results.

Source 05. Its package was called candidate-original: not independently refereed or formally verified; no named problem settled; the audit targeted; priority unverified. It omitted the Chen–Hamkins–Yang credit, which is restored here. Classical inputs (sign arithmetic, cardinal closure, well-founded recursion, quantifier elimination, and the simplicity rigidity of the automorphisms report) are credited. Regularity was its uniform hypothesis and it inferred no singular theorem; $H_\kappa \neq V_\kappa$ in general and $\mathcal{P}(\alpha)$ need not be a set in H_κ . The coordinate birthday is a convention, not a game birthday. No sharp product-birthday bound is used. The outer-model theorem needs transitive same-ordinal models; nonstandard models are handled internally. Its non-elementarity is witnessed with parameters, not by distinct parameter-free theories. The first new birthday is not asserted regular, and signs of birthday at most ω are not identified with reals. The bounded detector needs an enumeration bound. No small-subfield generalization and no different ordinal heights are covered. The embedding correspondence is a reduction, not an existence theorem, and claims no large-cardinal consistency result; exponential automorphisms are untouched, and the collection’s exponential faithfulness result is outside its proof dependencies. The detector formula is not claimed to be existential after expansion. Its Python checks are diagnostic only.

Source 06. Chen–Hamkins–Yang is prior work and the central coding idea is theirs; Jeřábek (2021) is prior for the conjugation obstruction; field closure and Prikry are imported. Novelty is provisional; not peer-reviewed or Lean-certified; no named open problem settled. The coordinate birthday is a convention, and the automorphism conclusions do not apply to the pure surcomplex field or classify its automorphisms. The $\beth_\omega$ theorem does not show that every singular κ fails Replacement, and internal ZFC does not imply external inaccessibility. The ZF boundary concerns this construction only, and internal Choice is not the range statement. Prikry gives relative consistency, not a ZFC theorem, and not $H_\kappa^N = V_\kappa^N$. A nonreal parameter does not give bi-interpretability with an unexpanded H_κ . Rigidity is not pointwise definability. Its reduct corollary is not a minimal-language claim. It left open noncardinal cutoffs (answered here for epsilon numbers), ill-founded models of a surreal-arithmetic theory, quantifier elimination for an order-and-simplicity language, and a universal embedding theorem without Global Choice. The finite tests do not verify the theorems. [Added 3 October 2026, batch 89: the last of these is answered in part by source 11; see the note after [question 21.7](#).]

Source 07. It omitted the Chen–Hamkins–Yang credit and presented the full-class bi-interpretation as a proposed contribution; that is corrected in [theorem 18.1](#), and it had no Jeřábek credit, now added. Its results were proposed contributions without priority certification, refereeing or Lean.

It claims no new proof of Kunen’s inconsistency, no rigidity of the pure field, no classification of bounded elementary self-embeddings, and no distinct parameter-free theories from non-elementarity; nothing about unrestricted field embeddings. Its plateau theorem is forward only, and its Choice remark says the construction is not a reconstruction of V without Choice. The Kunen transfer needs GBC or class-parameter hypotheses and does not forbid external self-maps of countable nonstandard models. Nonstandard models are handled internally. The reconstruction is a logical representation, not literal membership between chosen representatives; there is no canonical code, no decidability result, and no simple closed sign rules for arithmetic. Rangel–Mariano and Bertram are distinct programs whose categorical obstructions are not refuted. The finite checks are limited.

Source 09. The global result of Chen–Hamkins–Yang, the van den Dries–Ehrlich cutoff theorem with its erratum, and ordinary well-founded coding are prior work; the refinements may overlap unpublished details of the Chen–Hamkins–Yang project; no named conjecture is resolved; no Lean, no peer review. It did not credit Jeřábek; that credit is added. Cardinal rounding is not a rank cutoff. The principles **Et** and **Pack** are explicit schemes, not identified with any formulation of **SA**, and the independence examples are within this family only. The detector bound is sharp only for the old-power-set parameter; no quantifier-complexity claim is made across the translation; failure of the bound does not imply elementarity. Extension rigidity concerns standard cutoff structures and does not forbid abstract elementary extensions. The orientation result is a consequence of classical rigidity, not a new automorphism classification. Choice is used object by object and no ZF version of its theorems was claimed ([theorem 13.2](#) is source 06’s). Nonstandard models are internal; no class theory, Grothendieck universe or large cardinal is needed for the cutoff theorems. At ω the cutoff is a ring; non-epsilon cutoffs are not classified. The singular worldly construction is a standard illustration needing an inaccessible.

21 Open questions, re-scoped

Answered inside this report.

- Source 02’s first question, a sufficient criterion for elementary birthday inclusions: answered by source 09 ([theorem 9.3](#), [remark 9.4](#)).
- Source 02’s caution that a two-way theorem must name the cutoff: answered by source 09’s pointed bi-interpretation ([remark 6.5](#)).
- Source 05’s optimal cutoffs at singular cardinals: answered by sources 06 and 07; at noncardinal epsilon numbers: answered by source 09. Source 06’s and source 07’s noncardinal-cutoff questions: answered by source 09 for epsilon numbers.
- Source 09’s fourth target, the choice-free analogue: answered for the full-class graph coding by source 06 ([remark 13.4](#)).
- Source 06’s universal embedding theorem without Global Choice: answered in part, over ZF and with a ZFA counterexample, by source 11 ([Part VII](#)); see the note after [question 21.7](#). [Added 3 October 2026, batch 89.]

Still open.

Question 21.1 (Cutoffs that are not epsilon numbers). For which ordinals η that are not epsilon numbers, beyond the ring cutoffs of [proposition 8.1](#) (forward only), is there a uniform reconstruction from $\mathbf{No}_{<\eta}$ with its inherited operations, and what exactly is recovered? (Re-scoped from sources 05, 06 and 07.)

Question 21.2 (Minimal signature). How much of $\{0, 1, +, -, \cdot, <, b\}$ can be deleted while retaining a uniform interpretation of hereditary sets? [Corollary 12.6](#) shows that $<$, b and the graph of one ordinal pairing polynomial suffice; whether $<$ and b alone, or addition with birthday, suffice is open, as is whether multiplication is necessary. (Sources 02, 05, 07; partial progress by 06.)

Question 21.3 (Elementary embeddings and explicit criteria). Replace the condition $H_\kappa \prec H_\tau$ in [theorem 9.3](#) by useful explicit hypotheses in interesting families of cardinals (source 09). Classify the elementary self-embeddings of particular bounded $\mathcal{B}_\kappa$; by [theorems 9.10](#) and [19.5](#) this is the corresponding problem for H_κ (source 07).

Question 21.4 (Forcing and the bounded theory). Which changes of the complete bounded theory can occur under forcing that adds new bounded subsets, and how much of the cutoff predicate is first-order recoverable from an appropriately expanded H_κ (source 02)? The detectors of [section 16](#) show failure of elementarity with parameters; they do not decide changes of parameter-free theories.

Question 21.5 (Quantifier complexity). What is the least quantifier complexity of a uniform formula detecting all proper same-ordinal outer-model inclusions in the birthday language (source 05)?

Question 21.6 (Surcomplex reducts). What set-theoretic information is recoverable from $(\mathbf{No}_{<\kappa}[i]; 0, 1, +, -, \cdot, \text{cb})$ when neither conjugation nor i is named (source 05)? [Theorem 19.6](#) concerns the language with conjugation.

Question 21.7 (Axiomatics and formalization). Establish definitional equivalence, or its failure, between Et and Pack and particular published axiom formulations of surreal arithmetic, once finalized (source 09); analyse ill-founded models of such a theory, quantifier elimination for an order-and-simplicity language, and a universal embedding theorem without Global Choice (source 06); and formalize the size-certified interpretation, beginning with the prefix, bit and pairing lemmas, then the certificate verifier and its hereditary-size bound (sources 05, 06, 07, 09).

[Added 3 October 2026, batch 89: the third item, “a universal embedding theorem without Global Choice”, is answered in part by source 11 ([Part VII](#)). Over ZF, a set X carries an X -indexed algebraically independent family of positive infinite omnific integers, equivalently $\mathbb{R}(T_X)$ embeds in $\mathbf{No}$ over $\mathbb{R}$, iff X injects into the power set of an ordinal; so universality for the free fields on all sets is exactly the first Kinna–Wagner principle ([theorem 43.5](#)), with optimal set-sized real-closed targets U_κ ([theorem 44.3](#)). Every linear order on a well-orderable set order-embeds in $\mathbf{No}$ ([proposition 45.3](#)). In the ordered Mostowski permutation model, a model of ZFA, an ordered field whose finitely generated coordinate subfields all embed in order has no field embedding into the internal $\mathbf{No}$ ([theorem 49.1](#)). Still open: universality for prescribed orders and for general ordered fields without choice, and any transfer of the ZFA example to ZF ([questions 51.1](#), [51.2](#) and [51.5](#)). The other items of this question are not affected.] The collection’s Lean library already contains relevant sign-sequence infrastructure (prefix truncation, the failure of birthday precedence to be prefix simplicity, Hessenberg agreement of ordinal arithmetic), but none of the interpretation theorems of this report is formalized.

22 Provenance and verification

22.1 The sources’ repository and literature audits

All five sources inspected the repository read-only at the pinned commit

4f2645599121fa872c7104995e47f86f0382351f

and modified nothing. Source 02 inspected the tree, the root and documentation indexes, the catalogue source and the foundations report, and records its scope in `02-birthday-cutoffs-SOURCE_AUDIT.md`, kept byte-identical. Source 05 records its requested line ranges, some truncated, in `data/05-birthdays-recover-sets-repository_audit.json`. Source 06 inspected listings, the first 200 lines of the documentation catalogue, the first 150 lines of the foundations README, and the README and first 240 lines of the automorphisms article. Source 07 inspected the tree, READMEs, catalogue, formalization ledger, the foundations introduction and the automorphisms README. Source 09 inspected the catalogue, the relevant parts of the foundations report and the scope of the automorphisms report. The corrections these audits need are in [section 1.5](#).

The sources' literature checks were: the Chen–Hamkins–Yang announcements and slides (sources 02, 06, 09; source 06 cites slide pages 103, 107 and 122, and source 02 pages 96–122, which are compatible); van den Dries–Ehrlich and its erratum, Bournez–Guilmant, Poveda–Rinot–Sinapova, and Rangel–Mariano (sources 02, 06, 09); Freire–Hamkins, Ehrlich's paper on the simplicity hierarchy [18] (record and abstract only) and Ehrlich–Kaplan [19] (background, not an input) (source 05, whose session could not load the publisher PDF of the erratum); Bertram and Hamkins–Kirmayer–Perlmutter (source 07); Jeřábek (source 06). These external checks were not repeated for this merge.

22.2 Dependency and novelty ledger

Result	Main inputs	Status here
Global birthday-field bi-interpretation (theorem 18.1)	Pointed well-founded relation coding	Prior Chen–Hamkins–Yang announcement; written proof from source 07; not claimed new.
Real-closed epsilon cutoffs	Classical surreal field theory	Prior van den Dries–Ehrlich, with erratum.
Small arithmetic certificates, two routes	Option graphs (09); prefix-pair tables (05, 06, 07)	Explicit local proof ingredients; no priority claim for the general locality principle.
Cardinal rounding and cutoff recovery	Bounded pairing, rooted collapse, collector, sign-graph bridge	Proposed local refinement (09), with cases in 02, 05, 06, 07.
Elementary-inclusion spectrum and embeddings	Collector and coherent bi-interpretations	Proposed classification (09); embeddings (05, 06).
Power Set, eternity, packing, worldly, $\beth_\omega$	Hereditary cardinal bounds	Proposed calibrations (02, 06, 09); the underlying H_κ facts are standard.
ZF range	Collapse in ZF	Direct refinement of the coding (06); no claim about other interpretations.
Preservation, first new birthday	Sign/subset bijection	Direct consequences (02, 05, 06, 07), tools rather than isolated discoveries.

Result	Main inputs	Status here
Detectors, extension rigidity, $M = N$	Old-power-set parameter; row search	Proposed explicit refinements (05, 07, 09); outer-model reasoning is standard.
Prikry invisibility	Imported Prikry forcing	Proposed application under a measurable cardinal (02, 06).
Kunen transfer	Imported Kunen inconsistency (GBC)	Proposed application (07).
Surcomplex orientation, automorphisms	Real-axis recovery, rigidity, Jeřábek	Consequence of classical rigidity and the automorphisms report.
Two lifts, orientation threshold, full-class self-embeddings	Coordinates, rigidity, Kunen	Proved specializations (06, 07); the only surcomplex items new to the collection.

The places where the argument could not be replaced by a general slogan are the arithmetic certificates (saying that the global construction is definable does not prove that the same formula computes arithmetic in an H_κ without Replacement), and the collector (a noncardinal cutoff is not recovered merely by interpreting the unpointed hereditary-size universe).

22.3 Finite checks

Each source supplied a standard-library Python program checking finite analogues of its coding conventions. All five were rerun for this merge with Python 3.14.4 on copies of the files; each passed, and each recorded output was reproduced exactly, apart from line endings. The recorded runs are:

Source	Recorded finite checks
02	65,025 prefix comparisons and 1,538 bit positions through length 7; 173,880 natural-ordinal matrix addresses on 81 Cantor-normal-form samples; 687 relabellings of all 16 sets of V_4 ; 1,024 code pairs for equality and membership; 4 invalid codes rejected.
05	65,025 prefix comparisons through birthday 7, including the counterexample of remark 4.2 ; 64 acyclic graphs on four nodes, 4,096 graph pairs, 65,536 root-pair checks each for bisimulation equality and membership; 64 matrix round trips; 49 rectangle packings; 30 missing-row detector tests.
06	16,129 prefix and 642 bit instances on 127 words of length at most 6; pairing on 10,000 pairs; all 1,570 rooted relations on at most three vertices, 15 valid codes, 225 pairs each for equality and membership.
07	16,129 prefix comparisons, 642 sign checks, 819 radix addresses (3 radices), 42,849 graph pairs each for isomorphism and membership over 207 labeled graphs decoding 12 sets; the missing-range counterexample confirmed.
09	16,129 prefix pairs, 642 bit positions, 11,440 pairing checks; 530 relations and 1,570 rooted relations on at most three nodes, 15 valid codes, 225 pairs each for equality and membership; the counterexample of Appendix B confirmed.

These tests can expose a reversed edge, a missing root or range condition, a bad pairing index, or an incorrect finite sign formula. They do not prove transfinite well-foundedness, cardinal bounds, bi-interpretability, Replacement equivalences, forcing statements, or any large-cardinal assertion; those rest on the written proofs and the explicitly imported classical theorems.

[Added 3 October 2026, batch 89: the finite programs of sources 10 and 11 are described in [sections 37](#) and [52](#). Both were rerun with Python 3.14.4 on copies; both passed and reproduced their recorded

files exactly, apart from line endings: source 10 checked 8,256 endpoint intervals over the 127 sign words of length at most 6, 2,187 depth-two assignments (576 separated), 1,050 negative-code comparisons and 4,032 atom transpositions; source 11 passed 20,595 assertions with seed 20261003. Their provenance is in [sections 39.5](#) and [53.4](#).]

23 Conclusion

Birthday truncation has a precise set-theoretic effect. At an epsilon-number cutoff λ , rooted sign codes recover the hereditary-size universe at the cardinal ceiling of λ ; when λ is not itself a cardinal, its value survives as a definable interpreted set, which explains why noncardinal cutoff inclusions never become elementary even though all their ordered field reducts are elementary subfields. Packing a birthday fiber requires its full power-set cardinal to fit below the cutoff; bounding definable ordinal-indexed birthday growth is Replacement at cardinal cutoffs; together they detect worldly, not merely inaccessible, cardinals. Across universes, the birthday structure below a cardinal is unchanged exactly when no bounded subset is added, while any new set is detected by an explicit formula; yet a measurable cardinal can be singularized without changing the structure at all. The same information is available in surcomplex arithmetic once conjugation, coordinate birthday and an imaginary unit are specified; omitting the unit preserves mutual interpretation but leaves a two-element obstruction to bi-interpretation. The decisive structure is not the field alone: it is the field with a carefully specified memory of how its elements are born.

[Added 3 October 2026, batch 89: Parts VI and VII, which follow, add two further manuscripts: the weak-base form of the eternity–Replacement equivalence with the atom-support spectra of kernel models, and choiceless universality of the surreal numbers.]

Part VI

Replacement-free surreal cuts, definable families and atom-support spectra

24 Source 10: question, results, notation and relation to Part II

This part was added in batch 89. It prints one further manuscript, written independently of the five merged above: *Surreal Cuts, Replacement, and Atom-Support Spectra: Exact principles for definable families, weak set theories, and changes of language*, “Prepared for Vladimir Reshetnikov with ChatGPT”, 3 October 2026. It is batch-89 manuscript 03 and *source 10* in this report’s local numbering; its shipped files carry the prefix `10-cuts-replacement-`. It was pinned to repository commit `2b7b388ba` (“Catalogue batches 81 to 86”), arrived in commit `7c0f2d9f9` and was placed in commit `23adb85f9`. Its provenance, the mechanical changes made in printing it, and its finite checks are in [section 39](#). Inside this part, “this manuscript”, “this article”, “here” and “we” refer to source 10, and its own Theorems A–C are [theorems 24.1](#) to [24.3](#).

Status and attribution (source 10). This is an unrefereed research manuscript, not a claim of an independently certified breakthrough. Classical simplicity and small-support arguments are credited explicitly. The exact formulations, comparisons, and refinements are proved here, but their historical priority has not been established. No Lean or Rocq verification of the new manuscript is claimed.

Source 10’s summary. We investigate the intersection of Elliot Glazer’s work on weak set theories and reflection with the sign-sequence and foundational developments in the ProveIt repository. The central distinction is between an actual set of surreal bounds and a formula assigning a unique surreal to each member of a set. Over Zermelo set theory with Foundation, actual set cuts have unique simplest interpolants, without Replacement. In contrast, interpolation for definable set-indexed families is equivalent to ordinal bounding, to Replacement for surreal-valued functions, and even to strict upper bounding inside the single interval $(-1, 0)$. On a separately specified cumulative-hierarchy base, these schemes are equivalent to full Replacement and Collection.

A complementary atom-support analysis explains why the conclusion cannot simply be transferred to urelement theories. We give a uniform proof and an exact domain-size analysis for the classical small-kernel models M_κ^{ker} . All have the same pure sets and the same pure surreal structures. For a limit cardinal κ , Collection first fails at domain size $\text{cf } \kappa$; for a successor cardinal it holds without a domain-size restriction. Naming an external enumeration of κ atoms changes the picture: both Replacement and Collection first fail at $\text{cf } \kappa$, although Collection for pure-valued relations, including all surreal-valued relations, still holds. Complete mathematical proofs, a finite regression checker, an explicit $V_{\omega+\omega}$ countermodel, and a formalization plan are supplied.

24.1 Why this intersection is substantive

Glazer’s paper on global choice over Zermelo set theory demonstrates that the precise language and axiom scheme matter when Replacement is absent [20]. His joint 2026 paper with Bokai Yao separates Collection from several reflection principles in urelement set theory formulated with Replacement [21]. These provide two concrete prompts: which closure assertions actually require Replacement, and which failures remain invisible when one looks only at pure mathematical objects?

ProveIt has two relevant components: a surreal development organized around sign sequences, and an independent first-order set-theoretic development, including a Closure axiomatization equivalent to ZF [26, 27]. The repository also contains a substantial report on old surreal fields in larger universes [28]. The present question is different: rather than adding new pure sequences by passing between ZFC universes, we first weaken set-existence axioms and then vary atom supports while keeping *all pure sets fixed*.

The research question answered here is therefore:

What is the exact strength of surreal cut interpolation for definable set-indexed families, and how far can such a pure-structure principle detect failures of Replacement, Collection, or reflection in the ambient universe?

This is a question formulated and resolved within this investigation. We do not identify it with an already named, established open conjecture.

24.2 Three main results

The notation Z_f means Zermelo set theory with Foundation but without Replacement or Choice. The additional hierarchy axiom Hier (source 10's H, renamed in [convention 24.4](#)) will be stated explicitly in [section 25](#); it must not be suppressed.

Theorem 24.1 (A local surreal test for ordinal bounding; source 10's Theorem A; [theorems 28.2](#) and [29.1](#)). *Over Z_f , the following schemes are equivalent: ordinal bounding for definable set-indexed functions; existence of set graphs for such surreal-valued functions; surreal upper bounding; Collection for surreal-valued relations; interpolation for definable set-indexed surreal cuts; and strict upper bounding of definable families inside $(-1, 0)$. Over $Z_f + \text{Hier}$, each is also equivalent to full Replacement and full Collection.*

The construction detecting unbounded ordinals is elementary: a minus sign followed by plus signs, with birthday $\alpha + 2$, encodes the ordinal α inside $(-1, 0)$. The nontrivial foundational issue is showing that ordinary set-cut interpolation does *not* already use the principle being characterized.

Theorem 24.2 (Exact Collection spectra of cardinal-support models; source 10's Theorem B; [theorem 32.2](#) and [corollary 33.4](#)). *Work in a full urelement universe with Choice and a set A of atoms, where $|A| \geq \kappa$ and κ is infinite. Put*

$$M_\kappa^{\text{ker}} = \{x : |\text{ker}(x)| < \kappa\}.$$

Then M_κ^{ker} satisfies Replacement and Choice and contains all pure sets. If κ is a limit cardinal, including ω , Collection holds for every domain of size below $\text{cf } \kappa$ and fails for a domain of size exactly $\text{cf } \kappa$. If κ is a successor cardinal, full Collection holds.

The finite-support and countable-support mechanisms are classical; the exact uniform spectrum is the refinement developed in this presentation. The connection to established work is recorded in [section 24.3](#). The same spectrum, for the same models, is proved independently in the research-report collection ([remark 24.6](#)).

Theorem 24.3 (A language-sensitive separation invisible to surreals; source 10's Theorem C; [theorem 34.2](#) and [corollary 35.3](#)). *Let E name, as an external predicate, a bijection from κ onto a set of κ atoms. In $(M_\kappa^{\text{ker}}, E)$, Replacement and Collection both hold for all domains of size below $\text{cf } \kappa$ and both fail at size $\text{cf } \kappa$. Nevertheless, full Collection for pure-valued relations, and consequently every surreal principle in [theorem 24.1](#), holds even for formulas using E and impure parameters.*

Thus the restriction to pure objects is not merely a technicality. The same pure surreal universe can coexist with very different ambient set-existence principles.

24.3 What is established, and what is not

Conway's cut construction and Gonshor's sign-sequence presentation are established foundations of surreal theory [1, 2]; Bertram provides a recent related pure-set treatment [17]. The elementary simplicity theorem itself is not new. Here its construction is reorganized to exhibit a proof over a base without Replacement.

Small-kernel methods are also established. Hamkins and Yao discuss the finite-support model, crediting the Replacement argument to Lévy, and the countable-support model satisfying Collection

[22, §1.5]. Glazer–Yao’s Proposition 1 identifies the common-atom-base obstruction to Collection; their small-kernel permutation models provide much stronger reflection separations than are attempted here [21]. Yao’s earlier work treats the broader axiomatization and forcing setting [23].

The present contribution is a self-contained theorem package: an explicit weak-base equivalence, a fixed-interval ordinal detector, the sharp support-capacity classification with domain-size thresholds, and its language-sensitive comparison with pure surreal principles. These are proposed as useful research results and formalization targets. The sources consulted do not certify their priority; in particular, a generalization of a classical support argument must not automatically be advertised as a breakthrough.

Component	Evidential status
Ordinary surreal simplicity	Classical result; a bounded-container proof is supplied.
Weak-base equivalence and rank countermodel	Fully argued in this manuscript; not machine-checked.
Support-model mechanism	Classical method, explicitly credited; generalized proofs supplied.
Exact thresholds and predicate expansion	Proved in this manuscript; priority not independently established.
Finite verification	Executed regression tests, not a proof of transfinite statements.
Reflection separations of Glazer–Yao	External results used for context, not claimed as our discoveries.

24.4 Notation of this part

Convention 24.4 (Symbols of source 10). Source 10 was written without knowledge of this report, and several of its symbols already mean something else here. In this part:

Notion in source 10	Here	Do not confuse with
Zermelo set theory with Foundation, no Replacement, no Choice	Z_f	—
Hierarchy axiom (existence and coverage of the stages V_α)	Hier (source: H)	H_κ , the hereditary-size universe of Parts I–V
Set-graph scheme	Gr (source: SG)	SignGraph($\cdot$), the sign-graph operator of section 6 ; Gr asserts that a definable family has a <i>set</i> graph
Collection scheme, and its restriction to domains of size $< \tau$	Collection, $\text{Collection}_{<\tau}$	the collector $\text{Coll}(C)$ of theorem 6.2
Ordinal bounding, upper bounding, surreal Collection, definable-cut interpolation, bounding inside $(-1, 0)$	OB, UB, SC, DCI, NIB	—
Kernel (small-support) model $\{x : \ker(x) < \kappa\}$	$M_\kappa^{\ker}$ (source: M_κ)	the transitive models M, N of Part III and $\mathcal{B}_\kappa^M$; the full-class rank models of the report foundations-and-computation/surreal-well-orders , Part XV

Notion in source 10	Here	Do not confuse with
Rank-initial countermodel $V_{\omega+\omega}$	V_θ , $\theta = \omega + \omega$ (source: $M = V_\theta$)	the ground model M of Part III
Urelement set theory with Replacement	ZFU _R	theories named ZFU that include Collection
Ambient urelement universe; kernel; atom base	$\mathcal{U}$; $\ker(x)$; $V_\alpha(B)$	—
Generic ordinal variables	ζ (source: β), σ (source: δ)	β , reserved for the first new birthday, and δ , reserved for a code domain (convention 1.5)
Ordinal detector in $(-1, 0)$	ν_α	the hereditary cardinals ν_i in the proof of theorem 10.4
Birthday bound $\sup\{b(x) + 1\}$ of a cut	λ , any ordinal	the epsilon-number cutoffs λ of Parts I–V
A separator of a cut	c	the conjugation c of section 19
Sign container $\{s : b(s) \leq \sigma\}$	T_σ	TC
Birthday	$b(s)$, the length of s	the same function; here it is an ordinal, not a surreal ordinal

Only the order of sign sequences is used in this part: no field operation and no birthday *symbol* of a structure $\mathcal{B}_\lambda$ occurs. Typography only: source 10’s sans-serif theory names ZF, ZFC and ZFU_R are set as in the rest of this report.

24.5 Relation to Part II and to the rest of the collection

Remark 24.5 (The weak-base form of [Theorem 10.4](#)). Source 10 did not know this report. Its spine is the same principle as [theorem 10.4](#) (source 09’s): *Replacement is exactly the bounding of birthdays of definable families of surreals*. The two statements live on different bases, and neither contains the other.

- [Theorem 10.4](#) is a theorem of ZFC about two set structures at an uncountable cardinal κ : the eternity scheme Et ([section 10](#)), a scheme of the birthday-field language evaluated in $\mathcal{B}_\kappa$ for families indexed by embedded ordinals, holds iff the hereditary-size universe H_κ satisfies Replacement. Its proof uses the bi-interpretation of [Part I](#) and, in the converse direction, ambient Replacement and Choice.
- Source 10’s [theorems 28.2](#) and [29.1](#) are internal scheme equivalences over the weak theory Z_f in the ordinary membership language, with sign sequences as sets and only their order used. Over Z_f alone birthday bounding is ordinal bounding OB; it becomes full Replacement (and Collection) only over $Z_f + \text{Hier}$, and no Choice is used. [Corollary 28.3](#) is the per-family form, the counterpart of a single instance of Et. New beside Et is NIB: a strict bound can always be sought inside the one interval $(-1, 0)$.
- The bases differ: H_κ need not satisfy Power Set ([lemma 10.3](#) does not list it), which Z_f contains, and Hier can fail in H_κ . Where H_κ does satisfy $Z_f + \text{Hier}$, for example at an uncountable κ with $\kappa = \beth_\kappa$, where $H_\kappa = V_\kappa$, [theorem 29.1](#) applies inside H_κ , and together with [theorem 10.4](#) it gives: $\mathcal{B}_\kappa \models \text{Et}$ iff H_κ satisfies any one of the six schemes of [theorem 28.2](#), read in H_κ . This combination is an observation of this merge, not a statement of either source.

Source 10’s $V_{\omega+\omega}$ countermodel ([proposition 30.1](#)) has the same shape as the failures recorded in [section 10](#): every member of a definable family exists, but the family cannot be collected; compare

the $\sqsupset_\omega$ instance $\text{Rep}_\sqsupset$ (theorem 11.5) and the cofinal power-tower sequence at ε_0 that defeats Et. Source 10’s Replacement-free set-cut theorem (theorem 27.1) is a different statement from the Replacement-free arithmetic certificates of section 3: the latter compute field operations inside H_κ , the former constructs the simplest separator of a set cut from the order alone.

Remark 24.6 (The same Collection spectrum in two collections). Batch-89 manuscript 05, placed as the research-report collection’s new report `SetTheory/Cardinals/docs/reports/ordinal-s-and-order-types/naming-elementary-embeddings` (*Naming Elementary Embeddings: Atom Components, Replacement, and Reflection*, pinned to the same commit; AI-assisted, unrefereed, not formalized), works with the same kernel models, written $M_\kappa(A)$ there after Yao’s ideal construction. Its lemma “No new atoms in a unique base-language output” (Lemma 2.3, `nee:lem:new`) and its “Known kernel-ideal theorem, specialized” (Theorem 2.4, `nee:thm:base`) are lemma 32.1 and theorem 32.2. Its lemmas “Automatic small-domain Collection” (Lemma 7.1, `nee:lem:smallcoll`) and “The limit-cutoff obstruction is already bounded” (Lemma 7.2, `nee:lem:limitcoll`), with its corollary “Exact local spectrum at limit cutoffs” (Corollary 7.3, `nee:cor:limitspectrum`), prove lemma 33.1 and theorem 33.2 for the same models and for their expansions by named atom injections, with the added information that the failing instance at a limit cardinal has a Δ_0 matrix. Its “Full Collection criterion” (Theorem 7.5, `nee:thm:collection`) proves full Collection at successor cardinals, as in theorem 33.3, by a reservoir argument for named tuples. The two manuscripts were written independently on the same pin; neither is claimed first, and both credit the mechanism to Hamkins–Yao, Lévy, Glazer–Yao and Yao. This part prints source 10’s proofs. That report’s dated note after its Corollary 7.6 compares the two: its failing instance at a limit cardinal collects injection graphs with a Δ_0 matrix, where source 10 collects sets of atoms of prescribed cardinalities, and its successor-case reservoir consists of whole components, so that the permutations commute with the named maps. Source 10’s enumeration predicate E (section 34) and that report’s named lifts of atom injections are different expansions of the same models, and their theorems are different.

Remark 24.7 (Full-class rank models). Batch-89 manuscript 01, placed as Part XV of the report `foundations-and-computation/surreal-well-orders` (its source 31), computes for the two-sorted full-class rank models $(V_\lambda, \mathcal{P}(V_\lambda))$ at every limit $\lambda > \omega$ the domains on which class Replacement and Collection hold: exactly those of size below $\text{cf } \lambda$. At $\lambda = \omega + \omega$ this contains the failure of proposition 30.1 for the one definable family $n \mapsto \omega + n$. The failure of NIB there, and the weak-base equivalences, are source 10’s own.

25 The base theories and the meaning of a family

25.1 Zermelo set theory without hidden Replacement

Definition 25.1. Z_f consists of Extensionality, Empty Set, Pairing, Union, Power Set, Infinity, the full Separation scheme, and Foundation, in classical first-order logic. It does not contain Replacement or Choice.

We use ordinary set encodings of ordered pairs and functions. The usual finite constructions—Cartesian products, restrictions, domains and ranges of *set relations*, and finite tagged unions—are available in this theory. For example, $X \times Y$ is obtained by Separation from a sufficiently iterated power set of $X \cup Y$. No image of an arbitrary definable class function is presupposed.

All appearances of a class in the weak-base half of source 10 abbreviate a first-order formula with finitely many set parameters. An assertion quantified over formulas is an external scheme: there is one first-order assertion for each formula. There is no set of all formulas together with an

unrestricted truth predicate for the universe, and no quantification over arbitrary subclasses of a proper class.

Definition 25.2 (Definable set-indexed family). A formula $\varphi(i, s, p)$ presents a surreal family on a set I if

$$\forall i \in I \exists! s [s \in \mathbf{No} \wedge \varphi(i, s, p)].$$

We write s_i for its unique value. This notation does not assert that $\{(i, s_i) : i \in I\}$ is a set.

The distinction is essential. If a family is given *as a set function*, its range is a set in Z_f . If it is only given by a formula satisfying the displayed condition, forming its range is a Replacement-type step.

25.2 A separately named hierarchy hypothesis

Definition 25.3 (The hierarchy axiom Hier). The following two assertions constitute Hier. First, for every ordinal α there is a set function h with domain $\alpha + 1$ satisfying

$$h(0) = \emptyset, \quad h(\zeta + 1) = \mathcal{P}(h(\zeta)), \quad h(\lambda) = \bigcup_{\zeta < \lambda} h(\zeta)$$

for the relevant successor and nonzero limit indices. Second, every set x belongs to $h(\alpha)$ for some such α, h .

The unions in this definition are unions of ranges of restrictions of a *set function*. Uniqueness and coherence of the hierarchy segments follow by induction along an ordinal. Consequently V_α is unambiguously definable under Hier.

ZF proves Hier by its usual recursion and rank constructions. We do not assume that Hier follows from Z_f . Our characterization over plain Z_f is a characterization of ordinal bounding; only after adjoining Hier do we identify it with full Replacement. The model in [section 30](#) satisfies $Z_f + \text{Hier}$ but not Replacement, so Hier itself does not trivialize the result.

25.3 Two different ambient settings

The first half of source 10 concerns pure set theory and the sign-sequence order. The later support-model construction takes place externally in a full well-founded universe with atoms and Choice. We write ZFU_R for urelement set theory formulated with Replacement, rather than building Collection into its name. This prevents an ambiguity in the literature: some authors use “ZFU” for a theory already containing Collection.

In the atom setting, Extensionality applies to sets, and atoms have no members. A tagged interpretation in ordinary ZFC can be used to implement the background urelement universe. No new large-cardinal assumption is needed for the constructions here. When a class model is mentioned, satisfaction is used formula by formula through relativization; a universal class truth predicate is not introduced.

26 Sign sequences and bounded containers

Definition 26.1. A surreal sign sequence is a function $s : \alpha \rightarrow \{-, +\}$ for an ordinal α . Its birthday is $b(s) = \alpha$. The empty sequence is 0, and the one-sign sequence $(-)$ is denoted by -1 .

We distinguish the two signs from the numerical values of these named surreals. For comparison, use the order

$$- < \text{termination} < +.$$

Two unequal sequences are compared at the first position where their signs differ or where precisely one terminates. Such a position exists and has a least ordinal index. This defines a linear order. Write $p \preccurlyeq s$ if p is an initial segment of s , possibly equal to s .

Only this ordered tree is needed. We do not invoke the recursive definitions of surreal addition, multiplication, exponentiation, or real closedness in the weak base.

Lemma 26.2 (Bounded sign container). *For every ordinal σ , the class*

$$T_\sigma = \{s \in \mathbf{No} : \mathbf{b}(s) \leq \sigma\}$$

is a set in $\mathbf{Z}_f$.

Proof. Every such function is a subset of $\sigma \times \{-, +\}$. Separate from $\mathcal{P}(\sigma \times \{-, +\})$ the graphs which are functions whose domains are ordinals. These domains are subsets of σ , hence are at most σ . Conversely every sign sequence of length at most σ occurs. This uses Power Set and Separation, not Replacement. $\square$

Lemma 26.3 (Set families have bounded birthdays). *If S is a set of sign sequences, there is an ordinal γ with $\mathbf{b}(s) \leq \gamma$ for every $s \in S$. Moreover,*

$$\lambda = \sup\{\mathbf{b}(s) + 1 : s \in S\}$$

is an ordinal available in $\mathbf{Z}_f$, and $\mathbf{b}(s) < \lambda$ for all $s \in S$.

Proof. The union $\bigcup S$ is a set relation, so its domain exists. This domain is the union of the ordinal domains of the members of S , and is therefore an ordinal γ . Every $\mathbf{b}(s)$ is at most γ . To avoid assuming that the displayed successor image is a set, define directly

$$\lambda = \{\xi \in \gamma + 1 : \exists s \in S (\xi \leq \mathbf{b}(s))\}.$$

This is a downward-closed set of ordinals, equals the indicated supremum, and has the desired strict bound. When S is empty, it is 0. $\square$

Lemma 26.4 (All-plus ordinal codes). *For an ordinal α , let o_α be the all-plus sequence of length α . Then $\alpha < \zeta$ if and only if $o_\alpha < o_\zeta$. For every sign sequence s ,*

$$s \leq o_{\mathbf{b}(s)} < o_{\mathbf{b}(s)+1}.$$

In particular, $o_\alpha < s$ implies $\alpha < \mathbf{b}(s)$.

Proof. These comparisons follow at the first negative sign, or at the first termination position when no earlier negative sign occurs. Each code is constructed directly as a constant function on its given ordinal. There is no appeal to a set containing all ordinal codes. $\square$

Lemma 26.5 (One bounded interval contains an ordinal detector). *For each ordinal α , define ν_α on $\alpha + 2$ by*

$$\nu_\alpha(\xi) = \begin{cases} -, & \xi = 0, \\ +, & 0 < \xi < \alpha + 2. \end{cases}$$

Then

$$-1 < \nu_\alpha < 0, \quad \alpha < \zeta \iff \nu_\alpha < \nu_\zeta.$$

For every negative sign sequence c of birthday σ ,

$$c < \nu_\sigma.$$

Consequently $\nu_\alpha < c < 0$ implies $\alpha < b(c)$.

Proof. The first sign makes the sequence negative, and the plus at position 1 makes it greater than -1 . Adding more trailing plus signs increases its value. Right addition by the finite ordinal 2 is strictly increasing, giving the equivalence involving α and ζ .

A negative c has a minus at position 0. If it has another minus, then ν_σ is larger at the first such position. If it has only plus signs after position 0, then it is a proper initial segment of ν_σ and the latter has a plus at its termination position. Thus $c < \nu_\sigma$. If $\alpha \geq \sigma$, monotonicity would give $\nu_\alpha \geq \nu_\sigma > c$, proving the last assertion. $\square$

27 Actual set cuts: a Replacement-free simplicity proof

Theorem 27.1 (Set-cut interpolation with a birthday bound). *In Z_f , let L, R be sets of sign sequences with $L < R$, meaning $l < r$ for every $l \in L$ and $r \in R$. Empty sides are allowed. There is a unique simplest separator c satisfying*

$$L < c < R.$$

Every separator extends c as a sign sequence. In particular, c is the unique separator of least birthday, and

$$b(c) \leq \lambda, \quad \lambda = \sup\{b(x) + 1 : x \in L \cup R\}.$$

Proof. By lemma 26.3, λ is a set ordinal, and every endpoint has birthday below λ . The proof takes place inside the set T_λ of lemma 26.2.

For a prefix p , say it has a left obstruction if some $l \in L$ satisfies $p \leq l$, and a right obstruction if some $r \in R$ satisfies $r \leq p$. The two obstructions cannot occur together: they would yield $r \leq p \leq l < r$. If neither occurs, p is a separator. Otherwise prescribe the next sign to be plus for a left obstruction and minus for a right obstruction.

Call $p \in T_\lambda$ a *good path* if, for every $\zeta < b(p)$, the restriction $p \upharpoonright \zeta$ is not yet a separator and the sign $p(\zeta)$ is the prescribed sign at that restriction. Good paths form a set G by Separation. The empty path belongs to G .

Compatibility. Any two good paths are compatible. At a first disagreement they would have the same earlier prefix but different prescribed signs, contrary to the rule's uniqueness. Hence one extends the other.

The union $p_* = \bigcup G$ is therefore a sign sequence of length at most λ . It is itself good: each of its sign positions already belongs to some good path, whose earlier restrictions coincide with its own. Thus $p_* \in G$.

We verify the invariant

$$\begin{aligned} l \in L, p \not\leq l &\implies l < p, \\ r \in R, p \not\leq r &\implies p < r \end{aligned} \tag{27.1}$$

for every good path p . It is vacuous at the empty path. Suppose the prescribed extension is p^+ , obtained by appending plus. Any left endpoint not extending p^+ either was already separated from p , equals p , or takes minus at the next position; in every case it is below p^+ . A left obstruction ensures some $l \geq p$, and all right endpoints exceed this l . Therefore a right endpoint that still extends p cannot equal p or take minus next. Right endpoints not extending p remain separated at their earlier comparison position. This proves the invariant for p^+ . The minus case is symmetric.

At a limit path, an endpoint not extending that path already fails to extend some shorter prefix: use a position past its first disagreement or its termination. The invariant at that prefix persists in all later extensions. Induction along the set ordinal $\lambda + 1$ proves (27.1).

If $b(p_*) < \lambda$ and p_* were not a separator, appending its prescribed sign would yield a good path longer than the union of all good paths, a contradiction. If $b(p_*) = \lambda$, no endpoint can extend p_* , since every endpoint has smaller birthday. The invariant then says that p_* is a separator. Set $c = p_*$.

Finally let q be any separator. Induction along a good path shows that q extends that path. At a plus step, an extension of the current prefix that terminates or takes minus is at most the prefix and hence at most the obstructing left endpoint; it cannot be q . The minus step is symmetric. At limits, extension follows by union. Thus q extends c . Since c itself separates, it is the unique separator of least birthday. $\square$

Remark 27.2 (Why this is not an invocation of class recursion). The proof does not first build a class-length recursion and then take its image. It separates *all good approximations from one set container* and takes their union. Every induction is along a given set ordinal. The set-existence steps are precisely those available in $\mathbf{Z}_f$.

Proposition 27.3 (Sharpness). *The birthday bound in theorem 27.1 cannot be uniformly lowered. For a nonzero successor $\lambda = \zeta + 1$, the cut $(\{o_\zeta\}, \emptyset)$ has simplest separator o_λ . For a nonzero limit λ , the cut*

$$L = \{o_\zeta : \zeta < \lambda\}, \quad R = \emptyset$$

has simplest separator o_λ .

Proof. In the successor case, a separator of birthday at most ζ is excluded by lemma 26.4, and $o_{\zeta+1}$ works. In the limit case, L is a set by Separation in T_λ ; it is not being formed by an unbounded image operation. Any separator c must have birthday at least λ , because $o_\zeta < c$ gives $\zeta < b(c)$ for every $\zeta < \lambda$. Again o_λ works. The empty cut has simplest separator 0 and bound 0. $\square$

28 The equivalence theorem for definable surreal families

28.1 Six schemes

All formulas below may have arbitrary set parameters in the language under consideration.

Definition 28.1. The schemes are as follows.

- (1) **OB**: every definable function $i \mapsto \alpha_i$ from a set into the ordinals has an ordinal γ such that $\alpha_i < \gamma$ for all indices.
- (2) **Gr**: every definable set-indexed surreal family has a set graph.
- (3) **UB**: every definable set-indexed surreal family has a strict surreal upper bound.
- (4) **SC**: if I is a set and $\forall i \in I \exists s \in \mathbf{No} \varphi(i, s, p)$, there is a set $S \subseteq \mathbf{No}$ with $\forall i \in I \exists s \in S \varphi(i, s, p)$.

- (5) DCI: if two definable families $(l_i)_{i \in I}$ and $(r_j)_{j \in J}$ have set index domains and satisfy $l_i < r_j$ for all i, j , there is a surreal c with $l_i < c < r_j$ for all i, j .
- (6) NIB: every definable set-indexed family of elements of $(-1, 0)$ has a strict upper bound that also lies in $(-1, 0)$.

The right bound 0 is not an acceptable witness to NIB. This is why the last scheme is stronger than the tautological assertion that a family in $(-1, 0)$ is bounded in **No**.

Theorem 28.2 (Exact weak-base equivalence). *Over $\mathbf{Z}_f$,*

$$\boxed{\text{OB} \iff \text{Gr} \iff \text{UB} \iff \text{SC} \iff \text{DCI} \iff \text{NIB}.}$$

Proof. $\text{OB} \Rightarrow \text{Gr}$. Given a definable family s_i , its birthdays are uniquely defined ordinals. Choose γ bounding them. Every value lies in the set T_γ . The graph is now a subset of $I \times T_\gamma$, defined by the original formula, and exists by Separation.

$\text{Gr} \Rightarrow \text{UB}$. The range of the set graph is a set S . By lemma 26.3, all birthdays are at most an ordinal γ . The all-plus sequence $o_{\gamma+1}$ strictly bounds every member of S .

$\text{UB} \Rightarrow \text{OB}$. Apply UB to the definable family o_{α_i} . If c strictly bounds it, lemma 26.4 gives $\alpha_i < b(c)$ for every i .

$\text{OB} \Rightarrow \text{SC}$. For each i , let α_i be the least birthday of a surreal witness to $\varphi(i, s, p)$. The least ordinal exists without a choice of witnesses for the whole family: for each fixed i , one witness supplies an ordinal interval in which Separation and well-ordering locate a minimum. The definition of α_i is functional, so OB gives $\alpha_i < \gamma$ uniformly. The set T_γ is then a collecting set of possible witnesses. If desired, separate from it the witnesses relevant to at least one index. No choice function selecting a particular witness is needed.

$\text{SC} \Rightarrow \text{Gr}$. Apply SC to a functional surreal family. Its unique values all lie in a set S . Separate the graph inside $I \times S$.

$\text{Gr} \Rightarrow \text{DCI}$. Obtain set graphs and set ranges for the two families. Apply theorem 27.1 to these actual sets.

$\text{DCI} \Rightarrow \text{NIB}$. For a nonempty family in $(-1, 0)$, use it as the left family and use the constant right family $\{0\}$. Any separator is automatically greater than -1 . For an empty family, ν_0 works. Equivalently one can adjoin -1 to the left family using a finite tagged union of index sets.

$\text{NIB} \Rightarrow \text{OB}$. Given ordinal values α_i , apply NIB to ν_{α_i} . Its negative upper bound c yields $\alpha_i < b(c)$ by lemma 26.5. $\square$

Corollary 28.3 (The obstruction for an individual family). *For a given definable set-indexed surreal family, the following are equivalent in $\mathbf{Z}_f$: its graph is a set; its range is contained in a set of sign sequences; and its birthdays admit an ordinal bound. Mere numerical boundedness of its values is not equivalent to these conditions.*

Proof. A set graph has a set range and hence bounded birthdays. A birthday bound provides the container needed to separate the graph. The counterexample to the last possible equivalence is given in section 30. $\square$

Remark 28.4 (Restricted domains and restricted formulas). The proof preserves the index domain in most implications. The passage from a negative interval to a cut uses only a singleton right side. Thus the ordinal-detector implications already hold for a fixed domain such as ω . One must not conclude that the ω -restricted scheme implies Replacement on *all* domains. Similarly, the

transformations of formulas are explicit, but this article does not assert optimal preservation of their position in the Lévy hierarchy.

29 When the test becomes full Replacement

Theorem 29.1 (Ranked-base characterization). *Over $Z_f + \text{Hier}$, each of the six schemes in [theorem 28.2](#) is equivalent to full Replacement and to full Collection. No Choice is required.*

Proof. Replacement forms the set of ordinal outputs of a definable function and therefore gives OB.

Conversely assume OB and consider a total relation

$$\forall i \in I \exists y \varphi(i, y, p).$$

By Hier, for each i there is some ordinal α such that a suitable witness lies in V_α . Define

$$\alpha_i = \min\{\alpha : \exists y \in V_\alpha \varphi(i, y, p)\}.$$

This is a uniquely defined ordinal, even when the witness is not unique. As before, a least element of the nonempty ordinal class is obtained by bounding the minimization with one witness and using Separation. By OB, some γ bounds all α_i . Monotonicity of the cumulative hierarchy shows that V_γ collects witnesses for every i . This proves Collection.

Finally, for a functional relation, Collection puts the unique outputs inside a set, and Separation removes extraneous members and forms the graph. Thus Collection implies Replacement. Combining with [theorem 28.2](#) proves the assertion. $\square$

The precise conclusion. Actual set-cut interpolation is a theorem of Z_f . Definable set-indexed cut interpolation is equivalent to ordinal bounding over Z_f . It is equivalent to full Replacement over the *explicitly stronger base* $Z_f + \text{Hier}$.

This is an exact diagnosis of a common informal shortcut. The expression $\{s_i : i \in I\}$ cannot be inserted into the ordinary cut construction until its existence as a set has been justified. In a rank-complete pure universe, justifying that step for all formulas has the full strength of Replacement.

30 An explicit omitted cut in a rank-initial universe

We now work externally in ZFC. Let

$$\theta = \omega + \omega, \quad \text{and consider the structure } (V_\theta, \in).$$

All assertions about V_θ below refer to its internal membership structure. In particular, “a set” in V_θ means an element of V_θ .

Proposition 30.1. *V_θ satisfies $Z_f + \text{Hier}$ and Choice, but fails OB, Replacement, and NIB.*

Proof. The ordinal θ is a limit and is closed below itself under adding any fixed finite ordinal. Pairing, Union, and Power Set increase the ranks of the finitely many input objects by only a finite amount, so their required outputs remain below θ . Every subset of an element of V_θ is again in V_θ ,

giving Separation for formulas interpreted in V_θ . Infinity and Foundation hold, and Extensionality is inherited from transitivity.

For every $\alpha < \theta$, the hierarchy segment $\langle V_\zeta : \zeta \leq \alpha \rangle$ has rank at most α plus a fixed finite coding overhead, and therefore belongs to V_θ . Every member of V_θ belongs to a sufficiently high stage below θ . This gives **Hier**. An ambient well-ordering of any $x \in V_\theta$ is a relation on x of rank bounded by the rank of x plus a finite amount, so it also belongs to V_θ . Hence V_θ satisfies **Choice**.

The formula $n \mapsto \omega + n$, for $n \in \omega$, defines a total ordinal-valued family in V_θ . Every value exists there, but these values are cofinal in $\theta = \text{Ord}^{V_\theta}$. They have no internal ordinal bound, so **OB** fails. Replacement fails as well.

Finally define $s_n = \nu_{\omega+n}$. Each individual sign sequence has birthday $\omega + n + 2$ and, with ordinary pair coding, rank below θ . Thus it belongs to V_θ , and V_θ recognizes

$$\forall n \in \omega \quad -1 < s_n < 0.$$

Given any negative $c \in \mathbf{No}^{V_\theta}$, its birthday σ is below θ . Choose n with $\omega + n \geq \sigma$. By [lemma 26.5](#),

$$c < \nu_\sigma \leq \nu_{\omega+n} = s_n.$$

There is therefore no negative strict upper bound for this definable family. This refutes **NIB** and presents an omitted definable cut with right side $\{0\}$. $\square$

Remark 30.2 (What is and is not missing). The family is externally countable. Internally, however, it is only a formula assigning one existing object to each natural number; its graph is not an element of V_θ . Every *actual set cut* in V_θ still has an interpolant by [theorem 27.1](#). Also, 0 is an upper bound for the displayed family. What fails is an upper bound *strictly below* 0, not ordinary numerical boundedness.

This countermodel shows that the distinction survives **Choice**, the existence of all internal hierarchy stages, and the existence of each individual member of the family. The omitted resource is the ability to collect the entire definable family into one set.

[Added 4 October 2026, batch 89: the class form of this failure at every limit height, recorded in [remark 24.7](#). Part XV of [foundations-and-computation/surreal-well-orders](#) (its source 31, batch-89 manuscript 01) uses the same witness $n \mapsto \omega + n$ at $\lambda = \omega + \omega$ (its Section XV.11, [swo:gcz:sec:examples](#)). In the two-sorted full-class rank model $(V_\lambda, \mathcal{P}(V_\lambda))$, for every limit $\lambda > \omega$, class Replacement and class Collection on an internal set a hold exactly when $|a| < \text{cf } \lambda$ (its Lemma XV.9.2, [swo:gcz:lem:domain](#)), and full class Replacement holds exactly when λ is strongly inaccessible (its Theorem XV.9.3, [swo:gcz:thm:replacement](#)). The failure of Replacement above is the definable case at a height of cofinality ω ; the failure of **NIB** is source 10's own. Neither manuscript cites the other.]

31 Urelements: rank bounds versus atom-base bounds

For an object x in a full well-founded urelement universe, its kernel $\ker(x)$ is the set of atoms occurring in the transitive closure of $\{x\}$. An object is pure if its kernel is empty. For a set B of atoms define the hierarchy

$$V_0(B) = B, \quad V_{\alpha+1}(B) = \mathcal{P}(V_\alpha(B)) \cup B,$$

with unions at nonzero limit stages, and put $V(B) = \bigcup_{\alpha \in \text{Ord}} V_\alpha(B)$ as a class abbreviation. An object lies in $V(B)$ exactly when its kernel is contained in B .

There are now two independent resources: the height α and the atom base B . In the pure case the base is always empty. With atoms, an ordinal bound on heights need not control the supports of witnesses.

Proposition 31.1 (A two-resource witness criterion). *Suppose the required set hierarchies over atom bases and kernels exist. For a total relation on a set I , the following are equivalent:*

- (i) *there is a set collecting witnesses;*
- (ii) *there are a set B of atoms and an ordinal γ such that every index has a witness in $V_\gamma(B)$.*

If ordinal bounding is available, condition (ii) is equivalent to the existence of a single atom base B such that every index has some witness in $V(B)$.

Proof. A collecting set W has a kernel $B = \ker(W)$ and belongs to some $V_\eta(B)$. Its members then all lie in a sufficiently high common stage over B , giving (ii). Conversely $V_\gamma(B)$ itself is a collecting set.

For the last assertion, fix a common base B and assign to each index the least ordinal stage over B containing a witness. This assignment is functional. Ordinal bounding bounds all its values, producing a common γ . The reverse implication is immediate. $\square$

In ZFUR , Replacement supplies ordinal bounding. The resulting common-base characterization is precisely the mechanism highlighted by Glazer–Yao, Proposition 1 [21]. Our use of it is to separate what a pure surreal detector can see from what it cannot: birthdays measure ordinal height, whereas pure sign sequences carry no atom kernel at all.

32 Cardinal-support models and their basic axioms

Fix a full urelement universe $\mathcal{U}$ with Choice, containing a set A of all its atoms. Let κ be an infinite cardinal with $|A| \geq \kappa$, with cardinalities computed in $\mathcal{U}$. Define the class

$$M_\kappa^{\ker} = \{x \in \mathcal{U} : |\ker(x)| < \kappa\}.$$

It contains every atom individually and every pure set, but does not contain the set of all atoms. Thus its atoms form a proper class *internally*, even though A is a set in the ambient universe.

The restriction is on kernel cardinality, not on the cardinality or rank of the whole object. For a permitted base B , every stage $V_\alpha(B)$, at every ordinal height, belongs to $M_\kappa^{\ker}$.

Lemma 32.1 (Kernel confinement of a unique output). *Suppose $w, p \in M_\kappa^{\ker}$, $x \in w$, and a formula has a unique output $y \in M_\kappa^{\ker}$ for x, p , when interpreted in $M_\kappa^{\ker}$. Then*

$$\ker(y) \subseteq K, \quad K = \ker(w) \cup \ker(p).$$

Proof. Every permutation of A extends by rank recursion to an automorphism of $\mathcal{U}$ and preserves $M_\kappa^{\ker}$, because it preserves kernel cardinalities. Suppose $a \in \ker(y) \setminus K$. As κ is infinite, the union $K \cup \ker(y)$ has size below κ . Choose an atom b outside this union. The transposition swapping a and b fixes K pointwise, hence fixes x and p . It moves $\ker(y)$ and therefore moves y .

The image of y is another output satisfying the same formula in $M_\kappa^{\ker}$, by invariance under this automorphism. This contradicts uniqueness. $\square$

Theorem 32.2 (The base support model). *For every such infinite κ , $M_\kappa^{\ker}$ is a model of $\text{ZFUR} + \text{AC}$. It is transitive, has every ambient subset of each of its sets, and has exactly the same pure sets as $\mathcal{U}$.*

Proof. Membership can only decrease kernels, so the class is transitive. Subsets do not increase kernels; hence every ambient subset of an internal set is internal, and the ambient power set of an internal set is itself internal. Pairing and finite unions use only finite unions of kernels of size below an infinite κ , again of size below κ . Union similarly adds no atoms beyond those already in the input's kernel. Extensionality, Foundation, and Infinity follow immediately. Separation follows by relativizing a fixed formula to the definable class M_κ^{ker} and taking a subset of the given set.

For Replacement, suppose a formula assigns a unique internal output to every member of a set w , with parameter tuple coded by p . Ambient Replacement forms its range W ; relativization to M_κ^{ker} is a legitimate fixed ambient formula. By lemma 32.1, every output has kernel contained in $K = \ker(w) \cup \ker(p)$. Therefore $\ker(W) \subseteq K$ and $W \in M_\kappa^{\text{ker}}$. This proves every internal Replacement instance.

For Choice, an ambient well-ordering of a set $w \in M_\kappa^{\text{ker}}$ has kernel contained in $\ker(w)$, so that well-ordering belongs to M_κ^{ker} . Finally every pure set has empty kernel and is included; no impure object becomes pure by passing to this transitive model. $\square$

Lemma 32.3 (Agreement about cardinalities of internal sets). *For every $w \in M_\kappa^{\text{ker}}$, its cardinality as computed in M_κ^{ker} equals its ambient cardinality. Cofinalities of pure ordinals also agree.*

Proof. An ambient bijection between w and its initial ordinal has kernel contained in $\ker(w)$, so it is internal. Conversely an internal bijection is an ambient bijection. For cofinality, every function between pure ordinals is pure, so the two universes have exactly the same candidate cofinal maps. $\square$

33 The exact Collection spectrum

Write $\text{Collection}_{<\tau}$ for the scheme restricting index sets to cardinality below τ . Saying that Collection fails at τ means that one explicit formula has a total witness relation on a domain of cardinality τ with no collecting set. It is not a quantification over a set of all schemes.

Lemma 33.1 (The small-domain positive bound). *For every infinite κ , M_κ^{ker} satisfies $\text{Collection}_{<\text{cf } \kappa}$.*

Proof. Let $w \in M_\kappa^{\text{ker}}$ have size $\lambda < \text{cf } \kappa$, and suppose every $x \in w$ has an internal witness to a fixed formula. In the ambient universe, Collection followed by Choice gives a family of internal witnesses y_x . Each has a kernel of size below κ . A union of fewer than $\text{cf } \kappa$ sets, each of size below κ , has size below κ ; this is the usual cofinality calculation in the ambient universe with Choice. Consequently

$$|\ker(\{y_x : x \in w\})| \leq \left| \bigcup_{x \in w} \ker(y_x) \right| < \kappa.$$

The selected range is therefore an internal collecting set. By lemma 32.3, the domain-size hypothesis is the same internally and externally. $\square$

Theorem 33.2 (Limit-cardinal failure at the exact threshold). *Let κ be a limit cardinal, with ω included in this terminology, and set $\tau = \text{cf } \kappa$. Then*

$$M_\kappa^{\text{ker}} \models \text{Collection}_{<\tau}, \quad M_\kappa^{\text{ker}} \not\models \text{Collection}_\tau.$$

For every cardinal $\lambda \geq \tau$, Collection also fails on a domain of size λ .

Proof. The positive assertion is [lemma 33.1](#). Choose a pure sequence of cardinals $(\mu_i)_{i < \tau}$ below κ , cofinal in κ . For $\kappa = \omega$, use finite cardinals. The sequence and its domain belong to M_κ^{ker} because they are pure.

Consider the relation

$$\varphi(i, B) : \quad B \text{ is a set of atoms and } |B| = \mu_i.$$

Every $i < \tau$ has a witness: select an ambient set $B \subseteq A$ of size μ_i . Both B and a bijection witnessing its cardinality have kernels of size below κ , so they are internal.

If $W \in M_\kappa^{\text{ker}}$ collected such witnesses, put $\rho = |\ker(W)| < \kappa$. Some μ_i exceeds ρ . But every atom set $B \in W$ satisfies $B \subseteq \ker(W)$ and cannot have size μ_i . This is a contradiction.

For a larger domain, extend the same relation by giving a harmless fixed witness, such as the empty set, to the additional indices. A collecting set for the enlarged relation would collect witnesses for the original one. $\square$

Theorem 33.3 (The successor-cardinal case). *If $\kappa = \mu^+$ for an infinite cardinal μ , then M_κ^{ker} satisfies full Collection, with no restriction on the size of its index set.*

Proof. Let a formula with parameter p have a witness for every $x \in w$. Set $K = \ker(w) \cup \ker(p)$; its size is at most μ . Choose a set $B \subseteq A \setminus K$ of size μ . This is possible because $|A| \geq \mu^+$.

Given one internal witness y , its kernel outside K has size at most μ . There is a permutation of A fixing K pointwise and sending these remaining atoms into B . To justify extension to a permutation, choose an injection onto a subset of B and biject the two complementary parts of $A \setminus K$; both complements have cardinality $|A|$. The resulting automorphism fixes x, p and preserves M_κ^{ker} . Thus an appropriate witness exists in $V(K \cup B)$ for every x .

For each x take the least stage of this fixed-base hierarchy containing a witness. Replacement in M_κ^{ker} , already proved in [theorem 32.2](#), bounds these ordinal stages. A sufficiently high set $V_\gamma(K \cup B)$ belongs to M_κ^{ker} and collects all the required witnesses. $\square$

Corollary 33.4 (Complete capacity classification). *For an infinite cardinal κ , M_κ^{ker} satisfies full Collection if and only if κ is a successor cardinal. If it is not, the least domain cardinality at which Collection fails is $\text{cf } \kappa$.*

The distinction is not simply regular versus singular. For example, ω is regular but the finite-support model fails countable Collection. A regular uncountable limit cardinal has failure first at that cardinal. A successor cardinal, by contrast, has a largest permitted infinite support size, and the common-reservoir argument works for domains of arbitrary cardinality.

33.1 Prescribing the first failure

Corollary 33.5. *For every infinite regular cardinal τ , there is a cardinal-support model whose least Collection-failure domain has size τ , while its pure universe is unchanged from the ambient one.*

Proof. Choose $\kappa = \aleph_\tau$ and an ambient set of at least κ atoms. Since τ is an infinite regular ordinal, κ is a limit cardinal with $\text{cf } \kappa = \tau$. The latter equality follows from continuity and strict increase of the aleph sequence: an index-cofinal sequence gives the upper bound, while a shorter cofinal sequence of ordinals below $\aleph_\tau$ would yield a shorter cofinal sequence of indices below τ . Apply [theorem 33.2](#). $\square$

34 Naming an enumeration: exact failure of Replacement

Fix $A_0 \subseteq A$ of size κ , and an ambient bijection

$$e : \kappa \longrightarrow A_0.$$

Its graph has kernel A_0 , so it is not a set of M_κ^{ker} . Nevertheless it can be named as an external binary predicate E on the domain of that model. We analyze the expansion $(M_\kappa^{\text{ker}}, E)$ with *Separation and other schemes understood in the expanded language*.

This is a controlled instance of language sensitivity. It is not a reproof of Glazer’s global-choice theorem, which concerns a different pure-set situation [20]. The shared lesson is that adjoining a predicate can add functional definitions to which a scheme must apply.

Proposition 34.1 (Elementary axioms survive the expansion). *The expansion $(M_\kappa^{\text{ker}}, E)$ satisfies the urelement versions of $\mathbf{Z}_f$ and Choice, including full Separation for formulas using E .*

Proof. The structural axioms and Choice are unchanged. For an expanded-language formula, replace E by its ambient set graph and relativize quantifiers to M_κ^{ker} . Ambient Separation gives a subset of the specified internal set. Every such subset is internal by [theorem 32.2](#), so expanded Separation holds. $\square$

Theorem 34.2 (The exact expanded-language threshold). *For every infinite κ , let $\tau = \text{cf } \kappa$. In $(M_\kappa^{\text{ker}}, E)$, both Collection and Replacement hold for all domains of cardinality below τ , and both fail on a domain of cardinality exactly τ .*

Proof. The positive argument in [lemma 33.1](#) does not use invariance under atom permutations. It uses only ambient Collection and Choice to gather fewer than $\text{cf } \kappa$ witnesses, followed by a kernel-size calculation. It therefore works unchanged for formulas involving the ambient predicate E . This gives Collection, and hence Replacement, on all smaller domains.

Choose a pure strictly increasing cofinal sequence $(\alpha_i)_{i < \tau}$ in κ . The sequence is an internal set and may be used as a parameter. The expanded language defines, for each $i < \tau$, the unique set

$$Y_i = e \text{ “ } \alpha_i.$$

Each individual Y_i belongs to M_κ^{ker} because $|\alpha_i| < \kappa$. Its characterization uses E and the cofinal sequence, not an internally existing graph of e .

If the range $\{Y_i : i < \tau\}$ were an internal set, its union would be the set A_0 , since the α_i are cofinal in κ . But A_0 has kernel cardinality κ and is not internal. Replacement therefore fails for this functional family. A collecting set for these unique witnesses would have kernel containing their union A_0 , so Collection fails for the same family. $\square$

Example 34.3 (Countable supports). Let $\kappa = \aleph_1$. The reduct $M_{\aleph_1}^{\text{ker}}$ satisfies full Collection. After naming an enumeration of $\aleph_1$ atoms, all countable-domain Collection and Replacement instances still hold, but both schemes fail at domain size $\aleph_1$. No pure set, sign sequence, or ordinary set function between pure objects has changed.

Example 34.4 (Singular support capacity). Let $\kappa = \aleph_\omega$. In the reduct, Replacement holds on all domains but countable Collection fails. In the expansion, the sets $e \text{ “ } \aleph_n$ form a uniquely specified countable family with no internal range set. Thus countable Replacement now fails as well.

Capacity	Replacement in M_κ^{ker}	First Collection failure in M_κ^{ker}	First failure of both in $(M_\kappa^{\text{ker}}, E)$
ω	All domains	ω	ω
$\aleph_1$	All domains	None	$\aleph_1$
$\aleph_\omega$	All domains	ω	ω
$\aleph_{\omega_1}$	All domains	ω_1	ω_1
General limit κ	All domains	cf κ	cf κ
General successor κ	All domains	None	κ

Here “first failure” refers to the cardinality of the index domain. The table is an exact consequence of the preceding proofs, not a numerical experiment.

35 Why pure surreal structures cannot see the support obstruction

Proposition 35.1 (Identity of pure structures). *The pure universe of M_κ^{ker} is the pure universe of $\mathcal{U}$. Therefore their sign-sequence classes, birthdays, prefix relations, ordinal codes, and all set-coded relations between pure objects are identical. Any specified operation defined solely within the pure universe has the same interpretation in both.*

Proof. Every pure object has empty kernel and is internal, and membership on these objects is unchanged. Every set made entirely from pure objects is itself pure. Thus the two pure membership structures are identical. Induction on a formula whose quantifiers are restricted to pure objects proves agreement on all such definitions. Apply this to sign sequences and the stated relations. $\square$

This assertion is deliberately restricted to pure definitions. A formula with unrestricted quantification over impure objects can have different truth values in the two universes, even when its free variables happen to be pure.

Theorem 35.2 (Pure-valued Collection survives definable expansions). *In M_κ^{ker} , expanded by any fixed finite list of ambient-definable predicates, Collection holds for every total relation whose witnesses are required to be pure. Index sets and parameters may be impure. In particular, this holds in $(M_\kappa^{\text{ker}}, E)$.*

Proof. Let w be an internal set, and suppose that for each $x \in w$ some pure y satisfies a fixed formula in the expansion. Relativize that formula and the added predicates to the ambient universe. Ambient Collection produces a set of suitable witnesses. Separate away any extraneous nonpure members, obtaining a set S of pure witnesses covering every index. A set of pure objects is pure, so $S \in M_\kappa^{\text{ker}}$. This same S witnesses the original internal Collection instance. $\square$

Corollary 35.3 (All six surreal schemes survive). *Every scheme OB, Gr, UB, SC, DCI, NIB holds in $(M_\kappa^{\text{ker}}, E)$, including its instances using E and arbitrary internal parameters, despite the failure of full Replacement in [theorem 34.2](#).*

Proof. Surreal sign sequences and ordinals are pure. Hence [theorem 35.2](#) gives SC and collects ordinal witnesses. Set graphs can then be separated using expanded Separation; their kernels are contained in the kernel of the index domain. The pure range sets have pure ordinal birthday bounds and pure interpolants, all of which are internal. The arguments of [theorem 28.2](#) apply directly to these sign sequences. $\square$

There is no contradiction with [theorem 29.1](#). The hierarchy axiom used there is a cofinal *pure* hierarchy covering every object of an atom-free universe. No pure V_α contains an atom. In the urelement setting, the varying atom base is an additional resource not controlled by ordinal bounding.

Corollary 35.4 (No pure-surreal characterization of ambient Collection). *Across urelement models of the kind constructed here, no statement or scheme depending only on the pure surreal structure can characterize full ambient Collection. Even the enlarged package of all definable surreal-valued Collection and interpolation instances does not characterize it. Across the displayed predicate expansions, that package does not characterize full ambient Replacement either.*

Proof. Within an ambient universe with enough atoms, compare a successor-capacity model and a limit-capacity model. Their pure universes, and hence their pure surreal structures, are identical, while full Collection holds in the former and fails in the latter. Both have all the surreal schemes by Replacement or [theorem 35.2](#).

For Replacement, compare the same M_κ^{ker} with an empty interpretation of the extra predicate and with the enumeration interpretation E . The first expansion is definitionally equivalent to the reduct and satisfies full Replacement. The second fails it by [theorem 34.2](#), while satisfying all the surreal schemes by [corollary 35.3](#). $\square$

35.1 Reflection and the limit of the conclusion

Full first-order reflection to transitive sets implies Collection: reflect simultaneously the totality assertion for a witness relation and the relation itself, with the index set and parameters included. The reflecting set then supplies genuine witnesses for all indices. This standard implication is also explicit in Glazer–Yao’s analysis [21]. Thus our limit-capacity models fail this full reflection principle.

We do not obtain Glazer–Yao’s finer separations between partial reflection, supertransitive partial reflection, and full reflection. Nor do we exclude a useful *mixed* surreal–atom theory. Naming atom kernels or permitting atom-labelled objects can expose exactly the data that the pure structure forgets. The obstruction concerns the proposed pure detector, not the possibility of any formalization of urelement set theory.

36 A formalization route for ProveIt

36.1 Repository alignment and inspection boundary

The inspected tree snapshot was identified by commit

2b7b388ba81a3355b19a7c2d2fe797e92f572355.

The relevant repository locations are:

Location	Role in a future development
Algebra/SurrealNumbers/Surreal/Foundations/ /	Sign-sequence and cut infrastructure; connect the order-level result to the existing surreal representation.
SetTheory/ZF/	Object-language formulas, theories, and model semantics.
SetTheory/ClosureAxiomatization/	Example of maintaining both semantic directions and a deductive equivalence.
Algebra/SurrealNumbers/docs/foundations-and-computation/surreal-fields-across-universes/	Related but distinct universe-comparison report; useful for terminology and avoiding duplicate claims.

This was a targeted README and directory inspection, not a full source audit or a successful build of the repository. The repository’s existing kernel-audit claims do not transfer to the present manuscript. No purportedly checked declaration name is invented here.

[Added 3 October 2026, batch 89: the four locations exist unchanged at the placement commit 23adb85f9; the commit 2b7b388ba named above is the commit “Catalogue batches 81 to 86”. Source 10 did not inspect this report, whose Part II proves the cardinal-cutoff form of its spine (remark 24.5). Placement beside the Lean libraries confers no formal status: no statement of this part is formalized, and none of the module names below exists in the repository.]

36.2 Separate object-level strength from host-level resources

A theorem about ordinary Lean types can use host-level function types, images, ordinal recursion, and classical choice without recording whether these correspond to Replacement or Choice inside a weak object theory. Consequently proving a cut theorem for a universe-polymorphic surreal type is not by itself a formal proof that Z_f proves that theorem.

The proposed formalization should have two layers. The first is a conventional order-theoretic library proving the sign comparisons and the finite-container construction. The second interprets the same constructions in arbitrary models of the explicitly listed weak axioms. A function type in the host must not silently substitute for an internal set graph.

In particular, “ φ defines a functional family” and “there is an internal graph for φ ” should be different predicates. Their equivalence is one of the conclusions under investigation, not an implementation convenience.

36.3 Suggested modules and proof order

The following are proposed module names, not existing files.

1. **WeakSignSequences**: internal function graphs on ordinals, lexicographic comparison, prefix, bounded containers, and all-plus and negative ordinal codes.
2. **SetCutSimplicity**: the set of good paths, compatibility, its union, the separation invariant, and the unique simplest interpolant with its birthday bound.
3. **DefinableFamilySchemes**: syntactic transformations implementing the six implications, with explicit parameters and index domains.
4. **RankedReplacement**: the hierarchy axiom, least-stage witnesses, and the equivalence with Collection and Replacement.
5. **AtomKernelModels**: tagged urelement semantics, kernel operations, permutation action, and the proof of kernel confinement.

6. **CapacitySpectra**: the small-domain union bound, cofinal support-size obstruction, and the successor reservoir argument.
7. **PredicateExpansion**: interpretation of an external enumeration, expanded Separation, exact failure families, and pure-valued Collection.

The set-cut lemma should be completed before importing any full-ZF surreal construction into the weak-base argument. Its proof has a natural bounded interface: a fixed ordinal λ and the set T_λ .

36.4 Trust boundaries and quantification

A satisfactory formal artifact would explicitly audit at least four boundaries. First, it would specify the object theory rather than merely a list of host axioms. Second, it would distinguish each standard formula instance from a single internal assertion quantifying over formula codes. Third, it would not represent all subsets of a proper class by a host powerset. Fourth, it would distinguish the original language from its expansion by E when stating Separation and Replacement.

For the support models, one can formalize an external set-sized model and its interpreted submodel, or a definable class interpretation in a stronger metatheory. The theorem is relative: it constructs a model from an ambient model. It is not an assertion that a consistent first-order theory proves its own consistency.

37 Executable finite checks and reproducibility

The accompanying program, shipped as `code/10-cuts-replacement-verify_finite.py` (delivered as `code/verify_finite.py`), uses only the Python standard library. It represents a finite sign sequence by a tuple of -1 and $+1$, with termination compared as 0 . It implements the obstruction rule from [theorem 27.1](#), then independently enumerates candidates to check separation, unique minimum birthday, and the prefix property.

The recorded run, shipped as `data/10-cuts-replacement-finite_checks.json` (delivered as `data/finite_checks.json`), completed with the following results:

Check	Number
Sign words of birthday at most 6	127
Endpoint intervals, including empty sides	8 256
Assignments to left/right/neither at depth 2	2 187
Separated assignments among these	576
Negative-code comparisons and birthday implications	1 050
Atom-transposition checks	4 032

Every test passed. Endpoint intervals suffice for all finite sets of bounds from the depth-six endpoint universe, since a nonempty finite left set has a maximum and a nonempty finite right set has a minimum. The depth-two assignment check separately exercises the implementation on nonsingleton sets.

The atom test verifies the local combinatorial fact used by the confinement proof: a swap outside a fixed parameter support fixes that support while moving a witness support containing exactly one of the swapped atoms. It does not simulate a model of set theory or a singular cardinal.

In the delivered layout the run was

```
python3 code/verify_finite.py --output data/finite_checks.json
```

followed by two `pdflatex` passes of the delivered manuscript. With the shipped names, run the program on a copy and pass an explicit output path outside this directory (the report’s `README.md` gives the commands). For this merge it was rerun with Python 3.14.4 on a copy: it passed and reproduced the recorded file exactly, apart from line endings. The tests cannot establish a transfinite induction, an axiom-scheme equivalence, an internal cardinal calculation, or a relative-consistency claim. Those depend on the mathematical proofs above. The delivered archive also contained a `README`, a proof-status note and a build script; the proof-status note is shipped as `10-cuts-replacement-PROOF_STATUS.md` and the build script as `code/10-cuts-replacement-build.sh` (it builds the delivered manuscript in the delivered layout, not this report), while the delivery `README`, the manuscript and its PDF are not shipped ([section 39](#)).

38 Further research questions and proposed projects

The questions below are a research agenda arising from the proved results. Their status is “not resolved in this manuscript”; no claim is made that every formulation is globally new or currently open in the literature.

Question 38.1 (Remove or characterize the hierarchy hypothesis). Over plain Z_f , determine the exact relationship between OB and full Replacement without assuming Hier. Identify the weakest rank-coverage or tier-coverage principle that turns the equivalence in [theorem 28.2](#) into full Replacement, and give explicit models distinguishing the relevant choices of coverage axiom.

The point is not to omit Hier typographically. A useful answer must analyze the existence of ranks and hierarchy segments inside the weak model.

Question 38.2 (Formula-complexity accounting). Determine the optimal correspondence between restricted Σ_n or Π_n instances of ordinal bounding, surreal graph existence, and interval bounding. In particular, quantify the complexity increase caused by “least birthday of a witness” and by the good-path definition.

A precise answer would turn the full-scheme result into a hierarchy of proof-theoretic characterizations.

Question 38.3 (Domain spectra in weak pure models). For models of $Z_f + \text{Hier}$, classify the index-domain sizes on which the six equivalent principles hold. Which initial segments of cardinalities can occur, and how are they constrained by definable cofinal maps into the model’s ordinals?

The $V_{\omega+\omega}$ example gives failure already on ω , but the general domain spectrum deserves a separate analysis. Internal and external cardinalities must be kept distinct.

[Added 3 October 2026, batch 89: for the two-sorted *full-class* rank models $(V_\lambda, \mathcal{P}(V_\lambda))$ the domain spectrum of class Replacement and Collection is computed in Part XV of `foundations-and-computation/surreal-well-orders`: exactly the domains of size below $\text{cf } \lambda$ ([remark 24.7](#)). The question above, for the *definable* schemes in arbitrary models of $Z_f + \text{Hier}$, is not answered there and stays open.]

Question 38.4 (Beyond cardinality ideals). Replace $|\ker(x)| < \kappa$ by membership of the kernel in a permutation-invariant ideal $\mathcal{I}$ on the atoms. Give necessary and sufficient conditions on $\mathcal{I}$ for Replacement, for Collection on a specified class of domains, and for a common-reservoir argument.

The cardinal-capacity calculation suggests that the relevant invariants combine union closure, available support sizes, and the action of permutations fixing parameter kernels. An arbitrary ideal need not have the fresh-atom property used in [lemma 32.1](#).

Question 38.5 (Mixed surreal-atom witnesses). Design a natural two-sorted extension of the sign-sequence structure whose witnesses encode both a birthday and an atom base. Can a local interpolation or extension principle in this richer structure characterize full Collection without merely restating the two-resource criterion by definition?

A successful construction should remain mathematically recognizable and should explain how automorphisms act on the added data.

Question 38.6 (Admissible language expansions). Classify predicates P for which $(M_\kappa^{\text{ker}}, P)$ retains Replacement or full Collection. Can the first failure be characterized in terms of the support growth of uniquely P -definable outputs? How much of the permutation group must remain available to force kernel confinement?

The enumeration predicate supplies a sharp negative case. Predicates invariant under every atom permutation supply an immediate positive mechanism, but intermediate stabilizer groups are more interesting.

[Added 3 October 2026, batch 89: for one further family of predicates, the graphs of canonical lifts of finitely many atom injections, the report [SetTheory/Cardinals/docs/reports/ordinal-s-and-order-types/naming-elementary-embeddings](#) ([remark 24.6](#)) decides exactly when the expansion keeps Replacement and full Collection, by a criterion on the components of the named injections (Theorem 5.1, `nee:thm:profile`; Theorem 7.5, `nee:thm:collection`). This answers the question for that family only; the general classification stays open.]

Question 38.7 (Reflection spectra rather than Collection spectra). In controlled kernel models, determine which additional features separate partial reflection from full reflection while the pure surreal structure remains fixed. Seek a precise comparison with the more elaborate models of Glazer–Yao, rather than assuming the cardinal-capacity invariant alone controls reflection.

The present models are intentionally homogeneous and do not reproduce those finer separations. [Added 3 October 2026, batch 89: the same report shows, for the kernel models expanded by named lifts, that full transitive reflection is equivalent to Collection there (Theorem 7.7, `nee:thm:reflection`), and that partial transitive reflection of assertions $\forall x \in w \exists y \delta$ with Δ_0 matrix is Δ_0 Collection, which fails at limit cutoffs, although each single true $\exists y \delta$ reflects (Proposition 8.1, `nee:prop:partial`). It bears on this question but does not reproduce Glazer–Yao’s separations; the question stays open.]

Question 38.8 (Weaker power-set assumptions). The good-path proof uses the set $\mathcal{P}(\lambda \times \{-, +\})$. Determine whether a restricted binary-sequence existence axiom suffices in place of full Power Set, and whether interpolation over such a base becomes equivalent to additional Collection or recursion principles.

This asks for a genuinely different lower-strength construction, not a proof that secretly retains the same function-space container.

Question 38.9 (Constructive variants). Analyze which parts of the six-scheme equivalence survive intuitionistic logic. The current comparison rule uses a first disagreement and decidable alternatives between obstructions, while least-witness birthday selection also has classical content. A constructive version should specify which order and cut notions replace these choices.

Question 38.10 (A proof-assistant benchmark with semantic safeguards). Implement the weak-base and expanded-language results in ProveIt, using the same internal formula language for the positive and negative directions. Include regression tests that intentionally try to confuse a definable family with a set graph, or to use a reduct’s Replacement instance after adjoining E .

Such a benchmark would test more than symbolic algebra: it would test whether an automated proof pipeline preserves the intended theory and language across an argument.

39 Conclusion of source 10, ledger and provenance

39.1 Conclusion

The phrase “a set-indexed family of surreal numbers” conceals two different assertions. A family already represented by a set graph supports the ordinary simplicity construction in Zermelo set theory without Replacement. A merely definable functional family can fail to have that graph. Uniformly repairing this failure is equivalent to ordinal bounding, and on an explicit pure cumulative-hierarchy base it is equivalent to full Replacement and Collection. The obstruction is visible inside the single interval $(-1, 0)$.

Urelements expose a second direction of failure. Pure sign sequences retain ordinal height but forget atom bases. Cardinal-support models keep the entire pure universe unchanged while varying the exact domain threshold for Collection. An external atom enumeration can additionally destroy Replacement, with an exact cofinality threshold, even though every pure-valued witness relation still satisfies Collection.

These results provide a concrete intersection between weak-set-theoretic research and surreal formalization. Their most useful lesson is operational: record the language, the index domain, the birthday bound, and the atom support separately. Each is mathematically consequential, and none should be supplied silently by the notation or by the proof assistant’s host theory.

39.2 A compact dependency and limitation ledger

Result	Assumptions	Important exclusion
Set-cut simplicity	Z_f , actual sets of endpoints	No arbitrary definable images assumed.
Six-scheme equivalence	Z_f , all instances in one fixed language	No assertion of full Replacement without a hierarchy hypothesis.
Full Replacement characterization	$Z_f + \text{Hier}$	Pure, atom-free cumulative hierarchy.
$V_{\omega+\omega}$ example	External ZFC	The missing family is not an internal sequence.
M_κ^{ker} construction	Full ambient urelement universe with Choice; $ A \geq \kappa$; infinite κ	Internal proper-class atoms are an ambient set.
Collection spectrum	Same ambient assumptions	Restricted by domain size, not witness rank.
Enumeration expansion	Ambient set predicate E , not an internal graph	Reduct Replacement cannot be reused in the expansion.
Pure-valued Collection	Ambient-definable finite predicate expansion	Does not collect arbitrary impure witnesses.
Finite checks	Ordinary Python execution	No transfinite or kernel-level verification.

39.3 Bibliographic and repository audit notes

The research inspection used Glazer’s 2023/2024 global-choice paper, the February 2026 Glazer–Yao preprint, the Hamkins–Yao treatment of classical kernel models, Yao’s axiomatization paper, and the listed ProveIt pages. The live arXiv version of the Glazer–Yao paper inspected was version 1. No private professional information about Glazer was needed; the match of interests is established by his mathematical publications rather than inferred from a job title.

The limited search did not establish a published occurrence of the exact six-scheme formulation or of the combined spectrum-and-expansion package. That absence is not evidence sufficient to establish priority. Classical results and close precedents are credited in [section 24.3](#).

39.4 Non-claims of source 10

Every limitation stated by source 10, in its manuscript or in its proof-status note, is kept; most appear at their point of use. Collected: the manuscript is unrefereed; classical simplicity (Conway, Gonshor, Bertram) and the small-support arguments (Hamkins–Yao §1.5 with Lévy; Glazer–Yao, Proposition 1; Yao) are credited, and neither the six-scheme package nor the spectra is claimed to have priority; no Lean or Rocq development is included or claimed checked; the finite tests do not test infinite cardinals, class semantics or full axiom schemes; the full-Replacement conclusion in pure weak set theory assumes Hier; the first half uses only the sign-sequence order, not a field construction in the weak base; internal proper-class atoms can form a set in the ambient universe; cardinal computations in the support models use ambient Choice; Replacement for a reduct is not Replacement for an expanded language; pure-valued Collection is not Collection for impure witnesses; the models are relative interpretations, not consistency proofs; the expansion by E is not a reproof of Glazer’s global-choice theorem; Glazer–Yao’s reflection separations are background, not results here; the repository inspection was targeted at READMEs and directory organization, not a source audit or a build. No named conjecture is resolved.

39.5 Provenance of this part

The manuscript. Source 10 is batch-89 manuscript 03, delivered as the archive `Surreal_Cuts_Replacement_Atom_Support_Spectra.zip` in commit `7c0f2d9f9` and placed in commit `23adb85f9`. It names the repository commit `2b7b388ba` (“Catalogue batches 81 to 86”) as its inspected snapshot, and all the paths it names exist, unchanged, at the placement commit. Its manuscript (`article.tex`, 926 lines, 42 labels; a 23-page PDF), its README, its PDF and its checksum file `SHA256SUMS` (8 of 8 entries verified at placement) are not shipped; they survive in the arrival commit. Shipped, byte-identical: `10-cuts-replacement-PROOF_STATUS.md`, `code/10-cuts-replacement-verify_finite.py`, `code/10-cuts-replacement-build.sh`, `data/10-cuts-replacement-finite_checks.json` and `data/10-cuts-replacement-build_validation.json` (source 10’s own record of its 23-page build, not of this report). The Hilbert’s-tenth programme routed the batch-89 archives in commit `6aadaa4f2` (`Computability/HilbertTenthProblem/Papers/research-wip/native-stream-queue/review_topology_surreal_arrivals_20261003.md`): it read bounded ranges, ran nothing, made no correctness finding, and concluded that none of them gives a Diophantine compiler or an arithmetic saving.

How it is printed. The text from source 10’s title-page status box and abstract to its appendices is printed in full, in its order, as [section 24](#)–[section 39](#): the status box and the abstract open [section 24](#),

its Section 1 became the next three subsections, and its two appendices became subsections of this section. Every delivered label carries the prefix `hset:zr::`; its ten questions gained labels `hset:zr:q:...`; its Theorems A–C are numbered (theorems 24.1 to 24.3). The symbols renamed are listed in convention 24.4; the countermodel of section 30 is written V_θ instead of M ; delivered file names are replaced by the shipped ones; its references to Conway, Gonshor and Bertram use this report’s bibliography entries. Added for this report: sections 24.4 and 24.5, the bracketed dated notes, this subsection and the preceding one. No mathematical statement or proof of source 10 was changed.

Where the merge had to choose. (i) Placement: source 10 cites no report of the collection; it is placed here because its spine is the weak-base form of theorem 10.4 (remark 24.5). (ii) Its Collection spectrum for the kernel models is printed in full here, with its own proofs, although the research-report collection prints the same theorem from an independent manuscript (remark 24.6). (iii) Its hierarchy axiom and set-graph scheme were renamed rather than left to collide with H_κ and `SignGraph`.

Part VII

Surreal universality without choice

40 Source 11: question, results, notation and relation to the report

This part was also added in batch 89. It prints the manuscript *Surreal Universality Without Choice: Exact coding strength, set-sized universal fields, and finite-transcendence collapse*, “A research manuscript prepared for Vladimir Reshetnikov”, October 3, 2026. It is batch-89 manuscript 04 and *source 11* in this report’s local numbering; its shipped files carry the prefix `11-choicelless-universality-`. It names the repository commit `2b7b388ba` (“Catalogue batches 81 to 86”; source 11 calls it a “tree identifier”, but it is a commit), arrived in commit `7c0f2d9f9` and was placed in commit `23adb85f9`. Its provenance and the mechanical changes made in printing it are in section 53. Inside this part, “this manuscript”, “here”, “our” and “we” refer to source 11.

Research intersection (source 11). Choiceless set theory and permutation models, as in the work of Elliot Glazer and Bokai Yao, meet the sign-sequence, Hahn-series, and omnific-integer foundations developed in the ProveIt repository.

Status. The manuscript supplies complete mathematical arguments for the stated deductions from explicitly identified classical background. It is an AI-generated, unrefereed research draft, not a Lean- or Rocq-checked development. Historical novelty is not certified. It does not claim to solve the reflection problems studied by Glazer and Yao, nor to transfer an atom-model counterexample automatically to pure set theory.

Main message. Freely naming independent surreal coordinates costs exactly one powerset-of-an-ordinal code. Preserving a given order is a separate requirement. In symmetric atom models, finite support places a sharp finite bound on how many independent coordinates any one representation can retain.

Source 11’s summary. We calibrate several universality assertions for the surreal numbers over set theory without the axiom of choice. For every set X , an injection $X \rightarrow \mathbf{No}$ exists if and only if X injects into the powerset of an ordinal. An explicit double-monomial construction upgrades any such injection to algebraically independent positive infinite omnific integers. Consequently, the existence of embeddings of all free integer polynomial rings into $\mathbf{Oz}$, or all free real rational-function fields into $\mathbf{No}$, is exactly the first Kinna–Wagner principle.

For every infinite initial ordinal κ , we construct a set-sized real-closed subfield $U_\kappa \subseteq \mathbf{No}$, equinumerous with $\mathcal{P}(\kappa)$ without choice, which realizes precisely the free real rational-function fields whose variable sets inject into $\mathcal{P}(\kappa)$. The cardinal upper bound uses canonical fraction and root codes rather than an unjustified choice of quotient representatives.

An orbit-factorization theorem then yields sharp bounds in permutation models. A polynomial representation into the pure surreals supported by s atoms has image-field transcendence degree at most $s + 1$ in the basic Fraenkel model, and at most $2s + 1$ in the ordered Mostowski model. Both bounds are attained. The latter model contains an ordered rational-function field with no embedding into its surreals, although every finitely generated-over- $\mathbb{R}$ coordinate subfield embeds in order. Compatible local embeddings would produce a global one, so the missing ingredient is coherence, not a defect of the finite algebra.

We separate coding universality from prescribed-order universality, give an ordinal convex-resolution criterion for order embeddings, identify the precise foundations required by the proofs, and propose a formalization and research program.

40.1 The research question

The slogan that the surreal numbers are a universal ordered field suppresses several different quantifiers. Does one wish to assign distinct surreal coordinates to the elements of a set? To assign algebraically independent coordinates? To embed a field as a field, or as a field with its already specified order? Must the embedding belong to the universe in which the field was constructed? These questions coincide much less often when choice is unavailable.

This distinction provides a concrete meeting point between two research programs. Glazer and Yao’s *Reflection Principles in ZFU* studies how assertions that agree under choice can separate when urelements and permutation symmetries are present [21]. ProveIt’s surreal-number project develops sign-sequence foundations, normal forms, the real-closed surreal field, and the omnific-integer subring; it also explicitly distinguishes research claims from checked declarations [26]. Our subject is the choice cost of *representing labelled algebra inside those fields*.

The relevant weak-choice principle is classical. With the indexing convention of Karagila and Schilhan [24],

$$\text{KWP}_1 : \quad \forall X \exists \alpha \in \text{Ord} \quad X \hookrightarrow \mathcal{P}(\alpha). \quad (40.1)$$

Its coding formulation is not a new principle introduced here. Our contribution is an explicit algebraic calibration, a set-sized optimal target at each code level, and a quantitative analysis of how symmetry defeats the same constructions internally.

40.2 What is proved, and what is not claimed

The principal questions answered in this manuscript are as follows.

How much choice is required to realize arbitrary independent variables in No or Oz?
Exactly KWP_1 , for the free-algebra assertions specified in [theorem 43.5](#). The result is an equivalence over ZF, not merely an implication from choice.

Can one use an optimal set-sized real-closed target at a fixed ordinal code level?

Yes. The field U_κ of [theorem 44.3](#) has the same cardinality, in the sense of an actual bijection, as $\mathcal{P}(\kappa)$.

What does finite support force a polynomial representation to forget?

It identifies variables in the same stabilizer orbit. In two standard atom models this gives the sharp bounds $s + 1$ and $2s + 1$ of [theorems 48.1](#) and [48.3](#).

Does finite-stage embeddability ensure a global surreal embedding?

No, in the ordered permutation model. It does so precisely when the finite-stage maps can be assembled coherently, as formalized in [theorem 49.3](#).

These are natural questions formulated and answered here; they are not presented as named, previously published open problems. The elementary coding observation, finite normal-form identities, polynomial universal property, and standard permutation-model construction are established machinery. The possible novelty lies in the exact combined statements and the quantitative refinements. A targeted literature review did not establish historical priority for those formulations.

40.3 Reading paths

For the main positive results, read [sections 41](#) to [44](#). For the negative results, read [sections 47](#) to [49](#). The prescribed-order issue is isolated in [sections 45](#) and [46](#). The final sections discuss proof boundaries, formalization, and research questions. All displayed embeddings into a proper class have set-sized domains and set graphs.

40.4 Notation of this part

Convention 40.1 (Symbols of source 11). In this part:

Notion in source 11	Here	Do not confuse with
Birthday of a sign sequence	$b(s)$, an ordinal (source: b)	the same function; Parts I–V regard its value as a surreal ordinal
Sign sequences of length at most α	$\mathbf{No}_{\leq \alpha}$ ($= \mathbf{No}_{< \alpha+1}$)	the cutoff carrier $\mathbf{No}_{< \lambda}$ of Parts I–V: at an infinite initial κ , $\mathbf{No}_{\leq \kappa}$ also contains the sequences of length exactly κ
Sign sequences of length exactly κ	$\mathbb{S}_\kappa = \{-, +\}^\kappa$ (source: S_κ)	S_κ , sources 05 and 06's name for the carrier $\mathbf{No}_{< \kappa}$ (convention 1.5)
Free real field on $\mathbb{S}_\kappa$	$\mathcal{R}_\kappa$ (source: K_κ)	the collection's K , a complex field, and source 09's K_λ (renamed $\mathbf{No}_{< \lambda}$)
Sign-to-subset code	$\text{enc}_\alpha(s)$ (source: C_α)	the relation codes $C = (\delta, r, t)$ of section 5
Least code level, least birthday level	$\text{cl}(X)$, $\text{bl}(X)$ (source: $\rho(X)$, $\beta(X)$)	β , the first new birthday, and ϱ , the eternity bound
A group acting on variables	G_0 (source: H)	H_κ , the hereditary-size universe
Generic ordinal variable	ζ (source: β)	β (reserved)
An injection $X \rightarrow \mathcal{P}(\alpha)$ in theorem 42.1	c	the conjugation c of section 19

Notion in source 11	Here	Do not confuse with
Omnific integers; first Kinna–Wagner principle	Oz ; KWP_1	—
Length of a convex resolution; limit stages	θ ; λ	the pairing width θ of section 4 ; the epsilon-number cutoffs λ
Permutation model; atom support size	W ; s	s also denotes a sign sequence, as in the source

Source 11’s exponent sums E_m and the equivalence relations E_ξ of a convex resolution are its own two uses of E ; neither is the enumeration predicate E of [Part VI](#). Typography only: theory names (ZF, ZFA, AC, KWP) are set upright, as in the rest of this report.

40.5 Relation to the rest of the report and to the collection

Remark 40.2 (The open item of [Question 21.7](#), re-scoped). [Question 21.7](#) lists, as source 06’s open item, “a universal embedding theorem without Global Choice”. Source 11 answers it in part, over ZF: coordinate universality (every set of variables realized by algebraically independent surreals, or every free real rational-function field embedded in **No**) is exactly KWP_1 ([theorem 43.5](#)); at each code level the set-sized real-closed field U_κ is an optimal target ([theorem 44.3](#)); every linear order whose underlying set is well-orderable order-embeds in **No** ([proposition 45.3](#)), and order embeddability is characterized by convex resolutions ([theorem 46.2](#)); and in a ZFA permutation model an ordered field all of whose finitely generated coordinate subfields embed has no embedding ([theorem 49.1](#)). It leaves open universality for prescribed orders and for general ordered fields without choice (its [questions 51.1](#), [51.2](#) and [51.5](#)), and any ZF transfer of the ZFA counterexample. Source 11 did not know this report.

Remark 40.3 (Comparison with the choice-free [Section 13](#)). [Theorem 13.2](#) (source 06) identifies, over ZF, the range of the pointed ordinal-graph *interpretation* of sets: exactly HWO, the sets whose transitive closure is well-orderable. [Theorem 42.1](#) concerns *injections* of a set into **No**: $X \hookrightarrow \mathbf{No}$ iff $X \hookrightarrow \mathcal{P}(\alpha)$ for some α . The second class is the wider one. If $x \in \text{HWO}$ then $x \subseteq \text{TC}(\{x\})$ is well-orderable and so injects into an ordinal; but $\mathcal{P}(\omega)$ always injects into **No** (take $\alpha = \omega$), while in a model of ZF where $\mathcal{P}(\omega)$ is not well-orderable it is not in HWO ([proposition 13.3](#) uses the same hypothesis). Under Choice both classes are everything. A set coded by a surreal in the sense of [theorem 13.2](#) is recovered as a set; an injection into **No** only names its elements. This comparison is an observation of this merge.

Remark 40.4 (Other reports). (i) [Theorem 43.1](#) is the case $K = \mathbb{R}$, $G = \{0\}$ of [Lemma 8.1](#) (`dsn:lem:cosets`, “Disjoint support cosets”, that report’s source 04) of `foundations-and-computation/definable-surreals-and-omnific-integers`: a nonzero real has normal-form support $\{0\}$, and distinct Conway monomials e_x have no nonzero integer relation, so the family ω^{e_x} is algebraically independent over $\mathbb{R}$. Source 11’s proof is printed as its own; new beside that lemma are the omnific-integer conclusion, the free-algebra calibration and the ZF framing. (ii) The report `foundations-and-computation/surreal-well-orders` proves over GB without set choice that Global Choice holds iff **No** has a class well-order (its [Theorem 13.1](#), `swo:cf:choice`; see the dated note after [remark 13.4](#)); source 11 instead works with sets and no global principle, and calibrates coordinate universality by KWP_1 . That report also records the ZFC count $|\mathbf{No}_{<\kappa}| = 2^{<\kappa}$; source 11’s [proposition 42.3](#) is a ZF bijection at the closed bound $\mathbf{No}_{\leq\kappa}$. Its [Part XV](#) (batch-89 manuscript

01) studies countable global choice over Zermelo set theory: a related theme, different theorems. (iii) No other report of the collection treats Kinna–Wagner principles, permutation models or choiceless universality.

41 Foundations, notation, and the surreal interface

41.1 Set theory and classes

Unless otherwise specified, the positive results are theorems over ZF. We use classes only as definable predicates and definable operations on sets. Equivalently, one may work in a conservative class-theoretic presentation without adding set choice or global choice. The permutation-model sections work in ZFA: set theory with a *set* of atoms. This is a special setting within urelement theories, not the unrestricted proper-class-of-atoms setting of all results in [21].

A surreal number is represented by a function

$$s : \zeta \longrightarrow \{-, +\}, \quad \zeta \in \text{Ord}.$$

Its birthday is $b(s) = \zeta$. The totality **No** of such functions is a proper class. We write

$$\mathbf{No}_{\leq \alpha} = \bigcup_{\zeta \leq \alpha} \{-, +\}^\zeta.$$

This is a set. An expression $f : X \rightarrow \mathbf{No}$ means that f is a set of ordered pairs, its domain is the set X , and every value satisfies the defining predicate for **No**. Replacement therefore applies to the values and to their birthdays. We never take a powerset of **No**.

In ZFA, the signs, ordinals, and sign sequences are pure sets: their transitive closures contain no atoms. Atom permutations fix each surreal number pointwise. A function whose arguments involve atoms need not be pure, even if all its values are surreals. This difference drives the counterexamples.

41.2 The standard surreal facts used

We use the following classical interface, with no choice principle added to the ambient set theory. Conway’s foundational construction is the underlying reference [1]; Costin and Ehrlich explicitly develop their positive surreal theory in a class framework without set or class choice [25].

- (S1) **No** is an ordered real-closed field containing the canonically embedded ordered field $\mathbb{R}$.
- (S2) The Conway monomial map $x \mapsto \omega^x$ is strictly increasing, has positive values, and satisfies

$$\omega^{x+y} = \omega^x \omega^y, \quad \omega^0 = 1.$$

Here ω^x denotes Conway’s monomial, not the separate Gonshor exponential function.

- (S3) If $x_1 > \cdots > x_n$ and $r_i \in \mathbb{R} \setminus \{0\}$, the finite normal form

$$\sum_{i=1}^n r_i \omega^{x_i}$$

is nonzero and has the sign of r_1 . In particular, distinct monomials are linearly independent over $\mathbb{R}$.

- (S4) A surreal is omnific when its normal-form exponents are nonnegative and its constant coefficient is an ordinary integer. These numbers form the subring **Oz**. In particular, a monomial ω^e with $e > 0$ is a positive infinite omnific integer.

(S5) For set-sized $A, B \subseteq \mathbf{No}$ with $A < B$, there is a unique simplest surreal strictly between them. This is the usual set-cut construction.

Only finite normal forms are needed for the independence and symmetry theorems. The set-sized real-closed-target theorem additionally uses real closedness. No theorem about arbitrary rearrangements of infinite Hahn sums, surreal integration, or sharp product birthdays is used.

Remark 41.1 (Three meanings of support). A polynomial has a finite *variable support*; a Hahn or Conway series has a *normal-form support*; an object in a permutation model has an *atom support*. They are different notions. In particular, a polynomial homomorphism with finite atom support can have a domain with infinitely many variables.

41.3 Algebra on arbitrary sets of variables

For any set X , let $\mathbb{N}^{(X)}$ be the set of finitely supported functions $X \rightarrow \mathbb{N}$. The ring $k[T_X]$ consists of finitely supported coefficient functions on $\mathbb{N}^{(X)}$. Addition is pointwise and multiplication is finite convolution. This definition requires no ordering or enumeration of X . If k is a field, the resulting ring is a domain, and its fraction field is denoted $k(T_X)$.

An embedding of $\mathbb{R}$ -algebras fixes the specified copy of $\mathbb{R}$. A unital field homomorphism is automatically injective. A field embedding, unless explicitly called ordered, need not preserve a prescribed order on the source. Finite transcendence degrees below are ordinary nonnegative integers; no existence of an arbitrary infinite transcendence basis is assumed.

42 The exact coding strength of a surreal coordinate set

Theorem 42.1 (Ordinal-code equivalence). *For every set X , the following are equivalent over ZF:*

$$X \hookrightarrow \mathbf{No} \quad \Longleftrightarrow \quad \exists \alpha \in \text{Ord} \quad X \hookrightarrow \mathcal{P}(\alpha).$$

The same equivalence holds in ZFA with pure sign-sequence surreals.

Proof. Suppose first that $j : X \rightarrow \mathbf{No}$ is injective. By Replacement, the set of birthdays of its values exists. Let

$$\alpha = \sup\{b(j(x)) : x \in X\},$$

with $\alpha = 0$ if $X = \emptyset$. Every value belongs to $\mathbf{No}_{\leq \alpha}$.

For a sign sequence s of length $\zeta \leq \alpha$, put

$$\text{enc}_\alpha(s) = \{2 \cdot \xi + 1 : \xi < \zeta\} \cup \{2 \cdot \xi : \xi < \zeta \text{ and } s(\xi) = +\}. \quad (42.1)$$

This is a subset of the ordinal $2 \cdot \alpha$. Multiplication here is ordinal multiplication on the indicated side: $2 \cdot \alpha$ is the sum of α consecutive two-point blocks. The odd position of each block records whether the sign is defined; the even position records whether it is positive. Thus the code uniquely determines both the domain and every sign, so enc_α is injective. Composing with j proves the forward implication.

Conversely, given an injection $c : X \rightarrow \mathcal{P}(\alpha)$, let s_x be the sign sequence of exact length α defined by

$$s_x(\xi) = + \Longleftrightarrow \xi \in c(x).$$

Different codes give different surreals. This definition uses only Separation and Replacement and works unchanged when the members of X are atoms or sets involving atoms. $\square$

Remark 42.2 (Why definedness must be encoded). Coding a sign sequence solely by its positive positions loses information: the empty sequence and the one-symbol sequence – would both have empty code. The first half of (42.1) prevents that error. Padding by a third symbol, as in section 46, is an alternative.

Proposition 42.3 (Exact birthday capacity). *For every infinite initial ordinal κ , there is a bijection*

$$\mathbf{No}_{\leq \kappa} \simeq \mathcal{P}(\kappa)$$

in ZF. For a finite ordinal n ,

$$|\mathbf{No}_{\leq n}| = 2^{n+1} - 1.$$

Proof. Subsets of κ inject into the sign sequences of exact length κ . In the other direction, (42.1) maps $\mathbf{No}_{\leq \kappa}$ into $\mathcal{P}(2 \cdot \kappa)$. The ordinal $2 \cdot \kappa$ is equipotent with κ : finite products of an infinite well-orderable cardinal do not require choice. Transporting subsets along such a bijection and applying Cantor–Bernstein gives the result. At finite n , sum the 2^i sign sequences of each length $0 \leq i \leq n$. $\square$

Corollary 42.4 (Coincidence of least code and birthday budgets). *Let X be an infinite set admitting the embeddings in theorem 42.1. Define*

$$\text{cl}(X) = \min\{\alpha : X \hookrightarrow \mathcal{P}(\alpha)\}, \quad \text{bl}(X) = \min\{\alpha : X \hookrightarrow \mathbf{No}_{\leq \alpha}\}.$$

Then $\text{cl}(X) = \text{bl}(X)$, and their common value is an infinite initial ordinal.

Proof. Each minimum exists by taking the least ordinal below an existing witness. Neither is finite, since a set injecting into a finite set is finite. If an infinite witness ordinal α is not initial, replace it by its initial ordinal $\kappa < \alpha$. Subsets of α and of κ are equipotent. Moreover, the proof of proposition 42.3 shows $\mathbf{No}_{\leq \alpha} \hookrightarrow \mathcal{P}(\kappa) \hookrightarrow \mathbf{No}_{\leq \kappa}$. Thus a least witness is initial. At infinite initial ordinals the two target sets have equal capacity by proposition 42.3. $\square$

The finite restriction matters. A three-element set fits into $\mathbf{No}_{\leq 1}$, whereas it does not fit into $\mathcal{P}(1)$.

43 Double monomials and free-algebra universality

Theorem 43.1 (Uniform algebraic independence). *Let $j : X \rightarrow \mathbf{No}$ be an injection. Define*

$$e_x = \omega^{j(x)}, \quad t_x = \omega^{e_x}. \tag{43.1}$$

Then $(t_x)_{x \in X}$ is algebraically independent over $\mathbb{R}$. Every t_x is a positive infinite omnific integer. Evaluation gives embeddings

$$\mathbb{Z}[T_X] \hookrightarrow \mathbf{Oz}, \quad \mathbb{R}[T_X] \hookrightarrow \mathbf{No}, \quad \mathbb{R}(T_X) \hookrightarrow \mathbf{No}.$$

All these maps are definable from j and have set graphs.

Proof. The numbers e_x are distinct Conway monomials. By finite normal-form uniqueness they are linearly independent over $\mathbb{R}$, and in particular over $\mathbb{Z}$.

For $m \in \mathbb{N}^{(X)}$, define the finite sum

$$E_m = \sum_{x \in X} m(x)e_x.$$

If $m \neq n$, the finite linear combination $\sum_x (m(x) - n(x))e_x$ is nonzero. Hence $E_m \neq E_n$. The multiplication law for Conway monomials gives

$$\prod_x t_x^{m(x)} = \omega^{E_m}.$$

Now take a nonzero polynomial, written after collecting like monomials as

$$p = \sum_{m \in F} a_m T^m,$$

where F is finite and every $a_m \neq 0$. Evaluation produces

$$p(t) = \sum_{m \in F} a_m \omega^{E_m}. \quad (43.2)$$

The exponents are distinct. They can be sorted because they form a finite subset of a linear order. The resulting finite normal form is nonzero, proving algebraic independence.

Each $e_x > 0$, regardless of the sign or magnitude of $j(x)$. Thus $t_x = \omega^{e_x}$ has a single positive exponent and is a positive infinite element of $\mathbf{Oz}$. In (43.2), every nonconstant monomial has exponent $E_m > 0$. If the coefficients are integers, the only possible constant coefficient is an integer, so the value is omnific. Evaluation is therefore injective on both polynomial rings in the stated targets. Since nonzero denominators do not vanish, the real polynomial embedding extends uniquely to its fraction field. Replacement on the set of polynomials and fractions gives the graphs of the maps. $\square$

Example 43.2 (Why the second monomial layer is needed). An arbitrary injective assignment j does not make $\omega^{j(x)}$ algebraically independent. For labels $j(x_0) = 0$, $j(x_1) = 1$, and $j(x_2) = 2$, the assigned values are $1, \omega, \omega^2$, and two immediate polynomial relations hold. The second layer in (43.1) makes the *exponents* linearly independent before using them to build algebraically independent monomials.

Example 43.3 (A concrete leading-term calculation). With two labels $j(a) = 0$ and $j(b) = 1$, the double-monomial values are $t_a = \omega$ and $t_b = \omega^\omega$. For instance,

$$T_b^2 - 3T_a T_b + T_a^5 \quad \mapsto \quad \omega^{2\omega} - 3\omega^{\omega+1} + \omega^5 > 0.$$

The three displayed exponents are distinct and decreasing. This illustrates the proof mechanism: a formal monomial becomes a Conway monomial, and the greatest surviving exponent certifies nonvanishing. The products and sums in these exponents are surreal field operations, not the ordinal multiplication used in the coding formula.

Corollary 43.4 (Independent infinitesimal coordinates). *Under the hypotheses of theorem 43.1, the numbers*

$$u_x = t_x^{-1} = \omega^{-e_x}$$

are algebraically independent positive infinitesimals. They are not omnific integers.

Proof. They are infinitesimal because $-e_x < 0$. A polynomial relation among finitely many inverses, multiplied by a sufficiently large product of the corresponding t_x , would give a nonzero polynomial relation among the t_x . Their negative normal-form exponents exclude them from **Oz**. $\square$

Theorem 43.5 (Equivalent forms of coordinate universality). *For a set X , the following conditions are equivalent over ZF:*

- (i) $X \hookrightarrow \mathcal{P}(\alpha)$ for some ordinal α ;
- (ii) $X \hookrightarrow \mathbf{No}$;
- (iii) $X \hookrightarrow \mathbf{Oz}$;
- (iv) there is a unital ring embedding $\mathbb{Z}[T_X] \hookrightarrow \mathbf{Oz}$;
- (v) there is an $\mathbb{R}$ -algebra embedding $\mathbb{R}[T_X] \hookrightarrow \mathbf{No}$;
- (vi) there is an $\mathbb{R}$ -field embedding $\mathbb{R}(T_X) \hookrightarrow \mathbf{No}$;
- (vii) there is an X -indexed algebraically independent family over $\mathbb{R}$ consisting of positive infinite omnific integers.

Consequently, the universal closure over all sets X of any of these assertions is exactly KWP_1 .

Proof. The equivalence of (i) and (ii) is [theorem 42.1](#). From (ii), [theorem 43.1](#) gives (iv)–(vii) and also (iii). Conversely, (iii) implies (ii) by inclusion. An embedding in (iv), (v), or (vi) assigns distinct values to the distinct indeterminates: otherwise it kills the nonzero polynomial $T_x - T_y$. Thus each yields (ii). Algebraic independence likewise gives distinct values, so (vii) yields (ii). Universal closure and [\(40.1\)](#) prove the last statement. $\square$

Remark 43.6 (Not an equivalence with full choice). In pure ZF, KWP_1 is weaker than AC; Cohen's first model is a standard example satisfying the former and not the latter, as reviewed in [\[24\]](#). Thus these *free-algebra* universality assertions do not recover the well-ordering theorem. They also do not yet assert anything about preserving an independently prescribed order on X or on a field.

44 An optimal set-sized real-closed target

The preceding theorem places all realizations in the proper class **No**. It is useful, both mathematically and for formalization, to obtain a target that is an ordinary set and has exactly the required coding capacity.

Fix an infinite initial ordinal κ . Let

$$\mathbb{S}_\kappa = \{-, +\}^\kappa, \quad t_s = \omega^{\omega^s} \quad (s \in \mathbb{S}_\kappa),$$

where each s is identified with its surreal value. Set

$$\mathcal{R}_\kappa = \mathbb{R}(t_s : s \in \mathbb{S}_\kappa), \quad U_\kappa = \{u \in \mathbf{No} : u \text{ is algebraic over } \mathcal{R}_\kappa\}. \quad (44.1)$$

The notation on the right initially defines a class; the next proof shows that it is a set.

44.1 A choice-free coding lemma

Lemma 44.1 (Finite syntax at a powerset level). *Let κ be an infinite initial ordinal. Suppose that D and B inject into $\mathcal{P}(\kappa)$, that D is linearly ordered, and that B is a field with a fixed linear order on its underlying set. Then each of the following admits an injection into $\mathcal{P}(\kappa)$:*

$$B[T_D], \quad B(T_D), \quad \text{any ordered algebraic extension of } B(T_D).$$

For the last assertion the extension is a set, or is a definable class in which each polynomial has only finitely many roots and ordered root indexing applies.

Proof. Finite lists of subsets of κ can be encoded by a subset of κ : first tag the list length and each entry in $\omega \times \kappa$, then use a fixed bijection $\omega \times \kappa \simeq \kappa$. These are well-orderable sets, so this bijection needs no choice. Naturals and finite auxiliary flags can be accommodated by additional tags.

Every finite subset of D can be uniquely listed in increasing order. A monomial is therefore coded by an increasing finite variable list together with its positive integer exponents. Monomials are linearly ordered by comparing exponent vectors at the greatest variable where they differ. Every finite polynomial has a unique list of its nonzero terms in this order. Encoding these lists and their coefficients proves the assertion for polynomials.

A fraction requires more care. It would be invalid, without further argument, to infer that an arbitrary quotient of a $\mathcal{P}(\kappa)$ -codable set is again $\mathcal{P}(\kappa)$ -codable. Instead use the unique reduced representative

$$\frac{p}{q}, \quad \gcd(p, q) = 1, \quad \text{lc}(q) = 1, \quad (44.2)$$

with 0 represented as 0/1.

Here leading coefficients are taken with respect to the preceding largest-variable monomial order. They exist because a polynomial has finitely many terms. The polynomial ring in an arbitrary set of variables over a field is a unique factorization domain in the finite sense needed here: put p and q in the ring generated by their finite union of variables and use finite-variable factorization. A factor cannot introduce a new variable, since degree in that variable is additive in a polynomial domain. Gcds computed in the finite subring therefore remain gcds in the full ring. Dividing out a gcd and scaling by the leading coefficient of the denominator produces (44.2). Uniqueness follows from coprimality and cancellation in a UFD. These are existence-and-uniqueness arguments, not selections from an unstructured family of equivalence classes. Thus a fraction is injected into a pair of polynomial codes.

Finally, an algebraic element a of an ordered extension has a unique monic minimal polynomial over $B(T_D)$. Attach to it the position of a among the finitely many roots of that polynomial *in the extension*, in increasing order. Polynomial plus finite index uniquely determines a . This is an injection into finite lists of field elements with a natural-number tag, and hence into $\mathcal{P}(\kappa)$. $\square$

Remark 44.2. The order on D is not required to be a well-order, and the monomial order is not claimed to well-order all monomials. Only the maximum of each finite support and compatibility with addition of exponent vectors are used. Finite-variable gcd theory can use a separate finite enumeration when proving existence; the normalization in (44.2) makes the answer independent of that temporary enumeration.

Theorem 44.3 (Optimal code-level universal field). *Over ZF, for every infinite initial ordinal κ , the class U_κ in (44.1) is a set-sized real-closed subfield of $\mathbf{No}$ and*

$$U_\kappa \simeq \mathcal{P}(\kappa). \quad (44.3)$$

For every set X ,

$$X \hookrightarrow \mathcal{P}(\kappa) \iff \mathbb{R}(T_X) \hookrightarrow_{\mathbb{R}} U_\kappa, \quad (44.4)$$

$$X \hookrightarrow \mathcal{P}(\kappa) \iff \mathbb{Z}[T_X] \hookrightarrow U_\kappa \cap \mathbf{Oz}. \quad (44.5)$$

The embeddings in (44.4) do not have to preserve a separately prescribed order on the source.

Proof. By [theorem 43.1](#), the t_s are algebraically independent over $\mathbb{R}$. The set of finite rational expressions in them is a set, and their evaluation is definable. Replacement gives the set $\mathcal{R}_\kappa$, together with an isomorphism from $\mathbb{R}(T_{\mathbb{S}_\kappa})$.

For each nonzero polynomial $p \in \mathcal{R}_\kappa[Y]$, there are at most $\deg p$ distinct roots in the field **No**. Its definable class of roots is therefore a finite set. More explicitly, the assertion that a definable class has at most n members allows its members to be collected using finite induction; this is not a choice principle. The roots form a linearly ordered finite set. The function that assigns to (p, i) its i -th root, when present, and 0 otherwise is a definable function on a set. Replacement shows that the union of all these root sets is a set. It is exactly U_κ .

Algebraic elements are closed under field operations, so U_κ is an ordered field algebraic over $\mathcal{R}_\kappa$. If $a > 0$ belongs to it, its positive square root in **No** is algebraic over $\mathcal{R}_\kappa$ and thus lies in U_κ . If an odd-degree polynomial has coefficients in U_κ , take one of its roots in the real-closed field **No**. Its finitely many coefficients are algebraic over $\mathcal{R}_\kappa$, and algebraicity is transitive, so the root belongs to U_κ . The usual square-root/odd-degree criterion proves that U_κ is real closed. This is a real closure of $\mathcal{R}_\kappa$ inside the chosen ordered field **No**.

For the cardinal upper bound, $\mathbb{S}_\kappa \simeq \mathcal{P}(\kappa)$ by its definition and is linearly ordered by surreal order. The reals inject into $\mathcal{P}(\omega)$ by their rational Dedekind cuts, and thus into $\mathcal{P}(\kappa)$. Apply [lemma 44.1](#), using canonical reduced fractions and minimal-polynomial/root-index codes. It gives an injection $U_\kappa \hookrightarrow \mathcal{P}(\kappa)$. Conversely, $s \mapsto t_s$ injects $\mathbb{S}_\kappa$ into U_κ . Cantor–Bernstein proves (44.3).

Suppose now $c : X \hookrightarrow \mathcal{P}(\kappa)$. Convert each code to its length- κ sign sequence s_x , and send T_x to t_{s_x} . Algebraic independence gives the embedding into $\mathcal{R}_\kappa \subseteq U_\kappa$, and integer polynomial values are omnific. This proves the forward implications in (44.4) and (44.5). Either embedding sends distinct variables to distinct elements of U_κ ; compose with its established code to obtain $X \hookrightarrow \mathcal{P}(\kappa)$. $\square$

44.2 What optimality means here

The theorem is sharp in a precise coding sense. Any set-sized target that contains an injective realization of $\mathbb{S}_\kappa$ many independent variables must admit an injection from $\mathcal{P}(\kappa)$. Our field attains that lower bound and is real closed. We are *not* claiming that every ordered field whose underlying set injects into $\mathcal{P}(\kappa)$ order-embeds in U_κ .

Nor do we claim a sharp explicit birthday bound for U_κ . Such a bound exists by Replacement because U_κ is a set. This qualitative fact must not be confused with an optimal formula for birthdays under multiplication, inversion, or taking algebraic roots.

45 Prescribed order is a different universality problem

Let L be a linearly ordered set. The free abelian group $\mathbb{Z}^{(L)}$ has a largest-variable order: for distinct m, n , let ℓ be the greatest member of the finite set on which they differ, and declare

$$m < n \iff m(\ell) < n(\ell). \quad (45.1)$$

This is a translation-invariant linear order. Define an order on $\mathbb{R}[T_L]$ by saying that a nonzero polynomial is positive when the coefficient of its greatest monomial is positive. Extend it to the fraction field, denoted $F(L)$.

Lemma 45.1 (Largest-term ordering). *The preceding construction makes $F(L) = \mathbb{R}(T_L)$ an ordered field over $\mathbb{R}$, without requiring a well-order of L . Its variables are positive and satisfy*

$$\ell < \ell' \implies T_\ell < T_{\ell'}.$$

Every order automorphism of L induces an order automorphism of $F(L)$.

Proof. Translation invariance in (45.1) is immediate because adding the same vector leaves the greatest differing coordinate and the sign of the difference unchanged. A polynomial has a finite nonempty set of monomials, hence a greatest one. If m_0, n_0 are the greatest monomials of nonzero p, q , then $m_0 + n_0$ is the unique greatest exponent vector arising in their product: any strict decrease in either summand strictly decreases the sum. Consequently,

$$\text{lm}(pq) = \text{lm}(p) + \text{lm}(q), \quad \text{lc}(pq) = \text{lc}(p) \text{lc}(q).$$

This proves closure of positive elements under multiplication. For addition, unequal leading monomials retain the greater positive term, while equal leading monomials have positive coefficient sum. Exactly one of $p, -p$ is positive for nonzero p . These observations give an ordered domain, whose order extends uniquely to its fraction field. The largest variable in $T_{\ell'} - T_\ell$ is ℓ' , with coefficient 1. All definitions are preserved by an order automorphism of L . $\square$

Theorem 45.2 (Exact reduction of order universality). *For every linearly ordered set L , over ZF,*

$$L \hookrightarrow_{\text{ord}} \mathbf{No} \iff F(L) \hookrightarrow_{\text{ord}} \mathbf{No}.$$

In the forward direction the field embedding fixes $\mathbb{R}$.

Proof. Suppose $j : L \hookrightarrow_{\text{ord}} \mathbf{No}$. Use $t_\ell = \omega^{\omega^{j(\ell)}}$ as in theorem 43.1. For distinct exponent vectors m, n , the sign of

$$E_m - E_n = \sum_{\ell} (m(\ell) - n(\ell)) \omega^{j(\ell)}$$

is determined by the greatest ℓ where the coefficient is nonzero, because j preserves order and finite normal-form signs are determined by the greatest exponent. Thus $m < n$ in (45.1) exactly when $E_m < E_n$ in $\mathbf{No}$. The sign of an evaluated polynomial is its largest-term coefficient, so the resulting polynomial and fraction-field embeddings preserve order.

Conversely, restricting any ordered embedding $F(L) \rightarrow \mathbf{No}$ to the strictly increasing family of variables gives an ordered injection of L . $\square$

Proposition 45.3 (Well-orderable domains). *Every linear order whose underlying set is well-orderable order-embeds into $\mathbf{No}$ in ZF.*

Proof. Fix one well-ordering and its enumeration $(x_\xi)_{\xi < \theta}$. Recursively assign to x_ξ the simplest surreal between the images of earlier elements below it and the images of earlier elements above it. Those two collections are sets, are separated by the induction hypothesis, and the simplest choice is unique. Transfinite recursion and Replacement construct a set graph of length θ . No new choice is made at any stage. The result preserves all comparisons. $\square$

Remark 45.4 (A gap that must not be silently bridged). An injection of the underlying set of L into $\mathcal{P}(\alpha)$ gives it *some* surreal-induced order. It does not say that an already specified order on L is respected. Thus theorem 43.5 is not, by itself, a proof that KWP_1 implies ordered universality for all linear orders or all ordered fields. We leave that implication as a separately stated research question rather than conflating it with the proved equivalence.

46 A convex-resolution criterion for ordered embeddings

There is a choice-free combinatorial certificate for the missing prescribed-order requirement.

Definition 46.1. A *ternary convex resolution of length θ* of a linear order L is a sequence $(E_\xi)_{\xi \leq \theta}$ of equivalence relations on L such that:

- (a) $E_0 = L \times L$ and E_θ is equality;
- (b) the relations refine with increasing ξ , and every equivalence class is convex in L ;
- (c) at a limit $\lambda \leq \theta$, $E_\lambda = \bigcap_{\xi < \lambda} E_\xi$;
- (d) every E_ξ -class is partitioned into at most three $E_{\xi+1}$ -classes.

A binary resolution is defined by replacing three with two.

Theorem 46.2 (Order-coding certificate). *For a set-sized linear order L , the following are equivalent over ZF:*

- (i) L order-embeds into **No**;
- (ii) L has a ternary convex resolution of some ordinal length;
- (iii) L has a binary convex resolution of some ordinal length.

A ternary resolution of length θ produces an order embedding into sign sequences of exact length $2 \cdot \theta$.

Proof. Suppose $j : L \hookrightarrow_{\text{ord}} \mathbf{No}$. Bound the birthdays of its range by an ordinal θ . Pad each sign sequence to length θ with the symbol 0 after it ends, using $- < 0 < +$. Surreal comparison is precisely lexicographic comparison of these padded strings. Let $x E_\xi y$ mean that their first ξ symbols agree. Prefix classes in a lexicographic order are convex; each splits into at most three next-symbol classes, limit relations are intersections, and equality of the whole padded strings gives $x = y$. Thus (i) implies (ii).

Now suppose a ternary resolution is given. Inside each E_ξ -class, its finitely many $E_{\xi+1}$ -classes inherit a linear order. Label them by 0, 1, 2 according to their number of predecessors; unused labels do not matter. This requires no selection of class representatives. For $x \in L$, let $d_x(\xi)$ be the label of the child containing x .

If $x \neq y$, there is a least stage at which they cease to be equivalent. It is a successor stage: at a limit, the intersection condition would give an earlier separating stage. Thus their label strings first differ at the corresponding coordinate. Convexity says that $x < y$ precisely when the first differing label of x is smaller. Therefore $x \mapsto d_x$ order-embeds L into $\{0, 1, 2\}^\theta$ in lexicographic order.

Replace each digit by a two-sign block

$$0 \mapsto --, \quad 1 \mapsto -+, \quad 2 \mapsto ++.$$

Concatenating θ blocks gives a sign sequence of length $2 \cdot \theta$. The replacement preserves the first differing coordinate comparison, hence gives (i). It also gives a binary convex resolution by equality of prefixes of these fixed-length sign sequences. Thus (ii) implies (iii), and (iii) implies (ii) trivially. $\square$

The theorem does not identify an optimal resolution length. Its value is to isolate a checkable *order-sensitive* certificate, distinct from the unconstrained powerset coding of the underlying set.

47 Orbit factorization of polynomial representations

The negative theory begins with a general algebraic lemma that does not mention surreals, atoms, or choice principles.

Let a group G_0 act on a set X . It acts on $k[T_X]$ by permuting variables and fixing the coefficient field k . Let X/G_0 denote the set of its orbits. Put

$$I_{G_0} = (T_x - T_y : x, y \text{ belong to the same } G_0\text{-orbit}). \quad (47.1)$$

Theorem 47.1 (Orbit quotient and factorization). *There is a canonical k -algebra isomorphism*

$$k[T_X]/I_{G_0} \cong k[T_{X/G_0}].$$

If D is a k -algebra on which G_0 acts trivially and a k -algebra homomorphism $\phi : k[T_X] \rightarrow D$ satisfies

$$\phi(h \cdot p) = \phi(p) \quad (h \in G_0),$$

then it factors uniquely through this quotient. If X/G_0 has $q < \omega$ members and D is a field, then

$$\text{trdeg}_k \text{Frac}(\text{im } \phi) \leq q. \quad (47.2)$$

Equality holds precisely when the q orbit values are algebraically independent over k ; in that case $\ker \phi = I_{G_0}$.

Proof. The orbit map $x \mapsto [x]$ induces a surjective homomorphism

$$\pi : k[T_X] \longrightarrow k[T_{X/G_0}].$$

It kills I_{G_0} . Conversely, in $k[T_X]/I_{G_0}$ all variables from one orbit have the same residue. Assign to an orbit this uniquely determined residue; existence follows from the nonemptiness of the orbit, and uniqueness follows from the defining relations. This is a function defined by a unique-value condition, not a choice of representatives. The polynomial universal property gives an inverse to the induced map from the quotient. The isomorphism follows.

Invariance gives $\phi(T_x) = \phi(T_{hx})$, so $I_{G_0} \subseteq \ker \phi$ and the unique factorization follows. With q orbits the image field is generated over k by their q values. Its transcendence degree is at most q . By finite field theory it equals q exactly when those values are algebraically independent. Equivalently, the factored map from $k[T_{X/G_0}]$ is injective, which is exactly the assertion that the original kernel is I_{G_0} . $\square$

Corollary 47.2 (Obstruction for fraction fields). *If a G_0 -orbit in X has two members, there is no invariant unital field homomorphism $k(T_X) \rightarrow D$ to a field with trivial G_0 -action.*

Proof. Such a map would kill the nonzero element $T_x - T_y$ for two variables in that orbit. A unital homomorphism from a field has zero kernel. $\square$

Remark 47.3 (Coinvariants, not invariant polynomials). The quotient in [theorem 47.1](#) identifies variables. It is not the invariant subring $k[T_X]^{G_0}$, whose structure can be quite different. Confusing these two constructions would invalidate the transcendence bound.

48 Sharp support bounds in two permutation models

48.1 The model framework

Start with a ground universe of ZFA + AC with a set A of atoms. A permutation group G on A acts recursively on sets, fixing pure sets. Let the normal filter of subgroups be generated by the pointwise stabilizers

$$\text{Fix}_G(S) = \{g \in G : g(a) = a \text{ for every } a \in S\}$$

of finite subsets $S \subseteq A$. The hereditarily symmetric sets form the usual full Fraenkel–Mostowski model W . The permutation-model construction and its set-theoretic setting are standard; [21, Section 3.1] provides a recent primary-source presentation of this machinery.

Every set of W has some finite atom support S : it is fixed by every member of $\text{Fix}_G(S)$. Every pure ground set belongs to W , and the pure part is unchanged. In particular, the reals, ordinals, and surreals of W are pure. Each atom permutation fixes each of them individually.

A map ϕ in W with pure values and support S therefore satisfies

$$\phi(g \cdot x) = g \cdot \phi(x) = \phi(x) \quad (g \in \text{Fix}_G(S)). \quad (48.1)$$

For polynomial domains the induced action permutes indeterminates. This turns the algebraic factorization theorem into a precise internal obstruction.

48.2 The basic Fraenkel model

Take A infinite and $G = \text{Sym}(A)$, with finite supports.

Theorem 48.1 (The $s + 1$ bound). *In the basic Fraenkel model, let*

$$\phi : \mathbb{R}[T_A] \longrightarrow \mathbf{No}$$

be an internal $\mathbb{R}$ -algebra homomorphism with atom support S , where $|S| = s < \omega$. Then

$$\text{trdeg}_{\mathbb{R}} \text{Frac}(\text{im } \phi) \leq s + 1. \quad (48.2)$$

All variables indexed by $A \setminus S$ have the same image. For every finite S the bound is attained by a homomorphism supported by S . No such homomorphism is injective, and there is no internal unital field homomorphism $\mathbb{R}(T_A) \rightarrow \mathbf{No}$.

Proof. The stabilizer $G_0 = \text{Fix}_G(S)$ has exactly $s + 1$ orbits on A : the s singletons in S and the infinite complement $A \setminus S$. Any two atoms in that complement can be swapped while all of S is fixed. Equation (48.1) and theorem 47.1 prove the collapse and the bound. Two distinct atoms outside S give a nonzero difference in the kernel. The field assertion follows from corollary 47.2.

For sharpness, choose a listing $S = \{a_1, \dots, a_s\}$ of this one finite set. Assign distinct members $z_0, \dots, z_s$ of a pure algebraically independent family, for example

$$z_i = \omega^{\omega^i}, \quad 0 \leq i \leq s,$$

by sending T_{a_i} to z_i and every variable outside S to z_0 . This assignment and its polynomial evaluation graph are supported by S and hereditarily symmetric: each individual input polynomial involves only finitely many atoms, while every output is pure. Thus the map belongs to W . Its orbit values are algebraically independent, giving equality in (48.2). This construction uses only the finite choice of a listing of one given finite set, which is available without AC. $\square$

48.3 The ordered Mostowski model

Now take an atom set A with a dense linear order without endpoints externally isomorphic to $(\mathbb{Q}, <)$, and let

$$G = \text{Aut}(A, <)$$

with the same finite-support filter. The ordered set $(A, <)$ itself belongs to the symmetric model: its order relation is invariant under the whole group.

Lemma 48.2 (Stabilizer orbit count). *If $S = \{a_1 < \dots < a_s\} \subseteq A$, the pointwise stabilizer of S has exactly $2s + 1$ orbits:*

$$\{a_1\}, \dots, \{a_s\}, \quad (-\infty, a_1), (a_1, a_2), \dots, (a_s, \infty).$$

For $s = 0$, there is just the orbit A .

Proof. Order preservation and fixation of S prevent movement between the displayed pieces. Within any one interval, any chosen atom can be sent to any other by an order automorphism fixing all the endpoints in S . To justify existence, work in the ground universe with its fixed external countable presentation of the dense order. Extend the finite increasing partial map consisting of the fixed points of S and the proposed movement by the standard back-and-forth construction. At each alternating stage put the next unused domain or range point into the corresponding nonempty interval. Density and absence of endpoints ensure success. The resulting bijection is order preserving and fixes S . Thus each interval is a single orbit. This external argument computes the chosen permutation group; it does not put a countable enumeration of A into the symmetric model. $\square$

Theorem 48.3 (The $2s + 1$ bound). *In the ordered Mostowski model, every internal $\mathbb{R}$ -algebra homomorphism*

$$\phi : \mathbb{R}[T_A] \longrightarrow \mathbf{No}$$

with a support S of size s satisfies

$$\text{trdeg}_{\mathbb{R}} \text{Frac}(\text{im } \phi) \leq 2s + 1. \quad (48.3)$$

For every finite S , equality is attained by a map supported by S . No such map is injective, and there is no internal unital field homomorphism $\mathbb{R}(T_A) \rightarrow \mathbf{No}$.

Proof. Apply [theorem 47.1](#) to the $2s + 1$ orbits of [lemma 48.2](#), using [\(48.1\)](#). Every interval contains distinct atoms, so the kernel contains a nonzero variable difference; the field assertion follows as before.

The orbits themselves have a canonical finite order, alternating intervals and singletons. Assign to these ordered orbits the algebraically independent pure values $z_0, \dots, z_{2s}$ from the proof of [theorem 48.1](#). The resulting evaluation map is supported by S , belongs to the symmetric model, and has algebraically independent orbit values. This attains the bound. $\square$

Corollary 48.4 (Exact kernels at maximum dimension). *For each of the sharpness examples above, the entire kernel is generated by differences between variables in the same stabilizer orbit. There are no further polynomial relations among the orbit values.*

This follows from the last clause of [theorem 47.1](#). The upper bounds apply to any chosen support, not necessarily a smallest support. As the support size varies, maps of arbitrarily large finite image-field transcendence degree exist, but no one internal map has infinite image-field transcendence degree.

48.4 Why pure coordinates recover well-orderability in these models

Proposition 48.5. *In either full permutation model just described, for any internal set X the following are equivalent: X injects into a pure set; X is well-orderable internally; X injects into the internal $\mathbf{No}$.*

Proof. A pure set belongs to the unchanged pure ground universe, which satisfies choice. It has a pure well-order, and that well-order also belongs to the symmetric model. An internal injection into the pure set therefore pulls back an internal well-order of X . Conversely, a well-order identifies X with an ordinal, which is pure and embeds into $\mathbf{No}$.

An injection into $\mathbf{No}$ has a set-sized range; its range is a set of pure objects and is itself pure. Thus it gives an injection into a pure set. Alternatively, apply [theorem 42.1](#) and use the purity of $\mathcal{P}(\alpha)$ in the full permutation model. $\square$

There is no conflict with KWP_1 being strictly weaker than choice in pure ZF. The preceding proposition makes an additional assumption: the unchanged pure part of these particular atom models already satisfies choice. It is not a theorem about arbitrary models of ZFA.

49 An ordered field with all finite stages embeddable but no global embedding

Theorem 49.1 (Internal ordered-field counterexample). *In the ordered Mostowski model W , the set-sized ordered field*

$$F(A) = \mathbb{R}(T_a : a \in A)$$

with the largest-variable order of [lemma 45.1](#) has no unital field homomorphism into $\mathbf{No}^W$, even when order preservation is not required. Nevertheless, for every finite $J \subseteq A$, the subfield $F(J)$ has an internal ordered $\mathbb{R}$ -embedding into $\mathbf{No}^W$, and

$$F(A) = \bigcup_{J \in [A]^{<\omega}} F(J).$$

Proof. The polynomial ring, fraction-field equivalence relation, and field operations are invariant under every order automorphism of A . Each expression is hereditarily symmetric, since it uses only finitely many atoms and pure real coefficients. The largest-variable order is invariant by [lemma 45.1](#). Hence the ordered field is an object of W .

An internal field homomorphism into $\mathbf{No}^W$ would have a finite atom support. Restricting it to $\mathbb{R}[T_A]$ gives a map whose values are fixed by the stabilizer, whether or not the homomorphism fixes every real. In particular, two variables from one nonsingleton interval orbit have the same image. Their nonzero difference cannot be killed by a unital field homomorphism. This is a contradiction.

For a finite $J = \{a_1 < \dots < a_n\}$, the order-preserving map $a_i \mapsto i$ is a set map supported by J . The construction of [theorem 45.2](#) gives an ordered embedding $F(J) \rightarrow \mathbf{No}^W$. Its graph is supported by J and hereditarily symmetric, so it belongs to W . Every rational expression uses only finitely many variables, proving the union identity. $\square$

Remark 49.2 (Scope of the counterexample). This proves a failure of ordered-field universality in a model of ZFA. It is *not* asserted to be a counterexample in pure ZF. Removing atoms while preserving a statement about embeddings into the entire internal surreal class requires a separate transfer argument; none is supplied here. The external countability of A also causes no contradiction: an external enumeration can produce an external embedding, but such an enumeration and embedding do not belong to W .

49.1 The exact coherence obstruction

Theorem 49.3 (Gluing compatible coordinate embeddings). *Let L be a set-sized linear order in ZF or ZFA. A global ordered $\mathbb{R}$ -embedding $F(L) \rightarrow \mathbf{No}$ exists if and only if there is a set-indexed family*

$$(\phi_J : F(J) \hookrightarrow_{\text{ord}} \mathbf{No})_{J \in [L]^{<\omega}}$$

of ordered $\mathbb{R}$ -embeddings satisfying

$$J \subseteq K \implies \phi_K|_{F(J)} = \phi_J. \quad (49.1)$$

Proof. Restrict a global embedding to obtain the family; Replacement gives its set code. Conversely, take the union of the graphs of the ϕ_J . This union is a set. If an element lies in both $F(J)$ and $F(K)$, compare both maps with $\phi_{J \cup K}$ using (49.1); their values agree. Thus the union is a function on $F(L)$. Any finitely many arguments and their algebraic relations lie in one $F(J)$, so the union is an order-preserving field homomorphism. The same observation shows injectivity. $\square$

For the field in [theorem 49.1](#), each finite stage has an ordered embedding but no compatible family exists internally. This is not a claim that first-order compactness fails, and it does not concern the existence of a model of a finitely satisfiable first-order theory. The target is the particular internal class $\mathbf{No}$, and the obstruction is the demanded compatibility and internal existence of its coordinate map.

49.2 Finite polynomial tests can all succeed

Proposition 49.4 (Finite separation without a faithful representation). *In either of the two permutation models, every finite family of nonzero polynomials in $\mathbb{R}[T_A]$ is simultaneously nonzero under some internal $\mathbb{R}$ -algebra homomorphism to $\mathbf{No}$. Nevertheless, no one internal homomorphism is injective.*

Proof. Let J be the finite union of the variable supports of the given polynomials. Assign algebraically independent pure surreal values to the variables in J , and assign a fixed pure value, say 0, to every other variable. The evaluation map is supported by J and belongs to the model. Its restriction to $\mathbb{R}[T_J]$ is injective by algebraic independence, so it preserves the nonzeroness of every polynomial in the family. Nonexistence of an injective map follows from [theorems 48.1](#) and [48.3](#). $\square$

Equivalently, for each nonzero polynomial there exists a map not killing it, but no single map avoids all nonzero kernels. One may describe the intersection of all these kernels as zero, provided that phrase is read as a first-order quantification over maps, not as the formation of a set of all maps into a proper class.

50 Proof audit and relation to the existing repository

50.1 Dependency boundary

The main logical dependencies are short:

$$\begin{aligned} \text{sign sequences} + \text{Replacement} &\implies \text{ordinal-code equivalence,} \\ \text{finite normal forms} + \text{monomial multiplication} &\implies \text{free-algebra equivalence,} \\ \text{canonical fraction/root codes} + \text{real closedness} &\implies U_\kappa, \\ \text{polynomial orbit quotient} + \text{finite atom support} &\implies \text{sharp collapse bounds.} \end{aligned}$$

The negative construction does not assume any speculative research claim from ProveIt's other subjects. The positive algebraic construction uses only the standard surreal interface in [section 41](#). Thus an unrelated error in an unrefereed repository report would not affect the mathematical dependency chain here.

50.2 A ledger of theorem status

Result	Mathematical status in this manuscript	Novelty / checking status
Ordinal-code equivalence	Direct proof in ZF and ZFA; exact birthday capacity included	Elementary consequence of sign coding; no priority claim
Double-monomial family	Explicit finite-normal-form proof; free rings and fields calibrated	Classical ingredients; combined formulation developed here
Optimal field U_κ	Set existence, real closedness, and two cardinal injections proved without quotient-choice shortcut	Candidate synthesis/refinement; independent review needed
Convex resolution	Necessary and sufficient order-coding certificate proved	Related to ordered coding; historical novelty not established
Orbit factorization	Full algebraic proof, with finite-dimensional equality criterion	Standard universal-property mechanism
$s + 1$ and $2s + 1$ bounds	Exact stabilizer counts, upper bounds, and sharp constructions	Quantitative application developed here; priority unverified
Ordered-field failure	Internal counterexample in the stated ZFA permutation model	No automatic pure-ZF transfer asserted
Coherent gluing	Set-graph union proof; finite separation distinguished	Standard gluing mechanism, applied to fixed target

No theorem in this manuscript is presented as already kernel-checked. The supplied finite tests detect mistakes in finite encodings and algebraic identities but do not prove a theorem about all ordinals, arbitrary models, or proper classes.

50.3 The choice issue in a Lean audit

ProveIt’s surreal README reports a standard-axiom audit allowing `propext`, `Quot.sound`, and `Classical.choice`; it maps checked statements separately from research claims [26]. That is valuable evidence about the declared Lean foundation. It is not a certificate that a theorem is provable over *object-theoretic* ZF without choice.

There are two defensible routes for the present results. One can formalize enough constructive set and ordinal coding to avoid the choice axiom in the relevant dependency chains. Alternatively, one can work in Lean’s ordinary classical metatheory while explicitly formalizing a model of ZF or ZFA and proving that the relevant internal statements follow from the designated object-theoretic axioms. Metatheoretic choice then does not silently become internal choice; the semantic or syntactic theorem must enforce that distinction.

A theorem about every type in a universe is also not literally a theorem about the proper class of all sets. The birthday-bounded targets $\mathbf{No}_{\leq \alpha}$ and the fields U_κ offer useful set-sized interfaces, but a formalization must still record which universe contains each type and which internal sets it represents.

[Added 3 October 2026, batch 89: the description of `Algebra/SurrealNumbers/README.md` is accurate at the pinned commit `2b7b388ba` and at the placement commit `23adb85f9`; the README blob source `11` inspected, `18478d586`, is unchanged. No module named below exists in the repository, and no statement of this part is formalized; placement in this report confers no formal status.]

50.4 A proposed module decomposition

The following are suggested module responsibilities, not names of files already implemented in the repository.

OrdinalSignCoding

Define variable-length sign functions, the definedness code (42.1), its decoder, and fixed-length subset encoding. Prove the birthday-capacity theorem with explicit ordinal pairings.

DoubleMonomialIndependence

Abstract the finite-normal-form leading-term property. Prove that distinct monomials are linearly independent and that the double-monomial substitution is an injective multivariate-polynomial homomorphism. Derive the omnific range condition separately from the real-coefficient assertion.

CanonicalFieldCodes

Normalize multivariate fractions by gcd and leading coefficient, code algebraic roots by minimal polynomial and root index, and construct the set-sized relative real closure.

OrbitPolynomialQuotient

Formalize the group action on variables, the orbit-indexed polynomial algebra, the quotient isomorphism, and finite transcendence bounds. This module can be independent of surreal theory.

InternalSymmetryBridge

Formalize finite-support evaluation into a pure target inside a specified permutation model. The semantic statement that an internal map has a finite support is indispensable; a theorem only about an externally chosen group action is not yet the model-theoretic conclusion.

OrderedFreeFieldAndCoherence

Construct the largest-variable order using finite supports, prove the ordered embedding reduction, and formalize gluing of compatible finite-stage maps by a set union.

51 Further research questions and concrete next targets

The following are proposed research directions. Except where a cited paper explicitly supplies a status, they should not be read as claims that the literature has left each formulation open. A literature check and, where appropriate, contact with specialists should precede any priority claim.

Question 51.1 (Pure-set prescribed-order universality). What is the exact strength over ZF of the assertion that every set-sized linear order embeds in **No**? In particular, can one construct a pure-ZF analogue of the ordered-field counterexample in [theorem 49.1](#)?

The atom proof isolates the desired obstruction but does not transfer it. A useful first target is a pure symmetric extension containing a linearly ordered set with no ordinal-length convex resolution. One must analyze its *internal* sign sequences and maps rather than merely preserve a low-rank atom surrogate.

Question 51.2 (Does KWP_1 supply an order-sensitive code?). Does KWP_1 imply that every linear order admits a binary convex resolution of some ordinal length?

By [theorem 46.2](#), this is exactly the missing implication from coordinate universality to prescribed-order universality. A possible approach is to study a given order on a subset of $\mathcal{P}(\alpha)$ and ask whether membership coordinates can be converted into a continuous sequence of convex refinements. Arbitrary coordinate cuts need not be convex, which is the concrete obstruction to the naive argument.

Question 51.3 (Optimal convex-resolution height). For an order that embeds in **No**, how does its least binary resolution length compare with the least birthday bound of an ordered embedding?

The proof gives conversions with a factor of two at the ternary-to-binary step, but no optimality statement. Fixed-length and variable-length sign representations should be distinguished. Finite chains, ordinal orders, reversed ordinal orders, and lexicographic products provide initial test cases.

Question 51.4 (Explicit birthday bounds for U_κ). Find useful explicit ordinal bounds for the birthdays of all members of the field U_κ defined in [\(44.1\)](#).

Existence is immediate from its set-sizedness, but an effective bound requires controlling the two monomial operations and algebraic root formation. A conservative closure ordinal for a specified finite-expression grammar is a reasonable first goal; a sharp product-birthday formula should not be assumed.

Question 51.5 (More general fields at a fixed code level). Which additional hypotheses on a prescribed ordered field K with $K \hookrightarrow \mathcal{P}(\kappa)$ guarantee an ordered embedding into U_κ or into a canonically enlarged target of the same coding capacity?

Free rational-function fields with induced orders are handled here. Algebraic ordered extensions, valuations with order-embeddable value groups, and fields given by a coherent tower of simple ordered extensions are natural successive cases. The existence of a chosen transcendence basis must not be silently assumed.

Question 51.6 (Higher Kinna–Wagner levels). Is there a natural algebraic representation problem, not merely a relabelling of the definition, whose exact strength is KWP_n for $n > 1$?

Ordinary surreal elements have one-level ordinal subset codes. Higher levels might arise from unordered families of coordinates, quotient presentations, or structured moduli. The challenge is to exhibit algebraically meaningful invariants whose coding genuinely requires the extra powerset level.

Question 51.7 (Orbit profiles beyond the two examples). For a permutation group G on atoms, how precisely does the function

$$S \mapsto |A/\text{Fix}_G(S)|$$

control the transcendence spectra of pure-target polynomial representations?

The finite-orbit case is settled by [theorem 47.1](#). Infinite orbit sets introduce a new coding issue: a sharp realization requires enough independent pure coordinates indexed by the orbit set. Applying [theorem 43.5](#) to that quotient suggests an exact two-part criterion, combining orbit count and ordinal-code availability.

Question 51.8 (Minimal supports and essential dimension). Can one recover a lower bound on the least atom-support size of a map from the transcendence degree of its image, and when is the bound exact for every map?

Our inequalities immediately give $s \geq d - 1$ in the basic model and $s \geq \lceil (d - 1)/2 \rceil$ in the ordered model. They are sharp as existential bounds. A finer classification would distinguish maps with redundant support from those whose support is essential and relate this to the structure of their kernels.

Question 51.9 (Real closures under symmetry). What is the strongest ordered-algebraic enlargement of the source in [theorem 49.1](#) that can be constructed internally while retaining an explicit pure-target obstruction?

An externally selected real closure is not automatically an internal object with the desired action. A canonical construction through real-algebraic root codes, if it can be made equivariant over the ordered field, is a promising route. Any such extension would inherit nonembeddability by restriction to the original rational-function field.

Question 51.10 (Choice strength of compatible local embeddings). For which families of subfields does the existence of an individual embedding for every member imply a compatible family of embeddings into a fixed real-closed target?

The union theorem proves what compatibility would accomplish, not that it can be obtained. Separate chains with specified extension maps, directed systems indexed by finite subsets, and unrestricted systems. A useful target is an exact weak-choice principle for one sharply specified family rather than an undifferentiated appeal to compactness.

Question 51.11 (Residual separation and inverse limits of representations). Can the finite-separation phenomenon of [proposition 49.4](#) be organized as a failure of a natural inverse-limit existence principle, with a precise choice calibration?

For each finite variable set, consider its space of faithful coordinate assignments and restriction maps. The naive inverse system need not have surjective transition maps with a fixed birthday bound. Understanding that issue is essential before invoking any inverse-limit or compactness theorem.

Question 51.12 (Object-theoretic formal verification). Can the ZF equivalence and the ZFA counterexample be checked in one repository development with an explicit audit of internal versus metatheoretic choice?

The shortest first milestone is the orbit polynomial quotient plus the finite-normal-form independence lemma. The hard foundational milestone is a bridge from hereditarily symmetric internal maps to stabilizer invariance, together with set-coded surreal targets. Such a development would test the semantics of the statement, not just the algebra of the displayed proof.

52 Reproducibility and limits of the computational evidence

The accompanying Python program, shipped as `code/11-choiceless-universality-verify_finite.py` (delivered as `verify_finite.py`), uses only the standard library and a fixed random seed. It checks the finite definedness encoding, fixed-length ternary-to-binary order encoding, distinct exponent-vector substitution, translation invariance of largest-variable order, leading terms of polynomial products, and finite permutation-orbit quotient identities. It also checks simultaneous evaluation as a ring homomorphism by exact integer arithmetic. The supplied deterministic run passed 20,595 assertions; this count is evidence only about the finite checks just described.

These checks are deliberately finite. A finite linearly ordered set has only the identity order automorphism, so its automorphism group is *not* a valid finite simulation of the ordered Mostowski model. The program does not pretend otherwise. Infinite interval transitivity is proved by the ground-model back-and-forth argument in [lemma 48.2](#); ordinal recursion, class size, and model existence are mathematical arguments, not outputs of the script.

In the delivered layout the local checks were reproduced by

```
python3 verify_finite.py
```

followed by two `pdflatex` passes of the delivered manuscript. Without `-output` the program writes `verification_results.json` beside itself; with the shipped names, run it on a copy or pass an explicit output path outside this directory (the report's `README.md` gives the commands). The recorded run is shipped as `data/11-choiceless-universality-verification_results.json`. For this merge the program was rerun with Python 3.14.4 on a copy: it passed and reproduced the recorded file exactly, apart from line endings. The delivered archive contained the source, compiled PDF, verification script and its result file, a `README`, a source-and-status note and a checksum file; of these only the script, the result file and the source-and-status note (`11-choiceless-universality-SOURCES_AND_STATUS.md`) are shipped ([section 53](#)). No font files are distributed.

53 Conclusion of source 11, checklist and provenance

53.1 Conclusion

Surreal universality is not one assertion. For unconstrained independent variables, the exact obstruction is ordinal subset coding, and double Conway monomials convert that code into omnific algebra. This conversion can be carried out inside an optimally sized real-closed field at each infinite ordinal code level. When an existing order must be preserved, a convex resolution, rather than a mere injection of underlying sets, supplies the relevant certificate.

In permutation models the obstruction becomes quantitative. A pure-valued map supported by finitely many atoms cannot distinguish variables that its stabilizer moves into one another. The number of stabilizer orbits is therefore a sharp upper bound on the independent transcendence retained by the image. In the ordered model this produces an actual ordered field with no internal surreal embedding, while every finite coordinate stage embeds and every finite polynomial test can succeed.

The resulting program is concrete: distinguish coding from order, distinguish external from internal maps, and distinguish finite satisfiability from coherent realization. These distinctions are both mathematically substantive and well suited to a proof-assistant audit of choiceless surreal algebra.

53.2 A compact checklist for reviewing the main proofs

A referee or formalizer can focus first on the following points.

1. In [theorem 42.1](#), check both definedness and signs in the code, the side of ordinal multiplication, and the Replacement bound for the image birthdays.
2. In [theorem 43.1](#), check that the inner monomials are linearly independent, not merely distinct; this is what separates evaluated exponent vectors. Check the positive-exponent condition separately for the omnific conclusion.
3. In [lemma 44.1](#), check uniqueness of normalized fractions in the full arbitrary-variable polynomial ring. A surjection from a coded syntax set is not enough for the cardinal upper bound without a canonical code of each value.
4. In [theorem 44.3](#), verify collection of the algebraic roots as a set, and use the root index in the actual ordered extension. No choice of one root from each polynomial is made.
5. In [theorem 45.2](#), compare the two leading-term orders rather than assuming an arbitrary algebraic embedding preserves order.
6. In [theorem 46.2](#), use continuity at limits to justify that a first separation occurs at a successor stage.
7. In [theorem 47.1](#), distinguish the orbit quotient from the invariant subring, and avoid selecting one representative from every orbit.
8. In [theorems 48.1](#) and [48.3](#), apply atom support to the *graph of the map*; pure values alone do not constrain an external map.
9. In [theorem 49.1](#), verify that the source ordered field is an internal object and that the forbidden map is internal. Do not transfer the conclusion from ZFA to ZF without proof.
10. In [theorem 49.3](#), require a single set-coded compatible family; separate existence of each finite-stage map is weaker.

53.3 Non-claims of source 11

Every limitation stated by source 11, in its manuscript or in its source-and-status note, is kept; most appear at their point of use. Collected: the manuscript is AI-generated and unrefereed and is not a Lean- or Rocq-checked development; the novelty of its exact formulations is not established, and its twelve questions are proposed targets, not catalogued open problems; it claims no solution of Glazer and Yao’s reflection questions or of another named problem; the atom-model counterexample is a result in ZFA, and no ZF transfer is supplied or asserted; universality for prescribed orders does not follow from KWP₁ in what is proved; no sharp birthday bound for U_κ is claimed; KWP₁ and its coding form are classical (Karagila–Schilhan), as are sign coding, normal forms, the polynomial and

UFD facts, real-closed-field facts and the permutation-model construction; the finite script checks encodings and algebraic identities only, does not construct symmetric models, prove independence or check real closedness, and does not use finite chains as a model of $\text{Aut}(\mathbb{Q}, <)$; the proposed modules are a roadmap; the repository was not modified, built or verified.

53.4 Provenance of this part

The manuscript. Source 11 is batch-89 manuscript 04, delivered as the archive `Glazer_Surreal_Choice_Research.zip` in commit `7c0f2d9f9` and placed in commit `23adb85f9`. It records the README blob `18478d586` of `Algebra/SurrealNumbers/README.md` (unchanged at the placement commit) and the identifier `2b7b388ba`, which it calls a “main-tree” identifier, “not asserted to be a commit identifier”; it is in fact the commit “Catalogue batches 81 to 86”. Its bibliography linked the surreal README through the moving branch `main`; the entry [26] here gives the pinned commit. Its manuscript (`article.tex`, 911 lines, 52 labels; a 22-page PDF), README, PDF and checksum file `SHA256SUMS` (6 of 6 entries verified at placement) are not shipped; they survive in the arrival commit. Shipped, byte-identical: `11-choiceless-universality-SOURCES_AND_STATUS.md`, `code/11-choiceless-universality-verify_finite.py` and `data/11-choiceless-universality-verification_results.json`. The source-and-status note keeps the delivered wording about the identifier `2b7b388ba` and mentions a profile page given to the authoring session as an identifying pointer, without a URL. The Hilbert’s-tenth programme’s routing review of commit `6aadaa4f2` covers this archive too (section 39.5).

How it is printed. The text from source 11’s title-page boxes and abstract to its appendix is printed in full, in its order: the title-page boxes and the abstract open section 40, its Section 1 follows them, and its appendix became a subsection of this section. Every delivered label carries the prefix `hset:kw:`, and its twelve questions gained labels `hset:kw:q:...`. The symbols renamed are listed in convention 40.1; delivered file names are replaced by the shipped ones; its references to Glazer–Yao and to Conway use this report’s bibliography entries. Added for this report: sections 40.4 and 40.5, the bracketed dated note, this subsection and the preceding one. No mathematical statement or proof of source 11 was changed.

Where the merge had to choose. (i) Placement: source 11 cites no report of the collection; it is placed here because it answers in part the open item of question 21.7 and is the injection counterpart of theorem 13.2 (remarks 40.2 and 40.3). (ii) Theorem 43.1 is printed with its own proof although it is a case of a lemma of another report (remark 40.4). (iii) The two new parts follow Part V, so that no existing number changes; both are printed after this report’s conclusion and before its appendices.

A Formula expansion and size checks

A.1 What “first-order” means in the coding proof

Every abbreviation expands to a finite formula in $\{0, 1, +, -, \cdot, <, b\}$. Bounded ordinal quantifiers are shorthand for ordinary scalar quantifiers with guards $\text{Ord}(x)$ and $x < \delta$. A set variable in

an interpreted membership formula becomes a triple of scalar variables. Equality becomes $\sim$, membership becomes $\in_{\mathcal{I}}$, and quantifiers are relativized to Code:

$$(u = v)^{\mathcal{I}} = (C_u \sim C_v), \quad (u \in v)^{\mathcal{I}} = (C_u \in_{\mathcal{I}} C_v), \quad (\exists u \varphi)^{\mathcal{I}} = \exists \delta_u \exists r_u \exists t_u (\text{Code}(C_u) \wedge \varphi^{\mathcal{I}}),$$

commuting with the connectives. Induction on formulas proves $H_{\kappa(\lambda)} \models \varphi(x_{C_1}, \dots) \iff \mathcal{B}_\lambda \models \varphi^{\mathcal{I}}(C_1, \dots)$ for valid codes, independently of the representatives.

For example, a graph code f on $\delta \times \epsilon$, embedded in the square of width $\theta = \delta + \epsilon + 1$, uses

$$F(i, j) \iff i < \delta \wedge j < \epsilon \wedge \text{Bit}(f, p_\theta(i, j)).$$

It is a total function exactly when

$$\forall i < \delta \exists j < \epsilon (F(i, j) \wedge \forall k < \epsilon (F(i, k) \longrightarrow k = j)).$$

Injectivity is $\forall i, i' < \delta \forall j < \epsilon (F(i, j) \wedge F(i', j) \rightarrow i = i')$, and surjectivity is $\forall j < \epsilon \exists i < \delta F(i, j)$. Occurrences of $f(i)$ in the main text can therefore be eliminated by these unique-value clauses. Root preservation and edge correspondence add only finitely many quantifiers. The graph code's birthday is $\Lambda(\theta)$; all plus positions outside the relevant rectangle can be required absent, which makes canonical relation coding a definable operation when the intended relation is specified by a formula and its bit string exists.

The dependency order of the abbreviations is

Predicate	Earlier definitions needed	Quantification simulated
$\text{Ord}(a)$	$b(a) = a$	Embedded ordinals
$\text{Pre}(x, y)$	$<, b$	Literal prefix test
$\text{Bit}(s, a)$	$\text{Ord}, \text{Pre}, <, b$	One sign position
E_C	Bit, p_δ	Edges of a code
$\text{Code}(C)$	shape, clean bits, (5.1)–(5.3)	Pointed membership graphs; all subsets of one domain
$\sim, \in_{\mathcal{I}}$	Code, E , coded maps	Interpreted equality and membership
$\text{OC}(a), \text{VisOrd}$	$\text{Ord}, \text{Bit}, p_\theta, \sim$	Literal ordinal as a set code
Coll	Code, $\in_{\mathcal{I}}, \text{VisOrd}$	Collection of literal ordinals

The table follows source 02's ledger, with this report's names. A tuple variable is eliminated by replacing it with ordinary variables; “exactly one” has its usual two-quantifier expansion; variables are renamed before substitution to prevent capture. The specification is an ordinary finite formula, albeit a long one.

A.2 Every potentially large construction has a stated bound

Object	Bound used in the proof
Subsets tested in well-foundedness	Individual sign strings of birthday δ ; no collection of that whole fiber is required.
A relation on a code domain δ	Sign birthday $\Lambda(\delta) = \delta^2 + \delta < \lambda$.
An isomorphism or member embedding	Square width $\delta + \epsilon + 1 < \lambda$, with the corresponding Λ bound.

Object	Bound used in the proof
Collapse value of a rooted code	Transitive closure has exactly $ \delta $ members.
An arithmetic computation	At most μ word vertices or table cells, hereditary size μ , where μ bounds the inputs and is infinite.
Rank labels in that computation	Below μ^+ , hence each has cardinality at most μ .
Replacement image after eternity	One bound $\max(\aleph_0, \delta , \varrho) < \kappa$, not an arbitrary singular-cardinal union.
A packed birthday- α fiber	Exists exactly if $2^{ \alpha } < \kappa(\lambda)$.
The old-power-set parameter	Hereditary size $(2^{ \alpha })^M$ for infinite α .
Actual power set of a set of hereditary size μ	At most 2^μ ; this need not be below κ , and no earlier row needs it to be.

B An essential negative regression

In the membership definition, edge preservation alone is insufficient. Take the code for $x = \emptyset$ and the three-node chain coding $y = \{\{\emptyset\}\}$. Map the sole node of the first code to the middle node of the chain, which codes $\{\emptyset\}$. The image root is a child of the target root, and the one-node induced subgraph has no edges, just like the source. But $\emptyset \notin \{\{\emptyset\}\}$. The predecessor-full clause (5.6) rejects this false witness, because the middle node has a predecessor omitted from the image. The finite suites of sources 06, 07 and 09 confirm this counterexample.

C A small concrete membership code

Let $x = \{\emptyset, \{\emptyset\}\}$, the von Neumann ordinal 2 (source 02's example). Its transitive closure with the root included has the three nodes $0 \leftrightarrow \emptyset$, $1 \leftrightarrow \{\emptyset\}$, $2 \leftrightarrow x$. Take $\delta = 3$, root $t = 2$, and a relation word of length $\Lambda(3) = 12$. The edges are $(0, 1)$, $(0, 2)$ and $(1, 2)$; under $p_3(u, v) = 3u + v$ the plus positions are 1, 2, 5, and all other positions of the length-12 word are minus. Its collapse is the identity on the ordinal 3, with root value 2.

A different labeling can put x at label 0, the empty set at label 2, and $\{\emptyset\}$ at label 1. The relation word then changes, and the root is no longer the largest label. Nevertheless the isomorphism code identifies the two roots, so interpreted equality cannot be literal equality of the triples. The example also shows why label order must not be confused with membership order in the countable-ordinal overflow of corollary 5.6.

References

- [1] J. H. Conway, *On Numbers and Games*, second edition, A K Peters, 2001; first edition, Academic Press, 1976. Classical number forms and recursive arithmetic.
- [2] H. Gonshor, *An Introduction to the Theory of Surreal Numbers*, London Mathematical Society Lecture Note Series 110, Cambridge University Press, 1986. <https://doi.org/10.1017/CB09780511629143>.

- [3] L. van den Dries and P. Ehrlich, “Fields of surreal numbers and exponentiation,” *Fundamenta Mathematicae* **167** (2001), 173–188. <https://doi.org/10.4064/fm167-2-3>. Epsilon-cutoff closure: Proposition 4.7 and Corollary 4.9. Read with the following erratum.
- [4] L. van den Dries and P. Ehrlich, “Erratum to ‘Fields of surreal numbers and exponentiation’,” *Fundamenta Mathematicae* **168** (2001), 295–297. Bibliographic record: <https://eudml.org/doc/281975>.
- [5] O. Bournez and Q. Guilmant, “Surreal fields stable under exponential and logarithmic functions,” arXiv:2201.08199, 2022. <https://arxiv.org/abs/2201.08199>. Theorems 1.1–1.2 restate the corrected ring, field and real-closedness cutoff results.
- [6] J. D. Hamkins, reporting joint work with J. Chen and R. Yang, “The elementary theory of surreal arithmetic is bi-interpretable with set theory,” Kobe, September 2025; announcement dated August 20, 2025, with linked slides. <https://jdh.hamkins.org/theory-of-surreal-arithmetic-bi-interpretable-with-set-theory-japan-september-2025/>. The slides announce the global result and discuss the coding and growth principles; they are not a complete refereed paper.
- [7] J. D. Hamkins, reporting joint work with J. Chen and R. Yang, “Surreal arithmetic,” Notre Dame Logic Seminar; posted 12 November 2025, talk 18 November 2025, with linked slides. <https://jdh.hamkins.org/surreal-arithmetic-notre-dame-logic-seminar-nov-2025/>.
- [8] J. D. Hamkins, reporting joint work with J. Chen and R. Yang, “Surreal arithmetic is bi-interpretable with set theory,” CUNY Logic Workshop, 13 March 2026; announcement posted 4 February 2026. <https://jdh.hamkins.org/surreal-arithmetic-cuny-logic-workshop-march-2026/>; workshop page <https://nylogic.github.io/logic-workshop/2026/03/13/surreal-arithmetic-is-bi-interpretable-with-set-theory.html>; slides <https://jdh.hamkins.org/wp-content/uploads/Slides-Elementary-theory-of-surreal-arithmetic-Hamkins-CUNY-March-2026.pdf>. The slides describe joint work in progress, with details still being checked.
- [9] E. Jeřábek, answer to O. Cunningham’s question “Are these theories of real and complex number biinterpretable?”, MathOverflow, 13 September 2021. <https://mathoverflow.net/questions/403410/are-these-theories-of-real-and-complex-number-biinterpretable>. Original explanation of the named-imaginary-unit and automorphism-group obstruction; not a journal publication.
- [10] A. R. Freire and J. D. Hamkins, “Bi-interpretation in weak set theories,” *The Journal of Symbolic Logic* **86** (2021), 609–634. <https://doi.org/10.1017/jsl.2020.72>; arXiv:2001.05262.
- [11] A. Poveda, A. Rinot and D. Sinapova, Sigma-Prikry forcing I: the axioms, arXiv:1912.03335v2, 2020. <https://arxiv.org/abs/1912.03335>. Lemma 2.10 and Section 3.1: bounded-subset preservation and ordinary normal-measure Prikry forcing. Published version: <https://doi.org/10.4153/S0008414X20000425>.
- [12] J. D. Hamkins, G. Kirmayer and N. L. Perlmutter, “Generalizations of the Kunen inconsistency,” arXiv:1106.1951v2, 2012. <https://arxiv.org/abs/1106.1951v2>. Theorem 1 and the metamathematical discussion give the class-embedding form used here.
- [13] T. Jech, *Set Theory*, third millennium edition, revised and expanded, Springer Monographs in Mathematics, Springer, 2003. <https://doi.org/10.1007/3-540-44761-X>.

- [14] D. Marker, *Model Theory: An Introduction*, Graduate Texts in Mathematics 217, Springer, 2002. <https://doi.org/10.1007/b98860>.
- [15] A. Deloro, *Basic Model Theory of Algebraically Closed Fields*, mini-course, Moscow Higher School of Economics, June 2013, Sections 3.4–3.5. <https://webusers.imj-prg.fr/~adrien.deloro/teaching-archive/Moskva-ACF.pdf>.
- [16] D. R. Rangel and H. L. Mariano, “An algebraic (set) theory of surreal numbers, I,” arXiv:1911.12726, 2019. <https://arxiv.org/abs/1911.12726>. A distinct categorical foundational program, not a dependency of the proofs.
- [17] W. Bertram, “On Conway’s Numbers and Games, the Von Neumann Universe, and Pure Set Theory,” arXiv:2501.04412v2, 2025. <https://arxiv.org/abs/2501.04412v2>.
- [18] P. Ehrlich, “Conway names, the simplicity hierarchy and the surreal number tree,” *Journal of Logic and Analysis* **3** (2011), article 1, 1–26. <https://doi.org/10.4115/jla.2011.3.1>.
- [19] P. Ehrlich and E. Kaplan, “Surreal ordered exponential fields,” arXiv:2002.07739, 2020. <https://arxiv.org/abs/2002.07739>.
- [20] E. Glazer, “Global choice is not conservative over local choice for Zermelo set theory,” arXiv:2312.11902, 2023; version 3, 2024. <https://arxiv.org/abs/2312.11902>. Cited by source 10.
- [21] E. Glazer and B. Yao, “Reflection Principles in ZFU,” arXiv:2602.21970v1, 25 February 2026. <https://arxiv.org/abs/2602.21970>. Source 10 uses Proposition 1, Proposition 17 and the small-kernel-model discussion; source 11 uses the urelement and permutation-model framework (Section 3.1), not a claimed extension of their reflection theorems.
- [22] J. D. Hamkins and B. Yao, “Reflection in second-order set theory with abundant urelements bi-interprets a supercompact cardinal,” *The Journal of Symbolic Logic* **89** (2024), 1007–1043; arXiv:2204.09766. <https://arxiv.org/abs/2204.09766>. The elementary support examples used for comparison by source 10 are in §1.5.
- [23] B. Yao, “Axiomatization and Forcing in Set Theory with Urelements,” arXiv:2212.13627; *The Journal of Symbolic Logic*, DOI:10.1017/jsl.2024.58. <https://arxiv.org/abs/2212.13627>.
- [24] A. Karagila and J. Schilhan, “Intermediate models and Kinna–Wagner Principles,” arXiv:2409.07352v1, 11 September 2024. <https://arxiv.org/abs/2409.07352>. Used by source 11 for the convention and standard status of the Kinna–Wagner principles.
- [25] O. Costin and P. Ehrlich, “Integration on the Surreals,” arXiv:2208.14331v5, 4 July 2024. <https://arxiv.org/abs/2208.14331>. Its positive development adopts a class framework without set or class choice; source 11 needs only the elementary surreal interface, not its integration results.
- [26] V. Reshetnikov, ProveIt repository, surreal-number development and its status README Algebra/SurrealNumbers/README.md, at commit 2b7b388ba81a3355b19a7c2d2fe797e92f572355, inspected 3 October 2026 by sources 10 and 11. <https://github.com/VladimirReshetnikov/ProveIt/tree/2b7b388ba81a3355b19a7c2d2fe797e92f572355/Algebra/SurrealNumbers>. Source 11 recorded the README blob 18478d58670c384953fb85dcd1a6d931c644aa32 and linked the branch main.

- [27] V. Reshetnikov, ProveIt repository, Closure axiomatization of set theory, `SetTheory/ClosureAxiomatization`, README and project organization at commit `2b7b388ba`, inspected 3 October 2026 by source 10.
<https://github.com/VladimirReshetnikov/ProveIt/tree/2b7b388ba81a3355b19a7c2d2fe797e92f572355/SetTheory/ClosureAxiomatization>.
- [28] The collection's report `foundations-and-computation/surreal-fields-across-universes` (*Surreal Fields Across Set-Theoretic Universes*), README at commit `2b7b388ba`, inspected 3 October 2026 by source 10; unrefereed; used to distinguish its topic from the present one.