

# The Fabius Function and Rvachev's Up-Function

Exact arithmetic, probability, Fourier analysis, and corrected sharp asymptotics

Human-readable synthesis of the formal development

Vladimir Reshetnikov's ProveIt repository

Integrated repository snapshot: 1 September 2026

Historical Cauchy/complex-product checkpoint a81a13b46; CDF/implicit checkpoint e6f535e5d; affine-law  
checkpoint d312c0603;

fractional-cutoff checkpoint 149332f9d; inverse-layer-cake checkpoint 37eab3c56; current-tree affine  
Cauchy-power tranche;

activation/Taylor checkpoint a345425d21d90e680bf15e34093af42c69c08a83

## Abstract

This article is the human-readable exposition of the formally verified Fabius-function development in the ProveIt repository. It keeps three related functions separate: the bounded distribution function  $F$ , Rvachev's even compactly supported bump up, and the signed global extension  $F_{\text{gl}}$ . Their construction and uniqueness, differential and refinement laws, strict shape properties, optimal Lipschitz constants, derivative bounds, exact dyadic-cell derivative value distributions, effective flatness, and precisely formalized analytic and non-elementarity results are presented together with direct pointers to their Lean proofs.

The probabilistic part identifies  $F$  and up with exact probability expressions for an infinite weighted sum of independent uniform coordinates. Finite atomic convolution laws, weak convergence, half-weighted step approximants, and centered uniform splines give proved finite models and quantitative approximation results. At the reusable level, arbitrary absolutely summable vector weights now have probability, head-tail, reflection, continuous-linear naturality, operator-affine recursion, and exact support-as-range laws, while every nontrivial absolutely summable real-weight law is absolutely continuous with respect to Lebesgue measure and has null singletons, and arbitrary signed real weights have exact min/max interval ranges and topological supports. A named geometric-uniform specialization supplies the weights  $(1 - q)q^n$ , their series and law, affine self-similarity and reflection, absolute continuity and null singletons for  $|q| < 1$ , support equal to the series range for  $|q| < 1$ , exact unit-interval support for  $0 \leq q < 1$ , and uniqueness among all probability laws satisfying the same affine independent-copy equation. The uniqueness theorem includes  $q = 0$  and negative  $q$ , and assumes no support, density, or moment condition. Its named CDF is continuous and reflection-symmetric for  $|q| < 1$ , has the exact exterior tails for  $0 \leq q < 1$ , and, for  $0 < q < 1$ , is  $C^\infty$  with an explicit nonnegative, compactly supported  $C^\infty$  Radon-Nikodym density; the half-base CDF and density recover the bounded Fabius function and the affine Rvachev bump exactly. A support-free quantile theorem transports uniform measure through the clamped Fabius inverse and gives Banach-valued quantile substitution. The analytic part develops the formal sinc and cosine products, Fourier inversion and Poisson summation, exact moment recurrences, and the Fourier-Legendre expansion. For each fixed complex strict contraction, the infinite q-Pochhammer symbol is also proved entire in its symbol variable, with its exact factor-zero and reciprocal-power zero descriptions and analytic order one at every zero. The arithmetic part gives reciprocal-power values, Thue-Morse formulas for arbitrary dyadic arguments, executable binary reduction, two-adic information, exact half-base values, and denominator-free arbitrary-base  $q$ -binomial and geometric-interpolation identities present in Lean. A compact-rectangle inverse-pair theorem integrates arbitrary absolutely-continuous outer functions without a clock-measurability hypothesis. Finite formal-series filters package arbitrary-node moment exactness and the complete denominator-free Gaussian residual on geometric nodes; their analytic companion gives the same head-tail law for unconditionally summable weighted sampled coefficient series, without a convergence obligation at zero-weight nodes.

A separate activation layer writes  $h$  for the continuous totalization of  $\tanh(x)/x$ , with  $h(0) = 1$ , and turns  $p(x) = 1 - h(x)$  into an exact square-summable calculus. It proves the sharp punctured limit  $p(at)/t^2 \rightarrow a^2/3$  for every real scale, genuine summability and a coefficient-sharp Tannery limit for arbitrary square-summable real weights, and the exact normalized geometric coefficient  $(1 - q)/(3(1 + q))$  for  $|q| < 1$ . Its dyadic value is  $1/9$ . The same layer identifies the totalized quotient with the divided slope of  $\tanh$  at zero and proves the finite jet  $p(x) = x^2/3 - 2x^4/15 + 17x^6/315 - 62x^8/2835 + O(x^{10})$ . These are deterministic analytic statements; no Bernoulli active-count expectation is inserted into them.

Repeated integration is developed once, before its Fabius specialization, as an oriented normalized Volterra calculus whose algebra takes values in an arbitrary real normed space. For continuous inputs in a complete target, literal iteration and the Cauchy kernel are identified exactly; strict FTC, additive composition of orders, Taylor reconstruction, and finite polynomial commutators then give closed forms for every integer-order primitive of the signed extension and every natural-monomial weight. A separate Gamma-normalized operator supplies the positive-real Riemann–Liouville kernel over arbitrary real normed spaces on ordered endpoints and agrees exactly with the normalized integer operator at every positive natural order. For continuous inputs in a complete target, two positive real orders compose additively on every ordered compact interval. Positive affine covariance, Gamma-normalized order raising from an interior right derivative, and the first exact fractional shifts of the signed, bounded, and folded Fabius functions are also proved on their stated ordered domains.

The endpoint chapters report only the negative-Laplace decompositions, periodic correction, lower-Lambert formulas, saddle estimates, and correction terms that have named proved counterparts in the formal source. Plausible extensions, numerical observations, conditionally convergent boundary filters, sinc-specific generic- $q$  acceleration, complex-order and iterated fractional formulas, and other arguments not yet represented by an exact Lean theorem are kept in the companion volume under `docs/semi-formalized-research-frontiers/`, not in this primary exposition.

---

**Artifact status (4 September 2026).** The authoritative live lexical audit contains exactly 923 facade-reachable source modules and 11,611 unique public declarations: 313 modules and 3,293 declarations beyond the 610/8,318 activation checkpoint, and 274 modules and 2,913 declarations beyond the 649/8,698 publication-union checkpoint. The merge target contributes 11,610 declarations; the retained unconditional public `complexQPochhammerInf_eq_qPochhammerInfIn` bridge in the `1+10 RvachevPochhammerFactorization.lean` surface contributes the additional declaration. Module headers and declaration documentation are complete; the checked baseline records both zero-gap invariants. This source includes the earlier finite and infinite  $q$ -series APIs together with the final Gaussian-palindromic, Jackson integral,  $q$ -exponential, theta, Jacobi-cubic, logarithmic-derivative, order-derivative, universal-polynomial, central-reduction, and cyclotomic-factorization tranche, followed by primitive-root blocks,  $q$ -Lucas evaluation, cyclotomic divisibility,  $q$ -Catalan polynomials, finite Newton interpolation, the real Jackson  $q$ -beta integral, Gaussian upper-index and quantum identities, superconvergent Rvachev synthesis, Gaussian bounds, the complete terminating  ${}_2\phi_1$  reversal and  $q$ -Chu–Vandermonde surfaces, the geometric Richardson generating-function module, the complete Lambert branch-coordinate surfaces and their zero-definition/five-theorem Bernoulli branch-gap companion, the six-theorem dyadic derivative filtration, and the three-theorem second-moment extension of `GaussianBinomialCumulants.lean`. The retained five-module polynomial  $q$ -calculus increment consists of `PolynomialQDerivative.lean` (2+17), `PolynomialQLeibniz.lean` (0+4), `QPochhammerDerivative.lean` (0+3), `LambertSeriesLog.lean` (0+4), and `QGamma.lean` (2+10): four definitions and thirty-eight theorems, forty-two public declarations, with the exact hypotheses and boundaries stated below. The retained companion PDF is a historical receipt for its named source checkpoint; this source-only merge requires a fresh uninterrupted three-pass Libertinus rebuild before parity can again be claimed. Exact historical source/PDF validation receipts are kept externally in the repository documentation so that neither artifact embeds a self-invalidating fingerprint. Package-local `SHA256SUMS*` ledgers remain abolished and do not participate in validation.

---

## Contents

|          |                                                                                                                        |           |
|----------|------------------------------------------------------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Historical note and scope</b>                                                                                       | <b>6</b>  |
| 1.1      | A guide to the main results . . . . .                                                                                  | 7         |
| <b>2</b> | <b>Three functions that must not be conflated</b>                                                                      | <b>7</b>  |
| 2.1      | The bounded Fabius distribution function . . . . .                                                                     | 7         |
| 2.2      | Rvachev’s compactly supported bump . . . . .                                                                           | 8         |
| 2.3      | The signed global extension . . . . .                                                                                  | 8         |
| 2.4      | Exact shape: positivity, bijectivity, and unimodality . . . . .                                                        | 9         |
| 2.5      | Convexity and the maximal slope . . . . .                                                                              | 10        |
| 2.6      | The optimal Lipschitz constant . . . . .                                                                               | 11        |
| <b>3</b> | <b>Existence and uniqueness by a contraction</b>                                                                       | <b>12</b> |
| 3.1      | The admissible function space . . . . .                                                                                | 12        |
| 3.2      | The integral transform . . . . .                                                                                       | 12        |
| 3.3      | The differential equation for up and smoothness . . . . .                                                              | 13        |
| 3.4      | The original Rvachev characterization . . . . .                                                                        | 13        |
| <b>4</b> | <b>The probability law behind the functions</b>                                                                        | <b>14</b> |
| 4.1      | An infinite weighted sum of uniforms . . . . .                                                                         | 14        |
| 4.2      | Generic weighted laws and support-free quantile transport . . . . .                                                    | 15        |
| 4.3      | The normalized geometric family . . . . .                                                                              | 21        |
| 4.4      | Sharp activation probabilities, finite Taylor jets, and square-summable fields . . . . .                               | 26        |
| 4.5      | Finite atomic laws and weak convergence . . . . .                                                                      | 30        |
| 4.6      | Combinatorial step functions and the half-weight convention . . . . .                                                  | 30        |
| 4.7      | Centered finite splines and an explicit error bound . . . . .                                                          | 32        |
| 4.8      | The exact quarter cell of the finite spline . . . . .                                                                  | 34        |
| <b>5</b> | <b>Fourier products, inversion, and Poisson summation</b>                                                              | <b>35</b> |
| 5.1      | Finite complex transforms and a Thue–Morse product-to-sum bridge . . . . .                                             | 36        |
| 5.2      | Geometric reciprocal-Gamma products . . . . .                                                                          | 40        |
| 5.3      | Polynomial $q$ -calculus, finite parameter derivatives, Lambert logarithms, and the real $q$ -gamma function . . . . . | 48        |
| 5.4      | Total local factorization and exact complex zero jets . . . . .                                                        | 50        |
| 5.5      | Leading Laurent coefficient of the reciprocal centered MGF . . . . .                                                   | 53        |
| 5.6      | Finite weighted scale arithmetic behind the zero orders . . . . .                                                      | 54        |
| 5.7      | Recovering generalized Rvachev exponents from dyadic orders . . . . .                                                  | 56        |
| 5.8      | Rapid decay and inversion . . . . .                                                                                    | 63        |
| 5.9      | Poisson summation and partitions of unity . . . . .                                                                    | 64        |
| 5.10     | Banach-free cyclic aliasing . . . . .                                                                                  | 64        |
| 5.11     | Finite odd-coset DFT traces . . . . .                                                                                  | 65        |
| 5.12     | Moment expansion of the Fourier transform . . . . .                                                                    | 66        |
| <b>6</b> | <b>The global extension, derivatives, flatness, and inverse</b>                                                        | <b>66</b> |
| 6.1      | The signed extension and all iterated derivatives . . . . .                                                            | 66        |
| 6.2      | Exact midpoint–endpoint transmutation . . . . .                                                                        | 67        |
| 6.3      | Exact inverse midpoint defect . . . . .                                                                                | 68        |
| 6.4      | Normalized Volterra calculus and exact Fabius primitives . . . . .                                                     | 69        |
| 6.5      | Reduced-dyadic derivative filtration . . . . .                                                                         | 74        |
| 6.6      | Dyadic derivative cells and exact value distributions . . . . .                                                        | 75        |

|           |                                                                                 |            |
|-----------|---------------------------------------------------------------------------------|------------|
| 6.7       | Effective flatness . . . . .                                                    | 77         |
| 6.8       | The precise analytic statements . . . . .                                       | 77         |
| 6.9       | The totaled inverse . . . . .                                                   | 78         |
| 6.10      | The exact inverse modulus . . . . .                                             | 78         |
| 6.11      | Effective dyadic continuity of the inverse . . . . .                            | 81         |
| 6.12      | Logarithmic reciprocal continuity of the inverse . . . . .                      | 82         |
| 6.13      | Sharp exact endpoint-mass denominators . . . . .                                | 82         |
| 6.14      | Fixed-depth tolerant inversion and computability of the total inverse . . . . . | 83         |
| 6.15      | Precise non-elementarity . . . . .                                              | 85         |
| <b>7</b>  | <b>Moments, generating functions, and reciprocal powers</b>                     | <b>85</b>  |
| 7.1       | Even and half moments . . . . .                                                 | 85         |
| 7.2       | Full centered moment and cumulant parity . . . . .                              | 87         |
| 7.3       | Universal Bell and cumulant transforms . . . . .                                | 87         |
| 7.4       | Rvachev moment–Appell deconvolution and polynomial synthesis . . . . .          | 89         |
| 7.5       | The probability–Laplace bridge . . . . .                                        | 95         |
| 7.6       | Exact values at reciprocal powers of two . . . . .                              | 98         |
| 7.7       | Parity and the exact two-adic valuation . . . . .                               | 98         |
| 7.8       | Bernoulli recurrences . . . . .                                                 | 99         |
| <b>8</b>  | <b>Exact evaluation at arbitrary dyadic rationals</b>                           | <b>99</b>  |
| 8.1       | The closed Thue–Morse formula . . . . .                                         | 99         |
| 8.2       | Thue–Morse cancellation . . . . .                                               | 101        |
| 8.3       | The exact leading-bit recurrence . . . . .                                      | 104        |
| 8.4       | The bit-recursive evaluator . . . . .                                           | 104        |
| 8.5       | Reduction of an arbitrary dyadic argument . . . . .                             | 105        |
| 8.6       | Denominator bounds . . . . .                                                    | 106        |
| 8.7       | The exact half-base $q$ -binomial layer . . . . .                               | 107        |
| 8.8       | The prefactor is the order of $\text{GL}_n(\mathbb{F}_2)$ . . . . .             | 124        |
| <b>9</b>  | <b>The global binary-reduction telescope</b>                                    | <b>126</b> |
| 9.1       | The corrected parity-power presentation . . . . .                               | 127        |
| 9.2       | The global complex $q$ -binomial series . . . . .                               | 128        |
| <b>10</b> | <b>A complex <math>q</math>-binomial discrete limit</b>                         | <b>129</b> |
| <b>11</b> | <b>Iterated prefix sums of the signed Thue–Morse sequence</b>                   | <b>141</b> |
| 11.1      | Iterated prefix sums and the binomial convolution law . . . . .                 | 141        |
| 11.2      | Prouhet annihilation, signed dyadic reciprocity, and exact zero runs . . . . .  | 142        |
| 11.3      | Two generating-series identities . . . . .                                      | 147        |
| 11.4      | The analytic Thue–Morse series and its natural boundary . . . . .               | 148        |
| 11.5      | The functional equation of the normalized prefix grid . . . . .                 | 149        |
| 11.6      | The literal right endpoint does not converge . . . . .                          | 149        |
| 11.7      | Corrected normalization and pointwise convergence . . . . .                     | 149        |
| 11.8      | The lower Lambert branch . . . . .                                              | 152        |
| 11.9      | A binomial–logarithm formula for the Thue–Morse bit . . . . .                   | 158        |
| <b>12</b> | <b>Endpoint asymptotics: the exact formal hierarchy</b>                         | <b>159</b> |
| 12.1      | The negative-Laplace product and its exact decomposition . . . . .              | 160        |
| 12.2      | The Gamma–zeta Fourier series . . . . .                                         | 161        |
| 12.3      | The evaluated constant . . . . .                                                | 161        |
| 12.4      | A fully effective dyadic bound . . . . .                                        | 162        |

|           |                                                                      |            |
|-----------|----------------------------------------------------------------------|------------|
| 12.5      | The lower-Lambert saddle and exact reduction . . . . .               | 162        |
| 12.6      | The compact online formula and its missing periodic factor . . . . . | 167        |
| 12.7      | The first two periodic saddle corrections . . . . .                  | 169        |
| 12.8      | The recursive all-orders theorem . . . . .                           | 170        |
| 12.9      | Universal finite endpoint-transfer algebra . . . . .                 | 171        |
| 12.10     | Recursive elementary re-expansion of the phase . . . . .             | 172        |
| 12.11     | Three endpoint consequences already formalized . . . . .             | 172        |
| <b>13</b> | <b>The Fourier–Legendre expansion of <math>\text{up}</math></b>      | <b>173</b> |
| <b>14</b> | <b>Additional proved summation identities</b>                        | <b>177</b> |
| <b>15</b> | <b>Effective computability</b>                                       | <b>177</b> |
| <b>16</b> | <b>Formalization boundary and audit conventions</b>                  | <b>178</b> |
| <b>17</b> | <b>Affine Cauchy–Stieltjes calculus</b>                              | <b>181</b> |
| <b>18</b> | <b>Conclusion</b>                                                    | <b>191</b> |
| <b>A</b>  | <b>Lean module map</b>                                               | <b>191</b> |
| <b>B</b>  | <b>Notation crosswalk</b>                                            | <b>207</b> |

## 1 Historical note and scope

The probability distribution now attached to the name of Fabius is older than Fabius’s two-page note. It occurs in the 1935 work of Jessen and Wintner on infinite convolutions of independent random variables [1], was rediscovered several times, and entered a different mathematical culture through the compactly supported “finite function” of V. A. Rvachev. Fabius used it in 1966 as a particularly economical probabilistic construction of a nowhere analytic  $C^\infty$  function [2]. Independently, and for entirely different reasons, the Rvachevs introduced the same object in 1971 as the canonical compactly supported solution of a functional–differential equation [3], and V. A. Rvachev later developed it into the theory of *atomic functions* used in approximation theory and numerical analysis [4, 5, 6]. Rvachev’s normalization emphasizes the even compactly supported bump, whereas the Fabius normalization emphasizes its cumulative distribution function. Arias de Reyna’s 1982 paper gave a systematic treatment of the compactly supported function, including its Fourier product, probability interpretation, partitions of unity, derivatives, and dyadic values; an English translation appeared in 2017 [7]. His later arithmetic paper organized the exact rational sequences and denominator questions [8]. Haugland’s note [9] gives a practical algorithm for evaluating  $F$  at dyadic rationals, closely related to the bit-recursion of Section 8.

The present article is not merely a paraphrase of those papers. It follows the proof dependency graph of the Lean development in the `ProveIt` repository [22]. That development first constructs the bounded distribution function, then derives the Rvachev and signed-global forms, and finally builds the arithmetic and asymptotic layers on top. This order has two advantages. It prevents identities valid for the signed extension from being silently applied to the bounded distribution, and it makes the exact dyadic algorithms independent of any numerical approximation.

The formalization also settles several questions that are only implicit, conjectural, or incorrectly normalized in the source material. Most importantly, the sharp endpoint asymptotic contains a nonconstant periodic function of an exact lower-Lambert phase. The periodic term is not a cosmetic improvement: every nonzero Fourier mode is explicit and nonzero, and an elementary formula without it fails at the error scale usually claimed. The development further proves a full inverse-Lambert expansion, establishes an exact bit-recursive evaluator for every dyadic rational, and supplies an absolutely and uniformly convergent Fourier–Legendre expansion of  $\text{up}$ .

Arias de Reyna’s arithmetic paper [8] enumerates the independent occurrences of this function in chronological order, and two of them deserve mention beside Jessen–Wintner, Fabius, and the Rvachevs. In 1981 G. Kh. Kirov and G. A. Totkov arrived at it in a study of the distribution of the zeros of the derivatives of the function they wrote  $\lambda(x)$  (*Godishnik Vissh. Uchebn. Zaved. Prilozhna Mat.* **17** (1981), no. 4, 157–165, in Bulgarian with English and Russian summaries); related dyadic derivative phenomena are represented in the formal Taylor and exact-evaluation APIs used below. In 1985 R. Schnabl gave a further independent construction, “Über eine  $C^\infty$ -Funktion,” in *Zahlentheoretische Analysis*, Lecture Notes in Mathematics 1114, Springer, Berlin (1985), 134–142. Neither paper is used in the proofs below, which is why their data are given here rather than in the bibliography; they are recorded because a reader tracing the function through the literature will meet both names.

Our convention is fixed throughout:

- $F$  is the bounded cumulative distribution function, equal to 0 on  $(-\infty, 0]$  and to 1 on  $[1, \infty)$ ;
- $\text{up}$  is the even compactly supported bump on  $[-1, 1]$ ;
- $F_{\text{gl}}$  is the signed global extension satisfying  $F'_{\text{gl}}(x) = 2F_{\text{gl}}(2x)$  on all of  $\mathbb{R}$ .

All logarithms are natural logarithms. We write  $L = \log 2$  in the asymptotic sections,  $\text{wt}_2(n)$  for the sum of the binary digits of  $n$ , and  $\text{sinc}_{\text{rad}} z = \sin z/z$  with the removable value 1 at the origin.

## 1.1 A guide to the main results

The exposition follows the dependency order of the formal development. The opening chapters separate the bounded function  $F$ , the bump up, and the signed extension  $F_{\text{gl}}$ ; construct the canonical function; and record the exact uniqueness and shape theorems. The probability chapter then gives the weighted-uniform representation, the sharp square-summable activation calculus and its finite Taylor jet, and three finite approximations: atomic convolution measures, half-weighted step functions, and centered uniform splines. It also isolates the arbitrary-weight distribution laws and the support-free and compact inverse-CDF transport theorems before specializing them back to the Fabius quantile.

The Fourier and global-extension chapters collect the proved product, inversion, Poisson, derivative, normalized repeated-integration, exact dyadic-cell distribution, bound, and flatness APIs. The repeated-integration layer is first proved for arbitrary continuous Banach-valued inputs and polynomial weights, then specialized to the signed and bounded Fabius functions. These chapters lead to the exact analytic locus of  $F$ , the proved nonanalyticity regions for up and  $F_{\text{gl}}$ , the totalized inverse on  $[0, 1]$ , and the formal non-elementarity results. The moment and arithmetic chapters then derive the rational recurrences, reciprocal-power identities, path and composition formulas, parity and two-adic statements, exact dyadic evaluation, binary reduction, and the specialized half-base  $q$ -binomial formulas.

The later analytic chapters summarize the formal endpoint machinery: the exact negative-Laplace product and logarithmic decomposition, the periodic Fourier correction, the lower-Lambert coordinate, the proved saddle asymptotics and correction coefficients, and the rigorous obstructions to deleting the periodic term. The Fourier–Legendre, summation, and computability chapters finish the proved story. The final module guide names the Lean declarations supporting each retained theorem.

A statement appears here only when its complete mathematical content has an exact proved Lean counterpart. Material removed from earlier drafts because that test is not yet met is preserved in the research-frontier volume. Thus the division between this document and that volume is a proof-status boundary, not a judgment about mathematical plausibility or importance.

---

**One formal object, three global functions.** The bounded  $F$  is the cumulative distribution function used for probabilities and dyadic values. The compactly supported up is obtained by folding  $F$ , and its Fourier transform has the proved sinc and cosine product descriptions. The signed  $F_{\text{gl}}$  is the global Thue–Morse continuation and satisfies the pure dilation equation on all of  $\mathbb{R}$ . Domains and normalizations are stated explicitly whenever a theorem passes from one presentation to another.

---

## 2 Three functions that must not be conflated

The literature often uses the same letter  $F$  for three related, but globally different, functions. That economy of notation is harmless in a local calculation and dangerous in a long proof. We therefore begin by separating them.

### 2.1 The bounded Fabius distribution function

**Definition 2.1** (Bounded Fabius function). A function  $F : \mathbb{R} \rightarrow [0, 1]$  is called a *bounded Fabius function* if

$$F(x) = 0 \quad (x \leq 0), \quad (1)$$

$$F(x) = 1 \quad (x \geq 1), \quad (2)$$

$$F(1-x) = 1 - F(x) \quad (0 \leq x \leq 1), \quad (3)$$

$$F'(x) = 2F(2x) \quad (0 \leq x \leq \tfrac{1}{2}), \quad (4)$$

and  $F$  is infinitely differentiable on  $\mathbb{R}$ .

The derivative equation on the right half follows from symmetry:

$$F'(x) = 2F(2 - 2x), \quad \frac{1}{2} \leq x \leq 1. \quad (5)$$

In particular  $F'(x) > 0$  for  $0 < x < 1$ , once positivity in the interior is known. Thus  $F$  is a genuine cumulative distribution function, not merely a smooth interpolation between 0 and 1.

Because  $F$  is constant on both exterior half-lines and is  $C^\infty$ , every derivative of positive order vanishes at both endpoints:

$$F^{(m)}(0) = F^{(m)}(1) = 0 \quad (m \geq 1). \quad (6)$$

This flatness is the source of both its extraordinary endpoint decay and its failure to be analytic there.

## 2.2 Rvachev's compactly supported bump

**Definition 2.2** (Rvachev up-function). Given  $F$ , define

$$\text{up}(x) = \begin{cases} F(x + 1), & x \leq 0, \\ F(1 - x), & x > 0. \end{cases} \quad (7)$$

Equivalently,

$$\text{up}(x) = F(1 - |x|). \quad (8)$$

Here the right side is interpreted using the globally bounded function  $F$ .

It follows immediately that  $\text{up}$  is even, nonnegative, supported on  $[-1, 1]$ , and  $\text{up}(0) = 1$ . On the unit interval the two functions are simply translates:

$$F(x) = \text{up}(x - 1), \quad 0 \leq x \leq 1. \quad (9)$$

The graph of  $F$  is an infinitely flat sigmoid; the graph of  $\text{up}$  is its symmetric fold.

The primary exposition omits sampled plots: their decimal coordinates are not themselves Lean declarations. Exact point evaluation is developed below through rational dyadic formulas and executable evaluators.

## 2.3 The signed global extension

Let  $\text{wt}_2(n)$  denote the sum of the binary digits of  $n$ , and put

$$\varepsilon_n = (-1)^{\text{wt}_2(n)}. \quad (10)$$

The sequence  $(\varepsilon_n)$  is the  $\{\pm 1\}$  form of the Thue–Morse sequence [14].

**Definition 2.3** (Signed extension). Define

$$F_{\text{gl}}(x) = \sum_{n \geq 0} \varepsilon_n \text{up}(x - 2n - 1). \quad (11)$$

At a fixed  $x$  only finitely many summands can be nonzero, so this is a locally finite sum. On every block  $[2b, 2b + 2]$  only the  $b$ -th translate can be nonzero (and at the endpoints it too vanishes):

$$F_{\text{gl}}(x) = \varepsilon_b \text{up}(x - 2b - 1), \quad 2b \leq x \leq 2b + 2. \quad (12)$$

It vanishes for  $x \leq 0$ , and it agrees with the bounded  $F$  for every  $x \leq 1$  (both functions vanish on  $(-\infty, 0]$ ). It does *not* remain in  $[0, 1]$  to the right of that interval. This is the global object that satisfies the cleanest differential equation:

$$F'_{\text{gl}}(x) = 2F_{\text{gl}}(2x) \quad (x \in \mathbb{R}). \quad (13)$$

Arias de Reyna's arithmetic paper uses this extension. Statements about its values outside  $[0, 1]$  must therefore not be transferred blindly to the bounded cumulative distribution.



---

**Notation used in this article.**  $F$  always means the bounded distribution function;  $\text{up}$  always means the compactly supported even bump; and  $F_{\text{gl}}$  always means the signed global extension. We have  $F = F_{\text{gl}}$  throughout  $(-\infty, 1]$  (hence on  $[0, 1]$ ), but globally they are different functions.

---

## 2.4 Exact shape: positivity, bijectivity, and unimodality

The three defining equations (1)–(4) already determine the qualitative picture, but the statements that are actually used later are the sharp ones: not  $F \geq 0$  but  $F(x) > 0$  exactly for  $x > 0$ ; not  $\text{up} \leq 1$  but  $\text{up}(x) = 1$  exactly at the origin. We record them here, in the form of equivalences, because the strict versions are what make the convexity and Lipschitz theorems below sharp.

Everything rests on a single formula that merges (4) and (5) into one identity valid on the whole line.

**Lemma 2.4** (Unified derivative formula). *For every real  $x$ ,*

$$F'(x) = 2 \text{up}(2x - 1). \quad (14)$$

*Proof.* By (8),  $\text{up}(2x - 1) = F(1 - |2x - 1|)$ . For  $0 \leq x \leq \frac{1}{2}$  one has  $|2x - 1| = 1 - 2x$ , so  $\text{up}(2x - 1) = F(2x)$  and (14) is (4). For  $\frac{1}{2} \leq x \leq 1$  one has  $|2x - 1| = 2x - 1$ , so  $\text{up}(2x - 1) = F(2 - 2x)$  and (14) is (5). For  $x < 0$  or  $x > 1$  one has  $|2x - 1| > 1$ , hence  $1 - |2x - 1| < 0$  and  $\text{up}(2x - 1) = 0$  by (1), while  $F'(x) = 0$  because  $F$  is constant on each exterior half-line by (1) and (2).  $\square$

Since  $\text{up} \geq 0$ , formula (14) shows at once that  $F$  is nondecreasing on all of  $\mathbb{R}$ . It is also convenient to note that the reflection symmetry (3), stated on the unit interval, in fact holds globally:

$$F(1 - x) = 1 - F(x) \quad (x \in \mathbb{R}). \quad (15)$$

Indeed, for  $x < 0$  both sides equal 1 by (1) and (2), and for  $x > 1$  both sides equal 0. The evenness of  $\text{up}$  in (14) also gives the global first-jet reflection law

$$F'(1 - x) = F'(x) \quad (x \in \mathbb{R}), \quad (16)$$

formalized as `deriv_fabiusReal_one_sub` in `Differential.lean`.

**Theorem 2.5** (Positivity, strict monotonicity, bijectivity). *The bounded Fabius function satisfies the following.*

1.  $F(x) > 0$  if and only if  $x > 0$ .
2.  $F(x) < 1$  if and only if  $x < 1$ .
3.  $F'(x) > 0$  for  $0 < x < 1$ , and  $F$  is strictly increasing on  $[0, 1]$ .
4.  $F$  maps  $[0, 1]$  bijectively onto  $[0, 1]$ .

*Proof.* The heart of the matter is the following doubling step: if  $0 \leq x \leq \frac{1}{2}$  and  $F(x) = 0$ , then  $F(2x) = 0$ . To see it, observe that  $F \geq 0$  everywhere, so a zero of  $F$  is a global minimum of a function differentiable on all of  $\mathbb{R}$ ; Fermat's rule gives  $F'(x) = 0$ , and (4) turns this into  $2F(2x) = 0$ .

Suppose now that  $F(x) = 0$  for some  $x > 0$ . Since  $2^n x \rightarrow \infty$ , there is a smallest integer  $n \geq 0$  with  $2^n x \geq \frac{1}{2}$ . For  $0 \leq m < n$  we have  $2^m x < \frac{1}{2}$ , so the doubling step applies at  $2^m x$  and propagates the vanishing: from  $F(x) = 0$  we obtain  $F(2^m x) = 0$  for every  $m \leq n$ , and in particular  $F(2^n x) = 0$ . But  $F$  is nondecreasing and  $2^n x \geq \frac{1}{2}$ , so  $F(2^n x) \geq F(\frac{1}{2}) = \frac{1}{2}$ , where the midpoint value comes from (3) at  $x = \frac{1}{2}$ . This contradiction proves that  $F$  has no zero on the positive half-line; combined with (1) it proves (i).

Assertion (ii) follows from (i) and (15):  $F(x) < 1$  if and only if  $F(1 - x) > 0$ , that is, if and only if  $1 - x > 0$ .

For (iii), let  $0 < x < 1$ . Then  $2x - 1 \in (-1, 1)$ , so  $1 - |2x - 1| \in (0, 1]$ , and (i) gives  $\text{up}(2x - 1) = F(1 - |2x - 1|) > 0$ . Now (14) yields  $F'(x) > 0$ . A continuous function on  $[0, 1]$  whose derivative is positive throughout the interior is strictly increasing on  $[0, 1]$ .

For (iv),  $F$  is continuous with  $F(0) = 0$  and  $F(1) = 1$ , so the intermediate value theorem makes  $F([0, 1])$  contain  $[0, 1]$ ; since  $F$  takes values in  $[0, 1]$ , the image is exactly  $[0, 1]$ . Strict monotonicity from (iii) gives injectivity.  $\square$

The fold (8) now transfers the whole picture to Rvachev’s bump, and it does so in strict form.

**Theorem 2.6** (Strict unimodality of up). *Rvachev’s function satisfies the following.*

1. *up is strictly increasing on  $[-1, 0]$  and strictly decreasing on  $[0, 1]$ .*
2.  *$\text{up}(x) = 1$  if and only if  $x = 0$ ; the maximum value 1 is attained at the origin and nowhere else.*
3.  *$\text{up}(x) = 0$  if and only if  $x \notin (-1, 1)$ . Thus up is nonzero exactly on  $(-1, 1)$ , and  $\text{supp up} = [-1, 1]$ .*
4.  *$\text{up}'(x) > 0$  for  $-1 < x < 0$  and  $\text{up}'(x) < 0$  for  $0 < x < 1$ .*

*Proof.* By (8),  $\text{up}(x) = F(1 - x)$  for  $x \geq 0$ . As  $x$  increases through  $[0, 1]$  the argument  $1 - x$  decreases through  $[0, 1]$ , on which  $F$  is strictly increasing by theorem 2.5(iii); hence up is strictly decreasing on  $[0, 1]$ , and evenness gives (i) on  $[-1, 0]$ .

For (ii),  $\text{up}(x) = F(1 - |x|)$  and theorem 2.5(ii) say that  $\text{up}(x) < 1$  unless  $1 - |x| \geq 1$ , that is, unless  $x = 0$ ; and  $\text{up}(0) = F(1) = 1$ . For (iii),  $\text{up}(x) > 0$  if and only if  $1 - |x| > 0$  by theorem 2.5(i), and  $\text{up} \geq 0$ , so the zero set is exactly the complement of  $(-1, 1)$ ; the closed support is the closure  $[-1, 1]$ .

For (iv), the identity  $\text{up}(x) = F(1 - x)$  holds on the whole open half-line  $x > 0$ , hence may be differentiated there:  $\text{up}'(x) = -F'(1 - x)$ , which is negative for  $0 < x < 1$  by theorem 2.5(iii). Evenness gives  $\text{up}'(x) = -\text{up}'(-x)$ , so  $\text{up}' > 0$  on  $(-1, 0)$ .  $\square$

Parts (i) and (iii) are the classical description of the bump [6, 7]; the strict forms, and the equivalences in theorem 2.5, are exactly what the formal development records.

## 2.5 Convexity and the maximal slope

The derivative formula transfers the monotonicity of up to the derivative of  $F$ .

**Theorem 2.7** (Convexity, concavity, and the maximal slope). *The derivative  $F'$  is nondecreasing on  $(-\infty, \frac{1}{2}]$  and nonincreasing on  $[\frac{1}{2}, \infty)$ . It is strictly increasing on  $[0, \frac{1}{2}]$  and strictly decreasing on  $[\frac{1}{2}, 1]$ . Consequently  $F$  is convex on  $(-\infty, \frac{1}{2}]$ , concave on  $[\frac{1}{2}, \infty)$ , strictly convex on  $[0, \frac{1}{2}]$ , and strictly concave on  $[\frac{1}{2}, 1]$ . Moreover*

$$F'(\tfrac{1}{2}) = 2 = \max_{x \in \mathbb{R}} F'(x), \quad (17)$$

*and equality  $F'(x) = 2$  holds exactly at  $x = \frac{1}{2}$ .*

*Formal proof.* The derivative monotonicity statements are `monotoneOn_deriv_fabiusReal`, `antitoneOn_deriv_fabiusReal`, `strictMonoOn_deriv_fabiusReal_Icc`, and `strictAntitoneOn_deriv_fabiusReal_Icc`. The four convexity statements are `convexOn_fabiusReal_Iic_half`, `concaveOn_fabiusReal_Ici_half`, `strictConvexOn_fabiusReal_firstHalf`, and `strictConcaveOn_fabiusReal_secondHalf`. Finally `deriv_fabiusReal_half`, `deriv_fabiusReal_eq_two_iff`, and `isGreatest_deriv_fabiusReal` prove the exact maximal-slope statement. All are in `Convexity.lean`.  $\square$

The same module records the complete pointwise curvature profile. For every  $x \in \mathbb{R}$ ,

$$F''(x) = 8(\text{up}(4x - 1) - \text{up}(4x - 3)), \quad (18)$$

and therefore

$$\begin{aligned} F''(x) > 0 &\iff 0 < x < \frac{1}{2}, \\ F''(x) < 0 &\iff \frac{1}{2} < x < 1, \\ F''(x) = 0 &\iff x \notin (0, 1) \text{ or } x = \frac{1}{2}. \end{aligned} \quad (19)$$

The derivative-level statement and its pointwise corollary are `deriv_fabiusReal_hasDerivAt` and `deriv_deriv_fabiusReal`; the three equivalences are `deriv_deriv_fabiusReal_pos_iff`, `deriv_deriv_fabiusReal_neg_iff`, and `deriv_deriv_fabiusReal_eq_zero_iff`, with `deriv_deriv_fabiusReal_half` naming the midpoint specialization.

## 2.6 The optimal Lipschitz constant

The bound  $F' \leq 2$  of (17) is a Lipschitz estimate, and because the bound is attained the estimate cannot be improved.

**Theorem 2.8** (Two is the least Lipschitz constant). *Both  $F$  and  $\text{up}$  are 2-Lipschitz on  $\mathbb{R}$ :*

$$|F(x) - F(y)| \leq 2|x - y|, \quad |\text{up}(x) - \text{up}(y)| \leq 2|x - y| \quad (x, y \in \mathbb{R}). \quad (20)$$

Moreover 2 is the least admissible constant for each: if  $K \geq 0$  is such that  $|F(x) - F(y)| \leq K|x - y|$  for all  $x, y$ , or such that  $|\text{up}(x) - \text{up}(y)| \leq K|x - y|$  for all  $x, y$ , then  $K \geq 2$ .

*Proof.* By (14) and  $0 \leq \text{up} \leq 1$  we have  $0 \leq F' \leq 2$  on all of  $\mathbb{R}$ , so the mean value theorem gives the first estimate in (20). The map  $x \mapsto 1 - |x|$  is 1-Lipschitz, so the second estimate follows from the first through the fold (8).

For optimality, let  $K$  be admissible for  $F$ . Dividing  $|F(x) - F(\frac{1}{2})| \leq K|x - \frac{1}{2}|$  by  $|x - \frac{1}{2}|$  and letting  $x \rightarrow \frac{1}{2}$  gives  $K \geq |F'(\frac{1}{2})| = 2$  by (17). For  $\text{up}$  the same limit at the same point works: the proof of theorem 2.6(iv) gives  $\text{up}'(\frac{1}{2}) = -F'(\frac{1}{2}) = -2$ , so  $K \geq 2$ .  $\square$

The two extremal points are the same point in the two pictures:  $\frac{1}{2}$  is the centre of the sigmoid and the midpoint of the decreasing half of the bump, and the fold sends one to the other.

**Corollary 2.9** (Linear majorants at the endpoints). *For all admissible arguments,*

$$F(x) \leq 2x \quad (x \geq 0), \quad (21)$$

$$1 - F(x) \leq 2(1 - x) \quad (x \leq 1), \quad (22)$$

$$\text{up}(x) \leq 2(1 - |x|) \quad (|x| \leq 1). \quad (23)$$

*Proof.* Applying (20) with  $y = 0$  and using  $F(0) = 0$  gives (21). Replacing  $x$  by  $1 - x$  there and using (15) gives (22). Substituting  $1 - |x| \geq 0$  into (21) and using (8) gives (23).  $\square$

The content of (21) lies on  $[0, \frac{1}{2}]$ ; for  $x \geq \frac{1}{2}$  it is weaker than  $F \leq 1$ . Its value is that it holds with no hypothesis at all on the size of  $x$ , so it can be used as a linear majorant in any estimate near the left endpoint, and (23) says that the bump vanishes at least linearly at both ends of its support. These majorants and the optimality of the constant 2 are contributions of the formal development; the plain 2-Lipschitz bound is used implicitly throughout [7, 6].

### 3 Existence and uniqueness by a contraction

The most economical construction of  $F$  is a Banach fixed-point argument. It has an additional advantage: the contraction estimate later gives a clean uniqueness theorem, without presupposing Fourier analysis or probability.

#### 3.1 The admissible function space

Let  $I = [0, 1]$  and let  $C(I)$  carry the supremum norm. Define  $\mathcal{A} \subset C(I)$  by

$$\mathcal{A} = \{f \in C(I) : 0 \leq f \leq 1, f(1-x) = 1 - f(x)\}. \quad (24)$$

The set  $\mathcal{A}$  is closed in the Banach space  $C(I)$ , hence complete. It is nonempty; for example  $f(x) = x$  belongs to it.

Extend  $f \in \mathcal{A}$  to a continuous function  $\tilde{f}$  on  $\mathbb{R}$  by clamping the argument to  $I$ :

$$\tilde{f}(t) = \begin{cases} f(0), & t \leq 0, \\ f(t), & 0 \leq t \leq 1, \\ f(1), & t \geq 1. \end{cases} \quad (25)$$

Symmetry gives

$$\int_0^1 f(t) dt = \frac{1}{2}. \quad (26)$$

Indeed, replacing  $t$  by  $1-t$  and adding the two integrals gives  $2 \int_0^1 f = 1$ .

#### 3.2 The integral transform

For  $f \in \mathcal{A}$ , define  $Tf$  by

$$(Tf)(x) = \begin{cases} \int_0^{2x} \tilde{f}(t) dt, & 0 \leq x \leq \frac{1}{2}, \\ 1 - \int_0^{2-2x} \tilde{f}(t) dt, & \frac{1}{2} < x \leq 1. \end{cases} \quad (27)$$

The two definitions agree at  $x = 1/2$  by (26). Positivity and the upper bound  $f \leq 1$  show that  $0 \leq Tf \leq 1$ , and a direct substitution shows  $(Tf)(1-x) = 1 - (Tf)(x)$ . Hence  $T : \mathcal{A} \rightarrow \mathcal{A}$ .

The central estimate is stronger than the naive integral bound.

**Lemma 3.1** (Contraction estimate). *For  $f, g \in \mathcal{A}$ ,*

$$\|Tf - Tg\|_\infty \leq \frac{1}{2} \|f - g\|_\infty. \quad (28)$$

*Proof.* Write  $h = f - g$ . Since both  $f$  and  $g$  have integral  $1/2$ ,  $\int_0^1 h = 0$ . For  $0 \leq y \leq 1$  we may therefore estimate the primitive in two ways:

$$\begin{aligned} \left| \int_0^y h(t) dt \right| &\leq y \|h\|_\infty, \\ \left| \int_0^y h(t) dt \right| &= \left| \int_y^1 h(t) dt \right| \leq (1-y) \|h\|_\infty. \end{aligned}$$

Consequently

$$\left| \int_0^y h(t) dt \right| \leq \min(y, 1-y) \|h\|_\infty \leq \frac{1}{2} \|h\|_\infty.$$

Every value of  $Tf - Tg$  is one of these primitives, possibly with a minus sign. Taking the supremum proves (28).  $\square$

**Theorem 3.2** (Existence of the bounded Fabius function). *The transform  $T$  has a unique fixed point  $f_* \in \mathcal{A}$ . Extending  $f_*$  by 0 on  $(-\infty, 0]$  and by 1 on  $[1, \infty)$  produces the bounded Fabius function  $F$ .*

*Proof.* The space  $\mathcal{A}$  is complete and  $T$  is a strict contraction with constant  $1/2$ , so Banach’s fixed-point theorem gives a unique fixed point. On  $[0, 1/2]$  its fixed-point equation is

$$F(x) = \int_0^{2x} F(t) dt. \quad (29)$$

Differentiating gives  $F'(x) = 2F(2x)$ . The second half of (27) supplies symmetry and the matching right-hand formula. The endpoint values are immediate.  $\square$

### 3.3 The differential equation for up and smoothness

Folding (4) and (5) yields the refinement-type differential equation

$$\text{up}'(x) = 2(\text{up}(2x + 1) - \text{up}(2x - 1)) \quad (x \in \mathbb{R}). \quad (30)$$

At  $x = 0$  the two one-sided derivatives glue because  $\text{up}(1) = \text{up}(-1) = 0$ . Once  $\text{up}$  is continuously differentiable, (30) bootstraps regularity: if  $\text{up} \in C^r$ , then the right side is  $C^r$ , so  $\text{up} \in C^{r+1}$ . Induction gives  $\text{up} \in C^\infty$  and then  $F \in C^\infty$ .

**Proposition 3.3** (Basic shape). *The functions  $F$  and  $\text{up}$  have the following properties.*

1.  $F$  is strictly increasing on  $(0, 1)$  and satisfies  $0 < F(x) < 1$  there.
2.  $\text{up}$  is even, strictly positive on  $(-1, 1)$ , and zero outside  $[-1, 1]$ .
3.  $\int_{-1}^1 \text{up}(x) dx = 1$ .
4. Every derivative of  $\text{up}$  vanishes at  $x = \pm 1$ .

*Proof.* For  $0 < x \leq 1/2$ , (4) gives  $F'(x) = 2F(2x)$ . Positivity follows first near  $1/2$  from  $F(1/2) = 1/2$ , and then propagates toward zero by repeatedly doubling the argument; symmetry handles the right half. This proves strict monotonicity and, by (8), positivity of  $\text{up}$  in the interior of its support. Finally,

$$\int_{-1}^1 \text{up}(x) dx = 2 \int_0^1 F(t) dt = 1,$$

where the middle identity uses the fold and the last uses symmetry. Flatness at the support endpoints follows from smoothness and the fact that  $\text{up}$  is identically zero just outside the support.  $\square$

### 3.4 The original Rvachev characterization

The formal predicate `IsOriginalFabius` packages the source characterization exactly. For a function  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  and a scale  $k$ , it requires  $C^\infty$ -smoothness, topological support  $[-1, 1]$ , strict positivity on  $(-1, 1)$ , the normalization  $\varphi(0) = 1$ , positivity of  $k$ , and

$$\varphi'(x) = k(\varphi(2x + 1) - \varphi(2x - 1)) \quad (x \in \mathbb{R}). \quad (31)$$

The positivity of the scale is redundant: the smart constructor `IsOriginalFabius.mk_of_derivative_law` derives it from the other hypotheses.

**Theorem 3.4** (Original characterization). *If `IsOriginalFabius`  $\varphi$   $k$ , then*

$$k = 2, \quad \varphi = \text{up}. \quad (32)$$

*Conversely, the canonical  $\text{up}$  satisfies this predicate at scale 2; in particular there is exactly one original solution.*

*Formal proof.* The scale is fixed by `originalFabius_scale_eq_two`; the canonical witness is `canonical_isOriginalFabius`. Uniqueness and the displayed characterization are `originalFabius_eq_canonical`, `isOriginalFabius_iff_eq_canonical`, and `existsUnique_originalFabius` in `OriginalUniqueness.lean`.  $\square$

The fold loses only the unused right tail of a bounded candidate. Precisely,

$$\text{up}_G = \text{up}_H \iff G(t) = H(t) \text{ for every } t \leq 1. \quad (33)$$

Thus a bounded candidate is canonical exactly when its fold satisfies the original characterization at scale 2 and its right tail is constantly 1. These are the exact declarations `rvachevUp_eq_iff_eqOn_Iic_one`, `isFabius_iff_isOriginalFabius_rvachevUp_and_rightTail`, and `isOriginalFabius_iff_existsUnique_isFabius` in `OriginalUniqueness.lean`.

## 4 The probability law behind the functions

The fixed-point construction is analytic. A probabilistic construction explains the same self-similarity almost without calculation and makes monotonicity and positivity transparent.

### 4.1 An infinite weighted sum of uniforms

Let  $U_0, U_1, \dots$  be independent random variables, each uniformly distributed on  $[0, 1]$ , and define

$$X = \sum_{n=0}^{\infty} \frac{U_n}{2^{n+1}}. \quad (34)$$

The series converges absolutely for every outcome and satisfies  $0 \leq X \leq 1$ . Removing the first coordinate gives an independent copy  $X'$  of  $X$ , and

$$X = \frac{U_0 + X'}{2}. \quad (35)$$

Reflecting every coordinate,  $U_n \mapsto 1 - U_n$ , preserves the product measure and sends  $X$  to  $1 - X$ .

Let  $H(x) = \mathbb{P}(X \leq x)$  be the cumulative distribution function. Conditioning on  $U_0$  in (35) gives

$$H(y) = \int_0^1 H(2y - u) \, du. \quad (36)$$

For  $0 \leq y \leq 1/2$ , the integrand vanishes when  $u > 2y$ , so

$$H(y) = \int_0^{2y} H(t) \, dt. \quad (37)$$

Reflection gives  $H(1 - y) = 1 - H(y)$ . Thus the restriction of  $H$  to  $[0, 1]$  is a fixed point of the contraction  $T$  from [section 3](#).

**Theorem 4.1** (Global probabilistic representation). *For every  $x, y \in \mathbb{R}$ ,*

$$F(x) = \mathbb{P}\left(\sum_{n=0}^{\infty} \frac{U_n}{2^{n+1}} \leq x\right), \quad (38)$$

$$\text{up}(y) = \mathbb{P}\left(\sum_{n=0}^{\infty} \frac{U_n}{2^{n+1}} \leq 1 - |y|\right). \quad (39)$$

*The second identity includes both tails: if  $|y| \geq 1$ , both sides vanish.*

*Formal proof.* The product-space construction proves continuity and measurability of the uniformly convergent coordinate sum, identifies its pushforward distribution, establishes the head–tail self-similarity and the coordinate-reflection law, and invokes uniqueness of the bounded Fabius fixed point. The exact canonical specializations are `fabiusReal_eq_weightedSum_probability` and `rvachevUp_eq_weightedSum_probability_global` in `ProbabilityRepresentation.lean`. The same module makes the transition to the generic law literal: `ProbabilityRepresentation.weightedSumDistribution_eq_weightedUniformDistribution` identifies the dyadic `ProbabilityRepresentation.weightedSumDistribution` with the generic weighted-uniform law for  $w_n = 1/2/2^n = 2^{-n-1}$ . The additional bridges `ProbabilityRepresentation.weightedCoordinateSum_eq_geometricUniformSeries_one_half` and `ProbabilityRepresentation.weightedSumDistribution_eq_geometricUniformDistribution_one_half` identify the dyadic series and law with the named geometric specialization at  $q = 1/2$ . Its exact topological support is recorded by `ProbabilityRepresentation.weightedSumDistribution_support_eq_Icc`. The same `ProbabilityRepresentation.lean` ownership now includes the almost-everywhere carried support and restriction identity `ProbabilityRepresentation.ae_weightedSumDistribution_mem_Icc` and `ProbabilityRepresentation.weightedSumDistribution_restrict_Icc`.  $\square$

## 4.2 Generic weighted laws and support-free quantile transport

The probability mechanism does not depend on the dyadic weights. Let  $\Omega = [0, 1]^{\mathbb{N}_0}$ , let  $\mathbf{P}$  be product Lebesgue measure, and let  $\tau\omega = (\omega_1, \omega_2, \dots)$ . For a real normed vector space  $E$  equipped with a measurable structure and weights  $w : \mathbb{N}_0 \rightarrow E$ , put

$$S_w(\omega) := \sum_{n=0}^{\infty} \omega_n w_n, \quad \mu_w := (S_w)_* \mathbf{P}, \quad w_n^+ := w_{n+1}. \quad (40)$$

The series  $S_w$  and the pushforward  $\mu_w$  are totalized Lean definitions: they exist syntactically even without summability or measurability. The law statements below add precisely the hypotheses that make those definitions analytically meaningful.

**Theorem 4.2** (Head–tail product law and arbitrary weighted distributions). *For every measurable map  $g : \Omega \rightarrow A$  into an arbitrary measurable space,*

$$(\omega \mapsto (\omega_0, g(\tau\omega)))_* \mathbf{P} = \lambda_{[0,1]} \otimes g_* \mathbf{P}, \quad (41)$$

where  $\lambda_{[0,1]}$  is Lebesgue measure restricted to the unit interval.

Now let  $E$  be a real Banach space with its Borel measurable structure. If  $\sum_n \|w_n\| < \infty$ , then  $\mu_w$  is a probability measure. The weighted joint law is more general and independent of any parent sequence: for every separately chosen  $v : \mathbb{N}_0 \rightarrow E$  with  $\sum_n \|v_n\| < \infty$ ,

$$(\omega \mapsto (\omega_0, S_v(\tau\omega)))_* \mathbf{P} = \lambda_{[0,1]} \otimes \mu_v. \quad (42)$$

Under full summability, with  $W = \sum_n w_n$  and  $A_w(u, z) = u w_0 + z$ ,

$$\mu_w = (A_w)_* (\lambda_{[0,1]} \otimes \mu_{w^+}), \quad (43)$$

$$(x \mapsto W - x)_* \mu_w = \mu_w. \quad (44)$$

Thus every absolutely summable vector-weight law is centrally symmetric about  $W/2$ . If  $E = \mathbb{R}$  and at least one coefficient  $w_n$  is nonzero, then  $\mu_w$  is absolutely continuous with respect to Lebesgue measure and every singleton is  $\mu_w$ -null. No sign condition is imposed on the weights, and the nonzero coefficient may occur at an arbitrary index rather than at the head. More generally, its topological support is the exact range of the continuous series map:

$$\text{supp}(\mu_w) = S_w(\Omega). \quad (45)$$



If  $E = \mathbb{R}$ , with no sign restriction, put

$$L_w = \sum_n \min\{w_n, 0\}, \quad U_w = \sum_n \max\{w_n, 0\}.$$

Then cancellation and sign changes create no gaps:

$$S_w(\Omega) = \text{supp}(\mu_w) = [L_w, U_w]. \quad (46)$$

If every  $w_n \geq 0$ , this becomes the full interval  $[0, W]$ , including the degenerate case  $W = 0$ . Consequently

$$S_w(\Omega) = \text{supp}(\mu_w) = [0, W], \quad \mu_w([0, W]) = 1, \quad \mu_w(\mathbb{R} \setminus [0, W]) = 0. \quad (47)$$

Equation (41) is `ProbabilityRepresentation.uniformProduct_map_head_tail_comp`. Definition (40) uses the existing `ProbabilityRepresentation.weightedUniformSeries` and the new `ProbabilityRepresentation.weightedUniformDistribution`; its totalization boundary is part of the latter definition's statement. The probability and weighted head–tail results are `ProbabilityRepresentation.isProbabilityMeasure_weightedUniformDistribution` and `ProbabilityRepresentation.uniformProduct_map_head_tailWeightedUniformSeries`. The latter is the arbitrary- $v$  statement (42); the shifted choice  $v = w^+$  is made only in the subsequent split of  $\mu_w$ . Equations (43)–(44) are `ProbabilityRepresentation.weightedUniformDistribution_split` and `ProbabilityRepresentation.weightedUniformDistribution_reflection`; the two exact mass statements in (47) are `ProbabilityRepresentation.weightedUniformDistribution_Icc` and `ProbabilityRepresentation.weightedUniformDistribution_comp1_Icc`. Together these declarations give the probability, splitting, reflection, and carried-set layer of `WeightedUniformSeries.lean`.

The exact-support upgrade is isolated in `WeightedUniformSupport.lean`. Its ten public declarations are `ProbabilityRepresentation.isOpenPosMeasure_infinitePi`, `ProbabilityRepresentation.uniformProduct_isOpenPosMeasure`, `ProbabilityRepresentation.support_map_eq_closure_range_of_continuous`, `ProbabilityRepresentation.weightedUniformSeries_constCoordinates`, `ProbabilityRepresentation.weightedUniformDistribution_support_eq_range`, `ProbabilityRepresentation.range_weightedUniformSeries_eq_Icc_min_max`, `ProbabilityRepresentation.weightedUniformDistribution_support_eq_Icc_min_max`, `ProbabilityRepresentation.range_weightedUniformSeries_eq_Icc`, `ProbabilityRepresentation.weightedUniformDistribution_support_eq_Icc`, and `ProbabilityRepresentation.weightedUniformDistribution_support_eq_unitInterval`. The first three establish open positivity for arbitrary products and identify the support of a continuous pushforward with the closure of its range. Compactness of the coordinate cube then removes the closure in (45). In the real case, coordinates selecting each negative or positive coefficient attain the two endpoints, and connectedness fills the interval in (46); constant coordinate paths give the especially transparent nonnegative specialization.

The same module proves a more structural layer before any choice of geometric weights.

**Theorem 4.3** (Naturality and operator-affine recursion). *Let  $G$  and  $H$  be arbitrary measurable spaces, let  $g : \Omega \rightarrow G$  be measurable, and let  $\Phi : [0, 1] \times G \rightarrow H$  be measurable. If*

$$f(\omega) = \Phi(\omega_0, g(\tau\omega)) \quad \text{for } \mathbf{P}\text{-almost every } \omega,$$

*then no separate measurability hypothesis on  $f$  is needed and*

$$f_*\mathbf{P} = \Phi_*(\lambda_{[0,1]} \otimes g_*\mathbf{P}). \quad (48)$$

*There is also the immediate pointwise-recursion specialization.*



Let  $E$  be a real Banach space, let  $F$  be a real normed space, and let  $L : E \rightarrow F$  be continuous linear. The target  $F$  need not be complete. For  $\sum_n \|w_n\| < \infty$ , writing  $(Lw)_n = L(w_n)$ , one has

$$L(S_w(\omega)) = S_{Lw}(\omega). \quad (49)$$

If both spaces carry their Borel measurable structures, then

$$L_*\mu_w = \mu_{Lw}. \quad (50)$$

For scalar multiplication the pointwise identity

$$S_{cw}(\omega) = cS_w(\omega) \quad (51)$$

holds in every real normed space, with no completeness or summability assumption; the law-level identity  $(x \mapsto cx)_*\mu_w = \mu_{cw}$  retains completeness, Borel, and norm-summability hypotheses so that the totalized measure maps are genuine pushforwards.

Finally, let  $A : E \rightarrow E$  be continuous linear and suppose  $w_{n+1} = A(w_n)$  for every  $n$ . Then

$$S_w(\omega) = \omega_0 w_0 + A(S_w(\tau\omega)). \quad (52)$$

If  $E$  carries its Borel measurable structure, then also

$$A_*\mu_w = \mu_{w^+}, \quad (53)$$

$$\mu_w = (B_{A,w})_*(\lambda_{[0,1]} \otimes \mu_w), \quad B_{A,w}(u, z) = uw_0 + A(z). \quad (54)$$

The scalar case  $w_{n+1} = cw_n$  is a corollary of the operator statement with  $A = c \text{ id}$ . Here  $c \in \mathbb{R}$  is arbitrary: positivity and contraction are not built into either theorem; norm summability of the displayed weights is the analytic hypothesis.

Equation (48) and its pointwise specialization are `ProbabilityRepresentation.uniformProduct_map_of_ae_head_tail_recursion` and `ProbabilityRepresentation.uniformProduct_map_of_head_tail_recursion`. Equations (49)–(51) are `ProbabilityRepresentation.weightedUniformSeries_map` and `ProbabilityRepresentation.weightedUniformSeries_smul`; their law-level counterparts are `ProbabilityRepresentation.weightedUniformDistribution_map` and `ProbabilityRepresentation.weightedUniformDistribution_smul`. Equation (53) is `ProbabilityRepresentation.weightedUniformDistribution_map_eq_shift`. The pointwise and law recursions (52) and (54) are `ProbabilityRepresentation.weightedUniformSeries_split_of_shift_eq_map` and `ProbabilityRepresentation.weightedUniformDistribution_split_of_shift_eq_map`; their scalar corollaries are `ProbabilityRepresentation.weightedUniformSeries_geometric_split` and `ProbabilityRepresentation.weightedUniformDistribution_geometric_split`. These are exactly the eleven public declarations in the generic naturality tranche.

The compatibility layer also proves the exact barycenter in every complete Borel real normed space. If  $\sum_n \|w_n\| < \infty$ , then

$$\int_E x \, d\mu_w(x) = \frac{1}{2} \cdot \sum_{n=0}^{\infty} w_n. \quad (55)$$

This is `ProbabilityRepresentation.integral_id_weightedUniformDistribution`. It uses no sign or order on the weights, no coordinatewise independence expansion, and no termwise integration hypothesis: bounded-series integrability supplies the integral and central reflection gives total weight minus the barycenter equal to the barycenter.

The word “geometric” in the final two declaration names denotes only the conditional recursion  $w_{n+1} = cw_n$ . The named specialization is supplied separately, rather than being hidden in these generic theorems.

The refactored weighted-law surface also exposes seven compatibility declarations: `ProbabilityRepresentation.weightedUniformSeries_smul_weights` in `WeightedUniformSeries.lean`, together with `ProbabilityRepresentation.uniformProduct_map_head_tail_function`, `ProbabilityRepresentation.weightedUniformDistribution_isProbabilityMeasure`, `ProbabilityRepresentation.weightedUniformDistribution_smul_weights`, `ProbabilityRepresentation.uniformProduct_map_head_tail_weightedUniformSeries`, `ProbabilityRepresentation.weightedUniformDistribution_unitInterval`, and `ProbabilityRepresentation.weightedUniformDistribution_compl_unitInterval` in `WeightedUniformDistribution.lean`. The two scalar compatibility identities use the same totalized pointwise equality as (51); at law level the compatibility theorem is oriented as  $\mu_{cw} = (x \mapsto cx)_* \mu_w$ . Those two unit-interval compatibility declarations are carried-set statements; the separate exact support equalities are supplied by `WeightedUniformSupport.lean`.

The same compatibility module contains a four-declaration real absolute-continuity tranche. For an arbitrary s-finite measure  $\nu$  on  $\mathbb{R}$  and  $a \neq 0$ , `ProbabilityRepresentation.uniformScaledAdd_absolutelyContinuous` proves that the image of  $\lambda_{[0,1]} \otimes \nu$  under  $(u, x) \mapsto ua + x$  is absolutely continuous with respect to Lebesgue measure. The weighted head-coordinate workhorse is `ProbabilityRepresentation.weightedUniformDistribution_absolutelyContinuous_of_head_ne_zero`. The sharp theorem `ProbabilityRepresentation.weightedUniformDistribution_absolutelyContinuous` assumes only norm summability and  $\exists n, w_n \neq 0$ , and `ProbabilityRepresentation.weightedUniformDistribution_nullSingletonClass` packages the resulting zero mass of every singleton as `Mathlib's NullSingletonClass`. Together with the six compatibility declarations listed above and the barycenter theorem, these are exactly the eleven public theorems, and there are no public definitions, in `WeightedUniformDistribution.lean`.

The same measure language gives a clean inverse-CDF theorem without differentiating an inverse.

**Theorem 4.4** (Support-free and compact inverse-CDF transport). *Let  $\mu$  be an arbitrary probability measure on  $\mathbb{R}$ , with no support restriction, and let  $Q : \mathbb{R} \rightarrow \mathbb{R}$  be almost-everywhere measurable for  $\lambda_{[0,1]}$ . If, for every  $x \in \mathbb{R}$ ,*

$$Q(y) \leq x \iff y \leq \text{cdf}_\mu(x) \text{ for } \lambda_{[0,1]} \text{-almost every } y, \quad (56)$$

then

$$Q_* \lambda_{[0,1]} = \mu. \quad (57)$$

Thus neither  $Q([0, 1]) \subseteq [0, 1]$  nor compact support of  $\mu$  belongs to the foundational theorem; null-set choices of quantile convention are immaterial.

There is also a pointwise compact specialization. Suppose now that  $\mu$  is carried almost everywhere by  $[0, 1]$ , and:

1.  $Q([0, 1]) \subseteq [0, 1]$ ;
2. for every  $x, y \in [0, 1]$ ,

$$Q(y) \leq x \iff y \leq \text{cdf}_\mu(x).$$

Together with the same source-a.e. measurability, these hypotheses again imply (57). Both versions permit atoms.

The support-free theorem is exactly `map_quantile_eq`; its compact specialization is `map_inverseCDF_volume_restrict ICC`, both in `QuantileTransport.lean`. In both cases only the source measure is restricted to  $[0, 1]$ . The first theorem neither controls nor needs the values of  $Q$  outside a source-full-measure set.

**Corollary 4.5** (Fabius quantile substitution). *Let  $I = \text{fabiusInv}$  be the Fabius inverse obtained by first clamping its argument to  $[0, 1]$ , and let  $\mu_X$  be the dyadic weighted-sum distribution in (38). Then*

$$I_*\lambda_{[0,1]} = \mu_X. \quad (58)$$

Consequently, for every strongly measurable map  $\Phi : \mathbb{R} \rightarrow E$  into a real Banach space,

$$\int_{[0,1]} \Phi(I(y)) \, dy = \int_{\mathbb{R}} \Phi(x) \, d\mu_X(x). \quad (59)$$

No separate integrability hypothesis is imposed. Lean’s Bochner integral is totalized to zero in the non-integrable case, and the measure-map theorem identifies the two totalized integrals there as well.

The measure identity and Banach-valued integral identity are respectively `map_fabiusInv_restrict_Icc_eq_weightedSumDistribution` and `integral_comp_fabiusInv_restrict_Icc_eq_weightedSumDistribution` in `QuantileTransport.lean`. Together with the support-free and compact statements above, they are exactly the four public declarations of that module. The clamped definition of  $I$  is global, but both displayed transport identities deliberately use uniform measure restricted to  $[0, 1]$ . They do not by themselves prove the two-function integration identity; the separate absolutely-continuous theorem below supplies that full-interval result. The transport declarations assert neither complex Mellin continuation nor fractional-power substitutions.

A separate subgraph-Fubini argument does provide the exact first-order weighted inverse identity, without differentiating either clock.

**Theorem 4.6** (Weighted Fabius inverse layer cake). *Let  $\phi$  be interval-integrable on  $[0, 1]$ , with real or complex values, and let  $\psi$  be interval-integrable there with values in a compatible Banach space. Then*

$$\int_0^1 \phi(u) \left( \int_0^{I(u)} \psi(x) \, dx \right) du = \int_0^1 \left( \int_{F(x)}^1 \phi(u) \, du \right) \psi(x) \, dx. \quad (60)$$

If  $\Psi(0) = 0$ ,  $\psi$  is continuous on  $[0, 1]$ , and  $\Psi'(x) = \psi(x)$  pointwise there, the inner primitive on the left may be replaced by  $\Psi(I(u))$ . If instead  $\Psi : \mathbb{R} \rightarrow \mathbb{R}$  is merely absolutely continuous on  $[0, 1]$ , the same replacement is valid with  $\psi = \Psi'$  almost everywhere, for every real or complex interval-integrable weight  $\phi$ ; no separate integrability hypothesis on the right-hand side is needed.

The arbitrary-absolutely-continuous Fabius form is `intervalIntegral_mul_comp_fabiusInv_of_absolutelyContinuousOnInterval`. The explicit-primitive and pointwise- $C^1$  forms are respectively `intervalIntegral_smul_intervalIntegral_fabiusInv` and `intervalIntegral_smul_comp_fabiusInv` in `InverseLayerCake.lean`.

The generic theorem does not require the primitive to vanish at the left endpoint.

**Theorem 4.7** (Absolutely-continuous inverse layer cake on a rectangle). *Let  $a \leq b$  and  $c \leq d$ . Suppose  $C, Q : \mathbb{R} \rightarrow \mathbb{R}$  map  $[a, b]$  into  $[c, d]$  and  $[c, d]$  into  $[a, b]$ , respectively, and*

$$C(x) < u \iff x < Q(u) \quad (x \in [a, b], u \in [c, d]).$$

For  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ , let  $\phi : [c, d] \rightarrow \mathbb{K}$  be integrable and let  $\Psi : \mathbb{R} \rightarrow \mathbb{R}$  be absolutely continuous on  $[a, b]$ . Then

$$\int_c^d \phi(u) (\Psi(Q(u)) - \Psi(a)) \, du = \int_a^b \Psi'(x) \left( \int_{C(x)}^d \phi(u) \, du \right) \, dx. \quad (61)$$

When  $\Psi(a) = 0$ , the subtraction on the left disappears. No measurability hypothesis is imposed on either clock, and equality is allowed in either endpoint pair. Values of the clocks outside the two displayed intervals play no role.

The order-theoretic engine is `galoisConnection_Icc_restrict_of_lt_iff_lt`: the strict order equivalence makes the two interval restrictions a Galois connection and hence monotone. A canonical clamped extension of the restricted inverse is therefore measurable. Consequently, the generic explicit-primitive and pointwise- $C^1$  theorems `intervalIntegral_smul_intervalIntegral_of_lt_iff_lt` and `intervalIntegral_smul_comp_of_lt_iff_lt` no longer assume clock measurability. The subtraction and normalized forms of (61) are `intervalIntegral_mul_comp_sub_of_lt_iff_lt_of_absolutelyContinuousOnInterval` and `intervalIntegral_mul_comp_of_lt_iff_lt_of_absolutelyContinuousOnInterval`. The reusable arbitrary-s-finite-measure engine is `integral_smul_setIntegral_subgraph` in `SubgraphFubini.lean`. Finally, `intervalIntegral_mul_fabiusInv_eq` is the real weighted-area specialization; `intervalIntegral_fabiusInv_eq_intervalIntegral_rvachevUp` and `intervalIntegral_fabiusInv_eq_one_half` identify the inverse area with the unit-interval up area and with  $1/2$ , respectively.

The inverse-order relation also supports a genuinely two-function absolutely-continuous calculus. Here the important reusable point is that measurability of the clocks is a conclusion of the order structure on the rectangle, not an extra hypothesis.

**Theorem 4.8** (Absolutely-continuous calculus for an inverse pair). *Let  $a \leq b$  and  $c \leq d$ . Suppose  $C, Q : \mathbb{R} \rightarrow \mathbb{R}$  map  $[a, b]$  into  $[c, d]$  and  $[c, d]$  into  $[a, b]$ , respectively, and*

$$C(x) < u \iff x < Q(u) \quad (x \in [a, b], u \in [c, d]). \quad (62)$$

*If  $A, B : \mathbb{R} \rightarrow \mathbb{R}$ , with  $A$  absolutely continuous on  $[a, b]$  and  $B$  absolutely continuous on  $[c, d]$ , then*

$$\int_a^b A'(x)B(C(x)) \, dx + \int_c^d A(Q(u))B'(u) \, du = A(b)B(d) - A(a)B(c). \quad (63)$$

*Moreover, every  $y \in [a, b]$  cuts out the smaller inverse rectangle  $[a, y] \times [c, C(y)]$ , and*

$$\int_a^y A'(x)B(C(x)) \, dx + \int_c^{C(y)} A(Q(u))B'(u) \, du = A(y)B(C(y)) - A(a)B(c). \quad (64)$$

*No measurability, continuity, or differentiability assumption is imposed on  $C$  or  $Q$ . The strict-order equivalence forces both restrictions to be monotone; canonical clamped extensions are then measurable. Values of the two clocks outside their displayed compact intervals play no role. Equality of either endpoint pair is allowed. The stated orientation is essential: reversed endpoints require swapping the corresponding interval and its order data first.*

The complete and variable-upper statements are, respectively, `intervalIntegral_deriv_mul_comp_add_comp_mul_deriv_of_lt_iff_lt` and `intervalIntegral_deriv_mul_comp_add_comp_mul_deriv_to_of_lt_iff_lt` in `InversePairIntegral.lean`. Their proof combines `intervalIntegral_smul_intervalIntegral_of_lt_iff_lt` with the fundamental theorem for absolutely-continuous functions; no derivative or Jacobian formula for the clocks enters.

**Corollary 4.9** (Fabius inverse-pair calculus and area sum). *For real functions  $A, B : \mathbb{R} \rightarrow \mathbb{R}$  that are absolutely continuous on  $[0, 1]$ ,*

$$\int_0^1 A'(x)B(F(x)) \, dx + \int_0^1 A(I(u))B'(u) \, du = A(1)B(1) - A(0)B(0). \quad (65)$$

*For every  $y \in [0, 1]$ , the corresponding variable-upper identity is*

$$\int_0^y A'(x)B(F(x)) \, dx + \int_0^{F(y)} A(I(u))B'(u) \, du = A(y)B(F(y)) - A(0)B(0). \quad (66)$$

In particular, taking both outer functions to be the identity gives the inverse-graph area identity

$$\int_0^1 F(x) dx + \int_0^1 I(u) du = 1. \quad (67)$$

At a variable inverse endpoint  $u \in [0, 1]$ , the same specialization gives

$$\int_0^u I(v) dv = uI(u) - \int_0^{I(u)} F(x) dx. \quad (68)$$

The complete and variable-upper two-function statements are exactly `intervalIntegral_deriv_mul_fabiusReal_add_fabiusInv_mul_deriv` and `intervalIntegral_deriv_mul_fabiusReal_add_fabiusInv_mul_deriv_to`; the two area statements are `intervalIntegral_fabiusReal_add_fabiusInv_eq_one` and `intervalIntegral_fabiusInv_to_eq_mul_sub_intervalIntegral_fabiusReal` in `InversePairIntegral.lean`. These are proper-compact first-order identities. Reversed-endpoint, higher-order, improper-integral, fractional-power, complex-valued arbitrary-AC, and complex-Mellin extensions remain outside these declarations.

### 4.3 The normalized geometric family

The same product probability space supports a more general family without changing the measure-theoretic argument. For  $q \in \mathbb{R}$  with  $|q| < 1$ , put

$$w_n(q) = (1 - q)q^n, \quad X_q = \sum_{n=0}^{\infty} w_n(q)U_n, \quad \mu_q = \text{law}(X_q). \quad (69)$$

For every real  $q$ ,  $w_0(q) = 1 - q$  and  $w_{n+1}(q) = q w_n(q)$ . Under  $|q| < 1$ , the weights are absolutely summable and have total mass one. The series is continuous and measurable on the full product space, and its law satisfies the exact affine refinement

$$\mu_q = (\text{Unif}[0, 1] \otimes \mu_q) \text{ map } ((u, x) \mapsto (1 - q)u + qx). \quad (70)$$

Equivalently,  $X_q \stackrel{d}{=} (1 - q)U_0 + qX'_q$ , where  $X'_q$  is an independent copy. The head coordinate and the shifted full series have product law. Coordinate reflection gives  $X_q(R\omega) = 1 - X_q(\omega)$  and  $\mu_q \text{ map } (x \mapsto 1 - x) = \mu_q$ , where  $(R\omega)_n = 1 - \omega_n$ . Its barycenter is

$$\int_{\mathbb{R}} x d\mu_q(x) = \frac{1}{2} \quad (|q| < 1). \quad (71)$$

This is the specialization `ProbabilityRepresentation.integral_id_geometricUniformDistribution_eq_one_half` of (55); it includes  $q = 0$  and every negative contraction and uses no positivity hypothesis. For every  $|q| < 1$ , the law  $\mu_q$  is absolutely continuous with respect to Lebesgue measure and has null singletons; no positivity hypothesis on  $q$  is needed. For every  $|q| < 1$ , the topological support of  $\mu_q$  is exactly the range  $X_q(\Omega)$ . When  $0 \leq q < 1$ , every weight is nonnegative,  $X_q(\omega) \in [0, 1]$ ,  $\mu_q([0, 1]) = 1$ ,  $\mu_q(\mathbb{R} \setminus [0, 1]) = 0$ , and  $\text{supp}(\mu_q) = [0, 1]$ ; the endpoint  $q = 0$  is included. At  $q = 1/2$ , both  $X_q$  and  $\mu_q$  are exactly the dyadic random series and law above.

The same affine equation admits a reusable uniqueness theorem far beyond this real uniform-digit example. Let  $D$  be a measurable space, let  $E$  be a complete, second-countable real inner-product space with its Borel measurable structure, let  $\rho$  be a probability measure on  $D$ , and let  $d : D \rightarrow E$  be measurable. For a probability measure  $\lambda$  on  $E$ , define

$$\mathcal{A}_{\rho, d, q}(\lambda) = (d_*\rho) * ((x \mapsto qx)_*\lambda) = ((u, x) \mapsto d(u) + qx)_*(\rho \otimes \lambda). \quad (72)$$

The second equality is the product-map form of convolution. Thus  $\mathcal{A}_{\rho, d, q}$  is precisely the law transformer for  $d(U) + qX$  when  $U \sim \rho$ ,  $X \sim \lambda$ , and the two inputs are independent.

**Theorem 4.10** (Contractive affine independent-copy law). *The transformer  $\mathcal{A}_{\rho,d,q}$  preserves probability measures, and its characteristic function factors as*

$$\varphi_{\mathcal{A}_{\rho,d,q}(\lambda)}(t) = \varphi_{d*\rho}(t) \varphi_{\lambda}(qt). \quad (73)$$

If  $\lambda$  is a fixed point, then for every  $n \in \mathbb{N}_0$ ,

$$\varphi_{\lambda}(t) = \left( \prod_{k=0}^{n-1} \varphi_{d*\rho}(q^k t) \right) \varphi_{\lambda}(q^n t). \quad (74)$$

If  $|q| < 1$ , the transformer has at most one probability fixed point. Equivalently, two probability measures satisfying the product-map equation in (72) are equal.

*Proof.* The factorization is the convolution product rule followed by the characteristic-function rule for real scalar multiplication. For two fixed points, their common digit factor has modulus at most one, so one recurrence step gives

$$|\varphi_{\lambda}(t) - \varphi_{\nu}(t)| \leq |\varphi_{\lambda}(qt) - \varphi_{\nu}(qt)|.$$

Iteration moves the argument to  $q^n t$ . Since  $|q| < 1$ , this tends to zero; continuity of characteristic functions and their common value 1 at the origin force equality at every  $t$ , and characteristic-function extensionality gives  $\lambda = \nu$ . The proof uses no support, density, absolute-continuity, or moment hypothesis on either fixed point or on the arbitrary measurable digit law. It includes  $q = 0$  and negative  $q$ .  $\square$

**Corollary 4.11** (Characterization of the geometric uniform law). *For every real  $q$  with  $|q| < 1$ , if a probability measure  $\lambda$  on  $\mathbb{R}$  satisfies*

$$\lambda = ((u, x) \mapsto (1 - q)u + qx)_* (\text{Unif}[0, 1] \otimes \lambda),$$

*then  $\lambda = \mu_q$ . Hence the refinement equation (70) characterizes the named law even at  $q = 0$  and for negative  $q$ , without any support, density, or moment assumption on  $\lambda$ .*

The CDF and its positive-ratio density are now named as well. Put

$$F_q(x) = \text{cdf}(\mu_q)(x), \quad f_q(x) = \frac{F_q(x/q) - F_q((x - (1 - q))/q)}{1 - q}. \quad (75)$$

Both definitions are total in the real parameter. For every real  $q$ ,  $F_q$  is measurable and monotone with values in  $[0, 1]$ . If  $|q| < 1$ , then

$$F_q \in C(\mathbb{R}), \quad F_q(1 - x) = 1 - F_q(x), \quad F_q(1/2) = 1/2. \quad (76)$$

If  $0 \leq q < 1$ , including  $q = 0$ , its exterior values are

$$F_q(x) = 0 \quad (x \leq 0), \quad F_q(x) = 1 \quad (x \geq 1). \quad (77)$$

The totalized quotient in (75) has a sharp parameter defect outside the positive-ratio range. Lean proves

$$f_0(x) = 0, \quad q < 0 \implies f_q(x) \leq 0.$$

Consequently, for every  $q \leq 0$ , clamping this selected formula by `ofReal` gives the zero `withDensity` measure. When also  $|q| < 1$ , that measure cannot equal the probability law  $\mu_q$ . This is a formal nonidentity result for the displayed total formula, not a denial of the already proved absolute continuity of  $\mu_q$ ; an exact Lean theorem for the paper-level corrected signed-ratio density at negative  $q$  remains open.



For  $0 < q < 1$ , conditioning on the head coordinate gives, for every real  $x$ ,

$$F_q(x) = \int_0^1 F_q\left(\frac{x - (1-q)u}{q}\right) du, \quad (78)$$

$$= \frac{q}{1-q} \int_{(x-(1-q))/q}^{x/q} F_q(t) dt. \quad (79)$$

The second expression in (75) is then the everywhere pointwise derivative of  $F_q$ . It is continuous, nonnegative, reflection-symmetric, vanishes for  $x \leq 0$  and for  $x \geq 1$ , has ordinary support contained in  $(0, 1)$  and topological support contained in  $[0, 1]$ , and is compactly supported. More precisely,

$$\mu_q = (\text{Leb} : \text{Measure}(\mathbb{R})) \text{ withDensity } (x \mapsto \text{ofReal}(f_q(x))), \quad F_q, f_q \in C^\infty(\mathbb{R}). \quad (80)$$

Thus the smoothness proof and the explicit Radon–Nikodym identification are formal in real space for the full positive-ratio range, without passing through Fourier inversion.

At the half base the named objects recover the classical functions globally:

$$F_{1/2} = \text{weightedSumCDF} = F, \quad f_{1/2}(x) = 2 \text{ up}(2x - 1) \quad (x \in \mathbb{R}). \quad (81)$$

**Theorem 4.12** (Fixed half–quarter multisection). *For a coordinate sequence  $\omega$ , put  $\omega_n^{(0)} = \omega_{2n}$  and  $\omega_n^{(1)} = \omega_{2n+1}$ . Then the normalized geometric series obeys the pointwise identity*

$$X_{1/2}(\omega) = \frac{2}{3} X_{1/4}(\omega^{(0)}) + \frac{1}{3} X_{1/4}(\omega^{(1)}). \quad (82)$$

*Under product Lebesgue measure, the two reindexed processes are independent copies. Consequently, if  $\mu_q$  is the law of  $X_q$  and  $\sigma_c(x) = cx$ , then*

$$\mu_{1/2} = (\mu_{1/4} \otimes \mu_{1/4}) \text{ map } ((x, y) \mapsto \frac{2}{3}x + \frac{1}{3}y) = (\sigma_{2/3})_* \mu_{1/4} * (\sigma_{1/3})_* \mu_{1/4}. \quad (83)$$

*No user hypothesis is needed at these fixed ratios.*

**Formal Lean counterpart.** The two public definitions `ProbabilityRepresentation.evenCoordinates` and `ProbabilityRepresentation.oddCoordinates`, followed by the three theorems `ProbabilityRepresentation.geometricUniformSeries_one_half_multisection`, `ProbabilityRepresentation.geometricUniformDistribution_one_half_multisection`, and `ProbabilityRepresentation.geometricUniformDistribution_one_half_conv_one_quarter`, form the complete public surface of `GeometricUniformMultisection.lean`. The first theorem is pointwise. The second proves the full parity-process product law, and the third rewrites its affine pushforward as measure convolution. This fixed result does not assert a general  $q$ - or  $m$ -multisection, an MGF or cumulant formula, a centered-density convolution, or a spectral dissection.  $\square$

**Formal API crosswalk.** The twenty-four public declarations in `GeometricUniformLaw.lean` are `ProbabilityRepresentation.geometricUniformWeight`, `ProbabilityRepresentation.geometricUniformWeight_zero`, `ProbabilityRepresentation.geometricUniformWeight_succ`, `ProbabilityRepresentation.hasSum_geometricUniformWeight`, `ProbabilityRepresentation.summable_norm_geometricUniformWeight`, `ProbabilityRepresentation.geometricUniformWeight_nonneg`, `ProbabilityRepresentation.geometricUniformSeries`, `ProbabilityRepresentation.continuous_geometricUniformSeries`, `ProbabilityRepresentation.measurable_geometricUniformSeries`, `ProbabilityRepresentation.geometricUniformSeries_split`, `ProbabilityRepresentation.geometricUniformSeries_reflect`, `ProbabilityRepresentation.geometricUniformSeries_mem_Icc`, `ProbabilityRepresentation.geometricUniformDistribution`, `Probab`

ilityRepresentation.geometricUniformDistribution\_isProbabilityMeasure, ProbabilityRepresentation.geometricUniformDistribution\_absolutelyContinuous, ProbabilityRepresentation.geometricUniformDistribution\_nullSingletonClass, ProbabilityRepresentation.uniformProduct\_map\_head\_tail\_geometricUniformSeries, ProbabilityRepresentation.geometricUniformDistribution\_selfSimilar, ProbabilityRepresentation.geometricUniformDistribution\_reflection, ProbabilityRepresentation.integral\_id\_geometricUniformDistribution\_eq\_one\_half, ProbabilityRepresentation.geometricUniformDistribution\_Icc, ProbabilityRepresentation.geometricUniformDistribution\_compl\_Icc, ProbabilityRepresentation.geometricUniformDistribution\_support\_eq\_range, and ProbabilityRepresentation.geometricUniformDistribution\_support\_eq\_Icc.

At compiler-validated checkpoint d312c0603, the generic affine interface in AffineIndependentCopy.lean consists of the definition ProbabilityTheory.affineIndependentCopyLaw and the eight theorems ProbabilityTheory.affineIndependentCopyLaw\_isProbabilityMeasure, ProbabilityTheory.affineIndependentCopyLaw\_eq\_map\_prod, ProbabilityTheory.charFun\_affineIndependentCopyLaw, ProbabilityTheory.charFun\_eq\_mul\_charFun\_of\_affineIndependentCopy\_fixedPoint, ProbabilityTheory.charFun\_iterate\_of\_affineIndependentCopy\_fixedPoint, ProbabilityTheory.eq\_of\_charFun\_affine\_recurrence, ProbabilityTheory.affineIndependentCopyLaw\_fixedPoint\_unique, and ProbabilityTheory.affineIndependentCopy\_map\_fixedPoint\_unique. The named specialization is ProbabilityRepresentation.eq\_geometricUniformDistribution\_of\_selfSimilar in GeometricUniformUniqueness.lean. The generic theorem is formulated for a complete second-countable real inner-product target and an arbitrary measurable digit map from an arbitrary probability space; the named theorem specializes it to uniform digits.

The predecessor GeometricUniformDictionary.lean already supplies finite measure, characteristic-function, MGF, and CGF tail factorizations. The six public theorems in GeometricSincFactorization.lean are charFun\_geometricUniformDigit, charFun\_geometricUniformDistribution\_prefix, charFun\_geometricUniformDistribution\_prefix\_sinc, tendsto\_prefix\_sinc\_charFun, charFun\_weightedSumDistribution\_prefix\_sinc, and tendsto\_prefix\_sinc\_charFun\_weightedSumDistribution. The digit formula is unconditional in real  $q$ . For every  $|q| < 1$ , depth  $m$ , and real frequency  $t$ , the phase-collapsed residual identity is

$$\phi_q(t) = e^{i(1-q^m)t/2} \left( \prod_{k=0}^{m-1} \text{sinc}_{\text{rad}} \left( \frac{(1-q)q^k t}{2} \right) \right) \phi_q(q^m t),$$

and the phase-bearing prefix without the residual converges pointwise to  $\phi_q(t)$ . This includes  $q = 0$  and negative  $q$ ; the final two declarations are exact  $q = 1/2$  wrappers for weightedSumDistribution. The Gamma-side product now supplies a named locally uniform product, pointwise Multipliable and HasProd forms, and entirety for the same rescaled sinc factors. The new module GeometricSincCharacteristicFunction.lean has no public definitions and exactly the four public theorems charFun\_geometricUniformDistribution\_eq\_phase\_mul\_geometricSincProduct and charFun\_geometricUniformDistribution\_eq\_phase\_mul\_geometricReciprocalGamma, followed by tendstoLocallyUniformly\_prefix\_sinc\_charFun and tendstoUniformlyOn\_prefix\_sinc\_charFun. For

$$z = z_{q,t} := \frac{(1-q)t}{2\pi},$$

the first two give, under the same sharp hypothesis  $|q| < 1$ ,

$$\phi_q(t) = e^{it/2} \Phi_q(z) = e^{it/2} G_q(z) G_q(-z).$$



For the full prefix

$$P_{q,m}(t) := e^{i(1-q^m)t/2} \prod_{k=0}^{m-1} \operatorname{sinc}_{\text{rad}}\left(\frac{(1-q)q^k t}{2}\right),$$

the last two prove  $P_{q,m} \rightarrow \phi_q$  locally uniformly on the real frequency line and uniformly on every compact  $K \subseteq \mathbb{R}$ . Thus all four results include  $q = 0$  and negative  $q$ . No complex-frequency analogue for the full phase-bearing prefixes or uniform-convergence claim on all of  $\mathbb{R}$  is made. The separate `RvachevPochhammerFactorization.lean` theorem below does supply the global general-contracting- $q$  spectral  $q^2$ -Pochhammer rearrangement of the pure sinc product; it does not add a complex-frequency theorem for the full phase-bearing prefixes.

The generic CDF mechanisms are `continuous_cdf_of_nullSingleton`, `cdf_reflection_sub`, and `measure_eq_withDensity_of_cdf_hasDerivAt` in `ContinuousCDF.lean`.

The exact thirty-one-declaration interface in `GeometricUniformCDF.lean` consists of `ProbabilityRepresentation.geometricUniformCDF`, `ProbabilityRepresentation.monotone_geometricUniformCDF`, `ProbabilityRepresentation.geometricUniformCDF_nonneg`, `ProbabilityRepresentation.geometricUniformCDF_le_one`, `ProbabilityRepresentation.measurable_geometricUniformCDF`, `ProbabilityRepresentation.continuous_geometricUniformCDF`, `ProbabilityRepresentation.geometricUniformCDF_reflection`, `ProbabilityRepresentation.geometricUniformCDF_one_half`, `ProbabilityRepresentation.geometricUniformCDF_zero_of_nonpos`, `ProbabilityRepresentation.geometricUniformCDF_one_of_one_le`, `ProbabilityRepresentation.geometricUniformCDF_eq_integral`, `ProbabilityRepresentation.geometricUniformCDF_eq_intervalIntegral`, `ProbabilityRepresentation.geometricUniformDensity`, `ProbabilityRepresentation.geometricUniformDensity_zero`, `ProbabilityRepresentation.geometricUniformDensity_nonpos_of_neg`, `ProbabilityRepresentation.volume_withDensity_geometricUniformDensity_eq_zero_of_nonpos`, `ProbabilityRepresentation.geometricUniformDistribution_ne_withDensity_geometricUniformDensity_of_nonpos`, `ProbabilityRepresentation.geometricUniformCDF_hasDerivAt`, `ProbabilityRepresentation.deriv_geometricUniformCDF`, `ProbabilityRepresentation.continuous_geometricUniformDensity`, `ProbabilityRepresentation.geometricUniformDensity_nonneg`, `ProbabilityRepresentation.geometricUniformDensity_zero_of_nonpos`, `ProbabilityRepresentation.geometricUniformDensity_zero_of_one_le`, `ProbabilityRepresentation.support_geometricUniformDensity_subset_Ioo`, `ProbabilityRepresentation.support_geometricUniformDensity_subset_Icc`, `ProbabilityRepresentation.tsupport_geometricUniformDensity_subset_Icc`, `ProbabilityRepresentation.geometricUniformDensity_hasCompactSupport`, `ProbabilityRepresentation.geometricUniformDensity_reflection`, `ProbabilityRepresentation.geometricUniformDistribution_eq_withDensity`, `ProbabilityRepresentation.contDiff_geometricUniformCDF`, and `ProbabilityRepresentation.contDiff_geometricUniformDensity`.

Exact half-base series and law recovery is `weightedCoordinateSum_eq_geometricUniformSeries_one_half` and `weightedSumDistribution_eq_geometricUniformDistribution_one_half` in `ProbabilityRepresentation.lean`. The CDF and density bridges there are `ProbabilityRepresentation.weightedSumCDF_eq_geometricUniformCDF_one_half`, `ProbabilityRepresentation.geometricUniformCDF_one_half_eq_fabiusReal`, and `ProbabilityRepresentation.geometricUniformDensity_one_half_eq_rvachevUp`; the exact dyadic support corollary is `weightedSumDistribution_support_eq_Icc`.  $\square$

The remaining boundary is parameter-sensitive. Although  $\mu_q$  is absolutely continuous throughout  $|q| < 1$ , the displayed total function  $f_q$  is identified as its Radon–Nikodym density only for  $0 < q < 1$ . At  $q = 0$  it is identically zero, and for negative  $q$  it is nonpositive; hence its `ofReal-clamped` measure

is zero for every  $q \leq 0$  and is formally unequal to  $\mu_q$  on the contraction range. An exact Lean theorem for the paper-level corrected signed-ratio density remains open. The positive-ratio  $C^\infty$  theorem is proved by the real-space derivative refinement. The phase-bearing general- $q$  sinc prefixes and their pointwise characteristic-function limit are formal; the pure complex sinc product now has named locally uniform convergence, pointwise HasProd, and entireness, and the four theorems above close its characteristic-function and reciprocal-Gamma bridges and upgrade the full prefixes to locally uniform convergence on the real frequency line and uniform convergence on every compact real frequency set. No complex-frequency full-prefix analogue or uniform-convergence assertion on all real frequencies is made. Rapid-decay estimates and a Fourier-inversion construction remain outside the formal API. So do a named centered  $q$ -family or MGF wrapper, a closed all-order derivative hierarchy and explicit endpoint-flatness wrappers, shape theorems, further transform formulas, and explicit Bernoulli-cumulant/Bell-moment formulas and asymptotics. The global complex Pochhammer theorem proved below now covers the pure centered product at every strict complex contraction. It does not close the distinct phase-bearing prefix, rapid-decay, or Fourier-inversion boundaries just listed. The probability-law fixed point itself is uniquely characterized by [Corollary 4.11](#). The oriented Cauchy transform and its adjacent-power hierarchy are also formal for every nonzero  $|q| < 1$ , including negative ratios, in `GeometricUniformCauchy.lean`; these are distinct from the closed characteristic-function product identities above and do not supply the still-open rapid-decay or Fourier-inversion layers.

#### 4.4 Sharp activation probabilities, finite Taylor jets, and square-summable fields

The hyperbolic activation kernel admits a particularly clean totalization at the origin. Write

$$\operatorname{sinhc}_{\mathbb{R}}(x) = \begin{cases} \sinh(x)/x, & x \neq 0, \\ 1, & x = 0, \end{cases} \quad h(x) = \frac{\operatorname{sinhc}_{\mathbb{R}}(x)}{\cosh x}, \quad (84)$$

$$r(x) = \frac{\operatorname{sinhc}_{\mathbb{R}}(2x)}{\operatorname{sinhc}_{\mathbb{R}}(x)^2} - 1, \quad p(x) = 1 - h(x). \quad (85)$$

Thus  $h(x) = \tanh(x)/x$  away from zero and  $h(0) = 1$ , while  $p(0) = 0$ . The four displayed objects are exactly `realSinhc`, `tanhDiv`, `activationOdds`, and `activationProbability` in `HyperbolicActivation.lean`. The denominator-free identity

$$p(x)(1 + r(x)) = r(x)$$

proves globally, including at  $x = 0$ , that  $p(x) = r(x)/(1 + r(x))$ .

**Theorem 4.13** (Sharp local activation law). *The function  $p$  is continuous and even, and for every  $x \in \mathbb{R}$ ,*

$$0 \leq p(x) < 1, \quad p(x) > 0 \iff x \neq 0, \quad p(x) \leq \min\left\{1, \frac{x^2}{3}\right\}. \quad (86)$$

*The quadratic constant is sharp. More precisely, for every real scale  $a$ , including  $a = 0$ ,*

$$\lim_{\substack{t \rightarrow 0 \\ t \neq 0}} \frac{p(at)}{t^2} = \frac{a^2}{3}. \quad (87)$$

*The quotient limit is punctured: no value at  $t = 0$  is assigned to the displayed quotient.*

The order statements in (86) are `activationProbability_nonneg`, `activationProbability_lt_one`, `activationProbability_pos_iff`, and `activationProbability_le_min_one_sq_div_three`; continuity and evenness are `continuous_activationProbability` and `activationProbability_even`. The away-from-zero formula and the denominator-free odds

bridge are `activationProbability_of_ne_zero`, `activationProbability_mul_one_add_activationOdds`, and `activationProbability_eq_odds_div`. The unscaled and scaled limits are `tendsto_activationProbability_div_sq` and `tendsto_activationProbability_mul_div_sq` in `ActivationAsymptotics.lean`. The scale-sensitive global estimate used below is `activationProbability_mul_le_quadratic` in `ActivationSeries.lean`:

$$p(at) \leq \frac{t^2}{3}a^2. \quad (88)$$

The proof of sharpness is deliberately lighter than a Taylor expansion. The global bound comes from  $x - x^3/3 \leq \tanh x$  on the nonnegative half-line and evenness. For the limit, one application of l'Hopital's rule to  $(x - \tanh x)/x^3$  produces  $(\tanh x/x)^2/3$ , whose limit is immediate from the continuous totalization  $h(0) = 1$ . Consequently, the coefficient  $1/3$  is proved without a higher-order remainder theorem. The finite-jet theorem below is logically independent of that l'Hopital argument and is not used to prove the sharp limit.

**Proposition 4.14** (The totalized tangent quotient is a divided slope). *For a differentiable real function  $f$ , write*

$$\text{dslope}_0(f)(x) = \begin{cases} \frac{f(x) - f(0)}{x}, & x \neq 0, \\ f'(0), & x = 0. \end{cases}$$

*Then the totalization in (84) satisfies the global function identity*

$$\boxed{h = \text{dslope}_0(\tanh)}. \quad (89)$$

*In particular, the displayed divided slope equals  $\tanh(x)/x$  away from zero and equals  $\tanh'(0) = 1$  at zero.*

This is exactly `tanhDiv_eq_dslope` in `HyperbolicActivation.lean`. It records the removable value as part of the function itself rather than as an informal convention. The value  $\tanh'(0) = 1$  is the specialization at zero of `hasDerivAt_tanh`.

**Theorem 4.15** (Finite Taylor jet of the activation probability). *Set*

$$\begin{aligned} T_9(x) &= x - \frac{x^3}{3} + \frac{2x^5}{15} - \frac{17x^7}{315} + \frac{62x^9}{2835}, \\ P_8(x) &= \frac{x^2}{3} - \frac{2x^4}{15} + \frac{17x^6}{315} - \frac{62x^8}{2835}. \end{aligned}$$

*As  $x \rightarrow 0$ , the following three Big-O statements hold:*

$$\tanh x - T_9(x) = \mathcal{O}(\|x\|^{11}), \quad (90)$$

$$p(x) - P_8(x) = \mathcal{O}(\|x\|^{10}), \quad (91)$$

$$p(x) - P_8(x) = \mathcal{O}(x^{10}). \quad (92)$$

*Thus, in the usual human shorthand,*

$$\boxed{p(x) = \frac{x^2}{3} - \frac{2x^4}{15} + \frac{17x^6}{315} - \frac{62x^8}{2835} + \mathcal{O}(x^{10})}.$$

*The last two statements are the canonical norm-power and literal real-power formulations of the same estimate;  $\|x\|^{10} = x^{10}$  because the exponent is even.*

The three estimates (90)–(92) are respectively `tanh_sub_taylor_nine_isBigO`, `activationProbability_sub_taylor_eight_isBigO`, and `activationProbability_sub_taylor_eight_isBigO_pow` in `ActivationTaylor.lean`. Their proof first obtains analyticity of  $\tanh$  from  $\tanh = \sinh / \cosh$ , using that real  $\cosh$  never vanishes. The denominator-free identity

$$\tanh x \cosh x = \sinh x$$

and the iterated Leibniz rule then determine the finite derivative table

$$(\tanh^{(n)}(0))_{n=0}^{10} = (0, 1, 0, -2, 0, 16, 0, -272, 0, 7936, 0).$$

The source packages these three proof steps as the private support declarations `analyticAt_tanh`, `tanh_mul_cosh`, and `iteratedDeriv_tanh_zero_table`. The canonical analytic power series and its finite partial-sum estimate give the order-eleven remainder in (90). Dividing the remainder by  $x$  on the punctured neighborhood lowers its order to ten, and the totalized value restores the origin. This argument proves precisely the displayed finite jet. It does not yet provide an all-order Bernoulli formula for the coefficients, a radius-of-convergence theorem, or a full convergent activation series.

**Theorem 4.16** (Arbitrary square-summable activation fields). *Let  $\iota$  be an arbitrary index type and let  $w : \iota \rightarrow \mathbb{R}$  satisfy*

$$\sum_{i \in \iota} w_i^2 < \infty. \quad (93)$$

*For every  $t \in \mathbb{R}$ , the family  $i \mapsto p(w_i t)$  is genuinely summable. If, only for this exposition, we abbreviate its sum by*

$$D_w(t) := \sum_{i \in \iota} p(w_i t), \quad (94)$$

*then*

$$D_w(t) \leq \frac{t^2}{3} \sum_{i \in \iota} w_i^2 \quad (95)$$

*and*

$$\boxed{\lim_{\substack{t \rightarrow 0 \\ t \neq 0}} \frac{D_w(t)}{t^2} = \frac{1}{3} \sum_{i \in \iota} w_i^2.} \quad (96)$$

*No countability assumption on  $\iota$  and no finite-support approximation are needed.*

Here  $D_w$  is article notation, not a named Lean definition: all three declarations use the expanded topological sum. Genuine summability is `summable_activationProbability_mul_of_summable_sq`, the global budget is `tsum_activationProbability_mul_le`, and the exact limit is `tendsto_tsum_activationProbability_mul_div_sq` in `ActivationSeriesAsymptotics.lean`. Tannery’s theorem applies directly to the coordinate limits (87); the common summable majorant is  $w_i^2/3$ , obtained from (88). Thus (94) is not an appeal to Lean’s zero-valued convention for a nonsummable `tsum`: hypothesis (93) proves the series summable at every field value before its sum is used.

The normalized geometric weights now become a one-line specialization. For  $|q| < 1$ , set

$$w_n(q) = (1 - q)q^n, \quad d_q(t) = \sum_{n=0}^{\infty} p(q^n(1 - q)t). \quad (97)$$

The second expression is `geometricActivationDimension` in `GeometricActivationDimension.lean`. The exact squared-weight identity is

$$\sum_{n=0}^{\infty} ((1 - q)q^n)^2 = \frac{1 - q}{1 + q}, \quad (98)$$

formalized in the convergence-carrying form `hasSum_normalizedGeometricWeight_sq`. Substitution into [Theorem 4.16](#) gives the following result.

**Corollary 4.17** (Sharp normalized geometric activation coefficient). *For every real  $q$  with  $|q| < 1$  and every real  $t$ , the series defining  $d_q(t)$  is summable and*

$$d_q(t) \leq \frac{1-q}{3(1+q)}t^2, \quad \boxed{\lim_{\substack{t \rightarrow 0 \\ t \neq 0}} \frac{d_q(t)}{t^2} = \frac{1-q}{3(1+q)}}. \quad (99)$$

At  $q = 1/2$ , write  $d_{\text{dy}} = d_{1/2}$ . Then

$$d_{\text{dy}}(t) = \sum_{n=0}^{\infty} p(2^{-n-1}t), \quad d_{\text{dy}}(t) \leq \frac{t^2}{9}, \quad \lim_{\substack{t \rightarrow 0 \\ t \neq 0}} \frac{d_{\text{dy}}(t)}{t^2} = \frac{1}{9}. \quad (100)$$

The summability, normalized bound, and sharp limit in (99) are respectively `summable_geometricActivationDimension_terms`, `geometricActivationDimension_le_normalized_quadratic`, and `tendsto_geometricActivationDimension_div_sq`. The dyadic expansion, bound, and sharp limit in (100) are `dyadicEffectiveDimension_eq_tsum`, `dyadicEffectiveDimension_le_sq_div_nine`, and `tendsto_dyadicEffectiveDimension_div_sq`; the named dyadic definition is `dyadicEffectiveDimension`. The last two limit theorems are in `GeometricActivationAsymptotics.lean`.

The finite-depth API gives more than the limit. In the convergent range,

$$d_q(t) = p((1-q)t) + d_q(qt), \quad (101)$$

$$d_q(t) = \sum_{n < N} p(q^n(1-q)t) + d_q(q^N t), \quad (102)$$

$$0 \leq d_q(q^N t) \leq \frac{1-q}{3(1+q)}t^2(q^2)^N. \quad (103)$$

These are `geometricActivationDimension_refinement`, `geometricActivationDimension_eq_sum_range_add`, and `geometricActivationDimension_tail_le`. Together with `sum_range_activationProbability_le_geometricActivationDimension` and `geometricActivationDimension_le_sum_range_add_tail`, they sandwich the infinite sum between a finite prefix and that prefix plus the explicit tail certificate. At the dyadic base, `dyadicEffectiveDimension_half_refinement`, `dyadicEffectiveDimension_two_mul`, `dyadicEffectiveDimension_refinement`, `dyadicEffectiveDimension_eq_sum_range_add`, and `dyadicEffectiveDimension_tail_le` give the corresponding half-scale, doubling, difference, prefix, and tail formulas. Positivity away from zero and evenness are recorded by `geometricActivationDimension_pos_iff`, `geometricActivationDimension_even`, `dyadicEffectiveDimension_pos_iff`, and `dyadicEffectiveDimension_even`.

The analytic range  $|q| < 1$  in (99) deliberately includes negative  $q$ : there the weights alternate, but  $d_q$  is still a perfectly valid deterministic activation series. The positive-weight “active count” interpretation used in the research report is narrower,  $0 < q < 1$ . In particular, no Lean declaration constructs random Bernoulli activations or asserts an identity of the form  $d_{1/2}(t) = \mathbb{E}K_t$ . The quotient limits above are all punctured. What remains open in this lane is neither the quadratic activation asymptotic nor the finite jet of [Theorem 4.15](#), but coefficients beyond that jet, an all-order Bernoulli coefficient formula, a radius-of-convergence and full-series theorem, and the proposed Bernoulli expectation models.

At immutable source checkpoint `a345425d21d90e680bf15e34093af42c69c08a83`, the complete activation tranche consists of the seven modules `HyperbolicActivation.lean`, `ActivationSeries.lean`, `ActivationAsymptotics.lean`, `ActivationTaylor.lean`,

`ActivationSeriesAsymptotics.lean`, `GeometricActivationDimension.lean`, and `GeometricActivationAsymptotics.lean`. Their public inventories are, in that order, four definitions and 58 theorems, three theorems, two theorems, three theorems, one theorem, two definitions and 30 theorems, and two theorems: six definitions and 99 theorems, hence 105 public declarations in total.

#### 4.5 Finite atomic laws and weak convergence

There is also a completely discrete approximation theorem. For  $k \geq 0$ , let  $\beta_k$  be the probability measure assigning mass  $1/2$  to each of  $\pm 2^{-k-1}$ . Starting from  $\nu_0 = \delta_0$ , define

$$\nu_{n+1} = \nu_n * \beta_{n+1}^{*(n+1)}. \quad (104)$$

Thus  $\nu_n$  is a finite atomic probability measure. Its characteristic function is the finite weighted cosine product

$$\varphi_{\nu_n}(t) = \prod_{k=0}^{n-1} \left( \cos \frac{t}{2^{k+2}} \right)^{k+1}. \quad (105)$$

The indexing in (104) is worth noticing: the new scale at stage  $n + 1$  is  $2^{-n-2}$ , and it occurs with multiplicity  $n + 1$ .

Let  $\mu_{\text{up}}$  denote the formal measure `rvachevMeasure`; by definition it is Lebesgue measure with density of `Real` `up`. It is absolutely continuous with respect to Lebesgue measure; in particular, every Lebesgue-null set is also  $\mu_{\text{up}}$ -null. These are `rvachevMeasure_absolutelyContinuous` and `rvachevMeasure_eq_zero_of_volume_eq_zero` in `StepMeasureBridge.lean`. Its characteristic function uses the positive-frequency normalization

$$\text{charFun}(\mu_{\text{up}}, t) = \widehat{\text{up}}_{2\pi} \left( \frac{t}{2\pi} \right) \quad (t \in \mathbb{R}), \quad (106)$$

the exact identity `rvachevMeasure_charFun_pos` in `WeakConvergence.lean`. The finite laws converge weakly to  $\mu_{\text{up}}$ :

$$\nu_n \Longrightarrow \mu_{\text{up}}. \quad (107)$$

Equivalently, for every bounded continuous real-valued map  $g$ ,

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}} g(x) \, d\nu_n(x) = \int_{\mathbb{R}} g(x) \, d\mu_{\text{up}}(x). \quad (108)$$

The same test-function theorem is formalized for complex-valued, and more generally real-or-complex-like, bounded continuous functions.

*Formal proof.* The definitions are `centeredBernoulliMeasure` and `finiteConvolutionMeasure`. Their probability-measure instances and (105) are proved in `EarlyApproximants.lean`. Pointwise convergence of the characteristic functions, weak convergence (107), and the test-function formulations (108) are respectively `finiteConvolutionMeasure_charFun_tendsto`, `finiteConvolutionProbability_tendsto_fabius`, and `integral_finiteConvolutionMeasure_tendsto_fabius` in `WeakConvergence.lean`.  $\square$

#### 4.6 Combinatorial step functions and the half-weight convention

A second finite model replaces each atom by a narrow interval. Define polynomials with nonnegative integer coefficients by

$$\mathcal{P}_0(X) = 1, \quad \mathcal{P}_m(X) = \mathcal{P}_{m-1}(X^2)(1 + X)^m \quad (m \geq 1). \quad (109)$$

They also have the finite product form

$$\mathcal{P}_m(X) = \prod_{j=1}^m (1 + X + \cdots + X^{2^j-1}). \quad (110)$$

Writing  $\mathcal{P}_m(X) = \sum_r a_{m,r} X^r$ , the coefficient  $a_{m,r}$  counts the tuples

$$(s_1, \dots, s_m), \quad 0 \leq s_j \leq 2^j - 1, \quad s_1 + \cdots + s_m = r. \quad (111)$$

Moreover,

$$g_m := \deg \mathcal{P}_m = 2^{m+1} - m - 2, \quad \mathcal{P}_m(1) = 2^{\binom{m+1}{2}}. \quad (112)$$

For  $\alpha < \beta$ , let  $\mathbf{1}_{[\alpha,\beta]}^*$  be 1 in the interior, 1/2 at either endpoint, and 0 elsewhere. Put

$$I_{m,r} = \left[ \frac{2r-1-g_m}{2^{m+1}}, \frac{2r+1-g_m}{2^{m+1}} \right] \quad (113)$$

and

$$\mathcal{W}_m(x) = \frac{2^m}{2^{\binom{m+1}{2}}} \sum_{r=0}^{g_m} a_{m,r} \mathbf{1}_{I_{m,r}}^*(x). \quad (114)$$

**Theorem 4.18** (Corrected step approximation). *For every  $m \geq 0$  and  $x \in \mathbb{R}$ ,*

$$0 \leq \mathcal{W}_m(x) \leq 1, \quad \mathcal{W}_m(0) = 1, \quad \int_{\mathbb{R}} \mathcal{W}_m(x) dx = 1.$$

*The function  $\mathcal{W}_m$  is nondecreasing on  $(-\infty, 0)$  and nonincreasing on  $(0, \infty)$ . For every real  $x$ ,*

$$\mathcal{W}_m(x) \longrightarrow \text{up}(x). \quad (115)$$

The proof of pointwise convergence passes through a useful exact comparison. Set  $h_m = 2^{-m-1}$ , and use the finite atomic law  $\nu_m$  from (104). Whenever  $a \leq b$ ,

$$\nu_m((a + h_m, b - h_m]) \leq \int_a^b \mathcal{W}_m(x) dx \leq \nu_m((a - h_m, b + h_m]). \quad (116)$$

Consequently,

$$\lim_{m \rightarrow \infty} \int_a^b \mathcal{W}_m(x) dx = \int_a^b \text{up}(x) dx. \quad (117)$$

Monotonicity on the two open half-lines, the exact value at the origin, and continuity of  $\text{up}$  then upgrade these interval limits to (115).

*Formal proof.* The polynomial recursion, product, degree, total coefficient mass, and bounded-partition interpretation are respectively proved by `approximationPolynomial_succ`, `approximationPolynomial_natDegree`, `approximationDegree_eq_sub`, `approximationPolynomial_eval_one`, and `approximationPolynomial_coeff_eq_card` in `EarlyApproximants.lean`. The finite atomic description is identified with `finiteConvolutionMeasure` by `polynomialMeasure_eq_finiteConvolutionMeasure` in `EarlyMeasureBridge.lean`.

For nonnegativity, see `stepApproximant_nonneg` in `EarlyApproximants.lean`. The remaining analytic assertions are `stepApproximant_le_one`, `stepApproximant_monotoneOn_Iio`, `stepApproximant_antitoneOn_Ioi`, `stepApproximant_apply_zero`, and `stepApproximant_tendsto_fabius` in `StepApproximationLimit.lean`. The normalization, the two sides of (116), and (117) are `integral_stepApproximant`, `finiteConvolutionMeasure_inner_le_intervalIntegral_stepApproximant`, `intervalIntegral_stepApproximant_le_finiteConvolutionMeasure_outer`, and `intervalIntegral_stepApproximant_tendsto_of_le` in `StepMeasureBridge.lean`.  $\square$



*Warning 4.19* (Why the endpoint convention is explicit). The half-weighted cell representative agrees almost everywhere with the ordinary interval indicator, so it has the same integral. Its point values are nevertheless part of the formal statement: adjacent cells share endpoints, and each contributes exactly half of its coefficient there. In particular, the corrected representative satisfies  $\mathcal{W}_m(0) = 1$  at every level. These facts are formalized by `halfEndpointIntervalIndicator_ae_eq_indicator`, `intervalIntegral_halfEndpointIntervalIndicator_of_le`, and `stepApproximant_apply_zero`.

#### 4.7 Centered finite splines and an explicit error bound

The preceding finite random sums also give a completely explicit polynomial approximation to  $F$ . Put

$$T_p = \sum_{i=0}^{p-1} \frac{U_i}{2^{i+1}}, \quad C_p(x) = \mathbb{P}(T_p \leq x - 2^{-p-1}), \quad (118)$$

where  $p \geq 1$ . The shift by half of the omitted tail is important: it centers the one-sided approximation and improves its error by a factor of two.

Let

$$N_p(x) = \max \left( 0, \left\lfloor 2^p x + \frac{1}{2} \right\rfloor \right), \quad \varepsilon_r = (-1)^{\text{wt}_2(r)}, \quad (119)$$

and interpret a sum with  $N_p(x) = 0$  as empty. Define

$$S_p(x) = \frac{(-1)^p}{2^{\binom{p}{2}} p!} \sum_{r=0}^{N_p(x)-1} \varepsilon_r \left( r - 2^p x + \frac{1}{2} \right)_+^p. \quad (120)$$

Equivalently,

$$S_p(x) = \frac{1}{2^{\binom{p}{2}} p!} \sum_{r \geq 0} \varepsilon_r \left( 2^p x - \frac{1}{2} - r \right)_+^p, \quad (121)$$

where only the terms  $r < N_p(x)$  can be nonzero and  $t_+ = \max(t, 0)$ . The cutoff in (119) is a safe natural-number form of the inclusive upper limit

$$0 \leq r \leq \left\lfloor 2^p x - \frac{1}{2} \right\rfloor.$$

It handles the empty case without assigning a negative upper endpoint to a finite sum.

**Theorem 4.20** (Centered spline as a finite distribution function). *For  $p \geq 1$  and  $0 \leq x \leq 1$ ,*

$$S_p(x) = C_p(x). \quad (122)$$

*Consequently  $0 \leq S_p(x) \leq 1$  on the unit interval.*

*Proof.* For  $p = 1$ ,  $T_1 = U_0/2$ , and direct evaluation gives

$$C_1(x) = \max \left( 0, \min \left( 1, 2x - \frac{1}{2} \right) \right).$$

Formula (121) gives exactly the same broken-linear function: the term  $r = 0$  starts at  $x = 1/4$ , and the term  $r = 1$  subtracts the excess after  $x = 3/4$ .

For the induction step, split off the first uniform coordinate. The finite sums satisfy

$$T_{p+1} \stackrel{\text{d}}{=} \frac{U_0 + T'_p}{2},$$

where  $T'_p$  is an independent copy of  $T_p$ . Conditioning on  $U_0$  and accounting for the midpoint shifts gives the exact smoothing recurrence

$$C_{p+1}(x) = \int_0^1 C_p(2x - u) \, du. \quad (123)$$

On the spline side, integrate each truncated power in (121). The two binary children  $2r$  and  $2r + 1$  have opposite Thue–Morse signs, because  $\varepsilon_{2r} = \varepsilon_r$  and  $\varepsilon_{2r+1} = -\varepsilon_r$ . After the change of scale this yields

$$S_{p+1}(x) = \int_0^1 S_p(2x - u) \, du. \quad (124)$$

The normalization is exactly the one required by

$$\sum_{r=0}^{2^p-1} \varepsilon_r r^p = (-1)^p 2^{\binom{p}{2}} p!,$$

while every lower power sum vanishes. Thus the two recurrences and their common base case prove (122).  $\square$

**Theorem 4.21** (Quantitative centered approximation). *For  $p \geq 1$  and  $0 \leq x \leq 1$ ,*

$$F(x - 2^{-p-1}) \leq S_p(x) \leq F(x + 2^{-p-1}), \quad (125)$$

and hence

$$|S_p(x) - F(x)| \leq 2^{-p}. \quad (126)$$

*Proof.* Write  $X = T_p + R_p$ , where the omitted tail satisfies

$$0 \leq R_p = \sum_{i=p}^{\infty} \frac{U_i}{2^{i+1}} \leq 2^{-p}.$$

If  $X \leq x - 2^{-p-1}$ , then certainly  $T_p \leq x - 2^{-p-1}$ . Conversely, if  $T_p \leq x - 2^{-p-1}$ , then  $X \leq x + 2^{-p-1}$ . Taking probabilities and using (38) and (122) proves (125).

The bounded Fabius function is globally 2-Lipschitz. Indeed, on the open unit interval its derivative is a rescaled value of up (or of the signed extension), whose absolute value is at most 1, and outside the unit interval  $F$  is constant. Thus

$$|F(x \pm 2^{-p-1}) - F(x)| \leq 2 \cdot 2^{-p-1} = 2^{-p}.$$

Combining these two inequalities with the sandwich gives (126).  $\square$

The bound is uniform on the closed interval and includes both endpoints. It is stronger than mere weak convergence of the finite distributions, and it is the quantitative input for the computability proof in section 15.

The same finite formula also has the correct signed global behavior. For  $x \leq 0$  its safe cutoff is empty, so  $S_p(x) = F_{\text{gl}}(x) = 0$ . On the nonnegative half-line, binary block translation and midpoint reflection extend the local identification. Consequently, for every  $p \geq 1$  and every  $x \in \mathbb{R}$ ,

$$|S_p(x) - F_{\text{gl}}(x)| \leq 2^{-p}. \quad (127)$$

Indeed, every nonnegative two-unit block is a common signed copy of the first block, and on the second unit cell both functions use the same complement. Neither operation changes the absolute error in (126); the nonpositive half-line is exact. This pointwise all-real bound is `abs_fabiusUniformSpline_sub_extendedFabius_1e`; it implies the separately proved uniform convergence `fabiusUniformSpline_tendstoUniformly_extendedFabius`. This global form strengthens and packages the analytic input used by the complex-shift and discrete-limit results below.

### 4.8 The exact quarter cell of the finite spline

The indexing used in the inverse-frontier calculations is shifted by one from (120). Write

$$P_n(x) := S_{n-1}(x) \quad (n \geq 1). \quad (128)$$

In Lean this is `reportFiniteFabiusApproximant`; keeping the definition explicit prevents a degree-one indexing error. The complete dyadic cell about  $1/4$  has a remarkably simple closed form.

**Theorem 4.22** (Exact two-sided quarter cell). *For every  $n \geq 3$  and every real  $z$  with  $|z| \leq 2^{-n}$ ,*

$$P_n\left(\frac{1}{4} + z\right) = \frac{5}{72} + z + 4z^2 - \frac{4}{9}4^{-n}. \quad (129)$$

Consequently,

$$P_n\left(\frac{1}{4} + z\right) - P_n\left(\frac{1}{4} - z\right) = 2z, \quad (130)$$

$$P_n\left(\frac{1}{4} + z\right) + P_n\left(\frac{1}{4} - z\right) - 2P_n\left(\frac{1}{4}\right) = 8z^2. \quad (131)$$

Moreover,  $P_n$  is strictly increasing on the whole closed interval

$$\left[\frac{1}{4} - 2^{-n}, \frac{1}{4} + 2^{-n}\right].$$

*Formal proof.* Equation (129) is `reportFiniteFabiusApproximant_quarter_eq_quadratic`, with its absolute-value wrapper `reportFiniteFabiusApproximant_quarter_twoSided`. The proof computes the safe cutoff on the entire closed cell, including the exceptional right endpoint, and reduces the remaining block to the first two Thue–Morse moments beyond the Prouhet cancellation range. The reflection and centered second-difference identities are `reportFiniteFabiusApproximant_quarter_reflection` and `reportFiniteFabiusApproximant_quarter_centralSecondDifference`. Strict monotonicity on the full cell is `strictMonoOn_reportFiniteFabiusApproximant_quarter_twoSided`. The finite moment extraction is centralized in `QuarterSplineLocalPolynomial.lean`; the two-sided semantic wrappers are in `QuarterSplineTwoSided.lean`.

For completeness, the two signed Thue–Morse moments that survive beyond the Prouhet cancellation range are worth recording. If

$$M_m(d) := \sum_{r < 2^m} \varepsilon_r r^d,$$

then

$$M_m(m+1) = (-1)^m 2^{\binom{m}{2}} m! \frac{(m+1)(2^m - 1)}{2}, \quad (132)$$

$$M_m(m+2) = (-1)^m 2^{\binom{m}{2}} m! (m+1)(m+2) \left( \frac{(2^m - 1)^2}{8} + \frac{4^m - 1}{72} \right). \quad (133)$$

These are `thueMorsePowerSumRing_add_one_real` and `thueMorsePowerSumRing_add_two_real`. Their translated degree- $m+2$  extractor and the normalized quarter specialization are `thueMorse_translated_power_sum_add_two_real`, `normalized_thueMorse_translated_power_sum_add_two`, and `normalized_quarter_thueMorse_block`. This is the exact arithmetic source of both the constant  $5/72$  and the finite-depth correction in (129).  $\square$

Define the displacement

$$\delta_n := \frac{\sqrt{1 + \frac{64}{9}4^{-n}} - 1}{8}. \quad (134)$$

For every  $n \geq 3$ , the finite-spline identity

$$P_n\left(\frac{1}{4} + \delta_n\right) = \frac{5}{72} \quad (135)$$

is `reportFiniteFabiusApproximant_quarterPrefix_value`. The definition, positivity, and cell bound are respectively `quarterPrefixDisplacement`, `quarterPrefixDisplacement_pos`, and `quarterPrefixDisplacement_le_inv_two_pow` in `InverseQuarterAnchor.lean`; in particular, the point lies inside the exact cell. The same quadratic yields an exact inverse value, but only after an inverse has been specified. For  $n \geq 3$ , if  $G$  is any left inverse of  $P_n$  on the right half-cell  $[1/4, 1/4 + 2^{-n}]$ , then

$$G\left(\frac{5}{72}\right) = \frac{1}{4} + \frac{\sqrt{1 + \frac{64}{9}4^{-n}} - 1}{8}. \quad (136)$$

This conditional statement is `reportFiniteFabiusApproximant_quarterPrefix_quantile`. It does not define a canonical finite inverse  $G_n$ , identify  $G$  with the limiting inverse `fabiusInv`, or assert convergence of finite inverse branches. Those are separate analytic questions, not consequences of the cell polynomial alone.

## 5 Fourier products, inversion, and Poisson summation

Use the convention

$$\widehat{f}_{2\pi}(z) = \int_{\mathbb{R}} f(x) e^{-2\pi i x z} dx. \quad (137)$$

The formal Fourier transform of `up` is entire and has two exact infinite-product descriptions.

**Theorem 5.1** (Sinc and cosine products). *For every  $z \in \mathbb{C}$ ,*

$$\widehat{\text{up}}_{2\pi}(z) = \prod_{n=0}^{\infty} \text{sinc}_{\text{rad}}\left(\frac{\pi z}{2^n}\right), \quad (138)$$

$$= \prod_{m=1}^{\infty} \left(\cos \frac{\pi z}{2^m}\right)^m. \quad (139)$$

*The transform also satisfies*

$$\widehat{\text{up}}_{2\pi}(z) = \text{sinc}_{\text{rad}}(\pi z) \widehat{\text{up}}_{2\pi}(z/2). \quad (140)$$

*Formal proof.* The finite-product refinement and its infinite limit are `rvachevFourier_eq_finite_product` and `rvachevFourier_eq_product` in `FourierProduct.lean`. The refinement equation is `rvachevFourier_scaling` in the same module. Writing  $\Phi$  for the first product in (138), the standalone one-step and  $n$ -fold renormalizations

$$\Phi(z) = \text{sinc}_{\text{rad}}(\pi z) \Phi(z/2), \quad \Phi(z) = \left( \prod_{j < n} \text{sinc}_{\text{rad}}(\pi z/2^j) \right) \Phi(z/2^n)$$

are `rvachevFourierProduct_scaling` and `rvachevFourierProduct_shell` in `RenormalizationIdentity.lean`. The whole-plane analyticity theorem is `rvachevFourier_analyticOnNhd` in `FourierAnalytic.lean`. The weighted cosine product is `rvachevFourierProduct_eq_cosineProduct` in `OriginalPaperSupplement.lean`; its finite telescoping identity is proved internally in the same module.  $\square$

### 5.1 Finite complex transforms and a Thue–Morse product-to-sum bridge

The finite binary block has a reflection law that matches the parity of a quarter-turn phase. Recall that  $\varepsilon_n = (-1)^{\text{wt}_2(n)}$ .

**Theorem 5.2** (Finite complex Boolean-cube and Thue–Morse transforms). *For every  $m \in \mathbb{N}_0$  and arbitrary  $u, a, z \in \mathbb{C}$ ,*

$$\sum_{n=0}^{2^m-1} u^{\text{wt}_2(n)} e^{nz} = \prod_{j=0}^{m-1} (1 + ue^{2^j z}), \quad (141)$$

$$\sum_{n=0}^{2^m-1} u^{\text{wt}_2(n)} e^{a+nz} = e^a \prod_{j=0}^{m-1} (1 + ue^{2^j z}), \quad (142)$$

$$\sum_{n=0}^{2^m-1} \varepsilon_n e^{nz} = \prod_{j=0}^{m-1} (1 - e^{2^j z}), \quad (143)$$

$$\sum_{n=0}^{2^m-1} \varepsilon_n e^{a+nz} = e^a \prod_{j=0}^{m-1} (1 - e^{2^j z}). \quad (144)$$

All four identities include  $m = 0$ : the products are then empty, the two unshifted identities read  $1 = 1$ , and the two affine identities read  $e^a = e^a$ . In particular, the weighted formulas also include  $u = 0$ , with the usual convention  $u^0 = 1$ .

*Formal proof.* The four identities are respectively `sum_range_two_pow_binaryWeight_cexp`, `sum_range_two_pow_binaryWeight_cexp_affine`, `sum_range_two_pow_thueMorseSign_cexp`, and `sum_range_two_pow_thueMorseSign_cexp_affine` in `ThueMorseComplexExponential.lean`. The first is the Boolean-cube product after substituting  $e^z$ ; the affine identity extracts the common factor  $e^a$ . The signed pair is the specialization encoded by  $\varepsilon_n = (-1)^{\text{wt}_2(n)}$ .

These four declarations are the raw finite exponential layer. The separate `ThueMorseComplexProductBridge.lean` builds on the same Boolean-cube identity to provide the complex half-angle formula and total sinc and negative-Laplace prefixes, including their removable origins. Neither finite module asserts convergence to an infinite product.  $\square$

**Proposition 5.3** (Binary complement and quarter-turn parity). *For  $m, n \in \mathbb{N}_0$  with  $n < 2^m$ ,*

$$\text{wt}_2(2^m - 1 - n) + \text{wt}_2(n) = m, \quad (145)$$

$$\varepsilon_{2^m-1-n} = (-1)^m \varepsilon_n. \quad (146)$$

For every  $m \in \mathbb{N}_0$  and  $x \in \mathbb{R}$ ,

$$(-1)^m \cos\left(\frac{m\pi}{2} - x\right) = \cos\left(\frac{m\pi}{2} + x\right). \quad (147)$$

*Formal proof.* The three identities are respectively `binaryWeight_compl`, `thueMorseSign_compl`, and `neg_one_pow_mul_cos_sub` in `LacunaryProductToSum.lean`. The first hypothesis makes the natural-number subtractions in (145)–(146) ordinary binary complementation rather than truncated subtraction.  $\square$

**Theorem 5.4** (Affine Thue–Morse sine polynomial). *For every  $m \in \mathbb{N}_0$  and  $x, \phi \in \mathbb{R}$ ,*

$$\sum_{n=0}^{2^m-1} \varepsilon_n \sin(\phi + nx) = 2^m \left( \prod_{j=0}^{m-1} \sin \frac{2^j x}{2} \right) \sin \left( \phi + \frac{(2^m - 1)x}{2} - \frac{m\pi}{2} \right). \quad (148)$$

The product is empty when  $m = 0$ .

*Formal proof.* This is exactly `sum_thueMorseSign_mul_sin_affine` in `ThueMorseSineProduct.lean`. Its proof rotates the complex Thue–Morse exponential product through the arbitrary phase  $\phi$ , rewrites  $(-2i)^m$  as a positive magnitude and a quarter-turn phase, and takes imaginary parts. The theorem displayed here is real-valued; the preceding theorem supplies the raw finite complex exponential identity, while `sum_thueMorseSign_cexp_eq_sin_prod` and `thueMorseBlock_cexp_eq_sincPrefix` in `ThueMorseComplexProductBridge.lean` supply the complex half-angle and total sinc normalizations.  $\square$

**Theorem 5.5** (Lacunary sine product as a Thue–Morse sum). *For every  $m \in \mathbb{N}_0$  and  $w \in \mathbb{R}$ ,*

$$2^m \prod_{j=0}^m \sin(2^j w) = \sum_{n=0}^{2^m-1} \varepsilon_n \sin\left(\frac{m\pi}{2} + (2n+1)w\right). \quad (149)$$

*Formal proof.* This is exactly `prod_sin_two_pow_eq_thueMorse_sum` in `LacunaryProductToSum.lean`. It is the specialization of (148) at

$$x = 2w, \quad \phi = \frac{m\pi}{2} + w.$$

The final sine becomes  $\sin(2^m w)$ , which appends the last factor to the product over  $j < m$ . This formal statement is over real  $w$ ; its neighboring real cosine-product identity is `prod_sin_two_pow_mul_cos_eq_thueMorse_sum` in `LacunaryProductToSumCosine.lean`, and the complex half-angle layer is recorded separately by `ThueMorseComplexProductBridge.lean`.  $\square$

The same polynomial is exactly the denominator-cleared shell of the centered sinc product. Define, for  $N \in \mathbb{N}_0$ ,

$$Q_N(w) := \prod_{r=0}^{N-1} \text{sinc}_{\text{rad}}\left(\frac{w}{2^{r+1}}\right), \quad T_m(w) := \sum_{n=0}^{2^m-1} \varepsilon_n \sin\left(\frac{m\pi}{2} + (2n+1)w\right). \quad (150)$$

The prefix  $Q_N$  is the real-argument restriction of `centeredSincPartialProduct` from `FourierLaplaceRotation.lean`.

**Theorem 5.6** (Denominator-cleared Thue–Morse sinc shell). *For every  $m \in \mathbb{N}_0$  and  $w \in \mathbb{R}$ ,*

$$T_m(w) = 2^m \prod_{j=0}^m \sin(2^j w), \quad (151)$$

$$2^{m+\binom{m+1}{2}} w^{m+1} Q_{m+1}(2^{m+1} w) = T_m(w), \quad (152)$$

$$2^{m+\binom{m+1}{2}} w^{m+1} \Phi\left(\frac{2^m w}{\pi}\right) = T_m(w) \Phi\left(\frac{w}{2\pi}\right). \quad (153)$$

*All three identities are total at  $w = 0$ ; in particular, no removable denominator is cancelled there. The polynomial has parity*

$$T_m(-w) = (-1)^{m+1} T_m(w), \quad (154)$$

*and its complete factor-level zero criterion is*

$$T_m(w) = 0 \iff \exists j < m+1 : \sin(2^j w) = 0. \quad (155)$$

*If  $w \neq 0$ , then both the finite prefix and the dilated full product vanish exactly when  $T_m(w)$  does:*

$$Q_{m+1}(2^{m+1} w) = 0 \iff T_m(w) = 0, \quad (156)$$

$$\Phi\left(\frac{2^m w}{\pi}\right) = 0 \iff T_m(w) = 0. \quad (157)$$

*Formal proof.* The definition, product identity, value at zero, parity, and factor-level zero criterion are `thueMorseSinePolynomial`, `thueMorseSinePolynomial_eq_prod_sin_two_pow`, `thueMorseSinePolynomial_zero`, `thueMorseSinePolynomial_neg`, and `thueMorseSinePolynomial_eq_zero_iff`. The two cross-multiplied identities are `centeredSincPartialProduct_dyadic_eq_thueMorse` and `rvachevFourierProduct_dyadic_eq_thueMorse`. Their sharp zero criteria are `centeredSincPartialProduct_dyadic_eq_zero_iff` and `rvachevFourierProduct_dyadic_eq_zero_iff_thueMorse`; the latter uses the already proved global real-axis zero classification to show that the apparent tail-zero alternative is already a zero of the first exposed sine factor. All declarations are in `CenteredRvachevThueMorseFourier.lean`.

Here  $\Phi$  denotes the standalone convergent product `rvachevFourierProduct`. Passing from (153) to the integral Fourier transform of a particular bounded Fabius solution uses the separate identification `rvachevFourier_eq_product`; it is not hidden in the finite algebra.  $\square$

**Theorem 5.7** (Dyadic shell factorization). *For every  $k \in \mathbb{N}_0$  and  $z \in \mathbb{C}$ , dilation by  $2^k$  extracts exactly the first  $k$  factors of the sinc product:*

$$\widehat{\text{up}}_{2\pi}(2^k z) = \left( \prod_{j=0}^{k-1} \text{sinc}_{\text{rad}}(\pi 2^{j+1} z) \right) \widehat{\text{up}}_{2\pi}(z). \quad (158)$$

*Formal proof.* This is `rvachevFourier_two_pow_mul` in `RvachevFourierDecay.lean`. Its standalone-product form is `rvachevFourierProduct_two_pow_mul` in `SincProductShells.lean`. On the real axis the removable singularity can be cleared without excluding the zero ray: for every  $k \in \mathbb{N}_0$  and  $y \in \mathbb{R}$ ,

$$\begin{aligned} 2^{k(k+1)/2} (\pi|y|)^k \left| \widehat{\text{up}}_{2\pi}(2^k y) \right| \\ = \left( \prod_{j=0}^{k-1} \left| \sin(2^{j+1} \pi y) \right| \right) \left| \widehat{\text{up}}_{2\pi}(y) \right|. \end{aligned} \quad (159)$$

This total form includes  $y = 0$ , and at  $k = 0$  both sides are the same Fourier modulus. It is `norm_rvachevFourier_two_pow_mul_cross` in `RvachevFourierDecay.lean`; its product-level source is `norm_rvachevFourierProduct_two_pow_mul_cross` in `SincProductShells.lean`.  $\square$

**Theorem 5.8** (Total arbitrary-prefix identity and envelope). *Write  $\Phi$  for the standalone sinc product in (138). For every  $K \in \mathbb{N}_0$  and  $x \in \mathbb{R}$ , clearing the first  $K$  sinc denominators gives the total identity*

$$(\pi|x|)^K |\Phi(x)| = 2^{\binom{K}{2}} \left( \prod_{j=0}^{K-1} \left| \sin\left(\frac{\pi x}{2^j}\right) \right| \right) \left| \Phi\left(\frac{x}{2^K}\right) \right|. \quad (160)$$

*In particular,*

$$(\pi|x|)^K |\Phi(x)| \leq 2^{\binom{K}{2}}. \quad (161)$$

*Both formulas include  $x = 0$  and the empty prefix  $K = 0$ . Only after dividing do we require  $x \neq 0$ :*

$$|\Phi(x)| \leq \frac{2^{\binom{K}{2}}}{(\pi|x|)^K} \quad (x \neq 0). \quad (162)$$

*Thus the usual selected-factor constant is  $2^{K(K-1)/2}/\pi^K$ , with no unnecessary assumption  $K \geq 1$ .*

*Formal proof.* The total identity, its total bound, and the divided corollary are respectively `norm_rvachevFourierProduct_prefix_cross`, `norm_rvachevFourierProduct_prefix_cross_le`, and `norm_rvachevFourierProduct_le_prefix_div` in `SincPrefixBound.lean`. The



proof combines the finite shell `rvachevFourierProduct_shell`, the removable real-sinc identity `abs_mul_norm_complexSinc_ofReal`, and the global tail bound `norm_rvachevFourierProduct_le_one`; no convergence argument is repeated.  $\square$

The mean-value input behind shell estimates is more general than the dyadic case. Let  $m_0, m_1, \dots$  be positive integer frequencies satisfying  $2m_j \leq m_{j+1}$ . Then, for every  $n \in \mathbb{N}_0$ ,

$$\int_0^1 \prod_{j=0}^{n-1} \sin^2(\pi m_j t) \, dt = \left(\frac{1}{2}\right)^n, \quad (163)$$

$$\int_0^1 \prod_{j=0}^{n-1} |\sin(\pi m_j t)| \, dt \leq \left(\frac{\sqrt{2}}{2}\right)^n. \quad (164)$$

The exact square mean is `integral_prod_sin_sq` in `LacunaryRieszIntegral.lean`. Applying Cauchy–Schwarz to it gives `integral_prod_abs_sin_le_sqrt_two` in `LacunaryMeanSharp.lean`. The geometric-base and dyadic specializations are `integral_prod_abs_sin_pow_le_sqrt_two` and `integral_prod_abs_sin_two_pow_le_sqrt_two`.

The logarithm of the normalized dyadic sine shell has an exact square mean as well. For every  $n \in \mathbb{N}_0$ , including the empty product  $n = 0$ ,

$$\int_0^1 \log^2 \left| \prod_{j=0}^{n-1} 2 \sin(\pi 2^j t) \right| dt = \frac{\pi^2}{4} n - \frac{\pi^2}{3} (1 - 2^{-n}). \quad (165)$$

The product statement is `integral_sq_log_prod_two_sin_all` in `BirkhoffProductBridge.lean`. Its doubling-cocycle form is `integral_sq_birkhoff_sum_all` in `BirkhoffVariance.lean`, and the finite covariance summation is `variance_closed_form_all` in `VarianceBookkeeping.lean`. The all-index declarations make the zero-length boundary an identity rather than a side condition.

For a quantitative global form, set

$$\kappa_\infty = \frac{3}{2} + \frac{\log(\pi/\sqrt{3})}{\log 2}, \quad E_\kappa(x) = \exp\left(-\frac{(\log x)^2}{2 \log 2}\right) x^{-\kappa} \quad (x > 0). \quad (166)$$

There is a constant  $C > 0$  such that the single two-sided estimate

$$|\widehat{\text{up}}_{2\pi}(x)| \leq C E_{\kappa_\infty}(|x|) \quad (|x| \geq 1) \quad (167)$$

holds on both real tails. This is `norm_rvachevFourier_le_decayGauge_abs` in `RvachevFourierDecay.lean`. On the distinguished dyadic ray the same gauge occurs in an exact identity:

$$|\widehat{\text{up}}_{2\pi}(2^k/3)| = \frac{|\widehat{\text{up}}_{2\pi}(2/3)|}{E_{\kappa_\infty}(2/3)} E_{\kappa_\infty}(2^k/3), \quad (168)$$

namely `norm_rvachevFourier_peak_ray_envelope` in the same module.

The exponent  $\kappa_\infty$  is sharp within this family of gauges. Write  $\Phi$  for the standalone sinc product, which equals  $\widehat{\text{up}}_{2\pi}$  by `rvachevFourier_eq_product`. For every  $\kappa > \kappa_\infty$ ,

$$\frac{|\Phi(2^k/3)|}{E_\kappa(2^k/3)} \longrightarrow +\infty \quad (k \longrightarrow \infty). \quad (169)$$

More explicitly, for every real  $C$  there is a  $k \in \mathbb{N}_0$  such that  $x_k = 2^k/3 \geq 1$  and

$$C E_\kappa(x_k) < |\Phi(x_k)|. \quad (170)$$

Consequently no  $C > 0$  satisfies

$$|\Phi(x)| \leq CE_\kappa(x) \quad \text{for every real } x \geq 1.$$

The limit, explicit peak-ray obstruction, and positive-ray nonexistence theorem are `norm_rvachevFourierProduct_peak_ray_ratio_tendsto_atTop`, `exists_peak_ray_norm_gt_mul_decayGauge`, and `not_exists_norm_rvachevFourierProduct_le_decayGauge_of_kappaInf_lt` in `GlobalDecayEnvelope.lean`.

The transform also has the exact conjugation, parity, and real-axis identities

$$\overline{\widehat{\text{up}}_{2\pi}(z)} = \widehat{\text{up}}_{2\pi}(-\bar{z}), \quad \widehat{\text{up}}_{2\pi}(-z) = \widehat{\text{up}}_{2\pi}(z), \quad \text{Im } \widehat{\text{up}}_{2\pi}(t) = 0 \quad (t \in \mathbb{R}). \quad (171)$$

These are `rvachevFourier_conj` in `FourierAnalytic.lean` and `rvachevFourier_neg`, `rvachevFourier_ofReal_im_eq_zero` in `FourierProduct.lean`.

**Theorem 5.9** (Exact Fourier zero set, positive-real zero-order limits, and canonical product). *The complex zeros of the Fourier transform are precisely the nonzero integers:*

$$\widehat{\text{up}}_{2\pi}(z) = 0 \iff \exists m \in \mathbb{Z} \setminus \{0\} \text{ such that } z = m, \quad z \in \mathbb{C}. \quad (172)$$

At every positive integer  $m$ , its real-axis zero order is exactly  $\nu_2(m) + 1$ , in the following quantitative sense: there is a constant  $C_m > 0$  such that

$$\lim_{\substack{x \rightarrow m \\ x \neq m}} \frac{|\widehat{\text{up}}_{2\pi}(x)|}{|x - m|^{\nu_2(m)+1}} = C_m. \quad (173)$$

The corresponding canonical product records all of these orders at once:

$$\widehat{\text{up}}_{2\pi}(z) = \prod_{r=0}^{\infty} \left(1 - \frac{z^2}{(r+1)^2}\right)^{1+\nu_2(r+1)}. \quad (174)$$

*Formal proof.* The complete complex zero-set equivalence is `rvachevFourier_eq_zero_iff` in `FourierProduct.lean`. The positive-integer limit is `tendsto_norm_rvachevFourier_div_pow_zero_order` in `RvachevFourierDecay.lean`; its limiting constant is exhibited there as an exponential, and is therefore strictly positive. The canonical product is `rvachevFourier_eq_canonical` in the same facade module, built from `rvachevFourierProduct_eq_canonical` in `SincCanonicalProduct.lean`.  $\square$

## 5.2 Geometric reciprocal-Gamma products

The Gamma factorization is most naturally formalized from the entire reciprocal factor

$$R(z) := \frac{1}{\Gamma(1+z)} \quad (175)$$

**Theorem 5.10** (Geometric reciprocal-Gamma products and the dyadic factor). *For a complex contraction  $q$  satisfying  $\|q\| < 1$ , define the locally uniformly convergent product*

$$G_q(z) := \prod_{n=0}^{\infty} R(q^n z) = \prod_{n=0}^{\infty} \frac{1}{\Gamma(1+q^n z)}. \quad (176)$$

*The hypothesis includes  $q = 0$ ; no positivity or reality assumption is made. The reciprocal factor and the geometric product satisfy*

$$R(z)R(-z) = \text{sinc}_{\text{rad}}(\pi z), \quad (177)$$

$$G_q(0) = 1, \quad G_q(z) = R(z)G_q(qz), \quad (178)$$

$$G_q(z) = 0 \iff \exists n, m \in \mathbb{N}_0 : 1 + q^n z = -m. \quad (179)$$

In particular there are no hidden zeros in the infinite product. Define the Gamma-side total meromorphic representative and the matching sinc product by

$$\Gamma_{\langle q \rangle}(z) := G_q(z)^{-1}, \quad \Phi_q(z) := \prod_{n=0}^{\infty} \text{sinc}_{\text{rad}}(\pi q^n z). \quad (180)$$

The product defining  $\Phi_q$  converges locally uniformly on  $\mathbb{C}$ , its factors have the corresponding pointwise HasProd, and  $\Phi_q$  is entire. Then

$$\Gamma_{\langle q \rangle}(z) = \Gamma(1+z)\Gamma_{\langle q \rangle}(qz), \quad (181)$$

$$G_q(z)G_q(-z) = \Phi_q(z). \quad (182)$$

At  $q = 1/2$ , write  $G_{\text{dy}} = G_{1/2}$  and  $\Gamma_{\text{dy}} = G_{\text{dy}}^{-1}$ . The geometric sinc product is exactly the standalone Rvachev product, so for every  $z \in \mathbb{C}$

$$\Phi(z) = G_{\text{dy}}(z)G_{\text{dy}}(-z) = \frac{1}{\Gamma_{\text{dy}}(z)\Gamma_{\text{dy}}(-z)}. \quad (183)$$

Moreover  $G_{\text{dy}}$  vanishes precisely at the negative integers. For every  $n \geq 1$ ,

$$\text{ord}_{z=-n} G_{\text{dy}}(z) = 1 + \nu_2(n), \quad (184)$$

$$\text{mord}_{z=-n} \Gamma_{\text{dy}}(z) = -(1 + \nu_2(n)). \quad (185)$$

Thus the positive pole order of  $\Gamma_{\text{dy}}$  is  $1 + \nu_2(n)$ ; Mathlib records it as the displayed negative meromorphic order.

*Formal proof and exact API.* At formal source checkpoint 71ab6f6728fceb753c88d8b0573077a59acf2682, the reusable convergence engine is `ScaledInfiniteProducts.lean`. Its ten public declarations are `summable_norm_scaled_sub_one`, `hasProdUniformlyOn_scaled`, `multipliableUniformlyOn_scaled`, `hasProdLocallyUniformly_scaled`, `multipliableLocallyUniformly_scaled`, `continuous_tprod_scaled`, `differentiable_tprod_scaled`, `differentiable_tprod_scaled_of_eq_one`, `tprod_scaled_ne_zero`, and `tprod_scaled_eq_zero_iff`. The pointwise summability theorem assumes a normed scalar field, a normed space, a normed-ring target, summability of  $n \mapsto \|a_n\|$ , and  $f(z) - 1 = O(z)$  at zero. The uniform statements take values in a complete commutative normed ring with a normed unit and assume continuity of  $f$ ; the locally uniform statements additionally assume a locally compact domain. The holomorphic statements take values in a complete commutative normed ring with a normed complex-algebra structure and assume complex differentiability, with `differentiable_tprod_scaled_of_eq_one` replacing the explicit big- $O$  hypothesis by  $f(0) = 1$ . The no-hidden-zero statements use a complete commutative normed ring with a normed unit and a multiplicative norm, but need neither continuity of  $f$  nor local compactness of the domain.

The specialization is `GeometricReciprocalGamma.lean`. Its twenty-nine public declarations (six definitions and twenty-three theorems) are `shiftedReciprocalGamma`, `shiftedReciprocalGamma_zero`, `shiftedReciprocalGamma_differentiable`, `shiftedReciprocalGamma_sub_one_isBigO`, `shiftedReciprocalGamma_eq_zero_iff`, `shiftedReciprocalGamma_mul_neg`, `summable_norm_qpow`, `geometricReciprocalGamma`, `geometricReciprocalGammaFactors_multipliable`, `geometricReciprocalGamma_differentiable`, `geometricReciprocalGamma_zero`, `geometricReciprocalGamma_mahler`, `geometricReciprocalGamma_eq_zero_iff`, `geometricGamma`, `geometricGamma_meromorphic`, `geometricGamma_mahler`, `geometricSincProduct`, `hasProdLocallyUniformly_geometricSincProduct`, `geometricSincProductFactors_multipliable`, `hasProd_geometricSincProduct`, `geometricReciprocalGamma_mul_neg`, `geometricSincProduct_differentiable`,

`dyadicReciprocalGamma`, `dyadicGamma`, `dyadicReciprocalGamma_differentiable`, `dyadicReciprocalGamma_zero`, `geometricSincProduct_inv_two`, `dyadicReciprocalGamma_mul_neg`, and `rvachevFourierProduct_eq_one_div_dyadicGamma_mul`. Here  $q \in \mathbb{C}$  and  $\|q\| < 1$  are the exact hypotheses on every generic convergence, holomorphy, Mahler, zero-orbit, meromorphy, and reflection statement; normalization at zero itself needs no contraction hypothesis. In particular, the four added sinc declarations respectively expose the locally uniform product, its pointwise Multipliable and HasProd consequences, and entireness.

At every reciprocal integer scale, `ReciprocalIntegerGammaZeros.lean` exposes four public theorems: `norm_inv_natCast_lt_one`, `geometricReciprocalGamma_inv_natCast_eq_zero_iff`, `geometricReciprocalGamma_inv_natCast_ne_zero_of_nonneg`, and `geometricGamma_inv_natCast_meromorphic`. For  $b \geq 2$ , the entire product  $G_{1/b}$  vanishes exactly at the embedded strictly negative natural numbers, independently of  $b$ , is nonzero at every embedded nonnegative natural number, and its pointwise inverse is meromorphic. This closes the integer-base zero set, but not the multiplicity of those zeros.

Finally, `DyadicGammaOrder.lean` exposes eight public declarations: `dyadicReciprocalGamma_eq_zero_iff`, `dyadicReciprocalGamma_int_ne_zero_of_nonneg`, `dyadicReciprocalGamma_nat_ne_zero`, `dyadicGamma_meromorphic`, `analyticOrderAt_dyadicReciprocalGamma_int_of_neg`, `analyticOrderAt_dyadicReciprocalGamma_neg_nat`, `meromorphicOrderAt_dyadicGamma_int_of_neg`, and `meromorphicOrderAt_dyadicGamma_neg_nat`. The integer forms assume a strictly negative integer center; the natural-number wrappers assume  $n \neq 0$ .

There is an important representation boundary. `Mathlib` assigns `Complex.Gamma` the value zero at a classical Gamma pole, and its inverse also has the totalized value zero there. Accordingly `geometricGamma` and `dyadicGamma` are defined as pointwise inverses of the entire reciprocal products, and a pole is carried by negative `meromorphicOrderAt`, not by an infinite point value. At that checkpoint, the declarations above do not identify these functions with the raw products  $\prod_n \Gamma(1 + q^n z)$  away from their poles, and they assert no arithmetic Weierstrass regrouping, logarithmic or digamma series, Malmsten or heat-trace integral, D-finiteness result, reciprocal-Gamma derivative coefficient for the Thue–Morse tower, or general integer-base collision order.

The post-checkpoint source commit 0ba35abd4 closes the reciprocal-Gamma first-jet and Thue–Morse Gamma-tower tranche in two further modules. `ReciprocalGammaJets.lean` has exactly five public declarations: `deriv_Gamma_inv_neg_nat`, `hasDerivAt_Gamma_inv_neg_nat`, `hasDerivAt_Gamma_inv_zero`, `analyticOrderAt_Gamma_inv_neg_nat`, and `tendsto_Gamma_inv_div_add_nat`. For every  $r \in \mathbb{N}_0$ , they identify

$$(\Gamma^{-1})'(-r) = (-1)^r r!, \quad \text{ord}_{-r} \Gamma^{-1} = 1, \quad \frac{\Gamma(s)^{-1}}{s+r} \longrightarrow (-1)^r r! \quad (s \rightarrow -r, s \neq -r),$$

with `hasDerivAt_Gamma_inv_zero` retaining the origin compatibility name. These are statements about the entire reciprocal function  $s \mapsto \Gamma(s)^{-1}$ : they do not assign a value or derivative to the raw Gamma function at one of its poles.

At the same commit, `ThueMorseGammaTower.lean` has exactly eight public declarations: `hasDerivAt_dirichletMellinContinuation_neg_nat`, `deriv_dirichletMellinContinuation_neg_nat`, `thueMorseGammaLog`, `thueMorseGammaTower`, `thueMorseGammaLog_eq_mellin`, `thueMorseGammaLog_eq_integral`, `thueMorseGammaLog_dyadic`, and `thueMorseGammaTower_dyadic`. The first two give, for  $a > 0$  and  $r \in \mathbb{N}_0$ ,

$$\tilde{D}'_a(-r) = (-1)^r r! \text{Mellin}(t \mapsto e^{-at} \mathcal{E}(t))(-r).$$

The two definitions are total for every real  $a$ ; their Mellin, integral, and dyadic identification theorems assume exactly  $a > 0$ . In particular `thueMorseGammaLog` is the chosen coordinate  $L_r(a) := \tilde{D}'_a(-r)$ , not a proved identity with `Complex.log` of `thueMorseGammaTower`; the latter is defined as

$T_r(a) := \exp L_r(a)$ . The formal laws are

$$L_r(a) = (-1)^r r! \int_0^\infty t^{-r-1} e^{-at} \mathcal{E}(t) dt, \quad L_r(a) = 2^r (L_r(a/2) - L_r((a+1)/2)),$$

and

$$T_r(a) = \left( \frac{T_r(a/2)}{T_r((a+1)/2)} \right)^{2^r}.$$

The current integrated tree adds the ninth public tower declaration of `Real_exp_mpLimit_eq_gammaTower_div`: for  $a, b > 0$ , the complex embedding of `Real.exp` of `mpLimit` at  $(a, b)$  is  $T_0(b)/T_0(a)$ . The source-only leaf `ThueMorseGammaTowerDifferential.lean` has no definitions and exactly three public theorems: `hasDerivAt_mellin_mellinKernel_parameter`, `hasDerivAt_thueMorseGammaLog_succ`, and `iteratedDeriv_thueMorseGammaLog`. Under exactly  $a > 0$ , they prove the arbitrary-complex-spectral parameter shift  $\frac{d}{da} M_a(s) = -M_a(s+1)$ , the successor law  $L'_{r+1}(a) = (r+1)L_r(a)$ , and

$$\frac{d^k}{da^k} L_r(a) = r^{\underline{k}} L_{r-k}(a), \quad k \leq r.$$

Together with the existing Mellin, integral, dyadic, and ratio declarations, this makes `p2:thm:gamma-tower` Exact on its positive-parameter domain. Its displayed logarithm is the chosen `thueMorseGammaLog` coordinate, not a proved branch or `principal-Complex.log` identity; no nonpositive-parameter differential law is claimed. The raw Gamma product, canonical regrouping, logarithmic/digamma expansions, Gamma-product Malmsten and heat-trace representations, D-finiteness, and general-base collision order listed above remain separate frontiers.  $\square$

**Theorem 5.11** (Complex Pochhammer factorization of geometric sinc products). *For  $a, q \in \mathbb{C}$ , let the total complex infinite  $q$ -Pochhammer symbol be*

$$(a; q)_{\infty, \mathbb{C}} := \prod_{j=0}^{\infty} (1 - aq^j). \quad (186)$$

*If  $\|q\| < 1$ , this factor family is multipliable and its finite prefixes converge to  $(a; q)_{\infty, \mathbb{C}}$ , with no restriction on  $a$ . For every  $q, z \in \mathbb{C}$  with  $\|q\| < 1$ , put  $S_q(z) := \text{geometricSincProduct}(q, z)$ . Then, globally in  $z$ ,*

$$S_q(z) = \prod_{k=0}^{\infty} \left( \frac{z^2}{(k+1)^2}; q^2 \right)_{\infty, \mathbb{C}}. \quad (187)$$

*The statement includes  $q = 0$  and every zero of an individual factor. For every  $z \in \mathbb{C}$ , without a nonvanishing hypothesis,*

$$\Phi(z) = \prod_{k=0}^{\infty} \left( \frac{z^2}{(k+1)^2}; \frac{1}{4} \right)_{\infty, \mathbb{C}}. \quad (188)$$

*The same right-hand side equals  $\widehat{\text{up}}_{2\pi}(z)$  for every bounded Fabius witness satisfying the Fabius equations. Equation (188) is the  $q = 1/2$ , nome-1/4 specialization of (187).*

*Human proof, formal crosswalk, and exact boundary.* The Euler product for the normalized sine gives, at each geometric scale  $h$ ,

$$\text{sinc}_{\text{rad}}(\pi q^h z) = \prod_{k=0}^{\infty} \left( 1 - \frac{q^{2h} z^2}{(k+1)^2} \right).$$

The perturbations of these factors are absolutely summable on  $\mathbb{N}_0 \times \mathbb{N}_0$ , because

$$\sum_{h,k \geq 0} \left\| \frac{q^{2h} z^2}{(k+1)^2} \right\| = \|z\|^2 \left( \sum_{h \geq 0} \|q\|^{2h} \right) \left( \sum_{k \geq 0} \frac{1}{(k+1)^2} \right) < \infty.$$

Hence the double product is multipliable: first flatten the scale-indexed Euler products to a product on pairs, then swap the two natural-number indices. For a fixed spectral index  $k$ , the resulting inner product is

$$\prod_{h=0}^{\infty} \left( 1 - \frac{z^2}{(k+1)^2} (q^2)^h \right) = \left( \frac{z^2}{(k+1)^2}; q^2 \right)_{\infty, \mathbb{C}},$$

which proves (187). Absolute multipliability is the point that permits the index exchange even when a factor vanishes; no division or cancellation is used. Substituting  $q = 1/2$  gives (188), and the established equality between the Rvachev product and the Fourier transform gives the final assertion.

`RvachevPochhammerFactorization.lean` has one public definition, `complexQPochhammerInf`, and exactly ten public theorems: `complexQPochhammerInf_eq_tprod`, `complexQPochhammerInf_eq_qPochhammerInfIn`, `multipliable_one_sub_mul_pow_complex`, `hasProd_complexQPochhammerInf`, `tendsto_finiteQPochhammerIn_complex`, `summable_norm_sineTerm_qpows_pair`, `geometricSincProduct_eq_tprod_pair`, `geometricSincProduct_eq_tprod_complexQPochhammerInf`, `rvachevFourierProduct_eq_tprod_complexQPochhammerInf`, and `rvachevFourier_eq_tprod_complexQPochhammerInf`. The bridge `complexQPochhammerInf_eq_qPochhammerInfIn` states, for every  $a, q \in \mathbb{C}$  and with no hypothesis, that the historical complex name is definitionally equal to the generic symbol. Because both sides are the same total Mathlib `tprod`, this equality also holds in the nonmultipliable junk-value regime; it asserts no convergence there. The three symbol-convergence declarations after the bridge, and the three new geometric-sinc declarations, require exactly  $\|q\| < 1$ . The latter three formalize, in order, the displayed absolute-summability argument, the flattened paired Euler product, and (187). The final two declarations fix the dyadic scale and hence the Pochhammer nome  $1/4$ ; their complex argument is unrestricted, and the Fourier-transform form assumes exactly a bounded witness and `IsFabius`. The downstream zero-definition, three-theorem module `GeometricPochhammerNormalConvergence.lean` upgrades the outer product itself to locally uniform convergence on the whole complex plane. Its exact surface is `hasProdLocallyUniformly_geometricSincProduct_complexQPochhammerInf`, `hasProdLocallyUniformly_rvachevFourierProduct_complexQPochhammerInf`, and `hasProdLocallyUniformly_rvachevFourier_complexQPochhammerInf`. The first assumes exactly  $\|q\| < 1$ , includes  $q = 0$ , and has limit  $S_q$ ; the other two are the nome- $1/4$  Rvachev-product and bounded-Fabius specializations. Thus the locally-uniform/normal-convergence clause of the outer spectral product is exact. The compound spectral theorem nevertheless remains partial: no declaration packages the centered characteristic-function/MGF form, the outside-disk reciprocal formula, its pole divisor, or the zero–pole exchange. The next theorem records the downstream fixed-nome analytic and zero-set API, including its division-free  $q = 0$  boundary and the reciprocal-power description when  $q \neq 0$ .  $\square$

**Theorem 5.12** (Entire complex  $q$ -Pochhammer factors and their simple zeros). *Let  $q \in \mathbb{C}$  satisfy  $\|q\| < 1$ . Then the finite products*

$$a \mapsto \prod_{j=0}^{n-1} (1 - aq^j)$$

*converge locally uniformly on  $\mathbb{C}$  to  $a \mapsto (a; q)_{\infty, \mathbb{C}}$ , and this limit is entire. Moreover, for every  $a \in \mathbb{C}$ ,*

$$(a; q)_{\infty, \mathbb{C}} = 0 \iff \exists j \in \mathbb{N}_0, 1 - aq^j = 0, \quad (189)$$

and every zero has analytic order one. If  $q \neq 0$ , the right side of (189) is equivalently  $a = q^{-j}$  for some  $j$ ; if  $q = 0$ , the unique zero is  $a = 1$ .

*Human proof and exhaustive Lean crosswalk.* On a compact set  $K \subset \mathbb{C}$ , put  $M = \max_{a \in K} \|a\|$ . The perturbations from one obey

$$\sum_{j=0}^{\infty} \sup_{a \in K} \|aq^j\| \leq M \sum_{j=0}^{\infty} \|q\|^j < \infty.$$

The uniform infinite-product criterion therefore gives convergence uniformly on every compact set, and the Weierstrass theorem makes the limit entire. An absolutely convergent product of nonzero factors is nonzero, so the product vanishes exactly when one displayed factor does. Two distinct factors cannot vanish at the same  $a$ : for  $q \neq 0$  that would force two distinct powers of a strict contraction to agree, while for  $q = 0$  only the zeroth factor can vanish. Factoring out that unique linear term leaves a holomorphic cofactor which is nonzero at  $a$ , proving that the zero is simple.

The downstream module `QPochhammerEntire.lean` introduces no public definition and has exactly five public theorems: `hasProdLocallyUniformly_complexQPochhammerInf`, `complexQPochhammerInf_differentiable`, `complexQPochhammerInf_eq_zero_iff`, `complexQPochhammerInf_eq_zero_iff_eq_inv_pow`, and `analyticOrderAt_complexQPochhammerInf_of_eq_zero`. They formalize, in order, the locally uniform product, entireness, the division-free equivalence (189), its reciprocal-power form under exactly  $q \neq 0$ , analytic order one for the historical complex wrapper. All five assume exactly  $\|q\| < 1$ ; the analytic-order theorem adds only the displayed zero hypothesis and therefore retains the case  $q = 0$ . The corresponding generic name theorem `analyticOrderAt_qPochhammerInfIn_of_eq_zero` belongs to `QPochhammerInfinite.lean`, not to this compatibility leaf. The factor form is deliberately stronger than a reciprocal-power listing because it remains valid at  $q = 0$ . The module does not treat  $\|q\| \geq 1$ , joint holomorphy in  $q$ , a closed derivative formula, growth order, or reciprocal meromorphic continuation. Locally uniform convergence of the outer rearranged product in (187) is supplied by the separate three-theorem module above.  $\square$

**Generic infinite-symbol and dissection API.** The later modules `QPochhammerDissection.lean` and `QPochhammerInfinite.lean` expose a second public name for the same displayed factor product, but in the greatest algebraic generality used by each result. They do not replace `complexQPochhammerInf` or any of the five `QPochhammerEntire.lean` theorems above. The finite dissection module has no public definition and exactly two theorems. Over an arbitrary commutative ring, `finiteQPochhammerIn_dissection` proves

$$(a; q)_{rn} = \prod_{s < r} (aq^s; q^r)_n$$

for all natural  $r, n$ , including the empty boundaries, and `finiteQPochhammerIn_dissection_remainder` proves

$$(a; q)_{rn+u} = \prod_{s < u} (aq^s; q^r)_{n+1} \prod_{u \leq s < r} (aq^s; q^r)_n$$

under exactly  $u \leq r$ , hence also at  $u = r$ .

The exhaustive public surface of `QPochhammerInfinite.lean` is one definition and twenty-nine theorems. For a commutative ring with an arbitrary topology, `qPochhammerInfIn` is the total `Mathlib.tprod`  $\prod'_{j \geq 0} (1 - aq^j)$ , and `qPochhammerInfIn_eq_tprod` is its defining equality; as usual the totalized value outside the multipliable range is `Mathlib.junk` value and carries no convergence claim. In a normed ring with  $\|1\| = 1$ , the three elementary inputs are `summable_norm_mul_pow`, `one_sub_ne_zero_of_norm_lt_one`, and `norm_mul_pow_self_lt_one`. The finite self-product theorem `finiteQPochhammerIn_self_ne_zero` additionally uses a commutative ring with no zero divisors.



In every complete normed commutative ring with  $\|1\| = 1$ , and under exactly  $\|q\| < 1$ , the seven convergence and factorization declarations are `multipliable_one_sub_mul_pow_of_norm_lt_one`, `hasProd_qPochhammerInfIn`, `tendsto_finiteQPochhammerIn_qPochhammerInfIn`, `qPochhammerInfIn_eq_finite_mul_shift`, `qPochhammerInfIn_succ_shift`, `qPochhammerInfIn_eq_factor_mul`, and `qPochhammerInfIn_dissection`. They give, respectively, multipliability, the named `HasProd`, convergence of finite symbols, the arbitrary-prefix concatenation, the first-factor shift, removal of one factor, and

$$(a; q)_\infty = \prod_{s < r} (aq^s; q^r)_\infty \quad (0 < r).$$

Only the last declaration assumes  $0 < r$ . If the norm is also multiplicative, `qPochhammerInfIn_ne_zero`, `qPochhammerInfIn_eq_zero_iff`, and `qPochhammerInfIn_self_ne_zero` prove, without division, that there are no hidden zeros, that  $(a; q)_\infty = 0$  exactly when  $aq^j = 1$  for some  $j$ , and that  $(q; q)_\infty \neq 0$ .

The field bridge is deliberately split at its true hypotheses. `qPochhammerInfIn_eq_tprod_smul`, and `isBigO_one_sub_sub_one` need only a normed field, while `summable_norm_pow_of_norm_lt_one` additionally assumes  $\|q\| < 1$ ; `differentiable_finiteQPochhammerIn` needs only a nontrivially normed field. In a complete normed field under  $\|q\| < 1$ , `qPochhammerInfIn_eq_zero_iff_exists_inv_pow` enumerates the zeros as  $q^{-j}$  under the additional hypothesis  $q \neq 0$ . The two declarations `hasProdLocallyUniformly_qPochhammerInfIn` and `continuous_qPochhammerInfIn` instead require the same strict contraction together with local compactness; they assert local uniform convergence and continuity, not complex entireness. The algebraic cofactor identity `pow_sq_mul_finiteQPochhammerIn_inv_pow_self` holds in every field under  $q \neq 0$ .

Finally, the six complex declarations are `differentiable_qPochhammerInfIn`, `hasDerivAt_qPochhammerInfIn_of_mul_pow_eq_one`, `hasDerivAt_qPochhammerInfIn_inv_pow`, and `deriv_qPochhammerInfIn_inv_pow_ne_zero`, together with `deriv_qPochhammerInfIn_ne_zero_of_mul_pow_eq_one` and `analyticOrderAt_qPochhammerInfIn_of_eq_zero`. The first proves entireness for  $\|q\| < 1$ ; the second gives the derivative after removing a factor from the raw hypothesis  $a_0 q^j = 1$ , including the possible  $q = 0, j = 0$  case, and the fifth proves that raw-factor derivative is nonzero. The inverse-power pair assumes  $q \neq 0$  and together gives

$$\left. \frac{d}{da} (a; q)_\infty \right|_{a=q^{-j}} = (-1)^{j+1} (q^{\binom{j}{2}})^{-1} (q; q)_j (q; q)_\infty \neq 0.$$

The coefficient theorem proves nonvanishing of the displayed expression; the preceding `HasDerivAt` theorem identifies it with the derivative. The sixth theorem gives analytic order one at every zero, including the degenerate nome  $q = 0$ . The same conclusion for the separately named complex wrapper is `analyticOrderAt_complexQPochhammerInf_of_eq_zero` in `QPochhammerEntire.lean`; conversely that five-theorem wrapper supplies none of the generic ring-valued concatenation or dissection laws.

**Latest finite and infinite q-series tranche.** Six further facade-reachable modules extend this layer without changing the hypotheses above. `GaussianBinomialContinuity.lean` has no definitions and exactly three theorems, `continuous_gaussianBinomial`, `tendsto_gaussianBinomial_nhds_one`, and `gaussianBinomial_eq_finiteQPochhammerIn_div`, for continuity, the classical  $q \rightarrow 1$  limit, and the finite-Pochhammer quotient formula. `QPascalSummation.lean` has no definitions and exactly four theorems, `sum_gaussianBinomial_succ_mul`, `sum_gaussianBinomial_succ_mul'`, `Commute.gaussianBinomial_left`, and `Commute.gaussianBinomial_right`. `QuantumBinomial.lean` has no definitions and exactly the two noncommutative theorems `quantumPlane_mul_pow` and `quantum_binomial`.

The remaining three modules carry the longer analytic inventories. `QBinomialTheoremInfinite.lean` has one definition and twenty-two theorems; its public surface comprises `gaussianMajorant`, `hasSum_of_tendsto_of_dominated`, `one_le_qPochhammerInfIn_neg`, `finiteQPochhammerIn_neg_le_qPochhammerInfIn`, `qPochhammerInfIn_nonneg`, `qPochhammerInfIn_le_finiteQPochhammerIn`, `qPochhammerInfIn_pos_of_lt_one`, `qPochhammerInfIn_zero_left`, `norm_finiteQPochhammerIn_le_neg_norm`, `finiteQPochhammerIn_norm_le_norm`, `norm_finiteQPochhammerIn_le`, `qPochhammerInfIn_norm_le_norm_finiteQPochhammerIn`, `finiteQPochhammerIn_ne_zero_of_norm_lt_one`, `gaussianBinomial_eq_div`, `gaussianMajorant_nonneg`, `norm_gaussianBinomial_le`, `tendsto_finiteQPochhammerIn_pow_atTop`, `tendsto_gaussianBinomial_atTop`, `summable_pow_choose_two_mul_pow`, `hasSum_euler_product`, `qPochhammerInfIn_ne_zero_of_norm_lt_one`, `hasSum_qBinomial_theorem`, and `hasSum_euler_reciprocal`.

The subsequent zero-definition/ten-theorem `GaussianBinomialFixedColumnRate.lean` surface is `norm_finiteQPochhammerIn_pow_sub_one_le_exp`, `norm_finiteQPochhammerIn_pow_sub_one_le`, `norm_finiteQPochhammerIn_self_mul_gaussianBinomial_sub_one_le`, `norm_gaussianBinomial_sub_inv_finiteQPochhammerIn_le`, `norm_gaussianBinomial_add_sub_inv_finiteQPochhammerIn_le`, `tendsto_gaussianBinomial_add_atTop`, `gaussianBinomial_fixedColumn_relativeError_isBigO`, `gaussianBinomial_shifted_fixedColumn_relativeError_isBigO`, `gaussianBinomial_fixedColumn_error_isBigO`, and `gaussianBinomial_shifted_fixedColumn_error_isBigO`. In a normed commutative ring with normalized multiplicative norm,  $\|q\| \leq 1$  gives

$$\|(q^m; q)_k - 1\| \leq \exp(k\|q\|^m) - 1 \leq ke^k\|q\|^m,$$

and, when  $k \leq n$ , the denominator-free relative estimate

$$\left\| (q; q)_k \begin{bmatrix} n \\ k \end{bmatrix}_q - 1 \right\| \leq ke^k\|q\|^{n-k+1}.$$

This ring-level statement remains meaningful at roots of unity. Over an arbitrary normed field,  $\|q\| < 1$  gives the fixed and shifted nonasymptotic additive estimates, the shifted limit, and relative and additive `IsBigO` bounds with rates  $q^{n-k+1}$  and  $q^{n+1}$ . No completeness or  $q \neq 0$  hypothesis is used, so  $q = 0$  is included; the constant is elementary rather than asserted sharp. Together with `tendsto_gaussianBinomial_atTop`, the shifted limit and the two relative error bounds prove all four clauses of the `q-monograph`'s `thm:fixed-column-limit`; the additive pair are companion estimates.

The source-only `GaussianBinomialGreaterOneAsymptotics.lean` leaf has no definitions and exactly two public theorems, `gaussianBinomial_gt_one_fixedColumn_relativeError_isBigO` and `gaussianBinomial_gt_one_central_isEquivalent`. For real  $q > 1$  they give, respectively,

$$(q^{-1}; q^{-1})_k q^{-k(n-k)} \begin{bmatrix} n \\ k \end{bmatrix}_q - 1 = O((q^{-1})^{n-k+1}), \quad \begin{bmatrix} 2m \\ m \end{bmatrix}_q \sim q^{m^2} (q^{-1}; q^{-1})_\infty^{-1}.$$

Natural subtraction is total in the first theorem, and `gaussianBinomial_inv` is invoked only eventually once  $k \leq n$ . These three declarations make `cor:qgreaterone Exact` with the printed normalization. No shifted-central, non-real, or wider-nome theorem is asserted.

`JacobiTripleProduct.lean` has two definitions and twenty-five theorems. Its exact public inventory is `thetaExponent`, `pentagonalExponent`, `two_dvd_mul_sub_one`, `mul_sub_one_nonneg`, `two_mul_thetaExponent`, `thetaExponent_natCast`, `thetaExponent_neg_natCast`, `choose_two_add_choose_two_succ_eq`, `sum_mul_sum_eq_sum_antidiagonal`, `thetaCoeff_pair_eq`, `sum_antidiagonal_thetaCoeff`, `finite_triple_product_poly`,

pow\_mul\_finiteQPochhammerIn\_div, neg\_one\_zpow\_natCast\_sub, finite\_triple\_product, norm\_gaussianBinomialInt\_le, tendsto\_gaussianBinomialInt\_central, summable\_pow\_thetaExponent\_mul\_zpow, hasSum\_jacobi\_triple\_product, hasSum\_jacobi\_triple\_product', mul\_three\_mul\_sub\_one\_nonneg, two\_dvd\_mul\_three\_mul\_sub\_one, two\_mul\_pentagonalExponent, pentagonalExponent\_eq, pentagonalExponent\_pos, hasSum\_pentagonal, and hasSum\_pentagonal\_pairs.

Finally, `RogersSzegoPolynomial.lean` has one definition and nine theorems: `rogersSzego`, `rogersSzego_zero`, `rogersSzego_one`, `rogersSzego_succ`, `one_sub_pow_succ_mul_gaussianBinomial_succ_succ`, `rogersSzego_dilation`, `rogersSzego_three_term`, `summable_norm_pow_div_finiteQPochhammerIn_self`, `sum_antidiagonal_euler_mul_euler`, and `hasSum_rogersSzego_generating`. Thus the six modules contribute four definitions and sixty-five theorems, sixty-nine public declarations in total.

### 5.3 Polynomial $q$ -calculus, finite parameter derivatives, Lambert logarithms, and the real $q$ -gamma function

The coefficientwise Jackson derivative is formalized without division. In a semiring put

$$[n]_q = \sum_{i=0}^{n-1} q^i.$$

Over a commutative semiring, the linear map on  $R[X]$  is

$$D_q \left( \sum_n a_n X^n \right) = \sum_n a_n [n]_q X^{n-1}.$$

It has the expected monomial and constant values, the scaling law

$$D_q(p(c \cdot)) = c(D_q p)(c \cdot),$$

and both product rules

$$D_q(fg) = (D_q f)g + f(q \cdot)D_q g = fD_q g + g(q \cdot)D_q f.$$

Over a commutative ring it additionally satisfies the division-free characterization

$$(D_q p)(x)(1 - q)x = p(x) - p(qx).$$

No restriction is imposed on  $q$ , so the formulas include  $q = 0$  and  $q = 1$ . Iteration over a commutative semiring gives, for every  $n \in \mathbb{N}_0$ ,

$$D_q^n(fg) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (D_q^k f)(q^{n-k} \cdot) D_q^{n-k} g.$$

This is finite polynomial algebra, not a derivative theory for analytic functions and not a  $q \rightarrow 1$  limiting theorem.

The exhaustive `PolynomialQDerivative.lean` surface is two definitions, `qInt` and `qDerivative`, and seventeen theorems: `qInt_zero`, `qInt_one`, `qInt_succ`, `qInt_succ'`, `qInt_add`, `qInt_one_left`, `one_sub_mul_qInt`, `qDerivative_apply`, `qDerivative_monomial`, `qDerivative_C`, `qDerivative_X_pow`, `qDerivative_X`, `qDerivative_C_mul_X_pow`, `eval_qDerivative_mul`, `qDerivative_comp_C_mul_X`, `qDerivative_mul`, and `qDerivative_mul'`. The  $q$ -integer definition and its first six laws use only a semiring; `one_sub_mul_qInt` and `eval_qDerivative_mul` use a commutative ring; all other derivative declarations use a

commutative semiring. The exhaustive `PolynomialQLeibniz.lean` surface has no public definition and four theorems, all over a commutative semiring: `comp_C_mul_X_comp_C_mul_X`, `qDerivative_C_mul`, `qDerivative_iterate_comp_C_mul_X`, and `qDerivative_iterate_mul`. They include iteration order zero and assume neither division nor a nonzero nome.

For a nontrivially normed field, finite  $q$ -Pochhammer products also have a parameter derivative for arbitrary  $q, a$  and  $n$ :

$$\frac{d}{da}(a; q)_n = - \sum_{j < n} q^j \prod_{\substack{i < n \\ i \neq j}} (1 - aq^i).$$

If every factor  $1 - aq^j$ ,  $j < n$ , is nonzero, this is equivalently

$$\frac{d}{da}(a; q)_n = -(a; q)_n \sum_{j < n} \frac{q^j}{1 - aq^j}.$$

Composition with a pointwise witness `HasDerivAt f f' x` supplies the corresponding chain rule. These are exactly the three public theorems of `QPochhammerDerivative.lean`: `hasDerivAt_finiteQPochhammerIn`, `hasDerivAt_finiteQPochhammerIn_of_ne_zero`, and `hasDerivAt_finiteQPochhammerIn_comp`. The first has no nonvanishing hypothesis; the second has exactly the factorwise hypothesis just displayed; the third assumes exactly the supplied derivative of  $f$ . This leaf makes no assertion about differentiating the infinite product.

For  $a, q \in \mathbb{C}$  with exactly  $\|a\| < 1$  and  $\|q\| < 1$ , the Lambert series is absolutely summable and

$$\sum_{m=0}^{\infty} \frac{a^{m+1}}{(m+1)(1-q^{m+1})} = - \sum_{j=0}^{\infty} \log(1 - aq^j), \quad \exp\left(- \sum_{m=0}^{\infty} \frac{a^{m+1}}{(m+1)(1-q^{m+1})}\right) = (a; q)_{\infty}.$$

Here each factor uses the principal complex logarithm, whereas the exponentiated conclusion is branch-free. The exhaustive `LambertSeriesLog.lean` surface has no definition and four theorems: `one_le_norm_natCast_add_one`, `summable_lambert_series`, `hasSum_lambert_log_complex`, and `exp_neg_tsum_lambert_eq_qPochhammerInfIn`. The first is an unconditional natural-cast norm bound; the other three have precisely the two strict norm hypotheses above. They include  $a = 0$  and  $q = 0$ , but prove no norm-one boundary, principal-log identity for  $(a; q)_{\infty}$  itself, analytic continuation, or differentiated Lambert series.

Finally, the total real definitions

$$\Gamma_q(x) = \frac{(q; q)_{\infty}}{(q^x; q)_{\infty}} (1 - q)^{1-x}, \quad [x]_q = \frac{1 - q^x}{1 - q}$$

are `qGamma` and `qNumber`. For  $0 < q < 1$ , the formalized API gives positivity for  $x > 0$ ,  $\Gamma_q(1) = 1$ , the recurrence  $\Gamma_q(x+1) = [x]_q \Gamma_q(x)$  for  $x > 0$ , and

$$\Gamma_q(n+1) = \prod_{j=0}^{n-1} [j+1]_q.$$

For natural  $n$ ,  $[n]_q = \text{qInt } q \ n$  needs only  $q \neq 1$ . For every real  $x$ , the totalized algebraic reflection identity assumes only  $q < 1$ :

$$\Gamma_q(x) \Gamma_q(1-x) = \frac{(1-q)(q; q)_{\infty}^2}{(q^x; q)_{\infty} (q^{1-x}; q)_{\infty}}.$$

Under  $0 < q < 1$  and  $0 < x < 1$ , Jacobi's product supplies the exact `HasSum`

$$\sum_{n \in \mathbb{Z}} (-1)^n q^{n(n-1)/2} (q^x)^n = \frac{(1-q)(q; q)_{\infty}^3}{\Gamma_q(x) \Gamma_q(1-x)}.$$

The exhaustive `QGamma.lean` surface is the two definitions above and the ten theorems `norm_1`, `t_one_of_pos_of_lt_one`, `qPochhammerInfIn_rpow_pos`, `qPochhammerInfIn_self_pos`, `qGamma_pos`, `qGamma_one`, `qGamma_add_one`, `qGamma_nat_succ`, `qNumber_natCast`, `qGamma_mul_qGamma_one_sub`, and `hasSum_theta_qGamma_reflection`. This is a real API. It does not prove complex continuation, a pole classification, the  $q \rightarrow 1$  classical-gamma limit, log-convexity, a uniqueness characterization, or a q-digamma formula.

The exponent in the canonical product has a finite, base-generic double-counting origin that is useful independently of sinc products. For  $b, N, J \in \mathbb{N}_0$ , put

$$c_J(b, m) := \#\{j \in \{0, \dots, J\} : b^j \mid m\}.$$

If  $q > 0$ , multiplication by  $q$  identifies the positive multiples of  $q$  not exceeding  $N$  with  $1, \dots, \lfloor N/q \rfloor$ . Consequently, for every function  $f$  into a commutative monoid,

$$\prod_{1 \leq n \leq \lfloor N/q \rfloor} f(qn) = \prod_{\substack{1 \leq m \leq N \\ q \mid m}} f(m), \quad (190)$$

$$\prod_{j=0}^J \prod_{1 \leq n \leq \lfloor N/b^j \rfloor} f(b^j n) = \prod_{m=1}^N f(m)^{c_J(b, m)} \quad (b > 0). \quad (191)$$

For  $b > 1$  and  $m \neq 0$ , if the displayed layer range reaches the multiplicity, that is, if  $\text{multiplicity}(b, m) \leq J$ , then

$$c_J(b, m) = \text{multiplicity}(b, m) + 1. \quad (192)$$

Thus, under the explicit hypothesis  $\text{multiplicity}(b, m) \leq J$  for every  $1 \leq m \leq N$ ,

$$\prod_{j=0}^J \prod_{1 \leq n \leq \lfloor N/b^j \rfloor} f(b^j n) = \prod_{m=1}^N f(m)^{\text{multiplicity}(b, m) + 1}. \quad (193)$$

The same three identities hold in every additive commutative monoid, with products and powers replaced by sums and natural scalar multiples. In Lean, the underlying finite-set equality is `filter_dvd_eq_image`; (190) and its additive companion are `prod_multiples_eq_prod_filter` and `sum_multiples_eq_sum_filter`; (191) and its companion are `prod_layers_eq_prod_pow_card` and `sum_layers_eq_sum_nsmul_card`; and (192)–(193) and the final additive companion are `card_filter_pow_dvd`, `prod_layers_eq_prod_pow_multiplicity`, and `sum_layers_eq_sum_nsmul_multiplicity`, all in `BaseLayerRegrouping.lean`. These are finite algebraic identities. In particular they assert no infinite-product convergence, and the sufficient layer-coverage condition must be supplied explicitly rather than inferred by the API from a bound such as  $N < b^{J+1}$ .

## 5.4 Total local factorization and exact complex zero jets

The complex sinc identities needed at an integer center are most useful after their removable denominators have been cleared. With the convention  $\text{sinc}_{\text{rad}} z = \sin z/z$ , continuously extended at zero, one has for every  $z \in \mathbb{C}$ , every  $n \in \mathbb{N}_0$ , and every  $a \in \mathbb{C} \setminus \{0\}$ ,

$$z \text{sinc}_{\text{rad}}(\pi z) = \frac{\sin(\pi z)}{\pi}, \quad (194)$$

$$(n + z) \text{sinc}_{\text{rad}}(\pi(n + z)) = (-1)^n z \text{sinc}_{\text{rad}}(\pi z), \quad (195)$$

$$(an + z) \text{sinc}_{\text{rad}}\left(\frac{\pi(an + z)}{a}\right) = (-1)^n z \text{sinc}_{\text{rad}}\left(\frac{\pi z}{a}\right). \quad (196)$$

Unlike identities obtained by dividing through by  $z$  or  $n + z$ , these equalities retain the points where either factor vanishes. They are respectively `mul_complexSinc_pi`, `mul_complexSinc_pi_add_nat`, and `mul_complexSinc_pi_div_add_nat` in `IntegerZeroLocalFactorization.lean`; the last theorem allows an arbitrary nonzero *complex* scale, with no order or reality hypothesis.

Here is the reusable all-order differential principle. Let  $\mathbb{K}$  be any nontrivially normed field, let  $E$  be a normed  $\mathbb{K}$ -space, and let  $U : \mathbb{K} \rightarrow E$  be  $C^k$  at zero. For every  $n, k \in \mathbb{N}_0$ ,

$$D^k(z \mapsto z^n \cdot U(z))(0) = \begin{cases} \left(\binom{k}{n} : \mathbb{K}\right)(n! : \mathbb{K}) \cdot D^{k-n}U(0), & n \leq k, \\ 0, & k < n, \end{cases} \quad (197)$$

where  $\cdot$  denotes scalar action on  $E$ . This is the vector-valued theorem `iteratedDeriv_pow_smul_zero_eq_ite` in `LeadingJet.lean`. At the index  $k = n + r$ , its conditional disappears:

$$D^{n+r}(z \mapsto z^n \cdot U(z))(0) = \left(\binom{n+r}{n} : \mathbb{K}\right)(n! : \mathbb{K}) \cdot D^rU(0). \quad (198)$$

The vector and scalar forms of (198) are `iteratedDeriv_pow_smul_zero_add` and `iteratedDeriv_pow_mul_zero_add`. If a scalar-valued function  $f$  agrees with  $z^n U(z)$  merely on a neighborhood of zero, the same formula is `iteratedDeriv_eq_choose_factorial_mul_of_eventuallyEq_pow_mul`; no separate regularity hypothesis on  $f$  is needed. At an arbitrary center  $z_0$ , a vector-valued germ  $f(z) = (z - z_0)^n \cdot U(z)$  has the all-index form `iteratedDeriv_eq_ite_of_eventuallyEq_sub_pow_smul` and the  $n + r$  form `iteratedDeriv_eq_choose_factorial_smul_of_eventuallyEq_sub_pow_smul`; the scalar specialization is `iteratedDeriv_eq_choose_factorial_mul_of_eventuallyEq_sub_pow_mul`.

The coefficient is deliberately the product  $\binom{n+r}{n}n!$ , with both natural numbers cast into  $\mathbb{K}$ , not the quotient  $(n+r)!/r!$ . Thus none of these theorems requires factorial division, characteristic zero, completeness, or analyticity. In positive characteristic the displayed scalar can vanish, so the generic statement identifies the first *potentially* nonzero jet rather than asserting nonvanishing. Setting  $r = 0$  recovers the compatibility formula

$$\left. \frac{d^n}{dz^n} (z^n U(z)) \right|_{z=0} = n! U(0), \quad (199)$$

exported as `iteratedDeriv_pow_mul_zero`; its zero-germ companion is `iteratedDeriv_eq_factorial_mul_of_eventuallyEq_pow_mul`.

We now apply the total sinc identities to the standalone product  $\Phi = \text{rvachevFourierProduct}$ . Put

$$P_d(w) := \prod_{h=0}^{d-1} \text{sinc}_{\text{rad}}\left(\frac{\pi w}{2^h}\right), \quad P_0(w) = 1. \quad (200)$$

For arbitrary  $k, q \in \mathbb{N}_0$  and  $w \in \mathbb{C}$ , the global, denominator-cleared factorization at the center  $2^k q$  is

$$(2^k q + w)^{k+1} \Phi(2^k q + w) = (-1)^q w^{k+1} P_{k+1}(w) \Phi\left(\frac{q}{2} + \frac{w}{2^{k+1}}\right). \quad (201)$$

This is `rvachevFourierProduct_two_pow_mul_add_factorization` in `IntegerZeroLocalFactorization.lean`. It includes even  $q$ , when the tail can retain further zero multiplicity, and also the degenerate center  $q = 0$ . If  $q$  is odd, its sign is exactly  $-1$ ; the corresponding specialization is `rvachevFourierProduct_two_pow_mul_odd_add_factorization`.

For a positive integer  $m$ , let  $d = \nu_2(m) + 1$  and let  $q = m/2^{\nu_2(m)}$  be its canonical odd part. Then

$$(m + w)^d \Phi(m + w) = -w^d P_d(w) \Phi\left(\frac{q}{2} + \frac{w}{2^d}\right), \quad (202)$$

which is `rvachevFourierProduct_nat_add_factorization`. The fully signed form is just as clean. For  $m \in \mathbb{Z} \setminus \{0\}$ , set

$$d_m := \nu_2(|m|) + 1, \quad q_m := \frac{|m|}{2^{\nu_2(|m|)}}, \quad \sigma_m := \operatorname{sgn}(m). \quad (203)$$

Then  $q_m$  is odd and, for every complex  $w$ ,

$$(m + w)^{d_m} \Phi(m + w) = -w^{d_m} P_{d_m}(w) \Phi\left(\frac{q_m}{2} + \sigma_m \frac{w}{2^{d_m}}\right). \quad (204)$$

This is `rvachevFourierProduct_int_add_factorization`; the sign of the integer center occurs only in the tail coordinate. Equations (201)–(204) are polynomially cleared identities on all of  $\mathbb{C}$ , not punctured-neighborhood statements.

The remaining analytic argument can therefore be packaged into a single local unit. The product  $\Phi$  is entire by `rvachevFourierProduct_differentiable`, and, for every odd natural  $q$ ,

$$\Phi(q/2) \neq 0, \quad (205)$$

by `rvachevFourierProduct_nat_div_two_ne_zero_of_odd`. Define the total cofactor

$$U_m(w) := -\frac{P_{d_m}(w) \Phi(q_m/2 + \sigma_m w/2^{d_m})}{(m + w)^{d_m}}. \quad (206)$$

The Lean definition `integerZeroLocalCofactor` is total for every integer  $m$ . For  $m \neq 0$  and  $m + w \neq 0$ ,

$$\Phi(m + w) = w^{d_m} U_m(w), \quad (207)$$

by `rvachevFourierProduct_int_add_eq_pow_mul_cofactor`; the same equality holds as a germ at  $w = 0$  by `rvachevFourierProduct_int_add_eventuallyEq_pow_mul_cofactor`. In the original coordinate this is the centered germ

$$\Phi(z) = (z - m)^{d_m} U_m(z - m) \quad \text{near } z = m,$$

exported as `rvachevFourierProduct_int_eventuallyEq_sub_pow_mul_cofactor`.

For every integer parameter  $m$ , with no parity or nonzero hypothesis, the total quotient  $U_m$  is analytic at every point  $w_0$  satisfying  $m + w_0 \neq 0$ , by `integerZeroLocalCofactor_analyticAt_of_add_ne`. For nonzero  $m$ , its local-origin specialization is `integerZeroLocalCofactor_analyticAt`. Its exact central value is

$$U_m(0) = -\frac{\Phi(q_m/2)}{m^{d_m}} \neq 0. \quad (208)$$

These three facts are `integerZeroLocalCofactor_analyticAt`, `integerZeroLocalCofactor_or_zero`, and `integerZeroLocalCofactor_zero_ne`, respectively. More precisely, the middle theorem is the all-integer totalized identity

$$U_m(0) = -\frac{\Phi(\operatorname{divMaxPow}(|m|, 2)/2)}{m^{\operatorname{padicValNat}(2, |m|)+1}};$$

for  $m \neq 0$  it reduces to (208), and the last theorem makes it an analytic unit.

Combining the germ factorization, the nonzero unit, and (199) gives the exact complex multiplicity and its entire leading jet:

$$\operatorname{ord}_{z=m} \Phi(z) = d_m, \quad (209)$$

$$\Phi^{(j)}(m) = 0 \quad (0 \leq j < d_m), \quad (210)$$

$$\Phi^{(d_m)}(m) = -\frac{d_m! \Phi(q_m/2)}{m^{d_m}} \neq 0. \quad (211)$$



In fact the division-free all-order identity is, for every  $r \in \mathbb{N}_0$ ,

$$\Phi^{(d_m+r)}(m) = \binom{d_m+r}{d_m} d_m! U_m^{(r)}(0). \quad (212)$$

The order statement is `analyticOrderAt_rvachevFourierProduct_int`; the lower derivatives are `iteratedDeriv_rvachevFourierProduct_int_eq_zero_of_1t`; the displayed all-order formula is `iteratedDeriv_rvachevFourierProduct_int_add_order`; the displayed leading coefficient combines its  $r = 0$  case with `integerZeroLocalCofactor_zero` and is exported as `iteratedDeriv_rvachevFourierProduct_int`; and its nonvanishing is also exported directly as `iteratedDeriv_rvachevFourierProduct_int_ne_zero`, all in `IntegerZeroAnalyticOrder.lean`. This strengthens the real positive-integer limit in [Theorem 5.9](#): it is a signed-integer, complex-local theorem with the first surviving derivative evaluated exactly.

### 5.5 Leading Laurent coefficient of the reciprocal centered MGF

For a bounded Fabius solution  $F$ , define the centered Rvachev moment generating function in the manuscript normalization by

$$M_F(t) := \text{rvachevCenteredMGF } F \, t = \text{centeredComplexGeneratingFunction } F \, (2t). \quad (213)$$

The factor 2 corrects the half-scale coordinate of the repository’s pre-existing centered generating function. If  $h_F : \text{IsFabius } F$ , Fourier–Laplace rotation gives

$$M_F(t) = \Phi\left(\frac{it}{2\pi}\right), \quad (214)$$

which is `rvachevCenteredMGF_eq_rvachevFourierProduct`.

The reusable product-coordinate statement is worth recording first. For every nonzero  $m \in \mathbb{Z}$ , put  $d_m = \nu_2(|m|) + 1$ . Then

$$\lim_{\substack{z \rightarrow m \\ z \neq m}} (z - m)^{d_m} \Phi(z)^{-1} = U_m(0)^{-1}. \quad (215)$$

This is `tendsto_sub_pow_mul_inv_rvachevFourierProduct_int`; it follows directly from the local factorization in (207) and the fact that  $U_m$  is a nonvanishing analytic unit.

Now let  $n = 2^v u \neq 0$ , where  $v \in \mathbb{N}_0$  and  $u \in \mathbb{Z}$  is odd, and set  $T_n = 2\pi i n$ . Transporting (215) through (214) gives the manuscript-normalized leading Laurent law

$$\lim_{\substack{t \rightarrow T_n \\ t \neq T_n}} (t - T_n)^{v+1} M_F(t)^{-1} = -\frac{T_n^{v+1}}{M_F(\pi i u)}. \quad (216)$$

The odd-core value in the denominator is nonzero. The exact wrapper is `tendsto_rvachevCenteredMGF_laurent_two_pow_mul_odd`; the signed-integer form before choosing  $v, u$  is `tendsto_rvachevCenteredMGF_laurent_int`. Evenness converts the signed odd core to Lean’s positive `Nat.divMaxPow` representative through `rvachevCenteredMGF_pi_mul_I_int`, and `rvachevCenteredMGF_pi_mul_I_int_ne_zero_of_odd` proves nonvanishing.

Thus `RvachevLaurentLeading.lean` has the exhaustive public surface of one definition, `rvachevCenteredMGF`, and the six theorems `rvachevCenteredMGF_eq_rvachevFourierProduct`, `rvachevCenteredMGF_pi_mul_I_int`, `rvachevCenteredMGF_pi_mul_I_int_ne_zero_of_odd`, `tendsto_sub_pow_mul_inv_rvachevFourierProduct_int`, `tendsto_rvachevCenteredMGF_laurent_int`, and `tendsto_rvachevCenteredMGF_laurent_two_pow_mul_odd`. This makes `is_p2_thm : Laurent-leading Exact`. All three reciprocal limits use the punctured neighborhood, essentially: Lean totalizes inversion by  $0^{-1} = 0$ , so the displayed pole-cancelled expression does not have its limiting value at the pole itself. No lower Laurent coefficients, pole-shell summation, or downstream Appell-coefficient asymptotic is claimed.

## 5.6 Finite weighted scale arithmetic behind the zero orders

The valuation in (173) belongs to a more general finite additive calculus; neither primality of the scale base nor an analytic infinite product is needed for this layer. Let  $(A, +, 0)$  be an additive commutative monoid, let  $w : \mathbb{N}_0 \rightarrow A$ , and define the inclusive weighted prefix

$$P_w(n) := \sum_{h=0}^n w(h). \quad (217)$$

It satisfies

$$P_w(0) = w(0), \quad P_w(n+1) = P_w(n) + w(n+1), \quad (218)$$

$$P_w(m+n) = \sum_{h=0}^{m-1} w(h) + P_{w^{(m)}}(n), \quad w^{(m)}(h) := w(m+h). \quad (219)$$

Empty sums have their usual value. More generally, if  $S$  is finite,  $a : S \rightarrow \mathbb{N}_0$ , and  $a(x) < H$  for every  $x \in S$ , then

$$\sum_{x \in S} P_w(a(x)) = \sum_{h=0}^{H-1} \#\{x \in S : h \leq a(x)\} \odot w(h), \quad (220)$$

where  $q \odot y$  means the sum of  $q$  copies of  $y$ . This finite layer-cake identity requires no subtraction, order, or multiplication on  $A$ .

For a natural base  $b$ , put

$$\nu_b(n) := \text{padicValNat}(b, n), \quad W_{b,w}(n) := P_w(\nu_b(n)) = \sum_{h=0}^{\nu_b(n)} w(h). \quad (221)$$

Thus, for  $b > 1$ ,  $n > 0$ , and  $k \in \mathbb{N}_0$ ,

$$W_{b,w}(b^k n) = \sum_{h=0}^{k-1} w(h) + W_{b,w^{(k)}}(n), \quad (222)$$

$$W_{b,w}(bn) = w(0) + W_{b,w^{(1)}}(n). \quad (223)$$

For the divisibility form, the optimal algebraic hypothesis is only  $b \neq 1$ . If  $n > 0$  and  $\nu_b(n) < H$ , then

$$W_{b,w}(n) = \sum_{\substack{0 \leq h < H \\ b^h | n}} w(h). \quad (224)$$

Since  $\nu_b(n) < n$ , the canonical choice  $H = n$  gives

$$W_{b,w}(n) = \sum_{\substack{0 \leq h < n \\ b^h | n}} w(h). \quad (225)$$

Summing over the positive integers through  $N$  and exchanging the two finite sums yields

$$\sum_{m=0}^{N-1} W_{b,w}(m+1) = \sum_{h=0}^{N-1} (N \operatorname{div} b^h) \odot w(h) \quad (b \neq 1), \quad (226)$$

where  $\operatorname{div}$  denotes natural-number quotient. For  $b > 1$  this is the familiar floor  $\lfloor N/b^h \rfloor$ .

Pascal weights now give the hockey-stick identities

$$P_{h \mapsto \binom{h+r}{r}}(n) = \binom{n+r+1}{r+1}, \quad (227)$$

$$P_{h \mapsto \binom{h}{r}}(n) = \binom{n+1}{r+1}. \quad (228)$$

Consequently, with no restriction on  $b, n, r$ ,

$$W_{b, h \mapsto \binom{h}{r}}(n) = \binom{\nu_b(n) + 1}{r+1}, \quad (229)$$

and, for  $b \neq 1$ ,

$$\sum_{m=0}^{N-1} \binom{\nu_b(m+1) + 1}{r+1} = \sum_{h=0}^{N-1} (N \operatorname{div} b^h) \binom{h}{r}. \quad (230)$$

*Formal proof.* The definition (217), its zero and successor laws, and its shifted splitting law are `inclusivePrefixSum`, `inclusivePrefixSum_zero`, `inclusivePrefixSum_succ`, and `inclusivePrefixSum_add`. The monoid-valued Fubini identity (220) is `sum_inclusivePrefixSum_eq_sum_filter_card_nsmul`.

The scale definition (221) is `weightedScaleMultiplicity`; constant weights and unit natural weights are evaluated by `weightedScaleMultiplicity_const` and `weightedScaleMultiplicity_one_nat`. Equations (222) and (223) are `weightedScaleMultiplicity_base_pow_mul` and `weightedScaleMultiplicity_base_mul`. The bounded and canonical divisibility sums (224) and (225) are `weightedScaleMultiplicity_eq_sum_filter_pow_dvd_of_lt` and `weightedScaleMultiplicity_eq_sum_filter_pow_dvd`; the cumulative identity (226) is `sum_range_weightedScaleMultiplicity`.

Finally, (227) and (228) are `inclusivePrefixSum_add_choose` and `inclusivePrefixSum_choose`. Their valuation specialization and cumulative floor sum are `weightedScaleMultiplicity_choose` and `sum_range_padicValNat_choose`. All declarations are in `WeightedScaleMultiplicity.lean`.  $\square$

The finite arithmetic count has a sharp general-base form. Write

$$s_b(N) := \sum_{d \in \operatorname{digits}_b(N)} d.$$

For every integer  $b > 1$ , including a composite base, the logarithmic height already contains all layers:

$$\sum_{m=0}^{N-1} W_{b,w}(m+1) = \sum_{h=0}^{\lfloor \log_b N \rfloor} (N \operatorname{div} b^h) \odot w(h). \quad (231)$$

At unit natural weight this gives the exact digit recovery identities

$$(b-1) \sum_{n=1}^N (1 + \nu_b(n)) + s_b(N) = bN, \quad (232)$$

$$\sum_{n=1}^N (1 + \nu_b(n)) = \frac{bN - s_b(N)}{b-1}. \quad (233)$$

They include  $N = 0$ ; the quotient is natural-number division and the preceding additive identity proves its exactness.

*Formal proof.* The arbitrary-height and sharp logarithmic layer identities are `sum_range_weightedScaleMultiplicity_of_log_lt` and `sum_range_weightedScaleMultiplicity_log`. Splitting off the zeroth floor term is `sum_range_div_pow_log_eq_self_add_tail`; (232) and (233) are `sub_one_mul_sum_padicValNat_succ_add_digitSum` and `sum_range_padicValNat_succ_eq_sub_digitSum_div`. These five theorems are the exhaustive public surface of `BaseDigitMultiplicity.lean`.  $\square$

Lean totalizes the boundary cases:  $\nu_b(0) = \nu_0(n) = \nu_1(n) = 0$ , and natural division by zero is zero. Accordingly, the resulting  $W$ -values at those degenerate inputs are algebraic conventions. More importantly, this subsection proves finite arithmetic identities only. Formula (229) supplies the binomial expression proposed for higher-rank Rvachev products; the analytic bridge for admissible dyadic exponent sequences is now supplied by the following separate layer.

### 5.7 Recovering generalized Rvachev exponents from dyadic orders

Let  $a : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  satisfy  $\sum_{h \geq 0} a(h)/2^h < \infty$ , and set

$$\Phi_a(z) := \prod_{h \geq 0} \text{sinc}_{\text{rad}}\left(\frac{\pi z}{2^h}\right)^{a(h)}. \quad (234)$$

The complete generalized zero-divisor theorem and  $\nu_2(2^n) = n$  give the sparse but lossless sampling formula

$$\text{ord}_{z=2^n} \Phi_a(z) = W_{2,a}(2^n) = P_a(n) = \sum_{h=0}^n a(h). \quad (235)$$

These orders are finite. Hence Lean’s `ENat.toNat` converts them back to natural numbers, and the exponent decoder is

$$a(0) = \text{ENat.toNat}(\text{ord}_{z=1} \Phi_a(z)), \quad (236)$$

$$a(n+1) = \text{ENat.toNat}(\text{ord}_{z=2^{n+1}} \Phi_a(z)) - \text{ENat.toNat}(\text{ord}_{z=2^n} \Phi_a(z)). \quad (237)$$

In particular, for two admissible sequences  $a, c$ ,

$$\Phi_a = \Phi_c \iff a = c. \quad (238)$$

*Formal proof and exact recovery argument.* Equation (235) is `analyticOrderAt_generalizedRvachevProduct_two_pow`: the complete integer zero-divisor formula identifies the order at a positive integer with `weightedScaleMultiplicity`, and `padicValNat_base_pow` reduces its value at  $2^n$  to the inclusive prefix  $P_a(n)$ . At  $n = 0$  the sampled point is  $2^0 = 1$ , not the origin, and  $P_a(0) = a(0)$ ; every product in (234) instead has value one at the origin. Next,  $P_a(n+1) = P_a(n) + a(n+1)$ . Applying `ENat.toNat` to the finite natural-cast order formula and using natural-number cancellation gives (237).

The constructive formulas are `exponent_zero_eq_toNat_analyticOrderAt_generalizedRvachevProduct` and `exponent_succ_eq_toNat_analyticOrderAt_generalizedRvachevProduct`. Equality of all dyadic orders implies equality of the exponent sequences by `exponentSequence_eq_of_analyticOrderAt_two_pow_eq`; its reusable general-base cancellation lemma is `weightSequence_eq_of_weightedScaleMultiplicity_base_pow_eq`, valid for every  $b > 1$  in an additive cancellative commutative monoid. Finally, full function equality transports through `analyticOrderAt` at every dyadic point, so the decoder proves the forward implication in (238); the reverse implication is substitution. This is `generalizedRvachevProduct_eq_iff`. All six theorems, and no definitions, form `GeneralizedRvachevIdentifiability.lean`.  $\square$

The datum used here is the analytic divisor—the zero locations together with their multiplicities. The unweighted zero set alone is insufficient, and sampling the product values at the dyadic zeros is also insufficient: distinct positive exponent sequences can vanish at the same sampled points. The two summability hypotheses in the comparison theorem are essential because the underlying infinite product is totalized outside its admissible range. No spectral-zeta recovery, cumulant-sample recovery, or identifiability of an underlying probability law is asserted; those remain open interfaces. With the later concurrent source union, the real-MGF bridge formed the historical 905/11,458 checkpoint, the exhaustive complex moment-product leaf formed the historical 906/11,461 checkpoint, and the later exhaustive zero-definition/one-theorem `HalfQBinomialRootSimplicity.lean` leaf raised the merged-main checkpoint to 919/11,569. The exhaustive exterior reciprocal-germ leaf raises that checkpoint to 920/11,572, and the exhaustive sharp moment-polynomial degree leaf raises it to the historical 921/11,575 checkpoint. The exhaustive one-definition/six-theorem `RvachevLaurentLeading.lean` leaf then gives 922/11,582, and the eleven-definition/seventeen-theorem `FinitePrefixAppellRecovery.lean` leaf gives the target 923/11,610 census. The immediately preceding algebraic moment-polynomial checkpoint was 904/11,457, and the complex Bernoulli value-completion checkpoint 903/11,448 remains historical. The merged chronology also records the scaled-geometric checkpoint 914/11,555 and the earlier Appell/fixed-column/barycenter/Matrix checkpoint 910/11,525. The current TeX also contains the six-theorem prime-power companion-row API and the zero-definition/three-theorem outer Pochhammer normal-convergence API, plus the two-module/nine-declaration fixed-depth effective-inverse tranche, the eight-declaration rational-gap inverse module, the two-definition/ten-theorem sharp exact-dyadic inverse-modulus leaf, and six new q-calculus modules contributing 36 declarations, plus the two source-only central-reduction and cyclotomic-factorization modules contributing thirteen theorems, and the four-module root-of-unity, q-Lucas, cyclotomic-divisibility, and q-Catalan tranche contributing one definition and twenty-four theorems, plus the zero-definition/one-theorem half-base root-simplicity leaf, and the four-module Lambert branch-pairing, gap-bijection, symmetric-identity, and Bernoulli-series tranche contributing four definitions and thirty-seven theorems (forty-one public declarations). In its final leaf the complex Bernoulli series is summable exactly for  $|z| < 2\pi$ , hence diverges on and outside that boundary, and `hasSum_bernoulli_mul_pow_div_factorial_complex_iff` gives the canonical removable value  $\text{complexExp}(\text{m1Div}(z))^{-1}$  exactly on that disk, including value one at  $z = 0$ . This is not Lean’s literal totalized quotient at zero and asserts no holomorphy; the real quotient theorem retains  $z \neq 0$ , the branch theorem excludes both endpoints, and no remainder or higher-Puiseux claim is made. The union further contains the three-definition/seven-theorem geometric Richardson generating-function module, the three-theorem second-moment strengthening of the existing `GaussianBinomialCumulants.lean` module, the one-definition/seventeen-theorem `GeometricUniformRealization.lean` module, the two-definition/one-theorem `RegularCentralQBinomialSum.lean` leaf, and the zero-definition/ten-theorem `GaussianBinomialFixedColumnRate.lean` leaf, followed by the one-definition/fourteen-theorem `RvachevAppellHasse.lean` leaf, the one-definition/fourteen-theorem `RvachevLagrangeNodesOnly.lean` leaf, the zero-definition/two-theorem `GaussianBinomialGreaterOneAsymptotics.lean` leaf, the one-definition/eight-theorem `GeometricUniformMomentPolynomial.lean` module, the zero-definition/three-theorem `ThueMorseGammaTowerDifferential.lean` leaf, and the zero-definition/one-theorem `GeometricUniformMomentPolynomialBridge.lean` companion, the one-definition/two-theorem `GeometricUniformComplexMomentProduct.lean` complex-product leaf, the one-definition/two-theorem `GeometricUniformExteriorComplexMomentGerm.lean` exterior-germ leaf, the zero-definition/three-theorem `GeometricUniformMomentPolynomialDegree.lean` sharp-degree leaf, the one-definition/four-theorem `ThueMorseCornerIntegral.lean` and zero-definition/three-theorem `RvachevLegendreCentralSum.lean` leaves, and the sixteenth theorem in the zero-definition `FinitePolynomialFunctional.lean` module, followed by the zero-definition/one-theorem `HalfQBinomialRootSimplicity.lean`, one-definition/six-theorem `RvachevLaurentLeading.lean`, and eleven-definition/seventeen-theorem `FinitePrefixAppe`

11Recovery.lean leaves. The separately inventoried five-module polynomial  $q$ -calculus increment is retained in full: PolynomialQDerivative.lean (2+17), PolynomialQLeibniz.lean (0+4), QPochhammerDerivative.lean (0+3), LambertSeriesLog.lean (0+4), and QGamma.lean (2+10), totaling four definitions and thirty-eight theorems, forty-two public declarations, with the exact hypotheses and boundaries stated in their dedicated sections. Retaining the unconditional public complexQPochhammerInf\_eq\_qPochhammerInfIn bridge in the 1+10 RvachevPochhammerFactorization.lean surface gives the live semantic union of 923 modules and 11,611 unique public declarations, with no missing module header or declaration documentation. These are source-only overlays; the retained PDF is explicitly historical, and a fresh uninterrupted three-pass Libertinus parity rebuild remains pending. Exact historical source/PDF receipts are kept externally so this artifact embeds no mutable self-receipt.

The six  $q$ -calculus module counts are QPochhammerInfiniteBounds.lean (0+5), HeineTransformation.lean (2+5), QGaussSummation.lean (0+2), QPochhammerComplexOrder.lean (1+4), BasicHypergeometricSeries.lean (2+5), and QMultinomial.lean (1+9). They supply bounds, Heine and  $q$ -Gauss, complex order, general basic-hypergeometric summability, and the recursive  $q$ -multinomial API under their displayed hypotheses. The four subsequent module counts are GaussianBinomialPalindromic.lean (0+14), JacksonIntegral.lean (1+7), QExponential.lean (3+8), and ThetaQuasiPeriodicity.lean (1+6). Their five definitions and thirty-five theorems add Gaussian-polynomial degree, palindromicity, and the total linear-coefficient classifier,  $q$ -exponential and  $q$ -derivative laws, Jackson integration, and bilateral-theta quasi-periodicity and zeros, with all analytic hypotheses displayed.

The four newest module counts are GaussianBinomialPolynomialStructure.lean (0+5), JacobiCubic.lean (0+2), QPochhammerLogDerivative.lean (0+10), and QPochhammerOrderDerivative.lean (0+3). Their twenty theorems add universal Gaussian polynomial structure over  $\mathbb{N}_0[X]$ , Jacobi’s cubic identity, the  $q$ -Pochhammer logarithmic derivative and Lambert-series form, and the derivative with respect to complex order under the displayed hypotheses.

The newest moment-polynomial and product counts are GeometricUniformMomentPolynomial.lean (1+8) and GeometricUniformMomentPolynomialBridge.lean (0+1), followed by GeometricUniformComplexMomentProduct.lean (1+2) and GeometricUniformExteriorComplexMomentGerm.lean (1+2), and finally GeometricUniformMomentPolynomialDegree.lean (0+3). The algebraic module’s definition is geometricUniformMomentPolynomial; its theorems are geometricUniformMomentPolynomial\_zero, geometricUniformMomentPolynomial\_succ, geometricUniformMomentPolynomial\_natDegree\_le, geometricUniformMomentPolynomial\_eval\_zero, and the four explicit specializations geometricUniformMomentPolynomial\_one, geometricUniformMomentPolynomial\_two, geometricUniformMomentPolynomial\_three, and geometricUniformMomentPolynomial\_four. This supplies the recursive polynomial family, residual-product recurrence, triangular degree bound, value at zero, and the cases  $P_0, \dots, P_4$ . The companion’s exhaustive public surface is the single theorem geometricUniformMomentPolynomial\_eval2\_eq\_mgf\_taylorCoefficient. For every real  $q$  with  $|q| < 1$  and every  $n$ , it identifies the polynomial evaluation with the finite- $q$ -Pochhammer normalization of the  $n$ -th Taylor coefficient of the actual geometric-uniform moment generating function. Thus  $p7 : eq : Pn\text{-def}$  is Exact in this real contraction regime, including signed  $q$  and  $q = 0$ .

The complex leaf’s exhaustive public surface is the definition geometricUniformComplexMomentProduct and the two theorems hasProdLocallyUniformly\_geometricUniformComplexMomentProduct and geometricUniformMomentPolynomial\_eval2\_eq\_complexMomentProduct\_taylorCoefficient. For every complex  $q$  with  $\|q\| < 1$ , they give the actual infinite product, its locally uniform convergence on the complex plane, and the same finite- $q$ -Pochhammer normalization of its  $n$ -th Taylor coefficient. This is the complex analytic analogue of the real moment identity, not a complex probability-moment statement.

The exterior leaf’s exhaustive public surface is the definition geometricUniformExteri

orComplexMomentGerm and the two theorems analyticAt\_geometricUniformExteriorComplexMomentGerm and geometricUniformMomentPolynomial\_eval2\_eq\_exteriorComplexMomentGerm\_taylorCoefficient. For every complex  $q$  with  $1 < \|q\|$ , the definition is the actual reciprocal germ  $(\mathcal{A}_{q^{-1}}(-z))^{-1}$ , equivalently the manuscript’s reciprocal product indexed by  $j \geq 1$ ; the theorems prove analyticity at the origin and the same finite- $q$ -Pochhammer normalization of its  $n$ -th Taylor coefficient. The canonical  $q$ -monograph compound label `thm:qF-moment-polynomial` nevertheless remains Partial: no Lean `RatFunc` declaration packages one rational function  $a_n = d_n/n!$  across the inner and exterior regimes or proves the pole-clearing identity and resulting polynomial continuation at roots of unity. In particular, the exterior theorem does not assert a boundary value on  $\|q\| = 1$ .

The sharp-degree leaf’s exhaustive public surface consists of `coeff_geometricUniformMomentPolynomial_choose_two`, `coeff_geometricUniformMomentPolynomial_choose_two_sub_one`, and `geometricUniformMomentPolynomial_natDegree_eq`. With  $T_n = \binom{n}{2}$ , they prove

$$[q^{T_n}]P_n = \frac{B_n(1)}{n!}, \quad [q^{T_n-1}]P_n = -\frac{B_n(1)}{n!} + \frac{B_{n-1}(1)}{2(n-1)!} \quad (n \geq 2),$$

and

$$\deg P_n = \begin{cases} T_n, & n = 1 \text{ or } n \text{ is even,} \\ T_n - 1, & \text{otherwise.} \end{cases}$$

Here Lean’s `bernoulli`’ is the positive convention  $B_n(1) = (-1)^n B_n$ , with Mathlib’s  $B_1 = -1/2$ . Thus the canonical `prop:qF-P-degree-sharp` and the complete algebraic/analytic frontier theorem `p7:thm:Pn` are Exact. This does not fill the separate global `RatFunc` and root-continuation gap above. The later merged-main checkpoint was 919/11,569, the exterior  $1 + 2$  leaf raises it to 920/11,572, and this sharp  $0 + 3$  leaf raises it to the historical 921/11,575 checkpoint. The following `RvachevLaurentLeading.lean` (1 + 6) and `FinitePrefixAppellRecovery.lean` (11 + 17) leaves give the target checkpoints 922/11,582 and 923/11,610, respectively; retaining the unconditional public Pochhammer bridge gives the live 923/11,611 semantic union.

The final two module counts are `CentralQBinomialReduction.lean` (0+6) and `CyclotomicFactorization.lean` (0+7). Their thirteen division-free algebra theorems add sign pairing and even–odd finite  $q$ -Pochhammer dissection, the central Gaussian reduction, and the cyclotomic factorizations of  $(X; X)_n$  and  $[n, k]_X$ . The quotient reduction retains its two nonzero-denominator hypotheses, while the Gaussian cyclotomic factorization retains its integral-domain assumption.

The exhaustive six-theorem surface of `CentralQBinomialReduction.lean` is `finiteQPochhammerIn_mul_neg`, `finiteQPochhammerIn_two_mul`, `finiteQPochhammerIn_map_ringHom`, `central_gaussianBinomial_sq_mul_int`, `central_gaussianBinomial_sq_mul`, and `central_gaussianBinomial_sq_div`. The first five are polynomial or commutative-ring identities and use no division; the final quotient theorem is over a field and assumes exactly the two displayed finite  $q$ -Pochhammer denominators are nonzero. The exhaustive seven-theorem surface of `CyclotomicFactorization.lean` is `div_add_div_le_div`, `div_le_div_add_div_add_one`, `mem_range_and_mem_divisors_iff`, `finiteQPochhammerIn_X_eq_prod_cyclotomic`, `finiteQPochhammerIn_X_eq_gaussianBinomial_mul`, `prod_cyclotomic_pow_div_extend`, and `gaussianBinomial_X_eq_prod_cyclotomic`. The two floor inequalities show that each Gaussian cyclotomic exponent is zero or one. The shifted-factorial factorization and factorial identity work over every commutative ring; only the final cancellation to the Gaussian factorization requires an integral domain, and all Gaussian statements retain  $k \leq n$ .

The six post-factorization modules add exactly four definitions and forty-six theorems. The exhaustive surface of `PrimitiveRootBlock.lean` (0+3) is `gaussianBinomial_isPrimitiveRoot_eq_zero`, `neg_one_pow_mul_pow_choose_two`, and `finiteQPochhammerIn_isPrimitiveRoot`. They work over a commutative integral domain at a primitive  $d$ -th root: interior

Gaussian vanishing has exactly  $0 < k < d$ , while the phase and complete-block formulas explicitly assume  $0 < d$ .

The exhaustive public surface of `QLucas.lean` (0+7) is `add_mul_add_sub_one`, `choose_two_add`, `coeff_finiteQPochhammerIn_neg_X`, `finiteQPochhammerIn_neg_X_block`, `coeff_block_pow_mul`, `pow_choose_two_add_mul_eq`, and `gaussianBinomial_q_lucas`. The first two are natural-number identities; the two coefficient formulas are commutative-ring identities. The complete block, phase, and q-Lucas formulas use an integral domain, a primitive root, and  $0 < d$ ; q-Lucas additionally assumes  $b, s < d$ . Its formal endpoint is the evaluated identity  $[ad + b, rd + s]_\zeta = \binom{a}{r}[b, s]_\zeta$ , not a named polynomial congruence modulo  $\Phi_d$ . Its same-shaped `two_mul_choose_two` helper is private; the unique public theorem of that name is owned by `QChuVandermonde.lean`.

The exhaustive surface of `CyclotomicDivisibility.lean` (0+3) is `cyclotomic_exponent_eq_one_iff`, `cyclotomic_dvd_gaussianBinomial_iff`, and `gaussianBinomial_mul_isPrimitiveRoot`. The carry and divisibility criteria assume  $k \leq n$  and  $0 < d$ ; the divisibility theorem is specifically in  $\mathbb{Q}[X]$ . The multiple-index primitive-root value is over a commutative integral domain and assumes  $0 < n$ . These declarations, together with the earlier exponent bound, justify the squarefreeness row, but there is no separately named squarefree theorem; the Babbage-derivative row remains Partial because its value, but not its derivative, is formalized.

The exhaustive surface of `QCatalan.lean` (1+11) is `map_qInt`, `qInt_X_mononic`, `qInt_X_natDegree`, `X_sub_one_mul_qInt`, `qInt_X_eq_prod_cyclotomic`, `qInt_X_dvd_gaussianBinomial_rat`, `qInt_X_dvd_gaussianBinomial_int`, the definition `qCatalan`, and `qInt_X_mul_qCatalan`, `qCatalan_natDegree`, `qCatalan_eval_one_mul`, and `qCatalan_eval_one`. Ring maps already commute with q-integers at the semiring level; monicity and exact degree use a nontrivial commutative ring. The noncomputable  $\mathbb{Z}[X]$  quotient is defined for every  $n$ , including zero, and has degree  $n(n-1)$  and Catalan value at one. No coefficient-nonnegativity or unimodality theorem is asserted.

The exhaustive surface of `NewtonInterpolation.lean` (2+13) consists of the definitions `newtonCoeff` and `newtonInterpolant`, followed by `newtonCoeff_eq`, `newtonCoeff_zero`, `newtonCoeff_mul_prod`, `newtonPoly_succ`, `eval_newtonPoly`, `degree_newtonPoly_lt`, `newtonPoly_eq_interpolate`, `eq_newtonPoly_of_eval_eq`, `coeff_newtonPoly_self`, `newtonCoeff_eq_sum`, `nodal_range_pow`, `prod_erase_pow_sub_pow`, and `newtonCoeff_pow_eq_sum`. This is entirely finite interpolation over a field. Evaluation needs the relevant earlier-node difference product nonzero; uniqueness and divided differences use injectivity on the displayed finite range. The geometric basis assumes  $q \neq 0$ , and its coefficient sum assumes injectivity of  $j \mapsto q^j$ . The definition name `newtonInterpolant` preserves the established scalar-sequence `newtonPoly` API from `NewtonBasisGeneratingFunction.lean`. No topological or convergence claim is present.

Finally, `QBetaIntegral.lean` (1+8) has the total real definition `qBeta` and the theorems `qNumber_pos`, `qBeta_term_eq`, `qBeta_eq_prod`, `qBeta_eq_qGamma`, `qBeta_comm`, `qBeta_pos`, `qBeta_add_one_left`, and `qBeta_add_one_right`. The evaluation, q-gamma identity, symmetry, positivity, and recurrences assume exactly  $0 < q < 1$  and  $x, y > 0$ ; the term cancellation assumes  $y > 0$  with arbitrary real  $x$ . No complex continuation or classical-limit theorem is asserted. At this pre-upper-index checkpoint the 282 labelled forward rows were 80 Exact, 86 Partial, 108 None, and 8 N/A. In particular, q-Lucas is Partial: Lean proves the evaluated primitive-root identity, while the manuscript’s polynomial congruence modulo  $\Phi_d$  still needs a minimal-polynomial lift.

**Final q-calculus, theta, and derivative tranche.** Eight facade-reachable modules add exactly five definitions and fifty-five theorems. `GaussianBinomialPalindromic.lean` has no definitions and the fourteen theorems `reflect_add_of_natDegree_le`, `reflect_one'`, `gaussianBinomial_natDegree_le`, `gaussianBinomial_zero_left`, `gaussianBinomial_diag'`, `reflect_gaussianBinomial`, `coeff_gaussianBinomial_reflect`, `coeff_gaussi`



`anBinomial_zero`, `coeff_gaussianBinomial_top`, `gaussianBinomial_natDegree`, `gaussianBinomial_moniac`, and `two_mul_derivative_gaussianBinomial_eval_one`, `coeff_gaussianBinomial_one_of_pos_of_lt`, and `coeff_gaussianBinomial_one`. Over a commutative semiring they give the degree bound, reflection, coefficient symmetry, constant and top coefficients, the division-free first-moment identity, and the total linear-coefficient classifier (one exactly for  $0 < k < n$ , zero otherwise); exact degree and monicity additionally require a nontrivial coefficient type.

`QExponential.lean` has the three definitions `qDeriv`, `qExp`, and `qExpBig`, followed by the eight theorems `qFactorial_mul_one_sub_pow`, `qFactorial_ne_zero`, `qDeriv_mul`, `hasSum_qExp`, `hasSum_qExpBig`, `qExp_mul_qExpBig_neg`, `qDeriv_qExp`, and `qDeriv_qExpBig`. Its product rule is unconditional; its series and eigenfunction statements retain their strict-norm, argument-disc, and nonzero-point hypotheses. `JacksonIntegral.lean` has the definition `jacksonIntegral` and seven theorems: `qDeriv_jacksonIntegral`, `one_sub_mul_pow_mul_qDeriv`, `tendsto_jackson_sum_qDeriv`, `jacksonIntegral_qDeriv`, `tendsto_jackson_sum_parts`, `jackson_parts_of_tendsto`, and `jacksonIntegral_mul_qDeriv`. Its partial-sum telescoping statements are kept distinct from the summable fundamental-theorem and integration-by-parts forms.

`ThetaQuasiPeriodicity.lean` has the definition `bilateralTheta` and six theorems: `thetaExponent_add_one`, `pow_thetaExponent_add_one`, `hasSum_bilateralTheta`, `bilateralTheta_eq_prod`, `bilateralTheta_mul_left`, and `bilateralTheta_eq_zero_iff`. The quasi-periodicity and zero statements retain  $q \neq 0$  and  $z \neq 0$ ; the latter identifies exactly  $z = -q^m$ ,  $m \in \mathbb{Z}$ . `JacobiCubic.lean` has no definitions and exactly `two_mul_add_one_le_three_pow` and `hasSum_jacobi_cubic`, the latter under complex  $\|q\| < 1$ .

`QPochhammerLogDerivative.lean` has no definitions and ten theorems: `one_sub_le_norm_one_sub_mul_pow`, `summable_pow_div_one_sub_mul_pow`, `summable_log_one_sub_mul_pow`, `one_sub_mul_pow_ne_zero`, `qPochhammerInfIn_eq_cexp_tsum_log`, `hasDerivAt_tsum_log_one_sub_mul_pow`, `hasDerivAt_qPochhammerInfIn`, `hasDerivAt_lambert_series`, `tsum_neg_pow_div_one_sub_mul_pow_eq`, and `hasDerivAt_qPochhammerInfIn_lambert`. Its two derivative formulas use complex  $\|q\| < 1$  and  $\|a\| < 1$ . `QPochhammerOrderDerivative.lean` has no definitions and the three theorems `hasDerivAt_const_cpow`, `hasDerivAt_qPochhammerInfIn_mul_cpow`, and `hasDerivAt_qPochhammerC`; the order derivative retains  $q \neq 0$ ,  $\|q\| < 1$ , and  $\|aq^\alpha\| < 1$ .

The distinct `GaussianBinomialPolynomialStructure.lean` compatibility leaf has no definitions and five theorems: `natDegree_gaussianBinomial_universal`, `gaussianBinomial_universal_moniac`, `coeff_zero_gaussianBinomial_universal`, `gaussianBinomial_universal_reflect`, and `coeff_gaussianBinomial_universal_symm`. It works in  $\mathbb{N}_0[X]$  under  $k \leq n$  and supplies the exact universal degree, monicity, constant coefficient, reflection, and bounded-index coefficient symmetry.

The four root-of-unity and q-Catalan module counts are `CyclotomicDivisibility.lean` (0+3), `PrimitiveRootBlock.lean` (0+3), `QCatalan.lean` (1+11), and `QLucas.lean` (0+7). Their one definition and twenty-four theorems add the cyclotomic carry criterion, complete primitive-root q-Pochhammer blocks, q-Lucas over integral domains, and the integral q-Catalan polynomial with its exact degree and Catalan specialization.

The newest interpolation and q-integral module counts are `NewtonInterpolation.lean` (3+19) and `QBetaIntegral.lean` (1+8). Their four definitions and twenty-seven theorems construct triangular Newton coefficients and interpolants, prove evaluation, uniqueness, divided differences, and the geometric-grid formula, and evaluate the Jackson q-beta integral in product and q-gamma forms for  $0 < q < 1$  and positive arguments, with symmetry, positivity, and both successor recurrences. The Newton count includes the node-qualified family and all seven definitionally identical newton interpolant compatibility declarations. No endpoint or  $q \rightarrow 1$  limit is claimed.

The latest three q-series module counts are `GaussianBinomialInteger.lean` (1+10), `Ga`

`ussianBinomialComplexOrder.lean` (1+5), and `QPfaffSaalschutz.lean` (0+3). Their two definitions and eighteen theorems extend Gaussian coefficients to integer and principal-branch complex upper indices, prove the integer q-Pascal and negative-index reflection laws, derive the reciprocal finite and generalized complex-order q-binomial series, and establish the terminating balanced  ${}_3\phi_2$  q-Pfaff–Saalschütz summation over a field. Nonzero nomes and parameters, strict-contraction bounds, and every finite-product nonvanishing premise remain explicit. The q-Pfaff–Saalschütz surface is exactly `finiteQPochhammerIn_ne_zero_of_le`, `q_pfaff_saalschutz_term`, and `q_pfaff_saalschutz`. The term identity additionally assumes  $k \leq n$ , and no convergence or limiting claim is made.

The zero-definition/five-theorem module `QuantumMultinomial.lean` then decomposes natural tuple antidiagonals, transports Gaussian symmetry to arbitrary semirings, proves q-multinomial coefficient commutation, and establishes the ordered noncommutative q-multinomial theorem. Its hypotheses are exactly that the nome commute with each variable and that  $x_j x_i = q(x_i x_j)$  for  $i < j$ . The statement is finite and division-free, with no ambient commutativity or convergence assumption.

The final bound module `GaussianBinomialBounds.lean` has no public definitions and six theorems: `gaussianBinomial_inv`, `one_le_gaussianBinomial`, `finiteQPochhammerIn_pow_le_one`, `gaussianBinomial_le_inv_qPochhammerInfIn`, `pow_le_gaussianBinomial_of_one_lt`, and `gaussianBinomial_le_pow_div_of_one_lt`. It evaluates Gaussian reciprocity over fields, proves the constant-term lower bound and the uniform infinite-q-Pochhammer upper bound for nonnegative real strict contractions, and transfers them to reciprocal dimension-dominant two-sided bounds for every real  $Q > 1$ . The index condition  $k \leq n$ , nonzero nome for reciprocity, and all order and strict-contraction hypotheses remain explicit. Its proof reuses the stronger ordered-field theorem `finiteQPochhammerIn_self_pos` from `GeneralQConditionNumber.lean` rather than exporting a duplicate. The rigorous 282-row forward q-series ledger is therefore 177 Exact, 82 Partial, 15 None, and 8 interface rows; its 547-row source concordance is 99 Lean, 379 human, 60 N/A, and 9 conjecture rows. In particular q-Lucas remains Partial because Lean proves the evaluated primitive-root identity rather than the polynomial congruence modulo  $\Phi_d$ ; the Gaussian majorant row is Exact, and the  $Q > 1$  asymptotic row is Exact via the fixed-column and central equivalence theorems in `GaussianBinomialGreaterOneAsymptotics.lean`.

The terminating basic-hypergeometric closure is split exhaustively between `TwoPhiOneReversal.lean` (two definitions and twelve theorems) and `QChuVandermonde.lean` (zero definitions and ten theorems). The former’s definitions are `twoPhiOneFinite` and `twoPhiOneReflection`; its twelve theorems are `choose_two_add_succ_choose_two`, `finiteQPochhammerIn_sub_eq`, `finiteQPochhammerIn_reversal_ne_zero`, `finiteQPochhammerIn_inv_pow_self`, `twoPhiOneReflection_involutive`, `twoPhiOneFinite_reversal`, `twoPhiOneFinite_reversal_twice`, `twoPhiOneFinite_eq_sum_twoPhiOneTerm`, `twoPhiOne_eq_twoPhiOneFinite_inv_pow`, `twoPhiOne_reversal`, `twoPhiOne_reversal_twice`, and `twoPhiOne_one_eq_twoPhiOneFinite_zero`. Thus it identifies a terminating  ${}_2\phi_1$  with the actual `twoPhiOne_tsum`, defines the reflected parameter triple, and proves both involutivity and cancellation after applying the reversal twice.

The ten-theorem `QChuVandermonde.lean` surface is `two_mul_choose_two`, `mul_sub_one_eq_mul_sub_add`, `finiteQPochhammerIn_div_eq_sum_chu`, `q_chu_vandermonde_first`, `finiteQPochhammerIn_div_eq_sum_chu_second`, `twoPhiOneFinite_mul_finiteQPochhammerIn_eq_chu_second`, `q_chu_vandermonde_second`, `q_chu_vandermonde_second_by_reversal`, `twoPhiOne_q_chu_vandermonde_first`, and `twoPhiOne_q_chu_vandermonde_second`. It proves both q-Chu–Vandermonde sums, including a denominator-cleared second identity and the full-domain actual-tsum wrappers. Consequently the two displayed q-Chu formulas and the terminating reversal lemma are exact Lean crosswalks. The separate claim that the second formula follows by reversal on its entire displayed domain remains only partial: the compiled provenance theorem `q_chu_vandermonde_second_by_reversal` retains the auxiliary hypotheses  $C \neq 0$  and  $(A; q)_n \neq 0$ , whereas the full-domain proof uses finite q-Cauchy directly. No rational-continuation or

cleared commutative-ring extension is asserted here.

**Gaussian-binomial second cumulants.** The later three-theorem delta in `GaussianBinomialCumulants.lean`, whose complete public surface is now 2+24, is `eval_one_derivative_derivative_gaussianBinomial_X`, `twelve_mul_secondMoment_gaussianBinomial_eval_one`, and `twelve_mul_varianceNumerator_gaussianBinomial_eval_one`. Put  $P = [n, k]_X$ ,  $D = k(n - k)$ , and  $C = \binom{n}{k}$ . The first theorem is over a characteristic-zero field and assumes exactly  $k \leq n$ ; it gives

$$P''(1) = \frac{D(3D + n - 5)}{12} C.$$

The other two hold for every  $n, k$  over an arbitrary commutative semiring, with no division, nonvanishing, or characteristic hypothesis:

$$12(P''(1) + P'(1)) = D(3D + n + 1)C, \quad 12C(P''(1) + P'(1)) = 12P'(1)^2 + D(n + 1)C^2.$$

They are total above the Gaussian row by zero extension and include the zero and diagonal rows; the first displayed cleared identity is the unnormalized raw second moment, and the second is the variance numerator.

On the real axis the origin is the unique global point of unit modulus:

$$|\widehat{\text{up}}_{2\pi}(x)| = 1 \iff x = 0, \quad x \neq 0 \implies |\widehat{\text{up}}_{2\pi}(x)| < 1. \quad (239)$$

These are `norm_rvachevFourier_eq_one_iff` and `norm_rvachevFourier_lt_one_of_ne_zero`. On each positive real lobe the sign and peak are also exact. If  $m \in \mathbb{N}_0$  and  $m < x < m + 1$ , then

$$\widehat{\text{up}}_{2\pi}(x) = \varepsilon_m |\widehat{\text{up}}_{2\pi}(x)|, \quad (240)$$

by `rvachevFourier_eq_trueMorse_sign_mul_norm`. If moreover  $m \geq 1$ , there is exactly one point of  $(m, m + 1)$  maximizing the modulus over that open lobe and exactly one point of the reflected lobe  $(-(m + 1), -m)$  maximizing it there. These are `existsUnique_isMaxOn_rvachevFourier_lobe` and `existsUnique_isMaxOn_rvachevFourier_neg_lobe`. All five facade declarations are in `RvachevFourierDecay.lean`.

Globally, the origin is the unique real frequency of unit modulus:

$$|\widehat{\text{up}}_{2\pi}(x)| \leq 1, \quad |\widehat{\text{up}}_{2\pi}(x)| = 1 \iff x = 0. \quad (241)$$

Equivalently, the first inequality is strict whenever  $x \neq 0$ . These statements are `norm_rvachevFourier_le_one`, `norm_rvachevFourier_lt_one_of_ne_zero`, and `norm_rvachevFourier_eq_one_iff` in `RvachevFourierDecay.lean`.

## 5.8 Rapid decay and inversion

For every  $N \in \mathbb{N}_0$  there is a constant  $C_N > 0$  such that

$$|\xi|^N |\widehat{\text{up}}_{2\pi}(\xi)| \leq C_N \quad (\xi \in \mathbb{R}), \quad (242)$$

and the companion formal theorem gives the corresponding polynomially weighted bound for every iterated real-frequency derivative. Fourier inversion holds in the exact form

$$\text{up}(x) = \int_{\mathbb{R}} \widehat{\text{up}}_{2\pi}(\xi) e^{2\pi i x \xi} d\xi. \quad (243)$$

These are `rvachevFourier_real_rapidDecay`, `rvachevFourier_real_iteratedDeriv_rapidDecay`, and `rvachev_fourier_inversion_analytic` in `PoissonSummation.lean` and `FourierAnalytic.lean`.

### 5.9 Poisson summation and partitions of unity

For  $u > 0$  and  $t \in \mathbb{R}$ , the formal Poisson identity is

$$\sum_{k \in \mathbb{Z}} \text{up}(t + uk) = \frac{1}{u} \sum_{k \in \mathbb{Z}} \widehat{\text{up}}_{2\pi}(k/u) e^{2\pi i kt/u}. \quad (244)$$

It is `rvachev_poisson_summation`. Combining it with (172) gives, for every positive integer  $m$ ,

$$\sum_{k \in \mathbb{Z}} \text{up}\left(t + \frac{k}{m}\right) = m, \quad (245)$$

and in particular

$$\sum_{k \in \mathbb{Z}} \text{up}(t + k) = 1. \quad (246)$$

These are `rvachev_partition_one_over_nat` and `rvachev_partition_unity` in `PoissonSummation.lean`.

### 5.10 Banach-free cyclic aliasing

**Theorem 5.13** (Finite-range twists and cyclic aliasing). *The convergence mechanism behind a root-of-unity filter does not require a norm. Let  $R$  be any scalar type, let  $B$  be any index type, let  $I$  be finite, and let  $E$  be a complete uniform additive commutative group with a distributive  $R$ -action such that multiplication by each fixed scalar is continuous. If  $a : B \rightarrow E$  is summable,  $\ell : B \rightarrow I$ , and  $c : I \rightarrow R$ , then*

$$b \mapsto c(\ell(b)) a(b) \quad \text{is summable.} \quad (247)$$

*Thus a scalar multiplier taking only finitely many values preserves unconditional summability.*

*Separately, let  $E$  be a complete Hausdorff uniform additive commutative group, write  $\bar{n}$  for the class of  $n \in \mathbb{Z}$  in  $\mathbb{Z}\text{Mod}(q)$ , and define*

$$\mathcal{R}_r(a) := \sum_{\substack{n \in \mathbb{Z} \\ \bar{n} = r}} a(n). \quad (248)$$

*For every  $q \in \mathbb{N}_0$ , including Lean’s convention  $\mathbb{Z}\text{Mod}(0) = \mathbb{Z}$ , the family  $r \mapsto \mathcal{R}_r(a)$  has sum  $\sum_{n \in \mathbb{Z}} a(n)$ . In particular, for  $q > 0$ ,*

$$\sum_{r \in \mathbb{Z}/q\mathbb{Z}} \mathcal{R}_r(a) = \sum_{n \in \mathbb{Z}} a(n). \quad (249)$$

*Suppose in addition that  $E$  is a complex module and multiplication by each fixed complex scalar is continuous. Let  $\chi_q$  be the standard additive character of  $\mathbb{Z}/q\mathbb{Z}$ . For  $q > 0$ , put*

$$\mathcal{A}_k(a) := \sum_{n \in \mathbb{Z}} \chi_q(-\bar{n}k) a(n). \quad (250)$$

*Every twisted series in (250) is summable. With the unnormalized discrete Fourier transform*

$$(\mathcal{D}_q f)(k) := \sum_{r \in \mathbb{Z}/q\mathbb{Z}} \chi_q(-rk) f(r),$$

*one has the exact forward and inverse identities*

$$\mathcal{D}_q(\mathcal{R}_\bullet(a))(k) = \mathcal{A}_k(a), \quad (251)$$

$$\mathcal{R}_r(a) = \frac{1}{q} \sum_{k \in \mathbb{Z}/q\mathbb{Z}} \chi_q(kr) \mathcal{A}_k(a). \quad (252)$$

*Formal proof.* The finite-range convergence principle (247) is `summable_finite_range_smul`. The two sums are `intResidueTsum` and `intFourierTsum`; character twisting is `intFourierSummand_summable`. The fiberwise summation statement and its finite-modulus consequence are `hasSum_intResidueTsum` and `sum_intResidueTsum`. The forward transform and normalized inverse are `dft_intResidueTsum` and `intResidueTsum_eq_dft_average`. All eight declarations are in `SummableCyclicAlias.lean`.

The proof partitions the bilateral family into its residue fibers, observes that each cyclic character is constant on a fiber, and applies the finite DFT and its inverse. Summability of  $a$  is the only convergence hypothesis; norm-summability is not assumed.  $\square$

### 5.11 Finite odd-coset DFT traces

There is also a finite root-of-unity calculation behind the half-integer spectral frontier. It is useful to isolate it from every analytic alias or limiting argument.

**Theorem 5.14** (Odd-coset character filter and all-order trace). *Let  $N > 0$ , let  $R$  be a commutative domain, and let  $\zeta \in R$  be a primitive  $2N$ -th root of unity. Then, for every  $S \in \mathbb{N}_0$ ,*

$$\sum_{k=0}^{N-1} \zeta^{(2k+1)S} = \begin{cases} N(-1)^{S/N}, & N \mid S, \\ 0, & N \nmid S. \end{cases} \quad (253)$$

For samples  $f_0, \dots, f_{2N-1} \in R$ , put

$$A_k := \sum_{j=0}^{2N-1} f_j \zeta^{(2k+1)j}, \quad S(J) := \sum_{\ell=1}^m J_\ell, \quad (254)$$

where  $J = (J_1, \dots, J_m) \in \{0, \dots, 2N-1\}^m$ , and define

$$\mathcal{C}_{N,m}(f) := \sum_J \begin{cases} (-1)^{S(J)/N} \prod_{\ell=1}^m f_{J_\ell}, & N \mid S(J), \\ 0, & N \nmid S(J). \end{cases} \quad (255)$$

Then, for every  $m \in \mathbb{N}_0$ , including  $m = 0$ ,

$$\sum_{k=0}^{N-1} A_k^m = N \mathcal{C}_{N,m}(f). \quad (256)$$

Over a field, normalizing each coefficient by  $N$  gives

$$\sum_{k=0}^{N-1} \left( \frac{A_k}{N} \right)^m = \frac{N}{N^m} \mathcal{C}_{N,m}(f). \quad (257)$$

The labels also satisfy the period- $N$  identity  $A_{k+N} = A_k$ .

*Formal proof.* Equation (253) is `sum_odd_powers_eq_ramanujan`. The unnormalized and normalized trace identities are `oddDFTPowerTrace_eq_ramanujanConvolution` and `normalizedOddDFTPowerTrace_eq_ramanujanConvolution`; the coefficient, index-sum, and convolution definitions are `oddDFT`, `oddDFTIndexSum`, and `oddRamanujanConvolution`. The label identity is `oddDFT_add_period`. The tuple expansion before applying the root filter is separately exposed as `oddDFTPowerTrace_eq_indexSumExpansion`. These declarations are in `HalfIntegerOddDFT.lean`.

For arbitrary  $N$ , (253) sums the complete odd coset modulo  $2N$ . Passing from a full odd orbit to one representative from each reflection pair is available as `normalizedOddDFTPowerTrace_eq_two_mul_half`, but only under the exact reflection hypothesis supplied by its caller. No theorem here identifies the  $f_j$  with sampled Fabius data, proves an inverse DCT or a bilateral alias formula, or supplies an analytic error estimate.  $\square$

## 5.12 Moment expansion of the Fourier transform

With

$$c_n = \int_{-1}^1 x^{2n} \text{up}(x) \, dx. \quad (258)$$

the exact entire series is

$$\widehat{\text{up}}_{2\pi}(z) = \sum_{n=0}^{\infty} \frac{(-1)^n c_n}{(2n)!} (2\pi z)^{2n}. \quad (259)$$

This is `rvachevFourier_eq_momentSeries` in `AnalyticMoments.lean`. For later finite filters the individual mode is named

$$A_n(z) := \frac{(-1)^n c_n}{(2n)!} (2\pi z)^{2n} = \text{rvachevFourierMomentTerm } z \, n. \quad (260)$$

The declarations `rvachevFourierMomentTerm_zero` and `rvachevFourierMomentTerm_scale` prove

$$A_0(z) = 1, \quad A_n(cz) = (c^2)^n A_n(z) \quad (c, z \in \mathbb{C}, n \in \mathbb{N}_0).$$

More strongly than the equality of total sums in (259), `hasSum_rvachevFourierMomentTerm` proves `HasSum(A_•(z),  $\widehat{\text{up}}_{2\pi}(z)$ )` for every bounded Fabius solution. These four declarations are in `AnalyticMoments.lean`; the canonical product specialization used below is in `RvachevQBinomialFilter.lean`.

## 6 The global extension, derivatives, flatness, and inverse

### 6.1 The signed extension and all iterated derivatives

The translate series (11) is locally finite. It agrees with the bounded  $F$  on  $[0, 1]$ , vanishes on  $(-\infty, 0]$ , and satisfies

$$F'_{\text{gl}}(x) = 2F_{\text{gl}}(2x) \quad (x \in \mathbb{R}). \quad (261)$$

These facts are `extendedFabius_eq_fabiusReal`, `extendedFabius_eq_zero_of_nonpos`, and `extendedFabius_hasDerivAt` in `GlobalExtension.lean`.

Iterating the dilation law gives the exact identities

$$F_{\text{gl}}^{(m)}(x) = 2^{\binom{m+1}{2}} F_{\text{gl}}(2^m x), \quad (262)$$

$$\text{up}^{(m)}(x) = 2^{\binom{m+1}{2}} F_{\text{gl}}(2^m(x+1)) \quad (-1 \leq x \leq 1). \quad (263)$$

They are `iteratedDeriv_extendedFabius` and `iteratedDeriv_rvachev`. The global extension obeys

$$|F_{\text{gl}}(x)| \leq 1, \quad (264)$$

by `abs_extendedFabius_le_one`. Hence, for all  $m \in \mathbb{N}_0$  and  $x \in \mathbb{R}$ ,

$$|F^{(m)}(x)| \leq 2^{\binom{m+1}{2}}, \quad (265)$$

and this bound is attained at  $x = 2^{-m}$ :

$$F^{(m)}(2^{-m}) = 2^{\binom{m+1}{2}}. \quad (266)$$

The exact bound, attainment, and supremum statements are `abs_iteratedDeriv_fabiusReal_le`, `iteratedDeriv_fabiusReal_inv_two_pow_all`, and `isGreatest_abs_iteratedDeriv_fabiusReal_all` in `BoundedDerivatives.lean`.

In fact the single attaining point belongs to a complete matched-grid pattern. For  $k, m \in \mathbb{N}_0$  with  $0 \leq m < 2^k$ ,

$$F^{(k)}\left(\frac{m}{2^k}\right) = \begin{cases} 0, & m \text{ even,} \\ 2^{\binom{k+1}{2}} \varepsilon_{\lfloor m/2 \rfloor}, & m \text{ odd.} \end{cases} \quad (267)$$

Thus the even numerators are exactly the zeros, while every odd numerator attains the sharp absolute bound and the surviving signs form the Thue–Morse word. The exact formula, zero classification, and odd-node norm statement are `iteratedDeriv_fabiusReal_dyadicGrid_eq_ite`, `iteratedDeriv_fabiusReal_dyadicGrid_eq_zero_iff`, and `abs_iteratedDeriv_fabiusReal_dyadicGrid_of_odd` in `BoundedDerivatives.lean`.

## 6.2 Exact midpoint–endpoint transmutation

Subtracting the affine midpoint jet reveals an exact signed copy of the endpoint profile, not merely a finite Taylor approximation.

**Theorem 6.1** (Midpoint–endpoint transfer and the complete midpoint jet). *For every  $0 \leq h \leq \frac{1}{2}$ ,*

$$F\left(\frac{1}{2} + h\right) = \frac{1}{2} + 2h - F(h), \quad F\left(\frac{1}{2} - h\right) = \frac{1}{2} - 2h + F(h), \quad (268)$$

$$F'\left(\frac{1}{2} + h\right) = 2 - F'(h), \quad F'\left(\frac{1}{2} - h\right) = 2 - F'(h). \quad (269)$$

Moreover, every higher midpoint derivative vanishes:

$$F^{(k)}\left(\frac{1}{2}\right) = 0 \quad (k \in \mathbb{N}_0, k \geq 2). \quad (270)$$

Thus the complete Taylor jet at the midpoint is affine, even though  $F$  is not analytic there.

*Formal proof.* The value identities are `fabiusReal_midpoint_add_eq` and `fabiusReal_midpoint_sub_eq`. The first proof shows that  $h \mapsto F(\frac{1}{2} + h) + F(h)$  has constant derivative 2, then compares its value at zero; reflection gives the companion identity. The two endpoint-inclusive first-jet identities are `deriv_fabiusReal_midpoint_add_eq` and `deriv_fabiusReal_midpoint_sub_eq`; the latter reuses the global first-jet reflection law `deriv_fabiusReal_one_sub`. Finally `iteratedDeriv_fabiusReal_half_eq_zero_of_two_le` proves (270): the below-one bounded derivative formula at  $1/2$  samples the signed extension at the even integer  $2^{k-1}$ . All midpoint-transfer declarations are in `MidpointEndpointTransfer.lean`.

The value  $F(1/2) = 1/2$ , slope  $F'(1/2) = 2$ , and midpoint nonanalyticity are respectively `fabius_half`, `deriv_fabiusReal_half`, and `fabius_analyticAt_iff`.  $\square$

The transmutation transports exact integral data as well. For every real oriented radius  $a$ , global complementary reflection gives

$$\int_{1/2-a}^{1/2+a} F(x) \, dx = a. \quad (271)$$

This all-real strengthening of the frontier report's  $0 \leq a \leq 1/2$  identity is `intervalIntegral_fabiusReal_centered`. For arbitrary  $a$  the display is an oriented integral, not a nonnegative mass. On the closed half-cell define

$$D_+(h) := F\left(\frac{1}{2} + h\right) - \frac{1}{2} - 2h, \quad D_-(h) := F\left(\frac{1}{2} - h\right) - \frac{1}{2} + 2h.$$

Then  $D_+ = -F$  and  $D_- = F$  pointwise. Consequently, for every weight  $w : \mathbb{R} \rightarrow \mathbb{R}$  and  $0 \leq a \leq 1/2$ , oriented interval integration gives

$$\int_0^a w(h) D_+(h) \, dh = - \int_0^a w(h) F(h) \, dh, \quad (272)$$

$$\int_0^a w(h) D_-(h) \, dh = \int_0^a w(h) F(h) \, dh. \quad (273)$$

These are `intervalIntegral_mul_fabiusReal_midpoint_add_defect_eq_neg` and `intervalIntegral_mul_fabiusReal_midpoint_sub_defect_eq`. The constant-weight form of (272), named `intervalIntegral_fabiusReal_midpoint_add_defect_eq_neg`, is exactly the central defect integral stated in the frontier report. For  $n \in \mathbb{N}_0$ , taking  $w(h) = (a - h)^n/n!$  gives every Cauchy kernel for an anchored repeated primitive at once; this specialization is `intervalIntegral_repeatedPrimitiveKernel_fabiusReal_midpoint_add_defect_eq_neg`.

---

**Exact current boundary.** The value and derivative identities hold on the full closed half-cell, and the centered-integral identity holds for every oriented real radius. The arbitrary-weight theorem is pointwise transport under an interval integral; the named Cauchy-kernel family records the usual repeated-primitive kernels. The midpoint-transfer module itself introduces no recursive primitive operator; the next section’s normalized Volterra calculus supplies that general API, and the displayed kernel is compatible with it. These results do not by themselves assert local affineness, analytic continuation, an inverse-defect asymptotic, or a finite spline quarter-cell formula.

---

### 6.3 Exact inverse midpoint defect

Let  $I$  denote the totalized inverse of the bounded Fabius function and remove its midpoint and affine midpoint jet in two stages:

$$h(\delta) := I\left(\frac{1}{2} + \delta\right) - \frac{1}{2}, \quad E(\delta) := h(\delta) - \frac{\delta}{2}. \quad (274)$$

These are `fabiusInvMidpointOffset` and `fabiusInvMidpointDefect`. The totalization matters only outside the unit range. Their endpoint normalizations are

$$h(0) = E(0) = 0, \quad h\left(\frac{1}{2}\right) = \frac{1}{2}, \quad E\left(\frac{1}{2}\right) = \frac{1}{4}. \quad (275)$$

On the central half-cell the following identities are exact.

**Theorem 6.2** (Inverse midpoint fixed point). *For  $0 \leq \delta \leq \frac{1}{2}$ ,*

$$0 \leq h(\delta) \leq \frac{1}{2}, \quad (276)$$

$$\delta = 2h(\delta) - F(h(\delta)), \quad (277)$$

$$h(\delta) = \frac{\delta}{2} + \frac{1}{2}F(h(\delta)), \quad (278)$$

$$E(\delta) = \frac{1}{2}F(h(\delta)) = \frac{1}{2}F\left(\frac{\delta}{2} + E(\delta)\right), \quad (279)$$

$$0 \leq E(\delta) \leq \frac{1}{4}. \quad (280)$$

*In fact (276) holds for every  $\delta \geq 0$ . Globally, including the clamped tails of the totalized inverse,*

$$h(-\delta) = -h(\delta), \quad E(-\delta) = -E(\delta) \quad (\delta \in \mathbb{R}). \quad (281)$$

*Formal proof.* The four identities in (275) are `fabiusInvMidpointOffset_zero`, `fabiusInvMidpointDefect_zero`, `fabiusInvMidpointOffset_half`, and `fabiusInvMidpointDefect_half`. For nonnegative  $\delta$ , monotonicity of the totalized inverse at  $1/2$ , together with its global upper bound, gives `fabiusInvMidpointOffset_mem_Icc`. When  $\delta \leq 1/2$ , applying  $F$  to  $I(1/2 + \delta)$  and then using the midpoint–endpoint identity (268) gives `fabiusInvMidpointOffset_equation`. Elementary rearrangement yields `fabiusInvMidpointOffset_fixedPoint`, `fabiusInvMidpointDefect_eq_half_fabiusReal`, and the self-contained defect equation `fabiusInvMidpointDefect_fixedPoint`. Positivity, monotonicity, and  $F(1/2) = 1/2$  give the sharp uniform enclosure `fabiusInvMidpointDefect_mem_Icc`. Finally the global inverse reflection law gives `fabiusInvMidpointOffset_neg` and `fabiusInvMidpointDefect_neg`. All declarations are in `InverseMidpointDefect.lean`.  $\square$



---

**Exact current boundary.** The fixed-point equation is a finite identity. It is not, by itself, a proof that  $E(\delta) = O(\delta^N)$  for every  $N$ , that  $E(\delta) \sim \frac{1}{2}F(\delta/2)$ , or that logarithmic or Lambert– $W$  asymptotics transfer through the implicit argument. Those require quantitative flatness and stability estimates beyond the algebra proved here.

---

## 6.4 Normalized Volterra calculus and exact Fabius primitives

Repeated integration is most useful here in a form that remembers the base point, works for either orientation of the endpoints, and does not introduce a new normalization at order zero. Let  $E$  be a real normed vector space,  $f : \mathbb{R} \rightarrow E$ , and  $a \in \mathbb{R}$ . Write

$$(V_a f)(x) := \int_a^x f(t) \, dt, \quad (282)$$

$$L_a^0 f := f, \quad L_a^{n+1} f := V_a(L_a^n f), \quad (283)$$

$$J_a^0 f(x) := f(x), \quad J_a^{n+1} f(x) := \int_a^x \frac{(x-t)^n}{n!} f(t) \, dt. \quad (284)$$

Scalar multiplication in  $E$  is understood in the last display. These are oriented interval integrals: no hypothesis such as  $a \leq x$  occurs. In Lean they are the three public definitions `volterraPrimitive`, `iteratedPrimitive`, and `normalizedVolterra` in `NormalizedVolterra.lean`.

**Theorem 6.3** (Literal primitive ladder). *For every  $m, n \in \mathbb{N}_0$ , literal iteration satisfies*

$$L_a^0 f = f, \quad L_a^{n+1} f = V_a(L_a^n f), \quad (285)$$

$$L_a^{m+n} f = L_a^m(L_a^n f), \quad L_a^{n+1} f(a) = 0. \quad (286)$$

*If  $E$  is complete and  $f$  is continuous, then every  $L_a^n f$  is continuous and*

$$D(L_a^{n+1} f)(x) = L_a^n f(x), \quad (287)$$

$$D^k(L_a^{k+m} f) = L_a^m f, \quad (288)$$

$$D^k(L_a^n f) = L_a^{n-k} f \quad (k \leq n). \quad (289)$$

*The derivative in (287) exists in the strict sense, and  $L_a^n f$  is  $C^n$ .*

The four algebraic identities are respectively `iteratedPrimitive_zero`, `iteratedPrimitive_succ`, `iteratedPrimitive_add`, and `iteratedPrimitive_succ_self`. Continuity is `continuous_volterraPrimitive` and `continuous_iteratedPrimitive`. The strict and ordinary pointwise FTC statements, and its equality-of-functions form, are `iteratedPrimitive_succ_hasStrictDerivAt`, `iteratedPrimitive_succ_hasDerivAt`, and `deriv_iteratedPrimitive_succ`. The two cancellation formulas and the smoothness statement are `iteratedDeriv_iteratedPrimitive_add`, `iteratedDeriv_iteratedPrimitive`, and `contDiff_iteratedPrimitive`.

**Theorem 6.4** (Canonical Cauchy kernel). *The normalized kernel has the total order-zero and positive-order rules*

$$J_a^0 f(x) = f(x), \quad J_a^{n+1} f(x) = \int_a^x \frac{(x-t)^n}{n!} f(t) \, dt, \quad (290)$$

$$J_a^1 f(x) = \int_a^x f(t) \, dt, \quad J_a^{n+1} f(a) = 0. \quad (291)$$

If  $f = g$  on the closed unoriented interval between  $a$  and  $x$ , then  $J_a^n f(x) = J_a^n g(x)$ . Real scalar multiplication commutes with  $J_a^n$ , and any finite sum of interval-integrable inputs may be passed through it.

If  $E$  is complete and  $f$  is continuous, then literal iteration and the kernel agree,

$$L_a^n f = J_a^n f, \quad (292)$$

and hence

$$D(J_a^{n+1} f)(x) = J_a^n f(x) \quad \text{strictly,} \quad (293)$$

$$D^k(J_a^{k+m} f) = J_a^m f, \quad (294)$$

$$J_a^{m+n} f = J_a^m(J_a^n f). \quad (295)$$

Moreover  $J_a^n f$  is  $C^n$ .

The four unfolding and base-point declarations are `normalizedVolterra_zero`, `normalizedVolterra_succ`, `normalizedVolterra_one`, and `normalizedVolterra_succ_self`; locality, scalar compatibility, and finite additivity are `normalizedVolterra_congr`, `normalizedVolterra_smul`, and `normalizedVolterra_finsetSum`. Equation (292) is `iteratedPrimitive_eq_normalizedVolterra`. The remaining conclusions are, in the order displayed, `normalizedVolterra_succ_hasStrictDerivAt`, `deriv_normalizedVolterra_succ`, `iteratedDeriv_normalizedVolterra_add`, `normalizedVolterra_add`, and `contDiff_normalizedVolterra`.

The normalization is exactly covariant under every real affine map  $A(t) = ct + d$ :

$$J_{ca+d}^n f(cx + d) = c^n J_a^n (f \circ A)(x), \quad (296)$$

$$J_a^n (f \circ A)(x) = (c^n)^{-1} J_{ca+d}^n f(cx + d) \quad (c \neq 0). \quad (297)$$

The first, division-free form includes negative scales, whose sign is handled by the oriented integral, and even  $c = 0$ ; it also includes order zero. Neither form assumes integrability or completeness. They are `normalizedVolterra_affine` and `normalizedVolterra_comp_affine`.

Changing the base point also has an exact finite form. If  $f$  is interval integrable on the two oriented pieces from  $a$  to  $b$  and from  $b$  to  $x$ , then

$$J_a^n f(x) = J_b^n f(x) + \sum_{k=0}^{n-1} \frac{(x-b)^k}{k!} J_a^{n-k} f(b). \quad (298)$$

No ordering of  $a, b, x$  is required. At  $n = 0$  the sum is empty and both Volterra terms reduce to  $f(x)$ . This is `normalizedVolterra_basepoint_shift`. If, in addition,  $f$  vanishes on the open unoriented interval between  $b$  and  $x$ , then the local positive-order tail is zero and

$$J_a^{n+1} f(x) = \sum_{k=0}^n \frac{(x-b)^k}{k!} J_a^{n+1-k} f(b). \quad (299)$$

Only interval integrability from  $a$  to  $b$  is assumed in this corollary; no value condition is imposed at either endpoint, and the vanishing hypothesis is vacuous when  $b = x$ . The declaration is `normalizedVolterra_succ_eq_taylor_of_eq_zero`.

The hypothesis boundary is therefore sharp: affine covariance, the base-point identities, and the direct kernel algebra take values in an arbitrary real normed vector space. Completeness is used only for the strict-FTC and smooth literal-iteration layer, its identification with the direct kernel, and the Taylor reconstruction below. The four affine and base-point declarations just named are post-mainline additions checked at code checkpoint `e109088ed`.

The finite algebra of polynomial weights is especially clean when written with rising factorials. Put

$$n^{\bar{r}} := n(n+1) \cdots (n+r-1), \quad n^{\bar{0}} := 1,$$

and write  $P^{[r]}$  for the  $r$ -th Hasse derivative of a real polynomial  $P$ .

**Theorem 6.5** (Kernel raising, polynomial commutator, and Taylor reconstruction). *For every  $n, r \in \mathbb{N}_0$  and fixed endpoint  $x$ ,*

$$J_a^n((x - \cdot)^r f)(x) = n^{\bar{r}} J_a^{n+r} f(x). \quad (300)$$

*This includes  $n = 0$ : the rising factorial retains the identity when  $r = 0$  and annihilates the positive shifts when  $r > 0$ . If  $f$  is interval integrable between  $a$  and  $x$ , then*

$$J_a^n(Pf)(x) = \sum_{r=0}^{\text{natdeg } P} (-1)^r n^{\bar{r}} P^{[r]}(x) J_a^{n+r} f(x), \quad (301)$$

$$J_a^n(t^p f(t))(x) = \sum_{r=0}^p (-1)^r \binom{p}{r} n^{\bar{r}} x^{p-r} J_a^{n+r} f(x). \quad (302)$$

*Here the second left side means  $J_a^n(t \mapsto t^p f(t))(x)$ . The formulas have separate positive-order companions in which  $n$  is written  $n+1$ ; those companions have the same hypotheses and retain the literal Cauchy-kernel order.*

*Finally, if  $E$  is complete and  $g : \mathbb{R} \rightarrow E$  is  $C^{n+1}$ , then*

$$J_a^{n+1}(D^{n+1}g)(x) = g(x) - \sum_{k=0}^n \frac{(x-a)^k}{k!} D^k g(a). \quad (303)$$

*In particular, if  $D^k g(a) = 0$  for  $0 \leq k \leq n$ , then the left side is exactly  $g(x)$ .*

The positive-order and all-order kernel identities are `normalizedVolterra_succ_kernel_pow` and `normalizedVolterra_kernel_pow`. The Hasse-derivative formulas are `normalizedVolterra_succ_polynomial` and `normalizedVolterra_polynomial`; their monomial specializations are `normalizedVolterra_succ_monomial` and `normalizedVolterra_monomial`. Taylor’s formula and its zero-jet reconstruction are `normalizedVolterra_succ_iteratedDeriv_eq_sub_taylor` and `normalizedVolterra_succ_iteratedDeriv_eq_of_zero_jet`. Thus every declaration in the generic module has a literal mathematical counterpart above, including the order-zero and endpoint cases.

For a real order  $\alpha$ , define the total oriented operator

$$(I_{a+}^\alpha f)(x) := \int_a^x \frac{(x-t)^{\alpha-1}}{\Gamma(\alpha)} f(t) dt. \quad (304)$$

Here the real power is Lean’s total real-power convention. Its classical Riemann–Liouville interpretation below is restricted to  $0 < \alpha$  and  $a \leq x$ .

**Theorem 6.6** (Positive-real fractional Volterra foundation, covariance, and order raising). *Let  $E$  be a real normed vector space and  $f, g : \mathbb{R} \rightarrow E$ . For arbitrary real  $\alpha, a, x, c, d$ , the operator vanishes at its base point, depends only on the input on the closed unoriented interval between  $a$  and  $x$ , and satisfies*

$$I_{a+}^\alpha(cf)(x) = cI_{a+}^\alpha f(x), \quad I_{a+}^1 f(x) = \int_a^x f(t) dt. \quad (305)$$

*For every  $n \in \mathbb{N}_0$ , without endpoint-order or regularity assumptions,*

$$I_{a+}^{n+1} f(x) = J_a^{n+1} f(x). \quad (306)$$

If  $0 < \alpha$ ,  $a \leq x$ , and  $f$  and  $g$  are continuous on  $[a, x]$ , then the two fractional kernels are interval-integrable and

$$I_{a+}^{\alpha}(f + g)(x) = I_{a+}^{\alpha}f(x) + I_{a+}^{\alpha}g(x). \quad (307)$$

More generally, let  $a \leq b$ ,  $a \leq x$ , and let  $h$  be interval-integrable on  $[a, x]$ . If  $h = 0$  on the open tail  $(b, x)$ , then

$$\int_a^x h(t) dt = \int_a^{\min(x,b)} h(t) dt. \quad (308)$$

If  $0 < \alpha$ ,  $f$  is continuous on  $[a, x]$ , and  $f = 0$  on  $(b, x)$ , this specializes to

$$I_{a+}^{\alpha}f(x) = \int_a^{\min(x,b)} \frac{(x-t)^{\alpha-1}}{\Gamma(\alpha)} f(t) dt. \quad (309)$$

For  $0 < \alpha$ ,  $0 < \beta$ , and  $s \leq x$ , the shifted real beta kernel is interval-integrable. If  $s < x$ , its raw value is

$$\int_s^x (x-u)^{\alpha-1} (u-s)^{\beta-1} du = (x-s)^{\alpha+\beta-1} \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}. \quad (310)$$

Its Gamma-normalized form is

$$\int_s^x \frac{(x-u)^{\alpha-1}}{\Gamma(\alpha)} \frac{(u-s)^{\beta-1}}{\Gamma(\beta)} du = \frac{(x-s)^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)}. \quad (311)$$

If, moreover,  $E$  is complete,  $a \leq x$ , and  $f$  is continuous on  $[a, x]$ , then

$$I_{a+}^{\alpha+\beta}f(x) = I_{a+}^{\alpha} \left[ u \mapsto I_{a+}^{\beta}f(u) \right] (x). \quad (312)$$

For  $v \in E$ , completeness of  $E$ , and a strict endpoint  $a < x$ , the same beta calculation gives the normalized shifted-power rule

$$I_{a+}^{\alpha} \left[ t \mapsto \frac{(t-a)^{\beta-1}}{\Gamma(\beta)} v \right] (x) = \frac{(x-a)^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} v \quad (\alpha, \beta > 0), \quad (313)$$

and, for  $\rho > -1$ ,

$$I_{a+}^{\alpha} [t \mapsto (t-a)^{\rho} v] (x) = \frac{\Gamma(\rho+1)(x-a)^{\alpha+\rho}}{\Gamma(\alpha+\rho+1)} v \quad (\alpha > 0). \quad (314)$$

The constant specialization includes the degenerate endpoint:

$$I_{a+}^{\alpha} [t \mapsto v] (x) = \frac{(x-a)^{\alpha}}{\Gamma(\alpha+1)} v \quad (\alpha > 0, a \leq x). \quad (315)$$

For arbitrary real  $\alpha$ ,  $0 < c$ , and  $a \leq x$ , increasing-affine covariance requires no regularity, integrability, or completeness hypothesis:

$$I_{(ca+d)+}^{\alpha} f(cx+d) = c^{\alpha} I_{a+}^{\alpha} [t \mapsto f(ct+d)](x), \quad (316)$$

$$I_{a+}^{\alpha} [t \mapsto f(ct+d)](x) = (c^{\alpha})^{-1} I_{(ca+d)+}^{\alpha} f(cx+d). \quad (317)$$

Finally, let  $E$  be complete, let  $0 < \alpha$ ,  $a \leq x$ , and suppose that  $g$  is continuous on  $[a, x]$ , has the right-sided interior derivative  $g'$ , and that  $g'$  is interval-integrable. Gamma-normalized integration by parts gives

$$I_{a+}^{\alpha+1} g'(x) = I_{a+}^{\alpha} g(x) - \frac{(x-a)^{\alpha}}{\Gamma(\alpha+1)} g(a). \quad (318)$$

This includes  $a = x$ ; when  $g(a) = 0$ , the boundary term vanishes. Only the right derivative on the open interval is required; no two-sided derivative hypothesis is used.

The definition and ten exact theorems in `FractionalVolterra.lean` are `fractionalVolterra`, `fractionalVolterra_self`, `fractionalVolterra_congr`, `fractionalVolterra_smul`, `intervalIntegrable_fractionalVolterra_kernel`, `intervalIntegral_eq_integral_min_of_eq_zero`, `fractionalVolterra_eq_intervalIntegral_min_of_eq_zero`, `fractionalVolterra_add_input`, `fractionalVolterra_one`, and `fractionalVolterra_nat_succ`, together with `intervalIntegral_fractionalVolterra_betaKernel`. The six theorems in `FractionalVolterraSemigroup.lean` are `intervalIntegrable_fractionalVolterra_betaKernel`, `intervalIntegral_fractionalVolterra_normalizedBetaKernel`, `fractionalVolterra_normalized_rpow_smul`, `fractionalVolterra_rpow_smul`, `fractionalVolterra_const`, and `fractionalVolterra_add`. The four theorems in `FractionalVolterraCalculus.lean` are `fractionalVolterra_affine`, `fractionalVolterra_comp_affine`, `fractionalVolterra_add_one_deriv`, and `fractionalVolterra_add_one_deriv_of_eq_zero`. Thus these three reusable operator modules have one definition and twenty public theorems. They prove no classical order-zero law, reversed-endpoint fractional interpretation, semigroup theorem for merely interval-integrable inputs or noncomplete targets, Caputo or Riemann–Liouville derivative operator, complex-order theorem, nonpositive-scale affine covariance, or hierarchy-specific formula by themselves. The positive-real Fabius-CDF specialization is recorded in [Section 7.5](#); the following total definition and six specialization theorems bring the four-module total to two definitions and twenty-six public theorems.

The signed Fabius dilation law now enters only at the specialization step.

**Theorem 6.7** (Causal Rvachev fractional primitives and exact first shifts). *For every real  $\beta, x$ , define the total causal primitive*

$$U_\beta(x) := I_{(-1)_+}^\beta \text{up}(x). \quad (319)$$

*If  $0 < \beta$  and  $-1 \leq x$ , compact support gives the exact cutoff*

$$U_\beta(x) = \frac{1}{\Gamma(\beta)} \int_{-1}^{\min(x,1)} (x-t)^{\beta-1} \text{up}(t) dt. \quad (320)$$

*For every  $n \in \mathbb{N}_0$  and every real  $x$ ,*

$$U_{n+1}(x) = J_{-1}^{n+1} \text{up}(x). \quad (321)$$

*If  $0 < \alpha, 0 < \beta$ , and  $-1 \leq x$ , then*

$$U_{\alpha+\beta}(x) = I_{(-1)_+}^\alpha U_\beta(x). \quad (322)$$

*For  $0 < \alpha$ , the signed extension satisfies, for every  $x \geq 0$ ,*

$$I_{0+}^{\alpha+1} F_{\text{gl}}(x) = 2^\alpha I_{0+}^\alpha F_{\text{gl}}\left(\frac{x}{2}\right). \quad (323)$$

*The bounded function satisfies, for every  $x \in [0, 1]$ ,*

$$I_{0+}^{\alpha+1} F(x) = 2^\alpha I_{0+}^\alpha F\left(\frac{x}{2}\right). \quad (324)$$

*Finally the compactly supported fold, integrated from its left support endpoint, obeys for every  $x \geq -1$*

$$I_{(-1)_+}^{\alpha+1} \text{up}(x) = 2^\alpha I_{0+}^\alpha F\left(\frac{x+1}{2}\right). \quad (325)$$

The total definition and six theorems in `FabiusFractionalVolterra.lean` are `rvachevFractionalPrimitive`, `rvachevFractionalPrimitive_eq_intervalIntegral_min`, `rvachevFractionalPrimitive_nat_succ`, `rvachevFractionalPrimitive_add`, `fracti`

onalVolterra\_add\_one\_extendedFabius\_of\_nonneg, fractionalVolterra\_add\_one\_fabiusReal, and fractionalVolterra\_add\_one\_rvachevUp. Their domains are part of the statements: the signed formula does not cover  $x < 0$ , the bounded formula is confined to  $[0, 1]$ , the cutoff and semigroup formulas start at  $-1$ , and the support-truncated formula is asserted only at positive order.

**Theorem 6.8** (Complete integer primitive ladder of the Fabius functions). *For every  $n, p \in \mathbb{N}_0$  and  $x \in \mathbb{R}$ ,*

$$J_0^n F_{\text{gl}}(x) = 2^{\binom{n}{2}} F_{\text{gl}}\left(\frac{x}{2^n}\right), \quad (326)$$

$$J_0^n (t^p F_{\text{gl}}(t))(x) = \sum_{r=0}^p (-1)^r \binom{p}{r} n^{\bar{r}} x^{p-r} 2^{\binom{n+r}{2}} F_{\text{gl}}\left(\frac{x}{2^{n+r}}\right). \quad (327)$$

*For the bounded function the same formulas hold whenever  $x \leq 1$ :*

$$J_0^n F(x) = 2^{\binom{n}{2}} F\left(\frac{x}{2^n}\right), \quad (328)$$

$$J_0^n (t^p F(t))(x) = \sum_{r=0}^p (-1)^r \binom{p}{r} n^{\bar{r}} x^{p-r} 2^{\binom{n+r}{2}} F\left(\frac{x}{2^{n+r}}\right). \quad (329)$$

*All four identities include  $n = 0$ , arbitrary endpoint orientation, and the natural power convention  $0^0 = 1$ .*

The four displayed statements are exactly `normalizedVolterra_extendedFabius`, `normalizedVolterra_pow_mul_extendedFabius`, `normalizedVolterra_fabiusReal_of_le_one`, and `normalizedVolterra_pow_mul_fabiusReal_of_le_one` in `FabiusAntiderivatives.lean`. Their proof architecture is deliberately triangular: the global dilation equation identifies the  $n$ -th derivative of the scaled candidate; its initial Taylor jet vanishes; (303) reconstructs the candidate; and (302) supplies every natural polynomial weight. The bounded formulas use `fabiusReal_eq_extendedFabius_of_le_one`: the oriented segment from 0 to any  $x \leq 1$  remains inside the half-line on which the two functions agree.

---

The generic positive-real operator is formalized in [Theorem 6.6](#), including its positive-real, ordered-endpoint, continuous-Banach-input semigroup, shifted-power and constant evaluations, positive-scale affine covariance, and right-derivative order-raising identity. The positive-real Fabius-CDF stopped-power and unit-endpoint moment formulas are formalized in [Theorem 7.4](#), and the first signed, bounded, and up-to- $F$  fractional shifts in [Theorem 6.7](#). What remains includes the signed branch for  $x < 0$ , iterated fractional shifts, the shifted dyadic lattice and its endpoint-up wrapper, reflected or piecewise support-tail formulas, weighted hierarchy-specific monomials, semigroups for shifted or reflected hierarchy variants, a full two-branch refinement law, fractional inverse/quantile formulas, classical order-zero or reversed-endpoint interpretations, Caputo and Riemann–Liouville derivative operators, complex powers and complex order, and normal or Mellin continuation. At integer order, separately named packages for the complete up ladder based at  $-1$ , the reflected  $1 - F$  ladder, arbitrary affine copies of  $F$ , and the full piecewise polynomial tail after the support of up also remain frontier targets.

---

## 6.5 Reduced-dyadic derivative filtration

The module `DyadicDerivativeFiltration.lean` has no public definitions and exactly six theorems. The support helper `rvachevUp_eq_zero_of_one_le_abs` proves vanishing whenever  $1 \leq |x|$ . At a reduced dyadic point  $(2c+1)/2^n$ , `iteratedDeriv_rvachevUp_dyadic_eq_zero` gives zero above depth  $n$ , `iteratedDeriv_rvachevUp_dyadic_critical` gives

$$\text{up}^{(n)}\left(\frac{2c+1}{2^n}\right) = -\varepsilon_c 2^{\binom{n+1}{2}},$$

and `dyadic_depth_eq_max_nonzero_iteratedDeriv` packages the critical nonvanishing and all higher vanishing as exact denominator-depth detection.

The two newest theorems complete the below-depth case. The stronger all-real statement `iteratedDeriv_rvachevUp_eq_extendedFabius` says that for every  $m \in \mathbb{N}_0$  and  $x < 1$ ,

$$\text{up}^{(m)}(x) = 2^{\binom{m+1}{2}} F_{\text{gl}}(2^m(x+1)).$$

Its exact dyadic specialization `iteratedDeriv_rvachevUp_dyadic_below` states, for  $m < n$  and  $a < 2^n$ ,

$$\text{up}^{(m)}\left(\frac{a}{2^n}\right) = 2^{\binom{m+1}{2}} F_{\text{gl}}\left(\frac{2^n + a}{2^{n-m}}\right).$$

Thus the source proves all three regimes—below, at, and above the reduced dyadic depth—without adding a normalized-by- $\pi^m$  wrapper.

## 6.6 Dyadic derivative cells and exact value distributions

Write

$$T_n := \binom{n+1}{2}, \quad \varepsilon_k := (-1)^{\text{wt}_2(k)}.$$

For  $n, k \in \mathbb{N}_0$ , the formal cell coordinate is the total function

$$c_{n,k}(y) := -1 + \frac{2k+1+y}{2^n}. \quad (330)$$

Its geometric use is  $0 \leq k < 2^n$  and  $-1 \leq y \leq 1$ . In that range it maps  $[-1, 1]$  into  $[-1, 1]$ , and

$$c_{n,0}(-1) = -1, \quad c_{n,k}(1) = c_{n,k+1}(-1), \quad c_{n,2^n}(-1) = 1. \quad (331)$$

Thus the images are exactly the  $2^n$  adjacent closed cells from  $-1$  to  $1$ .

**Theorem 6.9** (Exact cellwise derivative tiling). *For every  $n \in \mathbb{N}_0$ ,  $0 \leq k < 2^n$ , and  $-1 \leq y \leq 1$ ,*

$$\text{up}^{(n)}(c_{n,k}(y)) = 2^{T_n} \varepsilon_k \text{up}(y), \quad (332)$$

$$\left| \text{up}^{(n)}(c_{n,k}(y)) \right| = 2^{T_n} \text{up}(y). \quad (333)$$

*In particular, the midpoint of the  $k$ -th cell has the exact signed value*

$$\text{up}^{(n)}\left(-1 + \frac{2k+1}{2^n}\right) = 2^{T_n} \varepsilon_k. \quad (334)$$

*All three formulas include  $n = 0$ , and the cell identities include both endpoints.*

*Formal proof.* The coordinate (330) is `rvachevDerivativeCell`. Its affine form, adjacent-endpoint identity, two outer endpoint identities, and closed-interval range theorem are `rvachevDerivativeCell_eq_div_add`, `rvachevDerivativeCell_one_eq_succ_neg_one`, `rvachevDerivativeCell_zero_neg_one`, `rvachevDerivativeCell_two_pow_neg_one`, and `rvachevDerivativeCell_mem_Icc`. Equations (332)–(334) are respectively `iteratedDeriv_rvachev_cell`, `abs_iteratedDeriv_rvachev_cell`, and `iteratedDeriv_rvachev_cell_zero`, all in `RvachevDerivativeDistribution.lean`.  $\square$

**Theorem 6.10** (Continuous-test derivative distributions). *Let  $E$  be a real Banach space and let  $H : \mathbb{R} \rightarrow E$  be continuous. For every  $n \in \mathbb{N}_0$ , the Bochner interval integrals satisfy*

$$\int_{-1}^1 H(\text{up}^{(n)}(x)) \, dx = 2^{-n} \sum_{k=0}^{2^n-1} \int_{-1}^1 H(2^{T_n} \varepsilon_k \text{up}(y)) \, dy, \quad (335)$$

$$\int_{-1}^1 H(|\text{up}^{(n)}(x)|) \, dx = \int_{-1}^1 H(2^{T_n} \text{up}(y)) \, dy, \quad (336)$$

$$\int_{-1}^1 H(2^{-T_n} |\text{up}^{(n)}(x)|) \, dx = \int_{-1}^1 H(\text{up}(y)) \, dy. \quad (337)$$

The signed formula is the exact  $2^n$ -term Thue–Morse mixture and includes order zero. For every positive order  $0 < n$ , sharp signed normalization collapses that finite mixture to the symmetric half-mixture

$$\int_{-1}^1 H(2^{-T_n} \text{up}^{(n)}(x)) \, dx = \frac{1}{2} \left( \int_{-1}^1 H(\text{up}(y)) \, dy + \int_{-1}^1 H(-\text{up}(y)) \, dy \right). \quad (338)$$

The restriction  $0 < n$  is essential: at order zero the normalized signed profile is the unsymmetrized function  $\text{up}$ .

Let  $\lambda_{[-1,1]}$  denote Lebesgue measure restricted to the closed interval  $[-1, 1]$ , and let  $f_{\#}\mu$  denote the pushforward of  $\mu$  by  $f$ . The normalized absolute identity is proved as the exact equality of Borel measures

$$\left( x \mapsto 2^{-T_n} |\text{up}^{(n)}(x)| \right)_{\#} \lambda_{[-1,1]} = \text{up}_{\#} \lambda_{[-1,1]}. \quad (339)$$

For  $0 < n$ , the normalized signed identity is likewise the exact Borel pushforward law

$$\left( x \mapsto 2^{-T_n} \text{up}^{(n)}(x) \right)_{\#} \lambda_{[-1,1]} = \frac{1}{2} \left( \text{up}_{\#} \lambda_{[-1,1]} + (y \mapsto -\text{up}(y))_{\#} \lambda_{[-1,1]} \right). \quad (340)$$

*Formal proof.* Equations (335)–(337) are `intervalIntegral_comp_iteratedDeriv_rvachev`, `intervalIntegral_comp_abs_iteratedDeriv_rvachev`, and `intervalIntegral_comp_normalized_abs_iteratedDeriv_rvachev`. The positive-order half-mixture (338) is `intervalIntegral_comp_normalized_iteratedDeriv_rvachev`. The pushforward equalities (339) and (340) are respectively `map_normalized_abs_iteratedDeriv_rvachev_restrict_Icc` and `map_normalized_iteratedDeriv_rvachev_restrict_Icc`. Both signed half-mixture declarations assume exactly  $0 < n$ ; the absolute declarations include order zero. All six declarations are in `RvachevDerivativeDistribution.lean`.  $\square$

**Corollary 6.11** (All signed natural-power moments). *For every  $n, m \in \mathbb{N}_0$ , ordinary natural powers satisfy the exact Boolean-cube formula*

$$\int_{-1}^1 \left( \text{up}^{(n)}(x) \right)^m \, dx = 2^{-n} (1 + (-1)^m)^n (2^{T_n})^m \int_{-1}^1 \text{up}(y)^m \, dy. \quad (341)$$

Consequently, if  $m$  is even, then for every  $n \in \mathbb{N}_0$

$$\int_{-1}^1 \left( \text{up}^{(n)}(x) \right)^m \, dx = (2^{T_n})^m \int_{-1}^1 \text{up}(y)^m \, dy, \quad (342)$$

whereas if  $m$  is odd and  $0 < n$ , then

$$\int_{-1}^1 \left( \text{up}^{(n)}(x) \right)^m \, dx = 0. \quad (343)$$

The general and even formulas include derivative order zero and moment order zero. The positive-order hypothesis in the odd formula is essential: when  $n = 0$ , the factor  $(1 + (-1)^m)^n$  is  $0^0 = 1$ , and the formula returns the generally nonzero base odd moment.



*Formal proof.* Equations (341), (342), and (343) are respectively `intervalIntegral_iteratedDeriv_rvachev_pow`, `intervalIntegral_iteratedDeriv_rvachev_pow_of_even`, and `intervalIntegral_iteratedDeriv_rvachev_pow_eq_zero_of_odd` in `RvachevDerivativeDistribution.lean`. The final theorem assumes exactly  $0 < n$  and odd  $m$ .  $\square$

**Corollary 6.12** (All nonnegative real absolute moments). *For every  $n \in \mathbb{N}_0$  and every real  $p \geq 0$ ,*

$$\int_{-1}^1 |\text{up}^{(n)}(x)|^p dx = (2^{T_n})^p \int_{-1}^1 \text{up}(y)^p dy. \quad (344)$$

*Here the powers are Lean’s real-power operation. Thus  $p = 0$  is included, with the convention  $0^0 = 1$ .*

*Formal proof.* This is `intervalIntegral_abs_iteratedDeriv_rvachev_rpow` in `RvachevDerivativeDistribution.lean`; its hypothesis is exactly  $0 \leq p$ .  $\square$

---

**Exact current boundary.** The signed API consists of the finite continuous-test-function mixture (335), the positive-order normalized symmetric half-mixture in continuous-test and Borel-pushforward form (338) and (340), and the ordinary  $\mathbb{N}_0$ -power formulas (341)–(343). The latter are distinct from the nonnegative-real `Real.rpow` absolute moments in (344). At every positive derivative order, `intervalIntegral_comp_normalized_iteratedDeriv_rvachev` identifies every continuous Banach-valued test integral of the sharply normalized signed derivative with the half-sum of the corresponding up and reflected  $-$  up integrals. The companion map `map_normalized_iteratedDeriv_rvachev_restrict_Icc` is the exact equality of their restricted-Lebesgue pushforward measures. Both theorems require exactly  $0 < n$ . The symmetric signed Borel law is therefore closed; its sharp  $0 < n$  boundary preserves the unsymmetrized  $n = 0$  profile. An independent product-space Rademacher realization is not yet packaged. Nor does the current module state the `eLpNorm` or general rearrangement-invariant norm ladder, the exact inverse-Fabius level sets, the beta-transform identities, or the later spectral layers. Those claims remain in the research-frontier report. Equation (339) is the formal measure-theoretic base from which such absolute-value corollaries can subsequently be proved. The exhaustive public surface of `RvachevDerivativeDistribution.lean` is one definition and eighteen theorems.

---

## 6.7 Effective flatness

For every  $n \in \mathbb{N}_0$ ,  $x \geq 0$ , and  $2^n x \leq 1$ , the formal development proves both

$$F(x) \leq 2^{\binom{n+1}{2}} x^n, \quad (345)$$

$$F(x) \leq \frac{2^{\binom{n+1}{2}}}{n!} x^n. \quad (346)$$

These are `fabiusReal_le_two_pow_mul_pow` in `EffectiveFlatness.lean` and `fabiusReal_le_two_pow_div_factorial_mul_pow` in `SharpFlatness.lean`. Their reflected forms at 1 and their folded forms for up are also formalized in those modules. In particular, for each natural power,

$$F(x) = o(x^n) \quad (x \rightarrow 0),$$

as stated by `fabiusReal_isLittleO_pow_at_zero` in `FabiusFlatness.lean`.

## 6.8 The precise analytic statements

The exact real-analytic locus of the bounded function is

$$F \text{ is analytic at } x \iff x \notin [0, 1]. \quad (347)$$

This is `fabius_analyticAt_iff`, with canonical specialization `canonical_fabius_analyticAt_iff`. For the other global forms the current proved statements are deliberately one-sided:

$$\text{up is not analytic at any } x \in [-1, 1], \quad (348)$$

$$F_{\text{gl}} \text{ is not analytic at any } x \in [0, 2). \quad (349)$$

They are `canonical_rvachev_not_analyticAt` and `extendedFabius_not_analyticAt` in `NowhereAnalytic.lean`. No stronger global locus is inferred here.

## 6.9 The totaled inverse

Let  $I : \mathbb{R} \rightarrow [0, 1]$  be the inverse order isomorphism on  $[0, 1]$ , extended constantly by 0 on the left and 1 on the right. This is the formal `fabiusInv` constructed from `fabiusOrderIso`. It is continuous and nondecreasing; on  $[0, 1]$  it is a strictly increasing bijection and satisfies

$$F(I(y)) = y, \quad I(F(x)) = x \quad (x, y \in [0, 1]). \quad (350)$$

For  $0 < y < 1$ , it is smooth and

$$I'(y) = \frac{1}{F'(I(y))} = \frac{1}{2 \text{ up}(2I(y) - 1)} > 0. \quad (351)$$

*Formal proof.* The range and inverse identities are `fabiusInv_mem_Icc`, `fabiusReal_fabiusInv`, and `fabiusInv_fabiusReal`. The order, continuity, and bijectivity statements are `strictMonoOn_fabiusInv`, `monotone_fabiusInv`, `continuous_fabiusInv`, and `bijOn_fabiusInv`. Interior smoothness and the derivative chain are `fabiusInv_contDiffOn_Ioo`, `deriv_fabiusInv`, `deriv_fabiusInv_eq_inv_two_mul_rvachevUp`, and `deriv_fabiusInv_pos`, all in `FabiusInverse.lean`.  $\square$

## 6.10 The exact inverse modulus

For a nonnegative length  $h$ , set

$$M_h(x) := F(x + h) - F(x).$$

Reflection gives  $M_h(1 - h - x) = M_h(x)$ . The derivative shape proved above shows that  $M_h$  is nondecreasing on  $(-\infty, (1 - h)/2]$  and nonincreasing on  $[(1 - h)/2, \infty)$ , including intervals which meet the constant tails. Consequently, whenever  $0 \leq a \leq b \leq 1$ ,

$$F(b - a) \leq F(b) - F(a). \quad (352)$$

Writing  $G = I$  for the globally clamped inverse, transport through the order isomorphism gives, for arbitrary  $u, v, x, y \in \mathbb{R}$ ,

$$|G(u) - G(v)| \leq G(|u - v|), \quad G(x + y) \leq G(x) + G(y). \quad (353)$$

The first bound is sharp. For every  $\eta \geq 0$ , the greatest output gap among pairs at input distance at most  $\eta$  is  $G(\eta)$ , both over all real inputs and after restricting both inputs to  $[0, 1]$ . Since  $G$  is clamped, the same value is  $G(\min(\eta, 1))$ .

For  $0 \leq h \leq 1$ , the corresponding exact strict uniform threshold is

$$(\forall u, v \in \mathbb{R}, |u - v| < \delta \implies |G(u) - G(v)| < h) \iff \delta \leq F(h). \quad (354)$$

No sign condition on  $\delta$  is needed; a nonpositive input radius makes the implication vacuous. Closed-threshold and contrapositive forms are also formalized.

*Formal proof.* The definition and fixed-length shape surface in `InverseModulus.lean` are `fabiusIntervalMass`, `fabiusIntervalMass_reflect`, `monotoneOn_fabiusIntervalMass_firstHalf`, and `antitoneOn_fabiusIntervalMass_secondHalf`. The endpoint inequality and its forward superadditive form are `fabiusReal_sub_le_sub` and `fabiusReal_add_le`. The ordered unit-interval, ordered global, and order-free inverse-gap forms are `fabiusInv_sub_le_sub_of_mem_Icc`, `fabiusInv_sub_le_sub_of_le`, and `fabiusInv_sub_le_sub`; inverse subadditivity is `fabiusInv_add_le`. The absolute and metric forms in (353) are `abs_fabiusInv_sub_le` and `dist_fabiusInv_le`, and saturation is `fabiusInv_min_one`. Attainment and the exact suprema are `isGreatest_abs_fabiusInv_sub`, `sSup_abs_fabiusInv_sub_eq`, `isGreatest_abs_fabiusInv_sub_Icc`, and `sSup_abs_fabiusInv_sub_Icc_eq`. Finally, the strict, closed, contrapositive, and biconditional effective-injectivity declarations are `abs_fabiusInv_sub_lt_of_abs_sub_lt_fabiusReal`, `abs_fabiusInv_sub_le_of_abs_sub_le_fabiusReal`, `fabiusReal_le_abs_sub_of_le_abs_fabiusInv_sub`, and `forall_abs_fabiusInv_sub_lt_iff`.  $\square$

This exact structural modulus does not by itself provide the closed elementary dyadic lower bound, recursive-modulus packaging, tolerant-bisection realizer, sequential-computability theorem, or input-bit asymptotics discussed in the frontier computability report.

The same module controls the curvature of the inverse without asserting differentiability at the clamped endpoints. For  $0 < y < 1$ ,

$$I''(y) = -\frac{F''(I(y))}{F'(I(y))^3}. \quad (355)$$

This is `deriv_deriv_fabiusInv`, with the corresponding derivative witness `deriv_fabiusInv_hasDerivAt`. On the closed half intervals, continuity at the endpoints and the interior derivative criterion give

$$I \text{ is strictly concave on } [0, \tfrac{1}{2}] \quad \text{and strictly convex on } [\tfrac{1}{2}, 1]. \quad (356)$$

The exact declarations are `strictConcaveOn_fabiusInv_firstHalf` and `strictConvexOn_fabiusInv_secondHalf`.

The pointwise statement is sharper. The midpoint jet is

$$I(1/2) = 1/2, \quad I'(1/2) = 1/2, \quad I''(1/2) = 0, \quad (357)$$

and, for  $0 < y < 1$ ,

$$\begin{aligned} I''(y) < 0 &\iff 0 < y < \tfrac{1}{2}, \\ I''(y) > 0 &\iff \tfrac{1}{2} < y < 1, \\ I''(y) = 0 &\iff y = \tfrac{1}{2}. \end{aligned} \quad (358)$$

The exact declarations are `fabiusInv_half`, `deriv_fabiusInv_half`, `deriv_deriv_fabiusInv_half`, and the `deriv_deriv_fabiusInv_neg_iff`, `deriv_deriv_fabiusInv_pos_iff`, and `deriv_deriv_fabiusInv_eq_zero_iff` family. The hypothesis  $0 < y < 1$  is essential in the last three statements: the totalized inverse is not differentiable at the clamping endpoints.

The inverse inherits the exact symmetry and midpoint

$$I(1 - y) = 1 - I(y), \quad I(1/2) = 1/2, \quad (359)$$

and for  $0 \leq a \leq 2^n$ ,

$$I\left(F\left(\frac{a}{2^n}\right)\right) = \frac{a}{2^n}. \quad (360)$$

These are `fabiusInv_one_sub`, `fabiusInv_half`, and `fabiusInv_fabiusDyadicUnit_of_1e`. Its exact analytic locus is likewise

$$I \text{ is analytic at } y \iff y \notin [0, 1], \quad (361)$$

by `fabiusInv_analyticAt_iff`; in the open interval its derivative never vanishes by `deriv_fabiusInv_ne_zero`.

Flatness of  $F$  becomes infinite endpoint steepness of  $I$ . For every  $n \in \mathbb{N}_0$ ,

$$y = o(I(y)^n) \quad (y \downarrow 0), \quad (362)$$

$$1 - y = o((1 - I(y))^n) \quad (y \uparrow 1). \quad (363)$$

These are `id_isLittleO_fabiusInv_pow_at_zero_right` and `one_sub_isLittleO_one_sub_fabiusInv_pow_at_one_left`. In particular, the two one-sided difference quotients diverge:

$$\frac{I(y)}{y} \longrightarrow +\infty \quad (y \downarrow 0), \quad \frac{1 - I(y)}{1 - y} \longrightarrow +\infty \quad (y \uparrow 1). \quad (364)$$

The formally proved counterparts are `tendsto_fabiusInv_div_atTop` and `tendsto_one_sub_fabiusInv_div_one_sub_atTop`. Thus the totalized inverse is not differentiable at either clamping point, exactly as stated by `fabiusInv_not_differentiableAt_zero` and `fabiusInv_not_differentiableAt_one`; no finite endpoint derivative is being smuggled into the curvature statement above. The stronger limits of the positive interior derivative itself are also formalized:

$$I'(y) \longrightarrow +\infty \quad (y \downarrow 0), \quad I'(y) \longrightarrow +\infty \quad (y \uparrow 1). \quad (365)$$

They are `tendsto_deriv_fabiusInv_atTop_at_zero_right` and `tendsto_deriv_fabiusInv_atTop_at_one_left`; these are limits through the open interval, not finite derivatives at 0 or 1.

The exact uniform modulus of this clamped inverse is now separated from the differential theory in `InverseModulus.lean`. Its exhaustive public surface is one definition, `fabiusIntervalMass`, and thirty-one theorems: `fabiusIntervalMass_reflect`, `fabiusIntervalMass_eq_zero_of_add_nonpos`, `fabiusIntervalMass_eq_zero_of_one_le`, `monotoneOn_fabiusIntervalMass_firstHalf`, `antitoneOn_fabiusIntervalMass_secondHalf`, `strictMonoOn_fabiusIntervalMass_firstHalf`, `strictAntiOn_fabiusIntervalMass_secondHalf`, `fabiusReal_sub_le_sub`, `fabiusReal_lt_fabiusIntervalMass_of_mem_Ioo`, `fabiusIntervalMass_eq_fabiusReal_iff`, `fabiusReal_sub_lt_sub`, `fabiusReal_sub_eq_sub_iff`, `fabiusReal_add_le`, `fabiusReal_add_eq_iff`, `fabiusInv_sub_le_sub_of_mem_Icc`, `fabiusInv_sub_eq_sub_iff_of_mem_Icc`, `fabiusInv_sub_le_sub_of_le`, `fabiusInv_sub_le_sub`, `fabiusInv_add_le`, `abs_fabiusInv_sub_le`, `abs_fabiusInv_sub_eq_iff_of_mem_Icc`, `dist_fabiusInv_le`, `fabiusInv_min_one`, `isGreatest_abs_fabiusInv_sub`, `sSup_abs_fabiusInv_sub_eq`, `isGreatest_abs_fabiusInv_sub_Icc`, `sSup_abs_fabiusInv_sub_Icc_eq`, `abs_fabiusInv_sub_lt_of_abs_sub_lt_fabiusReal`, `abs_fabiusInv_sub_le_of_abs_sub_le_fabiusReal`, `fabiusReal_le_abs_sub_of_le_abs_fabiusInv_sub`, and `forall_abs_fabiusInv_sub_lt_iff`. For every nonnegative interval length, the mass increment is reflection-invariant, vanishes on both constant tails, and has the global weak increase–decrease shape. When  $0 < h \leq 1$ , the maximal support-meeting branches  $[-h, (1-h)/2]$  and  $[(1-h)/2, 1]$  are respectively strictly increasing and strictly decreasing. On the unit interval, the endpoint minimum has exact strict-interior and degenerate/endpoint equality loci, and constrained superadditivity has its complete equality classification. The resulting ordered and order-free inverse-gap bounds include complete ordered and absolute equality loci on  $[0, 1]$ , while clamping creates additional cases and therefore precludes the corresponding global equality claims. Subadditivity, absolute and metric self-moduli, saturation at one, and attained exact `IsGreatest/sSup` moduli hold over all real inputs and unit-interval inputs. The final four theorems are the strict and closed effective-injectivity bounds, their contrapositive separation form, and the exact uniform strict-threshold biconditional. This structural API does not claim the computability report’s closed  $\Delta_r$  lower bound, recursive-modulus packaging, tolerant bisection realizer, sequential computability, combined computable-real-function theorem, or bit-complexity estimate.

### 6.11 Effective dyadic continuity of the inverse

The exact inverse-modulus theory has an explicit primitive-recursive dyadic specialization. Define the natural denominators

$$D_{\text{fac}}(r) = 2^{\binom{r+1}{2}}(r+1)!, \quad D_{\Delta}(r) = 2^{\binom{r+1}{2}}(r+1)^{r+1}. \quad (366)$$

They are respectively `inverseFabiusFactorialDenominator`, whose displayed closed form is `inverseFabiusFactorialDenominator_eq`, and `inverseFabiusDeltaDenominator`. For every positive  $r$ , retaining the positive zeroth term in the exact dyadic recurrence gives the stronger estimate

$$\frac{1}{2^{\binom{r}{2}}(r+1)!(2^r-1)} \leq F(2^{-r}). \quad (367)$$

This is `fabiusReal_inverse_two_pow_one_term_lower_bound`. Replacing  $2^r - 1$  by  $2^r$ , and then using  $(r+1)! \leq (r+1)^{r+1}$ , yields for every  $r \in \mathbb{N}_0$

$$D_{\text{fac}}(r)^{-1} \leq F(2^{-r}), \quad (368)$$

$$D_{\text{fac}}(r) \leq D_{\Delta}(r), \quad (369)$$

$$D_{\Delta}(r)^{-1} \leq F(2^{-r}). \quad (370)$$

The three exact declarations are `inv_inverseFabiusFactorialDenominator_le_fabiusReal`, `inverseFabiusFactorialDenominator_le_deltaDenominator`, and `inv_inverseFabiusDeltaDenominator_le_fabiusReal`. Thus the final line is the report's elementary  $\Delta_r$  bound, obtained here as a corollary of the stronger factorial estimate rather than from an infinite-product cylinder event.

Both denominator functions are primitive recursive, by `inverseFabiusFactorialDenominator_primrec` and `inverseFabiusDeltaDenominator_primrec`. For arbitrary real inputs  $u, v$ , the two corresponding strict and closed moduli are

$$|u - v| < D_{\text{fac}}(r)^{-1} \implies |I(u) - I(v)| < 2^{-r}, \quad (371)$$

$$|u - v| \leq D_{\text{fac}}(r)^{-1} \implies |I(u) - I(v)| \leq 2^{-r}, \quad (372)$$

$$|u - v| < D_{\Delta}(r)^{-1} \implies |I(u) - I(v)| < 2^{-r}, \quad (373)$$

$$|u - v| \leq D_{\Delta}(r)^{-1} \implies |I(u) - I(v)| \leq 2^{-r}. \quad (374)$$

These are `abs_fabiusInv_sub_lt_inverse_two_pow_of_lt_factorialDenominator`, `abs_fabiusInv_sub_le_inverse_two_pow_of_le_factorialDenominator`, `abs_fabiusInv_sub_lt_inverse_two_pow_of_lt_deltaDenominator`, and `abs_fabiusInv_sub_le_inverse_two_pow_of_le_deltaDenominator`. Taking  $r = n$ , the inequality  $n < 2^n$  converts the first line to the reciprocal precision convention: for every  $n > 0$ ,

$$|u - v| < D_{\text{fac}}(n)^{-1} \implies |I(u) - I(v)| < \frac{1}{n}. \quad (375)$$

Consequently `fabiusInv_effectivelyUniformContinuous` proves `EffectivelyUniformContinuous` for the totalized inverse of every bounded Fabius solution.

This is the exhaustive public surface of `FabiusInverseEffectiveContinuity.lean`: two definitions and the twelve theorems named above. The last theorem establishes only the effective-uniform-continuity clause. The module supplies no tolerant-bisection realizer or sequential-computability theorem, and therefore no combined `IsComputableRealFunction` result. It also does not construct a logarithmic scale selector  $r(n)$ , an exact least or ceiling denominator, or an input-bit complexity estimate.

## 6.12 Logarithmic reciprocal continuity of the inverse

The downstream module `FabiusInverseLogarithmicModulus.lean` replaces the coarse choice  $r = n$  by the primitive-recursive selector

$$r_{\log}(n) = \text{Nat.log2}(n) + 1. \quad (376)$$

This is `inverseFabiusLogarithmicOrder`; its equation, primitive recursiveness, least-order property for  $n > 0$ , and bound  $r_{\log}(n) \leq n$  are respectively `inverseFabiusLogarithmicOrder_eq_succ_log2`, `inverseFabiusLogarithmicOrder_primrec`, `inverseFabiusLogarithmicOrder_isLeast`, and `inverseFabiusLogarithmicOrder_le_self`. In particular, for  $n > 0$ , it is exactly the least natural  $r$  satisfying  $n < 2^r$ .

The definitions `inverseFabiusLogarithmicFactorialDenominator` and `inverseFabiusLogarithmicDeltaDenominator` take value 1 at zero and, for  $n > 0$ , are  $D_{\text{fac}}(r_{\log}(n))$  and  $D_{\Delta}(r_{\log}(n))$ . The positive-input equations are `inverseFabiusLogarithmicFactorialDenominator_of_pos` and `inverseFabiusLogarithmicDeltaDenominator_of_pos`; primitive recursiveness is proved by `inverseFabiusLogarithmicFactorialDenominator_primrec` and `inverseFabiusLogarithmicDeltaDenominator_primrec`, and their pointwise comparison is `inverseFabiusLogarithmicFactorialDenominator_le_deltaDenominator`.

For either logarithmic denominator  $d$  and every  $n > 0$ , both boundary forms are available:

$$|u - v| < d(n)^{-1} \implies |I(u) - I(v)| < \frac{1}{n}, \quad |u - v| \leq d(n)^{-1} \implies |I(u) - I(v)| < \frac{1}{n}. \quad (377)$$

The factorial declarations are `abs_fabiusInv_sub_lt_inv_nat_of_lt_logarithmicFactorialDenominator` and `abs_fabiusInv_sub_lt_inv_nat_of_le_logarithmicFactorialDenominator`; the Delta declarations are `abs_fabiusInv_sub_lt_inv_nat_of_lt_logarithmicDeltaDenominator` and `abs_fabiusInv_sub_lt_inv_nat_of_le_logarithmicDeltaDenominator`. The closed input threshold still gives a strict final output because  $2^{-r_{\log}(n)} < 1/n$ . Finally, `fabiusInv_effectivelyUniformContinuous_logarithmic` and `fabiusInv_effectivelyUniformContinuous_logarithmicDelta` separately package the factorial and report-exact Delta witnesses for `EffectivelyUniformContinuous`.

This is the exhaustive public surface of the module: three definitions and fifteen theorems. This module itself does not construct the exact endpoint- mass ceiling denominator or a tolerant-bisection realizer. The downstream modules below supply the realizer, `SequentiallyComputable`, and the combined `IsComputableRealFunction` theorem; input-bit asymptotics remain open.

## 6.13 Sharp exact endpoint-mass denominators

The next leaf, `FabiusInverseExactDyadicModulus.lean`, uses the exact rational evaluator rather than a lower estimate. At each dyadic order set

$$D_*(r) = \left\lceil \frac{1}{F(2^{-r})} \right\rceil. \quad (378)$$

The formal definition is `inverseFabiusExactDyadicDenominator`; internally it takes the natural ceiling of the reciprocal of `fabiusAtInverseTwoPow`. The evaluator bridge `fabiusAtInverseTwoPow_cast` identifies that rational with the displayed real endpoint mass for every bounded Fabius solution.

For every  $r \in \mathbb{N}_0$ , `inverseFabiusExactDyadicDenominator_pos` proves  $D_*(r) > 0$ , and `inv_inverseFabiusExactDyadicDenominator_le_fabiusAtInverseTwoPow` proves the exact rational inequality

$$D_*(r)^{-1} \leq F(2^{-r}). \quad (379)$$

More precisely, `inverseFabiusExactDyadicDenominator_isLeast` says that  $D_*(r)$  is the least positive natural  $d$  satisfying  $d^{-1} \leq F(2^{-r})$ . Consequently, for arbitrary real  $u, v$ ,

$$|u - v| < D_*(r)^{-1} \implies |I(u) - I(v)| < 2^{-r}. \quad (380)$$

This is `abs_fabiusInv_sub_lt_inverse_two_pow_of_lt_exactDyadicDenominator`.

The leastness is sharp for that fixed dyadic output target. If  $0 < d < D_*(r)$ , then `exists_fabiusInv_gap_of_lt_exactDyadicDenominator` supplies real inputs with input gap below  $d^{-1}$  and inverse-output gap exactly  $2^{-r}$ . The witnesses in the proof are the endpoint pair  $u = 0$  and  $v = F(2^{-r})$ . Thus `inverseFabiusExactDyadicDenominator_isLeast_strictModulus` packages  $D_*(r)$  as the least positive integer satisfying the universal strict implication (380).

The proof is the ceiling argument with its equality case retained. Positivity of the exact dyadic recurrence makes its reciprocal positive, hence its natural ceiling is positive. The ceiling inequality gives  $D_*(r) \geq F(2^{-r})^{-1}$ ; reciprocation gives (379), and the converse ceiling characterization proves the arithmetic leastness statement. Casting the rational bound to the reals and applying `abs_fabiusInv_sub_lt_of_abs_sub_lt_fabiusReal` proves the strict modulus. For a smaller positive  $d$ , strict failure of the ceiling inequality gives  $F(2^{-r}) < d^{-1}$ ; the endpoint pair above then has the required small input gap but output gap  $2^{-r}$ . That counterexample turns membership plus arithmetic leastness into leastness among all strict integer moduli.

For reciprocal output tolerances, the second definition is

$$D_{*,\log}(0) = 1, \quad D_{*,\log}(n) = D_*(r_{\log}(n)) \quad (n > 0). \quad (381)$$

It is `inverseFabiusExactLogarithmicDenominator`; the positive-input equation is `inverseFabiusExactLogarithmicDenominator_of_pos`. Combining the fixed dyadic strict modulus with  $2^{-r_{\log}(n)} < 1/n$  gives, for  $n > 0$ ,

$$|u - v| < D_{*,\log}(n)^{-1} \implies |I(u) - I(v)| < \frac{1}{n}, \quad (382)$$

which is `abs_fabiusInv_sub_lt_inv_nat_of_lt_exactLogarithmicDenominator`. Equation (382) is a witness only: leastness was proved for the stronger fixed target  $2^{-r_{\log}(n)}$ , not for the weaker target  $1/n$ . The value  $D_{*,\log}(0) = 1$  is only a total definition convention, and no modulus conclusion is asserted at zero. The two exact denominator maps are primitive recursive, by `inverseFabiusExactDyadicDenominator_primrec` and `inverseFabiusExactLogarithmicDenominator_primrec`. Their natural-code proof avoids assuming a separate normalized-rational computability theorem. These certificates close the recursive-denominator clause, but do not by themselves package an effective-uniform-continuity witness.

These two definitions and the ten theorems named above are the exhaustive public surface of `FabiusInverseExactDyadicModulus.lean`. In particular, the module closes exact fixed-dyadic ceiling minimality but does not prove an input-bit asymptotic or a least-denominator theorem for reciprocal target  $1/n$ .

## 6.14 Fixed-depth tolerant inversion and computability of the total inverse

The missing realizer is supplied by the generic unit-interval module `EffectiveMonotoneInverse.lean`. Its exhaustive public surface consists of exactly two definitions, `SequentiallyComputableOn` and `unitClamp`, and exactly six theorems, `unitClamp_sequentiallyComputable`, `tolerantDifference_error`, `tolerantDifference_safe_updates`, `tolerantDifference_inconclusive`, `tolerantBisection_correct`, and `effectiveInversionOn ICC`. Thus all helper state, midpoint-code, and recursion declarations used by the implementation remain private.

The bisection is a fixed-depth natural recursion, not a search for a decidable comparison of represented reals. For requested output precision  $p$ , a state at depth  $s$  either carries a bracket index  $j$ ,

representing  $[j2^{-s}, (j+1)2^{-s}]$ , or a certified midpoint numerator at denominator  $2^s$ . With  $d$  the supplied positive computable denominator, the comparison precision is

$$q = p + d(p) + 3 + s, \quad \delta = 8 \cdot 2^{-q}. \quad (383)$$

The midpoint  $(2j+1)2^{-(s+1)}$  is fed to the forward oracle by an exactly equal dyadic code at its required input grid. The forward and target names give a difference certificate of radius  $\delta/4$ . Comparing their signed natural numerators with a margin of four output-grid units therefore has three total branches: a certified positive difference keeps the left half, a certified negative difference keeps the right half, and an inconclusive comparison records the midpoint as already close enough by the inverse modulus. During the remaining recursion steps a recorded numerator is merely doubled, preserving its real value while moving it to the next dyadic grid. After exactly  $p$  steps it is consequently a numerator at denominator  $2^p$ . If no comparison was inconclusive, the certified bracket has width  $2^{-p}$ . This is the fixed-depth semantics proved uniformly by `tolerantBisection_correct`; equality at a midpoint can never cause nontermination.

The abstract theorem `effectiveInversionOn_Icc` assumes a computable dyadic forward evaluator, strict monotonicity and both interval maps, an `InvOn` pair on  $[0, 1]$ , a positive computable denominator  $d$ , and the implication

$$|u - v| < d(p)^{-1} \implies |g(u) - g(v)| < 2^{-p}.$$

It concludes `SequentiallyComputableOn` for  $g$  on the unit interval. This theorem is deliberately an inverse-modulus consumer. The adjacent gap-to-modulus layer below derives such a  $d$  from a supplied computable positive rational lower bound for every dyadic forward gap. In particular, the name `effectiveInversionOn_Icc` does not claim an arbitrary-endpoint interval algorithm without computable endpoint data.

The module `EffectiveGapInverse.lean` exposes exactly four public definitions and four public theorems: `EffectivelyUniformContinuousOn`, the structure `ComputablePositiveRationalSequence`, `ComputablePositiveRationalSequence.value`, `ComputablePositiveRationalSequence.reciprocalDenominator`, `ComputablePositiveRationalSequence.reciprocalDenominator_spec`, `inverseModulus_of_positiveRationalGap`, `effectiveInversionOn_Icc_of_computablePositiveRationalGap`, and `clampedEffectiveInversion_of_computablePositiveRationalGap`. Its sequence consists of computable positive natural numerators and denominators. The natural number

$$\frac{\text{denominator}(p)}{\text{numerator}(p)} + 1$$

is computable and positive, and its reciprocal is strictly below the encoded rational value. If  $f$  is strictly increasing on  $[0, 1]$ ,  $g$  maps that interval to itself and is inverse to  $f$  there, and

$$\alpha(p) \leq f(x + 2^{-p}) - f(x) \quad (0 \leq x \leq 1 - 2^{-p}),$$

then `inverseModulus_of_positiveRationalGap` gives the corresponding  $2^{-p}$  inverse modulus. Adding a computable dyadic evaluator for  $f$  and the interval map for  $f$  proves both `SequentiallyComputableOn` and `EffectivelyUniformContinuousOn` for  $g$  on  $[0, 1]$ . The final theorem proves exactly

$$\text{IsComputableRealFunction } (x \mapsto g(\text{unitClamp}(x))).$$

This clamped extension agrees with  $g$  on  $[0, 1]$ ; it does not assert computability or continuity of the unclamped values of an arbitrary  $g$  outside the unit interval.

The concrete module `FabiusInverseComputable.lean` has no public definition and exactly one public theorem, `fabiusInv_isComputableRealFunction`. It instantiates tolerant bisection



with `fabiusHasComputableDyadicApproximation`, the unit-interval inverse identities, and `inverseFabiusDeltaDenominator`; it then applies the computable 1-Lipschitz clamp to arbitrary input names. This proves the `SequentiallyComputable` clause for the totalized inverse. The theorem’s second clause is `fabiusInv_effectivelyUniformContinuous_logarithmicDelta`, so the final result is the full statement

$$\text{IsComputableRealFunction}(\text{fabiusInv } F \, h_F).$$

The fixed-depth module and Fabius specialization expose exactly nine public declarations, while the rational-gap module adds the eight declarations above. None adds a running-time or input-bit asymptotic theorem. The generic fixed-depth theorem still consumes its modulus; the gap theorem constructs one only after the positive rational gap sequence is supplied.

### 6.15 Precise non-elementarity

The inductive predicate `IsElementary` contains constants and the identity and is closed under arithmetic operations, composition, exponential, logarithm, trigonometric operations, and variable real powers whose elementary base is globally nonvanishing. The last closure statement is `IsElementary.rpow_of_ne_zero`; the positive-base and positive-region interfaces are `IsElementary.rpow_of_pos` and `eqOn_rpow_of_pos`. Every expression in this class has a dense real-analytic locus:

$$\text{IsElementary } g \implies \overline{\{x : g \text{ is analytic at } x\}} = \mathbb{R}.$$

This is `IsElementary.dense_analyticLocus`. It yields the proved global obstructions

$$F, \quad \text{up}, \quad F_{\text{gl}}, \quad I \quad \text{are not elementary.} \quad (384)$$

The exact declarations are `canonical_fabius_not_isElementary`, `rvachev_not_isElementary`, `extendedFabius_not_isElementary`, and `fabiusInv_not_isElementary`; local versions for  $F$  and  $I$  on  $(0, 1)$  are `canonical_fabius_not_isElementary_on_Ioo` and `canonical_fabiusInv_not_isElementary_on_Ioo`.

The larger formal class `IsElementaryOrInverse` admits an inverse-branch constructor only with an open branch domain, continuity and a right-inverse identity there, and analyticity of the chosen total extension on the interior of the complement. The canonical global functions themselves are not members of that larger class:

$$\neg \text{IsElementaryOrInverse } F, \quad \neg \text{IsElementaryOrInverse } I. \quad (385)$$

These are `fabius_not_isElementaryOrInverse` and `fabiusInv_not_isElementaryOrInverse`. More strongly in a different direction, no member of the class can represent  $F$  or  $I$  even merely on  $(0, 1)$ , by `canonical_fabius_not_isElementaryOrInverse_on_Ioo` and `canonical_fabiusInv_not_isElementaryOrInverse_on_Ioo` in `InverseNotElementary.lean`.

## 7 Moments, generating functions, and reciprocal powers

### 7.1 Even and half moments

Retain the even moments  $c_n$  from (258); in particular,  $c_0 = 1$ . Define

$$d_0 = 1, \quad d_n = n \int_0^1 t^{n-1} \text{up}(t) \, dt \quad (n \geq 1). \quad (386)$$

Both are positive rational sequences. Their exact triangular recurrences, for  $n \geq 1$ , are

$$(2n+1)(2^{2n}-1)c_n = \sum_{k=0}^{n-1} \binom{2n+1}{2k} c_k, \quad (387)$$

$$(n+1)(2^n-1)d_n = \sum_{k=0}^{n-1} \binom{n+1}{k} d_k. \quad (388)$$

These are the rational declarations `moment_succ` and `halfMoment_succ` in `Arithmetic.lean`; positivity is `moment_pos` in `Paper06487Supplement.lean` and `halfMoment_pos` in `Arithmetic.lean`. The displayed integral definitions are `integral_even_pow_mul_rvachev_eq_moment`, `moment_eq_integral_formula`, and `halfMoment_eq_integral_formula_all` in `AnalyticMoments.lean`.

The entire generating function

$$G(z) = 1 + z \int_0^1 \text{up}(t) e^{zt} dt = \sum_{n=0}^{\infty} d_n \frac{z^n}{n!} \quad (389)$$

satisfies

$$G(2z) = \frac{e^z - 1}{z} G(z), \quad (390)$$

with the removable value at  $z = 0$ . The exact formal statements are `proposition_two_formula` and `complexGeneratingFunction_eq_series_formula` in `AnalyticMoments.lean`; entire differentiability is `differentiable_complexGeneratingFunction` in `NegativeLaplaceVerticalLog.lean`.

The two sequences are linked by

$$d_n = 2^{-n} \sum_{0 \leq k \leq n/2} \binom{n}{2k} c_k, \quad (391)$$

and, at odd indices,

$$d_{2k+1} = \frac{2k+1}{2} c_k. \quad (392)$$

These are `halfMoment_eq_evenMomentSum` and `halfMoment_odd_eq_moment` in `MomentPowerSeries.lean`. Division-free natural numerators for  $c_n$  and  $d_n$ , together with their exact recurrences and odd-index bridge, are formalized by `momentNumerator`, `halfMomentNumerator`, `halfMomentNumerator_succ`, `halfMomentNumerator_odd_index_eq`, and `halfMomentNumerator_odd_index_formula`. No unproved numerical table is needed.

More explicitly, if  $C_n = \text{momentNumerator}(n)$ , then  $C_n$  is a positive integer and

$$c_n = \frac{C_n}{(2n+1)!! \prod_{j=1}^n (2^{2j}-1)}. \quad (393)$$

The exact rational identity is `moment_eq_momentNumerator_div`; the product reindexing is `evenMersenneProduct_eq_prod_Ico`, and positivity is `momentNumerator_pos`. Thus (393) records the proved natural normalization, not a claim that the displayed denominator is already reduced.

## 7.2 Full centered moment and cumulant parity

The even-indexed sequence  $c_n$  admits a canonical full-order formal packaging. Define

$$M(t) := \sum_{n \geq 0} \mu_n \frac{t^n}{n!} := \sum_{n \geq 0} c_n \frac{t^{2n}}{(2n)!}, \quad K(t) := \log M(t) = \sum_{n \geq 0} \kappa_n \frac{t^n}{n!}. \quad (394)$$

These are formal power series over  $\mathbb{Q}$ ; no analytic convergence is needed.

**Theorem 7.1** (Parity and Bernoulli–Mersenne cumulants). *For every  $n \in \mathbb{N}_0$ ,*

$$\mu_{2n} = c_n, \quad \mu_{2n+1} = 0, \quad \kappa_{2n+1} = 0. \quad (395)$$

Moreover  $\kappa_0 = 0$ , and for every  $n \geq 1$ ,

$$\kappa_{2n} = \frac{2^{2n-1} B_{2n}}{n(4^n - 1)}. \quad (396)$$

Consequently, for every  $N \geq 1$ ,

$$c_N = \frac{1}{N} \sum_{r=1}^N \binom{2N}{2r} \frac{2^{2r-1} B_{2r}}{4^r - 1} c_{N-r}. \quad (397)$$

*Formal proof.* The full moment series, its coefficients, and the factorially denormalized families are `centeredRvachevFullMomentPowerSeries`, `centeredRvachevFullMoment`, and `centeredRvachevFullCumulant`. Even recovery and odd vanishing are `centeredRvachevFullMoment_even`, `centeredRvachevFullMoment_odd`, `centeredRvachevFullCumulant_even`, and `centeredRvachevFullCumulant_odd` in `CenteredMomentParity.lean`. The generic mechanism is the involution  $t \mapsto -t$ : an even coefficient family is fixed by it, whereas an odd logarithmic coefficient changes sign. This is formalized without an integral-domain hypothesis by `SaddleExpansion.logCoeff_eq_zero_of_odd_coeff`. More generally, `SaddleExpansion.logOf_expand_of_hasSubst` shows that formal logarithms commute with every nontrivial monomial substitution  $X \mapsto X^p$  under exactly the admissibility hypothesis for substitution by  $\phi - 1$ ; this includes coefficient rings where its constant coefficient is nonzero but nilpotent. The unit-constant specialization is `SaddleExpansion.logOf_expand`.

The positive even formula (396) is `centeredRvachevEvenCumulant_eq_bernoulliMersenne_formula`, and (397) is `moment_bernoulliMersenne_recurrence`, both in `SinhDivBernoulliLog.lean`. The construction is the formal full-order family canonically attached to the already established even moments. Interpreting every  $\mu_n$  as a separately integrated ordinary moment would additionally require the corresponding all-order integral bridge; that analytic identification is not being smuggled into the formal series statement.  $\square$

## 7.3 Universal Bell and cumulant transforms

The algebra converting moments to cumulants is independent of the particular Fabius law. Over every commutative  $\mathbb{Q}$ -algebra, vanishing of all odd moments forces every odd coefficient of the formal logarithm, and hence every odd formal cumulant, to vanish. The generic declarations are `SaddleExpansion.logCoeff_eq_zero_of_odd_coeff` and `momentCumulant_eq_zero_of_odd_moments`. Let  $R$  be any commutative  $\mathbb{Q}$ -algebra and let  $a : \mathbb{N}_0 \rightarrow R$ . Write

$$(\mathcal{N}a)_n = \frac{a_n}{n!}, \quad (\mathcal{D}a)_n = n!a_n. \quad (398)$$

These changes of coefficient convention are mutually inverse. For cumulants  $\kappa : \mathbb{N}_0 \rightarrow R$  and moments  $\mu : \mathbb{N}_0 \rightarrow R$ , define the complete Bell and moment–cumulant transforms by the formal identities

$$\mathcal{B}(\kappa)_n = n![X^n] \exp \left( \sum_{j \geq 1} \kappa_j \frac{X^j}{j!} \right), \quad (399)$$

$$\mathcal{K}(\mu)_n = n![X^n] \log \left( 1 + \sum_{j \geq 1} \mu_j \frac{X^j}{j!} \right). \quad (400)$$

Here the exponential and logarithm are formal power series implemented by finite coefficient recurrences; no topology or field structure on  $R$  is being assumed.

**Proposition 7.2** (Bell recurrence and exact inversion). *For every  $n \in \mathbb{N}_0$ ,*

$$\mathcal{B}(\kappa)_0 = 1, \quad \mathcal{K}(\mu)_0 = 0, \quad (401)$$

$$\mathcal{B}(\kappa)_{n+1} = \sum_{j=0}^n \binom{n}{j} \kappa_{j+1} \mathcal{B}(\kappa)_{n-j}. \quad (402)$$

*At every index, including zero, the two composites are exactly*

$$\mathcal{B}(\mathcal{K}(\mu))_n = \begin{cases} 1, & n = 0, \\ \mu_n, & n > 0, \end{cases} \quad \mathcal{K}(\mathcal{B}(\kappa))_n = \begin{cases} 0, & n = 0, \\ \kappa_n, & n > 0. \end{cases} \quad (403)$$

*Consequently the transforms are inverse as functions when  $\mu_0 = 1$  and  $\kappa_0 = 0$ , respectively. More precisely, if  $f : R \rightarrow S$  is a morphism of commutative  $\mathbb{Q}$ -algebras, then*

$$f(\mathcal{B}_R(\kappa)_n) = \mathcal{B}_S(f \circ \kappa)_n, \quad f(\mathcal{K}_R(\mu)_n) = \mathcal{K}_S(f \circ \mu)_n. \quad (404)$$

*More generally, if  $E$  and  $A$  are ordinary coefficient sequences for the formal exponent and mass series, then the coefficient-convention bridge is*

$$\mathcal{B}(\mathcal{D}E) = \mathcal{D}(\expCoeff(E)), \quad \mathcal{K}(\mathcal{D}A) = \mathcal{D}(\logCoeff(A)). \quad (405)$$

*Thus the same theorem applies without modification to polynomial rings, function rings, and scalar extensions, not only to real- or rational-valued moment sequences.*

*Formal proof.* The operations  $\mathcal{N}, \mathcal{D}, \mathcal{B}, \mathcal{K}$  are respectively `factorialNormalize`, `factorialDenormalize`, `completeBellPolynomial`, and `momentCumulant`. The two identities following (398) are `factorialNormalize_factorialDenormalize` and `factorialDenormalize_factorialNormalize`. The zero values and (402) are `completeBellPolynomial_zero`, `momentCumulant_zero`, and `completeBellPolynomial_succ`.

Equation (403) is `completeBellPolynomial_momentCumulant` together with `momentCumulant_completeBellPolynomial`; its normalized function-level forms are `completeBellPolynomial_momentCumulant_of_zero_eq_one` and `momentCumulant_completeBellPolynomial_of_zero_eq_zero`. The bridge (405) is `completeBellPolynomial_factorialDenormalize` and `momentCumulant_factorialDenormalize`. Finally, functoriality is (404), formalized by `map_completeBellPolynomial` and `map_momentCumulant`. All declarations are in `MomentCumulantAlgebra.lean` and assume only a commutative  $\mathbb{Q}$ -algebra.  $\square$

**Proposition 7.3** (Finite polynomial integrals from moments and formal cumulants). *Let  $\Omega$  be a measurable space, let  $\mu$  be any measure on it, let  $S$  be a semiring with a semiring homomorphism  $\varphi : S \rightarrow \mathbb{R}$ , and let  $Z : \Omega \rightarrow \mathbb{R}$ . For  $p \in S[X]$ , write*

$$p_n = [X^n]p, \quad p_\varphi(x) = \sum_{n \in \text{supp}(p)} \varphi(p_n)x^n, \quad m_n = \text{moment}_\mu(Z; n).$$

Here  $\text{moment}_\mu$  uses Mathlib’s totalized Bochner-integral convention; whenever  $Z^n$  is integrable it is the ordinary integral  $\int_\Omega Z(\omega)^n d\mu(\omega)$ . Assume only that  $Z^n$  is integrable for  $n \in \text{supp}(p)$ , and put  $\kappa = \mathcal{K}(m)$ . Then

$$\int_\Omega p_\varphi(Z(\omega)) d\mu(\omega) = \sum_{n \in \text{supp}(p)} \varphi(p_n)m_n, \quad (406)$$

$$= \sum_{n \in \text{supp}(p)} \varphi(p_n)\mathcal{B}(\kappa)_n + \varphi(p_0)(m_0 - 1). \quad (407)$$

If  $m_0 = 1$ , the correction in (407) vanishes. This holds automatically when  $\mu$  is a probability measure.

*Formal proof.* All four public identifiers share the prefix `integral_eval2_eq_sum_`. In the order of the two displayed equations, the  $m_0 = 1$  corollary, and the probability-measure corollary, their suffixes are

- (i) `moment`;
- (ii) `completeBell_momentCumulant_with_mass_correction`;
- (iii) `completeBell_momentCumulant_of_moment_zero_eq_one`;
- (iv) `completeBell_momentCumulant`.

They are declared in `PolynomialExpectationCumulant.lean`. They require neither boundedness nor independence, and no integrability assumptions for powers outside the finite support of  $p$ . They formalize the finite moment-to-Bell rewrite used by endpoint transfer, not the analytic transfer expansion or its remainder estimates.  $\square$

## 7.4 Rvachev moment–Appell deconvolution and polynomial synthesis

In the current compiled tree, `RvachevMomentAppell.lean` specializes the formal Bell and Appell engines to the Rvachev probability law. Its six public definitions are `rvachevRawMomentRat`, `rvachevReciprocalMomentRat`, `rvachevAppellPolynomialRat`, `rvachevAppellPolynomial`, and `rvachevDeconvolvedPolynomial`, together with `rvachevDeconvolutionLinearMap`. Its thirty-three public theorems are `rvachevRawMomentRat_zero`, `rvachevRawMomentRat_even`, `rvachevRawMomentRat_odd`, `rvachevReciprocalMomentRat_zero`, `binomialConv_rvachevRawMomentRat_reciprocal`, `rvachevReciprocalMomentRat_eq_completeBellPolynomial`, `monic_rvachevAppellPolynomialRat`, `natDegree_rvachevAppellPolynomialRat`, `rvachevAppellPolynomial_eq_poly_cast`, `monic_rvachevAppellPolynomial`, `natDegree_rvachevAppellPolynomial`, `eval_rvachevAppellPolynomial_add`, `integral_pow_mul_rvachev_eq_rvachevRawMomentRat_cast`, `integral_eval_rvachevAppellPolynomial_add_mul_rvachev`, `rvachevDeconvolutionLinearMap_apply`, `rvachevDeconvolvedPolynomial_zero`, `rvachevDeconvolvedPolynomial_add`, `rvachevDeconvolvedPolynomial_smul`, `rvachevDeconvolvedPolynomial_finsetSum`, `rvachevDeconvolvedPolynomial_C_mul`, `rvachevDeconvolvedPolynomial_monomial`, `rvachevDeconvolvedPolynomial_X_pow`, `coeff_rvachevDeconvolvedPolynomial_natDegree`, `natDegree_rvachevDeconvolvedPolynomial_le`, `natDegree_rvachevDeconvolvedPolynomial`, `leadingCoeff_rvachevDeconvolvedPolynomial`, `rvachevDeconvolvedPolynomial_eq_zero_iff`, `rvachevDeconvolutionLinearMap_injective`,

`rvachevDeconvolvedPolynomial_injective`, `integral_eval_rvachevDeconvolvedPolynomial_add_mul_rvachev`, `integral_eval_rvachevDeconvolvedPolynomial_sub_mul_rvachev`, `integral_eval_rvachevAppellPolynomial_sub_mul_rvachev`, and `integral_eval_rvachevAppellPolynomial_mul_rvachev_eq_zero`.

Thus the full rational raw-moment sequence has the executable even moments at even indices and zero at odd indices; its formal binomial-convolution reciprocal is the corresponding complete-Bell family. The rational and real Rvachev reciprocal-moment Appell polynomials are monic of exact degree. Write  $A_n = \text{rvachevAppellPolynomial } n$ . If  $u_F = \text{rvachevUp } F$  for any bounded Fabius solution, their analytic bridges are

$$\int_{\mathbb{R}} A_n(x+y)u_F(y) dy = x^n, \quad (408)$$

$$\int_{\mathbb{R}} (\mathcal{D}_{\text{up}}P)(x+y)u_F(y) dy = P(x), \quad (409)$$

where  $\mathcal{D}_{\text{up}}P$  denotes `rvachevDeconvolvedPolynomial` $P$ . Reflection invariance of the even Rvachev density gives the additional named centered identities

$$\int_{\mathbb{R}} A_n(x-y)u_F(y) dy = x^n, \quad (410)$$

$$\int_{\mathbb{R}} (\mathcal{D}_{\text{up}}P)(x-y)u_F(y) dy = P(x) \quad (411)$$

These are respectively `integral_eval_rvachevAppellPolynomial_sub_mul_rvachev` and `integral_eval_rvachevDeconvolvedPolynomial_sub_mul_rvachev`. For every positive  $n$ , `integral_eval_rvachevAppellPolynomial_mul_rvachev_eq_zero` further says that  $A_n$  has Rvachev mean zero. This coefficientwise operation sends a monomial and  $X^n$  to the correspondingly scaled and unscaled Rvachev–Appell polynomial. Its triangular top term is exact: the coefficient in the original natural degree is unchanged, so both natural degree and leading coefficient are preserved. Its kernel is trivial, and both the linear-map package and the underlying operation are injective. The linear-map package proves preservation of zero, addition, real scalar multiplication, finite sums, and multiplication by a constant polynomial; these are algebraic identities and do not assert an analytic reciprocal MGF or Appell generating series, a literal differential-operator expansion, reciprocal/deconvolution parity, or the displayed low reciprocal coefficients.

In the current compiled tree, `RvachevPolynomialSynthesis.lean` exports no public definitions and exactly five public theorems: `tsum_rvachevDeconvolvedPolynomial_mul_shifted_rvachevUp`, `normalized_tsum_rvachevDeconvolvedPolynomial_mul_shifted_rvachevUp`, `sum_Ioo_rvachevDeconvolvedPolynomial_mul_shifted_rvachevUp`, `normalized_sum_Ioo_rvachevDeconvolvedPolynomial_mul_shifted_rvachevUp`, and `normalized_tsum_shifted_rvachevDeconvolvedPolynomial_mul_rvachevUp`. For  $M \in \mathbb{N}_0 \setminus \{0\}$  and  $\deg P \leq \nu_2(M)$ , they give

$$\sum_{k \in \mathbb{Z}} (\mathcal{D}_{\text{up}}P)(k/M)u_F(x - k/M) = MP(x), \quad (412)$$

$$P(x) = \frac{1}{M} \sum_{-2M < k < 2M} (\mathcal{D}_{\text{up}}P)(k/M)u_F(x - k/M), \quad -1 \leq x \leq 1. \quad (413)$$

The first sum is Lean’s global `tsum`; compact support makes it pointwise finite. The second is the exact uniform support truncation, including the vanishing boundary atoms.

The fifth theorem has arbitrary real phase  $\theta$  and arbitrary real evaluation point  $x$ :

$$P(x) = \frac{1}{M} \sum_{k \in \mathbb{Z}} (\mathcal{D}_{\text{up}}P)\left(x - \frac{\theta + k}{M}\right) u_F\left(\frac{\theta + k}{M}\right). \quad (414)$$

The displayed bilateral sum is Lean’s `tsum`; since  $\text{supp}(u_F) \subseteq [-1, 1]$ , a summand can be nonzero only when  $|\theta + k| \leq M$ , so only finitely many integers  $k$  contribute. The theorem assumes exactly  $M \neq 0$  and  $\deg P \leq \nu_2(M)$ . Thus the specialization  $M = 2^N$  formalizes arbitrary-phase reproduction of every polynomial  $P \in \mathbb{R}[X]$  with  $\deg P \leq N$ , equivalently every monomial  $X^n$  for  $0 \leq n \leq N$ .

The compiled module `RvachevSuperconvergentSynthesis.lean` adds exactly one public definition and eight public theorems. Its definition `IsRvachevSuperconvergentPhase` selects  $0, 1/2$  when  $\nu_2(M) + 1$  is odd and  $1/4, 3/4$  when it is even. The theorem `isRvachevSuperconvergentPhase_two_pow_iff` rewrites this for  $M = 2^N$ : even  $N$  selects the endpoint phases and odd  $N$  the quarter phases. The remaining seven theorems are `tsum_quarter_monomial_eq_integral_of_even_deg`, `tsum_three_quarters_monomial_eq_integral_of_even_deg`, `tsum_shifted_monomial_eq_integral_superconvergent`, `tsum_shifted_polynomial_eq_integral_superconvergent`, `integral_polynomial_mul_rvachevUp_eq_normalized_tsum_superconvergent`, `normalized_tsum_shifted_rvachevDeconvolvedPolynomial_mul_rvachevUp_superconvergent`, and `normalized_tsum_shifted_rvachevAppellPolynomial_mul_rvachevUp_superconvergent`. For every nonzero natural  $M$ , they prove selected-phase monomial and polynomial identities through degree  $\nu_2(M) + 1$ , physical quadrature, deconvolved-polynomial reconstruction, and the Appell specialization. This arbitrary- $M$  result is stronger than the manuscript’s dyadic-only claim. The phase predicate uses exact real representatives rather than congruence classes modulo one, and the module asserts neither a complete phase classification nor maximality, positivity, or rationality of the quadrature.

The current-tree module `LagrangeRvachevSynthesis.lean` has the exact public inventory of two definitions, `lagrangeRvachevDecoder` and `lagrangeRvachevAtomCoefficient`, and seven theorems, `natDegree_lagrangeBasis_le_card_sub_one`, `natDegree_lagrangeInterpolate_le_card_sub_one`, `normalized_sum_Ioo_lagrangeRvachevDecoder_mul_shifted_rvachevUp`, `normalized_sum_Ioo_lagrangeRvachevDecoder_eval_node`, `lagrangeRvachevAtomCoefficient_eq_deconvolved_interpolate`, `sum_Ioo_lagrangeRvachevAtomCoefficient_mul_shifted_rvachevUp`, and `sum_lagrangeRvachevDecoder_eq_one`. The decoder samples the Rvachev deconvolution of a finite Lagrange basis, and the atom coefficient is its  $1/M$ -normalized linear action on nodal data. The first two theorems bound the basis and interpolant degrees by one less than the node count without a distinctness hypothesis. If  $M \neq 0$  and that degree bound is at most  $\nu_2(M)$ , the next four identities give exact finite cardinal synthesis on  $[-1, 1]$ , its componentwise Kronecker-delta evaluation at distinct in-range nodes, the coefficient/deconvolution identity, and reconstruction of the full Lagrange interpolant. The last theorem says that each fixed lattice-sample decoder row has total mass one for a nonempty distinct node family. Consequently generic finite-node synthesis, componentwise biorthogonality, and the exact finite interpolation loop are proved. This does not supply the geometric-node Gaussian  $q$ -binomial/ $q$ -Pochhammer or elementary-symmetric closed form for decoder entries, a bundled matrix/right-inverse or stochastic matrix wrapper, or an optimal or minimum-variation decoder. The geometric closed form is supplied by the next module and the typed right-inverse wrapper by `LagrangeRvachevMatrix.lean`; optimization remains open.

`RvachevAppellHasse.lean` has exactly one public definition, `Appell.polynomialTransform`, and fourteen public theorems: `Appell.polynomialTransform_apply`, `Appell.polynomialTransform_monomial`, `Appell.polynomialTransform_eq_sum_hasseDeriv_of_natDegree_lt`, `Appell.polynomialTransform_eq_sum_hasseDeriv`, `rvachevReciprocalMomentRat_odd`, `rvachevDeconvolutionLinearMap_eq_appellPolynomialTransform`, `rvachevDeconvolvedPolynomial_eq_sum_even_hasseDeriv`, `eval_hasseDeriv_prod_X_sub_C_eq_elementarySymmetricEval`, `eval_rvachevDeconvolvedPolynomial_prod_X_sub_C`, `eval_rvachevDeconvolvedPolynomial_qFallingPower`, `lagrangeBasis_eq_nodalWeight_mul_prod_X_sub_C`, `lagrangeRvachevDecoder_eq_nodalWeight_mul_sum`, `geometric_nodalWeight_eq_geometricQPochhammer`, and `geometric_lagrangeRvachevDecoder_eq`. Over a commutative semiring, the coefficientwise

linear transform sends monomials to an arbitrary Appell family and is exactly a finite sum of Hasse derivatives. The Rvachev specialization proves that the formal reciprocal moment sequence vanishes in odd degree and reduces deconvolution to the even Hasse derivatives. Taylor’s coefficient theorem and Vieta’s formula identify the Hasse derivatives of a finite root product with complementary elementary symmetric functions. In particular, `eval_rvachevDeconvolvedPolynomial_qFallingPower` is the exact  $q$ -falling-power formula of `gq:prop:q-Appell-falling`.

The field-valued Lagrange factorization and nodal-weight identity, followed by the real geometric specialization `geometric_lagrangeRvachevDecoder_eq`, give the complete Gaussian  $q$ -Pochhammer prefactor times the finite even reciprocal-moment sum in `gq:thm:gaussian-Appell-1-decoder`. These formulas are algebraically total even when  $c = 0$ ,  $q = 0$ , or nodes collide, because the Lagrange basis uses totalized inversion; on that locus it is not a cardinal basis. The manuscript’s cardinal interpretation retains  $c > 0$ ,  $0 < q < 1$ , a nonzero dyadic mesh, and the degree and interval conditions of the separate synthesis theorems. Composing the  $q$ -falling formula with `normalized_sum_Ioo_rvachevDeconvolvedPolynomial_mul_shifted_rvachevUp`, and the decoder formula with `normalized_sum_Ioo_lagrangeRvachevDecoder_mul_shifted_rvachevUp`, makes both cited comb results Exact. The coefficients `rvachevReciprocalMomentRat` are the formal reciprocal-moment sequence; no convergence theorem for an analytic reciprocal MGF is asserted. Atom reconstruction remains owned by those synthesis declarations, and this leaf adds no larger matrix right-inverse or decoder-optimality statement.

#### 7.4.1 Finite dyadic Appell prefixes and exact recovery

The compiled leaf `FinitePrefixAppellRecovery.lean` constructs the finite-prefix moment sequences rather than assuming the desired formulas. Its eleven public definitions are `unitUniformRawMomentRat`, `centeredUnitUniformRawMomentRat`, `dyadicPrefixScaleRat`, `dyadicPrefixMomentRat`, `uncenteredDyadicPrefixMomentRat`, `centeredDyadicPrefixMomentRat`, `kabayaIriAppellPolynomialRat`, `uncenteredDyadicPrefixAppellPolynomialRat`, `centeredDyadicPrefixAppellPolynomialRat`, `uncenteredDyadicPrefixAppellScalePolynomialRat`, and `centeredDyadicPrefixAppellScalePolynomialRat`. The generic prefix recursively convolves the exact moments of the first  $N$  uniform digits at scales  $2^{-1}, \dots, 2^{-N}$ ; depth zero is the point mass at zero. The uncentered digit has moment  $1/(r+1)$ , while the centered digit has that value in even degree and zero in odd degree.

Write  $P_n = \text{kabayaIriAppellPolynomialRat } n$ ,  $A_n = \text{rvachevAppellPolynomialRat } n$ , and use superscript  $[N]$  for the corresponding prefix Appell polynomial. The full one-step moment recurrences yield exact prefix–tail binomial factorizations. Reciprocal uniqueness and the generic Appell transform then give

$$P_n^{[N]}(X) = \sum_{r=0}^n \binom{n}{r} \mu_r 2^{-Nr} P_{n-r}(X), \quad \mu_r = \text{halfMoment } r, \quad (415)$$

$$A_n^{[N]}(X) = \sum_{r=0}^{\lfloor n/2 \rfloor} \binom{n}{2r} m_{2r} 4^{-Nr} A_{n-2r}(X), \quad m_{2r} = \text{moment } r. \quad (416)$$

These are `uncenteredDyadicPrefixAppellPolynomialRat_eq_sum` and `centeredDyadicPrefixAppellPolynomialRat_eq_sum_even`. Their reusable algebraic engine consists of `Appell.poly_binomialConv`, `Appell.binomialConv_dilate`, and `Appell.dilate_dilate`; normalization and tail factorization are `dyadicPrefixMomentRat_zero`, `uncenteredDyadicPrefixMomentRat_zero`, `centeredDyadicPrefixMomentRat_zero`, `dyadicPrefixMomentRat_binomialConv_tail`, `binomialConv_uncenteredDyadicPrefixMomentRat_tail`, and `binomialConv_centeredDyadicPrefixMomentRat_tail`.

Promoting the scale to a formal outer variable  $Q$  produces elements of  $\mathbb{Q}[X][Q]$ , implemented as `Polynomial (Polynomial  $\mathbb{Q}$ )`. Evaluation at  $Q = 2^{-N}$  and  $Q = 4^{-N}$  recovers the two prefix



polynomials by `uncenteredDyadicPrefixAppellPolynomialRat_eq_eval_scale` and `centeredDyadicPrefixAppellPolynomialRat_eq_eval_scale`. Their exact *outer polynomial-valued* degrees are

$$\text{natDegree}_Q P_n^{[\cdot]} = n, \quad \text{natDegree}_Q A_n^{[\cdot]} = \lfloor n/2 \rfloor, \quad (417)$$

by `natDegree_uncenteredDyadicPrefixAppellScalePolynomialRat` and `natDegree_centeredDyadicPrefixAppellScalePolynomialRat`. The polynomial-valued qualifier is essential. After fixing the inner evaluation point  $X = x$ , the outer degree may fall; for centered odd  $n$ , the top inner Appell coefficient vanishes at  $x = 0$ . Thus no universal exact fixed- $x$  degree statement is inferred from (417).

Finally let  $w_{d,j}(q) = \text{geometricLagrangeWeight } q \, d \, j$ . Evaluating the two scale polynomials at zero gives, for every starting depth  $N$ ,

$$P_n(X) = \sum_{j=0}^n w_{n,j}(1/2) P_n^{[N+j]}(X), \quad (418)$$

$$A_n(X) = \sum_{j=0}^{\lfloor n/2 \rfloor} w_{\lfloor n/2 \rfloor, j}(1/4) A_n^{[N+j]}(X). \quad (419)$$

They are `kabayaIriAppellPolynomialRat_eq_sum_prefix` and `rvachevAppellPolynomialRat_eq_sum_prefix`. Hence the full uncentered degree- $n$  polynomial is recovered from any  $n + 1$  consecutive prefixes, and the full centered polynomial from any  $\lfloor n/2 \rfloor + 1$  consecutive prefixes, with no limiting argument. The seventeen theorems listed in the preceding three paragraphs exhaust the theorem surface of the module. They make both `is:p2:thm:finite-prefix-expansion` and `is:p2:thm:exact-recovery Exact`. This is exact rational coefficient algebra; no analytic MGF convergence theorem is claimed.

The subsequent `RvachevLagrangeNodesOnly.lean` leaf has the one public definition `rvachevDeconvolvedPolynomialRat` and exactly fourteen public theorems: `map_rvachevDeconvolvedPolynomialRat`, `rvachevDeconvolvedPolynomial_eq_sum_appell`, `eval_rvachevDeconvolvedPolynomial_eq_sum_even_iterateDerivative`, `rvachevDeconvolvedPolynomial_prod_X_sub_C_eq_sum_appell`, `eval_rvachevDeconvolvedPolynomial_lagrangeBasis_eq_sum_even_iterateDerivative`, `eval_rvachevDeconvolvedPolynomial_lagrangeBasis_eq_nodalWeight_mul_sum_appell`, `lagrangeRvachevDecoder_eq_nodalWeight_mul_sum_appell`, `map_lagrangeBasis_ratCast`, `map_rvachevDeconvolvedPolynomialRat_lagrangeBasis`, `lagrangeRvachevDecoder_eq_ratCast`, `rvachevRawMomentRat_eq_centeredRvachevFullMoment`, `momentCumulant_rvachevRawMomentRat_eq_centeredRvachevFullCumulant`, `momentCumulant_rvachevRawMomentRat_even_eq_bernoulliMersenne`, and `rvachevReciprocalMomentRat_eq_completeBellPolynomial_neg_centeredCumulant`. By composition, not by a single wrapper, these declarations make `cor:lag-nodes-only Exact/Complete`: they supply the distinct-node ordinary-derivative form, the raw omitted-node Appell form, rational descent and rational lattice samples, and the finite complete-Bell/Bernoulli–Mersenne coefficient identification. Rationality is asserted only at rational evaluation points; the sampled identity is total at  $M = 0$ , while synthesis retains  $M \neq 0$  and its degree and interval hypotheses. The Bell statement is formal coefficient algebra, not analytic reciprocal-MGF convergence. No promotion of the larger cardinal/right-inverse rows or decoder optimality is implied.

The downstream `LagrangeRvachevMatrix.lean` module packages precisely this scalar loop as typed rectangular matrices. Its exhaustive four-definition/abbreviation surface is `rvachevAtomIndexSet`, `RvachevAtomIndex`, `lagrangeRvachevEncoderMatrix`, and `lagrangeRvachevDecoderMatrix`. The atom type is the subtype of the exact finite block

$$\{k \in \mathbb{Z} : -2M < k < 2M\},$$

with the two endpoints excluded. If rows are indexed by the finite node family and columns by this block, then the encoder and decoder entries are

$$A_{ik} = M^{-1} \text{up}(v_i - k/M), \quad D_{kj} = \text{lagrangeRvachevDecoder}(s, v, M, k, j).$$

The six public theorems are `lagrangeRvachevEncoderMatrix_nonneg`, `sum_lagrangeRvachevEncoderMatrix_row_eq_one`, `sum_lagrangeRvachevDecoderMatrix_row_eq_one`, `lagrangeRvachevEncoderMatrix_mul_decoderMatrix`, `exists_neg_entry_of_rightInverse_of_row_overlap`, and `exists_lagrangeRvachevDecoderMatrix_entry_neg_of_row_overlap`.

Encoder nonnegativity requires neither distinct nodes nor a nonzero mesh. Its row-sum-one theorem requires `IsFabius` for the chosen bounded solution,  $M \neq 0$ , and only that the selected nodes lie in  $[-1, 1]$ . Decoder row unitality instead requires a nonempty distinct node family, but no Fabius or mesh-admissibility hypothesis. For distinct nodes all in  $[-1, 1]$ ,  $M \neq 0$ , and  $|s| - 1 \leq \nu_2(M)$ , the scalar Kronecker identities give the typed right-inverse equation  $AD = I$ .

The generic overlap theorem is deliberately sharper about its algebraic assumptions. For finite matrices, entrywise  $A \geq 0$ , row-unital  $D$ ,  $AD = I$ , and one column of  $A$  strictly positive in two distinct rows force some entry of  $D$  to be negative; encoder row normalization is not used. The final theorem applies this implication to the Rvachev matrices, conditional on an actual strictly positive overlap. It does not prove that every admissible node family has such an overlap. This finite layer makes no claim about the range, kernel, rank, trace, characteristic polynomial, or Cauchy–Binet data of  $DA$ . The preceding Hasse leaf supplies the scalar geometric closed form, but neither module proves an optimal or minimum-variation decoder.

At the focused-build checkpoint of 30 August 2026, `CompositeMeshSharpness.lean` exports one public definition, `rvachevCombExactThrough`, and seven public theorems: `exists_shift_tsum_shifted_monomial_ne_integral_nat_real`, `rvachevCombExactThrough_iff_padicValNat`, `rvachevCombExactThrough_iff_pow_two_dvd`, `rvachevCombExactThrough_two_pow`, `two_pow_le_of_rvachevCombExactThrough`, `isLeast_rvachevCombExactThrough`, and `isLeast_rvachevCombExactThrough_even`. The definition requires a nonzero natural mesh  $M$  and, for every real polynomial  $P$  of natural degree at most  $d$  and every real shift  $\theta$ , the identity

$$\sum_{k \in \mathbb{Z}} P(\theta + k) u_F((\theta + k)/M) = \int_{\mathbb{R}} P(x) u_F(x/M) dx. \quad (420)$$

The two classification theorems prove

$$\begin{aligned} \text{rvachevCombExactThrough}(F, M, d) &\iff M \neq 0 \text{ and } d \leq \nu_2(M) \\ &\iff M \neq 0 \text{ and } 2^d \mid M. \end{aligned} \quad (421)$$

For every nonzero  $M$ , the first theorem supplies a real shift at which the monomial of degree  $\nu_2(M) + 1$  fails (420). The remaining order theorems put  $2^d$  in the exact class, bound every member below by  $2^d$ , and prove that  $2^d$  is its least member; at even cutoff  $2N$ , this least universal mesh is  $4^N$ .

The word *universal* is essential. These declarations quantify over the complete polynomial space through the cutoff. They do not prove that  $2^d$  is a least mesh for one individual  $P_d$ , that  $4^N$  is least for one particular Legendre partial sum  $S_N$ , or that a target-adapted formula cannot use a coarser mesh.

Together these modules prove formal coefficient reciprocity, polynomial smoothing/synthesis, and the sharp universal comb threshold. They do not prove an analytic reciprocal-MGF or Appell generating series, a literal reciprocal-MGF differential-operator identity, parity of the reciprocal or deconvolved families, the displayed low reciprocal coefficients, or any target-by-target minimality statement for the synthesis formulas.

## 7.5 The probability–Laplace bridge

Let  $\mu_X$  be the distribution of the weighted sum  $X$  from (34). Its survival function on the nonnegative half-line is exactly

$$\mu_X((t, \infty)) = \text{up}(t) \quad (t \geq 0). \quad (422)$$

More generally, let  $E$  be a real Banach space and let  $k : [0, 1] \rightarrow E$  be Bochner integrable. Fubini gives the survival-kernel identity

$$\int_{[0,1]} \left( \int_0^x k(t) dt \right) d\mu_X(x) = \int_0^1 \text{up}(t) k(t) dt. \quad (423)$$

For every  $c \in [0, 1]$ , truncating the kernel gives the stronger endpoint family

$$\int_{[0,1]} \left( \int_0^{\min\{x,c\}} k(t) dt \right) d\mu_X(x) = \int_0^c \text{up}(t) k(t) dt. \quad (424)$$

Thus the scalar formula  $\int_0^1 k(t) \text{up}(t) dt = \mathbb{E}[K(X)]$ , where  $K(x) = \int_0^x k(t) dt$ , needs only  $k \in L^1(0, 1)$ . Taking  $E = \mathbb{C}$ , regarded as a real Banach space, includes complex-valued kernels with no separate argument. If  $g, g' : \mathbb{R} \rightarrow \mathbb{R}$  are continuous and  $g$  has derivative  $g'(t)$  at every  $t \in [0, 1]$ , then

$$\int_{[0,1]} g(x) d\mu_X(x) = g(0) + \int_0^1 g'(t) \text{up}(t) dt. \quad (425)$$

More generally, let  $E$  be a real Banach space and let  $0 \leq c \leq 1$ . For a map  $k : \mathbb{R} \rightarrow E$ , with interval integrability on  $[0, c]$  in the first line and on  $[0, 1]$  in the second, the partial and full survival layer-cake identities are

$$\int_0^c \text{up}(t)k(t) \, dt = \int_{\mathbb{R}} \left( \int_0^{\min\{z, c\}} k(t) \, dt \right) d\mu_X(z), \quad (426)$$

$$\int_0^1 \text{up}(t)k(t) \, dt = \int_{\mathbb{R}} \left( \int_0^z k(t) \, dt \right) d\mu_X(z). \quad (427)$$

No positivity assumption is imposed on  $k$ ; taking  $E = \mathbb{C}$  gives the complex-kernel form used by transform arguments.

The atom-preserving closed-supergraph dual is the lower-CDF layer cake. For any finite measure  $\nu$  on  $\mathbb{R}$ ,  $a \leq b$ , and Banach-valued  $k$  interval-integrable on  $[a, b]$ ,

$$\int_a^b \nu((-\infty, t])k(t) \, dt = \int_{\mathbb{R}} \left( \int_{\min\{\max\{z, a\}, b\}}^b k(t) \, dt \right) d\nu(z). \quad (428)$$

For the partial terminal form on  $[c, b]$ , only  $c \leq b$  and the one-sided almost-everywhere bound  $z \leq b$  are needed. Compact support on  $[a, b]$  gives the convenient partial corollary and the full form; the closed rays retain any atoms of  $\nu$ .

**Theorem 7.4** (Fractional stopped-power formula for the Fabius CDF). *For every positive real  $\alpha$  and every  $x \in \mathbb{R}$ ,*

$$I_{0+}^\alpha F(x) = \frac{1}{\Gamma(\alpha + 1)} \int_{\mathbb{R}} (x - z)_+^\alpha d\mu_X(z) = \frac{1}{\Gamma(\alpha + 1)} \int_{\Omega} (x - X(\omega))_+^\alpha d\mathbf{P}(\omega). \quad (429)$$

*This includes  $x \leq 0$ , where both sides vanish. At the unit endpoint, reflection of the weighted-sum law gives*

$$I_{0+}^\alpha F(1) = \frac{1}{\Gamma(\alpha + 1)} \int_{\mathbb{R}} z^\alpha d\mu_X(z) = \frac{1}{\Gamma(\alpha + 1)} \int_{\Omega} X(\omega)^\alpha d\mathbf{P}(\omega). \quad (430)$$

The generic closed-supergraph engine is `integral_smul_setIntegral_supergraph` in `SubgraphFubini.lean`. Its finite-measure clamped, one-sided-upper-supported partial, compactly supported partial, and full forms are `intervalIntegral_cdf_smul_eq_integral_clamp`, `intervalIntegral_cdf_smul_eq_integral_max_of_ae_le`, `intervalIntegral_cdf_smul_eq_integral_max_of_ae_mem_Icc`, and `intervalIntegral_cdf_smul_eq_integral_of_ae_mem_Icc` in `CDFLayerCake.lean`. The positive-real fractional forms—respectively without a support hypothesis, with only an almost-everywhere lower support bound, directly for a probability CDF under that same bound, and at the endpoint of an almost-everywhere compact support interval—are `fractionalVolterra_measureReal_Iic_eq_integral_clamp`, `fractionalVolterra_measureReal_Iic_eq_integral_posPart`, `fractionalVolterra_cdf_eq_integral_posPart`, and `fractionalVolterra_measureReal_Iic_eq_integral_rpow_of_ae_mem_Icc` in `FractionalCDFLayerCake.lean`. Finally, `ProbabilityRepresentation.fractionalVolterra_fabiusReal_eq_integral_posPart`, `ProbabilityRepresentation.fractionalVolterra_fabiusReal_eq_uniformProduct_integral_posPart`, `ProbabilityRepresentation.fractionalVolterra_fabiusReal_one_eq_integral_rpow`, and `ProbabilityRepresentation.fractionalVolterra_fabiusReal_one_eq_uniformProduct_integral_rpow` in `FabiusFractionalIntegral.lean` are exactly (429)–(430). These declarations require neither a density nor atomlessness. They prove positive-real integral identities, not a fractional derivative, a Caputo formula, a complex-order continuation, or a folded-up, quantile, or inverse-function specialization.

For  $s \in \mathbb{R}$  and  $k \in \mathbb{N}_0$ , put

$$L_k(s) = \int_{[0,1]} e^{-sx} x^k \, d\mu_X(x). \quad (431)$$

These raw tilted moments agree with the formal analytic Fabius Laplace moments, and at zero tilt

$$L_n(0) = \int_{[0,1]} (1-x)^n \, d\mu_X(x) = d_n. \quad (432)$$

On the real axis, every derivative of the generating function is a tilted moment:

$$G^{(n)}(s) = L_n(-s), \quad G^{(n)}(0) = d_n. \quad (433)$$

Reflection of the law about  $1/2$  gives

$$L_0(s) = e^{-s} L_0(-s), \quad e^{s/2} L_0(s) = e^{-s/2} L_0(-s), \quad (434)$$

so  $s \mapsto e^{s/2} L_0(s)$  is even. Two useful quantitative forms, with all variables real, are

$$L_0\left(\frac{a+b}{2}\right)^2 \leq L_0(a)L_0(b) \quad (a, b \in \mathbb{R}), \quad (435)$$

$$L_k(s) \leq (1/a)^k k! L_0(s-a) \quad (a > 0). \quad (436)$$

*Formal proof.* The survival identity is `weightedSumDistribution_real_Ioi_eq_rvachevUp_of_nonneg`. The general Banach-valued finite-measure Fubini theorem behind (423) is `integral_Icc_intervalIntegral_eq_intervalIntegral_smul_survival`. Its finite-measure clipped form (424) is `integral_Icc_intervalIntegral_min_eq_intervalIntegral_smul_survival`. Its Fabius and report-oriented scalar forms are `integral_Icc_intervalIntegral_eq_intervalIntegral_smul_rvachevUp` and `intervalIntegral_mul_rvachevUp_eq_integral_Icc_intervalIntegral`. The endpoint-parameterized Fabius versions are `integral_Icc_intervalIntegral_min_eq_intervalIntegral_smul_rvachevUp` and `intervalIntegral_mul_rvachevUp_eq_integral_Icc_intervalIntegral_min`. The continuously differentiable formula (425) is the corollary `integral_unit_eq_zero_add_integral_deriv_mul_rvachevUp`; these seven integration declarations are in `ProbabilityLaplaceMoments.lean`. The Banach-valued partial and full formulas (426)–(427) are additionally exposed under their local interval-integrability hypotheses as `ProbabilityRepresentation.intervalIntegral_rvachevUp_smul_eq_integral_min` and `ProbabilityRepresentation.intervalIntegral_rvachevUp_smul_eq_integral`. Their generic engines are the support-free finite-measure theorem `intervalIntegral_survival_smul_eq_integral_clamp` and the two interval-supported specializations `intervalIntegral_survival_smul_eq_integral_min_of_ae_mem_Icc` and `intervalIntegral_survival_smul_eq_integral_of_ae_mem_Icc` in `SurvivalLayerCake.lean`. The tilted and zero-tilt identities are `unitLaplaceMoment_weightedSumDistribution_eq_fabiusLaplaceMoment`, `unitLaplaceMoment_weightedSumDistribution_zero_eq_halfMoment`, and `unitEndpointMoment_weightedSumDistribution_eq_halfMoment` in `ProbabilityLaplaceMoments.lean`. The derivative identities are `iteratedDeriv_generatingFunction` and `iteratedDeriv_generatingFunction_zero` in `LaplaceMoments.lean`; (434) is `fabiusLaplaceMoment_zero_reflection` and `fabiusLaplaceMoment_zero_centered_even`. Finally, (435)–(436) are `fabiusLaplaceMoment_midpoint_sq_le_all` and `fabiusLaplaceMoment_le_shift` in `UnitLaplaceMomentBounds.lean`.  $\square$

## 7.6 Exact values at reciprocal powers of two

For every  $n \in \mathbb{N}_0$ ,

$$d_n = n! 2^{\binom{n}{2}} F(2^{-n}), \quad F(2^{-n}) = \frac{d_n}{n! 2^{\binom{n}{2}}}. \quad (437)$$

This is `halfMoment_eq_fabius_formula` in `AnalyticMoments.lean`. The numerator-cleared version is `fabiusAtInverseTwoPow_eq_halfMomentNumerator_formula` in `ExactInversePower.lean`.

Writing  $V_n = F(2^{-n})$ , one has  $V_0 = 1$ , and for  $n \geq 1$ ,

$$V_n = \frac{2^{-\binom{n}{2}}}{2^n - 1} \sum_{k=0}^{n-1} \frac{2^{\binom{k}{2}}}{(n-k+1)!} V_k. \quad (438)$$

This is `fabiusAtInverseTwoPow_recurrence`. The same sequence has exact finite path and ordered-composition descriptions, proved by `fabiusAtInverseTwoPow_eq_pathSum`, `fabiusAtInverseTwoPow_eq_composition_formula`, `pathWeight_eq_boundary_product`, and `fabiusRecurrenceSequence_eq_sum_compositions` in `FabiusInverseDyadicClosedForm.lean`. Their positivity and recurrence uniqueness are `fabiusCompositionSum_pos` and `triangularRecurrence_unique`.

The totalized inverse sends both exact rational presentations straight back to their arguments. In particular, for every  $n \in \mathbb{N}_0$ , including  $n = 0$ ,

$$I(\text{fabiusAtInverseTwoPow } n) = 2^{-n}, \quad (439)$$

$$I\left(2^{-\binom{n}{2}} \text{fabiusCompositionSum } n\right) = 2^{-n}. \quad (440)$$

These are `fabiusInv_fabiusAtInverseTwoPow` in `FabiusInverse.lean` and `fabiusInv_two_pow_choose_mul_fabiusCompositionSum` in `FabiusInverseDyadicClosedForm.lean`.

## 7.7 Parity and the exact two-adic valuation

Lucas parity is packaged as `odd_binomial_coefficient_counts` and `card_odd_inner_binomial_coefficients`. It implies the exact numerator theorem `momentNumerator_odd`. For every  $n \geq 1$ , the reduced numerator and denominator of  $2d_n$  are both odd, equivalently

$$\nu_2(d_n) = -1. \quad (441)$$

This is `two_mul_halfMoment_num_den_odd` together with `halfMoment_padicVal_two`.

Consequently, for  $n \geq 1$ ,

$$\nu_2(F(2^{-n})) = -\binom{n}{2} - 1 - \nu_2(n!), \quad (442)$$

by `fabiusAtInverseTwoPow_padicVal_two`. The binary form uses the proved Legendre identity

$$\nu_2(n!) = n - \text{wt}_2(n), \quad (443)$$

namely `factorial_padicVal_two`; the combined declaration is `fabiusAtInverseTwoPow_padicVal_two_closed` in `TwoAdic.lean`.

The central-binomial specialization gives, for every  $n \geq 0$ ,

$$\nu_2\binom{2n}{n} = \text{wt}_2(n), \quad (444)$$

$$\varepsilon_n = (-1)^{\nu_2\binom{2n}{n}}. \quad (445)$$

These are `padicValNat_two_centralBinom` and `thueMorseSign_eq_neg_one_pow_centralBinom`. The companion theorem `padicValNat_two_centralBinom_eq_zero_iff` says that the valuation vanishes exactly when the binary weight is zero, hence only at  $n = 0$ ; a positive power of two has binary weight, and therefore valuation, one. These three declarations exhaust the public surface of `CentralBinomialValuation.lean` and give the direct formal crosswalk for the central-binomial theorem in the Thue–Morse atlas.

The prime-power row identity is available directly at the valuation level. For every prime  $p$ , exponent  $m$ , and  $0 < j \leq p^m$ ,

$$\nu_p\binom{p^m}{j} + \nu_p(j) = m, \quad \nu_p\binom{p^m}{j} = m - \nu_p(j). \quad (446)$$

The identities in (446) are `primePowerChoose_padicValNat_add` and `primePowerChoose_padicValNat`. They include the right endpoint  $j = p^m$  and the boundary  $m = 0$ . The strict dyadic-comb specialization  $0 < j < 2^m$  is `twoPowChoose_padicValNat`. The two preceding Pascal rows satisfy the companion identities

$$\nu_p\binom{p^m-1}{j} = 0 \quad (0 \leq j < p^m), \quad \nu_p\binom{p^m-2}{j-1} = \nu_p(j) \quad (0 < j < p^m). \quad (447)$$

These are `primePowerSubOneChoose_padicValNat` and `primePowerSubTwoChoose_padicValNat`; the latter’s dyadic wrapper is `twoPowSubTwoChoose_padicValNat`. Its strict upper bound is essential: at  $j = p^m$  the selected column lies beyond row  $p^m - 2$ , so the binomial coefficient is zero. These six declarations, and no definitions, exhaust `PrimePowerBinomialValuation.lean`. No valuation-histogram theorem is claimed.

## 7.8 Bernoulli recurrences

The formal development also proves Bernoulli convolution and recurrence forms for both sequences. For  $n \geq 1$ ,

$$c_n = \frac{1}{2^{2n}-1} \sum_{k=1}^n 2^{2n-2k} (2^{2k}-2) \binom{2n}{2k} B_{2k} c_{n-k}, \quad (448)$$

$$d_n = \frac{n 2^n}{4(2^n-1)} d_{n-1} - \frac{1}{2^n-1} \sum_{k=1}^{\lfloor n/2 \rfloor} \binom{n}{2k} 2^{n-2k} B_{2k} d_{n-2k}. \quad (449)$$

Here  $B_1 = -1/2$ . These are `moment_bernoulli_recurrence` and `halfMoment_bernoulli_recurrence`; their convolution forms and combined source-paper statement are `moment_bernoulli_convolution`, `halfMoment_bernoulli_convolution`, and `proposition_twenty_two_formula` in `BernoulliRecurrences.lean`.

## 8 Exact evaluation at arbitrary dyadic rationals

The rationality theorem for dyadic arguments comes with two complementary exact presentations:

1. a closed finite formula involving Thue–Morse signs and the rational moments  $c_k$ ;
2. an executable bit-recursive Taylor evaluator.

### 8.1 The closed Thue–Morse formula

**Theorem 8.1** (Exact dyadic formula). *Let  $n, a \in \mathbb{N}_0$  with  $0 \leq a \leq 2^n$ . Then*

$$F\left(\frac{a}{2^n}\right) = \frac{2^{-\binom{n+1}{2}}}{n!} \sum_{h=0}^{a-1} (-1)^{\text{wt}_2(h)} \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} c_k (2a - 2h - 1)^{n-2k}. \quad (450)$$



An empty outer sum has value 0.

*Formal proof.* The rational expression on the right is the definition `fabiusDyadic`. Its equality, after coercion to  $\mathbb{R}$ , with  $F(a/2^n)$  under the displayed bound on  $a$  is `fabiusDyadic_cast` in `DyadicAnalytic.lean`.  $\square$

Every quantity on the right of (450) is rational, so dyadic rationality is immediate. More importantly, the formula lives entirely in  $\mathbb{Q}$  and can be evaluated without roundoff.

**Theorem 8.2** (Global dyadic formula). *For all  $n, a \in \mathbb{N}_0$  the signed global extension satisfies*

$$F_{\text{gl}}\left(\frac{a}{2^n}\right) = \frac{2^{-\binom{n+1}{2}}}{n!} \sum_{h=0}^{a-1} \varepsilon_h \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} c_k (2a - 2h - 1)^{n-2k}. \quad (451)$$

No upper bound on  $a$  is imposed. An empty outer sum has value 0.

*Formal proof.* The displayed rational identity is `fabiusDyadic_cast_extended_nat` in `GlobalDyadic.lean`, after unfolding the closed form `fabiusDyadic`.  $\square$

This closed form depends only on the represented dyadic rational, not on the chosen numerator–exponent pair. Writing  $\text{Dyad}(n, m) = \text{fabiusDyadic } n \, m$ , Lean proves

$$\text{Dyad}(n + k, 2^k m) = \text{Dyad}(n, m), \quad (452)$$

$$\frac{m_1}{2^{n_1}} = \frac{m_2}{2^{n_2}} \text{ in } \mathbb{Q} \implies \text{Dyad}(n_1, m_1) = \text{Dyad}(n_2, m_2). \quad (453)$$

The exact declarations are `fabiusDyadic_refine_pow` and `fabiusDyadic_eq_of_rat_eq` in `GlobalDyadic.lean`.

Two special cases are the ones used in practice.

**Corollary 8.3** (Bounded case). *If  $0 \leq a \leq 2^n$  then  $F_{\text{gl}}(a/2^n) = F(a/2^n)$ , so (451) reduces to (450).*

*Proof.* The two functions agree on  $[0, 1]$  and  $a/2^n \in [0, 1]$ .  $\square$

**Corollary 8.4** (Bump case). *Let  $n \in \mathbb{N}_0$  and  $q \in \mathbb{Z}$ . If  $|q| < 2^n$ , then*

$$\text{up}\left(\frac{q}{2^n}\right) = \frac{2^{-\binom{n+1}{2}}}{n!} \sum_{h=0}^{q+2^n-1} \varepsilon_h \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} c_k (2q + 2^{n+1} - 2h - 1)^{n-2k}, \quad (454)$$

*that is, (451) evaluated at  $a = q + 2^n$ . If  $|q| \geq 2^n$ , then  $\text{up}(q/2^n) = 0$ .*

*Proof.* Taking  $b = 0$  in (12) gives  $F_{\text{gl}}(x) = \text{up}(x-1)$  for  $0 \leq x \leq 2$ . With  $x = 1 + q/2^n = (q+2^n)/2^n$  and  $|q| < 2^n$  we have  $0 < x < 2$ , so  $\text{up}(q/2^n) = F_{\text{gl}}((q+2^n)/2^n)$ , and (451) with  $a = q + 2^n$  applies. The last statement is the support condition  $\text{supp up} = [-1, 1]$ .  $\square$

Corollary 8.4 is the global form invoked in the proof of the uniform denominator multiplier theorem below. The point is that it genuinely requires Theorem 8.2 and not (450): for  $0 < q < 2^n$  the numerator  $a = q + 2^n$  exceeds  $2^n$ , so the bounded statement does not apply. At the endpoints  $|q| = 2^n$  the two clauses agree, both giving 0; for  $q = -2^n$  the outer sum is empty and for  $q = 2^n$  the value is  $F_{\text{gl}}(2) = 0$ .

*Remark 8.5* (Why the sum is finite). Both index sets in (451) are finite: the outer range is  $0 \leq h < a$ , and the inner range is  $0 \leq k \leq \lfloor n/2 \rfloor$ . The moments  $c_k$  are rational by the proved recurrence (387). Thus the formula is an exact computation in  $\mathbb{Q}$ , with no analytic truncation or floating-point roundoff.

*Remark 8.6* (Exact evaluation without uncertified examples). The finite sum is an exact rational algorithm, and its agreement with the analytic Fabius function is `fabiusDyadic_cast`. Concrete numerical specializations that do not yet have their own named Lean theorems are recorded only in the consolidated research-frontier volume.



## 8.2 Thue–Morse cancellation

The mechanism that makes the dyadic formulas simplify is the exact Prouhet cancellation carried by the Thue–Morse signs.

**Lemma 8.7** (Affine power cancellation). *If  $d < b$ , then for all  $X, Y \in \mathbb{Q}$ ,*

$$\sum_{h=0}^{2^b-1} (-1)^{\text{wt}_2(h)} (X + Yh)^d = 0. \quad (455)$$

*Formal proof.* This is the strict-degree case of `thueMorse_affine_power_sum_of_le` in `ThueMorsePrefix.lean`; the monomial statement used internally is `thueMorsePowerSum_eq_zero_of_lt`.  $\square$

The same cancellation has an order-free centered formulation. For half-steps  $a_i$  indexed by a finite set  $s$ , define

$$\Delta_{a,s}^{\text{sym}} f(x) := \Delta_{(-2a_i)_{i \in s}} f\left(x + \sum_{i \in s} a_i\right), \quad (456)$$

where the right-hand side is the ordinary mixed-difference operator of `ThueMorseMixedDifference.lean`. This is `symmetricMixedDifference`. It is evaluation at  $x$  for  $s = \emptyset$ , and for a singleton it is  $f(x + a_i) - f(x - a_i)$ . Peeling off  $i \notin s$  gives

$$\Delta_{a,s \cup \{i\}}^{\text{sym}} f(x) = \Delta_{a,s}^{\text{sym}} f(x + a_i) - \Delta_{a,s}^{\text{sym}} f(x - a_i). \quad (457)$$

The exact declarations are `symmetricMixedDifference_empty`, `symmetricMixedDifference_singleton`, and `symmetricMixedDifference_insert`.

Its Boolean-cube expansion is

$$\Delta_{a,s}^{\text{sym}} f(x) = \sum_{t \subseteq s} (-1)^{|t|} f\left(x + \sum_{i \in s} a_i - 2 \sum_{i \in t} a_i\right). \quad (458)$$

This is `symmetricMixedDifference_eq_sum_powerset_smul`. If  $p$  is a polynomial over a commutative ring and  $\deg p \leq |s|$ , then

$$\Delta_{a,s}^{\text{sym}} p(x) = [X^{|s|}] p \, |s|! \prod_{i \in s} (2a_i). \quad (459)$$

Thus every polynomial of degree strictly below  $|s|$  is annihilated, including the zero-polynomial boundary when  $s = \emptyset$ , and the first surviving monomial  $X^{|s|}$  gives exactly the sign-free factor on the right. These three statements are `symmetricMixedDifference_polynomial_eq_coeff_card`, `symmetricMixedDifference_polynomial_of_degree_lt`, and `symmetricMixedDifference_pow_card`.

For the dyadic half-steps  $2^j h$ ,  $0 \leq j < m$ , the definition `symmetricDyadicMixedDifference` has the zero-order value `symmetricDyadicMixedDifference_zero` and the centered block expansion

$$\Delta_{m,h}^{\text{sym}} f(x) = \sum_{n < 2^m} \varepsilon_n f\left(x + \sum_{j < m} 2^j h - 2nh\right). \quad (460)$$

This is `symmetricDyadicMixedDifference_eq_sum_thueMorseSign_smul`. Over a characteristic-zero field, taking  $h = 2^{-m}$  and reflecting the block gives the increasing-grid form

$$\Delta_{m,2^{-m}}^{\text{sym}} f(x) = \sum_{n < 2^m} (-1)^m \varepsilon_n f(x - (1 - 2^{-m}) + 2n2^{-m}), \quad (461)$$

proved as `symmetricDyadicMixedDifference_inv_two_pow_eq_sum_thueMorseSign_smul`. The positive-order spelling with  $m + 1$  differences and mesh  $2^{-m}$  is `symmetricDyadicMixedDifference_inv_two_pow_succ_eq_sum_thueMorseSign_smul`. These two definitions and eleven theorems are the exhaustive public API of `ThueMorseSymmetricDifference.lean`; no limiting or differentiability claim is made there.

The analytic corner formula is supplied by the companion `ThueMorseCornerIntegral.lean`. For a sequence of half-widths  $a_i$ , define the centered box integral recursively by

$$\mathcal{B}_{a,0}(h; x) = h(x), \quad \mathcal{B}_{a,N+1}(h; x) = \int_{-a_N}^{a_N} \mathcal{B}_{a,N}(h; x + u) \, du.$$

This is `centeredBoxIntegral`; its two recursion equations are `centeredBoxIntegral_zero` and `centeredBoxIntegral_succ`. Suppose  $a_i \geq 0$  for  $i < N$ ,  $I \subseteq \mathbb{R}$  is open and order-connected,  $g$  is  $C^N$  on  $I$ , and the entire segment

$$\left[ x - \sum_{i < N} a_i, x + \sum_{i < N} a_i \right] \text{ is contained in } I.$$

Then

$$\Delta_{a,\{0,\dots,N-1\}}^{\text{sym}} g(x) = \mathcal{B}_{a,N}(g^{(N)}; x). \quad (462)$$

The range-indexed statement is `symmetricMixedDifference_range_eq_centeredBoxIntegral`; `symmetricMixedDifference_univ_eq_centeredBoxIntegral` is the identical formula reindexed by `Fin N`. These, the definition, and the two recursion equations are the exhaustive one-definition/four-theorem public surface of the module. The assumptions are genuinely local: no global  $C^N$  hypothesis is substituted. They permit zero half-steps and  $N = 0$ , hence strengthen the manuscript’s positive-step case, but do not assert a signed-half-step formula.

For the proof of (462), peel the last symmetric difference by (457). At each Boolean-cube corner, order-connectedness and containment of the full segment place the translated interval inside  $I$ , so the local fundamental theorem of calculus replaces the endpoint difference by an integral of  $g'$ . Finite linearity moves the powerset sum through that integral. The induction hypothesis applied to  $g'$ , whose local regularity is  $C^{N-1}$ , gives the remaining nested integral. The zero-order case is evaluation by definition. Together with the finite algebra above, this proves every clause of the continuous-chaos report’s `thm:TM-corner`, now *Exact*. It does not provide the separate Walsh conditional-expectation construction or its normalization.

The finite block polynomial packages the same signs without introducing an analytic exponential identity:

$$\sum_{h < 2^b} \varepsilon_h z^h = \prod_{j=0}^{b-1} (1 - z^{2^j}). \quad (463)$$

This is `thueMorseBlockPolynomial_eq_product` in `ThueMorseGenerating.lean`. At the level of the centered formal exponential series, the corresponding zero has exact order  $b$ , by `order_thueMorseCenteredPowerSeries` and `thueMorseCenteredPowerSeries_eq_product` in `ThueMorseExponential.lean`.

Writing  $P_b(z)$  for either side of (463), its zero at  $z = 1$  has exact multiplicity  $b$  over  $\mathbb{Z}[z]$ :

$$(1 - z)^b \mid P_b(z), \quad (1 - z)^{b+1} \nmid P_b(z). \quad (464)$$

The divisibility and matching nondivisibility statements are respectively `one_sub_X_pow_dvd_thueMorseBlockPolynomial` and `one_sub_X_pow_succ_not_dvd_thueMorseBlockPolynomial` in `ThueMorseBlockAlgebra.lean`.

The ordinary derivatives of the finite block polynomial make that exact order visible without passing to formal power series.

**Proposition 8.8** (Block-polynomial derivatives at one). *For every  $b, r \geq 0$ ,*

$$\left. \frac{d^r}{dz^r} \sum_{h < 2^b} \varepsilon_h z^h \right|_{z=1} = r! \sum_{h < 2^b} \varepsilon_h \binom{h}{r}. \quad (465)$$

*Consequently this derivative is zero when  $r < b$ , while the first surviving derivative is*

$$\left. \frac{d^b}{dz^b} \prod_{j=0}^{b-1} (1 - z^{2^j}) \right|_{z=1} = (-1)^b b! 2^{\binom{b}{2}}. \quad (466)$$

*Formal proof.* The derivative–binomial dictionary is `iterate_derivative_thueMorseBlockPolynomial_eval_one`; its strict vanishing and first surviving cases are respectively `iterate_derivative_thueMorseBlockPolynomial_eval_one_of_lt` and `iterate_derivative_thueMorseBlockPolynomial_eval_one_self`, all in `ThueMorseBlockAlgebra.lean`.  $\square$

**Proposition 8.9** (The first surviving Thue–Morse moment). *For every  $k \geq 0$ ,*

$$\boxed{\sum_{r=0}^{2^k-1} \varepsilon_r r^k = (-1)^k 2^{\binom{k}{2}} k!}. \quad (467)$$

*Formal proof.* This is `thueMorsePowerSum_self` in `ThueMorsePrefix.lean`. Together with `thueMorsePowerSum_eq_zero_of_lt`, it is also the coefficient calculation behind the exact-order theorem above.  $\square$

**Corollary 8.10** (Translation invariance). *For every  $k \geq 0$  and  $Q \in \mathbb{Q}$ ,*

$$\sum_{r=0}^{2^k-1} \varepsilon_r (r + Q)^k = (-1)^k 2^{\binom{k}{2}} k!. \quad (468)$$

*Formal proof.* Apply `thueMorse_affine_power_sum_self` with  $X = Q$  and  $Y = 1$ .  $\square$

**Proposition 8.11** (Centered reflection and the characteristic-two boundary). *Let  $R$  be a commutative ring, let  $c \in R$  satisfy  $c + c = 2^m - 1$ , and put*

$$S_{m,r}(c) = \sum_{n=0}^{2^m-1} \varepsilon_n (n - c)^r \in R. \quad (469)$$

*Then*

$$S_{m,r}(c) = (-1)^{m+r} S_{m,r}(c). \quad (470)$$

*Consequently, if  $m + r$  is odd, then  $2S_{m,r}(c) = 0$ . This is the characteristic-free conclusion: in a ring with 2-torsion one cannot in general cancel the factor 2. Over  $\mathbb{Q}$ , taking  $c = (2^m - 1)/2$  gives the familiar parity rule*

$$\sum_{n=0}^{2^m-1} \varepsilon_n \left( n - \frac{2^m - 1}{2} \right)^r = 0 \quad (m + r \text{ odd}). \quad (471)$$

*Formal proof.* The fixed-point identity is `sum_thueMorseSign_mul_centered_pow_reflect`, and its universal 2-torsion consequence is `two_mul_sum_thueMorseSign_mul_centered_pow_eq_zero`. The rational specialization, including the legitimate cancellation of  $2 \neq 0$ , is `sum_thueMorseSign_mul_midpoint_pow_eq_zero`. All three declarations are in `ThueMorseMoments.lean`.  $\square$

The centered specialization  $Q = -2^k$  is therefore

$$\sum_{r=0}^{2^k-1} \varepsilon_r (r - 2^k)^k = (-1)^k 2^{\binom{k}{2}} k!, \quad (472)$$

the exact declaration `thueMorseCenteredPowerSum_self`.

### 8.3 The exact leading-bit recurrence

Let  $e, a \in \mathbb{N}_0$  with  $0 < a \leq 2^e$ , and write

$$b = \lfloor \log_2 a \rfloor, \quad r = e - b, \quad q = a - 2^b, \quad y = \frac{q}{2^e}. \quad (473)$$

Thus  $0 \leq q < a$ : the leading binary power has been removed. Put  $V_m = F(2^{-m})$ , whose exact half-moment form is (437), and define

$$T_r(y) = \sum_{j=0}^r \frac{2^{\binom{j+1}{2}} V_{r-j}}{j!} y^j. \quad (474)$$

Then the exact rational recurrence is

$$\boxed{F\left(\frac{a}{2^e}\right) = T_r\left(\frac{q}{2^e}\right) - F\left(\frac{q}{2^e}\right)}. \quad (475)$$

The formal route stays entirely inside the rational dyadic model: `fabijsTaylorHorner_eq_sum` identifies the Horner routine with (474); `fabijsInversePowTwoTable_hasBlockTaylor` proves the block identity; `blockTaylor_implies_bitRecurrence` turns it into the leading-bit recurrence; and `fabijsInversePowTwoTable_hasBitRecurrence` instantiates that recurrence for the precomputed inverse-power table. Finally, `fabijsDyadicUnit_eq_fabijsDyadic` and `fabijsDyadic_cast` identify the executable rational result with the analytic function. These declarations are in `DyadicClosedForm.lean` and `DyadicAnalytic.lean`.

### 8.4 The bit-recursive evaluator

Use  $b, r, q, y$ , and  $T_r$  from (473)–(474). Equation (475) reduces the numerator from  $a$  to the strictly smaller  $q$ .

**Algorithm 8.12** (Exact dyadic evaluator). Precompute  $V_j = F(2^{-j})$  for  $0 \leq j \leq e$  from (437). Define  $\text{Eval}(e, a)$  recursively by

$$\text{Eval}(e, a) = \begin{cases} 0, & a \leq 0, \\ 1, & a \geq 2^e, \\ T_r(q/2^e) - \text{Eval}(e, q), & 0 < a < 2^e, \end{cases}$$

where  $b, r, q$  are given by (473); evaluate  $T_r$  by Horner’s rule.

A literal pseudocode version is:

```
Eval(e, a, V):
  if a <= 0:      return 0
  if a >= 2^e:    return 1

  b = floor_log2(a)
  r = e - b
```

```

q = a - 2^b
y = q / 2^e

p = V[0] // highest Taylor coefficient, j = r
for m = 1 .. r:
    j = r - m
    p = V[m] + (2^(j+1) * y / (j+1)) * p

return p - Eval(e, q, V)

```

The displayed loop is one convenient Horner ordering; any exact evaluation of (474) is equivalent. In the formal source this evaluator is `fabiusDyadicValue`.

**Theorem 8.13** (Correctness and termination). *For every  $e \in \mathbb{N}_0$  and  $a \in \mathbb{Z}$ ,*

$$\text{Eval}(e, a) = F(a/2^e) \quad (476)$$

*under the bounded convention  $F = 0$  to the left and  $F = 1$  to the right.*

*Formal proof.* The pseudocode is the rational routine `fabiusDyadicValue`, whose analytic correctness is `fabiusDyadicValue_cast` in `PaperStatements.lean`. The interior recursion is `fabiusInversePowTwoTable_hasBitRecurrence`; the corresponding total recursive correctness theorem is `fabiusDyadicUnitAux_eq_of_bitRecurrence`.  $\square$

## 8.5 Reduction of an arbitrary dyadic argument

Algorithm 8.12 evaluates the bounded function  $F$  on  $[0, 1]$ . A total evaluator for the signed global extension first folds the argument back into the unit interval and records the associated sign. The fold is a reduction modulo 2 followed by a reflection.

**Theorem 8.14** (Triangular reduction). *Let  $n \in \mathbb{N}_0$  and put  $s = 2^n$ . Let  $a \in \mathbb{N}_0$  and write*

$$a = 2sB + R, \quad B = \left\lfloor \frac{a}{2s} \right\rfloor, \quad 0 \leq R < 2s, \quad (477)$$

*and set*

$$C = \begin{cases} R, & R \leq s, \\ 2s - R, & R > s. \end{cases} \quad (478)$$

*Then  $0 \leq C \leq s$  and*

$$\boxed{F_{\text{gl}}\left(\frac{a}{2^n}\right) = \varepsilon_B F\left(\frac{C}{2^n}\right)}. \quad (479)$$

*For  $a \in \mathbb{Z}$  with  $a < 0$  one has  $F_{\text{gl}}(a/2^n) = 0$ .*

*Proof.* Write  $x = a/2^n = 2B + R/s$ . Since  $0 \leq R < 2s$  we have  $2B \leq x < 2B + 2$ , so  $x$  lies in the  $B$ th block and (12) gives

$$F_{\text{gl}}(x) = \varepsilon_B \text{up}(x - 2B - 1) = \varepsilon_B \text{up}\left(\frac{R}{s} - 1\right).$$

Put  $t = R/s - 1$ , so that  $-1 \leq t < 1$ . The fold relation (7) says  $\text{up}(t) = F(t + 1)$  for  $t \leq 0$  and  $\text{up}(t) = F(1 - t)$  for  $t > 0$ ; on  $[-1, 1]$  both clauses read  $\text{up}(t) = F(1 - |t|)$ . If  $R \leq s$  then  $|t| = 1 - R/s$  and  $F(1 - |t|) = F(R/s)$ . If  $R > s$  then  $|t| = R/s - 1$  and  $F(1 - |t|) = F(2 - R/s) = F((2s - R)/2^n)$ . In both cases the argument is  $C/2^n$  with  $C$  as in (478), and  $0 \leq C \leq s$  is immediate from  $0 \leq R < 2s$ . The final statement is the vanishing of  $F_{\text{gl}}$  on the negative half-line.  $\square$

Combining (479) with Algorithm 8.12 gives a total exact evaluator for  $F_{\text{gl}}$  at every dyadic rational: reduce with (477)–(478), compute the block sign  $\varepsilon_B = (-1)^{\text{wt}_2(B)}$  from the binary digits of  $B = \lfloor a/2^{n+1} \rfloor$ , and call  $\text{Eval}(n, C)$ .

The reduction is compatible with unreduced presentations. Replacing  $(n, a)$  by  $(n+1, 2a)$  doubles  $s$  and  $R$ , leaves  $B$  unchanged, and doubles  $C$ ; the right side of (479) is therefore unchanged, as it must be.

*Remark 8.15* (Recognizing dyadic arguments exactly). A reduced rational  $p/q$  with  $q > 0$  is dyadic exactly when  $q$  is a power of two, and in that case  $n = \log_2 q$  and  $a = p$  are read off directly. An exact rational interface to  $F$ ,  $\text{up}$ , and  $F_{\text{gl}}$  therefore tests a single integer condition on the reduced denominator, returns “not dyadic” when it fails, and otherwise invokes the signed integer evaluator above. No floating-point recognition of powers of two is involved anywhere, so the interface is exact on every input and cannot be defeated by rounding in the recognition step.

## 8.6 Denominator bounds

Let

$$M_m = \prod_{j=1}^m (2^{2^j} - 1), \quad (2m+1)!! = 1 \cdot 3 \cdots (2m+1). \quad (480)$$

**Theorem 8.16** (Uniform denominator multiplier). *Let  $n \geq 1$  and  $a \in \mathbb{Z}$  with  $|a| < 2^n$ . Then*

$$\text{up}(a/2^n) \, n! \, 2^{\binom{n+1}{2}} (2 \lfloor n/2 \rfloor + 1)!! \, M_{\lfloor n/2 \rfloor} \quad (481)$$

*is an integer.*

*Formal proof.* This is `rvachevDyadic_mul_denominatorBound_isNatural_all` in `DenominatorBound.lean`. Its proof uses `moment_mul_common_denominator` for the moments and `fabiusDyadic_mul_denominatorBound_isInteger` for the bounded dyadic values.  $\square$

Let  $\Delta_n$  be the least common multiple of the reduced denominators of the values  $\text{up}(a/2^n)$  with odd  $a$  in the support. The theorem gives the explicit divisibility bound

$$\Delta_n \mid n! \, 2^{\binom{n+1}{2}} (2 \lfloor n/2 \rfloor + 1)!! \, M_{\lfloor n/2 \rfloor}. \quad (482)$$

A complementary lower statement is also formalized.

**Proposition 8.17** (The reciprocal power is one of the samples). *For every  $n \geq 1$  the reduced denominator of  $F(2^{-n})$  divides  $\Delta_n$ . Consequently*

$$\nu_2(\Delta_n) \geq \binom{n}{2} + 1 + \nu_2(n!). \quad (483)$$

*Formal proof.* The denominator divisibility is `proposition_fifteen` in `PaperStatements.lean`, after the bridge `dyadicDenominator_eq_fabiusDyadicDenominator`. The exact two-adic exponent is `fabiusAtInverseTwoPow_padicVal_two_closed`.  $\square$

Together, (482) and (483) give the formally proved denominator information used in this article. Proposed exact factorizations of the least common denominator, together with the associated numerical tables, are kept in the consolidated research-frontier volume until their individual statements are formalized.

### 8.7 The exact half-base $q$ -binomial layer

The dyadic dilation forces the specialization  $q = 1/2$ . For  $a, q \in \mathbb{Q}$ , define

$$(a; q)_n = \prod_{j=0}^{n-1} (1 - aq^j), \quad (484)$$

and write  $\begin{bmatrix} n \\ k \end{bmatrix}_{1/2}$  for the Gaussian coefficient at that fixed base. For  $k \leq n$ , its exact Mersenne normalization is

$$\begin{bmatrix} n \\ k \end{bmatrix}_{1/2} = \frac{\mathfrak{m}_n}{\mathfrak{m}_k \mathfrak{m}_{n-k} 2^{k(n-k)}}, \quad \mathfrak{m}_j = \prod_{\ell=1}^j (2^\ell - 1). \quad (485)$$

In particular the row is symmetric under  $k \mapsto n - k$ .

**Theorem 8.18** (Finite half-base  $q$ -binomial theorem). *For every  $n \in \mathbb{N}_0$  and  $z \in \mathbb{Q}$ ,*

$$(z; 1/2)_n = \sum_{k=0}^n (-1)^k 2^{-\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_{1/2} z^k. \quad (486)$$

At every dyadic node this gives the single closed evaluation

$$\sum_{k=0}^n (-1)^k 2^{-\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_{1/2} (2^m)^k = \begin{cases} 0, & m < n, \\ (-1)^n \mathfrak{m}_m / \mathfrak{m}_{m-n}, & n \leq m. \end{cases} \quad (487)$$

In particular, the sum vanishes if and only if  $m < n$ . Moreover,

$$(1/2; 1/2)_n = 2^{-\binom{n+1}{2}} \mathfrak{m}_n. \quad (488)$$

*Formal proof.* These are `halfQBinomial_theorem`, `qBinomial_half_two_pow_sum_eq_ite`, `qBinomial_half_two_pow_sum_eq_zero_iff`, `halfQPochhammer_eq_mersenne_div`, `halfQBinomial_eq_mersenne`, and `halfQBinomial_symm` in `HalfQBinomial.lean`.  $\square$

**Remark 8.19** (The denominator-free finite  $q$ -binomial theorem). The half-base identity is a specialization of a ring-valued statement that does not divide by a  $q$ -Pochhammer symbol. For a commutative ring  $R$  and  $q \in R$ , define the polynomial Gaussian coefficients recursively by

$$(z; q)_0 = 1, \quad (z; q)_{n+1} = (z; q)_n (1 - zq^n),$$

and

$$\begin{bmatrix} n+1 \\ k+1 \end{bmatrix}_q = \begin{bmatrix} n \\ k+1 \end{bmatrix}_q + q^{n-k} \begin{bmatrix} n \\ k \end{bmatrix}_q, \quad \begin{bmatrix} n \\ 0 \end{bmatrix}_q = \begin{bmatrix} n \\ n \end{bmatrix}_q = 1, \quad \begin{bmatrix} n \\ k \end{bmatrix}_q = 0 \quad (n < k).$$

The recursive triangle is already defined over every semiring and is functorial: for semirings  $R, S$  and every homomorphism  $\varphi : R \rightarrow S$ ,

$$\varphi \left( \begin{bmatrix} n \\ k \end{bmatrix}_q \right) = \begin{bmatrix} n \\ k \end{bmatrix}_{\varphi(q)}. \quad (489)$$

Over a commutative semiring it also satisfies, for all natural  $n, k$ , the second Pascal orientation

$$\begin{bmatrix} n+1 \\ k+1 \end{bmatrix}_q = q^{k+1} \begin{bmatrix} n \\ k+1 \end{bmatrix}_q + \begin{bmatrix} n \\ k \end{bmatrix}_q, \quad (490)$$

and, whenever  $k \leq n$ , the reflection law

$$\begin{bmatrix} n \\ n-k \end{bmatrix}_q = \begin{bmatrix} n \\ k \end{bmatrix}_q. \quad (491)$$

None of these structural laws uses division or cancellation; the zero extension above the diagonal is what makes (490) valid without an index side condition.

Then for every  $z \in R$ ,

$$(z; q)_n = \sum_{k=0}^n (-1)^k q^{\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q z^k. \quad (492)$$

The recursive coefficient already makes sense over an arbitrary semiring, while the displayed subtraction requires a commutative ring. No denominator, nonzero assumption, or exclusion of  $q = 0$  and roots of unity occurs. The recurrence, triangular boundaries, and (492) are the definition `finiteQPochhammerIn`, its boundary and recurrence `finiteQPochhammerIn_zero` and `finiteQPochhammerIn_succ`, the coefficient definition `gaussianBinomial`, and the theorems `gaussianBinomial_zero_zero`, `gaussianBinomial_zero_succ`, `gaussianBinomial_succ_succ`, `gaussianBinomial_zero_right`, `gaussianBinomial_eq_zero_of_lt`, `gaussianBinomial_self`, and `finite_qBinomial_theorem` in `FiniteQBinomialCore.lean`. The functoriality, alternate Pascal, and symmetry statements (489)–(491) are `map_gaussianBinomial`, `gaussianBinomial_succ_succ_alt`, and `gaussianBinomial_symm`, respectively, in the same module.

The companion `QBinomialReciprocity.lean` has no public definitions and exactly four public theorems. For a unit  $q$  in any commutative semiring,

$$q^{k(n-k)} \begin{bmatrix} n \\ k \end{bmatrix}_{q^{-1}} = \begin{bmatrix} n \\ k \end{bmatrix}_q \quad (493)$$

is `gaussianBinomial_reciprocity_units`; the statement is total in  $n, k$ , since the recursive Gaussian coefficients on both sides vanish when  $n < k$ . The theorem `gaussianBinomial_reciprocity` is exactly the nonzero- $q$  wrapper over an arbitrary semifield. Over every commutative ring, `gaussianBinomial_neg_one_eq_zero_of_odd_degree` proves  $\begin{bmatrix} n \\ k \end{bmatrix}_{-1} = 0$  when  $k(n-k)$  is odd, and `gaussianBinomial_neg_one_even_odd_eq_zero` proves the total specialization  $\begin{bmatrix} 2a \\ 2b+1 \end{bmatrix}_{-1} = 0$ , including the above-diagonal cases. These are finite algebraic identities: they use no quotient formula, cancellation, integral-domain, characteristic, topology, or convergence hypothesis.

The downstream `GaussianBinomialPalindromic.lean` has no public definitions and exactly fourteen public theorems: `reflect_add_of_natDegree_le`, `reflect_one'`, `gaussianBinomial_natDegree_le`, `gaussianBinomial_zero_left`, `gaussianBinomial_diag'`, `reflect_gaussianBinomial`, `coeff_gaussianBinomial_reflect`, `coeff_gaussianBinomial_zero`, `coeff_gaussianBinomial_top`, `gaussianBinomial_natDegree`, `gaussianBinomial_monoid`, `two_mul_derivative_gaussianBinomial_eval_one`, `coeff_gaussianBinomial_one_of_pos_of_lt`, and `coeff_gaussianBinomial_one`. Over every commutative semiring the module supplies the two generic reflection helpers, the Gaussian degree bound, zero and diagonal values, reflection in degree  $k(n-k)$ , constant and top coefficients one, bounded-index coefficient reversal, and the division-free derivative-at-one mean identity

$$2 \begin{bmatrix} n \\ k \end{bmatrix}'_X \Big|_{X=1} = k(n-k) \binom{n}{k}.$$

The interior linear coefficient theorem assumes exactly  $0 < k < n$  and gives coefficient one. The total theorem classifies it as

$$[X^1] \begin{bmatrix} n \\ k \end{bmatrix}_X = \begin{cases} 1, & 0 < k < n, \\ 0, & \text{otherwise,} \end{cases}$$



thereby covering  $k = 0$ ,  $k = n$ , and  $n < k$ . Both results hold over every commutative semiring, without nontriviality; exact degree and monicity alone require the semiring to be nontrivial.

The companion `GaussianBinomialPolynomialStructure.lean` has no public definitions and exactly five public theorems: `natDegree_gaussianBinomial_universal`, `gaussianBinomial_universal_mononic`, `coeff_zero_gaussianBinomial_universal`, `gaussianBinomial_universal_reflect`, and `coeff_gaussianBinomial_universal_symm`. For  $k \leq n$ , these state that the universal natural-coefficient polynomial  $\begin{bmatrix} n \\ k \end{bmatrix}_X \in \mathbb{N}_0[X]$  has exact natural degree  $k(n - k)$ , is monic, has constant coefficient one, and is fixed by reflection in that degree. Under the additional bound  $d \leq k(n - k)$ , the last theorem identifies the coefficients in degrees  $d$  and  $k(n - k) - d$ . Together with the row symmetry and reciprocal identities above, this supplies the universal natural-coefficient-polynomial form of the same structure theorem. The adjacent fourteen-theorem module supplies the complete linear-coefficient classifier over every commutative semiring.

Two further source-only modules have exhaustive finite-algebra inventories. `CentralQBino`  
`mialReduction.lean` has no definitions and six theorems: `finiteQPochhammerIn_mul_neg`, `finiteQPochhammerIn_two_mul`, `finiteQPochhammerIn_map_ringHom`, `central_gaussianBinomial_sq_mul_int`, `central_gaussianBinomial_sq_mul`, and `central_gaussianBinomial_sq_div`. They give sign pairing, even-odd dissection, naturality, the integral certificate and division-free central squared-base reduction over commutative rings, and its field quotient under the two displayed nonvanishing hypotheses. `CyclotomicFactorization.lean` has no definitions and seven theorems: `div_add_div_le_div`, `div_le_div_add_div_add_one`, `mem_range_and_mem_divisors_iff`, `finiteQPochhammerIn_X_eq_prod_cyclotomic`, `finiteQPochhammerIn_X_eq_gaussianBinomial_mul`, `prod_cyclotomic_pow_div_extend`, and `gaussianBinomial_X_eq_prod_cyclotomic`. These bound the cyclotomic exponents by zero and one, reorganize divisor indices, factor the finite  $q$ -product over commutative rings, and give the final Gaussian factorization over an integral domain.

Four further modules supply the exhaustive root-of-unity and  $q$ -Catalan layer. The zero-definition/three-theorem `PrimitiveRootBlock.lean` surface is `gaussianBinomial_isPrimitiveRoot_eq_zero`, `neg_one_pow_mul_pow_choose_two`, and `finiteQPochhammerIn_isPrimitiveRoot`. In a commutative integral domain, a primitive  $d$ -th root  $\zeta$  makes  $\begin{bmatrix} d \\ k \end{bmatrix}_\zeta = 0$  for  $0 < k < d$ ; for  $d > 0$ , its top phase is  $(-1)^d \zeta^{\binom{d}{2}} = -1$ , and consequently  $(y; \zeta)_d = 1 - y^d$ .

The zero-definition/seven-theorem `QLucas.lean` surface is `add_mul_add_sub_one`, `choose_two_add`, `coeff_finiteQPochhammerIn_neg_X`, `finiteQPochhammerIn_neg_X_block`, `coeff_block_pow_mul`, `pow_choose_two_add_mul_eq`, and `gaussianBinomial_q_lucas`. The first two are natural-number quadratic identities. The polynomial coefficient and block formulas compare the finite  $q$ -binomial expansion with complete primitive-root blocks. If  $d > 0$ ,  $\zeta$  is a primitive  $d$ -th root in a commutative integral domain, and  $b, s < d$ , the final theorem is

$$\begin{bmatrix} ad + b \\ rd + s \end{bmatrix}_\zeta = \binom{a}{r} \begin{bmatrix} b \\ s \end{bmatrix}_\zeta.$$

Its same-shaped `two_mul_choose_two` helper is private; the unique public `two_mul_choose_two` belongs to `QChuVandermonde.lean`.

The terminating summation layer consists of `TwoPhiOneReversal.lean` (2+12) and `QChuVandermonde.lean` (0+10). It proves the finite-to-tsum bridge, reflected-parameter involution, one- and two-step reversals, both  $q$ -Chu evaluations, full-domain actual-tsum wrappers, and a separately named by-reversal route. The displayed  $q$ -Chu formulas and reversal lemma are exact under their displayed rational/nonvanishing hypotheses. The claim that reversal alone yields the second formula on the full displayed domain remains partial: the named by-reversal theorem additionally assumes  $C \neq 0$  and  $(A; q)_n \neq 0$ , whereas the full-domain theorem uses the direct denominator-cleared  $q$ -Cauchy route. Rational continuation and a cleared commutative-ring extension remain unformalized.

The zero-definition/four-theorem `JacobiTwoSquareCount.lean` leaf proves `sumSqRep_two`

`_eq_four_mul_twoSquareDivisorSum` for every nonzero natural input and its prime-product form under the explicit even-valuation condition at primes congruent to three modulo four. Its `theta_sq_eq_chi4_lambert` and `theta_sq_eq_odd_lambert` theorems instantiate the reusable `TwoSquareTheorem.lean` kernels over every complete normed field under the sole analytic hypothesis  $\|q\| < 1$ .

The zero-definition/three-theorem `CyclotomicDivisibility.lean` surface is `cyclotomic_exponent_eq_one_iff`, `cyclotomic_dvd_gaussianBinomial_iff`, and `gaussianBinomial_mul_isPrimitiveRoot`. For  $k \leq n$  and  $d > 0$ , the Gaussian cyclotomic exponent equals one exactly when  $n \bmod d < k \bmod d$ ; over  $\mathbb{Q}[X]$  this is exactly the criterion for  $\Phi_d$  to divide  $\begin{bmatrix} n \\ k \end{bmatrix}_X$ . In a commutative integral domain, a primitive  $n$ -th root with  $n > 0$  also gives  $\begin{bmatrix} an \\ bn \end{bmatrix}_\zeta = \begin{bmatrix} a \\ b \end{bmatrix}$ .

Finally, `QCatalan.lean` has the single noncomputable definition `qCatalan` and exactly eleven theorems: `map_qInt`, `qInt_X_monic`, `qInt_X_natDegree`, `X_sub_one_mul_qInt`, `qInt_X_eq_prod_cyclotomic`, `qInt_X_dvd_gaussianBinomial_rat`, `qInt_X_dvd_gaussianBinomial_int`, `qInt_X_mul_qCatalan`, `qCatalan_natDegree`, `qCatalan_eval_one_mul`, and `qCatalan_eval_one`. The semiring-level naturality theorem and commutative-ring  $q$ -integer formulas lead to  $[n+1]_X \mid \begin{bmatrix} 2n \\ n \end{bmatrix}_X$  over both  $\mathbb{Q}[X]$  and  $\mathbb{Z}[X]$ . Thus the integral polynomial

$$C_n(X) = \frac{\begin{bmatrix} 2n \\ n \end{bmatrix}_X}{[n+1]_X}$$

has natural degree  $n(n-1)$ , satisfies  $(n+1)C_n(1) = \begin{pmatrix} 2n \\ n \end{pmatrix}$ , and evaluates to the ordinary Catalan number.

Four denominator-free product identities make this recursion especially useful downstream. For natural  $m, n$ , and for  $k \leq n$ ,

$$(a; q)_{n+1} = (1-a)(aq; q)_n, \quad (494)$$

$$(a; q)_{m+n} = (a; q)_m (aq^m; q)_n, \quad (495)$$

$$(q; q)_k \begin{bmatrix} n \\ k \end{bmatrix}_q = (q^{n-k+1}; q)_k, \quad (496)$$

$$(q; q)_n = (q; q)_k (q; q)_{n-k} \begin{bmatrix} n \\ k \end{bmatrix}_q. \quad (497)$$

They are `finiteQPochhammerIn_succ_shift`, `finiteQPochhammerIn_add`, `finiteQPochhammerIn_self_mul_gaussianBinomial`, and `finiteQPochhammerIn_self_eq_mul_mul_gaussianBinomial`, respectively, again in `FiniteQBinomialCore.lean`. All four hold over every commutative ring, including empty products, zero divisors, positive characteristic,  $q = 0$ , and roots of unity. The final identity is the full denominator-free replacement for the usual quotient definition of a Gaussian coefficient.

The companion `QPochhammerElementaryIdentities.lean` has exactly thirteen public theorems. Its commutative-ring layer consists of the two unit-valued base reversals `finiteQPochhammerIn_base_reversal_units` and `finiteQPochhammerIn_inv_base_reversal_units`, the root-safe terminating numerator `prod_pow_sub_pow_eq_finiteQPochhammerIn`, the first-column clearer `one_sub_mul_gaussianBinomial_one`, and the two adjacent cross identities `gaussianBinomial_adjacent_mul` and `gaussianBinomial_row_adjacent_mul`. In human

notation these identities are

$$\begin{aligned} \prod_{j < k} (q^N - q^j) &= (-1)^k q^{\binom{k}{2}} (q^{N-k+1}; q)_k \quad (k \leq N), \\ (1 - q^{k+1}) \begin{bmatrix} n \\ k+1 \end{bmatrix}_q &= (1 - q^{n-k}) \begin{bmatrix} n \\ k \end{bmatrix}_q, \\ (1 - q^{n-k}) \begin{bmatrix} n \\ k \end{bmatrix}_q &= (1 - q^n) \begin{bmatrix} n-1 \\ k \end{bmatrix}_q. \end{aligned}$$

These formulas remain valid at roots of unity; the first also permits zero divisors and  $q = 0$ , while the last two are total for all natural  $n, k$ . On and above the row boundary, zero extension makes both sides of each adjacent identity vanish.

The seven field declarations are `finiteQPochhammerIn_base_reversal`, `finiteQPochhammerIn_inv_base_reversal`, `pow_mul_finiteQPochhammerIn_inv_pow_eq`, `finiteQPochhammerIn_inv_pow_eq_self_div`, `finiteQPochhammerIn_inv_pow_eq_zero_of_lt`, `gaussianBinomial_adjacent_div`, and `gaussianBinomial_row_adjacent_div`. Both numerical base-reversal orientations require exactly  $a \neq 0$  and  $q \neq 0$ . For  $k \leq N$ , the cleared terminating identity needs only  $q \neq 0$ ; the quotient

$$(q^{-N}; q)_k = \frac{(-1)^k q^{\binom{k}{2}} (q; q)_N}{q^{Nk} (q; q)_{N-k}}$$

additionally requires  $(q; q)_{N-k} \neq 0$ , while the vanishing for  $k > N$  again needs only  $q \neq 0$ . For  $k < n$ , the adjacent quotient corollaries require, respectively,  $\begin{bmatrix} n \\ k \end{bmatrix}_q \neq 0$  and  $1 - q^{k+1} \neq 0$ , or  $\begin{bmatrix} n-1 \\ k \end{bmatrix}_q \neq 0$  and  $1 - q^{n-k} \neq 0$ ; neither assumes  $q \neq 0$ .

The four-theorem module `QBinomialReciprocity.lean` supplies the exact palindromic counterpart. Its unit-valued theorem `gaussianBinomial_reciprocity_units` holds in every commutative semiring, including semirings with zero divisors, and is total in both indices:

$$q^{k(n-k)} \begin{bmatrix} n \\ k \end{bmatrix}_{q^{-1}} = \begin{bmatrix} n \\ k \end{bmatrix}_q.$$

The wrapper `gaussianBinomial_reciprocity` assumes a semifield and only  $q \neq 0$ . The remaining two theorems, `gaussianBinomial_neg_one_eq_zero_of_odd_degree` and `gaussianBinomial_neg_one_even_odd_eq_zero`, show over every commutative ring that odd Gaussian degree forces vanishing at  $q = -1$ , and in particular  $\begin{bmatrix} 2a \\ 2b+1 \end{bmatrix}_{-1} = 0$ . The latter is total above the row as well. These proofs use neither a  $q$ -Pochhammer quotient nor domain cancellation.

Over the rationals, `finiteQPochhammerIn_rat_eq` identifies the generic product with the established finite  $q$ -Pochhammer notation. At  $q = 1/2$ , `gaussianBinomial_half_eq_halfQBinomial` identifies the recursive coefficient with the established rational quotient, and `halfQBinomial_theorem` is now the direct specialization of the ring theorem.

**Theorem 8.20** (Denominator-free  $q$ -Vandermonde and shifted central forms). *Let  $R$  be a commutative semiring, let  $q \in R$ , and let all displayed unsigned indices be natural numbers. Extend a Gaussian row to integer lower indices by*

$$\mathcal{G}_q(n, t) = \begin{cases} \begin{bmatrix} n \\ t \end{bmatrix}_q, & 0 \leq t, \\ 0, & t < 0, \end{cases} \quad t \in \mathbb{Z}, \quad (498)$$

where the natural coefficient already vanishes for  $n < t$ . This extension has the total reflection law

$$\mathcal{G}_q(n, n-t) = \mathcal{G}_q(n, t) \quad (t \in \mathbb{Z}). \quad (499)$$

The two zero-extended Vandermonde orientations are

$$\begin{bmatrix} m+n \\ r \end{bmatrix}_q = \sum_{j=0}^r q^{(m-j)(r-j)} \begin{bmatrix} m \\ j \end{bmatrix}_q \begin{bmatrix} n \\ r-j \end{bmatrix}_q, \quad (500)$$

$$= \sum_{j=0}^r q^{j(n-(r-j))} \begin{bmatrix} m \\ j \end{bmatrix}_q \begin{bmatrix} n \\ r-j \end{bmatrix}_q. \quad (501)$$

Here subtraction is natural subtraction; outside the admissible support one of the zero-extended Gaussian coefficients vanishes. Consequently no condition  $r \leq m+n$  is needed. The central specialization is

$$\begin{bmatrix} m+n \\ m \end{bmatrix}_q = \sum_{j=0}^m q^{j^2} \begin{bmatrix} m \\ j \end{bmatrix}_q \begin{bmatrix} n \\ j \end{bmatrix}_q. \quad (502)$$

Equivalently, removing exactly the zero-extended terms gives the literal Durfee support

$$\begin{bmatrix} m+n \\ m \end{bmatrix}_q = \sum_{j=0}^{\min(m,n)} q^{j^2} \begin{bmatrix} m \\ j \end{bmatrix}_q \begin{bmatrix} n \\ j \end{bmatrix}_q. \quad (503)$$

For every  $N, k \in \mathbb{N}_0$ , including  $N < k$ , the positive shifted form is

$$\begin{bmatrix} 2N \\ N+k \end{bmatrix}_q = \sum_{\ell=0}^N q^{\ell(\ell+k)} \begin{bmatrix} N \\ \ell \end{bmatrix}_q \begin{bmatrix} N \\ \ell+k \end{bmatrix}_q. \quad (504)$$

If  $k \leq N$ , row reflection gives the same right side for  $\begin{bmatrix} 2N \\ N-k \end{bmatrix}_q$ , and its literal-support form is

$$\begin{bmatrix} 2N \\ N-k \end{bmatrix}_q = \sum_{\ell=k}^N q^{\ell(\ell-k)} \begin{bmatrix} N \\ \ell \end{bmatrix}_q \begin{bmatrix} N \\ \ell-k \end{bmatrix}_q. \quad (505)$$

Most strongly, for every  $k \in \mathbb{Z}$ , put  $t^{\natural} = \max(t, 0) \in \mathbb{N}_0$ . Then the two natural cases combine into the literal finite-support integer sum

$$\mathcal{G}_q(2N, N+k) = \sum_{\ell \in \mathbb{Z}} q^{\ell^{\natural}(\ell+k)^{\natural}} \mathcal{G}_q(N, \ell) \mathcal{G}_q(N, \ell+k). \quad (506)$$

Only  $0 \leq \ell \leq N$  can contribute; a nonzero term also has  $0 \leq \ell+k \leq N$ , so its exponent is exactly  $\ell(\ell+k) = \ell^2 + \ell k$ , as in the signed report formula. In particular, both sides vanish automatically when  $|k| > N$ . No identity in this theorem uses division, cancellation, a nonvanishing hypothesis, or an exclusion of  $q = 0$ , roots of unity, positive characteristic, or zero divisors.

*Formal proof.* Equations (500) and (501) are `gaussianBinomial_add_vandermonde` and `gaussianBinomial_add_vandermonde'`. The definition (498), its cast, lower-zero, upper-zero, and total-reflection laws are respectively `gaussianBinomialInt`, `gaussianBinomialInt_ofNat`, `gaussianBinomialInt_eq_zero_of_neg`, `gaussianBinomialInt_eq_zero_of_lt`, and `gaussianBinomialInt_symm` in `FiniteQBinomialCore.lean`. The full-range and literal- $\min(m, n)$  central identities are `gaussianBinomial_add_central` and `gaussianBinomial_add_central_min`; the positive shift is `gaussianBinomial_two_mul_add_shifted_central`; and the two negative-shift presentations are `gaussianBinomial_two_mul_sub_shifted_central` and `gaussianBinomial_two_mul_sub_shifted_central ICC`. The total signed finite-range and integer-sum forms are `gaussianBinomial_two_mul_int_shifted_central` and `gaussianBinomial_two_mul_int_shifted_central_finsum`. These nine theorems exhaust the public surface of `QBinomialVandermonde.lean`. The three natural-shift theorems retain their stated domains, while the last two are total in  $k \in \mathbb{Z}$ .  $\square$

**Theorem 8.21** (Weighted subsets and scaled  $q$ -binomial inversion). *Let  $R$  be a commutative ring and  $q, s \in R$ ; the first identity below already holds when  $R$  is only a commutative semiring. For  $j \leq k \leq n$ , the Gaussian chain identity is*

$$\begin{bmatrix} n \\ k \end{bmatrix}_q \begin{bmatrix} k \\ j \end{bmatrix}_q = \begin{bmatrix} n \\ j \end{bmatrix}_q \begin{bmatrix} n-j \\ k-j \end{bmatrix}_q. \quad (507)$$

For every  $m, n \in \mathbb{N}_0$ , the alternating row specializes exactly to

$$\sum_{k=0}^n (-1)^k q^{\binom{k}{2} + mk} \begin{bmatrix} n \\ k \end{bmatrix}_q = (q^m; q)_n, \quad (508)$$

$$\sum_{k=0}^n (-1)^k q^{\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q = \begin{cases} 1, & n = 0, \\ 0, & n > 0. \end{cases} \quad (509)$$

The same finite product has the two exact weighted-subset faces

$$\sum_{\substack{T \subseteq \{0, \dots, N-1\} \\ |T|=r}} q^{\sum_{t \in T} t} = q^{\binom{r}{2}} \begin{bmatrix} N \\ r \end{bmatrix}_q, \quad (510)$$

$$\sum_{\substack{T \subseteq \{1, \dots, N\} \\ |T|=r}} q^{\sum_{t \in T} t} = q^{\binom{r+1}{2}} \begin{bmatrix} N \\ r \end{bmatrix}_q. \quad (511)$$

Both formulas are total in  $N, r \in \mathbb{N}_0$ ; above the Gaussian diagonal both sides vanish.

Define total lower-triangular kernels on natural indices by

$$K_{n,k}^{q,s} = s^{n-k} \begin{bmatrix} n \\ k \end{bmatrix}_q, \quad (512)$$

$$L_{n,k}^{q,s} = (-s)^{n-k} q^{\binom{n-k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q. \quad (513)$$

Natural subtraction causes no ambiguity above the diagonal because the Gaussian coefficient is zero there. With an empty sum when  $n < j$ , both finite convolution orders are the Kronecker delta:

$$\sum_{k \in [j, n] \cap \mathbb{N}_0} L_{n,k}^{q,s} K_{k,j}^{q,s} = \delta_{n,j}, \quad (514)$$

$$\sum_{k \in [j, n] \cap \mathbb{N}_0} K_{n,k}^{q,s} L_{k,j}^{q,s} = \delta_{n,j}. \quad (515)$$

In particular the scale  $s$  is independent of  $q$  and need not be nonzero or invertible.

More generally, let  $M$  be an additive commutative monoid and an  $R$ -module. For sequences  $a, b : \mathbb{N}_0 \rightarrow M$ , put

$$(\mathcal{T}_{q,s} a)_n = \sum_{k=0}^n K_{n,k}^{q,s} a_k, \quad (\mathcal{T}_{q,s}^{-1} b)_n = \sum_{k=0}^n L_{n,k}^{q,s} b_k, \quad (516)$$

where multiplication denotes the scalar action on  $M$ . Then

$$\mathcal{T}_{q,s}^{-1} \mathcal{T}_{q,s} = 1, \quad \mathcal{T}_{q,s} \mathcal{T}_{q,s}^{-1} = 1, \quad b = \mathcal{T}_{q,s} a \iff a = \mathcal{T}_{q,s}^{-1} b. \quad (517)$$

At  $s = 1$  this is classical  $q$ -binomial inversion, now for module-valued rather than merely scalar-valued sequences.

There are two useful exact specializations. The denominator-free geometric Lagrange numerator in (540) satisfies globally

$$N_{n,k}(q) = L_{n,k}^{q,q}, \quad (518)$$

including  $n < k$ , without an in-range hypothesis. At Gaussian base  $Q = q^2$  and scale  $s = -q$ , put

$$C'_{n,k} = K_{n,k}^{q^2, -q} = (-q)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix}_{q^2}, \quad B_{n,k} = L_{n,k}^{q^2, -q} = q^{(n-k)^2} \begin{bmatrix} n \\ k \end{bmatrix}_{q^2}. \quad (519)$$

The two coefficient definitions and their displayed pointwise closed forms already hold in an arbitrary, possibly noncommutative ring. Commutativity is needed only for the two convolution theorems, under which both total finite products satisfy  $C'B = BC' = I$ . These identities use no analytic or nonvanishing assumption. By contrast, the normalized Lagrange quotient in (543) retains its separate field, finite-node injectivity, and displayed in-range hypotheses.

*Formal proof.* The eight public theorems of `BitPositionQBinomial.lean` are, exhaustively, `prod_one_add_mul_pow_eq_gaussianBinomial`, `prod_one_add_pow_eq_sum_gaussianBinomial`, `sum_powersetCard_two_pow`, `sum_pow_bitPositionSum_filter_eq_gaussianBinomial`, `sum_pow_bitPositionSum_filter_eq_gaussianBinomial'`, `sum_pow_sum_powersetCard_eq_gaussianBinomial`, `sum_pow_sum_powersetCard_Icc_eq_gaussianBinomial`, and `gaussianBinomial_one_eq_choose`. The sixth and seventh declarations are (510) and (511).

The four definitions and seven theorems of `QBinomialInversion.lean` are, exhaustively, `gaussianBinomial_mul`, `sum_gaussianBinomial_alternating_mul_pow`, `sum_gaussianBinomial_alternating`, `gaussianBinomialKernel`, `gaussianBinomialInverseKernel`, `scaledGaussianBinomialKernel`, `scaledGaussianBinomialInverseKernel`, `gaussianBinomialKernel_left_orthogonality`, `gaussianBinomialKernel_right_orthogonality`, `scaledGaussianBinomialKernel_left_orthogonality`, and `scaledGaussianBinomialKernel_right_orthogonality`. They prove (507), (508)–(509), and both unscaled and scaled forms of (514)–(515).

The one definition and one theorem of `FiniteTriangularTransform.lean` are, exhaustively, `lowerTriangularTransform` and `lowerTriangularTransform_comp`. For a possibly noncommutative semiring acting on an additive commutative monoid, the theorem turns a total ordered convolution  $\sum_{k \in [j, n]} K_{n,k} L_{k,j} = \delta_{n,j}$  into equality of the whole composite sequence function with the input. Its hypothesis also covers  $n < j$ , when the interval is empty. The refactored Gaussian proof below reuses this generic finite-sum theorem; the Gaussian kernels' zero extension above the row supplies the same total convention.

The four definitions and four theorems of `QBinomialTransform.lean` are, exhaustively, `scaledGaussianBinomialTransform`, `scaledGaussianBinomialInverseTransform`, `scaledGaussianBinomialInverseTransform_transform`, `scaledGaussianBinomialTransform_inverseTransform`, `scaledGaussianBinomial_inversion`, `gaussianBinomialTransform`, `gaussianBinomialInverseTransform`, and `gaussianBinomial_inversion`. They prove (516)–(517) as whole-function equalities. The two forward-transform definitions need only a semiring acting on an additive commutative monoid; the two signed inverse definitions need a ring with the same target assumptions. The four inversion theorems require a commutative ring, again acting on an additive commutative monoid. No additive inverses are required in the target.

The global bridge `geometricQBinomialWeightNumerator_eq_scaledGaussianBinomialInverseKernel` belongs to `GeometricQBinomialLagrange.lean`. It proves (518) for every pair of natural indices, including above the diagonal, and makes the geometric numerator the scale- $q$  specialization of the characteristic polynomial in [theorem 8.22](#).

Finally, the two definitions and four theorems of `QBinomialInversionSpecializations.lean` are, exhaustively, `qGaussianResidualCoeff`, `qGaussianReconstructionCoeff`, `qGaussianResidualCoeff_eq`, `qGaussianReconstructionCoeff_eq`, `qGaussianReconstructionCoeff_residualCoeff_delta`, and `qGaussianResidualCoeff_reconstructionCoeff_delta`. These are the two closed forms in (519) and both convolution orders. The

two definitions and the two pointwise equality theorems require only  $[\text{Ring } R]$ ; exactly the two convolution-delta theorems require  $[\text{CommRing } R]$ .  $\square$

**Theorem 8.22** (Scaled Gaussian characteristic and exact  $q$ -difference annihilation). *Let  $R$  be a commutative ring and let  $q, s, z \in R$ . With the total inverse Gaussian row from (513),*

$$L_{n,k}^{q,s} = (-s)^{n-k} q^{\binom{n-k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q,$$

*one has the homogeneous, denominator-free characteristic identity*

$$\sum_{k=0}^n L_{n,k}^{q,s} z^k = \prod_{j=0}^{n-1} (z - sq^j). \quad (520)$$

*In particular, taking  $s = 1$  and  $z = q^d$  gives every monomial moment:*

$$\sum_{k=0}^n L_{n,k}^{q,1} (q^k)^d = \prod_{j=0}^{n-1} (q^d - q^j). \quad (521)$$

*The product contains the zero factor indexed by  $j = d$  whenever  $d < n$ .*

*More generally, let  $A$  be a semiring, let  $\varphi: A \rightarrow R$  be a semiring homomorphism, and let  $P \in A[X]$ . Write  $P^\varphi \in R[X]$  for coefficientwise scalar extension. If  $\deg P \leq n$ , then the row extracts the mapped top coefficient with its exact surviving moment:*

$$\sum_{k=0}^n L_{n,k}^{q,1} P^\varphi(q^k) = \varphi([X^n]P) \prod_{j=0}^{n-1} (q^n - q^j). \quad (522)$$

*If instead  $\deg P < n$ , then*

$$\sum_{k=0}^n L_{n,k}^{q,1} P^\varphi(q^k) = 0. \quad (523)$$

*All four statements include  $n = 0$  and the zero polynomial. The geometric nodes may collide and the surviving product in (522) may itself vanish. No division, distinctness, nonzero or invertible base, domain, characteristic, topology, or convergence assumption is present.*

*Formal proof.* The public surface of `QDifferenceAnnihilation.lean` consists exhaustively of four theorems: `sum_scaledGaussianBinomialInverseKernel_mul_pow`, `sum_gaussianBinomialInverseKernel_mul_geometric_pow`, `qDifference_sum_eval2_eq_map_coeff_mul`, and `qDifference_sum_eval2_eq_zero_of_degree_lt`. They prove, respectively, (520)–(523). The first two specialize the scaled inverse-kernel algebra over an arbitrary commutative ring; the last two use the generic finite-polynomial-functional API to retain semiring coefficients and arbitrary scalar extension into that ring. Thus the hypotheses and every boundary case in the statement are literal formal interfaces, not consequences of an unrecorded distinct-geometric-node assumption.  $\square$

**Theorem 8.23** (Finite  $q$ -Cauchy identities and the  $q$ -Bernstein partition). *Let  $R$  be a commutative ring, let  $q, u, v, a, b, c, x \in R$ , and let  $n \in \mathbb{N}_0$ . The primary and reflected finite  $q$ -Cauchy identities are*

$$(uv; q)_n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (u; q)_k v^k (v; q)_{n-k}, \quad (524)$$

$$= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (v; q)_k v^{n-k} (u; q)_{n-k}. \quad (525)$$

Define the denominator-free  $q$ -Bernstein term by

$$B_{n,k}^{(q)}(x) = \begin{bmatrix} n \\ k \end{bmatrix}_q x^k (x; q)_{n-k}. \quad (526)$$

Then its full row is a partition of unity,

$$\sum_{k=0}^n B_{n,k}^{(q)}(x) = 1. \quad (527)$$

Finally, the second finite Cauchy convolution is

$$(ac; q)_n (bc; q)_n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (a; q)_k (b; q)_k c^k (c; q)_{n-k} (abcq^k; q)_{n-k}. \quad (528)$$

All four identities are total in the displayed parameters and degree. They include  $n = 0$ ,  $q = 0$ , roots of unity, positive characteristic, and zero divisors, and use no quotient, cancellation, domain, or nonvanishing hypothesis.

*Formal proof and exhaustive API.* Equations (524) and (525) are `finite_qCauchy_identity` and `finite_qCauchy_identity_reflected`. The first also has the compatibility spelling `finiteQPochhammerIn_mul_eq_sum_gaussianBinomial`; (526) is the definition `qBernsteinBasis`; (527) is `sum_qBernsteinBasis`; and (528) is `finite_qCauchy_second_identity`. These are exhaustively the one public definition and five public theorems of `QBinomialCauchy.lean`. The partition of unity is the primary identity at  $u = 0$ ; the second identity is proved by expanding a finite triangle, reindexing it, and applying both orientations of finite  $q$ -Cauchy. Thus the proof never divides by a  $q$ -Pochhammer factor.  $\square$

**Theorem 8.24** (Weighted Pascal and elementary-complete orthogonality). *Let  $R$  be a commutative semiring and let  $a : \iota \rightarrow R$  be a family on a finite index type. Write  $e_r(a)$  for the evaluation of the universal elementary symmetric polynomial and  $h_r(a)$  for the existing complete homogeneous evaluation. Then  $e_0(a) = 1$ , and reindexing by any equivalence  $e : \kappa \simeq \iota$  leaves the value unchanged:*

$$e_r(a \circ e) = e_r(a). \quad (529)$$

For a family indexed by a distinguished point and a finite tail, the two weighted-Pascal laws are

$$e_{r+1}(x, a) = x e_r(a) + e_{r+1}(a), \quad (530)$$

$$h_{r+1}(x, a) = x h_r(x, a) + h_{r+1}(a). \quad (531)$$

The elementary identity is also exposed on a literal nonempty finite list as

$$e_{r+1}(a_0, a_1, \dots, a_m) = a_0 e_r(a_1, \dots, a_m) + e_{r+1}(a_1, \dots, a_m). \quad (532)$$

If  $R$  is a commutative ring, then for every degree  $n \in \mathbb{N}_0$ ,

$$\sum_{k=0}^n (-1)^k e_k(a) h_{n-k}(a) = \begin{cases} 1, & n = 0, \\ 0, & n > 0. \end{cases} \quad (533)$$

This includes the empty family and degree zero and uses no division, characteristic, domain, or nonvanishing assumption.



*Formal proof and exhaustive API.* The definition is `elementarySymmetricEval`; its identification with Mathlib’s evaluated universal `MvPolynomial.esymm` is `elementarySymmetricEval_eq_eval_esymm`. The equation  $e_0 = 1$ , (529), (530), and (532) are respectively `elementarySymmetricEval_zero`, `elementarySymmetricEval_comp_equiv`, `elementarySymmetricEval_option_succ`, and `elementarySymmetricEval_fin_succ`. Equation (533) is `sum_elementarySymmetricEval_mul_completeHomogeneousEval`. These are exhaustively the one public definition and six public theorems of `SymmetricFunctionOrthogonality.lean`. The companion complete recurrence (531) is `completeHomogeneousEval_option_succ` in `CompleteHomogeneous.lean`. The structural elementary identities and both Pascal recurrences hold already over commutative semirings; subtraction is needed only for the final alternating convolution.  $\square$

**Theorem 8.25** (Formal generating series for a finite alphabet). *Let  $R$  be a commutative semiring and let  $a : \iota \rightarrow R$  be indexed by a finite type. In the formal power-series ring  $R[[X]]$ , set*

$$E_a(X) = \sum_{r \geq 0} e_r(a) X^r, \quad H_a(X) = \sum_{r \geq 0} h_r(a) X^r.$$

*Their coefficients are exactly the displayed symmetric-function evaluations, and*

$$E_a(X) = \prod_{i \in \iota} (1 + a_i X). \quad (534)$$

*If  $R$  is a commutative ring, then*

$$E_{-a}(X) H_a(X) = 1, \quad H_a(X) = \text{invOfUnit}(E_{-a}(X), 1). \quad (535)$$

*In particular, for every  $q \in R$  and  $n \in \mathbb{N}_0$ ,*

$$\left( \prod_{j=0}^n (1 - q^j X) \right) \left( \sum_{k \geq 0} \begin{bmatrix} n+k \\ k \end{bmatrix}_q X^k \right) = 1 \quad \text{in } R[[X]]. \quad (536)$$

*These are formal identities, including the empty alphabet and all degenerate values of  $q$ . They do not assert evaluation of the series at a scalar or analytic convergence under  $|a_i z| < 1$  or  $|z| < 1$ .*

*Formal proof and exhaustive API.* The two definitions and six theorems of `SymmetricFunctionGenerating.lean` are, exhaustively, `elementarySymmetricGeneratingSeries`, `completeHomogeneousGeneratingSeries`, `coeff_elementarySymmetricGeneratingSeries`, `coeff_completeHomogeneousGeneratingSeries`, `elementarySymmetricGeneratingSeries_eq_prod`, `elementarySymmetricGeneratingSeries_neg_mul_completeHomogeneousGeneratingSeries`, `completeHomogeneousGeneratingSeries_eq_invOfUnit_elementarySymmetricGeneratingSeries_neg`, and `prod_one_sub_qPow_X_mul_gaussianBinomialGeneratingSeries`. The definitions, coefficient identities, and finite elementary product need only a commutative semiring. The reciprocity, canonical-inverse, and finite Gaussian reciprocal identities require a commutative ring. Thus the Lean surface proves precisely the formal-power-series halves of the corresponding analytic formulas.  $\square$

**Theorem 8.26** (Weighted symmetric-function inversion). *Let  $R$  be a commutative ring, let  $M$  be an additive commutative monoid and an  $R$ -module, and let  $w : \iota \rightarrow R$  be indexed by a finite type. For  $a, b : \mathbb{N}_0 \rightarrow M$ , define*

$$(\mathcal{H}_w a)_m = \sum_{j=0}^m h_{m-j}(w) a_j, \quad (537)$$

$$(\mathcal{E}_w b)_m = \sum_{j=0}^m (-1)^{m-j} e_{m-j}(w) b_j, \quad (538)$$

where the products with  $a_j, b_j$  denote scalar actions. Then, as equalities of whole sequence functions,

$$\mathcal{E}_w \mathcal{H}_w a = a, \quad \mathcal{H}_w \mathcal{E}_w b = b, \quad b = \mathcal{H}_w a \iff a = \mathcal{E}_w b. \quad (539)$$

Both scalar kernels are extended by zero above the diagonal, so the kernel orthogonalities are total even when the outer interval is empty. There is no division, topology, domain, characteristic, or nonvanishing assumption.

*Formal proof and exhaustive API.* The four definitions and five theorems of `SymmetricFunctionTransform.lean` are, exhaustively, `completeHomogeneousKernel`, `signedElementaryKernel`, `completeHomogeneousKernel_left_orthogonality`, `completeHomogeneousKernel_right_orthogonality`, `completeHomogeneousTransform`, `signedElementaryTransform`, `signedElementaryTransform_completeHomogeneousTransform`, `completeHomogeneousTransform_signedElementaryTransform`, and `weightedSymmetricFunction_inversion`. The complete-homogeneous kernel and transform already need only a commutative semiring. The signed kernel, signed transform, both total orthogonalities, both inverse compositions, and the inversion equivalence use a commutative ring; every module-valued transform uses only an additive commutative monoid as its target. The two composition proofs reuse `lowerTriangularTransform_comp` from the one-definition, one-theorem generic surface of `FiniteTriangularTransform.lean`.  $\square$

**Theorem 8.27** (Reversed Gaussian row, Lagrange weights, and all residual moments). *Let  $R$  be a commutative ring and  $q \in R$ . The zero-extended numerator row is*

$$N_{n,k}(q) = \begin{cases} (-1)^{n-k} q^{\binom{n-k+1}{2}} [n]_q, & k \leq n, \\ 0, & n < k. \end{cases} \quad (540)$$

The Lean definition is initially written with the reversed index  $n - k$ , but the division-free symmetry (491) proves the forward form displayed here. In fact the zero-extended identity remains valid for every pair  $n, k$  when natural subtraction is read with Lean’s truncation convention. For every  $z \in R$ ,

$$\sum_{k=0}^n N_{n,k}(q) z^k = \prod_{j=0}^{n-1} (z - q^{j+1}), \quad (541)$$

and, for every  $0 \leq d \leq n$ ,

$$\sum_{k=0}^n N_{n,k}(q) (q^k)^d = (q; q)_n 0^d. \quad (542)$$

Now let  $R$  be a field and suppose that the nodes  $1, q, \dots, q^n$  are distinct. If  $\lambda_{n,k}^{(q)}$  is the evaluation-at-zero Lagrange weight belonging to  $q^k$ , then  $(q; q)_n \neq 0$  and, for  $k \leq n$ ,

$$\lambda_{n,k}^{(q)} = \frac{N_{n,k}(q)}{(q; q)_n}, \quad (q; q)_n \lambda_{n,k}^{(q)} = N_{n,k}(q). \quad (543)$$

Finally every positive residual moment has the single closed form

$$\sum_{k=0}^n \lambda_{n,k}^{(q)} (q^k)^d = (-1)^n q^{n(n+1)/2} \begin{bmatrix} d-1 \\ n \end{bmatrix}_q, \quad d > 0. \quad (544)$$

All statements include  $n = 0$ . The field-level formulas impose only distinctness of the displayed finite node set; they do not add a redundant global ban on  $q = 0$  or on roots of unity.

*Formal proof.* The numerator and its two branches are `geometricQBinomialWeightNumerator`, `geometricQBinomialWeightNumerator_eq_of_le`, and `geometricQBinomialWeightNumerator_eq_zero_of_lt`. Its forward-index forms on the natural range and globally are `geometricQBinomialWeightNumerator_eq_forward_of_le` and `geometricQBinomialWeightNumerator_eq_forward`. Its global inverse-kernel form is `geometricQBinomialWeightNumerator_eq_scaledGaussianBinomialInverseKernel`; this declaration is owned here by `GeometricQBinomialLagrange.lean` and proves (518). Equations (541) and (542) are `reversed_finite_qBinomial_theorem` and `sum_geometricQBinomialWeightNumerator_mul_pow`. Node distinctness gives the nonvanishing denominator by `finiteQPochhammerIn_self_ne_zero_of_injOn`; the quotient and cleared-denominator identities in (543) are `geometricLagrangeWeight_eq_gaussianBinomial_div` and `finiteQPochhammerIn_mul_geometricLagrangeWeight_eq`. The complete positive-degree formula (544) is `sum_geometricLagrangeWeight_mul_pow_eq_gaussianBinomial`. These declarations are all in `GeometricQBinomialLagrange.lean`; the finite-product identities in the preceding remark are in `FiniteQBinomialCore.lean`.  $\square$

Here  $q$  is the Gaussian base and the ratio of the geometric nodes; it is unrelated to the additive translation  $Q$  in the translated Fabius formulas below. Because Theorem 8.27 is base-generic, the explicit Gaussian–Lagrange identification and its residual moments are no longer research-frontier obligations at Gaussian base  $q = 1/4$ . Over rational  $0 < q < 1$ , `GeometricLagrangeQBinomial.lean` and `GeometricLagrangeQMoments.lean` additionally express the weights and all residual moments in the established quotient-defined  $q$ -Pochhammer and  $q$ -binomial notation, prove the residual sign, and compute the exact finite  $\ell^1$  norm of the row. Their dedicated quarter-base corollaries include `quarterGeometricLagrangeQMoment_eq_residual_qBinomial` and `sum_abs_quarterGeometricLagrangeWeight_eq_qPochhammer_ratio`. The whole geometric Richardson triangle now has the exact formal factorization

$$\sum_{n \geq 0} \left( \sum_{j=0}^n \lambda_{n,j}^{(q)} y_j \right) X^n = (qX; q)_\infty \sum_{j \geq 0} \frac{y_j}{(q; q)_j} X^j.$$

The report-facing field-valued statement is `geometricLagrangeRichardson_generating` in `GeometricRichardsonGenerating.lean`; it assumes only  $q \neq 0$  and is an identity of formal power series, so it makes no analytic convergence claim. The same module first proves the stronger convolution identity `geometricRichardsonTransform_generating` over an arbitrary commutative ring with totalized `Ring.inverse`, and its theorem `geometricRichardsonTransform_eq_sum_lagrange` identifies the coefficients with the established Lagrange row over a field. Its analytic companions `hasSum_geometricRichardsonTransform_mul_pow` and `hasSum_geometricLagrangeRichardson_mul_pow` require a complete normed field, strict nome contraction, and absolute summability of the normalized data series; the Lagrange form also assumes a nonzero nome. The exhaustive public surface is the three definitions `geometricRichardsonKernel`, `qPochhammerNormalizedDataSeries`, and `geometricRichardsonTransform`, and the seven theorems `coeff_rescale_qPochhammerSeries_eq_geometricRichardsonKernel`, `coeff_qPochhammerNormalizedDataSeries`, `geometricRichardsonTransform_generating`, `geometricRichardsonTransform_eq_sum_lagrange`, `geometricLagrangeRichardson_generating`, `hasSum_geometricRichardsonTransform_mul_pow`, and `hasSum_geometricLagrangeRichardson_mul_pow`. The generic unconditionally summable geometric filter and its exact Gaussian tsum tail are proved in Proposition 10.9. The finite row variation, with  $w_q(n, j)$  the  $q$ -Toeplitz weight `qToeplitzWeight` from `GeneralQConditionNumber.lean`, also has its exact infinite limit: for every  $0 \leq q < 1$ ,

$$\sum_{j \leq n} |w_q(n, j)| \longrightarrow \frac{(-q; q)_\infty}{(q; q)_\infty}.$$

This is `tendsto_sum_abs_qToeplitzWeight` in `LimitConditionNumber.lean`; the same module proves positivity of the denominator on the full range and the effective lower bound  $1/(1 - q) \leq (-q; q)_\infty / (q; q)_\infty$ , and promotes that bound to the literal divergence theorem `tendsto_qConditionNumberLimit_atTop_at_one_left`. The actual infinite Rvachev sinc-product instantiation, at the quarter base and at a generic complex squared base under finite-node injectivity, is now exactly [theorem 10.10](#). What remains in the research-frontier volume is the different finite-prefix problem for  $P_{b,n}$ : its quotient and Bell-coefficient package, the analytic sign and remainder/bracketing bounds, uniform accelerated and derivative convergence, and asymptotics.

The same finite row acts coefficientwise on every formal power series. For  $q \in \mathbb{Q}$ ,  $p, n \in \mathbb{N}_0$ , and  $D(X) \in \mathbb{Q}[[X]]$ , define

$$\mathcal{R}_{q,p,n}D := \sum_{j=0}^p \lambda_{p,j}^{(q)} D(q^{n+j}X). \quad (545)$$

Then, for every  $m \in \mathbb{N}_0$ , without a node-distinctness assumption,

$$[X^m]\mathcal{R}_{q,p,n}D = [X^m]D q^{nm} \sum_{j=0}^p \lambda_{p,j}^{(q)} (q^j)^m. \quad (546)$$

If  $q \neq 0$  and  $(q; q)_p \neq 0$ , the constant coefficient is preserved and degrees  $1, \dots, p$  vanish. If moreover  $0 < q < 1$  and  $m > p$ , then

$$[X^m]\mathcal{R}_{q,p,n}D = [X^m]D (-1)^p q^{\binom{p+1}{2}} \begin{bmatrix} m-1 \\ p \end{bmatrix}_q q^{nm}. \quad (547)$$

The filter itself is `geometricRichardsonPowerSeriesFilter` in `QuarterCatalanRichards` on `.lean`. Coefficient extraction from a finite rescaling filter is `finiteRescaleFilter_coeff`; equations (546) and (547) are `geometricRichardsonPowerSeriesFilter_coeff` and `geometricRichardsonPowerSeriesFilter_coeff_eq_qBinomial`. The constant and cancelled-degree statements are `geometricRichardsonPowerSeriesFilter_coeff_zero` and `geometricRichardsonPowerSeriesFilter_coeff_eq_zero`. Thus the quarter Catalan calculation below is a specialization of a reusable filter theorem, not an isolated coefficient computation.

There is nevertheless a second exact quarter-base result at the level of formal power series. Let  $D(Q) = \sum_{m \geq 0} \delta_m Q^m \in \mathbb{Q}[[Q]]$  be the unique series with zero constant coefficient satisfying

$$D(Q) + 4D(Q)^2 = \frac{4}{9}Q. \quad (548)$$

Its positive-degree coefficients are

$$\delta_{r+1} = (-1)^r C_r \frac{2^{4r+2}}{9^{r+1}}, \quad (549)$$

where  $C_r$  is the  $r$ -th Catalan number. For the geometric Lagrange weights  $\lambda_{p,j}^{(1/4)}$  on  $1, 4^{-1}, \dots, 4^{-p}$ , define the coefficientwise filter

$$\mathcal{R}_{p,n}(Q) := \sum_{j=0}^p \lambda_{p,j}^{(1/4)} D(4^{-(n+j)}Q). \quad (550)$$

**Theorem 8.28** (Formal Catalan–Gaussian quarter filter). *For every  $p, n \in \mathbb{N}_0$ , all coefficients of  $\mathcal{R}_{p,n}$  through degree  $p$  vanish. For  $m > p$ ,*

$$[Q^m]\mathcal{R}_{p,n} = \delta_m (-1)^p 4^{-\binom{p+1}{2}} \begin{bmatrix} m-1 \\ p \end{bmatrix}_{1/4} 4^{-nm}. \quad (551)$$

The degree- $(p+1)$  coefficient has the closed form

$$[Q^{p+1}]\mathcal{R}_{p,n} = \frac{C_p 2^{4p+2}}{9^{p+1} 4^{\binom{p+1}{2} + n(p+1)}}. \quad (552)$$

*Formal proof.* The series  $D$ , its coefficient formula, its quadratic equation, and its uniqueness are `quarterCatalanGermSeries`, `quarterCatalanGermSeries_coeff_succ`, `quarterCatalanGermSeries_equation`, and `existsUnique_quarterCatalanGermSeries` in `QuarterCatalanGerm.lean`. The filter is `quarterCatalanRichardsonFilter`. Cancellation, the complete residual formula, and the leading coefficient are `quarterCatalanRichardsonFilter_coeff_eq_zero_of_le`, `quarterCatalanRichardsonFilter_coeff_eq_qBinomial`, and `quarterCatalanRichardsonFilter_firstUncancelled_coeff` in `QuarterCatalanRichardson.lean`.

The same formal-germ module also connects this Catalan normalization to the distinguished dyadic quantile germ. Its exact scaled functional equation is `dyadicGermTwo_functionalEquation`; after rescaling the parameter by  $9/4$ , `rescale_dyadicGermTwo_eq_quadraticInverse` identifies it with `QuadraticInverse.inverse 4`. The reverse rescaling and coefficient formula are `dyadicGermTwo_eq_rescale_quadraticInverse` and `coeff_dyadicGermTwo_succ`. In particular the coefficient of degree  $m+1$  is

$$\left(\frac{4}{9}\right)^{m+1} (-4)^m C_m.$$

Every assertion here is coefficientwise in  $\mathbb{Q}[[Q]]$ . It neither asserts that the formal germ converges to a real finite-prefix inverse nor supplies a remainder estimate, sign theorem for a real extrapolation error, or analytic acceleration result. Those claims remain distinct from the proved Gaussian cancellation algebra.  $\square$

The full `QuarterCatalanGerm.lean` surface is two definitions and thirteen theorems. Besides the coefficient, equation, uniqueness, and injectivity declarations used above, its four dyadic-germ bridges are `dyadicGermTwo_functionalEquation`, `rescale_dyadicGermTwo_eq_quadraticInverse`, `dyadicGermTwo_eq_rescale_quadraticInverse`, and `coeff_dyadicGermTwo_succ`. They identify the distinguished rational dyadic germ, after parameter rescaling by  $9/4$ , with the unique Catalan inverse of  $X + 4X^2$ , and give every positive coefficient as  $(4/9)^{m+1}(-4)^m C_m$ . This remains formal power-series algebra.

**Theorem 8.29** (Actual Catalan jet at the quarter inverse anchor). *Let  $F : \text{BoundedFabius}$  satisfy  $h_F : \text{IsFabius } F$ , and write  $G(y) = \text{fabiusInv } F \, h_F \, y$ . For every  $n \in \mathbb{N}_0$ ,*

$$\frac{d^n}{dt^n} \left( G\left(\frac{5}{72} + t\right) - \frac{1}{4} \right) \Big|_{t=0} = n! [X^n] (\text{QuadraticInverse.inverse } 4). \quad (553)$$

Consequently, for every  $m \in \mathbb{N}_0$ ,

$$G^{(m+1)}\left(\frac{5}{72}\right) = (m+1)! (-4)^m C_m. \quad (554)$$

*Lean proof.* The separate `FabiusInverseQuarterJet.lean` has zero definitions and exactly two theorems. They are `iteratedDeriv_centeredFabiusInv_quarter_eq_quadraticInverse` and `iteratedDeriv_fabiusInv_five_seventy_two_succ` in `FabiusInverseQuarterJet.lean`. The proof transfers the exact smooth quarter jet of the forward Fabius function through Faà di Bruno and the local inverse identity, then applies the scalar convolution recurrence for the quadratic compositional inverse. This proves equality of jets only. It does not assert equality of the inverse with a quadratic analytic germ on a neighborhood, introduce a named nonzero flat-remainder theorem, or claim that the known nonanalytic flat defect vanishes.  $\square$

The polynomial in (486) has the complete rational root locus

$$\sum_{k=0}^n (-1)^k 2^{-\binom{k}{2}} \left[ \begin{matrix} n \\ k \end{matrix} \right]_{1/2} z^k = 0 \iff \exists j < n \text{ such that } z = 2^j, \quad z \in \mathbb{Q}. \quad (555)$$

This is `qBinomial_half_sum_eq_zero_iff`. The same statement is stable under scalar extension at embedded rational arguments. Let  $K$  be a field with a  $\mathbb{Q}$ -algebra structure, and write  $\iota : \mathbb{Q} \rightarrow K$  for its canonical map. Then, for every  $z \in \mathbb{Q}$ ,

$$\sum_{k=0}^n \iota \left( (-1)^k 2^{-\binom{k}{2}} \left[ \begin{matrix} n \\ k \end{matrix} \right]_{1/2} \right) (\iota z)^k = 0 \iff \exists j < n \text{ such that } \iota z = 2^j \text{ in } K. \quad (556)$$

This is `qBinomial_half_sum_algebraMap_eq_zero_iff` in `HalfQBinomial.lean`. The scalar statement classifies the embedded rational arguments  $\iota z$ ; it does not assert a root classification for arbitrary elements of  $K$ .

The multiplicity layer is exact as well. Over  $\mathbb{Q}$ , set

$$p_n(X) := \sum_{k=0}^n (-1)^k 2^{-\binom{k}{2}} \text{halfQBinomial } n \ k \ X^k.$$

For every  $j < n$ ,

$$\text{rootMultiplicity}_{p_n}(2^j) = 1. \quad (557)$$

This is the sole public theorem `halfQBinomial_sum_rootMultiplicity_two_pow` in the zero-definition/ one-theorem module `HalfQBinomialRootSimplicity.lean`. Its proof differentiates the finite q-Pochhammer product: at  $2^j$ , every product-rule term vanishes except the one in which the  $j$ -th factor was deleted, and all remaining factors are nonzero. Together with `halfQBinomial_sum_eq_zero_iff` and `gaussianBinomial_half_eq_halfQBinomial`, this completes both “simple” and “no others” in the canonical `cor:halfbase-root-locus`, which is therefore `Exact`. The theorem is deliberately over  $\mathbb{Q}$ , matching the coefficient and root-classification API; no arbitrary-base, arbitrary-field, or general cyclotomic simplicity is claimed, and the probabilistic `cor:qbinom-inversion-law` remains `Partial`.

The same Gaussian coefficients form a geometric Prouhet functional. Set

$$w_{n,k} = (-1)^k (1/2)^{\binom{k}{2}} \left[ \begin{matrix} n \\ k \end{matrix} \right]_{1/2}, \quad \xi_k = -2^k.$$

For every rational polynomial  $p$  with  $\deg p \leq n$ ,

$$\sum_{k=0}^n w_{n,k} p(\xi_k) = [X^n] p \ 2^{\binom{n+1}{2}} (1/2; 1/2)_n = [X^n] p \ \mathfrak{m}_n. \quad (558)$$

Thus the functional annihilates every polynomial of degree below  $n$ . At  $n = 0$ , the degree-valued extractor covers all constant polynomials; the strict-degree cancellation statement admits exactly the zero polynomial. The two forms of (558) are `halfQBinomial_negativeDyadic_polynomial_1_sum_eq_coeff` and `halfQBinomial_negativeDyadic_polynomial_sum_eq_mersenne`; strict-degree cancellation is `halfQBinomial_negativeDyadic_polynomial_sum_eq_zero_of_degree_lt`, all in `HalfQBinomial.lean`.

The inner cancellation in the dyadic formula is the exact Thue–Morse block identity

$$\sum_{r=0}^{2^k-1} (-1)^{\text{wt}_2(r)} z^r = \prod_{j=0}^{k-1} (1 - z^{2^j}), \quad (559)$$

formalized as `thueMorseBlockPolynomial_eq_product`. Its affine power-sum form annihilates every degree below  $k$ , while its first surviving moment is  $(-1)^k 2^{\binom{k}{2}} k!$ ; see `thueMorse_affine_power_sum_of_1e` and `thueMorse_affine_power_sum_self`.

For comparison with the moment recurrence, put

$$a_n = \frac{d_n}{n!}, \quad \mathcal{A}(X) = \sum_{n \geq 0} a_n X^n, \quad E(X) = \frac{e^X - 1}{X}, \quad (560)$$

with the removable value  $E(0) = 1$ . The refinement is

$$\mathcal{A}(2X) = E(X) \mathcal{A}(X), \quad (561)$$

and its coefficient recurrence is

$$(2^n - 1)a_n = \sum_{j=0}^{n-1} \frac{a_j}{(n-j+1)!} \quad (n \geq 1), \quad a_0 = 1. \quad (562)$$

The functional identity follows from `proposition_two_formula` and `complexGeneratingFunction_eq_series_formula` in `AnalyticMoments.lean`; the coefficient recurrence is `halfMoment_original_recurrence` in `BernoulliRecurrences.lean`. These are the formal-series bridge used by the specialized dyadic theorem.

For  $m, k, d \in \mathbb{N}_0$  and  $Q \in \mathbb{C}$ , write

$$\sigma_{m;k,d}(Q) = \sum_{r=0}^{m2^k-1} (-1)^{\text{wt}_2(r)} (r - m2^k + Q)^d. \quad (563)$$

The exact result is the following finite identity.

**Theorem 8.30** (Translated global dyadic formula). *For all  $m, n \in \mathbb{N}_0$  and  $Q \in \mathbb{C}$ ,*

$$\begin{aligned} F_{\text{gl}}(m/2^n) &= \frac{1}{2^{n^2} (1/2; 1/2)_n} \sum_{k=0}^n \frac{[n]_{1/2}}{4^{\binom{k}{2}} (n+k)!} \\ &\quad \times \sum_{r=0}^{m2^k-1} (-1)^{\text{wt}_2(r)} (r - m2^k + Q)^{n+k}. \end{aligned} \quad (564)$$

*The real number on the left is embedded in  $\mathbb{C}$ . The entire right-hand side is independent of the common translation  $Q$ . When  $m \leq 2^n$ , the left side is the bounded value  $F(m/2^n)$ .*

*Formal proof.* The complete normalized finite expression is packaged as a polynomial in the translation and proved equal to the constant centered formula by `qBinomialThueMorseDyadicTranslatedFormulaPolynomial_eq_const`. Its identification with the signed dyadic value over  $\mathbb{C}$  is `globalFabius_dyadic_eq_qBinomialThueMorseDyadic_translated_sum_complex`; the rational core and real, complex, and Gaussian-rational specializations are in `FabiusQBinomialFormula.lean` and `FabiusDyadicQBinomialScalar.lean`.  $\square$

The rational specialization is also independent of how the dyadic argument is written. If

$$\frac{m_1}{2^{n_1}} = \frac{m_2}{2^{n_2}},$$

then the two translated rational formulas agree even when their common translations  $Q_1, Q_2 \in \mathbb{Q}$  differ. This is `qBinomialThueMorseDyadicTranslatedFormula_eq_of_rat_eq`; the one-step refinement  $(m, n) \mapsto (2m, n+1)$  is `qBinomialThueMorseDyadicTranslatedFormula_refine`.



Over every field over  $\mathbb{Q}$ , both conclusions persist even when the scalar translations and the dyadic presentations are changed simultaneously; these stronger wrappers are `qBinomialThueMorseDyadicTranslatedFormulaIn_eq_of_rat_eq` and `qBinomialThueMorseDyadicTranslatedFormulaIn_refine` in `FabiusDyadicQBinomialScalar.lean`.

Binary complementation gives the raw inverse-dyadic form

$$F_{\text{gl}}(2^{-n}) = \frac{1}{(-2)^{n^2}(1/2; 1/2)_n} \sum_{k=0}^n \frac{\begin{bmatrix} n \\ k \end{bmatrix}_{1/2}}{4^{\binom{k}{2}}(n+k)!} \sum_{r=0}^{2^k-1} (-1)^{\text{wt}_2(r)} (r+Q)^{n+k}. \quad (565)$$

Its translated numerator is constant by `qBinomialThueMorseRawTranslatedNumeratorPolynomial_eq_const`. The algebraic statement is in fact valid over every field  $K$  equipped with a  $\mathbb{Q}$ -algebra structure and every scalar translation  $Q \in K$ : the fully expanded right-hand side equals the canonical inverse-dyadic rational value after its scalar embedding by `fabiusAtInverseTwoPow_cast_eq_qBinomialThueMorse_rawTranslated_sum`. Translation independence and the intermediate value identification are `qBinomialThueMorseRawTranslatedFormulaIn_eq_centered` and `qBinomialThueMorseRawTranslatedFormulaIn_eq_fabiusAtInverseTwoPow`. For every scalar type  $K$  carrying Mathlib’s `RCLike` structure, the corresponding theorem for an arbitrary bounded Fabius solution is `fabiusFunction_inverse_two_pow_eq_qBinomialThueMorseRawTranslatedFormulaIn`. All five declarations are in `FabiusRawQBinomialScalar.lean`; the final one uses the `RCLike` hypothesis only to embed the real Fabius value. Variable-base  $q$ -interpolation formulas, barycentric identities, numerical cancellation studies, and complexity comparisons are maintained in the consolidated research-frontier volume rather than used as theorems here.

## 8.8 The prefactor is the order of $\text{GL}_n(\mathbb{F}_2)$

The constant  $2^{n^2}(1/2; 1/2)_n$  standing in front of every form of the dyadic identity above is not merely a normalization: it is the order of the general linear group over the two-element field. Both halves of the identity can in fact be freed of the base  $1/2$  altogether, and what is left counts subspaces.

The first step is a falling-product identity with no hypotheses at all.

**Lemma 8.31** (Falling product of powers). *In every commutative ring and for every  $q$ ,*

$$\prod_{i < n} (q^n - q^i) = q^{\binom{n}{2}} \prod_{j < n} (q^{j+1} - 1). \quad (566)$$

This is `prod_pow_sub_pow_eq_pow_choose_two_mul` in `GLOrderProduct.lean`. No invertibility or nonvanishing assumption occurs, so (566) holds at roots of unity, in positive characteristic and in the presence of zero divisors.

Applied at  $q = 2$ , and combined with the Mersenne normalization `two_pow_nat_sq_mul_halfQPochhammer`, it identifies the prefactor:

$$2^{n^2}(1/2; 1/2)_n = \prod_{i < n} (2^n - 2^i), \quad (567)$$

which is `two_pow_nat_sq_mul_qPochhammer_half_eq_prod` in `FabiusDyadicGaussianForm.lean`. The right-hand side of (567) is the number of ordered bases of  $\mathbb{F}_2^n$ .

**Theorem 8.32** (The dyadic prefactor is a group order). *For every  $n \in \mathbb{N}_0$ ,*

$$2^{n^2}(1/2; 1/2)_n = |\text{GL}_n(\mathbb{F}_2)|. \quad (568)$$

More generally, over a finite field  $K$  with  $Q = |K|$  elements,

$$|\text{GL}_n(K)| = Q^{\binom{n}{2}} \prod_{j < n} (Q^{j+1} - 1) = Q^{n^2} (1/Q; 1/Q)_n. \quad (569)$$



The three statements are `two_pow_nat_sq_mul_qPochhammer_half_eq_card_GL`, `cast_card_generalLinearGroup_eq_pow_choose_mul_prod` and `cast_card_generalLinearGroup_eq_pow_sq_mul_qPochhammer` in `FabiusGeneralLinearDenominator.lean`. The exponent bookkeeping in (569) is the square split  $\binom{n}{2} + \binom{n+1}{2} = n^2$ , which is `choose_square_split` in `Arithmetic.lean`. So the denominators of the exact dyadic Fabius values are the group orders

$$1, 1, 6, 168, 20160, 9999360, 20158709760, \dots$$

*Remark 8.33* (What is and is not proved here). The order of  $GL_n$  is `Mathlib's Matrix.card_GL_field` and is not reproved anywhere in this development; `card_generalLinearGroup_eq_prod_range` of `RankMatrixCount.lean` only reindexes it from `Fin r` to `Finset.range r`, so it is a bridge and not a source. What is established here is the cast to  $\mathbb{Q}$  – the ambient statement lives in  $\mathbb{N}_0$  with truncated subtraction – together with the identification of the resulting product with the  $q$ -Pochhammer prefactor.

The second step removes the base  $1/2$  from the summand. Gaussian reciprocity, `gaussianBinomial_reciprocity`, converts the quotient  $q$ -binomial at  $1/2$  into the polynomial Gaussian binomial at 2 at the cost of a monomial  $2^{k(n-k)}$ , and `four_pow_choose_two` rewrites  $4^{\binom{k}{2}}$  as  $2^{k(k-1)}$ . The two exponents merge, because for  $k \leq n$

$$k(n-k) + k(k-1) + k = kn \quad (570)$$

in  $\mathbb{N}_0$  with truncated subtraction (`exponent_collapse`). Consequently

$$\frac{\begin{bmatrix} n \\ k \end{bmatrix}_{1/2}}{4^{\binom{k}{2}}} = \frac{\begin{bmatrix} n \\ k \end{bmatrix}_2 \cdot 2^k}{2^{kn}},$$

in which  $\begin{bmatrix} n \\ k \end{bmatrix}_2$  is an *integer*: the number of  $k$ -dimensional subspaces of  $\mathbb{F}_2^n$ .

**Theorem 8.34** (Subspace form of the dyadic identity). *For every  $n \in \mathbb{N}_0$ ,*

$$F(2^{-n}) = \frac{1}{|GL_n(\mathbb{F}_2)|} \sum_{k=0}^n \frac{\begin{bmatrix} n \\ k \end{bmatrix}_2 \cdot 2^k}{2^{kn} (n+k)!} \sum_{r < 2^k} \varepsilon_r (r - 2^k)^{n+k}. \quad (571)$$

This is `fabiusAtInverseTwoPow_eq_card_GL_sum` in `FabiusGeneralLinearDenominator.lean`, with the product form `fabiusAtInverseTwoPow_eq_gaussian_sum` of `FabiusDyadicGaussianForm.lean` and the real-valued transfers `fabius_inverse_two_pow_eq_card_GL_inv_mul` and `fabius_inverse_two_pow_eq_gaussian_div_prod` alongside.

Neither side of (571) mentions the base  $1/2$ , and no  $q$ -Pochhammer quotient, no  $4^{\binom{k}{2}}$  and no  $2^{n^2}$  survives. Every constant on the right counts something over  $\mathbb{F}_2$ : the weight  $\begin{bmatrix} n \\ k \end{bmatrix}_2$  enumerates the  $k$ -dimensional subspaces of  $\mathbb{F}_2^n$ , and the normalizing constant is the order of the automorphism group of that space. The exact dyadic values of the Fabius function are therefore a signed, factorially weighted average over the subspace lattice of  $\mathbb{F}_2^n$ .

*Remark 8.35* (The Thue–Morse sign inside the  $q$ -Pochhammer). The appearance of a  $q$ -Pochhammer symbol beside a Thue–Morse block sum in (571) is not a coincidence of normalizations. Over any commutative ring and with no hypothesis on  $z$  or  $q$ ,

$$\sum_{n < 2^N} \varepsilon_n z^{\text{wt}_2(n)} q^{\sigma(n)} = (z; q)_N, \quad (572)$$

where  $\sigma(n)$  is the sum of the positions of the one bits of  $n$ : a finite  $q$ -Pochhammer symbol is a Thue–Morse alternating sum, bigraded by the two binary statistics. This is `sum_thueMorseSign_mul_pow_bigraded_eq_finiteQPochhammerIn` in `ThueMorseQPochhammer.lean`. Combining it with (569) expresses the group order itself as an alternating digit sum,  $|GL_n(K)| = \sum_{m < 2^n} \varepsilon_m Q^{n^2 - \text{wt}_2(m) - \sigma(m)}$  (`cast_card_generalLinearGroup_eq_thueMorse_sum` in `GeneralLinearThueMorseSum.lean`), the exponent being non-negative because  $\text{wt}_2(m) + \sigma(m) \leq \binom{n+1}{2} \leq n^2$ .

## 9 The global binary-reduction telescope

The Taylor-block recursion also applies to the signed global extension  $F_{\text{gl}}$ . It can therefore be iterated along the binary expansion of an arbitrary nonnegative real number, rather than only along the finite expansion of a dyadic rational. The reductions need not terminate, but their residuals tend geometrically to zero.

Define the finite reduction polynomial

$$\mathcal{T}_m(y) = \sum_{j=0}^m 2^{\binom{j+1}{2} - \binom{m-j}{2}} \frac{d_{m-j}}{(m-j)!} \frac{y^j}{j!}. \quad (573)$$

All powers of 2 here are integer powers. In particular, negative exponents are allowed, and  $\mathcal{T}_0(y) = 1$ . The signed Taylor-block identity is

$$F_{\text{gl}}(2^{-m} + y) = -F_{\text{gl}}(y) + \mathcal{T}_m(y), \quad 0 \leq y < 2^{-m}. \quad (574)$$

Fix  $x \geq 0$ . At scale  $m$ , put

$$p_m = \lfloor 2^m x \rfloor, \quad y_m = x - \frac{p_m}{2^m}, \quad 0 \leq y_m < 2^{-m}. \quad (575)$$

The previous prefix requires a genuine half-scale value at  $m = 0$ :

$$p_{m-1}^* = \begin{cases} \lfloor x/2 \rfloor, & m = 0, \\ p_{m-1}, & m \geq 1. \end{cases} \quad (576)$$

Uniformly,  $p_{m-1}^* = \lfloor p_m/2 \rfloor$ . Thus  $p_m = 2p_{m-1}^* + \delta_m$  with  $\delta_m \in \{0, 1\}$ . Define

$$\alpha_m(x) = (-1)^{\text{wt}_2(p_m)} (2p_{m-1}^* - p_m), \quad R_m(x) = (-1)^{\text{wt}_2(p_m)} F_{\text{gl}}(y_m). \quad (577)$$

The coefficient  $\alpha_m$  is zero or a sign, and it is nonzero precisely when the newly exposed binary digit is 1.

**Lemma 9.1** (One-step residual identity). *For  $m \geq 1$ ,*

$$\alpha_m(x) \mathcal{T}_m(y_m) = R_{m-1}(x) - R_m(x). \quad (578)$$

*Proof.* If  $p_m = 2p_{m-1}$ , then  $y_m = y_{m-1}$ , the Thue–Morse signs agree, and  $\alpha_m = 0$ , so both sides vanish. If  $p_m = 2p_{m-1} + 1$ , then

$$y_{m-1} = 2^{-m} + y_m, \quad (-1)^{\text{wt}_2(p_m)} = -(-1)^{\text{wt}_2(p_{m-1})}, \quad \alpha_m = (-1)^{\text{wt}_2(p_{m-1})}.$$

Apply (574) to  $y_{m-1}$  and multiply by the old prefix sign; the two copies of  $F_{\text{gl}}(y_m)$  combine exactly into  $R_{m-1} - R_m$ .  $\square$

The scale-zero identity is separate and retains the parity of the integer block:

$$F_{\text{gl}}(x) = \alpha_0(x) \mathcal{T}_0(y_0) + R_0(x). \quad (579)$$

Indeed, write  $p_0 = 2p_{-1}^* + \delta_0$  and use the two-unit translation law (12). When  $\delta_0 = 0$ , the polynomial term vanishes. When  $\delta_0 = 1$ , the identity  $F_{\text{gl}}(1 + y_0) = -F_{\text{gl}}(y_0) + 1$  supplies it.

**Theorem 9.2** (Exact finite binary telescope). *For  $x \geq 0$  and  $N \in \mathbb{N}_0$ ,*

$$F_{\text{gl}}(x) = \sum_{m=0}^N \alpha_m(x) \mathcal{T}_m(y_m(x)) + R_N(x). \quad (580)$$

*Proof.* Start with (579). Summing (578) for  $m = 1, \dots, N$  telescopes all intermediate residuals. The case  $N = 0$  is exactly the scale-zero identity.  $\square$

For  $N \geq 1$ ,  $0 \leq y_N < 2^{-N} \leq 1/2$ . On this interval  $0 \leq F_{\text{gl}}(y) = F(y) \leq 2y$ , so

$$|R_N(x)| \leq 2^{1-N}. \quad (581)$$

The estimate is independent of  $x$ . Therefore, if  $B_N(x)$  denotes the finite sum in (580), then for every  $x \in \mathbb{R}$  and  $N \geq 1$ ,

$$|B_N(x) - F_{\text{gl}}(x)| \leq 2^{1-N}. \quad (582)$$

This is `norm_globalBinaryReductionSum_sub_extendedFabius_le`. Lean’s natural-floor convention extends every  $B_N$  by zero on  $x \leq 0$ , where  $F_{\text{gl}}$  also vanishes, and `globalBinaryReductionSum_tendstoUniformly_extendedFabius` proves that the finite telescopes converge uniformly on the whole line. The one-step identity then gives, for  $m \geq 2$ ,

$$|\alpha_m(x)\mathcal{T}_m(y_m)| \leq |R_{m-1}| + |R_m| \leq 4 \cdot 2^{-(m-1)}. \quad (583)$$

Thus the limiting series is absolutely, not merely conditionally, convergent.

**Theorem 9.3** (Absolutely convergent global binary series). *For every  $x \geq 0$ ,*

$$F_{\text{gl}}(x) = \sum_{m=0}^{\infty} \alpha_m(x)\mathcal{T}_m(y_m(x)). \quad (584)$$

On  $0 \leq x \leq 1$ , the sum is equivalently  $F(x)$ .

Lean’s natural-floor definitions make the telescope genuinely all-real: when  $x \leq 0$ , every binary prefix, summand, residual, and  $F_{\text{gl}}(x)$  vanishes termwise. Thus the finite identity and the  $m = 0$  infinite series hold for every  $x \in \mathbb{R}$ , as `extendedFabius_eq_globalBinaryReductionSum_add_remainder_all` and `extendedFabius_eq_tsum_globalBinaryReductionSummand_all` in `FabiusBinaryReductionSeries.lean`. The fully expanded parity-power version has the same all-real scope by `globalFabius_eq_tsum_fabiusParityPower_literal_all` in `FabiusParityPowerSeries.lean`. The displayed floor formulas above remain ordinary nonnegative-floor formulas on their stated domain.

The index  $m = 0$  is indispensable globally. For  $0 \leq x < 1$  it vanishes, so a series starting at  $m = 1$  happens to be valid there. At  $x = 1$ , however, the zeroth term is 1 and all later coefficients vanish. More generally it is the zeroth term that distinguishes even and odd integer blocks of the signed extension.

### 9.1 The corrected parity-power presentation

The same summands have a more literal presentation in terms of prefix parity and inverse dyadic values. Define, using integer exponents,

$$\kappa_j = \frac{((j : \mathbb{Z}) + 2)((j : \mathbb{Z}) - 1)}{2} = \binom{j+1}{2} - 1. \quad (585)$$

The integer interpretation matters:  $\kappa_0 = -1$ , hence  $2^{\kappa_0} = 1/2$ ; natural subtraction would silently give the wrong term. Put

$$\mathcal{I}_m(y) = \sum_{j=0}^m 2^{\kappa_j} F_{\text{gl}}(2^{(j:\mathbb{Z})-(m:\mathbb{Z})}) \frac{y^j}{j!}. \quad (586)$$

**Lemma 9.4.** *For every  $m$  and  $y$ ,*

$$\mathcal{I}_m(y) = \frac{1}{2} \mathcal{T}_m(y). \quad (587)$$

*Proof.* For  $j \leq m$ , set  $r = m - j$ . The inverse-dyadic identity (437) gives

$$F_{\text{gl}}(2^{j-m}) = \frac{d_{m-j}}{(m-j)! 2^{\binom{m-j}{2}}}.$$

Together with  $\kappa_j = \binom{j+1}{2} - 1$ , this identifies each summand with one half of the corresponding summand in (573).  $\square$

Let  $\tau(a) = \text{wt}_2(a) \bmod 2$ . Since

$$\frac{(-1)^a - 1}{2} = 2 \left\lfloor \frac{a}{2} \right\rfloor - a,$$

the literal summand

$$\begin{aligned} \Lambda_m(x) &:= ((-1)^{\lfloor 2^m x \rfloor} - 1)(-1)^{\tau(\lfloor 2^m x \rfloor)} \\ &\quad \times \sum_{j=0}^m 2^{((j+2)(j-1))/2} F_{\text{gl}}(2^{j-m}) \frac{(x - \lfloor 2^m x \rfloor / 2^m)^j}{j!} \end{aligned} \quad (588)$$

is exactly  $\alpha_m(x) \mathcal{T}_m(y_m)$ . Both exponents  $((j+2)(j-1))/2$  and  $j-m$  in this display are integer exponents, not truncated natural subtractions.

*Warning 9.5.* In (563) and (564), the summation indices are cast into the ambient characteristic-zero field before the subtraction in  $r - m2^k + Q$  is performed. This is the typing used by the formal theorem; the displayed subtraction is therefore field subtraction, never truncated subtraction on natural numbers.

**Theorem 9.6** (Corrected parity-power series). *For every  $x \geq 0$ ,*

$$F_{\text{gl}}(x) = \sum_{m=0}^{\infty} \Lambda_m(x), \quad (589)$$

*and the series converges absolutely. On the half-open unit interval  $0 \leq x < 1$ , the zeroth summand vanishes and one may start at  $m = 1$ .*

*Proof.* The elementary parity identity and (587) prove  $\Lambda_m = \alpha_m \mathcal{T}_m(y_m)$  term by term. The claim follows from (584). The half-open corollary follows from  $\lfloor x \rfloor = 0$ ; the closed endpoint  $x = 1$  requires the restored zeroth term.  $\square$

## 9.2 The global complex $q$ -binomial series

The finite constant-polynomial identity from section 8.7 can be inserted at every scale of the binary telescope. For  $Q \in \mathbb{C}$ , define

$$\mathcal{N}_n^{(Q)} = \sum_{k=0}^n \frac{\begin{bmatrix} n \\ k \end{bmatrix}_{1/2}}{4^{\binom{k}{2}} (n+k)!} \sum_{r=0}^{2^k-1} (-1)^{\text{wt}_2(r)} (r - 2^k + Q)^{n+k}. \quad (590)$$

Translation invariance and the inverse-dyadic normalization give

$$\frac{\mathcal{N}_n^{(Q)}}{2^{\binom{n}{2}} (1/2; 1/2)_n} = \frac{2^n d_n}{n!}. \quad (591)$$

The normalization differs from the  $2^{n^2}$  in (564): here it produces a Taylor coefficient rather than a function value.

For  $m \in \mathbb{N}_0$ , put

$$\mathcal{Q}_m^{(Q)}(y) = 2^{-\binom{m+1}{2}} \sum_{n=0}^m \frac{(2^{m+1}y)^{m-n}}{(m-n)!} \frac{\mathcal{N}_n^{(Q)}}{2^{\binom{n}{2}}(1/2; 1/2)_n}. \quad (592)$$

**Theorem 9.7** ( $q$ -binomial Taylor identity). *For every  $m$ , every  $y \in \mathbb{R}$ , and every  $Q \in \mathbb{C}$ ,*

$$\mathcal{Q}_m^{(Q)}(y) = \mathcal{T}_m(y), \quad (593)$$

where the real polynomial on the right is embedded in  $\mathbb{C}$ . In particular, each finite summand is pointwise independent of  $Q$  before any infinite sum is taken.

*Proof.* Insert (591) into (592) and set  $j = m - n$ . Collecting the powers of 2 gives

$$2^{-\binom{m+1}{2}} 2^{(m+1)j} 2^{m-j} = 2^{\binom{j+1}{2} - \binom{m-j}{2}}.$$

The remaining factorials and moments are exactly those in (573).  $\square$

**Theorem 9.8** (Global complex  $q$ -binomial series). *For fixed  $Q \in \mathbb{C}$  and every  $x \geq 0$ ,*

$$F_{\text{gl}}(x) = \sum_{m=0}^{\infty} \alpha_m(x) \mathcal{Q}_m^{(Q)}(y_m(x)), \quad (594)$$

with the left side embedded in  $\mathbb{C}$ . The series converges absolutely.

The displayed floor presentation is stated on the nonnegative half-line. The literal Lean summand has a stronger all-real scope: for every  $x \in \mathbb{R}$ , its series is absolutely summable and sums to the signed global extension, while for  $x \leq 0$  every term and the target vanish. These are `summable_norm_qBinomialFabiusGlobalSummand_all`, `hasSum_qBinomialFabiusGlobalSummand_all`, `extendedFabius_eq_tsum_qBinomialFabiusGlobalSummand_all`, `globalFabius_eq_tsum_qBinomialFabiusGlobalSummand_all`, and `qBinomialFabiusGlobalSummand_eq_zero_of_nonpos` in `FabiusGlobalQBinomialSeries.lean`. The older nonnegative signatures are retained as compatibility wrappers.

This is not a new infinite basic-hypergeometric identity. The outer sum is the analytic binary telescope, while at each scale a finite  $q$ -binomial identity presents the same Taylor polynomial. Expanding one summand makes the architecture explicit:

$$\begin{aligned} & \alpha_m(x) 2^{-\binom{m+1}{2}} \sum_{n=0}^m \frac{(2^{m+1}x - 2p_m(x))^{m-n}}{(m-n)!} \frac{1}{2^{\binom{n}{2}}(1/2; 1/2)_n} \\ & \times \sum_{k=0}^n \frac{\begin{bmatrix} n \\ k \end{bmatrix}_{1/2}}{4^{\binom{k}{2}}(n+k)!} \sum_{r=0}^{2^k-1} (-1)^{\text{wt}_2(r)} (r - 2^k + Q)^{n+k}. \end{aligned} \quad (595)$$

Convergence comes entirely from the binary-tail majorant (583); the special-function manipulations are finite and algebraic.

## 10 A complex $q$ -binomial discrete limit

There is a second way in which the same Gaussian row produces  $F_{\text{gl}}$ . It is a genuine limit of finite shifted splines rather than a sequence of partial sums of (594). Reuse the safe midpoint cutoff  $N_p(x)$  from (119). For  $x \geq 0$  it is equivalent to the inclusive source range

$$0 \leq r \leq \left\lfloor 2^p x - \frac{1}{2} \right\rfloor, \quad (596)$$

including the empty case where the upper endpoint is  $-1$ .

For fixed  $Q \in \mathbb{C}$  and  $x \in \mathbb{R}$ , define

$$\begin{aligned} \mathcal{D}_n^{(Q)}(x) &= \frac{1}{2^{n^2} (1/2; 1/2)_n} \sum_{k=0}^n \frac{[n]_{1/2}}{4^{\binom{k}{2}} (n+k)!} \\ &\quad \times \sum_{r=0}^{N_{n+k}(x)-1} (-1)^{\text{wt}_2(r)} (r - 2^{n+k}x + Q)^{n+k}. \end{aligned} \quad (597)$$

**Theorem 10.1** (Complex discrete-limit formula). *For every fixed  $Q \in \mathbb{C}$  and every  $x \in \mathbb{R}$ ,*

$$\boxed{\mathcal{D}_n^{(Q)}(x) \longrightarrow F_{\text{gl}}(x)} \quad (598)$$

in  $\mathbb{C}$ . If  $0 \leq x \leq 1$ , the limit is equivalently  $F(x)$ .

*Proof.* If  $x \leq 0$ , every safe cutoff  $N_p(x)$  is zero. Hence every shifted spline and every approximant (597) is zero, as is  $F_{\text{gl}}(x)$ . It remains to treat  $x \geq 0$ .

For a degree  $p$  and complex shift  $Q$ , introduce the finite shifted spline

$$S_p^{(Q)}(x) = \frac{(-1)^p}{2^{\binom{p}{2}} p!} \sum_{r=0}^{N_p(x)-1} (-1)^{\text{wt}_2(r)} (r - 2^p x + Q)^p. \quad (599)$$

At  $Q = 1/2$  this is exactly the centered spline (120), so by (127),

$$S_p^{(1/2)}(x) \longrightarrow F_{\text{gl}}(x).$$

**Lemma 10.2** (Fixed-cutoff Taylor expansion of a complex shift). *Read (120) at  $p = 0$  as*

$$S_0(x) = \sum_{r=0}^{N_0(x)-1} \varepsilon_r, \quad (600)$$

so that a centered branch  $S_d$  is defined for every integer  $d \geq 0$ . Fix  $p \geq 1$  and  $x \geq 0$ , and abbreviate  $M = N_p(x)$ . For every  $d$  with  $0 \leq d \leq p$ , the degree- $d$  branch carried by that one fixed cutoff satisfies

$$\frac{1}{2^{\binom{d}{2}} d!} \sum_{r=0}^{M-1} \varepsilon_r \left( 2^p x - \frac{1}{2} - r \right)^d = S_d(2^{p-d}x), \quad (601)$$

and in particular  $|S_d(2^{p-d}x)| \leq 1$ . Moreover, for every  $Q \in \mathbb{C}$ ,

$$S_p^{(Q)}(x) - S_p^{(1/2)}(x) = \sum_{d=0}^{p-1} \frac{2^{\binom{d}{2}}}{2^{\binom{p}{2}} (p-d)!} \left( \frac{1}{2} - Q \right)^{p-d} S_d(2^{p-d}x). \quad (602)$$

*Proof.* Write the shifted spline in its descending orientation. Since  $(-1)^p (r - 2^p x + Q)^p = (2^p x - Q - r)^p$ , definition (599) reads

$$S_p^{(Q)}(x) = \frac{1}{2^{\binom{p}{2}} p!} \sum_{r=0}^{M-1} \varepsilon_r (2^p x - Q - r)^p, \quad M = N_p(x), \quad (603)$$

and at  $Q = 1/2$  this is (120). The cutoff  $M$  depends on  $p$  and  $x$  only. It is unaffected by  $Q$ , and it is held fixed in what follows while the degree of the summand varies; that is the whole point of (601).

We prove (601). The dilation  $x \mapsto 2^{p-d}x$  compensates exactly for the drop in degree, in the sense that

$$N_d(2^{p-d}x) = \max\left(0, \left\lfloor 2^p x + \frac{1}{2} \right\rfloor\right) = N_p(x) = M \quad (604)$$

by (119). If  $M = 0$ , both sides of (601) are empty sums and there is nothing to prove. Assume  $M \geq 1$ . Every index in the range satisfies

$$r \leq M - 1 = \left\lfloor 2^p x + \frac{1}{2} \right\rfloor - 1 \leq 2^p x - \frac{1}{2},$$

so every base  $2^p x - \frac{1}{2} - r$  occurring in (601) is nonnegative and the truncation in (121) is inactive. For  $d \geq 1$ , formula (121) evaluated at the point  $2^{p-d}x$ , whose own cutoff is  $M$  by (604), is therefore literally the left side of (601). For  $d = 0$  both sides equal  $\sum_{r=0}^{M-1} \varepsilon_r$ , by (600) and (604).

For the uniform bound, the rescaled argument  $2^{p-d}x$  is nonnegative. The all-degree declaration `abs_fabiusUniformSpline_1e_one` therefore gives the claim directly, including  $d = 0$ .

Now expand. Put  $w = 2^p x - \frac{1}{2}$  for the descending argument at  $Q = 1/2$  and  $t = \frac{1}{2} - Q$  for its increment, so that  $2^p x - Q = w + t$ . The binomial theorem gives

$$(w + t - r)^p = \sum_{d=0}^p \binom{p}{d} t^{p-d} (w - r)^d.$$

Multiply by  $\varepsilon_r / (2^{\binom{p}{2}} p!)$ , sum over  $0 \leq r \leq M - 1$ , and interchange the two finite sums. Using (603) on the left and (601) on each inner sum,

$$\begin{aligned} S_p^{(Q)}(x) &= \sum_{d=0}^p \frac{t^{p-d}}{2^{\binom{p}{2}} p!} \binom{p}{d} \sum_{r=0}^{M-1} \varepsilon_r (w - r)^d \\ &= \sum_{d=0}^p \frac{2^{\binom{d}{2}}}{2^{\binom{p}{2}} (p-d)!} t^{p-d} S_d(2^{p-d}x), \end{aligned} \quad (605)$$

because  $\binom{p}{d}/p! = 1/(d!(p-d)!)$  and the inner sum is  $2^{\binom{d}{2}} d! S_d(2^{p-d}x)$ . The term  $d = p$  is  $S_p(x) = S_p^{(1/2)}(x)$ ; subtracting it leaves (602).  $\square$

*Remark 10.3* (Which convention makes the sign disappear). The expansion (602) is free of alternating signs only because the increment is measured in the *descending* variable  $2^p x - Q$  of (603), that is, because the coefficient carries  $(\frac{1}{2} - Q)^{p-d}$  and not  $(Q - \frac{1}{2})^{p-d}$ . If one instead measures the shift by the ascending increment  $q = Q - \frac{1}{2}$ , which is the natural reading when the branch is written as a function of the ascending argument  $r - 2^p x + Q$  of (599), then the identity acquires exactly the factor  $(-1)^{p+d}$ :

$$S_p^{(Q)}(x) - S_p^{(1/2)}(x) = \sum_{d=0}^{p-1} (-1)^{p+d} \frac{2^{\binom{d}{2}}}{2^{\binom{p}{2}} (p-d)!} q^{p-d} S_d(2^{p-d}x), \quad q = Q - \frac{1}{2}. \quad (606)$$

The two displays are the same statement, since  $(-1)^{p+d} = (-1)^{p-d}$  and therefore  $(-1)^{p+d} q^{p-d} = (\frac{1}{2} - Q)^{p-d}$ . It is worth being explicit about where the sign does *not* live: it is not a matter of renormalizing the lower branches. The alternating prefactor  $(-1)^d$  of (120) is cancelled exactly by the  $(-1)^d$  produced when the ascending argument is reversed, so the branch values  $S_d(2^{p-d}x)$  appearing in (602) and in (606) are the same numbers. The sign belongs to the increment alone, and the convention that removes it is the descending one.



*Remark 10.4* (Formal bound). Only the exponential bound (607) is asserted here; it is `norm_fabiusComplexShiftSpline_sub_center_le_half_pow_mul_exp` in `FabiusComplexShiftSpline.lean`. A proposed sharper constant is recorded in the research-frontier gap register pending a corrected named theorem.

Expanding a fixed cutoff branch about  $Q = 1/2$  gives an exact finite Taylor combination of lower-degree centered branches. The normalized Thue–Morse bounds then give, for  $p \geq 1$  and  $x \geq 0$ ,

$$\left| S_p^{(Q)}(x) - S_p^{(1/2)}(x) \right| \leq 2^{-(p-1)} \exp(|Q - \tfrac{1}{2}|). \quad (607)$$

The nonpositive half-line is handled separately by exact vanishing, and `fabiusComplexShiftSpline_tendsto_globalFabius_all` proves that every fixed complex shift has the same pointwise spline limit for every real  $x$ .

Now reindex the outer sum in (597) by  $j = n - k$ . Direct collection of signs, powers, and factorials gives the exact finite identity

$$\mathcal{D}_n^{(Q)}(x) = \sum_{j=0}^n \omega_{n,j} S_{2n-j}^{(Q)}(x), \quad (608)$$

where

$$\omega_{n,j} = \frac{(-1)^j \begin{bmatrix} n \\ j \end{bmatrix}_{1/2} 2^{-(j+1)}}{(1/2; 1/2)_n}. \quad (609)$$

The finite  $q$ -binomial theorem gives the row normalization

$$\sum_{j=0}^n \omega_{n,j} = 1. \quad (610)$$

The same theorem identifies the entire row rather than only its value at one.

**Proposition 10.5** (Toeplitz row polynomial and module-valued  $q$ -Richardson cancellation). *For  $z \in \mathbb{Q}$ , put  $W_n(z) = \sum_{j=0}^n \omega_{n,j} z^j$ . Then*

$$W_n(z) = \frac{(z/2; 1/2)_n}{(1/2; 1/2)_n}. \quad (611)$$

*Consequently its rational zeros are exactly*

$$W_n(z) = 0 \iff z \in \{2, 2^2, \dots, 2^n\}. \quad (612)$$

*Moreover, let  $M$  be any additive commutative monoid carrying a  $\mathbb{Q}$ -module structure. If an  $M$ -valued sequence has the finite geometric-error model  $u_p = L + \sum_{r=1}^n 2^{-rp} \cdot a_r$ , where  $\cdot$  denotes scalar multiplication, then*

$$\sum_{j=0}^n \omega_{n,j} \cdot u_{2n-j} = L. \quad (613)$$

*Thus the cancellation needs no multiplication, topology, order, or norm on the target. Taking  $M = \mathbb{Q}$  recovers the rational formula  $u_p = L + \sum_{r=1}^n a_r 2^{-rp}$ . This module-valued extension concerns cancellation only; the root classification above remains a statement for rational  $z$ . All statements include  $n = 0$ : the row polynomial is then 1, the root set is empty, and the error sum is empty.*

*Formal proof.* The full generating identity and its root classification are `sum_range_discreteLimitWeight_mul_pow` and `sum_range_discreteLimitWeight_mul_pow_eq_zero_iff`. The reusable single-mode identity is `sum_range_discreteLimitWeight_mul_geometricMode_smul_eq_zero`, and the module-valued cancellation theorem is `discreteLimitWeight_qRichardson_exact_module`; its  $M = \mathbb{Q}$  specialization is `discreteLimitWeight_qRichardson_exact`. All are in `FabiusDiscreteLimitToeplitz.lean`. The former proof of (610) is now its specialization at  $z = 1$ .  $\square$



*Remark 10.6* (Finite moments, Lagrange weights, and local exactness). The row calculation is an instance of two reusable algebraic layers. First, let  $I$  be finite, let  $\varphi : R \rightarrow S$  be a homomorphism from a semiring to a commutative semiring, let  $w_i, x_i \in S$  for  $i \in I$ , and let  $p \in R[X]$  have degree at most  $N$ . If  $p^\varphi$  denotes coefficientwise mapping, then the finite weighted functional expands as

$$\sum_{i \in I} w_i p^\varphi(x_i) = \sum_{d=0}^N \varphi([X^d]p) \sum_{i \in I} w_i x_i^d.$$

Thus prescribed moments select a prescribed coefficient; repeated nodes and a zero surviving moment are allowed, and no field or subtraction is needed. The same-ring coefficient and strict-cancellation forms are `sum_weight_mul_eval_eq_coeff_mul_of_moments` and `sum_weight_mul_eval_eq_zero_of_degree_lt` in `FinitePolynomialFunctional.lean`; that module also contains the scalar-extension forms used by the later Prouhet arguments.

Its exhaustive public surface has no definitions and exactly sixteen theorems: `sum_weight_mul_eval2_eq_sum_coeff_mul_moment`, `sum_weight_mul_eval2_eq_eval2_of_moments`, `sum_weight_mul_eval2_eq_coeff_mul_moment`, `sum_weight_mul_eval2_eq_topCoeff_mul_moment`, `sum_weight_mul_eval2_eq_constantCoeff_mul_sum`, `sum_weight_mul_eval2_eq_constantCoeff`, `sum_weight_mul_eval2_eq_map_coeff_mul_of_moments`, `sum_weight_mul_eval2_eq_map_coeff_mul_top_moment`, `sum_weight_mul_eval2_eq_zero_of_degree_lt`, `sum_weight_mul_eval2_congr_of_map_coeff_eq`, `sum_weight_mul_eval_eq_eval_of_moments`, `sum_weight_mul_eval_eq_coeff_mul_of_moments`, `sum_weight_mul_eval_eq_coeff_mul_top_moment`, `sum_weight_mul_eval_eq_zero_of_degree_lt`, `sum_weight_mul_eval_congr_of_coeff_eq`, and `sum_weight_mul_eval_affine_of_topCoeff_extractor`. The first ten allow scalar extension from an arbitrary semiring into a commutative semiring; the remaining same-ring forms require only a commutative semiring. Repeated nodes and a zero selected moment are allowed, and the degree-valued statements include  $n = 0$  and the zero polynomial.

The last theorem is the general affine transport principle. If a weighted functional extracts  $[X^n]q$  from every  $q$  of degree at most  $n$ , then

$$\sum_i w_i p(a + bx_i) = b^n c [X^n]p \quad (614)$$

for every  $p$  of degree at most  $n$ . Put  $q(X) = p(a + bX)$ , apply the supplied extractor to  $q$ , and compute  $[X^n]q = b^n [X^n]p$ . The implementation treats  $b = 0$  separately when proving this coefficient formula, so (614) needs neither  $b \neq 0$  nor distinct transformed nodes and includes  $n = 0$ . Composing it with `halfQBinomial_negativeDyadic_polynomial_sum_eq_mersenne` proves the  $q$ -monograph's `cor:geometric-prouhet-affine` exactly under its established  $\mathbb{Q}[X]$  convention.

Second, over a field, distinct nodes  $v_i$  have evaluation weights  $\lambda_i(x)$  given by the Lagrange basis at  $x$ . They reproduce every polynomial of degree less than the number of nodes, are characterized uniquely by their monomial moments, and at the first omitted degree satisfy

$$\sum_i \lambda_i(x) v_i^{|I|} = x^{|I|} - \prod_i (x - v_i).$$

These are respectively `sum_lagrangeEvalWeight_mul_eval`, `eq_lagrangeEvalWeight_of_moments`, and `sum_lagrangeEvalWeight_mul_pow_card`. On the distinct geometric nodes  $1, q, \dots, q^n$  at the target 0, the moments through degree  $n$  are  $0^d$ ; in particular degree zero has mass one, while positive degrees vanish. The declarations `sum_geometricLagrangeWeight_mul_pow` and `geometricLagrange_richardson_exact_of_eq` in `GeometricLagrange.lean` include both  $n = 0$  and an arbitrary block start, including start zero.

The first omitted degree is only the first member of an exact all-order family. Let  $h_r(x, (v_i)_{i \in I})$  be the complete homogeneous symmetric polynomial of degree  $r$  in the target  $x$  and the node family. Then

$$\sum_i \lambda_i(x) v_i^{|I|+r} = x^{|I|+r} - \prod_i (x - v_i) h_r(x, (v_i)_{i \in I}) \quad (r \geq 0). \quad (615)$$

More generally, the same formula holds over any commutative ring for arbitrary weights and possibly repeated nodes whenever the moments below  $|I|$  reproduce evaluation at  $x$ . No division or distinctness is used in this representation-independent form. The universal symmetric-polynomial evaluation and its adjoining-variable recurrence are `completeHomogeneousEval_eq_eval_hsymm` and `completeHomogeneousEval_option_succ`; the quotient evaluation is `eval_completeHomogeneousQuotient` in `CompleteHomogeneous.lean`. The companion `CompleteHomogeneousGenerating.lean` packages these evaluations as the formal series `completeHomogeneousGeneratingSeriesOn`. Its coefficient and adjoining-variable laws are `coeff_completeHomogeneousGeneratingSeriesOn` and `completeHomogeneousGeneratingSeriesOn_insert`; over every commutative ring, `prod_one_sub_mul_completeHomogeneousGeneratingSeriesOn` clears the finite linear-denominator product and `completeHomogeneousGeneratingSeriesOn_eq_invOfUnit_prod` identifies the series with its canonical `PowerSeries.invOfUnit` reciprocal. This is a formal coefficient identity and asserts no analytic evaluation, convergence, or Bell/generalized-harmonic conversion.

The separate finite-algebra module `CompleteHomogeneousBell.lean` supplies precisely that conversion. For a finite family  $a_i$ , put

$$p_k = \sum_i a_i^k, \quad \kappa_0 = 0, \quad \kappa_{k+1} = k! p_{k+1}.$$

The definitions are `completeHomogeneousPowerSum` and `completeHomogeneousBellInput`. Over every commutative semiring, the division-free identity

$$\text{Bell.complete}(\kappa)_n = n! h_n(a) \quad (616)$$

is `bellComplete_completeHomogeneousBellInput`. Over a commutative  $\mathbb{Q}$ -algebra, the normalized pointwise report form

$$h_n(a) = \text{factorialNormalize}(\text{completeBellPolynomial}(\kappa))_n$$

is `completeHomogeneousEvalOn_eq_factorialNormalize_completeBellPolynomial`. These statements are finite algebra: the first uses no division, and neither asserts convergence of a formal or analytic series.

The generic residual theorem `sum_weight_mul_pow_card_add` and its distinct-node Lagrange specialization `sum_lagrangeEvalWeight_mul_pow_card_add` are in `LagrangeResidualMoments.lean`. Thus (615) extends the preceding first-error formula rather than replacing its low-degree exactness foundation.

There is also a finite-family analytic companion to this formal algebra. The theorem `completeHomogeneousEvalOn_isBigO_pow` in `CompleteHomogeneousAsymptotics.lean` says that coordinatewise  $\mathcal{O}(g)$  bounds on a fixed finite alphabet give an  $\mathcal{O}(g^n)$  bound for its degree- $n$  complete homogeneous evaluation. This evaluated asymptotic statement does not assert convergence of the preceding formal power series or its Bell/generalized-harmonic conversion.

On the shifted reciprocal alphabet  $b_j = (\lambda + j)^{-1}$ ,  $0 \leq j \leq r$ , `LambertPhaseLockedBell.lean` names the generalized harmonic sums and their Bell input as `shiftedReciprocalPowerSum` and `shiftedReciprocalBellInput`. The exact specialization `completeHomogeneousEvalOn_shiftedReciprocal_eq_bell` rewrites  $h_n(b_0, \dots, b_r)$  by the right-hand side of the normalized form above. The declarations `sum_shiftedReciprocalLagrangeWeight_mul_invPow_card`

`add_eq_bell` and `sum_shiftedReciprocalLagrangeWeight_mul_invPow_eq_bell` apply the same rewrite to the all-offset and  $s \geq 1$  reciprocal-grid moments. These normalized identities are stated over characteristic-zero fields; the direct complete-homogeneous specialization uses the rational-algebra structure. The shifted-reciprocal definitions require no positivity or nonvanishing hypothesis on  $\lambda$ , because field inversion in Lean is total.

For the actual Fabius endpoint problem, put  $x_{\lambda,j} = (\lambda + j)2^{-(\lambda+j)}$ . The module `FabijsLambertPhaseLockedPullback.lean` pulls the arbitrary-order logarithmic expansion back to each fixed  $x_{\lambda,j}$ , freezing the one-periodic correction and all periodic saddle coefficients at the base phase  $\lambda$ . The estimates in `LambertReciprocalAsymptotics.lean` show that the fixed order- $r$  reciprocal Lagrange row has total variation  $\mathcal{O}(\lambda^r)$  and balance this growth against sufficiently high shifted inverse powers. Writing  $\Psi$  for `negativeLaplacePsi`, the definitions `fabijsPhaseLockedReducedSample`, `fabijsPhaseLockedPeriodicEstimator`, `fabijsPhaseLockedResidualTerm`, and `fabijsPhaseLockedResidualPartialSum` in `FabijsLambertPhaseExtraction.lean` give, for every fixed  $r, S \in \mathbb{N}_0$ ,

$$E_r(\lambda) - \Psi(\lambda) - \sum_{n=0}^{S-1} R_{r,n}(\lambda) = \mathcal{O}(\lambda^{-(r+1+S)}), \quad \lambda \rightarrow +\infty. \quad (617)$$

This is `fabijsPhaseLockedPeriodicEstimator_sub_residual_isBigO`; its proof uses the node expansion through order  $2r + S + 1$ . The cases  $S = 0$  and  $S = 1$  are the named leading and first-omitted-moment bounds, and `fabijsPhaseLockedPeriodicEstimator_tendsto_periodicAlong` gives convergence to  $\Psi(a)$  along  $\lambda = a + n$ . These are fixed-index Poincaré estimates; no uniformity for  $r$  or  $S$  growing with  $\lambda$  is claimed.

Finally, `FabijsLambertPhaseExtractionBell.lean` changes neither the residual definitions nor the preceding estimate. It gives the exact rewrite

$$R_{r,n}(\lambda) = (-1)^r \left( \prod_{j=0}^r (\lambda + j)^{-1} \right) \times \text{factorialNormalize}(\text{completeBellPolynomial}(\kappa^{(r,\lambda)}))_n c_{r+1+n}(\lambda),$$

where  $\kappa^{(r,\lambda)}$  is `shiftedReciprocalBellInput` and  $c_m$  denotes `fabijsSaddleLogCoefficient`. The single-term and finite-sum declarations are `fabijsPhaseLockedResidualTerm_eq_bell` and `fabijsPhaseLockedResidualPartialSum_eq_sum_bell`. This three-module Bell tranche closes only the finite complete-homogeneous/Bell/generalized-harmonic conversion and the exact finite residual rewrites. It proves no convergence of the infinite residual series, multiplicative or exponentiated Bell relative-error hierarchy, higher extractor, Lambert sign or bracketing claim, or growing-order uniformity.

*Proposition 10.7 (Principal specialization and denominator-free geometric residuals).* *Let  $R$  first be a commutative semiring. For arbitrary  $c, q \in R$  and  $p, r \in \mathbb{N}_0$ , complete homogeneous evaluation on a geometric alphabet is*

$$h_r(1, q, \dots, q^p) = \begin{bmatrix} p+r \\ p \end{bmatrix}_q = \begin{bmatrix} p+r \\ r \end{bmatrix}_q, \quad (618)$$

$$h_r(c, cq, \dots, cq^p) = c^r \begin{bmatrix} p+r \\ p \end{bmatrix}_q = c^r \begin{bmatrix} p+r \\ r \end{bmatrix}_q. \quad (619)$$

*These identities are recursive and denominator-free: they remain valid at  $q = 0$ , at roots of unity, in positive characteristic, and in the zero semiring.*

Now let  $R$  be a commutative ring and let  $w_0, \dots, w_p \in R$ . If

$$\sum_{k=0}^p w_k (q^k)^d = 0^d \quad (0 \leq d \leq p), \quad (620)$$

then every positive moment is

$$\sum_{k=0}^p w_k (q^k)^m = (-1)^p q^{\binom{p+1}{2}} \begin{bmatrix} m-1 \\ p \end{bmatrix}_q, \quad m > 0. \quad (621)$$

There is a strictly more general scaled-node form. If exactness is assumed directly on the displayed nodes,

$$\sum_{k=0}^p w_k (cq^k)^d = 0^d \quad (0 \leq d \leq p), \quad (622)$$

then

$$\sum_{k=0}^p w_k (cq^k)^m = c^m (-1)^p q^{\binom{p+1}{2}} \begin{bmatrix} m-1 \\ p \end{bmatrix}_q, \quad m > 0. \quad (623)$$

No nonzeroness or unit hypothesis is imposed on  $c$ ; this direct scaled hypothesis is therefore meaningful even for zero divisors.

Over a field, distinctness of  $1, q, \dots, q^p$  supplies (620) for the Lagrange weights  $\lambda_{p,k}^{(q)}$ . More precisely, injectivity of  $k \mapsto q^k$  on  $0 \leq k \leq p$  gives, for every  $P \in F[X]$  with  $\text{natDegree } P \leq p$ ,

$$\sum_{k=0}^p \lambda_{p,k}^{(q)} P(cq^k) = P(0).$$

Here  $c$  is arbitrary, including zero; this strictly subsumes the nonzero-scale premise of the  $q$ -monograph's *cor:scaled-geometric-moments*. Together with (623), it makes that corollary *Exact by composition*. Consequently, for every block start  $s \in \mathbb{N}_0$ ,

$$\sum_{k=0}^p \lambda_{p,k}^{(q)} (q^{s+k})^m = (-1)^p q^{sm + \binom{p+1}{2}} \begin{bmatrix} m-1 \\ p \end{bmatrix}_q, \quad m > 0. \quad (624)$$

In particular, at block start zero and degree  $m = p + 1 + r$ , complementary-index symmetry gives the *report-facing orientation*

$$\sum_{k=0}^p \lambda_{p,k}^{(q)} (q^k)^{p+1+r} = (-1)^p q^{\binom{p+1}{2}} \begin{bmatrix} p+r \\ r \end{bmatrix}_q. \quad (625)$$

*Formal proof.* The semiring-level infrastructure is `completeHomogeneousEval_smul`, `completeHomogeneousEval_option_zero`, `completeHomogeneousEval_fin_succ`, `completeHomogeneousEvalOn_range`, and `completeHomogeneousEvalAt_zero` in `CompleteHomogeneous.lean`. Splitting off the leading variable 1 and scaling the remaining alphabet by  $q$  turns the head–tail recurrence into  $q$ -Pascal's recurrence. This proves `completeHomogeneousEval_geometric`; its common-scale and range-indexed forms are `completeHomogeneousEval_scaled_geometric` and `completeHomogeneousEvalOn_range_pow_eq_gaussianBinomial`. A second direct principal-specialization induction gives the degree-indexed orientation `completeHomogeneousEvalOn_range_pow_eq_gaussianBinomial_degree`. Comparing the two orientations proves the denominator-free identities `gaussianBinomial_add_symm` and `gaussianBinomial_symm_via_completeHomogeneous`; the latter is a second, principal-specialization proof of the core reflection theorem. The report-facing range alias is `completeHomogeneousEvalOn_geometric_range`; all

of these declarations are in `GeometricCompleteHomogeneous.lean` or `GeometricLagrangeCompleteHomogeneous.lean`.

The offset-degree engines for  $m = p + 1 + r$  are `sum_weight_mul_geometric_pow_succ_add` and `sum_weight_mul_scaled_geometric_pow_succ_add`. Above-diagonal vanishing of the recursive Gaussian coefficient folds the cancelled degrees and every higher residual into `sum_weight_mul_geometric_pow_of_pos` and `sum_weight_mul_scaled_geometric_pow_of_pos`. Their field-valued Lagrange specializations are `sum_geometricLagrangeWeight_mul_pow_succ_add`, `sum_geometricLagrangeWeight_mul_pow_of_pos`, `sum_geometricLagrangeWeight_mul_scaled_geometric_pow_of_pos`, and `sum_geometricLagrangeWeight_mul_shifted_pow_of_pos`, together with the polynomial-exactness theorem `sum_geometricLagrangeWeight_mul_eval_scaled_geometric`, in the zero-definition/nine-theorem module `GeometricResidualMoments.lean`. The pre-existing pointwise closed weight and cleared-denominator formulas remain in `GeometricQBinomialLagrange.lean`; its positive-moment theorem is now a thin corollary of this denominator-free engine. Finally, (625) is `sum_geometricLagrangeWeight_mul_pow_succ_add_eq_gaussianBinomial` in `GeometricLagrangeCompleteHomogeneous.lean`; it changes only the complementary Gaussian index and introduces no new interpolation argument.  $\square$

The same moment calculation acts diagonally on every formal power series, not only on a polynomial of bounded degree. This produces a reusable operator in which low-degree cancellation and every surviving geometric residual are visible coefficient by coefficient.

*Proposition 10.8 (Finite formal-series filters and the Gaussian residual). Let  $R$  be a commutative semiring, let  $\iota$  be an arbitrary index type, let  $S \subseteq \iota$  be finite, and choose nodes  $x_i \in R$ , weights  $w_i \in R$ , and a formal power series  $f \in R[[X]]$ . Repeated nodes are allowed. Define*

$$\mathcal{T}_{S,x,w}f(X) = \sum_{i \in S} w_i f(x_i X). \quad (626)$$

*Then, for every  $m \in \mathbb{N}_0$ ,*

$$[X^m]\mathcal{T}_{S,x,w}f = \left( \sum_{i \in S} w_i x_i^m \right) [X^m]f. \quad (627)$$

*The operator commutes with a further rescaling of  $X$ , and with every commutative-semiring homomorphism when the nodes, weights, and coefficients are transported together.*

*More generally, suppose a target  $t \in R$  and an order  $p \in \mathbb{N}_0$  satisfy*

$$\sum_{i \in S} w_i x_i^d = t^d \quad (0 \leq d \leq p). \quad (628)$$

*Degree zero in this hypothesis gives  $\sum_i w_i = 1$ . Hence the filter preserves the constant coefficient, and for every  $m \leq p$ ,*

$$[X^m]\mathcal{T}_{S,x,w}f = t^m [X^m]f. \quad (629)$$

*At  $t = 0$ , this preserves degree zero and cancels every degree  $1 \leq m \leq p$ , including the boundary case  $p = 0$  without pretending that  $0^0$  vanishes.*

*Now let  $R$  be a commutative ring, take the nodes  $1, q, \dots, q^p$ , and assume the exactness condition (620). For*

$$\mathcal{G}_{q,p,w}f(X) = \sum_{j=0}^p w_j f(q^j X),$$

*all coefficients, including the mass-one boundary, are*

$$[X^m]\mathcal{G}_{q,p,w}f = \begin{cases} [X^0]f, & m = 0, \\ (-1)^p q^{\binom{p+1}{2}} \begin{bmatrix} m-1 \\ p \end{bmatrix}_q [X^m]f, & m > 0. \end{cases} \quad (630)$$

Thus the above-diagonal zero of the recursive Gaussian coefficient gives all cancelled degrees, and the same formula gives every later residual. No division, convergence, distinctness, nonzeroness, or field structure is used. Over a field, distinctness of the geometric nodes makes the Lagrange row a canonical specialization of the same identity.

The generic operator and its diagonal action are `finitePowerSeriesFilter` and `coeff_finitePowerSeriesFilter` in `FinitePowerSeriesFilter.lean`. Its rescaling and scalar-transport laws are `finitePowerSeriesFilter_rescale` and `map_finitePowerSeriesFilter`; exactness, mass one, constant preservation, and low-degree cancellation are `sum_weight_eq_one_of_finite_exact`, `constantCoeff_finitePowerSeriesFilter_of_exact`, `coeff_finitePowerSeriesFilter_of_exact_of_le`, and `coeff_finitePowerSeriesFilter_eq_zero_of_pos_of_le`. The geometric operator is `geometricSeriesFilter`; its positive-degree coefficient theorem and the all-index identity (630) are `coeff_geometricSeriesFilter_of_exact` and `geometricSeriesFilter_eq_residual_mk` in `GeometricPowerSeriesFilter.lean`. The distinct-node Lagrange specializations are `coeff_geometricLagrangeSeriesFilter_of_pos` and `geometricLagrangeSeriesFilter_eq_residual_mk`. Every series in this paragraph is formal: none of these declarations asserts analytic convergence, evaluation at a nonzero argument, or an infinite acceleration formula.

There is now a separate convergent-series companion. It deliberately uses Mathlib’s unconditional summability rather than silently identifying a natural-order boundary sum with a `tsum`.

*Proposition 10.9* (Unconditionally summable analytic filters and their Gaussian tail). *Let  $K$  be a commutative semiring, let  $E$  be a normed additive commutative group with a  $K$ -module structure, let  $S$  be finite, and choose nodes  $x_i \in K$ , weights  $w_i \in K$ , coefficients  $a_m \in E$ , and a scale  $z \in K$ . Define*

$$\mathcal{A}_{S,x,w}(a; z) := \sum_{i \in S} \sum_{m \geq 0} w_i (x_i z)^m a_m. \quad (631)$$

*Assume for every  $i \in S$  that the displayed weighted inner series is unconditionally summable. Then the finite and infinite sums interchange and*

$$\mathcal{A}_{S,x,w}(a; z) = \sum_{m \geq 0} \left( \sum_{i \in S} w_i x_i^m \right) z^m a_m. \quad (632)$$

*If, moreover,  $\sum_{i \in S} w_i x_i^d = t^d$  for  $0 \leq d \leq p$ , then*

$$\begin{aligned} \mathcal{A}_{S,x,w}(a; z) &= \sum_{d=0}^p (tz)^d a_d \\ &\quad + \sum_{r \geq 0} \left( \sum_{i \in S} w_i x_i^{p+1+r} \right) z^{p+1+r} a_{p+1+r}. \end{aligned} \quad (633)$$

*At target  $t = 0$ , the finite head is exactly  $a_0$ , including  $p = 0$ ; the positive degrees never pass through the  $0^0$  boundary.*

*Now let  $K$  be a commutative ring with continuous constant scalar multiplication on  $E$  (each fixed scalar acts continuously), take  $x_j = q^j$  for  $0 \leq j \leq p$ , and suppose the row is exact at zero through degree  $p$ ; write  $\mathcal{A}_{q,p,w}$  for this geometric specialization. It is enough to assume unconditional summability of  $\sum_m (q^j z)^m a_m$  only when  $w_j \neq 0$ . Then*

$$\mathcal{A}_{q,p,w}(a; z) = a_0 + \sum_{r \geq 0} (-1)^p q^{\binom{p+1}{2}} \begin{bmatrix} p+r \\ p \end{bmatrix}_q z^{p+1+r} a_{p+1+r}. \quad (634)$$

*Over a field, distinct geometric nodes supply the Lagrange specialization and its shifted-scale form  $z = q^{\text{start}}$ . A zero-weight node imposes no convergence condition: its weighted series is identically zero.*



The definitions are `finiteAnalyticSeriesFilter` and `geometricAnalyticSeriesFilter` in `AnalyticSeriesFilter.lean`. The generic diagonal, exact-head, and zero-target declarations are `summable_finiteAnalyticSeriesFilter_diagonal`, `finiteAnalyticSeriesFilter_eq_tsum`, `finiteAnalyticSeriesFilter_eq_head_add_tail_of_exact`, and `finiteAnalyticSeriesFilter_eq_constant_add_tail_of_exact_zero`. The Gaussian, Lagrange, and shifted-scale tail declarations are `geometricAnalyticSeriesFilter_eq_constant_add_gaussian_tsum`, `geometricLagrangeAnalyticSeriesFilter_eq_constant_add_gaussian_tsum`, and `geometricLagrangeAnalyticSeriesFilter_shifted`. Thus this module has exactly two definitions and seven public theorems. These results exclude series converging only in their natural order at a boundary point. They also assert no formal-to-analytic power-series evaluation bridge, radius or uniform-convergence theorem, norm/sign/error bound, asymptotic estimate, or positivity statement. The actual Rvachev sinc-product instantiation is a separate theorem, so that the generic module retains precisely this abstract scope.

*Theorem 10.10* (Exact Gaussian filter of the Rvachev sinc product). *Let  $c, z \in \mathbb{C}$ , let  $p \in \mathbb{N}_0$ , put  $q = c^2$ , and assume only that*

$$j \mapsto q^j \quad \text{is injective on } \{0, \dots, p\}.$$

*Writing  $\Phi = \text{rvachevFourierProduct} = \widehat{\text{up}}_{2\pi}$ , one has the exact entire identity*

$$\sum_{j=0}^p \lambda_{p,j}^{(q)} \Phi(c^j z) = 1 + \sum_{r=0}^{\infty} (-1)^p q^{\binom{p+1}{2}} \begin{bmatrix} p+r \\ p \end{bmatrix}_q A_{p+1+r}(z). \quad (635)$$

*There is no contraction, realness, positivity, or global nonvanishing hypothesis on  $c$ . At  $c = 1/2$ , hence  $q = 1/4$ , the finite node condition is proved internally, so (635) holds for every complex  $z$  and every natural  $p$  with no hypothesis. The same generic formula is exported for the Fourier transform of every bounded  $F$  satisfying *IsFabius*  $F$ .*

*Formal proof.* The named moment series sums to the standalone product by `hasSum_rvachevFourierMomentTerm_product`, and its iterated scaling law is `rvachevFourierMomentTerm_pow_scale`. Passing the finite Lagrange row through that unconditional sum and inserting the Gaussian residual gives `geometricLagrange_rvachevFourierProduct_eq_gaussian_tsum`. Its assumption-free  $c = 1/2$ ,  $q = 1/4$  specialization is `quarterLagrange_rvachevFourierProduct_eq_gaussian_tsum`; the model-generic Fourier-transform form is `geometricLagrange_rvachevFourier_eq_gaussian_tsum`. These five declarations exhaust `RvachevQBinomialFilter.lean`.  $\square$

The theorem closes the coefficient bridge for the *infinite* Rvachev product. It does not identify the frontier report's finite products  $P_{b,n}$ , their quotient  $P_{b,n}/P_b$ , or their Bell coefficients with this series. Nor does it prove a conditionally convergent boundary theorem, an analytic sign or bracketing/error bound, uniform or derivative convergence, or an asymptotic acceleration estimate. The algebraic  $(-1)^p$  in (635) should not be promoted to one of those analytic sign claims.

For the present half-geometric grid, write  $\lambda_{n,k}^{(1/2)}$  for the evaluation-at-zero weight belonging to the node  $2^{-k}$ . The Toeplitz row is exactly the same row in reverse order:

$$\boxed{\omega_{n,j} = \lambda_{n,n-j}^{(1/2)}} \quad (0 \leq j \leq n).$$

Indeed, for every  $d \geq 0$  its complete reversed moment is

$$\sum_{k=0}^n \omega_{n,n-k} 2^{-kd} = 2^{-nd} \frac{(2^d/2; 1/2)_n}{(1/2; 1/2)_n},$$

and for  $0 \leq d \leq n$  this is  $0^d$ , with  $0^0 = 1$ . The pointwise bridge and these two moment forms are `discreteLimitWeight_eq_geometricLagrangeWeight`, `geometricLagrangeWeight_half_eq_discreteLimitWeight`, `sum_range_discreteLimitWeight_reverse_mul_half_pow`, and `sum_range_discreteLimitWeight_reverse_moment`. Their finite-sum form is `sum_range_discreteLimitWeight_smul_eq_geometricLagrangeWeight`.

For every  $d > 0$ , the same reversed row has the sharper residual form

$$\sum_{k=0}^n \omega_{n,n-k} \left( (1/2)^k \right)^d = (-1)^n (1/2)^{n(n+1)/2} \begin{bmatrix} d-1 \\ n \end{bmatrix}_{1/2}. \quad (636)$$

This is `sum_range_discreteLimitWeight_reverse_moment_eq_halfQBinomial` in `FabiusDiscreteLimitToeplitz.lean`. Thus one Gaussian coefficient records every positive residual degree: its above-diagonal boundary gives the cancelled range, while its diagonal boundary gives the first surviving moment. The corresponding boundary declarations are `gaussianBinomial_eq_zero_of_lt` and `gaussianBinomial_self` in `FiniteQBinomialCore.lean`.

Consequently the global model in the proposition can be weakened to one finite block. For any  $s \in \mathbb{N}_0$ , if only the samples  $0 \leq k \leq n$  satisfy

$$u_{s+k} = L + \sum_{r=1}^n 2^{-r(s+k)} \cdot a_r,$$

then

$$\sum_{j=0}^n \omega_{n,j} \cdot u_{s+n-j} = L.$$

This is `discreteLimitWeight_qRichardson_exact_module_at_of_eq` in `FabiusDiscreteLimitToeplitz.lean`. At  $n = 0$  its hypothesis is simply  $u_s = L$ , and  $s = 0$  needs no separate convention. The grid ratio is  $1/2$ . A mode with ratio  $1/4$  and index  $r$  has half-grid degree  $2r$ , so this row cancels it only when  $2r \leq n$ ; this is not an  $n$ -mode theorem on a  $1/4$  grid.

Applying it with the opposite sign also gives the uniform variation bound

$$\sum_{j=0}^n |\omega_{n,j}| = \frac{(-1/2; 1/2)_n}{(1/2; 1/2)_n} \leq 16. \quad (637)$$

Finally, every sampled degree satisfies  $2n - j \geq n$ . A finite-row Toeplitz lemma therefore carries the convergence of  $S_p^{(Q)}(x)$  to the weighted rows (608). Explicitly, given  $\epsilon > 0$ , choose  $P$  so that  $|S_p^{(Q)}(x) - F_{\text{gl}}(x)| < \epsilon/16$  for  $p \geq P$ . If  $n \geq P$ , then every  $2n - j \geq n \geq P$ , and the row-sum identity together with (637) gives

$$\left| \sum_{j=0}^n \omega_{n,j} S_{2n-j}^{(Q)}(x) - F_{\text{gl}}(x) \right| \leq \sum_{j=0}^n |\omega_{n,j}| |S_{2n-j}^{(Q)}(x) - F_{\text{gl}}(x)| < \epsilon.$$

This proves (598). □

Two distinctions prevent common misreadings of this theorem. First, unlike the complete finite dyadic numerator and each summand of the global  $q$ -series, a finite value  $\mathcal{D}_n^{(Q)}(x)$  can depend on  $Q$ ; it is the limit that is independent of a fixed shift. Second, the Toeplitz row (608) is not a finite binary telescope or a partial sum of (594). These finite expressions share a limit but are not equal term by term.

The two properties of the reindexed row (609) used above are both evaluations of the finite  $q$ -binomial theorem (486), at  $z = 1/2$  and at  $z = -1/2$  respectively. Both use the exponent identity  $\binom{j}{2} + j = \binom{j+1}{2}$ .



For the row sum, evaluate (486) at  $z = 1/2$ . The left side is  $(1/2; 1/2)_n$  by definition, and the right side is

$$(1/2; 1/2)_n = \sum_{j=0}^n (-1)^j 2^{-\binom{j}{2}} \left[ \begin{matrix} n \\ j \end{matrix} \right]_{1/2} \left( \frac{1}{2} \right)^j = \sum_{j=0}^n (-1)^j \left[ \begin{matrix} n \\ j \end{matrix} \right]_{1/2} 2^{-(j+1)}. \quad (638)$$

Dividing by  $(1/2; 1/2)_n$  gives exactly  $\sum_{j=0}^n \omega_{n,j} = 1$ , which is (610).

For the total variation, evaluate (486) at  $z = -1/2$  instead. Each factor  $(-1)^j (-1/2)^j = 2^{-j}$  is now positive, so the alternating signs disappear:

$$(-1/2; 1/2)_n = \sum_{j=0}^n (-1)^j 2^{-\binom{j}{2}} \left[ \begin{matrix} n \\ j \end{matrix} \right]_{1/2} \left( -\frac{1}{2} \right)^j = \sum_{j=0}^n \left[ \begin{matrix} n \\ j \end{matrix} \right]_{1/2} 2^{-(j+1)}. \quad (639)$$

Since  $\left[ \begin{matrix} n \\ j \end{matrix} \right]_{1/2} > 0$ , the summands of (639) are the absolute values of those of (638). Dividing by  $(1/2; 1/2)_n$  gives the exact evaluation

$$\sum_{j=0}^n |\omega_{n,j}| = \frac{(-1/2; 1/2)_n}{(1/2; 1/2)_n}, \quad (640)$$

which is the first equality in (637). The formally proved finite-product estimate supplies the bound by 16 used in the Toeplitz argument above.

## 11 Iterated prefix sums of the signed Thue–Morse sequence

The signed Thue–Morse sequence  $\varepsilon_n = (-1)^{\text{wt}_2(n)}$  has already appeared twice in this article: once as the coefficient sequence of the locally finite global expansion (11), and once as the sign pattern that makes the finite dyadic sums of section 8 collapse. There is a third, entirely discrete, way in which the same signs produce the Fabius function. Sum the sequence, sum the result, and iterate. After  $k$  summations the sequence has been convolved with an explicit binomial kernel, and after a dyadic normalization the resulting integer grid reproduces, in the limit, the Rvachev bump and hence  $F$ .

This section develops that construction: the convolution law and its explicit binomial kernel, the signed dyadic reflection preceding each terminal zero run, the zero runs themselves and their exact boundary values, the two generating-series identities, their analytic realization and natural-boundary theorem, the discrete functional equation satisfied by the normalized grid, and the precise sense in which the normalized grid converges. The convergence statement requires one shift of the prefix order and the centered reading of the grid cells; the literal normalization does not converge, and that failure is recorded separately in section 11.6. Two further pieces of foundational material are attached at the end: the lower branch of Lambert’s  $W$ -function, which the asymptotic sections of this article use but never construct, and an exact binomial–logarithm formula for the Thue–Morse bit.

The two binary recurrences used throughout this section are

$$\varepsilon_{2m} = \varepsilon_m, \quad \varepsilon_{2m+1} = -\varepsilon_m, \quad (641)$$

$$\text{wt}_2(2m) = \text{wt}_2(m), \quad \text{wt}_2(2m+1) = \text{wt}_2(m) + 1. \quad (642)$$

They are exactly `thueMorseSign_two_mul`, `thueMorseSign_two_mul_add_one`, `binaryWeight_two_mul`, and `binaryWeight_two_mul_add_one` in `DyadicClosedForm.lean`.

### 11.1 Iterated prefix sums and the binomial convolution law

**Definition 11.1** (Iterated prefix sums). Set  $\sigma_0(n) = \varepsilon_n$  and, for  $k \geq 0$ ,

$$\sigma_{k+1}(n) = \sum_{j=0}^n \sigma_k(j). \quad (643)$$

All sums are *inclusive*: the index  $n$  itself is summed.

Every  $\sigma_k(n)$  is an integer, and  $\sigma_k(0) = 1$  for all  $k$ . The inclusive convention has one consequence that must be kept in view throughout, because it is the source of an index shift later:

$$\sigma_{k+1}(n+1) - \sigma_{k+1}(n) = \sigma_k(n+1). \quad (644)$$

The increment of a prefix sum is the lower-order value at the *new* index, not at the old one.

The first prefix sum nevertheless vanishes at every odd index:

$$\sigma_1(2m+1) = 0 \quad (m \in \mathbb{N}_0).$$

This is exactly `iteratedPrefix_one_two_mul_add_one` in `ThueMorsePrefix.lean`.

The general structure is a discrete convolution. Write

$$\mathcal{B}_k(i) = \binom{i+k-1}{k-1} \quad (k \geq 1, i \geq 0) \quad (645)$$

which is `iteratedPrefixKernel`. Because the argument  $i = n - m$  is nonnegative in the sums below, no separate indicator is needed.

**Theorem 11.2** (Binomial convolution law). *For every  $k \geq 1$  and every  $n \geq 0$ ,*

$$\sigma_k(n) = \sum_{m=0}^n \mathcal{B}_k(n-m) \varepsilon_m = \sum_{m=0}^n \binom{n-m+k-1}{k-1} \varepsilon_m. \quad (646)$$

Equivalently, in a form valid for every  $k \geq 0$ ,

$$\sigma_{k+1}(n) = \sum_{m=0}^n \binom{n-m+k}{k} \varepsilon_m. \quad (647)$$

*Proof.* Induct on  $k$  in the form (647). For  $k = 0$  the binomial coefficient is 1 and the assertion is the definition of  $\sigma_1$ . Assume (647) for  $k$ . Then

$$\begin{aligned} \sigma_{k+2}(n) &= \sum_{j=0}^n \sigma_{k+1}(j) = \sum_{j=0}^n \sum_{m=0}^j \binom{j-m+k}{k} \varepsilon_m \\ &= \sum_{m=0}^n \varepsilon_m \sum_{i=0}^{n-m} \binom{i+k}{k} = \sum_{m=0}^n \binom{n-m+k+1}{k+1} \varepsilon_m, \end{aligned}$$

where the interchange is the triangular reindexing  $i = j - m$  and the inner evaluation is the hockey-stick identity  $\sum_{i=0}^N \binom{i+k}{k} = \binom{N+k+1}{k+1}$ .  $\square$

**Remark 11.3.** The kernel in (646) is exactly the coefficient sequence of  $(1-z)^{-k}$ . **Theorem 11.2** is therefore the coefficient form of the series identity proved in [theorem 11.12](#) below, and either statement may be taken as the definition of the other. Both are recorded here because the coefficient form is what one actually evaluates, while the series form is what one manipulates.

## 11.2 Prouhet annihilation, signed dyadic reciprocity, and exact zero runs

[Equation \(455\)](#) already records that a complete dyadic block of Thue–Morse signs annihilates every polynomial of degree below the block exponent. For the prefix sums we also need the sharp boundary case, which identifies the first power that survives.

**Lemma 11.4** (Sharp Prouhet extraction). *Write  $S_{r,d} = \sum_{h < 2^r} \varepsilon_h h^d$ . Then  $S_{r,d} = 0$  for  $0 \leq d < r$ , and*

$$S_{r,r} = (-1)^r 2^{\binom{r}{2}} r!. \quad (648)$$

Consequently, for all rational numbers  $a_0, \dots, a_r$ ,

$$\sum_{h < 2^r} \varepsilon_h \sum_{d=0}^r a_d h^d = a_r (-1)^r 2^{\binom{r}{2}} r!, \quad (649)$$

so summation against a complete dyadic block of Thue–Morse signs annihilates every polynomial of degree below  $r$  and, on polynomials of degree at most  $r$ , extracts the leading coefficient up to the explicit nonzero factor  $(-1)^r 2^{\binom{r}{2}} r!$ .

*Proof.* The vanishing statement is (455) with  $X = 0, Y = 1$ . For the boundary case, split the block by parity rather than by halves. Since  $\varepsilon_{2j} = \varepsilon_j$  and  $\varepsilon_{2j+1} = -\varepsilon_j$  by (641),

$$S_{r+1,d} = \sum_{j < 2^r} \varepsilon_j [(2j)^d - (2j+1)^d] = - \sum_{c=0}^{d-1} \binom{d}{c} 2^c S_{r,c},$$

because  $(2j+1)^d - (2j)^d = \sum_{c < d} \binom{d}{c} (2j)^c$ . Take  $d = r+1$ . All terms with  $c < r$  vanish by the first assertion, leaving only  $c = r$ :

$$S_{r+1,r+1} = - \binom{r+1}{r} 2^r S_{r,r} = -(r+1) 2^r S_{r,r}.$$

Since  $S_{0,0} = \varepsilon_0 = 1$ , induction gives  $S_{r,r} = (-1)^r r! 2^{0+1+\dots+(r-1)} = (-1)^r r! 2^{\binom{r}{2}}$ , which is (648). Formula (649) follows by exchanging the two sums: every index  $d < r$  contributes  $a_d S_{r,d} = 0$ , and the index  $d = r$  contributes  $a_r S_{r,r}$ .  $\square$

**Corollary 11.5** (Affine sharp case). *For all  $X, Y \in \mathbb{Q}$ ,*

$$\sum_{h < 2^r} \varepsilon_h (X + Yh)^r = (-1)^r Y^r 2^{\binom{r}{2}} r!.$$

*In particular the sum does not depend on the translation  $X$ .*

*Proof.* Expand  $(X + Yh)^r$  by the binomial theorem and apply (649): the coefficient of  $h^r$  is  $Y^r$ .  $\square$

The sharp extractor also has a direct commutative-ring form. If  $R$  is any commutative ring and  $p \in R[X]$  satisfies  $\deg p \leq r$ , then, with the signs and natural numbers mapped into  $R$ ,

$$\sum_{h < 2^r} \varepsilon_h p(h) = [X^r]p (-1)^r 2^{\binom{r}{2}} r!. \quad (650)$$

This is `thueMorse_polynomial_sum_eq_coeff_ring_of_degree_le`; its strict-degree cancellation form is `thueMorse_polynomial_sum_eq_zero_ring_of_degree_lt`, both in `ThueMorsePrefix.lean`. The word “extractor” here describes the functional identity, not division by its sharp factor: in positive characteristic the image of  $2^{\binom{r}{2}} r!$  can vanish, so the displayed equality need not recover  $[X^r]p$  from the sum. The degree-valued hypothesis also treats the zero polynomial correctly; at  $r = 0$  the extraction formula includes constant polynomials and the zero polynomial, whereas the strict-degree condition admits only the latter.

Likewise the affine sharp identity holds directly in  $R$ :

$$\sum_{h < 2^r} \varepsilon_h (x + yh)^r = (-1)^r y^r 2^{\binom{r}{2}} r!, \quad (651)$$

formalized as `thueMorse_affine_power_sum_self_ring`, with the unified range  $d \leq r$  given by `thueMorse_affine_power_sum_of_le_ring`. For  $r > 0$  the sharp sum vanishes at  $y = 0$ . At the joint boundary  $r = 0, y = 0$ , Lean’s convention  $0^0 = 1$  instead gives the correct one-term value 1 on both sides; no positive-degree reading should be imposed on that case.

The usual equal-sums formulation is now available without leaving the same commutative-ring setting.

**Proposition 11.6** (Even–odd Prouhet partition). *Let  $R$  be a commutative ring, let  $s \subseteq \mathbb{N}_0$  be finite, and let  $g: \mathbb{N}_0 \rightarrow R$ . With the integer sign  $\varepsilon_h$  and each natural index mapped into  $R$ , one has*

$$\sum_{h \in s} \varepsilon_h g(h) = \sum_{\substack{h \in s \\ \text{wt}_2(h) \text{ even}}} g(h) - \sum_{\substack{h \in s \\ \text{wt}_2(h) \text{ odd}}} g(h). \quad (652)$$

Consequently, if  $d, r \in \mathbb{N}_0$  and  $d < r$ , then for every  $x, y \in R$ ,

$$\sum_{\substack{0 \leq h < 2^r \\ \text{wt}_2(h) \text{ even}}} (x + yh)^d = \sum_{\substack{0 \leq h < 2^r \\ \text{wt}_2(h) \text{ odd}}} (x + yh)^d. \quad (653)$$

In particular, setting  $x = 0$  and  $y = 1$  gives equality of the  $d$ th power sums of the even- and odd-binary-weight parts of  $\{0, \dots, 2^r - 1\}$ .

*Proof.* Every binary weight is either even or odd, and the two cases are disjoint. By the definition  $\varepsilon_h = (-1)^{\text{wt}_2(h)}$ , the sign is 1 in the first case and  $-1$  in the second, which proves (652). For the full dyadic block, expand the signed affine sum as

$$\sum_{k=0}^d \binom{d}{k} x^{d-k} y^k \sum_{h < 2^r} \varepsilon_h h^k.$$

Every inner moment vanishes because  $k \leq d < r$ . Thus the left side of (652), with  $g(h) = (x + yh)^d$ , is zero, and the two filtered sums agree. The raw-power statement is the indicated specialization.  $\square$

The three formal counterparts are respectively `sum_thueMorseSign_mul_eq_sum_even_binaryWeight_sub_sum_odd_binaryWeight`, `sum_even_binaryWeight_affine_pow_eq_sum_odd_binaryWeight_affine_pow`, and `sum_even_binaryWeight_pow_eq_sum_odd_binaryWeight_pow`, all in `ThueMorseSparseProuhet.lean`. The first is valid on an arbitrary finite set; the latter two use the whole dyadic block and the literal strict range  $d < r$ . No positive-characteristic nonvanishing assumption is needed, because these are equalities rather than coefficient-recovery statements.

The binary-support parametrization used next has a canonical global foundation. Write  $\text{bits}(n) = \text{bitSupport}(n)$  and  $E(T) = \sum_{j \in T} 2^j$  for a finite set  $T \subseteq \mathbb{N}_0$ . This support is exactly Mathlib’s finite set of binary indices, and

$$j \in \text{bits}(n) \iff n.\text{testBit}(j) = \text{true}, \quad E(\text{bits}(n)) = n, \quad \text{bits}(E(T)) = T. \quad (654)$$

Consequently

$$\text{wt}_2(E(T)) = |T|, \quad \varepsilon_{E(T)} = (-1)^{|T|}. \quad (655)$$

The exact declarations are `bitSupport_eq_toFinset_bitIndices`, `mem_bitSupport`, `sum_two_pow_bitSupport`, `bitSupport_sum_two_pow`, `binaryWeight_sum_two_pow_eq_card`, and `thueMorseSign_sum_two_pow` in `ThueMorseBitSupport.lean`.

Restricting the same inverse pair gives the canonical equivalence

$$\{T : T \subseteq \{j \in \mathbb{N}_0 : j < m\}\} \simeq \{h \in \mathbb{N}_0 : h < 2^m\}, \quad T \longmapsto E(T),$$

and hence, for every function into an additive commutative monoid,

$$\sum_{T \subseteq \{j \in \mathbb{N}_0 : j < m\}} f(E(T)) = \sum_{h=0}^{2^m-1} f(h). \quad (656)$$

These are `powersetRangeEquivFin` and `sum_powerset_two_pow`; the exact image statement is `image_sum_two_pow_powerset_range`. The singleton boundary at  $m = 0$  is included explicitly by `powersetRangeEquivFin_zero_apply`, all in `ThueMorseBitSupport.lean`.

The public Boolean-cube theorem uses the genuine degree  $\deg p \in \mathbb{N}_0 \cup \{-\infty\}$ , rather than the natural degree that assigns the zero polynomial degree zero. Over any commutative ring, for a finite set  $S$ , weights  $w_j$ , and a polynomial satisfying  $\deg p < |S|$ , it states

$$\sum_{T \subseteq S} (-1)^{|T|} p \left( x + \sum_{j \in T} w_j \right) = 0. \quad (657)$$

This is `sum_powerset_neg_one_pow_eval_of_degree_lt` in `ThueMorseSparseProuhet.lean`. The degree convention matters at the boundary  $S = \emptyset$ : the hypothesis then forces  $p = 0$ , so the theorem correctly includes the empty-support case instead of declaring it vacuous. With  $S$  the bit support of  $n$  and  $w_j = 2^j h$ , the corresponding sparse submask parametrization is

$$\sum_{T \subseteq \text{bits}(n)} (-1)^{|T|} p \left( x + h \sum_{j \in T} 2^j \right) = 0 \quad \text{when } \deg p < \text{wt}_2(n), \quad (658)$$

formalized as `sum_submask_neg_one_pow_eval_of_degree_lt`. It is valid also at  $n = 0$ , where again the only polynomial satisfying the strict degree bound is zero.

At the level of the centered formal exponential series, the corresponding zero has exact order  $r$ ; this is `order_thueMorseCenteredPowerSeries`. For the binomial kernel (645), the directly applicable formally proved form is instead the coefficient-extraction theorem `thueMorse_polynomial_sum_eq_coeff`, displayed in (649).

**Theorem 11.7** (Signed dyadic reciprocity). *Let  $k, r, d \in \mathbb{N}_0$  satisfy  $k \leq r$  and  $k + d < 2^r$ . Then the part of the order- $k$  prefix row immediately preceding its final  $k$  positions is the initial part of that row, read backwards and multiplied by one fixed sign:*

$$\sigma_k(2^r - k - 1 - d) = (-1)^{r-k} \sigma_k(d). \quad (659)$$

Consequently reflection preserves, rather than excludes, its zero locus:

$$\sigma_k(2^r - k - 1 - d) = 0 \iff \sigma_k(d) = 0. \quad (660)$$

The identity (659) and equivalence (660) are formalized respectively as `iteratedPrefix_dyadic_reverse_window` and `iteratedPrefix_dyadic_reverse_window_eq_zero_iff`. At  $k = 0$  the first identity is precisely the dyadic bit-complement law `thueMorseSign_dyadic_complement`. Its proof below uses the independently established left boundary in [theorem 11.8](#) as the seed for the prefix recurrence.

**Theorem 11.8** (Exact dyadic zero runs). *Let  $1 \leq k \leq r$ . Then*

$$\sigma_k(n) = 0 \quad \text{for } 2^r - k \leq n \leq 2^r - 1, \quad (661)$$

$$\sigma_k(2^r - k - 1) = (-1)^{r-k}, \quad (662)$$

$$\sigma_k(2^r) = -1. \quad (663)$$

Formula (662) also holds for  $k = 0$ , where it reduces to  $\varepsilon_{2^r-1} = (-1)^r$ , and (663) also holds for  $k = 0$ . Hence the zero run terminating at  $2^r - 1$  has length exactly  $k$ : it is bounded on the left by a value of modulus one and on the right by the value  $-1$ .

*Formal proof and Lean correspondence.* The terminal block is formalized by `iteratedPrefix_dyadic_zero_run`; its endpoint and reverse-indexed forms are `iteratedPrefix_dyadic_endpoint` and `iteratedPrefix_dyadic_zero_reverse`. Its left and right boundaries are `iteratedPrefix_before_dyadic_run` and `iteratedPrefix_at_dyadic`. All five declarations are in `ThueMorsePrefix.lean`. The mathematical argument behind them is as follows.

*Endpoint.* Put  $q = k - 1$ , so  $0 \leq q < r$ . By (646),

$$\sigma_k(2^r - 1) = \sum_{m=0}^{2^r-1} \binom{2^r-1-m+q}{q} \varepsilon_m = \frac{1}{q!} \sum_{m < 2^r} \varepsilon_m \prod_{i=1}^q (2^r - 1 - m + i).$$

The product is a polynomial in  $m$  of degree  $q < r$ , so the sum vanishes by (649).

*Rest of the run.* We prove, by induction on  $d$ , that  $\sigma_k(2^r - 1 - d) = 0$  whenever  $1 \leq k \leq r$  and  $0 \leq d < k$ . The case  $d = 0$  is the endpoint. Suppose the claim holds for  $d$  and that  $d + 1 < k$ , so  $k \geq 2$  and  $1 \leq k - 1 \leq r$ . The induction hypothesis applied at order  $k$  gives  $\sigma_k(2^r - 1 - d) = 0$ , and applied at order  $k - 1$ , which is legitimate because  $d < k - 1$ , gives  $\sigma_{k-1}(2^r - 1 - d) = 0$ . Now use (644) with  $n = 2^r - 1 - (d + 1)$ , so that  $n + 1 = 2^r - 1 - d$ :

$$0 - \sigma_k(n) = \sigma_k(n + 1) - \sigma_k(n) = \sigma_{k-1}(n + 1) = 0.$$

This proves (661), since the indices  $2^r - k, \dots, 2^r - 1$  are exactly the  $2^r - 1 - d$  with  $0 \leq d < k$ .

*Left boundary.* Induct on  $k$ . For  $k = 0$  the number  $2^r - 1$  has  $r$  binary digits, all equal to one, so  $\varepsilon_{2^r-1} = (-1)^r$ . Assume (662) at order  $k$  and let  $k + 1 \leq r$ . Put  $n = 2^r - (k + 1) - 1$ , so  $n + 1 = 2^r - k - 1$ . By the induction hypothesis  $\sigma_k(n + 1) = (-1)^{r-k}$ , and by (661) at order  $k + 1$  we have  $\sigma_{k+1}(n + 1) = \sigma_{k+1}(2^r - (k + 1)) = 0$ . Then (644) gives  $-\sigma_{k+1}(n) = (-1)^{r-k}$ , that is,  $\sigma_{k+1}(n) = (-1)^{r-(k+1)}$ .

*Right boundary.* Induct on  $k$ . For  $k = 0$ ,  $\text{wt}_2(2^r) = 1$ , so  $\varepsilon_{2^r} = -1$ . If (663) holds at order  $k$  and  $k + 1 \leq r$ , then by (644) and (661),  $\sigma_{k+1}(2^r) = \sigma_{k+1}(2^r - 1) + \sigma_k(2^r) = 0 - 1 = -1$ .  $\square$

*Proof of theorem 11.7.* Induct first on the prefix order  $k$ . At order zero,  $\sigma_0 = \varepsilon$ , and reflecting  $d$  in the block  $0, \dots, 2^r - 1$  complements its first  $r$  binary digits. Thus  $\varepsilon_{2^r-1-d} = (-1)^r \varepsilon_d$ , which is (659) for  $k = 0$ .

Now suppose the identity is known at order  $k$  and prove it at order  $k + 1$  by induction on  $d$ . For  $d = 0$ , equation (662) and  $\sigma_{k+1}(0) = 1$  give exactly the required identity. For the induction step, put

$$n = 2^r - (k + 1) - 1 - (d + 1), \quad s = (-1)^{r-(k+1)}.$$

The same-order induction hypothesis at offset  $d$  and the lower-order hypothesis at offset  $d + 1$  have the common reflected index  $n + 1$  and give

$$\sigma_{k+1}(n + 1) = s \sigma_{k+1}(d), \quad \sigma_k(n + 1) = (-1)^{r-k} \sigma_k(d + 1) = -s \sigma_k(d + 1).$$

Applying the forward-difference law (644) first at  $n$  and then at  $d$  therefore yields

$$\begin{aligned} \sigma_{k+1}(n) &= \sigma_{k+1}(n + 1) - \sigma_k(n + 1) \\ &= s(\sigma_{k+1}(d) + \sigma_k(d + 1)) = s \sigma_{k+1}(d + 1), \end{aligned}$$

which closes both inductions. Finally, the multiplier  $(-1)^{r-k}$  is nonzero, so (659) is equivalent to (660).  $\square$

*Remark 11.9.* At the level of statements, setting  $d = 0$  in (659) and using `iteratedPrefix_at_zero`, namely  $\sigma_k(0) = 1$ , recovers the left boundary (662). Reciprocity pairs any zeros that occur in the pre-run window; it does not claim that window is zero-free. Nor does it imply the terminal zero run (661) or the separate right boundary (663), both of which were proved independently above.

*Remark 11.10.* The right boundary value is not  $+1$ . Every iterated prefix sum, at every order  $k \leq r$ , takes the value  $-1$  at the index  $2^r$ . This single sign is what makes the literal normalization of section 11.5 fail at the right endpoint; see section 11.6.

### 11.3 Two generating-series identities

All series in this subsection are formal power series over  $\mathbb{Z}$ ; no convergence hypothesis on a complex variable is used or needed.

Let

$$\mathcal{K}_r(z) = \sum_{n < 2^r} \varepsilon_n z^n, \quad \mathcal{K}(z) = \sum_{n \geq 0} \varepsilon_n z^n, \quad \mathcal{S}_k(z) = \sum_{n \geq 0} \sigma_k(n) z^n. \quad (664)$$

**Proposition 11.11** (Block product and coefficient stabilization). *For every  $r \geq 0$ ,*

$$\mathcal{K}_r(z) = \prod_{j=0}^{r-1} (1 - z^{2^j}), \quad (665)$$

and, whenever  $n < 2^r$ , the coefficient of  $z^n$  in this finite product is  $\varepsilon_n$ . Hence each coefficient stabilizes once the finite product reaches a large enough dyadic block.

*Formal proof.* The finite product is `thueMorseBlockPolynomial_eq_product`; the stable coefficient identity is `coeff_finite_thueMorse_product`, both in `ThueMorseGenerating.lean`.  $\square$

**Theorem 11.12** (Prefix generating series). *For every  $k \geq 0$ ,*

$$\mathcal{S}_k(z) = \frac{\mathcal{K}(z)}{(1 - z)^k}, \quad (666)$$

or equivalently

$$(1 - z)^k \mathcal{S}_k(z) = \mathcal{K}(z). \quad (667)$$

For each fixed coefficient index, either side can be computed from a sufficiently long finite product (665).

*Formal proof.* The two identities are `iteratedPrefixSeries_eq` and `one_sub_X_pow_mul_iteratedPrefixSeries`; their finite coefficient bridge is `coeff_eq6_finite` in `ThueMorseGenerating.lean`.  $\square$

We use  $\prod_{j \geq 0} (1 - z^{2^j})$  only as shorthand for the coefficientwise stabilization of the finite products (665); no separate infinite-product object is asserted.

The two identities of [theorem 11.12](#) connect the prefix sums to the polynomial approximants used in [section 11.7](#). These are the combinatorial step polynomials already introduced in (110); their coefficient interpretation and exact degree were proved there. In the index convention of this section,

$$\mathcal{P}_n(z) = \prod_{i=1}^n (1 + z + \cdots + z^{2^i-1}), \quad g_n = \deg \mathcal{P}_n = \sum_{i=1}^n (2^i - 1) = 2^{n+1} - n - 2. \quad (668)$$

Since  $1 - z^{2^j} = (1 - z)(1 + z + \cdots + z^{2^j-1})$  and the factor with  $j = 0$  is trivial, (665) factors as

$$\mathcal{K}_{n+1}(z) = (1 - z)^{n+1} \mathcal{P}_n(z). \quad (669)$$

Comparing (669) with (667) at order  $k = n + 1$  and using stabilization below  $2^{n+1}$  gives the exact coefficient identification

$$\sigma_{n+1}(m) = [z^m] \mathcal{P}_n(z) \quad (0 \leq m < 2^{n+1}). \quad (670)$$

Note that (670) must fail at  $m = 2^{n+1}$ , because  $g_n < 2^{n+1}$  makes the right side zero while the left side is  $-1$  by (663). That single index is the exceptional right endpoint of [theorem 11.20](#).



### 11.4 The analytic Thue–Morse series and its natural boundary

The series  $\mathcal{K}$  in (664) was interpreted there as a formal power series. On the open unit disc the same coefficients define an analytic function, and we retain the notation  $\mathcal{K}$ :

$$\mathcal{K}(z) = \sum_{n=0}^{\infty} \varepsilon_n z^n, \quad |z| < 1. \quad (671)$$

**Proposition 11.13** (Analytic product and Mahler equation). *For every  $z \in \mathbb{C}$  with  $|z| < 1$ , the series (671) converges and*

$$\mathcal{K}(z) = \prod_{j=0}^{\infty} (1 - z^{2^j}). \quad (672)$$

The function  $\mathcal{K}$  is holomorphic on  $|z| < 1$  and satisfies

$$\mathcal{K}(z) = (1 - z)\mathcal{K}(z^2), \quad (673)$$

$$\mathcal{K}(z) = \left( \prod_{j=0}^{k-1} (1 - z^{2^j}) \right) \mathcal{K}(z^{2^k}) \quad (k \in \mathbb{N}_0). \quad (674)$$

On the real radius,

$$|\mathcal{K}(x)| \leq 1 - x \quad (0 \leq x < 1), \quad (675)$$

and  $\mathcal{K}(0) = 1$ .

*Formal proof.* The analytic function is `thueMorseDiscSeries`. Convergence, the product identity, and holomorphy are respectively `summable_thueMorseDiscSeries`, `thueMorseDiscSeries_eq_tprod`, and `thueMorseDiscSeries_differentiableOn`. Convergence of the product itself is `multipliable_one_sub_pow_two_pow_complex` in `ThueMorseInfiniteProduct.lean`. Equations (673) and (674) are `thueMorseDiscSeries_mahler` and `thueMorseDiscSeries_iterate`; (675) and the value at zero are `thueMorseDiscSeries_real_le` and `thueMorseDiscSeries_zero`. Except for the product-convergence declaration just cited, all these declarations are in `ThueMorseDiscSeries.lean`.  $\square$

**Theorem 11.14** (Dyadic radial zeros and the natural boundary). *If  $k \in \mathbb{N}_0$  and  $\zeta \in \mathbb{C}$  satisfy  $\zeta^{2^k} = 1$ , then*

$$\lim_{\rho \rightarrow 1^-} \mathcal{K}(\rho\zeta) = 0. \quad (676)$$

Moreover, for every  $z_0 \in \mathbb{C}$  with  $|z_0| = 1$  and every  $R > 0$ , there is no function  $g : \mathbb{C} \rightarrow \mathbb{C}$  which is holomorphic on  $\{z : |z - z_0| < R\}$  and satisfies

$$g(z) = \mathcal{K}(z) \quad \text{whenever} \quad |z - z_0| < R \quad \text{and} \quad |z| < 1. \quad (677)$$

Thus, in precisely this local-disc sense, every point of the unit circle is singular and the unit circle is a natural boundary for  $\mathcal{K}$ .

*Formal proof.* The radial limit is `tendsto_thueMorseDiscSeries_radial_zero`. More generally, if  $n \in \mathbb{N}_0$  is nonzero and  $\zeta^n = 1$ , then  $|\zeta| = 1$ ; this is `norm_eq_one_of_pow_eq_one`. The exact quantified nonextension statement (677), and hence the natural-boundary conclusion in the stated local-disc sense, is `thueMorse_natural_boundary`. All three declarations are in `ThueMorseNaturalBoundary.lean`.  $\square$



### 11.5 The functional equation of the normalized prefix grid

The natural dyadic normalization of the prefix sums is

$$\mathcal{V}_k(j) = \frac{\sigma_k(j)}{2^{\binom{k}{2}}}, \quad x_{k,j} = \frac{j}{2^k}, \quad (678)$$

so that  $\mathcal{V}_k$  is a rational-valued function on the level- $k$  dyadic grid. The exponent  $\binom{k}{2}$  is forced: it is the exponent that appears in the sharp Prouhet value (648), in the centered spline (120), and in the derivative scaling (262).

**Theorem 11.15** (Discrete functional equation). *Let  $q \geq 0$  and  $0 \leq j < 2^q$ , and put  $h = 2^{-(q+1)}$  and  $x = x_{q+1,j}$ . Then*

$$\frac{\mathcal{V}_{q+1}(j+1) - \mathcal{V}_{q+1}(j)}{h} = 2 \mathcal{V}_q(j+1), \quad (679)$$

and the abscissa attached to the right-hand value is

$$x_{q,j+1} = 2(x+h) = 2x_{q+1,j+1} \in [0, 1]. \quad (680)$$

The same identity holds verbatim for the shifted grid  $\tilde{\mathcal{V}}_k(j) = \sigma_{k+1}(j)/2^{\binom{k}{2}}$  of section 11.7.

*Proof.* Because  $\binom{q+1}{2} = \binom{q}{2} + q$ , the forward difference (644) gives

$$\mathcal{V}_{q+1}(j+1) - \mathcal{V}_{q+1}(j) = \frac{\sigma_{q+1}(j+1) - \sigma_{q+1}(j)}{2^{\binom{q}{2}+q}} = \frac{\sigma_q(j+1)}{2^{\binom{q}{2}}} \cdot \frac{1}{2^q} = \frac{\mathcal{V}_q(j+1)}{2^q}.$$

Dividing by  $h = 2^{-(q+1)}$  multiplies by  $2^{q+1}$  and gives (679). For (680),  $2(x+h) = 2(j+1)/2^{q+1} = (j+1)/2^q = x_{q,j+1}$ , and the hypothesis  $j < 2^q$  is exactly the condition  $j+1 \leq 2^q$  that keeps this abscissa in the unit interval. The shifted grid satisfies the same computation with  $\sigma$  replaced by  $\sigma_{\bullet+1}$  throughout.  $\square$

Equation (679) is the discrete counterpart of the global differential equation (261): a difference quotient at level  $q+1$  equals twice a level- $q$  value at the doubled abscissa. Two features of the discrete statement have no continuous analogue and are easy to lose. First, the right-hand side is evaluated at the *forward* node  $2(x+h)$  rather than at  $2x$ ; this is the inclusive-prefix convention (644). Second, the identity is an assertion about a level- $q$  value, so it is meaningful as a statement about  $F$  or up only when the doubled abscissa lies in the relevant interval, that is, only when  $j < 2^q$ . Both features are used in section 11.7.

### 11.6 The literal right endpoint does not converge

The unshifted normalization printed in the source formula cannot converge to a function whose right-endpoint value is 1: the sequence of literal values at the index  $2^k$  does not tend to 1. This exact negative statement is `paperPrefixGridValue_endpoint_not_tendsto_one` in `ThueMorseG` `enerating.lean`. The corrected normalization below is therefore a mathematical requirement, not a cosmetic reindexing.

### 11.7 Corrected normalization and pointwise convergence

We now identify the normalized prefix grid with the histogram approximants of the probabilistic construction, and deduce convergence. Two adjustments to the naive reading are required, and both are forced by identities already proved: the prefix order must be shifted by one, because (670) matches  $\sigma_{n+1}$  with  $\mathcal{P}_n$ ; and the grid index  $m$  must be read as labelling the *center* of a cell of width  $2^{-n}$ , not the left endpoint of a cell of width  $2^{-(n+1)}$ .

For the polynomial  $\mathcal{P}_n$  of (668), place the coefficient of the monomial  $z^m$  at

$$\xi_{n,m} = \frac{2m - g_n}{2^{n+1}}, \quad (681)$$

and spread it uniformly over the cell  $I_{n,m}$  of width  $2^{-n}$  centered at  $\xi_{n,m}$ . The consecutive centers are spaced  $2^{-n}$  apart, so these finitely many cells tile their support interval

$$\left[ -1 + \frac{n+1}{2^{n+1}}, 1 - \frac{n+1}{2^{n+1}} \right].$$

The resulting step approximant is

$$\mathcal{W}_n(x) = \frac{2^n}{2^{\binom{n+1}{2}}} \sum_{m=0}^{g_n} [z^m] \mathcal{P}_n(z) \cdot \mathbf{1}_{I_{n,m}}(x), \quad (682)$$

where a point shared by two adjacent cells is given the weight  $\frac{1}{2}$  in each, so that the two contributions add to the single value the formula is intended to have. This endpoint convention is not cosmetic: with ordinary closed-interval indicators the value at  $x = 0$  would be counted twice already for  $n = 1$ .

**Theorem 11.16** (Histogram approximants). *For every  $n \geq 0$  the function  $\mathcal{W}_n$  satisfies  $0 \leq \mathcal{W}_n \leq 1$ , is nondecreasing on  $(-\infty, 0)$ , is nonincreasing on  $(0, \infty)$ , and satisfies  $\mathcal{W}_n(0) = 1$ . Moreover  $\mathcal{W}_n(x) \rightarrow \text{up}(x)$  for every real  $x$ .*

*Proof.* For  $n \geq 1$ , this is exactly theorem 4.18, since (668) and (110) define the same polynomial, while (681) and (682) give the same cells and normalization as (113) and (114). At  $n = 0$  the empty product is  $\mathcal{P}_0 = 1$ ,  $g_0 = 0$ , and  $\mathcal{W}_0$  is the half-weighted indicator of  $[-1/2, 1/2]$ , so the stated order and normalization properties are immediate. The pointwise convergence of the sequence is already the final assertion of theorem 4.18.  $\square$

**Theorem 11.17** (Exact identification of the corrected grid). *Put  $\tilde{\mathcal{V}}_n(m) = \sigma_{n+1}(m)/2^{\binom{n}{2}}$ . For every  $n \geq 0$  and every  $m$  with  $0 \leq m < 2^{n+1}$ ,*

$$\tilde{\mathcal{V}}_n(m) = \mathcal{W}_n(\xi_{n,m}). \quad (683)$$

*Proof.* If  $m \leq g_n$ , the point  $\xi_{n,m}$  is the center of  $I_{n,m}$  and lies in no other cell, so (682) evaluates there to

$$\frac{2^n}{2^{\binom{n+1}{2}}} [z^m] \mathcal{P}_n(z).$$

Since  $\binom{n+1}{2} = \binom{n}{2} + n$ , this is  $[z^m] \mathcal{P}_n / 2^{\binom{n}{2}} = \tilde{\mathcal{V}}_n(m)$  by (670). If  $g_n < m < 2^{n+1}$ , then  $\xi_{n,m}$  lies strictly to the right of the finite support interval above, so  $\mathcal{W}_n(\xi_{n,m}) = 0$ ; meanwhile  $[z^m] \mathcal{P}_n = \sigma_{n+1}(m) = 0$  because  $\deg \mathcal{P}_n = g_n$ . Thus both sides again agree.  $\square$

Sampling at the dyadic floor now gives an approximation scheme. For  $x \in \mathbb{R}$  set

$$j_n(x) = \max(0, \lfloor 2^{n+1}x \rfloor), \quad \mathcal{J}_n(x) = \tilde{\mathcal{V}}_n(j_n(x)) = \frac{\sigma_{n+1}(j_n(x))}{2^{\binom{n}{2}}}. \quad (684)$$

**Lemma 11.18** (Moving cell centers). *For every  $x \in [0, 1)$ ,  $\xi_{n,j_n(x)} \rightarrow 2x - 1$  as  $n \rightarrow \infty$ .*

*Proof.* By (681),  $\xi_{n,j_n(x)} = 2 \cdot j_n(x) 2^{-(n+1)} - g_n 2^{-(n+1)}$ . The first term tends to  $2x$  because  $0 \leq 2^{n+1}x - j_n(x) < 1$ , and  $g_n 2^{-(n+1)} = 1 - (n+2)2^{-(n+1)} \rightarrow 1$ .  $\square$

**Lemma 11.19** (Stability under moving arguments). *If  $u_n \rightarrow y$  in  $\mathbb{R}$ , then  $\mathcal{W}_n(u_n) \rightarrow \text{up}(y)$ .*

*Proof.* Let  $\varepsilon > 0$ . Suppose first  $y > 0$ . By continuity of  $\text{up}$  choose  $d \in (0, y/2]$  with  $|\text{up}(y \pm d) - \text{up}(y)| < \varepsilon/2$ . For all large  $n$  we have  $u_n \in [y - d, y + d] \subset (0, \infty)$ , so the monotonicity of [theorem 11.16](#) on  $(0, \infty)$  gives

$$\mathcal{W}_n(y + d) \leq \mathcal{W}_n(u_n) \leq \mathcal{W}_n(y - d).$$

By pointwise convergence at the two fixed points  $y \pm d$ , for all large  $n$  both outer terms lie within  $\varepsilon/2$  of  $\text{up}(y \pm d)$ , hence within  $\varepsilon$  of  $\text{up}(y)$ . The case  $y < 0$  is identical using monotonicity on  $(-\infty, 0)$ . For  $y = 0$  the upper bound is  $\mathcal{W}_n(u_n) \leq 1 = \text{up}(0)$ , and the lower bound is  $\mathcal{W}_n(-d)$ ,  $\mathcal{W}_n(0) = 1$ , or  $\mathcal{W}_n(d)$  according to the sign of  $u_n$ ; each of the three converges to a value within  $\varepsilon/2$  of  $\text{up}(0)$ .  $\square$

**Theorem 11.20** (Corrected prefix approximation). *For every  $x \leq 1$ ,*

$$\mathcal{J}_n(x) \longrightarrow \text{up}(2x - 1), \quad (685)$$

*and on the whole nonpositive half-line the finite-stage values are exactly*

$$\mathcal{J}_n(x) = \mathcal{J}_n(0) = 2^{-\binom{n}{2}} \quad (x \leq 0). \quad (686)$$

*and consequently, for every  $x \in [0, 1]$ ,*

$$\mathcal{J}_n\left(\frac{x}{2}\right) \longrightarrow F(x). \quad (687)$$

*Proof.* For  $x \leq 0$ , the natural-floor clamp is exact: `correctedPrefixGridSample_eq_at_zero_of_nonpos` and `correctedPrefixGridSample_zero` give (686), while `correctedPrefixGridSample_tendsto_of_nonpos` gives the asserted limit. Now let  $0 < x < 1$ . Then  $2^{n+1}x < 2^{n+1}$ , so  $j_n(x) < 2^{n+1}$  and [theorem 11.17](#) applies:  $\mathcal{J}_n(x) = \mathcal{W}_n(\xi_{n,j_n(x)})$ . Combining [lemma 11.18](#) with [lemma 11.19](#) gives  $\mathcal{J}_n(x) \rightarrow \text{up}(2x - 1)$ .

At  $x = 1$  the identification fails, because  $j_n(1) = 2^{n+1}$  is exactly the first index not covered by (670). Here the value is computed instead from (663) with  $k = r = n + 1$ :

$$\mathcal{J}_n(1) = \frac{\sigma_{n+1}(2^{n+1})}{2^{\binom{n}{2}}} = -\frac{1}{2^{\binom{n}{2}}} \longrightarrow 0 = \text{up}(1),$$

since  $\binom{n}{2} \rightarrow \infty$  and  $\text{up}(1) = F(0) = 0$ . Altogether this is the public theorem `correctedPrefixGridSample_tendsto_rvachevUp_of_le_one` in `ThueMorseApproximation.lean`. For (687), apply it at  $x/2 \in [0, 1]$  and use  $2(x/2) - 1 = x - 1 \leq 0$ , so that  $\text{up}(x - 1) = F(x)$  by (7).  $\square$

**Warning 11.21.** [Theorem 11.20](#) is a pointwise statement and nothing more. No uniform rate is asserted, and none is proved. The proof passes through [lemma 11.19](#), whose only quantitative inputs are the shared monotonicity of the  $\mathcal{W}_n$  and the modulus of continuity of the limit  $\text{up}$  at the single point  $2x - 1$ ; the sandwich it uses degrades as that modulus degrades. A uniform bound of the kind supplied for the centered splines by (126) is available from [section 4.7](#), but it is a different approximation scheme, and the two should not be conflated.

**Remark 11.22.** The two corrections are independent of one another and both are necessary. Shifting the prefix order from  $k$  to  $k + 1$  is what makes (670) available; reading the index  $m$  as the center (681) of a cell of width  $2^{-n}$  rather than as an abscissa  $m2^{-k}$  is what supplies the affine change of variable  $x \mapsto 2x - 1$  between the unit interval and the support of  $\text{up}$ . In particular the corrected grid is a sample of the compactly supported bump, not of the distribution function; this is why its right-endpoint limit is 0 and not 1.

### 11.8 The lower Lambert branch

The asymptotic analysis of [section 12.5](#) is organized around the coordinate (730), where  $W_{-1}$  denotes the lower real branch of Lambert’s  $W$ -function [11]. That section, and the elementary expansions derived from it in [section 12.10](#), use three facts about  $W_{-1}$  without proving any of them: that the branch exists on  $(-e^{-1}, 0)$ , that it is characterized by being the solution below  $-1$ , and that it has the two-term expansion recorded below. We supply them here, since they are the foundational input for (732) and for every subsequent estimate in the Lambert variable.

**Proposition 11.23** (Existence, defining equation, and uniqueness). *For  $z \in (-e^{-1}, 0)$ , let  $W_{-1}(z)$  denote the totalized lower branch `lowerLambertW`. Then*

$$W_{-1}(z) < -1, \quad W_{-1}(z) e^{W_{-1}(z)} = z, \quad (688)$$

and  $W_{-1}(z)$  is the unique real  $w < -1$  with  $we^w = z$ .

*Formal proof.* These are respectively `lowerLambertW_lt_neg_one`, `lowerLambertW_mul_exp`, and `lowerLambertW_unique` in `LowerLambertW.lean`.  $\square$

On the smooth interior,  $W_{-1}$  is continuous and strictly decreasing, and its image is exactly  $(-\infty, -1)$ . At every  $z \in (-e^{-1}, 0)$ , it is differentiable with

$$\frac{d}{dz} W_{-1}(z) = \frac{W_{-1}(z)}{z(1 + W_{-1}(z))} < 0. \quad (689)$$

These assertions are `lowerLambertW_continuousOn`, `lowerLambertW_strictAntiOn`, `lowerLambertW_image`, `deriv_lowerLambertW`, and `deriv_lowerLambertW_neg`; the underlying derivative witness is `lowerLambertW_hasDerivAt`, all in `LowerLambertW.lean`.

The continuity statement now also includes the finite branch point: the raw lower branch is continuous on  $[-e^{-1}, 0)$ , by `lowerLambertW_continuousWithinAt_branchPoint` and `lowerLambertW_continuousOn_Ico`. Its principal companion has the corresponding right-continuity and full-domain continuity on  $[-e^{-1}, \infty)$ , namely `principalLambertW_continuousWithinAt_branchPoint` and `principalLambertW_continuousOn_Ici` in `PrincipalLambertW.lean`. At the regular point 0, that module also proves `deriv W_0(0) = 1` and  $W_0(z) \sim z$ , through `deriv_principalLambertW_zero` and `principalLambertW_isEquivalent_zero`. These endpoint-continuity declarations do not assert a derivative at the branch point. On the principal open domain, the nonsingular first-derivative identity is

$$W'_0(z) = \left[ e^{W_0(z)} (W_0(z) + 1) \right]^{-1}, \quad z > -e^{-1}. \quad (690)$$

The exact curvature formula is proved for both real branches. If  $j = 0$  and  $z > -e^{-1}$ , or if  $j = -1$  and  $-e^{-1} < z < 0$ , then

$$W''_j(z) = -\frac{e^{-2W_j(z)} (W_j(z) + 2)}{(W_j(z) + 1)^3}. \quad (691)$$

For the principal branch this is strictly negative everywhere on the open domain,  $W''_0(0) = -2$ , and  $W_0$  is strictly concave on the full closed natural domain  $[-e^{-1}, \infty)$ . The nonsingular first-derivative identity, second-derivative witness and formula, zero value, strict negativity, and strict concavity are respectively `deriv_principalLambertW`, `deriv_principalLambertW_hasDerivAt`, `deriv_deriv_principalLambertW`, `deriv_deriv_principalLambertW_zero`, `deriv_deriv_principalLambertW_neg`, and `strictConcaveOn_principalLambertW` in `LambertWCurvature.lean`.

For the lower branch, put  $z_* = -2e^{-2}$ , so that  $W_{-1}(z_*) = -2$ . Within  $(-e^{-1}, 0)$ ,

$$\begin{aligned} W_{-1}''(z) &> 0 &\iff z < z_*, \\ W_{-1}''(z) &= 0 &\iff z = z_*, \\ W_{-1}''(z) &< 0 &\iff z_* < z. \end{aligned} \tag{692}$$

Consequently  $W_{-1}$  is strictly convex on  $[-e^{-1}, z_*]$  and strictly concave on  $[z_*, 0)$ . The derivative witness and pointwise formula are `deriv_lowerLambertW_hasDerivAt` and `deriv_deriv_lowerLambertW`; the three equivalences are `deriv_deriv_lowerLambertW_pos_iff`, `deriv_deriv_lowerLambertW_eq_zero_iff`, and `deriv_deriv_lowerLambertW_neg_iff`; and the strict-shape statements are `strictConvexOn_lowerLambertW_left` and `strictConcaveOn_lowerLambertW_right`, all in `LambertWCurvature.lean`. The special value is `lowerLambertW_neg_two_mul_exp` in `LambertBranchDichotomy.lean`.

The strict-shape theorems include the finite branch point by continuity; they do not give a derivative or a second derivative there. The later branch-point leaves close the one-sided vertical geometry and the first square-root term, as follows.

### 11.8.1 The exact branch-gap coordinate and its symmetric laws

On the common open branch domain, put

$$-e^{-1} < x < 0, \quad a = W_0(x), \quad b = W_{-1}(x), \quad \delta = a - b > 0. \tag{693}$$

Then the two branches and their common argument are determined by the single positive coordinate  $\delta$ :

$$a = -\frac{\delta}{\exp(\delta) - 1}, \quad b = -\frac{\delta \exp(\delta)}{\exp(\delta) - 1} = -\frac{\delta}{1 - \exp(-\delta)}, \tag{694}$$

$$x = -\frac{\delta}{\exp(\delta) - 1} \exp\left(-\frac{\delta}{\exp(\delta) - 1}\right). \tag{695}$$

Conversely, every  $\delta > 0$  inserted into these formulas produces exactly one  $x \in (-e^{-1}, 0)$ , and the displayed  $a, b$  are respectively  $W_0(x), W_{-1}(x)$ . Thus  $x \mapsto W_0(x) - W_{-1}(x)$  is a bijection from  $(-e^{-1}, 0)$  onto  $(0, \infty)$ , with inverse given by (695). In the equivalent coordinate  $t = \exp(\delta) > 1$ ,

$$W_0(x) = -\frac{\log t}{t - 1}, \quad W_{-1}(x) = -\frac{t \log t}{t - 1}, \quad x = -\frac{\log t}{t - 1} t^{-1/(t-1)}. \tag{696}$$

Here is the elementary proof mirrored by the Lean decomposition. The two defining equations give  $a \exp(a) = b \exp(b)$ . Since neither branch value is zero, division yields  $b/a = \exp(a - b) = \exp(\delta)$ . Hence  $b = a \exp(\delta)$ , while  $\delta = a - b = a(1 - \exp(\delta))$ ; solving gives (694), and substitution into  $x = a \exp(a)$  gives (695). For the converse define  $a, b$  by the same formulas. The strict tangent inequality  $1 + \delta < \exp(\delta)$  gives  $-1 < a < 0$ , its negative-argument form gives  $b < -1$ , and direct algebra gives  $b = a - \delta$  and  $b \exp(b) = a \exp(a)$ . The two real-branch uniqueness theorems therefore identify these values with  $W_0, W_{-1}$ . They also prove both inverse laws, and hence the claimed bijection.

The same coordinate makes every symmetric identity immediate:

$$\frac{W_{-1}(x)}{W_0(x)} = \exp(\delta), \tag{697}$$

$$W_0(x) + W_{-1}(x) = -\delta \coth(\delta/2), \tag{698}$$

$$W_0(x)W_{-1}(x) = \frac{\delta^2}{4 \sinh^2(\delta/2)}. \tag{699}$$

Indeed, adding and multiplying (694) gives the exponential-rational forms; the definitions of  $\cosh$  and  $\sinh$  give (698)–(699). For  $y > 0$ ,  $\sinh y > y$  and  $(y \cosh y - \sinh y)' = y \sinh y > 0$ . With  $y = \delta/2$ , these facts give the strict interior bounds

$$W_0(x) + W_{-1}(x) < -2, \quad 0 < W_0(x)W_{-1}(x) < 1. \quad (700)$$

The four files have the following exhaustive public inventories. The forward file `LambertWBranchPairing.lean` has zero definitions and seven theorems: `principalLambertW_sub_lowerLambertW_pos`, `lowerLambertW_eq_principalLambertW_mul_exp_gap`, `principalLambertW_eq_neg_gap_div`, `lowerLambertW_eq_neg_gap_mul_exp_div`, `lowerLambertW_eq_neg_gap_div_one_sub_exp_neg`, `eq_neg_gap_div_mul_exp`, and `principalLambertW_lowerLambertW_eq_of_exp_gap`. The converse file `LambertWGapBijection.lean` has the four noncomputable definitions `gapPrincipal`, `gapLower`, `gapArg`, and `branchGap`, and exactly sixteen theorems: `gap_denominator_pos`, `gapPrincipal_mem_Ioo`, `gapLower_eq_mul_exp`, `gapLower_eq_sub`, `gapLower_lt_neg_one`, `gapLower_mul_exp`, `gapArg_mem_Ioo`, `principalLambertW_gapArg`, `lowerLambertW_gapArg`, `branchGap_gapArg`, `gapArg_branchGap`, `branchGap_invOn`, `branchGap_bijOn`, `principalLambertW_gapArg_log`, `lowerLambertW_gapArg_log`, and `gapArg_log`. Finally, `LambertWBranchSymmetry.lean` has zero definitions and nine theorems: `lowerLambertW_div_principalLambertW_eq_exp_branchGap`, `principalLambertW_add_lowerLambertW_eq_exp_branchGap`, `principalLambertW_add_lowerLambertW_eq_cosh_div_sinh_branchGap`, `principalLambertW_mul_lowerLambertW_eq_exp_branchGap`, `principalLambertW_mul_lowerLambertW_eq_sinh_sq_branchGap`, `principalLambertW_add_lowerLambertW_lt_neg_two`, `principalLambertW_mul_lowerLambertW_pos`, `principalLambertW_mul_lowerLambertW_lt_one`, and `principalLambertW_mul_lowerLambertW_mem_Ioo`. The final file `LambertWBranchGapBernoulli.lean` has zero definitions and exactly five theorems: `summable_norm_bernoulli_mul_pow_div_factorial`, `summable_bernoulli_mul_pow_div_factorial_iff`, `hasSum_bernoulli_mul_pow_div_factorial`, `hasSum_bernoulli_mul_pow_div_factorial_complex_iff`, and `principalLambertW_lowerLambertW_eq_bernoulliSeries`.

**Theorem 11.24** (Bernoulli exponential generating function in the gap coordinate). *For every  $w \in \mathbb{C}$ ,*

$$\sum_{n=0}^{\infty} \frac{B_n w^n}{n!} \text{ converges} \iff |w| < 2\pi.$$

*In particular, the series diverges at every point on or outside  $|w| = 2\pi$ . Writing*

$$E_{\exp}(w) = \begin{cases} (\exp(w) - 1)/w, & w \neq 0, \\ 1, & w = 0, \end{cases}$$

*the series has the canonical removable value*

$$\sum_{n=0}^{\infty} \frac{B_n w^n}{n!} = E_{\exp}(w)^{-1} \quad (|w| < 2\pi).$$

*For every real  $z$  with  $|z| < 2\pi$ ,*

$$\sum_{n=0}^{\infty} \left| \frac{B_n z^n}{n!} \right| < \infty. \quad (701)$$

*If also  $z \neq 0$ , then*

$$\sum_{n=0}^{\infty} \frac{B_n z^n}{n!} = \frac{z}{\exp(z) - 1}. \quad (702)$$

Consequently, for  $x \in (-e^{-1}, 0)$ , with  $\delta = W_0(x) - W_{-1}(x) < 2\pi$ ,

$$W_0(x) = - \sum_{n=0}^{\infty} \frac{B_n \delta^n}{n!}, \quad (703)$$

$$W_{-1}(x) = -\delta - \sum_{n=0}^{\infty} \frac{B_n \delta^n}{n!}. \quad (704)$$

*Formal coefficient bound and Cauchy-product proof.* Put  $q = |z|/(2\pi)$ , so  $0 \leq q < 1$ , and set

$$c_r = \frac{B_{2(r+1)}}{2(r+1)(2(r+1))!}.$$

The even-zeta estimate already proved for the Bernoulli logarithm gives

$$\left| c_r z^{2(r+1)} \right| \leq \frac{\pi^2}{6} q^{2(r+1)}.$$

All odd Bernoulli numbers beyond  $B_1$  vanish, while for the positive even indices

$$\left| \frac{B_{2(r+1)} z^{2(r+1)}}{(2(r+1))!} \right| = 2(r+1) \left| c_r z^{2(r+1)} \right| \leq 2(r+1) \frac{\pi^2}{6} q^{2(r+1)}.$$

Thus every term of index  $n \geq 2$  is bounded by  $n(\pi^2/6)q^n$ . The series  $\sum nq^n$  converges, and adding the two initial terms proves (701).

The same norm comparison at  $|w|$  proves convergence for complex  $w$  whenever  $|w| < 2\pi$ . Conversely, let

$$Z_{r+1} = \sum_{m=1}^{\infty} \frac{1}{m^{2(r+1)}} \geq 1.$$

The exact even-Bernoulli identity used in Lean gives

$$\left| \frac{B_{2(r+1)} w^{2(r+1)}}{(2(r+1))!} \right| = 2Z_{r+1} \left( \frac{|w|}{2\pi} \right)^{2(r+1)}.$$

If  $|w| \geq 2\pi$ , these even-index term norms are at least 2, so the terms do not tend to zero and the series cannot converge. This proves the exact complex convergence locus, including divergence on the boundary.

For the value of the sum away from the origin, the same argument works over  $\mathbb{C}$ . Write

$$f_n = \frac{B_n z^n}{n!}, \quad g_n = \frac{z^n}{n!} - \mathbf{1}_{\{n=0\}}.$$

The first sequence is absolutely summable by the preceding bound. The second is absolutely summable because the exponential series is absolutely summable and the subtracted sequence has finite support; its sum is  $\exp(z) - 1$ . Mathlib's formal power-series identity

$$\left( \sum_{n \geq 0} \frac{B_n}{n!} X^n \right) (\exp X - 1) = X$$

says coefficientwise that

$$\sum_{i+j=n} f_i g_j = \begin{cases} z, & n = 1, \\ 0, & n \neq 1. \end{cases}$$



Absolute summability therefore justifies the analytic Cauchy product and gives

$$\left( \sum_{n \geq 0} f_n \right) (\exp(z) - 1) = \sum_{n \geq 0} \sum_{i+j=n} f_i g_j = z.$$

When  $z \neq 0$ , the product identity itself makes  $\exp(z) - 1 \neq 0$ : a zero denominator would force  $z = 0$ . Division gives the complex quotient value. At  $z = 0$ , the Bernoulli series has only its constant term and sums to  $1 = E_{\exp}(0)^{-1}$ . Lean combines these cases with the exact convergence criterion in `hasSum_bernoulli_mul_pow_div_factorial_complex_iff`; restricting to real nonzero  $z$  proves (702). On the branch domain,  $\delta > 0$ , hence the extra hypothesis gives  $|\delta| < 2\pi$ . Substituting (702) into  $W_0 = -\delta/(\exp \delta - 1)$ , and then using  $W_{-1} = W_0 - \delta$ , proves (703)–(704). The five public declarations supporting these claims are respectively `summable_norm_bernoulli_mul_pow_div_factorial`, `summable_bernoulli_mul_pow_div_factorial_iff`, `hasSum_bernoulli_mul_pow_div_factorial`, `hasSum_bernoulli_mul_pow_div_factorial_complex_iff`, and `principalLambertW_lowerLambertW_eq_bernoulliSeries`.  $\square$

The forward pairing and symmetric identities have the strict input hypotheses in (693). The four converse-coordinate definitions are total, `gapLower_eq_mul_exp` is unconditional, the reconstruction theorems use  $\delta > 0$ , and the logarithmic-coordinate theorems use  $t > 1$ ; the inverse and bijection packages connect exactly the two open domains. At the branch point  $x = -e^{-1}$ , the gap is zero, the displayed quotients have zero denominators, and the limiting sum and product attain  $-2$  and  $1$ , so neither the parametrization as stated nor the strict inequalities include that endpoint. At  $x = 0$ , the classical lower branch has a singular extended endpoint, so it too is excluded; no claim is made about the all-real totalization there. The absolute-summability statement includes  $z = 0$ , but the quotient statement does not: its series has value  $B_0 = 1$  there, whereas Lean’s totalized  $0/(\exp 0 - 1)$  is zero. The complex `HasSum` equivalence gives the exact radius, proves divergence at every point on and outside  $|z| = 2\pi$ , and evaluates the series throughout the open disk as `complexExp1Div(z)^{-1}`. Thus the complex Bernoulli EGF is Exact under this canonical removable-origin convention, whose value at zero is one and which agrees with  $z/(\exp z - 1)$  away from zero; it is not a claim about Lean’s literal totalized quotient at zero, nor a holomorphy theorem for a named sum function. Its Lambert specialization still excludes both branch endpoints and additionally requires  $\delta < 2\pi$ . It is a convergent series in the gap coordinate, not a higher-order or full Puiseux expansion in  $x + e^{-1}$ , and it adds no logarithmic expansion at  $x = 0$ .

### 11.8.2 Branch-point vertical geometry and leading square-root law

Put

$$b = -\exp(-1), \quad \Delta(z) = z - b = z + \exp(-1), \quad \mathcal{S}(z) = \sqrt{2 \exp(1)(z + \exp(-1))}. \quad (705)$$

Every limit in this paragraph is one-sided, with  $z \downarrow b$ , equivalently along the filter of points strictly to the right of  $b$ . The two interior derivatives and the two endpoint secant slopes satisfy

$$W'_0(z) \longrightarrow +\infty, \quad W'_{-1}(z) \longrightarrow -\infty, \quad (706)$$

$$\frac{W_0(z) + 1}{z + \exp(-1)} \longrightarrow +\infty, \quad \frac{W_{-1}(z) + 1}{z + \exp(-1)} \longrightarrow -\infty. \quad (707)$$

Thus neither branch has a finite right derivative at  $b$ : formally, each fails `DifferentiableWithinAt` on the right-hand set  $(b, \infty)$ . Moreover, each totalized real branch fails to be differentiable at  $b$  as an all-real function.

The proof of (706) keeps the inverse derivative in the nonsingular form

$$W'_j(z) = \frac{1}{\exp(W_j(z))(W_j(z) + 1)}.$$



Endpoint continuity gives  $W_j(z) \rightarrow -1$ . On the principal branch the denominator therefore tends to zero through positive values, while on the lower branch it tends to zero through negative values; inversion gives the two signed infinite limits. Principal concavity gives  $W'_0(z) \leq (W_0(z) - W_0(b))/(z - b)$ , and lower convexity on the first shape interval gives  $(W_{-1}(z) - W_{-1}(b))/(z - b) \leq W'_{-1}(z)$ . These inequalities transfer the derivative limits to (707). Mathlib’s one-sided derivative-divergence obstruction then rules out a finite right derivative; ordinary differentiability would imply that forbidden within-set differentiability.

The exact eight-theorem crosswalk for this geometry layer is:

| Declaration in<br><code>LambertWBranchPointGeometry.lean</code>       | Formal content                                                |
|-----------------------------------------------------------------------|---------------------------------------------------------------|
| <code>tendsto_deriv_principalLambertW_branchPoint_atTop</code>        | Principal derivative tends to $+\infty$ from the right.       |
| <code>tendsto_deriv_lowerLambertW_branchPoint_atBot</code>            | Lower derivative tends to $-\infty$ from the right.           |
| <code>tendsto_principalLambertW_secantSlope_branchPoint_atTop</code>  | Principal endpoint secant slope tends to $+\infty$ .          |
| <code>tendsto_lowerLambertW_secantSlope_branchPoint_atBot</code>      | Lower endpoint secant slope tends to $-\infty$ .              |
| <code>principalLambertW_not_differentiableWithinAt_branchPoint</code> | No finite principal right derivative on $\text{loi}(b)$ .     |
| <code>lowerLambertW_not_differentiableWithinAt_branchPoint</code>     | No finite lower right derivative on $\text{loi}(b)$ .         |
| <code>principalLambertW_not_differentiableAt_branchPoint</code>       | The totalized principal branch is not differentiable at $b$ . |
| <code>lowerLambertW_not_differentiableAt_branchPoint</code>           | The totalized lower branch is not differentiable at $b$ .     |

For the first-order asymptotics, the forward map  $\phi(w) = w \exp w$  has a double contact at  $w = -1$ . A punctured-neighborhood L’Hôpital calculation gives the limit

$$\frac{\phi(w) + \exp(-1)}{(w + 1)^2} \longrightarrow \frac{\exp(-1)}{2} \quad (w \rightarrow -1, w \neq -1). \quad (708)$$

Indeed, after differentiating numerator and denominator, the quotient reduces away from  $-1$  to  $\exp(w)/2$ . Composing with either branch, using  $W_j(z) \exp(W_j(z)) = z$ , and then inverting the nonzero limit proves

$$\frac{(W_j(z) + 1)^2}{z + \exp(-1)} \longrightarrow 2 \exp(1), \quad j \in \{0, -1\}. \quad (709)$$

Equivalently, each squared displacement is asymptotic to  $2 \exp(1)(z + \exp(-1))$ . Finally, the strict signs  $W_0(z) + 1 > 0$  and  $W_{-1}(z) + 1 < 0$  select the two square roots:

$$W_0(z) + 1 \sim +\mathcal{S}(z), \quad W_{-1}(z) + 1 \sim -\mathcal{S}(z) \quad (z \downarrow -\exp(-1)). \quad (710)$$

The public inventory of `LambertWBranchPointAsymptotics.lean` is exactly one noncomputable definition and eight theorems:

| Declaration                                                        | Formal content                                                                      |
|--------------------------------------------------------------------|-------------------------------------------------------------------------------------|
| <code>lambertWBranchPointScale</code>                              | Definition of the exact scale<br>$\sqrt{2 \exp(1)(z + \exp(-1))}$ .                 |
| <code>lambertWBranchPointScale_pos</code>                          | Positivity of the scale for $b < z$ .                                               |
| <code>lambertWBranchPointScale_sq</code>                           | Exact square identity<br>$\mathcal{S}(z)^2 = 2 \exp(1)(z + \exp(-1))$ for $b < z$ . |
| <code>tendsto_principalLambertW_add_one_sq_div_branchPoint</code>  | Principal squared-ratio limit (709).                                                |
| <code>tendsto_lowerLambertW_add_one_sq_div_branchPoint</code>      | Lower squared-ratio limit (709).                                                    |
| <code>principalLambertW_add_one_sq_isEquivalent_branchPoint</code> | Principal squared-displacement asymptotic equivalence.                              |
| <code>lowerLambertW_add_one_sq_isEquivalent_branchPoint</code>     | Lower squared-displacement asymptotic equivalence.                                  |
| <code>principalLambertW_add_one_isEquivalent_branchPoint</code>    | The signed $+S$ principal law in (710).                                             |
| <code>lowerLambertW_add_one_isEquivalent_branchPoint</code>        | The signed $-S$ lower law in (710).                                                 |

The dependency order is exact. `PrincipalLambertW.lean` imports `LowerLambertW.lean`; `LambertBranchDichotomy.lean` imports the principal leaf; `LambertWCurvature.lean` imports that pair layer; `LambertWBranchPointGeometry.lean` imports the curvature leaf; and `LambertWBranchPointAsymptotics.lean` imports the geometry leaf together with `Mathlib`’s specific-asymptotics and `L’Hôpital` modules. The umbrella facade lists `LambertWCurvature.lean`, `LambertWBranchPointGeometry.lean`, and `LambertWBranchPointAsymptotics.lean` in this dependency order before the scaled power–exponential leaves.

This is a current-tree addendum compiled on 30 August 2026: its inventory is exactly eight geometry theorems and one asymptotic definition plus eight asymptotic theorems, with private proof helpers excluded. No aggregate commit hash is attributed to it. The result is only the leading square-root, or first Puiseux, equivalence. An  $O(\Delta)$  remainder, higher coefficients, and a convergent or full Puiseux expansion remain open. There are also no named generic power–exponential or Fabius endpoint vertical-tangent/secant/asymptotic wrappers yet, and the generic  $m \pm \sqrt{m}$  threshold and shape layer remains open.

**Corollary 11.25** (Solution of the saddle equation). *Let  $x > 0$  with  $Lx < e^{-1}$ , where  $L = \log 2$  as in (715), and put  $\lambda = -W_{-1}(-Lx)/L$  as in (731). Then  $\lambda 2^{-\lambda} = x$ , which is (732).*

*Formal proof.* The saddle equation is `paperLambertN_eq9` in `LowerLambertW.lean`. □

**Proposition 11.26** (Two-term expansion). *As  $\epsilon \downarrow 0$ ,*

$$W_{-1}(-\epsilon) - (\log \epsilon - \log |\log \epsilon|) \longrightarrow 0. \quad (711)$$

*Formal proof.* This is `tendsto_lowerLambertW_expansion` in `LowerLambertW.lean`. □

## 11.9 A binomial–logarithm formula for the Thue–Morse bit

The zero-one Thue–Morse bit is  $\tau(n) = \text{wt}_2(n) \bmod 2$ , so that  $\varepsilon_n = 1 - 2\tau(n)$ . Its primitive-recursive evaluator is `tmBitPR`; the theorems `tmBitPR_of_pos` and `tmBitPR_eq_thueMorseBit` prove its agreement with the binary-weight definition. The bit also admits a closed formula in which the binary structure is hidden inside a single row of Pascal’s triangle. Set

$$\mathcal{X}(n) = \sum_{k=0}^n (-1)^{\binom{n}{k}} \in \mathbb{Z}. \quad (712)$$

**Lemma 11.27** (Glaisher count). *The number of odd entries in the  $n$ th row of Pascal’s triangle is  $2^{\text{wt}_2(n)}$ .*

*Proof.* Let  $\nu(n)$  denote that number, so that  $\nu(n)$  is the number of nonzero coefficients of  $(1+z)^n$  in  $\mathbb{F}_2[z]$ . There  $(1+z)^2 = 1+z^2$ , hence

$$(1+z)^{2n} = (1+z^2)^n, \quad (1+z)^{2n+1} = (1+z^2)^n(1+z).$$

The substitution  $z \mapsto z^2$  does not change the number of nonzero coefficients, so  $\nu(2n) = \nu(n)$ ; and multiplying a polynomial in  $z^2$  by  $1+z$  produces two summands with disjoint supports, one even and one odd, so  $\nu(2n+1) = 2\nu(n)$ . Since  $\text{wt}_2(2n) = \text{wt}_2(n)$  and  $\text{wt}_2(2n+1) = \text{wt}_2(n) + 1$  by (642), and  $\nu(0) = 1 = 2^{\text{wt}_2(0)}$ , induction on the binary representation gives  $\nu(n) = 2^{\text{wt}_2(n)}$ .  $\square$

**Proposition 11.28** (The logarithm is exact). *For every  $n \geq 0$ ,*

$$n+1 - \mathcal{X}(n) = 2^{\text{wt}_2(n)+1}. \quad (713)$$

*In particular  $n+1 - \mathcal{X}(n)$  is a positive power of two, and  $\log_2(n+1 - \mathcal{X}(n)) = \text{wt}_2(n) + 1$  is an integer.*

*Proof.* Each summand of (712) is  $-1$  at an odd entry and  $+1$  at an even entry, and the row has  $n+1$  entries. Hence  $\mathcal{X}(n) = (n+1) - 2\nu(n)$  with  $\nu$  as in lemma 11.27, so  $n+1 - \mathcal{X}(n) = 2\nu(n) = 2^{\text{wt}_2(n)+1}$ .  $\square$

**Theorem 11.29** (Binomial–logarithm formula). *For every  $n \geq 0$ ,*

$$\tau(n) = \frac{1 + (-1)^{\log_2\left(n+1 - \sum_{k=0}^n (-1)^{\binom{n}{k}}\right)}}{2}, \quad \varepsilon_n = 1 - 2\tau(n). \quad (714)$$

*Proof.* By proposition 11.28 the exponent equals  $\text{wt}_2(n) + 1$ . If  $\text{wt}_2(n)$  is even then  $\tau(n) = 0$  and  $(-1)^{\text{wt}_2(n)+1} = -1$ , so the right side is 0; if  $\text{wt}_2(n)$  is odd then  $\tau(n) = 1$  and  $(-1)^{\text{wt}_2(n)+1} = 1$ , so the right side is 1. The second identity is  $\varepsilon_n = (-1)^{\text{wt}_2(n)} = 1 - 2(\text{wt}_2(n) \bmod 2)$ .  $\square$

*Remark 11.30.* The order of the two steps matters. As printed, (714) raises  $-1$  to a logarithm, an operation that is meaningful only because the argument of that logarithm has first been proved to be exactly a power of two. The formal development therefore proves (713) before it interprets the outer expression at all, evaluates the logarithm on the natural numbers, and reads the final division in  $\mathbb{Q}$ . A floating-point evaluation of the same display, or an interpretation of the logarithm as a real-valued function whose output is then rounded, is an approximation to a formula rather than the formula.

## 12 Endpoint asymptotics: the exact formal hierarchy

The behavior of  $F(x)$  as  $x \downarrow 0$  is governed by three nested structures: an exact dyadic product on the negative Laplace axis, a genuinely nonconstant periodic correction on logarithmic scale, and a lower-Lambert saddle coordinate. This section states the hierarchy in the form proved in Lean. Longer heuristic derivations, numerical experiments, and proposed refinements without exact theorem counterparts are collected in docs/semi-formalized-research-frontiers/semi-formalized-research-frontiers.tex.

Throughout, put

$$L = \log 2. \quad (715)$$

### 12.1 The negative-Laplace product and its exact decomposition

For  $s > 0$ , the entire generating function  $G$  from (389) satisfies

$$G(-s) = \prod_{j=1}^{\infty} \frac{1 - \exp(-s/2^j)}{s/2^j}. \quad (716)$$

This is the real specialization of `complexGeneratingFunction_neg_eq_tprod` in `NegativeLaplaceVertical.lean`. If

$$\kappa(u) = \log\left(\frac{1 - \exp(-u)}{u}\right), \quad \Lambda(s) = \log G(-s) = \sum_{j=1}^{\infty} \kappa(s/2^j),$$

then the series is absolutely summable for every  $s > 0$ ; the precise declarations are `summable_negativeLaplaceTerm`, `exp_negativeLaplaceLog_eq_generatingFunction_neg`, and `negativeLaplaceLog_eq_log_laplaceMoment`.

Let  $P$  denote `negativeLaplacePeriodicCorrection` and let  $E$  denote `negativeLaplaceTailError`. The exact decomposition is

$$\Lambda(s) = -\frac{(\log s)^2}{2L} + \frac{\log s}{2} + P(\log_2 s) + E(s) \quad (s > 0). \quad (717)$$

Moreover,

$$0 \leq E(s) \leq \frac{\exp(-s)}{(1 - \exp(-s))^2}, \quad E(s) \leq 4 \exp(-s) \quad (s \geq L). \quad (718)$$

These are exactly `negativeLaplaceLog_exact_periodic_decomposition`, `negativeLaplaceTailError_nonneg`, `abs_negativeLaplaceTailError_le`, and `abs_negativeLaplaceTailError_le_four_exp` in `NegativeLaplace.lean`. The correction  $P$  is  $C^\infty$  and one-periodic by `contDiff_infty_negativeLaplacePeriodicCorrection` and `negativeLaplacePeriodicCorrection_add_one`.

Write

$$\overline{P} = \int_0^1 P(u) \, du, \quad \Psi(t) = P(t) - \overline{P}. \quad (719)$$

Then  $\Psi$  is  $C^\infty$ , one-periodic, and has mean zero:

$$\Psi(t+1) = \Psi(t), \quad \int_0^1 \Psi(t) \, dt = 0.$$

The formal counterparts are `contDiff_infty_negativeLaplacePsi`, `negativeLaplacePsi_add_one`, and `integral_negativeLaplacePsi_zero`.

The first logarithmic derivative has a correspondingly sharp full-ray expansion. As  $s \rightarrow +\infty$ ,

$$\Lambda'(s) = \frac{-\log s/L + 1/2 + \Psi'(\log_2 s)/L}{s} + \mathcal{O}(s^{-2}). \quad (720)$$

Here  $\Lambda'$  is the formal `negativeLaplaceLogFirst`, identified with the derivative of  $\Lambda$  by `negativeLaplaceLog_hasDerivAt`. The exact  $\mathcal{O}(s^{-2})$  statement is `negativeLaplaceLogFirst_sub_periodic_main_isBigO_inv_sq_real` in `LaplacePeriodicSecondOrder.lean`. This promotes the proved derivative estimate without importing the longer unformalized contour and Mellin derivations from the frontier volume.

## 12.2 The Gamma–zeta Fourier series

For  $k \in \mathbb{Z}$ , use the Fourier convention

$$\widehat{\Psi}_{\text{FS}}[k] = \int_0^1 \Psi(t) \exp(-2\pi i k t) dt, \quad \chi_k = \frac{2\pi i k}{L}. \quad (721)$$

The zero mode vanishes, and every nonzero mode has the closed form

$$\widehat{\Psi}_{\text{FS}}[0] = 0, \quad \widehat{\Psi}_{\text{FS}}[k] = -\frac{\Gamma(-\chi_k)\zeta(1-\chi_k)}{L} \quad (k \neq 0). \quad (722)$$

Every coefficient with  $k \neq 0$  is nonzero, so  $\Psi$  is genuinely nonconstant. The coefficient sequence is absolutely summable and reconstructs  $\Psi$  pointwise:

$$\Psi(t) = \sum_{k \in \mathbb{Z}} \widehat{\Psi}_{\text{FS}}[k] \exp(2\pi i k t). \quad (723)$$

These assertions are `negativeLaplacePsiFourierCoeff_zero`, `negativeLaplacePsiFourierCoeff_eq_neg_gamma_zeta`, `negativeLaplacePsiFourierCoeff_ne_zero`, `negativeLaplacePsi_not_constant`, `summable_negativeLaplacePsiGammaZetaFourierCoeff`, and `hasSum_negativeLaplacePsi_gammaZeta_fourierSeries` in `PeriodicFourier.lean`. No numerical truncation or stronger coefficient-decay claim is needed here.

## 12.3 The evaluated constant

Let  $\gamma$  be Euler’s constant and let  $\gamma_1$  denote the formally defined first Stieltjes constant `firstStieltjesConstant`. The regularized Gamma–zeta combination

$$A_0 = \frac{\gamma^2}{2} + \gamma_1 - \frac{\pi^2}{12}$$

is `gammaZetaConstant`; its equivalent common-denominator form is proved by `gammaZetaConstant_eq_div_twelve`. The split, convergent finite-part Bose integral is evaluated as this same constant by `boseFinitePartIntegral_eq_gammaZetaConstant`.

The formal finite part is independent of where the convergent integral is split. More precisely, put

$$\kappa_0(x) = \frac{1}{x} \log\left(\frac{1 - \exp(-x)}{x}\right), \quad \kappa_\infty(x) = \frac{1}{x} \log(1 - \exp(-x)).$$

For every  $c > 0$ , both integrals below are convergent and

$$\boxed{\int_{(0,c]} \kappa_0(x) dx + \int_{(c,\infty)} \kappa_\infty(x) dx + \frac{(\log c)^2}{2} = A_0.} \quad (724)$$

If  $0 < a \leq b$ , changing the splitting point is governed by the exact transition identity

$$\int_{(a,b]} (\kappa_\infty(x) - \kappa_0(x)) dx = \frac{(\log b)^2 - (\log a)^2}{2}. \quad (725)$$

Here the two kernels are `boseFinitePartSmallKernel` and `boseFinitePartLargeKernel`; their integrability is `integrableOn_boseFinitePartSmallKernel_Ioc` and `integrableOn_boseFinitePartLargeKernel_Ioi_of_pos`. Equations (724) and (725) are respectively `boseFinitePartIntegral_eq_gammaZetaConstant_of_pos` and `integral_boseFinitePartLargeKernel_sub_small_Ioc` in `BoseFinitePartIntegral.lean`. Thus the constant is represented by ordinary convergent real integrals, without assigning a value to a divergent integral.

In turn, `negativeLaplacePeriodicMean_eq` gives

$$\bar{P} = \frac{A_0}{L} - \frac{L}{12}.$$

After the Gaussian normalization, the sharp Fabius constant is

$$C_F = \frac{6\gamma^2 + 12\gamma_1 - \pi^2}{12L} - \frac{7L}{12} - \frac{1}{2} \log \pi. \quad (726)$$

This is the exact identity `fabiusSharpAsymptoticConstant_eq` in `FabiusSharpConstant.lean`.

## 12.4 A fully effective dyadic bound

The dyadic specialization admits a proved numerical threshold and error constant. For  $n \geq 1$ , define

$$\begin{aligned} D_n = & -\frac{L}{2}n^2 - n \log n + \left(1 + \frac{L}{2}\right)n - \frac{(\log n)^2}{2L} + C_F + \Psi(\log_2 n) \\ & - \frac{n}{2}(\Lambda''(n) + \Lambda'(n)^2). \end{aligned} \quad (727)$$

Here the final term is exactly `dyadicEndpointSecondOrder`; the derivative identifications are `negativeLaplaceLog_hasDerivAt` and `negativeLaplaceLogFirst_hasDerivAt`. Then, for every  $n \geq 224043$ ,

$$\boxed{|\log F(2^{-n}) - D_n| \leq \frac{2512945}{12n}}. \quad (728)$$

This is `abs_log_fabius_dyadic_sub_explicitCumulantMain_le` in `FabiusDyadicSharpCumulant.lean`. The threshold is a sufficient certified threshold, not an optimality claim. Its proof uses the pointwise bounds `dyadicEndpointSecondOrder_sq_le`, `dyadicHigherLaplaceMoments_le`, and `abs_dyadicEndpointLaplaceLogError_add_secondOrder_le_unconditional` in `LaplaceMomentBounds.lean`, with respective constants  $256/n$ ,  $104448/n$ , and  $209408/n$  on their stated ranges.

## 12.5 The lower-Lambert saddle and exact reduction

On the endpoint-inclusive lower-branch domain

$$\mathcal{D}_L = \left(0, \frac{e^{-1}}{L}\right], \quad (729)$$

let

$$W = W_{-1}(-Lx), \quad (730)$$

and define

$$\lambda = \lambda(x) = -\frac{W}{L}, \quad r = r(x) = 2^\lambda. \quad (731)$$

The global Lean definitions are `fabiusLambertPhase` and `fabiusLambertRadius`; here we use them on  $\mathcal{D}_L$ . For every  $x \in \mathcal{D}_L$ , their exact identities are

$$\lambda 2^{-\lambda} = x, \quad \frac{\lambda}{r} = x, \quad rx = \lambda, \quad \log r = L\lambda. \quad (732)$$

The endpoint-inclusive quotient and product identities are `fabiusLambertPhase_div_radiu s_of_le` and `fabiusLambertRadius_mul_argument_of_le`; the first equality in (732) is the quotient identity rewritten using  $r = 2^\lambda$ . The logarithmic identity in fact holds for every real  $x$ ,

not only on  $\mathcal{D}_L$ ; this is `log_fabiusLambertRadius`. The corresponding strict-interior wrappers are `fabiusLambertPhase_div_radius` and `fabiusLambertRadius_mul_argument`. All five declarations are in `FabiusLambertSaddle.lean`.

This lower phase is also the exact specialization at  $(m, A, \beta) = (1, 1, L)$  of the generic scaled power–exponential inverse; this is `lowerPowerExponentialPhase_one_one_log_two` in `PowerExponentialLambertFabius.lean`. More generally, for  $m \in \mathbb{N}_{>0}$ ,  $A > 0$ , and  $\beta > 0$ , put

$$S_{m,A,\beta}(t) = At^m e^{-\beta t}, \quad P_{m,A,\beta} = A \left( \frac{m}{\beta} e^{-1} \right)^m.$$

Writing  $\lambda_0$  and  $\lambda_{-1}$  for the principal and lower Lambert phases, their exact images are

$$\lambda_0([0, P_{m,A,\beta}]) = \left[ 0, \frac{m}{\beta} \right], \quad \lambda_{-1}((0, P_{m,A,\beta}]) = \left[ \frac{m}{\beta}, \infty \right).$$

These are `principalPowerExponentialPhase_image_Icc` and `lowerPowerExponentialPhase_image_Ioc`; the companion declarations `principalPowerExponentialPhase_invOn` and `lowerPowerExponentialPhase_invOn` make each phase a two-sided `InvOn` inverse of  $S_{m,A,\beta}$  on the corresponding displayed interval pair. They are proved in `PowerExponentialLambertInverse.lean`. On the principal branch, the closed endpoint step uses the exact restriction  $W_0([-e^{-1}, 0]) = [-1, 0]$ , namely `principalLambertW_image_Icc` in `PrincipalLambertW.lean`.

The same hypotheses give continuity on the full natural profile-value domains, including the common finite branch point:

$$\lambda_0 \in C([0, P_{m,A,\beta}]), \quad \lambda_{-1} \in C((0, P_{m,A,\beta}]).$$

These are `principalPowerExponentialPhase_continuousOn_Icc` and `lowerPowerExponentialPhase_continuousOn_Ioc` in `PowerExponentialLambertCalculus.lean`; their proof composes the raw endpoint APIs above with the globally continuous normalized argument `powerExponentialLambertArgument_continuous`. Moreover, for  $x \in (0, P_{m,A,\beta}]$  and  $\mu \geq 0$ , Lean proves the exact dichotomy

$$S_{m,A,\beta}(\mu) = x \iff \mu = \lambda_0(x) \text{ or } \mu = \lambda_{-1}(x). \quad (733)$$

This is `powerExponentialSaddle_eq_iff_eq_principal_or_eq_lower`; on the strict interior  $0 < x < P_{m,A,\beta}$ , the two roots are unequal by `principalPowerExponentialPhase_ne_lowerPowerExponentialPhase`, both in `PowerExponentialLambertInverse.lean`. The nonnegativity assumption on  $\mu$  is essential: for even  $m$ , negative inputs can create additional positive profile values.

On that same strict profile-value interval, both inverse branches have the exact second derivative

$$\lambda_j''(x) = \frac{\lambda_j(x) [m - (m - \beta \lambda_j(x))^2]}{x^2 (m - \beta \lambda_j(x))^3}, \quad j \in \{0, -1\}. \quad (734)$$

The corresponding derivative witnesses are `deriv_principalPowerExponentialPhase_hasDerivAt` and `deriv_lowerPowerExponentialPhase_hasDerivAt`; the pointwise identities are `deriv_deriv_principalPowerExponentialPhase` and `deriv_deriv_lowerPowerExponentialPhase`, all in `PowerExponentialLambertCurvature.lean`. That generic module stops at the exact formula: it does not yet package the thresholds  $m \pm \sqrt{m}$ , their zero/sign classifications, or generic strict-convexity and strict-concavity corollaries.

The compiled small-input package uses the intrinsic positive scale

$$\epsilon_{m,A,\beta}(x) = \frac{\beta}{m} \left( \frac{x}{A} \right)^{1/m}. \quad (735)$$

With the same parameter hypotheses, as  $x \downarrow 0$ ,

$$\lambda_0(x) \sim \left(\frac{x}{A}\right)^{1/m}, \quad (736)$$

$$\lambda_{-1}(x) \longrightarrow +\infty, \quad (737)$$

$$\lambda_{-1}(x) - M_{m,A,\beta}(x) \longrightarrow 0, \quad (738)$$

where

$$M_{m,A,\beta}(x) = -\frac{m}{\beta} (\log \epsilon_{m,A,\beta}(x) - \log |\log \epsilon_{m,A,\beta}(x)|).$$

The exact Lean crosswalk is `powerExponentialLambertEpsilon`, `principalPowerExponentialPhase_isEquivalent_rpow`, `tendsto_lowerPowerExponentialPhase_nhdsGT_zero_atTop`, `lowerPowerExponentialPhaseIntrinsicMain`, and `lowerPowerExponentialPhase_sub_intrinsicMain_tendsto_zero` in `PowerExponentialLambertAsymptotics.lean`. This generic theorem stops at the intrinsic  $\epsilon$ -form with an  $o(1)$  remainder. A cleaned formula in  $\log(A/x)$ , further logarithmic terms, and a full series are not part of this package. The separate interior curvature formula (734) does not itself supply named generic-phase branch-point vertical-tangent, secant-slope, or leading square-root wrappers; the raw-branch results above have not yet been transported through the scaled coordinate.

For the classical parameters  $(m, A, \beta) = (1, 1, L)$ , the principal root is `fabiusPrincipalLambertPhase` and the lower root is `fabiusLambertPhase`. The specialized wrappers `fabiusSaddle_eq_iff_eq_principal_or_eq_lower`, `fabiusPrincipalLambertPhase_ne_fabiusLambertPhase`, `fabiusPrincipalLambertPhase_continuousOn_Icc`, and `fabiusLambertPhase_continuousOn_Ioc` give respectively the nonnegative-root dichotomy, strict-interior separation, and the two full-domain continuity statements. Finally `fabiusPrincipalLambertPhase_isEquivalent_id` gives  $\lambda_0(x) \sim x$  as  $x \downarrow 0$ . These declarations are in `PowerExponentialLambertFabius.lean`; the generic lower divergence and intrinsic expansion above specialize to the same lower Fabius phase.

On the same closed-upper domain, the phase has the sharp lower bound and endpoint characterization

$$\lambda(x) \geq \frac{1}{L}, \quad \lambda(x) = \frac{1}{L} \iff x = \frac{e^{-1}}{L}. \quad (739)$$

In particular,

$$\lambda\left(\frac{e^{-1}}{L}\right) = \frac{1}{L}. \quad (740)$$

Moreover,  $\lambda$  is strictly decreasing on  $\mathcal{D}_L$ , and its exact image is

$$\lambda(\mathcal{D}_L) = \left[\frac{1}{L}, \infty\right). \quad (741)$$

The bound, equality characterization, endpoint value, strict antitonicity, and image are respectively `fabiusLambertPhase_ge_inv_log_two`, `fabiusLambertPhase_eq_inv_log_two_iff`, `fabiusLambertPhase_branchPoint`, `fabiusLambertPhase_strictAntiOn_Ioc`, and `fabiusLambertPhase_image_Ioc` in `FabiusLambertSaddle.lean`.

Equivalently, under the hypotheses  $x > 0$  and  $Lx \leq e^{-1}$ , the same lower bound and the strict positivity  $0 < \lambda(x)$  are `one_div_log_two_le_fabiusLambertPhase` and `fabiusLambertPhase_pos_of_le`. Thus the algebraic saddle identities and continuity extend to the finite branch point even though the differentiable formulas below do not.

The differentiable API is deliberately restricted to the open interior  $0 < x < e^{-1}/L$ . There  $\lambda$  is strictly decreasing, its image is  $(1/L, \infty)$ , and it is differentiable with

$$\lambda'(x) = \frac{\lambda(x)}{x(1 - L\lambda(x))} < 0. \quad (742)$$



The interior continuity wrapper is `fabiusLambertPhase_continuousOn`; the remaining claims are `fabiusLambertPhase_strictAntiOn`, `fabiusLambertPhase_image`, `deriv_fabiusLambertPhase`, and `deriv_fabiusLambertPhase_neg` in `FabiusLambertSaddle.lean`; the derivative witness is `fabiusLambertPhase_hasDerivAt`, and pointwise continuity is `fabiusLambertPhase_continuousAt`.

The full curvature profile is also exact. Define

$$x_* := \frac{2e^{-2}}{L}. \quad (743)$$

Then  $0 < x_* < e^{-1}/L$ ,  $\lambda(x_*) = 2/L$ , and for  $0 < x < e^{-1}/L$ ,

$$\lambda''(x) = \frac{L\lambda(x)^2(2 - L\lambda(x))}{x^2(1 - L\lambda(x))^3}. \quad (744)$$

Moreover,

$$\begin{aligned} \lambda''(x) > 0 &\iff x < x_*, \\ \lambda''(x) = 0 &\iff x = x_*, \\ \lambda''(x) < 0 &\iff x_* < x. \end{aligned} \quad (745)$$

Thus the lower phase is strictly convex on  $(0, x_*]$  and strictly concave on  $[x_*, e^{-1}/L]$ . In Lean,  $x_*$  is `fabiusLambertInflectionInput`; its domain membership and phase value are `fabiusLambertInflectionInput_mem_Ioo` and `fabiusLambertPhase_inflectionInput`. The second derivative, its three sign loci, its exact zero, and the two strict-shape statements are respectively `deriv_deriv_fabiusLambertPhase`, `deriv_deriv_fabiusLambertPhase_pos_iff`, `deriv_deriv_fabiusLambertPhase_eq_zero_iff`, `deriv_deriv_fabiusLambertPhase_neg_iff`, `deriv_deriv_fabiusLambertPhase_inflectionInput`, `strictConvexOn_fabiusLambertPhase_left`, and `strictConcaveOn_fabiusLambertPhase_right` in `PowerExponentialLambertFabiusCurvature.lean`.

The principal Fabius phase has a complementary global result. For every real  $x$ ,

$$\lambda_0(x) = -\frac{W_0(-Lx)}{L}. \quad (746)$$

It is continuous on  $(-\infty, e^{-1}/L]$ ; for every  $x < e^{-1}/L$ ,

$$\lambda'_0(x) = \left[ e^{W_0(-Lx)} (W_0(-Lx) + 1) \right]^{-1}, \quad (747)$$

$$\lambda''_0(x) = \frac{Le^{-2W_0(-Lx)}(W_0(-Lx) + 2)}{(W_0(-Lx) + 1)^3} > 0. \quad (748)$$

In particular  $\lambda''_0(0) = 2L$ , and  $\lambda_0$  is strictly convex on the whole closed half-line  $(-\infty, e^{-1}/L]$ . The exact affine formula and continuity theorem are `fabiusPrincipalLambertPhase_eq_principalLambertW` and `fabiusPrincipalLambertPhase_continuousOn_Iic`. The first-derivative witness and identity are `fabiusPrincipalLambertPhase_hasDerivAt` and `deriv_fabiusPrincipalLambertPhase`; the second-derivative witness, identity, positivity, zero value, and whole-half-line strict convexity are `deriv_deriv_fabiusPrincipalLambertPhase_hasDerivAt`, `deriv_deriv_fabiusPrincipalLambertPhase`, `deriv_deriv_fabiusPrincipalLambertPhase_pos`, `deriv_deriv_fabiusPrincipalLambertPhase_zero`, and `strictConvexOn_fabiusPrincipalLambertPhase`, all in `PowerExponentialLambertFabiusCurvature.lean`.

Returning to the lower phase, at the small-argument endpoint  $x = 0$ , the phase and its reciprocal have the exact limits

$$\lambda(x) \longrightarrow +\infty, \quad \lambda(x)^{-1} \longrightarrow 0 \quad (x \downarrow 0). \quad (749)$$

These are `tendsto_fabiusLambertPhase_nhdsGT_zero_atTop` and `tendsto_inv_fabiusLambertPhase_nhdsGT_zero` in `FabiusLambertRates.lean`.

### 12.5.1 Phase-locked reciprocal Richardson algebra

The exact inverse relation also supplies a finite lattice on which every periodic coefficient is frozen, not merely asymptotically stable. Fix  $\lambda_0 > 1/L$  and set

$$x_j = (\lambda_0 + j)2^{-(\lambda_0+j)}, \quad h_j = (\lambda_0 + j)^{-1}. \quad (750)$$

Then  $x_j \in \mathcal{D}_L^\circ$  and

$$\lambda(x_j) = \lambda_0 + j. \quad (751)$$

Consequently every one-periodic function  $G$  satisfies  $G(\lambda(x_j)) = G(\lambda_0)$ .

For  $0 \leq j \leq r$ , let  $\omega_{r,j}$  be the evaluation-at-zero Lagrange weight on  $h_0, \dots, h_r$ . Its explicit real form is

$$\omega_{r,j} = \frac{(-1)^{r-j}}{r!} \binom{r}{j} (\lambda_0 + j)^r. \quad (752)$$

These weights obey

$$\sum_{j=0}^r \omega_{r,j} = 1, \quad (753)$$

$$\sum_{j=0}^r \omega_{r,j} (\lambda_0 + j)^{-m} = 0 \quad (1 \leq m \leq r), \quad (754)$$

$$\sum_{j=0}^r \omega_{r,j} G(\lambda(x_j)) = G(\lambda_0). \quad (755)$$

Every residual moment is explicit as well. For  $t \geq 0$ ,

$$\sum_{j=0}^r \omega_{r,j} (\lambda_0 + j)^{-(r+1+t)} = (-1)^r \left( \prod_{j=0}^r (\lambda_0 + j)^{-1} \right) h_t(h_0, \dots, h_r). \quad (756)$$

The node and reciprocal-row definitions are `lambertPhaseLockedNode`, `shiftedReciprocalNode`, and `shiftedReciprocalLagrangeWeight`. Node-domain membership, phase locking, and periodic preservation are `lambertPhaseLockedNode_mem_lowerBranch`, `fabiusLambertPhase_phaseLockedNode`, `Periodic.apply_fabiusLambertPhase_phaseLockedNode`, and `sum_shiftedReciprocalLagrangeWeight_mul_periodicPhaseLocked`. The weight formula, mass-one identity, cancellation, and all-residual identity are `shiftedReciprocalLagrangeWeight_eq_choose_of_pos`, `sum_shiftedReciprocalLagrangeWeight_eq_one`, `sum_shiftedReciprocalLagrangeWeight_mul_invPow_eq_zero`, and `sum_shiftedReciprocalLagrangeWeight_residual` in `LambertPhaseLockedRichardson.lean`.

This is exact finite interpolation algebra. It does not by itself extract the endpoint fluctuation  $\Psi$  from sampled values of  $F$ , justify applying an asymptotic expansion uniformly across the row, or prove a Richardson remainder estimate. Such analytic steps must be supplied separately.

The corrected main term is

$$\mathcal{M}(x) = -\frac{L}{2} \lambda(x)^2 + \left(1 + \frac{L}{2}\right) \lambda(x) - \frac{1}{2} \log \lambda(x) + C_F + \Psi(\lambda(x)). \quad (757)$$

This is definition `fabiusSharpLambertMain`. The exact saddle action itself is endpoint-inclusive: whenever  $x > 0$  and  $Lx \leq e^{-1}$ ,

$$r(x)x + \Lambda(r(x)) - \frac{1}{2} \log(2\pi\lambda(x)) = \mathcal{M}(x) + E(r(x)). \quad (758)$$

This is `fabiusLambertSaddleAction_eq_of_le` in `FabiusSharpLambertMain.lean`. If  $\mathcal{R}(x)$  denotes the normalized saddle ratio `fabiusSaddleRatio` at  $r(x)$  and  $\lambda(x)$ , then Lean proves the exact error separation

$$\log F(x) - \mathcal{M}(x) = \log \mathcal{R}(x) + E(r(x)). \quad (759)$$

The declaration is `log_fabius_sub_sharpLambertMain_eq_log_ratio_add_tail` in `FabiusSharpExactReduction.lean`.

The forward product tail in this separation is negligible at every algebraic order of the exact phase. Along  $x = 2^{-t}$ , for each  $N \in \mathbb{N}_0$ ,

$$E(r(2^{-t})) = o(\lambda(2^{-t})^{-N}) \quad (t \rightarrow +\infty). \quad (760)$$

This is `negativeLaplaceTailError_dyadicLambert_isLittleO_inv_pow` in `FabiusSharpLambertTransfer.lean`. At  $N = 0$  it says directly that the tail tends to zero; larger  $N$  show that it contributes to none of the inverse-phase coefficients.

The analytic input to that reduction is an exact Bromwich formula. For every  $r > 0$  and  $x \in \mathbb{R}$ , as an identity in  $\mathbb{C}$ ,

$$F(x) = \exp(rx) \int_{\mathbb{R}} \exp(2\pi i \xi x) \frac{G(-(r + 2\pi i \xi))}{r + 2\pi i \xi} d\xi. \quad (761)$$

This is `fabius_bromwich`; the dimensionless, Gaussian-normalized version used by the saddle machinery is `fabius_bromwich_scaled`. The supporting Laplace-transform identity is `generatingFunction_neg_div_eq_laplace`. All three occur in `LaplaceTransform.lean` and `FabiusBromwichInput.lean` with the positivity and integrability hypotheses shown there.

## 12.6 The compact online formula and its missing periodic factor

A January 2022 comment in a Mathematics Stack Exchange discussion asked whether the preceding calculation implied a particularly compact lower-Lambert formula for the endpoint behavior [20]. With the notation of (730), its logarithmic main term is

$$\mathcal{W}_{\text{MSE}}(x) = -\frac{7L}{12} - \frac{1}{2} \log(\pi x) + \frac{1}{L} \left( \gamma_1 + \frac{\gamma^2}{2} - \frac{\pi^2}{12} - W - \frac{W^2}{2} \right). \quad (762)$$

This is the human-readable form of definition `fabiusWikipediaLambertMain` in `FabiusWikipediaMain.lean`. Written in ordinary mathematical notation, the exponential of this logarithmic encoding is the expression printed in the discussion:

$$A_{\text{MSE}}(x) = \frac{1}{2^{7/12} \sqrt{\pi x}} \exp \left[ \frac{1}{L} \left( \gamma_1 + \frac{\gamma^2}{2} - \frac{\pi^2}{12} - W - \frac{W^2}{2} \right) \right]. \quad (763)$$

For  $x > 0$ , this factor is strictly positive. Lean names it `fabiusWikipediaLambertFactor`; `exp_fabiusWikipediaLambertMain_eq_WikipediaLambertFactor` identifies it with  $\exp(\mathcal{W}_{\text{MSE}}(x))$ , and `fabiusWikipediaLambertFactor_pos` proves its positivity. All three declarations occur in `FabiusWikipediaMain.lean`. The proposed symbol  $F(x) \sim A_{\text{MSE}}(x)$  means that the ratio should tend to 1 as  $x \downarrow 0$ .

That ratio-one claim is false. The answer on the cited page first derives a Gamma–zeta periodic term, then explicitly omits it from the subsequent approximation while noting that it must be retained for more precise asymptotics. The exact formal decomposition shows what is lost: on the positive lower-Lambert domain,

$$\mathcal{M}(x) = \mathcal{W}_{\text{MSE}}(x) + \Psi(\lambda(x)). \quad (764)$$

Here `fabiusCorrectedWikipediaMain_eq_WikipediaLambertMain_add` is the all-real identity between the two named compact definitions, while `fabiusCorrectedWikipediaMain_eq_sharpLambertMain` identifies the corrected one with  $\mathcal{M}$  on the stated domain. Consequently the proved ratio-one asymptotic is

$$F(x) \sim \frac{1}{2^{7/12}\sqrt{\pi x}} \exp\left[\frac{1}{L}\left(\gamma_1 + \frac{\gamma^2}{2} - \frac{\pi^2}{12} - W - \frac{W^2}{2}\right)\right] \times \exp(\Psi(-W/L)) \quad (x \downarrow 0). \quad (765)$$

This is the direct theorem `fabius_isEquivalentent_WikipediaLambertFactor_mul_periodic` in `FabiusSharpAsymptotic.lean`; the underlying exact positive-side factorization is `exp_fabiusCorrectedWikipediaMain_eq_WikipediaLambertFactor_mul_periodic` in `FabiusWikipediaMain.lean`. The plus sign in (764) is essential: the missing multiplier is  $\exp(\Psi(\lambda(x)))$ , with  $\lambda(x) = -W/L$ .

The logical bridge from logarithmic error to ratio-one equivalence is itself formalized. Along any filter, if  $f$  is eventually positive, then

$$f \sim \exp(g) \iff \log f - g \longrightarrow 0. \quad (766)$$

This is `isEquivalentent_exp_iff_tendsto_log_sub`. In the present case Lean proves both

$$\log F(x) - \mathcal{W}_{\text{MSE}}(x) \not\rightarrow 0 \quad (x \downarrow 0) \quad (767)$$

and that the same residual is not  $\mathcal{O}((-\log x)^{-1})$ . These are respectively `log_fabius_sub_WikipediaLambertMain_not_tendsto_zero` and `log_fabius_sub_WikipediaLambertMain_not_isBigO`. Applying (766) gives the direct theorem

$$F \not\sim A_{\text{MSE}}, \quad (768)$$

namely `fabius_not_isEquivalentent_exp_WikipediaLambertMain`; its direct literal-factor form is `fabius_not_isEquivalentent_WikipediaLambertFactor`. The criterion and all four obstruction declarations in this paragraph occur in `FabiusSharpAsymptotic.lean`. Thus the nonconvergence of the sampled periodic correction, rather than its numerical size, is decisive for ratio-one equivalence.

**Theorem 12.1** (Exact cluster interval for the online quotient). *For  $x > 0$ , put*

$$R_{\text{MSE}}(x) := \frac{F(x)}{A_{\text{MSE}}(x)}, \quad m_{\Psi} := \min_{0 \leq t \leq 1} \Psi(t), \quad M_{\Psi} := \max_{0 \leq t \leq 1} \Psi(t).$$

*Then  $m_{\Psi} < M_{\Psi}$ , and the quotient has the sharper additive asymptotic model*

$$R_{\text{MSE}}(x) - \exp(\Psi(\lambda(x))) \longrightarrow 0 \quad (x \downarrow 0). \quad (769)$$

*Write  $\text{Cl}_{0+}(R_{\text{MSE}})$  for the set of all  $z \in \mathbb{R}$  for which there is a sequence  $x_n \rightarrow 0$ , eventually positive, such that  $R_{\text{MSE}}(x_n) \rightarrow z$ . Its complete cluster set is*

$$\text{Cl}_{0+}(R_{\text{MSE}}) = [\exp(m_{\Psi}), \exp(M_{\Psi})]. \quad (770)$$

*In particular,*

$$\liminf_{x \downarrow 0} R_{\text{MSE}}(x) = \exp(m_{\Psi}) < \exp(M_{\Psi}) = \limsup_{x \downarrow 0} R_{\text{MSE}}(x). \quad (771)$$

*Consequently no real constant repairs the omitted factor: in Mathlib's filter-theoretic asymptotic-equivalence relation,*

$$\forall c \in \mathbb{R}, \quad F \not\sim c A_{\text{MSE}} \quad (x \downarrow 0). \quad (772)$$

*For  $c \neq 0$ , the last assertion is the usual failure of  $F(x)/(c A_{\text{MSE}}(x)) \rightarrow 1$ . The formal declaration also includes  $c = 0$ , without interpreting a quotient by zero.*

The online discussion supplies  $A_{\text{MSE}}$ ; the exact cluster interval is a consequence of the corrected formal asymptotic. The endpoint descriptions  $\exp(m_\Psi)$  and  $\exp(M_\Psi)$  are exact variational formulas in terms of the explicit periodic correction. No finite elementary evaluation, numerical certification, sign claim for the extrema, or uniqueness claim for their phases is asserted here.

The phase-sampling mechanism is reusable. Every value of a one-periodic map  $p$  is a cluster value of  $p(\lambda(x))$  as  $x \downarrow 0$ ; if  $p$  is continuous and its codomain is Hausdorff, the complete cluster set is exactly  $p(\mathbb{R})$ . These statements are `mapClusterPt_periodic_comp_fabiusLambertPhase` and `mapClusterPt_periodic_comp_fabiusLambertPhase_iff` in `FabiusWikipediaObstruction.lean`.

For the specialization above,  $R_{\text{MSE}}$  is `fabiusWikipediaLambertRatio`, while  $m_\Psi$  and  $M_\Psi$  are `negativeLaplacePsiPeriodMin` and `negativeLaplacePsiPeriodMax`. The exact global range, pointwise extremal bounds, and endpoint order are respectively `range_negativeLaplacePsi_eq_Icc_periodMin_periodMax`, `negativeLaplacePsiPeriodMin_le_value`, `negativeLaplacePsi_value_le_periodMax`, and `negativeLaplacePsiPeriodMin_le_periodMax`; attainment and strict separation are `exists_mem_Icc_negativeLaplacePsi_eq_periodMin`, `exists_mem_Icc_negativeLaplacePsi_eq_periodMax`, and `negativeLaplacePsiPeriodMin_lt_periodMax`. Exponentiation of the exact range is `range_exp_negativeLaplacePsi_eq_Icc_periodExtrema`.

The ratio-one comparison with the sampled periodic model is `fabiusWikipediaLambertRatio_isEquivalent_periodic`, and the stronger additive statement (769) is `tendsto_fabiusWikipediaLambertRatio_sub_periodic`. The interval identity (770) is `mapClusterPt_fabiusWikipediaLambertRatio_nhdsGT_zero_iff`; the two endpoint limits are `liminf_fabiusWikipediaLambertRatio_nhdsGT_zero_eq` and `limsup_fabiusWikipediaLambertRatio_nhdsGT_zero_eq`; and the universal obstruction is `fabius_not_isEquivalent_const_mul_WikipediaLambertFactor`. These declarations all occur in `FabiusSharpAsymptotic.lean`.

## 12.7 The first two periodic saddle corrections

Define

$$A_1(u) = \frac{1}{24} + \frac{1}{2L} - \frac{\Psi''(u) + \Psi'(u)^2}{2L^2}. \quad (773)$$

This is `fabiusFirstSaddleCorrection`. It is continuous, one-periodic, and bounded by `continuous_fabiusFirstSaddleCorrection`, `fabiusFirstSaddleCorrection_periodic`, and `isBounded_range_fabiusFirstSaddleCorrection`. The first corrected expansion is

$$\boxed{\log F(x) = \mathcal{M}(x) + \frac{A_1(\lambda(x))}{\lambda(x)} + \mathcal{O}(\lambda(x)^{-2})} \quad (x \downarrow 0). \quad (774)$$

It is the  $N = 2$  specialization of `log_fabius_sub_sharpLambertExpansion_isBigO`, together with `fabiusSharpLambertExpansion_two`.

The second coefficient has a compact closed form in the bounded exponent jets. For each  $m$ , write  $J_m(u)$  for the jet `negativeLaplaceBoundedExponentJet(m, u)`, and define

$$A_2(u) = \frac{1}{24} (8J_0^3 + 12J_0^2J_1 + 12J_0^2 + 48J_0J_1 + 12J_0J_2 + 6J_1^2 + 24J_1 + 20J_2 + 3J_3)(u). \quad (775)$$

This is exactly `fabiusSecondSaddleCorrection`; it is continuous, one-periodic, and bounded by `continuous_fabiusSecondSaddleCorrection`, `fabiusSecondSaddleCorrection_periodic`, and `isBounded_range_fabiusSecondSaddleCorrection`. The formally proved

second-correction theorem is

$$\log F(x) = \mathcal{M}(x) + \frac{A_1(\lambda(x))}{\lambda(x)} + \frac{A_2(\lambda(x))}{\lambda(x)^2} + \mathcal{O}(\lambda(x)^{-3}) \quad (x \downarrow 0). \quad (776)$$

Its exact declarations are `fabiusSharpLambertExpansion_three` and `log_fabius_sub_twoSaddleCorrections_isBigO` in `FabiusSecondSaddleExpansion.lean`; the companion theorem `log_fabius_sub_twoSaddleCorrections_isBigO_negLog` gives the literal remainder  $\mathcal{O}((-\log x)^{-3})$ .

## 12.8 The recursive all-orders theorem

For  $j \in \mathbb{N}_0$ , let  $a_j(u)$  denote `fabiusSaddleLogCoefficient( $j, u$ )`. Every  $a_j$  is continuous, one-periodic, and bounded, by `continuous_fabiusSaddleLogCoefficient`, `fabiusSaddleLogCoefficient_periodic`, and `isBounded_range_fabiusSaddleLogCoefficient`. The initial identifications are

$$a_0 = 0, \quad a_1 = A_1, \quad a_2 = A_2,$$

proved by `fabiusSaddleLogCoefficient_zero`, `fabiusSaddleLogCoefficient_one_eq_firstSaddleCorrection`, and `fabiusSaddleLogCoefficient_two_eq_secondSaddleCorrection`.

For every  $N \in \mathbb{N}_0$ , the complete exact-phase expansion is

$$\log F(x) = \mathcal{M}(x) + \sum_{j=0}^{N-1} \frac{a_j(\lambda(x))}{\lambda(x)^j} + \mathcal{O}(\lambda(x)^{-N}) \quad (x \downarrow 0). \quad (777)$$

The precise formal statement is `log_fabius_sub_sharpLambertMain_hasAsymptoticExpansion`; its finite-remainder forms are `log_fabius_sub_sharpLambertExpansion_isBigO` and `log_fabius_sub_sharpLambertExpansion_isBigO_negLog` in `FabiusFullAsymptoticExpansion.lean`.

The coefficients are constructive rather than existential. The formal development identifies the jet weights as coefficients of

$$\prod_{k=0}^{n-1} (X - (k+1)),$$

proves closed forms for the periodic and bounded exponent jets, and gives recursive exponent-polynomial identities. The exact declarations are `negativeLaplaceJetPolynomial_eq_prod`, `negativeLaplaceJetStirling_eq_coeff`, `negativeLaplacePeriodicJet_eq_closedForm`, `negativeLaplaceBoundedExponentJet_eq_closedForm`, `negativeLaplaceExponentCoefficient_succ_eq`, and `negativeLaplaceExponentPolynomial_succ_eq`. This coefficient-of-a-product description is the proved statement; no identification with an external Stirling-number API is asserted here.

The algebraic conversion between relative-mass and logarithmic coefficients is normalized at every index rather than only asymptotically. If `ExpCoeff` and `LogCoeff` denote the formal recurrences `expCoeff` and `logCoeff`, then

$$\begin{aligned} \text{ExpCoeff}(\text{LogCoeff}(a))_n &= \begin{cases} 1, & n = 0, \\ a_n, & n > 0, \end{cases} \\ \text{LogCoeff}(\text{ExpCoeff}(E))_n &= \begin{cases} 0, & n = 0, \\ \text{Expectation}_n, & n > 0. \end{cases} \end{aligned} \quad (778)$$

These are `expCoeff_logCoeff_eq_ite` and `logCoeff_expCoeff_eq_ite`; the corresponding identities of formal power series are `expSeries_logCoeff_eq_massSeries_normalize_zero` and `logSeries_expCoeff_eq_exponentSeries_normalize_zero` in `SaddleLogExpansionAlgebra.lean` and `SaddleLogExpansionPowerSeries.lean`.

The recurrence-defined logarithm is also multiplicative at the level of formal power series. Let  $R$  be a commutative  $\mathbb{Q}$ -algebra and define

$$\mathfrak{L}(C) := \text{SaddleExpansion.logOf}(C), \quad C \in R[[X]],$$

by applying `logSeries` to the coefficient family of  $C$ . Rebuilding a series from that family gives  $C$  itself by `SaddleExpansion.massSeries_coeff`; this reconstruction lemma needs only that  $R$  be a commutative ring. For every  $C$ , even before a unit-constant hypothesis is imposed,

$$[X^0]\mathfrak{L}(C) = 0, \tag{779}$$

which is `SaddleExpansion.constantCoeff_logOf`. If  $[X^0]C = 1$ , then

$$C \mathfrak{L}(C)' = C', \tag{780}$$

by `SaddleExpansion.mul_derivative_logOf`; conversely, `SaddleExpansion.logOf_eq_of` says that (779) and (780) characterize  $\mathfrak{L}(C)$  uniquely.

It follows that unit-constant series satisfy

$$\mathfrak{L}(AB) = \mathfrak{L}(A) + \mathfrak{L}(B), \quad \mathfrak{L}(1) = 0, \tag{781}$$

and, for every finite set  $S$  and family with  $[X^0]F_i = 1$ ,

$$\mathfrak{L}\left(\prod_{i \in S} F_i\right) = \sum_{i \in S} \mathfrak{L}(F_i). \tag{782}$$

These are `SaddleExpansion.logOf_mul`, `SaddleExpansion.logOf_one`, and `SaddleExpansion.logOf_prod` in `LogSeriesMultiplicative.lean`. They are identities in  $R[[X]]$ : no convergence, analytic logarithm, branch choice, or passage to an infinite product is asserted.

## 12.9 Universal finite endpoint-transfer algebra

The formal coefficient stage of endpoint transfer is independent of every probability and asymptotic estimate. To distinguish the universal endpoint polynomials from the periodic correction  $P$ , write  $P_r^{\text{end}}(X) = \text{endpointTransferPolynomial}(r)$ . In  $\mathbb{Q}[X]$ , define

$$e_0(X) = 0, \quad e_{n+1}(X) = -\frac{X^{n+2}}{n+2}, \quad P_r^{\text{end}}(X) = \text{ExpCoeff}(e)_r. \tag{783}$$

Then the recursive, formal-series, and unordered-partition descriptions agree:

$$P_{n+1}^{\text{end}}(X) = -\frac{1}{n+1} \sum_{j=0}^n \frac{j+1}{j+2} X^{j+2} P_{n-j}^{\text{end}}(X), \tag{784}$$

$$\sum_{r \geq 0} P_r^{\text{end}}(X) t^r = \exp\left(\sum_{r \geq 0} e_r(X) t^r\right) = \exp\left(-\sum_{j \geq 2} \frac{X^j t^{j-1}}{j}\right), \tag{785}$$

$$P_r^{\text{end}}(X) = \sum_{f \in \mathcal{P}_r} \prod_{j=1}^r \frac{e_j(X)^{f(j)}}{f(j)!},$$

$$\mathcal{P}_r = \left\{ f : \mathbb{N}_0 \rightarrow \mathbb{N}_0 \mid \supp(f) \subseteq \{1, \dots, r\}, \sum_{j=1}^r j f(j) = r \right\}. \tag{786}$$

The first coefficients are

$$P_0^{\text{end}} = 1, \quad P_1^{\text{end}} = -\frac{X^2}{2}, \quad P_2^{\text{end}} = \frac{X^4}{8} - \frac{X^3}{3}, \quad P_3^{\text{end}} = -\frac{X^6}{48} + \frac{X^5}{6} - \frac{X^4}{4}. \quad (787)$$

Because  $\mathbb{Q}[X]$  is the free rational algebra on one generator, evaluating  $X$  at any element  $z$  of a commutative  $\mathbb{Q}$ -algebra commutes both with each  $P_r^{\text{end}}$  and with the complete formal power series.

*Formal proof.* The definitions are `endpointTransferExponentCoefficient`, `endpointTransferPolynomial`, and `endpointTransferSeries` in `EndpointTransferPolynomials.lean`. Equations (784)–(786) are respectively `endpointTransferPolynomial_succ`, `endpointTransferSeries_eq_exp_subst`, and `endpointTransferPolynomial_eq_partitionExpSum`. The finite partition engine and its equality with the recursive exponential coefficients are `weightedPartitions`, `partitionExpSum`, and `partitionExpSum_eq_expCoeff` in `ExponentialPartition.lean` and `ExponentialBell.lean`. Evaluation and series naturality are `aeval_endpointTransferPolynomial` and `map_endpointTransferSeries`; the four identities in (787) are `endpointTransferPolynomial_zero`, `endpointTransferPolynomial_one`, `endpointTransferPolynomial_two`, and `endpointTransferPolynomial_three`.

All statements in this subsection are finite or formal. They do not identify the series analytically with  $\exp(X)(1 - Xt)^{1/t}$ , construct a tilted law, or prove a Taylor-window, tail, or endpoint-remainder estimate.  $\square$

## 12.10 Recursive elementary re-expansion of the phase

The exact Lambert phase can itself be expanded to arbitrary order. With

$$T(x) = -\log_2 x,$$

write  $\mathcal{L}_N(x)$  for `fabiusLambertLogarithmicApproximation(N, x)`. This approximation is built from the recursively defined polynomials `dyadicLambertDisplacementPolynomial`. For every  $N \in \mathbb{N}_0$ ,

$$\lambda(x) - \mathcal{L}_N(x) = \mathcal{O}(T(x)^{-N}) \quad (x \downarrow 0). \quad (788)$$

This is `fabiusLambertPhase_sub_logarithmicApproximation_isBigO`; the equivalent dyadic-scale form is `dyadicLambertAllOrderRemainder_isBigO`. Explicit coefficient polynomials are proved through index two, while higher orders are represented by the formal recursion.

The periodic term cannot simply be discarded from a sharp estimate. If  $\mathcal{M}_{\text{Wiki}}$  denotes the uncorrected elementary expression `fabiusWikipediaElementaryMain`, then Lean proves that  $x \mapsto \log F(x) - \mathcal{M}_{\text{Wiki}}(x)$  is not  $\mathcal{O}((-\log x)^{-1})$  as  $x \downarrow 0$ . This is `log_fabius_sub_WikipediaElementaryMain_not_isBigO` in `FabiusSharpAsymptotic.lean`. The obstruction is therefore a theorem, not a numerical diagnosis.

## 12.11 Three endpoint consequences already formalized

On the dyadic logarithmic scale, let  $g(t) = -\log F(2^{-t})$ . Then

$$\lim_{t \rightarrow \infty} \frac{g(t)}{t^2} = \frac{L}{2}. \quad (789)$$

This is `fabiusLogProfile_normalized_tendsto` in `FabiusLogSquaredAsymptotic.lean`. On the original small-argument scale, for every  $c > 0$ ,

$$\exp(-c/x) = o(F(x)) \quad (x \downarrow 0). \quad (790)$$

This is `exp_neg_div_isLittleO_fabius` in `FabiusDecayComparison.lean`. At the opposite endpoint, the exact global symmetry already proved in (15) gives only the identity

$$1 - F(1 - x) = F(x). \quad (791)$$

No additional right-endpoint asymptotic-transfer theorem is asserted here.



### 13 The Fourier–Legendre expansion of up

Let  $P_n$  be the Legendre polynomial normalized by  $P_n(1) = 1$ , and define

$$u_n = \frac{4n+1}{2} \int_{-1}^1 \text{up}(x) P_{2n}(x) dx. \quad (792)$$

Odd Legendre coefficients vanish by `rvachevFullLegendreCoefficient_odd_eq_zero`. The even coefficients have the exact finite formula

$$u_n = 4^{-n}(4n+1) \sum_{k=0}^n (-1)^{n+k} \binom{2n}{n+k} \binom{2n+2k}{2n} \\ \times (2k)! 2^{\binom{2k+1}{2}} F(2^{-2k-1}). \quad (793)$$

This is `rvachevLegendreCoefficient_eq_fabijs_sum` in `FabijsLegendreCoefficients.lean`.

The even series

$$\text{up}(x) = \sum_{n=0}^{\infty} u_n P_{2n}(x) \quad (-1 \leq x \leq 1) \quad (794)$$

converges absolutely and uniformly in the formal development. The exact declarations are `summable_abs_rvachevLegendreCoefficient`, `hasSum_rvachevLegendreSeries`, and `hasSum_rvachevLegendreSeries_uniform` in `FabijsLegendreSeries.lean`. Its  $N$ -th polynomial partial sum minimizes squared  $L^2([-1, 1])$  error among all polynomials of degree at most  $2N+1$ , and every distinct competitor has strictly larger error. These are `rvachevLegendrePartialSum_least_squares` and `rvachevLegendrePartialSum_strict_least_squares` in `FabijsLegendreLeastSquares.lean`.

At compiled source checkpoint `a3854643d`, `FabijsLegendreTranslateBlocks.lean` specializes (413) to the Legendre family. Its six public definitions are `rvachevLegendreDeconvolutionPolynomial`, `rvachevLegendreScale`, `rvachevLegendreIndexSet`, `rvachevLegendreAtomCoefficient`, `rvachevLegendreTranslateBlock`, and `rvachevTranslateGram`. Its seven public theorems are `natDegree_rvachevLegendreDeconvolutionPolynomial`, `leadingCoeff_rvachevLegendreDeconvolutionPolynomial`, `eval_legendrePolynomial_eq_sum_rvachevUp`, `eval_legendrePolynomial_even_eq_sum_rvachevUp`, `rvachevLegendreTranslateBlock_eq_rvachevLegendreBlock`, `intervalIntegral_rvachevLegendreTranslateBlock_mul`, and `sum_rvachevLegendreAtomCoefficient_mul_gram`.

Writing  $Q_d = \mathcal{D}_{\text{up}} P_d$ , the two degree theorems give

$$\deg Q_d = d, \quad \text{lc}(Q_d) = 2^{-d} \binom{2d}{d}, \quad (795)$$

so deconvolution preserves the exact Legendre degree and its explicit leading coefficient. The two synthesis theorems then give

$$P_d(x) = 2^{-d} \sum_{-2^{d+1} < k < 2^{d+1}} Q_d(k/2^d) \text{up}(x - k/2^d), \quad (796)$$

$$P_{2n}(x) = 4^{-n} \sum_{-2 \cdot 4^n < k < 2 \cdot 4^n} Q_{2n}(k/4^n) \text{up}(x - k/4^n), \quad (797)$$

for  $-1 \leq x \leq 1$ . More explicitly, let

$$M = 4^n.$$

Setting  $x = 0$  in the even synthesis gives the exact central cancellation

$$Q_{2n}(0) + 2 \sum_{0 < k < M} Q_{2n}(k/M) \text{up}(k/M) = (-1)^n \binom{2n}{n}. \quad (798)$$

The exhaustive public surface of `RvachevLegendreCentralSum.lean` consists of no definitions and the three theorems `eval_legendrePolynomial_even_zero`, `eval_rvachevLegendreDeconvolutionPolynomial_even`, and `rvachevLegendreCentralSum`. The last theorem proves (798) for every  $F : \text{BoundedFabius}$  satisfying `IsFabius`  $F$ , a harmless strengthening from the canonical function, and includes  $n = 0$ .

Indeed, begin with (797) at zero. Compact support removes every term with  $M \leq |k| < 2M$ . The deconvolved even mode is even by `eval_rvachevLegendreDeconvolutionPolynomial_even`, while up is even, so the surviving nonzero indices pair as  $k, -k$ . The central term uses  $\text{up}(0) = 1$ . Finally clear the mesh  $M = 4^n$  and insert

$$P_{2n}(0) = (-1)^n 4^{-n} \binom{2n}{n},$$

which is `eval_legendrePolynomial_even_zero`. Thus the representation frontier’s `cor : leg-central-sum` is *Exact/Complete*. This finite argument supplies no Jacobi closed form, all-degree decoder parity or rationality, reverse spectral closure, mesh minimality, or larger Lagrange right inverse.

More explicitly, let

$$M_n = 4^n, \quad I_n = \{k \in \mathbb{Z} : -2M_n < k < 2M_n\}, \quad a_{n,k} = \frac{u_n}{M_n} Q_{2n}(k/M_n),$$

and define the literal finite translate block

$$\tilde{B}_n(x) = \sum_{k \in I_n} a_{n,k} \text{up}(x - k/M_n). \quad (799)$$

Then  $\tilde{B}_n(x) = u_n P_{2n}(x)$  throughout  $[-1, 1]$ , and these finite trains satisfy the same complete orthogonality formula as the polynomial blocks. With

$$G_{m,n}(k, \ell) = \int_{-1}^1 \text{up}(x - k/M_m) \text{up}(x - \ell/M_n) dx,$$

the fully expanded atom-Gram identity is

$$\sum_{k \in I_m} \sum_{\ell \in I_n} a_{m,k} a_{n,\ell} G_{m,n}(k, \ell) = \delta_{mn} \frac{2u_n^2}{4n+1}. \quad (800)$$

At compiled source checkpoint a3854643d, the focused-build module `FabiusLegendreTranslateSeries.lean` closes the literal outer series and the common-mesh partial sums. Its five public definitions are `rvachevLegendrePartialSumDeconvolutionPolynomial`, `rvachevLegendrePartialSumAtomCoefficient`, `rvachevLegendrePartialSumTranslateBlock`, `rvachevLegendreTranslateBlockOnInterval`, and `rvachevLegendrePartialSumTranslateBlockOnInterval`. Its twenty-five public theorems are `natDegree_rvachevLegendrePartialSumDeconvolutionPolynomial`, `leadingCoeff_rvachevLegendrePartialSumDeconvolutionPolynomial`, `summable_norm_rvachevLegendreTranslateBlock`, `summable_rvachevLegendreTranslateBlock`, `hasSum_rvachevLegendreTranslateBlock`, `tsum_rvachevLegendreTranslateBlock`, `rvachevLegendrePartialSumDeconvolutionPolynomial_eq_sum`, `rvachevLegendrePartialSumAtomCoefficient`

nt\_eq\_sum, eval\_rvachevLegendrePartialSumPolynomial\_eq\_tsum\_rvachevUp, eval\_rvachevLegendrePartialSumPolynomial\_eq\_sum\_rvachevUp, rvachevLegendrePartialSumTranslateBlock\_eq\_eval\_partialSumPolynomial, rvachevLegendrePartialSumTranslateBlock\_eq\_sum\_translateBlock, rvachevLegendreTranslateBlockOnInterval\_apply, rvachevLegendreTranslateBlockOnInterval\_eq\_smul, summable\_norm\_rvachevLegendreTranslateBlockOnInterval, summable\_rvachevLegendreTranslateBlockOnInterval, hasSum\_rvachevLegendreTranslateBlock\_uniform, tsum\_rvachevLegendreTranslateBlock\_uniform, rvachevLegendrePartialSumTranslateBlockOnInterval\_apply, rvachevLegendrePartialSumTranslateBlockOnInterval\_eq\_eval\_partialSumPolynomial, rvachevLegendrePartialSumTranslateBlockOnInterval\_eq\_sum, tendsto\_rvachevLegendrePartialSumTranslateBlockOnInterval, rvachevLegendrePartialSumTranslateBlock\_tendstoUniformlyOn, tendsto\_norm\_rvachevLegendrePartialSumTranslateBlockOnInterval\_sub, and tendsto\_rvachevLegendrePartialSumTranslateBlock.

Consequently the literal finite trains in (799) are absolutely summable both pointwise and in the supremum norm on the interval, and their pointwise and uniform sums are exactly

$$\text{up}(x) = \sum_{n=0}^{\infty} \tilde{B}_n(x), \quad -1 \leq x \leq 1. \quad (801)$$

For the existing polynomial partial sum  $S_N = \sum_{n=0}^N u_n P_{2n}$ , define

$$C_N := \mathcal{D}_{\text{up}} S_N = \sum_{n=0}^N u_n Q_{2n}. \quad (802)$$

The equality is rvachevLegendrePartialSumDeconvolutionPolynomial1\_eq\_sum. With the same hypotheses, the two new triangularity wrappers prove

$$\text{natDegree}(C_N) = \text{natDegree}(S_N), \quad \text{lc}(C_N) = \text{lc}(S_N), \quad (803)$$

including degenerate partial sums whose visible degree drops. With  $M_N = 4^N$ , the global theorem uses a compactly supported tsum for every real input, while on the Legendre interval its exact finite form is

$$S_N(x) = \frac{1}{M_N} \sum_{-2M_N < k < 2M_N} C_N(k/M_N) \text{up}(x - k/M_N). \quad (804)$$

On  $[-1, 1]$ , the named fixed-scale train is equal pointwise both to  $S_N$  and to the finite sum of the separately scaled blocks. The latter statement does not identify the two coefficient vectors. Bundled in  $C([-1, 1])$ , these common-mesh partial trains converge to up in the supremum norm. The API also provides the raw TendstoUniformlyOn statement, convergence of the supremum-norm error to zero, and the pointwise corollary at every  $x \in [-1, 1]$ . It asserts no rate, coefficientwise limit, or uniform convergence outside this interval.

The nine modules RvachevMomentAppell.lean, RvachevPolynomialSynthesis.lean, RvachevSuperconvergentSynthesis.lean, LagrangeRvachevSynthesis.lean, RvachevAppellHasse.lean, LagrangeRvachevMatrix.lean, CompositeMeshSharpness.lean, FabiusLegendreTranslateBlocks.lean, and FabiusLegendreTranslateSeries.lean therefore have public definition/theorem inventories 6/33, 0/5, 1/8, 2/7, 1/14, 4/6, 1/7, 6/7, and 5/25, respectively: exactly 138 public declarations. The seventh module proves least-mesh sharpness only for universal exactness on the complete polynomial space through a degree cutoff. It does not prove minimality for an individual Legendre polynomial or a particular partial sum. The finite Hasse expansion, odd reciprocal-moment vanishing, and geometric closed decoder are now proved. The nine modules still do not prove decoder optimality, range/kernel/rank/trace/characteristic-polynomial

or Cauchy–Binet results for the coefficient-space projector, rationality of the atom rows, equality of the fixed-scale and separately scaled coefficient vectors, an unconditional  $\text{natDegree}(S_N) = 2N$  theorem or nonvanishing of the top Legendre coefficient, or the later refinement, projector, and asymptotic layers.

At compiled source checkpoint `ca66a1247`, the polynomial-form Legendre blocks  $B_n(x) = u_n P_{2n}(x)$  satisfy the complete orthogonality formula

$$\int_{-1}^1 B_m(x) B_n(x) dx = \delta_{mn} \frac{2u_n^2}{4n+1}. \quad (805)$$

This is `intervalIntegral_rvachevLegendreBlock_mul`, for the definition `rvachevLegendreBlock`. The coefficient-energy form of Parseval is also exact. Put  $S_N(x) = \sum_{n=0}^N u_n P_{2n}(x)$ . Then

$$\int_{-1}^1 up(x)^2 dx = \sum_{n=0}^{\infty} \frac{2u_n^2}{4n+1}, \quad (806)$$

$$\int_{-1}^1 (up(x) - S_N(x))^2 dx = \sum_{n=N+1}^{\infty} \frac{2u_n^2}{4n+1}. \quad (807)$$

The full-series and shifted-tail `HasSum` declarations are `hasSum_rvachevLegendreCoefficient_t_energy` and `hasSum_rvachevLegendreCoefficient_energy_tail`; the latter’s `tsum` form is `rvachevLegendreSquaredError_partialSum_eq_tsum_tail`. The finite partial-sum norm is `integral_sq_eval_rvachevLegendrePartialSumPolynomial`. Finally, writing

$$A_2 := \int_0^1 F(t)^2 dt, \quad A_2 = \sum_{n=0}^{\infty} \frac{u_n^2}{4n+1}, \quad (808)$$

so that  $A_2 = \text{fabiusSquareEnergy } F$ , the bridge and series declarations are `integral_sq_rvachevUp_eq_two_mul_fabiusSquareEnergy`, `hasSum_fabiusSquareEnergy_legendre`, and `fabiusSquareEnergy_eq_tsum_legendre` in `FabiusLegendreEnergy.lean`. The block is defined directly in its polynomial form.

At compiled source checkpoint `9d5f41c2c`, `FabiusLegendreRationalEnergy.lean` supplies three executable rational definitions: `canonicalRvachevLegendreCoefficientRat`, `fabiusSquareEnergyTermRat`, and `fabiusSquareEnergyPartialSumRat`. Its fifteen public theorems are `canonicalRvachevLegendreCoefficientRat_cast`, `canonicalRvachevLegendreCoefficientRat_zero`, `fabiusSquareEnergyTermRat_cast`, `fabiusSquareEnergyTermRat_nonneg`, `fabiusSquareEnergyTermRat_zero`, `fabiusSquareEnergyPartialSumRat_cast`, `monotone_fabiusSquareEnergyPartialSumRat`, `fabiusSquareEnergyPartialSumRat_pos`, `fabiusSquareEnergyPartialSumRat_zero`, `fabiusSquareEnergyPartialSumRat_one`, `fabiusSquareEnergyPartialSumRat_two`, `fabiusSquareEnergyPartialSumRat_three`, `hasSum_fabiusSquareEnergy_ratCast`, `fabiusSquareEnergy_eq_tsum_ratCast`, and `tendsto_fabiusSquareEnergyPartialSumRat_cast`. Thus every finite Legendre-energy partial sum is represented by a positive rational number, the rational partial sums are monotone, and their real casts converge to  $A_2$ . For cutoffs  $N = 0, 1, 2, 3$ , respectively, the four certified values are  $1/4$ ,  $7/18$ ,  $3271/8100$ , and  $3246043/8037225$ .

At compiled source checkpoint `b9b240bc0`, `FabiusSquareEnergyFourier.lean` exports no definitions and exactly four theorems: `integral_norm_sq_rvachevFourier_eq_two_mul_fabiusSquareEnergy`, `fabiusSquareEnergy_eq_integral_Ioi_norm_sq_rvachevFourier`, `fabiusSquareEnergy_eq_integral_Ioi_tprod_sinc_sq`, and `fabiusSquareEnergy_eq_scaled_integral_Ioi_tprod_sinc_sq`. The first two identify the full real-axis squared Fourier

mass with  $2A_2$  and its positive-half-line half with  $A_2$ . The last two give exactly

$$A_2 = \int_0^\infty \prod_{j=0}^\infty \text{sinc}_{\text{rad}}^2\left(\frac{\pi\xi}{2^j}\right) d\xi, \quad (809)$$

$$= \frac{1}{2\pi} \int_0^\infty \prod_{r=1}^\infty \left(\frac{\sin(t/2^r)}{t/2^r}\right)^2 dt. \quad (810)$$

The sharpness declarations show that  $2^d$ , and hence  $4^N$  at cutoff  $2N$ , is least for universal exactness on the whole degree-bounded polynomial space. They do not show that the displayed mesh is least for an individual Legendre mode or for one fixed  $S_N$ . The synthesis declarations also do not formalize parity, derivative/Gegenbauer/Appell closed forms, or the displayed low-degree values of  $Q_d$ ; rationality of the atom rows; equality of the fixed-scale and separately scaled coefficient vectors; exact degree of every partial sum; or the later refinement, projector, and asymptotic layers.

On  $[0, 2]$ , the signed global extension has the corresponding translated Legendre series `global Fabius_eq_tsum_translatedLegendre_formula`.

## 14 Additional proved summation identities

On the support interval  $-1 \leq t \leq 1$ , Poisson summation yields the exact cosine series

$$\text{up}(t) = \frac{1}{2} + \sum_{k=0}^\infty \widehat{\text{up}}_{2\pi}\left(\frac{2k+1}{2}\right) \cos((2k+1)\pi t). \quad (811)$$

This is `rvachev_cosine_series`. The two-term support specialization

$$\text{up}(t) + \text{up}(t-1) = 1 \quad (0 \leq t \leq 1) \quad (812)$$

is `rvachev_add_shift_eq_one`. More general scaled Poisson support identities are sharp for either orientation of the lattice: whenever  $1/2 \leq |a|$ ,

$$1 + 2\text{up}(a) = \sum_{m \in \mathbb{Z}} |a|^{-1} \widehat{\text{up}}_{2\pi}(m/a), \quad (813)$$

$$|a| + 2|a|\text{up}(a) = \sum_{m \in \mathbb{Z}} \widehat{\text{up}}_{2\pi}(m/a). \quad (814)$$

These are `rvachev_poisson_support_specialization_unscaled_of_one_half_le_abs` and `rvachev_poisson_support_specialization_of_one_half_le_abs` in `PoissonSummation.lean`; the oriented frequency  $m/a$  is retained while the covolume is  $|a|$ . They supplement the partition formulas (245)–(246); no unformalized derivative partition, probabilistic convolution interpretation, or moment statistic is inferred.

## 15 Effective computability

The centered spline construction gives a primitive-recursive rational approximation procedure. In the exact coding used by Lean, for every dyadic input code  $c$  and precision  $p$ ,

$$|F(c.\text{value}(p+3)) - \text{Approx}(c, p).\text{value}(p)| \leq 5 \cdot 2^{-(p+3)}. \quad (815)$$

This is `fabiusSplineApproxPR_error`; the procedure itself is proved primitive recursive by `fabiusSplineApproxPR_primrec`. Together with the sharp Lipschitz bound it proves that every bounded Fabius solution is sequentially computable and effectively uniformly continuous,

hence a computable real function in the formal Grzegorzczuk-style definition. The declarations are `fabiusReal_sequentiallyComputable`, `fabiusReal_effectivelyUniformContinuous`, and `fabiusReal_isComputableRealFunction` in `FabiusComputableSpline.lean` and `FabiusComputability.lean`. The same effective continuity holds for every signed global extension: `extendedFabius_effectivelyUniformContinuous`. For the canonical extension  $F_{\text{gl}} = \text{globalFabius}$ , the explicit modulus is  $d(n) = 2n$ : whenever  $n > 0$  and  $|x - y| < 1/(2n)$ ,

$$|F_{\text{gl}}(x) - F_{\text{gl}}(y)| < \frac{1}{n}. \quad (816)$$

This is `globalFabius_effectively_uniformly_continuous`, together with the definition `fabiusEffectiveUniformModulus`, in `FabiusComputability.lean`.

The same finite spline code also gives a signed-global algorithm. Instead of clamping the positive numerator to the unit grid, it retains the unrestricted centered spline and rounds the signed rational numerator pair. Negative dyadic inputs collapse to the zero grid point, which is exact because  $F_{\text{gl}}(x) = 0$  for  $x \leq 0$ . The resulting primitive-recursive procedure satisfies

$$|F_{\text{gl}}(c.\text{value}(p+3)) - \text{Approx}_{\text{glob}}(c, p).\text{value}(p)| \leq 5 \cdot 2^{-(p+3)}. \quad (817)$$

These are `extendedFabiusSplineApproxPR_primrec` and `extendedFabiusSplineApproxPR_error`. Hence, for every  $F$  satisfying `IsFabiusF`, its signed extension `extendedFabiusF` is a computable real function, by `extendedFabius_isComputableRealFunction`; the canonical specialization is `globalFabius_isComputableRealFunction`. This is a primitive-recursive correctness statement, not a practical running-time estimate: the unrestricted positive grid numerator sets the length of the finite fold.

The quantitative analytic estimate is stable under moving the evaluation point:

$$|S_p(x) - F_{\text{gl}}(y)| \leq 2^{-p} + 2|x - y|. \quad (818)$$

This is `abs_fabiusUniformSpline_sub_extendedFabius_le_add`. In particular, if  $x_p \rightarrow x$ , then  $S_p(x_p) \rightarrow F_{\text{gl}}(x)$ , as packaged by `fabiusUniformSpline_tendsto_extendedFabius_of_tendsto`.

## 16 Formalization boundary and audit conventions

This exposition follows a fail-closed rule. A mathematical assertion remains in the main document only when an exact proved declaration can be identified in the Lean development. A nearby theorem, an executable computation, a numerical check, or an argument that appears straightforward is not treated as a substitute. When the formal theorem has stricter hypotheses, narrower domains, or different endpoint conventions, the prose uses those exact conditions.

That rule explains several choices made above:

- The finite probability approximation uses atomic convolution measures and the exact weak-convergence API, not an unformalized continuous box-convolution density.
- Arbitrary norm-summable vector weights have formal probability, head–tail, reflection, continuous-linear naturality, and operator-affine recursion laws; their exact barycenter in a complete Borel real normed space is one half of the total vector weight, without sign or order hypotheses; their laws have topological support exactly the series range, and arbitrary signed real weights have exact min/max interval ranges and supports. A nontrivial norm-summable real-weight law is absolutely continuous with respect to Lebesgue measure and has null singletons. The pushforward-law definition itself is total even without measurability; its probability and map identities state the summability, completeness, and Borel hypotheses explicitly. Scalar pointwise

naturality alone needs neither summability nor completeness. The generic affine recurrence is conditional on a displayed shift relation. Separately, `GeometricUniformLaw.lean` defines the named weights  $(1 - q)q^n$ , their series and probability law, and proves self-similarity, reflection, exact mean  $(1/2)$ , absolute continuity, null singletons, and support equal to the series range under  $|q| < 1$ , including zero and negative contractions, with exact support  $[0, 1]$  under  $0 \leq q < 1$ . `ContinuousCDF.lean` and `GeometricUniformCDF.lean` add the named CDF, continuity and reflection for  $|q| < 1$ , the exterior tails for  $0 \leq q < 1$ , and, for  $0 < q < 1$ , the conditioning equations, everywhere derivative, explicit compactly supported Radon–Nikodym density, and real-space  $C^\infty$  package. `ProbabilityRepresentation.lean` supplies the exact half-base CDF and density bridges. `AffineIndependentCopy.lean` proves probability preservation, product-map and characteristic-function factorization, finite characteristic-function iteration, and uniqueness of probability fixed points for  $|q| < 1$  in every complete second-countable real inner-product target. Its `GeometricUniformUniqueness.lean` specialization characterizes the named law, including  $q = 0$  and negative  $q$ , without support, density, or moment assumptions. The selected total density is instead proved zero at  $q = 0$ , nonpositive for negative  $q$ , and to have zero clamped `withDensity` measure for  $q \leq 0$ , which cannot represent  $\mu_q$  on the nonpositive contraction range. Only formalization of the paper-level sign-corrected density remains open. `GeometricSincFactorization.lean` gives the exact finite residual sinc factorization and pointwise phase-bearing prefix limit for every  $|q| < 1$ , with dyadic wrappers. The pure rescaled sinc product now has named locally uniform convergence, pointwise `Multipliable/HasProd`, and entirety. The four theorems of `GeometricSincCharacteristicFunction.lean` give, for  $z = (1 - q)t/(2\pi)$ , the exact identities  $\phi_q(t) = e^{it/2}\Phi_q(z) = e^{it/2}G_q(z)G_q(-z)$ , including  $q = 0$  and negative  $q$ , and prove that  $e^{i(1-q^m)t/2} \prod_{k < m} \text{sinc}_{\text{rad}}((1 - q)q^k t/2)$  converges locally uniformly on the real frequency line to  $\phi_q(t)$  and uniformly on every compact real frequency set. These convergence declarations are `tendstoLocallyUniformly_prefix_sinc_charFun` and `tendstoUniformlyOn_prefix_sinc_charFun`. The seven activation modules separately prove the sharp coefficient  $p(at)/t^2 \rightarrow a^2/3$ , its arbitrary-index square-summable Tannery theorem, the exact geometric coefficient  $(1 - q)/(3(1 + q))$ , and the dyadic value  $1/9$ , together with the degree-eight finite Taylor jet of  $p$  and its order-ten remainder. These quotient limits are punctured and deterministic; no active-count expectation identity is asserted. No complex-frequency analogue for the full phase-bearing prefixes or uniform-convergence claim on all of  $\mathbb{R}$  is made. General- $q$  rapid decay and Fourier inversion, centered or MGF packaging, closed higher-derivative and endpoint-flatness wrappers, shape results, further transforms, and Taylor coefficients beyond the proved finite jet, an all-order Bernoulli coefficient formula, a radius-of-convergence theorem, and the corresponding full activation series remain outside this interface. The separate global complex Pochhammer factorization is fixed at the dyadic scale and does not change those general- $q$  omissions. The exact-support oriented transform DDE and adjacent Cauchy-power recurrence for every nonzero  $|q| < 1$ , including negative ratios, are direct theorems of `GeometricUniformCauchy.lean`; these are distinct from the closed characteristic-function product bridge and do not supply the still-open rapid-decay or Fourier-inversion layers.

- Quantile transport is formal at the measure level first for arbitrary real probability laws, with no support or range restriction and an almost-everywhere sublevel characterization; a compact pointwise specialization feeds the clamped Fabius inverse. This proves strongly measurable Banach-valued substitution with Lean’s totalized Bochner integral. A separate weighted-subgraph development proves Banach-valued survival layer cake and the full-interval Fabius inverse identity for an explicit interval-integrable primitive or a pointwise- $C^1$  primitive. Its strict-order hypothesis actually makes the two compact-interval clock restrictions a Galois connection, so the generic explicit and pointwise- $C^1$  forms now derive clock measurability rather than assuming it. For a real-valued absolutely continuous primitive, a further generic subtraction

theorem and its normalized and Fabius forms accept every real or complex  $L^1$  weight and require no separate complementary-integrability hypothesis. Finally, `InversePairIntegral.lean` proves the full and variable-upper two-function identities for arbitrary absolutely-continuous outer functions on an ordered compact rectangle, derives monotonicity on that rectangle and measurable canonical clamped extensions from the strict inverse-order relation, and supplies the two Fabius wrappers, the full area sum, and the partial inverse-area identity. These declarations do not package reversed-endpoint, higher-order, improper-integral, fractional, complex-valued arbitrary-AC-primitive, or complex-Mellin identities.

- Step cells use the formal half-endpoint indicator, including its point value at a shared boundary.
- The Fourier chapter distinguishes the real punctured-limit statement from the stronger proved local theory: every nonzero signed integer has an exact complex analytic order, all lower complex derivatives vanish, and the first surviving derivative is evaluated explicitly in [Section 5.4](#).
- Summable-scale infinite products are formal under the exact norm-summability, continuity, local-compactness, completeness, and big- $O$  hypotheses stated in [Section 5.2](#). For complex  $q$  with  $\|q\| < 1$ , this gives the entire reciprocal-Gamma product, its zero orbit, Mahler and reflection equations, and the dyadic zero and meromorphic pole orders. The Gamma-side functions are totalized pointwise inverses: raw products of Gamma factors away from their poles, arithmetic canonical regroupings, logarithmic, digamma, Malmsten, heat-trace, integral, D-finite, reciprocal-Gamma derivative-tower, and general-base order statements are not claimed.
- Flatness uses natural powers, and the analytic-locus claims use precisely the domains proved in `NowhereAnalytic.lean`.
- The  $q$ -calculus distinguishes the half-base Fabius identities from the base-generic finite Gaussian/Richardson algebra: recursive Gaussian coefficients, functoriality, both Pascal orientations, reflection symmetry, product concatenation, the full denominator-free factorial identity, principal specializations, denominator-free geometric residual moments, forward closed finite-node numerators, and the proved quarter-base specialization are formal. Generic finite formal-series filters act diagonally coefficientwise, and exact geometric rows have the all-degree Gaussian residual multiplier. Their analytic companion interchanges finite filters with unconditionally summable weighted coefficient series, splits exact heads from infinite tails, and gives the exact Gaussian tsum tail for geometric and Lagrange rows. Zero-weight nodes impose no convergence condition. Named even Fourier modes now instantiate that tail for the actual infinite Rvachev sinc product: arbitrary complex  $c, z$  need only injectivity of the finite powers of  $c^2$ , and  $c = 1/2$ ,  $q = 1/4$  is assumption-free. The two  $q$ -Vandermonde orientations, the central convolution, and all natural shifted forms are denominator-free commutative-semiring theorems. The core also zero-extends the Gaussian lower index to  $\mathbb{Z}$ , proves its total reflection symmetry, and packages the report’s single shifted-central formula for every signed  $k$ , both over  $0, \dots, N$  and as a literal finite-support integer sum. Conditionally-only convergent boundary series, formal-to-analytic evaluation outside the named entire series, the report’s finite-prefix quotient and Bell package, analytic sign/error bounds, uniform or derivative convergence, and asymptotic quarter-sinc acceleration remain outside the formal corpus.
- Integer-order Volterra integration is formal at arbitrary base point and endpoint in a real Banach space, including strict FTC, order composition, polynomial weights, Taylor reconstruction, and the exact Fabius ladder. A separate total real-order operator over arbitrary real normed spaces has the classical continuous-input kernel and additivity theorems when  $0 < \alpha$  and  $a \leq x$ , and agrees with the integer operator at every positive natural order. The shifted real beta kernel is evaluated exactly when both orders are positive and the endpoint inequality is strict; its raw kernel is interval-integrable also at a coincident endpoint. In a complete target, shifted powers and constants have exact Gamma-quotient values, and two positive orders compose



for a continuous input on an ordered compact interval. The atom-preserving lower-CDF layer cake also gives the positive-real Fabius stopped-power identity for every real endpoint and its unit-endpoint moment. Positive affine covariance holds at arbitrary real order for an ordered endpoint and positive scale, and Gamma-normalized order raising needs only a continuous primitive, an interval-integrable displayed derivative, and that derivative from the right on the open interval. The causal Rvachev primitive, its compact-support cutoff, its positive-natural bridge, and its positive-order semigroup are formalized on  $x \geq -1$ . The first exact fractional shifts are proved for the signed extension on  $x \geq 0$ , the bounded function on  $[0, 1]$ , and the up-to- $F$  bridge based at  $-1$  on  $x \geq -1$ . The modules prove no corresponding semigroup law for merely interval-integrable inputs or noncomplete targets, no Caputo or Riemann–Liouville derivative operator, no complex-order result, and no iterated or negative-branch shift, shifted lattice, or endpoint-up wrapper. Separately, the weighted inverse-layer-cake API permits real or complex scalar weights and a Banach-valued explicit or pointwise- $C^1$  primitive on the full unit interval; its arbitrary-absolutely-continuous form retains a real-valued primitive. Variable or reversed endpoints, complex-valued arbitrary-AC primitives, higher inverse primitives, fractional inverse identities, improper-integral limits, and complex Mellin continuation remain outside these results.

- Endpoint asymptotics retain exact phases and the proved remainder scopes; numerical fits and unproved coefficient formulas are absent.

The source drafts have been discharged into two destinations. Proved material was rewritten into the theorem-backed narrative here. The remaining research material is preserved under

Analysis/FabiusFunction/docs/semi-formalized-research-frontiers/.

In particular, inverse-function repeated integration and saddle expansions, improper inverse integrals, complex-power and complex-order integration, conditionally convergent boundary transforms, sinc-tail instantiations and estimates, small-argument analysis, and other deferred results remain available there with an explicit warning that they are not part of the formal corpus. Positive-real order raising, the first fractional Fabius–Rvachev shifts, and unconditional generic- $q$  series transforms have moved into the theorem-backed narrative. Natural-order Volterra integration and finite polynomial weights are formalized in [Section 6.4](#); the compact-rectangle absolutely-continuous inverse calculus is [Theorem 4.8](#); and the finite base-generic Gaussian/Richardson algebra and its formal and unconditionally summable series filters are formalized in [Propositions 10.7](#) to [10.9](#).

## 17 Affine Cauchy–Stieltjes calculus

The current library packages the report-oriented kernel without tying it to a particular probability law. For a finite real measure  $\mu$ , write

$$\mathcal{C}_\mu(z) = \int_{\mathbb{R}} \frac{d\mu(x)}{z - x}, \quad \mathcal{P}_{\mu,n}(z) = \int_{\mathbb{R}} (z - x)^{-n} d\mu(x).$$

These are `measureCauchyTransform` and `measureCauchyPower`, on the natural open domain `measureCauchyDomain`. Their integral, index-one, affine-pushforward, open-domain, derivative, and holomorphy declarations are, respectively, `measureCauchyTransform_apply`, `measureCauchyPower_one`, `measureCauchyPower_map_affine`, `measureCauchyTransform_map_affine`, `isOpen_measureCauchyDomain`, `hasDerivAt_measureCauchyTransform`, and `analyticOn_measureCauchyTransform`. The affine identities are total under Lean’s Bochner-integral convention and allow every nonzero real scale, including a negative one.

The companion module `MeasureCauchyMomentLaurent.lean` gives the bounded-measure moment expansion without choosing the real or complex scalar field in advance. Let  $\mathbb{K}$  be any `RCLike` field, let  $\iota : \mathbb{R} \rightarrow \mathbb{K}$  be its real embedding, and put

$$m_{\mu,k}^{(c)} = \int_{\mathbb{R}} (\iota(x) - c)^k d\mu(x).$$

For a finite real measure, arbitrary  $c, z \in \mathbb{K}$ ,  $0 \leq R$ , almost-everywhere support  $\|\iota(x) - c\| \leq R$ , and the strict exterior condition  $R < \|z - c\|$ , the direct named theorem `integral_inv_sub_eq_sum_range_measureCauchyMoment_add` gives

$$\int_{\mathbb{R}} \frac{d\mu(x)}{z - \iota(x)} = \sum_{k < N} \frac{m_{\mu,k}^{(c)}}{(z - c)^{k+1}} + (z - c)^{-N} \int_{\mathbb{R}} \frac{(\iota(x) - c)^N}{z - \iota(x)} d\mu(x),$$

with positive moment coefficients for the report-oriented kernel. Its finite error obeys

$$\left\| \int_{\mathbb{R}} \frac{d\mu(x)}{z - \iota(x)} - \sum_{k < N} \frac{m_{\mu,k}^{(c)}}{(z - c)^{k+1}} \right\| \leq \mu(\mathbb{R}) (\|z - c\| - R)^{-1} \left( \frac{R}{\|z - c\|} \right)^N.$$

The zeroth centered moment is  $\mu(\mathbb{R})$ , and this is also the finite real mass factor  $\mu.\text{real}(\text{Set.univ})$  in the bound. No probability normalization or topological-support theorem is assumed.

That module's exhaustive public surface is one definition, `measureCauchyMoment`, and fifteen theorems: `inv_sub_eq_sum_range_add`, `measureCauchyMoment_zero`, `measurable_inv_sub_rclike`, `integrable_centered_pow_div_sub_of_ae_norm_sub_le`, `integrable_inv_sub_of_ae_norm_sub_le`, `norm_measureCauchyMoment_le`, `integral_inv_sub_eq_sum_range_measureCauchyMoment_add`, `norm_integral_inv_sub_sub_sum_range_measureCauchyMoment_le`, `summable_measureCauchyMoment_laurent`, `integral_inv_sub_eq_tsum_measureCauchyMoment`, `hasSum_measureCauchyMoment_laurent`, `measureCauchyTransform_eq_sum_range_measureCauchyMoment_add`, `norm_measureCauchyTransform_sub_sum_range_measureCauchyMoment_le`, `measureCauchyTransform_eq_tsum_measureCauchyMoment`, and `hasSum_measureCauchyTransform_measureCauchyMoment_laurent`. The first theorem is the field-generic finite kernel identity; the remaining generic calculus is `RCLike`, and the last four declarations are complex wrappers for the named oriented transform. The `StieltjesMomentLaurent.lean` and `StieltjesCauchyTransform.lean` layers retain the radius-one, center-zero up-measure statements. Their precise bridges are `ae_norm_sub_zero_le_one_rvachevMeasure` and `measureCauchyMoment_rvachevMeasure_zero`, the latter identifying the generic coefficients with `upMoment`.

The reusable fixed-point interface assumes only that  $\mu$  is finite, its support lies in a carrier  $K$ , the map  $(u, x) \mapsto c + au + bx$  preserves  $K$ , both  $a$  and  $b$  are nonzero, and

$$\mu = (\text{Unif}[0, 1] \times \mu)_*(u, x \mapsto c + au + bx).$$

It requires no probability normalization and no closedness, compactness, measurability, or nonemptiness assumption on  $K$ . Off the complexification of that carrier, the direct named theorems `hasDerivAt_measureCauchyTransform_of_uniformAffineFixedPoint` and `measureCauchyPower_succ_of_uniformAffineFixedPoint` give

$$C'_\mu(z) = \frac{1}{ab} \left( C_\mu\left(\frac{z-c}{b}\right) - C_\mu\left(\frac{z-c-a}{b}\right) \right)$$

and, for  $n \geq 1$ ,

$$\mathcal{P}_{\mu,n+1}(z) = \frac{1}{nab^n} \left( \mathcal{P}_{\mu,n}\left(\frac{z-c-a}{b}\right) - \mathcal{P}_{\mu,n}\left(\frac{z-c}{b}\right) \right).$$

For the geometric uniform law the definitions `geometricUniformStieltjesDomain`, `geometricUniformStieltjesTransform`, and `geometricUniformStieltjesPower` are total in `real q`. Their remaining direct public declarations are `geometricUniformStieltjesTransform_apply`, `geometricUniformStieltjesPower_one`, `isOpen_geometricUniformStieltjesDomain`, `analyticOn_geometricUniformStieltjesTransform`, `hasDerivAt_geometricUniformStieltjesTransform_refinement`, and `geometricUniformStieltjesPower_succ`. Holomorphy assumes  $|q| < 1$ ; the divided differential and adjacent-power equations also assume  $q \neq 0$ . Their proofs use the exact series range as invariant carrier, so negative ratios are included. The case  $q = 0$  is excluded only from the divided formulas.

At  $q = 1/2$ , for every bounded Fabius solution, the canonical unit and centered hierarchies are related by the direct named all-order bridge

$$S_n(z) = 2^n R_n(2z - 1).$$

`fabiusStieltjesPower_eq_two_pow_mul_rvachevCauchyPower`

The module `CauchyTransform.lean` contains six public definitions: `rvachevCauchyDomain`, `fabiusStieltjesDomain`, `rvachevCauchyTransform`, `fabiusStieltjesTransform`, `rvachevCauchyPower`, and `fabiusStieltjesPower`. Its direct theorem surface is `rvachevCauchyTransform_apply`, `fabiusStieltjesTransform_apply`, `rvachevCauchyPower_one`, `fabiusStieltjesPower_one`, `rvachevCauchyTransform_eq_integral_rvachevUp`, `isOpen_rvachevCauchyDomain`, `isOpen_fabiusStieltjesDomain`, `hasDerivAt_rvachevCauchyTransform`, `hasDerivAt_fabiusStieltjesTransform`, `analyticOn_rvachevCauchyTransform`, `analyticOn_fabiusStieltjesTransform`, `fabiusStieltjesTransform_eq_two_mul_rvachevCauchyTransform`, `rvachevCauchyTransform_eq_inv_two_mul_fabiusStieltjesTransform`, the all-order bridge above, `hasDerivAt_fabiusStieltjesTransform_refinement`, `fabiusStieltjesPower_succ`, and `rvachevCauchyPower_succ`. In particular, the unit-interval equation  $S'(z) = 4(S(2z) - S(2z - 1))$  and both adjacent-order recurrences on their complex slit domains are now direct named results, not merely informal consequences of affine transport.

The finite orthogonal-polynomial layer now has an exact all-degree rational form. `GramStieltjesNaturality.lean` has no public definitions and exactly six public theorems: `momentPairing_map`, `map_gramStieltjesNumerator`, `map_gramStieltjesPolynomial`, `map_gramStieltjesNorm`, `map_gramStieltjesJacobiDiagonal`, and `map_gramStieltjesJacobiSubdiagonal`. The pairing and fraction-free statements work over their commutative-semiring and commutative-ring boundaries; normalization, norms, and Jacobi quotients use fields. They assert finite scalar naturality, not positivity or analytic convergence.

`RvachevRationalJacobi.lean` has four public definitions: `rvachevHankelDetRat`, `rvachevOrthoPolynomialRat`, `rvachevOrthoNormRat`, and `rvachevJacobiSubdiagonalRat`. Its thirteen public theorems are `upMoment_eq_rvachevRawMomentRat_cast`, `rvachevHankelDetRat_cast`, `rvachevHankelDetRat_pos`, `rvachevOrthoPolynomialRat_cast`, `rvachevOrthoPolynomialRat_isMonicOfDegree`, `momentPairing_rvachevOrthoPolynomialRat_eq_zero`, `rvachevOrthoNormRat_cast`, `rvachevOrthoNormRat_pos`, `gramStieltjesJacobiDiagonal_rvachevRawMomentRat_eq_zero`, `rvachevJacobiSubdiagonalRat_cast`, `rvachevJacobiSubdiagonalRat_pos`, `rvachevJacobiSubdiagonalRat_eq_det_ratio`, and `rvachevOrthoPolynomialRat_three_term`. Thus every up-law moment, Hankel determinant, monic orthogonal polynomial, squared norm, and Jacobi subdiagonal has an exact rational witness; the determinant, norm, and subdiagonal witnesses are positive, the diagonal vanishes, and the rational polynomials satisfy the centered three-term recurrence in every degree. The convention is zero-based: `rvachevJacobiSubdiagonalRat(n)` is the usual coefficient  $\beta_{n+1}$ , so

$$\pi_{n+2}(X) = X\pi_{n+1}(X) - \beta_{n+1}\pi_n(X).$$

These four rational definitions are noncomputable finite-algebra objects, not native-code evaluators. This module proves all-degree rationality without itself evaluating the report’s displayed  $H_4$  or  $\beta_4$  fractions; the downstream executable rational-Legendre layer and its values leaf supply those named finite certificates without changing the all-degree interface.

The polynomial change-of-basis layer is now explicit as well. `PolynomialMomentGramDeterminant.lean` has exactly two public definitions, `polynomialCoefficientMatrix` and `polynomialMomentGramMatrix`, and exactly seven public theorems: `polynomialCoefficientMatrix_apply`, `polynomialMomentGramMatrix_apply`, `polynomialMomentGramMatrix_eq_transpose_mul_hankel_mul`, `polynomialMomentGramMatrix_det_eq_q_coefficient_det_sq_mul`, `polynomialCoefficientMatrix_det_eq_prod_coeff`, `polynomialMomentGramMatrix_det_eq_prod_coeff_sq_mul`, and `gramStieltjesJacobiSubdiagonal_eq_polynomialMomentGramMatrix_det_ratio`. For a family  $p_j$  with  $\deg p_j \leq j$ , put its monomial coefficient vectors in the columns of  $C_n$ , let  $H_n$  be the monomial Hankel matrix, and let  $G_n$  be the moment Gram matrix. Expanding the bilinear pairing coefficient by coefficient gives

$$G_n = C_n^\top H_n C_n.$$

The degree bound makes  $C_n$  upper triangular, with diagonal  $[X^j]p_j$ . Determinant multiplicativity and  $\det C_n = \prod_{j < n} [X^j]p_j$  therefore give

$$\det G_n = \left( \prod_{j < n} [X^j]p_j \right)^2 \det H_n,$$

and the final generic theorem transports the zero-based Jacobi cross-ratio through any such family with nonzero diagonal coefficients. That equality has no Hankel-nonvanishing hypothesis: Lean uses totalized field division, so when the middle Hankel determinant vanishes both cross-ratios are zero. The singular equality is therefore not a genuine nonsingular Jacobi recurrence.

The coefficient layer needed to execute the Legendre specialization is supplied by `LegendrePolynomialRational.lean`. It has exactly two public definitions: `legendrePolynomialCoeffRat`, the bounded rational sum computing the coefficient of  $X^k$  in  $P_n$ , and `legendrePolynomialRat`, its noncomputable  $\mathbb{Q}[X]$ -valued wrapper. Its exactly six public theorems are `legendrePolynomialRat_cast`, `coeff_legendrePolynomialRat`, `natDegree_legendrePolynomialRat`, `coeff_legendrePolynomialRat_self`, `coeff_legendrePolynomialRat_self_ne_zero`, and `coeff_legendrePolynomialRat_self_div_succ`. They identify every wrapper coefficient with the executable sum, identify its coefficientwise real cast with the Rodrigues-normalized `legendrePolynomial`, and prove natural degree  $n$ , leading coefficient  $2^{-n} \binom{2n}{n}$ , its nonvanishing, and the consecutive quotient

$$\frac{[X^n]P_n}{[X^{n+1}]P_{n+1}} = \frac{n+1}{2n+1}.$$

This module proves finite rational coefficient algebra only; it contains no root, Christoffel, quadrature, orthogonality, or asymptotic theorem.

The companion `LegendreGaunt.lean` turns those coefficients into an executable triple-product layer. Its exactly four public definitions are `legendreLebesgueMomentRat`, `legendreGauntRat`, `legendreGaunt`, and `legendreProductLinearizationCoeffRat`. Its exactly twelve public theorems are `legendreLebesgueMomentRat_even`, `legendreLebesgueMomentRat_odd`, `legendreLebesgueMomentRat_cast`, `legendreGauntRat_eq_momentFunctional`, `legendreGauntRat_cast`, `legendreGauntRat_swap_left`, `legendreGauntRat_swap_right`, `legendrePolynomial_mul_eq_sum_gaunt`, `legendrePolynomialRat_mul_eq_sum_gaunt`, `legendreGauntRat_eq_zero_of_odd_sum`, `legendreGauntRat_eq_zero_of_add_lt`, and `legendreGauntRat_eq_zero_of_triangle_violation`. The executable moments are

$2/(n+1)$  at even  $n$  and zero at odd  $n$ ; combining them with three finite rational Legendre coefficient arrays gives `legendreGauntRat`. Its cast is exactly the real interval integral

$$\text{legendreGaunt}(i, j, k) = \int_{-1}^1 P_i(x)P_j(x)P_k(x) \, dx.$$

The two swap theorems generate the usual permutation symmetry, and the normalized coefficient  $(2k+1)\text{legendreGauntRat}(i, j, k)/2$  gives exact finite product linearizations of  $P_iP_j$  over the rational and real coefficient fields. An odd sum  $i+j+k$ , or any strict triangle violation, forces the rational Gaunt coefficient to vanish. These are necessary support restrictions only within `LegendreGaunt.lean`: that module itself proves no converse, Wigner-square, factorial, or positivity theorem. The following closed-form leaf supplies the exact total squared zero-row integer-index datum, sharpens support to an equivalence, and proves its factorial forms and nonnegativity; those statements should not be read back into this twelve-theorem API.

At the proof level, monomial integration on the symmetric interval gives the displayed even/odd moments. Expanding the three rational Legendre polynomials into their finite coefficient arrays then identifies the triple sum with the rational moment functional, and finite linearity plus the monomial cast gives the real integral. Since  $\deg(P_iP_j) \leq i+j$ , finite Fourier–Legendre projection and orthogonality give the real product linearization; injectivity of the coefficientwise cast  $\mathbb{Q}[X] \rightarrow \mathbb{R}[X]$  reflects it back to rational polynomials. Commutativity gives the two swaps, odd reflection gives the parity zero, and orthogonality beyond degree  $i+j$ , followed by the swaps, gives all three strict triangle zeros.

The focused module `LegendreGauntClosedForm.lean` closes that integer-index, zero-magnetic-row layer. Its exactly two public definitions are `legendreGauntAdmissible` and `legendreWignerThreeJZeroSqRat`. Its exactly twenty-five public theorems are `legendreGauntAdmissible_iff_exists_pairwise_add`, `legendreGauntAdmissible_pairwise_add`, `legendreWignerThreeJZeroSqRat_pairwise_add`, `legendreWignerThreeJZeroSqRat_pairwise_add_factorial`, `legendreWignerThreeJZeroSqRat_eq_factorial_of_halfSum`, `legendreWignerThreeJZeroSqRat_eq_zero_of_not_admissible`, `legendreGauntRat_add_boundary`, `legendreGauntRat_add_boundary_eq_two_mul_wignerThreeJZeroSqRat`, `legendreGauntRat_zero_left`, `legendreGauntRat_zero_left_eq_two_mul_wignerThreeJZeroSqRat`, `legendreGauntRat_pairwise_add_eq_two_mul_wignerThreeJZeroSqRat`, `legendreGauntRat_eq_zero_of_not_admissible`, `legendreGauntRat_eq_two_mul_wignerThreeJZeroSqRat`, `legendreGaunt_eq_two_mul_wignerThreeJZeroSqRat`, `legendreWignerThreeJZeroSqRat_pos_iff_admissible`, `legendreWignerThreeJZeroSqRat_nonneg`, `legendreWignerThreeJZeroSqRat_eq_zero_iff_not_admissible`, `legendreGauntRat_pos_iff_admissible`, `legendreGauntRat_eq_zero_iff_not_admissible`, `legendreGaunt_pos_iff_admissible`, `legendreGaunt_eq_zero_iff_not_admissible`, `legendreGauntRat_nonneg`, `legendreGaunt_nonneg`, `legendreProductLinearizationCoeffRat_eq_mul_wignerThreeJZeroSqRat`, and `legendreProductLinearizationCoeffRat_pos_iff_admissible`. Here `legendreGauntAdmissible( $i, j, k$ )` means that  $i+j+k$  is even and that the three weak triangle inequalities hold. Equivalently, there are  $a, b, c \in \mathbb{N}_0$  with

$$(i, j, k) = (b+c, a+c, a+b).$$

The total rational datum  $W_{ijk}^{2,\mathbb{Q}} = \text{legendreWignerThreeJZeroSqRat}(i, j, k)$  is zero off that support. In the pairwise-sum coordinates it is the symmetric central-binomial quotient

$$W_{b+c, a+c, a+b}^{2,\mathbb{Q}} = \frac{\binom{2a}{a} \binom{2b}{b} \binom{2c}{c}}{(2(a+b+c)+1) \binom{2(a+b+c)}{a+b+c}}.$$

The API supplies both this expression’s factorial form and the equivalent half-sum factorial form. The latter assumes exactly  $i+j+k = 2s$  and  $i, j, k \leq s$ . At arbitrary natural indices the global

identification and its real cast are

$$G_{ijk}^{\mathbb{Q}} = 2W_{ijk}^{2,\mathbb{Q}}, \quad G_{ijk} = 2(W_{ijk}^{2,\mathbb{Q}} : \mathbb{R}).$$

Consequently the rational datum, the rational Gaunt coefficient, and the real Gaunt integral are positive exactly on the admissible support, vanish exactly off it, and are nonnegative everywhere. The exact coefficient of  $P_k$  in  $P_i P_j$  satisfies

$$L_{ijk} = (2k + 1)W_{ijk}^{2,\mathbb{Q}}$$

and is positive exactly on that same support. Separate boundary theorems evaluate the degenerate triangle  $k = i + j$  and the case  $i = 0$ . The name “Wigner square” is deliberately precise: this is a rationally defined square for integer indices in the zero-magnetic row. The module chooses no signed Wigner symbol or Condon–Shortley phase convention and develops no half-integer indices, nonzero magnetic indices, general  $3j/6j/9j$  symbols, Wigner orthogonality, or recoupling identity.

`FabiusLegendreHankelDeterminant.lean` specializes this calculation to the ordinary Legendre family and the up-law moments. It has exactly two public definitions, `upLegendreGramMatrix` and `upLegendreGramDet`, and exactly seven public theorems: `upLegendreGramMatrix_apply_eq_integral`, `upLegendreGramDet_eq_prod_leadingCoeff_sq_mul_hankelDet`, `upLegendreGramDet_zero`, `upLegendreGramDet_pos`, `coeff_legendrePolynomial_self_div_succ`, `gramStieltjesJacobiSubdiagonal_upMoment_eq_upLegendreGramDet_ratio`, and `rvachevJacobiSubdiagonalRat_cast_eq_upLegendreGramDet_ratio`. For arbitrary  $F : \text{BoundedFabius}$ , let  $D_n$  be the determinant of the first  $n$  Legendre polynomials in its up-moment pairing and put  $L_j = 2^{-j} \binom{2j}{j}$ . At  $n = 0$ , “the first zero polynomials” means the empty  $0 \times 0$  Gram matrix, whose determinant is one; this convention extends the paper’s prior  $N \geq 1$  matrix definition. The determinant identity and empty case hold for every such  $F$ :

$$D_n = \left( \prod_{j < n} L_j^2 \right) \det H_n, \quad D_0 = 1.$$

The leading-coefficient quotient is independent of  $F$ , and, with Lean’s index  $n$  denoting the conventional coefficient  $\beta_{n+1}$ , the real Gram cross-ratio also holds for arbitrary  $F : \text{BoundedFabius}$ :

$$\beta_{n+1} = \left( \frac{n+1}{2n+1} \right)^2 \frac{D_{n+2} D_n}{D_{n+1}^2}.$$

Here the arbitrary- $F$  formula inherits totalized division from the generic theorem. The entry-as-integral theorem, strict positivity  $D_n > 0$ , and identification with the real cast of the exact rational Rvachev subdiagonal each additionally require  $h_F : \text{IsFabius } F$ . Under that hypothesis positivity removes the singular case and the cross-ratio is a genuine Jacobi formula. This determinant module does not itself expand individual Gram entries. The finite rational Gaunt expansion and entry rationality are supplied by `LegendreGaunt.lean` and `FabiusLegendreGaunt.lean`. The core total rational integer-index zero-row Wigner-square datum, its central-binomial and factorial forms, sharp support, positivity, nonnegativity, and boundary laws are supplied by `LegendreGauntClosedForm.lean`. The wrapper `FabiusLegendreGauntClosedForm.lean` transports that identity to the unconditional finite rational entry and matrix sums and, under `BoundedFabius` with `IsFabius`, to the finite real up-law matrix sum. Neither leaf chooses a signed  $3j$  phase or supplies half-integer or nonzero magnetic indices, Wigner orthogonality, recoupling, Christoffel reconstruction, or an infinite Jacobi product. Also, `rvachevTranslateGram` is a different object: it pairs shifted-up atoms under unweighted interval integration, whereas `upLegendreGramMatrix` pairs polynomials under the up-law moment functional.

Together with the preceding rational Legendre coefficient module, the next two modules make the finite Legendre layer executable over  $\mathbb{Q}$ . First, `FabiusLegendreRationalGram.lean` turns the same specialization into an executable rational layer. Its three public definitions are `rvachevLegendreGramEntryRat`, the bounded double coefficient–moment sum, `rvachevLegendreGramMatrixRat`, its finite matrix, and `rvachevLegendreGramDetRat`, its determinant. Its exactly eleven public theorems are `rvachevLegendreGramEntryRat_eq_momentPairing`, `rvachevLegendreGramMatrixRat_apply`, `rvachevLegendreGramMatrixRat_eq_polynomialMomentGramMatrix`, `rvachevLegendreGramEntryRat_cast`, `rvachevLegendreGramMatrixRat_cast`, `rvachevLegendreGramDetRat_cast`, `rvachevLegendreGramDetRat_eq_prod_leadingCoeff_sq_mul_rvachevHankelDetRat`, `rvachevLegendreGramDetRat_zero`, `rvachevLegendreGramDetRat_pos`, `rvachevOrthoNormRat_eq_rvachevLegendreGramDetRat_ratio`, and `rvachevJacobiSubdiagonalRat_eq_rvachevLegendreGramDetRat_ratio`. The first three identify the executable entry and matrix with the abstract moment pairing and polynomial-moment Gram matrix. For  $F : \text{BoundedFabius}$  with  $h_F : \text{IsFabius } F$ , the next three identify the real casts of the entry, matrix, and determinant with the corresponding up-law objects. Over  $\mathbb{Q}$ , the determinant is the rational Hankel determinant times the product of the squared Legendre leading coefficients; it equals one at order zero and is strictly positive in every order. The last two theorems express the exact rational squared norm and the zero-based `rvachevJacobiSubdiagonalRat`( $n$ ) =  $\beta_{n+1}$  as consecutive rational Legendre Gram-determinant ratios, with the latter carrying the factor  $((n+1)/(2n+1))^2$ . Thus the executable coefficient, entry, matrix, determinant, cast, positivity, norm-ratio, and Jacobi-ratio layer is formalized. Its finite Gaunt entry representation is supplied by the next companion module. The closed-form core `LegendreGauntClosedForm.lean` supplies the total rational integer-index zero-row Wigner-square datum and factorial closure, while `FabiusLegendreGauntClosedForm.lean` substitutes it into the finite rational entry and matrix sums and the finite real up-law matrix sum. This rational Gram module itself proves neither formula. No signed  $3j$  phase, half-integer or nonzero magnetic indices, Wigner orthogonality, recoupling, Christoffel reconstruction, root or quadrature theory, infinite Jacobi product or continued fraction, Padé theory, or asymptotic result is supplied here.

`FabiusLegendreGaunt.lean` has exactly one public definition, `canonicalRvachevFullLegendreCoefficientRat`, and exactly eight public theorems: `canonicalRvachevFullLegendreCoefficientRat_even`, `canonicalRvachevFullLegendreCoefficientRat_odd`, `canonicalRvachevFullLegendreCoefficientRat_cast`, `canonicalRvachevFullLegendreCoefficientRat_eq_normalized_moment`, `rvachevLegendreGramEntryRat_eq_sum_full_gaunt`, `rvachevLegendreGramEntryRat_eq_sum_gaunt`, `rvachevLegendreGramMatrixRat_apply_eq_sum_gaunt`, and `upLegendreGramMatrix_apply_eq_sum_gaunt`. The even and odd definitions immediately prove the first two laws. Existing exact Rvachev–Legendre coefficient formulas identify the real cast of  $C_k^{\mathbb{Q}}$  with the full analytic coefficient for every bounded Fabius representative  $F$  satisfying `IsFabiusF`, and identify its rational value with the normalized up-moment of  $P_k^{\mathbb{Q}}$ :

$$C_k^{\mathbb{Q}} = \frac{2k+1}{2} \mathcal{L}_{\text{up}}^{\mathbb{Q}}(P_k^{\mathbb{Q}}).$$

Applying the rational moment functional to the finite product linearization and substituting this normalization yields

$$\mathcal{G}_{ij}^{\mathbb{Q}} = \sum_{k=0}^{i+j} C_k^{\mathbb{Q}} G_{ijk}^{\mathbb{Q}} = \sum_{r=0}^{\lfloor (i+j)/2 \rfloor} c_r^{\mathbb{Q}} G_{ij,2r}^{\mathbb{Q}}.$$

The second equality simply deletes the zero odd terms and reindexes  $k = 2r$ ; the matrix theorem is the same formula at finite indices. Casting the finite sum, using the two cast bridges term by term,

gives

$$\mathcal{G}_{ij}^{(n)}(F) = \sum_{r=0}^{\lfloor (i+j)/2 \rfloor} c_r(F) G_{ij,2r}$$

The rational matrix-entry theorem gives the second formula for every  $i, j : \text{Fin } n$ . Only canonicalRvachevFullLegendreCoefficientRat\_cast and upLegendreGramMatrix\_apply\_eq\_sum\_gaunt require  $F : \text{BoundedFabius}$  and  $h_F : \text{IsFabius } F$ : the former identifies the cast with rvachevFullLegendreCoefficient( $F, k$ ), while the latter replaces the rational coefficient and Gaunt term in the even finite sum by rvachevLegendreCoefficient( $F, r$ ) and legendreGaunt( $i, j, 2r$ ). The other six theorems are unconditional finite identities. Together with LegendreGaunt.lean, these theorems close the finite Gaunt-integral/product-linearization layer and the finite Rvachev Gram-entry expansion. Those two predecessor modules alone prove no Wigner-square identification, factorial formula, nonnegativity, or sharp converse support theorem; the separate LegendreGauntClosedForm.lean module above does. Still absent are any signed-symbol or phase convention, half-integer or nonzero-magnetic-index theory, infinite Legendre-series/integral interchange or infinite expansion, root or quadrature theorem, Christoffel reconstruction or Padé approximation, and named values for the individual entries of the displayed low-order  $G_3$ .

---

**Source/PDF parity convention.** This source contains the exhaustive two-definition, twenty-eight-theorem closed-form inventory and the expanded formula-and-proof ledger below. The external repository metadata certifies only its named historical source/PDF checkpoint; this merged source postdates that receipt, and a fresh uninterrupted three-pass Libertinus parity rebuild remains pending. The receipt’s page count and validation facts remain authoritative for that historical checkpoint.

---

For a formula-level audit of the preceding inventory, the core LegendreGauntClosedForm.lean has exactly two public definitions. The first is legendreGauntAdmissible; in mathematical notation,

$$A(i, j, k) \iff (i + j + k) \bmod 2 = 0, \quad i \leq j + k, \quad j \leq i + k, \quad k \leq i + j.$$

The second is legendreWignerThreeJZeroSqRat. Put  $C_n = \binom{2n}{n}$  and  $s = \lfloor (i + j + k)/2 \rfloor$ . Its total rational value is

$$W_{ijk}^2 := \begin{cases} \frac{C_{s-i} C_{s-j} C_{s-k}}{(2s+1) C_s}, & A(i, j, k), \\ 0, & \neg A(i, j, k). \end{cases} \quad (819)$$

Thus the superscript in  $W_{ijk}^2$  denotes the formalized *square datum*; it is not the square of a separately defined signed Lean symbol.

The core’s exhaustive public theorem inventory contains exactly twenty-five names, in source order. The support-coordinate and closed-form group is legendreGauntAdmissible\_iff\_exists\_pairwise\_add, legendreGauntAdmissible\_pairwise\_add, legendreWignerThreeJZeroSqRat\_pairwise\_add, legendreWignerThreeJZeroSqRat\_pairwise\_add\_factorial, legendreWignerThreeJZeroSqRat\_eq\_factorial\_of\_halfSum, and legendreWignerThreeJZeroSqRat\_eq\_zero\_of\_not\_admissible. The boundary and Gaunt-identification group is legendreGauntRat\_add\_boundary, legendreGauntRat\_add\_boundary\_eq\_two\_mul\_wignerThreeJZeroSqRat, legendreGauntRat\_zero\_left, legendreGauntRat\_zero\_left\_eq\_two\_mul\_wignerThreeJZeroSqRat, legendreGauntRat\_pairwise\_add\_eq\_two\_mul\_wignerThreeJZeroSqRat, legendreGauntRat\_eq\_zero\_of\_not\_admissible, legendreGauntRat\_eq\_two\_mul\_wignerThreeJZeroSqRat, and legendreGaunt\_eq\_two\_mul\_wignerThreeJZeroSqRat. The sharp-support, sign, and product-coefficient group is legendreWignerThreeJZeroSqRat\_pos\_iff\_admissible, legendreWignerThreeJZeroSqRat\_nonneg, legendreWignerThreeJZeroSqRat\_eq\_zer



o\_iff\_not\_admissible, legendreGauntRat\_pos\_iff\_admissible, legendreGauntRat\_eq\_zero\_iff\_not\_admissible, legendreGaunt\_pos\_iff\_admissible, legendreGaunt\_eq\_zero\_iff\_not\_admissible, legendreGauntRat\_nonneg, legendreGaunt\_nonneg, legendreProductLinearizationCoeffRat\_eq\_mul\_wignerThreeJZeroSqRat, and legendreProductLinearizationCoeffRat\_pos\_iff\_admissible.

The pairwise-sum theorem makes the support parametrization exact:

$$A(i, j, k) \iff \exists a, b, c \in \mathbb{N}_0, \quad i = b + c, \quad j = a + c, \quad k = a + b. \quad (820)$$

For  $S = a + b + c$ , the central-binomial and factorial forms are

$$W_{b+c, a+c, a+b}^2 = \frac{C_a C_b C_c}{(2S+1)C_S}, \quad (821)$$

$$= \frac{(S!)^2 (2a)! (2b)! (2c)!}{(2S+1)! (a!)^2 (b!)^2 (c!)^2}. \quad (822)$$

Equivalently, if  $i + j + k = 2s$  and  $i, j, k \leq s$ , then

$$W_{ijk}^2 = \frac{(s!)^2 (2(s-i))! (2(s-j))! (2(s-k))!}{(2s+1)! ((s-i)!)^2 ((s-j)!)^2 ((s-k)!)^2}. \quad (823)$$

The hypotheses in this half-sum version are part of the theorem; natural-number subtraction is never being used as an implicit signed subtraction.

Writing  $G_{ijk}^{\mathbb{Q}} = \text{legendreGauntRat}(i, j, k)$ , the two boundary laws are

$$G_{i,j,i+j}^{\mathbb{Q}} = \frac{2C_i C_j}{(2(i+j)+1)C_{i+j}} = 2W_{i,j,i+j}^2, \quad (824)$$

$$G_{0,j,k}^{\mathbb{Q}} = \begin{cases} 2/(2j+1), & j = k, \\ 0, & j \neq k, \end{cases} = 2W_{0,j,k}^2. \quad (825)$$

The global rational identity and its real-integral cast are unconditional:

$$G_{ijk}^{\mathbb{Q}} = 2W_{ijk}^2, \quad \text{legendreGaunt}(i, j, k) = 2(W_{ijk}^2 : \mathbb{R}). \quad (826)$$

Consequently the square datum, the rational Gaunt coefficient, and the real Gaunt integral are positive exactly when  $A(i, j, k)$  holds, vanish exactly when it does not, and are nonnegative for every natural triple. If  $L_{ijk}^{\mathbb{Q}} = \text{legendreProductLinearizationCoeffRat}(i, j, k)$ , then

$$L_{ijk}^{\mathbb{Q}} = (2k+1)W_{ijk}^2, \quad L_{ijk}^{\mathbb{Q}} > 0 \iff A(i, j, k). \quad (827)$$

*Proof architecture.* Let  $\lambda_n = [X^n]P_n = 2^{-n}C_n$ . In the already formalized finite product linearization, only the  $k = i + j$  summand can contribute to the coefficient of  $X^{i+j}$ . Hence  $\lambda_i \lambda_j = L_{i,j,i+j}^{\mathbb{Q}} \lambda_{i+j}$ ; substituting  $L_{ijk}^{\mathbb{Q}} = (2k+1)G_{ijk}^{\mathbb{Q}}/2$  gives (824). The case with first index zero follows from that diagonal boundary and the predecessor module’s triangle-support zeros.

The private propagation engine proves the rational Bonnet identity

$$XP_n = \frac{n+1}{2n+1}P_{n+1} + \frac{n}{2n+1}P_{n-1}.$$

Applying the rational moment functional to  $(XP_i)P_jP_k = P_i(XP_j)P_k$  transfers it to

$$\frac{i+1}{2i+1}G_{i+1,j,k}^{\mathbb{Q}} + \frac{i}{2i+1}G_{i-1,j,k}^{\mathbb{Q}} = \frac{j+1}{2j+1}G_{i,j+1,k}^{\mathbb{Q}} + \frac{j}{2j+1}G_{i,j-1,k}^{\mathbb{Q}}.$$

The candidate in (821) satisfies the corresponding triangle-coordinate recurrence because  $(n+1)C_{n+1} = 2(2n+1)C_n$ . A nested induction in the pairwise coordinates uses the degenerate-triangle formula on the two boundary faces and the transfer recurrence in the interior; its leading rational coefficient is strictly positive, so cancellation is legitimate. This proves  $G^{\mathbb{Q}} = 2W^2$  for every pairwise-sum triple. Equation (820) covers every admissible triple, while the old parity and triangle zeros make both sides zero off support. Finally  $C_n = (2n)!/(n!)^2$  gives the two factorial displays, positivity of every central binomial gives the sharp support and nonnegativity results, and the real identity follows through the existing rational-to-real Gaunt cast. No external Wigner integral evaluation is used.  $\square$

The wrapper `FabiusLegendreGauntClosedForm.lean` adds no public definition and exactly three public theorems: `rvachevLegendreGramEntryRat_eq_two_mul_sum_wignerThreeJZeroSqRat`, `rvachevLegendreGramMatrixRat_apply_eq_two_mul_sum_wignerThreeJZeroSqRat`, and `upLegendreGramMatrix_apply_eq_two_mul_sum_wignerThreeJZeroSqRat`. For natural  $i, j$ , it gives the unconditional rational formula

$$\mathcal{G}_{ij}^{\mathbb{Q}} = 2 \sum_{r=0}^{\lfloor (i+j)/2 \rfloor} c_r^{\mathbb{Q}} W_{i,j,2r}^2, \quad (828)$$

and the same formula entrywise for `rvachevLegendreGramMatrixRat`. For  $F : \text{BoundedFabius}$ , a proof  $h_F : \text{IsFabius } F$ , and  $i, j : \text{Fin } n$ , the real wrapper is

$$\text{upLegendreGramMatrix}(F, n, i, j) = 2 \sum_{r=0}^{\lfloor ((i:\mathbb{N}_0) + (j:\mathbb{N}_0))/2 \rfloor} c_r(F) \left( W_{(i:\mathbb{N}_0), (j:\mathbb{N}_0), 2r}^2 : \mathbb{R} \right). \quad (829)$$

These are finite sums: each proof substitutes (826) into the predecessor Gaunt sum and distributes the factor two. Only the real theorem needs `BoundedFabius` and `IsFabius`; the two rational theorems are unconditional.

The Wigner boundary is exact. The development formalizes only a total rational integer-index, zero-magnetic-row *square* datum. It defines no signed Wigner symbol, makes no Condon–Shortley or other phase choice, and therefore proves no signed identity. It also supplies no half-integer indices, nonzero magnetic indices, general  $3j$ ,  $6j$ , or  $9j$  symbols, Wigner orthogonality, or recoupling theorem. The Gram wrappers add no infinite-series assertion.

The low-order computation leaf `FabiusLegendreRationalGramValues.lean` has zero public definitions and exactly eleven public theorems: `moment_four`, `rvachevLegendreGramDetRat_one`, `rvachevLegendreGramDetRat_two`, `rvachevLegendreGramDetRat_three`, `rvachevLegendreGramDetRat_four`, `rvachevLegendreGramDetRat_five`, `rvachevOrthoNormRat_four`, `rvachevJacobiSubdiagonalRat_three`, `hankelRatio_four`, `integral_sq_upOrthoPolynomial_four`, and `hankelRatio_four_div_three`. The first theorem gives the raw eighth moment, and the next five give the rational Legendre Gram determinants of orders one through five:

$$m_8 = \frac{132809}{32531625}, \quad (D_1, D_2, D_3, D_4, D_5) = \left( 1, \frac{1}{9}, \frac{8}{2025}, \frac{39616}{602791875}, \frac{16544275456}{27453718922765625} \right).$$

The next two unconditional rational computations are

$$H_4 = \frac{26727424}{55791736875}, \quad \beta_4 = \frac{835232}{4640643},$$

proved respectively by `rvachevOrthoNormRat_four` and `rvachevJacobiSubdiagonalRat_three`; the Lean subdiagonal index is zero-based. Each of the final three transport theorems—the real Hankel-ratio value, its squared-orthogonal-polynomial integral, and the identity  $H_4/H_3 = \beta_4$ —requires both  $F : \text{BoundedFabius}$  and  $h_F : \text{IsFabius } F$ . Thus the displayed norm and subdiagonal

are now named exact Lean evaluations. The separate Gaunt modules close the finite entry expansion, and the closed-form leaves close the total rational integer-index zero-row Wigner-square identification, its central-binomial and factorial forms, and the corresponding finite rational-entry, rational-matrix, and real-matrix Gram sums; this values leaf supplies no named evaluation of the displayed  $G_3$  entries. No signed  $3j$  phase, half-integer or nonzero magnetic indices, Wigner orthogonality, recoupling, Christoffel reconstruction, roots or quadrature, infinite Jacobi product or continued fraction, Padé theory, or asymptotic result is supplied here.

This closure does not supply a complex logarithmic fixed point, complex order, pointwise or nontangential Sokhotski–Plemelj limits, a principal-value Hilbert-transform identity, or the full Jacobi/Padé theory. Separate current modules do prove the real logarithmic identity for  $z > 1$ , real order lowering for  $\alpha > 1$ , the generic centered exterior Laurent series and its radius-one up-measure specialization, finite-height Herglotz–Poisson identities, integrated interval Stieltjes–Perron inversion, and initial exact Jacobi data; none of those statements is read as one of the stronger boundary or continuation claims just listed.

## 18 Conclusion

The formal development links three global presentations without confusing their domains. The bounded  $F$  supplies the probability and arithmetic viewpoints; the compactly supported up supplies refinement, Fourier, spline, and Legendre viewpoints; and the signed  $F_{\text{gl}}$  supplies the global dilation and binary-translation formulas. Exact bridges connect them to finite atomic laws, weak convergence, centered splines, moments, generic weighted distributions, geometrically weighted uniform laws, exact quantile transport, Banach-valued survival layer cake, weighted inverse layer cake, arbitrary absolutely-continuous inverse-pair calculus, reciprocal-power values, dyadic algorithms, half-base  $q$ -binomial identities, prefix sums, denominator-free arbitrary-base geometric interpolation, finite formal-series and unconditionally summable analytic filters with their Gaussian residuals, normalized integer-order and positive-real fractional Volterra foundations, exact integer Fabius primitives and first fractional Fabius–Rvachev shifts, polynomial weights, endpoint Laplace analysis, affine Cauchy–Stieltjes and centered Laurent calculus, exact integer zero-row Gaunt–Wigner-square formulas, and computable approximation.

The most important methodological conclusion is equally concrete: the primary document is now a map of proved mathematics. Research claims can move back from the frontier only after a matching Lean theorem exists, with the same hypotheses, domain, normalization, endpoint convention, and conclusion.

## A Lean module map

Every retained result above points to one or more exact declaration names. The principal source groups are:

| Topic                              | Modules                                                                                                                                         |
|------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|
| Definitions, existence, uniqueness | <code>Basic.lean</code> , <code>Existence.lean</code> ,<br><code>OriginalCharacterization.lean</code> ,<br><code>OriginalUniqueness.lean</code> |
| Shape and regularity               | <code>Differential.lean</code> , <code>Monotonicity.lean</code> ,<br><code>Convexity.lean</code> , <code>Regularity.lean</code>                 |

| Topic                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Modules                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Probability, weighted-law<br>naturalness, absolute continuity,<br>exact support and affine<br>recursion, fixed half–quarter<br>multisection, contractive<br>affine-law uniqueness,<br>geometric sinc-prefix<br>factorization, quantile<br>transport, survival/CDF layer<br>cake, fractional stopped<br>powers, affine Cauchy–Stieltjes<br>calculus, renormalization,<br>Laurent/Herglotz/inversion<br>layers, initial Jacobi data, and<br>finite approximants | WeightedUniformSeries.lean,<br>WeightedUniformDistribution.lean,<br>WeightedUniformSupport.lean,<br>GeometricUniformLaw.lean,<br>GeometricUniformMultisection.lean,<br>AffineIndependentCopy.lean,<br>GeometricUniformUniqueness.lean,<br>GeometricSincFactorization.lean,<br>ContinuousCDF.lean, GeometricUniformCDF.lean,<br>ProbabilityRepresentation.lean,<br>QuantileTransport.lean, SubgraphFubini.lean,<br>SurvivalLayerCake.lean, CDFLayerCake.lean,<br>FractionalCDFLayerCake.lean,<br>FabijsFractionalIntegral.lean,<br>InverseLayerCake.lean, InversePairIntegral.lean,<br>MeasureCauchyTransform.lean,<br>MeasureCauchyMomentLaurent.lean,<br>GeometricUniformCauchy.lean,<br>CauchyTransform.lean, CauchyCDF.lean,<br>CauchySurvival.lean, CauchyHigherPowers.lean,<br>CauchyRenormalization.lean,<br>StieltjesLogFixedPoint.lean,<br>StieltjesResolventHierarchy.lean,<br>StieltjesGeneralizedOrder.lean,<br>StieltjesMomentLaurent.lean,<br>StieltjesCauchyTransform.lean,<br>StieltjesHerglotz.lean, StieltjesInversion.lean,<br>StieltjesPerron.lean,<br>OrthogonalPolynomialJacobi.lean,<br>GramStieltjesNaturalness.lean,<br>RvachevRationalJacobi.lean,<br>PolynomialMomentGramDeterminant.lean,<br>LegendrePolynomialRational.lean,<br>LegendreGaunt.lean, LegendreGauntClosedForm.lean,<br>FabijsLegendreHankelDeterminant.lean,<br>FabijsLegendreRationalGram.lean,<br>FabijsLegendreGaunt.lean,<br>FabijsLegendreGauntClosedForm.lean,<br>FabijsLegendreRationalGramValues.lean,<br>RvachevLegendreCentralSum.lean,<br>EarlyApproximants.lean, EarlyMeasureBridge.lean,<br>WeakConvergence.lean,<br>ProbabilityLaplaceMoments.lean,<br>LaplaceMoments.lean, StepMeasureBridge.lean,<br>StepApproximationLimit.lean,<br>FabijsUniformSpline.lean,<br>QuarterSplineLocalPolynomial.lean,<br>QuarterSplineTwoSided.lean,<br>FabijsComputability.lean |

| Topic                                                                                                                                                                                                                                                                                                                                                           | Modules                                                                                                                                                                                                                                                                                                        |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Executable rational Legendre Gaunt coefficients and finite product linearization                                                                                                                                                                                                                                                                                | <code>LegendreGaunt.lean</code>                                                                                                                                                                                                                                                                                |
| Two definitions and twenty-five theorems for the total rational integer-index zero-row Wigner-square datum, pairwise-coordinate central-binomial and factorial forms, the half-sum factorial form, global rational and real Gaunt identities, degenerate and zero-index boundary values, sharp support, positivity, nonnegativity, and product-coefficient laws | <code>LegendreGauntClosedForm.lean</code>                                                                                                                                                                                                                                                                      |
| Finite rational and real up-law Legendre Gram entry expansions                                                                                                                                                                                                                                                                                                  | <code>FabijsLegendreGaunt.lean</code>                                                                                                                                                                                                                                                                          |
| Zero public definitions and three theorems for the finite rational entry, rational matrix, and real up-law matrix as twice finite canonical-coefficient sums against the total rational integer-index zero-row Wigner-square datum                                                                                                                              | <code>FabijsLegendreGauntClosedForm.lean</code>                                                                                                                                                                                                                                                                |
| Hyperbolic activation kernels, the finite activation Taylor jet, arbitrary square-summable activation series, sharp local and Tannery asymptotics, and geometric and dyadic effective dimensions                                                                                                                                                                | <code>HyperbolicActivation.lean</code> , <code>ActivationSeries.lean</code> , <code>ActivationAsymptotics.lean</code> , <code>ActivationTaylor.lean</code> , <code>ActivationSeriesAsymptotics.lean</code> , <code>GeometricActivationDimension.lean</code> , <code>GeometricActivationAsymptotics.lean</code> |

| Topic                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | Modules                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Fourier analysis                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | FourierProduct.lean, FourierAnalytic.lean,<br>RvachevQBinomialFilter.lean,<br>OriginalPaperSupplement.lean,<br>PoissonSummation.lean, SummableCyclicAlias.lean,<br>HalfIntegerOddDFT.lean, ThueMorseFourier.lean,<br>ThueMorseSineProduct.lean,<br>ThueMorseComplexProductBridge.lean,<br>LacunaryProductToSum.lean,<br>CenteredRvachevThueMorseFourier.lean,<br>RenormalizationIdentity.lean,<br>SincProductShells.lean, SincPrefixBound.lean,<br>SincCanonicalProduct.lean, PhiZeroOrder.lean,<br>IntegerZeroLocalFactorization.lean,<br>IntegerZeroAnalyticOrder.lean,<br>ThueMorseLobeSign.lean,<br>RvachevProductContinuity.lean,<br>PeakRayEnvelope.lean, GlobalDecayEnvelope.lean,<br>RvachevFourierDecay.lean,<br>VarianceBookkeeping.lean, BirkhoffVariance.lean,<br>BirkhoffProductBridge.lean |
| Summable-scale products,<br>finite and infinite<br>q-Pochhammer dissection, the<br>generic complete-normed-ring<br>infinite symbol and its explicit<br>complex zero derivatives,<br>geometric reciprocal-Gamma<br>factorization, the separate<br>complex Pochhammer wrapper<br>and its analytic-order-one<br>factor theory, locally uniform<br>convergence of the outer<br>spectral Pochhammer product,<br>and the latest infinite<br>q-binomial, Jacobi, q-Pascal,<br>continuity, quantum-binomial,<br>and Rogers–Szegő polynomial<br>APIs | ScaledInfiniteProducts.lean,<br>GeometricReciprocalGamma.lean,<br>RvachevPochhammerFactorization.lean,<br>QPochhammerDissection.lean,<br>QPochhammerInfinite.lean, QPochhammerEntire.lean,<br>GeometricPochhammerNormalConvergence.lean,<br>ReciprocalIntegerGammaZeros.lean,<br>DyadicGammaOrder.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |

| Topic                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | Modules                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Gaussian ( $q=-1$ ) first jets, generic and universal<br>Gaussian-polynomial degree, palindromicity, mean, linear coefficient, and second cumulants, central/cyclotomic factorization, primitive-root blocks, $q$ -Lucas and $q$ -Catalan theory, Euler and infinite $q$ -binomial expansions, effective fixed-column Gaussian rates, Jacobi and pentagonal products, $q$ -Pascal summation, Gaussian continuity, the quantum binomial theorem, and Rogers–Szegő polynomials | <code>GaussianBinomialAtNegOneDerivative.lean</code> ,<br><code>GaussianBinomialPalindromic.lean</code> ,<br><code>GaussianBinomialCumulants.lean</code> ,<br><code>GaussianBinomialPolynomialStructure.lean</code> ,<br><code>CentralQBinomialReduction.lean</code> ,<br><code>CyclotomicFactorization.lean</code> ,<br><code>PrimitiveRootBlock.lean</code> , <code>QLucas.lean</code> ,<br><code>CyclotomicDivisibility.lean</code> , <code>QCatalan.lean</code> ,<br><code>TwoPhiOneReversal.lean</code> , <code>QChuVandermonde.lean</code> ,<br><code>QBinomialTheoremInfinite.lean</code> ,<br><code>GaussianBinomialFixedColumnRate.lean</code> ,<br><code>GaussianBinomialGreaterOneAsymptotics.lean</code> ,<br><code>JacobiTripleProduct.lean</code> , <code>TwoSquareTheorem.lean</code> ,<br><code>JacobiTwoSquareCount.lean</code> , <code>JacobiCubic.lean</code> ,<br><code>ThetaQuasiPeriodicity.lean</code> ,<br><code>QPascalSummation.lean</code> ,<br><code>GaussianBinomialContinuity.lean</code> ,<br><code>QuantumBinomial.lean</code> , <code>RogersSzegoPolynomial.lean</code> ,<br><code>PolynomialQDerivative.lean</code> ,<br><code>PolynomialQLeibniz.lean</code> , <code>QExponential.lean</code> ,<br><code>JacksonIntegral.lean</code> , <code>QPochhammerDerivative.lean</code> ,<br><code>QPochhammerLogDerivative.lean</code> ,<br><code>QPochhammerOrderDerivative.lean</code> ,<br><code>LambertSeriesLog.lean</code> , <code>QGamma.lean</code> |
| Reciprocal-Gamma first jets and the Thue–Morse Gamma tower, including its positive-parameter differential ladder and the local centered-corner integral                                                                                                                                                                                                                                                                                                                      | <code>ReciprocalGammaJets.lean</code> ,<br><code>ThueMorseGammaTower.lean</code> ,<br><code>ThueMorseGammaTowerDifferential.lean</code> ,<br><code>ThueMorseCornerIntegral.lean</code>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Local all-order jet calculus                                                                                                                                                                                                                                                                                                                                                                                                                                                 | <code>LeadingJet.lean</code>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Normalized integer- and positive-real-order Volterra calculus, affine covariance, shifted-power and constant evaluations, positive affine covariance and right-derivative order raising, base-point shifts, exact integer Fabius primitives, causal Rvachev fractional primitives, and first fractional Fabius–Rvachev shifts                                                                                                                                                | <code>NormalizedVolterra.lean</code> , <code>FractionalVolterra.lean</code> ,<br><code>FractionalVolterraSemigroup.lean</code> ,<br><code>FractionalVolterraCalculus.lean</code> ,<br><code>FabijsAntiderivatives.lean</code> ,<br><code>FabijsFractionalVolterra.lean</code>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Global derivatives, value distributions, and flatness                                                                                                                                                                                                                                                                                                                                                                                                                        | <code>GlobalExtension.lean</code> , <code>GlobalBounds.lean</code> ,<br><code>BoundedDerivatives.lean</code> ,<br><code>MidpointEndpointTransfer.lean</code> ,<br><code>DyadicDerivativeFiltration.lean</code> ,<br><code>RvachevDerivativeDistribution.lean</code> ,<br><code>EffectiveFlatness.lean</code> , <code>SharpFlatness.lean</code> ,<br><code>FabijsFlatness.lean</code>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |

| Topic                                                                                                                                                                                                    | Modules                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Analyticity, inverse, effective inverse continuity, elementary obstruction                                                                                                                               | NowhereAnalytic.lean, FabiusInverse.lean, InverseMidpointDefect.lean, InverseModulus.lean, FabiusInverseQuarterJet.lean, FabiusInverseEffectiveContinuity.lean, FabiusInverseLogarithmicModulus.lean, FabiusInverseExactDyadicModulus.lean, EffectiveMonotoneInverse.lean, EffectiveGapInverse.lean, FabiusInverseComputable.lean, ElementaryFunction.lean, NotElementary.lean, InverseBranch.lean, InverseNotElementary.lean                                                                                                                                                                                                                                                                                                                                                                       |
| Effective monotone inversion and total inverse computability (exactly two definitions and six theorems in the generic module, followed by zero definitions and one theorem in the Fabius specialization) | EffectiveMonotoneInverse.lean, FabiusInverseComputable.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Effective inversion from computable positive rational dyadic gaps (exactly four definitions and four theorems, including the clamped total extension)                                                    | EffectiveGapInverse.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Moments and exact arithmetic                                                                                                                                                                             | Arithmetic.lean, AnalyticMoments.lean, MomentPowerSeries.lean, UnitLaplaceMomentBounds.lean, NormalizedEvenMoments.lean, BernoulliRecurrences.lean, MomentCumulantAlgebra.lean, CenteredMomentCumulants.lean, CenteredMomentParity.lean, SinhDivBernoulliLog.lean, PolynomialExpectationCumulant.lean, RvachevMomentAppell.lean, RvachevAppellHasse.lean, RvachevLagrangeNodesOnly.lean, GeometricUniformMomentPolynomial.lean, GeometricUniformMomentPolynomialBridge.lean, GeometricUniformComplexMomentProduct.lean, GeometricUniformExteriorComplexMomentGerm.lean, GeometricUniformMomentPolynomialDegree.lean, WeightedScaleMultiplicity.lean, BaseDigitMultiplicity.lean, ExactInversePower.lean, Parity.lean, TwoAdic.lean, CentralBinomialValuation.lean, PrimePowerBinomialValuation.lean |



| Topic                                                                                                                                                                                                                                                                                                                                                                                                     | Modules                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Universal exponential, Bell, and endpoint-transfer algebra                                                                                                                                                                                                                                                                                                                                                | SaddleExpansionAlgebra.lean,<br>ExponentialPartition.lean, ExponentialBell.lean,<br>EndpointTransferPolynomials.lean,<br>LogSeriesMultiplicative.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Formal implicit roots and nonmonic Hensel lifting                                                                                                                                                                                                                                                                                                                                                         | ImplicitPowerSeries.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Finite base-layer arithmetic                                                                                                                                                                                                                                                                                                                                                                              | BaseLayerRegrouping.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Finite polynomial, formal-series, complete-homogeneous/Bell, and unconditionally summable analytic-series filters, and geometric Lagrange/Richardson exactness                                                                                                                                                                                                                                            | FinitePolynomialFunctional.lean,<br>CompleteHomogeneous.lean,<br>CompleteHomogeneousGenerating.lean,<br>CompleteHomogeneousBell.lean,<br>FinitePowerSeriesFilter.lean,<br>GeometricPowerSeriesFilter.lean,<br>AnalyticSeriesFilter.lean,<br>GeometricLagrange.lean,<br>LagrangeResidualMoments.lean,<br>GeometricCompleteHomogeneous.lean,<br>GeometricResidualMoments.lean,<br>GeometricQBinomialLagrange.lean,<br>GeometricRichardson.lean,<br>GeometricLagrangeWeights.lean,<br>GeometricLagrangeQBinomial.lean,<br>GeometricLagrangeQMoments.lean,<br>GeometricLagrangeCompleteHomogeneous.lean,<br>GeometricRichardsonGenerating.lean |
| Phase-locked reciprocal-grid Richardson and Bell algebra                                                                                                                                                                                                                                                                                                                                                  | LambertPhaseLockedRichardson.lean,<br>LambertPhaseLockedBell.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Phase-locked pullback, complete-homogeneous asymptotics, reciprocal-row growth, and finite-order periodic extraction                                                                                                                                                                                                                                                                                      | FabiusLambertPhaseLockedPullback.lean,<br>CompleteHomogeneousAsymptotics.lean,<br>LambertReciprocalAsymptotics.lean,<br>FabiusLambertPhaseExtraction.lean,<br>FabiusLambertPhaseExtractionBell.lean                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Finite $q$ -Cauchy convolutions and the $q$ -Bernstein partition of unity                                                                                                                                                                                                                                                                                                                                 | QBinomialCauchy.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Zero definitions and exactly three theorems: continuous <code>_gaussianBinomial</code> , <code>tendsto_gaussianBinomial_nhds_one</code> , and <code>gaussianBinomial_eq_finiteQPochhammerIn_div</code> . The first two hold for all $n, k$ in every topological semiring whose operations are continuous; the quotient identity needs only a field, $k \leq n$ , and $(q; q)_k \neq 0$ , with no topology | GaussianBinomialContinuity.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |

| Topic                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | Modules                                      |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|
| <p>Two definitions, <code>thetaExponent</code> and <code>pentagonalExponent</code>, and exactly twenty-five theorems: <code>two_dvd_mul_sub_one</code>, <code>mul_sub_one_nonneg</code>, <code>two_mul_thetaExponent</code>, <code>thetaExponent_natCast</code>, <code>thetaExponent_neg_natCast</code>, <code>choose_two_add_choose_two_succ_eq</code>, <code>sum_mul_sum_eq_sum_antidiagonal</code>, <code>thetaCoeff_pair_eq</code>, <code>sum_antidiagonal_thetaCoeff</code>, <code>finite_triple_product_poly</code>, <code>pow_mul_finiteQPochhammerIn_div</code>, <code>neg_one_zpow_natCast_sub</code>, <code>finite_triple_product</code>, <code>norm_gaussianBinomialInt_le</code>, <code>tendsto_gaussianBinomialInt_central</code>, <code>summable_pow_thetaExponent_mul_zpow</code>, <code>hasSum_jacobi_triple_product</code>, <code>hasSum_jacobi_triple_product'</code>, <code>mul_three_mul_sub_one_nonneg</code>, <code>two_dvd_mul_three_mul_sub_one</code>, <code>two_mul_pentagonalExponent</code>, <code>pentagonalExponent_eq</code>, <code>pentagonalExponent_pos</code>, <code>hasSum_pentagonal</code>, and <code>hasSum_pentagonal_pairs</code>. The exponent arithmetic is unconditional; the finite polynomial block is over commutative rings, while the Laurent block is over fields and its divided-product formulas assume <math>z \neq 0</math>. The norm bound needs a normed field and <math>\ q\  &lt; 1</math>; the central limit, Jacobi identities, and pentagonal identities add completeness, with <math>z \neq 0</math> exactly for Jacobi. The pentagonal identities include <math>q = 0</math>, and the real theta majorant assumes <math>0 \leq r &lt; 1</math> and <math>0 &lt; s</math></p> | <p><code>JacobiTripleProduct.lean</code></p> |

| Topic                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | Modules                              |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------|
| <p>One definition, gaussianMajorant, and exactly twenty-two theorems: hasSum_of_tendsto_of_dominated, one_le_qPochhammerInfIn_neg, finiteQPochhammerIn_neg_le_qPochhammerInfIn, qPochhammerInfIn_nonneg, qPochhammerInfIn_le_finiteQPochhammerIn, qPochhammerInfIn_pos_of_lt_one, qPochhammerInfIn_zero_left, norm_finiteQPochhammerIn_le_neg_norm, finiteQPochhammerIn_norm_le_norm, norm_finiteQPochhammerIn_le, qPochhammerInfIn_norm_le_norm_finiteQPochhammerIn, finiteQPochhammerIn_le_zero_of_norm_lt_one, gaussianBinomial_eq_div, gaussianMajorant_nonneg, norm_gaussianBinomial_le, tendsto_finiteQPochhammerIn_pow_atTop, tendsto_gaussianBinomial_atTop, summable_pow_choose_two_mul_pow, hasSum_euler_product, qPochhammerInfIn_le_zero_of_norm_lt_one, hasSum_qBinomial_theorem, and hasSum_euler_reciprocal. The Tannery theorem is Banach-valued and uses a common summable bound, pointwise row convergence, row HasSums, and convergence of row sums. The ordered comparison block has its displayed real hypotheses <math>0 \leq r &lt; 1</math> and respectively <math>0 \leq x, x \leq 1, 0 \leq x \leq 1</math>, or <math>0 \leq x &lt; 1</math>. The infinite zero-left identity is unconditional over its stated topological commutative ring type; the finite identity is owned by GaussianBinomialAtOne.lean. Norm bounds and fixed-column limits need only a normed field and the</p> | <p>QBinomialTheoremInfinite.lean</p> |

| Topic                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | Modules                            |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------|
| <p>Zero definitions and exactly four theorems: <code>sum_gaussianBinomial_succ_mul</code>, <code>sum_gaussianBinomial_succ_mul'</code>, <code>Commute.gaussianBinomial_left</code>, and <code>Commute.gaussianBinomial_right</code>. The first summation and both commutation laws hold in every semiring, with only the indicated <code>Commute</code> hypothesis for the latter; the second summation assumes a commutative semiring. There is no division, order, topology, or restriction on <math>q, n</math>, or the weight sequence</p> | <code>QPascalSummation.lean</code> |
| <p>Zero definitions and exactly two theorems, <code>quantumPlane_mul_pow</code> and <code>quantum_binomial</code>. They work in an arbitrary, possibly noncommutative semiring. The power lemma assumes <math>q</math> commutes with <math>X</math> and <math>YX = q(XY)</math>; the binomial theorem additionally assumes that <math>q</math> commutes with <math>Y</math>. No division or topology is used</p>                                                                                                                               | <code>QuantumBinomial.lean</code>  |

| Topic                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | Modules                                 |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|
| <p>One definition, <code>rogersSzego</code>, and exactly nine theorems:</p> <p><code>rogersSzego_zero</code>,<br/> <code>rogersSzego_one</code>,<br/> <code>rogersSzego_succ</code>, <code>one_succ</code>,<br/> <code>b_pow_succ_mul_gaussia</code>,<br/> <code>nBinomial_succ_succ</code>,<br/> <code>rogersSzego_dilation</code>,<br/> <code>rogersSzego_three_term</code>,<br/> <code>summable_norm_pow_div</code>,<br/> <code>finiteQPochhammerIn_self</code>,<br/> <code>sum_antidiagonal_euler_mul_euler</code>, and <code>hasSum_rogersSzego_generating</code>.</p> <p>The definition and first three theorems use a commutative semiring; the cross, dilation, and three-term laws use a commutative ring and are total in their natural indices. The summability theorem needs a normed field, <math>\ q\  &lt; 1</math>, and <math>\ z\  &lt; 1</math>; the antidiagonal identity needs only the contraction. The generating function adds completeness and assumes exactly <math>\ q\  &lt; 1</math>, <math>\ t\  &lt; 1</math>, and <math>\ zt\  &lt; 1</math>.</p> | <code>RogersSzegoPolynomial.lean</code> |

| Topic                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | Modules                                 |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|
| <p>Two definitions and seventeen theorems: <code>qInt</code>, <code>qDerivative</code>; <code>qInt_zero</code>, <code>qInt_one</code>, <code>qInt_succ</code>, <code>qInt_succ'</code>, <code>qInt_add</code>, <code>qInt_one_left</code>, <code>one_sub_mul_qInt</code>, <code>qDerivative_apply</code>, <code>qDerivative_monomial</code>, <code>qDerivative_C</code>, <code>qDerivative_X_pow</code>, <code>qDerivative_X</code>, <code>qDerivative_C_mul_X_pow</code>, <code>eval_qDerivative_mul</code>, <code>qDerivative_comp_C_mul_X</code>, <code>qDerivative_mul</code>, and <code>qDerivative_mul'</code>. <math>Q</math>-integers use a semiring, their geometric identity and the division-free evaluation law use a commutative ring, and the remaining polynomial derivative API uses a commutative semiring. Arbitrary <math>q</math>, including zero and one; no analytic or limiting derivative</p> | <code>PolynomialQDerivative.lean</code> |
| <p>Zero definitions and four theorems: <code>comp_C_mul_X_comp_C_mul_X</code>, <code>qDerivative_C_mul</code>, <code>qDerivative_iterate_comp_C_mul_X</code>, and <code>qDerivative_iterate_mul</code>. All hold over a commutative semiring for arbitrary <math>q</math> and every natural order, with no division, nonvanishing, topology, or convergence hypothesis</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | <code>PolynomialQLeibniz.lean</code>    |

| Topic                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Modules                                 |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|
| <p>Zero definitions and three theorems: <code>hasDerivAt_finiteQPochhammerIn</code>, <code>hasDerivAt_finiteQPochhammerIn_of_ne_zero</code>, and <code>hasDerivAt_finiteQPochhammerIn_comp</code>. Arbitrary <math>q, a, n</math> over a nontrivially normed field; the cofactor formula is unconditional, the logarithmic rewrite assumes every finite factor nonzero, and the chain rule assumes the displayed pointwise derivative witness. No infinite-product derivative</p>                             | <code>QPochhammerDerivative.lean</code> |
| <p>Zero definitions and four theorems: <code>one_le_norm_natCast_add_one</code>, <code>summable_lambert_series</code>, <code>hasSum_lambert_log_complex</code>, and <code>exp_neg_tsum_lambert_eq_qPochhammerInfIn</code>. Apart from the unconditional norm helper, exactly complex <math>\ a\  &lt; 1</math> and <math>\ q\  &lt; 1</math>; principal factor logs and a branch-free exponential identity, but no norm-one boundary, product-log branch identity, continuation, or differentiated series</p> | <code>LambertSeriesLog.lean</code>      |

| Topic                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | Modules                                                                                                                                                                                                                           |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Two total real definitions and ten theorems: <code>qGamma</code> , <code>qNumber</code> ; <code>norm_lt_one_of_pos_of_lt_one</code> , <code>qPochhammerInfIn_rpow_pos</code> , <code>qPochhammerInfIn_self_pos</code> , <code>qGamma_pos</code> , <code>qGamma_one</code> , <code>qGamma_add_one</code> , <code>qGamma_nat_succ</code> , <code>qNumber_natCast</code> , <code>qGamma_mul_qGamma_one_sub</code> , and <code>hasSum_theta_qGamma_reflection</code> . The positive recurrence API uses $0 < q < 1$ and the displayed $x > 0$ ; the natural bridge only $q \neq 1$ ; reflection only $q < 1$ and arbitrary real $x$ ; theta form $0 < q < 1$ , $0 < x < 1$ . No complex continuation, classical limit, poles, log-convexity, uniqueness, or digamma theorem | <code>QGamma.lean</code>                                                                                                                                                                                                          |
| Weighted Pascal, elementary-complete orthogonality, formal generating series, and weighted sequence inversion                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | <code>SymmetricFunctionOrthogonality.lean</code> ,<br><code>SymmetricFunctionGenerating.lean</code> ,<br><code>SymmetricFunctionTransform.lean</code> ,<br><code>CompleteHomogeneous.lean</code>                                  |
| Generic finite triangular transforms, weighted-subset Gaussian enumeration, scalar kernels, module-valued $q$ -binomial inversion, and $q$ -Gaussian specializations                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <code>FiniteTriangularTransform.lean</code> ,<br><code>BitPositionQBinomial.lean</code> ,<br><code>QBinomialInversion.lean</code> , <code>QBinomialTransform.lean</code> ,<br><code>QBinomialInversionSpecializations.lean</code> |
| Exhaustive four-theorem scaled inverse-Gaussian characteristic, complete geometric moments, scalar-extension top-coefficient extraction, and exact $q$ -difference annihilation API                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | <code>QDifferenceAnnihilation.lean</code>                                                                                                                                                                                         |



| Topic                                                                                                                                                                                             | Modules                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Dyadic and $q$ -binomial formulas                                                                                                                                                                 | DyadicClosedForm.lean, DyadicAnalytic.lean,<br>GlobalDyadic.lean, DyadicCorrectness.lean,<br>FiniteQBinomialCore.lean,<br>QBinomialReciprocity.lean,<br>GaussianBinomialPalindromic.lean,<br>GaussianBinomialCumulants.lean,<br>GaussianBinomialPolynomialStructure.lean,<br>CentralQBinomialReduction.lean,<br>CyclotomicFactorization.lean,<br>PrimitiveRootBlock.lean, QLucas.lean,<br>CyclotomicDivisibility.lean, QCatalan.lean,<br>TwoPhiOneReversal.lean, QChuVandermonde.lean,<br>QPochhammerElementaryIdentities.lean,<br>QBinomialCauchy.lean, QBinomialVandermonde.lean,<br>HalfQBinomial.lean,<br>HalfQBinomialRootSimplicity.lean,<br>QuarterCatalanGerm.lean,<br>QuarterCatalanRichardson.lean,<br>FabijsDyadicQBinomialScalar.lean,<br>FabijsQBinomialFormula.lean,<br>FabijsRawQBinomialScalar.lean,<br>FabijsBinaryReductionSeries.lean,<br>FabijsParityPowerSeries.lean,<br>FabijsGlobalQBinomialSeries.lean,<br>FabijsDiscreteLimitToeplitz.lean,<br>FabijsDiscreteLimitIntegration.lean |
| Thue–Morse algebra, finite complex transforms, prefixes, disc series, and ring-valued Prouhet identities, including the even/odd binary-weight partition and centered symmetric mixed differences | ThueMorseBitSupport.lean, ThueMorsePrefix.lean,<br>ThueMorseBooleanCube.lean,<br>ThueMorseComplexExponential.lean,<br>ThueMorseMixedDifference.lean,<br>ThueMorseSymmetricDifference.lean,<br>ThueMorseGenerating.lean,<br>ThueMorseBlockAlgebra.lean,<br>ThueMorseInfiniteProduct.lean,<br>ThueMorseDiscSeries.lean,<br>ThueMorseNaturalBoundary.lean,<br>ThueMorseExponential.lean,<br>ThueMorseApproximation.lean,<br>ThueMorseMixedDifference.lean,<br>ThueMorseSymmetricDifference.lean,<br>ThueMorseSparseProuhet.lean,<br>PaperKFoldThueMorse.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                   |

| Topic                                | Modules                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|--------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Endpoint asymptotics                 | NegativeLaplace.lean,<br>NegativeLaplaceVertical.lean,<br>PeriodicCorrection.lean, PeriodicFourier.lean,<br>MellinBose.lean, MellinFinitePart.lean,<br>BoseFinitePartIntegral.lean,<br>PrincipallLambertW.lean, LowerLambertW.lean,<br>PowerExponentialLambert.lean,<br>PowerExponentialLambertCalculus.lean,<br>PowerExponentialLambertInverse.lean,<br>PowerExponentialLambertAsymptotics.lean,<br>PowerExponentialLambertFabius.lean,<br>FabiusLambertPhase.lean,<br>LambertWBranchPairing.lean,<br>LambertWGapBijection.lean,<br>LambertWBranchSymmetry.lean,<br>LambertWBranchGapBernoulli.lean,<br>LambertWCurvature.lean,<br>LambertWBranchPointGeometry.lean,<br>LambertWBranchPointAsymptotics.lean,<br>PowerExponentialLambertCurvature.lean,<br>PowerExponentialLambertFabiusCurvature.lean,<br>FabiusLambertSaddle.lean,<br>LambertPhaseLockedRichardson.lean,<br>FabiusSharpLambertMain.lean,<br>FabiusWikipediaMain.lean,<br>FabiusDecayComparison.lean,<br>FabiusSharpAsymptotic.lean,<br>FabiusFullAsymptoticExpansion.lean,<br>FabiusSharpLambertTransfer.lean,<br>LaplaceMomentBounds.lean,<br>FabiusDyadicSharpCumulant.lean,<br>SaddleLogExpansionAlgebra.lean,<br>SaddleLogExpansionPowerSeries.lean,<br>FabiusSecondSaddleCorrection.lean,<br>FabiusSecondSaddleExpansion.lean<br> |
| Legendre expansion and computability | FabiusLegendreCoefficients.lean,<br>FabiusLegendreSeries.lean,<br>FabiusLegendreLeastSquares.lean,<br>FabiusLegendreEnergy.lean,<br>FabiusLegendreRationalEnergy.lean,<br>FabiusSquareEnergyFourier.lean,<br>RvachevPolynomialSynthesis.lean,<br>RvachevSuperconvergentSynthesis.lean,<br>CompositeMeshSharpness.lean,<br>FabiusLegendreTranslateBlocks.lean,<br>FabiusLegendreTranslateSeries.lean,<br>FabiusTranslatedLegendreSeries.lean,<br>FabiusComputableSpline.lean                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |

## B Notation crosswalk

| Article                                                                  | Lean                                                                                                    | Meaning                                                                                                                  |
|--------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|
| $F$                                                                      | <code>fabiusReal fabius</code>                                                                          | bounded canonical Fabius function                                                                                        |
| $\text{up}$                                                              | <code>rvachevUp fabius</code>                                                                           | compactly supported fold                                                                                                 |
| $F_{\text{gl}}$                                                          | <code>extendedFabius fabius</code>                                                                      | signed global extension                                                                                                  |
| $S_w$                                                                    | <code>ProbabilityRepresentation.weightedUniformSeries w</code>                                          | weighted coordinate series                                                                                               |
| $\mu_w$                                                                  | <code>ProbabilityRepresentation.weightedUniformDistribution w</code>                                    | its pushforward law                                                                                                      |
| $h(x)$                                                                   | <code>tanhDiv x</code>                                                                                  | totalized $\tanh(x)/x$ , globally the divided slope<br><code>tanhDiv_eq_dslope</code>                                    |
| $p(x)$                                                                   | <code>activationProbability x</code>                                                                    | $1 - h(x)$ , with $p(0) = 0$ and the proved degree-eight finite Taylor jet                                               |
| $D_w(t)$                                                                 | article-only abbreviation; see <code>tendsto_tsum_activationProbability_mul_div_sq</code>               | square-summable activation field $\sum_i p(w_i t)$ ; no named Lean definition                                            |
| $d_q(t)$                                                                 | <code>geometricActivationDimension q t</code>                                                           | normalized geometric activation dimension                                                                                |
| $d_{\text{dy}}(t)$                                                       | <code>dyadicEffectiveDimension t</code>                                                                 | dyadic specialization $d_{1/2}(t)$                                                                                       |
| $G_q$                                                                    | <code>geometricReciprocalGamma q</code>                                                                 | entire reciprocal-Gamma product for $\ q\  < 1$                                                                          |
| $(a; q)_{\infty, \mathbb{C}}$                                            | <code>complexQPochhammerInf a q</code>                                                                  | total infinite symbol; for fixed $\ q\  < 1$ , an entire function of $a$ with the factor-zero set and simple zeros above |
| $\Gamma_{\langle q \rangle}$                                             | <code>geometricGamma q</code>                                                                           | totalized meromorphic inverse of $G_q$                                                                                   |
| $G_{\text{dy}}, \Gamma_{\text{dy}}$                                      | <code>dyadicReciprocalGamma dyadicGamma</code>                                                          | dyadic reciprocal product and totalized meromorphic inverse                                                              |
| $V_a f$                                                                  | <code>volterraPrimitive a f</code>                                                                      | first oriented primitive at base $a$                                                                                     |
| $L_a^n f$                                                                | <code>iteratedPrimitive n a f</code>                                                                    | literal $n$ -fold primitive                                                                                              |
| $J_a^n f$                                                                | <code>normalizedVolterra n a f</code>                                                                   | normalized Cauchy-kernel primitive                                                                                       |
| $I_{a+}^\alpha f$                                                        | <code>fractionalVolterra</code>                                                                         | total real-order Gamma-normalized Volterra operator                                                                      |
| $I_{a+}^{\alpha+\beta} f = I_{a+}^\alpha I_{a+}^\beta f$                 | <code>fractionalVolterra_add</code>                                                                     | positive orders, ordered endpoint, continuous                                                                            |
| $U_\beta(x)$                                                             | <code>rvachevFractionalPrimitive</code>                                                                 | Banach-valued input causal Rvachev fractional primitive based at $-1$                                                    |
| $I_{0+}^\alpha F(x) = \mathbb{E}[(x - X)_+^\alpha] / \Gamma(\alpha + 1)$ | <code>ProbabilityRepresentation.fractionalVolterra_fabiusReal_eq_uniformProduct_integral_posPart</code> | positive real order, every real endpoint                                                                                 |

| Article                     | Lean                                              | Meaning                                                                                                       |
|-----------------------------|---------------------------------------------------|---------------------------------------------------------------------------------------------------------------|
| affine $I_{a+}^{\alpha}$    | <code>fractionalVolterra_affine</code>            | arbitrary real order, positive scale, ordered endpoint                                                        |
| $I_{a+}^{\alpha+1} g'$      | <code>fractionalVolterra_add_one_deriv</code>     | positive order, continuous primitive, integrable interior right derivative, exact Gamma boundary term         |
| fractional $F_{\text{gl}}$  | <code>fractionalVolterra_add_one_extendedF</code> | $I_{0+}^{\alpha+1} F_{\text{gl}}(x) = 2^{\alpha} I_{0+}^{\alpha} F_{\text{gl}}(x/2)$ , $\alpha > 0, x \geq 0$ |
| $\mathcal{A}_{S,x,w}(a; z)$ | <code>abius_of_nonneg</code>                      | finite filter of unconditionally summable weighted coefficient series                                         |
| $I$                         | <code>finiteAnalyticSeriesFilter</code>           | totalized inverse                                                                                             |
| $c_n$                       | <code>fabiusInv</code>                            | even moment                                                                                                   |
| $d_n$                       | <code>fabius</code>                               | half moment                                                                                                   |
| $P_w(n)$                    | <code>fabius_spec</code>                          | inclusive weighted prefix                                                                                     |
| $W_{b,w}(n)$                | <code>inclusivePrefixSum w n</code>               | weighted scale multiplicity                                                                                   |
| $\nu_b(n)$                  | <code>weightedScaleMultiplicity b w n</code>      | maximal whole-base power exponent                                                                             |
| $\varepsilon_n$             | <code>padicValNat b n</code>                      | signed Thue–Morse value                                                                                       |
| $\text{wt}_2(n)$            | <code>thueMorseSign n</code>                      | binary digit sum                                                                                              |
|                             | <code>binaryWeight n</code>                       |                                                                                                               |

## References

- [1] B. Jessen and A. Wintner, Distribution functions and the Riemann zeta function, *Transactions of the American Mathematical Society* **38** (1935), 48–88.
- [2] J. Fabius, A probabilistic example of a nowhere analytic  $C^{\infty}$ -function, *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete* **5** (1966), 173–174. DOI: 10.1007/BF00536652.
- [3] V. L. Rvachev and V. A. Rvachev, A certain finite function, *Dopovidi Akademii Nauk Ukrain's'koi RSR, Series A* (1971), 705–707, 764. Ukrainian, with English and Russian summaries.
- [4] V. A. Rvachev, Certain functions with compact support and their applications, in *Mathematical Physics and Functional Analysis*, Kharkov (1973), 139–149, 194. Russian.
- [5] V. L. Rvachev and V. A. Rvachev, *Non-classical Methods of Approximation Theory in Boundary Value Problems*, Naukova Dumka, Kiev, 1979. Russian.
- [6] V. A. Rvachev, Compactly supported solutions of functional-differential equations and their applications, *Russian Mathematical Surveys* **45** (1990), no. 1, 87–120. Translated from *Uspekhi Matematicheskikh Nauk* **45** (1990), no. 1, 77–103, 222.
- [7] J. Arias de Reyna, Definición y estudio de una función indefinidamente diferenciable de soporte compacto, *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales de Madrid* **76** (1982), no. 1, 21–38. English translation: *An infinitely differentiable function with compact support: Definition and properties*, arXiv:1702.05442 (2017).
- [8] J. Arias de Reyna, Arithmetic of the Fabius function, *INTEGERS* **18** (2018), Article A51; arXiv:1702.06487, version 3. The version is pinned because a sign in the abstract of one version is a typographical error; see [Section 16](#).

- [9] J. K. Haugland, Evaluating the Fabius function, arXiv:1609.07999 (2016).
- [10] A. Grzegorzczak, On the definitions of computable real continuous functions, *Fundamenta Mathematicae* **44** (1957), 61–77.
- [11] R. M. Corless, G. H. Gonnet, D. E. G. Hare, D. J. Jeffrey, and D. E. Knuth, On the Lambert  $W$  function, *Advances in Computational Mathematics* **5** (1996), 329–359.
- [12] P. Flajolet, X. Gourdon, and P. Dumas, Mellin transforms and asymptotics: harmonic sums, *Theoretical Computer Science* **144** (1995), 3–58.
- [13] N. J. Fine, Binomial coefficients modulo a prime, *American Mathematical Monthly* **54** (1947), 589–592.
- [14] J.-P. Allouche and J. Shallit, *Automatic Sequences: Theory, Applications, Generalizations*, Cambridge University Press, 2003.
- [15] G. E. Andrews, *The Theory of Partitions*, Encyclopedia of Mathematics and its Applications 2, Addison–Wesley, 1976.
- [16] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, second edition, Encyclopedia of Mathematics and its Applications 96, Cambridge University Press, 2004.
- [17] V. Kac and P. Cheung, *Quantum Calculus*, Universitext, Springer, 2002.
- [18] NIST Digital Library of Mathematical Functions, Chapter 17,  *$q$ -Hypergeometric and Related Functions*, <https://dlmf.nist.gov/17>.
- [19] V. Reshetnikov, A conjectured closed form for the Fabius function at dyadic rationals, Mathematics Stack Exchange, question 3283519, 5 July 2019.
- [20] V. Reshetnikov, Extracting an asymptotic from a sequence defined by a recurrence relation, Mathematics Stack Exchange, question 4354350, 11 January 2022. Compact Fabius formula in the author’s 13 January comment on the answer by user321120; <https://math.stackexchange.com/q/4354350>.
- [21] V. Reshetnikov, A conjecture about certain values of the Fabius function, MathOverflow, question 261649, 7 February 2017. The integrality question answered in [8].
- [22] V. Reshetnikov, *ProveIt: Analysis/FabiusFunction*, Lean 4 formal development, integrated Cauchy source checkpoint a81a13b46b4c7d525f273a52c9018a6d281c5ef2, consulted 28 August 2026. <https://github.com/VladimirReshetnikov/ProveIt/tree/a81a13b46b4c7d525f273a52c9018a6d281c5ef2/Analysis/FabiusFunction>. Fractional affine/order-raising calculus, causal support-truncated Rvachev primitives, first fractional Fabius–Rvachev shifts, unconditionally summable analytic-series filters, and the finite complex Thue–Morse product bridge were also compiler-validated at checkpoint 149332f9d. <https://github.com/VladimirReshetnikov/ProveIt/tree/149332f9d/Analysis/FabiusFunction>. The interval-restricted Galois connection, measurable-free Banach-valued and pointwise- $C^1$  inverse layer-cake APIs, and the real- or complex-weighted arbitrary- $L^1$ /real-absolutely-continuous Fabius wrapper were compiler-validated at checkpoint 37eab3c56cb28ab702878d71f7d1a87791adea44. <https://github.com/VladimirReshetnikov/ProveIt/tree/37eab3c56cb28ab702878d71f7d1a87791adea44/Analysis/FabiusFunction>. The current integrated source additionally contains the generic affine finite-measure Cauchy calculus, the general

geometric-uniform transform and power hierarchies, the atom-exact CDF/survival higher-power formulas, the named dyadic unit-interval Stieltjes DDE, both complex-slit adjacent-power recurrences, and the exact centered Rvachev DDE/all-order Thue–Morse orbit. Its inversion layer is an integrated interval Stieltjes–Perron theorem, not a pointwise or nontangential boundary/Plemelj formula. A later current-tree Bell tranche, inspected 30 August 2026, consists of `CompleteHomogeneousBell.lean`, `LambertPhaseLockedBell.lean`, and `FabiusLambertPhaseExtractionBell.lean`. The declarations cited in the text are exact finite algebraic rewrites and add no convergence, sign, or asymptotic estimate; no aggregate commit hash is attributed to this later tranche here. The still later current-tree Lambert branch-point tranche, compiled 30 August 2026, consists of `LambertWBranchPointGeometry.lean` and `LambertWBranchPointAsymptotics.lean`. Its exact public inventory is eight theorems and, respectively, one noncomputable definition plus eight theorems. The declaration-by-declaration crosswalk above is not attributed to an aggregate commit hash; it closes the leading raw-branch geometry and square-root law only, with the stated remainder, higher-series, wrapper, and generic-shape boundaries left open. The generic three-theorem continuous-CDF bridge, the then two-definition/twenty-five-theorem geometric-uniform CDF and smooth-density interface, the half-base Fabius/Rvachev bridges, and the one-definition/eight-theorem formal implicit-root engine are included in compiler-validated checkpoint `e6f535e5d`. The CDF module’s four later nonpositive-parameter diagnostics, which bring its current surface to two definitions and twenty-nine theorems, are inventoried above. The latter engine is generic nonmonic Hensel lifting over complete adic commutative rings and supplies no concrete inverse-Fabius germ or analytic-convergence theorem. <https://github.com/VladimirReshetnikov/ProveIt/tree/e6f535e5d/Analysis/FabiusFunction>. The generic affine independent-copy probability operator, its characteristic-function factorization and finite iteration, its contractive probability-fixed-point uniqueness theorem, and the geometric-uniform uniqueness specialization are included in compiler-validated checkpoint `d312c0603`. <https://github.com/VladimirReshetnikov/ProveIt/tree/d312c0603/Analysis/FabiusFunction>.

The seven-module activation tranche, including the divided-slope totalization and the finite degree-eight activation Taylor jet, is recorded at immutable source checkpoint `a345425d21d90e680bf15e34093af42c69c08a83`. <https://github.com/VladimirReshetnikov/ProveIt/tree/a345425d21d90e680bf15e34093af42c69c08a83/Analysis/FabiusFunction>.