

# artemide ver.3.00

Alexey A. Vladimirov  
September 30, 2024

User manual for **artemide** package, which evaluated TMDs and related cross-sections.

**Manual is updating.**

This work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/> or send a letter to Creative Commons, PO Box 1866, Mountain View, CA 94042, USA.

If you use the **artemidev.1.??**, please, quote [? ].

If you use the **artemidev.2.??**, please, quote [? ].

If you use the **artemidev.2.06**, please, quote [? ].

If you find mistakes, have suggestions or questions, please, write to:

Alexey Vladimirov: [alexeyvl@ucm.es](mailto:alexeyvl@ucm.es) or [vladimirov.aleksey@gmail.com](mailto:vladimirov.aleksey@gmail.com)

**Stable Repository:** <https://github.com/VladimirovAlexey/artemide-public>  
**Dev Repository:** <https://github.com/VladimirovAlexey/artemide-development>

## Contents

|   |    |
|---|----|
| <b>Glossary</b>   | 4  |
| <b>I. General structure and user input of artemide</b>                                    | 5  |
| A. General concept  | 5  |
| B. Organization of TMD factorized cross-section and its implementation in <b>artemide</b> | 5  |
| C. User defined functions and options   | 7  |
| D. Installation   | 8  |
| E. Python interface: <b>harpy</b>   | 8  |
| F. Constants file and version compatibility   | 9  |
| <b>II. Theory</b>   | 11 |
| A. Definition of TMD distributions  | 11 |
| B. Evolution of TMD distributions   | 11 |
| 1. Equipotential lines & $\zeta$ -prescription  | 12 |
| 2. Exact solution for evolution to special line   | 13 |
| C. Expressions for evolution functions in <b>artemide</b>                                 | 14 |
| 1. Fixed order RAD:   | 14 |
| 2. Resummed RAD:  | 14 |
| 3. Fixed order $\zeta$ -line:   | 15 |
| 4. Resummed $\zeta$ -line:  | 15 |
| 5. Exact $\zeta$ -line:   | 16 |
| <b>III. aTMDe_setup module</b>  | 17 |
| A. aTMDe_Setup  | 18 |
| <b>IV. aTMDe_control module</b>   | 19 |
| A. Passing non-perturbative parameters  | 19 |
| <b>V. QCDeinput module</b>  | 21 |
| A. Initialization   | 21 |

|  |    |
|--|----|
| B. LHA_alpha module                                | 21 |
| C. LHA_PDF module                                  | 22 |
| <b>VI. EWinput module</b>                          | 23 |
| A. $\alpha_{\text{QED}}$                           | 23 |
| <b>VII. TMD evolution modules</b>                  | 24 |
| 1. INI-parameters                                  | 24 |
| 2. Functions                                       | 25 |
| A. TMD_AD module                                   | 26 |
| B. TMDR_model                                      | 27 |
| <b>VIII. Modules computing TMD distributions</b>   | 28 |
| 1. INI-parameters                                  | 29 |
| 2. Functions                                       | 30 |
| A. _model modules                                  | 33 |
| B. _OPE modules (tw2)                              | 33 |
| 1. Functions                                       | 34 |
| 2. Model parameters.                               | 35 |
| 3. Scale variation.                                | 35 |
| 4. Grid  | 35 |
| 5. Technical notes                                 | 36 |
| C. _OPE modules (tw3)                              | 38 |
| 1. Functions                                       | 38 |
| 2. Model parameters.                               | 39 |
| 3. Scale variation.                                | 39 |
| 4. Grid  | 39 |
| D. Fourier to $k_T$ space                          | 39 |
| E. TMM   | 42 |
| <b>IX. TMDs module</b>                             | 43 |
| A. Definition of low-scales                        | 45 |
| B. Evaluating TMDs                                 | 45 |
| C. Products of TMDs                                | 45 |
| D. Theoretical uncertainties                       | 46 |
| <b>X. TMDF module</b>                              | 47 |
| A. Initialization                                  | 47 |
| B. Evaluating Structure functions                  | 47 |
| C. Enumeration of structure functions              | 49 |
| D. Tests   | 54 |
| <b>XI. TMDX_DY module</b>                          | 56 |
| A. Initialization                                  | 57 |
| B. The parameters of cross-section                 | 57 |
| C. Cross-section evaluation                        | 58 |
| D. LeptonCutsDY                                    | 59 |
| E. Variation of scale                              | 59 |
| F. Options for evaluation of DY-like cross-section | 59 |
| 1. $\pi^2$ -resummation                            | 59 |
| 2. Power corrections                               | 59 |
| G. Enumeration of processes (LP case)              | 60 |
| H. Enumeration of processes (KPC case)             | 61 |
| <b>XII. TMDX_SIDIS module</b>                      | 63 |
| A. Cross-section evaluation                        | 64 |
| B. Enumeration of processes                        | 64 |
| C. Kinematic cuts                                  | 65 |
| D. Power corrections                               | 66 |

|   |    |
|---|----|
| E. Bin-integration routines                           | 66 |
| F. SIDIS theory                                       | 66 |
| 1. Kinematics   | 67 |
| 2. Cross-section                                      | 67 |
| <b>XIII. TMDs_inKT module</b>                         | 69 |
| A. Grids  | 71 |
| <b>XIV. KPC in TMD factorization</b>                  | 72 |
| A. Realization of convolution                         | 72 |
| B. Enumeration of factors processes                   | 72 |
| <b>XV. APPENDICES</b>                                 | 74 |
| A. Equivalence between process-enumeration in ver.< 3 | 74 |
| <b>XVI. Version history</b>                           | 77 |

## Glossary

**TMD = Transverse Momentum Dependent**

**PDF = Parton Distribution Function**

**FF = Fragmentation Function**

**NP = Non-perturbative**

**DY = Drell-Yan process**

**SIDIS = Semi-Inclusive Deep-Inelastic Scattering**

**LO = Leading order** (in QCD perturbation theory)

**NLO = Next-to-Leading order**, (in QCD perturbation theory)

**N<sup>2</sup>LO = Next-to-Next-to-Leading order (etc)**, (in QCD perturbation theory)

**LP = Leading power** (in power expansion of TMD factorization)

**NLP = Next-to-Leading power (etc)**, (in power expansion of TMD factorization)

**KPC = Kinematic power corrections**

## I. GENERAL STRUCTURE AND USER INPUT OF ARTEMIDE

### A. General concept

The **artemide** is a package of Fortran modules for calculation in TMD factorization framework. It has a modular structure, where each module is responsible for evaluation of some theory construct. For instance: a TMD distribution, a TMD evolution factor, cross-section. Each module produces a single function, which is a composite of integrals/products of lower-level functions. Thus, each level of operation can be used as is, in other programs. The highest-level task is the evaluation of cross-section within the TMD factorization theorem, including all needed integrations, and factors, i.e., such that it can be directly compared to the data. It also includes several tools for analysis of the obtained values, such as variation of scales, search for limiting parameters, etc. The theory structure of **artemide** is discussed in the next section. The dependency structure of modules is presented in fig.1.

Initially the **artemide** project was created for pure theoretical games. Since the beginning **artemide** appears to be successful also in the phenomenology, (and nowadays it is mostly used for it). Nonetheless, conceptually **artemide** is build as the theory playground, and will continue to be developed in this direction. That is why its architecture is not very optimal from the pure numerical point of view. In fact, there are several ways to optimize the code, melting together some modules and structures, but in this case **artemide** will lose its theoretical cleanness. Also **artemide** contains a lot of “unpractical” and rare options, and possibility to control each parameter. The positive side is the possibility to easily implement the new theory founding and check them, which is regularly done.

**Wide spectrum of application of artemide code makes it difficult to create a convenient interface.** Moreover, at the current stage of development, I prioritize the quality of computation, to the user interface. So, the interface is changing from version to version and often is not compatible with earlier versions. It slowly converges to the (almost) perfect shape – convenient for a wider community. If you have a particular task and not sure how to operate with **artemide** in this case, better write an e-mail.

**The main rule** (implemented in ver.2.00) each part encapsulates its theory parameters. It does not affect/change/interact with other modules, except requests for functions. A change of a parameter in a module can require a change in another module for consistency (however, I attempt to avoid such cases). Then each parameter must be changed individually in each module. However, the module **aTMDcontrol** does it automatically. So, I suggest to use **aTMDcontrol** to avoid possible inconsistencies.

**Historical note:** In versions before ver.2.00 this rule was not implemented. I tried to make connections between modules such that they automatically control consistency. However, at some moment (after inclusion of many hadrons and different types of cross-sections, and different orders) it became practically tough to keep such a system. So, I rearrange some modules (e.g. variation of  $c_4$  scale was in **TMDs**, while it related to definition of TMD itself), removed connections between modules (no link for change of NP parameters, etc.), and introduce **aTMDcontrol**.

### B. Organization of TMD factorized cross-section and its implementation in artemide

The ultimate goal of the **artemide** is to evaluate the observables in the TMD factorization framework, such as cross-section, asymmetries, etc. The general structure of the TMD factorized *fully differential* cross-section is

$$\frac{d\sigma}{dX} = d\sigma(q_T) = prefactor \times F, \quad (1.1)$$

where *prefactor* a process-dependent and experiment-dependent prefactor, and  $F$  is the reduced structure function. Example, for the photon induced Drell-Yan process one has for  $d\sigma/dq_T$

$$prefactor = \frac{4\pi}{9sQ^2} |C_V(Q, \mu_H)|^2 \mathcal{P}(\text{cuts}),$$

$$F = \int \frac{bdb}{2} J_0(bq_T) \sum_f |e_f|^2 F_f^f(x_A, b; \mu_H, \zeta_A) F_{\bar{f}}^f(x_B, b; \mu_H, \zeta_B),$$

where  $\mathcal{P}(\text{cuts})$  is the weighting factor for fiducial cuts. For specific expression we refer in corresponding sections of this text. The structure function  $F$  is generally defined as

$$F(q_T, x_1, x_2; \mu, \zeta_1, \zeta_2) = \int \frac{bdb}{2} b^n J_n(bq_T) \sum_{ff'} z_{ff'} F_1^f(x_1, b; \mu, \zeta_1) F_2^{f'}(x_2, b; \mu, \zeta_2), \quad (1.2)$$



FIG. 1: Modules of **artemide**: purpose and dependencies

where  $F_{1,2}$  are TMD distributions (of any origin and polarization),  $z_{ff'}$  is the process dependent flavor mixing factor. The number  $n$  is also process dependent, e.g. for unpolarized observables it is  $n = 0$ , while for SSA's it is  $n = 1$ . Evaluation structure functions  $F$  is performed in the module **TMDF**. The evaluation of cross-sections is performed in the modules **TMDX**. The TMD distributions are evaluated by the module **TMDs** and related submodules.

In this way, the computation of TMD-factorized cross-section can be naturally split into ordered parts. Each part is evaluated in corresponding module of **artemide**. See example, of evaluation scheme in fig.2.

### C. User defined functions and options

The **artemide** package has been created such that it allows to control **each aspect** of TMD factorization theorem. The TMD factorization has a large number of free, and “almost-free” parameters. It is a generally difficult task to provide a convenient interface for all these inputs. I do my best to make the interface convenient; however, some parts (e.g., setup of  $f_{NP}$ ) could not be simpler (at least within FORTRAN). Also, take care that **artemide** is evolving, and I try to keep back compatibility, but it is not the main option.

Starting from ver.2.0, **artemide** uses the text initialization file, which contains all required information on static parameters for a given setup. Throughout the text I call this file **constants-file**.

The user has to provide (or **use the default values**) the set of parameters, that control various aspects of evaluation. It includes PDF sets,  $f_{NP}$ , perturbative scales, parameters of numerics, non-QCD inputs, etc. There are three input sources for statical parameters.

**General parameters:** These are working parameters of **artemide**, such as amount of output, tolerance of integration routines, number of NP parameters, type of used evolution, gridding parameters, triggering of particular contributions, etc. There are many of them, and typically they are unchanged. These are set in **constants-file**. **Changes do not require recompilation.**

**External physics input:** It includes the definition of  $\alpha_s$ , collinear PDFs, and other distributions. Twist-2 distributions are taken from LHAPDF [? ], with routines defined in **QCDinput** module. For non-QCD parameters, e.g.  $\alpha_{QED}$ , SM parameters, there is a module **EWinput**. These are set in **constants-file**. **Changes do not require recompilation.**

**NP model:** The NP model consists in NP profiles of TMD distributions, NP model for large- $b$  evolution, selection of scales  $\mu$ , etc. These parameter and functions enter nearly each low level module. The code for corresponding functions is provided by user, in appropriate files, which are collected in the subdirectory **src/Model**. The name of files are shown on diagram in colored blobs adjusted to the related module. **Changes require recompilation.**

#### Comments:

- **IMPORTANT:** Each module is initialized individually via **constants-file**. So, each module can be used independently on the full package, given proper section of **constants-file** and submodules (see diagram). However, unless you understand what is going on, it is recommended to use **aTMDe\_setup** module for creation of **constant-file**, and **aTMDe\_control** module for proper control, initialization and operations of sub-modules.
- **constants-file** can be saved and used in future to reproduce setup. I try to keep compatibility between these files.
- **constants-file** is created and modified within **aTMDe\_setup** module. It could be also modified manually.
- NP functions are typically defined with a number of numeric parameters. The value of these parameters could be changed without restart (or recompilation) of the **artemide** by appropriate command. E.g. (**call TMDs\_SetNPParameters(lambda)**) on the level of **TMDs** module. See sections of corresponding modules.
- The number of parameters in the model for each module is set in **constants-file**.

- The directory `/Model` together with `constants-file` are convenient to keep as they are. They contain full information about particular evaluation, and thus results can be always reproduced (at least within the same version of `artemide`). I provide results of our extraction as such directories.
- Before ver.2.0, the interface was different and chaotic.

## D. Installation

Download and unpack `artemide`. The actual code is in the `/src`. Check the `makefile`. **You must provide options `FC` and `FOPT`, which are defined in the top of it.** `FC` is the FORTRAN compiler, `FOPT` is additional options for compiler (e.g. linking to LHAPDF library).

There is no actual installation procedure, there is just compilation. If model, inputs, etc. are set correctly (typical problem is linking to LHAPDF, be sure that it is installed correctly), then `make` compiles the library. The result are object files (`*.o`) (which are collected in `/obj`) and module files (`*.mod`) (which are collected in `/mod`).

The test of current compilation could be performed by `make test`. It compiles program `test.f90` from `/Prog` and run it (default test uses `NP31_nnlo_as_0118` set from LHAPDF, check that it is present in your LHAPDF installation). Program `test` runs some elementary code with minimum input. Output is shown later

Next, do your code, include appropriate modules of `artemide`, and compile it together with object-files (do not forget to add proper references to module files `-I/mod`). It should work! Linking could be done automatically if you call for `make program TARGET=...`, where ... is the name of the file with the code.

```
artemide.control: initialization done.
uTMDPDF: Grid is built ( 250 x 750) calc.time= 0.53s.
Calculating some values for cross-section one-by-one (DY around Z-boson peak, ATLAS 8TeV kinematics)
ptMin -- ptMax xSec
1.0000000000000000 -- 3.0000000000000000 48.499041273610260
3.0000000000000000 -- 5.0000000000000000 57.189916866533792
5.0000000000000000 -- 7.0000000000000000 49.321931214709110

Now the same by list
It must be faster since you use OPENMP
result: 48.499041273610260 57.189916866533792 49.321931214709110
-----
The programm evaluation took 8.8279999999999994 sec. (CPU time)
The programm evaluation took 6.1287177319172770 sec. (wallclock time)

If you do not like so many terminal messages check the parameter outputlevel in constants file.
Not forget to cite artemide [1706.01473]
-----
```

## E. Python interface: harpy

For simplicity of data analysis the `artemide` has a python interface, called `harpy` (linking is made by `f2py` library). It is not possible to interfacing the `artemide` directly to python since `artemide` is made on fortran95. `Artemide` uses some features of Fortran95, such as interfaces, and indirect list declarations, which are alien to python. Also I have not found any convenient way to include several dependent Fortran modules in `f2py` (if you have suggestion just tell me). Therefore, I made a wrap module `harpy.f90` that call some useful functions from `artemide` with simple declarations. It contains limited set of functions useful for phenomenology, and definitely cannot replace the FORTRAN interface for deep studies.

The compilation of `harpy` is slightly more complicated.

1. In the `makefile` check the variable `Fpath` (in the top part of the file). Put-in the full path for fortran compiler. It is needed by `f2py`.
2. (optional, does not work on Mac (?)) Run `make harpy-signature`. This will create an interface file (`artemide.pyf` for all functions in the `harpy.f90`. This file is already provided in the distribution, so if you did not change `harpy.f90`, you better skip this step. For some reason, it does not work on Mac.
3. Run `make harpy`. It compiles `harpy.f90` with interface `artemide.pyf` linking to `artemide`. The result is `artemide.so`.

All files are in `/harpy`. Link it to python and work. There is also extra python `harpy.py` which has several most important function.



## F. Constants file and version compatibility

Constants files contains the list of work-flow parameters, such as physical constants, LHA grid names, numerical setup, etc. Each constants file has version number (does not coincide with the version of artemide).

On the module initialization the constants file is read, parameter are setup. Thus, the version of constants file should be adjusted to version of artemide *if you run modules separately*.

The workflow is deferent of the initialization of **artemide** is made by the **artemide\_control**. In this case, the **artemide** creates a copy of the initial constants-file (**aTMDe\_temporary**), and (in the case the versions do not match) fill the absent options by default parameter. After it the lower-level modules are initialized by **aTMDe\_temporary**. Naturally, in this case the versions should match (if not something is wrong with installation).

You can update the **constants-file** to an actual version by

```
make update TARGET=...
```

where ... is the path to **constants-file** to be updated. The updated version will have all content of the original file, + new options setup by default setting.




FIG. 2: Evaluation of DY cross-section by artemide

## II. THEORY

**This section is under construction.**

In this section, the details of theoretical input coded in `artemide` is given. This section does not pretend to be a comprehensive review of TMD factorization. Only superficial details and references to particular realizations are given. For detailed and accurate description of the theory see specialized literature.

### A. Definition of TMD distributions

**TO BE WRITTEN.** In this section, there will be operator definition of TMD distributions, and details on way the perturbative matching and NP-modeling is realized in `artemide`.

$$F_f(x, b) = \int_x^1 \frac{dz}{z} C_{f \leftarrow f'}(z, b^*, c_4 \mu_{\text{OPE}}) f_{f'}\left(\frac{z}{x}, c_4 \mu_{\text{OPE}}\right) f_{NP}^f(x, z, b, \{\lambda\}), \quad (2.1)$$

where  $f_f(x, \mu)$  is PDF of flavor  $f$ ,  $C$  is the coefficient function in  $\zeta$ -prescription,  $f_{NP}$  is the non-perturbative function. The variable  $c_4$  is used to test the scale variation sensitivity of the TMD PDF. The NNLO coefficient functions used in the module were evaluated in [?] (please, cite it if use).

**PLAN TO ADD**

- Definition of TMD distributions, operators, Lorenz structures, coordinate and momentum space.
- Perturbative matching, and  $\zeta$ -prescription
- NP-modeling

### B. Evolution of TMD distributions

The detailed theory is given in the article [?]. The NLO rapidity anomalous dimension has been evaluated in [?]. The NNLO rapidity anomalous dimension has been evaluated in [? ?].

The evolution of TMD distribution of any kind is given by the following pair of equations

$$\mu^2 \frac{d}{d\mu^2} F_{f \leftarrow h}(x, b; \mu, \zeta) = \frac{\gamma_F^f(\mu, \zeta)}{2} F_{f \leftarrow h}(x, b; \mu, \zeta), \quad (2.2)$$

$$\zeta \frac{d}{d\zeta} F_{f \leftarrow h}(x, b; \mu, \zeta) = -\mathcal{D}^f(\mu, b) F_{f \leftarrow h}(x, b; \mu, \zeta), \quad (2.3)$$

where  $F_{f \leftarrow h}$  is the TMD distribution (TMDPDF or TMDFF) of the parton  $f$  in hadron  $h$ . The function  $\gamma_F(\mu, \zeta)$  is called the TMD anomalous dimension and contains both single and double logarithms. The function  $\mathcal{D}(\mu, b)$  is called the rapidity anomalous dimension. TMD and rapidity anomalous dimensions have not unified notation in the literature, for comparison of notation see table I in ref.[?]. The only important quantum number for TMDs is the color representation the initiating parton, which is tied to the parton flavor, namely, quark (fundamental representation) or gluon (adjoint representation). However, as the TMD evolution does not mix the flavors and for simplicity of notation, we omit the flavor index  $f$  in this section, unless it is important.

The uniqueness of solution for the coupled system (2.2)-(2.3) is guaranteed by *the integrability condition*

$$\zeta \frac{d}{d\zeta} \gamma_F(\mu, \zeta) = -\mu \frac{d}{d\mu} \mathcal{D}(\mu, b), \quad (2.4)$$

The mutual dependence can be worked out explicitly due to the fact that the ultraviolet divergences of the TMD operator partially overlap with the rapidity divergences, (see e.g.[? ?]),

$$\zeta \frac{d}{d\zeta} \gamma_F(\mu, \zeta) = -\Gamma(\mu), \quad (2.5)$$

$$\mu \frac{d}{d\mu} \mathcal{D}(\mu, b) = \Gamma(\mu), \quad (2.6)$$

where  $\Gamma$  is the (light-like) cusp anomalous dimension. The equation (2.5) entirely fixes the logarithm dependence of the TMD anomalous dimension, which reads

$$\gamma_F(\mu, \zeta) = \Gamma(\mu) \ln \left( \frac{\mu^2}{\zeta} \right) - \gamma_V(\mu). \quad (2.7)$$

The anomalous dimension  $\gamma_V$  refers to the finite part of the renormalization of the vector form factor. In contrast, the equation (2.6) cannot fix the logarithmic part of  $\mathcal{D}$  entirely, but only order by order in perturbation theory, because the parameter  $\mu$  is also responsible for the running of the coupling constant.

The solution of eq. (2.2)-(2.3) can be written as

$$F(x, b; \mu_f, \zeta_f) = R[b; (\mu_f, \zeta_f) \rightarrow (\mu_i, \zeta_i)] F(x, b; \mu_i, \zeta_i), \quad (2.8)$$

where  $R$  is the TMD evolution factor. The general form of the evolution factor is

$$R[b; (\mu_f, \zeta_f) \rightarrow (\mu_i, \zeta_i)] = \exp \left[ \int_P \left( \gamma_F(\mu, \zeta) \frac{d\mu}{\mu} - \mathcal{D}(\mu, b) \frac{d\zeta}{\zeta} \right) \right], \quad (2.9)$$

where  $(\mu_f, \zeta_f)$  and  $(\mu_i, \zeta_i)$  refer respectively to a final and initial set of scales. Here, the  $\int_P$  denotes the line integral along the path  $P$  in the  $(\mu, \zeta)$ -plane from the point  $(\mu_f, \zeta_f)$  to the point  $(\mu_i, \zeta_i)$ . The integration can be done on an arbitrary path  $P$ , and the solution is independent on it, thanks to the integrability condition eq. (2.4).

The TMD evolution factor  $R$  obeys the transitivity relation

$$R[b; (\mu_1, \zeta_1) \rightarrow (\mu_2, \zeta_2)] = R[b; (\mu_1, \zeta_1) \rightarrow (\mu_3, \zeta_3)] R[b; (\mu_3, \zeta_3) \rightarrow (\mu_2, \zeta_2)], \quad (2.10)$$

where  $(\mu_3, \zeta_3)$  is arbitrary point in  $(\mu, \zeta)$ -plane and the point inversion property

$$R[b; (\mu_1, \zeta_1) \rightarrow (\mu_2, \zeta_2)] = R^{-1}[b; (\mu_2, \zeta_2) \rightarrow (\mu_1, \zeta_1)]. \quad (2.11)$$

These equations are the cornerstones of the evolution mechanism, since they allow an universal definition of the non-perturbative distributions and the comparison of different experiments.

Different realizations of TMD evolution (if they are compatible with the factorization theorem) can be casted into selected of different paths of integration **ADD PICTURE**. All choices are equivalent, up to next-to-given order perturbative corrections, treatment of NP parts, and implementation of matching for TMD distributions. In **artemide** several paths are presented, however, the matching of TMD distribution to collinear distributions is done in  $\zeta$ -prescription.

### 1. Equipotential lines & $\zeta$ -prescription

The  $\zeta$ -prescription is based on consideration of equipotential lines for evolution equations (2.2)-(2.3) (or lines of null-evolution, for description of evolution potential and its properties see [? ]). An equipotential line  $\ell(b) = (\mu, \zeta)$  is determined by values of anomalous dimensions  $\gamma_F$  and  $\mathcal{D}$  at given  $b$ . The equation for line  $\ell(b)$  can be written in the form  $\zeta = \zeta(\mu, b)$ . In this parameterization the for function  $\zeta(\mu, b)$  is

$$-\Gamma(\mu) \ln \left( \frac{\zeta(\mu, b)}{\mu^2} \right) - \gamma_V(\mu) = 2\mathcal{D}(\mu, b) \frac{d \ln \zeta(\mu, b)}{d \ln \mu^2}. \quad (2.12)$$

Among set of equipotential lines there is a single special equipotential line, which passes thorough the saddle point of the evolution field. The saddle point  $(\mu_{\text{saddle}}, \zeta_{\text{saddle}})$  is defined by the equations

$$\mathcal{D}(\mu_{\text{saddle}}, b) = 0, \quad \gamma_V(\mu_{\text{saddle}}, \zeta_{\text{saddle}}) = 0. \quad (2.13)$$

Note, that the value of  $\mu_{\text{saddle}}$  depends on value of  $b$ , and on NP model for  $\mathcal{D}$ . In general it could be found only numerically. In typical models  $\mu_{\text{saddle}}$  is decreasing monotonous function of  $b$ . At certain large values of  $b$  it passes though the Landau pole, and its value could not be determined. This line we denote as

$$\text{special equipotential line} = \zeta_\mu(b), \quad (\mu_{\text{saddle}}, \zeta_{\text{saddle}}) \in \zeta_\mu(b). \quad (2.14)$$

By definition a TMD distribution is the same for all points of equipotential line. In particular it means that there is no dependence on variable  $\mu$ , since a variation of  $\mu$  is compensated by appropriate variation in  $\zeta(\mu)$ .

**Important note:** The presence of NP part in  $\mathcal{D}$  makes determination of special line involved. We distinguish three “realizations” of the special line. The exact which is determined by equation (2.12) and boundary conditions (2.14) (its values can be calculated by function **zetaSL**). The perturbative realization where NP part is dropped (its values can be calculated by functions **zetaMUp** and **zetaMUp**). And user-defined which is given by user in **TMDR\_model.f90** file.

## 2. Exact solution for evolution to special line

There is a possibility to define the evolution to the optimal equipotential line in perturbation theory for any NP input, at any  $b$ . Instead of problematic variable  $b$  one uses the function  $\mathcal{D}(\mu, b)$  as a variable. With the ansatz  $\zeta_\mu = \mu^2 \exp(-g(\mu, b))$  we rewrite equation (2.12) as

$$2\mathcal{D}(\mu, b) \left( 1 - \frac{dg(\mu, b)}{d \ln \mu^2} \right) - \Gamma(\mu)g(\mu, b) + \gamma_V(\mu) = 0. \quad (2.15)$$

or

$$2\mathcal{D} \left( 1 - \frac{\partial g(\mu, \mathcal{D})}{\partial \ln \mu^2} - \frac{\Gamma(\mu)}{2} g'(\mu, \mathcal{D}) \right) - \Gamma(\mu)g(\mu, \mathcal{D}) + \gamma_V(\mu) = 0, \quad (2.16)$$

where  $g' = dg/d\mathcal{D}$ . The boundary condition (2.13) is transformed to the condition that  $g$  is regular at  $\mathcal{D} \rightarrow 0$ . Note, that this condition does not depend on  $b$  and thus determines the function unambiguously at any  $b$ , even if saddle point is behind the Landau-pole values.

The equation (2.16) is simplified further with introduction of new function  $\tilde{g}(a_s, \mathcal{D}) = \mathcal{D}g(\mu, \mathcal{D})$ :

$$2\mathcal{D} + 2\beta(a_s) \frac{\partial \tilde{g}(a_s, \mathcal{D})}{\partial a_s} - \Gamma(a_s) \tilde{g}'(a_s, \mathcal{D}) + \gamma_V(a_s) = 0, \quad (2.17)$$

with boundary condition  $\tilde{g}(a_s, 0) = 0$ . The general solution reads

$$\tilde{g}(a_s, \mathcal{D}) = -\frac{\mathcal{D}}{2} \int^{a_s} \frac{da}{\beta(a)} - \int^{a_s} da \frac{\gamma_V(a)}{2\beta(a)} - \int^{a_s} da \frac{\Gamma(a)}{2\beta(a)} \int^a \frac{da'}{\beta(a')} + \Phi \left( \mathcal{D} + \int^{a_s} da \frac{\Gamma(a)}{2\beta(a)} \right), \quad (2.18)$$

where  $\Phi$  is a solution of the following transcendental equation

$$\Phi \left( \int^{a_s} da \frac{\Gamma(a)}{2\beta(a)} \right) = \int^{a_s} da \frac{\gamma_V(a)}{2\beta(a)} + \int^{a_s} da \frac{\Gamma(a)}{2\beta(a)} \int^a \frac{da'}{\beta(a')}. \quad (2.19)$$

Unfortunately, this equation is practically impossible to solve for higher than NNLL anomalous dimensions. It is straightforward to show that at large- $\mathcal{D}$  the solution is

$$\tilde{g}(a_s, \mathcal{D} \rightarrow \infty) = \mathcal{D} \int^{a_s} \frac{-1}{\beta(a)} da + \dots \quad (2.20)$$

Note that asymptotic is defined up to a constant.

The function  $g$  is convenient to derive as perturbative expansion

$$g(\mu, \mathcal{D}) = \frac{1}{a_s(\mu)} \sum_{n=0}^{\infty} a_s^n(\mu) g_n(\mathcal{D}). \quad (2.21)$$

We have found

$$g_0 = \frac{e^{-p} + p - 1}{\beta_0 p}, \quad (2.22)$$

$$g_1 = g_0 \left( \frac{\beta_1}{\beta_0} - \frac{\Gamma_1}{\Gamma_0} \right) + \frac{\gamma_1}{\gamma_0} - \frac{\beta_1}{2\beta_0^2} p, \quad (2.23)$$

$$g_2 = g_0 \frac{\beta_2 \Gamma_0 - \beta_1 \Gamma_1}{\beta_0 \Gamma_0} + \frac{chp - 1}{p} \frac{\beta_0 \Gamma_1^2 - \beta_0 \Gamma_0 \Gamma_2 + \beta_1 \Gamma_0 \Gamma_1 - \beta_2 \Gamma_0}{\beta_0^2 \Gamma_0^2} + \frac{e^p - 1}{p} \frac{\Gamma_0 \gamma_2 - \Gamma_1 \gamma_1}{\Gamma_0^2}, \quad (2.24)$$

where  $p = 2\beta_0 \mathcal{D} / \Gamma_0$ .

There is an additional problem that appears at NNNLO, namely, the coefficient  $g_3$  is negative. It results to a singular solution for  $\zeta_\mu$  at large- $\mathcal{D}$ . It indicates that the series has bad convergence properties. This problem occurs for very large  $\mathcal{D} > 0.9 - 1.1$ . Currently to by-pass this problem the 4-loop exact  $\zeta$ -line is not used (3-loop expression is used instead). For small- $\mathcal{D}$  the difference is negligible.

The solution for the evolution from  $(\mu, \zeta) \rightarrow (\mu, \zeta_\mu)$  is

$$R[b; (\mu, \zeta) \rightarrow (\mu, \zeta_\mu)] = \exp \left( -\mathcal{D}(\mu) \ln \left( \frac{\zeta}{\mu^2} \right) - \mathcal{D}(\mu, b) g(\mu, \mathcal{D}) \right). \quad (2.25)$$

### C. Expressions for evolution functions in artemide

Here are the expression for various evolution functions how they are written in **artemide** in TMDR-module.

#### 1. Fixed order RAD:

**Dpert(mu,bT,f):** returns  $\mathcal{D}^f(b, \mu)$  [real(dp)]  
**mu, bT** [real(dp)] scales in GeV, **f**[integer] flavour.  
 Order is controlled by **orderD** ( $= n_{\max}$ ). At **orderD=0**, **Dpert=0**.

Solving (2.6) we get

$$\mathcal{D}(b, \mu) = \sum_{n=1}^{\infty} a_s^n \sum_{k=0}^n \mathbf{L}_{\mu}^k d^{(n,k)}, \quad (2.26)$$

where  $d^{(n,0)}$  are obtained by calculation. The coefficients  $d^{(n,k)}$  are

$$\begin{aligned} d^{(1,1)} &= \frac{\Gamma_0}{2}, \\ d^{(2,2)} &= \frac{\Gamma_0 \beta_0}{4}, \\ d^{(2,1)} &= \frac{\Gamma_1}{2}, \\ d^{(3,3)} &= \frac{\Gamma_0 \beta_0^2}{6}, \\ d^{(3,2)} &= \frac{\Gamma_0 \beta_1 + 2\Gamma_1 \beta_0}{4}, \\ d^{(3,1)} &= \frac{\Gamma_2 + 4\beta_0 d^{(2,0)}}{2}, \end{aligned}$$

where flavor on both sides is the same.

#### 2. Resummed RAD:

**Dresum(mu,bT,f):** returns  $\mathcal{D}_{\text{resum}}^f(b, \mu)$  [real(dp)]  
**mu, bT** [real(dp)] scales in GeV, **f**[integer] flavour.  
 Order is controlled by **orderDresum** ( $= n_{\max}$ ). At **orderDresum=0**, **Dresum**  $= -\frac{\Gamma_0}{2\beta_0} \ln(1 - X)$ .

One can resum the logarithms in this expression and write it as

$$\mathcal{D}_{\text{resum}} = -\frac{\Gamma_0}{2\beta_0} \left[ \ln(1 - X) + \sum_{n=1}^{\infty} \frac{a_s^n}{(1 - X)^n} \sum_{k,l=0}^n X^k \ln^l(1 - X) d_r^{(n,k,l)} \right], \quad X = \beta_0 a_s \mathbf{L}_{\mu}, \quad (2.27)$$

the coefficients  $d_r$  are

$$\begin{aligned} d_r^{(1,0,1)} &= B_1, & d_r^{(1,1,0)} &= B_1 - G_1, & d_r^{(1,0,0)} &= 0, \\ d_r^{(2,0,2)} &= -\frac{B_1^2}{2}, & d_r^{(2,0,1)} &= B_1 G_1, \\ d_r^{(2,2,0)} &= \frac{1}{2} (G_2 - B_1 G_1 - B_2 + B_1^2) & d_r^{(2,1,0)} &= B_1 G_1 - G_2, \\ d_r^{(2,0,0)} &= -2 \frac{d^{(2,0)} \beta_0}{\Gamma_0}, & d_r^{(2,2,2)} &= d_r^{(2,1,1)} = 0, \end{aligned}$$

$$\begin{aligned}
d_r^{(3,0,3)} &= \frac{B_1^3}{3}, & d_r^{(3,0,2)} &= -\frac{B_1^3}{2} - B_1^2 G_1, & d_r^{(3,0,1)} &= B_1 G_2 + 4 \frac{d^{(2,0)} \beta_1}{\Gamma_0}, \\
d_r^{(3,1,1)} &= B_1 B_2 - B_1^3, & \text{other } d_r^{(3,k>0,l>0)} &= 0, \\
d_r^{(3,3,0)} &= \frac{1}{3} (B_1^3 - 2B_1 B_2 + B_3 - B_1^2 G_1 + B_2 G_1 + B_1 G_2 - G_3), \\
d_r^{(3,2,0)} &= -\frac{1}{2} B_1^3 + B_1 B_2 - \frac{B_3}{2} + B_1^2 G_1 - B_2 G_1 - B_1 G_2 + G_3, \\
d_r^{(3,1,0)} &= B_1 G_2 - G_3, & d_r^{(3,0,0)} &= -2 \frac{d^{(3,0)} \beta_0}{\Gamma_0}
\end{aligned}$$

Here,

$$G_i = \frac{\Gamma_i}{\Gamma_0}, \quad B_i = \frac{\beta_i}{\beta_0}$$

### 3. Fixed order $\zeta$ -line:

`zetaMUpert(mu,bT,f)`: returns  $\zeta^f(b, \mu)$  [real(dp)]  
`mu, bT` [real(dp)] scales in GeV, `f`[integer] flavour.  
Order is controlled by `orderZETA` ( $= n_{\max}$ ). At `orderZETA`i0, `zetaMUpert`=1.

The solution of (2.12) in PT can be written as

$$\zeta^f(b, \mu) = \frac{C_0 \mu}{b} e^{-v}, \quad v = \sum_{n=0}^{\infty} a_s^n \sum_{k=0}^{n+1} \mathbf{L}_\mu v^{(n,k)}. \quad (2.28)$$

The *perturbative boundary condition* is that  $v$  is regular at  $\mathbf{L} \rightarrow 0$ . The first coefficients are

$$\begin{aligned}
v^{(0,0)} &= g_1, & v^{(0,1)} &= 0, \\
v^{(1,2)} &= \frac{\beta_0}{12}, & v^{(1,1)} &= 0, & v^{(1,0)} &= \mathbf{d}^{(2,0)} - g_1 G_1 + g_2, \\
v^{(2,3)} &= \frac{\beta_0^2}{24}, & v^{(2,2)} &= \frac{\beta_0}{12} (B_1 + G_1), & v^{(2,1)} &= \frac{\beta_0}{2} \left( \frac{8}{3} \mathbf{d}^{(2,0)} - g_1 G_1 + g_2 \right), \\
v^{(2,0)} &= \mathbf{d}^{(3,0)} - \mathbf{d}^{(2,0)} G_1 + g_1 G_1^2 - g_2 G_1 - g_1 G_2 + g_3,
\end{aligned}$$

here

$$\mathbf{d}^{(n,k)} = \frac{d^{(n,k)}}{\Gamma_0}, \quad g_i = \frac{\gamma_V^{(i)}}{\Gamma_0}, \quad G_i = \frac{\Gamma_i}{\Gamma_0}, \quad B_i = \frac{\beta_i}{\beta_0}.$$

### 4. Resummed $\zeta$ -line:

`zetaMUresum(mu,bT,f)`: returns  $\zeta_{\text{resum}}^f(b, \mu)$  [real(dp)]  
`mu, bT` [real(dp)] scales in GeV, `f`[integer] flavour.  
Order is controlled by `orderZETA`.  
**REMOVED in v.2.03. TO BE REINTRODUCED LATER.**

5. Exact  $\zeta$ -line:

`zetaSL(mu,bT,f)`: returns  $\zeta_{\text{exact}}^f(b, \mu)$  [real(dp)]  
`mu`, `bT` [real(dp)] scales in GeV, `f`[integer] flavour.  
Order is controlled by `orderZETA`. At `orderZETA=0`, `zetaMUpert=1`.  
**Currently at the order =3, the result is NNLO. Due to instability of N<sup>3</sup>LO at very large  $\mathcal{D}$ .**

The solution for  $\zeta$  can be written as

$$\zeta(b, \mu) = \mu^2 \exp(-g(a_s, \mathcal{D})). \quad (2.29)$$

The function  $g$  is convenient to present as

$$g(a_s, \mathcal{D}) = \frac{\Omega(a_s, p)}{a_s \beta_0}, \quad p = \frac{2\beta_0 \mathcal{D}}{\Gamma_0}, \quad (2.30)$$

where  $\Omega$  is convenient to present in the following form

$$\Omega(a_s, p) = \sum_{n=0}^{\infty} a_s^n \Omega_n(p), \quad (2.31)$$

with

$$\begin{aligned} \Omega_0 &= z_1, & \Omega_1 &= (B_1 - G_1)z_1 + g_1 - \frac{B_1}{2}p, \\ \Omega_2 &= \frac{B_2 - B_1 G_1 + G_1^2 - G_2}{2} z_1 + \frac{-B_2 + B_1 G_1 + G_1^2 - G_2 - 2G_1 g_1 + 2g_2}{2} z_{-1} + g_2 - g_1 G_1 \\ \Omega_3 &= \frac{2B_3 + B_1^2 G_1 - B_2 G_1 - G_1^3 + 3G_1 G_2 - B_1 B_2 - B_1 G_2 - 2G_3}{6} z_1 \\ &+ \frac{G_1 - B_1}{2} (-B_2 + B_1 G_1 + G_1^2 - G_2 - 2G_1 g_1 + 2g_2) z_{-1} \\ &+ \left[ \frac{-B_3 + B_1^2 G_1 + 2B_2 G_1 - 4G_1^3 + 6G_1 G_2 - B_1 B_2 + B_1 G_2 - 2G_3}{12} + (2G_1 - B_1) \frac{G_1 g_1 - g_2}{2} - \frac{G_2 g_1 - g_3}{2} \right] z_{-2} \\ &+ G_1^2 g_1 - G_2 g_1 - G_1 g_2 + g_3. \end{aligned}$$

Here

$$B_i = \frac{\beta_i}{\beta_0}, \quad G_i = \frac{\Gamma_i}{\Gamma_0}, \quad g_i = \gamma^{(i)} \frac{\beta_0}{\Gamma_0}, \quad z_n = \frac{e^{-np} - 1 + np}{p}. \quad (2.32)$$

Note that  $\lim_{p \rightarrow 0} z_n \sim n^2 p / 2$ .



### III. ATMDE\_SETUP MODULE

The module `aTMDe_setup` creates and modifies the `constants-file`. It is a stand-alone module, which does not require the rest modules (however, it is called by `aTMDe_control`).

List of commands **optional parameters are shown in blue**.

| Command  | Sec.   | Short description  |
|--|--------|--|
| <code>artemide_Setup_Default(order)</code>   | III A  | The main command which initializes variables by default values corresponded to a particular order.   |
| <code>artemide_Setup_fromFile(file, <b>prefix</b>, <b>order</b>)</code>  | III A  | The main command which initialize variables by pre-saved values in <b>file</b> . <b>prefix</b> is path to the file. <b>order</b> is the order of default version of parameters (used if versions of files are incompatible).   |
| <code>CreateConstantsFile(file, <b>prefix</b>)</code>  | -      | Write a new <code>constants-file</code> according to current modification.   |
| <code>CheckConstantsFile(file, <b>prefix</b>)</code>   | -      | Function (logical). Compare the version of <code>constants-file</code> . Returns <code>.true.</code> if version of file $\geq$ version of <code>artemide</code> , <code>.false.</code> otherwise.  |
| <code>artemide_include(arg1, arg2, ..., arg10)</code>  | -      | Include the modules into the initialization procedure. <b>arg</b> is (string) with the module name.  |
| <code>Set_outputLevel(level, <b>numMessages</b>)</code>  | -      | Set the level of <code>artemide</code> -messages to (int) <b>level</b> . 0= only critical, 1+=warnings, 2+=module evaluation information, 3+= details. Default=2. (integer) <b>numMessages</b> set number of non-critical Warnings of the same type to show (prevent spamming by same messages). |
| <code>Set_uPDF(hadron, setName, <b>replica</b>)</code>   | V A    | Assign the hadron for uPDF with number <b>hadron</b> (int) a PDF set <b>setName</b> (string) in LHAPDF library. It will be initialized in with replica <b>replica</b> (int)(default =0).   |
| <code>Set_uFF(hadron, setName, <b>replica</b>)</code>  | V A    | Assign the hadron for uFF with number <b>hadron</b> (int) a FF set <b>setName</b> (string) in LHAPDF library. It will be initialized in with replica <b>replica</b> (int)(default =0).   |
| <code>Set_lpPDF(hadron, setName, <b>replica</b>)</code>  | V A    | Assign the hadron for (unpolarized)PDF that is used by <code>lpTMDPDF</code> with number <b>hadron</b> (int) a PDF set <b>setName</b> (string) in LHAPDF library. It will be initialized in with replica <b>replica</b> (int)(default =0).   |
| <code>Set_quarkMasses(mC, mB, mT)</code>   | -      | Set values for pole quark masses, charm, bottom and top (real), which determine $N_f$ -thresholds. Default <b>mC</b> =1.4, <b>mB</b> =4.75, <b>mT</b> =173.  |
| <code>Set_EWparameters(alphaInv, massZ, massW, widthZ, widthW, massH, widthH, vevHIGGS, sin2ThetaW, UD, US, UB, CD, CS, CB)</code> | -      | Set parameters of electro-weak theory. <b>alphaInv</b> =inverse $\alpha_{QED}(M_Z)$ . UD, US, UB, CD, CS, CB are elements of CKM matrix. Masses and widths are in GeV.   |
| <code>Set_TMDR_order(order)</code>   | ??     | Set the perturbative order of anomalous dimensions used for evolution. <b>order</b> =string(8)   |
| <code>Set_TMDR_evolutionType(num)</code>   | ??     | Set the type of evolution solution used to (int) <b>num</b> .  |
| <code>Set_TMDR_lengthNArray(num)</code>  | ??     | Set the length of $\lambda_{NP}$ for $\mathcal{D}_{NP}$ to <b>num</b> (int).   |
| <code>Set_uTMDPDF(hadron, setName, <b>replica</b>)</code>  | V A ?? | Assign the hadron for uTMDPDF with number <b>hadron</b> (int) a PDF set <b>setName</b> (string) in LHAPDF library. It will be initialized in with replica <b>replica</b> (int)(default =0). Automatically calls for <code>Set_uPDF</code> .  |
| <code>Set_uTMDPDF_order(order)</code>  | ??     | Set the perturbative order of convolution. <b>order</b> =string(8)   |
| <code>Set_uTMDPDF_gridEvaluation(prepareGrid, <b>includeGluon</b>)</code>  | ??     | Set the trigger to prepare the grid, and to include the gluons in the grid (default=.false.). Both <b>logical</b> .  |
| <code>Set_uTMDPDF_lengthNArray(num)</code>   | ??     | Set the length of $\lambda_{NP}$ for uTMDPDF NP model to <b>num</b> (int).   |
| <code>Set_uTMDFF(hadron, setName, <b>replica</b>)</code>   | V A ?? | Assign the hadron for uTMDFF with number <b>hadron</b> (int) a FF set <b>setName</b> (string) in LHAPDF library. It will be initialized in with replica <b>replica</b> (int)(default =0). Automatically calls for <code>Set_uFF</code> .   |

|  |        |   |
|--|--------|---|
| <code>Set_uTMDFF_order(order)</code>   | ??     | Set the perturbative order of convolution. <code>order=string(8)</code>   |
| <code>Set_uTMDFF_gridEvaluation</code><br>( <code>prepareGrid</code> , <code>includeGluon</code> )   | ??     | Set the trigger to prepare the grid, and to include the gluons in the grid (default=.false.). Both <code>logical</code> .   |
| <code>Set_uTMDFF_lengthNPararray(num)</code>   | ??     | Set the length of $\lambda_{NP}$ for uTMDFF NP model to <code>num(int)</code> .   |
| <code>Set_lpTMDPDF(hadron, setName, replica)</code>  | V A ?? | Assign the hadron for lpTMDPDF with number <code>hadron(int)</code> a PDF set <code>setName(string)</code> in LHAPDF library. It will be initialized in with replica <code>replica(int)</code> (default =0). Automatically calls for <code>Set_lpPDF</code> . |
| <code>Set_lpTMDPDF_order(order)</code>   | ??     | Set the perturbative order of convolution. <code>order=string(8)</code>   |
| <code>Set_lpTMDPDF_gridEvaluation</code><br>( <code>prepareGrid</code> , <code>includeGluon</code> ) | ??     | Set the trigger to prepare the grid, and to include the gluons in the grid (default=.true.). Both <code>logical</code> .  |
| <code>Set_lpTMDPDF_lengthNPararray(num)</code>   | ??     | Set the length of $\lambda_{NP}$ for lpTMDPDF NP model to <code>num(int)</code> .   |

### A. aTMDe\_Setup

TO BE WRITTEN




FIG. 3: Scheme of creation and modification of `constants-file` by `aTMDe_setup`.

#### IV. ATMDE\_CONTROL MODULE

The module `atMDe_setup` is used to coordinate the operation of other modules. It does not bring any new features, just operates other modules in proper order. It is only for convenience.

List of commands **optional parameters are shown in blue**.

| Command  | Sec.  | Short description   |
|--|-------|---|
| <code>artemide_Initialize(file, <b>prefix, order</b>)</code> | III A | The command which initialize modules according to <code>constants-file</code> created by <code>artemide_Setup_fromFile(file, <b>prefix, order</b>)</code> .   |
| <code>artemide_ShowStatistics()</code>                       |       | Shows some information.   |
| <code>artemide_SetNPparameters(lambdaNP)</code>              | IV A  | Receive a (real*8)list of NP parameters, split it according to current setup and passes NP parameters to appropriate modules. Reset modules counters.   |
| <code>artemide_SetNPparameters_TMDR(lambdaNP)</code>         | IV A  | Reset NP parameters of TMDR-module by (real*8) list <code>lambdaNP</code> . Reset modules counters.   |
| <code>artemide_SetNPparameters_uTMDPDF(lambdaNP)</code>      | IV A  | Reset NP parameters of uTMDPDF-module by (real*8) list <code>lambdaNP</code> . Reset modules counters.  |
| <code>artemide_SetNPparameters_uTMDFF(lambdaNP)</code>       | IV A  | Reset NP parameters of uTMDFF-module by (real*8) list <code>lambdaNP</code> . Reset modules counters.   |
| <code>artemide_SetNPparameters_lpTMDFF(lambdaNP)</code>      | IV A  | Reset NP parameters of lpTMDFF-module by (real*8) list <code>lambdaNP</code> . Reset modules counters.  |
| <code>artemide_SetReplica_TMDR(num)</code>                   | IV A  | Reset NP parameters of TMDR-module by values corresponding to replica (int) <code>num</code> . Reset modules counters.  |
| <code>artemide_SetReplica_uTMDPDF(num)</code>                | IV A  | Reset NP parameters of uTMDPDF-module by values corresponding to replica (int) <code>num</code> . Reset modules counters.   |
| <code>artemide_SetReplica_uTMDFF(num)</code>                 | IV A  | Reset NP parameters of uTMDFF-module by values corresponding to replica (int) <code>num</code> . Reset modules counters.  |
| <code>artemide_SetReplica_lpTMDPDF(num)</code>               | IV A  | Reset NP parameters of lpTMDPDF-module by values corresponding to replica (int) <code>num</code> . Reset modules counters.  |
| <code>artemide_SetScaleVariations(c1, c2, c3, c4)</code>     | -     | Change the value of scale-variation constants $c_1 - c_4$ .   |
| <code>artemide_GetReplicaFromFile(file, rep, array)</code>   | -     | Read the .rep file (path is <code>file</code> ), and search for the replica (int) <code>rep</code> . Return the (real*8, allocatable) <code>array</code> of NP parameters. <b>This command can work without initialization of artemide-control</b> , in this case, no check of consistency is performed and warning raised. |
| <code>artemide_NumOfReplicasInFile(file)</code>              | -     | Read the .rep file (path is <code>file</code> ), and return the number of replicas saved in it.   |

##### A. Passing non-perturbative parameters

The important part of the initialization is the number of NP parameters for each TMD distributions under consideration. Each TMD-evaluating module (say, uTMDPDF, uTMDFF, etc.) requires  $n_i$  number of parameters. The numbers are specified in `constants-file`. The number must be greater then zero  $n_i > 0$  for any used module, i.e.  $f_{NP}$  is at least 1-parametric (if it is not so, just do not use the parameter in the definition of  $f_{NP}$ , but keep  $n_i > 0$ ). These numbers are read during the initialization procedure, and allocate the memory.

The set of particular values of these parameters can be done by several ways.

**Option I:** Set all values in a single call  
call `artemide_SetNPParameters(lambda)`  
where

$\{\lambda_i\}$  real\*8(1: $\sum_i n_i$ ) The set of parameters which define the non-perturbative functions  $f_{NP}$  within modules. It is split into parts and send to corresponding modules. I.e.  $\text{lambda}(1:n_0) \rightarrow \text{uTMDR}$ ,  $\text{lambda}(n_0 + 1:n_0 + n_1) \rightarrow \text{uTMDPDF}$ ,  $\text{lambda}(n_0 + n_1 + 1:n_0 + n_1 + n_2) \rightarrow \text{uTMDFF}$  (fixed order).

**Option II:** Set value for particular function. For it call `artemide_SetNPparameters_???(lambdaNP)`, where ??? is module name, e.g.

`artemide_SetNPparameters_uTMDPDF(lambdaNP)`

will set parameters for unpolarized TMDPDF.

**Option III:** You can use presaved values of  $\lambda_{NP}$  for a given function. They are provided by model file (if provided). To set NP-input given by integer `num`, call `artemide_SetReplica_???(num)`, where ??? is module name, e.g.

`artemide_SetReplica_uTMDPDF(num)`

will set parameters for unpolarized TMDPDF.

## V. QCDINPUT MODULE

The module `QCDinput` gives an interface to external function provided by the user, such as PDF, FF, values of alpha-strong. By default the input for these functions is done via the LHAPDF tables. Originally, `artemide` used the LHAPDF library [? ], which however caused a number of inconveniences, such as difficulty in the installation, problems with simultaneous usage of many tables, and parallel access to the tables. Starting from the version 3.01 `artemide` is detached from the LHAPDF library, but continue to relay on LHAPDF-tables as input for the QCD parameters. The tables are read and interpolated by the original `artemide` sub-modules (`LHA_alpha` and `LHA_PDF`). The details on these modules are presented below.

For the technical reasons there is strict limit on the maximum number of hadrons. This limit can be changed but it requires modification of the code. The default limitation is the following: `uPDF` 3 hadrons, `uFF` 3 hadrons, `lpPDF` 1 hadron, `hPDF` 2 hadrons.

List of available commands [optional parameters are shown in blue](#).

| Command  | Description   |
|--|---|
| <code>QCDinput.Initialize(file, <a href="#">prefix</a>)</code> | Subroutine to initialize anything what is needed. (string) <code>file</code> is the name of <code>constants-file</code> , which contains initialization information. (string) <code>prefix</code> is appended to <code>file</code> if provided. |
| <code>As(Q)</code>   | Returns the (real*8) value of $\alpha_s(Q)/(4\pi)$ . $Q$ is (real*8).   |
| <code>xPDF(x, Q, hadron)</code>                                | Returns the (real*8(-5:5)) value of $xf(x, Q)$ for given hadron. $x$ , $Q$ are (real*8), <code>hadron</code> is (integer).  |
| <code>xFF(x, Q, hadron)</code>                                 | Returns the (real*8(-5:5)) value of $xd(x, Q)$ for given hadron. $x$ , $Q$ are (real*8), <code>hadron</code> is (integer).  |
| <code>x_hPDF(x, Q, hadron)</code>                              | Returns the (real*8(-5:5)) value of $xg_1(x, Q)$ for given hadron ( $g_1$ =helicity distribution). $x$ , $Q$ are (real*8), <code>hadron</code> is (integer).  |
| <code>mCHARM</code>  | (real*8)Constant for mass of charm quark.   |
| <code>mBOTTOM</code>   | (real*8)Constant for mass of bottom quark.  |
| <code>QCDinput.SetPDFreplica(num, h)</code>                    | Changes the unpolarized PDF replica number for hadron <code>h</code> to <code>rep</code> .  |
| <code>QCDinput.SetFFreplica(num, h)</code>                     | Changes the unpolarized FF replica number for hadron <code>h</code> to <code>rep</code> .   |
| <code>QCDinput.SetlpPDFreplica(num, h)</code>                  | Changes the replica of PDF associated with linearly polarized gluon PDF number for hadron <code>h</code> to <code>rep</code> .  |
| <code>QCDinput.SethPDFreplica(num, h)</code>                   | Changes the replica of helicity PDF number for hadron <code>h</code> to <code>rep</code> .  |

### A. Initialization

TOBE WRITTEN

### B. LHA\_alpha module

The module reads the info-file of the LHAPDF set and extracts the information about  $\alpha_s$ . For the moment the only possible input is with "`AlphaS.Type=ipol`". The interpolation is done by cubic-interpolation in logarithm scale, for each mass-threshold subgrid (in accordance of the description of LHAPDF library). Compared that the deviation from the original LHA library and `artemide` is better than  $10^{-6}$  (i.e. at the level of float-definition fluctuations).

Below the  $Q_{\min}$  I perform the extrapolation by the following formula

$$\alpha_s(Q) = \frac{\alpha_0 \alpha_1 \ln(Q_1/Q_0)}{(\alpha_0 - \alpha_1) \ln(Q/Q_0) + \alpha_1 \ln(Q_1/Q_0)}, \quad (5.1)$$

where  $Q_0 = Q_{\min}$  and  $Q_1$  is the next-to- $Q_{\min}$  node, and  $\alpha_{0,1} = \alpha_s(Q_{0,1})$ . This formula extrapolate  $1/\alpha$  logarithmically in the logarithm scale. It gives the values of  $\Lambda_{\text{eff}}$

$$\Lambda_{\text{eff}} = Q_0 \exp \left( -\frac{\alpha_1 \ln(Q_1/Q_0)}{\alpha_0 - \alpha_1} \right), \quad (5.2)$$

where the coupling constant diverges. The minimal possible  $Q$  is truncated to  $\min(0.4\text{GeV}, \Lambda_{\text{eff}} + 0.1\text{GeV})$  to avoid problems with low energy.

### C. LHA\_PDF module

The module reads info-file and the PDF-table and interpolates using the log-qubic interpolation (instead of cubic-Hermite interpolator from LHAPDF-library). There is a small-difference between interpolating results but they are smaller than  $10^{-4}$  (relative). I.e. far inside of the pdf uncertainties.

Below the  $Q_{\min} = Q_0$  the extrapolation is done by the formula

$$F(x, Q) = F(x, Q_0)(Q/Q_0)^{\Gamma(Q)}, \quad \Gamma(Q) = 2 + \sqrt{\frac{Q}{Q_0}}(\gamma(x) - 2), \quad (5.3)$$

where  $F(x, Q) = xf(x, Q)$ , and  $\gamma(x)$  is

$$\gamma(x) = \frac{\ln F(Q_1) - \ln F(Q_0)}{\ln(Q_1/Q_0)} \approx \frac{d \ln F(Q)}{d \ln Q}. \quad (5.4)$$

This is in the agreement of formulas (4,5) of ref.[? ]. Below 0.4GeV the extrapolation terminated and error is called.

## VI. EWINPUT MODULE

`EWinput` module contains definitions of various physical parameters of elector-weak interactions, such as masses of Z and W bosons, CKM matrix, code for evolution of  $\alpha_{QED}$  etc.

**TOBE WRITTEN**

### A. $\alpha_{QED}$

**This implementation has been added in ver.2.03. In earlier versions the code for  $\alpha_{QED}$  had a bug that generated slightly lower values (independently on the values of  $\mu$ ).**

The QED coupling constant is set by the one-loop QED evolution with LO matching at thresholds. In order to tune it finer, the value of  $\beta_0$  is modified by  $\beta_0 \rightarrow \beta_0 + \delta\beta$ , where  $\delta\beta$  is found by exact implementation of two boundary values,  $\alpha_{QED}(m_\tau)$  and  $\alpha_{QED}(m_Z)$ .

The LO evolution is

$$\alpha(\mu) = \frac{1}{\alpha_0^{-1} + 2\beta_0 \ln(\mu/\mu_0)}, \quad (6.1)$$

where  $\beta_0 = -(3\pi)^{-1}N_{eff}$  with

$$N_{eff} = \sum_{\text{active leptons}} 1 + N_c \sum_{\text{active quarks}} e_q^2. \quad (6.2)$$

In artemide I account the following thresholds

| $\mu$     | $\mu < m_e (\sim m_u, m_d)$ | $\mu < m_\mu (\sim m_s)$ | $\mu < m_c$ | $\mu < m_\tau$ | $\mu < m_b$    | $\mu < m_t$    | $\mu > m_t$ |
|-----------|-----------------------------|--------------------------|-------------|----------------|----------------|----------------|-------------|
| $N_{eff}$ | 0                           | $\frac{8}{3}$            | 4           | $\frac{16}{3}$ | $\frac{19}{3}$ | $\frac{20}{3}$ | 8           |

So the actual formula for  $\alpha_{QED}$  is

$$\alpha_{QED}(\mu) = \frac{1}{\alpha_Z^{-1} + 2\bar{\beta} \left[ N_{eff}(\mu) \ln \left( \frac{\mu}{m_{th}} \right) + \sum_{m_i \in m_{th}} N_{eff} \ln \left( \frac{m_i}{m_{i+1}} \right) \right]}, \quad (6.3)$$

where  $m_{th}$  are all thresholds from  $\mu$  till  $M_Z$ . The constant  $\bar{\beta}$  is defined by exact matching at  $M_Z$  and  $m_\tau$ . For default values  $\bar{\beta}/\beta_0 \sim 1.002\%$ , so it accounts for really tiny effects.

## VII. TMD EVOLUTION MODULES

The TMD evolution has several implementation. The artemide v3 (and higher) is hard-coded to use the fixed- $\mu$  evolution (see ref.[? ]). This choice is made after many test performed in earlier versions or artemide (where the type of evolution was selected by user), and the demonstration that they are equivalent within perturbative uncertainty.

The fixed- $\mu$  scale corresponds to evolution along  $\zeta$  is with fixed- $\mu$  solution to the special equi-potential line, and evolution along  $\mu$  along equi-potential line (this part is unity by definition). The evolution factor reads

$$R[b, (\mu_1, \zeta_1) \rightarrow (\mu_2, \zeta_2)] = \frac{R[b, (\mu_1, \zeta_1) \rightarrow (\mu_1, \zeta_{\mu_1})]}{R[b, (\mu_2, \zeta_2) \rightarrow (\mu_2, \zeta_{\mu_2})]}, \quad (7.1)$$

$$R[b, (\mu, \zeta) \rightarrow (\mu, \zeta_\mu)] = \exp \left( -\mathcal{D}(\mu, b) \ln \left( \frac{\zeta}{\zeta_\mu(b)} \right) \right), \quad (7.2)$$

where  $\mathcal{D}$  is the Collins-Soper kernel, and  $\zeta_\mu(b)$  is the special  $\zeta$ -line.

The Collins-Soper kernel is defined as

$$\mathcal{D}^f(\mu, b) = \mathcal{D}_{\text{pert}}^f(\mu_{\text{OPE}}, b) + \int_{\mu_{\text{OPE}}}^{\mu} \frac{d\mu'}{\mu'} \Gamma^f(\mu') + \mathcal{D}_{\text{NP}}^f(b), \quad (7.3)$$

where  $\mathcal{D}_{\text{pert}}^f$  is the high-scale perturbative part, and  $\mathcal{D}_{\text{NP}}$  is the non-perturbative model (e.g.  $g_K b^2$ )

The module for evolution includes three modules

- **TMD\_AD** this module contains collection of all anomalous dimensions and related quantities. Logically it is equivalent to `_OPE` modules.
- **TMDR\_model** this module contains the user-defined non-perturbative input.
- **TMDR** this module interface **TMD\_AD** and **TMDR\_model** into the evolution factor, and perform the general control.

The initialization is happened thorough the **TMDR**-module.

At small values of parameter  $Q = Q_0$  the point  $(Q, Q^2)$  crosses the  $\zeta$ -lines. The value of  $Q_0$  depends on  $b$ . The dangerous situation is then hard scale of the process  $Q$  is smaller then  $Q_0$  at large  $b = b_\infty$ . In this case the evolution kernel  $R[b_\infty, (Q, Q^2) \rightarrow \zeta_\mu] > 1$ , which is generally implies that it grows to infinity. However, it happens only at small values of  $Q$ . E.g. at NNLO the typical value of  $Q_0$  is  $\sim 1.5\text{GeV}$ . That should be taken into account during consideration of low-energy experiment and especially their error-band, since the point  $(c_2 Q, Q^2)$  could cross the point at larger values of  $Q$ .

The function **LowestQ()** returns the values (real\*8(1:3))  $\{Q_{-1}, Q_0, Q_{+1}\}$ , which are solution of equation  $Q^2 = \zeta_{cQ}(b)$ , for (fixed bu) large values of  $b$ .  $Q_{-1}$  corresponds to  $c = 0.5$ ,  $Q_0$  corresponds to  $c = 1$  and  $Q_{+1}$  corresponds to  $c = 2$ .

### 1. INI-parameters

**Parameters for TMD-evolution modules (section \*3).**



| Entry | Type | Description   |
|-------|------|---|
| A.p1  | str  | Order of the evolution <i>standardized</i> . Could be LO, NLO,.. (see table below for details)  |
| A.p2  | T/F  | The flag to override the <i>standard</i> evolution definition. If T, than instead of A.p1 the orders defined in A.p3 – A.p6 are used. |
| A.p3  | str  | Order of $\Gamma_{\text{cusp}}$ . LO= $a_s\Gamma_0$ , NLO= $a_s\Gamma_0 + a_s^2\Gamma_1$ , etc.                                       |
| A.p4  | str  | Order of $\gamma_V$ . LO=0, NLO= $a_s\gamma_1$ , etc.   |
| A.p5  | str  | Order of $\mathcal{D}_{\text{pert}}$ . LO=0, NLO= $a_s\mathbf{L} + d^{(1,0)}$ , etc.  |
| A.p6  | str  | Order of $\zeta_\mu$ (exact solution). LO= $a_s^{0*}(\text{exact solution})$ , etc.   |
| B.p1  | int  | Number of NP parameters.  |
| C.p1  | real | Tolerance general (used for check of variable modifications, etc. Does not affect integration precision). (default = $10^{-6}$ )      |
| C.p2  | real | Small-b value at which the evolution is frozen (default = $10^{-6}$ )   |
| C.p3  | real | Parameter for smoothing the small-b value of exact $\zeta$ -line (see below) (default =0.01).   |

The table of correspondence between **standard** definition and particular definitions

| order      | $\Gamma_{\text{cusp}}$ | $\gamma_V$ | $\mathcal{D}^*$ | $\mathcal{D}_{\text{resum}}$ | $\zeta_\mu^{**}$ |
|------------|------------------------|------------|-----------------|------------------------------|------------------|
| LO         | $a_s^1$                | $a_s^0$    | $a_s^0$         | $a_s^0$                      | $a_s^0$          |
| NLO        | $a_s^2$                | $a_s^1$    | $a_s^1$         | $a_s^1$                      | $a_s^2$          |
| NNLO=N2LO  | $a_s^3$                | $a_s^2$    | $a_s^2$         | $a_s^2$                      | $a_s^3$          |
| NNNLO=N3LO | $a_s^4$                | $a_s^3$    | $a_s^3$         | $a_s^2$                      | $a_s^4$          |
| N4LO       | $a_s^5$                | $a_s^4$    | $a_s^4$         | $a_s^3$                      | $a_s^{5***}$     |

\*  $\mathcal{D}_{\text{resum}}$  starts from  $a_s^0$ , it already contains  $\Gamma_0$ .

\*\* Definition of  $\zeta_\mu$  is correct only in the natural ordering, i.e. LO,NLO,NNLO. Proper definition in + orders would make the function too heavy. The resummed version has the same counting.

\*\*\* The finite part =0, since  $d^{(5,0)}$  is unknown.

#### Notes:

- For very small-values of  $b$  the value of  $b$  is fixed.
- For very large positive values of  $\mathcal{D}$  (i.e. for  $b > 25\text{GeV}^{-1}$  or so), the exact solution for  $\zeta$ -line diverges exponentially at 3- and 4- loops. It is because the coefficient in front of rising terms is negative. To prevent the divergence, the corresponding terms are expanded in  $a_s$  up to needed power ( $a_s^6$ ).

## 2. Functions

XX=name of module (e.g. uTMDPDF, uTMDFF, etc.)

| Entry                           | Type       | Description  |
|---------------------------------|------------|--|
| TMDR_Initialize(file,prefix)    | subroutine | Initialize the module from INI-file.   |
| TMDR_SetScaleVariation(c1)      | subroutine | Sets globally a new value for constant $c_1$ .   |
| TMDR_SetNPparameters(lambda)    | subroutine | Set a new vector of NP parameters $\lambda_{\text{NP}}$ . The new vector must be same size as declared at the setup. |
| TMDR_CurrentNPparameters()      | real(:)    | Returns the current vector for $\lambda_{\text{NP}}$ .   |
| CS_kernel(mu,b,f)               | real       | Returns the value Collins-Soper kernel at scale $\mu$ , point $b$ for flavor $f$ .                                   |
| zetaNP(mu,b,f)                  | real       | Returns the value $\zeta_\mu$ for give CS kernel at scale $\mu$ , point $b$ for flavor $f$ .                         |
| TMDR_Rzeta(b,muf,zetaf,f)       | real       | Return the factor $R$ for evolution from the point $(\mu, \zeta)$ down to the special evolution line.                |
| TMDR_R(b,muf,zetaf,mui,zetai,f) | real       | Return the factor $R$ for evolution from the point $(\mu_f, \zeta_f)$ to $(\mu_i, \zeta_i)$ .                        |
| LowestQ()                       | real       | Attempts to compute the lowest available scale $Q$ , for which the evolution converges.                              |

#### A. TMD\_AD module

The module TMD\_AD give access to various perturbative anomalous dimensions needed for TMD evolution. This module is essential part of TMDR module, and thus initialized by it.

List of available commands

| Command                            | Sec. | Short description  |
|------------------------------------|------|--|
| TMD_AD_Initialize(oC,oV,oD,oDr,oZ) | ??   | Initialization of module. The (integer) parameters (oC,oV,oD,oDr,oZ) are orders of definition for anomalous dimensions ( $\Gamma_{\text{cusp}}$ , $\gamma_V$ , $\mathcal{D}_{\text{pert}}$ , $\mathcal{D}_{\text{resum}}$ , $\zeta_{\text{pert}}$ ). |
| GammaCusp(mu,f)                    |      | $= \Gamma_{\text{cusp}}^f(\mu)$  |
| gammaV(mu,f)                       |      | $= \gamma_V^f(\mu)$  |
| Dpert(mu,bT,f)                     |      | $= \mathcal{D}_{\text{pert}}^f(b, \mu)$  |
| Dresum(mu,bT,f)                    |      | $= \mathcal{D}_{\text{resum}}^f(b, \mu)$   |
| zetaMUpert(mu,bT,f)                |      | $= \zeta_{\text{pert}}^f(b, \mu)$  |
| zetaSL(mu,rad,f)                   |      | $= \zeta_{\text{exact}}^f(\mathcal{D}, \mu)$ (Note, the argument.)   |
| RADEvolution(mu0,mu1,f)            |      | $= \int_{\mu_0}^{\mu_1} \frac{d\mu^2}{\mu^2} \frac{\Gamma(\mu)}{2}$ The evolution constant for $\mathcal{D}$   |

#### Notes:

- The integral over  $\mu$  is done using exact analytical formula by decomposition of  $\beta$ -function over roots.
- The exact expression for  $\zeta$ -line is numerically unstable at very small- $b$ . It is due to its double-exponential nature which requires exponentiation and cancellation of large-numbers. However, it could be computed analytically in this regime as expansion. Thus, to stabilize the computation of expression the following formula is used

$$\zeta_\mu = e^{-b^2/S} \zeta_{\text{pert}} + (1 - e^{-b^2/S}) \zeta_{\text{exact}}. \quad (7.4)$$

Here,  $S$  is the smoothing parameter. It should be small enough to guaranty non-impact of approximation (approx.  $S < 0.05$ ).

## B. TMDR\_model

This module gives the expression for  $\mathcal{D}$ . It could be modified by user, and one of modules that form a TMD model (together with models for TMD-distributions).

**Making this module follow the interface**

```
public:: ModelInitialization(NPlength)
public:: ModelUpdate(NParray)
real(dp),public:: DNP(b,f)
real(dp),public:: bSTAR(b)
real(dp),public:: muOPE(b,c1)
```

**ModelInitialization(NPlength):** this subroutine is called during the initialization of TMDR. The `integer,intent(in)::NParray(:)` is the length of array of NP-parameters. Use this subroutine to make preparation of the model.

**ModelUpdate(NParray):** this subroutine is called during the `TMDR_setNPparameters(NParray)`. The `real(dp),intent(in)::NParray(:)` is the array of NP-parameters passed to `TMDR_setNPparameters`. Use this subroutine to recompute model parameters (if needed), save NP-parameters (if needed), etc.

**DNP(b,f):** the function returns the value,  $\mathcal{D}_{\text{NP}}^f(b)$ . It contains only NP part.

**bSTAR(b):** the function returns the value,  $b^*(b)$  used instead of  $b$  within  $\mathcal{D}_{\text{pert}}$

**muOPE(b,c1):** the function returns the value,  $\mu_{\text{OPE}}$  used as reference scale of  $\mathcal{D}_{\text{pert}}$ . The parameter  $c_1$  is the prescription used during the scale variation.

**Important:** the module `TMDR_model` is compiled before TMDR, and thus can use only lower-level modules: `TMD_AD`, `QCDinput`, etc.

**Important:** There is an option to switch to the CSS-like computation (formula (2.14) in 1803.11089).

### VIII. MODULES COMPUTING TMD DISTRIBUTIONS

There are many TMD distributions each computed in a separate module. All TMD distributions are split in the following classes:

- **Pure Twist-2 OPE** These are `uTMDPDF`, `uTMDFF`, `lpTMDPDF`. These TMD distributions are defined by twist-2 collinear distribution at small- $b$
- **Pure Twist-3 OPE** These are `SiversTMDPDF`, `BoerMuldersTMDPDF`. These TMD distributions are defined by twist-3 collinear distribution at small- $b$
- **Twist-2 & Twist-3 OPE** These are `wgtTMDPDF`. These TMD distributions are defined by twist-2 (WW-term) and twist-3 collinear distribution at small- $b$

These lists are to be extended in the future.

Each module consists of several sub-modules, which have names as `TMDname_X`. The `X` corresponds to

- **OPE** this module computes the convolution of the twist-2 (or twist-3) coefficient function with corresponding PDF.
- **model** this module contains all information about NP-part and other free-parameters (such as  $\mu_{\text{OPE}}$  of the given TMD distribution. **These modules could be edited by user, to implement/update/fit TMDs.**

The most part of modules are identical. There are a few differences in the implementation. So, first the general structure is presented, and then particularities.

Different OPE's are known up to different orders. Consequently, the modules implement them at different orders. The orders are denoted as

$$\begin{aligned}
 \text{NA} & \rightarrow \text{no OPE} \\
 \text{LO} & \rightarrow C = \mathcal{O}(a_s^0) \\
 \text{NLO} & \rightarrow C = \mathcal{O}(a_s^1) \\
 \text{NNLO} = \text{N2LO} & \rightarrow C = \mathcal{O}(a_s^2) \\
 \text{NNNLO} = \text{N3LO} & \rightarrow C = \mathcal{O}(a_s^3).
 \end{aligned}$$

- The NA option is used to remove any relation to collinear distributions. In this case  $TMD(x, b) = f_{\text{NP}}(x, b)$
- If the coefficient function at LO/NLO/.. is zero, the convolution is also zero. E.g. linearly polarized gluons start at  $\sim a_s$ , the convolution is zero at LO.

List of available orders for different distributions

$$\begin{aligned}
 \text{uTMDPDF} & : \text{NA, LO, NLO, N2LO, N3LO} \\
 \text{uTMDFF} & : \text{NA, LO, NLO, N2LO, N3LO} \\
 \text{lpTMDPDF} & : \text{NA, LO, NLO, N2LO} \\
 \text{wgtTMDPDF}[tw2] & : \text{NA, LO, NLO} \\
 \text{wgtTMDPDF}[tw3] & : \text{NA} \\
 \text{SiversTMDPDF}[tw3] & : \text{NA} \\
 \text{BoerMuldersTMDPDF}[tw3] & : \text{NA}
 \end{aligned}$$

The functions are computed by the following rule

$$\text{uTMDPDF} = [C \times f]_{\text{tw2}}(x, b) F_{\text{NP}}(x, b), \quad (8.1)$$

$$\text{uTMDFF} = [C \times d]_{\text{tw2}}(x, b) F_{\text{NP}}(x, b), \quad (8.2)$$

$$\text{lpTMDPDF} = [C \times f]_{\text{tw2}}(x, b) F_{\text{NP}}(x, b), \quad (8.3)$$

$$\text{wgtTMDPDF} = [C \times h]_{\text{tw2}}(x, b) F_{\text{NP}}(x, b) + [C \times T]_{\text{tw3}}(x, b) F_{\text{NP}}^{\text{tw3}}(x, b), \quad (8.4)$$

$$\text{SiversTMDPDF} = [C \times T]_{\text{tw3}}(x, b) F_{\text{NP}}^{\text{tw3}}(x, b), \quad (8.5)$$

$$\text{BoerMuldersTMDPDF} = [C \times E]_{\text{tw3}}(x, b) F_{\text{NP}}^{\text{tw3}}(x, b), \quad (8.6)$$

where  $f, d, h$  are collinear distributions of twist-2, and  $[C \times h]_{\text{tw}2}$  is corresponding convolution;  $T$  and  $E$  are collinear distributions of twist-3, and  $[C \times T]_{\text{tw}3}, [C \times E]_{\text{tw}3}$  are corresponding convolution;  $F_{\text{NP}}(x, b)$  and  $F_{\text{NP}}^{\text{tw}3}(x, b)$  are non-perturbative functions to be fit. NP functions are defined in the `XX_model` modules, and share the same  $\lambda_{\text{NP}}$  (array of parameters). This structure correctly reproduces the small- $b$  properties of TMD-distributions. The convolutions are computed in separate modules see sec.VIII B, VIII C.

The modules computes the optimal TMDs, and also evolved TMDs. Generally a TMD distribution is given by the expression

$$F_f(x, b; \mu, \zeta) = R_f[b, (\mu, \zeta) \rightarrow (\mu, \zeta_\mu)] \tilde{F}_f(x, b), \quad (8.7)$$

where  $R$  is the TMD evolution kernel,  $\tilde{F}$  is an optimal TMD distribution.

#### 1. *INI-parameters*

**Parameters universal for all such modules which include only tw2 or tw3 OPE (mixed case see below).**

| Entry | Type    | Description   |
|-------|---------|---|
|       |         |   |
| A.p1  | T/F     | Include gluon into the computation. Usually gluon is computed by a separate rule, and switching off this option leads to $\sim 2$ increase in the computation velocity.         |
| A.p2  | in      | Number of hadrons. Specify maximum required number of hadrons.  |
| B.p1  | str     | Order of the coefficient function. Could be N/A, LO, NLO,.. (see table below for details)   |
| B.p2  | T/F     | Trigger to prepare the grid (T=prepare). Preparation is done on the initialization stage.   |
| B.p3  | T/F     | Trigger the test of the grid (T=run the test). Take the same time as computation of the grid (by a single processor). Requires A.p2=T. Example Result is shown in sec.VIII B 4. |
| C.p1  | int     | Number of NP parameters.  |
| C.p2  | real    | $B_{\text{MAX-ABS}}$ , a large number such that $b > B_{\text{MAX-ABS}}$ the TMD is returned 0. (default 100).  |
| D.p1  | real    | Tolerance for integration (default $=10^{-6}$ )   |
| D.p2  | real    | Tolerance general (used for check of variable modifications, etc. Does not affect integration precision). (default $=10^{-6}$ )   |
| D.p3  | int     | Maximum number of iteration for adaptive integration. (default $=10^4$ )  |
| E.p1  | real    | $x_{\min}$ for the grid (see (??))  |
| E.p2  | real    | $p_x$ for the grid (see (??))   |
| E.p3  | real    | $b_{\min}$ for the grid (see (??)). [It is not related to parameter $b_{\min}$ of $b^*$ -prescription!]   |
| E.p4  | real    | $b_{\max}$ for the grid (see (??)) [It is not related to parameter $b_{\max}$ of $b^*$ -prescription!]  |
| E.p5  | integer | Number of nodes in the x-grid   |
| E.p6  | integer | Number of nodes in the b-grid   |
| F.p1  | real    | Tolerance use in the Fourier integral to the $k_T$ -space (default $=10^4$ )  |
| F.p2  | real    | Step for the Ogata quadrature (default $=10^{-3}$ )   |
| F.p3  | real    | Minimal $k_T$ to compute. Below this number distribution is frozen. (default $=10^{-4}$ )   |
| G.p1  | real    | Tolerance use in the Fourier integral for TMM (default $=10^4$ )  |
| G.p2  | real    | Step for the Ogata quadrature for TMM computation (default $=10^{-3}$ )   |
| G.p3  | real    | Minimal $\mu$ to compute. Below this number the program terminates. (default $=0.8\text{GeV}$ )   |

- **wgtTMDPDF** has extra section F and G, which are analogous to sections B and C, but responsible for twist-3 convolution. The section F,G,I is correspondingly sections H,I.

## 2. Functions

XX=name of module (e.g. **uTMDPDF**, **uTMDFF**, etc.)

| Entry                      | Type       | Description  |
|----------------------------|------------|--|
| XX_Initialize(file,prefix) | subroutine | Initialize the module from INI-file.   |
| XX_SetPDFreplica(rep,h)    | subroutine | Passes to XX_OPE.SetPDFreplica   |
| XX_SetScaleVariation(c4)   | subroutine | Passes to XX_OPE.SetScaleVariation   |
| XX_SetLambdaNP(lambda)     | subroutine | Set a new vector of NP parameters $\lambda_{NP}$ . The new vector must be same size as declared at the setup.  |
| XX_CurrentLambdaNP()       | real(:)    | Returns the current vector for $\lambda_{NP}$ .  |
| XX_lowScale5(x,bT,h)       | real(-5:5) | <b>TO BE REMOVED</b> Returns the value of TMD distributions for given (x,b) (reals) and hadron h. The value is the vector over flavors. For $h < 0$ the quark-distributions are replaced by anti-quark distributions and vice versa. I.e. $F(-h) = F(h)(5 : -5 : -1)$  |
| XX_inB(x,bT,h)             | real(-5:5) | Returns the value of <b>optimal</b> TMD distributions for given (x,b) (reals) and hadron h. The value is the vector over flavors. For $h < 0$ the quark-distributions are replaced by anti-quark distributions and vice versa. I.e. $F(-h) = F(h)(5 : -5 : -1)$  |
| XX_inB(x,bT,mu,zeta,h)     | real(-5:5) | Returns the value of <b>evolved</b> TMD distributions (see (8.7)) for given (x,b, $\mu,\zeta$ ) (reals) and hadron h. The value is the vector over flavors. For $h < 0$ the quark-distributions are replaced by anti-quark distributions and vice versa. I.e. $F(-h) = F(h)(5 : -5 : -1)$ . $\mu$ is in GeV. $\zeta$ in $\text{GeV}^2$ .   |
| XX_inKT(x,kT,h)            | real(-5:5) | Returns the value of <b>optimal</b> TMD distributions <b>in momentum space</b> (see sec.VIII D) for given (x, $k_T$ ) (reals) and hadron h. The value is the vector over flavors. For $h < 0$ the quark-distributions are replaced by anti-quark distributions and vice versa. I.e. $F(-h) = F(h)(5 : -5 : -1)$  |
| XX_inKT(x,bT,mu,zeta,h)    | real(-5:5) | Returns the value of <b>evolved</b> TMD distributions (see (8.7)) <b>in momentum space</b> (see sec.VIII D) for given (x, $k_T,\mu,\zeta$ ) (reals) and hadron h. The value is the vector over flavors. For $h < 0$ the quark-distributions are replaced by anti-quark distributions and vice versa. I.e. $F(-h) = F(h)(5 : -5 : -1)$ . $\mu$ is in GeV. $\zeta$ in $\text{GeV}^2$ . |
| XX_TMM_G(x,mu,h)           | real(-5:5) | Returns the value of TMM $\mathcal{G}_{n,n}$ (see sec.VIII E) for given (x, $\mu$ ) (reals) and hadron h. The value is the vector over flavors. For $h < 0$ the quark-distributions are replaced by anti-quark distributions and vice versa.   |
| XX_TMM_X(x,mu,h)           | real(-5:5) | Returns the value of TMM $\mathcal{G}_{n+1,n}$ (see sec.VIII E) for given (x, $\mu$ ) (reals) and hadron h. The value is the vector over flavors. For $h < 0$ the quark-distributions are replaced by anti-quark distributions and vice versa.   |

### Functions unique for particular modules

| Entry   | Type                    | Description   |
|---|-------------------------|---|
| <code>uPDF_uPDF(x1,x2,<br/>bt,mu,zeta,h1,h2)</code> | <code>real(-5:5)</code> | Returns the value of product $f_q f_{\bar{q}}$ evolved unpolarized TMD distributions for given (x,b) (reals) and hadrons $h_1$ and $h_2$ . The value is the vector over flavors. If $h_{1,2}$ have the same sign the product is $f_q f_{\bar{q}}$ , if $h_{1,2}$ have different signs the product is $f_q f_q$ . This combination is used to compute Drell-Yan-type processes, and a bit faster than a direct multiplication. |



### A. `_model` modules

same for `uTMDFF_model` and `lpTMDPDF_model`.

This module gives the expression for  $f_{NP}$ ,  $\mu_{OPE}$  and  $b^*$ . It could be written by user, and one of modules that form a TMD model (together with models for TMD-evolution).

Making this module follow the interface

```
public:: ModelInitialization
public:: ModelUpdate
real(dp),public,dimension(-5:5):: FNP
real(dp),public:: bSTAR
real(dp),public:: muOPE
If there is a twist-3 part of small-b matching, there are extra functions
real(dp),public,dimension(-5:5):: FNP_tw3
real(dp),public:: bSTAR_tw3
real(dp),public:: muOPE_tw3
which contribute to the twist-3 term.
```

`ModelInitialization(NParray)`: this subroutine is called during the initialization of `XX` module. The `integer,intent(in)::NPlength(:)` is the size of the array of NP-parameters. Use this subroutine to make preparation of the model.

`ModelUpdate(NParray)`: this subroutine is called during the `XX_setNPparameters(NParray)`. The `real(dp),intent(in)::NParray(:)` is the array of NP-parameters passed to `TMDR_setNPparameters`. Use this subroutine to recompute model parameters (if needed), save NP-parameters (if needed), etc.

`FNP(x,b,hadron,lambdaNP)`: the function returns  $f_{NP}^{f,h}(x,b,\lambda)$ .

`bSTAR(b,x,y)`: the function returns  $b^*(b)$  function.

`mu_OPE(b,x,y,c4)`: the function returns  $\mu_{OPE}(z,b)$  function.

**Important:** the module `XX_model` is compiled before `XX`, and thus can use only lower-level modules: `TMD_AD`, `QCDinput`, etc.

### B. `_OPE` modules (`tw2`)

OPE modules compute the following convolution

$$[C \times f](x) = \int_x^1 \frac{dy}{y} C(y, \mathbf{L}) f\left(\frac{x}{y}, \mu_{OPE}\right) \quad (8.8)$$

$$= \frac{1}{x} \int_x^1 dy C(y, \mathbf{L}) F\left(\frac{x}{y}, \mu_{OPE}\right), \quad (8.9)$$

where  $F(x) = xf(x)$  and

$$\mathbf{L} = \ln \left( \frac{\mu_{OPE}^2 b^2}{4e^{-2\gamma_E}} \right). \quad (8.10)$$

OPE modules are initialized automatically from the corresponding TMD module.

#### Extra notes:

- The convolution is convergent, but the convolution kernel has singularity at  $y \rightarrow 1$ . To stabilize the integral, it is computed as

$$\int_x^1 dy(\dots) = \int_x^{y_C} dy(\dots) + \int_{y_C}^1 dy(\dots). \quad (8.11)$$

The first term is computed by the adaptive algorithm. The second term is computed by the approximate formulas

$$\begin{aligned}
\int_{y_C}^1 dy \frac{f(x/y) - f(x)}{1-y} &= f(x/y_C) - f(x), \\
\int_{y_C}^1 dy \frac{f(x/y) - f(x)}{1-y} \ln(1-y) &= (f(x/y_C) - f(x))(\ln(1-y_C) - 1), \\
\int_{y_C}^1 dy \frac{f(x/y) - f(x)}{1-y} \ln^2(1-y) &= (f(x/y_C) - f(x))(2 - \ln(1-y_C) + 2 \ln^2(1-y_C)), \\
\int_{y_C}^1 dy f(x/y) g(y) &= f(x) \int_{y_C}^1 dy g(y),
\end{aligned}$$

where  $g(y)$  is integrable, and the last integral is taken explicitly.

- The parameter  $y_C = 1 - 10^{-4}$  (**for a moment**) is fixed. It provides the accuracy of the last integral (6-7 digits), which is far better than.
- For  $x \sim y_C$  the accuracy drops. In this case, the values of  $y_C$  is recomputed by formula  $y_C = (100 + x)/101$  (i.e. it is 2-digits closer to 1 than  $x$ ).
- **The definition of TMDFF does not perfectly match the definition of collinear FF. There is a relative factor  $1/z^2$ .** So the convolution reads

$$D(z, b) = \int_z^1 \frac{dy}{y} \mathbb{C} \left( \frac{z}{y}, \mathbf{L}_\mu \right) \frac{d(y, \mu)}{y^2} = \frac{1}{z^3} \int_z^1 dy [y^2 C](y, \mathbf{L}_\mu) D\left(\frac{z}{y}, \mu\right), \quad (8.12)$$

where  $D(z) = zd(z)$ .

### 1. Functions

XX=name of module (e.g. uTMDPDF, uTMDFF, etc.)

| Entry                                       | Type          | Description  |
|---|---------------|--|
| <code>XX_OPE_Initialize(file,prefix)</code> | subroutine    | Initialize the module from INI-file.   |
| <code>XX_resetGrid()</code>                 | subroutine    | Force recomputation of the grid. Only if griding is ON.  |
| <code>XX_OPE_testGrid()</code>              | subroutine    | Run the test of grid (see sec.VIII B 4). Only if griding is ON.  |
| <code>XX_OPE_SetPDFreplica(rep,h)</code>    | subroutine    | Switch the PDF replica to <code>rep(int)</code> for hadron <code>h (int)</code> . Recompute the grid (if griding is ON). If the new replica is as before nothing is done.  |
| <code>XX_OPE_SetScaleVariation(c4)</code>   | subroutine    | Change the scale variation parameter $c_4$ (it enters the definition of $\mu_{\text{OPE}}$ ). Recompute the grid (if griding is ON). The parameter (real) <code>c4</code> is restricted to $0.1 < c_4 < 10$ .  |
| <code>XX_OPE_convolution(x,b,h,G)</code>    | dp_real(-5:5) | Returns (8.8) for $(x,b)=((\text{real})x, (\text{real})b)$ , and hadron number (int) <code>h</code> . In the case of <b>NA</b> returns 1. If grid is computed restore from the grid. The flag (T/F) <code>G</code> is optional. It indicates the inclusion of gluon into the result. Note, that if gluons were not included into the grid they are zero.   |
| <code>uTMDPDF_X0_AS(x,mu,mu0,h,G)</code>    | dp_real(-5:5) | Computes the $\text{AS}[\mathcal{X}_0[f_1]]$ as defined in [????????]. This is asymptotic term required to determine the second moment of TMDPDF. $(x,\mu,\mu_0)=((\text{real})x, (\text{real})\mu, (\text{real})\mu_0)$ , and hadron number (int) <code>h</code> . In the case of <b>NA</b> returns 0. The flag (T/F) <code>G</code> is optional. It indicates the inclusion of gluon into the result. <b>For a moment this function is done only for uTMDPDF, but it will be updated to all twist-TMDs</b> |

## 2. Model parameters.

The OPE is fully deterministic formula, but it contains the free scale, and often one implement  $b^*$ -modification. These parameters are written in `model`-module. Namely:

- `bSTAR(bt,x,y)` = is the function used instead of  $b$  in **L** in eqn.(8.8).
- `muOPE(bt,x,y,c4)` = is the function used for  $\mu_{\text{OPE}}$ .

In both cases, the variables  $x$  and  $y$  are defined as in eqn.(8.8). The parameter  $c_4$  is the scale variation parameter.

## 3. Scale variation.

TO BE WRITTEN

## 4. Grid

For fitting procedure one often needs to evaluate TMDs multiple times. For example, for fit performed in [?] the evaluation of single  $\chi^2$  entry requires  $\sim 16 \times 10^6$  calls of `uTMDPDF...`. So, every call of `uTMDPDF...` at NNLO order, requires  $\sim 200$  calls of pdfs, depending on  $x$ ,  $b$  and  $\lambda$ 's, in order to evaluate the integral over  $y$ . In such situations, it is much faster to make a grid of TMD distributions for a given set of non-pertrubative parameters (i.e. the grid is in  $x$  and  $b$ ), and then use this grid for the interpolation of TMD values. The calculation of a grid is not a very fast procedure, nonetheless, for large computation (number of TMD calls  $> 10^4 - 10^5$ ) is more efficient.




FIG. 4: Example of precision extraction from the grid. (left) with LHAPDF (right) with smooth input function (similar to PDF). Both cases the N<sup>3</sup>LO convolution with matching coefficient is included.

The grid used in the artemide 3.0 is Chebyshev grid similar to the one described in [2112.09703]. Along each direction the phase-space is split into subgrids  $[x_0, x_1, \dots, x_n]$  ( $x_n = 1$ ) and  $[b_0, b_1, \dots, b_k]$ . Within each subgrid there some amount of node, distributed as  $\ln(x)$  and  $\ln(b)$  over Chebyshev points. The interpolation is done by barycentric formula.

- For  $x < x_0$  and  $x > 1$  the error is returned.
- For  $b < b_0$  and  $b > b_k$  the value at  $b_0$  or  $b_k$  is returned.

All these parameters can be changed in `constants` file. However, we recommend to use default values, since they were checked. The check could be performed by the call `TestGrid()` subroutine of corresponding module. It computes values for central points of grids-cells from grid and from direct computation. This check can be also called by option in the `constants` file.

### WARNING!

LHAPDF produces the non-smooth interpolation (based on cubic splines), while the Chebychev interpolation is much smoother. Therefore, at some points the difference between Chebyshev and LHAPDF can be significant (up to 50% and more if absolute value of function is small). Practically, one cannot push average precision better than  $10^{-4}$ . In a bulk, it is acceptable. If the grid is tested at smooth input function one can easily reach precision  $\sim 10^{-8}$ .

#### Extra Notes:

- Grid is computed by parallel do (if OMP is enabled).

### 5. Technical notes

#### Convolution

The convolution integral reads

$$I(x) = \int_x^1 \frac{dz}{z} C(z) f\left(\frac{x}{z}\right) f_{NP}(x, z), \quad (8.13)$$

where the function  $C(z)$  has a general form

$$C(z) = C_0 \delta(1-z) + (C_1(z))_+ + C_2(z). \quad (8.14)$$

Here the plus-distribution is understood in the usual way

$$(C_1(z))_+ = C_1(z) - \delta(1-z) \int_0^1 dy C_1(y) dy. \quad (8.15)$$

In NNLO coefficient function there are only possible two  $(\cdot)_+$  terms,  $1/(1-z)$  and  $\ln(1-z)/(1-z)$ .

In order to simplify the integration we rewrite

$$I(x) = \frac{1}{x} \int_x^1 dz C(z) \tilde{f}\left(\frac{x}{z}\right) f_{NP}(x, z), \quad \tilde{f}(z) = z f(z). \quad (8.16)$$

Then the integral is split as

$$I(x) = \frac{1}{x} \left\{ I_2(x) + C_0 \tilde{f}(x) f_{NP}(x, 1) + \tilde{f}(x) f_{NP}(x, 1) \int_0^x dz C_1(z) \right\}, \quad (8.17)$$

where

$$I_2(x) = \int_x^1 dz \left[ C_1(z) \left( \tilde{f}\left(\frac{x}{z}\right) f_{NP}(x, z) - \tilde{f}(x) f_{NP}(x, 1) \right) + C_1(z) \tilde{f}\left(\frac{x}{z}\right) f_{NP}(x, z) \right] \quad (8.18)$$

Presumably, the term  $\sim C_0$  give the dominant contribution  $w$ , since it is  $\sim 1$  whereas the other terms  $\sim a_s$ . Therefore, it serves as the estimation of the integral value, with respect to which the integration convergence is calculated. The convolution integral is evaluated by the G7K15 rule adaptively with given tolerance with respect to  $w$ .

The integral  $I$  is calculated in the procedure `Common_lowScale50` and `Common_lowScale5`, which is common for all twist-2 terms. The integral  $I_2$  is calculated in the iterative procedures `MellinConvolutionVectorPart50` and `MellinConvolutionVectorPart5`, which is common for all twist-2 terms.

**In the case of TMDFF** we have the coefficient function with the structure  $C(z)/z^2$  (plus-distribution,  $\delta$ , etc, are multiplied by  $1/z^2$ ), and collinear function  $f(z)/z^2$ . In this case it is convenient to rewrite

$$I(x) = \int_x^1 \frac{dz}{z} \frac{C(z)}{z^2} \frac{f\left(\frac{x}{z}\right)}{(x/z)^2} f_{NP}(x, z) = \frac{1}{x^3} \int_x^1 dz C(z) \hat{f}\left(\frac{x}{z}\right) f_{NP}(x, z), \quad \hat{f}(z) = z f(z). \quad (8.19)$$

Since the common-code calculates the integral PDF-like convolution, I divide by factor  $1/x^2$  all output of the common-code.

### Coefficient functions

The coefficient function is spit to 6 terms.

$$C_{q \leftarrow q}, \quad C_{q \leftarrow g}, \quad C_{g \leftarrow q}, \quad C_{g \leftarrow g}, \quad C_{q \leftarrow \bar{q}}, \quad C_{q \leftarrow \bar{q}}, \quad C_{q \leftarrow q'}, \quad (8.20)$$

where  $C_{q \leftarrow q}$  and  $C_{q \leftarrow \bar{q}}$  contain *non-singlet* part only. The singlet contribution is given in  $C_{q \leftarrow q'}$ . So, the convolution for a quark (e.g.  $q = u$ ) reads

$$F_u = C_{q \leftarrow q} \otimes q_u + C_{q \leftarrow g} \otimes q_g + C_{q \leftarrow \bar{q}} \otimes q_{\bar{u}} + C_{q \leftarrow q'} \otimes \sum_{f \in q, \bar{q}} q_f. \quad (8.21)$$

For gluon

$$F_g = C_{g \leftarrow g} \otimes q_g + C_{g \leftarrow q} \otimes \sum_{f \in q, \bar{q}} q_f. \quad (8.22)$$

The kernels are split into regular, singular and  $\delta$ -parts, where singular contains  $(\ln(1-x)/(1-x))_+^n$  terms,  $\delta$  contains only  $\delta(1-x)$  terms, and regular is the rest. Singular and  $\delta$ -terms are present only for  $C_{q \leftarrow q}$  and  $C_{g \leftarrow g}$ .

The regular part is decomposed further, into singular (but integrable) terms at  $x \rightarrow 0$  and  $x \rightarrow 1$ , and the finite part. The singular terms are computed exactly, while the finite part is fitted to linear combination of several functions. The basis of fit is such that the expressions are exact for coefficients which do not contain PolyLog functions. These are NLO, leading log [at all orders], large-Nf [at all orders], and some others. The full basis for the regular part is

$$\left\{ \underbrace{\ln \bar{x}, \ln^2 \bar{x}, \ln^3 \bar{x}, \ln^4 \bar{x}, \ln^5 \bar{x}, \frac{1}{x}, \frac{\ln x}{x}, \frac{\ln^2 x}{x}}_{\text{singular-integrable exact}}, \ln x, \ln^2 x, \ln^3 x, \ln^4 x, \ln^5 x, \right. \quad (8.23)$$

$$\left. \underbrace{1, x}_{\text{exact}}, x\bar{x}, x \left( \frac{\ln x}{1-x} + 1 \right), \frac{x \ln^2 x}{1-x}, \frac{x \ln^3 x}{1-x}, x \ln x, x \ln^2 x, x \ln^3 x, x \ln^4 x, x\bar{x} \ln x, x\bar{x} \ln^2 x, x\bar{x} \ln^3 x, \right.$$

$$\left. \bar{x} \ln \bar{x}, \bar{x} \ln^2 \bar{x}, \bar{x} \ln^3 \bar{x}, x\bar{x} \ln \bar{x}, x\bar{x} \ln^2 \bar{x}, x\bar{x} \ln^3 \bar{x}, \ln x \ln \bar{x}, x \ln x \ln \bar{x}, x\bar{x} \ln x \ln \bar{x}, \frac{\ln \bar{x}(1-x^2+2 \ln x)}{1-x}, \frac{\bar{x}(\ln x + x)}{x} \right\}.$$

Decomposed in this basis the coefficient functions are very precise. The typical relative precision is  $10^{-8} - 10^{-10}$  for the finite part (other terms are exact). The worst precision happens at small  $x \sim 10^{-5}$  for  $g \leftarrow g$  channel where relative precision is  $\sim 10^{-5} - 10^{-6}$ .

Similar ansatz is used for uTMDFF, with some term removed (which are sensitive to small- $x$  region). Precision in the case of TMDFF is slightly worse  $10^{-6} - 10^{-8}$ . The worst precision is for N<sup>3</sup>LO  $q \rightarrow q'$  channel, where it is  $\sim 10^{-4} - 10^{-6}$ . However, it is very good precision which guarantees about 10-12 precise digits for the physical convolution.

### Computation of $\text{AS}[\mathcal{X}_0]$

The function  $\text{AS}\mathcal{X}_0$  is defined in [????????] and reads

$$\begin{aligned} \text{AS}[\mathcal{X}_0[F]] = & \alpha_s \left\{ P_1 + \alpha_s [P_2 + (C_1 - P_1) \otimes (P_1 - \beta_0)] \right. \\ & + \alpha_s^2 [P_3 + (C_1 - P_1) \otimes (P_2 - \beta_1) + (C_2 - P_2) \otimes (P_1 - 2\beta_0) \\ & \left. - (C_1 - P_1) \otimes (P_1 - \beta_0) \otimes (P_1 - 2\beta_0)] + \mathcal{O}(\alpha_s^3) \right\} \otimes f. \end{aligned} \quad (8.24)$$

It could be obtained by computation of ordinary optimal small- $b$  convolution with replacement of finite terms by 0, and logarithm terms by  $\mathbf{L}_b^k \rightarrow -k!$ . Thus, the function can be computed as ordinary convolution with the coefficient function  $X$  defined as (up to N<sup>3</sup>LO)

$$X = \frac{3}{4}C[\mathbf{L}_b = 0] - \frac{2}{3}C[\mathbf{L}_b = 1] - \frac{1}{12}C[\mathbf{L}_b = 4]. \quad (8.25)$$

### C. OPE modules (tw3)

OPE modules compute the following convolution

$$[C \times T](x) = \int [dy] C_{\text{tw3}}(y_{1,2,3}, \mathbf{L}) T(y_{1,2,3}, \mu_{\text{OPE}}), \quad (8.26)$$

where  $T$  is a twist-three function (qqq-correlator) and

$$\mathbf{L} = \ln \left( \frac{\mu_{\text{OPE}}^2 b^2}{4e^{-2\gamma_E}} \right). \quad (8.27)$$

OPE modules are initialized automatically from the corresponding TMD module.

**Twist-3 convolution is not implemented yet. All function are placeholders. The only working order is NA.**

#### 1. Functions

XX=name of module (e.g. uTMDPDF, uTMDFF, etc.)

| Entry                            | Type          | Description   |
|----------------------------------|---------------|---|
| XX_OPE.Initialize(file,prefix)   | subroutine    | Initialize the module from INI-file. (common with tw2-part)   |
| XX.tw3_resetGrid()               | subroutine    | Force recomputation of the grid. Only if griding is ON.   |
| XX_OPE.tw3_testGrid()            | subroutine    | Run the test of grid (see sec.VIII B 4). Only if griding is ON.   |
| XX_OPE.tw3_SetPDFreplica(rep,h)  | subroutine    | Switch the PDF replica to <code>rep(int)</code> for hadron <code>h (int)</code> . Recompute the grid (if griding is ON). If the new replica is as before nothing is done.   |
| XX_OPE.tw3_SetScaleVariation(c4) | subroutine    | Change the scale variation parameter $c_4$ (it enters the definition of $\mu_{\text{OPE}}$ ). Recompute the grid (if griding is ON). The parameter (real) <code>c4</code> is restricted to $0.1 < c_4 < 10$ .   |
| XX_OPE.tw3_convolution(x,b,h,G)  | dp_real(-5:5) | Returns (8.26) for $(x,b)=((\text{real})x, (\text{real})b)$ , and hadron number (int) <code>h</code> . In the case of <code>NA</code> returns 1. If grid is computed restore from the grid. The flag (T/F) <code>G</code> is optional. It indicates the inclusion of gluon into the result. Note, that if gluons were not included into the grid they are zero. |

## 2. Model parameters.

The OPE is fully deterministic formula, but it contains the free scale, and often one implement  $b^*$ -modification. These parameters are written in `model`-module. Namely:

- `bSTAR(bt,x,y)` = is the function used instead of  $b$  in  $\mathbf{L}$  in eqn.(8.8).
- `muOPE(bt,x,y,c4)` = is the function used for  $\mu_{\text{OPE}}$ .

In both cases, the variables  $x$  and  $y$  are defined as in eqn.(8.8). The parameter  $c_4$  is the scale variation parameter.

## 3. Scale variation.

**Not implemented yet**

## 4. Grid

**Not implemented yet**

## D. Fourier to $k_T$ space

We define

$$F(x, \mathbf{k}_T; \mu, \zeta) = \int \frac{d^2 \mathbf{b}}{(2\pi)^2} F(x, \mathbf{b}; \mu, \zeta) e^{-i(\mathbf{k}_T \mathbf{b})}. \quad (8.28)$$

Since all evaluated TMDs depends only on the modulus of  $\mathbf{b}$ , within the module we evaluate Hankel transformations. These transformations depends on the kind of TMD. Generally the integrals are

$$F(x, |\mathbf{k}_T|; \mu, \zeta) = \frac{M^{2n}}{n!} \int \frac{|b| d|\mathbf{b}|}{2\pi} \left( \frac{|\mathbf{b}|}{|\mathbf{k}_T|} \right)^n J_n(|\mathbf{b}| |\mathbf{k}_T|) F(x, |\mathbf{b}|; \mu, \zeta), \quad (8.29)$$

where the parameter  $M$  is the mass-scale (defined globally for all modules). The value of  $n$  depends on the type of the TMD distributions. Particularly, there are

- For `uTMDPDF`, `uTMDFF`

$$n = 0, \quad F(x, |\mathbf{k}_T|; \mu, \zeta) = \int \frac{d|\mathbf{b}|}{2\pi} |\mathbf{b}| J_0(|\mathbf{b}| |\mathbf{k}_T|) F(x, |\mathbf{b}|; \mu, \zeta),$$

- For `SiversTMDPDF`, `BoerMuldersTMDPDF`, `wgtTMDPDF`

$$n = 1, \quad F(x, |\mathbf{k}_T|; \mu, \zeta) = \frac{M^2}{k_T} \int \frac{d|\mathbf{b}|}{2\pi} |\mathbf{b}|^2 J_1(|\mathbf{b}| |\mathbf{k}_T|) F(x, |\mathbf{b}|; \mu, \zeta),$$

- For `lpTMDPDF`

$$n = 2, \quad F(x, |\mathbf{k}_T|; \mu, \zeta) = \frac{M^4}{2k_T^2} \int \frac{d|\mathbf{b}|}{2\pi} |\mathbf{b}|^3 J_2(|\mathbf{b}| |\mathbf{k}_T|) F(x, |\mathbf{b}|; \mu, \zeta),$$

**Clenshaw-Curtis-Levin algorithm.** The following algorithm provides sufficient precision in the large range of  $k_T$ . It is a mixture of Clenshaw-Curtis quadrature and Levin differential-equation approach. First split the integral into ranges

$$\{B_0, \dots, B_N\}, \quad (8.30)$$

where  $b_0$  is sufficiently small (default  $B_0 = 10^{-6}$ ) and  $B_n$  is sufficiently large (default  $B_n \sim 25 - 100$ ). Then the integral is split as

$$F(q) = \int_0^\infty db J_0(bq) f(b) = \sum_{n=1}^N \int_{B_{n-1}}^{B_n} db J_0(bq) f(b) + \Delta_0 + \Delta_N, \quad (8.31)$$

where

$$\Delta_0 = \int_0^{B_0} db J_0(bq) f(b), \quad \Delta_N = \int_{B_N}^\infty db J_0(bq) f(b).$$

Then for each sector one can estimate integral using different approaches. Here, I use the approaches that are quadratures over the nodes of Chebishev polynomials with

$$t_i = \cos \frac{i\pi}{K}, \quad (8.32)$$

where  $N$  is the size of the grid. So, for each interval  $[B_{n-1}, B_n]$  one defines a grid with  $(K+1)$  points (ultimate points enters two sub-grids). The points are distributed logarithmically. So

$$b_{nk} = \exp(\delta_n t_k + \bar{b}_n), \quad \delta_n = \frac{B_n - B_{n-1}}{2}, \quad \bar{b}_n = \frac{B_n + B_{n-1}}{2}, \quad (8.33)$$

where  $n \in [1, N]$  and  $k \in [0, K]$ . The integral is then defined as

$$\int_{B_{n-1}}^{B_n} db J_0(bq) f(b) = \sum_{k=0}^K w_{nk}(q) f(b_{nk}), \quad (8.34)$$

where  $w_{nk}(q)$  is precomputed by different algorithm.

The *Clenshaw-Curtis* quadrature leads to

$$\int_{B_{n-1}}^{B_n} db J_0(bq) f(b) = \int_{-1}^1 dt \delta_n b(t) J_0(b(t)) f(b(t)) = \sum_{k=0}^K w_{nk}^{\text{CC}}(q) f(b_{nk}), \quad (8.35)$$

where

$$w_{nk}^{\text{CC}}(q) = \delta_n b_{nk} J_0(b_{nk}) a_k, \quad a_k = \frac{4\beta_k}{K} \sum_{l=0,2,\dots}^K \frac{\beta_l}{1-l^2} \cos \frac{kl\pi}{K}, \quad (8.36)$$



with  $\beta_i = 1/2$  for  $i = 1$  or  $i = K$  and  $\beta_i = 1$  otherwise.

The *Levin* approach consists in the consideration of the differential system (*to be written later*), which can be bootstrapped on the grid, and leads to the system of linear equations. In case of present integral it requires the inversion of the Matrix

$$A(q) = \begin{pmatrix} \frac{D}{b\delta} & qI \\ -qI & \frac{D}{b\delta} - \frac{I}{b} \end{pmatrix}, \quad (8.37)$$

where  $D$  is the derivative matrix for the Chebyshev interpolation, and  $1/b$  is understood as division by  $b_i$  for corresponding row. The inverse of this matrix is made by Crout's method (the routine is derived from <https://github.com/Beliavsky/Matrix.Inversion>). Then the weights are

$$w_{nk}^L(q) = J_0(B_n q) A_{1k}^{-1} - J_0(B_{n-1} q) A_{K+1,k}^{-1} + J_1(B_n q) A_{K+2,k}^{-1} - J_1(B_{n-1} q) A_{2K+2,k}^{-1}. \quad (8.38)$$

Note, that it can be written as vector sum

$$\mathbf{w}_n^L(q) = J_0(B_n q) \mathbf{A}_1^{-1} - J_0(B_{n-1} q) \mathbf{A}_{K+1}^{-1} + J_1(B_n q) \mathbf{A}_{K+2}^{-1} - J_1(B_{n-1} q) \mathbf{A}_{2K+2}^{-1}, \quad (8.39)$$

where  $\mathbf{A}_k$  is  $[1, K]$  part of the  $k$ 'th row.

This approaches are used in different ranges.

- *Oscillatory integral* If the Bessel function is already oscillating in the range. In this case the integral is computed by Levin method.
- *Stable integral* If the Bessel function is not yet oscillating in the range. In this case the Clenshaw-Curtis quadrature is used.

Basically, for small-(bq) one should use CC quadrature, and for other cases Levin quadrature. To distinguish these case I use the inequality

$$w_{nk}(q) = \begin{cases} w_{nk}^{CC}(q), & \text{if } B_n q < j_0 \alpha, \\ w_{nk}^L(q), & \text{otherwise,} \end{cases} \quad (8.40)$$

where  $j_0$  is the first zero of Bessel function, and  $\alpha$  is a parameter of order 1 (default  $\alpha = 1$ ).

Finally, the correction to the integral could be estimated. The tale correction is known [cite]


$$\Delta_N \sim f(B_N), \quad (8.41)$$

so using sufficiently large  $B_N$  we get very small correction. The first point correction can be estimated assuming smooth and non-singular behavior of  $f(b)$

$$\Delta_0 = \int_0^{B_0} db J_0(bq) f(b) \simeq \int_0^{B_0} db J_0(bq) b \frac{f(B_0)}{B_0} = \frac{J_1(B_0 q)}{q} f(B_0) \simeq B_0 \frac{f(B_0)}{2}. \quad (8.42)$$

Practically, I add ( $n = 1$ ,  $k = K$  corresponds to  $B_0$  node)

$$w_{1K} \rightarrow w_{1K} + \frac{J_1(B_0 q)}{q}. \quad (8.43)$$



Example of Hankel transform. Test function is  $f(b) = be^{-2.1b^2}(1 + 6.3b^2 + 1.4b^6)$ . Shown is logarithm of ratio to the exact evaluation. Here the sub-grids are  $B = \{0.00001, 0.01, 0.2, 2., 8., 25.\}$  with  $K = 16$ .

### E. TMM

In ref.[2402.01836] the following TMMs are defined

$$\mathcal{G}_{n,n}[F] = \mathcal{G}[F] = \frac{1}{n!} \int_0^\infty db \mu \left( \frac{\mu b}{2} \right)^n J_{n+1}(\mu b) F(x, b) \quad (8.44)$$

$$\begin{aligned} \mathcal{G}_{n+1,n}[F] = \mathcal{X}[F] &= \frac{1}{2M^2 n!} \int_0^\infty db \mu^3 \left( \frac{\mu b}{2} \right)^n \frac{(n+1)J_{n+1}(\mu b) - J_{n+3}(\mu b)}{n+2} F(x, b) \\ &= \frac{1}{2M^2 n!} \int_0^\infty db \mu^3 \left( \frac{\mu b}{2} \right)^n \left( J_{n+1}(\mu b) - \frac{J_{n+2}(\mu b)}{(\mu b)} \right) F(x, b) \end{aligned} \quad (8.45)$$

The correspond to the cut-integral over TMD and particular TMMs. The value of n is defined by the type of TMD (see sec.VIII D).

## IX. TMDs MODULE

The module **TMDs** joins the lower modules and performs the evaluation of various TMD distributions in the  $\zeta$ -prescription. Generally a TMD distribution is given by the expression

$$F_f(x, b; \mu, \zeta) = R_f[b, (\mu, \zeta) \rightarrow (\mu_i, \zeta_{\mu_i})] \tilde{F}_f(x, b), \quad (9.1)$$

where  $R$  is the TMD evolution kernel,  $\tilde{F}$  is a TMD distribution at low scale. The scale  $\mu_i$  is dependent on the evolution type, and could be out of use. Note, that **TMDs** initializes the lower modules automatically. Therefore, no special initializations should be done.

List of available commands

| Command                          | Type           | Sec. | Short description   |
|----------------------------------|----------------|------|---|
| TMDs_Initialize(file,prefix)     | subrout.       | -    | Initialization of module. (string) <b>file</b> is the name of <b>constants-file</b> , which contains initialization information. (string) <b>prefix</b> is appended to <b>file</b> if provided. |
| TMDs_SetScaleVariations(c1,c3)   | subrout.       | IX D | Set new values for the scale-variation constants.   |
| uTMDPDF_5(x,b,mu,zeta,h)         | (real*8(-5:5)) | IX B | Unpolarized TMD PDF (gluon term undefined)  |
| uTMDPDF_50(x,b,mu,zeta,h)        | (real*8(-5:5)) | IX B | Unpolarized TMD PDF (gluon term defined)  |
| uTMDFF_5(x,b,mu,zeta,h)          | (real*8(-5:5)) | IX B | Unpolarized TMD FF (gluon term undefined)   |
| uTMDFF_50(x,b,mu,zeta,h)         | (real*8(-5:5)) | IX B | Unpolarized TMD FF (gluon term defined)   |
| lpTMDPDF_50(x,b,mu,zeta,h)       | (real*8(-5:5)) | IX B | Linearly polarized gluon TMD PDF (quarks are zero)  |
| SiversTMDPDF_5(x,b,mu,zeta,h)    | (real*8(-5:5)) | IX B | Sivers TMD PDF (gluon term undefined)   |
| SiversTMDPDF_50(x,b,mu,zeta,h)   | (real*8(-5:5)) | IX B | Sivers TMD PDF (gluon term defined)   |
| wgtTMDPDF_5(x,b,mu,zeta,h)       | (real*8(-5:5)) | IX B | worm-gear T TMD PDF (gluon term undefined)  |
| wgtTMDPDF_50(x,b,mu,zeta,h)      | (real*8(-5:5)) | IX B | worm-gear T TMD PDF (gluon term defined)  |
| uTMDPDF_5(x,b,h)                 | (real*8(-5:5)) | IX B | Unpolarized TMD PDF at optimal line (gluon term undefined)  |
| uTMDPDF_50(x,b,,h)               | (real*8(-5:5)) | IX B | Unpolarized TMD PDF at optimal line (gluon term defined)  |
| uTMDFF_5(x,b,h)                  | (real*8(-5:5)) | IX B | Unpolarized TMD FF at optimal line (gluon term undefined)   |
| uTMDFF_50(x,b,h)                 | (real*8(-5:5)) | IX B | Unpolarized TMD FF at optimal line (gluon term defined)   |
| lpTMDPDF_50(x,b,,h)              | (real*8(-5:5)) | IX B | Linearly polarized gluon TMD PDF at optimal line (quarks are zero)  |
| SiversTMDPDF_5(x,b,h)            | (real*8(-5:5)) | IX B | Sivers TMD PDF at optimal line (gluon term undefined)   |
| SiversTMDPDF_50(x,b,,h)          | (real*8(-5:5)) | IX B | Sivers TMD PDF at optimal line (gluon term defined)   |
| wgtTMDPDF_5(x,b,h)               | (real*8(-5:5)) | IX B | Worm gear T TMD PDF at optimal line (gluon term undefined)  |
| wgtTMDPDF_50(x,b,,h)             | (real*8(-5:5)) | IX B | Worm gear T TMD PDF at optimal line (gluon term defined)  |
| BoerMuldersTMDPDF_5(x,b,h)       | (real*8(-5:5)) | IX B | Sivers TMD PDF at optimal line (gluon term undefined)   |
| BoerMuldersTMDPDF_50(x,b,,h)     | (real*8(-5:5)) | IX B | Sivers TMD PDF at optimal line (gluon term defined)   |
| uPDF_uPDF(x1,x2,b,mu,zeta,h1,h2) | (real*8(-5:5)) | IX C | Product of Unpolarized TMD PDF $f_{q\leftarrow h_1}(x_1)f_{\bar{q}\leftarrow h_1}$ at the same scale (gluon term undefined)   |

List of functions which must be provided by a model code

| Input                  | Short description   |
|------------------------|---|
| <code>mu_LOW(b)</code> | The value of $\mu_i$ used in the evolutions of type 1 and 2 (improved $\mathcal{D}$ and $\gamma$ ). See [? ]. |
| <code>mu0(b)</code>    | The value of $\mu_0$ used in the evolution of type 1 (improved $\mathcal{D}$ ). See [? ].                     |

### A. Definition of low-scales

The low scales  $\mu_i$  and  $\mu_0$  are defined in the functions `mu_LOW(bt)` and `mu0(bt)` which can be found in the end of `TMDs.f90` code. Modify it if needed.

### B. Evaluating TMDs

The expression for unpolarized TMD PDF is obtained by the functions

`(real*8(-5:5))uTMDPDF_5(x,b,mu,zeta,h)`

where

`x` (real\*8) Bjorken- $x$  ( $0 < x < 1$ )

`b` (real\*8) Transverse distance ( $b > 0$ ) in GeV

`mu` (real\*8) The scale  $\mu_f$  in GeV. Typically,  $\mu_f = Q$ .

`zeta` (real\*8) The scale  $\zeta_f$  in GeV<sup>2</sup>. Typically,  $\zeta_f = Q^2$ .

`h` (integer) The hadron type. If  $h < 0$ , the hadron is anti-hadron.

This function return the vector `real*8(-5:5)` for  $\bar{b}, \bar{c}, \bar{s}, \bar{u}, \bar{d}, ?, d, u, s, c, b$ .

- Gluon contribution in `uTMDPDF_5` is undefined, but taken into account in the mixing contribution. The point is that evaluation of gluons slow down the procedure approximately by 40%, and often is not needed. To calculate the full flavor vector with the gluon TMD, call `uTMDPDF_50(x,b,mu,zeta,h)`, where all arguments defined in the same way.
- The other TMDs, such as unpolarized TMDFF, transversity, etc. are obtained by similar function see the table in the beginning of the section.
- Each function has version without parameters `mu` and `zeta`. It corresponds to the evaluation of a TMS at optimal line [? ]. Practically, it just transfers the outcome of corresponding TMD module, e.g. module `uTMDPDF`, see sec.??.
- For  $h < 0$  the quark-distributions are replaced by anti-quark distributions and vice versa. I.e.  $F(-h) = F(h)(5 : -5 : -1)$  [it is done in TMDs-Modules]

### C. Products of TMDs

The the evaluation of majority of cross-sections one needs the product of two TMDs at the same scale. There are set of functions which return these products. They are slightly faster then just evaluation and multiplication, due to the flavor blindness of the TMD evolution. The function have common interface

`(real*8(-5:5)) uPDF_uPDF(x1,x2,b,mu,zeta,h1,h2)`

where

`x1,x2` (real\*8) Bjorken- $x$ 's ( $0 < x < 1$ )

`b` (real\*8) Transverse distance ( $b > 0$ ) in GeV

`mu` (real\*8) The scale  $\mu_f$  in GeV. Typically,  $\mu_f = Q$ .

`zeta` (real\*8) The scale  $\zeta_f$  in GeV<sup>2</sup>. Typically,  $\zeta_f = Q^2$ .

`h1,h2` (integer) The hadron's types.

The function return a product of the form  $F_{f_1 \leftarrow h_1}(x_1, b; \mu, \zeta) F_{f_2 \leftarrow h_2}(x_2, b; \mu, \zeta)$ , where  $f_{1,2}$  and the type of TMDs depend on the function.

#### D. Theoretical uncertainties

`TMDs_SetScaleVariations(c1,c3,c4)` changes the scale multiplicative factors  $c_i$  (see [? ], sec.6). The default set of arguments is (1,1,1), i.e. the scales as they given in corresponding functions. This subroutine changes  $c1$  and  $c3$  constants and call corresponding subroutines for variation of  $c4$  in TMD defining packages. Note, that in some types of evolution particular variations absent.

## X. TMDF MODULE

This module evaluates the structure functions, that are universally defined as

$$F(Q^2, q_T, x_1, x_2, \mu, \zeta_1, \zeta_2) = \int_0^\infty \frac{bdb}{2} b^n J_n(bq_T) \sum_{ff'} z_{ff'}(Q^2) F_1^f(x_1, b; \mu, \zeta_1) F_2^{f'}(x_2, b; \mu, \zeta_2), \quad (10.1)$$

where

- $Q^2$  is hard scale.
- $q_T$  is transverse momentum in the factorization frame. It coincides with measured  $q_T$  in center-mass frame for DY, but  $q_T \sim p_T/z$  for SIDIS.
- $x_1$  and  $x_2$  are parts of collinear parton momenta. I.e. for DY  $x_{1,2} \simeq Qe^{\pm y}/\sqrt{s}$ , while for SIDIS  $x_2 \sim z$ . It can also obtain power correction, ala Nachmann variables.
- $\mu$  is the hard factorization scale  $\mu \sim Q$
- $\zeta_{1,2}$  are rapidity factorization scales. In the standard factorization scheme  $\zeta_1 \zeta_2 = Q^4$ . It can also be modified by power corrections.
- $f, f'$  are parton flavors.
- $z_{ff'}$  is the process related function. E.g. for photon DY on  $p + \bar{p}$ ,  $z_{ff'} = \delta_{ff'} |e_f|^2$ .
- $n$  The order of Bessel transformation is defined by structure function. E.g. for unpolarized DY  $n = 1$ . For SSA's  $n = 1$ . In general for angular modulation  $\sim \cos(n\theta)$ .
- $F_{1,2}^f$  TMD distribution (PDF or FF) of necessary polarization and flavor.

Some structure functions are multiplied by a constant with mass dimension (e.g. Siverts asymmetry). This constant (called global mass scale =  $M$ ) is defined globally in the constants file.

List of available commands

| Command  | Type     | Sec. | Short description   |
|--|----------|------|---|
| <code>TMDF_Initialize(file, prefix)</code>               | subrout. | X A  | Initialization of module. (string) <code>file</code> is the name of <code>constants-file</code> , which contains initialization information. (string) <code>prefix</code> is appended to <code>file</code> if provided. |
| <code>TMDF_ShowStatistic()</code>                        | subrout. | –    | Print current statistic on the number of calls.   |
| <code>TMDF_ResetCounters()</code>                        | subrout. | –    | Reset intrinsic counters. E.g. release convergence/lost state.  |
| <code>TMDF_F(Q2, qT, x1, x2, mu, zeta1, zeta2, N)</code> | (real*8) | X B  | Evaluates the structure function  |

### A. Initialization

Prior the usage module is to be initialized (once per run) by  
`call TMDF_Initialize(file)`  
 It reads `constants-file` and initialize it-self accordingly.

### B. Evaluating Structure functions

The value of the structure function is obtained by  
`(real*8)TMDF_F(Q2, qT, x1, x2, mu, zeta1, zeta2, N)`  
 where

Q2 (real\*8) hard scale in  $\text{GeV}^2$ .

qT (real\*8) modulus of transverse momentum in the factorization frame in  $\text{GeV}$ ,  $q_T > 0$

x1 (real\*8)  $x$  passed to the first TMD distribution ( $0 < x_1 < 1$ )

x2 (real\*8)  $x$  passed to the first TMD distribution ( $0 < x_2 < 1$ )

mu (real\*8) The hard scale  $\mu$  in  $\text{GeV}$ . Typically,  $\mu = Q$ .

zeta1 (real\*8) The scale  $\zeta_f$  in  $\text{GeV}^2$  for the first TMD distribution. Typically,  $\zeta_f = Q^2$ .

zeta2 (real\*8) The scale  $\zeta_f$  in  $\text{GeV}^2$  for the second TMD distribution. Typically,  $\zeta_f = Q^2$ .

N (integer) The number of process.

The function returns the value of

$$F^N(Q^2, q_T, x_1, x_2, \mu, \zeta_1, \zeta_2) = \int_0^\infty \frac{bdb}{2} b^n J_n(bq_T) \sum_{ff'} z_{ff'}^N(Q^2) F_1^f(x_1, b; \mu, \zeta_1) F_2^{f'}(x_2, b; \mu, \zeta_2). \quad (10.2)$$

The parameter  $n$  depends on the argument N and uniformly defined as

|         |                            |
|---------|----------------------------|
| $n = 0$ | for $N < 10000$            |
| $n = 1$ | for $N \in [10000, 20000]$ |
| $n = 2$ | for $N \in [20000, 30000]$ |
| $n = 3$ | for $N > 30000$            |

The particular values of  $z_{ff'}$  and  $F_{1,2}$  are given in the following table. User function can be implemented by code modification.

**Notes on the integral evaluation:**

- The integral is uniformly set to 0 for  $q_T < 10^{-7}$ .
- The integrand is uniformly set to 0 for  $b > 10^3$ .
- If for any element of evaluation (including TMD evolution factors and convolution integrals, and the integral of the structure function it-self) obtained divergent value. The trigger is set to ON. In this case, the integral returns uniformly large value  $10^{100}$  for all integrals without evaluation. The trigger is reset by new values of NP parameters. It is done in order to cut the improper values of NP parameters in the fastest possible way, which speed up fitting procedures.
- The Fourier is made by Ogata quadrature, which is double exponential quadrature. I.e.

$$\int_0^\infty \frac{bdb}{2} b^n I(b) J_n(q_T b) \simeq \frac{1}{q_T^{n+2}} \sum_{k=1}^\infty \tilde{\omega}_{nk} b_{nk}^{n+1} I\left(\frac{b_{nk}}{q_T}\right), \quad (10.3)$$

where

$$b_{nk} = \frac{\psi(\tilde{h}\tilde{\xi}_{nk})}{\tilde{h}} = \tilde{\xi}_{nk} \tanh\left(\frac{\pi}{2} \sinh(\tilde{h}\tilde{\xi}_{nk})\right)$$

$$\tilde{\omega}_{nk} = \frac{J_n(\tilde{b}_{nk})}{\tilde{\xi}_{nk} J_{n+1}^2(\tilde{\xi}_{nk})} \psi'(\tilde{h}\tilde{\xi}_{nk})$$

Here,  $\tilde{\xi}_{nk}$  is  $k$ 'th zero of  $J_n(x)$  function. Note, that  $\tilde{h} = h/\pi$  in the original Ogata's notation.

- The convergence properties of the quadrature essentially depends on  $h$ . I have found that  $h \sim q_T$ . For accurate evolution of integrals at different values of  $q_T$  I set  $h = h_0 s$ , where  $s$  is defined for intervals of  $q_T$ .
- The sum over  $k$  is restricted by  $N_{\max}$  where  $N_{\max}$  is hard coded number,  $N_{\max} = 200$ .



- The sum over  $k$  is evaluated until the sum of absolute values of last four terms is less than *tolerance*. If  $M > N_{\max}$  the integral declared divergent, and the trigger is set to ON.

### WARNING!

The error for Ogata quadrature is defined by parameters  $h$  and  $M$  (number of terms in the sum over  $k$ ). In the parameter  $M$  quadrature is double-exponential, i.e. converges fast as  $M$  approaches  $N_{\max}$ . And the convergence of the sum can be simply checked. In the parameter  $h$  the quadrature is quadratic (the convergence is rather poor). The convergence to the true value of integral is very expensive especially at large  $q_T$  (it requires the complete reevaluation of integral at all nodes). Unfortunately,  $N_{\max} \sim h^{-1}$ , and has to find the balance value for  $h$ . In principle, TMD functions decays rather fast, and suggested default value  $h = 0.005$  is trustful. **Nonetheless, we suggest to test other values (\*2) of  $h$  to validate the obtained values in your model.**

The adaptive check of convergence will implemented in the future versions.

### C. Enumeration of structure functions

List of enumeration of structures functions  
N<10000

| N    | $z_{ff'}$  | $F_1$ | $F_2$ | Short description  | Gluon | LP/KPC |
|------|--|-------|-------|--|-------|--------|
| 0    |  |       |       |  |       |        |
| 9999 | –  | –     | –     | Test cases (see X D)   | no    | LP/KPC |
| 9998 |  |       |       |  |       |        |
| 1    | $\delta_{\bar{f}f'}  e_f ^2$   | $f_1$ | $f_1$ | (unpol.) $h_1 + h_2 \rightarrow \gamma$  | no    | LP/KPC |
| 2    | $\delta_{\bar{f}f'} \frac{(1 - 2 ef s_w^2)^2 - 4e_f^2 s_w^4}{8s_w^2 c_w^2}$                  | $f_1$ | $f_1$ | (unpol.) $h_1 + h_2 \rightarrow Z$   | no    | LP/KPC |
| 3    | $\delta_{\bar{f}f'} \frac{z_{ll'} z_{ff'}}{\alpha_{\text{em}}^2}$ given in (2.8) of [?] ]    | $f_1$ | $f_1$ | (unpol.) $h_1 + h_2 \rightarrow Z + \gamma$  | no    | LP/KPC |
| 4    | $\frac{1}{4s_w^2} \frac{ V_{ff'} ^2}{4s_w^2} \frac{Q^4}{(Q^2 - M_W^2)^2 + \Gamma_W^2 M_W^2}$ | $f_1$ | $f_1$ | (unpol.) $h_1 + h_2 \rightarrow W^+$   | no    | LP/KPC |
| 5    | $\frac{1}{4s_w^2} \frac{ V_{ff'} ^2}{4s_w^2} \frac{Q^4}{(Q^2 - M_W^2)^2 + \Gamma_W^2 M_W^2}$ | $f_1$ | $f_1$ | (unpol.) $h_1 + h_2 \rightarrow W^-$   | no    | LP/KPC |
| 6    | $\frac{1}{4s_w^2} \frac{ V_{ff'} ^2}{4s_w^2} \frac{Q^4}{(Q^2 - M_W^2)^2 + \Gamma_W^2 M_W^2}$ | $f_1$ | $f_1$ | (unpol.) $h_1 + h_2 \rightarrow W^\pm$   | no    | LP/KPC |
| 7    | $\frac{ V_{ff'} ^2}{4s_w^2}$   | $f_1$ | $f_1$ | (unpol.) $h_1 + h_2 \rightarrow W^+$ (for narrow-width approx.)                        | no    | LP/KPC |
| 8    | $\frac{ V_{ff'} ^2}{4s_w^2}$   | $f_1$ | $f_1$ | (unpol.) $h_1 + h_2 \rightarrow W^-$ (for narrow-width approx.)                        | no    | LP/KPC |
| 9    | $\frac{ V_{ff'} ^2}{4s_w^2}$   | $f_1$ | $f_1$ | (unpol.) $h_1 + h_2 \rightarrow W^\pm$ (for narrow-width approx.)                      | no    | LP/KPC |
| 10   | $f_g(x_1) f_g(x_2) + h_1^\perp(x_1) h_1^\perp(x_2)$  | -     | -     | (unpol.) $h_1 + h_2 \rightarrow H$ Elementary Higgs production                         | yes   | LP/KPC |
| 11   | $f_g(x_1) f_g(x_2)$  | $f_1$ | $f_1$ | (unpol.) $h_1 + h_2 \rightarrow H$ Elementary Higgs production. Only unpol.TMDPDF term | yes   | LP/KPC |

|      |   |             |                 |   |     |        |
|------|---|-------------|-----------------|---|-----|--------|
| 12   | $h_{1g}^\perp(x_1)h_{1g}^\perp(x_2)$                                  | $h_1^\perp$ | $h_1^\perp$     | (unpol.) $h_1 + h_2 \rightarrow H$ Elementary Higgs production. Only linearly pol.TMDPDF term   | yes | LP/KPC |
| 101  | $R_{ff'}^{Cu}e_f e_{f'}$ see (3.1) of [?] ]                           | $f_1$       | $f_1^{Cu}$      | (unpol.) $h_1 + Cu \rightarrow \gamma$ where the isostructure of Cu is simulated from $hadron = 1$ , (used to describe E288 experiment)   | no  | LP/KPC |
| 102  | $R_{ff'}^{2H}e_f e_{f'}$ see (3.1) of [?] ] with $Z = 1$ and $A = 2$  | $f_1$       | $f_1^{2H}$      | (unpol.) $h_1 + {}^2H \rightarrow \gamma$ where the isostructure of ${}^2H$ is simulated from $hadron = 1$ , used to describe E772 experiment   | no  | LP/KPC |
| 103  | $R_{ff'}^W e_f e_{f'}$ see (3.1) of [?] ] with $Z = 74$ and $A = 183$ | $f_1$       | $f_1^W$         | (unpol.) $h_1 + W \rightarrow \gamma$ , where the isostructure of $W$ (tungsten) is simulated from $hadron = 1$ , used to describe E537 experiment  | no  | LP/KPC |
| 2001 | $\delta_{ff'} e_f ^2$   | $f_1$       | $d_1$           | (unpol) $h_1 + \gamma \rightarrow h_2$  | no  | LP     |
| 2002 | $\delta_{ff'} e_f ^2$   | $f_1^d$     | $d_1$           | (unpol) $d^{(h_1)} + \gamma \rightarrow h_2$ . Here, $d^{(h_1)}$ is the isoscalar target prepared from the hadron $h_1$ . I.e. $u, d \rightarrow (u + d)/2$ . For $h_1$ =proton, this simulates the deuteron target.                        | no  | LP     |
| 2003 | $\delta_{ff'} e_f ^2$   | $f_1^n$     | $d_1$           | (unpol) $n^{(h_1)} + \gamma \rightarrow h_2$ Here, $n^{(h_1)}$ is the iso-conjugated target prepared from the hadron $h_1$ . I.e. $u \leftrightarrow d$ . For $h_1$ =proton, this simulates the neutron target.                             | no  | LP     |
| 2101 | $\delta_{ff'} e_f ^2$   | $f_1$       | $d_1^{h_1+2}$   | (unpol) $h_1 + \gamma \rightarrow h_{1+2}$ , where $h_{1+2} = 1 + 2$ . Computed strictly with produced hadron 1 and 2 (accounts only the sign of $h_2$ argument).   | no  | LP     |
| 2102 | $\delta_{ff'} e_f ^2$   | $f_1$       | $d_1^{h_1+2+3}$ | (unpol) $h_1 + \gamma \rightarrow h_{1+2+3}$ , where $h_{1+2+3} = 1 + 2 + 3$ Computed strictly with produced hadron 1, 2 and 3 (accounts only the sign of $h_2$ argument).  | no  | LP     |
| 2103 | $\delta_{ff'} e_f ^2$   | $f_1^d$     | $d_1^{h_1+2}$   | (unpol) $d^{(h_1)} + \gamma \rightarrow h_{1+2}$ , where $h_{1+2} = 1 + 2$ and $d$ is isoscalar target $u, d \rightarrow (u + d)/2$ . Computed strictly with produced hadron 1 and 2 (accounts only the sign of $h_2$ argument).            | no  | LP     |
| 2104 | $\delta_{ff'} e_f ^2$   | $f_1^d$     | $d_1^{h_1+2+3}$ | (unpol) $d^{(h_1)} + \gamma \rightarrow h_{1+2+3}$ , where $h_{1+2+3} = 1 + 2 + 3$ and $d$ is isoscalar target $u, d \rightarrow (u + d)/2$ . Computed strictly with produced hadron 1, 2 and 3 (accounts only the sign of $h_2$ argument). | no  | LP     |
| 2105 | $\delta_{ff'} e_f ^2$   | $f_1^n$     | $d_1^{h_1+2}$   | (unpol) $n^{(h_1)} + \gamma \rightarrow h_{1+2}$ , where $h_{1+2} = 1 + 2$ and $n$ is neutron target ( $u \leftrightarrow d$ ). Computed strictly with produced hadron 1 and 2 (accounts only the sign of $h_2$ argument).                  | no  | LP     |
| 2106 | $\delta_{ff'} e_f ^2$   | $f_1^n$     | $d_1^{h_1+2+3}$ | (unpol) $n^{(h_1)} + \gamma \rightarrow h_{1+2+3}$ , where $h_{1+2+3} = 1 + 2 + 3$ and $n$ is neutron target ( $u \leftrightarrow d$ ). Computed strictly with produced hadron 1, 2 and 3 (accounts only the sign of $h_2$ argument).       | no  | LP     |

10000 ≤ N &lt; 20000

| N     | $z_{ff'}$  | $F_1$               | $F_2$               | Short description  | Gluon req. |
|-------|--|---------------------|---------------------|--|------------|
| 10000 |  |                     |                     |  |            |
| 19999 | –  | –                   | –                   | Test cases (see X D)   | no         |
| 19998 |  |                     |                     |  |            |
| 10001 | $-M\delta_{\bar{f}f'} e_f ^2$  | $-(f_{1T}^\perp)^p$ | $f_1$               | (Sivers) $h_1^\uparrow + h_2 \rightarrow \gamma$   | no         |
| 10003 | $-M\delta_{\bar{f}f'}\frac{z_{ll'}z_{ff'}}{\alpha_{\text{em}}^2}$ given in (2.8) of [?] ]    | $-(f_{1T}^\perp)^p$ | $f_1$               | (Sivers) $h_1^\uparrow + h_2 \rightarrow Z + \gamma$   | no         |
| 10004 | $-M\frac{1}{4s_w^2}\frac{ V_{ff'} ^2}{4s_w^2}\frac{Q^4}{(Q^2 - M_W^2)^2 + \Gamma_W^2 M_W^2}$ | $-(f_{1T}^\perp)^p$ | $f_1$               | (Sivers) $h_1^\uparrow + h_2 \rightarrow W^+$  | no         |
| 10008 | $-M\frac{1}{4s_w^2}\frac{ V_{ff'} ^2}{4s_w^2}\frac{Q^4}{(Q^2 - M_W^2)^2 + \Gamma_W^2 M_W^2}$ | $-(f_{1T}^\perp)^p$ | $f_1$               | (Sivers) $h_1^\uparrow + h_2 \rightarrow W^-$  | no         |
| 10011 | $+M\delta_{\bar{f}f'} e_f ^2$  | $f_1^h$             | $-(f_{1T}^\perp)^p$ | (Sivers) $h + p^\uparrow \rightarrow \gamma$ (with $h = 2$ )   | no         |
| 12001 | $-M\delta_{ff'} e_f ^2$  | $(f_{1T}^\perp)^p$  | $d_1^{h_1}$         | (Sivers) $p^\uparrow + \gamma \rightarrow h_1$   | no         |
| 1200N | $-M\delta_{ff'} e_f ^2$  | $(f_{1T}^\perp)^p$  | $d_1^{h_N}$         | (Sivers) $p^\uparrow + \gamma \rightarrow h_N$ , for $N = \{1, \dots, 9\}$ (including case 12001)  | no         |
| 12011 | $-M\delta_{ff'} e_f ^2$  | $(f_{1T}^\perp)^d$  | $d_1^{h_1}$         | (Sivers) $d^\uparrow + \gamma \rightarrow h_1$ . Here, d is isoscalar target $(p+n)/2$ .   | no         |
| 1201N | $-M\delta_{ff'} e_f ^2$  | $(f_{1T}^\perp)^d$  | $d_1^{h_N}$         | (Sivers) $d^\uparrow + \gamma \rightarrow h_N$ , for $N = \{1, \dots, 9\}$ (including case 12011). Here, d is isoscalar target $(p+n)/2$ .       | no         |
| 12021 | $-M\delta_{f\bar{f}'} e_f ^2$  | $(f_{1T}^\perp)^p$  | $d_1^{\bar{h}_1}$   | (Sivers) $p^\uparrow + \gamma \rightarrow \bar{h}_1$   | no         |
| 1202N | $-M\delta_{f\bar{f}'} e_f ^2$  | $(f_{1T}^\perp)^p$  | $d_1^{\bar{h}_N}$   | (Sivers) $p^\uparrow + \gamma \rightarrow \bar{h}_N$ , for $N = \{1, \dots, 9\}$ (including case 12021)  | no         |
| 12031 | $-M\delta_{f\bar{f}'} e_f ^2$  | $(f_{1T}^\perp)^d$  | $d_1^{\bar{h}_1}$   | (Sivers) $d^\uparrow + \gamma \rightarrow \bar{h}_1$ . Here, d is isoscalar target $(p+n)/2$ .   | no         |
| 1203N | $-M\delta_{f\bar{f}'} e_f ^2$  | $(f_{1T}^\perp)^d$  | $d_1^{\bar{h}_N}$   | (Sivers) $d^\uparrow + \gamma \rightarrow \bar{h}_N$ , for $N = \{1, \dots, 9\}$ (including case 12031). Here, d is isoscalar target $(p+n)/2$ . | no         |
| 1204N | $-M\delta_{ff'} e_f ^2$  | $(f_{1T}^\perp)^d$  | $d_1^{h_N}$         | (Sivers) $n^\uparrow + \gamma \rightarrow h_N$ , for $N = \{1, \dots, 9\}$ , n is neutron target $n = p(u \leftrightarrow d)$ .                  | no         |
| 1205N | $-M\delta_{ff'} e_f ^2$  | $(f_{1T}^\perp)^d$  | $d_1^{\bar{h}_N}$   | (Sivers) $n^\uparrow + \gamma \rightarrow \bar{h}_N$ , for $N = \{1, \dots, 9\}$ , n is neutron target $n = p(u \leftrightarrow d)$ .            | no         |
| 12101 | $-M\delta_{ff'} e_f ^2$  | $(f_{1T}^\perp)^p$  | $d_1^{h_{1+2}}$     | (Sivers) $p^\uparrow + \gamma \rightarrow h_{1+2}$ , where $h_{1+2} = h_1 + h_2$   | no         |

|  |                               |                    |                   |   |    |
|--|-------------------------------|--------------------|-------------------|---|----|
| 12102  | $-M\delta_{ff'} e_f ^2$       | $(f_{1T}^\perp)^p$ | $d_1^{h_1+2+3}$   | (Sivers) $p^\uparrow + \gamma \rightarrow h_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$  | no |
| 12103  | $-M\delta_{ff'} e_f ^2$       | $(f_{1T}^\perp)^d$ | $d_1^{h_1+2}$     | (Sivers) $d^\uparrow + \gamma \rightarrow h_{1+2}$ , where $h_{1+2} = h_1 + h_2$ and d is isoscalar target $(p+n)/2$ .                                  | no |
| 12104  | $-M\delta_{ff'} e_f ^2$       | $(f_{1T}^\perp)^d$ | $d_1^{h_1+2+3}$   | (Sivers) $d^\uparrow + \gamma \rightarrow h_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$ and d is isoscalar target $(p+n)/2$ .                        | no |
| 12105  | $-M\delta_{ff'} e_f ^2$       | $(f_{1T}^\perp)^n$ | $d_1^{h_1+2}$     | (Sivers) $n^\uparrow + \gamma \rightarrow h_{1+2}$ , where $h_{1+2} = h_1 + h_2$ and n is neutron target $n = p(u \leftrightarrow d)$ .                 | no |
| 12106  | $-M\delta_{ff'} e_f ^2$       | $(f_{1T}^\perp)^n$ | $d_1^{h_1+2+3}$   | (Sivers) $n^\uparrow + \gamma \rightarrow h_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$ and n is neutron target $n = p(u \leftrightarrow d)$ .       | no |
| 12111  | $-M\delta_{f\bar{f}'} e_f ^2$ | $(f_{1T}^\perp)^p$ | $d_1^{h_1+2}$     | (Sivers) $p^\uparrow + \gamma \rightarrow \bar{h}_{1+2}$ , where $h_{1+2} = h_1 + h_2$  | no |
| 12112  | $-M\delta_{f\bar{f}'} e_f ^2$ | $(f_{1T}^\perp)^p$ | $d_1^{h_1+2+3}$   | (Sivers) $p^\uparrow + \gamma \rightarrow \bar{h}_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$  | no |
| 12113  | $-M\delta_{f\bar{f}'} e_f ^2$ | $(f_{1T}^\perp)^d$ | $d_1^{h_1+2}$     | (Sivers) $d^\uparrow + \gamma \rightarrow \bar{h}_{1+2}$ , where $h_{1+2} = h_1 + h_2$ and d is isoscalar target $(p+n)/2$ .                            | no |
| 12114  | $-M\delta_{f\bar{f}'} e_f ^2$ | $(f_{1T}^\perp)^d$ | $d_1^{h_1+2+3}$   | (Sivers) $d^\uparrow + \gamma \rightarrow \bar{h}_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$ and d is isoscalar target $(p+n)/2$ .                  | no |
| 12115  | $-M\delta_{f\bar{f}'} e_f ^2$ | $(f_{1T}^\perp)^n$ | $d_1^{h_1+2}$     | (Sivers) $n^\uparrow + \gamma \rightarrow \bar{h}_{1+2}$ , where $h_{1+2} = h_1 + h_2$ and n is neutron target $n = p(u \leftrightarrow d)$ .           | no |
| 12116  | $-M\delta_{f\bar{f}'} e_f ^2$ | $(f_{1T}^\perp)^n$ | $d_1^{h_1+2+3}$   | (Sivers) $n^\uparrow + \gamma \rightarrow \bar{h}_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$ and n is neutron target $n = p(u \leftrightarrow d)$ . | no |
| $\Downarrow\Downarrow\Downarrow$ Enumeration for $F_{LT}$ in SIDIS is same as for $F_{UT}$ with $(12 \rightarrow 13)$ $\Downarrow\Downarrow\Downarrow$ |                               |                    |                   |   |    |
| 13001  | $+M\delta_{ff'} e_f ^2$       | $(g_{1T}^\perp)^p$ | $d_1^{h_1}$       | (wgt) $p^\uparrow + \gamma \rightarrow h_1$   | no |
| 1300N  | $+M\delta_{ff'} e_f ^2$       | $(g_{1T}^\perp)^p$ | $d_1^{h_N}$       | (wgt) $p^\uparrow + \gamma \rightarrow h_N$ , for $N = \{1, \dots, 9\}$ (including case 12001)  | no |
| 13011  | $+M\delta_{ff'} e_f ^2$       | $(g_{1T}^\perp)^d$ | $d_1^{h_1}$       | (wgt) $d^\uparrow + \gamma \rightarrow h_1$ . Here, d is isoscalar target $(p+n)/2$ .   | no |
| 1301N  | $+M\delta_{ff'} e_f ^2$       | $(g_{1T}^\perp)^d$ | $d_1^{h_N}$       | (wgt) $d^\uparrow + \gamma \rightarrow h_N$ , for $N = \{1, \dots, 9\}$ (including case 12011). Here, d is isoscalar target $(p+n)/2$ .                 | no |
| 13021  | $+M\delta_{f\bar{f}'} e_f ^2$ | $(g_{1T}^\perp)^p$ | $d_1^{\bar{h}_1}$ | (wgt) $p^\uparrow + \gamma \rightarrow \bar{h}_1$   | no |
| 1302N  | $+M\delta_{f\bar{f}'} e_f ^2$ | $(g_{1T}^\perp)^p$ | $d_1^{\bar{h}_N}$ | (wgt) $p^\uparrow + \gamma \rightarrow \bar{h}_N$ , for $N = \{1, \dots, 9\}$ (including case 12021)  | no |
| 13031  | $+M\delta_{f\bar{f}'} e_f ^2$ | $(g_{1T}^\perp)^d$ | $d_1^{\bar{h}_1}$ | (wgt) $d^\uparrow + \gamma \rightarrow \bar{h}_1$ . Here, d is isoscalar target $(p+n)/2$ .   | no |

|       |  |                    |                   |   |    |
|-------|--|--------------------|-------------------|---|----|
| 1303N | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^d$ | $d_1^{\bar{h}_N}$ | (wgt) $d^\dagger + \gamma \rightarrow \bar{h}_N$ , for $N = \{1, \dots, 9\}$ (including case 12031). Here, d is isoscalar target $(p+n)/2$ .        | no |
| 1304N | $+M\delta_{ff'} e_f ^2$  | $(g_{1T}^\perp)^d$ | $d_1^{h_N}$       | (wgt) $n^\dagger + \gamma \rightarrow h_N$ , for $N = \{1, \dots, 9\}$ , n is neutron target $n = p(u \leftrightarrow d)$ .                         | no |
| 1305N | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^d$ | $d_1^{\bar{h}_N}$ | (wgt) $n^\dagger + \gamma \rightarrow \bar{h}_N$ , for $N = \{1, \dots, 9\}$ , n is neutron target $n = p(u \leftrightarrow d)$ .                   | no |
| 13101 | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^p$ | $d_1^{h_{1+2}}$   | (wgt) $p^\dagger + \gamma \rightarrow h_{1+2}$ , where $h_{1+2} = h_1 + h_2$  | no |
| 13102 | $+M\delta_{ff'} e_f ^2$  | $(g_{1T}^\perp)^p$ | $d_1^{h_{1+2+3}}$ | (wgt) $p^\dagger + \gamma \rightarrow h_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$  | no |
| 13103 | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^d$ | $d_1^{h_{1+2}}$   | (wgt) $d^\dagger + \gamma \rightarrow h_{1+2}$ , where $h_{1+2} = h_1 + h_2$ and d is isoscalar target $(p+n)/2$ .                                  | no |
| 13104 | $+M\delta_{ff'} e_f ^2$  | $(g_{1T}^\perp)^d$ | $d_1^{h_{1+2+3}}$ | (wgt) $d^\dagger + \gamma \rightarrow h_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$ and d is isoscalar target $(p+n)/2$ .                        | no |
| 13105 | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^n$ | $d_1^{h_{1+2}}$   | (wgt) $n^\dagger + \gamma \rightarrow h_{1+2}$ , where $h_{1+2} = h_1 + h_2$ and n is neutron target $n = p(u \leftrightarrow d)$ .                 | no |
| 13106 | $+M\delta_{ff'} e_f ^2$  | $(f_{1T}^\perp)^n$ | $d_1^{h_{1+2+3}}$ | (wgt) $n^\dagger + \gamma \rightarrow h_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$ and n is neutron target $n = p(u \leftrightarrow d)$ .       | no |
| 13111 | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^p$ | $d_1^{h_{1+2}}$   | (wgt) $p^\dagger + \gamma \rightarrow \bar{h}_{1+2}$ , where $h_{1+2} = h_1 + h_2$  | no |
| 13112 | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^p$ | $d_1^{h_{1+2+3}}$ | (wgt) $p^\dagger + \gamma \rightarrow \bar{h}_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$  | no |
| 13113 | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^d$ | $d_1^{h_{1+2}}$   | (wgt) $d^\dagger + \gamma \rightarrow \bar{h}_{1+2}$ , where $h_{1+2} = h_1 + h_2$ and d is isoscalar target $(p+n)/2$ .                            | no |
| 13114 | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^d$ | $d_1^{h_{1+2+3}}$ | (wgt) $d^\dagger + \gamma \rightarrow \bar{h}_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$ and d is isoscalar target $(p+n)/2$ .                  | no |
| 13115 | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^n$ | $d_1^{h_{1+2}}$   | (wgt) $n^\dagger + \gamma \rightarrow \bar{h}_{1+2}$ , where $h_{1+2} = h_1 + h_2$ and n is neutron target $n = p(u \leftrightarrow d)$ .           | no |
| 13116 | $+M\delta_{f\bar{f}'} e_f ^2$  | $(g_{1T}^\perp)^n$ | $d_1^{h_{1+2+3}}$ | (wgt) $n^\dagger + \gamma \rightarrow \bar{h}_{1+2+3}$ , where $h_{1+2+3} = h_1 + h_2 + h_3$ and n is neutron target $n = p(u \leftrightarrow d)$ . | no |
| 13200 | $+M\delta_{\bar{f}f'} \frac{\bar{z}_{l'f} z_{ff'}}{\alpha_{\text{em}}^2}$ given in (2.8) of [? ] | $(g_{1T}^\perp)^p$ | $(-1)^f f_1$      | $G_{TU}^1$ cross-section in $h_1^\dagger + h_2 \rightarrow Z + \gamma$  | no |
| 13201 | $+M \frac{1}{4s_w^2} \frac{ V_{ff'} ^2}{4s_w^2} \frac{Q^4}{(Q^2 - M_W^2)^2 + \Gamma_W^2 M_W^2}$  | $(g_{1T}^\perp)^p$ | $(-1)^f f_1$      | $G_{TU}^1$ cross-section in $h_1^\dagger + h_2 \rightarrow W^+$   | no |
| 13202 | $+M \frac{1}{4s_w^2} \frac{ V_{ff'} ^2}{4s_w^2} \frac{Q^4}{(Q^2 - M_W^2)^2 + \Gamma_W^2 M_W^2}$  | $(g_{1T}^\perp)^p$ | $(-1)^f f_1$      | $G_{TU}^1$ cross-section in $h_1^\dagger + h_2 \rightarrow W^-$   | no |

$$20000 \leq N < 30000$$

| N     | $z_{ff'}$ | $F_1$ | $F_2$ | Short description    | Gluon req. |
|-------|-----------|-------|-------|----------------------|------------|
| 20000 |           |       |       |                      |            |
| 29999 | —         | —     | —     | Test cases (see X D) | no         |
| 29998 |           |       |       |                      |            |

| 30000 ≤ N |           |       |       |                      |            |
|-----------|-----------|-------|-------|----------------------|------------|
| N         | $z_{ff'}$ | $F_1$ | $F_2$ | Short description    | Gluon req. |
| 30000     |           |       |       |                      |            |
| 39999     | —         | —     | —     | Test cases (see X D) | no         |
| 39998     |           |       |       |                      |            |

**Notes on parameters and notation:**

$s_w^2$ ,  $M_Z$ ,  $\Gamma_Z$ , etc. are defined in **constants**

$f_1$ -unpolarized TMDPDF

$d_1$ -unpolarized TMDFF

$f_{1T}^\perp$ -Sivers function

$g_{1T}^\perp$ -wgt-function

$(-1)^f$  is (+1) for f=quark, and (-1) for f=anti-quark.

Everywhere proton is expected to be hadron number 1.

**D. Tests**

The test options are called by process number  $N + 0$ ,  $N + 9999$  and  $N + 9998$ . In these case, the integrand set by a fixed function. In the case  $N + 0$

$$N + 0 : \quad \sum z F F \rightarrow e^{-0.2b}. \quad (10.4)$$

The cases  $N + 9999$  and  $N + 9998$  are dependent on parameters of the function:

$$N + 9999 : \quad \sum z F F \rightarrow e^{-\mu b} (1 + x_1 b^2 + x_2 b^4), \quad (10.5)$$

$$N + 9998 : \quad \sum z F F \rightarrow e^{-\mu b^2} (1 + x_1 b^2 + x_2 b^4). \quad (10.6)$$

In all cases the Hankel integral could be evaluated. They read

$$9999 : \quad F = \frac{1}{2\mu^2(1+X)^{3/2}} \left( 1 + \frac{x_1}{\mu^2} \frac{6-9X}{(1+X)^2} + \frac{15x_2}{\mu^4} \frac{8-40X+15X^2}{(1+X)^4} \right), \quad (10.7)$$

$$19999 : \quad F = \frac{3\sqrt{X}}{2\mu^3(1+X)^{5/2}} \left( 1 + \frac{5x_1}{\mu^2} \frac{4-3X}{(1+X)^2} + \frac{105x_2}{\mu^4} \frac{8-20X+5X^2}{(1+X)^4} \right), \quad (10.8)$$

$$29999 : \quad F = \frac{15X}{2\mu^4(1+X)^{7/2}} \left( 1 + \frac{21x_1}{\mu^2} \frac{2-X}{(1+X)^2} + \frac{63x_2}{\mu^4} \frac{48-80X+15X^2}{(1+X)^4} \right), \quad (10.9)$$

$$39999 : \quad F = \frac{105X^{3/2}}{2\mu^5(1+X)^{9/2}} \left( 1 + \frac{9x_1}{\mu^2} \frac{8-3X}{(1+X)^2} + \frac{495x_2}{\mu^4} \frac{16-20X+3X^2}{(1+X)^4} \right), \quad (10.10)$$

where  $X = q_T^2/\mu^2$ .

$$9998 : \quad F = \frac{e^{-Y}}{4\mu} \left( 1 + \frac{x_1}{\mu}(1 - Y) + \frac{x_2}{\mu^2}(2 - 4Y + Y^2) \right), \quad (10.11)$$

$$19998 : \quad F = \frac{e^{-Y}\sqrt{Y}}{4\mu^{3/2}} \left( 1 + \frac{x_1}{\mu}(2 - Y) + \frac{x_2}{\mu^2}(6 - 6Y + Y^2) \right), \quad (10.12)$$

$$29998 : \quad F = \frac{e^{-Y}Y}{4\mu^2} \left( 1 + \frac{x_1}{\mu}(3 - Y) + \frac{x_2}{\mu^2}(12 - 8Y + Y^2) \right), \quad (10.13)$$

$$39998 : \quad F = \frac{e^{-Y}Y^{3/2}}{4\mu^{5/2}} \left( 1 + \frac{x_1}{\mu}(4 - Y) + \frac{x_2}{\mu^2}(20 - 10Y + Y^2) \right), \quad (10.14)$$

where  $Y = q_T^2/(4\mu)$ .

## XI. TMDX.DY MODULE

This module evaluates cross-sections with the Drell-Yan-like kinematics. I.e. it expects the following kinematic input,  $(s, Q, y)$  which defines the variables  $x$ 's, etc.

The general structure of the cross-section is

$$\Delta\sigma(q_T) = \int_{bin} dX \text{ prefactor2} \times \sum_n F_n, \quad (11.1)$$

where

$$dX = dQ^2 dy dq_T^2.$$

So, for a small bin

$$\Delta\sigma(q_T) \approx \Delta X \frac{d\sigma}{dX}, \quad \Delta X = (Q_{\max}^2 - Q_{\min}^2)(y_{\max} - y_{\min})(q_{T\max}^2 - q_{T\min}^2). \quad (11.2)$$

*Prefactor2* accumulates the definition of the phase-space, and general process. It has the form

$$\text{prefactor2} = (\text{phase-space Jacobian}) \times H(Q, \mu_H) \quad (11.3)$$

The  $F_n$  is a contribution specific for a channel. Some processes have several channels, which could be computed together or separately (e.g.  $\sim f_1 f_1$  and  $\sim h_1^\perp h_1^\perp$ ). Generally it has a form

$$F_n = (\text{cuts for lepton pair})_n \times F_n(Q, q_T, x_1, x_2, \mu, \zeta_1, \zeta_2), \quad (11.4)$$

where  $F$  is a product of TMDs and corresponding weight defined in (10.1) or (14.1). There are following feature of current implementation

- In the current version the scaling variables are set as

$$\mu^2 = \zeta_1 = \zeta_2 = Q^2. \quad (11.5)$$

Currently, it is hard coded and could not be easily modified. However, there is a possibility to vary the value of  $\mu$  as  $\mu = c_2 Q$ , where  $c_2$  is variation parameter (see sec.XI E).

- For the definition of (*cuts for lepton pair*)-function see [? ]. It is evaluated within module `LeptonCutsDY.f90`. The presence of cuts, and their parameters are set by the `SetCuts` subroutine.
- The expression for the hard factor  $H$  is taken from [? ]. It is the function of  $\ln(Q/\mu_H)$  and  $a_s(\mu_H)$ . Since in the current realization  $\mu_H = Q$ , the logarithm is replaced by  $\ln(c_2)$ , where  $c_2$  is the variation constant.

The cross-section can be computed by two different means. Using LP factorization or using KPC-resummed factorization. The selection is done by the parameter `A.p2`. One behavior totally exclude another.

This section is to be updated by definition of kinematics

List of available commands



| Command   | Type     | Sec. | Short description  |
|---|----------|------|--|
| TMDX_DY_Initialize(order)                         | subrout. | XI A | Initialization of module.  |
| TMDX_DY_ShowStatistic()                           | subrout. | –    | Print current statistic on the number of calls.                      |
| TMDX_DY_ResetCounters()                           | subrout. | –    | Reset intrinsic counters of the module.                              |
| TMDX_DY_SetScaleVariation(c2)                     | subrout. | –    | Set new value for the scale-variation constant $c_2$ .               |
| xSec_DY(X,proc,s,qT,Q,y,iC,CutP,Num)              | subrout. | XI C | Evaluates cross-section completely integrated the bin.               |
| xSec_DY_List(X,proc,s,qT,Q,y,iC,CutP,Num)         | subrout. | XI C | Evaluates cross-section completely integrated the bin over the list. |
| xSec_DY_List(X,proc,s,qT,Q,y,iC,CutP,Num)         | subrout. | XI C | Evaluates cross-section completely integrated the bin over the list. |
| xSec_DY_List_BINLESS(X,proc,s,qT,Q,y,iC,CutP,Num) | subrout. | XI C | Evaluates cross-section at the point over the list.                  |

### A. Initialization

Prior the usage module is to be initialized (once per run) by

```
call TMDX_DY_Initialize(file)
```

The order of used coefficient function is set in `constant-file` by 'LO'(a<sub>s</sub><sup>0</sup>), 'NLO'(a<sub>s</sub><sup>1</sup>), 'NNLO' or 'N2LO'(a<sub>s</sub><sup>2</sup>), 'N3LO'(a<sub>s</sub><sup>3</sup>), 'N4LO'(a<sub>s</sub><sup>4</sup>). The value in brackets shows the maximum included perturbative order in the hard coefficient function.

### B. The parameters of cross-section

**Process:** The process is encoded by integer numbers (/p1,p2,p3,p4,p5,.../) where

p1 (integer) Defines the *prefactor2* that contains the phase space elements and the universal part of factorization formula.

p2 (integer) Defines the type of hadron 1.

p3 (integer) Defines the type of hadron 2.

p4 (integer) Defines the structure function  $F$ . See sec.X C.

**Note:** In **artemide v.2.07 and earlier** the process was encoded by 3 numbers, and did not contained the types of hadrons.

**Kinematics:** Each bin is defined by  $s$  (Mandelstam variable, GeV<sup>2</sup>),  $Q$  (virtuality, GeV, [min,max]),  $y$  (rapidity, [min,max]) and  $q_T$  (transverse momentum, GeV, [min,max]). If the order of limits is inverted the result is 0.

**Lepton cuts:** The fiducial cuts on the lepton pair are defined by two variables: logical `iC` (If `inc=.true.` the evaluation of cuts will be done, otherwise it will be ignored.). And the array of four reals (`p1,p2,eta1,eta2`), such that the cuts are defined as

$$|l_T| < p1, \quad |l'_T| < p2, \quad \text{eta1} < \eta, \eta', \text{eta2}, \quad (11.6)$$

where  $l$  and  $l'$  are the momenta of produced leptons, with  $l_T$ 's being their transverse components and  $\eta$ 's being their rapidities. For accurate definition of the cut-function see sec.2.6 of [?] (particularly equations (2.40)-(2.42)).

### C. Cross-section evaluation

The main subroutine is called `xSec_DY` and it have the following interface:

$$\text{xSec\_DY}(X, \text{process}, s, qT, Q, y, \text{includeCuts}, \text{CutParameters}, \text{Num})$$

where

- $X$  is (real\*8) value of cross-section.
- `process` (integer)array (p1,p2,p3,p4).
- $s$  is Mandelshtam variable  $s$
- $qT$  is (real\*8)array (qtmin,qtmax)
- $Q$  is (real\*8)array (Qmin,Qmax)
- $y$  is (real\*8)array (ymin,ymax)
- `includeCuts` is logical
- `CutParameters` is (real\*8) array (k1,k2,eta1,eta2) **OPTIONAL**
- `Num` is even integer that determine number of section in  $q_T$  integral **OPTIONAL**

Note, that optional variables could be omit during the call.

**IMPORTANT:** Practically, the call of this function coincides with  $X$  evaluated by the following set of commands  
`call TMDX_DY_SetProcess(process)`  
`call TMDX_DY_XSetup(s, any, any)`  
`call SetCuts(includeCuts,k1,k2,eta1,eta2)`  
`call CalcXsec_DY_PTint_Qint_Yint (X,qtmin,qtmax,Qmin,Qmax,yMin,yMax,Num)`

**There is also an analogous subroutine that evaluate cross-section by list.** It is called `xSec_DY_List` and it have the following interface:

$$\text{xSec\_DY\_List}(X, \text{process}, s, qT, Q, y, \text{includeCuts}, \text{CutParameters}, \text{Num})$$

All variables are analogues to those of `xSec_DY`, but should come in lists, i.e.  $X$  is (1:n), `process` is (1:n,1:4),  $s$  is (1:n),  $qT, Q, y$  are (1:n,1:2), `includeCuts` is (1:n), `CutParameters` is (1:n,1:4), and `Num` is (1:n). **Only the `Num` argument is OPTIONAL arguments. The argument `CutParameters` must be presented.** This command compiled by OPENMP, so runs in parallel on multi-core computers.

**Finally, there is an also analogous subroutine that evaluate cross-section by list without bin-integration.** It is called `xSec_DY_List_BINLESS` and it have the following interface:

$$\text{xSec\_DY\_List}(X, \text{process}, s, qT, Q, y, \text{includeCuts}, \text{CutParameters}, \text{Num})$$

All variables are analogues to those of `xSec_DY_List`, but should come in lists of central values, i.e.  $X$  is (1:n), `process` is (1:n,1:4),  $s$  is (1:n),  $qT, Q, y$  are (1:n), `includeCuts` is (1:n), `CutParameters` is (1:n,1:4), and `Num` is (1:n). **Only the `Num` argument is OPTIONAL arguments. The argument `CutParameters` must be presented.** This command compiled by OPENMP, so runs in parallel on multi-core computers.

#### Extra notes:

- The integrations over  $Q$  and  $y$  are adaptive Simpsons. We have found that it is the fastest (adaptive) method for typical cross-sections with tolerance  $10^{-3} - 10^{-4}$ . Naturally, it is not suitable for higher precision, which however is not typically required.

- The integration over  $p_T$  is not adaptive, since typically  $p_T$ -bins are rather smooth and integral is already accurate if approximated by 4-8 points (default minimal value is set in INI-file, and is automatically increased for larger bins). For unexceptionally large  $q_T$ -bins, or very rapid behavior we suggest to use overloaded versions with manual set of  $N$  (number of integral sections).
- For very small value of  $q_T$  the cross-section is frozen at  $q_{T(abs\ min)}$ , which is defined in INI-file ( $10^{-4}$  by default). It is done to avoid badly converging integrals.
- The evaluation can be done in parallel. The list commands are parallel by default. Basically, in this case, individual values for the list of cross-sections are evaluated in parallel. Unfortunately the parallel scaling-rate is not very high, on 8 processors it is about 400%-500% only. The parallelization is made with OPENMP library. To make the parallel computation possible, add `-fopenmp` option in the compilation instructions. The maximum number of allowed threads is set `constants-file`.

#### D. LeptonCutsDY

The calculation of cut prefactor is made in `LeptonCutsDY.f90`. It has two public procedures: `SetCutParameters`, and `CutFactor`.

- The subroutine `SetCutParameters(kT,eta1,eta2)` set a default version of cut parameters:  $p_{1,2} < k_T$  and  $\eta_1 < \eta < \eta_2$ .
- The overloaded version of the subroutine `SetCutParameters(k1,k2,eta1,eta2)` set a default version of asymmetric cut parameters.  $p_1 < k_1, p_2 < k_2$  and  $\eta_1 < \eta < \eta_2$ .
- Function `CutFactor(qT,Q_in,y_in, CutParameters)` calculates the cut prefactor at the point  $q_T, Q, y$ . The argument `CutParameters` is **optional**. If it is not present, cut parameters are taken from default values (which are set by `SetCutParameters`). `CutParameters` is array (/ `k1,k2,eta1,eta2`/). The usage of global definition for `CutParameters` is not recommended, since it can result into running condition during parallel computation. This interface is left for compatibility with earlier version of `artemide`.

#### E. Variation of scale

TO BE WRITTEN

#### F. Options for evaluation of DY-like cross-section

##### 1. $\pi^2$ -resummation

The coefficient function of DY-like cross-section  $|C_V|^2$  is evaluated at  $(-q^2)$  (space-like momentum). Thus, it has contributions  $\sim \pi^2$ , which could be large, especially in the case of Higgs-boson production, see discussion in refs.Ahrens:2009cxz, Ahrens:2008qu. These corrections could be resummed by RG technique [? ]. So,

$$|C_V(-q^2)|^2 \rightarrow U_\pi |C_V(q^2)|^2, \quad (11.7)$$

where  $U_\pi$  is the resummation exponent for  $\pi a_s$ -correction. At LO it reads [? ]

$$U_\pi = \exp \left( \frac{\Gamma_0}{2a_s\beta_0^2} [2a \arctan a - \ln(1+a^2)] \right), \quad a = \pi a_s. \quad (11.8)$$

To use the  $\pi$ -resummed coefficient function switch the corresponded option in `constants-file` (Option 9.A.p3).

##### 2. Power corrections

There are many sources of power corrections. For a moment there is no systematic studies of power corrections for TMD factorization. Nonetheless I include some options in `artemide`, and plan to make more systematic treatment in the future.

**Exact values of  $x_{1,2}$  for DY:** The TMD distributions enter the cross-section with  $x_1$  and  $x_2$  equal to

$$\begin{aligned} x_1 &= \frac{q^+}{p_1^+} = \frac{Q}{\sqrt{s}} e^y \sqrt{1 + \frac{\mathbf{q}_T^2}{Q^2}} \simeq \frac{Q}{\sqrt{s}} e^y, \\ x_1 &= \frac{q^+}{p_1^+} = \frac{Q}{\sqrt{s}} e^{-y} \sqrt{1 + \frac{\mathbf{q}_T^2}{Q^2}} \simeq \frac{Q}{\sqrt{s}} e^{-y}. \end{aligned} \quad (11.9)$$

Traditionally, the corrections  $\sim q_T/Q$  are dropped, since they are power corrections. Nonetheless, they could be included since they have different sources in comparison to power corrections to the factorized cross-section. Usage of one or another version of  $x_{1,2}$  is switched in `constants-file` (Option 9.A.p2).

**Exact values for hard scale for DY:** The TMD hard coefficient function contains  $\ln(|2q^+q^-|/\mu^2)$  which is usually approximated by  $\ln(Q^2)$ . Similarly, the *zeta*-scales are normalized as  $\zeta\bar{\zeta} = (2q^+q^-)^2$ , which is usually approximated by  $Q^4$ . So, the exact values of scales are

$$\mu = \sqrt{Q^2 + \mathbf{q}_T^2}, \quad \zeta_1 = \zeta_2 = Q^2 + \mathbf{q}_T^2. \quad (11.10)$$

Traditionally, the corrections  $\sim q_T/Q$  are dropped, since they are power corrections. Nonetheless, they could be included since they have different sources in comparison to power corrections to the factorized cross-section. Usage of one or another version of *scales* is switched in `constants-file` (Option 9.A.p4).

## G. Enumeration of processes (LP case)

List of enumeration for p2

| p2 | prefactor2   | Short description  |
|----|--|--|
| 0  | 1  | Test case  |
| 1  | $\frac{4\pi}{9} \frac{\alpha_{\text{em}}^2(Q)}{sQ^2}  C_V^{DY}(c_2Q) ^2 R \times \text{cut}(q_T)$  | DY-cross-section $\frac{d\sigma}{dQ^2 dy d(q_T^2)}$ . $\text{cut}(q_T)$ is the weight for the lepton tensor with fiducial cuts (see sec.2.6 in [? ]). Corresponds to DY-cross-section $\frac{d\sigma}{dQ^2 dy d(q_T^2)}$ . The result is in pb/GeV <sup>2</sup>  |
| 2  | $\frac{2\sqrt{Q^2+q_T^2}}{\sqrt{s}} \cosh y \frac{4\pi}{9} \frac{\alpha_{\text{em}}^2(Q)}{sQ^2}  C_V^{DY}(c_2Q) ^2 R \times \text{cut}(q_T)$ | DY-cross-section $\frac{d\sigma}{dQ^2 dy d(q_T^2)}$ . $\text{cut}(q_T)$ is the weight for the lepton tensor with fiducial cuts (see sec.2.6 in [? ]). The result is in pb/GeV <sup>2</sup> . <b>Attention!:</b> this process corresponds to computation with the $dX = dx_F dQ^2 dq_T^2$ where $x_F = \frac{2\sqrt{Q^2+q_T^2}}{\sqrt{s}} \sinh y$ is Feynman $x$ . In this case the integration $y$ is preplaced by the integration over $x_F$ (i.e. limits of integration are treated as for $x_F$ ). |
| 3  | $\frac{4\pi^2}{3} \frac{\alpha_{\text{em}}(Q)}{s} Br_Z  C_V^{DY}(c_2Q) ^2 R \times \text{cut}(q_T)$  | DY-cross-section $\frac{d\sigma}{dQ^2 dy d(q_T^2)}$ for the Z-boson production in the narrow width approximation. $Br_Z = 0.03645$ is Z-boson branching ration to leptons. $\text{cut}(q_T)$ is the weight for the lepton tensor with fiducial cuts (see sec.2.6 in [? ]). The result is in pb/GeV <sup>2</sup>  |
| 4  | $\frac{4\pi^2}{3} \frac{\alpha_{\text{em}}(Q)}{s} Br_W  C_V^{DY}(c_2Q) ^2 R \times \text{cut}(q_T)$  | DY-cross-section $\frac{d\sigma}{dQ^2 dy d(q_T^2)}$ for the W-boson production in the narrow width approximation. $Br_W = 0.1086$ is W-boson branching ration to leptons. $\text{cut}(q_T)$ is the weight for the lepton tensor with fiducial cuts (see sec.2.6 in [? ]). Use together with p3=13,...,18. In this case, $Q$ is be $M_Z$ . The result is in pb/GeV <sup>2</sup>   |
| 5  | $\frac{2}{\pi} \frac{\pi m_H^2 a_s^2(Q)}{36sv^2} C_t^2(m_t, Q)  C_g(Q) ^2  A(x_t) ^2$  | Cross-section $\frac{d\sigma}{dy d(q_T^2)}$ for the exclusive Higgs-boson production. The result is in pb/GeV <sup>2</sup> . For definition of functions see [? ].   |

Here  $R = 0.3893379 \cdot 10^9$  the transformation factor from GeV to pb.

## H. Enumeration of processes (KPC case)

List of enumeration for p2

| p2 | <i>prefactor2</i>   | Short description  |
|----|---|--|
| 0  | 1   | Test case  |
| 1  | $\frac{2\pi^2}{9} \frac{\alpha_{em}^2(Q)}{s}  C_V^{DY}(c_2 Q) ^2 R \times \text{cut}(q_T)$  | DY-cross-section $\frac{d\sigma}{dQ^2 dy d(q_T^2)}$ . $\text{cut}(q_T)$ is the weight for the lepton tensor with fiducial cuts (see sec.2.6 in [? ]). Corresponds to DY-cross-section $\frac{d\sigma}{dQ^2 dy d(q_T^2)}$ . The result is in pb/GeV <sup>2</sup>  |
| 2  | $\frac{2\sqrt{Q^2+q_T^2}}{\sqrt{s}} \cosh y \frac{2\pi^2}{9} \frac{\alpha_{em}^2(Q)}{s}  C_V^{DY}(c_2 Q) ^2 R \times \text{cut}(q_T)$ | DY-cross-section $\frac{d\sigma}{dQ^2 dy d(q_T^2)}$ . $\text{cut}(q_T)$ is the weight for the lepton tensor with fiducial cuts (see sec.2.6 in [? ]). The result is in pb/GeV <sup>2</sup> . <b>Attention!:</b> this process corresponds to computation with the $dX = dx_F dQ^2 dq_T^2$ where $x_F = \frac{2\sqrt{Q^2+q_T^2}}{\sqrt{s}} \sinh y$ is Feynman $x$ . In this case the integration $y$ is preplaced by the integration over $x_F$ (i.e. limits of integration are treated as for $x_F$ ). |
| 3  | —   | Absent   |
| 4  | —   | Absent   |
| 5  | $\frac{\pi m_H^2 a_s^2(Q)}{36 s v^2} Q^2 C_t^2(m_t, Q)  C_g(Q) ^2  A(x_t) ^2$   | Cross-section $\frac{d\sigma}{dy d(q_T^2)}$ for the exclusive Higgs-boson production. The result is in pb/GeV <sup>2</sup> . For definition of functions see [? ]. <b>The correct formula is to be defined, this is LP formula multiplied by large-Q translation factor.</b>   |

Here  $R = 0.3893379 \cdot 10^9$  the transformation factor from GeV to pb.

## XII. TMDX\_SIDIS MODULE

This module evaluates cross-sections with the SIDIS-like kinematics. I.e. it expects the following kinematic input,  $(Q, x, z)$  or  $(Q, y, z)$  or  $(x, y, z)$ , that are equivalent.

The general structure of the cross-section is

$$\frac{d\sigma}{dX} = d\sigma(q_T) = \int [bin] prefactor2 \times F, \quad (12.1)$$

where

$$dX = dx dQ^2 d\mathbf{p}_T^2. \quad (12.2)$$

For further specifications of parameters see sec.XII F.

List of available commands

| Command                               | Type     | Sec.  | Short description   |
|---------------------------------------|----------|-------|---|
| TMDX_SIDIS_Initialize(file,prefix)    | subrout. | XI A  | * Initialization of module.   |
| TMDX_SIDIS_ResetCounters()            | subrout. | –     | * Reset intrinsic counters of module.                                     |
| TMDX_SIDIS_ShowStatistic()            | subrout. | –     | * Print current statistic on the number of calls.                         |
| TMDX_SIDIS_SetScaleVariation(c2)      | subrout. | –     | * Set new values for the scale-variation constants.                       |
| xSec_SIDIS(X,process,s,pT,z,x,Q)      | subrout. | XII A | Evaluates cross-section completely integrated over the bin.               |
| xSec_SIDIS_List(X,process,s,pT,z,x,Q) | subrout. | XII A | Evaluates cross-section completely integrated over the bin over the list. |

\*) The structure of the module repeats the structure of TMDX\_DY module, with the main change in the kinematic definition. Most part of routines has the same input and output with only replacement `_DY`→`_SIDIS`. We do not comment such commands. They marked by \* in the following table.

### A. Cross-section evaluation

The computation of SIDIS cross-section is called by the following subroutine:

$$\text{xSec\_SIDIS}(X, \text{process}, s, pT, z, x, Q, \text{incC}, \text{cuts}, \text{masses})$$

where

- $X$  is (real\*8) value of cross-section.
- **process** is (integer)array (p1,p2,p3,p4) (described below).
- $s$  is (real\*8) Mandelstam variable  $s$
- $pT$  is (real\*8) array (qtmin,qtmax)
- $z$  is (real\*8) array (zmin,zmax)
- $x$  is (real\*8) array (xmin,xmax)
- $Q$  is (real\*8) array (Qmin,Qmax)
- **incC** is (logical) flag to include cuts.
- **cuts** is (real\*8) array (ymin,ymax,W2min,W2max) that parameterizes kinematic cuts.
- **masses** is (real\*8) array (mt,mp) which is ( $M_{target}, m_{produced}$ ) in GeV **OPTIONAL**. If not specified, considered as zeros.

**There is also analogous subroutine that evaluate cross-section by list.** It is called `xSec_SIDIS_List` and it have the following interface:

$$\text{xSec\_SIDIS\_List}(X, \text{process}, s, pT, z, x, Q, \text{incC}, \text{cuts}, \text{masses})$$

All variables are analogues to those of `xSec_SIDIS`, but should come in lists, i.e.  $X$  is (1:n), **process** is (1:n,1:4),  $s$  is (1:n),  $pT, x, Q, z$  are (1:n,1:2). This command compiled by OPENMP, so runs in parallel on multi-core computers.

In addition to these main subroutines the following sub-routines are defined

$$\text{xSec\_SIDIS\_BINLESS}(\text{proc}, s, pT, z, x, Q, m1, m2)$$

Compute cross-section without integration over bin (and also ignoring any possible bin-cut options). All kinematic variables are (real\*8).

There harpy-interface prevents using optional variables. Thus there are separate copies of the function with argument **masses** being obligatory:

$$\begin{aligned} &\text{xSec\_SIDIS\_List\_forharpy}(xx, \text{process}, s, pT, z, x, Q, \text{doCut}, \text{Cuts}, \text{masses}) \\ &\text{xSec\_SIDIS\_BINLESS\_List\_forharpy}(xx, \text{process}, s, pT, z, x, Q, \text{masses}) \end{aligned}$$

### B. Enumeration of processes

The enumeration of processes for SIDIS is analogous to the DY case. Namely, each process is defined by 4 integer numbers

$$[p1, p2, p3, p4]$$

with

- $p_1$  indicates type of *prefactor2* to use. Depending on this prefactor also the integral over phase space may differ. See the table below for details.



- $p_2$  is the number of the target hadron (i.e. the hadron associated with TMDPDF)
- $p_3$  is the number of the produced hadron (i.e. the hadron associated with TMDFF)
- $p_4$  indicates the process (structure function). See the table below for details.

List of enumeration for *prefactor2*

p1

| p1 | <i>prefactor2</i>   | Short description   |
|----|---------------------|---|
| 0  | 1                   | For tests and cross-checks.   |
| 1  | $R_C$               | The cross-section $\frac{d\sigma}{dx dQ^2 dz d(p_\perp^2)}$ . The factor $R_C$ is defined below.                                      |
| 2  | $\frac{Q^2}{y} R_C$ | The cross-section $\frac{d\sigma}{dx dy dz d(p_\perp^2)}$ , in this case integration over $Q^2$ is replaced by integration over $y$ . |
| 3  | $\frac{x}{y} R_C$   | The cross-section $\frac{d\sigma}{dy dQ^2 dz d(p_\perp^2)}$ , in this case integration over $x$ is replaced by integration over $y$ . |

The factor  $R_C$  is standard normalization of the cross-section (however without factor  $1 + \gamma^2/2x$ , which is obsolete)

$$R_C = \frac{2\pi\alpha_{\text{em}}^2(Q)}{Q^4\sqrt{1-\gamma^2\rho_\perp^2}} \frac{y^2}{1-\varepsilon} |C_V^{SIDIS}(c_2Q)|^2 c_0^{(un)} R_t \text{ [pb/GeV}^2\text{]}, \quad (12.3)$$

with  $R_t = 0.3893379 \times 10^9$  the transformation factor from GeV to pb, and

$$c_0^{(un)} = 1 + \frac{\mathbf{p}_\perp^2}{(zQ)^2} \frac{\varepsilon - \frac{\gamma^2}{2}}{1 - \gamma^2\rho^2} = 1 + \frac{\mathbf{q}_T^2}{Q^2} \frac{\varepsilon - \frac{\gamma^2}{2}}{1 + \gamma^2}.$$

Note, that different parts of this formula can be switched on/off by parameters of the power-corrections.

### C. Kinematic cuts

Some experiments put extra constraints on the measurement. Typically, such constraint has the form

$$y_{\min} < y < y_{\max}, \quad W_{\min}^2 < W^2 < W_{\max}^2. \quad (12.4)$$

In this case the integration over  $x$  and  $Q^2$  has additional restrictions. Say, if one integrates over a bin:  $x_{\min} < x < x_{\max}$  and  $Q_{\min} < Q < Q_{\max}$  the boundaries of the bin should be modified as (here integration over  $x$  is before the integration over  $Q$ )

$$\hat{x}_{\min}(Q) < x < \hat{x}_{\max}(Q), \quad \hat{Q}_{\min}^2 < Q^2 < \hat{Q}_{\max}^2. \quad (12.5)$$

where

$$\begin{aligned} \hat{x}_{\min}(Q) &= \max\left\{x_{\min}, \frac{Q^2}{y_{\max}(s - M^2)}, \frac{Q^2}{Q^2 + W_{\max}^2 - M^2}\right\}, \\ \hat{x}_{\max}(Q) &= \min\left\{x_{\max}, \frac{Q^2}{y_{\min}(s - M^2)}, \frac{Q^2}{Q^2 + W_{\min}^2 - M^2}\right\}, \\ \hat{Q}_{\min}^2 &= \max\left\{Q_{\min}^2, x_{\min}y_{\min}(s - M^2), \frac{x_{\min}}{1 - x_{\min}}(W_{\min}^2 - M^2)\right\}, \\ \hat{Q}_{\max}^2 &= \min\left\{Q_{\max}^2, x_{\max}y_{\max}(s - M^2), \frac{x_{\max}}{1 - x_{\max}}(W_{\max}^2 - M^2)\right\}. \end{aligned}$$

If some cut is not required, then set corresponding parameter to limiting value, such as  $y_{\min} = 0$ ,  $y_{\max} = 1$ ,  $W_{\min}^2 = 0$ ,  $W_{\max}^2 \gg Q^2$ .

**Important:** If due to cuts the minimal and maximum values of parameter are inverted, the resulting cross-section is immediately set to zero.

### D. Power corrections

There are many sources of power corrections. For a moment there is no systematic studies of effect of power corrections for TMD factorization. Nonetheless we include some options in `artemide`, and plan to make systematic treatment in the future.

**Kinematic corrections to the variables/phase space.** There are three sources of kinematic corrections to the variables

$$\frac{p_{\perp}}{Q}, \quad \frac{M}{Q}, \quad \frac{m}{Q},$$

where  $M$  is the target mass,  $m$  is the mass of the fragmented hadron. These terms appear in many places of the SIDIS expression (see sec.XII F, for some minimal details), even without accounting for power-suppressed terms in the factorization formula. The nice feature of these correction is that they could be accounted exactly. The majority of these terms appears during of the Lorentz transformation of factorization frame (where the factorization is performed) to the laboratory frame (where the measurement is made).

**Exact values for hard scale for SIDIS:** The TMD hard coefficient function contains  $\ln(|2q^+q^-|/\mu^2)$  which is usually approximated by  $\ln(Q^2)$ . Similarly, the *zeta*-scales are normalized as  $\zeta\bar{\zeta} = (2q^+q^-)^2$ , which is usually approximated by  $Q^4$ . So, the exact values of scales are

$$\mu = \sqrt{Q^2 - q_T^2}, \quad \zeta_1 = \zeta_2 = Q^2 - q_T^2. \quad (12.6)$$

Traditionally, the corrections  $\sim q_T/Q$  are dropped, since they are power corrections. Nonetheless, they could be included since they have different sources in comparison to power corrections to the factorized cross-section. Usage of one or another version of *scales* is switched in `constants-file` (Option 10.A.p6).

The accounting of these corrections is switched on/off in `constants`.

### E. Bin-integration routines

The inclusion of bin-integration is essential for comparison with experiment. The integration over a bin is automatically included in the calculation of cross-section (although routines for unintegrated calculation are also presented). Typically, one does not need an extreme precision from such integration i.e. 4-5 digit precision is typically more than enough to reach the experimental precision. Also the cross-section is relatively slow-function (exception is Z-boson peak). Therefore, as a based method for bin-integration I have selected the adaptive Simpson integration rule. By default, the adaptive Simpson rule is used for integration over  $z$ ,  $x$ ,  $Q^2$ , and fixed-number-of-points Simpson method for  $p_T$ -integration. Note, one can switch to G7 method (faster but not precise) by modification of the pre-compile parameter in the beginning of the code.

The integration is made for phase-space element  $dQ^2 dx d\mathbf{p}_{\perp}^2$ . I.e.

$$\Delta\sigma = \int dz \int dQ^2 dx d\mathbf{p}_{\perp}^2 d\sigma = \int_{z_{\min}}^{z_{\max}} dz \int_{Q_{\min}}^{Q_{\max}} 2Q dQ \int_{x_{\min}}^{x_{\max}} dx \int_{p_{\min}}^{p_{\max}} 2|\mathbf{p}_{\perp}| d|\mathbf{p}_{\perp}| d\sigma. \quad (12.7)$$

For processes with  $dx \rightarrow dy$ , or  $dQ^2 \rightarrow dy$  the integration is done in the same manner but with limits of integration over  $Q$  or  $x$  re-computed at values of  $y$ .

#### Notes:

- If the limits of integration are inverted (i.e.  $\int_a^b$  with  $a > b$ ) the integral is set to zero. It is made intentionally, because the cuts may shift the limits upside-down, which corresponds to the completely-eliminated case.
- The central value is computed at the mean of the bin. I.e. for bin  $[a, b]$  the central value is computed at  $(a+b)/2$ .

### F. SIDIS theory

In many aspects the theory for SIDIS is more complicated than for Drell-Yan. Here I collect the main definition which were used in the `artemide`. For a more detailed description see ref.[? ]. Note, that some parts of definition were derived by me, since I have not found them in the literature.

### 1. Kinematics

The process is

$$H(P) + l(l) \rightarrow l(l') + h(p_h) + X, \quad (12.8)$$

with

$$P^2 = M^2, l^2 = l'^2 = m_l^2 \simeq 0, \quad p_h^2 = m^2. \quad (12.9)$$

The standard variables used for the description of SIDIS are

$$q = l - l' \quad \Rightarrow \quad Q^2 = -q^2, \quad x = \frac{Q^2}{2(Pq)}, \quad y = \frac{(Pq)}{(Pl)}, \quad z = \frac{(Pp_h)}{(Pq)}. \quad (12.10)$$

There are also a set of power variables used to define power-corrections

$$\gamma = \frac{2Mx}{Q}, \quad \rho = \frac{m}{zQ}, \quad \rho_\perp^2 = \frac{m^2 + \mathbf{p}_\perp^2}{z^2 Q^2}. \quad (12.11)$$

The variables  $x$ ,  $y$  and  $Q^2$  are dependent:  $xy(s - M^2) = Q^2$ . The phase-volume  $dxdQ^2$  can be expressed via  $dxdy$  or  $dydQ^2$ :

$$dxdQ^2 = \frac{Q^2}{y} dxdy = \frac{x}{y} dydQ^2. \quad (12.12)$$

A phase space point is totally characterized by the following numbers:

$$\{s, x, Q^2, z, \mathbf{p}_T^2\}. \quad (12.13)$$

Alternatively, one can replace  $x \leftrightarrow Q^2 \leftrightarrow y$ .

### 2. Cross-section

The cross-section for SIDIS has the general form

$$\frac{d\sigma}{dxdQ^2 dz d\psi d\phi_h d\mathbf{p}_{h\perp}^2} = \frac{\alpha_{\text{em}}^2(Q)}{Q^6} \frac{y^2}{8z} \frac{L_{\mu\nu} W^{\mu\nu}}{\sqrt{1 + \rho_\perp^2 \gamma^2}}, \quad (12.14)$$

where  $L_{\mu\nu}$  is the lepton tensor, and  $W_{\mu\nu}$  is the hadron tensor. The hadron tensor contains many term accompanied by polarization angles.

The TMD factorization is performed in the factorization frame with respect to  $\mathbf{q}_T^2 \ll Q^2$ . The factorization variable  $|\mathbf{q}_T|$  is related to the measured variable  $\mathbf{p}_{h\perp}$  as

$$|\mathbf{q}_T| = \frac{|\mathbf{p}_{h\perp}|}{z} \sqrt{\frac{1 + \gamma^2}{1 - \gamma^2 \rho^2}}, \quad \Leftrightarrow \quad |\mathbf{p}_{h\perp}| = z |\mathbf{q}_T| \sqrt{\frac{1 - \gamma^2 \rho^2}{1 + \gamma^2}}. \quad (12.15)$$

In the factorization frame, the unpolarized part of the hadron tensor

$$W^{\mu\nu} = \frac{-z_1 g_T^{\mu\nu}}{\pi} \int |\mathbf{b}| d|\mathbf{b}| J_0(|\mathbf{b}| |\mathbf{q}_T|) \sum_f e_f^2 |C_V(-Q^2, \mu^2)|^2 F_1^f(x_1, \mathbf{b}) D_1^f(z_1, \mathbf{b}) + \dots, \quad (12.16)$$

where dots denote, polarized terms, and power corrections. The LP variables  $x_1$  and  $z_1$  are

$$x_1 = -\frac{2x}{\gamma^2} \left( 1 - \sqrt{1 + \gamma^2 \left( 1 - \frac{\mathbf{q}_T^2}{Q^2} \right)} \right), \quad (12.17)$$

$$z_1 = -z \frac{1 - \sqrt{1 + \left( 1 - \frac{\mathbf{q}_T^2}{Q^2} \right) \gamma^2}}{\gamma^2} \frac{1 + \sqrt{1 - \rho^2 \gamma^2}}{1 - \frac{\mathbf{q}_T^2}{Q^2}}. \quad (12.18)$$

Making the convolution with unpolarized leptonic tensor, we get the following expression for the cross-section

$$\begin{aligned} \frac{d\sigma}{dx dQ^2 dz d\mathbf{p}_{h\perp}^2} &= 2\pi \frac{\alpha_{\text{em}}^2}{Q^4} \frac{1}{\sqrt{1+\boldsymbol{\rho}_\perp^2 \gamma^2}} \frac{y^2}{1-\varepsilon} \frac{z_1}{z} \left\{ 1 + \left( \varepsilon - \frac{\gamma^2}{2} \right) \frac{\boldsymbol{\rho}_\perp^2 - \rho^2}{1-\gamma^2 \rho^2} \right\} \\ &\quad \int \frac{|\mathbf{b}| d|\mathbf{b}|}{2} J_0 \left( \frac{|\mathbf{b}| |\mathbf{p}_{h\perp}|}{z} \sqrt{\frac{1+\gamma^2}{1-\gamma^2 \rho^2}} \right) \sum_f e_f^2 |C_V(-\overline{Q}^2, \mu^2)|^2 F_1^f(x_1, \mathbf{b}) D_1^f(z_1, \mathbf{b}), \end{aligned} \quad (12.19)$$

where

$$\varepsilon = \frac{1 - y - \frac{\gamma^2 y^2}{4}}{1 - y + \frac{y^2}{2} + \frac{y^2 \gamma^2}{4}}. \quad (12.20)$$

The value  $\overline{Q}^2 = -2q^+ q^-$ , note that  $\zeta \bar{\zeta} = \overline{Q}^4$ .

Clearly, this LP expression contains a set of power corrections. It is a standard problem of TMD factorization, where the limit of LP is not very standardly defined. Thus, there are options to switch off/on these corrections.

- **Transverse momentum corrections:** (option  $p_1$  in section 10.C) Include terms proportional to  $\boldsymbol{\rho}_\perp^2$ .
- **Target mass corrections:** (option  $p_2$  in section 10.C) Include terms proportional to  $\gamma^2$ .
- **Product mass corrections:** (option  $p_3$  in section 10.C) Include terms proportional to  $\rho^2$ .
- **Exact LP values of  $x_1$  and  $z_1$ :** (option  $p_4$  in section 10.C) Uses values of  $x_1$  and  $z_1$  (if option is False  $x_1 \rightarrow x$ ,  $z_1 \rightarrow z$ )
- **Exact LP value for factorization scale:** (option  $p_5$  in section 10.C) Uses exact LP value for  $\overline{Q}^2 = -2q^+ q^- = Q^2 - \mathbf{q}_T^2$  (if option is False  $\overline{Q}^2 = Q^2$ )

Note, that corrections are included in the order from top to bottom. For example, terms  $\sim \gamma^2 \rho^2 = 0$  if  $p_2 = \text{F}$  and  $p_3 = \text{T}$ .

### XIII. TMDs\_inKT MODULE

The module `TMDs_inKT` is derivative of the module `TMDs` that provides the TMD in the transverse momentum space. We define

$$F(x, \mathbf{k}_T; \mu, \zeta) = \int \frac{d^2 \mathbf{b}}{(2\pi)^2} F(x, \mathbf{b}; \mu, \zeta) e^{-i(\mathbf{k}_T \mathbf{b})}. \quad (13.1)$$

Since all evaluated TMDs depends only on the modulus of  $\mathbf{b}$ , within the module we evaluate Hankel transformations. These transformations depends on the kind of TMD. Generally the integrals are

$$F(x, |\mathbf{k}_T|; \mu, \zeta) = \int \frac{d|\mathbf{b}|}{2\pi} |\mathbf{b}|^{n+1} J_n(|\mathbf{b}||\mathbf{k}_T|) F(x, |\mathbf{b}|; \mu, \zeta). \quad (13.2)$$

On top of it there is could be extra prefactors specified in the table. For simplicity these cases indicated by numbers

$$n = 0, \quad F(x, |\mathbf{k}_T|; \mu, \zeta) = \int \frac{d|\mathbf{b}|}{2\pi} |\mathbf{b}| J_0(|\mathbf{b}||\mathbf{k}_T|) F(x, |\mathbf{b}|; \mu, \zeta), \quad (13.3)$$

$$n = 1, \quad F(x, |\mathbf{k}_T|; \mu, \zeta) = \int \frac{d|\mathbf{b}|}{2\pi} |\mathbf{b}|^2 J_1(|\mathbf{b}||\mathbf{k}_T|) F(x, |\mathbf{b}|; \mu, \zeta), \quad (13.4)$$

$$n = 2, \quad F(x, |\mathbf{k}_T|; \mu, \zeta) = - \int \frac{d|\mathbf{b}|}{2\pi} |\mathbf{b}|^3 J_2(|\mathbf{b}||\mathbf{k}_T|) F(x, |\mathbf{b}|; \mu, \zeta). \quad (13.5)$$

The module `TMDs_inKT` uses all functions from the module `TMDs`. In fact, all technical commands (like `Initialize`) just transfer the request to `TMDs`. Thus, the information on these commands can be found in the section IX. For the evaluation of Hankel integral we use the Ogata quadrature see X B (the parameters for it are set in corresponding section of `constants`).

List of available commands

| Command                                    | Type           | Sec. | Short description   |
|--|----------------|------|---|
| <code>TMDs_inKT_Initialize(order)</code>   | subrout.       | IX   | Initialization of module.   |
| <code>TMDs_inKT_ResetCounters()</code>     | subrout.       | -    | Reset intrinsic counters of modules                                     |
| <code>TMDs_inKT_ShowStatistic</code>       | subrout.       | -    | Print current statistic on the number of calls.                         |
| <code>testTMDPDF_kT(x,kT)</code>           | (real*8(-5:5)) | -    | Evaluates the Fourier of the text function (see below). $n = 0$         |
| <code>uTMDPDF_kT_5(x,kT,h)</code>          | (real*8(-5:5)) | IX B | Unpolarized TMD PDF at the optimal line (gluon term undefined). $n = 0$ |
| <code>uTMDPDF_kT_50(x,kT,h)</code>         | (real*8(-5:5)) | IX B | Unpolarized TMD PDF at the optimal line (gluon term defined). $n = 0$   |
| <code>uTMDPDF_kT_5(x,kT,mu,zeta,h)</code>  | (real*8(-5:5)) | IX B | Unpolarized TMD PDF (gluon term undefined). $n = 0$                     |
| <code>uTMDPDF_kT_50(x,kT,mu,zeta,h)</code> | (real*8(-5:5)) | IX B | Unpolarized TMD PDF (gluon term defined). $n = 0$                       |
| <code>uTMDFF_kT_5(x,kT,h)</code>           | (real*8(-5:5)) | IX B | Unpolarized TMD FF at the optimal line (gluon term undefined). $n = 0$  |
| <code>uTMDFF_kT_50(x,kT,h)</code>          | (real*8(-5:5)) | IX B | Unpolarized TMD FF at the optimal line (gluon term defined). $n = 0$    |

|   |                |      |   |
|---|----------------|------|---|
| <code>uTMDFF_kT_5(x,kT,mu,zeta,h)</code>        | (real*8(-5:5)) | IX B | Unpolarized TMD FF (gluon term undefined). $n = 0$  |
| <code>uTMDFF_kT_50(x,kT,mu,zeta,h)</code>       | (real*8(-5:5)) | IX B | Unpolarized TMD FF (gluon term defined). $n = 0$  |
| <code>lpTMDPDF_kT_50(x,kT,h)</code>             | (real*8(-5:5)) | IX B | Linearly polarized gluon TMD PDF at the optimal line (quark terms undefined). $n = 2$   |
| <code>lpTMDPDF_kT_50(x,kT,mu,zeta,h)</code>     | (real*8(-5:5)) | IX B | Linearly polarized gluon TMD PDF (quark terms undefined). $n = 2$   |
| <code>SiversTMDPDF_kT_5(x,kT,h)</code>          | (real*8(-5:5)) | IX B | Sivers TMD PDF at the optimal line (gluon term undefined). $n = 1$  |
| <code>SiversTMDPDF_kT_50(x,kT,h)</code>         | (real*8(-5:5)) | IX B | Sivers TMD PDF at the optimal line (gluon term defined). $n = 1$  |
| <code>SiversTMDPDF_kT_5(x,kT,mu,zeta,h)</code>  | (real*8(-5:5)) | IX B | Sivers TMD PDF (gluon term undefined). $n = 1$  |
| <code>SiversTMDPDF_kT_50(x,kT,mu,zeta,h)</code> | (real*8(-5:5)) | IX B | Sivers TMD PDF (gluon term defined). $n = 1$  |
| <code>Moment_G(n,x,mu,F,h)</code>               | (real*8(-5:5)) | IX B | Computed the moment $\mathcal{G}_n$ (see below). (int) <b>n,h</b> , are order of moment and hadonr, (real*8) <b>x,mu</b> collinear momentum fraction and scale. F is the name of optimal TMDPDF. <b>For a moment, <math>n = 0, 1, 2</math>, only.</b> |
| <code>Moment_X(n,x,mu,F,h)</code>               | (real*8(-5:5)) | IX B | Computed the moment $\mathcal{G}_n$ (see below). (int) <b>n,h</b> , are order of moment and hadonr, (real*8) <b>x,mu</b> collinear momentum fraction and scale. F is the name of optimal TMDPDF. <b>For a moment, <math>n = 0, 1</math>, only.</b>    |

#### Comments:

- The value of  $k_T$  is expected bigger 1MeV for smaller values the function evaluates at 1MeV.
- For gluonless TMDs (\_5) the gluon term is identically 0.
- The convergence of the integral is checked by convergence of each flavor (until all flavors are convergent).

The function `testTMDPDF_kT(x,kT)` returns the Fourier of the following test functions ( $x$  is the free parameter

$$\{b^3 e^{-xb}, b^2 e^{-xb}, b e^{-xb}, e^{-xb}, \frac{1}{(b^2 + x^2)^4}, \frac{1}{(b^2 + x^2)^2}, \frac{1}{b^2 + x^2}, b^2 e^{-xb^2}, e^{-xb^2}, \frac{J_0(x/b)}{b}, \frac{1}{b}\}. \quad (13.6)$$

The result of evaluation should be

$$\left\{ \begin{aligned} & \frac{3}{2\pi} \frac{3k_T^4 - 24k_T^2 x^2 + 8x^4}{(k_T^2 + x^2)^{9/2}}, \frac{3x}{2\pi} \frac{2x^2 - 3k_T^2}{(k_T^2 + x^2)^{7/2}}, \frac{1}{2\pi} \frac{2x^2 - k_T^2}{(k_T^2 + x^2)^{5/2}}, \frac{1}{2\pi} \frac{x}{(k_T^2 + x^2)^{3/2}}, \\ & \frac{k_T^3}{96\pi x^3} K_3(k_T x), \frac{k_T}{4\pi x} K_1(k_T x), \frac{K_0(k_T x)}{2\pi}, \\ & \frac{4x - k_T^2}{16\pi x^3} e^{-k_T^2/(4x)}, \frac{e^{-k_T^2/(4x)}}{4\pi x}, \frac{J_0(2\sqrt{xk_T})}{2\pi k_T}, \frac{1}{2\pi k_T} \end{aligned} \right\}. \quad (13.7)$$

The expressions for functions are selected such that the integral gives reasonable values for  $0.05 < x < 1$ .

Moments are defined as

$$\mathcal{G}_n[F] = \frac{1}{2^n} \int d^2 k_T \theta(|k_T| < \mu) \frac{k_T^{2n}}{M^{2n}} F(x, k_T) = \frac{\mu^{n+1}}{2^n n!} \int_0^\infty db b^n J_{n+1}(b\mu) \tilde{F}(x, b), \quad (13.8)$$

$$\begin{aligned} \mathcal{X}_n[F] &= \int^\mu d\mathbf{k}_T^2 \frac{(\mathbf{k}_T^2)^{n+1}}{2^n M^{2n}} F(x, k_T) \\ &= \frac{1}{2^n n!} \int_0^\infty db \mu^3 (\mu b)^n \frac{(n+1)J_{n+1}(\mu b) - J_{n+3}(\mu b)}{n+2} \tilde{F}(x, b). \end{aligned} \quad (13.9)$$

### A. Grids

The computation of KPC intensively uses the momentum space TMDs. Therefore, it is better to grid them, rather than evaluate each time. The grid is 3D, in  $(x, k_T, Q)$ . The grid used in the artemide 3.0 is:

**Arcosh grid with control parameter**

$$i = 0, \dots, N_x, \quad x_i = \frac{1}{\cosh^p(\Delta_x(i - N_x))}, \quad \Delta_x = \frac{\text{arccosh}(1/x_{\min}^{1/p})}{N_x}, \quad \text{inv}[x] = N_x - \frac{\text{arccosh}(1/x^{1/p})}{\Delta_x} \quad (13.10)$$

This grid is alike logarithmic at small- $x$ , but has extra points at  $x \rightarrow 1$ . Therefore, one does not need extra gridding at large- $x$ . The parameter  $p$  gives extra flexibility. At  $p_x > 1$  the density of points at  $x \rightarrow 1$  grows. Same as used in OPE part.

**Exponential grid with  $Q_{\min}$  and  $Q_{\max}$**

$$i = 0, \dots, N_Q, \quad Q_i = Q_{\min} e^{i\Delta_Q}, \quad \Delta_Q = \frac{\ln(Q_{\max}/Q_{\min})}{N_Q}, \quad \text{inv}[Q] = \frac{\ln(Q/Q_{\min})}{\Delta_Q}. \quad (13.11)$$

Usual logarithmic grid. The function is rather smooth in the  $Q$  direction.

**Exponential grid with  $k_{\min}$  and  $k_{\max}$**

$$i = 0, \dots, N_k, \quad k_i = k_{\min} e^{i\Delta_k}, \quad \Delta_k = \frac{\ln(k_{\max}/k_{\min})}{N_k}, \quad \text{inv}[k] = \frac{\ln(k/k_{\min})}{\Delta_k}. \quad (13.12)$$

Usual logarithmic grid. The function is rather smooth in the  $k$  direction once multiplied by  $k^2$ . The maximum value of  $k$  is defined as

$$k_{\max} = p_k Q_{\max}.$$

The parameter  $p_k$  must be bigger than  $\sqrt{2}$  ( $\sqrt{2}$  is reached by the utmost point in the integration for KPC). For values  $k > p_k Q$  (at given  $Q$  the expressions are not stored).

The interpolation is cubic. The grid is build for the finite domain of  $x \in (x_{\min}, 1)$  and  $k \in (k_{\min}, k_{\max})$  and  $Q \in (Q_{\min}, Q_{\max})$ .

- For  $x < x_{\min}$ ,  $k_T > k_{\max}(Q)$ ,  $Q < Q_{\min}$  and  $Q > Q_{\max}$  the program will be terminated (with an error).
- For  $k_T < k_{\min}$  the code returns the value at  $k_{\min}$ .

#### XIV. KPC IN TMD FACTORIZATION

The TMD factorization with KPCs is computed using different route. The primary problem is that one must compute these correction in the momentum space. It essentially complicates the computation, since the evolution is done in the position space.

The module `TMDfint_KPC_DY` realizes the computation of the following integral

$$\mathcal{C}_{KPC}[K, FF] = 4p_1^+ p_2^+ \int d\xi_1 d\xi_2 \int d^4 k_1 d^4 k_2 \delta^{(4)}(q - k_1 - k_2) \delta(k_1^2) \delta(k_2^2) \quad (14.1)$$

$$\delta(k_1^+ - \xi_1 p_1^+) \delta(k_2^- - \xi_2 p_2^-) K \times FF, \quad (14.2)$$

where  $K$  is a kinematic kernel, that is resulted from the convolution of tensors, and  $FF$  is the product of TMD distributions.

This function has the following interface

$$\text{KPC\_DYconv}(\mathbf{Q}, \mathbf{qT}, \mathbf{x1}, \mathbf{x2}, \mathbf{mu}, \mathbf{proc1}, \mathbf{proc2})$$

where  $\mathbf{Q}, \mathbf{qT}, \mathbf{x1}, \mathbf{x2}, \mathbf{mu}$  are usual variables, and  $\mathbf{proc1}$  is (int,int,int)-array with first item being the number of  $FF$ , the second and third items being number of hadrons;  $\mathbf{proc2}$  is the enumeration of factors  $K$ .

##### A. Realization of convolution

The integral for convolution can be presented as

$$\mathcal{PC}_{KPC}[K, FF] = \mathcal{P} \int_0^{\pi/2} d\alpha \int_0^\pi d\theta \frac{K \times FF}{2}, \quad (14.3)$$

where

$$\xi_1 = \frac{x_1}{2} (1 + S + \sqrt{\Lambda}), \quad (14.4)$$

$$\xi_2 = \frac{x_2}{2} (1 - S + \sqrt{\Lambda}), \quad (14.5)$$

$$\mathbf{k}_1^2 = \frac{\tau^2}{4} (1 - \Lambda + 2S + S^2), \quad (14.6)$$

$$\mathbf{k}_2^2 = \frac{\tau^2}{4} (1 - \Lambda - 2S + S^2), \quad (14.7)$$

with

$$S = \frac{(\mathbf{q}_T \Delta)}{\tau^2} = \sqrt{\sin \alpha} \frac{\mathbf{q}_T^2}{\tau^2} \cos \theta, \quad \Lambda = \frac{\lambda(\tau^2, \mathbf{k}_1^2, \mathbf{k}_2^2)}{\tau^4} = \left(1 - \frac{\mathbf{q}_T^2}{\tau^2}\right) (1 - \sin \alpha). \quad (14.8)$$

The factor  $\mathcal{P}$  is the fiducial-cut factor (multiplied within the `TMDX_DY` part).

##### B. Enumeration of factors processes

| N | $K$           | $FF$  | $\mathcal{P}$   | Short description                      |
|---|---------------|---|-----------------|--|
| 1 | 1             | $e_q^2 f_1 f_1$                                     | $\mathcal{P}_0$ | Unpolarized DY                         |
| 2 | 1 ( $= K_1$ ) | $\Delta^{GG'} z_{+\ell}^{GG'} z_{+f}^{GG'} f_1 f_1$ | $\mathcal{P}_0$ | Z-boson main term $\sim \mathcal{P}_0$ |



|     |   |   |                 |  |
|-----|---|---|-----------------|--|
| 20  | $1 - \frac{\tau^2 \Lambda}{Q^2}$  | $\Delta^{GG'} z_{+\ell}^{GG'} z_{+f}^{GG'} f_1 f_1$                               | $\mathcal{P}_0$ | $f_1 f_1$ part of angular coefficient $A_0$ . Also the part of Z-boson production $\sim \mathcal{P}_0$ .                                 |
| 21  | $\frac{\tau^2 S \sqrt{\Lambda}}{Q q_T}$   | $\Delta^{GG'} z_{+\ell}^{GG'} z_{+f}^{GG'} f_1 f_1$                               | $\mathcal{P}_1$ | $f_1 f_1$ part of angular coefficient $A_1$ . Also the part of Z-boson production $\sim \mathcal{P}_1$ .                                 |
| 22  | $\frac{Q^4 + \Lambda \tau^4 + Q^2 \tau^2 (-1 + 2S^2 - \Lambda)}{Q^2 q_T^2}$   | $\Delta^{GG'} z_{+\ell}^{GG'} z_{+f}^{GG'} f_1 f_1$                               | $\mathcal{P}_2$ | $f_1 f_1$ part of angular coefficient $A_2$ . Also the part of Z-boson production $\sim \mathcal{P}_2$ .                                 |
| 23  | $-2S \frac{\tau}{q_T}$  | $\Delta^{GG'} z_{-\ell}^{GG'} z_{-f}^{GG'} \{f_1 f_1\}_A$                         | $\mathcal{P}_3$ | $f_1 f_1$ part of angular coefficient $A_3$ . Also the part of Z-boson production $\sim \mathcal{P}_3$ .                                 |
| 24  | $-2\sqrt{\Lambda} \frac{\tau}{Q}$   | $\Delta^{GG'} z_{-\ell}^{GG'} z_{-f}^{GG'} \{f_1 f_1\}_A$                         | $\mathcal{P}_4$ | $f_1 f_1$ part of angular coefficient $A_4$ . Also the part of Z-boson production $\sim \mathcal{P}_4$ .                                 |
| 29  | $\frac{3}{2} - \frac{\tau^2 \Lambda}{2Q^2}$   | $\Delta^{GG'} z_{+\ell}^{GG'} z_{+f}^{GG'} f_1 f_1$                               | 1               | $f_1 f_1$ part of proportional to 1 in the cross-section ( $\Sigma_{UU} + \Sigma_0/2$ ). Used for the normalization of some asymmetries. |
| 30  | $-\frac{\tau^2}{4M^2 Q^2} (Q^2(-1 + S^2 + \Lambda) + (1 + S^2 - \Lambda)\Lambda\tau^2)$   | $\Delta^{GG'} z_{+\ell}^{GG'} r_{+f}^{GG'} h_1^\perp h_1^\perp$                   | $\mathcal{P}_0$ | $h_1 h_1$ part of angular coefficient $A_0$ . Also the part of Z-boson production $\sim \mathcal{P}_0$ .                                 |
| 31  | $\frac{\tau^2 S \sqrt{\Lambda}}{4M^2 Q q_T} (2Q^2 + (-1 + S^2 - \Lambda)\tau^2)$  | $\Delta^{GG'} z_{+\ell}^{GG'} r_{+f}^{GG'} h_1^\perp h_1^\perp$                   | $\mathcal{P}_1$ | $h_1 h_1$ part of angular coefficient $A_1$ . Also the part of Z-boson production $\sim \mathcal{P}_1$ .                                 |
| 32  | $\frac{\tau^2}{4M^2 Q^2 q_T^2} (Q^4(-1 + S^2 + \Lambda) + \tau^4 \Lambda(1 + S^2 - \Lambda) + Q^2 \tau^2 (2S^4 + (\Lambda - 1)^2 - 3S^2(1 + \Lambda)))$ | $\Delta^{GG'} z_{+\ell}^{GG'} r_{+f}^{GG'} h_1^\perp h_1^\perp$                   | $\mathcal{P}_2$ | $h_1 h_1$ part of angular coefficient $A_2$ . Also the part of Z-boson production $\sim \mathcal{P}_2$ .                                 |
| 35  | $-\frac{\tau^3 \sqrt{\Lambda}}{4M^2 Q q_T^2} (Q^2(1 + S^2 - \Lambda) + \tau^2(-1 + S^2 + \Lambda))$   | $i\Delta^{GG'} z_{+\ell}^{GG'} r_{-f}^{GG'} \{h_1^\perp h_1^\perp\}_A$            | $\mathcal{P}_5$ | $h_1 h_1$ part of angular coefficient $A_5$ . Also the part of Z-boson production $\sim \mathcal{P}_5$ .                                 |
| 36  | $-\frac{\tau^3 S}{4M^2 Q^2 q_T} (Q^2(-1 + S^2 - \Lambda) + 2\Lambda\tau^2)$   | $i\Delta^{GG'} z_{+\ell}^{GG'} r_{-f}^{GG'} \{h_1^\perp h_1^\perp\}_A (= F_{11})$ | $\mathcal{P}_6$ | $h_1 h_1$ part of angular coefficient $A_6$ . Also the part of Z-boson production $\sim \mathcal{P}_6$ .                                 |
| 101 | 1   | $R_{ff'}^{Cu} e_f e_{f'}$   | $\mathcal{P}_0$ | Unpolarized DY on $Cu$   |
| 102 | 1   | $R_{ff'}^{2H} e_f e_{f'}$   | $\mathcal{P}_0$ | Unpolarized DY on $^2H$ ( $Z = 1$ and $A = 2$ )  |
| 103 | 1   | $R_{ff'}^W e_f e_{f'}$  | $\mathcal{P}_0$ | Unpolarized DY on $W$ ( $Z = 74$ and $A = 183$ )   |
| 200 | $(X_{20} + X_{30})/X_2$   | —   | 1               | Angular coefficient $A_0$  |

|   |   |   |   |                               |
|---|---|---|---|-------------------------------|
| 201   | $(X_{21} + X_{31})/X_2$                               | – | 1 | Angular coefficient $A_1$     |
| 202   | $(X_{22} + X_{32})/X_2$                               | – | 1 | Angular coefficient $A_2$     |
| 203   | $X_{23}/X_2$  | – | 1 | Angular coefficient $A_3$     |
| 204   | $X_{24}/X_2$  | – | 1 | Angular coefficient $A_4$     |
| 205   | $X_{35}/X_2$  | – | 1 | Angular coefficient $A_5$     |
| 206   | $X_{36}/X_2$  | – | 1 | Angular coefficient $A_6$     |
| 210   | $1 - 2(X_{20} + X_{30})/(X_{29} + \frac{1}{2}X_{30})$ | – | 1 | Angular coefficient $\lambda$ |
| 211   | $(X_{21} + X_{31})/(X_{29} + \frac{1}{2}X_{30})$      | – | 1 | Angular coefficient $\mu$     |
| 212   | $(X_{22} + X_{32})/(X_{29} + \frac{1}{2}X_{30})$      | – | 1 | Angular coefficient $\nu$     |
| * The processes 200-299 are reserved for ratios. No common prefactors are included. |   |   |   |                               |
| * $X_n$ denotes the cross-section for corresponding N.                              |   |   |   |                               |

## XV. APPENDICES

### A. Equivalence between process-enumeration in ver.< 3

In the versions of artemide < 3, the enumeration of processes was done with 3 integers  $[\tilde{a}, \tilde{b}, \tilde{c}]$ , where  $\tilde{a}$  and  $\tilde{b}$  determined the prefactor, and  $\tilde{c}$  determined the actual process. In fact, first number was not used (extremely rarely), thus it was eliminated in version > 3. Instead, the numeration of hadrons was added. In artemide > 3 the process is enumerated by four integers  $[b, c, h_1, h_2]$  where  $b$  determines the prefactor,  $c$  the type of process and  $h_1$  and  $h_2$  the hadrons for scattering.

The correspondence of process numbering is the following

- $\tilde{a} = 1$ , other options absent
- $\tilde{b} = b$
- $\tilde{c}$  is replaced by the  $h_1, h_2, c$  according to the following table

List of available commands

| artemide < 3 | artemide > 3    | Process                          | Comment            |
|--------------|-----------------|----------------------------------|--------------------|
| $\tilde{c}$  | $[h_1, h_2, c]$ |                                  |                    |
| 0            | $[-,-,0]$       |                                  | Does not use TMDs. |
| 9999         | $[-,-,0]$       | Test cases                       | Do not use TMDs.   |
| 9998         | $[-,-,0]$       |                                  | Does not use TMDs. |
| 1            | $[1,1,1]$       | $p + p \rightarrow \gamma$       |                    |
| 2            | $[1,-1,1]$      | $p + \bar{p} \rightarrow \gamma$ |                    |

|      |           |                                      |                                |
|------|-----------|--------------------------------------|--------------------------------|
| 3    | [1,1,2]   | $p + p \rightarrow Z$                |                                |
| 4    | [1,-1,2]  | $p + \bar{p} \rightarrow Z$          |                                |
| 5    | [1,1,3]   | $p + p \rightarrow Z + \gamma$       |                                |
| 6    | [1,-1,3]  | $p + \bar{p} \rightarrow Z + \gamma$ |                                |
| 7    | [1,1,4]   | $p + p \rightarrow W^+$              |                                |
| 8    | [1,1,5]   | $p + p \rightarrow W^-$              |                                |
| 9    | [1,1,6]   | $p + p \rightarrow W^\pm$            |                                |
| 10   | [1,-1,4]  | $p + \bar{p} \rightarrow W^+$        |                                |
| 11   | [1,-1,5]  | $p + \bar{p} \rightarrow W^-$        |                                |
| 12   | [1,-1,6]  | $p + \bar{p} \rightarrow W^\pm$      |                                |
| 13   | [1,1,7]   | $p + p \rightarrow W^+$              | Same as 7 with $\Gamma_W = 0$  |
| 14   | [1,1,8]   | $p + p \rightarrow W^-$              | Same as 8 with $\Gamma_W = 0$  |
| 15   | [1,1,9]   | $p + p \rightarrow W^\pm$            | Same as 9 with $\Gamma_W = 0$  |
| 16   | [1,-1,7]  | $p + \bar{p} \rightarrow W^+$        | Same as 9 with $\Gamma_W = 0$  |
| 17   | [1,-1,8]  | $p + \bar{p} \rightarrow W^-$        | Same as 10 with $\Gamma_W = 0$ |
| 18   | [1,-1,9]  | $p + \bar{p} \rightarrow W^\pm$      | Same as 11 with $\Gamma_W = 0$ |
| 20   | [1,1,10]  | $p + p \rightarrow H$                |                                |
| 21   | [1,1,11]  | $p + p \rightarrow H$                | Only $f_1$                     |
| 22   | [1,1,12]  | $p + p \rightarrow H$                | Only $f_1^\perp$               |
| 101  | [2,1,1]   | $\pi^- + p \rightarrow \gamma$       |                                |
| 102  | [2,-1,1]  | $\pi^- + \bar{p} \rightarrow \gamma$ |                                |
| 103  | [-2,1,1]  | $\pi^+ + p \rightarrow \gamma$       |                                |
| 104  | [-2,-1,1] | $\pi^+ + \bar{p} \rightarrow \gamma$ |                                |
| 1001 | [1,1,101] | $p + Cu \rightarrow \gamma$          |                                |
| 1002 | [1,1,102] | $p + {}^2H \rightarrow \gamma$       |                                |

|              |               |   |         |
|--------------|---------------|---|---------|
| 1003         | [-1,1,103]    | $\bar{p} + W \rightarrow \gamma$              |         |
| 1004         | [2,1,103]     | $\pi^- + W \rightarrow \gamma$                |         |
| polarized DY |               |   |         |
| 10001        | [1,1,10001]   | $p^\uparrow + p \rightarrow \gamma$           |         |
| 10005        | [1,1,10003]   | $p^\uparrow + p \rightarrow \gamma + Z$       |         |
| 10007        | [1,1,10004]   | $p^\uparrow + p \rightarrow W^+$              |         |
| 10008        | [1,1,10005]   | $p^\uparrow + p \rightarrow W^-$              |         |
| 10101        | [2,1,10011]   | $\pi^+ + p^\uparrow \rightarrow \gamma$       |         |
| 10102        | [2,-1,10011]  | $\pi^+ + \bar{p}^\uparrow \rightarrow \gamma$ |         |
| 10103        | [-2,1,10011]  | $\pi^- + p^\uparrow \rightarrow \gamma$       |         |
| 10104        | [-2,-1,10011] | $\pi^- + \bar{p}^\uparrow \rightarrow \gamma$ |         |
| 13200        | [1,1,13200]   | $p^\uparrow + p \rightarrow Z + \gamma$       |         |
| 13201        | [1,1,13201]   | $p^\uparrow + p \rightarrow W^+$              |         |
| 13202        | [1,1,13202]   | $p^\uparrow + p \rightarrow W^-$              |         |
| SIDIS        |               |   |         |
| 200n         | [1,n,2001]    | $p + \gamma \rightarrow h_n$                  |         |
| 201n         | [1,n,2002]    | $d + \gamma \rightarrow h_n$                  |         |
| 202n         | [1,-n,2001]   | $p + \gamma \rightarrow \bar{h}_n$            |         |
| 203n         | [1,-n,2002]   | $d + \gamma \rightarrow \bar{h}_n$            |         |
| 204n         | [1,n,2003]    | $n + \gamma \rightarrow h_n$                  |         |
| 205n         | [1,-n,2003]   | $n + \gamma \rightarrow \bar{h}_n$            |         |
| 2101         | [1,a,2101]    | $p + \gamma \rightarrow h_{1+2}$              | $a > 0$ |
| 2102         | [1,a,2102]    | $p + \gamma \rightarrow h_{1+2+3}$            | $a > 0$ |
| 2103         | [1,a,2103]    | $d + \gamma \rightarrow h_{1+2}$              | $a > 0$ |
| 2104         | [1,a,2104]    | $d + \gamma \rightarrow h_{1+2+3}$            | $a > 0$ |
| 2105         | [1,a,2105]    | $n + \gamma \rightarrow h_{1+2}$              | $a > 0$ |

|      |             |  |         |
|------|-------------|--|---------|
| 2106 | [1,a,2106]  | $n + \gamma \rightarrow h_{1+2+3}$       | $a > 0$ |
| 2111 | [1,-a,2101] | $p + \gamma \rightarrow \bar{h}_{1+2}$   | $a > 0$ |
| 2112 | [1,?,2102]  | $p + \gamma \rightarrow \bar{h}_{1+2+3}$ | $a > 0$ |
| 2113 | [1,?,2103]  | $d + \gamma \rightarrow \bar{h}_{1+2}$   | $a > 0$ |
| 2114 | [1,?,2104]  | $d + \gamma \rightarrow \bar{h}_{1+2+3}$ | $a > 0$ |
| 2115 | [1,?,2105]  | $n + \gamma \rightarrow \bar{h}_{1+2}$   | $a > 0$ |
| 2116 | [1,?,2106]  | $n + \gamma \rightarrow \bar{h}_{1+2+3}$ | $a > 0$ |

## XVI. VERSION HISTORY

- Ver.3.01**
- **TMDX\_DY**: Fixed double-accounting of Jacobian in  $dx_F$ -integration
  - **TMDX\_SIDIS**: Fixed double-accounting of Jacobian in  $dy$ -integrations
  - **TMDX\_SIDIS**: Refactoring and updating into v3 paradigm.
  - **TMDR**: reintroduced the CSS-like evolution factor, as a “pre-compiler” option
  - **QCDinput**: removed connected to LHAPDF library. All changed to self-readers. Hardcore restriction on the number of hadrons.
  - **LHA\_alpha** & **LHA\_PDF**: implemented
- Ver.3.00**
- **DY\_KPC**: implemented
  - **MULTIPLE CHANGES. MASSIVE REORGANIZATION OF CODE**
- Ver.2.06**
- **harpy** Added functions for wgt-function and also for TMDR module
  - **common:tw2Convolution** Fixed bug, with incorrect estimation of the integral reminder at  $x_0 \rightarrow 1$
  - **All modules**: the options N2LO and N3LO can be used instead of NNLO and NNNLO
  - **uTMDFF:coeffFunc** 3-loop coefficient function added.
  - **uTMDPDF:coeffFunc** 3-loop coefficient function added.
  - **uTMDFF:coeffFunc** Corrected missprint in the NNLO coeff.function ( $\sim \pi^4 \delta(\bar{z})$  term)
  - **all .TMD**. Singular list updated to include  $(\ln^3 \bar{x}/(1-x))_+$  terms [appear at N<sup>3</sup>LO]
  - **TMD\_SIDIS** 4-loop coefficient function added.
  - **TMD\_DY** Added the check for the maximum size of the Q-bin size. If bin is too large, it is divided.
  - **TMD\_DY** 4-loop coefficient function added. Fixed a misprint in 3-loops (j0.01% numerically)
  - **TMD\_AD:AD\_Integral**, implemented **RADEvolution** function, and accompanying expressions (roots, Lagrange coefficients).
  - **TMDR**, Fixed mistake with counting of orders (order of  $\Gamma_{\text{cusp}}$  was higher by 1).
  - **TMD\_AD:AD\_atMu**, minor optimizations.
  - **TMD\_AD:AD\_secondary**, added 4-loop expressions for RAD, and zeta-lines.
  - **TMD\_AD:AD\_primary**, added 4-loop expressions for RAD, and anomalous dimensions, 5-loop  $\Gamma_{\text{cusp}}$ . The values are taken from [2202.04660, 2205.02249, 1812.11818]
  - **TMDF**, Added processes: 13001-13059, 13101-13106, 13111-13116, 13200-13202 (for  $A_{LT}$ )
  - **TMDs**, **TMDs.inKT**, **harpy** Added **wgtTMDPDF** routines.
  - **aTMDe\_control** Added **wgtTMDPDF** routines.
  - **aTMDe\_setup** Fixed bug with absent list of hadrons for Sivers.
  - **aTMDe\_setup** Modifications due to **QCDinput** and **wgtTMDPDF** updates.
  - **wgtTMDPDF** Implemented (Thanks to M.Horstmann).
  - **EWinput**: Fixed bug with the S-quark parameter for the interference term in Z-boson production (Thanks to S. Leal-Gomez).
  - **QCDinput**: Added a check for the changing of PDF replica. It is not updated if the PDF replica is same. Updated corresponding function ins TMD’s modules, such that they do not recompute grid if PDF remains the same.
  - **QCDinput**: Added individual function for changing PDF replica for each TMD module. Also added argument to specify hadron. Updated corresponding function ins TMD’s modules.
  - **QCDinput**: Helicity PDF input is added.
  - **QCDinput**: Bug in the enumeration of hadrons fixed.
  - **TMDX\_SIDIS**: Added option for exact factorization scales.
  - **TMDX\_DY**: Added option for exact factorization scales.

- Ver.2.05**
- **TMDX\_SIDIS**: Minor optimization.
  - **TMDf** Added the global mass scale used for normalization of TMD structure functions.
  - **TMDf** Processes unpolarized DY, 101-104, are changed to  $h + p$  (was  $p + h$ ).
  - **TMDf** Added processes for Sivers 10001,10005,10007,10008,10101-10104,12001-12059,12105,12106,12115,12116,
  - **TMDf** Added processes for unpolarized SIDIS 2105,2106,2115,2116,2041-2049.
  - **harpy** Update to python3.
  - **harpy** Added functions for values of uTMDFF, lpTMDPDF, SiversTMDPDF, and DNP, and for TMD in kT space.
  - **harpy** Added functions **Set\_NPparameters** for **lpTMDPDF** and **SiversTMDPDF**.
  - **IO\_functions** Added a check for errors in during the file-reading.
  - **aTMDe\_control** Added version check for replica files
  - **aTMDe\_control** Some code refactoring + **SiversTMDPDF** (fixed couple of bugs in lpTMD-part)
  - **aTMDe\_setup** Some code refactoring + **SiversTMDPDF**
  - **TMDs** Some code refactoring + **SiversTMDPDF**
  - **SiversTMDPDF** Implemented.
  - **\*TMD\*** Added NA-option for the order-definition.
  - **\*TMD\*** Grid-code is moved to a separate directory.
- Ver.2.04**
- **\*TMD\*&\*TMD\*\_model**: Deep refactoring of the code. In particular
    - \* **\*TMD\*\_model** turned to separate modules.
    - \* **\*TMD\*\_modelInterface** are removed. Their functionality is totally inside the model-module.
    - \* The code is distributed into separate text-files.
    - \* Few optimization twicks.
  - **TMDR&TMD\_AD**: Deep refactoring of the code. In particular
    - \* **TMDR\_model** turned to a separate module.
    - \* The definitions of  $\mathcal{D}$  and  $\zeta$  are moved to **TMD\_AD**-module
    - \* Removed evolution type-4, since it coincides with type-3 at  $\zeta = \zeta_{NP}$ .
    - \* Removed function **TMDR R toSL**, since it coincides with **TMDR Rzeta** at  $\zeta = \zeta_{NP}$
    - \* The code is distributed into separate text-files.
    - \* Few optimization twicks.
  - **aTMDe\_control**: Fixed bug in the check of length for  $\lambda_{NP}$  (Thanks to M.Echevarria).
  - **TMDs\_inKT**: The integration routine is updated. Changed the convergence check and added test cases.
  - **TMDX\_SIDIS**: Added integration method 'IO' for x and z integration.
  - **TMDX\_SIDIS**: Fixed bug in the evaluation of cross-section without bin integration and without cuts
- Ver.2.03**
- **TMDR**: Added **orderZETA=-1**, which is used in the LO case.
  - **TMDR**: **zeMuresum** temporary removed.
  - **QEDinput** **Changed routine for evaluation of  $\alpha_{QED}$ . Previous had incorrect determination of boundary value.**
  - **TMDf**: Added extra test for null-value integrals in OGATA quadrature.
  - **TMDf**: OGATA quadrature is updated, with different values of h for qT-intervals
  - **TMDf**: Added test processes N+9999, N+9998
  - **All**: Update of variables definition. Several “magic numbers” removed. Fixed several possible precision leaks.
  - **TMDR**: Functions for derived anomalous dimensions modified. No precalculated numbers, all precalculation is explicit and done inside **TMD\_AD**.

- **TMDR** and **T MDF**: Fixed minor leaks of precision, due to rounding errors in real to double real conversions.
- **TMDR**→**TMD\_AD**: definition of anomalous dimension migrated. Definitions redone in explicit way. Fixed bug in  $\Gamma_3$ .
- **TMD\_AD**: Implemented. This module contains definition of anomalous dimensions used in TMDR.
- **TMD\_numerics**: Implemented. This module contains definition of precision, and definition of various constants.
- **All**: Some refactoring of warning generation.
- **All TMD-modules and models**: Composite TMD option is added.

**Ver.2.02**

- **Common**: Added utility module **IO\_functions**, that contains common function for IO-control. Added added colored output.
- **All TMD-modules**: The initialization array is added to the initialization file. Now, after the initialization each module known the initial NP array.
- **aTMDe\_control**: The initialization procedure read initial values of NP arrays, and (after the initialization procedure) set them.
- **aTMDe\_control**: Added commands **artemide\_GetReplicaFromFile** and **artemide\_NumOfReplicasInFile**. The module commands **setReplica** are not used anymore.
- **aTMDe\_setup**: Added command **CheckConstantsFile**.
- **T MDF**: Added a check for input values of  $x, z, \dots$  Now, for  $x > 1$  the cross-section is evaluated to zero.
- **T MDF**: Added processes 2011-2039;2101-2104;2111-2114
- **TMDX\_DY**: Added additional checks for TMDX\_ConvergenceIsLost trigger. Now, each cross-section-evaluation routine returns  $10^9$  if convergence is lost.
- **TMDX\_SIDIS**: Added  $\alpha_s^3$ -term for hard-matching coefficient. NNNLO defined.
- **TMDX\_SIDIS**: Updated fiducial cut function. Now, it includes  $W_{\max}^2$  and needs 4-arguments.
- **TMDX\_SIDIS**: Added options for accounting of  $q_T$ -correction in  $x_1$  and  $z_1$  irrespectively kinematic  $q_T$ -corrections.
- **TMDX\_SIDIS**: Added options for selection of integration method for X- and Z- bins (SA and S5).
- **TMDX\_SIDIS**: Fixed an error is sign for produced mass-correction. Added missed factor  $(z_1/z)$ .
- **TMDX\_SIDIS**: Fixed rare bug in determination of cuts.
- **TMDX\_SIDIS**: Fixed a bug in the initialization routine (wrong line numbering for const-file.)

**Ver.2.01**

- **All modules**: Added a check for current input-file version. + Small corrections in messages, and bug reports.
- **TMDX\_DY**: Added coefficient functions for Higgs-boson production, and corresponding process 5.
- **TMDX\_DY**: Added option for coefficient function with  $\pi^2$ -resummation [? ].
- **TMDX\_DY**: Added a check: if  $y \notin [-y_0, y_0]$  the  $\sigma = 0$ .
- **lpTMDPDF**: Implemented.
- **TMDR**: Added expressions for gluon evolution.
- **TMDR**: Fixed numeric error in N<sup>3</sup>LO resummed expression for  $\mathcal{D}$  and NNLO+ expression for resummed  $\zeta_\mu$ .
- **TMDR&TMDs**: added evolution type 4. (Exact solution).
- **aTMDe\_control**: fixed a bug with change of NP-parameters for individual modules
- **QCDinput**: fixed a bug with resetting of the PDF-grid
- **T MDF**: Added processes 101-104,1004, 20-22.

**Ver.2.00**

- **TMDX\_DY**: fixed a memory leak in adaptive integration. It could cause to possible source of Segmentation fault error.
- **TMDX\_DY**: Added cut for very-small  $p_T$ .



- **uTMDPDF&uTMDF**: Changed priority of grid-calculation during scale-variation.
- **TMD-models&Twist2Convolution**: Added  $b^*$  parameter. **Model files are not compatible with earlier versions.**
- **Twist2Grid**: The routine for large- $b$  is changed. Now it is faster and more accurate.
- **aTMDe\_control&aTMDe\_setup**: Implemented.
- **ALL MODULES**: Total change of initialization routines, and interface subroutines.

- Ver.1.41**
- **TMDR&TMDs&TMDF**: fixed issues that arise with some older Fortran compilers (thanks to Wen-Chen Chang).
  - **TMDX\_SIDIS**: Totally rewritten: processes redefined, interface redefined, integration routines rewritten, masses correction, parallelization, etc.
  - **TMDX\_DY**: Added extra checks.
  - **uTMDF**: The factor  $z^2$  added as an external, see sec.VIII B 5. It should improve the convergence of "common"-convolution at small  $z$ .
  - **TMDF**: Added Ogata tables for  $\tilde{h} = h0.05$ . They are used for integrations at smaller  $q_T$ .
  - **TMDF**: Fixed potential bug in the initialization order.
  - **TMDF& higher**: Fixed misprint in the name of function **TMDF\_F**.
  - **TMDF**: Added processes 2002-2009.
  - **TMDF&TMDX\_DY**: Added processes [?,4,2013-2018]. W-boson in the narrow width approximation.
  - **TMDs**: Fixed error with the passing to NO parameters in the case of multiple distributions.
- Ver.1.4**
- **constants**: The format of EW input is changed. **Not compatible with older constants-file.**
  - **TMDF**: Added processes 7-12. Fixed a mistake in processes 1,2,2001 (thanks to L.Zoppi & D.Gutierrez-Reyes)
  - **TMDs, uTMDPDF & QCDinput**: Added **\_SetPDFreplica** routine.
  - **HARPY**: Implemented.
  - **TMDs and sub-modules**: Added function **SetReplica**.
  - **TMDs**: Added interface to optimal TMDs
  - **TMDs**: Added check for length of incoming  $\lambda_{NP}$
  - **TMDs**: Fixed bug with incorrect gluon TMDs in functions **\_50**.
  - **TMDs\_inKT**: Implemented.
  - **TMDX\_DY**: Added **xSec\_DY** subroutines.
  - **TMDX\_DY**: Encapsulated process, and cut-parameter variables.
  - **TMDX\_DY**: Defined **p1=2**, which corresponds to integration over  $x_F$ . Removed old functions for  $x_F$  integrations.
  - **TMDX\_DY**: Fixed a bug with variation of  $c_2$  (introduced in ver.1.3).
  - **TMDX\_DY**: Fixed a (potential) bug with  $y$ -symmetric processes.
  - **LeptonCutsDY** : Old version of function removed, cut-parameters encapsulated into single array variable.
  - **LeptonCutsDY** : asymmetric cuts in  $p_T$  are introduced.
  - **LeptonCutsDY** : New function **CutFactor4**, which is analogous to **CutFactor3** but with one integral integrated analytically. Thus, it is more accurate, and faster by 5-20%
  - **LeptonCutsDY** : some rearrangement of variables that makes **CutFactor4** and **CutFactor3** faster by 20%.
  - **uTMDPDF & uTMDF** :  $F_{NP}$  is now function of  $(x, z, b, h, \lambda)$ . For that reason **this version incompatible with earlier versions.**
  - **uTMDPDF & uTMDF**: The common block of the code is extracted into a separate files. It include calculation of Mellin convolution and Grid construction.

- Ver.1.32**
- **TMDX\_DY**: Added the routine with lists of y-bins, in addition to the lists of pt-bins.
  - **TMDX\_DY**: The implementation of parallel computation over the list of cross-sections.
  - **LeptonCutsDY**: The kinematic variables are encapsulated.
  - **TMDX\_DY**: The kinematic variables are encapsulated (by the cost of small reduction of performance).
  - **uTMDR**: Changed behavior at extremely small-b. Now values of b freeze at  $b = 10^{-6}$ .
  - **uTMDPDF** & **uTMDF**: Changed behavior at extremely small-b. Now values of b freeze at  $b = 10^{-6}$ .
  - **TMDR**: Added NNNLO evolution (only for quarks,  $\Gamma_3$  is from 1808.08981). Not tested.
  - **TMDs**: Functions **RuPDFuPDF** and **antiRuPDFuPDF** are added.
- Ver.1.31**
- **Global**: The module **TMDF** is split out from **TMDX...** modules.
  - **Global**: Constant tables are moved to the folder `\tables`.
  - **TMDX...**: Change the structure of process definition.
  - **TMDF**: Fixed bug with throwing exception for failed check of convergence of Ogata quadrature.
  - **TMDF**: Added possibility to vary the Ogata quadrature parameters.
  - **TMDX\_DY**: The structure of interface to integrated cross-section simplified.
  - **TMDX\_DY**: Added trigger for exact power-corrected values of  $x_{1,2}$ .
  - **uTMDPDF** & **uTMDF**: Fixed rare error for exceptional restoration of TMD distribution from grid, then  $f_{NP}$  evaluated to zero.
- Ver.1.3**
- **Global**: Complete change of interface. Interface update for all modules.
  - **uTMDPDF**: Added hadron dependence. **FNP** is now flavour and hadron dependent.
  - **uTMDPDF**: Renormalon correction is removed. As not used.
  - **TMDR**: The grid (and pre-grid) option is removed. Since it was incompatible with new interface. Also the new evolution (type 3) is faster any previous (with grids).
- Ver.1.2(unpub.)**
- **TMDR**: Older version is changed to **uTMDR1**. New evolution routine implemented.
  - **uTMDF**: Implemented.
  - **uTMDPDF**: Fixed bug in evaluation of gluon TMDs, within the evaluation of  $(..)_+$  part.
  - **Global**: Removed functions for the evaluation of only 3-flavours TMDs. As outdated and not used.
  - **Global**: Number of non-perturbative parameters is now read from 'constants'-file. Module **TMDs** initialize sub-modules with accordance to this set.
  - **Global**: Module **TMDX** is renamed into **TMDX\_DY**, also many functions in it renamed.
  - **uTMDX\_SIDIS**: Implemented.
  - **TMDs**: As an temporary solution introduced a rigid cut for  $TMD(\mu < m_q)$ .
  - **TMDX**: Update of Ogata quadrature, with more accurate estimation of convergence.
- Ver.1.1 hotfix** Bugs in **uTMDPDF** and **TMDR** related to the evaluation of gluon TMDs fixed (thanks to Valerio Bertone).
- Ver.1.1**
- **Global**: The physical, numerical and option constant are moved to the file **constants**, where they are read during the initialization stage.
  - **MakeGridsForTMDR**: Update of integration procedures to adaptive. Default grids accordingly updated (no significant effect).
  - **uTMDPDF**: Update of the integration procedure in **uTMDPDF**, to adaptive Gauss-Kronrod (G7-K15). with special treatment of the  $x \rightarrow 1$  singularity.
  - **uTMDPDF**: The procedure for evaluation of TMD for individual flavour (**uTMDPDF\_lowScale(f,x,b,mu)**) is removed, as outdated.
  - **uTMDPDF**: Removed argument  $\mu$ , from **uTMDPDF...**(**x,b**). Added function **mu\_OPE(b)**, which is used as  $\mu$ -definition for TMDs.
  - **uTMDPDF**: Optional griding of TMDs is added. See sec.??
  - **TMDR**: fixed potential error in the "close-to-Landau-pole" exception.

- **TMDs**: fixed potential error in the evaluation of the gluon evolution factor.
- **TMDX**: the name convention of subroutines `CalculateXsection...`, changed to `CalculateXsec...`, to shorten the name length.
- **TMDX**: added functions `CalculateXsec_PTint_Qint_YintComplete(X,qtMin,qtMax,QMin,QMax)` and `CalculateXsec_Qint_YintComplete(X,qtMin,qtMax,QMin,QMax)`.
- **TMDX&TMDs&uTMDPDF**: the independent variation constant  $c_4$  is added (in the ver.1 variation of  $c_3$  and  $c_4$  was simultaneous). The corresponding routines are updated.

**Ver.1** Release: **uTMDPDF**, **TMDR**, **TMDs** and **TMDX** modules. Only Drell-Yan-like cross-sections.