

Synopsis for extractions of unpolarized TMDs: SV19

Alexey A. Vladimirov

These notes contains some information on TMD distributions extracted in the fit SV19, and its later updates. The collection is not complete!, and mainly concerne the latest updates.

Reference: Scimemi:2019cmh

I. Scimemi and A. Vladimirov,

“Non-perturbative structure of semi-inclusive deep-inelastic and Drell-Yan scattering at small transverse momentum,”

JHEP **06** (2020), 137 doi:10.1007/JHEP06(2020)137

Required `artemide` version: >2.04

For the detailed description of the non-perturbative model see sec.2 of reference. NP RAD (2.88):

$$\mathcal{D}(b, \mu) = \mathcal{D}_{\text{resum}}(b^*(b), \mu) + c_0 b b^*(b), \quad b^*(b) = \frac{b}{\sqrt{1 + \frac{\mathbf{b}^2}{B_{\text{NP}}^2}}}. \quad (0.1)$$

NP part of unpolarized TMDPDF:

$$f_{NP}(x, b) = \exp \left(-\frac{\lambda_1 \bar{x} + \lambda_2 x + x \bar{x} \lambda_5}{\sqrt{1 + \lambda_3 x \lambda_4} \mathbf{b}^2} \right). \quad (0.2)$$

NP part of unpolarized TMDFF:

$$D_{NP}(z, b) = \exp \left(-\frac{\eta_1 z + \eta_2 \bar{z}}{\sqrt{1 + \eta_3 (\mathbf{b}/z)^2} z^2} \frac{\mathbf{b}^2}{z^2} \right) \left(1 + \eta_4 \frac{\mathbf{b}^2}{z^2} \right). \quad (0.3)$$

I. SV19_NNLO_M=0

Fit: DY+SIDIS (457+582=1039 points)

Perturbative order: NNLO

Number of replicas: 300

Constants-file: const-DY+SIDIS_NNP31+DSS_nnlo_m=0



Replica-file: SV19_nnlo_m=0.rep


Distinctive feature: Target and production mass corrections in SIDIS kinematics are neglected.

Additional restrictions: Parameter fixed: $B_{NP} = 2\text{GeV}$.

name	N	chiL ² /N	chiD ² /N	chi ² /N	sys.shift%
CDF1	33	0.538	0.100	0.639	7.095
CDF2	39	1.335	0.001	1.336	1.334
D01	16	0.762	0.005	0.768	-1.287
D02	8	1.437	0.000	1.437	0.000
D02m	3	0.421	0.300	0.721	0.293
A7-00y10	5	1.452	0.002	1.454	-0.036
A7-10y20	5	5.522	0.011	5.533	-0.085
A7-20y24	5	1.367	0.000	1.367	-0.010
A8-00y04	5	2.397	0.099	2.496	2.019
A8-04y08	5	2.586	0.464	3.050	2.018
A8-08y12	5	0.826	0.300	1.126	2.164
A8-12y16	5	1.515	0.268	1.782	2.750
A8-16y20	5	0.765	0.287	1.052	3.456
A8-20y24	5	1.716	0.908	2.624	4.143
A8-46Q66	3	0.243	0.085	0.328	-0.265
A8-116Q150	7	0.750	0.031	0.781	0.278
CMS7	8	1.248	0.000	1.248	0.000
CMS8	8	0.772	0.000	0.772	0.000
LHCb7	8	1.612	0.569	2.181	4.535
LHCb8	7	2.773	1.052	3.825	4.453
LHCb13	9	0.607	0.185	0.792	5.044
PHE200	3	0.284	0.000	0.284	-0.250
E228-200	43	1.034	0.043	1.077	33.958
E228-300	53	1.005	0.025	1.029	28.507
E228-400	76	0.834	0.008	0.842	19.414
E772	35	1.895	0.015	1.910	7.230
E605	53	0.513	0.034	0.547	20.057
Total	457	1.096	0.079	1.175	

name	N	chiL ² /N	chiD ² /N	chi ² /N	sys.shift%
hermes.p.vmsub.zxpt.pi+	24	2.371	0.237	2.607	1.983
hermes.p.vmsub.zxpt.pi-	24	1.277	0.044	1.321	0.853
hermes.d.vmsub.zxpt.pi+	24	0.680	0.023	0.703	0.683
hermes.d.vmsub.zxpt.pi-	24	0.880	0.029	0.909	0.772
hermes.p.vmsub.zxpt.k+	24	0.630	0.000	0.630	-0.039
hermes.p.vmsub.zxpt.k-	24	0.627	0.000	0.627	0.061
hermes.d.vmsub.zxpt.k+	24	0.474	0.000	0.474	0.038
hermes.d.vmsub.zxpt.k-	24	1.214	0.001	1.215	0.159
compass.d.h+	195	0.686	0.005	0.691	1.148
compass.d.h-	195	0.817	0.064	0.881	-4.102
Total	582	0.840	0.037	0.877	







Parameter vectors:

$$\text{Mean} = \{2., 0.0399, 0.186, 7.64, 492., 2.23, -2.49, 0., 0., 0.187, 0.476, 0.500, 0.428\} \quad (1.1)$$

$$\text{std} = \{0., 0.0036, 0.0088, 0.52, 70., 0.087, 0.74, 0., 0., 0.016, 0.015, 0.032, 0.027\} \quad (1.2)$$

Distribution of parameters over replicas





II. SV19_NNLO_ALL=0

Fit: DY+SIDIS (457+582=1039 points)

Perturbative order: NNLO

Number of replicas: 300

Constants-file: const-DY+SIDIS_NNP31+DSS_nnlo_all=0


Replica-file: SV19_nnlo_all=0.rep


Distinctive feature: Target and production mass corrections in SIDIS kinematics are neglected; q_T corrections in kinematics are neglected.

Additional restrictions: Parameter fixed: $B_{NP} = 2\text{GeV}$.

name	N	chiL ² /N	chiD ² /N	chi ² /N	sys.shift%
CDF1	33	0.463	0.065	0.529	5.724
CDF2	39	0.771	0.000	0.771	-0.091
D01	16	0.502	0.025	0.527	-2.790
D02	8	0.725	0.000	0.725	0.000
D02m	3	0.080	0.297	0.378	0.308
A7-00y10	5	4.477	0.005	4.483	-0.057
A7-10y20	5	7.977	0.007	7.984	-0.066
A7-20y24	5	1.737	0.000	1.737	0.012
A8-00y04	5	1.960	0.056	2.016	1.442
A8-04y08	5	4.798	0.556	5.355	1.444
A8-08y12	5	2.865	0.281	3.146	1.581
A8-12y16	5	1.795	0.177	1.973	2.062
A8-16y20	5	1.520	0.198	1.717	2.763
A8-20y24	5	1.460	0.618	2.078	3.344
A8-46Q66	3	0.539	0.184	0.724	-1.055
A8-116Q150	7	0.704	0.027	0.731	-0.406
CMS7	8	1.746	0.000	1.746	0.000
CMS8	8	0.918	0.000	0.918	0.000
LHCb7	8	1.840	0.294	2.134	3.259
LHCb8	7	2.673	0.645	3.318	3.488
LHCb13	9	0.727	0.102	0.829	3.741
PHE200	3	0.382	0.030	0.412	-5.055
E228-200	43	1.492	0.014	1.506	19.495
E228-300	53	1.409	0.007	1.416	15.134
E228-400	76	1.136	0.001	1.136	5.364
E772	35	2.237	0.001	2.237	-1.508
E605	53	0.562	0.010	0.572	10.936
Total	457	1.323	0.051	1.374	

name	N	chiL ² /N	chiD ² /N	chi ² /N	sys.shift%
hermes.p.vmsub.zxpt.pi+	24	2.591	0.324	2.916	2.320
hermes.p.vmsub.zxpt.pi-	24	1.472	0.061	1.533	1.007
hermes.d.vmsub.zxpt.pi+	24	0.860	0.072	0.932	1.207
hermes.d.vmsub.zxpt.pi-	24	1.028	0.048	1.077	0.991
hermes.p.vmsub.zxpt.k+	24	0.607	0.000	0.607	-0.011
hermes.p.vmsub.zxpt.k-	24	0.601	0.000	0.601	0.068
hermes.d.vmsub.zxpt.k+	24	0.427	0.000	0.427	0.074
hermes.d.vmsub.zxpt.k-	24	1.194	0.002	1.196	0.215
compass.d.h+	195	0.851	0.022	0.872	2.377
compass.d.h-	195	0.935	0.026	0.960	-2.589
Total	582	0.960	0.037	0.997	





Parameter vectors:

$$\text{Mean} = \{2., 0.0397, 0.185, 6.23, 580., 2.44, -2.53, 0., 0., 0.279, 0.460, 0.436, 0.551\} \quad (2.1)$$

$$\text{std} = \{0., 0.0037, 0.008, 0.45, 78., 0.10, 0.45, 0., 0., 0.013, 0.015, 0.025, 0.027\} \quad (2.2)$$

Distribution of parameters over replicas

