

Chapter 1.2

Predicates and Quantifiers

Propositional logic is not enough to express the meaning of all statements in Mathematics and natural language.

Ex: Is " $x > 1$ " True or false?

Is "x is a great tennis player" True or false?

⇒ Predicate

A predicate $P(x)$ is a sentence that contains a finite number of variables and becomes a proposition when specific values are substituted for the variable, where $P(x)$ is a propositional function and x is a predicate variable.

⇒ Domain

The domain of a predicate variable is the set of all possible values that may be substituted in place of variables.

Eg: "x is great tennis player"

[x is set of all human names]

classfellow

\Rightarrow Quantifier

Quantifiers are words that refer to quantities such as "some" or "all" and indicate how frequently a certain statement is true.

There are two types:-

i) Universal Quantifier

ii) Existential Quantifier

\Rightarrow Universal Quantifier

The phrase "for all" denoted by \forall is called universal quantifier.

Eg:-

Let "All students are smart"

Let $P(x)$ denote " x is smart"

Then the above sentence can be written as $\boxed{\forall x P(x)}$

\Rightarrow Existential Quantifier

The phrase "there exists" denoted by \exists is called existential Quantifier.

Eg:-

Let "There exists x such that $x^2 = 9$ "

Let $P(x) = x^2 = 9$

Then above sentence can be written as

$\boxed{\exists x P(x)}$

classfellow

Ques

Let $D = \{1, 2, 3, 4, \dots, 9\}$. Determine the truth value of each of the following statements.

$$(a) (\forall x \in D), x+4 < 15$$

\Rightarrow True, for every number in D satisfies $x+4 < 15$

$$(b) (\exists x \in D), x+4 \geq 10$$

\Rightarrow True, for if $x=6 \Rightarrow 6+4 \geq 10$

$$(c) (\forall x \in D), x+4 \leq 10$$

\Rightarrow False if $x=7 \Rightarrow 7+4 \geq 11$

$$x=8 \Rightarrow 8+4 \geq 12$$

$$x=9 \Rightarrow 9+4 \geq 13$$

$$(d) (\exists x \in D), x+4 \geq 15$$

\Rightarrow False $x=9 \Rightarrow 9+4 \geq 13 < 15$

False

\Rightarrow Negation of Quantified Statement

Consider the statement

"All students in the class have taken a course in Discrete Mathematics"

This statement can be written as

$$\boxed{\forall x P(x)}$$

where $P(x) \rightarrow$ "x has taken a course in discrete mathematics"

Its negation reads

"It is not the case that all students in the class have taken a course in discrete mathematics"

classfellow

This is equivalent to

"There is a student in the class who has not taken a course in discrete mathematics."

This is simply the existential quantification of the negation of the original propositional function, namely

$$\exists x \neg P(x)$$

This example illustrates the following equivalence

$$\neg \forall x P(x) \equiv \exists x \neg P(x) \quad \text{--- (1)}$$

Example 2:

$$\exists x P(x)$$

$$\neg \exists x P(x) \equiv \forall x \neg P(x) \quad \text{--- (2)}$$

(1) and (2) Negation of Quantified Statements

Statement	Negation
All true $\forall x F(x)$	$\exists x \neg F(x)$ At least 1 false
At least 1 false $\exists x [\neg F(x)]$	$\forall x F(x)$ All true
All false $\forall x [\neg F(x)]$	$\exists x F(x)$ At least 1 True
At least 1 True $\exists x F(x)$	$\forall x [\neg F(x)]$ All false

classfellow

\Rightarrow All human beings are mortal

$P(x)$: x is mortal

$$\Rightarrow \boxed{\forall x P(x)}$$

\Rightarrow Some human beings are mortal

$P(x)$: x is mortal

$$\boxed{\exists x P(x)}$$

\Rightarrow All students are successful

$P(x)$: x is successful

$$\boxed{\forall x P(x)}$$

\Rightarrow Some students are successful

$P(x)$: x is successful

$$\boxed{\exists x P(x)}$$

\Rightarrow Every person is precious

Rephrase: For every x , if x is a person then x is precious.

$M(x)$: x is a person

$A(x)$: x is precious

$$\forall x [M(x) \rightarrow A(x)]$$

\Rightarrow Every student is clever

$M(x)$: x is student

$A(x)$: x is clever

$$\forall x [M(x) \rightarrow A(x)]$$

$\forall x \rightarrow$ implies

$\exists x \rightarrow$ 1 and

classfellow

Page No. _____
Date: 1-1-2022

\Rightarrow All men are mortal

$M(x)$: x is man

$A(x)$: x is mortal

$\forall x [M(x) \rightarrow A(x)]$

\Rightarrow There exists a student

$P(x)$: x is a student

$\exists x P(x)$

\Rightarrow Some students are clever.

$M(x)$: x is a student

$P(x)$: x is clever

$\exists x [M(x) \wedge P(x)]$

\Rightarrow Some students are not successful

$M(x)$: x is a student

$P(x)$: x is successful

$\exists x [M(x) \wedge \sim P(x)]$

\Rightarrow Not all birds can fly

$M(x)$: x is a bird

$P(x)$: x can fly

~~$\exists x [M(x) \wedge P(x)]$~~

$\sim \forall x [M(x) \rightarrow P(x)]$

\Rightarrow There is a student who likes Maths but not Geography

$M(x)$: x is a student

$A(x)$: x likes Maths

$B(x)$: x does not like Geography,

$\exists x [M(x) \wedge A(x) \wedge \sim B(x)]$

classfellow

\Rightarrow Let $P(x)$: x can speak Tamil

$Q(x)$: x know the language C

- (a) There is a student who can speak Tamil and who knows C.

$$\exists x(P(x) \wedge Q(x))$$

- (b) There is a student who can speak Tamil but does not know C.

$$\exists x(P(x) \wedge \neg Q(x))$$

- (c) Every student can either speaks Tamil or knows C.

$$\forall x(P(x) \vee Q(x))$$

- (d) No student can speaks Tamil or knows C.

$$\sim [\forall x(P(x) \vee Q(x))]$$

\Rightarrow All birds can fly.

$B(x)$: x is a bird

$F(x)$: x can fly.

$$\forall x[B(x) \rightarrow F(x)]$$

\Rightarrow Some men are genius

$M(x)$: x is a man

$G(x)$: x is genius

$$\exists x(M(x) \wedge G(x))$$

classfellow

\Rightarrow Some numbers are not rational.

$N(x)$: x is a number

$R(x)$: x is rational

$\exists x [N(x) \wedge \sim R(x)]$

\Rightarrow All mammals are animals.

$M(x)$: x is a mammal

$A(x)$: x is an animal

$\forall x [M(x) \rightarrow A(x)]$

\Rightarrow No natural number is negative.

$N(x)$: x is a natural number

$A(x)$: x is negative

$\sim [\forall x [N(x) \rightarrow A(x)]]$

Negation Questions

\Rightarrow All integers are greater than 8

$P(x)$: x is an integer

$Q(x)$: x is greater than 8

$\forall x (P(x) \rightarrow Q(x))$

$= \forall x (\sim P(x) \vee Q(x))$

Negation: $\sim [\forall x (\sim P(x) \vee Q(x))]$

$= \exists x (P(x) \wedge \sim Q(x))$

Ans

classfellow

\Rightarrow Not all men are genius

$P(x)$: x is a man

$Q(x)$: x is genius

$$\neg \forall x(P(x) \rightarrow Q(x)) \Rightarrow \neg \forall x(\neg P(x) \vee Q(x)) \\ \Rightarrow \forall x \neg (\neg P(x) \vee Q(x))$$

Negation: $\neg [\forall x \neg (\neg P(x) \vee Q(x))]$

$$\Rightarrow \exists x(P(x) \rightarrow Q(x)) \text{ or } \exists x(\neg P(x) \vee Q(x))$$

Answ

$$\Rightarrow (\forall x P(x)) \vee (\exists y P(y))$$

$$\neg [(\forall x P(x)) \vee (\exists y P(y))]$$

$$\neg [\forall x P(x)] \wedge \neg [\exists y P(y)]$$

$$\exists x \neg P(x) \wedge \forall y \neg P(y)$$

$$\Rightarrow \forall x \in R \quad x > 5 \rightarrow x^3 > 125$$

$$\forall x \in R \neg (\neg(x > 5) \vee x^3 > 125)$$

Negation:-

$$\neg [\forall x \in R \quad \neg(x > 5) \vee x^3 > 125]$$

$$\exists x \in R \quad x > 5 \wedge x^3 \leq 125$$

Answ

\Rightarrow Some students are intelligent

$P(x)$: x is a student

$Q(x)$: x is intelligent

$$\exists x(P(x) \wedge Q(x))$$

Negation :- $\forall x(\neg P(x) \wedge \neg Q(x))$

classfellow

\Rightarrow Any integer is either positive or negative

$P(x)$: x is integer

$A(x)$: x is positive

$B(x)$: x is negative

$$\forall x (P(x) \rightarrow (A(x) \wedge B(x)))$$

$$\forall x [\sim P(x) \vee (A(x) \vee B(x))]$$

$$\forall x [P(x) \vee (A(x) \vee B(x))]$$

Negation:-

$$\exists x \sim [\sim P(x) \vee (A(x) \vee B(x))]$$

$$\exists x [P(x) \wedge \sim (A(x) \vee B(x))]$$

$$\exists x [P(x) \wedge \sim A(x) \wedge \sim B(x)]$$

\Rightarrow All flowers are red.

$P(x)$: x is a flower

$Q(x)$: x is red

$$\forall x (P(x) \rightarrow Q(x)) \Rightarrow \forall x [\sim P(x) \vee Q(x)]$$

$$\text{Negation: } \exists x [P(x) \wedge \sim Q(x)]$$

\Rightarrow Some boys are intelligent

$P(x)$: x is a boy

$Q(x)$: x is intelligent

$$\exists x [P(x) \wedge Q(x)]$$

$$\text{Negation: } \forall x [\sim P(x) \vee \sim Q(x)]$$

class fellow

\Rightarrow Every girl is not clever

$P(x)$: x is a girl

$Q(x)$: x is clever

$$\forall x(P(x) \rightarrow \neg Q(x)) \Rightarrow \forall x[\neg P(x) \vee \neg Q(x)]$$

Negation: $\exists x[P(x) \wedge Q(x)]$

\Rightarrow Some numbers are not irrational.

$P(x)$: x is a number

$Q(x)$: x is irrational

$$\exists x(P(x) \wedge \neg Q(x))$$

Negation: $\forall x[\neg P(x) \vee Q(x)]$

Rules of Inference

1. Addition

$$(a) \frac{P}{P \vee q} \quad (b) \frac{q}{P \vee q}$$

2. Simplification

$$(a) \frac{P \wedge q}{P} \quad (b) \frac{P \wedge q}{q}$$

3. Conjunction

$$\frac{\begin{array}{c} p \\ \hline q \end{array}}{p \wedge q}$$

4. Modus Ponens

$$\frac{\begin{array}{c} p \rightarrow q \\ p \end{array}}{q}$$

classfellow

5. Modus tollens

$$\begin{array}{c} p \rightarrow q \\ \neg q \\ \hline \neg p \end{array}$$

6. Hypothetical syllogism

$$\begin{array}{c} p \rightarrow q \\ q \rightarrow r \\ \hline p \rightarrow r \end{array}$$

7. Disjunctive syllogism

$$\begin{array}{c} p \vee q \\ \neg p \\ \hline q \end{array}$$

8. Constructive Dilemma $(p \rightarrow q) \wedge (r \rightarrow s)$

$$\begin{array}{c} p \vee r \\ \hline q \vee s \end{array}$$

9. Destructive Dilemma $(p \rightarrow q) \wedge (r \rightarrow s)$

$$\begin{array}{c} \neg p \vee \neg s \\ \neg p \vee \neg r \end{array}$$

classfellow

Questions on Rules of Inference

Q1. If Ram works hard, he will get a job. Ram works hard, therefore, he will get a job. Check validity.

Sol. p: Ram works hard
q: He will get a job

Premises: $p \rightarrow q, p$ Conclusion: q

1. $p \rightarrow q$ Premise

2. p Premise

3. q Modus Ponens using 1 and 2

The argument is valid.

Q2. If a man is a bachelor, he is unhappy. If a man is unhappy, he dies young. Therefore, bachelors die young. Check validity.

Sol. p: A man is a bachelor
q: A man is unhappy
r: He dies young

Premises: $p \rightarrow q, q \rightarrow r$ Conclusion: $p \rightarrow r$

1. $p \rightarrow q$ Premise

2. $q \rightarrow r$ Premise

3. $p \rightarrow r$ Hypothetical Syllogism using 1 and 2

The argument is valid.

classfellow

Q3. If this number is divisible by 6, then it is divisible by 3. This number is not divisible by 3. Therefore, this number is not divisible by 6.

Sol. p: The number is divisible by 6
q: The number is divisible by 3

Premises: $p \rightarrow q, \neg q$ Conclusion: $\neg p$

1. $p \rightarrow q$ Premises
2. $\neg q$ Premises
3. $\neg p$ Modus tollens using 1 and 2

The argument is valid.

Q4. Either Ram is not guilty or Shyam is telling the truth. Shyam is not telling the truth. Therefore, Ram is not guilty.

Sol. p: Ram is not guilty
q: Shyam is telling the truth

Premises: $p \vee q, \neg q$ Conclusion: $\neg p$

1. $p \vee q$ Premise
2. $\neg q$ Premise
3. $\neg p$ Disjunctive Syllogism using 1 and 2

The argument is valid.

classfellow

Pigeon Hole

If n pigeon hole are occupied by $k+1$ or more pigeons then atleast one pigeonhole is occupied by $k+1$ or more pigeons.

- Q1 Find the minimum numbers of teachers in a college to be sure that 4 of them are born in same month.

$$n=12 \quad k+1=4 \quad k=3$$

$$kn+1 = 12 \times 3 + 1 = 37 \text{ Ans.}$$

- Q2. A box contains 10 blue balls, 20 red balls, 8 green balls, 15 yellow balls and 25 white balls. How many balls must we choose to ensure that we have 12 balls of same colour.

$$n=5 \quad k+1=12 \quad k=11$$

$$kn+1 = 11 \times 5 + 1 = 56 \text{ Ans.}$$

- Q3. Prove that among 1,00,000 people there are two who are born on same time.

Let $P = \{P_1, P_2, \dots, P_{100000}\}$ be the set of people

Let $H = \{H_1, H_2, \dots, H_{24}\}$ be the set of all hours in a day.

By extended pigeon hole principle, there are atleast

$$\left[\frac{|P|}{|H|} \right] \text{ person during the same hour.}$$

$$= 100000 = 4167$$

24

Now $\Omega = \{\Omega_1, \Omega_2, \dots, \Omega_{4167}\}$ and $M = \{M_1, M_2, \dots, M_{60}\}$

classfellow

Then number of person born in same minute
are atleast

$$= \left[\frac{4167}{60} \right] = 70$$

Again $R = \{R_1, R_2, \dots, R_{70}\}$ and $S = \{S_1, S_2, \dots, S_{60}\}$

Then number of person born is same time
(same second) are atleast

$$= \left[\frac{70}{60} \right] = 2$$