Decision log: the embodied-fly path and its branch points

*Every crossroads on the way from the chip-to-animal study to the embodied fly, with the options that were open, the one taken, why, and where to branch from to explore another. Results are not restated here: each stage's numbers live in `report/ROADMAP_EMBODIED_FLY.md` §7, cited to `outputs/metrics.json`, and each stage's registration in `preregistration/`. Commits are on master unless marked otherwise. Decisions taken by the user are marked **(user)**; the rest were taken by the agent and are open to revision.*

How to explore another branch

Check out the commit named in the decision, create a branch, register the alternative in `preregistration/` before running it (fresh seeds if the original result has been seen), and run it with its own command or `python -m neuromps.fly.pipeline --only <step>`.

Decisions

D1. Which fly connectome release the model runs on

- **Options:** FlyWire 630 (the published model's release), FlyWire 783 (the current one).
- **Taken:** 783 for the studies, with a registered 630-vs-783 replication of the published sugar result. **Why:** the current release, with the published release kept as a check.
- **Revisit at:** `fly_chip_to_animal` registration; `neuromps.fly.connectome.load_630`.

D2. Which body simulator for the embodied fly

- **Options:** FlyGym / NeuroMechFly v2 (needs Python 3.12), the FlyMimic musculoskeletal leg loaded in plain MuJoCo, flybody.
- **Taken:** FlyMimic in plain MuJoCo (pinned `mujoco` as the optional `embodied` extra); flybody later for the proboscis. **Why:** muscle-driven, runs in the project's Python 3.11, no new framework. **Revisit:** a FlyGym environment would give six muscle-driven legs only if its musculoskeletal model covers them.

D3. After stage 1 failed: what to change first

- **Options:** rescale the male-CNS synaptic weight further; drive the male nerve cord from descending neurons with the brain removed; switch connectome.
- **Taken:** the nerve cord from descending neurons (stage 1c), then the FlyWire brain joined to it through type-matched descending neurons (stages 1e, 1f). **Why:** the male brain avalanches under the published parameters (`fly_spread`); each connectome is used where its model is validated.

D4. Stage 1e's precondition failed on an absolute rate line

- **Taken:** stage 1f, a relative precondition and depth of reach, on fresh seeds; 1e stays uninterpretable. **Why:** the absolute line had no basis.

D5. Stage 2a failed: how to look for a walking rhythm (user: sweep)

- **Options offered:** fix adaptation parameters from literature; sweep them and report the whole sweep. **Taken (user):** the sweep (stage 2c), with a registered selection rule and confirmation.
- **Revisit:** published adaptation values, if any appear for fly leg or premotor neurons.

D6. The glutamate sign rule (user: go ahead)

- **Taken by the agent:** two bounding rules (glutamate excitatory onto motor neurons only; and everywhere), each with the adaptation grid (stage 2d). **Unexplored:** under the corrected rhythm measure the selection rule would pick (glutamate everywhere, 16 mV, 100 ms); it was not confirmed on fresh seeds.

D7. A different neuron model (user: look at how others make flies walk)

- **Found:** the walking flies online use trained controllers (Eon Systems, flybody); only Pugliese et al. (2025) get a rhythm from nerve-cord wiring, with a firing-rate model.
- **Taken:** reimplement their model from the paper and configuration (their code is unlicensed).
- **Size stand-in (stage 2e):** total synapse count, because the male CNS has no volumes. This starved DNg100; stage 2f drove every command neuron to the same rate; stage 2g used their front-leg network and recruitment rule; stage 2h a geometric input search.
- **Unexplored:** neuron volumes (a future male-CNS release, or MANC volumes through the male-CNS-to-MANC matching); their own network files under a licence.

D8. A flaw in the registered rhythm measure

- **Taken:** a registered audit with a peak-requiring measure (prominence 0.1, calibrated on synthetic signals only), re-scoring stages 1d-2e on their original seeds; registered outcomes kept as recorded, the documents carry both. Stage 1d's D2 withdrawn.
- **Revisit:** prominence 0.05 or 0.2 give the same calibration results; a different peak criterion could be registered.

D9. Where to close the loop first (stage 3)

- **Options:** the front leg (muscle-driven, thinly traced proprioceptors), a middle or hind leg (well traced, but only position-servoed in flybody), the proboscis.
- **Taken:** the front leg through the claw neurons; the unknown flexion/extension tuning bracketed by two assignments and a sham; then a perturbation test (stage 3b).
- **Unexplored:** a hind-leg loop through flybody's servos and its 22 claw neurons; hook and club neurons; campaniform sensilla.

D10. Drugs in the loop (stage 4)

- **Taken:** each knob acts on every network of a trial; movement disruption uses

log2((x + 0.01 mm)/(x_vehicle + 0.01 mm))	with the parent's 0.5 line, 0.05 mm as sensitivity.

- **Disclosed issue:** at full GABA-A block the reach depth is negative, so that condition's index averages the two proboscis terms. **Revisit:** a registered rule for negative depths.

D11. Checking against real flies (stage 5)

- **Taken:** a search protocol registered before any search; every candidate classified.
- **Unexplored:** neuron-specific simulations of the identified-neuron studies (DSOG1, 13A/13B) instead of class-wide knobs; a satiety state for the feeding model.

D12. Release (stage 6)

- **Taken:** a pipeline that reruns every later fly study as its own process, wired into `run_all.py` as a full-run-only `embodied` stage. **Why:** one command reproduces every number; separate processes keep the connectomes from accumulating in memory.

D13. Working overnight without the user (user: continue, commit, push, update docs)

- **Taken:** "push" read as merging finished work into master through a pull request, because GitHub keeps master only; every crossroads tonight is added here.

D14. The precision a chip needs, at the movement level (overnight)

- **Options:** leave stage 4a's timing result as it stands; compute, as the parent study did at the neural level, the chip's change at the dose where movement first fails.
- **Taken:** the latter, as an exploratory analysis (both inputs had been seen), added to the pipeline as `stage4a_precision`. **Why:** the paper's assay-design message rests on the precision a chip needs at behavioural onset; this extends it to movement at no simulation cost.

D15. Confirming the cell the corrected measure selects (overnight)

- **Options:** leave stage 2d's alternative selection unconfirmed (as the audit registered); confirm it on fresh seeds.
- **Taken:** confirm it (registration `embodied_stage2d_confirm_corrected.json`, commit 8495b57), with the corrected measure throughout. **Why:** it is the one loose end the audit named, and it is cheap. **Not taken tonight:** a hind-leg closed loop through flybody's servos (D9), a volume-based size rule (D7), neuron-specific drug simulations (D11), each larger.

D16. Bringing the embodied fly into the technical report and a paper draft (overnight)

- **Taken:** Act VIII of `report/TECHNICAL_REPORT.md` (§27-§31, every number cited, the uncited-number budget unchanged) and `report/PAPER_DRAFT.md` (abstract to methods, every number cited). **Why:** the user asked for a paper and for the report to keep being retouched; the draft is a starting point for the team, not a submission.