



UNIVERSITAT<sub>DE</sub>  
BARCELONA

OPTIMIZATION

MASTER IN FUNDAMENTAL PRINCIPLES OF DATA SCIENCE

---

## OPTIMIZATION PROBLEM 6

CONJUGATE GRADIENT METHOD PROBLEM II

---



DATA SCIENCE @ UNIVERSITAT DE BARCELONA

### **Author**

Vladislav Nikolov Vasilev

FACULTY OF MATHEMATICS AND COMPUTER SCIENCE

ACADEMIC YEAR 2021-2022

## 1 Problem description

Consider the conjugate gradient method applied to the minimization of

$$f(\mathbf{x}) = \frac{1}{2} \mathbf{x}^T A \mathbf{x} - \mathbf{b}^T \mathbf{x}$$

where  $A$  is a positive definite and symmetric matrix.

Show that the iterate  $\mathbf{x}_k$  minimizes  $f$  over

$$\mathbf{x}_0 + \langle \mathbf{v}_0, A\mathbf{v}_0, \dots, A_{k-1}\mathbf{v}_0 \rangle$$

where  $\mathbf{v}_0 = \nabla f(\mathbf{x}_0)$ , and  $\langle \mathbf{v}_0, A\mathbf{v}_0, \dots, A_{k-1}\mathbf{v}_0 \rangle$  is the subspace generated by  $\mathbf{v}_0, A\mathbf{v}_0, \dots, A_{k-1}\mathbf{v}_0$ .

## 2 Solution

Let  $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_k \in \mathbb{R}^n$  with  $k \leq n$  be  $k$  mutually conjugate directions with respect to a given matrix  $A$ , which is symmetric and positive definite. Then,  $\mathbf{x}_k$ , which is a point defined from this directions, will minimize the function  $f$  over the affine set  $\mathbf{x}_0 + \langle \mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_k \rangle$ . Basically, what we want to prove is that  $\langle \mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_k \rangle = \langle \mathbf{v}_0, A\mathbf{v}_0, \dots, A_{k-1}\mathbf{v}_0 \rangle$ .

- If  $k = 1$ , we have that  $\mathbf{z}_1 = -\mathbf{v}_0 = -\nabla f(\mathbf{x})$ .