

# Reptile Formulary and Laboratory Normals\*

*James W. Carpenter, Eric Klaphake,  
and Paul M. Gibbons*

TABLE 1

## Antimicrobial, Antiviral, and Antifungal Agents<sup>a</sup>

Drug dosages are first listed by “most species,” followed by dosages in snakes, lizards, chelonians, and crocodilians; feed and water dosages are generally listed last.

Agent	Dosage	Species/Comments
Acyclovir	≥80 mg/kg PO q24h <sup>1</sup> 80 mg/kg PO q8h or 240 mg/kg/day PO <sup>2</sup>	Tortoises/PD; herpesvirus; poor oral absorption Tortoises/herpesvirus; uncertain efficacy
Amikacin	—  3.48 mg/kg IM once <sup>3</sup> 5 mg/kg IM then 2.5 mg/kg q72h <sup>4</sup>  5 mg/kg IM then 2.5 mg/kg q72h <sup>5,6</sup> 2.5-3 mg/kg IM q72h × 5 treatments <sup>7</sup> 5 mg/kg IM q48h <sup>8</sup> 2.25 mg/kg IM q72h <sup>9</sup>	Potentially nephrotoxic. Often used in combination with penicillins, cephalosporins or metronidazole Pythons/PD Gopher snakes/PD; house at high end of optimum temperature range during treatment Lizards Sea turtles Gopher tortoises/PD; 30°C (86°F) Alligators/PD
Amoxicillin	22 mg/kg PO q12-24h <sup>10,11</sup>	Most species
Amphotericin B	0.5 mg/kg IV q48-72h <sup>12</sup>  0.5-1 mg/kg IV, ICe q24-72h × 14-28 days <sup>10</sup> 1 mg/kg IT q24h × 14-28 days <sup>13</sup>  0.1 mg/kg intrapulmonary q24h × 28 days <sup>14</sup> 1 mg/kg q24h ICe × 2-4 wk <sup>15</sup> 5 mg/150 mL saline × 1 h nebulization q12h × 7 days <sup>16</sup>	Most species/nephrotoxic; can use in combination with ketoconazole; administer slowly Most species/aspergillosis Most species/respiratory infection; dilute with water or saline Greek tortoises/pneumonia Crocodilians Most species/pneumonia
Ampicillin	3-6 mg/kg PO, SC, IM q12-24h <sup>11,17</sup> 10-20 mg/kg SC, IM q12h <sup>18</sup> 20 mg/kg IM q24h <sup>19</sup> 50 mg/kg IM q12h <sup>20,21</sup>	Most species Most species Chelonians Tortoises
Azithromycin	10 mg/kg PO q2-7d <sup>22</sup>	Ball pythons/PD; single-dose study; may cause non-regenerative anemia; <i>Mycoplasma</i> , <i>Cryptosporidium</i> , <i>Giardia</i> ; location dictates dosage frequency: skin q3d; respiratory tract q5d; liver/kidneys, q7d
Carbenicillin	200 mg/kg IM q24h <sup>23</sup> 400 mg/kg IM q24h <sup>24</sup> 400 mg/kg SC, IM q24h <sup>5</sup> 200-400 mg/kg IM q48h <sup>19</sup> 400 mg/kg IM q48h <sup>25</sup>	Carpet pythons/PD Snakes/PD; 30°C (86°F) Lizards Chelonians: may cause skin sloughing in desert tortoises Chelonians/PD
Cefoperazone	100 mg/kg IM q96h <sup>20</sup> 125 mg/kg IM q24h <sup>20</sup>	Snakes/PD (24°C [75°F]) Lizards/PD (24°C [75°F])

\*All reptile formulary information has been modified from J. Carpenter. Exotic animal formulary. 4th ed. St Louis: Saunders, 2013, by permission.

TABLE 1—cont'd

Antimicrobial, Antiviral, and Antifungal Agents<sup>a</sup>

Agent	Dosage	Species/Comments
Cefotaxime	20-40 mg/kg IM q24h <sup>17,19</sup>	Most species
Ceftazidime	20-40 mg/kg SC, IM, q48-72h <sup>6,20,26</sup> 20 mg/kg SC, IM, IV q72h <sup>5,27</sup> 22 mg/kg IM, IV q72h <sup>28</sup>	Most species/in chameleons use q24h Snakes/PD; 30°C (86°F) Sea turtles
Ceftiofur sodium	2.2 mg/kg IM q48h <sup>10</sup> 5 mg/kg SC, IM q24h <sup>29</sup> 2.2 mg/kg IM q24h <sup>10</sup> 4 mg/kg IM q24h <sup>10,17</sup>	Snakes Lizards/PD Turtles Tortoises
Cefuroxime	100 mg/kg IM q24h <sup>10,17</sup>	Most species, snakes/30°C (86°F)
Cephalexin	20-40 mg/kg PO q12h <sup>20</sup>	Most species/unknown absorption
Cephalothin	20-40 mg/kg IM q12h <sup>17,20,30</sup>	Most species
Chloramphenicol	— 40 mg/kg PO, SC, IM q24h, or 20 mg/kg PO, SC, IM q12h <sup>19,20,31</sup> 40 mg/kg SC q24h <sup>32</sup> 50 mg/kg SC q12-72h <sup>20,33</sup>	Most species/public health concern Most species/20 mg/kg may be given q24h in larger crocodilians Snakes/PD (29°C [84°F]) Snakes/PD; q12h in indigo, rat, king snakes; q24h in boids, moccasin snakes; q48h in rattlesnakes; q72h in red-bellied water snakes
Chlortetracycline	200 mg/kg PO q24h <sup>20,30</sup>	Most species
Ciprofloxacin	10 mg/kg PO q48h <sup>10</sup> 11 mg/kg PO q48-72h <sup>34</sup>	Most species Pythons/PD
Clarithromycin	15 mg/kg PO q48-72h <sup>35,36</sup>	Tortoises/PD (mycoplasmosis)
Clindamycin	5 mg/kg PO q12h <sup>30</sup>	Most species
Danofloxacin	6 mg/kg SC, IM <sup>37</sup> 6 mg/kg SC q48h × 30d <sup>38</sup>	Loggerhead sea turtles Tortoises/mycoplasmosis
Doxycycline	5-10 mg/kg PO q24h × 10-45 days <sup>30,31</sup> 50 mg/kg IM, then 25 mg/kg q72h <sup>21,39</sup>	Most species/mycoplasmosis Tortoises 27°C (81°F)
Enrofloxacin	5-10 mg/kg q24h PO, SC, IM, ICe <sup>30</sup>  6.6 mg/kg IM q24h, or 11 mg/kg IM q48h <sup>34</sup> 10 mg/kg IM q48h <sup>40</sup> 5 mg/kg PO, IM q24h <sup>41</sup>  10 mg/kg IM q5d <sup>42</sup> 5 mg/kg IM q24-48h <sup>43</sup>  5 mg/kg IM q12-24h <sup>45</sup>  5 mg/kg IM q48h <sup>7</sup> 10 mg/kg IM q24h <sup>21</sup> 5 mg/kg IV q36-72h <sup>46,47</sup> Nasal flush 50 mg/250 mL sterile water; 1-3 mL/naris q24-48h <sup>31</sup>	Most species/IM is painful and may result in tissue necrosis/sterile abscesses; SC may cause skin discoloration/tissue necrosis Pythons/PD Snakes/PD Lizards/PD; marked pharmacokinetic variability with PO route Monitors/PD Chelonians, most species/PD; neurologic signs and diarrhea reported in a Galapagos tortoise <sup>44</sup> Chelonians/PD; q12h for <i>Pseudomonas</i> and <i>Citrobacter</i> ; q24h for other bacteria Sea turtles Chelonians/PD Crocodilians/PD; mycoplasmosis Tortoises/URT syndrome; use until no more discharge (5-10 days); may use concurrently with parenteral antibiotics
Fluconazole	5 mg/kg PO q24h <sup>48</sup> 21 mg/kg SC once, then 10 mg/kg SC 5 days later <sup>49,50</sup>	Lizards/dermatophytosis Loggerhead sea turtles/PD
Gentamicin	— 2.5 mg/kg IM q72h <sup>32,52</sup> 2.5-3 mg/kg IM, then 1.5 mg/kg q96h <sup>53</sup> 3 mg/kg IM q>96h <sup>54</sup>  6 mg/kg IM q72-96h <sup>55</sup> 1.75-2.25 mg/kg IM q72-96h <sup>9</sup>	Nephrotoxic, <sup>51</sup> especially in snakes Snakes/PD Snakes/PD Turtles/PD (29°C [84°F]); lower dose may be more appropriate Turtles/PD (24°C, [75°F]) Crocodilians/PD; respiratory infection
Griseofulvin	20-40 mg/kg PO q72h × five treatments <sup>56</sup> 15 mg/kg PO q72h <sup>57-59</sup>	Most species/dermatitis; limited success Most species

Continued

TABLE 1—cont'd

Antimicrobial, Antiviral, and Antifungal Agents<sup>a</sup>

Agent	Dosage	Species/Comments
Itraconazole	5 mg/kg PO q24h <sup>60</sup> 10 mg/kg PO q24h <sup>61</sup> 5 mg/kg PO q24h <sup>62</sup> 10 mg/kg PO q48h × 60 days <sup>63</sup> 23.5 mg/kg PO q24h <sup>64</sup> 5 mg/kg PO q24h or 15 mg/kg PO q72h <sup>65</sup>	Most species/some hepatotoxicity noted when used for <i>Chrysosporium</i> anamorph of <i>Nannizziopsis vriesii</i> Snakes Panther chameleons Chameleons (Parson's)/osteomyelitis Lizards/PD; after 3-day treatment, therapeutic plasma concentration persists for 6 days beyond peak concentration; treatment interval not determined Kemp's Ridley sea turtles
Kanamycin	10-15 mg/kg IM, IV q24h (or divided doses) <sup>10,11</sup>	Most species/24°C (75°F); nephrotoxic
Ketoconazole	— 15-30 mg/kg PO q24h × 14-28 days <sup>17</sup> 25 mg/kg PO q24h × 21 days <sup>66</sup> 15 mg/kg q72h PO <sup>57-59</sup> 15-30 mg/kg PO q24h × 14-28 days <sup>67,68</sup> 50 mg/kg PO q24h × 14-28 days <sup>30</sup>	May use antibiotics concomitantly to prevent bacterial overgrowth; may use concurrently with thiabendazole Most species Snakes, turtles Most species Chelonians/PD; systemic infection Crocodilians
Lincomycin	5 mg/kg IM q12-24h <sup>10</sup> 10 mg/kg PO q24h <sup>10</sup>	Most species; nephrotoxic Most species
Marbofloxacin	10 mg/kg PO q48h <sup>69</sup>	Ball pythons/PD
Metronidazole	20 mg/kg PO q48h × ≥7 days <sup>70</sup> 50 mg/kg PO q24h × 7-14 days <sup>34</sup> 20 mg/kg PO q48h <sup>71,72</sup> 20 mg/kg PO q24-48h <sup>73</sup>	Most species Most species; potential side effects at this dose; lower dose may be prudent Snakes/PD Iguanas/PD
Oxytetracycline	6-10 mg/kg PO, IM, IV q24h <sup>10,11</sup> 5-10 mg/kg IM q24h <sup>35</sup> 10 mg/kg IM, IV q5d <sup>74</sup>	Most species/may produce inflammation at injection site Tortoises/mycoplasmosis Crocodilians/PD (27°C [81°F]); mycoplasmosis
Penicillin, benzathine	10,000-20,000 U/kg IM q48-96h <sup>20</sup>	Most species
Penicillin G	10,000-20,000 U/kg SC, IM, IV, ICe q8-12h <sup>11</sup>	Most species/infrequently used
Piperacillin	50-100 mg/kg IM q24h <sup>10,11</sup> 50 mg/kg IM, then 25 mg/kg q24h <sup>10,30</sup> 100 mg/kg IM q48h <sup>75</sup> 100-200 mg/kg SC, IM q24-48h <sup>18</sup> 100 mg/10 mL saline × 30 min nebulization q12h <sup>76</sup>	Most species/nephrotoxic Snakes Snakes/PD Chameleons Most species/pneumonia
Streptomycin	10 mg/kg IM q12-24h <sup>11</sup>	Most species/nephrotoxic; avoid with hepatic dysfunction
Sulfadiazine	25 mg/kg PO q24h <sup>30</sup>	Most species/nephrotoxic
Sulfadimethoxine	90 mg/kg IM, then 45 mg/kg q24h <sup>11</sup>	Most species/nephrotoxic
Thiabendazole	50 mg/kg PO q24h × 14 days <sup>77</sup>	Chelonians/pneumonia; dermatitis
Ticarcillin	50-100 mg/kg IM q24h <sup>11</sup> 50-100 mg/kg IM, IV q24-48h <sup>78</sup>	Most species/nephrotoxic Loggerhead sea turtles/PD
Tobramycin	— 2.5 mg/kg IM q24-72h <sup>10,17</sup> 10 mg/kg IM q24-48h <sup>10</sup>	Nephrotoxic Most species Chelonians/can be given q48h in tortoises
Trimethoprim/sulfadiazine	— 15-25 mg/kg PO q24h <sup>30</sup> 20-30 mg/kg IM q24-48h <sup>34</sup> 30 mg/kg IM q24h × 2 days, then q48h <sup>10,11,20</sup>	Avoid in renal dysfunction; compound parenteral form Most species Most species Most species/can administer PO, SC
Trimethoprim/sulfamethoxazole	10-30 mg/kg PO q24h <sup>11</sup>	Most species/ Avoid in renal dysfunction
Tylosin	5 mg/kg IM q24h × 10-60 days <sup>10,17</sup>	Most species/mycoplasmosis
Voriconazole	10 mg/kg PO <sup>79</sup> 5 mg/kg SC <sup>80</sup>	Bearded dragons/no hepatotoxicity noted when used for <i>Chrysosporium</i> anamorph of <i>Nannizziopsis vriesii</i> Red-eared sliders/exceeded MIC only until 4 h after injection; 26°C (78°F)

ICe, Intracoelomic; IM, intramuscular; IT, intrathecal; IV, intravenous; MIC, minimum inhibitory concentration; PD, pharmacodynamic; PO, by mouth; SC, subcutaneous; URT, upper respiratory tract.

<sup>a</sup>Because reptiles are ectothermic, pharmacokinetics of drugs are influenced by ambient temperature. Antimicrobial therapy should be conducted at the upper end of the patient's preferred (selected) optimum temperature zone.



TABLE 2

## Antiparasitic Agents

Agent	Dosage	Species/Comments
Albendazole	50 mg/kg PO <sup>30</sup>	Most species/ascarids
Carbaryl powder (5%)	Lightly dust animal and environment; rinse after 1 h; repeat in 7 days <sup>20,81</sup>	Lizards; snakes/mites
Chloroquine	125 mg/kg PO q48h × 3 treatments <sup>30</sup>	Tortoises/hemoprotozoa
Emodepside (1.98%) + praziquantel (7.94%)	1.12 mL/kg <sup>82,83</sup>	Many species/PD; nematodes; cestodes; aquatic turtles must be kept dry 48 h after application; appears safe, but need more safety and efficacy data
Fenbendazole	—	May have antiprotozoan effect; may cause leukopenia, avoid with septicemia <sup>84</sup>
	25-100 mg/kg PO q14d × 1-4 treatments <sup>16,39,85</sup>	All species/nematodes
	50 mg/kg PO q24h × 3-5 days <sup>20,86,87</sup>	All species/nematodes (× 3 days); flagellates and giardia in chameleons (× 5 days)
	100 mg/kg once <sup>88</sup>	Tortoises/nematodes; ova still sheds for 30 days
Fipronil	Spray or wipe on then wash off in 5 min q7-10d prn <sup>20,81</sup>	Most species/mites, ticks; beware of reactions to alcohol carrier; needs safety evaluation <sup>89</sup>
Imidacloprid and moxidectin	0.2 mg/kg topical q14d × 3 treatments <sup>90</sup>	Lizards/eliminated hookworms and pinworms; needs safety and pharmacokinetic evaluation
Ivermectin	—	Do not use in chelonians, <sup>91</sup> crocodilians, indigo snakes, or skinks <sup>30,39,87</sup>
	0.2 mg/kg PO, SC, IM repeat in 14 days <sup>5,92,93</sup>	Snakes (except indigos), lizards (except skinks) <sup>39</sup> /nematodes, <sup>94</sup> mites; dilute with propylene glycol for oral use; colored animals may have skin discoloration at injection site; rare adverse effects reported in chameleons <sup>5</sup> ; do not use within 10 days of diazepam or tiletamine/zolazepam; rare death and occasional nervous system signs, lethargy, or inappetence; <sup>87</sup> pentastomids in monitors (with dexamethasone) <sup>92</sup>
	5-10 mg/L water topical spray q3-5d up to 28 days <sup>87</sup>	Snakes (except indigos), lizards (except skinks)/mites; less effective than fipronil; spray on skin and in newly cleaned cage, then allow to dry before replacing water dish
Metronidazole	—	Protozoan (i.e., flagellates, amoebae) overgrowth; may stimulate appetite; may cause severe neurologic signs at doses >200 mg/kg <sup>95</sup> ; death occurred in indigo and mountain king snakes at 100 mg/kg <sup>16</sup> ; for small patients, injectable form can be administered PO; oral liquid not available in the United States but can be compounded
	40-100 mg/kg PO, repeat in 10-14 days <sup>11</sup>	Most species/flagellate overgrowth
	20 mg/kg PO q48h <sup>71</sup>	Corn snakes/PD; 28°C (82°F); protozoa
	40 mg/kg PO, repeat in 14 days <sup>16,20</sup>	Uracoan rattler, milk, tricolor king, and indigo snakes/flagellates
	40-60 mg/kg PO q7d × 2-3 doses <sup>96</sup>	Chameleons/flagellates; amoebae
	40-200 mg/kg PO, repeat in 14 days <sup>97</sup>	Geckos/ocular lesions (40 mg/kg) and SC lesions (200 mg/kg) caused by <i>Trichomonas</i>
	50 mg/kg PO q24h × 2-5 days <sup>86</sup>	Chameleons/when accompanied by increased gastrointestinal symptoms
Paromomycin	20 mg/kg ICe q48h <sup>98</sup>	Red-eared sliders/PD; ICe administration not recommended; needs further safety evaluation
	25 mg/kg PO q24h × 5 days or 50 mg/kg PO q14 days prn <sup>99</sup>	Chelonians (tortoises)/amoebae; use 25 mg/kg dosage for clinically ill cases
	35-100 mg/kg PO q24h × ≤28 days <sup>11,16,93</sup>	Most species/amoebae
	100 mg/kg PO q24h × 7 days, then 2 ×/wk × 3 mo <sup>100</sup>	Snakes/cryptosporidia; reduced clinical signs and oocyte shedding; does not eliminate the organism
Permethrin	300-360 mg/kg PO q48h × 14 days <sup>101</sup>	Lizards (gila monsters)/cryptosporidia
	300-800 mg/kg PO q24h prn <sup>102</sup>	Geckos/cryptosporidia; reduced clinical signs; does not eliminate the organism
Permethrin	Environmental treatment, 1 s of spray/ft <sup>2</sup> ; wait until dry before returning animal to enclosure <sup>81</sup>	Lizards, snakes/mites; ticks; FDA approved; safe and effective; wash immediately if accidentally applied to skin
	Topical <sup>81</sup>	Tortoises/ticks

Continued

TABLE 2—cont'd

## Antiparasitic Agents

Agent	Dosage	Species/Comments
Ponazuril	30 mg/kg PO q48h × 2 treatments <sup>103,104</sup>	Bearded dragons/coccidiosis
Praziquantel (also see Emodepside)	8 mg/kg PO, SC, IM, repeat in 14 days <sup>5,18,30</sup>	Most species/cestodes, trematodes; higher dosages have been administered <sup>20</sup>
	5-10 mg/kg PO q14d <sup>86</sup>	Chameleons/flukes may best be left untreated if not causing a problem
	25-50 mg/kg PO q3h × 3 treatments <sup>105,106</sup>	Sea turtles (green, loggerhead)/PD; spirorchidiasis
Pyrantel pamoate	5 mg/kg PO, repeat in 14 days <sup>87</sup>	Most species/nematodes
	25 mg/kg PO q24h × 3 days; repeat in 3 wk <sup>20,107</sup>	Most species/ascarids, hookworms, pinworms
Pyrethrin spray (0.09%)	Topical q7d × 2-3 treatments <sup>81</sup>	Most species/use water-based labeled for kittens and puppies; apply with cloth; can also spray cage, wash out after 30 min; use sparingly and with caution; pyrethroids are safer
Quinacrine	19-100 mg/kg PO q48h × 14-21 days <sup>108</sup>	Most species/some hematozoa
Quinine sulfate	75 mg/kg PO q48h × 14-28 days <sup>108</sup>	Most species/some hematozoa; toxic at >100 mg/kg q24h; ineffective against exoerythrocytic forms
Resmethrin spray/shampoo	Topical, repeat q≥10d prn <sup>81</sup>	Most species/mites; pyrethroid; safer than pyrethrins; use with care; spray (0.35%) or shampoo entire animal, rinse off immediately in running, tepid water; protect eyes (other than snakes) with 1 drop of mineral oil; lightly spray environment, wipe off in 5-10 min
Spiramycin	160 mg/kg PO q24h × 10 days, then 2x/wk for 3 mo <sup>100</sup>	Snakes/cryptosporidia; may reduce clinical signs and oocyte shedding; does not eliminate
Sulfadiazine, sulfamerazine	—	Most species/coccidia; watch renal dysfunction <sup>95</sup>
	75 mg/kg PO, then 45 mg/kg q24h × 5 days <sup>11,86,108</sup>	Most species/coccidia
	25 mg/kg PO q24h × 21 days <sup>5,77,108</sup>	Snakes, lizards/coccidian
Sulfadimethoxine	50 mg/kg PO q24h × 3-5 days, then q48h prn <sup>87</sup>	Most species/coccidia; watch renal dysfunction
	90 mg/kg PO, IM, IV, then 45 mg/kg q24h × 5-7 days <sup>11,16,108</sup>	Most species/coccidia
Sulfamethazine	50 mg/kg PO q24h × 3 days, off 3 days, on 3 days <sup>108</sup>	Most species/coccidia
	25 mg/kg PO, IM q24h × 21 days <sup>109</sup>	Most species/coccidia
	75 mg/kg PO, IM, IV, then 40 mg/kg q24h × 5-7 days <sup>16,17</sup>	Most species/coccidia; ensure adequate hydration and renal function
Toltrazuril 5%	5-15 mg/kg q24h × 3 days <sup>110</sup>	Bearded dragons/coccidia
	15 mg/kg q48h × 10 days; discontinue for 2 wk; repeat q48h × 10 days prn <sup>111</sup>	Tortoises/intranuclear coccidiosis; needs safety, efficacy, and pharmacokinetic study
Trimethoprim/sulfa	—	Most species/coccidia; avoid potentiated sulfa drugs in cases of dehydration or renal dysfunction <sup>95</sup>
	30 mg/kg PO q24h × 2 days, then q48h × 21 days <sup>5,108</sup>	Most species/coccidia
	30 mg/kg IM q24h × 2 days, then 15 mg/kg IM q48h × 10-28 days <sup>108</sup>	Most species/coccidia

FDA, U.S. Food and Drug Administration; ICe, intracoelomic; IM, intramuscular; IV, intravenous; PD, pharmacodynamic; PO, by mouth; prn, as required; SC, subcutaneous.

TABLE 3

## Chemical Restraint/Anesthetic and Analgesic Agents

Agent	Dosage	Species/Comments
Acepromazine	0.05-0.25 mg/kg IM <sup>31</sup> 0.1-0.5 mg/kg IM <sup>112,113</sup>	Most species/preanesthetic with ketamine Most species/preanesthetic; reduce by 50% if used with barbiturates
Alphaxalone	— 6-9 mg/kg IV, or 9-15 mg/kg IM <sup>114</sup>  6-15 mg/kg IM, IV <sup>116</sup> 9 mg/kg IV <sup>117</sup>  15 mg/kg IM <sup>112</sup>  24 mg/kg ICe <sup>118</sup>	Not available in the United States Most species/good muscle relaxation; variable results; needs more evaluation; possible violent recovery; <sup>115</sup> avoid within 10 days of DMSO Most species Snakes, lizards/induction; not effective for blotched blue-tongued skinks Lizards, chelonians/induction, 35-40 min; duration 15-35 min; good muscle relaxation Chelonians (red-eared sliders)/surgical anesthesia with good relaxation
Atipamezole	Same volume SC, IV, IP as medetomidine or dexmedetomidine <sup>a,119,120</sup>	Most species/medetomidine and dexmedetomidine reversal; severe hypotension in gopher tortoises IV <sup>121</sup>
Atropine	0.01-0.04 mg/kg SC, IM, <sup>122</sup> IV, <sup>12</sup> ICe <sup>123</sup>  0.5 mg/kg IM, IV, IT, IO <sup>95</sup>	Most species/preanesthetic; rarely indicated; for profound bradycardia <sup>123</sup> ; may help prevent intracardiac shunting <sup>57</sup> ; ineffective dose in green iguanas <sup>124</sup> Most species/bradycardia, decrease secretions, CPR
Bupivacaine	1-2 mg/kg local q4-12h prn <sup>116</sup>	Most species; 4 mg/kg max dose
Buprenorphine	0.005-0.02 mg/kg IM q24-48h <sup>125</sup> 0.1-1 mg/kg IM <sup>126</sup>	Most species/analgesia Most species/analgesia
Butorphanol	—  0.4-1 mg/kg SC, IM <sup>123</sup> 0.5-2 mg/kg IM or 0.2-0.5 mg/kg IV, IO <sup>126</sup> 1-1.5 mg/kg SC, IM <sup>123</sup>  0.2 mg/kg IM <sup>125,130</sup> 20 mg/kg IM <sup>131</sup>	Combinations (see ketamine) follow; recent studies call into question use in providing analgesia in reptiles, including red-eared sliders, ball pythons, bearded dragons, and green iguanas; respiratory depression at high doses <sup>127-129</sup> Most species/analgesia; sedation; preanesthetic Most species/preanesthetic Lizards/administer 30 min before isoflurane for smooth, shorter induction Chelonians/tranquilizer Cornsnakes/potential analgesia
Butorphanol (B)/midazolam (M)	(B) 0.4 mg/kg + (M) 2 mg/kg IM <sup>132</sup>	Most species/preanesthetic; administer 20 min before induction
Carprofen	1-4 mg/kg PO, SC, IM, IV q24h, <sup>133</sup> follow with half the dose q24-72h <sup>134</sup>	Most species/analgesia; nonsteroidal antiinflammatory
Dexmedetomidine <sup>a</sup>	—	$\alpha_2$ agonist that has replaced medetomidine <sup>a</sup>
Diazepam	—  0.5 mg/kg IM, IV <sup>95</sup> 0.2-0.8 mg/kg IM <sup>123</sup>  0.2-2 mg/kg IM, IV <sup>116</sup> 2.5 mg/kg PO <sup>123</sup> 0.2-1 mg/kg IM <sup>116,123</sup>	Combinations follow; muscle relaxation; give 20 min before anesthesia; potentially reversible with flumazenil; interaction with ivermectin All species/seizures Snakes/use with ketamine for anesthesia with muscle relaxation Snakes, lizards Iguanas/reduce anxiety that often leads to aggression Chelonians/use with ketamine for anesthesia with muscle relaxation
Doxapram	5 mg/kg IM, IV <sup>132</sup> q10min prn  4-12 mg/kg IM, IV <sup>123</sup> 20 mg/kg IM, IV, IO <sup>95</sup>	Most species/respiratory stimulant; reduces recovery time; reported to partially "reverse" effects of dissociatives <sup>133</sup> Most species/respiratory stimulant Most species/respiratory stimulant
Epinephrine (1:1000)	0.5-1 mg/kg IV, IO, IT <sup>95</sup>	Most species/CPR, cardiac arrest
Fentanyl	2.5 $\mu$ g/h patch to caudodorsal lumbar skin <sup>135</sup>	Prehensile-tailed skinks/no side effects reported after 24 h when skink blood levels reached human therapeutic levels; environmental temperature can significantly affect absorption

Continued

TABLE 3—cont'd

## Chemical Restraint/Anesthetic and Analgesic Agents

Agent	Dosage	Species/Comments
Flunixin meglumine	0.1-0.5 mg/kg IM q12-24h <sup>133</sup> 1-2 mg/kg IM q24h × 2 treatments <sup>136,137</sup> 0.5-2 mg/kg IM q12-24h <sup>116</sup>	Most species/analgesia; use max of 3 days Lizards/nonsteroidal antiinflammatory; postsurgical analgesia Most species/nonsteroidal antiinflammatory; analgesia
Glycopyrrolate	0.01 mg/kg SC, <sup>122</sup> IM, IV <sup>132</sup>	Most species/preanesthetic; excess mucus; use with prolonged bradycardia; dose does not work in green iguanas <sup>124</sup>
Haloperidol	0.5-10 mg/kg IM q7-14d <sup>138</sup>	Boids/aggression management
Isoflurane	3%-5% induction <sup>13</sup> 1%-3% maintenance <sup>39</sup>	Most species/inhalation anesthetic of choice in reptiles
Ketamine	—  20-60 mg/kg IM, or 5-15 mg/kg IV <sup>31</sup>  22-44 mg/kg SC, IM <sup>115,139</sup> 55-88 mg/kg SC, IM <sup>139</sup>  20-60 mg/kg SC, IM <sup>122,140</sup> 60-80 mg/kg IM <sup>39</sup>  30-50 mg/kg SC, IM <sup>122,140</sup> 20-60 mg/kg IM <sup>113,140,141</sup>  25 mg/kg IM, IV <sup>7</sup> 38-71 mg/kg ICe <sup>142</sup>  60-90 mg/kg IM <sup>112,140</sup>  20-40 mg/kg SC, IM, ICe (sedation), to 40-80 mg/kg (anesthesia) <sup>143</sup>	Muscle relaxation and analgesia may be marginal; prolonged recovery with higher doses; larger reptiles require lower dose; painful at injection site; safety questionable in debilitated patients; avoid with renal dysfunction; snakes may be permanently aggressive after ketamine anesthesia <sup>115</sup> ; recommend only as preanesthetic before isoflurane for surgical anesthesia Most species/muscle relaxation improved with midazolam or diazepam Most species/sedation Most species/surgical anesthesia; induction, 10-30 min; recovery, 24-96 h Snakes/sedation; induction, 30 min; recovery, 2-48 h Snakes/light anesthesia; IPPV may be needed at higher doses Lizards/sedation; variable results Chelonians/sedation; induction, 30 min; recovery, ≥24 h; potentially dangerous in dehydrated and debilitated tortoises Sea turtles/sedation Green sea turtles/anesthesia; induction, 2-10 min; duration, 2-10 min; recovery, <30 min Chelonians/light anesthesia; induction, <30 min; recovery, hours to days; require higher doses than most other reptiles Crocodilians/induction, <30-60 min; recovery, hours to days; in larger animals, 12-15 mg/kg may permit tracheal intubation <sup>123</sup> ; not recommended alone in Nile crocodiles <sup>144</sup>
Ketamine (K)/butorphanol (B)	See (K) dosages + (B) ≤1.5 mg/kg IM <sup>123</sup> (K) 10-30 mg/kg + (B) 0.5-1.5 mg/kg IM <sup>123</sup>	Snakes/anesthesia with improved muscle relaxation Chelonians/minor surgical procedures (i.e., shell repair)
Ketamine (K)/diazepam (D)	See (K) dosages + (D) 0.2-0.8 mg/kg IM <sup>123</sup> (K) 60-80 mg/kg <sup>112</sup> + (D) 0.2-1 mg/kg IM <sup>123</sup>	Snakes/anesthesia with improved muscle relaxation Chelonians/anesthesia; muscle relaxation
Ketamine (K)/medetomidine (M) <sup>a</sup>	—  (K) 10 mg/kg + (M) 0.1-0.3 mg/kg IM <sup>145</sup> (K) 3-8 mg/kg + (M) 0.025-0.08 mg/kg IV <sup>146</sup> (K) 4 mg/kg + (M) 0.04 mg/kg IM <sup>147</sup> (K) 4-10 mg/kg + (M) 0.04-0.14 mg/kg IM <sup>148</sup>  (K) 5 mg/kg + (M) 0.05 mg/kg IV <sup>149</sup>  (K) 5-10 mg/kg IM + (M) 0.1-0.15 mg/kg IM, IV <sup>150</sup> (K) 10-20 mg/kg IM + (M) 0.15-0.3 mg/kg IM, IV <sup>150</sup>  (K) 5-10 mg/kg + (M) 0.1-0.15 mg/kg IM <sup>151</sup> (K) 10-15 mg/kg + (M) 0.15-0.25 mg/kg IM <sup>151</sup>	Medetomidine no longer commercially available, can be compounded <sup>a</sup> ; reverse with atipamezole Most species Giant tortoises (Aldabra) Green sea turtles Chelonians/sedation and muscle relaxation for shell repair Loggerhead sea turtles/induction of anesthesia for intubation Tortoises (small-medium) Turtles (fresh water) Alligators/adults Alligators/juveniles

TABLE 3—cont'd

## Chemical Restraint/Anesthetic and Analgesic Agents

Agent	Dosage	Species/Comments
Ketamine (K)/midazolam (M)	(K) 20-40 mg/kg + (M) $\leq$ 2 mg/kg IM <sup>152</sup> (K) 60-80 mg/kg <sup>112</sup> + (M) $\leq$ 2 mg/kg IM <sup>123</sup>	Chelonians/sedation; muscle relaxation Chelonians/anesthesia; muscle relaxation
Ketoprofen	2 mg/kg SC, IM q24h <sup>133</sup>	Most species/analgesia; in green iguanas, IM, IV q24h; administration at this dose may be too frequent based on PK study <sup>153</sup>
Lidocaine (0.5%-2%)	2-5 mg/kg; local or topical <sup>116,123</sup>	Most species/local analgesia; infiltrate to effect, often used in conjunction with chemical; 10 mg/kg maximum dosage immobilization
Medetomidine <sup>a</sup>	—	Medetomidine is no longer commercially available but can be compounded <sup>a</sup> ; poor immobilization alone, see combinations
Meloxicam	0.1-0.5 mg/kg PO, SC q24-48h <sup>57-59</sup> 0.5 mg/kg PO, IM or 0.22 mg/kg IV <sup>154</sup>  0.2 mg/kg PO, IV q24h <sup>155</sup> 0.1-0.2 mg/kg PO, IM q24h $\times$ 4-10 days <sup>148</sup>	Most species Red-eared sliders/PK; found better absorption IM vs. PO; <sup>154</sup> after IV administration, plasma levels decreased rapidly Green iguanas/PD Chelonians
Meperidine/Pethidine	5-10 mg/kg IM q12-24h <sup>125</sup>  20 mg/kg IM q12-24h <sup>133</sup> 2-4 mg/kg ICe q6-8h <sup>57</sup> 2-4 mg/kg ICe <sup>156</sup>	Most species/analgesia; no noticeable effect in snakes, even at 200 mg/kg Most species/analgesia Lizards Nile crocodiles/analgesia
Methohexital	5-20 mg/kg SC, <sup>139</sup> IV <sup>12</sup>  9-10 mg/kg SC, <sup>157</sup> ICe	Most species/induction, 5-30 min; recovery, 1-5 h; use at 0.125%-0.5% concentration; much species variability; decrease dose 20%-30% for young animals; avoid use in debilitated animals Colubrids/induction, $\geq$ 22 min; recovery, 2-5 h; does not produce soft tissue irritation seen with other barbiturates; may need to adjust dosage in obese snakes
Metomidate	10 mg/kg IM <sup>145,158</sup>	Snakes/profound sedation; not available in the United States
Midazolam	—  2 mg/kg IM <sup>115,139</sup>  0.5-2 mg/kg <sup>159</sup> 1.5 mg/kg IM <sup>160</sup>	See butorphanol, ketamine for combinations; can be reversed by flumazenil Most species/preanesthetic; increases efficacy of ketamine; effective in snapping turtles; not in painted turtles <sup>139</sup> Lizards Turtles (red-eared sliders)/sedation; onset, 5.5 min; duration, 82 min; recovery, 40 min; much individual variability
Morphine	—  1.5-6.5 mg/kg IM <sup>128,129,131,148</sup>  10 mg/kg IM <sup>129</sup>	No effective analgesic dose in cornsnakes <sup>129</sup> Red-eared sliders (long-lasting respiratory depression), freshwater crocodiles, <i>Anolis</i> lizards/may be effective thermal analgesia in bearded dragons Crocodilians/analgesia
Oxymorphone	0.025-0.1 mg/kg IV <sup>12</sup>  0.05-0.2 mg/kg SC, IM q12-48h <sup>125</sup>	Some species/analgesia; avoid in cases with hepatic or renal dysfunction; no noticeable effect in snakes, even at 1.5 mg/kg <sup>115</sup> See previous dosage comments within Oxymorphone
Pentazocine	2-5 mg/kg IM q6-24h <sup>125</sup>	Analgesia

Continued



TABLE 3—cont'd

## Chemical Restraint/Anesthetic and Analgesic Agents

Agent	Dosage	Species/Comments
Propofol	—  0.3-0.5 mg/kg/min IV, IO constant rate infusion or 0.5-1 mg/kg IV, IO periodic bolus <sup>116</sup> 5-10 mg/kg IV, IC <sup>158,162</sup> 3-5 mg/kg IV, IO <sup>150,163</sup>  5-10 mg/kg IV, IO <sup>164</sup>  2 mg/kg IV <sup>132</sup> 3-5 mg/kg IV <sup>148</sup> 10 mg/kg IV <sup>165</sup> 20 mg/kg IV <sup>165</sup> 10-15 mg/kg IV <sup>143</sup>	If administered in supravertebral sinus, be aware of potential submeningeal delivery <sup>161</sup> ; anesthesia; rapid, smooth induction; 15-25 min anesthesia and restraint in most species; rapid, excitement-free recovery; administered IV/IO slowly; reduce dosages by as much as 50% in premedicated animals; apnea and bradycardia; intubation and IPPV generally required Most species/maintenance anesthesia; must provide respiratory and thermal support Snakes Lizards/intubation and minor diagnostic procedures; may need additional dose in 3-5 min; less cardiopulmonary depression Iguanas/higher dose is recommended for induction for short duration procedures or intubation Giant tortoises Chelonians/sedation (i.e., shell repair) Red-eared sliders/40-85 min anesthesia Red-eared sliders/60-120 min anesthesia Crocodilians/duration, 0.5-1.5 h; maintain with gas anesthetics; experimental IM with hyaluronidase
Sevoflurane	To effect <sup>5,166</sup>	Most species/anesthesia; rapid induction and recovery when intubated
Tiletamine/zolazepam	—  4-5 mg/kg SC, IM <sup>139</sup>  5-10 mg/kg IM <sup>132</sup> 3 mg/kg IM <sup>150</sup>  10-30 mg/kg IM <sup>112</sup> to 20-40 mg/kg IM <sup>13,168</sup>  5-10 mg/kg IM, IV <sup>123</sup>  1-2 mg/kg IM <sup>15</sup> 2-10 mg/kg IM <sup>123</sup> 10-40 mg/kg SC, IM, ICe <sup>143</sup> 15 mg/kg IM <sup>170</sup>	Sedation, anesthesia; severe respiratory depression possible—IPPV; <sup>39</sup> variable results; prolonged recovery; lower dose in heavier species; good for muscle relaxation before intubation <sup>130,167</sup> Most species/sedation; induction, 9-15 min; recovery, 1-12 h; adequate for most noninvasive procedures Most species Snakes/facilitates handling and intubation of large snakes; induction, 30-45 min; prolongs recovery Snakes, lizards/induction, 8-20 min; recovery, 2-10 h; variable results; longer sedation and recovery times at 22°C (72°F) than at 30°C (86°F); <sup>169</sup> good sedation in boa constrictors at 25 mg/kg IM <sup>169</sup> ; generally need to supplement with inhalational agents for surgical anesthesia; some snakes died at 55 mg/kg Large tortoises/facilitates intubation; if light, mask with isoflurane rather than redosing Crocodilians/recovery takes several hours Large crocodilians/may permit intubation Crocodilians, anesthesia Alligators/induction, >20 min; minor procedures
Tramadol	11 mg/kg PO <sup>171</sup> 5-10 mg/kg PO <sup>172</sup>	Bearded dragons Red-eared sliders/thermal analgesia, higher doses may affect ventilation
Xylazine	—  0.1-1.25 mg/kg IM, IV <sup>12</sup> 0.1-1 mg/kg IM <sup>15</sup> 1-2 mg/kg IM <sup>113,143</sup>	Infrequently used; variable effects; potentially reversible with yohimbine Most species Crocodilians/atipamezole better reversal Nile crocodiles

CPR, Cardiopulmonary resuscitation; IC, intracardiac; ICe, intracoelomic; IM, intramuscular; IO, intraosseous; IP, intraperitoneal; IPPV, intermittent positive pressure ventilation; IT, intrathecal; IV, intravenous; PD, pharmacodynamic; PK, pharmacokinetic; PO, by mouth; prn, as required; SC, subcutaneous.

<sup>a</sup>Medetomidine is no longer commercially available, although it can be obtained from select compounding services. Dosages listed here are a guide for possible use with dexmedetomidine, an  $\alpha_2$  agonist that is the active optical enantiomer of the racemic compound medetomidine. It is used at half the dose of medetomidine but the same volume because of higher concentration. Although the same effects would be expected as with medetomidine, there are limited data on the efficacy and safety of dexmedetomidine in reptiles, and, to date, it appears to have been seldom used clinically in reptiles. The effects of the volume-to-volume use of the two drugs may not be equivalent, so the dose of dexmedetomidine may need to be adjusted based on clinical response.

TABLE 4

## Hormones and Steroids

Agent	Dosage	Species/Comments
Arginine vasotocin	0.01-1 µg/kg IV, ICe <sup>173</sup> q12-24h × 2-3 treatments	Most species/dystocias; give 30-60 min after par- enteral Ca; available for research; higher doses reported; 0.5 µg/kg commonly recommended
Calcitonin	1.5 U/kg SC q8h × 14-21 days prn <sup>12</sup> 50 U/kg IM repeat in 14 days <sup>30,174</sup> 50 U/kg q7d × 2-3 doses <sup>175,176</sup>	Most species (e.g., iguanas)/severe NSHP; administer after Ca; avoid if hypocalcemic Green iguanas/salmon calcitonin; do not give if hypocalcemic
Dexamethasone	0.6-1.25 mg/kg IM, IV <sup>12</sup> 2-4 mg/kg IM, IV q24h × 3 days <sup>177</sup>  0.3-1.5 mg/kg IM, IV, IO <sup>59</sup>	Most species/shock (septic/traumatic) Most species/inflammatory, noninfectious respiratory disease Chelonians/hyperthermia
Dexamethasone sodium phosphate	0.1-0.25 mg/kg SC, IM, IV <sup>17</sup>	Most species/shock (septic/traumatic)
Insulin	1-5 U/kg IM, ICe q24-72h <sup>178</sup>  5-10 U/kg IM, ICe q24-72h <sup>178</sup>	Snakes, chelonians/doses adjusted based on response to therapy and serial blood glucose; ICe may take 24-48 h before response Lizards, crocodilians/see previous dosage com- ments within Insulin
Leuprolide acetate	0.4 mg/kg IM <sup>179</sup>	Iguanas/did not suppress testosterone levels in males
Levothyroxine	0.02 mg/kg PO q48h <sup>180</sup>  0.025 mg/kg q24h in morning <sup>181</sup>	Tortoises/hypothyroidism; stimulates feeding in debilitated tortoises Tortoises/monitor T <sub>4</sub> levels
Methylprednisolone	1 mg/kg IV q24h <sup>59</sup>	Chelonians/ivermectin toxicity
Nandrolone	0.5-5 mg/kg IM q7-28d <sup>182</sup> 1 mg/kg IM q7-28d <sup>183</sup>	Most species/hepatic lipidosis Lizards/anabolic steroid; reduces protein catabolism; may stimulate erythropoiesis
Oxytocin	—  1-10 U/kg IM <sup>31,184</sup>  2 U/kg IM q4-6h × 1-3 treatments <sup>185</sup> 1-5 U/kg IM, <sup>186</sup> repeat in 1 h  1-2, <sup>187</sup> 2-20, <sup>17,67</sup> or 10-20 <sup>188</sup> U/kg IM	Dystocias; results variable; works well in chelonians, less in snakes and lizards; give 1 h after Ca administration Most species/higher dose is commonly used; may be repeated up to three treatments at 90 min intervals, increasing dosage Most species Lizards/alternatively, 5 U/kg by slow IV or IO over 4-8 h <sup>186</sup> Chelonians
Prednisolone	2-5 mg/kg PO, IM <sup>133</sup> 0.5 mg/kg q24h × 14 days, then q48h until PCV stable <sup>57</sup>	Most species/analgesia (chronic pain) Lizards/autoimmune hemolytic anemia
Prednisolone sodium succinate	5-10 mg/kg IM, IV, <sup>119</sup> IO <sup>183</sup>	Most species/shock; hyperthermia; may help reduce nephrocalcinosis
Prednisone	0.5-1 mg/kg PO, SC, IM, IV <sup>189</sup>  0.8 mg/kg q48h <sup>190</sup>	Most species/lymphoma, leukemia, myeloprolif- erative disease Most species/chronic T-lymphocytic leukemia; combine with chlorambucil, but monitor uric acid levels
Stanozolol	5 mg/kg IM q7d prn <sup>17</sup>	Most species/anabolic steroid; management of catabolic disease states

Ca, Calcium; ICe, intracoelomic; IM, intramuscular; IO, intraosseous; IV, intravenous; NSHPT, nutritional secondary hyperparathyroidism; PCV, packed cell volume; PO, by mouth; prn, as required; SC, subcutaneous; T<sub>4</sub>, thyroxine.

TABLE 5

## Nutritional/Mineral/Fluid Support\*

Agent	Dosage	Species/Comments
Calcium glubionate	10 mg/kg PO q12-24h prn <sup>70</sup> 360 mg/kg (1 mL/kg) PO q12-24h prn <sup>39,174</sup>	All species/NSHPT Most species/NSHPT; hypocalcemia; dystocia; ensure adequate UVB exposure and proper nutrition
Calcium gluconate	100 mg/kg SC, IM, ICe <sup>191,192</sup> q6-24h <sup>39,174</sup>	Most species/hypocalcemia (low ionized Ca); when patient is stable switch to oral Ca
Calcium gluconate/borogluconate	10-50 mg/kg SC, IM <sup>70</sup>	Most species/hypocalcemia; hypocalcemic dystocia
Calcium glycerophosphate/Calcium lactate	1-5 mg/kg SC, IM <sup>70</sup> 10-25 mg/kg SC, IM <sup>17</sup> 10 mg/kg SC, IM, ICe q24h × 1-7 days <sup>5,174</sup>	Most species/hypocalcemia; hypocalcemic dystocia Most species/hypocalcemia; dystocia Lizards (iguanas)/hypocalcemia
Electrolyte solutions	Voluntary drinking (whole body soak) <sup>193</sup>  10-20 mL/kg via gavage or esophagostomy tube q24h <sup>193</sup>	All species/oral fluid therapy; early treatment of anorexia; dilute 1:1 with water; caution against drowning All species/rehydration; when stable; first stage in supplemental nutrition
Hydroxyethyl starch	3-5 mL/kg slow IV or IO bolus prn <sup>193,194</sup>	All species/hypoalbuminemia; hypovolemic perfusion deficits; increased capillary permeability; use with crystalloids; reduce crystalloid volume 40%-60%; max volume 20 mL/kg <sup>195</sup>
Iodine	2-4 mg/kg PO q24h × 14-21 days, then q7d <sup>17</sup>	Herbivorous species/iodine deficiency (goiter); suggested daily dietary iodine 0.03 mg/kg BW <sup>196</sup>
Iron dextran	12 mg/kg IM 1-2 ×/wk × 45 days <sup>197</sup>	Crocodylians/iron deficiency; in other species for anemia <sup>99</sup>
Lactated Ringer's solution (LRS)	15-40 mL/kg SC, IV, IO prn <sup>194</sup>	Land turtles/fluid replacement; use extracoeleomically after warming the patient; avoid lactate if hepatic insufficiency
Metronidazole	12.5-50 mg/kg PO <sup>11</sup> 50-100 mg/kg PO <sup>18</sup>	Most species/appetite stimulant Chameleons/appetite stimulant
Polymerized bovine hemoglobin	3-5 mL/kg slow IV or IO bolus prn <sup>193,194</sup>	All species/hemoglobin polymer; hypoalbuminemia; hemorrhage; severe anemia; hypovolemic perfusion deficits; increased capillary permeability; use with crystalloids; reduce crystalloid volume 40%-60%; max volume 20 mL/kg <sup>195</sup>
Replacement crystalloid solutions	10-30 mL/kg q24h or divided into 2-3 boluses several hours apart <sup>198</sup> 15-25 mL/kg/day PO, SC, IV, IO, ICe, EpiCe prn <sup>136</sup>	All species/ongoing regurgitation or severe diarrhea All species/replacement fluid therapy; warm to 29°C (84°F) <sup>194</sup>
Ringer's solution for reptiles:	10-20 mL/kg q24h <sup>199</sup>	All species/hypertonic dehydration or to prevent aminoglycoside nephrotoxicity
1 part Normosol-R + 2 parts 2.5% dextrose in 0.45% saline <sup>197</sup> or	15 (large reptiles) to 25 (small reptiles) mL/kg q24h or divided into 2 doses per day <sup>136</sup>	All species/hypertonic dehydration; warm fluids to 28°C (82°F)
1 part Normosol-R + 1 part 5% dextrose + 1 part 0.9% saline	20 mL/kg q12h <sup>39</sup>	Chelonians/severe dehydration
Selenium	0.028 mg/kg IM <sup>5</sup>	Lizards/deficiency; myopathy
Vitamin A	—  1000-5000 U/kg IM q7-10d × 4 treatments <sup>17</sup> 2000 U/30 g BW PO once, repeat in 7 days <sup>201,202</sup> 200-300 U/kg <sup>196</sup> SC, IM	Overdose causes epidermal sloughing; greater risk with aqueous parenteral formulation <sup>196,200,201</sup> ; may help infectious stomatitis Most species/hypovitaminosis A Chameleons/eye swelling, respiratory disease, hemipenile plugs, dysecdysis Turtles/hypovitaminosis A; give in conjunction with PO vitamin A (2-8 U/g feed DM)

TABLE 5—cont'd

## Nutritional/Mineral/Fluid Support\*

Agent	Dosage	Species/Comments
Vitamins A, D <sub>3</sub> , E	0.15 mL/kg IM, repeat in 21 days <sup>31</sup>	Most species/hypovitaminosis A, D <sub>3</sub> , or E; contains alcohol and may sting when given; alcohol-free product can be compounded commercially
	0.3 mL/kg PO, then 0.06 mL/kg q7d × 3-4 treatments <sup>188</sup>	Box turtles/hypovitaminosis A; parenteral use may result in hypervitaminosis A and D; given PO may enhance Ca uptake
Vitamin B complex	0.3 mL/kg SC, IM q24h <sup>99</sup>	Most species/anorexia; hypovitaminosis B
	25 mg thiamine/kg PO q24h × 3-7 days <sup>185</sup>	Most species/appetite stimulant; hypovitaminosis B
Vitamin B <sub>1</sub> (thiamine)	50-100 mg/kg PO, SC, IM q24h <sup>203</sup>	Piscivores/thiamine deficiency from thawed fish
	30 g/kg feed fish PO <sup>17</sup>	Crocodylians/treat or prevent deficiency
Vitamin B <sub>12</sub> (cyanocobalamin)	0.05 mg/kg SC, IM <sup>17</sup>	Snakes, lizards/appetite stimulant
Vitamin C	10-20 mg/kg SC, IM q24h <sup>204,205</sup>	All species/hypovitaminosis C; stomatitis; skin slough in snakes; supportive therapy for bacterial infections
Vitamin D <sub>3</sub>	—	NSHPT; hypocalcemia; deficiency and excess may lead to soft tissue calcification
	1000 U/kg IM, repeat in 1 wk <sup>39</sup>	Most species/deficiency; use with oral calcium gluconate and carbonate, general dietary management, and UVB irradiation
	200 U/kg PO, IM q7d <sup>5,174</sup>	Lizards/PO safer than IM
	400 U/kg IM q7d × 3 treatments <sup>175</sup>	Green iguanas/NSHPT; use with calcitonin after normocalcemic; also give PO calcium
Vitamin E/selenium	1 U vitamin E/kg <sup>196</sup> IM	Piscivores/hypovitaminosis E; myopathy, anorexia, swollen subcutaneous nodules
	50 U vit E/kg + 0.025 mg selenium/kg IM <sup>206</sup>	Lizards/hypovitaminosis E (vitamin E/selenium)
Vitamin K <sub>1</sub>	0.25-0.5 mg/kg IM <sup>31</sup>	Most species/hypovitaminosis K <sub>1</sub> ; coagulopathies

BW, Body weight; Ca, calcium; DM, dry matter; ICe, intracoelomic; IM, intramuscular; IO, intraosseous; IV, intravenous; NSHPT, nutritional secondary hyperparathyroidism; PO, by mouth; prn, as required; SC, subcutaneous; UVB, ultraviolet B.

\*See also Table 10.

TABLE 6

## Miscellaneous Agents

Agent	Dosage	Species/Comments
Allopurinol	10-20 mg/kg PO q24h <sup>67,183,207</sup>	Most species/gout; decreases production of uric acid; <sup>208</sup> long-term therapy; tortoises may respond best
	25 mg/kg PO q24h <sup>209</sup>	Green iguanas
	50 mg/kg PO q24h × 30 days, then q72h <sup>210</sup>	Chelonians/hyperuricemia
Aluminum hydroxide	100 mg/kg PO q12-24h <sup>208</sup>	Most species/hyperphosphatemia; decreases intestinal absorption of phosphorus; use cautiously in patients with gastric outlet obstruction
Amidotrizoate	7.5 mL/kg PO <sup>211</sup>	Tortoises/gastrointestinal contrast agent; give via gavage; mean transit times: 2.6 h at 30.6°C (87°F); 6.6 h at 21.5°C (71°F)
Aminophylline	2-4 mg/kg IM <sup>12</sup>	Most species/bronchodilator
Atropine	0.01-0.04 mg/kg IM, IV q8-24h <sup>212</sup>	Most species/dries up excess mucous secretions with infectious stomatitis
	0.1-0.2 mg/kg IM prn <sup>31</sup>	Most species/organophosphate toxicity prn
	0.2 mg/kg SC, IM <sup>213</sup>	Most species/respiratory distress associated with excessive secretions
Barium sulfate	5-20 mL/kg PO <sup>214</sup>	Most species/gastrointestinal contrast studies

Continued



TABLE 6—cont'd

## Miscellaneous Agents

Agent	Dosage	Species/Comments
Calcium EDTA	10-40 mg/kg IM q12h <sup>191</sup>	Most species/heavy metal chelation; ensure hydration
Carboplatin	2.5-5 mg/kg IV, IC <sup>189</sup>	Most species/chemotherapy
Chlorambucil	0.1-0.2 mg/kg PO <sup>189</sup>	Most species/chemotherapy
Cimetidine	4 mg/kg PO, IM q8-12h <sup>17</sup>	Most species/gastric and duodenal ulceration; esophagitis; gastroesophageal reflux; may use in renal failure to increase phosphate loss
Cisapride	0.5-2 mg/kg PO q24h <sup>30</sup>	Most species/motility modifier; GI stasis; may be compounded in the United States; ineffective in desert tortoises at 1 mg/kg <sup>215</sup>
Cisplatin	0.5-1 mg/kg IV (prehydrate), IC, intralesional (in oil) <sup>189</sup>	Most species/chemotherapy
Cyclophosphamide	10 mg/kg SC, IM, IV, IC <sup>189</sup>	Most species/chemotherapy
Diocetyl sodium sulfosuccinate	1-5 mg/kg PO <sup>216</sup>	Most species/constipation; use 1:20 dilution
Doxorubicin	1 mg/kg IV q7d × 2 treatments, then q14d × 2 treatments, then q21d × 2 treatments <sup>217</sup>	Snakes/chemotherapy
Furosemide	2-5 mg/kg PO, IM, IV q12-24h <sup>17,31,57-59</sup>	Most species/diuretic for edema and pulmonary congestion, while lacking loop of Henle, may effect via other mechanisms
Hydrochlorothiazide	1 mg/kg q24-72h <sup>183</sup>	Lizards/promotes diuresis; monitor hydration status
Iodine compound	500 mg/kg IV, IO <sup>183</sup>	Lizards/IV urography; take radiographs 0, 5, 15, 30, and 60 min postinjection
Iohexol	5-20 mL/kg PO <sup>89</sup>	Most species/gastrointestinal contrast studies, nonionic, organic iodine solution; good alternative to barium; <sup>214</sup> faster transit time than barium; can be diluted 1:1 with water
Lactulose	0.5 mL/kg PO q24h <sup>57,59,218</sup>	Lizards, chelonians/hepatic lipidosis
L-Asparaginase	400 U/kg SC, IM, IC <sup>187</sup>	Most species/chemotherapy
Melphalan	0.05-0.1 mg/kg PO <sup>187</sup>	Most species/chemotherapy
Methimazole	2 mg/kg q24h × 30 days <sup>200</sup>	Snakes/excessive shedding from hyperthyroidism; limited effectiveness
Methotrexate	0.25 mg/kg IV, SC, PO <sup>187</sup>	Most species/chemotherapy
Metoclopramide	0.06 mg/kg PO q24h × 7 day <sup>30,219</sup> 1-10 mg/kg PO q24h <sup>220</sup>	Most species/stimulates gastric motility Tortoises/stimulates gastric motility; ineffective in desert tortoises at 1 mg/kg <sup>215</sup>
Milk thistle ( <i>Silybum marianum</i> )	4-15 mg/kg PO q8-12h <sup>57,59</sup>	Lizards, chelonians/ hepatoprotectant
Pentobarbital	60-100 mg/kg IV, ICe <sup>221,222</sup>	Euthanasia
Pimobendan	0.2 mg/kg PO q24h <sup>57</sup>	Lizards
Potassium chloride	2 mEq/kg IV, ICe <sup>174</sup>	Most species/euthanasia; cardioplegic; administer after a euthanasia solution
Probenecid	250 mg/kg PO q12h <sup>223</sup>	Most species/gout; increases uric acid excretion; can be increased prn
S-Adenosylmethionine	30 mg/kg PO q24h <sup>224</sup>	Savannah monitors/liver disease
Sodium bicarbonate	0.5-1 mg/kg IV <sup>30</sup>	Most species/hypoxic acidosis postanesthesia
Sucralfate	500-1000 mg/kg PO q6-8h <sup>12</sup>	Most species/oral, esophageal, gastric, and duodenal ulcers
Tamoxifen 60-day time-release pellets	Pellets containing 5 mg tamoxifen implants, ICe <sup>225</sup>	Leopard geckos/inhibition of follicular development for 60 days if implanted before vitellogenesis
Vincristine	0.025 mg/kg IV <sup>189</sup>	Most species/chemotherapy

GI, Gastrointestinal; IC, intracavitary; ICe, intracoelomic; IM, intramuscular; IV, intravenous; PO, by mouth; prn, as required; SC, subcutaneous.

TABLE 7

Hematologic and Serum Biochemical Values<sup>a</sup>

Measurement	Boa constrictor (Boa constrictor) <sup>226-228</sup>	Emerald tree boa (Corallus caninus) <sup>228</sup>	Rainbow boa (Epicrates cenchria) <sup>228</sup>	Blood python (Python curtus) <sup>228</sup>	South Asian pythons (Python molurus ssp) <sup>228</sup>	Ball python (Python regius) <sup>228,229</sup>	Green tree python (Chondropython viridis) <sup>228</sup>	Jungle carpet python (Morelia spilota cheynei) <sup>228,230</sup>	Reticulated python (Python reticulatus) <sup>228</sup>
<b>HEMATOLOGY</b>									
PCV (%)	29 (10-45)	22.6 (6-57)	28 (11-40)	25 (15-49)	28 (18-36)	22 (10-30)	17 (8-27)	25 (10-46)	24 (18-30)
RBC (10 <sup>6</sup> /μL)	0.73 (0.16-2.1)	2.16 (0.54-5.05)	0.92 (0.34-1.74)	0.65	1.44 (0.65-6.9)	0.82 (0.12-1.31)	0.85 (0.4-1.3)	0.92 (0.54-1.3)	0.72 (0.41-1.25)
Hgb (g/dL)	9 (2.6-15.3)	8.2 (6.1-11.4)	10.6 (8-13.1)	—	9.4 (7-11)	8.0 (5.5-9.6)	5.9 (4-7)	9.3 (4-15.5)	10.7 (5.2-30)
MCV (fL)	393 (159-625)	237 (37-360)	292 (175-534)	340	275 (52-384)	381 (211-917)	229 (208-250)	282 (178-414)	343 (176-429)
MCH (pg)	132 (84-208)	120 (113-128)	160	—	93 (16-127)	82-139	100	114 (67-159)	138 (60-186)
MCHC (g/dL)	32 (22-42)	34 (30-36)	36 (33-40)	—	32 (29-35)	31 (25-40)	36 (33-40)	40 (24-53)	37 (29-45)
WBC (10 <sup>3</sup> /μL)	8.54 (0.88-22.6)	4.35 (0.48-10.6)	7.99 (1-35.2)	11.7 (1.13-42.5)	9.94 (2-19.8)	9.74 (1-26)	11.3 (3.5-22.1)	11.93 (1.72-34.1)	7.6 (1.8-17.7)
Heterophils (10 <sup>3</sup> /μL)	2.46 (0.21-12.3)	1.61 (0.18-5.36)	2.85 (0.03-10)	1.82 (0.31-3.99)	2.36 (0.42-6.84)	2.86 (0.37-10.80)	3.89 (0.86-6.63)	2.78 (0.29-11.3)	2.03 (0.5-4.02)
Lymphocytes (10 <sup>3</sup> /μL)	4.01 (0.16-18.5)	2.48 (0.14-8.27)	3.91 (0.1-32.4)	6.71 (0.34-33.6)	5.78 (0.34-18.6)	4.18 (0.13-14.10)	3.44 (0.21-11.2)	6.11 (0.6-19.7)	3.77 (0.42-11.9)
Monocytes (10 <sup>3</sup> /μL)	0.68 (0.02-6.55)	0.45 (0.06-2.12)	0.9 (0.03-3.06)	0.62 (0.13-2.12)	1.016 (0.06-5.01)	1.00 (0.01-3.20)	0.74 (0.04-2.35)	2.21 (0.06-8.79)	1.72 (0.11-6.76)
Azurophils (10 <sup>3</sup> /μL)	1.75 (0-5.76)	0.71 (0.1-1.84)	1.08 (0.11-4.44)	2.82 (0.27-6.8)	1.81 (0.26-4.8)	3.4 (0.3-13.26)	5.42 (0.97-13.9)	3.46 (0.09-18.04)	3.73 (1.07-6.88)
Eosinophils (10 <sup>3</sup> /μL)	0.26 (0.03-1.22)	0.07 (0.06-0.08)	0.11 (0.04-0.22)	0.08	0.45 (0.1-1.4)	0.25 (0.12-0.37)	0.16 (0.1-0.22)	0.19 (0.08-0.34)	0.68 (0.04-1.95)
Basophils (10 <sup>3</sup> /μL)	0.46 (0.03-2.77)	0.07 (0.03-0.14)	0.1 (0.02-0.27)	0.93 (0.32-1.83)	0.25 (0.08-1.08)	0.38 (0.05-1.12)	0.55 (0.07-1.8)	0.26 (0.05-1.76)	0.06 (0.06-0.7)
<b>CHEMISTRIES</b>									
ALP (U/L)	287 (43-1342)	128 (61-323)	27 (14-37)	44 (8-56)	105 (7-728)	61 (13-153)	209 (112-349)	40 (13-60)	64 (4-144)
ALT (U/L)	14 (0-37)	5 (1-10)	4 (1-6)	10 (3-17)	19 (0-40)	14 (5-26)	43 (8-132)	17 (6-38)	8 (0-29)
Amylase (U/L)	22 (5-67)	293 (160-470)	—	—	3255	2490 (1611-3368)	902 (564-1240)	—	1690 (416-2963)
AST (U/L)	35 (3-331)	30 (5-92)	43 (9-136)	56 (6-209)	22 (1-205)	55 (2-118)	27 (11-75)	16 (5-46)	34 (7-105)
Bilirubin, total (mg/dL)	0.2 (0-0.6)	0.2 (0.2-0.3)	0.4 (0-0.8)	0.3 (0.2-0.5)	0.6 (0-2)	0.4 (0-2.1)	0.2	0.5	0.3
BUN (mg/dL)	5 (0-34)	2 (0-4)	2 (1-3)	1 (0-2)	4 (1-14)	1 (0-3)	2 (0-2)	3 (2-3)	4 (0-12)
Calcium (mg/dL)	15.9 (9-27)	12.7 (10.3-17.4)	13.4 (11.3-18.9)	14.7 (13.5-16.2)	19 (9.9-34)	15.3 (10.8-22.2)	23.4 (10.3-80.1)	14.4 (12.8-16.5)	23.5 (8.6-78)
Chloride (mEq/L)	125 (108-139)	130 (119-138)	115 (94-128)	131 (123-138)	118 (101-135)	120 (109-130)	125 (119-130)	115 (109-123)	118 (104-129)
Cholesterol (mg/dL)	131 (34-314)	206 (116-343)	206 (140-314)	214 (76-445)	289 (146-445)	182 (23-302)	204 (116-360)	315 (264-386)	309 (257-356)
Creatine kinase (U/L)	526 (53-1728)	592 (157-985)	154 (31-745)	668 (327-1009)	494 (42-3093)	1318 (93-3108)	614	398 (27-1350)	1818 (356-8342)
Creatinine (mg/dL)	0.4 (0.1-1.6)	0.6 (0.4-0.9)	0.4 (0.1-0.7)	0.9 (0.5-1.3)	1.4 (0.1-16.9)	0.2 (0-0.5)	0.2 (0.2-0.5)	1.3 (0.3-3.7)	0.2 (0.1-0.4)
GGT (U/L)	4 (0-10)	2 (1-2)	5	8 (0-16)	9 (0-26)	5	78 (5-223)	31 (16-71)	38 (14-64)
Glucose (mg/dL)	37 (9-85)	27 (5-56)	17 (2-46)	30 (13-74)	24 (1-83)	23 (2-43)	—	32 (9-55)	22
Iron (μg/dL)	113 (103-122)	—	—	207 (49-364)	456 (20-5525)	371 (77-782)	206	306 ± 160	313 (43-1048)
LDH (U/L)	235 (16-877)	632 (76-1680)	401 (141-661)	4.9	—	—	—	330 (48-547)	—
Lipase (U/L)	2730	—	—	3.7 (3.1-4.5)	4.7 (2.9-9.1)	3.8 (0.9-7.2)	11.8 (4-26.3)	3.4 (2.1-6.1)	7.2 (2.5-14.58)
Magnesium (mEq/L)	2.95 (2.9-3)	—	—	6.3 (3.3-11.2)	4.8 (3.8-6.3)	7.1 (4.3-10.5)	5.3 (5.2-5.3)	6.1 (4.9-9)	6.6 (4.4-10.2)
Osmolarity (mOsm/L)	306	—	—	6.2 (3.6-8.1)	7.3 (4.2-10.5)	6.7 (3.2-10.5)	5.4 (3.9-6.9)	7.6 (5.9-10.9)	7.5 (6.1-9.8)
Phosphorus (mg/dL)	4.7 (2.6-11.7)	4.1 (1.8-8)	5.4 (3.8-7.7)	2.3 (1.6-2.8)	2.7 (1.7-3.9)	2.5 (1-8.4)	2 (1.3-2.7)	3.1 (2.8-3.3)	3.9 (1.9-7.2)
Potassium (mEq/L)	5.4 (3-10)	5 (3-8.7)	3.2 (1.2-5)	4.1 (3.1-4.9)	4.9 (2.4-6.7)	4.6 (2.1-9)	4.4 (3.4-5)	4.9 (4.2-5.5)	5.4 (3.7-8)
Protein, total (g/dL)	7 (4.3-10.8)	4.5 (2.6-7.2)	6.2 (3.7-8)	160 (155-164)	158 (151-165)	159 (146-173)	158 (157-159)	153 (150-158)	157 (142-169)
Albumin (g/dL) <sup>b</sup>	3.1 (1.9-5.3)	2.6 (2-3.6)	2.6 (1.8-3.6)	16 (13-22)	114 (16-532)	31	—	30	45
Globulin (g/dL) <sup>b</sup>	4 (2.2-6.9)	2.8 (1.8-3.6)	4.1 (2.8-5.8)	4.3 (2.1-7.1)	4.3 (1.3-18)	7.6 (1.1-23)	7.7 (1.6-20.4)	6.5 (1.9-23.7)	7.8 (3.5-17.4)
Sodium (mEq/L)	158 (130-171)	157 (148-167)	155 (137-163)	—	—	—	—	—	—
Triglyceride (mg/dL)	103 (3-457)	24 (10-49)	72 (64-90)	—	—	—	—	—	—
Uric acid (mg/dL)	4.7 (0-11.9)	4.7 (1.4-19.2)	7.2 (2.1-27.5)	—	—	—	—	—	—

Continued

TABLE 7—cont'd

Hematologic and Serum Biochemical Values<sup>a</sup>

Measurement	Gopher snake ( <i>Pituophis catenifer</i> ) <sup>228,231</sup>	Cornsnake ( <i>Elaphe guttata</i> ) <sup>228</sup>	Yellow rat snake ( <i>Elaphe obsoleta quadrivittata</i> ) <sup>228,232</sup>	Common kingsnake ( <i>Lampropeltis getula</i> ) <sup>228</sup>	Milk snake ( <i>Lampropeltis triangulum</i> ) <sup>228</sup>	Indigo snake ( <i>Drymarchon corais</i> ) <sup>228,233</sup>	Panther chameleon ( <i>Furcifer pardalis</i> ) <sup>228</sup>	Spiny- tailed lizard ( <i>Uromastix</i> spp) <sup>228,234</sup>	Blue-tongued skink ( <i>Tiliqua scincoides</i> ) <sup>228</sup>
<b>HEMATOLOGY</b>									
PCV (%)	25 (15-38)	32 (21-52)	24 (9-46)	29 (12-45)	26 (8-38)	24 (15-34)	27 (16-35)	29 (4.9-44.5)	31 (22-46)
RBC (10 <sup>6</sup> /μL)	0.67 (0.14-1.4)	1.21 (0.62-1.85)	0.77 (0.21-1.34)	3.11 (0.35-14)	0.89 (0.49-2)	0.62 (0.43-0.76)	0.83 (0.42-1.6)	0.78 (0.33-4.1)	1.1 (0.73-1.36)
Hgb (g/dL)	9.7 (4.3-12.3)	11.5 (9.7-13.5)	8.3 (2.8-15.2)	—	10.4 (6.9-11.9)	9.2 (7.3-11.1)	—	9.9 (3.3-17.4)	10.4 (6-13)
MCV (fL)	578 (246-1571)	315 (171-404)	361 (198-765)	304 (27.9-500)	377 (135-615)	369 (221-558)	330 (200-418)	415 (119-614)	299 (266-354)
MCH (pg)	111 (81-132)	127 (110-143)	121 (90-175)	—	119 (89-164)	258	—	133 (1.2-203)	98 (44-173)
MCHC (g/dL)	33 (27.5-36)	35 (32-40)	32 (26-54)	—	34 (29-45)	40 (33-46)	—	33 (22-41)	33 (16-57)
WBC (10 <sup>3</sup> /μL)	6.36 (1.56-11.5)	11.3 (1.02-31.4)	9.32 (0.37-25)	12.7 (1.46-42.2)	9.17 (1.25-24.7)	10.8 (2-26.9)	—	958 (290-2290)	—
Heterophils (10 <sup>3</sup> /μL)	1.62 (0.18-9.43)	2.7 (0.21-8.35)	1.93 (0.06-10.5)	1.63 (0.19-4.8)	1.14 (0.05-4.46)	1.96 (0.35-8.1)	6.18 (0.47-18.6)	3.1 (1-8.1)	7.3 (2.2-19.6)
Lymphocytes (10 <sup>3</sup> /μL)	3.39 (0.24-8.03)	5.61 (0.41-22.9)	4.05 (0.14-14.4)	7.5 (0.13-33.3)	5.92 (0.47-19.2)	6.26 (0.28-16.7)	1.57 (0.09-4.17)	2 (0.59-5.36)	2.45 (0.54-6.24)
Monocytes (10 <sup>3</sup> /μL)	0.88 (0.05-3.07)	0.93 (0.04-1.75)	1.5 (0.01-9.72)	0.8 (0.05-5.83)	1.15 (0.07-3.6)	1.5 (0.04-4.84)	2.96 (0.21-9.67)	0.99 (0.27-4.05)	2.75 (0.32-10.9)
Azurophils (10 <sup>3</sup> /μL)	0.94 (0.45-1.67)	2 (0.15-5.34)	3.05 (0.06-8.33)	3.26 (0.21-8.87)	1.58 (0.03-5.93)	1.58 (0.5-4.69)	—	0.04 (0-0.5)	0.84 (0.35-1.49)
Eosinophils (10 <sup>3</sup> /μL)	0.08 (0.05-0.12)	0.1 (0.08-0.12)	0.1 (0.01-0.22)	0.11 (0.02-0.22)	0.1 (0.01-0.25)	0.23 (0.13-0.43)	1.25 (0.08-3.74)	—	0.34 (0-1.08)
Basophils (10 <sup>3</sup> /vL)	0.15 (0.02-0.51)	0.55 (0.07-1.44)	0.26 (0.05-0.66)	0.42 (0.1-1.3)	0.33 (0.04-1.82)	0.4 (0.08-0.74)	—	0.04 (0-0.2)	1.5 (0.03-2.96)
							0.1 (0.07-0.13)	0.03 (0-0.33)	0.98 (0.11-2.24)
<b>CHEMISTRIES</b>									
ALP (U/L)	58 (9-170)	54 (23-75)	142 (37-858)	88 (23-152)	108 (70-168)	123 (80-161)	—	31 (5.9-139)	71 (39-101)
ALT (U/L)	22 (11-65)	34 (4-62)	13 (2-72)	18 (8-25)	8 (3-17)	10 (3-16)	—	11 (2.4-35)	20 (5-34)
Amylase (U/L)	344 (214-473)	458 (366-574)	1337 (630-2626)	848	665	—	—	134	—
AST (U/L)	53 (16-127)	43 (10-224)	29 (2-142)	54 (5-249)	46 (0-178)	46 (6-163)	10 (0-31)	73 (29-172)	50 (7-106)
Bilirubin, total (mg/dL)	0.4 (0.3-0.6)	0.7 (0.1-1)	0.2 (0.1-0.8)	0.4 (0.1-0.7)	0.4 (0.1-0.9)	2.1 (0.6-3.5)	—	0.3 (0.1-0.7)	—
BUN (mg/dL)	2.2 (1-5)	2 (0-6)	5 (0-20)	2 (0-4)	5 (2-14)	7 (0-22)	—	0.56 (0-3)	1 (0-2)
Calcium (mg/dL)	14.5 (11.1-17.6)	16.1 (13.5-19.6)	15.9 (11.3-73.2)	26.4 (12.3-60)	14.6 (12.4-17.6)	33 (12-59) <sup>c,297</sup>	10.3 (8.9-11.7)	9.9 (7.2-13.2)	14.2 (10.4-20.6)
Chloride (mEq/L)	134 (109-148)	124 (109-137)	118 (68-140)	115 (88-136)	127 (111-134)	119 (100-129)	—	126 (111-135)	113 (111-115)
Cholesterol (mg/dL)	265 (152-493)	433 (314-572)	377 (101-745)	409 (117-1083)	357 (154-563)	278 (116-397)	—	161 (64-295)	183 (72-429)
Creatine kinase (U/L)	669 (175-1763)	699 (91-2460)	221 (73-646)	500 (108-1112)	202 (92-332)	644 (68-1923)	—	1780 (141-10k)	2517 (73-5832)
Creatinine (mg/dL)	0.3 (0.1-0.6)	0.6 (0.2-2)	0.5 (0.2-1.3)	0.6 (0-1.6)	0.5 (0.3-1.1)	0.3 (0.2-0.3)	—	0.4 (0.1-3)	0.3 (0.1-0.6)
GGT (U/L)	10 (0-34)	5 (0-13)	9 (1-35)	9	8 (3-13)	15	—	8	8
Glucose (mg/dL)	88 (24-129)	56 (32-88)	66 (26-117)	46 (8-82)	54 (15-76)	46 (28-89)	294 (219-341)	200 (68-356)	127 (63-160)
Iron (μg/dL)	98	—	—	190 (30-488)	816 (18-2807)	313 (13-1055)	—	209 (22-899)	735 (364-1106)
LDH (U/L)	76 (20-191)	182 (48-444)	419 (4-5665)	6.2 (0.7-15.2)	8.6 (2.9-25)	7.4 (5.3-9) <sup>c,297</sup>	—	3.48 (2.1-10.2)	364
Lipase (U/L)	8 (5-17)	—	4 (3-4)	5.3 (2.8-9.2)	5.5 (3.3-9.7)	6.5 (3.8-14.3)	8.2 (5.5-9.8)	4.5 (1.3-10)	5.7 (2.8-7.7)
Magnesium (mg/dL)	3.2	—	2.5	7.5 (3.6-12)	6.8 (3.8-11.6)	8.6 (4.6-12.3)	—	3.7 (3-4.6)	5.7 (4.3-8.6)
Phosphorus (mg/dL)	4.1 (2.5-5.7)	4.5 (2.8-5.7)	3.7 (1.7-14.7)	2.9 (1.6-6.8)	2.3 (1.8-3)	2.5 (1.7-4.6)	5.8 (4.7-7.8)	4 (2.6-7.4)	6.2 (5.3-7.6)
Potassium (mEq/L)	4.7 (3.3-6.5)	6.7 (3.3-16.6)	4.7 (1.6-8.5)	4 (1.8-5.8)	5.3 (3.3-9)	3.8 (3.3-4.4)	2.5 (1.8-3.2)	2 (1.2-3.1)	2.2 (1.3-2.9)
Protein, total (g/dL)	5.3 (3.2-7.4)	6.8 (4.6-10.8)	6.4 (3.3-8.9)	163 (132-184)	167 (157-178)	157 (143-170)	3.5 (2.9-4.6)	2.9 (2.2-4.6)	4 (3.3-5.2)
Albumin (g/dL) <sup>b</sup>	2.4 (1.6-3.2)	3.6 (2.5-7.6)	2.3 (1.6-3.8)	149 (66-278)	428 (68-1620)	92 (76-118)	—	173 ± 4	149 (142-158)
Globulin (g/dL) <sup>b</sup>	3.1 (1.9-3.2)	3.9 (3.2-4.8)	4.1 (2.6-6.4)	5.7 (2.4-14.7)	6.3 (2.1-35.6)	8.6 (2.2-17.1)	—	175 (111-238)	—
Sodium (mEq/L)	171 (155-187)	165 (154-174)	164 (151-177)	—	—	—	9.4 (4-16.1)	2.94 (0.3-7.3)	3.8 (0.7-8.5)
Triglyceride (mg/dL)	27 (16-37)	331 (47-1118)	195 (21-1017)	—	—	—	—	—	—
Uric acid (mg/dL)	6.7 (2-17.6)	7.2 (2.8-19.9)	7.6 (1.6-47)	—	—	—	—	—	—

Measurement	Bearded dragon ( <i>Pogona vitticeps</i> ) <sup>228,235</sup>	Chinese (Asian) water dragon ( <i>Physignathus cocincinus</i> ) <sup>236</sup>	Green iguana ( <i>Iguana iguana</i> ) <sup>228,237-242</sup>	Green iguana ( <i>Iguana iguana</i> ) male <sup>e,243,244</sup>	Green iguana ( <i>Iguana iguana</i> ) female <sup>e,243,244</sup>	Green iguana ( <i>Iguana iguana</i> ) juvenile <sup>e,243</sup>	Prehensile-tailed skink ( <i>Corucia zebrata</i> ) <sup>228,245</sup>	Tegu lizard ( <i>Tupinambis</i> spp) <sup>f,228,246</sup>	Green basilisk ( <i>Basiliscus plumifrons</i> ) <sup>228</sup>
<b>HEMATOLOGY</b>									
PCV (%)	30 (19-40)	35 (32-40)	25-38	34 (29-39)	38 (33-44)	38 (30-47)	35 (24-60)	25 ± 2.6	36 (29-41)
RBC (10 <sup>6</sup> /μL)	0.97 (0.68-1.21)	—	1-1.9	1.3 (1-1.7)	1.4 (1.2-1.8)	1.4 (1.3-1.6)	1.5 (0.66-3.28)	0.96 ± 0.14	—
Hgb (g/dL)	9.5 (6.7-12)	—	8-12	8.6 (6.7-10.2)	10.6 (9.1-12.2)	9.6 (9.2-10.1)	9.6 (7.4-11.6)	11.4 ± 1.6	8.9 (8.6-9.1)
MCV (fL)	311 (236-397)	—	165-305	266 (228-303)	270 (235-331)	—	263 (152-600)	261 ± 23	—
MCH (pg)	108 (81-140)	—	65-105	25 (23-28)	28 (25-31)	—	69 (42-111)	119 ± 12.5	—
MCHC (g/dL)	34 (24-45)	—	20-38	15 (11-25)	15 (8-25)	16 (8-22)	28 (17-56)	45.6 ± 3.4	26 (22-29)
WBC (10 <sup>3</sup> /μL)	8.14 (1.99-23)	13.5 (11.7-18.2)	3-10	3.6 (1-5.4)	3.2 (0.6-6.4)	2.2 (1-3.8)	12.4 (3.9-22.4)	16.8 ± 2.5	16.8 (6.1-31)
Heterophils (10 <sup>3</sup> /μL)	2.17 (0.35-4.99)	5.1 (3.9-6.9)	0.35-5.2	9.7 (5-16.5)	9.9 (5.2-14.4)	12.9 (6.2-17.2)	4.4 (1.02-6.4)	2.2 ± 0.45	8.9 (2.8-17.7)
Lymphocytes (10 <sup>3</sup> /μL)	4.68 (0.57-17)	7.2 (5.6-9.5)	0.5-5.5	1.3 (0.2-2.7)	1.2 (0.4-2.3)	0.4 (0.3-0.6)	2.7 (0.3-4.7)	7.5 ± 0.58	6.4 (1.8-10.7)
Monocytes (10 <sup>3</sup> /μL)	0.7 (0.03-2.72)	1.1 (0.4-1.9)	0-0.1	0.1- (0-0.3)	0.1 (0-0.2)	0.3 (0-0.4)	0.1 (0-0.1)	1 ± 0.41	1.5 (0.12-4)
Azurophils (10 <sup>3</sup> /μL)	0.53 (0.04-1.84)	0 (0-0.6)	0-1.7	0.4 (0.1-1)	0.5 (0.2-1.2)	0.5 (0.1-0.7)	2.8 (0.4-4.8)	1.8 ± 0.56	—
Eosinophils (10 <sup>3</sup> /μL)	0.15 (0.06-0.27)	0.2 (0.1-0.3)	0-1	100 (100-200)	100 (100-300)	100 (100-300)	0.6 (0-3)	4.1 ± 0.11	—
Basophils (10 <sup>3</sup> /μL)	0.39 (0.05-1.01)	0.5 (0.2-0.8)	0-0.5	—	—	—	1.9 (0.1-4.3)	0.4 ± 0.01	1.5
Fibrinogen (mg/dL)	180 (0-300)	—	0-300	—	—	—	—	133 (0-200)	—
<b>CHEMISTRIES</b>									
ALP (U/L)	151 (15-447)	—	40 (4-170)	39 (14-65)	59 (22-90)	—	154 (44-334)	160 ± 85	137 (77-230)
ALT (U/L)	12 (5-20)	—	21 (0-97)	32 (4-76)	45 (5-96)	—	9 (2-23)	33 ± 24	13 (5-21)
Amylase (U/L)	497	—	1815 (996-2988)	22 (12-30)	29 (19-41)	—	19 (0-76)	18 ± 14	60 (14-136)
AST (U/L)	27 (0-77)	16.5 (8-52)	52 (2-100)	33 (19-65)	40 (7-102)	41 (13-72)	0.2 (0-0.9)	0.3 ± 0.2	0.6 (0.5-0.8)
Bile acids (rest; μmol/L)	—	—	7.5 (2.6-30.3)	0.8 (0.1-1.4)	1.5 (0.3-3.1)	—	1 (0-2)	1 ± 1	5 (1-20)
Bile acids (7.5h; μmol/L)	—	—	32.5 (15.2-44.1)	11.3 (8.6-14.1)	12.5 (10.8-14)	14.3 (12.1-23.2)	13 (11-21)	12.2 ± 0.8	10.5 (9.8-11.2)
Bilirubin, total (mg/dL)	0.5 (0-3.7)	—	0.3 (0-4.9)	119 (115-124)	121 (113-129)	—	124 (123-129)	121 ± 7	127 (125-129)
BUN (mg/dL)	3 (1-7)	—	2 (0-10)	161 (82-214)	255 (204-347)	—	144 (11-252)	206 ± 67	957 (550-1393)
Calcium (mg/dL)	11.8 (8.6-27)	12.4 (11.6-13.3)	12 (6-18) <sup>d</sup>	19.9 (15.2-24.7)	19 (16-23)	—	210 (27-940)	641 ± 568	5355 (2691-9436)
Ionized Ca <sup>++</sup> (mmol/L)	—	—	1.01-1.62	79 (36-162)	270 (81-512)	—	0.2 (0-0.7)	0.3 ± 0.1	0.4 (0.2-0.8)
Chloride (mEq/L)	130 (104-160)	—	117 (102-130)	166 (70-244)	170 (105-258)	273 (131-335)	0 (0-1)	7	—
Cholesterol (mg/dL)	513 (230-900)	—	104-333 <sup>d</sup>	5.3 (3.2-7.6)	6.3 (2.8-9.3)	7.7 (4.3-9)	100 (70-122)	128 ± 30	174 (60-280)
Creatine kinase (U/L)	1211 (59-7000)	1747 (19-6630)	1876(174-8768) <sup>d</sup>	4 (2.8-6.1)	3.6 (2-5.8)	—	236 (42-1139)	540 ± 537	—
Creatinine (mg/dL)	0.3 (0.0-0.6)	—	0.5 (0.2-1.3)	5.4 (4.4-6.5)	6.1 (4.9-7.6)	5 (4.2-6.1)	361 (335-373)	—	—
GGT (U/L)	2 (1-2)	—	3 (0-10)	2 (1.3-3)	2.4 (1.5-3)	2.3 (2-2.8)	—	—	—
Glucose (mg/dL)	210 (139-291)	157 (112-243)	169-288	3.5 (2.5-4.4)	3.8 (2.8-5.2)	2.7 (2.2-3)	3.7 (2.8-6.7)	5.6 ± 2.1	9.5 (6.2-11.6)
Iron (μg/dL)	—	—	88-133	0.6 (0.4-0.9)	0.7 (0.3-1)	0.8 (0.7-0.9)	3.6 (1.4-5)	2.4 ± 1.4	2.9 (2.4-3.5)
LDH (U/L)	304 (35-628)	—	617 (36-7424) <sup>d</sup>	157 (152-162)	163 (156-172)	—	5.9 (3.8-9)	6.6 ± 1.3	5.7 (4-8.3)
Lipase (U/L)	—	—	21 (17-24)	10.2 (2.2-15.7)	0.26 (0.07-0.35)	—	2.4 (1.3-3.6)	3.6 ± 0.7	3.1 (2.6-3.5)
Magnesium (mEq/L)	—	—	2.4-4	2.7 (1.5-5.8)	3.6 (0.9-6.7)	3.3 (0.7-5.7)	4.8 (3.5-8)	—	—
Phosphorus (mg/dL)	5.7 (2.7-15.1)	5.7 (3.4-8.2)	5 (2.5-21) <sup>d</sup>	—	—	—	3.1 (1.9-5.4)	2.9 ± 1.2	2.9 (2.1-4.9)
Potassium (mEq/L)	3.6 (1-6.5)	4.2 (3.8-4.5)	1.3-3	—	—	—	158 (145-167)	159 ± 4	163 (145-172)
Protein, total (g/dL)	5 (3.6-6.4)	7 (6.6-7.5)	5.4 (4.1-7.4) <sup>d</sup>	—	—	—	93 (28-319)	31	285 (9-712)
Albumin (g/dL) <sup>b</sup>	2.6 (1.3-4.6)	2.2 (2.1-2.3)	2.1-2.8	—	—	—	1.6 (<0.3-3.1)	3.2 ± 2	10.9 (1-96.6)
Albumin (PEP; g/dL) <sup>b</sup>	—	—	1.8 (1.4-3.1)	—	—	—	—	—	—
Globulin (g/dL) <sup>b</sup>	2.3 (1-4.4)	4.7 (4.5-5.3)	2.5-4.3 <sup>d</sup>	—	—	—	—	—	—
α (PEP; g/dL) <sup>b</sup>	—	—	0.9 (0.4-1.2)	—	—	—	—	—	—
β (PEP; g/dL) <sup>b</sup>	—	—	2.2 (1.6-3.8) <sup>d</sup>	—	—	—	—	—	—
γ (PEP; g/dL) <sup>b</sup>	—	—	0.3 (0.1-0.4)	—	—	—	—	—	—
A/G Ratio	—	—	0.5 (0.41-0.78)	—	—	—	—	—	—
Sodium (mEq/L)	153 (137-190)	150 (147-153)	158-183	—	—	—	—	—	—
Triglyceride (mg/dL)	261 (93-437)	—	383 (7-1323) <sup>d</sup>	—	—	—	—	—	—
Uric acid (mg/dL)	5.2 (1.6-11.4)	2.3 (1.9-2.7)	2.6 (0-8.2) <sup>d</sup>	—	—	—	—	—	—
Vit D <sub>3</sub> (25-OH; nmol/L)	—	—	51-393 <sup>d</sup>	—	—	—	—	—	—

Continued



TABLE 7—cont'd

Hematologic and Serum Biochemical Values<sup>a</sup>

Measurement	Savannah monitor ( <i>Varanus exanthematicus</i> ) <sup>228</sup>	Water monitor ( <i>Varanus salvator</i> ) <sup>228</sup>	American alligator ( <i>Alligator mississippiensis</i> ) <sup>228</sup>	Dwarf caiman ( <i>Paleosuchus palpebrosus</i> ) <sup>228</sup>	Eastern box turtle ( <i>Terrapene carolina</i> ) <sup>228,247,248</sup>	Ornate box turtle ( <i>Terrapene ornata</i> ) <sup>228</sup>	Radiated tortoise ( <i>Astrochelys radiata</i> ) <sup>228,249,250</sup>	Red-footed tortoise ( <i>Chelonoidis carbonaria</i> ) <sup>228</sup>	Indian star tortoise ( <i>Geochelone elegans</i> ) <sup>228</sup>
<b>HEMATOLOGY</b>									
PCV (%)	31.2 (21-51)	33.2 (20-44)	24.7 (12-38)	22 (16-28)	26 (21-32)	26 (18-37)	10-51	29 (18-47)	21 (12-31)
RBC (10 <sup>6</sup> /μL)	1.23 (0.63-1.58)	0.98 (0.42-1.42)	0.57 (0.38-1.2)	0.66 (0.43-0.89)	0.49 (0.09-0.89)	0.62 (0.46-0.8)	0.3-1.1	2.1 (0.47-6.3)	0.37 (0.24-0.55)
Hgb (g/dL)	10.5 (6.2-13.2)	10.5 (9.8-11.5)	7.7 (5.7-11.3)	7.7 (6.2-8.8)	5.1	7.2 (6-9)	5.6 (4-8)	7.5 (7-7.9)	7.9 (6.9-8.5)
MCV (fL)	284 (229-382)	335 (227-595)	453 (230-762)	362 (180-535)	420 (229-1000)	408 (350-463)	454 (319-571)	347 (71-468)	—
MCH (pg)	94 (89-99)	140 (104-177)	146 (105-202)	98	102	122 (108-136)	108 (82-133)	136 (123-149)	—
MCHC (g/dL)	32 (26-38)	33 (30-40)	32 (23-44)	33 (23-38)	28 (26-29)	33 (31-33)	28 (26-33)	31 (29-32)	27 (26-28)
WBC (10 <sup>3</sup> /μL)	5.07 (1.2-11.3)	11.1 (2.9-26)	8.33 (1.75-20.8)	6 (2.7-13.5)	10.8 (8-13)	5.8 (2-11.1)	2.5-14	7.1 (2.2-13.4)	11.02 (0.75-31)
Heterophils (10 <sup>3</sup> /μL)	1.95 (0.38-4.06)	5.58 (2.09-11.1)	3.85 (0.65-16.7)	3 (1.4-6.2)	4.8 (1.7-9) <sup>f</sup>	1.8 (0.1-4.5)	0.7-8	1.8 (0.09-6.4)	4.09 (0.17-14.9)
Lymphocytes (10 <sup>3</sup> /μL)	2.12 (0.43-5.25)	3.43 (0.3-10.1)	3.18 (0.36-12.1)	2.57 (0.18-10.1)	4.9 (2.6-8.2) <sup>f</sup>	2.9 (1.5-5.6)	0.4-5.8	3.3 (0.23-6.1)	5.14 (0.16-17.6)
Monocytes (10 <sup>3</sup> /μL)	1.64 (0.06-6.67)	0.81 (0.06-3.38)	0.37 (0.04-0.84)	0.13 (0.03-0.23)	0.11 (0-0.4) <sup>f</sup>	0.06 (0.04-0.07)	0.02-0.5	0.18 (0.04-0.67)	0.38 (0.02-1.35)
Azurophils (10 <sup>3</sup> /μL)	0.11	2.26 (0.42-5.98)	0.64 (0.04-3.54)	0.52 (0-1.3)	—	0.25 (0.06-0.67)	0-0.82	0.98 (0.29-2.5)	0.72 (0.08-1.73)
Eosinophils (10 <sup>3</sup> /μL)	—	0.1	0.29 (0.04-1.02)	0.1 (0.05-0.18)	0.02 (0-0.1) <sup>f</sup>	0.96 (0.07-4.4)	0.03-0.82	0.5 (0.02-2.3)	0.96 (0.08-2.24)
Basophils (10 <sup>3</sup> /μL)	0.15 (0.07-0.28)	0.11 (0.06-0.14)	0.71 (0.04-3.23)	0.16 (0.05-0.48)	0.21 (0-0.4) <sup>f</sup>	0.42 (0.07-0.94)	0.1-2.5	1.4 (0.05-5.5)	0.76 (0.04-3)
Fibrinogen (mg/dL)	156 (100-300)	500 (200-700)	267 ± 115	100 (0-200)	—	—	117 (100-200)	—	—
<b>CHEMISTRIES</b>									
ALP (U/L)	89 (5-675)	187 (35-410)	46 ± 29	11 (6-16)	62 (29-102)	26 (15-37)	72-392	84 (39-173)	174 (38-379)
ALT (U/L)	70 (7-374)	36 (4-138)	49 ± 42	45 (24-71)	7 (2-14)	30 (25-33)	0-17	18 (4-63)	5 (0-15) 1235
Amylase (U/L)	—	1021 (265-1868)	471 ± 335	533	160	—	—	—	87 (12-296)
AST (U/L)	22 (1-78)	38 (5-239)	314 ± 158	78 (17-139)	124 (2-620)	62 (33-201)	25-348	238 (97-616)	—
Bilirubin, total (mg/dL)	0.1 (0-0.3)	0.1 (0-0.3)	0.2 ± 0.2	0.2 (0-0.6)	0.5 (0.1-1)	0.3 (0.1-0.4)	0.3-31.3	—	0.2 (0-0.4)
BUN (mg/dL)	1 (0-5)	3 (0-6)	3 ± 3	1 (0-2)	49 (20-102)	47 (4-65)	0-0.5	0.5 (0.1-1.1)	4 (0-11)
Calcium (mg/dL)	14.6 (11.9-17.5)	14.4 (8.5-22)	11.2 ± 2.2	10.7 (9.7-11.4)	13.6 (6.5-26.4)	11.1 (8-13.6)	2-34	17 (4-53)	12 (9.3-17.6)
Chloride (mEq/L)	116 (105-124)	110 (100-128)	109 ± 21	113 (99-126)	106 (101-112)	108 (96-115)	8.6-18	12.7 (9.5-15.8)	104 (90-112)
Cholesterol (mg/dL)	116 (49-231)	94 (22-291)	128 ± 73	147 (68-344)	240 (65-496)	155 (125-185)	91-112	100 (89-119)	128 (77-252)
Creatine kinase (U/L)	764 (150-3048)	772 (176-1818)	2663 ± 2493	1984 (37-9890)	463 (37-898)	918 (88-3100)	56-154	146 (47-284)	1496 (144-8518)
Creatinine (mg/dL)	8.7 (0-67)	0.5 (0-1)	0.4 ± 0.2	0.3 (0-0.4)	0.4	1 (0.2-2.4)	33-5666	360 (43-996)	0.3 (0.2-0.5)
GGT (U/L)	7 (1-11)	24 (7-48)	6 ± 7	84 (29-187)	84 (33-153)	71 (8-113)	0.1-0.5	0.4 (0.2-1.3)	4 (0-5)
Glucose (mg/dL)	118 (90-159)	115 (42-215)	92 ± 42	221 (80-485)	206 (111-313)	362 (300-424)	5 (0-11)	28 (7-130)	109 (39-199)
Iron (μg/dL)	—	242 (111-429)	—	—	3.5	—	21-93	94 (17-171)	—
LDH (U/L)	427 (29-3699)	693 (34-3588)	522 ± 541	303 (301-304)	—	—	60	107	438 (12-863)
Lipase (U/L)	—	—	12	3.8 (3.2-5.1)	4 (1.6-8.2)	3.6 (2.7-4.8)	213-6444	401 (242-534)	5
Magnesium (mEq/L)	3.1	2.5 (2.2-2.7)	—	4.1 (3.7-4.5)	5.6 (3-9.7)	7.2 (6-8.8)	5-50	—	4.1 (2.7-5.7)
Osmolarity (mOsmol/L)	332 (319-345)	—	—	5.1 (3.6-6.9)	5.6 (2.7-7.5)	4.9 (2.8-7.6)	2.5-7	3.8 (1.8-5.8)	5.2 (3.9-5.9)
Phosphorus (mg/dL)	4.6 (3.1-7.5)	5.8 (3-10)	4.3 ± 1.6	1.4 (1.1-1.7)	2.2 (1.2-3.2)	2.3 (1.6-3.4)	3.1-5.8	5.4 (3.7-6.8)	4.7 (3.9-5.9)
Potassium (mEq/L)	4.2 (3.2-5.7)	4.6 (3.5-6.5)	4.1 ± 0.9	2.2 (1.8-2.5)	—	—	3-6.6	5.2 (3.3-7.4)	2.1 (1.5-3.1)
Protein, total (g/dL)	6.7 (4.2-8.6)	7.6 (5.2-11.9)	5.4 ± 1.2	4.1 (3.5-5.2)	3.4 (2.5-4.7)	2.7 (1.2-4.2)	0.6-2.4	1.9 (1.3-3.4)	—
Albumin (g/dL) <sup>b</sup>	2.1 (1.5-3.5)	2.4 (1.2-3.4)	1.7 ± 0.3	147 (140-153)	144 (138-149)	140 (132-160)	0.9-2.4	—	2.7 (2.3-3.1)
Albumin (PEP; g/dL) <sup>b</sup>	3.2 (3.1-3.3)	3.1 (3-3.2)	—	92 (9-174)	—	—	1.4-3.2	3.1 (2-5.3)	—
Globulin (g/dL) <sup>b</sup>	4.9 (3.4-6.6)	4.9 (3.2-7.3)	3.7 ± 0.9	2.4 (0.7-4.5)	1.6 (0.5-3.1)	2.2 (0.6-5.6)	0.1-0.5	—	—
α-1 (PEP; g/dL) <sup>b</sup>	—	0.1	—	—	—	—	0.6-1.9	—	—
α-2 (PEP; g/dL) <sup>b</sup>	—	0.9 (0.8-1)	—	—	—	—	0.6-1.5	—	—
β (PEP; g/dL) <sup>b</sup>	—	0.9	—	—	—	—	0.4-0.9	—	128 (122-133)
γ (PEP; g/dL) <sup>b</sup>	—	4.7 (2.6-6.8)	—	—	—	—	121-146	131 (116-155)	60 (27-110)
Sodium (mEq/L)	156 (149-165)	156 (142-176)	144 ± 26	—	—	—	26-303	246 (28-480)	3.3 (1-8.1)
Triglyceride (mg/dL)	135 (17-476)	35 (6-78)	241 ± 360	—	—	—	0.3 (0-0.6)	0.8 (0.3-1.3)	—
Uric acid (mg/dL)	6.7 (1.2-18)	5.8 (1-16.6)	1.7 ± 1.1	—	—	—	—	—	—

Measurement	Desert tortoise ( <i>Gopherus agassizii</i> ) <sup>228, 251-254</sup>	Gopher tortoise ( <i>Gopherus polyphemus</i> ) <sup>255</sup>	Russian tortoise ( <i>Testudo horsfieldii</i> ) <sup>256,257</sup>	African spurred tortoise ( <i>Centrochelys sulcata</i> ) <sup>228</sup>	Leopard tortoise ( <i>Stigmochelys pardalis</i> ) <sup>228</sup>	Galapagos tortoise ( <i>Chelonoidis nigra</i> ) <sup>228</sup>	Sliders ( <i>Trachemys scripta ssp</i> ) <sup>228, 258-260</sup>	Painted turtle ( <i>Chrysemys picta</i> ) <sup>228,259,260</sup>	Green sea turtle ( <i>Chelonia mydas</i> ) <sup>228,261</sup>
<b>HEMATOLOGY</b>									
PCV (%)	15-39	23 (15-30)	23 (22-34)	28 (9-48)	22 (7-60)	19 (11-27)	29 (25-33)	28 (24-30)	37 (27-49)
RBC (10 <sup>6</sup> /μL)	0.28-1.34	0.54 (0.24-0.91)	—	0.9 (0.43-1.28)	0.82 (0.58-1.06)	0.39 (0.28-0.6)	0.84 (0.33-2.21)	0.57 (0.41-0.68)	1.94
Hgb (g/dL)	3.6-10.3	6.4 (4.2-8.6)	—	10.2 (6.4-15.7)	16.1 (8.8-28)	6 (4.1-9.9)	11.1 (10-12.2)	11.2 (10.7-11.7)	10.7
MCV (fL)	197-688	—	—	386 (201-575)	279 (179-379)	528 (280-667)	409 (179-697)	271 (183-365)	155
MCH (pg)	39-189	—	—	133 (91-165)	83	169 (111-222)	108	—	55
MCHC (g/dL)	19-35	—	—	35 (24-62)	44 (42-46)	31 (25-37)	30	—	36
WBC (10 <sup>3</sup> /μL)	0.97-10.9	15.7 (10-22)	8.5 (5-12.5)	7.97 (1.2-25.6)	10.23 (1.3-23)	7.72 (0.3-33.1)	13.7 (1.2-25.5)	6.7 (1.2-12.9)	11.7 (2.9-18)
Heterophils (10 <sup>3</sup> /μL)	0.49-7.3	4.7 (1-12.5) <sup>a</sup>	3.7 (1.3-4.6)	3.44 (0.32-9.62)	4.25 (0.21-12.8)	2.82 (0.27-20.4)	5 (0.95-14)	2.2 (0.17-4.26)	3.5 (0.9-6.6)
Lymphocytes (10 <sup>3</sup> /μL)	0-3.8	8.9 (3.2-17.4) <sup>a</sup>	4.7 (3.6-7.6)	3.68 (0.19-13.7)	2.7 (0.72-6.6)	4.24 (0.09-25.5)	3.3 (0.25-7.88)	1.6 (0.23-3.46)	6.7 (1.2-10.1)
Monocytes (10 <sup>3</sup> /μL)	0-0.57	1.1 (0.3-2.9) <sup>a</sup>	0.01 (0-0.02)	0.64 (0.06-2.25)	0.38	0.16 (0.01-0.58)	0.24 (0.14-0.38)	0.26	0.62 (0.11-1.3)
Azurophils (10 <sup>3</sup> /μL)	0-0.9	—	0.05 (0.03-0.12)	0.25 (0.02-0.54)	—	0.36 (0.03-1.97)	0.58 (0.23-1.33)	0.51 (0.25-0.77)	1.2 (0.17-2.7)
Eosinophils (10 <sup>3</sup> /μL)	0-0.95	—	0.05 (0.02-0.06)	0.35 (0.03-1.41)	0.1 (0.09-0.12)	0.42 (0.02-5.18)	1.53 (0.17-5.9)	0.55 (0.09-1.03)	0.42 (0.08-1.08)
Basophils (10 <sup>3</sup> /μL)	0-4.3	0.94 (0.2-2.4) <sup>a</sup>	0.05 (0.02-0.08)	0.3 (0.02-0.54)	0.21 (0.18-0.24)	0.54 (0.01-2.72)	3.8 (0.31-8)	2.17 (0.04-4.86)	0.18 (0.08-0.34)
<b>CHEMISTRIES</b>									
ALP (U/L)	43-176	39 (11-71)	498 (181-1188)	36 (12-59)	121 (54-173)	96 (30-302)	395 (201-677)	208	6-67
ALT (U/L)	21 (0-66)	15 (2-57)	1 (0-2)	13 (1-71)	8	7 (0-28)	14 (1-66)	—	31 (5-82)
AST (U/L)	41-106	136 (57-392)	20 (12-32)	—	—	167 (20-523)	493 (411-535)	152	534
Bile acids (μmol/L)	0-5.4	—	—	108 (34-401)	110 (41-344)	56 (19-187)	183 (108-350)	0.1	74-245
Bilirubin, total (mg/dL)	0-0.9	0.02 (0-0.1)	0.015 (0-0.09)	0.4 (0-2.6)	0.1	0.3 (0-1)	0.2 (0.1-0.5)	37	0.03-0.2
BUN (mg/dL)	0-4	30 (1-130)	12 (4-17)	3 (0-6)	41 (6-72)	14 (2-42)	25 (4-54)	10 (8.8-11.4)	93 (54-154)
Calcium (total; mg/dL)	9.3-14.7	12 (10-14)	13.2 (9.9-19.5)	12.1 (6.6-20)	12.2 (9.1-20.4)	12.6 (4.8-41)	13.4 (10-18)	85	8-8.8
Ionized calcium (mmol/L)	—	—	1.28 (1-1.6)	106 (93-123)	—	96 (83-119)	102 (81-107)	—	101-121
Chloride (mEq/L)	94-112	102 (35-128)	—	160 (69-394)	189 (164-237)	225 (78-506)	162 (106-227)	—	221 (142-354)
Cholesterol (mg/dL)	56-233	76 (19-150)	109 (25-210)	1518 (291-6205)	359 (223-704)	1262 (216-8090)	3095 (392-4856)	—	326-2729
Creatine kinase (U/L)	2262 (944-3880)	160 (32-628)	123 (6-344)	0.3 (0.1-0.4)	0.6	0.2 (0.1-0.5)	0.3 (0.2-0.5)	76 (76-84)	0.1 (0-0.4)
Creatinine (mg/dL)	0.11-0.37	0.3 (0.1-0.4)	—	14 (3-19)	—	6 (0-10)	7 (0-21)	—	6 (0-21)
GLDH (U/L)	—	—	1 (0.6-1.5)	142 (54-277)	91 (48-132)	144 (38-445)	84 (52-138)	412	67-178
Glucose (mg/dL)	92-165	75 (55-128)	59 (40-86)	81 (80-82)	—	73 (8-593)	—	—	362 (117-600)
LDH (U/L)	25-250	273 (18-909)	—	1315 (258-1980)	446 (346-546)	719 (197-1884)	1713 (371-5763)	4.8	75-477
Magnesium (mEq/L)	2.1 (1.8-2.4)	4.1 (3.3-4.8)	—	3.9 (1.5-7.8)	4 (2.5-6.6)	4.6 (2-10.3)	6 (1-15)	3.1 (2.8-3.1)	—
Phosphorus (mg/dL)	1-6.3	2.1 (1-3.1)	2.6 (1.3-3.9)	5 (3.6-8.5)	6.7 (5.4-7.4)	5.3 (3.2-8.1)	2.2	2.9 (2.6-3.2)	4.8-12.2
Potassium (mEq/L)	3.5-4.7	5 (2.9-7.)	5.3 (1.9-7.2)	3.9 (1.6-7)	3.2 (1.6-5.9)	4.8 (2.4-8.3)	5.2 (3.7-7.5)	2.4 (2.4-4.4)	4.9-11.1
Protein, total (g/dL)	3-4.6	3.1 (1.3-4.6)	3 (2.5-4.6)	1.5 (0.6-2)	1.1 (0.5-1.5)	1.8 (0.8-3.1)	4.1 (3.5-4.7)	1.2	3-7.1
Albumin (g/dL) <sup>b</sup>	1.2-2.2	1.5 (0.5-2.6)	1.6 (1.2-2.3)	2 (1.0-2.9)	1.5 (1.1-1.8)	3.2 (1.2-5.7)	4.6 (2.8-6.1)	1.2	2.1-6.2
Albumin (PEP; g/dL) <sup>b</sup>	—	—	—	138 (121-155)	141 (135-149)	128 (114-141)	2 (1.6-2.5)	143	0.7-1.8

Continued

TABLE 7—cont'd

Hematologic and Serum Biochemical Values<sup>a</sup>

Measurement	Desert tortoise ( <i>Gopherus agassizii</i> ) <sup>228, 251-254</sup>	Gopher tortoise ( <i>Gopherus polyphemus</i> ) <sup>255</sup>	Russian tortoise ( <i>Testudo horsfieldii</i> ) <sup>256,257</sup>	African spurred tortoise ( <i>Centrochelys sulcata</i> ) <sup>228</sup>	Leopard tortoise ( <i>Stigmochelys pardalis</i> ) <sup>228</sup>	Galapagos tortoise ( <i>Chelonoidis nigra</i> ) <sup>228</sup>	Sliders ( <i>Trachemys scripta ssp</i> ) <sup>228, 258-260</sup>	Painted turtle ( <i>Chrysemys picta</i> ) <sup>228,259,260</sup>	Green sea turtle ( <i>Chelonia mydas</i> ) <sup>228,261</sup>
<b>CHEMISTRIES—cont'd</b>									
Globulin (g/dL) <sup>b</sup>	1.2-2.6	—	1.4 (1.3-2.3)	—	—	2 (1-2)	2.6 (1.2-3.7)	—	1.5-4.7
α-1 Glob (PEP; g/dL) <sup>b</sup>	1	—	—	163 (53-388)	—	271 (29-1345)	137 (124-144)	2 (1.2-2)	139-158
α-2 Glob (PEP; g/dL) <sup>b</sup>	1	—	—	—	—	29	304 (30-664)	—	492 (124-932)
β Glob (PEP; g/dL) <sup>b</sup>	0.6	—	—	5.2 (2.1-10.5)	3.2 (0.5-15.3)	2.3 (0.5-7.4)	1.2 (0.6-3.2)	—	1 (0.2-1.5)
γ Glob (PEP; g/dL) <sup>b</sup>	—	—	—	—	—	—	—	—	—
Sodium (mEq/L)	122-139	138 (127-148)	138 (131-149)	—	—	—	—	—	—
Triglyceride (mg/dL)	0-425	—	—	—	—	—	—	—	—
Uric acid (mg/dL)	2.7-7.2	3.5 (0.9-8.5)	1.2 (0.8-3.9)	—	—	—	—	—	—
Vitamin A (μg/mL)	0.2-0.6	—	—	—	—	—	—	—	—
Zinc (ppm)	0.4-3.7	—	—	—	—	—	—	—	—

*A/G ratio*, Albumin/globulin ratio; *ALP*, alkaline phosphatase; *ALT*, alanine aminotransferase; *AST*, aspartate aminotransferase; *BUN*, blood urea nitrogen; *GGT*, gamma-glutamyl transpeptidase; *Hgb*, hemoglobin; *LDH*, lactate dehydrogenase; *MCH*, mean corpuscular hemoglobin; *MCHC*, mean corpuscular hemoglobin concentration; *MCV*, mean corpuscular volume; *PCV*, packed cell volume; *PEP*, protein electrophoresis; *RBC*, red blood cell; *WBC*, white blood cell.

<sup>a</sup>Listed values are mean (min-max), unless a single value indicates n = 1 or a range that is not enclosed in parentheses indicates a reported reference range.

<sup>b</sup>Albumin is measured by colorimetry (e.g., bromocresol green), and globulin value is calculated unless otherwise indicated.

<sup>c</sup>Remarkably high reference ranges for Ca (mean, 159 mg/dL; range, 30-337 mg/dL) and P (mean, 35 mg/dL; range, 8-69) have also been reported.<sup>233</sup>

<sup>d</sup>Can be elevated in gravid females; vitamin D<sub>3</sub> is higher in female green iguanas.<sup>242</sup>

<sup>e</sup>These data were obtained from iguanas housed outdoors with unfiltered sunlight.

<sup>f</sup>Adults.

<sup>g</sup>Calculated from data.

TABLE 8

Environmental, Dietary, and Reproductive Characteristics of Reptiles<sup>137,199,262-267</sup>

Species	Environmental Preference		Diet <sup>d</sup>	Method of Reproduction <sup>e</sup>	Gestation/ Incubation Period (Days) <sup>f</sup>
	Temperature <sup>a-c</sup>	RH (%)			
SNAKES					
Ball (Royal) python ( <i>Python regius</i> )	25-30°C (77-86°F)	70-80 (use humidity box) <sup>g</sup>	C	Ov	90
Boa constrictor ( <i>Boa constrictor</i> )	28-34°C (82-93°F)	50-70 (use humidity box) <sup>g</sup>	C	V	120-240
Garter snake ( <i>Thamnophis sirtalis</i> )	22-30°C (72-86°F)	60-80 (use humidity box) <sup>g</sup>	C	V	90-110
King snake ( <i>Lampropeltis getula</i> )	23-30°C (73-86°F)	50-70 (use humidity box) <sup>g</sup>	Op/c	Ov	50-60
Sand boa ( <i>Eryx</i> sp)	25-30°C (77-86°F)	20-30	C	V	120-180
LIZARDS					
Bearded dragon ( <i>Pogona vitticeps</i> )	27-30°C (80-85°F)	— (use humidity box) <sup>g</sup>	I: young; H: adult	Ov	65-90
Green anole ( <i>Anolis carolinensis</i> )	23-29°C (73-84°F)	70-80	I/c	Ov	60-90
Green iguana ( <i>Iguana iguana</i> )	29-38°C (84-100°F)	60-80	H	Ov	73
Jackson's chameleon ( <i>Chamaeleo jacksonii</i> )	21-27°C (70-81°F)	50-70	I	V	90-180
Leopard ground gecko ( <i>Eublepharis macularius</i> )	25-30°C (77-86°F)	20-30 (use humidity box) <sup>g</sup>	I	Ov	55-60
Australian Water dragon ( <i>Physignathus lesueurii</i> )	25-34°C (77-93°F)	80-90; Need water with filter <sup>h</sup>	I/om	Ov	90
CHELONIANS					
Common box turtle ( <i>Terrapene carolina</i> )	24-29°C (75-84°F)	60-80 (use humidity box) <sup>g</sup>	C/f	Ov	50-90
Desert tortoise ( <i>Gopherus agassizii</i> )	25-30°C (77-86°F)	— (use humidity box) <sup>g</sup>	H	Ov	84-120
Greek tortoise ( <i>Testudo graeca</i> )	20-27°C (68-81°F)	30-50 (use humidity box) <sup>g</sup>	H/om	Ov	60
Painted turtle ( <i>Chrysemys picta</i> )	23-28°C (73-82°F)	80-90; Need water with filter <sup>h</sup>	H/I/o	Ov	47-99
Red-eared slider ( <i>Trachemys scripta elegans</i> )	22-30°C (72-86°F)	80-90; Need water with filter <sup>h</sup>	C	Ov	59-93
Russian tortoise ( <i>Testudo horsfieldii</i> )	21-32°C (70-90°F)	— (use humidity box) <sup>g</sup>	H	Ov	56-84
CROCODILIAN					
American alligator ( <i>Alligator mississippiensis</i> )	30-35°C (86-95°F)	80-90; Need water with filter <sup>h</sup>	C/p	Ov	62-65

C, Carnivorous; F, frugivorous; H, herbivorous; I, insectivorous; O, molluscivorous; Om, omnivorous; Op, ophiophagus; Ov, oviparous; P, piscivorous; RH, relative humidity; V, viviparous.

<sup>a</sup>Temperatures shown are ideal ambient daytime temperature gradients. These should be allowed to fall by approximately 5°C (9°F) during the night. "Hot-spot" temperatures should generally be 5°C (9°F) greater than the highest temperature shown.

<sup>b</sup>Preferred daytime temperature range for other commonly housed captive snakes are rosy boa (*Lichanura trivirgata*), 27-29.5°C (81-85°F); green tree python (*Morelia viridis*), 24-28°C (75-82°F); carpet python (*Morelia spilota*), 27-29.5°C (81-85°F); cornsnake (*Elaphe guttata*), 25-30°C (77-86°F); black rat snake (*Elaphe obsoleta*), 25-29°C (77-84°F); gopher/bullsnake (*Pituophis melanoleucus*), 25-29°C (77-84°F).

<sup>c</sup>Preferred daytime temperature range for other commonly housed captive lizards are day gecko (*Phelsuma* sp), 29.5°C (85°F); chameleons (montane) (*Chamaeleo* spp), 21-27°C (70-81°F); chameleons (lowland) (*Chamaeleo* spp), 27-29°C (81-84°F); blue-tongued skink (*Tiliqua* sp), 27-29.5°C (81-85°F); monitor lizards (*Varanus* spp), 29-31°C (84-88°F); tegu (*Tupinambis* spp), 27-30°C (81-86°F).

<sup>d</sup>Uppercase letters denote principal dietary requirements. Lowercase denotes secondary preference.

<sup>e</sup>Temperature-dependent.

<sup>f</sup>Can have long hatch times dependent on incubation parameters.

<sup>g</sup>This simulates humid underground burrow. Use dark colored plastic container with cut entrance, moistened paper towels, or sphagnum moss.

<sup>h</sup>Need to set up water component like fish tank with proper filter (use one for koi or turtles), pump, water quality testing, dechlorinator.



TABLE 9

Urinalysis Values of Chelonians<sup>267-270</sup>

Measurement	Normal Values	Abnormal Values
Specific gravity	1.003-1.014 (mean, 1.008)	Up to 1.034
pH	Herbivores: alkaline Omnivores: 5-8	Acidic <sup>a</sup>
Color	Colorless to pale yellow with white urates	Dark yellow, yellow-brown, yellow-green
Turbidity	Clear	Cloudy
Protein	Trace proteinuria	Increased proteinuria
Glucose	Glucosuria up to 30 mg/dL	Glucosuria can be higher than 50 mg/dL with anorexia
Renal casts	None	Various types present
Calcium, phosphorus, ammonia, urea, creatinine	Detectable in urine	Significantly increased in urine of <i>Testudo</i> spp. with renal disease
AST, CK, LDH	Detectable in urine	Significantly increased in urine of <i>Testudo</i> spp. with renal disease
Crystals	Amorphous urates/ammonium biurates	Many other crystals found in renal failure; uric acid crystals in gout; bilirubin and tyrosine crystals in liver disease

AST, Aspartate aminotransferase; CK, creatine kinase; LDH, lactate dehydrogenase.

<sup>a</sup>May be associated with hibernation, anorexia, and improper diet.

TABLE 10

Selected Products and Guidelines Used in Force-Feeding Anorectic or Debilitated Reptiles<sup>a,b</sup>

Agent	Guidelines	Species/Comments
Alfalfa pellets (e.g., iguana or rabbit pellets) or powder (Alfalfa Powder, NOW Foods)	Blend (1:4) with electrolyte solution or water; 20 mL/kg PO q48h (lizards) to q84h (chelonians) <sup>29,271</sup>	Herbivorous reptiles/administer via gavage; may clog feeding tube; for iguanas, may gavage equal volume of water on alternate days until patient is stable and eating <sup>271</sup> ; soaked pellets can also be hand-fed (especially by owner)
Baby foods	Vegetable; blend in with other food sources Meat (small amount); blend in with other food sources	Herbivorous reptiles/administer via gavage; for some species, some fruit baby food can be added Omnivorous species/administer via gavage
Dog/cat food, canned (a/d, Hill's; Maximum-Calorie Iams; Nutritional Recovery Formula, Eukanuba)	30 mL/kg PO q7-14d <sup>38,159</sup>	Carnivorous species/administer via gavage; although low protein (8.5%), some concern over high purine and vitamin A levels (probably OK unless concurrent renal disease); in dehydrated animals, dilute 1:1 with physiologic solution, oral human electrolyte solutions (pediatric: Pedialyte, Ross; Gatorade); once stabilized, small whole animals (lubricated with egg white) can be force-fed
Electrolyte solutions (Pedialyte, Ross; Gatorade)	15-25 mL/kg PO q24h	Most species
High protein powders (Carnivore Care, Oxbow Pet Products; Emerald Carnivore, Lafeber)	Mix as labeled, generally to pancake batter consistency; feed small amount to start (see bag suggestion) once daily	Insectivorous and carnivorous species/once reconstituted, can be mixed 1:1 with an alfalfa or timothy product for true omnivorous reptiles; administer via gavage
Timothy hay-based powder (Herbivore Critical Care Fine Grind, Oxbow Pet Products; Emerald Herbivore, Lafeber)	Mix as labeled, generally to pancake batter consistency; feed small amount to start (see bag suggestion) once daily	Herbivorous species/once reconstituted, can be mixed 1:1 with an alfalfa or timothy product for true omnivorous reptiles; administer via gavage

<sup>a</sup>General guidelines for force-feeding: generally provide nutrition after rehydration of patient; needs may vary with specific disease (e.g., low protein with renal disease); force-feeding volumes are frequently started at a low/modest level and gradually brought up to the desired level (for patients with severe disease/cachexia, transition should be very gradual); concurrent to force-feeding and hydrating a patient, highly palatable food items should be provided for voluntary food intake.

<sup>b</sup>Dietary fiber supplements (alfalfa pellets or powder; barley powder; purified cellulose) should be an integral part of enteral therapy for herbivorous reptiles.

TABLE 11

Selected Sources of Diets and Other Commercial Products for Reptiles<sup>a,b</sup>**FOODS AND SUPPLEMENTS**

Fluker Farms	800-735-8537	<a href="http://www.flukerfarms.com">www.flukerfarms.com</a>
Drs Foster & Smith	800-381-7179	<a href="http://www.drsfostersmith.com">www.drsfostersmith.com</a>
Kaytee	800-529-8331	<a href="http://www.kaytee.com">www.kaytee.com</a>
Mazuri	800-227-8941	<a href="http://www.mazuri.com">www.mazuri.com</a>
Oxbow Animal Health	800-249-0366	<a href="http://www.oxbowanimalhealth.com">www.oxbowanimalhealth.com</a>
Pretty Pets	800-356-5020	<a href="http://www.prettybird.com">www.prettybird.com</a>
Reliable Protein Products	480-361-3940	<a href="http://www.zoofood.com">www.zoofood.com</a>
Repashy Superfoods	855-737-2749	<a href="http://www.store.repashy.com">www.store.repashy.com</a>
Rep-Cal	800-406-6446	<a href="http://www.repcal.com">www.repcal.com</a>
San Francisco Bay Brand	510-792-7200	<a href="http://sfbb.com">http://sfbb.com</a>
Sticky Tongue Farms	951-244-3434	<a href="http://www.stickytonguefarms.com">www.stickytonguefarms.com</a>
Tetra Fauna	800-423-6458	<a href="http://www.tetra-fish.com">www.tetra-fish.com</a>
T-Rex Products	800-991-8739	<a href="http://www.t-rexproducts.com">www.t-rexproducts.com</a>
Zoo Med Laboratories	888-496-6633	<a href="http://www.zoomed.com">www.zoomed.com</a>
ZuPreem	800-345-4767	<a href="http://www.zupreem.com">www.zupreem.com</a>

**LIVE/FROZEN FOODS FOR CARNIVORES**

American Rodent Supply	317-899-1599	<a href="http://www.americanrodent.com">www.americanrodent.com</a>	Frozen mice, rats
Big Cheese Rodents	800-887-0921	<a href="http://www.bigcheeserodents.com">www.bigcheeserodents.com</a>	Frozen mice, rats
Carolina Mouse Farm	864-944-6192	—	Frozen mice, rats
The Gourmet Rodent	352-472-9189	<a href="http://www.gourmetrodent.com">www.gourmetrodent.com</a>	Frozen mice, rats, rabbits, chicks
Hoosier Mouse Supply	317-831-1219	<a href="http://www.hoosiermousesupply.com">www.hoosiermousesupply.com</a>	Live (local) and frozen mice, rats
Komodo Reptiles	914-788-8722	<a href="http://www.komodoreptiles.com">www.komodoreptiles.com</a>	Live (local) and frozen mice, rats
Mouse Factory	800-720-0076	—	Live and frozen mice, rats
Perfect Pets Inc	800-366-8794	<a href="http://www.perfectpet.net">www.perfectpet.net</a>	Frozen mice, rats, hamsters, gerbils, guinea pigs, rabbits, chicks
Rodent Pro	812-867-7598	<a href="http://www.rodentpro.com">www.rodentpro.com</a>	Frozen mice, rats, rabbits, guinea pigs, chicks, quail

**LIVE FOODS FOR INSECTIVORES**

Arbico Organics	800-827-2847	<a href="http://www.arbico-organics.com">www.arbico-organics.com</a>	Crickets, fly pupae, mealworms, waxworms, superworms
Bassett's Cricket Ranch	800-634-2445	<a href="http://www.bccricket.com">www.bccricket.com</a>	Crickets, mealworms
The Drosophila Co	954-227-3966	<a href="http://www.jtresser.com/flypaper.html">http://www.jtresser.com/flypaper.html</a>	Fruit flies
Fluker Farms	800-735-8537	<a href="http://www.flukerfarms.com">www.flukerfarms.com</a>	Crickets, mealworms, fruit flies, hissing cockroaches
Ghann's Cricket Farm	800-476-2248	<a href="http://www.ghann.com">www.ghann.com</a>	Crickets, soldier fly larvae
Grubco	800-222-3563	<a href="http://www.grubco.com">www.grubco.com</a>	Crickets, superworms, mealworms, fly larvae
Knutson's	800-248-9318	<a href="http://www.knutsonlivebait.com">www.knutsonlivebait.com</a>	Night crawlers, crickets, mealworms
Komodo Reptiles	914-788-8722	<a href="http://www.komodoreptiles.com">www.komodoreptiles.com</a>	Crickets, superworms, goliath worms, fruit flies, mealworms, nightcrawlers
Millbrook Cricket Farm	800-654-3506	<a href="http://www.millbrookcrickets.com">www.millbrookcrickets.com</a>	Crickets, mealworms, superworms
Mulberry Farms	760-731-6088	<a href="http://www.mulberryfarms.com">www.mulberryfarms.com</a>	Silkworm larvae
The Phoenix Worm Store	—	<a href="http://www.phoenixworm.com">www.phoenixworm.com</a>	Soldier fly larvae
Rainbow Mealworms	800-777-9676	<a href="http://www.rainbowmealworms.net">www.rainbowmealworms.net</a>	Crickets, mealworms, cockroaches
Russell's Cricket Farm	234-738-3663	<a href="http://www.livecrickets.com">www.livecrickets.com</a>	Crickets, mealworms, superworms
Sunshine Mealworms	800-322-1100	—	Mealworms, crickets, superworms
Top Hat Cricket Farm	800-638-2555	<a href="http://www.tophatcrickets.com">www.tophatcrickets.com</a>	Crickets, mealworms, superworms, hornworms, waxworms
Topline Whlse Dist Co.	888-922-0464	—	Nightcrawlers, redworms, hissing cockroaches

**LIGHTS**

Duro-Test Lighting	800-289-3876	—	Vita-Lite fluorescent
Fluker Farms	800-735-8537	<a href="http://www.flukerfarms.com">www.flukerfarms.com</a>	Incandescent, heat
General Electric	800-435-4448	<a href="http://www.gelighting.com">www.gelighting.com</a>	Incandescent, heat
Hikari Sales USA	800-621-5619	<a href="http://www.hikariusa.com">www.hikariusa.com</a>	UVB fluorescent
Mac Industries, Inc	252-241-4584	<a href="http://www.reptileuv.com">www.reptileuv.com</a>	Mega-ray UVB
Philips	800-555-0050	<a href="http://www.lighting.philips.com">www.lighting.philips.com</a>	Incandescent, heat
Sylvania	978-777-1900	<a href="http://www.sylvania.com">www.sylvania.com</a>	350BL blacklights
T-Rex Products	800-991-8739	<a href="http://www.t-rexproducts.com">www.t-rexproducts.com</a>	Mercury vapor UVB, incandescent, heat

Continued

TABLE 11—cont'd

Selected Sources of Diets and Other Commercial Products for Reptiles<sup>a,b</sup>

Zilla	888-255-4527	<a href="http://www.zilla-rules.com">www.zilla-rules.com</a>	Incandescent, heat, UVB fluorescent
Zoo Med Laboratories	888-496-6633	<a href="http://www.zoomed.com">www.zoomed.com</a>	Incandescent, heat, mercury vapor UVB, fluorescent UVB
<b>HEATING DEVICES</b>			
Avitec	800-646-2473	<a href="http://www.avitec.com">www.avitec.com</a>	Conical ceramic heat emitters, thermostats
The Bean Farm	877-708-5882	<a href="http://www.beanfarm.com">www.beanfarm.com</a>	Heat tape, heat pads
Big Apple Pet Supply	800-922-7753	<a href="http://www.bigappleherp.com">www.bigappleherp.com</a>	Thermostats
Fluker Farms	800-735-8537	<a href="http://www.flukerfarms.com">www.flukerfarms.com</a>	Under-cage heat pads
Helix Controls	760-726-4464	<a href="http://www.helixcontrols.com">www.helixcontrols.com</a>	Thermostats, heat tape, heat panels
LLL Reptile	888-547-3784	<a href="http://lllreptile.com">http://lllreptile.com</a>	Pearlco conical ceramic heat emitters
Zilla	888-255-4527	<a href="http://www.zilla-rules.com">www.zilla-rules.com</a>	Conical ceramic heat emitters, thermostats
Zoo Med Laboratories	888-496-6633	<a href="http://www.zoomed.com">www.zoomed.com</a>	Thermostats, rheostats, heat pads, tape, cables
<b>HUMIDITY DEVICES</b>			
Exo Terra (Hagen)	800-724-2436	<a href="http://www.exo-terra.com">www.exo-terra.com</a>	Ultrasonic fogger, Monsoon rainfall
Humidifirst	561-752-1936	<a href="http://www.humidifirst.com">www.humidifirst.com</a>	Mist Pac ultrasonic humidifiers
Zoo Med Laboratories	888-496-6633	<a href="http://www.zoomed.com">www.zoomed.com</a>	Ultrasonic fogger, Repti fogger, Habba mist, Hygro-Therm humidity controller
<b>ENVIRONMENTAL SENSING AND MONITORING DEVICES</b>			
Exo Terra (Hagen)	800-724-2436	<a href="http://www.exo-terra.com">www.exo-terra.com</a>	Remote digital thermometers, hygrometers
FLIR	866-477-3687	<a href="http://www.flir.com">www.flir.com</a>	Infrared thermography
Onset Computer Corp.	800-564-4377	<a href="http://www.onsetcomp.com">www.onsetcomp.com</a>	HOB0 data loggers
Raytek	800-227-8074	<a href="http://www.raytek.com">www.raytek.com</a>	Digital infrared thermometer
Solartech	800-798-3311	<a href="http://www.solarmeter.com">www.solarmeter.com</a>	Solarmeter 6.2 UVB meter
Zilla	888-255-4527	<a href="http://www.zilla-rules.com">www.zilla-rules.com</a>	Digital infrared thermometer
Zoo Med Laboratories	888-496-6633	<a href="http://www.zoomed.com">www.zoomed.com</a>	Hygro-Therm humidity/heat monitor and controller

<sup>a</sup>Many pet stores sell live and frozen food for reptiles, as well as many of the products listed.<sup>b</sup>Numerous sources of information were used to compile this table, particularly Internet sources.

## REFERENCES

- Gaio C, Rossi T, Villa R, et al. Pharmacokinetics of acyclovir after a single oral administration in marginated tortoises, *Testudo marginata*. *J Herpetol Med Surg* 2007;17:8-11.
- McArthur S. Problem solving approach to common diseases of terrestrial and semi-aquatic chelonians. In: McArthur S, Wilkinson R, Meyer J, eds. *Medicine and surgery of tortoises and turtles*. Oxford, UK: Blackwell Publishing Ltd, 2004;309-377.
- Johnson JH, Jensen JM, Brumbaugh GW, et al. Amikacin pharmacokinetics and the effects of ambient temperature on the dosage regimen in ball pythons (*Python regius*). *J Zoo Wildl Med* 1997;28:80-88.
- Mader DR, Conzelman GM, Baggot JD. Effects of ambient temperature on half life and dosage regimen of amikacin in the gopher snake. *J Am Vet Med Assoc* 1985;187:1134-1136.
- Barten SL. The medical care of iguanas and other common pet lizards. *Vet Clin North Am Small Anim Pract* 1993;23:1213-1249.
- Stahl SJ. Pet lizard conditions and syndromes. *Semin Avian Exot Pet Med* 2003;12:162-182.
- Whitaker BR, Krum H. Medical management of sea turtles in aquaria. In: Fowler ME, Miller ME, eds. *Zoo & wild animal medicine: current therapy*. 4th ed. Philadelphia: WB Saunders Co, 1999;217-231.
- Caligiuri R, Kollias GV, Jacobson E, et al. The effects of ambient temperature on amikacin pharmacokinetics in gopher tortoises. *J Vet Pharm Therapeut* 1990;13:287-291.
- Jacobson ER, Brown MP, Chung M, et al. Serum concentration and disposition kinetics of gentamicin and amikacin in juvenile American alligators. *J Zoo Anim Med* 1988;19:188-194.
- Diver SJ. Empirical doses of antimicrobial drugs commonly used in reptiles. *Exot DVM* 1998;1:23.
- Frye FL. *Reptile care: an atlas of diseases and treatments*. Neptune City, NJ: TFH Publications, Inc, 1991;1-637.
- Frye FL. *Reptile clinician's handbook*. Malabar, Fla: Krieger Publishing, 1994.
- Jenkins JR. A formulary for reptile and amphibian medicine, in *Proceedings*. 4th Annu Avian/Exot Anim Med Symp, University of California, Davis, 1991;24-27.
- Hernandez-Divers SJ. Pulmonary candidiasis caused by *Candida albicans* in a Greek tortoise (*Testudo graeca*) and treatment with intrapulmonary amphotericin B. *J Zoo Wildl Med* 2001;32:352-359.
- Lloyd M. Crocodilia. In: Fowler ME, Miller RE, eds. *Zoo and wild animal medicine*. 5th ed. Philadelphia: Elsevier Saunders; 2003;59-70.
- Jacobson ER. Use of chemotherapeutics in reptile medicine. In: Jacobson ER, Kollias GV Jr, eds. *Exotic animals*. New York: Churchill Livingstone, 1988;35-48.
- Gauvin J. Drug therapy in reptiles. *Semin Avian Exot Pet Med* 1993;2:48-59.
- Jenkins JR. Husbandry and diseases of Old World chameleons. *J Small Exot Anim Med* 1992;1:166-171.

19. Page CD, Mautino M. Clinical management of tortoises. *Compend Cont Educ Pract Vet* 1990;12:79-85.
20. Funk RS. A formulary for lizards, snakes, and crocodilians. *Vet Clin North Am Exot Anim Pract* 2000;3:333-358.
21. Spörl H, Gobel T, Schildger B. Blood levels of some anti-infectives in the spur-thighed tortoise (*Testudo hermanni*), in *Proceedings*. 4th Intl Colloq Path Med Rept Amphib 1991;120-128.
22. Coke RL, Hunter RP, Isaza R, et al. Pharmacokinetics and tissue concentrations of azithromycin in ball pythons (*Python regius*). *Am J Vet Res* 2003;64:225-228.
23. Holz PH, Burger JP, Baker R, et al. Effect of injection site on carbenicillin pharmacokinetics in the carpet python, *Morelia spilota*. *J Herpetol Med Surg* 2002;12:12-16.
24. Lawrence K, Needham JR, Palmer GH, et al. A preliminary study on the use of carbenicillin in snakes. *J Vet Pharmacol Therapeut* 1984;7:119-124.
25. Lawrence K, Palmer GH, Needham JR. Use of carbenicillin in 2 species of tortoise (*Testudo graeca* and *Testudo hermanni*). *Res Vet Sci* 1986;40:413-415.
26. White SD, Bourdeau P, Bruet V, et al. Reptiles with dermatological lesions: a retrospective study of 301 cases at two university veterinary teaching hospitals (1992-2008). *Vet Dermatol* 2010;22:150-161.
27. Lawrence K, Muggleton PW, Needham JR. Preliminary study on the use of ceftazidime, a broad spectrum cephalosporin antibiotic, in snakes. *Res Vet Sci* 1984;36:16-20.
28. Stamper MA, Papich MG, Lewbart GA, et al. Pharmacokinetics of ceftazidime in loggerhead sea turtles (*Caretta caretta*) after single intravenous and intramuscular injections. *J Zoo Wildl Med* 1999;30:32-35.
29. Benson KG, Tell LA, Young LA, et al. Pharmacokinetics of ceftiofur sodium after intramuscular or subcutaneous administration in green iguanas (*Iguana iguana*). *Am J Vet Res* 2003;64:1278-1282.
30. Stein G. Reptile and amphibian formulary. In: Mader DR, ed. *Reptile medicine and surgery*. Philadelphia: WB Saunders Co, 1996;465-472.
31. Jenkins JR. Medical management of reptile patients. *Compend Cont Educ Pract Vet* 1991;13:980-988.
32. Bush M, Smeller JM, Charache PN, et al. Preliminary study of antibiotics in snakes, in *Proceedings*. Annu Conf Am Assoc Zoo Vet 1976;50-54.
33. Clark CH, Rogers ED, Milton JL. Plasma concentrations of chloramphenicol in snakes. *Am J Vet Res* 1985;46:2654-2657.
34. Klingenberg RJ. Therapeutics. In: Mader DR, ed. *Reptile medicine and surgery*. Philadelphia: WB Saunders Co, 1996;299-321.
35. Johnson JD, Mangone B, Jarchow JL. A review of mycoplasmosis infections in tortoises and options for treatment, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 1998;89-92.
36. Wimsatt J, Johnson J, Mangone BA, et al. Clarithromycin pharmacokinetics in the desert tortoise (*Gopherus agassizii*). *J Zoo Wildl Med* 1999;30:36-43.
37. Marin P, Bayon A, Fernandez-Varon E, et al. Pharmacokinetics of danofloxacin after single dose intravenous, intramuscular, and subcutaneous administration to loggerhead turtles *Caretta caretta*. *Dis Aquat Organ* 2008;82:231-236.
38. Mader DR: Pers. com., 2011.
39. Boyer TH. *Essentials of reptiles: a guide for practitioners*. Lakewood, Colo: AAHA Press, 1998;1-253.
40. Young LA, Schumacher J, Papich MG, et al. Disposition of enrofloxacin and its metabolite ciprofloxacin after intravascular injection in juvenile Burmese pythons (*Python molurus bivittatus*). *J Zoo Wildl Med* 1997;28:71-79.
41. Maxwell LK, Jacobson ER. Preliminary single-dose pharmacokinetics of enrofloxacin after oral and intramuscular administration in green iguanas (*Iguana iguana*), in *Proceedings*. Annu Conf Am Assoc Zoo Vet 1997;25.
42. Hungerford C, Spelman L, Papich M. Pharmacokinetics of enrofloxacin after oral and intramuscular administration in savannah monitors (*Varanus exanthematicus*), in *Proceedings*. Annu Conf Am Assoc Zoo Vet 1997;89-92.
43. Prezant RM, Isaza R, Jacobson ER. Plasma concentrations and disposition kinetics of enrofloxacin in gopher tortoises (*Gopherus polyphemus*). *J Zoo Wildl Med* 1994;25:82-87.
44. Casares M, Enders F. Enrofloxacin side effects in a Galapagos tortoise (*Geochelone elephantopus nigra*), in *Proceedings*. Annu Conf Am Assoc Zoo Vet 1996;446-448.
45. Raphael BL, Papich M, Cook RA. Pharmacokinetics of enrofloxacin after a single intramuscular injection in Indian star tortoises (*Geochelone elegans*). *J Zoo Wildl Med* 1994;25:88-94.
46. Helmick KE, Papich MG, Vliet KA, et al. Pharmacokinetics of enrofloxacin after single-dose oral and intravenous administration in the American alligator (*Alligator mississippiensis*). *J Zoo Wildl Med* 2004;35:333-340.
47. Martelli P, Lai OR, Krishnasamy K, et al. Pharmacokinetic behavior of enrofloxacin in estuarine crocodile (*Crocodylus porosus*) after single intravenous, intramuscular, and oral doses. *J Zoo Wildl Med* 2009;40:696-704.
48. Wissman MA, Parsons B. Dermatophytosis of green iguanas (*Iguana iguana*). *J Small Exot Anim Med* 1993;2:133-136.
49. Harms CA, Lewbart GA, Beasley J. Medical management of mixed nocardial and unidentified fungal osteomyelitis in a Kemp's Ridley sea turtle, *Lepidochelys kempii*. *J Herpetol Med Surg* 2002;12:21-26.
50. Mallo KM, Harms CA, Lewbart GA, et al. Pharmacokinetics of fluconazole in loggerhead sea turtles (*Caretta caretta*) after single intravenous and subcutaneous injections, and multiple subcutaneous injections. *J Zoo Wildl Med* 2002;33:29-35.
51. Montali RJ, Bush M, Smeller JM. Pathology of nephrotoxicity of gentamicin in snakes: model for reptilian gout. *Vet Pathol* 1979;16:108-115.
52. Bush M, Smeller JM, Charache P, et al. Biological half-life of gentamicin in gopher snakes. *Am J Vet Res* 1978;39:171-173.
53. Hilf M, Swanson D, Wagner R, et al. A new dosing schedule for gentamicin in blood pythons (*Python curtus*): a pharmacokinetic study. *Res Vet Sci* 1991;50:127-130.
54. Beck K, Loomis M, Lewbart G, et al. Preliminary comparison of plasma concentrations of gentamicin injected into the cranial and caudal limb musculature of the eastern box turtle (*Terrapene carolina carolina*). *J Zoo Wildl Med* 1995;26:265-268.
55. Raphael B, Clark CH, Hudson RJr. Plasma concentration of gentamicin in turtles. *J Zoo Anim Med* 1985;16:136-139.
56. Rossi J. Practical reptile dermatology, in *Proceedings*. North Am Vet Conf 1995;648-649.
57. Jepson L. Lizards. In: *Exotic animal medicine: a quick reference guide*. Philadelphia: Saunders/Elsevier, 2009;268-314.
58. Jepson L. Snakes. In: *Exotic animal medicine: a quick reference guide*. Philadelphia: Saunders/Elsevier, 2009;315-357.
59. Jepson L. Turtles and tortoises. In: *Exotic animal medicine: a quick reference guide*. Philadelphia: Saunders/Elsevier, 2009;358-411.
60. Martel A, Hellebuyck T, Van Waeyenbergh L. Treatment of infections with *Nannizziopsis vriesii*, an emergent reptilian dermatophyte, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2009;69-70.
61. Mitchell M. Ophidia. In: Fowler ME, Miller RE, eds. *Zoo and wild animal medicine*. 5th ed. Philadelphia: Saunders/Elsevier, 2003;82-91.
62. Heatley J, Mitchell M, Williams J, et al. Fungal periodontal osteomyelitis in a chameleon, *Furcifer pardalis*. *J Herpetol Med Surg* 2001;11:7-12.
63. Bicknese E, Pessier A, Boedeker N. Successful treatment of fungal osteomyelitis in a Parson's chameleon (*Calumma parsonii*) using surgical and anti-fungal treatments, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2008;86.
64. Gamble KC, Alvarado TP, Bennett CL. Itraconazole plasma and tissue concentrations in the spiny lizard (*Sceloporus* sp.) following once-daily dosing. *J Zoo Wildl Med* 1997;28:89-93.
65. Manire CA, Rhinehart HL, Pennick GJ, et al. Steady-state plasma concentrations of itraconazole after oral administration in Kemp's ridley sea turtles, *Lepidochelys kempi*. *J Zoo Wildl Med* 2003;34:171-178.



66. Jacobson ER. Antimicrobial drug use in reptiles. In: Prescott JF, Baggot JD, eds. *Antimicrobial therapy in veterinary medicine*. Ames: Iowa State University Press, 1993;543-552.
67. Mautino M, Page CD. Biology and medicine of turtles and tortoises. *Vet Clin North Am Small Anim Pract* 1993;23:1251-1270.
68. Page CD, Mautino M, Derendorf H, et al. Multiple dose pharmacokinetics of ketoconazole administered orally to gopher tortoises (*Gopherus polyphemus*). *J Zoo Wildl Med* 1991;22:191-198.
69. Coke RL, Isaza R, Koch DE, et al. Preliminary single-dose pharmacokinetics of marbofloxacin in ball pythons (*Python regius*). *J Zoo Wildl Med* 2006;37:6-10.
70. Funk RS, Diethelm G. Reptile formulary. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St. Louis: Saunders/Elsevier, 2006;1119-1139.
71. Bodri MS, Rambo TM, Wagner RA, et al. Pharmacokinetics of metronidazole administered as a single oral bolus to red rat snakes, *Elaphe guttata*. *J Herpetol Med Surg* 2006;16:15-19.
72. Kolmstetter CM, Cox S, Ramsay EC. Pharmacokinetics of metronidazole in the yellow rat snake, *Elaphe obsoleta quadrivittata*. *J Herpetol Med Surg* 2001;11:4-8.
73. Kolmstetter CM, Frazier D, Cox S, et al. Pharmacokinetics of metronidazole in the green iguana, *Iguana iguana*. *Bull Assoc Rept Amphib Vet* 1998;8:4-7.
74. Helmick KE, Papich MG, Vliet KA, et al. Pharmacokinetic disposition of a long-acting oxytetracycline formulation after single-dose intravenous and intramuscular administrations in the American alligator (*Alligator mississippiensis*). *J Zoo Wildl Med* 2004;35:341-346.
75. Hilf M, Swanson D, Wagner R, et al. Pharmacokinetics of piperacillin in blood pythons (*Python curtus*) and in vitro evaluation of efficacy against aerobic gram-negative bacteria. *J Zoo Wildl Med* 1991;22:199-203.
76. Murray MJ. Pneumonia and lower respiratory tract disease. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St. Louis: Saunders/Elsevier, 2006;865-877.
77. Jacobson E, Kollias GV Jr, Peters LJ. Dosages for antibiotics and parasitides used in exotic animals. *Compend Cont Educ Pract Vet* 1983;5:315-324.
78. Manire CA, Hunter RP, Koch DE, et al. Pharmacokinetics of ticarcillin in the loggerhead sea turtle (*Caretta caretta*) after single intravenous and intramuscular injections. *J Zoo Wildl Med* 2005;36:44-53.
79. Van Waeyenberghe L, Baert K, Pasmans F, et al. Voriconazole, a safe alternative for treating infections caused by the *Chrysosporium anamorph* of *Nannizziopsis vriesii* in bearded dragons (*Pogona vitticeps*). *Med Mycol* 2010;48:880-885.
80. Innis C, Young D, Wetzlich S, et al. Plasma voriconazole concentrations in four red-eared slider turtles (*Trachemys scripta elegans*) after a single subcutaneous injection, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2008;72.
81. Fitzgerald KT, Vera R. Acariasis. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St. Louis: Saunders/Elsevier, 2006;720-738.
82. Mehlhorn H, Schmahl G, Frese M, et al. Effects of a combination of emodepside and praziquantel on parasites of reptiles and rodents. *Parasitol Res* 2005;97(Suppl 1):S65-S69.
83. Schilliger L, Betremieux O, Rochet J, et al. Absorption and efficacy of a spot-on combination containing emodepside plus praziquantel in reptiles. *Rev Med Vet (Toulouse)* 2009;160:557-561.
84. Neiffer DL, Lydick D, Burks K, et al. Hematologic and plasma biochemical changes associated with fenbendazole administration in Hermann's tortoises (*Testudo hermanni*). *J Zoo Wildl Med* 2005;36:661-672.
85. Klingenberg RJ. A comparison of fenbendazole and ivermectin for the treatment of nematode parasites in ball pythons, *Python regius*. *Bull Assoc Rept Amphib Vet* 1992;2:5-6.
86. Klingenberg RJ. Diagnosing parasites of Old World chameleons. *Exot DVM* 2000;1(6):17-21.
87. Klingenberg RJ. *Understanding reptile parasites*. Irvine, Calif: Advanced Vivarium Systems, 2007;1-200.
88. Giannetto S, Brianti E, Poglayen G, et al. Efficacy of oxfendazole and fenbendazole against tortoise (*Testudo hermanni*) oxyurids. *Parasitol Res* 2007;100:1069-1073.
89. Carpenter JW. Pers. observation, 2011.
90. Groza A, Mederle N, Darabus G. Advocate-therapeutical solution in parasitological infestation in frillneck lizard (*Chlamydosaurus kingii*) and bearded dragon (*Pogona vitticeps*). *Lucrari Stiintifice—Universitatea Stiinte Agricole Banatului Timisoara, Med Vet* 2009;42:105-108.
91. Teare JA, Bush M. Toxicity and efficacy of ivermectin in chelonians. *J Am Vet Med Assoc* 1983;183:1195-1197.
92. Flach EJ, Riley J, Mutlow AG, et al. Pentastomiasis in Bosc's monitor lizards (*Varanus exanthematicus*) caused by an undescribed *Sambonia* species. *J Zoo Wildl Med* 2000;31:91-95.
93. Wilson SC, Carpenter JW. Endoparasitic diseases of reptiles. *Semin Avian Exot Pet Med* 1996;5:64-74.
94. Luppi M, Costa M, Malta M, et al. Treatment of *Rhabdias labiata* with levamisole and ivermectin in boa constrictor (*Boa constrictor amarali*). *Veterinaria Noticias* 2007;13:61-65.
95. Mitchell MA. Therapeutics. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St. Louis: Saunders/Elsevier, 2006;631-664.
96. Stahl SJ. Clinician's approach to the chameleon patient, in *Proceedings*. North Am Vet Conf 2006;1667-1670.
97. Miller HA, Brandt PJ, Frye FL, et al. *Trichomonas* associated with ocular and subcutaneous lesions in geckos, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 1994;102-107.
98. Innis C, Papich M, Young D. Pharmacokinetics of metronidazole in the red-eared slider turtle (*Trachemys scripta elegans*) after single intracoelomic injection. *J Vet Pharm Therapeut* 2007;30:168-171.
99. Gibbons PM. Pers. observation, 2011.
100. Cranfield MR, Graczyk TK. Cryptosporidiosis. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St. Louis: Saunders/Elsevier, 2006;756-762.
101. Paré JA, Crawshaw GJ, Barta JR. Treatment of cryptosporidiosis in Gila monsters (*Heloderma suspectum*) with paromomycin, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 1997;23.
102. Coke RL, Tristan TE. *Cryptosporidium* infection in a colony of leopard geckos, *Eublepharis macularius*, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 1998;157-163.
103. Bogoslavsky B. The use of ponazuril to treat coccidiosis in eight inland bearded dragons (*Pogona vitticeps*), in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2007;8-9.
104. Mitchell MA. Ponazuril. *J Exot Pet Med* 2008;17:228-229.
105. Adnyana W, Ladds PW, Blair D. Efficacy of praziquantel in the treatment of green sea turtles with spontaneous infection of cardiovascular flukes. *Aust Vet J* 1997;75:405-407.
106. Jacobson E, Harman G, Laille E, et al. Plasma concentrations of praziquantel in loggerhead sea turtles (*Caretta caretta*) following oral administration of single and multiple doses, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2002;37-39.
107. McFarlen J. Commonly occurring reptilian internal parasites, in *Proceedings*. Annu Conf Am Assoc Zoo Vet 1991;120-127.
108. Willette-Frahm M, Wright KM, Thode BC. Select protozoal diseases in amphibians and reptiles. *Bull Assoc Rept Amphib Vet* 1995;5:19-29.
109. Girling SJ, Raiti P. *BSAVA manual of reptiles*. 2nd ed. Quedgeley, Gloucester, UK: British Small Animal Veterinary Association, 2004.
110. Doneley B. Caring for the bearded dragon, in *Proceedings*. North Am Vet Conf 2006;1607-1611.
111. Prashag P, Gibbons P, Boyer T, et al. An outbreak of intranuclear coccidiosis in *Pyxis* spp. Tortoises. *Proc 8th Annu Symp Conserv Biol Tortoises Freshwater Turtles* 2010;42-43.
112. Millichamp NJ. Surgical techniques in reptiles. In: Jacobson ER, Kollias GV Jr, eds. *Exotic animals*. New York: Churchill Livingstone, 1988;49-74.
113. Page CD. Current reptilian anesthesia procedures. In: Fowler ME, ed. *Zoo & wild animal medicine: current therapy*. 3rd ed. Philadelphia: WB Saunders Co, 1993;140-143.

114. Lawton MPC. Anaesthesia. In: Benyon PH, Lawton MPC, Cooper JE, eds. *Manual of reptiles*. Ames, Iowa: Iowa State University Press, 1992;170-183.
115. Bennett RA. A review of anesthesia and chemical restraint in reptiles. *J Zoo Wildl Med* 1991;22:282-303.
116. Schumacher J, Yelen T. Anesthesia and analgesia. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St Louis: Saunders/Elsevier, 2006;442-452.
117. Scheelings TF, Holz P, Haynes L, et al. A preliminary study of the chemical restraint of selected squamate reptiles with alfaxalone, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2010;114-115.
118. Hackenbroich C, Failing K, Axt-Findt U, et al. Alphaxalone-alphadolone anesthesia in *Trachemys scripta elegans* and its influence on respiration, circulation and metabolism, in *Proceedings*. 2nd Conf Euro Assoc Zoo Wildl Vet 1998;431-436.
119. Fleming GJ. Capture and chemical immobilization of the Nile crocodile (*Crocodylus niloticus*) in South Africa, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 1996;63-66.
120. Smith JA, McGuire NC, Mitchell MA. Cardiopulmonary physiology and anesthesia in crocodilians, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 1998;17-21.
121. Dennis PM, Heard DJ. Cardiopulmonary effects of a medetomidine-ketamine combination administered intravenously in gopher tortoises. *J Am Vet Med Assoc* 2002;220:1516-1519.
122. Boyer TH. Clinical anesthesia of reptiles. *Bull Assoc Rept Amphib Vet* 1992;2:10-13.
123. Schumacher J. Reptiles and amphibians. In: Thurman JC, Tranquilli WJ, Benson GJ, eds. *Lumb and Jones' veterinary anesthesia*. 3rd ed. Baltimore, Md: Williams & Wilkins, 1996;670-685.
124. Pace L, Mader DR. Atropine and glycopyrrolate, route of administration and response in the green iguana (*Iguana iguana*), in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2002;79.
125. Heard DJ. Principles and techniques of anesthesia and analgesia for exotic practice. *Vet Clin North Am Small Anim Pract* 1993;23:1301-1327.
126. Bennett RA, Divers SJ, Schumacher J, et al. Roundtable: anesthesia. *Bull Assoc Rept Amphib Vet* 1999;9:20-27.
127. Schnellbacher R. Butorphanol. *J Exot Pet Med* 2010;19:192-195.
128. Sladky KK, Kinney ME, Johnson SM. Effects of opioid receptor activation on thermal antinociception in red-eared slider turtles (*Trachemys scripta*). *Am J Vet Res* 2009;70:1072-1078.
129. Sladky KK, Miletic V, Paul-Murphy J, et al. Analgesic efficacy and respiratory effects of butorphanol and morphine in turtles. *J Am Vet Med Assoc* 2007;230:1356-1362.
130. Raphael BL. Chelonians. In: Fowler ME, Miller RE, eds. *Zoo and wild animal medicine*. 5th ed. Philadelphia: Saunders/Elsevier, 2003;48-58.
131. Sladky KK, Kinney ME, Johnson SM. Analgesic efficacy of butorphanol and morphine in bearded dragons and corn snakes. *J Am Vet Med Assoc* 2008;233:267-273.
132. Bennett RA. Reptile anesthesia. *Semin Avian Exot Pet Med* 1998;7:30-40.
133. Lawton MPC. Pain management after surgery, in *Proceedings*. North Am Vet Conf 1999;782.
134. McArthur SDJ, Wilkinson RJ, Barrows MG. Tortoises and turtles. In: Meredith A, Redrobe S, eds. *BSAVA manual of exotic pets*. 4th ed. Gloucestershire, GB: British Small Animal Veterinary Association, 2002;208-222.
135. Gamble KC. Plasma fentanyl concentrations achieved after transdermal fentanyl patch application in prehensile-tailed skinks, *Corucia zebrata*. *J Herpetol Med Surg* 2008;18:81-85.
136. Boyer TH. Emergency care of reptiles. *Vet Clin North Am Exot Anim Pract* 1998;1:191-206.
137. Stahl SJ. Reproductive disorders of the green iguana, in *Proceedings*. North Am Vet Conf 1998;810-813.
138. Stegman N, Heatley JJ. The use of haloperidol in mitigation of human-directed aggression in boid snakes, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2010;113.
139. Bennett RA. Anesthesia. In: Mader DR, ed. *Reptile medicine and surgery*. Philadelphia: WB Saunders Co, 1996;241-247.
140. Johnson JH. Anesthesia, analgesia, and euthanasia of reptiles and amphibians, in *Proceedings*. Annu Conf Am Assoc Zoo Vet 1991;132-138.
141. Holz P, Holz RM. Evaluation of ketamine, ketamine/xylazine, and ketamine midazolam anesthesia in red-eared sliders (*Trachemys scripta elegans*). *J Zoo Wildl Med* 1994;25:531-537.
142. Wood FE, Critchley KH, Wood JR. Anesthesia in green the sea turtle, *Chelonia mydas*. *Am J Vet Res* 1982;43:1882.
143. Lloyd ML. Crocodilian anesthesia. In: Fowler ME, Miller RE, eds. *Zoo & wild animal medicine: current therapy*. 4th ed. Philadelphia: WB Saunders Co, 1999;205-216.
144. Lane T. Crocodilians. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St. Louis: Saunders Elsevier, 2006;100-117.
145. Divers SJ. Anesthetics in reptiles. *Exot DVM* 1999;1(3):7-8.
146. Lock BA, Heard DJ, Dennis P. Preliminary evaluation of medetomidine/ketamine combinations for immobilization and reversal with atipamezole in three tortoise species. *Bull Assoc Rept Amphib Vet* 1998;8:6-9.
147. Helmick KE, Bennett RA, Ginn P, et al. Intestinal volvulus and stricture associated with a leiomyoma in a green turtle (*Chelonia mydas*). *J Zoo Wildl Med* 2000;31:221-227.
148. Fleming G. Clinical technique: chelonian shell repair. *J Exot Pet Med* 2008;17:246-258.
149. Chittick EJ, Stamper MA, Beasley JF, et al. Medetomidine, ketamine, and sevoflurane for anesthesia of injured loggerhead sea turtles: 13 cases (1996-2000). *J Am Vet Med Assoc* 2002;221:1019-1025.
150. Heard DJ. Advances in reptile anesthesia, in *Proceedings*. North Am Vet Conf 1999;770.
151. Heaton-Jones TG, Ko J, Heaton-Jones DL. Evaluation of medetomidine-ketamine anesthesia with atipamezole reversal in American alligators (*Alligator mississippiensis*). *J Zoo Wildl Med* 2002;33:36-44.
152. Bienzle D, Boyd CJ. Sedative effects of ketamine and midazolam in snapping turtles. *J Zoo Wildl Med* 1992;23:201-204.
153. Tuttle AD, Papich M, Lewbart GA, et al. Pharmacokinetics of ketoprofen in the green iguana (*Iguana iguana*) following single intravenous and intramuscular injections. *J Zoo Wildl Med* 2006;37:567-570.
154. Rojo-Solis C, Ros-Rodriguez JM, Valls M. Pharmacokinetics of meloxicam (Metacam) after intravenous, intramuscular, and oral administration in red-eared slider turtles (*Trachemys scripta elegans*), in *Proceedings*. Joint Conf Am Assoc Zoo Vet/Am Assoc Wildl Vet 2009;228.
155. Divers SJ, Papich MG, McBride M, et al. Pharmacokinetics of meloxicam following intravenous and oral administration in green iguanas (*Iguana iguana*). *Am J Vet Res* 2010;71:1277-1283.
156. Allen DG, Pringle JK, Smith D. *Handbook of Veterinary Drugs*. Philadelphia: JB Lippincott Co, 1993;534-567.
157. Nichols DK, Lamirande EW. Use of methohexital sodium as an anesthetic in two species of colubrid snakes, in *Proceedings*. Joint Conf Am Assoc Zoo Vet/Assoc Rept Amphib Vet 1994;161-162.
158. Schaeffer DO. Anesthesia and analgesia in nontraditional laboratory animal species. In: Kohn DF, Wixson SK, White WJ, et al, eds. *Anesthesia and analgesia in laboratory animals*. New York: Academic Press, 1997;338-378.
159. Schumacher J. Lacertilia. In: Fowler ME, Miller RE, eds. *Zoo and wild animal medicine*. 5th ed. Philadelphia: Saunders/Elsevier, 2003;73-81.
160. Oppenheim YC, Moon PF. Sedative effects of midazolam in red-eared sliders (*Trachemys scripta elegans*). *J Zoo Wildl Med* 1995;26:409-413.
161. Quesada RJ, Aitken-Palmer C, Conley K, et al. Accidental submeningeal injection of propofol in gopher tortoises (*Gopherus polyphemus*). *Vet Rec* 2010;167:494-495.
162. Anderson NL, Wack RF, Calloway L. Cardiopulmonary effects and efficacy of propofol as an anesthetic agent in brown tree snakes, *Boiga irregularis*. *Bull Assoc Rept Amphib Vet* 1999;9:9-15.

163. Heard DJ. Advances in reptile anesthesia and medicine, in *Proceedings. Annu Conf Assoc Avian Vet/Avian Specialty Advanced Prog/Small Mam Rept Prog* 1998;113-119.
164. Bennett RA, Schumacher J, Hedjazi-Haring K, et al. Cardiopulmonary and anesthetic effects of propofol administered intraosseously to green iguanas. *J Am Vet Med Assoc* 1998;212:93-98.
165. Ziolo M, Bertelsen M. Effects of propofol administered via the supravertebral sinus in red-eared sliders. *J Am Vet Med Assoc* 2009;234:390-393.
166. Rooney MB, Levine G, Gaynor J, et al. Sevoflurane anesthesia in desert tortoises (*Gopherus agassizii*). *J Zoo Wildl Med* 1999;30:64-69.
167. Fleming GJ. Crocodilian anesthesia. *Vet Clin North Am Exot Anim Pract* 2001;4:119-145.
168. Schobert E. Telazol use in wild and exotic animals. *Vet Med* 1987;82:1080-1088.
169. Stirl R, Bonath KH. Tiletamine/zolazepam sedation in boa constrictors and its influence on respiration, circulation, and metabolism, in *Proceedings. Conf Euro Assoc Zoo Wildl Vet* 1996;115-119.
170. Clyde VL, Cardeilhac PT, Jacobson ER. Chemical restraint of American alligators (*Alligator mississippiensis*) with atracurium or tiletamine-zolazepam. *J Zoo Wildl Med* 1994;25:525-530.
171. Greenacre CB, Massi K, Schumacher JP, et al. Comparative antinociception of various opioids and non-steroidal anti-inflammatory medications versus saline in the bearded dragon (*Pogona vitticeps*) using electrostimulation, in *Proceedings. Annu Conf Rept Amphib Vet* 2008;87.
172. Baker BB, Sladky KK, Johnson SM. Evaluation of the analgesic effects of oral and subcutaneous tramadol administration in red-eared slider turtles. *J Am Vet Med Assoc* 2011;238:220-227.
173. Lloyd ML. Reptilian dystocias review—causes, prevention, management, and comments on the synthetic hormone vasotocin, in *Proceedings. Annu Conf Am Assoc Zoo Vet* 1990;290-296.
174. Bennett RA. Management of common reptile emergencies, in *Proceedings. Annu Conf Assoc Rept Amphib Vet* 1998;67-72.
175. Mader DR. Metabolic bone diseases. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St Louis: Saunders/Elsevier, 2006; 841-851.
176. Mader DR. IME—Use of calcitonin in green iguanas, *Iguana iguana*, with metabolic bone disease. *Bull Assoc Rept Amph Vet* 1993;3:5.
177. Schumacher J. Respiratory diseases of reptiles. *Semin Avian Exot Pet Med* 1997;6:209-215.
178. Stahl SJ. Diseases of the reptile pancreas. *Vet Clin North Am Exot Anim Pract* 2003;6:191-212.
179. Kirchgeßner M, Mitchell M, Domenzain L, et al. Evaluating the effect of leuprolide acetate on testosterone levels in captive male green iguanas (*Iguana iguana*). *J Herpetol Med Surg* 2009;19:128-131.
180. Norton TM, Jacobson ER, Caligiuri R, et al. Medical management of a Galapagos tortoise (*Geochelone elephantopus*) with hypothyroidism. *J Zoo Wildl Med* 1989;20:212-216.
181. Franco KH, Hoover JP. Levothyroxine as a treatment for presumed hypothyroidism in an adult male African spurred tortoise (*Centrochelys* [formerly *Geochelone*] *sulcata*). *J Herpetol Med Surg* 2009;19:42-44.
182. Hernandez-Divers SJ, Cooper JE. Hepatic lipidosis. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St. Louis: Saunders/Elsevier, 2006;806-813.
183. Divers SJ. Clinician's approach to renal disease in lizards, in *Proceedings. Annu Conf Assoc Rept Amphib Vet* 1997;5-11.
184. Faulkner JE, Archambault A. Anesthesia and surgery in the green iguana. *Semin Avian Exot Pet Med* 1993;2:103-108.
185. Antinoff N, Bauck L, Boyer TH, et al. *Exotic animal formulary*. 2nd ed. Lakewood, Colo: AAHA Press, 1999.
186. Divers SJ. Medical and surgical treatment of pre-ovulatory ova stasis and post-ovulatory egg stasis in oviparous lizards, in *Proceedings. Annu Conf Assoc Rept Amphib Vet* 1996;119-123.
187. Boyer TH. Emergency care of reptiles. *Semin Avian Exot Pet Med* 1994;3:210-216.
188. Boyer TH. Common problems of box turtles (*Terrapene* spp) in captivity. *Bull Assoc Rept Amphib Vet* 1992;2:9-14.
189. Mauldin GN, Done LB. Oncology. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St Louis: Saunders/Elsevier, 2006;299-322.
190. Georoff TA, Stacy NI, Newton AN, et al. Diagnosis and treatment of chronic T-lymphocytic leukemia in a green tree monitor (*Varanus prasinus*). *J Herpetol Med Surg* 2009;19:106-114.
191. Norton TM. Chelonian emergency and critical care. *Semin Avian Exot Pet Med* 2005;14:106-130.
192. Rivera S. Health assessment of the reptilian reproductive tract. *J Exot Pet Med* 2008;17:259-266.
193. Gibbons PM. Critical care nutrition and fluid therapy in reptiles, in *Proceedings. 15th Annu Intl Vet Emerg Crit Care Symp* 2009;91-94.
194. Mader DR, Rudloff E. Emergency and critical care. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St Louis: Saunders/Elsevier, 2006;533-548.
195. Wellehan JFX, Gunkel CI. Emergent diseases in reptiles. *Semin Avian Exot Pet Med* 2004;13:160-174.
196. Donoghue S. Nutrition. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St. Louis: Saunders/Elsevier, 2006;251-298.
197. Suedmeyer WK. Iron deficiency in a group of American alligators: diagnosis and treatment. *J Small Exot Anim Med* 1991;1:69-72.
198. Kirchgeßner M, Mitchell MA. Chelonians. In: Mitchell MA, Tully TN Jr, eds. *Manual of exotic pet practice*. St. Louis: Saunders/Elsevier, 2009;207-249.
199. Jarchow JL. Hospital care of the reptile patient. In: Jacobson ER, Kollias GV Jr, eds. *Exotic animals*. New York: Churchill Livingstone, 1988;19-34.
200. Millichamp NJ. Ophthalmology. In: Girling SJ, Raiti P, eds. *BSAVA manual of reptiles*. 2nd ed. Quedgeley, Gloucester: British Small Animal Veterinary Association, 2004;199-209.
201. Stahl SJ. Captive management, breeding, and common medical problems of the veiled chameleon (*Chamaeleo calytratus*), in *Proceedings. Annu Conf Assoc Rept Amphib Vet* 1997;29-40.
202. Harkewicz KA. Dermatologic problems of reptiles. *Semin Avian Exot Pet Med* 2002;11:151-161.
203. Calvert I. Nutritional problems. In: Girling SJ, Raiti P, eds. *BSAVA manual of reptiles*. 2nd ed. Quedgeley, Gloucester: British Small Animal Veterinary Association, 2004;289-308.
204. Fraser MA, Girling SJ. Dermatology. In: Girling SJ, Raiti P, eds. *BSAVA manual of reptiles*. 2nd ed. Quedgeley, Gloucester: British Small Animal Veterinary Association, 2004;184-198.
205. Mehler SJ, Bennett RA. Upper alimentary tract disease. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St Louis: Saunders/Elsevier, 2006;924-930.
206. Farnsworth RJ, Brannian RE, Fletcher KC, et al. A vitamin E-selenium responsive condition in a green iguana. *J Zoo Anim Med* 1986;17:42-43.
207. Raiti P. Veterinary care of the common kingsnake, *Lampropeltis getula*. *Bull Assoc Rept Amphib Vet* 1995;5:11-18.
208. Mader DR. Gout. In: Mader DR, ed. *Reptile medicine and surgery*. Philadelphia: WB Saunders Co, 1996;374-379.
209. Hernandez-Divers SJ, Martinez-Jimenez D, Bush S, et al. Effects of allopurinol on plasma uric acid levels in normouricaemic and hyperuricaemic green iguanas (*Iguana iguana*). *Vet Rec* 2008;162:112-115.
210. Koelle P. Efficacy of allopurinol in European tortoises with hyperuricemia, in *Proceedings. Annu Conf Assoc Rept Amphib Vet* 2001;185-186.
211. Meyer J. Gastrographin as a gastrointestinal contrast agent in the Greek tortoise (*Testudo hermanni*). *J Zoo Wildl Med* 1998;29:183-189.
212. Messonnier S. Formulary for exotic pets. *Vet Forum* 1996;Aug:46-49.
213. Rossi JV. Emergency medicine of reptiles, in *Proceedings. North Am Vet Conf* 1998;799-801.
214. Carpenter JW. Radiographic imaging of reptiles, in *Proceedings. North Am Vet Conf* 1998;873-875.
215. Tohill A, Johnson J, Branvold H, et al. Effect of cisapride, erythromycin, and metoclopramide on gastrointestinal transit time in the desert tortoise, *Gopherus agassizii*. *J Herpetol Med Surg* 2000;10:16-20.



216. Gillespie D. Reptiles. In: Birchard SJ, Sherding RG, eds. *Saunders manual of small animal practice*. Philadelphia: WB Saunders Co, 1994;1390-1411.
217. Rosenthal K. Chemotherapeutic treatment of a sarcoma in a corn snake, in *Proceedings*. Joint Conf Am Assoc Zoo Vet/Assoc Rept Amphib Vet 1994;46.
218. Stahl SJ. Medical management of bearded dragons, in *Proceedings*. North Am Vet Conf 1999;789-792.
219. Divers SJ. Constipation in snakes with particular reference to surgical correction in a Burmese python (*Python molurus bivittatus*), in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 1996;67-69.
220. Wright KM. Common medical problems of tortoises, in *Proceedings*. North Am Vet Conf 1997;769-771.
221. American Veterinary Medical Association Web site. AVMA guidelines on euthanasia. Available at: [www.avma.org/issues/animal\\_welfare/euthanasia.pdf](http://www.avma.org/issues/animal_welfare/euthanasia.pdf). Accessed May 6, 2011.
222. Burns RB, McMahan W. Euthanasia methods for ectothermic vertebrates. In: Bonagura JD, ed. *Kirk's current veterinary therapy XII: small animal practice*. Philadelphia: WB Saunders Co, 1995;1379-1381.
223. Plumb DC. Probenecid. In: Plumb DC, ed. *Plumb's veterinary drug handbook*. 6th ed. Ames, Iowa: Blackwell Publishing, 2008;1028-1030.
224. Naples LM, Langan JN, Mylniczenko ND, et al. Islet cell tumor in a Savannah monitor (*Varanus exanthematicus*). *J Herpetol Med Surg* 2009;19:97-105.
225. DeNardo DF, Helmski G. Birth control in lizards? Therapeutic inhibition of reproduction, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2000;65-66.
226. Chiodini RJ, Sundberg JP. Blood chemical values of the common boa constrictor (*Constrictor constrictor*). *Am J Vet Res* 1982;43:1701-1702.
227. Machado CC, Silva LFN, Ramos PRR, et al. Seasonal influence on hematologic values and hemoglobin electrophoresis in Brazilian *Boa constrictor amarali*. *J Zoo Wildl Med* 2006;37:487-491.
228. Teare JA. Physiological data reference values project. International Species Information System (ISIS), Apple Valley, Minn, 2010.
229. Johnson JH, Benson PA. Laboratory reference values for a group of captive ball pythons (*Python regius*). *Am J Vet Res* 1996;57:1304-1307.
230. Centini R, Klaphake E. Hematologic values and cytology in a population of captive jungle carpet pythons, *Morelia spilota cheynei*, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2002;107-111.
231. Mader DR, Horvath CC, Paul-Murphy J. The hematocrit and serum profile of the gopher snake (*Pituophis melanoleucas catenifer*). *J Zoo Anim Med* 1985;16:139-140.
232. Ramsay EC, Dotson TK. Tissue and serum enzyme activities in the yellow rat snake (*Elaphe obsoleta quadrivittata*). *Am J Vet Res* 1995;56:423-428.
233. Drew ML. Hypercalcemia and hyperphosphatemia in indigo snakes (*Drymarchon corais*) and serum biochemical reference values. *J Zoo Wildl Med* 1994;25:48-52.
234. Naldo JL, Libanan NL, Samour JH. Health assessment of a spiny-tailed lizard (*Uromastyx* spp.) population in Abu Dhabi, United Arab Emirates. *J Zoo Wildl Med* 2009;40:445-452.
235. Ellman MM. Hematology and plasma chemistry of the inland bearded dragon, *Pogona vitticeps*. *Bull Assoc Rept Amphib Vet* 1997;7:10-12.
236. Mayer J. Characterizing the hematologic and plasma chemistry profiles of captive Chinese water dragons, *Physignathus cocincinus*. *J Herpetol Med Surg* 2005;15:45-52.
237. Dennis PM, Bennett RA, Harr KE, et al. Plasma concentration of ionized calcium in healthy iguanas. *J Am Vet Med Assoc* 2001;219:326-328.
238. Divers SJ, Redmayne G, Aves EK. Haematological and biochemical values of 10 green iguanas (*Iguana iguana*). *Vet Rec* 1996;138:203-205.
239. Gimenez M, Saco Y, Pato R, et al. Plasma protein electrophoresis of *Trachemys scripta* and *Iguana iguana*. *Vet Clin Pathol* 2010;39:227-235.
240. Hernandez-Divers SJ, Knott CD, MacDonald J. Diagnosis and surgical treatment of thyroid adenoma-induced hyperthyroidism in a green iguana (*Iguana iguana*). *J Zoo Wildl Med* 2001;32:465-475.
241. McBride M, Hernandez-Divers SJ, Koch T, et al. Preliminary evaluation of pre- and post-prandial 3[alpha]-hydroxy bile acids in the green iguana, *Iguana iguana*. *J Herpetol Med Surg* 2006;16:129-134.
242. Nevarez JG, Mitchell MA, Le Blanc C, et al. Determination of plasma biochemistries, ionized calcium, vitamin D<sub>3</sub>, and hematocrit values in captive green iguanas (*Iguana iguana*) from El Salvador, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2002;87-91.
243. Harr KE, Alleman AR, Dennis PM, et al. Morphologic and cytochemical characteristics of blood cells and hematologic and plasma biochemical reference ranges in green iguanas. *J Am Vet Med Assoc* 2001;218:915-921.
244. Judd HL, Laughlin GA, Bacon JP, et al. Circulating androgen and estrogen concentrations in lizards (*Iguana iguana*). *Gen Comp Endocrinol* 1976;30:391-395.
245. Wright KM, Skeba S. Hematology and plasma chemistries of captive prehensile-tailed skinks (*Corucia zeburata*). *J Zoo Wildl Med* 1992;23:429-432.
246. Troiano J, Gould E, Gould I. Hematological reference intervals in Argentine lizard *Tupinambis merianae*. *Comp Clin Pathol* 2008;17:169-174.
247. Diethelm G, Stein G. Hematologic and blood chemistry values in reptiles. In: Mader DR, ed. *Reptile medicine and surgery*. 2nd ed. St. Louis: Saunders/Elsevier, 2006;1103-1118.
248. Fudge AM. Laboratory reference ranges for selected avian, mammalian, and reptilian species. In: Fudge AM, ed. *Laboratory medicine: avian and exotic pets*. Philadelphia: WB Saunders Co, 2000;375-400.
249. Marks SK, Citino SB. Hematology and serum chemistry of the radiated tortoise (*Testudo radiata*). *J Zoo Wildl Med* 1990;21:342-344.
250. Zaais J, Norton T, Fickel A, et al. Biochemical and hematologic values for 18 clinically healthy radiated tortoises (*Geochelone radiata*) on St. Catherine's Island, Georgia. *Vet Clin Pathol* 2006;35:321-325.
251. Alleman AR, Jacobson ER, Raskin RE. Morphologic and cytochemical characteristics of blood cells from the desert tortoise (*Gopherus agassizii*). In *Proceedings*. Annu Conf Assoc Rept Amphib Vet 1996;51-55.
252. Christopher MM, Berry KH, Wallis IR, et al. Reference intervals and physiologic alterations in hematologic and biochemical values of free-ranging desert tortoises in the Mojave Desert. *J Wildl Dis* 1999;35:212-238.
253. Dickinson VM, Jarchow JL, Trueblood MH. Hematology and plasma biochemistry reference range values for free-ranging desert tortoises in Arizona. *J Wildl Dis* 2002;38:143-153.
254. Gottdenker NL, Jacobson ER. Effect of venipuncture sites on hematologic and clinical biochemical values in desert tortoises (*Gopherus agassizii*). *Am J Vet Res* 1995;56:19-21.
255. Taylor RW Jr, Jacobson ER. Hematology and serum chemistry of the gopher tortoise, *Gopherus polyphemus*. *Comp Biochem Physiol A* 1982;72:425-428.
256. Knotkova Z, Doubek J, Knotek Z, et al. Blood cell morphology and plasma biochemistry in Russian tortoises (*Agrynemys horsfieldi*). *Acta Vet Brno* 2002;71:191-198.
257. Mathes KA, Holz A, Fehr M. Blood reference values of terrestrial tortoises (*Testudo* spp.) kept in Germany. *Tierarztl Prax Ausg K Klientiere Heimtiere* 2006;34:268-274.
258. Crawshaw GJ, Holz P. Comparison of plasma biochemical values in blood and blood-lymph mixtures from red-eared sliders, *Trachemys scripta elegans*. *Bull Assoc Rept Amphib Vet* 1996;6:7-9.
259. Gaumer A, Goodnight CJ. Some aspects of the hematology of turtles as related to their activity. *Am Midland Nat* 1957;58:332-340.
260. Jacobson ER. Evaluation of the reptile patient. In: Jacobson ER, Kollias GV Jr, eds. *Exotic Animals*. New York: Churchill Livingstone, 1988;1-18.
261. Flint M, Morton JM, Limpus CJ, et al. Development and application of biochemical and haematological reference intervals to identify unhealthy green sea turtles (*Chelonia mydas*). *Vet J* 2010;185:299-304.



262. Barrie MT. Chameleon medicine. In: Fowler ME, Miller RE, eds. *Zoo & wild animal medicine: current therapy*. 4th ed. Philadelphia: WB Saunders Co, 1999;200-205.
263. Cunningham AA, Gill C. Management in captivity. In: Benyon PH, Lawton MPC, Cooper JE, eds. *Manual of reptiles*. Ames: Iowa State University Press, 1992;14-31.
264. Dundee HA, Rossman DA. *The amphibians and reptiles of Louisiana*. Baton Rouge: Louisiana State University Press, 1989;270-272.
265. Harless M, Morlock H. *Turtles—perspectives and research*. New York: John Wiley & Sons, 1979;356-366.
266. McKeown S. General husbandry and management. In: Mader DR, ed. *Reptile medicine and surgery*. Philadelphia: WB Saunders Co, 1996;9-19.
267. Gibbons PM, Horton SJ, Brandl SR. Urinalysis in box turtles, *Terapene* spp, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2000;161-168.
268. Innis CJ. Observations on urinalysis of clinically normal captive tortoises, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 1997;109-112.
269. Koelle P. Urinalysis in tortoises, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2000;111-113.
270. Koelle P, Hoffmann R. Urinalysis in European tortoises—part II, in *Proceedings*. Annu Conf Assoc Rept Amphib Vet 2002;117.
271. Stahl SJ. Common Diseases of the Green Iguana. *Proc North Am Vet Conf* 1998;806-809.