

Original Article

Petrel extinction in Macaronesia (North-East Atlantic Ocean): the case of the genus *Pterodroma* (Aves: Procellariiformes: Procellariidae)

Juan C. Rando^{1, ID}, Harald Pieper², Fernando Pereira^{3, ID}, Enric Torres-Roig^{4, ID} and Josep Antoni Alcover^{4,5, *, ID}

¹Departamento de Biología Animal, Universidad de La Laguna (ULL), E-38206 La Laguna, Tenerife—Canary Islands, Spain

²Ulmstraße 21, D-24223 Schwentinental, Germany

³Centre for Ecology, Evolution and Environmental Changes (Ce3C) and Azorean Biodiversity Group, Universidade dos Açores, Rua Capitão João d'Ávila, Pico da Urze, 9700-042 Angra do Heroísmo, Terceira, Açores, Portugal

⁴Institut Mediterrani d'Estudis Avançats (CSIC-UIB, Center of Excellence María de Maeztu), Ctra. Valldemossa km 7.5, 07122 Palma – Balearic Islands, Spain

⁵Affiliated Researcher ad Honorem of CSIC, Institut Mediterrani d'Estudis Avançats (CSIC-UIB, Center of Excellence María de Maeztu), Ctra. Valldemossa km 7.5, 07122 Palma – Balearic Islands, Spain

*Corresponding author. Institut Mediterrani d'Estudis Avançats (CSIC-UIB), Ctra. Valldemossa km 7.5, 07122 Palma—Balearic Islands, Spain.

E-mail: jaalcover@imedea.csic-uib.es

ABSTRACT

The Late Quaternary fossil record indicates that formerly in the North Atlantic volcanic Macaronesian archipelagos (Azores, Madeira, Selvagens, Canary Islands, and Cape Verde) there was a higher avian diversity, including numerous now extinct species. Currently, only three gadfly petrels (genus *Pterodroma*) remain in two archipelagos: the Fea's petrel, *Pt. feae*, in Cape Verde (islands of Santiago, Fogo, Santo Antão, and São Nicolau); the Zino's petrel of Madeira, *Pt. madeira*, on the island of Madeira; and the Desertas petrel, *Pt. deserta*, on the islet of Bugio (Desertas Islands, Madeira Archipelago). Herein we describe the former distribution of the genus in Macaronesia based on the palaeontological record. However, the original specific diversity cannot be accurately established through the biometry of their fossil bones but the fossil record of *Pterodroma* in Macaronesia indicates: (i) its former presence in all Macaronesian archipelagos; (ii) the extinction of at least 16 island populations (73% of its original distribution on these islands); and (iii) the extinction of, at least, one species from Azores, here named *Pterodroma zinorum* sp. nov.. Radiocarbon dates indicate that this species was still alive at sometime among 1104 and 1672 CE, documenting its extinction in the last millennium.

Keywords: extinction; gadfly petrels; Macaronesian Islands; new species; radiocarbon dates; sea birds

INTRODUCTION

Quaternary extinctions have had a dramatic effect on island biotas, with a high number of vanished insular land and sea birds during the Holocene. During the prehistoric period, in the Pacific islands, the number of terrestrial non-passerine extinct bird species was 983 (95% CI: 731–1332) (Duncan *et al.* 2013). Seabirds and passerines from these islands also suffered a high number of extinctions (Steadman 2006). Parallel extinction took place in other worldwide archipelagos like the Canary Islands or the Caribe (Hume and Walters 2012). A subsequent wave of historic extinctions followed this episode of prehistoric

extinctions at these same archipelagos and on other islands that had remained uninhabited by humans until then, such as the Mascarene Islands, Madeira, and the Azores archipelago (Curnutt and Pimm 2001, Steadman 2006, Russell and Kuefner 2019, Sayol *et al.* 2020). Thus, extinct insular birds represent 10%–20% of the original avian diversity (Fromm and Meiri 2021). Currently, the number of extinct terrestrial native birds widely overpasses that of the extant species in some archipelagos (e.g. Hawaii, 71% vs. 29%, respectively; Easter Island, 100% vs. 0%) (Steadman 1995, Boyer 2008). This loss of native insular vertebrates is a direct consequence of human activities and the

Received 2 May 2024; revised 19 August 2024; accepted 11 September 2024

[Version of Record, first published online 25 October 2024, with fixed content and layout in compliance with Art. 8.1.3.2 IZN. <http://zoobank.org/urn:lsid:zoobank.org:pub:22E1B731-EFAB-4791-ASF3-C32775EB6AE0>]

© 2024 The Linnean Society of London.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.

introduction of alien species (Olson and James 1982, Worthy and Holdaway 2002, Blackburn *et al.* 2004, Steadman 2006, Turvey 2009, Hume and Walters 2012, Cooke *et al.* 2023).

Among seabirds, Procellariiformes are highly vulnerable to human settlement and to the introduction of alien mammals, because they mainly breed on islands free of terrestrial predators, they have a strong breeding-site fidelity, and they lack effective anti-predator behaviour (Warham 1996). This vulnerability could be why at least 56% of the species of Procellariiformes lost populations during the Holocene (Scofield 2009), and that more than 15 species have become extinct (Tyrberg 2009).

Within Procellariiformes, the family Procellariidae contains 16 genera and 98 extant species (Winkler *et al.* 2020). Uncertainty arose from the difficulties in establishing the species' limits, mainly within the two most diversified genera, *Puffinus* (shearwaters) and *Pterodroma* (gadfly petrels). Limits among species of both genera are controversial (Ferrer Obiol *et al.* 2023) as different recognized species can present similar or only slightly different coloration patterns, body size, and proportions, or can be subtly or strongly differentiated by their voices, breeding behaviour, and/or migration patterns. Some species are highly variable along their distribution range, while others are more homogeneous. Different genetic criteria have been used to establish their relationships. There are populations prone to be differentiated more morphologically than genetically in both genera, while other populations are more similar morphologically than genetically. The great philopatry exhibited by these different populations, occasionally together with the breeding allochrony and geographic disparity, probably favours their evolutionary differentiation (Ferrer Obiol *et al.* 2023). All these factors create challenges to resolving phylogenetic relationships within both genera.

The gadfly petrels are small to medium-sized (26–46 cm and 150–800 g) found in all oceans but are more diverse in the subtropical South Atlantic. Their short, robust bills are adapted for soft prey captured dipping at the sea surface. They only come to land to breed in colonies on islands, with one white egg laid in a burrow or on open ground. They are nocturnal in the breeding colonies (Warham 1996, Harrison *et al.* 2021).

In this paper we focus on a group of the Northern Atlantic gadfly petrels (genus *Pterodroma*), the species found in the volcanic Macaronesian archipelagos (Azores, Madeira, Selvagens, Canary Islands, and Cape Verde), located in the north-east Atlantic Ocean (15°N–39°N and 10°W–30°W), and between ~100 km (Canary Islands) and ~1350 km (Azores) from Africa and Europe, respectively (Fig. 1). Three species of gadfly petrels (genus *Pterodroma*) are currently breeding in two archipelagos: *Pt. feae* (Salvadori, 1899) in Cape Verde (islands of Santiago, Fogo, Santo Antão, and São Nicolau), *Pt. madeira* Mathews, 1934 on Madeira Island, and *Pt. deserta* Mathews, 1934 on the islet of Bugio (Desertas Islands, Madeira Archipelago) (Hazevoet 1995, Jesus *et al.* 2009, Ramos *et al.* 2016). Initially, these three populations were considered to belong to *Pt. mollis* (Gould, 1844) due to their morphological similarities, later as one distinct species (*Pt. feae*) with three subspecies, or as two species (one in Madeira Island, *Pt. madeira*, and the other shared by Bugio and Cape Verde, *Pt. feae*), and recently as three species (see: Zino *et al.* 2008, Jesus *et al.* 2009, Gangloff *et al.* 2013).

The palaeontological record indicates a wider distribution of the genus in the past of Macaronesia. The island of El Hierro (Canary Island) had a population of this genus in former times, which probably went extinct after human colonization less than 2000 years ago (Rando 2002), and *Pterodroma* bones have been found in the archipelago of Selvagens (see: Gangloff *et al.* 2013) (Fig. 1).

Our palaeontological survey provided bones of gadfly petrels (genus *Pterodroma*) in both archipelagos of Madeira and the Azores. Specifically, in the islands of Madeira, Bugio, and Porto Santo in the former, and in São Miguel, Terceira, Pico, Faial, São Jorge, and Graciosa in the latter.

This paper provides information on the original diversity and distribution of the genus *Pterodroma* in these archipelagos before human settlement, including a description of a new extinct endemic species from the Azores, and discusses the chronology and possible causes for regression and extinction to explain its current narrow distribution in Macaronesia.

MATERIALS AND METHODS

Skeletons used for comparisons are from the Museu de História Natural do Funchal (MMF, Madeira, Portugal), the Institut Mediterrani d'Estudis Avançats (IMEDEA, Mallorca, Spain), and Departamento de Biología Animal, Geología y Edafología de la Universidad de La Laguna (DZUL, Canary Islands, Spain). The modern specimens used as comparative material are listed in the Appendix.

The material was collected from volcanic tubes in Madeira Island and São Miguel, Terceira, Pico, Faial, São Jorge, and Graciosa (Azores Islands, Portugal), at the sand dunes on Porto Santo, and at sedimentary sites on Madeira and Bugio (Madeira Archipelago). Information about the Azorean caves is available in Pereira *et al.* (2015).

The fossil specimens described herein are housed in the following collections: (i) Museu de História Natural do Funchal (MMF), Funchal, Madeira, Portugal; (ii) Museu de História Natural do Jardim Botânico da Madeira, Funchal, Portugal; (iii) Museu Carlos Machado (MCMa), São Miguel, Azores, Portugal; (iv) Museu dos Montanheiros, Angra do Heroísmo, Terceira, Azores, Portugal (without collection acronym); and (v) Institut Mediterrani d'Estudis Avançats (IMEDEA), Mallorca, Balearic Islands, Spain.

Measurements were taken with electronic digital callipers to 0.01 mm, and rounded to the nearest 0.1 mm. We followed the criteria of Rando (2002) for taking measurements of the bones. Osteological terminology follows that of Livezey and Zusi (2006). The lengths of the main bones, beaks, and jaws were measured to determine biometrical differences species of *Pterodroma* performing multivariate analysis of variances (MANOVA) and one-way analysis of variance (one-way ANOVA). The pairwise comparisons were made through *t*-test and elements known only from three specimens were pairwise compared with a nonparametric test (Mann–Whitney U-tests).

The collagen of selected bones was ¹⁴C-dated using Accelerator Mass Spectrometry (AMS) to obtain chronological information about the material. Two samples from Pico (Azores) were subjected to an improved collagen extraction (van Strydonck




Figure 1. Map of the Macaronesian Islands. The three species of endemics *Pterodroma* and the islands where they currently breed are shown by solid forms (*Pt. madeira* in Madeira; *Pt. deserta* in Bugio; and *Pt. feae* in Fogo, Santiago, São Nicolau and Santo Antão). The islands where bones of this genus have been found are shown by empty silhouettes (two empty silhouettes in the islands where two size classes of *Pterodroma* were found).

et al. 2005) by the modified Longin method (Brown et al. 1998) at the Koninklijk Instituut voor Het Kunstmuseum (Brussels). AMS measurements were performed on extracts

of ultrafiltered bone gelatine from a sample from Porto Santo (Madeira Archipelago) at the Radiocarbon Dating Laboratory of The University of Waikato (Hamilton, New Zealand).

The ^{14}C ages are given by the laboratories in years before present (BP) and are expressed as 2σ intervals (i.e., $p = 95.45\%$). Their interpretation is based exclusively on the extreme values of these intervals indicating that the true age of the dated material is more recent than the lower extreme value of the 2σ interval and, independently, it is more ancient than the upper extreme value of the 2σ interval (see e.g. Zilhão 2001). Dates calculated from calibration (cal) are given as ‘CE’ or ‘BCE’, CE being ‘current era’ and BCE ‘before current era’.

Radiocarbon samples from species that obtain their carbon from a source (or reservoir) different to atmospheric carbon, such as seabirds, will yield radiocarbon dates excessively old and require the application of a correction factor that varies with location due to complexities in ocean circulation. This regional difference from the average global marine reservoir correction is designated as ΔR and is assumed to be a constant for a given region and is calculated from the difference in ^{14}C years of known age marine samples and the marine model age for that calendar age (Stuiver and Braziunas 1993). The ages were calibrated using OxCal v.4.4 (Bronk Ramsey 2009) and a marine calibration curve (Marine20) (Heaton et al. 2020). Other calibrations were performed with a reservoir correction of $\Delta\text{R} = 18 \pm 141$ for a sample from Porto Santo (Madeira Archipelago), and $\Delta\text{R} = 38 \pm 139$ for two samples from Pico (Azores). Reservoir corrections were obtained from the 20 samples from the coast of mainland Portugal closest to the dated samples using the Marine Reservoir Correction Database (<http://calib.org/marine/>).

RESULTS

Systematic palaeontology

Order Procellariiformes Fürbringer, 1888

Family Procellariidae Leach, 1820

Genus *Pterodroma* Bonaparte, 1856

The material presented herein is included in the genus *Pterodroma* because of the following combination of characters: the dorsal distal outline of the premaxilla shows a deep concave kink in lateral view, the lachrymal is perfectly fused to the frontal and ectethmoid without any noticeable remaining suture between them forming part of the anterior wall of the eye socket (Warham 1996).

The bones are larger than Hydrobatidae and Oceanitidae with different proportions to those of Oceanitidae as they have short, stout humeri, ulnae, and femora, and greatly elongated tibiotarsi and tarsometatarsi (Olson 1985). These bones show no tendency toward the deep-diving adaptations of *Puffinus* or *Pelecanoides* in either the wing (flattened humerus and ulna) or hindlimbs (enlargement of crista cnemialis of the tibiotarsi and flattened tarsometatarsi) (Warham 1996). The cranium lacks the elongation of the area between the ectethmoid and the nasofrontal hinge with anterior elongation of the lacrimal typical of Oceanitidae (Olson and James 1991).

The fusion of the lachrymal to the frontal and ectethmoid is absent in most Procellariiformes (Olson 1975b, Wharham 1996), but is present in the close genus *Lugensa*. The present material can be differentiated from *Lugensa* by the presence of a fenestrated fossa glandulae nasalis in the latter (Maurício 2014).

The shape of the fossil humeri also differs from the very slender and cylindrical humeri of *Bulweria* (Kuroda, 1983). In addition, the shape of these bones and the limb proportions are like other *Pterodroma* species (*Pt. madeira*, *Pt. feae*, *Pt. cahow*, and *Pt. deserta*) and different from those of *Bulweria* (Olson, 1975b).

These anatomical features have been checked with skeletons of *Pt. madeira*, *Pt. feae*, *Pt. cahow*, and *Pt. deserta*, *Puffinus puffinus* (Brünnich, 1764), *Pu. baroli* Bonaparte, 1856, and *Bulweria bulwerii* (Jardine and Selby, 1828) (Procellariidae), *Pelagodroma marina* (Latham, 1790) (Oceanitidae), and *Hydrobates pelagicus* (Linnaeus, 1758) (Hydrobatidae).

Pterodroma zinorum Rando et al. sp. nov.

urn:lsid:zoobank.org:act:69B8A7C7-875A-4D23-B3E8-414785A6EED4.

Holotype: MCMA 23020.023, a partial associated skeleton, including almost complete skull with premaxilla lacking palatinum and pterygoideum, left and right quadratum, 16 vertebrae, synsacrum, clavica, a proximal fragment of the sternum, left and right coracoids, left and right scapula lacking distal part, right humerus, left and right ulna, left and right radius, right carpometacarpus, left phalanx proximalis digitii majoris, left phalanx distalis digitii majoris, left femur, distal fragment of right femur, proximal fragment of the right tibiotarsus, left tibiotarsus, left tarsometatarsus, and four pedal phalanges (Fig. 2; Supporting Information, Figs S1–S8).

Measurements of the holotype (mm): skull length: 66.3; skull height: 19.7; interorbital width: 8.7; culmen length: 16.5; coracoid length: 26; coracoid width: 17.4; right humerus length: 74.2; left ulna length: 76.5; right carpometacarpus length: 37; left femur length: 27; left tibiotarsus length: 49; left tarsometatarsus length: 30.2.

Type locality: Furna das Torres, Pico Island (Azores).

Horizon: Late Quaternary–Recent.

Distribution: The Azores archipelago (at least the islands of Graciosa, São Jorge, Pico, Terceira, and São Miguel).

Status: Extinct.

Etymology: The species’ name honours both the late Paul Alexander Zino and his son Francis Zino for their effort and contributions to the knowledge and conservation of *Pterodroma* species from Macaronesia.

Paratypes: MCMA 23021.023, a partial associated skeleton including an almost complete skull with premaxilla lacking palatinum and pterygoideum, three fragments of the mandible, eight vertebrae, incomplete sternum, right humerus, left ulna, left radius, a proximal fragment of right radius, right carpometacarpus, left phalanx proximalis digitii majoris, right femur, right tibiotarsus, right tarsometatarsus, and two pedal phalanges.

MCMA 23022.023, a partial associated skeleton, including an almost complete skull with premaxilla lacking pterygoideum, left quadratum, three vertebrae, synsacrum, a proximal fragment of the sternum, left coracoid, a distal fragment of the right coracoid,




Figure 2. Bones of *Pterodroma zinorum* holotype from Pico Island (Azores) (left) and *Pt. madeira* from Madeira island (right). A, skull right lateral view. B, skull dorsal view. C, humerus, caudal view. D, ulna, dorsal view. E, carpometacarpus, dorsal view. F, coracoid, dorsal view. G, femur, caudal view. H, tarsometatarsus, dorsal view. All paired bones are of the left side except D. A-H (right series) MMF 42034; A-D, F (left) MCMA 23020.023; E, H (left) MCMA 23021.023; G (left) MCMA 23022.023.

left scapula, right humerus, right ulna, a distal fragment of the left ulna, a distal fragment of the left radius, right carpometacarpus, a distal fragment of the left carpometacarpus, right femur, left tibiotarsus, a distal fragment of the right tibiotarsus, left tarsometatarsus, and a proximal fragment of the right tarsometatarsus.

Other material belonging to Pterodroma zinorum: See [Supporting Information, Table S1](#).

Suggested English name: Azorean Little Gadfly Petrel.

Suggested Portuguese name: Freira pequena dos Açores.

Diagnosis: A small species of *Pterodroma* similar in size to *Pt. madeira* from Madeira Island but with shorter forelimb bones (specially the ulna) and tarsometatarsus, but with a wider fossa glandulae nasalis in *Pt. zinorum* than in *Pt. madeira*. In addition, the distal extreme of the rostri maxillae of the new species shows a dorsal side higher and more curved than in *Pt. madeira* (Fig. 2).

Much smaller than the other Atlantic species of this genus (Tables 2–4; Figs 2–7; [Supporting Information, S1–S8](#)),

including the South Atlantic *Pt. incerta* (Schlegel, 1863) ([Carboneras et al. 2020a](#)), but with a humerus longer than the small Pacific Ocean species of this genus, *Pt. pycrofti* Falla, 1933, *Pt. cookii* (Gray, 1843), *Pt. leucoptera* (Gould, 1844), *Pt. nigripennis* (Rothschild, 1893), and *Pt. axillaris* (Salvin, 1893) [see [Table 3](#) and [Cooper and Tennyson \(2008\)](#)].

Remarks: The ^{14}C date of the complete right ulna of one specimen of *Pt. zinorum* sp. nov. from the type locality (RICH-21397) 1181 ± 30 BP was calibrated with 38 ± 139 ($\Delta R \pm SD$) providing the interval 1104–1672 CE ([Table 1](#)). Although there is overlap in the length of the hindlimb bones (femur, tibiotarsus, and tarsometatarsus) of *Pt. zinorum* and *Pt. madeira*, forelimb bones referable to the former (humerus, ulna, and carpometacarpus) are smaller than in *Pt. madeira* ([Tables 2–4](#); Figs 2–4, 6, 7; [Supporting Information, Figs S1–S8](#)), so *Pt. zinorum* is the smallest Atlantic species of the genus. The morphology of the postcranial bones of the new species is similar to that of *Pt. madeira* with differences in the bone length between the two species. The ulna, tarsometatarsus (Wilks's

Table 1. Radiocarbon ages (yr BP) of gadfly petrels (genus *Pterodroma*) and two 2σ calibration intervals (cal BCE/cal CE). The two 2σ calibration intervals have been performed (i) using the Marine20 calibration curve and, (ii) using the Marine20 calibration curve with a $\Delta R \pm SD$ calculated from the 20 closer samples to each archipelago (18 ± 141 for Madeira and 38 ± 139 for Azores) through the Marine Reservoir Correction Database (<http://calib.org/marine/>). (A), Azores; (M), Madeira Archipelago.

Lab code	Island	Sample	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	Radiocarbon age	2σ	2σ $\Delta R \pm SD$
RICH-21397	Pico (A)	<i>Pt. zinorum</i>	-13.8	+13.5	1181 ± 30 BP	1254–1467 cal CE	1104–1672 cal CE
RICH-21398	Pico (A)	<i>Pterodroma</i> sp. B	-14.0	+14.6	1654 ± 30 BP	760–1045 cal CE	665–1263 cal CE
WK-28566	Porto Santo (M)	<i>Pterodroma</i> sp. A	-13.4		7113 ± 37 BP	5607–5328 cal BCE	5739–5115 cal BCE

Table 2. Skull and mandible measurements of current species of *Pterodroma* (*Pt. madeira* from Madeira Island, *Pt. deserta* from Bugio Islet, Madeira Archipelago, *Pt. feae* from Cape Verde Archipelago, and *Pt. cahow* from Bermuda Island) and *Pt. zinorum* from Pico Island (Azores Archipelago), *Pterodroma* sp. A from North Dunes of Porto Santo (Madeira Archipelago), and *Pterodroma* sp. from Canary Islands. Sample size, mean length \pm standard error (mm), and range are given. Measurements as in Rando (2002).

	<i>Pt. madeira</i> Madeira	<i>Pt. deserta</i> Bugio	<i>Pt. feae</i> Cape Verde	<i>Pt. cahow</i> Bermuda	<i>Pt. zinorum</i> sp. nov. Pico	<i>Pterodroma</i> sp. A Porto Santo	<i>Pterodroma</i> sp. Canary Islands
Skull							
Length	(1) 66.7	(3) 74.2 ± 1.5 76.3–72.7	(1) 71.3	(1) 71.7	(3) 64.8 ± 1.8 66.4–62.3	(1) 72.2	(1) 76.7
Height	(1) 19.7	(3) 20.4 ± 0.1 20.3–20.4	(1) 20.2	(1) 20.1	(5) 20.3 ± 0.2 20–20.4		(2) 23.1 ± 0.7 23.6–22.6
Width	(1) 26.7	(3) 29.5 ± 0.7 30.2–28.6	(1) 28	(1) 28.8	(1) 26.3		(1) 30.1
Interorbital width	(1) 9.6	(3) 10.1 ± 0.1 10.1–10	(1) 10.1	(1) 8.5	(5) 9.7 ± 0.6 10.3–8.7	(2) 9.6 ± 0.1 9.6–9.5	(2) 10 ± 0.9 9.3–10.7
Maxilla dorsal length	(4) 32.6 ± 0.6 33.6–31.8	(3) 36.8 ± 0.4 37.1–36.2	(1) 35.3	(1) 35.2	(2) 31.4 ± 1.2 32.5–30.2	(1) 34.7	(1) 39.5
Maxilla length (from jugal articulation)	(3) 29.1 ± 0.4 29.6–28.6	(3) 31.5 ± 1 32.9–30.8	(1) 31.6	(1) 30.9	(2) 26.2 ± 0.2 26.4–26	(1) 29.9	(1) 33.1
Culmen length (from anterior edge of narina)	(6) 18.5 ± 0.4 19.2–18	(3) 20.2 ± 0.8 21.2–19.5	(1) 19.6	(1) 19.6	(3) 16.9 ± 0.9 17.9–15.8	(1) 18.8	(1) 21.3
Culmen width	(3) 11.1 ± 0.7 11.8–10.1	(3) 13.2 ± 0.5 13.8–12.7	(1) 13		(1) 11.1		(2) 14 ± 0.5 14.4–13.7
Width of the nasofrontal joint	(4) 8.8 ± 0.4 9.1–8.1	(3) 10.8 ± 0.2 11–10.6	(1) 10.4	(1) 10.1	(5) 8.5 ± 0.3 8.9–8.1	(1) 8.9	(2) 11.6 11.6–11.6
Mandible							
Length	(1) 52.35	(3) 58.4 ± 0.6 59.2–57.6	(1) 56.9	(1) 60.2			-

$\Lambda = 0.203$; d.f. = 4, 52; $P < .001$), and carpometacarpus ($U = -2.683$; $P < .007$, $U = -2.761$; $P < .006$) are smaller than *Pt. madeira* (tests performed between current bones of *Pt. madeira* and material from both Graciosa and Pico Islands, respectively). The new species has a shorter culmen ($U = -2.121$; $P = .034$) and a narrower nasofrontal joint ($U = -2.236$; $P = .025$) (tests performed between current bones of *Pt. madeira* and material from Pico) (Tables 2–4; Fig. 2).

Pterodroma deserta

Material and collection information: IMEDEA 109596, fragmented humerus. IMEDEA 109597, left complete ulna. IMEDEA 109595, right complete tarsometatarsus.

Horizon: Holocene–Recent.

Distribution: Bugio Islet (Desertas Islands, Madeira Archipelago).

Remarks: The material, probably very recent, is biometrically and morphologically indistinguishable from those of the current population of *Pt. deserta* that breed on this small islet.

Pterodroma sp. A

The fossil bones included under this name are slightly larger than those of current *Pt. madeira*, specifically the skull length (Tables 2, 4). This fact, together with its breeding habitat in sand dunes close to the coast, which is very different from that of *Pt. madeira* that breeds inland 1600 m above sea level on Madeira Island (Carboneras et al. 2020b), suggests that it could be a new species. However, its ascription of these materials to other current species cannot be ruled out due to the close morphology and

Table 3. Measurements of *Pterodroma* long bones from the Azores. Data of *Pterodroma zinorium* and *Pterodroma* sp. B from the islands of Graciosa, São Jorge, Pico, São Miguel and Terceira. Sample size, mean length ± standard error (mm), and range are given. Measurements as in Rando (2002).

	<i>Pt. zinorium</i> Graciosa	<i>Pt. zinorium</i> São Jorge	<i>Pt. zinorium</i> Pico	<i>Pt. zinorium</i> São Miguel	<i>Pterodroma</i> sp. A Graciosa	<i>Pterodroma</i> sp. B São Jorge	<i>Pterodroma</i> sp. B Pico	<i>Pterodroma</i> sp. B São Miguel	<i>Pterodroma</i> sp. B Terceira
Humerus length	(23) 74.3 ± 1.9 78.0–70.1	(22) 72.9 ± 1.2 75.8–70.5	(1) 74.1 (13) 75.5 ± 1.6 (1) 72.7	(1) 73.9 (1) 76.8	(8) 84.9 ± 1.8 87.2–81.4	(3) 88.4 ± 0.9 89.4–87.3	(4) 88.4 ± 0.6 89.0–87.4	(3) 89.6 ± 0 89.7–89.6	(1) 89.9
Ulna length	(9) 76.0 ± 1.8 77.8–72.2	(16) 73.3 ± 0.7 38.2–35.2	(22) 36.6 ± 0.8 (10) 26.6 ± 0.7 27.5–25.6	(1) 37.5 (2) 37.9 ± 0.1 37.9–37.8	(5) 86.4 ± 1.9 89.7–84.0	(6) 92.1 ± 0.8 93.3–91.0	(4) 92.4 ± 1 94.1–91.3	(2) 92.5 ± 1.2 93.7–91.3	(2) 92.5 ± 1.2 93.7–91.3
Carpometacarpus length	(16) 27.6 ± 0.8 29.2–26.1	(11) 49.3 ± 1.2 52.3–49.5	(10) 26.6 ± 0.7 27.5–25.6	(2) 42.3 ± 1 44.8–40.7	(3) 44.1 ± 1.1 45.5–42.8	(8) 44.2 ± 0.5 44.9–43.3	(2) 44.9 ± 0.3 45.2–44.6	(2) 44.5 ± 1.4 45.9–43.1	(2) 44.5 ± 1.4 45.9–43.1
Femur length	(13) 30.2 ± 0.6 31.3–28.7	(13) 29.7 ± 0.5 30.7–28.5	(1) 31.3 (1) 31.3	(2) 31.6 ± 0.3 32.0–31.3	(1) 30.4 (1) 30.4	(2) 32.5 ± 0.4 32.9–32.1	(2) 32.5 ± 0.4 32.9–32.1	(1) 29.9	(1) 29.9
Tibiotarsus length	(5) 50.7 ± 0.9 52.3–49.5	(13) 47.4 51.9–47.4	(11) 49.3 ± 1.2 51.9–47.4	(3) 56.7 ± 1.4 58.2–54.8	(1) 57.6 (1) 57.6	(1) 56.7 (1) 56.7	(1) 56.7 (1) 56.7	(1) 56.1	(1) 56.1
Tarsometatarsus length	(13) 30.2 ± 0.6 31.3–28.7	(13) 29.7 ± 0.5 30.7–28.5	(1) 31.3 37.0–37.0	(2) 37.5 ± 0.5 38.0–37.0	(1) 37.3 37.0–36.4	(2) 36.7 ± 0.3 37.0–36.4	(1) 34.3 (1) 34.3	(1) 34.3	(1) 35.4

Table 4. Measurements of *Pterodroma* long bones from the Madeiran Archipelago. *Pterodroma madeirensis* from Madeira Island (material of current specimens), *Pterodroma* sp. A and B from Madeira Island and Porto Santo (Madeira Archipelago) (fossil material), *Pt. deserta* from Bugio (Madeira Archipelago) (current and fossil material), *Pt. feae* from Cape Verde (material of current specimens), *Pt. cahow* from Bermuda (material of current specimens), and *Pterodroma* sp. from Canary Islands (fossil material). Sample size, mean length ± standard error (mm), and range are given. Measurements as in Rando (2002).

	<i>Pt. madeirensis</i> Madeira current	<i>Pterodroma</i> sp. B Madeira	<i>Pterodroma</i> sp. B Porto Santo	<i>Pterodroma</i> sp. A Porto Santo	<i>Pterodroma</i> sp. A North Dunes	<i>Pt. deserta</i> Bugio	<i>Pt. deserta</i> Bugio fossil	<i>Pt. feae</i> C.Verde current	<i>Pt. cahow</i> Bermuda current	<i>Pterodroma</i> sp. Canary Islands
Humerus length	(2) 79.3 ± 3.3 82.6–75.9	(22) 83.2 ± 1.8 86.4–79.3	(11) 87.5 ± 1.8 90.4–84.2	(6) 82.4 ± 2.2 85.1–79.6	(12) 82.2 ± 1.4 85.7–80.0	(4) 85.5 ± 1.4 87.6–83.9	(1) 84.6 87.6–83.9	(1) 84.9 87.8–85.3	(4) 86.6 ± 0.9 87.8–85.3	(2) 87.9 ± 0.3 88.2–87.7
Ulna length	(8) 83.6 ± 2.6 88.2–80	(27) 86.7 ± 1.8 89.4–82.7	(13) 89.4 ± 2.5 93.9–83.4	(6) 85.0 ± 2 87.8–81.8	(5) 84.4 ± 0.5 85.8–83.7	(4) 88.4 ± 1 90.2–87.5	(1) 89.1 91.9–84.3	(1) 87.3 91.9–84.3	(4) 88.6 ± 2.7 91.2–90.1	(2) 90.6 ± 0.5 91.2–90.1
Carpometacarpus length	(3) 40.1 ± 1 41.2–38.7	(34) 41.9 ± 1.1 43.5–39.2	(36) 43.8 ± 1 46–41.5	(12) 41.9 ± 1.1 43.6–39.9	(19) 42.2 ± 0.8 43.7–41	(4) 44.2 ± 0.5 45–43.6	(1) 40.7 46.1–41.0	(4) 43.3 ± 1.9 46.1–41.0	(3) 44.5 ± 0.5 45.1–44.0	(3) 44.5 ± 0.5 45.1–44.0
Femur Length	(1) 27.1 30.9–27.6	(12) 29.5 ± 1 30.3–29	(8) 29.6 ± 0.4 30.3–29	(1) 29.1 29.5–27.5	(11) 28.6 ± 0.6 30.9–29.8	(3) 30.4 ± 0.5 30.9–29.8	(1) 30.8 30.8–28.8	(3) 29.5 ± 0.9 30.8–28.8	(1) 32.3 30.8–28.8	(1) 32.3 30.8–28.8
Tibiotarsus length	(1) 49.9 56.5–51.2	(5) 54.0 ± 2.3 55.8–53.5	(9) 54.9 ± 0.7 52.5–51.2	(3) 51.7 ± 0.6 54.4–51.2	(8) 52.9 ± 0.9 55.7–53.0	(3) 54.5 ± 1.1 56.4–55.2	(1) 55.3 57.4–55.2	(3) 56.5 ± 1 57.4–55.2	(1) 56 57.4–55.2	(1) 56 57.4–55.2
Tarsometatarsus length	(12) 31.9 ± 0.6 33.0–30.7	(11) 35.3 ± 0.9 36.1–33.3	(18) 34.3 ± 0.9 35.9–32.1	(3) 32.7 ± 0.9 33.7–31.5	(14) 33.7 ± 0.8 34.6–32.1	(3) 35.2 ± 0.9 36.4–34.3	(1) 35.2 36.3–34.3	(3) 35.3 ± 0.8 36.3–34.3	(1) 35.4 36.3–34.3	(1) 35.4 36.3–34.3



Figure 3. Forelimb bones of *Pterodroma*. Humeri (caudal view), ulnae (ventral view) and carpometacarpi (ventral view), upper (u), middle (m) and lower (l) row respectively. Current species: *Pt. cahow* from Bermuda (A), *Pt. feae* from Cape Verde (B), *Pt. deserta* from Bugio (Madeira Archipelago) (C) and *Pt. madeira* from Madeira Island (D). Fossil material from Macaronesia: *Pterodroma* sp. A from Porto Santo (Madeira Archipelago) (E), and *Pt. zinorum* from Azores Islands, São Miguel (F), Graciosa (G), Pico (H) and Terceira (I), *Pterodroma* sp. B from Madeira Island (J), Porto Santo (K), and Azores Islands, São Miguel (L), Graciosa (M), Pico (N) and Terceira (O). Bones A (u), C (u), D (u), K (u), M (u), O (u), B (m), E (m), F (m), H (m), I (m), J (m), K (m), L (m), N (m), O (m), A (l), B (l), C (l), F (l), H (l), I (l) and M (l) are from the right side, the other are from the left. A, IMEDEA 94430; B, IMEDEA 103568; C, IMEDEA 94424; D (u and l) MMF 42034, D (m) IMEDEA 94425; E (u) MHMN 45597, E (m) MHMN 45466, E (l) MHMN 45644; F (u) MCMA 2271.017, F (m) MCMA 2270.017, F (l) MCMA 2263.017; G (u) MCMA 2011.017, G (m) MCMA 2037.017, G (l) MCMA 2062.017; H (u) MCMA 23021.023, H (m) MCMA 2302.017, H (l) MCMA 2306.017; I (u) MCMA 2258.017, I (m) MCMA 2260.017, I (l) 2260.017; J (u) MMF 45261, J (m) MMF 45333, J (l) MMF 45305; K (u) MHMN 46087, K (m) MHMN 46061, K (l) MHMN 46022; L (u and m) MCMA 2265017, L (l) MCMA 2264.017; M (u) MCMA 2287.017, M (m) MCMA 2171.017, M (l) MCMA 2184.017; N (u) MCMA 2282.017, N (m) MCMA 2287.017, N (l) MCMA 2283.017; O (u) MCMA 2255.017, O (m) MCMA 2254.017, O (l) MCMA 2253.017.

size that show the species inside this genus (Figs 3–7; Tables 2, 4; Supporting Information, Figs S1–S8).

Material and collection information: See Supporting Information, Table S1.

Horizon: Quaternary–Recent.

Distribution: Sandy sites of Porto Santo (Fonte da Areia and North Dunes) (Madeira Archipelago).

Remarks: The ¹⁴C date of the left ulna, radius, and carpometacarpus of one specimen of *Pterodroma* sp. A from North Dunes (WK-28566, 7113 ± 37 BP) was calibrated with 18 ± 141 ($\Delta R \pm SD$)

to provide the interval 5739–5115 cal BCE (Table 1). Some of the material overlaps with several extant North Atlantic species of *Pterodroma* (Tables 2, 4) but is closer to those of *Pt. madeira* than to the other species of bigger size (Fig. 3). No differences were found between ulna lengths from the current *Pt. madeira* and from North Dunes ($U = -0.645$; $P = .519$) and from Fonte da Areia ($U = 0$; $P = 1$) but the carpometacarpus length of the current *Pt. madeira* is smaller than those of North Dunes ($U = -2.021$; $P = .043$) and Fonte da Areia ($U = -2.44$; $P = .015$).

Material from Fonte da Areia and North Dunes (humerus, ulna, carpometacarpus, femur, tibiotarsus, and tarsometatarsus) is longer than those of *Pt. zinorum* from the Azores (Wilks'




Figure 4. Hindlimb bones of *Pterodroma*. Femora (caudal view), tibiotarsi (cranial view) and tarsometatarsi (dorsal view), upper (u), middle (m) and lower (l) row respectively. Current species: *Pt. cahow* from Bermuda (A), *Pt. feae* from Cape Verde (B), *Pt. deserta* from Bugio (Madeira Archipelago) (C) and *Pt. madeira* from Madeira Island (D). Fossil material from Macaronesia: *Pterodroma*, sp. A from Porto Santo (E), *Pt. zinorum* from Azores Islands, São Miguel (F), Graciosa (G), Pico (H), *Pterodroma*, sp. B from Madeira Island (J), Porto Santo (K), and the Azores São Miguel (L), Graciosa (M), Pico (N) and Terceira (O). Bones B (u), F (u), G (u), H (u), M (u), N (u), A (m), G (m), H (m), J (m), L (m), N (m), C (l), E (l), F (l), G (l), H (l), J (l), M (l) and N (l) are from the right side, the other are from the left. A, IMEDEA 94430; B, IMEDEA 103568; C IMEDEA 94424; D MMF 42034; E (u) MHMN 45781, E (m) MHMN 45981, E (l) MHMN 45611; F (u) MCMA 2270.017, F (l) MCMA 2270.017; G (u) MCMA 2114.017, G (m) MCMA 2127.017, G (l) MCMA 2137.017; H (u, m and l) MCMA 23020.023; J (u) MMF 45282, J (m) MMF 45423, J (l) MMF 45323; K (u) MMF 46151, K (m) MMF 46074, K (l) MMF 46153; L (m and l) MCMA 2273.017; M (u) MCMA 2209.017, M (m) MCMA 2210.017, M (l) MCMA 2214.017; N (u, m and l) MCMA 2292.017; O (u and m) MCMA 2253.017.

Lambda = 0.53; d.f. = 6, 17; $P < .001$) but many are smaller than those of *Pt. deserta* and *Pt. cahow* (Nichols and Mowbray, 1916) (humerus and ulna, Wilk's Lambda = 0.28; d.f. = 4, 30; $P = .001$; tibiotarsus, $U = -2.102$; $P = .036$ and $U = -2.57$; $P = .01$; and tarsometatarsus, $U = -2.488$; $P = .013$ in both tests).

The MANOVA performed with the humerus, ulna and carpometacarpus lengths from Porto Santo (Fonte da Areia and North Dunes) and fossil material from Madeira revealed differences (Wilk's Lambda = 0.455; d.f. = 6, 36; $P = .021$) with the ulnae from the Madeiran fossils being longer than those from North Dunes ($P = .009$) and Fonte da Areia ($P = .002$). However, there were no differences between the humerus

($P = .644$ and $P = .881$) and carpometacarpus ($P = .571$ and $P = .276$) of fossil material from Madeira and these sites of Porto Santo. The ANOVA performed between the femur length of fossil material from Madeira and Fonte da Areia indicates a longer size of the former ($F_{1,21} = 6.154$; $P = .022$). The tibiotarsus length of the Madeira fossils is similar to those from Fonte da Areia ($F_{1,11} = 1.283$; $P = .281$) and North Dunes ($U = -1.043$; $P = .297$), whereas the tarsometatarsus is longer in the fossil material from Madeira than Fonte da Areia ($F_{1,23} = 32.55$; $P < .001$) and North Dunes ($U = -2.413$; $P = .016$).

The bones from the Porto dos Frades site (Porto Santo) are longer than those from Fonte da Areia and North Dunes.



Figure 5. Articulated skeleton of *Pterodroma* sp. A. from North Dunes site in Porto Santo (Madeira Archipelago). Note the remains of an egg in the posterior side of the bird, documenting that *Pterodroma* sp. A was a burrow nesting species. Specimen exposed at Museu de História Natural do Jardim Botânico da Madeira, Funchal (without collection number).

The MANOVA performed with the humerus, ulna, and carpometacarpus lengths from North Dunes, Fonte da Areia, and Porto dos Frades (all sites from Porto Santo) show differences (Wilks's Lambda = 0.194; d.f. = 6, 34; $P < .001$) between the material from Porto dos Frades and Fonte da Areia ($P < .001$) and between Porto dos Frades and North Dunes ($P < .001$). The femurs from Porto dos Frades are longer than those of Fonte da Areia ($F_{1,17} = 14.63$; $P = .001$). The tibiotarsus from Porto dos Frades are longer than both from North dunes ($U = -2,49$; $P = .013$) and Fonte da Areia ($F_{1,15} = 23.25$; $P < .001$), like the tarsometatarsus ($U = -2211$; $P = .027$, and $F_{1,30} = 10.12$; $P < .003$, respectively).

The material from Fonte da Areia and North Dunes is smaller than those from the Azores (excluding those of *Pt. zinorum* n. sp.) (see Figs 3, 4, 6, 7; Tables 2–4). In addition, the *Pterodroma* bones from El Hierro (Canary Islands) are larger than those from Fonte da Areia ($U = -2727$; $P = .006$) and North Dunes ($U = -2598$; $P = .009$) (test performed with carpometacarpus lengths).

Pterodroma sp. B

This name includes material that is similar in morphology and close in size to those of the current *Pt. deserta*, *Pt. feae*, and *Pt. cahow*, so they could be included in any of these species.




Figure 6. Plot of mean \pm SD of forelimb bones length (humerus, ulna and carpometacarpus) and hindlimb bones length (femur, tibiotarsus and tarsometatarsus) of *Pterodroma* found in the fossil record of the Macaronesian islands (*Pt. zinorum*, *Pterodroma* sp. A, *Pterodroma* sp. B and *Pterodroma* sp. from El Hierro, Canary Islands. Green squares) and of current species (*Pt. madeira*, *Pt. deserta*, *Pt. feae*, *Pt. cahow* and *Pt. mollis*. Red circles). In the case of the material from Porto Santo (Madeira Archipelago) information about the site is indicated, for the other fossil material information on the island and archipelago is given. Data from Tables 3 and 4. Measurements of *Pt. rupinarum* and *Pt. mollis* are from Olson (1975) and Cooper & Tennyson (2008) respectively.

Nevertheless, it also remains possible (even probable) that they represent one or more cryptic unknown species (Figs 3, 4, 6, 7; Tables 2, 3, 4; Supporting Information, Figs S1–S8).

Material and collection information: See Supporting Information, Table S1.

Horizon: Late Quaternary–Recent.

Distribution: Porto dos Frades (Porto Santo, Madeira Archipelago), Madeira Island, and the Azores (São Miguel, Terceira, Graciosa, São Jorge, and Pico).

Remarks: The ^{14}C date of two fragments of ulna (proximal right and distal left) of one specimen of *Pterodroma* sp. B from Furnas das Torres, Pico (Azores): RICH-21398 and 1654 ± 30 BP, was calibrated with 38 ± 139 ($\Delta R \pm \text{SD}$) providing the interval 665–1263 CE (Table 1). These bones are longer than those of the current *Pt. madeira* (Tables 2–4; Figs 3, 4, 6, 7). The ANOVA performed with the ulna lengths of the current *Pt. madeira* and those from these material [from Madeira Island, Porto dos Frades (Porto Santo), and Graciosa, Pico, and São Miguel (Azores)] indicate a smaller size of *Pt. madeira* bones ($F_{5,62} = 18.526$; $P < .001$). The carpometacarpus from Madeira Island, Porto dos Frades (Porto Santo), Graciosa, and Pico are longer than those of the current *Pt. madeira* ($U = -2.170$; $P = .03$, $U = -2.846$; $P = .004$, $U = -2.536$; $P = .011$, and $U = -2.449$; $P = .014$, respectively). In addition, the

ANOVA performed with the tarsometatarsus lengths of the current *Pt. madeira* and those of *Pterodroma* sp. B from the island of Madeira sites and from Porto Santo (Porto dos Frades) indicate a small size of the former ($F_{2,41} = 53.326$; $P < .001$).

The material of this genus found in the Canary Islands (El Hierro) (Rando 2002) is of a similar size to those of this group (Tables 2, 4; Figs 3, 4, 6, 7). No differences were found in the carpometacarpus lengths between the fossil bones from El Hierro, the fossils from Pico ($U = -0.816$; $P = .414$), São Jorge ($U = -0.655$; $P = .513$), Porto dos Frades ($U = -1.528$; $P = .126$), and the bones of the current specimens of *Pt. cahow* ($U = -1.061$; $P = .289$) and *Pt. deserta* ($U = -0.707$; $P = .48$).

DISCUSSION

The material from Macaronesia and the fossil and archaeological records from northern Europe (Serjeantson 2005) and Gibraltar (Cooper 2005) indicate a previous wider distribution of *Pterodroma* in the North Atlantic Ocean. These records show at least two species of *Pterodroma* co-inhabiting several Macaronesian islands (Madeira, Porto Santo, São Miguel, Terceira, Pico, Faial, São Jorge, and Graciosa) in contrast to the current situation where the living species are the only one of this genus in the islands where they bred (Ramos et al. 2016).

Some current species of *Pterodroma* are impossible to discriminate through morphometric characters of their bones. For instance, the MANOVA of the humerus, ulna, and carpometacarpus




Figure 7. Plot of forelimb bones length (humerus+ulna+carpometacarpus) vs. hindlimb bones length (femur+tibiotarsus+tarsometatarsus) of *Pterodroma* found in the fossil record of the Macaronesian islands (*Pt. zinorum*, *Pterodroma* sp. A and *Pterodroma* sp. B, green squares) and of current species (*Pt. madeira*, *Pt. deserta*, *Pt. feae*, *Pt. cahow* and *Pt. mollis*, red circles). In the case of the material from Porto Santo (Madeira Archipelago) information about the site is given, for the other fossil material information on the island and archipelago is given in brackets. Data from Tables 3 and 4. Measurements of *Pt. rupinarum* and *Pt. mollis* are from Olson (1975) and Cooper & Tennyson (2008) respectively.

lengths of *Pt. deserta* (Bugio Islet, Madeira Archipelago) and *Pt. cahow* (Bermuda) did not reveal any differences (Wilks' Lambda = 0.376; d.f. = 3, 4; $P = .23$). Also, the femur ($U = -1.091$; $P = .275$), the tibiotarsus ($U = -1.528$; $P = .127$), and the tarsometatarsus lengths ($U = -0.218$; $P = .827$) of these species are not different. In addition, the lengths of limb bones of the solely available specimen of *Pt. feae* (Fogo, Cape Verde islands) fall inside, or close to, the range of variation of both species (Table 2, 4), thus it is difficult to determine the original diversity of this genus in the Atlantic through bone biometry. Nonetheless, the fossil records of *Pterodroma* in Macaronesia indicate the extinction of at least one species and a former much wider distribution of this genus in the Atlantic Ocean.

Considering that the fossil remains obtained of each identified entity (*Pterodroma* sp. A, *Pterodroma* sp. B, and *Pt. zinorum*) on each island document at least an ancient population, at least 11 populations of this genus have disappeared from the Azores: *Pt. zinorum* and *Pterodroma* sp. B in São Miguel, Terceira, Pico, São Jorge, and Graciosa, plus another population of *Pterodroma* sp. B in Faial. One population has disappeared from the island of Madeira (*Pterodroma* sp. B) plus two from Porto Santo (*Pterodroma* sp. A and B), whereas in the Canary Islands and in the Selvagens the *Pterodroma* fossils have been found only in one island on each archipelago (Rando 2002, Gangloff et al. 2013).

The current *Pterodroma* species are present on six Macaronesian islands (Ramos et al. 2016) (Fig. 1). The data indicate that of the 22 former populations of *Pterodroma* in four archipelagos, only six populations (27%) survive today in two archipelagos (Cape Verde and Madeira). This approach is very conservative, so it is probable that both the number of extinct populations and species is higher.

Due to their breeding-site fidelity (Ramos et al. 2016), the extinction of many populations indicates a loss of an important part of the original genetic diversity of *Pterodroma* in Macaronesia in contrast to *Calonectris*, another Procellariiform genus that breeds on all Macaronesian islands (Gómez-Díaz et al. 2006). In the Canary Islands, *C. borealis* (Cory, 1881) exhibits high values of genetic diversity with similarities between the ancient and the modern populations (Ramírez et al. 2013).

Gadfly petrels are currently absent as breeders in latitudes above 33°N (Madeira archipelago) (Fig. 1) showing a gap in its distribution in the North Atlantic Ocean (Ramos et al. 2017), so *Pterodroma* species may either have been exterminated by introduced predators or overlooked in the Azores, especially if they nest, as they often do elsewhere, in the winter (Bourne 1965, 1983). In 1990, a living *Pterodroma* was caught in a seabird colony in the Azores with similar measurements to those found on Bugio (*Pt. deserta*) but with a longer tail more like that of *Pt. cahow* and

a moderately developed brood patch, so it seems quite likely that this bird breeds in the Azores (Bibby and Nevo 1991). In 1993, another *Pterodroma* caught on an islet was identified as *Pt. feae* either as a vagrant from Bugio (currently *Pt. deserta*) or the Cape Verde Islands or as an Azores' bird from an unknown breeding population (Monteiro and Furness 1995). This bird was closely similar in measurements to the specimen caught by Bibby and Nevo (1991), except for a shorter tail length similar to that of the birds of Bugio (Monteiro and Furness 1995). The authors obtained a louse, identified as *Halipeurus theresiae* Timmermann, 1969, from this bird. This species has been previously found in *Pterodroma* from Bugio (Zonfrillo 1993), so it is possible that the Azores' population of *Pterodroma*, if it exists, may already be close to extinction as a consequence of rats (Monteiro and Furness 1995). The dimensions of the two birds from the Azores are larger than *Pt. madeira* and similar to *Pt. deserta*, *Pt. feae*, and *Pt. cahow*. The *Pterodroma* sp. B. remains from the Azores are similar to the latter three species (Figs 6, 7) so these captures may represent a vestigial population of this taxon from islets where rodents have not been introduced.

Within the genus *Pterodroma*, the current Macaronesian species are a monophyletic clade (Jesus *et al.* 2009), *Pt. madeira* the ancestral taxon, and *Pt. feae* and *Pt. deserta* sister-taxa that diverged later (Gangloff *et al.* 2013). The divergence of the three populations occurred without gene flow and very recently, the split between *Pt. feae*–*Pt. deserta* was estimated at 32 Kya (9–153 Kya) and the split between the ancestor *Pt. feae*–*Pt. deserta* and *Pt. madeira* occurred at 153 Kya (47–359 Kya). The populations of Madeira and Bugio Islands in the Madeira Archipelago (the closest populations) (Fig. 1) have been intensively monitored over the last three decades with no interbreeding between *Pt. madeira* and *Pt. deserta* observed (Gangloff *et al.* 2013).

Additionally, mitochondrial DNA sequences from *Pterodroma* bones from the Hebrides (Scotland), and from *Pt. rupinarum* from the island of Saint Helena indicate that both are part of a very young evolutionary radiation with a former wider distribution during the last millennia (Serjeantson 2005, Brace *et al.* 2014, Welch *et al.* 2014).

The bones of *Pt. feae* and *Pt. deserta* are similar morphologically but *Pt. madeira* is smaller ($F_{1,10} = 10.36$, $P = .009$ for ulna; $U = -2.121$, $P = .034$ for carpometacarpus; and $U = -2.598$, $P = .009$ for tarsometatarsus lengths) (Tables 2, 4; Figs 3, 4; Supporting Information, Figs S1–S8). The findings of recent extinct populations of *Pterodroma* close in size to both *Pt. feae*–*deserta*, and *Pt. madeira*, and the new species *Pt. zinorum*, smaller than *Pt. madeira*, show a more complex scenario in the phylogeography of this genus in Macaronesia.

The fossil record of *Pterodroma* in the Azores, with at least two species, seems to parallel the current situation in the Madeira archipelago where two current species of different sizes are breeding (*Pt. madeira* and *Pt. deserta* breeding on Madeira and Bugio Islands, respectively). As in the Madeira archipelago, the Azores were inhabited by gadfly petrels of two different body sizes. Both size classes are close in both archipelagos, so it is possible a close phylogenetic relationship between the species belonging to the same size class, like occur between the petrels of Bugio (*Pt. deserta*) and Cape Verde (*Pt. feae*) (Jesus *et al.* 2009, Gangloff *et al.* 2013). If this hypothesis is true, the larger petrels from the Azores could be part of the recent radiation *Pt. deserta*–*Pt. feae* in the Atlantic (Brace *et al.* 2014, Welch *et al.* 2014), and

Pt. zinorum could be a sister-taxon of the smaller *Pt. madeira*; only through ancient DNA techniques will it be possible to validate this hypothesis.

Radiocarbon data indicate that *Pt. zinorum* survived, at least, until the 12th century (Table 1), thus this species is the third extinct seabird during the last millennium in the western Palaearctic, together with the great auk *Pinguinus impennis* (Linnaeus, 1758) (Thomas *et al.* 2019) and the lava shearwater *Puffinus olsoni* McMinn *et al.*, 1990 (Rando and Alcover 2008). The 2σ calibration interval, of the dated bones of *Pt. zinorum* performed using the Marine20 calibration curve with 38 ± 139 ($\Delta R \pm SD$) (1104–1672 cal CE) overlaps with the Portuguese presence on the archipelago in the first half of the 15th century (Newitt 2005). The other radiocarbon dates of *Pterodroma* bones from Macaronesia indicate a late last record for *Pterodroma* sp. B from Pico (665–1263 cal CE) and a Mid-Holocene (5739–5115 cal BCE) last record for *Pterodroma* sp. A from Porto Santo (Madeira). These recent dates, especially the two former, point to historical events linked to human activities and to the introduction of alien mammals as the most probable cause for the extinction, like other Procellariiformes in Macaronesia and other islands around the world (Blackburn *et al.* 2004, Rando and Alcover 2008, 2010, Scofield 2009).

SUPPLEMENTARY DATA

Supplementary data are available at *Zoological Journal of the Linnean Society* online.

ACKNOWLEDGEMENTS

J.C. Illera (Oviedo), E. Blanco (Tenerife), M. Trias (Mallorca), and J. Femenias (Mallorca) assisted in the fieldwork. H.P. thanks his wife, O. Runze, for all her indefatigable help during the visits to Madeira and the Azores, including collecting bones included in this study, and the remaining authors also want to acknowledge her assistance also. Mr K. Groh (Hackenheim-Germany) provided several petrel bones used in this study. We received great logistic support from P. Borges (Universidade dos Açores in Terceira), S. Ávila, and all his team (Universidade dos Açores in São Miguel), Os Amigos dos Açores (especially D. Caetano in São Miguel and J. Melo in Santa Maria), and Os Montanheiros (Terceira, Pico and São Jorge). We are indebted to the curators of the Museu de História Natural do Funchal (MMF, Funchal), M. Bischoito, and the Museu Carlos Machado (MCMA, São Miguel), J.P. Constança, for their facilities and patience. A. Díaz (formerly at Institut Mediterrani d'Estudis Avançats) helped to catalogue the material. We are very grateful to three anonymous reviewers and the associate editor for their valuable contributions to this work.

AUTHOR CONTRIBUTIONS

Fieldwork and material curation J.C.R., H.P., F.P., E.T.R., and J.A.A.; data analysis and writing original draft J.C.R. and J.A.A.; review and editing J.C.R., H.P., F.P., E.T.R., and J.A.A.; project administration J.A.A.. All authors approved the final submitted version for publication.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest in relation to this work.

FUNDING

This paper is a result of the Research Projects CGL2012-38087 and CGL2016-79795R, 'Cambios holocénicos en la biodiversidad animal de las islas de la Macaronesia y de Baleares I and II' funded by the Agencia Estatal de Investigación (Ministerio de Economía, Industria y Competitividad) Fondo Europeo de Desarrollo Económico y Regional (Dirección General de Investigación Científica y Técnica, Ministerio de Economía y Competitividad). In addition, J.C.R. was funded by a research grant from the Spanish Ministry of Science, Innovation and Universities, and the European Regional Development Fund (PID2022-140091NB-I00).

DATA AVAILABILITY

Data will be made available on reasonable request.

APPENDIX

CURRENT COMPARATIVE MATERIAL EXAMINED

Pterodroma madeira (most of the available material are partial skeletons of birds depredated by rats and collected in breeding areas on the Island of Madeira): MMF 26304 (1–4), 26306, 30500, 31492, 31596, 31599, 31600, 31603 (1–2), 42033, 42034, 42037 (1–5), 42038, 42039, 42041, 42042.

Pterodroma deserta (Bugio Islet, Madeiran Archipelago): MMF 33794; IMEDEA 94424, 94426, 94427.

Pterodroma feae (Fogo Island, Cape Verde): IMEDEA 103568.

Pterodroma cahow (partial skeletons from Bermuda Island): MMF 2922-2; IMEDEA 94430 (1–30).

Pelagodroma marina (Selvagens Islands): DZUL 2396, 2397, 2405.

Hydrobates pelagicus (Canary Islands): DZUL 2843, 2847, 2849.

Bulweria bulwerii (Canary Islands): DZUL 393, 1379, 1832.

Puffinus puffinus (Canary Islands): DZUL 392, 2756.

Puffinus baroli (Canary Islands): DZUL 384, 826, 2255.

REFERENCES

- Bibby CJ, del Nevo AJ. A first record of *Pterodroma feae* from the Azores. *Bulletin of the British Ornithological Club* 1991;111:183–186.
- Blackburn TM, Cassey P, Duncan RP et al. Avian extinction and mammalian introductions on Oceanic Islands. *Science* 2004;305:1955–8. <https://doi.org/10.1126/science.1101617>
- Bourne WRP. The missing petrels. *Bulletin of the British Ornithologists' Club* 1965;85:97–105.
- Bourne WRP. The Soft-plumaged Petrel, the Gon-gon and the Freira, *Pterodroma mollis*, *P. feae* and *P. madeira*. *Bulletin of the British Ornithologists' Club* 1983;103:52–8.
- Boyer AG. Extinction patterns in the avifauna of the Hawaiian Islands. *Diversity and Distributions* 2008;14:509–17. <https://doi.org/10.1111/j.1472-4642.2007.00459.x>
- Brace S, Barnes I, Kitchener AC et al. Late Holocene range collapse in a former British seabird species. *Journal of Biogeography* 2014;41:1583–9. <https://doi.org/10.1111/jbi.12305>
- Bronk Ramsey C. Bayesian analysis of radiocarbon dates. *Radiocarbon* 2009;51:337–60. <https://doi.org/10.1017/s003382200033865>
- Brown TA, Vogel JS, Southon JR. Improved collagen extraction by modified Longin method. *Radiocarbon* 1998;30:171–7.
- Carboneras C, Jutglar F, Kirwan GM. Atlantic Petrel (*Pterodroma incerta*), version 1.0. In: del Hoyo J, Elliott A, Sargatal J et al. (eds), *Birds of the World*. Ithaca, NY, USA: Cornell Lab of Ornithology, 2020a. <https://doi.org/10.2173/bowatlpet1.01> (16 August 2024, date last accessed).
- Carboneras C, Jutglar F, Kirwan GM. Zino's Petrel (*Pterodroma madeira*), version 1.0. In: del Hoyo J, Elliott A, Sargatal J et al. (eds), *Birds of the World*. Ithaca, NY, USA: Cornell Lab of Ornithology, 2020b. <https://doi.org/10.2173/bowmadpet.01> (16 August 2024, date last accessed).
- Cooke R, Sayol F, Andermann T et al. Undiscovered bird extinctions obscure the true magnitude of human-driven extinction waves. *Nature Communications* 2023;14:8116. <https://doi.org/10.1038/s41467-023-43445-2>
- Cooper JH. Pigeons and pelagics: interpreting the Late Pleistocene avifaunas of the continental 'Island' of Gibraltar. *Monografies de la Societat d'Historia Natural de les Balears* 2005;12:101–12.
- Cooper JH, Tennyson JD. Wrecks and residents: the fossil gadfly petrels (*Pterodroma* spp.) of the Chatham Islands, New Zealand. *Oryctos* 2008;7:227–48.
- Curnutt J, Pimm SL. How many bird species in Hawaii and the Central Pacific before first contact? *Studies in Avian Biology* 2001;22:15–30.
- Duncan RP, Boyer AG, Blackburn TM. Magnitude and variation of prehistoric bird extinctions in the Pacific. *Proceedings of the National Academy of Sciences of the United States of America* 2013;110:6436–41. <https://doi.org/10.1073/pnas.1216511110>
- Ferrer Obiol J, Herranz JM, Paris JR et al. Species delimitation using genomic data to resolve taxonomic uncertainties in a speciation continuum of pelagic seabirds. *Molecular Phylogenetics and Evolution* 2023;179:107671. <https://doi.org/10.1016/j.ympev.2022.107671>
- Fromm A, Meiri S. Big, flightless, insular and dead: characterising the extinct birds of the Quaternary. *Journal of Biogeography* 2021;48:2350–9. <https://doi.org/10.1111/jbi.14206>
- Gangloff B, Zino F, Shirihai H et al. The evolution of north-east Atlantic gadfly petrels using statistical phylogeography. *Molecular Ecology* 2013;22:495–507. <https://doi.org/10.1111/mec.12119>
- Gómez-Díaz E, González-Solís J, Peinado MA et al. Phylogeography of the *Calonectris* shearwaters using molecular and morphometric data. *Molecular Phylogenetics and Evolution* 2006;41:322–32. <https://doi.org/10.1016/j.ympev.2006.05.006>
- Harrison P, Perrow M, Larsson H. *Seabirds. The New Identification Guide*. Barcelona: Lynx Editions, 2021.
- Hazevoet CJ. *The Birds of the Cape Verde Islands*. Tring: British Ornithologists' Union, 1995.
- Heaton T, Köhler P, Butzin M et al. Marine20—the marine radiocarbon age calibration curve (0–55 000 cal BP). *Radiocarbon* 2020;62:779–820.
- Hume JP, Walters M. *Extinct Birds*. London: T & A.D. Poyser, 2012.
- Jesus J, Menezes D, Gomes S et al. Phylogenetic relationships of gadfly petrels *Pterodroma* spp. from the Northeastern Atlantic Ocean: molecular evidence for specific status of Bugio and Cape Verde petrels and implications for conservation. *Bird Conservation International* 2009;19:199–214. <https://doi.org/10.1017/s0959270909008296>
- Kuroda N. Some osteological notes on Procellariiformes. *Japanese Journal of Ornithology* 1983;32:41–61. <https://doi.org/10.3838/jjo1915.32.41>
- Livezey BC, Zusi RL. Phylogeny of Neornithes. *Bulletin of Carnegie Museum of Natural History* 2006;37:1–544.
- Monteiro LR, Furness RS. Fea's petrel *pterodroma feae* in the Azores. *Bulletin of the British Ornithological Club* 1995;115:9–14.
- Newitt M. *A History of Portuguese Overseas Expansion, 1400–1668*. London: Routledge, 2005.
- Olson SL. Remarks on the generic characters of *Bulweria*. *Ibis* 1975b;117:111–3. <https://doi.org/10.1111/j.1474-919x.1975.tb04194.x>
- Olson SL. Early Pliocene Procellariiformes (Aves) from Langebaanweg, south-western Cape Province, South Africa. *Annals of the South African Museum* 1985;95:123–14.
- Olson SL, James HF. Fossil birds from the Hawaiian Islands: evidence for a wholesale extinction by man before Western contact. *Science* 1982;217:633–5. <https://doi.org/10.1126/science.217.4560.633>

- Olson SL, James HF. Descriptions of thirty-two new species of birds from the Hawaiian Islands: Part I, Non-Passeriformes. *Ornithological Monographs* 1991;45:1–88. <https://doi.org/10.2307/40166794>
- Pereira F, Nunes JC, Borges PAV, et al. *Catálogo das Cavidades Vulcânicas dos Açores (Algares Vulcânicos, Grutas Lávicas e de Erosão Marinha)*. Angra do Heroísmo: Associação Os Montanheiros, 2015.
- Ramírez O, Gómez-Díaz E, Olalde I et al. Population connectivity buffers genetic diversity loss in a seabird. *Frontiers in Zoology* 2013;10:28. <https://doi.org/10.1186/1742-9994-10-28>
- Ramos R, Ramírez I, Paiva VH et al. Global spatial ecology of three closely-related gadfly petrels. *Scientific Reports* 2016;6:23447. <https://doi.org/10.1038/srep23447>
- Ramos R, Carlile N, Madeiros J et al. It is the time for oceanic seabirds: tracking year-round distribution of gadfly petrels across the Atlantic Ocean. *Diversity and Distributions* 2017;23:794–805. <https://doi.org/10.1111/ddi.12569>
- Rando JC. New data of fossil birds from El Hierro (Canary Islands): probable causes of extinction and some biogeographical considerations. *Ardeola* 2002;49:39–49.
- Rando JC, Alcover JA. Evidence for a second western Palaearctic seabird extinction during the last millennium: the Lava Shearwater *Puffinus olsoni*. *Ibis* 2008;150:188–92. <https://doi.org/10.1111/j.1474-919x.2007.00741.x>
- Rando JC, Alcover JA. On the extinction of the Dune Shearwater (*Puffinus holeae*) from the Canary Islands. *Journal of Ornithology* 2010;151:365–9. <https://doi.org/10.1007/s10336-009-0463-6>
- Russell JC, Kueffer C. Island Biodiversity in the Anthropocene. *Annual Review of Environment and Resources* 2019;44:31–60. <https://doi.org/10.1146/annurev-environ-101718-033245>
- Sayol F, Steinbauer MJ, Blackburn TM et al. Anthropogenic extinctions conceal widespread evolution of flightlessness in birds. *Science Advances* 2020;6:eabb6095. <https://doi.org/10.1126/sciadv.abb6095>
- Scofield RP. Procellariiform extinctions in the Holocene: threat processes and wider ecosystem-scale implications. In: Turvey ST (ed.), *Holocene Extinctions*. Oxford: Oxford University Press, 2009, 151–66.
- Serjeantson D. Archaeological records of a gadfly petrel *Pterodroma* sp. from Scotland in the first millennium A.D. *Documenta Archaeobiologiae* 2005;3:235–46.
- Steadman DW. Prehistoric extinctions of pacific island birds: biodiversity meets zooarchaeology. *Science* 1995;267:1123–31. <https://doi.org/10.1126/science.267.5201.1123>
- Steadman DW. *Extinction and Biogeography of Tropical Pacific Birds*. Chicago: University Chicago Press, 2006.
- Stuiver M, Braziunas TF. Sun, ocean climate and atmospheric $^{14}\text{CO}_2$: an evaluation of causal relationships. *The Holocene* 1993;3:289–305. <https://doi.org/10.1177/095968369300300401>
- Thomas JE, Carvalho GR, Haile J et al. Demographic reconstruction from ancient DNA supports rapid extinction of the great auk. *eLife* 2019;8:e47509. <https://doi.org/10.7554/eLife.47509>
- Turvey ST. *Holocene Extinctions*. Oxford: Oxford University Press, 2009.
- Tyrberg T. Holocene avian extinctions. In: Turvey ST (ed.), *Holocene Extinctions*. Oxford: Oxford University Press, 2009, 63–106.
- van Strydonck M, Boudin M, Ervynck A. Humans and *Myotragus*: the issue of sample integrity in radiocarbon dating. *Monografies de la Societat d'Història Natural de les Balears* 2005;12:369–76.
- Warham, J. *The Behaviour, Population Biology and Physiology of the Petrels*. London: Academic Press, 1996.
- Welch AJ, Storrs LO, Fleischer RC. Phylogenetic relationships of the extinct St Helena petrel, *Pterodroma rupinarum* Olson, 1975 (Procellariiformes: Procellariidae), based on ancient DNA. *Zoological Journal of the Linnean Society* 2014;170:494–505.
- Winkler DW, Billerman SM, Lovette IJ. Shearwaters and petrels (Procellariidae), version 1.0. In: Billerman SM, Keeney BK, Rodewald PG et al. (eds), *Birds of the World*. Ithaca: Cornell Lab of Ornithology, 2020 (6 August 2024, date last accessed).
- Worthy TH, Holdaway RH. *Prehistoric Life of New Zealand. The Lost World of the Moa*. Bloomington: Indiana University Press, 2002.
- Zilhão J. Radiocarbon evidence for maritime pioneer colonization at the origins of farming in west Mediterranean Europe. *Proceedings of the National Academy of Sciences of the United States of America* 2001;98:14180–5. <https://doi.org/10.1073/pnas.241522898>
- Zino F, Brown R, Bischoff M. The separation of *Pterodroma madeira* (Zino's petrel) from *Pterodroma feae* (Feae's petrel) (Aves: Procellariidae). *Ibis* 2008;150:326–34. <https://doi.org/10.1111/j.1474-919x.2007.00794.x>
- Zonfrillo B. The relationships of the *Pterodroma* petrels from the Madeira archipelago inferred from their feather lice. *Boletim do Museu Municipal do Funchal* 1993;2:325–31.