

INTRODUCTION TO STOCHASTIC CALCULUS WITH APPLICATIONS

THIRD EDITION


Fima C Klebaner

Imperial College Press

INTRODUCTION TO
STOCHASTIC CALCULUS
WITH APPLICATIONS

THIRD EDITION

This page intentionally left blank



INTRODUCTION TO STOCHASTIC CALCULUS WITH APPLICATIONS

THIRD EDITION

Fima C Klebaner

Monash University, Australia

Published by

Imperial College Press
57 Shelton Street
Covent Garden
London WC2H 9HE

Distributed by

World Scientific Publishing Co. Pte. Ltd.
5 Toh Tuck Link, Singapore 596224
USA office: 27 Warren Street, Suite 401-402, Hackensack, NJ 07601
UK office: 57 Shelton Street, Covent Garden, London WC2H 9HE

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library.

INTRODUCTION TO STOCHASTIC CALCULUS WITH APPLICATIONS

Third Edition

Copyright © 2012 by Imperial College Press

All rights reserved. This book, or parts thereof, may not be reproduced in any form or by any means, electronic or mechanical, including photocopying, recording or any information storage and retrieval system now known or to be invented, without written permission from the Publisher.

For photocopying of material in this volume, please pay a copying fee through the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, USA. In this case permission to photocopy is not required from the publisher.

ISBN-13 978-1-84816-831-2
ISBN-10 1-84816-831-4
ISBN-13 978-1-84816-832-9 (pbk)
ISBN-10 1-84816-832-2 (pbk)

Typeset by Stallion Press
Email: enquiries@stallionpress.com

Printed in Singapore.

Contents

Preface	xi
1. Preliminaries From Calculus	1
1.1 Functions in Calculus	1
1.2 Variation of a Function	4
1.3 Riemann Integral and Stieltjes Integral	9
1.4 Lebesgue's Method of Integration	14
1.5 Differentials and Integrals	14
1.6 Taylor's Formula and Other Results	15
2. Concepts of Probability Theory	21
2.1 Discrete Probability Model	21
2.2 Continuous Probability Model	28
2.3 Expectation and Lebesgue Integral	33
2.4 Transforms and Convergence	37
2.5 Independence and Covariance	39
2.6 Normal (Gaussian) Distributions	41
2.7 Conditional Expectation	43
2.8 Stochastic Processes in Continuous Time	47
3. Basic Stochastic Processes	55
3.1 Brownian Motion	56
3.2 Properties of Brownian Motion Paths	63
3.3 Three Martingales of Brownian Motion	65
3.4 Markov Property of Brownian Motion	67

3.5	Hitting Times and Exit Times	69
3.6	Maximum and Minimum of Brownian Motion	71
3.7	Distribution of Hitting Times	73
3.8	Reflection Principle and Joint Distributions	74
3.9	Zeros of Brownian Motion — Arcsine Law	75
3.10	Size of Increments of Brownian Motion	78
3.11	Brownian Motion in Higher Dimensions	81
3.12	Random Walk	81
3.13	Stochastic Integral in Discrete Time	83
3.14	Poisson Process	86
3.15	Exercises	88
4.	Brownian Motion Calculus	91
4.1	Definition of Itô Integral	91
4.2	Itô Integral Process	100
4.3	Itô Integral and Gaussian Processes	103
4.4	Itô's Formula for Brownian Motion	106
4.5	Itô Processes and Stochastic Differentials	108
4.6	Itô's Formula for Itô Processes	112
4.7	Itô Processes in Higher Dimensions	118
4.8	Exercises	121
5.	Stochastic Differential Equations	123
5.1	Definition of Stochastic Differential Equations (SDEs)	123
5.2	Stochastic Exponential and Logarithm	129
5.3	Solutions to Linear SDEs	131
5.4	Existence and Uniqueness of Strong Solutions	134
5.5	Markov Property of Solutions	136
5.6	Weak Solutions to SDEs	137
5.7	Construction of Weak Solutions	139
5.8	Backward and Forward Equations	144
5.9	Stratonovich Stochastic Calculus	146
5.10	Exercises	148
6.	Diffusion Processes	151
6.1	Martingales and Dynkin's Formula	151
6.2	Calculation of Expectations and PDEs	155

6.3	Time-Homogeneous Diffusions	159
6.4	Exit Times from an Interval	163
6.5	Representation of Solutions of ODES	167
6.6	Explosion	168
6.7	Recurrence and Transience	170
6.8	Diffusion on an Interval	171
6.9	Stationary Distributions	172
6.10	Multi-dimensional SDEs	175
6.11	Exercises	183
7.	Martingales	185
7.1	Definitions	185
7.2	Uniform Integrability	187
7.3	Martingale Convergence	189
7.4	Optional Stopping	191
7.5	Localization and Local Martingales	197
7.6	Quadratic Variation of Martingales	200
7.7	Martingale Inequalities	203
7.8	Continuous Martingales — Change of Time	205
7.9	Exercises	211
8.	Calculus For Semimartingales	213
8.1	Semimartingales	213
8.2	Predictable Processes	214
8.3	Doob–Meyer Decomposition	215
8.4	Integrals with Respect to Semimartingales	217
8.5	Quadratic Variation and Covariation	220
8.6	Itô’s Formula for Continuous Semimartingales	222
8.7	Local Times	224
8.8	Stochastic Exponential	226
8.9	Compensators and Sharp Bracket Process	230
8.10	Itô’s Formula for Semimartingales	236
8.11	Stochastic Exponential and Logarithm	238
8.12	Martingale (Predictable) Representations	239
8.13	Elements of the General Theory	242
8.14	Random Measures and Canonical Decomposition	246
8.15	Exercises	249

9.	Pure Jump Processes	251
9.1	Definitions	251
9.2	Pure Jump Process Filtration	252
9.3	Itô's Formula for Processes of Finite Variation	253
9.4	Counting Processes	254
9.5	Markov Jump Processes	261
9.6	Stochastic Equation for Jump Processes	264
9.7	Generators and Dynkin's Formula	265
9.8	Explosions in Markov Jump Processes	267
9.9	Exercises	268
10.	Change of Probability Measure	269
10.1	Change of Measure for Random Variables	269
10.2	Change of Measure on a General Space	273
10.3	Change of Measure for Processes	276
10.4	Change of Wiener Measure	281
10.5	Change of Measure for Point Processes	283
10.6	Likelihood Functions	284
10.7	Exercises	287
11.	Applications in Finance: Stock and FX Options	289
11.1	Financial Derivatives and Arbitrage	289
11.2	A Finite Market Model	295
11.3	Semimartingale Market Model	299
11.4	Diffusion and the Black–Scholes Model	304
11.5	Change of Numeraire	312
11.6	Currency (FX) Options	315
11.7	Asian, Lookback, and Barrier Options	318
11.8	Exercises	321
12.	Applications in Finance: Bonds, Rates, and Options	325
12.1	Bonds and the Yield Curve	325
12.2	Models Adapted to Brownian Motion	327
12.3	Models Based on the Spot Rate	328
12.4	Merton's Model and Vasicek's Model	329
12.5	Heath–Jarrow–Morton (HJM) Model	333
12.6	Forward Measures — Bond as a Numeraire	338
12.7	Options, Caps, and Floors	341

12.8	Brace–Gatarek–Musiela (BGM) Model	343
12.9	Swaps and Swaptions	347
12.10	Exercises	349
13.	Applications in Biology	353
13.1	Feller’s Branching Diffusion	353
13.2	Wright–Fisher Diffusion	357
13.3	Birth–Death Processes	359
13.4	Growth of Birth–Death Processes	363
13.5	Extinction, Probability, and Time to Exit	366
13.6	Processes in Genetics	369
13.7	Birth–Death Processes in Many Dimensions	375
13.8	Cancer Models	377
13.9	Branching Processes	379
13.10	Stochastic Lotka–Volterra Model	386
13.11	Exercises	393
14.	Applications in Engineering and Physics	395
14.1	Filtering	395
14.2	Random Oscillators	402
14.3	Exercises	408
	Solutions to Selected Exercises	411
	References	429
	Index	435

This page intentionally left blank

Preface

Preface to the Third Edition

The third edition is revised and expanded in the direction of biological applications. New material on Birth–Death processes and their application in genetics and to cancer modelling is added. The focus, as throughout the text, is kept on the methods of Stochastic Calculus. When dealing with applications one quickly realizes that standard theory often cannot be applied directly, and certain extensions are needed. For example, the Branching Diffusion in biology and the Constant Elasticity of Variance (CEV) model in finance don't satisfy the conditions of the Theorem on existence and uniqueness of solutions and require extra analysis. Such extensions are developed in this edition.

It is anticipated that an accompanying text on Problems and Solutions in Stochastic Calculus will be published in 2013, which will enhance the present text. I am grateful to Kais Hamza, the co-author of the forthcoming text, for joining the project and helping with the present edition. I would like to acknowledge many readers, especially Jicheng Liu of Huazhong University of Science and Technology, China, who contributed comments and corrections. I hope that ideas of analysis of random processes by methods of Stochastic Calculus are made clear in the new applications.

Melbourne, 2012

Preface to the Second Edition

The second edition is revised, expanded and enhanced. This is now a more complete text in Stochastic Calculus, from both a theoretical and

an applications point of view. Changes came about as a result of using this book for teaching courses in Stochastic Calculus and Financial Mathematics over a number of years. All topics, starting from more elementary results on Normal correlation and optimal estimation, to the more advanced such as the change of time and the change of measure, are expanded with more worked out examples and exercises. Specific techniques, such as the change of numeraire and the stochastic logarithm, used in options pricing, are introduced. More advanced applications, such as pricing of exotic options, are covered in greater detail. A new chapter on bonds and interest rates contains derivations of the main pricing models, including currently used market models. The presentation of Applications in Finance is now more comprehensive and self-contained. The power of Stochastic Calculus is also demonstrated on models in biology introduced in the new edition: the age-dependent branching process and a stochastic model for competition of species. The mathematical theory of filtering is based on the methods of Stochastic Calculus. In the new edition, we derive stochastic equations for a non-linear filter first and then obtain the linear case, the celebrated Kalman–Bucy filter, as a corollary. It is important to note that all models arising in applications are treated rigorously and show in detail how to apply theoretical results presented earlier to the problem at hand. This approach might not make certain places easy reading, however, by using this book, the reader will acquire a working knowledge of Stochastic Calculus.

Preface to the First Edition

This book aims at providing a concise presentation of Stochastic Calculus with some of its applications in finance, engineering, and science.

During the past 20 years there has been an increasing demand for tools and methods of Stochastic Calculus in various disciplines. One of the greatest demands has come from the growing area of mathematical finance, where Stochastic Calculus is used for pricing and hedging of financial derivatives, such as options. In engineering, Stochastic Calculus is used in filtering and control theory. In physics stochastic Calculus is used to study the effects of random excitations on various physical phenomena. In biology, Stochastic Calculus is used to model the effects of stochastic variability in reproduction and environment on populations.

From an applied perspective Stochastic Calculus can be loosely described as a field of mathematics that is concerned with infinitesimal

calculus on non-differentiable functions. The need for this calculus comes from the necessity to include unpredictable factors into modelling. This is where probability comes in and the result is a calculus for random functions or stochastic processes.

This is a mathematical text that builds on theory of functions and probability and develops the martingale theory, which is highly technical. This text is aimed at gradually taking the reader from a fairly low technical level to a sophisticated one. This is achieved by making use of many solved examples. Every effort has been made to keep presentation as simple as possible while mathematically rigorous. Simple proofs are presented, but more technical proofs are left out and replaced by heuristic arguments with references to other more complete texts. This allows the reader to arrive at advanced results sooner. These results are required in applications. For example, the change of measure technique is needed in options pricing; calculations of conditional expectations with respect to a new filtration is needed in filtering. It turns out that completely unrelated applied problems have their solutions rooted in the same mathematical result. For example, the problem of pricing an option and the problem of optimal filtering of a noisy signal, both rely on the martingale representation property of Brownian motion.

This text presumes less initial knowledge than most texts on the subject (Métivier (1982), Dellacherie and Meyer (1982), Protter (1992), Liptser and Shiryaev (1989), Jacod and Shiryaev (1987), Karatzas and Shreve (1988), Stroock and Varadhan (1979), Revuz and Yor (2001), Rogers and Williams (1990)), however it still presents a fairly complete and mathematically rigorous treatment of Stochastic Calculus for both continuous processes and processes with jumps.

A brief description of the contents follows (for more details see the Table of Contents). The first two chapters describe the basic results in Calculus and Probability needed for further development. These chapters have examples but no exercises. Some more technical results in these chapters may be skipped and referred to later when needed.

In Chapter 3 the two main stochastic processes used in Stochastic Calculus are given: Brownian motion (for calculus of continuous processes) and Poisson process (for calculus of processes with jumps). Integration with respect to Brownian motion and closely related processes (Itô processes) is introduced in Chapter 4. It allows one to define a stochastic differential equation. Such equations arise in applications when random noise is introduced into ordinary differential equations. Stochastic differential equations are treated in Chapter 5. Diffusion processes arise as solutions to

stochastic differential equations, they are presented in Chapter 6. As the name suggests, diffusions describe a real physical phenomenon, and are met in many real-life applications. Chapter 7 contains information about martingales, examples of which are provided by Itô processes and compensated Poisson processes, introduced in earlier chapters. The martingale theory provides the main tools of Stochastic Calculus. These include optional stopping, localization and martingale representations. These are abstract concepts, but they arise in applied problems where their use is demonstrated. Chapter 8 gives a brief account of calculus for most general processes, called semimartingales. Basic results include Itô's formula and stochastic exponential. The reader has already met these concepts in Brownian motion calculus given in Chapter 4. Chapter 9 treats Pure Jump processes, where they are analyzed by using compensators. The change of measure is given in Chapter 10. This topic is important in options pricing, and for inference for stochastic processes. Chapters 11–14 are devoted to applications of Stochastic Calculus. Applications in Finance are given in Chapters 11 and 12, stocks and currency options (Chapter 11); bonds, interest rates and their options (Chapter 12). Applications in biology are given in Chapter 13. They include diffusion models, Birth–Death processes, age-dependent (Bellman–Harris) branching processes, and a stochastic version of the Lotka–Volterra model for competition of species. Chapter 14 gives applications in engineering and physics. Equations for a non-linear filter are derived and applied to obtain the Kalman–Bucy filter. Random perturbations to two-dimensional differential equations are given as an application in Physics. Exercises are given at the end of each chapter.

This text can be used for a variety of courses in Stochastic Calculus and financial mathematics. The application to finance is extensive enough to use it for a course in mathematical finance and for self-study. This text is suitable for advanced undergraduate students, graduate students as well as research workers and practitioners.

Acknowledgments

Thanks to Robert Liptser and Kais Hamza, who provided most valuable comments. Thanks to my colleagues and students from universities and banks. Thanks also to my family for being supportive and understanding.

Fima C. Klebaner
Melbourne, 2004

Chapter 1

Preliminaries From Calculus

Stochastic calculus deals with functions of time t , $0 \leq t \leq T$. In this chapter some concepts of the infinitesimal calculus used in the sequel are given.

1.1 Functions in Calculus

Continuous and Differentiable Functions

A function g is called continuous at the point $t = t_0$ if the increment of g over small intervals is small,

$$\Delta g(t) = g(t) - g(t_0) \rightarrow 0 \quad \text{as } \Delta t = t - t_0 \rightarrow 0.$$

If g is continuous at every point of its domain of definition, it is simply called continuous.

g is called differentiable at the point $t = t_0$ if at that point

$$\Delta g \sim C \Delta t \quad \text{or} \quad \lim_{\Delta t \rightarrow 0} \frac{\Delta g(t)}{\Delta t} = C,$$

this constant C is denoted by $g'(t_0)$. If g is differentiable at every point of its domain, it is called differentiable.

An important application of the derivative is a theorem on finite increments.

Theorem 1.1 (Mean Value Theorem): *If f is continuous on $[a, b]$ and has a derivative on (a, b) , then there is c , $a < c < b$, such that*

$$f(b) - f(a) = f'(c)(b - a). \tag{1.1}$$

Clearly, differentiability implies continuity, but not the other way around, as continuity states that the increment Δg converges to zero together with Δt , whereas differentiability states that this convergence is at the same rate or faster.

Example 1.1: The function $g(t) = \sqrt{t}$ is not differentiable at 0, as at this point

$$\frac{\Delta g}{\Delta t} = \frac{\sqrt{\Delta t}}{\Delta t} = \frac{1}{\sqrt{\Delta t}} \rightarrow \infty$$

as $t \rightarrow 0$.

It is surprisingly difficult to construct an example of a continuous function which is not differentiable at *any* point.

Example 1.2: An example of a continuous, nowhere differentiable function was given by Weierstrass in 1872: for $0 \leq t \leq 2\pi$

$$f(t) = \sum_{n=1}^{\infty} \frac{\cos(3^n t)}{2^n}. \quad (1.2)$$

We do not give a proof of these properties; a justification for continuity is given by the fact that if a sequence of continuous functions converges uniformly, then the limit is continuous and a justification for non-differentiability can be provided in some sense by differentiating term by term, which results in a divergent series.

To save repetition the following notations are used: *a continuous function f is said to be a C function; a differentiable function f with continuous derivative is said to be a C¹ function; a twice differentiable function f with continuous second derivative is said to be a C² function; etc.*

Right- and Left-Continuous Functions

We can rephrase the definition of a continuous function: a function g is called continuous at the point $t = t_0$ if

$$\lim_{t \rightarrow t_0} g(t) = g(t_0), \quad (1.3)$$

it is called right-continuous (left-continuous) at t_0 if the values of the function $g(t)$ approach $g(t_0)$ when t approaches t_0 from the right (left)

$$\lim_{t \downarrow t_0} g(t) = g(t_0), \quad (\lim_{t \uparrow t_0} g(t) = g(t_0)). \quad (1.4)$$

If g is continuous it is, clearly, both right- and left-continuous.

The left-continuous version of g , denoted by $g(t-)$, is defined by taking the left limit at each point,

$$g(t-) = \lim_{s \uparrow t} g(s). \quad (1.5)$$

From the definitions we have: g is left-continuous if $g(t) = g(t-)$.

The concept of $g(t+)$ is defined similarly,

$$g(t+) = \lim_{s \downarrow t} g(s). \quad (1.6)$$

If g is a right-continuous function then $g(t+) = g(t)$ for any t , so that $g_+ = g$.

Definition 1.2: A point t is called a discontinuity of the first kind or a jump point if both limits $g(t+)$ and $g(t-)$ exist and are not equal. The jump at t is defined as $\Delta g(t) = g(t+) - g(t-)$. Any other discontinuity is said to be of the second kind.

Example 1.3: The function $\sin(1/t)$ for $t \neq 0$ and 0 for $t = 0$ has discontinuity of the second kind at zero, because the limits from the right or the left do not exist.

An important result is that a function can have at most countably many jump discontinuities (see for example Hobson (1921)).

Theorem 1.3: *A function defined on an interval $[a, b]$ can have no more than countably many jumps.*

A function, of course, can have more than countably many discontinuities, but then they are not all jumps, i.e. they would not have limits from right or left.

Another useful result is that a derivative cannot have jump discontinuities at all.

Theorem 1.4: *If f is differentiable with a finite derivative $f'(t)$ in an interval, then at all points $f'(t)$ is either continuous or has a discontinuity of the second kind.*

PROOF. If t is such that $f'(t+) = \lim_{s \downarrow t} f'(s)$ exists (finite or infinite), then by the mean value theorem the same value is taken by the derivative from the right

$$f'(t) = \lim_{\Delta \downarrow 0} \frac{f(t + \Delta) - f(t)}{\Delta} = \lim_{\Delta \downarrow 0, t < c < t + \Delta} f'(c) = f'(t+).$$

Similarly for the derivative from the left, $f'(t) = f'(t-)$. Hence $f'(t)$ is continuous at t . The result follows. \square

This result explains why functions with continuous derivatives are sought as solutions to ordinary differential equations.

Functions Considered in Stochastic Calculus

Functions considered in stochastic calculus are functions without discontinuities of the second kind, that is, functions that have both right and left limits at any point of the domain and have one-sided limits at the boundary. These functions are called *regular* functions. It is often agreed to identify functions if they have the same right and left limits at any point.

The class $D = D[0, T]$ of right-continuous functions on $[0, T]$ with left limits has a special name, *càdlàg* functions (which is the abbreviation of “right continuous with left limits” in French). Sometimes these processes are called R.R.C. for regular right continuous. Note that this class of processes includes C , the class of continuous functions.

Let $g \in D$ be a *càdlàg* function, then by definition all the discontinuities of g are jumps. According to Theorem 1.3 such functions have no more than countably many discontinuities.

Remark 1.1: In stochastic calculus $\Delta g(t)$ usually stands for the size of the jump at t . In standard calculus $\Delta g(t)$ usually stands for the increment of g over $[t, t + \Delta]$, $\Delta g(t) = g(t + \Delta) - g(t)$. The meaning of $\Delta g(t)$ will be clear from the context.

1.2 Variation of a Function

If g is a function of real variable, its variation over the interval $[a, b]$ is defined as

$$V_g([a, b]) = \sup \sum_{i=1}^n |g(t_i^n) - g(t_{i-1}^n)|, \quad (1.7)$$

where the supremum is taken over partitions:

$$a = t_0^n < t_1^n < \cdots < t_n^n = b. \quad (1.8)$$

Clearly, (by the triangle inequality) the sums in (1.7) increase as new points are added to the partitions. Therefore variation of g is

$$V_g([a, b]) = \lim_{\delta_n \rightarrow 0} \sum_{i=1}^n |g(t_i^n) - g(t_{i-1}^n)|, \quad (1.9)$$

where $\delta_n = \max_{1 \leq i \leq n} (t_i - t_{i-1})$. If $V_g([a, b])$ is finite then g is said to be a function of finite variation on $[a, b]$. If g is a function of $t \geq 0$, then the variation function of g as a function of t is defined by

$$V_g(t) = V_g([0, t]).$$

Clearly, $V_g(t)$ is a non-decreasing function of t .

Definition 1.5: g is of finite variation if $V_g(t) < \infty$ for all t . g is of bounded variation if $\sup_t V_g(t) < \infty$, in other words, if for all t , $V_g(t) < C$, a constant independent of t .

Example 1.4:

1. If $g(t)$ is increasing then for any i , $g(t_i) > g(t_{i-1})$ resulting in a telescoping sum, where all the terms excluding the first and the last cancel out, leaving

$$V_g(t) = g(t) - g(0).$$

2. If $g(t)$ is decreasing then, similarly,

$$V_g(t) = g(0) - g(t).$$

Example 1.5: If $g(t)$ is differentiable with continuous derivative $g'(t)$, $g(t) = \int_0^t g'(s)ds$, and $\int_0^t |g'(s)|ds < \infty$, then

$$V_g(t) = \int_0^t |g'(s)|ds.$$

This can be seen by using the definition and the mean value theorem. $\int_{t_{i-1}}^{t_i} g'(s)ds = g'(\xi_i)(t_i - t_{i-1})$, for some $\xi_i \in (t_{i-1}, t_i)$. Thus $|\int_{t_{i-1}}^{t_i} g'(s)ds| = |g'(\xi_i)|(t_i - t_{i-1})$, and

$$\begin{aligned} V_g(t) &= \lim \sum_{i=1}^n |g(t_i) - g(t_{i-1})| = \lim \sum_{i=1}^n \left| \int_{t_{i-1}}^{t_i} g'(s)ds \right| \\ &= \sup \sum_{i=1}^n |g'(\xi_i)|(t_i - t_{i-1}) = \int_0^t |g'(s)|ds. \end{aligned}$$

The last equality is due to the last sum being a Riemann sum for the final integral.

Alternatively, the result can be seen from the decomposition of the derivative into the positive and negative parts,

$$g(t) = \int_0^t g'(s)ds = \int_0^t [g'(s)]^+ ds - \int_0^t [g'(s)]^- ds.$$

Notice that $[g'(s)]^-$ is zero when $[g'(s)]^+$ is positive, and the other way around. Using this one can see that the total variation of g is given by the sum of the variation of the above integrals. However, these integrals are monotone functions with the value zero at zero. Hence

$$\begin{aligned} V_g(t) &= \int_0^t [g'(s)]^+ ds + \int_0^t [g'(s)]^- ds \\ &= \int_0^t ([g'(s)]^+ + [g'(s)]^-) ds = \int_0^t |g'(s)|ds. \end{aligned}$$

Example 1.6: (Variation of a pure jump function)

If g is a regular right-continuous (*càdlàg*) function or regular left-continuous (*càglàd*), and changes only by jumps,

$$g(t) = \sum_{0 \leq s \leq t} \Delta g(s),$$

then it is easy to see from the definition that

$$V_g(t) = \sum_{0 \leq s \leq t} |\Delta g(s)|.$$

Example 1.7: The function $g(t) = t \sin(1/t)$ for $t > 0$, and $g(0) = 0$ is continuous on $[0, 1]$, differentiable at all points except zero, but has infinite variation on any interval that includes zero. Take the partition $1/(2\pi k + \pi/2), 1/(2\pi k - \pi/2)$, $k = 1, 2, \dots$.

The following theorem gives necessary and sufficient conditions for a function to have finite variation.

Theorem 1.6 (Jordan Decomposition): *Any function $g : [0, \infty) \rightarrow \mathbb{R}$ of finite variation can be expressed as the difference of two increasing functions*

$$g(t) = a(t) - b(t).$$

One such decomposition is given by

$$a(t) = V_g(t) \quad b(t) = V_g(t) - g(t). \quad (1.10)$$

It is easy to check that $b(t)$ is increasing and $a(t)$ is obviously increasing. The representation of a function of finite variation as the difference of two increasing functions is not unique. Another decomposition is

$$g(t) = \frac{1}{2}(V_g(t) + g(t)) - \frac{1}{2}(V_g(t) - g(t)).$$

The sum, the difference, and the product of functions of finite variation are also functions of finite variation. This is also true for the ratio of two functions of finite variation provided the modulus of the denominator is larger than a positive constant.

The following result follows by Theorem 1.3 and its proof is easy.

Theorem 1.7: *A finite variation function can have no more than countably many discontinuities. Moreover, all discontinuities are jumps.*

PROOF. It is enough to establish the result for monotone functions, since a function of finite variation is a difference of two monotone functions.

A monotone function has left and right limits at any point, therefore any discontinuity is a jump. The number of jumps of size greater or equal to $\frac{1}{n}$ is no more than $(g(b) - g(a))n$. The set of all jump points is a union of the sets of jump points with the size of the jumps greater than or equal to $\frac{1}{n}$. Since each such set is finite, the total number of jumps is at most countable. \square

A sufficient condition for a *continuous* function to be of finite variation is given by the following theorem, the proof of which is outlined in Example 1.5.

Theorem 1.8: *If g is continuous, g' exists and $\int |g'(t)|dt < \infty$ then g is of finite variation.*

Theorem 1.9 (Banach): *Let $g(t)$ be a continuous function on $[0, 1]$, and denote by $s(a)$ the number of t 's with $g(t) = a$. Then the variation of g is $\int_{-\infty}^{\infty} s(a)da$.*

Continuous and Discrete Parts of a Function

Let $g(t)$, $t \geq 0$, be a right-continuous increasing function. Then it can have at most countably many jumps, moreover, the sum of the jumps is finite over finite time intervals. Define the discontinuous part g^d of g by

$$g^d(t) = \sum_{s \leq t} (g(s) - g(s-)) = \sum_{0 < s \leq t} \Delta g(s), \quad (1.11)$$

and the continuous part g^c of g by

$$g^c(t) = g(t) - g^d(t). \quad (1.12)$$

Clearly, g^d changes only by jumps, g^c is continuous, and $g(t) = g^c(t) + g^d(t)$. Since a finite variation function is the difference of two increasing functions, the decomposition (1.12) holds for functions of finite variation. Although representation as the difference of increasing functions is not unique, decomposition (1.12) is essentially unique, in the sense that any two such decompositions differ by a constant. Indeed, if there were another such decomposition $g(t) = h^c(t) + h^d(t)$, then $h^c(t) - g^c(t) = g^d(t) - h^d(t)$, implying that $h^d - g^d$ is continuous. Hence h^d and g^d have the same set of jump points, and it follows that $h^d(t) - g^d(t) = c$ for some constant c .

Quadratic Variation

If g is a function of real variable, define its quadratic variation over the interval $[0, t]$ as the limit (when it exists)

$$[g](t) = \lim_{\delta_n \rightarrow 0} \sum_{i=1}^n (g(t_i^n) - g(t_{i-1}^n))^2, \quad (1.13)$$

where the limit is taken over partitions: $0 = t_0^n < t_1^n < \dots < t_n^n = t$, with $\delta_n = \max_{1 \leq i \leq n} (t_i^n - t_{i-1}^n)$.

Remark 1.2: Similarly to the concept of variation, there is a concept of Φ -variation of a function. If $\Phi(u)$ is a positive function, increasing monotonically with u , then the Φ -variation of g on $[0, t]$ is

$$V_\Phi[g] = \sup \sum_{i=1}^n \Phi(|g(t_i^n) - g(t_{i-1}^n)|), \quad (1.14)$$

where supremum is taken over *all* partitions. Functions with finite Φ -variation on $[0, t]$ form a class V_Φ . With $\Phi(u) = u$ one obtains the class VF of functions of finite variation, with $\Phi(u) = u^p$ one obtains the class of functions of p -th finite variation, VF_p . If $1 \leq p < q < \infty$, then finite p -variation implies finite q -variation.

The stochastic calculus definition of quadratic variation is different to the classical one with $p = 2$ (unlike for the first variation $p = 1$, when they are the same). In stochastic calculus the limit in (1.13) is taken over *shrinking* partitions with $\delta_n = \max_{1 \leq i \leq n} (t_i^n - t_{i-1}^n) \rightarrow 0$, and not over all possible partitions. We shall use only the stochastic calculus definition.

Quadratic variation plays a major role in stochastic calculus, but is hardly ever met in standard calculus due to the fact that smooth functions have zero quadratic variation.

Theorem 1.10: *If g is continuous and of finite variation then its quadratic variation is zero.*

PROOF.

$$\begin{aligned} [g](t) &= \lim_{\delta_n \rightarrow 0} \sum_{i=0}^{n-1} (g(t_{i+1}^n) - g(t_i^n))^2 \\ &\leq \lim_{\delta_n \rightarrow 0} \max_i |g(t_{i+1}^n) - g(t_i^n)| \sum_{i=0}^{n-1} |g(t_{i+1}^n) - g(t_i^n)| \\ &\leq \lim_{\delta_n \rightarrow 0} \max_i |g(t_{i+1}^n) - g(t_i^n)| V_g(t). \end{aligned}$$

Since g is continuous, it is uniformly continuous on $[0, t]$, hence $\lim_{\delta_n \rightarrow 0} \max_i |g(t_{i+1}^n) - g(t_i^n)| = 0$, and the result follows. \square

Note that there are functions with zero quadratic variation and infinite variation (called functions of *zero energy*).

Define the quadratic covariation (or simply covariation) of f and g on $[0, t]$ by the following limit (when it exists)

$$[f, g](t) = \lim_{\delta_n \rightarrow 0} \sum_{i=0}^{n-1} (f(t_{i+1}^n) - f(t_i^n)) (g(t_{i+1}^n) - g(t_i^n)), \quad (1.15)$$

when the limit is taken over partitions $\{t_i^n\}$ of $[0, t]$ with $\delta_n = \max_i (t_{i+1}^n - t_i^n)$.

The same proof as for Theorem 1.10 works for the following result.

Theorem 1.11: *If f is continuous and g is of finite variation, then their covariation is zero, $[f, g](t) = 0$.*

Let f and g be such that their quadratic variation is defined. By using simple algebra, one can see that covariation satisfies the following result.

Theorem 1.12 (Polarization Identity):

$$[f, g](t) = \frac{1}{2}([f + g, f + g](t) - [f, f](t) - [g, g](t)). \quad (1.16)$$

It is obvious that covariation is symmetric, $[f, g](t) = [g, f](t)$, it follows from (1.16) that it is linear, that is, for any constants α and β

$$[\alpha f + \beta g, h](t) = \alpha [f, h](t) + \beta [g, h](t). \quad (1.17)$$

Due to symmetry it is bilinear, that is, linear in both arguments. Thus the quadratic variation of the sum can be opened similarly to multiplication of sums $(\alpha_1 f + \beta_1 g)(\alpha_2 h + \beta_2 k)$. It follows from the definition of quadratic variation that it is a non-decreasing function in t and consequently it is of finite variation. According to the polarization identity, covariation is also of finite variation. More about quadratic variation is given in Chapters 4, 7, and 8.

1.3 Riemann Integral and Stieltjes Integral

Riemann Integral

The Riemann integral of f over interval $[a, b]$ is defined as the limit of Riemann sums

$$\int_a^b f(t) dt = \lim_{\delta \rightarrow 0} \sum_{i=1}^n f(\xi_i^n) (t_i^n - t_{i-1}^n), \quad (1.18)$$

where t_i^n 's represent partitions of the interval,

$$a = t_0^n < t_1^n < \cdots < t_n^n = b, \delta = \max_{1 \leq i \leq n} (t_i^n - t_{i-1}^n), \quad \text{and} \quad t_{i-1}^n \leq \xi_i^n \leq t_i^n.$$

It is possible to show that the Riemann integral is well-defined for continuous functions, and by splitting up the interval it can be extended to functions which are discontinuous at finitely many points.

Calculation of integrals is often done by using the antiderivative, and is based on the following result.

Theorem 1.13 (The Fundamental Theorem of Calculus): *If f is differentiable on $[a, b]$ and f' is Riemann integrable on $[a, b]$ then*

$$f(b) - f(a) = \int_a^b f'(s)ds.$$

In general, this result cannot be applied to discontinuous functions, see Example 1.8 below. For such functions a jump term must be added, see (1.20).

Example 1.8: Let $f(t) = 2$ for $1 \leq t \leq 2$, $f(t) = 1$ for $0 \leq t < 1$. Then $f'(t) = 0$ at all $t \neq 1$. $\int_0^t f'(s)ds = 0 \neq f(t) - f(0)$. f is continuous and is differentiable at all points but one, the derivative is integrable, but the function does not equal the integral of its derivative.

The main tools for calculations of Riemann integrals are change of variables and integration by parts. These are reviewed below in the more general framework of the Stieltjes integral.

Stieltjes Integral

The Stieltjes integral is an integral of the form $\int_a^b f(t)dg(t)$, where g is a function of finite variation. Since a function of finite variation is a difference of two increasing functions, it is sufficient to define the integral with respect to monotone functions.

Stieltjes Integral with Respect to Monotone Functions

The Stieltjes integral of f with respect to a monotone function g over an interval $[a, b]$ is defined as

$$\int_a^b f dg = \int_a^b f(t)dg(t) = \lim_{\delta \rightarrow 0} \sum_{i=1}^n f(\xi_i^n)(g(t_i^n) - g(t_{i-1}^n)), \quad (1.19)$$

with the quantities appearing in the definition being the same as those used above for the Riemann integral. This integral is a generalization of the Riemann integral, which is recovered when we take $g(t) = t$. This integral is also known as the Riemann–Stieltjes integral.

Particular Cases

If $g'(t)$ exists, and $g(t) = g(0) + \int_0^t g'(s)ds$, then it is possible to show that

$$\int_a^b f(t)dg(t) = \int_a^b f(t)g'(t)dt.$$

If $g(t) = \sum_{k=a}^{[t]} h(k)$ (a integer, and $[t]$ stands for the integer part of t) then

$$\int_a^b f(t)dg(t) = \sum_{a+1}^b f(k)h(k).$$

This property allows us to represent sums as integrals.

Example 1.9:

$$1. \quad g(t) = 2t^2 \quad \int_a^b f(t)dg(t) = 4 \int_a^b t f(t)dt.$$

$$2. \quad g(t) = \begin{cases} 0 & t < 0 \\ 2 & 0 \leq t < 1 \\ 3 & 1 \leq t < 2 \\ 5 & 2 \leq t \end{cases}$$

$$\int_{-\infty}^{\infty} f(t)dg(t) = 2f(0) + f(1) + 2f(2).$$

If, for example, $f(t) = t$ then $\int_{-\infty}^{\infty} t dg(t) = 5$. If $f(t) = (t+1)^2$ then $\int_{-\infty}^{\infty} (t+1)^2 dg(t) = 2 + 4 + 18 = 24$.

Let g be a function of finite variation and

$$g(t) = a(t) - b(t)$$

with $a(t) = V_g(t)$, $b(t) = V_g(t) - g(t)$, which are non-decreasing functions.

If

$$\int_0^t |f(s)|da(s) = \int_0^t |f(s)|dV_g(s) := \int_0^t |f(s)||dg(s)| < \infty$$

then f is Stieltjes integrable with respect to g and its integral is defined by

$$\int_{(0,t]} f(s)dg(s) = \int_{(0,t]} f(s)da(s) - \int_{(0,t]} f(s)db(s).$$

Notation: $\int_a^b f(s)dg(s) = \int_{(a,b]} f(s)dg(s)$.

Note: $\int_{(0,t]} dg(s) = g(t) - g(0)$ and $\int_{(0,t)} dg(s) = g(t-) - g(0)$.

If f is Stieltjes integrable with respect to a function g of finite variation, then the variation of the integral is

$$V(t) = \int_0^t |f(s)||dg(s)| = \int_0^t |f(s)|dV_g(s).$$

Impossibility of a Direct Definition of an Integral with Respect to Functions of Infinite Variation

In stochastic calculus we need to consider integrals with respect to functions of infinite variation. Such functions arise, for example, as models of stock prices. Integrals with respect to a function of infinite variation cannot be defined as a usual limit of approximating sums. The following result Theorem 1.14 explains, see for example Protter (1992).

Theorem 1.14: Let $\delta_n = \max_i(t_i^n - t_{i-1}^n)$ denote the largest interval in the partition of $[a, b]$. If

$$\lim_{\delta_n \rightarrow 0} \sum_{i=1}^n f(t_{i-1}^n)[g(t_i^n) - g(t_{i-1}^n)]$$

exists for any continuous function f then g must be of finite variation on $[a, b]$.

This shows that if g has infinite variation then the limit of the approximating sums does not exist for some functions f .

Integration by Parts

Let f and g be functions of finite variation. Denote here $\Delta g(s) = g(s) - g(s-)$, then (with integrals on $(a, b]$)

$$\begin{aligned} f(b)g(b) - f(a)g(a) &= \int_a^b f(s-)dg(s) + \int_a^b g(s-)df(s) + \sum_{a < s \leq b} \Delta f(s)\Delta g(s) \\ &= \int_a^b f(s-)dg(s) + \int_a^b g(s)df(s). \end{aligned} \quad (1.20)$$

The last equation is obtained by putting together the sum of jumps with one of the integrals.

Note that although the sum in (1.20) is written over uncountably many values $a < s \leq b$, it has at most countably many non-zero terms. This is because a finite variation function can have at most a countable number of jumps.

If f is continuous so that $f(s-) = f(s)$ for all s then the formula simplifies and in this case we have the familiar integration by parts formula

$$f(b)g(b) - f(a)g(a) = \int_a^b f(s)dg(s) + \int_a^b g(s)df(s).$$

Example 1.10: Let $g(s)$ be of finite variation, $g(0) = 0$, and consider $g^2(s)$. Using the integration by parts with $f = g$, we have

$$g^2(t) = 2 \int_0^t g(s-)dg(s) + \sum_{s \leq t} (\Delta g(s))^2.$$

In other words,

$$\int_0^t g(s-)dg(s) = \frac{g^2(t)}{2} - \frac{1}{2} \sum_{s \leq t} (\Delta g(s))^2.$$

Now using the formula (1.20) we also have

$$\int_0^t g(s)dg(s) = g^2(t) - \int_0^t g(s-)dg(s) = \frac{g^2(t)}{2} + \frac{1}{2} \sum_{s \leq t} (\Delta g(s))^2.$$

Thus it follows that

$$\int_0^t g(s-)dg(s) \leq \frac{g^2(t)}{2} \leq \int_0^t g(s)dg(s).$$

Change of Variables

Let f have a continuous derivative ($f \in C^1$) and g be of finite variation and continuous, then

$$f(g(t)) - f(g(0)) = \int_0^t f'(g(s))dg(s) = \int_{g(0)}^{g(t)} f'(u)du.$$

If g is of finite variation, has jumps, and is right-continuous, then

$$\begin{aligned} f(g(t)) - f(g(0)) &= \int_0^t f'(g(s-))dg(s) \\ &\quad + \sum_{0 < s \leq t} \left(f(g(s)) - f(g(s-)) - f'(g(s-))\Delta g(s) \right), \end{aligned}$$

where $\Delta g(s) = g(s) - g(s-)$ denotes the jump of g at s . This is known in stochastic calculus as Itô's formula.

Example 1.11: Take $f(x) = x^2$, then we obtain

$$g^2(t) - g^2(0) = 2 \int_0^t g(s-)dg(s) + \sum_{s \leq t} (\Delta g(s))^2.$$

Remark 1.3: Note that for a continuous f and finite variation g on $[0, t]$ the approximating sums converge as $\delta = \max_i(t_{i+1}^n - t_i^n) \rightarrow 0$,

$$\sum_i f(g(t_i^n))(g(t_{i+1}^n) - g(t_i^n)) \rightarrow \int_0^t f(g(s-))dg(s).$$

Remark 1.4: One of the shortcomings of Riemann or Stieltjes integrals is that they do not preserve the monotone convergence property, that is, for a sequence of functions $f_n \uparrow f$ does not necessarily follow that their integrals converge. The Lebesgue (or Lebesgue–Stieltjes) integral preserves this property.

1.4 Lebesgue’s Method of Integration

While Riemann sums are constructed by dividing the domain of integration on the x -axis and the interval $[a, b]$, into smaller subintervals, Lebesgue sums are constructed by dividing the range of the function on the y -axis, the interval $[c, d]$, into smaller subintervals $c = y_0 < y_1 < \dots < y_k < \dots < y_n = d$, and forming sums

$$\sum_{k=0}^{n-1} y_k \text{length}(\{t : y_k \leq f(t) < y_{k+1}\}).$$

The Lebesgue integral is the limit of the above sums as the number of points in the partition increases. It turns out that the Lebesgue integral is more general than the Riemann integral, and preserves convergence. This approach also allows integration of functions in abstract probability spaces more general than \mathbb{R} or \mathbb{R}^n ; it requires additional concepts and is made more precise in Chapter 2 (see Section 2.3).

Remark 1.5: In folklore the following analogy is used: imagine that money is spread out on a floor. In the Riemann method of integration, you collect the money as you progress in the room. In the Lebesgue method, first you collect \$100 bills everywhere you can find them, then \$50 bills, etc.

1.5 Differentials and Integrals

The differential $df(t)$ of a differentiable function f at t is defined as the *linear in Δt* part of the increment at t , $f(t + \Delta t) - f(t)$. If the differential of the independent variable is denoted $dt = \Delta t$, then $f(t + dt) - f(t) = df(t) +$ smaller order terms, and it follows from the existence of the derivative at t , that

$$df(t) = f'(t)dt. \quad (1.21)$$

If g is also a differentiable function of t , then $f(g(t))$ is differentiable, and

$$df(g(t)) = f'(g(t))g'(t)dt = f'(g(t))dg(t), \quad (1.22)$$

which is known as the *chain rule*.

Differential calculus is important in applications because many physical problems can be formulated in terms of differential equations. The main relation between the integral and the differential (or derivative) is given by the fundamental theorem of calculus, Theorem 1.13.

For differentiable functions, differential equations of the form

$$df(t) = \varphi(t)dw(t)$$

can be written in the integral form

$$f(t) = f(0) + \int_0^t \varphi(s)dw(s).$$

In stochastic calculus stochastic differentials do not formally exist and the random functions $w(t)$ are not differentiable at any point. By introducing a new (stochastic) integral, stochastic differential equations can be defined, and, by definition, solutions to these equations are given by the solutions to the corresponding stochastic integral equations.

1.6 Taylor's Formula and Other Results

This section contains Taylor's formula and conditions on functions used in results on differential equations. It may be treated as an appendix and referred to only when needed.

Taylor's Formula for Functions of One Variable

If we consider the increment of a function $f(x) - f(x_0)$ over the interval $[x_0, x]$, then provided $f'(x_0)$ exists, the differential at x_0 is the linear part in $(x - x_0)$ of this increment and it provides the first approximation to the increment. Taylor's formula gives a better approximation by taking higher order terms of powers of $(x - x_0)$ provided higher derivatives of f at x_0 exist. If f is a function of x with derivatives up to order $n + 1$, then

$$\begin{aligned} f(x) - f(x_0) &= f'(x_0)(x - x_0) + \frac{1}{2}f''(x_0)(x - x_0)^2 + \frac{1}{3!}f^{(3)}(x_0)(x - x_0)^3 \\ &\quad + \cdots + \frac{1}{n!}f^{(n)}(x_0)(x - x_0)^n + R_n(x, x_0), \end{aligned}$$

where R_n is the remainder, and $f^{(n)}$ is the derivative of $f^{(n-1)}$. The remainder can be written in the form

$$R_n(x, x_0) = \frac{1}{(n+1)!}f^{(n+1)}(\theta_n)(x - x_0)^{n+1}$$

for some point $\theta_n \in (x_0, x)$.

In our applications we shall use this formula with two terms.

$$f(x) - f(x_0) = f'(x)(x - x_0) + \frac{1}{2}f''(\theta)(x - x_0)^2, \quad (1.23)$$

for some point $\theta \in (x_0, x)$.

Taylor's Formula for Functions of Several Variables

Similarly to the one-dimensional case, Taylor's formula gives successive approximations to the increment of a multivariable function. A function of n real variables $f(x_1, x_2, \dots, x_n)$ is differentiable at point $\mathbf{x} = (x_1, x_2, \dots, x_n)$ if the increment at this point can be approximated by a linear part, which is the differential of f at \mathbf{x} .

$$\Delta f(\mathbf{x}) = \sum_{i=1}^n C_i \Delta x_i + o(\rho), \quad \text{when } \rho = \sqrt{\sum_{i=1}^n (\Delta x_i)^2} \quad \text{and} \quad \lim_{\rho \rightarrow 0} \frac{o(\rho)}{\rho} = 0. \quad (1.24)$$

If f is differentiable at $\mathbf{x} = (x_1, x_2, \dots, x_n)$, then in particular it is differentiable as a function of any one variable x_i at that point, when all the other coordinates are kept fixed. The derivative with respect to x_i is called the partial derivative $\partial f / \partial x_i$. Unlike in the one-dimensional case, the existence of all partial derivatives $\partial f / \partial x_i$ at \mathbf{x} is necessary but not sufficient for differentiability of f at \mathbf{x} . However, if all partial derivatives exist and are continuous at that point, then f is differentiable at that point, moreover, C_i in (1.24) is given by the value of $\partial f / \partial x_i$ at \mathbf{x} . If we define the differential of the independent variable as its increment $dx_i = \Delta x_i$, then we have

Theorem 1.15: *For f to be differentiable at a point, it is necessary that f has partial derivatives at that point, and it is sufficient that it has continuous partial derivatives at that point. If f is differentiable at \mathbf{x} , then its differential at \mathbf{x} is given by*

$$df(x_1, x_2, \dots, x_n) = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x_1, x_2, \dots, x_n) dx_i. \quad (1.25)$$

The first approximation of the increment of a differentiable function is the differential,

$$\Delta f(\mathbf{x}) \approx df(\mathbf{x}).$$

If f possesses higher order partial derivatives, then further approximation is possible and it is given by Taylor's formula. In stochastic calculus the second order approximation plays an important role.

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be C^2 , ($f(x_1, x_2, \dots, x_n)$ has continuous partial derivatives up to order two), $\mathbf{x} = (x_1, x_2, \dots, x_n)$, $\mathbf{x} + \Delta\mathbf{x} = (x_1 + \Delta x_1, x_2 + \Delta x_2, \dots, x_n + \Delta x_n)$. Then by considering the function of one variable $g(t) = f(\mathbf{x} + t\Delta\mathbf{x})$ for $0 \leq t \leq 1$, the following result is obtained.

$$\begin{aligned} & \Delta f(x_1, x_2, \dots, x_n) \\ &= f(\mathbf{x} + \Delta\mathbf{x}) - f(\mathbf{x}) \approx \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x_1, x_2, \dots, x_n) dx_i \\ &+ \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \frac{\partial^2 f}{\partial x_i \partial x_j}(x_1 + \theta \Delta x_1, \dots, x_n + \theta \Delta x_n) dx_i dx_j, \end{aligned} \quad (1.26)$$

where just like in the case of one variable the second derivatives are evaluated at some “middle” point, $(x_1 + \theta \Delta x_1, \dots, x_n + \theta \Delta x_n)$ for some $\theta \in (0, 1)$, and $dx_i = \Delta x_i$.

Lipschitz and Hölder Conditions

Lipschitz and Hölder conditions describe subclasses of continuous functions. They appear as conditions on the coefficients in the results on the existence and uniqueness of solutions of ordinary and stochastic differential equations.

Definition 1.16: f satisfies a Hölder condition (Hölder continuous) of order α , $0 < \alpha \leq 1$, on $[a, b]$ (\mathbb{R}) if there is a constant $K > 0$, so that for all $x, y \in [a, b]$ (\mathbb{R})

$$|f(x) - f(y)| \leq K|x - y|^\alpha. \quad (1.27)$$

A Lipschitz condition is a Hölder condition with $\alpha = 1$,

$$|f(x) - f(y)| \leq K|x - y|. \quad (1.28)$$

It is easy to see that a Hölder continuous of order α function on $[a, b]$ is also Hölder continuous of any lesser order.

Example 1.12: The function $f(x) = \sqrt{x}$ on $[0, \infty)$ is Hölder continuous with $\alpha = 1/2$ but is not Lipschitz, since its derivative is unbounded near zero. To see that it is Hölder, it is enough to show that for all $x, y \geq 0$ the following ratio is bounded,

$$\frac{|\sqrt{x} - \sqrt{y}|}{\sqrt{|x - y|}} \leq K. \quad (1.29)$$

It is an elementary exercise to establish that the left-hand side is bounded by dividing through by \sqrt{y} (if $y = 0$, then the bound is obviously one), and applying l'Hôpital's rule. Similarly $|x|^r$, $0 < r < 1$ is Hölder of order r .

A simple sufficient condition for a function to be Lipschitz is to be continuous and piecewise smooth; precise definitions follow.

Definition 1.17: f is smooth on $[a, b]$ if it possesses a continuous derivative f' on (a, b) such that the limits $f'(a+)$ and $f'(b-)$ exist.

Definition 1.18: f is piecewise continuous on $[a, b]$ if it is continuous on $[a, b]$ except possibly a finite number of points at which right-hand and left-hand limits exist.

Definition 1.19: f is piecewise smooth on $[a, b]$ if it is piecewise continuous on $[a, b]$ and f' exists and is also piecewise continuous on $[a, b]$.

Growth Conditions

The linear growth condition also appears in the results on existence and uniqueness of solutions of differential equations. $f(x)$ satisfies the linear growth condition if

$$|f(x)| \leq K(1 + |x|). \quad (1.30)$$

This condition describes the growth of a function for large values of x and states that f is bounded for small values of x .

Example 1.13: It can be shown that if $f(0, t)$ is a bounded function of t , $|f(0, t)| \leq C$ for all t , and $f(x, t)$ satisfies the Lipschitz condition in x uniformly in t , $|f(x, t) - f(y, t)| \leq K|x - y|$, then $f(x, t)$ satisfies the linear growth condition in x , $|f(x, t)| \leq K_1(1 + |x|)$.

The polynomial growth condition on f is the condition of the form

$$|f(x)| \leq K(1 + |x|^m), \quad \text{for some } K, m > 0. \quad (1.31)$$

Theorem 1.20 (Gronwall's Inequality): Let $g(t)$ and $h(t)$ be regular non-negative functions on $[0, T]$, then for any regular $f(t) \geq 0$ satisfying the inequality for all $0 \leq t \leq T$

$$f(t) \leq g(t) + \int_0^t h(s)f(s)ds \quad (1.32)$$

we have

$$f(t) \leq g(t) + \int_0^t h(s)g(s) \exp\left(\int_s^t h(u)du\right) ds. \quad (1.33)$$

This form is taken from Dieudonné (1960). In particular, if g is non-decreasing, the integral above simplifies to give

$$f(t) \leq g(t)e^{\int_0^t h(s)ds}. \quad (1.34)$$

In its simplest form when $g = A$ and $h = B$ are constants,

$$f(t) \leq Ae^{Bt}. \quad (1.35)$$

Solution of First Order Linear Differential Equations

Linear differential equations, by definition, are linear in the unknown function and its derivatives. A first order linear equation, in which the coefficient of $\frac{dx(t)}{dt}$ does not vanish, can be written in the form

$$\frac{dx(t)}{dt} + g(t)x(t) = k(t). \quad (1.36)$$

These equations are solved by using the integrating factor method. The integrating factor is the function $e^{G(t)}$, where $G(t)$ is chosen by $G'(t) = g(t)$. After multiplying both sides of the equation by $e^{G(t)}$, integrating, and solving for $x(t)$, we have

$$x(t) = e^{-G(t)} \int_0^t \left(e^{G(s)} k(s) \right) ds + x(0)e^{G(0)-G(t)}. \quad (1.37)$$

The integrating factor $G(t)$ is determined up to a constant, but it is clear from (1.37), that the solution $x(t)$ remains the same.

Further Results on Functions and Integration

Results given here are not required to understand subsequent material. Some of these involve the concepts of a set of zero Lebesgue measure. This is given in the next chapter (see Section 2.2); any countable set has Lebesgue measure zero, but there are also uncountable sets of zero Lebesgue measure. A partial converse to Theorem 1.8 also holds (see for example Saks (1964), and Freedman (1983), for the following results).

Theorem 1.21 (Lebesgue): *A finite variation function g on $[a, b]$ is differentiable almost everywhere on $[a, b]$.*

In what follows, sufficient conditions for a function to be Lipschitz and not to be Lipschitz are given.

1. If f is continuously differentiable on a finite interval $[a, b]$, then it is Lipschitz. Indeed, since f' is continuous on $[a, b]$, it is bounded, $|f'| \leq K$. Therefore

$$|f(x) - f(y)| = \left| \int_x^y f'(t) dt \right| \leq \int_x^y |f'(t)| dt \leq K|x - y|. \quad (1.38)$$

2. If f is continuous and piecewise smooth, then it is Lipschitz, the proof is similar to the above.
3. A Lipschitz function does not have to be differentiable, for example $f(x) = |x|$ is Lipschitz, but it is not differentiable at zero.

4. It follows from the definition of a Lipschitz function (1.28) that if it is differentiable, then its derivative is bounded by K .
5. A Lipschitz function has finite variation on finite intervals, since for any partition $\{x_i\}$ of a finite interval $[a, b]$,

$$\sum |f(x_{i+1}) - f(x_i)| \leq K \sum (x_{i+1} - x_i) = K(b - a). \quad (1.39)$$

6. As functions of finite variation have derivatives almost everywhere (with respect to Lebesgue measure), a Lipschitz function is differentiable almost everywhere.

(Note that functions of finite variation have derivatives which are integrable with respect to Lebesgue measure, but the function does not have to be equal to the integral of the derivative.)

7. A Lipschitz function multiplied by a constant, and a sum of two Lipschitz functions are Lipschitz functions. The product of two bounded Lipschitz functions is again a Lipschitz function.
8. If f is Lipschitz on $[0, N]$ for any $N > 0$ but with the constant K depending on N , then it is called locally Lipschitz. For example, x^2 is Lipschitz on $[0, N]$ for any finite N , but it is not Lipschitz on $[0, +\infty)$, since its derivative is unbounded.
9. If f is a function of two variables $f(x, t)$ and it satisfies the Lipschitz condition in x for all t , $0 \leq t \leq T$, with same constant K independent of t , it is said that f satisfies the Lipschitz condition in x uniformly in t , $0 \leq t \leq T$.

A necessary and sufficient condition for a function f to be Riemann integrable was given by Lebesgue (see for example Saks (1964), Freedman (1983)).

Theorem 1.22 (Lebesgue): *A necessary and sufficient condition for a function f to be Riemann integrable on a finite closed interval $[a, b]$ is that f is bounded on $[a, b]$ and almost everywhere continuous on $[a, b]$, that is, continuous at all points except possibly on a set of Lebesgue measure zero.*

Remark 1.6: (The following is not used anywhere in the book and is directed only to readers with knowledge of functional analysis).

Continuous functions on $[a, b]$ with the supremum norm $\|h\| = \sup_{x \in [a, b]} |h(x)|$ is a Banach space, denoted $C([a, b])$. By a result in functional analysis, any linear functional on this space can be represented as $\int_{(a,b)} h(x)dg(x)$ for some function g of finite variation. In this way, the Banach space of functions of finite variation on $[a, b]$ with the norm $\|g\| = V_g([a, b])$ can be identified with the space of linear functionals on the space of continuous functions, in other words, the dual space of $C([a, b])$.

Chapter 2

Concepts of Probability Theory

In this chapter, we give fundamental definitions of probabilistic concepts. Since the theory is more transparent in the discrete case, it is presented first. The most important concepts not met in elementary courses are the models for information, its flow, and conditional expectation. This is only a brief description and a more detailed treatment can be found in many books on probability theory, for example, Breiman (1968), Loeve (1978), and Durret (1991). Conditional expectation with its properties is central for further development, but some of the material in this chapter may be treated as an appendix.

2.1 Discrete Probability Model

A probability model consists of a filtered probability space on which variables of interest are defined. In this section we introduce a discrete probability model by using an example of discrete trading in stock.

Filtered Probability Space

A filtered probability space consists of the following: a sample space of elementary events, a field of events, a probability defined on that field, and a filtration of increasing subfields.

Sample Space

Here, we describe a finite sample space. Consider, for example, a single stock with price S_t at time $t = 1, 2, \dots, T$. Denote by Ω the set of all possible

values of stock during these times.

$$\Omega = \{\omega : \omega = (S_1, S_2, \dots, S_T)\}.$$

If we assume that the stock price can go up by a factor u and down by a factor d , then the relevant information reduces to the knowledge of the movements at each time.

$$\Omega = \{\omega : \omega = (a_1, a_2, \dots, a_T)\} \quad a_t = u \text{ or } d.$$

To model uncertainty about the price in the future, we “list” all possible future prices, and call them possible states of the world. The unknown future is just one of many possible outcomes called the true state of the world. As time passes more and more information is revealed about the true state of the world. At time $t = 1$ we know prices S_0 and S_1 . Thus the true state of the world lies in a smaller set, a subset of Ω , $A \subset \Omega$. After observing S_1 we know which prices did not happen at time 1. Therefore we know that the true state of the world is in A and not in $\Omega \setminus A = \bar{A}$.

Fields of Events

Define by \mathcal{F}_t the information available to investors at time t , which consists of stock prices before and at time t .

For example, when $T = 2$, at $t = 0$ we have no information about S_1 and S_2 , and $\mathcal{F}_0 = \{\emptyset, \Omega\}$; all we know is that a true state of the world is in Ω . Consider the situation at $t = 1$. Suppose at $t = 1$ stock went up by u . Then we know that the true state of the world is in A , and not in its complement \bar{A} , where

$$A = \{(u, S_2), S_2 = u \text{ or } d\} = \{(u, u), (u, d)\}.$$

Thus our information at time $t = 1$ is

$$\mathcal{F}_1 = \{\emptyset, \Omega, A, \bar{A}\}.$$

Note that $\mathcal{F}_0 \subset \mathcal{F}_1$, since we do not forget the previous information.

At time t investors know which part of Ω contains the true state of the world. \mathcal{F}_t is called a *field* or *algebra* of sets.

\mathcal{F} is a field if

1. $\emptyset, \Omega \in \mathcal{F}$.
2. $A \in \mathcal{F}$, and $B \in \mathcal{F}$ then $A \cup B \in \mathcal{F}$, $A \cap B \in \mathcal{F}$, $A \setminus B \in \mathcal{F}$.

Example 2.1: (Examples of fields)

It is easy to verify that any of the following is a field of sets:

1. $\{\Omega, \emptyset\}$ is called the trivial field \mathcal{F}_0 .
2. $\{\emptyset, \Omega, A, \bar{A}\}$ is called the field generated by set A and denoted by \mathcal{F}_A .
3. $\{A : A \subseteq \Omega\}$ is the field of all the subsets of Ω . It is denoted by 2^Ω .

A partition of Ω is a collection of exhaustive and mutually exclusive subsets,

$$\{D_1, \dots, D_k\}, \text{ such that } D_i \cap D_j = \emptyset, \quad \text{and} \quad \bigcup_i D_i = \Omega.$$

The field generated by the partition is the collection of all finite unions of D_j s and their complements. These sets are like the basic building blocks for the field. If Ω is finite, then any field is generated by a partition.

If one field is included in the other, $\mathcal{F}_1 \subset \mathcal{F}_2$, then any set from \mathcal{F}_1 is also in \mathcal{F}_2 . In other words, a set from \mathcal{F}_1 is either a set or a union of sets from the partition generating \mathcal{F}_2 . This means that the partition that generates \mathcal{F}_2 has “finer” sets than the ones that generate \mathcal{F}_1 .

Filtration

A filtration \mathbb{F} is the collection of fields,

$$\mathbb{F} = \{\mathcal{F}_0, \mathcal{F}_1, \dots, \mathcal{F}_t, \dots, \mathcal{F}_T\} \quad \mathcal{F}_t \subset \mathcal{F}_{t+1}.$$

\mathbb{F} is used to model a flow of information. As time passes, an observer knows more and more detailed information, that is, finer and finer partitions of Ω . In the example of the price of stock, \mathbb{F} describes how the information about prices is revealed to investors.

Example 2.2: $\mathbb{F} = \{\mathcal{F}_0, \mathcal{F}_A, 2^\Omega\}$ is an example of filtration.

Stochastic Processes

If Ω is a finite sample space, then a function X defined on Ω attaches numerical values to each $\omega \in \Omega$. Since Ω is finite, X takes only finitely many values x_i , $i = 1, \dots, k$.

If a field of events \mathcal{F} is specified, then any set in it is called a *measurable* set. If $\mathcal{F} = 2^\Omega$, then any subset of Ω is measurable.

A function X on Ω is called \mathcal{F} -measurable or a *random variable* on (Ω, \mathcal{F}) if all the sets $\{X = x_i\}$, $i = 1, \dots, k$, are members of \mathcal{F} . This means that if we have the information described by \mathcal{F} , that is, we know which event in \mathcal{F} has occurred, then we know which value of X has occurred. Note that if $\mathcal{F} = 2^\Omega$, then any function on Ω is a random variable.

Example 2.3: Consider the model for trading in stock, $t = 1, 2$, where at each time the stock can go up by the factor u or down by the factor d .

$\Omega = \{\omega_1 = (u, u), \omega_2 = (u, d), \omega_3 = (d, u), \omega_4 = (d, d)\}$. Take $A = \{\omega_1, \omega_2\}$, which is the event that at $t = 1$ the stock goes up. $\mathcal{F}_1 = \{\emptyset, \Omega, A, \bar{A}\}$, and $\mathcal{F}_2 = 2^\Omega$ contains all 16 subsets of Ω . Consider the following functions on Ω : $X(\omega_1) = X(\omega_2) = 1.5$, $X(\omega_3) = X(\omega_4) = 0.5$. X is a random variable on \mathcal{F}_1 . Indeed, the set $\{\omega : X(\omega) = 1.5\} = \{\omega_1, \omega_2\} = A \in \mathcal{F}_1$. Also, $\{\omega : X(\omega) = 0.5\} = \bar{A} \in \mathcal{F}_1$.

If $Y(\omega_1) = (1.5)^2$, $Y(\omega_2) = 0.75$, $Y(\omega_3) = 0.75$, and $Y(\omega_4) = 0.5^2$, then Y is not a random variable on \mathcal{F}_1 , it is not \mathcal{F}_1 -measurable. Indeed, $\{\omega : Y(\omega) = 0.75\} = \{\omega_2, \omega_3\} \notin \mathcal{F}_1$. Y is \mathcal{F}_2 -measurable.

Definition 2.1: A stochastic process is a collection of random variables $\{X(t)\}$. For any fixed t , $t = 0, 1, \dots, T$, $X(t)$ is a random variable on (Ω, \mathcal{F}_T) .

A stochastic process is called adapted to filtration \mathbb{F} if for all $t = 0, 1, \dots, T$, $X(t)$ is a random variable on \mathcal{F}_t , that is, if $X(t)$ is \mathcal{F}_t -measurable.

Example 2.4: (Example 2.3 continued)

$X_1 = X$, $X_2 = Y$ is a stochastic process adapted to $\mathbb{F} = \{\mathcal{F}_1, \mathcal{F}_2\}$. This process represents stock prices at time t under the assumption that the stock can appreciate or depreciate by 50% in a unit of time.

Field Generated by a Random Variable

Let $(\Omega, 2^\Omega)$ be a sample space with the field of all events, and X be a random variable with values x_i , $i = 1, 2, \dots, k$. Consider sets

$$A_i = \{\omega : X(\omega) = x_i\} \subseteq \Omega.$$

These sets form a partition of Ω , and the field generated by this partition is called the field generated by X . It is the smallest field that contains all the sets of the form $A_i = \{X = x_i\}$ and it is denoted by \mathcal{F}_X or $\sigma(X)$. The field generated by X represents the information we can extract about the true state ω by observing X .

Example 2.5: (Example 2.3 continued)

$$\{\omega : X(\omega) = 1.5\} = \{\omega_1, \omega_2\} = A, \{\omega : X(\omega) = 0.5\} = \{\omega_3, \omega_4\} = \bar{A}.$$

$$\mathcal{F}_X = \mathcal{F}_1 = \{\emptyset, \Omega, A, \bar{A}\}.$$

Filtration Generated by a Stochastic Process

Given (Ω, \mathcal{F}) and a stochastic process $\{X(t)\}$, let $\mathcal{F}_t = \sigma(\{X_s, 0 \leq s \leq t\})$ be the field generated by random variables X_s , $s = 0, \dots, t$. It is all the information available from the observation of the process up to time t . Clearly, $\mathcal{F}_t \subseteq \mathcal{F}_{t+1}$, so that these fields form a filtration. This filtration is called the natural filtration of the process $\{X(t)\}$.

If $A \in \mathcal{F}_t$, then by observing the process from 0 to t we know at time t whether the true state of the world is in A or not. We illustrate this in our financial example.

Example 2.6: Take $T = 3$ and assume that at each trading time the stock can go up by the factor u or down by d .

$$\begin{array}{c} \boxed{\begin{matrix} u & u & u \\ u & u & d \end{matrix}} \quad \boxed{\begin{matrix} d & u & u \\ d & u & d \end{matrix}} \quad \bar{B} \\ \Omega = \boxed{\begin{matrix} u & d & u \\ u & d & d \end{matrix}} \quad \boxed{\begin{matrix} d & d & u \\ d & d & d \end{matrix}} \quad B \\ A \quad \bar{A} \quad | \end{array}$$

Look at the sets generated by information about S_1 . This is a partition of Ω , $\{A, \bar{A}\}$. Together with the empty set and the whole set, this is the field \mathcal{F}_1 . Sets generated by information about S_2 are B and \bar{B} . Thus the sets formed by knowledge of S_1 and S_2 are the partition of Ω , consisting of all intersections of the above sets. Together with the empty set and the whole set this is the field \mathcal{F}_2 . Clearly, any set in \mathcal{F}_1 is also in \mathcal{F}_2 , for example, $A = (A \cap B) \cup (A \cap \bar{B})$. Similarly, if we add information about S_3 we obtain all the elementary sets, ω 's, and hence all subsets of Ω , $\mathcal{F}_3 = 2^\Omega$. In particular, we will know the true state of the world when $T = 3$.

$\mathcal{F}_0 \subset \mathcal{F}_1 \subset \mathcal{F}_2 \subset \mathcal{F}_3$ is the filtration generated by the price process $\{S_t, t = 1, 2, 3\}$.

Predictable Processes

Suppose that a filtration $\mathbb{F} = (\mathcal{F}_0, \mathcal{F}_1, \dots, \mathcal{F}_t, \dots, \mathcal{F}_T)$ is given. A process H_t is called *predictable* (with respect to this filtration) if for each t , H_t is \mathcal{F}_{t-1} -measurable, that is, the value of the process H at time t is determined by the information up to and including time $t - 1$. For example, the number of shares held at time t is determined on the basis of information up to and including time $t - 1$. Thus this process is predictable with respect to the filtration generated by the stock prices.

Stopping Times

τ is called a random time if it is a non-negative random variable, which can also take value ∞ on (Ω, \mathcal{F}_T) . Suppose that a filtration $\mathbb{F} = (\mathcal{F}_0, \mathcal{F}_1, \dots, \mathcal{F}_t, \dots, \mathcal{F}_T)$ is given. τ is called a *stopping time* with respect to this filtration if for each $t = 0, 1, \dots, T$ the event

$$\{\tau \leq t\} \in \mathcal{F}_t. \tag{2.1}$$

This means that by observing the information contained in \mathcal{F}_t we can decide whether the event $\{\tau \leq t\}$ has or has not occurred. If the filtration \mathbb{F} is generated by $\{S_t\}$, then by observing the process up to time t ,

S_0, S_1, \dots, S_t , we can decide whether the event $\{\tau \leq t\}$ has or has not occurred.

Probability

If Ω is a finite sample space, then we can assign to each outcome ω a probability, $P(\omega)$, that is, the likelihood of it occurring. This assignment can be arbitrary. The only requirement is that $P(\omega) \geq 0$ and $\sum P(\omega) = P(\Omega) = 1$.

Example 2.7: Take $T = 2$ in our basic example 2.3. If the stock goes up or down independently of its value and if, say, the probability to go up is 0.4, then

$$\begin{array}{llll} \Omega = & \{(u,u); (u,d); (d,u); (d,d)\} \\ P(\omega) & 0.16 & 0.24 & 0.24 & 0.36 \end{array}.$$

Distribution of a Random Variable

Since a random variable X is a function from Ω to \mathbb{R} , and Ω is finite, X can take only finitely many values, as the set $X(\Omega)$ is finite. Denote these values by x_i , $i = 1, 2, \dots, k$. The collection of probabilities p_i of sets $\{X = x_i\} = \{\omega : X(\omega) = x_i\}$ is called the probability distribution of X ; for $i = 1, 2, \dots, k$

$$p_i = P(X = x_i) = \sum_{\omega: X(\omega) = x_i} P(\omega).$$

Expectation

If X is a random variable on (Ω, \mathcal{F}) and P is a probability, then the expectation of X with respect to P is

$$E_P X = \sum X(\omega)P(\omega),$$

where the sum is taken over all elementary outcomes ω . It can be shown that the expectation can be calculated by using the probability distribution of X ,

$$E_P X = \sum_{i=1}^k x_i P(X = x_i).$$

Of course, if the probability P is changed to another probability Q , then the same random variable X may have a different probability distribution $q_i = Q(X = x_i)$, and a different expected value, $E_Q X = \sum_{i=1}^k x_i q_i$. When the probability P is fixed, or it is clear from the context with respect to which probability P the expectation is taken, then the reference to P is dropped from the notation, and the expectation is denoted simply by $E(X)$ or EX .

Conditional Probabilities and Expectations

Let $(\Omega, 2^\Omega, P)$ be a finite probability space, and \mathcal{G} be a field generated by a partition of Ω , $\{D_1, \dots, D_k\}$ such that $D_i \cap D_j = \emptyset$, and $\cup_i D_i = \Omega$. Recall that if D is an event of positive probability, $P(D) > 0$, then the conditional probability of an event A given the event D is defined by

$$P(A|D) = \frac{P(A \cap D)}{P(D)}.$$

Suppose that all the sets D_i in the partition have a positive probability. The conditional probability of the event A given the field \mathcal{G} is the random variable that takes values $P(A|D_i)$ on D_i , $i = 1, \dots, k$. Let I_D denote the indicator of the set D , that is, $I_D(\omega) = 1$ if $\omega \in D$, and $I_D(\omega) = 1$ if $\omega \in \bar{D}$. Using this notation, the conditional probability can be written as

$$P(A|\mathcal{G})(\omega) = \sum_{i=1}^k P(A|D_i) I_{D_i}(\omega). \quad (2.2)$$

For example, if $\mathcal{G} = \{\emptyset, \Omega\}$ is the trivial field, then

$$P(A|\mathcal{G}) = P(A|\Omega) = P(A).$$

Let Y now be a random variable that takes values y_1, \dots, y_k , then the sets $D_i = \{\omega : Y(\omega) = y_i\}$, $i = 1, \dots, k$, form a partition of Ω . If \mathcal{F}_Y denotes the field generated by Y , then the conditional probability given \mathcal{F}_Y is denoted by

$$P(A|\mathcal{F}_Y) = P(A|Y).$$

It has been assumed so far that all the sets in the partition have a positive probability. If the partition contains a set of zero probability, call it N , then the conditional probability is not defined on N by formula (2.2). It can be defined for an $\omega \in N$ arbitrarily. Consequently, any random variable which is defined by formula (2.2) and is defined arbitrarily on N is a version of the conditional probability. Any two versions only differ on N , which is a set of probability zero.

Conditional Expectation

In this section, let X take values x_1, \dots, x_p and $A_1 = \{X = x_1\}, \dots, A_p = \{X = x_p\}$. Let the field \mathcal{G} be generated by a partition $\{D_1, D_2, \dots, D_k\}$ of Ω . Then the conditional expectation of X given \mathcal{G} is defined by

$$E(X|\mathcal{G}) = \sum_{i=1}^p x_i P(A_i|\mathcal{G}).$$

Note that $E(X|\mathcal{G})$ is a linear combination of random variables so that it is a random variable. It is clear that $P(A|\mathcal{G}) = E(I_A|\mathcal{G})$, and $E(X|\mathcal{F}_0) = EX$, where $\mathcal{F}_0 = \{\emptyset, \Omega\}$ is the trivial field.

By the definition of measurability, X is \mathcal{G} -measurable if, and only if, for any i , $\{X = x_i\} = A_i$ is a member of \mathcal{G} , which means that it is either one of the D_j 's or a union of some of the D_j 's. Since $X(\omega) = \sum_{i=1}^p x_i I_{A_i}(\omega)$, a \mathcal{G} -measurable X can be written as $X(\omega) = \sum_{j=1}^k x_j I_{D_j}(\omega)$, where some x_j 's may be equal. Now it is easy to see that

$$\text{if } X \text{ is } \mathcal{G}\text{-measurable, then } E(X|\mathcal{G}) = X.$$

Note that since the conditional probabilities are defined up to a null set, so is the conditional expectation.

If X and Y are random variables both taking a finite number of values, then $E(X|Y)$ is defined as $E(X|\mathcal{G})$, where $\mathcal{G} = \mathcal{F}_Y$ is the field generated by the random variable Y . In other words, if X takes values x_1, \dots, x_p and Y takes values y_1, \dots, y_k , and $P(Y = y_i) > 0$ for all $i = 1, \dots, k$, then $E(X|Y)$ is a random variable, which takes values $\sum_{j=1}^p x_j P(X = x_j | Y = y_i)$ on the set $\{Y = y_i\}$ $i = 1, \dots, k$. These values are denoted by $E(X|Y = y_i)$. It is clear from the definition that $E(X|Y)$ is a function of Y ,

$$E(X|Y)(\omega) = E(X|\mathcal{F}_Y)(\omega) = \sum_{i=1}^k \left(\sum_{j=1}^p x_j P(X = x_j | Y = y_i) \right) I_{\{Y=y_i\}}(\omega).$$

2.2 Continuous Probability Model

In this section, we define similar probabilistic concepts for a continuous sample space. We start with general definitions.

σ -Fields

A σ -field is a field which is closed with respect to countable unions and countable intersections of its members; it is a collection of subsets of Ω that satisfies

1. $\emptyset, \Omega \in \mathcal{F}$.
2. $A \in \mathcal{F} \Rightarrow \bar{A} \in \mathcal{F}$.
3. $A_1, A_2, \dots, A_n, \dots \in \mathcal{F}$ then $\bigcup_{n=1}^{\infty} A_n \in \mathcal{F}$ (and then also $\bigcap_{n=1}^{\infty} A_n \in \mathcal{F}$).

Any subset B of Ω that belongs to \mathcal{F} is called a measurable set.

Borel σ -Field

Borel σ -field is the most important example of a σ -field that is used in the theory of functions, Lebesgue integration, and probability. Consider the σ -field \mathcal{B} on \mathbb{R} ($\Omega = \mathbb{R}$) generated by the intervals. It is obtained by taking all the intervals first and then all the sets obtained from the intervals by forming countable unions; countable intersections and their complements are included in the collection, and countable unions and intersections of these sets are included, etc. It can be shown that we end up with the smallest σ -field which contains all the intervals. A rigorous definition follows. One can show that the intersection of σ -fields is again a σ -field. Take the intersection of all σ -fields containing the collection of intervals. It is the smallest σ -field containing the intervals, the Borel σ -field on \mathbb{R} . In this model, a measurable set is a set from \mathcal{B} , a Borel set.

Probability

A probability P on (Ω, \mathcal{F}) is a non-negative set function on \mathcal{F} , such that

1. $P(\Omega) = 1$.
2. If $A \in \mathcal{F}$, then $P(\bar{A}) = 1 - P(A)$.
3. If $A_1, A_2, \dots, A_n, \dots \in \mathcal{F}$, are mutually exclusive, then $P(\bigcup_{n=1}^{\infty} A_n) = \sum_{n=1}^{\infty} P(A_n)$. This is known as countable additivity (σ -additivity).

The σ -additivity property is equivalent to finite additivity plus the continuity property of probability, which states: if $A_1 \supseteq A_2 \supseteq \dots \supseteq A_n \supseteq \dots \supseteq A = \bigcap_{n=1}^{\infty} A_n \in \mathcal{F}$, then

$$\lim_{n \rightarrow \infty} P(A_n) = P(A).$$

A similar property holds for an increasing sequence of events.

How can one define a probability on a σ -field? It is not hard to see that it is impossible to assign probabilities to all individual ω 's since there are too many of them and $P(\{\omega\}) = 0$. On the other hand, it is difficult to assign probabilities to sets in \mathcal{F} directly since in general we do not even know what a set from \mathcal{F} looks like. The standard way is to define the probability on a field which generates the σ -field, and then extend it to the σ -field.

Theorem 2.2 (Caratheodory Extension Theorem): *If a set function P is defined on a field \mathcal{F} , satisfies $P(\Omega) = 1$, $P(\bar{A}) = 1 - P(A)$, and is countably additive, then there is a unique extension of P to the σ -field generated by \mathcal{F} .*

Lebesgue Measure

As an application of Theorem 2.2 we define the Lebesgue measure on $[0, 1]$. Let $\Omega = [0, 1]$, and take for \mathcal{F} the class of all finite unions of disjoint intervals contained in $[0, 1]$. It is clearly a field. Define the probability $P(A)$ on \mathcal{F} by the length of A . It is not hard to show that P is σ -additive on \mathcal{F} . Thus there is a unique extension of P to \mathcal{B} , the Borel σ -field generated by \mathcal{F} . This extension is the Lebesgue measure on \mathcal{B} . It is also a probability on \mathcal{B} , since the length of $[0, 1]$ is one.

Any point has Lebesgue measure zero. Indeed, $\{x\} = \cap_n(x - 1/n, x + 1/n)$. Therefore $P(\{x\}) = \lim_{n \rightarrow \infty} 2/n = 0$. By countable additivity it follows that any countable set has Lebesgue measure zero. In particular the set of rationals on $[0, 1]$ is of zero Lebesgue measure. The set of irrationals on $[0, 1]$ has Lebesgue measure one.

The term “almost everywhere” (for “almost all x ”) means everywhere (for all x) except, perhaps, a set of Lebesgue measure zero.

Random Variables

A random variable X on (Ω, \mathcal{F}) is a measurable function from (Ω, \mathcal{F}) to $(\mathbb{R}, \mathcal{B})$, where \mathcal{B} is the Borel σ -field on the line. This means that for any Borel set $B \in \mathcal{B}$ the set $\{\omega : X(\omega) \in B\}$ is a member of \mathcal{F} . Instead of verifying the definition for all Borel sets, it is enough to have that for all real x the set $\{\omega : X(\omega) \leq x\} \in \mathcal{F}$. In simple words, for a random variable we can assign probabilities to sets of the form $\{X \leq x\}$, and $\{a < X \leq b\}$.

Example 2.8: Take $\Omega = \mathbb{R}$ with the Borel σ -field \mathcal{B} . A measurable function on \mathbb{R} is usually understood to be a \mathcal{B} -measurable function, that is, a random variable on $(\mathbb{R}, \mathcal{B})$. In order to define a probability on \mathcal{B} , take $f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$ and define $P(A) = \int_A f(x)dx$ for any interval A . It is easy to show that P so defined is a probability on the algebra containing the intervals, and it is continuous at the \emptyset . Thus it extends uniquely to \mathcal{B} . The function $X(x) = x$ on this probability space is called the standard normal random variable.

An important question is how to describe measurable functions, and how to decide whether a given function is measurable. It is easy to see that indicators of measurable sets are measurable. (An indicator of a set A is defined as $I_A(\omega) = 1$ if, and only if, $\omega \in A$.) Conversely, if an indicator of a set is measurable, then the set is measurable, $A \in \mathcal{F}$. A simple function (simple random variable) is a finite linear combination of indicators of measurable sets. By definition, a simple function takes finitely many values and is measurable.

Theorem 2.3: X is a random variable on (Ω, \mathcal{F}) if, and only if, it is a simple function or a limit of simple functions.

The sufficiency part follows from example 2.9. The necessity is not hard to prove by establishing that the supremum and infimum of random variables is a random variable.

Example 2.9: The following sequence of simple functions approximates a random variable X .

$$X_n(\omega) = \sum_{k=-n^{2^n}}^{n^{2^n}-1} \frac{k}{2^n} I_{[\frac{k}{2^n}, \frac{k+1}{2^n})}(X(\omega)).$$

These variables are constructed by taking the interval $[-n, n]$ and dividing it into n^{2^n+1} equal parts. X_n is zero on the set where $X \geq n$ or $X < -n$. On the set where the values of X belong to the interval $[\frac{k}{2^n}, \frac{k+1}{2^n})$, X is replaced by $\frac{k}{2^n}$, its smallest value on that set. Note that all the sets $\{\omega : \frac{k}{2^n} \leq X(\omega) < \frac{k+1}{2^n}\}$ are measurable by definition of a random variable. It is easy to see that the X_n 's are increasing and therefore converge to a limit, and for all ω , $X(\omega) = \lim_{n \rightarrow \infty} X_n(\omega)$.

This example is due to Lebesgue, who gave it for non-negative functions, demonstrating that a measurable function X is a limit of a monotone sequence of simple functions X_n , $X_{n+1} \geq X_n$.

The next result states that a limit of random variables and a composition of random variables is again a random variable.

Theorem 2.4:

1. If X_n are random variables on (Ω, \mathcal{F}) and $X(\omega) = \lim_{n \rightarrow \infty} X_n(\omega)$, then X is also a random variable.
2. If X is a random variable on (Ω, \mathcal{F}) and g is a \mathcal{B} -measurable function, then $g(X)$ is also a random variable.

Remark 2.1: In the above theorem, the requirement that the limit $X(\omega) = \lim_{n \rightarrow \infty} X_n(\omega)$ exists for all ω can be replaced by its existence for all ω outside a set of probability zero, and on a subsequence. Such a limit is in probability and it is introduced later.

σ -Field Generated by a Random Variable

The σ -field generated by a random variable X is the smallest σ -field containing sets of the form $\{\omega : a \leq X(\omega) \leq b\}$, for any $a, b \in \mathbb{R}$.

Distribution of a Random Variable

The probability distribution function of X is defined as

$$F(x) = F_X(x) = P(X \leq x).$$

It follows from the properties of probability that F is non-decreasing and right-continuous. Due to monotonicity, it is of finite variation.

Joint Distribution

If X and Y are random variables defined on the same space, then their joint distribution function is defined as

$$F(x, y) = \mathbb{P}(X \leq x, Y \leq y),$$

for any choice of real numbers x and y .

The distribution of X is recovered from the joint distribution of X and Y by $F_X(x) = F(x, \infty)$, and similarly, the distribution of Y is given by $F(\infty, y)$; they are called the marginal distributions.

The joint distribution of n random variables X_1, X_2, \dots, X_n is defined by

$$\mathbb{P}(X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n).$$

The collection of random variables $\mathbf{X} = (X_1, X_2, \dots, X_n)$ is referred to as a random vector \mathbf{X} , and the joint distribution as a multivariate probability distribution of \mathbf{X} . One can consider \mathbf{X} an \mathbb{R}^n -valued random variable, and it is possible to prove that \mathbf{X} is an \mathbb{R}^n -valued random variable if, and only if, all of its components are random variables.

Transformation of Densities

A random vector \mathbf{X} has a density $f(\mathbf{x}) = f(x_1, x_2, \dots, x_n)$ if for any set B (a Borel subset of \mathbb{R}^n),

$$\mathbb{P}(\mathbf{X} \in B) = \int_{\mathbf{x} \in B} f(\mathbf{x}) dx_1 dx_2 \dots dx_n. \quad (2.3)$$

If \mathbf{X} is transformed into \mathbf{Y} by a transformation $\mathbf{y} = \mathbf{y}(\mathbf{x})$, i.e.

$$y_1 = y_1(x_1, x_2, \dots, x_n)$$

$$y_2 = y_2(x_1, x_2, \dots, x_n)$$

...

$$y_n = y_n(x_1, x_2, \dots, x_n),$$

then, provided this transformation is one-to-one, and the inverse transformation has a non-vanishing Jacobian

$$J = \det \begin{bmatrix} \frac{\partial x_1}{\partial y_1} & \frac{\partial x_1}{\partial y_2} & \cdots & \frac{\partial x_1}{\partial y_n} \\ \frac{\partial x_2}{\partial y_1} & \frac{\partial x_2}{\partial y_2} & \cdots & \frac{\partial x_2}{\partial y_n} \\ \cdots & \cdots & \cdots & \cdots \\ \frac{\partial x_n}{\partial y_1} & \frac{\partial x_n}{\partial y_2} & \cdots & \frac{\partial x_n}{\partial y_n} \end{bmatrix},$$

\mathbf{Y} has a density given by

$$f_{\mathbf{Y}}(\mathbf{y}) = f(x_1(\mathbf{y}), x_2(\mathbf{y}), \dots, x_n(\mathbf{y}))|J(\mathbf{y})|. \quad (2.4)$$

This is easily established by using the change of variables in multiple integrals (see Example 2.15 for calculation of the bivariate density).

2.3 Expectation and Lebesgue Integral

Let X be a random variable on (Ω, \mathcal{F}) , and P be a probability on \mathcal{F} . Recall that in the discrete case the expectation of X is defined as $\sum_{\omega} X(\omega)P(\omega)$. The expectation in the continuous model is defined by the means of an integral

$$EX = \int_{\Omega} X(\omega)dP(\omega).$$

The expectation is defined for positive random variables first. The general case is obtained by using the decomposition $X = X^+ - X^-$, where $X^+ = \max(X, 0)$ and $X^- = \max(-X, 0)$, and letting $EX = EX^+ - EX^-$, provided both EX^+ and EX^- are finite.

If $X \geq 0$ is a random variable, then it can be approximated by simple random variables (see Theorem 2.3 and Example 2.9). The expectation of a simple random variable is defined as a sum, that is, if

$$X = \sum_{k=1}^n c_k I_{A_k}, \quad \text{then} \quad EX = \sum_{k=1}^n c_k P(A_k).$$

Note that for a simple random variable $X \geq 0$ implies $EX \geq 0$. This in turn implies that if $X \geq Y$, where X and Y are simple random variables, then $EX \geq EY$.

Any positive random variable X can be approximated by an increasing sequence X_n of simple random variables; such approximation is given in Example 2.9. It now follows that since X_n is an increasing sequence, EX_n is also an increasing sequence, hence it has a limit. The limit of EX_n is taken to be EX . It can be shown that this limit does not depend on the choice of the approximating sequence of simple random variables, so that the expectation is defined unambiguously.

Definition 2.5: A random variable X is called integrable if both EX^+ and EX^- are finite. In this case, $EX = EX^+ - EX^-$.

Note that for X to be integrable both EX^+ and EX^- must be finite, which is the same as $E|X| = EX^+ + EX^- < \infty$.

Lebesgue–Stieltjes Integral

A distribution function on \mathbb{R} is a non-decreasing right-continuous function which approaches 0 at $-\infty$ and 1 at $+\infty$. Such a distribution function, say F , defines uniquely a probability on the Borel σ -field \mathcal{B} by setting $P((a, b]) = F(b) - F(a)$.

Now take $(\Omega, \mathcal{F}) = (\mathbb{R}, \mathcal{B})$ and a probability on \mathcal{B} given by a distribution function $F(x)$. A random variable on this space is a measurable function $f(x)$. Its expected value is written as $\int_{\mathbb{R}} f(x)F(dx)$ and is called the Lebesgue–Stieltjes integral of f with respect to F .

The distribution function F can be replaced by any function of finite variation, giving rise to the general Lebesgue–Stieltjes integral.

The probability distribution of a random variable X on (Ω, \mathcal{F}) is the probability on \mathcal{B} carried from \mathcal{F} by X : for any $B \in \mathcal{B}$,

$$P_X(B) = P(X \in B). \quad (2.5)$$

The distribution function is related to this probability by $F(x) = P_X((-\infty, x])$. Equation (2.5) gives the relation between the expectations of indicator functions,

$$\int_{\Omega} I(X(\omega) \in B)dP(\omega) = \int_{-\infty}^{\infty} I(x \in B)P_X(dx).$$

This can be extended from indicators to measurable functions using an approximation by simple functions, and we have the following result.

Theorem 2.6: *If X is a random variable with distribution function $F(x)$, and h is a measurable function on \mathbb{R} , such that $h(X)$ is integrable, then*

$$Eh(X) := \int_{\Omega} h(X(\omega))dP(\omega) = \int_{-\infty}^{\infty} h(x)P_X(dx) := \int_{-\infty}^{\infty} h(x)F(dx). \quad (2.6)$$

Lebesgue Integral on the Line

The Lebesgue–Stieltjes integral with respect to $F(x) = x$ is known as the Lebesgue integral.

Example 2.10: Let $\Omega = [0, 1]$, its elements ω are real numbers x , and take for probability the Lebesgue measure. Take $X(\omega) = X(x) = x^2$. Then $EX = \int_0^1 x^2 dx = 1/3$. Construct an approximating sequence of simple functions and verify the value of the above integral.

Similarly, for any continuous function $f(x)$ on $[0, 1]$, $X(\omega) = X(x) = f(x)$ is a random variable (using that a continuous function is measurable) with expectation $EX = Ef = \int_0^1 f(x)dx$.

Theorem 2.7: If f is Riemann integrable on $[a, b]$, then it is Lebesgue integrable on $[a, b]$ and the integrals coincide.

On the other hand, there are functions which are Lebesgue integrable but not Riemann integrable. Recall that for a function to be Riemann integrable, it must be continuous at all points except for a set of Lebesgue measure zero. Some everywhere discontinuous functions are Lebesgue integrable.

Example 2.11: $\Omega = [0, 1]$, and probability is given by the Lebesgue measure. Take $X(x) = I_Q(x)$ to be the indicator function of the set Q of all rationals. Q has Lebesgue measure zero. As the expectation of an indicator is the probability of its set, $EX = \int_0^1 I_Q(x)dx = 0$. However, $I_Q(x)$ is discontinuous at every point, so that the set of discontinuities of $I_Q(x)$ is $[0, 1]$ which has Lebesgue measure one, therefore $I_Q(x)$ is not Riemann integrable.

The next result is the fundamental theorem for the Lebesgue integral on the line.

Theorem 2.8: If f is Lebesgue integrable on $[a, b]$ then the derivative of the integral exists for almost all $x \in (a, b)$, and

$$\frac{d}{dx} \int_a^x f(t)dt = f(x). \quad (2.7)$$

Properties of Expectation (Lebesgue Integral)

It is not hard to show that the expectation (Lebesgue integral) satisfies the following properties:

1. Linearity. If X and Y are integrable and α and β are constants, then $E(\alpha X + \beta Y) = \alpha EX + \beta EY$.
2. If random variables X and Y satisfy $|X| \leq Y$ and Y is integrable, then X is also integrable and $E|X| \leq EY$.
3. If a random variable $X \geq 0$, then $EX = 0$ if, and only if, $P(X = 0) = 1$.

Jumps and Probability Densities

The jump of F at x gives the probability $P(X = x)$, $F(x) - F(x-) = P(X = x)$. Since F is right-continuous it has at most countably many jumps.

Definition 2.9: F is called discrete if it changes only by jumps.

If $F(x)$ is continuous at x then $P(X = x) = 0$.

Definition 2.10: $F(x)$ is called absolutely continuous if there is a function $f(x) \geq 0$, such that for all x , $F(x)$ is given by the Lebesgue integral $F(x) = \int_{-\infty}^x f(t)dt$. In this case, $F'(x) = f(x)$ for almost all x (Theorem 2.8).

f is called the probability density function of X . It follows from the definition that for any $a < b$

$$P(a \leq X \leq b) = \int_a^b f(x)dx.$$

There are plenty of examples in any introductory book on probability or statistics of continuous random variables with densities that are normal, exponential, uniform, gamma, Cauchy, etc.

The random variables X and Y with the joint distribution $F(x, y)$ possess a density $f(x, y)$ (with respect to the Lebesgue measure) if for any x, y

$$F(x, y) = \int_{-\infty}^x \int_{-\infty}^y f(u, v)dudv,$$

and then for almost all (with respect to the Lebesgue measure on the plane) x, y ,

$$f(x, y) = \frac{\partial^2 F}{\partial x \partial y}(x, y).$$

A density for an n -dimensional random vector is defined similarly.

Decomposition of Distributions and FV Functions

Any distribution function can be decomposed into a continuous part and a jump part. Continuous distribution functions can be decomposed further (Lebesgue decomposition).

If F is a continuous distribution function, then it can be decomposed into the sum of two continuous distribution functions, the absolutely continuous part and the singular part, i.e. for some $0 \leq a \leq 1$

$$F = aF_{ac} + (1 - a)F_{sing}. \quad (2.8)$$

F_{ac} is characterized by its density that exists at almost all points. For the singular part, $F'_{sing}(x)$ exists for almost all x and is zero. An example of such a function is the Cantor function (see Example 2.13), where the distribution function is a constant between the points of the Cantor set. In most applications in statistics continuous distributions are absolutely continuous with zero singular part.

Example 2.12: (Cantor set)

Consider the set $\{x : x = \sum_{n=1}^{\infty} \alpha_n/3^n, \alpha_n \in \{0, 2\}\}$. It is possible to show that this set does not have isolated points (a perfect set), that is, any point of the set is a limit of other points. Indeed, for a given sequence of α_n 's that contains infinitely many 2s, consider a new sequence which is the same up to the m -th term, with all the rest being zeros. The distance between these two numbers is given by $\sum_{n=m+1}^{\infty} \alpha_n/3^n < 3^{-m}$,

which can be made arbitrarily small as m increases. For a number with finitely many 2s, say k , the numbers with the first k same places, and the rest zeros except the m -th place which is 2, approximate it. Indeed, the distance between these two numbers is $2/3^m$. It is also not hard to see that this set is uncountable (by the diagonal argument) and that it has Lebesgue measure zero. Although the Cantor set seems to be artificially constructed, Cantor type sets arise naturally in the study of Brownian motion; for example, the set of zeros of Brownian motion is a Cantor type set.

Example 2.13: (Cantor distribution)

The distribution function F of the random variable $X = \sum_{n=1}^{\infty} \alpha_n / 3^n$, where α_n are independent identically distributed random variables taking values 0 and 2 with probability 1/2, is continuous and its derivative is zero almost everywhere.

It takes a rather pathological example to construct a continuous singular distribution in one dimension. In dimension two, such examples can be simple.

Example 2.14: (Continuous singular distributions on the plane)

Take F such that $\frac{\partial^2 F}{\partial x \partial y} = 0$ almost everywhere on the plane. If F is a linear function in x and y , or a distribution function that does not depend on one of the variables x or y , then it is singular. For example, $0 \leq X, Y \leq 1$ such that their joint distribution is determined by $F(x, y) = \frac{1}{2}(x + y)$, for x, y satisfying $0 \leq x, y \leq 1$. In this case, only sets that have non-empty intersection with the axis have positive probability.

Functions of finite variations have a similar structure to distribution functions. They can be decomposed into a continuous part and a jump part, and the continuous part can be decomposed further into an absolutely continuous part and a singular part.

2.4 Transforms and Convergence

If X^k is integrable, $E|X|^k < \infty$, then the k -th moment of X is defined as $E(X^k)$. The moment generating function of X is defined as

$$m(t) = E(e^{tX}),$$

provided e^{tX} is integrable, for t in a neighbourhood of 0.

Using the series expansion for the exponential function

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!},$$

we can formally write, by interchanging summation and the expectation,

$$m(t) = Ee^{tX} = E \sum_{n=0}^{\infty} \frac{t^n X^n}{n!} = \sum_{n=0}^{\infty} \frac{t^n}{n!} E(X^n). \quad (2.9)$$

Thus $E(X^n)$ can be obtained from the power series expansion of the moment generating function.

The characteristic function of X is defined as

$$\phi(t) = E(e^{itX}) = E(\cos(tX)) + iE(\sin(tX)),$$

where $i = \sqrt{-1}$. The characteristic function determines the distribution uniquely; so does the moment generating function when it exists on an interval containing zero. The advantage of the characteristic function over the moment generating function is that it exists for any random variable X , since the functions $\cos(tx)$ and $\sin(tx)$ are bounded for any t on the whole line, whereas e^{tx} is unbounded and the moment generating function need not exist. Existence of the moment generating function around zero implies existence of all the moments. If X does not have all the moments, then its moment generating function does not exist.

Convergence of Random Variables

There are four main concepts of convergence of a sequence of random variables. We give the definitions of progressively stronger concepts and some results on their relations.

Definition 2.11: $\{X_n\}$ converge in distribution to X , if their distribution functions $F_n(x)$ converge to the distribution function $F(x)$ at any point of continuity of F .

It can be shown that $\{X_n\}$ converge in distribution to X if, and only if, their characteristic functions (or moment generating functions) converge to that of X . Convergence in distribution is also equivalent to the requirement that $Eg(X_n) \rightarrow Eg(X)$ as $n \rightarrow \infty$ for all bounded continuous functions g on \mathbb{R} .

Definition 2.12: $\{X_n\}$ converge in probability to X if for any $\epsilon > 0$ $P(|X_n - X| > \epsilon) \rightarrow 0$ as $n \rightarrow \infty$.

Definition 2.13: $\{X_n\}$ converge almost surely (a.s.) to X if for any ω outside a set of zero probability $X_n(\omega) \rightarrow X(\omega)$ as $n \rightarrow \infty$.

Almost sure convergence implies convergence in probability, which in turn implies convergence in distribution. It is also not hard to see that convergence in distribution to a constant is the same as the convergence in probability to the same constant. Convergence in probability implies the almost sure convergence on a subsequence, namely, if $\{X_n\}$ converge in probability to X , then there is a subsequence n_k that converges almost surely to the same limit.

L^r -convergence (convergence in the r -th mean), $r \geq 1$, is defined as follows.

Definition 2.14: $\{X_n\}$ converge to X in L^r if for any n $E(|X_n|^r) < \infty$, and $E(|X_n - X|^r) \rightarrow 0$ as $n \rightarrow \infty$.

Using the concept of uniform integrability (Chapter 7), convergence in L^r is equivalent to convergence in probability and uniform integrability of $|X_n|^r$ (see for example, Loeve (1978), p. 164).

The following result, which is known as the Slutskii theorem, is frequently used in applications.

Theorem 2.15: *If X_n converges to X and Y_n converges to Y , then $X_n + Y_n$ converges to $X + Y$, for any type of stochastic convergence, except for convergence in distribution. However, if $Y = 0$ or X_n and Y_n are independent, then the result is also true for convergence in distribution.*

Convergence of Expectations

Theorem 2.16 (Monotone Convergence): *If $X_n \geq 0$ and X_n are increasing to a limit X , which may be infinite, then $\lim_{n \rightarrow \infty} EX_n = EX$.*

Theorem 2.17 (Fatou's Lemma): *If $X_n \geq 0$ (or $X_n \geq c > -\infty$), then $E(\liminf_n X_n) \leq \liminf_n EX_n$.*

Theorem 2.18 (Dominated Convergence): *If $\lim_{n \rightarrow \infty} X_n = X$ in probability and for all n $|X_n| \leq Y$ with $EY < \infty$, then $\lim_{n \rightarrow \infty} EX_n = EX$.*

2.5 Independence and Covariance

Independence

Two events A and B are called independent if $P(A \cap B) = P(A)P(B)$.

A collection of events A_i , $i = 1, 2, \dots$ is called independent if for any finite n and any choice of indices i_k , $k = 1, 2, \dots, n$

$$P\left(\bigcap_{k=1}^n A_{i_k}\right) = \prod_{k=1}^n P(A_{i_k}).$$

Two σ -fields are called independent if for any choice of sets from each of them these sets are independent.

Two random variables X and Y are independent if the σ -fields they generate are independent. It follows that their joint distribution is given

by the product of their marginal distributions (since the sets $\{X \leq x\}$ and $\{Y \leq y\}$ are in the respective σ -fields)

$$\mathrm{P}(X \leq x, Y \leq y) = \mathrm{P}(X \leq x)\mathrm{P}(Y \leq y),$$

and it can be seen that it is an equivalent property.

One can formulate the independence property in terms of expectations by writing the above in terms of indicators

$$\mathrm{E}(I(X \leq x)I(Y \leq y)) = \mathrm{E}(I(X \leq x))\mathrm{E}(I(Y \leq y)).$$

Since it is possible to approximate indicators by continuous bounded functions, X and Y are independent if, and only if, for any bounded continuous functions f and g ,

$$\mathrm{E}(f(X)g(Y)) = \mathrm{E}(f(X))\mathrm{E}(g(Y)).$$

X_1, X_2, \dots, X_n are called independent if for any choice of random variables $X_{i_1}, X_{i_2}, \dots, X_{i_k}$ their joint distribution is given by the product of their marginal distributions (alternatively, if the σ -fields they generate are independent).

Covariance

The covariance of two integrable random variables X and Y is defined, provided XY is integrable, by

$$\mathrm{Cov}(X, Y) = \mathrm{E}(X - \mathrm{E}X)(Y - \mathrm{E}Y) = \mathrm{E}(XY) - \mathrm{E}X\mathrm{E}Y. \quad (2.10)$$

The variance of X is the covariance of X with itself, $\mathrm{Var}(X) = \mathrm{Cov}(X, X)$.

The Cauchy–Schwarz inequality

$$(\mathrm{E}|XY|)^2 \leq \mathrm{E}(X^2)\mathrm{E}(Y^2), \quad (2.11)$$

assures that covariance exists for square integrable random variables. Covariance is symmetric,

$$\mathrm{Cov}(X, Y) = \mathrm{Cov}(Y, X),$$

and is linear in both variables (bilinear)

$$\mathrm{Cov}(aX + bY, Z) = a\mathrm{Cov}(X, Z) + b\mathrm{Cov}(Y, Z).$$

Using this property with $X + Y$ we obtain the formula for the variance of the sum. The following property of the covariance holds

$$\mathrm{Var}(X + Y) = \mathrm{Var}(X) + \mathrm{Var}(Y) + 2\mathrm{Cov}(X, Y). \quad (2.12)$$

Random variables X and Y are called *uncorrelated* if $\mathrm{Cov}(X, Y) = 0$. It is easy to see from the definitions that for independent random variables

$$\mathrm{E}(XY) = \mathrm{E}X\mathrm{E}Y,$$

which implies that they are uncorrelated. The opposite implication is not true in general. The important exception is the Gaussian case.

Theorem 2.19: If the random variables have a joint Gaussian distribution, then they are independent if, and only if, they are uncorrelated.

Definition 2.20: The covariance matrix of a random vector $\mathbf{X} = (X_1, X_2, \dots, X_n)$ is the $n \times n$ matrix with the elements $Cov(X_i, X_j)$.

2.6 Normal (Gaussian) Distributions

The normal (Gaussian) probability density is given by

$$f(x; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}.$$

It is completely specified by its mean μ and its standard deviation σ . The normal family $N(\mu, \sigma^2)$ is obtained from the standard normal distribution, $N(0, 1)$ by a linear transformation.

If X is $N(\mu, \sigma^2)$, then $Z = \frac{X - \mu}{\sigma}$ is $N(0, 1)$ and $X = \mu + \sigma Z$.

An important property of the normal family is that a linear combination of independent normal variables results in a normal variable, that is, if $X_1 \sim N(\mu_1, \sigma_1^2)$ and $X_2 \sim N(\mu_2, \sigma_2^2)$ are independent then $\alpha X_1 + \beta X_2 \sim N(\alpha\mu_1 + \beta\mu_2, \alpha^2\sigma_1^2 + \beta^2\sigma_2^2)$. The moment generating function of X with $N(\mu, \sigma^2)$ distribution is given by

$$m(t) = Ee^{tX} = \int_{-\infty}^{\infty} e^{tx} f(x; \mu, \sigma^2) dx = e^{\mu t} e^{(\sigma t)^2/2} = e^{\mu t + (\sigma t)^2/2}.$$

A random vector $\mathbf{X} = (X_1, X_2, \dots, X_n)$ has an n -variate normal (Gaussian) distribution with mean vector μ and covariance matrix Σ if there exists an $n \times n$ matrix A such that its determinant $|A| \neq 0$, and $\mathbf{X} = \boldsymbol{\mu} + A\mathbf{Z}$, where $\mathbf{Z} = (Z_1, Z_2, \dots, Z_n)$ is the vector with independent standard normal components, and $\Sigma = AA^T$. Vectors are taken as column vectors here and in the sequel.

The probability density of \mathbf{Z} is obtained by using the independence of its components, for independent random variables the densities multiply. Then performing a change of variables in the multiple integral we find the probability density of \mathbf{X}

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{(2\pi)^{n/2} |\Sigma|^{1/2}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})\Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu})^T}.$$

Example 2.15: Let a bivariate normal have $\boldsymbol{\mu} = \mathbf{0}$ and $\Sigma = \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}$. Let $\mathbf{X} = (X, Y)$ and $\mathbf{x} = (x, y)$. Then \mathbf{X} can be obtained from \mathbf{Z} by the transformation $\mathbf{X} = A\mathbf{Z}$ with $A = \begin{bmatrix} 1 & 0 \\ \rho & \sqrt{1-\rho^2} \end{bmatrix}$.

Since $\begin{cases} x = z_1 \\ y = \rho z_1 + \sqrt{1 - \rho^2} z_2 \end{cases}$, the inverse transformation

$\begin{cases} z_1 = x \\ z_2 = (y - \rho x) / \sqrt{1 - \rho^2} \end{cases}$ has the Jacobian

$$J = \det \begin{bmatrix} \frac{\partial z_1}{\partial x} & \frac{\partial z_1}{\partial y} \\ \frac{\partial z_2}{\partial x} & \frac{\partial z_2}{\partial y} \end{bmatrix} = \det \begin{bmatrix} 1 & 0 \\ \frac{1}{\sqrt{1-\rho^2}} & \end{bmatrix} = \frac{1}{\sqrt{1-\rho^2}}.$$

The density of \mathbf{Z} is given by the product of standard normal densities, by independence, $f_{\mathbf{Z}}(z_1, z_2) = \frac{1}{2\pi} e^{-\frac{1}{2}(z_1^2 + z_2^2)}$. Using formula (2.4) we obtain the joint density of the bivariate normal

$$f_{\mathbf{X}}(x, y) = \frac{1}{2\pi\sqrt{1-\rho^2}} e^{-\frac{1}{2(1-\rho^2)}[x^2 - 2\rho xy + y^2]}.$$

It follows from the definition that if \mathbf{X} has a multivariate normal distribution and \mathbf{a} is a non-random vector, then $\mathbf{a}\mathbf{X} = \mathbf{a}(\boldsymbol{\mu} + A\mathbf{Z}) = \mathbf{a}\boldsymbol{\mu} + \mathbf{a}A\mathbf{Z}$. Since a linear combination of independent normal random variables is a normal random variable, $\mathbf{a}A\mathbf{Z}$ is a normal random variable. Hence $\mathbf{a}\mathbf{X}$ has normal distribution with mean $\mathbf{a}\boldsymbol{\mu}$ and variance $(\mathbf{a}A)(\mathbf{a}A)^T = \mathbf{a}\boldsymbol{\Sigma}\mathbf{a}^T$. Thus we have

Theorem 2.21: *A linear combination of jointly Gaussian random variables is a Gaussian random variable.*

Similarly it can be shown that if $\mathbf{X} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ and B is a non-random matrix, then $B\mathbf{X} \sim N(B\boldsymbol{\mu}, B\boldsymbol{\Sigma}B^T)$.

The moment generating function of a vector \mathbf{X} is defined as

$$E(e^{\mathbf{t}\mathbf{X}}) = E(e^{\sum_{i=1}^n t_i X_i}),$$

where $\mathbf{t} = (t_1, t_2, \dots, t_n)$, and $\mathbf{t}\mathbf{X}$ is the scalar product of vectors \mathbf{t} and \mathbf{X} .

It is not hard to show that the moment generating function of a Gaussian vector $\mathbf{X} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ is given by

$$M_{\mathbf{X}}(\mathbf{t}) = e^{\boldsymbol{\mu}\mathbf{t}^T - \frac{1}{2}\mathbf{t}\boldsymbol{\Sigma}\mathbf{t}^T}.$$

Definition 2.22: A collection of random variables is called a Gaussian process if the joint distribution of any finite number of its members is Gaussian.

Theorem 2.23: *Let $X(t)$ be a process with independent Gaussian increments, that is, for any $s < t$, $X(t) - X(s)$ has a normal distribution, and is independent of the values $X(u)$, $u \leq s$ (the σ -field \mathcal{F}_s generated by the process up to time s). Then $X(t)$ is a Gaussian process.*

See Chapter 3.1, Example 3.3, for the proof.

2.7 Conditional Expectation

Conditional Expectation and Conditional Distribution

The conditional distribution function of X given $Y = y$ is defined by

$$P(X \leq x | Y = y) = \frac{P(X \leq x, Y = y)}{P(Y = y)},$$

provided $P(Y = y) > 0$. However, such an approach fails if the event we condition on has zero probability, $P(Y = y) = 0$. This difficulty can be overcome if X, Y have a joint density $f(x, y)$. In this case it follows that both X and Y possess densities $f_X(x)$ and $f_Y(y)$; $f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy$ and $f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx$. The conditional distribution of X given $Y = y$ is defined by the conditional density

$$f(x|y) = \frac{f(x, y)}{f_Y(y)},$$

at any point where $f_Y(y) > 0$. It is easy to see that so defined $f(x|y)$ is indeed a probability density for any y , as it is non-negative and integrates to unity.

The expectation of this distribution, when it exists, is called the conditional expectation of X given $Y = y$,

$$E(X|Y = y) = \int_{-\infty}^{\infty} xf(x|y) dx. \quad (2.13)$$

The conditional expectation $E(X|Y = y)$ is a function of y . Let g denote this function, $g(y) = E(X|Y = y)$, then by replacing y by Y we obtain a random variable $g(Y)$, which is the conditional expectation of X given Y , $E(X|Y) = g(Y)$.

Example 2.16: Let X and Y have a standard bivariate normal distribution with parameter ρ . Then $f(x, y) = \frac{1}{2\pi\sqrt{1-\rho^2}} \exp\left\{-\frac{1}{2(1-\rho^2)}[x^2 - 2\rho xy + y^2]\right\}$ and $f_Y(y) = \frac{1}{\sqrt{2\pi}} e^{-y^2/2}$, so that $f(x|y) = \frac{f(x,y)}{f_Y(y)} = \frac{1}{\sqrt{2\pi(1-\rho^2)}} \exp\left\{-\frac{(x-\rho y)^2}{2(1-\rho^2)}\right\}$, which is the $N(\rho y, 1 - \rho^2)$ distribution. Its mean is ρy , therefore $E(X|Y = y) = \rho y$ and $E(X|Y) = \rho Y$.

Similarly, it can be seen that in the multivariate normal case the conditional expectation is also a linear function of Y .

The conditional distribution and the conditional expectation are defined only at the points where $f_Y(y) > 0$. Both can be defined arbitrarily on the set $\{y : f_Y(y) = 0\}$. Since there are many functions which agree on the set $\{y : f_Y(y) > 0\}$, any one of them is called a version of the conditional

distribution (the conditional expectation) of X given $Y = y$. The different versions of $f(x|y)$ and $E(X|Y = y)$ differ only on the set $\{y : f_Y(y) = 0\}$, which has zero probability under the distribution of Y ; $f(x|y)$ and $E(X|Y = y)$ are defined as uniquely Y -almost surely.

General Conditional Expectation

The conditional expectation in a more general form is defined as follows. Let X be an integrable random variable. $E(X|Y) = G(Y)$, a function of Y such that for any bounded function h ,

$$E(Xh(Y)) = E(G(Y)h(Y)), \quad (2.14)$$

or $E((X - G(Y))h(Y)) = 0$. Existence of such a function is assured by the Radon–Nikodym theorem from functional analysis. However, uniqueness is easy to prove. If there are two such functions, G_1, G_2 , then $E((G_1(Y) - G_2(Y))h(Y)) = 0$. Take $h(y) = \text{sign}(G_1(y) - G_2(y))$. Then we have $E|G_1(Y) - G_2(Y)| = 0$. Thus $P(G_1(Y) = G_2(Y)) = 1$, and they coincide with (Y) probability one.

A more general conditional expectation of X given a σ -field \mathcal{G} , $E(X|\mathcal{G})$ is a \mathcal{G} -measurable random variable such that for any bounded \mathcal{G} -measurable ξ

$$E(\xi E(X|\mathcal{G})) = E(\xi X). \quad (2.15)$$

In the literature, $\xi = I_B$ is taken as an indicator function of a set $B \in \mathcal{G}$, which is an equivalent condition: for any set $B \in \mathcal{G}$

$$\int_B X dP = \int_B E(X|\mathcal{G}) dP, \quad \text{or} \quad E(XI(B)) = E(E(X|\mathcal{G})I(B)). \quad (2.16)$$

The Radon–Nikodym theorem (see Theorem 10.6) implies that such a random variable exists and is almost surely unique, in the sense that any two versions differ only on a set of probability zero.

The conditional expectation $E(X|Y)$ is given by $E(X|\mathcal{G})$ with $\mathcal{G} = \sigma(Y)$, the σ -field generated by Y . Often the Equations (2.15) or (2.16) are not used because easier calculations are possible for various specific properties, but they are used to establish the fundamental properties given below. In particular, the conditional expectation defined in (2.13) by using densities satisfies (2.15) or (2.16).

Properties of Conditional Expectation

Conditional expectations are random variables. Their properties are stated as equalities of two random variables. Random variables X and Y , defined on the same space, are equal if $P(X = Y) = 1$. This is also written $X = Y$

a.s. If not stated otherwise, whenever the equality of random variables is used it is intended in the “almost sure” sense, and often writing “almost surely” is omitted.

1. If \mathcal{G} is the trivial field $\{\emptyset, \Omega\}$, then

$$\mathbb{E}(X|\mathcal{G}) = EX. \quad (2.17)$$

2. If X is \mathcal{G} -measurable, then

$$\mathbb{E}(XY|\mathcal{G}) = X\mathbb{E}(Y|\mathcal{G}). \quad (2.18)$$

This means that if \mathcal{G} contains all the information about X , then given \mathcal{G} , X is known, and therefore it is treated as a constant.

3. If $\mathcal{G}_1 \subset \mathcal{G}_2$ then

$$\mathbb{E}(\mathbb{E}(X|\mathcal{G}_2)|\mathcal{G}_1) = \mathbb{E}(X|\mathcal{G}_1). \quad (2.19)$$

This is known as the smoothing property of conditional expectation. In particular, by taking \mathcal{G}_1 to be the trivial field we obtain the law of double expectation

$$\mathbb{E}(\mathbb{E}(X|\mathcal{G})) = \mathbb{E}(X). \quad (2.20)$$

4. If $\sigma(X)$ and \mathcal{G} are independent, then

$$\mathbb{E}(X|\mathcal{G}) = EX, \quad (2.21)$$

that is, if the information we know provides no clues about X , then the conditional expectation is the same as the expectation. The next result is an important generalization.

5. If $\sigma(X)$ and \mathcal{G} are independent, \mathcal{F} and \mathcal{G} are independent, and $\sigma(\mathcal{F}, \mathcal{G})$ denotes the smallest σ -field containing both of them, then

$$\mathbb{E}(X|\sigma(\mathcal{F}, \mathcal{G})) = \mathbb{E}(X|\mathcal{F}). \quad (2.22)$$

6. Jensen's inequality. If $g(x)$ is a convex function on I , that is, for all $x, y \in I$ and $\lambda \in (0, 1)$

$$g(\lambda x + (1 - \lambda)y) \leq \lambda g(x) + (1 - \lambda)g(y),$$

and X is a random variable with range I , then

$$g(\mathbb{E}(X|\mathcal{G})) \leq \mathbb{E}(g(X)|\mathcal{G}). \quad (2.23)$$

In particular, with $g(x) = |x|$

$$|\mathbb{E}(X|\mathcal{G})| \leq \mathbb{E}(|X||\mathcal{G}|). \quad (2.24)$$

7. Monotone convergence. If $0 \leq X_n$ and $X_n \uparrow X$ with $E|X| < \infty$, then

$$E(X_n|\mathcal{G}) \uparrow E(X|\mathcal{G}). \quad (2.25)$$

8. Fatou's lemma. If $0 \leq X_n$, then

$$E\left(\liminf_n X_n|\mathcal{G}\right) \leq \liminf_n E(X_n|\mathcal{G}). \quad (2.26)$$

9. Dominated convergence. If $\lim_{n \rightarrow \infty} X_n = X$ almost surely and $|X_n| \leq Y$ with $EY < \infty$, then

$$\lim_{n \rightarrow \infty} E(X_n|\mathcal{G}) = E(X|\mathcal{G}). \quad (2.27)$$

Most of the properties of conditional expectation are easily verified using the definition (2.15). For example, we show the law of double expectation, which is very frequently used, $E(E(X|\mathcal{G})) = E(X)$. Let $\hat{X} = (E(X|\mathcal{G}))$. Then for all bounded \mathcal{G} -measurable ξ we have $E(\hat{X}\xi) = E(X\xi)$. Since Ω belongs to any σ -field (by definition of a σ -field), I_Ω , the indicator of Ω , is \mathcal{G} -measurable, and we have $E(\hat{X}) = E(\hat{X}I_\Omega) = E(XI_\Omega) = E(X)$.

For results on conditional expectations see, for example, Breiman (1968). The conditional probability $P(A|\mathcal{G})$ is defined as the conditional expectation of the indicator function,

$$P(A|\mathcal{G}) = E(I_A|\mathcal{G}),$$

and it is a \mathcal{G} -measurable random variable defined P -almost surely.

The following results are often used.

Theorem 2.24: *Let X and Y be two independent random variables and $\phi(x, y)$ be such that $E|\phi(X, Y)| < +\infty$. Then*

$$E(\phi(X, Y)|Y) = G(Y),$$

where $G(y) = E(\phi(X, y))$.

Theorem 2.25: *Let (\mathbf{X}, \mathbf{Y}) be a Gaussian vector. Then the conditional distribution of \mathbf{X} given \mathbf{Y} is also Gaussian. Moreover, provided the matrix $Cov(\mathbf{Y}, \mathbf{Y})$ is non-singular (has the inverse),*

$$E(\mathbf{X}|\mathbf{Y}) = E(\mathbf{X}) + Cov(\mathbf{X}, \mathbf{Y})Cov^{-1}(\mathbf{Y}, \mathbf{Y})(\mathbf{Y} - E(\mathbf{Y})).$$

In the case when $Cov(\mathbf{Y}, \mathbf{Y})$ is singular, the same formula holds with the inverse replaced by the generalized inverse, the Moore–Penrose pseudoinverse matrix.

If we want to predict/estimate X by using observations on Y , then a predictor is some function of Y . For a square integrable X , $E(X^2) < \infty$,

the best predictor \hat{X} , by definition, minimizes the mean square error. It is easy to show the following theorem:

Theorem 2.26 (Best Estimator/Predictor): *Let \hat{X} be such that for any Y -measurable random variable Z ,*

$$\mathbb{E}(X - \hat{X})^2 \leq \mathbb{E}(X - Z)^2.$$

Then $\hat{X} = \mathbb{E}(X|Y)$.

2.8 Stochastic Processes in Continuous Time

The construction of a mathematical model of uncertainty and of flow of information in continuous time follows the same ideas as in discrete time, but it is much more complicated. Consider constructing a probability model for a process $S(t)$ when time changes continuously between 0 and T . Take for the sample space the set of all possibilities of movement of the process. If we make a simplifying assumption that the process changes continuously, we obtain the set of all continuous functions on $[0, T]$ denoted by $C[0, T]$. This is a very rich space. In a more general model it is assumed that the observed process is a right-continuous function with left limits (R.R.C., *càdlàg*) function.

Let the sample space $\Omega = D[0, T]$ be the set of all R.R.C. functions on $[0, T]$. An element of this set, ω is a R.R.C. function from $[0, T]$ into \mathbb{R} . First we must decide what kind of sets of these functions are measurable. The simplest sets for which we would like to calculate the probabilities are sets of the form $\{a \leq S(t_1) \leq b\}$ for some t_1 . If $S(t)$ represents the price of a stock at time t , then the probability of such a set gives the probability that the stock price at time t_1 is between a and b . We are also interested in how the price of stock at time t_1 affects the price at another time t_2 . Thus we need to talk about the joint distribution of stock prices $S(t_1)$ and $S(t_2)$. This means that we need to define probability on the sets of the form $\{S(t_1) \in B_1, S(t_2) \in B_2\}$ where B_1 and B_2 are intervals on the line. More generally we would like to have all *finite-dimensional distributions* of the process $S(t)$, that is, probabilities of the sets: $\{S(t_1) \in B_1, \dots, S(t_n) \in B_n\}$, for any choice of $0 \leq t_1 \leq t_2, \dots \leq t_n \leq T$. The sets of the form $\{\omega(\cdot) \in D[0, T] : \omega(t_1) \in B_1, \dots, \omega(t_n) \in B_n\}$, where B_i 's are intervals on the line, are called *cylinder sets* or *finite-dimensional rectangles*. The stochastic process $S(t)$ on this sample space is just $s(t)$, the value of the function s at t . Probability is defined first on the cylinder sets, and then extended to the σ -field \mathcal{F} generated by the cylinders, that is, the smallest σ -field containing all cylinder sets. One needs to be careful with consistency

of probability defined on cylinder sets, so that when one cylinder contains another no contradiction of probability assignment is obtained. The result that shows that a consistent family of distributions defines a probability function continuous at \emptyset on the field of cylinder sets is known as Kolmogorov's extension theorem. A probability defined on a field of cylinder sets can be extended in a unique way (by Caratheodory's theorem) to \mathcal{F} (see for example, Breiman (1968), Durrett (1991), or Dudley (1989) for details).

It follows immediately from this construction that: (a) for any choice of $0 \leq t_1 \leq t_2, \dots \leq t_n \leq T$, $S(t_1), S(t_2), \dots, S(t_n)$ is a random vector; and (b) that the process is determined by its finite-dimensional distributions.

Continuity and Regularity of Paths

As discussed in the previous section, a stochastic process is determined by its finite-dimensional distributions. In studying stochastic processes it is often natural to think of them as random functions in t . Let $S(t)$ be defined for $0 \leq t \leq T$, then for a fixed ω it is a function in t called the sample path or a realization of S . Finite-dimensional distributions do not determine the continuity property of sample paths. The following example illustrates this.

Example 2.17: Let $X(t) = 0$ for all t , $0 \leq t \leq 1$, and τ be a uniformly distributed random variable on $[0, 1]$. Let $Y(t) = 0$ for $t \neq \tau$ and $Y(t) = 1$ if $t = \tau$. Then for any fixed t , $P(Y(t) \neq 0) = P(\tau = t) = 0$, hence $P(Y(t) = 0) = 1$, so that all one-dimensional distributions of $X(t)$ and $Y(t)$ are the same. Similarly, all finite-dimensional distributions of X and Y are the same. However, the sample paths of the process X , that is, the functions $X(t)_{0 \leq t \leq 1}$ are continuous in t , whereas every sample path $Y(t)_{0 \leq t \leq 1}$ has a jump at the point τ . Notice that $P(X(t) = Y(t)) = 1$ for all t , $0 \leq t \leq 1$.

Definition 2.27: Two stochastic processes are called versions (modifications) of one another if

$$P(X(t) = Y(t)) = 1 \quad \text{for all } t, 0 \leq t \leq T.$$

Thus the two processes in Example 2.17 are versions of one another; one has continuous sample paths and the other does not. If we agree to pick any version of the process we want, then we can pick the continuous version when it exists. In general, we choose the smoothest possible version of the process.

For two processes X and Y , denote by $N_t = \{X(t) \neq Y(t)\}$, $0 \leq t \leq T$. In Example 2.17, $P(N_t) = P(\tau = t) = 0$ for any t , $0 \leq t \leq 1$. However, $P(\bigcup_{0 \leq t \leq 1} N_t) = P(\tau = t \text{ for some } t \text{ in } [0, 1]) = 1$. Although each of N_t is a P -null set, the union $N = \bigcup_{0 \leq t \leq 1} N_t$ contains uncountably many null sets, and in this case it is a set of probability one.

If it happens that $P(N) = 0$, then N is called an *evanescent* set, and the processes X and Y are called *indistinguishable*. Note that in this case

$P(\{\omega : \exists t : X(t) \neq Y(t)\}) = P(\bigcup_{0 \leq t \leq 1} \{X(t) \neq Y(t)\}) = 0$, and $P(\bigcap_{0 \leq t \leq 1} \{X(t) = Y(t)\}) = P(X(t) = Y(t) \text{ for all } t \in [0, T]) = 1$. It is clear that if the time is discrete, then any two versions of the process are indistinguishable. It is also not hard to see that if $X(t)$ and $Y(t)$ are versions of one another and they both are right-continuous, then they are indistinguishable.

Conditions for the existence of the continuous and the regular (paths with only jump discontinuities) versions of a stochastic process are given below.

Theorem 2.28: $S(t)$, $0 \leq t \leq T$ is an \mathbb{R} -valued stochastic process.

1. If there exist $\alpha > 0$ and $\epsilon > 0$, so that for any $0 \leq u \leq t \leq T$,

$$E|S(t) - S(u)|^\alpha \leq C(t - u)^{1+\epsilon}, \quad (2.28)$$

for some constant C , then there exists a version of S with continuous sample paths, which are Hölder continuous of order $h < \epsilon/\alpha$.

2. If there exist $C > 0$, $\alpha_1 > 0$, $\alpha_2 > 0$ and $\epsilon > 0$, so that for any $0 \leq u \leq v \leq t \leq T$,

$$E(|S(v) - S(u)|^{\alpha_1} |S(t) - S(v)|^{\alpha_2}) \leq C(t - u)^{1+\epsilon}, \quad (2.29)$$

then there exists a version of S with paths that may have discontinuities of the first kind only (which means that at any interior point both right and left limits exist, and one-sided limits exist at the boundaries).

Note that the above result allows us to decide on the existence of the continuous (regular) version by means of the joint bivariate (trivariate) distributions of the process. The same result applies when the process takes values in \mathbb{R}^d , except that the Euclidean distance replaces the absolute value in the above conditions.

Functions without discontinuities of the second kind are considered to be the same if at all points of the domain they have the same right and left limits. In this case it is possible to identify any such function with its right-continuous version.

The following result gives a condition for the existence of a regular right-continuous version of a stochastic process.

Theorem 2.29: If the stochastic process $S(t)$ is right-continuous in probability (that is, for any t the limit in probability $\lim_{u \downarrow t} S(u) = S(t)$) and it does not have discontinuities of the second kind, then it has a right-continuous version.

Other conditions for the regularity of path can be given if we know some particular properties of the process. For example later we give such conditions for processes that are martingales and supermartingales.

σ -Field Generated by a Stochastic Process

$\mathcal{F}_t = \sigma(S_u, u \leq t)$ is the smallest σ -field that contains sets of the form $\{a \leq S_u \leq b\}$ for $0 \leq u \leq t$, $a, b \in \mathbb{R}$. It is the information available to an observer of the process S up to time t .

Filtered Probability Space and Adapted Processes

A filtration \mathbb{F} is a family $\{\mathcal{F}_t\}$ of increasing σ -fields on (Ω, \mathcal{F}) , $\mathcal{F}_t \subset \mathcal{F}$. \mathbb{F} specifies how the information is revealed in time. The property that a filtration is increasing corresponds to the fact that the information is not forgotten.

If we have a set Ω , a σ -field of subsets of Ω , \mathcal{F} , a probability P defined on elements of \mathcal{F} , and a filtration \mathbb{F} such that

$$\mathcal{F}_0 \subset \mathcal{F}_t \subset \cdots \subset \mathcal{F}_T = \mathcal{F},$$

then $(\Omega, \mathcal{F}, \mathbb{F}, P)$ is called a filtered probability space.

A stochastic process on this space $\{S(t), 0 \leq t \leq T\}$ is called *adapted* if for all t , $S(t)$ is \mathcal{F}_t -measurable, that is, if for any t , \mathcal{F}_t contains all the information about $S(t)$ (and may contain extra information).

The Usual Conditions

Filtration is called right-continuous if $\mathcal{F}_{t+} = \mathcal{F}_t$, where

$$\mathcal{F}_{t+} = \bigcap_{s>t} \mathcal{F}_s.$$

The standard assumption (referred to as the usual condition) is that filtration is right-continuous, for all t , $\mathcal{F}_t = \mathcal{F}_{t+}$. It has the following interpretation: any information known immediately after t is also known at t .

Remark 2.2: Note that if $S(t)$ is a process adapted to \mathbb{F} , then we can always take a right-continuous filtration to which $S(t)$ is adapted by taking $\mathcal{G}_t = \mathcal{F}_{t+} = \bigcap_{s>t} \mathcal{F}_s$. Then S_t is \mathcal{G}_t adapted.

The assumption of right-continuous filtration has a number of important consequences. For example, it allows us to assume that martingales, submartingales, and supermartingales have a regular right-continuous version.

It is also assumed that any set which is a subset of a set of zero probability is \mathcal{F}_0 -measurable. Of course, such a set must have zero probability. *A priori* such sets need not be measurable and we enlarge the σ -fields to include such sets. This procedure is called the completion by the null sets.

Martingales, Supermartingales, Submartingales

Definition 2.30: A stochastic process $\{X(t), t \geq 0\}$ adapted to a filtration \mathbb{F} is a supermartingale (submartingale) if for any t it is integrable,

$E|X(t)| < \infty$, and for any $s < t$

$$E(X(t)|\mathcal{F}_s) \leq X(s), \quad (E(X(t)|\mathcal{F}_s) \geq X(s)).$$

If $E(X(t)|\mathcal{F}_s) = X(s)$, then the process $X(t)$ is called a martingale.

An example of a martingale is given by the following

Theorem 2.31 (Doob–Levy Martingale): *Let Y be an integrable random variable, that is, $E|Y| < \infty$, then*

$$M(t) = E(Y|\mathcal{F}_t) \tag{2.30}$$

is a martingale.

PROOF. By the law of double expectation

$$E(M(t)|\mathcal{F}_s) = E(E(Y|\mathcal{F}_t)|\mathcal{F}_s) = E(Y|\mathcal{F}_s) = M(s).$$

□

Using the law of double expectation, it is easy to see that the mean of a martingale is a constant in t , the mean of a supermartingale is non-increasing in t , and the mean of a submartingale is non-decreasing in t .

If $X(t)$ is a supermartingale, then $-X(t)$ is a submartingale, directly from the definition.

We have the following result for the existence of the right-continuous version for super or submartingales without the assumption of continuity in probability imposed on the process (see for example Liptser and Shiryaev (1974)).

Theorem 2.32: *Let the filtration \mathbb{F} be right-continuous and each of the σ -fields \mathcal{F}_t be completed by the P-null sets from \mathcal{F} . In order that the supermartingale $X(t)$ has a right-continuous version it is necessary and sufficient that its mean function $EX(t)$ is right-continuous. In particular, any martingale with right-continuous filtration admits a regular right-continuous version.*

In view of these results, it will often be assumed that the version of the process under consideration is regular and right-continuous (*càdlàg*).

Stopping Times

Definition 2.33: A non-negative random variable τ , which is allowed to take the value ∞ , is called a *stopping time* (with respect to filtration \mathbb{F}) if for each t , the event

$$\{\tau \leq t\} \in \mathcal{F}_t.$$

It is immediately apparent that for all t , the complementary event $\{\tau > t\} \in \mathcal{F}_t$. If $\tau < t$, then for some n , $\tau \leq t - 1/n$. Thus

$$\{\tau < t\} = \bigcup_{n=1}^{\infty} \{\tau \leq t - 1/n\}.$$

The event $\{\tau \leq t - 1/n\} \in \mathcal{F}_{t-1/n}$. Since \mathcal{F}_t are increasing, $\{\tau \leq t - 1/n\} \in \mathcal{F}_t$, therefore $\{\tau < t\} \in \mathcal{F}_t$.

Introduce

$$\mathcal{F}_{t-} := \bigvee_{s < t} \mathcal{F}_s := \sigma\left(\bigcup_{s < t} \mathcal{F}_s\right).$$

The above argument shows that $\{\tau < t\} \in \mathcal{F}_{t-}$.

Theorem 2.34: *If filtration is right-continuous, then τ is a stopping time if, and only if, for each t the event $\{\tau < t\} \in \mathcal{F}_t$.*

PROOF. One direction has just been established, the other one is seen as follows.

$$\{\tau \leq t\} = \bigcap_{n=1}^{\infty} \{\tau < t + 1/n\}.$$

Since $\{\tau < t + 1/n\} \in \mathcal{F}_{t+1/n}$, by right-continuity of \mathbb{F} , $\{\tau \leq t\} \in \mathcal{F}_t$. \square

The assumption of right-continuity of \mathbb{F} is important when studying exit times and hitting times of a set by a process. If $S(t)$ is a random process on \mathbb{R} adapted to \mathbb{F} , then the *hitting time* of set A is defined as

$$T_A = \inf\{t \geq 0 : S(t) \in A\}. \quad (2.31)$$

The first exit time from a set D is defined as

$$\tau_D = \inf\{t \geq 0 : S(t) \notin D\}. \quad (2.32)$$

Note that $\tau_D = T_{\mathbb{R} \setminus D}$.

Theorem 2.35: *Let $S(t)$ be continuous and adapted to \mathbb{F} . If $D = (a, b)$ is an open interval, or any other open set on \mathbb{R} (a countable union of open intervals), then τ_D is a stopping time. If $A = [a, b]$ is a closed interval, or any other closed set on \mathbb{R} (its complement is an open set), then T_A is a stopping time. If in addition the filtration \mathbb{F} is right-continuous, then also for closed sets D and open sets A , τ_D and T_A are stopping times.*

PROOF. $\{\tau_D > t\} = \{S(u) \in D, \text{ for all } u \leq t\} = \cap_{0 \leq u \leq t} \{S(u) \in D\}$. This event is an uncountable intersection over all $u \leq t$ of events in \mathcal{F}_u . The point of the proof is to represent this event as a countable intersection. Due to the continuity of $S(u)$ and D being open, for any irrational u with $S(u) \in D$ there is a rational q with $S(q) \in D$. Therefore

$$\bigcap_{0 \leq u \leq t} \{S(u) \in D\} = \bigcap_{0 \leq q - \text{rational} \leq t} \{S(q) \in D\},$$

which is now a countable intersection of the events from \mathcal{F}_t , and hence is itself in \mathcal{F}_t . This shows that τ_D is a stopping time. Since for any closed set A , $\mathbb{R} \setminus A$ is open, and $T_A = \tau_{\mathbb{R} \setminus A}$, T_A is also a stopping time.

Assume now that filtration is right-continuous. If $D = [a, b]$ is a closed interval, then $D = \cap_{n=1}^{\infty} (a - 1/n, b + 1/n)$. If $D_n = (a - 1/n, b + 1/n)$, then τ_{D_n} is a stopping time, and the event $\{\tau_{D_n} > t\} \in \mathcal{F}_t$. It is easy to see that $\cap_{n=1}^{\infty} \{\tau_{D_n} > t\} = \{\tau_D \geq t\}$, hence $\{\tau_D \geq t\} \in \mathcal{F}_t$, and also $\{\tau_D < t\} \in \mathcal{F}_t$ as its complementary, for any t . The rest of the proof follows by Theorem 2.34. \square

For general processes the following result holds.

Theorem 2.36: *Let $S(t)$ be regular right-continuous and adapted to \mathbb{F} , and \mathbb{F} be right-continuous. If A is an open set on \mathbb{R} , then T_A is a stopping time. If A is a closed set, then $\inf\{t > 0 : S(t) \in A, \text{ or } S(t-) \in A\}$ is a stopping time.*

It is possible, although much harder, to show that the hitting time of a Borel set is a stopping time.

The following results give basic properties of stopping times.

Theorem 2.37: *Let S and T be two stopping times, then $\min(S, T)$, $\max(S, T)$, $S + T$ are all stopping times.*

σ -Field \mathcal{F}_T

If T is a stopping time, events observed before or at time T are described by σ -field \mathcal{F}_T , defined as the collection of sets

$$\mathcal{F}_T = \{A \in \mathcal{F} : \text{for any } t, A \cap \{T \leq t\} \in \mathcal{F}_t\}.$$

Theorem 2.38: *Let S and T be two stopping times. The following properties hold. If $A \in \mathcal{F}_S$, then $A \cap \{S = T\} \in \mathcal{F}_T$, consequently $\{S = T\} \in \mathcal{F}_S \cap \mathcal{F}_T$. If $A \in \mathcal{F}_S$, then $A \cap \{S \leq T\} \in \mathcal{F}_T$, consequently $\{S \leq T\} \in \mathcal{F}_S \cap \mathcal{F}_T$.*

Fubini's Theorem

Fubini's theorem allows us to interchange integrals (sums) and expectations. We give a particular case of Fubini's theorem; it is formulated in the way we use it in applications.

Theorem 2.39: *Let $X(t)$ be a stochastic process $0 \leq t \leq T$ (for all t $X(t)$ is a random variable), with regular sample paths (for all ω at any point t , $X(t)$ has left and right limits). Then*

$$\int_0^T \mathbb{E}|X(t)|dt = \mathbb{E}\left(\int_0^T |X(t)|dt\right).$$

Furthermore, if this quantity is finite, then

$$\mathbb{E}\left(\int_0^T X(t)dt\right) = \int_0^T \mathbb{E}(X(t))dt.$$

Chapter 3

Basic Stochastic Processes

This chapter is mainly about Brownian motion. It is the main process in the calculus of continuous processes. The Poisson process is the main process in the calculus of processes with jumps. Both processes give rise to functions of positive quadratic variation. For stochastic calculus only Chapter 3.1–3.5 are needed, but in applications other sections are also used.

Introduction

Observations of prices of stocks, positions of a diffusing particle, and many other processes observed in time are often modelled by a stochastic process. A stochastic process is an umbrella term for any collection of random variables $\{X(t)\}$ depending on time t . Time can be discrete, for example, $t = 0, 1, 2, \dots$, or continuous, $t \geq 0$. Calculus is suited more to continuous time processes. At any time t , the observation is described by a random variable which we denote by X_t or $X(t)$. A stochastic process $\{X(t)\}$ is frequently denoted by X or with a slight abuse of notation also by $X(t)$.

In practice, we typically observe only a single realization of this process, a single path, out of a multitude of possible paths. Any single path is a function of time t , $x_t = x(t)$, $0 \leq t \leq T$, and the process can also be seen as a random function. In order to describe the distribution and to be able to do probability calculations about the uncertain future, one needs to know the so-called finite-dimensional distributions. Namely, we need to specify how to calculate: probabilities of the form $P(X(t) \leq x)$ for any time t , i.e. the probability distribution of the random variable $X(t)$; probabilities of the form $P(X(t_1) \leq x_1, X(t_2) \leq x_2)$ for any times t_1, t_2 , i.e. the joint bivariate distributions of $X(t_1)$ and $X(t_2)$; and probabilities of the form

$$P(X(t_1) \leq x_1, X(t_2) \leq x_2, \dots, X(t_n) \leq x_n), \quad (3.1)$$

for any choice of time points $0 \leq t_1 < t_2 \dots < t_n \leq T$, and any $n \geq 1$ with $x_1, \dots, x_n \in \mathbb{R}$. Often one does not write the formula for (3.1), but merely points out how to compute it.

3.1 Brownian Motion

Botanist R. Brown described the motion of a pollen particle suspended in fluid in 1828. It was observed that a particle moved in an irregular, random fashion. Einstein, in 1905, argued that the movement is due to bombardment of the particle by the molecules of the fluid; he obtained the equations for Brownian motion. In 1900 L. Bachelier used the Brownian motion as a model for movement of stock prices in his mathematical theory of speculation. The mathematical foundation for Brownian motion as a stochastic process was laid by N. Wiener in 1923, and this process is also called the Wiener process.

The Brownian motion process $B(t)$ serves as a basic model for the cumulative effect of pure noise. If $B(t)$ denotes the position of a particle at time t , then the displacement $B(t) - B(0)$ is the effect of the purely random bombardment by the molecules of the fluid, or the effect of noise over time t .

Defining Properties of Brownian Motion

Brownian motion $\{B(t)\}$ is a stochastic process with the following properties:

1. (Independence of increments) $B(t) - B(s)$, for $t > s$, is independent of the past, that is, of B_u , $0 \leq u \leq s$, or of \mathcal{F}_s , the σ -field generated by $B(u)$, $u \leq s$.
2. (Normal increments) $B(t) - B(s)$ has normal distribution with mean 0 and variance $t - s$. This implies (taking $s = 0$) that $B(t) - B(0)$ has $N(0, t)$ distribution.
3. (Continuity of paths) $B(t)$, $t \geq 0$ are continuous functions of t .

The initial position of Brownian motion is not specified in the definition. When $B(0) = x$, then the process is Brownian motion started at x . Properties 1 and 2 above determine all the finite-dimensional distributions (see (3.4)) and it is possible to show (see Theorem 3.3) that all of them are Gaussian. P_x denotes the probability of events when the process starts at x . The time interval on which Brownian motion is defined is $[0, T]$ for some $T > 0$, which is allowed to be infinite.

We do not prove here that a Brownian motion exists; it can be found in many books on stochastic processes, and one construction is outlined in

Chapter 5.7. However, we can deduce continuity of paths by using normality of increments and appealing to Theorem 2.28. Since

$$\mathbb{E}(B(t) - B(s))^4 = 3(t-s)^2,$$

a continuous version of Brownian motion exists.

Remark 3.1: A definition of Brownian motion in a more general model (that contains extra information) is given by a pair $\{B(t), \mathcal{F}_t\}$, $t \geq 0$, where \mathcal{F}_t is an increasing sequence of σ -fields (a filtration) and $B(t)$ is an adapted process, i.e. $B(t)$ is \mathcal{F}_t measurable, such that properties 1–3 hold.

An important representation used for calculations in processes with independent increments is that for any $s \geq 0$

$$B(t+s) = B(s) + (B(t+s) - B(s)), \quad (3.2)$$

where two variables are independent. An extension of this representation is the process version.

Let $W(t) = B(t+s) - B(s)$. Then for a fixed s , as a process in t , $W(t)$ is a Brownian motion started at zero. This is seen by verifying the defining properties.

Other examples of Brownian motion processes constructed from other processes are given below as well as in the exercises.

Example 3.1: Although $B(t) - B(s)$ is independent of the past, $2B(t) - B(s)$ or $B(t) - 2B(s)$ is not, as, for example, $B(t) - 2B(s) = (B(t) - B(s)) - B(s)$, is a sum of two variables, with only one independent of the past and $B(s)$.

The following example illustrates calculations of some probabilities for Brownian motion.

Example 3.2: Let $B(0) = 0$.

We calculate $P(B(t) \leq 0$ for $t = 2)$ and $P(B(t) \leq 0$ for $t = 0, 1, 2)$.

Since $B(2)$ has normal distribution with mean zero and variance 2,

$$P(B(t) \leq 0 \text{ for } t = 2) = \frac{1}{2}.$$

Since $B(0) = 0$, $P(B(t) \leq 0 \text{ for } t = 0, 1, 2) = P(B(1) \leq 0, B(2) \leq 0)$. Note that $B(2)$ and $B(1)$ are not independent, therefore this probability cannot be calculated as a product $P(B(1) \leq 0)P(B(2) \leq 0) = 1/4$. Using the decomposition $B(2) = B(1) + (B(2) - B(1)) = B(1) + W(1)$, where the two random variables are independent, we have

$$\begin{aligned} P(B(1) \leq 0, B(2) \leq 0) &= P(B(1) \leq 0, B(1) + W(1) \leq 0) \\ &= P(B(1) \leq 0, W(1) \leq -B(1)). \end{aligned}$$

By conditioning and by using Theorem 2.24 and (2.20)

$$P(B(1) \leq 0, W(1) \leq -B(1)) = \int_{-\infty}^0 P(W(1) \leq -x) f(x) dx = \int_{-\infty}^0 \Phi(-x) d\Phi(x),$$

where $\Phi(x)$ and $f(x)$ denote the distribution and the density functions of the standard normal distribution. By changing variables in the last integral, we obtain

$$\int_0^\infty \Phi(x)f(-x)dx = \int_0^\infty \Phi(x)d\Phi(x) = \int_{1/2}^1 ydy = \frac{3}{8}.$$

Transition Probability Functions

If the process is started at x , $B(0) = x$, then $B(t)$ has the $N(x, t)$ distribution. More generally, the conditional distribution of $B(t+s)$ given that $B(s) = x$ is $N(x, t)$. The transition function $P(y, t, x, s)$ is the cumulative distribution function of this distribution,

$$P(y, t, x, s) = P(B(t+s) \leq y | B(s) = x) = P_x(B(t) \leq y).$$

The density function of this distribution is the transition probability density function of Brownian motion,

$$p_t(x, y) = \frac{1}{\sqrt{2\pi t}} e^{-\frac{(y-x)^2}{2t}}. \quad (3.3)$$

The finite-dimensional distributions can be computed with the help of the transition probability density function by using independence of increments in a way similar to that exhibited in the above example.

$$\begin{aligned} & P_x(B(t_1) \leq x_1, B(t_2) \leq x_2, \dots, B(t_n) \leq x_n) \\ &= \int_{-\infty}^{x_1} p_{t_1}(x, y_1) dy_1 \int_{-\infty}^{x_2} p_{t_2-t_1}(y_1, y_2) dy_2 \dots \int_{-\infty}^{x_n} p_{t_n-t_{n-1}}(y_{n-1}, y_n) dy_n. \end{aligned} \quad (3.4)$$

Space Homogeneity

It is easy to see that the one-dimensional distributions of Brownian motion satisfy $P_0(B(t) \in A) = P_x(B(t) \in x + A)$, where A is an interval on the line.

If $B^x(t)$ denotes Brownian motion started at x , then it follows from (3.4) that all finite-dimensional distributions of $B^x(t)$ and $x + B^0(t)$ are the same. Thus $B^x(t) - x$ is Brownian motion started at 0, and $B^0(t) + x$ is Brownian motion started at x , in other words

$$B^x(t) = x + B^0(t). \quad (3.5)$$

The property (3.5) is called the *space-homogeneous* property of Brownian motion.

Definition 3.1: A stochastic process is called space-homogeneous if its finite-dimensional distributions do not change with a shift in space, namely if

$$\begin{aligned} & P(X(t_1) \leq x_1, X(t_2) \leq x_2, \dots, X(t_n) \leq x_n | X(0) = 0) \\ &= P(X(t_1) \leq x_1 + x, X(t_2) \leq x_2 + x, \dots, X(t_n) \leq x_n + x | X(0) = x). \end{aligned}$$

Four realizations of Brownian motion $B = B(t)$ started at zero are shown in Figure 3.1. Although it is a process governed by pure chance with zero mean, it has regions where motion looks like it has “trends”.




Fig. 3.1: Four realizations or paths of Brownian motion $B(t)$.

Brownian Motion as a Gaussian Process

Recall that a process is called Gaussian if all its finite-dimensional distributions are multivariate normal.

Example 3.3: Let random variables X and Y be independent normal with distributions $N(\mu_1, \sigma_1^2)$ and $N(\mu_2, \sigma_2^2)$. Then the distribution of $(X, X+Y)$ is bivariate normal with mean vector $(\mu_1, \mu_1 + \mu_2)$ and covariance matrix $\begin{bmatrix} \sigma_1^2 & \sigma_1^2 \\ \sigma_1^2 & \sigma_1^2 + \sigma_2^2 \end{bmatrix}$.

To see this let $\mathbf{Z} = (Z_1, Z_2)$ have standard normal components, then it is easy to see that

$$(X, X+Y)^T = \boldsymbol{\mu}^T + A\mathbf{Z}^T,$$

where T means the transpose, $\boldsymbol{\mu} = (\mu_1, \mu_1 + \mu_2)$, and matrix $A = \begin{bmatrix} \sigma_1 & 0 \\ \sigma_1 & \sigma_1 + \sigma_2 \end{bmatrix}$. The result follows by the definition of the general normal distribution as a linear transformation of standard normals (see Chapter 2.6).

Similarly to the previous example, the following representation

$$(B(t_1), B(t_2), \dots, B(t_n))$$

$$= (B(t_1), B(t_1) + (B(t_2) - B(t_1)), \dots, B(t_{n-1}) + (B(t_n) - B(t_{n-1})))$$

shows that this vector is a linear transformation of the standard normal vector, hence it has a multivariate normal distribution.

Let $Y_1 = B(t_1)$, and for $k > 1$, $Y_k = B(t_k) - B(t_{k-1})$. Then by the property of independence of increments of Brownian motion, Y_k 's are independent. They also have normal distribution, $Y_1 \sim N(0, t_1)$, and $Y_k \sim N(0, t_k - t_{k-1})$. Thus $(B(t_1), B(t_2), \dots, B(t_n))$ is a linear transformation of (Y_1, Y_2, \dots, Y_n) . But $Y_1 = \sqrt{t_1}Z_1$, and $Y_k = \sqrt{t_k - t_{k-1}}Z_k$, where Z_k 's are independent standard normal. Thus $(B(t_1), B(t_2), \dots, B(t_n))$ is a linear transformation of (Z_1, \dots, Z_n) . Finding the matrix A of this transformation is left as an exercise (Exercise 3.7).

Definition 3.2: The *covariance function* of the process $X(t)$ is defined by

$$\begin{aligned}\gamma(s, t) &= \text{Cov}(X(t), X(s)) = E(X(t) - EX(t))(X(s) - EX(s)) \\ &= E(X(t)X(s)) - EX(t)EX(s).\end{aligned}\quad (3.6)$$

The next result characterizes Brownian motion as a particular Gaussian process.

Theorem 3.3: *A Brownian motion started at zero is a Gaussian process with zero mean function, and covariance function $\min(t, s)$. Conversely, a Gaussian process with zero mean function, and covariance function $\min(t, s)$ is a Brownian motion.*

PROOF. Since the mean of the Brownian motion is zero,

$$\gamma(s, t) = \text{Cov}(B(t), B(s)) = E(B(t)B(s)).$$

If $t < s$, then $B(s) = B(t) + B(s) - B(t)$, and

$$E(B(t)B(s)) = EB^2(t) + E(B(t)(B(s) - B(t))) = EB^2(t) = t,$$

where we used the independence of increments. Similarly, if $t > s$, $E(B(t)B(s)) = s$. Therefore

$$E(B(t)B(s)) = \min(t, s).$$

To show the converse, let t be arbitrary and $s \geq 0$. $X(t)$ is a Gaussian process, thus the joint distribution of $X(t), X(t+s)$ is a bivariate normal, and by conditions has zero mean. Therefore the vector $(X(t), X(t+s) - X(t))$ is also

bivariate normal. For any $u \leq t$, the variables $X(u)$ and $X(t+s) - X(t)$ are uncorrelated, using $\text{Cov}(X(t), X(s)) = \min(t, s)$,

$$\begin{aligned} & \text{Cov}(X(u), X(t+s) - X(t)) \\ &= \text{Cov}(X(u), X(t+s)) - \text{Cov}(X(u), X(t)) = u - u = 0. \end{aligned}$$

A property of the bivariate normal distribution implies that these variables are independent. Thus the increment $X(t+s) - X(t)$ is independent of $X(u)$, $u \leq t$, and has $N(0, s)$ distribution. Therefore it is a Brownian motion. \square

Example 3.4: We find the distribution of $B(1) + B(2) + B(3) + B(4)$.

Consider the random vector $\mathbf{X} = (B(1), B(2), B(3), B(4))$. Since Brownian motion is a Gaussian process, all its finite-dimensional distributions are normal, in particular \mathbf{X} has a multivariate normal distribution with mean vector zero and covariance matrix given by $\sigma_{ij} = \text{Cov}(X_i, X_j)$. For example, $\text{Cov}(X_1, X_3) = \text{Cov}((B(1), B(3)) = 1$.

$$\Sigma = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}$$

Now, let $a = (1, 1, 1, 1)$. Then

$$\mathbf{a}\mathbf{X} = X_1 + X_2 + X_3 + X_4 = B(1) + B(2) + B(3) + B(4).$$

$\mathbf{a}\mathbf{X}$ has a normal distribution with mean zero and variance $\mathbf{a}\Sigma\mathbf{a}^T$, and in this case the variance is given by the sum of the elements of the covariance matrix. Thus $B(1) + B(2) + B(3) + B(4)$ has a normal distribution with mean zero and variance 30. Alternatively, we can calculate the variance of the sum by the formula

$$\text{Var}(X_1 + X_2 + X_3 + X_4)$$

$$= \text{Cov}(X_1 + X_2 + X_3 + X_4, X_1 + X_2 + X_3 + X_4) = \sum_{i,j} \text{Cov}(X_i, X_j) = 30.$$

Example 3.5: In order to illustrate the use of scaling, we find the distribution of $B(\frac{1}{4}) + B(\frac{1}{2}) + B(\frac{3}{4}) + B(1)$. Consider the random vector $\mathbf{Y} = (B(\frac{1}{4}), B(\frac{1}{2}), B(\frac{3}{4}), B(1))$. It is easy to see that \mathbf{Y} and $\frac{1}{2}\mathbf{X}$, where $\mathbf{X} = (B(1), B(2), B(3), B(4))$ have the same law. Therefore its covariance matrix is given by $\frac{1}{4}\Sigma$, with Σ as above. Consequently, $\mathbf{a}\mathbf{Y}$ has a normal distribution with mean zero and variance $30/4$.

Example 3.6: We find the probability $P(\int_0^1 B(t)dt > \frac{2}{\sqrt{3}})$.

Note first that since Brownian motion has continuous paths, the Riemann integral $\int_0^1 B(t)dt$ is well-defined for any random path as we integrate path by path. In order to find the required probability we need to know the distribution of $\int_0^1 B(t)dt$. This can be obtained as a limit of the distributions of the approximating sums,

$$\sum B(t_i)\Delta,$$

where points t_i partition $[0, 1]$ and $\Delta = t_{i+1} - t_i$. If, for example, $t_i = i/n$, then for $n = 4$ the approximating sum is $\frac{1}{4}(B(\frac{1}{4}) + B(\frac{1}{2}) + B(\frac{3}{4}) + B(1))$, the distribution of which was found in Example 3.5 to be $N(0, \frac{15}{32})$. Similarly, the distribution of all of the approximating sums is normal with zero mean. It can be shown that the limit of Gaussian distributions is a Gaussian distribution. Thus $\int_0^1 B(t)dt$ has a normal distribution with zero mean. Therefore it only remains to compute its variance.

$$\begin{aligned} \text{Var}\left(\int_0^1 B(t)dt\right) &= \text{Cov}\left(\int_0^1 B(t)dt, \int_0^1 B(s)ds\right) \\ &= \mathbb{E}\left(\int_0^1 B(t)dt \int_0^1 B(s)ds\right) = \int_0^1 \int_0^1 \mathbb{E}(B(t)B(s)) dt ds \\ &= \int_0^1 \int_0^1 \text{Cov}(B(t), B(s)) dt ds = \int_0^1 \int_0^1 \min(t, s) dt ds = 1/3. \end{aligned}$$

Exchanging the integrals and expectation is justified by Fubini's theorem since

$$\int_0^1 \int_0^1 \mathbb{E}|B(t)B(s)| dt ds \leq \int_0^1 \int_0^1 \sqrt{ts} dt ds < 1.$$

Thus $\int_0^1 B(t)dt$ has $N(0, 1/3)$ distribution, and the desired probability is approximately 0.025. Later we shall prove that the distribution of the integral $\int_0^a B(t)dt$ is normal $N(0, a^3/3)$ by considering a transformation to the Itô integral (see Example 6.4).

Brownian Motion as a Random Series

The process

$$\xi_0 \frac{t}{\sqrt{\pi}} + \sqrt{\frac{2}{\pi}} \sum_{j=1}^{\infty} \frac{\sin(jt)}{j} \xi_j, \quad (3.7)$$

where ξ_j 's $j = 0, 1, \dots$, are independent standard normal random variables, is Brownian motion on $[0, \pi]$. Convergence of the series is understood almost surely. This representation resembles the example of a continuous but nowhere differentiable function (see Example 1.2). One can prove the assertion by showing that the partial sums converge uniformly, and by verifying that the process in (3.7) is Gaussian, has zero mean, and covariance $\min(s, t)$ (see for example Breiman (1968), Itô and McKean (1965)).

Remark 3.2: A similar, more general representation of a Brownian motion is given by using an orthonormal sequence of functions on $[0, T]$, $h_j(t)$. $B(t) = \sum_{j=0}^{\infty} \xi_j H_j(t)$, where $H_j(t) = \int_0^t h_j(s)ds$, is a Brownian motion on $[0, T]$.

3.2 Properties of Brownian Motion Paths

An occurrence of Brownian motion observed from time 0 to time T , is a random function of t on the interval $[0, T]$. It is called a *realization*, a *path*, or *trajectory*.

Quadratic Variation of Brownian Motion

The quadratic variation of Brownian motion $[B, B](t)$ is defined as

$$[B, B](t) = [B, B]([0, t]) = \lim \sum_{i=1}^n |B(t_i^n) - B(t_{i-1}^n)|^2, \quad (3.8)$$

where the limit is taken over all shrinking partitions of $[0, t]$, with $\delta_n = \max_i(t_{i+1}^n - t_i^n) \rightarrow 0$ as $n \rightarrow \infty$. It is remarkable that although the sums in definition (3.8) are random, their limit is non-random, as the following result shows.

Theorem 3.4: *Quadratic variation of a Brownian motion over $[0, t]$ is t .*

PROOF. We give the proof for a sequence of partitions for which $\sum_n \delta_n < \infty$. An example of this is when the interval is divided into two, then each subinterval is divided into two, etc. Let $T_n = \sum_i |B(t_i^n) - B(t_{i-1}^n)|^2$. It is easy to see that

$$\mathbb{E}(T_n) = \mathbb{E} \sum_i |B(t_i^n) - B(t_{i-1}^n)|^2 = \sum_{i=1}^n (t_i^n - t_{i-1}^n)^2 = t - 0 = t.$$

By using the fourth moment of $N(0, \sigma^2)$ distribution is $3\sigma^4$, we obtain the variance of T_n

$$\begin{aligned} \text{Var}(T_n) &= \text{Var} \left(\sum_i |B(t_i^n) - B(t_{i-1}^n)|^2 \right) = \sum_i \text{Var}(B(t_i^n) - B(t_{i-1}^n))^2 \\ &= \sum_i 2(t_i^n - t_{i-1}^n)^2 \leq 2 \max(t_i^n - t_{i-1}^n)t = 2t\delta_n. \end{aligned}$$

Therefore $\sum_{n=1}^{\infty} \text{Var}(T_n) < \infty$. Using the monotone convergence theorem, we find $\mathbb{E} \sum_{n=1}^{\infty} (T_n - \mathbb{E} T_n)^2 < \infty$. This implies that the series inside the expectation converges almost surely. Hence its terms converge to zero, and $T_n - \mathbb{E} T_n \rightarrow 0$ almost surely, consequently $T_n \rightarrow t$ almost surely.

It is possible to show that $T_n \rightarrow t$ almost surely for any sequence of partitions which are successive refinements and satisfy $\delta_n \rightarrow 0$ as $n \rightarrow \infty$ (see for example Loeve (1978) for the proof, or Breiman (1968)). \square

Varying t , the quadratic variation process of Brownian motion is t . Note that the classic quadratic variation of Brownian paths (defined as the supremum over all partitions of sums in Equation (3.8) (see Chapter 1) is infinite (see for example Freedman (1971)).

Properties of Brownian Paths

$B(t)$'s as functions of t have the following properties. Almost every sample path $B(t)$, $0 \leq t \leq T$ is

1. A continuous function of t .
2. Not monotone in any interval, no matter how small the interval is.
3. Not differentiable at any point.
4. Has infinite variation on any interval, no matter how small it is.
5. Has quadratic variation on $[0, t]$ equal to t , for any t .

Properties 1 and 3 of Brownian motion paths state that although any realization $B(t)$ is a continuous function of t , it has increments $\Delta B(t)$ over an interval of length Δt much larger than Δt as $\Delta t \rightarrow 0$. Since $E(\underline{B}(t + \Delta t) - B(t))^2 = \Delta t$, it suggests that the increment is roughly like $\sqrt{\Delta t}$. This is made precise by the quadratic variation property 5.

Note that by Theorem 1.10, a positive quadratic variation implies infinite variation so that property 4 follows from property 5. Since a monotone function has finite variation, property 2 follows from property 4.

According to Theorem 1.8 a continuous function with a bounded derivative is of finite variation. Therefore it follows from property 4 that $B(t)$ can not have a bounded derivative on any interval, no matter how small the interval is. It is not yet the non-differentiability at any point, but it is close to it. For the proof of the result that with probability one Brownian motion paths are nowhere differentiable (due to Dvoretzki, Erdős and Kakutani) see Breiman (1968). Here we show a simple statement.

Theorem 3.5: *For any t almost all trajectories of Brownian motion are not differentiable at t .*

PROOF. Consider $\frac{B(t+\Delta)-B(t)}{\Delta} = \frac{\sqrt{\Delta}Z}{\Delta} = \frac{Z}{\sqrt{\Delta}}$, for some standard normal random variable Z . Thus the ratio converges to ∞ in distribution, since $P(|\frac{Z}{\sqrt{\Delta}}| > K) \rightarrow 1$ for any K , as $\Delta \rightarrow 0$, precluding existence of the derivative at t . \square

To realize the above argument on a computer take, for example, $\Delta = 10^{-20}$. Then $\Delta B(t) = 10^{-10}Z$, and $\Delta B(t)/\Delta = 10^{10}Z$, which is very large in absolute value with overwhelming probability.

3.3 Three Martingales of Brownian Motion

In this section three main martingales associated with Brownian motion are given. Recall the definition of a martingale.

Definition 3.6: A stochastic process $\{X(t), t \geq 0\}$ is a martingale if for any t it is integrable, $E|X(t)| < \infty$, and for any $s > 0$

$$E(X(t+s)|\mathcal{F}_t) = X(t), \quad \text{a.s.} \quad (3.9)$$

where \mathcal{F}_t is the information about the process up to time t , and the equality holds almost surely.

The martingale property means that if we know the values of the process up to time t , and $X(t) = x$, then the expected future value at any future time is x .

Remark 3.3: \mathcal{F}_t represents information available to an observer at time t . A set $A \in \mathcal{F}_t$ if, and only if, by observing the process up to time t one can decide whether or not A has occurred. Formally, $\mathcal{F}_t = \sigma(X(s), 0 \leq s \leq t)$ denotes the σ -field (σ -algebra) generated by the values of the process up to time t .

Remark 3.4: As the conditional expectation given a σ -field is defined as a random variable (see for example, Chapter 2.7), all the relations involving conditional expectations such as equalities and inequalities must be understood in the almost sure sense. This will always be assumed, and the almost sure “a.s.” specification will be frequently dropped.

Examples of martingales constructed from Brownian motion are given in the next result.

Theorem 3.7: *Let $B(t)$ be Brownian Motion. Then*

1. $B(t)$ is a martingale.
2. $B(t)^2 - t$ is a martingale.
3. For any u , $e^{uB(t) - \frac{u^2}{2}t}$ is a martingale.

PROOF. The key idea in establishing the martingale property is that for any function g , the conditional expectation of $g(B(t+s) - B(t))$ given \mathcal{F}_t is equal to the unconditional one,

$$E(g(B(t+s) - B(t))|\mathcal{F}_t) = E(g(B(t+s) - B(t))), \quad (3.10)$$

due to independence of $B(t+s) - B(t)$ and \mathcal{F}_t . The latter expectation is just $Eg(X)$, where X is a normal $N(0, s)$ random variable.

1. By definition, $B(t) \sim N(0, t)$, so that $B(t)$ is integrable with $E(B(t)) = 0$.

$$\begin{aligned} E(B(t+s)|\mathcal{F}_t) &= E(B(t) + (B(t+s) - B(t))|\mathcal{F}_t) \\ &= E(B(t)|\mathcal{F}_t) + E(B(t+s) - B(t)|\mathcal{F}_t) \\ &= B(t) + E(B(t+s) - B(t)) = B(t). \end{aligned}$$

2. By definition, $E(B^2(t)) = t < \infty$, therefore $B^2(t)$ is integrable. Since

$$\begin{aligned} B^2(t+s) &= (B(t) + B(t+s) - B(t))^2 \\ &= B^2(t) + 2B(t)(B(t+s) - B(t)) + (B(t+s) - B(t))^2, \\ E(B^2(t+s)|\mathcal{F}_t) &= B^2(t) + 2E(B(t)(B(t+s) - B(t))|\mathcal{F}_t) \\ &\quad + E((B(t+s) - B(t))^2|\mathcal{F}_t) \\ &= B^2(t) + s, \end{aligned}$$

where we used that $B(t+s) - B(t)$ is independent of \mathcal{F}_t and has mean 0, and (3.10) with $g(x) = x^2$. Subtracting $(t+s)$ from both sides gives the martingale property of $B^2(t) - t$.

3. Consider the moment generating function of $B(t)$,

$$E(e^{uB(t)}) = e^{tu^2/2} < \infty,$$

since $B(t)$ has the $N(0, t)$ distribution. This implies integrability of $e^{uB(t)-tu^2/2}$, moreover

$$E(e^{uB(t)-tu^2/2}) = 1.$$

The martingale property is established by using (3.10) with $g(x) = e^{ux}$.

$$\begin{aligned} E(e^{uB(t+s)}|\mathcal{F}_t) &= E(e^{uB(t)+u(B(t+s)-B(t))}|\mathcal{F}_t) \\ &= e^{uB(t)}E(e^{u(B(t+s)-B(t))}|\mathcal{F}_t) \quad (\text{since } B(t) \text{ is } \mathcal{F}_t\text{-measurable}) \\ &= e^{uB(t)}E(e^{u(B(t+s)-B(t))}) \quad (\text{since increment is independent of } \mathcal{F}_t) \\ &= e^{\frac{u^2}{2}s}e^{uB(t)}. \end{aligned}$$

The martingale property of $e^{uB(t)-tu^2/2}$ is obtained by multiplying both sides by $e^{-\frac{u^2}{2}(t+s)}$. \square

Remark 3.5: All three martingales have a central place in the theory. The martingale $B^2(t) - t$ provides a characterization (Levy's characterization) of Brownian motion. It will be seen later that if a process $X(t)$ is a continuous martingale such that $X^2(t) - t$ is also a martingale, then $X(t)$ is Brownian motion. The martingale $e^{uB(t)-tu^2/2}$ is known as the exponential martingale, and as it is related to the moment generating function, it is used for establishing distributional properties of the process.

3.4 Markov Property of Brownian Motion

The Markov property states that if we know the present state of the process, then the future behaviour of the process is independent of its past. The process $X(t)$ has the Markov property if the conditional distribution of $X(t+s)$ given $X(t) = x$, does not depend on the past values (but it may depend on the present value x). The process “does not remember” how it got to the present state x . Let \mathcal{F}_t denote the σ -field generated by the process up to time t .

Definition 3.8: X is a Markov process if for any t and $s > 0$, the conditional distribution of $X(t+s)$ given \mathcal{F}_t is the same as the conditional distribution of $X(t+s)$ given $X(t)$, that is,

$$\mathrm{P}(X(t+s) \leq y | \mathcal{F}_t) = \mathrm{P}(X(t+s) \leq y | X(t)), \quad \text{a.s.} \quad (3.11)$$

Theorem 3.9: *Brownian motion $B(t)$ possesses the Markov property.*

PROOF. It is easy to see by using the moment generating function that the conditional distribution of $B(t+s)$ given \mathcal{F}_t is the same as that given $B(t)$. Indeed,

$$\begin{aligned} \mathrm{E}(e^{uB(t+s)} | \mathcal{F}_t) &= e^{uB(t)} \mathrm{E}(e^{u(B(t+s)-B(t))} | \mathcal{F}_t) \\ &= e^{uB(t)} \mathrm{E}(e^{u(B(t+s)-B(t))}) \quad (\text{since } e^{u(B(t+s)-B(t))} \text{ is} \\ &\quad \text{independent of } \mathcal{F}_t) \\ &= e^{uB(t)} e^{u^2 s/2} \quad (\text{since } B(t+s) - B(t) \text{ is } N(0, s)) \\ &= e^{uB(t)} \mathrm{E}(e^{u(B(t+s)-B(t))} | B(t)) = \mathrm{E}(e^{uB(t+s)} | B(t)). \quad \square \end{aligned}$$

The transition probability function of a Markov process X is defined as

$$P(y, t, x, s) = \mathrm{P}(X(t) \leq y | X(s) = x),$$

the conditional distribution function of the process at time t , given that it is at point x at time $s < t$. It is possible to choose these functions so that for any

fixed x they are true probabilities on the line. In the case of Brownian motion it is given by the distribution function of the normal $N(x, t - s)$ distribution

$$P(y, t, x, s) = \int_{-\infty}^y \frac{1}{\sqrt{2\pi(t-s)}} e^{\frac{(u-x)^2}{2(t-s)}} du.$$

The transition probability function of Brownian motion satisfies $P(y, t, x, s) = P(y, t - s, x, 0)$. In other words,

$$P(B(t) \leq y | B(s) = x) = P(B(t - s) \leq y | B(0) = x). \quad (3.12)$$

For fixed x and t , $P(y, t, x, 0)$ has the density $p_t(x, y)$ which is given by (3.3). The property (3.12) states that Brownian motion is time-homogeneous, that is, its distributions do not change with a shift in time. For example, the distribution of $B(t)$ given $B(s) = x$ is the same as that of $B(t - s)$ given $B(0) = x$. It follows from property (3.12) and function (3.4) that all finite-dimensional distributions of Brownian motion are time-homogeneous.

In what follows, P_x denotes the conditional probability given $B(0) = x$. More information on transition functions is given in Chapter 5.5.

Stopping Times and Strong Markov Property

Definition 3.10: A random time T is called a stopping time for $B(t)$, $t \geq 0$, if for any t it is possible to decide whether T has occurred or not by observing $B(s)$, $0 \leq s \leq t$. More rigorously, for any t the sets $\{T \leq t\} \in \mathcal{F}_t$, the σ -field generated by $B(s)$, $0 \leq s \leq t$.

Example 3.7: Examples of stopping times and random times.

1. Any non-random time T is a stopping time. Formally, $\{T \leq t\}$ is either the \emptyset or Ω , which are members of \mathcal{F}_t for any t .
2. Let T be the first time $B(t)$ takes value (hits) 1. Then T is a stopping time. Clearly, if we know $B(s)$ for all $s \leq t$, then we know whether the Brownian motion took value 1 before or at t or not. Thus we know that $\{T \leq t\}$ has occurred or not just by observing the past of the process prior to t . Formally, $\{T > t\} = \{B(u) < 1, \text{ for all } u \leq t\} \in \mathcal{F}_t$.
3. Similarly, the first passage time of level a , $T_a = \inf\{t > 0 : B(t) = a\}$ is a stopping time.
4. Let T be the time when Brownian motion reaches its maximum on the interval $[0, 1]$. Then clearly, in order to decide whether $\{T \leq t\}$ has occurred or not, it is not enough to know the values of the process prior to t ; one needs to know all the values on the interval $[0, 1]$ so that T is not a stopping time.
5. Let T be the last zero of Brownian motion before time $t = 1$. Then T is not a stopping time since if $T \leq t$, then there are no zeros in $(t, 1]$, which is the event that is decided by observing the process up to time 1, and this set does not belong to \mathcal{F}_t .

The strong Markov property is similar to the Markov property, except that in the definition a fixed time t is replaced by a stopping time T .

Theorem 3.11: *Brownian motion $B(t)$ has the strong Markov property: for any finite stopping time T the regular conditional distribution of $B(T+t), t \geq 0$ given \mathcal{F}_T is $P_{B(T)}$, that is,*

$$P(B(T+t) \leq y | \mathcal{F}_T) = P(B(T+t) \leq y | B(T)) \quad \text{almost surely.}$$

Corollary 3.12: *Let T be a finite stopping time. Define the new process in $t \geq 0$ by*

$$\hat{B}(t) = B(T+t) - B(T). \quad (3.13)$$

Then $\hat{B}(t)$ is a Brownian motion which started at zero and is independent of \mathcal{F}_T .

We do not give the proof of the strong Markov property here, but it can be found for example in Rogers and Williams (1994), and can be done by using the exponential martingale and the optional stopping theorem given in Chapter 7.

Note that the strong Markov property applies only when T is a stopping time. If T is just a random time, then $B(T+t) - B(T)$ need not be Brownian motion.

3.5 Hitting Times and Exit Times

Let T_x denote the first time $B(t)$ hits level x , $T_x = \inf\{t > 0 : B(t) = x\}$. Denote the time to exit an interval (a, b) by $\tau = \min(T_a, T_b)$.

Theorem 3.13: *Let $a < x < b$, and $\tau = \min(T_a, T_b)$. Then $P_x(\tau < \infty) = 1$ and $E_x \tau < \infty$.*

PROOF. $\{\tau > 1\} = \{a < B(s) < b, \text{ for all } 0 \leq s \leq 1\} \subset \{a < B(1) < b\}$. Therefore we have

$$P_x(\tau > 1) \leq P_x(B(1) \in (a, b)) = \frac{1}{\sqrt{2\pi}} \int_a^b e^{-(x-y)^2/2} dy.$$

The function $P_x(B(1) \in (a, b))$ is continuous in x on $[a, b]$, hence it reaches its maximum $\theta < 1$. By using the strong Markov property we can show that $P_x(\tau > n) \leq \theta^n$. For any non-negative random variable $X \geq 0$, $EX \leq \sum_{n=0}^{\infty} P(X > n)$ (see Exercise 3.2). Therefore,

$$E_x \tau \leq \sum_{n=0}^{\infty} \theta^n = \frac{1}{1-\theta} < \infty.$$

The bound on $P_x(\tau > n)$ is established as follows.

$$\begin{aligned}
 P_x(\tau > n) &= P_x(B(s) \in (a, b), 0 \leq s \leq n) \\
 &= P_x(B(s) \in (a, b), 0 \leq s \leq n-1, B(s) \in (a, b), n-1 \leq s \leq n) \\
 &= P_x(\tau > n-1, B(s) \in (a, b), n-1 \leq s \leq n) \\
 &= P_x(\tau > n-1, B(n-1) + \hat{B}(s) \in (a, b), 0 \leq s \leq 1) \quad \text{by (3.13)} \\
 &= E_x(I(\tau > n-1)I(B(n-1) + \hat{B}(s) \in (a, b), 0 \leq s \leq 1)) \\
 &= E_x(E_x(I(\tau > n-1)I(B(n-1) + \hat{B}(s) \in (a, b), 0 \leq s \leq 1)) | \mathcal{F}_{n-1}) \\
 &= E_x(I(\tau > n-1)E_x(I(B(n-1) + \hat{B}(s) \in (a, b), 0 \leq s \leq 1)) | \mathcal{F}_{n-1}).
 \end{aligned}$$

The conditional expectation inside is the conditional probability $P_x(B(n-1) + \hat{B}(s) \in (a, b), 0 \leq s \leq 1) | \mathcal{F}_{n-1}$, i.e. the probability of not exiting during time 1, which by earlier assertion does not exceed θ ,

$$\begin{aligned}
 &P_x(B(n-1) + \hat{B}(s) \in (a, b), 0 \leq s \leq 1) | \mathcal{F}_{n-1} \\
 &\leq \max_{y \in (a, b)} P_y(\hat{B}(s) \in (a, b), 0 \leq s \leq 1) = \theta.
 \end{aligned}$$

Proceeding, we have

$$P_x(\tau > n) \leq \theta P_x(\tau > n-1) \leq \theta^n. \quad \square$$

The next result gives the recurrence property of Brownian motion.

Theorem 3.14:

$$P_a(T_b < \infty) = 1, \quad P_a(T_a < \infty) = 1.$$

PROOF. The second statement follows from the first, since

$$P_a(T_a < \infty) \geq P_a(T_b < \infty)P_b(T_a < \infty) = 1.$$

We show now that $P_0(T_1 < \infty) = 1$, for other points the proof is similar. Observe firstly that by the previous result and by symmetry, for any a and b

$$P_{\frac{a+b}{2}}(T_a < T_b) = \frac{1}{2}.$$

Hence $P_0(T_{-1} < T_1) = \frac{1}{2}$, $P_{-1}(T_{-3} < T_1) = \frac{1}{2}$, $P_{-3}(T_{-7} < T_1) = \frac{1}{2}$, etc. Consider now $P_0(T_{-(2^n-1)} < T_1)$. Since the paths of Brownian motion are continuous, in order to reach $-(2^n - 1)$ the path must reach -1 first, then it must go from -1 to -3 , etc. Hence we obtain

$$\begin{aligned}
 &P_0(T_{-(2^n-1)} < T_1) \\
 &= P_0(T_{-1} < T_1)P_{-1}(T_{-3} < T_1) \dots P_{-(2^{n-1}-1)}(T_{-(2^n-1)} < T_1) = \frac{1}{2^n}.
 \end{aligned}$$

If A_n denotes the event that Brownian motion hits $-(2^n - 1)$ before it hits 1, then we showed that $P(A_n) = 2^{-n}$. Note that $A_n \subset A_{n-1}$, as if Brownian motion hits $-(2^n - 1)$ before 1, it also hits the points bigger than $-(2^n - 1)$. Thus

$$\bigcap_{i=1}^n A_i = A_n$$

and

$$P\left(\bigcap_{i=1}^{\infty} A_i\right) = \lim_{n \rightarrow \infty} P(A_n) = \lim_{n \rightarrow \infty} 2^{-n} = 0.$$

This implies that

$$P\left(\bigcup_{n=1}^{\infty} A_n^c\right) = 1.$$

In other words, with probability 1, one of the events complementary to A_n occurs, that is, there is n such that Brownian motion hits 1 before it hits $-(2^n - 1)$. This implies that $P_0(T_1 < \infty) = 1$. Another proof of this fact that uses properties of martingales is given in Chapter 7. \square

3.6 Maximum and Minimum of Brownian Motion

In this section we establish the distribution of the maximum and the minimum of Brownian motion on $[0, t]$,

$$M(t) = \max_{0 \leq s \leq t} B(s) \quad \text{and} \quad m(t) = \min_{0 \leq s \leq t} B(s),$$

as well as the distribution of the first hitting (passage) time of x , $T_x = \inf\{t > 0 : B(t) = x\}$.

Theorem 3.15: *For any $x > 0$,*

$$P_0(M(t) \geq x) = 2P_0(B(t) \geq x) = 2\left(1 - \Phi\left(\frac{x}{\sqrt{t}}\right)\right),$$

where $\Phi(x)$ stands for the standard normal distribution function.

PROOF. Notice that the events $\{M(t) \geq x\}$ and $\{T_x \leq t\}$ are the same. Indeed, if the maximum at time t is greater than x , then at some time before t Brownian motion took value x , and if Brownian motion took value x at some time before t , then the maximum will be at least x . Since

$$\{B(t) \geq x\} \subset \{T_x \leq t\}$$

$$P(B(t) \geq x) = P(B(t) \geq x, T_x \leq t).$$

As $B(T_x) = x$,

$$\mathbb{P}(B(t) \geq x) = \mathbb{P}(T_x \leq t, B(T_x + (t - T_x)) - B(T_x) \geq 0).$$

According to Theorem 3.14, T_x is a finite stopping time, and by the strong Markov property (3.13) the random variable $\hat{B}(s) = B(T_x + s) - B(T_x)$ is independent of \mathcal{F}_{T_x} and has a normal distribution; we therefore have

$$\mathbb{P}(B(t) \geq x) = \mathbb{P}(T_x \leq t, \hat{B}(t - T_x) \geq 0). \quad (3.14)$$

If we had s independent of T_x , then

$$\begin{aligned} \mathbb{P}(T_x \leq t, \hat{B}(s) \geq 0) &= \mathbb{P}(T_x \leq t) \mathbb{P}(\hat{B}(s) \geq 0) \\ &= \mathbb{P}(T_x \leq t) \frac{1}{2} = \mathbb{P}(M(t) \geq x) \frac{1}{2}, \end{aligned} \quad (3.15)$$

and we are done. However, in (3.14) $s = t - T_x$, and is clearly dependent on T_x . It is not easy to show that

$$\begin{aligned} \mathbb{P}(B(t) \geq x) &= \mathbb{P}(T_x \leq t, \hat{B}(t - T_x) \geq 0) \\ &= \mathbb{P}(T_x \leq t) \frac{1}{2} = \mathbb{P}(M(t) \geq x) \frac{1}{2}. \end{aligned}$$

The proof can be found, for example, in Dudley (1989). \square

A simple application of the result is given in Example 3.8, from which it follows that Brownian motion changes sign in $(0, \varepsilon)$, for any ε however small.

Example 3.8: We find the probability $\mathbb{P}(B(t) \leq 0)$ for all $t, 0 \leq t \leq 1$. Note that the required probability involves uncountably many random variables: all $B(t)$'s are less than or equal to zero, $0 \leq t \leq 1$; we want to know the probability that the entire path from 0 to 1 will stay below 0. We could calculate the desired probability for n values of the process and then take the limit as $n \rightarrow \infty$. However, it is simpler in this case to express this probability as a function of the whole path. All $B(t)$'s are less than or equal zero if, and only if, their maximum is less than or equal to zero.

$$\{B(t) \leq 0 \text{ for all } t, 0 \leq t \leq 1\} = \{\max_{0 \leq t \leq 1} B(t) \leq 0\},$$

and consequently these events have the same probabilities. Now,

$$\mathbb{P}(\max_{0 \leq t \leq 1} B(t) \leq 0) = 1 - \mathbb{P}(\max_{0 \leq t \leq 1} B(t) > 0).$$

According to the law of the maximum of Brownian motion,

$$\mathbb{P}(\max_{0 \leq t \leq 1} B(t) > 0) = 2\mathbb{P}(B(1) > 0) = 1.$$

Hence $\mathbb{P}(B(t) \leq 0)$ for all $t, 0 \leq t \leq 1) = 0$.

In order to find the distribution of the minimum of Brownian motion $m(t) = \min_{0 \leq s \leq t} B(s)$, we use the symmetry argument, and that

$$-\min_{0 \leq s \leq t} B(s) = \max_{0 \leq s \leq t} (-B(s)).$$

Theorem 3.16: *If $B(t)$ is a Brownian motion with $B(0) = 0$, then $\hat{B}(t) = -B(t)$ is also a Brownian motion with $\hat{B}(0) = 0$.*

PROOF. The process $\hat{B}(t) = -B(t)$ has independent and normally distributed increments. It also has continuous paths, therefore it is Brownian motion. \square

Theorem 3.17: *For any $x < 0$*

$$P_0\left(\min_{0 \leq s \leq t} B(s) \leq x\right) = 2P_0(B(t) \geq -x) = 2P_0(B(t) \leq x).$$

The proof is straightforward and is left as an exercise.

3.7 Distribution of Hitting Times

T_x is finite by Theorem 3.14. The result below gives the distribution of T_x and establishes that T_x has infinite mean.

Theorem 3.18: *The probability density of T_x is given by*

$$f_{T_x}(t) = \frac{|x|}{\sqrt{2\pi}} t^{-\frac{3}{2}} e^{-\frac{x^2}{2t}},$$

which is the inverse gamma density with parameters $\frac{1}{2}$ and $\frac{x^2}{2}$. $E_0 T_x = +\infty$.

PROOF. Take $x > 0$. The events $\{M(t) \geq x\}$ and $\{T_x \leq t\}$ are the same, so that

$$\begin{aligned} P(T_x \leq t) &= P(M(t) \geq x) \\ &= 2P(B(t) \geq x) = \int_x^\infty \sqrt{\frac{2}{\pi t}} e^{-\frac{y^2}{2t}} dy. \end{aligned}$$

The formula for the density of T_x is obtained by differentiation after the change of variables $u = \frac{y}{\sqrt{t}}$ in the integral. Finally,

$$E_0 T_x = \frac{|x|}{\sqrt{2\pi}} \int_0^\infty t^{-\frac{1}{2}} e^{-\frac{x^2}{2t}} dt = \infty, \quad \text{since } t^{-\frac{1}{2}} e^{-\frac{x^2}{2t}} \sim 1/\sqrt{t}, \quad t \rightarrow \infty.$$

For $x < 0$ the proof is similar. \square

Remark 3.6: The property $P(T_x < \infty) = 1$ is called the *recurrence* property of Brownian motion. Although $P(T_x < \infty) = 1$, $E(T_x) = \infty$, even though x is visited with probability one, the expected time for it to happen is infinite.

The next result looks at hitting times T_x as a process in x .

Theorem 3.19: *The process of hitting times $\{T_x\}$, $x \geq 0$, has increments independent of the past, that is, for any $0 < a < b$, $T_b - T_a$ is independent of $B(t)$, $t \leq T_a$, and the distribution of the increment $T_b - T_a$ is the same as that of T_{b-a} and it is given by the density*

$$f_{T_b - T_a}(t) = \frac{b-a}{\sqrt{2\pi}} t^{-\frac{3}{2}} e^{-\frac{(b-a)^2}{2t}}.$$

PROOF. By the strong Markov property $\hat{B}(t) = B(T_a + t) - B(T_a)$ is Brownian motion started at zero, and independent of the past $B(t)$, $t \leq T_a$. $T_b - T_a = \inf\{t \geq 0 : \hat{B}(t) = b - a\}$. Hence $T_b - T_a$ is the same as the first hitting time of $b - a$ by \hat{B} . \square

3.8 Reflection Principle and Joint Distributions

Let $B(t)$ be a Brownian motion started at x , and $\hat{B}(t) = -B(t)$. Then $\hat{B}(t)$ is a Brownian motion started at $-x$. The proof is straightforward by checking the defining properties. This is the simplest form of the reflection principle. Here the Brownian motion is reflected about the horizontal axis.

In greater generality, the process that is obtained by reflection of a Brownian motion about the horizontal line passing through $(T, B(T))$, for a stopping time T , is also a Brownian motion.

Note that for $t \geq T$ the reflected path is, $\hat{B}(t) - B(T) = -(B(t) - B(T))$, giving $\hat{B}(t) = 2B(T) - B(t)$.

Theorem 3.20 (Reflection Principle): *Let T be a stopping time. Define $\hat{B}(t) = B(t)$ for $t \leq T$, and $\hat{B}(t) = 2B(T) - B(t)$ for $t \geq T$. Then \hat{B} is also Brownian motion.*

The proof is beyond the scope of this book, but a heuristic justification is that for $t \geq T$ the process $-(B(t) - B(T))$ is also a Brownian motion by the strong Markov property, so that \hat{B} , constructed from the Brownian motion before the stopping time, and another Brownian motion after the stopping time, is again Brownian motion. For a rigorous proof see Freedman (1971). Using the reflection principle, the joint distribution of the Brownian motion with its maximum can be obtained.

Theorem 3.21: *The joint distribution of $(B(t), M(t))$ has the density*

$$f_{B,M}(x, y) = \sqrt{\frac{2}{\pi}} \frac{(2y - x)}{t^{3/2}} e^{-\frac{(2y-x)^2}{2t}}, \quad \text{for } y \geq 0, x \leq y. \quad (3.16)$$

PROOF. Let, for $y > 0$ and $y > x$, $\hat{B}(t)$ be $B(t)$ reflected at T_y . Then

$$\begin{aligned} P(B(t) \leq x, M(t) \geq y) &= P(T_y \leq t, B(t) \leq x) \quad (\text{since } \{M(t) \geq y\} = \{T_y \leq t\}) \\ &= P(T_y \leq t, \hat{B}(t) \geq 2y - x) \quad (\text{on } \{T_y \leq t\}, \quad \hat{B}(t) = 2y - B(t)) \\ &= P(T_y \leq t, B(t) \geq 2y - x) \quad (\text{since } T_y \text{ is the same for } B \text{ and } \hat{B}) \\ &= P(B(t) \geq 2y - x) \quad (\text{since } y - x > 0, \text{ and } \{B(t) \geq 2y - x\} \subset \{T_y \leq t\}) \\ &= 1 - \Phi\left(\frac{2y - x}{\sqrt{t}}\right). \end{aligned}$$

The density is obtained by differentiation. \square

It is possible to show (see for example Karatzas and Shreve 1988) that $|B(t)|$ and $M(t) - B(t)$ have the same distribution.

Theorem 3.22: *The two processes $|B(t)|$ and $M(t) - B(t)$ are both Markov processes with transition probability density function $p_t(x, y) + p_t(x, -y)$, where $p_t(x, y) = \frac{1}{\sqrt{2\pi t}} e^{-\frac{(y-x)^2}{2t}}$ is the transition probability function of Brownian motion. Consequently, they have the same finite-dimensional distributions.*

The next result gives the joint distribution of $B(t)$, $M(t)$ and $m(t)$; for a proof see Freedman (1971).

Theorem 3.23:

$$P(a < m(t) \leq M(t) < b, \quad \text{and} \quad B(t) \in A) = \int_A k(y) dy, \quad (3.17)$$

where $k(y) = \sum_{n=-\infty}^{\infty} p_t(2n(b-a), y) - p_t(2a, 2n(b-a) + y)$ for $t > 0$ and $a < 0 < b$.

Remark 3.7: Joint distributions given above are used in the pricing of the so-called barrier options (see Chapter 11).

3.9 Zeros of Brownian Motion — Arcsine Law

A time point τ is called a zero of Brownian motion if $B(\tau) = 0$. As an application of the distribution of the maximum we obtain the following information about zeros of Brownian motion. In the following $\{B^x(t)\}$ denotes Brownian motion started at x .

Theorem 3.24: For any $x \neq 0$ the probability that $\{B^x(t)\}$ has at least one zero in the time interval $(0, t)$ is given by

$$\frac{|x|}{\sqrt{2\pi}} \int_0^t u^{-\frac{3}{2}} e^{-\frac{x^2}{2u}} du.$$

PROOF. If $x < 0$, then due to continuity of $B^x(t)$, (draw a picture of this event)

$$P(B^x \text{ has at least one zero between } 0 \text{ and } t) = P(\max_{0 \leq s \leq t} B^x(s) \geq 0).$$

Since $B^x(t) = B(t) + x$, where $B(t)$ is Brownian motion started at zero at time zero,

$$\begin{aligned} P_x(B \text{ has a zero between } 0 \text{ and } t) &= P(\max_{0 \leq s \leq t} B^x(s) \geq 0) \\ &= P_0(\max_{0 \leq s \leq t} B(s) + x \geq 0) = P_0(\max_{0 \leq s \leq t} B(s) \geq -x) \\ &= 2P_0(B(t) \geq -x) = P_0(T_x \leq t) \\ &= \int_0^t f_{T_x}(u) du = \frac{-x}{\sqrt{2\pi}} \int_0^t u^{-\frac{3}{2}} e^{-\frac{x^2}{2u}} du. \end{aligned}$$

For $x > 0$ the proof is similar, and is based on the distribution of the minimum of Brownian motion. \square

Using this result we can establish

Theorem 3.25: The probability that Brownian motion $B(t)$ started at zero has at least one zero in the time interval (a, b) is given by

$$\frac{2}{\pi} \arccos \sqrt{\frac{a}{b}}.$$

PROOF. Denote by $h(x) = P(B \text{ has at least one zero in } (a, b)|B_a = x)$. By the Markov property $P(B \text{ has at least one zero in } (a, b)|B_a = x)$ is the same as $P(B^x \text{ has at least one zero in } (0, b-a))$. By conditioning

$$\begin{aligned} P(B \text{ has at least one zero in } (a, b)) &= \int_{-\infty}^{\infty} P(B \text{ has at least one zero in } (a, b)|B_a = x) P(B_a \in dx) \\ &= \int_{-\infty}^{\infty} h(x) P(B_a \in dx) = \sqrt{\frac{2}{\pi a}} \int_0^{\infty} h(x) e^{-\frac{x^2}{2a}} dx. \end{aligned}$$

Putting in the expression for $h(x)$ from the previous example and performing the necessary calculations we obtain the result. \square

The arcsine law now follows:

Theorem 3.26: *The probability that Brownian motion $\{B(t)\}$ has no zeros in the time interval (a, b) is given by $\frac{2}{\pi} \arcsin \sqrt{\frac{a}{b}}$.*

The next result gives distributions of the last zero before t , and the first zero after t . Let

$$\gamma_t = \sup\{s \leq t : B(s) = 0\} = \text{last zero before } t. \quad (3.18)$$

$$\beta_t = \inf\{s \geq t : B(s) = 0\} = \text{first zero after } t. \quad (3.19)$$

Note that β_t is a stopping time, but γ_t is not.

Theorem 3.27:

$$P(\gamma_t \leq x) = \frac{2}{\pi} \arcsin \sqrt{\frac{x}{t}}. \quad (3.20)$$

$$P(\beta_t \geq y) = \frac{2}{\pi} \arcsin \sqrt{\frac{t}{y}}. \quad (3.21)$$

$$P(\gamma_t \leq x, \beta_t \geq y) = \frac{2}{\pi} \arcsin \sqrt{\frac{x}{y}}. \quad (3.22)$$

PROOF. All of these follow from the previous result. For example,

$$P(\gamma_t \leq x) = P(B \text{ has no zeros in } (x, t)).$$

$$P(\gamma_t \leq x, \beta_t \geq y) = P(B \text{ has no zeros in } (x, y)). \quad \square$$

Since Brownian motion is continuous, and it has no zeros on the interval (γ_t, β_t) it keeps the same sign on this interval, either positive or negative. When Brownian motion is entirely positive or entirely negative on an interval, it is said that it is an *excursion* of Brownian motion. Thus the previous result states that excursions have the arcsine law. In order to picture a Brownian path consider for every realization $B = \{B(t), 0 \leq t \leq 1\}$, the set of its zeros on the interval $[0, 1]$, that is, the random set $L_0 = L_0(B) = \{t : B(t) = 0, 0 \leq t \leq 1\}$.

Theorem 3.28: *The set of zeros of Brownian motion is a random uncountable closed set without isolated points and has Lebesgue measure zero.*

PROOF. According to Example 3.8, the probability that Brownian motion stays below zero on the interval $[0, 1]$ is zero. Therefore it changes sign on this interval. This implies, since Brownian motion is continuous, that it has a zero inside $[0, 1]$. The same reasoning leads to the conclusion that for any positive t , the probability that Brownian motion has the same sign on the interval $[0, t]$ is zero. Therefore it has a zero inside $[0, t]$ for any t , no matter how small it is. This implies that the set of zeros is an infinite set, moreover time $t = 0$ is a limit of zeros from the right.

Observe next that the set of zeros is closed, that is, if $B(\tau_n) = 0$ and $\lim_{n \rightarrow \infty} \tau_n = \tau$, then $B(\tau) = 0$. This is true since $B(t)$ is a continuous function of t .

By using the strong Markov property, it is possible to see that any zero of Brownian motion is a limit of other zeros. If $B(\tau) = 0$, and τ is a stopping time, then by (3.13) $\hat{B}(t) = B(t + \tau) - B(\tau) = B(t + \tau)$ is again Brownian motion started anew at time τ . Therefore time $t = 0$, for the new Brownian motion \hat{B} is a limit from the right of zeros of \hat{B} , but $\hat{B}(t) = B(t + \tau)$, so that τ is a limit from the right of zeros of B . However, not every zero of Brownian motion is a stopping time. For example, for a fixed t , γ_t , the last zero before t is not a stopping time. Nevertheless, using a more intricate argument, one can see that any zero is a limit of other zeros. A sketch is given below. If τ is the first zero after t , then τ is a stopping time. Thus the set of all sample paths such that τ is a limit point of zeros from the right has probability one. The intersection of such sets over all rational t 's is again a set of probability one. Therefore for almost all sample paths the first zero that follows any rational number is a limit of zeros from the right. This implies that any point of L_0 is a limit of points from L_0 (it is a perfect set). A general result from the set theory, which is not hard to prove, states that if an infinite set coincides with the set of its limit points, then it is uncountable.

Although uncountable, L_0 has Lebesgue measure zero. This is seen by writing the Lebesgue measure of L_0 as $|L_0| = \int_0^1 I(B(t) = 0)dt$. It is a non-negative random variable. Taking the expectation and interchanging the integrals by Fubini's theorem

$$\mathbb{E}|L_0| = \mathbb{E} \int_0^1 I(B(t) = 0)dt = \int_0^1 \mathbb{P}(B(t) = 0)dt = 0.$$

This implies $\mathbb{P}(|L_0| = 0) = 1$. □

Theorem 3.29: *Any level set $L_a = \{t : B(t) = a, 0 \leq t \leq 1\}$ has the same properties as L_0 .*

PROOF. Let T_a be the first time with $B(t) = a$. Then by the strong Markov property, $\hat{B}(t) = B_{T_a+t} - B_{T_a} = B_{T_a+t} - a$ is a Brownian motion. The set of zeros of \hat{B} is the level a set of B . □

3.10 Size of Increments of Brownian Motion

Increments over large time intervals satisfy the law of large numbers and the law of the iterated logarithm. For proofs see, for example, Karatzas and Shreve (1988).

Theorem 3.30 (Law of Large Numbers):

$$\lim_{t \rightarrow \infty} \frac{B(t)}{t} = 0 \quad a.s.$$

A more precise result is provided by the law of iterated logarithm.

Theorem 3.31 (Law of the Iterated Logarithm):

$$\limsup_{t \rightarrow \infty} \frac{B(t)}{\sqrt{2t \ln \ln t}} = 1, \quad a.s.$$

$$\liminf_{t \rightarrow \infty} \frac{B(t)}{\sqrt{2t \ln \ln t}} = -1 \quad a.s.$$

In order to obtain the behaviour for small t near zero the process $W(t) = tB(1/t)$ is considered, which is also Brownian motion.

Example 3.9: Let $B(t)$ be Brownian motion. The process $W(t)$ defined as $W(t) = tB(1/t)$, for $t > 0$, and $W(0) = 0$, is also Brownian motion. Indeed, $W(t)$ has continuous paths. Continuity at zero follows from the law of large numbers. It is clearly a Gaussian process and has zero mean. Its covariance is given by

$$\text{Cov}(W(t), W(s)) = E(W(t)W(s)) = tsE(B(1/t)B(1/s)) = ts(1/t) = s, \quad \text{for } s < t.$$

Since $W(t)$ is a Gaussian process with zero mean and the covariance of Brownian motion, it is Brownian motion.

This result allows us to transfer results on the behaviour of paths of Brownian motion for large t to that of small t . For example, we have immediately the law of the iterated logarithm near zero from the same law near infinity.

Graphs of Some Functions of Brownian Motion

Graphs of some functions of Brownian motion are given in order to visualize these processes. In order to obtain these, 1,000 independent normal random variables with mean zero and variance 0.001 were generated. Time is taken to be discrete (as any other variable on a computer) varying from 0 to 1 with steps of 0.001. The first two pictures in Figure 3.2 are realizations of white noise. Pictures of $0.1B(t) + t$ and $B(t) + 0.1t$ (on the second row of Figure 3.2) demonstrate that when noise is small in comparison with drift, the drift dominates, and if drift is small, then the noise dominates in the behaviour of the process. The next two (on the third row) are realizations of the martingale $B^2(t) - t$ which has zero mean. By the recurrence property of Brownian motion, $B(t)$ will always come back to zero. Thus $B^2(t) - t$ will always come back to $-t$ in the long run. The last two (on the fourth row) are realizations of the exponential martingale $e^{B(t)-t/2}$. Although this martingale has mean 1, $\lim_{t \rightarrow \infty} e^{B(t)-t/2} = 0$, which can be seen by the law of large numbers. Therefore a realization of this martingale will approach zero in the long run.




Fig. 3.2: White noise and functions of Brownian motion.

3.11 Brownian Motion in Higher Dimensions

Definition 3.32: Define Brownian motion in dimension two and higher as a random vector $\mathbf{B}(t) = (B^1(t), B^2(t), \dots, B^n(t))$ with all coordinates $B^i(t)$ being independent one-dimensional Brownian motions.

Alternatively, Brownian motion in \mathbb{R}^n can be defined as a process with independent multivariate Gaussian increments. It follows from the definitions, similarly to the one-dimensional case, that Brownian motion in \mathbb{R}^n is a Markov–Gaussian process homogeneous both in space and time. Its transition probability density is given by

$$p_t(\mathbf{x}, \mathbf{y}) = \frac{1}{\sqrt{2\pi t}} e^{-(|\mathbf{x}-\mathbf{y}|^2)/2t}, \quad (3.23)$$

where \mathbf{x}, \mathbf{y} are n -dimensional vectors and $|\mathbf{x}|$ is the length of \mathbf{x} .

Remark 3.8: In dimensions one and two Brownian motion is recurrent, that is, it will come back to a neighbourhood, however small, of any point infinitely often. In dimensions three and higher Brownian motion is transient, it will leave a ball, however large, around any point never to return (Polya (1921)).

3.12 Random Walk

The analogue of the Brownian motion process in discrete time $t = 0, 1, 2, \dots, n, \dots$ is the random walk process. Brownian motion can be constructed as the limit of random walks when step sizes get smaller and smaller. Random walks occur in many applications, including insurance, finance, and biology.

A model of pure chance is served by an ideal coin being tossed with equal probabilities for heads and tails to come up. Introduce a random variable ξ taking values $+1$ (heads) and -1 (tails) with probability $\frac{1}{2}$. If the coin is tossed n times, then a sequence of random variables $\xi_1, \xi_2, \dots, \xi_n$ describes this experiment. All ξ_i have exactly the same distribution as ξ_1 , moreover they are all independent. The process S_n is a random walk, defined by $S_0 = 0$ and

$$S_n = \xi_1 + \xi_2 + \dots + \xi_n. \quad (3.24)$$

S_n gives the amount of money after n plays when betting \$1 on the outcomes of a coin when \$1 is won if heads come up, but lost otherwise.

Since $E(\xi_i) = 0$, and $Var(\xi_i) = E(\xi^2) = 1$, the mean and the variance of the random walk are given by

$$E(S_n) = E(\xi_1 + \xi_2 + \cdots + \xi_n) = E(\xi_1) + E(\xi_2) + \cdots + E(\xi_n) = 0,$$

$$Var(S_n) = Var(\xi_1) + Var(\xi_2) + \cdots + Var(\xi_n) = nVar(\xi_1) = n,$$

as the variance of a sum of independent variables equals the sum of variances.

More generally, a random walk is the process

$$S_n = S_0 + \sum_{i=1}^n \xi_i, \quad (3.25)$$

where ξ_i 's are independent and identically distributed random variables (i.i.d.). In particular, this model contains gambling on the outcomes of a biased coin $P(\xi_i = 1) = p$, $P(\xi_i = -1) = q = 1 - p$.

Martingales in Random Walks

Some interesting questions about random walks, such as ruin probabilities and the like, can be answered with the help of martingales.

Theorem 3.33: *The following are martingales:*

1. $S_n - \mu n$, where $\mu = E(\xi_1)$. In particular, if the random walk is unbiased ($\mu = 0$), then it is itself a martingale.
2. $(S_n - \mu n)^2 - \sigma^2 n$, where $\sigma^2 = E(\xi_1 - \mu)^2 = Var(\xi_1)$.
3. For any u , $e^{uS_n - nh(u)}$, where $h(u) = \ln E(e^{u\xi_1})$. In particular, in the case $P(\xi_1 = 1) = p$, $P(\xi_1 = -1) = q = 1 - p$, $(\frac{q}{p})^{S_n}$ is a martingale.

PROOF.

1. Since, by the triangle inequality,

$$\begin{aligned} E|S_n - n\mu| &= E|S_0 + \sum_{i=1}^n \xi_i - n\mu| \leq E|S_0| + \sum_{i=1}^n E|\xi_i| + n|\mu| \\ &= E|S_0| + n(E|\xi_1| + |\mu|), \end{aligned}$$

$S_n - n\mu$ is integrable provided $E|\xi_1| < \infty$ and $E|S_0| < \infty$. In order to establish the martingale property consider for any n

$$E(S_{n+1}|S_n) = S_n + E(\xi_{n+1}|S_n).$$

Since ξ_{n+1} is independent of the past, and S_n is determined by the first n variables, ξ_{n+1} is independent of S_n . Therefore $E(\xi_{n+1}|S_n) = E(\xi_{n+1})$. It now follows that

$$E(S_{n+1}|S_n) = S_n + E(\xi_{n+1}|S_n) = S_n + \mu,$$

and, subtracting $(n + 1)\mu$ from both sides of the equation, the martingale property is obtained,

$$\mathbb{E}(S_{n+1} - (n + 1)\mu | S_n) = S_n - n\mu.$$

2. This is left as an exercise.

3. Put $M_n = e^{uS_n - nh(u)}$. Since $M_n \geq 0$, $\mathbb{E}|M_n| = \mathbb{E}(M_n)$, which is given by

$$\begin{aligned} \mathbb{E}(M_n) &= \mathbb{E}e^{uS_n - nh(u)} = e^{-nh(u)} \mathbb{E}e^{uS_n} = e^{-nh(u)} \mathbb{E}e^{u(S_0 + \sum_{i=1}^n \xi_i)} \\ &= e^{uS_0} e^{-nh(u)} \mathbb{E} \prod_{i=1}^n e^{u\xi_i} = e^{uS_0} e^{-nh(u)} \prod_{i=1}^n \mathbb{E}(e^{u\xi_i}) \quad \text{by independence} \\ &= e^{uS_0} e^{-nh(u)} \prod_{i=1}^n e^{h(u)} = e^{uS_0} < \infty. \end{aligned}$$

The martingale property is shown by using the fact that

$$S_{n+1} = S_n + \xi_{n+1}, \tag{3.26}$$

with ξ_{n+1} independent of S_n and of all previous ξ_i 's $i \leq n$, or independent of \mathcal{F}_n . Using the properties of conditional expectation we have

$$\begin{aligned} \mathbb{E}(e^{uS_{n+1}} | \mathcal{F}_n) &= \mathbb{E}(e^{uS_n + u\xi_{n+1}} | \mathcal{F}_n) \\ &= e^{uS_n} \mathbb{E}(e^{u\xi_{n+1}} | \mathcal{F}_n) = e^{uS_n} \mathbb{E}(e^{u\xi_{n+1}}) \\ &= e^{uS_n + h(u)}. \end{aligned}$$

Multiplying both sides of the above equation by $e^{-(n+1)h(u)}$, the martingale property is obtained, $\mathbb{E}(M_{n+1} | \mathcal{F}_n) = M_n$.

In the special case when $P(\xi_i = 1) = p$, $P(\xi_i = -1) = q = 1 - p$ choosing $u = \ln(q/p)$ in the previous martingale, we have $e^{u\xi_1} = (q/p)^{\xi_1}$ and $\mathbb{E}(e^{u\xi_1}) = 1$. Thus $h(u) = \ln \mathbb{E}(e^{u\xi_1}) = 0$, and $e^{uS_n - nh(u)} = (q/p)^{S_n}$. Alternatively in this case, the martingale property of $(q/p)^{S_n}$ is easy to verify directly. \square

3.13 Stochastic Integral in Discrete Time

Let S_n be an unbiased random walk representing the capital of a player when betting on a fair coin. Let H_n be the amount of money (the number of betting units) a gambler will bet at time n . This can be based on the outcomes of the game at times $1, \dots, n-1$, but not on the outcome at time n . This is an example of a predictable process. The concept of predictable processes plays a most important role in stochastic calculus. The process $\{H_n\}$ is called predictable if H_n can be predicted with certainty from the information available at time $n-1$. Rigorously, let \mathcal{F}_{n-1} be the σ -field generated by S_0, S_1, \dots, S_{n-1} .

Definition 3.34: $\{H_n\}$ is called predictable if for all $n \geq 1$, H_n is \mathcal{F}_{n-1} measurable.

If a betting strategy $\{H_n\}_{n=0}^t$ is used in the game of coin tossing, then the gain at time t is given by

$$(H \cdot S)_t = \sum_{n=1}^t H_n(S_n - S_{n-1}), \quad (3.27)$$

since $S_n - S_{n-1} = +1$ or -1 when the n -th toss results in a win or loss respectively. More generally, $S_n - S_{n-1} = \xi_n$ represents the amount of money lost or won on one betting unit on the n -th bet. If H_n units are placed, then the amount of money lost or won on the n -th bet is $H_n(S_n - S_{n-1})$. The gain at time t is obtained by adding up monies lost and won on all the bets.

Definition 3.35: The stochastic integral in discrete time of a predictable process H with respect to the process S is defined by

$$(H \cdot S)_t := H_0 S_0 + \sum_{n=1}^t H_n(S_n - S_{n-1}). \quad (3.28)$$

The stochastic integral gives the gain in a game of chance when betting on S and the betting strategy H is used. For a martingale the stochastic integral (3.28) is also called a martingale transform. The next result states that a betting system used on a martingale will result again in a martingale.

Theorem 3.36: If M_n is a martingale, H_n is predictable and the random variables $(H \cdot M)_t$ are integrable, then $(H \cdot M)_t$ is a martingale.

PROOF.

$$\begin{aligned} E[(H \cdot M)_{t+1} | \mathcal{F}_t] &= E[(H \cdot M)_t | \mathcal{F}_t] + E[H_{t+1}(M_{t+1} - M(t)) | \mathcal{F}_t] \\ &= (H \cdot M)_t + H_{t+1}E[(M_{t+1} - M(t)) | \mathcal{F}_t] = (H \cdot M)_t. \square \end{aligned}$$

As a corollary we obtain

Theorem 3.37: If M_n is a martingale and H_n is predictable and bounded, then $(H \cdot M)_t$ is a martingale.

PROOF. The assumption of bounded H_n implies that

$$\begin{aligned} E|(H \cdot M)_t| &= E|\sum_{n=1}^t H_n(M_n - M_{n-1})| \\ &\leq \sum_{n=1}^t E|H_n(M_n - M_{n-1})| \leq 2C \sum_{n=1}^t E|M_n| < \infty. \end{aligned}$$

So that $(H \cdot M)_t$ is integrable and the condition of Theorem 3.36 is fulfilled. \square

Stopped Martingales

Let (M_n, \mathcal{F}_n) be a martingale and τ be a stopping time. Recall the definition

Definition 3.38: A random time is a stopping time if for any n , $\{\tau > n\} \in \mathcal{F}_n$.

Take $H_n = 1$ if $n \leq \tau$, and $H_n = 0$ if $n > \tau$, in other words, $H_n = I(\tau \geq n)$. Then H_n is predictable, because $\{\tau \geq n\} = \{\tau > n - 1\} \in \mathcal{F}_{n-1}$. The stochastic integral gives the martingale stopped at τ ,

$$\begin{aligned}(H \cdot M)_n &= H_0 M_0 + H_1(M_1 - M_0) + \cdots + H_n(M_n - M_{n-1}) \\ &= M_{\tau \wedge n} = M_\tau I(\tau \leq n) + M_n I(\tau > n).\end{aligned}$$

Since $H_n = I(\tau \geq n)$ is bounded by 1, Theorem 3.37 implies that the process $(H \cdot M)_n = M_{\tau \wedge n}$ is a martingale. Thus we have shown

Theorem 3.39: A martingale stopped at a stopping time τ , $M_{\tau \wedge n}$ is a martingale. In particular,

$$EM_{\tau \wedge n} = EM_0. \quad (3.29)$$

Note here that Theorem 3.39 holds also in continuous time, see Theorem 7.14. It is a basic stopping result, which is harder to prove.

Example 3.10: (Doubling bets strategy)

Consider the doubling strategy when betting on heads in tosses of a fair coin. Bet $H_1 = 1$. If heads comes up then stop. The profit is $G_1 = 1$. If the outcome is tails, then bet $H_2 = 2$ on the second toss. If the second toss comes up heads, then stop. The profit is $G_2 = 4 - 3 = 1$. If the game continues for n steps, (meaning that the $n - 1$ tosses did not result in a win), then bet $H_n = 2^{n-1}$ on the n -th toss. If the n -th toss comes up heads then stop. The profit is $G_n = 2 \times 2^{n-1} - (1 + 2 + \dots + 2^{n-1}) = 2^n - (2^n - 1) = 1$. The probability that the game will stop at a finite number of steps is one minus the probability that heads never come up. Probability of only tails on the first n tosses is, by independence, 2^{-n} . The probability that heads never comes up is the limit $\lim_{n \rightarrow \infty} 2^{-n} = 0$, thus the game will stop for sure. For any non-random time T the gain process G_t , $t \leq T$ is a martingale with zero mean. The doubling strategy does not contradict the result above, because the strategy uses an unbounded stopping time, the first time one dollar is won.

Further information on discrete time martingales and on their stopping is given in Chapter 7.

3.14 Poisson Process

If the Brownian motion process is a basic model for cumulative small noise present continuously, the Poisson process is a basic model for cumulative noise that occurs as a shock.

Let $\lambda > 0$. A random variable X has a Poisson distribution with parameter λ , denoted $Pn(\lambda)$, if it takes non-negative integer values $k \geq 0$ with probabilities

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}, \quad k = 0, 1, 2, \dots \quad (3.30)$$

The moment generating function of this distribution is given by

$$E(e^{uX}) = e^{\lambda(e^u - 1)}. \quad (3.31)$$

Defining Properties of Poisson Process

A Poisson process $N(t)$ is a stochastic process with the following properties

1. (Independence of increments) $N(t) - N(s)$ is independent of the past, that is, of \mathcal{F}_s , the σ -field generated by $N(u), u \leq s$.
2. (Poisson increments) $N(t) - N(s), t > s$, has a Poisson distribution with parameter $\lambda(t-s)$. If $N(0) = 0$, then $N(t)$ has the $Pn(\lambda t)$ distribution.
3. (Step function paths) The paths $N(t), t \geq 0$, are increasing functions of t changing only by jumps of size 1.

Remark 3.9: A definition of a Poisson process in a more general model (that contains extra information) is given by a pair $\{N(t), \mathcal{F}_t\}, t \geq 0$, where \mathcal{F}_t is an increasing sequence of σ -fields (a filtration), $N(t)$ is an adapted process, i.e. $N(t)$ is \mathcal{F}_t measurable, such that properties 1–3 above hold.

Consider a model for occurrence of independent events. Define the rate λ as the average number of events per unit of time. Let $N(t)$ be the number of events that occur up to time t , i.e. in the time interval $(0, t]$. Then $N(t) - N(s)$ gives the number of events that occur in the time interval $(s, t]$.

A Poisson process $N(t)$ can be constructed as follows. Let τ_1, τ_2, \dots , be independent random variables with the exponential $\exp(\lambda)$ distribution, that is, $P(\tau_1 > t) = e^{-\lambda t}$. τ 's represent the times between the occurrence of successive events. Let $T_n = \sum_{i=1}^n \tau_i$ be the time of the n -th event. Then

$$N(t) = \sup\{n : T_n \leq t\}$$

counts the number of events up to time t . It is not hard to verify the defining properties of the Poisson process for this construction. $N(t)$ has the Poisson distribution with parameter λt . Consequently,

$$\begin{aligned} \Pr(N(t) = k) &= e^{-\lambda t} \frac{(\lambda t)^k}{k!}, \quad k = 0, 1, 2, \dots, \\ EN(t) &= \lambda t, \quad \text{and} \quad \text{Var}(N(t)) = \lambda t. \end{aligned}$$

Variation and Quadratic Variation of the Poisson Process

Let $0 = t_0^n < t_1^n < \dots < t_n^n = t$ be a partition of $[0, t]$. Then it is easy to see that variation of a Poisson path is

$$V_N(t) = \lim \sum_{i=1}^n |N(t_i^n) - N(t_{i-1}^n)| = N(t) - N(0) = N(t), \quad (3.32)$$

where the limit is taken when $\delta_n = \max_i(t_i^n - t_{i-1}^n) \rightarrow 0$ and $n \rightarrow \infty$. Recall that variation of a pure jump function is the sum of absolute values of the jumps, see Example 1.6. Since the Poisson process has only positive jumps of size one, (3.32) follows.

In order to calculate its quadratic variation, observe that $N(t_i^n) - N(t_{i-1}^n)$ takes only two values 0 and 1 for small $t_i^n - t_{i-1}^n$, hence it is the same as its square, $N(t_i^n) - N(t_{i-1}^n) = (N(t_i^n) - N(t_{i-1}^n))^2$. Thus the quadratic variation of N is the same as its variation

$$[N, N](t) = \lim \sum_{i=1}^n (N(t_i^n) - N(t_{i-1}^n))^2 = N(t) - N_0 = N(t).$$

Thus for a Poisson process both the variation and quadratic variation are positive and finite.

Poisson Process Martingales

The process $N(t)$ is increasing, hence it can not be a martingale. However, the compensated process $N(t) - \lambda t$ is a martingale. This martingale is analogous to the Brownian motion.

Theorem 3.40: *The following are martingales.*

1. $N(t) - \lambda t$.
2. $(N(t) - \lambda t)^2 - \lambda t$.
3. $e^{\ln(1-u)N(t)+u\lambda t}$, for any $0 < u < 1$.

PROOF. The martingale property follows from independence of increments, the Poisson distribution of increments, and the expressions for the mean and the variance of Poisson distribution. We show the martingale property for the exponential martingale.

$$\begin{aligned}
 & \mathbb{E}(e^{\ln(1-u)N(t+s)} | \mathcal{F}_t) \\
 &= \mathbb{E}(e^{\ln(1-u)N(t)+\ln(1-u)(N(t+s)-N(t))} | \mathcal{F}_t) \\
 &= e^{\ln(1-u)N(t)} \mathbb{E}(e^{\ln(1-u)(N(t+s)-N(t))} | \mathcal{F}_t) \text{ (since } N(t) \text{ is } \mathcal{F}_t\text{-measurable)} \\
 &= e^{\ln(1-u)N(t)} \mathbb{E}(e^{\ln(1-u)(N(t+s)-N(t))}) \text{ (increment is independent of } \mathcal{F}_t) \\
 &= e^{\ln(1-u)N(t)} e^{-u\lambda s} \quad \text{by (3.31), since } N(t+s) - N(t) \text{ is Poisson}(\lambda s).
 \end{aligned}$$

Multiplying both sides by $e^{u\lambda(t+s)}$, the martingale property follows. \square

Using the exponential martingale, it can be shown that the Poisson process has the strong Markov property.

Notes: Materials in this chapter can be found in Chung (1982), Durrett (1991), and Grimmet and Stirzaker (1987).

3.15 Exercises

Exercise 3.1: Derive the moment generating function of the multivariate normal distribution $N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$.

Exercise 3.2: Show that for a non-negative random variable X , $\mathbb{E}X = \int_0^\infty \mathbb{P}(X \geq x)dx$. Hint: use $\mathbb{E}X = \int_0^\infty x dF(x) = \int_0^\infty \int_0^x dt dF(x)$ and change the order of integration.

Exercise 3.3: Show that if $X \geq 0$, $\mathbb{E}X \leq \sum_{n=0}^\infty \mathbb{P}(X > n)$.

Exercise 3.4: Let $B(t)$ be a Brownian motion. Show that the following processes are Brownian motions on $[0, T]$.

1. $X(t) = -B(t)$.
2. $X(t) = B(T-t) - B(T)$, where $T < +\infty$.
3. $X(t) = cB(t/c^2)$, where $T \leq +\infty$.
4. $X(t) = tB(1/t)$, $t > 0$, and $X(0) = 0$.

Hint: Check the defining properties. Alternatively, show that the process is a Gaussian process with correlation function $\min(s, t)$. Alternatively, show that the process is a continuous martingale with quadratic variation t (this is Levy's characterization, and will be proven later).

Exercise 3.5: Let $B(t)$ and $W(t)$ be two independent Brownian motions. Show that $X(t) = (B(t) + W(t))/\sqrt{2}$ is also a Brownian motion. Find correlation between $B(t)$ and $X(t)$.

Exercise 3.6: Let $\mathbf{B}(t)$ be an n -dimensional Brownian motion, and \mathbf{x} be a non-random vector in \mathbb{R}^n with length 1, $|\mathbf{x}|^2 = 1$. Show that $W(t) = \mathbf{x} \cdot \mathbf{B}(t)$ is a (one-dimensional) Brownian motion.

Exercise 3.7: Let $B(t)$ be a Brownian motion and $0 \leq t_1, \dots \leq t_n$. Give a matrix A , such that $(B(t_1), B(t_2), \dots, B(t_n))^T = A(Z_1, \dots, Z_n)^T$, where Z_i 's are standard normal variables. Hence give the covariance matrix of $(B(t_1), B(t_2), \dots, B(t_n))$. Here T stands for transpose, and the vectors are column vectors.

Exercise 3.8: Let $B(t)$ be a Brownian motion and $0 \leq s < t$. Show that the conditional distribution of $B(s)$ given $B(t) = b$ is normal and give its mean and variance.

Exercise 3.9: Show that the random variables $M(t)$ and $|B(t)|$ have the same distribution.

Exercise 3.10: Show that the moments of order r of the hitting time T_x are finite $E(T_x^r) < \infty$ if, and only if, $r < 1/2$.

Exercise 3.11: Derive the distribution of the maximum $M(t)$ from the joint distribution of $(B(t), M(t))$.

Exercise 3.12: By considering $-B(t)$, derive the joint distribution of $B(t)$ and $m(t) = \min_{s \leq t} B(s)$.

Exercise 3.13: Show that the random variables $M(t)$, $|B(t)|$, and $M(t) - B(t)$ have the same distributions, cf. Exercise 3.9.

Exercise 3.14: The first zero of Brownian motion started at zero is 0. What is the second zero?

Exercise 3.15: Let T be the last time before time 1 a Brownian motion visits 0. Explain why $X(t) = B(t+T) - B(T) = B(t+T)$ is not a Brownian motion.

Exercise 3.16: Formulate the law of large numbers and the law of the iterated logarithm for Brownian motion near zero.

Exercise 3.17: Let $B(t)$ be Brownian motion. Show that $e^{-\alpha t} B(e^{2\alpha t})$ is a Gaussian process. Find its mean and covariance functions.

Exercise 3.18: Let $X(t)$ be a Gaussian process with zero mean and covariance $\gamma(s, t) = e^{-\alpha|t-s|}$. Show that X has a version with continuous paths.

Exercise 3.19: Show that in a normal random walk $S_n = S_0 + \sum_{i=1}^n \xi_i$, when ξ_i 's are standard normal random variables, $e^{uS_n - nu^2/2}$ is a martingale.

Exercise 3.20: Let $S_n = S_0 + \sum_{i=1}^n \xi_i$ be a random walk, with $P(\xi_1 = 1) = p$, $P(\xi_1 = -1) = 1 - p$. Show that for any λ , $e^{\gamma S_n - \lambda n}$ is a martingale for the appropriate value of γ .

Exercise 3.21: The process X_t is defined for discrete times $t = 1, 2, \dots$. It can take only three values 1, 2, and 3. Its behaviour is defined by the rule: from state 1 it goes to 2, from 2 it goes to 3, and from 3 it goes back to 1. X_1 takes values 1, 2, and 3 with equal probabilities. Show that this process is Markov. Show also that

$$P(X_3 = 3 | X_2 = 1 \text{ or } 2, X_1 = 3) \neq P(X_3 = 3 | X_2 = 1 \text{ or } 2).$$

This demonstrates that to apply the Markov property we must know the present state of the process exactly, it is not enough to know that it can take one of the two (or more) possible values.

Exercise 3.22: A discrete time process $X(t)$, $t = 0, 1, 2, \dots$, is said to be autoregressive of order p (AR(p)) if there exists $a_1, \dots, a_p \in \mathbb{R}$, and a white noise $Z(t)$ ($E(Z(t)) = 0$, $E(Z^2(t)) = \sigma^2$, and for $s > 0$, $E(Z(t)Z(t+s)) = 0$) such that

$$X(t) = \sum_{s=1}^p a_s X(t-s) + Z(t).$$

1. Show that $X(t)$ is Markovian if, and only if, $p = 1$.
2. Show that if $X(t)$ is AR(2), then $\mathbf{Y}(t) = (X(t), X(t+1))$ is Markovian.
3. Suppose that $Z(t)$ is a Gaussian process. Write the transition probability function of an AR(1) process $X(t)$.

Exercise 3.23: The distribution of a random variable τ has the lack of memory property if $P(\tau > a + b | \tau > a) = P(\tau > b)$. Verify the lack of memory property for the exponential $\exp(\lambda)$ distribution. Show that if τ has the lack of memory property and a density, then it has an exponential distribution.

Chapter 4

Brownian Motion Calculus

In this chapter stochastic integrals with respect to Brownian motion are introduced and their properties are given. They are also called Itô integrals, and the corresponding calculus Itô calculus.

4.1 Definition of Itô Integral

Our goal is to define the stochastic integral $\int_0^T X(t)dB(t)$, also denoted $\int X dB$ or $X \cdot B$. This integral should have the property that if $X(t) = 1$, then $\int_0^T dB(t) = B(T) - B(0)$. Similarly, if $X(t)$ is a constant c , then the integral should be $c(B(T) - B(0))$. In this way we can integrate constant processes with respect to B . The integral over $(0, T]$ should be the sum of integrals over two subintervals $(0, a]$ and $(a, T]$. Thus if $X(t)$ takes two values, c_1 on $(0, a]$, and c_2 on $(a, T]$, then the integral of X with respect to B is easily defined. In this way the integral is defined for *simple* processes, that is, processes which are constant on finitely many intervals. According to the limiting procedure the integral is then defined for more general processes.

Itô Integral of Simple Processes

Consider first integrals of a *non-random* simple process $X(t)$, which is a function of t and does not depend on $B(t)$. By definition a simple non-random process $X(t)$ is a process for which there exist times $0 = t_0 < t_1 < \dots < t_n = T$ and constants c_0, c_1, \dots, c_{n-1} , such that

$$X(t) = c_0 I_0(t) + \sum_{i=0}^{n-1} c_i I_{(t_i, t_{i+1}]}(t). \quad (4.1)$$

The Itô integral $\int_0^T X(t)dB(t)$ is defined as a sum

$$\int_0^T X(t)dB(t) = \sum_{i=0}^{n-1} c_i (B(t_{i+1}) - B(t_i)). \quad (4.2)$$

It is easy to see by using the independence property of Brownian increments that the integral, which is the sum in (4.2), is a Gaussian random variable with mean zero and variance

$$\begin{aligned} Var\left(\int_0^T X(t)dB(t)\right) &= Var\left(\sum_{i=0}^{n-1} c_i (B(t_{i+1}) - B(t_i))\right) \\ &= \sum_{i=0}^{n-1} Var(c_i(B(t_{i+1}) - B(t_i))) = \sum_{i=0}^{n-1} c_i^2(t_{i+1} - t_i). \end{aligned}$$

Example 4.1: Let $X(t) = -1$ for $0 \leq t \leq 1$, $X(t) = 1$ for $1 < t \leq 2$, and $X(t) = 2$ for $2 < t \leq 3$. Then (note that $t_i = 0, 1, 2, 3$, $c_i = X(t_{i+1})$, $c_0 = -1$, $c_1 = 1$, $c_2 = 2$)

$$\begin{aligned} \int_0^3 X(t)dB(t) &= c_0(B(1) - B(0)) + c_2(B(2) - B(1)) + c_3(B(3) - B(2)) \\ &= -B(1) + (B(2) - B(1)) + 2(B(3) - B(2)) = 2B(3) - B(2) - 2B(1). \end{aligned}$$

Its distribution is $N(0, 6)$, either directly as a sum of independent $N(0, 1) + N(0, 1) + N(0, 4)$, or by using the result above.

By taking limits of simple non-random processes, more general but still only non-random processes can be integrated with respect to Brownian motion. We shall see later (Chapter 6) that, as in Example 4.1, the Itô integral of a deterministic process $X(t)$ is a normal random variable with mean zero and variance $\int_0^T X^2(t)dt$, $\int_0^T X(t)dB(t) \sim N(0, \int_0^T X^2(t)dt)$.

In order to integrate random processes, it is important to allow for constants c_i in (4.1) to be random. If c_i 's are replaced by random variable ξ_i 's, then, in order to have convenient properties of the integral, the random variable ξ_i 's are allowed to depend on the values of $B(t)$ for $t \leq t_i$, but not on future values of $B(t)$ for $t > t_i$. If \mathcal{F}_t is the σ -field generated by Brownian motion up to time t , then ξ_i is \mathcal{F}_{t_i} -measurable. The approach of defining the integral by approximation can be carried out for the class of *adapted* processes $X(t)$, $0 \leq t \leq T$.

Definition 4.1: A process X is called adapted to the filtration $\mathbb{F} = (\mathcal{F}_t)$, if for all t , $X(t)$ is \mathcal{F}_t -measurable.

Remark 4.1: In order that the integral has desirable properties, in particular that the expectation and the integral can be interchanged (by Fubini's theorem), the requirement that X is adapted is too weak, and a stronger condition, that of a *progressive* (progressively measurable) process is needed. X is progressive if it is a measurable function in the pair of variables (t, ω) , i.e. $\mathcal{B}([0, t]) \times \mathcal{F}_t$ measurable as a map from $[0, t] \times \Omega$ into \mathbb{R} . It can be seen that every adapted right-continuous with left limits or left-continuous with right limits (regular, *càdlàg*) process is progressive. Since it is easier to understand what is meant by a regular adapted process, we use “regular adapted” terminology without further reference to progressive or measurable in (t, ω) processes.

Definition 4.2: A process $X = \{X(t), 0 \leq t \leq T\}$ is called a simple adapted process if there exist times $0 = t_0 < t_1 < \dots < t_n = T$ and random variables $\xi_0, \xi_1, \dots, \xi_{n-1}$, such that ξ_0 is a constant, ξ_i is \mathcal{F}_{t_i} -measurable (this depends on the values of $B(t)$ for $t \leq t_i$, but not on values of $B(t)$ for $t > t_i$), and $E(\xi_i^2) < \infty$, $i = 0, \dots, n-1$; such that

$$X(t) = \xi_0 I_0(t) + \sum_{i=0}^{n-1} \xi_i I_{(t_i, t_{i+1})}(t). \quad (4.3)$$

For simple adapted processes the Itô integral $\int_0^T X dB$ is defined as a sum

$$\int_0^T X(t) dB(t) = \sum_{i=0}^{n-1} \xi_i (B(t_{i+1}) - B(t_i)). \quad (4.4)$$

Note that when ξ_i 's are random, the integral need not have a normal distribution, as in the case of non-random c_i 's.

Remark 4.2: Simple adapted processes are defined as left-continuous step functions. One can also take right-continuous functions. However, when the stochastic integral is defined with respect to general martingales, other than the Brownian motion, only left-continuous functions are taken.

Properties of the Itô Integral of Simple Adapted Processes

Here we establish the main properties of the Itô integral of simple processes. These properties carry over to the Itô integral of general processes.

1. Linearity. If $X(t)$ and $Y(t)$ are simple processes and α and β are some constants then

$$\int_0^T (\alpha X(t) + \beta Y(t)) dB(t) = \alpha \int_0^T X(t) dB(t) + \beta \int_0^T Y(t) dB(t).$$

2. For the indicator function of an interval $I_{(a,b]}(t)$ ($I_{(a,b]}(t) = 1$ when $t \in (a, b]$, and zero otherwise)

$$\begin{aligned} \int_0^T I_{(a,b]}(t) dB(t) &= B(b) - B(a), \quad \int_0^T I_{(a,b]}(t) X(t) dB(t) \\ &= \int_a^b X(t) dB(t). \end{aligned}$$

3. Zero mean property. $E \int_0^T X(t) dB(t) = 0$.

4. Isometry property.

$$E \left(\int_0^T X(t) dB(t) \right)^2 = \int_0^T E(X^2(t)) dt. \quad (4.5)$$

PROOF. Properties 1 and 2 are verified directly from the definition. Proof of linearity of the integral follows from the fact that a linear combination of simple processes is again a simple process, and so is $I_{(a,b]}(t)X(t)$.

Since ξ_i 's are square integrable, then by the Cauchy-Schwarz inequality

$$E|\xi_i(B(t_{i+1}) - B(t_i))| \leq \sqrt{E(\xi_i^2)E(B(t_{i+1}) - B(t_i))^2} < \infty,$$

which implies that

$$E \left| \sum_{i=0}^{n-1} \xi_i (B(t_{i+1}) - B(t_i)) \right| \leq \sum_{i=0}^{n-1} E|\xi_i(B(t_{i+1}) - B(t_i))| < \infty, \quad (4.6)$$

and the stochastic integral has expectation. By the martingale property of Brownian motion, using that ξ_i 's are \mathcal{F}_{t_i} -measurable

$$E(\xi_i(B(t_{i+1}) - B(t_i))|\mathcal{F}_{t_i}) = \xi_i E((B(t_{i+1}) - B(t_i))|\mathcal{F}_{t_i}) = 0, \quad (4.7)$$

and it follows that $E(\xi_i(B(t_{i+1}) - B(t_i))) = 0$, which implies property 3.

In order to prove property 4, write the square as the double sum

$$\begin{aligned} E \left(\sum_{i=0}^{n-1} \xi_i (B(t_{i+1}) - B(t_i)) \right)^2 &= \sum_{i=0}^{n-1} E(\xi_i^2 (B(t_{i+1}) - B(t_i))^2) \\ &\quad + 2 \sum_{i < j} E(\xi_i \xi_j (B(t_{i+1}) - B(t_i))(B(t_{j+1}) - B(t_j))). \end{aligned} \quad (4.8)$$

Using the martingale property of Brownian motion,

$$\begin{aligned} \sum_{i=0}^{n-1} E(\xi_i^2 (B(t_{i+1}) - B(t_i))^2) &= \sum_{i=0}^{n-1} E E(\xi_i^2 (B(t_{i+1}) - B(t_i))^2 | \mathcal{F}_{t_i}) \\ &= \sum_{i=0}^{n-1} E(\xi_i^2 E((B(t_{i+1}) - B(t_i))^2 | \mathcal{F}_{t_i})) = \sum_{i=0}^{n-1} E(\xi_i^2)(t_{i+1} - t_i). \end{aligned}$$

The last sum is exactly $\int_0^T \mathbf{E}(X^2(t))dt$, since $X(t) = \xi_i$ on (t_i, t_{i+1}) . By conditioning, in a similar way, we obtain for $i < j$,

$$\mathbf{E}(\xi_i \xi_j (B(t_{i+1}) - B(t_i))(B(t_{j+1}) - B(t_j))) = 0,$$

so that the sum $\sum_{i < j}$ in (4.8) vanishes, and property 4 is proved. \square

Itô Integral of Adapted Processes

Let $X^n(t)$ be a sequence of simple processes convergent in probability to the process $X(t)$. Then, under some conditions, the sequence of their integrals $\int_0^T X^n(t)dB(t)$ also converges in probability to a limit J . The random variable J is taken to be the integral $\int_0^T X(t)dB(t)$.

Example 4.2: We find $\int_0^T B(t)dB(t)$.

Let $0 = t_0^n < t_1^n < t_2^n < \dots < t_n^n = T$ be a partition of $[0, T]$, and let

$$X^n(t) = \sum_{i=0}^{n-1} B(t_i^n) I_{(t_i^n, t_{i+1}^n]}(t).$$

Then for any n , $X^n(t)$ is a simple adapted process. (Here $\xi_i^n = B(t_i^n)$). By the continuity of $B(t)$, $\lim_{n \rightarrow \infty} X^n(t) = B(t)$ almost surely as $\max_i(t_{i+1}^n - t_i^n) \rightarrow 0$. The Itô integral of the simple function $X^n(t)$ is given by

$$\int_0^T X^n(t)dB(t) = \sum_{i=0}^{n-1} B(t_i^n)(B(t_{i+1}^n) - B(t_i^n)).$$

We show that this sequence of integrals converges in probability to $J = \frac{1}{2}B^2(T) - \frac{1}{2}T$. Adding and subtracting $B^2(t_{i+1}^n)$, we obtain

$$B(t_i^n)(B(t_{i+1}^n) - B(t_i^n)) = \frac{1}{2}(B^2(t_{i+1}^n) - B^2(t_i^n) - (B(t_{i+1}^n) - B(t_i^n))^2),$$

and

$$\begin{aligned} \int_0^T X^n(t)dB(t) &= \frac{1}{2} \sum_{i=0}^{n-1} (B^2(t_{i+1}^n) - B^2(t_i^n)) - \frac{1}{2} \sum_{i=0}^{n-1} (B(t_{i+1}^n) - B(t_i^n))^2 \\ &= \frac{1}{2}B^2(T) - \frac{1}{2}B^2(0) - \frac{1}{2} \sum_{i=0}^{n-1} (B(t_{i+1}^n) - B(t_i^n))^2, \end{aligned}$$

since the first sum is a telescopic one. According to the definition of the quadratic variation of Brownian motion the second sum converges in probability to T . Therefore $\int_0^T X^n(t)dB(t)$ converges in probability to the limit J

$$\int_0^T B(t)dB(t) = J = \lim_{n \rightarrow \infty} \int_0^T X^n(t)dB(t) = \frac{1}{2}B^2(T) - \frac{1}{2}T. \quad (4.9)$$

Remark 4.3:

- If $X(t)$ is a differentiable function (more generally, a function of finite variation), then the stochastic integral $\int_0^T X(t)dB(t)$ can be defined by formally using the integration by parts:

$$\int_0^T X(t)dB(t) = X(T)B(T) - X(0)B(0) - \int_0^T B(t)dX(t),$$

(see Paley (1933).)

However, this approach fails when $X(t)$ depends on $B(t)$.

- Brownian motion has no derivative, but it has a generalized derivative as a Schwartz distribution. It is defined by the following relation. For a smooth function g with a compact support (zero outside a finite interval)

$$\int g(t)B'(t)dt := - \int B(t)g'(t)dt.$$

However, this approach fails when $g(t)$ depends on $B(t)$.

- For simple processes the Itô integral is defined for each ω , path by path, but in general this is not possible. For example, $\int_0^1 B(\omega, t)dB(\omega, t)$ is not defined, whereas $(\int_0^1 B(t)dB(t))(\omega) = J(\omega)$ is defined as a limit in probability of integrals (sums) of simple processes.

Theorem 4.3: Let $X(t)$ be a regular adapted process such that with probability one $\int_0^T X^2(t)dt < \infty$. Then the Itô integral $\int_0^T X(t)dB(t)$ is defined and has the following properties.

- Linearity. If Itô integrals of $X(t)$ and $Y(t)$ are defined and α and β are some constants then

$$\int_0^T (\alpha X(t) + \beta Y(t))dB(t) = \alpha \int_0^T X(t)dB(t) + \beta \int_0^T Y(t)dB(t).$$

2.

$$\int_0^T X(t)I_{(a,b]}(t)dB(t) = \int_a^b X(t)dB(t).$$

The following two properties hold when the process satisfies an additional assumption

$$\int_0^T \mathbb{E}(X^2(t))dt < \infty. \quad (4.10)$$

3. *Zero mean property.* If condition (4.10) holds then

$$\mathbb{E} \left(\int_0^T X(t) dB(t) \right) = 0. \quad (4.11)$$

4. *Isometry property.* If condition (4.10) holds then

$$\mathbb{E} \left(\int_0^T X(t) dB(t) \right)^2 = \int_0^T \mathbb{E}(X^2(t)) dt. \quad (4.12)$$

PROOF. Only an outline of the proof is given. Firstly, it is shown that the Itô integral is well-defined for adapted processes that satisfy an additional assumption (4.10). Such processes can be approximated by simple processes

$$X^n(t) = X(0) + \sum_{i=0}^{n-1} X(t_i^n) I_{(t_i^n, t_{i+1}^n]}(t), \quad (4.13)$$

where $\{t_i^n\}$ is a partition of $[0, T]$ with $\delta_n = \max_i(t_{i+1}^n - t_i^n) \rightarrow 0$ as $n \rightarrow \infty$. In this sum only one term is different from zero, corresponding to the interval in the partition containing t . Now,

$$\int_0^T (X^n(t))^2 dt = \sum_{k=0}^{n-1} X^2(t_i^n)(t_{i+1}^n - t_i^n) \longrightarrow \int_0^T X^2(t) dt, \quad (4.14)$$

$$\int_0^T \mathbb{E}(X^n(t))^2 dt = \sum_{k=0}^{n-1} \mathbb{E} X^2(t_i^n)(t_{i+1}^n - t_i^n) \longrightarrow \int_0^T \mathbb{E} X^2(t) dt, \quad (4.15)$$

as $n \rightarrow \infty$, because the sums are Riemann sums for the corresponding integrals. Moreover, it is possible to show that

$$\lim_{n \rightarrow \infty} \int_0^T \mathbb{E}(X^n(t) - X(t))^2 dt = 0. \quad (4.16)$$

Denote the Itô integral of the simple process by

$$J_n = \int_0^T X^n(t) dB(t) = \sum_{k=0}^{n-1} X(t_i^n)(B(t_{i+1}^n) - B(t_i^n)). \quad (4.17)$$

The condition (4.10) for J_n holds, so that $\mathbb{E}(J_n) = 0$, and by the isometry property $\mathbb{E}(J_n^2)$ is given by the sum in (4.15). Using the isometry property

(4.16) implies that

$$\begin{aligned}
 E(J_n - J_m)^2 &= E\left(\int_0^T X^m(t)dB(t) - \int_0^T X^n(t)dB(t)\right)^2 \\
 &= E\left(\int_0^T (X^m(t) - X^n(t))dB(t)\right)^2 = E\int_0^T (X^m(t) - X^n(t))^2 dt \\
 &\leq 2E\int_0^T (X^m(t) - X(t))^2 dt + 2E\int_0^T (X^n(t) - X(t))^2 dt \longrightarrow 0,
 \end{aligned} \tag{4.18}$$

as $n, m \rightarrow \infty$. The space L^2 of random variables with zero mean, finite second moments and convergence in the mean square is complete, and (4.18) shows that J_n form a Cauchy sequence in this space. This implies that there is an element J , such that $J_n \rightarrow J$ in L^2 . This limit J is taken to be the Itô integral $\int_0^T X(t)dB(t)$. If we were to use another approximating sequence, then it is not hard to check, this limit does not change.

Now consider adapted processes with finite integral $\int_0^T X^2(t)dt$, but not necessarily of finite expectation. It can be shown, using the previous result, that such processes can be approximated by simple processes by taking limit in probability rather than in mean square. The sequence of corresponding Itô integrals is a Cauchy sequence in probability. It converges in probability to a limit $\int_0^T X(t)dB(t)$.

For details of this proof see, for example, Gihman and Skorohod (1972), Liptser and Shiryaev (1974), and Karatzas and Shreve (1988). \square

Note that Itô integrals need not have mean and variance, but when they do, the mean is zero and the variance is given by (4.12).

Corollary 4.4: *If X is a continuous adapted process then the Itô integral $\int_0^T X(t)dB(t)$ exists. In particular, $\int_0^T f(B(t))dB(t)$, where f is a continuous function on \mathbb{R} is well defined.*

PROOF. Since any path of $X(t)$ is a continuous function, $\int_0^T X^2(t)dt < \infty$, and the result follows by Theorem 4.3. If f is continuous on \mathbb{R} , then $f(B(t))$ is continuous on $[0, T]$. \square

Remark 4.4: It follows from the proof that the sums (4.17) approximate the Itô integral $\int_0^T X(t)dB(t)$

$$\sum_{k=0}^{n-1} X(t_i^n)(B(t_{i+1}^n) - B(t_i^n)).$$

In an approximation of the Stieltjes integral by sums, the function f on the interval $[t_i, t_{i+1}]$ is replaced by its value at some middle point $\theta_i \in [t_i, t_{i+1}]$, whereas in the above approximations for the Itô integral, the left most point must be taken for $\theta_i = t_i$, otherwise the process may not be adapted.

It is possible to define an integral (different to an Itô integral) when θ_i is chosen to be an interior point of the interval, $\theta_i = \lambda t_i + (1 - \lambda)t_{i+1}$, for some $\lambda \in (0, 1)$. The resulting integral may depend on the choice of λ . When $\lambda = 1/2$, the Stratonovich stochastic integral results. Calculus with such integrals is closely related to the Itô calculus.

Remark 4.5: Note that the Itô integral does not have the monotonicity property: $X(t) \leq Y(t)$ does not imply $\int_0^T X(t)dB(t) \leq \int_0^T Y(t)dB(t)$. A simple counter-example is $\int_0^1 1 \times dB(t) = B(1)$. With probability half this is smaller than 0, the Itô integral of 0.

We give examples of Itô integrals of the form $\int_0^1 f(B(t))dB(t)$ with and without the first two moments.

Example 4.3: Take $f(t) = e^t$. $\int_0^1 e^{B(t)}dB(t)$ is well-defined as e^x is continuous on \mathbb{R} . Since $E(\int_0^1 e^{2B(t)}dt) = \int_0^1 E(e^{2B(t)})dt = \int_0^1 e^{2t}dt = \frac{1}{2}(e^2 - 1) < \infty$, $E(\int_0^1 e^{B(t)}dB(t)) = 0$ and $E(\int_0^1 e^{B(t)}dB(t))^2 = \frac{1}{2}(e^2 - 1)$.

Example 4.4: Take $f(t) = t$, that is, consider $\int_0^1 B(t)dB(t)$. Then the condition (4.10) is satisfied since $\int_0^1 E(B^2(t))dt = \int_0^1 tdt = 1/2 < \infty$. Thus $\int_0^1 B(t)dB(t)$ has mean zero and variance $1/2$.

Example 4.5: Take $f(t) = e^{t^2}$, that is, consider $\int_0^1 e^{B^2(t)}dB(t)$. Although this integral is well-defined, the condition (4.10) fails, as $\int_0^1 E(e^{2B^2(t)})dt = \infty$, due to the fact that $E(e^{2B^2(t)}) = \int_R e^{2x^2} \frac{1}{\sqrt{2\pi t}} e^{-\frac{x^2}{2t}} dx = \infty$ for $t \geq 1/4$. Therefore we can not claim that this Itô integral has finite moments. By using martingale inequalities given in the sequel, it can be shown that the expectation of the Itô integral does not exist.

Example 4.6: Let $J = \int_0^1 t dB(t)$. We calculate $E(J)$ and $Var(J)$.

Since $\int_0^1 t^2 dt < \infty$, the Itô integral is defined. Since the integrand t is non-random, condition (4.10) holds and the integral has the first two moments, $E(J) = 0$, and $E(J^2) = \int_0^1 t^2 dt = 1/3$.

Example 4.7: For what values of α is the integral $\int_0^1 (1-t)^{-\alpha} dB(t)$ defined? For the Itô integral to be defined it must have $\int_0^1 (1-t)^{-2\alpha} dt < \infty$. This gives $\alpha < 1/2$.

A consequence of the isometry property is the expectation of the product of two Itô integrals.

Theorem 4.5: Let $X(t)$ and $Y(t)$ be regular adapted processes, such that $\mathbb{E} \int_0^T X(t)^2 dt < \infty$ and $\mathbb{E} \int_0^T Y(t)^2 dt < \infty$. Then

$$\mathbb{E} \left(\int_0^T X(t) dB(t) \int_0^T Y(t) dB(t) \right) = \int_0^T \mathbb{E}(X(t)Y(t)) dt. \quad (4.19)$$

PROOF. Denote the Itô integrals $I_1 = \int_0^T X(t) dB(t)$, $I_2 = \int_0^T Y(t) dB(t)$. Write their product by using the identity $I_1 I_2 = (I_1 + I_2)^2 / 2 - I_1^2 / 2 - I_2^2 / 2$. Then use the isometry property. \square

4.2 Itô Integral Process

Let X be a regular adapted process, such that $\int_0^T X^2(s) ds < \infty$ with probability one, so that $\int_0^t X(s) dB(s)$ is defined for any $t \leq T$. Since it is a random variable for any fixed t , $\int_0^t X(s) dB(s)$ as a function of the upper limit t defines a stochastic process

$$Y(t) = \int_0^t X(s) dB(s). \quad (4.20)$$

It is possible to show that there is a version of the Itô integral $Y(t)$ with continuous sample paths. It is always assumed that the continuous version of the Itô integral is taken. It will be seen later in this section that the Itô integral has a positive quadratic variation and infinite variation.

Martingale Property of the Itô Integral

It is intuitively clear from the construction of Itô integrals that they are adapted. In order to see this more formally, Itô integrals of simple processes are clearly adapted, and also continuous. Since $Y(t)$ is a limit of integrals of simple processes, it is itself adapted.

Suppose that in addition to the condition $\int_0^T X^2(s) ds < \infty$, condition (4.10) holds, $\int_0^T \mathbb{E} X^2(s) ds < \infty$. (The latter implies the former by Fubini's theorem.) Then $Y(t) = \int_0^t X(s) dB(s)$, $0 \leq t \leq T$, is defined and possesses the first two moments. It can be shown, first for simple processes and then in general, that for $s < t$,

$$\mathbb{E} \left(\int_s^t X(u) dB(u) | \mathcal{F}_s \right) = 0.$$

Thus

$$\mathbb{E}(Y(t) | \mathcal{F}_s) = \mathbb{E} \left(\int_0^t X(u) dB(u) | \mathcal{F}_s \right)$$

$$\begin{aligned}
&= \int_0^s X(u) dB(u) + \mathbb{E} \left(\int_s^t X(u) dB(u) | \mathcal{F}_s \right) \\
&= \int_0^s X(u) dB(u) = Y(s).
\end{aligned}$$

Therefore $Y(t)$ is a martingale. The second moments of $Y(t)$ are given by the isometry property,

$$\mathbb{E} \left(\int_0^t X(s) dB(s) \right)^2 = \int_0^t \mathbb{E} X^2(s) ds. \quad (4.21)$$

This shows that $\sup_{t \leq T} \mathbb{E}(Y^2(t)) = \int_0^T \mathbb{E} X^2(s) ds < \infty$.

Definition 4.6: A martingale is called *square integrable* on $[0, T]$ if its second moments are bounded.

Thus we have

Theorem 4.7: Let $X(t)$ be a regular adapted process such that $\int_0^T \mathbb{E} X^2(s) ds < \infty$. Then $Y(t) = \int_0^t X(s) dB(s)$, $0 \leq t \leq T$, is a continuous zero mean square integrable martingale.

Remark 4.6: If $\int_0^T \mathbb{E} X^2(s) ds = \infty$, then the Itô integral $\int_0^t X(s) dB(s)$ may fail to be a martingale, but it is always a *local martingale* (see Chapter 7) for definition and properties.

Theorem 4.7 provides a way of constructing martingales.

Corollary 4.8: For any bounded function f with only discontinuities of the first kind on \mathbb{R} , $\int_0^t f(B(s)) dB(s)$ is a square integrable martingale.

PROOF. $X(t) = f(B(t))$ is adapted and is regular. Since $|f(x)| < K$, for some constant $K > 0$, $\int_0^T \mathbb{E} f^2(B(s)) ds \leq KT$. The result follows by Theorem 4.7. \square

Quadratic Variation and Covariation of Itô Integrals

The Itô integral $Y(t) = \int_0^t X(s) dB(s)$, $0 \leq t \leq T$, is a random function of t . It is continuous and adapted. The quadratic variation of Y is defined by (see (1.13))

$$[Y, Y](t) = \lim \sum_{i=0}^{n-1} (Y(t_{i+1}^n) - Y(t_i^n))^2, \quad (4.22)$$

where for each n , $\{t_i^n\}_{i=0}^n$, is a partition of $[0, t]$, and the limit is in probability, taken over all partitions with $\delta_n = \max_i(t_{i+1}^n - t_i^n) \rightarrow 0$ as $n \rightarrow \infty$.

Theorem 4.9: *The quadratic variation of the Itô integral $\int_0^t X(s)dB(s)$ is given by*

$$\left[\int_0^t X(s)dB(s), \int_0^t X(s)dB(s) \right] (t) = \int_0^t X^2(s)ds. \quad (4.23)$$

It is easy to verify the result for simple processes (see Example 4.8). The general case can be proved by approximations by simple processes.

Example 4.8: For simplicity, suppose that X takes only two different values on $[0, 1]$: ξ_0 on $[0, 1/2]$, and ξ_1 on $(1/2, 1]$

$$X_t = \xi_0 I_{[0, 1/2]}(t) + \xi_1 I_{(1/2, 1]}(t).$$

It is easy to see that

$$Y(t) = \int_0^t X(s)dB_s = \begin{cases} \xi_0 B(t) & \text{if } t \leq 1/2 \\ \xi_0 B(1/2) + \xi_1 (B(t) - B(1/2)) & \text{if } t > 1/2. \end{cases}$$

Thus for any partition of $[0, t]$,

$$Y(t_{i+1}^n) - Y(t_i^n) = \begin{cases} \xi_0 (B(t_{i+1}^n) - B(t_i^n)) & \text{if } t_i^n < t_{i+1}^n \leq 1/2 \\ \xi_1 (B(t_{i+1}^n) - B(t_i^n)) & \text{if } 1/2 \leq t_i^n < t_{i+1}^n. \end{cases}$$

Including $1/2$ in a partition, one can verify the following: for $t \leq 1/2$

$$\begin{aligned} [Y, Y](t) &= \lim \sum_{i=0}^{n-1} (Y(t_{i+1}^n) - Y(t_i^n))^2 \\ &= \xi_0^2 \lim \sum_{i=0}^{n-1} (B(t_{i+1}^n) - B(t_i^n))^2 = \xi_0^2 [B, B](t) = \xi_0^2 t = \int_0^t X^2(s)ds; \end{aligned}$$

and for $t > 1/2$

$$\begin{aligned} [Y, Y](t) &= \lim \sum_{i=0}^{n-1} (Y(t_{i+1}^n) - Y(t_i^n))^2 \\ &= \xi_0^2 \lim_{t_i^n < 1/2} (B(t_{i+1}^n) - B(t_i^n))^2 + \xi_1^2 \lim_{t_i^n > 1/2} (B(t_{i+1}^n) - B(t_i^n))^2 \\ &= \xi_0^2 [B, B](1/2) + \xi_1^2 [B, B]((1/2, t]) = \int_0^t X^2(s)ds. \end{aligned}$$

The limits above are limits in probability when $\delta_n = \max_i \{(t_{i+1}^n - t_i^n)\} \rightarrow 0$. In the same way (4.23) is verified for any simple function.

Example 4.9: Using the formula (4.23), quadratic variation of the Itô integral

$$\left[\int_0^{\cdot} B(s)dB(s) \right] (t) = \int_0^t B^2(s)ds.$$

Corollary 4.10: If $\int_0^t X^2(s)ds > 0$, for all $t \leq T$, then the Itô integral $Y(t) = \int_0^t X(s)dB(s)$ has infinite variation on $[0, t]$ for all $t \leq T$.

PROOF. If $Y(t)$ were of finite variation, its quadratic variation would be zero, leading to a contradiction. \square

Like Brownian motion, the Itô integral $Y(t)$ is a continuous but nowhere differentiable function of t .

Let now $Y_1(t)$ and $Y_2(t)$ be Itô integrals of $X_1(t)$ and $X_2(t)$ with respect to the same Brownian motion $B(t)$. Then, clearly, the process $Y_1(t) + Y_2(t)$ is also an Itô integral of $X_1(t) + X_2(t)$ with respect to $B(t)$.

Quadratic covariation of Y_1 and Y_2 on $[0, t]$ is defined by

$$[Y_1, Y_2](t) = \frac{1}{2}([Y_1 + Y_2, Y_1 + Y_2](t) - [Y_1, Y_1](t) - [Y_2, Y_2](t)). \quad (4.24)$$

According to (4.23) it follows that

$$[Y_1, Y_2](t) = \int_0^t X_1(s)X_2(s)ds. \quad (4.25)$$

It is clear that $[Y_1, Y_2](t) = [Y_2, Y_1](t)$, and it can be seen that quadratic covariation is given by the limit in probability of products of increments of the processes Y_1 and Y_2 when partitions $\{t_i^n\}$ of $[0, t]$ shrink,

$$[Y_1, Y_2](t) = \lim \sum_{i=0}^{n-1} (Y_1(t_{i+1}^n) - Y_1(t_i^n))(Y_2(t_{i+1}^n) - Y_2(t_i^n)).$$

4.3 Itô Integral and Gaussian Processes

We have seen in Chapter 4.1 that the Itô integral of simple non-random processes is a normal random variable. It is easy to see by using moment generating functions (see Exercise 4.3) that a limit in probability of such a sequence is also Gaussian. This implies the following result.

Theorem 4.11: If $X(t)$ is non-random such that $\int_0^T X^2(s)ds < \infty$, then its Itô integral $Y(t) = \int_0^t X(s)dB(s)$ is a Gaussian process with zero mean and covariance function given by

$$\text{Cov}(Y(t), Y(t+u)) = \int_0^t X^2(s)ds, \quad u \geq 0. \quad (4.26)$$

Moreover, $Y(t)$ is a square integrable martingale.

PROOF. Since the integrand is non-random, $\int_0^t \mathbf{E} X^2(s) ds = \int_0^t X^2(s) ds < \infty$. According to the zero mean property of the Itô integral, Y has zero mean. In order to compute the covariance function write \int_0^{t+u} as $\int_0^t + \int_t^{t+u}$ and use the martingale property of $Y(t)$ to obtain

$$\mathbf{E} \left(\int_0^t X(s) dB(s) \mathbf{E} \left(\int_t^{t+u} X(s) dB(s) \middle| \mathcal{F}_t \right) \right) = 0.$$

Hence

$$\begin{aligned} \text{Cov}(Y(t), Y(t+u)) &= \mathbf{E} \left(\int_0^t X(s) dB(s) \int_0^{t+u} X(s) dB(s) \right) \\ &= \mathbf{E} \left(\int_0^t X(s) dB(s) \right)^2 = \int_0^t \mathbf{E} X^2(s) ds = \int_0^t X^2(s) ds. \end{aligned} \quad \square$$

A proof of normality of integrals of non-random processes will be done later by using Itô's formula.

Example 4.10: According to Theorem 4.11 $J = \int_0^t s dB(s)$ has a normal $N(0, t^3/3)$ distribution.

Example 4.11: Let $X(t) = 2I_{[0,1]}(t) + 3I_{(1,3)}(t) - 5I_{(3,4)}(t)$. Give the Itô integral $\int_0^4 X(t) dB(t)$ as a sum of random variables, give its distribution, mean, and variance. Show that the process $M(t) = \int_0^t X(s) dB(s)$, $0 \leq t \leq 4$, is a Gaussian process and a martingale.

$$\begin{aligned} \int_0^4 X(t) dB(t) &= \int_0^1 X(t) dB(t) + \int_1^3 X(t) dB(t) + \int_3^4 X(t) dB(t) \\ &= \int_0^1 2dB(t) + \int_1^3 3dB(t) + \int_3^4 (-5)dB(t) \\ &= 2(B(1) - B(0)) + 3(B(3) - B(1)) - 5(B(4) - B(3)). \end{aligned}$$

The Itô integral is a sum of three independent normal random variables (by independence of increments of Brownian motion), $2N(0, 1) + 3N(0, 2) - 5N(0, 1)$. Its distribution is $N(0, 47)$.

The martingale property and the Gaussian property of $M(t) = \int_0^t X(s) dB(s)$, $0 \leq t \leq 4$, follow from the independence of the increments of $M(t)$, zero mean increments, and the normality of the increments. $M(t) - M(s) = \int_s^t X(u) dB(u)$. Take for example $0 < s < t < 1$, then $M(t) - M(s) = \int_s^t X(u) dB(u) = 2(B(t) - B(s))$, which is independent of the Brownian motion up to time s , and has $N(0, 4(t-s))$ distribution.

If $0 < s < 1 < t < 3$, then $M(t) - M(s) = \int_s^t X(u) dB(u) = \int_s^1 X(u) dB(u) + \int_1^t X(u) dB(u) = 2(B(1) - B(s)) + 3(B(t) - B(1))$, which is independent of the

Brownian motion up to time s , $B(u), u \leq s$ (and also $M(u), u \leq s$), and has $N(0, 4(1-s) + 9(t-1))$ distribution. Other cases are similar. According to Theorem 2.23 the process $M(t)$ is Gaussian.

Independence of increments plus zero mean of increments imply the martingale property of $M(t)$. For example, If $0 < s < 1 < t < 3$, $E(M(t)|M(u), u \leq s) = E(M(s) + M(t) - M(s)|M(u), u \leq s) = M(s) + E(M(t) - M(s)|M(u), u \leq s) = M(s)$.

If $Y(t) = \int_0^t X(t, s) dB(s)$ where $X(t, s)$ depends on the upper integration limit t , then $Y(t)$ need not be a martingale, but remains a Gaussian process for non-random $X(t, s)$.

Theorem 4.12: For any $t \leq T$, let $X(t, s)$ be a regular non-random function with $\int_0^t X^2(t, s) ds < \infty$. Then the process $Y(t) = \int_0^t X(t, s) dB(s)$ is a Gaussian process with mean zero and covariance function $t, u \geq 0$

$$Cov(Y(t), Y(t+u)) = \int_0^t X(t, s) X(t+u, s) ds. \quad (4.27)$$

PROOF. For a fixed t , the distribution of $Y(t)$, as that of an Itô integral of a non-random function, is normal with mean 0 and variance $\int_0^t X^2(t, s) ds$. We do not prove the process is Gaussian (it can be seen by approximating $X(t, s)$ by functions of the form $f(t)g(s)$), but calculate the covariance. For $u > 0$

$$Y(t+u) = \int_0^t X(t+u, s) dB(s) + \int_t^{t+u} X(t+u, s) dB(s).$$

Since $X(t+u, s)$ is non-random, the Itô integral $\int_t^{t+u} X(t+u, s) dB(s)$ is independent of \mathcal{F}_t . Therefore

$$E\left(\int_0^t X(t, s) dB(s) \int_t^{t+u} X(t+u, s) dB(s)\right) = 0,$$

and

$$\begin{aligned} Cov(Y(t), Y(t+u)) &= E(Y(t)Y(t+u)) \\ &= E\left(\int_0^t X(t, s) dB(s) \int_0^t X(t+u, s) dB(s)\right) \\ &= \int_0^t X(t, s) X(t+u, s) ds, \end{aligned} \quad (4.28)$$

where the last equality is obtained by the expectation of a product of Itô integrals (Equation (4.19)). \square

4.4 Itô's Formula for Brownian Motion

Itô's formula, also known as the *change of variable* and the *chain rule*, is one of the main tools of stochastic calculus. It gives rise to many others, such as the Dynkin, Feynman–Kac, and integration by parts formulae.

Theorem 4.13: *If $B(t)$ is a Brownian motion on $[0, T]$, and $f(x)$ is a twice continuously differentiable function on \mathbb{R} , then for any $t \leq T$*

$$f(B(t)) = f(0) + \int_0^t f'(B(s))dB(s) + \frac{1}{2} \int_0^t f''(B(s))ds. \quad (4.29)$$

PROOF. Note first that both integrals in (4.29) are well-defined, the Itô integral by Corollary 4.4. Let $\{t_i^n\}$ be a partition of $[0, t]$. Clearly,

$$f(B(t)) = f(0) + \sum_{i=0}^{n-1} (f(B(t_{i+1}^n)) - f(B(t_i^n))).$$

Apply Taylor's formula to $f(B(t_{i+1}^n)) - f(B(t_i^n))$ to obtain

$$\begin{aligned} f(B(t_{i+1}^n)) - f(B(t_i^n)) &= f'(B(t_i^n))(B(t_{i+1}^n) - B(t_i^n)) \\ &\quad + \frac{1}{2} f''(\theta_i^n)(B(t_{i+1}^n) - B(t_i^n))^2, \end{aligned}$$

where $\theta_i^n \in (B(t_i^n), B(t_{i+1}^n))$. Thus

$$\begin{aligned} f(B(t)) &= f(0) + \sum_{i=0}^{n-1} f'(B(t_i^n))(B(t_{i+1}^n) - B(t_i^n)) \\ &\quad + \frac{1}{2} \sum_{i=0}^{n-1} f''(\theta_i^n)(B(t_{i+1}^n) - B(t_i^n))^2. \end{aligned} \quad (4.30)$$

Taking limits as $\delta_n \rightarrow 0$, the first sum in (4.30) converges with the Itô integral $\int_0^t f'(B(s))dB(s)$. According to Theorem 4.14 the second sum in (4.30) converges to $\int_0^t f''(B(s))ds$ and the result follows. \square

Theorem 4.14: *If g is a continuous function and $\{t_i^n\}$ represents partitions of $[0, t]$, then for any $\theta_i^n \in (B(t_i^n), B(t_{i+1}^n))$, the limit in probability*

$$\lim_{\delta_n \rightarrow 0} \sum_{i=0}^{n-1} g(\theta_i^n) (B(t_{i+1}^n) - B(t_i^n))^2 = \int_0^t g(B(s))ds. \quad (4.31)$$

PROOF. Take first $\theta_i^n = B(t_i^n)$ to be the left end of the interval $(B(t_i^n), B(t_{i+1}^n))$. We show that the sums converge in probability

$$\sum_{i=0}^{n-1} g(B(t_i^n)) (B(t_{i+1}^n) - B(t_i^n))^2 \rightarrow \int_0^t g(B(s)) ds. \quad (4.32)$$

By continuity of $g(B(t))$ and definition of the integral it follows that

$$\sum_{i=0}^{n-1} g(B(t_i^n)) (t_{i+1}^n - t_i^n) \rightarrow \int_0^t g(B(s)) ds. \quad (4.33)$$

Next we show that the difference between the sums converges to zero in L^2 ,

$$\sum_{i=0}^{n-1} g(B(t_i^n)) (B(t_{i+1}^n) - B(t_i^n))^2 - \sum_{i=0}^{n-1} g(B(t_i^n)) (t_{i+1}^n - t_i^n) \rightarrow 0. \quad (4.34)$$

With $\Delta B_i = B(t_{i+1}^n) - B(t_i^n)$, and $\Delta t_i = t_{i+1}^n - t_i^n$, by using conditioning it is seen that the cross-product term in the following expression vanishes and

$$\begin{aligned} E\left(\sum_{i=0}^{n-1} g(B(t_i^n)) ((\Delta B_i)^2 - \Delta t_i)\right)^2 &= E \sum_{i=0}^{n-1} g^2(B(t_i^n)) E(((\Delta B_i)^2 - \Delta t_i)^2 | \mathcal{F}_{t_i}) \\ &= 2E \sum_{i=0}^{n-1} g^2(B(t_i^n)) (\Delta t_i)^2 \leq \delta 2E \sum_{i=0}^{n-1} g^2(B(t_i^n)) \Delta t_i \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \end{aligned}$$

It follows that

$$\sum_{i=0}^{n-1} g(B(t_i^n)) ((\Delta B_i)^2 - \Delta t_i) \rightarrow 0,$$

in the square mean (L^2), implying (4.34) and that both sums in (4.33) and (4.32) have the same limit, and (4.32) is established. Now for any choice of θ_i^n we have as $\delta_n \rightarrow 0$,

$$\begin{aligned} &\sum_{i=0}^{n-1} (g(\theta_i^n) - g(B(t_i^n))) (B(t_{i+1}^n) - B(t_i^n))^2 \\ &\leq \max_i (g(\theta_i^n) - g(B(t_i^n))) \sum_{i=0}^{n-1} (B(t_{i+1}^n) - B(t_i^n))^2 \rightarrow 0. \quad (4.35) \end{aligned}$$

The first term converges to zero almost surely by continuity of g and B , and the second converges in probability to the quadratic variation of Brownian motion, t , implying convergence to zero in probability in (4.35). This implies that both sums $\sum_{i=0}^{n-1} g(\theta_i^n)(\Delta B_i)^2$ and $\sum_{i=0}^{n-1} g(B(t_i^n))(\Delta B_i)^2$ have the same limit in probability, and the result follows by (4.32). \square

Example 4.12: Taking $f(x) = x^m$, $m \geq 2$ we have

$$B^m(t) = m \int_0^t B^{m-1}(s) dB(s) + \frac{m(m-1)}{2} \int_0^t B^{m-2}(s) ds.$$

With $m = 2$,

$$B^2(t) = 2 \int_0^t B(s) dB(s) + t.$$

Rearranging we recover the result on the stochastic integral

$$\int_0^t B(s) dB(s) = \frac{1}{2} B^2(t) - \frac{1}{2} t.$$

Example 4.13: Taking $f(x) = e^x$ we have

$$e^{B(t)} = 1 + \int_0^t e^{B(s)} dB(s) + \frac{1}{2} \int_0^t e^{B(s)} ds.$$

4.5 Itô Processes and Stochastic Differentials

Definition of Itô Processes

An Itô process has the form

$$Y(t) = Y(0) + \int_0^t \mu(s) ds + \int_0^t \sigma(s) dB(s), \quad 0 \leq t \leq T, \quad (4.36)$$

where $Y(0)$ is \mathcal{F}_0 -measurable and processes $\mu(t)$ and $\sigma(t)$ are \mathcal{F}_t -adapted, such that $\int_0^T |\mu(t)| dt < \infty$ and $\int_0^T \sigma^2(t) dt < \infty$.

It is said that the process $Y(t)$ has the *stochastic differential* on $[0, T]$

$$dY(t) = \mu(t) dt + \sigma(t) dB(t), \quad 0 \leq t \leq T. \quad (4.37)$$

We emphasize that a representation (4.37) only has meaning by way of (4.36), and no other.

Note that the processes μ and σ in (4.36) may (and often do) depend on $Y(t)$ or $B(t)$ as well, or even on the whole past path of $B(s)$, $s \leq t$; for example, they may depend on the maximum of Brownian motion $\max_{s \leq t} B(s)$.

Example 4.14: Example 4.12 shows that

$$B^2(t) = t + 2 \int_0^t B(s) dB(s). \quad (4.38)$$

In other words, with $Y(t) = B^2(t)$ we can write $Y(t) = \int_0^t ds + \int_0^t 2B(s) dB(s)$. Thus $\mu(s) = 1$ and $\sigma(s) = 2B(s)$. The stochastic differential of $B^2(t)$

$$d(B^2(t)) = 2B(t) dB(t) + dt.$$

The only meaning this has is the integral relation (4.38).

Example 4.15: Example 4.13 shows that $Y(t) = e^{B(t)}$ has stochastic differential

$$de^{B(t)} = e^{B(t)}dB(t) + \frac{1}{2}e^{B(t)}dt,$$

or

$$dY(t) = Y(t)dB(t) + \frac{1}{2}Y(t)dt.$$

Itô's formula (4.29) in differential notation becomes for a C^2 function f

$$d(f(B(t))) = f'(B(t))dB(t) + \frac{1}{2}f''(B(t))dt. \quad (4.39)$$

Example 4.16: We find $d(\sin(B(t)))$.

$f(x) = \sin(x)$, $f'(x) = \cos(x)$, $f''(x) = -\sin(x)$. Thus

$$d(\sin(B(t))) = \cos(B(t))dB(t) - \frac{1}{2}\sin(B(t))dt.$$

Similarly,

$$d(\cos(B(t))) = -\sin(B(t))dB(t) - \frac{1}{2}\cos(B(t))dt.$$

Example 4.17: We find $d(e^{iB(t)})$ with $i^2 = -1$.

The application of Itô's formula to a complex-valued function means its application to the real and complex parts of the function. A formal application by treating i as another constant gives the same result. Using the above example, we can calculate $d(e^{iB(t)}) = d\cos(B(t)) + id\sin(B(t))$, or directly by using Itô's formula with $f(x) = e^{ix}$, we have $f'(x) = ie^{ix}$, $f''(x) = -e^{ix}$ and

$$d(e^{iB(t)}) = ie^{iB(t)}dB(t) - \frac{1}{2}e^{iB(t)}dt.$$

Thus $X(t) = e^{iB(t)}$ has stochastic differential

$$dX(t) = iX(t)dB(t) - \frac{1}{2}X(t)dt.$$

Quadratic Variation of Itô Processes

Let $Y(t)$ be an Itô process

$$Y(t) = Y(0) + \int_0^t \mu(s)ds + \int_0^t \sigma(s)dB(s), \quad (4.40)$$

where it is assumed that μ and σ are such that the integrals in question are defined. Then by the properties of the integrals, $Y(t)$, $0 \leq t \leq T$, is a (random) continuous function, the integral $\int_0^t \mu(s)ds$ is a continuous function of t and is of finite variation (it is differentiable almost everywhere),

and the Itô integral $\int_0^t \sigma(s) dB(s)$ is continuous. Quadratic variation of Y on $[0, t]$ is defined by (see (1.13))

$$[Y](t) = [Y, Y]([0, t]) = \lim_{\delta_n \rightarrow 0} \sum_{i=0}^{n-1} (Y(t_{i+1}^n) - Y(t_i^n))^2, \quad (4.41)$$

where for each n , $\{t_i^n\}$, is a partition of $[0, t]$, and the limit is in probability taken over partitions with $\delta_n = \max_i (t_{i+1}^n - t_i^n) \rightarrow 0$ as $n \rightarrow \infty$, and is given by

$$\begin{aligned} [Y](t) &= \left[\int_0^{\cdot} \mu(s) ds + \int_0^{\cdot} \sigma(s) dB(s) \right] (t) = \left[\int_0^{\cdot} \mu(s) ds \right] (t) \\ &\quad + 2 \left[\int_0^{\cdot} \mu(s) ds, \int_0^{\cdot} \sigma(s) dB(s) \right] (t) + \left[\int_0^{\cdot} \sigma(s) dB(s) \right] (t). \end{aligned}$$

The result on the covariation, Theorem 1.11, states that the quadratic covariation of a continuous function with a function of finite variation is zero. This implies that the quadratic covariation of the integral $\int_0^t \mu(s) ds$ with terms above is zero, and we obtain it by using the result on the quadratic variation of Itô integrals (Theorem 4.9)

$$[Y](t) = \left[\int_0^{\cdot} \sigma(s) dB(s) \right] (t) = \int_0^t \sigma^2(s) ds. \quad (4.42)$$

If $Y(t)$ and $X(t)$ have stochastic differentials with respect to the same Brownian motion $B(t)$, then clearly process $Y(t) + X(t)$ also has a stochastic differential with respect to the same Brownian motion. It follows that covariation of X and Y on $[0, t]$ exists and is given by

$$[X, Y](t) = \frac{1}{2} ([X + Y, X + Y](t) - [X, X](t) - [Y, Y](t)). \quad (4.43)$$

Theorem 1.11 has an important corollary

Theorem 4.15: *If X and Y are Itô processes and X is of finite variation, then covariation $[X, Y](t) = 0$.*

Example 4.18: Let $X(t) = \exp(t)$, $Y(t) = B(t)$, then $[X, Y](t) = [\exp, B](t) = 0$.

Introduce a convention that allows a formal manipulation with stochastic differentials,

$$dY(t)dX(t) = d[X, Y](t), \quad (4.44)$$

and in particular

$$(dY(t))^2 = d[Y, Y](t). \quad (4.45)$$

Since $X(t) = t$ is a continuous function of finite variation and $Y(t) = B(t)$ is continuous with quadratic variation t , the following rules follow

$$dB(t)dt = 0, \quad (dt)^2 = 0, \quad (4.46)$$

but

$$(dB(t))^2 = d[B, B](t) = dt. \quad (4.47)$$

Remark 4.7: In some texts, for example, Protter (1992), quadratic variation is defined by adding the value $Y^2(0)$ to (4.41). The definition given here gives a more familiar looking formula for integration by parts, and it is used in many texts, for example, Rogers and Williams (1987) p. 59, Metivier (1982) p. 175.

Integrals With Respect to Itô Processes

It is necessary to extend integration with respect to processes obtained from Brownian motion. Let the Itô integral process $Y(t) = \int_0^t X(s)dB(s)$ be defined for all $t \leq T$, where $X(t)$ is an adapted process, such that $\int_0^T X^2(s)ds < \infty$ with probability one. Let an adapted process $H(t)$ satisfy $\int_0^T H^2(s)X^2(s)ds < \infty$ with probability one. Then the Itô integral process $Z(t) = \int_0^t H(s)X(s)dB(s)$ is also defined for all $t \leq T$. In this case one can formally write by identifying $dY(t)$ and $X(t)dB(t)$,

$$Z(t) = \int_0^t H(s)dY(s) := \int_0^t H(s)X(s)dB(s). \quad (4.48)$$

In Chapter 8 integrals with respect to $Y(t)$ will be introduced in a direct way, but the result agrees with the one above.

More generally, if Y is an Itô process satisfying

$$dY(t) = \mu(t)dt + \sigma(t)dB(t), \quad (4.49)$$

and H is adapted and satisfies $\int_0^t H^2(s)\sigma^2(s)ds < \infty$, $\int_0^t |H(s)\mu(s)|ds < \infty$, then $Z(t) = \int_0^t H(s)dY(s)$ is defined as

$$Z(t) = \int_0^t H(s)dY(s) := \int_0^t H(s)\mu(s)ds + \int_0^t H(s)\sigma(s)dB(s). \quad (4.50)$$

Example 4.19: If $a(t)$ denotes the number of shares held at time t , then the gain from trading in shares during the time interval $[0, T]$ is given by $\int_0^T a(t)dS(t)$.

4.6 Itô's Formula for Itô Processes

Theorem 4.16 (Itô's Formula for $f(X(t))$): Let $X(t)$ have a stochastic differential for $0 \leq t \leq T$

$$dX(t) = \mu(t)dt + \sigma(t)dB(t). \quad (4.51)$$

If $f(x)$ is twice continuously differentiable (C^2 function), then the stochastic differential of the process $Y(t) = f(X(t))$ exists and is given by

$$\begin{aligned} df(X(t)) &= f'(X(t))dX(t) + \frac{1}{2}f''(X(t))d[X, X](t) \\ &= f'(X(t))dX(t) + \frac{1}{2}f''(X(t))\sigma^2(t)dt \\ &= \left(f'(X(t))\mu(t) + \frac{1}{2}f''(X(t))\sigma^2(t) \right) dt + f'(X(t))\sigma(t)dB(t). \end{aligned} \quad (4.52)$$

The meaning of the above is

$$f(X(t)) = f(X(0)) + \int_0^t f'(X(s))dX(s) + \frac{1}{2} \int_0^t f''(X(s))\sigma^2(s)ds, \quad (4.53)$$

where the first integral is an Itô integral with respect to the stochastic differential. Existence of the integrals in the formula (4.53) is assured by the arguments following Theorem 4.13. The proof also follows the same ideas as Theorem 4.13, and is omitted. Proofs of Itô's formula can be found in Liptser and Shiryaev (2001), Revuz and Yor (2001), Protter (1992), and Rogers and Williams (1990).

Example 4.20: Let $X(t)$ have stochastic differential

$$dX(t) = X(t)dB(t) + \frac{1}{2}X(t)dt. \quad (4.54)$$

We find a process X satisfying (4.54). Let us look for a *positive* process X . Using Itô's formula for $\ln X(t)$ ($(\ln x)' = 1/x$ and $(\ln x)'' = -1/x^2$),

$$\begin{aligned} d\ln X(t) &= \frac{1}{X(t)}dX(t) - \frac{1}{2X_t^2}dt \text{ by using } \sigma(t) = X(t) \\ &= dB(t) + \frac{1}{2}dt - \frac{1}{2}dt = dB(t). \end{aligned}$$

So that $\ln X(t) = \ln X(0) + B(t)$, and we find

$$X(t) = X(0)e^{B(t)}. \quad (4.55)$$

Using Itô's formula we verify that this $X(t)$ indeed satisfies (4.54). We do not claim at this stage that (4.55) is the only solution.

Integration by Parts

We give a representation of the quadratic covariation $[X, Y](t)$ of two Itô processes $X(t)$ and $Y(t)$ in terms of Itô integrals. This representation gives rise to the integration by parts formula.

Quadratic covariation is a limit over decreasing partitions of $[0, t]$,

$$[X, Y](t) = \lim_{\delta_n \rightarrow 0} \sum_{i=0}^{n-1} (X(t_{i+1}^n) - X(t_i^n))(Y(t_{i+1}^n) - Y(t_i^n)). \quad (4.56)$$

The sum on the right of (4.56) can be written as

$$\begin{aligned} &= \sum_{i=0}^{n-1} \left(X(t_{i+1}^n)Y(t_{i+1}^n) - X(t_i^n)Y(t_i^n) \right) \\ &\quad - \sum_{i=0}^{n-1} X(t_i^n)(Y(t_{i+1}^n) - Y(t_i^n)) - \sum_{i=0}^{n-1} Y(t_i^n)(X(t_{i+1}^n) - X(t_i^n)) \\ &= X(t)Y(t) - X(0)Y(0) \\ &\quad - \sum_{i=0}^{n-1} X(t_i^n)(Y(t_{i+1}^n) - Y(t_i^n)) - \sum_{i=0}^{n-1} Y(t_i^n)(X(t_{i+1}^n) - X(t_i^n)). \end{aligned}$$

The last two sums converge in probability to Itô integrals $\int_0^t X(s)dY(s)$ and $\int_0^t Y(s)dX(s)$, compare with Remark (4.4). Thus the following expression is obtained:

$$[X, Y](t) = X(t)Y(t) - X(0)Y(0) - \int_0^t X(s)dY(s) - \int_0^t Y(s)dX(s). \quad (4.57)$$

The formula for integration by parts (stochastic product rule) is given by

$$X(t)Y(t) - X(0)Y(0) = \int_0^t X(s)dY(s) + \int_0^t Y(s)dX(s) + [X, Y](t). \quad (4.58)$$

In differential notation this reads

$$d(X(t)Y(t)) = X(t)dY(t) + Y(t)dX(t) + d[X, Y](t). \quad (4.59)$$

If

$$dX(t) = \mu_X(t)dt + \sigma_X(t)dB(t), \quad (4.60)$$

$$dY(t) = \mu_Y(t)dt + \sigma_Y(t)dB(t), \quad (4.61)$$

then, as seen earlier, their quadratic covariation can be obtained formally by multiplication of dX and dY , namely

$$\begin{aligned} d[X, Y](t) &= dX(t)dY(t) \\ &= \sigma_X(t)\sigma_Y(t)(dB(t))^2 = \sigma_X(t)\sigma_Y(t)dt, \end{aligned}$$

leading to the formula

$$d(X(t)Y(t)) = X(t)dY(t) + Y(t)dX(t) + \sigma_X(t)\sigma_Y(t)dt.$$

Note that if one of the processes is of finite variation, then the covariation term is zero. Thus for such processes the stochastic product rule is the same as usual.

The integration by parts formula (4.59) can be established rigorously by making the argument above more precise, or by using Itô's formula for the function of two variables xy , or by approximations by simple processes.

Formula (4.57) provides an alternative representation for quadratic variation

$$[X, X](t) = X^2(t) - X^2(0) - 2 \int_0^t X(s)dX(s). \quad (4.62)$$

For Brownian motion this formula was established in Example 4.2.

It follows from the definition of quadratic variation that it is a non-decreasing process in t , and consequently it is of finite variation. It is also obvious from (4.62) that it is continuous. According to the polarization identity, covariation is also continuous and is of finite variation.

Example 4.21: $X(t)$ has stochastic differential

$$dX(t) = B(t)dt + dB(t), \quad X(0) = 0.$$

We find $X(t)$ and give its distribution, its mean, and covariance. $X(t) = tB(t)$ satisfies the above equation since the product rule for stochastic differentials is the same as usual, when one of the processes is continuous and of finite variation. Thus $X(t) = tB(t)$ is Gaussian, with mean zero, and covariance function

$$\begin{aligned} \gamma(t, s) &= \text{Cov}(X(t), X(s)) = \mathbb{E}(X(t)X(s)) \\ &= ts\mathbb{E}(B(t)B(s)) = ts\text{Cov}(B(t)B(s)) = ts\min(t, s). \end{aligned}$$

Example 4.22: Let $Y(t)$ have stochastic differential

$$dY(t) = \frac{1}{2}Y(t)dt + Y(t)dB(t), \quad Y(0) = 1.$$

Let $X(t) = tB(t)$. We find $d(X(t)Y(t))$.

$Y(t)$ is a geometric Brownian motion $e^{B(t)}$ (see Example 4.17). For $d(X(t)Y(t))$ use the product rule. We need the expression for $d[X, Y](t)$.

$$\begin{aligned} d[X, Y](t) &= dX(t)dY(t) = (B(t)dt + dB(t)) \left(\frac{1}{2}Y(t)dt + Y(t)dB(t) \right) \\ &= \frac{1}{2}B(t)Y(t)(dt)^2 + \left(B(t)Y(t) + \frac{1}{2}tY(t) \right) dB(t)dt \\ &\quad + tY(t)(dB(t))^2 = tY(t)dt, \end{aligned}$$

as $(dB(t))^2 = dt$ and all the other terms are zero. Thus

$$\begin{aligned} d(X(t)Y(t)) &= X(t)dY(t) + Y(t)dX(t) + d[X, Y](t) \\ &= X(t)dY(t) + Y(t)dX(t) + tY(t)dt, \end{aligned}$$

and substituting the expressions for X and Y the answer is obtained.

Example 4.23: Let f be a C^2 function and $B(t)$ Brownian motion. We find quadratic covariation $[f(B), B](t)$.

We find the answer by doing formal calculations. Using Itô's formula

$$df(B(t)) = f'(B(t))dB(t) + \frac{1}{2}f''(B(t))dt,$$

and the convention

$$d[f(B), B](t) = df(B(t))dB(t),$$

we have

$$\begin{aligned} d[f(B), B](t) &= df(B(t))dB(t) = f'(B(t))(dB(t))^2 \\ &\quad + \frac{1}{2}f''(B(t))dB(t)dt = f'(B(t))dt. \end{aligned}$$

Here we used $(dB)^2 = dt$, and $dBdt = 0$. Thus

$$[f(B), B](t) = \int_0^t f'(B(s))ds.$$

In a more intuitive way, from the definition of the covariation, taking limits over shrinking partitions

$$\begin{aligned} [f(B), B](t) &= \lim \sum_{i=0}^{n-1} (f(B(t_{i+1}^n)) - f(B(t_i^n))) (B(t_{i+1}^n) - B(t_i^n)) \\ &= \lim \sum_{i=0}^{n-1} \frac{f(B(t_{i+1}^n)) - f(B(t_i^n))}{B(t_{i+1}^n) - B(t_i^n)} (B(t_{i+1}^n) - B(t_i^n))^2 \\ &\approx \lim \sum_{i=0}^{n-1} f'(B(t_i^n)) (B(t_{i+1}^n) - B(t_i^n))^2 = \int_0^t f'(B(s))ds, \end{aligned}$$

where we have used Theorem 4.14 in the last equality.

Example 4.24: Let $f(t)$ be an increasing differentiable function, and let $X(t) = B(f(t))$. We show that

$$[X, X](t) = [B(f), B(f)](t) = [B, B]f(t) = f(t). \quad (4.63)$$

By taking limits over shrinking partitions

$$\begin{aligned}[X, X](t) &= \lim \sum_{i=0}^{n-1} (B(f(t_{i+1}^n)) - B(f(t_i^n)))^2 \\&= \lim \sum_{i=0}^{n-1} (f(t_{i+1}^n) - f(t_i^n)) \left(\frac{B(f(t_{i+1}^n)) - B(f(t_i^n))}{\sqrt{f(t_{i+1}^n) - f(t_i^n)}} \right)^2 \\&= \lim \sum_{i=0}^{n-1} (f(t_{i+1}^n) - f(t_i^n)) Z_i^2 = \lim T_n,\end{aligned}$$

where $Z_i = \frac{B(f(t_{i+1}^n)) - B(f(t_i^n))}{\sqrt{f(t_{i+1}^n) - f(t_i^n)}}$ are standard normal, and independent, by the properties of Brownian motion, and $T_n = \sum_{i=0}^{n-1} (f(t_{i+1}^n) - f(t_i^n)) Z_i^2$. Then for any n ,

$$\begin{aligned}\mathbb{E}(T_n) &= \sum_{i=0}^{n-1} (f(t_{i+1}^n) - f(t_i^n)) = f(t). \\Var(T_n) &= Var \left(\sum_{i=0}^{n-1} (f(t_{i+1}^n) - f(t_i^n)) Z_i^2 \right) = 3 \sum_{i=0}^{n-1} (f(t_{i+1}^n) - f(t_i^n))^2,\end{aligned}$$

by independence, and $Var(Z^2) = 3$. The last sum converges to zero, since f is of finite variation and continuous, implying that

$$\mathbb{E}(T_n - f(t))^2 \rightarrow 0.$$

This means that the limit in L^2 of T_n is $f(t)$, which implies that the limit in probability of T_n is $f(t)$, and

$$[B(f), B(f)] = f(t).$$

Itô's Formula for Functions of Two Variables

If two processes X and Y both possess a stochastic differential with respect to $B(t)$, and $f(x, y)$ has continuous partial derivatives up to order two, then $f(X(t), Y(t))$ also possesses a stochastic differential. In order to find its form consider formally the Taylor expansion of order two,

$$\begin{aligned}df(x, y) &= \frac{\partial f(x, y)}{\partial x} dx + \frac{\partial f(x, y)}{\partial y} dy \\&\quad + \frac{1}{2} \left(\frac{\partial^2 f(x, y)}{\partial x^2} (dx)^2 + \frac{\partial^2 f(x, y)}{\partial y^2} (dy)^2 + 2 \frac{\partial^2 f(x, y)}{\partial x \partial y} dxdy \right).\end{aligned}$$

Now, $(dX(t))^2 = dX(t)dX(t) = d[X, X](t) = \sigma_X^2(t)dt$,

$(dY(t))^2 = d[Y, Y]_t = \sigma_Y^2(t)dt$, and $dX(t)dY(t) = d[X, Y]_t$

$= \sigma_X(t)\sigma_Y(t)dt$, where $\sigma_X(t)$, and $\sigma_Y(t)$ are the diffusion coefficients of X and Y respectively. So we have

Theorem 4.17: Let $f(x, y)$ have continuous partial derivatives up to order two (a C^2 function), and X, Y be Itô processes, then

$$\begin{aligned} df(X(t), Y(t)) &= \frac{\partial f}{\partial x}(X(t), Y(t))dX(t) + \frac{\partial f}{\partial y}(X(t), Y(t))dY(t) \\ &\quad + \frac{1}{2} \frac{\partial^2 f}{\partial x^2}(X(t), Y(t))\sigma_X^2(t)dt + \frac{1}{2} \frac{\partial^2 f}{\partial y^2}(X(t), Y(t))\sigma_Y^2(t)dt \\ &\quad + \frac{\partial^2 f}{\partial x \partial y}(X(t), Y(t))\sigma_X(t)\sigma_Y(t)dt. \end{aligned} \quad (4.64)$$

The proof is similar to that of Theorem 4.13, and is omitted. It is stressed that differential formulae have meaning only through their integral representation.

Example 4.25: If $f(x, y) = xy$, then we obtain a differential of a product (or the product rule) which gives the integration by parts formula.

$$d(X(t)Y(t)) = X(t)dY(t) + Y(t)dX(t) + \sigma_X(t)\sigma_Y(t)dt.$$

An important case of Itô's formula is for functions of the form $f(X(t), t)$.

Theorem 4.18: Let $f(x, t)$ be twice continuously differentiable in x , and continuously differentiable in t (a $C^{2,1}$ function), and let X be an Itô process, then

$$\begin{aligned} df(X(t), t) &= \frac{\partial f}{\partial x}(X(t), t)dX(t) + \frac{\partial f}{\partial t}(X(t), t)dt \\ &\quad + \frac{1}{2}\sigma_X^2(t) \frac{\partial^2 f}{\partial x^2}(X(t), t)dt. \end{aligned} \quad (4.65)$$

This formula can be obtained from Theorem 4.17 by taking $Y(t) = t$ and by observing that $d[Y, Y] = 0$ and $d[X, Y] = 0$.

Example 4.26: We find a stochastic differential of $X(t) = e^{B(t)-t/2}$.

Use Itô's formula with $f(x, t) = e^{x-t/2}$. $X(t) = f(B(t), t)$ satisfies

$$\begin{aligned} dX(t) &= df(B(t), t) = \frac{\partial f}{\partial x}dB(t) + \frac{\partial f}{\partial t}dt + \frac{1}{2} \frac{\partial^2 f}{\partial x^2}dt \\ &= f(B(t), t)dB(t) - \frac{1}{2}f(B(t), t)dt + \frac{1}{2}f(B(t), t)dt \\ &= f(B(t), t)dB(t) = X(t)dB(t). \end{aligned}$$

So that

$$dX(t) = X(t)dB(t).$$

4.7 Itô Processes in Higher Dimensions

Let $\mathbf{B}(t) = (B_1(t), B_2(t), \dots, B_d(t))$ be Brownian motion in \mathbb{R}^d , that is, all coordinates $B_i(t)$ are independent one-dimensional Brownian motions. Let \mathcal{F}_t be the σ -field generated by $\mathbf{B}(s)$, $s \leq t$. Let $\mathbf{H}(t)$ be a regular adapted d -dimensional vector process, i.e. each of its coordinates is such. If for each j , $\int_0^T H_j^2(t)dt < \infty$, then the Itô integrals $\int_0^T H_j(t)dB_j(t)$ are defined. A single equivalent condition in terms of the length of the vector $|\mathbf{H}|^2 = \sum_{i=1}^d H_i^2$ is

$$\int_0^T |\mathbf{H}(t)|^2 dt < \infty.$$

It is customary to use a scalar product notation (even suppressing \cdot)

$$\begin{aligned} \mathbf{H}(t) \cdot d\mathbf{B}(t) &= \sum_{j=1}^d H_j(t)dB_j(t), \quad \text{and} \\ \int_0^T \mathbf{H}(t) \cdot d\mathbf{B}(t) &= \sum_{j=1}^d \int_0^T H_j(t)dB_j(t). \end{aligned} \quad (4.66)$$

If $b(t)$ is an integrable function, then the process

$$dX(t) = b(t)dt + \sum_{j=1}^d H_j(t)dB_j(t)$$

is well defined. It is a scalar Itô process driven by a d -dimensional Brownian motion. More generally, we can have any number n of processes driven by a d -dimensional Brownian motion, (the vector $\mathbf{H}_i = (\sigma_{i1}, \dots, \sigma_{id})$)

$$dX_i(t) = b_i(t)dt + \sum_{j=1}^d \sigma_{ij}(t)dB_j(t), \quad i = 1, \dots, n \quad (4.67)$$

where σ is $n \times d$ matrix valued function, \mathbf{B} is d -dimensional Brownian motion, \mathbf{X}, \mathbf{b} are n -dim vector-valued functions, and the integrals with respect to Brownian motion are Itô integrals. Then \mathbf{X} is called an Itô process. In vector form (4.67) becomes

$$d\mathbf{X}(t) = \mathbf{b}(t)dt + \sigma(t)d\mathbf{B}_t. \quad (4.68)$$

The dependence of $\mathbf{b}(t)$ and $\sigma(t)$ on time t can be via the whole path of the process up to time t , path of \mathbf{B}_s , $s \leq t$. The only restriction is that this dependence results in the following:

for any $i = 1, 2, \dots, n$, $b_i(t)$ is adapted and $\int_0^T |b_i(t)|dt < \infty$ almost surely
 for any $i = 1, 2, \dots, n$, and $j = 1, 2, \dots, d$, $\sigma_{ij}(t)$ is adapted and $\int_0^T \sigma_{ij}^2(t)dt < \infty$ almost surely, which assure existence of the required integrals.

An important case is when this dependence is of the form $\mathbf{b}(t) = \mathbf{b}(\mathbf{X}(t), t)$, $\sigma(t) = \sigma(\mathbf{X}(t), t)$. In this case the stochastic differential is written as

$$d\mathbf{X}(t) = \mathbf{b}(\mathbf{X}(t), t)dt + \sigma(\mathbf{X}(t), t)d\mathbf{B}(t), \quad (4.69)$$

and $\mathbf{X}(t)$ is then a diffusion process (see Chapters 5 and 6).

For Itô's formula we need the quadratic variation of a multi-dimensional Itô process. It is not hard to see that quadratic covariation of two independent Brownian motions is zero.

Theorem 4.19: *Let $B_1(t)$ and $B_2(t)$ be independent Brownian motions. Then their covariation process exists and is identically zero.*

PROOF. Let $\{t_i^n\}$ be a partition of $[0, t]$ and consider

$$T_n = \sum_{i=0}^{n-1} (B_1(t_{i+1}^n) - B_1(t_i^n))(B_2(t_{i+1}^n) - B_2(t_i^n)).$$

Using independence of B_1 and B_2 , $E(T_n) = 0$. Since increments of Brownian motion are independent, the variance of the sum is the sum of variances, and we have

$$\begin{aligned} Var(T_n) &= \sum_{i=0}^{n-1} E(B_1(t_{i+1}^n) - B_1(t_i^n))^2 E(B_2(t_{i+1}^n) - B_2(t_i^n))^2 \\ &= \sum_{i=0}^{n-1} (t_{i+1}^n - t_i^n)^2 \leq \max_i (t_{i+1}^n - t_i^n)t. \end{aligned}$$

Thus $Var(T_n) = E(T_n^2) \rightarrow 0$ as $\delta_n = \max_i (t_{i+1}^n - t_i^n) \rightarrow 0$. This implies that $T_n \rightarrow 0$ in probability, and the result is proved. \square

Thus for $k \neq l$, $k, l = 1, 2, \dots, d$,

$$[B_k, B_l](t) = 0. \quad (4.70)$$

Using (4.70), and the bi-linearity of covariation, it is easy to see from (4.67)

$$d[X_i, X_j](t) = dX_i(t)dX_j(t) = a_{ij}dt, \quad \text{for } i, j = 1, \dots, n. \quad (4.71)$$

where a , called the diffusion matrix, is given by

$$a = \sigma\sigma^{Tr}, \quad (4.72)$$

with σ^{Tr} denoting the transposed matrix of σ .

Itô's Formula for Functions of Several Variables

If $\mathbf{X}(t) = (X_1(t), X_2(t), \dots, X_n(t))$ is a vector Itô process, and $f(x_1, x_2, \dots, x_n)$ is a C^2 function of n variables, then $f(X_1(t), X_2(t), \dots, X_n(t))$ is also an Itô process, moreover, its stochastic differential is given by

$$\begin{aligned} df(X_1(t), X_2(t), \dots, X_n(t)) \\ = \sum_{i=1}^n \frac{\partial}{\partial x_i} f(X_1(t), X_2(t), \dots, X_n(t)) dX_i(t) \\ + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \frac{\partial^2}{\partial x_i \partial x_j} f(X_1(t), X_2(t), \dots, X_n(t)) d[X_i, X_j](t). \end{aligned} \quad (4.73)$$

When there is only one Brownian motion, $d = 1$, this formula is a generalization of Itô's formula for a function of two variables (Theorem 4.17).

For examples and applications see multi-dimensional diffusions in Chapters 5 and 6. We comment here on the integration by parts formula.

Remark 4.8 (Integration by Parts):

Let $X(t)$ and $Y(t)$ be two Itô processes that are adapted to *independent* Brownian motions B_1 and B_2 . Take $f(x, y) = xy$ and note that only one of the second derivatives is different from zero, $\frac{\partial^2 xy}{\partial x \partial y}$, but then the term it multiplies is zero, $d[B_1, B_2](t) = 0$ by Theorem 4.19. So the covariation of $X(t)$ and $Y(t)$ is zero, and one obtains from (4.73)

$$d(X(t)Y(t)) = X(t)dY(t) + Y(t)dX(t), \quad (4.74)$$

which is the usual integration by parts formula.

Remark 4.9: In some applications correlated Brownian motions are used. These are obtained by a linear transformation of independent Brownian motions. If B_1 and B_2 are independent, then the pair of processes B_1 and $W = \rho B_1 + \sqrt{1 - \rho^2} B_2$ are correlated Brownian motions. It is easy to see that W is indeed a Brownian motion, and that $d[B_1, W](t) = \rho dt$.

More results about Itô processes in higher dimensions are given in Chapter 6.

Remark 4.10: Itô's formula can be generalized to functions less smooth than C^2 , in particular for $f(x) = |x|$. Itô's formula for $f(x) = |x|$ becomes Tanaka's formula, and leads to the concept of local time. This development requires additional concepts, which are given later (see Chapter 8.7 in the general theory for semimartingales).

Notes: Material in this chapter can be found in Chung and Williams (1983), Gihman and Skorohod (1972), Liptser and Shiryaev (2001), (1989), Karatzas and Shreve (1988), Gard (1988), and Rogers and Williams (1990), (1994).

4.8 Exercises

Exercise 4.1: Give values of α for which the following process is defined $Y(t) = \int_0^t (t-s)^{-\alpha} dB(s)$. (This process is used in the definition of the so-called fractional Brownian motion.)

Exercise 4.2: Show that if X is a simple bounded adapted process, then $\int_0^t X(s) dB(s)$ is continuous.

Exercise 4.3: Let X_n be a Gaussian sequence convergent in distribution to X . Show that the distribution of X is either normal or degenerate. Deduce that if $EX_n \rightarrow \mu$ and $Var(X_n) \rightarrow \sigma^2 > 0$, then the limit is $N(\mu, \sigma^2)$. Since convergence in probability implies convergence in distribution, deduce convergence of Itô integrals of simple non-random processes to a Gaussian limit.

Exercise 4.4: Show that if $X(t)$ is non-random (does not depend on $B(t)$) and is a function of t and s with $\int_0^t X^2(t, s) ds < \infty$, then $\int_0^t X(t, s) dB(s)$ is a Gaussian random variable $Y(t)$. The collection $Y(t)$, $0 \leq t \leq T$, is a Gaussian process with zero mean and covariance function for $u \geq 0$ given by $Cov(Y(t), Y(t+u)) = \int_0^t X(t, s)X(t+u, s) ds$.

Exercise 4.5: Show that a Gaussian martingale on a finite time interval $[0, T]$ is a square integrable martingale with independent increments. Deduce that if X is non-random and $\int_0^t X^2(s) ds < \infty$, then $Y(t) = \int_0^t X(s) dB(s)$ is a Gaussian square integrable martingale with independent increments.

Exercise 4.6: Obtain the alternative relation for the quadratic variation of Itô processes, Equation (4.62), by applying Itô's formula to $X^2(t)$.

Exercise 4.7: $X(t)$ has a stochastic differential with $\mu(x) = bx + c$ and $\sigma^2(x) = 4x$. Assuming $X(t) \geq 0$, find the stochastic differential for the process $Y(t) = \sqrt{X(t)}$.

Exercise 4.8: A process $X(t)$ on $(0, 1)$ has a stochastic differential with coefficient $\sigma(x) = x(1-x)$. Assuming $0 < X(t) < 1$, show that the process defined by $Y(t) = \ln(X(t)/(1-X(t)))$ has a constant diffusion coefficient.

Exercise 4.9: $X(t)$ has a stochastic differential with $\mu(x) = cx$ and $\sigma^2(x) = x^a$, $c > 0$. Let $Y(t) = X(t)^b$. What choice of b will give a constant diffusion coefficient for Y ?

Exercise 4.10: Let $X(t) = tB(t)$ and $Y(t) = e^{B(t)}$. Find $d\left(\frac{X(t)}{Y(t)}\right)$.

Exercise 4.11: Obtain the differential of a ratio formula $d\left(\frac{X(t)}{Y(t)}\right)$ by taking $f(x, y) = x/y$. Assume that the process Y stays away from 0.

Exercise 4.12: Find $d(M(t))^2$, where $M(t) = e^{B(t)-t/2}$.

Exercise 4.13: Let $M(t) = B^3(t) - 3tB(t)$. Show that M is a martingale, first directly and then by using Itô integrals.

Exercise 4.14: Show that $M(t) = e^{t/2} \sin(B(t))$ is a martingale by using Itô's formula.

Exercise 4.15: For a function of n variables and n -dimensional Brownian motion, write Itô's formula for $f(B_1(t), \dots, B_n(t))$ by using gradient notation

$$\nabla f = \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n} \right).$$

Exercise 4.16: $\Phi(x)$ is the standard normal distribution function. Show that for a fixed $T > 0$ the process $\Phi\left(\frac{B(t)}{\sqrt{T-t}}\right)$, $0 \leq t \leq T$ is a martingale.

Exercise 4.17: Let $X(t) = (1-t) \int_0^t \frac{dB(s)}{1-s}$, where $0 \leq t < 1$. Find $dX(t)$.

Exercise 4.18: Let $X(t) = tB(t)$. Find its quadratic variation $[X, X](t)$.

Exercise 4.19: Let $X(t) = \int_0^t (t-s) dB(s)$. Find $dX(t)$ and its quadratic variation $[X, X](t)$. Compare to the quadratic variation of Itô integrals.

Chapter 5

Stochastic Differential Equations

Differential equations are used to describe the evolution of a system. Stochastic differential equations (SDEs) arise when a random noise is introduced into ordinary differential equations (ODEs). In this chapter we define two concepts of solutions of SDEs: the strong solution and the weak solution. We give a connection between SDEs and random ODEs, solutions to linear SDEs, stochastic exponential and logarithm, methods of solutions to some SDEs, and results on existence and uniqueness of solutions. Construction of weak solutions on the canonical space, as well as solutions to martingale problems, are more advanced, and can be skipped without impact on further reading.

5.1 Definition of Stochastic Differential Equations (SDEs)

Ordinary Differential Equations (ODEs)

If $x(t)$ is a differentiable function defined for $t \geq 0$, $\mu(x, t)$ is a function of x and t , and the following relation is satisfied for all t , $0 \leq t \leq T$

$$\frac{dx(t)}{dt} = x'(t) = \mu(x(t), t), \quad \text{and} \quad x(0) = x_0, \quad (5.1)$$

then $x(t)$ is a solution of the ODE with the initial condition x_0 . Usually, the requirement that $x'(t)$ is continuous is added (see also Theorem 1.4).

The above equation can be written in other forms as follows:

$$dx(t) = \mu(x(t), t) dt$$

and (by continuity of $x'(t)$)

$$x(t) = x(0) + \int_0^t \mu(x(s), s) ds.$$

Before we give a rigorous definition of SDEs, we show how they arise as randomly perturbed ODEs and give a physical interpretation.

White Noise and SDEs

The white noise process $\xi(t)$ is formally defined as the derivative of the Brownian motion,

$$\xi(t) = \frac{dB(t)}{dt} = B'(t). \quad (5.2)$$

It does not exist as a function of t in the usual sense, since a Brownian motion is nowhere differentiable.

If $\sigma(x, t)$ is the intensity of the noise at point x at time t , then it is agreed that $\int_0^T \sigma(X(t), t)\xi(t)dt = \int_0^T \sigma(X(t), t)B'(t)dt = \int_0^T \sigma(X(t), t)dB(t)$, where the integral is the Itô integral.

SDEs arise, for example, when the coefficients of ordinary equations are perturbed by white noise.

Example 5.1: The Black–Scholes–Merton model for growth with uncertain rate of return. $x(t)$ is the value of \$1 after time t , invested in a savings account. According to the definition of compound interest, it satisfies the ODE $dx(t)/x(t) = rdt$, or $dx(t)/dt = rx(t)$, (r is called the interest rate). If the rate is uncertain, it is taken to be perturbed by noise, $r + \sigma\xi(t)$, and the following SDE is obtained:

$$\frac{dX(t)}{dt} = (r + \sigma\xi(t))X(t),$$

meaning

$$dX(t) = rX(t)dt + \sigma X(t)dB(t).$$

Case $\sigma = 0$ corresponds to no noise, and recovers the deterministic equation. The solution of the deterministic equation is easily obtained by separating variables as $x(t) = e^{rt}$. The solution to the above SDE is given by a geometric Brownian motion, as can be verified by Itô's formula (see Example 5.5)

$$X(t) = e^{(r-\sigma^2/2)t+\sigma B(t)}. \quad (5.3)$$

Example 5.2: Population growth. If $x(t)$ denotes the population density, then the population growth can be described by the ODE $dx(t)/dt = ax(t)(1 - x(t))$. The growth is exponential with birth rate a when this density is small, and slows down when the density increases. Random perturbation of the birth rate results in the equation $dX(t)/dt = (a + \sigma\xi(t))X(t)(1 - X(t))$, or the SDE

$$dX(t) = aX(t)(1 - X(t))dt + \sigma X(t)(1 - X(t))dB(t).$$

A Physical Model of Diffusion and SDEs

The physical phenomena which give rise to the mathematical model of diffusion (and of Brownian motion) are the microscopic motions of a particle suspended in a fluid. Molecules of the fluid move with various velocities and collide with the particle from every possible direction, thus producing a constant bombardment. As a result of this bombardment, the particle exhibits an ever present erratic movement. This movement intensifies with an increase in the temperature of the fluid. Denote by $X(t)$ the displacement of the particle in one direction from its initial position at time t . If $\sigma(x, t)$ measures the effect of temperature at point x at time t , then the displacement due to bombardment during time $[t, t + \Delta]$ is modelled as $\sigma(x, t)(B(t + \Delta) - B(t))$. If the velocity of the fluid at point x at time t is $\mu(x, t)$, then the displacement of the particle due to the movement of the fluid during $[t, t + \Delta]$ is $\mu(x, t)\Delta$. Thus the total displacement from its position x at time t is given by

$$X(t + \Delta) - x \approx \mu(x, t)\Delta + \sigma(x, t)(B(t + \Delta) - B(t)). \quad (5.4)$$

Thus we obtain from this equation that the mean displacement from x during short time Δ is given by

$$\mathbb{E}((X(t + \Delta) - X(t))|X(t) = x) \approx \mu(x, t) \cdot \Delta, \quad (5.5)$$

and the second moment of the displacement from x during short time Δ is given by

$$\mathbb{E}((X(t + \Delta) - X(t))^2|X(t) = x) \approx \sigma^2(x, t)\Delta. \quad (5.6)$$

The above relations show that for small intervals of time both the mean and the second moment (and variance) of the displacement of a diffusing particle at time t at point x are proportional to the length of the interval, with coefficients $\mu(x, t)$ and $\sigma^2(x, t)$ respectively.

It can be shown that, taken as asymptotic relations as $\Delta \rightarrow 0$, that is, replacing \approx sign by the equality and adding terms $o(\Delta)$ to the right-hand sides, these two requirements characterize diffusion processes.

Assuming that $\mu(x, t)$ and $\sigma(x, t)$ are smooth functions, heuristic Equation (5.4) also indicates that for small intervals of time Δ , diffusions are approximately Gaussian processes. Given $X(t) = x$, $X(t + \Delta) - X(t)$ is approximately normally distributed, $N(\mu(x, t)\Delta, \sigma^2(x, t)\Delta)$. Of course, for large intervals of time, diffusions are not Gaussian unless the coefficients are non-random.

A stochastic differential equation is obtained heuristically from the relation (5.4) by replacing Δ by dt , and $\Delta B = B(t + \Delta) - B(t)$ by $dB(t)$, and $X(t + \Delta) - X(t)$ by $dX(t)$.

Stochastic Differential Equations

Let $B(t)$, $t \geq 0$, be a Brownian motion process. An equation of the form

$$dX(t) = \mu(X(t), t)dt + \sigma(X(t), t)dB(t), \quad (5.7)$$

where functions $\mu(x, t)$ and $\sigma(x, t)$ are given and $X(t)$ is the unknown process, is called a stochastic differential equation (SDE) driven by Brownian motion. The functions $\mu(x, t)$ and $\sigma(x, t)$ are called respectively the drift and the diffusion coefficient.

Definition 5.1: A process $X(t)$ is called a strong solution of the SDE (5.7) if for all $t > 0$ the integrals $\int_0^t \mu(X(s), s)ds$ and $\int_0^t \sigma(X(s), s)dB(s)$ exist, with the second being an Itô integral, and

$$X(t) = X(0) + \int_0^t \mu(X(s), s)ds + \int_0^t \sigma(X(s), s)dB(s). \quad (5.8)$$

Remark 5.1:

1. A strong solution is some function (functional) $F(t, (B(s), s \leq t))$ of the given Brownian motion $B(t)$.
2. When $\sigma = 0$, the SDE becomes an ODE.
3. Another interpretation of (5.7), called the weak solution, is a solution in distribution which will be given later.

Equations of the form (5.7) are called diffusion-type SDEs. More general SDEs have the form

$$dX(t) = \mu(t)dt + \sigma(t)dB(t), \quad (5.9)$$

where $\mu(t)$ and $\sigma(t)$ can depend on t and the whole past of the processes $X(t)$ and $B(t)$ ($X(s), B(s), s \leq t$), that is, $\mu(t) = \mu((X(s), s \leq t), t)$, $\sigma(t) = \sigma((X(s), s \leq t), t)$. The only restriction on $\mu(t)$ and $\sigma(t)$ is that they must be adapted processes, with respective integrals defined. Although many results (such as existence and uniqueness results) can be formulated for general SDEs, we concentrate here on diffusion-type SDEs.

Example 5.3: We have seen that $X(t) = \exp(B(t) - t/2)$ is a solution of the stochastic exponential SDE $dX(t) = X(t)dB(t)$, $X(0) = 1$.

Example 5.4: Consider the process $X(t)$ satisfying $dX(t) = a(t)dB(t)$, where $a(t)$ is non-random. Clearly, $X(t) = X(0) + \int_0^t a(s)dB(s)$. We can represent this as a function of the Brownian motion by integrating by parts, $X(t) = X(0) + a(t)B(t) - \int_0^t B(s)a'(s)ds$, assuming $a(t)$ is differentiable. In this case, the function $F(t, (x(s), s \leq t)) = X(0) + a(t)x(t) - \int_0^t x(s)a'(s)ds$.

The following two examples demonstrate how to find a strong solution by using Itô's formula and integration by parts.

Example 5.5: (Example 5.1 continued)

Consider the SDE

$$dX(t) = \mu X(t)dt + \sigma X(t)dB(t), \quad X(0) = 1. \quad (5.10)$$

Take $f(x) = \ln x$, then $f'(x) = 1/x$ and $f''(x) = -1/x^2$.

$$\begin{aligned} d(\ln X(t)) &= \frac{1}{X(t)}dX(t) + \frac{1}{2}\left(-\frac{1}{X^2(t)}\right)\sigma^2 X^2(t)dt \\ &= \frac{1}{X(t)}\left(\mu X(t)dt + \sigma X(t)dB(t)\right) - \frac{1}{2}\sigma^2 dt \\ &= (\mu - \frac{1}{2}\sigma^2)dt + \sigma dB(t), \end{aligned}$$

so that $Y(t) = \ln X(t)$ satisfies

$$dY(t) = (\mu - \frac{1}{2}\sigma^2)dt + \sigma dB(t).$$

Its integral representation gives

$$Y(t) = Y(0) + (\mu - \frac{1}{2}\sigma^2)t + \sigma B(t),$$

and

$$X(t) = X(0)e^{(\mu - \frac{1}{2}\sigma^2)t + \sigma B(t)}. \quad (5.11)$$

Example 5.6: (Langevin equation and Ornstein–Uhlenbeck process)

Consider the SDE

$$dX(t) = -\alpha X(t)dt + \sigma dB(t), \quad (5.12)$$

where α and σ are some non-negative constants.

Note that in the case $\sigma = 0$, the solution to the ODE is $x_0 e^{-\alpha t}$, in other words, $x(t)e^{\alpha t}$ is a constant. In order to solve the SDE consider the process $Y(t) = X(t)e^{\alpha t}$. Use the differential of the product rule, and note that the co-variation of $e^{\alpha t}$ with $X(t)$ is zero, as it is a differentiable function ($d(e^{\alpha t})dX(t) = \alpha e^{\alpha t} dt dX(t) = 0$), we have $dY(t) = e^{\alpha t} dX(t) + \alpha e^{\alpha t} X(t)dt$. Using the SDE for $dX(t)$, we obtain $dY(t) = \sigma e^{\alpha t} dB(t)$. This gives $Y(t) = Y(0) + \int_0^t \sigma e^{\alpha s} dB(s)$. Now the solution for $X(t)$ is

$$X(t) = e^{-\alpha t}(X(0) + \int_0^t \sigma e^{\alpha s} dB(s)). \quad (5.13)$$

The process $X(t)$ in (5.12) is known as the Ornstein–Uhlenbeck process.

We can also find the functional dependence of the solution on the Brownian motion path. Performing integration by parts, we find the function giving the strong solution

$$X(t) = F(t, (B(s), 0 \leq s \leq t)) = e^{-\alpha t} X(0) + \sigma B(t) - \sigma \alpha \int_0^t e^{-\alpha(t-s)} B(s)ds.$$

A more general equation is

$$dX(t) = (\beta - \alpha X(t))dt + \sigma dB(t), \quad (5.14)$$

with the solution

$$X(t) = \frac{\beta}{\alpha} + e^{-\alpha t} \left(X(0) - \frac{\beta}{\alpha} + \int_0^t \sigma e^{\alpha s} dB(s) \right). \quad (5.15)$$

Using Itô's formula, it is easy to verify that (5.15) is indeed a solution.

Example 5.7: Consider the SDE $dX(t) = B(t)dB(t)$.

Clearly, $X(t) = X(0) + \int_0^t B(s)dB(s)$, and using integration by parts (or Itô's formula), we obtain

$$X(t) = X(0) + \frac{1}{2}(B^2(t) - t).$$

Remark 5.2: If a strong solution exists, then by definition it is adapted to the filtration of the given Brownian motion, and as such it is intuitively clear that it is a function on the path $(B(s), s \leq t)$. Results of Yamada and Watanabe (1971), and Kallenberg (1996) state that provided the conditions of the existence and uniqueness theorem are satisfied, then there exists a function F such that the strong solution is given by $X(t) = F(t, (B(s), s \leq t))$. Not much is known about F in general. Often it is not easy to find this function even for Itô integrals $X(t) = \int_0^t f(B(s))dB(s)$, e.g. $X(t) = \int_0^t |B(s)|^{1/2}dB(s)$. For a representation of Itô integrals as functions of Brownian motion paths, see for example Rogers and Williams (1990).

Stochastic and Random Ordinary Differential Equations (ODEs)

Consider an SDE with a unit diffusion coefficient,

$$X(t) = X(0) + \int_0^t \mu(X(s), s)ds + B(t).$$

Note that if we denote $Y(t) = X(t) - B(t)$, then $X(t) = Y(t) + B(t)$, and $Y(t)$ satisfies the following equation:

$$Y(t) = Y(0) + \int_0^t \mu(Y(s) + B(s), s)ds. \quad (5.16)$$

However, this is an ODE, which involves Brownian motion $B(t)$

$$\frac{dY(t)}{dt} = \mu(Y(t) + B(t), t).$$

In the other direction, if $Y(t)$ solves the above random ODE, then $X(t) = Y(t) + B(t)$ solves the stochastic one. In this way, it is possible to solve a

stochastic equation by solving the ordinary one. Some stochastic equations with a more general diffusion coefficient $\sigma(x)$ can be transformed into the equation with the unit diffusion coefficient and the above method applied. This transformation is related to the Doss–Sussman method (Rogers and Williams, 1990; Karatzas and Shreve, 1988). For examples of solutions of SDEs see Klebaner and Azmy (2010).

Remark 5.3: Only some classes of SDEs admit a closed form solution. When a closed form solution is hard to find, an existence and uniqueness result is important because without it, it is not clear what exactly the equation means. When a solution exists and is unique, then numerical methods can be employed to compute it. Similarly to ordinary differential equations, linear SDEs can be solved explicitly.

5.2 Stochastic Exponential and Logarithm

Let X have a stochastic differential, and U satisfy

$$dU(t) = U(t)dX(t), \quad \text{and} \quad U(0) = 1, \quad \text{or} \quad U(t) = 1 + \int_0^t U(s)dX(s). \quad (5.17)$$

Then U is called the stochastic exponential of X , and is denoted by $\mathcal{E}(X)$. If $X(t)$ is of finite variation, then the solution to (5.17) is given by $U(t) = e^{X(t)}$. For Itô processes the solution is given by

Theorem 5.2: *The only solution of (5.17) is given by*

$$U(t) = \mathcal{E}(X)(t) := e^{X(t)-X(0)-\frac{1}{2}[X,X](t)}. \quad (5.18)$$

PROOF. The proof of existence of a solution to (5.17) consists of verification, by using Itô's formula, of (5.18). Write $U(t) = e^{V(t)}$, with $V(t) = X(t) - X(0) - \frac{1}{2}[X, X](t)$. Then

$$d\mathcal{E}(X)(t) = dU(t) = d(e^{V(t)}) = e^{V(t)}dV(t) + \frac{1}{2}e^{V(t)}d[V, V](t).$$

Since $[X, X](t)$ is of finite variation, and $X(t)$ is continuous, $[X, [X, X]](t) = 0$, and $[V, V](t) = [X, X](t)$. Using this with the expression for $V(t)$, we obtain

$$d\mathcal{E}(X)(t) = e^{V(t)}dX(t) - \frac{1}{2}e^{V(t)}d[X, X](t) + \frac{1}{2}e^{V(t)}d[X, X](t) = e^{V(t)}dX(t),$$

and (5.17) is established. The proof of uniqueness is done by assuming that there is another process satisfying (5.17), say $U_1(t)$, and showing by integration by parts that $d(U_1(t)/U(t)) = 0$. It is left as an exercise. \square

If U satisfies $dU(t) = U(t)dX(t)$ with arbitrary $U(0)$, then the solution is

$$U(t) = U(0)\mathcal{E}(X)(t).$$

Note that unlike in the case of the usual exponential $g(t) = \exp(f)(t) = e^{f(t)}$, the stochastic exponential $\mathcal{E}(X)$ requires the knowledge of all the values of the process up to time t , since it involves the quadratic variation term $[X, X](t)$.

Example 5.8: The stochastic exponential of Brownian motion $B(t)$ is given by $U(t) = \mathcal{E}(B)(t) = e^{B(t)-\frac{1}{2}t}$, and it satisfies for all t , $dU(t) = U(t)dB(t)$ with $U(0) = 1$.

Example 5.9: (Application in finance: stock process and its return process)
Let $S(t)$ denote the price of stock and assume that it is an Itô process, i.e. it has a stochastic differential. The process of the return on stock $R(t)$ is defined by the relation

$$dR(t) = \frac{dS(t)}{S(t)}.$$

In other words,

$$dS(t) = S(t)dR(t) \quad (5.19)$$

and the stock price is the stochastic exponential of the return. Returns are usually easier to model from first principles. For example, in the Black–Scholes model it is assumed that the returns over non-overlapping time intervals are independent and have finite variance. This assumption leads to the model for the return process $R(t) = \mu t + \sigma B(t)$. The stock price is then given by

$$\begin{aligned} S(t) &= S(0)\mathcal{E}(R)_t = S_0e^{R(t)-R(0)-\frac{1}{2}[R,R](t)} \\ &= S(0)e^{(\mu-\frac{1}{2}\sigma^2)t+\sigma B(t)}. \end{aligned} \quad (5.20)$$

Stochastic Logarithm

If $U = \mathcal{E}(X)$, then the process X is called the stochastic logarithm of U , denoted $\mathcal{L}(U)$. This is the inverse operation to the stochastic exponential. For example, the stochastic exponential of Brownian motion $B(t)$ is given by $e^{B(t)-\frac{1}{2}t}$. So $B(t)$ is the stochastic logarithm of $e^{B(t)-\frac{1}{2}t}$.

Theorem 5.3: Let U have a stochastic differential and not take value 0. Then the stochastic logarithm of U satisfies the SDE

$$dX(t) = \frac{dU(t)}{U(t)}, \quad X(0) = 0, \quad (5.21)$$

moreover

$$X(t) = \mathcal{L}(U)(t) = \ln \left(\frac{U(t)}{U(0)} \right) + \int_0^t \frac{d[U, U](S)}{2U^2(S)}. \quad (5.22)$$

PROOF. The SDE for the stochastic logarithm $\mathcal{L}(U)$ is by the definition of $\mathcal{E}(X)$. The solution (5.22) and uniqueness are obtained by Itô's formula. \square

Example 5.10: Let $U(t) = e^{B(t)}$. We find its stochastic logarithm $\mathcal{L}(U)$ directly and then verify (5.22). $dU(t) = e^{B(t)}dB(t) + \frac{1}{2}e^{B(t)}dt$. Hence

$$dX(t) = d\mathcal{L}(U)(t) = \frac{dU(t)}{U(t)} = dB(t) + \frac{1}{2}dt.$$

Thus

$$X(t) = \mathcal{L}(U)(t) = B(t) + \frac{1}{2}t.$$

Now, $d[U, U](t) = dU(t)dU(t) = e^{2B(t)}dt$, so that

$$\mathcal{L}(U)(t) = \ln U(t) + \int_0^t \frac{e^{2B(S)}dS}{2e^{2B(S)}} = B(S) + \int_0^t \frac{1}{2}dS = B(t) + \frac{1}{2}t,$$

which verifies (5.22).

Remark 5.4: The stochastic logarithm is useful in financial applications (see Kallsen and Shiryaev (2002)).

5.3 Solutions to Linear SDEs

Linear SDEs form a class of SDEs that can be solved explicitly. Consider a general linear SDE in one dimension

$$dX(t) = (\alpha(t) + \beta(t)X(t))dt + (\gamma(t) + \delta(t)X(t))dB(t), \quad (5.23)$$

where functions $\alpha, \beta, \gamma, \delta$ are *given* adapted processes, and are continuous functions of t . Examples considered in Chapter 5.2 are particular cases of linear SDEs.

Stochastic Exponential SDEs

Consider finding solutions in the case when $\alpha(t) = 0$ and $\gamma(t) = 0$. The SDE becomes

$$dU(t) = \beta(t)U(t)dt + \delta(t)U(t)dB(t). \quad (5.24)$$

This SDE is of the form

$$dU(t) = U(t)dY(t), \quad (5.25)$$

where the Itô process $Y(t)$ is defined by

$$dY(t) = \beta(t)dt + \delta(t)dB(t).$$

The SDE (5.24) is the stochastic exponential of Y (see Chapter 5.2). The stochastic exponential of Y is given by

$$\begin{aligned} U(t) &= U(0)\mathcal{E}(Y)(t) \\ &= U(0) \exp \left(Y(t) - Y(0) - \frac{1}{2}[Y, Y](t) \right) \\ &= U(0) \exp \left(\int_0^t \beta(s)ds + \int_0^t \delta(s)dB(s) - \frac{1}{2} \int_0^t \delta^2(s)ds \right) \\ &= U(0) \exp \left(\int_0^t \left(\beta(s) - \frac{1}{2}\delta^2(s) \right) ds + \int_0^t \delta(s)dB(s) \right), \end{aligned} \quad (5.26)$$

where $[Y, Y](t)$ is obtained from calculations $d[Y, Y](t) = dY(t)dY(t) = \delta^2(t)dt$.

General Linear SDEs

To find a solution for Equation (5.23) in the general case, look for a solution of the form

$$X(t) = U(t)V(t), \quad (5.27)$$

where

$$dU(t) = \beta(t)U(t)dt + \delta(t)U(t)dB(t), \quad (5.28)$$

and

$$dV(t) = a(t)dt + b(t)dB(t). \quad (5.29)$$

Set $U(0) = 1$ and $V(0) = X(0)$. Note that U is given by (5.26). Taking the differential of the product, it is easy to see that we can choose coefficients $a(t)$ and $b(t)$ in such a way that relation $X(t) = U(t)V(t)$ holds. The desired coefficients $a(t)$ and $b(t)$ turn out to satisfy equations

$$b(t)U(t) = \gamma(t), \quad \text{and} \quad a(t)U(t) = \alpha(t) - \delta(t)\gamma(t). \quad (5.30)$$

Using the expression for $U(t)$, $a(t)$ and $b(t)$ are then determined. Thus $V(t)$ is obtained, and $X(t)$ is found to be

$$X(t) = U(t) \left(X(0) + \int_0^t \frac{\alpha(s) - \delta(s)\gamma(s)}{U(s)} ds + \int_0^t \frac{\gamma(s)}{U(s)} dB(s) \right). \quad (5.31)$$

Langevin-Type SDE

Let $X(t)$ satisfy

$$dX(t) = a(t)X(t)dt + dB(t), \quad (5.32)$$

where $a(t)$ is a given adapted and continuous process. When $a(t) = -\alpha$, the equation is the Langevin equation (see Example 5.6).

We solve the SDE in two ways: by using the formula (5.31), and directly, similarly to Langevin's SDE.

Clearly, $\beta(t) = a(t)$, $\gamma(t) = 1$, and $\alpha(t) = \delta(t) = 0$. In order to find $U(t)$ we must solve $dU(t) = a(t)U(t)dt$, which gives $U(t) = e^{\int_0^t a(s)ds}$. Thus from (5.31) $X(t) = e^{\int_0^t a(s)ds}(X(0) + \int_0^t e^{-\int_0^u a(s)ds}dB(u))$.

Consider the process $e^{-\int_0^t a(s)ds}X(t)$ and use integration by parts. The process $e^{-\int_0^t a(s)ds}$ is continuous and is of finite variation. Therefore it has zero covariation with $X(t)$, hence

$$\begin{aligned} d\left(e^{-\int_0^t a(s)ds}X(t)\right) &= e^{-\int_0^t a(s)ds}dX(t) - a(t)e^{-\int_0^t a(s)ds}X(t)dt \\ &= e^{-\int_0^t a(s)ds}dB(t). \end{aligned}$$

Integrating we obtain

$$e^{-\int_0^t a(s)ds}X(t) = X(0) + \int_0^t e^{-\int_0^u a(s)ds}dB(u),$$

and finally

$$X(t) = X(0)e^{\int_0^t a(s)ds} + e^{\int_0^t a(s)ds} \int_0^t e^{-\int_0^u a(s)ds}dB(u). \quad (5.33)$$

Brownian Bridge

The Brownian bridge, or pinned Brownian motion, is a solution to the following SDE:

$$dX(t) = \frac{b - X(t)}{T - t}dt + dB(t), \quad \text{for } 0 \leq t < T, \quad X(0) = a. \quad (5.34)$$

This process is a transformed Brownian motion with fixed values at each end of the interval $[0, T]$, $X(0) = a$ and $X(T) = b$. The above SDE is a linear SDE with

$$\alpha(t) = \frac{b}{T - t}, \quad \beta(t) = -\frac{1}{T - t}, \quad \gamma(t) = 1, \quad \text{and } \delta(t) = 0.$$

Identifying $U(t)$ and $V(t)$ in (5.31) we obtain

$$X(t) = a\left(1 - \frac{t}{T}\right) + b\frac{t}{T} + (T - t) \int_0^t \frac{1}{T - s}dB(s), \quad \text{for } 0 \leq t < T. \quad (5.35)$$

Since the function under the Itô integral is deterministic, and for any $t < T$, $\int_0^t ds/(T-s)^2 < \infty$, the process $\int_0^t \frac{1}{T-s} dB(s)$ is a martingale. Moreover, it is Gaussian, by Theorem 4.11. Thus $X(t)$, on $[0, T]$ is a Gaussian process with initial value $X(0) = a$. The value at T , which is $X(T) = b$, is determined by continuity, see Example 5.11 below.

Thus a Brownian bridge is a continuous Gaussian process on $[0, T]$ with mean function $a(1-t/T)+bt/T$, and covariance function $\text{Cov}(X(t), X(s)) = \min(s, t) - st/T$.

Example 5.11: We show that $\lim_{t \uparrow T} (T-t) \int_0^t \frac{1}{T-s} dB(s) = 0$ almost surely. Using integration by parts, which is the same as the standard formula due to zero covariation between the deterministic term and Brownian motion, for any $t < T$

$$\int_0^t \frac{1}{T-s} dB(s) = B(t)/(T-t) - \int_0^t \frac{B(s)}{(T-s)^2} ds,$$

and

$$(T-t) \int_0^t \frac{1}{T-s} dB(s) = B(t) - (T-t) \int_0^t \frac{B(s)}{(T-s)^2} ds. \quad (5.36)$$

It is an exercise in calculus (by changing variables $u = 1/(t-s)$ or by considering integrals $\int_0^{T-\delta}$ and $\int_{T-\delta}^t$) to see that for any continuous function $g(s)$,

$$\lim_{t \uparrow T} (T-t) \int_0^t \frac{g(s)}{(T-s)^2} ds = g(T).$$

Applying this with $g(s) = B(s)$ shows that the limit in (5.36) is zero.

5.4 Existence and Uniqueness of Strong Solutions

Let $X(t)$ satisfy

$$dX(t) = \mu(X(t), t)dt + \sigma(X(t), t)dB(t). \quad (5.37)$$

Theorem 5.4 (Existence and Uniqueness): *If the following three conditions are satisfied*

1. *Coefficients are locally Lipschitz in x uniformly in t , that is, for every T and N there is a constant K depending only on T and N , such that for all $|x|, |y| \leq N$ and all $0 \leq t \leq T$*

$$|\mu(x, t) - \mu(y, t)| + |\sigma(x, t) - \sigma(y, t)| < K|x - y|. \quad (5.38)$$

2. Coefficients satisfy the linear growth condition

$$|\mu(x, t)| + |\sigma(x, t)| \leq K(1 + |x|). \quad (5.39)$$

3. $X(0)$ is independent of $(B(t), 0 \leq t \leq T)$, and $E X^2(0) < \infty$.

Then there exists a unique strong solution $X(t)$ of the SDE (5.37). $X(t)$ has continuous paths, moreover,

$$E\left(\sup_{0 \leq t \leq T} X^2(t)\right) < C(1 + E X^2(0)), \quad (5.40)$$

where constant C depends only on K and T .

The proof of existence is carried out by successive approximations similar to that for ordinary differential equations (Picard iterations). It can be found in Friedman (1975), Gihman and Skorohod (1972), and Rogers and Williams (1990). It is not hard to see, by using Gronwall's lemma, that the Lipschitz condition implies uniqueness.

The Lipschitz condition (5.38) holds if, for example, partial derivatives $\frac{\partial \mu}{\partial x}(t, x)$ and $\frac{\partial \sigma}{\partial x}(t, x)$ are bounded for $|x|, |y| \leq N$ and all $0 \leq t \leq T$, which in turn is true if the derivatives are continuous (see Chapter 1).

Less Stringent Conditions for Strong Solutions

The next result is specific for one-dimensional SDEs. It is given for the case of time-independent coefficients. A similar result holds for time-dependent coefficients, see for example, Ethier and Kurtz (1986), p. 298, Rogers and Williams (1990), p. 265.

Theorem 5.5 (Yamada–Watanabe): Suppose that $\mu(x)$ satisfies the Lipschitz condition and $\sigma(x)$ satisfies a Hölder condition of order α , $\alpha \geq 1/2$, that is, there is a constant K such that

$$|\sigma(x) - \sigma(y)| < K|x - y|^\alpha. \quad (5.41)$$

Then if a strong solution exists it is unique.

Example 5.12: (Girsanov's SDE)

$dX(t) = |X(t)|^r dB(t)$, $X(0) = 0$, $1/2 \leq r < 1$. Note that for such r , $|x|^r$ is Hölder but not Lipschitz (see Chapter 1). $X(t) \equiv 0$ is a strong solution. Since the conditions of Theorem 5.5 are satisfied, $X(t) \equiv 0$ is the only solution.

5.5 Markov Property of Solutions

The Markov property asserts that given the present state of the process, the future is independent of the past. This can be stated as follows: if \mathcal{F}_t denotes the σ -field generated by the process up to time t , then for any $0 \leq s < t$

$$\mathrm{P}(X(t) \leq y | \mathcal{F}_s) = \mathrm{P}(X(t) \leq y | X(s)) \quad \text{almost surely} \quad (5.42)$$

It is intuitively clear from the heuristic Equation (5.4) that solutions to SDEs should have the Markov property. For a small Δ , given $X(t) = x$, $X(t + \Delta)$ depends on $B(t + \Delta) - B(t)$, which is independent of the past.

We do not prove that strong solutions possess the Markov property. However, by the construction of the solution on the canonical space (a weak solution), it can be seen that the Markov property holds.

Transition Function

Markov processes are characterized by the transition probability function. Denote by

$$P(y, t, x, s) = \mathrm{P}(X(t) \leq y | X(s) = x) \quad (5.43)$$

the conditional distribution function of the random variable $X(t)$, given that $X(s) = x$, i.e. the distribution of the values at time t , given that the process was in the state x at time s .

Theorem 5.6: *Let $X(t)$ be a solution to the SDE (5.37). Then $X(t)$ has the Markov property.*

Using the law of total probability, by conditioning on all possible values z of the process at time u , for $s < u < t$, we obtain that the transition probability function $P(y, t, x, s)$ in (5.43) satisfies the Chapman–Kolmogorov equation

$$P(y, t, x, s) = \int_{-\infty}^{\infty} P(y, t, z, u) P(dz, u, x, s), \quad \text{for any } s < u < t. \quad (5.44)$$

In fact, any function that satisfies this equation and is a distribution function in y for fixed values of the other arguments, is a transition function of some Markov process.

Example 5.13: If $P(y, t, x, s) = \int_{-\infty}^y \frac{1}{\sqrt{2\pi(t-s)}} e^{\frac{(u-x)^2}{2(t-s)}} du$ is the cumulative distribution function of the normal $N(x, t-s)$ distribution, then the corresponding diffusion process is Brownian motion. Indeed, $\mathrm{P}(B(t) \leq y | \mathcal{F}_s) = \mathrm{P}(B(t) \leq y | B(s))$, and the conditional distribution of $B(t)$ given $B(s) = x$ is $N(x, t-s)$.

Example 5.14: Let $X(t)$ solve SDE $dX(t) = \mu X(t)dt + \sigma X(t)dB(t)$ for some constants μ and σ . We know (see Example 5.5) that $X(t) = X(0)e^{(\mu-\sigma^2/2)t+\sigma B(t)}$. Hence $X(t) = X(s)e^{(\mu-\sigma^2/2)(t-s)+\sigma(B(t)-B(s))}$ and its transition probability function $P(X(t) \leq y|X(s) = x) = P(X(s)e^{(\mu-\sigma^2/2)(t-s)+\sigma(B(t)-B(s))} \leq y|X(s) = x) = P(xe^{(\mu-\sigma^2/2)(t-s)+\sigma(B(t)-B(s))} \leq y|X(s) = x)$. Using independence of $B(t) - B(s)$ and $X(s)$, the conditional probability is given by $P(xe^{(\mu-\sigma^2/2)(t-s)+\sigma(B(t)-B(s))} \leq y) = P(e^{(\mu-\sigma^2/2)(t-s)+\sigma(B(t)-B(s))} \leq y/x)$. Thus $P(y, t, x, s) = \Phi(\frac{\ln(y/x) - (\mu - \sigma^2/2)(t-s)}{\sigma\sqrt{t-s}})$.

Remark 5.5: Introduce a useful representation, which requires us to keep track of when and where the process starts. Denote by $X_s^x(t)$ the value of the process at time t when it starts at time s from the point x . It is clear that for $0 \leq s < t$, $X_0^x(t) = X_s^{X_0^x(s)}(t)$. The Markov property states that conditionally on $X_s^{x_0}(t) = x$, the processes $X_s^{x_0}(u)$, $s \leq u \leq t$, and $X_t^x(u)$, $t \leq u$ are independent.

Definition 5.7: A process has the strong Markov property if the relation (5.42) holds when a non-random time s is replaced by a finite stopping time τ .

Solutions to SDEs have also the strong Markov property, meaning that given the history up to a stopping time τ , the behaviour of the process at some future time t is independent of the past (see also Chapter 3.4).

If an SDE has a strong solution $X(t)$, then $X(t)$ has a transition probability function $P(y, t, x, s)$. This function can be found as a solution to the forward or the backward partial differential equations (see Chapter 5.8).

A transition probability function $P(y, t, x, s)$ may exist for SDEs without a strong solution. This function in turn determines a Markov process uniquely (all finite-dimensional distributions). This process is known as a weak solution to an SDE. In this way, one can define a solution for an SDE under less stringent conditions on its coefficients. The concepts of the weak solution are considered next.

5.6 Weak Solutions to SDEs

The concept of weak solutions allows us to give a meaning to an SDE when strong solutions do not exist. Weak solutions are solutions in distribution, they can be realized (defined) on some other probability space, and they exist under less stringent conditions on the coefficients of the SDE.

Definition 5.8: If there exists a probability space with a filtration, a Brownian motion $\hat{B}(t)$, and a process $\hat{X}(t)$ adapted to that filtration, such that:

$\hat{X}(0)$ has the given distribution, for all t the integrals below are defined, and $\hat{X}(t)$ satisfies

$$\hat{X}(t) = \hat{X}(0) + \int_0^t \mu(\hat{X}(u), u) du + \int_0^t \sigma(\hat{X}(u), u) d\hat{B}(u), \quad (5.45)$$

then $\hat{X}(t)$ is called a weak solution to the SDE

$$dX(t) = \mu(X(t), t) dt + \sigma(X(t), t) dB(t). \quad (5.46)$$

Definition 5.9: A weak solution is called unique if any two solutions (possibly on different probability spaces) with the same distributions have the same finite dimensional distributions.

Clearly, by definition, a strong solution is also a weak solution. Uniqueness of the strong solution (pathwise uniqueness) implies uniqueness of the weak solution (a result of Yamada and Watanabe (1971)). In the next example a strong solution does not exist, but a weak solution exists and is unique.

Example 5.15: (Tanaka's SDE)

$$dX(t) = \text{sign}(X(t)) dB(t), \quad (5.47)$$

where

$$\text{sign}(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ -1 & \text{if } x < 0. \end{cases}$$

Since $\sigma(x) = \text{sign}(x)$ is discontinuous, it is not Lipschitz, and conditions for the strong existence fail. It can be shown that a strong solution to Tanaka's SDE does not exist (see for example Gihman and Skorohod (1972), and Rogers and Williams (1990)). We show that the Brownian motion is the unique weak solution of Tanaka's SDE. Let $X(t)$ be some Brownian motion. Consider the process

$$Y(t) = \int_0^t \frac{1}{\text{sign}(X(s))} dX(s) = \int_0^t \text{sign}(X(s)) dX(s).$$

$\text{Sign}(X(t))$ is adapted, $\int_0^T (\text{sign}(X(t)))^2 dt = T < \infty$, and $Y(t)$ is well defined and is a continuous martingale.

$$[Y, Y](t) = \int_0^t \text{sign}^2(X(s)) d[X, X](s) = \int_0^t ds = t.$$

By Levy's theorem (which is proven later), $Y(t)$ is a Brownian motion, call it $\hat{B}(t)$,

$$\hat{B}(t) = \int_0^t \frac{dX(s)}{\text{sign}(X(s))}.$$

Rewrite the last equality in the differential notation to obtain Tanaka's SDE. Levy's characterization theorem implies also that *any* weak solution is a Brownian motion.

Example 5.16: (Girsanov's SDE)

The equation

$$dX(t) = |X(t)|^r dB(t), \quad (5.48)$$

$r > 0, t \geq 0$ has a strong solution $X(t) \equiv 0$. For $r \geq 1/2$, this is the only strong solution by Theorem 5.5. Therefore there are no weak solutions other than zero. For $0 < r < 1/2$, the SDE has infinitely many weak solutions (Rogers and Williams (1990)). Therefore there is no strong uniqueness in this case, otherwise it would have only one weak solution. Compare this to the non-uniqueness of the solution of the equation $dx(t) = 2\sqrt{|x(t)|}dt$, which has solutions $x(t) = 0$ and $x(t) = t^2$.

5.7 Construction of Weak Solutions

In this section we give results on the existence and uniqueness of weak solutions to SDEs. Construction of weak solutions requires more advanced knowledge, and this section can be skipped.

Theorem 5.10: *If for each $t > 0$, functions $\mu(x, t)$ and $\sigma(x, t)$ are bounded and continuous, then the SDE (5.46) has at least one weak solution starting at time s at point x , for all s , and x . If in addition their partial derivatives with respect to x up to order two are also bounded and continuous, then the SDE (5.46) has a unique weak solution starting at time s at point x . Moreover, this solution has the strong Markov property.*

These results are proved in Stroock and Varadhan (1979), Chapter 6. However, better conditions are available; Stroock and Varadhan (1979), Corollary 6.5.5 (see also Pinsky (1995), Theorem 1.10.2).

Theorem 5.11: *If $\sigma(x, t)$ is positive and continuous, and for any $T > 0$ there is K_T such that for all $x \in \mathbb{R}$*

$$|\mu(x, t)| + |\sigma(x, t)| \leq K_T(1 + |x|), \quad (5.49)$$

then there exists a unique weak solution to SDE (5.46) starting at any point $x \in \mathbb{R}$ at any time $s \geq 0$; moreover, it has the strong Markov property.

Canonical Space for Diffusions

Solutions to SDEs or diffusions can be realized on the probability space of continuous functions. We indicate (a) how to define probability on this space by means of a transition function, (b) how to find the transition function from the given SDE, and (c) how to verify that the constructed process indeed satisfies the given SDE.

Probability Space $(\Omega, \mathcal{F}, \mathbb{P})$

Weak solutions can be constructed on the canonical space $\Omega = C([0, \infty))$ of continuous functions from $[0, \infty)$ to \mathbb{R} . The Borel σ -field on Ω is the one generated by the open sets. Open sets, in turn, are defined with the help of a metric, for example, an open ball of radius ϵ centered at ω is the set $D_\epsilon(\omega) = \{\omega' : d(\omega, \omega') < \epsilon\}$. The distance between two continuous functions ω_1 and ω_2 is taken as

$$d(\omega_1, \omega_2) = \sum_{n=1}^{\infty} \frac{1}{2^n} \frac{\sup_{0 \leq t \leq n} |\omega_1(t) - \omega_2(t)|}{1 + \sup_{0 \leq t \leq n} |\omega_1(t) - \omega_2(t)|}.$$

Convergence of the elements of Ω in this metric is the uniform convergence of functions on bounded closed intervals $[0, T]$. Diffusions on a finite interval $[0, T]$ can be realized on the space $C([0, T])$ with the metric

$$d(\omega_1, \omega_2) = \sup_{0 \leq t \leq T} |\omega_1(t) - \omega_2(t)|.$$

The canonical process $X(t)$ is defined by $X(t, \omega) = \omega(t)$, $0 \leq t < \infty$. It is known (see for example Dudley (1989)) that the Borel σ -field \mathcal{F} on $C([0, \infty))$ is given by $\sigma(X(t), 0 \leq t < \infty)$. The filtration is defined by the σ -fields $\mathcal{F}_t = \sigma(X(s), 0 \leq s \leq t)$.

Probability Measure

We outline the construction of probability measures from a given transition function $P(y, t, x, s)$. In particular, this construction gives the Wiener measure that corresponds to the Brownian motion process.

For any fixed $x \in \mathbb{R}$ and $s \geq 0$, a probability $\mathbb{P} = \mathbb{P}_{x,s}$ on (Ω, \mathcal{F}) can be constructed by using the following properties:

1. $\mathbb{P}(X(u) = x, 0 \leq u \leq s) = 1$.
2. $\mathbb{P}(X(t_2) \in B | \mathcal{F}_{t_1}) = P(B, t_2, X(t_1), t_1)$.

The second property asserts that for any Borel sets $A, B \subset \mathbb{R}$, we have

$$\begin{aligned} \mathbb{P}_{t_1, t_2}(A \times B) &= \mathbb{P}(X(t_1) \in A, X(t_2) \in B) \\ &= \mathbb{E}(\mathbb{P}(X(t_2) \in B | \mathcal{F}_{t_1}) I(X(t_1) \in A)) \\ &= \mathbb{E}(\mathbb{P}(B, t_2, X(t_1), t_1) I(X(t_1) \in A)) \\ &= \int_A \int_B \mathbb{P}(dy_2, t_2, y_1, t_1) \mathbb{P}_{t_1}(dy_1), \end{aligned}$$

where $P_{t_1}(C) = P(X(t_1) \in C)$. This extends to the n -dimensional cylinder sets $\{\omega \in \Omega : (\omega(t_1), \dots, \omega(t_n)) \in J_n\}$, where $J_n \subset \mathbb{R}^n$, by

$$P_{t_1, \dots, t_{n+1}}(J_{n+1}) = \int_{J_{n+1}} P(dy_{n+1}, t_{n+1}, y_n, t_n) P_{t_1, \dots, t_n}(dy_1 \times \dots \times dy_n).$$

These probabilities give the finite dimensional distributions $P((\omega(t_1), \dots, \omega(t_n)) \in J_n)$. Consistency of these probabilities is a consequence of the Chapman–Kolmogorov equation for the transition function. Thus by Kolmogorov's extension theorem P can be extended in a unique way to \mathcal{F} . This probability measure $P = P_{x,s}$ corresponds to the Markov process started at x at time s , denoted earlier by $X_s^x(t)$. Thus any transition function defines a probability so that the canonical process is a Markov process. We described in particular a construction of the Wiener measure, or Brownian motion.

Transition Function

Under appropriate conditions on the coefficients $\mu(x, t)$ and $\sigma(x, t)$, $P(y, t, x, s)$ is determined from a partial differential equation (PDE),

$$\frac{\partial u}{\partial s}(x, s) + L_s u(x, s) = 0, \quad (5.50)$$

called the backward PDE, involving a second order differential operator L_s ,

$$L_s f(x, s) = (L_s f)(x, s) = \frac{1}{2} \sigma^2(x, s) \frac{\partial^2 f}{\partial x^2}(x, s) + \mu(x, s) \frac{\partial f}{\partial x}(x, s). \quad (5.51)$$

It follows from the key property of the transition function, that

$$f(X(t)) - \int_s^t (L_u f)(X(u)) du \quad (5.52)$$

is a martingale under $P_{x,s}$ with respect to \mathcal{F}_t for $t \geq s$, for any twice continuously differentiable function f vanishing outside a finite interval (with compact support), $f \in C_K^2(\mathbb{R})$.

SDE on the Canonical Space is Satisfied

Extra concepts (namely that of local martingales and their integrals) are needed to prove the claim rigorously. The main idea is as follows. Suppose that (5.52) holds for functions $f(x) = x$ and $f(x) = x^2$. (Although they do not have a compact support, they can be approximated by C_K^2 functions on any finite interval.) Applying (5.52) to the linear function, we obtain that

$$Y(t) = X(t) - \int_s^t \mu(X(u), u) du \quad (5.53)$$

is a martingale. Applying (5.52) to the quadratic function, we obtain that

$$X^2(t) - \int_s^t (\sigma^2(X(u), u) + 2\mu(X(u), u)X(u)) du \quad (5.54)$$

is a martingale. According to the characterization property of quadratic variation for continuous martingales, $Y^2(t) - [Y, Y](t)$ is a martingale, and it follows from the above relations that $[Y, Y](t) = \int_s^t \sigma^2(X(u), u)du$. One can define the Itô integral process $B(t) = \int_s^t dY(u)/\sigma(X(u), u)$. From the properties of stochastic integrals it follows that $B(t)$ is a continuous local martingale with $[B, B](t) = t$. Thus by Levy's theorem $B(t)$ is Brownian motion. Putting all of the above together and using differential notation, the required SDE is obtained. For details see, for example, Rogers and Williams (1990), and also Stroock and Varadhan (1979).

Weak Solutions and the Martingale Problem

Taking the relation (5.52) as primary, Stroock and Varadhan defined a weak solution to the SDE

$$dX(t) = \mu(X(t), t)dt + \sigma(X(t), t)dB(t) \quad (5.55)$$

as a solution to the so-called martingale problem.

Definition 5.12: The martingale problem for the coefficients, or the operator L_s , is as follows. For each $x \in \mathbb{R}$ and $s > 0$, find a probability measure $P_{x,s}$ on Ω, \mathcal{F} such that

1. $P_{x,s}(X(u) = x, 0 \leq u \leq s) = 1$.
2. For any twice continuously differentiable function f vanishing outside a finite interval, the following process is a martingale under $P_{x,s}$ with respect to \mathcal{F}_t :

$$f(X(t)) - \int_s^t (L_u f)(X(u))du. \quad (5.56)$$

In the case when there is exactly one solution to the martingale problem, it is said that the martingale problem is *well-posed*.

Example 5.17: Brownian motion $B(t)$ is a solution to the martingale problem for the Laplace operator $L = \frac{1}{2} \frac{d^2}{dx^2}$, that is, for a twice continuously differentiable function f vanishing outside a finite interval

$$f(B(t)) - \int_0^t \frac{1}{2} f''(B(s))ds$$

is a martingale. Since Brownian motion exists and is determined by its distribution uniquely, the martingale problem for L is well-posed.

Remark 5.6: Note that if a function vanishes outside a finite interval, then its derivatives also vanish outside that interval. Thus for a twice continuously differentiable vanishing outside a finite interval function f ($f \in C_K^2$), $(L_s f)$ exists, is continuous, and vanishes outside that interval. This assures that the expectation of the process in (5.56) exists. If one demands only that f is twice continuously differentiable with bounded derivatives ($f \in C_b^2$), then $(L_s f)$ exists but may not be bounded, and expectation in (5.56) may not exist. If one takes ($f \in C_b^2$), then one seeks solutions to the local martingale problem, and any such solution makes the process in (5.56) into a local martingale. Local martingales are covered in Chapter 7.

As there are two definitions of weak solutions, Definition 5.8 and Definition 5.12, we show that they are the same.

Theorem 5.13: *Weak solutions in the sense of Definition 5.8 and in the sense of Definition 5.12 are equivalent.*

PROOF. We already indicated the proof in one direction, that if the martingale problem has a solution, then the solution satisfies the SDE. The other direction is obtained by using Itô's formula. Let $X(t)$ be a weak solution in the sense of Definition 5.8. Then there is a space supporting Brownian motion $B(t)$ so that

$$X(t) = X(s) + \int_s^t \mu(X(u), u) du + \int_s^t \sigma(X(u), u) dB(u), \text{ and } X(s) = x \quad (5.57)$$

is satisfied for all $t \geq s$. Let f be twice continuously differentiable with compact support. Applying Itô's formula to $f(X(t))$, we have

$$f(X(t)) = f(X(s)) + \int_s^t (L_u f)(X(u)) du + \int_s^t f'(X(u)) \sigma(X(u), u) dB(u). \quad (5.58)$$

Thus

$$f(X(t)) - \int_s^t (L_u f)(X(u)) du = f(X(s)) + \int_s^t f'(X(u)) \sigma(X(u), u) dB(u). \quad (5.59)$$

Since f and its derivatives vanish outside an interval, say $[-K, K]$, the functions $f'(x)\sigma(x, u)$ also vanish outside this interval for any u . Assuming that $\sigma(x, u)$ are bounded in x on finite intervals with the same constant for all u it follows that $|f'(x)\sigma(x, u)| < K_1$. Thus the integral $\int_s^t f'(X(u)) \sigma(X(u), u) dB(u)$ is a martingale in t , for $t \geq s$; thus the martingale problem has a solution. \square

5.8 Backward and Forward Equations

In many applications, such as physics, engineering, and finance, the importance of diffusions lies in their connection to PDEs, and often diffusions are specified by a PDE called the Fokker–Planck equation (introduced below (5.63), see for example Soize (1994)). Although PDEs are hard to solve in closed form, they can be easily solved numerically. In practice it is often enough to check that conditions of the existence and uniqueness result are satisfied, and then the solution can be computed by a PDE solver to the desired degree of accuracy.

This section outlines how to obtain the transition function that determines the weak solution to an SDE

$$dX(t) = \mu(X(t), t)dt + \sigma(X(t), t)dB(t), \quad \text{for } t \geq 0. \quad (5.60)$$

The results below are the main results from the theory of PDEs which are used for construction of diffusions (see for example Friedman (1975), Stroock and Varadhan (1979)).

Define the differential operator L_s , $0 \leq s \leq T$ by

$$L_s f(x, s) = (L_s f)(x, s) = \frac{1}{2} \sigma^2(x, s) \frac{\partial^2 f}{\partial x^2}(x, s) + \mu(x, s) \frac{\partial f}{\partial x}(x, s). \quad (5.61)$$

The operator L_s acts on twice continuously differentiable in x functions $f(x, s)$, and the result of its action on $f(x, s)$ is another function, denoted by $(L_s f)$, the values of which at point (x, s) are given by (5.61).

Definition 5.14: A fundamental solution of the PDE

$$\frac{\partial u}{\partial s}(x, s) + L_s u(x, s) = 0 \quad (5.62)$$

is a non-negative function $p(y, t, x, s)$ with the following properties:

1. It is jointly continuous in y, t, x, s , twice continuously differentiable in x , and satisfies Equation (5.62) with respect to s and x .
2. For any bounded continuous function $g(x)$ on \mathbb{R} , and any $t > 0$,

$$u(x, s) = \int_{\mathbb{R}} g(y) p(y, t, x, s) dy$$

is bounded, satisfies Equation (5.62) and $\lim_{s \uparrow t} u(x, s) = g(x)$, for $x \in \mathbb{R}$.

Theorem 5.15: Suppose that $\sigma(x, t)$ and $\mu(x, t)$ are bounded and continuous functions such that

- (A1) $\sigma^2(x, t) \geq c > 0$,
 (A2) $\mu(x, t)$ and $\sigma^2(x, t)$ satisfy a Hölder condition with respect to x and t , that is, for all $x, y \in \mathbb{R}$ and $s, t > 0$
 $|\mu(y, t) - \mu(x, s)| + |\sigma^2(y, t) - \sigma^2(x, s)| \leq K(|y - x|^\alpha + |t - s|^\alpha)$.

Then the PDE (5.62) has a fundamental solution $p(y, t, x, s)$, which is unique, and is strictly positive.

If in addition $\mu(x, t)$ and $\sigma(x, t)$ have two partial derivatives with respect to x , which are bounded and satisfy a Hölder condition with respect to x , then $p(y, t, x, s)$ as a function in y and t , satisfies the PDE

$$-\frac{\partial p}{\partial t} + \frac{1}{2} \frac{\partial^2}{\partial y^2} (\sigma^2(y, t)p) - \frac{\partial}{\partial y} (\mu(y, t)p) = 0. \quad (5.63)$$

Theorem 5.16: Suppose coefficients of L_s in (5.61) satisfy conditions (A1) and (A2) of Theorem 5.15. Then PDE (5.62) has a unique fundamental solution $p(y, t, x, s)$. The function $P(y, t, x, s) = \int_{-\infty}^y p(u, t, x, s)du$ uniquely defines a transition probability function. Moreover, this function has the property that for any bounded function $f(x, t)$ twice continuously differentiable in x and once continuously differentiable in t ($f \in C_b^{2,1}(\mathbb{R} \times [0, t])$)

$$\int_{\mathbb{R}} f(y, t)P(dy, t, x, s) - f(x, s) = \int_s^t \int_{\mathbb{R}} \left(\frac{\partial}{\partial u} + L_u \right) f(y, u)P(dy, u, x, s)du \quad (5.64)$$

for all $0 \leq s < t$, $x \in \mathbb{R}$.

The transition function $P(y, t, x, s)$ in the above theorem defines uniquely a Markov process $X(t)$, that is, for all x, y and $0 \leq s \leq t$

$$P(y, t, x, s) = \mathbb{P}(X(t) \leq y | X(s) = x). \quad (5.65)$$

Equation (5.62) is a PDE in the backward variables (x, s) and is therefore called the *backward equation*, also known as Kolmogorov's backward equation. Equation (5.63) is a PDE in the forward variables (y, t) and is therefore called the *forward equation*, also known as *Fokker–Planck equation*, *diffusion equation*, or Kolmogorov's forward equation.

The process $X(t)$ is called a diffusion, the differential operator L_s is called its generator. The property (5.64) implies that $X(t)$ satisfies the SDE (5.60).

Remark 5.7: A weak solution exists and is unique, possesses a strong Markov property, and has density under the conditions of Theorem 5.11, much weaker than those of Theorem 5.15.

5.9 Stratonovich Stochastic Calculus

Stochastic integrals in applications are often taken in the sense of Stratonovich calculus. This calculus is designed in such a way that its basic rules, such as the chain rule and integration by parts, are the same as in the standard calculus (e.g. Rogers and Williams (1990)). Although the rules of manipulations are the same, the calculi are still very different. The processes need to be adapted, just as in Itô calculus. Since Stratonovich stochastic integrals can be reduced to Itô integrals, the standard SDE theory can be used for Stratonovich stochastic differential equations. Note also that the Stratonovich integral is more suited to generalizations of stochastic calculus on manifolds (see Rogers and Williams (1990)).

A direct definition of the Stratonovich integral, denoted $\int_0^t Y(s) \partial X(s)$, is done as a limit in mean square (L^2) of Stratonovich approximating sums

$$\sum_{i=0}^{n-1} \frac{1}{2} (Y(t_{i+1}^n) + Y(t_i^n)) (X(t_{i+1}^n) - X(t_i^n)), \quad (5.66)$$

when partitions $\{t_i^n\}$ become finer and finer. In Stratonovich approximating sums the average value of Y on the interval (t_i^n, t_{i+1}^n) , $\frac{1}{2}(Y(t_{i+1}^n) + Y(t_i^n))$, is taken, whereas in the Itô integral the left most value of $Y(t_i^n)$ is taken. An alternative definition of the Stratonovich integral is given by using the Itô integral below.

Definition 5.17: Let X and Y be continuous adapted processes, such that the stochastic integral $\int_0^t Y(s) dX(s)$ is defined. The Stratonovich integral is defined by

$$\int_0^t Y(s) \partial X(s) = \int_0^t Y(s) dX(s) + \frac{1}{2}[Y, X](t). \quad (5.67)$$

The Stratonovich differential is defined by

$$Y(t) \partial X(t) = Y(t) dX(t) + \frac{1}{2} d[Y, X](t). \quad (5.68)$$

Integration by Parts: Stratonovich Product Rule

Theorem 5.18: *Provided all terms below are defined,*

$$X(t)Y(t) - X(0)Y(0) = \int_0^t X(s) \partial Y(s) + \int_0^t Y(s) \partial X(s), \quad (5.69)$$

$$\partial(X(t)Y(t)) = X(t) \partial Y(t) + Y(t) \partial X(t). \quad (5.70)$$

PROOF. The proof is the direct application of the stochastic product rule,

$$\begin{aligned} d(X(t)Y(t)) &= X(t)dY(t) + Y(t)dX(t) + d[X, Y](t) \\ &= X(t)dY(t) + \frac{1}{2}[X, Y](t) + Y(t)dX(t) + \frac{1}{2}[X, Y](t) \\ &= X(t)\partial Y(t) + Y(t)\partial X(t). \end{aligned}$$

□

Change of Variables: Stratonovich Chain Rule

Theorem 5.19: Let X be continuous and f three times continuously differentiable (in C^3), then

$$\begin{aligned} f(X(t)) - f(X(0)) &= \int_0^t f'(X(s))\partial X(s), \\ \partial f(X(t)) &= f'(X(t))\partial X(t). \end{aligned} \tag{5.71}$$

PROOF. According to Itô's formula $f(X(t))$ is a semimartingale, and by definition of the stochastic integral $f(X(t)) - f(X(0)) = \int_0^t df(X(s))$. According to Itô's formula

$$df(X(t)) = f'(X(t))dX(t) + \frac{1}{2}f''(X(t))d[X, X](t).$$

Let $Y(t) = f'(X(t))$. Then according to (5.68) it is enough to show that $d[Y, X](t) = f''(X(t))d[X, X](t)$. However this follows by Itô's formula as

$$dY(t) = df'(X(t)) = f''(X(t))dX(t) + \frac{1}{2}f'''(X(t))d[X, X](t),$$

and

$$d[Y, X](t) = dY(t)dX(t) = f''(X(t))dX(t)dX(t) = f''(X(t))d[X, X](t),$$

as needed. □

Example 5.18: If $B(t)$ is Brownian motion, then its Stratonovich stochastic differential is

$$\partial B^2(t) = 2B(t)\partial B(t),$$

as compared to its Itô differential

$$dB^2(t) = 2B(t)dB(t) + dt.$$

Conversion of Stratonovich SDEs into Itô SDEs

Theorem 5.20: Suppose that $X(t)$ satisfies the following SDE in the Stratonovich sense

$$dX(t) = \mu(X(t))dt + \sigma(X(t))\partial B(t), \quad (5.72)$$

with $\sigma(x)$ twice continuously differentiable. Then $X(t)$ satisfies the Itô SDE

$$dX(t) = \left(\mu(X(t)) + \frac{1}{2}\sigma'(X(t))\sigma(X(t)) \right)dt + \sigma(X(t))dB(t). \quad (5.73)$$

Thus the infinitesimal drift coefficient in Itô diffusion is $\mu(x) + \frac{1}{2}\sigma'(x)\sigma(x)$, and the diffusion coefficient is the same $\sigma(x)$.

PROOF. According to the definition of the Stratonovich integral $X(t)$ satisfies

$$dX(t) = \mu(X(t))dt + \sigma(X(t))dB(t) + \frac{1}{2}d[\sigma(X), B](t). \quad (5.74)$$

Since $[\sigma(X), B](t)$ is a finite variation process, it follows that $X(t)$ solves a diffusion type SDE with the same diffusion coefficient $\sigma(X(t))$. Computing formally the bracket, we have

$$d[\sigma(X), B](t) = d\sigma(X(t))dB(t).$$

Applying Itô's formula

$$d\sigma(X(t)) = \sigma'(X(t))dX(t) + \frac{1}{2}\sigma''(X(t))d[X, X](t).$$

It follows from (5.74) that

$$d[X, B](t) = dX(t)dB(t) = \sigma(X(t))dt,$$

therefore

$$d[\sigma(X), B](t) = d\sigma(X(t))dB(t) = \sigma'(X(t))dX(t)dB(t) = \sigma'(X(t))\sigma(X(t))dt.$$

Equation (5.73) now follows from (5.74). \square

Notes: Proofs and other details can be found in Dynkin (1965), Friedman (1975), (1976), Karatzas and Shreve (1988), and Stroock and Varadhan (1979).

5.10 Exercises

Exercise 5.1: (Gaussian diffusions)

Show that if $X(t)$ satisfies the SDE $dX(t) = a(t)dt + b(t)dB(t)$, with deterministic bounded coefficients $a(t)$ and $b(t)$, such that $\int_0^T |a(t)|dt < \infty$, and $\int_0^T b^2(t)dt < \infty$, then $X(t)$ is a Gaussian process with independent Gaussian increments.

Exercise 5.2: Give the SDEs for $X(t) = \cos(B(t))$ and $Y(t) = \sin(B(t))$.

Exercise 5.3: Solve the SDE $dX(t) = B(t)X(t)dt + B(t)X(t)dB(t)$, $X(0) = 1$.

Exercise 5.4: Solve the SDE $dX(t) = X(t)dt + B(t)dB(t)$, $X(0) = 1$. Comment on whether it is a diffusion type SDE.

Exercise 5.5: Find $d(\mathcal{E}(B)(t))^2$.

Exercise 5.6: Let $X(t)$ satisfy $dX(t) = X^2(t)dt + X(t)dB(t)$, $X(0) = 1$. Show that $X(t)$ satisfies $X(t) = e^{\int_0^t (X(s)-1/2)ds + B(t)}$.

Exercise 5.7: By definition, the stochastic logarithm satisfies $\mathcal{L}(\mathcal{E}(X)) = X$. Show that, provided $U(t) \neq 0$ for any t , $\mathcal{E}(\mathcal{L}(U)) = U$.

Exercise 5.8: Find the stochastic logarithm of $B^2(t) + 1$.

Exercise 5.9: Let $\mathbf{B}(t)$ be a d -dimensional Brownian motion, and $\mathbf{H}(t)$ a d -dimensional regular adapted process. Show that

$$\mathcal{E}\left(\int_0^\cdot \mathbf{H}(s)d\mathbf{B}(s)\right)(t) = \exp\left(\int_0^t \mathbf{H}(s)d\mathbf{B}(s) - \frac{1}{2} \int_0^t |\mathbf{H}(s)|^2 ds\right).$$

Exercise 5.10: Find the transition probability function $P(y, t, x, s)$ for Brownian motion with drift $B(t) + t$.

Exercise 5.11: Show that under the assumptions of Theorem 5.15 the transition function $P(y, t, x, s)$ satisfies the backward equation. Give also the forward equation for $P(y, t, x, s)$ and explain why it requires extra smoothness conditions on the coefficients $\mu(x, t)$ and $\sigma(x, t)$ for it to hold.

Exercise 5.12: Let $X(t)$ satisfy the following SDE for $0 \leq t \leq T$, $dX(t) = \sqrt{X(t) + 1}dB(t)$, and $X(0) = 0$. Assuming that Itô integrals are martingales, find $\mathbb{E}X(t)$, and $\mathbb{E}(X^2(t))$. Let $m(u, t) = \mathbb{E}e^{uX(t)}$ be the moment generating function of $X(t)$. Show that it satisfies the PDE

$$\frac{\partial m}{\partial t} = \frac{u^2}{2} \frac{\partial m}{\partial u} + \frac{u^2}{2} m.$$

Exercise 5.13: Solve the following Stratonovich SDE $\partial U = U\partial B$, $U(0) = 1$, where $B(t)$ is Brownian motion.

This page intentionally left blank

Chapter 6

Diffusion Processes

In this chapter various properties of solutions of stochastic differential equations (SDEs) are studied. The approach taken here relies on martingales obtained by means of Itô's formula. Relationships between SDEs and partial differential equations (PDEs) are given, but no prior knowledge of PDEs is required. Solutions to SDEs are referred to as diffusions.

6.1 Martingales and Dynkin's Formula

Itô's formula provides a source for construction of martingales. Let $X(t)$ solve the SDE

$$dX(t) = \mu(X(t), t)dt + \sigma(X(t), t)dB(t), \text{ for } t \geq 0, \quad (6.1)$$

and L_t be the generator of $X(t)$, that is, the second order differential operator associated with SDE (6.1),

$$L_t f(x, t) = (L_t f)(x, t) = \frac{1}{2}\sigma^2(x, t)\frac{\partial^2 f}{\partial x^2}(x, t) + \mu(x, t)\frac{\partial f}{\partial x}(x, t). \quad (6.2)$$

Itô's formula (4.65) takes a compact form.

Theorem 6.1: *For any twice continuously differentiable in x , and once in t function $f(x, t)$*

$$df(X(t), t) = \left(L_t f(X(t), t) + \frac{\partial f}{\partial t}(X(t), t) \right) dt + \frac{\partial f}{\partial x}(X(t), t)\sigma(X(t), t)dB(t). \quad (6.3)$$

Since under appropriate conditions the Itô integral is a martingale (see Theorem 4.7), by isolating the Itô integral martingales are obtained.

In order to illustrate this simple idea, let f have a bounded (by K) derivative, and use Itô's formula for $f(B(t))$. Then

$$f(B(t)) = f(0) + \int_0^t \frac{1}{2} f''(B(s))ds + \int_0^t f'(B(s))dB(s).$$

The Itô integral $\int_0^t f'(B(s))dB(s)$ is a martingale on $[0, T]$ because condition (4.10) holds, $\int_0^T (f'(B(s)))^2 ds < K^2 T < \infty$. Thus $f(B(t)) - \int_0^t \frac{1}{2} f''(B(s))ds$ is a martingale. The next result is more general.

Theorem 6.2: *Let $X(t)$ be a solution to SDE (6.1) with coefficients satisfying conditions of Theorem 5.4, that is, $\mu(x, t)$ and $\sigma(x, t)$ are Lipschitz in x with the same constant for all t , and satisfy the linear growth condition (1.30), $|\mu(x, t)| + |\sigma(x, t)| \leq K(1 + |x|)$. If $f(x, t)$ is a twice continuously differentiable in x and once in t function ($C^{2,1}$) with bounded first derivative in x , then the process*

$$M_f(t) = f(X(t), t) - \int_0^t \left(L_u f + \frac{\partial f}{\partial t} \right) (X(u), u) du \quad (6.4)$$

is a martingale.

PROOF. According to Itô's formula

$$M_f(t) = \int_0^t \frac{\partial f}{\partial x} (X(u), u) \sigma(X(u), u) dB(u). \quad (6.5)$$

By assumption, $\frac{\partial f}{\partial x}(x, u)$ is bounded for all x and u , $\left(\frac{\partial f}{\partial x}(x, u)\right)^2 < K_1$. Therefore

$$\int_0^T E\left(\frac{\partial f}{\partial x}(X(u), u) \sigma(X(u), u)\right)^2 du \leq K_1 \int_0^T E\left(\sigma^2(X(u), u)\right) du. \quad (6.6)$$

Using the linear growth condition,

$$\int_0^T E\left(\frac{\partial f}{\partial x}(X(u), u) \sigma(X(u), u)\right)^2 du \leq 2K_1 K^2 T \left(1 + E\left(\sup_{u \leq T} X^2(u)\right)\right). \quad (6.7)$$

However, $E\left(\sup_{u \leq T} X^2(u)\right) < \infty$ by the existence and uniqueness result, Theorem 5.4, therefore the expression in (6.7) is finite. Thus the Itô integral is a martingale by Theorem 4.7. \square

The condition of bounded partial derivative of f can be replaced by the exponential growth condition (see for example Pinsky (1995)).

Theorem 6.3: Let $X(t)$ satisfy conditions of Theorem 6.2. If $|X(0)|$ possesses moment generating function $\mathbb{E}e^{u|X(0)|} < \infty$, for all real u , then so does $|X(t)|$, $\mathbb{E}e^{u|X(t)|} < \infty$ for all $t \geq 0$. In this case

$$M_f(t) = f(X(t), t) - \int_0^t \left(L_u f + \frac{\partial f}{\partial t} \right) (X(u), u) du \quad (6.8)$$

is a martingale for all $f(x, t) \in C^{2,1}$, satisfying the following condition: for any t , there exist constants c_t and k_t such that for all x , all $t > 0$, and all $0 \leq u \leq t$

$$\max \left(\left| \frac{\partial f(x, u)}{\partial t} \right|, \left| \frac{\partial f(x, u)}{\partial x} \right|, \left| \frac{\partial^2 f(x, t)}{\partial x^2} \right| \right) \leq c_t e^{k_t |x|}. \quad (6.9)$$

PROOF. The proof is given in Pinsky (1995) for bounded coefficients of the SDE, but it can be extended for this case. We give the proof when the diffusion $X(t) = B(t)$ is Brownian motion. Let $X(t) = B(t)$, then by Itô's formula $M_f(t)$ is given by

$$M_f(t) = \int_0^t \frac{\partial f(B(s), s)}{\partial x} dB(s). \quad (6.10)$$

By the bound on $|\frac{\partial f(x, s)}{\partial x}|$, for $s \leq t$

$$\mathbb{E} \left(\frac{\partial f(B(s), s)}{\partial x} \right)^2 \leq c_t^2 \mathbb{E} \left(e^{2k_t |B(s)|} \right).$$

Writing the last expectation as an integral with respect to the density of the $N(0, t)$ distribution, it is evident that it is finite, and its integral over $[0, t]$ is finite,

$$\int_0^t \mathbb{E} \left(\frac{\partial f(B(s), s)}{\partial x} \right)^2 ds < \infty. \quad (6.11)$$

According to the martingale property of Itô integrals, (6.11) implies that the Itô integral (6.10) is a martingale. One can prove the result without use of Itô's formula by doing calculations of integrals with respect to normal densities (see for example Rogers and Williams (1987)). \square

Corollary 6.4: Let $f(x, t)$ solve the backward equation

$$L_t f(x, t) + \frac{\partial f}{\partial t}(x, t) = 0, \quad (6.12)$$

and the conditions of either of the two theorems above hold. Then $f(X(t), t)$ is a martingale.

Example 6.1: Let $X(t) = B(t)$, then $(Lf)(x) = \frac{1}{2}f''(x)$. Solutions to $Lf = 0$ are linear functions $f(x) = ax + b$. Hence $f(B(t)) = aB(t) + b$ is a martingale, which is also obvious from the fact that $B(t)$ is a martingale.

Example 6.2: Let $X(t) = B(t)$. The function $f(x, t) = e^{x-t/2}$ solves the backward equation

$$\frac{1}{2} \frac{\partial^2 f}{\partial x^2}(x, t) + \frac{\partial f}{\partial t}(x, t) = 0. \quad (6.13)$$

Therefore, by Corollary 6.4 we recover the exponential martingale of Brownian motion $e^{B(t)-t/2}$.

Corollary 6.5 (Dynkin's Formula): *Let $X(t)$ satisfy (6.1). If the conditions of either of the above theorems hold, then for any t , $0 \leq t \leq T$,*

$$\mathbb{E}f(X(t), t) = f(X(0), 0) + \mathbb{E} \int_0^t \left(L_u f + \frac{\partial f}{\partial t} \right) (X(u), u) du. \quad (6.14)$$

The result is also true if t is replaced by a bounded stopping time τ , $0 \leq \tau \leq T$.

PROOF. The bounds on the growth of the function and its partial derivatives are used to establish integrability of $f(X(t), t)$ and other terms in (6.14). Since $M_f(t)$ is a martingale, the result follows by taking expectations. For bounded stopping times the result follows by the optional stopping theorem, given in Chapter 7. \square

Remark 6.1: If f and $Lf + \frac{\partial f}{\partial t}$ are bounded, then their difference is a bounded martingale, for which the optional stopping theorem can be applied and Dynkin's formula holds. Remember that continuous functions on compacts are bounded.

Example 6.3: We show that $J = \int_0^T f(t) dB(t)$ has $N(0, \int_0^T f^2(t) dt)$ distribution. We carry out calculations for $J = \int_0^1 s dB(s)$ by finding its moment generating function $m(u) = \mathbb{E}(e^{uJ})$. The case of general f is similar. Consider the Itô integral $X(t) = \int_0^t s dB(s)$, $t \leq 1$, and note that $J = X(1)$. As $dX(t) = t dB(t)$, $X(t)$ is an Itô process with $\mu(x, t) = 0$ and $\sigma(x, t) = t$. Take $f(x, t) = f(x) = e^{ux}$. This function satisfies conditions of Theorem 6.3. It is easy to see that $L_t f(x, t) = \frac{1}{2}t^2 u^2 e^{ux}$, note that $\frac{\partial f}{\partial t} = 0$. Therefore by Dynkin's formula

$$\mathbb{E}\left(e^{uX(t)}\right) = 1 + \frac{1}{2}u^2 \int_0^t s^2 \mathbb{E}\left(e^{uX(s)}\right) ds.$$

Denote $h(t) = \mathbb{E}(e^{uX(t)})$, then differentiation with respect to t leads to a simple equation

$$h'(t) = \frac{1}{2}u^2 t^2 h(t), \text{ with } h(0) = 1.$$

By separating variables, $\log h(t) = \frac{1}{2}u^2 \int_0^t s^2 ds = \frac{1}{2}u^2 \frac{t^3}{3}$. Thus $h(t) = e^{\frac{1}{2}u^2 \frac{t^3}{3}}$, which corresponds to the $N(0, \frac{t^3}{3})$ distribution. Thus $X(t) = \int_0^t s dB(s)$ has $N(0, \frac{t^3}{3})$ distribution, and the result follows.

Example 6.4: We prove that $\int_0^1 B(t)dt$ has $N(0, 1/3)$ distribution (see also Example 3.6). Using integration by parts $\int_0^1 B(t)dt = B(1) - \int_0^1 t dB(t) = \int_0^1 dB(t) - \int_0^1 t dB(t) = \int_0^1 (1-t)dB(t)$, and the result follows.

6.2 Calculation of Expectations and PDEs

Results in this section provide a method for the calculation of expectations of a function or a functional of a diffusion process on the boundary. This expectation can be computed by using a solution to the corresponding partial differential equation with a given boundary condition. This connection shows that solutions to PDEs can be represented as functions (functionals) of the corresponding diffusion.

Let $X(t)$ be a diffusion satisfying the SDE for $t > s \geq 0$,

$$dX(t) = \mu(X(t), t)dt + \sigma(X(t), t)dB(t), \text{ and } X(s) = x. \quad (6.15)$$

Backward PDE and $E(g(X(T))|X(t) = x)$

We give results on $E(g(X(T))|X(t) = x)$. Observe first that $g(X(T))$ must be integrable ($E|g(X(T))| < \infty$) for this to make sense. Of course, if g is bounded, then this is true. Observe next that by the Markov property of $X(t)$,

$$E(g(X(T))|X(t)) = E(g(X(T))|\mathcal{F}_t).$$

The latter is a martingale, by Theorem 2.31. The last ingredient is Itô's formula, which connects this to the PDE. Again, care is taken for Itô's formula to produce a martingale term, for which assumptions on the function and its derivatives are needed (see Theorem 6.3). Apart from these requirements, the results are elegant and easy to derive.

Theorem 6.6: Let $f(x, t)$ solve the backward equation, with L_t given by (6.2),

$$L_t f(x, t) + \frac{\partial f}{\partial t}(x, t) = 0, \quad \text{with } f(x, T) = g(x). \quad (6.16)$$

If $f(x, t)$ satisfies the conditions of Corollary 6.4, then

$$f(x, t) = E(g(X(T))|X(t) = x). \quad (6.17)$$

PROOF. According to Corollary 6.4, $f(X(t), t)$, $s \leq t \leq T$, is a martingale. The martingale property gives

$$\mathbb{E}(f(X(T), T) | \mathcal{F}_t) = f(X(t), t).$$

On the boundary $f(x, T) = g(x)$, so that $f(X(T), T) = g(X(T))$, and

$$f(X(t), t) = \mathbb{E}(g(X(T)) | \mathcal{F}_t).$$

According to the Markov property of $X(t)$, $f(X(t), t) = \mathbb{E}(g(X(T)) | X(t))$, and the result follows. \square

It is tempting to show that the expectation $\mathbb{E}(g(X(T)) | X(t) = x) = f(x, t)$ satisfies the backward PDE, thus establishing the existence of its solutions. Assume for a moment that we can use Itô's formula with $f(X(t), t)$, then

$$f(X(t), t) = f(X(0), 0) + \int_0^t \left(L_s f + \frac{\partial f}{\partial s} \right) (X(s), s) ds + M(t),$$

where $M(t)$ is a martingale. As noted earlier, $f(X(t), t) = \mathbb{E}(g(X(T)) | \mathcal{F}_t)$ is a martingale. It follows that $\int_t^T (L_s f + \frac{\partial f}{\partial s})(X(s), s) ds$ is a martingale as a difference of two martingales. Since the integral with respect to ds is a function of finite variation, and a martingale is not, the latter can only be true if the integral is zero, implying the backward Equation (6.16).

In order to make this argument precise one needs to establish the validity of Itô's formula, i.e. smoothness of the conditional expectation. This can be seen by writing the expectation as an integral with respect to the density function,

$$f(x, t) = \mathbb{E}(g(X(T)) | X(t) = x) = \int g(y) p(y, T, x, t) dy, \quad (6.18)$$

where $p(y, T, x, t)$ is the transition probability density. So x is now in the transition function and the smoothness in x follows. For Brownian motion $p(y, T, x, t) = \frac{1}{\sqrt{T-t}} e^{-\frac{(y-x)^2}{2\sqrt{T-t}}}$ is differentiable in x (infinitely many times) and the result follows by differentiating under the integral. For other Gaussian diffusions the argument is similar. It is harder to show this in the general case (see Theorem 5.16).

Remark 6.2: Theorem 6.6 shows that any solution of the backward equation with the boundary condition given by $g(x)$ is given by the integral of g with respect to the transition probability density, Equation (6.18), affirming that the transition probability density is a fundamental solution (see Definition 5.14).

A result similar to Theorem 6.6 is obtained when zero on the right-hand side (rhs) of the backward equation is replaced by a known function $-\phi$.

Theorem 6.7: Let $f(x, t)$ solve

$$L_t f(x, t) + \frac{\partial f}{\partial t}(x, t) = -\phi(x), \quad \text{with } f(x, T) = g(x). \quad (6.19)$$

Then

$$f(x, t) = \mathbb{E} \left(\left(g(X(T)) + \int_t^T \phi(X(s)) ds \right) \middle| X(t) = x \right). \quad (6.20)$$

The proof is similar to Theorem 6.6 and is left as an exercise.

Feynman–Kac Formula

A result more general than Theorem 6.6 is given by the Feynman–Kac formula.

Theorem 6.8 (Feynman–Kac Formula): For given bounded functions $r(x, t)$ and $g(x)$ let

$$C(x, t) = \mathbb{E} \left(e^{-\int_t^T r(X(u), u) du} g(X(T)) \middle| X(t) = x \right). \quad (6.21)$$

Assume that there is a solution to

$$\frac{\partial f}{\partial t}(x, t) + L_t f(x, t) = r(x, t)f(x, t), \quad \text{with } f(x, T) = g(x). \quad (6.22)$$

Then the solution is unique and $C(x, t)$ is that solution.

PROOF. We give a sketch of the proof by using Itô’s formula coupled with solutions of a linear SDE. Take a solution to (6.22) and apply Itô’s formula

$$df(X(t), t) = \left(\frac{\partial f}{\partial t}(X(t), t) + L_t f(X(t), t) \right) dt + \frac{\partial f}{\partial x}(X(t), t)\sigma(X(t), t)dB(t).$$

The last term is a martingale term, so write it as $dM(t)$. Now use (6.22) to obtain

$$df(X(t), t) = r(X(t), t)f(X(t), t)dt + dM(t).$$

This is a linear SDE of Langevin type for $f(X(t), t)$ where $B(t)$ is replaced by $M(t)$. Integrating this SDE between t and T , and using $T \geq t$ as a time variable and t as the origin (see (5.33) in Section 5.23) we obtain

$$\begin{aligned} f(X(T), T) &= f(X(t), t)e^{\int_t^T r(X(u), u) du} \\ &\quad + e^{\int_t^T r(X(u), u) du} \int_t^T e^{\int_t^s r(X(u), u) du} dM(s). \end{aligned}$$

However, $f(X(T), T) = g(X(T))$, and rearranging, we obtain

$$g(X(T))e^{-\int_t^T r(X(u), u) du} = f(X(t), t) + \int_t^T e^{\int_t^s r(X(u), u) du} dM(s).$$

As the last term is an integral of a bounded function with respect to the martingale, it is itself a martingale with zero mean. Taking the expectation given $X(t) = x$, we obtain that $C(x, t) = f(x, t)$. For other proofs see for example Friedman (1975), and Pinsky (1995). \square

Remark 6.3: The expression $e^{-r(T-t)}\mathbb{E}(g(X(T))|X(t) = x)$ occurs in finance as a discounted expected payoff, where r is a constant. The discounting results in the term rf in the rhs of the backward PDE.

Example 6.5: Give a probabilistic representation of the solution $f(x, t)$ of the PDE

$$\frac{1}{2}\sigma^2 x^2 \frac{\partial^2 f}{\partial x^2} + \mu x \frac{\partial f}{\partial x} + \frac{\partial f}{\partial t} = rf, \quad 0 \leq t \leq T, \quad f(x, T) = x^2, \quad (6.23)$$

where σ , μ , and r are positive constants. Solve this PDE using the solution of the corresponding stochastic differential equation.

The SDE corresponding to L is $dX(t) = \mu X(t)dt + \sigma X(t)dB(t)$. Its solution is $X(t) = X(0)e^{(\mu - \sigma^2/2)t + \sigma B(t)}$. By the Feynman–Kac formula

$$f(x, t) = E\left(e^{-r(T-t)}X^2(T) \middle| X(t) = x\right) = e^{-r(T-t)}E(X^2(T)|X(t) = x).$$

Using $X(T) = X(t)e^{(\mu - \sigma^2/2)(T-t) + \sigma(B(T) - B(t))}$, we obtain
 $E(X^2(T)|X(t) = x) = x^2 e^{(2\mu + \sigma^2)(T-t)}$, giving $f(x, t) = x^2 e^{(2\mu + \sigma^2 - r)(T-t)}$.

The following result shows that $f(x, t) = \mathbb{E}(g(X(T))|X(t) = x)$ satisfies the backward PDE, and can be found in Gihman and Skorohod (1972), and Friedman (1975). (See also Theorem 6.5.3 in Friedman (1975) for $C(x, t)$.)

Theorem 6.9 (Kolmogorov's Equation): *Let $X(t)$ be a diffusion with generator L_t . Assume that the coefficients $\mu(x, t)$ and $\sigma(x, t)$ of L_t are locally Lipschitz and satisfy the linear growth condition (see (1.30)). Assume in addition that they possess continuous partial derivatives with respect to x up to order two, and that they have at most polynomial growth (see (1.31)). If $g(x)$ is twice continuously differentiable and satisfies together with its derivatives a polynomial growth condition, then the function $f(x, t) = \mathbb{E}(g(X(T))|X(t) = x)$ satisfies*

$$\frac{\partial f}{\partial t}(x, t) + L_t f(x, t) = 0, \text{ in the region } 0 \leq t < T, \quad x \in \mathbb{R}, \quad (6.24)$$

with boundary condition $f(x, T) = \lim_{t \uparrow T} f(x, t) = g(x)$.

6.3 Time-Homogeneous Diffusions

The case of time-independent coefficients in SDEs corresponds to the so-called time-homogeneous diffusions,

$$dX(t) = \mu(X(t))dt + \sigma(X(t))dB(t). \quad (6.25)$$

Theorem 6.10: *Assume that there is a unique weak solution to (6.25). Then the transition probability function of the solution $P(y, t, x, s) = P(y, t - s, x, 0)$ depends only on $t - s$.*

PROOF. Denote by (X, B) the weak solution to (6.25). According to the definition of the transition function $P(y, t, x, s) = P(X(t) \leq y | X(s) = x) = P(X_s^x(t) \leq y)$, where the process $X_s^x(t)$ satisfies $X_s^x(s) = x$ and for $t > 0$

$$X_s^x(s+t) = x + \int_s^{s+t} \mu(X_s^x(u))du + \int_s^{s+t} \sigma(X_s^x(u))dB(u). \quad (6.26)$$

Let $Y(t) = X_s^x(s+t)$, and $B_1(t) = B(s+t) - B(s)$, $t \geq 0$. Then $B_1(t)$ is a Brownian motion and from the above equation, $Y(t)$ satisfies for $t \geq 0$

$$Y(t) = x + \int_0^t \mu(Y(v))dv + \int_0^t \sigma(Y(v))dB_1(v), \quad \text{and} \quad Y(0) = x. \quad (6.27)$$

Put $s = 0$ in (6.26) to obtain

$$X_0^x(t) = x + \int_0^t \mu(X_0^x(v))dv + \int_0^t \sigma(X_0^x(v))dB(v), \quad \text{and} \quad X_0^x(0) = x. \quad (6.28)$$

Thus $Y(t)$ and $X_0^x(t)$ satisfy the same SDE. Hence $Y(t)$ and $X_0^x(t)$ have the same distribution. Therefore for $t > s$

$$\begin{aligned} P(y, t, x, s) &= P(X_s^x(t) \leq y) = P(Y(t-s) \leq y) \\ &= P(X_0^x(t-s) \leq y) = P(y, t-s, x, 0). \end{aligned}$$
□

Since the transition function of a homogeneous diffusion depends on t and s only through $t - s$, it is denoted as

$$P(t, x, y) = P(y, t+s, x, s) = P(y, t, x, 0) = P(X(t) \leq y | X(0) = x), \quad (6.29)$$

and it gives the probability for the process to go from x to $(-\infty, y)$ during time t . Its density $p(t, x, y)$, when it exists, is the density of the conditional distribution of $X(t)$ given $X(0) = x$.

The generator L of a time-homogeneous diffusion is given by

$$Lf(x) = \frac{1}{2}\sigma^2(x)f''(x) + \mu(x)f'(x). \quad (6.30)$$

Under appropriate conditions (conditions (A1) and (A2) of Theorem 5.15) $p(t, x, y)$ is the fundamental solution of the backward Equation (5.62), which becomes in this case

$$\frac{\partial p}{\partial t} = Lp = \frac{1}{2}\sigma^2(x)\frac{\partial^2 p}{\partial x^2} + \mu(x)\frac{\partial p}{\partial x}. \quad (6.31)$$

If, moreover, $\sigma(x)$ and $\mu(x)$ have derivatives, $\sigma'(x)$, $\mu'(x)$, and $\sigma''(x)$, which are bounded and satisfy a Hölder condition, then $p(t, x, y)$ satisfies the forward equation in t and y for any fixed x , which becomes

$$\frac{\partial p}{\partial t}(t, x, y) = \frac{1}{2}\frac{\partial^2}{\partial y^2}\left(\sigma^2(y)p(t, x, y)\right) - \frac{\partial}{\partial y}\left(\mu(y)p(t, x, y)\right). \quad (6.32)$$

In terms of the generator, the backward and the forward equations are written as

$$\frac{\partial p}{\partial t} = Lp, \quad \frac{\partial p}{\partial t} = L^*p, \quad (6.33)$$

where

$$(L^*f)(y) = \frac{1}{2}\frac{\partial^2}{\partial y^2}\left(\sigma^2(y)f(y)\right) - \frac{\partial}{\partial y}\left(\mu(y)f(y)\right),$$

denotes the operator appearing in Equation (6.32), and is known as the adjoint operator to L . (The adjoint operator is defined by the requirement that whenever the following integrals exist, $\int g(x)Lf(x)dx = \int f(x)L^*g(x)dx$.)

Example 6.6: The generator of Brownian motion is $L = \frac{1}{2}\frac{d^2}{dx^2} \cdot \Delta = \frac{d^2}{dx^2}$ is called the Laplacian. The backward equation for the transition probability density is

$$\frac{\partial p}{\partial t} = Lp = \frac{1}{2}\frac{\partial^2 p}{\partial x^2}. \quad (6.34)$$

Since the distribution of $B(t)$, when $B(0) = x$, is $N(x, t)$, the transition probability density is given by the density of $N(x, t)$, and is the fundamental solution of PDE (6.34)

$$p(t, x, y) = \frac{1}{\sqrt{2\pi t}}e^{-\frac{(y-x)^2}{2t}}.$$

The adjoint operator L^* is the same as L , so that L is self-adjoint. A stronger result than Theorem 6.9 holds. It is possible to show that if $\int e^{-ay^2}|g(y)|dy < \infty$ for some $a > 0$, then $f(x, t) = E_x g(B(t))$ for $t < 1/2a$ satisfies the heat equation with initial condition $f(0, x) = g(x)$, $x \in \mathbb{R}$ (see for example) Karatzas and Shreve (1988). A result of Widder (1944), states that any non-negative solution to the heat equation can be represented as $\int p(t, x, y)dF(y)$ for some non-decreasing function F .

Example 6.7: The Black–Scholes SDE

$$dX(t) = \mu X(t)dt + \sigma X(t)dB(t)$$

for constants μ and σ . The generator of this diffusion is

$$Lf(x) = \frac{1}{2}\sigma^2 x^2 f''(x) + \mu x f'(x). \quad (6.35)$$

Its density is the fundamental solution of the PDE

$$\frac{\partial p}{\partial t} = \frac{1}{2}\sigma^2 x^2 \frac{\partial^2 p}{\partial x^2} + \mu x \frac{\partial p}{\partial x}.$$

The transition probability function of $X(t)$ was found in Example 5.14. Its density is $p(t, x, y) = \frac{\partial}{\partial y} \Phi\left(\frac{\ln(y/x) - (\mu - \sigma^2/2)(t-s)}{\sigma\sqrt{t-s}}\right)$.

Itô's Formula and Martingales

If $X(t)$ is a solution of (6.25), then Itô's formula takes the form: for any twice continuously differentiable $f(x)$

$$df(X(t)) = Lf(X(t))dt + f'(X(t))\sigma(X(t))dB(t). \quad (6.36)$$

Theorem 6.2 and Theorem 6.3 for time-homogeneous diffusions become

Theorem 6.11: *Let X be a solution to SDE (6.25) with coefficients satisfying conditions of Theorem 5.4, that is, $\mu(x)$ and $\sigma(x)$ are Lipschitz and satisfy the linear growth condition $|\mu(x)| + |\sigma(x)| \leq K(1 + |x|)$. If $f(x)$ is twice continuously differentiable in x , with derivatives growing not faster than the exponential satisfying condition (6.9), then the following process is a martingale:*

$$M_f(t) = f(X(t)) - \int_0^t Lf(X(u))du. \quad (6.37)$$

Weak solutions to (6.25) are defined as solutions to the martingale problem, by requiring existence of a filtered probability space, with an adapted process $X(t)$, so that

$$f(X(t)) - \int_0^t Lf(X(u))du \quad (6.38)$$

is a martingale for any twice continuously differentiable f vanishing outside a finite interval (see Chapter 5.8). Equation (6.38) also allows us to identify generators.

Remark 6.4: The concept of the generator is a central concept in studies of Markov processes. The generator of a time-homogeneous Markov process (not necessarily a diffusion process) is a linear operator defined by

$$Lf(x) = \lim_{t \rightarrow 0} \frac{E(f(X(t))|X_0 = x) - f(x)}{t}. \quad (6.39)$$

If the above limit exists, we say that f is in the domain of the generator. If $X(t)$ solves (6.25) and f is bounded and twice continuously differentiable, then from (6.39) the generator for a diffusion is obtained. This can be seen by interchanging the limit and the expectation (dominated convergence), using Taylor's formula. Generators of pure jump processes, such as birth–death processes, are given later. For the theory of construction of Markov processes from their generators and their studies see, for example, Dynkin (1965), Ethier and Kurtz (1986), Stroock and Varadhan (1979), and Rogers and Williams (1990).

The result on existence and uniqueness of weak solutions (Theorem 5.11) becomes

Theorem 6.12: *If $\sigma(x)$ is positive and continuous, and for any $T > 0$ there is K_T such that for all $x \in \mathbb{R}$*

$$|\mu(x)| + |\sigma(x)| \leq K_T(1 + |x|), \quad (6.40)$$

then there exists a unique weak solution to SDE (6.25) starting at any point $x \in \mathbb{R}$; moreover, the solution has the strong Markov property.

The following result is specific for one-dimensional homogeneous diffusions and does not carry over to higher dimensions.

Theorem 6.13 (Engelbert–Schmidt): *The SDE*

$$dX(t) = \sigma(X(t))dB(t)$$

has a weak solution for every initial value $X(0)$ if, and only if, for all $x \in \mathbb{R}$ the condition

$$\int_{-a}^a \frac{dy}{\sigma^2(x+y)} = \infty \quad \text{for all } a > 0$$

implies $\sigma(x) = 0$. The weak solution is unique if the above condition is equivalent to $\sigma(x) = 0$.

Corollary 6.14: *If $\sigma(x)$ is bounded away from zero, then the above SDE has a unique weak solution.*

Example 6.8: According to Corollary 6.14, Tanaka's SDE, Example 5.15,

$$dX(t) = \text{sign}(X(t))dB(t), \quad X(0) = 0,$$

has a unique weak solution.

6.4 Exit Times from an Interval

The main tool for studying various properties of diffusions is the result on exit times from an interval. Define $\tau_{(a,b)}$ to be the first time the diffusion exits (a, b) , $\tau_{(a,b)} = \inf\{t > 0 : X(t) \notin (a, b)\}$. Since $X(t)$ is continuous, $X(\tau_{(a,b)}) = a$ or b . It was shown in Theorem 2.35 that τ is a stopping time; moreover, since the filtration is right-continuous, $\{\tau < t\}$ and $\{\tau \geq t\}$ are in \mathcal{F}_t for all t . In this section results on $\tau_{(a,b)}$ are given. As the interval (a, b) is fixed, denote in this section $\tau_{(a,b)} = \tau$.

The fact that the process started in (a, b) and remains in (a, b) for all $t < \tau$ allows us to construct martingales, without additional assumptions on functions and coefficients. The following important result for analyzing diffusions, which is also known as Dynkin's formula, is established first. Introduce T_a and T_b as the hitting times of a and b , $T_a = \inf\{t > 0 : X(t) = a\}$, with the convention that the infimum of an empty set is infinity. Clearly, $\tau = \min(T_a, T_b) = T_a \wedge T_b$. The next result is instrumental for obtaining properties of τ .

Theorem 6.15: *Let $X(t)$ be a diffusion with a continuous $\sigma(x) > 0$ on $[a, b]$ and $X(0) = x$, $a < x < b$. Then for any twice continuously differentiable function $f(x)$ on \mathbb{R} the following process is a martingale*

$$f(X(t \wedge \tau)) - \int_0^{t \wedge \tau} Lf(X(s))ds. \quad (6.41)$$

Consequently,

$$\mathbb{E}_x \left(f(X(t \wedge \tau)) - \int_0^{t \wedge \tau} Lf(X(s))ds \right) = f(x). \quad (6.42)$$

PROOF. Using Itô's formula and replacing t by $t \wedge \tau$

$$f(X(t \wedge \tau)) - \int_0^{t \wedge \tau} Lf(X(s))ds = f(x) + \int_0^{t \wedge \tau} f'(X(s))\sigma(X(s))dB(s). \quad (6.43)$$

Write the Itô integral as $\int_0^t I(s \leq \tau) f'(X(s))\sigma(X(s))dB(s)$. According to Theorem 2.35 $\{\tau \geq s\}$ are in \mathcal{F}_s for all s . Thus $I(s \leq \tau)$ is adapted. Now, for any $s \leq \tau$, $X(s) \in [a, b]$. Since $f'(x)\sigma(x)$ is continuous on $[a, b]$, it is bounded on $[a, b]$, say by K . Thus for any $s \leq t$, $|I(s \leq \tau) f'(X(s))\sigma(X(s))| \leq K$, and expectation $\mathbb{E} \int_0^t I(s \leq \tau) (f'(X(s))\sigma(X(s)))^2 ds < K^2 t$ is finite. Therefore the Itô integral $\int_0^t I(s \leq \tau) f'(X(s))\sigma(X(s))dB(s)$ is a martingale for $t \leq T$, and any T . Since it is a martingale it has a constant mean, and taking expectations in (6.43), formula (6.42) is obtained. \square

The next result establishes in particular that τ has a finite expectation, consequently it is finite with probability one.

Theorem 6.16: Let $X(t)$ be a diffusion with generator L with continuous $\sigma(x) > 0$ on $[a, b]$ and $X(0) = x$, $a < x < b$. Then $E_x(\tau) = v(x)$ satisfies the following differential equation

$$Lv = -1, \quad (6.44)$$

with $v(a) = v(b) = 0$.

PROOF. Take $v(x)$ satisfying (6.44). According to Theorem 6.15

$$E_x\left(v(X(t \wedge \tau)) - \int_0^{t \wedge \tau} Lv(X(s))ds\right) = v(x). \quad (6.45)$$

However, $Lv = -1$, therefore

$$E_x\left(v(X(t \wedge \tau))\right) + E_x(t \wedge \tau) = v(x), \quad (6.46)$$

and

$$E_x(t \wedge \tau) = v(x) - E_x(v(X(t \wedge \tau))). \quad (6.47)$$

$(t \wedge \tau)$ increases to τ as $t \rightarrow \infty$. Since the functions v and X are continuous, $v(X(t \wedge \tau)) \rightarrow v(X(\tau))$ as $t \rightarrow \infty$. $X(t \wedge \tau) \in (a, b)$ for any t and $v(x)$ is bounded on $[a, b]$, say by K , therefore $E_x(v(X(t \wedge \tau))) \leq K$. It follows from Equation (6.47) by Fatou's lemma that $E_x \liminf_{t \rightarrow \infty} (t \wedge \tau) \leq \liminf_{t \rightarrow \infty} E_x(t \wedge \tau) \leq 2K < \infty$. Hence τ is almost surely finite, moreover, by dominated convergence $E_x(v(X(t \wedge \tau))) \rightarrow E_x(v(X(\tau)))$. However, $X(\tau) = a$ or b , so that $v(X(\tau)) = 0$. Thus from (6.47) $E_x(\tau) = v(x)$. \square

The probability that the process reaches b before it reaches a , that is, $P_x(T_b < T_a)$, is used to obtain further properties. This probability is calculated with the help of the function $S(x)$, which is a solution to the equation

$$\frac{1}{2}\sigma^2(x)S''(x) + \mu(x)S'(x) = 0, \quad \text{or} \quad LS = 0. \quad (6.48)$$

Any solution to (6.48) is called a *harmonic function* for L . Only positive harmonic functions are of interest, and, ruling out constant solutions, it is easy to see (see Example 6.9) that for any such function

$$S'(x) = C \exp\left(-\int_{x_0}^x \frac{2\mu(s)}{\sigma^2(s)} ds\right). \quad (6.49)$$

$S'(x)$ is either positive for all x , if $C > 0$, or negative for all x if $C < 0$. Consequently, if S is not identically constant, $S(x)$ is monotone. Assume

that $\sigma(x)$ is continuous and positive, and $\mu(x)$ is continuous. Then any L -harmonic function comes from the general solution to (6.48), which is given by

$$S(x) = \int^x \exp \left(- \int^u \frac{2\mu(y)}{\sigma^2(y)} dy \right) du, \quad (6.50)$$

and involves two undetermined constants.

Example 6.9: We show that harmonic functions for L are given by (6.50). S must solve

$$\frac{1}{2} \sigma^2(x) S''(x) + \mu(x) S'(x) = 0. \quad (6.51)$$

This equation (with $h = S'$) leads to

$$h'/h = -2\mu(x)/\sigma^2(x),$$

and provided $\mu(x)/\sigma^2(x)$ is integrable,

$$S''(x) = e^{- \int^x \frac{2\mu(y)}{\sigma^2(y)} dy}.$$

Integrating again we find $S(x)$.

Theorem 6.17: Let $X(t)$ be a diffusion with generator L with continuous $\sigma(x) > 0$ on $[a, b]$. Let $X(0) = x$, $a < x < b$. Then

$$P_x(T_b < T_a) = \frac{S(x) - S(a)}{S(b) - S(a)}, \quad (6.52)$$

where $S(x)$ is given by (6.50).

PROOF. According to Theorem 6.15

$$E_x \left(S(X(t \wedge \tau)) - \int_0^{t \wedge \tau} LS(X(s)) ds \right) = S(x). \quad (6.53)$$

However, $LS = 0$, therefore

$$E_x(S(X(t \wedge \tau))) = S(x). \quad (6.54)$$

Since τ is finite it takes values T_b with probability $P_x(T_b < T_a)$ and T_a with the complementary probability. It is not a bounded stopping time, but by taking the limit as $t \rightarrow \infty$, we can assert by dominated convergence, that

$$ES(X(\tau)) = ES(X(0)) = S(x).$$

Expanding the expectation on the left and rearranging gives the result. \square

Remark 6.5: Note that the ratio in (6.52) remains the same no matter what non-constant solution $S(x)$ to Equation (6.50) is used.

Note that although the proof is given under the assumption of continuous drift $\mu(x)$, the result holds true for $\mu(x)$ bounded on finite intervals (see for example Pinsky (1995)).

Theorem 6.17 has a number of far reaching corollaries.

Corollary 6.18: Let $X(t)$ be a diffusion with zero drift on (a, b) , $X(0) = x$, $a < x < b$. Then

$$P_x(T_b < T_a) = \frac{x - a}{b - a}. \quad (6.55)$$

PROOF. If $\mu(x) = 0$ on (a, b) , then $S(x)$ is a linear function on (a, b) and the result follows from (6.52). \square

Diffusion $S(X(t))$ has zero drift by Itô's formula and (6.48). The exit probabilities from an interval are proportional to the distances from the end points. This explains why the function $S(x)$ is called the *scale* function. The diffusion $S(X(t))$ is said to be on the *natural scale*.

Example 6.10: We specify $P_x(T_b < T_a)$ for Brownian motion and Ornstein–Uhlenbeck processes. Brownian motion is in natural scale, since $\mu(x) = 0$. Thus $S(x) = x$, and $P_x(T_b < T_a) = (x - a)/(b - a)$.

For Ornstein–Uhlenbeck process with parameters $\mu(x) = -\alpha x$, $\sigma^2(x) = \sigma^2$,

$$S(x) = \int_a^x \exp\left(\frac{\alpha}{\sigma^2} y^2\right) dy.$$

Consequently,

$$P_x(T_b < T_a) = \frac{\int_a^x \exp\left(\frac{\alpha}{\sigma^2} y^2\right) dy}{\int_a^b \exp\left(\frac{\alpha}{\sigma^2} y^2\right) dy}.$$

Under standard assumptions there is a positive probability for the diffusion process to reach any point from any starting point.

Corollary 6.19: Let $X(t)$ be a diffusion satisfying the assumptions of Theorem 6.17. Then for any $x, y \in (a, b)$

$$P_x(T_y < \infty) > 0. \quad (6.56)$$

Indeed, for $x < y$, $T_y < T_b$, and $P_x(T_y < \infty) > P_x(T_b < \infty) > P_x(T_b < T_a) > 0$, and similarly for $y < x$.

As an application of the properties of the scale function, a better result for the existence and uniqueness of strong solutions is obtained. The transformation $Y(t) = S(X(t))$ results in a diffusion with no drift which allows us to waive assumptions on the drift.

Theorem 6.20 (Zvonkin): Suppose that $\mu(x)$ is bounded and $\sigma(x)$ is Lipschitz and is bounded away from zero. Then the strong solution exists and is unique. In particular, any SDE of the form

$$dX(t) = \mu(X(t))dt + \sigma dB(t), \quad \text{and} \quad X(0) = x_0, \quad (6.57)$$

with any bounded $\mu(x)$ and constant σ has a unique strong solution.

PROOF. If $Y(t) = S(X(t))$, then $dY(t) = \sigma(X(t))S'(X(t))dB(t)$. Notice that $S(x)$ is strictly increasing, therefore $X(t) = h(Y(t))$, where h is the inverse to S . Thus SDE for $Y(t)$ is $dY(t) = \sigma(h(Y(t)))S'(h(Y(t)))dB(t) = \sigma_Y(Y(t))dB(t)$. The rest of the proof consists of verification that $\sigma_Y(x) = \sigma(h(x))S'(h(x))$ is locally Lipschitz and, under the stated assumptions, satisfies conditions of the existence and uniqueness result. \square

6.5 Representation of Solutions of ODEs

Solutions to some PDEs have stochastic representations. Such representations are given by Theorems 6.6 and 6.8. Here we show that if a solution to an ODE satisfying a given boundary condition exists, then it has a representation as the expectation of a diffusion process stopped at the boundary.

Theorem 6.21: Let $X(t)$ be a diffusion with generator L with time-independent coefficients, $L = \frac{1}{2}\sigma^2(x)\frac{d^2}{dx^2} + \mu(x)\frac{d}{dx}$, continuous $\sigma(x) > 0$ on $[a, b]$, and $X(0) = x$, $a < x < b$. If f is twice continuously differentiable in (a, b) , and continuous on $[a, b]$ and solves

$$Lf = -\phi \text{ in } (a, b), \quad f(a) = g(a), \quad f(b) = g(b) \quad (6.58)$$

for some bounded functions g and ϕ , then f has the representation

$$f(x) = \mathbb{E}_x(g(X(\tau))) + \mathbb{E}_x\left(\int_0^\tau \phi(X(s))ds\right), \quad (6.59)$$

where τ is the exit time from (a, b) . In particular if $\phi \equiv 0$, the solution of (6.58) is given by

$$f(x) = \mathbb{E}_x(g(X(\tau))). \quad (6.60)$$

PROOF. The proof is immediate from Theorem 6.15. Indeed, by (6.42)

$$\mathbb{E}_x\left(f(X(t \wedge \tau)) - \int_0^{t \wedge \tau} Lf(X(u))du\right) = f(x).$$

Since τ is finite, by taking limits as $t \rightarrow \infty$ by dominated convergence

$$\mathbb{E}_x(f(X(\tau))) = f(x) + \mathbb{E}_x\left(\int_0^\tau Lf(X(u))du\right).$$

However, $Lf(X(u)) = -\phi(X(u))$ for any $u < \tau$, and $X(\tau)$ is in the boundary $\{a, b\}$, where $f(x) = g(x)$, and the result follows. \square

Example 6.11: Let $X = B$ be Brownian motion. Consider the solution of the problem

$$\frac{1}{2}f''(x) = 0 \text{ in } (a, b), \quad f(a) = 0, \quad f(b) = 1.$$

Here we solve this problem directly and verify the result of Theorem 6.21. Clearly, the solution is a linear function, and from boundary conditions it follows that it must be $f(x) = (x - a)/(b - a)$. According to (6.60) this solution has the representation

$$\begin{aligned} f(x) &= \mathbb{E}_x(g(B(\tau))) \\ &= g(a)\mathbb{P}_x(T_a < T_b) + g(b)(1 - \mathbb{P}_x(T_a < T_b)) = \mathbb{P}_x(T_b < T_a). \end{aligned}$$

As we know, for Brownian motion, $\mathbb{P}_x(T_b < T_a) = (x - a)/(b - a)$, and the result is verified.

6.6 Explosion

Explosion refers to the situation when the process reaches infinite values in finite time. For example, the function $1/(1-t)$, $t < 1$, explodes at $t = 1$. Similarly, the solution $x(t)$ to the ordinary differential equation

$$dx(t) = (1 + x^2(t))dt, \quad x(0) = 0$$

explodes. Indeed, consider $x(t) = \tan(t)$, which approaches infinity as t approaches $\pi/2$. The time of explosion is $\pi/2$. Similar situations can occur with solutions to SDEs, except the time of explosion will be random. Solutions can be considered until the time of explosion.

Let diffusion $X(t)$ satisfy SDE on \mathbb{R}

$$dX(t) = \mu(X(t))dt + \sigma(X(t))dB(t), \quad \text{and} \quad X(0) = x. \quad (6.61)$$

Let $D_n = (-n, n)$ for $n = 1, 2, \dots$, and $\tau_n = \tau_{D_n}$ be the first time the process has absolute value n . Since a diffusion process is continuous, it must reach level n before it reaches level $n+1$. Therefore τ_n are non-decreasing, hence they converge to a limit $\tau_\infty = \lim_{n \rightarrow \infty} \tau_n$. Explosion occurs on the set $\{\tau_\infty < \infty\}$, because on this set, by continuity of $X(t)$, $X(\tau_\infty) = \lim_{n \rightarrow \infty} X(\tau_n)$. Thus $|X(\tau_\infty)| = \lim_{n \rightarrow \infty} |X(\tau_n)| = \lim_{n \rightarrow \infty} n = \infty$, and infinity is reached in finite time on this set.

Definition 6.22: Diffusion started from x explodes if $P_x(\tau_\infty < \infty) > 0$.

Note that under appropriate conditions on the coefficients, if diffusion explodes when started at some x_0 , then it explodes when started at *any* $x \in \mathbb{R}$. Indeed, if for any x, y , $P_x(T_y < \infty) > 0$ (see Corollary 6.19), then $P_y(\tau_\infty < \infty) \geq P_y(T_x < \infty)P_x(\tau_\infty < \infty) > 0$.

The result below gives the necessary and sufficient conditions for explosions. It is known as Feller's test for explosions.

Theorem 6.23: Suppose $\mu(x)$, $\sigma(x)$ are bounded on finite intervals, and $\sigma(x) > 0$ and is continuous. Then the diffusion process explodes if, and only if, one of the two following conditions holds. There exists x_0 such that

1. $\int_{-\infty}^{x_0} \exp\left(-\int_{x_0}^x \frac{2\mu(s)}{\sigma^2(s)} ds\right) \left(\int_x^{x_0} \frac{\exp\left(\int_{x_0}^y \frac{2\mu(s)}{\sigma^2(s)} ds\right)}{\sigma^2(y)} dy \right) dx < \infty.$
2. $\int_{x_0}^{\infty} \exp\left(-\int_{x_0}^x \frac{2\mu(s)}{\sigma^2(s)} ds\right) \left(\int_{x_0}^x \frac{\exp\left(\int_{x_0}^y \frac{2\mu(s)}{\sigma^2(s)} ds\right)}{\sigma^2(y)} dy \right) dx < \infty.$

The proof relies on the analysis of exit times τ_n with the aid of Theorem 6.16 and the Feynman–Kac formula (see for example Gihman and Skorohod (1972) and Pinsky (1995)).

If the drift coefficient $\mu(x) \equiv 0$, then both conditions in Theorem 6.23 fail, since $\int_x^{x_0} \sigma^{-2}(y) dy \not\rightarrow 0$ as $x \rightarrow -\infty$, hence the following result.

Corollary 6.24: SDEs of the form $dX(t) = \sigma(X(t))dB(t)$ do not explode.

Example 6.12: Consider the SDE $dX(t) = cX^r(t)dt + dB(t)$, $c > 0$. Solutions of $dx(t) = cx^r(t)dt$ explode if, and only if, $r > 1$ (see Exercise 6.11). Here, $\sigma(x) = 1$ and $\mu(x) = cx^r$, $c > 0$, and $D = (\alpha, \beta) = (0, \infty)$. It is clear that this diffusion drifts to $+\infty$ due to the positive drift for any $r > 0$. However, explosion occurs only in the case of $r > 1$, that is, $P_x(\tau_D < \infty) > 0$ if $r > 1$, and $P_x(\tau_D < \infty) = 0$ if $r \leq 1$. The integral in condition 2 of Theorem 6.23 is

$$\int_{x_0}^{\infty} \frac{\int_{x_0}^x \exp\left(\frac{2c}{r+1} y^{r+1}\right) dy}{\exp\left(\frac{2c}{r+1} x^{r+1}\right)} dx.$$

Using the l'Hôpital rule, it can be seen that the function under the integral is of order x^{-r} as $x \rightarrow \infty$. Since $\int_{x_0}^{\infty} x^{-r} dx < \infty$ if, and only if, $r > 1$, the result is established (see Pinsky (1995)).

Example 6.13: Consider the SDE $dX(t) = X^2(t)dt + X^r(t)dB(t)$. Using the integral test it can be seen that if $r < 3/2$ there is no explosion, and if $r > 3/2$ there is an explosion.

6.7 Recurrence and Transience

Let $X(t)$ be a diffusion on \mathbb{R} . There are various definitions of recurrence and transience in the literature, however, under the imposed assumptions on the coefficients they are all equivalent.

Definition 6.25: A point x is called recurrent for diffusion $X(t)$ if the probability of the process coming back to x infinitely often is one, that is,

$$P_x(X(t) = x \text{ for a sequence of } t\text{'s increasing to infinity}) = 1.$$

Definition 6.26: A point x is called transient for diffusion $X(t)$ if

$$P_x(\lim_{t \rightarrow \infty} |X(t)| = \infty) = 1.$$

If all points of a diffusion are recurrent, the diffusion itself is called recurrent.

If all points of a diffusion are transient, the diffusion itself is called transient.

Theorem 6.27: Let $X(t)$ be a diffusion on \mathbb{R} satisfying the assumptions of the existence and uniqueness result Theorem 5.11, that is, $\mu(x)$ and $\sigma(x)$ are bounded on finite intervals, $\sigma(x)$ is continuous and positive, and $\mu(x)$, $\sigma(x)$ satisfy the linear growth condition. Then

1. If there is one recurrent point, then all points are recurrent.
2. If there are no recurrent points, then the diffusion is transient.

In order to prove this result two fundamental properties of diffusions are used. The first is the strong Markov property, and the second is the *strong Feller* property, which states that for any bounded function $f(x)$, $E_x f(X(t))$ is a continuous function in x for any $t > 0$. Both these properties hold under the stated conditions. It can also be seen that the recurrence is equivalent to the property: for any x, y $P_x(T_y < \infty) = 1$, where T_y is the hitting time of y . According to Theorem 6.27, transience is equivalent to the property: for any x, y $P_x(T_y < \infty) < 1$ (see for example Pinsky (1995)). In order to decide whether $P_x(T_y < \infty) < 1$, the formula (6.52) for the probability of exit from one end of an interval in terms of the scale function is used. If a diffusion does not explode, then the hitting time of infinity is defined as $T_\infty = \lim_{b \rightarrow \infty} T_b = \infty$ and $T_{-\infty} = \lim_{a \rightarrow -\infty} T_a = \infty$. Remember, that $S(x) = \int_{x_0}^x \exp\left(-\int_{x_0}^u \frac{2\mu(s)}{\sigma^2(s)} ds\right) du$, and that by (6.52) $P_x(T_b < T_a) = (S(x) - S(a))/(S(b) - S(a))$. Take any $y > x$, then

$$P_x(T_y < \infty) = \lim_{a \rightarrow -\infty} P_x(T_y < T_a) = \lim_{a \rightarrow -\infty} \frac{S(x) - S(a)}{S(y) - S(a)}. \quad (6.62)$$

Thus if $S(-\infty) = \lim_{a \rightarrow -\infty} S(a) = \infty$, then $P_x(T_y < \infty) = 1$.

Similarly, for $y < x$,

$$P_x(T_y < \infty) = \lim_{b \rightarrow \infty} P_x(T_y < T_b) = \lim_{b \rightarrow \infty} \frac{S(x) - S(b)}{S(y) - S(b)}.$$

Thus if $S(\infty) = \lim_{b \rightarrow \infty} S(b) = \infty$, then $P_x(T_y < \infty) = 1$. Thus for any y , $P_x(T_y < \infty) = 1$, which is the recurrence property.

If one of the values $S(-\infty)$ or $S(\infty)$ is finite, then for some y , $P_x(T_y < \infty) < 1$, which is the transience property. Thus the necessary and sufficient conditions for recurrence and transience are given by

Theorem 6.28: Let operator $L = \frac{1}{2}\sigma^2(x)\frac{d^2}{dx^2} + \mu(x)\frac{d}{dx}$ have coefficients that satisfy assumptions of Theorem 6.27. Denote for a fixed x_0 ,

$$I_1 = \int_{-\infty}^{x_0} \exp\left(-\int_{x_0}^u \frac{2\mu(s)}{\sigma^2(s)} ds\right) du \text{ and } I_2 = \int_{x_0}^{\infty} \exp\left(-\int_{x_0}^u \frac{2\mu(s)}{\sigma^2(s)} ds\right) du.$$

The diffusion corresponding to L is recurrent if, and only if, both I_1 and I_2 are infinite, and transient otherwise, that is, when one of I_1 or I_2 is finite.

6.8 Diffusion on an Interval

Consider diffusion on an interval (α, β) , where one of the ends is finite. The main difference between this case and diffusion on the whole line is that the finite end of the interval may be attained in finite time, the situation analogous to explosion in the former case. Take α finite, that is $-\infty < \alpha < \beta \leq \infty$. The case of finite β is similar. Write the scale function in the form

$$S(x) = \int_{x_1}^x \exp\left(-\int_{x_0}^u \frac{2\mu(s)}{\sigma^2(s)} ds\right) du, \quad (6.63)$$

where $x_0, x_1 \in (\alpha, \beta)$. By using stopping of $S(X(t))$, a martingale is obtained, and by the same argument as before the probabilities of exit from an interval $(a, b) \subset (\alpha, \beta)$ are given by (6.52)

$$P_x(T_a < T_b) = (S(b) - S(x))/(S(b) - S(a)).$$

If $S(\alpha) = -\infty$, then the above probability can be made arbitrarily small by taking $a \rightarrow \alpha$. This means that $P_x(T_\alpha < T_b) = 0$, and the boundary α is not attained before b for any b . If $S(\alpha) > -\infty$, then $P_x(T_\alpha < T_b) > 0$. A result similar to Theorem 6.28 holds.

Theorem 6.29: Let $L_1 = \int_{\alpha}^b \exp\left(-\int_b^u \frac{2\mu(s)}{\sigma^2(s)} ds\right) du$. If $L_1 = \infty$, then the diffusion attains the point b before α , for any initial point $x \in (\alpha, b)$. If $L_1 < \infty$, then let $L_2 = \int_{\alpha}^b \frac{1}{\sigma^2(y)} \int_{\alpha}^y \exp\left(-\int_b^x \frac{2\mu(s)}{\sigma^2(s)} ds\right) dx \exp\left(\int_b^y \frac{2\mu(s)}{\sigma^2(s)} ds\right) dy$.

1. If $L_2 < \infty$, then for all $x \in (\alpha, b)$ the diffusion exits (α, b) in finite time; moreover, $P_x(T_\alpha < \infty) > 0$.
2. If $L_2 = \infty$, then either the exit time of (α, b) is infinite and $\lim_{t \rightarrow \infty} X(t) = \alpha$, or the exit time of (α, b) is finite and $P_x(T_b < T_\alpha) = 1$.

Example 6.14: Consider a diffusion given by the SDE

$$dX(t) = ndt + 2\sqrt{X(t)}dB(t), \quad (6.64)$$

where n is a positive integer. It will be seen later that $X(t)$ is the squared distance from the origin of the Brownian motion in n dimensions (see (6.82)). If T_0 is the first visit of zero, we show that if $n \geq 2$, then $P_x(T_0 = \infty) = 1$. This means that $X(t)$ never visits zero, that is, $P(X(t) > 0, \text{ for all } t \geq 0) = 1$. However, for $n = 1$ $P_x(T_0 < \infty) = 1$. For $n = 2$ the scale function $S(x)$ is given by $S(x) = \ln x$, so that for any $b > 0$ $P_1(T_0 < T_b) = (S(1) - S(b))/(S(0) - S(b)) = 0$, hence $P_1(T_0 < \infty) = 0$. For $n \geq 3$, the scale function $S(x) = (1 - x^{-n/2+1})/(1 - n/2)$. Therefore $P_1(T_0 < \infty) = (S(1) - S(\infty))/(S(0) - S(\infty)) = 0$. Thus for any $n \geq 2$, $P_1(T_0 = \infty) = 1$. Directly, $\alpha = 0$, by Theorem 6.29 $L_1 = \infty$, and the result follows. When $n = 1$ calculations show that $L_1 < \infty$, and also $L_2 < \infty$, thus $P_1(T_0 < \infty) > 0$.

Remark 6.6: There is a classification of boundary points depending on the constants L_1 , L_2 , and L_3 , where $L_3 = \int_\alpha^b \frac{1}{\sigma^2(y)} \exp \left(\int_{x_0}^y \frac{2\mu(s)}{\sigma^2(s)} ds \right) dy$. The boundary α is called

1. Natural, if $L_1 = \infty$.
2. Attracting, if $L_1 < \infty$, $L_2 = \infty$.
3. Absorbing, if $L_1 < \infty$, $L_2 < \infty$, $L_3 = \infty$.
4. Regular, if $L_1 < \infty$, $L_2 < \infty$, $L_3 < \infty$.

See for example Gihman and Skorohod (1972).

6.9 Stationary Distributions

Consider the diffusion process given by the SDE

$$dX(t) = \mu(X(t))dt + \sigma(X(t))dB(t),$$

with $X(0)$ having a distribution $\nu(x) = P(X(0) \leq x)$. The distribution $\nu(x)$ is called *stationary* or *invariant* for the diffusion process $X(t)$ if for any t the distribution of $X(t)$ is the same as $\nu(x)$. If $P(t, x, y)$ denotes the transition probability function of the process $X(t)$, that is, $P(t, x, y) = P(X(t) \leq y | X(0) = x)$,

then an invariant $\nu(x)$ satisfies

$$\nu(y) = \int P(t, x, y) d\nu(x). \quad (6.65)$$

In order to justify (6.65) use the total probability formula and the fact that the stationary distribution is the distribution of $X(t)$ for all t ,

$$P(X_0 \leq y) = P(X_t \leq y) = \int P(X_t \leq y | X_0 = x) d\nu(x). \quad (6.66)$$

If the stationary distribution has a density, $\pi(x) = d\nu(x)/dx$, then $\pi(x)$ is called a stationary or invariant density. If $p(t, x, y) = \partial P(t, x, y)/\partial y$ denotes the density of $P(t, x, y)$, then a stationary π satisfies

$$\pi(y) = \int p(t, x, y) \pi(x) dx. \quad (6.67)$$

Under appropriate conditions on the coefficients (μ and σ are twice continuously differentiable with second derivatives satisfying a Hölder condition) an invariant density exists if, and only if, the following two conditions hold:

1. $\int_{-\infty}^{x_0} \exp\left(-\int_{x_0}^x \frac{2\mu(s)}{\sigma^2(s)} ds\right) dx = \int_{x_0}^{\infty} \exp\left(-\int_x^{x_0} \frac{2\mu(s)}{\sigma^2(s)} ds\right) dx = \infty.$
2. $\int_{-\infty}^{\infty} \frac{1}{\sigma^2(x)} \exp\left(\int_{x_0}^x \frac{2\mu(s)}{\sigma^2(s)} ds\right) dx < \infty.$

Furthermore, if an invariant density is twice continuously differentiable, then it satisfies the ordinary differential equation

$$L^* \pi = 0, \text{ that is, } \frac{1}{2} \frac{\partial^2}{\partial y^2} \left(\sigma^2(y) \pi \right) - \frac{\partial}{\partial y} \left(\mu(y) \pi \right) = 0. \quad (6.68)$$

Moreover, any solution of this equation with finite integral defines an invariant probability density. In order to see this, apply Dynkin's formula to a function f that vanishes outside a compact interval,

$$Ef(X(t)) = Ef(X(0)) + \int_0^t ELf(X(s)) ds.$$

It follows by arbitrariness of t that $ELf(X(t)) = \int Lf(x) \pi(x) dx = 0$. Using integration by parts, we obtain that $\int f(x) L^* \pi(x) dx = 0$. Now, by arbitrariness of f it follows that $L^* \pi = 0$.

For a rigorous proof see, for example, Pinsky (1995). Another heuristic justification of Equation (6.68) comes from the forward equation. Remember, that under appropriate conditions the density of $X(t)$ satisfies

the forward (Fokker–Planck) Equation (5.63). If the system is in a stationary regime, its distribution does not change with time, which means that the derivative of the density with respect to t is zero, resulting in Equation (6.68).

Equation (6.68) can be solved, as it can be reduced to a first order differential equation (see (1.36)). Using the integrating factor $\exp\left(-\int_a^x \frac{2\mu(y)}{\sigma^2(y)} dy\right)$, we find that the solution is given by

$$\pi(x) = \frac{C}{\sigma^2(x)} \exp\left(\int_{x_0}^x \frac{2\mu(y)}{\sigma^2(y)} dy\right), \quad (6.69)$$

where C is found from $\int \pi(x) dx = 1$.

Example 6.15: For Brownian motion, condition 1 above is true, but condition 2 fails. Thus no stationary distribution exists. The forward equation for the invariant distribution is

$$\frac{1}{2} \frac{\partial^2 p}{\partial x^2} = 0,$$

which has for its solutions linear functions of x , and none of these has a finite integral.

Example 6.16: The forward equation for the Ornstein–Uhlenbeck process is

$$\frac{\partial p}{\partial t} = L^* p = \frac{1}{2} \sigma^2 \frac{\partial^2 p}{\partial x^2} - \alpha \frac{\partial}{\partial x}(xp),$$

and the equation for the invariant density is

$$\frac{1}{2} \sigma^2 \frac{\partial^2 p}{\partial x^2} - \alpha \frac{\partial}{\partial x}(xp) = 0.$$

The solution is given by

$$\pi(x) = \frac{C}{\sigma^2} \exp\left(\int_0^x \frac{-2\alpha y}{\sigma^2} dy\right) = \frac{C}{\sigma^2} \exp\left(-\frac{\alpha}{\sigma^2} x^2\right). \quad (6.70)$$

This shows that if α is negative, no stationary distribution exists, and if α is positive, then the stationary density is normal $N(0, \sigma^2/(2\alpha))$. The fact that $N(0, \sigma^2/(2\alpha))$ is a stationary distribution can be easily verified directly from representation (5.13).

Remark 6.7: The Ornstein–Uhlenbeck process has the following properties: it is a Gaussian process with continuous paths, it is Markov, and it is stationary, provided the initial distribution is the stationary distribution $N(0, \frac{\sigma^2}{2\alpha})$. Stationarity means that finite-dimensional distributions do not change with shifts in time. For Gaussian processes stationarity is equivalent to the covariance function to be a function of $|t - s|$ only, i.e. $\text{Cov}(X(t), X(s)) = h(|t - s|)$ (see Exercise (6.3)). The Ornstein–Uhlenbeck process is the only process that is simultaneously Gaussian, Markov, and stationary (see for example Breiman (1968)).

Invariant Measures

A measure ν is called invariant for $X(t)$ if it satisfies the equation $\nu(B) = \int_{-\infty}^{\infty} P(t, x, B)d\nu(x)$ for all intervals B . In Equation (6.65) intervals of the form $B = (-\infty, y]$ were used. The general equation reduces to (6.65) if $C = \nu(\mathbb{R}) < \infty$. In this case ν can be normalized to a probability distribution. If $C = \infty$ this is impossible. Densities of invariant measures, when they exist and are smooth enough, satisfy Equation (6.67). Conversely, any positive solution to (6.67) is a density of an invariant measure.

For Brownian motion $\pi(x) = 1$ is a solution of the Equation (6.67). This is seen as follows. Since $p(t, x, y)$ is the density of the $N(x, t)$ distribution, $p(t, x, y) = \frac{1}{\sqrt{2\pi t}} \exp(-(y-x)^2/(2t))$. Note that for a fixed y , as a function of x , it is also the density of the $N(y, t)$ distribution. Therefore it integrates to unity, $\int_{\mathbb{R}} p(t, x, y)dx = 1$. Thus $\pi(x) = 1$ is a positive solution of the Equation (6.67). In this case the density 1 corresponds to the Lebesgue measure, which is an invariant measure for Brownian motion. Since $\int_{\mathbb{R}} 1dx = \infty$, it cannot be normalized to a probability density. Note also that since the mean of the $N(y, t)$ distribution is y , we have

$$\int_{-\infty}^{\infty} xp(t, x, y)dx = y.$$

So that $\pi(x) = x$ is also a solution of the Equation (6.67), but it is not a positive solution, and therefore is not a density of an invariant measure.

An interpretation of the invariant measure which is not a probability measure may be given by the density of a large number (infinite number) of particles with locations corresponding to the invariant measure, all diffusing according to the diffusion equation. Then at any time the density of the particles at any location will be preserved.

6.10 Multi-dimensional SDEs

We cover the concepts very briefly, relying on analogy with the one-dimensional case, but state the differences arising due to the increase in dimension. Let $\mathbf{X}(t)$ be a diffusion in n dimensions, described by the multi-dimensional SDE

$$d\mathbf{X}(t) = \mathbf{b}(\mathbf{X}(t), t)dt + \sigma(\mathbf{X}(t), t)d\mathbf{B}(t), \quad (6.71)$$

where σ is an $n \times d$ matrix valued function, \mathbf{B} is d -dimensional Brownian motion (see Chapter 4.7), and \mathbf{X}, \mathbf{b} are n -dimensional vector valued functions. In coordinate form this reads

$$dX_i(t) = b_i(\mathbf{X}(t), t)dt + \sum_{j=1}^d \sigma_{ij}(\mathbf{X}(t), t)dB_j(t), \quad i = 1, \dots, n, \quad (6.72)$$

and it means that for all $t > 0$ and $i = 1, \dots, n$

$$X_i(t) = X_i(0) + \int_0^t b_i(\mathbf{X}(u), u) du + \sum_{j=1}^d \int_0^t \sigma_{ij}(\mathbf{X}(u), u) dB_j(u). \quad (6.73)$$

The coefficients of the SDE are the vector $\mathbf{b}(\mathbf{x}, t)$, and the matrix $\sigma(\mathbf{x}, t)$.

An existence and uniqueness result for strong solutions, under the assumption of locally Lipschitz coefficients, holds in the same form (see Theorem 5.4), except for absolute values that should be replaced by the norms. The norm of the vector is its length, $|\mathbf{b}| = \sqrt{\sum_{i=1}^n b_i^2}$. The norm of the matrix σ is defined by $|\sigma|^2 = \text{trace}(\sigma\sigma^{Tr})$, with σ^{Tr} being the transposed of σ . The trace(a) = $\sum_{i=1}^n a_{ii}$. The matrix $a = \sigma\sigma^{Tr}$ is called the diffusion matrix.

Theorem 6.30: *If the coefficients are locally Lipschitz in \mathbf{x} with a constant independent of t , that is, for every N there is a constant K depending only on T and N such that for all $|\mathbf{x}|, |\mathbf{y}| \leq N$ and all $0 \leq t \leq T$*

$$|\mathbf{b}(\mathbf{x}, t) - \mathbf{b}(\mathbf{y}, t)| + |\sigma(\mathbf{x}, t) - \sigma(\mathbf{y}, t)| < K|\mathbf{x} - \mathbf{y}|, \quad (6.74)$$

then for any given $\mathbf{X}(0)$ the strong solution to SDE (6.71) is unique. If in addition to condition (6.74) the linear growth condition holds

$$|\mathbf{b}(\mathbf{x}, t)| + |\sigma(\mathbf{x}, t)| \leq K_T(1 + |\mathbf{x}|),$$

$\mathbf{X}(0)$ is independent of \mathbf{B} , and $E|\mathbf{X}(0)|^2 < \infty$, then the strong solution exists and is unique on $[0, T]$; moreover,

$$E\left(\sup_{0 \leq t \leq T} |\mathbf{X}(t)|^2\right) < C(1 + E|\mathbf{X}(0)|^2), \quad (6.75)$$

where constant C depends only on K and T .

Note that unlike in the one-dimensional case, the Lipschitz condition on σ can not be weakened in general to a Hölder condition, i.e. there is no Yamada–Watanabe-type result for multi-dimensional SDEs.

The quadratic covariation is easy to work out from (6.72) by taking into account that independent Brownian motions have zero quadratic covariation.

$$d[X_i, X_j](t) = dX_i(t)dX_j(t) = a_{ij}(\mathbf{X}(t), t)dt. \quad (6.76)$$

It can be shown that if $\mathbf{X}(t)$ is a solution to (6.71), then

$$\begin{aligned} E(X_i(t + \Delta) - x_i | \mathbf{X}(t) = \mathbf{x}) &= b_i(\mathbf{x}, t)\Delta + o(\Delta) \\ E\left((X_i(t + \Delta) - x_i)(X_j(t + \Delta) - x_j) | \mathbf{X}(t) = \mathbf{x}\right) &= a_{ij}(\mathbf{x}, t)\Delta + o(\Delta), \end{aligned}$$

as $\Delta \rightarrow 0$. Thus $\mathbf{b}(\mathbf{x}, t)$ is the coefficient in the infinitesimal mean of the displacement from point \mathbf{x} at time t , and $a(\mathbf{x}, t)$ is approximately the coefficient in the infinitesimal covariance of the displacement.

Weak solutions can be defined as solutions to the martingale problem. Let the operator L_t , acting on twice continuously differentiable functions from \mathbb{R}^n to \mathbb{R} , be

$$L_t = \sum_{i=1}^n b_i(\mathbf{x}, t) \frac{\partial}{\partial x_i} + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n a_{ij}(\mathbf{x}, t) \frac{\partial^2}{\partial x_i \partial x_j}. \quad (6.77)$$

Note that L_t depends on σ only through a . Then $\mathbf{X}(t)$ is a weak solution started at \mathbf{x} at time s , if

$$f(\mathbf{X}(t)) - \int_s^t (L_u f)(\mathbf{X}(u)) du \quad (6.78)$$

is a martingale for any twice continuously differentiable function f vanishing outside a compact set in \mathbb{R}^n . This process is called a diffusion with generator L_t . In the case of time-independent coefficients, the process is a time-homogeneous diffusion with generator L .

Theorem 6.31: Assume that $a(\mathbf{x}, t)$ is continuous and satisfies condition (A)

$$(A) \sum_{i,j=1}^n a_{ij}(\mathbf{x}, t) v_i v_j > 0, \text{ for all } \mathbf{x} \in \mathbb{R}^n \text{ and } \mathbf{v} \neq \mathbf{0},$$

and $\mathbf{b}(\mathbf{x}, t)$ is bounded on bounded sets. Then there exists a unique weak solution up to the time of explosion. If, in addition, the linear growth condition is satisfied, that is, for any $T > 0$ there is K_T such that for all $x \in \mathbb{R}$

$$|\mathbf{b}(\mathbf{x}, t)| + |a(\mathbf{x}, t)| \leq K_T(1 + |\mathbf{x}|), \quad (6.79)$$

then there exists a unique weak solution to the martingale problem (6.78) starting at any point $x \in \mathbb{R}$ at any time $s \geq 0$; moreover, this solution has the strong Markov property.

Since the weak solution is defined in terms of the generator, which itself depends on σ only through a , the weak solution to (6.71) can be constructed using a single Brownian motion provided the matrix a remains the same. If a single SDE is equivalent to a number of SDEs, heuristically it means that there is as much randomness in a d -dimensional Brownian motion as there is in a single Brownian motion. Replacement of a system of SDEs by a single one is shown in detail for the Bessel process.

Note that the equation $\sigma\sigma^{Tr} = a$ has many solutions for σ , the matrix square root is non-unique. However, if $a(\mathbf{x}, t)$ is non-negative definite for all \mathbf{x} and t , and has for entries twice continuously differentiable functions of \mathbf{x} and t , then it has a locally Lipschitz square root $\sigma(\mathbf{x}, t)$ of the same dimension as $a(\mathbf{x}, t)$ (see for example Friedman (1975)).

Bessel Process

Let $\mathbf{B}(t) = (B_1(t), B_2(t), \dots, B_d(t))$ be the d -dimensional Brownian motion, $d \geq 2$. Denote by $R(t)$ its squared distance from the origin, that is,

$$R(t) = \sum_{i=1}^d B_i^2(t). \quad (6.80)$$

The SDE for $R(t)$ is given by (using $d(B^2(t)) = 2B(t)dB(t) + dt$)

$$dR(t) = d dt + 2 \sum_{i=1}^d B_i(t) dB_i(t). \quad (6.81)$$

In this case we have one equation driven by d independent Brownian motions. Clearly, $b(x) = d$, $\sigma(x)$ is $(1 \times d)$ matrix $2(B_1(t), B_2(t), \dots, B_d(t))$, so that $a(X(t)) = \sigma(X(t))\sigma^{Tr}(X(t)) = 4 \sum_{i=1}^d B_i^2(t) = 4R(t)$ is a scalar. Thus the generator of $X(t)$ is given by $L = d \frac{d}{dx} + \frac{1}{2}(4x) \frac{d^2}{dx^2}$. But the same generator corresponds to the process $X(t)$ satisfying the SDE below driven by a single Brownian motion

$$dX(t) = d dt + 2\sqrt{X(t)} dB(t). \quad (6.82)$$

Therefore the squared distance process $R(t)$ in (6.80) satisfies SDE (6.82). This SDE was considered in Example 6.14. The Bessel process is defined as the distance from the origin, $Z(t) = \sqrt{\sum_{i=1}^d B_i^2(t)} = \sqrt{R(t)}$. Since $R(t)$ has the same distribution as $X(t)$ given by (6.82), by Itô's formula, $Z(t)$ satisfies

$$dZ(t) = \frac{d-1}{2Z(t)} dt + dB(t). \quad (6.83)$$

Using the one-dimensional SDE (6.82) for $R(t)$ we can decide on the recurrence, transience, and attainability of $\mathbf{0}$ of Brownian motion in dimensions two and higher. It follows from Example 6.14 that in one and two dimensions Brownian motion is recurrent, but in dimensions three and higher it is transient. It was also shown in Example 6.14 that in dimension two and above Brownian motion never visits zero. See also Karatzas and Shreve (1988).

Itô's Formula and Dynkin's Formula

Let $\mathbf{X}(t) = (X_1(t), \dots, X_n(t))$ be a diffusion in \mathbb{R}^n with generator L , (the general case is similar, but in what follows, the time-homogeneous case will be considered). Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a twice continuously differentiable (C^2)

function. Then Itô's formula states that

$$df(\mathbf{X}(t)) = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(\mathbf{X}(t))dX_i(t) + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n a_{ij}(\mathbf{X}(t)) \frac{\partial^2 f}{\partial x_i \partial x_j}(\mathbf{X}(t))dt. \quad (6.84)$$

It can be regarded as a Taylor's formula expansion where

$$dX_i dX_j \equiv d[X_i, X_j] \quad \text{and} \quad dX_i dX_j dX_k \equiv 0.$$

Itô's formula can be written with the help of the generator as

$$df(\mathbf{X}(t)) = (Lf)(\mathbf{X}(t)) + \sum_{i=1}^n \sum_{j=1}^d \frac{\partial f}{\partial x_i}(\mathbf{X}(t)) \sigma_{ij}(\mathbf{X}(t)) dB_j(t). \quad (6.85)$$

The analogues of Theorems 6.3 and 6.15 hold. It is clear from (6.85) that if partial derivatives of f are bounded, and $\sigma(\mathbf{x})$ is bounded, then

$$f(\mathbf{X}(t)) - \int_0^t Lf(\mathbf{X}(u))du \quad (6.86)$$

is a martingale. (Without the assumption of functions being bounded, it is a local martingale.)

Theorem 6.32: Suppose that the assumptions of Theorem 6.31 hold. Let $D \subset \mathbb{R}^n$ be a bounded domain (an open and simply connected set) in \mathbb{R}^n . Let $\mathbf{X}(0) = \mathbf{x}$ and denote by τ the exit time from D , $\tau_D = \inf\{t > 0 : \mathbf{X}(t) \in \partial D\}$. Then for any twice continuously differentiable f ,

$$f(\mathbf{X}(t \wedge \tau_D)) - \int_0^{t \wedge \tau_D} Lf(\mathbf{X}(u))du \quad (6.87)$$

is a martingale.

It can be shown that under conditions of Theorem 6.32

$$\sup_{\mathbf{x} \in D} E_{\mathbf{x}}(\tau_D) < \infty. \quad (6.88)$$

As a corollary the following is obtained:

Theorem 6.33: Suppose that the assumptions of Theorem 6.31 hold. If f is twice continuously differentiable in D , continuous on ∂D , and solves

$$Lf = -\phi \quad \text{in } D \quad \text{and} \quad f = g \quad \text{on } \partial D \quad (6.89)$$

for some bounded functions g and ϕ , then $f(\mathbf{x})$, $\mathbf{x} \in D$, has representation

$$f(\mathbf{x}) = \mathbb{E}_{\mathbf{x}}(g(\mathbf{X}(\tau_D))) + \mathbb{E}_{\mathbf{x}}\left(\int_0^{\tau_D} \phi(\mathbf{X}(s))ds\right). \quad (6.90)$$

In particular, if $\phi \equiv 0$, the solution has representation as

$$f(\mathbf{x}) = \mathbb{E}_{\mathbf{x}}(g(\mathbf{X}(\tau_D))).$$

The proof is exactly the same as for Theorem 6.21 in one dimension.

Definition 6.34: A function $f(\mathbf{x})$ is said to be L -harmonic on D if it is twice continuously differentiable on D and $Lf(\mathbf{x}) = 0$ for $\mathbf{x} \in D$.

The following result follows from (6.86).

Corollary 6.35: For any bounded L -harmonic function on D with bounded derivatives, $f(\mathbf{X}(t \wedge \tau_D))$ is a martingale.

Example 6.17: Denote by $\Delta = \sum_{i=1}^3 \frac{\partial^2}{\partial x_i^2}$ the three-dimensional Laplacian. This operator is the generator of three-dimensional Brownian motion $\mathbf{B}(t)$, $L = \frac{1}{2}\Delta$. Let $D = \{\mathbf{x} : |\mathbf{x}| > r\}$. Then $f(\mathbf{x}) = 1/|\mathbf{x}|$ is harmonic on D , that is, $Lf(\mathbf{x}) = 0$ for all $\mathbf{x} \in D$. In order to see this perform differentiation and verify that $\sum_{i=1}^3 \frac{\partial^2}{\partial x_i^2} \left(\frac{1}{\sqrt{x_1^2 + x_2^2 + x_3^2}} \right) = 0$ at any point $\mathbf{x} \neq \mathbf{0}$. Note that in one dimension all harmonic functions for the Laplacian are linear functions, whereas in higher dimensions there are many more. It is easy to see that $1/|\mathbf{x}|$ and its derivatives are bounded on D , consequently if $\mathbf{B}(0) = \mathbf{x} \neq \mathbf{0}$, then $1/|\mathbf{B}(t \wedge \tau_D)|$ is a martingale.

The backward (Kolmogorov's) equation in higher dimensions is the same as in one dimension, with the obvious replacement of the state variable $\mathbf{x} \in \mathbb{R}^n$. We have seen that solutions to the backward equation can be expressed by means of diffusion, Theorem 6.33. However, it is a formidable task to prove that if $\mathbf{X}(t)$ is a diffusion, and $g(\mathbf{x})$ is a smooth function on D , then $f(\mathbf{x}) = \mathbb{E}_{\mathbf{x}}(g(\mathbf{X}(t)))$ solves the backward equation (see for example Friedman (1975)).

Theorem 6.36: Let $g(\mathbf{x})$ be a function with two continuous derivatives satisfying a polynomial growth condition, that is, the function and its derivatives in absolute value do not exceed $K(1 + |\mathbf{x}|^m)$ for some constants $K, m > 0$. Let $\mathbf{X}(t)$ satisfy (6.71). Assume that coefficients $\mathbf{b}(\mathbf{x}, t)$, $\sigma(\mathbf{x}, t)$ are Lipschitz in \mathbf{x} uniformly in t , satisfy the linear growth condition, and their two derivatives satisfy a polynomial growth condition. Let

$$f(\mathbf{x}, t) = \mathbb{E}(g(\mathbf{X}(T)) | \mathbf{X}(t) = \mathbf{x}). \quad (6.91)$$

Then f has continuous derivatives in \mathbf{x} , which can be computed by differentiating (6.91) under the expectation sign. Moreover, f has a continuous derivative in t , and solves the backward PDE

$$\begin{aligned} L_t f + \frac{\partial f}{\partial t} &= 0, \text{ in } \mathbb{R}^n \times [0, T) \\ f(\mathbf{x}, T) &\rightarrow g(\mathbf{x}), \text{ as } t \uparrow T. \end{aligned} \quad (6.92)$$

The fundamental solution of (6.92) gives the transition probability function of the diffusion (6.71).

Remark 6.8 (Diffusions on Manifolds):

The PDEs above can also be considered when the state variable x belongs to a manifold, rather than \mathbb{R}^n . The fundamental solution then corresponds to the diffusion on the manifold and represents the way heat propagates on that manifold. It turns out that various geometric properties of the manifold can be obtained from the properties of the fundamental solution (Molchanov (1975)).

Remark 6.9: The Feynman–Kac formula holds also in the multi-dimensional case in the same way as in one dimension. If zero in the right hand side of the PDE (6.92) is replaced by rf , for a bounded function r , then the solution satisfying a given boundary condition $f(\mathbf{x}, T) = g(\mathbf{x})$ has a representation

$$f(\mathbf{x}, t) = \mathbb{E} \left(e^{-\int_t^T r(\mathbf{X}(u), u) du} g(\mathbf{X}(T)) \mid \mathbf{X}(t) = \mathbf{x} \right).$$

See Karatzas and Shreve (1988).

Recurrence, Transience, and Stationary Distributions

Properties of recurrence and transience of multi-dimensional diffusions, solutions to (6.71), are defined similarly to the one-dimensional case. However, in higher dimensions a diffusion $\mathbf{X}(t)$ is recurrent if for any starting point $\mathbf{x} \in \mathbb{R}^n$ the process will visit a ball around any point $\mathbf{y} \in \mathbb{R}$ of radius ϵ , $D_\epsilon(\mathbf{y})$, however small, with probability one.

$$\mathbb{P}_{\mathbf{x}}(\mathbf{X}(t) \in D_\epsilon(\mathbf{y}) \text{ for a sequence of } t \text{'s increasing to infinity}) = 1.$$

A diffusion $\mathbf{X}(t)$ on \mathbb{R}^n is transient if for any starting point $\mathbf{x} \in \mathbb{R}^n$ the process will leave any ball, however large, never to return. It follows by a diffusion analysis of the squared lengths of the multi-dimensional Brownian motion (see Example 6.14) that in dimensions one and two Brownian motion is recurrent, but it is transient in dimensions three and higher.

For time-homogeneous diffusions under conditions of Theorem 6.31 on the coefficients, recurrence is equivalent to the property that the process started

at any point \mathbf{x} hits the closed ball $\bar{D}_\epsilon(\mathbf{y})$ around any point \mathbf{y} in finite time. Under these conditions, there is a dichotomy, a diffusion is either transient or recurrent. Invariant measures are defined in exactly the same way as in one dimension. Stationary distributions are finite invariant measures; they may exist only if a diffusion is recurrent. A diffusion is recurrent and admits a stationary distribution if, and only if, the expected hitting time of $\bar{D}_\epsilon(\mathbf{y})$ from \mathbf{x} is finite. When this property holds, diffusion is also called *ergodic* or *positive recurrent*.

In general there are no necessary and sufficient conditions for recurrence and ergodicity for multi-dimensional diffusions, however, there are various tests for these properties. The method of Lyapunov functions, developed by R.Z. Hasminskii, consists of finding a suitable function f , such that $Lf \leq 0$ outside a ball around zero. If $\lim_{|\mathbf{x}| \rightarrow \infty} f(\mathbf{x}) = \infty$, then the process is transient. If f is ultimately decreasing, then the process is recurrent. If $Lf \leq -\epsilon$ for some $\epsilon > 0$ outside a ball around zero, with $f(\mathbf{x})$ bounded from below in that domain, then a diffusion is positive recurrent. Proofs consist of an application of Itô's formula coupled with the martingale theory (convergence property of supermartingales). For details see Bhattacharya (1978), Hasminskii (1980), and Pinsky (1995).

Higher Order Random Differential Equations

Similarly to ODEs higher order random differential equations have interpretations as multi-dimensional SDEs. For example, a second order random differential equation of the form

$$\ddot{x} + h(x, \dot{x}) = \dot{B}, \quad (6.93)$$

where $\dot{x}(t) = dx(t)/dt$, $\ddot{x}(t) = d^2x(t)/dt^2$, and \dot{B} denotes the white noise, has interpretation as the following two-dimensional SDE by letting

$$x_1(t) = x(t), \quad (6.94)$$

$$x_2(t) = dx_1(t)/dt. \quad (6.95)$$

$$dX_1(t) = X_2(t)dt, \quad (6.96)$$

$$dX_2(t) = -h(X_1(t), X_2(t)) + dB(t). \quad (6.97)$$

Such equations are considered in Chapter 14.2.

Higher n -th order random equations are interpreted in a similar way: by letting $X_1(t) = X(t)$ and $dX_i(t) = X_{i+1}(t)dt$, $i = 1, \dots, n-1$, an n -dimensional SDE is obtained.

Notes: Most of the material can be found in Friedman (1975), Gihman and Skorohod (1972), Stroock and Varadhan (1979), Karatzas and Shreve (1988), Durrett (1984), Rogers and Williams (1990), and Pinsky (1995).

6.11 Exercises

Exercise 6.1: Show that for any u , $f(x, t) = \exp(ux - u^2t/2)$ solves the backward equation for Brownian motion. Take derivatives, first, second, etc., of $\exp(ux - u^2t/2)$ with respect to u , and set $u = 0$, to obtain that functions x , $x^2 - t$, $x^3 - 3tx$, $x^4 - 6tx^2 + 3t^2$, etc. also solve the backward Equation (6.13). Deduce that $B^2(t) - t$, $B(t)^3 - 3tB(t)$, $B^4(t) - 6tB^2(t) + 3t^2$ are martingales.

Exercise 6.2: Find the generator for the Ornstein–Uhlenbeck process, write the backward equation, and give its fundamental solution. Verify that it satisfies the forward equation.

Exercise 6.3: Let $X(t)$ be a stationary process. Show that the covariance function $\gamma(s, t) = \text{Cov}(X(s), X(t))$ is a function of $|t - s|$ only. Hint: take $k = 2$. Deduce that for Gaussian processes stationarity is equivalent to the requirements that the mean function is a constant and the covariance function is a function of $|t - s|$.

Exercise 6.4: $X(t)$ is a diffusion with coefficients $\mu(x) = cx$ and $\sigma(x) = 1$. Give its generator and show that $X^2(t) - 2c \int_0^t X^2(s)ds - t$ is a martingale.

Exercise 6.5: $X(t)$ is a diffusion with $\mu(x) = 2x$ and $\sigma^2(x) = 4x$. Give its generator L . Solve $Lf = 0$ and give a martingale M_f . Find the SDE for the process $Y(t) = \sqrt{X(t)}$, and give the generator of $Y(t)$.

Exercise 6.6: Find $f(x)$ such that $f(B(t) + t)$ is a martingale.

Exercise 6.7: $X(t)$ is a diffusion with coefficients $\mu(x, t), \sigma(x, t)$. Find a differential equation for $f(x, t)$ such that $Y(t) = f(X(t), t)$ has infinitesimal diffusion coefficient equal to 1.

Exercise 6.8: Show that the mean exit time of a diffusion from an interval, which (by Theorem 6.16) satisfies the ODE (6.44) is given by

$$v(x) = - \int_a^x 2G(y) \int_a^y \frac{ds}{\sigma^2(s)G(s)} dy + \int_a^b 2G(y) \int_a^y \frac{ds}{\sigma^2(s)G(s)} dy \frac{\int_a^x G(s)ds}{\int_a^b G(s)ds}, \quad (6.98)$$

where $G(x) = \exp \left(- \int_a^x \frac{2\mu(s)}{\sigma^2(s)} ds \right)$.

Exercise 6.9: Find $P_x(T_b < T_a)$ for Brownian motion with drift when $\mu(x) = \mu$ and $\sigma^2(x) = \sigma^2$.

Exercise 6.10: Give a probabilistic representation of the solution $f(x, t)$ of the PDE

$$\frac{1}{2} \frac{\partial^2 f}{\partial x^2} + \frac{\partial f}{\partial t} = 0, \quad 0 \leq t \leq T, \quad f(x, T) = x^2.$$

Solve this PDE using the solution of the corresponding stochastic differential equation.

Exercise 6.11: Show that the solution of the following ordinary differential equation $dx(t) = cx^r(t)dt$, $c > 0$, $x(0) = x_0 > 0$, explodes if and only if $r > 1$.

Exercise 6.12: Investigate for explosions the following process:

$$dX(t) = X^2(t)dt + \sigma X^\alpha(t)dB(t).$$

Exercise 6.13: Show that Brownian motion $B(t)$ is recurrent. Show that $B(t) + t$ is transient.

Exercise 6.14: Show that the Ornstein–Uhlenbeck process is positively recurrent. Show that the limiting distribution for the Ornstein–Uhlenbeck process (5.6) exists, and is given by its stationary distribution. Hint: the distribution of $\sigma e^{-\alpha t} \int_0^t e^{\alpha s} dB_s$ is normal, find its mean and variance, and take limits.

Exercise 6.15: Show that the square of the Bessel process $X(t)$ in (6.64) comes arbitrarily close to zero when $n = 2$, that is, $P(T_y < \infty) = 1$ for any small $y > 0$, but when $n \geq 3$, $P(T_y < \infty) < 1$.

Exercise 6.16: Let diffusion $X(t)$ have $\sigma(x) = 1$, $\mu(x) = -1$ for $x < 0$, $\mu(x) = 1$ for $x > 0$ and $\mu(0) = 0$. Show that $\pi(x) = e^{-|x|}$ is a stationary distribution for X .

Exercise 6.17: Let diffusion on (α, β) be such that the transition probability density $p(t, x, y)$ is symmetric in x and y , $p(t, x, y) = p(t, y, x)$ for all x, y , and t . Show that if (α, β) is a finite interval, then the uniform distribution is invariant for the process $X(t)$.

Exercise 6.18: Investigate for absorption at zero the following process (used as a model for interest rates, the square root model of Cox, Ingersoll, and Ross). $dX(t) = b(a - X(t))dt + \sigma \sqrt{X(t)}dB(t)$, where parameters b, a , and σ are constants.

Chapter 7

Martingales

Martingales play a central role in the modern theory of stochastic processes and stochastic calculus. Martingales constructed from a Brownian motion were considered in Chapter 3.3, and martingales arising in diffusions in Chapter 6.1. Martingales have a constant expectation, which remains the same under random stopping. Martingales converge almost surely. Stochastic integrals are martingales. These are the most important properties of martingales, which hold under some conditions.

7.1 Definitions

The main ingredient in the definition of a martingale is the concept of conditional expectation (see Chapter 2 for its definition and properties).

Definition 7.1: A stochastic process $M(t)$, where time t is continuous $0 \leq t \leq T$, or discrete $t = 0, 1, \dots, T$, adapted to a filtration $\mathbb{F} = (\mathcal{F}_t)$, is a martingale if for any t , $M(t)$ is integrable, that is, $E|M(t)| < \infty$ and for any t and s with $0 \leq s < t \leq T$,

$$E(M(t)|\mathcal{F}_s) = M(s) \text{ almost surely.} \quad (7.1)$$

$M(t)$ is a martingale on $[0, \infty)$ if it is integrable and the martingale property (7.1) holds for any $0 \leq s < t < \infty$.

Definition 7.2: A stochastic process $X(t)$, $t \geq 0$ adapted to a filtration \mathbb{F} is a supermartingale (submartingale) if $X(t)$ is integrable, and for any t and s , $0 \leq s < t$

$$E(X(t)|\mathcal{F}_s) \leq X(s), \quad (E(X(t)|\mathcal{F}_s) \geq X(s)) \text{ almost surely.}$$

If $X(t)$ is a supermartingale, then $-X(t)$ is a submartingale. The mean of a supermartingale is non-increasing with t , the mean of a submartingale is non-decreasing in t , and the mean of a martingale is constant in t . This property is used to test if a super(sub)martingale is a true martingale.

Theorem 7.3: *A supermartingale $M(t)$, $0 \leq t \leq T$, is a martingale if, and only if, $\text{EM}(T) = \text{EM}(0)$.*

PROOF. If M is a martingale, then $\text{EM}(T) = \text{EM}(0)$ follows by the martingale property with $s = 0$ and $t = T$. Conversely, suppose $M(t)$ is a supermartingale and $\text{EM}(T) = \text{EM}(0)$. If for some t and s we have a strict inequality, $\text{E}(M(t)|\mathcal{F}_s) < M(s)$ on a set of positive probability, then by taking expectations, we obtain $\text{EM}(t) < \text{EM}(s)$. Since the expectation of a supermartingale is non-increasing, $\text{EM}(T) \leq \text{EM}(t) < \text{EM}(s) \leq \text{EM}(0)$. However, this contradicts the condition of the theorem $\text{EM}(T) = \text{EM}(0)$. Thus for all t and s the inequality $\text{E}(M(t)|\mathcal{F}_s) \leq M(s)$ must be an equality almost surely. \square

We refer to Theorem 2.32 on the existence of the regular right-continuous version for supermartingales. Regular right-continuous versions of processes will be taken.

Square Integrable Martingales

A special role in the theory of integration is played by square integrable martingales.

Definition 7.4: A random variable X is square integrable if $\text{E}(X^2) < \infty$. A process $X(t)$ on the time interval $[0, T]$, where T can be infinite, is square integrable if $\sup_{t \in [0, T]} \text{E}X^2(t) < \infty$ ($\sup_{t \geq 0} \text{E}X^2(t) < \infty$), i.e. second moments are bounded.

Example 7.1:

1. Brownian motion $B(t)$ on a finite time interval $0 \leq t \leq T$ is a square integrable martingale, since $\text{E}B^2(t) = t < T < \infty$. Similarly, $B^2(t) - t$ is a square integrable martingale. They are not square integrable when $T = \infty$.
2. If $f(x)$ is a bounded and continuous function on \mathbb{R} , then Itô integrals $\int_0^t f(B(s))dB(s)$ and $\int_0^t f(s)dB(s)$ are square integrable martingales on any finite time interval $0 \leq t \leq T$. Indeed, by (4.7), an Itô integral is a martingale, and since $|f(x)| \leq K$,

$$\text{E} \left(\int_0^t f(B(s))dB(s) \right)^2 = \text{E} \left(\int_0^t f^2(B(s))ds \right) \leq K^2 t \leq K^2 T < \infty.$$

If, moreover, $\int_0^\infty f^2(s)ds < \infty$, then $\int_0^t f(s)dB(s)$ is a square integrable martingale on $[0, \infty)$.

7.2 Uniform Integrability

In order to appreciate the definition of uniform integrability of a process, recall what is meant by integrability of a random variable X . It is called integrable if $E|X| < \infty$. It is easy to see that this holds if and only if

$$\lim_{n \rightarrow \infty} E(|X|I(|X| > n)) = 0. \quad (7.2)$$

Indeed, if X is integrable, then (7.2) holds by the dominated convergence, since $\lim_{n \rightarrow \infty} |X|I(|X| > n) = 0$ and $|X|I(|X| > n) \leq |X|$. Conversely, let n be large enough for the lhs in (7.2) to be finite. Then

$$E|X| = E(|X|I(|X| > n)) + E(|X|I(|X| \leq n)) < \infty,$$

since the first term is finite by (7.2), and the second is bounded by n .

Definition 7.5: A process $X(t)$, $0 \leq t \leq T$ is called uniformly integrable if $a_{t,n} = E(|X(t)|I(|X(t)| > n))$ converges to zero as $n \rightarrow \infty$ uniformly in t , that is,

$$\lim_{n \rightarrow \infty} \sup_t E(|X(t)|I(|X(t)| > n)) = 0, \quad (7.3)$$

where the supremum is over $[0, T]$ in the case of a finite time interval and $[0, \infty)$ if the process is considered on $0 \leq t < \infty$.

Example 7.2: We show that if $X(t)$, $0 \leq t \leq T$ is uniformly integrable, then it is integrable, that is, $\sup_t E|X(t)| < \infty$. Indeed,

$$\sup_t E|X(t)| < \sup_t E(|X(t)|I(|X(t)| > n)) + n.$$

Since $X(t)$ is uniformly integrable, the first term converges to zero as $n \rightarrow \infty$, in particular, it is bounded, and the result follows.

Sufficient conditions for uniform integrability are given next.

Theorem 7.6: If the process X is dominated by an integrable random variable, $|X(t)| \leq Y$ and $E(Y) < \infty$, then it is uniformly integrable. In particular, if $E(\sup_t |X(t)|) < \infty$, then it is uniformly integrable.

PROOF. $E(|X(t)|I(|X(t)| > n)) < E(|Y|I(|Y| > n)) \rightarrow 0$, as $n \rightarrow \infty$. \square

Note that there are uniformly integrable processes (martingales) which are not dominated by an integrable random variable, so that the sufficient condition for uniform integrability $E(\sup_t |X(t)|) < \infty$ is not necessary for uniform integrability. Another sufficient condition for uniform integrability is given by the following result, see, for example, Protter (1992), and Liptser and Shiryaev (2001).

Theorem 7.7 (Vallee–Poussin): *If for some positive, increasing, convex function $G(x)$ on $(0, \infty)$ such that $\lim_{x \rightarrow \infty} G(x)/x = \infty$,*

$$\sup_{t \leq T} \mathbb{E}(G(|X(t)|)) < \infty, \quad (7.4)$$

then $X(t)$, $t \leq T$ is uniformly integrable.

We omit the proof. In practice, the above result is used with $G(x) = x^r$ for $r > 1$, and uniform integrability is checked by using moments. For second moments $r = 2$, we have: square integrability implies uniform integrability.

Corollary 7.8: *If $X(t)$ is square integrable, that is, $\sup_t \mathbb{E}X^2(t) < \infty$, then it is uniformly integrable.*

In view of this, examples of uniformly integrable martingales are provided by square integrable martingales given in Example 7.1. The following result provides a construction of uniformly integrable martingales.

Theorem 7.9 (Doob's, Levy's Martingale): *Let Y be an integrable random variable, that is, $\mathbb{E}|Y| < \infty$, and define*

$$M(t) = \mathbb{E}(Y|\mathcal{F}_t). \quad (7.5)$$

Then $M(t)$ is a uniformly integrable martingale.

PROOF. It is easy to see that $M(t)$ is a martingale. Indeed, by the law of double expectation, $\mathbb{E}(M(t)|\mathcal{F}_s) = \mathbb{E}(\mathbb{E}(Y|\mathcal{F}_t)|\mathcal{F}_s) = \mathbb{E}(Y|\mathcal{F}_s) = M(s)$. The proof of uniform integrability is more involved. It is enough to establish the result for $Y \geq 0$, as the general case will follow by considering Y^+ and Y^- . If $Y \geq 0$, then $M(t) \geq 0$ for all t . We show next that $M^* = \sup_t M(t) < \infty$. Take a sequence of stopping times, $\tau_n = \inf\{t \geq 0 : M(t) > n\}$, or ∞ if this set is empty. On the set $\{M^* = \infty\}$, $\tau_n < \infty$ for all n , in other words, $\{M^* = \infty\} \subset \cap_n \{\tau_n < \infty\}$. Suppose that $\mathbb{P}(M^* = \infty) > 0$. Then $\mathbb{E}M(\tau_n) \geq n\mathbb{P}(\tau_n < \infty) \geq n\mathbb{P}(M^* = \infty)$ can be made arbitrarily large. However, this is a contradiction because by the rule of double expectation $\mathbb{E}M(\tau_n) = \mathbb{E}\mathbb{E}(Y|\mathcal{F}_{\tau_n}) = \mathbb{E}Y$. Next, by the general definition of conditional expectation, see (2.16), $\mathbb{E}(M(t)I(M(t) > n)) = \mathbb{E}(YI(M(t) > n))$. Since $\{M(t) > n\} \subseteq \{M^* > n\}$, $\mathbb{E}(YI(M(t) > n)) \leq \mathbb{E}(YI(M^* > n))$. Thus $\mathbb{E}(M(t)I(M(t) > n)) \leq \mathbb{E}(YI(M^* > n))$. Since the right-hand side does not depend on t , $\sup_{t \leq T} \mathbb{E}(M(t)I(M(t) > n)) \leq \mathbb{E}(YI(M^* > n))$. However, this converges to zero as $n \rightarrow \infty$, because M^* is finite and Y is integrable. \square

The martingale in (7.5) is said to be *closed* by Y . An immediate corollary is as follows.

Corollary 7.10: *Any martingale $M(t)$ on a finite time interval $0 \leq t \leq T < \infty$ is uniformly integrable and is closed by $M(T)$.*

It will be seen in the following section that a uniformly integrable martingale on $[0, \infty)$ is also of the form (7.5). That is, there exists a random variable, called $M(\infty)$, such that the martingale property holds for all $0 \leq s < t$, including $t = \infty$.

7.3 Martingale Convergence

In this section martingales on the infinite time interval $(0, \infty)$ are considered.

Theorem 7.11 (Martingale Convergence Theorem): *If $M(t)$, $0 \leq t < \infty$, is an integrable martingale (supermartingale or submartingale), that is, if $\sup_{t \geq 0} E|M(t)| < \infty$, then there exists an almost sure limit $\lim_{t \rightarrow \infty} M(t) = Y$, and Y is an integrable random variable.*

The proof of this result is due to Doob and it is too involved to be given here. If $M(t)$ is a martingale, then the condition $\sup_{t \geq 0} E|M(t)| < \infty$ is equivalent to any of the following conditions:

1. $\lim_{t \rightarrow \infty} E|M(t)| < \infty$. This is because $|x|$ is a convex function, implying that $|M(t)|$ is a submartingale, and expectation of a submartingale is an increasing function of t . Hence the supremum is the same as the limit.
2. $\lim_{t \rightarrow \infty} EM^+(t) < \infty$. This is because $E|M(t)| = EM^+(t) + EM^-(t)$. If $EM(t) = c$, then $EM(t) = EM^+(t) - EM^-(t) = c$, and $EM^+(t) = EM^-(t) + c$.
3. $\lim_{t \rightarrow \infty} EM^-(t) < \infty$.

If $M(t)$ is a submartingale it is enough to demand $\sup_t EM^+(t) < \infty$, and if it is a supermartingale it is enough to demand $\sup_t EM^-(t) < \infty$, for the existence of a finite limit.

Corollary 7.12:

1. *Uniformly integrable martingales converge almost surely.*
2. *Square integrable martingales converge almost surely.*
3. *Positive martingales converge almost surely.*
4. *Submartingales bounded from above (negative) converge almost surely.*
5. *Supermartingales bounded from below (positive) converge almost surely.*

PROOF. Since uniformly integrable martingales are integrable, they converge. Since square integrable martingales are uniformly integrable, they converge. If $M(t)$ is positive, then $|M(t)| = M(t)$, and $E|M(t)| = EM(t) = EM(0) < \infty$. \square

Note that expectations $EM(t)$ may or may not converge to the expectation of the limit EY (see Example 7.3). The case when $EY = \lim_{t \rightarrow \infty} EM(t)$ is precisely when $M(t)$ is uniformly integrable. The next result, given without proof, shows that uniformly integrable martingales have form (7.5).

Theorem 7.13: *If $M(t)$ is a uniformly integrable martingale, then it converges to a random variable Y almost surely and in L^1 . Conversely, if $M(t)$ is a martingale that converges in L^1 to a random variable Y , then $M(t)$ is uniformly integrable, and it converges almost surely to Y . In any case $M(t) = E(Y|\mathcal{F}_t)$.*

Example 7.3: (Exponential martingale of Brownian motion)

Let $M(t) = e^{B(t)-t/2}$. Then $M(t)$, $t \geq 0$ is a martingale. Since it is positive, it converges by Corollary 7.12 almost surely to a limit Y . According to the law of large numbers for Brownian motion $B(t)/t$ converges almost surely to zero. Thus $M(t) = e^{t(B(t)-t/2)} \rightarrow 0$ as $t \rightarrow \infty$. Thus $Y = 0$ almost surely. Therefore $M(t)$ is not uniformly integrable, as $EY = 0 \neq 1 = EM(t)$.

Example 7.4: Let $f(s)$ be non-random such that $\int_0^\infty f^2(s)ds < \infty$. We show that $M(t) = \int_0^t f(s)dB(s)$ is a uniformly integrable martingale, and find a representation for the closing random variable.

Since $\int_0^\infty f^2(s)ds < \infty$, $\int_0^t f(s)dB(s)$ is defined for all $t > 0$ and is a martingale. Since $\sup_{t>0} E(M^2(t)) = \sup_{t>0} \int_0^t f^2(s)ds = \int_0^\infty f^2(s)ds < \infty$, $M(t)$ is uniformly integrable. Thus it converges almost surely to Y . Convergence is also in L^1 , that is, $E|M(t) - Y| \rightarrow 0$ as $t \rightarrow \infty$. Denote $Y = M(\infty) = \int_0^\infty f(s)dB(s)$, then we have shown that $Y - M(t) = \int_t^\infty f(s)dB(s)$ converges to zero almost surely and in L^1 . Y is the closing variable. Indeed,

$$E(Y|\mathcal{F}_t) = E(M(\infty)|\mathcal{F}_t) = E\left(\int_0^\infty f(s)dB(s)|\mathcal{F}_t\right) = \int_0^t f(s)dB(s) = M(t).$$

Example 7.5: A bounded positive martingale $M(t) = E(I(Y > 0)|\mathcal{F}_t)$ with $Y = \int_0^\infty f(s)dB(s)$, where $f(s)$ is non-random and $\int_0^\infty f^2(s)ds < \infty$, from the previous example.

$$\begin{aligned} M(t) &= E(I(Y > 0)|\mathcal{F}_t) = P(Y > 0|\mathcal{F}_t) \\ &= P\left(\int_t^\infty f(s)dB(s) > -\int_0^t f(s)dB(s)\Big|\mathcal{F}_t\right) \\ &= \Phi\left(\frac{\int_0^t f(s)dB(s)}{\sqrt{\int_t^\infty f^2(s)ds}}\right), \end{aligned} \tag{7.6}$$

where the last equality is due to the normality of the Itô integral for a non-random f .

By taking f to be zero on (T, ∞) , a result is obtained for martingales of the form $E(I(\int_0^T f(s)dB(s) > 0)|\mathcal{F}_t)$. In particular, by taking $f(s) = 1_{[0,T]}(s)$, we obtain that

$$\Phi(B(t)/\sqrt{T-t})$$

is a positive bounded martingale on $[0, T]$. Its distribution for $t = T$ is left as an exercise.

7.4 Optional Stopping

In this section we consider results on stopping martingales at random times. Recall that a random time τ is called a stopping time if for any $t > 0$ the sets $\{\tau \leq t\} \in \mathcal{F}_t$. For filtrations generated by a process X , τ is a stopping time if it is possible to decide whether τ has occurred or not by observing the process up to time t . A martingale stopped at a random time τ is the process $M(t \wedge \tau)$. A basic stopping result, given here without proof, states that a martingale stopped at a stopping time is a martingale, in particular $EM(\tau \wedge t) = EM(0)$. This equation is used most frequently.

Theorem 7.14: *If $M(t)$ is a martingale and τ is a stopping time, then the stopped process $M(\tau \wedge t)$ is a martingale. Moreover,*

$$EM(\tau \wedge t) = EM(0). \quad (7.7)$$

This result was proved in discrete time (see Theorem 3.39). We refer to (7.7) as the basic stopping equation.

Remark 7.1: We stress that in this theorem $M(\tau \wedge t)$ is a martingale with respect to the original filtration \mathcal{F}_t . Since it is adapted to $\mathcal{F}_{\tau \wedge t}$, it is also an $\mathcal{F}_{\tau \wedge t}$ -martingale (see Exercise 7.1).

Example 7.6: (Exit of Brownian motion from an interval)

Let $B(t)$ be Brownian motion started at x , and let τ be the first time when $B(t)$ exits the interval (a, b) , $a < x < b$, that is, $\tau = \inf\{t : B(t) = a \text{ or } b\}$. Clearly, τ is a stopping time. By the basic stopping result (7.7), $EB(t \wedge \tau) = B(0) = x$. By definition of τ , $|B(t \wedge \tau)| \leq \max(|a|, |b|)$. Thus one can take $t \rightarrow \infty$, and use dominated convergence to obtain $EB(\tau) = x$. However, $B(\tau) = b$ with probability p and $B_\tau = a$ with probability $1 - p$. From these equations we obtain that $p = (x - a)/(b - a)$ is the probability that Brownian motion reaches b before it reaches a .

If M is a martingale, then $EM(t) = EM(0)$. If τ is a stopping time, then $EM(\tau)$ may be different to $EM(0)$, as Example 7.7 shows.

Example 7.7: Let $B(t)$ be Brownian motion started at 0 and τ is the hitting time of 1. Then by definition $B(\tau) = 1$ and $EB(\tau) = 1 \neq 0 = EB(0)$.

However, under some additional assumptions on the martingale or on the stopping time, the random stopping does not alter the expected value. The following result gives sufficient conditions for optional stopping to hold.

Theorem 7.15 (Optional Stopping): *Let $M(t)$ be a martingale.*

1. *If $\tau \leq K < \infty$ is a bounded stopping time, then $EM(\tau) = EM(0)$.*
2. *If $M(t)$ is uniformly integrable, then for any stopping time τ , $EM(\tau) = EM(0)$.*

Applied to gambling this shows that when betting on a martingale, on average no loss or gain is made, even if a clever stopping rule is used, provided optional stopping holds.

The first statement follows from the basic stopping result by taking $t > K$ and applying (7.7), $EM(0) = EM(t \wedge \tau) = EM(\tau)$. It is important to note that the second statement holds for any stopping time even if it takes infinite value with a positive probability. We do not give a proof of this statement, the difficult point is in showing that $M(\tau)$ is integrable, see Dellacherie and Meyer (1982). For finite stopping times we show

Theorem 7.16: *Let $M(t)$ be a martingale and τ a finite stopping time. If $E|M(\tau)| < \infty$, and*

$$\lim_{t \rightarrow \infty} E(M(t)I(\tau > t)) = 0, \quad (7.8)$$

then $EM(\tau) = EM(0)$.

PROOF. Write $M(\tau \wedge t)$ as

$$M(\tau \wedge t) = M(t)I(t < \tau) + M(\tau)I(t \geq \tau). \quad (7.9)$$

Using the basic stopping result (7.7), $EM(\tau \wedge t) = EM(0)$. Taking expectations in (7.9), we have

$$EM(0) = E(M(t)I(t < \tau)) + E(M(\tau)I(t \geq \tau)). \quad (7.10)$$

Now take the limit in (7.10) as $t \rightarrow \infty$. Since τ is finite, $I(t \geq \tau) \rightarrow I(\tau < \infty) = 1$. $|M(\tau)|I(t \geq \tau) \leq |M(\tau)|$, integrable. Hence $E(M(\tau)I(t \geq \tau)) \rightarrow EM(\tau)$ by dominated convergence. It is assumed that $E(M(t)I(t < \tau)) \rightarrow 0$ as $t \rightarrow \infty$, and the result follows. \square

The basic stopping result or optional stopping are used to find the distribution of stopping times for Brownian motion and random walks.

Example 7.8: (Hitting times of Brownian motion)

We derive the Laplace transform of hitting times, from which it also follows that they are finite. Let $B(t)$ be a Brownian motion starting at 0, and $T_b = \inf\{t : B(t) = b\}$, $b > 0$. Consider the exponential martingale of Brownian motion $e^{uB(t)-u^2t/2}$, $u > 0$, stopped at T_b , $e^{uB(t\wedge T_b)-(t\wedge T_b)u^2/2}$. Using the basic stopping result (7.7)

$$\mathbb{E}e^{uB(t\wedge T_b)-(t\wedge T_b)u^2/2} = 1.$$

The stopped martingale is bounded from above by e^{ub} and it is positive. If we take it as already proven that T_b is finite, $\mathbb{P}(T_b < \infty) = 1$, then we obtain by taking $t \rightarrow \infty$ that $\mathbb{E}e^{ub-T_b u^2/2} = 1$. Replacing u by $\sqrt{2u}$, we obtain the Laplace transform of T_b

$$\psi(u) = \mathbb{E}\left(e^{-uT_b}\right) = e^{-b\sqrt{2u}}. \quad (7.11)$$

We now show the finiteness of T_b . Write the expectation of the stopped martingale

$$\mathbb{E}(e^{ub-T_b u^2/2} I(T_b \leq t)) + \mathbb{E}(e^{uB(t)-tu^2/2} I(T_b > t)) = 1. \quad (7.12)$$

The term $\mathbb{E}(e^{uB(t)-tu^2/2} I(T_b > t)) \leq \mathbb{E}(e^{ub-tu^2/2} I(T_b > t)) \leq e^{ub-tu^2/2} \rightarrow 0$, as $t \rightarrow \infty$. Thus taking limits in the above Equation (7.12), $\mathbb{E}(e^{ub-T_b u^2/2} I(T_b \leq t)) \rightarrow \mathbb{E}(e^{ub-T_b u^2/2} I(T_b < \infty)) = 1$. Therefore $\mathbb{E}(e^{-T_b u^2/2} I(T_b < \infty)) = e^{-ub}$. However, $e^{-T_b u^2} I(T_b = \infty) = 0$, therefore, by adding this term, we can write $\mathbb{E}(e^{-T_b u^2/2}) = e^{-ub}$. It follows in particular that $\mathbb{P}(T_b < \infty) = \lim_{u \downarrow 0} \psi(u) = 1$, and T_b is finite. Hence (7.11) is proved. The distribution of T_b corresponding to the transform (7.11) is given in Theorem 3.18.

The following result is in some sense the converse to the optional stopping theorem.

Theorem 7.17: *Let $X(t)$, $t \geq 0$, be such that for any bounded stopping time τ , $X(\tau)$ is integrable and $\mathbb{E}X(\tau) = \mathbb{E}X(0)$. Then $X(t)$, $t \geq 0$ is a martingale.*

PROOF. The proof consists of checking the martingale property by using appropriate stopping times. Since a deterministic time t is a stopping time, $X(t)$ is integrable. Without loss of generality take $X(0) = 0$. Next we show that for $t > s$, $\mathbb{E}(X(t)|\mathcal{F}_s) = X(s)$. In other words, we need to show that for any $s < t$ and any set $B \in \mathcal{F}_s$

$$\mathbb{E}(X(t)I(B)) = \mathbb{E}(X(s)I(B)). \quad (7.13)$$

Fix a set $B \in \mathcal{F}_s$ and for any $t > s$, define a stopping time $\tau = sI(B) + tI(B^c)$. We have $\mathbb{E}(X(\tau)) = \mathbb{E}(X(s)I(B)) + \mathbb{E}(X(t)I(B^c))$. Since $\mathbb{E}X(\tau) = 0$, $\mathbb{E}(X(s)I(B)) = \mathbb{E}X(\tau) - \mathbb{E}(X(t)I(B^c)) = -\mathbb{E}(X(t)I(B^c))$. As the right hand side of the above equality does not depend on s , it follows that (7.13) holds. \square

The following result is sometimes known as the optional sampling theorem (see for example Rogers and Williams (1990)).

Theorem 7.18 (Optional Sampling): *Let $M(t)$ be a uniformly integrable martingale, and $\tau_1 \leq \tau_2 \leq \infty$ two stopping times. Then*

$$\mathbb{E}(M(\tau_2)|\mathcal{F}_{\tau_1}) = M(\tau_1), \text{ almost surely.} \quad (7.14)$$

Optional Stopping of Discrete Time Martingales

We consider next the case of discrete time $t = 0, 1, 2, \dots$, and martingales arising in a random walk.

Gambler's Ruin

Consider a game played by two people betting on the outcomes of tosses of a coin. You win \$1 if heads come up and lose \$1 if tails come up. The game stops when one party has no money left. You start with x , and your opponent with b dollars. Then S_n , the amount of money you have at time n , is a random walk (see Chapter 3.12). The gambler's ruin problem is to find the probabilities of ruin of the players.

In this game the loss of one person is the gain of the other (a zero sum game). Assuming that the game will end in a finite time τ (this fact will be shown later), it follows that the ruin probabilities of the players add up to one.

Consider first the case of the fair coin. Then

$$S_n = x + \sum_{i=1}^n \xi_i, \quad P(\xi_i = 1) = \frac{1}{2}, \quad P(\xi_i = -1) = \frac{1}{2},$$

is a martingale (see Theorem 3.33). Let τ be the time when the game stops, the first time the amount of money you have is equal to 0 (your ruin) or $x + b$ (your opponent's ruin). Then τ is a stopping time. Denote by u the probability of your ruin. It is the probability of you losing your initial capital x before winning b dollars. Thus

$$P(S_\tau = 0) = u \text{ and } P(S_\tau = x + b) = 1 - u. \quad (7.15)$$

Formally applying the optional stopping theorem

$$\mathbb{E}(S_\tau) = S_0 = x. \quad (7.16)$$

However,

$$\mathbb{E}(S_\tau) = (x + b) \times (1 - u) + 0 \times u = (x + b)(1 - u).$$

These equations give

$$u = \frac{b}{x+b}, \quad (7.17)$$

so that the ruin probabilities are given by a simple calculation using martingale stopping.

We now justify the steps. S_n is a martingale, and τ is a stopping time. According to Theorem 7.14 the stopped process $S_{n \wedge \tau}$ is a martingale. It is non-negative and bounded by $x+b$, by the definition of τ . Thus $S_{n \wedge \tau}$ is a uniformly integrable martingale. Hence it converges almost surely to a finite limit Y , (with $EY = x$), $\lim_{n \rightarrow \infty} S_{n \wedge \tau} = Y$. According to Theorem 3.33 $S_n^2 - n$ is a martingale, and so is $S_{n \wedge \tau}^2 - n \wedge \tau$. Thus for all n by taking expectation

$$E(S_{n \wedge \tau}^2) = E(n \wedge \tau) + E(S_0^2). \quad (7.18)$$

By dominated convergence, the lhs has a finite limit, therefore there is a finite limit $\lim_{n \rightarrow \infty} E(n \wedge \tau)$. Expanding this, $E(n \wedge \tau) \geq nP(\tau > n)$, we can see that for a limit to exist it must be

$$\lim_{n \rightarrow \infty} P(\tau > n) = 0, \quad (7.19)$$

so that $P(\tau < \infty) = 1$, and τ is finite. (Note that a standard proof of finiteness of τ is done by using Markov chain theory, the property of the recurrence of states in a random walk.) Writing $E(S_{n \wedge \tau}) = x$ and taking limits as $n \rightarrow \infty$, the Equation (7.16) is obtained (alternatively, the conditions of the optional stopping theorem hold). This concludes a rigorous derivation of the ruin probability in an unbiased random walk.

We now consider the case when the random walk is biased, $p \neq q$.

$$S_n = x + \sum_{i=1}^n \xi_i, \quad P(\xi_i = 1) = p, \quad P(\xi_i = -1) = q = 1 - p.$$

In this case, the exponential martingale of the random walk $M_n = (q/p)^{S_n}$ is used (see Theorem 3.33). Stopping this martingale, we obtain the ruin probability

$$u = \frac{(q/p)^{b+x} - (q/p)^x}{(q/p)^{b+x} - 1}. \quad (7.20)$$

Justification of the equation $E(M_\tau) = M_0$ is similar to the previous case.

Hitting Times in Random Walks

Let S_n denote a random walk on the integers started at $S_0 = x$, $S_n = S_0 + \sum_{i=1}^n \xi_i$, $P(\xi_i = 1) = p$, $P(\xi_i = -1) = q = 1 - p$, with arbitrary p , and

T_b the first hitting time of b , $T_b = \inf\{n : S_n = b\}$ (infimum of an empty set is infinity). Without loss of generality take the starting state $x = 0$, otherwise consider the process $S_n - x$. Consider hitting the level $b > 0$, for $b < 0$ consider the process $-S_n$.

We find the Laplace transform of T_b , $\psi(\lambda) = E(e^{-\lambda T_b})$, $\lambda > 0$, by stopping the exponential martingale of the random walk $M_n = e^{uS_n - nh(u)}$, where $h(u) = \ln E(e^{u\xi_1})$, and u is arbitrary (see Chapter 3.12).

$$E(M_{n \wedge T_b}) = E(e^{uS_{n \wedge T_b} - (n \wedge T_b)h(u)}) = 1. \quad (7.21)$$

Take u , so that $h(u) = \lambda > 0$. Write the expectation in (7.21) as

$$E(e^{uS_{n \wedge T_b} - (n \wedge T_b)h(u)}) = E(e^{uS_{T_b} - T_b h(u)} I(T_b \leq n)) + E(e^{uS_n - nh(u)} I(T_b > n)).$$

The first term equals $E(e^{ub - T_b h(u)} I(T_b \leq n))$. The second term converges to zero, because by definition of T_b ,

$$E(e^{uS_n - nh(u)} I(T_b > n)) \leq E(e^{ub - nh(u)} I(T_b > n)) \leq e^{ub - nh(u)} \rightarrow 0.$$

Now taking limits in (7.21), using dominated convergence we obtain

$$E(e^{-h(u)T_b} I(T_b < \infty)) = e^{-ub}.$$

Note that $e^{-h(u)T_b} I(T_b = \infty) = 0$, therefore by adding this term, we can write

$$E(e^{-h(u)T_b}) = e^{-ub}. \quad (7.22)$$

This is practically the Laplace transform of T_b , it remains to replace $h(u)$ by λ , by taking $u = h^{(-1)}(\lambda)$, with $h^{(-1)}$ being the inverse of h . Thus the Laplace transform of T_b is given by

$$\psi(\lambda) = E(e^{-\lambda T_b}) = e^{-h^{(-1)}(\lambda)b}. \quad (7.23)$$

In order to find $h^{(-1)}(\lambda)$, solve $h(u) = \lambda$, which is equivalent to $E(e^{u\xi_1}) = e^\lambda$, or $pe^u + (1-p)e^{-u} = e^\lambda$. There are two values for $e^u = (e^\lambda \pm \sqrt{e^{2\lambda} - 4p(1-p)})/(2p)$, but only one corresponds to a Laplace transform, (7.22). Thus we have

$$\psi(\lambda) = E(e^{-\lambda T_b}) = \left(\frac{2p}{e^\lambda + \sqrt{e^{2\lambda} - 4p(1-p)}} \right)^b. \quad (7.24)$$

Using a general result on the Laplace transform of a random variable,

$$P(T_b < \infty) = \lim_{\lambda \downarrow 0} \psi(\lambda) = \left(\frac{2p}{1 + |1 - 2p|} \right)^b. \quad (7.25)$$

It now follows that the hitting time T_b of b is finite if, and only if, $p \geq 1/2$. For $p < 1/2$, there is a positive probability that level b is never reached, $P(T_b = \infty) = 1 - (\frac{p}{1-p})^b$.

When the hitting time of level b is finite, it may or may not have a finite expectation. If $p \geq 1/2$, we have

$$E(T_b) = -\psi'(0) = \begin{cases} \frac{b}{2p-1} & \text{if } p > 1/2 \\ \infty & \text{if } p = 1/2. \end{cases}$$

Thus we have shown that when $p \geq 1/2$ any positive state will be reached from 0 in a finite time, but when $p = 1/2$ the average time for it to happen is infinite.

The results obtained above are known as transience ($p \neq 1/2$) and recurrence ($p = 1/2$) of the random walk, and are usually obtained by Markov chain theory.

Example 7.9: (Optional stopping of discrete time martingales)

Let $M(t)$ be a discrete time martingale and τ be a stopping time such that $E|M(\tau)| < \infty$.

1. If $E\tau < \infty$ and $|M(t+1) - M(t)| \leq K$, then $EM(\tau) = EM(0)$.
2. If $E\tau < \infty$ and $E(|M(t+1) - M(t)| | \mathcal{F}_t) \leq K$, then $EM(\tau) = EM(0)$.

PROOF. We prove the first statement. $M(t) = M(0) + \sum_{i=0}^{t-1} (M(i+1) - M(i))$. This together with the bound on increments gives

$$M(t) \leq |M(0)| + \sum_{i=0}^{t-1} |M(i+1) - M(i)| \leq |M(0)| + Kt.$$

Take for simplicity non-random $M(0)$. Then

$$EM(t)I(\tau > t) \leq |M(0)|P(\tau > t) + KtP(\tau > t).$$

The last term converges to zero, $tP(\tau > t) \leq E(\tau I(\tau > t)) \rightarrow 0$, by dominated convergence due to $E(\tau) < \infty$. Thus condition (7.8) holds, and the result follows. The proof of the second statement is similar and is left as an exercise. \square

7.5 Localization and Local Martingales

As was seen earlier in Chapter 4, Itô integrals $\int_0^t X(s)dB(s)$ are martingales under the additional condition $E \int_0^T X^2(s)ds < \infty$. In general, stochastic integrals with respect to martingales are only local martingales rather than true martingales. This is the main reason for introducing local martingales. We have also seen that for the calculation of expectations stopping and truncations are often used. These ideas give rise to the following.

Definition 7.19: A property of a stochastic process $X(t)$ is said to hold locally if there exists a sequence of stopping times τ_n , called the localizing sequence, such that $\tau_n \uparrow \infty$ as $n \rightarrow \infty$, and for each n the stopped processes $X(t \wedge \tau_n)$ have this property.

For example, the uniform integrability property holds locally for any martingale. According to Theorem 7.13 a martingale convergent in L^1 is uniformly integrable. Here $M(t \wedge n) = M(n)$ for $t > n$, and therefore $\tau_n = n$ is a localizing sequence.

Local martingales are defined by localizing the martingale property.

Definition 7.20: An adapted process $M(t)$ is called a local martingale if there exists a sequence of stopping times τ_n such that $\tau_n \uparrow \infty$ and for each n the stopped processes $M(t \wedge \tau_n)$ are uniformly integrable martingales in t .

As we have just seen, any martingale is a local martingale. Examples of local martingales which are not martingales are given below.

Example 7.10: $M(t) = 1/|\mathbf{B}(t)|$, where $\mathbf{B}(t)$ is the three-dimensional Brownian motion, $\mathbf{B}(0) = \mathbf{x} \neq \mathbf{0}$. We have seen in Example 6.17 that if D_r is the complementary set to the ball of radius r centered at the origin, $D_r = \{\mathbf{z} : |\mathbf{z}| > r\}$, then $f(\mathbf{z}) = 1/|\mathbf{z}|$ is a harmonic function for the Laplacian on D_r . Consequently, $1/|\mathbf{B}(t \wedge \tau_{D_r})|$ is a martingale, where τ_{D_r} is the time of exit from D_r . Take now τ_n to be the exit time from $D_{1/n}$, that is, $\tau_n = \inf\{t > 0 : |\mathbf{B}(t)| = 1/n\}$. Then for any fixed n , $1/|\mathbf{B}(t \wedge \tau_n)|$ is a martingale. τ_n increases to, say, τ and by continuity, $\mathbf{B}(\tau) = \mathbf{0}$. As Brownian motion in three dimensions never visits the origin (see Example 6.14) it follows by continuity that τ is infinite. Thus $M(t)$ is a local martingale. In order to see that it is not a true martingale, recall that in three dimensions Brownian motion is transient and $|\mathbf{B}(t)| \rightarrow \infty$ as $t \rightarrow \infty$. Therefore $EM(t) \rightarrow 0$, whereas $EM(0) = 1/|\mathbf{x}|$. Since the expectation of a martingale is constant, $M(t)$ is not a martingale.

Example 7.11: (Itô integrals)

Let $M(t) = \int_0^t e^{B^2(s)} dB(s)$, $t > 1/4$, where B is Brownian motion in one dimension with $B(0) = 0$. Let $\tau_n = \inf\{t > 0 : e^{B^2(t)} = n\}$. Then for $t \leq \tau_n$, the integrand is bounded by n . According to the martingale property of Itô integrals, $M(t \wedge \tau_n)$ is a martingale in t for any n . Thus $\tau_n \rightarrow \tau = \infty$, as according to continuity, $\exp(B^2(\tau)) = \infty$. Therefore $M(t)$ is a local martingale. In order to see that it is not a martingale notice that for $t > 1/4$, $E(e^{2B^2(t)}) = \infty$, implying that $M(t)$ is not integrable.

Remark 7.2: Note that it is not enough for a local martingale to be integrable in order to be a true martingale. For example, positive local martingales are integrable, but in general they are not martingales but only supermartingales (see Theorem 7.23). Even uniformly integrable local martingales may not be martingales. However, if a local martingale is dominated by an integrable random variable then it is a martingale.

Theorem 7.21: Let $M(t)$, $0 \leq t < \infty$, be a local martingale such that $|M(t)| \leq Y$, with $EY < \infty$. Then M is a uniformly integrable martingale.

PROOF. Let τ_n be a localizing sequence. Then for any n and $s < t$

$$E(M(t \wedge \tau_n) | \mathcal{F}_s) = M(s \wedge \tau_n). \quad (7.26)$$

M is clearly integrable, since $E|M(t)| \leq EY < \infty$. Since $\lim_{n \rightarrow \infty} M(t \wedge \tau_n) = M(t)$, by dominated convergence of conditional expectations $\lim_{n \rightarrow \infty} E(M(t \wedge \tau_n) | \mathcal{F}_s) = E(M(t) | \mathcal{F}_s)$. Since $\lim_{n \rightarrow \infty} M(s \wedge \tau_n) = M(s)$, the martingale property is established by taking limits in (7.26). If a martingale is dominated by an integrable random variable then it is uniformly integrable (see Theorem 7.6). \square

Corollary 7.22: Let $M(t)$, $0 \leq t < \infty$, be a local martingale such that for all t , $E(\sup_{s \leq t} |M(s)|) < \infty$. Then it is a martingale, and as such it is uniformly integrable on any finite interval $[0, T]$. If in addition $E(\sup_{t \geq 0} |M(t)|) < \infty$, then $M(t)$, $t \geq 0$, is uniformly integrable on $[0, \infty)$.

In financial applications we meet positive local martingales.

Theorem 7.23: A non-negative local martingale $M(t)$, $0 \leq t \leq T$, is a supermartingale, that is, $EM(t) < \infty$, and for any $s < t$, $E(M(t) | \mathcal{F}_s) \leq M(s)$.

PROOF. Let τ_n be a localizing sequence. Then, since $M(t \wedge \tau_n) \geq 0$, by Fatou's lemma

$$E\left(\liminf_{n \rightarrow \infty} M(t \wedge \tau_n)\right) \leq \liminf_{n \rightarrow \infty} E(M(t \wedge \tau_n)). \quad (7.27)$$

Since the limit exists, the lower limit is the same, that is, $\lim_{n \rightarrow \infty} M(t \wedge \tau_n) = M(t)$ implies $\liminf_{n \rightarrow \infty} M(t \wedge \tau_n) = M(t)$. However, $EM(t \wedge \tau_n) = EM(0 \wedge \tau_n) = EM(0)$ by the martingale property of $M(t \wedge \tau_n)$. Therefore by taking limits, $EM(t) \leq EM(0)$, so that M is integrable. The supermartingale property is established similarly. Using Fatou's lemma for conditional expectations,

$$E\left(\liminf_{n \rightarrow \infty} M(t \wedge \tau_n) | \mathcal{F}_s\right) \leq \liminf_{n \rightarrow \infty} E(M(t \wedge \tau_n) | \mathcal{F}_s) = \liminf_{n \rightarrow \infty} M(s \wedge \tau_n), \quad (7.28)$$

and we obtain $E(M(t) | \mathcal{F}_s) \leq M(s)$ almost surely. \square

From this result and Theorem 7.3 we obtain

Theorem 7.24: A non-negative local martingale $M(t)$, $0 \leq t \leq T$, is a martingale if, and only if, $EM(T) = M(0)$.

For a general local martingale a necessary and sufficient condition to be a uniformly integrable martingale is described in terms of the property of Dirichlet class (D). This class of processes also arises in other areas of calculus and is given in the following section.

Dirichlet Class (D)

Definition 7.25: A process X is of Dirichlet class (D) if the family $\{X(\tau) : \tau \text{ a finite stopping time}\}$ is uniformly integrable.

Any uniformly integrable martingale M is of class (D). Indeed, by Theorem 7.13, M is closed by $Y = M(\infty)$, $M(\tau) = E(Y|\mathcal{F}_\tau)$, and the last family is uniformly integrable.

Using localization one can show the other direction, and we have a theorem.

Theorem 7.26: A local martingale M is a uniformly integrable martingale if, and only if, it is of class (D).

PROOF. Suppose that M is a local martingale of class (D).

Let τ_n be a localizing sequence so that $M(t \wedge \tau_n)$ is a uniformly integrable martingale in t . Then for $s < t$,

$$M(s \wedge \tau_n) = E(M(t \wedge \tau_n) | \mathcal{F}_s). \quad (7.29)$$

The martingale property of M is obtained by taking $n \rightarrow \infty$ in both sides of the equation.

Since $\tau_n \rightarrow \infty$, $M(s \wedge \tau_n) \rightarrow M(s)$ almost surely. $s \wedge \tau_n$ is a finite stopping time, and because M is in (D), the sequence of random variables $\{M(s \wedge \tau_n)\}_n$ is uniformly integrable. Thus $M(s \wedge \tau_n) \rightarrow M(s)$ also in L^1 , that is,

$$E|M(s \wedge \tau_n) - M(s)| \rightarrow 0. \quad (7.30)$$

Using the properties of conditional expectation,

$$\begin{aligned} E|E(M(t \wedge \tau_n) | \mathcal{F}_s) - E(M(t) | \mathcal{F}_s)| &= E|E(M(t \wedge \tau_n) - M(t) | \mathcal{F}_s)| \\ &\leq E(E|M(t \wedge \tau_n) - M(t)| | \mathcal{F}_s) \\ &= E|M(t \wedge \tau_n) - M(t)|. \end{aligned}$$

The latter converges to zero by (7.30). This implies $E(M(t \wedge \tau_n) | \mathcal{F}_s) \rightarrow E(M(t) | \mathcal{F}_s)$ as $n \rightarrow \infty$. Taking limits in (7.29) as $n \rightarrow \infty$ establishes the martingale property of M . Since it is in (D), by taking $\tau = t$, it is uniformly integrable. \square

7.6 Quadratic Variation of Martingales

Quadratic variation of a process $X(t)$ is defined as a limit in probability

$$[X, X](t) = \lim \sum_{i=1}^n (X(t_i^n) - X(t_{i-1}^n))^2, \quad (7.31)$$

where the limit is taken over partitions:

$$0 = t_0^n < t_1^n < \cdots < t_n^n = t,$$

with $\delta_n = \max_{0 \leq i \leq n} (t_i^n - t_{i-1}^n) \rightarrow 0$. If $M(t)$ is a martingale, then $M^2(t)$ is a submartingale, and its mean increases (unless $M(t)$ is a constant). By compensating $M^2(t)$ by some increasing process, it is possible to make it into a martingale. The process which compensates $M^2(t)$ to a martingale turns out to be the quadratic variation process of M . It can be shown that quadratic variation of martingales exists and is characterized by the above property.

Theorem 7.27:

1. Let $M(t)$ be a martingale with finite second moments, $E(M^2(t)) < \infty$ for all t . Then its quadratic variation process $[M, M](t)$ defined in (7.31) exists; moreover, $M^2(t) - [M, M](t)$ is a martingale.
2. If M is a local martingale, then $[M, M](t)$ exists; moreover, $M^2(t) - [M, M](t)$ is a local martingale.

PROOF. We outline the proof of the first statement only. The second follows for locally square integrable martingales by localization. For local martingales the result follows from representation of quadratic variation by means of stochastic integrals. For a full proof see, for example, Liptser and Shiryaev (1989).

$$E(M(t)M(s)) = EE(M(t)M(s)|\mathcal{F}_s) = E(M(s)E(M(t)|\mathcal{F}_s)) = E(M^2(s)). \quad (7.32)$$

Using this it is easy to obtain

$$E(M(t) - M(s))^2 = E(M^2(t)) - E(M^2(s)). \quad (7.33)$$

It is easy to see that the sums in the definition of quadratic variation $[M, M](t)$ have constant mean, that of $E(M^2(t)) - E(M^2(0))$. It is possible, but not easy, to prove that these sums converge in probability to the limit $[M, M](t)$. Now using property (7.33), we can write

$$\begin{aligned} E(M^2(t) - M^2(s)|\mathcal{F}_s) &= E((M(t) - M(s))^2|\mathcal{F}_s) \\ &= E\left(\sum_{i=0}^{n-1} (M(t_{i+1}) - M(t_i))^2|\mathcal{F}_s\right), \end{aligned} \quad (7.34)$$

where $\{t_i\}$ is a partition of $[s, t]$. Taking the limit as the size of the partition goes to zero, we obtain

$$E(M^2(t) - M^2(s)|\mathcal{F}_s) = E([M, M](t) - [M, M](s)|\mathcal{F}_s). \quad (7.35)$$

Rearranging, we obtain the martingale property of $M^2(t) - [M, M](t)$. \square

For the next result note that if M is a martingale, then for any t

$$\mathbb{E}(M(t) - M(0))^2 = \mathbb{E}(M^2(t)) - \mathbb{E}(M^2(0)),$$

which shows that $\mathbb{E}(M^2(t)) > \mathbb{E}(M^2(0))$ unless $M(t) = M(0)$ almost surely. Thus M^2 cannot be a martingale on $[0, t]$ unless $M(t) = M(0)$. If $M(t) = M(0)$, then for all $s < t$, $M(s) = \mathbb{E}(M(t)|\mathcal{F}_s) = M(0)$, and M is a constant on $[0, t]$.

Theorem 7.28: *Let M be a martingale with $M(0) = 0$. If for some t , $M(t)$ is not identically zero, then $[M, M](t) > 0$. Conversely, if $[M, M](t) = 0$, then $M(s) = 0$ almost surely for all $s \leq t$. The result also holds for local martingales.*

PROOF. We prove the result for square integrable martingales; for local martingales it can be shown by localization. Suppose that $[M, M](t) = 0$ for some $t > 0$. Then, since $[M, M]$ is non-decreasing, $[M, M](s) = 0$ for all $s \leq t$. According to Theorem 7.27 $M^2(s)$, $s \leq t$, is a martingale. In particular, $\mathbb{E}(M^2(t)) = 0$. This implies that $M(t) = 0$ almost surely, which is a contradiction. Therefore $[M, M](t) > 0$.

Conversely, if $[M, M](t) = 0$, the same argument shows that $M(t) = 0$ almost surely, and by the martingale property $M(s) = 0$ for all $s \leq t$. \square

It also follows from the proof that M and $[M, M]$ have the same intervals of constancy. This theorem implies remarkably that a continuous martingale which is not a constant has infinite variation on any interval.

Theorem 7.29: *Let M be a continuous local martingale and fix any t . If $M(t)$ is not identically equal to $M(0)$, then M has infinite variation over $[0, t]$.*

PROOF. $M(t) - M(0)$ is a martingale, null at zero, with its value at time t not equal identically to zero. According to Theorem 7.28 M has a positive quadratic variation on $[0, t]$, $[M, M](t) > 0$. According to Theorem 1.10 a continuous process of finite variation on $[0, t]$ has zero quadratic variation over this interval. Therefore M must have infinite variation over $[0, t]$. \square

Corollary 7.30: *If a continuous local martingale has finite variation over an interval, then it must be a constant over that interval.*

Remark 7.3: Note that there are martingales with finite variation, but by the previous result they cannot be continuous. An example of such a martingale is the Poisson process martingale $N(t) - t$.

7.7 Martingale Inequalities

$M(t)$ denotes a martingale or a local martingale on the interval $[0, T]$ with possibly $T = \infty$.

Theorem 7.31: *If $M(t)$ is a martingale (or a positive submartingale), then for $p \geq 1$*

$$P(\sup_{s \leq t} |M(s)| \geq a) \leq a^{-p} \sup_{s \leq t} E(|M(s)|^p). \quad (7.36)$$

If $p > 1$, then

$$E\left(\sup_{s \leq t} |M(s)|^p\right) \leq \left(\frac{p}{p-1}\right)^p E(|M(t)|^p). \quad (7.37)$$

The case of $p = 2$ is called Doob's inequality for martingales.

$$E\left(\sup_{s \leq T} M(s)^2\right) \leq 4E(M^2(T)). \quad (7.38)$$

As a consequence, if for $p > 1$, $\sup_{t \leq T} E(|M(t)|^p) < \infty$, then $M(t)$ is uniformly integrable (This is a particular case of Theorem 7.7).

Theorem 7.32: *If M is a locally square integrable martingale with $M(0)=0$, then*

$$P(\sup_{t \leq T} |M(t)| > a) \leq a^{-2} E([M, M](T)). \quad (7.39)$$

Theorem 7.33 (Davis Inequality): *There are constants $c > 0$ and $C < \infty$ such that for any local martingale $M(t)$, null at zero,*

$$cE\left(\sqrt{[M, M](T)}\right) \leq E\left(\sup_{t \leq T} |M(t)|\right) \leq CE\left(\sqrt{[M, M](T)}\right). \quad (7.40)$$

Theorem 7.34 (Burkholder–Gundy Inequality): *There are constants c_p and C_p depending only on p , such that for any local martingale $M(t)$, null at zero,*

$$c_p E([M, M](T)^{p/2}) \leq E\left(\left(\sup_{t \leq T} |M(t)|\right)^p\right) \leq C_p E([M, M](T)^{p/2}), \quad (7.41)$$

for $1 \leq p < \infty$. If, moreover, $M(t)$ is continuous, then the result holds also for $0 < p < 1$.

The above inequalities hold when T is a stopping time.

Proofs of these inequalities involve concepts of stochastic calculus for general processes and can be found, for example, in Protter (1992), Rogers and Williams (1990), and Liptser and Shiryaev (1989).

We use the above inequalities to give sufficient conditions for a local martingale to be a true martingale.

Theorem 7.35: Let $M(t)$ be a local martingale, null at zero, such that $E(\sqrt{[M, M](t)}) < \infty$ for all t . Then $M(t)$ is a uniformly integrable martingale on $[0, T]$, for any finite T .

If, moreover, $E[M, M](t) < \infty$, then $M(t)$ is a martingale with $EM^2(t) = E[M, M](t) < \infty$ for all t .

If $\sup_{t < \infty} E[M, M](t) < \infty$, then $M(t)$ is a square integrable martingale.

PROOF. According to the Davis inequality, $\sup_{t \leq T} |M(t)|$ is an integrable random variable, $E(\sup_{t \leq T} |M(t)|) \leq CE(\sqrt{[M, M](T)}) < \infty$. Thus $M(t)$ is dominated by an integrable random variable on any finite time interval. Therefore it is a uniformly integrable martingale by Theorem 7.21, and the first claim is proved.

The condition $E[M, M](t) < \infty$ implies the previous condition $E(\sqrt{[M, M](t)}) < \infty$, as for $X \geq 0$, $E(X) \geq (E(\sqrt{X}))^2$, due to $Var(\sqrt{X}) \geq 0$. Thus M is a martingale. Alternatively, use the Burkholder–Gundy inequality with $p = 2$ which also shows that $M(t)$ is square integrable for each t . Next, recall that by Theorem 7.27, if $M(t)$ is a martingale with $E(M^2(t)) < \infty$, then $M^2(t) - [M, M](t)$ is a martingale. In particular, for any finite t

$$E(M^2(t)) = E[M, M](t),$$

and the second statement is proved. In order to prove the third, notice that since both sides in the above equation are non-decreasing, they have a limit. Since by assumption $\lim_{t \rightarrow \infty} E[M, M](t) < \infty$, $\sup_{t < \infty} EM^2(t) < \infty$, and $M(t)$, $0 \leq t < \infty$ is a square integrable martingale. \square

Application to Itô Integrals

Let $X(t) = \int_0^t H(s)dB(s)$. Being an Itô integral, X is a local martingale. Its quadratic variation is given by $[X, X](t) = \int_0^t H^2(s)ds$. The Burkholder–Gundy inequality with $p = 2$ gives $E(\sup_{t \leq T} X^2(t)) \leq CE([X, X](T)) = E \int_0^T H^2(s)ds$. If $E \left(\int_0^T H^2(s)ds \right) < \infty$, then $X(t)$ is a square integrable martingale. Thus from Theorem 7.35 we recover the known fact that $E(X^2(t)) = E \left(\int_0^t H^2(s)ds \right)$.

The Davis inequality gives

$$E \left(\sup_{t \leq T} \left| \int_0^t H(s)dB(s) \right| \right) \leq CE \left(\sqrt{\int_0^T H^2(s)ds} \right).$$

Thus the condition

$$E \left(\sqrt{\int_0^T H^2(s)ds} \right) < \infty \tag{7.42}$$

is a sufficient condition for the Itô integral to be a martingale and, in particular, to have zero mean. This condition, however, does not assure second moments.

7.8 Continuous Martingales — Change of Time

Brownian motion is the basic continuous martingale from which all continuous martingales can be constructed, either by random change of time, given in this section, or by stochastic integration, as will be seen in Chapter 8. The starting point is a result that characterizes a Brownian motion.

Levy's Characterization of Brownian Motion

Theorem 7.36 (Levy): *A process M with $M(0) = 0$ is a Brownian motion if, and only if, it is a continuous local martingale with quadratic variation process $[M, M](t) = t$.*

PROOF. If M is a Brownian motion, then it is a continuous martingale with $[M, M](t) = t$.

Let $M(t)$ be a continuous local martingale with $[M, M](t) = t$. Then $uM(t)$ is a continuous local martingale with $[uM, uM](t) = u^2t$. We show that

$$U(t) = e^{uM(t)-u^2t/2} = e^{uM(t)-[uM, uM](t)/2} \quad (7.43)$$

is a martingale. Once this is established, the rest of the proof follows by an application of the martingale property.

The general theory of integration with respect to martingales is required to show the martingale property of $U(t)$. It is an easy corollary of a general result on stochastic exponential martingales (Theorem 8.17, Corollary 8.18). Writing the martingale property, we have

$$\mathbb{E}(e^{uM(t)-u^2t/2} | \mathcal{F}_s) = e^{uM(s)-u^2s/2}, \quad (7.44)$$

from which it follows that

$$\mathbb{E}(e^{u(M(t)-M(s))} | \mathcal{F}_s) = e^{u^2(t-s)/2}. \quad (7.45)$$

Since the right hand side of (7.45) is non-random, it follows that $M(t)$ has independent increments. Taking expectation in (7.45), we obtain

$$\mathbb{E}(e^{u(M(t)-M(s))}) = e^{u^2(t-s)/2}, \quad (7.46)$$

which shows that the increment of the martingale $M(t) - M(s)$ has normal distribution with mean zero and variance $(t - s)$. Therefore M is a

continuous process with independent Gaussian increments, hence it is Brownian motion. \square

Example 7.12: Any solution of Tanaka's SDE in Example 5.15 is a Brownian motion (weak uniqueness).

$dX(t) = \text{sign}(X(t))dB(t)$, where $\text{sign}(x) = 1$ if $x \geq 0$ and -1 if $x < 0$. $X(0) = 0$. $X(t) = \int_0^t \text{sign}(X(s))dB(s)$. Since it is an Itô integral, it is a local martingale (even a martingale, as the condition for it to be a martingale holds). It is continuous, and its quadratic variation is given by $[X, X](t) = \int_0^t \text{sign}^2(X(s))ds = t$. Therefore it is a Brownian motion.

Change of Time for Martingales

The main result below states that a continuous martingale M is a Brownian motion with a change of time, where time is measured by the quadratic variation $[M, M](t)$, namely, there is a Brownian motion $B(t)$, such that $M(t) = B([M, M](t))$. This $B(t)$ is constructed from $M(t)$. Define

$$\tau_t = \inf\{s : [M, M](s) > t\}. \quad (7.47)$$

If $[M, M](t)$ is strictly increasing, then τ_t is its inverse.

Theorem 7.37 (Dambis–Dubins–Schwarz (DDS)): Let $M(t)$ be a continuous martingale, null at zero, such that $[M, M](t)$ is non-decreasing to ∞ , and τ_t defined by (7.47). Then the process $B(t) = M(\tau_t)$ is a Brownian motion with respect to the filtration \mathcal{F}_{τ_t} . Moreover, $[M, M](t)$ is a stopping time with respect to this filtration, and the martingale M can be obtained from the Brownian motion B by the change of time $M(t) = B([M, M](t))$. The result also holds when M is a continuous local martingale.

We outline the idea of the proof; for details see, for example, Rogers and Williams (1990), Karatzas and Shreve (1988), Protter (1992), and Revuz and Yor (2001).

PROOF. Let $M(t)$ be a local martingale. τ_t defined by (7.47) are finite stopping times since $[M, M](t) \rightarrow \infty$. Thus \mathcal{F}_{τ_t} are well defined (see Chapter 2 for the definition of \mathcal{F}_τ). Note that $\{[M, M](s) \leq t\} = \{\tau_t \geq s\}$. This implies that $[M, M](s)$ are stopping times for \mathcal{F}_{τ_t} . Since $[M, M](s)$ is continuous $[M, M](\tau_t) = t$. Let $X(t) = M(\tau_t)$. Then it is a continuous local martingale since M and $[M, M]$ have the same intervals of constancy (see the comment following Theorem 7.28). Using Theorem 7.27 we obtain $E[X^2(t)] = E[X, X](t) = E[M, M](\tau_t) = t$. Thus X is a Brownian motion by Levy's characterization in Theorem 7.36. The second part is proven as follows. Recall that M and $[M, M]$ have the same intervals of constancy. Thus $X([M, M](t)) = M(\tau_{[M, M](t)}) = M(t)$. \square

Example 7.13: Let $M(t) = \int_0^t f(s)dB(s)$, with f continuous and non-random. Then M is a Gaussian martingale. Its quadratic variation is given by $[M, M](t) = \int_0^t f^2(s)ds$. For example with $f(s) = s$, $M(t) = \int_0^t s dB(s)$ and $[M, M](t) = \int_0^t s^2 ds = t^3/3$. In this example $[M, M](t)$ is non-random and increasing. τ_t is given by its inverse, $\tau_t = (3t)^{1/3}$. Let $X(t) = M(\tau_t) = \int_0^{\sqrt[3]{3t}} s dB(s)$. Then, clearly, X is continuous, as a composition of continuous functions. It is also a martingale with quadratic variation $\tau_t^3/3 = t$. Hence, by Levy's theorem, it is a Brownian motion, $X(t) = \hat{B}(t)$. According to Theorem 7.37, $M(t) = \hat{B}(t^3/3)$.

Example 7.14: If $M(t) = \int_0^t H(s)dB(s)$ is an Itô integral, then it is a local martingale with quadratic variation $[M, M](t) = \int_0^t H^2(s)ds$. If $\int_0^\infty H^2(s)ds = \infty$, then $M(t) = \hat{B}(\int_0^t H^2(s)ds)$, where $\hat{B}(t)$ is Brownian motion and can be recovered from $M(t)$ with the appropriate change of time.

Example 7.15: (Brownian bridge as time changed Brownian motion)

The SDE for the Brownian bridge (5.35) contains as its only stochastic term $y(t) = \int_0^t \frac{1}{T-s} dB(s)$. Since for any $t < T$, it is a continuous martingale with quadratic variation $[Y, Y](t) = \int_0^t \frac{1}{(T-s)^2} ds = \frac{t}{T(T-t)}$, it follows by the DDS theorem

$$Y(t) = \hat{B}\left(\frac{t}{T(T-t)}\right),$$

for some Brownian motion \hat{B} . Therefore SDE (5.35) has the following representation:

$$X(t) = a\left(1 - \frac{t}{T}\right) + b\frac{t}{T} + (T-t)\hat{B}\left(\frac{t}{T(T-t)}\right), \quad \text{for } 0 \leq t \leq T. \quad (7.48)$$

In this representation $t = T$ is allowed and understood by continuity since the limit of $tB(1/t)$ as $t \rightarrow 0$ is zero by the law of large numbers for Brownian motion.

Change of Time in SDEs

We use the change of time (DDS) theorem for constructing weak solutions of some SDEs. Let

$$X(t) = \int_0^t \sqrt{f'(s)} dB(s), \quad (7.49)$$

where $f(t)$ is an adapted, positive, increasing, differentiable process, null at zero. It is a local martingale with quadratic variation $[X, X](t) = \int_0^t f'(s)ds = f(t)$. Thus $\tau_t = f^{(-1)}(t)$, the inverse of f , and according to the change of time theorem, the process $X(f^{(-1)}(t)) = \hat{B}(t)$ is a Brownian motion (with respect to \mathcal{F}_{τ_t}), and

$$X(t) = \hat{B}(f(t)). \quad (7.50)$$

Thus from Equations (7.49) and (7.50) we have

Theorem 7.38: *Let $f(t)$ be an adapted, positive, increasing, differentiable process, and*

$$dX(t) = \sqrt{f'(t)}dB(t). \quad (7.51)$$

Then the process $\hat{B}(f(t))$ is a weak solution.

We can write Equation (7.51) as follows: for a Brownian motion B , and a function f , there is a Brownian motion \hat{B} such that

$$d\hat{B}(f(t)) = \sqrt{f'(t)}dB(t). \quad (7.52)$$

In the case of non-random change of time in Brownian motion $B(f(t))$, it is easy to check directly that $M(t) = B(f(t))$ is a martingale (with respect to the filtration $\mathcal{F}_{f(t)}$). The quadratic variation of $B(f(t))$ is $[M, M](t) = [B(f), B(f)](t) = f(t)$, was calculated directly in Example 4.24, Equation (4.63).

Example 7.16: (Ornstein–Uhlenbeck process as time changed Brownian motion) With $f(t) = \sigma^2(e^{2\alpha t} - 1)/(2\alpha)$, the process $B(\sigma^2(e^{2\alpha t} - 1)/(2\alpha))$ is a weak solution to the SDE

$$dX(t) = \sigma e^{\alpha t} dB(t).$$

Consider $U(t) = e^{-\alpha t} X(t)$. Integrating by parts, $U(t)$ satisfies

$$dU(t) = -\alpha U(t)dt + \sigma dB(t). \quad (7.53)$$

Recall that the solution to this SDE is given by (5.13). Thus $U(t)$ is an Ornstein–Uhlenbeck process (see Example 5.6). Thus an Ornstein–Uhlenbeck process has representation

$$U(t) = e^{-\alpha t} \hat{B}(\sigma^2(e^{2\alpha t} - 1)/(2\alpha)). \quad (7.54)$$

In order to have $U(0) = x$, take $B(t)$ to be a Brownian motion started at x . Note that in Equations (7.53) and (7.54), $B(t)$ denotes different Brownian motions.

Next we construct a weak solution to the SDEs of the form

$$dX(t) = \sigma(X(t))dB(t)$$

with $\sigma(x) > 0$ such that

$$G(t) = \int_0^t \frac{ds}{\sigma^2(B(s))}$$

is finite for finite t , and increases to infinity, $\int_0^\infty \frac{ds}{\sigma^2(B(s))} = \infty$ almost surely. Then $G(t)$ is adapted, continuous, and strictly increasing to $G(\infty) = \infty$. Therefore it has inverse

$$\tau_t = G^{(-1)}(t). \quad (7.55)$$

Note that for each fixed t , τ_t is a stopping time, as it is the first time the process $G(s)$ hits t , and that τ_t is increasing.

Theorem 7.39: *The process $X(t) = B(\tau_t)$ is a weak solution to the SDE*

$$dX(t) = \sigma(X(t))dB(t). \quad (7.56)$$

PROOF. $X(t) = B(\tau_t) = B(G^{(-1)}(t))$. Using Equation (7.52) with $f = G^{(-1)}$, we obtain

$$dB(G^{(-1)}(t)) = \sqrt{(G^{(-1)})'(t)}d\hat{B}(t).$$

$$(G^{(-1)})'(t) = \frac{1}{G'(G^{(-1)}(t))} = \frac{1}{1/\sigma^2(B(G^{(-1)}(t)))} = \sigma^2(B(\tau_t)). \quad (7.57)$$

Thus we obtain

$$dB(\tau_t) = \sigma^2(B(\tau_t))d\hat{B}(t),$$

and the result is proved. \square

Another proof, by considering the martingale problem, is given next. Note that (7.57) gives $d\tau_t = \sigma^2(B(\tau_t))dt$.

PROOF. The diffusion operator for the SDE (7.56) is given by $Lf(x) = \frac{1}{2}\sigma^2(x)f''(x)$. We show that $X(t) = B(\tau_t)$ is a solution to the martingale problem for L . Indeed, we know (see Example 5.17) that for any twice continuously differentiable function f vanishing outside a compact interval, the process $M(t) = f(B(t)) - \int_0^t \frac{1}{2}f''(B(s))ds$ is a martingale. Since τ_t are increasing stopping times it can be shown (by using the optional stopping Theorem 7.18) that the process $M(\tau_t)$ is also a martingale, that is, $f(B(\tau_t)) - \int_0^{\tau_t} \frac{1}{2}f''(B(s))ds$ is a martingale. Now perform the change of variable $s = \tau_u$, and observe from (7.57) that $d\tau_t = \sigma^2(B(\tau_t))dt$, to obtain that the process $f(B(\tau_t)) - \int_0^t \frac{1}{2}\sigma^2(B(\tau_u))f''(B(\tau_u))du$ is a martingale. However, since $X(t) = B(\tau_t)$, this being the same as $f(X(t)) - \int_0^t \frac{1}{2}\sigma^2(X(u))f''(X(u))du$ is a martingale, and $X(t)$ solves the martingale problem for L . \square

An application of Theorem 7.37 gives a result on uniqueness of the solution of SDE (7.56). This result is weaker than Theorem 6.13 of Engelbert–Schmidt.

Theorem 7.40: *Let $\sigma(x)$ be a positive function bounded away from zero, $\sigma(x) \geq \delta > 0$. Then the stochastic differential Equation (7.56) has a unique weak solution.*

PROOF. Let $X(t)$ be a weak solution to (7.56). Then $X(t)$ is a local martingale and there is a Brownian motion $\beta(t)$, such that $X(t) = \beta([X, X](t))$. Now

$$[X, X](t) = \int_0^t \sigma^2(X(s))ds = \int_0^t \sigma^2(\beta([X, X](s)))ds.$$

Thus $[X, X](t)$ is a solution to the ODE $da(t) = \sigma^2(\beta(a(t)))dt$. Since the solution to this ODE is unique, the solution to (7.56) is unique. \square

A more general change of time is done for the SDE

$$dX(t) = \mu(X(t))dt + \sigma(X(t))dB(t). \quad (7.58)$$

Let $g(x)$ be a positive function for which $G(t) = \int_0^t g(X(s))ds$ is finite for finite t and increases to infinity almost surely. Define $\tau_t = G^{(-1)}(t)$.

Theorem 7.41: *Let $X(t)$ be a solution to (7.58) and define $Y(t) = X(\tau_t)$. Then $Y(t)$ is a weak solution to the SDE*

$$dY(t) = \frac{\mu(Y(t))}{g(Y(t))}dt + \frac{\sigma(Y(t))}{\sqrt{g(Y(t))}}dB(t), \quad \text{with } Y(0) = X(0). \quad (7.59)$$

One can use the change of time on an interval $[0, T]$, for a stopping time T .

Example 7.17: (Lamperti's change of time)

Let $X(t)$ satisfy the SDE (Feller's branching diffusion)

$$dX(t) = \mu X(t)dt + \sigma \sqrt{X(t)}dB(t), \quad X(0) = x > 0, \quad (7.60)$$

with positive constants μ and σ . Lamperti's change of time is $G(t) = \int_0^t X(s)ds$. Here $g(x) = x$. Then $Y(t) = X(\tau_t)$ satisfies the SDE

$$\begin{aligned} dY(t) &= \frac{\mu Y(t)}{Y(t)}dt + \frac{\sigma \sqrt{Y(t)}}{\sqrt{Y(t)}}dB(t), \\ &= \mu dt + \sigma dB(t) \quad \text{with } Y(0) = x, \end{aligned}$$

and

$$Y(t) = x + \mu t + \sigma B(t). \quad (7.61)$$

In other words, with a random change of time, the branching diffusion is a Brownian motion with drift. At the (random) point where $G(t)$ stops increasing its inverse τ_t , defined as the right-inverse, $\tau_t = \inf\{s : G(s) = t\}$, also remains the same. This happens at the point of time when $X(t) = 0$. It can be seen that once the process is at zero, it stays at zero forever. Let $T = \inf\{t : X(t) = 0\}$. T is a stopping time, and $Y(t)$ is the Brownian motion stopped at that time.

The other direction is also true, a branching diffusion can be obtained from a Brownian motion with drift. Let $Y(t)$ satisfy (7.61), and let $T = \inf\{t : Y(t) = 0\}$. $Y(t) > 0$ for $t \leq T$. Define

$$G(t) = \int_0^{t \wedge T} \frac{1}{Y(s)}ds,$$

and let τ_t be the inverse of G , which is well-defined on $(0, T)$. Then $X(t) = Y(\tau_t)$ satisfies the SDE (7.60) stopped when it hits zero.

Remark 7.4: Any solution to an SDE with time-independent coefficients can be obtained from Brownian motion by using change of variables and random time change (Gihman and Skorohod (1972)).

There are three main methods used for solving SDEs: change of state space, that is, change of variable (Itô's formula); change of time; and change of measure. We have seen examples of SDEs solved by using change of variables and change of time. The change of measure approach will be covered later.

Notes: Material for this chapter is based on Protter (1992), Rogers and Williams (1990), Gihman and Skorohod (1972), Liptser and Shiryaev (1974), (1989), and Revuz and Yor (2001).

7.9 Exercises

Exercise 7.1: Let $M(t)$ be an \mathcal{F}_t -martingale and denote its natural filtration by \mathcal{G}_t . Show that $M(t)$ is a \mathcal{G}_t -martingale.

Exercise 7.2: Show that an increasing integrable process is a submartingale.

Exercise 7.3: Show that if $X(t)$ is a submartingale and g is a non-decreasing convex function such that $E|g(X(t))| < \infty$, then $g(X(t))$ is a submartingale.

Exercise 7.4: Show that $M(t)$ is a square integrable martingale if, and only if, $M(t) = E(Y|\mathcal{F}_t)$, where Y is square integrable, $E(Y^2) < \infty$.

Exercise 7.5: (Expected exit time of Brownian motion from (a, b))
Let $B(t)$ be a Brownian motion started at $x \in (a, b)$, and $\tau = \inf\{t : B(t) = a \text{ or } b\}$. By stopping the martingale $M(t) = B(t)^2 - t$, show that $E_x(\tau) = (x - a)(b - x)$.

Exercise 7.6: Find the probability of $B(t) - t/2$ reaching a before it reaches b when started at x , $a < x < b$. Hint: use the exponential martingale $M(t) = e^{B(t)-t/2}$.

Exercise 7.7: Find the expected length of the game in gambler's ruin, when

1. Betting is done on a fair coin.
2. Betting is done on a biased coin.

Exercise 7.8: Give the probability of ruin when playing a game of chance against an infinitely rich opponent (with initial capital $b \rightarrow \infty$).

Exercise 7.9: (Ruin probability in insurance)

A discrete time risk model for the surplus U_n of an insurance company at the end of year n , $n = 1, 2, \dots$ is given by $U_n = U_0 + cn - \sum_{k=1}^n X_k$, where c is the total annual premium and X_k is the total (aggregate) claim in year k . The time of ruin T is the first time when the surplus becomes negative, $T = \min\{n : U_n < 0\}$, with $T = \infty$ if $U_n \geq 0$ for all n . Assume that $\{X_k, k = 1, 2, \dots\}$ are i.i.d. random variables, and there exists a constant $R > 0$ such that $E(e^{-R(c-X_1)}) = 1$. Show that for all n , $P_x(T \leq n) \leq e^{-Rx}$, where $U_0 = x$ the initial funds, and the ruin probability $P_x(T < \infty) \leq e^{-Rx}$. Hint: show that $M_n = e^{-RU_n}$ is a martingale, and use the optional stopping theorem.

Exercise 7.10: (Ruin probability in insurance continued)

Find the bound on the ruin probability when the aggregate claims have $N(\mu, \sigma^2)$ distribution. Give the initial amount x required to keep the ruin probability below level α .

Exercise 7.11: Let $B(t)$ be a Brownian motion starting at zero and T be the first exit time from $(-1, 1)$, that is, the first time when $|B|$ takes value 1. Use the Davis inequality to show that $E(\sqrt{T}) < \infty$.

Exercise 7.12: Let $B(t)$ be a Brownian motion, $X(t) = \int_0^t \text{sign}(B(s)) dB(s)$. Show that X is also a Brownian motion.

Exercise 7.13: Let $M(t) = \int_0^t e^s dB(s)$. Find $g(t)$ such that $M(g(t))$ is a Brownian motion.

Exercise 7.14: Let $B(t)$ be a Brownian motion. Give an SDE for $e^{-\alpha t} B(e^{2\alpha t})$.

Exercise 7.15: Prove the change of time result in SDEs, Theorem 7.41.

Exercise 7.16: Let $X(t)$ satisfy SDE $dX(t) = \mu(t)dt + \sigma(t)dB(t)$ on $[0, T]$. Show that $X(t)$ is a local martingale if, and only if, $\mu(t) = 0$ a.e.

Exercise 7.17: $f(x, t)$ is differentiable in t and twice in x . It is known that $X(t) = f(B(t), t)$ is of finite variation. Show that f is a function of t alone.

Exercise 7.18: Let $Y(t) = \int_0^t B(s)dB(s)$ and $W(t) = \int_0^t \text{sign}(B(s))dB(s)$. Show that $dY(t) = \sqrt{t + 2Y(t)}dW(t)$. Show uniqueness of the weak solution of the above SDE.

Chapter 8

Calculus For Semimartingales

In this chapter rules of calculus are given for the most general processes for which stochastic calculus is developed, called semimartingales. A semimartingale is a process consisting of a sum of a local martingale and a finite variation process. Integration with respect to semimartingales involves integration with respect to local martingales, and these integrals generalize the Itô integral where integration is done with respect to a Brownian motion. Important concepts, such as compensators and the sharp bracket processes, are introduced, and Itô's formula in its general form is given.

8.1 Semimartingales

In stochastic calculus only regular processes are considered. These are either continuous processes, right-continuous with left limits, or left-continuous with right limits. The regularity of the process implies that it can have at most countably many discontinuities, and all of them are jumps (Chapter 1). The definition of a semimartingale presumes a given filtration and processes which we consider are adapted to it. Following the classical approach (see, for example, Métivier (1982), Liptser and Shiryaev (1989) p.85) a semimartingale is a local martingale plus a process of finite variation. More precisely,

Definition 8.1: A regular right-continuous with left limits (*càdlàg*) adapted process $S(t)$ is a semimartingale if it can be represented as a sum of two processes: a local martingale $M(t)$ and a process of finite variation $A(t)$, with $M(0) = A(0) = 0$, and

$$S(t) = S(0) + M(t) + A(t). \tag{8.1}$$

Example 8.1: (Semimartingales)

1. $S(t) = B^2(t)$, where $B(t)$ is a Brownian motion, is a semimartingale. $S(t) = M(t) + t$, where $M(t) = B^2(t) - t$ is a martingale and $A(t) = t$ is a finite variation process.
2. $S(t) = N(t)$, where $N(t)$ is a Poisson process with rate λ , is a semimartingale as it is a finite variation process.
3. One way to obtain semimartingales from known semimartingales is by applying a twice continuously differentiable (C^2) transformation. If $S(t)$ is a semimartingale and f is a C^2 function, then $f(S(t))$ is also a semimartingale. The decomposition of $f(S(t))$ into martingale part and finite variation part is given by Itô's formula, given later. In this way we can assert that, for example, the geometric Brownian motion $e^{\sigma B(t)+\mu t}$ is a semimartingale.
4. A right-continuous with left limits (*càdlàg*) deterministic function $f(t)$ is a semimartingale if, and only if, it is of finite variation. Thus $f(t) = t \sin(1/t)$, $t \in (0, 1]$, $f(0) = 0$ is continuous, but not a semimartingale (see Example 1.7).
5. A diffusion, that is, a solution to a stochastic differential equation with respect to Brownian motion, is a semimartingale. Indeed, the Itô integral with respect to $dB(t)$ is a local martingale, and the integral with respect to dt is a process of finite variation.
6. Although the class of semimartingales is rather large, there are processes which are not semimartingales. Examples are: $|B(t)|^\alpha$, $0 < \alpha < 1$, where $B(t)$ is the one-dimensional Brownian motion; $\int_0^t (t-s)^{-\alpha} dB(s)$, $0 < \alpha < 1/2$. It requires analysis to show that the above processes are not semimartingales.

For a semimartingale X , the process of jumps ΔX is defined by

$$\Delta X(t) = X(t) - X(t-), \quad (8.2)$$

and represents the jump at point t . If X is continuous, then, of course, $\Delta X = 0$.

8.2 Predictable Processes

In this section we describe the class of predictable processes. This class of processes has a central role in the theory. In particular, only predictable processes can be integrated with respect to a semimartingale. Recall that in discrete time a process H is predictable if H_n is \mathcal{F}_{n-1} measurable, that is, H is known with certainty at time n on the basis of information up to time $n-1$. Predictability in continuous time is harder to define. We recall some general definitions of processes starting with the class of adapted processes.

Definition 8.2: A process X is called adapted to filtration $\mathbb{F} = (\mathcal{F}_t)$, if for all t , $X(t)$ is \mathcal{F}_t -measurable.

In the construction of the stochastic integral $\int_0^t H(u)dS(u)$, processes H and S are taken to be adapted to \mathbb{F} . For a general semimartingale S , the requirement that H is adapted is too weak, it fails to assure measurability of some basic constructions. H must be *predictable*. The exact definition of predictable processes involves σ -fields generated on $\mathbb{R}^+ \times \Omega$ and is given later in Chapter 8.13. Note that left-continuous processes are predictable, in the sense that $H(t) = \lim_{s \uparrow t} H(s) = H(s-)$. So that if the values of the process before t are known, then the value at t is determined by the limit. For our purposes it is enough to describe a subclass of predictable processes which can be defined constructively.

Definition 8.3: H is predictable if it is one of the following:

1. A left-continuous adapted process, in particular, a continuous adapted process.
2. A limit (almost sure, in probability) of left-continuous adapted processes.
3. A regular right-continuous process such that, for any stopping time τ , H_τ is $\mathcal{F}_{\tau-}$ -measurable, the σ -field generated by the sets $A \cap \{T < t\}$, where $A \in \mathcal{F}_t$.
4. A Borel-measurable function of a predictable process.

Example 8.2: Poisson process $N(t)$ is right-continuous and is obviously adapted to its natural filtration. It can be shown, see Example 8.31, that it is not predictable. Its left-continuous modification $N(t-) = \lim_{s \uparrow t} N(s)$ is predictable because it is adapted and left-continuous by (1). Any measurable function (even right-continuous) of $N(t-)$ is also predictable by (2).

Example 8.3: Right-continuous adapted processes may not be predictable, even though they can be approached by left-continuous processes, for example $X(t) = \lim_{\epsilon \rightarrow 0} X((t + \epsilon)-)$.

Example 8.4: Let τ be a stopping time. This means that for any t , the set $\{\tau \leq t\} \in \mathcal{F}_t$. Consider the indicator of random interval process $X(t) = I_{[0,\tau]}(t)$. $X(t) = 1$ if, and only if, $\omega \in \{t \leq \tau\} = \{\tau < t\}^c \in \mathcal{F}_t$ (see Chapter 2). Hence it is adapted. $X(t)$ is also left-continuous. Thus it is a predictable process by (1).

Example 8.5: It will be seen later that when filtration is generated by Brownian motion, then any right-continuous adapted process is predictable. This is why in the definition of the Itô integral right-continuous functions are allowed as integrands.

8.3 Doob–Meyer Decomposition

Recall that a process is a submartingale if for all $s < t$, $E(X(t)|\mathcal{F}_s) \geq X(s)$ almost surely.

Theorem 8.4: *If X is a submartingale or a local submartingale, then there exists a local martingale $M(t)$ and a unique increasing predictable process $A(t)$, locally integrable, such that*

$$X(t) = X(0) + M(t) + A(t). \quad (8.3)$$

If $X(t)$ is a submartingale of Dirichlet class (D) (see Definition 7.25), then the process A is integrable, that is, $\sup_t \mathbb{E}A(t) < \infty$, and $M(t)$ is a uniformly integrable martingale.

Example 8.6:

1. Let $X(t) = B^2(t)$ on a finite interval $t \leq T$. $X(t)$ is a submartingale. Decomposition (8.3) holds with $M(t) = B^2(t) - t$ and $A(t) = t$. Since the interval is finite M is uniformly integrable and A is integrable.
2. Let $X(t) = B^2(t)$ on the infinite interval $t \geq 0$. Then (8.3) holds with $M(t) = B^2(t) - t$ and $A(t) = t$. Since the interval is infinite M is a martingale and A is locally integrable (take for example the localizing sequence $\tau_n = n$).
3. Let $X(t) = N(t)$ be a Poisson process with intensity λ . Then X is a submartingale. Decomposition (8.3) holds with $M(t) = N(t) - \lambda t$ and $A(t) = \lambda t$.

Proof of Theorem 8.4 can be found in Rogers and Williams (1990), Delacherie (1972), and Meyer (1966).

Doob's Decomposition

For processes in discrete time, decomposition (8.3) is due to Doob, and it is simple to obtain. Indeed, if X_n is a submartingale, then clearly, $X_{n+1} = X_0 + \sum_{i=0}^n (X_{i+1} - X_i)$. By adding and subtracting $\mathbb{E}(X_{i+1}|\mathcal{F}_i)$ we obtain the Doob decomposition

$$X_{n+1} = X_0 + \sum_{i=0}^n ((X_{i+1} - \mathbb{E}(X_{i+1}|\mathcal{F}_i)) + \sum_{i=0}^n (\mathbb{E}(X_{i+1}|\mathcal{F}_i) - X_i), \quad (8.4)$$

where martingale and an increasing process are given by

$$M_{n+1} = \sum_{i=0}^n (X_{i+1} - \mathbb{E}(X_{i+1}|\mathcal{F}_i)) \quad \text{and} \quad A_{n+1} = \sum_{i=0}^n (\mathbb{E}(X_{i+1}|\mathcal{F}_i) - X_i). \quad (8.5)$$

A_n is increasing due to the submartingale property, $\mathbb{E}(X_{i+1}|\mathcal{F}_i) - X_i \geq 0$ for all i . It is also predictable because $\mathbb{E}(X_{n+1}|\mathcal{F}_n)$ and all other terms are \mathcal{F}_n measurable.

It is much harder to prove decomposition (8.3) in continuous time and this was done by Meyer.

8.4 Integrals With Respect to Semimartingales

In this section the stochastic integral $\int_0^T H(t)dS(t)$ is defined, where $S(t)$ is a semimartingale. Due to representation $S(t) = S(0) + M(t) + A(t)$ the integral with respect to $S(t)$ is the sum of two integrals, one with respect to a local martingale $M(t)$ and the other with respect to a finite variation process $A(t)$. The integral with respect to $A(t)$ can be done path by path as the Stieltjes integral, since $A(t)$, although random, is of finite variation.

The integral with respect to the martingale $M(t)$ is new, it is the stochastic integral $\int_0^T H(t)dM(t)$. When $M(t)$ is Brownian motion $B(t)$, it is the Itô integral, defined in Chapter 4. But now martingales are allowed to have jumps and this makes the theory more complicated. The key property used in the definition of the Itô integral is that on finite intervals Brownian motion is a square integrable martingale. This property in its local form plays an important role in the general case. Conditions for the existence of the integral with respect to a martingale involves the martingale's quadratic variation, which was introduced in Section 7.6.

Stochastic Integral With Respect to Martingales

For a simple predictable process $H(t)$, given by

$$H(t) = H(0)I_0 + \sum_{i=0}^{n-1} H_i I_{(T_i, T_{i+1}]}(t), \quad (8.6)$$

where $0 = T_0 \leq T_1 \leq \dots \leq T_n = T$ are stopping times and H_i 's are \mathcal{F}_{T_i} -measurable, the stochastic integral is defined as the sum

$$\int_0^T H(t)dM(t) = \sum_{i=0}^{n-1} H_i (M(T_{i+1}) - M(T_i)). \quad (8.7)$$

If $M(t)$ is a locally square integrable martingale, then by the L^2 theory (Hilbert space theory) one can extend the stochastic integral from simple predictable processes to the class of predictable processes H such that

$$\int_0^T H^2(t)d[M, M](t) < \infty \text{ almost surely,} \quad (8.8)$$

and $\int_0^T H^2(t)d[M, M](t)$ is locally integrable, i.e. there is a sequence of stopping times $\tau_n \rightarrow \infty$ such that

$$\mathbb{E} \int_0^{T \wedge \tau_n} H^2(t)d[M, M](t) < \infty.$$

Properties of Stochastic Integrals With Respect to Martingales

1. Local martingale property. If $M(t)$ is a local martingale, the integral $\int_0^t H(s)dM(s)$ is a local martingale.
2. Isometry property. If $M(t)$ is a square integrable martingale, and H satisfies

$$\mathbb{E}\left(\int_0^T H^2(s)d[M, M](s)\right) < \infty, \quad (8.9)$$

then $\int_0^t H(s)dM(s)$ is a square integrable martingale with zero mean and variance

$$\mathbb{E}\left(\int_0^t H(s)dM(s)\right)^2 = \mathbb{E}\left(\int_0^t H^2(s)d[M, M](s)\right). \quad (8.10)$$

3. If a local martingale $M(t)$ is of finite variation, then the stochastic integral is indistinguishable from the Stieltjes integral.

Example 8.7: Consider Itô integrals with respect to Brownian motion. Since $B(t)$ is a square integrable martingale on $[0, T]$, with $[B, B](t) = t$, we recover that for a predictable H , such that $\mathbb{E} \int_0^T H^2(s)ds < \infty$, $\int_0^t H(s)dB(s)$ is a square integrable martingale with zero mean and variance $\int_0^t \mathbb{E}H^2(s)ds$.

Stochastic Integrals With Respect to Semimartingales

Let S be a semimartingale with representation

$$S(t) = S(0) + M(t) + A(t), \quad (8.11)$$

where M is a local martingale and A is a finite variation process. Let H be a predictable process such that conditions (8.12) and (8.8) hold.

$$\int_0^T |H(t)|dV_A(t) < \infty, \quad (8.12)$$

where $V_A(t)$ is the variation process of A . Then the stochastic integral is defined as the sum of integrals,

$$\int_0^t H(t)dS(t) = \int_0^T H(t)dM(t) + \int_0^T H(t)dA(t). \quad (8.13)$$

Since a representation of a semimartingale (8.11) is not unique, one should check that the stochastic integral does not depend on the representation used. Indeed, if $S(t) = S(0) + M_1(t) + A_1(t)$ is another representation,

then $(M - M_1)(t) = -(A - A_1)(t)$. So that $M - M_1$ is a local martingale of finite variation. However, for such martingales stochastic and Stieltjes integrals are the same, and it follows that $\int H(t)dM_1(t) + \int H(t)dA_1(t) = \int H(t)dM(t) + \int H(t)dA(t) = \int H(t)dS(t)$.

Since the integral with respect to a local martingale is a local martingale, and the integral with respect to a finite variation process is a process of finite variation, it follows that a stochastic integral with respect to a semimartingale is a semimartingale.

For details see Liptser and Shiryaev (1989).

Example 8.8: Let $N(t)$ be a Poisson process. $N(t)$ is of finite variation and the integral $\int_0^T N(t)dN(t)$ is well-defined as a Stieltjes integral, $\int_0^T N(t)dN(t) = \sum_{\tau_i \leq T} N(\tau_i)$, where τ_i 's are the jumps of $N(t)$. However, $\int_0^T N(t)dN(t)$ is not the stochastic integral since $N(t)$ is not predictable, but $\int_0^T N(t-)dN(t)$ is. It is indistinguishable from the integral in the sense of Stieltjes $\int_0^T N(t-)dN(t) = \sum_{\tau_i \leq T} N(\tau_{i-1})$.

Properties of Stochastic Integrals With Respect to Semimartingales

Let X be a semimartingale and H a predictable process such that the stochastic integral exists for $0 \leq t \leq T$, and denote

$$(H \cdot X)(t) := \int_0^t H(s)dX(s).$$

Then the stochastic integral $H \cdot X$ has the following properties:

1. The jumps of the integral occur at the points of jumps of X , and $\Delta(H \cdot X)(t) = H(t)\Delta X(t)$. In particular, a stochastic integral with respect to a continuous semimartingale is continuous.
2. If τ is a stopping time, then the stopped integral is the integral with respect to the stopped semimartingale,

$$\int_0^{t \wedge \tau} H(s)dX(s) = \int_0^t H(s)I(s \leq \tau)dX(s) = \int_0^t H(s)dX(s \wedge \tau).$$

3. If X is of finite variation, then $\int_0^t H(s)dX(s)$ is indistinguishable from the Stieltjes integral computed path by path.
4. Associativity. If $Y(t) = \int_0^t H(s)dX(s)$ is a semimartingale, and if K is a predictable process such that $(K \cdot Y)(t) = \int_0^t K(s)dY(s)$ is defined, then $K \cdot Y = K \cdot (H \cdot X) = (KH) \cdot X$, that is,

$$\int_0^t K(s)dY(s) = \int_0^t K(s)H(s)dX(s).$$

8.5 Quadratic Variation and Covariation

If X, Y are semimartingales on the common space, then the quadratic covariation process, also known as the *square bracket* process and denoted $[X, Y](t)$, is defined, as usual, by

$$[X, Y](t) = \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} (X(t_{i+1}^n) - X(t_i^n))(Y(t_{i+1}^n) - Y(t_i^n)), \quad (8.14)$$

where the limit is taken over shrinking partitions $\{t_i^n\}_{i=0}^n$ of the interval $[0, t]$ when $\delta_n = \max_i(t_{i+1}^n - t_i^n) \rightarrow 0$ and is in probability. Taking $Y = X$ we obtain the quadratic variation process of X .

Example 8.9: We have seen that quadratic variation of Brownian motion $B(t)$ is $[B, B](t) = t$ and of Poisson process $N(t)$ is $[N, N](t) = N(t)$.

Properties of Quadratic Variation

We give the fundamental properties of the quadratic variation process with some explanations, but omit the proofs.

1. If X is a semimartingale, then $[X, X]$ exists and is an adapted process.
2. It is clear from the definition that quadratic variation over non-overlapping intervals is the sum of the quadratic variation over each interval. As such, $[X, X](t)$ is a non-decreasing function of t . Consequently $[X, X](t)$ is a function of finite variation.
3. It follows from the definition (8.14) that $[X, Y]$ is bilinear and symmetric, that is, $[X, Y] = [Y, X]$ and

$$[\alpha X + Y, \beta U + V] = \alpha\beta[X, U] + \alpha[X, V] + \beta[Y, U] + [Y, V]. \quad (8.15)$$

4. Polarization identity.

$$[X, Y] = \frac{1}{2} \left([X + Y, X + Y] - [X, X] - [Y, Y] \right). \quad (8.16)$$

This property follows directly from the previous one.

5. $[X, Y](t)$ is a regular right-continuous (*càdlàg*) function of finite variation. This follows from the polarization identity, as $[X, Y]$ is the difference of two increasing functions.
6. The jumps of the quadratic covariation process occur only at points where *both* processes have jumps,

$$\Delta[X, Y](t) = \Delta X(t)\Delta Y(t). \quad (8.17)$$

7. If one of the processes X or Y is of finite variation, then

$$[X, Y](t) = \sum_{s \leq t} \Delta X(s) \Delta Y(s). \quad (8.18)$$

Note that although the summation is taken over all s not exceeding t , there are at most countably many terms different from zero.

The following property is frequently used, it follows directly from (8.18).

Corollary 8.5: *If $X(t)$ is a continuous semimartingale of finite variation, then it has zero quadratic covariation with any other semimartingale $Y(t)$.*

Quadratic Variation of Stochastic Integrals

The quadratic covariation of stochastic integrals has the following property:

$$\left[\int_0^{\cdot} H(s) dX(s), \int_0^{\cdot} K(s) dY(s) \right](t) = \int_0^t H(s) K(s) d[X, Y](s). \quad (8.19)$$

In particular, the quadratic variation of a stochastic integral is given by

$$\left[\int_0^{\cdot} H(s) dX(s), \int_0^{\cdot} H(s) dX(s) \right](t) = \int_0^t H^2(s) d[X, X](s), \quad (8.20)$$

and

$$\begin{aligned} \left[\int_0^{\cdot} H(s) dX(s), Y \right](t) &= \left[\int_0^{\cdot} H(s) dX(s), \int_0^{\cdot} 1 dY(s) \right](t) \\ &= \int_0^t H(s) d[X, Y](s). \end{aligned} \quad (8.21)$$

Quadratic variation has a representation in terms of the stochastic integral.

Theorem 8.6: *Let X be a semimartingale null at zero. Then*

$$[X, X](t) = X^2(t) - 2 \int_0^t X(s-) dX(s). \quad (8.22)$$

For a partition $\{t_i^n\}$ of $[0, t]$, consider

$$v_n(t) = \sum_{i=0}^{n-1} (X(t_{i+1}^n) - X(t_i^n))^2.$$

For any fixed t the sequence $v_n(t)$ converges in probability to a limit $[X, X](t)$ as $\max_{i \leq n} (t_{i+1}^n - t_i^n) \rightarrow 0$. Moreover, there is a subsequence n_k such that the processes $v_{n_k}(t)$ converge uniformly on any bounded time interval to the process $X^2(t) - 2 \int_0^t X(s-) dX(s)$.

We justify (8.22) heuristically. By opening the brackets in the sum for $v_n(t)$, adding and subtracting $X^2(t_i)$, we obtain

$$v_n(t) = \sum_{i=0}^{n-1} (X^2(t_{i+1}^n) - X^2(t_i^n)) - 2 \sum_{i=0}^{n-1} X(t_i^n)(X(t_{i+1}^n) - X(t_i^n)).$$

The first sum is $X^2(t)$. The second has for its limit in probability $\int_0^t X(s-)dX(s)$. This argument can be made into a rigorous proof, see for example Métivier (1982). Alternatively, (8.22) can be established by using Itô's formula.

Using the polarization identity, it is easy to see

Corollary 8.7: *For semimartingales X, Y the quadratic covariation process is given by*

$$[X, Y](t) = X(t)Y(t) - X(0)Y(0) - \int_0^t X(s-)dY(s) - \int_0^t Y(s-)dX(s). \quad (8.23)$$

This is also known as the integration by parts or product rule formula.

8.6 Itô's Formula for Continuous Semimartingales

If $X(t)$ is a *continuous* semimartingale and f is a twice continuously differentiable function, then $Y(t) = f(X(t))$ is a semimartingale and admits the following representation:

$$f(X(t)) - f(X(0)) = \int_0^t f'(X(s))dX(s) + \frac{1}{2} \int_0^t f''(X(s))d[X, X](s). \quad (8.24)$$

In differential form this is written as

$$df(X(t)) = f'(X(t))dX(t) + \frac{1}{2}f''(X(t))d[X, X](t). \quad (8.25)$$

It follows, in particular, that $f(X(t))$ is also a semimartingale, and its decomposition into the martingale part and the finite variation part can be obtained from Itô's formula by splitting the stochastic integral with respect to $X(t)$ into the integral with respect to a local martingale $M(t)$ and a finite variation process $A(t)$.

We have given a justification of Itô's formula and examples of its use in Chapters 4 and 6.

Remark 8.1:

1. The differentiability properties of f may be relaxed. If, for example, X is of finite variation, then f needs to be only once continuously differentiable. f can be defined only on an open set rather than a whole line, but then X must take its values almost surely in this set. For example, if X is a positive semimartingale, then Itô's formula can be used with $f = \ln$.
2. Itô's formula holds for convex functions (Protter (1992)), and more generally, for functions which are the difference of two convex functions. This is the Meyer–Itô (Itô–Tanaka) formula (see for example Protter (1992), Rogers and Williams (1990), and Revuz and Yor (2001)). In particular, if f is a convex function on \mathbb{R} and $X(t)$ is a semimartingale, then $f(X(t))$ is also a semimartingale. See also Chapter 8.7.
3. It follows from Itô's formula that if a semimartingale X is continuous with nil quadratic variation $[X, X](t) = 0$, then the differentiation rule is the same as in the ordinary calculus. If $X(t)$ is a Brownian motion, then $d[X, X](t) = dt$, and we recover formulae (4.39) and (4.53). If X has jumps, then the formula has an extra term (see Chapter 8.10).

The following result is a direct corollary to Itô's formula.

Corollary 8.8: *Let $X(t)$ be a continuous semimartingale and f be twice continuously differentiable. Then*

$$[f(X), f(X)](t) = \int_0^t (f'(X(s)))^2 d[X, X](s). \quad (8.26)$$

PROOF. Since $[X, X]$ is of finite variation it follows from (8.24)

$$[f(X), f(X)](t) = \left[\int_0^\cdot f'(X(s))dX(s), \int_0^\cdot f'(X(s))dX(s) \right](t). \quad \square$$

Itô's Formula for Functions of Several Variables

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be C^2 , and let $\mathbf{X}(t) = (X_1(t), \dots, X_n(t))$ be a *continuous* semimartingale in \mathbb{R}^n , that is, each X_i is a continuous semimartingale. Then $f(\mathbf{X})$ is a semimartingale and has the following representation:

$$\begin{aligned} f(\mathbf{X}(t)) - f(\mathbf{X}(0)) &= \int_0^t \sum_{i=1}^n \frac{\partial f}{\partial x_i}(\mathbf{X}(s))dX_i(s) \\ &\quad + \frac{1}{2} \int_0^t \sum_{i=1}^n \sum_{j=1}^n \frac{\partial^2 f}{\partial x_i \partial x_j}(\mathbf{X}(s))d[X_i, X_j](s). \end{aligned} \quad (8.27)$$

8.7 Local Times

Let $X(t)$ be a continuous semimartingale. Consider $|X(t) - a|$, $a \in \mathbb{R}$. The function $|x - a|$ is not differentiable at a , but at any other point its derivative is given by $\text{sign}(x - a)$, where $\text{sign}(x) = 1$ for $x > 0$ and $\text{sign}(x) = -1$ for $x \leq 0$. It is possible to extend Itô's formula for this case and prove Tanaka's formula (see Rogers and Williams (1990), and Protter (1992)).

Theorem 8.9 (Tanaka's Formula): *Let $X(t)$ be a continuous semimartingale. Then for any $a \in \mathbb{R}$ there exists a continuous non-decreasing adapted process $L^a(t)$, called the local time at a of X , such that*

$$|X(t) - a| = |X(0) - a| + \int_0^t \text{sign}(X(s) - a) dX(s) + L^a(t). \quad (8.28)$$

As a function in a , $L^a(t)$ is right-continuous with left limits. For any fixed a as a function in t $L^a(t)$ increases only when $X(t) = a$, that is, $L^a(t) = \int_0^t I(X(s) = a) dL^a(s)$. Moreover, if $X(t)$ is a continuous local martingale, then $L^a(t)$ is jointly continuous in a and t .

Remark 8.2: Heuristically, Tanaka's formula can be justified by a formal application of Itô's formula to the function $\text{sign}(x)$. The derivative of $\text{sign}(x)$ is zero everywhere but at zero, where it is not defined. However, it is possible to define the derivative as a generalized function or a Schwartz distribution, in which case it is equal to 2δ , where δ is the Dirac delta function. Thus the second derivative of $|x - a|$ is $2\delta(x - a)$ in the generalized function sense. The local time at a of X is defined as $L^a(t) = \int_0^t \delta(X(s) - a) ds$. Formal use of Itô's formula gives (8.28).

Theorem 8.10 (Occupation Times Formula): *Let $X(t)$ be a continuous semimartingale with local time $L^a(t)$. Then for any bounded measurable function $g(x)$*

$$\int_0^t g(X(s)) d[X, X](s) = \int_{-\infty}^{\infty} g(a) L^a(t) da. \quad (8.29)$$

In particular,

$$[X, X](t) = \int_{-\infty}^{\infty} L^a(t) da. \quad (8.30)$$

Example 8.10: Let $X(t) = B(t)$ be Brownian motion. Then its local time at zero process, $L^0(t)$ satisfies (Tanaka's formula)

$$L^0(t) = |B(t)| - \int_0^t \text{sign}(B(s)) dB(s). \quad (8.31)$$

The occupation times formula (8.29) becomes

$$\int_0^t g(B(s))ds = \int_{-\infty}^{\infty} g(a)L^a(t)da. \quad (8.32)$$

The time Brownian motion spends in a set $A \subset \mathbb{R}$ up to time t is given by (with $g(x) = I_A(x)$)

$$\int_0^t I_A(B(s))ds = \int_{-\infty}^{\infty} I_A(a)L^a(t)da = \int_A L^a(t)da. \quad (8.33)$$

Remark 8.3: Taking $A = (a, a+da)$ and $g(x) = I_{(a,a+da)}(x)$ its indicator in (8.32), $L^a(t)da$ is the time Brownian motion spends in $(a, a+da)$ up to time t , which explains the name “local time”. The time Brownian motion spends in a set A is $\int_A L^a(t)da$, hence the name “occupation times density” formula (8.33). For a continuous semimartingale the formula (8.29) is the “occupation times density” formula relative to the random “clock” $d[X, X](s)$.

Example 8.11: $X(t) = |B(t)|$ is a semimartingale, since $|x|$ is a convex function. Its decomposition into the martingale and finite variation parts is given by Tanaka’s formula (8.31). $\sqrt{|B(t)|}$ is not a semimartingale (see Protter (1992)).

Example 8.12: The function $(x-a)^+$ is important in financial application as it gives the payoff of a financial stock option. The Meyer–Tanaka’s formula for $(x-a)^+$

$$(X(t) - a)^+ = (X(0) - a)^+ + \int_0^t I(X(s) > a)dX(s) + \frac{1}{2}L_t^a. \quad (8.34)$$

Theorem 8.11: Let $L^a(t)$ be the local time of Brownian motion at a , and $f_t(a)$ the density of $N(0, t)$ at a . Then

$$E(L^a(t)) = \int_0^t f_s(a)ds \quad \text{hence} \quad \frac{dE(L^a(t))}{dt} = f_t(a). \quad (8.35)$$

PROOF. Taking expectation in both sides of Equation (8.32) and changing the order of integration, we obtain for any positive and bounded g

$$\int_{-\infty}^{\infty} g(a) \int_0^t f_s(a)ds da = \int_{-\infty}^{\infty} g(a) E(L^a(t)) da.$$

The result follows since g is arbitrary. \square

A similar result can be established for continuous semimartingales by using Equation (8.29) (e.g. Klebaner (2002)).

Remark 8.4: Local times can also be defined for discontinuous semimartingales. For any fixed a , $L^a(t)$ is a continuous non-decreasing function in t , and it increases only at points of continuity of X where it is equal to a , that is, $X(t-) = X(t) = a$. The formula (8.29) holds with quadratic variation $[X, X]$ replaced by its continuous part $[X, X]^c$ (see, for example, Protter (1992)).

8.8 Stochastic Exponential

The stochastic exponential (also known as the semimartingale, or Doléans–Dade exponential) is a stochastic analogue of the exponential function. Recall that if $f(t)$ is a smooth function, then $g(t) = e^{f(t)}$ is the solution to the differential equation $dg(t) = g(t)df(t)$. The stochastic exponential is defined as a solution to a similar stochastic equation. The stochastic exponential of Itô processes was introduced in Chapter 5.2. For a semimartingale X , its stochastic exponential $\mathcal{E}(X)(t) = U(t)$ is defined as the unique solution to the equation

$$U(t) = 1 + \int_0^t U(s-)dX(s) \quad \text{or} \quad dU(t) = U(t-)dX(t); \quad \text{with } U(0) = 1. \quad (8.36)$$

As an application of Itô’s formula and the rules of stochastic calculus we prove

Theorem 8.12: *Let X be a continuous semimartingale. Then its stochastic exponential is given by*

$$U(t) = \mathcal{E}(X)(t) = e^{X(t) - X(0) - \frac{1}{2}[X, X](t)}. \quad (8.37)$$

PROOF. Write $U(t) = e^{V(t)}$, with $V(t) = X(t) - X(0) - \frac{1}{2}[X, X](t)$. Then

$$dU(t) = d(e^{V(t)}) = e^{V(t)}dV(t) + \frac{1}{2}e^{V(t)}d[V, V](t).$$

Using the fact that $[X, X](t)$ is a continuous process of finite variation, we obtain $[X, [X, X]](t) = 0$, and $[V, V](t) = [X, X](t)$. Using this, we obtain

$$dU(t) = e^{V(t)}dX(t) - \frac{1}{2}e^{V(t)}d[X, X](t) + \frac{1}{2}e^{V(t)}d[X, X](t) = e^{V(t)}dX(t),$$

or $dU(t) = U(t)dX(t)$. Thus $U(t)$ defined by (8.37) satisfies (8.36). In order to show uniqueness, let $U_1(t)$ be another solution to (8.36), and consider $U_1(t)/U(t)$. Integration by parts

$$d\left(\frac{U_1(t)}{U(t)}\right) = U_1(t)d\left(\frac{1}{U(t)}\right) + \frac{1}{U(t)}dU_1(t) + d\left[U_1, \frac{1}{U}\right](t).$$

Itô’s formula, using that $U(t)$ is continuous and satisfies (8.36)

$$d\left(\frac{1}{U(t)}\right) = -\frac{1}{U(t)}dX(t) + \frac{1}{U(t)}d[X, X](t),$$

which leads to

$$\begin{aligned} d\left(\frac{U_1(t)}{U(t)}\right) &= -\frac{U_1(t)}{U(t)}dX(t) + \frac{U_1(t)}{U(t)}dX(t) \\ &\quad + \frac{U_1(t)}{U(t)}d[X, X](t) - \frac{U_1(t)}{U(t)}d[X, X](t) = 0. \end{aligned}$$

Thus $U_1(t)/U(t) = \text{const.} = U_1(0)/U(0) = 1$. □

Properties of the stochastic exponential are given by the following result.

Theorem 8.13: *Let X and Y be semimartingales on the same space. Then*

1. $\mathcal{E}(X)(t)\mathcal{E}(Y)(t) = \mathcal{E}(X + Y + [X, Y])(t)$.
2. If X is continuous, $X(0) = 0$, then $(\mathcal{E}(X)(t))^{-1} = \mathcal{E}(-X + [X, X])(t)$.

The proof uses the integration by parts formula and is left as an exercise.

Example 8.13: (Stock process and its return process)

An application in finance is provided by the relation between the stock process and its return process. The return is defined by $dR(t) = dS(t)/S(t-)$. Hence the stock price is the stochastic exponential of the return, $dS(t) = S(t-)dR(t)$, and $S(t) = S(0)\mathcal{E}(R)(t)$.

Stochastic Exponential of Martingales

Stochastic exponential $U = \mathcal{E}(M)$ of a martingale, or a local martingale, $M(t)$ is a stochastic integral with respect to $M(t)$. Since stochastic integrals with respect to martingales or local martingales are local martingales, $\mathcal{E}(M)$ is a local martingale. In applications it is important to have conditions for $\mathcal{E}(M)$ to be a true martingale.

Theorem 8.14 (Martingale Exponential): *Let $M(t), 0 \leq t \leq T < \infty$ be a continuous local martingale null at zero. Then its stochastic exponential $\mathcal{E}(M)$ is given by $e^{M(t)-\frac{1}{2}[M,M](t)}$ and it is a continuous positive local martingale. Consequently, it is a supermartingale, it is integrable, and has a finite non-increasing expectation. It is a martingale if any of the following conditions hold.*

1. $E(e^{M(T)-\frac{1}{2}[M,M](T)}) = 1$.
2. For all $t \geq 0$, $E(\int_0^t e^{2M(s)-[M,M](s)} d[M, M](s)) < \infty$.
3. For all $t \geq 0$, $E(\int_0^t e^{2M(s)} d[M, M](s)) < \infty$.

Moreover, if the expectations above are bounded by $K < \infty$, then $\mathcal{E}(M)$ is a square integrable martingale.

PROOF. According to Theorem 8.12, $\mathcal{E}(M)(t) = e^{M(t)-\frac{1}{2}[M,M](t)}$, therefore it is positive. Being a stochastic integral with respect to a martingale, it is a local martingale. Thus $\mathcal{E}(M)$ is a supermartingale, as a positive local martingale, see Theorem 7.23. A supermartingale has a non-increasing expectation, and is a martingale if, and only if, its expectation at T is the same as at 0 (see Theorem 7.3). This gives the first condition. The second condition follows from Theorem 7.35, which states that if a local martingale

has finite expectation of its quadratic variation, then it is a martingale. So that if $E[\mathcal{E}(M), \mathcal{E}(M)](t) < \infty$, then $\mathcal{E}(M)$ is a martingale. According to the quadratic variation of an integral

$$[\mathcal{E}(M), \mathcal{E}(M)](t) = \int_0^t e^{2M(s)-[M, M](s)} d[M, M](s). \quad (8.38)$$

The third condition follows from the second since $[M, M]$ is positive and increasing. The last statement follows by Theorem 7.35 since the bound implies $\sup_{0 \leq t} E[\mathcal{E}(M), \mathcal{E}(M)](t) < \infty$. \square

Theorem 8.15 (Kazamaki's Condition): *Let M be a continuous local martingale with $M(0) = 0$. If $e^{\frac{1}{2}M(t)}$ is a submartingale, then $\mathcal{E}(M)$ is a martingale.*

The result is proven in Revuz and Yor (2001).

Theorem 8.16: *Let M be a continuous martingale with $M(0) = 0$. If*

$$E(e^{\frac{1}{2}M(T)}) < \infty, \quad (8.39)$$

then $\mathcal{E}(M)$ is a martingale on $[0, T]$.

PROOF. According to Jensen's inequality (see Exercise 7.3) if g is a convex function, and $E|g(M(t))| < \infty$ for $t \leq T$, then $Eg(M(t)) \leq Eg(M(T))$ and $g(M(t))$ is a submartingale. Since $e^{x/2}$ is convex the result follows by Kazamaki's condition. \square

Theorem 8.17 (Novikov's Condition): *Let M be a continuous local martingale with $M(0) = 0$. Suppose that for each $t \leq T$*

$$E(e^{\frac{1}{2}[M, M](t)}) < \infty. \quad (8.40)$$

Then $\mathcal{E}(M)$ is a martingale with mean one. In particular, if for each t there is a constant K_t such that $[M, M](t) < K_t$, then $\mathcal{E}(M)(t)$, $t \leq T$ is a martingale.

PROOF. The condition (8.40) implies that $[M, M](t)$ has moments. According to the Burkholder–Davis–Gundy (BDG) inequality (7.41), $\sup_{t \leq T} M(t)$ is integrable, therefore $M(t)$ is a martingale. Next, using the formula for $\mathcal{E}(M)$ and the Cauchy–Schwarz inequality, $E\sqrt{XY} \leq \sqrt{EXEY}$,

$$E(e^{\frac{1}{2}M(T)}) = E\sqrt{\mathcal{E}(M)(T)e^{\frac{1}{2}[M, M](T)}} \leq \sqrt{E\mathcal{E}(M)(T)E(e^{\frac{1}{2}[M, M](T)})} < \infty. \quad (8.41)$$

The last expression is finite since $E\mathcal{E}(M)(T) < \infty$, as $\mathcal{E}(M)$ is a supermartingale, and by the condition of the theorem. This shows that condition (8.39) holds and the result follows. \square

Another proof can be found in Karatzas and Shreve (1988).

Now we can complete the missing step in the proof of Levy's theorem on the characterization of Brownian motion (Theorem 7.36).

Corollary 8.18: *If $M(t)$ is a continuous local martingale with $[M, M](t) = t$, then $U(t) = \mathcal{E}(uM)(t) = e^{uM(t)-u^2t/2}$ is a martingale.*

PROOF. Clearly, $uM(t)$ is a continuous local martingale with quadratic variation u^2t . The result now follows by Novikov's condition.

However, it is possible to give a direct proof. For ease of notation take $u = 1$. Since U is a stochastic exponential it satisfies the SDE

$$U(t) = 1 + \int_0^t U(s)dM(s).$$

Hence $U(t) - 1 = \int_0^t U(s)dM(s)$ is a local martingale with quadratic variation

$$\left[\int_0^t U(s)dM(s), \int_0^t U(s)dM(s) \right] = \int_0^t U^2(s)d[M, M](s) = \int_0^t U^2(s)ds. \quad (8.42)$$

According to the BDG inequality with $p = 2$

$$E\left(\sup_{t' \leq t} \left| \int_0^{t'} U(s)dM(s) \right| \right)^2 \leq CE \left(\int_0^t U^2(s)ds \right).$$

Replacing supremum by the last point makes it smaller, and we have $(\sup_{t' \leq t} |\int_0^{t'} U(s)dM(s)|)^2 \geq (\int_0^t U(s)dM(s))^2 = (U(t) - 1)^2$. Thus from above we have

$$E(U(t) - 1)^2 \leq C \int_0^t EU(s)^2 ds.$$

By an elementary inequality $(a+1)^2 \leq 2(a^2+1)$ we obtain with $U(s) - 1 = a$ that $f(t) = E(U(t) - 1)^2$ satisfies the following inequality:

$$f(t) \leq 2Ct + 2C \int_0^t f(s)ds.$$

Gronwall's inequality (Theorem 1.20) implies that $f(t) \leq 2Cte^{2Ct} < \infty$. This implies that $\int_0^T E(U^2(s))dt < \infty$. A sufficient condition for a local martingale to be a martingale is finiteness of the expectation of its quadratic variation, Theorem 7.35, which by (8.42) is now established. \square

The next section gives more information and tools for processes with jumps.

8.9 Compensators and Sharp Bracket Process

A process N is called increasing if all of its realizations $N(t)$ are non-decreasing functions of t . A process N is of finite variation if all of its realizations $N(t)$ are functions of finite variation, $V_N(t) < \infty$ for all t , where $V_N(t)$ is the variation of N on $[0, t]$.

Definition 8.19: An increasing process N , $t \geq 0$, is called integrable if $\sup_{t \geq 0} E N(t) < \infty$.

A finite variation process N is of integrable variation if its variation process is integrable, $\sup_{t \geq 0} E V_N(t) < \infty$.

A finite variation process N is of locally integrable variation if there is a sequence of stopping times τ_n such that $\tau_n \uparrow \infty$ so that $N(t \wedge \tau_n)$ is of integrable variation, that is, $\sup_{t \geq 0} E V_N(t \wedge \tau_n) < \infty$.

Example 8.14: A Poisson process $N(t)$ with parameter λ is of finite, but not integrable, variation since for any t , $V_N(t) = N(t) < \infty$, but $\sup_{t \geq 0} E V_N(t) = \infty$. It is of locally integrable variation, since $\sup_{t \geq 0} E V_N(t \wedge n) = \lambda n < \infty$. Here $\tau_n = n$.

Example 8.15: It can be seen that a finite variation process $N(t)$ with bounded jumps $|\Delta N(t)| \leq c$ is of locally integrable variation. If $\tau_n = \inf\{t : V_N(t) \geq n\}$, then $N(t \wedge \tau_n)$ has variation bounded by $n + c$. τ_n are stopping times, as first times of boundary crossing.

Definition 8.20: Let $N(t)$ be an adapted process of integrable or locally integrable variation. Its compensator $A(t)$ is the unique predictable process such that $M(t) = N(t) - A(t)$ is a local martingale.

Existence of compensators is assured by the Doob–Meyer decomposition.

Theorem 8.21: Let $N(t)$ be an adapted process of integrable or locally integrable variation. Then its compensator exists. Moreover, it is locally integrable.

PROOF. As a finite variation process is a difference of two increasing processes, it is enough to establish the result for increasing processes. According to localization it is possible to assume that it is integrable. However, an increasing integrable process is a submartingale, and the result follows by the Doob–Meyer decomposition Theorem 8.4. \square

Remark 8.5: The condition $M = N - A$ is a local martingale is equivalent to the condition (see Liptser and Shiryaev (1989))

$$E \int_0^\infty H(s) dN(s) = E \int_0^\infty H(s) dA(s), \quad (8.43)$$

for any positive predictable process H . Sometimes this integral condition (8.43) is taken as the definition of the compensator, e.g. Karr (1986).

The compensator of N is also called the dual predictable projection of N (Rogers and Williams (1990), and Liptser and Shiryaev (1989)).

Note that the compensator is unique with respect to the given filtration and probability. If the filtration or probability are changed, then the compensator will also change.

Recall that the quadratic variation process $[X, X](t)$ of a semimartingale X exists and is non-decreasing. Consider now semimartingales with integrable $(\sup_{t \geq 0} E[X, X](t) < \infty)$ or locally integrable quadratic variation.

Definition 8.22: The sharp bracket (or angle bracket, or predictable quadratic variation) $\langle X, X \rangle(t)$ process of a semimartingale X is the compensator of $[X, X](t)$. That is, it is the unique predictable process that makes $[X, X](t) - \langle X, X \rangle(t)$ into a local martingale.

Example 8.16: Let N be a Poisson process. It is of finite variation and changes only by jumps (pure jump process), which are of size 1, $\Delta N(t) = 0$ or 1, and $(\Delta N(t))^2 = \Delta N(t)$. Its quadratic variation is the process $N(t)$ itself,

$$[N, N](t) = \sum_{0 \leq s \leq t} (\Delta N(s))^2 = \sum_{0 \leq s \leq t} \Delta N(s) = N(t).$$

Clearly, $\sup_{0 \leq t \leq T} E[N, N](t) = T$. Thus N is of integrable variation on $[0, T]$. t is non-random, therefore predictable. Since $[N, N](t) - t = N(t) - t$ is a martingale,

$$\langle N, N \rangle(t) = t.$$

Example 8.17: Let B be a Brownian motion. Its quadratic variation is $[B, B](t) = t$, and since it is non-random it is predictable. Hence $\langle B, B \rangle(t) = t$, and the martingale part in the Doob–Meyer decomposition of $[B, B](t)$ is trivial, $M(t) \equiv 0$.

Example 8.17 generalizes to any continuous semimartingale.

Theorem 8.23: *If $X(t)$ is a continuous semimartingale with integrable quadratic variation, then $\langle X, X \rangle(t) = [X, X](t)$, and there is no difference between the sharp and the square bracket processes.*

PROOF. The quadratic variation jumps at the points of jumps of X and $\Delta[X, X](s) = (\Delta X(s))^2$. Since X has no jumps, $[X, X](t)$ is continuous. $[X, X](t)$ is predictable as a continuous and adapted process, the martingale part in the Doob–Meyer decomposition of $[X, X](t)$ is trivial, $M(t) \equiv 0$, and $\langle X, X \rangle(t) = [X, X](t)$. \square

Example 8.18: Let X be a diffusion solving the SDE $dX(t) = \mu(X(t))dt + \sigma(X(t))dB(t)$. Then $[X, X](t) = \int_0^t \sigma^2(X(s))ds = \langle X, X \rangle(t)$.

Sharp Bracket for Square Integrable Martingales

Let M be a square integrable martingale, that is, $\sup_t E M^2(t) < \infty$. Recall that the quadratic variation of M has the property

$$M^2(t) - [M, M](t) \text{ is a martingale.} \quad (8.44)$$

M^2 is a submartingale since x^2 is a convex function. Using the Doob–Meyer decomposition for submartingales, Theorem 8.4, we can prove the following:

Theorem 8.24: *Let M be a square integrable martingale. Then the sharp bracket process $\langle M, M \rangle(t)$ is the unique predictable increasing process for which*

$$M^2(t) - \langle M, M \rangle(t) \text{ is a martingale.} \quad (8.45)$$

PROOF. According to the definition of the sharp bracket process, $[M, M](t) - \langle M, M \rangle(t)$ is a martingale. As a sum of two martingales, $M^2(t) - \langle M, M \rangle(t)$ is also a martingale. Since $\langle M, M \rangle(t)$ is predictable and increasing, and $M^2(t)$ is a submartingale, uniqueness follows by the Doob–Meyer decomposition. \square

By taking expectations in (8.45) we obtain a useful corollary.

Corollary 8.25: *Let $M(t)$ be a square integrable martingale with $M(0) = 0$. Then*

$$E(M^2(t)) = E[M, M](t) = E\langle M, M \rangle(t). \quad (8.46)$$

This result allows us to use Doob's martingale inequality with the sharp bracket

$$E\left(\left(\sup_{s \leq T} M(s)\right)^2\right) \leq 4E(M^2(T)) = 4E(\langle M, M \rangle(T)). \quad (8.47)$$

By using localization in the proof of Theorem 8.24 one can show

Theorem 8.26: *Let M be a locally square integrable martingale, then the predictable quadratic variation $\langle M, M \rangle(t)$ is the unique predictable process for which $M^2(t) - \langle M, M \rangle(t)$ is a local martingale.*

The next result allows us to decide when a local martingale is a martingale by using the predictable quadratic variation.

Theorem 8.27: *Let $M(t)$, $0 \leq t \leq T < \infty$ be a local martingale, null at zero, such that for all t , $E\langle M, M \rangle(t) < \infty$. Then M is a square integrable martingale, moreover, $EM^2(t) = E[M, M](t) = E\langle M, M \rangle(t)$. If $T = \infty$, and $\sup_{t < \infty} E\langle M, M \rangle(t) < \infty$, then $M(t)$ is a square integrable martingale on $[0, \infty)$.*

PROOF. $[M, M](t) - \langle M, M \rangle(t)$ is a local martingale. Let τ_n be a localizing sequence. Then $E[M, M](t \wedge \tau_n) = E\langle M, M \rangle(t \wedge \tau_n)$. Since both sides are non-decreasing they converge to the same limit as $n \rightarrow \infty$. However, $\lim_{n \rightarrow \infty} E\langle M, M \rangle(t \wedge \tau_n) = E\langle M, M \rangle(t) < \infty$. Therefore $E[M, M](t) = E\langle M, M \rangle(t) < \infty$. Thus the conditions of Theorem 7.35 are satisfied and the result follows. \square

Since a continuous local martingale is locally square integrable we obtain

Corollary 8.28: *The sharp bracket process (predictable quadratic variation) for a continuous local martingale exists.*

Continuous Martingale Component of a Semimartingale

A function of finite variation has a decomposition into continuous and discrete parts. A semimartingale is a sum of a process of finite variation and a local martingale. It turns out that one can decompose any local martingale into a continuous local martingale and a purely discontinuous one. Such decomposition requires a different approach to the case of finite variation processes.

Definition 8.29: A local martingale is purely discontinuous if it is orthogonal to any continuous local martingale. Local martingales M and N are orthogonal if MN is a local martingale.

Example 8.19: A compensated Poisson process $\bar{N}(t) = N(t) - t$ is a purely discontinuous martingale. Let $M(t)$ be any continuous local martingale. Then by the integration by parts formula (8.23)

$$M(t)\bar{N}(t) = \int_0^t M(s-)d\bar{N}(s) + \int_0^t \bar{N}(s-)dM(s) + [M, \bar{N}](t).$$

Since \bar{N} is of finite variation, by the property (8.18) of quadratic covariation $[M, \bar{N}](t) = \sum_{s \leq t} \Delta M(s)\Delta \bar{N}(s)$. However, M is continuous, $\Delta M(s) = 0$, and $[M, \bar{N}](t) = 0$. Therefore $M(t)\bar{N}(t)$ is a sum of two stochastic integrals with respect to local martingales, and itself is a local martingale.

It is possible to show that any local martingale M has a unique decomposition

$$M = M^c + M^d,$$

where M^c is a continuous and M^d a purely discontinuous local martingale (see for example Liptser and Shiryaev (1989), Protter (1992), and Jacod and Shiryaev (1987)).

If X is a semimartingale with representation

$$X(t) = X(0) + M(t) + A(t), \quad (8.48)$$

with a local martingale M , then M^c is called the continuous martingale component of X and is denoted by X^{cm} . Even if the above representation of X is not unique, the continuous martingale component of X is the same for all representations. Indeed, if $X(t) = X(0) + M_1(t) + A_1(t)$ is another representation, then $(M - M_1)(t) = -(A - A_1)(t)$. Hence $(M - M_1)$ is a martingale of finite variation. Hence its continuous component is also a martingale of finite variation. However, a continuous martingale of finite variation is a constant. This implies that $M^c - M_1^c = 0$. Thus $X^{cm} = M^c$ is the same for all representations. If X is of finite variation, then the martingale part is zero and by the uniqueness of X^{cm} we have

Corollary 8.30: *If X is a semimartingale of finite variation, then $X^{cm} \equiv 0$.*

For example, the compensated Poisson process $N(t) - t$ has zero continuous martingale component.

It can be shown that

$$\langle X^{cm}, X^{cm} \rangle = [X, X]^c, \quad (8.49)$$

where $[X, X]^c$ is the continuous part of the finite variation process $[X, X]$. Of course, because X^{cm} is continuous $\langle X^{cm}, X^{cm} \rangle = [X^{cm}, X^{cm}]$.

Let $\Delta X(s) = X(s) - X(s-)$ and put $X(0-) = 0$ and $[X, X]^c(0) = 0$. Since the jumps of quadratic variation satisfy (see (8.17))

$$\begin{aligned} \Delta[X, X](s) &= (\Delta X(s))^2, \\ [X, X](t) &= [X, X]^c(t) + \sum_{0 < s \leq t} \Delta[X, X](s) = [X, X]^c(t) + \sum_{0 < s \leq t} (\Delta X(s))^2 \\ &= \langle X^{cm}, X^{cm} \rangle + \sum_{0 < s \leq t} (\Delta X(s))^2. \end{aligned} \quad (8.50)$$

Since the quadratic variation $[X, X]$ for a semimartingale exists, and the predictable quadratic variation $\langle X^{cm}, X^{cm} \rangle$ exists (Corollary 8.28) we obtain

Corollary 8.31: *If X is a semimartingale, then for each t*

$$\sum_{s \leq t} (\Delta X(s))^2 < \infty. \quad (8.51)$$

Conditions for Existence of a Stochastic Integral

The class of processes H for which the stochastic integral with respect to a martingale M can be defined depends in an essential way on the properties of the predictable quadratic variation $\langle M, M \rangle$ of M . Consider integrals with

respect to a locally square integrable martingale M , possibly discontinuous. The stochastic integral $\int_0^T H(s)dM(s)$ can be defined for predictable processes H such that

$$\int_0^T H^2(t)d\langle M, M \rangle(t) < \infty, \quad (8.52)$$

and in this case the integral $\int_0^t H(s)dM(s)$, $0 \leq t \leq T$ is a local martingale.

The class of processes H that can be integrated against M is wider when $\langle M, M \rangle(t)$ is continuous, and wider still when $\langle M, M \rangle(t)$ is absolutely continuous (can be represented as an integral with respect to dt). These classes and conditions are given, for example, in Liptser and Shiryaev (2001).

Example 8.20: Let filtration \mathbb{F} be generated by a Brownian motion $B(t)$ and a Poisson process $N(t)$. The process $N(t-)$ is a left-continuous modification of $N(t)$. By definition, $N(t-) = \lim_{s \uparrow t} N(s)$. Being left-continuous, it is predictable. Condition (8.52) holds. The integral $\int_0^t N(s-)dB(s)$ is a well-defined stochastic integral of a predictable process with respect to a martingale B .

Properties of the Predictable Quadratic Variation

The predictable quadratic variation (the sharp bracket process) has similar properties to the quadratic variation (the square bracket) process. We list them without proof. All the processes below are assumed to be semimartingales with locally integrable quadratic variation.

1. $\langle X, X \rangle(t)$ is increasing in t .
2. $\langle X, Y \rangle$ is bilinear and symmetric,

$$\langle \alpha X + Y, \beta U + V \rangle = \alpha \beta \langle X, U \rangle + \alpha \langle X, V \rangle + \beta \langle Y, U \rangle + \langle Y, V \rangle. \quad (8.53)$$

3. Polarization identity. $\langle X, Y \rangle = \frac{1}{4}(\langle X + Y, X + Y \rangle - \langle X - Y, X - Y \rangle).$
4. $\langle X, Y \rangle$ is a predictable process of finite variation.
5. $\langle X, Y \rangle = 0$ if X or Y is of finite variation and one of them is continuous.
6. The sharp bracket process of stochastic integrals $\langle H \cdot X, K \cdot Y \rangle(t)$.

$$\left\langle \int_0^\cdot H(s)dX(s), \int_0^\cdot K(s)dY(s) \right\rangle(t) = \int_0^t H(s)K(s)d\langle X, Y \rangle(s), \quad (8.54)$$

in particular $\left\langle \int_0^\cdot H(s)dX(s), \int_0^\cdot H(s)dX(s) \right\rangle(t) = \int_0^t H^2(s)d\langle X, X \rangle(s)$, $\left\langle \int_0^\cdot H(s)dX(s), Y \right\rangle(t) = \left\langle \int_0^\cdot H(s)dX, \int_0^\cdot dY \right\rangle(t) = \int_0^t H(s)d\langle X, Y \rangle(s)$.

Recall that stochastic integrals with respect to local martingales are again local martingales. Using the sharp bracket of the integral together with Theorem 8.27 we obtain

Theorem 8.32: *Let $M(t)$, $0 \leq t \leq T$ be a local martingale and $H(t)$ be a predictable process such that $E(\int_0^T H^2(s)d\langle M, M\rangle(s)) < \infty$. Then $\int_0^t H(s)dM(s)$ is a square integrable martingale, moreover,*

$$\left\langle \int_0^\cdot H(s)dM(s), \int_0^\cdot H(s)dM(s) \right\rangle(t) = \int_0^t H^2(s)d\langle M, M\rangle(s). \quad (8.55)$$

Using this result we obtain the isometry property for stochastic integrals in terms of the sharp bracket process.

$$E\left(\int_0^t H(s)dM(s)\right)^2 = E\left(\int_0^t H^2(s)d\langle M, M\rangle(s)\right). \quad (8.56)$$

Example 8.21: Let $M(t)$ be the compensated Poisson process, $M(t) = N(t) - t$, and H be predictable, satisfying $E\int_0^T H^2(t)dt < \infty$. Then $\int_0^t H(s)dM(s)$ is a martingale, moreover,

$$E\int_0^t H(s)dM(s) = 0, \text{ and } E\left(\int_0^t H(s)dM(s)\right)^2 = E\int_0^t H^2(s)ds.$$

8.10 Itô's Formula for Semimartingales

Let $X(t)$ be a semimartingale and f be a C^2 function. Then $f(X(t))$ is a semimartingale, and Itô's formula holds

$$\begin{aligned} f(X(t)) - f(X(0)) &= \int_0^t f'(X(s-))dX(s) + \frac{1}{2} \int_0^t f''(X(s-))d[X, X](s) \\ &\quad + \sum_{s \leq t} \left(f(X(s)) - f(X(s-)) - f'(X(s-))\Delta X(s) \right. \\ &\quad \left. - \frac{1}{2}f''(X(s-))(\Delta X(s))^2 \right). \end{aligned} \quad (8.57)$$

The quadratic variation $[X, X]$ jumps at the points of jumps of X and its jumps $\Delta[X, X](s) = (\Delta X(s))^2$. Thus the jump part of the integral $\int_0^t f''(X(s-))d[X, X](s)$ is given by $\sum_{s \leq t} f''(X(s-))(\Delta X(s))^2$, leading to an equivalent form of the formula

$$\begin{aligned} f(X(t)) - f(X(0)) &= \int_0^t f'(X(s-))dX(s) + \frac{1}{2} \int_0^t f''(X(s-))d[X, X]^c(s) \\ &\quad + \sum_{s \leq t} (f(X(s)) - f(X(s-)) - f'(X(s-))\Delta X(s)), \end{aligned} \quad (8.58)$$

where $[X, X]^c$ is the continuous component of the finite variation function $[X, X]$. Using the relationship between the square and the sharp brackets (8.50) we can write Itô's formula with the sharp bracket process of X , provided the sharp bracket exists,

$$\begin{aligned} f(X(t)) - f(X(0)) &= \int_0^t f'(X(s-))dX(s) \\ &\quad + \frac{1}{2} \int_0^t f''(X(s-))d\langle X^{cm}, X^{cm} \rangle(s) \\ &\quad + \sum_{s \leq t} (f(X(s)) - f(X(s-)) - f'(X(s-))\Delta X(s)), \end{aligned} \tag{8.59}$$

where X^{cm} denotes the continuous martingale part of X .

Example 8.22: Let $N(t)$ be a Poisson process. We calculate $\int_0^t N(s-)dN(s)$. The answer can be derived from the integration by parts formula (8.23), but now we use (8.59). Since $(N(t) - t)^{cm} = 0$ (by Corollary 8.30)

$$N^2(t) = 2 \int_0^t N(s-)dN(s) + \sum_{s \leq t} (N^2(s) - N^2(s-) - 2N(s-) \Delta N(s)).$$

Since $N(s) = N(s-) + \Delta N(s)$, $(N(s-) + \Delta N(s))^2 - N^2(s-) - 2N(s-) \Delta N(s) = (\Delta N(s))^2 = \Delta N(s)$, and the sum simplifies to $\sum_{s \leq t} \Delta N(s) = N(t)$. Thus we obtain

$$\int_0^t N(s-)dN(s) = \frac{1}{2}(N^2(t) - N(t)).$$

A formula (8.59) for a function of n variables reads: $\mathbf{X}(t) = (X^1(t), \dots, X^n(t))$ is a semimartingale and f is a C^2 function of n variables,

$$\begin{aligned} f(\mathbf{X}(t)) - f(\mathbf{X}(0)) &= \sum_{i=1}^n \int_0^t \frac{\partial f}{\partial x_i}(\mathbf{X}(s-))dX^i(s) \\ &\quad + \frac{1}{2} \sum_{i,j=1}^n \int_0^t \frac{\partial^2 f}{\partial x_i \partial x_j}(\mathbf{X}(s-))d\langle X^{i,cm}, X^{j,cm} \rangle(s) \\ &\quad + \sum_{s \leq t} \left(f(\mathbf{X}(s)) - f(\mathbf{X}(s-)) - \sum_{i=1}^n \frac{\partial}{\partial x_i} f(\mathbf{X}(s-)) \Delta X^i(s) \right). \end{aligned} \tag{8.60}$$

Itô's formula can be found in many texts, see, for example, Protter (1992), Rogers and Williams (1990), Liptser and Shiryaev (1989), Métivier (1982), and Dellacherie and Meyer (1982).

8.11 Stochastic Exponential and Logarithm

As an application of Itô's formula and the rules of stochastic calculus we outline a proof of the following result.

Theorem 8.33: *Let X be a semimartingale. Then the stochastic equation*

$$U(t) = 1 + \int_0^t U(s-)dX(s) \quad (8.61)$$

has a unique solution, called the stochastic exponential of X , and this solution is given by

$$U(t) = \mathcal{E}(X)(t) = e^{X(t)-X(0)-\frac{1}{2}\langle X, X \rangle^c(t)} \prod_{s \leq t} (1 + \Delta X(s)) e^{-\Delta X(s)}. \quad (8.62)$$

Formula (8.62) can be written by using quadratic variation as follows:

$$\mathcal{E}(X)(t) = e^{X(t)-X(0)-\frac{1}{2}[X, X](t)} \prod_{s \leq t} (1 + \Delta X(s)) e^{(-\Delta X(s))+\frac{1}{2}(\Delta X(s))^2}. \quad (8.63)$$

PROOF. Let $Y(t) = X(t) - X(0) - \frac{1}{2}\langle X, X \rangle^c(t)$ and $V(t) = \prod_{s \leq t} (1 + \Delta X(s)) e^{-\Delta X(s)}$. Note that although the product is taken for all $s \leq t$, there are at most countably many points at which $\Delta X(s) \neq 0$ (by the regularity property of the process), hence there are at most countably many elements different from 1 in the product. We show that the product converges. Since by (8.51) $\sum_{s \leq t} (\Delta X(s))^2 < \infty$, there are only finitely many points s at which $|\Delta X(s)| > 0.5$, which give a finite non-zero contribution to the product. Taking the product with over s at which $|\Delta X(s)| \leq 1/2$, and taking logarithm, it is enough to show that $\sum_{s \leq t} |\ln(1 + \Delta X(s)) - \Delta X(s)|$ converges. However, this follows from the inequality $|\ln(1 + \Delta X(s)) - \Delta X(s)| \leq (\Delta X(s))^2$ by (8.51). In order to see that $U(t)$ defined by (8.62) satisfies (8.61), use Itô's formula applied to the function $f(Y(t), V(t))$ with $f(x_1, x_2) = e^{x_1} x_2$. For the uniqueness of the solution of (8.61) and other details see Liptser and Shiryaev (1989). \square

Example 8.23: The stochastic exponential (8.62) of a Poisson process is easily seen to be $\mathcal{E}(N)(t) = 2^{N(t)}$.

If $U = \mathcal{E}(X)$ is the stochastic exponential of X , then $X = \mathcal{L}(U)$ is the stochastic logarithm of U , satisfying Equation (8.61)

$$dX(t) = \frac{dU(t)}{U(t-)}, \text{ or } \mathcal{L}(\mathcal{E}(X)) = X.$$

For Itô processes an expression for $X(t)$ is given in Theorem 5.2, for the general case see Exercise 8.17.

8.12 Martingale (Predictable) Representations

In this section we give results on the representation of martingales by stochastic integrals of predictable processes, also called *predictable* representations. Let $M(t)$ be a martingale, $0 \leq t \leq T$, adapted to the filtration $\mathbb{F} = (\mathcal{F}_t)$, and $H(t)$ be a predictable process satisfying $\int_0^T H^2(s)d\langle M, M\rangle(s) < \infty$ with probability one. Then $\int_0^t H(s)dM(s)$ is a local martingale. The predictable representation property means that the converse is also true. Let $\mathbb{F}^M = (\mathcal{F}_t^M)$ denote the natural filtration of M .

Definition 8.34: A local martingale M has the predictable representation property if for any \mathbb{F}^M -local martingale X there is a predictable process H such that

$$X(t) = X(0) + \int_0^t H(s)dM(s). \quad (8.64)$$

This definition is different to the classical one for martingales with jumps, see Remark 8.7 below, but is the same for continuous martingales.

Brownian motion has the predictable representation property (see, for example, Revuz and Yor (2001) p. 209, Liptser and Shiryaev (2001) I p. 170).

Theorem 8.35 (Brownian Martingale Representation):

Let $X(t)$, $0 \leq t \leq T$ be a local martingale adapted to the Brownian filtration $\mathbb{F}^B = (\mathcal{F}_t)$. Then there exists a predictable process $H(t)$ such that $\int_0^T H^2(s)ds < \infty$ with probability one, and Equation (8.65) holds:

$$X(t) = X(0) + \int_0^t H(s)dB(s). \quad (8.65)$$

Moreover, if Y is an integrable \mathcal{F}_T -measurable random variable, $E|Y| < \infty$, then

$$Y = EY + \int_0^T H(t)dB(t). \quad (8.66)$$

If, in addition, Y and B have jointly a Gaussian distribution, then the process $H(t)$ in (8.66) is deterministic.

PROOF. We do not prove the representation of a martingale, but only the representation for a random variable based on it.

Take $X(t) = E(Y|\mathcal{F}_t)$. Then $X(t)$, $0 \leq t \leq T$ is a martingale (see Theorem 7.9). Hence by the martingale representation there exists H such that $X(t) = X(0) + \int_0^t H(s)dB(s)$. Taking $t = T$ gives the result. \square

Remark 8.6: A functional of the path of the Brownian motion $B_{[0,T]}$ is a random variable Y , \mathcal{F}_T -measurable. Theorem 8.35 states that under the above assumptions any functional of Brownian motion has the form (8.66).

Since Itô integrals are continuous, and any local martingale of a Brownian filtration is an Itô integral, it follows that all local martingales of a Brownian filtration are continuous. In fact we have the following result (the second statement is not straightforward)

Corollary 8.36:

1. All local martingales of the Brownian filtration are continuous.
2. All right-continuous adapted processes are predictable.

Corollary 8.37:

Let $X(t)$, $0 \leq t \leq T$, be a square integrable martingale adapted to the Brownian filtration \mathbb{F} . Then there exists a predictable process $H(t)$ such that $E \int_0^T H^2(s)ds < \infty$ and representation (8.65) holds. Moreover,

$$\langle X, B \rangle(t) = \int_0^t H(s)ds \quad \text{and} \quad H(t) = \frac{d\langle X, B \rangle(t)}{dt}. \quad (8.67)$$

Equation (8.67) follows from (8.65) by the rule of the sharp bracket for integrals.

Example 8.24: (Representation of martingales)

1. $X(t) = B^2(t) - t$. Then $X(t) = \int_0^t 2B(s)dB(s)$. Here $H(t) = 2B(t)$, which can also be found by using (8.67).
2. Let $X(t) = f(B(t), t)$ be a martingale. According to Itô's formula $dX(t) = \frac{\partial f}{\partial x}(B(t), t)dB(t)$. Thus $H(t) = \frac{\partial f}{\partial x}(B(t), t)$. This also shows that

$$\frac{d\langle f(B, t), B \rangle(t)}{dt} = \frac{\partial f}{\partial x}(B(t), t).$$

Example 8.25: (Representation of random variables)

1. If $Y = \int_0^T B(s)ds$, then $Y = \int_0^T (T-s)dB(s)$.
2. $Y = B^2(1)$. Then $M(t) = E(B^2(1)|\mathcal{F}_t) = B^2(t) + (1-t)$. Using Itô's formula for $M(t)$ we obtain $B^2(1) = 1 + 2 \int_0^1 B(t)dB(t)$.

Similar results hold for the Poisson process filtration.

Theorem 8.38 (Poisson Martingale Representation):

Let $M(t)$, $0 \leq t \leq T$, be a local martingale adapted to the Poisson filtration. Then there exists a predictable process $H(t)$ such that

$$M(t) = M(0) + \int_0^t H(s)d\bar{N}(s), \quad (8.68)$$

where $\bar{N}(t) = N(t) - t$ is the compensated Poisson process.

When a filtration is larger than the natural filtration of a martingale, then there is the following result (see Revuz and Yor (2001), Liptser and Shiryaev (2001)).

Theorem 8.39: *If $M(t)$, $0 \leq t \leq T$, is any continuous local martingale, and X a continuous \mathbb{F}^M -local martingale, then X has a representation*

$$X(t) = X(0) + \int_0^t H(s)dM(s) + Z(t), \quad (8.69)$$

where H is predictable and $\langle M, Z \rangle = 0$; (consequently $\langle X - Z, Z \rangle = 0$).

Example 8.26: Let \mathbb{F} be generated by two independent Brownian motions B and W , and let $M(t) = \int_0^t W(s)dB(s)$. It is a martingale, as a stochastic integral satisfying $E \int_0^t W^2(s)ds < \infty$. We show that M does not have the predictable representation property. $\langle M, M \rangle(t) = \int_0^t W^2(s)ds$. Hence $W^2(t) = \frac{d\langle M, M \rangle(t)}{dt}$, which shows that $W^2(t)$ is \mathcal{F}_t^M -measurable. Hence the martingale $X(t) = W^2(t) - t$ is adapted to \mathcal{F}_t^M , but it is not an integral of a predictable process with respect to M . According to Itô's formula $X(t) = 2 \int_0^t W(s)dW(s)$. Hence $\langle X, M \rangle(t) = \int_0^t W^2(s)d\langle W, B \rangle(s) = 0$. Suppose there is H , such that $X(t) = \int_0^t H(u)dM(u)$. Then by (8.67) $H(t) = \frac{d\langle X, M \rangle(t)}{d\langle M, M \rangle(t)} = 0$, implying that $X(t) = 0$, which is a contradiction.

This example has an application in finance, it shows non-completeness of a stochastic volatility model.

Example 8.27: Let \mathbb{F} be generated by a Brownian motion B and a Poisson process N , and let $M(t) = B(t) + N(t) - t = B(t) + \bar{N}(t)$, where $\bar{N}(t) = N(t) - t$. M is a martingale, as a sum of two martingales. We show that M does not have the predictable representation property. $[M, M](t) = [B, B](t) + [N, N](t) = t + N(t)$. This shows that $N(t) = [M, M](t) - t$ is \mathcal{F}_t^M -measurable. Hence $B(t) = M(t) - N(t) + t$ is \mathcal{F}_t^M -measurable. Thus the martingale $X(t) = \int_0^t N(s-)dB(s)$ is \mathcal{F}_t^M -measurable, but it does not have a predictable representation. If it did, $\int_0^t N(s-)dB(s) = \int_0^t H(s)dB(s) + \int_0^t H(s)d\bar{N}(s)$, and $\int_0^t (N(s-) - H(s))dB(s) = \int_0^t H(s)d\bar{N}(s)$. Since the integral on the rhs is of finite variation, $H(s) = N(s-)$, for almost all s . Thus $\int_0^t H(s)d\bar{N}(s) = 0$. This is the same as $\int_0^t N(s-)dN(s) = \int_0^t N(s-)ds$, but this is impossible. In order to see a contradiction, let $t = T_2$ be the time of the second jump of N . Then $\int_0^{T_2} N(s-)dN(s) = 1$ and $\int_0^{T_2} N(s-)ds = T_2 - T_1$.

This example shows the non-completeness of models of stock prices with jumps. It can be generalized to a model that supports a martingale with a jump component.

Remark 8.7: Definition 8.34 of the predictable representation property given here agrees with the standard definition given for continuous

martingales, but it is different to the definition for predictable representation with respect to semimartingales given in Liptser and Shiryaev (1989), Jacod and Shiryaev (1987), and Protter (1992). The general definition allows for different predictable functions h and H to be used in the integrals with respect to the continuous martingale part M^c and the discrete martingale part M^d of M ,

$$X(t) = X(0) + \int_0^t h(s)dM^c(s) + \int_0^t H(s)dM^d(s).$$

In this definition, the martingale M in Example 8.27 has the predictable representation property.

The definition given here is more suitable for financial applications. According to the financial mathematics theory an option can be priced if it can be replicated, which means that it is an integral of a predictable process H with respect to the discounted stock price process M , which is a martingale. The process H represents the number of shares bought/sold so it does not make sense to have H consist of two different components.

8.13 Elements of the General Theory

The basic setup consists of the probability space (Ω, \mathcal{F}, P) , where \mathcal{F} is a σ -field on Ω , and P is a probability on \mathcal{F} . A stochastic process is a map from $\mathbb{R}^+ \times \Omega$ to \mathbb{R} , namely $(t, \omega) \rightarrow X(t, \omega)$. \mathbb{R}^+ has the Borel σ -field of measurable sets, and \mathcal{F} is the σ -field of measurable sets on Ω . Only measurable processes are considered, that is, for any $A \in \mathcal{B}$

$$\{(t, \omega) : X(t, \omega) \in A\} \in \mathcal{B}(\mathbb{R}^+) \times \mathcal{F}.$$

Theorem 8.40 (Fubini): *Let $X(t)$ be a measurable stochastic process. Then*

1. *P-almost surely the functions $X(t, \omega)$ (trajectories) are (Borel) measurable.*
2. *If $E X(t)$ exists for all t , then it is measurable as a function of t .*
3. *If $\int_a^b E|X(t)|dt < \infty$ P-almost surely for all $a < b$ then almost all trajectories $X(t)$ are integrable and $\int_a^b E X(t)dt = E \int_a^b X(t)dt$.*

Let \mathbb{F} be a filtration of increasing σ -fields on Ω . Important classes of processes are introduced via measurability with respect to various σ -fields of subsets of $\mathbb{R}^+ \times \Omega$: adapted processes, progressively measurable processes, optional processes, and predictable processes, given in the order of inclusion.

X is adapted if, for all t , $X(t)$ is \mathcal{F}_t measurable. X is progressively measurable if, for any t , $\{(s \leq t, \omega) : X(s, \omega) \in A\} \in \mathcal{B}([0, t]) \times \mathcal{F}_t$. Any progressively measurable process is clearly adapted. It can be shown that any right- or left-continuous process is progressively measurable.

Definition 8.41:

1. The σ -field generated by the adapted left-continuous processes is called the predictable σ -field \mathcal{P} .
2. The σ -field generated by the adapted right-continuous processes is called the optional σ -field \mathcal{O} .
3. A process is called predictable if it is measurable with respect to the predictable σ -field \mathcal{P} ; it is called optional if it is measurable with respect to the optional σ -field \mathcal{O} .

Remarks:

1. The predictable σ -field \mathcal{P} is also generated by the adapted continuous processes.
2. Define $\mathcal{F}_{t-} = \sigma(\cup_{s < t} \mathcal{F}_s)$ the smallest σ -field containing \mathcal{F}_s for all $s < t$, and $\mathcal{F}_{0-} = \mathcal{F}_0$. \mathcal{F}_{t-} represents the information available prior to t . Then the predictable σ -field \mathcal{P} is generated by the sets $[s, t) \times A$ with $s < t$ and $A \in \mathcal{F}_{t-}$.
3. Predictable and optional σ -fields are also generated respectively by simple adapted left-continuous processes and simple adapted right-continuous processes.
4. Since a left-continuous adapted process $H(t)$ can be approximated by right-continuous adapted processes, ($H(t) = \lim_{\epsilon \rightarrow 0} H((t - \epsilon) +)$), any predictable process is also optional. Therefore $\mathcal{P} \subseteq \mathcal{O}$.
5. The Poisson process is right-continuous and it can be shown that it cannot be approximated by left-continuous adapted processes. Therefore there are optional processes which are not predictable, $\mathcal{P} \subset \mathcal{O}$.
6. In discrete time optional is the same as adapted.

Stochastic Sets

Subsets of $\mathbb{R}^+ \times \Omega$ are called stochastic sets. If A is a stochastic set, then “its projection on Ω ” $\pi_A = \{\omega : \exists t \text{ such that } (t, \omega) \in A\}$. A is called *evanescent* if $P(\pi_A) = 0$.

Two processes $X(t)$ and $Y(t)$ are called indistinguishable if the stochastic set $A = \{(t, \omega) : X(t, \omega) \neq Y(t, \omega)\}$ is an evanescent set. A process indistinguishable from zero is called P-negligible.

If τ_1 and τ_2 are stopping times, a closed stochastic interval is defined as $[[\tau_1, \tau_2]] = \{(t, \omega) : \tau_1(\omega) \leq t \leq \tau_2(\omega)\}$. Similarly, half-closed $[[\tau_1, \tau_2[$, $]\tau_1, \tau_2]]$, and open $]\tau_1, \tau_2[$ stochastic intervals are defined. Double brackets are used to emphasize that the intervals are subsets of $\mathbb{R}^+ \times \Omega$ and to distinguish them from intervals on \mathbb{R}^+ .

The stochastic interval $[[\tau, \tau]] = \{(t, \omega) : \tau(\omega) = t\}$ is called the graph of the stopping time τ .

A stochastic set A is called *thin* if there are stopping times τ_n such that $A = \cup_n [[\tau_n]]$.

Example 8.28: Let $N(t)$ be a Poisson process with rate λ , and $A = \{\Delta N \neq 0\}$. Then $A = \cup_n [[\tau_n]]$, where τ_n is the time of the n -th jump. Hence A is a thin set.

It can be shown that for any regular right-continuous process the stochastic set of jumps $\{\Delta X \neq 0\}$ is a thin set (Liptser and Shiryaev (1989)).

Classification of Stopping Times

Recall that τ is a stopping time with respect to filtration \mathbb{F} if for all $t \geq 0$ the event $\{\tau \leq t\} \in \mathcal{F}_t$. If \mathbb{F} is right-continuous then also $\{\tau < t\} \in \mathcal{F}_t$. Events observed before or at time τ are described by the σ -field \mathcal{F}_τ , defined as the collection of sets $\{A \in \mathcal{F} : \text{for any } t \ A \cap \{\tau \leq t\} \in \mathcal{F}_t\}$. Events observed before time τ are described by the σ -field $\mathcal{F}_{\tau-}$, the σ -field generated by \mathcal{F}_0 and the sets $A \cap \{\tau > t\}$, where $A \in \mathcal{F}_t$, $t > 0$.

There are three types of stopping times that are used in stochastic calculus:

1. Predictable stopping times.
2. Accessible stopping times.
3. Totally inaccessible stopping times.

τ is a predictable stopping time if there exists a sequence of stopping times τ_n , $\tau_n < \tau$, and $\lim_n \tau_n = \tau$. In this case it is said that the sequence τ_n announces τ .

Example 8.29: If τ is a stopping time, then for any constant $a > 0$, $\tau + a$ is a predictable stopping time. Indeed, it can be approached by $\tau_n = \tau + a - 1/n$.

Example 8.30: Let $B(t)$ be Brownian motion started at 0, \mathbb{F} its natural filtration, and τ the first hitting time of 1, that is, $\tau = \inf\{t : B(t) = 1\}$. τ is a predictable stopping time, since $\tau_n = \inf\{t : B(t) = 1 - 1/n\}$ converge to τ .

τ is an accessible stopping time if it is possible to announce τ , but with different sequences on different parts of Ω , that is, $[[\tau]] \subset \cup_n [[\tau_n]]$, where τ_n are predictable stopping times. All other types of stopping times are called totally inaccessible.

Example 8.31: Let $N(t)$ be a Poisson process, \mathbb{F} its natural filtration, and τ the time of the first jump, $\tau = \inf\{t : N(t) = 1\}$. τ is a totally inaccessible stopping time. Any predictable stopping time $\tau_n < \tau$ is a constant, since $\mathcal{F}_t \cap \{t < \tau\}$ is trivial. However, τ has a continuous distribution (exponential), thus it cannot be approached by constants.

The optional σ -field is generated by the stochastic intervals $[[0, \tau[$, where τ is a stopping time. The predictable σ -field is generated by the stochastic intervals $[[0, \tau]]$.

A set A is called predictable if its indicator is a predictable process, $I_A \in \mathcal{P}$.

The following results allow us to decide on predictability.

Theorem 8.42: *Let $X(t)$ be a predictable process and τ be a stopping time. Then*

1. $X(\tau)I(\tau < \infty)$ is $\mathcal{F}_{\tau-}$ measurable.
2. The stopped process $X(t \wedge \tau)$ is predictable.

For a proof see, for example, Liptser and Shiryaev (1989).

Theorem 8.43: *An adapted regular process is predictable if, and only if, for any predictable stopping time τ the random variable $X(\tau)I(\tau < \infty)$ is $\mathcal{F}_{\tau-}$ measurable and for each totally inaccessible stopping time τ one of the following two conditions hold:*

1. $X(\tau) = X(\tau-)$ on $\tau < \infty$.
2. The set $\{\Delta X \neq 0\} \cap [[\tau]]$ is P-evanescent.

For a proof see, for example, Liptser and Shiryaev (1989).

Theorem 8.44: *A stopping time τ is predictable if, and only if, for any bounded martingale M , $E(M(\tau)I(\tau < \infty)) = E(M(\tau-)I(\tau < \infty))$.*

Theorem 8.45: *The compensator $A(t)$ is continuous if, and only if, the jump times of the process $X(t)$ are totally inaccessible.*

See for example Liptser and Shiryaev (2001) for the proof.

Example 8.32: The compensator of the Poisson process is t , which is continuous. According to the above result the jump times of the Poisson process are totally inaccessible. This was shown in Example 8.31.

It can be shown that for Brownian motion and diffusions any stopping time is predictable. This implies that the class of optional processes is the same as the class of predictable processes.

Theorem 8.46: *For Brownian motion filtration any martingale (local martingale) is continuous and any positive stopping time is predictable. Any optional process is also predictable, $\mathcal{O} = \mathcal{P}$.*

A similar result holds for diffusions (see, for example, Rogers and Williams (1990)).

Remark 8.8: It can be shown that $\langle X, X \rangle$ is the conditional quadratic variation of $[X, X]$ conditioned on the predictable events \mathcal{P} .

8.14 Random Measures and Canonical Decomposition

The canonical decomposition of semimartingales with jumps uses the concepts of a random measure and its compensator, as well as integrals, with respect to random measures. We do not use this material elsewhere in the book. However, the canonical decomposition is often met in research papers.

Random Measure for a Single Jump

Let ξ be a random variable. For a Borel set $A \subset \mathbb{R}$ define

$$\mu(\omega, A) = I_A(\xi(\omega)) = I(\xi(\omega) \in A). \quad (8.70)$$

Then μ is a random measure, meaning that for each $\omega \in \Omega$, $\mu(\omega, A)$ is a measure when A varies, $A \in \mathcal{B}(\mathbb{R})$. Its (random) distribution function has a single jump of size 1 at ξ . The following random Stieltjes integrals consist of a single term:

$$\int_{\mathbb{R}} x \mu(\omega, dx) = \xi(\omega), \text{ and for a function } h, \int_{\mathbb{R}} h(x) \mu(\omega, dx) = h(\xi(\omega)). \quad (8.71)$$

There is a special notation for this random integral

$$h * \mu := \int_{\mathbb{R}} h(x) \mu(dx). \quad (8.72)$$

Random Measure of Jumps and its Compensator in Discrete Time

Let X_0, \dots, X_n, \dots be a sequence of random variables, adapted to \mathcal{F}_n , and let $\xi_n = \Delta X_n = X_n - X_{n-1}$. Let $\mu_n = I_A(\xi_n)$ be jump measures, and let

$$\nu_n(A) = E(\mu_n(A)|\mathcal{F}_{n-1}) = E(I_A(\xi_n)|\mathcal{F}_{n-1}) = P(\xi_n \in A|\mathcal{F}_{n-1})$$

be the conditional distributions, $n = 1, 2, \dots$. Define

$$\mu((0, n], A) = \sum_{i=1}^n \mu_i(A), \quad \text{and} \quad \nu((0, n], A) = \sum_{i=1}^n \nu_i(A). \quad (8.73)$$

Then for each A the sequence $\mu((0, n], A) - \nu((0, n], A)$ is a martingale. The measure $\mu((0, n], A)$ is called the measure of jumps of the sequence X_n , and $\nu((0, n], A)$ its compensator (A does not include 0). Clearly, the measure $\mu = \{\mu((0, n])\}_{n \geq 1}$ admits representation

$$\mu = \nu + (\mu - \nu), \quad (8.74)$$

where $\nu = \{\nu((0, n])\}_{n \geq 1}$ is predictable, and $\mu - \nu = \{\mu((0, n]) - \nu((0, n])\}_{n \geq 1}$ is a martingale. This is Doob's decomposition for random measures. With notation (8.72)

$$X_n = X_0 + (x * \mu)_n. \quad (8.75)$$

Regular conditional distributions exist, and for a function $h(x)$ the conditional expectations can be written as integrals with respect to these

$$E(h(\xi_n)|\mathcal{F}_{n-1}) = \int_{\mathbb{R}} h(x)\nu_n(dx),$$

provided $h(\xi_n)$ is integrable, $E|h(\xi_n)| < \infty$.

Assume now that ξ_n are integrable, then it's Doob's decomposition (8.4)

$$X_n = X_0 + \sum_{i=1}^n E(\xi_i|\mathcal{F}_{i-1}) + \sum_{i=1}^n (\xi_i - E(\xi_i|\mathcal{F}_{i-1})) = X_0 + A_n + M_n. \quad (8.76)$$

Using random measures and their integrals, we can express A_n and M_n as

$$A_n = \sum_{i=1}^n E(\xi_i|\mathcal{F}_{i-1}) = \sum_{i=1}^n \int_{\mathbb{R}} x\nu_i(dx) = (x * \nu)_n, \quad (8.77)$$

$$M_n = \sum_{i=1}^n (\xi_i - E(\xi_i|\mathcal{F}_{i-1})) = \sum_{i=1}^n \int_{\mathbb{R}} x(\mu_i(dx) - \nu_i(dx)) = (x * (\mu - \nu))_n.$$

Thus the semimartingale decomposition of X is given by using the random measure and its compensator

$$X_n = X_0 + (x * \nu)_n + (x * (\mu - \nu))_n. \quad (8.78)$$

However, the jumps of X , $\xi_n = \Delta X_n$, may not be integrable. Then the term $(x * \nu)_n$ is not defined. In this case a truncation function is used, such as $h(x) = xI(|x| \leq 1)$, and a similar decomposition is achieved, called the canonical decomposition,

$$\begin{aligned} X_n &= X_0 + \sum_{i=1}^n E(h(\xi_i) | \mathcal{F}_{i-1}) + \sum_{i=1}^n (h(\xi_i) - E(h(\xi_i) | \mathcal{F}_{i-1})) \\ &\quad + \sum_{i=1}^n (\xi_i - h(\xi_i)) \\ &= X_0 + (h * \nu)_n + (h * (\mu - \nu))_n + ((x - h(x)) * \mu)_n. \end{aligned} \quad (8.79)$$

The above representation has well-defined terms, and in addition it has another advantage that carries over to the continuous time case.

Random Measure of Jumps and its Compensator

Let X be a semimartingale. For a fixed t consider the jump $\Delta X(t)$. Taking $\xi = \Delta X(t)$ we obtain the measure of the jump at t

$$\mu(\{t\}, A) = I_A(\Delta X(t)), \quad \text{with } \int_{\mathbb{R}} x \mu(\{t\}, dx) = \Delta X(t). \quad (8.80)$$

Now consider the measure of jumps of X (in $\mathbb{R}^+ \times \mathbb{R}^0$, with $\mathbb{R}^0 = \mathbb{R} \setminus 0$)

$$\mu((0, t] \times A) = \sum_{0 < s \leq t} I_A(\Delta X(s)) \quad (8.81)$$

for a Borel set A that does not include 0 (there are no jumps of size 0; if $\Delta X(t) = 0$, then t is a point of continuity of X). It is possible to define the compensator of μ such that $\mu((0, t] \times A) - \nu((0, t] \times A)$ is a local martingale.

For the canonical decomposition of semimartingales, similar to (8.79), firstly large jumps are taken out, then the small jumps are compensated as follows. Consider

$$(x - h(x)) * \mu(t) = \int_0^t \int_{\mathbb{R} \setminus 0} (x - h(x)) \mu(ds, dx) = \sum_{s \leq t} (\Delta X_s - h(\Delta X_s)),$$

where $h(x)$ is a truncation function. This is a sum over “large” jumps with $|\Delta X(s)| > 1$ (since $x - h(x) = 0$ for $|x| \leq 1$). Since the sum of squares of jumps is finite (Corollary 8.31, (8.51)) the above sum has only finitely many terms, hence it is finite. Thus the following canonical decomposition of a semimartingale is obtained

$$\begin{aligned} X(t) &= X(0) + A(t) + X^{cm}(t) + (h(x) * (\mu - \nu))(t) + ((x - h(x)) * \mu)(t), \\ &= X(0) + A(t) + X^{cm}(t) + \int_0^t \int_{|x| \leq 1} x d(\mu - \nu) + \int_0^t \int_{|x| > 1} x d\mu, \end{aligned}$$

where A is a predictable process of finite variation, X^{cm} is a continuous martingale component of X , μ is the measure of jumps of X , and ν its compensator. For a proof see Liptser and Shiryaev (1989) p. 188, and Shiryaev (1999) p. 663.

Let $C = \langle X^{cm}, X^{cm} \rangle$ (it always exists for continuous processes). The following three processes appearing in the canonical decomposition (A, C, ν) are called the *triplet of predictable characteristics* of the semimartingale X .

Notes: Material for this chapter is based on Protter (1992), Rogers and Williams (1990), Métivier (1982), Liptser and Shiryaev (1989), and Shiryaev (1999).

8.15 Exercises

Exercise 8.1: Let $\tau_1 < \tau_2$ be stopping times. Show that $I_{(\tau_1, \tau_2]}(t)$ is a simple predictable process.

Exercise 8.2: Let $H(t)$ be a regular adapted process, not necessarily left-continuous. Show that for any $\delta > 0$, $H(t - \delta)$ is predictable.

Exercise 8.3: Show that a continuous process is locally integrable. Show that a continuous local martingale is locally square integrable.

Exercise 8.4: M is a local martingale and $E \int_0^T H^2(s)d[M, M](s) < \infty$. Show that $\int_0^t H(s)dM(s)$ is a square integrable martingale.

Exercise 8.5: Find the variance of $\int_0^1 N(t-)dM(t)$, where M is the compensated Poisson process $M(t) - t$.

Exercise 8.6: If S and T are stopping times, show that

1. $S \wedge T$ and $S \vee T$ are stopping times.
2. The events $\{S = T\}$, $\{S \leq T\}$ and $\{S < T\}$ are in \mathcal{F}_S .
3. $\mathcal{F}_S \cap \{S \leq T\} \subset \mathcal{F}_T \cap \{S \leq T\}$.

Exercise 8.7: Let U be a positive random variable on a probability space (Ω, \mathcal{F}, P) , and let \mathcal{G} be a sub- σ -field of \mathcal{F} .

1. Let $t \geq 0$. Show that $\mathcal{F}_t := \{A \in \mathcal{F} : \exists B \in \mathcal{G} \text{ such that } A \cap \{U > t\} = B \cap \{U > t\}\}$ is a σ -field.
2. Show that \mathcal{F}_t is a right-continuous filtration on (Ω, \mathcal{F}, P) .
3. Show that U is a stopping time for \mathcal{F}_t .
4. What are \mathcal{F}_0 , \mathcal{F}_{U-} , and \mathcal{F}_U equal to?

Exercise 8.8: Let U_1, U_2, \dots be (strictly) positive random variables on a probability space (Ω, \mathcal{F}, P) , and $\mathcal{G}_1, \mathcal{G}_2, \dots$ be sub- σ -fields of \mathcal{F} . Suppose that for all n , U_1, U_2, \dots, U_n are \mathcal{G}_n -measurable and denote by T_n the random variable $\sum_{i=1}^n U_i$. Set $\mathcal{F}_t = \bigcap_n \{A \in \mathcal{F} : \exists B_n \in \mathcal{G}_n \text{ such that } A \cap \{T_n > t\} = B_n \cap \{T_n > t\}\}$.

1. Show that \mathcal{F}_t is a right-continuous filtration on (Ω, \mathcal{F}, P) .
2. Show that for all n , T_n is a stopping time for \mathcal{F}_t .
3. Suppose that $\lim_n T_n = \infty$ almost surely. Show that $\mathcal{F}_{T_n} = \mathcal{G}_{n+1}$ and $\mathcal{F}_{T_n-} = \mathcal{G}_n$.

Exercise 8.9: Let $B(t)$ be a Brownian motion and $H(t)$ be a predictable process. Show that $M(t) = \int_0^t H(s)dB(s)$ is a Brownian motion if, and only if, $\text{Leb}(\{t : |H(t)| \neq 1\}) = 0$ almost surely.

Exercise 8.10: Let T be a stopping time. Show that the process $M(t) = 2B(t \wedge T) - B(t)$ obtained by reflecting $B(t)$ at time T , is a Brownian motion.

Exercise 8.11: Let $B(t)$ and $N(t)$ be respectively a Brownian motion and a Poisson process on the same space. Denote by $\bar{N}(t) = N(t) - t$ the compensated Poisson process. Show that the following processes are martingales: $B(t)\bar{N}(t)$, $\mathcal{E}(B)(t)\bar{N}(t)$, and $\mathcal{E}(\bar{N})(t)B(t)$.

Exercise 8.12: $X(t)$ solves the SDE $dX(t) = \mu X(t)dt + aX(t-)dN(t) + \sigma X(t)dB(t)$. Find the condition for $X(t)$ to be a martingale.

Exercise 8.13: Find the predictable representation for $Y = B^5(1)$.

Exercise 8.14: Find the predictable representation for the martingale $e^{B(t)-t/2}$.

Exercise 8.15: Let $Y = \int_0^1 \text{sign}(B(s))dB(s)$. Show that Y has a normal distribution. Show that there is no deterministic function $H(s)$ such that $Y = \int_0^1 H(s)dB(s)$. This shows that the assumption that Y, B are jointly Gaussian in Theorem 8.35 is indispensable.

Exercise 8.16: Find the quadratic variation of $|B(t)|$.

Exercise 8.17: (Stochastic logarithm)

Let U be a semimartingale such that $U(t)$ and $U(t-)$ are never zero. Show that there exists a unique semimartingale X with $X(0) = 0$, ($X = \mathcal{L}(U)$) such that $dX(t) = \frac{dU(t)}{U(t-)}$, and

$$\begin{aligned} X(t) &= \ln \left| \frac{U(t)}{U(0)} \right| + \frac{1}{2} \int_0^t \frac{d \langle U^c, U^c \rangle(s)}{U^2(s-)} \\ &\quad - \sum_{s \leq t} \left(\ln \left| \frac{U(s)}{U(s-)} \right| + 1 - \frac{U(s)}{U(s-)} \right), \end{aligned} \tag{8.82}$$

see Kallsen and Shiryaev (2002).

Chapter 9

Pure Jump Processes

In this chapter we consider pure jump processes, that is, processes that change only by jumps. Counting processes and Markov jump processes are defined and their semimartingale representation is given. This representation allows us to see the process as a solution to a stochastic equation driven by discontinuous martingales.

9.1 Definitions

A counting process is determined by a sequence of non-negative random variables T_n , satisfying $T_n < T_{n+1}$ if $T_n < \infty$ and $T_n = T_{n+1}$ if $T_n = \infty$. T_n can be considered as the time of the n -th occurrence of an event, and they are often referred to as arrival times. $N(t)$ counts the number of events that occurred up to time t , that is,

$$N(t) = \sum_{n=1}^{\infty} I(T_n \leq t), \quad N(0) = 0. \quad (9.1)$$

$N(t)$ is piece-wise constant and has jumps of size one at the points T_n . Such processes are also known as simple point processes to distinguish them from more general *marked point processes*, which are described by a sequence (T_n, Z_n) for some random variables Z_n . Z_n , for example, may describe the size of jump at T_n .

The pure jump process X is defined as follows:

$$X(t) = X(0) + \sum_{n=1}^{\infty} I(T_n \leq t)Z_n. \quad (9.2)$$

Note that X in (9.2) is right-continuous, piece-wise constant with the time of the n -th jump at T_n , and $Z_n = X(T_n) - X(T_n-) = X(T_n) - X(T_{n-1})$ is the size of the jump at T_n .

9.2 Pure Jump Process Filtration

The filtration \mathbb{F} considered here is the natural filtration of the process. We recall related definitions. For a process X its natural filtration is defined by (the augmentation of) σ -fields $\mathcal{F}_t = \sigma(X(s), 0 \leq s \leq t)$ and represents the information obtained by observing the process on $[0, t]$. The strict past is the information obtained by observing the process on $[0, t)$ and is denoted by $\mathcal{F}_{t-} = \sigma(X(s), 0 \leq s < t)$.

A non-negative random variable τ , which is allowed to be infinity, is a stopping time if $\{\tau \leq t\} \in \mathcal{F}_t$ for every t . Thus by observing the process on $[0, t]$ it is possible to deduce whether τ has occurred.

Information obtained from observing the process up to a stopping time τ is \mathcal{F}_τ , defined by $\mathcal{F}_\tau = \{A \in \mathcal{F}: \text{for any } t, A \cap \{\tau \leq t\} \in \mathcal{F}_t\}$. The strict past of X at τ is described by the σ -field $\mathcal{F}_{\tau-} = \sigma(A \cap \{t < \tau\} : t \geq 0, A \in \mathcal{F}_t) \vee \mathcal{F}_0$. Note that $\tau \in \mathcal{F}_{\tau-}$ (take $A = \{\tau > t\} \in \mathcal{F}_t$). Clearly, $\mathcal{F}_{\tau-} \subset \mathcal{F}_\tau$.

Clearly, the arrival times T_n are stopping times for \mathbb{F} . They are usually taken as a localizing sequence. Note that $\mathcal{F}_{T_n} = \sigma((T_i, Z_i), i \leq n)$ and that $X(T_n-) = X(T_{n-1})$, since for t satisfying $T_{n-1} \leq t < T_n$, $X(t) = X(T_{n-1})$, and this value is kept until the next jump at time T_n . As $T_n \in \mathcal{F}_{T_n-}$, $\mathcal{F}_{T_n-} = \sigma((T_i, Z_i), i \leq n-1, T_n)$. Thus Z_n , the jump size at T_n , is the only information in \mathcal{F}_{T_n} not available in \mathcal{F}_{T_n-} .

It can be shown that \mathcal{F}_t is right-continuous, that is, $\mathcal{F}_t = \mathcal{F}_{t+}$, as well as the following result, which is essential in finding compensators.

Theorem 9.1: *If τ is a stopping time, then for each n there is a random variable ζ_n which is \mathcal{F}_{T_n} -measurable such that*

$$\tau \wedge T_{n+1} = (T_n + \zeta_n) \wedge T_{n+1} \quad \text{on } \{T_n \leq \tau\}. \quad (9.3)$$

Proofs of the above results can be found, for example, in Liptser and Shiryaev (1974), and Karr (1986).

Since compensators are predictable processes, it is important to have some criteria to decide on predictability. By definition, any adapted left-continuous or continuous process is predictable. The following construction, which is often met in calculations, results in a predictable process.

Theorem 9.2: *Let T_n be the arrival times in a pure jump process, and for all $n = 0, 1, \dots$, $Y_n(t)$ be an adapted process such that for any $t \in (T_n, T_{n+1})$ it is \mathcal{F}_{T_n} -measurable. Then the process $X(t) = \sum_{n=0}^{\infty} Y_n(t)I(T_n < t \leq T_{n+1})$ is predictable.*

PROOF. We outline the proof. The process $I_n(t) = I(T_n < t \leq T_{n+1})$ is predictable. Indeed, $I_n(t) = I(T_n < t \leq T_{n+1}) = I(t \leq T_{n+1}) - I(t \leq T_n)$. Since for each n , T_n is a stopping time, $\{T_n \geq t\} \in \mathcal{F}_t$ and therefore $I_n(t)$ is adapted. However, it is left-continuous, hence it is predictable. Since $Y_n(t)$ is “known” when $T_n < t \leq T_{n+1}$, $X(t)$ is predictable. \square

Assumptions

$T_\infty = \lim_{n \rightarrow \infty} T_n$ exists since T_n 's are non-decreasing. The results given below hold for $t < T_\infty$, and in order to avoid repetitions we assume throughout that $T_\infty = \infty$, unless stated otherwise. In the case of $T_\infty < \infty$ there are infinitely many jumps on the finite time interval $(0, T_\infty)$, and it is said that explosion occurs. We assume that there are no explosions. Sufficient conditions for Markov jump processes are given later.

Assume that the jumps are integrable, $E|Z_n| < \infty$ for all n . Under this assumption X is locally integrable, since $E|X(t \wedge T_n)| \leq \sum_{i=1}^n E|Z_i| < \infty$, and therefore it has a uniquely defined compensator A .

$M(t) = X(t) - A(t)$ is the local martingale associated with X , also called the innovation martingale.

9.3 Itô's Formula for Processes of Finite Variation

If a semimartingale X is of finite variation, then its continuous martingale part $X^{cm} = 0$, consequently $\langle X, X \rangle^c(t) = \langle X^{cm}, X^{cm} \rangle(t) = 0$. Therefore the term containing the second derivative f'' disappears in (8.59). Moreover, since X is of finite variation its continuous part X^c satisfies $dX^c(t) = dX(t) - \Delta X(t)$, and Itô's formula takes the form: for any C^1 function f

$$f(X(t)) - f(X(0)) = \int_0^t f'(X(s-))dX^c(s) + \sum_{s \leq t} (f(X(s)) - f(X(s-))). \quad (9.4)$$

If the continuous part X^c is zero, then the formula is an identity, representing a function as the sum of its jumps. A similar formula holds for a function of n variables.

Stochastic Exponential

The stochastic exponential (8.62) of finite variation processes simplifies to

$$\mathcal{E}(X)(t) = e^{X(t) - X(0)} \prod_{s \leq t} (1 + \Delta X(s)) e^{-\Delta X(s)} = e^{X^c(t)} \prod_{s \leq t} (1 + \Delta X(s)), \quad (9.5)$$

where X^c is the continuous part of X . The last equality is due to $X^c(t) = X(t) - X(0) - \sum_{s \leq t} \Delta X(s)$.

Example 9.1: Let $X(t)$ be a process with jumps of size one (counting process), so that $\Delta X(s) = 0$ or 1. Its stochastic exponential is given by

$$\mathcal{E}(X)(t) = \prod_{s \leq t} (1 + \Delta X(s)) = 2^{X(t) - X(0)}. \quad (9.6)$$

Integration by Parts for Processes of Finite Variation

The integration by parts formula is obtained directly from the integral representation of the quadratic covariation (8.23) and (1.20). Recall that if X or Y are of finite variation, then their quadratic covariation is given by

$$\begin{aligned} [X, Y](t) &= \sum_{0 < s \leq t} \Delta X(s) \Delta Y(s) \quad \text{and} \\ [X](t) &= [X, X](t) = \sum_{0 < s \leq t} (\Delta X(s))^2. \end{aligned} \quad (9.7)$$

Using (8.23) we obtain

$$\begin{aligned} X(t)Y(t) - X(0)Y(0) \\ = \int_0^t X(s-)dY(s) + \int_0^t Y(s-)dX(s) + \sum_{0 < s \leq t} \Delta X(s) \Delta Y(s). \end{aligned} \quad (9.8)$$

Remark 9.1: The following formula holds for finite variation processes:

$$\sum_{s \leq t} \Delta X(s) \Delta Y(s) = \int_0^t \Delta X(s) dY(s). \quad (9.9)$$

Indeed, by letting $Y^c(t) = Y(t) - \sum_{s \leq t} \Delta Y(s)$ be the continuous part of Y , we have

$$\int_0^t \Delta X(s) dY(s) - \sum_{s \leq t} \Delta X(s) \Delta Y(s) = \int_0^t \Delta X(s) dY^c(s) = 0, \quad (9.10)$$

since $Y^c(s)$ is continuous and $\Delta X(s)$ is different from zero at mostly countably many points.

9.4 Counting Processes

Let N be a counting process, then it is a pure jump process with jumps of size one. Thus

$$[N, N](t) = \sum_{s \leq t} (\Delta N(s))^2 = N(t). \quad (9.11)$$

Theorem 9.3: *The compensator A and the sharp bracket process of N are the same, $A = \langle N, N \rangle$.*

PROOF. N is of locally integrable variation since $N(t \wedge T_n) \leq n$ and T_n is a localizing sequence. A is the unique predictable process such that $N(t) - A(t)$ is a local martingale. $\langle N, N \rangle$ is the unique predictable process such that $[N, N](t) - \langle N, N \rangle(t)$ is a local martingale. The result follows from (9.11) and the uniqueness of the compensator. \square

Heuristically the compensator for a counting process is given by

$$dA(t) = E(dN(t)|\mathcal{F}_{t-}), \quad (9.12)$$

where \mathcal{F}_{t-} denotes the information available prior to time t by observing the process over $[0, t)$, and $dN(t) = N(t + dt) - N(t)$. $dM(t) = d(N - A)(t) = dN(t) - dA(t)$ is that part of $dN(t)$ that cannot be foreseen from the observations of N over $[0, t)$.

The next result shows the relation between the sharp bracket of the martingale and the compensator in a counting process. Since the proof is a straight application of stochastic calculus rules it is given below. It is useful for the calculation of the variance of M , indeed by Theorem 8.24 $EM^2(t) = E\langle M, M \rangle(t)$.

Theorem 9.4: *Let $M = N - A$. Then $\langle M, M \rangle(t) = \int_0^t (1 - \Delta A(s))dA(s)$. In particular, if A is continuous, then $\langle M, M \rangle(t) = A(t)$.*

PROOF. By integration by parts (9.8) we have

$$M^2(t) = 2 \int_0^t M(s-)dM(s) + \sum_{s \leq t} (\Delta M(s))^2. \quad (9.13)$$

Use $\Delta M(s) = \Delta N(s) - \Delta A(s)$ and expand the sum to obtain

$$\sum_{s \leq t} (\Delta M(s))^2 = \sum_{s \leq t} \Delta N(s) - 2 \sum_{s \leq t} \Delta N(s) \Delta A(s) + \sum_{s \leq t} (\Delta A(s))^2,$$

where we used that since N is a simple process $(\Delta N(s))^2 = \Delta N(s)$. Thus we obtain by using formula (9.9) and $N = M + A$,

$$\begin{aligned} M^2(t) &= 2 \int_0^t M(s-)dM(s) + N(t) - 2 \int_0^t \Delta A(s)dN(s) + \int_0^t \Delta A(s)dA(s) \\ &= \int_0^t (2M(s-) + 1 - 2\Delta A(s))dM(s) + \int_0^t (1 - \Delta A(s))dA(s). \end{aligned}$$

The process in the first integral is predictable because $M(s-)$ is adapted and left-continuous, $A(s)$ is predictable, so that $\Delta A(s)$ is also predictable. Therefore the first integral is a local martingale. The second integral is predictable since A is predictable. Thus the above equation is the Doob–Meyer decomposition of M^2 . The result now follows by the uniqueness of the Doob–Meyer decomposition. \square

We give examples of processes and their compensators next.

Point Process of a Single Jump

Let T be a random variable and define the process N as a single jump of size 1 at time T , that is, $N(t) = I(T \leq t)$. Let F be the distribution function of T .

Theorem 9.5: *The compensator $A(t)$ of $N(t) = I(T \leq t)$ is given by*

$$A(t) = \int_0^{t \wedge T} \frac{dF(s)}{1 - F(s-)}. \quad (9.14)$$

PROOF. $A(t)$ is clearly predictable. In order to show that $N(t) - A(t)$ is a martingale, by Theorem 7.17 it is enough to show that $E(N(S)) = EA(S)$ for any stopping time S . According to Theorem 9.1 there exists an \mathcal{F}_0 -measurable (i.e. almost surely constant) random variable ζ such that

$$\{S \geq T\} = \{S \wedge T = T\} = \{\zeta \wedge T = T\} = \{T \leq \zeta\}.$$

Therefore

$$\begin{aligned} EN(S) &= P(S \geq T) = P(T \leq \zeta) \\ &= \int_0^\zeta dF(t) = \int_0^\zeta \frac{P(T \geq t)}{1 - F(t-)} dF(t) \\ &= E \left(\int_0^\zeta \frac{I(T \geq t)}{1 - F(t-)} dF(t) \right) = E \left(\int_0^{\zeta \wedge T} \frac{dF(t)}{1 - F(t-)} \right) \\ &= E \left(\int_0^{S \wedge T} \frac{dF(t)}{1 - F(t-)} \right). \end{aligned}$$

Thus (9.14) is obtained. \square

Corollary 9.6: *If F has a density f , then the compensator*

$$A(t) = \int_0^{t \wedge T} h(s) ds, \quad (9.15)$$

where

$$h(t) = \frac{f(t)}{1 - F(t)}, \quad (9.16)$$

called the hazard function, and gives the likelihood of the occurrence of the jump at t given that the jump has not occurred before t .

Compensators of Counting Processes

The next result gives an explicit form of the compensator of a general counting process. Since the proof uses the same ideas as above it is omitted.

Theorem 9.7: *Let N be a counting process generated by the sequence T_n . Denote by $U_{n+1} = T_{n+1} - T_n$ the inter-arrival times, $T_0 = 0$. Let $F_n(t) = P(U_{n+1} \leq t | T_1, \dots, T_n)$ denote the regular conditional distributions, and $F_0(t) = P(T_1 \leq t)$. Then the compensator $A(t)$ is given by*

$$A(t) = \sum_{i=0}^{\infty} \int_0^{t \wedge T_{i+1} - t \wedge T_i} \frac{dF_i(s)}{1 - F_i(s-)}. \quad (9.17)$$

Note that if the conditional distributions F_n in Theorem 9.7 are continuous with $F_n(0) = 0$, then by changing variables we have

$$\int_0^a \frac{dF_n(s)}{1 - F_n(s-)} = \int_0^a \frac{dF_n(s)}{1 - F_n(s)} = -\log(1 - F_n(a)), \quad (9.18)$$

and Equation (9.17) can be simplified accordingly.

Renewal Process

A renewal process N is a point process in which all inter-arrival times are i.i.d, that is, $T_1, T_2 - T_1, \dots, T_{n+1} - T_n$ are i.i.d. with distribution function $F(x)$. In this case all the conditional distributions F_n in Theorem 9.17 are given by F . As a result of Theorem 9.7 we obtain the following corollary.

Corollary 9.8: *Assume that the inter-arrival distribution is continuous and $F(0) = 0$. Then the compensator of the renewal process is given by*

$$A(t) = - \sum_{n=1}^{\infty} \log(1 - F(t \wedge T_n - t \wedge T_{n-1})). \quad (9.19)$$

Stochastic Intensity

Definition 9.9: If $A(t) = \int_0^t \lambda(s)ds$, where $\lambda(t)$ is a positive predictable process, then $\lambda(t)$ is called the stochastic intensity of N .

Note that if $A(t)$ is deterministic and differentiable with derivative $A'(t)$, and $A(t) = \int_0^t A'(s)ds$, then $\lambda(t) = A'(t)$ is predictable and is the stochastic intensity. If $A(t)$ is random and differentiable with derivative $A'(t)$, satisfying $A(t) = \int_0^t A'(s)ds$, then $\lambda(t) = A'(t-)$. Indeed, $A(t) = \int_0^t A'(s-)ds$.

If the stochastic intensity exists, then by definition of the compensator, $N(t) - \int_0^t \lambda(s)ds$ is a local martingale. For counting processes a heuristic interpretation of the intensity is given by

$$\lambda(t)dt = dA(t) = E(dN(t)|\mathcal{F}_{t-}) = P(dN(t) = 1|\mathcal{F}_{t-}). \quad (9.20)$$

If the stochastic intensity exists, then the compensator is continuous and by Theorem 9.4 the sharp bracket of the martingale $M = N - A$ is given by

$$\langle M, M \rangle(t) = \int_0^t \lambda(s)ds. \quad (9.21)$$

Example 9.2:

1. A deterministic point process is its own compensator, so it does not have stochastic intensity.
2. Stochastic intensity for the renewal process with continuous inter-arrival distribution F is given by $h(V(t-))$, where h is the hazard function and $V(t) = t - T_{N(t)}$, called the age process. This can be seen by differentiating the compensator in (9.19).
3. Stochastic intensity for the renewal process with a discrete inter-arrival distribution F does not exist.

Non-Homogeneous Poisson Processes

Theorem 9.10: Let $N(t)$ be point process with a continuous deterministic compensator $A(t)$. Then it has independent Poisson distributed increments, that is, the distribution of $N(t) - N(s)$ is Poisson with parameter $A(t) - A(s)$, $0 \leq s < t$.

If $A(t)$ has a density $\lambda(t)$, that is, $A(t) = \int_0^t \lambda(s)ds$, then $N(t)$ is called the non-homogeneous Poisson process with rate $\lambda(s)$.

PROOF. We prove the result by an application of the stochastic exponential. $M(t) = N(t) - A(t)$ is a martingale. For a fixed $0 < u < 1$, $-uM(t)$ is

also a martingale. Consider $\mathcal{E}(uM)(t)$. According to (9.5)

$$\begin{aligned}\mathcal{E}(-uM)(t) &= e^{uA(t)} \prod_{s \leq t} (1 - u\Delta M(s)) \\ &= e^{uA(t)} \prod_{s \leq t} (1 - u\Delta N(s)) = e^{uA(t)}(1-u)^{N(t)} \\ &= e^{uA(t)+N(t)\log(1-u)},\end{aligned}\tag{9.22}$$

where we have used that $\Delta N(s)$ is zero or one. The stochastic exponential of a martingale is always a local martingale, but in this case it is also a true martingale by Theorem 7.21. Indeed, since $A(t)$ is deterministic and non-decreasing,

$$\mathbb{E} \sup_{t \leq T} e^{uA(t)+N(t)\log(1-u)} \leq e^{uA(T)} \mathbb{E} \sup_{t \leq T} e^{N(t)\log(1-u)} \leq e^{uA(T)} < \infty,$$

and the condition of Theorem 7.21 is satisfied. Taking expectations in (9.22) we obtain the moment generating function of $N(t)$,

$$\mathbb{E}((1-u)^{N(t)}) = e^{-uA(t)}$$

or with $v = 1 - u$,

$$\mathbb{E}(v^{N(t)}) = e^{(v-1)A(t)}.\tag{9.23}$$

This shows that $N(t)$ is a Poisson random variable with parameter $A(t)$. If we take conditional expectation in (9.22) and use the martingale property, we obtain in the same way that for all $s < t$

$$\mathbb{E}(v^{N(t)-N(s)} | \mathcal{F}_s) = e^{(v-1)(A(t)-A(s))},\tag{9.24}$$

which shows that the distribution of $N(t) - N(s)$ does not depend on the past and is Poisson. \square

A similar result holds if the compensator is deterministic but discontinuous. The proof is more involved and can be found, for example, in Liptser and Shiryaev (1974) p. 279, where the form of the distribution of the increments is also given.

Theorem 9.11: *Let $N(t)$ be a point process with a deterministic compensator $A(t)$. Then it has independent increments.*

The following result states that a point process with a continuous, but possibly random, compensator can be transformed into a Poisson process by a random change of time. Compare this result to change of time for continuous martingales, Dambis–Dubins–Schwarz Theorem 7.37.

Theorem 9.12: Let a counting process $N(t)$ have a continuous compensator $A(t)$ and $\lim_{t \rightarrow \infty} A(t) = \infty$. Define $\rho(t) = \inf\{s \geq 0 : A(s) = t\}$. Let $K(t) = N(\rho(t))$ and $\mathcal{G}_t = \mathcal{F}_{\rho(t)}$. Then the process $K(t)$ with respect to filtration \mathcal{G}_t is Poisson with rate 1.

The proof can be found in Liptser and Shiryaev (1974) p. 280, and (2001), and is not given here. In order to convince ourselves of the result, consider the case when $A(t)$ is strictly increasing, then $\rho(t)$ is the usual inverse, that is, $A(\rho(t)) = t$. Then $EK(t) = EN(\rho(t)) = EA(\rho(t)) = t$ so that K has the right mean.

For more information on point processes see, for example, Liptser and Shiryaev (2001), (1989) and Karr (1986).

Compensators of Pure Jump Processes

Let for all $t \geq 0$

$$X(t) = X(0) + \sum_{n=1}^{\infty} Z_n I(T_n \leq t), \quad (9.25)$$

be a pure jump point process generated by the sequence (T_n, Z_n) . By using the same arguments as in the proof of Theorem 9.7 we can show the following result:

Theorem 9.13: Let $F_n(t) = P(T_{n+1} - T_n \leq t | \mathcal{F}_{T_n})$ denote the regular conditional distributions of inter-arrival times, $F_0(t) = P(T_1 \leq t)$, and $m_n = E(Z_{n+1} | \mathcal{F}_{T_n}) = E(X(T_{n+1}) - X(T_n) | \mathcal{F}_{T_n})$ denote the conditional expectation of the jump sizes. Then the compensator $A(t)$ is given by

$$A(t) = \sum_{n=0}^{\infty} m_n \int_0^{t \wedge T_{n+1} - t \wedge T_n} \frac{dF_n(s)}{1 - F_n(s-)}. \quad (9.26)$$

The following observation is frequently used in the calculus of pure jump processes:

Theorem 9.14: If $X(t)$ is a pure jump process, then for a function f , $f(X(t))$ is also a pure jump process with the same jump times T_n . The size of the jump at T_n is given by $Z'_n = f(X(T_n)) - f(X(T_n-))$. Consequently, if f is such that $E|Z'_n| < \infty$, then the compensator of $f(X)$ is given by (9.26) with m_n replaced by $m'_n = E(Z'_{n+1} | \mathcal{F}_{T_n})$.

Theorem 9.15: Let $X(t)$ be a pure jump point process generated by the sequence (T_n, Z_n) . Assume conditions and notations of Theorem 9.13 and that the compensator $A(t)$ is continuous. Suppose in addition that $EZ_n^2 < \infty$ and $v_n = E(Z_{n+1}^2 | \mathcal{F}_{T_n})$. Let $M(t) = X(t) - A(t)$, then $\langle M, M \rangle(t)$ is given by (9.26) with m_n replaced by v_n .

PROOF. Since A is assumed to be continuous (and it is always of finite variation)

$$[M, M](t) = [X - A, X - A](t) = [X, X](t) = \sum_{0 < s \leq t} (\Delta X(s))^2. \quad (9.27)$$

Thus $[M, M]$ is a pure jump process with jump times T_n and jump sizes $(\Delta X(T_n))^2 = Z_n^2$. Thus $[M, M](t)$ is a pure jump process generated by the sequence (T_n, Z_n^2) . $\langle M, M \rangle(t)$ is its compensator. The result follows by Theorem 9.13. \square

A particular case of pure jump processes is the class of processes with exponentially distributed inter-arrival times and independent jump sizes. This is the class of Markov jump processes.

9.5 Markov Jump Processes

Definitions

Let for all $t \geq 0$

$$X(t) = X(0) + \sum_{n=1}^{\infty} Z_n I(T_n \leq t), \quad (9.28)$$

where T_n and Z_n have the following conditional distributions. Given that $X(T_n) = x$, $T_{n+1} - T_n$ is exponentially distributed with mean $\lambda^{-1}(x)$, and independent of the past; and the jump $Z_{n+1} = X(T_{n+1}) - X(T_n)$ is independent of the past and has a distribution that depends only on x .

$$F_n(t) = P(T_{n+1} - T_n \leq t | \mathcal{F}_{T_n}) = 1 - e^{-\lambda(X(T_n))t}, \quad (9.29)$$

and for some family of distribution functions $K(x, \cdot)$

$$P(X(T_{n+1}) - X(T_n) \leq t | \mathcal{F}_{T_n}) = K(X(T_n), t), \quad (9.30)$$

$$E(X(T_{n+1}) - X(T_n) | \mathcal{F}_{T_n}) = m(X(T_n)) = m_n.$$

It is intuitively clear that X defined in this way possesses the Markov property due to the lack of memory of the exponential distribution. We omit the proof.

Heuristically, Markov jump processes can be described as follows. If the process is in state x then it stays there for an exponential length of time with mean $\lambda^{-1}(x)$ (parameter $\lambda(x)$) after which it jumps from x to a new state $x + \xi(x)$, where $P(\xi(x) \leq t) = K(x, t)$ denotes the distribution of the jump from x . The parameters of the process are: the function $\lambda(x)$ (the holding time parameter) and distributions $K(x, \cdot)$ (distributions of jump sizes).

$\lambda(x)$ is always non-negative. If for some x , $\lambda(x) = 0$, then once the process gets into x it stays in x forever, in this case the state x is called absorbing. We shall assume that there are no absorbing states. If $\lambda(x) = \infty$, then the process leaves x instantaneously. We assume that $\lambda(x)$ is finite on finite intervals so that there are no instantaneous states.

For construction, classification, and properties of Markov jump processes see, for example, Breiman (1968), Chung (1960), and Ethier and Kurtz (1986).

The Compensator and the Martingale

We derive the compensator heuristically first and then give the precise result. Suppose that $X(t) = x$. According to the lack of memory property of the exponential distribution it does not matter how long the process has already spent in x , the jump from x will still occur after exponentially distributed (with parameter $\lambda(x)$) random time. Therefore the conditional probability that a jump occurs in $(t, t + dt)$, given that it has not occurred before, is (with U having exponential $\exp(\lambda(x))$ distribution)

$$P(U \leq dt) = 1 - \exp(-\lambda(x)dt) \approx \lambda(x)dt.$$

When the jump occurs its size is $\xi(x)$, with mean $m(x) = E(\xi(x))$. Therefore

$$dA(t) = E(dX(t)|\mathcal{F}_{t-}) = \lambda(X(t))m(X(t))dt. \quad (9.31)$$

Assume that $\lim_{n \rightarrow \infty} T_n = T_\infty = \infty$ (such processes are called regular, and sufficient conditions for this are given in a later section). If $T_\infty < \infty$, then the compensator is given for times $t < T_\infty$.

Theorem 9.16: *Let X be a Markov jump process such that for all x , the holding time parameter is positive, $\lambda(x) > 0$, and the size of the jump from x is integrable with mean $m(x)$.*

1. *The compensator of X is given by*

$$A(t) = \int_0^t \lambda(X(s))m(X(s))ds. \quad (9.32)$$

2. *Suppose the second moments of the jumps are finite, $v(x) = E\xi^2(x) < \infty$. Then the sharp bracket of the local martingale $M(t) = X(t) - A(t)$ is given by*

$$\langle M, M \rangle(t) = \int_0^t \lambda(X(s))v(X(s))ds. \quad (9.33)$$

PROOF. The formula (9.32) follows from (9.26). Indeed, using the exponential form of F_n (9.29), we have

$$A(t) = \sum_{i=0}^{\infty} m(X(T_i))\lambda(X(T_i))(t \wedge T_{i+1} - t \wedge T_i). \quad (9.34)$$

Note that since the process X has a constant value $X(T_i)$ on the time interval (T_i, T_{i+1}) , we have for a function f

$$\int_0^t f(X(s))ds = \sum_{i=0}^{\infty} f(X(T_i))(t \wedge T_{i+1} - t \wedge T_i), \quad (9.35)$$

and taking $f(x) = \lambda(x)m(x)$ gives the result. The formula (9.33) follows from Theorem 9.15. \square

More generally, using Theorems 9.14 and 9.15 in a similar way we have the compensator of the process $f(X(t))$ and quadratic variation of the compensated process.

Theorem 9.17:

- Denote $m_f(x) = E(f(x + \xi(x)) - f(x))$, then the compensator of $f(X(t))$ is given by

$$A^f(t) = \int_0^t \lambda(X(s))m_f(X(s))ds. \quad (9.36)$$

- Suppose the second moments of the jumps of $f(X)$ are finite, $v_f(x) = E(f(x + \xi(x)) - f(x))^2 < \infty$. Then the sharp bracket of the local martingale $M^f(t) = f(X(t)) - A^f(t)$ is given by

$$\langle M, M \rangle(t) = \int_0^t \lambda(X(s))v_f(X(s))ds. \quad (9.37)$$

PROOF. The proof of these formulae follows from Theorems 9.14 and 9.16. The process $f(X(t))$ is a pure jump process with the same arrival times T_n , and jumps of size $Z'_n = f(X(T_n)) - f(X(T_n-))$. The jumps are bounded since f is bounded. The mean of the jump from x is given by $E(Z'_{n+1} | \mathcal{F}_{T_n}, X(T_n) = x) = E(f(x + \xi(x)) - f(x)) = m_f(x)$. Therefore the compensator of $f(X(t))$ is, from Theorem 9.16, $\int_0^t \lambda(X(s))m_f(X(s))ds$. This implies that M^f is a local martingale. Expression (9.37) follows from Theorem 9.15. \square

9.6 Stochastic Equation for Jump Processes

It is sometimes useful to have an equation for the process itself, especially when dealing with approximations. It is clear from Theorem 9.16 that a Markov jump process X has a semimartingale representation

$$X(t) = X(0) + A(t) + M(t) = X(0) + \int_0^t \lambda(X(s))m(X(s))ds + M(t). \quad (9.38)$$

This is the integral form of the stochastic differential equation for X

$$dX(t) = \lambda(X(t))m(X(t))dt + dM(t), \quad (9.39)$$

and is driven by the purely discontinuous, finite variation martingale M . By analogy with diffusions the infinitesimal mean is $\lambda(x)m(x)$ and the infinitesimal variance is $\lambda(x)v(x)$, where $m(x)$ and $v(x)$ are the first and the second moments of the jump size from x respectively.

It is useful to have conditions assuring that the local martingale M in the representation (9.38) is a martingale. Such conditions are given in the next result.

Theorem 9.18: *Suppose that for all x*

$$\lambda(x)\mathbb{E}(|\xi(x)|) \leq C(1 + |x|), \quad (9.40)$$

then representation (9.38) holds with a zero mean martingale M . If in addition

$$\lambda(x)v(x) \leq C(1 + x^2), \quad (9.41)$$

then M is square integrable with

$$\langle M, M \rangle(t) = \int_0^t \lambda(X(s))v(X(s))ds. \quad (9.42)$$

In particular, we have the following corollary:

Corollary 9.19: *Suppose that the conditions of Theorem 9.18 hold. Then*

$$\mathbb{E}X(t) = \mathbb{E}X(0) + \mathbb{E} \int_0^t \lambda(X(s))m(X(s))ds, \quad (9.43)$$

and

$$\mathbb{E}X^2(t) = \mathbb{E}X^2(0) + \mathbb{E} \int_0^t \lambda(X(s)) \left(v(X(s)) + 2X(s)m(X(s)) \right) ds. \quad (9.44)$$

The proof of the result can be found in Hamza and Klebaner (1995b).

An application of the stochastic equation approach to the model of birth-death processes is given in Chapter 13.

Remark 9.2: Markov jump processes with countably or finitely many states are called Markov chains. The jump variables have discrete distributions in a Markov chain, whereas they can have a continuous distribution in a Markov process. Markov jump processes are also known as Markov chains with general state space.

Remark 9.3: A model for a randomly evolving population can be served by a Markov chain on non-negative integers. In this case the states of the process are the possible values for the population size. The traditional way of defining a Markov chain is by the infinitesimal probabilities: for integer i and j and small δ

$$P(X(t + \delta) = j | X(t) = i) = \lambda_{ij}\delta + o(\delta),$$

where $\lim_{\delta \rightarrow 0} o(\delta)/\delta = 0$. In this section we presented an almost sure, path by path representation of a Markov jump process. Another representation related to the Poisson process can be found in Ethier and Kurtz (1986).

9.7 Generators and Dynkin's Formula

Let X be the Markov jump process described by (9.28), i.e. it stays in a state x for a random time distributed exponentially with parameter $\lambda(x)$ and then jumps to $x + \xi(x)$, where $\xi(x)$ is the random variable of the jump from x . Define the following linear operator L acting on bounded functions by

$$Lf(x) = \lambda(x)E(f(x + \xi(x)) - f(x)) = \lambda(x)m_f(x). \quad (9.45)$$

We recognize the compensator of $f(X(t))$ as $A^f(t)$ in Theorem 9.17. The results can be re-stated in terms of operators.

Theorem 9.20: Let L be as above then $M^f(t)$

$$M^f(t) = f(X(t)) - f(X(0)) - \int_0^t Lf(X(s))ds \quad (9.46)$$

is a local martingale. Moreover,

$$\langle M^f, M^f \rangle(t) = \int_0^t \Gamma f(X(s))ds, \quad (9.47)$$

where the operator Γ is given by

$$\Gamma f(x) = Lf^2(x) - 2f(x)Lf(x). \quad (9.48)$$

PROOF. Only the second statement is new. Expanding (9.37) gives (9.47). Alternatively, calculating Lf^2 directly $f^2(X(t)) - \int_0^t Lf^2(X(s))ds = f^2(X(t)) - \int_0^t \Gamma f(X(s))ds - 2 \int_0^t f(X(s))Lf(X(s))ds$ is a martingale. On the other hand, since the difference $f(X) - M^f$ is continuous and of finite variation, $[M^f, M^f] = [f(X), f(X)]$. According to Theorem 8.6 $[f(X), f(X)](t) = f^2(X(t)) - 2 \int_0^t f(X(s-))df(X(s)) = f^2(X(t)) - 2 \int_0^t f(X(s-))dM^f(s) - 2 \int_0^t f(X(s-))Lf(X(s))ds$. Taking the difference of the two expressions we obtain that $[M^f, M^f](t) - \int_0^t \Gamma f(X(s))ds$ is a martingale. Since $\int_0^t \Gamma f(X(s))ds$ is continuous (9.47) is proved. \square

In the case when the local martingale M^f above (9.46) is a true martingale, one can take expectations, and Dynkin's formula is obtained.

Theorem 9.21 (Dynkin's Formula): *Suppose that $\lambda(x)$ is bounded, that is, $\sup_x \lambda(x) < \infty$. Then M^f is a martingale, and consequently*

$$\mathbb{E}f(X(t)) = f(X(0)) + \mathbb{E} \int_0^t Lf(X(s))ds. \quad (9.49)$$

The same holds when t is replaced by a stopping time τ .

PROOF. If $\lambda(x)$ is bounded, then since f is also bounded, it follows that $Lf(x)$ is bounded by a constant. Hence M^f is bounded by a constant Ct on any finite time interval $[0, t]$. Since a bounded local martingale is a martingale, Corollary 7.22, M^f is a martingale. Moreover, it is square integrable on any finite time interval. Since the optional stopping theorem applies, the result holds for stopping times. \square

The local martingale M^f is a true martingale under less restrictive conditions than bounded functions. For example, Theorem 9.18 generalizes Dynkin's formula for unbounded λ and f , such as $f(x) = x$ and $f(x) = x^2$. These are particular cases of a more general theorem that extends Dynkin's formula to unbounded functions.

Theorem 9.22: *Assume that there are no explosions, that is, $T_n \uparrow \infty$. Suppose that for an unbounded function f there is a constant C such that for all x*

$$\lambda(x)\mathbb{E}|f(x + \xi(x)) - f(x)| \leq C(1 + |f(x)|). \quad (9.50)$$

Suppose also that $\mathbb{E}|f(X(0))| < \infty$. Then for all t , $0 \leq t < \infty$, $\mathbb{E}|f(X(t))| < \infty$, moreover, M^f in (9.46) is a martingale and (9.49) holds.

For the proof see Hamza and Klebaner (1995b).

The linear operator L in (9.45) is called the generator of X . It can be shown that with its help one can define probabilities on the space of right-continuous functions so that the coordinate process is the Markov process (compare with the solution to the martingale problem for diffusions).

9.8 Explosions in Markov Jump Processes

Let $X(t)$ be a Markov jump process taking values in \mathbb{R} . If the process is in states x , then it stays there for an exponential length of time with mean $\lambda^{-1}(x)$ after which it jumps from x . If $\lambda(x) \rightarrow \infty$ for a set of values x , then the expected duration of stay in state x , $\lambda^{-1}(x) \rightarrow 0$, and the time spent in x becomes shorter and shorter. It can happen that the process jumps infinitely many times in a finite time interval, that is, $\lim_{n \rightarrow \infty} T_n = T_\infty < \infty$. This phenomenon is called explosion. The terminology comes from the case when the process takes only integer values and $\lambda(x)$ can tend to infinity only when $x \rightarrow \infty$. In this case there are infinitely many jumps only if the process reaches infinity in finite time.

When the process does not explode it is called *regular*. In other words, the regularity assumption is

$$T_\infty = \lim_{n \rightarrow \infty} T_n = \infty \text{ almost surely.} \quad (9.51)$$

If $P(T_\infty < \infty) > 0$, then it is said that the process explodes on the set $\{T_\infty < \infty\}$.

Theorem 9.23: *A necessary and sufficient condition for non-explosion is given by*

$$\sum_0^\infty \frac{1}{\lambda(X(T_n))} = \infty \text{ almost surely.} \quad (9.52)$$

PROOF. We sketch the proof (see Breiman (1968) for details). Let ν_n be a sequence of independent exponentially distributed random variables with parameter λ_n . Then it is easy to see (by taking a Laplace transform) that $\sum_0^n \nu_i < \infty$ converges in distribution if, and only if, $\sum_0^\infty \frac{1}{\lambda_n} < \infty$. Using the result that a series of *independent* random variables converges almost surely if, and only if, it converges in distribution, we have $\sum_0^\infty \nu_n < \infty$ almost surely if, and only if, $\sum_0^\infty \frac{1}{\lambda_n} < \infty$. Condition (9.52) now follows since the conditional distribution of $T_{n+1} - T_n$, given the information up to the n -th jump, \mathcal{F}_{T_n} , is exponential with parameter $\lambda(X(T_n))$. \square

Clearly, if $\lambda(x)$ is bounded, then (9.52) holds. However, condition (9.52) is hard to check in general since it involves the variables $X(T_n)$. A simple condition, which is essentially in terms of the function $\lambda(x)m(x)$ (the drift), follows.

Theorem 9.24: *Assume that $X \geq 0$, and there exists a monotone function $f(x)$ such that $\lambda(x)m(x) \leq f(x)$ and $\int_0^\infty \frac{dx}{f(x)} = \infty$. Then the process X does not explode, that is, $P(X(t) < \infty \text{ for all } t, 0 \leq t < \infty) = 1$.*

According to this, the first condition of the result on integrability, Theorem 9.18, guarantees that the process does not explode. Proof of Theorem 9.24 and other sharp sufficient conditions for regularity in terms of the parameters of the process are given in Kersting and Klebaner (1995), (1996).

Notes: Material for this chapter is based on Liptser and Shiryaev (1974), (1989), Karr (1986), and the research papers quoted in the text.

9.9 Exercises

Exercise 9.1: Let $U \geq 0$ and denote $h(x) = \int_0^x \frac{dF(s)}{1-F(s)}$, where $F(x)$ is the distribution function of U . Show that for $a > 0$, $Eh(U \wedge a) = \int_0^a dF(x)$.

Exercise 9.2: Show that when the distribution of inter-arrival times is exponential the formula (9.19) gives the compensator λt , hence $N(t)$ is a Poisson process.

Exercise 9.3: Show that when the distribution of inter-arrival times is geometric then $N(t)$ has a binomial distribution.

Exercise 9.4: Prove the formula for the compensator in (9.26).

Exercise 9.5: Let $X(t)$ be a pure jump process with first k moments. Let $m_k(x) = E\xi^k(x)$ and assume for all $i = 1, \dots, k$, $\lambda(x)E|\xi(x)|^i \leq C|x|^i$. Show that

$$EX^k(t) = EX^k(0) + \sum_{i=0}^{k-1} \binom{k}{i} \int_0^t E(\lambda(X(s))X^i(s)m_{k-i}(X(s)))ds.$$

Chapter 10

Change of Probability Measure

In this chapter we describe what happens to random variables and processes when the original probability measure is changed to an equivalent one. Change of measure for processes is done by using Girsanov's theorem.

10.1 Change of Measure for Random Variables

Change of Measure on a Discrete Probability Space

We start with a simple example. Let $\Omega = \{\omega_1, \omega_2\}$ with probability measure P given by $P(\omega_1) = p$, $P(\omega_2) = 1 - p$.

Definition 10.1: Q is equivalent to P ($Q \sim P$) if they have the same null sets, i.e. $Q(A) = 0$ if, and only if, $P(A) = 0$.

Let Q be a new probability measure equivalent to P . In this example this means that $Q(\omega_1) > 0$ and $Q(\omega_2) > 0$ (or $0 < Q(\omega_1) < 1$). Put $Q(\omega_1) = q$, $0 < q < 1$.

Now let $\Lambda(\omega) = \frac{Q(\omega)}{P(\omega)}$, that is, $\Lambda(\omega_1) = \frac{Q(\omega_1)}{P(\omega_1)} = \frac{q}{p}$, and $\Lambda(\omega_2) = \frac{1-q}{1-p}$.

By definition of Λ , for all ω

$$Q(\omega) = \Lambda(\omega)P(\omega). \quad (10.1)$$

Now let X be a random variable. The expectation of X under the probability P is given by

$$\mathbb{E}_P(X) = X(\omega_1)P(\omega_1) + X(\omega_2)P(\omega_2) = pX(\omega_1) + (1 - p)X(\omega_2)$$

and under the probability Q

$$E_Q(X) = X(\omega_1)Q(\omega_1) + X(\omega_2)Q(\omega_2).$$

From (10.1)

$$E_Q(X) = X(\omega_1)\Lambda(\omega_1)P(\omega_1) + X(\omega_2)\Lambda(\omega_2)P(\omega_2) = E_P(\Lambda X). \quad (10.2)$$

Take $X = 1$, then

$$E_Q(X) = 1 = E_P(\Lambda). \quad (10.3)$$

On the other hand, take any random variable $\Lambda > 0$, such that $E_P(\Lambda) = 1$, and define Q by (10.1).

Then Q is a probability because $Q(\omega_i) = \Lambda(\omega_i)P(\omega_i) > 0$, $i = 1, 2$, and

$$\begin{aligned} Q(\Omega) &= Q(\omega_1) + Q(\omega_2) = \Lambda(\omega_1)P(\omega_1) + \Lambda(\omega_2)P(\omega_2) \\ &= E_P(\Lambda) = 1. \end{aligned}$$

Q is equivalent to P since Λ is strictly positive.

Thus we have shown that for any equivalent change of measure $Q \sim P$ there is a positive random variable Λ , such that $E_P(\Lambda) = 1$, and $Q(\omega) = \Lambda(\omega)P(\omega)$. This is the simplest version of the general result, the Radon–Nikodym theorem. The expectation under Q of a random variable X is given by,

$$E_Q(X) = E_P(\Lambda X). \quad (10.4)$$

By taking indicators $I(X \in A)$ we obtain the distribution of X under Q

$$Q(X \in A) = E_P(\Lambda I(X \in A)).$$

These formulae, obtained here for a simple example, hold also in general.

Change of Measure for Normal Random Variables

Consider first a change of a normal probability measure on \mathbb{R} . Let μ be any real number, $f_\mu(x)$ denote the probability density of $N(\mu, 1)$ distribution, and P_μ the $N(\mu, 1)$ probability measure on \mathbb{R} ($\mathcal{B}(\mathbb{R})$). Then it is easy to see that

$$f_\mu(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(x-\mu)^2} = f_0(x)e^{\mu x - \mu^2/2}. \quad (10.5)$$

Put

$$\Lambda(x) = e^{\mu x - \mu^2/2}. \quad (10.6)$$

Then the above equation reads

$$f_\mu(x) = f_0(x)\Lambda(x). \quad (10.7)$$

According to the definition of a density function the probability of a set A on the line is the integral of the density over this set,

$$P(A) = \int_A f(x)dx = \int_A dP. \quad (10.8)$$

In infinitesimal notations this relation is written as

$$dP = P(dx) = f(x)dx. \quad (10.9)$$

Hence the relation between the densities (10.7) can be written as a relation between the corresponding probability measures

$$f_\mu(x)dx = f_0(x)\Lambda(x)dx, \quad \text{and} \quad P_\mu(dx) = \Lambda(x)P_0(dx). \quad (10.10)$$

According to a property of the expectation (integral) (Chapter 2.3), if a random variable $X \geq 0$ then $EX = 0$ if, and only if, $P(X = 0) = 1$,

$$P_\mu(A) = \int_A \Lambda(x)P_0(dx) = E_{P_0}(I_A\Lambda) = 0$$

implies

$$P_0(I_A\Lambda = 0) = 1.$$

Since $\Lambda(x) > 0$ for all x this implies $P_0(A) = 0$. Using that $\Lambda < \infty$, the other direction follows: if $P_0(A) = 0$, then $P_\mu(A) = E_{P_0}(I_A\Lambda) = 0$, which proves that these measures have the same null sets and are equivalent.

Another notation for Λ is

$$\Lambda = \frac{dP_\mu}{dP_0}, \quad \frac{dP_\mu}{dP_0}(x) = e^{\mu x - \mu^2/2}. \quad (10.11)$$

This shows that any $N(\mu, 1)$ probability is obtained by an equivalent change of probability measure from the $N(0, 1)$ distribution.

We now give the same result in terms of changing the distribution of a random variable.

Theorem 10.2: *Let X have $N(0, 1)$ under P , and $\Lambda(X) = e^{\mu X - \mu^2/2}$. Define measure Q by*

$$Q(A) = \int_A \Lambda(X)dP = E_P(I_A\Lambda(X)) \quad \text{or} \quad \frac{dQ}{dP}(X) = \Lambda(X). \quad (10.12)$$

Then Q is an equivalent probability measure, and X has $N(\mu, 1)$ distribution under Q .

PROOF. We show first that Q defined by (10.12) is indeed a probability. $Q(\Omega) = E_Q(1) = E_P(1\Lambda(X)) = E_P(e^{\mu X - \mu^2/2}) = e^{-\mu^2/2}E_P(e^{\mu X}) = 1$. The last equality is because $E_P(e^{\mu X}) = e^{\mu^2/2}$ for the $N(0, 1)$ random variable. Other properties of probability Q follow from the corresponding properties of the integral. Consider the moment generating function of X under Q . $E_Q(e^{uX}) = E_P(e^{uX}\Lambda(X)) = E_P(e^{(u+\mu)X - \mu^2/2}) = e^{-\mu^2/2}E_P(e^{(u+\mu)X}) = e^{u\mu + u^2/2}$, which corresponds to the $N(\mu, 1)$ distribution. \square

Remark 10.1: If P is a probability such that X is $N(0, 1)$, then $X + \mu$ has $N(\mu, 1)$ distribution under the same P . This is an operation on the outcomes x . When we change the probability measure P to $Q = P_\mu$ we leave the outcomes as they are but assign different probabilities to them (more precisely to sets of outcomes). Under the new measure the same X has $N(\mu, 1)$ distribution.

Example 10.1: (Simulations of rare events)

Change of probability measure is useful in simulation and estimation of probabilities of rare events. Consider estimation of $P(N(6, 1) < 0)$ by simulations. This probability is about 10^{-10} . Direct simulation is done by the expression

$$P(N(6, 1) < 0) \approx \frac{1}{n} \sum_{i=1}^n I(x_i < 0), \quad (10.13)$$

where x_i 's are the observed values from the $N(6, 1)$ distribution. Note that in a million runs, $n = 10^6$, we should expect no values below zero, and the estimate is zero.

Let P be the $N(6, 1)$ distribution on \mathbb{R} . Consider P as changed from $N(0, 1) = Q$, as follows

$$\frac{dP}{dQ}(x) = \frac{dN(6, 1)}{dN(0, 1)}(x) = \Lambda(x) = e^{\mu x - \mu^2/2} = e^{6x - 18}.$$

So we have

$$P(A) = N(6, 1)(A) = E_Q(\Lambda(x)I(A)) = E_{N(0, 1)}(e^{6x - 18}I(A)).$$

In our case $A = (-\infty, 0]$.

Thus we are led to the following estimate:

$$P(N(6, 1) < 0) = P(A) \approx \frac{1}{n} \sum_{i=1}^n e^{6x_i - 18} I(x_i < 0) = e^{-12} \frac{1}{n} \sum_{i=1}^n e^{6(x_i - 1)} I(x_i < 0), \quad (10.14)$$

with x_i generated from $N(0, 1)$ distribution. Of course, about half of the observations x_i 's will be negative, resulting in a more precise estimate, even for a small number of runs n .

Next we give the result for random variables, set up in a way similar to GirSANOV's theorem for processes, changing the measure to remove the mean.

Theorem 10.3 (Removal of the Mean): Let X have $N(0, 1)$ distribution under P , and $Y = X + \mu$. Then there is an equivalent probability $Q \sim P$, such that Y has $N(0, 1)$ under Q . $(dQ/dP)(X) = \Lambda(X) = e^{-\mu X - \mu^2/2}$.

PROOF. Similarly to the previous proof, Q is a probability measure, with the same null sets as P .

$$\begin{aligned} E_Q(e^{uY}) &= E_P(e^{u(X+\mu)} \Lambda(X)) \\ &= E_P(e^{(u-\mu)X+u\mu-\frac{1}{2}\mu^2}) = e^{u\mu-\frac{1}{2}\mu^2} E_P(e^{(u-\mu)X}). \end{aligned}$$

Using the moment generating function of $N(0, 1)$, $E_P(e^{(u-\mu)X}) = e^{(u-\mu)^2/2}$, which gives $E_Q(e^{uY}) = e^{u^2/2}$, establishing $Y \sim N(0, 1)$ under Q . \square

Finally, we give the change of one normal probability to another.

Theorem 10.4: Let X have $N(\mu_1, \sigma_1^2)$ distribution, call it P . Define Q by

$$(dQ/dP)(X) = \Lambda(X) = \frac{\sigma_1}{\sigma_2} e^{\frac{(X-\mu_1)^2}{2\sigma_1^2} - \frac{(X-\mu_2)^2}{2\sigma_2^2}}. \quad (10.15)$$

Then X has $N(\mu_2, \sigma_2^2)$ distribution under Q .

PROOF. The form of Λ is easily verified as the ratio of normal densities. This proves the statement for P and Q on \mathbb{R} . On a general space the proof is similar to the above by working out the moment generating function of X under Q . \square

10.2 Change of Measure on a General Space

Let two probability measures P and Q be defined on the same space.

Definition 10.5: Q is called absolutely continuous with respect to P , written as $Q \ll P$, if $Q(A) = 0$ whenever $P(A) = 0$. P and Q are called equivalent if $Q \ll P$ and $P \ll Q$.

Theorem 10.6 (Radon–Nikodym): If $Q \ll P$, then there exists a random variable Λ such that $\Lambda \geq 0$, $E_P \Lambda = 1$, and

$$Q(A) = E_P(\Lambda I(A)) = \int_A \Lambda dP \quad (10.16)$$

for any measurable set A . Λ is P -almost surely unique. Conversely, if there exists a random variable Λ with the above properties and Q is defined by (10.16), then it is a probability measure and $Q \ll P$.

The random variable Λ in Theorem 10.6 is called the Radon–Nikodym derivative or the density of Q with respect to P , and is denoted by dQ/dP . It follows from (10.16) that if $Q \ll P$, then expectations under P and Q are related by

$$E_Q X = E_P(\Lambda X), \quad (10.17)$$

for any random variable X integrable with respect to Q .

Calculations of expectations are sometimes made easier by using a change of measure.

Example 10.2: (A lognormal calculation for a financial option)

$E(e^X I(X > a))$, where X has $N(\mu, 1)$ distribution (P). Take $\Lambda(X) = e^X / Ee^X = e^{X-\mu-\frac{1}{2}}$, and $dQ = \Lambda(X)dP$. Then $E\Lambda(X) = 1$, $0 < \Lambda(X) < \infty$, so $Q \sim P$.

$$\begin{aligned} E_P(e^X I(X > a)) &= e^{\mu+\frac{1}{2}} E_P(\Lambda(X)I(X > a)) \\ &= e^{\mu+\frac{1}{2}} E_Q(I(X > a)) = e^{\mu+\frac{1}{2}} Q(X > a). \\ E_Q(e^{uX}) &= E_P(e^{uX} \Lambda(X)) = E_P(e^{(u+1)X-\mu-\frac{1}{2}}) = e^{u\mu+u+u^2/2}, \end{aligned}$$

which is the transform of $N(\mu + 1, 1)$ distribution. Thus the Q distribution of X is $N(\mu + 1, 1)$, and $E(e^X I(X > a)) = e^{\mu+\frac{1}{2}} Q(X > a) = e^{\mu+\frac{1}{2}} \Pr(N(\mu + 1, 1) > a) = e^{\mu+\frac{1}{2}} (1 - \Phi)(a - \mu - 1)$.

We give the definition of the “opposite” concept to the absolute continuity of two measures.

Definition 10.7: Two probability measures P and Q defined on the same space are called singular (P_Q) if there exist a set A such that $P(A) = 0$ and $Q(A) = 1$.

Singularity means that by observing an outcome we can decide with certainty on the probability model.

Example 10.3:

1. Let $\Omega = \mathbb{R}^+$, P is the exponential distribution with parameter 1, and Q is the Poisson distribution with parameter 1. Then P and Q are singular. Indeed the set of non-negative integers has Q probability 1, and P probability 0.
2. We shall see later that the probability measures induced by processes $\sigma B(t)$, where B is a Brownian motion, are singular for different values of σ .

Expectations for an absolutely continuous change of probability measure are related by the formula (10.17). Conditional expectations are related by an equation, similar to Bayes formula.

Theorem 10.8 (General Bayes Formula): *Let \mathcal{G} be a sub- σ -field of \mathcal{F} on which two probability measures Q and P are defined. If $Q \ll P$ with $dQ = \Lambda dP$ and X is Q -integrable, then ΛX is P -integrable and Q -almost surely*

$$E_Q(X|\mathcal{G}) = \frac{E_P(X\Lambda|\mathcal{G})}{E_P(\Lambda|\mathcal{G})}. \quad (10.18)$$

PROOF. We check the definition of conditional expectation given \mathcal{G} , (2.15) for any bounded \mathcal{G} -measurable random variable ξ

$$\mathbb{E}_Q(\xi X) = \mathbb{E}_Q(\xi \mathbb{E}_Q(X|\mathcal{G})).$$

Clearly the rhs of (10.18) is \mathcal{G} -measurable.

$$\begin{aligned}\mathbb{E}_Q\left(\frac{\mathbb{E}_P(X\Lambda|\mathcal{G})}{\mathbb{E}_P(\Lambda|\mathcal{G})}\xi\right) &= \mathbb{E}_P\left(\Lambda\frac{\mathbb{E}_P(X\Lambda|\mathcal{G})}{\mathbb{E}_P(\Lambda|\mathcal{G})}\xi\right) \\ &= \mathbb{E}_P\left(\mathbb{E}_P(\Lambda|\mathcal{G})\frac{\mathbb{E}_P(X\Lambda|\mathcal{G})}{\mathbb{E}_P(\Lambda|\mathcal{G})}\xi\right) \\ &= \mathbb{E}_P(\mathbb{E}_P(X\Lambda|\mathcal{G})\xi) \\ &= \mathbb{E}_P(X\Lambda\xi) = \mathbb{E}_Q(X\xi).\end{aligned}$$

□

In the rest of this section we address the general case of two measures not necessarily equivalent. Let P and Q be two measures on the same probability space. Consider the measure $\nu = \frac{P+Q}{2}$. Then, clearly, $P \ll \nu$ and $Q \ll \nu$. Therefore, by the Radon–Nikodym theorem, there exist Λ_1 and Λ_2 such that $dP/d\nu = \Lambda_1$ and $dQ/d\nu = \Lambda_2$. Introduce the notation $x^{(+)} = 1/x$, if $x \neq 0$, and 0 otherwise, so that $x^{(+)}x = I(x \neq 0)$. We have

$$\begin{aligned}Q(A) &= \int_A \Lambda_2 d\nu = \int_A \Lambda_2 (\Lambda_1^{(+)} \Lambda_1 + 1 - \Lambda_1^{(+)} \Lambda_1) d\nu \\ &= \int_A \Lambda_2 \Lambda_1^{(+)} \Lambda_1 d\nu + \int_A (1 - \Lambda_1^{(+)} \Lambda_1) \Lambda_2 d\nu \\ &= \int_A \Lambda_2 \Lambda_1^{(+)} dP + \int_A (1 - \Lambda_1^{(+)} \Lambda_1) dQ \\ &:= Q^c(A) + Q^s(A),\end{aligned}$$

where $Q^c(A) = \int_A \Lambda_2 \Lambda_1^{(+)} dP$ is absolutely continuous with respect to P , and $Q^s(A) = \int_A (1 - \Lambda_1^{(+)} \Lambda_1) dQ$. The set $A = \{\Lambda_1 > 0\}$ has P probability one because the integral over A and Ω are the same, and $\mathbb{E}_\nu \Lambda_1 = 1$

$$P(A) = \int_A \Lambda_1 d\nu = \int_\Omega \Lambda_1 d\nu = P(\Omega) = 1,$$

but has measure zero under Q^s because on this set $\Lambda_1^{(+)} \Lambda_1 = 1$ and $Q^s(A) = 0$. This shows that Q^s is singular with respect to P . Thus we have

Theorem 10.9 (Lebesgue Decomposition): *Let P and Q be two measures on the same probability space. Then $Q(A) = Q^c(A) + Q^s(A)$, where $Q^c \ll P$ and $Q^s \perp P$.*

10.3 Change of Measure for Processes

Since realizations of a Brownian motion are continuous functions, the probability space is taken to be the set of continuous functions on $[0, T]$, $\Omega = C([0, T])$. Because we have to describe the collection of sets to which the probability is assigned, the concepts of an open (closed) set is needed. These are defined in the usual way with the help of the distance between two functions taken as $\sup_{t \leq T} |w_1(t) - w_2(t)|$. The σ -field is the one generated by open sets. Probability measures are defined on the measurable subsets of Ω . If $\omega = w_{[0, T]}$ denotes a continuous function, then there is a probability measure P on this space such that the “coordinate” process $B(t, \omega) = B(t, w_{[0, T]}) = w(t)$ is a Brownian motion, P is the Wiener measure, see Chapter 5.7. Note that although a Brownian motion can be defined on $[0, \infty]$, the equivalent change of measure is defined only on finite intervals $[0, T]$. A random variable on this space is a function $X : \Omega \rightarrow \mathbb{R}$, and $X(\omega) = X(w_{[0, T]})$, also known as a functional of Brownian motion. Since probabilities can be obtained as expectation of indicators, $P(A) = E I_A$, it is important to know how to calculate expectations. $E(X)$ is given as an integral with respect to the Wiener measure

$$E(X) = \int_{\Omega} X(w_{[0, T]}) dP. \quad (10.19)$$

In particular, if $X(w_{[0, T]}) = h(w(T))$, for some function of real argument h , then

$$E(X) = \int_{\Omega} h(w(T)) dP = E(h(B(T))), \quad (10.20)$$

which can be evaluated by using the $N(0, T)$ distribution of $B(T)$. Similarly, if X is a function of finitely many values of w , $X(w_{[0, T]}) = h(w(t_1), \dots, w(t_n))$, then the expectation can be calculated by using the multivariate normal distribution of $B(t_1), \dots, B(t_n)$. However, for a functional which depends on the whole path $w_{[0, T]}$, such as integrals of $w(t)$ or $\max_{t \leq T} w(t)$, the distribution of the functional is required, see (10.21). It can be calculated by using the distribution of X , F_X , as for any random variable

$$E(X) = \int_{\Omega} X(w_{[0, T]}) dP = \int_{\mathbb{R}} x dF_X(x). \quad (10.21)$$

If P and Q are equivalent, that is, they have the same null sets, then there exists a random variable Λ (the Radon–Nikodym derivative $\Lambda = dQ/dP$) such that the probabilities under Q are given by $Q(A) = \int_A \Lambda dP$. Girsanov’s theorem gives the form of Λ .

First we state a general result, that follows directly from Theorem 10.8, for the calculation of expectations and conditional expectations under an absolutely continuous change of measure. It is also known as the general Bayes formula.

Theorem 10.10: Let $\Lambda(t)$ be a positive P -martingale such that $E_P(\Lambda(T)) = 1$. Define the new probability measure Q by the relation $Q(A) = \int_A \Lambda(T)dP$ (so that $dQ/dP = \Lambda(T)$). Then Q is absolutely continuous with respect to P and for a Q -integrable random variable X

$$E_Q(X) = E_P(\Lambda(T)X). \quad (10.22)$$

$$E_Q(X|\mathcal{F}_t) = E_P\left(\frac{\Lambda(T)}{\Lambda(t)}X|\mathcal{F}_t\right), \quad (10.23)$$

and if X is \mathcal{F}_t -measurable, then for $s \leq t$

$$E_Q(X|\mathcal{F}_s) = E_P\left(\frac{\Lambda(t)}{\Lambda(s)}X|\mathcal{F}_s\right). \quad (10.24)$$

PROOF. It remains to show (10.24). $E_Q(X|\mathcal{F}_s) = E_P\left(\frac{\Lambda(T)}{\Lambda(s)}X|\mathcal{F}_s\right) = E_P\left(E_P\left(\frac{\Lambda(T)}{\Lambda(s)}X|\mathcal{F}_t\right)|\mathcal{F}_s\right) = E_P\left(X\frac{1}{\Lambda(s)}E_P(\Lambda(T)|\mathcal{F}_t)|\mathcal{F}_s\right) = E_P\left(\frac{\Lambda(t)}{\Lambda(s)}X|\mathcal{F}_s\right)$. \square

The following result follows immediately from (10.24):

Corollary 10.11: A process $M(t)$ is a Q -martingale if, and only if, $\Lambda(t)M(t)$ is a P -martingale.

By taking $M(t) = 1/\Lambda(t)$ we obtain a result that is used in financial applications.

Theorem 10.12: Let $\Lambda(t)$ be a positive P -martingale such that $E_P(\Lambda(T)) = 1$, and $dQ/dP = \Lambda(T)$. Then $1/\Lambda(t)$ is a Q -martingale.

We show next that convergence in probability is preserved under an absolutely continuous change of measure.

Theorem 10.13: Let $X_n \rightarrow X$ in probability P and $Q \ll P$. Then $X_n \rightarrow X$ in probability Q .

PROOF. Denote $A_n = \{|X_n - X| > \varepsilon\}$. Then convergence in probability of X_n to X means $P(A_n) \rightarrow 0$. However, $Q(A_n) = E_P(\Lambda I_{A_n})$. Since Λ is P -integrable the result follows by dominated convergence. \square

Corollary 10.14: The quadratic variation of a process does not change under an absolutely continuous change of the probability measure.

PROOF. The sums $\sum_{i=0}^n (X(t_{i+1}) - X(t_i))^2$ approximating the quadratic variation converge in P probability to $[X, X](t)$. According to the above result they converge to the same limit under an equivalent to P probability Q . \square

Theorem 10.15 (Girsanov's Theorem for Brownian Motion):

Let $B(t)$, $0 \leq t \leq T$, be a Brownian motion under probability measure P . Consider the process $W(t) = B(t) + \mu t$. Define the measure Q by

$$\Lambda = \frac{dQ}{dP}(B_{[0,T]}) = e^{-\mu B(T) - \frac{1}{2}\mu^2 T}, \quad (10.25)$$

where $B_{[0,T]}$ denotes a path of Brownian motion on $[0, T]$. Then Q is equivalent to P , and $W(t)$ is a Q -Brownian motion.

$$\frac{dP}{dQ}(W_{[0,T]}) = \frac{1}{\Lambda} = e^{\mu W(T) - \frac{1}{2}\mu^2 T}. \quad (10.26)$$

PROOF. The proof uses Levy's characterization of Brownian motion, as a continuous martingale with quadratic variation process t . Quadratic variation is the same under P and Q by Theorem 10.13. Therefore (with a slight abuse of notation) using the fact that μt is smooth and does not contribute to the quadratic variation,

$$[W, W](t) = [B(t) + \mu t, B(t) + \mu t] = [B, B](t) = t.$$

It remains to establish that $W(t)$ is a Q -martingale. Let $\Lambda(t) = E_P(\Lambda | \mathcal{F}_t)$. According to Corollary 10.11 it is enough to show that $\Lambda(t)W(t)$ is a P -martingale. This is done by direct calculations.

$$E_P(W(t)\Lambda(t) | \mathcal{F}_s) = E_P\left((B(t) + \mu t)e^{-\mu B(t) - \frac{1}{2}\mu^2 t} | \mathcal{F}_s\right) = W(s)\Lambda(s). \quad \square$$

It turns out that a drift of the form $\int_0^t H(s)ds$ with $\int_0^T H^2(s)ds < \infty$ can be removed similarly by a change of measure.

Theorem 10.16 (Girsanov's Theorem for Removal of Drift): Let $B(t)$ be a P -Brownian motion, and $H(t)$ is such that $X(t) = -\int_0^t H(s)dB(s)$ is defined, moreover, $\mathcal{E}(X)$ is a martingale. Define an equivalent measure Q by

$$\Lambda = \frac{dQ}{dP}(B) = e^{-\int_0^T H(s)dB(s) - \frac{1}{2}\int_0^T H^2(s)ds} = \mathcal{E}(X)(T). \quad (10.27)$$

Then the process

$$W(t) = B(t) + \int_0^t H(s)ds \quad \text{is a } Q\text{-Brownian motion.} \quad (10.28)$$

PROOF. The proof is similar to the previous one and we only sketch it. We show that $W(t) = B(t) + \int_0^t H(s)ds$ is a continuous Q -local martingale with quadratic variation t . The result then follows by Levy's characterization.

The quadratic variation of $W(t) = B(t) + \int_0^t H(s)ds$ is t , since the integral is continuous and is of finite variation, $[W, W]_Q(t) = [B, B]_Q(t) = t$ by Corollary 10.14. $W(t)$ is clearly continuous. In order to establish that $W(t)$ is a Q-local martingale, by Corollary 10.11, it is enough to show that $\Lambda(t)W(t) = \Lambda(t)(B(t) + \int_0^t H(s)ds)$ is a P-local martingale, with $\Lambda(t) = E_P(\Lambda | \mathcal{F}_t)$. To this end we calculate $d(\Lambda(t)W(t))$ and show that it has no dt term. Since $\Lambda(t)$ is a martingale exponential of martingale $X(t)$, it satisfies

$$\begin{aligned} d\Lambda(t) &= \Lambda(t)dX(t), \quad \text{with } dX_t = -H(t)dB(t). \\ d(\Lambda(t)W(t)) &= \Lambda(t)dW(t) + W(t)d\Lambda(t) + d\Lambda(t)dW(t). \end{aligned}$$

Now,

$$d\Lambda(t)dW(t) = \Lambda(t)dX(t)(dB(t) + H(t)dt) = -\Lambda(t)H(t)dt.$$

Proceeding we see that

$$d(\Lambda(t)W(t)) = \Lambda(t)dB(t) + W(t)d\Lambda(t),$$

implying that $\Lambda(t)W(t)$ is a P-local martingale and $W(t)$ is a Q-local martingale. \square

Girsanov's theorem holds also in n dimensions.

Theorem 10.17: *Let \mathbf{B} be a P n -dimensional Brownian motion and $\mathbf{W} = (W^1(t), \dots, W^n(t))$, where*

$$W^i(t) = B^i(t) + \int_0^t H^i(s)ds,$$

with $\mathbf{H}(t) = (H^1(t), H^2(t), \dots, H^n(t))$ a regular adapted process satisfying $\int_0^T |\mathbf{H}(s)|^2 ds < \infty$. Let

$$X(t) = -\mathbf{H} \cdot \mathbf{B} := -\sum_{i=1}^n \int_0^t H^i(s)dB^i(s),$$

and assume that $\mathcal{E}(X)_T$ is a martingale. Then there is an equivalent probability measure Q such that \mathbf{W} is a Q-Brownian motion. Q is determined by

$$\frac{dQ}{dP}(\mathbf{B}_{[0,T]}) = \Lambda(\mathbf{B}_{[0,T]}) = \mathcal{E}(X)(T).$$

Comment that a sufficient condition for $\mathcal{E}(X)$ to be a martingale is Theorem 8.17, $E(e^{\frac{1}{2} \int_0^T H^2(s)ds}) < \infty$, and for $\mathcal{E}(\mathbf{H} \cdot \mathbf{B})$ to be a martingale is $E(e^{\frac{1}{2} \int_0^T |\mathbf{H}(s)|^2 ds}) < \infty$.

The proof for n dimensions is similar to one dimension, using calculations from Exercise 10.5.

We give a version of Girsanov's theorem for martingales. The proof is similar to the one above and is not given.

Theorem 10.18: *Let $M_1(t)$, $0 \leq t \leq T$ be a continuous P-martingale. Let $X(t)$ be a continuous P-martingale such that $\mathcal{E}(X)$ is a martingale. Define a new probability measure Q by*

$$\frac{dQ}{dP} = \Lambda = \mathcal{E}(X)(T) = e^{X(T) - \frac{1}{2}[X, X](T)}. \quad (10.29)$$

Then

$$M_2(t) = M_1(t) - [M_1, X](t) \quad (10.30)$$

is a continuous martingale under Q.

A sufficient condition for $\mathcal{E}(X)$ to be a martingale is Novikov's condition $E(e^{\frac{1}{2}[X, X]_T}) < \infty$, or Kazamaki's condition (8.39).

Change of Drift in Diffusions

Let $X(t)$ be a diffusion so that with a P-Brownian motion $B(t)$, $X(t)$ satisfies the following stochastic differential equation with $\sigma(x, t) > 0$,

$$dX(t) = \mu_1(X(t), t)dt + \sigma(X(t), t)dB(t). \quad (10.31)$$

Let

$$H(t) = \frac{\mu_1(X(t), t) - \mu_2(X(t), t)}{\sigma(X(t), t)}, \quad (10.32)$$

and define Q by $dQ = \Lambda dP$ with

$$\Lambda = \frac{dQ}{dP} = \mathcal{E}\left(-\int_0^t H(s)dB(s)\right)(T) = e^{-\int_0^T H(s)dB(s) - \frac{1}{2}\int_0^T H^2(s)ds}. \quad (10.33)$$

According to Girsanov's theorem, provided the process $\mathcal{E}(H \cdot B)$ is a martingale, the process $W(t) = B(t) + \int_0^t H(s)ds$ is a Q-Brownian motion. However,

$$dW(t) = dB(t) + H(t)dt = dB(t) + \frac{\mu_1(X(t), t) - \mu_2(X(t), t)}{\sigma(X(t), t)}dt. \quad (10.34)$$

Rearranging, we obtain the equation for $X(t)$

$$dX(t) = \mu_2(X(t), t)dt + \sigma(X(t), t)dW(t), \quad (10.35)$$

with a Q-Brownian motion $W(t)$. Thus the change of measure for a Brownian motion given above results in the change of the drift in the SDE.

Example 10.4: (Maximum of arithmetic Brownian motion)

Let $W(t) = \mu t + B(t)$, where $B(t)$ is P -Brownian motion, and $W^*(t) = \max_{s \leq t} W(s)$. When $\mu = 0$ the distribution of the maximum as well as the joint distribution are known, Theorem 3.21. We find these distributions when $\mu \neq 0$.

Let $B(t)$ be a P -Brownian motion, then $W(t)$ is a Q -Brownian motion with $\Lambda = dP/dQ$ given by (10.26). Let $A = \{W(T) \in I_1, W^*(T) \in I_2\}$, where I_1, I_2 are intervals on the line. Then we obtain from (10.26)

$$P(A) = \int_A e^{\mu W(T) - \frac{1}{2}\mu^2 T} dQ = E_Q(e^{\mu W(T) - \frac{1}{2}\mu^2 T} I(A)). \quad (10.36)$$

Denote by $q_{W,W^*}(x, y)$ and $p_{W,W^*}(x, y)$ the joint density of $(W(T), W^*(T))$ under Q and P respectively. Then it follows from (10.36)

$$\int_{\{x \in I_1, y \in I_2\}} p_{W,W^*}(x, y) dx dy = \int_{\{x \in I_1, y \in I_2\}} q_{W,W^*}(x, y) e^{\mu x - \frac{1}{2}\mu^2 T} dx dy. \quad (10.37)$$

Thus

$$p_{W,W^*}(x, y) = e^{\mu x - \frac{1}{2}\mu^2 T} q_{W,W^*}(x, y). \quad (10.38)$$

The density q_{W,W^*} is given by (3.16). The joint density $p_{W,W^*}(x, y)$ can be computed (see Exercise 10.6), and the distribution of the maximum is found by $\int p_{W,W^*}(x, y) dx$.

10.4 Change of Wiener Measure

Girsanov's theorem 10.16 states that if the Wiener measure P is changed to Q by $dQ/dP = \Lambda = \mathcal{E}(\int_0^t H(s) dB(s))$ where $B(t)$ is a P -Brownian motion, and $H(t)$ is some predictable process, then Q is equivalent to P , and $B(t) = W(t) + \int_0^t H(s) ds$ for a Q -Brownian motion W . In this section we prove the converse that the Radon–Nikodym derivative of any measure Q , equivalent to the Wiener measure P , is a stochastic exponential of $\int_0^t q(s) dB(s)$ for some predictable process q . Using the predictable representation property of Brownian martingales we prove the following result first.

Theorem 10.19: *Let \mathbb{F} be Brownian motion filtration and Y be a positive random variable. If $EY < \infty$, then there exists a predictable process $q(t)$ such that $Y = (EY)e^{\int_0^T q(t) dB(t) - \frac{1}{2} \int_0^T q^2(t) dt}$.*

PROOF. Let $M(t) = E(Y|\mathcal{F}_t)$. Then $M(t)$, $0 \leq t \leq T$ is a positive uniformly integrable martingale. According to Theorem 8.35 $M(t) = EY + \int_0^t H(s) dB(s)$. Define $q(t) = H(t)/M(t)$. Then we have

$$dM(t) = H(t) dB(t) = M(t)q(t) dB(t) = M(t)dX(t), \quad (10.39)$$

with $X(t) = \int_0^t q(s) dB(s)$. Thus M is a semimartingale exponential of X , $M(t) = M(0)e^{X(t) - \frac{1}{2}[X,X](t)}$ and the result follows. \square

It remains to show that q is properly defined. We know that for each t , $M(t) > 0$ with probability one. However, there may be an exceptional set N_t , of probability zero, on which $M(t) = 0$. As there are uncountably many t 's the union of N_t 's may have a positive probability (even probability one), precluding q being finite with a positive probability. The next result shows that this is impossible.

Theorem 10.20: *Let $M(t)$, $0 \leq t \leq T$, be a martingale, such that for any t , $P(M(t) > 0) = 1$. Then $M(t)$ never hits zero on $[0, T]$.*

PROOF. Let $\tau = \inf\{t : M(t) = 0\}$. Using the basic stopping equation $EM(\tau \wedge T) = EM(T)$. So we have

$$\begin{aligned} EM(T) &= EM(\tau \wedge T) \\ &= EM(T)I(\tau > T) + EM(\tau)I(\tau \leq T) = EM(T)I(\tau > T). \end{aligned}$$

It follows that $EM(T)(1 - I(\tau > T)) = 0$. Since the random variables under the expectation are non-negative, this implies $M(T)(1 - I(\tau > T)) = 0$ almost surely. Thus $I(\tau > T) = 1$ almost surely. Thus there is a null set N such that $\tau > T$ outside N , or $P(M(t) > 0, \text{ for all } t \leq T) = 1$. Since T was arbitrary the argument implies that $\tau = \infty$ and zero is never hit. \square

Remark 10.2: Note that if for some stopping time τ a non-negative martingale is zero, $M(\tau) = 0$, and optional stopping holds, for all $t > \tau$ $E(M(t)|\mathcal{F}_\tau) = M(\tau) = 0$, then $M(t) = 0$ almost surely for all $t > \tau$.

Corollary 10.21: *Let P be the Wiener measure, $B(t)$ be a P -Brownian motion, and Q be equivalent to P . Then there exists a predictable process $q(t)$, such that*

$$\Lambda(B_{[0,T]}) = \frac{dQ}{dP}(B_{[0,T]}) = e^{\int_0^T q(t)dB(t) - \frac{1}{2} \int_0^T q^2(t)dt}. \quad (10.40)$$

Moreover, $B(t) = W(t) + \int_0^t q(s)ds$, where $W(t)$ is a Q -Brownian motion.

PROOF. According to the Radon–Nikodym theorem, $(dQ/dP) = \Lambda$ with $0 < \Lambda < \infty$. Since $P(\Lambda > 0) = 1$ and $E_P \Lambda = 1$, existence of the process $q(t)$ follows by Theorem 10.19. Representation for $B(t)$ as a Brownian motion with drift under Q follows from (10.27) in Girsanov's theorem. \square

(Corollary 10.21) is used in finance, where $q(t)$ denotes the market price for risk.

10.5 Change of Measure for Point Processes

Consider a point process $N(t)$ with intensity $\lambda(t)$ (see Chapter 9 for definitions). This presumes a probability space, which can be taken as the space of right-continuous non-decreasing functions with unit jumps, and a probability measure P so that N has intensity $\lambda(t)$ under P . Girsanov's theorem asserts that there is a probability measure Q , equivalent to P , under which $N(t)$ is a Poisson process with the unit rate. Thus an equivalent change of measure corresponds to a change in the intensity. If we look upon the compensator $\int_0^t \lambda(s)ds$ as the drift, then an equivalent change of measure results in a change of the drift.

Theorem 10.22 (Girsanov's Theorem for Poisson Processes): *Let $N(t)$ be a Poisson process with rate 1 under P , $0 \leq t \leq T$. For a constant $\lambda > 0$ define Q by*

$$\frac{dQ}{dP} = e^{(1-\lambda)T+N(T)\ln\lambda}. \quad (10.41)$$

Then under Q , N is a Poisson process with rate λ .

PROOF. Let

$$\Lambda(t) = e^{(1-\lambda)t+N(t)\ln\lambda} = e^{(1-\lambda)t}\lambda^{N(t)}. \quad (10.42)$$

It is easy to see that $\Lambda(t)$ is a P -martingale with respect to the natural filtration of the process $N(t)$. Indeed, using independence of increments of the Poisson process we have

$$\begin{aligned} E_P(\Lambda(t)|\mathcal{F}_s) &= e^{(1-\lambda)t}E_P(\lambda^{N(t)}|\mathcal{F}_s) \\ &= e^{(1-\lambda)t}\lambda^{N(s)}E_P(\lambda^{N(t)-N(s)}|\mathcal{F}_s) \\ &= e^{(1-\lambda)t}\lambda^{N(s)}E(\lambda^{N(t)-N(s)}) \\ &= e^{(1-\lambda)t}\lambda^{N(s)}e^{(\lambda-1)(t-s)} = e^{(1-\lambda)s}\lambda^{N(s)} = \Lambda(s). \end{aligned}$$

We have used that P -distribution of $N(t) - N(s)$ is Poisson with parameter $(t-s)$, hence $E_P(\lambda^{N(t)-N(s)}) = e^{(\lambda-1)(t-s)}$. We show that under Q the increments $N(t) - N(s)$ are independent of the past and have the Poisson $\lambda(t-s)$ distribution, establishing the result. Fix $u > 0$. The conditional expectation under the new measure, Theorem 10.10, and the definition of $\Lambda(t)$ (10.42) yield

$$\begin{aligned} E_Q\left(e^{u(N(t)-N(s))}|\mathcal{F}_s\right) &= E_P\left(e^{u(N(t)-N(s))}\frac{\Lambda(t)}{\Lambda(s)}|\mathcal{F}_s\right) \\ &= e^{(1-\lambda)(t-s)}E_P\left(e^{(u+\log\lambda)(N(t)-N(s))}|\mathcal{F}_s\right) \\ &= e^{(1-\lambda)(t-s)}E_P\left(e^{(u+\log\lambda)(N(t)-N(s))}\right) \\ &= e^{\lambda(t-s)(e^u-1)}. \end{aligned}$$

□

Theorem 10.23: Let $N(t)$, $0 \leq t \leq T$, be a Poisson process with rate λ under Q . Define P by

$$\frac{dP}{dQ} = e^{(\lambda-1)T - N(T)\ln \lambda}. \quad (10.43)$$

Then under P , $N(t)$ is a Poisson process with rate 1.

As a corollary we obtain that the measures of Poisson processes with constant rates are equivalent with the likelihood ratio given by the following theorem.

Theorem 10.24: Let N be a Poisson process with rate $\lambda > 0$ under the probability P_λ . Then it is a Poisson process with rate μ under the equivalent measure P_μ defined by

$$\frac{dP_\lambda}{dP_\mu} = e^{(\mu-\lambda)T + N(T)(\ln \lambda - \ln \mu)}. \quad (10.44)$$

Note that the key point in changing the measure was the martingale property (under the original measure) of the likelihood $\Lambda(t)$. This property was established directly in the proofs. Observe that the likelihood is the stochastic exponential of the point process martingale $M(t) = N(t) - A(t)$. For example, in changing the rate from 1 to λ , $\Lambda = \mathcal{E}((\lambda - 1)M)$ with $M(t) = N(t) - t$. It turns out that a general point process $N(t)$ with intensity $\lambda(t)$ can be obtained from a Poisson process with rate 1 by a change of measure. In fact, Theorem 10.24 holds for general point processes with stochastic intensities. The form of stochastic exponential $\mathcal{E}((\lambda - 1) \cdot M)$ for non-constant λ is not hard to obtain, see Exercise 10.8,

$$\Lambda = \mathcal{E}\left(\int (\lambda(s) - 1)dM(s)\right)(T) = e^{\int_0^T (1 - \lambda(s))ds + \int_0^T \ln \lambda(s)dN(s)}. \quad (10.45)$$

The next result establishes that a point process is a Poisson process (with a constant rate) under a suitable change of measure.

Theorem 10.25: Let $N(t)$ be a Poisson process with rate 1 under P , and $M(t) = N(t) - t$. If for a predictable process $\lambda(s)$, $\mathcal{E}(\int (\lambda(s) - 1)dM(s))$ is a martingale for $0 \leq t \leq T$, then under the probability measure Q defined by (10.45) $dQ/dP = \Lambda$, N is a point process with the stochastic intensity $\lambda(t)$. Conversely, if Q is absolutely continuous with respect to P , then there exists a predictable process $\lambda(t)$, such that under Q , N is a point process with the stochastic intensity $\lambda(t)$.

10.6 Likelihood Functions

When observations X are made from competing models described by the probabilities P and Q , the likelihood is the Radon–Nikodym derivative $\Lambda = dQ/dP$.

Likelihood for Discrete Observations

Suppose that we observe a discrete random variable X , and there are two competing models for X : it can come from distribution P or Q. For the observed number x the likelihood is given by

$$\Lambda(x) = \frac{Q(X=x)}{P(X=x)}. \quad (10.46)$$

Small values of $\Lambda(x)$ provide evidence for model P, and large values for model Q. If X is a continuous random variable with densities $f_0(x)$ under P and $f_1(x)$ under Q, then $dP = f_0(x)dx$, $dQ = f_1(x)dx$, and the likelihood is given by

$$\Lambda(x) = \frac{f_1(x)}{f_0(x)}. \quad (10.47)$$

If the observed data are a finite number of observations x_1, x_2, \dots, x_n , then similarly with $\mathbf{x} = (x_1, x_2, \dots, x_n)$

$$\Lambda(\mathbf{x}) = \frac{Q(\mathbf{X} = \mathbf{x})}{P(\mathbf{X} = \mathbf{x})}, \quad \text{or} \quad \Lambda(\mathbf{x}) = \frac{f_1(\mathbf{x})}{f_0(\mathbf{x})}, \quad (10.48)$$

depending on whether the models are discrete or continuous. Note that if one model is continuous and the other is discrete, then the corresponding measures are singular and the likelihood does not exist.

Likelihood Ratios for Diffusions

Let X be a diffusion solving the SDE with a P-Brownian motion $B(t)$

$$dX(t) = \mu_1(X(t), t)dt + \sigma(X(t), t)dB(t). \quad (10.49)$$

Suppose that it satisfies another equation with a Q-Brownian motion $W(t)$

$$dX(t) = \mu_2(X(t), t)dt + \sigma(X(t), t)dW(t). \quad (10.50)$$

The likelihood, as we have seen, is given by (10.33)

$$\Lambda(X_{[0,T]}) = \frac{dQ}{dP} = e^{\int_0^T \frac{\mu_2(X(t), t) - \mu_1(X(t), t)}{\sigma(X(t), t)} dB(t) - \frac{1}{2} \int_0^T \frac{(\mu_2(X(t), t) - \mu_1(X(t), t))^2}{\sigma^2(X(t), t)} dt}. \quad (10.51)$$

Since $B(t)$ is not observed directly the likelihood should be expressed as a function of the observed path $X_{[0,T]}$. Using Equation (10.49) we obtain $dB(t) = \frac{dX(t) - \mu_1(X(t), t)dt}{\sigma(X(t), t)}$, and putting this into the likelihood, we obtain

$$\Lambda(X)_T = \frac{dQ}{dP} = e^{\int_0^T \frac{\mu_2(X(t), t) - \mu_1(X(t), t)}{\sigma^2(X(t), t)} dX(t) - \frac{1}{2} \int_0^T \frac{\mu_2^2(X(t), t) - \mu_1^2(X(t), t)}{\sigma^2(X(t), t)} dt}. \quad (10.52)$$

Using the likelihood a decision can be made as to what model is more appropriate for X .

Example 10.5: (Hypotheses testing)

Suppose that we observe a continuous function x_t , $0 \leq t \leq T$, and we want to know whether it is just white noise, the null hypothesis corresponding to probability measure P or whether it is some signal contaminated by noise, the alternative hypothesis corresponding to probability measure Q .

$$H_0 : \text{noise: } dX(t) = dB(t), \quad \text{and,} \quad H_1 : \text{signal + noise: } dX(t) = h(t)dt + dB(t).$$

Here we have $\mu_1(x) = 0$, $\mu_2(x) = h(t)$, $\sigma(x) = 1$. The likelihood is given by

$$\Lambda(X)_T = \frac{dQ}{dP} = e^{\int_0^T h(t)dX(t) - \frac{1}{2} \int_0^T h^2(t)dt}. \quad (10.53)$$

This leads to the likelihood ratio test of the form: conclude the presence of noise if $\Lambda \geq k$, where k is determined from setting the probability of the type one error to α .

$$P(e^{\int_0^T h(t)dB(t) - \frac{1}{2} \int_0^T h^2(t)dt} \geq k) = \alpha. \quad (10.54)$$

Example 10.6: (Estimation in the Ornstein–Uhlenbeck model)

Consider estimation of the friction parameter α in the Ornstein–Uhlenbeck model on $0 \leq t \leq T$,

$$dX(t) = -\alpha X(t)dt + \sigma dB(t).$$

Denote by P_α the measure corresponding to $X(t)$ so that P_0 corresponds to $\sigma B(t)$, $0 \leq t \leq T$. The likelihood is given by

$$\Lambda(\alpha, X_{[0,T]}) = \frac{dP_\alpha}{dP_0} = \exp \left(\int_0^T \frac{-\alpha X(t)}{\sigma^2} dX(t) - \frac{1}{2} \int_0^T \frac{\alpha^2 X^2(t)}{\sigma^2} dt \right).$$

Maximizing the log likelihood we find

$$\hat{\alpha} = -\frac{\int_0^T X(t)dX(t)}{\int_0^T X^2(t)dt}. \quad (10.55)$$

Remark 10.3: Let $X(t)$ and $Y(t)$ satisfy the stochastic differential equations for $0 \leq t \leq T$

$$dX(t) = \mu_X(X(t), t)dt + \sigma_X(X(t), t)dW(t), \quad (10.56)$$

and

$$dY(t) = \mu_Y(Y(t), t)dt + \sigma_Y(Y(t), t)dW(t). \quad (10.57)$$

Consider probability measures induced by these diffusions on the space of continuous functions on $[0, T]$, $C[0, T]$. It turns out that if $\sigma_X \neq \sigma_Y$, then P_X and P_Y are *singular* (“live” on different sets). It means that by observing

a process continuously over an interval of time we can decide precisely from which equation it comes. This identification can be made with the help of the quadratic variation process,

$$d[X, X](t) = \sigma^2(X(t), t)dt, \quad \text{and} \quad \sigma^2(X(t), t) = \frac{d[X, X](t)}{dt}, \quad (10.58)$$

and $[X, X]$ is known exactly if a path is observed continuously.

If $\sigma_X = \sigma_Y = \sigma$ then P_X and P_Y are equivalent. The Radon–Nikodym derivatives (the likelihoods) are given by

$$\frac{dP_Y}{dP_X}(X_{[0,T]}) = e^{\int_0^T \frac{\mu_Y(X(t), t) - \mu_X(X(t), t)}{\sigma^2(X(t), t)} dX(t) - \frac{1}{2} \int_0^T \frac{\mu_Y^2(X(t), t) - \mu_X^2(X(t), t)}{\sigma^2(X(t), t)} dt}, \quad (10.59)$$

and

$$\frac{dP_X}{dP_Y}(Y_{[0,T]}) = e^{-\int_0^T \frac{\mu_Y(Y(t), t) - \mu_X(Y(t), t)}{\sigma^2(Y(t), t)} dY(t) + \frac{1}{2} \int_0^T \frac{\mu_Y^2(Y(t), t) - \mu_X^2(Y(t), t)}{\sigma^2(Y(t), t)} dt}. \quad (10.60)$$

Notes: Material for this chapter is based on Karatzas and Shreve (1988), Liptser and Shiryaev (1974), and Lamberton and Lapeyre (1996).

10.7 Exercises

Exercise 10.1: Let P be $N(\mu_1, 1)$ and Q be $N(\mu_2, 1)$ on \mathbb{R} . Show that they are equivalent and that the Radon–Nikodym derivative $dQ/dP = \Lambda$ is given by $\Lambda(x) = e^{(\mu_2 - \mu_1)x + \frac{1}{2}(\mu_1^2 - \mu_2^2)}$. Give also dP/dQ .

Exercise 10.2: Show that if X has $N(\mu, 1)$ distribution under P , then there is an equivalent measure Q such that X has $N(0, 1)$ distribution under Q . Give the likelihood dQ/dP and also dP/dQ . Give the Q -distribution of $Y = X - \mu$.

Exercise 10.3: Y has a lognormal $LN(\mu, \sigma^2)$ distribution. Using a change of measure calculate $EYI(Y > K)$. Hint: change measure from $N(\mu, \sigma^2)$ to $N(\mu + \sigma^2, \sigma^2)$.

Exercise 10.4: Let $X(t) = B(t) + \sin t$ for a P -Brownian motion $B(t)$. Let Q be an equivalent measure to P such that $X(t)$ is a Q -Brownian motion. Give $\Lambda = dQ/dP$.

Exercise 10.5: Let \mathbf{B} be an n -dim Brownian motion and \mathbf{H} an adapted regular process. Let $\mathbf{H} \cdot \mathbf{B}(T) = \sum_{i=1}^n \int_0^T H^i(s) dB^i(s)$ be a martingale. Show that the martingale exponential is given by $\exp(\mathbf{H} \cdot \mathbf{B}(T) - \frac{1}{2} \int_0^T |\mathbf{H}(s)|^2 ds)$. Hint: show that quadratic variation of $\mathbf{H} \cdot \mathbf{B}$ is given by $\int_0^T |\mathbf{H}(s)|^2 ds$, where $|\mathbf{H}(s)|^2$ denotes the length of vector $\mathbf{H}(s)$.

Exercise 10.6: Let $W_\mu(t) = \mu t + B(t)$, where $B(t)$ is a P-Brownian motion. Show that

$$P(\max_{t \leq T} W_\mu(t) \leq y | W_\mu(T) = x) = 1 - e^{\frac{-2y(y-x)}{T}}, \quad x \leq y,$$

and

$$P(\min_{t \leq T} W_\mu(t) \geq y | W_\mu(T) = x) = 1 - e^{\frac{-2y(y-x)}{T}}, \quad x \geq y.$$

Hint: use the joint distributions (10.38) and (3.16).

Exercise 10.7: Prove Theorem 10.23.

Exercise 10.8: Let $N(t)$ be a Poisson process with rate 1 and $\bar{N}(t) = N(t) - t$. Show that for an adapted, continuous, and bounded process $H(t)$, the process $M(t) = \int_0^t H(s)d\bar{N}(s)$ is a martingale for $0 \leq t \leq T$. Show that

$$\mathcal{E}(M)(t) = e^{-\int_0^t H(s)ds + \int_0^t \ln(1+H(s))dN(s)}.$$

Exercise 10.9: (Estimation of parameters)

Find the likelihood corresponding to different values of μ of the process $X(t)$ given by $dX(t) = \mu X(t)dt + \sigma X(t)dB(t)$ on $[0, T]$. Give the maximum likelihood estimator.

Exercise 10.10: Verify the martingale property of the likelihood occurring in the change of rate in a Poisson process.

Exercise 10.11: Let $dQ = \Lambda dP$ on \mathcal{F}_T , $\Lambda(t) = \mathbb{E}_P(\Lambda | \mathcal{F}_t)$ is continuous. For a P-martingale $M(t)$ find a finite variation process $A(t)$ such that $M'(t) = M(t) - A(t)$ is a Q-local martingale.

Exercise 10.12: Let B and N be respectively a Brownian motion and a Poisson process on the same space $(\Omega, \mathcal{F}, \mathbb{F}, P)$, $0 \leq t \leq T$. Define $\Lambda(t) = e^{(\ln 2)N(t)-t}$ and $dQ = \Lambda(T)dP$. Show that B is a Brownian motion under Q.

Exercise 10.13:

1. Let B and N be respectively a Brownian motion and a Poisson process on the same space $(\Omega, \mathcal{F}, \mathbb{F}, P)$, $0 \leq t \leq T$, and $X(t) = B(t) + N(t)$. Give an equivalent probability measure Q_1 such that $B(t) + t$ and $N(t) - t$ are Q_1 -martingales. Deduce that $X(t)$ is a Q_1 -martingale.
2. Give an equivalent probability measure Q_2 such that $B(t) + 2t$ and $N(t) - 2t$ are Q_2 -martingales. Deduce that $X(t)$ is a Q_2 -martingale.
3. Deduce that there are infinitely many equivalent probability measures Q such that $X(t) = B(t) + N(t)$ is a Q -martingale.

Chapter 11

Applications in Finance: Stock and FX Options

In this chapter the fundamentals of the mathematics of option pricing are given. The concept of arbitrage is introduced, and a martingale characterization of models that do not admit arbitrage is given, the first fundamental theorem of asset pricing. The theory of pricing by no arbitrage is presented first in the finite market model, and then in a general semimartingale model, where the martingale representation property is used. Change of measure and its application as the change of numeraire are given as a corollary to Girsanov's theorem and the general Bayes formula for expectations. They represent the main techniques used for pricing foreign exchange options, exotic options (asian, lookback, barrier options), and interest rates options.

11.1 Financial Derivatives and Arbitrage

A financial derivative or a contingent claim on an asset is a contract that allows purchase or sale of this asset in the future on terms that are specified in the contract. An option on stock is a basic example.

Definition 11.1: A call option on stock is a contract that gives its holder the right to buy this stock in the future at the price K written in the contract, called the exercise price or the strike.

A European call option allows the holder to exercise the contract (that is, to buy this stock at K) at a particular date T , called the maturity or the expiration date. An American option allows the holder to exercise the contract at any time before or at T .

A contract that gives its holder the right to sell stock is called a *put* option. There are two parties to a contract, the seller (writer) and the buyer (holder). The holder of the options has the right, but no obligation in exercising the contract. The writer of the options has the obligation to abide by the contract, for example, they must sell the stock at price K if the call option is exercised. Denote the price of the asset (e.g. stock) at time t by $S(t)$. A contingent claim on this asset has its value at maturity specified in the contract, and it is some function of the values $S(t)$, $0 \leq t \leq T$. Simple contracts depend only on the value at maturity $S(T)$.

Example 11.1: (Value at maturity of European call and put options)

If you hold a call option, then you can buy one share of stock at T for K . If at time T , $S(T) < K$, you will not exercise your option as you can buy stock cheaper than K , thus this option is worthless. If at time T , $S(T) \geq K$, then you can buy one share of stock for K and sell it immediately for $S(T)$ making a profit of $S(T) - K$. Thus a European call option has the value at time T

$$C(T) = (S(T) - K)^+ = \max(0, S(T) - K). \quad (11.1)$$

Similar considerations reveal that the value of the European put at maturity is

$$P(T) = (K - S(T))^+ = \max(0, K - S(T)). \quad (11.2)$$

Example 11.2: (Exotic options)

The value at maturity of exotic options depends on the prices of the asset on the whole time interval before maturity, $S(t)$, $t \leq T$.

1. Lookback Options. Lookback call pays at T , $X = (S(T) - S_{\min})^+ = S(T) - S_{\min}$ and lookback put $X = S_{\max} - S(T)$, where S_{\min} and S_{\max} denote the smallest and the largest values of $S(t)$ on $[0, T]$.
2. Barrier Options. The call that gets knocked out when the price falls below a certain level H (down and out call) pays at T

$$X = (S(T) - K)^+ I(\min_{0 \leq t \leq T} S(t) \geq H), \quad S(0) > H, \quad K > H.$$

3. Asian Options. Payoff at T depends on the average price $\bar{S} = \frac{1}{T} \int_0^T S(u) du$ during the life of the option. Average call pays $X = (\bar{S} - K)^+$, and average put $X = (K - \bar{S})^+$. Random strike option pays $X = (S(T) - \bar{S})^+$.

Whereas the value of a financial claim at maturity can be obtained from the terms specified in the contract, it is not so clear how to obtain its value prior to maturity. This is done by the pricing theory, which is also used to manage financial risk.

If the payoff of an option depends only on the price of stock at expiration, then it is possible to graph the value of an option at expiration against the underlying price of stock. For example, the payoff function of a call option is $(x - K)^+$. Some payoff functions are given in Exercise 11.1.

Arbitrage and Fair Price

Arbitrage is defined in finance as a strategy that allows one to make a profit *out of nothing* without taking any risk. Mathematical formulation of arbitrage is given later in the context of a market model.

Example 11.3: (Futures, or forward)

Consider a contract that gives the holder 1 share of stock at time T , so its value at time T is the market price $S(T)$. Denote the price of this contract at time 0 by $C(0)$. The following argument shows that $C(0) = S(0)$, as any other value results in an arbitrage profit.

If $C(0) > S(0)$, then sell the contract and receive $C(0)$; buy the stock and pay $S(0)$. The difference $C(0) - S(0) > 0$ can be invested in a risk-free bank account with interest rate r . At time T you have the stock to deliver, and make arbitrage profit of $(C(0) - S(0))e^{rT}$. If $C(0) < S(0)$, the reverse strategy results in arbitrage profit: you buy the contract and pay $C(0)$; sell the stock and receive $S(0)$ (selling the stock when you do not hold it is called short selling and it is allowed). Invest the difference $S(0) - C(0) > 0$ in a risk-free bank account. At time T exercise the contract by buying back stock at $S(T)$. The profit is $(S(0) - C(0))e^{rT}$. Thus any price $C(0)$ different from $S(0)$ results in arbitrage profit. The only case which does not result in arbitrage profit is $C(0) = S(0)$.

The fair price in a game of chance with a profit X (negative profit is loss) is the expected profit EX . The above example also demonstrates that financial derivatives are not priced by their expectations, and in general, their arbitrage-free value is different to their fair price.

The following example uses a two-point distribution for the stock on expiration to illustrate that prices for options cannot be taken as expected payoffs, and all the prices but one lead to arbitrage opportunities.

Example 11.4: The current price of the stock is \$10. We want to price a call option on this stock with the exercise price $K = 10$ and expiration in one period. Suppose that the stock price at the end of the period can have only two values \$8 and \$12 per share, and that the riskless interest rate is 10%. Suppose the call is priced at \$1 per share. Consider the strategy: buy call option on 200 shares and sell 100 shares of stock

	$S_T = 12$	$S_T = 8$	
Buy option on 200 shares	-200	400	0
Sell (short) 100 shares	1000	-1200	-800
Savings account	800	880	880
Profit	0	+80	+80

We can see that in either case $S_T = 8$ or $S_T = 12$ an arbitrage profit of \$80 is realized. It can be seen, by reversing the above strategy, that any price above \$1.36 will result in arbitrage. The price that does not lead to arbitrage is \$1.36.

Equivalence Portfolio. Pricing by No Arbitrage

The main idea in pricing by no-arbitrage arguments is to replicate the payoff of the option at maturity by a portfolio consisting of stock and bond (cash). In order to avoid arbitrage, the price of the option at any other time must equal the value of the replicating portfolio, which is valued by the market. Consider the one period case $T = 1$ first, and assume that in one period stock price moves up by factor u or down by d , $d < 1 < u$, and we want to price a claim C that has the values C_u and C_d on maturity. Schematically the following trees for prices of the stock and the option are drawn.

uS	C_u	if the price goes up
S	C	
dS	C_d	if the price goes down.

Note that the values of the option C_u and C_d are known since the values of the stock uS and dS are known by assumption. A portfolio that replicates the options consists of a shares of stock and b of bond (cash in a savings account). After one period the value of this portfolio is

$$aS_T + br = \begin{cases} auS + br & \text{if } S_T = uS \\ adS + br & \text{if } S_T = dS. \end{cases}$$

Since this portfolio is equivalent to the option, by matching the payoff, we have a system of two linear equations for a and b .

$$\left. \begin{array}{l} auS + br = C_u \\ adS + br = C_d \end{array} \right\}.$$

$$a = \frac{C_u - C_d}{(u - d)S}, \quad b = \frac{uC_d - dC_u}{(u - d)r}. \quad (11.3)$$

The price of the option must equal that of the replicating portfolio

$$C = aS + b, \quad (11.4)$$

with a and b from (11.3). In order to prove that the price is given by (11.4) consider cases when the price is above and below that value C . If an option is priced above C , then selling the option and buying the portfolio specified by a and b results in arbitrage. If the option is priced below C then buying the option and selling the portfolio results in arbitrage. It will be seen later that there are no arbitrage strategies when the option is priced at C .

Example 11.5: (continued) $K = 10$, $C_T = (S_T - K)^+$, $r = 1.1$.

$$\begin{array}{lll} 12 & u = 1.2 & 2 = C_u \\ 10 & & C \\ 8 & d = 0.8 & 0 = C_d \end{array}$$

$a = 0.5$, $b = -3.64$, from (11.3). Thus this option is replicated by the portfolio consisting of 0.5 shares and borrowing \$3.64. The initial value of this portfolio is $C = 0.5 \cdot 10 - 3.64 = 1.36$, which gives the no-arbitrage price for the call option.

The formula for the price of the option C (11.4) can be written as

$$C = aS + b = \frac{1}{r}(pC_u + (1-p)C_d), \quad (11.5)$$

with

$$p = \frac{r - d}{u - d}. \quad (11.6)$$

It can be viewed as *the discounted expected payoff of the claim, with probability p of up and $(1-p)$ down movements*. The probability p , calculated from the given returns of the stock by (11.6), is called the arbitrage-free or risk-neutral probability, and has nothing to do with subjective probabilities of the market going up or down.

In the above example $p = \frac{1.1-0.8}{1.2-0.8} = 0.75$. So that $C = \frac{1}{1.1}2 \cdot 0.75 = 1.36$.

Binomial Model

The one-period formula can be applied recursively to price a claim when trading is done one period after another. The tree for options' prices in a two-period model is given by

$$\begin{array}{c} & & C_{uu} \\ & C_u & \\ & & C_{ud} \\ C & & \\ & C_{du} & \\ C_d & & \\ & C_{dd}. \end{array}$$

The final prices of the option are known from the assumed model for stock prices. The prices prior to expiration are obtained recursively, by using the one-period formula (11.5),

$$C_u = \frac{1}{r}(pC_{uu} + (1-p)C_{ud}), \quad C_d = \frac{1}{r}(pC_{du} + (1-p)C_{dd}).$$

Using formula (11.5) again

$$C = \frac{1}{r} (pC_u + (1-p)C_d) = \frac{1}{r^2} (p^2 C_{uu} + p(1-p)C_{ud} + (1-p)pC_{du} + (1-p)^2 C_{dd}).$$

In the n period model, $T = n$, if $C_{udu\dots du} = C_{u\dots ud\dots d}$, then continuing by induction

$$C = \frac{1}{r^n} E(C_n) = \frac{1}{r^n} \sum_{i=0}^n \binom{n}{i} p^i (1-p)^{n-i} \underbrace{C_{u\dots u}}_i \underbrace{d\dots d}_{n-i} \quad (11.7)$$

is today's price of a claim which is to be exercised n periods from now. C can be seen as the expected payoff, expressed in the present dollar value, of the final payoff C_n , when the probability of the market going up on each period is the arbitrage-free probability p . For a call option

$$\underbrace{C_{u\dots u}}_i \underbrace{d\dots d}_{n-i} = (u^i d^{n-i} S - K)^+,$$

and C can be written by using the complementary binomial cumulative probability distribution function $\text{Bin}(j; n, p) = P(\text{Bin}(n, p) > j)$

$$C = S \text{Bin}(j; n, p') - r^{-n} K \text{Bin}(j; n, p),$$

where $j = \lceil \frac{\ln(K/Sd^n)}{\ln(u/d)} \rceil$ and $p' = \frac{u}{r} p$.

It is possible to obtain the option pricing formula of Black and Scholes from the binomial formula by taking limits as the length of the trading period goes to zero and the number of trading periods n goes to infinity.

Pricing by No Arbitrage

Given a general model these are the questions we ask.

1. If we have a model for the evolution of prices, *how can we tell if there are arbitrage opportunities?* Not finding any is a good start, but not a proof that there are none.
2. If we know that there are no arbitrage opportunities in the market, *how do we price a claim, such as an option?*
3. *Can we price any option, or are there some that cannot be priced by arbitrage arguments?*

The answers are given by two main results called the fundamental theorems of asset pricing. In what follows we outline the mathematical theory of pricing of claims in finite and general market models.

11.2 A Finite Market Model

Consider a model with one stock with price $S(t)$ at time t , and a riskless investment (bond, or cash in a savings account) with price $\beta(t)$ at time t . If the riskless rate of investment is a constant $r > 1$, then $\beta(t) = r^t \beta(0)$.

A market model is called finite if $S(t), t = 0, \dots, T$ take finitely many values. A portfolio $(a(t), b(t))$ is the number of shares of stock and bond units held during $[t-1, t)$. The information available after observing prices up to time t is denoted by the σ -field \mathcal{F}_t . The portfolio is decided on the basis of information at time $t-1$, in other words $a(t), b(t)$ are \mathcal{F}_{t-1} measurable, $t = 1, \dots, T$, or, in our terminology, they are predictable processes. The change in market value of the portfolio at time t is the difference between its value after it has been established at time $t-1$ and its value after prices at t are observed, namely

$$a(t)S(t) + b(t)\beta(t) - a(t)S(t-1) - b(t)\beta(t-1) = a(t)\Delta S(t) + b(t)\Delta\beta(t).$$

A portfolio (trading strategy) is called *self-financing* if all the changes in the portfolio are due to gains realized on investment, that is, no funds are borrowed or withdrawn from the portfolio at any time,

$$V(t) = V(0) + \sum_{i=1}^t (a(i)\Delta S(i) + b(i)\Delta\beta(i)), \quad t = 1, 2, \dots, T. \quad (11.8)$$

The initial value of the portfolio $(a(t), b(t))$ is $V(0) = a(1)S(0) + b(1)\beta(0)$ and subsequent values $V(t)$, $t = 1, \dots, T$ are given by

$$V(t) = a(t)S(t) + b(t)\beta(t). \quad (11.9)$$

$V(t)$ represents the value of the portfolio just before time t transactions after time t price was observed. Since the market value of the portfolio $(a(t), b(t))$ at time t after $S(t)$ is announced is $a(t)S(t) + b(t)\beta(t)$, and the value of the newly set up portfolio is $a(t+1)S(t) + b(t+1)\beta(t)$, a self-financing strategy must satisfy

$$a(t)S(t) + b(t)\beta(t) = a(t+1)S(t) + b(t+1)\beta(t). \quad (11.10)$$

A strategy is called *admissible* if it is self-financing and the corresponding value process is non-negative.

A contingent claim is a non-negative random variable X on (Ω, \mathcal{F}_T) . It represents an agreement which pays X at time T , for example, for a call with strike K , $X = (S(T) - K)^+$.

Definition 11.2: A claim X is called attainable if there exists an admissible strategy replicating the claim, that is, $V(t)$ satisfies (11.8), $V(t) \geq 0$ and $V(T) = X$.

Definition 11.3: An arbitrage opportunity is an admissible trading strategy such that $V(0) = 0$, but $\text{EV}(T) > 0$.

Note that since $V(T) \geq 0$, $\text{EV}(T) > 0$ is equivalent to $P(V(T) > 0) > 0$. The following result is central to the theory.

Theorem 11.4: Suppose there is a probability measure Q such that the discounted stock process $Z(t) = S(t)/\beta(t)$ is a Q -martingale. Then for any admissible trading strategy the discounted value process $V(t)/\beta(t)$ is also a Q -martingale.

Such Q is called an equivalent martingale measure (EMM) or a risk-neutral probability measure.

PROOF. Since the market is finite the value process $V(t)$ takes only finitely many values, therefore $\text{EV}(t)$ exist. The martingale property is verified as follows:

$$\begin{aligned} E_Q \left(\frac{V(t+1)}{\beta(t+1)} \middle| \mathcal{F}_t \right) &= E_Q(a(t+1)Z(t+1) + b(t+1) | \mathcal{F}_t) \\ &= a(t+1)E_Q(Z(t+1) | \mathcal{F}_t) + b(t+1) \text{ since } a(t) \\ &\quad \text{and } b(t) \text{ are predictable} \\ &= a(t+1)Z(t) + b(t+1) \text{ since } Z(t) \text{ is a martingale} \\ &= a(t)Z(t) + b(t) \text{ since } (a(t), b(t)) \text{ is self-financing (11.10)} \\ &= \frac{V(t)}{\beta(t)}. \end{aligned}$$

□

This result is “nearly” the condition for no-arbitrage since it states that if we start with zero wealth, then positive wealth cannot be created if probabilities are assigned by Q . Indeed, by the above result $V(0) = 0$ implies $E_Q(V(T)) = 0$, and $Q(V(T) = 0) = 1$. However, in the definition of an arbitrage strategy the expectation is taken under the original probability measure P . In order to establish the result for P , equivalence of probability measures is used. Recall the definition from Chapter 10.

Definition 11.5: Two probability measures P and Q are called equivalent if they have the same null sets, that is, for any set A with $P(A) = 0$, $Q(A) = 0$ and vice versa.

Equivalent probability measures in a market model reflect the fact that investors agree on the space of all possible outcomes, but assign different probabilities to the outcomes. The following result gives a probabilistic condition to assure that the model does not allow for arbitrage opportunities. It states that no arbitrage is equivalent to the existence of EMM.

Theorem 11.6 (First Fundamental Theorem): *A market model does not have arbitrage opportunities if, and only if, there exists a probability measure Q , equivalent to P , such that the discounted stock process $Z(t) = S(t)/\beta(t)$ is a Q -martingale.*

PROOF. Proof of sufficiency. Suppose there exists a probability measure Q , equivalent to P such that the discounted stock process $Z(t) = S(t)/\beta(t)$ is a Q -martingale. Then there are no arbitrage opportunities. According to Theorem 11.4 any admissible strategy with $V(0) = 0$ must have $Q(V(T) > 0) = 0$. Since Q is equivalent to P , $P(V(T) > 0) = 0$, but then $E_P(V(T)) = 0$. Thus there are no admissible strategies with $V(0) = 0$ and $E_P(V(T)) > 0$, in other words, there is no arbitrage.

A proof of necessity requires additional concepts (see for example Harrison and Pliska (1981)). \square

Claims are priced by the replicating portfolios.

Theorem 11.7 (Pricing by No Arbitrage): *Suppose that the market model does not admit arbitrage, and X is an attainable claim with maturity T . Then $C(t)$, the arbitrage-free price of X at time $t \leq T$, is given by $V(t)$, the value of a portfolio of any admissible strategy replicating X . Moreover,*

$$C(t) = V(t) = E_Q \left(\frac{\beta(t)}{\beta(T)} X | \mathcal{F}_t \right), \quad (11.11)$$

where Q is an equivalent martingale probability measure.

PROOF. Since X is attainable it is replicated by an admissible strategy with the value of the replicating portfolio $V(t)$, $0 \leq t \leq T$, and $X = V(T)$. Fix one such strategy. In order to avoid arbitrage, the price of X at any time $t < T$ must be given by the value of this portfolio $V(t)$, otherwise arbitrage profit is possible.

Since the model does not admit arbitrage a martingale probability measure Q exists by Theorem 11.6. The discounted value process $V(t)/\beta(t)$ is a Q -martingale by Theorem 11.4, hence by the martingale property

$$\frac{V(t)}{\beta(t)} = E_Q \left(\frac{V(T)}{\beta(T)} | \mathcal{F}_t \right). \quad (11.12)$$

However, $V(T) = X$, and we have

$$\frac{V(t)}{\beta(t)} = E_Q \left(\frac{1}{\beta(T)} X | \mathcal{F}_t \right). \quad (11.13)$$

\square

Note that $\frac{\beta(t)}{\beta(T)}X$ represents the value of the claim X in dollars at time t , and the price of X at time $t = 0$ is given by

$$C(0) = E_Q \left(\frac{1}{\beta(T)} X \right). \quad (11.14)$$

Remark 11.1: The lhs of Equation (11.13) $V(t)/\beta(t) = a(t)Z(t) + b(t)$ is determined by the portfolio $a(t), b(t)$, but its rhs $E_Q \left(\frac{1}{\beta(T)} X | \mathcal{F}_t \right)$ is determined by the measure Q and has nothing to do with a chosen portfolio. This implies that for a given martingale measure Q and any t , all self-financing portfolios replicating X have the same value, moreover, this common value is also the same for different martingale probability measures. Thus Equation (11.11) provides an unambiguous price for the claim X at time t .

However, if a claim X is not attainable its expectation may vary with the measure Q (see Example 11.7 below).

Now we know how to price attainable claims. If all the claims in the market are attainable, then we can price any claim.

Definition 11.8: Market models in which any claim is attainable are called complete.

The following result characterizes complete models in terms of the martingale measure. The proof can be found in Harrison and Kreps (1979), and Harrison and Pliska (1983).

Theorem 11.9 (Completeness): *The market model is complete if, and only if, the martingale probability measure Q is unique.*

Example 11.6: (A complete model)

The one-step binomial model ($t = 0, 1$) can be described by the payoff vector of the stock (d, u) and of the bond (r, r) . A claim X is a vector (x_1, x_2) representing the payoff of the claim when the market goes down, x_1 , and up, x_2 . As the two vectors (r, r) and (d, u) span \mathbb{R}^2 , any vector is a linear combination of those two, hence any claim can be replicated by a portfolio, and the model is complete. In order to find a martingale probability Q , we solve $E_Q Z(1) = Z(0)$, or $\frac{1}{r}(pu + (1-p)d) = 1$. The unique solution for $p = \frac{r-d}{u-d}$. It is a probability if, and only if, $d < r < u$. Thus the existence and uniqueness of the martingale measure is verified. In this model any claim can be priced by no-arbitrage considerations.

Example 11.7: (An incomplete model)

This is the model where a stock's payoff can take three possible values $(d, 1, u)$ and the bond with payoff $(1, 1, 1)$ is not complete. These vectors span a subspace of \mathbb{R}^3 , therefore not all possible returns can be replicated. For example, the claim that pays \$1 when the stock goes up and nothing in any other case has the payoff

$(0, 0, 1)$ and cannot be replicated. We now verify that there are many martingale measures. In order to find them we must solve $E_Q Z(1) = Z(0)$.

$$\left. \begin{aligned} \frac{1}{r}[p_u u + p_n + p_d d] &= 1 \\ p_u + p_n + p_d &= 1 \end{aligned} \right\}$$

Any solution of this system makes $Z(t)$, $t = 0, 1$ into a martingale.

$$p_u = \frac{r - 1}{u - 1} + \frac{1 - d}{u - 1} p_d, \quad p_n = \frac{u - r}{u - 1} - \frac{u - d}{u - 1} p_d, \quad 0 \leq p_u, p_n, p_d \leq 1.$$

11.3 Semimartingale Market Model

Arbitrage in Continuous Time Models

In continuous time there are different versions of the no-arbitrage concept. The main premise is the same as in discrete time, it should be impossible to make “something” out of nothing without taking risk. The difference between different versions of no arbitrage is in the kinds of allowable (admissible) self-financing strategies that define how “something” is made.

Let $a(t)$ and $b(t)$ denote the number of shares and bond units respectively held at time t . The market value of the portfolio at time t is given by

$$V(t) = a(t)S(t) + b(t)\beta(t). \quad (11.15)$$

The change in the value of the portfolio due to change in the price of assets during dt is $a(t)dS(t) + b(t)d\beta(t)$.

Definition 11.10: A portfolio $(a(t), b(t))$, $0 \leq t \leq T$, is called self-financing if the change in value comes only from the change in the prices of the assets,

$$dV(t) = a(t)dS(t) + b(t)d\beta(t), \quad (11.16)$$

$$V(t) = V(0) + \int_0^t a(u)dS(u) + \int_0^t b(u)d\beta(u). \quad (11.17)$$

It is assumed that $S(t)$ and $\beta(t)$ are semimartingales. The processes $a(t)$ and $b(t)$ must be predictable processes satisfying a certain condition for the stochastic integral to be defined, see (8.12) and (8.52).

In a general situation, when both $S(t)$ and $\beta(t)$ are stochastic, the rhs of (11.17) can be seen as a stochastic integral with respect to the vector process $(S(t), \beta(t))$. Such integrals extend the standard definition of scalar stochastic integrals and can be defined for a larger class of integrands due to possible interaction between the components. We do not go into details as they are rather complicated, and refer to Shiryaev (1999), and Jacod and

Shiryayev (1987). The following concepts describe no arbitrage: no arbitrage (NA), no free lunch (NFL), no free lunch with bounded risk (NFLBR), no free lunch with vanishing risk (NFLVR), no feasible free lunch with vanishing risk (NFFLVR) (see Shiryaev (1999), Kreps (1981), Delbaen and Schachermayer (1994), and Kabanov (2001)). We do not aim to present the arbitrage theory in continuous time and consider a simpler formulation, following Harrison and Pliska (1981).

We consider a model in continuous time $0 \leq t \leq T$ consisting of two assets, a semimartingale $S(t)$ representing the stock price process, and the savings account (or bond) $\beta(t)$, $\beta(0) = 1$. We assume $\beta(t)$ is continuous and of finite variation. The following is a central result.

Theorem 11.11: *($a(t), b(t)$) is self-financing if, and only if, the discounted value process $\frac{V(t)}{\beta(t)}$ is a stochastic integral with respect to the discounted price process*

$$\frac{V(t)}{\beta(t)} = V(0) + \int_0^t a(u)dZ(u), \quad (11.18)$$

where $Z(t) = S(t)/\beta(t)$.

PROOF. Using the assumption that the bond process is continuous and of finite variation, we have

$$\begin{aligned} d\left(\frac{V(t)}{\beta(t)}\right) &= \frac{1}{\beta(t-)}dV(t) + V(t-)d\left(\frac{1}{\beta(t)}\right) + d[V, \frac{1}{\beta}](t) \\ &= \frac{1}{\beta(t)}dV(t) + V(t-)d\left(\frac{1}{\beta(t)}\right). \end{aligned}$$

Using the self-financing property

$$\begin{aligned} d\left(\frac{V(t)}{\beta(t)}\right) &= \frac{1}{\beta(t)}(a(t)dS(t) + b(t)d\beta(t)) + (a(t)S(t- \\ &\quad + b(t)\beta(t))d\left(\frac{1}{\beta(t)}\right) = a(t)\underbrace{\left(\frac{dS(t)}{\beta(t-)} + S(t-)d\left(\frac{1}{\beta(t)}\right)\right)}_{dZ(t)=d(S(t)/\beta(t))} \\ &\quad + b(t)\underbrace{\left(\frac{1}{\beta(t)}d\beta(t) + \beta(t)d\left(\frac{1}{\beta(t)}\right)\right)}_{d(\beta(t)\cdot\frac{1}{\beta(t)})=0} = a(t)dZ(t). \end{aligned}$$

The other direction. Assume (11.18). From $V(t) = a(t)S(t) + b(t)\beta(t)$ we find $b(t) = V(t)/\beta(t) - a(t)Z(t)$. Using (11.18)

$$b(t) = V(0) + \int_0^t a(u)dZ(u) - a(t)Z(t). \quad (11.19)$$

Using (11.18) $V(t) = V(0)\beta(t) + \beta(t) \int_0^t a(u)dZ(u)$. Hence

$$\begin{aligned} dV(t) &= V(0)d(\beta(t)) + \int_0^t a(u)dZ(u)d(\beta(t)) + \beta(t)a(t)dZ(t) \\ &= V(0)d(\beta(t)) + (b(t) - V(0) + a(t)Z(t))d(\beta(t)) + \beta(t)a(t)dZ(t) \\ &= b(t)d(\beta(t)) + a(t)Z(t)d(\beta(t)) + \beta(t)a(t)dZ(t) \\ &= b(t)d(\beta(t)) + a(t)d(Z(t)\beta(t)) = b(t)d(\beta(t)) + a(t)dS(t), \end{aligned}$$

and self-financing property of $V(t)$ is established. \square

The basis for the mathematical formulation of arbitrage is the existence of the EMM, the equivalent martingale probability measure. In a general model existence of such a measure is introduced as an assumption.

EMM Assumption

There exists a martingale probability measure Q which is equivalent to the original measure P such that the discounted price process $Z(t) = S(t)/\beta(t)$ is a Q -martingale.

We give examples of models where the EMM assumption does and does not hold.

Example 11.8: $dS(t) = .04S(t)dt$. $d\beta(t) = 0.03\beta(t)dt$. $S(t) = S(0)e^{0.04t}$, $\beta(t) = e^{0.03t}$, $S(t)e^{-0.03t} = e^{0.01t}$. Since $e^{0.01t}$ is a deterministic non-constant function, there is no probability measure Q that would make it into a martingale.

Example 11.9: $S(t) = S(0) + \int_0^t B(s)ds$, where $B(s)$ is a P -Brownian motion. According to Corollary 10.21 an equivalent change of measure results in $B(t)$ being transformed into $B(t) + q(t)$, for some process $q(t)$. Thus under the EMM Q , $S(t) = S(0) + \int_0^t B(s)ds + \int_0^t q(s)ds$. Since $S(t)$ has finite variation it cannot be a martingale because a continuous martingale is either a constant or has infinite variation (see Theorem 7.29). Hence there is no EMM in this model.

Example 11.10: (Bachelier model)

$$S(t) = S(0) + \mu t + \sigma B(t), \quad (11.20)$$

for positive constants μ, σ . $\beta(t) = 1$. The EMM exists (and is unique) by Girsanov's theorem.

Example 11.11: (Black–Scholes model)

$$dS(t) = \mu S(t)dt + \sigma S(t)dB(t), \quad \beta(t) = e^{rt}. \quad (11.21)$$

Solving the SDE for $S(t)$, $Z(t) = S(t)e^{-rt} = S(0)e^{(\mu - r - \frac{1}{2}\sigma^2)t + \sigma B(t)}$. This process is a martingale if, and only if, $\mu = r$; when $\mu = r$ it is the exponential martingale of

$\sigma B(t)$, when $\mu \neq r$ it is a martingale times a non-constant deterministic function, easily seen as not a martingale. Writing $dS(t) = \sigma S(t)(\frac{\mu}{\sigma}dt + dB(t))$ and using the change of drift in diffusions, there is a (unique) Q such that $\frac{\mu}{\sigma}dt + dB(t) = \frac{r}{\sigma}dt + dW(t)$ for a Q Brownian motion $W(t)$. So $\sigma B(t) = rt + \sigma W(t) - \mu t$. Thus the equation for $Z(t)$ in terms of $W(t)$ is

$$Z(t) = S(0)e^{(\mu - r - \frac{1}{2}\sigma^2)t + \sigma B(t)} = S(0)e^{-\frac{1}{2}\sigma^2t + \sigma W(t)},$$

verifying that Q is the EMM.

Admissible Strategies

The discounted value of a replicating self-financing portfolio $V(t)/\beta(t)$ is a stochastic integral with respect to the Q -martingale $Z(t)$, Theorem 11.11. We would like it to be a martingale because then all its values can be determined by its final value, which is matched to the claim X . The martingale property implies $V(t)/\beta(t) = E_Q(V(T)/\beta(T)|\mathcal{F}_t) = E_Q(X/\beta(T)|\mathcal{F}_t)$.

However, a stochastic integral with respect to a martingale is only a local martingale. Thus the discounted value process of a self-financing portfolio in (11.18) is a local martingale under Q . Since it is non-negative, it is a supermartingale (Theorem 7.23). Supermartingales have non-increasing expectations. In particular, there are strategies (called suicidal) that can turn the initial investment into nothing. Adding such a strategy to any other self-financing portfolio will change the initial value without changing the final value. Thus there are self-financing strategies with the same final value but different initial values. This phenomenon is precisely the difference in the situation between the finite market model and the general model. Note that, similar to the finite market model, a self-financing strategy cannot create something out of nothing since the expectations of such strategies are non-increasing. In order to eliminate the undesirable strategies from consideration only martingale strategies are admissible. We follow Harrison and Pliska (1981).

Fix a reference EMM, equivalent martingale probability measure Q , so that the discounted stock price process $Z(t)$ is a Q -martingale; expectations are taken with respect to this measure Q .

Definition 11.12: A predictable and self-financing strategy $(a(t), b(t))$ is called admissible if

$$\sqrt{\int_0^t a^2(u)d[Z, Z](u)} \text{ is finite and locally integrable } 0 \leq t \leq T. \quad (11.22)$$

Moreover, $V(t)/\beta(t)$ is a non-negative Q -martingale.

Note that condition (11.22) is needed to define the stochastic integral $\int_0^t a(u)dZ(u)$, see condition (8.8). If (11.22) holds, then $\int_0^t a(u)dZ(u)$ and consequently $V(t)/\beta(t)$ are local martingales. If, moreover, $V(t)/\beta(t) \geq 0$, then it is a supermartingale.

Pricing of Claims

A claim is a non-negative random variable. It is attainable if it is integrable, $EX < \infty$, and there exists an admissible trading strategy such that at maturity T , $V(T) = X$. In order to avoid arbitrage the value of an attainable claim at time $t < T$ must be the same as that of the replicating portfolio at t .

Theorem 11.13: *The price $C(t)$ at time t of an attainable claim X is given by the value of an admissible replicating portfolio $V(t)$, moreover,*

$$C(t) = E_Q \left(\frac{\beta(t)}{\beta(T)} X | \mathcal{F}_t \right). \quad (11.23)$$

The proof follows by the martingale property of $V(t)/\beta(t)$ in exactly the same way as in the finite model case.

$$\begin{aligned} C(t) &= V(t) = \beta(t)E_Q(V(T)/\beta(T)|\mathcal{F}_t) \\ &= \beta(t)E_Q \left(\frac{V(T)}{\beta(T)} | \mathcal{F}_t \right) = \beta(t)E_Q \left(\frac{X}{\beta(T)} | \mathcal{F}_t \right). \end{aligned}$$

Since attainable claims can be priced, the natural question is “how can one tell whether a claim is attainable”? The following result gives an answer using the predictable representation property of the discounted stock price.

Theorem 11.14: *Let X be an integrable claim and let $M(t) = E_Q \left(\frac{X}{\beta(T)} | \mathcal{F}_t \right)$, $0 \leq t \leq T$. Then X is attainable if, and only if, $M(t)$ can be represented in the form*

$$M(t) = M(0) + \int_0^t H(u)dZ(u)$$

for some predictable process H . Moreover, $V(t)/\beta(t) = M(t)$ is the same for any admissible portfolio that replicates X .

PROOF. Suppose X is attainable, and that $(a(t), b(t))$ replicates it. According to the previous result $V(t)/\beta(t) = M(t)$. It follows by (11.18) that the desired representation holds with $H(t) = a(t)$. Conversely, if $M(t) = M(0) + \int_0^t H(u)dZ(u)$, take $a(t) = H(t)$, and $b(t) = M(0) + \int_0^t H(u)dZ(u) - H(t)Z(t)$. This gives a self-financing strategy. \square

Completeness of a Market Model

A market model is *complete* if any integrable claim is attainable, in other words, can be replicated by a self-financing portfolio. If a model is complete, then any claim can be priced by no-arbitrage considerations.

We know by Theorem 11.14 that for a claim to be attainable the martingale $M(t) = \mathbb{E}_Q(\frac{X}{\beta(T)} | \mathcal{F}_t)$ should have a predictable representation with respect to the Q-martingale $Z(t) = S(t)/\beta(t)$. Recall that the martingale $Z(t)$ has the predictable representation property if any other martingale can be represented as a stochastic integral with respect to it (see Definition 8.34). For results on predictable representations see Chapter 8.12. In particular, if the martingale $Z(t)$ has the predictable representation property, then all claims in the model are attainable. It turns out that the opposite is also true, moreover, there is a surprising characterization, the equivalent martingale measure Q is unique.

Theorem 11.15 (Second Fundamental Theorem): *The following are equivalent:*

1. *The market model is complete.*
2. *The martingale $Z(t)$ has the predictable representation property.*
3. *The EMM Q that makes $Z(t) = S(t)/\beta(t)$ into a martingale is unique.*

11.4 Diffusion and the Black–Scholes Model

In this section we apply general results to the diffusion models of stock prices. In a diffusion model the stock price is assumed to satisfy

$$dS(t) = \mu(S(t))dt + \sigma(S(t))dB(t), \quad (11.24)$$

where $B(t)$ is P-Brownian motion. Bond price is assumed to be deterministic and continuous $\beta(t) = \exp(\int_0^t r(u)du)$. According to Theorem 11.13 pricing of claims is done under the martingale probability measure Q that makes $Z(t) = S(t)/\beta(t)$ into a martingale.

Theorem 11.16: *Let $H(t) = \frac{\mu(S(t)) - r(t)S(t)}{\sigma(S(t))}$. Suppose that $\mathcal{E}(\int_0^t H(u)du | \mathcal{F}_t)$ is a martingale. Then the EMM exists and is unique. It is defined by*

$$\Lambda = \frac{dQ}{dP} = \mathcal{E}\left(-\int_0^t H(u)dB(u)\right)(T) = e^{-\int_0^T H(u)dB(u) - \frac{1}{2} \int_0^T H^2(u)dt}. \quad (11.25)$$

The SDE for $S(t)$ under Q with a Brownian motion W is

$$dS(t) = r(t)S(t)dt + \sigma(S(t))dW(t). \quad (11.26)$$

PROOF. It is easy to see by Itô's formula that $Z(t)$ satisfies

$$\begin{aligned} dZ(t) &= d\left(\frac{S(t)}{\beta(t)}\right) = \frac{\mu(S(t)) - r(t)S(t)}{\beta(t)}dt + \frac{\sigma(S(t))}{\beta(t)}dB(t). \\ &= \frac{\sigma(S(t))}{\beta(t)} \left(\frac{\mu(S(t)) - r(t)S(t)}{\sigma(S(t))} dt + dB(t) \right). \end{aligned}$$

For $Z(t)$ to be a martingale it must have zero drift coefficient. Define Q by (11.25). Using (10.34), $\frac{\mu(S(t)) - r(t)S(t)}{\sigma(S(t))}dt + dB(t) = dW(t)$ with a Q Brownian motion $W(t)$. This gives the SDE

$$dZ(t) = d\left(\frac{S(t)}{\beta(t)}\right) = \frac{\sigma(S(t))}{\beta(t)}dW(t).$$

Using integration by parts we obtain the SDE (11.26) under the EMM Q for $S(t)$. \square

In order to price claims expectations should be calculated by using Equation (11.26), and not the original one in (11.24). The effect of the change of measure is the change in the drift: $\mu(x)$ is changed into rx , where r is the riskless interest rate.

Black–Scholes Model

The Black–Scholes model is the commonly accepted model for pricing of claims in the financial industry. The main assumptions of the model are: the riskless interest rate is a constant r , $\sigma(S(t)) = \sigma S(t)$, where the constant σ is called “volatility”. The stock price processes $S(t)$ satisfies SDE

$$dS(t) = \mu S(t)dt + \sigma S(t)dB(t). \quad (11.27)$$

Using Itô's formula with $f(x) = \ln x$ (see Example 5.5) we find that the solution is given by

$$S(t) = S(0)e^{(\mu - \frac{\sigma^2}{2})t + \sigma B(t)}. \quad (11.28)$$

This model corresponds to the simplest random model for the return $R(t)$ on stock

$$dR(t) = \frac{dS(t)}{S(t)} = \mu dt + \sigma dB(t). \quad (11.29)$$

$S(t)$ is the stochastic exponential of $R(t)$, $dS(t) = S(t)dR(t)$. According to Theorem 8.12 $S(t) = S(0)\mathcal{E}(R)_t = S(0)e^{R(t) - \frac{1}{2}[R, R](t)}$, giving (11.28).

The EMM Q makes $S(t)e^{-rt}$ into a martingale. According to Theorem 11.16 it exists and is unique. It is obtained by letting $\frac{\mu}{\sigma}dt + dB(t) = \frac{r}{\sigma}dt + dW(t)$, for a Q-Brownian motion $W(t)$. In this case $H(t) = \frac{\mu-r}{\sigma}$,

$\int_0^t H(t)dB(t) = \frac{\mu-r}{\sigma}B(t)$ and $\mathcal{E}\left(-\int_0^t H(t)dB(t)\right)(T) = \mathcal{E}\left(-\frac{\mu-r}{\sigma}B\right)(T)$ is the exponential martingale of Brownian motion. The SDE (11.26) under the EMM Q for $S(t)$ is

$$dS(t) = rS(t)dt + \sigma S(t)dW(t). \quad (11.30)$$

It has the solution

$$S(t) = S(0)e^{(r-\frac{\sigma^2}{2})t+\sigma W(t)}. \quad (11.31)$$

Thus under the equivalent martingale measure Q, $S(T)$ has a lognormal distribution, $LN\left((r - \frac{\sigma^2}{2})T + \ln S(0), \sigma^2 T\right)$. The price of a claim X at time T is given by

$$C(t) = e^{-r(T-t)}\mathbb{E}_Q(X|\mathcal{F}_t). \quad (11.32)$$

If $X = g(S(T))$, then by the Markov property of $S(t)$, $C(t) = \mathbb{E}_Q(g(S(T))|\mathcal{F}_t) = \mathbb{E}_Q(g(S(T))|S(t))$. The conditional distribution under Q given \mathcal{F}_t is obtained from the equation (using (11.31))

$$S(T) = S(t)e^{(r-\frac{\sigma^2}{2})(T-t)+\sigma(W(T)-W(t))}. \quad (11.33)$$

Hence it is lognormal, $LN\left((r - \frac{\sigma^2}{2})(T - t) + \ln S(t), \sigma^2(T - t)\right)$.

Pricing a Call Option

A call option pays $(S(T) - K)^+$ at time T . Recall that its price at time t is given by (Theorem 11.13)

$$C(t) = e^{-r(T-t)}\mathbb{E}_Q((S(T) - K)^+|\mathcal{F}_t).$$

As we have just seen above, the conditional Q-distribution of S_T given \mathcal{F}_t is lognormal with mean $\mu = (r - \frac{\sigma^2}{2})(T - t) + \ln S(t)$ and $\sigma_1^2 = \sigma^2(T - t)$, write it as e^Y where Y has $N(\mu, \sigma_1^2)$ distribution. Thus the expectation above is the same as $\mathbb{E}(e^Y - K)^+$. Writing now P and E for the probabilities and expectation under $N(\mu, \sigma_1^2)$ distribution we have

$$\mathbb{E}(e^Y - K)^+ = \mathbb{E}((e^Y - K)I(e^Y > K)) = \mathbb{E}(e^Y I(e^Y > K)) - KP(e^Y > K).$$

The second term equals $K(1 - \Phi((\log K - \mu)/\sigma_1))$. The first term can be calculated by a direct integration, or by changing the measure, as in Example 10.2. We recall the change of measure calculations. Take $dQ_1/dP = \Lambda = e^Y/\mathbb{E}e^Y$. Then

$$\begin{aligned} \mathbb{E}(e^Y I(e^Y > K)) &= \mathbb{E}e^Y \mathbb{E}((e^Y/\mathbb{E}e^Y)I(e^Y > K)) = \mathbb{E}e^Y \mathbb{E}(\Lambda I(e^Y > K)) \\ &= \mathbb{E}e^Y \mathbb{E}_{Q_1} I(e^Y > K) = \mathbb{E}e^Y Q_1(e^Y > K). \end{aligned}$$

Since $dQ_1/dP = e^{Y-\mu-\sigma_1^2/2}$ it follows by Theorem 10.4 that Q_1 is in fact the $N(\mu + \sigma_1^2, \sigma_1^2)$ distribution. Therefore

$$\mathbb{E}(e^Y I(e^Y > K)) = e^{\mu+\sigma_1^2/2} (1 - \Phi((\ln K - \mu - \sigma_1^2)/\sigma_1)).$$

Using the property $1 - \Phi(x) = \Phi(-x)$ we obtain

$$\mathbb{E}(e^Y - K)^+ = e^{\mu+\sigma^2/2} \Phi\left(\frac{\mu + \sigma_1^2 - \ln K}{\sigma_1}\right) - K \Phi\left(\frac{\mu - \ln K}{\sigma_1}\right). \quad (11.34)$$

Finally, replacing μ and σ_1 by their values $\mu = (r - \frac{\sigma^2}{2})(T-t) + \ln S(t)$ and $\sigma_1^2 = \sigma^2(T-t)$ we obtain that $C(t)$ is given by the Black–Scholes formula

$$C(t) = S(t)\Phi(h(t)) - Ke^{-r(T-t)}\Phi\left(h(t) - \sigma\sqrt{T-t}\right), \quad (11.35)$$

where

$$h(t) = \frac{\ln \frac{S(t)}{K} + (r + \frac{1}{2}\sigma^2)(T-t)}{\sigma\sqrt{T-t}}. \quad (11.36)$$

Pricing of Claims by a PDE. Replicating Portfolio

Let X be a claim of the form $X = g(S(T))$. Since the stock price satisfies SDE (11.30), by the Markov property of $S(t)$ it follows from (11.32) that the price of X at time t

$$C(t) = e^{-r(T-t)}\mathbb{E}_Q(g(S(T))|\mathcal{F}_t) = e^{-r(T-t)}\mathbb{E}_Q(g(S(T))|S(t)). \quad (11.37)$$

According to the Feynman–Kac formula (Theorem 6.8),

$$C(x, t) = e^{-r(T-t)}\mathbb{E}_Q(g(S(T))|S(t) = x)$$

solves the following PDE

$$\frac{1}{2}\sigma^2 x^2 \frac{\partial^2 C(x, t)}{\partial x^2} + rx \frac{\partial C(x, t)}{\partial x} + \frac{\partial C(x, t)}{\partial t} - rC = 0. \quad (11.38)$$

The boundary condition is given by the value of the claim at maturity

$$C(x, T) = g(x), \quad x \geq 0,$$

with $g(x)$ as the value of the claim at time T when the stock price is x . When $x = 0$, Equation (11.38) gives

$$C(0, t) = e^{-r(T-t)}g(0), \quad 0 \leq t \leq T,$$

(a portfolio of only a bond has its value at time t as the discounted final value). For a call option $g(x) = (x - K)^+$. This partial differential equation was solved by Black and Scholes with the Fourier transform method. The solution is the Black–Scholes formula (11.35).

Next we give an argument to find the replicating portfolio, which is also used to re-derive the PDE (11.38). It is clear that $C(t)$ in (11.37) is a function of $S(t)$ and t . Let $(a(t), b(t))$ be a replicating portfolio. Since it is self-financing its value process satisfies

$$dV(t) = a(t)dS(t) + b(t)d\beta(t). \quad (11.39)$$

If X is an attainable claim, that is, $X = V(T)$, then the arbitrage-free value of X at time $t < T$ is the value of the portfolio

$$V(t) = a(t)S(t) + b(t)\beta(t) = C(S(t), t). \quad (11.40)$$

Thus we have by (11.39)

$$dC(S(t), t) = dV(t) = a(t)dS(t) + b(t)d\beta(t). \quad (11.41)$$

Assume that $C(x, t)$ is smooth enough to apply Itô's formula. Then from the SDE for the stock price (11.30) (from which $d[S, S](t) = \sigma^2 S^2(t)dt$) we have

$$dC(S(t), t) = \frac{\partial C(S(t), t)}{\partial x}dS(t) + \frac{\partial C(S(t), t)}{\partial t}dt + \frac{1}{2} \frac{\partial^2 C(S(t), t)}{\partial x^2}\sigma^2 S^2(t)dt. \quad (11.42)$$

By equating the two expressions above we have

$$\left(a(t) - \frac{\partial C(S(t), t)}{\partial x} \right) dS(t) = \left(\frac{\partial C(S(t), t)}{\partial t} + \frac{1}{2} \frac{\partial^2 C(S(t), t)}{\partial x^2}\sigma^2 S^2(t) \right) dt - b(t)d\beta(t). \quad (11.43)$$

The lhs has a positive quadratic variation unless $a(t) - \frac{\partial C(S(t), t)}{\partial x} = 0$, and the rhs has zero quadratic variation. For them to be equal we must have for all t

$$a(t) = \frac{\partial C}{\partial x}(S(t), t), \quad (11.44)$$

and consequently

$$b(t)d\beta(t) = \left(\frac{\partial C(S(t), t)}{\partial t} + \frac{1}{2} \frac{\partial^2 C(S(t), t)}{\partial x^2}\sigma^2 S^2(t) \right) dt. \quad (11.45)$$

By putting the values of $a(t)$ and $b(t)\beta(t)$ into Equation (11.40), taking into account $d\beta(t) = r\beta(t)dt$, and replacing $S(t)$ by x , we obtain PDE (11.38). The replicating portfolio is given by (11.44) and (11.45).

Using the Black–Scholes formula, as the solution of this PDE, the replicating portfolio is obtained from (11.44)

$$a(t) = S(t)\Phi(h(t)), \quad b(t) = K\Phi(h(t) - \sigma\sqrt{T-t}), \quad (11.46)$$

where $h(t)$ is given by (11.36).

Another way to derive the Black–Scholes PDE, useful in other models, is given in Exercise 11.8.

Validity of the Assumptions

The simulated time series of a geometric Brownian motion (with $\sigma = 1$ and various μs (265 points)) look similar to the time series of observed stock prices (daily stock closing prices for the period 5 August 1991–7 August 1992), see Figure 11.1. This shows that the suggested model is capable of producing realistic looking price processes, but, of course, does not prove that this is the correct model. The assumption of normality can be checked by looking at histograms and using formal statistical tests. The histogram of the BHP stock returns points to normality (see Figures 11.2 and 11.3). However, the assumption of constant volatility does not seem to be true, and this issue is addressed next.

Implied Volatility

Denote by C^m the observed market price of an option with strike K and expiration T . The implied volatility $I_t(K, T)$ is defined as the value of the volatility parameter σ in the Black–Scholes formula (11.35) that matches the observed price, namely

$$C^{\text{BS}}(I_t(K, T)) = C_t^m(K, T), \quad (11.47)$$

where C^{BS} is given by (11.35). It has been observed that the implied volatility as a function of strike K (and of term T) is not a constant, and has a graph that looks like a smile. Models with stochastic volatility are able to reproduce the observed behaviour of implied volatilities (see Fouque *et al.* (2000)).

Stochastic Volatility Models

A class of models in which the volatility parameter is not a constant but a stochastic process itself is known as stochastic volatility models. These models were introduced to explain the smile in the implied volatility. An example of such is the Heston (1993) model, in which the stock price $S(t)$ and the volatility $v(t)$ satisfy the following SDEs under the EMM,

$$\begin{aligned} dS(t) &= rS(t)dt + \sqrt{v(t)}S(t)dB(t) \\ dv(t) &= \alpha(\mu - v(t))dt + \delta\sqrt{v(t)} dW(t), \end{aligned} \quad (11.48)$$




Fig. 11.1: Simulated and real stock prices.




Fig. 11.2: Lognormal $\text{LN}(0,1)$ and normal $\text{N}(0,1)$ densities.




Fig. 11.3: Histogram of BHP daily returns over 30 days.

where the Brownian motions B and W are correlated. Stochastic volatility models are incomplete (see Example 8.26). Therefore a replicating portfolio involves a stock and another option. Pricing by using an EMM or replicating by a portfolio with a stock and another option leads to a PDE for the price of an option by using the Feynman–Kac formula with a two-dimensional diffusion process (see Fouque *et al.* (2000)). Heston (1993) derived a PDE and its solution for the transform of the option price. The price itself is obtained by the inversion of the transform.

11.5 Change of Numeraire

A numeraire is the asset in which values of other assets are measured. The no-arbitrage arguments imply that absence of arbitrage opportunities can be stated in terms of the existence of the equivalent martingale probability measure, EMM (risk-neutral measure), that makes the price process measured in a chosen numeraire into a martingale. If the savings account $\beta(t) = e^{rt}$ is a numeraire, then the price $S(t)$ is expressed in units of the savings account, and Q makes $Z(t) = S(t)/e^{rt}$ into a martingale. However, we can choose the stock price to be the numeraire, then the pricing probability measure, EMM Q_1 , is the one that makes $\beta(t)/S(t)$ into a martingale. Change of numeraire is used for currency options and interest rate options.

The price of an attainable claim paying X at T , $C = E_Q(\frac{X}{\beta(T)})$ can be calculated also as $C = E_{Q_1}(\frac{X}{S(T)})$. The way to change measures and express the prices of claims under different numeraires are given by the next results. Note that both assets $S(t)$ and $\beta(t)$ can be stochastic, the only requirement is that $S(t)/\beta(t)$, $0 \leq t \leq T$ is a positive martingale.

Theorem 11.17 (Change of Numeraire): *Let $S(t)/\beta(t)$, $0 \leq t \leq T$ be a positive Q -martingale. Define Q_1 by*

$$\frac{dQ_1}{dQ} = \Lambda(T) = \frac{S(T)/S(0)}{\beta(T)/\beta(0)}. \quad (11.49)$$

Then under Q_1 , $\beta(t)/S(t)$ is a martingale. Moreover, the price of an attainable claim X at time t is related under the different numeraires by the formula

$$C(t) = E_Q \left(\frac{\beta(t)}{\beta(T)} X | \mathcal{F}_t \right) = E_{Q_1} \left(\frac{S(t)}{S(T)} X | \mathcal{F}_t \right). \quad (11.50)$$

PROOF. $\Lambda(t) = \frac{S(t)/S(0)}{\beta(t)/\beta(0)}$ is a positive Q -martingale with $E_Q(\Lambda(T)) = 1$. Therefore by Theorem 10.12

$$\frac{1}{\Lambda(t)} = \frac{\beta(t)/\beta(0)}{S(t)/S(0)} \text{ is a } Q_1\text{-martingale.} \quad (11.51)$$

According to the general Bayes formula Theorem 10.10 (10.23)

$$\begin{aligned} E_Q\left(\frac{\beta(t)}{\beta(T)}X|\mathcal{F}_t\right) &= \frac{E_{Q_1}\left(\frac{1}{\Lambda(T)}\frac{\beta(t)}{\beta(T)}X|\mathcal{F}_t\right)}{E_{Q_1}\left(\frac{1}{\Lambda(T)}|\mathcal{F}_t\right)} \\ &= \frac{S(0)E_{Q_1}\left(\frac{\beta(t)}{S(T)}X|\mathcal{F}_t\right)}{\beta(0)\frac{1}{\Lambda(t)}} = E_{Q_1}\left(\frac{S(t)}{S(T)}X|\mathcal{F}_t\right). \end{aligned}$$

□

A General Option Pricing Formula

As a corollary we obtain the price of a call option in a general setting.

$$E_Q\left(\frac{1}{\beta(T)}(S(T) - K)^+\right) = E_Q\left(\frac{S(T)}{\beta(T)}I(S(T) > K)\right) - E_Q\left(\frac{K}{\beta(T)}I(S(T) > K)\right).$$

The first term is evaluated by changing the numeraire,

$$E_Q\left(\frac{S(T)}{\beta(T)}I(S(T) > K)\right) = E_{Q_1}\left(\frac{1}{\Lambda_T}\frac{S(T)}{\beta(T)}I(S(T) > K)\right) = \frac{S(0)}{\beta(0)}Q_1(S(T) > K).$$

The second term is $KQ(S(T) > K)/\beta(T)$, when $\beta(t)$ is deterministic. Thus

$$C = \frac{S(0)}{\beta(0)}Q_1(S(T) > K) - \frac{K}{\beta(T)}Q(S(T) > K). \quad (11.52)$$

This is a generalization of the Black–Scholes formula (Geman *et al.* (1995), Björk (1998), and Klebaner (2002)). One can verify that the Black–Scholes formula is also obtained when the stock is used as the numeraire (Exercise 11.12).

If $\beta(t)$ is stochastic, then by using a T -bond as a numeraire,

$$E_Q\left(\frac{K}{\beta(T)}I(S(T) > K)\right) = P(0, T)KQ_T(S(T) > K),$$

where $P(0, T) = E_Q(1/\beta(T))$, $\Lambda(T) = \frac{1}{P(0, T)\beta(T)}$. To evaluate expectations above, we need to find the distribution under the new measure. This can be done by using the SDE under Q_1 .

SDEs Under a Change of Numeraire

Let $S(t)$ and $\beta(t)$ be positive processes. Let Q and Q_1 be respectively the equivalent martingale measures when $\beta(t)$ and $S(t)$ are numeraires so that $S(t)/\beta(t)$ is a Q -martingale, and $\beta(t)/S(t)$ a Q_1 -martingale. Suppose that under Q

$$\begin{aligned} dS(t) &= \mu_S(t)dt + \sigma_S(t)dB(t), \\ d\beta(t) &= \mu_\beta(t)dt + \sigma_\beta(t)dB(t), \end{aligned}$$

where B is a Q -Brownian motion, and coefficients are adapted processes. Let $X(t)$ have the SDEs under Q and Q_1 , with a Q_1 -Brownian motion B^1

$$\begin{aligned} dX(t) &= \mu_0(X(t))dt + \sigma(X(t))dB(t), \\ dX(t) &= \mu_1(X(t))dt + \sigma(X(t))dB^1(t). \end{aligned}$$

Since these measures are equivalent the diffusion coefficient for X is the same, and the drift coefficients are related by

Theorem 11.18:

$$\mu_1(X(t)) = \mu_0(X(t)) + \sigma(X(t)) \left(\frac{\sigma_S(t)}{S(t)} - \frac{\sigma_\beta(t)}{\beta(t)} \right). \quad (11.53)$$

PROOF. We know from Chapter 10.3, that Q_1 and Q are related by

$$\Lambda(T) = \frac{dQ_1}{dQ} = \mathcal{E} \left(\int_0^T H(t) dB(t) \right) (T),$$

with $H(t) = \frac{\mu_1(X(t)) - \mu_0(X(t))}{\sigma(X(t))}$. On the other hand, by Theorem 11.17 $\frac{dQ_1}{dQ} = \Lambda(T) = \frac{S(T)/S(0)}{\beta(T)/\beta(0)}$. Moreover, $\Lambda(t) = E_Q(\Lambda(T)|\mathcal{F}_t) = \frac{S(t)/S(0)}{\beta(t)/\beta(0)}$ by the martingale property of (S/β) . Using the exponential SDE,

$$d\Lambda(t) = \Lambda(t)H(t)dB(t) = d \left(\frac{S(t)/S(0)}{\beta(t)/\beta(0)} \right),$$

it follows that

$$\frac{\mu_1(X(t)) - \mu_0(X(t))}{\sigma(X(t))} dB(t) = \frac{\beta(t)}{S(t)} d \left(\frac{S(t)}{\beta(t)} \right).$$

Using the SDEs for S and β , and that S/β has no dt term, the result follows. \square

Corollary 11.19: *The SDE for $S(t)$ under Q_1 , when it is a numeraire, is*

$$dS(t) = \left(\mu_S(t) + \frac{\sigma_S^2(t)}{S(t)} - \frac{\sigma_S(t)\sigma_\beta(t)}{\beta(t)} \right) dt + \sigma_S(t)dB^1(t), \quad (11.54)$$

where $B^1(t)$ is a Q_1 -martingale. Moreover, in the Black–Scholes model when $\mu_S = \mu S$, and $\sigma_S = \sigma S$, the new drift coefficient is $(\mu + \sigma^2)S$.

For other results on change of numeraire see Geman *et al.* (1995).

11.6 Currency (FX) Options

Currency options involve at least two types of market, domestic and foreign. For simplicity's sake we consider just one foreign market. For details on arbitrage theory in foreign market derivatives see Musiela and Rutkowski (1998).

The foreign and domestic interest rates in riskless accounts are denoted by r_f and r_d (subscripts f and d denoting foreign and domestic respectively). The EMMs exist in both markets, Q_f and Q_d . Let $U(t)$ denote the value of a foreign asset in foreign currency at time t , say the Japanese Yen (JPY). Assume that $U(t)$ evolves according to the Black–Scholes model, and write its equation under the EMM (risk-neutral) measure,

$$dU(t) = r_f U(t)dt + \sigma_U U(t)dB_f(t), \quad (11.55)$$

where B_f is a Q_f -Brownian motion. A similar equation holds for assets in the domestic market under its EMM Q_d . The price process discounted by $e^{r_d t}$ in the domestic market is a Q_d -martingale, and an option on an asset in the domestic market which pays $C(T)$ (in domestic currency) at T has its time t price

$$C(t) = e^{-r_d(T-t)} \mathbb{E}_{Q_d}(C(T)|\mathcal{F}_t).$$

A link between the two markets is provided by the exchange rate. Take $X(t)$ to be the price in the domestic currency of one unit of foreign currency at time t . Note that $X(t)$ itself is not an asset, but it gives the price of a foreign asset in domestic currency when multiplied by the price of the foreign asset. We assume that $X(t)$ follows the Black–Scholes SDE under the domestic EMM Q_d

$$dX(t) = \mu_X X(t)dt + \sigma_X X(t)dB_d(t),$$

and show that μ_X must be $r_d - r_f$. Take the foreign savings account $e^{r_f t}$, its value in domestic currency is $X(t)e^{r_f t}$. Thus $e^{-r_d t}X(t)e^{r_f t}$ should be a Q_d -martingale, as a discounted price process in the domestic market. However, from (11.31)

$$e^{-r_d t}X(t)e^{r_f t} = X(0)e^{(r_f - r_d + \mu_X - \frac{1}{2}\sigma_X^2)t + \sigma_X B_d(t)},$$

implying $\mu_X = r_d - r_f$. Thus the equation for $X(t)$ under the domestic EMM Q_d is

$$dX(t) = (r_d - r_f)X(t)dt + \sigma_X X(t)dB_d(t), \quad (11.56)$$

with a Q_d -Brownian motion $B_d(t)$. In general, the Brownian motions B_d and B_f are correlated, i.e. for some $|\rho| < 1$, $B_f(t) = \rho B_d(t) + \sqrt{1 - \rho^2}W(t)$, with independent Brownian motions B_d and W . $d[B_d, B_f](t) = \rho dt$. (The model can also be set up by using two-dimensional independent Brownian motions, a (column) vector Q_d -Brownian motion $[B_d, W]$, and a matrix σ_U , the row vector $\sigma_U = [\rho, \sqrt{1 - \rho^2}]$.)

The value of the foreign asset in domestic currency at time t is given by $U(t)X(t) = \tilde{U}(t)$. Therefore $e^{-r_d t}\tilde{U}(t)$, as a discounted price process, is a Q_d -martingale. One can easily see from (11.55) and (11.56) that

$$e^{-r_d t}\tilde{U}(t) = X(0)U(0)e^{-\frac{1}{2}(\sigma_U^2 + \sigma_X^2)t + \sigma_U B_f(t) + \sigma_X B_d(t)}.$$

For it to be a Q_d -martingale it must hold that $dB_f(t) + \rho\sigma_X dt = d\tilde{B}_d(t)$ is a Q_d -Brownian motion (correlated with B_d). This is accomplished by Girsanov's theorem, and the SDE for $U(t)$ under the domestic EMM Q_d becomes

$$dU(t) = (r_f - \rho\sigma_U\sigma_X)U(t)dt + \sigma_U U(t)d\tilde{B}_d(t). \quad (11.57)$$

Now it is also easy to see that the process $\tilde{U}(t)$ (the foreign asset in domestic currency) under the domestic EMM Q_d has the SDE

$$d\tilde{U}(t) = r_d\tilde{U}(t)dt + (\sigma_U d\tilde{B}_d(t) + \sigma_X dB_d(t))\tilde{U}(t). \quad (11.58)$$

Since

$$\sigma_U d\tilde{B}_d(t) + \sigma_X dB_d(t) = \bar{\sigma} d\tilde{W}_d(t),$$

for some Q_d -Brownian motion $\tilde{W}_d(t)$, with

$$\bar{\sigma}^2 = \sigma_U^2 + \sigma_X^2 + 2\rho\sigma_U\sigma_X, \quad (11.59)$$

the volatility of $\tilde{U}(t)$ is given by the formula that combines the foreign asset volatility and the exchange rate volatility (11.59).

Options on Foreign Currency

Consider a call option on one unit of foreign currency, say on JPY, with the strike price K_d in Australian Dollars (AUD). Its payoff at T in AUD is $(X(T) - K_d)^+$. Thus its price at time t is given by

$$C(t) = e^{-r_d(T-t)} \mathbb{E}_{Q_d}((X(T) - K_d)^+ | \mathcal{F}_t).$$

The conditional Q_d -distribution of $X(T)$, given $X(t)$ is lognormal, by using (11.56) and standard calculations, give the Black–Scholes currency formula (Garman and Kohlhagen (1983))

$$C(t) = X(t)e^{-r_f(T-t)}\Phi(h(t)) - K_d e^{-r_d(T-t)}\Phi(h(t) - \sigma_X \sqrt{T-t}),$$

$$h(t) = \frac{\ln \frac{X(t)}{K_d} + (r_d - r_f + \sigma_S^2/2)(T-t)}{\sigma_X \sqrt{T-t}}.$$

Other options on currency are priced similarly by using Equation (11.56).

Options on Foreign Assets Struck in Foreign Currency

Consider options on a foreign asset denominated in the domestic currency. A call option pays at time T the amount $X(T)(U(T) - K_f)^+$. Since the amount is specified in the domestic currency the pricing should be done under the domestic EMM Q_d . The following argument achieves the result. In the foreign market options are priced by the Black–Scholes formula. The price in domestic currency is obtained by multiplying by $X(t)$. A call option is priced by

$$C(t) = U(t) \left(\Phi(h(t)) - K_f e^{-r_f(T-t)} \Phi \left(h(t) - \sigma_U \sqrt{T-t} \right) \right),$$

where

$$h(t) = \frac{\ln \frac{U(t)}{K_f} + (r_f + \frac{1}{2}\sigma_U^2)(T-t)}{\sigma_U \sqrt{T-t}}.$$

Calculations of the price under Q_d are left as an exercise in change of numeraire, Exercise 11.13.

Options on Foreign Assets Struck in Domestic Currency

Consider a call option on the foreign asset $U(t)$ with strike price K_d in AUD. Its payoff at T in AUD is $(X(T)U(T) - K_d)^+ = (\tilde{U}(T) - K_d)^+$. Thus its price at time t is given by

$$C(t) = e^{-r_d(T-t)} \mathbb{E}_{Q_d}((\tilde{U}(T) - K_d)^+ | \mathcal{F}_t).$$

The conditional distribution of $\tilde{U}(T)$ given $\tilde{U}(t)$ is lognormal, by Equation (11.58), with volatility given by $\bar{\sigma}^2 = \sigma_U^2 + \sigma_X^2 + 2\rho\sigma_U\sigma_X$ from (11.59). Thus

$$C(t) = \tilde{U}(t)\Phi(h(t)) - K_d e^{-r_d(T-t)} \Phi(h(t) - \bar{\sigma}\sqrt{T-t}),$$

$$h(t) = \frac{\ln \frac{\tilde{U}(t)}{K_d} + (r_d + \frac{\bar{\sigma}^2}{2})(T-t)}{\bar{\sigma}\sqrt{T-t}}.$$

Guaranteed Exchanged Rate (Quanto) Options

A quanto option pays in domestic currency the foreign payoff converted at a predetermined rate. For example, a quanto call pays at time T the amount $X(0)(U(T) - K_f)^+$ (K_f is in foreign currency). Thus its time t value is given by

$$C(t) = X(0)e^{-r_d(T-t)} \mathbb{E}_{Q_d}((U(T) - K_f)^+ | \mathcal{F}_t).$$

The conditional distribution of $U(T)$ given $U(t)$ is lognormal, by Equation (11.57), with volatility σ_U . Standard calculations give

$$C(t) = X(0)e^{-r_d(T-t)} \left(U(t)e^{\delta(T-t)} \Phi(h(t)) - K_f \Phi(h(t) - \sigma_U \sqrt{T-t}) \right),$$

where $\delta = r_f - \rho\sigma_U\sigma_X$ and $h(t) = \frac{\ln \frac{U(t)}{K_f} + (\delta + \frac{\sigma_U^2}{2})(T-t)}{\sigma_U \sqrt{T-t}}$.

11.7 Asian, Lookback, and Barrier Options

Asian Options

We assume that $S(t)$ satisfies the Black–Scholes SDE, which we write under the EMM Q

$$dS(t) = rS(t)dt + \sigma S(t)dB(t).$$

Asian options pay at time T , the amount $C(T)$ given by $(\frac{1}{T} \int_0^T S(u)du - K)^+$ (fixed strike K) or $(\frac{1}{T} \int_0^T S(t)dt - KS(T))^+$ (floating strike $KS(T)$). Both kinds involve the integral average of the stock price $\bar{S} = \frac{1}{T} \int_0^T S(u)du$. The average of lognormals is not an analytically tractable distribution, therefore direct calculations do not lead to a closed form solution for the price of the option. Pricing by using PDEs was done by Rogers and Shi (1995), and Vecer (2002). We present Vecer’s approach, which rests on the idea that it is possible to replicate the integral average of the stock \bar{S} by a self-financing portfolio. Let

$$\begin{aligned} a(t) &= \frac{1}{rT}(1 - e^{-r(T-t)}) \quad \text{and} \\ dV(t) &= a(t)dS(t) + r(V(t) - a(t)S(t))dt \\ &= rV(t)dt + a(t)(dS(t) - rS(t)dt), \end{aligned} \tag{11.60}$$

with $V(0) = a(0)S(0) = \frac{1}{rT}(1 - e^{-rT})S(0)$. It is easy to see that $V(t)$ is a self-financing portfolio (Exercise 11.14). Solving the SDE (11.60) (by looking at $d(V(t)e^{-rt})$) for $V(t)$ and using integration by parts we obtain

$$\begin{aligned} V(T) &= e^{rT}V(0) + a(T)S(T) - e^{rT}a(0)S(0) - \int_0^T e^{r(T-t)}S(t)da(t) \\ &= \frac{1}{T} \int_0^T S(t)dt, \end{aligned} \tag{11.61}$$

because of

$$\begin{aligned} d(e^{r(T-t)}S(t)a(t)) &= e^{r(T-t)}a(t)dS(t) - re^{r(T-t)}a(t)S(t)dt \\ &\quad + e^{r(T-t)}S(t)da(t), \end{aligned}$$

and

$$a(T)S(T) - e^{rT}a(0)S(0) = \int_0^T e^{r(T-t)}a(t)(dS(t) - rS(t)dt) + \int_0^T e^{r(T-t)}S(t)da(t).$$

The self-financing portfolio $V(t)$ consists of $a(t)$ shares and $b(t) = \frac{e^{-rT}}{T} \int_0^t S(u)du$ cash and has the value \bar{S} at time T .

Consider next pricing the option with the payoff $(\bar{S} - K_1 S(T) - K_2)^+$, which encompasses both the fixed and the floating kinds (by taking one of the K 's as 0). In order to replicate such an option hold at time t $a(t)$ of the stock, start with initial wealth $V(0) = a(0)S(0) - e^{-rT}K_2$, and follow the self-financing strategy (11.60). The terminal value of this portfolio is $V(T) = \bar{S} - K_2$. The payoff of the option is

$$(\bar{S} - K_1 S(T) - K_2)^+ = (V(T) - K_1 S(T))^+.$$

Thus the price of the option at time t is

$$C(t) = e^{-r(T-t)} \mathbb{E}_Q(V(T) - K_1 S(T))^+ | \mathcal{F}_t.$$

Let $Z(t) = V(t)/S(t)$. Then proceeding,

$$C(t) = e^{-r(T-t)} \mathbb{E}_Q(S(T)(Z(T) - K_1)^+ | \mathcal{F}_t).$$

The multiplier $S(T)$ inside is dealt with by the change of numeraire, and the remaining expectation is that of a diffusion $Z(t)$, which satisfies the backward PDE. Using $S(t)$ as a numeraire (Theorem 11.17, (11.50)) we have

$$C(t) = S(t) \mathbb{E}_{Q_1}((Z_T - K_1)^+ | \mathcal{F}_t). \quad (11.62)$$

According to Ito's formula the SDE for $Z(t)$ under Q , when e^{rt} is a numeraire, is

$$dZ(t) = \sigma^2(Z(t) - a(t))dt + \sigma(a(t) - Z(t))dB(t).$$

According to Theorem 11.18 the SDE for $Z(t)$ under the EMM Q_1 , when $S(t)$ is a numeraire, is

$$dZ(t) = -\sigma(Z(t) - a(t))dB^1(t),$$

where $B^1(t)$ is a Q_1 -Brownian motion, $(B^1(t) = B(t) - \sigma t)$. Now we can write a PDE for the conditional expectation in (11.62).

$$\mathbb{E}_{Q_1}((Z(T) - K_1)^+ | \mathcal{F}_t) = \mathbb{E}_{Q_1}((Z(T) - K_1)^+ | Z(t)),$$

by the Markov property. Hence

$$u(x, t) = \mathbb{E}_{Q_1}((Z(T) - K_1)^+ | Z(t) = x)$$

satisfies the PDE (see Theorem 6.6)

$$\frac{\sigma^2}{2}(x - a(t))^2 \frac{\partial^2 u}{\partial x^2} + \frac{\partial u}{\partial t} = 0,$$

subject to the boundary condition $u(x, T) = (x - K_1)^+$. Finally, the price of the option at time zero is given by using (11.62)

$$V(0, S(0), K_1, K_2) = S(0)u(Z(0), 0),$$

with $Z(0) = V(0)/S(0) = \frac{1}{rT}(1 - e^{-rT}) - e^{-rT}K_2/S(0)$.

Lookback Options

A lookback call pays $X = S(T) - S_*$ and a lookback put $X = S^* - S(T)$, where S_* and S^* denote the smallest and the largest values of stock on $[0, T]$. The price of a lookback put is given by

$$C = e^{-rT} \mathbb{E}_Q(S^* - S(T)) = e^{-rT} \mathbb{E}_Q(S^*) - e^{-rT} \mathbb{E}_Q(S(T)), \quad (11.63)$$

where \mathbb{Q} is the martingale probability measure.

Since $S(t)e^{-rt}$ is a \mathbb{Q} -martingale, $e^{-rT} \mathbb{E}_Q(S(T)) = S(0)$. In order to find $\mathbb{E}_Q(S^*)$, the \mathbb{Q} -distribution of S^* is needed. The equation for $S(t)$ is given by $S(t) = S(0)e^{(r-\frac{1}{2}\sigma^2)t+\sigma B(t)}$ with a \mathbb{Q} -Brownian motion $B(t)$ (see (11.31)). Clearly, S^* satisfies

$$S^* = S(0)e^{((r-\frac{1}{2}\sigma^2)t+\sigma B(t))^*}.$$

The distribution of the maximum of a Brownian motion with drift is found by a change of measure (see Exercise 10.6 and Example 10.4 (10.38)). According to Girsanov's theorem $(r - \frac{1}{2}\sigma^2)t + \sigma B(t) = \sigma W(t)$, for a \mathbb{Q}_1 -Brownian motion $W(t)$, with $\frac{d\mathbb{Q}}{d\mathbb{Q}_1}(W_{[0,T]}) = e^{cW(T)-\frac{1}{2}c^2T}$, $c = (r - \frac{1}{2}\sigma^2)/\sigma$. Clearly, $S^* = S(0)e^{\sigma W^*}$. The distribution of W^* under \mathbb{Q}_1 is the distribution of the maximum of Brownian motion, and its distribution under \mathbb{Q} is obtained as in Example 10.4 (see (10.38)). Therefore

$$e^{-rT} \mathbb{E}_Q(S^*) = e^{-rT} S(0) \int_{-\infty}^{\infty} e^{\sigma y} f_{W^*}(y) dy, \quad (11.64)$$

where $f_{W^*}(y)$ is obtained from (10.38) (see also Exercise 11.15). Lookback calls are priced similarly. The price of a lookback call is given by $S(0)$ times

$$(1 + \frac{\sigma^2}{2r}) \Phi \left(\frac{\ln(K/S(0)) + \frac{\sigma^2 T}{2}}{\sigma \sqrt{T}} \right) - (1 - \frac{\sigma^2}{2r}) e^{-rT} \Phi \left(\frac{\ln(K/S(0)) - \frac{\sigma^2 T}{2}}{\sigma \sqrt{T}} \right) - \frac{\sigma^2}{2r}. \quad (11.65)$$

Barrier Options

Examples of barrier options are down-and-out calls or down-and-in calls. There are also corresponding put options. A down-and-out call gets knocked out when the price falls below a certain level H , it has the payoff $(S(T) - K)^+ I(S_* \geq H)$, $S(0) > H$, $K > H$. A down-and-in call has the payoff $(S(T) - K)^+ I(S_* \leq H)$. In order to price these options the joint distribution of $S(T)$ and S_* is needed under \mathbb{Q} . For example, the price of a down-and-in call is given by

$$C = e^{-rT} \mathbb{E}_Q((S(T) - K)^+ I(S_* \leq H))$$

$$= e^{-rT} \int_{\frac{\ln(K/S(0))}{\sigma}}^{\infty} \int_{-\infty}^{\frac{\ln(H/S(0))}{\sigma}} (S(0)e^{\sigma x} - K) g(x, y) dx dy,$$

where $g(x, y)$ is the probability density of $(W(T), W_*(T))$ under the martingale measure Q . It is found by changing the measure as described above and in Example 10.4 (see also Exercise 11.16). Double barrier options have payoffs depending on $S(T)$, S_* , and S^* . An example is a call which pays only if the price never goes below a certain level H_1 or above a certain level H_2 , with the payoff

$$X = (S(T) - K)^+ I(H_1 < S_* < S^* < H_2).$$

The joint distribution of Brownian motion and its minimum and maximum is given by Theorem 3.23. By using the change of measure described above, the joint distribution of $S(T)$, S_* , S^* can be found and double barrier options can be priced. Calculations are based on the following result.

Let $W_\mu(t) = \mu t + B(t)$. Then the probability of hitting barriers a and b on the interval $[0, t]$, given the values at the end points $W_\mu(0) = x$ and $W_\mu(t) = y$

$$P(a < \min_{0 \leq S \leq t} W_\mu(S) \leq b | W_\mu(0) = x, W_\mu(t) = y),$$

does not depend on μ , and is given by (e.g. Borodin and Salminen (1996))

$$\begin{aligned} P(a, b, x, y, t) = & \exp\left\{ \frac{(y-x)^2}{2t} \right\} \times \sum_{n=-\infty}^{\infty} \left(\exp\left\{ \frac{(y-x+2n(b-a))^2}{-2t} \right\} \right. \\ & \left. - \exp\left\{ \frac{(y-x+2n(b-a)+2(x-a))^2}{-2t} \right\} \right) \end{aligned} \quad (11.66)$$

if $a \leq x \leq b$, $a \leq y \leq b$, and zero otherwise. This formula becomes simpler when one of the barriers is infinite, namely if $a = -\infty$ (single high barrier) we get

$$P(-\infty, b, x, y, t) = 1 - \exp\left\{ -2 \frac{(b-x)(b-y)}{t} \right\} \quad (11.67)$$

if $x \leq b$, $y \leq b$, and zero otherwise. If $b = \infty$ (single low barrier) we get

$$P(a, \infty, x, y, t) = 1 - \exp\left\{ -2 \frac{(a-x)(a-y)}{t} \right\} \quad (11.68)$$

if $a \leq x$, $a \leq y$, and zero otherwise.

11.8 Exercises

Exercise 11.1: (Payoff functions and diagrams)

Graph the following payoffs:

1. A *straddle* consists of buying a call and a put with the same exercise price and expiration date.
2. A *butterfly spread* consists of buying two calls with exercise prices K_1 and K_3 and selling a call with exercise price K_2 , $K_1 < K_2 < K_3$.

Exercise 11.2: (Binomial pricing model)

1. Give the probability space for the binomial model.
2. Show that the stock price can be expressed as $S(t+1) = S(t)\xi_{t+1}$, where ξ_{t+1} is the return in period $t+1$, $t = 0, 1, \dots, n-1$, and the variables ξ_n are independent and identically distributed with values u and d .
3. Show that the binomial model does not admit arbitrage if, and only if, $d < r < u$.
4. Describe the arbitrage-free probability Q , and show that the discounted stock price S_t/r^t , $t = 0, 1, \dots, n$ is a Q -martingale.
5. Show that this model is complete.
6. Show that if the price of an option is given by (11.7), then arbitrage strategies do not exist.

Exercise 11.3: Verify that in the model given in Example 11.7 any attainable claim has the same price under any of the martingale measures. Give an example of an unattainable claim X and show that $E_Q(X)$ is different for different martingale measures Q .

Exercise 11.4: Show that if Q is equivalent to P and $X \geq 0$, then $E_P(X) > 0$ implies $E_Q(X) > 0$ and vice versa.

Exercise 11.5: (Pricing in incomplete markets)

1. Show that if $M(t)$, $0 \leq t \leq T$, is a martingale under two different probability measures Q and P , then for $s < t$ $E_Q(M(t)|\mathcal{F}_s) = E_P(M(t)|\mathcal{F}_s)$ almost surely. If in addition $M(0)$ is non-random, then $E_P M(t) = E_Q M(t)$.
2. Show that the price of an attainable claim X , $C(t) = \beta(t)E_Q(X/\beta(T)|\mathcal{F}_t)$ is the same for all martingale measures.

Exercise 11.6: (Non-completeness in mixed models)

In this exercise the price of an asset is modelled as a sum of a diffusion and a jump process. Take for example $X(t) = W(t) + N(t)$, with Brownian motion W and Poisson process N . Give at least two equivalent probability measures Q_1 and Q_2 such that X is a Q_i -martingale, $i = 1, 2$ (see Exercise 10.13).

Exercise 11.7: Give the Radon–Nikodym derivative Λ in the change to the EMM Q in the Black–Scholes model.

Exercise 11.8: A way to derive a PDE for the option price is based on the fact that $C(t)e^{-rt} = V(t)e^{-rt}$ is a Q -local martingale. Obtain the Black–Scholes PDE for the price of the option using the Black–Scholes model for the stock price. Hint: expand $d(C(S(t), t)e^{-rt})$ and equate the coefficient of dt to zero.

Exercise 11.9: Derive the PDE for the price of the option in Heston’s model.

Exercise 11.10: Show that the expected return on stock under the martingale probability measure Q is the same as on the riskless asset. This is the reason why the martingale measure Q is also called the “risk-neutral” probability measure.

Exercise 11.11: Assume $S(t)$ evolves according to the Black–Scholes model. Show that under the EMM Q_1 , when $S(t)$ is a numeraire, $d(e^{rt}/S(t)) = \sigma(e^{rt}/S(t))dW_t$, where $W(t) = B(t) - \sigma t$ is a Q_1 -Brownian motion. Give the likelihood dQ_1/dQ . Give the SDE for $S(t)$ under Q_1 .

Exercise 11.12: Derive the Black–Scholes formula by using the stock price as the numeraire.

Exercise 11.13: A call option on an asset in the foreign market pays at time T , $S(T)(U(T) - K)^+$ in the domestic currency, and its time t price $C(t) = e^{-r_d(T-t)}\mathbb{E}_{Q_d}(S(T)(U(T) - K)^+ | \mathcal{F}_t)$. Taking numeraire based on the Q_d -martingale $S(t)e^{-(r_d - r_f)t}$ obtain the formula for $C(t)$.

Exercise 11.14: Let $V(t) = a(t)S(t) + b(t)e^{rt}$ be a portfolio, $0 \leq t \leq T$. Show that it is self-financing if, and only if, $dV(t) = a(t)dS(t) + r(V(t) - a(t)S(t))dt$.

Exercise 11.15: Derive the price of a lookback call.

Exercise 11.16: Show that the price of a down-and-in call is given by

$$e^{-rT} \left(\frac{H}{S(0)} \right)^{2r/\sigma^2 - 1} \left(F\Phi \left(\frac{\ln(F/K) + \frac{\sigma^2 T}{2}}{\sigma\sqrt{T}} \right) - K\Phi \left(\frac{\ln(F/K) - \frac{\sigma^2 T}{2}}{\sigma\sqrt{T}} \right) \right),$$

where $F = e^{rT}H^2/S(0)$.

Exercise 11.17: Assume that $S(T)/S$ does not depend on S , where $S(T)$ is the price of stock at T and $S = S(0)$. Let T be the exercise time and K the exercise price of the call option. Show that the price of this option satisfies the following PDE:

$$C = S \frac{\partial C}{\partial S} + K \frac{\partial C}{\partial K}.$$

You may assume all the necessary differentiability. Hence show that the delta of the option $\frac{\partial C}{\partial S}$ in the Black–Scholes model is given by $\Phi(h(t))$ with $h(t)$ given by (11.36).

Chapter 12

Applications in Finance: Bonds, Rates, and Options

Money invested for different terms T yield a different return corresponding to the rate of interest $R(T)$. This function is called the yield curve, or the term structure of interest rates. Every day this curve changes, the time t curve denoted by $R(t, T)$. However, the rates are not traded directly, they are derived from prices of bonds traded on the bond market. This leads to the construction of models for bonds and no-arbitrage pricing for bonds and their options. We present the main models used in the literature and in applications, treating in detail the Merton, Vasicek's, Heath–Jarrow–Morton (HJM), and Brace–Gatarek–Musiela (BGM) models. In our treatment we concentrate on the main mathematical techniques used in such models without going into the details of their calibration. For this and other details see Brigo and Mercurio (2006).

12.1 Bonds and the Yield Curve

A \$1 bond with maturity T is a contract that guarantees the holder \$1 at T . Sometimes bonds also pay a certain amount, called a coupon, during the life of the bond, but for the theory it suffices to consider only bonds without coupons (zero-coupon bonds). Denote by $P(t, T)$ the price at time t of the bond paying \$1 at T , $P(T, T) = 1$. The yield to maturity of the bond is defined as

$$R(t, T) = -\frac{\ln P(t, T)}{T - t}, \quad (12.1)$$

and as a function in T , is called the yield curve at time t . Assume also that a savings account paying at t instantaneous rate $r(t)$, called the spot

(or short) rate, is available. If \$1 is invested until time t it will result in

$$\beta(t) = e^{\int_0^t r(s)ds}. \quad (12.2)$$

In order to avoid arbitrage between bonds and savings accounts a certain relation must hold between bonds and the spot rate. If there were no uncertainty, then to avoid arbitrage the following relation must hold:

$$P(t, T) = e^{-\int_t^T r(s)ds}, \quad (12.3)$$

since investing either of these amounts at time t results in \$1 at time T . When the rate is random, then $\int_t^T r(s)ds$ is also random and in the future of t , whereas the price $P(t, T)$ is known at time t , and the above relation holds only “on average” (Equation (12.5)).

We assume a probability model with a filtration $\mathbb{F} = \{\mathcal{F}_t\}$, $0 \leq t \leq T^*$, and adapted processes $P(t, T)$, $t \leq T \leq T^*$, and $\beta(t)$. For the extension of the no-arbitrage theory see Artzner and Delbaen (1989), Lamberton and Lapeyre (1996), Björk (1998), Musiela and Rutkowski (1998), and Shiryaev (1999). In addition to the number of no-arbitrage concepts in continuous time (see Chapter 11.3) the continuum of bond maturities T makes the market model have infinitely many assets and produces further complication. There are different approaches, including finite portfolios, where at each time only finitely many bonds are allowed, and infinite, measure-valued portfolios. In all of the approaches the no-arbitrage condition is formulated with the help of the following assumption.

EMM Assumption

There is a probability Q (called the equivalent martingale measure), equivalent to P (the original “real-world” probability), such that simultaneously for all $T \leq T^*$, the process in t , $P(t, T)/\beta(t)$ is a martingale, $0 \leq t \leq T$.

The martingale property implies that

$$\mathbb{E}_Q\left(\frac{1}{\beta(T)}|\mathcal{F}_t\right) = \mathbb{E}_Q\left(\frac{P(T, T)}{\beta(T)}|\mathcal{F}_t\right) = \frac{P(t, T)}{\beta(t)}, \quad (12.4)$$

where \mathcal{F}_t denotes the information available from the bond prices up to time t . Since $P(T, T) = 1$ we obtain the expression for the price of the bond

$$P(t, T) = \mathbb{E}_Q\left(\frac{\beta(t)}{\beta(T)}|\mathcal{F}_t\right) = \mathbb{E}_Q\left(e^{-\int_t^T r(s)ds}|\mathcal{F}_t\right). \quad (12.5)$$

It shows that the bond can be seen as a derivative on the short rate.

12.2 Models Adapted to Brownian Motion

Here we derive the SDE for bond $P(t, T)$ under Q starting only with the EMM assumption. The SDE under P is then derived as a consequence of the predictable representation property. As usual this property points out the existence of certain processes but does not say how to find them.

Consider a probability space with a P -Brownian motion $W(t)$ and its filtration \mathcal{F}_t . Assume that the spot rate process $r(t)$ generates \mathcal{F}_t , and that the bond processes $P(t, T)$, for any $T < T^*$, are adapted. The EMM assumption gives the price of the bond by the Equation (12.5). The martingale

$$\frac{P(t, T)}{\beta(t)} = \mathbb{E}_Q \left(e^{-\int_0^T r(s)ds} \middle| \mathcal{F}_t \right) \quad (12.6)$$

is adapted to the Brownian filtration. According to Theorem 10.19 there exists an adapted process $X(t) = \int_0^t \sigma(S, T)dB(S)$, where $B(t)$ is a Q -Brownian motion such that

$$d \left(\frac{P(t, T)}{\beta(t)} \right) = \left(\frac{P(t, T)}{\beta(t)} \right) dX(t) = \sigma(t, T) \left(\frac{P(t, T)}{\beta(t)} \right) dB(t).$$

Opening $d(P/\beta)$ we obtain the SDE for $P(t, T)$ under the EMM Q

$$\frac{dP(t, T)}{P(t, T)} = r(t)dt + \sigma(t, T)dB(t). \quad (12.7)$$

This is the pricing equation for bonds and their options.

Note that the return on the savings account satisfies $d\beta(t)/\beta(t) = r(t)dt$, and the return on the bond has an extra term with a Brownian motion. This makes bonds $P(t, T)$ riskier than the (also random) savings account $\beta(t)$.

We find the SDE for the bond under the original probability measure P next. Since Q is equivalent to the Wiener measure, by Corollary 10.21 there is an adapted process $q(t)$ such that

$$W(t) = B(t) + \int_0^t q(s)ds \quad (12.8)$$

is a P -Brownian motion, with $dQ/dP = e^{\int_0^T q(s)dW(s) - \frac{1}{2} \int_0^T q^2(s)dt}$. Substituting $dB(t) = dW(t) - q(t)dt$ into SDE (12.7) we obtain the SDE under P

$$\frac{dP(t, T)}{P(t, T)} = (r(t) - \sigma(t, T)q(t))dt + \sigma(t, T)dW(t). \quad (12.9)$$

Remark 12.1: It follows from Equation (12.9) that $-q(t)$ is the excess return on the bond above the riskless rate, expressed in standard units; it is known as “the market price of risk” or “risk premium”. The most common assumption is that $q(t) = q$ is a constant.

12.3 Models Based on the Spot Rate

A model for term structure and pricing can be developed from a model for the spot rate. These models are specified under the real world probability measure P , and $r(t)$ is assumed to satisfy

$$dr(t) = m(r(t))dt + \sigma(r(t))dW(t), \quad (12.10)$$

where $W(t)$ is a P -Brownian motion, m and σ are functions of a real variable. The bond price $P(t, T)$ satisfies (12.5)

$$P(t, T) = E_Q\left(e^{-\int_t^T r(s)ds} \middle| \mathcal{F}_t\right).$$

The expectation is under Q , and the model (12.10) is specified under P . Therefore the SDE for $r(t)$ under Q is needed. Note that we could express the above expectation in terms of E_P by

$$E_P\left(e^{-\int_t^T r(s)ds + \int_t^T q(s)dW(s) - \frac{1}{2}\int_t^T q^2(s)ds} \middle| \mathcal{F}_t\right),$$

but this expectation seems to be intractable even in simple models.

We move between P and the EMM Q by using (12.8) expressed as $dB(t) = dW(t) - q(t)dt$. Thus under Q

$$dr(t) = (m(r(t)) + \sigma(r(t))q(t))dt + \sigma(r(t))dB(t). \quad (12.11)$$

The process $r(t)$ is also a diffusion under Q , therefore by the Markov property

$$E_Q\left(e^{-\int_t^T r(s)ds} \middle| \mathcal{F}_t\right) = E_Q\left(e^{-\int_t^T r(s)ds} \middle| r(t)\right).$$

The last expression satisfies a PDE by the Feynman–Kac formula (Theorem 6.8). Fix T , and denote by

$$C(x, t) = E_Q\left(e^{-\int_t^T r(s)ds} \middle| r(t) = x\right),$$

then by (6.22) it satisfies

$$\frac{1}{2}\sigma^2(x)\frac{\partial^2 C}{\partial x^2}(x, t) + (m(x) + \sigma(x)q(t))\frac{\partial C}{\partial x}(x, t) + \frac{\partial C}{\partial t}(x, t) - xC(x, t) = 0, \quad (12.12)$$

with the boundary condition $C(x, T) = 1$. The price of the bond is obtained from this function by

$$P(t, T) = C(r(t), t).$$

A similar PDE with suitable boundary conditions holds for options on bonds.

We list some of the well-known models for the spot rate.
The Merton model:

$$dr(t) = \mu dt + \sigma dW(t). \quad (12.13)$$

The Vasicek model:

$$dr(t) = b(a - r(t))dt + \sigma dW(t). \quad (12.14)$$

The Dothan model:

$$dr(t) = \mu r(t)dt + \sigma dW(t). \quad (12.15)$$

The Cox–Ingersoll–Ross (CIR) model:

$$dr(t) = b(a - r(t))dt + \sigma \sqrt{r(t)}dW(t). \quad (12.16)$$

The Ho–Lee model:

$$dr(t) = \mu(t)dt + \sigma dW(t). \quad (12.17)$$

The Black–Derman–Toy model:

$$dr(t) = \mu(t)r(t)dt + \sigma(t)dW(t). \quad (12.18)$$

The Hull–White model:

$$dr(t) = b(t)(a(t) - r(t))dt + \sigma(t)dW(t). \quad (12.19)$$

The Black–Karasinski model:

$$dr(t) = r(t)(a(t) - b(t) \ln r(t))dt + \sigma(t)r(t)dW(t). \quad (12.20)$$

The functions $m(r)$ and $\sigma(r)$ involve parameters that need to be estimated. They are chosen in such a way that the values of bonds and options agree as closely as possible with the values observed in the market. This process is called calibration, and we do not address it.

In what follows we derive prices of bonds and their options for some models by using probability calculations rather than solving PDEs.

12.4 Merton's Model and Vasicek's Model

Merton's Model

The spot rate in the Merton model satisfies SDE (12.13). Its solution is

$$r(t) = r_0 + \mu t + \sigma W(t). \quad (12.21)$$

The savings account is given by

$$\beta(t) = e^{\int_0^t r(s)ds} = e^{r_0 t + \mu t^2/2} e^{\sigma \int_0^t W_s ds}.$$

Assume the constant price for risk $q(t) = q$, then by (12.11) the SDE for $r(t)$ under Q is

$$dr(t) = (\mu + \sigma q)dt + \sigma dB(t).$$

The price of the bond $P(t, T)$ is given by $P(t, T) = \mathbb{E}_Q(e^{-\int_t^T r(s)ds} | \mathcal{F}_t)$. Since the conditional expectation given \mathcal{F}_t is needed, use the decomposition of $r(s)$ into an \mathcal{F}_t -measurable part, and \mathcal{F}_t -independent part.

$$\begin{aligned} r(s) &= r(t) + (\mu + \sigma q)(s - t) + \sigma(W(s) - W(t)) \\ &= r(t) + (\mu + \sigma q)(s - t) + \sigma \hat{W}(s - t), \end{aligned}$$

with $\hat{W}(s - t)$ independent of \mathcal{F}_t . Then

$$\begin{aligned} P(t, T) &= e^{-r(t)(T-t) - (\mu + \sigma q)(T-t)^2/2} \mathbb{E}_Q(e^{-\sigma \int_t^T \hat{W}(s-t)ds} | \mathcal{F}_t) \\ &= e^{-r(t)(T-t) - (\mu + \sigma q)(T-t)^2/2 + \sigma^2(T-t)^3/6}, \end{aligned} \quad (12.22)$$

where we used that the distribution of the random integral $\int_0^{T-t} \hat{W}(u)du$ is $N(0, (T-t)^3/3)$ (see Example 3.6). The yield curve is given by

$$R(t, T) = -\frac{\ln P(t, T)}{T-t} = r(t) + \frac{1}{2}(\mu + \sigma q)(T-t) - \frac{1}{6}\sigma^2(T-t)^2.$$

Since $r(t)$ has a normal distribution, so does $R(t, T)$, so that $P(t, T)$ is lognormal. The pricing of a call option on the bond in this model is covered in Exercise 12.2.

Note that the yields for different maturities differ by a deterministic quantity. Therefore $R(t, T_1)$ and $R(t, T_2)$ are perfectly correlated. This is taken as a shortcoming of the model.

Vasicek's Model

The spot rate in Vasicek's model satisfies SDE (12.14). Its solution is

$$r(t) = a - e^{-bt}(a - r(0)) + \sigma \int_0^t e^{-b(t-s)} dW(s). \quad (12.23)$$

We derive the solution below, but note that it is easy to check that (12.23) is indeed a solution (see also the Langevin SDE Example 5.6, Equation (5.15)). Writing the SDE (12.14) in the integral form and taking expectations (it is easy to see that the Itô integral has zero mean and

interchanging the integral and the expectation is justified by Fubini's theorem), we have

$$\text{Er}(t) - \text{Er}(0) = \int_0^t b(a - \text{Er}(s))ds. \quad (12.24)$$

Put $h(t) = \text{Er}(t)$. Differentiating we obtain

$$h'(t) = b(a - h(t)).$$

This equation is solved by separating variables. Integrating from 0 to t , and performing the change of variable $u = h(s)$ we obtain

$$\text{Er}(t) = a - e^{-bt}(a - r(0)). \quad (12.25)$$

Now let

$$X(t) = r(t) - \text{Er}(t) = r(t) - h(t). \quad (12.26)$$

$X(t)$ clearly satisfies

$$dX(t) = dr(t) - dh(t) = -bX(t)dt + \sigma dW(t), \quad (12.27)$$

with the initial condition $X(0) = 0$. However, this is the equation for the Ornstein–Uhlenbeck process. According to (5.13), $X(t) = \sigma \int_0^t e^{-b(t-s)} dW(s)$, and (12.23) follows from Equations (12.25) and (12.26).

Make two observations next. First, the long-term mean is a ,

$$\lim_{t \rightarrow \infty} \text{Er}(t) = a.$$

Second, the process $X(t) = r(t) - \text{Er}(t)$ reverts to zero, hence $r(t)$ reverts to its mean: if $r(t)$ is above its mean, then the drift is negative, making $r(t)$ decrease; and if $r(t)$ is below its mean, then the drift is positive, making $r(t)$ increase. Mean reversion is a desirable property in the modelling of rates.

In order to proceed to the calculation of bond prices $P(t, T) = \mathbb{E}_Q(e^{-\int_t^T r(s)ds} | \mathcal{F}_t)$, further assumptions on the market price of risk $q(t)$ are needed. Assume $q(t) = q$ is a constant.

We move between P and the EMM Q by using (12.8), which states that $dB(t) = dW(t) - qdt$. Therefore the equation for $r(t)$ under Q is given by (12.14) with a replaced by

$$a^* = a + \frac{\sigma q}{b}. \quad (12.28)$$

In order to calculate the Q-conditional distribution of $\int_t^T r(s)ds$ given \mathcal{F}_t needed for the bond price, observe that by the Markov property of the solution, for $s > t$ the process starts at $r(t)$ and runs for $s - t$, giving

$$r(s) = a^* - e^{-b(s-t)}(a^* - r(t)) + \sigma e^{-b(s-t)} \int_0^{s-t} e^{bu} dB(u), \quad (12.29)$$

with a Q-Brownian motion $B(u)$, independent of \mathcal{F}_t . Thus $r(s)$ is a Gaussian process since the function in the Itô integral is deterministic (see Theorem 4.11). The conditional distribution of $\int_t^T r(s)ds$ given \mathcal{F}_t is the same as that given $r(t)$ and is a normal distribution. Thus the calculation of the bond price involves the expectation of a lognormal, $E_Q(e^{-\int_t^T r(s)ds}|r(t))$. Thus

$$P(t, T) = e^{-\mu_1 + \sigma_1^2/2}, \quad (12.30)$$

where μ_1 and σ_1^2 are the conditional mean and variance of $\int_t^T r(s)ds$ given $r(t)$. Using (12.29) and calculating directly $h = a^* - h'/b$, we have

$$\begin{aligned} \mu_1 &= E_Q \left(\int_t^T r(s)ds \middle| r(t) \right) = \int_t^T E_Q(r(s)|r(t))ds \\ &= a^*(T-t) - \frac{1}{b}(a^* - r(t))(1 - e^{-b(T-t)}). \end{aligned} \quad (12.31)$$

In order to calculate σ_1^2 use the representation for $r(s)$ conditional on $r(t)$, Equation (12.29)

$$\sigma_1^2 = \text{Cov} \left(\int_t^T r(s)ds, \int_t^T r(u)du \right) = \int_t^T \int_t^T \text{Cov}(r(s), r(u))dsdu.$$

Now it is not hard to calculate, by the formula for the covariance of Itô Gaussian integrals (4.26) or (4.27), that for $s > u$

$$\text{Cov}(r(s), r(u)) = \frac{\sigma^2}{2b} \left(e^{b(u-s)} - e^{-b(u+s-2t)} \right).$$

Putting this expression (and a similar one for when $s < u$) in the double integral, we obtain

$$\sigma_1^2 = \frac{\sigma^2}{b^2} (T-t) - \frac{\sigma^2}{2b^3} (3 - 4e^{-b(T-t)} + e^{-2b(T-t)}).$$

Thus denoting $R(\infty) = a^* - \frac{\sigma^2}{2b^2}$,

$$P(t, T) = e^{\frac{1}{b}(1-e^{-b(T-t)})(R(\infty)-r(t))-(T-t)R(\infty)-\frac{\sigma^2}{4b^3}(1-e^{-b(T-t)})^2}. \quad (12.32)$$

From the formula (12.32) the yield to maturity is obtained by (12.1). Vasicek (1977) obtained the formula (12.32) by solving the PDE (12.12).

Since the price of bonds is lognormal with known mean and variance, a closed form expression for the price of an option on the bond can be obtained.

12.5 Heath–Jarrow–Morton (HJM) Model

The class of models suggested by Heath, Jarrow, and Morton (1992) is based on modelling the forward rates. These rates are implied by bonds with different maturities. By definition, forward rates $f(t, T)$, $t \leq T \leq T^*$ are defined by the relation

$$P(t, T) = e^{-\int_t^T f(t, u) du}. \quad (12.33)$$

Thus the forward rate $f(t, T)$, $t \leq T$, is the (continuously compounding) rate at time T as seen from time t ,

$$f(t, T) = -\frac{\partial \ln P(t, T)}{\partial T}.$$

The spot rate $r(t) = f(t, t)$. Consequently, the savings account $\beta(t)$ grows according to

$$\beta(t) = e^{\int_0^t f(s, s) ds}. \quad (12.34)$$

The assumption of the HJM model is that for a fixed T , the forward rate $f(t, T)$ is a diffusion in t variable, namely

$$df(t, T) = \alpha(t, T) dt + \sigma(t, T) dW(t), \quad (12.35)$$

where $W(t)$ is P -Brownian motion and processes $\alpha(t, T)$ and $\sigma(t, T)$ are adapted and continuous. $\alpha(t, T)$, $\sigma(t, T)$, and the initial conditions $f(0, T)$, are the parameters of the model.

EMM Assumption

There exists an equivalent martingale probability measure (EMM), $Q \sim P$ such that for all $T \leq T^*$, $\frac{P(t, T)}{\beta(t)}$ is a Q -martingale. Assuming the existence of Q we find equations for the bonds and the rates under Q .

Bonds and Rates Under Q and the No-Arbitrage Condition

The EMM assumption implies that $\alpha(t, T)$ is determined by $\sigma(t, T)$ when SDE for forward rates is considered under Q .

Theorem 12.1: *Assume the forward rates satisfy SDE (12.35), the EMM assumption holds, $\frac{P(t, T)}{\beta(t)}$ is a Q -martingale, and all the conditions on the coefficients of the SDE (12.35) needed for the analysis below. Let*

$$\tau(t, T) = \int_t^T \sigma(t, u) du. \quad (12.36)$$

Then

$$\alpha(t, T) = \sigma(t, T)\tau(t, T). \quad (12.37)$$

Moreover, under the EMM Q the forward rates satisfy the SDE with a Q-Brownian motion B

$$df(t, T) = \sigma(t, T)\tau(t, T)dt + \sigma(t, T)dB(t). \quad (12.38)$$

Conversely, if the forward rates satisfy the SDE (12.38) then $\frac{P(t, T)}{\beta(t)}$ is a Q-local martingale. If the appropriate integrability conditions hold, then it is a Q-martingale.

PROOF. The idea is simple: find $d\left(\frac{P(t, T)}{\beta(t)}\right)$ and equate the coefficient of dt to zero. Let

$$X(t) = \ln P(t, T) = - \int_t^T f(t, u)du. \quad (12.39)$$

According to Itô's formula

$$d\left(\frac{P(t, T)}{\beta(t)}\right) = \frac{P(t, T)}{\beta(t)} \left(dX(t) + \frac{1}{2}d[X, X](t) - r(t)dt \right). \quad (12.40)$$

It is not hard to show, see Example 12.1, that

$$dX(t) = -d\left(\int_t^T f(t, u)du\right) = -A(t, T)dt - \tau(t, T)dW(t), \quad (12.41)$$

where $A(t, T) = -r(t) + \int_t^T \alpha(t, u)du$. Thus

$$d\left(\frac{P(t, T)}{\beta(t)}\right) = \frac{P(t, T)}{\beta(t)} \left(\left(- \int_t^T \alpha(t, u)du \right) dt + \frac{1}{2}\tau^2(t, T)dt - \tau(t, T)dW(t) \right). \quad (12.42)$$

According to Girsanov's theorem

$$dW(t) + \left(\frac{\int_t^T \alpha(t, u)du}{\tau(t, T)} - \frac{1}{2}\tau(t, T) \right) dt = dB(t), \quad (12.43)$$

for a Q-Brownian motion $B(t)$. This gives the SDE for the discounted bond under Q

$$d\left(\frac{P(t, T)}{\beta(t)}\right) = -\frac{P(t, T)}{\beta(t)}\tau(t, T)dB(t). \quad (12.44)$$

Thus if the model is specified under Q, $W(t) = B(t)$, then

$$\int_t^T \alpha(t, u)du = \frac{1}{2}\tau^2(t, T). \quad (12.45)$$

Differentiating in T gives the condition (12.37). The SDE (12.38) follows from (12.37). \square

Corollary 12.2: *The bonds satisfy the following equations under Q for $t \leq T$:*

$$dP(t, T) = P(t, T)(r(t)dt - \tau(t, T)dB(t)), \quad \text{and} \quad (12.46)$$

$$P(t, T) = P(0, T)e^{-\int_0^t \tau(s, T)dB(s) - \frac{1}{2} \int_0^t \tau^2(s, T)ds + \int_0^t r(s)ds}. \quad (12.47)$$

PROOF. Equation (12.44) shows that $\frac{P(t, T)}{\beta(t)}$ is the stochastic exponential of $-\int_0^t \tau(s, T)dB(s)$. Hence

$$\begin{aligned} \frac{P(t, T)}{\beta(t)} &= \frac{P(0, T)}{\beta(0)} \mathcal{E}\left(-\int_0^t \tau(s, T)dB(s)\right) \\ &= \frac{P(0, T)}{\beta(0)} e^{-\int_0^t \tau(s, T)dB(s) - \frac{1}{2} \int_0^t \tau^2(s, T)ds}. \end{aligned} \quad (12.48)$$

Since $\beta(t) = e^{\int_0^t r(s)ds}$ the bond's price is given by (12.47). The SDE (12.46) follows, as the stochastic exponential SDE of $-\int_0^t \tau(s, T)dB(s) + \int_0^t r(s)ds$. \square

Using (12.47) for T_1 and T_2 we obtain for $t \leq T_1 \leq T_2$ by eliminating $\int_0^t r(s)ds$.

Corollary 12.3: *A relation between bonds with different maturities is given by*

$$P(t, T_2) = \frac{P(0, T_2)}{P(0, T_1)} e^{-\int_0^t (\tau(s, T_2) - \tau(s, T_1))dB(s) - \frac{1}{2} \int_0^t (\tau^2(s, T_2) - \tau^2(s, T_1))ds} P(t, T_1). \quad (12.49)$$

Remark 12.2:

1. Equation (12.37) is known as the no-arbitrage or EMM condition.
2. The effect of the change to the EMM Q is in the change of drift in the SDE for the forward rates, from (12.35) to (12.38).
3. The volatility of the bond is $\tau(t, T) = \int_t^T \sigma(t, s)ds$, the integrated forward volatilities by (12.46).
4. The expression (12.47) for $P(t, T)$ includes the Itô integral $\int_0^t \tau(s, T)dB(s)$, which is not observed directly. It can be obtained from $\int_t^T f(t, u)du$ using (12.38) and interchanging integrals. Integrated processes and interchanging integrals can be justified rigorously, see Heath *et al.* (1992), and in greater generality Hamza *et al.* (2005).

5. The vector case of $W(t)$ and $\sigma(t, T)$ in (12.35) gives similar formulae by using notation $\sigma(t, T)dW(t) = \sum_{i=1}^d \sigma_i(t, T)dW_i(T)$, the scalar product, and replacing σ^2 by the norm $|\sigma|^2$, see Exercise 5.9.

Example 12.1: Differentiation of an integral of the form $\int_t^T f(t, s)ds$. We show that (12.41) holds. Introduce $G(u, t) = \int_u^T f(t, s)ds$. We are interested in $dG(t, t)$; it is found by

$$dG(t, t) = \left(\frac{\partial}{\partial u} G(t, t) + \frac{\partial}{\partial t} G(t, t) \right) dt,$$

or

$$d \left(\int_t^T f(t, v)dv \right) = -f(t, t)dt + \int_t^T df(t, v).$$

Now $f(t, t) = r(t)$ and, using the model for $df(t, v)$, we obtain

$$d \left(\int_t^T f(t, v)dv \right) = -r(t)dt + \left(\int_t^T \alpha(t, v)dv \right) dt + \left(\int_t^T \sigma(t, v)dv \right) dW(t).$$

Exchange of the integrals is justified in Exercise 12.3.

Example 12.2: (Ho–Lee model)

Consider the SDE for forward rates (12.38) with the simplest choice of constant volatilities $\sigma(t, T) = \sigma = \text{const}$. Then $\tau(t, T) = \int_t^T \sigma(t, u)du = \sigma(T - t)$. Thus

$$\begin{aligned} df(t, T) &= \sigma^2(T - t)dt + \sigma dB(t), \\ f(t, T) &= f(0, T) + \sigma^2 t(T - \frac{t}{2}) + \sigma B(t), \quad \text{and} \\ r(t) &= f(t, t) = f(0, t) + \sigma^2 \frac{t^2}{2} + \sigma B(t). \end{aligned}$$

They contain the Brownian motion, which is not observed directly. Eliminating it, $\sigma B(t) = r(t) - f(0, t) - \sigma^2 \frac{t^2}{2}$, we have

$$f(t, T) = r(t) + \sigma^2 t(T - t) + f(0, T) - f(0, t).$$

This equation shows that forward rates $f(t, T_1) - f(t, T_2) = f(0, T_1) - f(0, T_2) + \sigma^2 t(T_1 - T_2)$ differ by a deterministic term, therefore they are also perfectly correlated. $r(t)$ and $f(t, T)$ are also perfectly correlated. This seems to contradict observations in financial markets.

The bond in terms of the forward rates (12.33)

$$P(t, T) = e^{- \int_t^T f(t, u)du} = e^{- \int_t^T f(0, u)du - \sigma^2 \int_t^T t(u - \frac{t}{2})du - \sigma B(t)(T - t)}.$$

Using $-\int_t^T f(0, u)du = \ln P(0, T) - \ln P(0, t)$,

$$P(t, T) = \frac{P(0, T)}{P(0, t)} e^{-\sigma^2 t T (T-t)/2 - \sigma B(t)(T-t)}.$$

Eliminating $B(t)$ we obtain the equation for the bond in terms of the spot rate, from which the yield curve is determined

$$P(t, T) = \frac{P(0, T)}{P(0, t)} e^{-(T-t)r(t)-\sigma^2 t(T-t)^2/2+(T-t)f(0,t)}.$$

Figures 12.1 and 12.2 illustrate possible curves produced by the Ho–Lee model.




Fig. 12.1: Forward rates and yield curves.




Fig. 12.2: Bonds $P(t, T_i)$ for two maturities T_1 and T_2 one year apart.

12.6 Forward Measures — Bond as a Numeraire

Options on a Bond

An attainable claim is one that can be replicated by a self-financing portfolio of bonds. The predictable representation property gives the arbitrage-free price at time t of an attainable European claim with payoff X at maturity S , $t \leq S \leq T$,

$$C(t) = E_Q \left(\frac{\beta(t)}{\beta(S)} X | \mathcal{F}_t \right) = E_Q (e^{- \int_t^S r(u) du} X | \mathcal{F}_t). \quad (12.50)$$

For example, the price at time t of a European put with maturity S and strike K on a bond with maturity T (the right to sell the bond at time S for K) is given by

$$C(t) = E_Q \left(\frac{\beta(t)}{\beta(S)} (K - P(S, T))^+ | \mathcal{F}_t \right). \quad (12.51)$$

This option is in effect a cap on the interest rate over $[S, T]$ (see Section 12.7).

Forward Measures

Taking the bond, rather than the savings account, as a numeraire allows us to simplify the option pricing formulae.

Definition 12.4: The measure Q_T , called the T -forward measure, is obtained when the T -bond is a numeraire, i.e. $\beta(t)/P(t, T)$, $t \leq T$ is a Q_T -martingale.

Theorem 12.5: The forward measure Q_T is defined by

$$\Lambda(T) = \frac{dQ_T}{dQ} = \frac{1}{P(0, T)\beta(T)}. \quad (12.52)$$

The price of an attainable claim X at time t under different numeraire is related by the formula,

$$C(t) = E_Q \left(\frac{\beta(t)}{\beta(T)} X | \mathcal{F}_t \right) = P(t, T) E_{Q_T}(X | \mathcal{F}_t). \quad (12.53)$$

PROOF. It follows directly from the change of numeraire Theorem 11.17. According to the EMM assumption $P(t, T)/\beta(t)$, $t \leq T$, is a positive Q -martingale. $Q_T := Q_1$, with $\frac{dQ_T}{dQ} = \Lambda(T) = \frac{P(T, T)/P(0, T)}{\beta(T)/\beta(0)} = \frac{1}{P(0, T)\beta(T)}$, which is (12.52). According to Equation (11.50)

$$C(t) = E_Q \left(\frac{\beta(t)}{\beta(T)} X | \mathcal{F}_t \right) = E_{Q_T} \left(\frac{P(t, T)}{P(T, T)} X | \mathcal{F}_t \right) = P(t, T) E_{Q_T}(X | \mathcal{F}_t). \quad \square$$

Remark 12.3: $E_{Q_T}(X|\mathcal{F}_t) = F(t)$ is called the forward price at t for the date T of an attainable claim X . These prices are Q_T -martingales, see the Doob-Levy martingale, Theorem 7.9.

Forward Measure in HJM

Theorem 12.6: *The process*

$$dB^T(t) = dB(t) + \tau(t, T)dt \quad (12.54)$$

is a Q_T -Brownian motion. Moreover, the forward rates $f(t, T)$ and the bond $P(t, T)$ satisfy the following SDEs under Q_T

$$df(t, T) = \sigma(t, T)dB^T(t). \quad (12.55)$$

$$dP(t, T) = P(t, T)((r(t) + \tau^2(t, T))dt - \tau(t, T)dB^T(t)). \quad (12.56)$$

PROOF. The T -forward martingale measure Q_T is obtained by using $\Lambda(T) = \frac{1}{P(0, T)\beta(T)}$. Then

$$\Lambda_t = E(\Lambda|\mathcal{F}_t) = \frac{P(t, T)}{P(0, T)\beta(t)} = \mathcal{E}\left(-\int_0^t \tau(s, T)dB(s)\right).$$

Therefore by Girsanov's theorem $dB^T(t) = dB(t) + \tau(t, T)dt$ is a Q_T -Brownian motion. Under Q the forward rates satisfy the SDE (12.38)

$$df(t, T) = \sigma(t, T)\tau(t, T)dt + \sigma(t, T)dB(t).$$

The SDE under Q_T is obtained by replacing $B(t)$ with $B^T(t)$,

$$df(t, T) = \sigma(t, T)(\tau(t, T)dt + dB(t)) = \sigma(t, T)dB^T(t).$$

The SDE under Q_T for the bond prices is obtained similarly from the SDE (12.46) under Q . \square

The above result shows that under the forward measure $f(t, T)$ is a Q_T -local martingale. It also shows that $\tau(t, T)$ is the volatility of $P(t, T)$ under the forward measure Q_T .

Distributions of the Bond in HJM with Deterministic Volatilities

The following result, needed for option pricing, gives the conditional distribution of $P(T, T + \delta)$. As a corollary, distributions of $P(t, T)$ are obtained.

Theorem 12.7: Suppose that the forward rates satisfy the HJM model with a deterministic $\sigma(t, T)$. Then for any $0 \leq t < T$ and any $\delta > 0$, the Q -conditional distribution of $P(T, T + \delta)$ given \mathcal{F}_t is lognormal with mean

$$E_Q(\ln P(T, T + \delta) | \mathcal{F}_t) = \ln \frac{P(t, T + \delta)}{P(t, T)} - \frac{1}{2} \int_t^T (\tau^2(s, T + \delta) - \tau^2(s, T)) ds,$$

and variance

$$\gamma^2(t) = \text{Var}(\ln P(T, T + \delta) | \mathcal{F}_t) = \int_t^T (\tau(s, T + \delta) - \tau(s, T))^2 ds. \quad (12.57)$$

PROOF. By letting $t = T_1 = T$ and $T_2 = T + \delta$ in Equation (12.49), the expression for $P(T, T + \delta)$ is obtained,

$$P(T, T + \delta) = \frac{P(0, T + \delta)}{P(0, T)} e^{-\int_0^T (\tau(s, T + \delta) - \tau(s, T)) dB(s) - \frac{1}{2} \int_0^T (\tau^2(s, T + \delta) - \tau^2(s, T)) ds}. \quad (12.58)$$

In order to find its conditional distribution given \mathcal{F}_t , separate the \mathcal{F}_t -measurable term,

$$\begin{aligned} P(T, T + \delta) &= \frac{P(0, T + \delta)}{P(0, T)} e^{-\int_0^t (\tau(s, T + \delta) - \tau(s, T)) dB(s) - \frac{1}{2} \int_0^t (\tau^2(s, T + \delta) - \tau^2(s, T)) ds} \\ &\quad \times e^{-\int_t^T (\tau(s, T + \delta) - \tau(s, T)) dB(s) - \frac{1}{2} \int_t^T (\tau^2(s, T + \delta) - \tau^2(s, T)) ds} \\ &= \frac{P(t, T + \delta)}{P(t, T)} e^{-\int_t^T (\tau(s, T + \delta) - \tau(s, T)) dB(s) - \frac{1}{2} \int_t^T (\tau^2(s, T + \delta) - \tau^2(s, T)) ds}. \end{aligned} \quad (12.59)$$

Since $\tau(t, T)$ is non-random the exponential term is independent of \mathcal{F}_t . Hence the desired conditional distribution given \mathcal{F}_t is lognormal with the mean and variance as stated. \square

Corollary 12.8: The conditional distribution of $P(T, T + \delta)$ given \mathcal{F}_t under the forward measure $Q_{T+\delta}$ is lognormal with mean

$$E_{Q_{T+\delta}}(\ln P(T, T + \delta) | \mathcal{F}_t) = \ln \frac{P(t, T + \delta)}{P(t, T)} + \frac{1}{2} \int_t^T (\tau(s, T + \delta) - \tau(s, T))^2 ds,$$

and variance $\gamma^2(t)$ (12.57).

PROOF. Use Equation (12.59) for the conditional representation together with (12.54), a $Q_{T+\delta}$ -Brownian motion $dB^{T+\delta}(t) = dB(t) + \tau(t, T + \delta)dt$, to have

$$P(T, T + \delta) = \frac{P(t, T + \delta)}{P(t, T)} e^{-\int_t^T (\tau(s, T + \delta) - \tau(s, T)) dB^{T+\delta}(s) + \frac{1}{2} \int_t^T (\tau^2(s, T + \delta) - \tau^2(s, T)) ds}. \quad (12.60)$$

\square

By taking $T = t$, $t = 0$, $T + \delta = T$ in Theorem 12.7 and its corollary we obtain the result.

Corollary 12.9: *In the case of constant volatilities the Q-distribution and Q_T -distribution of $P(t, T)$ are lognormal.*

The lognormality of $P(t, T)$ can also be obtained directly from Equations (12.47) and (12.38).

12.7 Options, Caps, and Floors

We give common options on interest rates and show how they relate to options on bonds.

Cap and Caplets

A *cap* is a contract that gives its holder the right to pay the smaller rate of interest of the two, the floating rate, and rate k , specified in the contract. A party holding the cap will never pay a rate exceeding k , the rate of payment is capped at k . Since the payments are done at a sequence of payments dates T_1, T_2, \dots, T_n , called a *tenor*, with $T_{i+1} = T_i + \delta$ (e.g. $\delta = \frac{1}{4}$ of a year), the rate is capped over intervals of time of length δ . Thus a cap is a collection of caplets (see Figure 12.3).




Fig. 12.3: Payment dates and simple rates.

Consider a caplet over $[T, T + \delta]$. Without the caplet the holder of a loan must pay at time $T + \delta$ an interest payment of $f\delta$, where f is the floating, simple rate over the interval $[T, T + \delta]$. If $f > k$, then a caplet allows the holder to pay $k\delta$. Thus the caplet is worth $f\delta - k\delta$ at time $T + \delta$. If $f < k$, then the caplet is worthless. Therefore, the caplet's worth to the holder is $(f - k)^+\delta$. In other words, a caplet pays to its holder the amount $(f - k)^+\delta$ at time $T + \delta$. Therefore a caplet is a call option on the rate f , and its price at time t , as any other option, is given by the expected discounted payoff at maturity under the EMM Q,

$$\text{Caplet}(t) = \mathbb{E}_Q \left(\frac{\beta(t)}{\beta(T + \delta)} (f - k)^+\delta \mid \mathcal{F}_t \right). \quad (12.61)$$

In order to evaluate this expectation we need to know the distribution of f under Q. One way to find the distribution of the simple floating rate is to relate it to the bond over the same time interval. By definition,

$\frac{1}{P(T, T+\delta)} = 1 + f\delta$. This relation is justified as the amounts obtained at time $T + \delta$ when \$1 invested at time T in the bond and in the investment account with a simple rate f . Thus

$$f = \frac{1}{\delta} \left(\frac{1}{P(T, T+\delta)} - 1 \right). \quad (12.62)$$

This is a basic relation between the rates that appear in option contracts and bonds. It gives the distribution of f in terms of that of $P(T, T + \delta)$.

Caplet as a Put Option on the Bond

We show next that a caplet is in effect a put option on the bond. From the basic relation (EMM) $P(T, T + \delta) = E(\frac{\beta(T)}{\beta(T+\delta)} | \mathcal{F}_T)$. Proceeding from (12.61) by the law of double expectation, with $E = E_Q$

$$\begin{aligned} \text{Caplet}(t) &= E \left(E \left(\frac{\beta(t)\beta(T)}{\beta(T)\beta(T+\delta)} \left(\frac{1}{P(T, T+\delta)} - 1 - k\delta \right)^+ | \mathcal{F}_T \right) | \mathcal{F}_t \right) \\ &= E \left(\frac{\beta(t)}{\beta(T)} \left(\frac{1}{P(T, T+\delta)} - 1 - k\delta \right)^+ E \left(\frac{\beta(T)}{\beta(T+\delta)} | \mathcal{F}_T \right) | \mathcal{F}_t \right) \\ &= (1 + k\delta) E \left(\frac{\beta(t)}{\beta(T)} \left(\frac{1}{(1+k\delta)} - P(T, T+\delta) \right)^+ | \mathcal{F}_t \right). \end{aligned} \quad (12.63)$$

Thus a caplet is a put option on $P(T, T + \delta)$ with strike $\frac{1}{(1+k\delta)}$ and exercise time T . In practical modelling, as in the HJM model with deterministic volatilities, the distribution of $P(T, T + \delta)$ is lognormal, giving rise to the Black–Scholes type formula for a caplet, Black's (1976) formula.

The price of a caplet is easier to evaluate by using a forward measure (Theorem 12.5). Take $P(t, T + \delta)$ as a numeraire, which corresponds to $T + \delta$ -forward measure $Q_{T+\delta}$,

$$\begin{aligned} \text{Caplet}(t) &= P(t, T + \delta) E_{Q_{T+\delta}}((f - k)^+ \delta | \mathcal{F}_t) \\ &= P(t, T + \delta) E_{Q_{T+\delta}} \left(\left(\frac{1}{P(T, T+\delta)} - 1 - k\delta \right)^+ | \mathcal{F}_t \right). \end{aligned} \quad (12.64)$$

Caplet Pricing in the HJM Model

The price of a caplet, which caps the rate at k over the time interval $[T, T + \delta]$, at time t is given by Equation (12.63) under the EMM Q , and by (12.64) under the forward EMM $Q_{T+\delta}$. These can be evaluated in closed form when the volatilities are non-random.

For evaluating the expectation in the caplet formula (12.64) note that if X is lognormal $e^{N(\mu, \sigma^2)}$, then $1/X = e^{N(-\mu, \sigma^2)}$ is also lognormal. This

allows us to price the caplet by doing standard calculations for $E(X - K)^+$ giving Black's caplet formula.

The price of a cap is then found by using forward measures

$$\text{Cap}(t) = \sum_{i=1}^n \text{Caplet}_i(t) = \sum_{i=1}^n P(t, T_i) E_{Q_{T_i}}((f_{i-1} - k)^+ \delta | \mathcal{F}_t). \quad (12.65)$$

The cap pricing formula is given in Exercise 12.6.

A floor is a contract that gives its holder the right to receive the larger of the two rates, the rate specified in the contract k , and the floating simple rate f . Floors are priced similarly to caps with floorlets being $(k - f_{i-1})^+$.

12.8 Brace–Gatarek–Musiela (BGM) Model

In financial markets Black–Scholes-like formulae are used for everything: bonds, rates, etc. In order to make the practice consistent with the theory, Brace, Gatarek, and Musiela introduced a class of models, which can be seen as a subclass of HJM models, where instead of the forward rates $f(t, T)$, the London Inter-Bank Offer Rates (LIBORs) are modelled (Brace *et al.* (1997), Musiela and Rutkowski (1998)). In BGM models the rates are lognormal under forward measures, a fact that implies option pricing formulae are consistent with market use.

LIBOR

The time t forward δ -LIBOR is the simple rate of interest on $[T, T + \delta]$

$$L(t, T) = \frac{1}{\delta} \left(\frac{P(t, T)}{P(t, T + \delta)} - 1 \right); \quad (12.66)$$

note that $L(T, T) = f$ is the rate that gets capped, see (12.62).

Theorem 12.10: *The SDE for $L(t, T)$ under the forward $Q_{T+\delta}$ measure, when $P(t, T + \delta)$ is the numeraire, is*

$$dL(t, T) = L(t, T) \left(\frac{1 + L(t, T)\delta}{L(t, T)\delta} \right) (\tau(t, T + \delta) - \tau(t, T)) dB^{T+\delta}(t). \quad (12.67)$$

PROOF. According to Corollary 12.3, Equation (12.49),

$$\frac{P(t, T)}{P(t, T + \delta)} = \frac{P(0, T)}{P(0, T + \delta)} e^{- \int_0^t (\tau(s, T) - \tau(s, T + \delta)) dB(s) - \frac{1}{2} \int_0^t (\tau^2(s, T) - \tau^2(s, T + \delta)) ds}. \quad (12.68)$$

$dB^{T+\delta}(t) = dB(t) + \tau(t, T + \delta)dt$, is a $Q_{T+\delta}$ -Brownian motion (12.54), giving

$$\begin{aligned} \frac{P(t, T)}{P(t, T + \delta)} &= \frac{P(0, T)}{P(0, T + \delta)} e^{-\int_0^t (\tau(s, T) - \tau(s, T + \delta)) dB^{T+\delta}(s) - \frac{1}{2} \int_0^t (\tau(s, T) - \tau(s, T + \delta))^2 ds} \\ &= \frac{P(0, T)}{P(0, T + \delta)} \mathcal{E}\left(\int_0^t (\tau(s, T + \delta) - \tau(s, T)) dB^{T+\delta}(s)\right). \end{aligned} \quad (12.69)$$

Using the stochastic exponential equation it follows that

$$d\left(\frac{P(t, T)}{P(t, T + \delta)}\right) = \left(\frac{P(t, T)}{P(t, T + \delta)}\right) (\tau(t, T + \delta) - \tau(t, T)) dB^{T+\delta}(t). \quad (12.70)$$

Finally, using the definition of $L(t, T)$, the SDE (12.67) is obtained. \square

Now choose the HJM volatility $\sigma(t, s)$ such that $\gamma(t, T)$ is deterministic

$$\gamma(t, T) = \left(\frac{1 + L(t, T)\delta}{L(t, T)\delta}\right) (\tau(t, T + \delta) - \tau(t, T)) = \frac{1 + L(t, T)\delta}{L(t, T)\delta} \int_T^{T+\delta} \sigma(t, s) ds. \quad (12.71)$$

Corollary 12.11: Let $\gamma(t, T)$ be deterministic such that $\int_0^T \gamma^2(s, T) ds < \infty$. Then $L(t, T)$ for $t \leq T$ solves the SDE

$$dL(t, T) = L(t, T) \gamma(t, T) dB^{T+\delta}(t), \quad (12.72)$$

with the initial condition $L(0, T) = \frac{1}{\delta}(\frac{P(0, T)}{P(0, T + \delta)} - 1)$. Thus $L(t, T)$ is lognormal under the forward measure $Q_{T+\delta}$, moreover, it is a martingale.

$$L(T, T) = L(t, T) e^{\int_t^T \gamma(s, T) dB^{T+\delta}(s) - \frac{1}{2} \int_t^T \gamma^2(s, T) ds}, \quad (12.73)$$

and the conditional distribution of $L(T, T)$ given \mathcal{F}_t is lognormal with mean $\ln L(t, T) - \frac{1}{2} \int_t^T \gamma^2(s, T) ds$ and variance $\int_t^T \gamma^2(s, T) ds$.

We now prove that the choice of forward volatilities specified above is possible.

Theorem 12.12: Let $\gamma(t, T)$, $t \leq T$ be given such that the Itô integral $\int_0^T \gamma(s, T) dB(s)$ is defined. Then there exist forward rates volatilities $\sigma(t, T)$ such that the integrated volatility $\int_T^{T+\delta} \sigma(s, u) du$ is determined uniquely, and (12.72) holds.

PROOF. (12.72) is equivalent to

$$L(t, T) = L(0, T) \mathcal{E}\left(\int_0^t \gamma(s, T) dB^{T+\delta}(s)\right), \quad t \leq T. \quad (12.74)$$

According to the definition of $L(t, T)$ and (12.69) we have

$$\begin{aligned} L(t, T) &= \frac{1}{\delta} \left(\frac{P(t, T)}{P(t, T + \delta)} - 1 \right) \\ &= \frac{1}{\delta} \left(\frac{P(0, T)}{P(0, T + \delta)} \mathcal{E} \left(\int_0^t (\tau(s, T + \delta) - \tau(s, T)) dB^{T+\delta}(s) \right) - 1 \right). \end{aligned}$$

Equating this to (12.74) we obtain the equation for stochastic exponentials

$$\mathcal{E} \left(\int_0^t (\tau(s, T + \delta) - \tau(s, T)) dB^{T+\delta}(s) \right) = (1 - c) \mathcal{E} \left(\int_0^t \gamma(s, T) dB^{T+\delta}(s) \right) + c, \quad (12.75)$$

with $c = \frac{P(0, T + \delta)}{P(0, T)}$. Now, using the stochastic logarithm (Theorem 5.3)

$$\int_0^t \left(\int_T^{T+\delta} \sigma(s, u) du \right) dB^{T+\delta}(s) = \mathcal{L} \left((1 - c) \mathcal{E} \left(\int_0^t \gamma(s, T) dB^{T+\delta}(s) \right) + c \right), \quad (12.76)$$

from which the integrated volatility $\int_T^{T+\delta} \sigma(s, u) du$ and a suitable process $\sigma(t, T)$ can be found (see Exercise 12.7). \square

Caplet in BGM

The caplet is a call on LIBOR, it pays $\delta(L(T, T) - k)^+$ at $T + \delta$. According to (12.53) its price at time t

$$C(t) = P(t, T + \delta) \mathbb{E}_{Q_{T+\delta}}(\delta(L(T, T) - k)^+ | \mathcal{F}_t). \quad (12.77)$$

Using that $L(T, T)$ is lognormal under the forward measure $Q_{T+\delta}$ (12.73), the caplet is priced by Black's caplet formula (agrees with the market) by evaluating under the forward measure.

$$\begin{aligned} C(t) &= P(t, T + \delta) (L(t, T) \Phi(h_1) - k \Phi(h_2)), \\ h_{1,2} &= \frac{\ln \frac{L(t, T)}{k} \pm \frac{1}{2} \int_t^T \gamma^2(s, T) ds}{\sqrt{\int_t^T \gamma^2(s, T) ds}}. \end{aligned} \quad (12.78)$$

SDEs for Forward LIBOR Under Different Measures

Consider now a sequence of dates T_0, T_1, \dots, T_n , and denote by Q_{T_k} the forward measure corresponding to the numeraire $P(t, T_k)$. Corollary 12.11 states that $L(t, T_{k-1})$ for $t \leq T_{k-1}$ satisfies the SDE

$$dL(t, T_{k-1}) = \gamma(t, T_{k-1}) L(t, T_{k-1}) dB^{T_k}(t), \quad (12.79)$$

where $B^{T_k}(t)$ is a Q_{T_k} -Brownian motion.

An SDE for all of the rates under a single measure is sometimes required (for a swaption), it is given by

Theorem 12.13: *For a given i , the SDE for the forward LIBOR $L(t, T_{k-1})$ on $[T_{k-1}, T_k]$ under the forward measure Q_{T_i} , is given by (12.79) for $i = k$, by (12.80) for $i > k$, and (12.81) $i < k$.*

$$\begin{aligned} dL(t, T_{k-1}) &= -L(t, T_{k-1}) \sum_{j=k}^{i-1} \frac{\delta\gamma(t, T_{k-1})\gamma(t, T_j)L(t, T_j)}{1 + \delta L(t, T_j)} dt \\ &\quad + \gamma(t, T_{k-1})L(t, T_{k-1})dB^{T_i}(t). \end{aligned} \quad (12.80)$$

$$\begin{aligned} dL(t, T_{k-1}) &= L(t, T_{k-1}) \sum_{j=i}^{k-1} \frac{\delta\gamma(t, T_{k-1})\gamma(t, T_j)L(t, T_j)}{1 + \delta L(t, T_j)} dt \\ &\quad + \gamma(t, T_{k-1})L(t, T_{k-1})dB^{T_i}(t). \end{aligned} \quad (12.81)$$

PROOF. We establish the relationship between different forward measures as well as corresponding Brownian motions. According to (12.54) the Brownian motion under the forward measure Q_{T_k} satisfies

$$dB^{T_k}(t) = dB(t) + \tau(t, T_k)dt. \quad (12.82)$$

Hence, using this with $k-1$, we obtain that B^{T_k} and $B^{T_{k-1}}$ are related by

$$dB^{T_k}(t) = dB^{T_{k-1}}(t) + (\tau(t, T_k) - \tau(t, T_{k-1}))dt.$$

According to (12.71), from the choice of γ ,

$$\tau(t, T_k) - \tau(t, T_{k-1}) = \gamma(t, T_{k-1}) \frac{L(t, T_{k-1})\delta}{1 + L(t, T_{k-1})\delta},$$

giving the relationship for the SDEs for LIBOR

$$dB^{T_k}(t) = dB^{T_{k-1}}(t) + \gamma(t, T_{k-1}) \frac{L(t, T_{k-1})\delta}{1 + L(t, T_{k-1})\delta} dt.$$

Now fix $i > k$. Using the above relation iteratively from i to k we obtain

$$dB^{T_i}(t) = dB^{T_k}(t) + \sum_{j=k}^{i-1} \gamma(t, T_j) \frac{L(t, T_j)\delta}{1 + L(t, T_j)\delta} dt. \quad (12.83)$$

Replace B^{T_k} in the SDE (12.79) for $L(t, T_{k-1})$ under Q_{T_k} ,

$dL(t, T_{k-1}) = \gamma(t, T_{k-1})L(t, T_{k-1})dB^{T_k}(t)$, by B^{T_i} with the drift from (12.83) to obtain (12.80). The case $i < k$ is proved similarly. \square

Another proof can be done by using the result on the SDE under a new numeraire, Theorem 11.18, with $\beta(t) = P(t, T_k)$ and $S(t) = P(t, T_i)$.

Choice of Bond Volatilities

We comment briefly on the choice of bond volatilities. For a sequence of dates Equation (12.71) implies from the choice of γ ,

$$\tau(t, T_k) - \tau(t, T_{k-1}) = \gamma(t, T_{k-1}) \frac{L(t, T_{k-1})\delta}{1 + L(t, T_{k-1})\delta}. \quad (12.84)$$

Therefore the bond volatilities $\tau(t, T_k)$ can be obtained recursively,

$$\begin{aligned} \tau(t, T_k) &= \tau(t, T_{k-1}) + \gamma(t, T_{k-1}) \frac{L(t, T_{k-1})\delta}{1 + L(t, T_{k-1})\delta} \\ &= \tau(t, T_0) + \sum_{j=0}^{k-1} \gamma(t, T_j) \frac{L(t, T_j)\delta}{1 + L(t, T_j)\delta}. \end{aligned} \quad (12.85)$$

In practice $\gamma(t, T)$ is taken to be a function of T only (Rebonato (2002), Brace (1998)), for example, $\gamma(t, T) = (a + be^{cT})$.

12.9 Swaps and Swaptions

A basic instrument in the interest rate market is the payer swap, in which the floating rate is swapped in arrears (at T_i) against a fixed rate k at n intervals of length $\delta = T_i - T_{i-1}$, $i = 1, 2, \dots, n$. The other party in the swap enters a receiver swap, in which a fixed rate is swapped against the floating. By a swap we shall mean only a payer swap. A swap value at time t , by the basic relation (12.50), is given by

$$\text{Swap}(t, T_0, k) = E_Q \sum_{i=1}^n \delta \left(\frac{\beta(t)}{\beta(T_i)} (L(T_{i-1}, T_{i-1}) - k) \middle| \mathcal{F}_t \right).$$

This can be written using forward measures (Theorem 12.5)

$$\begin{aligned} \text{Swap}(t, T_0, k) &= \delta \sum_{i=1}^n P(t, T_i) E_{Q_{T_i}} \left((L(T_{i-1}, T_{i-1}) - k) \middle| \mathcal{F}_t \right) \\ &= \delta \sum_{i=1}^n P(t, T_i) (L(t, T_{i-1}) - k), \end{aligned} \quad (12.86)$$

as under the forward measure Q_{T_i} , $L(t, T_{i-1})$ is a martingale (Corollary 12.11).

A swap agreement is worthless at initiation. The forward swap rate is that fixed rate of interest which makes a swap worthless. Namely, the swap rate at time t for the date T_0 solves, by definition, $\text{Swap}(t, T_0, k) = 0$. Hence

$$k(t, T_0) = \frac{\sum_{i=1}^n P(t, T_i) L(t, T_{i-1})}{\sum_{i=1}^n P(t, T_i)}. \quad (12.87)$$

Thus the t -value of the swap in terms of this rate is

$$\text{Swap}(t, T_0, k) = \delta \left(\sum_{i=1}^n P(t, T_i) \right) (k(t, T_0) - k). \quad (12.88)$$

Other expressions for the swap are given in Exercises 12.9 and 12.10.

A payer swaption maturing (with exercise) at $T = T_0$ gives its holder the right to enter a payer swap at time T_0 . A swaption, $\text{Swaption}(t, T_0)$, delivers at time T_0 a swap, $\text{Swap}(T_0, T_0)$, when the value of that swap is positive. This shows the value of the swaption at maturity is $(\text{Swap}(T_0, T_0))^+$. Thus its value at time $t \leq T_0$ is

$$\begin{aligned} \text{Swaption}(t, T_0) &= E_Q \left(\frac{\beta(t)}{\beta(T_0)} (\text{Swap}(T_0, T_0))^+ \middle| \mathcal{F}_t \right) \\ &= \delta E_Q \left(\frac{\beta(t)}{\beta(T_0)} \left(\sum_{i=1}^n P(T_0, T_i) \right) (k(T_0, T_0) - k)^+ \middle| \mathcal{F}_t \right). \end{aligned} \quad (12.89)$$

Consider taking $\sum_{i=1}^n P(t, T_i)$ as the numeraire instead of $\beta(t)$. The process $\sum_{i=1}^n P(t, T_i)/\beta(t)$ is a Q-martingale, as a sum of martingales. According to Theorem 11.17, the new measure \hat{Q}_{T_0} defined by $\Lambda(T) = \frac{\sum_{i=1}^n P(T_0, T_i) / \sum_{i=1}^n P(0, T_i)}{\beta(T_0)}$, as its Radon–Nikodym derivative with respect to Q gives (a call on the swap rate)

$$\text{Swaption}(t, T_0) = \delta \left(\sum_{i=1}^n P(t, T_i) \right) E_{\hat{Q}_{T_0}} \left((k(T_0, T_0) - k)^+ \middle| \mathcal{F}_t \right). \quad (12.90)$$

The distribution of the swap rate under the swap-rate measure \hat{Q}_{T_0} is approximately lognormal. Differentiating, simplifying, and approximating $k(t, T_0)$ in (12.87) leads to the SDE for the swap rate

$$dk(t, T_0) = \hat{\sigma}(t, T_0) k(t, T_0) d\hat{B}_{T_0},$$

where \hat{B}_{T_0} is a \hat{Q}_{T_0} -Brownian motion, and

$$\hat{\sigma}(t, T_0) = \sum_{i=1}^n w_i \gamma(t, T_{i-1}), \quad w_i = \frac{\frac{P(0, T_i)}{P(0, T_0)} L(0, T_{i-1})}{\sum_{i=1}^n \frac{P(0, T_i)}{P(0, T_0)} L(0, T_{i-1})}.$$

The expression for the swaption (12.90) integrates to the Black swaption formula as used in the market.

Another way to evaluate a swaption is by simulation. For details on analytic approximations and simulations see Brace *et al.* (1997), Brace (1998), and Musiela and Rutkowski (1998).

Remark 12.4: We have presented a brief mathematical treatment of models for rates and bonds based on diffusion processes. There is a large amount of literature on models based on processes with jumps. For jumps in the spot rate see Borovkov *et al.* (2003) and references therein. HJM with jumps were studied by Shirakawa (1991), and more generally by Björk, Kabanov, and Runggaldier (BKR) (1996), and Björk, Di Masi, Kabanov, and Runggaldier (BDMKR) (1997). Kennedy (1994) considered a Gaussian Markov field model. HJM and BDMKR models can be criticized for being an infinite-dimensional diffusion driven by a finite number of independent noise processes. Cont (1998) suggests modelling the forward curves by an infinite-dimensional diffusion driven by an infinite-dimensional Brownian motion. This approach is included in random fields models, such as Brownian and Poisson sheet models, also known as models with space-time noise. The most general model that allows for the existence of EMM is given in Hamza *et al.* (2005); it includes Gaussian and Poisson random field models.

12.10 Exercises

Exercise 12.1: Show that a European call option on the T -bond is given by $C(t) = P(t, T)Q_T(P(s, T) > K|\mathcal{F}_t) - KP(t, s)Q_s(P(s, T) > K|\mathcal{F}_t)$, where s is the exercise time of the option and Q_s , Q_T are s and T -forward measures.

Exercise 12.2: Show that a European call option on the bond in the Merton model is given by

$$P(t, T)\Phi\left(\frac{\ln \frac{P(t, T)}{KP(t, s)} + \frac{\sigma^2(T-s)^2(s-t)}{2}}{\sigma(T-s)\sqrt{s-t}}\right) - KP(t, s)\Phi\left(\frac{\ln \frac{P(t, T)}{KP(t, s)} - \frac{\sigma^2(T-s)^2(s-t)}{2}}{\sigma(T-s)\sqrt{s-t}}\right).$$

Exercise 12.3: (Stochastic Fubini theorem)

Let $H(t, s)$ be continuous $0 \leq t, s \leq T$, and for any fixed s , $H(t, s)$ as a process in t , $0 \leq t \leq T$ is adapted to the Brownian motion filtration \mathcal{F}_t . Assume $\int_0^T H^2(t, s)dt < \infty$ so that for each s the Itô integral $X(s) = \int_0^T H(t, s)dW(t)$ is defined. Since $H(t, s)$ is continuous, $Y(t) = \int_0^T H(t, s)ds$ is defined and it is continuous and adapted. Assume

$$\int_0^T \mathbb{E}\left(\int_0^T H^2(t, s)dt\right) ds < \infty.$$

1. Show that $\int_0^T \mathbb{E}|X(s)|ds \leq \int_0^T \left(\mathbb{E} \int_0^T H^2(t, s)dt\right)^{1/2} ds < \infty$, consequently $\int_0^T X(s)ds$ exists.

2. If $0 = t_0 < t_1 < \dots < t_n = T$ is a partition of $[0, T]$, show that

$$\int_0^T \left(\sum_{i=0}^{n-1} H(t_i, s)(W(t_{i+1}) - W(t_i)) \right) ds = \sum_{i=0}^{n-1} \left(\int_0^T H(t_i, s)ds \right) (W(t_{i+1}) - W(t_i)).$$

3. By taking the limits as the partition shrinks, show that

$$\int_0^T X(s)ds = \int_0^T Y(t)dW(t),$$

in other words, the order of integration can be interchanged

$$\int_0^T \left(\int_0^T H(t, s)dW(t) \right) ds = \int_0^T \left(\int_0^T H(t, s)ds \right) dW(t).$$

Exercise 12.4: (One factor HJM)

Show that the correlation between the forward rates $f(t, T_1)$ and $f(t, T_2)$ in the HJM model with deterministic volatilities $\sigma(t, T)$ is given by

$$\rho(T_1, T_2) = \text{Corr}(f(t, T_1), f(t, T_2)) = \frac{\int_0^t \sigma(s, T_1)\sigma(s, T_2)ds}{\sqrt{\int_0^t \sigma^2(s, T_1)ds \int_0^t \sigma^2(s, T_2)ds}}.$$

Give the class of volatilities for which the correlation is one.

Exercise 12.5: Find the forward rates $f(t, T)$ in Vasicek's model. Give the price of a cap.

Exercise 12.6: (Caps pricing market formula)

Show that in the HJM model with a deterministic $\sigma(t, T)$ the price of a cap with trading dates $T_i = T + i\delta$, $i = 1, \dots, n$, and strike rate k is given by

$$\sum_{i=1}^n (P(t, T_{i-1})\Phi(-h_{i-1}(t)) - (1 + K\delta)P(t, T_i)\Phi(-h_{i-1}(t) - \gamma_{i-1}(t))),$$

where $\gamma_{i-1}^2(t) = \text{Var}(\ln P(T_{i-1}, T_i)) = \int_t^{T_{i-1}} |\tau(s, T_i) - \tau(s, T_{i-1})|^2 ds$, with $\tau(t, T) = \int_t^T \sigma(t, s)ds$ and $h_{i-1}(t) = \frac{1}{\gamma_{i-1}(t)} \left(\ln \frac{(1+k\delta)P(t, T_i)}{P(t, T_{i-1})} - \frac{1}{2}\gamma_{i-1}^2(t) \right)$.

Exercise 12.7: Let $0 < c < 1$, and $\gamma(t)$ a bounded deterministic function be given. Show that there is a process $\beta(s)$ such that

$$c + (1 - c)\mathcal{E}\left(\int_0^t \gamma(s)dB(s)\right) = \mathcal{E}\left(\int_0^t \beta(s)dB(s)\right).$$

Hence deduce the existence of forward rates volatilities $\sigma(t, T)$ in HJM from specification of the forward LIBOR volatilities $\gamma(t, T)$ in BGM.

Exercise 12.8: (Two-factor and higher HJM models)

A two-factor HJM is given by the SDE

$$df(t, T) = \alpha(t, T)dt + \sigma_1(t, T)dW_1(t) + \sigma_2(t, T)dW_2(t),$$

where W_1 and W_2 are independent Brownian motions.

1. Give the stochastic differential equation for the log of the bond prices and show that

$$\frac{P(t, T)}{\beta(t)} = P(0, T)e^{-\int_0^t A_1(u, T)du - \int_0^t \tau_1(u, T)dW_1(u) - \int_0^t \tau_2(u, T)dW_2(u)},$$

with $\tau_i(t, T) = \int_t^T \sigma_i(t, s)ds$, $i = 1, 2$, and $A_1(t, T) = \int_t^T \alpha(t, s)ds$.

2. Using the same proof as in the one-factor model show that the no-arbitrage condition is given by

$$\alpha(t, T) = \sigma_1(t, T) \int_t^T \sigma_1(t, s)ds + \sigma_2(t, T) \int_t^T \sigma_2(t, s)ds.$$

Exercise 12.9: Show that a swap can be written as

$$\text{Swap}(t, T_0, k) = P(t, T_0) - P(t, T_n) - k\delta \sum_{i=1}^n P(t, T_i).$$

Exercise 12.10: Denote by $b(t) = \delta \sum_{i=1}^n P(t, T_i)$. Show that for $0 < t \leq T_0$, $\text{Swap}(t, T_0, k) = P(t, T_0) - P(t, T_n) - kb(t)$, and that the swap rate

$$k(t) = \frac{P(t, T_0) - P(t, T_n)}{b(t)}.$$

Exercise 12.11: (Jamshidian (1996) swaptions pricing formula)

Assume that the swap rate $k(t) > 0$, and that $v^2(t) = \int_t^T \frac{1}{k^2(s)} d[k, k](s)$ is deterministic. Show that

$$\text{Swaption}(t) = b(t) (\alpha_+(t)k(t) - k\alpha_-(t)),$$

where

$$\alpha_{\pm}(t) = \Phi \left(\frac{\ln k(t)/k}{v(t)} \pm \frac{v(t)}{2} \right).$$

This page intentionally left blank

Chapter 13

Applications in Biology

In this chapter applications of stochastic calculus to population models are given. Diffusion models, Markov jump processes, non-Markov models (age-dependent), and stochastic models for competition of species are presented. Diffusion models are used in various areas of biology as models for population growth and genetic evolution. Birth–death processes are random walks in continuous time and many models in genetics, cell biology, and cancer are based on them. For example, the Moran model is a genetic model for competition of traits, it is also used for modelling cancer. Other models include random perturbation of deterministic laws found in kinetics. First we present diffusion models, and jump models afterwards, as their treatment, although similar, requires some extra ideas. A novel approach to the age-dependent branching (Bellman–Harris) process is given by treating it as a simple measure-valued process. The stochastic model for interacting populations that generalizes the Lotka–Volterra prey–predator model is treated by using a semimartingale representation. It is possible to formulate these models as stochastic differential equations. We demonstrate how results on stochastic differential equations and martingales presented in earlier chapters are used to analyze the above mentioned models.

13.1 Feller’s Branching Diffusion

A simple branching process is a model in which individuals reproduce independently of each other and of the history of the process. The continuous approximation to branching processes is the branching diffusion. It is given by the stochastic differential equation for the population size $X(t)$, $X(0) > 0$,

$$dX(t) = \alpha X(t)dt + \sigma \sqrt{X(t)}dB(t). \quad (13.1)$$

The diffusion coefficient $\sigma\sqrt{x}$ is not Lipschitz continuous at zero, see Example 1.12. Thus it is not obvious that a solution exists. However, it is possible to construct a solution by approximations using equations with Lipschitz continuous coefficients. We shall show that a unique solution exists, and is non-negative, so that there is no problem with the square root in the diffusion coefficient. The corresponding forward (Kolmogorov or Fokker–Plank) equation for the probability density of $X(t)$ is

$$\frac{\partial p(t, x)}{\partial t} = -\alpha \frac{\partial(xp(t, x))}{\partial x} + \frac{\sigma^2}{2} \frac{\partial^2(xp(t, x))}{\partial x^2}.$$

Feller (1951) analyzed this process by solving the partial differential Equation (13.1) (and thus constructing a weak solution). Here we demonstrate the stochastic calculus approach by obtaining the information directly from the stochastic Equation (13.1). First we prove that there is a unique strong solution to (13.1), which is always non-negative, moreover, once it hits 0 it stays at 0 forever (0 is absorbing). Below we use notation $a \vee b = \max(a, b)$.

Theorem 13.1: *The stochastic differential Equation (13.1) with $X(0) > 0$ possesses a unique strong non-negative solution. Furthermore, if $X(t)$ is that solution and $\tau = \inf\{t : X(t) = 0\}$, then $X(t) = 0$ for all $t \geq \tau$.*

PROOF. We follow Abramov *et al.* (2011). Consider the following stochastic differential equation

$$dX^n(t) = \alpha X^n(t)dt + \sigma \sqrt{\frac{1}{n} \vee X^n(t)} dB(t), \quad X^n(0) = X(0),$$

with $n > 1/X(0)$. Since the diffusion coefficient $\sigma\sqrt{n^{-1} \vee x}$ is Lipschitz continuous this equation has a unique strong solution.

Denote $\tau_0 = 0$, and $\tau_n = \inf\{t > 0 : X^n(t) = n^{-1}\}$, with $\tau_n = \infty$ if no such t exists. Note that for $t < \tau_n$ all solutions with index greater than n coincide, as they all solve the same Equation (13.1)

$$X^{n+1}(t) = X^n(t).$$

It is clear that $\tau_{n+1} > \tau_n$. Hence there is a limit (possibly infinite) $\tau = \lim_{n \rightarrow \infty} \tau_n$. Define $X(t \wedge \tau) = X^{n+1}(t)$ if $\tau_n \leq t < \tau_{n+1}$, and $X(t) = 0$ for $t > \tau$, if $\tau < \infty$. Then $X(t)$ is a solution to (13.1). Yamada–Watanabe’s theorem, Theorem 5.5, guarantees the uniqueness of the strong solution to (13.1) because the diffusion parameter is Hölder continuous with parameter $\frac{1}{2}$. \square

The next result describes the exponential growth of the population.

Theorem 13.2: Let $X(t)$ solve (13.1) and $X(0) > 0$. Then $EX(t) = X(0)e^{\alpha t}$. $X(t)e^{-\alpha t}$ is a non-negative martingale which converges almost surely to a non-degenerate limit as $t \rightarrow \infty$ if $\alpha > 0$.

Before we write a formal equation for the expectation, we must show that it is finite for any t . For this we need an auxiliary lemma.

Lemma 13.3: Let $g(t) \geq 0$ be a non-negative function defined for $t \geq 0$. Then for any positive t and T

$$\int_0^{t \wedge T} g(s)ds \leq \int_0^t g(s \wedge T)ds. \quad (13.2)$$

PROOF. (Of the lemma) If $t \leq T$, then both integrals are the same. If $t > T$, then

$$\int_0^t g(s \wedge T)ds = \int_0^T g(s)ds + \int_T^t g(T)ds \geq \int_0^T g(s)ds = \int_0^{t \wedge T} g(s)ds. \quad \square$$

PROOF. (Of the theorem) First we show that $X(t)$ is integrable. Since Itô integrals are local martingales, $\int_0^t \sqrt{X(s)}dB(s)$ is a local martingale. Let $T_n = \inf\{t : X(t) \geq n\}$. It is a localizing sequence so that $\int_0^{t \wedge T_n} \sqrt{X(s)}dB(s)$ is a martingale in t for any fixed n . Then using (13.1) we can write

$$X(t \wedge T_n) = X(0) + \alpha \int_0^{t \wedge T_n} X(s)ds + \sigma \int_0^{t \wedge T_n} \sqrt{X(s)}dB(s).$$

Since for $t \leq T_n$, $X(t) \leq n$, we can take expectation to obtain

$$EX(t \wedge T_n) = X(0) + \alpha E \int_0^{t \wedge T_n} X(s)ds.$$

Using (13.2) and Fubini's theorem we obtain the inequality

$$EX(t \wedge T_n) \leq X(0) + \alpha \int_0^t EX(s \wedge T_n)ds.$$

Using Gronwall's inequality (Theorem 1.20) with the function $EX(t \wedge T_n)$ we obtain the bound

$$EX(t \wedge T_n) \leq X(0)e^{\alpha t}.$$

Since $X(t \wedge T_n) \rightarrow X(t)$ as $n \rightarrow \infty$, we obtain by Fatou's lemma (noting that if a limit exists, then $\liminf = \lim$)

$$\begin{aligned} EX(t) &= E \lim_{n \rightarrow \infty} X(t \wedge T_n) = E \liminf_{n \rightarrow \infty} X(t \wedge T_n) \\ &\leq \liminf_{n \rightarrow \infty} EX(t \wedge T_n) = \lim_{n \rightarrow \infty} EX(t \wedge T_n) \leq X(0)e^{\alpha t}. \end{aligned}$$

Now we can show that the local martingale $\int_0^t \sqrt{X(s)} dB(s)$ is a true martingale. Consider its quadratic variation

$$\mathbb{E}\left[\int_0^{\cdot} \sqrt{X(s)} dB(s), \quad \int_0^{\cdot} \sqrt{X(s)} dB(s)\right](t) = \mathbb{E} \int_0^t X(s) ds < Ce^{\alpha t},$$

where we have used that $EX(s) \leq X(0)e^{\alpha s}$, and C is a constant. Thus by Theorem 7.35, $\int_0^t \sqrt{X(s)} dB(s)$ is a martingale. Now we can take expectations in (13.1). Differentiating with respect to t and solving the resulting equation we find $EX(t) = X(0)e^{\alpha t}$.

In order to show that $X(t)e^{-\alpha t}$ is a martingale use integration by parts

$$U(t) = X(t)e^{-\alpha t} = X(0) + \sigma \int_0^t e^{-\alpha s} \sqrt{X(s)} dB(s),$$

and $U(t)$ is a local martingale.

$$\mathbb{E}[U, U](t) = \sigma^2 \mathbb{E} \int_0^t e^{-2\alpha s} X(s) ds = \sigma^2 \int_0^t e^{-\alpha s} ds.$$

Hence for any finite T , $\mathbb{E}[U, U](T) < \infty$. Therefore $U(t)$ is a square integrable martingale on $[0, T]$. If $\alpha > 0$ then $\mathbb{E}[U, U](\infty) < \infty$, and $U(t)$ is a square integrable martingale on $[0, \infty]$. Since it is uniformly integrable it converges to a non-degenerate limit. \square

The next result uses diffusion theory to find the probability of extinction.

Theorem 13.4: *Let $X(t)$ solve (13.1) and $X(0) = x > 0$. Then the probability of ultimate extinction is $e^{-\frac{2\alpha}{\sigma^2}x}$ if $\alpha > 0$, and 1 if $\alpha \leq 0$.*

PROOF. Let $T_0 = \tau = \inf\{t : X(t) = 0\}$ be the first hitting time of zero, and T_b the first hitting time of $b > 0$. According to the formula (6.52) on exit probabilities for $0 < x < b$

$$P_x(T_0 < T_b) = \frac{S(b) - S(x)}{S(b) - S(0)},$$

where $S(x)$ is the scale function, see (6.50). In this example

$$S(x) = \int_{x_1}^x e^{-\int_{x_0}^u \frac{2\alpha}{\sigma^2} dy} du,$$

where x_0 and x_1 are arbitrary positive constants. Simplifying we obtain

$$P_x(T_0 < T_b) = \frac{e^{-cx} - e^{-cb}}{1 - e^{-cb}},$$

with $c = 2\alpha/\sigma^2$. The probability of ultimate extinction is obtained by taking the limit as $b \rightarrow \infty$, that is,

$$P_x(T_0 < T_\infty) = \lim_{b \rightarrow \infty} P_x(T_0 < T_b) = \begin{cases} e^{-\frac{2\alpha}{\sigma^2}x} & \text{if } \alpha > 0, \\ 1 & \text{if } \alpha \leq 0, \end{cases}$$

where $T_\infty = \lim_{b \rightarrow \infty} T_b$ is the explosion time, which is infinite if explosion does not occur. Using the test for explosion Theorem 6.23, there is no explosion and $T_\infty = \infty$. \square

By a result of Lamperti, Example 7.17, a branching diffusion is a time-changed Brownian motion with the drift stopped at the time it hits zero.

13.2 Wright–Fisher Diffusion

In population dynamics frequencies of genes or alleles are studied. It is assumed for simplicity that the population size N is fixed and individuals are of two types: A and a . If individuals of type A mutate to type a with the rate γ_1/N , and individuals of type a mutate to type A with the rate γ_2/N , then it is possible to approximate the frequency of type A individuals $X(t)$ by the Wright–Fisher diffusion, given by the stochastic equation

$$dX(t) = (-\gamma_1 X(t) + \gamma_2(1 - X(t)))dt + \sqrt{X(t)(1 - X(t))}dB(t),$$

with $0 < X(0) < 1$. For a complete description of the process, its behaviour at the boundaries, 0 and 1, should also be specified. Here we have that when $X(t) = 0$ or 1, then all individuals at time t are of the same type. Consider first the case of no mutation: $\gamma_1 = \gamma_2 = 0$. Then the equation for $X(t)$ is

$$dX(t) = \sqrt{X(t)(1 - X(t))}dB(t),$$

with $0 < X(0) = x < 1$. The scale function is given by $S(x) = x$, consequently for $0 \leq a < x < b \leq 1$

$$P_x(T_b < T_a) = \frac{x - a}{b - a}.$$

The process is stopped when it reaches 0 or 1 because at that time, and at all future times, all individuals are of the same type. This phenomenon is called fixation. The probability that fixation at A occurs having started with a proportion x of type A individuals is x , and with the complementary probability fixation at a occurs.

The expected time to fixation is found by Theorem 6.16 as the solution to $Lv = -1$ with boundary conditions $v(0) = v(1) = 0$, and $Lv = \frac{x(1-x)}{2}v''$. Solving this equation (using $\int \ln x = x \ln x - x$), we obtain that the expected

time to fixation having started with proportion x of type A individuals is given by

$$v(x) = \mathbb{E}_x \tau = -2((1-x)\ln(1-x) + x\ln x).$$

In the model with one-way mutation when, for example, $\gamma_2 = 0$, $\gamma_1 = \gamma > 0$, $X(t)$ satisfies

$$dX(t) = -\gamma X(t)dt + \sqrt{X(t)(1-X(t))}dB(t),$$

with $0 < X(0) < 1$. The process is stopped once it reaches the boundaries 0 or 1. In this case the scale function is given by $S(x) = (1 - (1-x)^{1-2\gamma})/(1-2\gamma)$ if $\gamma \neq 1/2$ and $S(x) = -\log(1-x)$ when $\gamma = 1/2$. Note that by continuity of paths $T_b \uparrow T_1$ as $b \uparrow 1$, and it is easy to see that if $\gamma \geq 1/2$ then

$$\mathbb{P}_x(T_1 < T_0) = \lim_{b \uparrow 1} \mathbb{P}_x(T_b < T_0) = 0.$$

It is possible to see by Theorem 6.16 that the expected time to fixation is finite. Thus fixation at type a occurs with probability 1. If $\gamma < 1/2$, then the expected time to fixation is finite, but there is a positive probability that fixation at type A also occurs.

In the model with two-way mutation, $\gamma_1, \gamma_2 > 0$. Analysis of this model is done using the diffusion processes techniques described in Chapter 6, but it is too involved to be given here in detail. The important feature of this model is that fixation does not occur and $X(t)$ admits a stationary distribution. Stationary distributions can be found by formula (6.69). We find

$$\pi(x) = \frac{C}{\sigma^2(x)} \exp\left(\int_{x_0}^x \frac{2\mu(y)}{\sigma^2(y)} dy\right) = C(1-x)^{2\gamma_1-1}x^{2\gamma_2-1},$$

which is the density of a beta distribution. $C^{-1} = \Gamma(2\gamma_1)\Gamma(2\gamma_2)/\Gamma(2\gamma_1+2\gamma_2)$.

Diffusion models find frequent use in population genetics, see, for example, Ewens (1979). For more information on Wright–Fisher diffusion see, for example, Karlin and Taylor (1981).

Remark 13.1: The theory of weak convergence is used to show that a diffusion approximation to allele frequencies is valid. This theory is not presented here but can be found in many texts, see, for example, Liptser and Shiryaev (1989), and Ethier and Kurtz (1986). More detailed analysis of weak convergence will be given in an accompanying text (Hamza and Klebaner (forthcoming 2013)).

13.3 Birth–Death Processes

Definition

A birth–death process is a model for a developing-in-time population of particles. Each particle lives for a random length of time at the end of which it splits into two particles or dies. If there are x particles in the population, then the lifespan of a particle has an exponential distribution with parameter $a(x)$, the split into two occurs with probability $p(x)$ and with the complementary probability $1 - p(x)$ a particle dies producing no offspring.

Denote by $X(t)$ the population size at time t . The change in the population occurs only at the death of a particle, and from state x the process jumps to $x+1$ if a split occurs or to $x-1$ if a particle dies without splitting. Thus the jump variables $\xi(x)$ take values 1 and -1 with probabilities $p(x)$ and $1-p(x)$ respectively. Using the fact that the minimum of exponentially distributed random variables is exponentially distributed, we can see that the process stays at x for an exponentially distributed length of time with parameter $\lambda(x) = xa(x)$.

Usually, the birth–death process is described in terms of birth and death rates; in a population of size x , a particle is born at rate $b(x)$ and dies at the rate $d(x)$. These relate to the infinitesimal probabilities of population increasing and decreasing by one, namely for an integer $x \geq 1$

$$\mathrm{P}(X(t+\delta) = x+1 | X(t) = x) = b(x)\delta + o(\delta), \quad (13.3)$$

$$\mathrm{P}(X(t+\delta) = x-1 | X(t) = x) = d(x)\delta + o(\delta), \quad (13.4)$$

and with the complementary probability that no births or deaths happen in $(t, t+\delta)$ and the process remains unchanged

$$\mathrm{P}(X(t+\delta) = x | X(t) = x) = 1 - (b(x) + d(x))\delta + o(\delta).$$

Here $o(\delta)$ denotes a function (that may depend on x) such that $\lim_{\delta \rightarrow 0} o(\delta)/\delta = 0$. Note that $b(x)$ and $d(x)$ are rates, not probabilities. The state 0 is usually taken to be absorbing ($b(0) = d(0) = 0$), but in some models transition from 0 to 1 ($b(0) > 0$, $d(0) = 0$) is possible.

Stochastic Equation

It can be seen that these assumptions lead to the model of a Markov jump process (see Section 9.5) that stays at x for an exponential time with parameter $\lambda(x)$

$$\lambda(x) = b(x) + d(x), \quad (13.5)$$

and jump from x according to

$$\xi(x) = \begin{cases} +1 & \text{with probability } \frac{b(x)}{b(x)+d(x)} \\ -1 & \text{with probability } \frac{d(x)}{b(x)+d(x)}. \end{cases} \quad (13.6)$$

The first two moments of the jumps are easily computed to be

$$m(x) = \frac{b(x) - d(x)}{b(x) + d(x)} \quad \text{and} \quad v(x) = 1. \quad (13.7)$$

According to Theorem 9.16 the compensator of $X(t)$ is given in terms of

$$\lambda(x)m(x) = b(x) - d(x),$$

which is the survival rate in the population. Thus the birth–death process $X(t)$ can be written as a stochastic equation

$$X(t) = X(0) + \int_0^t (b(X(s)) - d(X(s))) ds + M(t), \quad (13.8)$$

where $M(t)$ is a local martingale with sharp bracket

$$\langle M, M \rangle(t) = \int_0^t (b(X(s)) + d(X(s))) ds. \quad (13.9)$$

Since $|\xi(x)| = 1$, it follows by Theorem 9.18 that if the linear growth condition

$$b(x) + d(x) \leq C(1 + x), \quad (13.10)$$

holds then the local martingale M in representation (13.8) is a square integrable martingale. Remark here that square integrability requires $\lambda(x) \leq C(1 + x^2)$, which is weaker than (13.10) for integer valued processes.

It follows from Theorem 9.18 and its corollary that the first two moments of the process are given by

$$EX(t) = EX(0) + E \int_0^t (b(X(s)) - d(X(s))) ds, \quad (13.11)$$

and

$$\begin{aligned} EX^2(t) &= EX^2(0) + E \int_0^t \lambda(X(s))(v(X(s)) + 2X(s)m(X(s))) ds \\ &= EX^2(0) + E \int_0^t (\lambda(X(s)) + 2X(s)\lambda(X(s))m(X(s))) ds. \end{aligned} \quad (13.12)$$

Remark 13.2: Sometimes it is convenient to describe the model in terms of the *individual* birth and death rates. In a population of size x , each particle is born at the rate $\beta(x)$ and dies at the rate $\gamma(x)$. The population rates relate to the individual rates by multiplication by the size of the population x ,

$$b(x) = x\beta(x), \quad \text{and} \quad d(x) = x\gamma(x). \quad (13.13)$$

Introduce the individual survival rate $\alpha(x) = \beta(x) - \gamma(x)$. Then the stochastic Equation (13.8) becomes

$$X(t) = X(0) + \int_0^t \alpha(X(s))X(s)ds + M(t).$$

Generator of a Birth-Death Process

Let us write a representation for $f(X(t))$ for a bounded function f as a sum of its compensator and a martingale. According to Theorem 9.20

$$f(X(t)) = f(X(0)) + \int_0^t Lf(X(s))ds + M^f(t), \quad (13.14)$$

where

$$\begin{aligned} Lf(x) &= \lambda(x)\mathbb{E}(f(x + \xi(x)) - f(x)) \\ &= b(x)f(x + 1) + d(x)f(x - 1) - (b(x) + d(x))f(x), \end{aligned} \quad (13.15)$$

is the generator of $X(t)$ and $M^f(t)$ is a local martingale with quadratic variation given by

$$\langle M^f, M^f \rangle(t) = \int_0^t \left(Lf^2(X(s)) - 2f(X(s))Lf(X(s)) \right) ds.$$

Under the linear growth condition on the rates (13.10), by Theorem 9.22 M^f is also a martingale. It follows directly from the above equation that if both f and Lf are bounded, then M^f is bounded (on bounded time intervals) as a difference of bounded functions, and as such is a martingale. If f is compactly supported (is zero outside a finite interval), then Lf is also compactly supported and M^f is a martingale. When M^f is a martingale Dynkin's formula holds

$$\begin{aligned} \mathbb{E}f(X(t)) &= \mathbb{E}f(X(0)) + \mathbb{E} \int_0^t Lf(X(s))ds \\ &= \mathbb{E}f(X(0)) + \int_0^t \mathbb{E}(Lf(X(s)))ds, \end{aligned} \quad (13.16)$$

where the expectation is taken inside the integral by Fubini's theorem. In this formula, a fixed time t can be replaced by a bounded stopping time.

The above equations describe the dynamics of the process. They are useful for approximations (limit theorems) and for finding martingales. Using appropriate martingales one can find exit probabilities, mean exit times from intervals, similarly to the case of diffusions. In this case, instead of second order differential equations we have second order difference equations, as shown below.

Forward Kolmogorov Equations

These are equations for the probability $P_i(X(t) = j) = P(X(t) = j | X(0) = i)$ that the process $X(t)$ will be in state j at time t when started at time zero in state i . Consider the function $f_j(x) = I(x = j)$, the indicator of $\{j\}$. Then

$$\begin{aligned} Lf_j(x) &= b(x)f_j(x+1) + d(x)f_j(x-1) - (b(x) + d(x))f_j(x) \\ &= b(j-1)I(x=j-1) + d(j+1)I(x=j+1) - (b(j) + d(j))I(x=j). \end{aligned}$$

For $j = 0$, the above also holds by taking $b(-1) = 0$; $d(0) = 0$ by definition. Since $f_j(x)$ and Lf_j are bounded, Dynkin's formula holds. Using it and noting that the expectation of the indicator is the probability, $Ef_j(X(t)) = E(I(X(t) = j)) = P(X(t) = j)$, we have

$$\begin{aligned} P(X(t) = j) &= P(X(0) = j) + \int_0^t \left(b(j-1)P(X(s) = j-1) \right. \\ &\quad \left. + d(j+1)P(X(s) = j+1) - (b(j) + d(j))P(X(s) = j) \right) ds. \end{aligned} \tag{13.17}$$

With notation $P_{ij}(t) = P_i(X(t) = j)$ we have the following system of equations

$$P_{ij}(t) = P_{ij}(0) + \int_0^t \left(b(j-1)P_{i,j-1}(s) + d(j+1)P_{i,j+1}(s) - (b(j) + d(j))P_{ij}(s) \right) ds.$$

In differential form, these equations are known as the Kolmogorov forward equations.

$$P'_{ij}(t) = b(j-1)P_{i,j-1}(t) + d(j+1)P_{i,j+1}(t) - (b(j) + d(j))P_{ij}(t).$$

For $j = 0$, take $b(-1) = 0$ and $d(0) = 0$.

Stationary Distribution

The distribution of $X(0)$ together with the transition probabilities define a probability measure on the space of paths. For given transition probabilities

an initial distribution is said to be stationary (or invariant) if for all $t > 0$, $X(t)$ has the same distribution as $X(0)$. Let π be the distribution of $X(0)$ and assume that π is a stationary distribution. Recall Equation (13.17)

$$\begin{aligned} \text{P}(X(t) = j) &= \text{P}(X(0) = j) + \int_0^t \left(b(j-1)\text{P}(X(s) = j-1) \right. \\ &\quad \left. + d(j+1)\text{P}(X(s) = j+1) - (b(j) + d(j))\text{P}(X(s) = j) \right) ds. \end{aligned}$$

Using stationarity, $\text{P}(X(t) = j) = \pi(j)$ for all t , and it follows that the expression under the integral must be zero. Hence the following system of equations is obtained

$$b(j-1)\pi(j-1) + d(j+1)\pi(j+1) - (b(j) + d(j))\pi(j) = 0.$$

Here again $b(-1) = 0$. This system is solved by induction.

$$\pi(j) = \prod_{x=0}^{j-1} \frac{b(x)}{d(x+1)} \pi(0). \quad (13.18)$$

($\prod_{x=0}^{-1} = 1$). $\pi(0)$ is found by equating the sum of the probabilities $\pi(j)$ to 1.

$$\pi(0) = \left(\sum_{i=0}^{\infty} \prod_{x=0}^{i-1} \frac{b(x)}{d(x+1)} \right)^{-1}. \quad (13.19)$$

This shows that when stationary distribution exists then the series of products above must necessarily converge.

13.4 Growth of Birth–Death Processes

Growth of Birth–Death Processes with Linear Rates

Suppose that the *individual* birth and death rates β and γ are constants. This corresponds to linear rates $b(x) = \beta x$, $d(x) = \gamma x$ (see (13.13)). The linear growth condition (13.10) is satisfied, and we have from Equation (13.8)

$$X(t) = X(0) + \alpha \int_0^t X(s)ds + M(t),$$

where $\alpha = \beta - \gamma$ is the survival rate and $M(t)$ is a martingale. Moreover, by (13.9)

$$\langle M, M \rangle(t) = (\beta + \gamma) \int_0^t X(s)ds.$$

Hence by taking expectations we obtain that

$$EX(t) = X(0) + \alpha \int_0^t EX(s)ds.$$

Solving this equation we find

$$EX(t) = X(0)e^{\alpha t}.$$

Using integration by parts (note that $e^{\alpha t}$ has zero covariation with $X(t)$), we have

$$d\left(X(t)e^{-\alpha t}\right) = e^{-\alpha t}dX(t) - \alpha e^{-\alpha t}X(t-)dt.$$

Since $X(t-)dt$ can be replaced by $X(t)dt$, we obtain

$$d\left(X(t)e^{-\alpha t}\right) = e^{-\alpha t}dX(t) - \alpha e^{-\alpha t}X(t)dt = e^{-\alpha t}dM(t).$$

Thus

$$X(t)e^{-\alpha t} = X(0) + \int_0^t e^{-\alpha s}dM(s).$$

Using the rule for the sharp bracket of the integral we find

$$\begin{aligned} \left\langle \int_0^\cdot e^{-\alpha s}dM(s), \int_0^\cdot e^{-\alpha s}dM(s) \right\rangle(t) &= \int_0^t e^{-2\alpha s}d\langle M, M \rangle(s) \\ &= (\beta + \gamma) \int_0^t e^{-2\alpha s}X(s)ds. \end{aligned}$$

Since $EX(t) = X(0)e^{\alpha t}$, by taking expectations it follows for $\alpha > 0$ that

$$E \left\langle \int_0^\cdot e^{-\alpha s}dM(s), \int_0^\cdot e^{-\alpha s}dM(s) \right\rangle(\infty) < \infty.$$

Consequently, $X(t)e^{-\alpha t}$ is a square integrable martingale on $[0, \infty]$.

$$\begin{aligned} Var(X(t)e^{-\alpha t}) &= E \left\langle \int_0^\cdot e^{-\alpha s}dM(s), \int_0^\cdot e^{-\alpha s}dM(s) \right\rangle(t) \\ &= X(0) \frac{\beta + \gamma}{\alpha} (1 - e^{-\alpha t}). \end{aligned}$$

Since $X(t)e^{-\alpha t}$ is a square integrable martingale on $[0, \infty]$, it converges almost surely as $t \rightarrow \infty$ to a non-degenerate limit.

Growth of Processes with Stabilizing Reproduction

Consider the case when the rates stabilize as population size increases, namely $\beta(x) \rightarrow \beta$ and $\gamma(x) \rightarrow \gamma$ as $x \rightarrow \infty$. Then, clearly, $\alpha(x) = \beta(x) - \gamma(x) \rightarrow \alpha = \beta - \gamma$. Depending on the value of α radically distinct modes of behaviour occur. If $\alpha < 0$, then the process dies out with probability one. If $\alpha > 0$, then the process tends to infinity with a positive probability. If the rate of convergence of $\alpha(x)$ to α is fast enough, then exponential growth persists as in the classical case. Indeed, under some technical conditions the following condition is necessary and sufficient for exponential growth,

$$\int_1^\infty \frac{|\alpha(x) - \alpha|}{x} dx < \infty.$$

If it holds, then $X(t)e^{-\alpha t}$ converges almost surely and in the mean square to a non-degenerate limit.

The case $\alpha = 0$ provides examples of slowly growing populations, such as those with a linear rate of growth. Let $\alpha(x) > 0$ and $\alpha(x) \downarrow 0$. Among such processes there are some that become extinct with probability one, but there are others that have a positive probability of growing to infinity. Consider the case when $\alpha(x) = c/x$ when $x \geq 1$. It is possible to show that if $c > 1/2$, then $q = P(X(t) \rightarrow \infty) > 0$. The stochastic equation for $X(t)$ becomes

$$X(t) = X(0) + c \int_0^t I(X(s) > 0) ds + M(t).$$

By taking the expectations above one can show that $\lim_{t \rightarrow \infty} EX(t)/t = cq$, and the mean of such processes grows linearly. By using Theorem 9.22 and Exercise 9.5 it is possible to show that for any k , $E(X^k(t))/t^k$ converges to the k -th moment of a gamma distribution multiplied by q . This implies that on the set $\{X(t) \rightarrow \infty\}$, $X(t)/t$ converges in distribution to a gamma distribution. In other words, such processes grow linearly, and not exponentially. By changing the rate of convergence of $\alpha(x)$ other polynomial growth rates of the population can be obtained.

Similar results are available for population-dependent Markov branching processes that generalize birth-death processes by allowing a random number of offspring at the end of a particle's life. For details of the stochastic equation approach to Markov jump processes see Klebaner (1994), and Hamza and Klebaner (1995a), where such processes were studied as randomly perturbed linear differential equations.

13.5 Extinction, Probability, and Time to Exit

Probability of Extinction

Let $X(t)$ be a birth-death process with zero being an absorbing state, $b(0) = 0$, and all other states having positive rates. Let $T = T_0 = \inf\{t \geq 0 : X(t) = 0\}$ be the time to hit zero; if there is no such t then we take $T = \infty$. Since $b(0) = 0$, when zero is reached the process will stay there forever after. This event is called extinction. Denote by $u(x) = P_x(T < \infty)$, the probability to be absorbed when the process starts at state x . By definition, $u(0) = 1$.

Theorem 13.5: *Let $X(t)$ be a birth-death process with generator L , $X(0) = x$. Then $u(x) = P_x(T < \infty)$ solves $Lu = 0$, $u(0) = 1$. Furthermore, either $u(x) = 1$ for all x , or else $u(x) < 1$ for all $x \geq 1$, u is strictly decreasing and $u(\infty) = \lim_{x \rightarrow \infty} u(x) = 0$.*

PROOF. Conditioning on the first transition from $x \geq 1$ we obtain that

$$u(x) = \frac{b(x)}{b(x) + d(x)}u(x+1) + \frac{d(x)}{b(x) + d(x)}u(x-1). \quad (13.20)$$

According to the formula for the generator (13.15) this is the same as $Lu(x) = 0$. Re-arranging this formula we obtain for all $x \geq 1$

$$u(x+1) - u(x) = \frac{d(x)}{b(x)}(u(x) - u(x-1)). \quad (13.21)$$

If $u(1) = 1$, then from the above equation $u(2) - u(1) = 0$, and by induction, $u(x+1) = u(x) = 1$ for all x . If $u(1) < u(0) = 1$, then $u(2) - u(1) < 0$, and by induction, $u(x+1) < u(x)$ for all x . In this case we consider $u(X(t))$. Using Equation (13.14) we have $u(X(t)) = u(X(0)) + M^u(t)$. Hence $Y(t) = u(X(t))$ is a local martingale. Since it is bounded, it is a true martingale. According to the martingale convergence theorem it converges as $t \rightarrow \infty$ to a finite limit. Since u is strictly decreasing it is one to one, and it follows that $X(t)$ also converges, possibly to ∞ . Since $X(t)$ is integer valued, convergence to a finite value means that $X(t)$ assumes some value and remains there forever, after some time. This implies that the only possible states for the limit are the absorbing states. Since by assumption only 0 is absorbing, the limit of $X(t)$ is either 0 or ∞ . Thus $u(X(t))$ converges to $u(0) = 1$ or $u(\infty)$. Now using the optional stopping theorem for the martingale $u(X(t))$ we have $u(x) = Eu(X(T)) = u(0)P_x(T < \infty) + u(\infty)P_x(T = \infty) = u(x) + u(\infty)P_x(T = \infty)$. Since $P_x(T = \infty) = 1 - u(x) > 0$, we must have $u(\infty) = 0$. \square

It is easy to solve the recurrence relation (13.21) and we have

Corollary 13.6: If $\sum_{j=1}^{\infty} \prod_{i=1}^j \frac{d(i)}{b(i)} < \infty$, then $u(x) = \frac{\sum_{j=x}^{\infty} \prod_{i=1}^j \frac{d(i)}{b(i)}}{1 + \sum_{j=1}^{\infty} \prod_{i=1}^j \frac{d(i)}{b(i)}}$, and if the series diverges, then $u(x) = 1$ for all x .

PROOF. Iterations of (13.21) give (with $k(i) = d(i)/b(i)$)

$$\begin{aligned} u(x+1) &= u(x) + (u(1) - u(0)) \prod_{i=1}^x k(i) \\ &= u(1) + (u(1) - u(0)) \sum_{j=1}^x \prod_{i=1}^j k(i). \end{aligned} \quad (13.22)$$

Here we have $u(0) = 1$. If the series $\sum_{j=1}^{\infty} \prod_{i=1}^j k(i)$ converges, then we use $u(\infty) = 0$ to find $u(1)$ and then $u(x)$. If the series diverges, then the above equation shows that $u(1)$ must be 1 to have finite $u(x)$. This implies $u(x) = 1$ for all x . \square

Example 13.1: In the case of linear rates, when $b(x) = bx$ and $d(x) = dx$, for positive constants b, d , the extinction probabilities $u(x) = 1$ if $b \leq d$, and $u(x) = \left(\frac{d}{b}\right)^x$ if $b > d$, $x = 0, 1, \dots$

Mean Exit Time

Let $X(t)$ be a birth-death process and m, n two integers. Denote by τ the exit time from the interval (m, n) , when $X(0) = x \in (m, n)$, $\tau = \inf\{t \geq 0 : X(t) \notin (m, n)\}$.

Theorem 13.7: Let L be the generator of $X(t)$, and assume that equation $Lv = -1$, for integers $x \in [m, n]$, with boundary conditions $v(m) = v(n) = 0$ has a solution, $v(x)$. Then $v(x) = E_x \tau$.

PROOF. Using Equation (13.14) with v and that $Lv = -1$, we obtain

$$v(X(t)) = v(X(0)) + \int_0^t Lv(X(s))ds + M^v(t) = v(x) - t + M^v(t),$$

where $M^v(t)$ is a local martingale. Applying the above to $t \wedge \tau$, we get

$$v(X(t \wedge \tau)) = v(x) - t \wedge \tau + M^v(t \wedge \tau).$$

Since $X(t \wedge \tau)$ takes finitely many values within $[m, n]$, so does $v(X(t \wedge \tau))$. Denote a bound for v by K . As $t \wedge \tau \leq t$ we obtain from the above that $M^v(t \wedge \tau)$ is bounded (by $2K + t$). Hence it is a martingale and taking expectations in the above equation we have

$$E_x v(X(t \wedge \tau)) = v(x) - E_x(t \wedge \tau).$$

Now, since $\lim_{t \rightarrow \infty} t \wedge \tau = \tau$ (possibly infinite), by Fatou's lemma

$$\mathbb{E}_x \tau = \mathbb{E}_x \liminf_{t \rightarrow \infty} (t \wedge \tau) \leq \liminf_{t \rightarrow \infty} \mathbb{E}_x (t \wedge \tau) \leq 2K < \infty,$$

because of the previous equation. Hence $\tau < \infty$. By taking limits in the previous equation, and using that $t \wedge \tau$ becomes τ , $X(t \wedge \tau)$ becomes $X(\tau)$, and $v(X(\tau)) = 0$, we obtain the result, by dominated convergence. \square

A situation when the exit time from (m, n) is infinite occurs when for some j and k , $m < j < x < k < n$, $d(j) = b(k) = 0$. In this case the process is trapped between j and k . Assumptions of the following theorem prevent this from happening.

Theorem 13.8: *Provided either $b(x) > 0$ for all $x \in [m, n]$ or $d(x) > 0$ for all $x \in [m, n]$, there is a unique solution to the equation $Lv = -1$ with $v(m) = v(n) = 0$. For the case $b(x) > 0$ it is given by*

$$v(x) = \sum_{l=m}^{x-1} \left[w \prod_{i=m}^l \frac{d(i)}{b(i)} - \sum_{j=m}^l \frac{1}{b(j)} \prod_{i=j+1}^l \frac{d(i)}{b(i)} \right], \quad (13.23)$$

$$w = \sum_{l=m}^{n-1} \sum_{j=m}^l \frac{1}{b(j)} \prod_{i=j+1}^l \frac{d(i)}{b(i)} \Bigg/ \sum_{l=m}^{n-1} \prod_{i=m}^l \frac{d(i)}{b(i)}.$$

PROOF. Assume that $b(x) > 0$, the case $d(x) > 0$ is similar. $Lv = -1$ can be written for $x \geq 1$

$$b(x)(v(x+1) - v(x)) = d(x)(v(x) - v(x-1)) - 1,$$

which holds for $x = m, m+1, \dots, n$. Letting $w(x) = v(x) - v(x-1)$ we have with $k(x) = d(x)/b(x)$,

$$w(x+1) = k(x)w(x) - 1/b(x).$$

The statement follows from solving a linear difference equation with given coefficients. \square

Exit Probabilities

Let T_m denote the first hitting of m . Then as shown in Theorems 13.7 and 13.8 the process exits (m, n) with probability one, so that τ , the exit time from the interval (m, n) is $\tau = T_m \wedge T_n$ is finite with probability one.

Theorem 13.9: Let $X(t)$ be a birth-death process with generator L and S solve $LS = 0$. Then for $x \in [m, n]$

$$P_x(T_n < T_m) = \frac{S(x) - S(m)}{S(n) - S(m)} = \frac{\sum_{j=m+1}^x \prod_{i=1}^j \frac{d(i)}{b(i)}}{\sum_{j=m+1}^n \prod_{i=1}^j \frac{d(i)}{b(i)}}.$$

PROOF. The general solution to $LS = 0$ is given by (13.22)

$$S(x) = S(1) + (S(1) - S(0)) \sum_{j=1}^x \prod_{i=1}^j \frac{d(i)}{b(i)}. \quad (13.24)$$

Now, by using representation (13.14) and the fact that $LS = 0$, we obtain

$$S(X(t \wedge \tau)) = S(x) + M^S(t \wedge \tau).$$

Since $X(t \wedge \tau)$ takes finitely many values, $S(X(t \wedge \tau))$ is bounded and $M^S(t \wedge \tau)$ is a martingale. Taking expectations we get $E_x S(X(t \wedge \tau)) = S(x)$. Taking limits as $t \rightarrow \infty$ and using that $X(\tau) = m$ or n with probabilities that add up to one, we obtain

$$P_x(T_n < T_m) = \frac{S(x) - S(m)}{S(n) - S(m)}.$$

The result now follows from the expression for $S(x)$. \square

13.6 Processes in Genetics

The main models in genetics are the Wright–Fisher and the Moran models. They both consider a population consisting of N individuals, which can be of one of two types. The types correspond to alleles competing for dominance. Many models in cell biology and cancer research are based on the Moran model.

Wright–Fisher Model

The change in frequency of the two types of individuals comprising the population is known as the genetic drift. The Wright–Fisher model describes the genetic drift in a finite population assuming random mating. The time is taken to be discrete so that generations do not overlap. If $X(t)$ is the number of type 1 individuals at time t , then due to random mating $X(t+1)$ has the binomial distribution with parameters N and $X(t)/N$. Thus if for some τ , $X(\tau) = 0$ or N , then for all future times $t > \tau$, $X(t) = 0$ or N . In other words, states 0 and N are absorbing and correspond to the event of fixation, when the whole

population becomes of the same type, type 1 ($X(t) = N$) or type 2 ($X(t) = 0$). It is of interest to find the probability of fixation.

Since the mean of the binomial $\text{Bin}(N, p)$ distribution is Np ,

$$\mathbb{E}(X(t+1)|X(t)) = N \frac{X(t)}{N} = X(t).$$

Hence $X(t)$ is a bounded martingale. $\mathbb{E}X(t) = X(0)$, moreover, $X(t)$ converges as $t \rightarrow \infty$ almost surely to a random variable V . According to dominated convergence $\mathbb{E}V = X(0)$. Next, by using that the second moment of the $\text{Bin}(N, p)$ distribution is $Np(1-p) + (Np)^2$, we obtain by conditioning

$$\mathbb{E}X^2(t+1) = X(0) + \left(1 - \frac{1}{N}\right)\mathbb{E}X^2(t).$$

Since $\mathbb{E}V^2 = \lim_{t \rightarrow \infty} \mathbb{E}X^2(t)$ we obtain by taking the limit in the above equation that $\mathbb{E}V^2 = NX(0)$. Hence $\mathbb{E}(N-V) = 0$. Since $V(N-V) \geq 0$, it follows that $\mathbb{P}(V(N-V) = 0) = 1$. Thus with probability one, $V = 0$ or N . Now note that a limit of integer valued variables is again integer valued, and convergence to an integer valued variable means that from some time onwards, the value is actually assumed. Therefore it follows that fixation is certain, moreover, it occurs in finite time τ . Using the optional stopping theorem we have $\mathbb{E}X(\tau) = X(0)$. However, $X(\tau) = N$ or 0, so that $\mathbb{E}X(\tau) = qN$, where q is the probability of fixation in type 1. Hence the probability that type 1 takes over the population is $q = X(0)/N$.

Moran Model in Discrete Time

The Moran (1958) model assumes overlapping generations. At each time step one individual is chosen to reproduce and one individual is chosen to die. Two individuals are chosen at random, each with probability $1/N$ (the same individual can be chosen for death and for reproduction in the same step). This guarantees that the total size of the population remains constant. All events at a given step are taken to be independent so that the probability that they occur together is given by the product of their probabilities. The rules of reproduction of the new individual determine the process. Denote the number of individuals of type 1 at time t by $X(t)$. First we take time to be discrete, giving a model of a (state-dependent) random walk, and then continuous, giving a model of a birth-death process.

It is clear from the description that the process $X(t)$ is Markov, and that $X(t+1)$ can only take values $X(t) \pm 1$ and $X(t)$. In other words, $X(t)$ is a random walk on integers $0, 1, \dots, N$ with state-dependent probabilities.

Neutral drift

In this model, when an individual is selected for reproduction it produces its own type. Hence the number of type 1 increases by 1 when type 2 is chosen to die and type 1 to reproduce,

$$P(X(t+1) = X(t) + 1 | X(t)) = \left(1 - \frac{X(t)}{N}\right) \frac{X(t)}{N}.$$

$X(t)$ decreases by 1 when type 1 is chosen to die and type 2 to reproduce, hence

$$P(X(t+1) = X(t) - 1 | X(t)) = \frac{X(t)}{N} \left(1 - \frac{X(t)}{N}\right).$$

With the complementary probability, the number of type 1 does not change. It is clear from the above probabilities that if for some t , $X(t) = 0$ or $X(t) = N$, then for all future times the process stays at 0 or N . Thus these states are absorbing. The event of absorption into each of these states is called fixation, and corresponds to the event when the total population becomes of a single type. The fixation of a type 1 individual is the event $X(t) = N$ for some t . By conditioning on the first transition, the probability of this event can be found by writing a difference equation similar to (13.20). Here we show how to find this probability by martingale arguments. Note that the process $X(t)$ is a martingale,

$$\begin{aligned} E(X(t+1) | X(t)) &= (X(t) + 1) \frac{X(t)}{N} \left(1 - \frac{X(t)}{N}\right) + (X(t) - 1) \frac{X(t)}{N} \left(1 - \frac{X(t)}{N}\right) \\ &\quad + X(t) \left(1 - 2 \frac{X(t)}{N} \left(1 - \frac{X(t)}{N}\right)\right) = X(t). \end{aligned}$$

Since it is a bounded martingale, between 0 and N , it converges almost surely to a limit V . Calculations for the second moment give that

$$E(X(t+1))^2 = 2E\frac{X(t)}{N} \left(1 - \frac{X(t)}{N}\right) + E(X(t))^2.$$

Since $X(t)$ is bounded, by dominated convergence $E(X(t))^2$ converges to EV^2 . Taking the limit in the above equation we obtain that $E(X(t)(N - X(t)))$ converges to 0 as $t \rightarrow \infty$. Hence

$$E(V(N - V)) = \lim_{t \rightarrow \infty} E(X(t)(N - X(t))) = 0.$$

Since $V(N - V) \geq 0$, it implies $P(V(N - V) = 0) = 1$, or that $V = 0$ or N . As V is a limit of integer-valued random variables the limit is attained at some finite random time. Thus fixation is certain, and it happens in finite time. The probability of fixation in state N when started in state $X(0) = i$, is obtained from the martingale property of the limit $EV = NP(V = N) = X(0)$. Hence $P_i(V = N) = i/N$.

Mutation

Mutation is when one type produces the other. In the model with mutation each individual produces its own type with probability $1 - \gamma_i$, and the other type with probability γ_i , $i = 1, 2$. The event that $X(t)$ becomes $X(t) + 1$ can happen in two ways. One is when type two is chosen to die and type 1 is chosen to reproduce faithfully (no mutation). The probability of this event is $(1 - \frac{X(t)}{N}) \frac{X(t)}{N} (1 - \gamma_1)$. Another is when type 2 is chosen to die and type 2 to reproduce, and it reproduces type 1 (mutation). The probability of this event is $(1 - \frac{X(t)}{N})^2 \gamma_2$. Thus $X(t)$ increases by 1 with probability

$$\left(1 - \frac{X(t)}{N}\right) \frac{X(t)}{N} (1 - \gamma_1) + \left(1 - \frac{X(t)}{N}\right)^2 \gamma_2.$$

Similarly, $X(t)$ decreases by 1 with probability

$$\frac{X(t)}{N} \left(1 - \frac{X(t)}{N}\right) (1 - \gamma_2) + \left(\frac{X(t)}{N}\right)^2 \gamma_1,$$

i.e. when type 1 was chosen to die and type 2 was chosen to reproduce without mutation, or when type 1 was chosen to die and type 1 to reproduce and mutation occurred. It can be seen that all other combinations of replacements, births, and mutations do not change $X(t)$, so that with the complementary probability the process does not change its state.

When both mutation rates are positive, $\gamma_i > 0$, the process visits all states infinitely often, and there is a stationary distribution. It can be found explicitly by (13.18).

If only one mutation rate is positive and the other one is zero, for example $\gamma_1 = 0$, then state N is the only absorbing state, and the process will be absorbed at this state with probability one. This fact can also be recovered by solving a difference equation, obtained by conditioning on the first transition.

Moran Model in Continuous Time

Here we allow each individual to live for an exponential time with parameter 1. Then the smallest of N such independent lifespans has exponential distribution with parameter N . Therefore $X(t)$, the number of type 1 individuals, is a birth-death process with rates

$$\begin{aligned} b(x) &= N \left[\left(1 - \frac{x}{N}\right) \frac{x}{N} (1 - \gamma_1) + \left(1 - \frac{x}{N}\right)^2 \gamma_2 \right], \\ d(x) &= N \left[\frac{x}{N} \left(1 - \frac{x}{N}\right) (1 - \gamma_2) + \left(\frac{x}{N}\right)^2 \gamma_1 \right]. \end{aligned} \tag{13.25}$$

Note that all states $0 < x < N$ have positive rates (unless $\gamma_1 = 1 - \gamma_2 = 0$ or $\gamma_2 = 1 - \gamma_1 = 0$), and if $\gamma_1, \gamma_2 > 0$, the states 0 and N are reflecting, $b(0) > 0$, $d(0) = 0$, $b(N) = 0$, $d(N) > 0$.

If we assume, as in Karlin and Taylor (1975), that the mutation rates decay with N , $\gamma_1 N \rightarrow \kappa_1$, and $\gamma_2 N \rightarrow \kappa_2$, then we can show convergence to the Wright–Fisher diffusion (see Chapter 13.2).

Stochastic Equation and Its Limit

The equation for $X(t)$ is given by (13.8) with rates given by (13.25).

$$\begin{aligned} X(t) &= X(0) + \int_0^t (b(X(s)) - d(X(s))) ds + M(t) \\ &= X(0) + N \int_0^t \left(\gamma_2 - (\gamma_1 + \gamma_2) \frac{X(s)}{N} \right) ds + M(t). \end{aligned} \quad (13.26)$$

$M(t)$ is a martingale with the sharp bracket

$$\begin{aligned} \langle M, M \rangle(t) &= \int_0^t (b(X(s)) + d(X(s))) ds \\ &= N \int_0^t \left[(2 - \gamma_1 - \gamma_2) \frac{X(s)}{N} \left(1 - \frac{X(s)}{N} \right) + \gamma_2 \left(1 - \frac{X(s)}{N} \right)^2 + \gamma_1 \left(\frac{X(s)}{N} \right)^2 \right] ds. \end{aligned}$$

The next result gives an approximation which assumes a genetic scale Nt for time. Convergence is in the sense of the weak convergence of processes. This is beyond the scope of this book, and more on this topic will be given in the forthcoming accompanying publication by Hamza and Klebaner (forthcoming 2013). We just state here that this convergence implies convergence of finite dimensional distributions, and in particular, for any fixed t , the distribution of $X(Nt)$ can be approximated by using the limiting diffusion at time t .

Theorem 13.10: *Let $X(0)/N$ converge to y in distribution. Then the sequence of processes $X(Nt)/N$ converges in distribution to the Wright–Fisher diffusion, with drift parameter $\mu(x) = \kappa_2 - (\kappa_1 + \kappa_2)x$ and diffusion parameter $\sigma^2(x) = 2x(1 - x)$.*

PROOF. While we do not give a complete proof, the intuition behind this approximation lies in the following calculation. Consider the equation for the process $X(Nt)/N$. We obtain from Equation (13.26)

$$\frac{X(Nt)}{N} = \frac{X(0)}{N} + \int_0^{Nt} \left(\gamma_2 - (\gamma_1 + \gamma_2) \frac{X(s)}{N} \right) ds + \frac{1}{N} M(Nt), \quad (13.27)$$

with quadratic variation of the martingale given by

$$\begin{aligned} \left\langle \frac{1}{N} M, \frac{1}{N} M \right\rangle(Nt) &= \frac{1}{N^2} \langle M, M \rangle(Nt) \\ &= \frac{1}{N} \int_0^{Nt} \left[(2 - \gamma_1 - \gamma_2) \frac{X(s)}{N} \left(1 - \frac{X(s)}{N} \right) + \gamma_2 \left(1 - \frac{X(s)}{N} \right)^2 + \gamma_1 \left(\frac{X(s)}{N} \right)^2 \right] ds. \end{aligned}$$

Now make the change of variables $s = Nu$. Then we have with the re-scaled process $Y^N(t) = \frac{X(Nt)}{N}$

$$Y^N(t) = Y^N(0) + \int_0^t N \left(\gamma_2 - (\gamma_1 + \gamma_2)Y^N(u) \right) du + \frac{1}{N} M(Nt) \quad (13.28)$$

with quadratic variation $\langle Y^N, Y^N \rangle(t)$ that of the martingale $\frac{1}{N} M(Nt)$,

$$\int_0^t \left[(2 - \gamma_1 - \gamma_2)Y^N(u) \left(1 - Y^N(u) \right) + \gamma_2 \left(1 - Y^N(u) \right)^2 + \gamma_1 \left(Y^N(u) \right)^2 \right] du.$$

Since $N\gamma_i \rightarrow \kappa_i$, if we formally take the limit as $N \rightarrow \infty$, we obtain that the drift of $Y^N(t)$ converges to $\int_0^t (\kappa_2 - (\kappa_1 + \kappa_2)Y(u)) du$ and its quadratic variation to $\int_0^t 2Y(u)(1 - Y(u))du$, where Y is the limit of Y^N . It can be shown that processes $Y^N(t)$ converge weakly as $N \rightarrow \infty$ to the diffusion $Y(t)$ given by the following stochastic equation:

$$dY(t) = (\kappa_2 - (\kappa_1 + \kappa_2)Y(t))dt + 2Y(t)(1 - Y(t))dB(t), \quad Y(0) = y. \quad \square$$

Moran Model with Selective Advantage

If one individual in the model has a fitness advantage over the other it will be more likely to be chosen for reproduction. If we let the fitness of type 1 be r and that of type 2 be 1, then the probability that a type 1 individual is chosen for reproduction is $rx/(rx + N - x)$ rather than x/N , and the probability that a type 2 individual is chosen is the complementary probability $(N - x)/(N - x + rx)$. Type 1 has selective advantage if $r > 1$, type 2 if $r < 1$, and when $r = 1$ selection is neutral. This leads to the model of a birth-death process with rates

$$b(x) = \lambda \left[\left(1 - \frac{x}{N} \right) \frac{rx(1 - \gamma_1)}{N - x + rx} + \left(1 - \frac{x}{N} \right) \frac{(N - x)\gamma_2}{N - x + rx} \right],$$

$$d(x) = \lambda \left[\frac{x}{N} \frac{(N - x)(1 - \gamma_2)}{N - x + rx} + \frac{x}{N} \frac{rx\gamma_1}{N - x + rx} \right].$$

When there are no mutations there are two absorbing states 0 and N . It is not hard to show that as $t \rightarrow \infty X(t) \rightarrow V$ with $P(V = 0) + P(V = N) = 1$. The fixation probability can be calculated for this model, and starting in state i ,

$$P_i(V = N) = \frac{1 - r^{-i}}{1 - r^{-N}}.$$

Suppose that type 1 is a new mutant, then the probability that it will take over the whole population is the probability of its fixation, and it is given by

$$\rho = P_1(V = N) = \frac{1 - r^{-1}}{1 - r^{-N}}.$$

When $r = 1$ this probability is given by $1/N$ (the limit of the above expression as $r \rightarrow 1$). This is used in cancer models, e.g. Komarova (2007).

13.7 Birth–Death Processes in Many Dimensions

These are Markov processes taking values $\mathbf{x}(t) = (x_1(t), \dots, x_d(t))$ with each $x_i(t) \geq 0$ and integer valued. At each such state the process remains for an exponentially distributed length of time with parameter $\lambda(\mathbf{x})$ after which it jumps to a new state in which a coordinate is decreased or increased by 1. That is, the jump from \mathbf{x} , $\mathbf{w}(\mathbf{x}) = \pm \mathbf{e}_i$ the i -th d -dimensional coordinate vector. As in the one-dimensional case, the probabilities $P(\mathbf{w}(\mathbf{x}) = \pm \mathbf{e}_i)$ are given by the ratio of the rates, $b_i(\mathbf{x})$ (the i -th birth rate, the rate at which the i -th coordinate increases by 1) and $d_i(\mathbf{x})$ (the i -th death rate, the rate at which the i -th coordinate decreases by 1) and $\lambda(\mathbf{x}) = \sum_{i=1}^d (b_i(\mathbf{x}) + d_i(\mathbf{x}))$. Birth–death processes in many dimensions are hard to treat analytically, and in applications are studied mostly by simulations. The above description of the process is convenient for simulations. The process stays in a state \mathbf{x} for an exponentially distributed time with parameter $\lambda(\mathbf{x})$. Then the chance for each coordinate to go up or down is $b_i(\mathbf{x})/\lambda(\mathbf{x})$ or $d_i(\mathbf{x})/\lambda(\mathbf{x})$. This procedure is either repeated until a certain event is observed, or it is repeated for a given length of time.

Here we set up the stochastic equation for such processes. These equations are useful for certain analytical derivations, and especially for studying the limiting behaviour (approximations).

The generator of the process L acts on scalar functions of d variables (from \mathbb{R}^d into \mathbb{R}) and is defined by

$$\begin{aligned} Lf(\mathbf{x}) &= \lambda(\mathbf{x})E\left(f(\mathbf{x} + \mathbf{w}(\mathbf{x})) - f(\mathbf{x})\right) \\ &= \sum_{i=1}^d (f(\mathbf{x} + \mathbf{e}_i) - f(\mathbf{x}))b_i(\mathbf{x}) + (f(\mathbf{x} - \mathbf{e}_i) - f(\mathbf{x}))d_i(\mathbf{x}). \end{aligned} \tag{13.29}$$

For a bounded f we can write

$$f(\mathbf{X}(t)) = f(\mathbf{X}(0)) + \int_0^t Lf(\mathbf{X}(s))ds + M^f(t), \tag{13.30}$$

where $M^f(t)$ is a local martingale, and it is a martingale for a compactly supported f . It can be shown that, just like in one dimension, the quadratic variation of M^f is given by

$$\langle M^f, M^f \rangle(t) = \int_0^t \Gamma(f, f)(\mathbf{X}(s))ds, \quad (13.31)$$

where

$$\Gamma(f, f)(\mathbf{x}) = Lf^2(\mathbf{x}) - 2f(\mathbf{x})Lf(\mathbf{x}) = \lambda(\mathbf{x})E\left(f(\mathbf{x} + \mathbf{w}(\mathbf{x})) - f(\mathbf{x})\right)^2. \quad (13.32)$$

More generally, for two functions f and g

$$\langle M^f, M^g \rangle(t) = \int_0^t \Gamma(f, g)(\mathbf{X}(s))ds, \quad (13.33)$$

where

$$\begin{aligned} \Gamma(f, g)(\mathbf{x}) &= Lfg(\mathbf{x}) - f(\mathbf{x})Lg(\mathbf{x}) - g(\mathbf{x})Lf(\mathbf{x}) \\ &= \lambda(\mathbf{x})E\left((f(\mathbf{x} + \mathbf{w}(\mathbf{x})) - f(\mathbf{x}))(g(\mathbf{x} + \mathbf{w}(\mathbf{x})) - g(\mathbf{x}))\right) \\ &= \sum_{i=1}^d (f(\mathbf{x} + \mathbf{e}_i) - f(\mathbf{x}))(g(\mathbf{x} + \mathbf{e}_i) - g(\mathbf{x}))b_i(\mathbf{x}) \quad (13.34) \\ &\quad + \sum_{i=1}^d (f(\mathbf{x} - \mathbf{e}_i) - f(\mathbf{x}))(g(\mathbf{x} - \mathbf{e}_i) - g(\mathbf{x}))d_i(\mathbf{x}). \end{aligned}$$

Formula (13.30) also holds for some unbounded functions provided a growth condition holds, just like in the one-dimensional case (9.50) of Theorem 9.22. Taking $f(\mathbf{x}) = x_j$, the j -th coordinate function, we obtain the equation for $X_j(t)$

$$X_j(t) = X_j(0) + \int_0^t (b^j(\mathbf{X}(s)) - d^j(\mathbf{X}(s)))ds + M_j(t), \quad (13.35)$$

$$\langle M^j, M^j \rangle(t) = \int_0^t (b^j(\mathbf{X}(s)) + d^j(\mathbf{X}(s)))ds. \quad (13.36)$$

The above equations can be written in vector form

$$\mathbf{X}(t) = \mathbf{X}(0) + \int_0^t (\mathbf{b}(\mathbf{X}(s)) - \mathbf{d}(\mathbf{X}(s)))ds + \mathbf{M}(t), \quad (13.37)$$

with $\langle M^i, M^i \rangle(t)$ given as above. Note that it follows from (13.34) $\langle M^i, M^j \rangle(t) = 0$ for $j \neq i$; this signifies the fact that two coordinates cannot jump simultaneously.

When the local martingale M_t^f in (13.30) is a true martingale (similar to conditions in Theorem 9.22) Dynkin's formula holds

$$\mathbb{E}f(\mathbf{X}(t)) = \mathbb{E}f(\mathbf{X}(0)) + \mathbb{E} \int_0^t Lf(\mathbf{X}(s))ds. \quad (13.38)$$

Birth–death processes in one and many dimensions are used in modelling the evolution of cell populations, and in particular, the development of cancer.

13.8 Cancer Models

There are many stochastic models that address various aspects of cancer formation and development. The literature on the subject is vast, see, for example, Tan and Hanin (2008). Here we give a very brief introduction to models where stochastic calculus methods can be applied.

Cancer Development

Statistical analysis has lead to the multi-stage theory of cancer, which states that cancer forms as a result of accumulated mutations to a cell's DNA (Armitage and Doll (1954), (1957), Fisher (1958), Nordling (1953), and Knudson (1971)). The two-hit hypothesis led to the discovery of tumor suppressor genes. These genes allow a cell to function normally after one mutation. Only after a second mutation does a cell become malignant. It is now accepted that for some cancers to develop a number of mutations must occur. Mathematical models of this mechanism are given in Moolgavkar and Knudson (1981), Michor *et al.* (2006), Michor and Iwasa (2006), Komarova (2007), Haeno *et al.* (2007), and Durrett *et al.* (2009), amongst others.

Oncogene Model

This model comes from the theory that some cancers develop as a result of mutation that gives rise to a gene responsible for malignancy, an oncogene. The population of cells is taken to be of fixed size N , initially all healthy. The evolution is described by the Moran model, where the two types of individuals are healthy and mutated cells. Since it is a birth–death model, approximations and time taken for the population to become malignant can be calculated.

Two-Hit Model

In this model (we follow Haeno *et al.* (2007)) there are three types of cells: type 0 (healthy), type 1 (one mutation), and type 2 (two mutations). The

growth rates of these cells are r , r_1 , and r_2 , and their death rates are δ , δ_1 , and δ_2 . Type 0 mutates into type 1 at the rate u_1 , and type 1 mutates into type 2 at the rate u_2 . Hence at state $\mathbf{x} = (x, y, z)^T$ the birth rates (the rates at which coordinates increase by 1) are $b_1(x, y, z) = rx(1 - u_1)$, $b_2(x, y, z) = rxu_1 + r_1y(1 - u_2)$, and $b_3(x, y, z) = r_1yu_2 + r_2z$, and the death rates (the rates at which coordinates decrease by 1) are $d_1(x, y, z) = \delta x$, $d_2(x, y, z) = \delta_1y$, and $d_3(x, y, z) = \delta_2z$. The sum of these rates gives the holding time parameter $\lambda(x, y, z) = (r + \delta)x + (r_1 + \delta_1)y + (r_2 + \delta_2)z$. Stochastic equations for the process are

$$\begin{aligned} X(t) &= X(0) + \int_0^t (r(1 - u_1) - \delta_1)X(s)ds + M_1(t), \\ Y(t) &= Y(0) + \int_0^t (ru_1X(s) + r_1(1 - u_2)Y(s) - \delta_1Y(s))ds + M_2(t), \\ Z(t) &= Z(0) + \int_0^t (r_1u_2Y(s) + r_2Z(s) - \delta_2Z(s))ds + M_3(t), \end{aligned}$$

with initial conditions $X(0) = 1$, $Y(0) = Z(0) = 0$.

Expectations $EX(t)$, $EY(t)$, $EZ(t)$ are easily found by taking expectations in the above equations and solving the resulting differential equations recursively. Using Dynkin's formula for the function $f(x, y, z) = e^{u_1x+u_2y+u_3z}$, the joint moment generating function of the coordinates can be found by solving the resulting partial differential equations. More complicated quantities, such as the probability that there is a type 2 cell by the time the total population reaches a given size, is found by simulations.

Tumor–Host Interaction

This class of models is based on the kinetic interaction between cells. The kinetic model of Garay and Lefever (1978) assumes that normal cells transform into malignant ones at some constant rate, the transformed cells replicate with some rate, and there is the immunological interaction with the host, which follows the Michaelis–Menten kinetics. After some simplification the following differential equation for the *concentration* $x(t)$ of malignant cells is obtained

$$\frac{dx(t)}{dt} = (1 - \theta x(t))x(t) - \beta \frac{x(t)}{1 + x(t)}, \quad (13.39)$$

$0 \leq \theta \leq 1$, $\beta > 0$. The parameters of this equation are affected by random fluctuations, giving rise to stochastic models. There are different ways of putting noise into the equation. Ochab-Marcinek and Gudowska-Nowak

(2006) add white noise, leading to the stochastic equation

$$dX(t) = \left((1 - \theta X(t))X(t) - \beta \frac{X(t)}{1 + X(t)} \right) dt + \sigma dB(t). \quad (13.40)$$

Rosenkranz (1985) considered σ to be state-dependent, and arrived at its form by using a limit of branching processes. By adding white noise the coefficient β transforms an ODE into an SDE, with $\sigma(x) = \sigma \frac{x}{1+x}$. Once the equation is written down its source becomes irrelevant, and ideas for solutions may come from a different field, physics in this case. The model without noise is seen as

$$\frac{dx(t)}{dt} = -U'(x(t)),$$

where the function U is called the potential. The noisy model is given by the Langevin equation

$$\frac{dX(t)}{dt} = -U'(X(t)) + \xi(t),$$

where $\xi(t)$ is white noise. Of course, we write this equation as an Itô SDE. The process $X(t)$ can be viewed as a motion of a particle in a potential $U(x)$, whose shape varies in time due to the noisy parameter. The potential in the Garay–Lefever model (13.39) is

$$U(x) = -\frac{x^2}{2} + \frac{\theta x^3}{3} + \beta x - \beta \ln(x+1).$$

A minima of the potential corresponds to stable states of the population. In a deterministic system these are stationary (fixed) points where the derivative is zero and the cell concentration does not grow or decay. One of the minima is always at $x = 0$, i.e. the state with no cancer cells is a stable one. The other minimum corresponds to a fixed-size cell population. Transition between those stable states may be interpreted as a spontaneous extinction of cancer due to environmental fluctuations, as well as due to random variations in the immune response efficiency. Time taken to escape from the equilibrium is often studied by simulations, however, in one-dimensional diffusions and birth–death processes it is possible to give analytical approximations. One way to study such stochastic systems is by looking at their stationary distributions; this corresponds to the fixed point analysis in deterministic systems, compare with Bezen and Klebaner (1996). As most studies are done by simulations of such systems we do not pursue this further here.

13.9 Branching Processes

In this section we look at a population model known as the age-dependent Bellman–Harris branching process, see Harris (1963), and Athreya and

Ney (1972). This model generalizes the birth–death process model in two respects: firstly, the lifespan of individuals need not have the exponential distribution, and secondly, more than one particle can be born. In the age-dependent model a particle lives for a random length of time with distribution G , and at the end of its life leaves a random number of children Y , with mean $m = EY$. All particles reproduce and die independently of each other. Denote by $Z(t)$ the number of particles alive at time t . It can be seen that unless the lifetime distribution is exponential, the process $Z(t)$ is not Markovian, and its analysis is usually done by using renewal theory (Harris (1963)). Here we apply stochastic calculus to obtain a limiting result for the population size counted in a special way (by reproductive value).

Consider a collection of individuals with ages (a^1, \dots, a^z) . It is convenient to look at the vector of ages (a^1, \dots, a^z) as a counting measure A , defined by $A(B) = \sum_{i=1}^z 1_B(a^i)$, for any Borel set B in \mathbb{R}^+ . For a function f on \mathbb{R} the following notations are used:

$$(f, A) = \int f(x)A(dx) = \sum_{i=1}^z f(a^i).$$

Denote by A_t the measure corresponding to the ages of individuals alive at time t . Then the population size process in this notation is $Z(t) = (1, A_t)$. In this approach the process of ages A_t is a measure-valued process, although simpler than studied, for example, in Dawson (1993), because the diffusion part corresponding to Laplacian is absent. In order to convert a measure-valued process into a scalar-valued one, test functions are used. Test functions used on the space of measures are of the form $F((f, \mu))$, where F and f are functions on \mathbb{R} . Let $h(a) = \frac{G'(a)}{1-G(a)}$ be the rate of dying at age a . E with and without the subscript A denotes the expectation when the processes start with individuals aged $(a^1, \dots, a^z) = A$. The following result is obtained by direct calculations.

Theorem 13.11: *For a bounded differentiable function F on \mathbb{R} and a continuously differentiable function f on \mathbb{R}^+ , the following limit exists:*

$$\lim_{t \rightarrow 0} \frac{1}{t} E_A \left\{ F((f, A_t)) - F((f, A)) \right\} = \mathcal{G}F((f, A)), \quad (13.41)$$

$$\begin{aligned} \mathcal{G}F((f, A)) &= F'((f, A))(f', A) \\ &\quad + \sum_{j=1}^z h(a^j) \{ E_A (F(Y_j f(0) + (f, A) - f(a^j)) - F((f, A))) \}. \end{aligned}$$

The operator \mathcal{G} defines a generator of a measure-valued branching process in which the movement of the particles is deterministic, namely shift. The following result gives the integral equation (SDE) for the age process under the test functions.

Theorem 13.12: For a bounded C^1 function F on \mathbb{R} and a C^1 function f on \mathbb{R}^+

$$F((f, A_t)) = F((f, A_0)) + \int_0^t \mathcal{G}F((f, A_s))ds + M_t^{F,f}, \quad (13.42)$$

where $M_t^{F,f}$ is a local martingale with the sharp bracket given by

$$\langle M^{F,f}, M^{F,f} \rangle_t = \int_0^t \mathcal{G}F^2((f, A_s))ds - 2 \int_0^t F((f, A_s))\mathcal{G}F((f, A_s))ds.$$

Consequently,

$$\mathbb{E}_A(M_t^{F,f})^2 = \mathbb{E}_A \left(\int_0^t \mathcal{G}F^2((f, A_s))ds - 2 \int_0^t F((f, A_s))\mathcal{G}F((f, A_s))ds \right),$$

provided $\mathbb{E}_A(M_t^{F,f})^2$ exists.

PROOF. The first statement is obtained by Dynkin's formula. The expression for quadratic variation is obtained by letting $U_t = F((f, A_t))$ and an application of the following result. \square

Theorem 13.13: Let U_t be a locally square integrable semimartingale such that $U_t = U_0 + A_t + M_t$, where A_t is a predictable process of locally finite variation and M_t is a locally square integrable local martingale, $A_0 = M_0 = 0$. Let $U_t^2 = U_0^2 + B_t + N_t$, where B_t is a predictable process and N_t is a local martingale. Then

$$\langle M, M \rangle_t = B_t - 2 \int_0^t U_{s-} dA_s - \sum_{s \leq t} (A_s - A_{s-})^2.$$

Of course, if A is continuous (as in our applications) the last term in the above formula vanishes.

PROOF. According to the definition of the quadratic variation process (or integration by parts)

$$\begin{aligned} U_t^2 &= U_0^2 + 2 \int_0^t U_{s-} dU_s + [U, U]_t \\ &= U_0^2 + 2 \int_0^t U_{s-} dA_s + 2 \int_0^t U_{s-} dM_s + [U, U]_t. \end{aligned}$$

Using the representation for U_t^2 given in the conditions of Theorem 13.13, we obtain that $[U, U]_t - B_t + 2 \int_0^t U_{s-} dA_s$ is a local martingale. Since $[A, A]_t = \sum_{s \leq t} (A_s - A_{s-})^2$ the result follows. \square

Let $v^2 = \mathbb{E}(Y^2)$ be the second moment of the offspring distribution. Applying Dynkin's formula to the function $F(u) = u$ (and writing M^f for $M^{1,f}$), we obtain the following result. Recall that h is the hazard function of G , $h = G'/(1 - G)$.

Theorem 13.14: *For a C^1 function f on \mathbb{R}^+*

$$(f, A_t) = (f, A_0) + \int_0^t (Lf, A_s) ds + M_t^f, \quad (13.43)$$

where the linear operator L is defined by

$$Lf = f' - hf + mh f(0), \quad (13.44)$$

and M_t^f is a square integrable local martingale with the sharp bracket given by

$$\langle M^f, M^f \rangle_t = \int_0^t (f^2(0)v^2h + hf^2 - 2f(0)mhf, A_s) ds. \quad (13.45)$$

PROOF. The first statement is Dynkin's formula for $F(u) = u$. This function is unbounded and the standard formula cannot be applied directly. However, it can be applied by taking smooth bounded functions that agree with u on bounded intervals, $F_n(u) = u$ for $u \leq n$, and the sequence of stopping times $T_n = \inf\{(f, A_t) > n\}$ as a localizing sequence. The form of the operator L follows from (13.42). Similarly, (13.45) follows by taking $F(u) = u^2$. \square

By taking f to be a constant, $f(u) = 1$, Theorem 13.14 yields the following corollary for the population size at time t , $Z(t) = (1, A_t)$.

Corollary 13.15: *The compensator of $Z(t)$ is given by $(m-1) \int_0^t (h, A_s) ds$.*

It is useful to be able to take expectations in Itô's formula. The next result gives a sufficient condition, i.e. conditions for Dynkin's formula to hold.

Theorem 13.16: *Let $f \geq 0$ be a C^1 function on \mathbb{R}^+ that satisfies*

$$|(Lf, A)| \leq C(1 + (f, A)), \quad (13.46)$$

for some constant C and any A , and assume that (f, A_0) is integrable. Then (f, A_t) and M_t^f in (13.43) are also integrable with $\mathbb{E}M_t^f = 0$.

PROOF. Let T_n be a localizing sequence for M^f , then from (13.43)

$$(f, A_{t \wedge T_n}) = (f, A_0) + \int_0^{t \wedge T_n} (Lf, A_s) ds + M_{t \wedge T_n}^f, \quad (13.47)$$

where $M_{t \wedge T_n}^f$ is a martingale. Taking expectations we have

$$\mathbb{E}(f, A_{t \wedge T_n}) = \mathbb{E}(f, A_0) + \mathbb{E} \int_0^{t \wedge T_n} (Lf, A_s) ds. \quad (13.48)$$

According to Condition (13.46)

$$\begin{aligned} \left| \mathbb{E} \int_0^{t \wedge T_n} (Lf, A_s) ds \right| &\leq \mathbb{E} \int_0^{t \wedge T_n} |(Lf, A_s)| ds \\ &\leq Ct + C\mathbb{E} \int_0^{t \wedge T_n} (f, A_s) ds. \end{aligned}$$

Thus we have from (13.48) by using (13.2)

$$\begin{aligned} \mathbb{E}(f, A_{t \wedge T_n}) &\leq \mathbb{E}(f, A_0) + Ct + C\mathbb{E} \int_0^{t \wedge T_n} (f, A_s) ds \\ &\leq \mathbb{E}(f, A_0) + Ct + C\mathbb{E} \int_0^t I(s \leq T_n)(f, A_s) ds \\ &\leq \mathbb{E}(f, A_0) + Ct + C \int_0^t \mathbb{E}(f, A_{s \wedge T_n}) ds. \end{aligned}$$

It now follows by Gronwall's inequality (Theorem 1.20) that

$$\mathbb{E}(f, A_{t \wedge T_n}) \leq \mathbb{E}(f, A_0) + Ct + te^{Ct} < \infty. \quad (13.49)$$

Taking $n \rightarrow \infty$ we conclude that $\mathbb{E}(f, A_t) < \infty$ by Fatou's lemma. Thus (f, A_t) is integrable as it is non-negative. Now by Condition (13.46)

$$\mathbb{E} \left| \int_0^t (Lf, A_s) ds \right| \leq \int_0^t \mathbb{E} |(Lf, A_s)| ds \leq \int_0^t C(1 + \mathbb{E}(f, A_s)) ds < \infty. \quad (13.50)$$

It follows from (13.50) that $\int_0^t (Lf, A_s) ds$ and its variation process $\int_0^t |(Lf, A_s)| ds$ are both integrable, and from (13.43) that

$$M_t^f = (f, A_t) - (f, A_0) - \int_0^t (Lf, A_s) ds \quad (13.51)$$

is integrable with zero mean. \square

For simplicity we assume that $G(u) < 1$ for all $u \in \mathbb{R}^+$. Equation (13.43) can be analyzed through the eigenvalue problem for the operator L .

Theorem 13.17: Let L be the operator in (13.44). Then the equation

$$Lq = rq \quad (13.52)$$

has a solution q_r for any r . The corresponding eigenfunction (normed so that $q(0) = 1$) is given by

$$q_r(u) = \frac{me^{ru}}{1 - G(u)} \left(1 - \int_0^u e^{-rs} dG(s) \right). \quad (13.53)$$

PROOF. Since eigenfunctions are determined up to a multiplicative constant we can take $q(0) = 1$. Equation (13.52) is a first order linear differential equation, and solving it we obtain the solution (13.53). \square

Theorem 13.18: Let q_r be a positive eigenfunction of L corresponding to the eigenvalue r . Then $Q_r(t) = e^{-rt}(q_r, A_t)$ is a positive martingale.

PROOF. Using (13.43) and the fact that q_r is an eigenfunction for L we have

$$(q_r, A_t) = (q_r, A_0) + r \int_0^t (q_r, A_s) ds + M_t^{q_r}, \quad (13.54)$$

where $M_t^{q_r}$ is a local martingale. The functions q_r clearly satisfy condition (13.46). Therefore (q_r, A_t) is integrable, and it follows from (13.54) by taking expectations that

$$\mathbb{E}(q_r, A_t) = e^{rt} \mathbb{E}(q_r, A_0). \quad (13.55)$$

Using integration by parts for $e^{-rt}(q_r, A_t)$ we obtain from (13.54) that

$$dQ_r(t) = d(e^{-rt}(q_r, A_t)) = e^{-rt} dM_t^{q_r},$$

and

$$Q_r(t) = (q_r, A_0) + \int_0^t e^{-rs} dM_s^{q_r} \quad (13.56)$$

is a local martingale as an integral with respect to the local martingale M^{q_r} . Since a positive local martingale is a supermartingale, and $Q_r(t) \geq 0$, $Q_r(t)$ is a supermartingale. However, from (13.55) it follows that $Q_r(t)$ has a constant mean. Thus the supermartingale $Q_r(t)$ is a martingale. \square

The Malthusian parameter α is defined as the value of r which satisfies

$$m \int_0^\infty e^{-ru} dG(u) = 1. \quad (13.57)$$

We assume that the Malthusian parameter α exists and is *positive*, in this case the process is called supercritical.

Theorem 13.19: *There is only one bounded positive eigenfunction V , the reproductive value function, corresponding to the eigenvalue α which is the Malthusian parameter,*

$$V(u) = \frac{me^{\alpha u}}{1 - G(u)} \int_u^\infty e^{-\alpha s} dG(s). \quad (13.58)$$

PROOF. It follows that for $r > \alpha$, $m \int_0^\infty e^{-ru} dG(u) < 1$ and the eigenfunction q_r in (13.53) is positive and grows exponentially fast or faster. For $r < \alpha$, the eigenfunction q_r takes negative values. When $r = \alpha$, $q^\alpha = V$ in (13.58). In order to see that it is bounded by m , replace $e^{-\alpha s}$ by its largest value $e^{-\alpha u}$. \square

This is the main result of this section.

Theorem 13.20: *$W_t = e^{-\alpha t}(V, A_t)$ is a positive square integrable martingale, and therefore converges almost surely and in L^2 to a non-degenerate limit $W \geq 0$, $\text{EW} = (V, A_0) > 0$ and $\text{P}(W > 0) > 0$.*

PROOF. That W_t is a martingale follows by Theorem 13.18. It is positive, therefore it converges almost surely to a limit $W \geq 0$ by the martingale convergence Theorem 7.11. In order to show that the limit is not identically zero we show that convergence is also in L^2 , i.e. the second moments (and the first moments) converge, implying that $\text{EW} = \lim_{t \rightarrow \infty} \text{EW}_t = \text{EW}_0 = (V, A_0)$.

It follows from (13.45) that

$$\langle M^V, M^V \rangle_t = \int_0^t \left((\sigma^2 + (m - V)^2)h, A_s \right) ds.$$

According to (13.56) we obtain

$$W_t = (V, A_0) + \int_0^t e^{-\alpha s} dM_s^V,$$

and that

$$\begin{aligned} \langle W, W \rangle_t &= \int_0^t e^{-2\alpha s} d\langle M^V, M^V \rangle_s \\ &= \int_0^t e^{-2\alpha s} \left((\sigma^2 + (m - V)^2)h, A_s \right) ds. \end{aligned} \quad (13.59)$$

Now it is easy to check that there is a constant C and r , $\alpha \leq r < 2\alpha$ such that

$$\left((\sigma^2 + (m - V)^2)h, A_s \right) \leq C(q_r, A_s),$$

and using Theorem 13.18 we get

$$\mathbb{E} \int_0^\infty e^{-2\alpha s} \left((\sigma^2 + (m - V)^2) h, A_s \right) ds < C \int_0^\infty e^{(r-2\alpha)s} ds < \infty.$$

This implies from (13.59) that $\mathbb{E}\langle W, W \rangle_\infty < \infty$. Therefore W_t is a square integrable martingale (see Theorem 8.27) and the result follows. \square

The martingale $\{W_t\}$ is given in Harris (1963). The convergence of the population numbers $Z(t)e^{-\alpha t}$ is obtained from that of W_t by using a Tauberian theorem. For details and extensions of the model to the population-dependent case see Jagers and Klebaner (2000).

13.10 Stochastic Lotka–Volterra Model

Deterministic Lotka–Volterra System

The Lotka–Volterra system of ordinary differential equations (Lotka (1925), Volterra (1926))

$$\begin{aligned} \dot{x}_t &= \mathbf{a}x_t - \mathbf{b}x_ty_t \\ \dot{y}_t &= \mathbf{c}x_ty_t - \mathbf{d}y_t, \end{aligned} \tag{13.60}$$

with positive x_0, y_0 , and positive parameters $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ describes a behaviour of a prey–predator system in terms of the prey and predator “intensities” x_t and y_t . Here, \mathbf{a} is the rate of increase of prey in the absence of predators, \mathbf{d} is a rate of decrease of predators in the absence of prey, while the rate of decrease in prey is proportional to the number of predators $\mathbf{b}y_t$, and similarly the rate of increase in predators is proportional to the number of prey $\mathbf{c}x_t$. Unlike in previous sections, here we denote scalar coefficients in bold font. The system (13.60) is one of the simplest non-linear systems.

Since the population numbers are discrete a description of the predator–prey model in terms of continuous intensities x_t, y_t is based implicitly on a natural assumption that the numbers of both populations are large, and the intensities are obtained by a normalization of population numbers by a large parameter K . Thus (13.60) is an approximation, an asymptotic description of the interaction between the predator and the prey. Although this model may capture some essential elements in that interaction, it is not suitable for answering questions of extinction of populations, as the extinction never occurs in the deterministic model, see Figure 13.1 for the pair x_t, y_t in the phase plane.

We introduce here a probabilistic model which has as its limit the deterministic Lotka–Volterra model, evolves in continuous time according to

the same local interactions, and allows us to evaluate asymptotically the time for extinction of prey species.

There is a vast amount of literature on the Lotka–Volterra model and a history of research on stochastic perturbations of this system exact, approximate, and numerical.

The system (13.60) possesses the first integral which is a closed orbit in the first quadrant of phase plane x, y . It is given by

$$r(x, y) = \mathbf{c}x - \mathbf{d} \log x + \mathbf{b}y - \mathbf{a} \log y + r_0, \quad (13.61)$$

where r_0 is an arbitrary constant. It depends only on the initial point (x_0, y_0) (see Figure 13.1).




Fig. 13.1: First integral $r(x, y)$, $x_0 = 0.3$, $y_0 = 3$, $\mathbf{a} = 5$, $\mathbf{b} = 1$, $\mathbf{c} = 5$, $\mathbf{d} = 1$.

Stochastic Lotka–Volterra System

Let X_t and Y_t be numbers of prey and predators at time t . We start with simple equations for prey–predator populations

$$\begin{aligned} X_t &= X_0 + \pi'_t - \pi''_t \\ Y_t &= Y_0 + \widehat{\pi}'_t - \widehat{\pi}''_t \end{aligned} \quad (13.62)$$

where π'_t is the number of prey born up to time t , π''_t is the number of prey killed up to time t , $\widehat{\pi}'_t$ is the number of predators born up to time t , and $\widehat{\pi}''_t$ is the number of predators that have died up to time t .

We assume that $\pi'_t, \pi''_t, \widehat{\pi}'_t, \widehat{\pi}''_t$ are Poisson processes with the following state-dependent random rates $\mathbf{a}X_t$, $\frac{\mathbf{b}}{K}X_tY_t$, $\frac{\mathbf{c}}{K}X_tY_t$, $\mathbf{d}Y_t$ respectively and disjoint jumps (the latter assumption reflects the fact that in a short time interval $(t, t + \delta t)$ or only one prey might be born and only one might be

killed, or only one predator might be born and only one might die, with the above mentioned intensities. Moreover, all these events are disjoint in time).

Assume $X_0 = Kx_0$ and $Y_0 = Ky_0$ for some fixed positive x_0, y_0 and a large integer parameter K . Introduce the prey and predator populations normalized by K

$$x_t^K = \frac{X_t}{K}, \quad \text{and} \quad y_t^K = \frac{Y_t}{K}.$$

In terms of x_t^K and y_t^K the introduced intensities for Poisson processes can be written as $\mathbf{a}Kx_t^K, \mathbf{b}Kx_t^Ky_t^K, \mathbf{c}Kx_t^Ky_t^K, \mathbf{d}Ky_t^K$.

Existence

In this section we show that a random process (X_t, Y_t) can be constructed to satisfy Equations (13.62). To this end we introduce four independent sequences of Poisson processes:

$$\begin{aligned}\Pi_t^{\mathbf{a}} &= (\Pi_t^{\mathbf{a}}(1), \Pi_t^{\mathbf{a}}(2), \dots), \\ \Pi_t^{\mathbf{b}/K} &= (\Pi_t^{\mathbf{b}/K}(1), \Pi_t^{\mathbf{b}/K}(2), \dots), \\ \Pi_t^{\mathbf{c}/K} &= (\Pi_t^{\mathbf{c}/K}(1), \Pi_t^{\mathbf{c}/K}(2), \dots), \\ \Pi_t^{\mathbf{d}} &= (\Pi_t^{\mathbf{d}}(1), \Pi_t^{\mathbf{d}}(2), \dots).\end{aligned}$$

Each of them is a sequence of i.i.d. Poisson processes with rates $\mathbf{a}, \frac{\mathbf{b}}{K}, \frac{\mathbf{c}}{K}, \mathbf{d}$ respectively. Define the processes (X_t, Y_t) by the system of Itô equations

$$\begin{aligned}X_t &= X_0 + \int_0^t \sum_{n \geq 1} I(X_{s-} \geq n) d\Pi_s^{\mathbf{a}}(n) - \int_0^t \sum_{n \geq 1} I(X_{s-} - Y_{s-} \geq n) d\Pi_s^{\mathbf{b}/K}(n), \\ Y_t &= Y_0 + \int_0^t \sum_{n \geq 1} I(X_{s-} - Y_{s-} \geq n) d\Pi_s^{\mathbf{c}/K}(n) - \int_0^t \sum_{n \geq 1} I(Y_{s-} \geq n) d\Pi_s^{\mathbf{d}}(n),\end{aligned}\tag{13.63}$$

governed by these Poisson processes, which obviously has a unique solution until the time of explosion, on the time interval $[0, T_\infty]$, where

$$T_\infty = \inf\{t > 0 : X_t \vee Y_t = \infty\}.$$

The Poisson processes with state-dependent rates in (13.62) are obtained as follows:

$$\begin{aligned}\pi'_t &= \int_0^t \sum_{n \geq 1} I(X_{s-} \geq n) d\Pi_s^{\mathbf{a}}(n), \quad \pi''_t = \int_0^t \sum_{n \geq 1} I(X_{s-} - Y_{s-} \geq n) d\Pi_s^{\mathbf{b}/K}(n), \\ \widehat{\pi}'_t &= \int_0^t \sum_{n \geq 1} I(X_{s-} - Y_{s-} \geq n) d\Pi_s^{\mathbf{c}/K}(n), \quad \widehat{\pi}''_t = \int_0^t \sum_{n \geq 1} I(Y_{s-} \geq n) d\Pi_s^{\mathbf{d}}(n).\end{aligned}$$

It is easy to see that $\pi'_t, \pi''_t, \widehat{\pi}'_t, \widehat{\pi}''_t$ have the required properties. Their jumps are of size one. Since all Poisson processes are independent their jumps are disjoint, so that the jumps of $\pi'_t, \pi''_t, \widehat{\pi}'_t, \widehat{\pi}''_t$ are disjoint as well. In order to describe their intensities introduce a stochastic basis $(\Omega, \mathcal{F}, \mathbb{F} = (\mathcal{F}_t)_{t \geq 0}, P)$, the filtration \mathbb{F} is generated by all Poisson processes and satisfies the general conditions. Then, obviously, the random process

$$A'_t = \int_0^t \sum_{n \geq 1} I(X_s \geq n) \mathbf{a} ds$$

is adapted and continuous, hence predictable. Using $\int_0^t f(s-) ds = \int_0^t f(s) ds$,

$$\begin{aligned} \pi'_t - A'_t &= \int_0^t \sum_{n \geq 1} I(X_{s-} \geq n) d\Pi_s^{\mathbf{a}}(n) - \int_0^t \sum_{n \geq 1} I(X_{s-} \geq n) \mathbf{a} ds \\ &= \int_0^t \left(\sum_{n \geq 1} I(X_{s-} \geq n) \right) d(\Pi_s^{\mathbf{a}}(n) - \mathbf{a} s). \end{aligned}$$

Thus $\pi'_t - A'_t$ is an integral with respect to a martingale, hence it is a local martingale. Therefore A'_t is the compensator of π'_t . Since X_s is an integer-valued random variable, $\sum_{n \geq 1} I(X_s \geq n) = X_s$, giving that $A'_t = \int_0^t \mathbf{a} X_s ds$ and the intensity of π'_t is $\mathbf{a} X_t$, as claimed. Analogously, other compensators are

$$A''_t = \int_0^t \frac{\mathbf{b}}{K} X_s Y_s ds, \quad \widehat{A}'_t = \int_0^t \frac{\mathbf{c}}{K} X_s Y_s ds, \quad \widehat{A}''_t = \int_0^t \mathbf{d} Y_s ds,$$

and thus all other intensities have the required form.

We now show that the process $(X_t, Y_t)_{t \geq 0}$ does not explode.

Theorem 13.21:

$$\mathbb{P}(T_\infty = \infty) = 1.$$

PROOF. Set $T_n^X = \inf\{t > 0 : X_t \geq n\}$, $n \geq 1$ and denote by $T_\infty^X = \lim_{n \rightarrow \infty} T_n^X$. Using (13.63) we obtain

$$\mathbb{E} X_{t \wedge T_n^X} \leq X_0 + \int_0^t \mathbf{a} \mathbb{E} X_{s \wedge T_n^X} ds.$$

According to Gronwall's inequality, $\mathbb{E} X_{T_n^X \wedge T} \leq X_0 e^{\mathbf{a} T}$ for every $T > 0$. Hence by Fatou's lemma $\mathbb{E} X_{T_\infty^X \wedge T} \leq X_0 e^{\mathbf{a} T}$. Consequently, for all $T > 0$

$$\mathbb{P}(T_\infty^X \leq T) = 0.$$

Set $T_\ell^Y = \inf\{t : Y_t \geq \ell\}$, $\ell \geq 1$ and denote by $T_\infty^Y = \lim_{\ell \rightarrow \infty} T_\ell^Y$. Using (13.63) we obtain

$$\begin{aligned}\text{E}Y_{t \wedge T_n^X \wedge T_\ell^Y} &\leq Y_0 + \int_0^t \frac{\mathbf{c}}{K} \text{E}(X_{s \wedge T_n^X \wedge T_\ell^Y} Y_{s \wedge T_n^X \wedge T_\ell^Y}) ds \\ &\leq Y_0 + \int_0^t \frac{\mathbf{c}}{K} n \text{E}Y_{s \wedge T_n^X \wedge T_\ell^Y} ds.\end{aligned}$$

Hence, by Gronwall's inequality, for every $T > 0$, $\text{E}Y_{T \wedge T_n^X \wedge T_m^Y} \leq Y_0 e^{\frac{\mathbf{c}}{K} n T}$ and by Fatou's lemma,

$$\text{E}Y_{T \wedge T_n^X \wedge T_\infty^Y} \leq Y_0 e^{\frac{\mathbf{c}}{K} n T}, \quad n \geq 1.$$

Consequently, $\text{P}(T_\infty^Y \leq T_n^X \wedge T) = 0, \forall T > 0, n \geq 1$ and, since $T_n^X \nearrow \infty$, as $n \rightarrow \infty$, we obtain

$$\text{P}(T_\infty^Y \leq T) = 0.$$

Since $T_\infty = T_\infty^X \wedge T_\infty^Y$, $\text{P}(T_\infty \leq T) \leq \text{P}(T_\infty^X \leq T) + \text{P}(T_\infty^Y \leq T)$, and we have $\text{P}(T_\infty \leq T) = 0$ for any $T > 0$. \square

Corollary 13.22: For $T_n = \inf\{t : X_t \vee Y_t \geq n\}$, and for all $T > 0$

$$\lim_{n \rightarrow \infty} \text{P}(T_n \leq T) = 0.$$

The above description of the model allows us to claim that (X_t, Y_t) is a continuous time pure jump Markov process with jumps of two possible sizes in both coordinates: 1 and -1 and infinitesimal transition probabilities (as $\delta t \rightarrow 0$)

$$\begin{aligned}\text{P}\left(X_{t+\delta t} = X_t + 1 \mid X_t, Y_t\right) &= \mathbf{a}X_t \delta t + o(\delta t) \\ \text{P}\left(X_{t+\delta t} = X_t - 1 \mid X_t, Y_t\right) &= \frac{\mathbf{b}}{K} X_t Y_t \delta t + o(\delta t) \\ \text{P}\left(Y_{t+\delta t} = Y_t + 1 \mid X_t, Y_t\right) &= \frac{\mathbf{c}}{K} X_t Y_t \delta t + o(\delta t) \\ \text{P}\left(Y_{t+\delta t} = Y_t - 1 \mid X_t, Y_t\right) &= \mathbf{d}Y_t \delta t + o(\delta t).\end{aligned}$$

Semimartingale Decomposition for (x_t^K, y_t^K)

Let $A'_t, A''_t, \widehat{A}'_t, \widehat{A}''_t$ be the compensators of $\pi'_t, \pi''_t, \widehat{\pi}'_t, \widehat{\pi}''_t$ defined above. Introduce martingales

$$M'_t = \pi'_t - A'_t, \quad M''_t = \pi''_t - A''_t, \quad \widehat{M}'_t = \widehat{\pi}'_t - \widehat{A}'_t, \quad \widehat{M}''_t = \widehat{\pi}''_t - \widehat{A}''_t,$$

and also normalized martingales

$$m_t^K = \frac{M'_t - M''_t}{K} \quad \text{and} \quad \hat{m}_t^K = \frac{\widehat{M}'_t - \widehat{M}''_t}{K}. \quad (13.64)$$

Then from (13.63) it follows that the process (x_t^K, y_t^K) admits the semi-martingale decomposition

$$\begin{aligned} x_t^K &= x_0 + \int_0^t [\mathbf{a}x_s^K - \mathbf{b}x_s^K y_s^K] ds + m_t^K, \\ y_t^K &= y_0 + \int_0^t [\mathbf{c}x_s^K y_s^K - \mathbf{d}y_s^K] ds + \hat{m}_t^K, \end{aligned} \quad (13.65)$$

which is a stochastic analogue (in integral form) of the Equations (13.60).

In the sequel we need quadratic variations of the martingales in (13.65). According to Theorem 9.3 all martingales are locally square integrable and possess the predictable quadratic variations

$$\langle M', M' \rangle_t = A'_t, \quad \langle M'', M'' \rangle_t = A''_t \quad \text{and} \quad \langle \widehat{M}', \widehat{M}' \rangle_t = \widehat{A}'_t, \quad \langle \widehat{M}'', \widehat{M}'' \rangle_t = \widehat{A}''_t, \quad (13.66)$$

and zero covariations $\langle M', M'' \rangle_t \equiv 0, \dots, \langle \widehat{M}', \widehat{M}'' \rangle_t \equiv 0$, since the jumps of $\pi'_t, \pi''_t, \widehat{\pi}'_t, \widehat{\pi}''_t$ are disjoint. Hence we obtain the sharp brackets of the martingales in Equations (13.65),

$$\begin{aligned} \langle m^K, \hat{m}^K \rangle_t &\equiv 0, \\ \langle m^K, m^K \rangle_t &= \frac{1}{K} \int_0^t (\mathbf{a}x_s^K + \mathbf{b}x_s^K y_s^K) ds, \\ \langle \hat{m}^K, \hat{m}^K \rangle_t &= \frac{1}{K} \int_0^t (\mathbf{c}x_s^K y_s^K + \mathbf{d}y_s^K) ds. \end{aligned} \quad (13.67)$$

Note that the stochastic equations above do not satisfy the linear growth condition in x_t^K, y_t^K . Nevertheless, the solution exists for all t .

Deterministic (Fluid) Approximation

We now show that the Lotka–Volterra Equations (13.60) describe a limit (also known as fluid approximation) for the family (x_t^K, y_t^K) as parameter $K \rightarrow \infty$. Results on the fluid approximation for discontinuous Markov processes can be found in Kurtz (1981).

Theorem 13.23: *For any $T > 0$ and $\eta > 0$*

$$\lim_{K \rightarrow \infty} \mathbb{P} \left(\sup_{t \leq T} (|x_t^K - x_t| + |y_t^K - y_t|) > \eta \right) = 0.$$

PROOF. Set

$$T_n^K = \inf\{t : x_t^K \vee y_t^K \geq n\}. \quad (13.68)$$

According to Corollary 13.22, since $T_n^K = T_{nK}$,

$$\lim_{n \rightarrow \infty} \limsup_{K \rightarrow \infty} P(T_n^K \leq T) = 0. \quad (13.69)$$

Hence it suffices to show that for every $n \geq 1$,

$$\lim_{K \rightarrow \infty} P \left(\sup_{t \leq T_n^K \wedge T} (|x_t^K - x_t| + |y_t^K - y_t|) > \eta \right) = 0. \quad (13.70)$$

Since $\sup_{t \leq T_n^K \wedge T} (x_t^K \vee y_t^K) \leq n+1$, there is a constant L_n , depending on n and T , such that for $t \leq T_n^K \wedge T$

$$\begin{aligned} |(\mathbf{a}x_t^K - \mathbf{b}x_t^K y_t^K) - (\mathbf{a}x_t - \mathbf{b}x_t y_t)| &\leq L_n (|x_t^K - x_t| + |y_t^K - y_t|) \\ |(\mathbf{c}x_t^K y_t^K - \mathbf{d}y_t^K) - (\mathbf{c}x_t y_t - \mathbf{d}y_t)| &\leq L_n (|x_t^K - x_t| + |y_t^K - y_t|). \end{aligned}$$

These inequalities, (13.60), and (13.65) imply

$$\begin{aligned} &|x_{T_n^K \wedge T}^K - x_{T_n^K \wedge T}| + |y_{T_n^K \wedge T}^K - y_{T_n^K \wedge T}| \\ &\leq 2L_n \int_0^t (|x_{s \wedge T_n^K}^K - x_{s \wedge T_n^K}| + |y_{s \wedge T_n^K}^K - y_{s \wedge T_n^K}|) ds \\ &+ \sup_{t \leq T_n^K \wedge T} |m_t^K| + \sup_{t \leq T_n^K \wedge T} |\hat{m}_t^K|. \end{aligned}$$

Now by Gronwall's inequality we find

$$\sup_{t \leq T_n^K \wedge T} (|x_t^K - x_t| + |y_t^K - y_t|) \leq e^{2L_n T} \left(\sup_{t \leq T_n^K \wedge T} |m_t^K| + \sup_{t \leq T_n^K \wedge T} |\hat{m}_t^K| \right).$$

Therefore (13.70) holds if both $\sup_{t \leq T_n^K \wedge T} |m_t^K|$ and $\sup_{t \leq T_n^K \wedge T} |\hat{m}_t^K|$ converge in probability to zero as $K \rightarrow \infty$. According to (13.67) and definition (13.68) of T_n^K ,

$$E\langle m^K \rangle_{T_n^K \wedge T} \leq \frac{1}{K} (\mathbf{a}(n+1) + \mathbf{b}(n+1)^2) T.$$

Thus by Doob's inequality for martingales (8.47)

$$E \left(\left(\sup_{t \leq T_n^K \wedge T} |m_t^K| \right)^2 \right) \leq 4E(\langle m^K, m^K \rangle_{T_n^K \wedge T}) \rightarrow 0,$$

as $K \rightarrow \infty$. This implies $\sup_{t \leq T_n^K \wedge T} |m_t^K| \rightarrow 0$ as $K \rightarrow \infty$. The second term $\sup_{t \leq T_n^K \wedge T} |\hat{m}_t^K|$ is treated similarly, and the proof is complete. \square

Using the stochastic Lotka–Volterra model one can evaluate asymptotically, when K is large, the time to extinction of prey species, as well as the likely trajectory to extinction. One such trajectory is given in Figure 13.2. This analysis is done by using the large deviations theory (see Klebaner and Liptser (2001)).




Fig. 13.2: A likely path to extinction.

13.11 Exercises

Exercise 13.1: Find the expected time to extinction $E T_0$ in the branching diffusion with $\alpha < 0$. Hint: use Theorem 6.16 and formula (6.98) to find $E(T_0 \wedge T_b)$. $E T_0 = \lim_{b \rightarrow \infty} E(T_0 \wedge T_b)$ by monotone convergence.

Exercise 13.2: Let $X(t)$ be a branching diffusion satisfying SDE (13.1).

1. Let $c = 2\alpha/\sigma^2$. Show that e^{-cX_t} is a martingale.
2. Let T be the time to extinction, $T = \inf\{t : X_t = 0\}$, where $T = \infty$ if $X_t > 0$ for all $t \geq 0$, and let $q_t(x) = P_x(T \leq t)$ be the probability that extinction occurs by time t when the initial population size is x . Prove that $q_t(x) \leq e^{-cx}$.

Exercise 13.3: A model for population growth is given by the following SDE $dX(t) = 2X(t)dt + \sqrt{X(t)}dB(t)$, and $X(0) = x > 0$. Find the probability that the population doubles its initial size x before it becomes extinct. Show that when the initial population size x is small, then this probability is approximately $1/2$, but when x is large this probability is nearly one.

Exercise 13.4: Derive the stationary distribution for the Wright–Fisher diffusion.

Exercise 13.5: Let a deterministic growth model be given by the differential equation

$$dx(t) = g(x(t))dt, \quad x_0 > 0, \quad t \geq 0,$$

for a positive function $g(x)$ and consider its stochastic analogue

$$dX(t) = g(X(t))dt + \sigma(X(t))dB(t), \quad X(0) > 0.$$

One way to analyze the stochastic equation is by comparison with the deterministic solution.

1. Find $G(x)$ such that $G(x(t)) = G(x(0)) + t$ and consider $Y(t) = G(X(t))$.
Find $dY(t)$.
2. Let $g(x) = x^r$, $0 \leq r < 1$, and $\sigma(x)/x^r \rightarrow 0$ as $x \rightarrow \infty$. Give conditions on $g(x)$ and $\sigma^2(x)$ so that the law of large numbers holds for $Y(t)$, that is, $Y(t)/t \rightarrow 1$, as $t \rightarrow \infty$ on the set $\{Y(t) \rightarrow \infty\}$.

A systematic analysis of this model is given in Keller *et al.* (1984).

Exercise 13.6: Let L_1, L_2, \dots, L_x be independent exponentially distributed random variables with parameter $a(x)$ (they represent the lifespan of particles in the population of size x). Show that $\min(L_1, L_2, \dots, L_x)$ has an exponential distribution with parameter $xa(x)$.

Exercise 13.7: (Birth–death processes mean exit time)

1. Find the expected time to exit from (a, b) by using Theorem 13.8.
2. Find the expected time to extinction by taking $a = 0$ and $b \rightarrow \infty$ in the result above.

Exercise 13.8: Find the variance of the proportion of type 1 particles $X(t)/N$ in the Moran model.

Exercise 13.9: (Birth–death processes: stochastic representation)

Let $N_k^\lambda(t)$ and $N_k^\mu(t)$, $k \geq 1$ be two independent sequences of independent Poisson processes with rates λ and μ . Let $X(0) > 0$ and for $t > 0$ $X(t)$ satisfies

$$X(t) = X(0) + \int_0^t \sum_{k \geq 1} I(X(s-) \geq k) dN_k^\lambda(s) - \int_0^t \sum_{k \geq 1} I(X(s-) \geq k) dN_k^\mu(s).$$

1. Show that $X(t)$ is a birth–death process and identify the rates.
2. Give the semimartingale decomposition for $X(t)$.

Chapter 14

Applications in Engineering and Physics

In this chapter methods of stochastic calculus are applied to the filtering problem in engineering and random oscillators in physics. The filtering problem consists of finding the best estimator of a signal when observations are contaminated by noise. For a number of classical equations of motions in physics we find stationary densities when the motion is subjected to random excitations.

14.1 Filtering

The filtering problem is the problem of estimation of a signal contaminated by noise. Let $Y(t)$ be the observation process, and \mathcal{F}_t^Y denote the information available by observing the process up to time t , that is, $\mathcal{F}_t^Y = \sigma(Y(s), s \leq t)$. The observation $Y(t)$ at time t is the result of a deterministic transformation of the signal process $X(s)$, $s \leq t$, typically a linear transformation, to which a random noise is added. The filtering problem is to find the “best” estimate $\pi_t(X)$ of the signal $X(t)$ on the basis of all the observations $Y(s)$, $s \leq t$, or \mathcal{F}_t^Y . The “best” is understood in the sense of the smallest estimation error $E((X(t) - Z(t))^2 | \mathcal{F}_t^Y)$, when $Z(t)$ varies over all \mathcal{F}_t^Y -measurable processes. Denote for an adapted process $h(t)$

$$\pi_t(h) = E(h(t) | \mathcal{F}_t^Y). \quad (14.1)$$

Then by Theorem 2.26 (see also Exercise 14.2) the filtering problem is in finding $\pi_t(X)$.

Two main results of stochastic calculus are used to solve the filtering problem: Levy's characterization of Brownian motion, and the predictable representation property of Brownian filtration.

General Non-linear Filtering Model

Let (Ω, \mathcal{F}, P) be a complete probability space, and let (\mathcal{F}_t) , $0 \leq t \leq T$, be a non-decreasing family of right-continuous σ -algebras of \mathcal{F} , satisfying the usual conditions, and supporting adapted processes $X(t)$ and $Y(t)$.

We aim at finding $\pi_t(h)$ for an adapted process $h(t)$. In particular, we can find $E(g(X(t))|\mathcal{F}_t^Y)$, for a function of a real variable g . When g ranges over a set of test functions the conditional distribution of $X(t)$ given \mathcal{F}_t^Y is obtained. Moreover, by taking $g(x) = x$, we obtain the best estimator $\pi_t(X)$. We assume that the process $h(t)$, which we want to filter, and the observation process $Y(t)$ satisfy the SDEs

$$\begin{aligned} dh(t) &= H(t)dt + dM(t), \\ dY(t) &= A(t)dt + B(Y(t))dW(t), \end{aligned} \tag{14.2}$$

where:

1. The process $M(t)$ is a \mathcal{F}_t -martingale.
2. The process $W(t)$ is a \mathcal{F}_t -Brownian motion.
3. The processes $H(t)$ and $A(t)$ are random, satisfying with probability one $\int_0^T |H(t)|dt < \infty$, $\int_0^T |A(t)|dt < \infty$.
4. $\sup_{t \leq T} E(h^2(t)) < \infty$, $\int_0^T EH^2(t)dt < \infty$, $\int_0^T EA^2(t)dt < \infty$.
5. The diffusion coefficient of the observation process $Y(t)$ is a function $B(y)$ of $Y(t)$ only, and not X . $B^2(y) \geq C > 0$, and B^2 satisfies Lipschitz and the linear growth conditions.

Theorem 14.1: *For each t , $0 \leq t \leq T$,*

$$d\pi_t(h) = \pi_t(H)dt + (\pi_t(D) + \frac{\pi_t(hA) - \pi_t(h)\pi_t(A)}{B(Y(t))})d\bar{W}(t), \tag{14.3}$$

where $d\bar{W}(t) = \frac{dY(t) - \pi_t(Adt)}{B(Y(t))}$ is an \mathcal{F}_t^Y -Brownian motion and $D(t) = \frac{d\langle M, W \rangle(t)}{dt}$.

$\bar{W}(t)$ is called the innovation process.

We outline the main ideas of the proof. According to (14.2)

$$h(t) = h(0) + \int_0^t H(s)ds + M(t) \quad \text{and} \tag{14.4}$$

$$E(h(t) | \mathcal{F}_t^Y) = E(h(0) | \mathcal{F}_t^Y) + E\left(\int_0^t H(s)ds | \mathcal{F}_t^Y\right) ds + E(M(t) | \mathcal{F}_t^Y).$$

The difficulty in calculation of the conditional expectations given \mathcal{F}_t^Y is that the processes, as well as the σ -fields, both depend on t .

Theorem 14.2: Let $A(t)$ be an \mathcal{F}_t -adapted process such that $\int_0^T E(|A(t)|)dt < \infty$, and $V(t) = \int_0^t A(s)ds$. Then

$$\pi_t(V) - \int_0^t \pi_s(A)ds = E\left(\int_0^t A(s)ds \mid \mathcal{F}_t^Y\right) - \int_0^t E(A(s) \mid \mathcal{F}_s^Y)ds$$

is a \mathcal{F}_t^Y -martingale.

PROOF. Let $\tau \leq T$ be an \mathcal{F}_t^Y -stopping time. According to Theorem 7.17 it is enough to show that $E(\pi_\tau(V)) = E(\int_0^\tau \pi_s(A)ds)$.

$$E(\pi_\tau(V)) = E(V(\tau)) \text{ by the law of double expectation}$$

$$\begin{aligned} &= E\left(\int_0^\tau A(s)ds\right) = \int_0^\tau E(I(s \leq \tau)A(s))ds \\ &= \int_0^\tau E(I(s \leq \tau)\pi_s(A))ds \text{ since } I(s \leq \tau) \text{ is } \mathcal{F}_s^Y\text{-measurable} \\ &= E\left(\int_0^\tau \pi_s(A)ds\right). \end{aligned}$$

□

Note that by the conditional version of Fubini's theorem,

$$E\left(\int_0^t A(s)ds \mid \mathcal{G}\right) = \int_0^t E(A(s) \mid \mathcal{G})ds. \quad (14.5)$$

Verification of the next result is straightforward and is left as an exercise (Exercise 14.4).

Theorem 14.3: Let $M(t)$ be a \mathcal{F}_t -martingale. Then $\pi_t(M)$ is an \mathcal{F}_t^Y -martingale.

Corollary 14.4:

$$\pi_t(h) = \pi_0(h) + \int_0^t \pi_s(H)ds + M_1(t) + M_2(t) + M_3(t), \quad (14.6)$$

where $M_i(t)$ are martingales null at zero; $M_1(t) = E(h(X(0))|\mathcal{F}_t^Y) - h(0)$, $M_2(t) = E\left(\int_0^t H(s)ds \mid \mathcal{F}_t^Y\right) ds - \int_0^t \pi_s(H)ds$, $M_3(t) = E(M(t) \mid \mathcal{F}_t^Y)$.

PROOF. Using (14.4), the first term $E(h(X(0))|\mathcal{F}_t^Y)$ is a (Doob–Levy) martingale. The second term is a martingale by Theorem 14.2, the third by Theorem 14.3. □

We want to use a representation of \mathcal{F}_t^Y martingales. This is done by using the innovation process.

Theorem 14.5: *The innovation process $\bar{W}(t) = \int_0^t \frac{dY(s) - \pi_s(A)ds}{B(Y(s))}$ is an \mathcal{F}_t^Y -Brownian motion. Moreover,*

$$dY(t) = \pi_A(t)dt + B(Y(t))d\bar{W}(t). \quad (14.7)$$

PROOF. By using (14.2)

$$\bar{W}(t) = \int_0^t \frac{A(s) - \mathbb{E}(A(s)|\mathcal{F}_s^Y)}{B(Y(s))} ds + W(t). \quad (14.8)$$

For $t > t'$ write

$$\begin{aligned} \mathbb{E}\left(\bar{W}(t) - \bar{W}(t')|\mathcal{F}_{t'}^Y\right) &= \mathbb{E}\left(W(t) - W(t')|\mathcal{F}_{t'}^Y\right) \\ &\quad + \int_{t'}^t \mathbb{E}\left(\frac{A(s) - \mathbb{E}(A(s)|\mathcal{F}_s^Y)}{B(Y(s))} \mid \mathcal{F}_{t'}^Y\right) ds. \end{aligned}$$

The rhs is zero, the first term by Theorem 14.3, and the second by the conditional version of Fubini's theorem. Thus $\bar{W}(t)$ is an \mathcal{F}_t^Y -martingale. It is clearly continuous. It follows from (14.8) that $[\bar{W}, \bar{W}](t) = [W, W](t) = t$. According to Levy's characterization theorem the claim follows. \square

Now, if

$$\mathcal{F}_t^{\bar{W}} = \mathcal{F}_t^Y, \quad (14.9)$$

then conditional expectations given \mathcal{F}_t^Y are the same given $\mathcal{F}_t^{\bar{W}}$, and Theorem 8.35 on the representation of martingales with respect to a Brownian filtration can be used for the martingales $M_i(t)$ in (14.6). Observe that for (14.9) to hold it is sufficient that the SDE for $Y(t)$ (14.7) has unique weak solution. Theorem 8.35 and its corollary (8.67) give that there are predictable processes $g_i(s)$ such that

$$M_i(t) = \int_0^t g_i(s)d\bar{W}(s), \quad \text{with} \quad g_i(t) = \frac{d\langle M_i, \bar{W} \rangle(t)}{dt}.$$

It is possible to show that

$$g_1(t) + g_2(t) + g_3(t) = \pi_t(hA) + \frac{\pi_t(hA) - \pi_t(h)\pi_t(A)}{B(Y(t))},$$

and the result follows from (14.6). For details see Liptser and Shiryaev (2001).

Filtering of Diffusions

Let $(X(t), Y(t))$, $0 \leq t \leq T$, be a diffusion process with respect to independent Brownian motions $W_i(t)$, $i = 1, 2$; $\mathcal{F}_t = \sigma(W_1(s), W_2(s), s \leq t)$, and

$$\begin{aligned} dX(t) &= a(X(t))dt + b(X(t))dW_1(t) \\ dY(t) &= A(X(t))dt + B(Y(t))dW_2(t). \end{aligned} \quad (14.10)$$

Assume that the coefficients satisfy the Lipschitz condition, for any of the functions a, A, b, B , e.g. $|a(x') - a(x'')| \leq K|x' - x''|$; and $B^2(y) \geq C > 0$.

Let $h = h(X(t))$. The function h is assumed to be twice continuously differentiable. We apply Theorem 14.1 to $h(X(t))$. According to Itô's formula, Theorem 6.1, we have

$$h(X(t)) = h(X(0)) + \int_0^t Lh(X(s))ds + \int_0^t h'(X(s))b(X(s))dW_1(s),$$

where

$$Lh(x) = h'(x)a(x) + \frac{1}{2}h''(x)b^2(x).$$

According to Theorem 14.1

$$\pi_t(h) = \pi_0(h) + \int_0^t \pi_s(Lh)ds + \int_0^t \frac{\pi_s(Ah) - \pi_s(A)\pi_s(h)}{B(Y(s))}d\bar{W}(s), \quad (14.11)$$

where

$$\bar{W}(t) = \int_0^t \frac{dY(s) - \pi_s(A)ds}{B(Y(s))}.$$

The linear case can be solved and is given in the next section.

Kalman–Bucy Filter

Assume that the signal and the observation processes satisfy linear SDEs with time-dependent non-random coefficients

$$dX(t) = a(t)X(t)dt + b(t)dW_1(t), \quad (14.12)$$

$$dY(t) = A(t)X(t)dt + B(t)dW_2(t), \quad (14.13)$$

with two independent Brownian motions (W_1, W_2) , and initial conditions $X(0), Y(0)$. Due to linearity this case admits a closed form solution for the processes X and Y and also for the optimal estimator of $X(t)$ given \mathcal{F}_t^Y .

In the linear case it is easy to solve the above SDEs and verify that the processes $X(t)$, $Y(t)$ are jointly Gaussian. It is convenient in this case to use notation $\hat{X}(t) = \pi_t(X) = E(X(t)|\mathcal{F}_t^Y)$.

Theorem 14.6: Suppose that the signal $X(t)$ and the observation $Y(t)$ are given by (14.12) and (14.13). Then the best estimator $\hat{X}(t) = E(X(t)|\mathcal{F}_t^Y)$ satisfies the following SDE:

$$d\hat{X}(t) = \left(a(t) - v(t) \frac{A^2(t)}{B^2(t)} \right) \hat{X}(t) dt + v(t) \frac{A(t)}{B^2(t)} dY(t), \quad (14.14)$$

where $v(t) = E(X(t) - \hat{X}(t))^2$ is the squared estimation error. It satisfies the Riccati ordinary differential equation

$$\frac{dv(t)}{dt} = 2a(t)v(t) + b^2(t) - \frac{A^2(t)v^2(t)}{B^2(t)}, \quad (14.15)$$

with initial conditions $X(0)$ and $v(0) = Var(X(0)) - \frac{Cov^2(X(0), Y(0))}{Var(Y(0))}$.

PROOF. Apply Theorem 14.1 with $h(x) = x$ (see also Equation (14.11)) to have

$$d\hat{X}(t) = a(t)\hat{X}(t)dt + \frac{A(t)}{B^2(t)} \left(\pi_t(h^2) - (\pi_t(h))^2 \right) (dY(t) - A(t)\hat{X}(t)dt). \quad (14.16)$$

Note that

$$\pi_t(h^2) - (\pi_t(h))^2 = E((X(t) - \hat{X}(t))^2 | \mathcal{F}_t^Y).$$

Because the processes X and Y are jointly Gaussian, $\hat{X}(t)$ is the orthogonal projection, and $X(t) - \hat{X}(t)$ is orthogonal (uncorrelated) to $Y(s)$, $s \leq t$. However, if Gaussian random variables are uncorrelated, they are independent (see Theorem 2.19). Thus $X(t) - \hat{X}(t)$ is independent of \mathcal{F}_t^Y and we have that $v(t)$ is deterministic,

$$v(t) = E((X(t) - \hat{X}(t))^2 | \mathcal{F}_t^Y) = E(X(t) - \hat{X}(t))^2.$$

The initial value $v(0)$ is obtained by the Theorem 2.25 on normal correlation, as stated in the theorem. Let $\delta(t) = X(t) - \hat{X}(t)$. Then $v(t) = E(\delta^2(t))$. The SDE for $\delta^2(t)$ is obtained from SDEs (14.12) and (14.16) as follows:

$$d\delta(t) = a(t)\delta(t)dt + b(t)dW_1(t) - \frac{A^2(t)v(t)}{B^2(t)}\delta(t)dt - \frac{A(t)v(t)}{B(t)}dW_2(t).$$

Applying Itô's formula to $\delta^2(t)$ we now find

$$\begin{aligned} d\delta^2(t) &= \left(2 \left(a(t) - \frac{A^2(t)v(t)}{B^2(t)} \right) \delta^2(t) + \left(b^2(t) + \frac{A^2(t)v^2(t)}{B^2(t)} \right) \right) dt \\ &\quad + 2\delta(t) \left(b(t)dW_1(t) - \frac{A(t)v(t)}{B(t)} dW_2(t) \right). \end{aligned} \quad (14.17)$$

Writing (14.17) in integral form and taking expectation

$$\begin{aligned} v(t) &= v(0) + \int_0^t \left(2 \left(a(s) - \frac{A^2(s)v(s)}{B^2(s)} \right) v(s) + \left(b^2(s) + \frac{A^2(s)v^2(s)}{B^2(s)} \right) \right) ds \\ &= v(0) + \int_0^t \left(2a(s)v(s) + b^2(s) - \frac{A^2(s)v^2(s)}{B^2(s)} \right) ds, \end{aligned} \quad (14.18)$$

which establishes (14.15). \square

The multi-dimensional case is similar, with the only difference being that the Equation (14.15) is the matrix Riccati equation for the estimation error covariance matrix.

The Kalman–Bucy filter allows on-line implementation of the above equations, which are used recursively to compute $\hat{X}(t + \Delta t)$ from the previous values of $\hat{X}(t)$ and $v(t)$.

Example 14.1: (Model with constant coefficients)

Consider the case of constant coefficients,

$$\begin{aligned} dX(t) &= aX(t)dt + dW_1(t), \\ dY(t) &= cX(t)dt + dW_2(t). \end{aligned}$$

In this case

$$d\hat{X}(t) = (a - v(t)c^2) \hat{X}(t)dt + cv(t)dY(t), \quad (14.19)$$

and $v(t)$ satisfies the Riccati equation

$$\frac{dv(t)}{dt} = 2av(t) + 1 - c^2v^2(t). \quad (14.20)$$

This equation has an explicit solution:

$$v(t) = \frac{\gamma\alpha e^{\lambda t} + \beta}{\gamma e^{\lambda t} + 1}, \quad (14.21)$$

where α and β are the roots of $1 + 2ax - c^2x^2$, assumed to be $\alpha > 0$, $\beta < 0$, $\lambda = c^2(\alpha - \beta)$, and $\gamma = (\sigma^2 - \beta)/(\alpha - \sigma^2)$, with $\sigma^2 = \text{Var}(X(0))$. Using $v(t)$ above the optimal estimator $\hat{X}(t)$ is found from (14.19).

For more general results see, for example, Liptser and Shiryaev (2001), Rogers and Williams (1990), and Oksendal (1995). There is a vast amount of literature on filtering; see, for example, Kallianpur (1980), Krishnan (1984), and references therein.

14.2 Random Oscillators

Second order differential equations

$$\ddot{x} + h(x, \dot{x}) = 0$$

are used to describe a variety of physical phenomena, and oscillations are one of them.

Example 14.2: (Harmonic oscillator)

The autonomous vibrating system is governed by $\ddot{x} + x = 0$, $x(0) = 0$, $\dot{x}(0) = 1$, where $x(t)$ denotes the displacement from the static equilibrium position. It has the solution $x(t) = \sin(t)$. The trajectories of this system in the phase space $x_1 = x$, $x_2 = \dot{x}$ are closed circles.

Example 14.3: (Pendulum)

The undamped pendulum is governed by $\ddot{x} + a \sin x = 0$, where $x(t)$ denotes the angular displacement from the equilibrium. Its solution cannot be obtained in terms of elementary functions, but can be given in the phase plane, $(\dot{x})^2 = 2a \cos x + C$.

Example 14.4: (Van der Pol oscillator)

In some systems the large oscillations are damped, whereas small ones are boosted (negative damping). Such motion is governed by the Van der Pol equation

$$\ddot{x} - a(1 - x^2)\dot{x} + x = 0, \quad (14.22)$$

where $a > 0$. Its solution cannot be obtained in terms of elementary functions, even in the phase plane.

Example 14.5: (Rayleigh oscillator)

$$\ddot{x} - a(1 - \dot{x}^2)\dot{x} + x = 0, \quad (14.23)$$

where $a > 0$.

We consider random excitations of such systems by white noise (in applied language) of the form $\sum_{i=1}^2 f_i(x, \dot{x})\dot{W}_i(t)$, where $\dot{W}_i(t)$, $i = 1, 2$, are white noises with delta-type correlation functions

$$E(\dot{W}_i(t)\dot{W}_j(t + \tau)) = 2\pi K_{ij}\delta(\tau)dt.$$

Thus the randomly perturbed equation has the form

$$\ddot{X} + h(x, \dot{X}) = \sum_i f_i(X, \dot{X})\dot{W}_i(t).$$

The white noise is formally the derivative of Brownian motion. Since Brownian motion is nowhere differentiable the above system has only a

formal meaning. A rigorous meaning to such equations is given by a system of first order Itô stochastic equations (a single vector valued equation). The representation as a system of two first order equations follows the same idea as in the deterministic case by letting $x_1 = x$ and $x_2 = \dot{x}$. Using Theorem 5.20 on the conversion of Stratonovich SDEs into Itô SDEs we end up with the following Itô system of stochastic differential equations:

$$\begin{aligned} dX_1 &= X_2 dt, \\ dX_2 &= \left(-h(X_1, X_2) + \pi \sum_{j,k} K_{jk} f_j(X_1, X_2) \frac{\partial f_k}{\partial x_2}(X_1, X_2) \right) dt \\ &\quad + \sum_{i=0}^2 f_i(X_1, X_2) dW_i(t). \end{aligned}$$

The above system is a two-dimensional diffusion, and it has the same generator as the system below driven by a single Brownian motion $B(t)$,

$$dX_1 = X_2 dt, \tag{14.24}$$

$$dX_2 = A(X_1, X_2) dt + G(X_1, X_2) dB(t),$$

where

$$\begin{aligned} A(x_1, x_2) &= -h(x_1, x_2) + \pi \sum_{j,k} K_{jk} f_j(x_1, x_2) \frac{\partial f_k(x_1, x_2)}{\partial x_2}, \\ G(x_1, x_2) &= (2\pi)^{1/2} \left(\sum_{j,k} K_{jk} f_j(x_1, x_2) f_k(x_1, x_2) \right)^{1/2}. \end{aligned}$$

The system (14.24) is a rigorous mathematical model of random oscillators, and this form is the starting point of our analysis. Solutions to the resulting Fokker–Planck equations give densities of invariant measures or stationary distributions for such random systems. The corresponding Fokker–Planck equation has the form

$$x_2 \frac{\partial}{\partial x_1} p_s + \frac{\partial}{\partial x_2} (A p_s) - \frac{1}{2} \frac{\partial^2}{\partial x_2^2} (G^2 p_s) = 0,$$

where $p_s(x_1, x_2)$ is the density of the invariant measure. When a solution to the Fokker–Planck equation has a finite integral, then it is a stationary distribution of the process. However, they may not exist, especially in systems with trajectories approaching infinity. In such cases invariant measures which are not probability distributions may exist and provide information about the underlying dynamics.

In order to illustrate this consider a stochastic differential equation of order two with no noise in \dot{x}

$$\ddot{X} + h(X, \dot{X}) = \sigma \dot{B}.$$

The corresponding Itô system is given by

$$\begin{aligned} dX_1 &= X_2 dt, \\ dX_2 &= -h(X_1, X_2)dt + \sigma dB(t). \end{aligned}$$

In the notation of Chapter 6.10 the dimension $n = 2$ and there is only one Brownian motion, $d = 1$, so that $\mathbf{X}(t) = (X_1(t), X_2(t))^T$, $b(\mathbf{X}(t)) = (b_1(\mathbf{X}(t)), b_2(\mathbf{X}(t)))^T = (X_2(t), -h(X_1(t), X_2(t)))^T$, $\sigma(\mathbf{X}(t)) = (0, \sigma)^T$. The diffusion matrix

$$a = \sigma\sigma^T = \begin{bmatrix} 0 & 0 \\ 0 & \sigma^2 \end{bmatrix},$$

hence the generator is given by

$$\begin{aligned} L &= \sum_{i=1}^2 b_i \frac{\partial}{\partial x_i} + \frac{1}{2} \sum_{i=1}^2 \sum_{j=1}^2 a_{ij} \frac{\partial^2}{\partial x_i \partial x_j} \\ &= x_2 \frac{\partial}{\partial x_1} - h(x_1, x_2) \frac{\partial}{\partial x_2} + \frac{1}{2} \sigma^2 \frac{\partial^2}{\partial x_2^2}. \end{aligned} \quad (14.25)$$

An important example is provided by linear equations

$$\ddot{X} + a\dot{X} + bX = \sigma \dot{B}, \quad (14.26)$$

with constant coefficients. Solutions to these equations can be written in a general formula:

$$\mathbf{X}(t) = (\exp(Ft)) \left(\mathbf{X}(0) + \int_0^t (\exp(-Fs))(0, \sigma)^T dB(s) \right),$$

where $F = \begin{bmatrix} 0 & 1 \\ b & -a \end{bmatrix}$ and $\exp(Ft)$ stands for the matrix exponential (see for example Gard (1988)).

Solutions to the Fokker–Planck equation corresponding to some linear systems (14.26) are given in Bezen and Klebaner (1996).

Non-linear Systems

For some systems the Fokker–Planck equation can be solved by the method of detailed balance. We give examples of such systems, for details see Bezen and Klebaner (1996). In what follows, K_0, K_1 denote scaling constants.

Duffing Equation

Consider additive stochastic perturbations of the Duffing equation:

$$\ddot{X} + a\dot{X} + X + bX^3 = \dot{W}.$$

$$p_s(x_1, x_2) = \exp\left(-\frac{a}{2\pi K_0}(x_2^2 + \frac{1}{2}bx_1^4 + x_1^2)\right).$$

Typical phase portraits and densities of invariant measures are shown in Figure 14.1.

Random Oscillator

Consider the following oscillator with additive noise

$$\ddot{X} - a(1 - X^2 - \dot{X}^2)\dot{X} + X = \dot{W}.$$

The density of the invariant measure is given by

$$p_s = \exp\left(-\frac{a(x_1^2 + x_2^2)^2}{4\pi K_0}\right).$$

It follows that when $a = 0$ the surface representing the invariant density is a plane. When $a > 0$ a fourth order surface with respect to x_1, x_2 has its maximum in a curve representing the limit cycle of the deterministic equation.

Consider now the same equation with parametric noise of the form

$$\ddot{X} - a(1 - X^2 - \dot{X}^2)\dot{X} + X = \dot{W}_0 + (X^2 + \dot{X}^2)\dot{W}_1.$$

The density of the invariant measure is given by

$$p_s = (K_0 + 2K_1x_1^2x_2^2 + K_1x_1^4 + K_1x_2^4)^{-\frac{\sqrt{K_0}(a+2\pi K_1)}{4}}$$

$$\exp\left(\frac{2a \arctan \frac{\sqrt{K_1}(x_1^2+x_2^2)}{\sqrt{K_0}}}{4\pi\sqrt{K_0K_1}}\right).$$

Typical phase portraits and densities of the invariant measures are shown in Figure 14.2.

A System with a Cylindric Phase Plane

Consider random perturbations to a system with a cylindric phase $-\pi \leq x < \pi$, $-\infty < \dot{x} < \infty$,

$$\ddot{X} + a\dot{X} + b + \sin(X) = \dot{W}.$$




Fig. 14.1: The Duffing equation.




Fig. 14.2: Random oscillator.




Fig. 14.3: The system with a cylindric phase plane.

Its invariant density is given by

$$p_s = \exp\left(-\frac{a(x_2^2 + 2bx_1 - 2\cos x_1)}{2\pi K_0}\right).$$

Typical phase portraits and densities of invariant measures are shown in Figure 14.3.

For applications of stochastic differential equations see, for example, Soong (1973). The Fokker–Planck equation was studied, for example, in Soize (1994).

14.3 Exercises

Exercise 14.1: Let X be a square integrable random variable. Show that the value of the constant c for which $E(X - c)^2$ is the smallest is given by EX .

Exercise 14.2: Let X, Y be square integrable random variables. Show that

$$\mathrm{E}(X - \mathrm{E}(X|Y))^2 \leq \mathrm{E}(X - Z)^2$$

for any \mathcal{F}^Y -measurable random variable Z . Hint: show that $X - \mathrm{E}(X|Y)$ and Z are uncorrelated, and write $X - Z = (X - \mathrm{E}(X|Y)) + (\mathrm{E}(X|Y) - Z)$.

Exercise 14.3: Let $M(t)$ be an \mathcal{F}_t -martingale and σ -fields $\mathcal{G}_t \subset \mathcal{F}_t$. Show that if $M(t)$ is \mathcal{G}_t -measurable, then it is a \mathcal{G}_t -martingale.

Exercise 14.4: Let $M(t)$ be an \mathcal{F}_t -martingale and σ -fields $\mathcal{G}_t \subset \mathcal{F}_t$. Show that $\hat{M}(t) = \mathrm{E}(M(t)| \mathcal{G}_t)$ is a \mathcal{G}_t -martingale.

Exercise 14.5: Let $W(t)$ be an \mathcal{F}_t -Brownian motion, and $\mathcal{G}_t \subset \mathcal{F}_t$. According to Exercise 14.4 $\hat{W}(t) = \mathrm{E}(W(t)| \mathcal{G}_t)$ is a \mathcal{G}_t -martingale. Give an example of \mathcal{G}_t such that $\hat{W}(t)$ is not a Brownian motion.

Exercise 14.6: (Observation of a constant)

Let the signal be a constant $X(t) = c$, for all t , and the observation process satisfy $dY(t) = X(t)dt + dW(t)$. Give the Kalman–Bucy filter and find $\hat{X}(t)$.

Exercise 14.7: (Observation of Brownian motion)

Let the signal be a Brownian motion $X(t) = W_1(t)$, and the observation process satisfy $dY(t) = X(t)dt + dW_2(t)$. Give the Kalman–Bucy filter and find $\hat{X}(t)$.

Exercise 14.8: (Filtering of indirectly observed stock prices)

Let the signal follow the Black–Scholes model $X(t) = X(0) \exp(\sigma W_1(t) + \mu t)$, and the observation process satisfy $dY(t) = X(t)dt + dW_2(t)$. Give the Kalman–Bucy filter and find $\hat{X}(t)$.

This page intentionally left blank

Solutions to Selected Exercises

Exercises to Chapter 3

Exercise 3.1: $X = \mu + AZ$ for the vector μ and the vector of independent standard normal random variables Z . For $t = (t_1, \dots, t_n)$

$$E(e^{itX}) = E(e^{it(\mu+AZ)}) = e^{it\mu} E(e^{it(AZ)}) = e^{it\mu} E(e^{i(tA)Z}) = e^{it\mu} \varphi(tA),$$

where φ is the characteristic function of the vector Z . By independence $\varphi(u) = E(e^{iuZ}) = \prod_{j=1}^n E(e^{iu_j Z_j}) = \prod_{j=1}^n e^{-u_j^2/2} = e^{-\frac{1}{2} \sum_{j=1}^n u_j^2}$. Hence $\varphi(tA) = e^{-\frac{1}{2} \sum_{j=1}^n (tA)_j^2} = e^{-\frac{1}{2}(tA)(tA)^T} = e^{-\frac{1}{2}tAA^Tt^T} = e^{-\frac{1}{2}t\Sigma t^T}$. Finally, $E(e^{itX}) = e^{it\mu} e^{-\frac{1}{2}t\Sigma t^T}$.

Exercise 3.2: $EX = \int_0^\infty x dF(x) = \int_0^\infty \int_0^x dt dF(x) = \int_0^\infty \int_t^\infty dF(x) dt = \int_0^\infty (1 - F(t)) dt$.

Exercise 3.3: If $f(t)$ is non-increasing, then $\int_0^\infty f(t) dt = \sum_{n=0}^\infty \int_n^{n+1} f(t) dt \leq \sum_{n=0}^\infty f(n)$. Now apply the previous result.

Exercise 3.9: For $x \geq 0$, by using the distribution of $M(t) = \max_{s \leq t} B(s)$ $P(|B(t)| > x) = P(B(t) > x) + P(B(t) < -x) = 2P(B(t) > x) = P(M(t) > x)$.

Exercise 3.10: According to Theorem 3.18, $E(T_x^r) = \int_0^\infty t^r f(t) dt = \frac{|x|}{\sqrt{2\pi}} \int_0^\infty t^{r-\frac{3}{2}} e^{-\frac{x^2}{2t}} dt = \frac{|x|}{\sqrt{2\pi}} \int_0^\infty s^{r-\frac{1}{2}} e^{-\frac{x^2}{2}s} ds$. The integral converges at infinity for any r . At zero it converges only for $r + \frac{1}{2} < 1$.

Exercise 3.11: $f_M(y) = \int_{-\infty}^\infty f_{B,M}(x, y) dx = \int_{-\infty}^y \sqrt{\frac{2}{\pi}} \frac{(2y-x)}{t^{3/2}} e^{-\frac{(2y-x)^2}{2t}} dx = \sqrt{\frac{2}{\pi}} \frac{1}{t^{1/2}} \int_{-\infty}^y de^{-\frac{(2y-x)^2}{2t}} = \sqrt{\frac{2}{\pi t}} e^{-\frac{y^2}{2t}} = 2f_B(y)$, by (3.16).

Exercise 3.12: $\min_{s \leq t} B(s) = -\max_{s \leq t} -B(s)$. Let $W(t) = -B(t)$, then it is also a Brownian motion and we have $P(B(t) \geq x, \min_{s \leq t} B(s) \leq y) = P(W(t) \leq -x, \max_{s \leq t} W(s) \geq -y) = 1 - \Phi(\frac{-2y+x}{\sqrt{t}})$. With $\phi = \Phi'$, for $y \leq 0, x \geq y$, $f_{B,M}(x,y) = -\frac{\partial^2}{\partial x \partial y} P(B(t) \geq x, m(t) \leq y) = \frac{2}{t} \phi'(\frac{-2y+x}{\sqrt{t}}) = \frac{2}{t} \frac{1}{\sqrt{2\pi}} e^{-\frac{(-2y+x)^2}{2t}} (\frac{-2y+x}{\sqrt{t}}) = \sqrt{\frac{2}{\pi}} \frac{(x-2y)}{t^{3/2}} e^{-\frac{-(2y-x)^2}{2t}}$.

Exercise 3.13: Let $a > 0$. Let $D = \{(x,y) : y - x > a\}$. Then we have $P(M(t) - B(t) > a) = \int \int_D f_{B,M}(x,y) dx dy = \int_0^\infty \int_{-\infty}^{y-a} f_{B,M}(x,y) dx dy = \int_0^\infty \int_{-\infty}^{y-a} \sqrt{\frac{2}{\pi}} \frac{(2y-x)}{t^{3/2}} e^{-\frac{-(2y-x)^2}{2t}} dx dy = \sqrt{\frac{2}{t\pi}} \int_0^\infty (-\int_{-\infty}^{y-a} (\frac{\partial}{\partial x} e^{-\frac{-(2y-x)^2}{2t}}) dx) dy = \sqrt{\frac{2}{t\pi}} \int_0^\infty e^{-\frac{(y+a)^2}{2t}} dy = 2 \int_0^\infty \frac{1}{\sqrt{2\pi t}} e^{-\frac{(y+a)^2}{2t}} dy = 2 \int_a^\infty \frac{1}{\sqrt{2\pi t}} e^{-\frac{u^2}{2t}} du = 2P(B(t) > a)$.

Exercise 3.14: $T_2 = \inf\{t > 0 : B(t) = 0\}$. Since any interval $(0, \epsilon)$ contains a zero of Brownian motion, $T_2 = 0$, the second zero is also zero.

Exercise 3.15: According to the above argument, any zero of Brownian motion is a limit from the right of other zeroes. According to the definition of T , it is a zero of X , but is not a limit of other zeroes. Thus X is not a Brownian motion.

Exercise 3.16: $\limsup_{t \rightarrow \infty} \frac{tB(1/t)}{\sqrt{2t \ln \ln t}} = 1$. $\frac{tB(1/t)}{\sqrt{2t \ln \ln t}} = \frac{B(1/t)}{\sqrt{2(1/t) \ln(-\ln(1/t))}}$. Thus with $\tau = 1/t$, $\limsup_{\tau \rightarrow 0} \frac{B(\tau)}{\sqrt{2\tau \ln(-\ln \tau)}} = 1$. Similarly for \liminf .

Exercise 3.17: $(B(e^{2\alpha t_1}), B(e^{2\alpha t_2}), \dots, B(e^{2\alpha t_n}))$ is a Gaussian vector. The finite-dimensional distributions of $X(t) = e^{-\alpha t} B(e^{2\alpha t})$ are obtained by multiplying this vector by a non-random diagonal matrix A , with the diagonal elements $(e^{-\alpha t_1}, e^{-\alpha t_2}, \dots, e^{-\alpha t_n})$. Therefore finite-dimensional distributions of X are multivariate normal. The mean is zero. Let $s < t$ $Cov(X(s), X(t)) = e^{-\alpha s} e^{-\alpha t} E(B(e^{2\alpha s}) B(e^{2\alpha t})) = e^{-\alpha(s+t)} e^{2\alpha s} = e^{-\alpha(t-s)}$. Note that this Gaussian process has correlated increments.

Exercise 3.18: The process $X(t) = e^{-\alpha t} B(e^{2\alpha t})$ has the given mean and covariance functions and is continuous. Since a Gaussian process is determined by these two functions this is the required version.

Exercise 3.19: $E(e^{uS_{n+1}}|S_n) = E(e^{uS_n} e^{u\xi_{n+1}}|S_n) = E(e^{u\xi_{n+1}})E(e^{uS_n}|S_n) = e^{u^2/2} e^{uS_n}$. Multiply both sides by $e^{-(n+1)u^2/2}$ for the martingale property.

Exercise 3.21: If $X_1 = 3$, then X_2 must be 1, implying that $X_3 = 2$, and cannot be 3, so that $P(X_3 = 3|X_2 = 1 \text{ or } 2, X_1 = 3) = 0$. Using standard calculations of conditional probabilities we can see that $P(X_3 = 3|X_2 = 1 \text{ or } 2) = 1/2$.

Exercise 3.22:

1. If $p = 1$ then $X(t) = aX(t-1) + Z(t)$. $P(X(t) \in A|\mathcal{F}_{t-1}) = P(aX(t-1) + Z(t) \in A|\mathcal{F}_{t-1}) = P(aX(t-1) + Z(t) \in A|X(t-1))$. If $p \geq 2$ then the distribution of $X(t)$ depends on both $X(t-1)$ and $X(t-2)$.
3. The one-step transition probability function is found by $P(X(t) \leq y|X(t-1) = x) = P(aX(t-1) + Z(t) \leq y|X(t-1) = x) = P(ax + Z(t) \leq y|X(t-1) = x) = P(Z(t) \leq y - ax) = \Phi(\frac{y-ax}{\sigma})$. The one-step transition probability density function is $\frac{\partial}{\partial y} \Phi(\frac{y-ax}{\sigma}) = \frac{1}{\sigma} f(\frac{y-ax}{\sigma})$.

Exercises to Chapter 4

Exercise 4.1: The necessary condition is $\int_0^t (t-s)^{-2\alpha} ds < \infty$. This holds if, and only if, $\alpha < 1/2$.

Exercise 4.2: According to (4.3), $X(t) = \sum_{i=0}^n \xi_i I_{(t_i, t_{i+1}]}(t)$ for $t > 0$. Hence $\int_0^t X(s) dB(s) = \int_0^T X(s) I_{(0,t]}(s) dB(s) = \sum_{i=0}^n \xi_i (B(t_{i+1} \wedge t) - B(t_i \wedge t))$, where $t_i \wedge t = \min(t_i, t)$. The Itô integral is continuous as a sum of continuous functions.

Exercise 4.3: The first statement follows by using moment generating functions. Convergence of $e^{\mu_n t + \sigma_n^2 t^2/2}$ implies that $\mu_n \rightarrow \mu$ and $\sigma_n^2 \rightarrow \sigma^2 \geq 0$. If $\sigma^2 = 0$, then the limit is a constant μ , otherwise the limit is $N(\mu, \sigma^2)$.

An Itô integral of a non-random function is a limit of approximating Itô integrals of simple non-random functions $X_n(t)$. Since $X_n(t)$ takes finitely many non-random values, $\int_0^T X_n(t) dB(t)$ has a normal distribution with mean zero and variance $\int_0^T X_n^2(t) dt$. According to the first statement $\int_0^T X(t) dB(t)$ has normal distribution with mean zero and variance $\int_0^T X^2(t) dt$. An alternative derivation of this fact is done by using the martingale exponential of the martingale $u \int_0^t X(s) dB(s)$.

Exercise 4.5: $M(T)$ has a normal distribution, $E M^2(T) < \infty$. By Jensen's inequality for conditional expectation (p. 45) with $g(x) = x^2$, $E(M^2(T)|\mathcal{F}_t) \geq (E(M(T)|\mathcal{F}_t))^2 = M^2(t)$. Therefore $E(M^2(t)) \leq E(M^2(T))$ and $M(t)$ is a square integrable martingale. The covariance between

$M(s)$ and $M(t) - M(s)$ for $s < t$ is zero because by the martingale property $\mathbb{E}(M(t) - M(s)|\mathcal{F}_s) = 0$ and $\mathbb{E}(M(s)(M(t) - M(s)) = \mathbb{E}\mathbb{E}(M(s)(M(t) - M(s))|\mathcal{F}_s) = \mathbb{E}(M(s)\mathbb{E}(M(t) - M(s)|\mathcal{F}_s)) = 0$. Now, if jointly Gaussian variables are uncorrelated they are independent. Thus the increment $M(t) - M(s)$ is independent of $M(s)$.

Let $M(t) = \int_0^t X(s)dB(s)$. Since $X(s)$ is non-random, $\mathbb{E} \int_0^T X^2(s)ds = \int_0^T X^2(s)ds < \infty$ by assumption. Thus the Itô integral $M(t)$ is a martingale. It is also Gaussian by Exercise 4.3. Thus it is a square integrable martingale with independent increments.

Exercise 4.6: Take $f(x) = x^2$, then Itô's formula gives $dX^2(t) = 2X(t)dX(t) + d[X, X](t), [X, X](t) = X^2(t) - X^2(0) - 2 \int_0^t X(s)dX(s)$.

Exercise 4.7: Let $f(x) = \sqrt{x}$, then Itô's formula gives $df(X(t)) = f'(X(t))dX(t) + \frac{1}{2}f''(X(t))d[X, X](t) = \frac{1}{2\sqrt{X(t)}}dX(t) - \frac{1}{8X(t)\sqrt{X(t)}}\sigma^2(X(t))dt$. Rearranging we obtain $dY(t) = \frac{1}{2} \left[bY(t) + \frac{c-1}{Y(t)} \right] dt + dB(t)$.

Exercise 4.10: Let $f(x, t) = txe^{-x}$. Then, if $x = B(t)$, we have $X(t)/Y(t) = f(B(t), t)$. Using Itô's formula (with partial derivatives denoted by subscripts) $df(B(t), t) = f_x(B(t), t)dB(t) + \frac{1}{2}f_{xx}(B(t), t)dt + d[B, B](t) + f_t(B(t), t)dt = t(1 - B(t))e^{-B(t)}dB(t) + \frac{1}{2}t(B(t) - 2)e^{-B(t)}dt + B(t)e^{-B(t)}dt$.

Exercise 4.13: $\mathbb{E}|M(t)| = \mathbb{E}|B^3(t) - 3tB(t)| \leq \mathbb{E}|B^3(t)| + 3t|B(t)| < \infty$, since a normal distribution has all moments. Use the expansion $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ with decomposition $B(t+s) = B(t) + (B(t+s) - B(t))$. Take $a = B(t)$, $b = B(t+s) - B(t)$, and use the fact that $\mathbb{E}(B(t+s) - B(t))^3 = 0$, $\mathbb{E}(B(t+s) - B(t))^2 = s$ to get the martingale property. Using Itô's formula $dM(t) = d(B^3(t) - 3tB(t)) = 3B^2(t)dB(t) + \frac{1}{2}6B^2(t)dt - 3B(t)dt - 3t dB(t) = (3B^2(t) - 3t)dB(t)$. Since $\int_0^T \mathbb{E}(3B^2(t) - 3t)^2 dt < \infty$, $M(t)$ is a martingale on $[0, T]$ for any T .

Exercise 4.15: $df(B_1(t), \dots, B_n(t)) = \sum \frac{\partial f}{\partial x_i}(B)dB_i(t) + \frac{1}{2} \sum_i \frac{\partial^2 f}{\partial x_i^2}(B)dt = \nabla f \cdot dB + \frac{1}{2} \nabla \cdot \nabla f dt$, where “ \cdot ” is the scalar product of vectors and $B = (B_1(t), \dots, B_n(t))$ with $dB = (dB_1(t), \dots, dB_n(t))$. The operator $\nabla \cdot \nabla = \Delta = \sum_i \frac{\partial^2}{\partial x_i^2}$ is the Laplacian so that Itô's formula becomes $df(B(t)) = \frac{1}{2}\Delta f(B(t))dt + \nabla f \cdot dB$.

Exercise 4.16: Denote $\phi(x) = \Phi'(x)$, then $\frac{\partial}{\partial x} \Phi(\frac{x}{\sqrt{T-t}}) = \phi(\frac{x}{\sqrt{T-t}}) \frac{1}{\sqrt{T-t}}$, $\frac{\partial^2}{\partial x^2} \Phi(\frac{x}{\sqrt{T-t}}) = \phi'(\frac{x}{\sqrt{T-t}}) \frac{1}{(T-t)} = -\frac{x}{(T-t)^{3/2}} \phi(\frac{x}{\sqrt{T-t}})$, using $\phi'(x) = -x\phi(x)$

and $\frac{\partial}{\partial t} \Phi(\frac{x}{\sqrt{T-t}}) = -\frac{1}{2} \phi(\frac{x}{\sqrt{T-t}}) \frac{x}{(T-t)^{3/2}}$. Thus by Itô's formula for all $0 \leq t < T$, $d\Phi(\frac{B(t)}{\sqrt{T-t}}) = \phi(\frac{B(t)}{\sqrt{T-t}}) \frac{1}{\sqrt{T-t}} dB(t)$ and $\Phi(\frac{B(t)}{\sqrt{T-t}}) = \frac{1}{2} + \int_0^t \phi(\frac{B(s)}{\sqrt{T-s}}) \frac{1}{\sqrt{T-s}} dB(s)$. Since $\phi(\frac{B(s)}{\sqrt{T-s}}) \frac{1}{\sqrt{T-s}} \leq \frac{1}{\sqrt{T-s}}$ the Itô integral above is a martingale. Thus for all $t < T$ and $s < t$, $X(t) = \Phi(\frac{B(t)}{\sqrt{T-t}})$ satisfies $E(X(t)|\mathcal{F}_s) = X(s)$.

Next, as $t \rightarrow T$, $X(t) \rightarrow Y = I(B(T) > 0) + \frac{1}{2}I(B(T) = 0)$. The martingale property holds also for $t = T$ by dominated convergence.

Exercise 4.18: $dX(t) = dB(t) + B(t)dt$, $d[X, X](t) = (dX(t))^2 = t^2 dt$, $[X, X](t) = \int_0^t s^2 ds = t^3/3$.

Exercise 4.19: $X(t) = tB(t) - \int_0^t s dB(s) = \int_0^t s dB(s) + \int_0^t B(s)ds - \int_0^t s dB(s) = \int_0^t B(s)ds$. Thus $X(t)$ is differentiable and of finite variation. Hence $[X, X](t) = 0$. Itô integrals of the form $\int_0^t h(s)dB(s)$ have a positive quadratic variation. In this case the Itô integral is of the form $\int_0^t h(t, s)dB(s)$.

Exercises to Chapter 5

Exercise 5.1: According to Example 4.5, $\int_0^t b(s)dB(s)$ is a Gaussian process. $X(t) = \int_0^t a(s)ds + \int_0^t b(s)dB(s)$ is also Gaussian as $\int_0^t a(s)ds$ is non-random.

Exercise 5.3: $dX(t) = X(t)(B(t)dt + B(t)dB(t))$. $X(t) = \mathcal{E}(R)(t)$, where $R(t) = \int_0^t B(s)ds + \int_0^t B(s)dB(s)$. Thus $X(t) = e^{\int_0^t (B(s) - \frac{1}{2}B^2(s))ds + \int_0^t B(s)dB(s)}$.

Exercise 5.4: Let $dM(t) = B(t)dB(t)$. Then $dX(t) = X(t)dt + dM(t)$, which is a Langevin type SDE. Solving similarly to Example 5.6 we obtain $X(t) = e^t(1 + \int_0^t e^{-s}dM(s)) = e^t(1 + \int_0^t e^{-s}B(s)dB(s))$. The SDE is not of the diffusion type as $\sigma(t) = B(t)$. By introducing $Y(t) = B(t)$ it is a diffusion in two dimensions.

Exercise 5.5: Let $U(t) = \mathcal{E}(B)(t)$. Then $dU(t) = U(t)dB(t)$. $dU^2(t) = 2U(t)dU(t) + d[U, U](t) = 2U^2(t)dB(t) + U^2(t)dt = U^2(t)d(2B(t) + t)$. So that $U^2(t) = \mathcal{E}(2B(t) + t)$.

Exercise 5.6: $dX(t) = X(t)(X(t)dt + dB(t))$. If $dY(t) = X(t)dt + dB(t)$, then $X(t) = \mathcal{E}(Y)(t)$.

Exercise 5.10: By definition, $P(y, t, x, s) = P(B(t) + t \leq y | B(s) + s = x) = P(B(t) - B(s) + t - s \leq y - x | B(s) + s = x) = P(B(t) - B(s) + t - s \leq y - x)$ by independence of increments. $B(t) - B(s)$ has $N(0, t - s)$ distribution, and $P(y, t, x, s) = \Phi\left(\frac{y-x-t+s}{\sqrt{t-s}}\right)$.

Exercise 5.12: $X(t) = \int_0^t \sqrt{X(s) + 1} dB(s)$. Assuming it is a martingale $\mathbb{E}X(t) = 0$, $\mathbb{E}X^2(t) = \mathbb{E}\left(\int_0^t \sqrt{X(s) + 1} dB(s)\right)^2 = \mathbb{E}\left(\int_0^t (X(s) + 1) ds\right) = t$. $de^{uX(t)} = ue^{uX(t)} dX(t) + \frac{1}{2}u^2 e^{uX(t)} d[X, X](t)$. $d[X, X](t) = (X(t) + 1)dt$, and $de^{uX(t)} = ue^{uX(t)} \sqrt{X(t) + 1} dB(t) + \frac{1}{2}u^2 e^{uX(t)} (X(t) + 1)dt$. Taking expectation $m(t) = 1 + \frac{1}{2}u^2 \mathbb{E}\left(\int_0^t e^{uX(s)} (X(s) + 1) ds\right) = 1 + \frac{1}{2}u^2 \int_0^t \mathbb{E}(e^{uX(s)} X(s)) ds + \frac{1}{2}u^2 \int_0^t \mathbb{E}(e^{uX(s)}) ds$. Thus $\frac{\partial m}{\partial t} = \frac{u^2}{2} \mathbb{E}(e^{uX(t)} X(t)) + \frac{u^2}{2} \mathbb{E}e^{uX(t)}$ and the desired PDE follows.

Exercises to Chapter 6

Exercise 6.1: Denote $L = \frac{1}{2} \frac{\partial^2}{\partial x^2}$ and $f(x, t) = e^{ux - u^2 t/2}$. Then $Lf = \frac{1}{2}u^2 f$ and $\frac{\partial f}{\partial t} = -\frac{1}{2}u^2 f$. Thus $Lf + \frac{\partial f}{\partial t} = 0$. According to Itô's formula and Corollary 6.4 $f(B(t), t) = e^{uB(t) - u^2 t/2}$ is a martingale. Since f solves $Lf + \frac{\partial f}{\partial t} = 0$ for any fixed u , take partial derivative $\frac{\partial f}{\partial u}$ and interchange the order of differentiation to have that for any fixed u , $\frac{\partial}{\partial u} Lf + \frac{\partial}{\partial u} \frac{\partial f}{\partial t} = L\left(\frac{\partial f}{\partial u}\right) + \frac{\partial}{\partial t} \frac{\partial f}{\partial u} = 0$, so that $\frac{\partial f}{\partial u}$ solves the backward equation for any fixed u and in particular for $u = 0$. Calculating the derivatives we get the result.

Exercise 6.2: The generator is given by (6.30) $L = \frac{\sigma^2}{2} \frac{d^2}{dx^2} - \alpha x \frac{d}{dx}$. The backward equation is $\frac{\sigma^2}{2} \frac{\partial^2 f}{\partial x^2} - \alpha x \frac{\partial f}{\partial x} = \frac{\partial f}{\partial t}$. The fundamental solution is given by the probability density of the solution to the SDE $X(t)$. $p(t, x, y) = \frac{\partial}{\partial y} P(t, x, y)$, where $P(t, x, y) = P(X(t) \leq y | X(0) = x)$. According to (5.13) $P(t, x, y) = P(xe^{-\alpha t} + e^{-\alpha t} \int_0^t e^{\alpha s} dB(s) \leq y) = P\left(\int_0^t e^{\alpha s} dB(s) \leq ye^{\alpha t} - x\right) = \Phi\left(\frac{\sqrt{\alpha}(ye^{\alpha t} - x)}{\sqrt{e^{\alpha t} - 1}}\right)$ since $\int_0^t e^{\alpha s} dB(s)$ has $N(0, \frac{1}{\alpha}(e^{\alpha t} - 1))$ distribution. Thus $p(t, x, y) = \frac{\partial}{\partial y} P(t, x, y) = \phi\left(\frac{\sqrt{\alpha}(ye^{\alpha t} - x)}{\sqrt{e^{\alpha t} - 1}}\right) \frac{\sqrt{\alpha}e^{\alpha t}}{\sqrt{e^{\alpha t} - 1}}$ with ϕ denoting the density of $N(0, 1)$.

Exercise 6.4: $L = \frac{1}{2} \frac{\partial^2}{\partial x^2} + cx \frac{\partial}{\partial x}$. Take $f(x) = x^2$, then $Lf(x) = 1 + 2cx^2$ and by Theorem 6.11 $X^2(t) - \int_0^t (Lf + \frac{\partial f}{\partial t})(X(s)) ds = X^2(t) - t - 2c \int_0^t X^2(s) ds$ is a martingale.

Exercise 6.6: Using Corollary 6.4 or Itô's formula we have $df(B(t) + t) = f'(B(t) + t)dB(t) + f'(B(t) + t)dt + \frac{1}{2}f''(B(t) + t)dt$. A necessary condition

for $f(B(t) + t)$ to be a martingale is that the dt term is zero. This gives $f'(x) + \frac{1}{2}f''(x) = 0$. For example, take $f(x) = e^{-2x}$ and check directly that $e^{-2(B(t)+t)}$ is a martingale. ($e^{-2(B(t)+t)} = \mathcal{E}(2B)(t)$).

Exercise 6.7: $df(X(t), t) = \frac{\partial f}{\partial x}dX(t) + \frac{1}{2}\frac{\partial^2 f}{\partial x^2}d[X, X](t) + \frac{\partial f}{\partial t}dt$. The term with $dB(t)$ is given by $\sigma(X(t), t)\frac{\partial f}{\partial x}(X(t), t)dB(t)$. Thus the PDE for f is $\sigma(x, t)\frac{\partial f}{\partial x}(x, t) = 1$.

Exercise 6.8: By letting $v' = y$ we obtain a first order differential equation $y' + \frac{2\mu}{\sigma^2}y + \frac{2}{\sigma^2} = 0$. Use the integrating factor $V(x) = \exp\left(\int_a^x \frac{2\mu(s)}{\sigma^2(s)}ds\right)$ to obtain $(yV)' = -\frac{2}{\sigma^2}V$ and $y = -\frac{1}{V} \int \frac{2}{\sigma^2}V$. The result follows.

Exercise 6.9: The scale function (6.50) $S'(x) = e^{-\int^x \frac{2\mu}{\sigma^2}ds} = e^{-\frac{2\mu}{\sigma^2}x}$ gives $S(x) = Ce^{-\frac{2\mu}{\sigma^2}x}$. $P_x(T_b < T_a) = \frac{S(x)-S(a)}{S(b)-S(a)} = (e^{-\frac{2\mu}{\sigma^2}x} - e^{-\frac{2\mu}{\sigma^2}a})/(e^{-\frac{2\mu}{\sigma^2}b} - e^{-\frac{2\mu}{\sigma^2}a})$.

Exercise 6.10: The generator of Brownian motion is $\frac{1}{2}\frac{\partial^2 f}{\partial x^2}$. According to Theorem 6.6 $f(x, t) = \mathbb{E}(B^2(T)|B(t) = x) = \mathbb{E}((x + B(T) - B(t))^2|B(t) = x)$. By independence of increments it is the same as $\mathbb{E}(x + B(T) - B(t))^2 = x^2 + T - t$.

Exercise 6.12: Check condition 2, Theorem 6.23, to decide convergence of the integral $\int_1^\infty \exp\left(-\int_1^x \frac{2s^2}{\sigma^2 s^{2\alpha}} ds\right) \left(\int_1^x \frac{1}{\sigma^2 y^{2\alpha}} \exp\left(\int_1^y \frac{2s^2}{\sigma^2 s^{2\alpha}} ds\right) dy\right) dx$. $\int_1^\infty \frac{2s^2}{\sigma^2 s^{2\alpha}} ds = \frac{2}{\sigma^2} \frac{x^{3-2\alpha}-1}{3-2\alpha}$. When $3-2\alpha > 0$, then $\exp(-x^{3-2\alpha}/\sigma^2)$ converges. Consequently, the process explodes. When $3-2\alpha < 0$, then the integral diverges and there is no explosion. The case $\alpha = 3/2$ needs further analysis.

Exercise 6.13: Check the conditions of Theorem 6.28. $\mu(x) = 0$, $\sigma(x) = 1$. $I_1 = \int_{-\infty}^{x_0} 1du = \infty$, $I_2 = \int_{x_0}^\infty 1du = \infty$. Hence $B(t)$ is recurrent. For the process $B(t) + t$, $\mu(x) = 1$ and $\sigma(x) = 1$. $I_2 = \int_{x_0}^\infty \exp\left(-2(u-x_0)\right) du = \frac{1}{2} < \infty$. Thus $B(t) + t$ is transient. The Ornstein–Uhlenbeck process is left to the reader.

Exercise 6.15: For $n = 2$, $S(x) = \ln x$. So that for $0 < y < x$, $P_x(T_y < T_b) = \frac{\ln x - \ln b}{\ln y - \ln b}$. Since $X(t)$ does not explode $T_b \uparrow \infty$ as $b \uparrow \infty$. Therefore $P_x(T_y < \infty) = \lim_{b \rightarrow \infty} P_x(T_y < T_b) = \lim_{b \rightarrow \infty} \frac{\ln x - \ln b}{\ln y - \ln b} = 1$. For $n \geq 3$,

$S(x) = \frac{1-x^{1-n/2}}{1-n/2}$. $P_x(T_y < T_b) = \frac{b^{1-n/2}-x^{1-n/2}}{b^{1-n/2}-y^{1-n/2}} \rightarrow (\frac{y}{x})^{n/2-1} < 1$. The explosion test shows that $X(t)$ does not explode, and $\lim_{b \rightarrow \infty} T_b = \infty$.

Exercise 6.17: Check (6.67). Since $1 = \int_{\alpha}^{\beta} p(t, x, y) dy = \int_{\alpha}^{\beta} p(t, y, x) dy$, for any C , $C = \int_{\alpha}^{\beta} Cp(t, y, x) dy$. Since C is a distribution on (α, β) , it must be the uniform density.

Exercise 6.18: In order to classify 0 as a boundary calculate L_1 , L_2 , and L_3 , Remark 6.6, see also Theorem 6.29. $\mu(x) = b(a-x)$, $\sigma^2(x) = \sigma^2 x$. For constants C_1, C_2 $C_1 \int_0^c u^{-2ba/\sigma^2} du \leq L_1 \leq C_2 \int_0^c u^{-2ba/\sigma^2} du$, which shows that L_1 converges if $2ba/\sigma^2 < 1$. Thus 0 is a natural boundary if, and only if, $2ba/\sigma^2 \geq 1$. $L_2 < \infty$. $L_3 < \infty$ if $ba > 0$. So if $0 < 2ba/\sigma^2 < 1$, 0 is a regular boundary point and if $ba < 0$, 0 is an absorbing boundary.

Exercises to Chapter 7

Exercise 7.1: $\mathcal{G}_t \subseteq \mathcal{F}_t$. Let $s < t$. According to the smoothing property of conditional expectation 3, p. 45 (double expectation), $E(M(t)|\mathcal{G}_s) = E(E(M(t)|\mathcal{F}_s)|\mathcal{G}_s) = E(M(s)|\mathcal{G}_s) = M(s)$.

Exercise 7.3: By convexity, property 6 p. 45 of conditional expectation, $E(g(X(t))|\mathcal{F}_s) \geq g(E(X(t)|\mathcal{F}_s))$. By the submartingale property $E(X(t)|\mathcal{F}_s) \geq X(s)$. Since g is non-decreasing the result follows.

Exercise 7.4: Since square-integrable martingales are uniformly integrable (Corollary 7.8) there is Y such that $M(t) = E(Y|\mathcal{F}_t)$. $Y = \lim_{t \rightarrow \infty} M(t)$. According to Fatou's lemma (p. 39) $E(Y^2) \leq \liminf_{t \rightarrow \infty} E(M^2(t)) < \infty$.

Exercise 7.5: Let τ be the first time $B(t)$ hits a or b when $B(0) = x$, $a < x < b$. τ is finite by Theorem 3.13. By stopping martingale $B(t)$ we find that $P(B(\tau) = b) = (x-a)/(b-a)$, $P(B(\tau) = a) = 1 - P(B(\tau) = b)$, Example 7.6. Stopping $M(t)$, $EM(\tau \wedge t) = M(0) = x^2$, or $EB^2(\tau \wedge t) - E(\tau \wedge t) = x^2$. Take $t \rightarrow \infty$. $\tau \wedge t \rightarrow \tau$, $B^2(\tau \wedge t) \rightarrow B^2(\tau)$, and by dominated convergence $EB^2(\tau) - E(\tau) = x^2$. However, $EB^2(\tau) = a^2P(B(\tau) = a) + b^2P(B(\tau) = b) = a^2 \frac{b-x}{b-a} + b^2 \frac{x-a}{b-a}$. Thus $E(\tau) = EB^2(\tau) - x^2$, and the result follows.

Exercise 7.6: Let $\tau = \inf\{t : B(t) - t/2 = a \text{ or } b\}$. As in the previous exercise we obtain $EM(\tau) = M(0) = e^x$ or $e^a P(M(\tau) = e^a) + e^b (1 - P(M(\tau) = e^b)) = e^x$. This gives $P(B(\tau) - \tau/2 = a) = P(M(\tau) = e^a) = \frac{e^b - e^x}{e^b - e^a}$.

Exercise 7.8: When the game is fair $p = q = 1/2$, use formula (7.17) to see that $u \rightarrow 1$. If $p \neq q$, use (7.20) to see that $u \rightarrow 1$ when $p < q$, and $u \rightarrow (q/p)^x$ when $p > q$.

Exercise 7.12: $\int_0^T \text{sign}^2(B(s))ds = T < \infty$. Thus $X(t)$ is a martingale. $[X, X](t) = \int_0^t \text{sign}^2(B(s))ds = t$. According to Levy's theorem X is a Brownian motion.

Exercise 7.13: $[M, M](t) = \int_0^t e^{2s}ds = \frac{1}{2}(e^{2t} - 1)$. Its inverse function is $g(t) = \frac{1}{2}\ln(2t + 1)$. $M(g(t))$ is a Brownian motion by the DDS Theorem 7.37.

Exercise 7.16: $X(t) = X(0) + A(t) + M(t) = X(0) + \int_0^t \mu(s)ds + \int_0^t \sigma(s)dB_s$ is a local martingale. Therefore $A(t) = X(t) - M(t)$ is a local martingale as a difference of two local martingales. $A(t)$ is continuous. By using Corollary 7.30 $A(t)$ has infinite variation unless it is a constant. Since $A(t)$ is of finite variation it must be zero. The result follows.

Exercise 7.17: By Itô's formula $f(B(t), t) - \int_0^t \left(\frac{\partial f}{\partial t}(B(s), s) + \frac{1}{2} \frac{\partial^2 f}{\partial x^2}(B(s), s) \right) ds = \int_0^t \frac{\partial f}{\partial x}(B(s), s) dB(s)$. The rhs is a continuous local martingale, and the lhs is of finite variation. This can only happen when the local martingale is a constant. Thus $\frac{\partial f}{\partial x} = 0$, and f is a constant in x , hence a function of t alone.

Exercise 7.18: We know $Y(t) = \frac{1}{2}B^2(t) - \frac{1}{2}t$, and $B^2(t) = 2Y(t) + t$. Thus $B(t) = \text{sign}(B(t))\sqrt{2Y(t) + t}$. Therefore $dY(t) = \text{sign}(B(t))\sqrt{2Y(t) + t}dB(t)$ or $dY(t) = \sqrt{2Y(t) + t}dW(t)$, where $dW(t) = \text{sign}(B(t))dB(t)$, W is a Brownian motion. Weak uniqueness follows by Theorem 5.11. Alternatively, one can prove it directly by calculating of moments of $Y(t)$, $\mu_n(t) = EY^n(t)$ (by using Ito's formula $\mu_n(t) = \frac{n(n-1)}{2} \int_0^t (s\mu_{n-2}(s) + 2\mu_{n-1}(s))ds$) and checking Carleman's condition $\sum(\mu_n)^{-1/n} = \infty$, see Feller (1971).

Exercises to Chapter 8

Exercise 8.1: $I_{(\tau_1, \tau_2]}(t) = I_{[0, \tau_2]}(t) - I_{[0, \tau_1]}(t)$ is a sum of two left-continuous adapted functions. Let τ be a stopping time. Then $X(t) = I_{[0, \tau]}(t)$ is adapted. $\{X(t) = 0\} = \{t > \tau\} = \bigcup_{n=1}^{\infty} \{\tau \leq t - 1/n\} \in \mathcal{F}_t$ since $\{\tau \leq t - 1/n\} \in \mathcal{F}_{t-1/n} \subset \mathcal{F}_t$. See p. 52.

Exercise 8.2: The left-continuous modification of the process $H(t - \delta + \epsilon)$ is adapted and as $\epsilon \rightarrow 0$ tends to $H(t - \delta)$.

Exercise 8.4: Let $X(t) = \int_0^t H(s)dM(s)$, then it is a local martingale as a stochastic integral with respect to a local martingale.

$[X, X](t) = \int_0^t H^2(s)d[M, M](s)$. Thus $E[X, X](T) < \infty$. According to Theorem 7.35 this condition implies X is a square integrable martingale.

Exercise 8.5: $E(\int_0^1 N(t-)dM(t)) = 0$, so that $Var(\int_0^1 N(t-)dM(t)) = E(\int_0^1 N(t-)dM(t))^2 = E\int_0^1 N^2(t-)d[M, M](t)$. $[M, M](t) = [N, N](t) = N(t)$ so that $Var(\int_0^1 N(t-)dM(t)) = E\int_0^1 N^2(t-)dN(t) = E(\sum_{\tau_i \leq 1} N^2(\tau_{i-1}))$, where τ_i denotes the time of the i -th jump of N . However, $N(\tau_i) = i$ so that $Var(\int_0^1 N(t-)dM(t)) = E(\sum_{\tau_i \leq 1} (i-1)^2) = E(\sum_{i=1}^{N(1)} (i-1)^2) = E(\sum_{k=0}^{N(1)-1} k^2) = E(2N^3(1) - 3N^2(1) + N(1))/6$, since $\tau_i \leq 1$ is equivalent to $i \leq N(1)$, and $\sum_{k=0}^n k^2 = (2(n+1)^3 - 3(n+1)^2 + n+1)/6$. Alternatively, $\int_0^1 N^2(t-)dN(t)$ can be obtained by using formula (1.20). Compute moments using the mgf $m(s) = Ee^{sN(1)} = e^{-1}e^{e^s}$. $m'(0) = EN(1) = 1$, $m''(0) = EN^2(1) = 2$, $m^{(3)}(0) = EN^3(1) = 5$, giving $Var(\int_0^1 N(t-)dM(t)) = 5/6$.

Exercise 8.6:

1. $S \wedge T$ and $S \vee T$ are stopping times because $\{S \wedge T > t\} = \{S > t\} \cap \{T > t\} \in \mathcal{F}_t$, $\{S \vee T \leq t\} = \{S \leq t\} \cap \{T \leq t\} \in \mathcal{F}_t$.
2. The events $\{S = T\}$, $\{S \leq T\}$, and $\{S < T\}$ are in \mathcal{F}_S .
3. $\mathcal{F}_S \cap \{S \leq T\} \subset \mathcal{F}_T \cap \{S \leq T\}$. See Theorem 2.38.

Exercise 8.9: $M(t)$ is a continuous martingale, $[M, M](t) = \int_0^t H^2(s)ds = t$. According to Levy's theorem it is a Brownian motion. If $M(t)$ is a Brownian motion, then $[M, M](t) = \int_0^t H^2(s)ds = t$ for all t . Taking derivatives $H^2(t) = 1$ Lebesgue almost surely.

Exercise 8.10: The proof is not easy if done from basic properties. However, it is easy when using Levy's theorem. $M(t)$ can be written as $B(t) = \int_0^t dB(s)$, $B(t \wedge T) = \int_0^t I_{[0,T]}(s)dB(s)$, $M(t) = \int_0^t (2I_{[0,T]}(s) - 1)dB(s)$. Exercise 8.1 shows that $I_{[0,T]}(s)$ is predictable. Clearly, $M(t)$ is a continuous local martingale as an Itô integral. It is also a martingale, as $E\int_0^t (2I_{[0,T]}(s) - 1)^2 ds \leq 9t < \infty$. $[M, M](t) = \int_0^t (2I_{[0,T]}(s) - 1)^2 ds$.

If $T \geq t$, then for all $s \leq t$, $I_{[0,T]}(s) = 1$ implying $[M, M](t) = t$. If $T < t$, then $[M, M](t) = \int_0^T (2I_{[0,T]}(s) - 1)^2 ds + \int_T^t (-1)^2 ds = T + (t - T) = t$. Thus $[M, M](t) = t$, for any T , and M is a Brownian motion by Levy's theorem.

Exercise 8.11: $M = \bar{N}$. $d(B(t)M(t)) = B(t)dM(t) + M(t)dB(t) + d[B, M](t)$. Since M is of finite variation, by property 7 of quadratic variation (8.18) $[B, M](t) = \sum_{s \leq t} \Delta B(s)\Delta M(s) = 0$ since by continuity of Brownian motion $\Delta B(s) = 0$. Thus $B(t)M(t) = \int_0^t B(s)dM(s) + \int_0^t M(s)dB(s)$. Both stochastic integrals satisfy the condition to be a martingale, i.e. $E \int_0^T B^2(s)d[M, M](s) = E \int_0^T B^2(s)dN(s) = E \sum_{\tau_i \leq T} B^2(\tau_i) = E \sum_{i=1}^{N(T)} \tau_i < TEN(T) = T^2 < \infty$. $E \int_0^T M^2(s)d[B, B](s) = E \int_0^T M^2(s)ds = \int_0^T Var(N(s))ds = \int_0^T sds < \infty$. The other two processes are martingales by similar arguments, verifying that a purely discontinuous martingale is orthogonal to any continuous martingale, i.e. their product is a martingale, p. 233.

Exercise 8.12: $dX(t) = \mu X(t)dt + aX(t-)\lambda dt + aX(t-)(dN(t) - \lambda dt) + \sigma X(t)dB(t)$. $aX(t-)(dN(t) - \lambda dt) + \sigma X(t)dB(t) = dM(t)$ is a martingale, as a sum of stochastic integrals with respect to martingales. $\int X(t-)dt = \int X(t)dt$. Thus $X(t)$ is a martingale when $\mu = -a\lambda$.

Exercise 8.13: $B^5(1) = \int_0^1 (5B^4(t) + 30(1-t)B^2(t) + 15(1-t)^2) dB(t)$. Use Itô's formula for $B^5(t)$ and that the following functions $x^5 - 10tx^3 + 15t^2x$ and $x^3 - 3xt$ produce martingales.

Exercise 8.15: $\int_0^1 (\text{sign}(B(s)) - H(s))dB(s) = 0$. The proposition follows from $E \int_0^1 (\text{sign}(B(s)) - H(s))^2 ds = 0$.

Exercise 8.17: We show the continuous case, the case with jumps is similar. $\mathcal{E}(X)(t) = e^{X(t)-\frac{1}{2}[X, X](t)}$. According to (8.82) $X(t) = \ln \left| \frac{U(t)}{U(0)} \right| + \frac{1}{2} \int_0^t \frac{d[U, U](s)}{U^2(s)}$. According to Itô's formula for $\ln U(t)$ we obtain $[X, X](t) = \int_0^t \frac{d[U, U](s)}{U^2(s)}$. Thus $\mathcal{E}(X)(t) = U(t)$.

Exercises to Chapter 9

Exercise 9.1: Change the order of integration.

Exercise 9.3: $P(T_1 = x) = p(1-p)^{x-1}$, $x = 1, 2, \dots$. Use the lack of memory property of geometric distribution, $P(T_1 \leq t + c | T_1 \geq t) = P(T_1 \leq c)$ to see that the compensator of $N(t)$ is $p[t]$, where $[t]$ stands for the integer part of t .

Exercise 9.4: Repeat the proof of Theorem 9.7 and condition on \mathcal{F}_{T_n-} .

Exercise 9.5: Apply Theorem 9.21 to $f(x) = x^k$.

Exercises to Chapter 10

Exercise 10.2: According to Theorem 10.3 the equivalent measure Q is given by $dQ/dP = e^{-\mu X + \mu^2/2}$ and $dP/dQ = e^{\mu X - \mu^2/2}$.

Exercise 10.3: Since $Y > 0$ and $E(Y/EY) = 1$, let $\Lambda = Y/EY$ and define $dQ/dP = \Lambda = e^{X-\mu-\sigma^2/2}$. According to Theorem 10.4, the Q distribution of Y is $N(\mu + \sigma^2, \sigma^2)$. Thus $EYI(Y > K) = EYE\Lambda I(Y > K) = EYE_Q I(Y > K) = e^{\mu+\sigma^2/2}Q(Y > K)$. Finally, $EYI(Y > K) = e^{\mu+\sigma^2/2}\Phi((\mu + \sigma^2 - K)/\sigma)$, using $1 - \Phi(x) = \Phi(-x)$.

Exercise 10.4: $X(t) = B(t) + \int_0^t \cos s ds$. Thus by Girsanov's Theorem 10.16 $dQ/dP = e^{-\int_0^T \cos s dB(s) - \frac{1}{2} \int_0^T \cos^2 s ds}$.

Exercise 10.5: With $X = \mathbf{H} \cdot \mathbf{B}$, $\mathcal{E}(X)(T) = e^{X(T)-X(0)-\frac{1}{2}[X,X](T)}$. $[\mathbf{H} \cdot \mathbf{B}, \mathbf{H} \cdot \mathbf{B}](T) = \sum_{i,j=1}^d [H^i \cdot B^i, H^j \cdot B^j] = \sum_{i,j=1}^d \int_0^T H^i(t) H^j(t) d[B^i, B^j](t)$. $\sum_{i=1}^d \int_0^T (H^i(t))^2 dt = \int_0^T |\mathbf{H}(s)|^2 ds$, since $[B^i, B^j](t) = 0$ for $i \neq j$. Thus $\mathcal{E}(\mathbf{H} \cdot \mathbf{B})(T) = e^{\sum_{i=1}^d \int_0^T H^i(s) dB^i(s) - \frac{1}{2} \sum_{i=1}^d \int_0^T (H^i(s))^2 ds} = e^{\int_0^T \mathbf{H}(s) d\mathbf{B}(s) - \frac{1}{2} \int_0^T |\mathbf{H}(s)|^2 ds}$.

Exercise 10.8: Since $H(t)$ is bounded $E \int_0^T H^2(t) d[\bar{N}, \bar{N}](t) = E \int_0^T H^2(t) dt$ is finite and $M(t)$ is a martingale. According to (9.5) $\mathcal{E}(M)(t) = e^{\int_0^t H(s) d\bar{N}(s)} \prod_{s \leq t} (1 + \Delta(H \cdot \bar{N})(s)) e^{-\Delta(H \cdot \bar{N})(s)}$. However, the jumps of the integral occur at the points of jumps of \bar{N} and $\Delta \bar{N} = \Delta N$ so that $\Delta(H \cdot N)(s) = H(s) \Delta N(s)$. Next, $\prod_{s \leq t} e^{-\Delta(H \cdot \bar{N})(s)} = e^{-\sum_{s \leq t} H(s) \Delta N(s)} = e^{-\int_0^t H(s) dN(s)}$. Proceeding, $\prod_{s \leq t} (1 + \Delta(H \cdot \bar{N})(s)) = e^{\sum_{s \leq t} \ln(1+H(s)) \Delta N(s)} = e^{\int_0^t \ln(1+H(s)) dN(s)}$, and the result follows.

Exercise 10.9: Using (10.52) or (10.51) with $\mu_i(x, t) = \mu_i x$, $i = 1, 2$, $\sigma(x, t) = \sigma x$, P corresponds to μ_1 , $B(t)$ is a P -Brownian motion. $\Lambda(X)_T = \frac{dQ}{dP} = \mathcal{E}(\frac{\mu_2 - \mu_1}{\sigma} B)(T) = e^{\frac{\mu_2 - \mu_1}{\sigma} B(T) - \frac{(\mu_2 - \mu_1)^2}{2\sigma^2} T}$. Replace $B(T)$ by its expression as a function of X . In this case $B(T) = (\ln(\frac{X(T)}{X(0)}) - (\mu_1 - \frac{1}{2}\sigma^2)T)/\sigma$.

Exercise 10.11: According to Corollary 10.11 $M'(t)$ is a Q -martingale if, and only if, $M'(t)\Lambda(t)$ is a P -martingale. $d(M'(t)\Lambda(t)) = M'(t)d\Lambda(t) + \Lambda(t)dM'(t) + d[M', \Lambda](t) = M'(t)d\Lambda(t) + \Lambda(t)dM(t) - \Lambda(t)dA(t) + d[M', \Lambda](t)$. The first two terms are P (local)-martingales, as stochastic integrals with respect to martingales ($\Lambda(t)$ is a P -martingale). Thus $-\Lambda(t)dA(t) + d[M', \Lambda](t) = 0$. However, $[A, \Lambda](t) = 0$ since A is continuous and of finite variation. Thus $dA(t) = d[M, \Lambda](t)/\Lambda(t)$.

Exercise 10.13:

- Using Girsanov's Theorem 10.15 there exists an equivalent measure Q_1 (defined by $\Lambda = dQ_1/dP = e^{-T/2-B(T)}$) such that $W(t) = B(t) + t$ is a Q_1 -martingale. We show that under Q_1 , $N(t)$ remains to be a Poisson process with rate 1. This follows by independence of $N(t)$ and $B(t)$ under P , $E_Q e^{uN(t)} = E_P(\Lambda e^{uN(t)}) = E_P(e^{-T/2-B(T)} e^{uN(t)}) = E_P(e^{-T/2-B(T)}) E_P(e^{uN(t)}) = E_P(e^{uN(t)})$, since $E_P \Lambda = 1$. This shows that one-dimensional distributions of $N(t)$ are Poisson (1). Similarly, the increments of $N(t)$ under Q_1 are Poisson and independent of the past. The statement follows.
- Let $\Lambda = \Lambda(T) = \frac{dQ_2}{dP} = \Lambda_1 \Lambda_2 = e^{-T-2B(T)} e^{-T+N(T)\ln 2} = e^{-2T-2B(T)+N(T)\ln 2}$. Then $B(t) + 2t$ is a Q_2 -Brownian motion, and $N(t)$ is a Q -Poisson process with rate 2. In order to see this $E_Q(X) = E_P(\Lambda X) = E_P(\Lambda_1 \Lambda_2 X)$, for any X and $E_P \Lambda_1 = E_P \Lambda_2 = 1$. So under Q_2 , $N(t) - 2t$ is a martingale, and $B(t) + 2t$ is a martingale, thus $X(t) = B(t) + N(t)$ is a Q_2 -martingale.
- The probability measures defined as follows for $a > 0$ are equivalent to P , $dQ_a = e^{-a^2 T/2 - aB(T)} e^{(1-a)T + N(T)\ln a} = e^{(1-a-a^2/2)T - aB(T) + N(T)\ln a} dP$, and make the processes $B(t) + at$ into a Q_a -Brownian motion, and $N(t)$ into a Q_a -Poisson process with rate a .

Exercises to Chapter 11

Exercise 11.3: If $X = aS_1 + b$ for some a and b then $E_Q(X/r) = aE_Q(S_1/r) + b = aS_0 + b$ the same for any Q . Note that since the vectors S and β are not collinear, the representation of X by a portfolio (a, b) is unique. Take the claim that pays \$1 when the stock goes up and nothing in any other case, then this claim is unattainable. $E_Q(X) = 1.5(0.2 + p_d)$ depends on the choice of p_d .

Exercise 11.5:

- $E_Q(M(t)|\mathcal{F}_s) = E_P(M(t)|\mathcal{F}_s) = M(s)$ almost surely. Take $s = 0$, $E_P M(t) = E_Q M(t) = M(0)$.
- If a claim is attainable, $X = V(T)$, where $V(t)/\beta(t)$ is a Q -martingale. Its price at time t is $C(t) = V(t)$, which does not depend on the choice of Q .

According to the martingale property $V(t) = \beta(t)\mathbb{E}_Q(V(T)/\beta(T)|\mathcal{F}_t)$. This shows that $C(t) = \beta(t)\mathbb{E}_Q(X/\beta(T)|\mathcal{F}_t)$ is the same for all Qs.

Exercise 11.10: Consider continuous time. $d(\frac{S(t)}{\beta(t)}) = \frac{dS(t)}{\beta(t)} + S(t)d(\frac{1}{\beta(t)})$. $dR(t) = \frac{dS(t)}{S(t)} = \frac{\beta(t)}{S(t)}d(\frac{S(t)}{\beta(t)}) - \beta(t)d(\frac{1}{\beta(t)})$. $R(t) = \int_0^t \frac{\beta(u)}{S(u)}d(\frac{S(u)}{\beta(u)}) + \int_0^t \frac{d\beta(s)}{\beta(s)}$. The first integral is a martingale, as a stochastic integral with respect to a Q-martingale. Thus $\mathbb{E}_Q R(t) = \int_0^t \frac{d\beta(s)}{\beta(s)}$, which is the risk-free return. In discrete time the statement easily follows from the martingale property of $S(t)/\beta(t)$.

Exercise 11.11: Let $X(t) = S(t)e^{-rt}$. Then it is easy to see that $X(t)$ satisfies the SDE $dX(t) = \sigma X(t)dB(t)$, for a Q-Brownian motion $B(t)$. According to Itô's formula $d(\frac{1}{X(t)}) = -\sigma \frac{1}{X(t)}(dB(t) - \sigma dt) = -\sigma \frac{1}{X(t)}dW(t)$, with $dW(t) = dB(t) - \sigma dt$. $W(t)$ is a Q-Brownian motion with drift. Using the change of measure $dQ_1/dQ = e^{\sigma B(T) - \sigma^2 T/2}$, by Girsanov's theorem $W(t)$ is a Q_1 -Brownian motion. Finally, $-W(t)$ is also a Q_1 -Brownian motion, and we can remove the minus sign in the SDE.

Exercise 11.12: According to Theorem 11.17, $C(t) = \mathbb{E}_{Q_1}(\frac{S(t)}{S(T)}(S_T - K)^+|\mathcal{F}_t) = S_t \mathbb{E}_{Q_1}((1 - \frac{K}{S(T)})^+|\mathcal{F}_t)$. The conditional distribution of $1/S(T)$ given \mathcal{F}_t under Q_1 is obtained by using the SDE for $1/S(t)$. With $Y(t) = 1/X(t) = e^{rt}/S(t)$, we have from above $dY(t) = \sigma Y(t)dW(t)$ and $1/S(t) = Y(t)e^{-rt}$. Using the product rule $d(\frac{1}{S(t)}) = \frac{1}{S(t)}(-rdt + \sigma dW(t))$. Thus $1/S(t)$ is a stochastic exponential, and $1/S(T) = (1/S(t))e^{(T-t)(-r-\sigma^2/2)+\sigma(W(T)-W(t))}$. Thus, given \mathcal{F}_t , the Q_1 conditional distribution of $1/S(T)$ is lognormal with mean and variance $-\ln S(t) - (T-t)(r + \sigma^2/2)$ and $\sigma^2(T-t)$. Using calculations for the $E(1-X)^+$ similar to p. 313, we recover the Black–Scholes formula.

Exercise 11.14: From the first equation $b(t) = (V(t) - a(t)S(t))e^{-rt}$. Putting it in the self-financing condition $dV(t) = a(t)dS(t) + b(t)d(e^{rt})$ we get the SDE for $V(t)$.

For the other direction, let $b(t)$ be as above. Then $V(t) = a(t)S(t) + b(t)e^{rt}$, moreover, the SDE for $V(t)$ gives the self-financing condition.

Exercise 11.17: Let $F(y) = e^{-rT}E((S_T/S - y)^+)$, then $C = SF(K/S)$. Now, $\partial C/\partial S = F(K/S) - KF'(K/S)/S$ and $\partial C/\partial K = F'(K/S)$. Thus $S\partial C/\partial S + K\partial C/\partial K = SF(K/S) = C$. The expression for $\frac{\partial C}{\partial S}$ in the Black–Scholes model follows from the Black–Scholes formula.

Exercises to Chapter 12

Exercise 12.1: Use Theorem 12.5, $C(t) = E_Q(\frac{\beta(t)}{\beta(s)}(P(s, T) - K)^+ | \mathcal{F}_t) = E_Q(\frac{\beta(t)}{\beta(s)}P(s, T)I(P(s, T) > K) | \mathcal{F}_t) - KE_Q(\frac{\beta(t)}{\beta(s)}I(P(s, T) > K) | \mathcal{F}_t)$. For the first term take Λ_2 based on the Q-martingale $P(s, T)/\beta(s)$, $s \leq T$, which corresponds to numeraire $P(t, T)$, or T -forward measure. Since the expectation must be unity, $\Lambda_2 = \frac{P(s, T)}{\beta(s)P(0, T)}$ (Formula (12.53)). Then the first term is $E_Q(\frac{P(s, T)\beta(t)}{\beta(s)}X | \mathcal{F}_t) = P(t, T)E_{Q_2}(X | \mathcal{F}_t)$. For the second term consider $\Lambda_1 = 1/(P(0, s)\beta(s))$ based on the martingale $P(t, s)/\beta(t)$, $t \leq s$, which corresponds to numeraire $P(t, s)$, or s -forward measure. Then for any X , $E_Q(\frac{\beta_t}{\beta_s}X | \mathcal{F}_t) = P(t, s)E_{Q_1}(X | \mathcal{F}_t)$. This gives for the second term $KP(t, s)Q_1(P(s, T) > K | \mathcal{F}_t)$, and the result follows.

Exercise 12.3:

1. Follows by $E|X| \leq \sqrt{EX^2}$.
2. By additivity of the integral $\int_0^T (\sum_{i=0}^{n-1} H(t_i, s)(W(t_{i+1}) - W(t_i)))ds = \sum_{i=0}^{n-1} (\int_0^T H(t_i, s)ds)(W(t_{i+1}) - W(t_i))$.
3. Let $X_n(s) = \sum_{i=0}^{n-1} H(t_i, s)(W(t_{i+1}) - W(t_i))$. Then $X_n(s)$ is an approximation to the Itô integral $X(s) = \int_0^T H(t, s)dW(t)$. $EX^2(s) = \int_0^T EH^2(t, s)dt < \infty$, and under the stated conditions $\int_0^T E(X_n(s) - X(s))^2 ds$ converges to zero. This implies convergence in L^2 and in probability of $\int_0^T X_n(s)ds$ to $\int_0^T X(s)ds$. The rhs $\sum_{i=0}^{n-1} (\int_0^T H(t_i, s)ds)(W(t_{i+1}) - W(t_i))$ is an approximation of the Itô integral of $Y(t)$, and convergence to it in probability.

Exercise 12.5: Use the formula for the bond (12.32). $f(t, T) = -\frac{\partial \ln P(t, T)}{\partial T}$. A cap is a sum of caplets, which are priced by (12.64).

Exercise 12.7: Let $Y(t) = \int_0^t \gamma(s)dB(s)$ and $X(t) = \int_0^t \beta(s)dB(s)$ for some $\beta(s)$ to be determined. Then $\mathcal{E}(X)(t) = c + (1-c)\mathcal{E}(Y)(t)$. Thus

$$dX(t) = \frac{d\mathcal{E}(X)(t)}{\mathcal{E}(X)(t)} = \frac{(1-c)d\mathcal{E}(Y)(t)}{c + (1-c)\mathcal{E}(Y)(t)} = \frac{(1-c)\mathcal{E}(Y)(t)\gamma(t)}{c + (1-c)\mathcal{E}(Y)(t)} dB(t), \text{ and}$$

$$\beta(t) = \frac{(1-c)\mathcal{E}(Y)(t)\gamma(t)}{c + (1-c)\mathcal{E}(Y)(t)} = \frac{(1-c)\gamma(t)e^{\int_0^t \gamma(s)dB(s) - \frac{1}{2}\int_0^t \gamma^2(s)ds}}{c + (1-c)e^{\int_0^t \gamma(s)dB(s) - \frac{1}{2}\int_0^t \gamma^2(s)ds}}.$$

The existence of forward rates volatilities follows from Equation (12.76).

Exercise 12.9: At T_i the following exchange is made: the amount received is $f_{i-1}(T_i - T_{i-1})$ and paid out $k(T_i - T_{i-1})$. The resulting amount at time T_i is $1/P(T_{i-1}, T_i) - 1 - k\delta$, using $1/P(T_{i-1}, T_i) = 1 + f_{i-1}(T_i - T_{i-1})$. Thus the value at time t of the swap is

$$\text{Swap}(t, T_0, k) = \sum_{i=1}^n \mathbb{E}_Q \left(\frac{\beta(t)}{\beta(T_i)} \left(\frac{1}{P(T_{i-1}, T_i)} - 1 - k\delta \right) \mid \mathcal{F}_t \right).$$

Using the martingale property of the discounted bonds (12.4) we obtain that $\mathbb{E}_Q \left(\frac{\beta(t)}{\beta(T_i)} \mid \mathcal{F}_{T_{i-1}} \right) = P(T_{i-1}, T_i)$. The result is obtained by conditioning on $\mathcal{F}_{T_{i-1}}$ in the above sum.

Exercise 12.10: Follows from the previous exercise.

Exercise 12.11: Consider the portfolio of bonds that at any time $t < T$ is long $\alpha_+(t)$ of the T_0 -bond, short $\alpha_-(t)$ of the T_n -bond, and for each i , $1 \leq i \leq n$ short $\alpha_-(t)\delta k$ of the T_i -bond. The value of this portfolio at time $t < T$ is $C(t) = \alpha_+(t)(P(t, T_0) - P(t, T_n)) - k\alpha_-(t)b(t)$, with $b(t) = \delta \sum_{i=1}^n P(t, T_i)$. By using the expression for the swap rate $C(t) = b(t)(\alpha_+(t)k(t) - k\alpha_-(t))$. It can be seen that this portfolio has the correct final value and is self-financing, i.e. $dC(t) = \alpha_+(t)d(P(t, T_0) - P(t, T_n)) - k\alpha_-(t)db(t)$.

Exercises to Chapter 13

Exercise 13.1: Use Theorem 6.16 and formula (6.98) to find $E(T_0 \wedge T_b)$. $ET_0 = \lim_{b \rightarrow \infty} E(T_0 \wedge T_b)$ by monotone convergence.

Exercise 13.2:

1. $d(e^{-cX(t)}) = -ce^{-cX(t)}dX(t) + \frac{c^2}{2}e^{-cX(t)}d[X, X](t) = -ce^{-cX(t)}\sigma\sqrt{X}(t)dB(t)$. According to Theorem 13.1 $X(t) \geq 0$ for all t . The function $e^{-cx}\sqrt{x}$ is bounded for $x \geq 0$. Thus $e^{-cX(t)}$ is a martingale as an Itô integral of a bounded process, Theorem 4.7.
2. Let $\tau = T \wedge t$, then τ is a bounded stopping time. Applying optional stopping we have $e^{-cx} = \mathbb{E}e^{-cX(0)} = \mathbb{E}e^{-cX(\tau)} = \mathbb{E}(e^{-cX(\tau)}I(\tau = T)) + \mathbb{E}(e^{-cX(\tau)}I(\tau = t)) \geq \mathbb{E}(e^{-cX(\tau)}I(\tau = T)) = \mathbb{P}(T \leq t)$.

Exercise 13.5:

1. Let $G(x) = \int_a^x du/g(u)$. Then $G(x(t)) = G(x(0)) + t$. Now $dY(t) = dt + \frac{\sigma(X(t))}{g(X(t))}dB(t) - \frac{\sigma^2(X(t))g'(X(t))}{g^2(X(t))}dt$.

2. $G(x) = x^{1-r}/(c(1-r))$ and $Y(t) = t - \int_0^t \frac{\sigma^2(X(s))r}{c(X(s))^{r+1}} ds + \int_0^t \frac{\sigma(X(s))}{c(X(s))^r} dB(s)$. According to the growth condition $\mathbb{E}(\int_0^t \frac{\sigma(X(s))}{cX(s))^r} dB(s))^2 = \int_0^t \mathbb{E}(\frac{\sigma(X(s))}{cX(s))^r})^2 ds \leq Ct$. It follows $\mathbb{E}(\frac{1}{t} \int_0^t \frac{\sigma(X(s))}{cX(s))^r} dB(s))^2 \rightarrow 0$, and implies $\frac{1}{t} \int_0^t \frac{\sigma(X(s))}{cX(s))^r} dB(s) \rightarrow 0$ in probability. It follows by the l'Hôpital rule that $\frac{1}{t} \int_0^t \frac{\sigma^2(X(s))r}{c(X(s))^{r+1}} ds \rightarrow 0$ on the set $\{X(t) \rightarrow \infty\}$. The law of large numbers (LLN) for $Y(t)$ now follows.

Exercise 13.6: $P(\min(L_1, L_2, \dots, L_x) > t) = P(L_1 > t, L_2 > t, \dots, L_x > t) = (P(L_1 > t))^x = (e^{-a(x)t})^x = e^{-xa(x)t}$.

Exercise 13.9:

1. $\pi_1(t) = \int_0^t \sum_{k \geq 1} I(X(s-) \geq k) dN_k^\lambda(s)$ has jumps of size one. Verify that $A_t^1 = \int_0^t \lambda X_s ds$ is its compensator. Similarly for the process $\pi_2(t) = \int_0^t \sum_{k \geq 1} I(X(s-) \geq k) dN_k^\mu(s)$.
2. $X(t) = X(0) + A_1(t) - A_2(t) + \pi_1(t) - A_1(t) - \pi_2(t) + A_2(t)$.

Exercises to Chapter 14

Exercise 14.3: $\mathbb{E}(M(t)|\mathcal{G}_s) = \mathbb{E}(\mathbb{E}(M(t)|\mathcal{F}_s)|\mathcal{G}_s) = \mathbb{E}(M(s)|\mathcal{G}_s) = M(s)$.

Exercise 14.5: Let \mathcal{G}_t be the trivial σ -field, then $\widehat{W}(t) = \mathbb{E}(W(t)|\mathcal{G}_t) = 0$.

Exercise 14.6: Formally using Theorem 14.6 with $a(t) = b(t) = 0$ and $A(t) = B(t) = 1$, $d\widehat{X}(t) = -v(t)\widehat{X}(t)dt + v(t)dY(t)$, and $dv(t) = -v^2(t)dt$. Hence $d\widehat{X}(t) = -(1/t)\widehat{X}(t)dt + (1/t)dY(t)$. We obtain $v(t) = 1/t$ and $\widehat{X}(t) = Y(t)/t$. These can be obtained directly using $Y(t) = ct + W(t)$.

Exercise 14.8: $dX(t) = (\mu + \sigma^2/2)X(t)dt + \sigma X(t)dW(t)$. Apply Theorem 14.6 with $a(t) = \mu + \sigma^2/2$, $b(t) = \sigma$, $A(t) = B(t) = 1$, see Example 14.1.

This page intentionally left blank

References

- Abramov, V., Klebaner, F.C. and Liptser, R.S. (2011). The Euler-Maruyama approximations for the CEV model, *Discrete and Continuous Dynamical Systems B*, **16**, 1–14.
- Armitage, P. and Doll, R. (1954). The age distribution of cancer and a multi-stage theory of carcinogenesis, *British J. of Cancer*, **8**, 1–12.
- Armitage, P. and Doll, R. (1957). A two-stage theory of carcinogenesis in relation to the age distribution of human cancer, *British J. of Cancer*, **11**, 161–169.
- Artzner, P. and Delbaen, F. (1989). Term structure of interest rates: The martingale approach, *Adv. Appl. Math.*, **10**, 95–129.
- Athreya, K.B. and Key, P.E. (1972). *Branching Processes*, Springer, New York.
- Bachelier, L. (1990). *Théorie de la Spéculation*, Gauthier–Villars, Paris.
- Bezen, A. and Klebaner, F.C. (1996). Stationary solutions and stability of second order random differential equations, *Physica A*, **233/234**, 809–823.
- Bhattacharya, R.N. (1978). Criteria for recurrence and existence of invariant measures for multi-dimensional diffusions, *Ann. Probab.*, **6**, 541–553.
- Björk, T. (1998). *Arbitrage Pricing Theory in Continuous Time*, Oxford University Press, Oxford.
- Björk, T., Di Masi, G., Kabanov, Y. and Runggalidier, W. (1997). Toward a general theory of bond markets, *Finance and Stochastics*, **2**, 141–174.
- Björk, T., Kabanov, Y. and Rungaldier, W. (1996). Bond market structure in the presence of marked point processes, *Math. Finance*, **7**, 211–239.
- Black, F. (1976). The pricing of commodity contracts, *J. Finan. Econ.* **3**, 167–179.
- Borodin, A.N. and Salminen, P. (1996). *Handbook of Brownian Motion*, Birkhäuser, Basel.
- Borovkov, K., Klebaner, F.C. and Virág, E. (2003). Random step functions model for interest rates, *Finance and Stochastics*, **7**, 123–143.
- Brace, A. (1998). Swaprate Volatilities in BGM, FMMA notes.
- Brace, A., Gatarek, D. and Musiela, M. (1997). The market model of interest rate dynamics, *Math. Finance*, **7**(2), 127–156.
- Breiman, L. (1968). *Probability*, Addison-Wesley, Reading, MA.

- Brigo, D. and Mercurio, F. (2006). *Interest Rate Models: Theory and Practice*, 2nd ed., Springer-Verlag, Berlin.
- Brown, R. (1828). A brief account of microscopical observations made in the months of June, July and August 1827, on the particles contained in the pollen of plants, and on the general existence of active molecules in organic and inorganic bodies, *The Philosophical Magazine*, **4**, 161–173, London: R. Taylor.
- Chung, K.L. (1960). *Markov Chains with Stationary Transition Probabilities*, Springer-Verlag, Berlin.
- Chung, K.L. (1982). *Lectures from Markov Processes to Brownian Motion*, Springer-Verlag, Berlin.
- Chung, K.L. and Williams, R.J. (1983). *Introduction to Stochastic Integration*, Birkhäuser, Basel.
- Cont, R. (1998). Modeling term structure dynamics: An infinite dimensional approach, *Lyapunov Institute Workshop on Mathematical Finance*, Rocquencourt.
- Dawson, D.A. (1993). *Measure-Valued Markov Processes*, Lecture Notes in Math., Springer-Verlag, Berlin.
- Delbaen, F. and Schachermayer, W. (1994). A general version of the fundamental theorem of asset pricing, *Math. Annalen*, **300**, 463–520.
- Dellacherie, C. and Meyer, P.A. (1982). *Probabilities and Potential B*, North-Holland, Amsterdam.
- Dieudonné, J. (1960). *Foundations of Modern Analysis*, Academic Press, New York.
- Dudley, R.M. (1989). *Real Analysis and Probability*, Wadsworth, Belmont.
- Durrett, R. (1984). *Brownian Motion and Martingales in Analysis*, Wadsworth, Belmont.
- Durrett, R. (1991). *Probability: Theory and Example*, Wadsworth, Belmont.
- Durrett, R., Schmidt, D. and Schweinsberg, J. (2009). A waiting time problem arising from the study of multistage carcinogenesis, *Ann. Appl. Probab.*, **19**, 676–718.
- Dynkin, E.B. (1965). *Markov Processes*, Springer-Verlag, Berlin.
- Einstein, A. (1905). Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in rehenden Flüssigkeiten suspendierten Teilchen, *Annalen der Physik*, **322**(8), 549–560.
- Ethier, S.N. and Kurtz, T.G. (1986). *Markov Processes*, Wiley, New York.
- Ewens, W.J. (1979). *Mathematical Population genetics*, Springer-Verlag, Berlin.
- Feller, W. (1951). Diffusion processes in genetics, *Second Berkeley Symposium on Math. Stat. and Probab.*, 227–246.
- Feller, W. (1971). *An Introduction to Probability Theory and Its Applications*, Vol. 2, Wiley, New York.
- Fisher, J.C. (1958). Multiple mutation theory of carcinogenesis, *Nature*, **181**, 651–652.
- Fouque, J.-P., Papanicolaou, G. and Sircar, K.R. (2000). *Derivatives in Financial Markets with Stochastic Volatility*, Cambridge University Press, Cambridge.
- Freedman, D. (1971). *Brownian Motion and Diffusion*, Springer-Verlag, Berlin.

- Freedman, D. (1983). *Brownian Motion and Diffusion*, Springer-Verlag, Berlin.
- Friedman, A. (1975). *Stochastic Differential Equations and Applications*, Vol. 1, Academic Press, New York.
- Friedman, A. (1976). *Stochastic Differential Equations and Applications*, Vol. 2, Academic Press, New York.
- Garay, R.P. and Lefever, R. (1978). A kinetic approach to the immunology of cancer. Stationary state properties of effector-target cell reactions, *J. Theor. Biol.*, **73**, 417–438.
- Gard, T.C. (1988). *Introduction to Stochastic Differential Equations*, Marcel Dekker, New York.
- Garman, M.B. and Kohlhagen S.W. (1983). Foreign currency option values, *J. Int. Money Finance*, **2**, 231–237.
- Geman, H., El Karoui, N. and Rochet, J.C. (1995). Changes of numeraire, changes of probability measure and option pricing, *J. Appl. Probab.*, **32**, 443–458.
- Gihman, I.I. and Skorohod, A.V. (1972). *Stochastic Differential Equations*, Springer-Verlag, Berlin.
- Grimmet, G. and Stirzaker, D. (1987). *Probability and Random Processes*, Oxford University Press, New York.
- Haeno, H., Iwasa, Y. and Michor, F. (2007). The evolution of two mutations during clonal expansion, *Genetics*, **177**, 2209–2221.
- Hamza, K., Jacka, S.D. and Klebaner, F.C. (2005). The EMM conditions in a general model for interest rates, *Adv. Appl. Probab.*, **37**, 415–434.
- Hamza, K. and Klebaner, F.C. (1995a). Representation of Markov chains as stochastic differential equations, *Lecture Notes in Statistics*, **99**, 144–151.
- Hamza, K. and Klebaner, F.C. (1995b). Conditions for integrability of Markov Chains, *J. Appl. Probab.*, **32**, 541–547.
- Hamza, K. and Klebaner, F.C. (Forthcoming 2013). *Problems and Solutions in Stochastic Calculus*, Imperial College Press, London.
- Harris, T.E. (1963). *The Theory of Branching Processes*, Springer-Verlag, Berlin.
- Harrison, J.M. and Kreps, D. (1979). Martingales and arbitrage in multiperiod security markets, *J. Econ. Theory*, **20**, 381–408.
- Harrison, J.M. and Pliska, S.R. (1981). Martingales and stochastic integrals in the theory of continuous trading, *Stoch. Proc. Appl.*, **11**, 215–260.
- Harrison, J.M. and Pliska, S.R. (1983). A stochastic model of continuous trading: Complete markets, *Stoch. Proc. Appl.*, **15**, 313–316.
- Hasminskii, R.Z. (1980). *Stochastic Stability of Differential Equations*, Sijthoff and Noordhoff, Alphen aan den Rijn.
- Heath, D., Jarrow, R. and Morton, A. (1992). Bond pricing and the term structure of interest rates: A new methodology for contingent claims valuation, *Econometrica*, **60**, 77–105.
- Heston, S. (1993). A closed-form solution for options with stochastic volatility with applications to bond and currency options, *Review of Financial Studies*, **6**(2), 327–343.
- Hobson, E.W. (1921). *The Theory of Functions of a Real Variable*, Cambridge University Press, Cambridge.

- Ito, K. and McKean, H.P. (1965). *Diffusion Processes and Their Sample Paths*, Springer-Verlag, Berlin.
- Jacob, J. and Shiryaev, A.N. (1987). *Limit Theorems for Stochastic Processes*, Springer-Verlag, Berlin.
- Jagers, P. and Klebaner, F.C. (2000). Population-size-dependent and age-dependent branching processes, *Stoch. Proc. Appl.*, **87**, 235–254.
- Jamshidian, F. (1996). Sorting out swaptions, *Risk*, **9**(3), 59–60.
- Kabanov, Y.M. (2001). Arbitrage Theory, in Jouini, E., Cvitanic, J. and Musiela, M. eds., *Option Pricing, Interest Rates and Risk Management*, Cambridge University Press, Cambridge, pp. 3–42.
- Kallenberg, O. (1996). On the existence of universal functional solutions to classical SDEs, *Ann. Probab.*, **24**, 196–205.
- Kallianpur, G. (1980). *Stochastic Filtering Theory*, Springer-Verlag, Berlin.
- Kallsen, J. and Shiryaev, A.N. (2002). The cumulant process and Escher's change of measure, *Finance and Stochastics*, **6**, 397–428.
- Karatzas, I. and Shreve, S.E. (1988). *Brownian Motion and Stochastic Calculus*, Springer-Verlag, London.
- Karlin, S. and Taylor, H.M. (1975). *A First Course in Stochastic Processes*, Academic Press, New York.
- Karlin, S. and Taylor, H.M. (1981). *A Second Course in Stochastic Processes*, Academic Press, New York.
- Karr, A.F. (1986). *Point Processes and Their Statistical Inference*, Marcel Dekker, New York.
- Keller, G., Kersting, G. and Rösler, U. (1984). On the asymptotic behaviour of solutions of stochastic differential equations, *Z. Wahrscheinlichkeitsth.*, **68**, 163–189.
- Kennedy, P.D. (1994). The term structure of interest rates as a Gaussian Markov field, *Math. Finance*, **4**, 247–258.
- Kersting, G. and Klebaner, F.C. (1995). Sharp conditions for nonexplosions and explosions in Markov jump processes, *Ann. Probab.*, **23**, 268–272.
- Kersting, G. and Klebaner, F.C. (1996). Explosions in Markov jump processes and submartingale convergence, *Lecture Notes in Statistics* **114**, 127–136, Springer-Verlag, Berlin.
- Klebaner, F.C. (1994). Asymptotic behaviour of Markov population processes with asymptotically linear rate of change, *J. Appl. Probab.*, **31**, 614–625.
- Klebaner, F.C. (2002). Option price when the stock is a semimartingale, *Elect. Comm. in Probab.*, **7**, 79–83. Correction, **8**.
- Klebaner, F.C. and Azmy, E. (2010). Solutions and simulations of some one-dimensional stochastic differential equations, *Asia Pacific Finan. Markets*, **17**, 365–372.
- Klebaner, F.C. and Liptser, R. (2001). Asymptotic analysis and extinction in randomly perturbed Lotka–Volterra model, *Ann. Appl. Probab.*, **11**(4), 1263–1291.
- Knudson, A.G.J. (1971). Mutation and cancer: Statistical study of retinoblastoma, *Proc. Natl. Acad. Sci. USA*, **68**(4), 820–823.

- Komarova, N.L. (2007). Loss- and gain-of-function mutations in cancer: Mass action, spatial and hierarchical models. *J. Statist. Physics*, **128**, 413–446.
- Kreps, D.M. (1981). Arbitrage and equilibrium in economies with infinitely many commodities, *J. Math. Economics*, **8**, 15–35.
- Krishnan, V. (1984). *Nonlinear Filtering and Smoothing: An Introduction to Martingales, Stochastic Integrals, and Estimation*, Wiley, New York.
- Kurtz, T.G. (1981). *Approximation of Population Processes*, Society for Industrial and Applied Mathematics, Philadelphia.
- Lamberton, D. and Lapeyre, B. (1996). *Introduction to Stochastic Calculus Applied to Finance*, Chapman and Hall, London.
- Liptser, R.S. and Shiryaev, A.N. (1974). *Statistics of Random Processes I and II*, Springer-Verlag, Berlin.
- Liptser, R.S. and Shiryaev, A.N. (1989). *Theory of Martingales*, Kluwer, Dordrecht.
- Liptser, R.S. and Shiryaev, A.N. (2001). *Statistics of Random Processes I and II, 2nd ed.*, Springer-Verlag, Berlin.
- Loeve, M. (1978). *Probability Theory I and II*, Springer-Verlag, Berlin.
- Lotka, A.J. (1925). *Elements of Physical Biology*, Williams and Wilkins, Baltimore.
- Métivier, M. (1982). *Semimartingales*, de Gruyter, Berlin.
- Meyer, P.A. (1966). *Probability and Potentials*, Baidell Publishing Co., Waltham.
- Michor, F., Nowak, M.A. and Iwasa, Y. (2006). Stochastic dynamics of metastasis formation, *J. Theor. Biol.*, **240**, 521–530.
- Michor, F. and Iwasa, Y. (2006). Dynamics of metastasis suppressor gene inactivation, *J. Theor. Biol.*, **241**, 676–689.
- Molchanov, S.A. (1975). Diffusion processes and Riemannian geometry, *Russian Math. Surveys*, **30**, 1–63.
- Moolgavkar, S.H. and Knudson, A.G.J. (1981). Mutation and cancer: A model for human carcinogenesis, *J. Nat. Cancer Inst.*, **66**, 1037–1052.
- Moran, P.A.P. (1958). Random processes in genetics, *Proc. Cambridge Philos. Soc.*, **54**, 60–71.
- Musiela, M. and Rutkowski, M. (1998). *Martingale Methods in Financial Modelling*, Springer, New York.
- Nordling, C.O. (1953). A new theory of cancer-inducing mechanism, *Brit. J. Cancer*, **7**, 68–72.
- Ochab-Marcinek, A. and Gudowska-Nowak, E. (2006). Coexistence of resonant activation and noise enhanced stability in a model of tumor–host interaction: Statistics of extinction times, *Acta Physica Polonica B*, **37**, 1651–1666.
- Oksendal, B. (1995). *Stochastic Differential Equations*, Springer-Verlag, Heidelberg, New York.
- Paley, R.E.A.C., Wiener, N. and Zygmund, A. (1933). Notes on random functions, *Math. Z.*, **37**, 647–668.
- Pinsky, R. (1995). *Positive Harmonic Functions and Diffusion*, Cambridge University Press, Cambridge.
- Polya, G. (1921). Über eine Aufgabe der Wahrscheinlichkeitsrechnung betreffend die Irrfahrt im Strassennetz, *Math. Ann.*, **84**, 149–160.

- Protter, P. (1992). *Stochastic Integration and Differential Equations*, Springer-Verlag, Berlin.
- Rebonato, R. (2002). *Modern Pricing of Interest-Rate Derivatives*, Princeton University Press, Princeton.
- Revuz, D. and Yor, M. (1991). *Continuous Martingales and Brownian Motion*, Springer-Verlag, Berlin.
- Revuz, D. and Yor, M. (2001). *Continuous Martingales and Brownian Motion*, 3rd ed., Springer-Verlag, Berlin.
- Rogers, L.C.G. and Shi, Z. (1995). The value of an Asian option, *J. Appl. Probab.*, **32**, 1077–1088.
- Rogers, L.C.G. and Williams, D. (1987). *Diffusions, Markov Processes, and Martingales, Vol. 2, Ito Calculus*, Wiley, New York.
- Rogers, L.C.G. and Williams, D. (1990). *Diffusions, Markov Processes, and Martingales, Vol. 2, Ito Calculus*, reprint, Wiley, New York.
- Rogers, L.C.G. and Williams, D. (1994). *Diffusions, Markov Processes, and Martingales, Vol. 1, Foundations*, Wiley, New York.
- Rosenkranz, G. (1985). Growth models with stochastic differential equations. An example from tumor immunology, *Math. Biosci.*, **75**, 175–186.
- Saks, S. (1964). *Theory of the Integral*, Dover, New York.
- Shirakawa, H. (1991). Interest rate option pricing with Poission–Gaussian forward rates curve processes. *Math. Finance*, **1**, 77–94.
- Shiryayev, A.N. (1999). *Essentials of Stochastic Finance*, World Scientific, Singapore.
- Soize, C. (1994). *The Fokker–Planck Equation for Stochastic Dynamical Systems and its Explicit Steady State Solutions*, World Scientific, Singapore.
- Soong, T.T. (1973). *Random Differential Equations in Science and Engineering*, Academic Press, New York.
- Stroock, D. and Varadhan, S.R.S. (1979). *Multidimensional Diffusion Processes*, Springer-Verlag, Berlin.
- Tan, W.Y. and Hanin, L. (2008). *Handbook of Cancer Models with Applications*, World Scientific, Singapore.
- Vasicek, O. (1977). An equilibrium characterization of the term structure, *J. Fin. Econ.*, **5**, 177–188.
- Vecer, J. (2002). Unified Asian pricing, *Risk*, **15**(6), 113–116.
- Volterra, V. (1926). Variazioni e fluttuazioni del numero d'individui in specie d'animali conviventi. *Mem. Acad. Lincei*, **2**, 31–113.
- Weierstrass, K. (1895). Über continuirliche Functionen eines reellen Arguments, die für keinen Werth des letzteren einen bestimmten Differentialquotienten besitzen, in *Königlich Preussischen Akademie der Wissenschaften, Mathematische Werke von Karl Weierstrass*, vol. 2, Mayer & Mueller, Berline, 71–74.
- Widder, D.V. (1944). Positive temperatures on an infinite rod, *Trans. Amer. Math. Soc.*, **55**, 85–95.
- Wiener, N. (1923). Differential space, *Journal Math. Phys.*, **2**, 131–174.
- Yamada, T. and Watanabe, S. (1971). On the uniqueness of solutions of stochastic differential equations, *J. Math. Kyoto Univ.*, **11**, 155–167.

Index

- absorbtion probabilities, 366
adapted process, 50, 243
admissible strategy, 295, 302
arbitrage, 291
arcsine law, 75
Asian options, 318
attainable claim, 295, 298
- backward equation, 180
barrier options, 320
Bayes formula, 276
BDG inequality, 203
Bessel process, 178
best estimator, 47
BGM model, 343
binomial model, 293
birth–death process, 359, 375
bivariate normal, 42
Black’s formula, 345, 350
Black–Scholes, 301, 305
Black–Scholes PDE, 307, 328
bond, 325
Borel σ -field, 29
bracket process, 220
branching diffusion, 353
branching process, 379
Brownian bridge, 133, 207
Brownian motion, 56, 63, 75, 133,
 160, 168, 175, 184, 280, 285
càdlàg, 4, 213
call option, 289
- cancer models, 377
cap, 350
caplet, 341, 342, 345
chain rule, 14
change of drift, 280
change of measure, 269, 273, 281
change of numeraire, 312
change of time, 207
change of variables, 13, 222
Chapman–Kolmogorov equation, 136
characteristic function, 38
compensator, 230, 231, 247, 259,
 262, 382
complete market model, 298, 304
conditional distribution, 43
conditional expectation, 27, 43
continuity of paths, 48
continuous, 1
continuous martingales, 205
convergence almost surely, 38
convergence in L^r , 38
convergence in distribution, 38
convergence in probability, 38
convergence of expectations, 39
convergence of random variables, 38
correlated Brownian motions, 89,
 120, 312
counting process, 251, 254
covariance, 40, 103, 114, 121
cross-currency options, 317
currency options, 315

- Davis' inequality, 203
 decomposition of FV functions, 36
 diffusion, 304, 358
 diffusion coefficient, 116, 126
 diffusion operator, 144
 Dirichlet class (D), 200
 discrete time risk model, 212
 distribution, 26, 31
 dominated convergence, 46
 Doob's decomposition, 216, 247
 Doob's inequality, 203
 Doob–Levy martingale, 51, 397
 Doob–Meyer decomposition, 216
 Doss–Sussman method, 129
 drift, 305
 drift coefficient, 126
 dual predictable projection, 231
 Dynkin's formula, 151, 265, 361, 380
- EMM assumption, 333
 ergodic, 182
 evanescent, 48
 exit time, 69, 163, 164, 367
 exotic options, 290
 expectation, 26, 33
 explosion, 168, 267, 389
 extinction, 366
 extinction probability, 356
- Fatou's lemma, 39, 46, 368
 Feller property, 170
 Feynman–Kac formula, 157, 328
 field, 22, 24
 filtered probability space, 21, 50
 filtering equation, 396
 filtration, 23, 24, 242
 finite variation, 4
 fluid approximation, 391
 Fokker–Planck equation, 145
 forward equation, 145, 362
 forward measure, 338
 forward rate, 333
 fractional Brownian motion, 121
 Fubini's theorem, 53, 242
 fundamental solution of PDE, 144
 FX options, 315
- gambler's ruin, 194
 Gaussian distribution, 41
 Gaussian process, 42, 103, 134, 148,
 332
 gene frequencies, 357
 generator, 160, 265, 267, 361, 380
 Gronwall's inequality, 18, 383
 growth conditions, 18
- Hölder condition, 17, 135
 harmonic function, 164, 180
 hedging, 303
 higher order SDE, 182
 hitting time, 69, 73, 195, 356
 HJM model, 333
 Ho–Lee model, 336
- implied volatility, 309
 increment, 1
 indicator, 30
 integration by parts, 12, 113, 222, 254
 invariant distribution, 172, 362
 invariant measure, 405
 isometry, 94, 97, 218, 236
 Itô integral, 91, 93, 95, 103, 111, 204
 Itô's formula, 109, 112, 116, 178, 222,
 236, 237, 253
- joint distribution, 32
 Jordan decomposition, 6
 jump, 3
- Kalman–Bucy filter, 399
- Lamperti's change of time, 210
 Langevin equation, 127, 379
 law of iterated logarithm, 79
 law of large numbers, 79
 Lebesgue decomposition, 36
 Lebesgue integral, 33, 34
 Lebesgue Measure, 30
 Lebesgue–Stieltjes Integral, 34
 left-continuous, 2
 Levy's characterization, 205
 LIBOR, 343
 likelihood, 285, 286

- linear rates, 363
linear SDE, 132, 404
Lipschitz conditions, 17
local martingale, 101, 197
local time, 224
localization, 198, 382
lookback options, 320
Lotka–Volterra model, 386
- marked point processes, 251
Markov process, 162, 265
Markov property, 67
martingale, 50, 65, 100, 180, 185
martingale exponential, 82, 90
martingale measure, 298
martingale representation, 239
martingale square integrable, 186
martingale transform, 84
martingales in random walks, 82
maximum of Brownian motion, 71
mean exit time, 164
measure of jumps, 246
Merton’s model, 329
Meyer–Tanaka formula, 225
minimum of Brownian motion, 73
moment generating function, 38, 41
moments, 37
monotone convergence, 46
Moran model, 369
mutation, 372
- natural scale, 166
non-linear filter, 396
normal distribution, 41
numeraire, 313
- occupation time, 224
option pricing formula, 313
optional, 243
optional stopping, 191
Ornstein–Uhlenbeck, 127, 174, 286
- PDE, 145
Poisson process, 86, 87, 258, 388
polarization identity, 9
population dependence, 365, 387
- positive recurrent, 182
predictable, 25, 83, 214, 230–232, 235, 236, 243, 246
predictable quadratic variation, 231
predictable representation, 239
prey–predator, 387
price, 303
pricing by PDE, 307
pricing equation, 304
probability, 26, 29
probability density, 32, 35
product rule, 113, 222
progressively measurable, 243
pure jump process, 260
put option, 290
- quadratic variation, 8, 63, 101, 109, 200, 202, 220, 221, 278, 308
Quanto options, 317
- Radon–Nikodym derivative, 273
random measure, 246
random ODE, 128
random oscillator, 405
random variable, 23, 26, 30
random walk, 81
recurrence, 74, 81
reflection principle, 74
regular functions, 4, 213
renewal process, 257
Riemann integral, 9
right-continuous, 2, 49
ruin probability, 195, 212
- scale function, 166
SDE, 126, 176, 359
self-financing portfolio, 295, 300
semimartingale, 213, 390
semimartingale exponential, 226
sharp bracket, 231, 235, 237, 381
sigma-field, 28, 31, 49
simple processes, 91, 217
simulation, 272
simulation birth–death processes, 375
space-homogeneous, 58
spot rate, 328–330

- spread, 322
square bracket, 221, 231
square integrable martingale, 101,
 186, 232, 233
stationary distribution, 172, 358,
 362, 405
Stieltjes integral, 10
stochastic differential, 108
stochastic differential equations, 123
stochastic exponential, 129, 131, 226,
 227, 238, 258
stochastic integral, 234
stochastic integral discrete time, 84
stochastic intensity, 258, 388
stochastic logarithm, 129, 130,
 238, 250
stochastic processes, 23, 47
stochastic sets, 243
stochastic volatility, 309
stopping time, 25, 51, 68, 244
straddle, 322
Stratonovich calculus, 146
strong Markov property, 68
strong solution, 126, 167
super(sub)martingale, 50, 185
swap, 347
swaption, 347, 351
Tanaka's formula, 224
Taylor's formula, 179
term structure, 328
transformation, 32
transforms, 37
transient, 81
transition function, 136
transition probabilities, 67
uniform integrability, 187
usual conditions, 50
variation, 4, 6, 8
variation of martingales, 202
Vasicek's model, 330
weak solution, 137, 138, 209
white noise, 124
Wright–Fisher model, 369
Yamada–Watanabe, 135, 176
yield curve, 325, 332