ON BRANCHED MINIMAL IMMERSIONS OF SURFACES BY FIRST EIGENFUNCTIONS

DONATO CIANCI, MIKHAIL KARPUKHIN, AND VLADIMIR MEDVEDEV

ABSTRACT. It was proved by Montiel and Ros that for each conformal structure on a compact surface there is at most one metric which admits a minimal immersion into some unit sphere by first eigenfunctions. We generalize this theorem to the setting of metrics with conical singularities induced from branched minimal immersions by first eigenfunctions into spheres. Our primary motivation is the fact that metrics realizing maxima of the first non-zero Laplace eigenvalue are induced by minimal branched immersions into spheres. In particular, we show that the properties of such metrics induced from \mathbb{S}^2 differ significantly from the properties of those induced from \mathbb{S}^m with m>2. This feature appears to be novel and needs to be taken into account in the existing proofs of the sharp upper bounds for the first non-zero eigenvalue of the Laplacian on the 2-torus and the Klein bottle. In the present paper we address this issue and give a detailed overview of the complete proofs of these upper bounds following the works of Nadirashvili, Jakobson-Nadirashvili-Polterovich, El Soufi-Giacomini-Jazar, Nadirashvili-Sire and Petrides.

1. Introduction

Let (Σ, g) denote a closed, connected Riemannian surface where the metric g is induced from a minimal isometric immersion into a round sphere of radius r. That is, $\Phi \colon (\Sigma, g) \to (\mathbb{S}_r^n, g_{\operatorname{can}})$ is a minimal isometric immersion. By a well known result of Takahashi [Tak66, Theorem 3], the coordinate functions of such minimal immersions Φ are given by eigenfunctions for the Laplace-Beltrami operator on (Σ, g) with corresponding eigenvalue $\frac{2}{r^2}$. However, not all immersions are by first eigenfunctions. The following theorem shows that each conformal class of Σ admits at most one metric induced from an immersion into a sphere by first eigenfunctions:

Theorem 1.1 ([MR86, ESI86]). For each conformal structure on a compact surface, there exists at most one metric which admits an isometric immersion into some unit sphere by first eigenfunctions.

In this article we generalize Theorem 1.1 to the setting of branched minimal immersions into round spheres by first eigenfunctions (see Theorem 1.4 for a precise statement). Branched minimal immersions are given by smooth maps $\Phi \colon \Sigma \to \mathbb{S}^n$ which are minimal immersions except at finitely many points at which Φ becomes

The first author was supported by a CRM-Laval Postdoctoral Fellowship while some of this research was conducted.

The second author was partially supported by a Schulich Fellowship.

singular. In this situation, the pullback metric $\Phi^*g_{\rm can}$ on Σ will possess conical singularities at the singular points of Φ . Branched minimal immersions into spheres by first eigenfunctions occur in the study of metrics which maximize the first non-zero Laplace eigenvalue, denoted λ_1 , among all metrics of area one. Indeed, in [MS17], Matthiesen and Siffert proved that for any closed surface Σ there exists a metric \hat{g} of area one, smooth except for possibly finitely many points which correspond to conical singularities, that maximizes λ_1 among all other unit-area metrics on Σ . These maximal metrics are induced from branched minimal immersions into a round sphere by first eigenfunctions and do in general possess conical singularities (see [NS18]). Therefore, it is natural to study Theorem 1.1 in the context of branched minimal immersions.

A technical difficulty unique to the branched immersion case is that one can have branched minimal immersions by first eigenfunctions whose images are an equatorial 2-sphere. Indeed, the conclusion of Theorem 1.1 is valid only with the restriction that the image of the branched minimal immersion is not an equatorial 2-sphere. This restriction indicates that the branched minimal immersions by first eigenfunctions into \mathbb{S}^2 are in a way special. Moreover, we show that if a conformal class has a metric induced by a branched minimal immersion by first eigenfunctions to \mathbb{S}^2 then it does not have a metric induced by a non-trivial branched minimal immersion by first eigenfunctions to a higher-dimensional sphere.

Theorem 1.1 has been applied to help classify certain metrics which maximize λ_1 (see the discussion in the next section). However, our generalization of Theorem 1.1 presents a novel feature that needs to be taken into account in this classification. In the present article we address this issue by proving that there are no branched minimal immersions of a torus or a Klein bottle to \mathbb{S}^2 . In order to precisely state our results, we give a more detailed version of the previous discussion.

1.1. Maximization of the first eigenvalue on surfaces and minimal immersions. After fixing a surface Σ , let $\mathcal{R}(\Sigma)$ be the collection of Riemannian metrics on Σ . We have the following homothety invariant functional on $\mathcal{R}(\Sigma)$:

$$\bar{\lambda}_1: \mathcal{R}(\Sigma) \to \mathbb{R}_{\geq 0}; \quad \bar{\lambda}_1(\Sigma, \cdot): g \mapsto \lambda_1(g) \operatorname{vol}(\Sigma, g),$$

where $\lambda_1(g)$ is the first non-zero Laplace-Beltrami eigenvalue of (Σ, g) and $\operatorname{vol}(\Sigma, g)$ is the area of (Σ, g) .

Using the notion of *conformal volume* (see Section 2), Li and Yau [LY82] established the following upper bound for $\bar{\lambda}_1(\Sigma, g)$ when Σ is orientable and has genus γ (see also [YY80]):

(1)
$$\bar{\lambda}_1(\Sigma, g) \le 8\pi \left| \frac{\gamma + 3}{2} \right|,$$

where the bracket denotes the integer part of the number inside. Modifying the ideas of Li and Yau, the second author [Kar16, Theorem 1] proved the following upper

bound for non-orientable surfaces (of genus γ):

(2)
$$\bar{\lambda}_1(\Sigma, g) \le 16\pi \left| \frac{\gamma + 3}{2} \right|.$$

Here the genus of a non-orientable surface is defined to be the genus of its orientable double cover.

Thus, $\bar{\lambda}_1(\Sigma, g)$ is bounded above on $\mathcal{R}(\Sigma)$. Naturally, one is interested in finding sharp upper bounds for $\bar{\lambda}_1(\Sigma, g)$ for a given surface and also characterizing the maximal metrics.

Definition 1.1. Let Σ be a closed surface. A metric g_0 on Σ is said to be maximal for the functional $\bar{\lambda}_1(\Sigma, g)$ if

$$\bar{\lambda}_1(\Sigma, g_0) = \sup_{g \in \mathcal{R}(\Sigma)} \bar{\lambda}_1(\Sigma, g).$$

Throughout, we will denote the value of $\sup_{g \in \mathcal{R}(\Sigma)} \bar{\lambda}_1(\Sigma, g)$ by $\Lambda_1(\Sigma)$. Additionally we set $\Lambda_1(\Sigma, [g])$ to be $\sup_{g \in [g]} \bar{\lambda}_1(\Sigma, g)$, where [g] denotes the conformal class of a metric g. The following theorem guarantees the existence of a maximal metric on Σ , modulo finitely many points at which the metric may have conical singularities.

Theorem 1.2 ([MS17]). For any closed surface Σ , there is a metric g on Σ , smooth away from finitely many conical singularities, achieving $\Lambda_1(\Sigma)$, i.e.

$$\Lambda_1(\Sigma) = \bar{\lambda}_1(\Sigma, g).$$

- Remark 1.2. (i) The proof of Theorem 1.2 uses results of Nadirashvili and Sire [NS15] and Petrides [Pet14] on the maximization of $\Lambda_1(\Sigma, g)$ in a conformal class.
 - (ii) Nayatani and Shoda [NS18] recently proved that Λ_1 is maximized by a metric on the Bolza surface with constant curvature one and six conical singularities (this metric was proposed to be maximal in [JLN+05]). Thus, Theorem 1.2 is optimal in regards to the regularity of a maximal metric.

As the next theorem shows, these maximal metrics for $\bar{\lambda}_1$ are induced from branched minimal immersions into round spheres. It was first proved by Nadirashvili in [Nad96] for the particular case of $\bar{\lambda}_1$. Later, the theorem was generalized to maximal metrics for higher Laplace eigenvalues in [ESI08]. As noted in [NP18], the theorem also holds for metrics with conical singularities (in part because the variational characterization of λ_1 is the same whether considering metrics with conical singularities or smooth metrics). Together with Theorem 1.2, Theorem 1.3 is our motivation for studying branched minimal immersions by first eigenfunctions.

Theorem 1.3 ([Nad96],[ESI08],[Kok14]). Let g_0 be a metric on a closed surface Σ , possibly with conical singularities. Moreover, suppose that:

$$\Lambda_1(\Sigma) = \bar{\lambda}_1(\Sigma, g_0).$$

Then g_0 is induced from a (possibly branched) minimal isometric immersion into a sphere by first eigenfunctions.

We briefly review some results regarding $\bar{\lambda}_1$ -maximal metrics (for results regarding extremal metrics, see the survey [Pen13a] and the papers [ESI00, Kar13, Kar14, Kar15, Lapo8, Pen12, Pen13b, Pen15]). By Theorem 1.3, any $\bar{\lambda}_1$ -maximal metric is induced by a (possibly branched) minimal immersion into a sphere. Hersch proved in 1970 that $\Lambda_1(\mathbb{S}^2)$ is achieved by any constant curvature metric [Her70]. By the work of Li and Yau [LY82], $\bar{\lambda}_1(\mathbb{RP}^2, g) \leq 12\pi$ for any metric g with equality for constant curvature metrics. Indeed, the metric of constant curvature one on \mathbb{RP}^2 can be realized as the induced metric from a minimal embedding into S⁴ called the Veronese embedding. Since there is only one conformal class of metrics on \mathbb{RP}^2 , Theorem 1.1 shows that $\bar{\lambda}_1(\mathbb{RP}^2,q) < 12\pi$ with equality only if the metric is a constant curvature metric. In [Nad96], Nadirashvili proved the existence of maximal metrics on the 2-torus (see also [Gir09]) and outlined a proof of existence for metrics on the Klein bottle. In the next section we discuss the cases of the 2-torus and the Klein bottle in more detail. Finally, the maximal metric is known for Σ_2 , the orientable surface of genus 2. Nayatani and Shoda proved in [NS18] that the metric on the Bolza surface proposed in [JLN+05] is maximal. As a result, $\Lambda_1(\Sigma_2) = 16\pi$.

1.2. **Main results.** We prove the following generalization of Theorem 1.1 to the setting of *branched* minimal immersions.

Theorem 1.4. Let Σ be a closed surface endowed with a conformal class c. Then c belongs to exactly one of the following categories:

- 1) There does not exist $g \in c$ such that g admits a branched minimal immersion to a sphere by first eigenfunctions;
- 2) There exists a unique $g \in c$ such that g admits a branched minimal immersion by first eigenfunctions to \mathbb{S}^m whose image is not an equatorial 2-sphere;
- 3) There exists $g \in c$ such that g admits a branched minimal immersion by first eigenfunctions to \mathbb{S}^2 . In this case any two such immersions differ by a post-composition with a conformal automorphism of \mathbb{S}^2 .

Remark 1.3. If Σ is not orientable, then $\Phi(\Sigma)$ can never be an equatorial 2-sphere. Indeed, this would make $\Phi \colon \Sigma \to \mathbb{S}^2$ a branched cover, which is impossible. In Proposition 3.8 we also prove that if Σ is a 2-torus the image of a branched minimal immersion by first eigenfunctions cannot be an equatorial 2-sphere. Thus, category 3) in Theorem 1.4 is not possible in these cases.

Remark 1.4. Theorem 1.4 allows us to construct an example where a maximal metric for $\bar{\lambda}_1$ cannot be induced by a branched minimal immersion whose components form a basis in the λ_1 -eigenspace. Indeed, in [NS18] the authors showed that on a surface of genus 2 there exists a family of maximal metrics for $\bar{\lambda}_1$ induced from a branched minimal immersion to \mathbb{S}^2 . Moreover, there are metrics in the family such that the multiplicity of the first eigenvalue is equal to 5. At the same time by Theorem 1.4 such a metric can only be induced by a 3-dimensional family of eigenfunctions.

The following theorem was proved by Nadirashvili in [Nad96] for the case of \mathbb{T}^2 , and the same paper contains an outline of the proof for the Klein bottle. Later, Girouard completed some of the steps of this outline in [Gir09].

Theorem 1.5. The maximal values $\Lambda_1(\mathbb{T}^2)$ and $\Lambda_1(\mathbb{KL})$ are achieved by smooth Riemannian metrics.

Let us discuss the proof of Theorem 1.5 presented in [Nad96]. Nadirashvili first shows that Theorem 1.2 holds for the torus, i.e. there exists a maximal metric possibly with conical singularities. Then he applies Theorem 1.1 to conclude that the maximal metric is flat. However, as we see from Theorem 1.4 the conclusion of Theorem 1.1 does not hold for branched immersions and special care is needed if the maximal metric happens to be induced by a branched minimal immersion to \mathbb{S}^2 . Our contribution to Theorem 1.5 is that we show that there are no branched minimal immersions by first eigenfunctions to \mathbb{S}^2 from either the 2-torus or the Klein bottle. While the case of the Klein bottle is elementary, see Remark 1.3, additional considerations are required to settle the case of the 2-torus, see Proposition 3.8.

Once Theorem 1.5 is proved, Nadirashvili's argument shows that the maximal metric on \mathbb{T}^2 is flat. Let us recall it in more detail. It follows by Theorem 1.1 that any conformal transformation of a smooth maximal metric is an isometry. Since any metric on the 2-torus has a transitive group of conformal transformations, then any smooth maximal metric must have a transitive group of isometries and is therefore flat. It follows that a smooth maximal metric is a scalar multiple of the flat metric on the equilateral torus. In the same paper [Nad96], Nadirashvili used a similar argument to deduce that any smooth maximal metric on the Klein bottle must be a surface of revolution. Later, Jakobson, Nadirashvili, and I. Polterovich [JNP06] found a candidate for a smooth maximal metric for the Klein bottle (by proving the existence of a metric of revolution that was extremal for $\bar{\lambda}_1$). The metric they found corresponded to a bipolar surface of Lawson's $\tau_{3,1}$ -torus. Then, in [ESGJ06], El Soufi, Giacomini, and Jazar proved that this metric was the only smooth extremal metric on the Klein bottle.

Remark 1.5. Note that Theorem 1.5 can be obtained by combining Theorem 1.4 and Proposition 3.8 with the recent work of Matthiesen and Siffert [MS17]. For completeness, we prove Theorem 1.5 without the result of Matthiesen and Siffert and instead combine results of Girouard [Gir09] with the fact that the functional $\Lambda_1(\Sigma, [g])$ is continuous on the moduli space of conformal classes of metrics on Σ . The latter fact seems to be well-known but we were unable to find a reference. We present its proof in Section 4.

As a result of the previous discussion we have the following corollaries of Theorem 1.5.

Corollary 1.6 ([Nad96]). The maximum for the functional $\bar{\lambda}_1(\mathbb{T}^2, g)$ on the space of Riemannian metrics on a 2-torus \mathbb{T}^2 is attained if and only if the metric g is homothetic to the flat metric g_{eq} on the equilateral torus and has the following value:

$$\Lambda_1(\mathbb{T}^2) = 8\pi^2/\sqrt{3}.$$

Corollary 1.7 ([JNP06, ESGJ06]). The maximum for the functional $\bar{\lambda}_1(\mathbb{KL}, g)$ on the space of Riemannian metrics on a Klein bottle \mathbb{KL} is attained if and only if the metric g is homothetic to a metric of revolution:

$$g_0 = \frac{9 + (1 + 8\cos^2 v)^2}{1 + 8\cos^2 v} \left(du^2 + \frac{dv^2}{1 + 8\cos^2 v} \right),$$

 $0 \le u < \pi/2, 0 < v \le \pi$ and has the following value:

$$\Lambda_1(\mathbb{KL}) = 12\pi E(2\sqrt{2}/3) \approx 13.365\pi,$$

where $E(\cdot)$ is a complete elliptic integral of the second kind.

There are inconsistencies in the literature regarding whether there is a complete proof that the extremal metric on the Klein bottle found in [JNP06] is indeed maximal. See, for instance, Remark 1.1 in [ESGJ06]. One of the goals of the present article is to eliminate this inconsistency.

Paper outline. The rest of the paper is organized as follows: In Section 2 we provide the necessary background for studying branched minimal immersions into spheres and recall the definition of conformal volume. Section 3 contains the proofs of Theorem 1.4 and Proposition 3.8. In Section 4 we prove Theorem 1.5 and that the conformal spectrum is continuous on the moduli space of conformal classes of metrics on Σ .

Acknowledgements. The authors are grateful to I. Polterovich and A. Girouard for suggesting this problem. The authors are also grateful to I. Polterovich, A. Girouard, G. Ponsinet, A. Penskoi, G. Kokarev, and Alex Wright for fruitful discussions and especially to A. Girouard for a careful first reading of this manuscript. This research is part of the third author's PhD thesis at the Université de Montréal under the supervision of Iosif Polterovich.

2. Background

2.1. Branched immersions and conical singularities. Given a surface Σ endowed with a conformal structure one defines a metric g with conical singularities by declaring that at finitely many points $\{p_1,...,p_N\} \subset \Sigma$ (which are referred to as conical points) the metric has the following form in conformal coordinates centered at p_i : $\rho_i(z)|z|^{2\beta_i}|dz|^2$, where $\beta_i > -1$ and $\rho_i(z) > 0$ is smooth. The metric is singular in the sense that it becomes degenerate at the conical points. This approach is taken, for instance, in [Tro91]. One can check that if $\rho = 1$ near the conical points then g is isometric to a cone with cone angle $2\pi(\beta_i + 1)$. In this article we are primarily concerned with metrics with conical singularities that arise from branched minimal immersions into spheres. A good introductory reference for branched minimal immersions is [GOR73].

Fix a compact surface Σ equipped with a smooth Riemannian metric g_0 (without conical points). Let $\Phi \colon (\Sigma, g_0) \to (\mathbb{S}^n, g_{\operatorname{can}})$ be a smooth map that is harmonic and

conformal away from points at which $D_p\Phi=0$. We will refer to points p at which $D_p\Phi=0$ as branch points and call Φ a branched conformal immersion. Note that away from the branch points the quadratic form $g=\Phi^*g_{\rm can}$ is actually an inner product on the tangent space and makes Φ a minimal immersion. Thus, we say that $\Phi\colon (\Sigma,g)\to (\mathbb{S}^n,g_{\rm can})$ is a branched minimal immersion into a sphere. We will see that g possesses conical singularities at the branch points.

In a neighborhood of a branch point we can choose conformal coordinates $z = z_1 + iz_2$ on Σ centered at p and coordinates $x_1, ..., x_n$ centered at $\Phi(p)$ such that $\Phi(z)$ takes the form:

$$x_1 + ix_2 = z^{m+1} + \sigma(z)$$

$$x_k = \phi_k(z); \ k \ge 3,$$

for $m \geq 1$ such that $\sigma(z)$ and $\phi_k(z)$ are $o(|z|^{m+1})$ and $\frac{\partial \sigma}{\partial z_j}$ and $\frac{\partial \phi_k}{\partial z_j}$ are $o(|z|^m)$ as $z \to 0$ (that this is possible follows from the discussion found in [GOR73, Section 2]). The integer m is referred to as the *order* of the branch point. Moreover, there exist $C^{1,\alpha}$ -coordinates (see Lemma 1.3 of [GOR73]) \widetilde{z} , which we will refer to as distinguished parameters, in which the map Φ takes the form:

$$x_1 + ix_2 = \widetilde{z}^{m+1}$$
$$x_k = \psi_k(z); \ k \ge 3,$$

with $\psi_k(z)$ possessing the same asymptotics as $\phi_k(z)$ as $z \to 0$. Note that the distinguished parameters are not an admissible coordinate system for the smooth structure on Σ since \widetilde{z} is related to z by $\widetilde{z} = z \left[1 + z^{-(m+1)}\sigma(z)\right]^{1/(m+1)}$. By looking at the form Φ takes in distinguished parameters it is clear that $D_p\Phi \neq 0$ in a punctured neighborhood of a branch point. Thus, branch points are isolated. Moreover, since regular points form an open set, Σ can only posses finitely many branch points. From the previous discussion we see that in conformal coordinates centered at a branch point the metric is of the form $\rho(z)|z|^{2m}|dz|^2$, with $\rho(z)>0$ smooth. In other words, near the branched point the metric is conformal to the Euclidean cone of total angle $2\pi(m+1)$. We will refer to m as the order of the conical singularity and will also refer to the branch points p as conical points.

We recall the following lemma, which allows one to define the tangent space to $\Phi(\Sigma)$ at the image of a conical point $\Phi(p)$. For simplicity, we state the lemma in the setting of branched conformal immersions into a round sphere. However, it holds in greater generality (see Lemma 3.1, 3.2 and the remark on page 771 of [GOR73] for the proof).

Lemma 2.1. Let $\Phi: \Sigma \to \mathbb{S}^n$ be a branched minimal immersion into a round sphere with a branch point at $p \in \Sigma$. Let w and x be distinguished parameters at p and $\Phi(p)$, respectively. Define the tangent space to $\Phi(p)$ in distinguished parameters as the x_1, x_2 -plane.

(i) If $\{p_n\}$ is a sequence in Σ such that $p_n \to p$ and Φ is regular at p_n , then the tangent plane to $\Phi(\Sigma)$ at $\Phi(p_n)$ tends to the x_1, x_2 -plane in distinguished

parameters. Consequently, the Gauss map, which assigns to each point $q \in \Sigma$ the tangent plane to $\Phi(\Sigma)$ at $\Phi(q) \in \mathbb{S}^n$ is continuous on all of Σ .

(ii) The definition of the tangent space to $\Phi(p)$ does not depend on the choice of distinguished parameters.

Let g be a metric on Σ with conical singularities. Thus, $g = fg_0$, where g_0 is a smooth Riemannian metric on Σ and f is a smooth function on Σ that is nonzero except at possibly finitely many points. One can define the first Laplace eigenvalue corresponding to this singular metric using the variational characterization:

(3)
$$\lambda_1(g) = \inf_{\substack{u \in H^1(\Sigma,g) \\ u+1}} R(u,g),$$

where:

$$R(u,g) = \frac{\int_{\Sigma} |\nabla u|^2 dV(g_0)}{\int_{\Sigma} u^2 dV(g)}$$

is the Rayleigh quotient and $H^1(\Sigma, g)$ is the completion of the set:

$$\left\{ u \in L^2(\Sigma, dV(g)); \quad \int_{\Sigma} |\nabla u|^2 dV(g_0) < \infty \right\}$$

with respect to the norm:

$$||u||_{H^{1}(g)}^{2} = \int_{\Sigma} u^{2} dV(g) + \int_{\Sigma} |\nabla u|^{2} dV(g_{0}).$$

When a metric g_0 is smooth, we will regard $H^1(\Sigma, g_0)$ as the usual Sobolev space. Note that, essentially by the conformal invariance of the Dirichlet energy, $H^1(\Sigma, g) = H^1(\Sigma, g_0)$, meaning that they are equal as sets, and the norms define the same topology (see [Tro91, Proposition 3]). A function $u \in H^1(\Sigma, g)$ for which the infimum of the Rayleigh quotient in (3) is achieved is called a first eigenfunction. For metrics with conical singularities, first eigenfunctions exist (see [Kok14] Proposition 3.1). By the usual elliptic regularity argument (see, for instance, [GT01, Corollary 8.11]), the first eigenfunctions are smooth and satisfy the following equation:

$$\Delta_{g_0} u = \lambda_1(g) f u.$$

2.2. Conformal volume. The notion of conformal volume was introduced by Li and Yau to prove bounds on λ_1 that depend only on the genus [LY82]. It will be used in our poof of Theorem 1.4. Throughout, let G_n denote the group of conformal diffeomorphisms of the n-sphere with its canonical metric and let $\Phi \colon \Sigma \to \mathbb{S}^n$ be a conformal immersion with possible branch points.

Definition 2.1. (i) The conformal n-volume of Φ is given by:

$$\operatorname{vol}_c(n,\Phi) := \sup_{\gamma \in G_n} \operatorname{vol}(\Sigma, (\gamma \circ \Phi)^* g_{\operatorname{can}}).$$

(ii) The conformal n-volume of Σ , denoted $\operatorname{vol}_c(n,\Sigma)$, is the infimum of $\operatorname{vol}_c(n,\Phi)$ over all branched conformal immersions $\Phi \colon \Sigma \to \mathbb{S}^n$.

Remark 2.2. In the recent preprint [Kok17] Kokarev used conformal volume to obtain bounds for higher eigenvalues λ_k .

3. Proofs of main results

To prove Theorem 1.4 we follow the same steps used by El Soufi and Ilias to prove the analog of Theorem 1.4 for minimal immersions without branch points (see Corollary 3.3 in [ESI86]). While some propositions easily generalize to the setting of branched minimal immersions (compare Proposition 3.2 with [ESI86, Theorem 2.2]) others do not generalize completely (compare Proposition 3.3 with [ESI86, Proposition 3.1]).

Proposition 3.1. Let (Σ, g) be a compact Riemannian surface possibly with conical singularities and suppose that there exists a branched minimal isometric immersion Φ of (Σ, g) into a round sphere of dimension n, then:

$$\operatorname{vol}(\Sigma, g) = \operatorname{vol}_c(n, \Phi) \ge \operatorname{vol}_c(n, \Sigma).$$

Moreover, if $\Phi(\Sigma)$ is not an equatorial 2-sphere, then $\operatorname{vol}(\Sigma, g) > \operatorname{vol}(\Sigma, (\gamma \circ \Phi)^* g_{\operatorname{can}})$ for every $\gamma \in G \setminus O(n+1)$.

Proof. Given a unit-vector $a \in \mathbb{R}^{n+1}$, let A denote the projection of the vector onto the tangent space of each point \mathbb{S}^n . Then A is the gradient vector field of the function $u = \langle \cdot, a \rangle$. Let $(\gamma_t^a)_t$ be the time-t map for the flow associated to A. Then $(\gamma_t^a)^* g_{\text{can}} = e^{2f} g_{\text{can}}$, with f a smooth function on the sphere. Recall that for every $\gamma \in G_n$ there exists $r \in O(n+1)$ and γ_t^a such that $\gamma = r \circ \gamma_t^a$ (see the Lemma on page 259 of [ESI86]). Thus, it suffices to show that:

$$\operatorname{vol}(\Sigma, (\gamma_t^a \circ \Phi)^* g_{\operatorname{can}}) \leq \operatorname{vol}(\Sigma, \Phi^* g_{\operatorname{can}}) \text{ for every } a \in \mathbb{S}^n \text{ and } t \geq 0.$$

First, we need to verify that we can make use of the first variation formula for the area of Σ . Let $\{p_1, ..., p_N\}$ denote the branch points of Φ . Then

$$\widehat{\Sigma} := \Sigma \backslash \{p_1, ..., p_N\}$$

is an (open) Riemannian manifold and Φ induces a minimal isometric immersion of $\widehat{\Sigma}$ into \mathbb{S}^n . For convenience, we will often identify $\widehat{\Sigma}$ and its image under Φ . In coordinates centered at a branch point of order m, the volume form is given by:

$$dV((\gamma_t^a \circ \Phi)^* g_{can})(z) = \rho(t, z)|z|^{2m}|dz \wedge d\overline{z}|,$$

where $\rho(t,z)$ is a smooth positive function. Thus, it is clear that the volume form is differentiable (smooth) in t and the derivative with respect to t is identically zero at the branch points. Set $\gamma = \gamma_{t_0}^a$. Away from the singular points, we have the usual expression for the derivative of the volume form:

(4)
$$\frac{d}{dt}\Big|_{t=t_0} dV((\gamma_t^a \circ \Phi)^* g_{\text{can}})(x) = -\left\langle A_{\gamma(x)}, H_{\gamma(x)}^{\gamma(\widehat{\Sigma})} \right\rangle dV((\gamma \circ \Phi)^* g_{\text{can}})(x) + div_{\gamma(\widehat{\Sigma})} (A_{\gamma(x)}^{\top}) dV((\gamma \circ \Phi)^* g_{\text{can}})(x),$$

where $H_{\gamma(x)}^{\gamma(\widehat{\Sigma})}$ is the mean curvature vector for $\gamma(\widehat{\Sigma})$ and A^{\top} is the projection of A onto the tangent space of $\gamma(\widehat{\Sigma})$. Since Φ is minimal away from the branch points, one can compute the expression for the mean curvature vector explicitly away from the branch points (see page 260 of [ESI86]):

$$H_{\gamma(x)}^{\gamma(\widehat{\Sigma})} = -2e^{-2f}D\gamma((\nabla f^{\perp})_x).$$

Moreover, by Lemma 2.1, ∇f^{\perp} extends to a continuous vector field on all of $\Phi(\Sigma)$. Thus, $H^{\gamma(\Sigma)}$ extends to a continuous vector field on the branch points. It follows that the first term in the right hand side of (4) is zero at a branch point. Thus, the second term in the right hand side extends to something continuous and zero at the branch points.

Now we integrate both sides of (4) to recover the usual first variation formula. However, since A_x^{\top} only extends to a continuous vector field on $\Phi(\Sigma)$, some care is required to show that the integral of the second term in the right hand side of (4) is zero. Notice that the integrals of the left hand side and the first term in the right hand side of (4) converge as improper integrals. Thus, it suffices to exhibit an exhaustion of $\gamma(\widehat{\Sigma})$ by compact sets with smooth boundary such that the integral of the second term on the right hand side of (4) converges to zero. Let $\{\Omega_n\}_{n=1}^{\infty}$ be an exhaustion of $\widehat{\Sigma}$ by compact sets with smooth boundary such that for each connected component of $\partial\Omega_n$ there exist distinguished parameters (z_1, z_2) centered at a singular point such that the image of the connected component of $\partial\Omega_n$ is given by $z_1^2 + z_2^2 = \frac{1}{n^2}$, for n large enough. Then $\{\gamma(\Omega_n)\}$ is an exhaustion of $\gamma(\widehat{\Sigma})$. The Divergence Theorem yields:

$$\int_{\gamma(\Omega_n)} \operatorname{div}_{\gamma(\Omega_n)} \left(A_x^{\top} \right) dV(g_{\operatorname{can}}) = \int_{\partial \gamma(\Omega_n)} \langle A_x^{\top}, N_x \rangle ds
= \int_{\partial \Omega_n} \langle A_{\gamma(x)}^{\top}, N_{\gamma(x)} \rangle e^{2f} ds
= \int_{\partial \Omega_n} \langle D\gamma(A_x)^{\top}, N_{\gamma(x)} \rangle e^{2f} ds
= \int_{\partial \Omega_n} \langle D\gamma(A_x^{\top}), N_{\gamma(x)} \rangle e^{2f} ds,$$

where N is the outward pointing unit normal vector field along $\partial \gamma(\Omega_n)$ and ds is the length element along $\partial \gamma(\Omega_n)$. Again by Lemma 2.1, A_x^{\top} extends to a continuous vector field on $\Phi(\Sigma)$, then the Cauchy-Schwartz inequality shows that the integral is $O((1/n)^{2m+1})$. Thus, we see that, as an improper integral, $\int_{\Sigma} \operatorname{div}_{\gamma(\Sigma)}(A_{\gamma(x)}^{\top}) \operatorname{dV}((\gamma \circ \Phi)^* g_{\operatorname{can}}) = 0$. Integrating both sides of (4) yields the usual first variation formula:

$$\frac{d}{dt}\bigg|_{t=t_0} \operatorname{vol}(\Sigma, (\gamma_t^a \circ \Phi)^* g_{\operatorname{can}}) = -\int_{\Sigma} \langle H_{\gamma(x)}^{\gamma(\widehat{\Sigma})}, A_{\gamma(x)} \rangle \, dV \left((\gamma \circ \Phi)^* g_{\operatorname{can}} \right).$$

At this point, the calculation done in the proof of Theorem 1.1 of [ESI86] applies:

(5)
$$\frac{d}{dt}\Big|_{t=t_0} \operatorname{vol}(\Sigma, (\gamma_t^a \circ \Phi)^* g_{\operatorname{can}}) = 2 \int_{\Sigma} \frac{u - u \circ \gamma}{1 - u^2} |A^{\perp}|^2 \, dV((\gamma \circ \Phi)^* g_{\operatorname{can}}),$$

where the integrand is extended by continuity at the branch points. Since $u(x) \le u(\gamma(x))$, we conclude that:

$$\frac{d}{dt}\Big|_{t=t_0} \operatorname{vol}(\gamma_t^a(\widehat{\Sigma})) \le 0.$$

Thus, $\operatorname{vol}(\gamma_t^a(\Phi(\Sigma)))$ is non-increasing.

Now suppose that there exists a and $t_0 > 0$ such that $\operatorname{vol}(\gamma_{t_0}^a(\Phi(\Sigma))) = \operatorname{vol}(\Phi(\Sigma))$. Then $\frac{d}{dt}|_{t=s}\operatorname{vol}(\gamma_t^a(\Phi(\Sigma))) = 0$ for $0 \le s \le t_0$. From this observation and (5) we conclude that $A^{\perp} = 0$ on $\Phi(\Sigma)$. Thus, A restricts to a vector field on $\widehat{\Sigma}$. Observe that the integral curves of A are great circles inside \mathbb{S}^n connecting a and -a. Therefore, $a, -a \in \Phi(\Sigma)$. If a is a regular value of Φ , then $\Phi(\Sigma)$ is just given by the image of $T_a\Phi(\Sigma)$ under the Riemannian exponential map of \mathbb{S}^n based at a. So $\Phi(\Sigma)$ is an equatorial 2-sphere.

Now suppose that a corresponds to a singular value of Φ . Again, by Lemma 2.1 we may define the tangent space to $\Phi(\Sigma)$ at a in $T_a\mathbb{S}^n$. Let V denote this subspace. Given $p \in \Phi(\widehat{\Sigma})$ sufficiently close to a, let $\alpha \colon [0,1] \to \mathbb{S}^n$ be the minimizing geodesic connecting p and a. Then $\alpha((0,1))$ is contained in $\Phi(\widehat{\Sigma})$. Moreover, since $\alpha'(t)$ is in the tangent space to $\Phi(\widehat{\Sigma})$ for every $t \in (0,1)$, then $\alpha'(1) \in V$. This shows that $\Phi(\Sigma)$ is again the image of V under the Riemannian exponential map of \mathbb{S}^n at a. Thus, $\Phi(\Sigma)$ is a 2-sphere.

The following proposition is a generalization of Theorem 2.2 in [ESI86] to the setting of branched minimal immersions. See also Theorem 1 in [LY82]. The proof of Theorem 2.2 in [ESI86] carries through without changes to the setting of branched minimal immersions.

Proposition 3.2. Suppose (Σ, g) is a Riemannian surface with possible conical singularities. For all n such that the conformal n-volume is defined, we have:

$$\bar{\lambda}_1(\Sigma, g) \leq 2 \operatorname{vol}_c(n, \Sigma).$$

Equality holds if and only if (Σ, g) admits, up to homothety, a branched minimal immersion into a sphere by first eigenfunctions.

We generalize Proposition 3.1 of [ESI86] to the setting of branched minimal immersions. However, the statement is complicated by the fact that the image of a branched minimal immersion can be an equatorial 2-sphere.

Proposition 3.3. Let (Σ, g) be a surface with possible conical singularities. Moreover, suppose that the metric g is induced from a branched minimal immersion Φ into a

12

sphere. Then every metric with possible conical singularities \tilde{g} conformal to g satisfies the following:

$$\bar{\lambda}_1(\Sigma, \widetilde{g}) \leq 2 \operatorname{vol}(\Sigma, g).$$

Criteria for equality are as follows:

- If the image of Φ is not an equatorial 2-sphere, then equality holds if and only if the components of Φ are first eigenfunctions and \widetilde{g} is homothetic to g.
- If the image of Φ is an equatorial 2-sphere, then equality holds if and only if there exists a conformal automorphism γ of \mathbb{S}^2 such that \widetilde{g} is homothetic to $(\gamma \circ \Phi)^* g_{\operatorname{can}}$ and the components of $\gamma \circ \Phi$ are first eigenfunctions.

Remark 3.1. Note that in the second case the coordinates of Φ are not necessarily first eigenfunctions, see Example 3.2 below.

Proof. Let $\tilde{g} \in [g]$ be another metric with possible conical singularities from the conformal class of the metric g. Let Φ_i denote the i-th component function corresponding to Φ (as a map from Σ to $\mathbb{S}^n \subset \mathbb{R}^{n+1}$). According to the proof of Theorem 1 in [LY82], there exists a conformal automorphism of \mathbb{S}^n , denoted γ , such that for every i we have:

(6)
$$\int_{\Sigma} (\gamma \circ \Phi)_i dV(\widetilde{g}) = 0.$$

Set $\Psi = \gamma \circ \Phi$. Then using the conformal invariance of the Dirichlet energy and the variational characterization of Laplace eigenvalues, we have:

(7)
$$\lambda_1(\widetilde{g}) \le \frac{\sum_i \int_{\Sigma} |d\Psi_i|_{\widetilde{g}}^2 \, dV(\widetilde{g})}{\sum_i \int_{\Sigma} \Psi_i^2 \, dV(\widetilde{g})}$$

(8)
$$= \frac{2\operatorname{vol}(\Sigma, (\gamma \circ \Phi)^* g_{\operatorname{can}})}{\operatorname{vol}(\Sigma, \widetilde{g})}$$

(9)
$$\leq 2 \operatorname{vol}(\Sigma, g) \operatorname{vol}(\Sigma, \widetilde{g})^{-1},$$

where the second inequality follows from Proposition 3.1. The desired inequality follows.

Now assume that the equality is achieved, i.e. inequalities (7) and (9) turn into equalities and one has

(10)
$$\bar{\lambda}_1(\Sigma, \widetilde{g}) = 2 \operatorname{vol}(\Sigma, g).$$

By Proposition 3.2 $\bar{\lambda}_1(\Sigma, \tilde{g}) \leq 2 \operatorname{vol}_c(n, \Sigma)$. Combining with equality (10), this yields $\operatorname{vol}(\Sigma, q) \leq \operatorname{vol}_c(n, \Sigma)$.

At the same time, by Proposition 3.1, the reverse inequality is true. Therefore, one has an equality.

Assume that the image is not an equatorial 2-sphere. Then by Proposition 3.1 γ is an isometry. Since any isometry of the sphere is linear, equality (6) is satisfied

with $\gamma = \text{id}$. Together with equality (7) this yields that coordinates of Φ are first eigenfunctions for the metric \tilde{g} . If $\tilde{g} = e^{2\omega}g$ then

(11)
$$\lambda_1(\widetilde{g})\Phi = \Delta_{\widetilde{g}}\Phi = e^{-2\omega}\Delta_g\Phi = e^{-2\omega}\lambda_1(g)\Phi.$$

We conclude that in this case ω is constant and \widetilde{g} is homothetic to g.

Now suppose that the image is an equatorial 2-sphere. In this case we cannot conclude that γ is an isometry. Nevertheless, equalities in (6) and (7) imply that coordinates of Ψ are first eigenfunctions for \tilde{g} . Setting $g' = \Psi^* g_{\text{can}}$ and $\tilde{g} = e^{2\omega} g'$ we obtain similarly to (11),

$$\lambda_1(\widetilde{g})\Psi = \Delta_{\widetilde{g}}\Psi = e^{-2\omega}\Delta_{g'}\Psi = 2e^{-2\omega}\Psi.$$

We conclude that in this case ω is constant and \tilde{g} is homothetic to g'.

Let us prove the converse to the equality statements. If the components of Φ are first eigenfunctions then $\lambda_1(g) = 2$. Since \tilde{g} is homothetic to g one has

$$\bar{\lambda}_1(\Sigma, \widetilde{g}) = \bar{\lambda}_1(\Sigma, g) = 2 \operatorname{vol}(\Sigma, g).$$

Suppose that the image of Φ is an equatorial 2-sphere. Set $\Psi = \gamma \circ \Phi$ then after rescaling we may assume $\tilde{g} = \Psi^* g_{\text{can}}$ and $\text{vol}(\Sigma, g) = 4\pi |\deg \Phi| = 4\pi |\deg \Psi| = \text{vol}(\Sigma, \tilde{g})$, since conformal transformations preserve the absolute value of the degree. If the components of Ψ are first eigenfunctions then $\lambda_1(\tilde{g}) = 2$ and one has

$$\bar{\lambda}_1(\Sigma, \widetilde{g}) = 8\pi |\deg \Psi| = 8\pi |\deg \Phi| = 2\operatorname{vol}(\Sigma, g).$$

The following proposition is proved in [MR91, Theorem 6]. We reprove it here using a slightly different approach.

Proposition 3.4. Suppose that $\Phi \colon \Sigma \to \mathbb{S}^2$ is a branched minimal immersion by first eigenfunctions. Then

- (i) For any other conformal map $\Psi \colon \Sigma \to \mathbb{S}^2$ one has $|\deg \Psi| \ge |\deg \Phi|$;
- (ii) If $|\deg \Psi| = |\deg \Phi|$ then there exists a conformal transformation γ such that $\Psi = \gamma \circ \Phi$.

Proof. This proposition is a direct corollary of Proposition 3.3. To prove (i), we apply Proposition 3.3 for metrics $\tilde{g} = \Phi^* g_{\text{can}}$ and $g = \Psi^* g_{\text{can}}$. Then $\lambda_1(\tilde{g}) = 2$ and we conclude

$$8\pi |\deg \Phi| = \bar{\lambda}_1(\Sigma, \widetilde{g}) \le 2\operatorname{vol}(\Sigma, g) = 8\pi |\deg \Psi|.$$

If $|\deg \Phi| = |\deg \Psi|$, then we switch the roles of g and \tilde{g} and observe that we have an equality, i.e. by Proposition 3.3 there exists a conformal automorphism γ_0 such that

$$(12) \qquad (\gamma_0 \circ \Phi)^* g_{\operatorname{can}} = \Psi^* g_{\operatorname{can}}.$$

We would like to show that it implies the existence of an isometry I of \mathbb{S}^2 such that $\Psi = I \circ \gamma_0 \circ \Phi$.

Lemma 3.5. Let f and h be two holomorphic maps $\Sigma \to \mathbb{S}^2$ (i.e. meromorphic functions) such that for any choice of local coordinates one has $|f_z| = |h_z|$. Then there exists $\alpha \in \mathbb{R}$ and $c \in \mathbb{C}$ such that $f = e^{i\alpha}h + c$.

Proof. First, note that the condition of the lemma implies that that f and h have the same singular sets. Let $p \in \Sigma$ be any regular point of f and h, i.e. $df(p) \neq 0$ and $dh(p) \neq 0$. Let z be local coordinates, then there exists a real-valued function $\alpha(z)$ such that $f_z = e^{i\alpha(z)}h_z$. Taking $\partial_{\bar{z}}$ of both parts we obtain

$$i(\partial_{\bar{z}}\alpha)e^{i\alpha}h_z=0.$$

Since $h_z \neq 0$ in a neighborhood of p, one concludes that α is a real-valued holomorphic function. Thus, α is a constant. Integrating the equality $f_z = e^{i\alpha}h_z$, we obtain an equality $f = e^{i\alpha}h + c$ valid in a neighborhood of p. Since it is an equality between two meromorphic functions, by unique continuation it is valid everywhere on Σ . \square

By taking conjugates if necessary, we can assume that $\gamma_0 \circ \Phi$ and Ψ are both holomorphic. Equality (12) guarantees that we can apply the previous lemma to these functions. The conclusion of the lemma then provides a desired isometry I. Setting $\gamma = I \circ \gamma_0$ concludes the proof.

Example 3.2. In this example we demonstrate that the application of conformal transformations does not in general preserve the property "coordinate functions are the first eigenfunctions."

Let S be a Bolza surface and let $\Pi: S \to \mathbb{S}^2$ be the corresponding hyperelliptic projection. By [NS18], Π is given by first eigenfunctions. Let us consider instead

 $\Pi_t = \gamma_t \circ \Pi$, where $\gamma_t = \frac{z+it}{1-itz}$, $t \in [0,1)$ is a conformal transformation moving the points of \mathbb{S}^2 towards the point i along the shortest geodesic (the point -i does not move). We claim that for t close to 1 the first eigenvalue $\lambda_1(\mathcal{S}, \Pi_t^* g_{\operatorname{can}})$ is close to 0. Informally, it can be explained in the following way. As t tends to 1 the images of the branch points of Π_t are getting closer and closer together. As a result, for large t the surface $(\Sigma, \Pi_t^* g_{\operatorname{can}})$ looks like two spheres glued together with three small cylinders. To make this argument precise, we prove the following proposition.

Proposition 3.6. Suppose that $\Phi: \Sigma \to \mathbb{S}^2$ is a holomorphic map of degree d such that the images of all the branch points lie in an open disk D_r of radius r. Then $\lambda_{d-1}(\Sigma, \Phi^*g_{\operatorname{can}}) = o(1)$ as $r \to 0$.

Proof. Let p be a center of D_r and let π be a stereographic projection onto $\mathbb C$ from -p. Then $\pi(D_r)$ is a Euclidean ball $B_\rho(0)$ of radius $\rho=2\tan\frac{r}{2}$ with center at 0. Note that $\rho=O(r)$ as $r\to 0$. Moreover, the variational capacity of $B_\rho(0)$ in $B_1(0)$ is $2\pi|\ln\rho|^{-1}$. Therefore, there exists a function $f_r\in H^1_0(B_1(0))$ such that $f_r\equiv 1$ on $B_\rho(0)$ and the Dirichlet energy of f_r is o(1). Let $h_r=1-\pi^*f_r$. Then $h_r\equiv 0$ on D_r , $h_r\equiv 1$ on a hemisphere and by conformal invariance of the Dirichlet energy

$$\int_{\mathbb{S}^2} |\nabla h_r|^2 \, dV(g_{\text{can}}) = o(r).$$

Outside $\Phi^{-1}(D_r)$ the map Φ is a covering map. Since $\mathbb{S}^2 \backslash D_r$ is a topological disk, all its covering spaces are trivial. Therefore $\Phi^{-1}(D_r)$ coincides with d copies of $\mathbb{S}^2 \backslash D_r$. Let $h_{1,r}, \ldots, h_{d,r}$ be functions h_r considered as functions on their own copy of $\mathbb{S}^2 \backslash D_r$. We can extend them by zero and consider as functions on Σ . Then their support is disjoint and

$$\frac{\int_{\Sigma} |\nabla h_{i,r}|^2 \, dV(\Phi^* g_{\text{can}})}{\int_{\Sigma} h_{i,r}^2 \, dV(\Phi^* g_{\text{can}})} = \frac{\int_{\mathbb{S}^2} |\nabla h_r|^2 \, dV(g_{\text{can}})}{\int_{\mathbb{S}^2} h_r^2 \, dV(g_{\text{can}})} \le \frac{o(1)}{2\pi} = o(1).$$

The standard argument with min-max characterization of the eigenvalues concludes the proof. \Box

We see that for t close to 1 all branch points of Π_t will concentrate in a small disk around i. Since deg $\Pi_t = 2$ the previous proposition yields that $\lambda_1(\mathcal{S}, \Pi_t^* g_{\operatorname{can}}) \to 0$ as $t \to 1$. Note that this argument works with the point i replaced by an arbitrary point distinct from the branch point of Π .

Proposition 3.7. Let $\Psi \colon \Sigma \to \mathbb{S}^2$ be a conformal map. Suppose $\Phi \colon \Sigma \to \mathbb{S}^n$ is such that $\Phi^*g_{\operatorname{can}} = \Psi^*g_{\operatorname{can}}$. Then the image of Φ lies in an equatorial 2-sphere.

Proof. Let $g = \Psi^* g_{\text{can}} = \Phi^* g_{\text{can}}$. Let II denote the second fundamental form of Φ . Then Gauss' equation implies that at any regular point

$$1 = K + \frac{1}{2}|II|_g^2.$$

At the regular point of g one has K=1, therefore, II=0, i.e. $\Phi(\Sigma)$ is totally geodesic in a neighborhood of any smooth point. Thus, that neighborhood gets mapped to a piece of an equatorial 2-sphere. The conclusion follows from the following standard open-closed argument.

Fix a regular point $p \in \Sigma$. Since Φ is totally geodesic in a neighbourhood U_p of p there exists a 3-dimensional subspace E_p such that $D\Phi(TU_p) \subset E_p$. Let $\Sigma_{\text{reg}} \subset \Sigma$ be the set of regular points. Define V_p to be the set of $q \in \Sigma_{\text{reg}}$ such that $D\Phi(T_q\Sigma) \subset E_p$. Then V_p possesses the following properties.

Non-empty. Indeed, $U_p \subset V_p$.

Open. Indeed, let $q \in V_p$. On one hand, it means that $D\Phi(T_q\Sigma) \subset E_p$. On the other, it is always true that $D\Phi(T_q\Sigma) \subset E_q$. Since $D\Phi(T_q\Sigma)$ is 2-dimensional, it follows that $E_p = E_q$. Therefore, $U_q \subset V_p$.

Closed. Indeed, the complement to V_p has the form $\cup V_q$ for some $q \in \Sigma_{\text{reg}}$. Therefore, it is open.

Finally, we remark that Σ_{reg} is Σ with finitely many points removed. Thus, it is connected. Therefore, $V_p = \Sigma_{\text{reg}}$ and by continuity $\Phi(\Sigma) \subset E_p$.

Proof of Theorem 1.4. Assume that Φ_1 and Φ_2 are branched minimal immersions by first eigenfunctions. Then from Proposition 3.2 we have:

$$2 \operatorname{vol}(\Sigma, g_1) = 2 \operatorname{vol}(\Sigma, g_2),$$

where $g_1 := \Phi_1^* g_{\text{can}}$ and $g_2 := \Phi_2^* g_{\text{can}}$. However, if neither of the images of Φ_1 and Φ_2 are equatorial 2-spheres then by Proposition 3.3 this is only possible if the metrics g_1 and g_2 are homothetic. Since their volumes coincide, they are equal.

Suppose that the image of the map Φ_1 lies in a 2-sphere and the image of the map Φ_2 does not. Let g_1 and g_2 be the corresponding induced metrics. First, note that $\lambda_1(g_1) = \lambda_1(g_2) = 2$. Second, by Proposition 3.2 one has $\bar{\lambda}_1(\Sigma, g_1) = \bar{\lambda}_1(\Sigma, g_2)$. Then by Proposition 3.3 applied to Φ_2 we conclude that g_1 is homothetic to g_2 . Moreover, they have the same first eigenvalue, therefore, $g_1 = g_2$. The conclusion follows from Proposition 3.7.

The aim of the following proposition is to show that there is no conformal class which falls into category 3) of Theorem 1.4 when the surface is a 2-torus.

Proposition 3.8. Let Σ be a 2-torus and $\Phi: \Sigma \to \mathbb{S}^n$ be a non-constant branched minimal immersion by first eigenfunctions. Then the image of Φ cannot be an equatorial 2-sphere.

Proof. This proposition is stated as obvious in Montiel and Ros [MR91, Corollary 8(b)]. However, we were unable to come up with an obvious explanation of this fact. Instead, we provide a proof below. Suppose that there exists a branched minimal map $\Phi \colon \mathbb{T}^2 \to \mathbb{S}^2$ by first eigenfunctions of the metric g. After possibly taking a conjugate, we may assume that Φ is holomorphic, i.e. is given by a meromorphic function f.

Claim 1. $\deg f = 2$.

By inequality (1), deg $f \leq 2$. At the same time, any meromorphic function of degree one is invertible which is impossible for f since $\mathbb{T}^2 \not\approx \mathbb{S}^2$.

Claim 2. For any two meromorphic functions f, h of degree 2 there exists a holomorphic automorphism γ of \mathbb{S}^2 such that $f = \gamma \circ h$.

This immediately follows from Proposition 3.4.

Claim 3. For any point $p \in \mathbb{T}^2$ there exists a meromorphic function f_p of degree 2 such that its only pole has degree 2 and is located at p.

Let Λ be a full rank lattice in \mathbb{C} and suppose that g is conformal to the flat metric on the torus \mathbb{C}/Λ . Then we may take f_p to be $\wp(x-p)$, where \wp is the Weierstrass elliptic function corresponding to Λ (for a definition of the Weierstrass elliptic function, see [Don11, Section 6.2]).

Let $p \neq q$ and let $\gamma(z) = \frac{az+b}{cz+d}$ be such that $f_p = \gamma \circ f_q$. Then

$$(13) f_p(cf_q + d) = af_q + b$$

Claim 4. The divisor of $h = cf_q + d$ is 2p - 2q, i.e. the only zero of h is p, its order is 2; and the only pole of h is q, its order is 2.

The function f_p has a pole of order 2 at p but the left hand side of (13) is finite at p. Therefore, h has a zero of order at least 2 at p. At the same time, deg $h \le 2$, so p is the unique zero and is of order exactly 2. Similarly, h^{-1} has a unique zero of order 2 at q.

By Abel's Theorem (see [Jos02, Section 5.9]), there exists h such that (h) = 2p - 2q iff 2p - 2q = 0 as points in \mathbb{C}/Λ . We arrive at a contradiction since p and q were chosen arbitrary.

4. Application to the 2-torus and the Klein bottle

4.1. Conformal degeneration on the 2-torus and maximal metrics. It is well-known that any metric on the 2-torus is conformally equivalent to a flat one obtained from the Euclidean metric on \mathbb{C} under factorization by some lattice $\Gamma \subset \mathbb{C}$ generated by 1 and $a + ib \in \mathcal{M}$, where

$$\mathcal{M} := \{ a + ib \in \mathbb{C} | 0 \le a \le 1/2, a^2 + b^2 \ge 1, b > 0 \}.$$

Thus, conformal classes are encoded by points of \mathcal{M} (the moduli space of flat tori).

We point out the following action of a subgroup of the group of conformal diffeomorphisms isomorphic to \mathbb{S}^1 . Let \mathbb{C}/Γ where Γ is generated by 1 and $a+ib\in \mathcal{M}$. For $\theta\in\mathbb{R}$ we have an action on \mathbb{C} via translation: $s_{\theta}(x+iy)=x+\theta+iy$. This \mathbb{R} -action on \mathbb{C} induces an \mathbb{S}^1 -action on \mathbb{C}/Γ that has no fixed points. A metric in the conformal class corresponding to $a+ib\in\mathcal{M}$ is given by $f(x+iy)(dx^2+dy^2)$ where f(z) is a smooth positive function that is invariant under the action of Γ . Since s_{θ} is a translation we have: $s_{\theta}^*(f(x+iy)(dx^2+dy^2)) = f(x+\theta+iy)(dx^2+dy^2)$. Thus, s_{θ} acts by conformal diffeomorphisms. We recall the following result concerning maximization of λ_1 and conformal degeneration.

Theorem 4.1 ([Gir09]). Let (g_n) be a sequence of metrics of area one on the 2-torus such that the corresponding sequence $(a_n + ib_n) \in \mathcal{M}$ satisfies $\lim_{n\to\infty} b_n = \infty$, then

$$\lim_{n\to\infty} \lambda_1(g_n) \le 8\pi.$$

4.2. Conformal degeneration on the Klein bottle and maximal metrics. We define the Klein bottle as the quotient of \mathbb{C} under the action of the group G_b , generated by the following elements:

$$t_b(x+iy) = x + i(y+b); \quad \tau(x+iy) = x + \pi - iy.$$

As a consequence of the Uniformization Theorem, any metric on the Klein bottle is conformal to a flat metric on $K_b := \mathbb{C}/G_b$ for some b > 0. Thus, the moduli space of conformal classes of metrics on the Klein bottle is encoded by the positive real numbers. Similar to the case of the 2-torus there is a group of conformal diffeomorphisms isomorphic to \mathbb{S}^1 . Indeed, translations of the form $x + iy \mapsto x + \theta + iy$ induce an action of $\mathbb{R}/\pi\mathbb{Z}$ on K_b without fixed points. Just as above this action induces an action by conformal diffeomorphisms. We recall the following result:

Theorem 4.2 ([Gir09]). Let $(g_n) \subset \mathcal{R}(\mathbb{KL})$ be a sequence of metrics of area one on the Klein bottle.

- (i) If $\lim_{n\to\infty} b_n = 0$, then $\lim \sup_{n\to\infty} \lambda_1(g_n) \le 8\pi$.
- (ii) If $\lim_{n\to\infty} b_n = \infty$, then $\lim \sup_{n\to\infty} \lambda_1(g_n) \le 12\pi$.

Roughly speaking, Theorems 4.1 and 4.2 prove that the maximal metrics for the functional $\bar{\lambda}_1$ on the 2-torus and the Klein bottle must be in a conformal class which corresponds to a fundamental domain which cannot be too "long and skinny."

4.3. Continuity results. One of the classical distances considered on the moduli space of complex structures is the $Teichmüller\ distance$. Naturally, this distance induces a distance d_T on the space of conformal classes. In this section we show that the conformal eigenvalues

$$\Lambda_k(\Sigma, [g]) := \sup_{g' \in [g]} \lambda_k(g') \operatorname{vol}(\Sigma, g')$$

are continuous on the space of conformal classes. This fact should be well-known but we were not able to find a reference.

Here we follow [FM12]. First, we define the Teichmüller distance for orientable surfaces. We define a notion of dilatation. Let $f: \Sigma_1 \to \Sigma_2$ be an orientation-preserving homeomorphism between two Riemann surfaces which is a diffeomorphism outside a finite set of points. The function $k_f(p)$ of f at p is defined in local coordinates as $k_f(p) = \frac{|f_z| + |f_{\bar{z}}|}{|f_z| - |f_{\bar{z}}|}$. It is defined only at points where f is smooth and does not depend on the choice of coordinates. One defines the dilatation of f by the formula $K_f = ||k_f||_{\infty}$. If $K_f < \infty$ then one says that f is K_f -quasiconformal.

For any holomorphic quadratic differential q_1 on a Riemann surface Σ_1 its absolute value $|q_1|$ defines a flat metric with conical singularities at zeroes of q_1 compatible with the complex structure. For any non-singular point $p_0 \in \Sigma_1$ one can define natural coordinates $\eta = \int_{p_0}^p \sqrt{q_1}$ such that $q_1 = d\eta^2$ and $|q_1| = |d\eta|^2$, i.e. the metric is Euclidean in these coordinates. Natural coordinates are defined up to a sign and a translation.

A homeomorphism $f: \Sigma_1 \to \Sigma_2$ between Riemann surfaces is called a *Teichmüller mapping* if there exist holomorphic differentials q_1 on Σ_1 and q_2 on Σ_2 and a real number K > 1 such that

- (i) f maps zeroes of q_1 to zeroes of q_2 ;
- (ii) If p is not a zero of q_1 then with respect to a set of natural coordinates for q_1 and q_2 centered at p and f(p) respectively, the mapping f can be written as

$$f(z) = \frac{1}{2} \left(\frac{K+1}{\sqrt{K}} z + \frac{K-1}{\sqrt{K}} \bar{z} \right),$$

or, equivalently,

$$f(x+iy) = \sqrt{K}x + i\frac{1}{\sqrt{K}}y$$

In particular, a Teichmüller map has dilatation K and is smooth outside of zeroes of q_1 . Moreover, in natural coordinates $|q_1| = dx^2 + dy^2$ and $f^*|q_2| = Kdx^2 + \frac{1}{K}dy^2$.

Theorem 4.3 (Teichmüller's Theorem). Given an orientation preserving homeomorphism $f: \Sigma_1 \to \Sigma_2$ between non-isomorphic Riemann surfaces there exists a Teichmüller mapping homotopic to f. It is unique provided $\chi(\Sigma) < 0$. If $\chi(\Sigma) = 0$

then a Teichmüller mapping is affine and is unique up to a translation, therefore the dilatation is independent of the choice of the mapping.

Definition 4.1. Let Σ be an orientable surface. Consider two different complex structures on Σ making it into Riemann surfaces Σ_1 and Σ_2 . Then one defines the *Teichmüller distance* between Σ_1 and Σ_2 as follows,

$$d_T(\Sigma_1, \Sigma_2) = \frac{1}{2} \inf_f \log(K_f),$$

where f ranges over all Teichmüller mappings $f: \Sigma_1 \to \Sigma_2$.

Remark 4.2. The fact that d_T is indeed a distance function is not obvious and relies on proper discontinuity of the action of the mapping class group on the Teichmüller space.

Teichmüller distance d_T on the moduli space of complex structures induces a distance function on the moduli space of conformal classes. Indeed, let $[g_1]$ and $[g_2]$ be two conformal classes. Choose complex structures Σ_i compatible with $[g_i]$ inducing the same orientation on Σ . Then one sets $d_T([g_1], [g_2]) = d_T(\Sigma_1, \Sigma_2)$.

Up until now we considered orientable surfaces Σ . Let us now address the case of non-orientable Σ . Denote by $\pi: \hat{\Sigma} \to \Sigma$ an orientable double cover and by σ a corresponding involution exchanging the leaves of π . Let $[g_1]$ and $[g_2]$ be two conformal classes on Σ . Choose two complex structures $\hat{\Sigma}_1$ and $\hat{\Sigma}_2$ on $\hat{\Sigma}$ compatible with $[\pi^*g_1]$ and $[\pi^*g_2]$. Then one defines

$$d_T([g_1], [g_2]) = \frac{1}{2} \inf_f \log(K_f),$$

where f ranges over all Teichmüller mappings $f: \hat{\Sigma}_1 \to \hat{\Sigma}_2$ commuting with σ .

Remark 4.3. This definition is implicitly making use of the equivariant version of Teichmüller's Theorem. If in Theorem 4.3 $f \circ \sigma$ is homotopic to $\sigma \circ f$ then the Teichmüller mapping can be chosen σ -equivariant. Indeed, suppose that h is the Teichmüller mapping for f. Then $h \circ \sigma$ is the Teichmüller map for $f \circ \sigma$. Similarly, $\sigma \circ h$ is the Teichmüller map for $\sigma \circ f$. If $\sigma \circ f$ is homotopic to $f \circ \sigma$, then $h \circ \sigma$ must be homotopic to $\sigma \circ h$, and by the uniqueness part of Teichmüller's Theorem we obtain $\sigma \circ h = h \circ \sigma$.

Proposition 4.4. The conformal eigenvalues $\Lambda_k(\Sigma, [g])$ are continuous in the distance d_T .

Proof. We follow the notation of [Kok14]. Namely, given a conformal class c of metrics and a measure μ on Σ we define the Rayleigh quotient

$$R_c(u,\mu) = \frac{\int_{\Sigma} |\nabla u|_g^2 \, dV(g)}{\int_{\Sigma} u^2 \, d\mu}$$

and the eigenvalues $\lambda_k(c,\mu)$ as critical values of the Rayleigh quotient. For a comprehensive study of eigenvalues in this context, including a proof of the existence of eigenfunctions, see [Kok14].

We start with the case of an orientable Σ . Let $[g_1]$ and $[g_2]$ be two conformal classes and let Σ_1 and Σ_2 be the corresponding Riemann surfaces. Denote by $f: \Sigma_1 \to \Sigma_2$ any Teichmüller map, let q_1 and q_2 be the corresponding quadratic differentials and suppose that S_1 and S_2 are their zeroes respectively. By property (ii) at any point of $\Sigma_1 \backslash S_1$ one has

(14)
$$\frac{1}{K}f^*|q_2| \le |q_1| \le Kf^*|q_2|.$$

At this point we use the conformal invariance of the Dirichlet energy. Let $K_i \subset K_{i+1}$ be a compact exhaustion of $\Sigma_2 \backslash S_2$. Similarly, $f^{-1}(K_i)$ form a compact exhaustion of $\Sigma_1 \backslash S_1$. Then for any $u \in H^1(\Sigma_2)$ one has

$$\int_{\Sigma_1 \setminus f^{-1}(K_i)} |\nabla(f^*u)|_{f^*|q_2|}^2 \, dV(f^*|q_2|) = \int_{\Sigma_2 \setminus K_i} |\nabla u|_{|q_2|}^2 \, dV(|q_2|).$$

Combining this with inequality (14), one obtains

$$\frac{1}{K} \int_{\Sigma_1 \setminus f^{-1}(K_i)} |\nabla(f^*u)|_{f^*|q_1|}^2 \, dV(f^*|q_1|) \le \int_{\Sigma_2 \setminus K_i} |\nabla u|_{|q_2|}^2 \, dV(|q_2|)
\le K \int_{\Sigma_1 \setminus f^{-1}(K_i)} |\nabla(f^*u)|_{f^*|q_1|}^2 \, dV(f^*|q_1|)$$

Passing to the limit $i \to \infty$ and using conformal invariance of the Dirichlet energy leads to

(15)
$$\frac{1}{K} \int_{\Sigma_1} |\nabla(f^*u)|_{g_1}^2 \, dV(g_1) \le \int_{\Sigma_2} |\nabla u|_{g_2}^2 \, dV(g_2) \le K \int_{\Sigma_1} |\nabla(f^*u)|_{g_1}^2 \, dV(g_1).$$

Let $h_2 \in [g_2]$ be a smooth metric on Σ_2 . Then $\mu = (f^{-1})_* v_{h_2}$ defines a measure on Σ_1 such that

$$\int_{\Sigma_1} f^* u \, d\mu = \int_{\Sigma_2} u \, dV(h_2).$$

In particular, $vol(\Sigma_1, \mu) = vol(\Sigma_2, h_2)$.

Putting these bounds together, we obtain that

$$\frac{1}{K}R_{[g_1]}(f^*u,\mu) \le R_{[h_2]}(u) \le KR_{[g_1]}(f^*u,\mu).$$

However, since the Teichmüller mapping f is not smooth at zeroes of q_1 , the measure μ is not a volume measure of a smooth Riemannian metric. In the last step of this proof we show that there exists a sequence of metrics $\rho_n \in [g_1]$ such that $\lambda_1(\rho_n) \to \lambda_1([g_1], \mu)$. In order to do that we first obtain a local expression for μ close to the singular points.

Let s be a zero of q_1 and let z_i , i=1,2 be local complex coordinates in the neighborhood of s and f(s) respectively such that $q_i=z_i^kdz_i^2$. In cones with vertices at s and f(s) respectively one can introduce the coordinates $\zeta_i=\frac{k+2}{2}\int_0^{z_i}\sqrt{q_i}=z_i^{\frac{k+2}{2}}$.

Then in coordinates ζ_i the mapping f is linear, i.e. in appropriately chosen cones the mapping f in z_i -coordinates takes form

(16)
$$f(z_1) = \left(\tilde{f}(z_1^{\frac{k+2}{2}})\right)^{\frac{2}{k+2}},$$

where \tilde{f} is linear and the branch of the root function is chosen so that in the coordinate cone $z_i = \zeta_i^{\frac{2}{k+2}}$.

Now suppose that $dV(h_2) = \alpha(z)dz_2d\bar{z}_2$. Then using (16) one obtains

$$d\mu = \alpha(f(z_1))df(z_1)d\overline{f(z_1)}$$

$$=\alpha(f(z_1))\left|z_1^{-\frac{k+2}{2}}\tilde{f}(z_1^{\frac{k+2}{2}})\right|^{-\frac{k}{k+2}}\left(|\tilde{f}_z(z_1^{\frac{k+2}{2}})|^2-|\tilde{f}_{\bar{z}}(z_1^{\frac{k+2}{2}})|^2\right)dz_1d\bar{z}_1.$$

As \tilde{f} is linear, we conclude that $d\mu = \beta dz_1 d\bar{z}_1$ where $\beta \in L^{\infty}(\Sigma)$. At this point an appropriate approximation can be constructed using the following lemma and a standard regularization procedure.

Lemma 4.5. Let g be a Riemannian metric on a surface Σ . Suppose that $\{\rho_{\varepsilon}\}\subset L^{\infty}(\Sigma)$ is an equibounded sequence such that $\rho_{\varepsilon}\to\rho$ as $\varepsilon\to 0$ dV(g)-a.e. Then one has for every k>0

$$\lambda_k([g], \rho_{\varepsilon} dV(g)) \to \lambda_k([g], \rho dV(g)).$$

Therefore, taking the supremum over all h_2 yields

$$\Lambda_k(\Sigma, [g_2]) \le K\Lambda_k(\Sigma, [g_1]).$$

Switching the role of Σ_1 and Σ_2 and considering f^{-1} instead of f in the previous argument completes the proof in the orientable case.

The proof in the non-orientable case is easily reduced to the orientable case using the following construction. For any metric g on Σ the metric π^*g on $\hat{\Sigma}$ is σ -invariant. Thus, its eigenvalues are split into σ -even and σ -odd. Moreover, σ -even eigenvalues coincide with eigenvalues of (Σ, g) . Since the Teichmüller map in this case preserves σ -even functions, one can repeat the proof of the orientable case, restricting oneself to even eigenvalues. This completes the proof, modulo the proof of the lemma. \square

Proof of Lemma 4.5. The proof of this lemma follows the proof of a similar statement for Steklov eigenvalues found in Lemma 3.1 of [Kar18]. For completeness, we provide the proof. First, we observe there is a constant C > 0 that does not depend on ϵ such that $\lambda_k([g], \rho_{\epsilon} dV(g)) \leq Ck$. This follows from the Theorem A_k on the top of page 7 of [Kok14]. Moreover, by Proposition 1.1 of [Kok14] we also have

$$\limsup \lambda_k([g], \rho_{\epsilon} dV(g)) \le \lambda_k([g], \rho dV(g)).$$

Thus, it suffices to prove that $\lambda_k([g], \rho dV(g)) \leq \liminf \lambda_k([g], \rho_{\epsilon} dV(g))$. Let u_{ϵ} be an eigenfunction corresponding to $\lambda_k([g], \rho_{\epsilon} dV(g))$ normalized so that $\|u_{\epsilon}\|_{L^2(\rho_{\epsilon} dV(g))} = 1$. We will show that the $L^2(dV(g))$ and $H^1(\Sigma, dV(g))$ -norms of the u_{ϵ} are bounded uniformly in ϵ , for $\epsilon > 0$ sufficiently small. We recall the following proposition:

22

Proposition 4.6. ([AH96] Lemma 8.3.1) Let (M, g) be a Riemannian manifold. Then there exists a constant C > 0 such that for all $L \in H^{-1}(M)$ with L(1) = 1 one has

$$||u - L(u)||_{L^2(M)} \le C||L||_{H^{-1}(M)} \left(\int_M |\nabla u|_g^2 dV_g\right)^{1/2}$$

for all $u \in H^1(M)$.

We will apply Proposition 4.6 with $L_{\epsilon}(u) = \int_{\Sigma} u \rho_{\epsilon} \, dV(g)$. First, we compute the norm of L_{ϵ} . We have:

$$\left| \int_{\Sigma} u \rho_{\epsilon} dV(g) \right| \leq C \int_{\Sigma} |u| dV(g) \leq C ||u||_{L^{2}(dV(g))} \leq C ||u||_{H^{1}(\Sigma,g)},$$

where we used in order the uniform boundedness of ρ_{ϵ} , the Cauchy-Schwarz inequality, and the compact embedding of $H^1(\Sigma, g)$ into $L^2(\mathrm{dV}(g))$. Thus, the family of operators L_{ϵ} are uniformly bounded in $H^{-1}(\Sigma)$. Applying Proposition 4.6 to L_{ϵ} and u_{ϵ} yields:

$$||u_{\epsilon}||_{L^{2}(dV(g))} \leq C \left(\int_{\Sigma} |\nabla u_{\epsilon}|^{2} dV(g) \right)^{1/2} = C \sqrt{\lambda_{k}([g], \rho_{\epsilon} dV(g))},$$

since $L_{\epsilon}(u_{\epsilon}) = 0$. Thus, we see that u_{ϵ} are uniformly bounded in $H^{1}(\Sigma, dV(g))$ and $L^{2}(dV(g))$.

We will now show that the family u_{ϵ} is uniformly bounded with respect to ϵ . Indeed, each u_{ϵ} satisfies $\Delta_g u_{\epsilon} = \lambda_k([g], \rho_{\epsilon} dV(g)) \rho_{\epsilon} u_{\epsilon}$ in a weak sense. Since Δ_g is a second order elliptic differential operator by the Sobolev Embedding Theorem and elliptic regularity we have:

$$||u_{\epsilon}||_{\infty} \leq C||u_{\epsilon}||_{H^{2}(dV(g))} \leq C(||u_{\epsilon}||_{L^{2}(dV(g))} + ||\lambda_{k}([g], \rho_{\epsilon}dV(g))\rho_{\epsilon}u_{\epsilon}||_{L^{2}(dV(g))})$$

$$\leq C(1 + \lambda_{k}([g], \rho_{\epsilon}dV(g))),$$

where the last inequality comes from the fact that the L^2 -norms of the u_{ϵ} and the L^{∞} -norms of ρ_{ϵ} are uniformly bounded. The claim follows since the eigenvalues are uniformly bounded.

Now we show that if u_{ϵ} and v_{ϵ} are $\rho_{\epsilon} dV(g)$ -orthogonal eigenfunctions then:

$$\int_{\Sigma} u_{\epsilon}^{2} \rho \, dV(g) \to 1 \text{ and } \int_{\Sigma} u_{\epsilon} v_{\epsilon} \rho \, dV(g) \to 0.$$

Since $||u_{\epsilon}||_{L^2(\rho_{\epsilon}dV(g))} = 1$ we have:

$$\left| \int_{\Sigma} u_{\epsilon}^2 \rho \, dV(g) - 1 \right| \le \int_{\Sigma} |u_{\epsilon}|^2 |\rho - \rho_{\epsilon}| \, dV(g).$$

Since the u_{ϵ} are uniformly bounded, the first claim follows. Since $\int_{\Sigma} u_{\epsilon} v_{\epsilon} \rho_{\epsilon} \, dV(g) = 0$ a similar argument shows that $\int_{\Sigma} u_{\epsilon} v_{\epsilon} \rho \, dV(g) \to 0$.

Finally, let $E_{k+1}(\epsilon)$ be a direct sum of the first k eigenspaces for $([g], \rho_{\epsilon} dV(g))$ with ρ_{ϵ} -orthonormal basis given by $\{u_{\epsilon}^{i}\}_{i=1}^{k+1}$. Any function in $E_{k+1}(\epsilon)$ can be written as

 $\sum_{i=1}^{k+1} c_i u_{\epsilon}^i$. Plugging this into the Rayleigh quotient yields:

$$\begin{split} \lambda_k([g], \rho \mathrm{dV}(g)) &\leq \frac{\int_{\Sigma} |\sum_i c_i \nabla u_{\epsilon}^i|^2 \, \mathrm{dV}(g)}{\int_{\Sigma} (\sum_i c_i u_{\epsilon}^i)^2 \rho \, \mathrm{dV}(g)} \\ &= C_{\epsilon} \frac{\sum_i c_i^2 \int_{\Sigma} |\nabla u_{\epsilon}^i|^2 \, \mathrm{dV}(g)}{\sum_i c_i^2 \int_{\Sigma} (u_{\epsilon}^i)^2 \rho \, \mathrm{dV}(g)} \\ &\leq C_{\epsilon} \max_i \frac{\int_{\Sigma} |\nabla u_{\epsilon}^i|^2 \, \mathrm{dV}(g)}{\int_{\Sigma} (u_{\epsilon}^i)^2 \rho \, \mathrm{dV}(g)}, \end{split}$$

where

$$C_{\epsilon} = \frac{\sum_{i} c_{i}^{2} \int_{\Sigma} (u_{\epsilon}^{i})^{2} \rho \, dV(g)}{\sum_{i} c_{i}^{2} \int_{\Sigma} (u_{\epsilon}^{i})^{2} \rho \, dV(g) + \sum_{i \leq j} 2c_{i}c_{j} \int_{\Sigma} u_{\epsilon}^{i} u_{\epsilon}^{i} \rho \, dV(g)}$$

and in the last inequality we made use of the inequality $\frac{x_1+x_2}{y_1+y_2} \leq \max\left(\frac{x_1}{y_1}, \frac{x_2}{y_2}\right)$, for positive real numbers x_1, x_2, y_1 and y_2 . By our previous observations $C_{\epsilon} \to 1$ while the numerator of the last term in the inequality is $\lambda_k([g], \rho_{\epsilon} dV(g))$ and the denominator goes to one. Thus, we have $\lambda_k([g], \rho dV(g)) \leq \liminf \lambda_k([g], \rho_{\epsilon} dV(g))$, which completes the proof.

4.4. **Proof of Theorem 1.5.** The proof contains two short steps. First, we prove that the values $\Lambda_1(\mathbb{T}^2)$ and $\Lambda_1(\mathbb{KL})$ are achieved by metrics smooth away from finitely many conical singularities. Second, we apply Theorem 1.4 to prove that these metrics cannot have conical points, i.e. they must be smooth everywhere.

Proof. **Step 1.** As we discussed in Sections 4.1 and 4.2 the space of conformal classes \mathbb{T}^2 and \mathbb{KL} can be identified with subsets of \mathbb{R} and \mathbb{C} respectively. Moreover, the induced topology coincides with the topology generated by Tecihmüller distance, see [FM12].

Let Σ denote either \mathbb{T}^2 or \mathbb{KL} and $\{g_n\}$ be a sequence of metrics on Σ such that $\lim \bar{\lambda}_1(\Sigma, g_n) \to \Lambda_1(\Sigma)$. From [Nad96] and [JNP06] we know that:

$$\Lambda_1(\mathbb{T}^2) > 8\pi$$

and

$$\Lambda_1(\mathbb{KL}) \ge 12\pi E(2\sqrt{2}/3) > 12\pi.$$

Therefore, by Theorems 4.1, 4.2 the conformal classes $[g_n]$ belong to a compact subset of the space of conformal classes. Thus, the sequence $\{[g_n]\}$ has a limit point [g]. By Proposition 4.4 one has $\Lambda_1(\Sigma, [g]) = \Lambda_1(\Sigma)$. It was proved by Petrides [Pet14] that for any conformal class [h] there exists a metric $\tilde{h} \in [h]$, smooth except maybe at a finite number of singular points corresponding to conical singularities, such that $\Lambda_1(\Sigma, [h]) = \bar{\lambda}_1(\Sigma, \tilde{h})$. We conclude that there exists a metric $\tilde{g} \in [g]$ such that $\Lambda_1(\Sigma) = \bar{\lambda}_1(\Sigma, \tilde{g})$. Moreover, by Theorem 1.3 and Remark 1.3 \tilde{g} is induced from a (possibly branched) minimal immersion by first eigenfunctions Φ of Σ into a round sphere of dimension at least three.

Step 2. Suppose that \tilde{g} has a conical point. From Theorem 1.3 it follows that this metric is induced from a branched minimal isometric immersion into a round sphere. As it was observed in sections 4.1 and 4.2, on Σ there exist natural \mathbb{S}^1 -actions s_{θ} by conformal diffeomorphisms without fixed points. Then the mapping $\Phi \circ s_{\theta}$ is again a branched minimal immersion. The metric induced by this immersion is $s_{\theta}^* \tilde{g}$. Therefore, since the components of Φ are the first eigenfunctions of (Σ, \tilde{g}) , then the components of $s_{\theta}^* \Phi = \Phi \circ s_{\theta}$ are the first eigenfunctions of $(\Sigma, s_{\theta}^* \tilde{g})$. By Theorem 1.4 the metrics $s_{\theta}^* \tilde{g}$ and \tilde{g} must be equal. Thus, a $\bar{\lambda}_1$ -maximal metric must be a metric of revolution. Under this \mathbb{S}^1 -action the conical point forms a 1-dimensional singular set, which contradicts Step 1 (the set of conical points of Λ_1 -maximal metric is at most finite). This completes the proof of the theorem.

References

- [AH96] David R. Adams and Lars Inge Hedberg. Function spaces and potential theory, volume 314 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1996.
- [Don11] Simon Donaldson. Riemann surfaces, volume 22 of Oxford Graduate Texts in Mathematics. Oxford University Press, Oxford, 2011.
- [ESGJ06] Ahmad El Soufi, Hector Giacomini, and Mustapha Jazar. A unique extremal metric for the least eigenvalue of the Laplacian on the Klein bottle. *Duke Math. J.*, 135(1):181–202, 2006
- [ESI86] Ahmad El Soufi and Saïd Ilias. Immersions minimales, première valeur propre du Laplacien et volume conforme. *Math. Ann.*, 275(2):257–267, 1986.
- [ESI00] Ahmad El Soufi and Saïd Ilias. Riemannian manifolds admitting isometric immersions by their first eigenfunctions. *Pacific Journal of Mathematics*, 195(1):91–99, 2000.
- [ESI08] Ahmad El Soufi and Saïd Ilias. Laplacian eigenvalue functionals and metric deformations on compact manifolds. J. Geom. Phys., 58(1):89–104, 2008.
- [FM12] Benson Farb and Dan Margalit. A primer on mapping class groups, volume 49 of Princeton Mathematical Series. Princeton University Press, Princeton, NJ, 2012.
- [Gir09] Alexandre Girouard. Fundamental tone, concentration of density, and conformal degeneration on surfaces. *Canad. J. Math.*, 61(3):548–565, 2009.
- [GOR73] R. D. Gulliver, II, R. Osserman, and H. L. Royden. A theory of branched immersions of surfaces. *Amer. J. Math.*, 95:750–812, 1973.
- [GT01] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [Her70] Joseph Hersch. Quatre propriétés isopérimétriques de membranes sphériques homogènes. C. R. Acad. Sci. Paris Sér. A-B, 270:A1645–A1648, 1970.
- [JLN⁺05] Dmitry Jakobson, Michael Levitin, Nikolai Nadirashvili, Nilima Nigam, and Iosif Polterovich. How large can the first eigenvalue be on a surface of genus two? *Int. Math. Res. Not.*, (63):3967–3985, 2005.
- [JNP06] Dmitry Jakobson, Nikolai Nadirashvili, and Iosif Polterovich. Extremal metric for the first eigenvalue on a Klein bottle. *Canad. J. Math.*, 58(2):381–400, 2006.
- [Jos02] Jürgen Jost. Compact Riemann surfaces. Springer-Verlag, Berlin, second edition, 2002. An introduction to contemporary mathematics.
- [Kar13] Mikhail A. Karpukhin. Nonmaximality of known extremal metrics on torus and Klein bottle. *Sbornik: Mathematics*, 204(12):1728, 2013.
- [Kar14] Mikhail A. Karpukhin. Spectral properties of bipolar surfaces to Otsuki tori. *J. Spectr. Theory*, 4(1):87–111, 2014.

- [Kar15] Mikhail A. Karpukhin. Spectral properties of a family of minimal tori of revolution in five-dimensional sphere. *Canad. Math. Bull.*, 58(2):285–296, 2015.
- [Kar16] Mikhail A. Karpukhin. Upper bounds for the first eigenvalue of the Laplacian on non-orientable surfaces. Int. Math. Res. Not. IMRN, (20):6200-6209, 2016.
- [Kar18] Mikhail A. Karpukhin. Maximal metrics for the first Steklov eigenvalue on surfaces. arXiv:1801.06914, January 2018.
- [Kok14] Gerasim Kokarev. Variational aspects of Laplace eigenvalues on Riemannian surfaces. *Adv. Math.*, 258:191–239, 2014.
- [Kok17] Gerasim Kokarev. Conformal volume and eigenvalue problems. arXiv:1712.08150, December 2017.
- [Lap08] Hugues Lapointe. Spectral properties of bipolar minimal surfaces in \mathbb{S}^4 . Differential Geometry and its Applications, 26(1):9–22, 2008.
- [LY82] Peter Li and Shing Tung Yau. A new conformal invariant and its applications to the Will-more conjecture and the first eigenvalue of compact surfaces. *Invent. Math.*, 69(2):269–291, 1982.
- [MR86] Sebastián Montiel and Antonio Ros. Minimal immersions of surfaces by the first eigenfunctions and conformal area. *Invent. Math.*, 83(1):153–166, 1986.
- [MR91] Sebastián Montiel and Antonio Ros. Schrödinger operators associated to a holomorphic map. In *Global differential geometry and global analysis (Berlin, 1990)*, volume 1481 of *Lecture Notes in Math.*, pages 147–174. Springer, Berlin, 1991.
- [MS17] Henrik Matthiesen and Anna Siffert. Existence of metrics maximizing the first eigenvalue on closed surfaces. arXiv:1703.01264, March 2017.
- [Nad96] Nikolai Nadirashvili. Berger's isoperimetric problem and minimal immersions of surfaces. Geom. Funct. Anal., 6(5):877–897, 1996.
- [NP18] Nikolai S. Nadirashvili and Alexei V. Penskoi. An isoperimetric inequality for the second non-zero eigenvalue of the Laplacian on the projective plane. Geometric and Functional Analysis, Jul 2018.
- [NS15] Nikolai Nadirashvili and Yannick Sire. Conformal spectrum and harmonic maps. *Mosc. Math. J.*, 15(1):123–140, 182, 2015.
- [NS18] Shin Nayatani and Toshihiro Shoda. Metrics on a closed surface of genus two which maximize the first eigenvalue of the Laplacian. arXiv:1704.06384, May 2018.
- [Pen12] Alexei V. Penskoi. Extremal spectral properties of Lawson tau-surfaces and the Lamé equation. *Mosc. Math. J.*, 12(1):173–192, 216, 2012.
- [Pen13a] Alexei V. Penskoi. Extremal metrics for the eigenvalues of the Laplace-Beltrami operator on surfaces. *Uspekhi Mat. Nauk*, 68(6(414)):107–168, 2013.
- [Pen13b] Alexei V Penskoi. Extremal spectral properties of Otsuki tori. *Mathematische Nachrichten*, 286(4):379–391, 2013.
- [Pen15] Alexei V Penskoi. Generalized Lawson tori and Klein bottles. *The Journal of Geometric Analysis*, 25(4):2645–2666, 2015.
- [Pet14] Romain Petrides. Existence and regularity of maximal metrics for the first Laplace eigenvalue on surfaces. *Geom. Funct. Anal.*, 24(4):1336–1376, 2014.
- [Tak66] Tsunero Takahashi. Minimal immersions of Riemannian manifolds. J. Math. Soc. Japan, 18:380–385, 1966.
- [Tro91] Marc Troyanov. Prescribing curvature on compact surfaces with conical singularities. Trans. Amer. Math. Soc., 324(2):793–821, 1991.
- [YY80] Paul C. Yang and Shing Tung Yau. Eigenvalues of the Laplacian of compact Riemann surfaces and minimal submanifolds. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4), 7(1):55–63, 1980.

University of Michigan, Department of Mathematics, 530 Church Street, Ann Arbor, MI 48109

 $E ext{-}mail\ address: cianci@umich.edu}$

University of California, Irvine, Department of Mathematics, 340 Rowland Hall, Irvine, CA 92697-3875

 $E ext{-}mail\ address: mkarpukh@uci.edu}$

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, PAVILLON ANDRÉ-AISENSTADT, UNI-VERSITÉ DE MONTRÉAL, MONTRÉAL, QC, CANADA, H3C 3J7

E-mail address: medvedevv@dms.umontreal.ca