

# Partially Optimal Cubic Subspace Clustering

Research Project Machine Learning

Volodymyr Drobitko

Technische Universität Dresden

21.07.2025

- 1 Introduction
- 2 Partial Optimality for Cubic Clique Partition Problem
- 3 Cubic Subspace Instance Construction
- 4 Experiments and Evaluation
- 5 Conclusion

# Problem Statement (1)

Finite sample set  $S$ , cost function  $c: \binom{S}{3} \rightarrow \mathbb{R}$ .

Instance of the **Cubic Clique Partition Problem**:

$$\min_{y: \binom{S}{2} \rightarrow \{0,1\}} \sum_{abc \in \binom{S}{3}} c_{abc} y_{ab} y_{bc} y_{ac}$$

subject to  $y_{ab} + y_{bc} - 1 \leq y_{ac}$  for all distinct  $a, b, c \in S$ .

# Problem Statement (1)

Finite sample set  $S$ , cost function  $c: \binom{S}{3} \rightarrow \mathbb{R}$ .

Instance of the **Cubic Clique Partition Problem**:

$$\min_{y: \binom{S}{2} \rightarrow \{0,1\}} \sum_{abc \in \binom{S}{3}} c_{abc} y_{ab} y_{bc} y_{ac}$$

subject to  $y_{ab} + y_{bc} - 1 \leq y_{ac}$  for all distinct  $a, b, c \in S$ .

Find a **partially optimal solution**, i.e. fix some labels  $y_{ab}$  for distinct  $a, b \in S$


$$\begin{cases} y_{ab} = 1 & \text{join } a, b \\ y_{ab} = 0 & \text{cut } a, b \\ y_{ab} = ? & \text{unknown} \end{cases}$$

in such way that there still exists an optimal solution.

## Problem Statement (2)

**Subspace Instances** of the Cubic Clique Partition Problem

Samples  $S$ : points  $S \subset \mathbb{R}^3$




(a) Samples  $S$

# Problem Statement (2)


## Subspace Instances of the Cubic Clique Partition Problem

Samples  $S$ : points  $S \subset \mathbb{R}^3$

Point generation: 3 distinct planes containing the origin, noise  $\sigma$



(a) Samples  $S$



(b) Optimal labeling  $y^*$


# Problem Statement (2)

## Subspace Instances of the Cubic Clique Partition Problem


Samples  $S$ : points  $S \subset \mathbb{R}^3$

Point generation: 3 distinct planes containing the origin, noise  $\sigma$

Optimal labeling  $y^*$ : original planes



(a) Samples  $S$



(b) Optimal labeling  $y^*$

# Problem Statement (2)


## Subspace Instances of the Cubic Clique Partition Problem

Samples  $S$ : points  $S \subset \mathbb{R}^3$


Point generation: 3 distinct planes containing the origin, noise  $\sigma$

Optimal labeling  $y^*$ : original planes

Cost function  $c$ ? (no concrete plane information given)



(a) Samples  $S$



(b) Optimal labeling  $y^*$



## **Related Work:**

- Stein, Di Gregorio, and Andres, "Partial Optimality in Cubic Correlation Clustering"
- Lange, Karrenbauer, and Andres, "Partial Optimality and Fast Lower Bounds for Weighted Correlation Clustering"
- Lange, Andres, and Swoboda, "Combinatorial persistency criteria for multicut and max-cut"

## Related Work:

- Stein, Di Gregorio, and Andres, “Partial Optimality in Cubic Correlation Clustering”
- Lange, Karrenbauer, and Andres, “Partial Optimality and Fast Lower Bounds for Weighted Correlation Clustering”
- Lange, Andres, and Swoboda, “Combinatorial persistency criteria for multicut and max-cut”

## Tasks and Solutions:

- 1 Read Stein, Di Gregorio, and Andres, “Partial Optimality in Cubic Correlation Clustering”, implement the partial optimality algorithm  
→ implementation in C++

## Related Work:

- Stein, Di Gregorio, and Andres, “Partial Optimality in Cubic Correlation Clustering”
- Lange, Karrenbauer, and Andres, “Partial Optimality and Fast Lower Bounds for Weighted Correlation Clustering”
- Lange, Andres, and Swoboda, “Combinatorial persistency criteria for multicut and max-cut”

## Tasks and Solutions:

- 1 Read Stein, Di Gregorio, and Andres, “Partial Optimality in Cubic Correlation Clustering”, implement the partial optimality algorithm  
→ implementation in C++
- 2 Construct subspace instances of increasing difficulty  
→ point generation, appropriate **cost function c**

## Related Work:

- Stein, Di Gregorio, and Andres, “Partial Optimality in Cubic Correlation Clustering”
- Lange, Karrenbauer, and Andres, “Partial Optimality and Fast Lower Bounds for Weighted Correlation Clustering”
- Lange, Andres, and Swoboda, “Combinatorial persistency criteria for multicut and max-cut”

## Tasks and Solutions:

- 1 Read Stein, Di Gregorio, and Andres, “Partial Optimality in Cubic Correlation Clustering”, implement the partial optimality algorithm  
→ implementation in C++
- 2 Construct subspace instances of increasing difficulty  
→ point generation, appropriate **cost function c**
- 3 Apply algorithm to the subspace instances, assess partial optimality, accuracy and computation time  
→ experiments and evaluation

- 1 Introduction
- 2 Partial Optimality for Cubic Clique Partition Problem
- 3 Cubic Subspace Instance Construction
- 4 Experiments and Evaluation
- 5 Conclusion

# Partial Optimality for Cubic Clique Partition Problem

Extended cost function  $c: \binom{S}{3} \cup \binom{S}{2} \cup \emptyset \rightarrow \mathbb{R}$

# Partial Optimality for Cubic Clique Partition Problem

Extended cost function  $c: \binom{S}{3} \cup \binom{S}{2} \cup \emptyset \rightarrow \mathbb{R}$

Instance of the extended cubic clique partition problem:

$$\min_{y: \binom{S}{2} \rightarrow \{0,1\}} \sum_{abc \in \binom{S}{3}} c_{abc} y_{ab} y_{bc} y_{ac} + \sum_{ab \in \binom{S}{2}} c_{ab} y_{ab} + c_{\emptyset}$$

subject to  $y_{ab} + y_{bc} - 1 \leq y_{ac}$  for all distinct  $a, b, c \in S$ .

# Partial Optimality for Cubic Clique Partition Problem

Extended cost function  $c: \binom{S}{3} \cup \binom{S}{2} \cup \emptyset \rightarrow \mathbb{R}$

Instance of the extended cubic clique partition problem:

$$\min_{y: \binom{S}{2} \rightarrow \{0,1\}} \sum_{abc \in \binom{S}{3}} c_{abc} y_{ab} y_{bc} y_{ac} + \sum_{ab \in \binom{S}{2}} c_{ab} y_{ab} + c_{\emptyset}$$

subject to  $y_{ab} + y_{bc} - 1 \leq y_{ac}$  for all distinct  $a, b, c \in S$ .

Construct **Improving Maps** for the labeling  $y$



# Partial Optimality for Cubic Clique Partition Problem

Extended cost function  $c: \binom{S}{3} \cup \binom{S}{2} \cup \emptyset \rightarrow \mathbb{R}$

Instance of the extended cubic clique partition problem:

$$\min_{y: \binom{S}{2} \rightarrow \{0,1\}} \sum_{abc \in \binom{S}{3}} c_{abc} y_{ab} y_{bc} y_{ac} + \sum_{ab \in \binom{S}{2}} c_{ab} y_{ab} + c_{\emptyset}$$

subject to  $y_{ab} + y_{bc} - 1 \leq y_{ac}$  for all distinct  $a, b, c \in S$ .

Construct **Improving Maps** for the labeling  $y$

→ **Partial Optimality Conditions:**

# Partial Optimality for Cubic Clique Partition Problem

Extended cost function  $c: \binom{S}{3} \cup \binom{S}{2} \cup \emptyset \rightarrow \mathbb{R}$

Instance of the extended cubic clique partition problem:

$$\min_{y: \binom{S}{2} \rightarrow \{0,1\}} \sum_{abc \in \binom{S}{3}} c_{abc} y_{ab} y_{bc} y_{ac} + \sum_{ab \in \binom{S}{2}} c_{ab} y_{ab} + c_{\emptyset}$$

subject to  $y_{ab} + y_{bc} - 1 \leq y_{ac}$  for all distinct  $a, b, c \in S$ .

Construct **Improving Maps** for the labeling  $y$

→ **Partial Optimality Conditions:**

- ① Subproblem-CUT-condition (cut subset from its complement)

# Partial Optimality for Cubic Clique Partition Problem

Extended cost function  $c: \binom{S}{3} \cup \binom{S}{2} \cup \emptyset \rightarrow \mathbb{R}$

Instance of the extended cubic clique partition problem:

$$\min_{y: \binom{S}{2} \rightarrow \{0,1\}} \sum_{abc \in \binom{S}{3}} c_{abc} y_{ab} y_{bc} y_{ac} + \sum_{ab \in \binom{S}{2}} c_{ab} y_{ab} + c_{\emptyset}$$

subject to  $y_{ab} + y_{bc} - 1 \leq y_{ac}$  for all distinct  $a, b, c \in S$ .

Construct **Improving Maps** for the labeling  $y$

→ **Partial Optimality Conditions:**

- 1 Subproblem-CUT-condition (cut subset from its complement)
- 2 CUT-conditions (cut pairs and triples)

# Partial Optimality for Cubic Clique Partition Problem

Extended cost function  $c: \binom{S}{3} \cup \binom{S}{2} \cup \emptyset \rightarrow \mathbb{R}$

Instance of the extended cubic clique partition problem:

$$\min_{y: \binom{S}{2} \rightarrow \{0,1\}} \sum_{abc \in \binom{S}{3}} c_{abc} y_{ab} y_{bc} y_{ac} + \sum_{ab \in \binom{S}{2}} c_{ab} y_{ab} + c_{\emptyset}$$

subject to  $y_{ab} + y_{bc} - 1 \leq y_{ac}$  for all distinct  $a, b, c \in S$ .

Construct **Improving Maps** for the labeling  $y$

→ **Partial Optimality Conditions:**

- 1 Subproblem-CUT-condition (cut subset from its complement)
- 2 CUT-conditions (cut pairs and triples)
- 3 JOIN-conditions (join subsets, pairs and triples)

# Partial Optimality for Cubic Clique Partition Problem

Extended cost function  $c: \binom{S}{3} \cup \binom{S}{2} \cup \emptyset \rightarrow \mathbb{R}$

Instance of the extended cubic clique partition problem:

$$\min_{y: \binom{S}{2} \rightarrow \{0,1\}} \sum_{abc \in \binom{S}{3}} c_{abc} y_{ab} y_{bc} y_{ac} + \sum_{ab \in \binom{S}{2}} c_{ab} y_{ab} + c_{\emptyset}$$

subject to  $y_{ab} + y_{bc} - 1 \leq y_{ac}$  for all distinct  $a, b, c \in S$ .

Construct **Improving Maps** for the labeling  $y$

→ **Partial Optimality Conditions:**

- 1 Subproblem-CUT-condition (cut subset from its complement)
- 2 CUT-conditions (cut pairs and triples)
- 3 JOIN-conditions (join subsets, pairs and triples)

CUT-conditions can be applied simultaneously

JOIN-conditions cannot be applied simultaneously!

# Partial Optimality for Cubic Clique Partition Problem

Extended cost function  $c: \binom{S}{3} \cup \binom{S}{2} \cup \emptyset \rightarrow \mathbb{R}$

Instance of the extended cubic clique partition problem:

$$\min_{y: \binom{S}{2} \rightarrow \{0,1\}} \sum_{abc \in \binom{S}{3}} c_{abc} y_{ab} y_{bc} y_{ac} + \sum_{ab \in \binom{S}{2}} c_{ab} y_{ab} + c_{\emptyset}$$

subject to  $y_{ab} + y_{bc} - 1 \leq y_{ac}$  for all distinct  $a, b, c \in S$ .

Construct **Improving Maps** for the labeling  $y$

→ **Partial Optimality Conditions:**

- 1 Subproblem-CUT-condition (cut subset from its complement)
- 2 CUT-conditions (cut pairs and triples)
- 3 JOIN-conditions (join subsets, pairs and triples)

CUT-conditions can be applied simultaneously

JOIN-conditions cannot be applied simultaneously!

Apply partial optimality conditions → solve subproblems

# Partial Optimality Algorithm

## **Partial Optimality Algorithm:**

**Input:** labeling  $y$  without fixed labels

**while** condition applied **do**

    apply subproblem-CUT-condition exhaustively

    apply one of JOIN-conditions (in effective order)

**end while**

    apply CUT-conditions exhaustively

**Output:** partially optimal labeling  $y$  with some fixed labels

# Partial Optimality Algorithm

## Partial Optimality Algorithm:

**Input:** labeling  $y$  without fixed labels

**while** condition applied **do**

    apply subproblem-CUT-condition exhaustively

    apply one of JOIN-conditions (in effective order)

**end while**

apply CUT-conditions exhaustively

**Output:** partially optimal labeling  $y$  with some fixed labels

## Reduction to subproblems:

- 1 Subproblem-CUT-condition: fix CUT labels for element pairs from different sample subsets; solve each subset as an independent problem and accumulate the results in  $c_\emptyset$ ;



# Partial Optimality Algorithm

## Partial Optimality Algorithm:

**Input:** labeling  $y$  without fixed labels

**while** condition applied **do**

    apply subproblem-CUT-condition exhaustively

    apply one of JOIN-conditions (in effective order)

**end while**

apply CUT-conditions exhaustively

**Output:** partially optimal labeling  $y$  with some fixed labels

## Reduction to subproblems:

- 1 Subproblem-CUT-condition: fix CUT labels for element pairs from different sample subsets; solve each subset as an independent problem and accumulate the results in  $c_\emptyset$ ;
- 2 JOIN-Conditions: fix JOIN labels for elements of the sample subset; add the join-cost to  $c_\emptyset$ ; solve the problem where the subset is considered as one sample;

## Subproblem-CUT and Subset-JOIN

**Subproblem-CUT:** cut sample subsets  $R_1, R_2, \dots, R_k$  that are only connected via non-negative costs (applied if  $k > 1$ )

# Subproblem-CUT and Subset-JOIN


**Subproblem-CUT:** cut sample subsets  $R_1, R_2, \dots, R_k$  that are only connected via non-negative costs (applied if  $k > 1$ )

**Subset-JOIN:** join sample subset  $R$  with only non-positive costs if its worst bipartition joining cost is less than or equal to the reward of joining  $R$  with  $\bar{R}$  (applied if  $|R| > 1$ )  
(The worst bipartition joining cost  $\approx$  min-cut)

# Subproblem-CUT and Subset-JOIN

**Subproblem-CUT:** cut sample subsets  $R_1, R_2, \dots, R_k$  that are only connected via non-negative costs (applied if  $k > 1$ )

**Subset-JOIN:** join sample subset  $R$  with only non-positive costs if its worst bipartition joining cost is less than or equal to the reward of joining  $R$  with  $\bar{R}$  (applied if  $|R| > 1$ )  
(The worst bipartition joining cost  $\approx$  min-cut)




$$c_{\emptyset} = 0$$


# Subproblem-CUT and Subset-JOIN

**Subproblem-CUT:** cut sample subsets  $R_1, R_2, \dots, R_k$  that are only connected via non-negative costs (applied if  $k > 1$ )

**Subset-JOIN:** join sample subset  $R$  with only non-positive costs if its worst bipartition joining cost is less than or equal to the reward of joining  $R$  with  $\bar{R}$  (applied if  $|R| > 1$ )  
(The worst bipartition joining cost  $\approx$  min-cut)



$$c_{\emptyset} = 0$$




$$c_{\emptyset} = -15$$


# Subproblem-CUT and Subset-JOIN

**Subproblem-CUT:** cut sample subsets  $R_1, R_2, \dots, R_k$  that are only connected via non-negative costs (applied if  $k > 1$ )


**Subset-JOIN:** join sample subset  $R$  with only non-positive costs if its worst bipartition joining cost is less than or equal to the reward of joining  $R$  with  $\bar{R}$  (applied if  $|R| > 1$ )  
(The worst bipartition joining cost  $\approx$  min-cut)



$$c_{\emptyset} = 0$$



$$c_{\emptyset} = -15$$




$$c_{\emptyset} = -75$$


# Subproblem-CUT and Subset-JOIN

**Subproblem-CUT:** cut sample subsets  $R_1, R_2, \dots, R_k$  that are only connected via non-negative costs (applied if  $k > 1$ )


**Subset-JOIN:** join sample subset  $R$  with only non-positive costs if its worst bipartition joining cost is less than or equal to the reward of joining  $R$  with  $\bar{R}$  (applied if  $|R| > 1$ )  
(The worst bipartition joining cost  $\approx$  min-cut)



$$c_{\emptyset} = 0$$



$$c_{\emptyset} = -15$$




$$c_{\emptyset} = -75$$


# Subproblem-CUT and Subset-JOIN

**Subproblem-CUT:** cut sample subsets  $R_1, R_2, \dots, R_k$  that are only connected via non-negative costs (applied if  $k > 1$ )


**Subset-JOIN:** join sample subset  $R$  with only non-positive costs if its worst bipartition joining cost is less than or equal to the reward of joining  $R$  with  $\bar{R}$  (applied if  $|R| > 1$ )  
(The worst bipartition joining cost  $\approx$  min-cut)



$$c_{\emptyset} = 0$$



$$c_{\emptyset} = -15$$



$$c_{\emptyset} = -75$$



$$c_{\emptyset} = -76$$





## Other JOIN-conditions

**Pair-JOIN-1:** join samples  $i, j$  if their overall joining reward  $\geq$  the sum of rewards and penalties for joining some subset  $R$  with  $i \in R$  and  $\bar{R}$  with  $j \in \bar{R}$  ( $\approx$  i-j min-cut)

## Other JOIN-conditions

**Pair-JOIN-1:** join samples  $i, j$  if their overall joining reward  $\geq$  the sum of rewards and penalties for joining some subset  $R$  with  $i \in R$  and  $\bar{R}$  with  $j \in \bar{R}$  ( $\approx$  i-j min-cut)

**Pair-JOIN-2:** join samples  $i, k$  if there exists a sample triple  $ijk$  that fulfills 3 conditions ( $\approx$  i-jk min-cut,  $\approx$  ij-k min-cut, 1 explicit condition)

## Other JOIN-conditions

**Pair-JOIN-1:** join samples  $i, j$  if their overall joining reward  $\geq$  the sum of rewards and penalties for joining some subset  $R$  with  $i \in R$  and  $\bar{R}$  with  $j \in \bar{R}$  ( $\approx$  i-j min-cut)

**Pair-JOIN-2:** join samples  $i, k$  if there exists a sample triple  $ijk$  that fulfills 3 conditions ( $\approx$  i-jk min-cut,  $\approx$  ij-k min-cut, 1 explicit condition)

**Pair-JOIN-3:** join samples  $i, j$  if  $c_{ij} \leq$  the sum of reward costs for joining pairs and triples containing  $i$  or  $j$

## Other JOIN-conditions

**Pair-JOIN-1:** join samples  $i, j$  if their overall joining reward  $\geq$  the sum of rewards and penalties for joining some subset  $R$  with  $i \in R$  and  $\bar{R}$  with  $j \in \bar{R}$  ( $\approx$  i-j min-cut)

**Pair-JOIN-2:** join samples  $i, k$  if there exists a sample triple  $ijk$  that fulfills 3 conditions ( $\approx$  i-jk min-cut,  $\approx$  ij-k min-cut, 1 explicit condition)

**Pair-JOIN-3:** join samples  $i, j$  if  $c_{ij} \leq$  the sum of reward costs for joining pairs and triples containing  $i$  or  $j$

**Pair-JOIN-4:** join samples  $i, k$  if there exists a sample triple  $ijk$  such that 7 explicit conditions hold

## Other JOIN-conditions

**Pair-JOIN-1:** join samples  $i, j$  if their overall joining reward  $\geq$  the sum of rewards and penalties for joining some subset  $R$  with  $i \in R$  and  $\bar{R}$  with  $j \in \bar{R}$  ( $\approx$  i-j min-cut)

**Pair-JOIN-2:** join samples  $i, k$  if there exists a sample triple  $ijk$  that fulfills 3 conditions ( $\approx$  i-jk min-cut,  $\approx$  ij-k min-cut, 1 explicit condition)

**Pair-JOIN-3:** join samples  $i, j$  if  $c_{ij} \leq$  the sum of reward costs for joining pairs and triples containing  $i$  or  $j$


**Pair-JOIN-4:** join samples  $i, k$  if there exists a sample triple  $ijk$  such that 7 explicit conditions hold

**Triple-JOIN:** join samples  $i, j, k$  if the condition holds (similar to Pair-JOIN-1) ( $\approx$  i-jk min-cut)

# Pyramid Instance and CUT-conditions

$$c_{bcd} = 10$$



$$c_{abc} = c_{abd} = c_{acd} = -50$$



# Pyramid Instance and CUT-conditions

$$c_{bcd} = 10$$

$$c_{abc} = c_{abd} = c_{acd} = -50$$




Pair-JOIN-2


# Pyramid Instance and CUT-conditions

$$c_{bcd} = 10$$

$$c_{abc} = c_{abd} = c_{acd} = -50$$



Pair-JOIN-2



Subset-JOIN




$$c_{\emptyset} = -140$$




# Pyramid Instance and CUT-conditions

$$c_{bcd} = 10$$

$$c_{abc} = c_{abd} = c_{acd} = -50$$



Pair-JOIN-2



Subset-JOIN




$$c_{\emptyset} = -140$$

**Pair-CUT:** cut samples  $i, j$  if the direct joining penalty  $\geq$  the sum of rewards for joining some subset  $R$  with  $i \in R$  and  $\bar{R}$  with  $j \in \bar{R}$  ( $\approx$  i-j min-cut)


# Pyramid Instance and CUT-conditions

$$c_{bcd} = 10$$

$$c_{abc} = c_{abd} = c_{acd} = -50$$



Pair-JOIN-2



Subset-JOIN



$$c_{\emptyset} = -140$$


**Pair-CUT:** cut samples  $i, j$  if the direct joining penalty  $\geq$  the sum of rewards for joining some subset  $R$  with  $i \in R$  and  $\bar{R}$  with  $j \in \bar{R}$  ( $\approx$  i-j min-cut)

**Triple-CUT:** cut samples  $i, j, k$  if the condition holds (similar to Pair-CUT) ( $\approx$  i-jk min-cut)


# Pyramid Instance and CUT-conditions

$$c_{bcd} = 10$$

$$c_{abc} = c_{abd} = c_{acd} = -50$$



Pair-JOIN-2



Subset-JOIN



$$c_{\emptyset} = -140$$


**Pair-CUT:** cut samples  $i, j$  if the direct joining penalty  $\geq$  the sum of rewards for joining some subset  $R$  with  $i \in R$  and  $\bar{R}$  with  $j \in \bar{R}$  ( $\approx$  i-j min-cut)

**Triple-CUT:** cut samples  $i, j, k$  if the condition holds (similar to Pair-CUT) ( $\approx$  i-jk min-cut)

Samples in the pyramid with  $c_{bcd} = 100$  are unjoinable!

Triple-CUT is applied to the triple  $bcd$

# Program Structure




- 1 Introduction
- 2 Partial Optimality for Cubic Clique Partition Problem
- 3 Cubic Subspace Instance Construction
- 4 Experiments and Evaluation
- 5 Conclusion

# Plane and Point Generation

## Plane Generation:


- generate 3 planes  
as distinct normal vectors  
 $\vec{n}_1, \vec{n}_2, \vec{n}_3$  (normalized)



# Plane and Point Generation

## Plane Generation:


- generate 3 planes as distinct normal vectors  $\vec{n}_1, \vec{n}_2, \vec{n}_3$  (normalized)
- compute a  $\vec{r}_{i,1}$  (normalized) orthogonal to  $\vec{n}_i$  ( $i \in \{1, 2, 3\}$ )



# Plane and Point Generation

## Plane Generation:

- generate 3 planes as distinct normal vectors  $\vec{n}_1, \vec{n}_2, \vec{n}_3$  (normalized)
- compute a  $\vec{r}_{i,1}$  (normalized) orthogonal to  $\vec{n}_i$  ( $i \in \{1, 2, 3\}$ )
- compute the  $\vec{r}_{i,2}$  (normalized) orthogonal to  $\vec{n}_i$  and  $\vec{r}_{i,1}$






# Plane and Point Generation

## Plane Generation:

- generate 3 planes as distinct normal vectors  $\vec{n}_1, \vec{n}_2, \vec{n}_3$  (normalized)
- compute a  $\vec{r}_{i,1}$  (normalized) orthogonal to  $\vec{n}_i$  ( $i \in \{1, 2, 3\}$ )
- compute the  $\vec{r}_{i,2}$  (normalized) orthogonal to  $\vec{n}_i$  and  $\vec{r}_{i,1}$



**Point Generation** on the plane  $(\vec{n}, \vec{r}_1, \vec{r}_2)$ , parameters  $(D, \sigma)$ :

- random variables  $k_1, k_2 \in [-D, D]$  (uniform distribution)
- random variable  $k_n$  (normal distribution based on  $\sigma$ )
- generate point  $p = k_1 \vec{r}_1 + k_2 \vec{r}_2 + k_n \vec{n}$



# Cost Function (1)

**Triangle**  $abc \in \binom{S}{3}$

# Cost Function (1)

**Triangle**  $abc \in \binom{S}{3}$


- (1) Smallest side  $s < D/2$   
 $\rightarrow c_{abc} = 0$



# Cost Function (1)

**Triangle**  $abc \in \binom{S}{3}$

- (1) Smallest side  $s < D/2$   
 $\rightarrow c_{abc} = 0$
- (2) Largest angle  $\alpha > 150^\circ$   
 $\rightarrow c_{abc} = 0$



# Cost Function (1)

**Triangle**  $abc \in \binom{S}{3}$

(1) Smallest side  $s < D/2$

$$\rightarrow c_{abc} = 0$$


(2) Largest angle  $\alpha > 150^\circ$

$$\rightarrow c_{abc} = 0$$

(3)  $h_a, h_b, h_c$ : distances to the best fitted plane containing the origin;

$$h_a + h_b + h_c > 3\sigma + 10^{-6}$$


$$\rightarrow c_{abc} = \frac{(h_a + h_b + h_c) - (3\sigma + 10^{-6})}{3D}$$



## Cost Function (1)


**Triangle**  $abc \in \binom{S}{3}$

- (1) Smallest side  $s < D/2$   
 $\rightarrow c_{abc} = 0$
- (2) Largest angle  $\alpha > 150^\circ$   
 $\rightarrow c_{abc} = 0$
- (3)  $h_a, h_b, h_c$ : distances  
to the best fitted plane  
containing the origin;  
 $h_a + h_b + h_c > 3\sigma + 10^{-6}$   
 $\rightarrow c_{abc} = \frac{(h_a + h_b + h_c) - (3\sigma + 10^{-6})}{3D}$
- (4)  $h_o$ : distance from the origin  
to the triangle plane;  
 $h_o > \frac{10}{\#points}\sigma + 10^{-6}$   
 $\rightarrow c_{abc} = 0$




## Cost Function (2)

- (5) for all points  $p$ :  
 $h_p$ : distance to the best fitted plane containing the origin;



## Cost Function (2)

- (5) for all points  $p$ :  
 $h_p$ : distance to the best fitted  
plane containing the origin;  
choose  $p$  if  
 $h_p < \sigma + 10^{-6} \wedge |\vec{p}| > 0.3D$





## Cost Function (2)



(5) for all points  $p$ :

$h_p$ : distance to the best fitted plane containing the origin;

choose  $p$  if

$$h_p < \sigma + 10^{-6} \wedge |\vec{p}| > 0.3D$$

$h'_p$ : distance to the best fitted plane to all chosen points  $p$  containing the origin;



## Cost Function (2)

(5) for all points  $p$ :

$h_p$ : distance to the best fitted plane containing the origin;



choose  $p$  if

$$h_p < \sigma + 10^{-6} \wedge |\vec{p}| > 0.3D$$

$h'_p$ : distance to the best fitted plane to all chosen points  $p$  containing the origin;

$$\delta_p := \frac{hp' - (\sigma + 10^{-6})}{D}$$

$$M := \{p \mid p \text{ chosen} \wedge \delta_p < 0\}$$



# Cost Function (2)

(5) for all points  $p$ :

$h_p$ : distance to the best fitted plane containing the origin;

choose  $p$  if



$$h_p < \sigma + 10^{-6} \wedge |\vec{p}| > 0.3D$$

$h'_p$ : distance to the best fitted plane to all chosen points  $p$  containing the origin;

$$\delta_p := \frac{hp' - (\sigma + 10^{-6})}{D}$$

$$M := \{p \mid p \text{ chosen} \wedge \delta_p < 0\}$$

$$|M| \leq 3 \rightarrow c_{abc} = 0$$



# Cost Function (2)

(5) for all points  $p$ :

$h_p$ : distance to the best fitted plane containing the origin;

choose  $p$  if

$$h_p < \sigma + 10^{-6} \wedge |\vec{p}| > 0.3D$$



$h'_p$ : distance to the best fitted plane to all chosen points  $p$  containing the origin;

$$\delta_p := \frac{h'_p - (\sigma + 10^{-6})}{D}$$

$$M := \{p \mid p \text{ chosen} \wedge \delta_p < 0\}$$


$$|M| \leq 3 \rightarrow c_{abc} = 0$$

$$\rightarrow c_{abc} = 2^{|M|-4} \cdot \sum_{p \in M} \delta_p$$





- 1 Introduction
- 2 Partial Optimality for Cubic Clique Partition Problem
- 3 Cubic Subspace Instance Construction
- 4 Experiments and Evaluation
- 5 Conclusion

# Cost Function Evaluation (3x15 points, $\sigma = 1$ )



# Cost Function Evaluation (3x15 points, $\sigma = 1$ )



# Experiments

- WSL2 Ubuntu, Intel Core i7-11370H (3.30 GHz), 16 GB RAM



# Experiments

- WSL2 Ubuntu, Intel Core i7-11370H (3.30 GHz), 16 GB RAM
- Apply the algorithm to the random cubic subspace instances with  $D = 100$  and fixed  $\sigma = 0, 1, 2, 3, 4, 5$ :
  - 3x7 points (solve 15 instances)
  - 3x10 points (solve 15 instances)
  - 3x15 points (solve 7 instances)
  - 3x20 points (solve 1 instance)


# Experiments

- WSL2 Ubuntu, Intel Core i7-11370H (3.30 GHz), 16 GB RAM
- Apply the algorithm to the random cubic subspace instances with  $D = 100$  and fixed  $\sigma = 0, 1, 2, 3, 4, 5$ :
  - 3x7 points (solve 15 instances)
  - 3x10 points (solve 15 instances)
  - 3x15 points (solve 7 instances)
  - 3x20 points (solve 1 instance)
- Track
  - computation time (s)
  - partial optimality (%)
  - accuracy (%) with respect to the truth (correct labeling)


# Experiments

- WSL2 Ubuntu, Intel Core i7-11370H (3.30 GHz), 16 GB RAM
- Apply the algorithm to the random cubic subspace instances with  $D = 100$  and fixed  $\sigma = 0, 1, 2, 3, 4, 5$ :
  - 3x7 points (solve 15 instances)
  - 3x10 points (solve 15 instances)
  - 3x15 points (solve 7 instances)
  - 3x20 points (solve 1 instance)
- Track
  - computation time (s)
  - partial optimality (%)
  - accuracy (%) with respect to the truth (correct labeling)
- Capture
  - 1.quartile (Q1)
  - median (Q2)
  - 3.quartile(Q3)
  - the worst computation time


## Partial Optimality (3x7 points)




## Accuracy (3x7 points)




## Partial Optimality (3x10 points)



## Accuracy (3x10 points)




## Partial Optimality (3x15 points)







## Accuracy (3x15 points)




# Partial Optimality



# Partial Optimality



# Computation Time (worst case)



- 1 Introduction
- 2 Partial Optimality for Cubic Clique Partition Problem
- 3 Cubic Subspace Instance Construction
- 4 Experiments and Evaluation
- 5 Conclusion

- ① Implementation of the partial optimality algorithm:
  - arbitrary sample type
  - sparse cost representation
  - pair labeling and triple cuts
  - reasonable adjustments of the partial optimality conditions
  - self-explaining logs

# Conclusion

- ① Implementation of the partial optimality algorithm:
  - arbitrary sample type
  - sparse cost representation
  - pair labeling and triple cuts
  - reasonable adjustments of the partial optimality conditions
  - self-explaining logs
- ② Subspace instance generation using linear algebra methods, geometric cost function  $c$ :
  - high accuracy (over 75%)
  - significant noise tolerance ( $\sigma \geq 1$ )
  - $O(k \cdot n^6)$  for  $n = \# \text{points}$  and a small  $k$

# Conclusion

- ① Implementation of the partial optimality algorithm:
  - arbitrary sample type
  - sparse cost representation
  - pair labeling and triple cuts
  - reasonable adjustments of the partial optimality conditions
  - self-explaining logs
- ② Subspace instance generation using linear algebra methods, geometric cost function  $c$ :
  - high accuracy (over 75%)
  - significant noise tolerance ( $\sigma \geq 1$ )
  - $O(k \cdot n^6)$  for  $n = \# \text{points}$  and a small  $k$

## Future Work:




- optimize and parallelize the algorithm
- overcome the partial optimality loss for  $c$
- determine better parameters for  $c$
- update  $c$  with advanced criteria



## My program, scripts and presentation:

<https://github.com/Vovsanka/ResearchProjectML>

## Bibliography:

-  Lange, Jan-Hendrik, Bjoern Andres, and Paul Swoboda. “Combinatorial persistency criteria for multicut and max-cut”. In: *CVPR* (2019).
-  Lange, Jan-Hendrik, Andreas Karrenbauer, and Bjoern Andres. “Partial Optimality and Fast Lower Bounds for Weighted Correlation Clustering”. In: *ICML* (2018).
-  Stein, David, Silvia Di Gregorio, and Bjoern Andres. “Partial Optimality in Cubic Correlation Clustering”. In: *ICML* (2023).