

PHY 4210-01 Senior Lab
Lab N4: Rutherford Scattering

Sarah Arends
Jacquelyne Miksanek
Ryan Wojtyla

Instructor: Dr. Marcus Hohlmann

March 14, 2019

Abstract

Contents

1	Objective of the Experiment	3
2	Theory of the Experiment	3
3	Equipment Utilized	4
4	Procedure	4
4.1	Procedural Modifications	4
5	Data Analysis	4
5.1	Data Analysis I: Gold	4
5.2	Data Analysis II: Aluminum	4
6	Results	5
6.1	Results I: Gold	5
6.2	Results II: Aluminum	5
7	Conclusion	5
8	Appendices	5
8.1	Appendix A: Data	5
8.2	Appendix B: Source Code	5

1 Objective of the Experiment

2 Theory of the Experiment

When an alpha particle with impact parameter b approaches a nucleus, it is scattered at an angle θ . If the impact parameter is given an infinitesimal range of $[b, b + db]$, the resulting scattering angle then has a range of $[\theta - d\theta, \theta]$; the impact parameter and scattering angle are inversely proportional.

Because the alpha particle can be incident within a defined range at any angle relative to the nucleus, a ring of possible incident locations is created in front of the nucleus. This ring is illustrated in Figure 1.




Figure 1: The ring whose area represents the possible region in which alpha particles may be incident on a target nucleus.

The area of this ring is found as the area of any ring is found:

$$\begin{aligned} A &= \pi(b + db)^2 - \pi b^2 \\ &= \pi(b^2 + 2bdb + db^2) - \pi b^2 \\ &= \pi b^2 + 2\pi bdb + \pi db^2 - \pi b^2 \\ &= 2\pi bdb + \pi db^2 \end{aligned}$$

Since db is infinitesimally small, it can be approximated to be zero. Therefore, the area of the incident ring, $\Delta\sigma$, is

$$\Delta\sigma = 2\pi bdb$$

Since $\Delta\sigma$ is the area of the full incident ring, an infinitesimal angular portion of that ring, $d\phi$, may be represented by $\Delta\sigma(\phi) = bdbd\phi$. Additionally, because the impact parameter b is related to the scattering angle θ , $\Delta\sigma$ is also a function of θ . Hence,

$$\Delta\sigma(\theta, \phi) = bdbd\phi$$

Since the impact parameter b is directly proportional to the size of the cross section and the scattering angle θ is inversely proportional to the impact parameter, the size of the cross section decreases as the scattering angle increases. Therefore, the cross section experiences a negative rate of change as θ increases. Hence,

$$\Delta\sigma(\theta, \phi) = -d\sigma(\theta, \phi) \tag{1}$$

3 Equipment Utilized

- List equipment and specifications

Figure 2: Description of schematic here

4 Procedure

4.1 Procedural Modifications

5 Data Analysis

5.1 Data Analysis I: Gold

5.2 Data Analysis II: Aluminum

6 Results

6.1 Results I: Gold

6.2 Results II: Aluminum

7 Conclusion

8 Appendices

8.1 Appendix A: Data

8.2 Appendix B: Source Code