

The role of letrozole in in vitro fertilization treatment: new remedy or old mirage?

Nathalie Bülow, M.D., Ph.D.,^a and Nick Macklon, Ph.D.^b

^a The Fertility Clinic, Department of Gynecology, Fertility and Obstetrics, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark; ^b London Women's Clinic, London, United Kingdom

Aromatase inhibitors, particularly letrozole (LZ), are now established as first-line ovulation induction agents, offering an effective ovarian stimulation strategy to enhance outcomes of intrauterine insemination. In recent years, they have also emerged as potentially valuable adjuvants to gonadotropin ovarian stimulation for in vitro fertilization, particularly in fertility preservation in women with estrogen-responsive cancers. Their primary mechanism of action is to reduce the circulating estrogen levels by inhibiting androgen aromatization. Recent studies have provided evidence that this property may confer therapeutic advantages in other patients undergoing in vitro fertilization. In this study, evidence supporting the role of adjuvant LZ in poor responders, as a moderator of ovarian hyperstimulation syndrome symptoms, and an agent for improving the luteal phase after ovarian stimulation is reviewed. The use of LZ for endometrial preparation in the frozen-thawed embryo transfer cycle is also considered. (Fertil Steril® 2025;123:41–9. ©2024 by American Society for Reproductive Medicine.)

Key Words: Letrozole, aromatase inhibitor, artificial reproductive technology, IVF, ovulatory disorders

Modern fertility treatments now offer real hope to millions of couples struggling with infertility. Although ovarian stimulation regimens are now mature, opportunities to further improve outcomes by ameliorating detrimental effects associated with supraphysiological endocrine responses require exploration. Cotreatment with aromatase inhibitors, such as letrozole (LZ), represents a promising strategy. This article reviews how LZ is developing a promising role in fertility treatments beyond ovulation induction (OI), considers the clinical potential of LZ as a “multipurpose” adjuvant agent in in vitro fertilization (IVF), and presents recent evidence suggesting it to, indeed, represent a “new remedy.”

MECHANISM OF ACTION

Over the past 2 decades, LZ, a third-generation nonsteroidal aromatase

inhibitor, has proven to be a drug of considerable utility for OI and other fertility treatments (1, 2). Its primary mechanism involves reducing circulating estrogens by reversibly binding to the heme group of the cytochrome p450 subunit of the CYP19 aromatase enzyme, thereby potentially blocking the conversion of androstenedione to estrone and testosterone to estradiol (E2). The aromatase enzyme is highly expressed in the granulosa cells of the ovaries, especially the preovulatory follicles, but is also located in the bone, placenta, brain, and adipose tissue (3–5) (Fig. 1).

The decrease in circulating estrogens results in diminished negative feedback on the hypothalamic-pituitary axis. Consequently, there is an increase in the secretion of endogenous follicle-stimulating hormone (FSH) and luteinizing hormone (LH), which stimulates follicular growth and

enhances ovulation. Letrozole also increases intraovarian androgen levels by inhibiting the aromatization of testosterone and androstenedione, working synergistically with the elevated LH levels to stimulate androgen production from pregnenolone in the theca cells (6–9). This is evident from the midfollicular phase, whereas the dehydroepiandrosterone sulfate levels remained unaffected, indicating a less significant adrenal contribution (10). The overall increase in androgen levels, along with elevated gonadotropin level and enhanced FSH receptor (FSHR) expression, likely promotes earlier follicular recruitment (11, 12).

PHARMACOKINETICS OF LZ

Letrozole is generally available as tablets (2.5 mg) that are fully absorbed from the intestine regardless of food intake. Letrozole is primarily metabolized in the liver and has a half-life of approximately 2–4 days, which is significantly shorter than that of other antiestrogens, such as clomiphene citrate (CC) (13). Once steady state is achieved (after approximately 14

Received August 30, 2024; revised October 22, 2024; accepted October 24, 2024; published online October 31, 2024.

Correspondence: Nathalie Bülow, M.D., Ph.D., The Fertility Clinic, Department of Gynecology, Fertility and Obstetrics, Copenhagen University Hospital, Rigshospitalet, Section 4071, Blegdamsvej 9, DK-2100, Copenhagen, Denmark (E-mail: nsbulow@dadlnet.dk).

Fertil Steril® Vol. 123, No. 1, January 2025 0015-0282/\$36.00

Copyright ©2024 American Society for Reproductive Medicine, Published by Elsevier Inc.
<https://doi.org/10.1016/j.fertnstert.2024.10.040>

days), the effect of LZ is estimated to last 9–10 days after discontinuation. However, this duration is likely shorter in fertility treatments, where the treatment duration often does not reach a steady state. An effect from follicular phase treatment on serum E2 and progesterone (P4) levels has been observed midluteal phase. However, this effect does not extend into subsequent cycles (10, 14, 15).

THERAPEUTIC INDICATIONS

Before LZ emerged as a fertility treatment option, it was primarily used in women with breast cancer as a chemotherapeutic agent because it was shown to slow or halt the growth of estrogen receptor-positive breast tumors because of its potent antiestrogenic effects (16, 17). This remains the primary context in which LZ is officially approved for use by major regulatory agencies such as the U.S. Food and Drug Administration or the European Medicines Agency. Although clinical applications in fertility have largely been adopted in an off-label setting, this has not prevented the recommendation of its use by national and international guidelines, and LZ is now an established agent across a range of fertility treatments (18, 19).

OI AND OVARIAN STIMULATION FOR INTRAUTERINE INSEMINATION

It is now almost 25 years since Mitwally and Casper (20) first proposed LZ as an OI agent acting through suppressing E2 negative feedback on the hypothalamic-pituitary axis and, thus, enhancing endogenous FSH release and stimulating follicular growth. Since these early studies, a large body of research has served to determine the place of LZ among the therapeutic repertoire for treating anovulatory or oligo-ovulatory infertility, and it is now considered by the European Society of Human Reproduction and Embryology and international guideline consensus groups to represent the first-line OI agent for anovulatory infertility (18, 21).

In couples with unexplained infertility and no clear ovulatory problem, outcomes after intrauterine insemination (IUI) or timed intercourse supported by LZ treatment and empirical LZ treatment have been studied in 14 randomized controlled trials (RCTs) and numerous observational studies. In a network meta-analysis, similar reproductive outcomes were shown when comparing LZ with CC and FSH/human menopausal gonadotropin (hMG) (22), and a similar conclusion was drawn by a Cochrane review of eight studies (23). However, mild stimulation with LZ tends to confer a lower risk of multiple pregnancies than FSH/hMG in couples with unexplained infertility (22). Although the use of LZ in these clinical contexts is not the focus of this study, its established role in OI and IUI has provided the confidence to explore the potential benefits that it may offer in IVF.

ADJUVANT LZ TO OVARIAN STIMULATION WITH GONADOTROPINS FOR IVF

In recent years, interest has grown in the role of LZ in IVF treatment. Adjuvant LZ has been proposed to mitigate the negative impact of the supraphysiological serum E2 levels with a possible positive impact on the oocyte, endometrium,

and implantation (24). This property has led to its introduction as an adjuvant to FSH ovarian stimulation in women with estrogen-sensitive tumors, including breast cancer, who are undergoing fertility preservation treatment (25).

In contrast to tamoxifen or CC, which are also sometimes used as adjuvants in this context, the decreased E2 levels with LZ are obtained without affecting estrogen receptors in peripheral tissues, thus avoiding any detrimental effects on endometrium (13, 26). Letrozole has previously been shown in IVF to alter the sex steroid profile in both follicular fluid and serum, resulting in lower E2 and higher androgen levels in LZ-exposed women (10, 27). This shift in sex steroids toward higher androgens may result in increased FSHR expression (8). Furthermore, the suppressed E2 level reduces negative feedback on the hypothalamic-pituitary axis and increased the secretion of both FSH and LH (28). This may act in synergy to stimulate follicular growth and increase luteal phase endogenous LH, FSH-induced LH receptors, and P4 production (10, 14, 15). However, disrupting the endocrine milieu could have detrimental effects on reproductive functions and clinical outcomes, and most studies, thus far, have been underpowered to detect these effects.

A ROLE IN POOR RESPONDERS?

In the search for effective strategies to improve outcomes in low responders, adjuvant LZ has attracted attention. It has been proposed that by increasing precursor androgens and, hence, up-regulating follicular FSHR expression, LZ cotreatment may render recalcitrant follicles more sensitive to stimulation (8, 11, 28). Moreover, the higher endogenous FSH production generated by LZ from early in the follicular phase contributes to the total serum FSH levels and lower FSH consumption, as shown in 3 RCTs and a retrospective cohort study (10, 29–31). This endogenous production includes the acidic isoform of FSH, absent in the exogenous recombinant FSH often used (9). These mechanisms provide a rationale for adding LZ as an alternative to increasing FSH doses in poor responders (29). In our recent systematic review and meta-analysis of 20 studies (11 RCTs) in poor responders, which included 2,596 women, evidence emerged to support this notion (32). Although the included studies varied in the specific protocol applied, cotreatment with LZ was associated with a significant increase in the live birth rate (LBR) (7%; 95% confidence interval, 1%–13%) (Fig. 2) (32). There was no difference in the number of oocytes retrieved or cycle cancellations; however, there was a significant reduction in FSH consumption and stimulation days (32). A further meta-analysis found slightly improved reproductive outcomes with an increased clinical pregnancy rate with an odds ratio of 1.57 (95% confidence interval, 1.00–2.44) and reduced FSH consumption (33). A large retrospective study reported an 11% increase in LBR in younger low responders (<35 years) treated with adjuvant LZ but no difference in older women (34). To obtain a consistent suppression of the E2 levels, LZ (5 mg daily) concomitantly with FSH for at least 5 days or continued until the day of triggering final oocyte maturation can be recommended with gonadotropin release hormone (GnRH) antagonist cotreatment.

CAN LZ COTREATMENT ENHANCE THE FOLLICULAR PHASE OF EXPECTED NORMAL AND HIGH RESPONDERS?


Our recent RCT of applying adjuvant LZ to gonadotropin ovarian stimulation revealed certain novel insights into the endocrine effects of LZ cotreatment. Those randomized to receive the intervention in this double-blind placebo-controlled trial revealed higher androgen levels and endogenous FSH production during the follicular phase (Fig. 3) (10). In a subcohort, we also observed an enhanced follicular recruitment in those cotreated with LZ but no significant improvement in reproductive outcomes, although the study was underpowered for this endpoint (10, 15). This finding was consistent with our systematic review showing that the addition of LZ in normal responders resulted in nearly two more oocytes per cycle but did not significantly affect other outcomes (32). This meta-analysis, which included a limited number of studies on high responders, showed no significant effect of LZ on LBR or miscarriage rates and provided no data on other outcomes. On the basis of current evidence, there is insufficient evidence to recommend adjuvant LZ in IVF for expected normal or high responders to improve clinical outcomes (32).

CAN FOLLICULAR PHASE LZ TREATMENT ACT TO SUPPORT THE LUTEAL PHASE?

The impact of ovarian stimulation on the luteal phase has been a subject of much conjecture and research since the early

days of IVF (35). Proposed mechanisms for the “inadequate luteal phase” and the need for support in IVF included prolonged suppression of the pituitary gland by GnRH agonists and the removal of granulosa cells at oocyte retrieval. However, in an intervention study published some years ago, Beckers et al. (36) provided compelling evidence to implicate high late follicular phase E2 levels as the primary disruptor. In this study, the duration of the unsupported luteal phase was found to be negatively correlated to the late follicular E2 blood level. Our group hypothesized that by preventing the supraphysiological estrogen levels from increasing, this may avoid suppression of corpus luteal function and, hence, limit the detrimental effect on luteal phase endocrinology. In turn, cotreatment with the LZ in the follicular phase may reduce the need for aggressive luteal phase support. Initial evidence supporting this hypothesis was provided by our RCT of LZ as an adjuvant for women undergoing IVF with luteal support (10, 15). Those who had received LZ cotreatment throughout the follicular phase were found to maintain high blood LH, P4, and androgen levels in the midluteal phase (Fig. 3). We then tested this hypothesis in a placebo-controlled RCT in oocyte donors who were triggered with GnRH agonist after antagonist cotreatment but received no luteal support. This study showed that when a GnRH agonist is employed to trigger final oocyte maturation, follicular phase cotreatment with LZ resulted in an extended duration of the unsupported luteal phase and a marked and significant increase in the midluteal phase P4 levels (Fig. 4) (14). Indeed, this increase was observed


FIGURE 1



Letrozole mechanism of action. Created with BioRender. FSH = follicle-stimulating hormone; FSHR = follicle-stimulating hormone receptor; GnRH = gonadotropin-releasing hormone; LH = luteinizing hormone; LHR = luteinizing hormone receptor.

Bülow. Letrozole in IVF. Fertil Steril 2025.

FIGURE 2



to be 30-fold compared with that in the group that did not receive LZ cotreatment (14). These striking findings suggest that adjuvant LZ mitigates the adverse effects of ovarian stimulation on the luteal phase and potentially decreases the necessity for supplementary luteal hormonal support in cases where a GnRH agonist is used for triggering. Clinical studies are now warranted to test this further.

OOCYTE AND EMBRYO QUALITY

It has been suggested that the relative decrease in the E2 level within the follicular compartment, which could arise in response to LZ cotreatment during ovarian stimulation, may be detrimental to oocyte development (37, 38). However, a number of studies challenge this (39, 40). Studies in fertility preservation using adjuvant LZ report not only more mature oocytes and frozen embryos but also higher abnormal fertilization rates (41, 42). A similar number and quality of oocytes and markers of embryo development were reported in both a recent systematic review and subsequent RCT (43, 44). Other studies have shown adjuvant LZ to be associated with an increase in oocytes obtained and the quality of the embryos (31, 45–48). Thus, the effect of adjuvant LZ on oocyte development in IVF treatment remains unclear. Our recent systematic review on adjuvant LZ found insufficient studies for a qualified comparison of embryo quality between LZ-exposed and nonexposed oocytes, indicating the need for further research (32).

DOES LZ OFFER A “BEST OF BOTH WORLDS” APPROACH TO ENDOMETRIAL PREPARATION FOR FROZEN-THAWED EMBRYO TRANSFER CYCLES?

Endometrial preparation before frozen-thawed embryo transfer (FET) can be achieved through various approaches. In women with ovulatory cycles, a natural cycle FET can be


employed, whereas stimulated FET using LZ or FSH/hMG can be used for women with oligoanovulation to induce follicular growth and ensure formation of a corpus luteum. Artificial cycle (AC) FET, involving E2 followed by combined E2 and P4 treatment, is widely used because of the convenience of scheduling (2, 49). However, recent studies have highlighted adverse obstetric and perinatal outcomes associated with AC FET in the absence of a corpus luteum (49, 50) suggesting that this approach should perhaps be restricted to women with premature ovarian failure undergoing oocyte donation.

The comparison of LZ vs. AC FET for women with polycystic ovary syndrome or oligoanovulation has been studied extensively in cohort studies showing an increased LBR in most cases (51–54). Perinatal outcomes with hypertensive disorders of pregnancy are mostly found lower or similar (52–55), as well as the risk of having a newborn with birth weight larger for gestational age (52, 53, 55, 56). However, the improved LBRs or ongoing pregnancy rates are not confirmed in the only two RCTs, where the LBRs/ongoing pregnancy rates are similar in the two groups (57, 58). In conclusion, LZ FET shows potential benefits in reducing perinatal complications and may emerge as an optimal means of supporting the natural cycle in oligoanovulatory women while retaining corpus luteum function. Clinical studies addressing this possibility are ongoing.

DOES COTREATMENT WITH LZ REDUCE THE RISK OF OVARIAN HYPERSTIMULATION SYNDROME?

Restricting the supraphysiological estrogen levels with LZ has been proposed as a strategy to reduce the risk of clinical consequences of ovarian hyperstimulation syndrome (OHSS). In a systematic review of the incidence of OHSS with adjuvant LZ in a long GnRH agonist protocol that included eight studies, a significantly reduced overall incidence of early OHSS in two

FIGURE 3




Follicular and supported luteal phase hormonal changes with adjuvant letrozole vs. placebo. Median and interquartile range. *P<0.05. FSH = follicle-stimulating hormone; LH = luteinizing hormone; OPU = oocyte pick up; stim = stimulation. (Modified from Bülow et al. (10) with permission.)

Bülow. Letrozole in IVF. Fertil Steril 2025.

studies of 566 women and pooled moderate and severe OHSS in eight studies of 1,551 women were reported. However, no difference was found when patients were categorized by mild, moderate, or severe OHSS. Because of significant heterogeneity among the included studies, LZ cotreatment

for OHSS prevention is not recommended by recently updated OHSS guidelines for the American Society for Reproductive Medicine (59, 60). ClinicaTrials.gov shows that an RCT of LZ's role in OHSS risk reduction is being performed, and the evidence base may, therefore, shift in the future.

FIGURE 4

SIDE EFFECTS OF LZ

LZ is associated with mild and infrequent side effects, such as gastrointestinal issues, fatigue, headaches, and dizziness (61), and some reports suggest that it has a similar side effect profile to CC; however, CC is more commonly linked to a higher incidence of hot flashes (62, 63). The side effects listed in the product information for LZ primarily stem from its long-term use in postmenopausal women, making it challenging to directly apply these data to younger women who use the medication for shorter durations.

NEONATAL SAFETY

The safety of LZ treatment concerning neonatal outcomes has been a topic of concern since 2005 when a study of 150 newborns indicated an increased risk of locomotor disabilities and congenital cardiovascular malformations, despite no overall increase in malformations (64). This study faced significant criticism because of its high-risk of bias: the control group was from a tertiary hospital and excluded high-risk births, and the women were spontaneously pregnant and significantly younger and included only one twin pregnancy compared with 19 twin pregnancies in the LZ group. This led to widespread abandonment of LZ use by health authorities, restricting its use to controlled trials.

Subsequent research provided more reassuring results. A large retrospective study involving 911 children and a

systematic review of 4,697 children born after LZ treatment for IUI found no increase in major congenital malformations compared with other fertility treatments (65, 66). Furthermore, published data of FET cycles using LZ show no increase in the proportion of congenital malformations, with a total of 5 of 1,181 newborns after LZ treatment, which was not significantly different from the AC preparation before FET, although the numbers in both groups are suspected to be underreported compared with the congenital malformation rates in naturally conceived pregnancies (52, 54, 56, 67). Despite the first study of LZ in IVF being published in 2004, only one retrospective study of adjuvant LZ in IVF in obese women has specifically addressed neonatal malformations and found no malformations in either group (36 LZ-treated women vs. 21 controls) (68). This forces clinicians to rely on IUI and FET data for safety extrapolations. Although these findings are promising, the dose and timing of adjuvant LZ in IVF differ, underscoring the need for further research on neonatal outcomes in this specific context.

Participants in our recent RCT (10) consented to the screening of their resulting offspring's medical histories after birth. The findings showed that all children had a normal weight for gestational age, and no neonatal malformations were recorded within 1 month after birth, although these data remain unpublished. This provides preliminary evidence supporting the safety of adjuvant LZ in IVF but highlights the necessity for more comprehensive studies to confirm these outcomes.

TABLE 1

Potential roles of letrozole in assisted reproductive technology.

Patient groups	Insemination	FET cycles
Ovulatory women	Higher LBR with LZ than with CC, equal to FSH	No benefit compared with natural cycles
Oligoanovulatory women	Higher LBR with LZ than with CC, equal to FSH	Higher LBR with LZ than with AC FET
		Lower HDP and lower risk of LGA children with LZ than with AC FET
	Adjuvant letrozole to FSH for ovarian stimulation for IVF	
Poor responders	Higher LBR, the highest benefit in younger women	
Normal responders	Similar LBR	
	Evidence for improved luteal phase	
High responders	Similar LBR	
	Awaiting a larger study of OHSS risk reduction	

Note: AC = artificial cycle; CC = clomiphene citrate; FET = frozen-thawed embryo transfer; FSH = follicle-stimulating hormone; HDP = hypertensive disorders of pregnancy; IVF = in vitro fertilization; LBR = live birth rate; LGA = large for gestational age; LH = luteinizing hormone; LZ = letrozole; OHSS = ovarian hyperstimulation syndrome.

Bülow. Letrozole in IVF. Fertil Steril 2025.

COST-EFFECTIVENESS OF LZ

Fertility treatments can be financially burdensome, and the cost of LZ is significantly lower than that of other fertility medications; hence, LZ offers a more affordable alternative for OI. Additionally, the observed increase in FSH levels and decreased FSH consumption with adjuvant LZ have the potential to reduce costs in IVF. This could be particularly advantageous for patients with a poor ovarian response (13). The considered indications and current evidence for the comparative efficacy of LZ are briefly summarized in Table 1.

CONCLUSION

Aromatase inhibitors, particularly LZ, represent an important therapeutic option in the management of ovulatory disorders and unexplained infertility with insemination. In addition, LZ can induce follicular growth, producing E2 for endometrial preparation before FET and after ovulation, securing a corpus luteum during pregnancy with a similar LBR and lower perinatal complication. Much of this review has focused on the potential advantages of incorporating adjuvant LZ into ovarian stimulation regimens for IVF. Recently published evidence indicates that it may have a more central place in treatment than hitherto recognized. Benefits appear to include an increased LBR in poor responders and a means of normalizing the luteal phase after ovarian stimulation without the need for burdensome luteal support. Further clinical studies will elucidate whether this is a mirage or, indeed, a new remedy.

CRediT Authorship Contribution Statement

Nathalie Bülow: Software; Investigation, Writing – original draft. Nick Macklon: Writing – review & editing, Supervision, Conceptualization.

Declaration of Interests

N.B. has nothing to disclose. N.M. has nothing to disclose.

REFERENCES

1. Franik S, Eltrop SM, Kremer JA, Kiesel L, Farquhar C. Aromatase inhibitors (letrozole) for subfertile women with polycystic ovary syndrome. *Cochrane Database Syst Rev* 2018;5:CD010287.
2. Løssl K, Spangmose AL, Asserhøj LL, Dam TV, Pinborg A. The future of frozen-thawed embryo transfer in hormone replacement therapy cycles. *Curr Opin Obstet Gynecol* 2023;35:200–9.
3. Simpson ER, Mahendroo MS, Means GD, Kilgore MW, Hinshelwood MM, Graham-Lorence S, et al. Aromatase cytochrome P450, the enzyme responsible for estrogen biosynthesis. *Endocr Rev* 1994;15:342–55.
4. Jeppesen JV, Kristensen SG, Nielsen ME, Humaidan P, Dal Canto M, Fadini R, et al. LH-receptor gene expression in human granulosa and cumulus cells from antral and preovulatory follicles. *J Clin Endocrinol Metab* 2012;97:E1524–31.
5. Bulun SE. Chapter 17 Physiology and pathology of the female reproductive axis. In: *William textbook of endocrinology*. Amsterdam, Netherlands: Elsevier; 2016: 589–663.
6. Garcia-Velasco JA, Moreno L, Pacheco A, Guillén A, Duque L, Requena A, et al. The aromatase inhibitor letrozole increases the concentration of intra-ovarian androgens and improves in vitro fertilization outcome in low responder patients: a pilot study. *Fertil Steril* 2005;84:82–7.
7. Fleming R, Jenkins J. The source and implications of progesterone rise during the follicular phase of assisted reproduction cycles. *Reprod Biomed Online* 2010;21:446–9.
8. Nielsen ME, Rasmussen IA, Kristensen SG, Christensen ST, Mollgard K, Wreford Andersen E, et al. In human granulosa cells from small antral follicles, androgen receptor mRNA and androgen levels in follicular fluid correlate with FSH receptor mRNA. *Mol Hum Reprod* 2011;17:63–70.
9. Yding Andersen C. Inhibin-B secretion and FSH isoform distribution may play an integral part of follicular selection in the natural menstrual cycle. *Mol Hum Reprod* 2017;23:16–24.
10. Bülow NS, Skouby SO, Warzecha AK, Udengaard H, Andersen CY, Holt MD, et al. Impact of letrozole co-treatment during ovarian stimulation with gonadotrophins for IVF: a multicentre, randomized, double-blinded placebo-controlled trial. *Hum Reprod* 2022;37:309–21.
11. Weil S, Vendola K, Zhou J, Bondy CA. Androgen and follicle-stimulating hormone interactions in primate ovarian follicle development. *J Clin Endocrinol Metab* 1999;84:2951–6.
12. Løssl K, Andersen AN, Loft A, Freiesleben NL, Bangsbo S, Andersen CY. Androgen priming using aromatase inhibitor and hCG during early-

- follicular-phase GnRH antagonist down-regulation in modified antagonist protocols. *Hum Reprod* 2006;21:2593–600.
13. Buzdar AU, Robertson JF, Eiermann W, Nabholz JM. An overview of the pharmacology and pharmacokinetics of the newer generation aromatase inhibitors anastrozole, letrozole, and exemestane. *Cancer* 2002;95:2006–16.
14. Dreyer Holt M, Skouby SO, Bülow NS, Englund ALM, Birch Petersen K, Macklon NS. The impact of suppressing estradiol during ovarian stimulation on the unsupported luteal phase: a randomized controlled trial. *J Clin Endocrinol Metab* 2022;107:e3633–43.
15. Poulsen LC, Warzecha AK, Bülow NS, Bungum L, Macklon NS, Yding Andersen C, et al. Effects of letrozole cotreatment on endocrinology and follicle development in women undergoing ovarian stimulation in an antagonist protocol. *Hum Reprod* 2022;37:1557–71.
16. Ellis MJ, Coop A, Singh B, Mauriac L, Llombert-Cussac A, Jänicke F, et al. Letrozole is more effective neoadjuvant endocrine therapy than tamoxifen for ErbB-1- and/or ErbB-2-positive, estrogen receptor-positive primary breast cancer: evidence from a phase III randomized trial. *J Clin Oncol* 2001;19:3808–16.
17. Reddy J, Oktay K. Ovarian stimulation and fertility preservation with the use of aromatase inhibitors in women with breast cancer. *Fertil Steril* 2012;98:1363–9.
18. Teede HJ, Tay CT, Laven J, Dokras A, Moran LJ, Piltonen TT, et al. Recommendations from the 2023 international evidence-based guideline for the assessment and management of polycystic ovary syndrome. *Fertil Steril* 2023;120:767–93.
19. Dahlberg ES, Andersen D, Andersson PS, Birch K, Colmon L, Holt MD, et al. Letrozol I Fertilitetsbehandling. *Ugeskr Laeger* 2023;185:V05230315.
20. Mitwally M, Casper R. Use of an aromatase inhibitor for induction of ovulation in patients with an inadequate response to clomiphene citrate. *Fertil Steril* 2001;75:305–9.
21. Gadalla MA, Norman RJ, Tay CT, Hiam DS, Melder A, Pundir J, et al. Medical and surgical treatment of reproductive outcomes in polycystic ovary syndrome: an overview of systematic reviews. *Int J Fertil Steril* 2020;13:257–70.
22. Danhof NA, Wang R, van Wely M, van der Veen F, Mol BWJ, Mochtar MH. IUI for unexplained infertility—a network meta-analysis. *Hum Reprod Update* 2020;26:1–15.
23. Cantineau AE, Rutten AG, Cohlen BJ. Agents for ovarian stimulation for intrauterine insemination (IUI) in ovulatory women with infertility. *Cochrane Database Syst Rev* 2021;11:CD005356.
24. Macklon NS, Stouffer RL, Giudice LC, Fauser BC. The science behind 25 years of ovarian stimulation for in vitro fertilization. *Endocr Rev* 2006;27:170–207.
25. Oktay K, Hourvitz A, Sahin G, Oktem O, Safo B, Cil A, et al. Letrozole reduces estrogen and gonadotropin exposure in women with breast cancer undergoing ovarian stimulation before chemotherapy. *J Clin Endocrinol Metab* 2006;91:3885–90.
26. Vanden Bossche HV, Moereels H, Koymans LM. Aromatase inhibitors—mechanisms for non-steroidal inhibitors. *Breast Cancer Res Treat* 1994;30:43–55.
27. Andersen CY, Løssl K. Increased intrafollicular androgen levels affect human granulosa cell secretion of anti-Müllerian hormone and inhibin-B. *Fertil Steril* 2008;89:1760–5.
28. Løssl K, Freiesleben NC, Wissing ML, Birch Petersen K, Holt MD, Mamsen LS, et al. Biological and clinical rationale for androgen priming in ovarian stimulation. *Front Endocrinol* 2020;11:627.
29. Bastu E, Buyru F, Ozurmeli M, Demiral I, Dogan M, Yeh J. A randomized, single-blind, prospective trial comparing three different gonadotropin doses with or without addition of letrozole during ovulation stimulation in patients with poor ovarian response. *Eur J Obstet Gynecol Reprod Biol* 2016;203:30–4.
30. Ecmis T, Tasci Y, Caglar GS. Controlled ovarian hyperstimulation with sequential letrozole co-treatment in normo/high responders. *Gynecol Endocrinol* 2016;32:206–9.
31. Eftekhar M, Saeed L. Effect of adding letrozole to gonadotropin on in vitro fertilization outcomes: an RCT. *Int J Reprod Biomed* 2020;18:287–94.
32. Bülow NS, Dreyer Holt M, Skouby SO, Birch Petersen K, Englund ALM, Pinborg A, et al. Co-treatment with letrozole during ovarian stimulation for IVF/ICSI: a systematic review and meta-analysis. *Reprod Biomed Online* 2022;44:717–36.
33. Qin Y. Effects of using letrozole in combination with the GnRH antagonist protocol for patients with poor ovarian response: a meta-analysis. *J Gynecol Obstet Hum Reprod* 2021;50:102139.
34. Kahraman A, Tulek F. Effects of letrozole co-treatment on outcomes of gonadotropin-releasing hormone antagonist cycles in POSEIDON groups 3 and 4 expected poor responders. *Arch Gynecol Obstet* 2022;306:1313–9.
35. Macklon NS, Fauser BC. Impact of ovarian hyperstimulation on the luteal phase. *J Reprod Fertil Suppl* 2000;55:101–8.
36. Beckers NG, Macklon NS, Eijkemans MJ, Ludwig M, Felberbaum RE, Diedrich K, et al. Nonsupplemented luteal phase characteristics after the administration of recombinant human chorionic gonadotropin, recombinant luteinizing hormone, or gonadotropin-releasing hormone (GnRH) agonist to induce final oocyte maturation in in vitro fertilization patients after ovarian stimulation with recombinant follicle-stimulating hormone and GnRH antagonist cotreatment. *J Clin Endocrinol Metab* 2003;88:4186–92.
37. Revelli A, Delle Piane L, Casano S, Molinari E, Massobrio M, Rinaudo P. Follicular fluid content and oocyte quality: from single biochemical markers to metabolomics. *Reprod Biol Endocrinol* 2009;7:40.
38. Carpintero NL, Suárez OA, Mangas CC, Varea CG, Rioja RG. Follicular steroid hormones as markers of oocyte quality and oocyte development potential. *J Hum Reprod Sci* 2014;7:187–93.
39. Rosenbusch B, Djalali M, Sterzik K. Is there any correlation between follicular fluid hormone concentrations, fertilizability, and cytogenetic analysis of human oocytes recovered for in vitro fertilization? *Fertil Steril* 1992;57:1358–60.
40. Costa LO, Mendes MC, Ferriani RA, Moura MD, Reis RM, Silva de Sá MF. Estradiol and testosterone concentrations in follicular fluid as criteria to discriminate between mature and immature oocytes. *Braz J Med Biol Res* 2004;37:1747–55.
41. Shulman Y, Almog B, Kalma Y, Fouks Y, Azem F, Cohen Y. Effects of letrozole or tamoxifen coadministered with a standard stimulation protocol on fertility preservation among breast cancer patients. *J Assist Reprod Genet* 2021;38:743–50.
42. Turan V, Bedoschi G, Emirdar V, Moy F, Oktay K. Ovarian stimulation in patients with cancer: impact of letrozole and BRCA mutations on fertility preservation cycle Outcomes. *Reprod Sci* 2018;25:26–32.
43. Bonardi B, Massarotti C, Bruzzone M, Goldrat O, Mangili G, Anserini P, et al. Efficacy and safety of controlled ovarian stimulation with or without letrozole co-administration for fertility preservation: a systematic review and meta-analysis. *Front Oncol* 2020;10:574669.
44. Bülow NS, Warzecha AK, Nielsen MV, Andersen CY, Holt MD, Petersen MR, et al. Impact of letrozole co-treatment during ovarian stimulation on oocyte yield, embryo development, and live birth rate in women with normal ovarian reserve: secondary outcomes from the RIOT trial. *Hum Reprod* 2023;38:2154–65.
45. Davar R, Oskouian H, Ahmadi S, Firouzabadi RD. GnRH antagonist/letrozole versus microdose GnRH agonist flare protocol in poor responders undergoing in vitro fertilization. *Taiwan J Obstet Gynecol* 2010;49:297–301.
46. Lee VC, Chan CC, Ng EH, Yeung WS, Ho PC. Sequential use of letrozole and gonadotrophin in women with poor ovarian reserve: a randomized controlled trial. *Reprod Biomed Online* 2011;23:380–8.
47. Liu S, Kuang Y, Wu Y, Feng Y, Lyu Q, Wang L, et al. High oestradiol concentration after ovarian stimulation is associated with lower maternal serum beta-HCG concentration and neonatal birth weight. *Reprod Biomed Online* 2017;35:189–96.
48. Nabati A, Peivandi S, Khalilian A, Mirzaeaird S, Hashemi SA. Comparison of GnRh agonist microdose flare up and GnRh antagonist/letrozole in treatment of poor responder patients in intra cytoplasmic sperm injection: randomized clinical trial. *Glob J Health Sci* 2015;8:166–71.
49. von Versen-Höyneck F, Griesinger G. Should any use of artificial cycle regimen for frozen-thawed embryo transfer in women capable of ovulation be abandoned: yes, but what's next for FET cycle practice and research? *Hum Reprod* 2022;37:1697–703.
50. Busnelli A, Schirripa I, Fedele F, Bulfoni A, Levi-Setti PE. Obstetric and perinatal outcomes following programmed compared to natural frozen-thawed embryo transfer cycles: a systematic review and meta-analysis. *Hum Reprod* 2022;37:1619–41.

51. Atkinson M, Crittenden J, Smith H, Sjoblom C. Retrospective cohort study on preparation regimens for frozen embryo transfer. *Reprod Fertil* 2021;2:308–16.
52. Zhang W, Liu Z, Zhang J, Ren B, Liu M, Li J, et al. Comparison of perinatal outcomes of letrozole-induced ovulation and hormone replacement therapy protocols in patients with abnormal ovulation undergoing frozen-thawed embryo transfer: a propensity score matching analysis. *Front Endocrinol (Lausanne)* 2022;13:837731.
53. Wang X, Li Y, Tan H, Cai S, Ma S, Peng Y, et al. Letrozole-stimulated endometrial preparation protocol is a superior alternative to hormone replacement treatment for frozen embryo transfer in women with polycystic ovary syndrome, a cohort study. *Reprod Biol Endocrinol* 2023;21:101.
54. Wen W, Shi J, Liu X. What is the optimal means of preparing the endometrium in frozen-thawed embryo transfer cycles among anovulatory women? A retrospective cohort study. *Eur J Obstet Gynecol Reprod Biol* 2024;294:79–83.
55. Zhang Y, Fu X, Gao S, Gao S, Gao S, Ma J, et al. Letrozole use in vitrified single-blastocyst transfer cycles is associated with lower risk of large for gestational age infants in patients with polycystic ovary syndrome. *J Assist Reprod Genet* 2023;40:2885–94.
56. Zhang J, Li Z, Sun L, Guan Y, Du M. Comparison of pregnancy and neonatal outcomes of single frozen blastocyst transfer between letrozole-induction and HRT cycles in patients with abnormal ovulation. *Front Endocrinol (Lausanne)* 2021;12:664072.
57. Hosseini-Najarkolaee A, Moini A, Kashani L, Farid Mojtahedi M, Hosseini-Najarkolaee E, Salehi E. The effect of letrozole versus artificial hormonal endometrial preparation on pregnancy outcome after frozen-thawed embryos transfer cycles: a randomized clinical trial. *Reprod Biol Endocrinol* 2020;18:115.
58. Yuan Y, Chang Q, Wen Y, Gao J, Huang S, Xu Y, et al. Letrozole during frozen embryo transfer in women with polycystic ovarian syndrome: a randomized controlled trial. *Obstet Gynecol* 2023;142:1087–95.
59. Practice Committee of the American Society for Reproductive Medicine. Electronic address:asrm@asrm.org; Practice Committee of the American Society for Reproductive Medicine. Testing and interpreting measures of ovarian reserve: a committee opinion. *Fertil Steril* 2020;114:1151–7.
60. Zhao J, Xu B, Huang X, Yan Y, Li Y. Whether letrozole could reduce the incidence of early ovary hyperstimulation syndrome after assisted reproductive technology? A systematic review and meta-analysis. *Reprod Health* 2020;17:181.
61. Casper RF, Mitwally M. A historical perspective of aromatase inhibitors for ovulation induction. *Fertil Steril* 2012;98:1352–5.
62. Legro RS, Brzyski RG, Diamond MP, Coutifaris C, Schlaff WD, Casson P, et al. Letrozole versus clomiphene for infertility in the polycystic ovary syndrome. *N Engl J Med* 2014;371:119–29.
63. Al-Thuwainee S, Swadi AAJ. Comparing efficacy and safety of stair step protocols for clomiphene citrate and letrozole in ovulation induction for women with polycystic ovary syndrome (PCOS): a randomized controlled clinical trial. *J Med Life* 2023;16:725–30.
64. Biljan MM, Hemmings R, Brassard N. The outcome of 150 babies following the treatment with letrozole or letrozole and gonadotropins. *Fertil Steril* 2005;84:S95.
65. Tulandi T, Martin J, Al-Fadhli R, Kabli N, Forman R, Hitkari J, et al. Congenital malformations among 911 newborns conceived after infertility treatment with letrozole or clomiphene citrate. *Fertil Steril* 2006;85:1761–5.
66. Pundir J, Achilli C, Bhide P, Sabatini L, Legro RS, Rombauts L, et al. Risk of foetal harm with letrozole use in fertility treatment: a systematic review and meta-analysis. *Hum Reprod Update* 2021;27:474–85.
67. Niu Y, Zhao D, Wang Y, Suo L, Zou J, Wei D. Ovulation induction regimens are associated with a higher rate of livebirth after frozen single-blastocyst transfer among women with polycystic ovary syndrome. *Front Endocrinol (Lausanne)* 2022;13:987813.
68. Lin J, Wu F, Zhu Y, Wang B, Cao Z, Lin J. Letrozole co-treatment in an antagonist protocol for overweight women undergoing IVF treatment: a retrospective study. *BMC Pregnancy Childbirth* 2024;24:592.