

Applied Graph Theory

Contents

List of Symbols	2
1 Introduction	3
2 Cartesian Products	3
3 Subgraphs and Complements	3
4 Walks, Paths, and Cycles	4
5 Distances	5
6 Connectedness	6
7 Graph Isomorphism	6
8 Self-Complementary Graphs	7
9 Bipartite Graphs	8
10 Trees	9
11 Blocks	10
12 Line Graphs	11
13 Adjacency Matrices	12
14 Incidence Matrices	13

List of Symbols

$V(G)$	Vertex set of graph G
$E(G)$	Edge set of graph G
\sim	is adjacent to
$\deg(v)$	Degree of vertex v
$\text{ecc}(v)$	Eccentricity of vertex v
$\text{rad}(G)$	Radius of graph G
$\text{diam}(G)$	Diameter of graph G
$A(G)$	Adjacency matrix graph G
$G - v$	Graph obtained from G by deleting vertex v and all edges incident with v
$G - e$	Graph obtained from G by deleting edge e
K_n	Complete graph of order n (size $\binom{n}{2}$)
P_n	Path graph of order n (size $n - 1$)
C_n	Cycle graph of order n (size n)

1 Introduction

A **graph** is an ordered pair $G = (V, E)$ where V is a nonempty set called the **vertex set**, whose elements are the **vertices** of G , and E is a set of unordered pairs of distinct vertices of G , whose elements are the **edges** of G . The **order** of G is its number of vertices and the **size** of G is its number of edges.

Two vertices u and v of G are **adjacent**, denoted by $u \sim v$, if $(u, v) \in E$ – we may also denote the edge (u, v) as simply uv . Note that $uv = vu$. If there is no edge between u and v , then they are **nonadjacent**, denoted by $u \not\sim v$. The vertex v and the edge uv are said to be **incident** with each other. The **degree** of a vertex v is the number of edges incident with it, or equivalently, the number of vertices adjacent with it, and is denoted by $\deg(v)$.

Lemma 1.1 (Handshaking Lemma). *The sum of the degrees of all vertices of a graph is twice its size.*

Proof. Exercise. □

Corollary 1.2. *The number of vertices of odd degree in a graph is always even.*

Proof. Exercise. □

A **regular graph** is a graph in which all vertices have the same degree. This common value of the degree is the **regularity** of the graph. A regular graph of regularity k is a **k -regular graph**. A **cubic graph** is a 3-regular graph.

Exercise 1.1. Prove that any cubic graph has even order. For what other order(s) will this statement hold?

2 Cartesian Products

The **Cartesian product** of two graphs G and H is the graph denoted by $G \times H$, whose vertex set is $V(G) \times V(H)$ (the Cartesian product of the vertex sets $V(G)$ and $V(H)$, consisting of all ordered pairs (u, v) where u is a vertex of G and v is a vertex of H), in which two vertices (u_1, v_1) and (u_2, v_2) are adjacent if and only if either $u_1 = u_2$ and $v_1 \sim v_2$ or $u_1 \sim u_2$ and $v_1 = v_2$.

Exercise 2.1. Prove that if G is a graph of order p_1 and size q_1 , and H is a graph of order p_2 and size q_2 , then $G \times H$ is a graph of order $p_1 p_2$ and size $p_1 q_2 + p_2 q_1$.

Exercise 2.2. Show that if G is an r -regular graph and H is an s -regular graph, then $G \times H$ is a regular graph of regularity $r + s$.

3 Subgraphs and Complements

A **subgraph** of a graph G is a graph H whose vertex and edge sets are, respectively, subsets of the vertex and edge sets of G – i.e., $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. A **spanning subgraph** of G is a subgraph containing all the vertices of G . A subgraph H of a graph G is an **induced subgraph** if for all vertices $u, v \in V(H)$, $uv \in E(H)$ whenever $uv \in E(G)$ – i.e., every edge of G that joins two vertices of H is present in H .

Example 3.1. Fig. 1 shows a graph G and three subgraphs H_1 , H_2 , and H_3 of G . The subgraph H_1 is neither a spanning subgraph (e.g. it does not contain v_3), nor an induced subgraph (e.g. it contains vertices v_2 and v_5 of G but not the edge v_2v_5). H_2 is a spanning subgraph and H_3 is an induced subgraph of G .




Figure 1: A graph with three of its subgraphs

A graph is **complete** if all its vertices are adjacent to one another. The complete graph of order n is denoted by K_n . As there is an edge between every pair of distinct vertices of K_n , the size of K_n is $\binom{n}{2}$. The **totally disconnected graph** of order n , denoted by $\overline{K_n}$ is the graph of order n that has no edges.

The **complement** of a graph G is the graph \overline{G} with the same vertices as G , in which any two vertices u and v are adjacent if and only if they are non-adjacent in G . Thus, if G is an (n, m) -graph, then \overline{G} is an $(n, \binom{n}{2} - m)$ -graph. Observe that $\overline{K_n}$ is (as the notation suggests) the complement of K_n .

Exercise 3.1. If G is a graph of order n , and $\deg_G v = k$, what is $\deg_{\overline{G}} v$?

Exercise 3.2. Show that if G and \overline{G} are regular graphs of order n with the same regularity, then the regularity is even and $n = 4t + 1$ for some integer t .


4 Walks, Paths, and Cycles

A **walk** in a simple graph G is a sequence of vertices v_1, v_2, \dots, v_k such that $v_i \sim v_{i+1}$ for $i = 1, 2, \dots, k-1$. This is a walk **from** v_1 **to** v_k or **between** v_1 **and** v_k , having **length** $k-1$. We may also say that this is a walk of length $k-1$ starting at v_1 and ending at v_k .


A walk in which no vertex is repeated is a **path**. Note that the length of a path is the number of edges in it.

A **closed walk** in G is a walk starting and ending at the same vertex. A closed walk in which no vertex is repeated is a **cycle**. The length of a cycle is also equal to the number of vertices in it. A cycle of the form v_1, \dots, v_{k-1}, v_k is usually referred to as “the cycle v_1, \dots, v_{k-1} ”.


Example 4.1. Consider the graph G shown below.




A walk of length 5 from b to c in G is b, f, e, a, e, c .



This walk is not a path, since the vertex e is repeated. An example of a path from b to c in the same graph G is b, f, e, c . The length of this path is 3.



A cycle of length 4 in G is b, f, e, c, b .



5 Distances

The **distance** between two vertices u and v of a graph G is the length of a shortest path between u and v – such a shortest path between u and v is a **geodesic** from u to v . Thus, the distance between u and v in G , denoted by $d_G(u, v)$ is the length of a geodesic from u to v in G . When the graph G is clear from the context, we will write $d_G(u, v)$ as $d(u, v)$.

The **eccentricity** of a vertex v , denoted by $\text{ecc}(v)$, is the maximum of all distances from v to any other vertex. That is,

$$\text{ecc}(v) = \max_{u \in V(G)} d(u, v).$$

The **diameter** of G is the maximum of the eccentricities of vertices of G , and the **radius** of G is the minimum of the eccentricities of vertices of G . They are denoted by $\text{diam}(G)$ and $\text{rad}(G)$ respectively. Thus,

$$\text{diam}(G) = \max_{v \in V(G)} \text{ecc}(v) = \max_{u, v \in V(G)} d(u, v)$$

$$\text{rad}(G) = \min_{v \in V(G)} \text{ecc}(v) = \min_{v \in V(G)} \max_{u \in V(G)} d(u, v).$$

The set of all vertices of G having minimum eccentricity, i.e., the set of all $v \in V(G)$ such that $\text{ecc } v = \text{rad } G$, is the **centre** of G .

6 Connectedness

A graph is **connected** if there is a path between every two of its vertices. Otherwise, it is **disconnected**. A **(connected) component** of a graph G is a maximal connected subgraph of G . Thus, a graph G is connected if and only if it has exactly one component.

Theorem 6.1. *For any graph G , either G or \overline{G} is connected.*

Proof. Consider a graph G , and suppose that G is disconnected. We shall show that \overline{G} is connected, by showing that there is a path between every two vertices of \overline{G} .

Let u and v be any two vertices of \overline{G} . If they are not adjacent in G , then they are adjacent in \overline{G} , and hence there is a path uv from u to v in \overline{G} . If u and v are adjacent in G , then they belong to the same component of G . As G is disconnected, it has at least one more component containing at least one vertex, say w , which is necessarily non-adjacent to both u and v . Hence, in \overline{G} , w is adjacent to both u and v . Thus, there is a path uwv from u to v in \overline{G} . Therefore, \overline{G} is connected. \square

Theorem 6.2. *For any connected graph G , if $\text{diam } G \geq 3$, then $\text{diam } \overline{G} \leq 3$.*

Proof. Consider a graph G of diameter at least 3. Then there exist vertices u and v in G such that $d_G(u, v) = 3$. This implies that u and v are non-adjacent in G , and hence adjacent in \overline{G} .

Now, consider any two vertices x and y other than u and v . In G , x can be adjacent to at most one of u and v , for otherwise, $d_G(u, v) = 2$. Hence, x is adjacent to at least one of u and v in \overline{G} . Similarly, y is adjacent to at least one of u and v in \overline{G} . Thus, there is a path of length at most 3 from x to y in \overline{G} (namely xuy , xvy , $xuvy$, or $xvuy$), which implies that $d_{\overline{G}}(x, y) \leq 3$. Therefore, $\text{diam } \overline{G} \leq 3$. \square

7 Graph Isomorphism

An **isomorphism** from a graph G to a graph H is a bijective¹ function $f: V(G) \rightarrow V(H)$ such for all vertices u and v of G , $u \sim_G v$ if and only if $f(u) \sim_H f(v)$. In other words, a graph isomorphism is a bijection from the vertex set of the first graph to that of the second, that preserves both edges and non-edges. If there exists an isomorphism from G to H , then G and H are **isomorphic**. Then we write $G \cong H$.

Isomorphic graphs have exactly the same structure – i.e., all properties of the graph that do not depend on the labelling or drawing will be shared by isomorphic graphs. For example, isomorphic graphs have the same order, size, degree sequence, diameter, and radius. Indeed, if f is an isomorphism from G to H , and v is a vertex of G , then the neighbours of $f(v)$ in H are the images of the neighbours of v in G , and $\deg_G v = \deg_H f(v)$. Similarly, if u and v are two vertices of G , then $d_G(u, v) = d_H(f(u), f(v))$.

Theorem 7.1. *Graph isomorphism is an equivalence relation on the class of all graphs.*

Proof. To show that graph isomorphism is an equivalence relation, we need to show that \cong is a reflexive, symmetric, transitive relation between graphs. First, observe that the identity map on the vertex set of a graph is an isomorphism from the graph to itself – for, if G is a graph and id denotes the identity map on its vertex set then for any two vertices u and v of G , $u \sim v$ if and only if $\text{id}(u) \sim \text{id}(v)$, as $\text{id}(u) = u$ and $\text{id}(v) = v$. Hence, \cong is reflexive.


¹injective and surjective; i.e., 1-1 and onto

Next, suppose that $G \cong H$. Then there exists an isomorphism f from G to H . Since $f: V(G) \rightarrow V(H)$ is bijective, it has an inverse $f^{-1}: V(H) \rightarrow V(G)$. We claim that f^{-1} is an isomorphism from H to G . We know that f^{-1} is bijective. To see that it preserves edges and non-edges, observe that if x and y are two vertices of H , then $x = f(f^{-1}(x))$ and $y = f(f^{-1}(y))$, which implies that $x \sim_H y$ if and only if $f^{-1}(x) \sim_G f^{-1}(y)$ (since f is an isomorphism from G to H). Therefore, f^{-1} is an isomorphism from H to G . This shows that $H \cong G$. Thus, \cong is symmetric.

Finally, suppose that $G \cong H$ and $H \cong K$. Then there exist isomorphisms f from G to H and g from H to K . We claim that $g \circ f$ is an isomorphism from G to K . Indeed, $g \circ f$ is a function from $V(G)$ to $V(K)$, and being a composition of bijections, is itself a bijection. Now, suppose u and v are two vertices of G . Then $u \sim_G v$ if and only if $f(u) \sim_H f(v)$ (since f is an isomorphism from G to H) if and only if $g(f(u)) \sim_K g(f(v))$ (since g is an isomorphism from H to K). Thus, $g \circ f$ is an isomorphism from G to K . Therefore, \cong is transitive. \square

8 Self-Complementary Graphs

A graph is **self-complementary** if it is isomorphic to its complement – i.e., G is self-complementary if $G \cong \overline{G}$. For example, K_1 , P_4 and C_5 are self-complementary graphs of orders 1, 4, and 5 respectively. There is one more self-complementary graph of order 5, namely the **bull graph**, which can be constructed by adding one new vertex to P_4 and making it adjacent to the two non-pendant vertices of the path. This graph is shown below.



Theorem 8.1. *The order of any self-complementary graph is $4k$ or $4k + 1$, for some non-negative integer k .*

Proof. Let G be a self-complementary graph of order n and size m . Then the size of \overline{G} is $\binom{n}{2} - m$. Since $G \cong \overline{G}$, $m = \binom{n}{2} - m$, which implies that $m = \frac{1}{2} \binom{n}{2} = \frac{n(n-1)}{4}$. As m is an integer, this implies that 4 divides $n(n-1)$, which in turn implies that either one of n and $n-1$ is divisible by 4, or both n and $n-1$ are even. Since the latter is not possible, it follows that $n = 4k$ or $n-1 = 4k$, i.e., $n = 4k$ or $4k + 1$ for some integer k . \square

From Theorem 6.1, we know that a graph and its complement cannot both be disconnected. Thus, if G is a disconnected graph, then \overline{G} must be connected, and therefore it cannot be isomorphic to G . This implies that a self-complementary graph is necessarily connected.

Corollary 8.2. *Every self-complementary graph is connected.* \square

Similarly, from Theorem 6.2, we obtain the following corollary about the diameter of self-complementary graphs.

Corollary 8.3. *Every non-trivial self-complementary graph has diameter 2 or 3.*

Proof. Consider a non-trivial, self-complementary graph G , and suppose that it has diameter strictly greater than 3. Then, by Theorem 6.2, $\text{diam } \overline{G} \leq 3$, which implies that $\text{diam } \overline{G} \neq \text{diam } G$, a contradiction, since $G \cong \overline{G}$. Hence, $\text{diam } G \leq 3$. On the other hand, a non-trivial graph cannot have diameter 0, hence $\text{diam } G \geq 1$. But if $\text{diam } G = 1$, then G is a non-trivial complete graph, whose complement is a totally disconnected graph, which is again impossible. Hence, $\text{diam } G = 2$ or 3. \square

Exercise 8.1. If G is a regular self-complementary graph of order n , show that $n = 4k + 1$, for some integer k , and $\text{diam } G = 2$.

Exercise 8.2. Let G be a self-complementary graph, and let H be the graph obtained by taking the disjoint union of H and P_4 , and making every vertex of H adjacent to the two non-pendant vertices of this P_4 . Then show that H is also self-complementary. What is the graph H obtained in this manner if $G = K_1$?

Exercise 8.3. Construct a self-complementary graph of order 9.

Exercise 8.4. Show that for each positive integer n of the form $4k$ or $4k + 1$, where k is an integer, there exists at least one self-complementary graph of order n .

9 Bipartite Graphs

A graph $G = (V, E)$ is **bipartite** if its vertex set V can be partitioned into two subsets V_1 and V_2 such that no two vertices in V_i are adjacent, for $i = 1, 2$. That is, there exist two non-empty, disjoint subsets $V_1, V_2 \subseteq V$ such that $V_1 \cup V_2 = V$, and every edge of G (if any) joins a vertex in V_1 with a vertex in V_2 . Then we say that (V_1, V_2) is a **bipartition** of G .

The following theorem characterises bipartite graphs in terms of its cycles. A cycle is **even** if its length is even and **odd** if its length is odd.

Theorem 9.1. A graph is bipartite if and only if it contains no odd cycles.

Proof. First, suppose that G is a bipartite graph with bipartition (V_1, V_2) , and let v_1, v_2, \dots, v_k be a cycle of length k in G . Without loss of generality, suppose that $v_1 \in V_1$. Then, since $v_2 \sim v_1$, $v_2 \in V_2$, which in turn implies that $v_3 \in V_1$, as $v_2 \sim v_3$. Proceeding similarly, we see $v_i \in V_1$ if i is odd and $v_i \in V_2$ if i is even. But $v_k \sim v_1$ and $v_1 \in V_1$ implies that $v_k \in V_2$. Therefore, k (the length of the cycle) is even.

Conversely, suppose G is a graph that has no odd cycles. Without loss of generality, assume that G is connected – otherwise, apply the argument to each component. Let v be any vertex of G . Define subsets V_1 and V_2 of $V(G)$ as follows:

$$\begin{aligned} V_1 &= \{ u \in V \mid d(u, v) \text{ is even} \} \\ V_2 &= \{ u \in V \mid d(u, v) \text{ is odd} \}. \end{aligned}$$

We claim that (V_1, V_2) is a bipartition of G .

Suppose not, and let x and y be adjacent vertices of G , both belonging to V_i for $i = 1$ or 2. Let P and Q be shortest paths from v to x and y respectively. Let w be the last vertex common to both P and Q when traversing P from v to x . Then the portion of P from w to x , the edge (x, y) , and the portion of Q from y to w forms a cycle in G , having length, say l , given by

$$\begin{aligned} l &= d(w, x) + d(w, y) + 1 \\ &= [d(v, x) - d(v, w)] + [d(v, y) - d(v, w)] + 1 \\ &= d(v, x) + d(v, y) - 2d(v, w) + 1. \end{aligned}$$

If $x, y \in V_1$, then both $d(v, x)$ and $d(v, y)$ are even, and if $x, y \in V_2$, then both these numbers are odd. In either case, $d(v, x) + d(v, y)$ is even, and hence l is odd, which is a contradiction. Hence, (V_1, V_2) is a bipartition of G as claimed. \square

10 Trees

A **tree** is a connected, acyclic graph. There are several well known characterisations or alternative definitions of trees. We take the given definition as the basic one and prove its equivalence to some others.

Theorem 10.1. *A graph T is a tree if and only if there is a unique path joining every two vertices of T .*

Proof. First, suppose that T is a tree, and let u and v be vertices of T . Since T is connected, there is a path, say P_1 , joining u and v . Now we must show that this path is unique. Assume to the contrary that there exists another path P_2 from u to v . When traversing P_1 from u to v , let w be the first vertex that is present on P_1 but not P_2 . Let x be the vertex on P_1 preceding w , and note that x is on P_2 as well. Let y be the next vertex common to both P_1 and P_2 when traversing P_1 from x to v . Then the portion of P_1 from x to y together with the portion of P_2 from y to x forms a cycle in the tree T , which is a contradiction. Thus, P_1 is the unique path joining u and v .

Conversely, suppose that T is a graph in which there is a unique path joining any two vertices. Clearly, T is connected. To show that T is acyclic, suppose that v_1, v_2, \dots, v_n is a cycle in T . Then we get two different paths joining v_1 and v_n , namely the path v_1, v_2, \dots, v_n and the path v_1, v_n (since $v_1 \sim v_n$ in the cycle). This contradicts our assumption. Thus, T must be acyclic and hence is a tree. \square

The next two results show that the size of a tree is always one less than its order, and that conversely, this property together with either connectedness or acyclicity implies that the graph is a tree.

Theorem 10.2. *A (p, q) -graph T is a tree if and only if it is connected and $p = q + 1$.*

Proof. Let T be a tree with p vertices and q edges. Then T is connected. We prove that $p = q + 1$ by induction. This is clearly true when $p = 1$. Assume it to be true for all trees of order less than p . Now in T , we know that every two vertices are joined by a unique path. Thus, if e is any edge of T , then the graph $T - \{e\}$ obtained by deleting e has exactly two components, say T_1 and T_2 . Each one is a tree, since it is connected and acyclic. Let T_i have p_i vertices and q_i edges, $i = 1, 2$. Then by the hypothesis, $p_i = q_i + 1$ (since $p_i < p$). But $p = p_1 + p_2$ and $q = q_1 + q_2 + 1$ (since the size of $T - \{e\}$ is one less than that of T). Thus, $p = q_1 + q_2 + 2 = q + 1$.

For the converse, suppose that T is a connected (p, q) -graph with $p = q + 1$. We must show that T is acyclic. Suppose to the contrary that T has a cycle C with k vertices. Then C has k edges as well. Since T is connected, there is a path from every vertex not on C to some vertex of C . The shortest path from each vertex v not on C to a vertex on C has a unique edge incident with v , which is not part of C . Since there are $p - k$ vertices in T not on C , there are $p - k$ such edges. Thus $q \geq (p - k) + k = p$, which contradicts our assumption that $p = q + 1$. Thus, T must be acyclic. \square

In the following theorem, the proof of the direct part is identical to that of Theorem 10.2, except for the assertion being about acyclicity rather than connectedness. The proof of the converse part is entirely different.

Theorem 10.3. *A (p, q) -graph T is a tree if and only if it is acyclic and $p = q + 1$.*

Proof. Let T be a tree with p vertices and q edges. Then T is acyclic. We prove that $p = q + 1$ by induction. This is clearly true when $p = 1$. Assume it to be true for all trees of order less than p . Now in T , we know that every two vertices are joined by a unique path. Thus, if e is any edge of T , then the graph $T - \{e\}$ obtained by deleting e has exactly two components, say T_1 and T_2 . Each one is a tree, since it is connected and acyclic. Let T_i have p_i vertices and q_i edges, $i = 1, 2$. Then by the hypothesis, $p_i = q_i + 1$ (since $p_i < p$). But $p = p_1 + p_2$ and $q = q_1 + q_2 + 1$ (since the size of $T - \{e\}$ is one less than that of T). Thus, $p = q_1 + q_2 + 2 = q + 1$.

Conversely, suppose that T is an acyclic (p, q) -graph with $p = q + 1$. To show that T is connected, we need to prove that it is connected – i.e., it has only one component. Let T have k components T_1, \dots, T_k . Each one is acyclic, and being connected, is a tree. Thus from the first part of the theorem, we know that if p_i and q_i are respectively the order and size of the component T_i , $p_i = q_i + 1$. Now $p = p_1 + \dots + p_k = (q_1 + 1) + \dots + (q_k + 1) = q + k$. But we know that $p = q + 1$. Therefore, $k = 1$. Thus, T is a tree. \square

Exercise 10.1. A **pendant vertex** of a graph is a vertex of degree 1. Prove that every non-trivial tree contains at least two pendant vertices.

Hint: Observe that a non-trivial tree cannot have a vertex of degree zero. Use Handshaking Lemma and assume every degree is at least 2 to get a contradiction.

Exercise 10.2. The **centre** of a graph G is the set of all vertices of G with minimum eccentricity – i.e., the set of all vertices v of G with $\text{ecc } v = \text{rad } v$. Show that every tree has a centre consisting of either exactly one vertex or exactly two adjacent vertices.


Hint: Observe that deleting all pendant vertices of a tree results in a new tree with the same centre.

Exercise 10.3. If G and H are two trees of orders n and m respectively, what is the size of $G \times H$?

11 Blocks

A **cutvertex** of a graph is a vertex whose removal increases the number of components, i.e., a vertex v of G such that $G - v$ has more components than G . If G is connected, we can equivalently say that v is a cutvertex if $G - v$ is disconnected. Similarly, a **cutedge** or **bridge** of a graph whose removal increases the number of components. A **nonseparable** graph is a connected, non-trivial graph with no cutvertices. A maximal nonseparable subgraph of a graph is a **block** of the graph. A nonseparable graph is itself said to be a block as well.

Example 11.1. The graph shown below has 6 blocks and 3 cutvertices (v_1, v_2, v_3).




The 6 blocks of this graph are shown below.

- (i) v is a cutvertex of G .
- (ii) There exist vertices u and w of G , distinct from v , such that every u - w path passes through v .
- (iii) There exists a partition of $V(G) - v$ into two non-empty subsets U and W such that for all $u \in U$ and $w \in W$, every u - w path passes through v .

(ii) \Rightarrow (i). Consider the graph $G - v$. As every $u-w$ path passes through v , none of them is present in $G - v$, and therefore $G - v$ is disconnected. Hence, v is a cutvertex of G . \square

Example 12.1. Fig. 2 shows a graph G with eight edges and its line graph $L(G)$.




11

13 Adjacency Matrices

The **adjacency matrix** of a graph G of order n , with vertex set $V = \{v_1, \dots, v_n\}$, is the $n \times n$ matrix $A = A(G)$ whose (i, j) -entry is

$$a_{ij} = \begin{cases} 1, & v_i \sim v_j \\ 0, & v_i \not\sim v_j. \end{cases}$$

Example 13.1. The adjacency matrix of the graph



is

$$A = \begin{bmatrix} 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Observe that, as the graphs we discuss are simple graphs and therefore have no self-loops on vertices, no vertex is adjacent to itself – i.e., $a_{ii} = 0$ for all $i = 1, \dots, n$. Also, since the graphs are undirected, $v_i \sim v_j$ if and only if $v_j \sim v_i$ – i.e., $a_{ij} = a_{ji}$. Thus, we have the following observation.

Observation 13.2. *The adjacency matrix of a (simple, undirected) graph is a symmetric, zero-diagonal, 0-1 matrix.*

In the i^{th} row of the adjacency matrix, for each j , the j^{th} entry is 1 if v_j is adjacent to v_i , and 0 otherwise. That is, the number of 1s in the i^{th} row is the number of vertices adjacent to v_i , or in other words, the degree of v_i . Thus, the row sums of A are the vertex degrees. Observe that $A\mathbb{1}$ is the vector of row sums, where $\mathbb{1}$ is the vector (of suitable size) with all entries equal to 1. For instance, with the matrix A given in Example 13.1,

$$A = \begin{bmatrix} 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 6 \\ 2 \\ 6 \\ 2 \\ 2 \\ 2 \\ 2 \end{bmatrix}.$$

From the Handshaking Lemma and the preceding observation, it follows that the sum of all entries of A is twice the number of edges of the graph.

Exercise 13.1. Show that the (i, j) -entry of A^2 is the number of walks of length 2 from v_i to v_j . Hence show that $\text{tr}(A^2) = 2|E(G)|$.

Hint: Recall that if A is any $n \times n$ matrix, then the (i, j) -entry of A^2 is $\sum_{k=1}^n a_{ik}a_{kj}$. As A is a 0-1 matrix, each term in this summation is 1 or 0, with the former if and only if $a_{ik} = a_{kj} = 1$. What does this imply about the vertices v_i , v_k , and v_j ? Then, as k varies from 1 to n , what does the value of the sum imply about v_i and v_j ?

The following result (which generalises the statement in Exercise 13.1) shows that the adjacency matrix can be used to obtain certain information about walks in the graph.

Theorem 13.3. Let A be the adjacency matrix of a graph G with vertex set $\{v_1, \dots, v_n\}$. Then the (i, j) -entry of A^m , for any positive integer m , is the number of walks of length k from v_i to v_j .

Proof. We prove the result by induction on m . For $m = 1$, the (i, j) -entry of $A^1 = A$ is a_{ij} , which is 1 if and only if v_i is adjacent to v_j , i.e., if and only if there is a walk of length 1 (namely, an edge) from v_i to v_j . Thus, the result holds for $m = 1$.

Now suppose, for the sake of induction, that the result holds for some $m \geq 1$, and consider A^{m+1} . The (i, j) -entry of A^{m+1} is


$$(A^{m+1})_{ij} = \sum_{k=1}^n (A^m)_{ik}a_{kj}.$$

First, note that $a_{kj} = 1$ if and only if $v_k \sim v_j$. Therefore, the above sum is equal to the sum of all $(A^m)_{ik}$ where $v_k \sim v_j$. Now, by the induction hypothesis, $(A^m)_{ik}$ is the number of walks of length m from v_i to v_k . If v_k is adjacent to v_j , then each walk of length m from v_i to v_k , together with the edge from v_k to v_j , forms a walk of length $m + 1$ from v_i to v_j . Thus, for each k such that $v_k \sim v_j$, $(A^m)_{ik}a_{kj} = (A^m)_{ik}$ is the number of walks of length $m + 1$ from v_i to v_j that pass through k . Summing over all k , this gives the total number of walks of length $m + 1$ from v_i to v_j . Hence the result follows by induction. \square

14 Incidence Matrices

Consider an (n, m) -graph G having vertex set $\{v_1, \dots, v_n\}$ and edge set $\{e_1, \dots, e_m\}$. The **incidence matrix** of G is the $n \times m$ matrix $B = B(G)$ whose (i, j) -entry is 1 if the vertex v_i is incident with the edge e_j , and 0 otherwise.

Example 14.1. The incidence matrix of the graph



is

$$B = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Each column of the incidence matrix corresponds to an edge of the graph, and the only non-zero entries in the column correspond to the end-vertices of the edge – thus, each column contains exactly two 1s. In a simple graph, there is at most one edge between a given pair of vertices, and hence no two columns of the incidence matrix can be equal. We therefore have the following observation.

Observation 14.2. *The incidence matrix of a (simple, undirected) graph is a 0-1 matrix in which every column has exactly two 1s, and no two columns are equal.*