

MARK SCHEME for the May/June 2011 question paper
for the guidance of teachers

9709 MATHEMATICS

9709/11

Paper 1, maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	11

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol $\checkmark$ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	11

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only often written by a ‘fortuitous’ answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through $\sqrt{}$ ” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	11

1 ${}^7C_2 x^5 \left(\frac{2}{x^2}\right)^2$ SOI and leading to final answer 84 or $84x$ as final answer	B2 B1 [3]	B1 for 2/3 parts correct leading to ans. If no answer; $84x$ seen scores B2, else ${}^7C_2 x^5 \left(\frac{2}{x^2}\right)^2$ scores SCB1 only
2 $\left(\frac{dv}{dr}\right)4\pi r^2$ $= 4\pi \times 10^2$ $\frac{dr}{dt} = \frac{dv}{dt} \frac{dv}{dr}$ OE used $\frac{50}{4\pi \times 10^2} = \frac{1}{8\pi}$ or 0.0398	M1 A1 M1 A1 [4]	SOI at any point Correct link between differentials with $\frac{dr}{dt}$ finally as subject Allow $\frac{50}{400\pi}$. Non-calculus methods $\frac{0}{4}$
3 (i) Correct shape – touching positive x-axis (ii) $(\pi) \int (x-2)^4 dx$ $(\pi) \left[\frac{(x-2)^5}{5} \right]$ $(\pi)[0 - (-32)/5]$ $\frac{32\pi}{5}$ or 6.4π	B1 [1] M1 A1 M1 A1 [4]	Ignore intersections with axes Use $(\pi) \int y^2 dx$ & attempt integrate but expansion before integ needs 5 terms Use of limits 0, 2 on <i>their</i> $(\pi) \int y^2 dx$ cao Rotation about y-axis max 1/5
4 (i) $\overrightarrow{CP} = -6\mathbf{i} + 6\mathbf{j} - 2\mathbf{k}$ $\overrightarrow{CQ} = -6\mathbf{i} + 6\mathbf{j} + 3\mathbf{k}$ (ii) Scalar product = $36 + 36 - 6$ $66 = \overrightarrow{CP} \overrightarrow{CQ} \cos \theta$ $\overrightarrow{CP} = \sqrt{76}$, $\overrightarrow{CQ} = \sqrt{81}$ Angle $PCQ = 32.7^\circ$ (or 0.571 rad)	B1 B1 [2] M1 M1 M1 A1 [4]	Use of $x_1x_2 + y_1y_2 + z_1z_2$ Linking everything correctly Correct magnitude for either cao 147.3° converted to 32.7° gets A0

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	11

5 (i) $\frac{2\sin^2 \theta \sin^2 \theta}{1 - \sin^2 \theta} = 1$ $2\sin^4 \theta + \sin^2 \theta - 1 = 0$ AG (ii) $(2\sin^2 \theta - 1)(\sin^2 \theta + 1) = 0$ $\sin \theta = \frac{(\pm)1}{\sqrt{2}}$ $\theta = 45^\circ, 135^\circ$ $\theta = 225^\circ, 315^\circ$	M1 A1 [2] M1 A1 A1 A1 [4]	Equation as function of $\sin \theta$ Or use formula on quadratic in $\sin^2 \theta$ Provided no excess solutions in range
6 (i) $z = 3x + 2\left(\frac{600}{x}\right)$ or $x\frac{(z-3x)}{2} = 600$ OE → AG (ii) $\frac{dz}{dx} = 3 - \frac{1200}{x^2}$ or $\frac{dz}{dy} = 2 - \frac{1800}{y^2}$ $= 0 \rightarrow x = 20$ or $= 0 \rightarrow y = 30$ $z = 60 + \frac{120}{20} = 120$ $\frac{d^2z}{dx^2} = \frac{2400}{x^3}$ $> 0 \Rightarrow$ minimum	B1 [1] B1 M1A1 A1√ B1√ B1 [6]	Set to 0 & attempt to solve. Allow ± 20 Ft from <i>their</i> x provided positive Or other valid method Dep. on $\frac{d^2z}{dx^2} = \frac{k}{x^3}$ ($k > 0$) or other valid method.
7 (i) $\frac{3(1+2x)^{-1}}{-1} + (c)$ $y = \frac{3(1+2x)^{-1}}{-2} + (c)$ Sub (1, (1/2)) $\frac{1}{2} = \frac{3}{-6} + c \Rightarrow c = 1$ (ii) $(1+2x)^2(>)9$ or $4x^2 + 4x - 8(>)0$ OE 1, -2 $x > 1, x < -2$ ISW	B1 B1(indep) M1 A1 [4] M1 A1 A1 [3]	Division by 2 $y =$ necessary Dependent on c present Use of $y = mx + c$ etc. gets 0/4
8 (i) 1000, 2000, 3000... or 50, 100, 150... $\frac{40}{2(1000 + 40000)}$ or $\frac{40}{2(2000 + 39000)}$ × 5% of attempt at valid sum 41000 (ii) 1000, 1000×1.1, $1000 \times 1.1^2 + \dots$ or with $a = 50$ $\frac{1000(1.1^{40} - 1)}{1.1 - 1}$ 22100	M1 M1 M1 A1 [4] M1 M1 A1 [3]	Recognise series, correct a/d (or 3 terms) Correct use of formula Can be awarded in either (i) or (ii) cao Recognise series, correct a/r (or 3 terms) Correct use of formula. Allow e.g. $r = 0.1$ Or answers rounding to this

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	11

9 (i) $AS = r \tan \theta$ Area $OAB = r^2 \tan \theta$ or $(OAS) = \frac{1}{2} r^2 \tan \theta$ Area of sector $= \frac{1}{2} r^2 \times 2\theta (= r^2 \theta)$ Shaded area $= r^2 (\tan \theta - \theta)$ OE (ii) $\cos \frac{\pi}{3} = \frac{6}{OA} \Rightarrow OA = 12$ $AP = 6$ $AS = 6 \tan \frac{\pi}{3} (\Rightarrow AB = 12\sqrt{3})$ Arc $(PST) = 12 \frac{\pi}{3}$ Perimeter $= 12 + 12\sqrt{3} + 4\pi$	M1 A1 B1 A1 [4] M1 A1 B1 B1 A1 [5]	Or $(AB) = 2r \tan \theta$ or $(AO) = \frac{r}{\cos \theta}$ Or $OAB = \frac{1}{2} \frac{r^2}{\cos 2\theta} \sin 2\theta$ Or area sector $(OPS) = \frac{1}{2} r^2 \theta$ Allow e.g. $r^2 \tan \theta - \frac{1}{2} r^2 2\theta$ Or arc $(PS) = 6 \frac{\pi}{3}$ or arc $(ST) = 6 \frac{\pi}{3}$ Allow unsimplified 4π
10 (i) $2(x-1)^2 - 1$ OR $a = 2, b = -1, c = -1$ $A = (1, -1)$ (ii) $2x^2 - 5x - 3 = 0 \Rightarrow (2x+1)(x-3) = 0$ OE in y $x = -\frac{1}{2}, y = 3\frac{1}{2}$ (iii) Mid-point of $AP = (2, 3)$ Gradient of line $= \frac{\frac{1}{2}}{\frac{-5}{2}} = \frac{-1}{5}$ Equation is $y - 3 = \frac{-1}{5}(x - 2)$ OE	B1, B1, B1 B1✓ [4] M1, M1 A1 [3] B1✓ B1 B1 [3]	Allow alt. method for final mark Complete elim & simplify, attempt soln. Additional (3, 7) not penalised Follow through on <i>their A</i> Or $y - 3\frac{1}{2} = -\frac{1}{5(x + \frac{1}{2})}$
11 (i) $fg(x) = 2x^2 - 3, gf(x) = 4x^2 + 4x - 1$ (ii) $2a^2 - 3 = 4a^2 + 4a - 1 \Rightarrow 2a^2 + 4a + 2 = 0$ $(a+1)^2 = 0$ $a = -1$ (iii) $b^2 - b - 2 = 0 \rightarrow (b+1)(b-2) = 0$ $b = 2$ Allow $b = -1$ in addition (iv) $f^{-1}(x) = \frac{1}{2(x-1)}$ $f^{-1}g(x) = \frac{1}{2(x^2-3)}$ (v) $x = (\pm)\sqrt{y+2}$ $h^{-1}(x) = -\sqrt{x+2}$	B1, B1 [2] M1 M1 A1 [3] M1 A1 [2] B1 B1✓ [2] M1 A1 [2]	fg & gf clearly transposed gets B0B0 Dep. quadratic. Allow x for all 3 marks Allow marks in (ii) if transposed in (i) Allow in terms of x for M1 only Correct answer without working B2 Must be simplified. Ft from <i>their</i> f^{-1}