




mod1 sharks
(n=1130 surveyed, 305 with MaxN>0)




mod2 pelagic-piscivores
(n=1130 surveyed, 908 with MaxN>0)




mod3 snappers
(n=1130 surveyed, 1011 with MaxN>0)




mod4 gag-0
(n=1130 surveyed, 39 with MaxN>0)




mod5 gag-1
(n=1130 surveyed, 145 with MaxN>0)




mod6 gag-2
(n=1130 surveyed, 190 with MaxN>0)




mod7 gag-3
(n=1130 surveyed, 192 with MaxN>0)




mod8 gag-4
(n=1130 surveyed, 176 with MaxN>0)




mod9 gag-5-
(n=1130 surveyed, 342 with MaxN>0)




mod10 red-grouper-0
(n=1130 surveyed, 108 with MaxN>0)




mod11 red-grouper-1
(n=1130 surveyed, 10 with MaxN>0)




mod12 red-grouper-2
(n=1130 surveyed, 41 with MaxN>0)




mod13 red-grouper-3
(n=1130 surveyed, 162 with MaxN>0)




mod14 red-grouper-4
(n=1130 surveyed, 262 with MaxN>0)




mod15 red-grouper-5-
(n=1130 surveyed, 695 with MaxN>0)




mod16 large-reef-fish
(n=1130 surveyed, 614 with MaxN>0)




mod17 small-reef-fish
(n=1130 surveyed, 1101 with MaxN>0)



mod18 demersal-fish
(n=1130 surveyed, 734 with MaxN>0)



mod19 small-pelagics
(n=1130 surveyed, 715 with MaxN>0)



availability coverage

- good
- partial
- poor (e.g. mud — not identifiable)

- constrained ratio used for affinity ($MUD=0$, sand < rock & gravel)
- family divider (reef | bottom); A is scaled by one global maximum