




mod1 sharks
(n=1131 surveyed, 305 with MaxN>0)




mod2 pelagic-piscivores
(n=1131 surveyed, 908 with MaxN>0)




mod3 snappers
(n=1131 surveyed, 1011 with MaxN>0)




mod4 gag-0
(n=1131 surveyed, 42 with MaxN>0)




mod5 gag-1
(n=1131 surveyed, 144 with MaxN>0)




mod6 gag-2
(n=1131 surveyed, 179 with MaxN>0)







mod7 gag-3
(n=1131 surveyed, 183 with MaxN>0)




mod8 gag-4
(n=1131 surveyed, 186 with MaxN>0)







mod13 red-grouper-3
(n=1131 surveyed, 153 with MaxN>0)




mod14 red-grouper-4
(n=1131 surveyed, 266 with MaxN>0)




mod15 red-grouper-5-
(n=1131 surveyed, 691 with MaxN>0)




mod16 large-reef-fish
(n=1131 surveyed, 614 with MaxN>0)




mod17 small-reef-fish
(n=1131 surveyed, 1101 with MaxN>0)



mod18 demersal-fish
(n=1131 surveyed, 734 with MaxN>0)



mod19 small-pelagics
(n=1131 surveyed, 715 with MaxN>0)



availability coverage

- good
- partial
- poor (e.g. mud — not identifiable)

- constrained ratio used for affinity ($MUD=0$, sand < rock & gravel)
- family divider (reef | bottom); A is scaled by one global maximum