




mod1 sharks
(426 stations with MaxN>0)




mod2 pelagic piscivores
(4663 stations with MaxN>0)




mod3 snappers
(6999 stations with MaxN>0)




mod4 gag 0 [pooled: gag 1]
(159 stations with MaxN>0)




mod5 gag 1
(139 stations with MaxN>0)




mod6 gag 2
(189 stations with MaxN>0)




mod7 gag 3
(170 stations with MaxN>0)




mod8 gag 4
(162 stations with MaxN>0)




mod9 gag 5+
(478 stations with MaxN>0)




mod10 red grouper 0
(159 stations with MaxN>0)




mod11 red grouper 1 [pooled: red grouper 0]
(165 stations with MaxN>0)




mod12 red grouper 2 [pooled: red grouper 1]
(190 stations with MaxN>0)




mod13 red grouper 3
(185 stations with MaxN>0)




mod14 red grouper 4
(373 stations with MaxN>0)




mod15 red grouper 5+
(2433 stations with MaxN>0)




mod16 large reef fish
(2071 stations with MaxN>0)




mod17 small reef fish
(9322 stations with MaxN>0)



mod18 demersal fish
(2407 stations with MaxN>0)



mod19 small pelagics
(2214 stations with MaxN>0)



depth/region adjustment

- stratified by SPACE_STRAT (design-based)
- depth as continuous covariate (quasi-Poisson GLM, G-computation)

bars agreeing across both adjustments = habitat signal; disagreeing = depth artifact

..... family divider – affinity is normalized within reef and within substrate