




**mod1 sharks**  
(426 stations with MaxN>0)




**mod2 pelagic piscivores**  
(4663 stations with MaxN>0)




**mod3 snappers**  
(6999 stations with MaxN>0)




**mod4 gag 0 [pooled: gag 1]**  
(153 stations with MaxN>0)




**mod5 gag 1**  
(139 stations with MaxN>0)




**mod6 gag 2**  
(183 stations with MaxN>0)




**mod7 gag 3**  
(173 stations with MaxN>0)




**mod8 gag 4**  
(158 stations with MaxN>0)




**mod9 gag 5+**  
(507 stations with MaxN>0)




**mod10 red grouper 0**  
(159 stations with MaxN>0)




**mod11 red grouper 1 [pooled: red grouper 0]**  
(163 stations with MaxN>0)




**mod12 red grouper 2**  
(34 stations with MaxN>0)




**mod13 red grouper 3**  
(157 stations with MaxN>0)




**mod14 red grouper 4**  
(381 stations with MaxN>0)




**mod15 red grouper 5+**  
(2430 stations with MaxN>0)




**mod16 large reef fish**  
(2071 stations with MaxN>0)




**mod17 small reef fish**  
(9322 stations with MaxN>0)



**mod18 demersal fish**  
(2407 stations with MaxN>0)



**mod19 small pelagics**  
(2214 stations with MaxN>0)



depth/region adjustment

- stratified by SPACE\_STRAT (design-based)
- depth as continuous covariate (quasi-Poisson GLM, G-computation)

bars agreeing across both adjustments = habitat signal; disagreeing = depth artifact

..... family divider – affinity is normalized within reef and within substrate