

th the largest gain in samples included, then the 8 with samples >= 10,000 cells/L but no such cell (page 1 of 3): samples | MODIS−clipped (the 6 Oct 2020

2 in the titles = bloom area >= 10,000 cells/L (+ = added by the hybrid). Orange = cells the hybrid changed. Purple shading in the hull column = the mask5 set (VIIRS no−data mask or within 5 km of the coa
mples: fill = FWC class (viridis, light = background to dark = > 1,000,000 cells/L); red border = a water sample >= 10,000 cells/L outside the MODIS footprint. Maps: pipeline palette, white below 10,000 cells


2006−07: FWC samples (fit fitted, 19 nFLH polygons)




before (MODIS−clipped): 9,702 km2




after (hybrid mask5): 10,615 km2 (+913)




hull−clipped: 7,310 km2 (set shaded)




2003−02: FWC samples (fit fitted, 51 nFLH polygons)




before (MODIS−clipped): 11,741 km2




after (hybrid mask5): 12,278 km2 (+537)




hull−clipped: 3,769 km2 (set shaded)




2003−08: FWC samples (fit fitted, 41 nFLH polygons)




before (MODIS−clipped): 18,709 km2




after (hybrid mask5): 19,854 km2 (+1,145)




hull−clipped: 10,798 km2 (set shaded)




2002−08: FWC samples (fit fitted, 49 nFLH polygons)




before (MODIS−clipped): 11,509 km2




after (hybrid mask5): 12,422 km2 (+913)




hull−clipped: 6,840 km2 (set shaded)




2003−07: FWC samples (fit fitted, 45 nFLH polygons)




before (MODIS−clipped): 14,306 km2




after (hybrid mask5): 15,366 km2 (+1,060)




hull−clipped: 14,048 km2 (set shaded)




2007−01: FWC samples (fit fitted, 27 nFLH polygons)




before (MODIS−clipped): 17,902 km2




after (hybrid mask5): 18,360 km2 (+458)




hull−clipped: 6,878 km2 (set shaded)




2003−05: FWC samples (fit fitted, 31 nFLH polygons)




before (MODIS−clipped): 11,896 km2




after (hybrid mask5): 12,888 km2 (+992)




hull−clipped: 8,250 km2 (set shaded)




2005−09: FWC samples (fit fitted, 32 nFLH polygons)




before (MODIS−clipped): 79,367 km2



after (hybrid mask5): 80,330 km2 (+963)




hull−clipped: 48,234 km2 (set shaded)




th the largest gain in samples included, then the 8 with samples >= 10,000 cells/L but no such cell (page 2 of 3): samples | MODIS−clipped (the 6 Oct 2020

2 in the titles = bloom area >= 10,000 cells/L (+ = added by the hybrid). Orange = cells the hybrid changed. Purple shading in the hull column = the mask5 set (VIIRS no−data mask or within 5 km of the coa
mples: fill = FWC class (viridis, light = background to dark = > 1,000,000 cells/L); red border = a water sample >= 10,000 cells/L outside the MODIS footprint. Maps: pipeline palette, white below 10,000 cells


2003−06: FWC samples (fit fitted, 59 nFLH polygons)




before (MODIS−clipped): 11,631 km2




after (hybrid mask5): 12,545 km2 (+914)




hull−clipped: 8,379 km2 (set shaded)




2006−08: FWC samples (fit fitted, 37 nFLH polygons)




before (MODIS−clipped): 14,754 km2




after (hybrid mask5): 15,593 km2 (+839)




hull−clipped: 12,888 km2 (set shaded)




2003−04: FWC samples (fit fitted, 30 nFLH polygons)




before (MODIS−clipped): 13,049 km2




after (hybrid mask5): 14,191 km2 (+1,142)




hull−clipped: 13,829 km2 (set shaded)




2025−01: FWC samples (fit fitted, 38 nFLH polygons)




before (MODIS−clipped): 23,519 km2




after (hybrid mask5): 24,427 km2 (+908)




hull−clipped: 19,030 km2 (set shaded)




2006−01: FWC samples (fit fitted, 28 nFLH polygons)




before (MODIS−clipped): 0 km2




after (hybrid mask5): 0 km2 (+0)




hull−clipped: 0 km2 (set shaded)




2008−02: FWC samples (fit none, 32 nFLH polygons)




before (MODIS−clipped): 0 km2




after (hybrid mask5): 0 km2 (+0)




hull−clipped: 0 km2 (set shaded)




2008−03: FWC samples (fit none, 35 nFLH polygons)




before (MODIS−clipped): 0 km2




after (hybrid mask5): 0 km2 (+0)




hull−clipped: 0 km2 (set shaded)




2008−04: FWC samples (fit fitted, 27 nFLH polygons)




before (MODIS−clipped): 0 km2



after (hybrid mask5): 0 km2 (+0)




hull−clipped: 0 km2 (set shaded)




th the largest gain in samples included, then the 8 with samples >= 10,000 cells/L but no such cell (page 3 of 3): samples | MODIS−clipped (the 6 Oct 2020

2 in the titles = bloom area >= 10,000 cells/L (+ = added by the hybrid). Orange = cells the hybrid changed. Purple shading in the hull column = the mask5 set (VIIRS no−data mask or within 5 km of the coa
mples: fill = FWC class (viridis, light = background to dark = > 1,000,000 cells/L); red border = a water sample >= 10,000 cells/L outside the MODIS footprint. Maps: pipeline palette, white below 10,000 cells


2009−04: FWC samples (fit fitted, 27 nFLH polygons)




before (MODIS−clipped): 0 km2




after (hybrid mask5): 153 km2 (+153)




hull−clipped: 229 km2 (set shaded)




2010−01: FWC samples (fit fitted, 71 nFLH polygons)




before (MODIS−clipped): 0 km2




after (hybrid mask5): 0 km2 (+0)




hull−clipped: 0 km2 (set shaded)




2010−02: FWC samples (fit fitted, 80 nFLH polygons)




before (MODIS−clipped): 0 km2




after (hybrid mask5): 0 km2 (+0)




hull−clipped: 0 km2 (set shaded)




2011−08: FWC samples (fit fitted, 27 nFLH polygons)



before (MODIS−clipped): 0 km2



after (hybrid mask5): 0 km2 (+0)



hull−clipped: 0 km2 (set shaded)

