


1985-09 | source used: hull | VIIRS layer: no coverage

samples 42, positives 9, $\geq 10,000$ cells/L: 5, $\geq 100,000$: 4 (max 930,000 cells/L) | final bloom area = 100% of the hull-clipped one | 5 of 5 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,093 km², max 363,485 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 4,199 km²)
bloom area 2,338 km²



1985-10 | source used: hull | VIIRS layer: no coverage

samples 43, positives 22, $\geq 10,000$ cells/L: 11, $\geq 100,000$: 9 (max 1,200,000 cells/L) | final bloom area = 100% of the hull-clipped one | 11 of 11 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,221 km2, max 820,865 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 7,178 km2)
bloom area 5,104 km2




1985-11 | source used: hull | VIIRS layer: no coverage

samples 53, positives 12, >= 10,000 cells/L: 5, >= 100,000: 4 (max 195,000 cells/L) | final bloom area = 100% of the hull-clipped one | 5 of 5 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 159,148 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 5,319 km2)
bloom area 4,104 km2




1986-09 | source used: hull | VIIRS layer: no coverage

samples 105, positives 53, $\geq 10,000$ cells/L: 37, $\geq 100,000$: 27 (max $9e+05$ cells/L) | final bloom area = 100% of the hull-clipped one | 37 of 37 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 1,506,650 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 7,786 km²)
bloom area 6,842 km²




1986-10 | source used: hull | VIIRS layer: no coverage

samples 88, positives 56, $\geq 10,000$ cells/L: 44, $\geq 100,000$: 41 (max 7,530,000 cells/L) | final bloom area = 100% of the hull-clipped one | 43 of 43 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 288,327 km², max 2,129,701 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 8,449 km²)
bloom area 4,702 km²




1986-11 | source used: hull | VIIRS layer: no coverage

samples 26, positives 9, >= 10,000 cells/L: 4, >= 100,000: 2 (max 9e+05 cells/L) | final bloom area = 100% of the hull-clipped one | 3 of 3 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,525 km2, max 396,991 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 3,431 km2)
bloom area 686 km2




1986-12 | source used: hull | VIIRS layer: no coverage

samples 57, positives 22, >= 10,000 cells/L: 6, >= 100,000: 2 (max 210,000 cells/L) | final bloom area = 100% of the hull-clipped one | 6 of 6 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,754 km2, max 51,010 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 3,561 km2)
bloom area 2,124 km2




1987-01 | source used: hull | VIIRS layer: no coverage

samples 109, positives 24, $\geq 10,000$ cells/L: 6, $\geq 100,000$: 2 (max 530,000 cells/L) | final bloom area = 100% of the hull-clipped one | 6 of 6 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 2,359 km², max 57,730 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 5,727 km²)
bloom area 1,597 km²




1987-02 | source used: hull | VIIRS layer: no coverage

samples 121, positives 23, $\geq 10,000$ cells/L: 8, $\geq 100,000$: 4 (max 1,850,000 cells/L) | final bloom area = 100% of the hull-clipped one | 8 of 8 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 314,481 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 7,198 km²)
bloom area 6,396 km²



1987-05 | source used: hull | VIIRS layer: no coverage

samples 147, positives 61, $\geq 10,000$ cells/L: 32, $\geq 100,000$: 13 (max 5×10^7 cells/L) | final bloom area = 100% of the hull-clipped one | 30 of 30 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 1,806,128 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 11,236 km²)
bloom area 9,879 km²




1987-06 | source used: hull | VIIRS layer: no coverage

samples 138, positives 30, $\geq 10,000$ cells/L: 13, $\geq 100,000$: 4 (max 150,000 cells/L) | final bloom area = 100% of the hull-clipped one | 12 of 12 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 39,229 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 6,418 km²)
bloom area 5,396 km²




1987-08 | source used: hull | VIIRS layer: no coverage

samples 96, positives 14, $\geq 10,000$ cells/L: 2, $\geq 100,000$: 0 (max 67,000 cells/L) | final bloom area = -% of the hull-clipped one | 2 of 2 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 3,388 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 3,444 km²)
bloom area 0 km²




1987-09 | source used: hull | VIIRS layer: no coverage

samples 133, positives 24, $\geq 10,000$ cells/L: 6, $\geq 100,000$: 1 (max 277,000 cells/L) | final bloom area = 100% of the hull-clipped one | 6 of 6 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 19,418 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 3,977 km²)
bloom area 2,892 km²




1988-10 | source used: hull | VIIRS layer: no coverage

samples 118, positives 40, $\geq 10,000$ cells/L: 16, $\geq 100,000$: 9 (max 970,000 cells/L) | final bloom area = 100% of the hull-clipped one | 16 of 16 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 184,304 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 9,207 km²)
bloom area 7,777 km²




1988-11 | source used: hull | VIIRS layer: no coverage

samples 115, positives 22, $\geq 10,000$ cells/L: 12, $\geq 100,000$: 6 (max 4,100,000 cells/L) | final bloom area = 100% of the hull-clipped one | 12 of 12 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 1,204,796 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 8,564 km²)
bloom area 8,310 km²




1989-03 | source used: hull | VIIRS layer: no coverage

samples 82, positives 16, >= 10,000 cells/L: 11, >= 100,000: 3 (max 503,000 cells/L) | final bloom area = 100% of the hull-clipped one | 10 of 10 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,057 km2, max 378,921 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 3,381 km2)
bloom area 2,196 km2




1989-04 | source used: hull | VIIRS layer: no coverage

samples 106, positives 30, >= 10,000 cells/L: 14, >= 100,000: 7 (max 850,000 cells/L) | final bloom area = 100% of the hull-clipped one | 9 of 9 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 165,752 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 3,043 km2)
bloom area 2,271 km2



1989-05 | source used: hull | VIIRS layer: no coverage

samples 71, positives 6, >= 10,000 cells/L: 0, >= 100,000: 0 (max 6,700 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 1,027 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (4 footprint(s), 1,545 km2)
bloom area 0 km2




1990-02 | source used: hull | VIIRS layer: no coverage

samples 75, positives 11, >= 10,000 cells/L: 5, >= 100,000: 4 (max 233,000 cells/L) | final bloom area = 100% of the hull-clipped one | 3 of 3 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 299,247 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 2,378 km2)
bloom area 1,893 km2




1990-07 | source used: hull | VIIRS layer: no coverage

samples 66, positives 13, >= 10,000 cells/L: 5, >= 100,000: 0 (max 30,000 cells/L) | final bloom area = 100% of the hull-clipped one | 5 of 5 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 3,179 km2, max 16,568 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (5 footprint(s), 6,441 km2)
bloom area 303 km2




1990-08 | source used: hull | VIIRS layer: no coverage

samples 69, positives 10, >= 10,000 cells/L: 0, >= 100,000: 0 (max 9,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 2,840 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (8 footprint(s), 2,970 km2)
bloom area 0 km2




1990-09 | source used: hull | VIIRS layer: no coverage

samples 83, positives 35, $\geq 10,000$ cells/L: 19, $\geq 100,000$: 1 (max 106,000 cells/L) | final bloom area = 100% of the hull-clipped one | 19 of 19 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 20,794 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 64,007 km²)
bloom area 62,472 km²




1991-01 | source used: hull | VIIRS layer: no coverage

samples 123, positives 37, $\geq 10,000$ cells/L: 13, $\geq 100,000$: 0 (max 87,000 cells/L) | final bloom area = 100% of the hull-clipped one | 9 of 9 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 15,898 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 3,211 km²)
bloom area 2,279 km²




1991-02 | source used: hull | VIIRS layer: no coverage

samples 60, positives 6, >= 10,000 cells/L: 0, >= 100,000: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 9,202 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (6 footprint(s), 1,884 km2)
bloom area 0 km2



1991-07 | source used: hull | VIIRS layer: no coverage

samples 83, positives 19, $\geq 10,000$ cells/L: 18, $\geq 100,000$: 18 (max 1,800,000 cells/L) | final bloom area = 100% of the hull-clipped one | 14 of 14 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 1,690,810 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 2,109 km²)
bloom area 1,370 km²



1991-08 | source used: hull | VIIRS layer: no coverage

samples 292, positives 186, $\geq 10,000$ cells/L: 145, $\geq 100,000$: 116 (max 27,300,000 cells/L) | final bloom area = 100% of the hull-clipped one | 117 of 117 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,207 km², max 1,053,226 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 11,204 km²)
bloom area 9,967 km²




1991-09 | source used: hull | VIIRS layer: no coverage

samples 132, positives 63, >= 10,000 cells/L: 48, >= 100,000: 24 (max 925,000 cells/L) | final bloom area = 100% of the hull-clipped one | 33 of 33 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,765 km2, max 190,234 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 4,243 km2)
bloom area 1,902 km2




1991-10 | source used: hull | VIIRS layer: no coverage

samples 93, positives 18, $\geq 10,000$ cells/L: 10, $\geq 100,000$: 0 (max 55,000 cells/L) | final bloom area = 100% of the hull-clipped one | 4 of 4 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 22,467 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 3,399 km²)
bloom area 2,280 km²




1992-09 | source used: hull | VIIRS layer: no coverage

samples 280, positives 152, >= 10,000 cells/L: 129, >= 100,000: 109 (max 1.05e+08 cells/L) | final bloom area = 100% of the hull-clipped one | 104 of 104 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 10,899,744 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 8,753 km2)
bloom area 7,777 km2




1992-10 | source used: hull | VIIRS layer: no coverage

samples 106, positives 23, $\geq 10,000$ cells/L: 11, $\geq 100,000$: 5 (max 1,900,000 cells/L) | final bloom area = 100% of the hull-clipped one | 10 of 10 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 233,693 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 3,693 km²)
bloom area 2,436 km²




1992-11 | source used: hull | VIIRS layer: no coverage

samples 162, positives 61, $\geq 10,000$ cells/L: 45, $\geq 100,000$: 39 (max 2,900,000 cells/L) | final bloom area = 100% of the hull-clipped one | 28 of 28 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 2,128,102 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 5,032 km²)
bloom area 3,961 km²




1992-12 | source used: hull | VIIRS layer: no coverage

samples 126, positives 23, $\geq 10,000$ cells/L: 7, $\geq 100,000$: 5 (max 1,160,000 cells/L) | final bloom area = 100% of the hull-clipped one | 6 of 6 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 124,890 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 4,804 km²)
bloom area 3,586 km²



1994-09 | source used: hull | VIIRS layer: no coverage

samples 189, positives 101, $\geq 10,000$ cells/L: 91, $\geq 100,000$: 54 (max 4.8×10^7 cells/L) | final bloom area = 100% of the hull-clipped one | 75 of 75 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 25,904,724 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 3,085 km²)
bloom area 2,130 km²



1994-10 | source used: hull | VIIRS layer: no coverage

samples 209, positives 120, >= 10,000 cells/L: 88, >= 100,000: 51 (max 5,400,000 cells/L) | final bloom area = 100% of the hull-clipped one | 56 of 56 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 802,170 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 7,073 km2)
bloom area 5,561 km2




1994-11 | source used: hull | VIIRS layer: no coverage

samples 200, positives 111, $\geq 10,000$ cells/L: 81, $\geq 100,000$: 36 (max 68,500,000 cells/L) | final bloom area = 100% of the hull-clipped one | 48 of 48 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 253,281 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 3,552 km²)
bloom area 2,512 km²




1994-12 | source used: hull | VIIRS layer: no coverage

samples 186, positives 129, >= 10,000 cells/L: 114, >= 100,000: 90 (max 3.58e+08 cells/L) | final bloom area = 100% of the hull-clipped one | 76 of 76 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,907 km2, max 7,139,824 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 4,138 km2)
bloom area 2,893 km2




1995-01 | source used: hull | VIIRS layer: no coverage

samples 117, positives 33, $\geq 10,000$ cells/L: 13, $\geq 100,000$: 5 (max 1,480,000 cells/L) | final bloom area = 100% of the hull-clipped one | 11 of 11 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,375 km², max 1,094,972 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 7,636 km²)
bloom area 6,110 km²




1995-02 | source used: hull | VIIRS layer: no coverage

samples 81, positives 12, >= 10,000 cells/L: 7, >= 100,000: 4 (max 280,000 cells/L) | final bloom area = 100% of the hull-clipped one | 7 of 7 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 469,478 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 1,294 km2)
bloom area 1,004 km2




1995-04 | source used: hull | VIIRS layer: no coverage

samples 206, positives 106, $\geq 10,000$ cells/L: 85, $\geq 100,000$: 59 (max 81,700,000 cells/L) | final bloom area = 100% of the hull-clipped one | 52 of 52 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 1,305,792 cells/L




VIIRS-clipped



MODIS nFLH-clipped



USED: Hull-clipped (1 footprint(s), 3,406 km²)
bloom area 2,589 km²



1995-05 | source used: hull | VIIRS layer: no coverage

samples 304, positives 193, >= 10,000 cells/L: 175, >= 100,000: 138 (max 17,240,000 cells/L) | final bloom area = 100% of the hull-clipped one | 100 of 100 water samples >= 10,000 cells/L inside the final footprint






Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 1,408,146 cells/L

VIIRS-clipped

MODIS nFLH-clipped

USED: Hull-clipped (1 footprint(s), 6,769 km2)
bloom area 5,009 km2


cells/L



1995-06 | source used: hull | VIIRS layer: no coverage

samples 254, positives 137, $\geq 10,000$ cells/L: 106, $\geq 100,000$: 45 (max 51,930,000 cells/L) | final bloom area = 100% of the hull-clipped one | 66 of 66 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,285 km², max 1,174,509 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 11,230 km²)
bloom area 9,798 km²



1995-07 | source used: hull | VIIRS layer: no coverage

samples 173, positives 65, $\geq 10,000$ cells/L: 56, $\geq 100,000$: 18 (max 4,170,000 cells/L) | final bloom area = 100% of the hull-clipped one | 34 of 34 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 464,421 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 4,881 km²)
bloom area 3,641 km²




1995-08 | source used: hull | VIIRS layer: no coverage

samples 314, positives 193, $\geq 10,000$ cells/L: 140, $\geq 100,000$: 77 (max 64,843,000 cells/L) | final bloom area = 100% of the hull-clipped one | 93 of 93 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,395 km², max 630,438 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 15,063 km²)
bloom area 11,718 km²




1995-09 | source used: hull | VIIRS layer: no coverage

samples 388, positives 230, $\geq 10,000$ cells/L: 170, $\geq 100,000$: 101 (max 20,417,000 cells/L) | final bloom area = 100% of the hull-clipped one | 127 of 127 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 1,205,908 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 11,711 km²)
bloom area 8,976 km²




1995-10 | source used: hull | VIIRS layer: no coverage

samples 354, positives 102, $\geq 10,000$ cells/L: 44, $\geq 100,000$: 8 (max 480,000 cells/L) | final bloom area = 100% of the hull-clipped one | 26 of 26 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 273,856 km², max 50,588 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 7,735 km²)
bloom area 3,149 km²




1995-11 | source used: hull | VIIRS layer: no coverage

samples 167, positives 65, $\geq 10,000$ cells/L: 21, $\geq 100,000$: 3 (max 150,000 cells/L) | final bloom area = 100% of the hull-clipped one | 12 of 12 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 20,417 km², max 23,735 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 5,580 km²)
bloom area 2,514 km²




1995-12 | source used: hull | VIIRS layer: no coverage

samples 158, positives 63, $\geq 10,000$ cells/L: 33, $\geq 100,000$: 9 (max 35,391,000 cells/L) | final bloom area = 100% of the hull-clipped one | 23 of 23 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 98,743 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 5,250 km²)
bloom area 3,812 km²




1996-01 | source used: hull | VIIRS layer: no coverage

samples 84, positives 10, >= 10,000 cells/L: 9, >= 100,000: 3 (max 471,000 cells/L) | final bloom area = 100% of the hull-clipped one | 3 of 3 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 107,655 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 1,161 km2)
bloom area 835 km2




1996-02 | source used: hull | VIIRS layer: no coverage

samples 179, positives 112, $\geq 10,000$ cells/L: 92, $\geq 100,000$: 62 (max 197,656,000 cells/L) | final bloom area = 100% of the hull-clipped one | 64 of 64 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 52,343,756 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 8,023 km²)
bloom area 5,494 km²



1996-03 | source used: hull | VIIRS layer: no coverage

samples 234, positives 173, $\geq 10,000$ cells/L: 140, $\geq 100,000$: 78 (max 4,700,000 cells/L) | final bloom area = 100% of the hull-clipped one | 97 of 97 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,134 km², max 3,324,951 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 10,458 km²)
bloom area 7,478 km²




1996-04 | source used: hull | VIIRS layer: no coverage

samples 346, positives 174, $\geq 10,000$ cells/L: 108, $\geq 100,000$: 53 (max 1,770,000 cells/L) | final bloom area = 100% of the hull-clipped one | 82 of 82 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 617,817 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 9,102 km²)
bloom area 6,408 km²




1996-05 | source used: hull | VIIRS layer: no coverage

samples 321, positives 143, >= 10,000 cells/L: 81, >= 100,000: 30 (max 3,171,000 cells/L) | final bloom area = 100% of the hull-clipped one | 62 of 62 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 477,131 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 9,417 km2)
bloom area 7,302 km2




1996-06 | source used: hull | VIIRS layer: no coverage

samples 335, positives 137, $\geq 10,000$ cells/L: 80, $\geq 100,000$: 33 (max 967,000 cells/L) | final bloom area = 100% of the hull-clipped one | 64 of 64 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 127,472 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 12,788 km²)
bloom area 10,338 km²




1996-07 | source used: hull | VIIRS layer: no coverage

samples 218, positives 14, >= 10,000 cells/L: 1, >= 100,000: 0 (max 40,000 cells/L) | final bloom area = -% of the hull-clipped one | 1 of 1 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 9,120 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 5,467 km2)
bloom area 0 km2




1996-08 | source used: hull | VIIRS layer: no coverage

samples 197, positives 16, >= 10,000 cells/L: 2, >= 100,000: 1 (max 333,000 cells/L) | final bloom area = 100% of the hull-clipped one | 2 of 2 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 11,488 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 3,707 km2)
bloom area 2,652 km2




1996-09 | source used: hull | VIIRS layer: no coverage

samples 139, positives 8, >= 10,000 cells/L: 6, >= 100,000: 0 (max 80,000 cells/L) | final bloom area = 100% of the hull-clipped one | 6 of 6 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 285,846 km2, max 47,431 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 1,956 km2)
bloom area 1,225 km2




1996-10 | source used: hull | VIIRS layer: no coverage

samples 177, positives 40, $\geq 10,000$ cells/L: 18, $\geq 100,000$: 3 (max 567,000 cells/L) | final bloom area = 100% of the hull-clipped one | 18 of 18 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 284,265 km², max 71,538 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (6 footprint(s), 4,450 km²)
bloom area 2,181 km²



1996-11 | source used: hull | VIIRS layer: no coverage

samples 206, positives 82, $\geq 10,000$ cells/L: 20, $\geq 100,000$: 8 (max 933,000 cells/L) | final bloom area = 100% of the hull-clipped one | 18 of 18 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 6,121 km², max 50,132 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 14,416 km²)
bloom area 4,166 km²




1996-12 | source used: hull | VIIRS layer: no coverage

samples 124, positives 25, $\geq 10,000$ cells/L: 12, $\geq 100,000$: 2 (max 790,000 cells/L) | final bloom area = 100% of the hull-clipped one | 12 of 12 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,759 km², max 424,771 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 6,554 km²)
bloom area 4,731 km²




1997-01 | source used: hull | VIIRS layer: no coverage

samples 134, positives 29, $\geq 10,000$ cells/L: 17, $\geq 100,000$: 9 (max 8,800,000 cells/L) | final bloom area = 100% of the hull-clipped one | 17 of 17 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 5,270,520 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 4,993 km²)
bloom area 4,729 km²




1997-02 | source used: hull | VIIRS layer: no coverage

samples 129, positives 10, $\geq 10,000$ cells/L: 3, $\geq 100,000$: 1 (max $2e+05$ cells/L) | final bloom area = 100% of the hull-clipped one | 3 of 3 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 5,513 km², max 65,741 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 4,535 km²)
bloom area 1,909 km²




1997-08 | source used: hull | VIIRS layer: no coverage

samples 164, positives 11, $\geq 10,000$ cells/L: 7, $\geq 100,000$: 1 (max 233,000 cells/L) | final bloom area = 100% of the hull-clipped one | 7 of 7 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 133,503 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 2,292 km²)
bloom area 2,228 km²




1997-09 | source used: hull | VIIRS layer: no coverage

samples 159, positives 10, $\geq 10,000$ cells/L: 2, $\geq 100,000$: 2 (max 190,000 cells/L) | final bloom area = 100% of the hull-clipped one | 2 of 2 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 11,117 km2, max 188,198 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 2,178 km2)
bloom area 374 km2




1997-10 | source used: hull | VIIRS layer: no coverage

samples 125, positives 12, >= 10,000 cells/L: 7, >= 100,000: 4 (max 147,000 cells/L) | final bloom area = 100% of the hull-clipped one | 7 of 7 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 142,019 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 3,237 km2)
bloom area 2,365 km2



1997-11 | source used: hull | VIIRS layer: no coverage

samples 266, positives 84, $\geq 10,000$ cells/L: 39, $\geq 100,000$: 17 (max 683,000 cells/L) | final bloom area = 100% of the hull-clipped one | 36 of 36 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,286 km², max 260,350 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 13,635 km²)
bloom area 11,457 km²




1997-12 | source used: hull | VIIRS layer: no coverage

samples 163, positives 31, >= 10,000 cells/L: 7, >= 100,000: 4 (max 506,000 cells/L) | final bloom area = 100% of the hull-clipped one | 7 of 7 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 2,625 km2, max 90,208 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 5,621 km2)
bloom area 903 km2




1998-01 | source used: hull | VIIRS layer: no coverage

samples 124, positives 16, $\geq 10,000$ cells/L: 12, $\geq 100,000$: 4 (max 1,300,000 cells/L) | final bloom area = 100% of the hull-clipped one | 12 of 12 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 380,969 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 2,051 km²)
bloom area 1,680 km²




1998-02 | source used: hull | VIIRS layer: no coverage

samples 102, positives 14, $\geq 10,000$ cells/L: 10, $\geq 100,000$: 4 (max 68,333,333 cells/L) | final bloom area = 100% of the hull-clipped one | 10 of 10 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 48,392,456 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (4 footprint(s), 8,392 km²)
bloom area 7,934 km²




1998-03 | source used: hull | VIIRS layer: no coverage

samples 104, positives 6, $\geq 10,000$ cells/L: 2, $\geq 100,000$: 1 (max 288,333 cells/L) | final bloom area = 100% of the hull-clipped one | 2 of 2 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 64,582 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 1,755 km²)
bloom area 1,752 km²




1998-05 | source used: hull | VIIRS layer: no coverage

samples 78, positives 12, >= 10,000 cells/L: 2, >= 100,000: 1 (max 133,000 cells/L) | final bloom area = 100% of the hull-clipped one | 2 of 2 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 2,139 km2, max 17,489 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 6,390 km2)
bloom area 1,986 km2




1998-09 | source used: hull | VIIRS layer: no coverage

samples 138, positives 12, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 1 (max $2e+05$ cells/L) | final bloom area = -% of the hull-clipped one | 1 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 6,755 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 758 km²)
bloom area 0 km²




1998-10 | source used: hull | VIIRS layer: no coverage

samples 119, positives 9, >= 10,000 cells/L: 0, >= 100,000: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 902 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 1,558 km2)
bloom area 0 km2




1998-11 | source used: hull | VIIRS layer: no coverage

samples 292, positives 139, $\geq 10,000$ cells/L: 86, $\geq 100,000$: 25 (max 780,000 cells/L) | final bloom area = 100% of the hull-clipped one | 86 of 86 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 612 km², max 106,180 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 7,183 km²)
bloom area 612 km²




1998-12 | source used: hull | VIIRS layer: no coverage

samples 292, positives 96, $\geq 10,000$ cells/L: 64, $\geq 100,000$: 34 (max 34,430,000 cells/L) | final bloom area = 100% of the hull-clipped one | 62 of 62 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,524 km², max 3,257,894 cells/L




VIIRS-clipped



MODIS nFLH-clipped



USED: Hull-clipped (1 footprint(s), 14,616 km²)
bloom area 12,694 km²



1999-01 | source used: hull | VIIRS layer: no coverage

samples 341, positives 156, >= 10,000 cells/L: 80, >= 100,000: 44 (max 3,630,000 cells/L) | final bloom area = 100% of the hull-clipped one | 76 of 76 water samples >= 10,000 cells/L inside the final footprint






Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,444 km2, max 2,297,256 cells/L

VIIRS-clipped

MODIS nFLH-clipped

USED: Hull-clipped (1 footprint(s), 10,648 km2)
bloom area 6,893 km2


cells/L




1999-02 | source used: hull | VIIRS layer: no coverage

samples 335, positives 163, $\geq 10,000$ cells/L: 100, $\geq 100,000$: 48 (max 3.3×10^7 cells/L) | final bloom area = 100% of the hull-clipped one | 65 of 65 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 288,134 km², max 312,218 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 8,430 km²)
bloom area 4,962 km²




1999-03 | source used: hull | VIIRS layer: no coverage

samples 226, positives 21, $\geq 10,000$ cells/L: 2, $\geq 100,000$: 2 (max 169,000 cells/L) | final bloom area = 100% of the hull-clipped one | 2 of 2 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 1,305 km², max 52,207 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 5,617 km²)
bloom area 537 km²




1999-04 | source used: hull | VIIRS layer: no coverage

samples 161, positives 13, >= 10,000 cells/L: 2, >= 100,000: 0 (max 26,700 cells/L) | final bloom area = -% of the hull-clipped one | 1 of 1 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 5,969 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 2,046 km2)
bloom area 0 km2




1999-08 | source used: hull | VIIRS layer: no coverage

samples 242, positives 36, $\geq 10,000$ cells/L: 8, $\geq 100,000$: 1 (max 933,000 cells/L) | final bloom area = 100% of the hull-clipped one | 7 of 7 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 13,052 km², max 21,358 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 4,206 km²)
bloom area 1,352 km²




1999-09 | source used: hull | VIIRS layer: no coverage

samples 574, positives 284, $\geq 10,000$ cells/L: 191, $\geq 100,000$: 102 (max 15,500,000 cells/L) | final bloom area = 100% of the hull-clipped one | 174 of 174 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 280,351 km2, max 782,733 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (4 footprint(s), 38,855 km2)
bloom area 28,430 km2



1999-10 | source used: hull | VIIRS layer: no coverage

samples 342, positives 107, >= 10,000 cells/L: 54, >= 100,000: 36 (max 9,040,000 cells/L) | final bloom area = 100% of the hull-clipped one | 49 of 49 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 248,084 km2, max 6,448,972 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 18,704 km2)
bloom area 9,961 km2



1999-11 | source used: hull | VIIRS layer: no coverage

samples 135, positives 47, $\geq 10,000$ cells/L: 35, $\geq 100,000$: 26 (max 11,700,000 cells/L) | final bloom area = 100% of the hull-clipped one | 22 of 22 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,213 km², max 550,331 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 2,434 km²)
bloom area 1,264 km²




1999-12 | source used: hull | VIIRS layer: no coverage

samples 307, positives 90, >= 10,000 cells/L: 47, >= 100,000: 19 (max 1,430,000 cells/L) | final bloom area = 100% of the hull-clipped one | 35 of 35 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 287,473 km2, max 401,351 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 17,564 km2)
bloom area 12,434 km2




2000-01 | source used: hull | VIIRS layer: no coverage

samples 312, positives 98, $\geq 10,000$ cells/L: 45, $\geq 100,000$: 15 (max 2,230,000 cells/L) | final bloom area = 100% of the hull-clipped one | 33 of 33 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 287,514 km², max 270,552 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 16,909 km²)
bloom area 14,064 km²




2000-02 | source used: hull | VIIRS layer: no coverage

samples 266, positives 49, $\geq 10,000$ cells/L: 30, $\geq 100,000$: 12 (max 1,830,000 cells/L) | final bloom area = 100% of the hull-clipped one | 21 of 21 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 288,388 km², max 460,810 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 6,140 km²)
bloom area 3,291 km²




2000-03 | source used: hull | VIIRS layer: no coverage

samples 293, positives 44, $\geq 10,000$ cells/L: 29, $\geq 100,000$: 9 (max 2,400,000 cells/L) | final bloom area = 100% of the hull-clipped one | 17 of 17 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 297,622 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 8,542 km²)
bloom area 7,951 km²




2000-04 | source used: hull | VIIRS layer: no coverage

samples 266, positives 31, $\geq 10,000$ cells/L: 10, $\geq 100,000$: 2 (max 6×10^5 cells/L) | final bloom area = 100% of the hull-clipped one | 7 of 7 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,284 km², max 47,264 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 2,728 km²)
bloom area 2,062 km²




2000-05 | source used: hull | VIIRS layer: no coverage

samples 279, positives 9, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 98,000 cells/L) | final bloom area = -% of the hull-clipped one | 1 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 5,383 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 7,131 km²)
bloom area 0 km²




2000-08 | source used: hull | VIIRS layer: no coverage

samples 174, positives 8, $\geq 10,000$ cells/L: 3, $\geq 100,000$: 1 (max 247,000 cells/L) | final bloom area = 100% of the hull-clipped one | 3 of 3 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 1,040 km², max 221,744 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (5 footprint(s), 3,631 km²)
bloom area 668 km²




2000-09 | source used: hull | VIIRS layer: no coverage

samples 811, positives 396, $\geq 10,000$ cells/L: 254, $\geq 100,000$: 168 (max 2,010,000 cells/L) | final bloom area = 100% of the hull-clipped one | 246 of 246 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 15,152 km², max 605,794 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 30,157 km²)
bloom area 6,159 km²




2000-10 | source used: hull | VIIRS layer: no coverage

samples 537, positives 289, $\geq 10,000$ cells/L: 209, $\geq 100,000$: 160 (max 19,900,000 cells/L) | final bloom area = 100% of the hull-clipped one | 205 of 205 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 20,309 km², max 1,442,639 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (4 footprint(s), 26,424 km²)
bloom area 13,064 km²




2000-11 | source used: hull | VIIRS layer: no coverage

samples 240, positives 70, $\geq 10,000$ cells/L: 14, $\geq 100,000$: 1 (max 367,000 cells/L) | final bloom area = 100% of the hull-clipped one | 9 of 9 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 296 km², max 13,668 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 15,832 km²)
bloom area 296 km²




2000-12 | source used: hull | VIIRS layer: no coverage

samples 65, positives 11, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 14,000 cells/L) | final bloom area = -% of the hull-clipped one | 1 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 3,866 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (5 footprint(s), 4,145 km²)
bloom area 0 km²




2001-01 | source used: hull | VIIRS layer: no coverage

samples 105, positives 14, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 9,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,840 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (4 footprint(s), 4,084 km²)
bloom area 0 km²




2001-05 | source used: hull | VIIRS layer: no coverage

samples 140, positives 10, >= 10,000 cells/L: 0, >= 100,000: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 1,365 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 2,958 km2)
bloom area 0 km2




2001-08 | source used: hull | VIIRS layer: no coverage

samples 427, positives 183, $\geq 10,000$ cells/L: 117, $\geq 100,000$: 69 (max 43,430,000 cells/L) | final bloom area = 100% of the hull-clipped one | 102 of 102 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,905 km², max 5,895,738 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 7,532 km²)
bloom area 6,259 km²



2001-09 | source used: hull | VIIRS layer: no coverage

samples 743, positives 500, $\geq 10,000$ cells/L: 422, $\geq 100,000$: 340 (max 1.05×10^8 cells/L) | final bloom area = 100% of the hull-clipped one | 396 of 396 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 18,840,108 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 17,833 km²)
bloom area 16,284 km²




2001-10 | source used: hull | VIIRS layer: no coverage

samples 426, positives 307, $\geq 10,000$ cells/L: 229, $\geq 100,000$: 151 (max 25,900,000 cells/L) | final bloom area = 100% of the hull-clipped one | 215 of 215 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,207 km², max 2,853,565 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 24,770 km²)
bloom area 22,090 km²




2001-11 | source used: hull | VIIRS layer: no coverage

samples 379, positives 260, $\geq 10,000$ cells/L: 212, $\geq 100,000$: 126 (max 10,830,000 cells/L) | final bloom area = 100% of the hull-clipped one | 198 of 198 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 280,913 km², max 1,852,299 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 18,347 km²)
bloom area 13,690 km²




2001-12 | source used: hull | VIIRS layer: no coverage

samples 398, positives 233, $\geq 10,000$ cells/L: 153, $\geq 100,000$: 54 (max 1,800,000 cells/L) | final bloom area = 100% of the hull-clipped one | 133 of 133 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 281,169 km², max 211,740 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 16,537 km²)
bloom area 10,500 km²




2002-01 | source used: hull | VIIRS layer: no coverage

samples 301, positives 154, $\geq 10,000$ cells/L: 86, $\geq 100,000$: 32 (max 9,400,000 cells/L) | final bloom area = 100% of the hull-clipped one | 83 of 83 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 284,890 km², max 429,356 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 17,144 km²)
bloom area 12,195 km²




2002-02 | source used: hull | VIIRS layer: no coverage

samples 322, positives 207, $\geq 10,000$ cells/L: 146, $\geq 100,000$: 94 (max 2,920,000 cells/L) | final bloom area = 100% of the hull-clipped one | 119 of 119 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 4,115,524 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 16,709 km²)
bloom area 14,733 km²




2002-03 | source used: hull | VIIRS layer: no coverage

samples 237, positives 66, $\geq 10,000$ cells/L: 33, $\geq 100,000$: 13 (max 1,930,000 cells/L) | final bloom area = 100% of the hull-clipped one | 30 of 30 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 285,793 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 8,351 km²)
bloom area 5,873 km²




2002-04 | source used: hull | VIIRS layer: no coverage

samples 158, positives 36, $\geq 10,000$ cells/L: 20, $\geq 100,000$: 5 (max 430,000 cells/L) | final bloom area = 100% of the hull-clipped one | 15 of 15 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 277,689 km², max 80,953 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 5,255 km²)
bloom area 2,062 km²




2002-05 | source used: hull | VIIRS layer: no coverage

samples 305, positives 65, $\geq 10,000$ cells/L: 23, $\geq 100,000$: 0 (max 82,000 cells/L) | final bloom area = 100% of the hull-clipped one | 5 of 5 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 153 km², max 13,779 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 5,439 km²)
bloom area 153 km²




2002-06 | source used: hull | VIIRS layer: no coverage

samples 175, positives 45, $\geq 10,000$ cells/L: 16, $\geq 100,000$: 4 (max 203,000 cells/L) | final bloom area = 100% of the hull-clipped one | 16 of 16 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 22,319 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 5,610 km²)
bloom area 4,947 km²




2002-07 | source used: modis | VIIRS layer: no coverage

samples 245, positives 68, $\geq 10,000$ cells/L: 25, $\geq 100,000$: 10 (max 1,100,000 cells/L) | final bloom area = 207% of the hull-clipped one | 11 of 18 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 95,540 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (36 polygon(s))
bloom area 10,685 km²




Hull-clipped (2 footprint(s), 6,408 km²)
bloom area 5,151 km²




2002-08 | source used: modis | VIIRS layer: no coverage | FLAGS: samples_outside

samples 271, positives 111, $\geq 10,000$ cells/L: 55, $\geq 100,000$: 17 (max 2,880,000 cells/L) | final bloom area = 168% of the hull-clipped one | 7 of 47 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,209 km², max 104,171 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (49 polygon(s))
bloom area 11,509 km²




Hull-clipped (2 footprint(s), 8,145 km²)
bloom area 6,840 km²




2002-09 | source used: modis | VIIRS layer: no coverage

samples 274, positives 121, $\geq 10,000$ cells/L: 48, $\geq 100,000$: 9 (max 1,570,000 cells/L) | final bloom area = 343% of the hull-clipped one | 33 of 39 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 40,254 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (30 polygon(s))
bloom area 32,038 km²




Hull-clipped (3 footprint(s), 10,997 km²)
bloom area 9,350 km²




2002-10 | source used: modis | VIIRS layer: no coverage

samples 137, positives 29, $\geq 10,000$ cells/L: 6, $\geq 100,000$: 1 (max 367,000 cells/L) | final bloom area = 653% of the hull-clipped one | 4 of 6 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 279,357 km², max 28,374 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (22 polygon(s))
bloom area 25,391 km²




Hull-clipped (3 footprint(s), 10,999 km²)
bloom area 3,890 km²




2002-11 | source used: modis | VIIRS layer: no coverage

samples 195, positives 85, $\geq 10,000$ cells/L: 34, $\geq 100,000$: 2 (max 302,000 cells/L) | final bloom area = 132% of the hull-clipped one | 32 of 33 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 4,970 km², max 31,135 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (41 polygon(s))
bloom area 3,824 km²




Hull-clipped (2 footprint(s), 10,605 km²)
bloom area 2,906 km²




2002-12 | source used: modis | VIIRS layer: no coverage

samples 107, positives 22, $\geq 10,000$ cells/L: 9, $\geq 100,000$: 4 (max 1,318,000 cells/L) | final bloom area = 731% of the hull-clipped one | 9 of 9 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 285,700 km², max 576,764 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (39 polygon(s))
bloom area 30,250 km²




Hull-clipped (1 footprint(s), 8,076 km²)
bloom area 4,138 km²



2003-01 | source used: modis | VIIRS layer: no coverage

samples 309, positives 154, $\geq 10,000$ cells/L: 87, $\geq 100,000$: 35 (max 9,300,000 cells/L) | final bloom area = 320% of the hull-clipped one | 50 of 81 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,209 km², max 1,432,337 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (33 polygon(s))
bloom area 34,539 km²




Hull-clipped (1 footprint(s), 13,615 km²)
bloom area 10,801 km²




2003-02 | source used: modis | VIIRS layer: no coverage | FLAGS: samples_outside

samples 218, positives 105, $\geq 10,000$ cells/L: 52, $\geq 100,000$: 14 (max 980,000 cells/L) | final bloom area = 312% of the hull-clipped one | 1 of 41 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 281,354 km², max 94,223 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (51 polygon(s))
bloom area 11,741 km²




Hull-clipped (2 footprint(s), 10,591 km²)
bloom area 3,769 km²




2003-03 | source used: modis | VIIRS layer: no coverage

samples 247, positives 136, $\geq 10,000$ cells/L: 95, $\geq 100,000$: 58 (max 4,030,000 cells/L) | final bloom area = 152% of the hull-clipped one | 57 of 78 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 288,849 km², max 2,755,672 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (37 polygon(s))
bloom area 14,042 km²



Hull-clipped (2 footprint(s), 12,570 km²)
bloom area 9,258 km²




cells/L


2003-04 | source used: modis | VIIRS layer: no coverage

samples 339, positives 224, $\geq 10,000$ cells/L: 140, $\geq 100,000$: 56 (max 9,610,000 cells/L) | final bloom area = 94% of the hull-clipped one | 96 of 126 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 495,165 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (30 polygon(s))
bloom area 13,049 km²




Hull-clipped (4 footprint(s), 16,300 km²)
bloom area 13,829 km²




2003-05 | source used: modis | VIIRS layer: no coverage

samples 344, positives 157, $\geq 10,000$ cells/L: 94, $\geq 100,000$: 34 (max 3,800,000 cells/L) | final bloom area = 144% of the hull-clipped one | 46 of 80 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 275,903 km², max 215,695 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (31 polygon(s))
bloom area 11,896 km²




Hull-clipped (3 footprint(s), 10,892 km²)
bloom area 8,250 km²




2003-06 | source used: modis | VIIRS layer: no coverage | FLAGS: samples_outside

samples 263, positives 92, $\geq 10,000$ cells/L: 44, $\geq 100,000$: 9 (max 665,000 cells/L) | final bloom area = 139% of the hull-clipped one | 17 of 35 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 100,660 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (59 polygon(s))
bloom area 11,631 km²




Hull-clipped (2 footprint(s), 9,607 km²)
bloom area 8,379 km²




2003-07 | source used: modis | VIIRS layer: no coverage | FLAGS: samples_outside

samples 330, positives 165, $\geq 10,000$ cells/L: 87, $\geq 100,000$: 27 (max 1,380,000 cells/L) | final bloom area = 102% of the hull-clipped one | 33 of 70 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,208 km², max 187,043 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (45 polygon(s))
bloom area 14,306 km²




Hull-clipped (5 footprint(s), 16,084 km²)
bloom area 14,048 km²




2003-08 | source used: modis | VIIRS layer: no coverage | FLAGS: samples_outside

samples 358, positives 157, >= 10,000 cells/L: 91, >= 100,000: 36 (max 2,020,000 cells/L) | final bloom area = 173% of the hull-clipped one | 38 of 79 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km2, max 270,735 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (41 polygon(s))
bloom area 18,709 km2




Hull-clipped (2 footprint(s), 12,563 km2)
bloom area 10,798 km2




2003-09 | source used: modis | VIIRS layer: no coverage

samples 338, positives 130, $\geq 10,000$ cells/L: 71, $\geq 100,000$: 32 (max 2,560,000 cells/L) | final bloom area = 150% of the hull-clipped one | 38 of 58 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 624,008 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (31 polygon(s))
bloom area 36,166 km²




Hull-clipped (3 footprint(s), 26,998 km²)
bloom area 24,180 km²




2003-10 | source used: modis | VIIRS layer: no coverage

samples 158, positives 38, $\geq 10,000$ cells/L: 14, $\geq 100,000$: 2 (max 139,000 cells/L) | final bloom area = 463% of the hull-clipped one | 6 of 6 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 18,476 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (22 polygon(s))
bloom area 34,324 km²




Hull-clipped (2 footprint(s), 9,327 km²)
bloom area 7,413 km²




2003-11 | source used: modis | VIIRS layer: no coverage

samples 246, positives 110, $\geq 10,000$ cells/L: 37, $\geq 100,000$: 3 (max 187,000 cells/L) | final bloom area = 72% of the hull-clipped one | 28 of 34 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 4,879 km², max 71,092 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (30 polygon(s))
bloom area 2,976 km²




Hull-clipped (3 footprint(s), 19,084 km²)
bloom area 4,116 km²




2003-12 | source used: modis | VIIRS layer: no coverage | FLAGS: samples_outside

samples 202, positives 45, $\geq 10,000$ cells/L: 20, $\geq 100,000$: 3 (max 213,000 cells/L) | final bloom area = 573% of the hull-clipped one | 9 of 19 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 54,726 km², max 48,441 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (61 polygon(s))
bloom area 20,915 km²




Hull-clipped (3 footprint(s), 8,734 km²)
bloom area 3,647 km²




2004-01 | source used: modis | VIIRS layer: no coverage

samples 296, positives 126, $\geq 10,000$ cells/L: 76, $\geq 100,000$: 36 (max 543,000 cells/L) | final bloom area = 104% of the hull-clipped one | 34 of 61 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,210 km², max 477,141 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (41 polygon(s))
bloom area 12,224 km²




Hull-clipped (1 footprint(s), 13,614 km²)
bloom area 11,731 km²




2004-02 | source used: modis | VIIRS layer: no coverage

samples 240, positives 139, $\geq 10,000$ cells/L: 81, $\geq 100,000$: 35 (max 2,010,000 cells/L) | final bloom area = 357% of the hull-clipped one | 48 of 52 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,979 km², max 358,426 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (58 polygon(s))
bloom area 15,493 km²




Hull-clipped (1 footprint(s), 6,309 km²)
bloom area 4,337 km²




2004-03 | source used: modis | VIIRS layer: no coverage

samples 312, positives 7, >= 10,000 cells/L: 0, >= 100,000: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 1,571 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (22 polygon(s))
bloom area 0 km2




Hull-clipped (1 footprint(s), 4,659 km2)
bloom area 0 km2




2004-04 | source used: modis | VIIRS layer: no coverage

samples 206, positives 9, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 3,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,203 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (26 polygon(s))
bloom area 0 km²




Hull-clipped (1 footprint(s), 2,555 km²)
bloom area 0 km²




2004-10 | source used: modis | VIIRS layer: no coverage

samples 186, positives 40, $\geq 10,000$ cells/L: 4, $\geq 100,000$: 2 (max 118,000 cells/L) | final bloom area = 100% of the hull-clipped one | 4 of 4 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 153 km², max 32,933 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (19 polygon(s))
bloom area 153 km²




Hull-clipped (2 footprint(s), 4,057 km²)
bloom area 153 km²




2004-11 | source used: modis | VIIRS layer: no coverage

samples 202, positives 61, $\geq 10,000$ cells/L: 42, $\geq 100,000$: 32 (max 1,080,000 cells/L) | final bloom area = 720% of the hull-clipped one | 37 of 42 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 283,719 km², max 2,097,662 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (30 polygon(s))
bloom area 38,753 km²




Hull-clipped (1 footprint(s), 12,144 km²)
bloom area 5,385 km²




2004-12 | source used: modis | VIIRS layer: no coverage

samples 142, positives 32, $\geq 10,000$ cells/L: 22, $\geq 100,000$: 7 (max 593,000 cells/L) | final bloom area = 849% of the hull-clipped one | 22 of 22 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 282,337 km², max 145,922 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (39 polygon(s))
bloom area 48,476 km²




Hull-clipped (1 footprint(s), 9,848 km²)
bloom area 5,711 km²




2005-01 | source used: modis | VIIRS layer: no coverage

samples 195, positives 92, $\geq 10,000$ cells/L: 65, $\geq 100,000$: 44 (max 6×10^7 cells/L) | final bloom area = 410% of the hull-clipped one | 53 of 58 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 284,373 km², max 5,834,068 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (33 polygon(s))
bloom area 25,542 km²




Hull-clipped (3 footprint(s), 8,545 km²)
bloom area 6,234 km²




2005-02 | source used: modis | VIIRS layer: no coverage

samples 198, positives 142, $\geq 10,000$ cells/L: 113, $\geq 100,000$: 69 (max 151,100,000 cells/L) | final bloom area = 166% of the hull-clipped one | 85 of 92 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,902 km², max 2,112,130 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (37 polygon(s))
bloom area 15,545 km²




Hull-clipped (2 footprint(s), 11,198 km²)
bloom area 9,392 km²




2005-03 | source used: modis | VIIRS layer: no coverage

samples 259, positives 134, $\geq 10,000$ cells/L: 100, $\geq 100,000$: 51 (max 4,402,000 cells/L) | final bloom area = 436% of the hull-clipped one | 73 of 77 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,293 km², max 4,628,416 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (17 polygon(s))
bloom area 19,543 km²




Hull-clipped (1 footprint(s), 6,732 km²)
bloom area 4,487 km²




2005-04 | source used: modis | VIIRS layer: no coverage | FLAGS: samples_outside

samples 269, positives 56, $\geq 10,000$ cells/L: 14, $\geq 100,000$: 4 (max 3,270,000 cells/L) | final bloom area = 361% of the hull-clipped one | 1 of 5 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 27,206 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (25 polygon(s))
bloom area 13,439 km²




Hull-clipped (4 footprint(s), 4,776 km²)
bloom area 3,727 km²




2005-05 | source used: modis | VIIRS layer: no coverage

samples 164, positives 83, $\geq 10,000$ cells/L: 59, $\geq 100,000$: 38 (max 49,400,000 cells/L) | final bloom area = 434% of the hull-clipped one | 25 of 27 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 2,140,817 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (24 polygon(s))
bloom area 7,236 km²




Hull-clipped (2 footprint(s), 2,278 km²)
bloom area 1,665 km²




2005-06 | source used: modis | VIIRS layer: no coverage

samples 208, positives 116, $\geq 10,000$ cells/L: 106, $\geq 100,000$: 79 (max 152,210,000 cells/L) | final bloom area = 371% of the hull-clipped one | 65 of 73 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 2,379,490 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (28 polygon(s))
bloom area 12,116 km²




Hull-clipped (1 footprint(s), 5,035 km²)
bloom area 3,262 km²



2005-07 | source used: modis | VIIRS layer: no coverage

samples 320, positives 189, $\geq 10,000$ cells/L: 154, $\geq 100,000$: 84 (max 64,200,000 cells/L) | final bloom area = 370% of the hull-clipped one | 114 of 118 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,285 km², max 1,916,334 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (47 polygon(s))
bloom area 22,694 km²




Hull-clipped (1 footprint(s), 8,052 km²)
bloom area 6,130 km²




2005-08 | source used: modis | VIIRS layer: no coverage

samples 637, positives 414, $\geq 10,000$ cells/L: 283, $\geq 100,000$: 142 (max 4,600,000 cells/L) | final bloom area = 172% of the hull-clipped one | 169 of 256 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 958,657 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (35 polygon(s))
bloom area 29,365 km²




Hull-clipped (5 footprint(s), 20,359 km²)
bloom area 17,079 km²




2005-09 | source used: modis | VIIRS layer: no coverage

samples 617, positives 493, $\geq 10,000$ cells/L: 392, $\geq 100,000$: 239 (max 162,188,000 cells/L) | final bloom area = 164% of the hull-clipped one | 321 of 337 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,064 km², max 8,447,494 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (32 polygon(s))
bloom area 79,367 km²




Hull-clipped (1 footprint(s), 53,617 km²)
bloom area 48,234 km²




2005-10 | source used: modis | VIIRS layer: no coverage

samples 504, positives 265, $\geq 10,000$ cells/L: 134, $\geq 100,000$: 58 (max 3,907,000 cells/L) | final bloom area = 534% of the hull-clipped one | 105 of 111 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,208 km², max 1,741,901 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (30 polygon(s))
bloom area 72,526 km²




Hull-clipped (3 footprint(s), 22,471 km²)
bloom area 13,585 km²




2005-11 | source used: modis | VIIRS layer: no coverage

samples 460, positives 285, $\geq 10,000$ cells/L: 163, $\geq 100,000$: 81 (max 7,170,000 cells/L) | final bloom area = 267% of the hull-clipped one | 136 of 139 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,795 km², max 2,277,396 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (24 polygon(s))
bloom area 54,794 km²




Hull-clipped (4 footprint(s), 28,871 km²)
bloom area 20,510 km²




2005-12 | source used: modis | VIIRS layer: no coverage

samples 259, positives 93, $\geq 10,000$ cells/L: 40, $\geq 100,000$: 20 (max 4,700,000 cells/L) | final bloom area = 291% of the hull-clipped one | 29 of 30 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,134 km², max 758,756 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (28 polygon(s))
bloom area 38,022 km²




Hull-clipped (4 footprint(s), 17,889 km²)
bloom area 13,060 km²




2006-01 | source used: modis | VIIRS layer: no coverage

samples 471, positives 85, $\geq 10,000$ cells/L: 9, $\geq 100,000$: 3 (max 205,000 cells/L) | final bloom area = -% of the hull-clipped one | 7 of 8 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 5,891 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (28 polygon(s))
bloom area 0 km²



Hull-clipped (3 footprint(s), 12,050 km²)
bloom area 0 km²




cells/L


2006-02 | source used: modis | VIIRS layer: no coverage

samples 396, positives 7, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 5,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 4,947 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (32 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 5,964 km²)
bloom area 0 km²




2006-05 | source used: modis | VIIRS layer: no coverage

samples 473, positives 8, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 4,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 854 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (28 polygon(s))
bloom area 0 km²




Hull-clipped (6 footprint(s), 2,030 km²)
bloom area 0 km²




2006-06 | source used: modis | VIIRS layer: no coverage | FLAGS: samples_outside

samples 334, positives 33, >= 10,000 cells/L: 10, >= 100,000: 4 (max 455,000 cells/L) | final bloom area = 667% of the hull-clipped one | 3 of 9 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 230,682 km2, max 57,162 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (24 polygon(s))
bloom area 9,715 km2




Hull-clipped (2 footprint(s), 8,244 km2)
bloom area 1,456 km2




2006-07 | source used: modis | VIIRS layer: no coverage

samples 565, positives 188, $\geq 10,000$ cells/L: 124, $\geq 100,000$: 73 (max 1.25×10^8 cells/L) | final bloom area = 133% of the hull-clipped one | 67 of 104 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 3,511,317 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (19 polygon(s))
bloom area 9,702 km²




Hull-clipped (6 footprint(s), 9,563 km²)
bloom area 7,310 km²




2006-08 | source used: modis | VIIRS layer: no coverage

samples 675, positives 385, $\geq 10,000$ cells/L: 265, $\geq 100,000$: 134 (max 55,625,000 cells/L) | final bloom area = 114% of the hull-clipped one | 166 of 210 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 3,348,804 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (37 polygon(s))
bloom area 14,754 km²




Hull-clipped (2 footprint(s), 14,920 km²)
bloom area 12,888 km²




2006-09 | source used: modis | VIIRS layer: no coverage

samples 631, positives 484, $\geq 10,000$ cells/L: 393, $\geq 100,000$: 252 (max 45,090,000 cells/L) | final bloom area = 131% of the hull-clipped one | 283 of 319 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 279,487 km², max 4,995,852 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (25 polygon(s))
bloom area 21,711 km²



Hull-clipped (3 footprint(s), 20,103 km²)
bloom area 16,569 km²




cells/L


2006-10 | source used: modis | VIIRS layer: no coverage

samples 673, positives 454, $\geq 10,000$ cells/L: 365, $\geq 100,000$: 224 (max 22,660,000 cells/L) | final bloom area = 161% of the hull-clipped one | 273 of 279 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,357 km², max 1,628,582 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (23 polygon(s))
bloom area 32,576 km²




Hull-clipped (2 footprint(s), 23,255 km²)
bloom area 20,259 km²




2006-11 | source used: modis | VIIRS layer: no coverage

samples 666, positives 338, $\geq 10,000$ cells/L: 219, $\geq 100,000$: 80 (max 5,600,000 cells/L) | final bloom area = 174% of the hull-clipped one | 177 of 187 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,055 km², max 1,403,588 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (25 polygon(s))
bloom area 35,649 km²




Hull-clipped (1 footprint(s), 24,040 km²)
bloom area 20,441 km²



2006-12 | source used: modis | VIIRS layer: no coverage

samples 374, positives 89, $\geq 10,000$ cells/L: 31, $\geq 100,000$: 7 (max 1,453,333 cells/L) | final bloom area = 335% of the hull-clipped one | 22 of 27 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 283,731 km², max 165,604 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (40 polygon(s))
bloom area 20,486 km²




Hull-clipped (1 footprint(s), 12,476 km²)
bloom area 6,117 km²



2007-01 | source used: modis | VIIRS layer: no coverage

samples 356, positives 161, $\geq 10,000$ cells/L: 98, $\geq 100,000$: 32 (max 2,378,000 cells/L) | final bloom area = 260% of the hull-clipped one | 35 of 65 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 268,823 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (27 polygon(s))
bloom area 17,902 km²




Hull-clipped (1 footprint(s), 8,700 km²)
bloom area 6,878 km²




2007-02 | source used: modis | VIIRS layer: no coverage | FLAGS: samples_outside

samples 347, positives 30, $\geq 10,000$ cells/L: 10, $\geq 100,000$: 3 (max 1,870,000 cells/L) | final bloom area = 97% of the hull-clipped one | 3 of 9 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 44,622 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (52 polygon(s))
bloom area 12,319 km²




Hull-clipped (2 footprint(s), 13,522 km²)
bloom area 12,742 km²




2007-03 | source used: modis | VIIRS layer: no coverage

samples 424, positives 49, $\geq 10,000$ cells/L: 15, $\geq 100,000$: 5 (max 480,000 cells/L) | final bloom area = 80% of the hull-clipped one | 12 of 15 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 383 km², max 86,799 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (43 polygon(s))
bloom area 306 km²




Hull-clipped (3 footprint(s), 4,242 km²)
bloom area 383 km²




2007-06 | source used: modis | VIIRS layer: no coverage

samples 412, positives 11, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 913 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (30 polygon(s))
bloom area 0 km2




Hull-clipped (1 footprint(s), 5,512 km2)
bloom area 0 km2




2007-08 | source used: modis | VIIRS layer: no coverage

samples 528, positives 8, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,600 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 515 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (29 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 981 km²)
bloom area 0 km²




2007-09 | source used: modis | VIIRS layer: no coverage

samples 389, positives 38, $\geq 10,000$ cells/L: 25, $\geq 100,000$: 21 (max 1,372,000 cells/L) | final bloom area = 719% of the hull-clipped one | 22 of 24 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,300 km², max 1,165,641 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (34 polygon(s))
bloom area 25,327 km²




Hull-clipped (3 footprint(s), 5,164 km²)
bloom area 3,522 km²




2007-10 | source used: modis | VIIRS layer: no coverage

samples 826, positives 214, $\geq 10,000$ cells/L: 146, $\geq 100,000$: 92 (max 14,059,000 cells/L) | final bloom area = 213% of the hull-clipped one | 135 of 138 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 286,417 km², max 1,168,410 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (40 polygon(s))
bloom area 33,243 km²




Hull-clipped (2 footprint(s), 23,262 km²)
bloom area 15,627 km²



2007-11 | source used: modis | VIIRS layer: no coverage

samples 593, positives 85, $\geq 10,000$ cells/L: 32, $\geq 100,000$: 17 (max 3,530,000 cells/L) | final bloom area = 229% of the hull-clipped one | 15 of 21 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 235,364 km², max 348,828 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (28 polygon(s))
bloom area 13,200 km²



Hull-clipped (2 footprint(s), 18,625 km²)
bloom area 5,770 km²




cells/L


2007-12 | source used: modis | VIIRS layer: no coverage

samples 418, positives 30, $\geq 10,000$ cells/L: 8, $\geq 100,000$: 3 (max 130,000 cells/L) | final bloom area = 503% of the hull-clipped one | 1 of 2 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 23,577 km², max 52,330 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (26 polygon(s))
bloom area 5,723 km²




Hull-clipped (5 footprint(s), 12,042 km²)
bloom area 1,137 km²




2008-01 | source used: modis | VIIRS layer: no coverage

samples 407, positives 9, >= 10,000 cells/L: 0, >= 100,000: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = FALSE)
bloom area 0 km2, max 1,000 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (37 polygon(s))
bloom area 0 km2




Hull-clipped (3 footprint(s), 3,141 km2)
bloom area 0 km2




2008-04 | source used: modis | VIIRS layer: no coverage

samples 444, positives 17, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 33,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 7,900 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (27 polygon(s))
bloom area 0 km2




Hull-clipped (2 footprint(s), 6,839 km2)
bloom area 0 km2




2008-10 | source used: modis | VIIRS layer: no coverage

samples 722, positives 128, $\geq 10,000$ cells/L: 51, $\geq 100,000$: 10 (max 755,000 cells/L) | final bloom area = 100% of the hull-clipped one | 51 of 51 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 1,147 km², max 71,739 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (33 polygon(s))
bloom area 1,071 km²




Hull-clipped (2 footprint(s), 8,308 km²)
bloom area 1,071 km²




2008-11 | source used: modis | VIIRS layer: no coverage

samples 311, positives 24, $\geq 10,000$ cells/L: 4, $\geq 100,000$: 1 (max 135,000 cells/L) | final bloom area = 420% of the hull-clipped one | 3 of 4 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 9,027 km², max 19,774 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (24 polygon(s))
bloom area 6,792 km²




Hull-clipped (4 footprint(s), 5,089 km²)
bloom area 1,618 km²




2008-12 | source used: modis | VIIRS layer: no coverage

samples 380, positives 15, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,168 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (55 polygon(s))
bloom area 0 km²




Hull-clipped (6 footprint(s), 8,094 km²)
bloom area 0 km²




2009-01 | source used: modis | VIIRS layer: no coverage

samples 339, positives 8, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 806 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (30 polygon(s))
bloom area 0 km2




Hull-clipped (3 footprint(s), 1,349 km2)
bloom area 0 km2




2009-04 | source used: modis | VIIRS layer: no coverage

samples 404, positives 9, $\geq 10,000$ cells/L: 2, $\geq 100,000$: 0 (max 32,666 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 229 km2, max 13,216 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (27 polygon(s))
bloom area 0 km2




Hull-clipped (2 footprint(s), 8,360 km2)
bloom area 229 km2




2009-05 | source used: modis | VIIRS layer: no coverage

samples 358, positives 14, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 941 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (22 polygon(s))
bloom area 0 km2




Hull-clipped (3 footprint(s), 3,138 km2)
bloom area 0 km2




2009-06 | source used: modis | VIIRS layer: no coverage

samples 432, positives 13, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 7,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,734 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (29 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 2,373 km²)
bloom area 0 km²




2009-07 | source used: modis | VIIRS layer: no coverage

samples 462, positives 21, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 3,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,059 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (28 polygon(s))
bloom area 0 km²



Hull-clipped (4 footprint(s), 3,426 km²)
bloom area 0 km²




cells/L


2009-09 | source used: modis | VIIRS layer: no coverage

samples 440, positives 19, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 1,082 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (27 polygon(s))
bloom area 0 km2



Hull-clipped (1 footprint(s), 4,359 km2)
bloom area 0 km2




cells/L


2009–10 | source used: modis | VIIRS layer: no coverage

samples 878, positives 153, $\geq 10,000$ cells/L: 66, $\geq 100,000$: 22 (max 5,609,300 cells/L) | final bloom area = 219% of the hull-clipped one | 55 of 66 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 287,491 km², max 128,270 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (25 polygon(s))
bloom area 20,061 km²



Hull-clipped (1 footprint(s), 12,486 km²)
bloom area 9,178 km²




cells/L


2009-11 | source used: modis | VIIRS layer: no coverage | FLAGS: samples_outside

samples 389, positives 65, $\geq 10,000$ cells/L: 28, $\geq 100,000$: 21 (max 20,100,000 cells/L) | final bloom area = 1024% of the hull-clipped one | 10 of 28 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,758 km², max 71,396,480 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (45 polygon(s))
bloom area 29,006 km²




Hull-clipped (1 footprint(s), 6,677 km²)
bloom area 2,834 km²




2009–12 | source used: modis | VIIRS layer: no coverage

samples 396, positives 97, $\geq 10,000$ cells/L: 24, $\geq 100,000$: 2 (max 172,000 cells/L) | final bloom area = 1850% of the hull-clipped one | 17 of 17 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,447 km², max 18,510 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (76 polygon(s))
bloom area 46,598 km²




Hull-clipped (2 footprint(s), 4,412 km²)
bloom area 2,519 km²




2010-01 | source used: modis | VIIRS layer: no coverage

samples 423, positives 40, >= 10,000 cells/L: 10, >= 100,000: 1 (max 122,000 cells/L) | final bloom area = -% of the hull-clipped one | 10 of 10 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 8,760 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (71 polygon(s))
bloom area 0 km2




Hull-clipped (1 footprint(s), 10,503 km2)
bloom area 0 km2




2010-02 | source used: modis | VIIRS layer: no coverage

samples 353, positives 11, $\geq 10,000$ cells/L: 3, $\geq 100,000$: 0 (max 15,700 cells/L) | final bloom area = -% of the hull-clipped one | 2 of 2 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 6,597 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (80 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 2,977 km²)
bloom area 0 km²




2010-04 | source used: modis | VIIRS layer: no coverage

samples 356, positives 8, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,600 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 901 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (26 polygon(s))
bloom area 0 km2




Hull-clipped (3 footprint(s), 2,015 km2)
bloom area 0 km2




2010-05 | source used: modis | VIIRS layer: no coverage

samples 471, positives 10, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 533 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (20 polygon(s))
bloom area 0 km2




Hull-clipped (4 footprint(s), 4,112 km2)
bloom area 0 km2




2010-06 | source used: modis | VIIRS layer: no coverage

samples 479, positives 7, >= 10,000 cells/L: 0, >= 100,000: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 470 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (30 polygon(s))
bloom area 0 km2




Hull-clipped (4 footprint(s), 1,514 km2)
bloom area 0 km2




2010-09 | source used: modis | VIIRS layer: no coverage

samples 371, positives 7, >= 10,000 cells/L: 0, >= 100,000: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 1,531 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (30 polygon(s))
bloom area 0 km2




Hull-clipped (2 footprint(s), 1,406 km2)
bloom area 0 km2




2010-10 | source used: modis | VIIRS layer: no coverage

samples 584, positives 12, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 5,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 2,054 cells/L




VIIRS-clipped




USED: MODIS nFLH-clipped (25 polygon(s))
bloom area 0 km²



Hull-clipped (3 footprint(s), 2,155 km²)
bloom area 0 km²




cells/L




2011-02 | source used: modis | VIIRS layer: no coverage

samples 385, positives 7, >= 10,000 cells/L: 0, >= 100,000: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 1,155 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (37 polygon(s))
bloom area 0 km2




Hull-clipped (4 footprint(s), 1,507 km2)
bloom area 0 km2




2011-03 | source used: modis | VIIRS layer: no coverage

samples 430, positives 25, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,141 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (27 polygon(s))
bloom area 0 km²



Hull-clipped (2 footprint(s), 9,486 km²)
bloom area 0 km²




cells/L


2011-04 | source used: modis | VIIRS layer: no coverage

samples 435, positives 22, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 7,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 1,587 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (24 polygon(s))
bloom area 0 km2




Hull-clipped (3 footprint(s), 10,707 km2)
bloom area 0 km2




2011-05 | source used: modis | VIIRS layer: no coverage

samples 453, positives 6, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 1,108 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (26 polygon(s))
bloom area 0 km2




Hull-clipped (1 footprint(s), 2,164 km2)
bloom area 0 km2




2011-06 | source used: modis | VIIRS layer: no coverage

samples 476, positives 10, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,300 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 701 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (22 polygon(s))
bloom area 0 km2




Hull-clipped (2 footprint(s), 3,370 km2)
bloom area 0 km2




2011-07 | source used: modis | VIIRS layer: no coverage

samples 457, positives 13, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,100 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (28 polygon(s))
bloom area 0 km²




Hull-clipped (3 footprint(s), 4,620 km²)
bloom area 0 km²




2011-08 | source used: modis | VIIRS layer: no coverage

samples 470, positives 7, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 17,410 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 2,625 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (27 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 5,680 km²)
bloom area 0 km²




2011-09 | source used: modis | VIIRS layer: no coverage

samples 437, positives 41, $\geq 10,000$ cells/L: 14, $\geq 100,000$: 7 (max 2,450,000 cells/L) | final bloom area = 558% of the hull-clipped one | 8 of 9 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 147,435 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (23 polygon(s))
bloom area 19,558 km²




Hull-clipped (1 footprint(s), 4,680 km²)
bloom area 3,503 km²




2011-10 | source used: modis | VIIRS layer: no coverage

samples 569, positives 152, $\geq 10,000$ cells/L: 102, $\geq 100,000$: 65 (max 22,275,000 cells/L) | final bloom area = 605% of the hull-clipped one | 98 of 98 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,133 km², max 4,478,020 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (38 polygon(s))
bloom area 24,969 km²




Hull-clipped (1 footprint(s), 5,330 km²)
bloom area 4,130 km²



2011-11 | source used: modis | VIIRS layer: no coverage

samples 471, positives 112, $\geq 10,000$ cells/L: 65, $\geq 100,000$: 46 (max 30,448,000 cells/L) | final bloom area = 543% of the hull-clipped one | 57 of 57 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 2,757,693 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (29 polygon(s))
bloom area 22,888 km²




Hull-clipped (1 footprint(s), 6,371 km²)
bloom area 4,213 km²



2011-12 | source used: modis | VIIRS layer: no coverage

samples 511, positives 163, $\geq 10,000$ cells/L: 130, $\geq 100,000$: 95 (max 23,936,000 cells/L) | final bloom area = 558% of the hull-clipped one | 113 of 114 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,750 km², max 7,715,438 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (30 polygon(s))
bloom area 23,115 km²



Hull-clipped (1 footprint(s), 5,672 km²)
bloom area 4,146 km²




cells/L


2012-01 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 438, positives 91, $\geq 10,000$ cells/L: 34, $\geq 100,000$: 17 (max 1,319,300 cells/L) | final bloom area = 149% of the hull-clipped one | 13 of 29 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,596 km², max 177,518 cells/L




USED: VIIRS-clipped (48 positive VIIRS cells)
bloom area 6,482 km²



MODIS nFLH-clipped (37 polygon(s))
bloom area 17,017 km²




Hull-clipped (4 footprint(s), 6,549 km²)
bloom area 4,354 km²




2012-02 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 376, positives 12, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 5,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 746 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (61 polygon(s))
bloom area 0 km²




Hull-clipped (3 footprint(s), 7,549 km²)
bloom area 0 km²




2012-03 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 458, positives 41, $\geq 10,000$ cells/L: 14, $\geq 100,000$: 8 (max 513,300 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 13 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 8,657 km², max 177,829 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (35 polygon(s))
bloom area 2,318 km²




Hull-clipped (5 footprint(s), 14,023 km²)
bloom area 5,098 km²




2012-04 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 439, positives 20, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 11,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,520 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (22 polygon(s))
bloom area 0 km²




Hull-clipped (3 footprint(s), 4,042 km²)
bloom area 0 km²




2012-05 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 386, positives 8, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 989 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (27 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 1,561 km²)
bloom area 0 km²




2012-06 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 469, positives 42, $\geq 10,000$ cells/L: 2, $\geq 100,000$: 0 (max 80,300 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 3,827 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (26 polygon(s))
bloom area 0 km²




Hull-clipped (5 footprint(s), 7,924 km²)
bloom area 0 km²




2012-08 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 359, positives 23, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 6,300 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 3,880 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (40 polygon(s))
bloom area 0 km²




Hull-clipped (3 footprint(s), 7,184 km²)
bloom area 0 km²




2012-09 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 496, positives 163, $\geq 10,000$ cells/L: 20, $\geq 100,000$: 0 (max 58,000 cells/L) | final bloom area = 50% of the hull-clipped one | 5 of 13 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 152 km², max 17,329 cells/L




USED: VIIRS-clipped (47 positive VIIRS cells)
bloom area 76 km²



MODIS nFLH-clipped (35 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 9,988 km²)
bloom area 152 km²




2012-10 | source used: viirs | VIIRS layer: positive

samples 737, positives 485, $\geq 10,000$ cells/L: 309, $\geq 100,000$: 164 (max 68,420,000 cells/L) | final bloom area = 160% of the hull-clipped one | 184 of 243 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 1,705,330 cells/L




USED: VIIRS-clipped (104 positive VIIRS cells)
bloom area 15,494 km²



MODIS nFLH-clipped (41 polygon(s))
bloom area 32,084 km²




Hull-clipped (3 footprint(s), 12,352 km²)
bloom area 9,680 km²




2012-11 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 737, positives 499, $\geq 10,000$ cells/L: 359, $\geq 100,000$: 147 (max 1.5×10^8 cells/L) | final bloom area = 30% of the hull-clipped one | 66 of 290 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,207 km², max 1,423,496 cells/L




USED: VIIRS-clipped (18 positive VIIRS cells)
bloom area 2,913 km²



MODIS nFLH-clipped (37 polygon(s))
bloom area 22,096 km²




Hull-clipped (1 footprint(s), 13,713 km²)
bloom area 9,554 km²




2012-12 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 497, positives 247, $\geq 10,000$ cells/L: 102, $\geq 100,000$: 40 (max 8,420,000 cells/L) | final bloom area = 33% of the hull-clipped one | 24 of 75 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,133 km², max 519,385 cells/L




USED: VIIRS-clipped (15 positive VIIRS cells)
bloom area 2,527 km²



MODIS nFLH-clipped (48 polygon(s))
bloom area 22,421 km²




Hull-clipped (2 footprint(s), 10,744 km²)
bloom area 7,697 km²




2013-01 | source used: viirs | VIIRS layer: positive

samples 697, positives 490, $\geq 10,000$ cells/L: 369, $\geq 100,000$: 227 (max 63,506,600 cells/L) | final bloom area = 83% of the hull-clipped one | 182 of 269 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,035 km², max 5,020,582 cells/L




USED: VIIRS-clipped (36 positive VIIRS cells)
bloom area 5,269 km²



MODIS nFLH-clipped (31 polygon(s))
bloom area 18,488 km²




Hull-clipped (2 footprint(s), 9,869 km²)
bloom area 6,338 km²




2013-02 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 625, positives 403, $\geq 10,000$ cells/L: 297, $\geq 100,000$: 174 (max 3,984,300 cells/L) | final bloom area = 50% of the hull-clipped one | 93 of 201 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 288,780 km², max 2,371,374 cells/L




USED: VIIRS-clipped (23 positive VIIRS cells)
bloom area 3,217 km²



MODIS nFLH-clipped (29 polygon(s))
bloom area 19,335 km²




Hull-clipped (1 footprint(s), 9,798 km²)
bloom area 6,434 km²




2013-03 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 493, positives 173, $\geq 10,000$ cells/L: 69, $\geq 100,000$: 15 (max 853,300 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 49 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 10,402 km², max 130,662 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (39 polygon(s))
bloom area 763 km²




Hull-clipped (2 footprint(s), 6,437 km²)
bloom area 2,292 km²




2013-04 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, samples_outside

samples 683, positives 133, $\geq 10,000$ cells/L: 9, $\geq 100,000$: 2 (max 352,600 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 9 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 6,387 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (29 polygon(s))
bloom area 0 km²




Hull-clipped (1 footprint(s), 9,856 km²)
bloom area 0 km²




2013-05 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 568, positives 85, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 5,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,308 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (25 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 9,697 km²)
bloom area 0 km²




2013-06 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 459, positives 19, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 5,667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 3,050 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (23 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 8,186 km²)
bloom area 0 km²




2013-07 | source used: viirs | VIIRS layer: positive

samples 380, positives 15, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 19,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 1,990 cells/L




USED: VIIRS-clipped (4 positive VIIRS cells)
bloom area 0 km2



MODIS nFLH-clipped (39 polygon(s))
bloom area 0 km2




Hull-clipped (1 footprint(s), 1,686 km2)
bloom area 0 km2




2013-08 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 449, positives 6, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = FALSE)
bloom area 0 km², max 1,023 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (45 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 2,385 km²)
bloom area 0 km²




2013-09 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 472, positives 20, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 11,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,896 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (30 polygon(s))
bloom area 0 km²



Hull-clipped (1 footprint(s), 6,058 km²)
bloom area 0 km²




cells/L


2013-10 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, samples_outside

samples 615, positives 113, $\geq 10,000$ cells/L: 13, $\geq 100,000$: 2 (max 133,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 9 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 6,915 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (24 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 8,668 km²)
bloom area 0 km²




2013-11 | source used: viirs | VIIRS layer: positive

samples 578, positives 215, $\geq 10,000$ cells/L: 83, $\geq 100,000$: 30 (max 1,821,000 cells/L) | final bloom area = 73% of the hull-clipped one | 63 of 78 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 286,178 km², max 417,324 cells/L




USED: VIIRS-clipped (21 positive VIIRS cells)
bloom area 2,222 km²



MODIS nFLH-clipped (62 polygon(s))
bloom area 21,236 km²




Hull-clipped (1 footprint(s), 7,835 km²)
bloom area 3,062 km²




2013-12 | source used: viirs | VIIRS layer: positive

samples 427, positives 13, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 895 cells/L




USED: VIIRS-clipped (7 positive VIIRS cells)
bloom area 0 km2



MODIS nFLH-clipped (47 polygon(s))
bloom area 0 km2




Hull-clipped (3 footprint(s), 3,658 km2)
bloom area 0 km2




2014-01 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 403, positives 15, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 3,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,117 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (30 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 3,108 km²)
bloom area 0 km²




2014-02 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 499, positives 16, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 644 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (31 polygon(s))
bloom area 0 km²




Hull-clipped (3 footprint(s), 5,464 km²)
bloom area 0 km²




2014-03 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 418, positives 17, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 532 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (29 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 6,853 km²)
bloom area 0 km²




2014-04 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 503, positives 27, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 526 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (31 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 12,810 km²)
bloom area 0 km²




2014-05 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 450, positives 17, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 658 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (27 polygon(s))
bloom area 0 km²




Hull-clipped (1 footprint(s), 6,092 km²)
bloom area 0 km²




2014-06 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 409, positives 15, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,300 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,179 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (37 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 5,376 km²)
bloom area 0 km²




2014-07 | source used: viirs | VIIRS layer: positive

samples 455, positives 34, $\geq 10,000$ cells/L: 16, $\geq 100,000$: 10 (max 20,265,000 cells/L) | final bloom area = 107% of the hull-clipped one | 12 of 16 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 280,205 km², max 2,765,506 cells/L




USED: VIIRS-clipped (52 positive VIIRS cells)
bloom area 7,797 km²



MODIS nFLH-clipped (48 polygon(s))
bloom area 14,894 km²




Hull-clipped (4 footprint(s), 12,116 km²)
bloom area 7,308 km²




2014-08 | source used: viirs | VIIRS layer: positive

samples 836, positives 256, $\geq 10,000$ cells/L: 152, $\geq 100,000$: 80 (max 1,462,667 cells/L) | final bloom area = 118% of the hull-clipped one | 140 of 152 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 28,438 km², max 1,096,336 cells/L




USED: VIIRS-clipped (129 positive VIIRS cells)
bloom area 15,692 km²



MODIS nFLH-clipped (39 polygon(s))
bloom area 6,380 km²




Hull-clipped (7 footprint(s), 24,250 km²)
bloom area 13,310 km²




2014-09 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 810, positives 122, $\geq 10,000$ cells/L: 60, $\geq 100,000$: 30 (max 1,972,667 cells/L) | final bloom area = 54% of the hull-clipped one | 14 of 60 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 19,663 km², max 1,543,536 cells/L




USED: VIIRS-clipped (55 positive VIIRS cells)
bloom area 3,809 km²



MODIS nFLH-clipped (48 polygon(s))
bloom area 6,271 km²




Hull-clipped (2 footprint(s), 23,558 km²)
bloom area 7,014 km²




2014-10 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 620, positives 59, $\geq 10,000$ cells/L: 18, $\geq 100,000$: 10 (max 2,300,000 cells/L) | final bloom area = 38% of the hull-clipped one | 0 of 18 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 16,378 km², max 226,387 cells/L




USED: VIIRS-clipped (36 positive VIIRS cells)
bloom area 1,051 km²



MODIS nFLH-clipped (30 polygon(s))
bloom area 3,736 km²




Hull-clipped (4 footprint(s), 12,438 km²)
bloom area 2,739 km²




2014-11 | source used: viirs | VIIRS layer: positive

samples 521, positives 91, $\geq 10,000$ cells/L: 40, $\geq 100,000$: 16 (max 5,547,500 cells/L) | final bloom area = 248% of the hull-clipped one | 33 of 36 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 247,162 km², max 603,791 cells/L




USED: VIIRS-clipped (88 positive VIIRS cells)
bloom area 8,932 km²



MODIS nFLH-clipped (34 polygon(s))
bloom area 25,439 km²




Hull-clipped (3 footprint(s), 9,312 km²)
bloom area 3,609 km²




2015-03 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 474, positives 11, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 560 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (18 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 3,073 km²)
bloom area 0 km²




2015-06 | source used: viirs | VIIRS layer: positive

samples 530, positives 6, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 500 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 360 cells/L




USED: VIIRS-clipped (3 positive VIIRS cells)
bloom area 0 km2




MODIS nFLH-clipped (31 polygon(s))
bloom area 0 km2



Hull-clipped (6 footprint(s), 1,884 km2)
bloom area 0 km2




cells/L




2015-09 | source used: viirs | VIIRS layer: positive

samples 470, positives 84, $\geq 10,000$ cells/L: 24, $\geq 100,000$: 5 (max 10,883,333 cells/L) | final bloom area = 76% of the hull-clipped one | 11 of 18 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 9,004 km², max 145,125 cells/L




USED: VIIRS-clipped (31 positive VIIRS cells)
bloom area 3,117 km²



MODIS nFLH-clipped (44 polygon(s))
bloom area 2,157 km²



Hull-clipped (2 footprint(s), 14,411 km²)
bloom area 4,083 km²




cells/L


2015-10 | source used: viirs | VIIRS layer: positive

samples 715, positives 284, $\geq 10,000$ cells/L: 163, $\geq 100,000$: 92 (max 32,560,000 cells/L) | final bloom area = 59% of the hull-clipped one | 87 of 130 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,517 km², max 4,962,534 cells/L




USED: VIIRS-clipped (123 positive VIIRS cells)
bloom area 16,484 km²



MODIS nFLH-clipped (16 polygon(s))
bloom area 22,626 km²




Hull-clipped (6 footprint(s), 32,469 km²)
bloom area 27,919 km²




2015-11 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 571, positives 312, $\geq 10,000$ cells/L: 231, $\geq 100,000$: 162 (max 41,900,000 cells/L) | final bloom area = 98% of the hull-clipped one | 78 of 158 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 2,451,805 cells/L




USED: VIIRS-clipped (81 positive VIIRS cells)
bloom area 11,670 km²



MODIS nFLH-clipped (19 polygon(s))
bloom area 24,983 km²




Hull-clipped (4 footprint(s), 15,673 km²)
bloom area 11,945 km²




2015-12 | source used: viirs | VIIRS layer: positive

samples 600, positives 361, $\geq 10,000$ cells/L: 237, $\geq 100,000$: 128 (max 7,841,667 cells/L) | final bloom area = 118% of the hull-clipped one | 89 of 171 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,677 km², max 2,667,769 cells/L




USED: VIIRS-clipped (96 positive VIIRS cells)
bloom area 13,253 km²



MODIS nFLH-clipped (58 polygon(s))
bloom area 23,613 km²




Hull-clipped (2 footprint(s), 17,656 km²)
bloom area 11,187 km²




2016-01 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 595, positives 349, $\geq 10,000$ cells/L: 270, $\geq 100,000$: 158 (max 35,267,000 cells/L) | final bloom area = 54% of the hull-clipped one | 37 of 206 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 23,059 km², max 670,774 cells/L




USED: VIIRS-clipped (27 positive VIIRS cells)
bloom area 3,977 km²



MODIS nFLH-clipped (85 polygon(s))
bloom area 12,024 km²




Hull-clipped (4 footprint(s), 15,806 km²)
bloom area 7,391 km²




2016-02 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 615, positives 324, $\geq 10,000$ cells/L: 209, $\geq 100,000$: 95 (max 21,400,000 cells/L) | final bloom area = 40% of the hull-clipped one | 25 of 136 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 382,835 cells/L




USED: VIIRS-clipped (23 positive VIIRS cells)
bloom area 3,910 km²



MODIS nFLH-clipped (34 polygon(s))
bloom area 25,058 km²




Hull-clipped (2 footprint(s), 13,331 km²)
bloom area 9,911 km²




2016-03 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 649, positives 133, $\geq 10,000$ cells/L: 30, $\geq 100,000$: 2 (max 219,333 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 27 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 2,208 km², max 78,879 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (41 polygon(s))
bloom area 1,144 km²




Hull-clipped (5 footprint(s), 15,424 km²)
bloom area 1,601 km²




2016-04 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 768, positives 246, $\geq 10,000$ cells/L: 57, $\geq 100,000$: 10 (max 1,610,000 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 41 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 1,211 km², max 39,961 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (23 polygon(s))
bloom area 227 km²




Hull-clipped (2 footprint(s), 13,394 km²)
bloom area 985 km²




2016-05 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 501, positives 79, >= 10,000 cells/L: 5, >= 100,000: 0 (max 51,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 3 water samples >= 10,000 cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 4,172 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km2



MODIS nFLH-clipped (18 polygon(s))
bloom area 0 km2




Hull-clipped (3 footprint(s), 12,600 km2)
bloom area 0 km2




2016-08 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 636, positives 36, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 689 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (31 polygon(s))
bloom area 0 km²




Hull-clipped (1 footprint(s), 11,951 km²)
bloom area 0 km²




2016-09 | source used: viirs | VIIRS layer: positive

samples 555, positives 165, $\geq 10,000$ cells/L: 92, $\geq 100,000$: 56 (max 48,604,333 cells/L) | final bloom area = 100% of the hull-clipped one | 34 of 65 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 2,804,972 cells/L




USED: VIIRS-clipped (64 positive VIIRS cells)
bloom area 9,253 km²



MODIS nFLH-clipped (19 polygon(s))
bloom area 16,283 km²




Hull-clipped (3 footprint(s), 11,596 km²)
bloom area 9,261 km²




2016-10 | source used: viirs | VIIRS layer: positive

samples 732, positives 448, $\geq 10,000$ cells/L: 330, $\geq 100,000$: 222 (max 60,409,023 cells/L) | final bloom area = 197% of the hull-clipped one | 172 of 256 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 288,915 km², max 2,720,364 cells/L




USED: VIIRS-clipped (132 positive VIIRS cells)
bloom area 17,557 km²



MODIS nFLH-clipped (20 polygon(s))
bloom area 21,559 km²




Hull-clipped (5 footprint(s), 12,800 km²)
bloom area 8,917 km²




2016-11 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 631, positives 391, $\geq 10,000$ cells/L: 308, $\geq 100,000$: 195 (max 23,600,000 cells/L) | final bloom area = 111% of the hull-clipped one | 100 of 218 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 1,335,938 cells/L




USED: VIIRS-clipped (141 positive VIIRS cells)
bloom area 18,235 km²



MODIS nFLH-clipped (21 polygon(s))
bloom area 21,780 km²




Hull-clipped (2 footprint(s), 20,630 km²)
bloom area 16,474 km²




2016-12 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 649, positives 299, $\geq 10,000$ cells/L: 203, $\geq 100,000$: 111 (max 17,680,000 cells/L) | final bloom area = 67% of the hull-clipped one | 53 of 136 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 1,221,185 cells/L




USED: VIIRS-clipped (34 positive VIIRS cells)
bloom area 6,053 km²



MODIS nFLH-clipped (32 polygon(s))
bloom area 16,862 km²




Hull-clipped (2 footprint(s), 12,572 km²)
bloom area 9,060 km²




2017-01 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 554, positives 274, $\geq 10,000$ cells/L: 161, $\geq 100,000$: 95 (max 5,767,962 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 116 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,289 km², max 602,578 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (41 polygon(s))
bloom area 21,891 km²




Hull-clipped (2 footprint(s), 8,395 km²)
bloom area 4,181 km²




2017-02 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 606, positives 320, $\geq 10,000$ cells/L: 192, $\geq 100,000$: 85 (max 5,944,400 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 129 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 260,962 km², max 826,661 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (26 polygon(s))
bloom area 10,044 km²




Hull-clipped (3 footprint(s), 12,955 km²)
bloom area 7,244 km²




2017-03 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 688, positives 258, $\geq 10,000$ cells/L: 98, $\geq 100,000$: 17 (max 638,094 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 59 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 281,758 km², max 121,434 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (29 polygon(s))
bloom area 8,376 km²




Hull-clipped (3 footprint(s), 10,452 km²)
bloom area 6,417 km²




2017-04 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 603, positives 168, $\geq 10,000$ cells/L: 34, $\geq 100,000$: 5 (max 167,000 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 21 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 457 km², max 17,926 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (26 polygon(s))
bloom area 0 km²




Hull-clipped (6 footprint(s), 10,058 km²)
bloom area 457 km²




2017-05 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 707, positives 79, $\geq 10,000$ cells/L: 3, $\geq 100,000$: 0 (max 15,500 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 2,101 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (33 polygon(s))
bloom area 0 km²




Hull-clipped (3 footprint(s), 9,944 km²)
bloom area 0 km²




2017-06 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 699, positives 37, $\geq 10,000$ cells/L: 3, $\geq 100,000$: 0 (max 10,500 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 2 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,950 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (48 polygon(s))
bloom area 0 km²




Hull-clipped (1 footprint(s), 4,376 km²)
bloom area 0 km²




2017-07 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 672, positives 17, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 465 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (33 polygon(s))
bloom area 0 km²




Hull-clipped (5 footprint(s), 7,276 km²)
bloom area 0 km²




2017-08 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 847, positives 8, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,333 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 632 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (48 polygon(s))
bloom area 0 km²




Hull-clipped (5 footprint(s), 2,437 km²)
bloom area 0 km²




2017-10 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 689, positives 70, $\geq 10,000$ cells/L: 9, $\geq 100,000$: 0 (max 32,667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 9 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 4,932 cells/L




USED: VIIRS-clipped (4 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (34 polygon(s))
bloom area 0 km²




Hull-clipped (5 footprint(s), 12,219 km²)
bloom area 0 km²




2017-11 | source used: viirs | VIIRS layer: positive

samples 654, positives 124, $\geq 10,000$ cells/L: 53, $\geq 100,000$: 29 (max 2,968,560 cells/L) | final bloom area = 235% of the hull-clipped one | 27 of 40 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 11,491 km², max 1,055,268 cells/L




USED: VIIRS-clipped (72 positive VIIRS cells)
bloom area 7,360 km²



MODIS nFLH-clipped (26 polygon(s))
bloom area 5,213 km²



Hull-clipped (3 footprint(s), 7,598 km²)
bloom area 3,136 km²




cells/L


2017-12 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 629, positives 224, $\geq 10,000$ cells/L: 109, $\geq 100,000$: 54 (max 6,479,760 cells/L) | final bloom area = 158% of the hull-clipped one | 37 of 80 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 259,340 km², max 922,532 cells/L




USED: VIIRS-clipped (75 positive VIIRS cells)
bloom area 10,051 km²



MODIS nFLH-clipped (21 polygon(s))
bloom area 19,703 km²




Hull-clipped (3 footprint(s), 13,559 km²)
bloom area 6,349 km²




2018-01 | source used: viirs | VIIRS layer: positive | FLAGS: low_vs_hull, samples_outside

samples 612, positives 116, $\geq 10,000$ cells/L: 67, $\geq 100,000$: 24 (max 1,307,600 cells/L) | final bloom area = 14% of the hull-clipped one | 7 of 46 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 284,379 km², max 312,623 cells/L




USED: VIIRS-clipped (9 positive VIIRS cells)
bloom area 1,456 km²



MODIS nFLH-clipped (76 polygon(s))
bloom area 25,290 km²




Hull-clipped (2 footprint(s), 15,330 km²)
bloom area 10,150 km²




2018-02 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 627, positives 82, $\geq 10,000$ cells/L: 38, $\geq 100,000$: 15 (max 881,073 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 31 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 249,480 km², max 230,908 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (25 polygon(s))
bloom area 8,992 km²




Hull-clipped (2 footprint(s), 14,016 km²)
bloom area 7,808 km²




2018-03 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 605, positives 236, $\geq 10,000$ cells/L: 153, $\geq 100,000$: 68 (max 2,450,000 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 96 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 55,038 km², max 763,526 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (36 polygon(s))
bloom area 5,322 km²




Hull-clipped (4 footprint(s), 12,713 km²)
bloom area 7,061 km²




2018-04 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 700, positives 190, $\geq 10,000$ cells/L: 79, $\geq 100,000$: 36 (max 4,117,733 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 60 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 272,276 km², max 581,517 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (22 polygon(s))
bloom area 6,373 km²




Hull-clipped (2 footprint(s), 11,260 km²)
bloom area 4,982 km²




2018-05 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 623, positives 116, $\geq 10,000$ cells/L: 24, $\geq 100,000$: 5 (max 890,633 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 23 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 1,150 km², max 49,815 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (41 polygon(s))
bloom area 536 km²




Hull-clipped (5 footprint(s), 12,467 km²)
bloom area 766 km²




2018-06 | source used: viirs | VIIRS layer: positive

samples 797, positives 297, $\geq 10,000$ cells/L: 206, $\geq 100,000$: 119 (max 27,746,720 cells/L) | final bloom area = 88% of the hull-clipped one | 74 of 142 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 276,253 km², max 2,837,807 cells/L




USED: VIIRS-clipped (32 positive VIIRS cells)
bloom area 4,973 km²



MODIS nFLH-clipped (32 polygon(s))
bloom area 6,049 km²




Hull-clipped (2 footprint(s), 14,250 km²)
bloom area 5,662 km²




2018-07 | source used: viirs | VIIRS layer: positive

samples 719, positives 286, $\geq 10,000$ cells/L: 188, $\geq 100,000$: 126 (max 38,803,126 cells/L) | final bloom area = 99% of the hull-clipped one | 113 of 131 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 16,069,707 cells/L




USED: VIIRS-clipped (64 positive VIIRS cells)
bloom area 8,892 km²



MODIS nFLH-clipped (34 polygon(s))
bloom area 11,935 km²




Hull-clipped (3 footprint(s), 11,227 km²)
bloom area 8,992 km²




2018-08 | source used: viirs | VIIRS layer: positive

samples 1,033, positives 614, $\geq 10,000$ cells/L: 487, $\geq 100,000$: 364 (max $9e+07$ cells/L) | final bloom area = 160% of the hull-clipped one | 253 of 356 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 268,178 km², max 24,706,926 cells/L




USED: VIIRS-clipped (155 positive VIIRS cells)
bloom area 20,714 km²



MODIS nFLH-clipped (26 polygon(s))
bloom area 19,808 km²




Hull-clipped (4 footprint(s), 19,515 km²)
bloom area 12,980 km²




2018-09 | source used: viirs | VIIRS layer: positive

samples 1,026, positives 525, $\geq 10,000$ cells/L: 369, $\geq 100,000$: 234 (max 51,063,420 cells/L) | final bloom area = 206% of the hull-clipped one | 210 of 290 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 7,746,619 cells/L




USED: VIIRS-clipped (360 positive VIIRS cells)
bloom area 46,765 km²



MODIS nFLH-clipped (38 polygon(s))
bloom area 28,309 km²




Hull-clipped (2 footprint(s), 26,892 km²)
bloom area 22,738 km²




2018-10 | source used: viirs | VIIRS layer: positive

samples 1,007, positives 449, $\geq 10,000$ cells/L: 282, $\geq 100,000$: 188 (max 26,135,589 cells/L) | final bloom area = 183% of the hull-clipped one | 174 of 217 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 288,710 km², max 14,611,601 cells/L




USED: VIIRS-clipped (447 positive VIIRS cells)
bloom area 55,881 km²



MODIS nFLH-clipped (23 polygon(s))
bloom area 48,062 km²




Hull-clipped (4 footprint(s), 36,358 km²)
bloom area 30,534 km²




2018-11 | source used: viirs | VIIRS layer: positive

samples 782, positives 500, $\geq 10,000$ cells/L: 388, $\geq 100,000$: 289 (max 67,254,526 cells/L) | final bloom area = 187% of the hull-clipped one | 206 of 294 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,908 km², max 10,097,242 cells/L




USED: VIIRS-clipped (488 positive VIIRS cells)
bloom area 62,071 km²



MODIS nFLH-clipped (41 polygon(s))
bloom area 50,918 km²




Hull-clipped (2 footprint(s), 40,400 km²)
bloom area 33,223 km²




2018-12 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 494, positives 145, $\geq 10,000$ cells/L: 74, $\geq 100,000$: 21 (max 3,462,998 cells/L) | final bloom area = 284% of the hull-clipped one | 12 of 38 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 12,797 km², max 95,374 cells/L




USED: VIIRS-clipped (170 positive VIIRS cells)
bloom area 8,451 km²



MODIS nFLH-clipped (50 polygon(s))
bloom area 10,814 km²




Hull-clipped (4 footprint(s), 10,378 km²)
bloom area 2,970 km²




2019-01 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 766, positives 165, $\geq 10,000$ cells/L: 75, $\geq 100,000$: 38 (max 186,266,667 cells/L) | final bloom area = 26% of the hull-clipped one | 11 of 57 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,241 km², max 17,102,862 cells/L




USED: VIIRS-clipped (20 positive VIIRS cells)
bloom area 3,762 km²



MODIS nFLH-clipped (20 polygon(s))
bloom area 37,458 km²




Hull-clipped (2 footprint(s), 16,220 km²)
bloom area 14,312 km²




2019-02 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 633, positives 11, $\geq 10,000$ cells/L: 4, $\geq 100,000$: 2 (max 403,000 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 4 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 9,578 km², max 259,872 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (31 polygon(s))
bloom area 3,242 km²




Hull-clipped (5 footprint(s), 2,992 km²)
bloom area 1,233 km²




2019-04 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 671, positives 6, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 540 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (24 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 1,338 km²)
bloom area 0 km²




2019-05 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 672, positives 49, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 9,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 847 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (27 polygon(s))
bloom area 0 km²




Hull-clipped (3 footprint(s), 5,093 km²)
bloom area 0 km²




2019-06 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 744, positives 14, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 636 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (37 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 11,314 km²)
bloom area 0 km²




2019-07 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 728, positives 6, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 496 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (34 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 1,636 km²)
bloom area 0 km²




2019-08 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 598, positives 13, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 11,333 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 6,153 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (31 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 9,258 km²)
bloom area 0 km²




2019-09 | source used: viirs | VIIRS layer: positive

samples 703, positives 47, $\geq 10,000$ cells/L: 9, $\geq 100,000$: 2 (max 320,333 cells/L) | final bloom area = 72% of the hull-clipped one | 4 of 6 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 537 km², max 44,322 cells/L




USED: VIIRS-clipped (72 positive VIIRS cells)
bloom area 384 km²



MODIS nFLH-clipped (24 polygon(s))
bloom area 77 km²




Hull-clipped (3 footprint(s), 13,470 km²)
bloom area 537 km²




2019-10 | source used: viirs | VIIRS layer: positive

samples 762, positives 196, $\geq 10,000$ cells/L: 143, $\geq 100,000$: 104 (max 10,470,772 cells/L) | final bloom area = 44% of the hull-clipped one | 58 of 102 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 284,236 km², max 4,995,715 cells/L




USED: VIIRS-clipped (49 positive VIIRS cells)
bloom area 7,744 km²



MODIS nFLH-clipped (17 polygon(s))
bloom area 22,737 km²




Hull-clipped (3 footprint(s), 24,601 km²)
bloom area 17,728 km²




2019-11 | source used: viirs | VIIRS layer: positive

samples 876, positives 405, $\geq 10,000$ cells/L: 329, $\geq 100,000$: 271 (max 53,724,050 cells/L) | final bloom area = 76% of the hull-clipped one | 167 of 231 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 279,909 km², max 11,696,372 cells/L




USED: VIIRS-clipped (117 positive VIIRS cells)
bloom area 14,573 km²



MODIS nFLH-clipped (32 polygon(s))
bloom area 28,517 km²




Hull-clipped (3 footprint(s), 24,405 km²)
bloom area 19,156 km²




2019-12 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 533, positives 108, $\geq 10,000$ cells/L: 39, $\geq 100,000$: 16 (max 611,866 cells/L) | final bloom area = 114% of the hull-clipped one | 10 of 29 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,221 km², max 228,973 cells/L




USED: VIIRS-clipped (42 positive VIIRS cells)
bloom area 7,006 km²



MODIS nFLH-clipped (37 polygon(s))
bloom area 30,888 km²




Hull-clipped (2 footprint(s), 9,179 km²)
bloom area 6,130 km²




2020-01 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 573, positives 56, $\geq 10,000$ cells/L: 5, $\geq 100,000$: 0 (max 72,333 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 5 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 3,397 km², max 55,607 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (23 polygon(s))
bloom area 3,397 km²




Hull-clipped (5 footprint(s), 18,086 km²)
bloom area 1,694 km²




2020-02 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 449, positives 14, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 26,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,445 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (37 polygon(s))
bloom area 0 km²




Hull-clipped (3 footprint(s), 4,353 km²)
bloom area 0 km²




2020-03 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 348, positives 14, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 9,333 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 968 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (33 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 2,483 km²)
bloom area 0 km²




2020-05 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 304, positives 9, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 3,667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,412 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (25 polygon(s))
bloom area 0 km²




Hull-clipped (6 footprint(s), 2,085 km²)
bloom area 0 km²




2020-08 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 680, positives 9, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 11,667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,218 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (37 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 4,274 km²)
bloom area 0 km²




2020-09 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 569, positives 13, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,333 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 492 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (42 polygon(s))
bloom area 0 km²




Hull-clipped (2 footprint(s), 4,277 km²)
bloom area 0 km²




2020-10 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 579, positives 8, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,333 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,112 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (41 polygon(s))
bloom area 0 km²




Hull-clipped (8 footprint(s), 2,512 km²)
bloom area 0 km²




2020-11 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 460, positives 37, $\geq 10,000$ cells/L: 8, $\geq 100,000$: 0 (max 60,500 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 5 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 687 km², max 17,231 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (30 polygon(s))
bloom area 534 km²




Hull-clipped (2 footprint(s), 5,796 km²)
bloom area 458 km²




2020-12 | source used: viirs | VIIRS layer: positive

samples 652, positives 193, $\geq 10,000$ cells/L: 106, $\geq 100,000$: 71 (max 7,699,414 cells/L) | final bloom area = 202% of the hull-clipped one | 60 of 81 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 281,089 km², max 5,355,940 cells/L




USED: VIIRS-clipped (82 positive VIIRS cells)
bloom area 10,856 km²



MODIS nFLH-clipped (33 polygon(s))
bloom area 22,658 km²




Hull-clipped (2 footprint(s), 10,632 km²)
bloom area 5,363 km²




2021-01 | source used: viirs | VIIRS layer: positive

samples 684, positives 169, $\geq 10,000$ cells/L: 121, $\geq 100,000$: 72 (max 5,102,720 cells/L) | final bloom area = 201% of the hull-clipped one | 62 of 89 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,521 km², max 1,152,965 cells/L




USED: VIIRS-clipped (95 positive VIIRS cells)
bloom area 11,557 km²



MODIS nFLH-clipped (30 polygon(s))
bloom area 17,804 km²




Hull-clipped (1 footprint(s), 8,074 km²)
bloom area 5,749 km²




2021-02 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 681, positives 147, $\geq 10,000$ cells/L: 72, $\geq 100,000$: 24 (max 958,181 cells/L) | final bloom area = 66% of the hull-clipped one | 24 of 59 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 179,080 cells/L




USED: VIIRS-clipped (26 positive VIIRS cells)
bloom area 3,852 km²



MODIS nFLH-clipped (38 polygon(s))
bloom area 14,523 km²




Hull-clipped (2 footprint(s), 7,728 km²)
bloom area 5,809 km²




2021-03 | source used: viirs | VIIRS layer: positive

samples 753, positives 226, $\geq 10,000$ cells/L: 131, $\geq 100,000$: 39 (max 5,916,333 cells/L) | final bloom area = 33% of the hull-clipped one | 63 of 99 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 328,675 cells/L




USED: VIIRS-clipped (12 positive VIIRS cells)
bloom area 2,067 km²



MODIS nFLH-clipped (25 polygon(s))
bloom area 9,197 km²




Hull-clipped (1 footprint(s), 8,907 km²)
bloom area 6,202 km²




2021-04 | source used: viirs | VIIRS layer: positive | FLAGS: low_vs_hull, samples_outside

samples 873, positives 273, $\geq 10,000$ cells/L: 124, $\geq 100,000$: 38 (max 6,893,933 cells/L) | final bloom area = 0% of the hull-clipped one | 2 of 75 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 275,457 km2, max 914,571 cells/L




USED: VIIRS-clipped (7 positive VIIRS cells)
bloom area 0 km2



MODIS nFLH-clipped (26 polygon(s))
bloom area 6,105 km2




Hull-clipped (3 footprint(s), 11,994 km2)
bloom area 4,132 km2




2021-05 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 749, positives 298, $\geq 10,000$ cells/L: 135, $\geq 100,000$: 41 (max 11,526,144 cells/L) | final bloom area = 38% of the hull-clipped one | 40 of 103 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 267,749 km², max 638,460 cells/L




USED: VIIRS-clipped (18 positive VIIRS cells)
bloom area 2,979 km²



MODIS nFLH-clipped (19 polygon(s))
bloom area 11,635 km²




Hull-clipped (4 footprint(s), 13,272 km²)
bloom area 7,786 km²




2021-06 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, low_vs_hull, samples_outside

samples 1,047, positives 376, $\geq 10,000$ cells/L: 230, $\geq 100,000$: 111 (max 7,964,888 cells/L) | final bloom area = 0% of the hull-clipped one | 0 of 188 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 285,799 km², max 2,020,226 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (47 polygon(s))
bloom area 9,106 km²




Hull-clipped (2 footprint(s), 15,116 km²)
bloom area 7,361 km²




2021-07 | source used: viirs | VIIRS layer: positive

samples 904, positives 467, $\geq 10,000$ cells/L: 376, $\geq 100,000$: 266 (max 46,600,000 cells/L) | final bloom area = 93% of the hull-clipped one | 162 of 280 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,366 km², max 7,545,742 cells/L




USED: VIIRS-clipped (63 positive VIIRS cells)
bloom area 9,253 km²



MODIS nFLH-clipped (31 polygon(s))
bloom area 12,278 km²




Hull-clipped (1 footprint(s), 14,206 km²)
bloom area 9,984 km²




2021-08 | source used: viirs | VIIRS layer: positive

samples 1,020, positives 390, $\geq 10,000$ cells/L: 258, $\geq 100,000$: 175 (max 42,900,000 cells/L) | final bloom area = 144% of the hull-clipped one | 153 of 190 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 10,606,419 cells/L




USED: VIIRS-clipped (116 positive VIIRS cells)
bloom area 15,925 km²



MODIS nFLH-clipped (35 polygon(s))
bloom area 19,796 km²




Hull-clipped (1 footprint(s), 13,922 km²)
bloom area 11,058 km²




2021-09 | source used: viirs | VIIRS layer: positive

samples 987, positives 325, $\geq 10,000$ cells/L: 208, $\geq 100,000$: 110 (max 21,700,000 cells/L) | final bloom area = 88% of the hull-clipped one | 152 of 171 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,441 km², max 8,068,618 cells/L




USED: VIIRS-clipped (239 positive VIIRS cells)
bloom area 33,308 km²



MODIS nFLH-clipped (45 polygon(s))
bloom area 38,134 km²




Hull-clipped (4 footprint(s), 41,506 km²)
bloom area 37,724 km²




2021-10 | source used: viirs | VIIRS layer: positive

samples 1,174, positives 582, $\geq 10,000$ cells/L: 380, $\geq 100,000$: 197 (max 38,425,200 cells/L) | final bloom area = 155% of the hull-clipped one | 207 of 309 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 7,636,488 cells/L




USED: VIIRS-clipped (482 positive VIIRS cells)
bloom area 63,315 km²



MODIS nFLH-clipped (30 polygon(s))
bloom area 50,708 km²




Hull-clipped (3 footprint(s), 46,414 km²)
bloom area 40,846 km²




2021-11 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 960, positives 226, $\geq 10,000$ cells/L: 91, $\geq 100,000$: 28 (max 1,374,000 cells/L) | final bloom area = 112% of the hull-clipped one | 29 of 75 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 336,253 cells/L




USED: VIIRS-clipped (146 positive VIIRS cells)
bloom area 20,425 km²



MODIS nFLH-clipped (47 polygon(s))
bloom area 35,295 km²




Hull-clipped (2 footprint(s), 26,085 km²)
bloom area 18,257 km²




2021-12 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 649, positives 32, $\geq 10,000$ cells/L: 1, $\geq 100,000$: 0 (max 74,667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 1 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 3,521 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (27 polygon(s))
bloom area 0 km²




Hull-clipped (5 footprint(s), 12,862 km²)
bloom area 0 km²




2022-05 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 698, positives 13, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 6,667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 4,311 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (22 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 8,029 km²)
bloom area 0 km²




2022-06 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 785, positives 10, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,333 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 614 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (28 polygon(s))
bloom area 0 km²




Hull-clipped (3 footprint(s), 6,859 km²)
bloom area 0 km²




2022-08 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 838, positives 6, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = FALSE)
bloom area 0 km², max 661 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (41 polygon(s))
bloom area 0 km²




Hull-clipped (6 footprint(s), 1,884 km²)
bloom area 0 km²




2022-10 | source used: viirs | VIIRS layer: positive

samples 721, positives 126, $\geq 10,000$ cells/L: 58, $\geq 100,000$: 23 (max 18,400,000 cells/L) | final bloom area = 187% of the hull-clipped one | 41 of 43 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 284,307 km², max 552,662 cells/L




USED: VIIRS-clipped (113 positive VIIRS cells)
bloom area 13,247 km²



MODIS nFLH-clipped (26 polygon(s))
bloom area 28,250 km²




Hull-clipped (4 footprint(s), 10,400 km²)
bloom area 7,077 km²




2022-11 | source used: viirs | VIIRS layer: positive

samples 726, positives 363, $\geq 10,000$ cells/L: 290, $\geq 100,000$: 183 (max 388,400,000 cells/L) | final bloom area = 139% of the hull-clipped one | 119 of 193 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 3,543,436 cells/L




USED: VIIRS-clipped (133 positive VIIRS cells)
bloom area 17,950 km²



MODIS nFLH-clipped (28 polygon(s))
bloom area 28,273 km²



Hull-clipped (4 footprint(s), 15,754 km²)
bloom area 12,918 km²




cells/L


2022-12 | source used: modis | VIIRS layer: missing

samples 610, positives 260, $\geq 10,000$ cells/L: 168, $\geq 100,000$: 96 (max 36,002,150 cells/L) | final bloom area = 184% of the hull-clipped one | 118 of 122 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,054 km², max 4,789,492 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (24 polygon(s))
bloom area 22,861 km²




Hull-clipped (4 footprint(s), 16,363 km²)
bloom area 12,403 km²




2023-01 | source used: viirs | VIIRS layer: positive

samples 818, positives 363, $\geq 10,000$ cells/L: 185, $\geq 100,000$: 66 (max 5,514,014 cells/L) | final bloom area = 82% of the hull-clipped one | 97 of 149 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 284,119 km², max 545,300 cells/L




USED: VIIRS-clipped (67 positive VIIRS cells)
bloom area 9,482 km²



MODIS nFLH-clipped (34 polygon(s))
bloom area 16,766 km²




Hull-clipped (3 footprint(s), 17,841 km²)
bloom area 11,602 km²




2023-02 | source used: viirs | VIIRS layer: positive

samples 793, positives 478, $\geq 10,000$ cells/L: 363, $\geq 100,000$: 214 (max 55,655,327 cells/L) | final bloom area = 91% of the hull-clipped one | 163 of 268 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 4,241,206 cells/L




USED: VIIRS-clipped (89 positive VIIRS cells)
bloom area 13,450 km²



MODIS nFLH-clipped (25 polygon(s))
bloom area 19,170 km²




Hull-clipped (4 footprint(s), 18,284 km²)
bloom area 14,804 km²




2023-03 | source used: viirs | VIIRS layer: positive

samples 916, positives 475, $\geq 10,000$ cells/L: 298, $\geq 100,000$: 150 (max 12,085,102 cells/L) | final bloom area = 68% of the hull-clipped one | 119 of 210 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 1,026,155 cells/L




USED: VIIRS-clipped (58 positive VIIRS cells)
bloom area 9,120 km²



MODIS nFLH-clipped (16 polygon(s))
bloom area 12,802 km²




Hull-clipped (4 footprint(s), 17,365 km²)
bloom area 13,386 km²




2023-04 | source used: viirs | VIIRS layer: positive | FLAGS: samples_outside

samples 802, positives 270, $\geq 10,000$ cells/L: 93, $\geq 100,000$: 19 (max 1,023,333 cells/L) | final bloom area = 37% of the hull-clipped one | 26 of 65 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 31,111 cells/L




USED: VIIRS-clipped (20 positive VIIRS cells)
bloom area 3,512 km²



MODIS nFLH-clipped (27 polygon(s))
bloom area 10,336 km²




Hull-clipped (3 footprint(s), 12,888 km²)
bloom area 9,448 km²




2023-05 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit, samples_outside

samples 938, positives 171, $\geq 10,000$ cells/L: 20, $\geq 100,000$: 1 (max 184,500 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 9 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 5,824 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (32 polygon(s))
bloom area 0 km²




Hull-clipped (3 footprint(s), 14,326 km²)
bloom area 0 km²




2023-06 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 873, positives 13, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 420 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (26 polygon(s))
bloom area 0 km²




Hull-clipped (8 footprint(s), 6,491 km²)
bloom area 0 km²




2023-10 | source used: viirs | VIIRS layer: zero | FLAGS: zeroed_fit

samples 774, positives 33, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 4,500 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 967 cells/L




USED: VIIRS-clipped (0 positive VIIRS cells)
bloom area 0 km²



MODIS nFLH-clipped (37 polygon(s))
bloom area 0 km²




Hull-clipped (5 footprint(s), 9,575 km²)
bloom area 0 km²




2024-09 | source used: viirs | VIIRS layer: positive

samples 609, positives 58, $\geq 10,000$ cells/L: 13, $\geq 100,000$: 1 (max 126,333 cells/L) | final bloom area = 503% of the hull-clipped one | 12 of 13 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 291,361 km², max 10,927 cells/L




USED: VIIRS-clipped (163 positive VIIRS cells)
bloom area 21,384 km²



MODIS nFLH-clipped (40 polygon(s))
bloom area 26,624 km²



Hull-clipped (1 footprint(s), 5,588 km²)
bloom area 4,248 km²




cells/L


2024-10 | source used: viirs | VIIRS layer: positive

samples 758, positives 172, $\geq 10,000$ cells/L: 92, $\geq 100,000$: 35 (max 2,091,450 cells/L) | final bloom area = 145% of the hull-clipped one | 80 of 83 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 278,457 km², max 641,485 cells/L




USED: VIIRS-clipped (232 positive VIIRS cells)
bloom area 28,437 km²



MODIS nFLH-clipped (65 polygon(s))
bloom area 56,097 km²




Hull-clipped (4 footprint(s), 24,421 km²)
bloom area 19,581 km²




2024-11 | source used: viirs | VIIRS layer: positive

samples 595, positives 244, $\geq 10,000$ cells/L: 139, $\geq 100,000$: 40 (max 1,013,000 cells/L) | final bloom area = 137% of the hull-clipped one | 63 of 97 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,983 km², max 293,029 cells/L




USED: VIIRS-clipped (202 positive VIIRS cells)
bloom area 25,696 km²



MODIS nFLH-clipped




Hull-clipped (2 footprint(s), 20,823 km²)
bloom area 18,778 km²




2024-12 | source used: viirs | VIIRS layer: positive

samples 666, positives 375, $\geq 10,000$ cells/L: 197, $\geq 100,000$: 65 (max 11,948,368 cells/L) | final bloom area = 118% of the hull-clipped one | 103 of 147 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 290,981 km², max 2,487,012 cells/L




USED: VIIRS-clipped (153 positive VIIRS cells)
bloom area 21,280 km²



MODIS nFLH-clipped




Hull-clipped (3 footprint(s), 21,481 km²)
bloom area 18,016 km²




2025-01 | source used: modis | VIIRS layer: no coverage

samples 589, positives 293, $\geq 10,000$ cells/L: 165, $\geq 100,000$: 70 (max 2,952,000 cells/L) | final bloom area = 124% of the hull-clipped one | 97 of 112 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 289,547 km², max 1,711,895 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (38 polygon(s))
bloom area 23,519 km²




Hull-clipped (2 footprint(s), 24,193 km²)
bloom area 19,030 km²




2025-02 | source used: modis | VIIRS layer: no coverage

samples 709, positives 187, $\geq 10,000$ cells/L: 69, $\geq 100,000$: 33 (max 9,419,152 cells/L) | final bloom area = 91% of the hull-clipped one | 40 of 57 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 268,807 km², max 752,799 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (29 polygon(s))
bloom area 6,671 km²




Hull-clipped (4 footprint(s), 14,645 km²)
bloom area 7,329 km²




2025-03 | source used: modis | VIIRS layer: no coverage

samples 666, positives 6, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 703 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (26 polygon(s))
bloom area 0 km2




Hull-clipped (5 footprint(s), 1,570 km2)
bloom area 0 km2




2025-05 | source used: modis | VIIRS layer: no coverage

samples 541, positives 7, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 500 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 458 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (22 polygon(s))
bloom area 0 km2




Hull-clipped (4 footprint(s), 3,013 km2)
bloom area 0 km2




2025-09 | source used: modis | VIIRS layer: no coverage

samples 617, positives 10, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 2,667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 793 cells/L




VIIRS-clipped



USED: MODIS nFLH-clipped (27 polygon(s))
bloom area 0 km²




Hull-clipped (4 footprint(s), 2,788 km²)
bloom area 0 km²




2025-10 | source used: hull | VIIRS layer: no coverage

samples 637, positives 58, $\geq 10,000$ cells/L: 33, $\geq 100,000$: 19 (max 4,850,063 cells/L) | final bloom area = 100% of the hull-clipped one | 26 of 26 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 253,469 km², max 1,268,534 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 15,862 km²)
bloom area 4,075 km²




2025-11 | source used: hull | VIIRS layer: no coverage

samples 586, positives 71, $\geq 10,000$ cells/L: 33, $\geq 100,000$: 19 (max 2,502,780 cells/L) | final bloom area = 100% of the hull-clipped one | 23 of 23 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 7,359 km², max 1,881,018 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (5 footprint(s), 6,767 km²)
bloom area 1,705 km²




2025-12 | source used: hull | VIIRS layer: no coverage

samples 619, positives 108, $\geq 10,000$ cells/L: 55, $\geq 100,000$: 17 (max 620,786 cells/L) | final bloom area = 100% of the hull-clipped one | 43 of 43 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 283,635 km², max 242,070 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (2 footprint(s), 4,887 km²)
bloom area 1,704 km²




2026-01 | source used: hull | VIIRS layer: no coverage

samples 533, positives 91, $\geq 10,000$ cells/L: 36, $\geq 100,000$: 10 (max 1,000,505 cells/L) | final bloom area = 100% of the hull-clipped one | 30 of 30 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 2,222 km², max 145,399 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 8,748 km²)
bloom area 1,258 km²




2026-02 | source used: hull | VIIRS layer: no coverage

samples 537, positives 27, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 9,333 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km², max 1,601 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (3 footprint(s), 4,714 km²)
bloom area 0 km²




2026-03 | source used: hull | VIIRS layer: no coverage

samples 630, positives 22, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 499 cells/L




VIIRS-clipped



MODIS nFLH-clipped




USED: Hull-clipped (1 footprint(s), 6,632 km2)
bloom area 0 km2




2026-04 | source used: hull | VIIRS layer: no coverage

samples 654, positives 14, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 667 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 356 cells/L




VIIRS-clipped



MODIS nFLH-clipped



USED: Hull-clipped (2 footprint(s), 10,021 km2)
bloom area 0 km2




cells/L


2026-05 | source used: hull | VIIRS layer: no coverage

samples 562, positives 9, $\geq 10,000$ cells/L: 0, $\geq 100,000$: 0 (max 1,000 cells/L) | final bloom area = -% of the hull-clipped one | 0 of 0 water samples $\geq 10,000$ cells/L inside the final footprint


Unclipped sdmTMB (fitted, converged = TRUE)
bloom area 0 km2, max 739 cells/L



VIIRS-clipped



MODIS nFLH-clipped



USED: Hull-clipped (5 footprint(s), 2,131 km2)
bloom area 0 km2

