

Ebola Virus Disease mathematical models and epidemiological parameters: a systematic review and meta-analysis

Pathogen Epidemiology Review Group

*MRC Centre for Global Infectious Disease Analysis &
WHO Collaborating Centre for Infectious Disease Modelling,
Jameel Institute, School of Public Health,
Imperial College London*

Pathogen Epidemiology Review Group


MRC

Centre for
Global Infectious
Disease Analysis

Imperial College
London



- deadly **filovirus**
- transmitted through **close contact** and **bodily fluids**
 - especially during traditional burials and caregiving.
- **38 known outbreaks** since its discovery in 1976.
 - Most in **Central and Western Africa**,
- West African (WA) Ebola epidemic, caused over 11,000 reported deaths between 2013 and 2016 mainly across Guinea, Liberia and Sierra Leone.
- Four species of Ebola Virus are known to affect humans: Zaire, Bundibugyo, Sudan and Tai Forest.
- Reston, is only known to cause disease in non-human primates and Bombali, was identified in samples taken from bats in Sierra Leone.





First search until 8th March 2019. Updated until 7th July 2023.

Ebola AND ((transmissi* OR epidemiolog*) OR (model* NOT imag*) OR (severity OR "case fatality ratio*" OR CFR OR "case fatality rate*" OR "mortality rate*" OR "attack rate*") OR ("infectious period*" OR "serial interval*" OR "incubation period*" OR "generation time*" OR "generation interval*" OR "latent period*" OR latency) OR (heterogeneit* OR superspread* OR "super spread*" OR super- spread* OR overdispersion OR overdispersed OR over-dispersion OR over-dispersed OR "over dispersion" OR "over dispersed") OR (infectivity OR infectiousness OR "growth rate*" OR "reproduction number*" OR "reproductive number*" OR R0 OR "reproduction ratio*" OR "reproductive rate*") OR ("pre-existing immunity" OR serological OR serology OR serosurvey*) OR (evolution* OR mutation* OR substitution*) OR (outbreak* OR cluster* OR epidemic*) OR ("risk factor*"))).

Inclusion	Exclusion
Measures/estimates of human: Reproduction numbers (R , R_0 , R_t , r , R_e), growth rate (r), doubling times, generation time, serial interval, incubation/latent period, case fatality ratio (CFR), attack rate, mutation rate (e.g. from phylogenetic study), overdispersion, risk factors (risk and the measure). Mathematical or statistical model of transmission.	Non-English language publication .
Measures of seroprevalence and negative seroprevalence in humans.	Studies of co-infections. (local, regional, national, international).
Nature letters.	Animal studies.
	Qualitative studies, e.g., KAP studies.
	Pathogen not the primary focus of study.
	Duplicates.
	Does not match any of the inclusion criteria.
	In-vitro studies.
	Non-peer reviewed publications, conference proceedings, abstracts, posters, letters to the editor.

PRISMA flow chart








- Data package to host curated and annotated data
- Install from github
- **version 1.1.0** releases data from Lassa and Ebola review

Basic Reproduction Number

MRC


Centre for
Global Infectious
Disease Analysis

Imperial College
London




serial interval

A




incubation period

B




time from symptom onset to death

C



time from symptom onset to recovery

D




Ebola species analysis

MRC



Centre for
Global Infectious
Disease Analysis

Imperial College
London

Onset to death



Onset to recovery



Case Fatality Ratio

MRC

Centre for
Global Infectious
Disease Analysis

Imperial College
London



- We extracted 295 models.
 - 71% compartmental,
 - 6% branching process,
 - 6% agent-based models,
 - 17% other model types or combinations of models.
- Various assumptions were made in the models including homogenous mixing, heterogeneity in transmission rates between groups or over time and the latent period being the same as the incubation period.
- Only 13% of models (n=37) have any publicly available code associated with them, limiting re-usability.

- Much is already known about the Ebola Zaire species; however, our review highlights a critical lack of evidence for other species such as Sudan, Bundibugyo and Taï forest.
- Initial analyses suggest statistically significant differences in key parameters between species such as the incubation period and delays from onset to death or recovery.
- There is high variability across studies, likely driven by differences in epidemic context.

pre-print on medRxiv

