

List of Supplementary Tables

Table 1 – Meta-Analyses Evaluating Risk of HIV Transmission with Depot medroxyprogesterone acetate (DMPA)	2
Table 2 – Individual Studies of the Effects of DMPA HIV Transmission	2
Table 3 – Breast Cancer (Cohort Studies).....	5
Table 4 – Breast Cancer (Case Control Studies)	7
Table 5 – Breast Cancer (Meta-Analyses).....	10
Table 6 – Cervical Cancer	16
Table 7 – Individual Studies of the Effects of COCs on the Development of Crohn’s Disease	20
Table 8 – Individual Studies of the Effects of COCs on the Development of Ulcerative Colitis	24
Table 9 – Individual Studies of the Effects of COCs on the Development of Systemic Lupus Erythematosus	28
Table 10 – Studies of Chemical Contraceptives and Depression, Mood Disorders and Suicides.....	30
Table 11 – Individual Studies of the Effects of COCs on the Development of Multiple Sclerosis	35
Table 12 – Individual Studies of the Effects of Contraceptives on the Development of Osteoporotic Fractures	37
Table 13 – Effect of Chemical Contraceptives on Weight Gain	43
Table 14 – Relative Risk of Venous Thromboembolism in Current Users of Different Combined Hormonal Contraceptives as Compared with Nonusers Unless Otherwise Specified	46
Table 15 – Relative Risk of Thrombotic Stroke and Myocardial Infarction among Users of Selected Types of Combined Oral Contraception with Ethinyl Estradiol at a Dose of 30 to 40 µg, as Compared with Nonusers, According to Duration of Use (from Lidegaard 2012).	50

Table 1 – Meta-Analyses Evaluating Risk of HIV Transmission with Depot medroxyprogesterone acetate (DMPA)

Meta-analysis	# Included studies	Pooled Adj. OR or HR (95% CI)
Ralph et al. 2015	10 (longitudinal)	HR 1.40 (1.16–1.69)
Morrison et al. 2015	18 (longitudinal)	HR 1.50 (1.24–1.83)
Brind et al. 2015	8 (cross-sectional)	OR 1.41 (1.15–1.73)
Brind et al. 2015	16 (longitudinal)	HR 1.49 (1.28–1.73)

Table 2 – Individual Studies of the Effects of DMPA HIV Transmission

Study	Yr.(s) of study	Pop. size	Nation and locale	Subject source	Months of follow-up	Follow-up interval (months)	Type of data shown	HR or IRR (95% CI)	Weight (%)
Crook 2014	2005–2009	8,663	S Africa, Uganda, Tanzania, Zambia	Microbicide trial sero-disc. couples	12	1	Inv. Prob. W'ted HR	1.45 (1.09–1.93)	16.39
McCoy 2013	2003–2007	4,913	South Africa, Zimbabwe	Diaphragm/gel HIV prev. trial	24	3	MV HR	1.22 (0.85–1.76)	13.20
Morrison 2012	2004–2007	5,567	South Africa	General population	9–24	3	MSM HR	1.27 (0.93–1.73)	15.32
Wand 2012	Not reported	2,236	Durban, S. Africa	>90% from microbicide trial	Not reported	3	MV HR	2.02 (1.37–2.99)	12.22
Heffron 2012	2004–2010	3,790	7 African nations	Sero-discordant couples	12–24	3	MSM HR	3.93 (1.38–11.21)	2.81
Morrison 2007	1999–2004	6,109	Uganda, Zimbabwe, Thailand	Family planning clinics	21.5	3	MSM HR	1.25 (0.88–1.77)	13.86
Myer 2007	2000–2004	4,073	Cape Town, So. Africa	General population	24	6, 6, & 12	MV IRR	0.75 (0.33–1.69)	4.36
Kleinschmidt 2007	1999–2001	551	Orange Farm, So. Africa	Family planning clinic	12	3	MV HR	0.46 (0.06–3.66)	0.78
Baeten 2007	1993–1997	779	Mombasa, Kenya	CSW	120	1	MV HR	1.73 (1.28–2.34)	15.69
Kiddugavu 2003	1994–1999	5,117	Rakai, Uganda	General population	31	10	IRR, MLR	0.84 (0.41–1.72)	5.37

Risk of HIV Transmission References

Baeten JM, Benki S, Chohan V, Lavreys L, McClelland RS, Mandaliya K, Ndinya-Achola JO, Jaoko W, and Overbaugh J. Hormonal contraceptive use, herpes simplex virus infection, and risk of HIV-1 acquisition among Kenyan women. *AIDS* 2007; 21:1771–1777.

Brind J, Condly SJ, Mosher SW, Morse AR, and Kimball J. Risk of HIV infection in Depot Medroxyprogesterone Acetate (DMPA) users: a systematic review and meta-analysis. *Issues Law Med* 2015; 30:129–139.

Crook AM, Ford D, Gafos M, Hayes R, Kamali A, Kapiga S, Nunn A, Chisembele M, Ramjee G, Rees H, and McCormack S. Injectable and oral contraceptives and risk of HIV acquisition in women: an analysis of data from the MDP301 trial. *Hum Reprod* 2014; 29:1810–1817.

Govender Y, Avenant C, Verhoog NJ, Ray RM, Grantham NJ, Africander D, and Hapgood JP. The injectable-only contraceptive medroxyprogesterone acetate, unlike norethisterone acetate and progesterone, regulates inflammatory genes in endocervical cells via the glucocorticoid receptor. *PLoS One* 2014; 9:e96497.

Hapgood JP and Tomasicchio M. Modulation of HIV-1 virulence via the host glucocorticoid receptor: towards further understanding the molecular mechanisms of HIV-1 pathogenesis. *Arch Virol* 2010; 155:1009-1019. Doi: 10.1007/s00705-010-0678-0.

Hapgood JP, Ray RM, Govender Y, Avenant C, and Tomasicchio M. Differential glucocorticoid receptor-mediated effects on immunomodulatory gene expression by progestin contraceptives: implications for HIV-1 pathogenesis. *Am J Reprod Immunol* 2014; 71(6):505–512. Doi: 10.1111/aji.12214.

Heffron R, Donnell D, Rees H, Celum C, Mugo N, Were E, de Bruyn G, Nakku-Joloba E, Ngure K, Kiarie J, Coombs RW, Baeten JM; Partners in Prevention HSV/HIV Transmission Study Team. Use of hormonal contraceptives and risk of HIV-1 transmission: A prospective cohort study. *Lancet Infect Dis* 2012; 12:19–26.

Huijbregts RP, Michel KG and Hel Z. Effect of progestins on immunity: medroxyprogesterone but not norethisterone or levonorgestrel suppresses the function of T cells and pDCs. *Contraception* 2014 Aug; 90(2):123–129. Doi: 10.1016/j.contraception.2014.02.006.

Kiddugavu M, Makumbi F, Wawer MJ, Serwadda D, Sewankambo NK, Wabwire-Mangen F, Lutalo T, Meehan M, Xianbin, Gray RH; Rakai Project Study Group. Hormonal contraceptive use and HIV-1 infection in a population-based cohort in Rakai, Uganda. *AIDS* 2003; 17:233–240.

Kleinschmidt I, Rees H, Delany S, Smith D, Dinat N, Nkala B, and McIntyre JA. Injectable progestin contraceptive use and risk of HIV infection in a South African family planning cohort. *Contraception* 2007; 75:461–467.

Maritz MF, Ray RM, Bick AJ, Tomasicchio M, Woodland JG, Govender Y, Avenant C, and Hapgood JP. Medroxyprogesterone acetate, unlike norethisterone, increases HIV-1 replication in human peripheral blood mononuclear cells and an indicator cell line, via mechanisms involving the glucocorticoid receptor, increased CD4/CD8 ratios and CCR5 levels. *PLoS One* 2018 <https://doi.org/10.1371/journal.pone.0196043>.

- McCoy SI, Zheng W, Montgomery ET, Blanchard K, van der Straten A, de Bruyn G, and Padian NS. Oral and injectable contraception use and risk of HIV acquisition among women in sub-Saharan Africa. *AIDS* 2013; 27:1001–1009.
- Morrison CS, Richardson BA, Mmiro F, Chipato T, Celentano DD, Luoto J, Mugerwa R, Padian N, Rugpao S, Brown JM, Cornelisse P, Salata RA; Hormonal Contraception and the Risk of HIV Acquisition (HC-HIV) Study Group. Hormonal contraception and the risk of HIV acquisition. *AIDS* 2007; 21:85–95.
- Morrison CS, Skoler-Karpoﬀ S, Kwok C, Chen PL, van de Wijgert J, Gehret-Plagianos M, Patel S, Ahmed K, Ramjee G, Friedland B, and Lahteenmaki P.. Hormonal contraception and the risk of HIV acquisition among women in South Africa. *AIDS* 2012; 26:497–504.
- Morrison CS, Chen P-L, Kwok C, Baeten JM, Brown J, Crook AM, Van Damme L, Delany-Moretlwe S, Francis SC, Friedland BA, Hayes RJ, Heffron R, Kapiga S, Karim QA, Karpoﬀ S, Kaul R, McClelland RS, McCormack S, McGrath N, Myer L, Rees H, van der Straten A, Watson-Jones D, van de Wijgert JH, Stalter R, and Low N. Hormonal contraception and the risk of HIV acquisition: an individual participant data meta-analysis. *PLoS Med* 2015; 12:e1001778. doi: 10.1371/journal.pmed.1001778.
- Myer L, Denny L, Wright TC, and Kuhn L. Prospective study of hormonal contraception and women's risk of HIV infection in South Africa. *Int J Epidemiol* 2007; 36:166–174.
- Ralph LJ, McCoy SI, Shiu K and Padian NS. Hormonal contraceptive use and women's risk of HIV acquisition: a meta-analysis of observational studies. *Lancet Infect Dis* 2015; 15:181–189. Doi: 10.1016/S1473-3099(14)71052–1057.
- Schindler AE, Campagnoli C, Druckmann R, Huber J, Pasqualini JR, Schweppe KW, and Thijssen JH. Classification and pharmacology of progestins. *Maturitas* 2003; 46 (Suppl 1):S7-S16. Doi: 10.1016/j.maturitas.2003.09.014.
- Tomasicchio, Michele, Avenant C, Du Toit A, Ray RM, and Hapgood JP. The Progestin-Only Contraceptive Medroxyprogesterone Acetate, but Not Norethisterone Acetate, Enhances HIV-1 Vpr-Mediated Apoptosis in Human CD4+ T Cells through the Glucocorticoid Receptor. *PLoS One* 2013; 8:e62895.
- Wand H and Ramjee G. The effects of injectable hormonal contraceptives on HIV seroconversion and on sexually transmitted infections. *AIDS* 2012; 26:375–380.

Table 3 – Breast Cancer (Cohort Studies)

Study	Study Design	OR ¹ Ever Use	RR ² Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Mørch et al. 2017	Cohort		1.2 ³ (1.14–1.26)					1,797,932	* ⁴	100%
Heikkinen et al. 2016	Cohort		1.37 (1.12–1.68)					7,000	20,000	100%
Lund et al. 2007	Cohort		1.33 (1.11–1.59)					11,777	23,676	96%
Poosari et al. 2014	Cohort		1.31 (0.65–2.65)					70	11,344	92%
Phipps et al. 2011	* ⁵		0.80 ⁶ (0.68–0.94)					5,194		92%
Brohet et al. 2007 ⁷	Cohort		1.47 (1.16–1.87)					846	747	88%
Thorbjarnardottir et al. 2014	Cohort		1.32 (1.02–1.70)					654	16,928	84%
Samson et al. 2017	Cohort		1.80 ⁸ (1.29–2.55)					4816		83%
Rosenberg et al. 2010	Cohort		1.65 (1.19–2.30)					789	53,848	83%
Silvera et al. 2005	Cohort		0.88 ⁹ (0.73–1.07)					1,707	25,611	78%
Hunter et al. 2010	Cohort		1.12 (0.95–1.33)		1.33 (1.03–1.73)			1,344	115,264	73%
			1.42 ¹⁰ (1.05–1.94)							

¹ OR = odds ratio (95 % confidence interval).

² RR = relative risk (95 % confidence interval).

³ Initiation before age 20, greater than 10 years of use and evaluation within 5 yrs. of stopping further increased the risk.

⁴ Entire population of Denmark was the cohort.

⁵ Concurrent randomized clinical trials and an observational study.

⁶ Hazard ratio shown. Note that women started COCs after age 25, had been off COCs for many years.

⁷ Evaluation in patients carrying BRCA mutations. Hazard ratios shown.

⁸ Hazard ratio shown.

⁹ Hazard ratio shown.

¹⁰ Eight or more years of use.

Study	Study Design	OR ¹ Ever Use	RR ² Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
			3.05 ¹¹ (2.00–4.66)							
Trivers et al. 2007 ¹²	Cohort			1.57 (0.95–2.61)				292 ¹³	1,264 ¹⁴	67%

¹¹ Levonorgestrel containing combined oral contraceptives.

¹² Looked at mortality in patients with breast cancer over 8-10 years depending on whether they were on COCs at the time of diagnosis or within one year.

¹³ Deaths.

¹⁴ Total cohort.

Table 4 – Breast Cancer (Case Control Studies)

Study	Study Design	OR ¹⁵ Ever Use	RR ¹⁶ Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Dolle et al. 2009	Case control	2.5 (0.9-5.24)		4.2 (1.9-9.3)				898	961	100%
Lee et al. 2008	Case Case ¹⁷	0.68 (0.33-1.38)						94	444	100%
Sweeney et al. 2007	Case control	1.27 (0.99-1.63)						2,318	2,515	100%
Beaber et al. 2014b	Case control	1.5 (1.1-2.2)						985	882	100%
Li et al. 2012 ¹⁸	Case control	2.2 (1.2-4.2)						1,028	919	96%
Beaber et al. 2014a	Case control			1.5 ¹⁹ (1.3-1.9)				1,102	21,952	96%
Ichida et al. 2015	Case control			0.45 (0.22-0.90)				155	12,333	96%
Ma et al. 2010	Case control	2.87 ²⁰ (1.44-5.74)						1,197	2,015	95%
Folger et al. 2007	Case control	1.0 ²¹ (0.8-1.1)						4575	4682	92%
Jernstrom et al. 2005	Case control					2.10 (1.32-3.33)		245	745	92%
Kotsopoulos et al. 2014 ²²	Case control	1.45 ²³ (1.20-1.75)						2,492	2,492	88%
		1.19 ²⁴ (0.99-1.42)								
Figueiredo et al. 2010 ²⁵	Case control					2.38 (0.72-7.83)		705	1,398	86%

¹⁵ OR = odds ratio (95 % confidence interval).

¹⁶ RR = relative risk (95 % confidence interval).

¹⁷ BRCA1 and BRCA2 carriers with breast cancer.

¹⁸ Population-based case-control of women 20-44 yo with recent DMPA use for at least 12 months.

¹⁹ Use within the past year of COCs increases risk of breast cancer.

²⁰ Triple negative breast cancer if less than 18 yo on COCs.

²¹ Evaluated short-term use only.

²² Study of BRCA+ patients.

²³ <20 years old.

²⁴ 20-25 years old.

²⁵ Evaluation of BRCA1 and BRCA2 carriers; controls with unilateral breast cancer compared with contralateral cases.

Study	Study Design	OR ¹⁵ Ever Use	RR ¹⁶ Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Veneroso et al. 2008	Case Case ²⁶	1.12 (1.03-1.23)						116	99	86%

Study	Study Design	OR ²⁷ Ever Use	RR ²⁸ Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Ma et al. 2006	Case control	1.27 ²⁹ (0.75-2.14)				0.76 (0.49-1.18)		1,366	440	84%
		0.76 ³⁰ (0.49-1.18)								
Rosenberg et al. 2008	Case control	1.5 ³¹ (1.2-1.8)						907	1,711	83%
Haile et al. 2006	Case control	0.77 ³² (0.53-1.12)						195	497	83%
		1.62 ³³ (0.90-2.92)						128	307	
Milne et al. 2005	Case control	1.52 (1.22-1.91)						1156	815	83%
Amadou et al. 2013	Case control	1.68 (0.67-4.21)						1,000	1,074	75%
Ozmen et al. 2009	Case control	0.60 (0.48-0.74)						1,492	2,167	74%
Delort et al. 2007	Population based ³⁴	1.84 ³⁵ (1.38-2.44)						934		71%
Beji et al. 2006	Case control	1.98 (1.38-2.85)						405	1,050	63%
Veisy et al. 2015	Case control	2.11 (1.44-3.08)						235	235	63%
Tehrani et al. 2010	Case control	2.83 (1.87-4.24)						321	321	58%

²⁶ Comparison of more aggressive with less aggressive cases.

²⁷ OR = odds ratio (95 % confidence interval).

²⁸ RR = relative risk (95 % confidence interval).

²⁹ ER-/PR-

³⁰ ER+/PR+

³¹ OR for 5+ years of use.

³² BRCA1+ patients.

³³ BRCA2+ patients.

³⁴ Population-based study of early onset breast cancer.

³⁵ OR for developing breast cancer 2 years earlier than non-users.

Study	Study Design	OR ²⁷ Ever Use	RR ²⁸ Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Lumachi et al. 2010	Retrospective Review	2.06 (1.14-3.70)						404	408	33%

Table 5 – Breast Cancer (Meta-Analyses)

Study	Study Design	OR ³⁶ Ever Use	RR ³⁷ Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Kahlenborn et al. 2006 ³⁸	Meta-analysis	1.19 (1.09-1.29)						18,406	27,677	91%
		1.29 ³⁹ (1.20-1.40)								
		1.24 ⁴⁰ (0.92-1.67)								
		1.44 ⁴¹ (1.28-1.62)								
Bethea et al. 2015	Meta-analysis	1.46 ⁴² (1.18-1.81)						1,848	10,044	85%
		1.57 ⁴³ (1.22-1.43)						1,043	10,044	
		1.78 ⁴⁴ (1.25-2.53)						494	10,044	
Zhu et al. 2012	Meta-analysis	1.08 ⁴⁵ (0.99-1.17)								54%
Friebel et al. 2014 ⁴⁶	Meta-analysis	1.36 ⁴⁷ (0.99-1.88)								27%
		1.51 ⁴⁸ (1.10-2.08)								
Moorman et al. 2013	Meta-analysis	1.21 ⁴⁹ (0.93-1.58)								

³⁶ OR = odds ratio (95 % confidence interval).

³⁷ RR = relative risk (95 % confidence interval).

³⁸ Limited to case-control studies from 1980-2004.

³⁹ Parous women.

⁴⁰ Nulliparous women.

⁴¹ Use before first full term pregnancy among parous women.

⁴² ER+

⁴³ ER-

⁴⁴ Triple negative.

⁴⁵ For each 5 years on COCs the risk increased by 7%, but statistical significance not achieved.

⁴⁶ Study limited to BRCA1 and BRCA2 mutation carriers.

⁴⁷ 1-3 years of use.

⁴⁸ >3 years of use.

⁴⁹ 8 studies on BRCA1+ or BRCA2+ patients and breast cancer risk with CSC use.

Breast Cancer References

- Adams-Campbell LL, Makambi KH, Frederick WA, Gaskins M, Dewitty RL, and McCaskill-Stevens W. Breast cancer risk assessments comparing Gail and CARE models in African-American women. *Breast J*. 15 Suppl 2009; 1:S72–75.
- Amadou A, Fabre A, Torres-Mejía G, Ortega-Olvera C, Angeles-Llerenas A, McKenzie F, Biessy C, Hainaut P, and Romieu I. Hormonal therapy and risk of breast cancer in Mexican women. *PLoS One* 2013; 8:e79695.
- Beaber EF, Buist DS, Barlow WE, Malone KE, Reed SD, and Li CI. Recent oral contraceptive use by formulation and breast cancer risk among women 20 to 49 years of age. *Cancer Res* 2014a; 74:4078–4089.
- Beaber EF, Malone KE, Tang MT, Barlow WE, Porter PL, Daling JR, and Li CI. Oral contraceptives and breast cancer risk overall and by molecular subtype among young women. *Cancer Epidemiol Biomarkers Prev* 2014b; 23:755–764.
- Beji NK and Reis N. Risk factors for breast cancer in Turkish women: a hospital-based case-control study. *Eur J Cancer Care (Engl)*. 2007; 16:178–184.
- Bethea TN, Rosenberg L, Hong CC, Troester MA, Lunetta KL, Bandera EV, Schedin P, Kolonel LN, Olshan AF, Ambrosone CB, and Palmer JR. A case-control analysis of oral contraceptive use and breast cancer subtypes in the African American Breast Cancer Epidemiology and Risk Consortium. *Breast Cancer Res* 2015; 17:22. Doi: 10.1186/s13058-015-0535-x.
- Brohet RM, Goldgar DE, Easton DF, Antoniou AC, Andrieu N, Chang-Claude J, Peock S, Eeles RA, Cook M, Chu C, Noguès C, Lasset C, Berthet P, Meijers-Heijboer H, Gerdes AM, Olsson H, Caldes T, van Leeuwen FE, and Rookus MA. Oral contraceptives and breast cancer risk in the international BRCA1/2 carrier cohort study: a report from EMBRACE, GENEPSO, GEO-HEBON, and the IBCCS Collaborating Group. *J Clin Oncol* 2007; 25:3831–3836.
- Delort L, Kwiatkowski F, Chalabi N, Satih S, Bignon YJ, and Bernard-Gallon DJ. Risk factors for early age at breast cancer onset—the “COSA program” population-based study. *Anticancer Res* 2007; 27:1087–1094.
- Dolle JM, Daling JR, White E, Brinton LA, Doody DR, Porter PL, and Malone KE. Risk factors for triple-negative breast cancer in women under the age of 45 years. *Cancer Epidemiol Biomarkers Prev* 2009; 18:1157–1166.
- Figueiredo JC, Haile RW, Bernstein L, Malone KE, Largent J, Langholz B, Lynch CF, Bertelsen L, Capanu M, Concannon P, Borg A, Børresen-Dale AL, Diep A, Teraoka S, Torngren T, Xue S, and Bernstein JL. Oral contraceptives and postmenopausal hormones and risk of contralateral breast cancer among BRCA1 and BRCA2 mutation carriers and noncarriers: the WECARE Study. *Breast Cancer Res Treat* 2010; 120:175–183.
- Folger SG, Marchbanks PA, McDonald JA, Bernstein L, Ursin G, Berlin JA, Daling JR, Norman SA, Strom BL, Weiss LK, Simon MS, Burkman RT, Malone KE, and Spirtas R. Risk of breast cancer associated with short-term use of oral contraceptives. *Cancer Causes Control* 2007; 18:189–198.
- Friebel TM, Domchek SM, and Rebbeck TR. Modifiers of cancer risk in BRCA1 and BRCA2 mutation carriers: systematic review and meta-analysis. *J Natl Cancer Inst* 2014; 106(6):dju091. Doi: 10.1093/jnci/dju091.

Haile RW, Thomas DC, McGuire V, Felberg A, John EM, Milne RL, Hopper JL, Jenkins MA, Levine AJ, Daly MM, Buys SS, Senie RT, Andrulis IL, Knight JA, Godwin AK, Southey M, McCredie MR, Giles GG, Andrews L, Tucker K, Miron A, Apicella C, Tesoriero A, Bane A, Pike MC; kConFab Investigators; Ontario Cancer Genetics Network Investigators, and Whittemore AS. BRCA1 and BRCA2 mutation carriers, oral contraceptive use, and breast cancer before age 50. *Cancer Epidemiol Biomarkers Prev* 2006; 15:1863–1870.

Heikkinen S, Koskenvuo M, Malila N, Sarkeala T, Pukkala E, and Pitkaniemi J. Use of exogenous hormones and the risk of breast cancer: results from self-reported survey data with validity assessment. *Cancer Causes Control* 2016; 27:249–258.

Hunter DJ, Colditz GA, Hankinson SE, Malspeis S, Spiegelman D, Chen W, Stampfer MJ, and Willett WC. Oral contraceptive use and breast cancer: a prospective study of young women. *Cancer Epidemiol Biomarkers Prev* 2010; 19:2496–2502.

Ichida M, Kataoka A, Tsushima R, and Taguchi T. No increase in breast cancer risk in Japanese women taking oral contraceptives: a case-control study investigating reproductive, menstrual and familial risk factors for breast cancer. *Asian Pac J Cancer Prev* 2015; 16:3685–3690.

Jernström H, Loman N, Johannsson OT, Borg A, and Olsson H. Impact of teenage oral contraceptive use in a population-based series of early-onset breast cancer cases who have undergone BRCA mutation testing. *Eur J Cancer* 2005; 41:2312–2320.

Kahlenborn C, Modugno F, Potter DM, and Severs WB. Oral Contraceptive Use as a Risk Factor for Premenopausal Breast Cancer: A Meta-analysis. *Mayo Clin Proc* 2006; 81:1290–1302.

Kotsopoulos J, Lubinski J, Moller P, Lynch HT, Singer CF, Eng C, Neuhausen SL, Karlan B, Kim-Sing C, Huzarski T, Gronwald J, McCuaig J, Senter L, Tung N, Ghadirian P, Eisen A, Gilchrist D, Blum JL, Zakalik D, Pal T, Sun P, and Narod SA; Hereditary Breast Cancer Clinical Study Group. Timing of oral contraceptive use and the risk of breast cancer in BRCA1 mutation carriers. *Breast Cancer Res Treat* 2014; 143:579–586.

Lee E, Ma H, McKean-Cowdin R, Van Den Berg D, Bernstein L, Henderson BE, and Ursin G. Effect of reproductive factors and oral contraceptives on breast cancer risk in BRCA1/2 mutation carriers and noncarriers: results from a population-based study. *Cancer Epidemiol Biomarkers Prev* 2008; 17:3170–3178.

Li CI, Beaber EF, Tang MT, Porter PL, Daling JR, and Malone KE. Effect of depo-medroxyprogesterone acetate on breast cancer risk among women 20 to 44 years of age. *Cancer Res* 2012; 72:2028–2035.

Lumachi F, Frigo AC, Basso U, Tombolan V, and Ermani M. Estrogen therapy and risk of breast cancer in postmenopausal women: a case-control study and results of a multivariate analysis. *Menopause* 2010; 17:524–528.

Lund E, Bakken K, Dumeaux V, Andersen V, and Kumle M. Hormone replacement therapy and breast cancer in former users of oral contraceptives--The Norwegian Women and Cancer study. *Int J Cancer* 2007; 121:645–648.

Ma H, Bernstein L, Ross RK, and Ursin G. Hormone-related risk factors for breast cancer in women under age 50 years by estrogen and progesterone receptor status: results from a case-control and a case-case comparison. *Breast Cancer Res* 2006; 8:R39.

Ma H, Wang Y, Sullivan-Halley J, Weiss L, Marchbanks PA, Spirtas R, Ursin G, Burkman RT, Simon MS, Malone KE, Strom BL, McDonald JA, Press MF, and Bernstein L. Use of four biomarkers to evaluate the risk of breast cancer subtypes in the women's contraceptive and reproductive experiences study. *Cancer Res* 2010; 70:575–587.

Mehrgou A and Akouchekian M. The importance of BRCA1 and BRCA2 genes mutations in breast cancer development. *Med J Islam Repub Iran* 2016; 30:369.

Milne RL, Knight JA, John EM, Dite GS, Balbuena R, Ziogas A, Andrulis IL, West DW, Li FP, Southey MC, Giles GG, McCredie MR, Hopper JL, and Whittemore AS. Oral contraceptive use and risk of early-onset breast cancer in carriers and noncarriers of BRCA1 and BRCA2 mutations. *Cancer Epidemiol Biomarkers Prev* 2005; 14:350–356.

Moorman PG, Havrilesky LJ, Gierisch JM, Coeytaux RR, Lowery WJ, Peragallo Urrutia R, Dinan M, McBroom AJ, Hasselblad V, Sanders GD, and Myers ER. Oral contraceptives and risk of ovarian cancer and breast cancer among high-risk women: a systematic review and meta-analysis. *J Clin Oncol* 2013;31(33): 4188–4198. Doi: 10.1200/JCO.2013.48.9021.

Mørch LS, Skovlund CW, Hannaford PC, Iversen L, Fielding S, and Lidegaard Ø. Contemporary hormonal contraception and the risk of breast cancer. *N Engl J Med* 2017; 377:2228–2239.

Ozmen V, Ozcinar B, Karanlik H, Cabioglu N, Tukenmez M, Disci R, Ozmen T, Igci A, Muslumanoglu M, Kecer M, and Soran A. Breast cancer risk factors in Turkish women—a University Hospital based nested case control study. *World J Surg Oncol* 2009; 7:37.

Parkin DM. Cancers attributable to exposure to hormones in the UK in 2010. *British Journal of Cancer* 2011; 105: S42–S48.

Phipps AI, Chlebowski RT, Prentice R, McTiernan A, Wactawski-Wende J, Kuller LH, Adams-Campbell LL, Lane D, Stefanick ML, Vitolins M, Kabat GC, Rohan TE, and Li CI. Reproductive history and oral contraceptive use in relation to risk of triple-negative breast cancer. *J Natl Cancer Inst* 2011; 103:470–477.

Poosari A, Promthet S, Kamsa-ard S, Suwanrungruang K, Longkul J, and Wiangnon S. Hormonal contraceptive use and breast cancer in Thai women. *J Epidemiol* 2014; 24:216–220.

Rosenberg L, Zhang Y, Coogan PF, Strom BL, and Palmer JR. A case-control study of oral contraceptive use and incident breast cancer. *Am J Epidemiol* 2008; 169:473–479.

Rosenberg L, Boggs DA, Wise LA, Adams-Campbell LL, and Palmer JR. Oral contraceptive use and estrogen/progesterone receptor-negative breast cancer among African American women. *Cancer Epidemiol Biomarkers Prev* 2010; 19:2073–2079.

Samson ME, Adams SA, Mulatya CM, Zhang J, Bennett CL, Hebert J, and Steck SE. Types of oral contraceptives and breast cancer survival among women enrolled in Medicaid: A competing-risk model. *Maturitas* 2017; 95:42–49.

Silvera SA, Miller AB, and Rohan TE. Oral contraceptive use and risk of breast cancer among women with a family history of breast cancer: a prospective cohort study. *Cancer Causes Control*. 2005; 16:1059–1063.

Soini T, Hurskainen R, Grénman S, Mäenpää J, Paavonen J, and Pukkala E. Cancer risk in women using the levonorgestrel-releasing intrauterine system in Finland. *Obstet Gynecol* 2014; 124 (2 Pt 1): 292–299.

Sweeney C, Giuliano AR, Baumgartner KB, Byers T, Herrick JS, Edwards SL, and Slattery ML. Oral, injected and implanted contraceptives and breast cancer risk among U.S. Hispanic and non-Hispanic white women. *Int J Cancer* 2007; 121:2517–2523.

Thorbjarnardottir T, Olafsdottir EJ, Valdimarsdottir UA, Olafsson O, and Tryggvadottir L. Oral contraceptives, hormone replacement therapy and breast cancer risk: a cohort study of 16 928 women 48 years and older. *Acta Oncol* 2014; 53:752–758.

Trivers KF, Gammon MD, Abrahamson PE, Lund MJ, Flagg EW, Moorman PG, Kaufman JS, Cai J, Porter PL, Brinton LA, Eley JW, and Coates RJ. Oral contraceptives and survival in breast cancer patients aged 20 to 54 years. *Cancer Epidemiol Biomarkers Prev* 2007; 16:1822–1827.

Veisy A, Lotfinejad S, Salehi K, and Zhian F. Risk of breast cancer in relation to reproductive factors in North-West of Iran, 2013-2014. *Asian Pac J Cancer Prev* 2015; 16:451–455.

Veneroso C, Siegel R, and Levine PH. Early age at first childbirth associated with advanced tumor grade in breast cancer. *Cancer Detect Prev* 2008; 32:215–223.

Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for reporting observational studies. *PLoS Medicine* 2007; 4:1623–27 (e296). Doi: 10.1371/journal.pmed. 0040296.

Williams WV, Carlson K, Mitchell LA, and Raviele K. Association of Combined Estrogen-Progestogen and Progestogen Only Contraceptives with the Development of Cancer. *The Linacre Quarterly* 2018; 85(4): 412–452.

World Health Organization International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risk to Humans. Combined Estrogen–Progestogen Contraceptives and Combined Estrogen–Progestogen Menopausal Therapy. 2007; <http://monographs.iarc.fr/ENG/Monographs/vol91/mono91.pdf>.

World Health Organization International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risk to Humans_Combined Estrogen–Progestogen Contraceptives. *IARC Monographs* 2012; 100A:283–318; <http://monographs.iarc.fr/ENG/Monographs/vol100A/mono100A-19.pdf>.

Yager JD and Davidson NE. Estrogen Carcinogenesis in Breast Cancer. *N Engl J Med* 2006; 354:270-282.

Zhu H, Lei X, Feng J, and Wang Y. Oral contraceptive use and risk of breast cancer: a meta-analysis of prospective cohort studies. *Eur J Contracept Reprod Health Care* 2012; 17:402–414.

Table 6 – Cervical Cancer

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Roura et al. 2016	Cohort Study		1.1 ¹ (0.9–1.3)		1.8 ¹⁰ (1.4–2.4)		1 ¹⁰ (0.9–1.3)	1,065	306,971	94%
			1.6 ² (1.1–2.3)		2.2 ⁸ (1.3–4.0)		1.6 ⁸ (1.1–2.2)	261	306,971	
Leslie et al. 2014	Case Control Study	1.35 ³ (0.99–1.85)						219	2,300	87%
McFarlane-Anderson et al. 2008	Case Control Study	1.59 ⁴ (0.87–2.82)						240	102	83%
		2.48 ⁵ (1.30–4.74)								
Vanakankovit et al. 2008	Case Control Study	1.49 (0.79–2.64)						60	180	76%
Wilson et al. 2013	Case Control Study	1.22 (0.96–1.56)						724	3,479	76%
Matos et al. 2005	Case Control Study	1.3 (0.8–3.1)						140	157	47%
International Collaboration 2007 ⁶	Meta-analysis	1.05 ⁷ (1.04–1.07)						16,573	35,509	97%
	<5 years of use	0.96 (0.04) ⁸								
	5–9 years of use	1.2 (0.05) ⁵								
	10+ years of use	1.56 (0.08) ⁵								
	<5 years of use	1.07 (0.08) ⁹						7,227	19,335	
	5+ years of use	1.22 (0.11) ⁶								
Moreno 2002 ¹⁰	Meta-analysis							1676	255	95%
	Invasive cervical cancer (ICC)	1.29 (0.88–1.91)								
	ICC 5+ years of use	4.01 (2.01–8.02)								

¹ Includes Cervical Intraepithelial Neoplasia Grade 3, carcinoma in situ and invasive cervical cancer.

² Analysis limited to invasive cervical cancer.

³ Study limited to HIV+ women.

⁴ Combined hormonal contraceptives.

⁵ Progesterone only contraceptives.

⁶ Meta-analysis of 24 studies (15 cohort and 9 case-control studies).

⁷ Relative risk per year of use for current users of combined hormonal contraceptives.

⁸ Floating standard error shown for users of combined hormonal contraceptives.

⁹ Progestin only contraceptives. Floating standard error shown. The 95% CI for 5+ years of use is 1.01–1.46.

¹⁰ Pooled data from 8 case-control studies of invasive cervical cancer and 2 of carcinoma in situ, analyzing only the subset positive for Human Papilloma Virus DNA in cervical cells.

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
	In situ carcinoma (ISC)	1.42 (0.99-2.04)								
	ISC 5+ years of use	3.42 (2.13-5.48)								

Cervical Cancer References

World Health Organization International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risk to Humans. Combined Estrogen–Progestogen Contraceptives and Combined Estrogen–Progestogen Menopausal Therapy. 2007; <http://monographs.iarc.fr/ENG/Monographs/vol91/mono91.pdf>.

World Health Organization International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risk to Humans. Combined Estrogen–Progestogen Contraceptives. *IARC Monographs* 2012; 100A:283–318; <http://monographs.iarc.fr/ENG/Monographs/vol100A/mono100A-19.pdf>.

International Collaboration of Epidemiological Studies of Cervical Cancer, Appleby P, Beral V, Berrington de González A, Colin D, Franceschi S, Goodhill A, Green J, Peto J, Plummer M, and Sweetland S. Cervical cancer and hormonal contraceptives: collaborative reanalysis of individual data for 16,573 women with cervical cancer and 35,509 women without cervical cancer from 24 epidemiological studies. *Lancet* 2007; 370:1609–1621.

Leslie HH, Karasek DA, Harris LF, Chang E, Abdulrahim N, Maloba M, and Huchko MJ. Cervical cancer precursors and hormonal contraceptive use in HIV-positive women: application of a causal model and semi-parametric estimation methods. *PLoS One* 2014; 9:e101090.

Matos A, Moutinho J, Pinto D, and Medeiros R. The influence of smoking and other cofactors on the time to onset to cervical cancer in a southern European population. *Eur J Cancer Prev* 2005; 14:485–491.

McFarlane-Anderson N, Bazuaye PE, Jackson MD, Smikle M, and Fletcher HM. Cervical dysplasia and cancer and the use of hormonal contraceptives in Jamaican women. *BMC Womens Health* 2008; 8:9.

Moreno V, Bosch FX, Muñoz N, Meijer CJ, Shah KV, Walboomers JM, Herrero R, Franceschi S; International Agency for Research on Cancer. Multicentric Cervical Cancer Study Group. Effect of oral contraceptives on risk of cervical cancer in women with human papillomavirus infection: the IARC multicentric case-control study. *Lancet* 2002; 359(9312):1085–1092.

Roura E, Travier N, Waterboer T, de Sanjosé S, Bosch FX, Pawlita M, Pala V, Weiderpass E, Margall N, Dillner J, Gram IT, Tjønneland A, Munk C, Palli D, Khaw KT, Overvad K, Clavel-Chapelon F, Mesrine S, Fournier A, Fortner RT, Ose J, Steffen A, Trichopoulou A, Lagiou P, Orfanos P, Masala G, Tumino R, Sacerdote C, Polidoro S, Mattiello A, Lund E, Peeters PH, Bueno-de-Mesquita HB, Quirós JR, Sánchez MJ, Navarro C, Barricarte A, Larrañaga N, Ekström J, Lindquist D, Idahl A, Travis RC, Merritt MA, Gunter MJ, Rinaldi S, Tommasino M, Franceschi S, Riboli E, and Castellsagué X. The Influence of Hormonal Factors on the Risk of Developing Cervical Cancer and Pre-Cancer: Results from the EPIC Cohort. *PLoS One* 2016; 11:e0147029.

Vanakankovit N and Taneepanichskul S. Effect of oral contraceptives on risk of cervical cancer. *J Med Assoc Thai* 2008; 91:7–12.

Wilson JC, O'Rourke MA, Cooper JA, Murray LJ, Hughes CM, Gormley GJ, and Anderson LA. Non-steroidal anti-inflammatory drug use and cervical cancer risk: a case-control study using the Clinical Practice Research Datalink. *Cancer Epidemiol* 2013; 37:897–904.

Table 7 – Individual Studies of the Effects of COCs on the Development of Crohn’s Disease

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Khalili et al. 2013 ¹	Cohort		1.43 (1.08–1.90)		2.82 (1.65–4.82)		1.39 (1.05–1.85)	315	117,060	93%
García Rodríguez et al. 2005 ²	Cohort				1.94 (0.85–4.45)		1.04 (0.50–2.17)	171	10,000	88%
Logan and Kay 1989	Cohort		1.7 (0.88–3.2)					42	45,958	54%
Vessey et al. 1986 ³	Cohort				1.33			18	17,014	46%
Boyko et al. 1994	Case-control		2 (1.0–3.7)					91	169	94%
Katschinski 1993 ⁴	Case-control				2.5 (0.75–4.6)					93%
Katschinski 1993 ⁵	Case-control				3.1 (1.1–6.7)					93%
Lashner et al. 1989	Case-control	1 (0.46–2.16)		0.73 (0.34–1.59)		1.8 (0.61–5.29)		51	51	88%
Lesko et al. 1985 ⁶	Case-control		1.7 (1.0–3.2)					57	2189	83%
Sandler et al. 1992	Case-control		1.49 (0.99–2.26)					184	217	81%
Persson et al. 1993	Case-control		1.7 (0.9–3.2)					152	305	81%
Halfvarson et al. 2006 ⁷	Case-control				1.5 (0.4–5.3)			102	102	75%
Lowe et al. 2009 ⁸	Case-control		1.05					21,172	754,6131	74%
Ng et al. 2012 ⁹	Case-control	4						125	125	74%

¹ Hazard ratios (RR adjusted for time).

² OR increased with duration of use.

³ Authors’ calculation adjusted for smoking.

⁴ Adjusted RR for 1–3 years prior to disease onset.

⁵ Adjusted RR for >3 years prior to disease onset.

⁶ RR is from multiple logistic regression analysis.

⁷ Monozygotic and dizygotic twins.

⁸ Adjusted incidence rate ratio.

⁹ Twins study.

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
		(1.1–14.2)								
Ng et al. 2012 ¹⁰	Case-control	9.04 (1.11–73.6)								74%
Sicilia et al. 2001	Case-control	2.8 (1.01–7.77)						103	103	71%
Corrao et al. 1998	Case-control ever use			3.4 (1.0–11.9)		1.8 (0.4–7.3)		225	225	67%
Katschinski 1993 ¹¹	Case-control		4.3 (1.3–14.4)					83	83	57%
Han et al. 2010	Case-control		0.66 (0.38–1.15)					315	536	52%
Calkins et al. 1986 ¹²	Case-control	1.14 (0.44–2.96)						66	67	42%
Calkins et al. 1986 ¹³	Case-control	1.6 (0.59–4.37)						66	71	42%
Vcev et al. 2015	Case-control	0.28 (0.03–2.46)						11	42	31%
Cornish et al. 2008	Meta-analysis				1.46 (1.26–1.70)		1.04 (0.816–.340)	1251	74,564	91%
Cornish et al. 2008 ¹⁴	Meta-analysis				1.58 (1.07–2.40)					91%
Godet et al. 1995 ¹⁵	Meta-analysis		1.44 (1.12–1.86)					531	49,156	82%

¹⁰ Multivariate analysis.

¹¹ RR for use >3 years.

¹² Hospital controls.

¹³ Neighborhood controls.

¹⁴ High quality studies.

¹⁵ Adjusted for smoking.

Crohn's Disease References

- Boyko EJ, Theis MK, Vaughan TL, and Nicol-Blades B. Increased risk of inflammatory bowel disease associated with oral contraceptive use. *American Journal of Epidemiology* 1994; 140:268–278.
- Calkins BM, Mendeloff AI, and Garland C. Inflammatory bowel disease in oral contraceptive users. *Gastroenterology* 1986; 91: 523–524.
- Cornish JA, Tan E, Simillis C, Clark SK, Teare J, and Tekkis PP. The risk of oral contraceptives in the etiology of inflammatory bowel disease: a meta-analysis. *American Journal of Gastroenterology* 2008; 103:2394–2400.
- Corrao G, Tragnone A, Caprilli R, Trallori G, Papi C, Andreoli A, Di Paolo M, Riegler G, Rigo GP, Ferraù O, Mansi C, Ingrosso M, and Valpiani D. Risk of inflammatory bowel disease attributable to smoking, oral contraception and breastfeeding in Italy: a nationwide case-control study. Cooperative Investigators of the Italian Group for the Study of the Colon and the Rectum (GISC). *International Journal of Epidemiology* 1998; 27:397–404.
- García Rodríguez LA, González-Pérez A, Johansson S, and Wallander MA. Risk factors for inflammatory bowel disease in the general population. *Alimentary Pharmacology & Therapeutics* 2005; 22:309–315.
- Godet PG, May GR, and Sutherland LR. Meta-analysis of the role of oral contraceptive agents in inflammatory bowel disease. *Gut* 1995; 37:668–673.
- Halfvarson J, Jess T, Magnuson A, Montgomery SM, Orholm M, Tysk C, Binder V, and Järnerot G. Environmental factors in inflammatory bowel disease: a co-twin control study of a Swedish-Danish twin population. *Inflammatory Bowel Disease* 2006; 12:925–933.
- Han DY, Fraser AG, Dryland P, and Ferguson LR. Environmental factors in the development of chronic inflammation: a case-control study on risk factors for Crohn's disease within New Zealand. *Mutation Research* 2010; 690:116–122.
- Katschinski B. [Smoking and ovulation inhibitor in inflammatory bowel diseases]. *Medizinische Klinik (Munich, Germany)* 88 Suppl 1993; 1:5–8.
- Khalili H, Higuchi LM, Ananthakrishnan AN, Richter JM, Feskanich D, Fuchs CS, and Chan AT. Oral contraceptives, reproductive factors and risk of inflammatory bowel disease. *Gut* 2013; 62:1153–1159.
- Lashner BA, Kane SV, and Hanauer SB. Lack of association between oral contraceptive use and Crohn's disease: a community-based matched case-control study. *Gastroenterology* 1989; 97:1442–1447.
- Lesko SM, Kaufman DW, Rosenberg L, Helmrich SP, Miller DR, Stolley PD, and Shapiro S. Evidence for an increased risk of Crohn's disease in oral contraceptive users. *Gastroenterology* 1985; 89:1046–1049.
- Logan RF and Kay CR. Oral contraception, smoking and inflammatory bowel disease—findings in the Royal College of General Practitioners Oral Contraception Study. *International Journal of Epidemiology* 1989; 18: 105–107.
- Lowe AM, Roy PO, Poulin M, Michel P, Bitton A, St-Onge L, and Brassard P. Epidemiology of Crohn's disease in Quebec, Canada. *Inflammatory Bowel Disease* 2009; 15:429–435.

Ng SC, Woodrow S, Patel N, Subhani J, and Harbord M. Role of genetic and environmental factors in British twins with inflammatory bowel disease. *Inflammatory Bowel Disease* 2012; 18:725–736.

Persson PG, Leijonmarck CE, Bernell O, Hellers G, and Ahlbom A. Risk indicators for inflammatory bowel disease. *International Journal of Epidemiology* 1993; 2(2):268–272.

Sandler RS, Wurzelmann JI, and Lyles CM. Oral contraceptive use and the risk of inflammatory bowel disease. *Epidemiology* 1992; 3:374–378.

Sicilia, B., C. López Miguel, F. Arribas, J. López Zaborras, E. Sierra, and F. Gomollón. Environmental risk factors and Crohn's disease: A population-based, case-control study in Spain. *Digestive and Liver Disease* 2001; 33: 762–67.

Vcev A, Pezerovic D, Jovanovic Z, Nakic D, Vcev I, and Majnarić L. A retrospective, case-control study on traditional environmental risk factors in inflammatory bowel disease in Vukovar-Srijem County, north-eastern Croatia 2010. *Wiener Klinische Wochenschrift* 2015; 127:345–354.

Vessey M, Jewell D, Smith A, Yeates D, and McPherson K. Chronic inflammatory bowel disease, cigarette smoking, and use of oral contraceptives: findings in a large cohort study of women of childbearing age. *British Medical Journal (Clinical Research Edition)* 1986; 292:1101–1113.

Table 8 – Individual Studies of the Effects of COCs on the Development of Ulcerative Colitis

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Khalili et al. 2013 ¹	Cohort		1.18 (0.92–1.52)		1.22 (0.74–2.07)		1.18 (0.91–1.52)	392	116,983	93%
García Rodríguez et al. 2005	Cohort				1.58 (0.71–3.52)		0.67 (0.32–1.39)	222	10,000	88%
Logan and Kay 1989	Cohort		1.3 (0.82–2.0)					78	45,922	54%
Vessey et al. 1986 ²	Cohort				2.1			31	17,001	46%
Boyko et al 1994	Case-control		1.7 (1.1–2.7)					211	341	94%
Lashner et al. 1990	Case-control	0.86 (0.40–1.85)		0.7 (0.27–1.83)		1.14 (0.41–.15)		46	46	81%
Sandler et al. 1992 ³	Case-control		1.1 (0.65–1.85)					89	217	81%
Persson et al. 1993	Case-control		1.7 (0.8–3.3)					145	305	81%
Halfvarson et al. 2006 ⁴	Case-control				0.6 (0.1–2.5)			125	125	75%
Ng et al. 2012 ⁵	Case-control	0.43 (0.11–1.66)						125	125	74%
Parrello et al. 1997 ⁶	Case-control	3.11 (1.54–6.3)						536	755	67%
Corrao et al. 1998	Case-control			1.6 (0.9–3.0)		1.3 (0.6–2.8)		594	594	67%
Calkins et al. 1986 ⁷	Case-control	0.62 (0.11–3.42)						35	32	42%
Calkins et al. 1986 ⁸	Case-control	0.57 (0.11–2.88)						35	38	42%

¹ Hazard ratios (RR adjusted for time).

² Authors' calculation, adjusted for smoking.

³ Interaction with smoking notes, higher RR in smokers (2.49).

⁴ Monozygotic and dizygotic twins.

⁵ Twins studies.

⁶ Unclear how the calculation was done.

⁷ Hospital controls.

⁸ Neighborhood controls.

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Vcev et al. 2015	Case-control	0.75 (0.30–1.88)						62	42	31%
Cornish et al. 2008	Meta-analysis				1.28 (1.06–1.54)		1.07 (0.702–1.640)	883	74,932	91%
Cornish et al. 2008 ⁹	Meta-analysis				1.24 (0.999–1.54)					91%
Godet et al. 1995 ¹⁰	Meta-analysis		1.29 (0.94–1.77)					851	49,875	82%

⁹ High quality studies.

¹⁰ Adjusted for smoking.

Ulcerative Colitis References

- Boyko EJ, Theis MK, Vaughan TL, and Nicol-Blades B. Increased risk of inflammatory bowel disease associated with oral contraceptive use. *American Journal of Epidemiology* 1994; 140:268–278.
- Calkins BM, Mendeloff AI, and Garland C. Inflammatory bowel disease in oral contraceptive users. *Gastroenterology* 1986; 91:523–524.
- Cornish JA, Tan E, Simillis C, Clark SK, Teare J, and Tekkis PP. The risk of oral contraceptives in the etiology of inflammatory bowel disease: a meta-analysis. *American Journal of Gastroenterology* 2008; 103:2394–2400.
- Corrao G, Tragnone A, Caprilli R, Trallori G, Papi C, Andreoli A, Di Paolo M, Riegler G, Rigo GP, Ferraù O, Mansi C, Ingrosso M, and Valpiani D. Risk of inflammatory bowel disease attributable to smoking, oral contraception and breastfeeding in Italy: a nationwide case-control study. Cooperative Investigators of the Italian Group for the Study of the Colon and the Rectum (GISC). *International Journal of Epidemiology* 1998; 27:397–404.
- García Rodríguez LA, González-Pérez A, Johansson S, and Wallander MA. Risk factors for inflammatory bowel disease in the general population. *Alimentary Pharmacology & Therapeutics* 2005; 22:309–315.
- Godet PG, May GR, and Sutherland LR. Meta-analysis of the role of oral contraceptive agents in inflammatory bowel disease. *Gut* 1995; 37:668–673.
- Halfvarson J, Jess T, Magnuson A, Montgomery SM, Orholm M, Tysk C, Binder V, and Järnerot G. Environmental factors in inflammatory bowel disease: a co-twin control study of a Swedish-Danish twin population. *Inflammatory Bowel Disease* 2006; 12:925–933.
- Khalili H, Higuchi LM, Ananthakrishnan AN, Richter JM, Feskanich D, Fuchs CS, and Chan AT. Oral contraceptives, reproductive factors and risk of inflammatory bowel disease. *Gut* 2013; 62:1153–1159.
- Lashner BA, Kane SV, and Hanauer SB. Lack of association between oral contraceptive use and ulcerative colitis. *Gastroenterology* 1990; 99:1032–36.
- Logan RF and Kay CR. Oral contraception, smoking and inflammatory bowel disease—findings in the Royal College of General Practitioners Oral Contraception Study. *International Journal of Epidemiology* 1989; 18:105–107.
- Ng SC, Woodrow S, Patel N, Subhani J, and Harbord M. Role of genetic and environmental factors in British twins with inflammatory bowel disease. *Inflammatory Bowel Disease* 2012; 18:725–736.
- Parrello T, Pavia M, Angelillo IF, Monteleone G, Riegler G, Papi G, D'Inca R, Annese V, Tonelli F, Caprilli R, and Pallone F. Appendectomy is an independent protective factor for ulcerative colitis: results of a multicentre case control study. The Italian Group for the Study of the Colon and Rectum (GISC). *Italian Journal of Gastroenterology and Hepatology* 1997; 29:208–211.
- Persson PG, Leijonmarck CE, Bernell O, Hellers G, and Ahlbom A. Risk indicators for inflammatory bowel disease. *International Journal of Epidemiology* 1993; 22:268–272.
- Sandler RS, Wurzelmann JI, and Lyles CM. Oral contraceptive use and the risk of inflammatory bowel disease. *Epidemiology* 1992; 3:374–378.

Vcev A, Pezerovic D, Jovanovic Z, Nakic D, Vcev I, and Majnarić L. A retrospective, case-control study on traditional environmental risk factors in inflammatory bowel disease in Vukovar-Srijem County, north-eastern Croatia, 2010. *Wiener Klinische Wochenschrift* 2015; 127:345–354.

Vessey M, Jewell D, Smith A, Yeates D, and McPherson K. Chronic inflammatory bowel disease, cigarette smoking, and use of oral contraceptives: findings in a large cohort study of women of childbearing age. *British Medical Journal (Clinical Research Edition)* 1986; 292:1101–1113.

Table 9 – Individual Studies of the Effects of COCs on the Development of Systemic Lupus Erythematosus

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Costenbader et al. 2007 ¹	Cohort		1.5 (1.1–2.1)				1.7 (1.2–2.3)	262	238,046	96%
Costenbader et al. 2007 ²	Cohort		1.6 (1.1–2.2)				1.6 (1.1–2.2)	164	102,882	96%
Costenbader et al. 2007 ³	Cohort		2.3 (1.0–5.0)				2.3 (1.1–5.2)	98	107,854	96%
Bernier et al. 2009	Cohort		1.19 (0.98–1.45)		1.54 (1.15–2.07)		1.06 (0.85–1.33)	786	7817	96%
Bernier et al. 2009 ⁴	Cohort				2.52 (1.14–5.57)			786	7817	96%
Bernier et al. 2009 ⁵	Cohort				1.45 (1.06–1.99)			786	7817	96%
Sanchez-Guerrero et al. 1997	Cohort		1.4 (0.9–2.1)					99	121,546	88%
Sanchez-Guerrero et al. 1997 ⁶	Cohort		1.9 (1.1–3.3)					58	121,587	88%
Cooper et al. 2002	Case-control			1.5 (0.8–2.7)		1.3 (0.8–2.0)		240	321	92%
Strom et al. 1994	Case-control	0.8 (0.5–1.4)						195	143	73%
Zonana-Nacach et al. 2002 ⁷	Case-control	2.1 (1.18–3.6)						130	130	61%
Grimes et al. 1985	Case-control			0.5 (0.11–2.3)				109	109	58%

¹ Pooled RR from the Nurses' Health Study (NHS) and NHS II.

² RR from the NHS (data collection through 1976).

³ RR from NHS II (data collection through 1989).

⁴ RR for short term use (starting COCs within ≤3 months).

⁵ RR for long term use (starting COCs over 3 months previously with current use ongoing).

⁶ Using most stringent definition of systemic lupus erythematosus.

⁷ Paper written in Spanish. OR is for use of oral contraceptives for more than one year.

Systemic Lupus Erythematosus References

Bernier MO, Mikaeloff Y, Hudson M, and Suissa S. Combined oral contraceptive use and the risk of systemic lupus erythematosus. *Arthritis and Rheumatism* 2009; 61:476–481.

Cooper GS, Dooley MA, Treadwell EL, St Clair EW, and Gilkeson GS. Hormonal and reproductive risk factors for development of systemic lupus erythematosus: results of a population-based, case-control study. *Arthritis and Rheumatism* 2002; 46:1830–1839.

Costenbader KH, Feskanich D, Stampfer MJ, and Karlson EW. Reproductive and menopausal factors and risk of systemic lupus erythematosus in women. *Arthritis and Rheumatism* 2007; 56:1251–1262.

Grimes DA, LeBolt SA, Grimes KR, and Wingo PA. Systemic lupus erythematosus and reproductive function: A case control study. *American Journal of Obstetrics and Gynecology* 1985; 153:179–186.

Sanchez-Guerrero J, Karlson EW, Liang MH, Hunter DJ, Speizer FE, and Colditz GA. Past use of oral contraceptives and the risk of developing systemic lupus erythematosus. *Arthritis and Rheumatism* 1997; 40: 804–808.

Strom BL, Reidenberg MM, West S, Snyder ES, Freundlich B, and Stolley PD. Shingles, allergies, family medical history, oral contraceptives, and other potential risk factors for systemic lupus erythematosus. *American Journal of Epidemiology* 1994; 140:632–642.

Zonana-Nacach A, Rodríguez-Guzmán LM, Jiménez-Balderas FJ, Camargo-Coronel A, Escobedo-de la Peña J, and Fraga A. [Risk factors associated with systemic lupus erythematosus in a Mexican population]. *Salud Pública de México* 2002; 44:213–218.

Table 10 – Studies of Chemical Contraceptives and Depression, Mood Disorders and Suicides

	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	Cases	Controls/Cohort Size
Skovlund 2016 incl /Worley	Prospective Cohort Incident Depression – COCs		1.1 ⁹² (1.08-1.14)				1,061,997
	Incident Depression – POCs		1.2 ⁹³ (1.04-1.31)				
	First use of Antidepressants – COCs		1.23 ⁹⁴ (1.22-1.25)				
	First use of Antidepressants – POCs		1.3 ⁹⁵ (1.27-1.40)				
Skovlund 2018 incl /Worley	Prospective Cohort						475,802
	Prospective Cohort Suicide attempts		1.97 ⁹⁶ (1.85-2.10)				
	Suicides		3.08 ⁹⁷ (1.34-7.08)				
Gregory 2018	NCHA survey					146,938	202,759
	Ever Diagnosed with Depression	1.558 (1.506- 1.612)					
	Academic performance affected by depression	1.282 (1.245- 1.321)					
Keyes 2013	COC reduced depression among women 25-34 years of age. ⁹⁸ 4 waves of L-Hanes			-1.04 ⁹⁹ (-1.73 - -0.35)		3224	1219
	Suicide attempts			0.38 (0.15-0.97)			
Toffol 2011	Population/choice			-0.988 ¹⁰¹ (-1.917 – -0.059)			2,310

⁹² First diagnosis of depression for combined oral contraceptive users.

⁹³ First diagnosis of depression for all progestin-only method users.

⁹⁴ First use of an antidepressant for combined oral contraceptive users.

⁹⁵ First use of an antidepressant for all progestin-only method users.

⁹⁶ Hazard ratio for suicide attempts; all hormonal contraceptives.

⁹⁷ Hazard ratio for suicides; all hormonal contraceptives.

⁹⁸ “The presence of depressive symptoms during the past 7 days was assessed in all waves using the Center for Epidemiologic Studies Depression Scale (CES-D).”

⁹⁹ β statistic shown.

¹⁰¹ β statistic shown for the Beck Depression Inventory (BDI). None of the other parameters assessed was statistically significant (including any psychiatric diagnosis, alcohol dependence, major depressive episode or disorder, dysthymic disorder, or anxiety disorder).

	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	Cases	Controls/Cohort Size
	Cross sectional 30-54 yrs. of age ¹⁰⁰						
Toffol 2012	Population-based cross-sectional study ¹⁰²			-0.42 (-1.79 - -0.04) ¹⁰³			8,586
Svendal 2012 ¹⁰⁴	Population-based cross-sectional study					40	458
	POC Use – mood disorder			3.0 (1.1-7.8)			
	COC Use – mood disorder			0.3 (0.1-0.9)			
Horibe 2018	Retrospective ¹⁰⁵					253	6,157,897
	Post-partum depression w/ levonorgestrel			12.5 (8.7-18)			
	Post-partum depression w/ etonogestrel			14.0 (8.5-22.8)			
	Post-partum depression w/ sertraline & drospirenone			5.4 (2.7-10.9)			
Singata-Madliki 2016	Single-blind randomized controlled trial of post-partum DMPA vs. copper IUD			106		111 ¹⁰⁷	117 ¹⁰⁸

¹⁰⁰ “The associations between the current use of COCs and the LNG-IUS, and their duration versus mood symptoms [Beck Depression Inventory (BDI)], psychological well-being [(General Health Questionnaire-12 (GHQ-12))] and recent psychiatric diagnoses [(Composite International Diagnostic Interview (CIDI))] were examined among women who participated in the Finnish-population-based Health 2000 study.” “Overall, hormonal contraception was well tolerated with few significant effects on psychological well-being.”

¹⁰² Data were collected in the context of the National FINRISK Study Survey, a cross-sectional population-based health survey carried out in Finland every 5 years since 1972. For the purpose of this study, data collected in the years 1997, 2002 and 2007 were analyzed for ages 25–54. OC vs. LNG. inconsistent questions between surveys, BDI, recall bias, etc. “Presence of somatic and psychological symptoms was assessed by asking the participants how often (often, sometimes, not at all) in the previous month they had had one or more out of 13 symptoms.” Also administered the Beck Depression Inventory-13. “A negative association between the current use of COCs and Beck Depression Inventory-13 (BDI-13) score was found. Some other negative associations, all characterized by a small effect size, were detected between current use of COCs and the BDI items feelings of dissatisfaction, feelings of uselessness, irritability, lost interest in people and lost appetite.”

¹⁰³ Results for the BDI-13 shown. Other parameters (including BDI-21, low mood last year, anhedonia last year, recent diagnosis of depression and recent other psychiatric diagnosis) did not reach statistical significance.

¹⁰⁴ Women in Australia 20-50 years of age. Evaluated for the occurrence of mood disorders, including major depressive disorder (MDD), minor depression, bipolar disorder, dysthymia, mood disorder due to a general medical condition and substance induced mood disorder.

¹⁰⁵ Data is from the FDA Adverse Event Reporting System (FAERS) database. Reporting Odds Ratios (ROR) are shown.

¹⁰⁶ Beck Depression Inventory (BDI-II) and the Edinburgh Postnatal Depression Scale (EPDS) evaluated. The one-month EPDS depression scores were statistically significantly higher in the DMPA arm compared with the IUD arm ($p=0.04$). Three-month BDI-II scores were significantly higher in the DMPA arm than in the IUD arm ($p=0.002$) and, according to the BDI-II but not the EPDS, more women in the DMPA arm had major depression at this time-point (8 vs 2; $p=0.05$).

¹⁰⁷ 111 randomized to DMPA.

¹⁰⁸ 117 randomized to IUDs.

	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	Cases	Controls/Cohort Size
Kulkarni 2005 ¹⁰⁹	Case-control pilot study COCs vs non-users			p=0.001 depression for all scales ¹¹⁰		26	32
Roberts 2017	Retrospective cohort study ¹¹¹			With Dx of depression ¹¹²	w/anti depressant use ¹¹³	31,506 ¹¹⁴	44,022 ¹¹⁵
	Norethindrone-only pills			0.56 (0.49-0.64)	0.58 (0.52-0.64)		
	Levonorgestrel intrauterine system			0.65 (0.52-0.82)	1.01 (0.87-1.18)		
	Etonogestrel subdermal implant			1.01 (0.83-1.22)	1.22 (1.06-1.41)		
	Ethinyl estradiol/ norgestimate (pill)			0.89 (0.70-1.14)	1.02 (0.85-1.22)		
	Ethinyl estradiol/norethindrone (pill)			0.82 (0.59-1.12)	0.88 (0.69-1.13)		
	Ethinyl estradiol/etonogestrel (ring)			1.09 (0.80-1.50)	1.45 (1.16-1.80)		
Tsai 2010	Retrospective chart review ¹¹⁶	DMPA	Controls			55	192
	Mean EPDS scores at 6 weeks postpartum	5.02	6.17				
Griksiene 2011	Case-control study ¹¹⁷	¹¹⁸				23 ¹¹⁹	20 ¹²⁰

¹⁰⁹ Assessment tools included three depression rating scales: Beck Depression Inventory (BDI), Hamilton Rating Scale for Depression (HAM-D) and Montgomery-Asberg Depression Rating Scale (MADRS); also used the Global Assessment of Functioning (GAF) Scale.

¹¹⁰ ANOVA of GAF, BDI, HAM-D & MADRS scales all significantly different.

¹¹¹ Post-partum depression with hormonal contraception.

¹¹² Adjusted hazard ratios shown.

¹¹³ Adjusted hazard ratios shown.

¹¹⁴ Number on hormonal contraceptives.

¹¹⁵ Number not on hormonal contraceptives.

¹¹⁶ Depot medroxyprogesterone in the immediate post-partum period and depression. Evaluated the Edinburgh Postnatal Depression Scale (EPDS).

¹¹⁷ Verbal fluency and mental rotation (spatial perception) are affected by progestins w/androgenic or antiandrogenic properties.

¹¹⁸ Naturally cycling women performed better on verbal fluency task as compared to OC users. Subjects who used the third generation (androgenic) COCs generated significantly fewer words as compared to new generation (anti-androgenic) OC users and non-users. The third generation OC users demonstrated significantly longer RT in MRT task as compared to non-users. The MRT, verbal fluency and mood parameters did not depend on the phase of menstrual cycle.

¹¹⁹ Women on hormonal contraception.

¹²⁰ Control women not on hormonal contraception.

Depression, Mood Disorders, and Suicide References

- Del Rio JP, Allende MI, Molina N, Serrano FG, Molina S, and Vigil P. Steroid Hormones and their Action in Women's Brains: The Importance of Hormonal Balance. *Frontiers in Public Health* 2018; May(6) art. 141:1–15.
- Gregory Sean T, Hall K, Quast T, Gatto A, Bleck J, Storch EA, and DeBate R. Hormonal contraception, depression and Academic Performance among females attending college in the United States. *Psychiatry Research* 2018; 270:111–116.
- Griksiene R, Ruksenas O. Effects of hormonal contraceptives on mental rotation and verbal fluency. *Psychoneuroendocrinology*. 2011; 36(8):1239-48.
- Horibe M, Hane Y, Abe J, Matsui T, Kato Y, Ueda N, Sasaoka S, Motooka Y, Hatahira H, Hasegawa S, Kinosada Y, Hara H, Nakamura M. Contraceptives as possible risk factors for postpartum depression: A retrospective study of the food and drug administration adverse event reporting system, 2004-2015. *Nurs Open*. 2018; 5(2):131-138.
- Keyes Katherine T, Cheslack-Postava K, Westhoff C, Heim CM, Haloossim M, Walsh K, and Koenen K. Association of Hormonal Contraception Use with Reduced levels of Depressive Symptoms: A National Study of Sexually Active women in the United States. *Am J. Epidemiol* 2013; 178(9):1378–1388.
- Kulkarni J, Liew J, Garland KA. Depression associated with combined oral contraceptives--a pilot study. *Aust Fam Physician*. 2005; 34(11):990.
- Roberts TA, Hansen S. Association of Hormonal Contraception with depression in the postpartum period. *Contraception*. 2017; 96(6):446-452.
- Singata-Madliki M, Hofmeyr GJ, Lawrie TA. The effect of depot medroxyprogesterone acetate on postnatal depression: a randomised controlled trial. *J Fam Plann Reprod Health Care*. 2016; 42(3):171-6.
- Skovlund CW, Mørch LS, Kessling LV, and Lidegaard O. Association of Hormonal Contraception with Depression. *JAMA Psychiatry* 2016; 73(11):1154–1162.
- Skovlund CW, Mørch LS, Kessling LV, Lange T, and Lidegaard, O. Association of Hormonal Contraception with Suicide Attempts and Suicides. *Am. J Psychiatry* 2018; 175(4):336–342.
- Svendal G, Berk M, Pasco JA, and Jacka FN. The use of hormonal contraceptive agents and mood disorders in women. *J Affective Disorders* 2012; 140:92–96.
- Toffol E, Heikinheimo O, Koponen P, Luoto R, Partonen T. Further evidence for lack of negative associations between hormonal contraception and mental health. *Contraception*. 2012; 86(5):470-80.
- Toffol E, Heikinheimo, Koponen P, Luoto R, and Partonen T. Hormonal contraception and mental health: results of a population based study. *Human Reproduction* 2011; 26(11):3085–3093.
- Tsai R, Schaffir J. Effect of depot medroxyprogesterone acetate on postpartum depression. *Contraception*. 2010; 82(2):174-7.

Worly Brett L, Gur TL, and Schaffir J. The relationship between progestin hormonal contraception and depression: a systematic review. *Contraception* 2018; 97:478–489.

Young EA, Kornstein SG, Harvey AT, Wisniewski SR, Barkin J, Fava M, Trivedi MH, and Rush AJ. Influences of Hormone-Based Contraception in Depressive symptoms in Premenopausal Women with Major Depression. *Psychoneuroendocrinology* 2007; 32(7):843–853.

Table 11 – Individual Studies of the Effects of COCs on the Development of Multiple Sclerosis

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Hernán et al. 2000 ¹²¹	Cohort		1.1 (0.9-1.5)		1 (0.6-1.6)		1.2 (0.9-1.5)	313	237,318	90%
Thorogood et al. 1998 ¹²²	Cohort				1.2 (0.7-2.0)		1.3 (0.9-2.0)	114	46,000	75%
Villard-Mackintosh et al. 1993	Cohort		0.8 (0.5-1.4)					63	16,969	65%
Hellwig et al. 2016	Case-control	1.51 (1.12-2.03)		1.47 1.05-2.05		1.55 (1.20-2.00)		400	3804	92%
Kotzamani et al. 2012	Case-control	1.6 (1.1-2.4)						254	314	81%
Alonso et al. 2005 ¹²³	Case-control	0.6 (0.4-1.0)		0.5 (0.3-1.2)		0.6 (0.4-1.0)		106	1001	77%

¹²¹ NHS I and II cohorts.

¹²² Funded by drug companies that make HCs.

¹²³ OC use over the 3 years prior to the index date. Limited to women ≤50 years of age.

Multiple Sclerosis References

Alonso A, Jick SS, Olek MJ, Ascherio A, Jick H, and Hernán MA. Recent use of oral contraceptives and the risk of multiple sclerosis. *Archives of Neurology* 2005; 62:1362–1365.

Hellwig K, Chen LH, Stanczyk FZ, and Langer-Gould AM. Oral Contraceptives and Multiple Sclerosis/Clinically Isolated Syndrome Susceptibility. *PLoS One* 2016; 11:e0149094. Doi:10.1371/journal.pone.0149094.

Hernán MA, Hohol MJ, Olek MJ, Spiegelman D, and Ascherio A. Oral contraceptives and the incidence of multiple sclerosis. *Neurology* 2000; 55:848–854.

Kotzamani D, Panou T, Mastorodemos V, Tzagournissakis M, Nikolakaki H, Spanaki C, and Plaitakis A. Rising incidence of multiple sclerosis in females associated with urbanization. *Neurology* 2012; 78:1728–1735.

Thorogood M, and Hannaford PC. The influence of oral contraceptives on the risk of multiple sclerosis. *British Journal of Obstetrics and Gynaecology* 1998; 105:1296–1299.

Villard-Mackintosh L, and Vessey MP. Oral contraceptives and reproductive factors in multiple sclerosis incidence. *Contraception* 1993; 47:161–168.

Table 12 – Individual Studies of the Effects of Contraceptives on the Development of Osteoporotic Fractures

Study	Study Design	Intervention	OR	RR		Cases	Controls or Cohort Size	Outcome
Cooper 1993 ¹²⁴	Cohort	COCs		1.20 (1.08-1.34)		1365	46,000	All fractures
Vessey 1998 ¹²⁵	Cohort	COCs		1.5 (1.1-2.1)		1308	17,032	First fracture: radius or ulna
Vessey 1998 ¹²⁶	Cohort	COCs		1.2 (1.1-1.4)				First fracture: all sites
Vessey 1998 ¹²⁷	Cohort	COCs		2.5 (1.5-4.0)				First fracture: radius or ulna
Vessey 1998 ¹²⁸	Cohort	COCs		1.3 (1.1-1.5)				First fracture: all sites
Vessey 1998 ¹²⁹	Cohort	COCs		5.7 (p=0.017)				First fracture: radius or ulna
Vessey 1998 ¹³⁰	Cohort	COCs		11.2 (p<0.001)				First fracture: all sites
Barad 2005 ¹³¹	Cohort	OCs ¹³²		1.07 (1.01–1.15)		4,674	80,947	First fracture
Barad 2005 ¹³³	Cohort	OCs		1.15 (1.04-1.27)		4,674	80,947	First fracture
Barad 2005 ¹³⁴	Cohort	OCs		1.09 (0.97–1.23)		4,674	80,947	First fracture
Lanza 2013 ¹³⁵	Retrospective cohort study	DMPA ¹³⁶		1.41 (1.35–1.47)		11,822	312,395	Incident fractures

¹²⁴ From the Royal College of General Practitioners (RCGP) Oral Contraception Study.

¹²⁵ OC use > 97 months vs no use. Recruited age 25 to 39 years; followed to 45 years.

¹²⁶ OC use > 97 months vs no use. Recruited age 25 to 39 years; followed to 45 years.

¹²⁷ Interval since use: 73 to 96 months vs no use (radius or ulna). Recruited age 25 to 39 years; followed to 45 years.

¹²⁸ < 12 months vs no use (all fractures). Recruited age 25 to 39 years; followed to 45 years.

¹²⁹ χ^2 trend.

¹³⁰ χ^2 trend.

¹³¹ Recruited age 50 to 74 years; OC use: any vs none.

¹³² The patients were asked about oral contraceptive use, which likely was predominantly COCs but was not broken down with regard to COCs or POCs.

¹³³ Among women without any postmenopausal hormone treatment, past OC use for 5 years or less.

¹³⁴ Among women without any postmenopausal hormone treatment, past OC use for more than 5 years.

¹³⁵ They note that, “Although DMPA users experienced more fractures than nonusers, this association may be the result of confounding by a pre-existing higher risk for fractures in women who chose DMPA for contraception.” However, this is based on analysis of relatively few fractures prior to DMPA use.

¹³⁶ Depot medroxyprogesterone acetate = DMPA.

Study	Study Design	Intervention	OR	RR		Cases	Controls or Cohort Size	Outcome
	Past use ¹³⁷	DMPA		1.32 (1.24–1.41)				Incident fractures
	Recent use ¹³⁸	DMPA		1.41 (1.31–1.50)				Incident fractures
	Current use ¹³⁹	DMPA		1.51 (1.41–1.61)				Incident fractures
Tuppurainen 1993 ¹⁴⁰	Case-control	OCs	1.21 (0.93-1.57)			629	13,100	All fractures
Tuppurainen 1993 ¹⁴¹	Case-control	OCs	1.35 (0.88-2.05)			210	13,100	Wrist fractures
O'Neill 1996	Case-control	OCs	0.3 (0.1-0.9)			62	116	Distal forearm fractures only Population controls
O'Neill 1996	Case-control	OCs	0.7 (0.2-2.4)			62	50	Distal forearm fractures only Fall controls
Michaëlsson 1999 ¹⁴²	Case-control	Any ¹⁴³	0.75 (0.59–0.96)			1327	3312	Hip fractures
Vestergaard 2006 ¹⁴⁴	Case-control	OCs	<0.3 DDD/day	0.3–0.99 DDD/day	1+ DDD/day	64,548	193,641	Any fracture in the year 2000
	<25 years ¹⁴⁵	OCs	0.97 (0.91–1.03)	0.96 (0.92–1.01)	0.92 (0.86–0.98)			Any fracture in the year 2000
	25-49 years	OCs	0.91 (0.82–1.00)	0.90 (0.77–1.05)	0.87 (0.64–1.18)			Any fracture in the year 2000
	50+ years	OCs	0.92 (0.77–1.10)	0.69 (0.45–1.05)	0.62 (0.27–1.41)			Any fracture in the year 2000
Vestergaard 2008a ¹⁴⁶	Case-control	OCs	<0.3 DDD/day	0.3–0.99 DDD/day	1+ DDD/day	64,548	193,641	Any fracture in the year 2000
	<15	OCs	1.02 (0.75–1.37)	1.17 (1.01–1.37)	0.97 (0.85–1.11)			Any fracture in the year 2000

¹³⁷ Active DMPA use based on the interleaving of active 90-day exposures generated by each injection.

¹³⁸ Recent exposure is 640 or fewer days after the last active exposure.

¹³⁹ Past exposure begins after “recent” exposure (641 or more days after the last active exposure).

¹⁴⁰ Oral contraceptive use for 6+ years.

¹⁴¹ Oral contraceptive use for 6+ years.

¹⁴² No significant correlation was seen with duration of use, time since last use or time between last use and menopause.

¹⁴³ Any type of chemical contraceptive was evaluated, not separated as COCs or POCs.

¹⁴⁴ “The exposure time for oral contraceptives may thus maximally have spanned 5 years (from January 1, 1996, to December 31, 2000).” This and the other Vestergaard study are not useful as they do not take into account remote use or cumulative lifetime use. ORs shown.

¹⁴⁵ Defined daily dosages = DDD.

¹⁴⁶ Similar to Vestergaard 2006; only looked at use within the past 5 years. A younger group examined here. ORs shown.

Study	Study Design	Intervention	OR	RR		Cases	Controls or Cohort Size	Outcome
	15.1-17	OCs	1.22 (1.02–1.47)	1.14 (1.00–1.30)	1.04 (0.90–1.19)			Any fracture in the year 2000
	17.1-19	OCs	0.97 (0.87–1.09)	0.93 (0.84–1.02)	1.02 (0.89–1.18)			Any fracture in the year 2000
	>19	OCs	0.99 (0.93–1.05)	1.00 (0.93–1.08)	0.88 (0.78–0.99)			Any fracture in the year 2000
Vestergaard 2008b ¹⁴⁷	Case-control	DMPA	1.44 (1.01–2.06)			64,548	193,641	Any fracture in the year 2000 DMPA use
Wei 2011 ¹⁴⁸	Cross-sectional		<5 years of use	5-10 years of use	>10 years of use		491	
		OCs	0.85 (0.45–1.58)	0.45 (0.21–0.93)	0.75 (0.36–1.54)			Presence of vertebral deformity
		OCs	0.96 (0.62–1.48)	0.63 (0.37–1.07)	0.94 (0.56–1.56)			Number of vertebral deformities
Meier 2010 ¹⁴⁹	Case-control		Current Use	Past Use		17,527	70,130	Incident fracture
	1-2 DMPA Scripts	DMPA	1.18 (0.93–1.49)	1.17 (1.07–1.29)				Incident fracture
	3-9 DMPA scripts	DMPA	1.36 (1.15–1.60)	1.23 (1.11–1.36)				Incident fracture
	10+ DMPA scripts	DMPA	1.54 (1.33–1.78)	1.30 (1.09–1.55)				Incident fracture
	1-2 COC Scripts	COCs	1.01 (0.87–1.18)	1.00 (0.95–1.07)				Incident fracture
	3-9 COC scripts	COCs	1.01 (0.94–1.09)	0.99 (0.94–1.04)				Incident fracture
	10+ COC scripts	COCs	1.09 (1.03–1.16)	1.03 (0.97–1.10)				Incident fracture
Memon 2011 ¹⁵⁰	Case-control	COCs	1.05 (0.86–1.29)			651	1302	Any fracture
Kyvernitis 2017 ¹⁵¹	Case-control		OR Current Use	OR Past Use		4189	4189	First-time fracture diagnosis
	1-2 DMPA scripts	DMPA	0.97 (0.51–1.86)	0.96 (0.73–1.26)				
	3-9 DMPA scripts	DMPA	2.41 (1.42–4.08)	1.14 (0.86–1.51)				
	10+ DMPA scripts	DMPA	1.46 (0.96–2.23)	1.55 (1.07–2.27)				

¹⁴⁷ Similar to Vestergaard 2006; only looked at use within the past 5 years. DMPA examined here. ORs shown.

¹⁴⁸ Small cross-sectional study. ORs shown.

¹⁴⁹ Females aged 20–44 years with an incident fracture diagnosis between 1995 and 2008.

¹⁵⁰ Nested case-control study of the Cooper study from the Royal College of General Practitioners (RCGP) Oral Contraception Study. Last OC use > 10 years vs never.

¹⁵¹ Women between 20 and 44 years of age with a first-time fracture diagnosis, matched with random controls using the Disease Analyzer database.

Study	Study Design	Intervention	OR	RR		Cases	Controls or Cohort Size	Outcome
	1-2 COC scripts	COCs	0.98 (0.73–1.31)	0.90 (0.77–1.05)				
	3-9 COC scripts	COCs	1.39 (1.12–1.73)	0.90 (0.78–1.03)				
	10+ COC scripts	COCs	1.07 (0.88–1.30)	1.04 (0.90–1.21)				

Osteoporotic Fracture References

- Barad D, Kooperberg C, Wactawski-Wende J, Liu J, Hendrix SL, and Watts NB. Prior oral contraception and postmenopausal fracture: a Womens' Health Initiative observational cohort study. *Fertility and Sterility* 2005; 84:374–383.
- Cooper C, Hannaford P, Croft P, and Kay CR. Oral contraceptive pill use and fractures in women: a prospective study. *Bone* 1993; 14(1):41–45.
- Curtis KM and Martins SL. Progestin-only contraception and bone mineral density: a systematic review. *Contraception* 2006; 73:470–487.
- FDA 2004: U.S. Food and Drug Administration. FDA Talk Paper. Black box Warning Added Concerning Long-term Use of Depo-Provera Contraceptive Injection. <http://web.archive.org/web/20070809090332/http://www.fda.gov/bbs/topics/ANSWERS/2004/ANS01325.html> (accessed 2019 February 23).
- Kyvernitakis I, Kostev K, Nassour T, Thomasius F, Hadji P. The impact of depot medroxyprogesterone acetate on fracture risk: a case-control study from the UK. *Osteoporos Int*. 2017; 28(1):291-297.
- Lanza LL, McQuay LJ, Rothman KJ, Bone HG, Kaunitz AM, Harel Z, Ataher Q, Ross D, Arena PL, and Wolter KD. Use of depot medroxyprogesterone acetate contraception and incidence of bone fracture. *Obstetrics and Gynecology* 2013; 121(3):593–600.
- Lopez LM, Chen M, Mullins Long S, Curtis KM, and Helmerhorst FM. Steroidal contraceptives and bone fractures in women: evidence from observational studies (Review). *Cochrane Database of Systematic Reviews* 2015; Issue 7. Art. No.: CD009849. DOI: 10.1002/14651858.CD009849.pub3.
- Meier C, Brauchli YB, Jick SS, Kraenzlin ME, and Meier CR. Use of depot medroxyprogesterone acetate and fracture risk. *Journal of Clinical Endocrinology and Metabolism* 2010; 95(11):4909–4916.
- Memon S, Iversen L, and Hannaford PC. Is the oral contraceptive pill associated with fracture in later life? New Evidence from the Royal College of General Practitioners Oral Contraception Study. *Contraception* 2011; 84(1):40–47.
- Michaëlsson K, Baron JA, Farahmand BY, and Ljunghall S. Influence of parity and lactation on hip fracture risk. *American Journal of Epidemiology* 2001; 153(12):1166–1172.
- Michaëlsson K, Baron JA, Farahmand BY, Persson I, and Ljunghall S. Oral-contraceptive use and risk of hip fracture: a case-control study. *Lancet* 1999; 353(9163):1481–1484.
- O'Neill TW, Marsden D, Adams JE, and Silman AJ. Risk factors, falls, and fracture of the distal forearm in Manchester, UK. *Journal of Epidemiology and Community Health* 1996; 50(3):288–292.
- Parazzini G, Tavani A, Ricci E, and La Vecchia C. Menstrual and reproductive factors and hip fractures in post menopausal women. *Maturitas* 1996; 24:191–196.

Sirola J, Rikkinen T, Tuppurainen M, Honkanen R, and Kroger H. Should risk of bone fragility restrict weight control for other health reasons in postmenopausal women?—A ten year prospective study. *Maturitas* 2012; 71(2):162–168.

Tuppurainen M, Honkanen R, Kröger H, Saarikoski S, and Alhava E. Osteoporosis risk factors, gynaecological history and fractures in perimenopausal women—the results of the baseline postal enquiry of the Kuopio Osteoporosis Risk Factor and Prevention Study. *Maturitas* 1993; 17(2):89–100.

Vessey M, Mant J, and Painter R. Oral contraception and other factors in relation to hospital referral for fracture. Findings in a large cohort study. *Contraception* 1998; 57(4):231–235.

Vestergaard P, Rejnmark L, and Mosekilde L. Fracture risk in very young women using combined oral contraceptives. *Contraception* 2008a; 78(5):358–364.

Vestergaard P, Rejnmark L, and Mosekilde L. Oral contraceptive use and risk of fractures. *Contraception* 2006; 73(6):571–576.

Vestergaard P, Rejnmark L, and Mosekilde L. The effects of depot medroxyprogesterone acetate and intrauterine device use on fracture risk in Danish women. *Contraception* 2008b; 78(6):459–464.

Wei S, Venn A, Ding C, Foley S, Laslett L, and Jones G. The association between oral contraceptive use, bone mineral density and fractures in women aged 50-80 years. *Contraception* 2011; 84(4):357–362.

Women's Health Initiative Study Group. Design of the Women's Health Initiative clinical trial and observational study. *Controlled Clinical Trials* 1998; 19(1):61–109.

Table 13 – Effect of Chemical Contraceptives on Weight Gain

Study	Design	Comparison	N	Time	Weight change (Kg)	Fat mass change	Lean mass change	Comments
Pantoja 2010	Retrospec.	DMPA 150 vs CuIUC	758	1yr	1.76 vs -0.42*			Largest differences noted in normal and overweight BMI subgroups, minimal differences in obese BMI subgroup
				2yr	3.1 vs 0.4*			
				3yr	3.9 vs 0.8*			
Modesto 2015	Retrospec.	DMPA150 vs CuIUC	1277	1yr	1.3 vs 0.2*			Adjusted for years of school & # children. 20% loss @4yrs 84% @ 10yr.
				4yr	3.5 vs 1.9*			
				10yr	6.6 vs 4.9*			
Taneepanichskul 1998	Retrospec.	DMPA 150 vs CuIUC	100	10yr	10.9 vs 11.2			Included women 37-50 years (no younger women)
Vickery 2013	Prospec.	DMPA 150 vs CuIUC	167	1yr	2.2 vs 0.16			CHOICE study subgroup
Dal'Ava 2014	Prospec.	DMPA 150 vs CuIUC	110	1yr	1.9 vs 1.1	1.6 vs -0.9 (Kg)	0.3 vs 1.2 (kg)	Paired by age (+/-2yr) & weight (+/-2kg)
Dos Santos 2014	Prospec.	DMPA 150 vs CuIUC	71	1yr	1.4 vs 0.3	1.57 vs 0.52 (kg)	(0.31) vs (0.26) (kg)	Matched by age & BMI
								()= negative value
Studies comparing LNG IUC to non-hormonal contraceptive								
Study	Design	Comparison	N	Time	Weight change (Kg)	Total body fat	Lean body mass	
Dal'Ava 2012	Prospec.	LNG-IUC vs non-hormonal IUC	76	1yr	2.9 vs 1.4	2.5% vs -1.3%*	(1.4%) vs 1.0%*	Paired by age & BMI
Napolitano 2015	Prospec.	LNG IUC vs no method	60	1yr	0.6 vs (0.2)	1.1% vs (0.5%)*	(1.1%) vs 0.5*	
Vickery 2013	Prospec.	LNG-IUC vs Cu IUC	230	1yr	1.03 vs 0.16	nd	nd	

Modesto 2015	Retrospec.	LNG-IUC vs CulUC	1204	1yr	0.7 vs 0.2	nd	nd	
				4yr	2.7 vs 1.9			
				10yr	4.0 vs 4.9			
Studies comparing progestin-only COCs to non-hormonal								
Study	Design	Comparison	N	Time	Weight change (Kg)	Total body fat	Lean body mass	
Napolitano 2015	Prospec.	Desogestrel 75ug vs no hormonal	68	1yr	0.3 vs -0.2	1.1% vs -0.5%*	(2.8%) vs 0.5%*	
Studies comparing combined COCs to non-hormonal								
None found-								
Abstract from 2014 Cochrane review of combined oral contraceptives on weight gain:								
<p>"We found 49 trials that met our inclusion criteria. The trials included 85 weight change comparisons for 52 distinct contraceptive pairs (or placebos). <i>The four trials with a placebo or no intervention group did not find evidence supporting a causal association</i> between combination oral contraceptives or a combination skin patch and weight change. Most comparisons of different combination contraceptives showed no substantial difference in weight. In addition, discontinuation of combination contraceptives because of weight change did not differ between groups where this was studied.</p> <p>Gallo MF, Lopez LM, Grimes DA, Carayon F, Schulz KF, Helmerhorst FM. Combination contraceptives: effects on weight. Cochrane Database of Systematic Reviews 2014, Issue 1. Art. No.: CD003987.</p>								

* Significant difference (p<0.05).

Impact of Contraceptives on Body Mass References

Dal'Ava N, Bahamondes L, Bahamondes MV, Bottura BF, and Monteiro I. Body weight and body composition of depot medroxyprogesterone acetate users. *Contraception* 2014; 90(2):182–187.

Dal'Ava N, Bahamondes L, Bahamondes MV, de Oliveira Santos A, and Monteiro I. Body weight and composition in users of levonorgestrel-releasing intrauterine system. *Contraception* 2012; 86(4):350–353.

Dos Santos, PdeNS, Modesto WO, Dal'Ava N, Bahamondes M, Pavin E, and Fernandes A. Body composition and weight gain in new users of the three-monthly injectable contraceptive, depot-medroxyprogesterone acetate, after 12 months of follow-up. *European Journal of Contraception and Reproductive Health Care* 2014; 19(6):432–438.

Gallo MF, Lopez LM, Grimes DA, Carayon F, Schulz KF, and Helmerhorst FM. Combination contraceptives: effects on weight. *Cochrane Database of Systematic Reviews* 2014; Issue 1. Art. No.: CD003987.

Lopez LM, Ramesh S, Chen M, Edelman A, Otterness C, Trussell J, and Helmerhorst FM. Progestin-only contraceptives: effects on weight (Review). *Cochrane Database of Systematic Reviews* 2016; Issue 8. Art. No.: CD008815.

Lopez LM, Ramesh S, Chen M, Edelman A, Otterness C, Trussell J, and Helmerhorst FM. Progestin-only contraceptives: effects on weight. *Cochrane Database Syst Reviews* 2016a Aug 28;(8):CD008815. Doi: 10.1002/14651858.

Modesto W, dos Santos PdeNS, Correia VM, Borges L, and Bahamondes L. Weight variation in users of depot-medroxyprogesterone acetate, the levonorgestrel-releasing intrauterine system and a copper intrauterine device for up to ten years of use. *European Journal of Contraception and Reproductive Health Care* 2015; 20(1):57–63.

Napolitano A, Zanin R, Palma F, Romani C, Grandi G, Di Carlo C, and Cagnacci A. Body composition and resting metabolic rate of perimenopausal women using continuous progestogen contraception. *European Journal of Contraception and Reproductive Health Care* 2015 Aug 25 [Epub ahead of print]:1–8.

Pantoja M, Medeiros T, Baccarin MC, Morais SS, Bahamondes L, and Fernandes AM. Variations in body mass index of users of depot-medroxyprogesterone acetate as a contraceptive. *Contraception* 2010; 81(2):107–111. Taneepanichskul S, Reinprayoon D, Khaosaad P. Comparative study of weight change between long-term DMPA and IUD acceptors. *Contraception* 1998; 58(3):149–151.

Vickery Z, Madden T, Zhao Q, Secura GM, Allsworth JE, and Peipert JF. Weight change at 12 months in users of three progestin-only contraceptive methods. *Contraception* 2013; 88(4):503–508.

Table 14 – Relative Risk of Venous Thromboembolism in Current Users of Different Combined Hormonal Contraceptives as Compared with Nonusers Unless Otherwise Specified

Study	Data Sampling Period	VTE (number)	COCs with levonorgestrel RR (95% CI)	COCs with desogestrel/gestodene RR (95% CI)	COCs with drospirenone RR (95% CI)
Blomenkamp 1995	1988 - 1992	126	3.8 (1.7 - 8.4)	8.7 (3.9 - 19.3)	-
WHO 1995a, 1995b	1989 - 1993	433	3.6 (2.5 - 5.1)	7.4 (4.2 - 12.9)	-
Jick 1995	1991 - 1994	80	1 (reference)	1.8 (1.0 - 3.2)	-
Spitzer 1996	1991 - 1995	471	3.7 (2.2 - 6.2)	6.7 (3.4 - 13.0)	-
Lewis 1999	1993 - 1995	502	2.9 (1.9 - 4.2)	2.3 (1.5 - 3.5)	-
Farmer 1997	1991 - 1995	85	3.1‡ (2.1 - 4.5)	5.0‡ (3.7 - 6.5)	-
Todd 1999	1992 - 1997	99	1 (reference)	1.4 (0.7 - 2.8)	-
Bloemenkamp 1999	1994 - 1998	185	3.7 (1.9 - 7.2)	5.6 (not given)	-
Parkin 2000	1990 - 1998	26	5.1 (1.2 - 21.4)	14.9 (3.5 - 64.3)	-
Lidegaard 2002	1994 - 1998	987	2.9 (2.2 - 3.8)	4.0 (3.2 - 4.9)	-
Dinger 2007	2000 - 2004	118	1 (reference)	1.3 (NA)	1.0 (0.6 - 1.8)
Vlieg 2009	1999 - 2004	1524	3.6 (2.9 - 4.6)	7.3 (5.3 - 10.0)/5.6 (3.7 - 8.4)	6.3 (2.9 - 13.7)
Lidegaard 2009	1995 - 2005	4213	2.0 (1.8 - 2.3)	3.6 (3.3 - 3.8)	4.0 (3.3 - 4.9)
Dinger 2010	2002 - 2008	680	1 (reference)	NA	1.0 (0.6 - 1.8)
Parkin 2011	2002 - 2009	61	1 (reference)	NA	2.7 (1.5 - 4.7)
Jick 2011	2002 - 2008	186	1 (reference)	NA	2.8 (2.1 - 3.8)
Lidegaard 2011	2001 - 2009	4246	2.2 (1.7 - 2.8)	4.2 (3.6 - 4.9)	4.5 (3.9 - 5.1)
Confirmed only	2001 - 2009	2707	2.9 (2.2 - 3.8)	6.8 (5.7 - 8.1)	6.3 (5.4 - 7.5)
FDA Kaiser 2011	2001 - 2007	625	1 (reference)	NA	1.5 (1.2 - 1.9)
Gronich 2011	2002 - 2008	518	1 (reference)	1.4 (0.9 - 2.1)	1.7 (1.0 - 2.7)
Lidegaard 2012	2001 - 2010	5287	3.2 (2.7 - 3.8)	6.5 (4.7 - 8.9)*	NA
Dinger 2014	2005 - 2010	162	1 (reference)	NA	0.8 (0.5 - 1.6)

‡ Absolute risk per 10,000 years.

* Vaginal ring with the third-generation progestin etonogestrel.

Venous Thromboembolism References

Bloemenkamp KWM, Rosendaal FR, Büller HR, Helmerhorst FM, Colly LP, and Vandenbroucke JP. Risk of venous thrombosis with use of current low-dose oral contraceptives is not explained by diagnostic suspicion and referral bias. *Arch Intern Med* 1999; 159:65–70.

Bloemenkamp KWM, Rosendaal FR, Helmerhorst FM, Büller HR, and Vandenbroucke JP. Enhancement by factor V Leiden mutation of risk of deep-vein thrombosis associated with oral contraceptives containing a third- generation progestagen. *Lancet* 1995; 346:1593–1596.

Dinger J, Assmann A, M€ohner S, and Minh TD. Risk of venous thromboembolism and the use of dienogest- and drospirenone-containing oral contraceptives: results from a German case-control study. *J Fam Plann Reprod Health Care* 2010; 36:123–129.

Dinger J, Bardenheuer K, and Heinemann K. Cardiovascular and general safety of a 24-day regimen of drospirenonecontaining combined oral contraceptives: final results from the International Active surveillance Study of Women Taking Oral Contraceptives. *Contraception* 2014; 89(4):253–263.

Dinger JC, Heinemann LAJ, and Ku“hl-Habich D. The safety of a drospirenone-containing oral contraceptive: final results from the European Active Surveillance study on oral contraceptives based on 142,475 women years of observation. *Contraception* 2007; 75:344–354.

Farmer RDT, Lawrenson RA, Thompson CR, Kennedy JG, and Hambleton IR. Population-based study of risk of venous thromboembolism associated with various oral contraceptives. *Lancet* 1997; 349:83–88.

Food and Drug Administration, Office of surveillance and epidemiology. Combined hormonal contraceptives (CHCs) and the risk of cardiovascular disease endpoints. FDA. 2011; <http://www.fda.gov/downloads/Drugs/DrugSafety/UCM277384.pdf>.

Gomer K. Women, Birth Control Pills, and Thrombophilia: An Analysis of Risk Communication Kerry Gomer, Clemson University. 2009; https://tigerprints.clemson.edu/cgi/viewcontent.cgi?referer=&httpsredir=1&article=1573&context=all_theses.

Gronich N, Lavi I, and Rennert G. Higher risk of venous thrombosis associated with drospirenone-containing oral contraceptives: a population-based cohort study. *CMAJ* 2011; 183(18):E1319-25.

Jick H, Jick SS, Gurewich V, Myers MW, and Vasilakis C. Risk of ideopathic cardiovascular death and nonfatal venous thromboembolism in women using oral contraceptives with differing progestagen components. *Lancet* 1995; 346:1589-1593.

Jick SS, and Hernandez RK. Risk of non-fatal venous thromboembolism in women using oral contraceptives containing drospirenone compared with women using oral contraceptives containing levonorgestrel: case-control study using United States claims data. *British Medical Journal* 2011; 340:d2151.

Keenan L, Kerr T, Duane M, and Van Gundy K. Systematic Review of Hormonal Contraception and Risk of Venous Thrombosis. *The Linacre Quarterly* 2018; 85(4):470–477.

Le Moigne E, Delluc A, Tromeur C, Nowak E, Mottier D, Lacut K, and Le Gal G. Risk of recurrent venous thromboembolism among young women after a first event while exposed to combined oral contraception versus not exposed to: a cohort study. *Thromb Res* 2013; 132:51–55.

Lewis MA, MacRae KD, Kúhl-Habich D, Bruppacher R, Heinemann LA, and Spitzer WO. The differential risk of oral contraceptives: the impact of full exposure history. *Hum Re prod* 1999; 14:1493–1499.

Lidegaard Ø, Edström B, and Kreiner S. Oral contraceptives and venous thromboembolism. A five-year national case-control study. *Contraception* 2002; 65:187–196.

Lidegaard Ø, Løkkegaard E, Svendsen AL, and Agger C. Hormonal contraception and risk of venous thromboembolism: national follow-up study. *British Medical Journal* 2009; 339:b2890.

Lidegaard Ø, Nielsen LH, Skovlund CW, and Løkkegaard E. Venous thrombosis in users of non-oral hormonal contraception: follow-up study, Denmark 2001–10. *British Medical Journal* 2012; 344:e2990.

Lidegaard Ø, Nielsen LH, Skovlund CW, Skjeldestad FE, and Løkkegaard E. Risk of venous thromboembolism from use of oral contraceptives containing different progestogens and estrogen doses: Danish cohort study 2001–2009. *British Medical Journal* 2011; 343:d6423.

Lidegaard Ø. Hormonal contraception, thrombosis and age. *Expert Opin Drug Saf* 2014; 13:1353–360.

Parkin L, Sharples K, Hernandez RK, and Jick SS. Risk of venous thromboembolism in users of oral contraceptives containing drospirenone or levonorgestrel: nested case-control study based on UK General Practice Research Database. *British Medical Journal* 2011; 340:d2139.

Parkin L, Skegg DCG, Wilson M, Herbison GP, and Paul C. Oral contraceptives and fatal pulmonary embolism. *Lancet* 2000; 355:2133–2134.

Peck R and Norris CW. Significant Risks of Oral Contraceptives (OCPs) *Linacre Q* 2012; 79(1): 41–56. Published online 2012 Feb 1. Doi: 10.1179/002436312803571447.

Phillippe HM, Hornsby LB, Treadway S, Armstrong EM, and Bellone JM. Inherited thrombophilia. *J Pharm Pract* 2014; 27:227–233.

Spitzer WO, Lewis MA, Heinemann LAJ, Thorogood M, and MacRae KD. Third generation oral contraceptives and risk of venous thromboembolic disorders: an international case-control study. *British Medical Journal* 1996; 312:83-88.

Todd J-C, Lawrenson R, Farmer RDT, Williams TJ, and Leydon GM. Venous thromboembolic disease and combined oral contraceptives: a re-analysis of the MediPlus database. *Hum Reprod* 1999; 14:1500–1505.

Van Hylckama VA, Helmerhorst FM, Vandenbroucke JP, Doggen CJ, and Rosendaal FR. The Venous Thrombotic Risk of Oral Contraceptives, Effects of Estrogen Dose and Progestogen Type: Results of the MEGA Case-Control Study. *British Medical Journal* 2009; 339:b2921. Doi:10.1136/bmj.b2921.

Vinogradova Y, Coupland C and Hippisley-Cox J. Use of combined oral contraceptives and risk of venous thromboembolism: nested case-control studies using the QResearch and CPRD databases. *British Medical Journal* 2015; 350:h2135. Doi: <https://doi.org/10.1136/bmj.h2135> (Published 26 May 2015).

World Health Organisation Collaborative Study on Cardiovascular Disease and Steroid Hormone Contraception. Effect of different progestogens in low estrogen oral contraceptives on venous thromboembolic disease. *Lancet* 1995; 346:1582–1588.

World Health Organization Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. Venous thromboembolic disease and combined oral contraceptives: results of international multicentre case-control study. *Lancet* 1995; 346:1575-1582.

Yasmin Prescribing Information: Highlights of Yasmin Prescription Information, https://www.accessdata.fda.gov/drugsatfda_docs/label/2012/021098s019lbl.pdf.

Table 15 – Relative Risk of Thrombotic Stroke and Myocardial Infarction among Users of Selected Types of Combined Oral Contraception with Ethinyl Estradiol at a Dose of 30 to 40 µg, as Compared with Nonusers, According to Duration of Use (from Lidegaard 2012).

Duration of use	No. of person-yrs.	Thrombotic Stroke		Myocardial infarction	
		No. of events	Relative Risk (95% CI)	No. of events	Relative Risk (95% CI)
<1 year	987,564	213	1.90 (1.64–2.20)	86	1.85 (1.48–2.31)
1-4 years	992,825	194	1.55 (1.33–1.80)	108	1.99 (1.63–2.43)
>4 years	399,461	173	1.93 (1.65–2.26)	91	2.11 (1.70–2.62)

Atherosclerosis and Cardiovascular Events References

Bagdade JD and Subbaiah PV. Serum from Oral Contraceptive Users Stimulates Growth of Arterial Smooth Muscle Cells. *AHA Journals, Arteriosclerosis* 1982; 2(2):170–176.

Beral V. Cardiovascular disease mortality trends and oral contraceptive use in young women. *Lancet* 1976; 2:1047–1052.

Fallah S, Nouroozi V, Seifi M, Samadikuchaksaraei A, and Aghdashi EM. Influence of oral contraceptive pills on homocysteine and nitric oxide levels: as risk factors for cardiovascular disease. *J Clin Lab Anal* 2012; 26:120–123.

Hennekens CH, and MacMahon B. Oral contraceptive and myocardial infarction. *N Engl J Med* 1977; 296:1166–1167.

Hickson SS, Miles KL, McDonnell BJ, Yasmin, Cockcroft JR, Wilkinson IB, McEniery CM; ENIGMA Study Investigators. Use of the oral contraceptive pill is associated with increased large artery stiffness in young women: the ENIGMA study. *J Hypertens* 2011; 29(6):1155–1159.

Heidarzadeh Z, Asadi B, Saadatnia M, Ghorbani A, and Fatehi F. The Effect of Low-dose Combined Oral Contraceptive Pills on Brachial Artery Endothelial Function and Common Carotid Artery Intima–Media Thickness. *Journal of Stroke and Cerebrovascular Diseases* 2014; 23:675–680.

Lidegaard Ø, Løkkegaard E, Jensen A, Skovlund CW, and Keiding N. Thrombotic stroke and myocardial infarction with hormonal contraception. *N Engl J Med* 2012; 366:2257–2266.

Priest SE, Shenouda N, and MacDonald MJ. Effect of sex, menstrual cycle phase, and monophasic oral contraceptive pill use on local and central arterial stiffness in young adults. *Am J Physiol Heart Circ Physiol* 2018; 315(2):H357–H365.

Norouzi V, Seifi M, Fallah S, Korani M, and Samadikuchaksaraei A. Effect of oral contraceptive therapy on homocysteine and C-reactive protein levels in women: an observational study. *Anadolu Kardiyol Derg* 2011; 11:698–702.

Rietzschel E, De Buyzere M, De Baquer D, Bekaert S, Segers P, Cassiman P, Verdonck P, De Backer G, Gillebert T and ASKLEPIOS Investigators. Anticonceptive Drug Use And Increased Carotid and Femoral Plaque Prevalence: Population Data From Asklepios. *Circulation*. 2007; 116:820

Rietzschel E, De Buyzere M, De Baquer D, Langlois M, Bekaert S, Segers P, Van Damme P, Verdonck P, De Backer G, Gillebert T and ASKLEPIOS Investigators. Oral contraceptives cause major C-reactive protein rises in the female general population. *Circulation* 2018; 116:800–801.

Rietzschel E, De Buyzere M, Segers P, Bekaert S, De Baquer D, De Backer G, Gillebert T. Long Term Oral Contraceptive Use is an Independent Risk Factor for Arterial Stiffening. *Circulation*. 2008; 118:S_803–S_804

Vessey M, Doll R, Peto R, Johnson B, and Wiggins P. A longterm follow-up study of women using different methods of contraception—an Interim report. *J Biosoc Sci* 1976; 8:373–427.

Yu A, Giannone T, Scheffler P, Doonan RJ, Egiziano G, Gomez YH, Papaioannou TG, and Daskalopoulou SS. The effect of oral contraceptive pills and the natural menstrual cycle on arterial stiffness and hemodynamic (CYCLIC). *J Hypertens* 2014; 32(1):100–107.

Zoler ML. Oral Contraceptive Use Linked to Arterial Plaque. *Internal Medicine News* October 15, 2007; 32.