

Electronic Supplementary Material

Table S1. Baseline patient characteristics (n = 1608)

Characteristics	Total population (N=1608)
Mean age, yr (SD)	46.3 (17.3)
Female, no (%)	1036 (64)
Disease severity rating, no (%)	
- Mild	857 (53)
- Moderate	722 (45)
- Severe	29 (2)
Comorbidities, no (%)*	
- Cardiovascular disease	269 (17)
- Diabetes	106 (7)
- Chronic respiratory condition	159 (10)
- Hepatic, hematologic, neurologic or neurodevelopmental condition	51 (3)
- Obesity	221 (14)
Country, no (%)	
- Belgium	54 (3)
- Germany	146 (9)
- Spain	48 (3)
- France	7 (<1)
- Georgia	437 (27)
- Greece	248 (15)
- Hungary	205 (13)
- Ireland	134 (8)
- Poland	147 (9)
- United-Kingdom	182 (11)
Number of days with CA-ARTI	
- Before inclusion in the trial, mean (SE)	5.97 (0.10)
- After inclusion in the trial, mean (SE)	5.55 (0.13)

Abbreviations: no = number; SD = Standard Deviation; yr = years. *Patients could have multiple comorbidities.




Figure S1. EQ-5D-5L dimension differences between “day inclusion” and “day return” (n = 1608).

Sub-group analyses (based on the Dutch value set)

Table S2. CA-ARTI disutility by comorbidity - cardiovascular disease

Cardiovascular disease	Mean disutility [95% confidence interval]	Standard Error	Number of patients
Yes	-0.203 [-0.227, -0.179]	0.012	269
No	-0.198 [-0.207, -0.188]	0.005	1339

Table S3. CA-ARTI disutility by comorbidity - diabetes

Diabetes	Mean disutility [95% confidence interval]	Standard Error	Number of patients
Yes	-0.196 [-0.237, -0.156]	0.021	106
No	-0.199 [-0.208, -0.190]	0.005	1502

Table S4. CA-ARTI disutility by comorbidity - chronic respiratory condition

Chronic respiratory condition	Mean disutility [95% confidence interval]	Standard Error	Number of patients
Yes	-0.172 [-0.195, -0.149]	0.012	159
No	-0.202 [-0.211, -0.192]	0.005	1449

Table S5. CA-ARTI disutility by comorbidity - hepatic, hematologic, neurologic or neurodevelopmental condition

Hepatic, hematologic, neurologic or neurodevelopmental condition	Mean disutility [95% confidence interval]	Standard Error	Number of patients
Yes	-0.208 [-0.270, -0.147]	0.032	51
No	-0.198 [-0.207, -0.189]	0.005	1557

Table S6. CA-ARTI disutility by comorbidity – obesity

Obesity	Mean disutility [95% confidence interval]	Standard Error	Number of patients
Yes	-0.205 [-0.229, -0.181]	0.012	221
No	-0.198 [-0.207, -0.188]	0.005	1387

Table S7. CA-ARTI disutility by age base on Dutch value set.

Age group	Mean disutility [95% confidence interval]	Standard error	Number of patients
15-19	0.260 [-0.307, -0.213]	0.024	102
20-24	0.214 [-0.250, -0.178]	0.018	118
25-29	0.201 [-0.235, -0.166]	0.018	133
30-34	0.215 [-0.242, -0.188]	0.014	148
35-39	0.214 [-0.244, -0.183]	0.016	142
40-44	0.198 [-0.226, -0.171]	0.014	168
45-49	0.180 [-0.207, -0.153]	0.014	143
50-54	0.205 [-0.231, -0.178]	0.013	136

55-59	0.220 [-0.253, -0.186]	0.017	133
60-64	0.155 [-0.184, -0.126]	0.015	125
65-69	0.159 [-0.188, -0.130]	0.015	110
70-74	0.155 [-0.193, -0.118]	0.019	76
75-79	0.178 [-0.213, -0.143]	0.018	46
80-84	0.163 [-0.217, -0.108]	0.028	24
85-90	0.448 [-0.768, -0.128]	0.163	4