

a
Full
Mean ρ 0.257
Sequence
Mean ρ 0.269
Frequency
Mean ρ 0.111
Paired Δρ
CPEB4
FMR1
FUS
FXR1
FXR2
HNRNPA1
HNRNPC
HNRNPL
IGF2BP2
IGF2BP3
KHDRBS1
MATR3
PABPC4
PABPN1
PCBP1
PCBP2
QKI
RBM5
SFPQ
SRSF1
SRSF7
SRSF9
TARDBP
TIA1
U2AF2
RNAcompete Spearman ρ
Full − Frequency
Mean paired Δρ
+0.146
95% CI
[0.038, 0.248]
b QKI
Z5
ΔZ5
Δlog count
F
S
H
K
Δlog count
replacement−native
c HNRNPA1
UGAGG → UAGGG
RNAcompete Z5
1.32
6.37
Native 5-mer count
244
143
HepG2
K562
Δlog count (UAGGG − UGAGG)
d
Model accuracy (%)
Frequency ordering accuracy (%)
Full−Freq +3.78 pp
[0.90, 6.73]
e
chr8:1,844,440 C>T · K562
REF CUGGACGACAG
ALT CUGGAUGACAG
Φ fixed
Center -3.50
ASB: REF
ASB+ · exposed · n=396
preferred−other · asinh
Full Δlog count · asinh
Sequence Δlog count · asinh
f
Full
Sequence
Flat Φ
Perm. Φ
REF rule
52.04
58.77
53.46
52.36
64.77
RBP-equal accuracy (%, 95% CI)
Full · Φ rank slope (95% CI)
Direction
|Δlog count|
g ELP1
LP · 2 stars
c.3460+1G>A
chr9:108880051 C>T
exon
intron
REF AGGUCUGGGUGAGUAUC
ALT AGGUCUGGAUGAGUAUC
Φ
+1
HepG2
ASB: REF
AQR
PRPF8
ALT−REF
Δlog count · asinh
Full
Sequence
Flat Φ
Perm Φ
h
8 sites · 13 exposed ASB events
Site
RNA REF
ASL G>A GCCGGUU
ELP1 C>T UGGGUGA
EIF3F G>T GAGGUGA
HSP90B1 A>T UGAAGAA
MAP2K3 C>T AGCCGAC
MAP2K3 G>A ACCGUUU
C3 C>T AAGGUGA
ASXL1 C>A UUACCAG
ClinVar P/LP
TCGA
H: HepG2 K: K562
REF Φ · 101 nt
RBP/cell
RBM15 K
AQR H
PRPF8 H
AQR K
RBM5 H
G3BP1 H
G3BP1 H
AQR H
PRPF8 H
RBFOX2 H
BUD13 K
RBM15 K
ZNF622 K
ASB: REF
ALT−REF Δlog count · asinh