

ECE 2321 Signals and Systems

Discrete Signals



Martin Wafula

Multimedia University of Kenya

These slides are adapted from Prof Alejandro Ribeiro, UPenn

Discrete signals

Inner products and energy

Discrete complex exponentials

Orthogonality of Discrete Complex Exponentials

Appendix: Plots of Discrete Complex Exponentials


- ▶ We consider a discrete and finite time index set $\Rightarrow n = 0, 1, \dots, N - 1 \equiv [0, N - 1]$.
- ▶ A discrete signal x is a function mapping the time index set $[0, N - 1]$ to a set of real values $x(n)$

$$x : [0, N - 1] \rightarrow \mathbb{R}$$

- ▶ The values that the signal takes at time index n is $x(n)$
- ▶ Sometimes, it makes sense to talk about complex signals $\Rightarrow x : [0, N - 1] \rightarrow \mathbb{C}$
 - \Rightarrow The values $x(n) = x_R(n) + j x_I(n)$ the signal takes are complex numbers
- ▶ The space of all possible signals is the space of vectors with N components $\Rightarrow \mathbb{R}^N$ (or \mathbb{C}^N)


- ▶ The discrete delta function $\delta(n)$ is a spike at (initial) time $n = 0$

$$\delta(n) = \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{else} \end{cases}$$



- ▶ The shifted delta function $\delta(n - n_0)$ has a spike at time $n = n_0$


$$\delta(n - n_0) = \begin{cases} 1 & \text{if } n = n_0 \\ 0 & \text{else} \end{cases}$$



- ▶ This is not a new definition. Just a time shift of the previous definition


- ▶ A constant function $x(n)$ has the same value c for all n

$$x(n) = c, \quad \text{for all } n$$




- ▶ A square pulse of width M , $\Pi_M(n)$, equals one for the first M values

$$\Pi_M(n) = \begin{cases} 1 & \text{if } 0 \leq n < M \\ 0 & \text{if } M \leq n \end{cases}$$



- ▶ Can consider shifted pulses $\Pi_M(n - n_0)$, with $n_0 < N - M$

- ▶ The **Sampling time** T_s is the clock time elapsed between time indexes n and $n + 1$
- ▶ The sampling frequency $f_s := 1/T_s$ is the inverse of the sampling time
- ▶ Discrete time index n represents clock (actual) time $t = nT_s$




- ▶ Total signal duration is $T = NT_s \Rightarrow$ We “hold” the last sample for T_s time units

- For a signal of duration N define (assume N is even):

⇒ Discrete cosine of discrete frequency $k \Rightarrow x(n) = \cos(2\pi kn/N)$

⇒ Discrete sine of discrete frequency $k \Rightarrow x(n) = \sin(2\pi kn/N)$

Cosine $x(n) = \cos(2\pi kn/N)$ and sine $x(n) = \sin(2\pi kn/N)$. Frequency $k = 2$ and number of samples $N = 32$.




- Frequency k is discrete. I.e., $k = 0, 1, 2, \dots$


⇒ Have an integer number of complete oscillations

- ▶ Discrete frequency $k = 0$ is a constant
- ▶ Discrete frequency $k = 1$ is a complete oscillation
- ▶ Frequency $k = 2$ is two oscillations, for $k = 3$ three oscillations ...


Frequency $k = 0$. Number of samples $N = 32$




Frequency $k = 1$. Number of samples $N = 32$





Frequency $k = 2$. Number of samples $N = 32$



Frequency $k = 3$. Number of samples $N = 32$




- ▶ Frequency k represents k complete oscillations
- ▶ Although for large k the oscillations may be difficult to see



- ▶ Do note that we can't have more than $N/2$ oscillations
 - ⇒ Indeed $1 \rightarrow -1 \rightarrow 1, \rightarrow -1, \dots$
 - ⇒ Frequency $N/2$ is the last one with physical meaning
- ▶ Larger frequencies replicate frequencies between $k = 0$ and $k = N/2$

- Frequencies k and $N - k$ represent the same cosine




- Actually, if $k + l = N$, cosines of frequencies k and l are equivalent
- Not true for sines, but almost. The signals have opposite signs

- ▶ What is the **discrete frequency k** of a cosine of **frequency f_0** ?
- ▶ Depends on sampling time T_s , frequency $f_s = \frac{1}{T_s}$, duration $T = NT_s$
- ▶ Period of discrete cosine of frequency k is T/k (k oscillations)
- ▶ Thus, regular frequency of said cosine is $\Rightarrow f_0 = \frac{k}{T} = \frac{k}{NT_s} = \frac{k}{N}f_s$
- ▶ A cosine of **frequency f_0** has discrete **frequency $k = (f_0/f_s)N$**
- ▶ Only frequencies up to $N/2 \leftrightarrow f_s/2$ have physical meaning
- ▶ **Sampling frequency f_s** \Rightarrow **Cosines up to frequency $f_0 = f_s/2$**

- ▶ Generate $N = 32$ samples of an A note with sampling frequency $f_s = 1,760\text{Hz}$
- ▶ The frequency of an A note is $f_0 = 440\text{Hz}$. This entails a discrete frequency

$$k = \frac{f_0}{f_s} N = \frac{440\text{Hz}}{1,760\text{Hz}} 32 = 8$$

The A note observed during $T = NT_s = 18.2\text{ms}$ with a sampling rate $f_s = 1,760\text{Hz}$



- ▶ Alternatively $\Rightarrow x(n) = \cos \left[2\pi k n / N \right] = \cos \left[2\pi (f_0 / f_s) N n / N \right]$
- ▶ Which simplifies to $\Rightarrow x(n) = \cos \left[2\pi (f_0 / f_s) n \right] = \cos \left[2\pi f_0 (n T_s) \right]$

- ▶ The frequency k does not need to be an integer. In that case we talk of sampled cosines and sines
 - ⇒ **Sampled** cosine ⇒ $x(n) = \cos(2\pi kn/N)$ with arbitrary, **not necessarily integer** k
 - ⇒ **Sampled** sine ⇒ $x(n) = \sin(2\pi kn/N)$ with arbitrary, **not necessarily integer** k
- ▶ Sampled sines and cosines have fractional oscillations (k not integer)
- ▶ Discrete sines and cosines have complete oscillations (k is integer)
 - ⇒ **Discrete** sines and cosines are used to define **Fourier transforms** (As we will see later)

Discrete signals

Inner products and energy

Discrete complex exponentials

Orthogonality of Discrete Complex Exponentials


Appendix: Plots of Discrete Complex Exponentials

- ▶ Given two signals x and y with components $x(n)$ and $y(n)$ define the **inner product** of x and y as

$$\begin{aligned}\langle x, y \rangle &:= \sum_{n=0}^{N-1} x(n)y^*(n) \\ &= \sum_{n=0}^{N-1} x_R(n)y_R(n) - \sum_{n=0}^{N-1} x_I(n)y_I(n) + j \sum_{n=0}^{N-1} x_I(n)y_R(n) + j \sum_{n=0}^{N-1} x_R(n)y_I(n)\end{aligned}$$

- ▶ This is the same as the inner product between vectors x and y . Just with different notation
- ▶ The Inner product is a linear operations $\Rightarrow \langle x, y + z \rangle = \langle x, y \rangle + \langle x, z \rangle$
- ▶ Reversing the order of the factor results in conjugation $\Rightarrow \langle y, x \rangle = \langle x, y \rangle^*$

- ▶ The inner product $\langle x, y \rangle$ is the projection of the signal (vector) y on the signal (vector) x
- ▶ The value of $\langle x, y \rangle$ is how much of y falls in x direction
 - ⇒ How much y resembles x . How much x predicts y . Knowing x , how much of y we know
 - ⇒ Very importantly, if $\langle x, y \rangle = 0$ the signals are **orthogonal**. They are “unrelated”




- ▶ Define the **norm** of signal x as $\Rightarrow \|x\| := \left[\sum_{n=0}^{N-1} |x(n)|^2 \right]^{1/2} = \left[\sum_{n=0}^{N-1} |x_R(n)|^2 + \sum_{n=0}^{N-1} |x_I(n)|^2 \right]^{1/2}$
- ▶ Define the **energy** as the norm squared $\Rightarrow \|x\|^2 := \sum_{n=0}^{N-1} |x(n)|^2 = \sum_{n=0}^{N-1} |x_R(n)|^2 + \sum_{n=0}^{N-1} |x_I(n)|^2$
- ▶ The energy of x is the inner product of x with itself $\Rightarrow \|x\|^2 = \langle x, x \rangle$
- ▶ Recall that for complex numbers we have $x(n)x^*(n) = |x_R(n)|^2 + |x_I(n)|^2 = |x(n)|^2$

- ▶ Inner product can't exceed the product of the norms $\Rightarrow -\|x\| \|y\| \leq \langle x, y \rangle \leq \|x\| \|y\|$
- ▶ Inner product squared can't exceed product of energies $\Rightarrow \langle x, y \rangle^2 \leq \|x\|^2 \|y\|^2$
- ▶ If you prefer explicit expressions $\Rightarrow \sum_{n=0}^{N-1} x(n)y^*(n) \leq \left[\sum_{n=0}^{N-1} |x(n)|^2 \right] \left[\sum_{n=0}^{N-1} |y(n)|^2 \right]$
- ▶ The equalities hold if and only if the signals (vectors) x and y are collinear (aligned)

- ▶ The unit energy square pulse is the signal $\Pi_M(n)$ that takes values

$$\begin{aligned}\Pi_M(n) &= \frac{1}{\sqrt{M}} & \text{if } 0 \leq n < M \\ \Pi_M(n) &= 0 & \text{if } M \leq n\end{aligned}$$




- ▶ To compute energy of the pulse we just evaluate the definition

$$\|\Pi_M\|^2 := \sum_{n=0}^{N-1} |\Pi_M(n)|^2 = \sum_{n=0}^{M-1} \left| (1/\sqrt{M}) \right|^2 = \frac{M}{M} = 1$$

- ▶ As name indicates, the unit energy square pulse has unit energy . If pulse height is 1, energy is M .

- Shift pulse by modifying argument $\Rightarrow \Pi_M(n - K) \Rightarrow$ Pulse is now centered at K




- If the pulse support is disjoint ($K \geq M$), the inner product of two pulses is zero

$$\langle \Pi_M(n), \Pi_M(n - K) \rangle := \sum_{n=0}^{N-1} \Pi_M(n) \Pi_M(n - K) = 0$$

- Pulses are orthogonal \Rightarrow They are “unrelated.” One pulse does not predict the other

- If $K < M$ the pulses overlap. They overlap between $n = K$ and $n = M - 1$. Thus, the inner product is

$$\langle \Pi_M(n), \Pi_M(n - K) \rangle := \sum_{n=0}^{N-1} \Pi_M(n) \Pi_M(n - K) = \sum_{n=K}^{M-1} \left(1/\sqrt{M}\right) \left(1/\sqrt{M}\right) = \frac{M - K}{M} = 1 - \frac{K}{M}$$



- Inner product proportional to relative overlap \Rightarrow How much the pulses are “related” to each other

Discrete signals

Inner products and energy

Discrete complex exponentials

Orthogonality of Discrete Complex Exponentials


Appendix: Plots of Discrete Complex Exponentials

- ▶ Discrete complex exponential of discrete frequency k and duration N


$$e_{kN}(n) = \frac{1}{\sqrt{N}} e^{j2\pi kn/N} = \frac{1}{\sqrt{N}} \exp(j2\pi kn/N)$$

- ▶ The complex exponential function is $\Rightarrow e^{j2\pi kn/N} = \cos(2\pi kn/N) + j \sin(2\pi kn/N)$
- ▶ The Real part is a discrete cosine. The imaginary part a discrete sine. An oscillation

$\text{Re}(e^{j2\pi kn/N})$, with $k = 2$ and $N = 32$



$\text{Im}(e^{j2\pi kn/N})$, with $k = 2$ and $N = 32$



[P1] For frequency $k = 0$, the exponential $e_{kN}(n) = e_{0N}(n)$ is a constant $\Rightarrow e_{kN}(n) = \frac{1}{\sqrt{N}} = \frac{1}{\sqrt{N}} 1$

[P2] For frequency $k = N$, the exponential $e_{kN}(n) = e_{NN}(n)$ is a constant. True for any multiple $k \in \dot{N}$

$$e_{N\dot{N}}(n) = \frac{e^{j2\pi \dot{N}n/N}}{\sqrt{N}} = \frac{(e^{j2\pi})^n}{\sqrt{N}} = \frac{(1)^n}{\sqrt{N}} = \frac{1}{\sqrt{N}}$$

[P3] For $k = \frac{N}{2}$, the exponential $e_{kN}(n) = e_{N/2N}(n) = (-1)^n / \sqrt{N}$. Fastest possible oscillation with N samples

$$e_{N/2N}(n) = \frac{e^{j2\pi (N/2)n/N}}{\sqrt{N}} = \frac{(e^{j\pi})^n}{\sqrt{N}} = \frac{(-1)^n}{\sqrt{N}}$$

That $e^{j2\pi} = 1$ follows from $e^{j\pi} = -1$. Which follows from $e^{j\pi} + 1 = 0$. Relates five most important constants in mathematics.

Theorem

If the *frequency difference is* $k - l = N$ the signals $e_{kN}(n)$ and $e_{lN}(n)$ coincide for all n , i.e.,

$$e_{kN}(n) = \frac{e^{j2\pi kn/N}}{\sqrt{N}} = \frac{e^{j2\pi ln/N}}{\sqrt{N}} = e_{lN}(n)$$

- **Exponentials** with frequencies k and l are **equivalent** if the frequency difference is $k - l = N$

Proof.

- ▶ We prove by showing that the ratio $e_{kN}(n)/e_{lN}(n) = 1$. Combine exponents

$$\frac{e_{kN}(n)}{e_{lN}(n)} = \frac{e^{j2\pi kn/N}}{e^{j2\pi ln/N}} = e^{j2\pi(k-l)n/N}$$

- ▶ By hypothesis we have that $k - l = N$. Therefore, the latter simplifies to

$$\frac{e_{kN}(n)}{e_{lN}(n)} = e^{j2\pi Nn/N} = \left[e^{j2\pi} \right]^n = 1^n = 1$$

□

- ▶ Canonical set \Rightarrow Suffice to look at N consecutive frequencies, e.g., $k = 0, 1, \dots, N - 1$

$$\begin{array}{cccc} -N, & -N + 1, & \dots, & -1 \\ 0, & 1, & \dots, & N - 1 \\ N, & N + 1, & \dots, & 2N - 1 \end{array}$$

- ▶ Another canonical choice is to make $k = 0$ a center frequency

$$\begin{array}{ccccccc} -N/2 + 1, & \dots, & -1, & 0, & \dots, & N/2 \\ N/2 + 1, & \dots, & N - 1, & N, & \dots, & 3N/2 \end{array}$$

- ▶ With N even (as usual) we use $N/2$ positive frequencies and $N/2 - 1$ negative frequencies
- ▶ From one canonical set to the other \Rightarrow Chop and shift

Theorem

Opposite frequencies k and $-k$ yield conjugate signals: $e_{-kN}(n) = e_{kN}^*(n)$

Proof.

- ▶ Just use the definitions to write the chain of equalities

$$e_{-kN}(n) = \frac{e^{j2\pi(-k)n/N}}{\sqrt{N}} = \frac{e^{-j2\pi kn/N}}{\sqrt{N}} = \left[\frac{e^{j2\pi kn/N}}{\sqrt{N}} \right]^* = e_{kN}^*(n)$$

□

- ▶ Opposite frequencies \Rightarrow Same real part. Opposite imaginary part

\Rightarrow The cosine is the same, the sine changes sign

- ▶ Of N canonical frequencies, only $\frac{N}{2} + 1$ are distinct. No more than $\frac{N}{2}$ oscillations in N samples

$$0, \quad \begin{matrix} 1, & \dots, & N/2 - 1 \\ -1, & \dots, & -N/2 + 1 \\ N - 1, & \dots, & N/2 + 1 \end{matrix} \quad N/2$$

- ▶ The frequencies 0 and $N/2$ do not have a conjugate counterpart. All Others do
- ▶ The canonical set $-N/2 + 1, \dots, -1, 0, 1, \dots, N/2$ is easier to interpret
 - ⇒ Positive frequencies ranging from 0 to $N/2 \leftrightarrow f_s/2$ have physical meaning
 - ⇒ The negative frequencies are conjugates of the corresponding positive frequencies

Discrete signals

Inner products and energy

Discrete complex exponentials

Orthogonality of Discrete Complex Exponentials

Appendix: Plots of Discrete Complex Exponentials

Theorem

Complex exponentials with nonequivalent frequencies are orthogonal. I.e.

$$\langle e_{kN}, e_{lN} \rangle = 0$$

when $k - l < N$. E.g., when $k = 0, \dots, N - 1$, or $k = -N/2 + 1, \dots, N/2$.

- ▶ Signals of canonical sets are “unrelated.” Different rates of change
- ▶ Also note that the energy is $\|e_{kN}\|^2 = \langle e_{kN}, e_{kN} \rangle = 1$
- ▶ Exponentials with frequencies $k = 0, 1, \dots, N - 1$ are orthonormal

$$\langle e_{kN}, e_{lN} \rangle = \delta(l - k)$$

- ▶ They are an orthonormal basis of signal space with N samples

Proof.

- Use definitions of inner product and discrete complex exponential to write

$$\langle e_{kN}, e_{lN} \rangle = \sum_{n=0}^{N-1} e_{kN}(n) e_{lN}^*(n) = \sum_{n=0}^{N-1} \frac{e^{j2\pi kn/N}}{\sqrt{N}} \frac{e^{-j2\pi ln/N}}{\sqrt{N}}$$

- Regroup terms to write as geometric series

$$\langle e_{kN}, e_{lN} \rangle = \frac{1}{N} \sum_{n=0}^{N-1} e^{j2\pi(k-l)n/N} = \frac{1}{N} \sum_{n=0}^{N-1} \left[e^{j2\pi(k-l)/N} \right]^n$$

- Geometric series with basis a sums to $\sum_{n=0}^{N-1} a^n = (1 - a^N)/(1 - a)$. Thus,

$$\langle e_{kN}, e_{lN} \rangle = \frac{1}{N} \frac{1 - \left[e^{j2\pi(k-l)/N} \right]^N}{1 - e^{j2\pi(k-l)/N}} = \frac{1}{N} \frac{1 - 1}{1 - e^{j2\pi(k-l)/N}} = 0$$

- Completed proof by noting $\left[e^{j2\pi(k-l)/N} \right]^N = e^{j2\pi(k-l)} = \left[e^{j2\pi} \right]^{(k-l)} = 1$

□

Discrete signals


Inner products and energy

Discrete complex exponentials

Orthogonality of Discrete Complex Exponentials


Appendix: Plots of Discrete Complex Exponentials

- ▶ When signal durations is $N = 2$ only frequencies $k = 0$ and $k = 1$ represent distinct signals




- ▶ The signals are real, they have no imaginary parts

- When $N = 4$, $k = 0, 1, 2$ are distinct. $k = -1$ is conjugate of $k = 1$




- Can also use $k = 3$ as canonical instead of $k = -1$ (conjugate of $k = 1$)

- Frequencies from $k = 1$ to $k = 4$ represent distinct signals



- Frequencies $k = -1$ to $k = -3$ are conjugate signals of $k = 1$ to $k = 3$



- All other frequencies represent one of the signals above

- There are 9 distinct frequencies and 7 conjugates (not shown). Some look like actual oscillations

