

A Practical Guide to Theatre-to-ICU Obstetric Handover

Residents' Academic Committee^{a,b,*}

^a*Department of Anaesthesia and Intensive Care, Aminu Kano Teaching Hospital, Kano, Nigeria,*

^b*Department of Anaesthesia and Intensive Care, Aminu Kano Teaching Hospital,*

Abstract

Handover between theatre and the intensive care unit is a recognised point of risk, particularly for obstetric patients who may have ongoing haemorrhage risk, pre-eclampsia, or cardiovascular instability. This guide sets out a structured, repeatable handover sequence intended for use by residents rotating between obstetric theatre and the ICU, drawing on the wider patient-safety-checklist literature.

Keywords: handover, obstetric anaesthesia, patient safety, ICU admission

Departmental guide

Prepared by the Residents' Academic Committee

Department of Anaesthesia and Intensive Care, Aminu Kano Teaching Hospital

Summary. Handover between theatre and the intensive care unit is a recognised point of risk, particularly for obstetric patients who may have ongoing haemorrhage risk, pre-eclampsia, or cardiovascular instability. This guide sets out a structured, repeatable handover sequence intended for use by residents rotating between obstetric theatre and the ICU.

KEYWORDS · HANDOVER · OBSTETRIC ANAESTHESIA · PATIENT SAFETY · ICU ADMISSION

1. Why a structured handover matters

Verbal handovers that aren't structured tend to omit items unpredictably — often exactly the detail that matters for the next few hours of care. A short,

*Corresponding author

Email address: anaesthesia@akth.org.ng (Residents' Academic Committee)

consistent structure reduces that risk without adding much time to a busy transfer. This mirrors the logic behind the WHO Surgical Safety Checklist, which was shown to reduce post-operative morbidity and mortality across a range of hospital settings when used consistently [1, 2].

2. Suggested handover structure

1. **Patient identification & background** — name, age, parity, key obstetric history, indication for surgery.
2. **Procedure performed** — type of anaesthesia, surgical procedure, any intra-operative complications.
3. **Haemodynamic course** — blood loss, fluids and blood products given, vasopressor use, any instability and how it was managed.
4. **Airway and respiratory status** — extubation status, oxygen requirement, any airway concerns for the ICU team to be aware of.
5. **Analgesia plan** — what has been given, what is prescribed next, and any regional technique in place (e.g. spinal opioid, wound catheter).
6. **Outstanding concerns and thresholds for escalation** — specifically what should prompt the ICU team to call anaesthesia back.

3. Using this guide

This is a teaching resource for residents, not a substitute for local protocol. Departments should adapt the sequence to match their own documentation and handover tools, and residents should confirm current local practice with their supervising consultant.

Have a guide, protocol summary, or teaching resource to contribute? Email the department to have it added here.

References

- [1] A. B. Haynes, T. G. Weiser, W. R. Berry, S. R. Lipsitz, A.-H. S. Breizat, E. P. Dellinger, T. Herbosa, S. Joseph, P. L. Kibatala, M. C. M. Lapitan, A. F. Merry, K. Moorthy, R. K. Reznick, B. Taylor, A. A. Gawande, A surgical safety checklist to reduce morbidity and mortality in a global population, *New England Journal of Medicine* 360 (5) (2009) 491–499. [doi:10.1056/NEJMs0810119](https://doi.org/10.1056/NEJMs0810119).
- [2] Safe Surgery Saves Lives Study Group, [WHO Guidelines for Safe Surgery 2009: Safe Surgery Saves Lives](https://www.who.int/publications/i/item/9789241598552), World Health Organization, Geneva, 2009. URL <https://www.who.int/publications/i/item/9789241598552>