

THE 3-ADIC MIXING RATE OF THE SYRACUSE RANDOM VARIABLE: REDUCTIONS, A BOUNDARY-LAYER MECHANISM, AND A SHARP-RATE CONJECTURE

HIKARU WAKAURA

ABSTRACT. Let **Syrac** be the Syracuse random variable, the 3-adic limit of $\sum_{j \geq 1} 3^{j-1} 2^{-A_j}$ with A_j the partial sums of i.i.d. $\text{Geom}(1/2)$ exponents, and let $\delta_m = \max_{3 \nmid \xi} |\mathbb{E} e(\xi \mathbf{Syrac}/3^m)|$ be its maximal nontrivial Fourier coefficient — the quantity whose decay drives Tao’s almost-all result for the Collatz map and any effective version thereof. Tao proved $\delta_m \ll_A m^{-A}$ and explicitly flagged the exponential improvement as possible but unpursued; no later work supplies it. We prove: (1) since 2 is a primitive root mod 3^m , the Fourier problem is exactly a one-dimensional lattice recursion over the exponent of the frequency, with δ_m non-increasing and exactly stationary after m steps; (2) a first-passage renormalization identity with the certified bound $|F(d, r)| \leq 2^{-r} \delta_{d-1}$; (3) a weighted contraction lemma whose optimal rate is exactly 2^{-I_0} , $I_0 = (1 - H(\theta))/\theta = 0.0793186\dots$, $\theta = \log_3 2$, reducing the upper-bound problem to the control of frequencies near and above the critical line $2^k \approx 3^m$; (4) the real Syracuse variable is a Kesten chain with tail exponent exactly 1; (5) an exact 2-adic law: over full multiplicative periods, the 2-adic valuations of the signed residues of $2^K \bmod 3^m$ inside a symmetric ball are geometrically distributed, by an elementary two-line identity. Assembling (1)–(5) with an audited weighted- ℓ^1 bookkeeping and sharp product bounds along Tao’s update process, we obtain a complete conditional proof of $\delta_m \leq \text{poly}(m) 2^{-0.0504 m}$ whose single unproven input is a local equidistribution statement for the orbit $\{2^e \bmod 3^n\}$ in windows of length $O(\log q)$ — a finite-orbit quantitative form of Furstenberg $\times 2 \times 3$ rigidity that we show lies outside current methods. For each fixed m the input is decidable, and we certify the resulting unconditional bounds at $m = 64, 128, 256$, with the certified rate improving toward 0.0504. Certified lattice computations to $m = 1200$ support the sharp-rate conjecture $\delta_m = \Theta(2^{-I_0 m} m^{-c})$: the extremal frequency tracks the cycle-resonance line $2^{k^*} \approx 3^m$ at constant deficit $k^* = m \log_2 3 - (7.0 \pm 0.3)$, and the extremal amplitude is generated by a single pinned boundary-layer bridge which already attains $2^{-I_0 m} m^{-1.6}$ with only polynomial phase loss.

1. INTRODUCTION

The Collatz $(3x + 1)$ conjecture concerns the map on positive integers that sends even n to $n/2$ and odd n to $3n + 1$. The strongest known result, due to Tao [1], is that almost all orbits (in logarithmic density) attain almost bounded values. At the analytic heart of that proof sits a single quantitative object: the 3-adic equidistribution of the *Syracuse random variable*, the random 3-adic integer

$$\mathbf{Syrac} = \sum_{j \geq 1} 3^{j-1} 2^{-A_j}, \quad A_j = a_1 + \dots + a_j, \quad a_i \text{ i.i.d., } \mathbb{P}(a_i = a) = 2^{-a},$$

which describes the residues mod 3^m of the accelerated Collatz iteration started from a random odd integer. Writing $e(x) = e^{2\pi i x}$, the quantity that controls everything is the maximal nontrivial

Date: September 1, 2026.

2020 Mathematics Subject Classification. Primary 11B83; Secondary 11K16, 37P10, 60H25.

Key words and phrases. $3x + 1$ problem, Syracuse random variable, Fourier decay, 3-adic equidistribution, Kesten chain, Furstenberg $\times 2 \times 3$ rigidity, experimental mathematics.

Fourier coefficient

$$\delta_m = \max_{\xi \in \mathbb{Z}/3^m\mathbb{Z}, 3 \nmid \xi} |\mathbb{E} e(\xi \mathbf{Syrac}/3^m)|.$$

Tao's Proposition 1.17 proves $\delta_m \ll_A m^{-A}$ for every fixed A — superpolynomial decay — and Section 1.4 of [1] states explicitly that an improvement to $O(\exp(-cm))$ may be possible but is not needed there and not pursued. A survey of the citing literature (Section 9) shows that no later work supplies the exponential improvement. Its value would be more than aesthetic: the rate of δ_m is the rate-limiting constant in any *effective* version of Tao's theorem.

However, the exact decay rate of δ_m — even its numerical value — appears never to have been determined. Here we determine it numerically with certified lattice computations, identify its mechanism, prove the structural theorems that reduce the upper-bound problem to a single isolated arithmetic statement, and certify unconditional exponential bounds for fixed levels $m \leq 256$. Our results are of four kinds, and we keep the four statuses — *theorem*, *computational fact* (exact or certified computation), *conditional theorem*, and *conjecture* — explicitly labeled throughout.

Theorems. (i) Since 2 is a primitive root mod 3^m , every nontrivial frequency is a power of 2, and the two-variable Fourier problem collapses, with no assumptions, to a one-dimensional recursion over the exponent lattice, with δ_m non-increasing in m and the underlying residue chain exactly stationary after m steps (Theorem 2.1). (ii) A first-passage renormalization identity expresses g_m in terms of shorter chains, with the certified kernel bound $|F(d, r)| \leq 2^{-r} \delta_{d-1}$ and an exact identification of the phase-free region with the characteristic function of the real Syracuse variable (Theorem 3.1). (iii) A weighted contraction lemma: with weight $2^{\lambda u}$ in the coordinate $u = k - m \log_2 3$, the phase-free weighted row sum equals $f(\lambda) = 3^\lambda / (2^{1+\lambda} - 1)$, and

$$\min_{\lambda} f(\lambda) = 2^{-I_0}, \quad I_0 = \frac{1 - H(\theta)}{\theta} = D(\text{Geom}(\theta) \parallel \text{Geom}(1/2)) = 0.0793186 \dots,$$

$\theta = \log_3 2$, H the binary entropy (Theorem 4.1) — below the critical line no phase cancellation is needed to contract at exactly the conjectured rate, so the upper-bound problem reduces to the near-line and above-line frequencies. (iv) The real Syracuse variable is a Kesten chain whose Kesten equation $\mathbb{E}[(3 \cdot 2^{-a})^\kappa] = 1$ holds at $\kappa = 1$ *exactly* (the identity $3 = 2^2 - 1$), placing it at the Cauchy-type boundary with log-periodic corrections (Theorem 5.1). (v) An exact 2-adic law on full periods: the identity $\nu_2(r(K)) \geq \sigma \iff |r(K - \sigma)| \leq M/2^{\sigma+1}$ for signed residues $r(K)$ of $2^K \bmod M = 3^m$ converts 2-adic valuation into Archimedean size, and yields that within any symmetric ball the valuation law over a full period is geometric with explicit error (Theorem 6.2).

Conditional theorem. Assembling (i)–(v) with a capped weighted- ℓ^1 bookkeeping (audited exactly on all 399 computed levels, with zero violations) and sharp product bounds along Tao's two-dimensional update process, we obtain a complete proof architecture for

$$\delta_m \leq \text{poly}(m) \cdot 2^{-0.0504 m}$$

— the qualitative form of the improvement flagged in [1] — whose *single* unproven input (Hypothesis 9.4) is a local equidistribution statement: entries of the orbit $\{2^e \bmod 3^n\}$ into a short symmetric interval have geometrically decaying 2-adic valuations, uniformly over windows of length $O(\log q)$, $q = 3^n$. We show (Section 9) that this statement is a finite-orbit quantitative form of Furstenberg's $\times 2 \times 3$ phenomenon [8], that the effective results of Bourgain–Lindenstrauss–Michel–Venkatesh [9] do not reach it, and that the window length $O(\log q)$ is exponentially below the reach of the known exponential-sum technology; it is, however, decidable in time $O(m^4)$ for each fixed m .

Computational facts. Certified exact-arithmetic computations give: full-frequency scans for $m \leq 15$; rescaled window computations to $m = 1200$ showing that the extremal frequency exponent satisfies $k^* = m \log_2 3 - (7.0 \pm 0.3)$ with the deficit *constant* over $m = 200$ –1200, and a decay rate in $[0.0790, 0.0800]$ bits per level approaching I_0 one-sidedly; a boundary-layer structure theory (Section 8); and unconditional certified instances

$$\delta_{64} \leq C 2^{-0.0350 \cdot 64}, \quad \delta_{128} \leq C 2^{-0.0406 \cdot 128}, \quad \delta_{256} \leq C 2^{-0.0457 \cdot 256},$$

with all band constraints closing at positive slack and the certified rate increasing toward 0.0504 (Section 10).

Conjecture. The sharp rate (Conjecture 11.1): $\delta_m = \Theta(2^{-I_0 m} m^{-c})$. Equivalently, the 3-adic mixing bottleneck of Syracuse is the cycle-resonance line $2^A \approx 3^k$, and the mixing rate coincides, after clock conversion, with the classical stopping-time density exponent $1 - H(\theta)$: one constant read in three clocks.

Method. This paper practices experimental mathematics with pre-registered decision criteria: for each major numerical discrimination we declared acceptance/rejection thresholds before computing, and we report the outcomes — including a rejected candidate rate — in Appendix A, which doubles as a calibration log. All certified quantities use exact integer or interval-safe arithmetic; scripts and frozen data accompany the paper. A companion paper [14] studies the 2-adic (certificate-complexity) side of the same constant.

2. SETUP AND THE ONE-DIMENSIONAL REDUCTION

Fix $m \geq 1$ and let w_0 be arbitrary in $\mathbb{Z}/3^m\mathbb{Z}$. The Syracuse residue chain is

$$(1) \quad w_t = (3w_{t-1} + 1) \cdot 2^{-a_t} \bmod 3^m, \quad a_t \text{ i.i.d., } \mathbb{P}(a_t = a) = 2^{-a} \ (a \geq 1),$$

where 2^{-1} denotes the inverse of $2 \bmod 3^m$. This is the law of the residues $\bmod 3^m$ of the accelerated Collatz map $n \mapsto (3n + 1)/2^{\nu_2(3n+1)}$ along the odd iterates, in the sense made precise in [1, §1]; the stationary law of (1) is the law of **Syrac** $\bmod 3^m$.

Theorem 2.1 (one-dimensional reduction). (a) *2 is a primitive root mod 3^m for every $m \geq 1$: $\text{ord}(2 \bmod 3^m) = 2 \cdot 3^{m-1} = |(\mathbb{Z}/3^m\mathbb{Z})^\times|$.*

(b) *Consequently every ξ with $3 \nmid \xi$ is $\pm$ a power of 2 mod 3^m , and with $g_m(k) := \mathbb{E} \mathfrak{e}(2^k \mathbf{Syrac}/3^m)$ (exponents read mod $\text{ord}(2)$) one has $\delta_m = \max_k |g_m(k)|$ and the exact lattice recursion*

$$(2) \quad g_m(k) = \sum_{a \geq 1} 2^{-a} \mathfrak{e}\left(\frac{2^{k-a} \bmod 3^m}{3^m}\right) g_{m-1}(k - a).$$

(c) *The chain (1) is exactly stationary after m steps: for $t \geq m$ the law of w_t equals the law of **Syrac** $\bmod 3^m$, for every initial w_0 .*

(d) $\delta_m \leq \delta_{m-1}$ for all $m \geq 2$.

Proof. (a) From $2^2 = 1 + 3$ and the lifting-the-exponent identity $(1 + 3^s u)^3 \equiv 1 + 3^{s+1} u \pmod{3^{s+2}}$ one gets $\nu_3(2^{2 \cdot 3^t} - 1) = t + 1$ exactly, so the order of $2 \bmod 3^m$ divides $2 \cdot 3^{m-1}$ and is divisible by 3^{m-1} ; and $2^{3^j} \equiv (-1)^{3^j} = -1 \not\equiv 1 \pmod{3}$ shows the order is even. Hence $\text{ord}(2) = 2 \cdot 3^{m-1}$.

(b) By (a) the cyclic group generated by 2 is the whole unit group, so the powers of 2 exhaust all ξ with $3 \nmid \xi$, and the maximum defining δ_m is attained on the exponent lattice. The recursion follows by conditioning (1) on the last step: with $\xi = 2^k$,

$$\mathfrak{e}(\xi(3w + 1)2^{-a}/3^m) = \mathfrak{e}(2^{k-a} \cdot 3w/3^m) \mathfrak{e}(2^{k-a}/3^m) = \mathfrak{e}(2^{k-a}w/3^{m-1}) \mathfrak{e}(2^{k-a}/3^m),$$

and taking expectations over $w \sim \mathbf{Syrac} \bmod 3^{m-1}$ and a gives (2).

(c) By induction on j : $w_t \bmod 3^j$ depends only on $(a_{t-j+1}, \dots, a_t)$, because $(3w+1) \bmod 3^j$ depends only on $w \bmod 3^{j-1}$ and multiplication by the unit 2^{-a} preserves this. For $j = m$, $t \geq m$, the law of w_t is a fixed function of the last m steps, independent of w_0 ; it is therefore the unique stationary law.

(d) Bound (2) by the triangle inequality: $|g_m(k)| \leq \sum_a 2^{-a} |g_{m-1}(k-a)| \leq \delta_{m-1}$, using that every frequency 2^{k-a} is again a unit $\bmod 3^{m-1}$. $\square$

Remark 2.2. Theorem 2.1(b) upgrades to a theorem what would otherwise be a heuristic restriction to special frequencies: *every* nontrivial frequency lives on the exponent lattice, so the recursion (2) loses nothing. Full-frequency exhaustive scans for $m \leq 15$ confirm that the maximizer is always a power of 2 in a narrow window of exponents (Section 7).

3. FIRST-PASSAGE RENORMALIZATION

Let $Y_m = \sum_{j=1}^m 3^{j-1} 2^{-A_j}$ denote the *real* Syracuse variable (the same series read in $\mathbb{R}$), and write $x = 2^k/3^m$.

Theorem 3.1 (renormalization identity). *For all m, k :*

- (a) (*phase-free region*) *On the event $\{A_m \leq k\}$ the 3-adic phase coincides exactly with the real one: $e(2^k \mathbf{Syrac}_m/3^m) = e(2^k Y_m/3^m)$ pathwise, where $\mathbf{Syrac}_m$ is the level- m truncation. Hence $\mathbb{E}[e(2^k \mathbf{Syrac}_m/3^m); A_m \leq k] = \mathbb{E}[e(xY_m); A_m \leq k]$.*
- (b) (*first-passage decomposition*) *With $\text{amp}_j(A) = \mathbb{E}[e(xY_j); A_j = A]$ and*

$$F(d, r) = \sum_{a>r} 2^{-a} e\left(\frac{2^{r-a} \bmod 3^d}{3^d}\right) V_{d-1}(2^{r-a}), \quad V_{d-1}(\xi) = \mathbb{E} e(\xi \mathbf{Syrac}_{d-1}/3^{d-1}),$$

one has the exact identity

$$g_m(k) = \mathbb{E}[e(xY_m); A_m \leq k] + \sum_{j<m} \sum_{A \leq k} \text{amp}_j(A) F(m-j, k-A),$$

obtained by splitting at the first step where the valuation sum exceeds k .

- (c) (*kernel bound*) $|F(d, r)| \leq 2^{-r} \delta_{d-1}$.

Proof. (a) On $\{A_m \leq k\}$ every exponent $k - A_j$ ($j \leq m$) is non-negative, so $2^k \cdot (2^{-A_j} \bmod 3^m) \equiv 2^{k-A_j} \bmod 3^m$ with 2^{k-A_j} a genuine integer, and $e(\sum_j 3^{j-1} 2^{k-A_j}/3^m)$ is simultaneously the 3-adic and the real phase.

(b) Decompose according to the first index $j+1$ with $A_{j+1} > k$ (if none, we are in the event of (a)). Conditioning on $A_j = A \leq k$ and on the overshooting step $a_{j+1} = a > k - A$, the remaining $m-j-1$ steps form a fresh chain at level $m-j-1$, and collecting phases gives the stated kernel with $r = k - A$; the identity is exact. (It was additionally verified numerically to relative error 10^{-14} across levels.)

(c) $|F(d, r)| \leq \sum_{a>r} 2^{-a} |V_{d-1}(2^{r-a})| \leq \delta_{d-1} \sum_{a>r} 2^{-a} = 2^{-r} \delta_{d-1}$, again because the frequencies are units. $\square$

Remark 3.2 (why absolute values do not suffice). It is tempting to iterate Theorem 3.1(b)–(c) with absolute values throughout. This provably fails to close: the phase-free main term must then be bounded by the large-deviation probability $\mathbb{P}(A_m \leq k)$, and near the critical line the inductive inequality loses a constant factor per level that destroys any exponent. The failure is structural — on the good event the final phase still rotates by at least $1/3$ of a turn, so positivity-based lower bounds are equally impossible, and the modulus of g_m is carried by cancellation among crossing terms (Section 8). This motivates the weighted approach of the next section.

4. WEIGHTED CONTRACTION AND THE REDUCTION OF THE UPPER BOUND

Write $\alpha = \log_2 3$ and use the line coordinate $u = k - m\alpha$.

Theorem 4.1 (weighted contraction). *For $\lambda > -1$ define the phase-free weighted row sum of the recursion (2) with weight $2^{\lambda u}$:*

$$f(\lambda) = \sum_{a \geq 1} 2^{-a} 2^{\lambda(\alpha-a)} = \frac{3^\lambda}{2^{1+\lambda} - 1}.$$

Then f is strictly log-convex on $(-1, \infty)$ and

$$\min_{\lambda} f(\lambda) = 2^{-I_0}, \quad I_0 = \frac{1 - H(\theta)}{\theta}, \quad \text{attained at } 2^{1+\lambda^*} = \frac{\alpha}{\alpha - 1}, \quad \lambda^* = 0.4380326 \dots$$

Applying the weight to (2) and bounding phases by 1 wherever the weight decays, one obtains for the weighted norm $\|g_m\|_\lambda = \sup_k 2^{\lambda u} |g_m(k)|$ the one-step inequality $\|g_m\|_\lambda \leq f(\lambda) \|g_{m-1}\|_\lambda +$ (boundary terms supported near and above $u = 0$): with no phase input at all, the below-line region contracts at exactly the conjectured rate, and the upper-bound problem for δ_m reduces to the control of frequencies with $u \gtrsim 0$. The precise audited form of this reduction is the capped functional described after the proof.

Proof. The row sum is a geometric series: $\sum_a 2^{-a(1+\lambda)} = 1/(2^{1+\lambda} - 1)$, times $2^{\lambda\alpha} = 3^\lambda$. For the minimum, substitute $t = 2^{1+\lambda} > 1/2$: then $3^\lambda = t^\alpha/3$ and $f = t^\alpha/(3(t-1))$, with critical point $\alpha(t-1) = t$, i.e. $t^* = \alpha/(\alpha-1)$, and

$$\min f = \frac{(t^*)^\alpha}{3(t^* - 1)} = \frac{\alpha^\alpha}{3(\alpha - 1)^{\alpha-1}}.$$

Taking $-\log_2$ and using $\log_2 3 = \alpha$: $-\log_2 \min f = \alpha - \alpha \log_2 \alpha + (\alpha - 1) \log_2(\alpha - 1)$. On the other hand, with $\theta = 1/\alpha$,

$$\frac{1 - H(\theta)}{\theta} = \alpha \left(1 - \frac{1}{\alpha} \log_2 \alpha - \left(1 - \frac{1}{\alpha} \right) \log_2 \frac{\alpha}{\alpha-1} \right) = \alpha - \alpha \log_2 \alpha + (\alpha - 1) \log_2(\alpha - 1),$$

which is the same expression. (The identity was additionally checked to 50 digits.) Log-convexity holds because $f(\lambda)$ is the moment generating function $\mathbb{E} 2^{\lambda(\alpha-a)}$ of the step variable. $\square$

Two consequences organize everything that follows. First, I_0 is simultaneously a Kullback–Leibler divergence:

$$(3) \quad I_0 = D(\text{Geom}(\theta) \parallel \text{Geom}(1/2)), \quad \theta I_0 = 1 - H(\theta),$$

the second equality being a one-line identity; the tilted step law at the optimum is exactly $\text{Geom}(\theta)$, whose mean step is $1/\theta = \alpha$ — the drift that tracks the critical line. Second, the reduction can be run with a *free* weight exponent. For $\mu < 0$ define

$$(4) \quad r(\mu) = \mu \alpha + \log_2(2^{1-\mu} - 1),$$

the contraction rate of the capped weighted- ℓ^1 functional $N_\mu(m) = \sum_k \min(2^{\mu u}, 1) |g_m(k)|$; one checks $\max_\mu r(\mu) = I_0$ at $\mu = -\lambda^*$, and smaller $|\mu|$ trades rate for weaker demands on the above-line frequencies. The functional N_μ has no above-line escape route: the weight decays upward, and, unlike the sup norm, the ℓ^1 sum sees every frequency.

Computational Fact 4.2 (audited bookkeeping). The one-step inequality $N_\mu(m) \leq 2^{-r(\mu)} N_\mu(m-1)$ with $\mu = -0.16$, $r(\mu) = 0.0504$, was audited against the exact lattice data on all 399 computed levels: zero violations, with no injection terms required within the window. The remaining bookkeeping (seam and wrap-around of the exponent circle) is exactly the constraint system used in the certified instances of Section 10.

5. KESTEN STRUCTURE OF THE REAL VARIABLE

Theorem 5.1. *The real Syracuse variable $Y = \sum_{j \geq 1} 3^{j-1} 2^{-A_j}$ satisfies the stochastic fixed-point equation $Y \stackrel{d}{=} 3 \cdot 2^{-a} Y + 2^{-a}$ ($a \sim \text{Geom}(1/2)$ independent of Y on the right), and its Kesten equation $\mathbb{E}[(3 \cdot 2^{-a})^\kappa] = 1$ holds exactly at $\kappa = 1$:*

$$\mathbb{E}[3 \cdot 2^{-a}] = \sum_{a \geq 1} 2^{-a} \cdot 3 \cdot 2^{-a} = 3 \sum_{a \geq 1} 4^{-a} = 3 \cdot \frac{1}{3} = 1.$$

The multiplier $\log(3 \cdot 2^{-a}) = \log 3 - a \log 2$ is arithmetically lattice, and the Goldie conditions hold; hence $\mathbb{P}(Y > t) = h(\log t)/t (1 + o(1))$ with h log-periodic, $\mathbb{E}Y = \infty$, and the tail exponent is exactly 1.

Proof. The fixed-point equation restates the definition after one step. The Kesten computation at $\kappa = 1$ is the displayed line — $3^\kappa = 2^{\kappa+1} - 1$ has the integer solution $\kappa = 1$ because $3 = 2^2 - 1$. The moment and lattice conditions of implicit renewal theory [6, 7] are immediate ($\mathbb{E}[(3 \cdot 2^{-a}) \log^+(3 \cdot 2^{-a})] < \infty$), giving the tail with log-periodic prefactor in the lattice case. $\square$

This places Syracuse at the *boundary* Cauchy-type case: power corrections in $1/m$ and log-periodic constants are forced, which is exactly what the numerics of Section 7 display, and the polynomial factor in Conjecture 11.1 should be regarded as this mechanism's signature rather than as numerical noise.

6. AN EXACT 2-ADIC LAW ON FULL PERIODS

The conditional route of Section 9 needs one probabilistic input about the deterministic orbit $\{2^K \bmod 3^m\}$: entries into a short symmetric interval should have geometrically distributed 2-adic valuations. On *full periods* this is a theorem, by an elementary identity that converts the 2-adic condition into an Archimedean one.

For odd $M \geq 3$ let $r(K) \in (-M/2, M/2]$ denote the signed residue of $2^K \bmod M$.

Lemma 6.1 (valuation–size identity). *For all K and $\sigma \geq 0$:*

$$\nu_2(r(K)) \geq \sigma \quad \Longleftrightarrow \quad |r(K - \sigma)| \leq \frac{M}{2^{\sigma+1}}.$$

Proof. ($\Rightarrow$) If $2^\sigma \mid r(K)$ then $r(K)/2^\sigma$ is an integer congruent to $2^{K-\sigma} \bmod M$ with absolute value at most $M/2^{\sigma+1} < M/2$, hence it is the signed residue $r(K - \sigma)$. ($\Leftarrow$) If $|r(K - \sigma)| \leq M/2^{\sigma+1}$ then $2^\sigma r(K - \sigma)$ is congruent to $2^K \bmod M$ with absolute value at most $M/2$, hence equals $r(K)$, which is thus divisible by 2^σ . $\square$

Theorem 6.2 (geometric law on cycles). *Let $M = 3^m$ and let K range over a full period of the orbit $\{2^K \bmod 3^m\}$ (length $2 \cdot 3^{m-1}$, all units, by Theorem 2.1(a)). Then for every $\varepsilon \in (0, 1/2]$ and every $\sigma \geq 0$,*

$$\mathbb{P}_K\left(\nu_2(r(K)) \geq \sigma \mid |r(K)| \leq \varepsilon 3^m\right) = 2^{-\sigma} + O\left(\frac{2^\sigma}{\varepsilon 3^m}\right),$$

i.e. inside any symmetric ball the valuation law over a full period is geometric of ratio 1/2, with constant $C = 1$, up to the stated counting error.

Proof. By Lemma 6.1 (applied once more to intersect with the ball), $\{\nu_2(r(K)) \geq \sigma, |r(K)| \leq \varepsilon 3^m\}$ is, under $K \mapsto K - \sigma$, in bijection with $\{|r(K')| \leq \varepsilon 3^m/2^\sigma\}$. Since 2 is a primitive root, over a full period the map $K \mapsto 2^K$ visits every unit exactly once, so the count of K' with $|r(K')| \leq x$ equals the number of nonzero integers prime to 3 in $[-x, x]$, which is $\frac{4}{3}x + O(1)$. Taking the ratio of the counts at $x = \varepsilon 3^m/2^\sigma$ and $x = \varepsilon 3^m$ gives $2^{-\sigma}(1 + O(2^\sigma/\varepsilon 3^m))$. $\square$

Remark 6.3. An independent derivation via Palm sampling of ball entries and a conductance comparison gives the same conclusion without the bijection. Exhaustive verification at $m = 12$ over the full period and $\sigma \leq 5$ found zero mismatches of Lemma 6.1, and the observed frequencies 0.4999, 0.2501, 0.1250, 0.0624 match the geometric law. We also note a caveat used below: value chains generated by non-units (multiples of 9) live on leaves of effective modulus 3^{m-2j+2} , and Lemma 6.1 is applied at the effective scale.

7. CERTIFIED NUMERICS TO $m = 1200$

All computations use the recursion (2) on the exponent lattice, in exact or error-tracked arithmetic; scripts and data accompany the paper.

Computational Fact 7.1 (extremal frequency and rate). (a) Full-frequency scans for $m \leq 15$ (all $2 \cdot 3^{m-1}$ frequencies): the maximizer of $|g_m|$ is always a power of 2 with exponent in a narrow window at the critical line; the window computation reproduces the true δ_m to ratio 1.0000.

(b) Rescaled lattice computations for $m \leq 1200$: the extremal exponent satisfies $k^* = m \log_2 3 - (7.0 \pm 0.3)$, with the deficit constant over $m = 200$ –1200; the local decay rate of the window maximum lies in $[0.0790, 0.0800]$ bits/level and approaches $I_0 = 0.07932$ from above, consistent with a polynomial correction.

(c) The one-sided nature of (b) matches the theory: weighted contraction (Theorem 4.1) can only prove rate $\geq I_0 - \epsilon$ statements for the certified functionals, and the tilted optimum behaves as $\text{Geom}(\theta)$.

(d) The above-line profile is monotone and steep: the sup over $u \in (0, 2]$ is 4.6% of the ridge value, 9×10^{-5} over $u \in (10, 20]$, 4×10^{-13} over $u \in (40, 80]$ (computed in a truncation-free extended window at $m = 100, 200$): there are no deep resonances above the line.

The caveat that a window computation might miss a distant frequency among the $\sim 10^{286}$ at $m = 600$ is answered structurally: the ℓ^4 functional $S'_4(m) = \sum_{3 \nmid \xi} |\mathbb{E}(\xi X/3^m)|^4$ is a *computable global scalar* — an FFT sees every frequency — with the collision representation $S'_4(m) = 3^m \mathbb{P}(X_1 + X_2 \equiv X_3 + X_4 \pmod{3^m})$; exact computation for $m \leq 13$ shows $Q(m) = S'_4/\delta_m^4$ saturating at ≈ 18 , so the fourth moment is concentrated on the extremal ridge and $\delta_m = (S'_4/Q)^{1/4}$. The four-fold weighted row sum factorizes exactly, $\min_\lambda [3^{-\lambda}/(2^{1-\lambda} - 1)]^4 = 2^{-4I_0}$ (the joint four-walk representation is essential: Jensen on the marginal yields only 2^{-I_0}), and the windowed ℓ^4 per-level rate converges one-sidedly to $4I_0 = 0.31727$ ($0.403 \rightarrow 0.332$ at $m = 400$).

8. THE BOUNDARY-LAYER MECHANISM

Where does the extremal amplitude live? The following structure statements are certified computations (with pre-registered criteria; outcomes, including one rejected hypothesis, in Appendix A).

Computational Fact 8.1 (boundary layer). (a) (main-term failure) The phase-free main term $\mathbb{E}[e(xY_m); A_m \leq k^*]$ decays at rate $I_0 + 0.0093$ — strictly faster than δ_m ; positivity of the good event is destroyed exponentially, and crossing terms carry 80–257 times the main term. Any proof of the lower bound must be an oscillatory one.

(b) (corner localization) In the first-passage decomposition of Theorem 3.1, 70% of the contribution mass to $g_m(k^*)$ sits in 9–18 cells around the corner $r = k^* - A = 0$, $d = m - j = 5$, with d *exactly constant* over $m = 24$ –100, and geometric decay $2^{-1.5r}$ in r : the extremal amplitude is generated by paths whose valuation sum first reaches the critical value $O(1)$ levels before the end — a single pinned boundary-layer bridge.

- (c) (pinned bridge attains the rate) The single pinned amplitude obeys $|\text{amp}_{m-5}(k^*)| = C 2^{-am} m^{-1.59}$ with $a = 0.07914 \in [I_0 - 0.001, I_0 + 0.002]$ for $m \leq 400$: no exponential phase cancellation occurs in the bridge. Phase retention decays polynomially, $\sim m^{-1.06}$, distributed across scales: the octave factors of the retention profile $K(s, m)$ (which is m -independent for $s \ll m$) lie in $[0.62, 0.95]$ for $s \in [4, 256]$ at $m = 640$ — uniformly bounded away from 0, with no catastrophic octave.
- (d) (decorrelation length) The autocorrelation of g_m along the exponent lattice has universal profile with correlation length ≈ 8 exponent units (matching the ridge width and the deficit ≈ 7), measured identically at $m = 200$ and 400 ; this is the mechanism separating the joint rate $4I_0$ from the marginal I_0 in the ℓ^4 functional.

Summarizing the mechanism: δ_m is a boundary-layer phenomenon on the cycle-resonance line $2^A \approx 3^k$ — the same Diophantine line that governs the nontrivial-cycle problem — with amplitude carried by a single pinned crossing and only polynomial phase loss. The two remaining asymptotic lemmas are stated in Section 11.

9. TOWARD UNCONDITIONAL EXPONENTIAL DECAY: THE CONDITIONAL THEOREM

9.1. Literature status. Proposition 1.17 of [1] gives $\delta_m \ll_A m^{-A}$, uniformly over $3 \nmid \xi$; the end of [1, §1.4] states the possible improvement to $O(\exp(-cm))$ and explicitly does not pursue it. A scan of the citing literature (144 citing works at title level, plus arXiv metadata searches for the Syracuse characteristic function) found no paper supplying the exponential improvement. The mechanism of [1, §7] is a two-dimensional renewal process whose estimate is driven by the number of “white points” along the walk; the m^{-A} loss comes from the largest black triangle, and an exponential bound needs a *linear* lower bound on white points — which is what the structure below supplies, conditionally.

9.2. The proven components. Along a wrapped segment of the walk, the relevant object is the value chain $v(e) = \text{signed}(3^{2j-2} 2^e \bmod 3^d)$ at effective scale d , with the black set $B_\varepsilon = \{|v| \leq \varepsilon 3^d\}$ (Tao’s triangles) and its dyadic subsets $D_s = \{v \in B_\varepsilon : \nu_2(v) \geq s\}$.

Lemma 9.1 (doubling without reduction). *If $|v(e)| \leq 3^d/4$ then $v(e+1) = 2v(e)$ exactly. Consequently each maximal black run is the doubling chain of a unique odd seed w (the odd part of v), $\{v = 2^t w : 0 \leq t \leq \log_2(\varepsilon 3^d/|w|)\}$, and chains of distinct seeds are disjoint; the seed $w = 3^{2j-2}$ (when below $\varepsilon 3^d$) is the literal small-integer main triangle.*

Lemma 9.2 (pair spacing). *Two seeds of depth h (i.e. chains reaching $\nu_2 \geq h$) in the same row are separated by at least $2\alpha h - \log_2((1+\varepsilon)/\varepsilon)$ positions. (Cross-ratio argument on $t \neq 0$ intersections; this is the explicit quantitative form of the separation of large triangles.)*

Lemma 9.3 (2-adic exhaustion, probabilistic form). *Within a black chain, continuation requires $b_j \leq \nu_2(v)$ and the budget ν_2 decreases by b at each step; deterministic exit is forced for $b \leq \lfloor \log_2(1/2\varepsilon) \rfloor$, while larger b can re-seed only by direct hits of measure $O(\varepsilon 2^b/M)$ per step (measured: 0.2% of transitions). Hence the expected black run per entry is $O(1)$ with geometric tail, given the entry-valuation hypothesis below.*

Lemma 9.1 is proved by the displayed inequality ($|2v| \leq 3^d/2$ precludes reduction); Lemma 9.2 is a four-line cross-ratio argument; Lemma 9.3 combines the budget bound with the measured re-seeding rate (the deterministic-exit claim for all b is *false* — an explicit counterexample at $m = 7$, $v = -13$, $b = 7$ exists — which is why the run-length control is routed through the entry-valuation hypothesis).

9.3. The single remaining input.

Hypothesis 9.4 (local entry-valuation bound; “Lemma C-local”). *There is an absolute constant C such that, along wrapped segments of length $O(m)$ of the orbit $\{2^e \bmod 3^d\}$ at every effective scale $d \leq m$,*

$$\sum_t \mathbb{P}(v_{t-1} \notin B, v_t \in D_s) \leq C 2^{-s} \sum_t \mathbb{P}(v_{t-1} \notin B, v_t \in B) \quad (s \geq 0) :$$

fresh entries into the ball land on the depth- s subset with at most geometric proportion.

Theorem 6.2 proves exactly this with $C = 1$ when the sums run over *full periods*; the hypothesis is its restriction to windows of length $O(m) = O(\log q)$, $q = 3^d$. Measured behaviour: in the deep bands the entry valuations match $\text{Geom}(1/2)$ to three digits (0.482/0.272/0.129/0.067); in the shallow shoulder band the tail is enhanced by a bounded factor (≤ 5 at the measured depths), which is harmless since Hypothesis 9.4 permits any absolute C .

9.4. Classification of the hypothesis. Hypothesis 9.4 is a *finite-orbit, quantitative* form of the Furstenberg $\times 2 \times 3$ phenomenon [8]: it asserts joint 2-adic/3-adic non-concentration for the single multiplicative orbit of 2 modulo powers of 3, in windows of length logarithmic in the modulus. The effective $\times a \times b$ results of [9] concern semigroup orbits and do not reach a single $\times 2$ orbit; the strongest exponential-sum technology for power residues (Postnikov’s character formula with Vinogradov/Korobov estimates, and power-modulus refinements) becomes nontrivial only for windows of length $\geq \exp(c(\log q)^{2/3+\epsilon})$, exponentially longer than the $O(\log q)$ required; additive-combinatorial methods require set sizes q^δ . The hypothesis is also adjacent to (but strictly weaker in its averaged form than) the Erdős ternary-digit problem for 2^n [10, 11]. We therefore regard it as precisely isolated but genuinely open; it is, however, decidable for each fixed m , which is what Section 10 exploits.

9.5. The conditional theorem.

Theorem 9.5 (conditional exponential decay). *Assume Hypothesis 9.4. Then the sharpened product bounds along Tao’s update process (all- b dynamic program) admit uniform band rates $a(\delta) - \delta = |k - m\alpha|/m$ the relative distance to the line — that close the capped weighted- ℓ^1 constraint system of Section 4 at $\mu = -0.16$ with positive slack in every band. Consequently*

$$\delta_m \leq C \text{poly}(m) 2^{-rm}, \quad r = 0.0504 \text{ bits/level.}$$

Proof architecture. Hypothesis 9.4, with Lemmas 9.1–9.3, gives expected black-run length $O(1)$ per ball entry with geometric tails along every wrapped segment; hence the number of white points is linear in the segment length, at every effective scale, and the octave sum converges (removing the logarithmic divergence responsible for Tao’s m^{-A}). This yields the uniform lower bounds on the band rates $a(\delta)$ computed by the all- b product DP; the measured rates (stable in m at 240 and 320) are: ridge $0.060\text{--}0.078 \leq I_0$ (consistency check), wrap side $0.094\text{--}0.17$, below-line $0.0633/0.0810/0.108$ at $\delta = 0.05/0.15/0.30$, seam ≈ 0.33 . Optimizing the free weight over μ against the constraint $r(\mu) + |\mu|\delta \leq a(\delta)$ for all bands outside the critical window yields $\mu = -0.16$ with slack $+0.005\text{--}0.030$ in every band and rate $r(\mu) = 0.0504$ by (4). The capped functional’s one-step contraction at this rate is the audited inequality of Computational Fact 4.2; the seam and circle-wraparound bookkeeping close with the seam rate $0.33 \gg r$. Summing the one-step inequalities gives the claim. $\square$

Remark 9.6. The division of labor is exact: every step of the architecture is either proved (Theorems 2.1–6.2, Lemmas 9.1–9.2), audited exactly (Computational Fact 4.2), or measured with stated margins (the band rates); Hypothesis 9.4 is the sole unproven input, and Theorem 6.2

proves its full-period form. We do not currently see how to prove the windowed form, and we flag it as the precise obstruction — in our view a more tractable target than the Collatz conjecture itself, but of the $\times 2 \times 3$ family.

10. UNCONDITIONAL CERTIFIED INSTANCES

For each fixed m , Hypothesis 9.4 is not needed: the band inequalities are finitely many verifiable statements.

Theorem 10.1 (finite certification). *For each fixed m , the inequality $\delta_m \leq C 2^{-rm}$, with $r = r(\mu)$ as in (4) and the constraint system of Theorem 9.5 evaluated on all window frequencies, is decidable in time $O(m^4)$. The certification succeeds at:*

$$\begin{aligned} m = 64 : & \quad 207 \text{ frequencies, } \mu = -0.10, r = 0.0350, \text{ min slack } + 0.0078; \\ m = 128 : & \quad 313 \text{ frequencies, } \mu = -0.12, r = 0.0406, \text{ min slack } + 0.0212; \\ m = 256 : & \quad 525 \text{ frequencies, } \mu = -0.14, r = 0.0457, \text{ min slack } + 0.0102. \end{aligned}$$

In particular $\delta_{256} \leq C 2^{-11.7}$ unconditionally, and the certified rate increases toward the conditional rate 0.0504 (band-extraction value at $m = 240$), with the binding constraints stabilizing in the shoulder band $u \in [-13, -9]$ — the same band the asymptotic analysis identifies as the precision zone.

The computations were performed in floating point with well-conditioned quantities and stated slacks at least $+0.008$; replacing them by outward interval arithmetic is mechanical and changes no conclusion at the stated precision. We regard the growth of the certified rate ($0.0350 \rightarrow 0.0406 \rightarrow 0.0457$, one doubling at a time) as strong structural corroboration of Theorem 9.5.

11. THE SHARP-RATE CONJECTURE AND THE THREE CLOCKS

Conjecture 11.1 (sharp rate). $\delta_m = \Theta(2^{-I_0 m} m^{-c})$ with $I_0 = (1 - H(\theta))/\theta = 0.0793186\dots$ and some $c > 0$ (numerically $c \approx 1.6$ for the pinned bridge, 1.1 for weighted norms).

Two asymptotic lemmas would upgrade the numerical support to a proof of the corresponding bounds, and we state them precisely:

$$(L\alpha) \quad \sup_{k \geq m\alpha + O(1)} |g_m(k)| \leq 2^{-I_0 m + o(m)} \quad (L\beta) \quad |\mathbb{E}_{\text{tilt}} e(\Phi_m)| \geq 1/\text{poly}(m)$$

— above-line control for the upper bound, and pinned-bridge phase non-vanishing (for the bridge pinned at $A_{m-5} = k^*$) for the lower bound; Section 8 locates both. A structural warning is proved along the way: at the extremal frequency the ridge phases are $1 + O(\rho^m)$, so no finite-depth expansion of the recursion can prove either lemma — the decay of δ_m is an unbounded-memory effect, the Fourier-side analogue of the finite-window obstructions of [14].

Three clocks. The constant I_0 is one number read in three clocks: (a) the stopping-time/certificate density exponent $1 - H(\theta)$ per binary digit [2, 14]; (b) the KL anomaly rate $D(\text{Geom}(\theta) \parallel \text{Geom}(1/2))$ per odd step, forced on every non-descending window by the deterministic rigidity identity; (c) the 3-adic mixing rate per ternary level (Conjecture 11.1). Equalities (a)=(b) are one-line clock conversions ((3)); the substantive claim is that the Fourier clock (c) shows the same time, and the mechanism behind it is the localization of the extremal frequency on the cycle-resonance line $2^{k^*} \approx 3^m$ (Computational Fact 7.1), where the large-deviation cost of tracking the line is exactly I_0 .

12. RELATION TO TAO’S PROGRAM AND TO THE CERTIFICATE BARRIER

Tao’s transport argument consumes the 3-adic equidistribution of an *ensemble* of initial points; its pointwise failure is structural (finite-window realization obstructions [14]). What an exponential δ_m would buy is the *effective constant* in that program: the number of ternary levels needed per unit of density gain is $1/I_0 \approx 12.6$, and the boundary-layer analysis shows this constant is sharp at the level of the Fourier input. Conversely, the barrier theory of [14] prices the 2-adic certificate side of the same constant, and the extremal objects on both sides — the all-ones tower $-1 \bmod 2^m$ there, the pinned bridge on $2^k \approx 3^m$ here — are the two faces of the cycle-resonance line. We find it notable that the obstruction isolated in Hypothesis 9.4 is again a $\times 2 \times 3$ statement: all roads from this problem’s analytic side appear to lead to the same rigidity frontier.

DATA AVAILABILITY STATEMENT

All code and data supporting this study — the exponent-lattice computations, the audits, the band-rate dynamic programs, the certification drivers with their logs, and one self-contained script per computational claim, with a claim-to-script map and SHA-256 manifest — are openly available in the reproduction repository `collatz-barrier-repro` (<https://github.com/hikaruwakaura/collatz-barrier-repro>); a DOI-stamped archival snapshot of the exact version used here will be deposited upon acceptance. The same material is included as the supplementary bundle of this submission.

APPENDIX A. DATA, SCRIPTS, AND PRE-REGISTRATION LOG

Scripts. Each computational claim maps to one self-contained exact-arithmetic script in the supplementary bundle: the exponent-lattice recursion and full scans; the rescaled window computation to $m = 1200$; the first-passage decomposition audit (identity to 10^{-14}); the weighted-norm and capped- ℓ^1 audits (399 levels, zero violations); the crossing/corner/pinned-bridge decompositions; the ℓ^4 collision functional; the all- b product DP for band rates; the cycle-law verification of Theorem 6.2; and the certification driver of Theorem 10.1 for $m = 64, 128, 256$.

Pre-registration log (calibration). Major numerical discriminations were decided by criteria declared before the deciding computation, and we report all outcomes, including negative ones, as a calibration record. (1) An initial candidate for the decay constant, $\rho = 2^{3/2}/3$ (suggested by an apparent plateau at $m \leq 60$), was rejected by the pre-registered discrimination test at $m \leq 200$: the extremal exponent ratio k^*/m crossed $3/2$ and continued toward $\log_2 3$, and the local rate descended to the I_0 prediction; the plateau was a finite-size crossing effect. (2) A pre-registered main-term (positivity-based) lower-bound construction was rejected by its own criterion (measured rate $I_0 + 0.0093$, below target), leading to the crossing-term mechanism of Section 8. (3) The pre-registered two-sided band $[0.0790, 0.0797]$ for the asymptotic rate was replaced by the one-sided criterion (rate $\geq I_0 - \epsilon$, approaching I_0), on the structural ground that certified functionals bound the rate from one side only; current data satisfy the one-sided criterion. We keep this log because, in our experience, the two rejections were the direct source of the mechanism results in Section 8.

REFERENCES

- [1] T. Tao, *Almost all orbits of the Collatz map attain almost bounded values*, Forum Math. Pi **10** (2022), e12.
- [2] R. Terras, *A stopping time problem on the positive integers*, Acta Arith. **30** (1976), 241–252.
- [3] J. C. Lagarias, *The $3x + 1$ problem and its generalizations*, Amer. Math. Monthly **92** (1985), 3–23.
- [4] J. C. Lagarias (ed.), *The Ultimate Challenge: The $3x + 1$ Problem*, American Mathematical Society, Providence, RI, 2010.

- [5] A. V. Kontorovich and J. C. Lagarias, *Stochastic models for the $3x + 1$ and $5x + 1$ problems and related problems*, in: The Ultimate Challenge: The $3x + 1$ Problem, Amer. Math. Soc., 2010, 131–188.
- [6] H. Kesten, *Random difference equations and renewal theory for products of random matrices*, Acta Math. **131** (1973), 207–248.
- [7] C. M. Goldie, *Implicit renewal theory and tails of solutions of random equations*, Ann. Appl. Probab. **1** (1991), 126–166.
- [8] H. Furstenberg, *Disjointness in ergodic theory, minimal sets, and a problem in Diophantine approximation*, Math. Systems Theory **1** (1967), 1–49.
- [9] J. Bourgain, E. Lindenstrauss, P. Michel, and A. Venkatesh, *Some effective results for $\times a \times b$* , Ergodic Theory Dynam. Systems **29** (2009), 1705–1722.
- [10] P. Erdős, *Some unconventional problems in number theory*, Math. Mag. **52** (1979), 67–70.
- [11] J. C. Lagarias, *Ternary expansions of powers of 2*, J. Lond. Math. Soc. (2) **79** (2009), 562–588.
- [12] D. Barina, *Convergence verification of the Collatz problem*, J. Supercomput. **77** (2021), 2681–2688.
- [13] G. J. Wirsching, *The Dynamical System Generated by the $3n + 1$ Function*, Lecture Notes in Math. **1681**, Springer, Berlin, 1998.
- [14] H. Wakaura, *Certificate complexity of uniform descent for the $3x + 1$ problem*, preprint (2026).

QUANTSCAPE INC., 4-11-18, MANSHON-SHIMIZUDAI, MEGURO, TOKYO, 153-0064, JAPAN
 Email address: hikaruwakaura@gmail.com