

# Lab 6 - La Quinta is Spanish for next to Denny's, Pt. 1

## Visualizing spatial data

Walaa Ali

Student ID: 2201002583

Load the needed packages. Note: dsbox is not yet on CRAN. For now, you need to install it before you load the library. Uncomment, by deleting # from, the two lines of code below.

```
install.packages("devtools")
devtools::install_github("rstudio-education/dsbox")
```

Load the libraries here.

```
library(tidyverse)
library(dsbox)
```

To help with our analysis we will also use a dataset on US states, which is located in your repository's data folder.

```
states <- read_csv("data/states.csv")
```

## Exercises

1. What are the dimensions of the Denny's dataset? (Hint: Use inline R code and functions like `nrow` and `ncol` to compose your answer.) What does each row in the dataset represent? What are the variables?

row = 1643 , column = 6 the row it's about the restaurant address (more information about it ) column ( address , city , state , zip , longitude , latitude)

```
dim(dennys)
```

```
## [1] 1643    6
```

```
head(dennys)
```

```
## # A tibble: 6 x 6
##   address          city      state zip  longitude latitude
##   <chr>          <chr>    <chr> <chr>    <dbl>    <dbl>
## 1 2900 Denali    Anchorage AK   99503   -150.    61.2
## 2 3850 Debarr Road Anchorage AK   99508   -150.    61.2
## 3 1929 Airport Way Fairbanks AK   99701   -148.    64.8
## 4 230 Connector Dr Auburn    AL   36849   -85.5    32.6
## 5 224 Daniel Payne Drive N Birmingham AL   35207   -86.8    33.6
## 6 900 16th St S, Commons on Gree Birmingham AL   35294   -86.8    33.5
```

The 'dennys' dataset has 1,643 rows and 6 columns. Each row represents location information about a denny's restaurant.

2. What are the dimensions of the La Quinta's dataset? What does each row in the dataset represent? What are the variables?

```
row = 909 , column = 6
```

the row it's about the hotels address (more information about it ) column ( address , city , state , zip , longitude , latitude)

```
dim(laquinta)
```

```
## [1] 909 6
```

```
head(laquinta)
```

```
## # A tibble: 6 x 6
##   address                city      state zip longitude latitude
##   <chr>                 <chr>   <chr> <chr>   <dbl>   <dbl>
## 1 793 W. Bel Air Avenue  "\nAberdeen" MD    21001   -76.2    39.5
## 2 3018 CatClaw Dr       "\nAbilene"  TX    79606   -99.8    32.4
## 3 3501 West Lake Rd     "\nAbilene"  TX    79601   -99.7    32.5
## 4 184 North Point Way   "\nAcworth"  GA    30102   -84.7    34.1
## 5 2828 East Arlington Street "\nAda"     OK    74820   -96.6    34.8
## 6 14925 Landmark Blvd   "\nAddison"  TX    75254   -96.8    33.0
```

This dataset has 909 rows and 6 columns. Each row represents location information about laquinta's hotel.

Knit, commit, and push your changes to GitHub with an appropriate commit message. Make sure to commit and push all changed files so that your Git pane is cleared up afterwards.

3. Add a country variable to the Denny's and Laquinta's datasets and set all observations equal to "United States". Remember, you can use the `mutate` function for adding a variable. Make sure to save the result of this as `dn` and `lq`, respectively, so that the stored data frame contains the new variable going forward.

For Denny's

```
dn <- dennys %>%
  mutate(country = "United States")
```

For La Quinta

```
lq <- laquinta %>%
  mutate(country = "United States")
```

Knit, commit, and push your changes to GitHub with an appropriate commit message. Make sure to commit and push all changed files so that your Git pane is cleared up afterwards.

4. Which states have the most and fewest Denny's locations? What about La Quinta? Is this surprising? Why or why not?

the most = CA and the fewest = DE

For Denny's - Most = CA

La Quinta the most = TX and the fewest = AG

```
dn %>%
  count(state) %>%
  arrange(desc(n))
```

```
## # A tibble: 51 x 2
##   state      n
##   <chr> <int>
## 1 CA      403
## 2 TX      200
```

```
## 3 FL      140
## 4 AZ      83
## 5 IL      56
## 6 NY      56
## 7 WA      49
## 8 OH      44
## 9 MO      42
## 10 PA     40
## # ... with 41 more rows
```

CA, which stands for California has the most Denny's locations. It is not surprising because it is the largest state in the US by population.

For Denny's - Fewest = DE

```
dn %>%
  count(state) %>%
  arrange(n)
```

```
## # A tibble: 51 x 2
##   state      n
##   <chr> <int>
## 1 DE      1
## 2 DC      2
## 3 VT      2
## 4 AK      3
## 5 IA      3
## 6 NH      3
## 7 SD      3
## 8 WV      3
## 9 LA      4
## 10 MT     4
## # ... with 41 more rows
```

DE, Delaware has the fewest locations. It is not surprising because it is one of the smallest US states by population.

For La Quinta - Most

```
lq %>%
  count(state) %>%
  arrange(desc(n))
```

```
## # A tibble: 59 x 2
##   state      n
##   <chr> <int>
## 1 TX     237
## 2 FL     74
## 3 CA     56
## 4 GA     41
## 5 TN     30
## 6 OK     29
## 7 LA     28
## 8 CO     27
## 9 NM     19
## 10 NY     19
## # ... with 49 more rows
```

TX, Texas has the most La Quinta locations. The surprising thing is that Texas has more locations than California. This despite the fact that California has more population and has stronger economy than Texas

For La Quinta - Fewest

```
lq %>%
  count(state) %>%
  arrange(n)
```

```
## # A tibble: 59 x 2
##   state      n
##   <chr> <int>
## 1 AG         1
## 2 ANT         1
## 3 BC          1
## 4 CH          1
## 5 FM          1
## 6 ME          1
## 7 ON          1
## 8 QR          1
## 9 SL          1
## 10 VE         1
## # ... with 49 more rows
```

#### 10 states with only one La Quinta Locations

5. Which states have the most Denny's locations per thousand square miles? What about La Quinta?

`\textcolor{red}{We are interested in the number of locations per state, which we could find in Denny's or La Quinta's datasets, and the area of each state, which is only available in the states dataset, which we imported earlier from the Data folder using the code states <- read_csv("data/states.csv").}`

Joining Denny's and states

```
dn %>%
  count(state) %>%
  inner_join(states, by = c("state" = "abbreviation"))
```

```
## # A tibble: 51 x 4
##   state      n name          area
##   <chr> <int> <chr>          <dbl>
## 1 AK         3 Alaska      665384.
## 2 AL         7 Alabama      52420.
## 3 AR         9 Arkansas      53179.
## 4 AZ        83 Arizona     113990.
## 5 CA       403 California   163695.
## 6 CO        29 Colorado     104094.
## 7 CT        12 Connecticut    5543.
## 8 DC         2 District of Columbia    68.3
## 9 DE         1 Delaware       2489.
## 10 FL       140 Florida     65758.
## # ... with 41 more rows
```

Joining La Quinta's and states

```
lq %>%
  count(state) %>%
  inner_join(states, by = c("state" = "abbreviation"))
```

```
## # A tibble: 48 x 4
##   state      n name      area
##   <chr> <int> <chr>    <dbl>
## 1 AK         2 Alaska  665384.
## 2 AL        16 Alabama  52420.
## 3 AR        13 Arkansas  53179.
## 4 AZ        18 Arizona  113990.
## 5 CA        56 California 163695.
## 6 CO        27 Colorado  104094.
## 7 CT         6 Connecticut  5543.
## 8 FL        74 Florida   65758.
## 9 GA        41 Georgia   59425.
## 10 IA         4 Iowa     56273.
## # ... with 38 more rows
```

We need to add an identifier variable which we'll call `establishment` and set the value to "Denny's" and "La Quinta" for the `dn` and `lq` data frames, respectively.

```
dn <- dn %>%
  mutate(establishment = "Denny's")
```

```
lq <- lq %>%
  mutate(establishment = "La Quinta")
```

Since the two data frames (`dn` and `lq`) have the same columns, we can easily bind them with the `bind_rows` function:

```
dn_lq <- bind_rows(dn, lq)
```

We can plot the locations of the two establishments using a scatter plot, and color the points by the establishment type. Note that the latitude is plotted on the x-axis and the longitude on the y-axis.

```
ggplot(dn_lq, mapping = aes(x = longitude, y = latitude, color = establishment)) +
  geom_point()
```



The graph figure shows that indeed there are many La Quinta hotels that located next to Denny's restaurants. We could also see from the figure that there is the presence of the two chains across the US as the figure almost depict the exact map of the country. The figure also indicates the presence of these chains in Alaska (top-left), Hawaii (bottom-left), and some cities in Central America(bottom-right)

*Knit, commit, and push your changes to GitHub with an appropriate commit message. Make sure to commit and push all changed files so that your Git pane is cleared up afterwards and review the md document on GitHub to make sure you're happy with the final state of your work.*