

Mathematical Methods

# Problem Sheet 3

Partial Differential Equations

Xin, Wenkang

June 19, 2024

# Partial Differential Equations

## 1 Laplace equation in two dimensions

(a)

(b)

(c) Consider the Laplacian in polar coordinates:

$$\nabla^2 = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2}{\partial \phi^2} \quad (1)$$

Assuming a solution of the form  $V(r, \phi) = R(r)\Phi(\phi)$ , we have the separated equations:

$$\begin{aligned} \Phi'' + k^2 \Phi &= 0 \\ r^2 R'' + r R' - k^2 R &= 0 \end{aligned} \quad (2)$$

First consider non-zero  $k$ . The first equation has the solution:

$$\Phi(\phi) = A \cos k\phi + B \sin k\phi \quad (3)$$

The natural boundary condition  $\Phi(0) = \Phi(2\pi)$  demands  $k \in \mathbb{Z}$ . The second equation can be solved with the Ansatz  $R(r) = r^\lambda$  that leads to  $\lambda = \pm k$  and the general solution:

$$R(r) = Cr^k + Dr^{-k} \quad (4)$$

In the case of  $k = 0$ , we have  $\Phi$  is some constant and  $R(r) = C \ln r + D$ . The most general solution to the Laplace equation is then:

$$V(r, \phi) = \frac{a_0}{2} + \frac{\tilde{a}_0}{2} \ln r + \sum_{k=1}^{\infty} (a_k r^k + \tilde{a}_k r^{-k}) \cos k\phi + \sum_{k=1}^{\infty} (b_k r^k + \tilde{b}_k r^{-k}) \sin k\phi \quad (5)$$

where we have rewritten the arbitrary constants and taken the general solution as a sum of the solutions for  $k \in \mathbb{Z}$ .

(d) For the given boundary condition, we need:

$$\frac{a_0}{2} + \frac{\tilde{a}_0}{2} \ln a_{\pm} + \sum_{k=1}^{\infty} (a_k a_{\pm}^k + \tilde{a}_k a_{\pm}^{-k}) \cos k\phi + \sum_{k=1}^{\infty} (b_k a_{\pm}^k + \tilde{b}_k a_{\pm}^{-k}) \sin k\phi = g_{\pm}(\phi) \quad (6)$$

This is a Fourier series of  $g_{\pm}(\phi)$  if we identify the coefficients:

$$c_{0,\pm} + \sum_{k=1}^{\infty} (c_{k,\pm} \cos k\phi + d_{k,\pm} \sin k\phi) = g_{\pm}(\phi) \quad (7)$$

where the coefficients are given by:

$$c_{k,\pm} = a_k a_{\pm}^k + \tilde{a}_k a_{\pm}^{-k} \quad \text{and} \quad d_{k,\pm} = b_k a_{\pm}^k + \tilde{b}_k a_{\pm}^{-k} \quad (8)$$

The coefficients can be solved by the orthogonality of the trigonometric functions:

$$\begin{aligned} c_{k,\pm} &= \frac{1}{\pi} \int_0^{2\pi} g_{\pm}(\phi) \cos k\phi \, d\phi \\ d_{k,\pm} &= \frac{1}{\pi} \int_0^{2\pi} g_{\pm}(\phi) \sin k\phi \, d\phi \end{aligned} \quad (9)$$

These leads to equations that can be solved for  $a_k$ ,  $\tilde{a}_k$ ,  $b_k$ , and  $\tilde{b}_k$ .

•

## 2 Laplace equation in three dimensions

(a) Without loss of generality let the y-axis align with the vector pointing from the origin to the charge  $q$  at  $(0, y, 0)$ . We seek another charge  $q'$  at  $(0, d, 0)$  such that the potential at the ball surface is zero. A direct calculation gives:

$$q' = -\frac{b}{y}q \quad (10)$$

and:

$$d = \frac{b^2}{y} \quad (11)$$

Let us define  $\mathbf{d} = b^2 \mathbf{y}/y^3$ , then the solution to the Laplace equation is:

$$V = q \left( \frac{1}{|\mathbf{r} - \mathbf{y}|} - \frac{b/y}{|\mathbf{r} - \mathbf{d}|} \right) \quad (12)$$

Generalising to the case of  $n$  charges  $q_i$  each at  $\mathbf{y}_i$ , we introduce image charges  $q'_i = -bq_i/y_i$  at  $\mathbf{d}_i = b^2 \mathbf{y}_i/y_i^3$  and the solution is:

$$V = \sum_{i=1}^n q_i \left( \frac{1}{|\mathbf{r} - \mathbf{y}_i|} - \frac{b/y_i}{|\mathbf{r} - \mathbf{d}_i|} \right) \quad (13)$$

(b) Consider the Laplacian in spherical coordinates:

$$\nabla^2 = \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \quad (14)$$

Assuming a solution of the form  $V(r, \theta, \phi) = R(r)\Theta(\theta)\Phi(\phi)$ , we have the separated equations:

$$\begin{aligned} \frac{d}{dr} \left( r^2 \frac{dR}{dr} \right) &= l(l+1)R \\ \frac{d^2 \Phi}{d\phi^2} &= -m^2 \Phi \\ \frac{1}{\sin \theta} \frac{d}{d\theta} \left( \sin \theta \frac{d\Theta}{d\theta} \right) + \left[ l(l+1) - \frac{m^2}{\sin^2 \theta} \right] \Theta &= 0 \end{aligned} \quad (15)$$

The radial solution can be solved with the Ansatz  $R(r) = r^\lambda$  that leads to  $\lambda = l$  or  $\lambda = -l - 1$ . The solution is then:

$$R(r) = Ar^l + Br^{-l-1} \quad (16)$$

The solution to  $\Phi$  is:

$$\Phi(\phi) = e^{im\phi} \quad (17)$$

where we have dropped the arbitrary constant and the natural boundary condition  $m \in \mathbb{Z}$ .

The  $\Theta$  equation is the associated Legendre equation with the solution:

$$\Theta(\theta) = P_l^m(\cos \theta) \quad (18)$$

Combining the solutions, we have:

$$V(r, \theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l (A_{lm}r^l + B_{lm}r^{-l-1}) Y_{lm}(\theta, \phi) \quad (19)$$

where  $Y_{lm}(\theta, \phi) = P_l^m(\cos \theta)e^{im\phi}$  are called the spherical harmonics.

If azimuthal symmetry can be assumed, then  $m = 0$  and the solution is:

$$V(r, \theta) = \sum_{l=0}^{\infty} (A_l r^l + B_l r^{-l-1}) P_l(\cos \theta) \quad (20)$$

(c) For the boundary condition  $V(a, \theta, \phi) = \phi_0(1 + \cos \theta)$ , we have azimuthal symmetry and may discard  $B_l$  terms as they diverge at  $r = 0$ . We need:

$$\sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta) = \phi_0(1 + \cos \theta) \quad (21)$$

Since  $P_l(\cos \theta)$  are orthogonal, we have  $A_0 = \phi_0$ ,  $A_1 = \phi_0/a$  and  $A_l = 0$  for  $l \geq 2$ . The solution is then:

$$V(r, \theta) = \phi_0 \left( 1 + \frac{r}{a} \cos \theta \right) \quad (22)$$

(d) For the boundary condition  $V(a, \theta, \phi) = \phi_0 \sin^2 \theta$ , we still have azimuthal symmetry and may discard  $A_l$  terms as they diverge at  $r \rightarrow \infty$ . We need:

$$\sum_{l=0}^{\infty} B_l r^{-l-1} P_l(\cos \theta) = \phi_0 \sin^2 \theta = \phi_0(1 - \cos^2 \theta) \quad (23)$$

Comparing the coefficients, we have  $B_0 = 2\phi_0 b/3$ ,  $B_1 = 0$ ,  $B_2 = -2\phi_0 b^3/3$  and  $B_l = 0$  for  $l \geq 3$ . The solution is then:

$$V(r, \theta) = \phi_0 \left[ \frac{2b}{3r} - \frac{b^3}{3r^3} (3 \cos^2 \theta - 1) \right] \quad (24)$$

•

### 3 Multiple expansion

(b) The first few spherical harmonics are:

$$\begin{aligned} Y_0^0 &= \sqrt{\frac{1}{4\pi}} \\ Y_1^0 &= \sqrt{\frac{3}{4\pi}} \cos \theta \\ Y_1^1 &= -\sqrt{\frac{3}{8\pi}} \sin \theta e^{i\phi} \\ Y_1^{-1} &= \sqrt{\frac{3}{8\pi}} \sin \theta e^{-i\phi} \end{aligned} \quad (25)$$

(c) First consider the case of  $l = 0$ . The term  $q_{00}$  can be written as:

$$q_{00} = \int_{\mathbb{R}^3} \sqrt{\frac{1}{4\pi}} \rho(\mathbf{r}') d^3 r' = \sqrt{\frac{1}{4\pi}} Q \quad (26)$$

so that the monopole term is:

$$4\pi \frac{q_{00}}{r} Y_0^0 = \frac{Q}{r} \quad (27)$$

Next consider the case of  $l = 1$ . The terms  $q_{10}$ ,  $q_{11}$ , and  $q_{1-1}$  can be written as:

$$q_{10} = \int_{\mathbb{R}^3} \sqrt{\frac{3}{4\pi}} \cos \theta' \mathbf{r}' \rho(\mathbf{r}') d^3 r' \quad (28)$$

•

## 4 Strings

(a) Consider the wave equation:

$$\frac{\partial^2 \psi}{\partial x^2} = \frac{\partial^2 \psi}{\partial t^2} \quad (29)$$

Assuming a solution of the form  $\psi(x, t) = X(x)T(t)$ , separation of variables leads to the general solution:

$$\psi(t, x) = \sum_{\omega} C_{\omega} \sin(\omega x + \phi_x) \sin(\omega t + \phi_t) \quad (30)$$

The boundary condition  $\psi(t, 0) = \psi(t, a) = 0$  demands  $\omega = n\pi/a$  for  $n \in \mathbb{Z}$  and  $\phi_x = 0$ . Indexing via  $n$ , the solution is:

$$\psi(t, x) = \sum_{n=1}^{\infty} C_n \sin\left(\frac{n\pi x}{a}\right) \sin\left(\frac{n\pi t}{a} + \phi_n\right) \quad (31)$$

(b) Given the initial condition  $\psi(0, x) = \psi_0 \sin(\pi x/a)$  and  $\dot{\psi}(0, x) = 0$ , we need:

$$\dot{\psi}(0, x) = \sum_{n=1}^{\infty} C_n \frac{n\pi}{a} \sin\left(\frac{n\pi x}{a}\right) \cos \phi_n = 0 \quad (32)$$

which gives  $\phi_n = \pi/2$ .

We may as well write the temporal part as cosine functions  $\cos(n\pi t/a)$  with the same coefficients. The position initial condition gives:

$$\psi(0, x) = \sum_{n=1}^{\infty} C_n \sin\left(\frac{n\pi x}{a}\right) = \psi_0 \sin\left(\frac{\pi x}{a}\right) \quad (33)$$

so that  $C_n = 0$  except for  $C_1 = \psi_0$ .

The solution is then:

$$\psi(t, x) = \psi_0 \sin\left(\frac{\pi x}{a}\right) \cos\left(\frac{\pi t}{a}\right) \quad (34)$$

(c) Given the initial condition  $\psi(0, x) = 0$  and  $\dot{\psi}(0, x) = \psi_0 \sin(2\pi x/a)$ , we need:

$$\psi(0, x) = \sum_{n=1}^{\infty} C_n \sin\left(\frac{n\pi x}{a}\right) \sin \phi_n = 0 \quad (35)$$

which gives  $\phi_n = 0$ .

The velocity initial condition gives:

$$\dot{\psi}(0, x) = \sum_{n=1}^{\infty} C_n \frac{n\pi}{a} \sin\left(\frac{n\pi x}{a}\right) = \psi_0 \sin\left(\frac{2\pi x}{a}\right) \quad (36)$$

so that  $C_n = 0$  except for  $C_2 = \psi_0 a/2\pi$ .

The solution is then:

$$\psi(t, x) = \frac{\psi_0 a}{2\pi} \sin\left(\frac{2\pi x}{a}\right) \cos\left(\frac{2\pi t}{a}\right) \quad (37)$$

(d) Consider the skewed triangle wave initial condition:

$$\psi(0, x) = \begin{cases} \frac{hx}{b} & 0 \leq x \leq b \\ \frac{h(a-x)}{a-b} & b \leq x \leq a \end{cases} \quad (38)$$

and  $\dot{\psi}(0, x) = 0$ . From the velocity initial condition, we write the temporal part as cosine functions:

$$\psi(t, x) = \sum_{n=1}^{\infty} C_n \sin\left(\frac{n\pi x}{a}\right) \cos\left(\frac{n\pi t}{a}\right) \quad (39)$$

Then from the position initial condition, we have a Fourier sine series with the coefficients given by:

$$\begin{aligned} C_n &= \frac{2}{a} \left[ \int_0^b \frac{hx}{b} \sin\left(\frac{n\pi x}{a}\right) dx + \int_b^a \frac{h(a-x)}{a-b} \sin\left(\frac{n\pi x}{a}\right) dx \right] \\ &= \frac{2ha^2}{b(a-b)(n\pi)^2} \sin\left(\frac{n\pi b}{a}\right) \end{aligned} \quad (40)$$

Consider  $C_n/h$  as a function of  $(b/a) \equiv \lambda$ :

$$\frac{C_n}{h} = \frac{2}{\lambda(1-\lambda)(n\pi)^2} \sin(n\pi\lambda) \quad (41)$$

•

## 5 Membranes

Consider the two dimensional wave equation:

$$\frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} = \frac{\partial^2 \psi}{\partial t^2} \quad (42)$$

Separation of variables leads to the general solution:

$$\psi = \sum_{\omega, k} C_{\omega, k} \sin(\omega t + \phi_t) \sin(kx + \phi_x) \sin(\sqrt{\omega^2 - k^2}y + \phi_y) \quad (43)$$

where we demand  $\omega^2 > k^2$ .

For the boundary conditions, we need  $\phi_x = 0$  and  $ka = n\pi$  for  $n \in \mathbb{Z}$ , which means  $k = n\pi/a$ . We also need  $\phi_y = 0$  and  $\sqrt{\omega^2 - k^2}b = m\pi$  for  $m \in \mathbb{Z}$ , which means  $\omega^2 = (m\pi/b)^2 + (n\pi/a)^2$ . Indexing via  $n$  and  $m$ , the general solution is:

$$\psi(x, y, t) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} C_{m,n} \sin\left(\frac{n\pi x}{a}\right) \sin\left(\frac{m\pi y}{b}\right) \sin(\omega_{m,n}t + \phi_{m,n}) \quad (44)$$

where we have defined the frequencies  $\omega_{m,n} = \sqrt{(m\pi/b)^2 + (n\pi/a)^2}$ .

For a square membrane with  $a = b$ , we have  $\omega_{m,n} = \pi\sqrt{m^2 + n^2}/a$ . The ratio of the two lowest frequencies is:

$$\frac{\omega_{1,1}}{\omega_{0,1}} = \sqrt{2} \quad (45)$$



(b) For a triangular membrane with  $a = b$ , we impose the further condition that  $\psi = 0$  along the line  $y = x$ . This means  $m = n$  and  $\omega_{m,n} = \pi\sqrt{2}m/a$ . The ratio of the two lowest frequencies is:

•

## 6 Eigenfunctions of the Laplacian

(a) Assuming a solution of the form  $V(r, \theta, \phi) = R(r)\Theta(\theta)\Phi(\phi)$ , with the eigenvalue problem  $-\nabla^2 V = EV$ , we have the separated equations:

$$\begin{aligned} \frac{d}{dr} \left( r^2 \frac{dR}{dr} \right) + Er^2 R &= l(l+1)R \\ \frac{d^2 \Phi}{d\phi^2} &= -m^2 \Phi \\ \frac{1}{\sin \theta} \frac{d}{d\theta} \left( \sin \theta \frac{d\Theta}{d\theta} \right) + \left[ l(l+1) - \frac{m^2}{\sin^2 \theta} \right] \Theta &= 0 \end{aligned} \quad (46)$$

The angular parts are the same as the homogeneous Laplace equation. The solutions are the spherical harmonics  $Y_{lm}(\theta, \phi) = P_l^m(\cos \theta)e^{im\phi}$ . The radial part satisfies the differential equation:

$$r^2 R'' + 2rR' + [Er^2 - l(l+1)]R = 0 \quad (47)$$

(b) Consider the change of variables  $\rho = \sqrt{E}r$  and  $u(\rho) = \sqrt{\rho}R(\rho)$ . We have the relationships:

$$\begin{aligned} R' &= \frac{d}{d\rho} \left( \frac{u}{\sqrt{\rho}} \right) = \frac{u'}{\sqrt{\rho}} - \frac{u}{2\rho^{3/2}} \\ R'' &= \frac{d}{d\rho} \left( \frac{u'}{\sqrt{\rho}} - \frac{u}{2\rho^{3/2}} \right) = \frac{u''}{\sqrt{\rho}} - \frac{u'}{\rho^{3/2}} + \frac{3u}{4\rho^{5/2}} \end{aligned} \quad (48)$$

so that the radial equation becomes:

$$\begin{aligned} \frac{\rho^2}{E} \left( \frac{u''}{\sqrt{\rho}} - \frac{u'}{\rho^{3/2}} + \frac{3u}{4\rho^{5/2}} \right) + 2\frac{\rho}{\sqrt{E}} \left( \frac{u'}{\sqrt{\rho}} - \frac{u}{2\rho^{3/2}} \right) + [\rho^2 - l(l+1)] \frac{u}{\sqrt{\rho}} &= 0 \\ \rho^2 u'' + \rho u' + \left[ \rho^2 - \left( l + \frac{1}{2} \right)^2 \right] u &= 0 \end{aligned} \quad (49)$$

which is the Bessel equation of order  $l + 1/2$ .

(c) Assuming integer  $l$ , the solution to  $R(\rho)$  is the Bessel functions:

$$R(\rho) = \frac{1}{\sqrt{\rho}} \sum_{l=0}^{\infty} A_l J_{l+1/2}(\rho) + B_l J_{-l-1/2}(\rho) \quad (50)$$

The finiteness of  $R(\rho)$  at  $\rho = 0$  demands  $B_l = 0$ . We can now write the general solution as:

$$V(r, \theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l A_{l,m} \frac{J_{l+1/2}(\sqrt{E_l}r)}{(\sqrt{E_l}r)^{1/2}} Y_{lm}(\theta, \phi) \quad (51)$$

We further require  $V(r = a) = 0$  so that:

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l A_{l,m} J_{l+1/2}(\sqrt{E_l}a) = 0 \quad (52)$$

We invoke without proof the Bourget's hypothesis, which says that the Bessel functions (differing by integers) do not share common zeros except at the origin. This means  $A_{a,b} = 0$  except for some  $a = l$  and  $b = m$  such that  $J_{l+1/2}(\sqrt{E_l}a) = 0$ . Thus given some  $E_l$  as the eigenvalue, we can uniquely determine all  $A_{l,m}$ .

The solution is then:

$$V(r, \theta, \phi) = \sum_{m=-l}^l A_l \frac{J_{l+1/2}(\sqrt{E_l}r)}{(\sqrt{E_l}r)^{1/2}} Y_{lm}(\theta, \phi) \quad (53)$$

## 7 Heat equation

(a) The boundary conditions and the eigenfunction relation are trivial to show. To demonstrate orthogonality, consider the inner product of  $q_k$  and  $q_l$ :

$$\begin{aligned} \langle q_k, q_l \rangle &= \frac{2}{a} \int_0^a \sin \left[ \frac{\pi}{a}(k + 1/2)x \right] \sin \left[ \frac{\pi}{a}(l + 1/2)x \right] dx \\ &= \frac{2}{\pi} \int_0^\pi \sin [(k + 1/2)y] \sin [(l + 1/2)y] dy \\ &= \delta_{kl} \end{aligned} \quad (54)$$

where at the second step the substitution  $y = \pi x/a$  is used and the last step follows from the orthogonality of sine functions.

(c) We write the general solution as:

•