

# Chapter 1

## Congruency Theorems in Neutral Geometry

**Note on notation:** For a vertex  $A$  in  $\triangle ABC$ , we may refer to the interior angle  $\angle CAB$  as  $\angle a$ . We denote the length of a segment  $\overline{AB}$  as  $|\overline{AB}|$ . We do not rely on any definition of length except to say that  $|\overline{AB}| = |\overline{CD}| \iff \overline{AB} \cong \overline{CD}$ . Everywhere a statement about equality of lengths is made, it is equivalent to a statement about congruence of line segments or radii of circles, but we have chosen this notation for clarity and consistency with modern geometry. We assume here that angles have a measurement such that they can be related in equality and added. In particular, we assume the cancellation law for measurements of angles under addition, namely  $\angle a + \angle b = \angle a + \angle c \iff \angle b = \angle c$ .

Theorems are indexed  $X.Y$  and constructions are indexed  $X.Y.Z$ .

### 1.1 Side-Angle-Side Congruence

Two triangles are congruent if two sides and the included angle of one are congruent respectively to two sides and the included angle of the other.

*Proof.* Let  $\triangle ABC$  and  $\triangle A'B'C'$  be constructed st.  $\overline{AB} \cong \overline{A'B'}$ ,  $\overline{BC} \cong \overline{B'C'}$ , and  $\angle b \cong \angle b'$ . Move  $\triangle A'B'C'$  so that  $\overline{A'B'}$  coincides with  $\overline{AB}$ , and so that  $C$  and  $C'$  are on the same side of  $\overline{AB}$ . By the definition of congruent angles,  $\overline{BC}$  and  $\overline{B'C'}$ , starting from the same point  $B$  and extending out along  $\angle b$  and  $\angle b'$  respectively, coincide with each other. Since  $\overline{BC} \cong \overline{B'C'}$ , we know they have a common endpoint at  $C$ . Then,  $\overline{AC}$  and  $\overline{A'C'}$  are line segments connecting coinciding points  $A, A'$  to coinciding points  $C, C'$ , so they must also coincide. Since we have now that each of the segments in  $\triangle ABC$  coincides with a segment in  $\triangle A'B'C'$ , the two triangles are congruent.

## 1.2 Angle-Side-Angle Congruence

Two triangles are congruent if two angles and the included side of one are congruent respectively to two angles and the included side of the other.


*Proof.* Let  $\triangle ABC$  and  $\triangle A'B'C'$  be constructed st.  $\angle a \cong \angle a'$ ,  $\overline{AB} \cong \overline{A'B'}$ , and  $\angle b \cong \angle b'$ . Move  $\triangle A'B'C'$  so that  $\overline{A'B'}$  coincides with  $\overline{AB}$ , and so that  $C$  and  $C'$  are on the same side of  $\overline{AB}$ . By the definition of congruent angles,  $\overline{BC}$  and  $\overline{B'C'}$ , starting from the same point  $B$  and extending out along  $\angle b$  and  $\angle b'$  respectively, coincide with each other. Similarly, by the definition of congruent angles,  $\overline{AC}$  and  $\overline{A'C'}$ , starting from the same point  $A$  and extending out along  $\angle a$  and  $\angle a'$  respectively, coincide with each other. By the definition of triangle, we know that  $\overline{AC}$  and  $\overline{BC}$  intersect at point  $C$ . Since  $\overline{A'C'}$  and  $\overline{B'C'}$  coincide with  $\overline{AC}$ ,  $\overline{BC}$  respectively, they must intersect at the point  $C'$  which coincides with  $C$ . Since  $A, B, C$  all coincide with  $A', B', C'$ , the two triangles are congruent.

### 1.2.1 Construction of an Equilateral Triangle

We will construct a triangle with three equal sides on a given base  $\overline{AB}$ .

Draw a circle  $\alpha$  with radius  $|\overline{AB}|$  centered at  $A$ , and a circle  $\beta$  with radius  $|\overline{BA}|$  centered at  $B$  [Euclid Post. 3]. Label a point of intersection of these circles  $C$ . Then  $\triangle ABC$  is equilateral, since all points on  $\alpha$ , including  $C$ , have distance  $|\overline{AB}|$  from  $A$ , so  $|\overline{AC}| = |\overline{AB}|$ . We can conclude with an identical argument  $|\overline{BC}| = |\overline{AB}|$ , thus showing  $\triangle ABC$  is equilateral.


Note that this method of construction allows for two different equilateral triangles, depending on the choice of which intersection of  $\alpha$  and  $\beta$  to use.



### 1.2.2 Construction of a Given Length

Given a line segment  $\overline{AB}$ , and an arbitrary point  $C$ , we will construct a circle of radius  $|\overline{AB}|$  at  $C$ .

Connect  $A$  and  $C$  with a line, and with base  $\overline{AC}$  construct the equilateral triangle  $\triangle ACD$  (the choice of intersection for  $D$  does not matter). Draw a circle,  $\alpha$ , with a center at  $A$  with a radius of  $\overline{AB}$ . Continue the line segments  $\overline{DA}$  and  $\overline{DC}$  indefinitely. Label the point of intersection between the line  $\overline{DA}$  and  $\alpha$   $E$ . Then draw a cir-




cle,  $\beta$  with center  $D$  and radius  $\overline{DE}$ . Label the intersection of  $\overline{DC}$  and  $\beta$   $F$ . Then, we claim  $|\overline{AE}| = |\overline{CF}|$ .  $|\overline{DE}| = |\overline{DF}|$ , as they are both radii of  $\beta$ . We then have  $|\overline{DE}| = |\overline{DA}| + |\overline{AE}| = |\overline{DC}| + |\overline{CF}| = |\overline{DF}|$ . Since  $|\overline{DA}| = |\overline{DC}|$ , as they are both sides of the same equilateral triangle, we get  $|\overline{AE}| = |\overline{CF}|$  from our former equality.

### 1.2.3 Angle Bisection

This will illustrate how to construct a line bisecting a given angle.


Given an angle formed by two rays extending from a point  $A$ , we draw a circle of arbitrary radius [Euclid Post. 3] and label its points of intersection on the two rays  $B$  and  $C$ . Now, with  $\overline{BC}$  as its base, we construct the equilateral triangle  $\triangle BCD$ , choosing point  $D$  as the further one from  $A$ . We then consider  $\triangle ABD$  and  $\triangle ACD$ . They share the side  $\overline{AD}$ . We have that  $\overline{BD} \cong \overline{CD}$  as they are both part of the same equilateral triangle. Finally,  $\overline{BC} \cong \overline{AC}$  as they are radii of the same circle centered at  $A$ . By [1.3],  $\triangle ABD \cong \triangle ACD$ . From this we have  $\angle BAD \cong \angle DAC$ , meaning  $\overline{AD}$  bisects  $\angle BAC$ .



## 1.3 Isosceles Angle Congruence


The angles opposite the congruent sides of an isosceles triangle are congruent.

*Proof.* Let  $\triangle ABC$  be isosceles, and let  $\overline{AB} \cong \overline{AC}$ . Construct the angle bisector of  $\angle A$  [1.2.3] and extend it to intersect with  $\overline{BC}$ , labeling the point of intersection  $D$ . Consider  $\triangle ADC$  and  $\triangle BDC$ . We know since  $\overline{AD}$  is the angle bisector of  $\angle A$ , so  $\angle BAD \cong \angle CAD$ . The triangles have  $\overline{AD}$  in common, and have  $\overline{AB} \cong \overline{AC}$ . By [1.1],  $\triangle ADC \cong \triangle BDC$ , giving that  $\angle ABC \cong \angle ACB$ .



## 1.4 Isosceles Side Congruence

If two angles of a triangle are congruent, the sides opposite those angles are congruent (and so the triangle is isosceles). Note that this is the converse of [1.3].




*Proof.* Let  $\triangle ABC$  have  $\angle b \cong \angle c$ . With base  $\overline{BC}$  we construct the equilateral triangle  $\triangle BCD$ , choosing  $D$  as the further point from  $A$  (see the note to [1.2.1]). Draw the line segment  $\overline{AD}$ . Consider the triangles  $\triangle ABD$  and  $\triangle ACD$ . These have  $\overline{AD}$  in common, have  $\angle ADB \cong \angle ADC$  since they are the angles to an equilateral triangle, and have  $\overline{BD} \cong \overline{DC}$ , since these are the sides of an equilateral triangle. By [1.1] we have  $\triangle ABD \cong \triangle ACD$ , which gives  $\overline{AB} \cong \overline{AC}$ .

## 1.5 Side-Side-Side Congruence

Two triangles are congruent if the three sides of one are congruent respectively to the three sides of the other.

*Proof.* Let  $\triangle ABC$  and  $\triangle A'C'D$  have  $|\overline{AC}| = |\overline{A'C'}|$ ,  $|\overline{AB}| = |\overline{A'D}|$ , and  $|\overline{BC}| = |\overline{DC'}|$ . Move  $\triangle A'C'D$  so that  $\overline{A'C'}$  coincides with  $\overline{AC}$ , and so that  $B$  and  $D$  are on different sides of  $\overline{AC}$ . Draw  $\overline{BD}$ . Now,  $\triangle ABD$  is isosceles, and so  $|\angle ABD| = |\angle ADB|$ .  $\triangle CBD$  is also isosceles, and so  $|\angle CBD| = |\angle CDB|$ . Since  $|\angle ABC| = |\angle ABD| + |\angle CBD|$ , and  $|\angle ADC| = |\angle ADB| + |\angle CDB|$ , substituting our previous angle equalities gives  $\angle ABC \cong \angle ADC$ . Then we have  $\triangle ABC \cong \triangle A'C'D$  by [1.1].




### Equilateral triangles have equal angles

*Proof.* This is a corollary of [1.5], but it is worth noting as it will be used later.

### 1.5.1 Construction of a Perpendicular Line off a Given Point

Given a line segment  $\overline{AB}$ , and a point  $C$  on  $\overline{AB}$ , we will construct a line extending from  $C$  that is perpendicular to  $\overline{AB}$  (makes right angles with  $\overline{AB}$ ).

Centered at  $C$ , draw the circle  $\alpha$  with the smaller of  $|\overline{AC}|$ ,  $|\overline{CB}|$  as its radius. WLOG, we will assume it is of radius  $|\overline{AC}|$ . Let  $D$  be the point of intersection of  $\alpha$  with  $\overline{AB}$  that is not  $A$ . Then, draw an equilateral triangle  $\triangle ADE$  with base  $|\overline{AD}|$ . Construct the line  $|\overline{EC}|$ . We claim that




$\overrightarrow{EC}$  is perpendicular to  $\overline{AB}$ .

Clearly  $\angle ACE$  and  $\angle DCE$  are supplementary, and [1.5] shows that  $\triangle ACE$  and  $\triangle DCE$  are congruent. Since  $\angle ACE$  and  $\angle DCE$  are supplementary angles congruent to each other, they are right by definition.

## 1.6 Supplementary Angle Congruence

Supplementary angles are congruent to two right angles. Recall that supplementary angles are the angles formed by the intersection of two lines which share a common line segment.



*Proof.* Consider the supplementary angles  $\angle AFD$  and  $\angle DFB$  formed by the intersection of  $\overline{AB}$  with  $\overline{CD}$ , labeling their point of intersection  $F$ . From  $F$  construct a line perpendicular to  $\overline{AB}$  [1.5.1]. Label a point on this line  $E$ . If  $\angle AFD \cong \angle AFE$ , the proof is trivial, so we will assume that  $\overline{FE}$  and  $\overline{FD}$  do not coincide, WLOG having  $\angle AFE$  within  $\angle AFD$ . Then,  $\angle AFD = \angle AFE + \angle EFD$ , and  $\angle EFB = \angle EFD + \angle DFB$ . Adding these equalities gives:

$$\angle AFD + \angle EFD + \angle DFB = \angle AFE + \angle EFD + \angle EFB$$


Which may simplify to

$$\angle AFD + \angle DFB = \angle AFE + \angle EFB$$

Which is exactly our theorem, as  $\angle AFE + \angle EFB$  are two right angles.

## 1.7 Vertical Angle Congruence

Vertical angles are congruent. Recall that vertical angles (also called opposite angles) are the angles formed by the intersection of two lines which do not share a common line segment.



*Proof.* Consider the vertical angles  $\angle AEC$  and  $\angle DEB$  formed by the intersection of  $\overline{AB}$  with  $\overline{CD}$ , labeling their point of intersection  $E$ . Let  $\alpha$  represent the measure of two right angles. By [1.6], we have  $\angle AEC + \angle AED = \alpha$ , and by [1.6] we have  $\angle DEB + \angle AED = \alpha$ . From this we have  $\angle AEC + \angle AED = \angle DEB + \angle AED$ , and applying the cancellation property gives  $\angle AEC = \angle DEB$ .